A descriptive, diagnostic and practical essay on disorders of the digestive organs and general health, and particularly on their numerous forms and complications, contrasted with some acute and insidious diseases: being an attempt to prosecute the views of Dr. Hamilton and Mr. Abernethy, and a second edition of the essay On the mimoses, with additions / by Marshall Hall.

Contributors

Hall, Marshall, 1790-1857. Hall, Marshall, 1790-1857. On the mimoses. University of Leeds. Library

Publication/Creation

London: Longman, Hurst, Rees, Orme, and Brown, 1820.

Persistent URL

https://wellcomecollection.org/works/cfh5z7ur

Provider

Leeds University Archive

License and attribution

This material has been provided by This material has been provided by The University of Leeds Library. The original may be consulted at The University of Leeds Library. where the originals may be consulted.

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org AN

ESSAY

ON


DISORDERS

OF THE

DIGESTIVE ORGANS AND GENERAL HEALTH,

AND ON THEIR

Complications.



DESCRIPTIVE, DIAGNOSTIC AND PRACTICAL

ESSAY



ON

DISORDERS

OF THE

DIGESTIVE ORGANS AND GENERAL HEALTH,

AND PARTICULARLY ON THEIR NUMEROUS

Forms and Complications,

CONTRASTED WITH SOME

ACUTE AND INSIDIOUS DISEASES;

BEING AN ATTEMPT TO PROSECUTE THE VIEWS OF DR. HAMILTON AND MR. ABERNETHY, AND A

SECOND EDITION

OF THE ESSAY ON THE MIMOSES,
WITH ADDITIONS.



RY

MARSHALL HALL, M. D. F. R. S. E.

&c. &c.

LONDON:

PRINTED FOR LONGMAN, HURST, REES, ORME, AND BROWN.

PATER-NOSTER ROW.

1820.

W. Wheelhouse, Printer, Nottingham.

MATTHEW BAILLIE, M.D.

F. R. SS. L. AND E.

PHYSICIAN EXTRAORDINARY TO THE KING,

ETC. ETC. ETC.

SIR,

Your indulgent expression of approbation of my former Work, and your condescending acceptance of the dedication of the present one, constitute at once the most ample reward and the greatest encouragement of my efforts to contribute to the advancement of Diagnosis and the History of Diseases.

With every sentiment which the most profound respect can inspire,

I remain, SIR,

Your most obliged Servant,

MARSHALL HALL.

Nottingham, October 21st, 1820.

MATTHEW BALLURE M. D.

Kinne I am & A

ONLY SELLO BELLOURS NAMED AND ASSESSED OF THE WIND

and me and

the second of the continues of the second of the continues of the continues of the continues of the continues of the second of t

because the other properties distinct the distance of the ASPA

Andres and

Transin Sun

the new years around the seasons of

TREEL TRANSPORT

Spiralphon, Orgolas Stat. 2008.

ON THE MIMOSES.

CHAPTER I.

ON THE MIMOSIS ACUTA.

SECTION I. DESCRIPTION, WITH CASES.

The General Character,—of the Severe, and Less } 24, 39
Lists of Complications,
Description,
Cases,
History,
Weight of Patients,p. 37, a note

SECRETORY II
SECTION II. THE DIAGNOSIS, WITH CASES.
I. The Diagnosis of the General Affection,
1. From Fevers,
2. From Organic Disease,
3. From Strumous Disease of the Mesentery in
3. From Strumous Disease of the Mesentery in Adults,
4. From an Effect of Drunkenness,
II. The Diagnosis of the Complications,
1. With Affection of the HEAD,
1. of an Apoplectic,
2. of an Epileptic character
3. Pain and some other Affections,89-94
2. With Affection of the Chest,
1. Oppressive Dyspnæa,96—98
2. Cough with Expectoration,99—105
3. With Affection of the Heart,
1. Fluttering, Palpitation, irregular action,
2. Fainting,
4. With Affections of the Stomach and Bowels
1. Hiccough,
2. Hæmatemesis, Melæna,
3. Pain,
4. Constipation, Diarrhæa,126—127
5. With Icterus,
6. With Symptoms of affection of the Bladder, 132
7. With Pain resembling the Tic Douloureux,133
8. With Convulsive and Spasmodic Affections 135—144
9. With Paralytic Weakness,
10. With a singular Tumidity of the Face and General Surface
General Surface
III. The Local Complications,
1. An Arid and Exfoliating state of the Prolabia,151 2. An Ulcerated state of the Tongue,152
2. All officiated state of the Torone,
IV. A Cachectic Form of Mimosis,

SECTION III. THE CAUSES AND TREATMENT.
1. Of the General Affection,
2. Of the Complications,
OF THE PERSON OF
CITA DEPUT IT
CHAPTER II.
ON THE MIMOSIS CHRONICA.
SECTION I. THE DESCRIPTION.
The General Character,185
The Description,
The Description,
SECTION II. THE DIAGNOSIS.
1. From the Less Severe Form of the Mimosis Acuta, 196
2. From Insidious Organic Disease,
SECTION III. THE CAUSES AND TREATMENT, 198-203
and the Producted Plans of the stop.
CITA DIED TIX
CHAPTER III.
ON THE MIMOSIS DECOLOR.
ON THE MIMOSIS DECOLOR.
SECTION I. THE DESCRIPTION, WITH CASES.
The General Character,
Description of the Inci-
pient, Confirmed, and 208-214; 215-222; 223, 250 Inveterate Forms or
Inveterate Forms or (203-214, 213-222, 223, 230
Stages,
And of the Chronic Form,
Of some Varieties,
As it occurs in Married Persons,
And in the Male Sex,
List of Complications,
9

Pathological Remark	28
Cases	
History,	
Remedies,	35
Dissection,	
AGE,	
SECTION II. THE DIAGNOSIS, WITH CASES.	
I. The Diagnosis of the General Affection,	
1. From Insidious Organic Disease,	38
2. From Disease of the LIVER,	
Park College College Contract of Contract	
II. The Diagnosis of the Complications,	
1. With Sudden Affection of the HEAD,271-27	
2. With Continued Affection of the HEAD, 276-27	
3. With Sudden or Continued Cough and Dyspnæa,	30
Dyspnœa,	
4. With Palpitation of the Heart,	31
With Sudden Pain and Tenderness of the Side, 282, 28	33
With Protracted Pain of the Side,284, 28	
With Sudden, or Protracted, Pain of the Abdomen,	88
With Melæna,	
With Menorrhagia and Leucorrhæa,	
SECTION III. THE CAUSES AND TREATMENT.	
The Course Character of the Character of	
CHAP. IV.	
nee to a representative and the second discountry of	
ON THE MIMOSIS URGENS.	
SECTION I. THE DESCRIPTION.	
The General Character,	24
Of the Mild, Severe, and Inveterate Forms, 310; 311—322, 329; 323, 33	30
Copious List of the other Forms,32	25

SECTION II. THE DIAGNOSIS.

	1. Of the Convulsion,	3
	2. Of the Affections of the Head,—Pain, Deli- rium, Stupor, &c	R
	rium, Stupor, &c	0
	3. Of Pain of the Chest,	9
	4. Of Urgent Dyspnæa,34	0
	5. Of Fits of Coughing34	1
	6. Of Suspended Respiration,34	3
	7. Of a painful Affection of the Diaphragm,344-34	
	8. Of the Imitation of CROUP,	7
	9. Of the Palpitation or Syncope,34	
	10. Of the Pain of the Abdomen,349, 35	
	11. Of the Hiccough or Retching,	1
	12. Of the Dysury,	2
	13. Of the Partial Paralysis,	
	14. Of the Trismus, Tetanus, Clenched Hand, Contracted Foot, Twisted Legs, &c	
	Contracted Foot, Twisted Legs, &c	00
	Remark,35	7
S	ECTION III. THE CAUSES AND TREATMENT, 358-36	1

POSTSCRIPT.

ON THE MIMOSIS INQUIETA.

ADVERTISEMENT.

I have changed the Title-page of this little work, but have retained the term Mimosis in the Text. The reader will adopt or reject it as he deems most proper. I still think it the most appropriate in itself and the most useful in its practical tendency.

The present Essay describes the various affections of the General Health and of different Parts, which arise from the more continued action of Derangement of the Digestive Organs. The Essay alluded to in the Postscript, p. 183, displays the more acute and alarming effects of Intestinal Load and Irritation, and of Exhaustion from Loss of Blood, or other Causes. These two Classes of Morbid Affections are, as the reader will observe, allied in many respects.

M. H.

MIMOSES.

- 1. THERE is a Class of Disorders, each of which consists of a more GENERAL morbid affection, usually combined with some TOPICAL symptom or symptoms.
- 2. The general affection is complex and various; the complications are multiform and changeable, and, by their incidental predominance, frequently imitate other diseases widely different in their nature.
- 3. These affections have been variously and perhaps too exclusively attributed, by some authors, to a state of derangement in one or more of the chylopoetic viscera; and by others, to an unequal and undue distribution of the blood, by which a state of arterial excitement or of venous congestion is induced

in some particular organ, or in some particular part of the sanguiferous system. I have scarcely ventured, in this work, to enter into any speculation relative to the pathology of the affections of which it treats: for this part of medicine, notwithstanding the ingenuity of some late theorists, seems scarcely to have advanced from the state of conjecture and uncertainty described by Celsus, whose words* are still, in every sense, but too admissible.

- 4. As the real nature and connexion of the general and topical affections in these complaints, may thus frequently be dubious, I have deemed it advisable to appropriate some new term, which might, without implying any opinion on the subject, sufficiently express a prominent and important feature of this class of morbid affections. The denomination MIMOSIS, from the Greek word µ1µ105, imitator, will at once denote a remarkable peculiarity,—the multiform character,—of these disorders, and will serve to impress the mind with the necessity of distinguishing, in
- * Cum hæc per multa volumina, perque magnæ contentionis disputationes, a medicis sæpe tractata sint atque tractentur; subjiciendum est, quæ proxima vero videri possint. Ea neque addicta alterutri opinioni sunt, neque ab utraque nimium abhorrentia; media quodammodo inter diversas sententias: quod in plurimis contentionibus deprehendere licet, sine ambitione verum scrutantibus, ut in hac ipsa re. Nam quæ demum causæ, vel secundam valetudinem praestent, vel morbos excitent, ne sapientiae quidem professores scientia comprehendunt, sed conjectura persequuntur. Cujus autem rei non est certa notitia, ejus opinio certum reperire remedium non potest. Verumque est, ad ipsam curandi rationem nihil plus conferre, quam experientiam.—Celsi Præf.

Local Affections, between those which belong to the present Class, and others which are either primary, or have a different origin,—a point of great practical importance.

- 5. For a similar reason, I have discarded the terms bilious, spasmodic, &c. as denominations for diseases; and have reserved them only to denote certain symptoms of morbid affections. In the latter sense, their import is generally understood, and may, perhaps, be sufficiently definite; but in the former, they could only serve to satisfy the mind with vague conceptions of the affection, and to check the investigation of its particular and individual nature.*
- 6. Of the Mimoses I have observed FIVE FORMS;
 the Mimosis Acuta, the Mimosis Chronica, the
 Mimosis Decolor, the Mimosis Urgens, and the Mimosis Inquieta.
- I. The Mimosis Acuta appears to me not to have been before fully described in medical writings, unless this affection be, as I suspect, identical with the Scorbutus of WILLIS and other writers of his day.†
- II. The Mimosis Chronica is the Dyspepsia or Hypochondriasis of medical authors.

^{*} See the author's Treatise on Diagnosis, pp. x, 2-3.

[†] See the Edinb. Med. and Surg. Journal, Vol 16, p. 204; on the acceptation of the Term Scoreutus.

- III. The Mimosis Decolor embraces the Chlorosis of medical writers, together with some other morbid affections, in which the complexion and general surface suffer materially, and which have not, I think, been hitherto sufficiently, if at all, distinctly described.
- IV. The Mimosis Urgens is the Hysteria of authors. Its numerous forms are, I think, more amply collected and described in this than in any former publication.
- V. The Mimosis Inquieta is, I believe, described in this work for the first time. It appears, in general, to combine the effects of intestinal irritation and of exhaustion from loss of blood or other causes.*
- 7. The COMPLICATIONS of the Mimoses are apt to be mistaken and mistreated for different Inflammatory and other Local diseases, and appear to me to constitute a Class of morbid affections scarcely less frequent or less important, and requiring to be distinguished with the utmost care. For an enumeration of these complications the reader is referred to the table of Contents.
- 8. The object of this Essay is to establish this Class of general and local morbid affections more dis-
- * Since the appearance of the first edition of this Essay, I have published a separate tract on this subject, with the following title:—Cases of a Serious Morbid Affection, chiefly occurring after Delivery, Miscarriage, &c. from various causes of Irritation and Exhaustion; and of a Similar Affection, unconnected with the Puerperal State.

tinctly and extensively than before,—to collect and embody the system of facts which belong to this part of pathology,—to present accurate **DESCRIPTIONS** of the different forms, and to trace the **DIAGNOSIS** of the numerous **COMPLICATIONS** of these disorders.

- 9. I have thus been very explicit as to the design and pretensions of this little Essay. With regard to the NATURE of these affections, I beg to observe that I consider them all as similar. The Mimosis Acuta differs from the Mimosis Chronica, in affecting the process of assimilation, the strength, and the general health, more seriously. The Mimosis Decolor presents some extraordinary states of the complexion, the general surface, and the capillary circulation, the cause remaining probably similar. The Mimosis Urgens is not necessarily connected with any of these affections of the assimilation or of the surface, and yet the cause may still be, in a general sense, the same. The Mimosis Inquieta conjoins, with this cause, the state of exhaustion.
- 10. Too much praise cannot be conferred on those members of the profession, who have so well elucidated the nature and treatment of some of the subjects of the preceding pages. There is no doubt, indeed, that this investigation of the Mimoses was suggested to me by what I have learnt from the invaluable labours of Dr. Hamilton, Mr. Abernethy, and other respectable writers. My situation in Nottingham, how-

ever, has been a principal cause of fixing my attention on a Class of Disorders, of which the usual causes are sedentariness and confinement. This town, so celebrated for its manufactories of cotton stockings and lace, embraces a very extensive population, a great majority of which,—men, women, and children,—are engaged from morning till evening, in the numerous sedentary occupations which these manufactories imply,—deprived of the salutary influence of pure air and gentle exercise.

- 11. To these numbers, which are peculiar to my situation, must be added the sedentary amongst the remaining part of the population,—the literary, persons of a delicate mode of life, females in general, tailors, mantua-makers, and the youthful inhabitants of the schools.
- 12. Nor is sedentariness the only cause of the Mimoses, the operation of which is peculiarly frequent amongst the poor of Nottingham. I have noticed the frequent occurrence of the Mimosis Decolor in cooks and housemaids; and the same remark applies equally to those persons who are much engaged in 'ironing,' and, of consequence, much confined to an atmosphere overheated by stoves for the purpose of quickly drying the articles subjected to this process.
- 13. To this view of the causes of the Mimoses peculiar to a manufacturing town, may be added the

baneful influence of a confined and impure atmosphere, and an indigestible and poor diet,—an influence which we learn to estimate more perfectly, by adverting to the impediment they afford to our attempts to cure, and to the beneficial effects of the country air and exercise, with a proper and nutritious diet.

- 14. A certain activity of the body would appear to be necessary to insure the peristaltic movements of the intestines, and, in consequence, the propulsion of their contents. During sedentariness, these movements are probably retarded, the alvine evacuation becomes more scanty or less frequent, and the intestines remain loaded.
- 15. From this loaded state of the bowels, their functions, and those of ALL the chylopoetic viscera, most probably become deranged. The alvine contents become disordered merely by delay; and their presence induces in its turn, a disordered state of the functions,—or actions,—of all the organs contributory to digestion, and at length of other organs more remotely situated in the animal frame.
- 16. The functions of the parts within the MOUTH become obviously disordered. The secretions become morbid; the tongue becomes loaded and swollen; the gums red and tumid; the breath tainted; and the saliva sometimes profuse and offensive. The complexion and the skin become morbid, and there

or the Mimosis Decolor, and frequently ædema. This condition of the complexion and skin varies with the state of the original disorder, and with that of the tongue and internal mouth, of which it affords indeed an INDEX. With the state of the mouth and skin, that of the secretions and other functions of the whole course of the ALIMENTARY CANAL and the contributory digestive organs,—the LIVER, the PANCREAS, &c. may be presumed to be all morbidly affected. Digestion is variously deranged; the contents of the bowels become unnatural; and thus reciprocally. According to the state of things, nutrition is impaired, or the sensations are uneasy and painful.

- 17. To term these disorders stomachic, intestinal, hepatic, or bilious, would alike afford partial and inadequate views of this comprehensive subject. As co-existent or subsequent links of this chain of sympathies, the functions of the BRAIN, HEART, RESPIRATION, STOMACH, INTESTINES, UTERUS, BLADDER, ETC. become variously affected. The MUSCULAR SYSTEM and the SENSES also suffer in different instances. And nutrition, absorption, and secretion are impeded or impaired.
- 18. From this view of the subject, the character of the Mimoses may be deduced. And the recurrence of this word, leads me once more to apologize for the introduction of a new denomination for these

diseases. I have been induced to adopt this term, first, to prevent a great deal of circumlocution; and, secondly, because I could find no other in use, which was not objectionable from implying some hypothetical view of the subject. These motives, I still trust, will appear sufficient to justify the innovation. I can at least assert in the words of Morgagni,—longe mihi potior cura est veritatis quam novitatis.

- 19. It was originally intended to publish the following Essay in a larger form, accompanied by representations of the Complexion, Tongue, Tinge of Surface and of the Hands. It is now found necessary, however, to leave the task of procuring plates to some one more fortunately situated, or, at least, to a subsequent period. In the mean time, the text, it is hoped, will be found a faithful portrait from Nature, not unacceptable to the reader of practical medicine.
- 20. Before I conclude these few preliminary remarks, I may be allowed to observe, that there is in the profession a two-fold prejudice, respecting medical investigations:—whilst one class of practitioners seem inclined to refer too many morbid affections to a deranged state of the chylopoetic organs, another,—'contrarium errorem errantes,'—turn their attention too exclusively to such diseases as leave traces under the scalpel of the anatomist. Both parties appear to be in the wrong; each of these topics deserves and demands its due share of attention. The important study

of DIAGNOSIS seems to be that by which alone we may be safely piloted through these opposite dangers.—I trust the following pages will serve to illustrate the force and truth of this remark.

CHAP. I.

THE MIMOSIS ACUTA.

SECTION I. DESCRIPTION, WITH CASES

- 21. FROM the great diversity and complication of the symptoms of this form of Mimosis, and from the occasional prevalence of one particular symptom over the rest, it is probable that it has sometimes been mistaken for some other affections. With the view of assisting the distinction of this case in future, it will be my object, first to detail the symptoms which characterize the complaint in general, and, in the second place, to enumerate those particular symptoms which are apt, in certain instances, to engross the attention of the patient and of his friends, and even to occasion some difficulty and embarrasment in the diagnosis, to the medical practitioner.
- 22. In order to facilitate the description, the Mimosis Acuta will be considered under two forms, differing only in degree,—in the greater or less severity of its symptoms in general,—and in the presence or absence of some of the severer symptoms. These two forms, of course, admit of all intermediate shades;

they are also, each, apt to be continued in a protracted form.

- 23. The Mimosis Acuta, even in its severer form, comes on insidiously, and the patient gradually becomes incapacitated for business or his usual exercise: the less severe form occurs more gradually and insidiously still.
- 24. The severer form of the Mimosis Acuta is early and principally characterized and distinguished by the concurrence of the following symptoms,—namely, Weakness, Tremor, Headach, Vertigo, Fluttering, Faintishness, Tendency to Perspiration, susceptibility to Hurry and Agitation, Weariness, and loss of flesh.
- 25. The countenance is rather pale and thin; the lips are pale, and, with the chin, frequently tremulous, especially on speaking; the surface of the face is generally affected with an appearance of oily, clammy, and swarthy perspiration, especially near the nose; there is a loss of colour, and usually a degree of sallowness and darkness of the complexion in general, but principally about the eyes. The face is often bloated, and the skin coarse, at first; afterwards, there is some degree of emaciation.
- 26. The tongue is almost invariably loaded:—sometimes only slightly, whilst its edges are clean

and red; -in severe cases, a load has formed over the tongue, and has, almost at once, peeled off, leaving the surface morbidly red, smooth, and tender; -- at other times it is more loaded, swollen, and ædematous, formed into deep sulci or plaits, and marked by pressure against the contiguous teeth,—the inside of the cheeks being also impressed in the same manner; the papillæ of the tongue are numerous and enlarged; the gums are red and swollen, and occasionally bleed; the teeth and the mouth are in general foul, and the breath loaded and fœtid; in a fourth instance, the tongue may, however, be clean, but lobulated, whilst the internal mouth and breath are little affected. The first and second states of the tongue are observed, when the affection has not continued long; the third, when its accession has been particularly slow and gradual; and the fourth, when a similar but chronic state of disorder has long subsisted, and has, at length, been succeeded by the Mimosis Acuta.

- 27. There is a tendency to perspiration, on slight exertion, or any surprise, and, sometimes, in the night or early in the morning; the skin is, in general, cool, rather moist, and clammy; in some protracted cases, it has become dry and harsh. The hands and feet are apt to be very cold, and the nails occasionally assume a lilac hue.
- 28. The patient is usually affected with great tremor, observed sometimes in a quivering of the lip, or

dimpling of the chin, but more usually, on holding out the hand, or in carrying a cup of tea, for instance to the mouth, on attempting to stand erect or walk, or on being fatigued or hurried. The tremor, in some protracted cases, has formed the most remarkable feature of the affection; in others, it has been much less observed, but it is rarely, if ever, entirely absent. The patient is liable to experience faintishness in the upright position, if sustained for a little time. And he feels unaccountably feeble and weary, and suffers from a sense of aching, after slight exertion.

- 29. There is an early and daily loss of flesh. This, as well as the restoration of flesh, during recovery, may be ascertained by WEIGHING, as exemplified in some of the cases of this affection, to be given hereafter.
- 30. The patient experiences head-ach and vertigo, and he is nervous, and easily hurried and agitated. There is sometimes heaviness for sleep; sometimes great wakefulness and restlessness; sometimes incubus, rarely delirium; sometimes loss of memory and absence of mind. There is almost universally a peculiar sense of fluttering about the heart and pit of the stomach. And there is frequently an acute pain in some part of the course of the colon.
- 31. The pulse is generally nearly natural, but it is often rather frequent, and usually easily accelerat-

- ed. There are often great irregularity and intermittence of the pulse.
- 32. The appetite is generally much impaired, and there is often loathing of food; but sometimes the appetite is even greater than natural, and there is almost constant craving. The digestion is various, being sometimes quick, and at other times attended with great sense of load, distention, flatus, eructation, hiccough, and even rumination or vomiting of food. The bowels are at first constipated; afterwards constipation and diarrhæa alternate, and sometimes the latter symptom becomes nearly permanent: the motions, during the constipation, are small, during the diarrhæa, scanty, extremely fætid, dark coloured, often accompanied by blood, and frequently attended by tenesmus.
- 33. The urine is extremely loaded in the commencement of the Mimosis Acuta, but may become perfectly transparent during its continuance; it is often high coloured; and like the other symptoms the appearance of this secretion is very liable to change. A continued state of Mimosis seems to lead to the formation of gravel.
- 34. Besides the symptoms just enumerated, there are others which prevail more or less in almost every case; but they are, on the whole, less constant and more diversified; and of these one sometimes predo-

minates so much over the rest, as to engross the attention of the patient and sometimes of the practitioner, too exclusively. The secondary affection is then considered as idiopathic, and the symptom is apt to be treated as the disease. It is therefore of the utmost importance to present the reader with the following distinct enumeration of these symptoms:—

- 1. HEADACH; VERTIGO; STUPOR; ETC.
- 2. COUGH; VISCID EXPECTORATION.
- 3. PAROXYSMS OF OPPRESSIVE DYSPNŒA.
- 4. PALPITATION OF THE HEART; FLUTTERING; FAINTISHNESS; IRREGULARITY AND FREQUENCY OF THE PULSE.
- 5. FREQUENT AND VIOLENT HICCOUGH; VOMITING OF FOOD.
- 6. SOME CONVULSIVE AND SPASMODIC AFFECTIONS.
- 7. PAIN IN THE EPIGASTRIC, OR ONE OR BOTH OF THE HYPOCHON-DRIAC, OR CHONDILIAC REGIONS.
- 8. CONSTIPATION; DIARRHŒA; TENESMUS.
- 9. HÆMATEMESIS; MELÆNA.
- 10. ICTERUS.
- 11. SEVERE PAINS OF SOME OF THE LIMBS.
- 12. SUDDEN TUMEFACTION OF THE INTEGUMENTS, ESPECIALLY OF THE FACE.*
- 35. The diagnosis in these cases will be attempted in the section appropriated to this subject. At present it is of importance to remark that in most instances, even where one of these symptoms is particularly marked and severe, several concur, and are experienced in a mitigated form, affording a characteristic feature of this disorder and a principal source of dis-

^{*} See Section II. and the Table of Contents.

crimination; for whilst most local diseases are denoted by being simple, and definite, this affection is distinguished by its multiplicity, and by apparently conjoining many or all disorders in one,—Ουχ ΈΝ τι των κακων φαινεται, αλλ' ετιν ότε ΠΟΛΛΑ, η και ΠΑΝΤΑ.

- 36. This form of Mimosis is also characterized, although less so perhaps than the more chronic and continued forms of this affection to be described hereafter, by being variable,—better and worse,—with this or that prevailing feeling or symptom,—even during a general recovery;—changes chiefly induced by bodily fatigue, mental agitation, errors in diet, or constipation.
- 37. It is also to be remarked that, although the usual accession of this affection is insidious, as stated § 23, yet when the causes* inducing it are severe, or the patient subject or predisposed to this affection, it may appear in a still more acute form; and in this case it assumes, in some instances, even the character of a Febrile disorder; and in others, that of a serious morbid affection, soon depriving the patient both of strength and of flesh in an alarming degree.
- 38. By an attention to the circumstances detailed § 34, and by a reference to the general symptoms of the Mimosis Acuta, §§ 24—33, the different se-

condary affections may be generally distinguished and identified, in their connexion with the primary disorder. It should, however, ever be borne in mind that an idiopathic and organic affection of some part may co-exist with the Mimosis Acuta. The diagnosis is then often very difficult. It is frequently only ascertained when the state of disorder constituting Mimosis, is removed. The prognosis should therefore be given with caution.—Another circumstance suggesting caution in the prognosis, is the possibility of the transition of the state of DISORDER into that of DISEASE, which is not unusual in those parts and organs which are, at first, only affected in a secondary and functional manner.

- 39. In the less severe form of the Mimosis Acuta, the debility, tremor, loss of flesh, and tendency to faintishness and perspiration, § 24, are less observed, although perhaps not altogether absent.
- 40. In this form of the Mimosis Acuta the countenance is rather sallow, and its surface is more or less affected as in the severer form described above. The tongue and the internal mouth are often affected in the severer degree described § 26. The patient is incapable of pursuing any laborious employment. He is prone to perspire from slight exertion or agitation. He perhaps experiences some loss of flesh. He is low spirited and listless. The appetite is sometimes impaired, but sometimes craving. And he suffers from

the symptoms described §§ 30—33, and from the complications enumerated § 34, only in a milder and more protracted form than the subject of the severer cases of Mimosis Acuta.

- 41. Besides the symptoms enumerated § 34, the less severe but more continued form of the Mimosis Acuta, is sometimes attended with one of the following affections:—
 - 1. FURUNCULI; PARONYCHIA; HORDEOLA.
 - 2. ERYSIPELAS; ERYTHEMA NODOSUM; URTICARIA CHRONICA; LICHEN.
 - 3. PURPURA.
 - 4. ULCERATIONS AND PUSTULES OF THE CONJUNCTIVA.
 - 5. DECAY OF THE TEETH; A MORBID STATE OF THE GUMS; CHRONIC SORE THOAT.
 - 6. SOME AFFECTIONS RESEMBLING SYPHILIS.*
- 42. Persons who have long laboured under a state of Mimosis appear liable to decline, at length, into Phthisis Pulmonalis, or to become the subjects of other organic disease, and especially of the Liver.

^{*} Sce Section II. and the Table of Contents.

- 43. IT may now be proper to confirm the description of the Mimosis Acuta just given, by a selection from the CASES from which it has been taken:—
- 44. Case I. The first case of the Mimosis Acuta which I was enabled to distinguish, was that of Mr. M. aged 25.—He had, as I understood, been treated for Fever, principally by the pulvis antimonialis, for several weeks.—He was, on my first visit, affected with tremor, debility, tendency to perspiration, and had suffered a considerable loss of flesh. The tongue was affected with large papillæ, indented, and much loaded; the breath was extremely fætid. The pulse was frequent. He complained of pain in the left hypochondre.—These complaints were soon removed by a course of gentle purgatives.—This patient was employed in the sedentary occupation of the lace-frame.
- 45. Case II. The second case occurred soon after the former, and was noted rather more particularly.—Mr. S. aged 25, had been indisposed several weeks when I first saw him, and had left Nottingham for the country; he had taken saline powders principally, and I therefore judge that his complaint had been deemed Fever; it had made very considerable progress in inducing a loss of flesh and strength. The countenance, especially the lips and chin, was affected with a tremulous movement on speaking; there was

an expression of great debility; the prolabium of each lip was palish. He was scarcely able to walk or stand erect, in attempting which he tottered and trembled; and there was a manifestation of great feebleness in the general manner. The skin was moist. tongue was affected with enlarged papillæ, much loaded, and indented; the mouth in general was very disagreeable, and the breath tainted and fœtid. pulse was frequent, and easily and greatly accelerated. He was extremely nervous, and easily agitated. He had no appetite. The bowels were costive, with scanty and offensive stools. The urine deposited a most copious sediment.—The friends of this patient now suspected an attack of Consumption.—The affection yielded very soon to the administration of purgative medicines.

46. Case III. J. S. aged 30. He had been ill during three months before his application to me, having been affected at first with constipation of the bowels, which yielded to a state of diarrhea, with discharges of blood by stool. He had suffered much loss of flesh and strength. I made the following short note on first seeing this patient;—the countenance is rather pale, with a little appearance of oily or clammy perspiration; the tongue is much loaded; he is much thinner than when in health; feeble; tremulous on holding out the hand; he becomes faint when standing upright; he is nervous and easily flurried and agitated; he has suffered from headach and vertigo;

he has experienced much sense of fluttering in the region of the heart; he has no cough, and has not had hiccough.—He became soon better, in every respect, from taking the pil. hydrag. and magnesia with rhubarb and pimento.

- 47. Case IV. Mrs F. aged 24. She was married five months ago and supposes herself pregnant. She began to be indisposed seven weeks ago, becoming affected with sickness, load and wind at the stomach, hiccough, lowness, weakness, and loss of flesh .- The following note was taken on visiting this patient:-The countenance is rather pale and the eye-lids dark. She has lost much flesh. Complains of weakness. She is easily hurried by surprise, and she then complains of a sense of fluttering at the pit of the stomach. There is no sweating. She suffers from headach sometimes. She is sleepless during the night. She sighs frequently. There is no cough. Much hiccough, a sense of load and wind at the stomach, eructation, and sometimes sickness and vomiting. The appetite is very bad. The bowels costive. The catamenia are suppressed and the mammæ enlarged .- This patient proved to be pregnant.—She recovered slowly and gradually, from the use of purgative medicines,
- 48. The two following cases present the appearances in the Mimosis Acuta, in its less severe form:—
- 49. Case V. July 8th, 1818. Mr. S. F. aged 22. He had worked during one year in the lace frame,

principally during the night, when he became affected with the following complaint, about four months He first experienced an unusual degree of weakness, which incapacitated him for his employment; he then complained of a propensity to sweating on any slight exertion or emotion, with pain of the head, loss of appetite, constipation, &c. At first too, he lost his flesh rather rapidly, to the amount, as he supposes, of about 14lb. The countenance became palish and sallow, and he was told that he had a Liver Complaint. The general weakness increased, and a degree of trembling was observed when he lifted his cup of tea to his mouth, and when he attempted to walk or underwent any fatigue.—At present the prolabia are rather pale, the face near the nose is affected with an oily perspiration, the eye-lids are dark, and the general complexion is of a palish, sallow appearance. The albuginea is perfectly white. The tongue is white, loaded, and clammy. He perspires more than usual on any exertion. He has no headach or vertigo now, and has had no cough at any time. There is a sense of fluttering about the heart and stomach, especially on lying down. He says that he is low spirited, and does not like to be long alone or still. His appetite is still impaired; he complains of a sense of load at the stomach, with eructation, but he has no hiccough. The bowels are open. The urine was at first much loaded, but it is less so now. He complains of shifting pains about the limbs .-These complaints have been nearly stationary lately,

the progress they appeared to be making at first having been arrested.—The patient became gradually better, after the date of this report, by taking opening medicines, bathing, and using gentle exercise with a change of air.

50. Case VI. Mr. E. J. aged 28, a tailor. This patient has felt indisposed for a year or more, from recurrent pain of the head, and in the left hypochondriac region; with an irregular state of the bowels,constipation alternating with diarrhea, -the stools being extremely feetid, scanty and dark-coloured .-Five weeks ago he became more seriously indisposed and incapacitated for work, from faintishness, weakness, tremor, tendency to sweating, looseness of the bowels, a loaded state of the tongue, mouth, and breath, and dryness of the throat, but without any manifest loss of flesh.—At present the countenance appears rather bloated, dark in complexion, with a clammy oily moisture. The tongue, gums, and internal mouth are considerably affected, and there is a degree of salivation, which has been observed for a month or two. The skin is cool, and there is a tendency to perspiration on any exertion of body or agitation of mind. He complains of weakness, trembling, and faintishness; and of sweating whenever he is surprised or hurried, which readily happens. His sleep is sometimes disturbed by dreams. The appetite has always been tolerable. The bowels have generally

been loose, especially in the night, and the stools scanty, offensive, and dark-coloured .- The following is the patient's own account of his complaints:- 'I have long had occasional pain of the head, and in the left side under the ribs; my hands tremble very much when I have been at work; I feel faint and weak; I sweat much; I have suffered much from purging during the last four months, having sometimes had six or seven motions in the course of a night; I suffer much from a sour and disagreeable taste in my mouth, and experience dryness with phlegm in my throat, and a sensation of hairs on my tongue.'-This patient's complaints were ascribed to a too assiduous attention to his sedentary business, which he was induced to pay from feeling the pressure of a young and numerous family.—He is already much better from employing, only for a short time, mild purgative medicines, gentle exercise, daily bathing in the trent, and a light nutritious diet; but the cure is now less rapidly progressive than at first, a circumstance very usual in this affection.

51. Case VII. This case is remarkable for the degree of tremor by which it was so long attended. E. W. a tailor, aged 36. August 3rd, 1819. He has been indisposed one year. During the heat of last summer, he became affected with pain of the head which was relieved by a spontaneous diarrhæa, and he lost his flesh and strength, and became affected with great tremor from which he has never recovered.

He remained feeble, tremulous, and thin, and subject to vertigo, fluttering at the scrobiculus cordis, nervousness, impaired appetite, and tendency to constipation,-when I first saw him in February. The degree of tremor was very remarkable; on being asked to write down his complaints, he gave me the following list; - ' weakness, trembling, being easily overcome, and made to perspire, and frequently fainty and fluttered;'-the tongue was clammy and slightly indented, the bowels irregular-constipation alternating with diarrhea. Rhubarb, sulphat of magnesia, calcined magnesia, &c. were prescribed, and the country air recommended. Under this procedure he gained flesh and strength, and his general health improved. He has, however, again experienced some of the preceding symptoms since he returned to his sedentary occupation.

dinary degree of nervousness which sometimes attends this morbid affection. I have given one report only, for the sake of brevity; the subsequent ones were only repetitions of the same symptoms with their usual variation.—Mr. H. T. aged 27. June 25th, 1819. He states that he had been weak and low for some time, when seven weeks ago, after reading the account of the assassination of Mr. Perceval, he was taken with great alarm, and has since experienced the following complaints.—He has lost his colour, flesh, and strength. The countenance is palish, and inclined to

sallowness; the prolabia are however of good colour. He thinks he has lost a stone of flesh. He breaks out into perspiration on the slightest exertion; the least thing induces alarm, hurry, trembling, fluttering about the heart, nervousness, &c. The tongue is rather white; the breath fœtid; the mouth clammy and disagreeable. There are headach, heavy pain at the occiput, vertigo, loss of memory, fear of insanity, sleepiness; oppression about the chest, with sighing breathing; the pulse is nearly natural; there was at first, loss of appetite, but this is now rather removed from taking opening medicine; the bowels were costive; the urine is different at different times, being high-coloured and pale alternately, and sometimes fœtid.

- 53. Having thus given a condensed view of several cases of the Mimosis Acuta, in both its forms, I shall now proceed to state the particulars of some other cases more in detail:—
- 54. Case IX. Mrs. K. aged 32. She was affected by a complaint similar to the one about to be described, two years ago. There were tremor, loss of flesh, and great frequency of the pulse. The affection was deemed incipient *Phthisis* by an eminent practitioner. The complaint receded, but returned in August, 1817. The first symptoms then observed were a yellowness and darkness of complexion, inducing some apprehension of *Jaundice*; loss of flesh; weak-

ness in walking; a tendency to stooping in walking or sitting; great tremulousness of the countenance in speaking, and of the hand when held out; tendency to perspiration both during the night and day. There were occasionally, and especially at first, violent headach, and at other times, vertigo. Twice on going to bed, there was violent palpitation of the heart, rendering the erect position necessary, and attended with coldness of the feet. The pulse varied from 120 to 130. The appetite was greater than natural. bowels are said to have been regular.-This patient residing at a considerable distance from Nottingham, and the nature of her complaint being unknown to me, I corresponded with Mr. K. and in order to ascertain whether a journey to Nottingham were really necessary, I begged that a register might be kept of the WEIGHT of the patient, from time to time. Of this register the following is a copy:-

On August the 18th, 1817, the weight was I	06lb.
On25th,1	$105\frac{1}{2}lb$.
A journey having been taken at this tir	me,
On September the 1st, the weight was1	05 <i>lb</i> .
On8th,1	$05\frac{1}{2}lb.$
On October6th,	06lb.
The patient having returned home,	
On November the 3rd, the weight was	03½lb.
On7th,	01 <i>lb</i> .
On17th,	98 <u>1</u> lb.
On26th,	96½lb.
	93½lb.
On 6th,	92½lb.

At this period it was judged expedient that Mrs. K. should come to Nottingham .- The countenance then appeared thin, rather sallow, and with a dark ring round the eyes. The tongue was clean and moist, but lobulated, somewhat in the manner of some parts of the cerebellum. There was considerable tremor in speaking and in holding out the hand. There was manifest emaciation. Still some headach and vertigo. The pulse was about 130, sometimes even 140. Some days before there had been palpitation of the heart on going to bed. No cough. No shivering. The appetite was great. The bowels said to be regular.-Mrs. K. was now recommended a course of purgative medicines.-The countenance gradually improved, both in flesh and colour; the strength returned; there was a greater capability of exertion; the flesh was restored in the following manner:-

The pulse was often as low as 114 and 116, but was easily accelerated. At this period Mrs K. returned into Lincolnshire. During the week before January 16, 1818, she had gained two pounds. The amendment continued progressive, and in July the weight was stationary at 106lb. The pulse was nearly natural, and the tremor and debility had disappeared. The interval had been occupied in travelling. The use of the purgative medicines had been continued, and a little ale and a nutritive diet had been allowed.

—The affection had been previously deemed a Disease of the Heart, and had been treated with the digitalis.

- 56. This case was thus so far modified in its different periods as to assume the characters of three different diseases, Phthisis, Icterus, and Disease of the Heart. It therefore affords an interesting illustration of the general character of this affection, and of the propriety of the denomination adopted for it.
- 57. Case X. Mr. D. S. aged 53. He was indisposed nearly in the manner about to be described, in May, 1817. The complaint had come on slowly and gradually. It was removed by pills composed of the pil. hydrarg. and rhubarb .-- The affection returned, and made at first a slow, and afterwards a more rapid progress. In November last, he became incapable of supporting the fatigue of an easy occupation. About Christmas his complaint made serious advances, and I saw him again on January the 5th. On this day he was affected with an emaciated countenance, a tremulous state of the lips, a darkness of complexion, and a clammy perspiration on the cheeks near the nose. The tongue was loaded, especially at the back part, with bright red papillæ at the point, and redness of its sides, and it exhaled a halituous moisture; the breath was extremely fœtid. He was extremely feeble, almost incapable of walking, and affected with great and constant tremor, and with a sense of weari-

ness. There had been great loss of flesh: in September he weighed 9st. 7lb. and on this day only 8st. 11lb. There was great perspiration during the night, but no febrile heat. The finger nails had assumed a pale lilac hue. There was little headach, but great vertigo, and faintishness when upright. Restless and sleepless nights. No cough. Frequently rapid, violent. and irregular palpitation of the heart, often without external exciting cause. The pulse was frequent and extremely irregular. There has been much hiccough; at a previous period this symptom had been violent, painful, and almost constant, during fifteen days. Twice there had been vomiting. The appetite was much impaired. The bowels were early laxative; but the stools were scanty, dark coloured, and fætid. -To this date, January 31, 1818, the symptoms gradually declined:-the countenance is improved, the strength much restored, the tremor diminished, and he has this day walked a distance of three miles to Nottingham; the appetite is better, and the hiccough has ceased. The stools are more natural in colour, but still offensive. The following is the register of this patient's weight to this period:-

In September	.9st.	716.
On January the 5th	.8st.	1116.
On12th		
On19th	.8st.	6lb.
On26th	8st.	916.

So that by the use of purgative medicines, the progress of the emaciation has been successively dimi-

nished and arrested, and a degree of flesh has been restored.

February 23. It has been stated that on the day of the last report, the patient walked three miles; the fatigue of this exertion was too great for his strength although much restored; and from this cause, or from partaking greedily of celery, or from having neglected his purgatives, he experienced a most serious relapse.—The ensuing week he lost 3lb. in weight. He became affected with constant and violent hiccough, which continued night and day. There was a complete interruption to sleep and repose. The debility, weariness, and tremor returned. The head was affected with vertigo. The pulse was extremely irregular. - The purgatives were repeated; pills, containing opium and hyosciamus were prescribed; and afterwards an enema with eighty drops of the tinctura opii.—During the second week, he again regained 2lb. of flesh; the hiccough ceased; rest was procured; and he again became renewed in strength.—This improvement still continued, and on

6lb.
816.
916.
8lb.
916.
12116.
016.
4lb.

It is remarkable however that this patient's improvement in flesh was not attended with a proportionate amendment in strength and general health. He still remained feeble and tremulous; he complained extremely of pain, lameness, and weariness in the loins, arms, and legs; he had once more a severe attack of hiccough, which was much relieved by taking white wine vinegar in doses of a teaspoonful.-He continued his opening medicine, which consisted principally of rhubarb, and he has all along taken ale. -The following is the patient's own list of his complaints during the course of his indisposition: -"trembling, weakness, weariness, faintishness, loss of flesh, cold and greasy perspirations, sleepless nights, pain of the head, dizziness, palpitation of the heart, hiccough, disagreeable breath, loss of appetite, a continued lax, weakness and bearing-down of the bowels, much pain and lameness of the limbs."

July 27. This patient has lost 2lb. of flesh during the last month;

 On July the 13th, he weighed.....9st.
 3lb.

 On.......27th,......9st.
 2lb.

 On March....14th, 1819, still......9st.
 2lb.

but he states that he has gained strength notwithstanding, and that the muscles appear to be firmer on his limbs. The tremor is very much less. His lameness, principally in the shoulders, is much diminished and he has now no pain. The tongue is still loaded. The nights still restless. The bowels less loose. The appetite improved. There has been no hiccough lately. There is still some degree of the sense of fluttering about the heart, and the pulse is 84, and not quite regular.—He considers himself better, and intends to resume his usual occupation in a few days. But he is evidently in a state of health altogether precarious.

August 31. My patient has continued to improve in appearance, strength, flesh, and general health, and there is now a hope of permanency in this amendment. He has continued to take rhubarb principally, with a little ale, and a nutritious diet; he is enjoined daily exercise, bathing during the warm season, and sponging when it becomes cold.

58. Case XI. The last case which I shall detail in this place, will afford another instance of the extreme diversity in the history of this affection.—Miss M. A. aged 19, employed at the tambour. Two years and a half ago she became affected with loss of flesh, great weakness, faintishness, palpitation, fluttering about the heart, vertigo, headach, and general nervousness; she was at the same time easily startled and affected with agitation and trembling. Four years ago she was affected with Melæna, and again two years and a half ago, together with the symptoms just detailed.—She seemed to recover from these complaints and was in better health during the sum-

mer of 1817. In October she again became affected with the symptoms of the Mimosis Acuta, and she had again melæna with the other symptoms. She took the pil. hydrarg. with rhubarb, and has never since had any discharge of blood. She recovered rapidly during several weeks. Her complaint then became stationary, and afterwards worse. She took to bed, and continued almost constantly in bed for twelve weeks. She continued her pills, and united the most nourishing diet she could obtain, with a little ale. A month ago she rose from her bed, and has since gradually improved in strength and somewhat in flesh .- At present the countenance is pale; she is very feeble; she however trembles less; she has some appetite; the bowels are kept open by pills of rhubarb; the tongue is whitish. She seems to be recovering, and complains most of vertigo and pain of the head, of fluttering, of occasional palpitation, of faintishness, weakness, &c .- Feb. 22 1819. This patient has gradually and imperceptibly recovered; but still remains feeble, tremulous, at times, and pale. The tongue is whitish, the bowels costive; she complains of pain of the head and back; two weeks ago she again parted with three-fourths of a teacupful of blood, and again ten days and a week ago, with pain in the bowels; without hæmorrhoids.-April 12th. She is still more recovered .- May 3. She now looks very well, but complains of headach.

- 59. The extreme variety in the history, the form, the course and the duration, of the Mimosis Acuta, will be sufficiently evident from the cases thus detailed. The accession of this affection is usually slow and insidious, as before stated; but it is now and then developed, and apparently occasioned, by the occurrence of some other indisposition, or of an accident.
- 60. It is a question of great interest, how far the original affection, the complications, and the changes, in this multiform complaint, are to be viewed in the light of causes and of effects. But I dare not enter into this discussion at present. I shall only observe that this connexion seems often to be distinctly observable; and that the functional derangements have appeared, in some instances, to have undergone the changes denoted by the term metastasis, the affection of one organ having been nearly forgotten by the patient, whilst the derangement of another has engrossed his attention; -in one case the patient had laboured from symptoms of organic disease of the heart for several years; this was at length almost forgotten, and his chief complaint was an alarming degree of vertigo.
- 61. The amendment from the employment of the course of gentle tonic purgatives to be noticed more particularly hereafter, is often rapid, and in cases of short duration, and especially in youth, the cure is soon effected; but in cases of longer duration, when

the constitution has suffered materially, and when the age of the patient is more advanced, the progress towards amendment is slow, and apt to be interrupted, —especially by errors in diet, imprudent exertions, and accidental agitations of mind. The severer form of the Mimosis Acuta is rapid in its course; the less severe form may continue, or be protracted, for several years; the two forms may pass into each other respectively; and each may be variously complicated at different periods.

- 62. Some of the accessory affections too, may assume the serious and alarming character of DISEASE, especially those of the head, heart, lungs, liver, stomach, or bowels.
- 63. The chance of cure is in an inverse ratio to the age of the patient, to the previous duration of the complaint, and to its effect in undermining the general system, or in inducing local organic darangement.
- 64. In conclusion of the History of the Mimosis Acuta, I must regret my ignorance of the morbid anatomy of this affection. This subject will not, however, be neglected if opportunities for dissection should be offered to me hereafter.
- I have noticed the Weight of patients affected with the Mimosis Acuta. I intend hereafter to pursue this subject in relation to
 Chronic Disorders and Diseases in general. The investigation would
 appear, a priori, to lead to some useful information relative to the Diagnosis and Prognosis. The result of this investigation will be given
 in another work,—the Principles of Diagnosis.

SECTION II. THE DIAGNOSIS.

1. THE DIAGNOSIS OF THE GENERAL AFFECTION.

- 65. THE Mimosis Acuta, in its severer but simpler form, has, I think, been generally mistaken for Fever; in its less severe form it is necessary to distinguish this affection from obscure and Insidious Organic Disease; and in its complications, the Mimosis Acuta must be carefully discriminated from an original and local disease of the part symptomatically affected.
- 66. From every kind of Idiopathic Fever, with which the Mimosis Acuta agrees in apparently affecting the whole system or many organs at once, this affection may be distinguished by an attention to the HISTORY of the case, and to the PARTICULAR CONCURRENCE OF SYMPTOMS.
- 67. From the Februs Brevis* the Mimosis Acuta differs entirely in its History; but it is probable that the nature of these two affections is similar. The former is quick or sudden in its accession, and of short duration; the latter is insidious and gradual in its

^{*} See the treatise on DIAGNOSIS, Part II, § 1.

formation, and of more or less considerable continuance. The mistake, however, may be made, if reference be not had to this circumstance; for the general aspect of the patient, the countenance, the tongue, the surface, and some of the symptoms, are very similar in both affections.

- is at once distinguished by the absence of the suffusion and tumidity of the countenance, and of the heat, tumidity, softness, injection, and dryness of the general surface, so characteristic of that kind of Fever. In the Febris Acuta there is more urgency and acuteness of complaint; the patient is obliged to betake himself to bed; the head is more, or more constantly affected; and the breathing more anxious and hurried, whilst there is an absence of the symptoms detailed, § 24. The History of the two affections is also very dissimilar, the Febris Acuta generally following some manifest exciting cause, and being formed quickly, and thus affording a striking contrast with the insidious formation of the Mimosis Acuta.
- 69. From the Februs Lenta† the Mimosis Acuta is distinguished by an attention to the state of the countenance, of the general surface, and of the tongue; to the peculiar concurrence of symptoms; and to the History of the affection.—In the Februs Lenta the countenance is often partially flushed, the surface be-

^{*} See the treatise on DIAGNOSIS, Part II. § 2. + Ibid. § 4.

comes dry, with a burning heat, and the skin harsh and exfoliating, a state sufficiently contrasted with the cool and clammy condition of the countenance and general surface, and the tendency to perspiration, observed in the Mimosis Acuta; in the Febris Lenta, the tongue is at first whitish, and afterwards clean, perhaps morbidly clean, red, smooth, and dry, whilst the breath is little tainted, a sufficient contrast to the state of tongue in the Mimosis Acuta, § 26. In the Febris Lenta the patient suffers perhaps from vertigo. but there is little faintishness or fluttering, and, in general, no tendency to perspiration; the patient is rather torpid than nervous, and less alive to those circumstances which are apt to induce so much hurry and agitation in the Mimosis Acuta, and the Fever is altogether a less multiform and multiplex affection. The patient affected with the Febris Lenta perhaps sits up for a time; then becomes gradually worse and obliged to remain in bed and in the horizontal position; he is in this manner for a longer or shorter period in an inactive state, and at length as gradually recovers or sinks; in the Mimosis Acuta the patient generally remains all along out of bed, or at least much longer than in the former case; and the affection changes and varies more in its severity and in its form, both in its beginning and deline.

70. The Februs Gravis* is distinguished from the Mimosis Acuta by the peculiar aspect and appear-

^{*} See the treatise on Diagnosis, Part II, §§ 6, 7.

ance of the patient, which are generally recognized by those who have noticed them with attention:there is a look of great weakness and oppression, and the patient appears almost incapable of supporting or retaining the erect position, or of undergoing the slightest fatigue. The Febris Gravis is early attended with dulness, pain, or vertigo about the head, and with peculiar aching pains of the limbs and loins. There are sometimes burning heat and a degree of harshness, sometimes a cool moisture, of the general surface. The tongue is at first loaded but generally soon becomes parched, dry, and cracked. The bowels are torpid.—The Febris Gravis is generally formed rapidly, and the patient usually betakes himself early to bed .- These circumstances sufficiently distinguish the Febris Gravis from the Mimosis Acuta.

71. I shall now proceed to draw the Diagnosis between the less severe form of the Mimosis Acuta and Insidious Organic Disease. In the first place the latter affection presents a complete contrast with the character of the Mimosis in general, stated § 2;* it is distinguished by its definite form, and by a regular, slow, progressive, and almost unvaried course, during which the patient becomes gradually affected with paleness, debility, and emaciation, with comparatively little tremor.—The countenance becomes pale and thin, with long and deep wrinkles, but generally remains free from tremor, except in phthisis pulmo-

^{*} See also the treatise on DIAGNOSIS, Part II, § 182.

nalis, and it is peculiarly expressive of disease. The muscular strength becomes gradually much impaired; but this loss of strength differs from that observed in the Mimosis Acuta, in being less attended with the remarkable degree of tremor observed in the latter affection. The general surface is without the peculiarity observed in the Mimosis Acuta, § 27. There is generally an absence of the headach or vertigo, the faintishness, the sense of fluttering, the nervousness, the tendency to cold perspiration, and of the complexity, observed in the Mimosis Acuta, §§ 24-34. The tongue, internal mouth, and breath are at first comparatively little affected, but afterwards frequently become covered with aphthæ. There are chilliness, spontaneous chills, and flushes, and early-morning perspirations.—At length the emaciation becomes extreme; the ankles become ædematous; and anasarca, or ascites may supervene.

- 72. The reader will bear in mind, at the same time, the possibility of the transition of the Mimosis from the state of Disorder to that of Organic Disease, already adverted to, § 38.
- 73. Before I quit the subject of Insidious Organic Diseases, I shall digress somewhat from the proper object of this work, in order to call the reader's attention, in a particular manner, to the characteristics of a disease of this nature, the Diagnosis of which is not always, I think, established in practice;—namely,

the STRUMOUS AFFECTION OF THE MESENTERY in Adults,—a subject which appears to me to have been, in some degree, neglected by practical writers.

74. This disease is particularly characterized by its very slow and insidious accession and progress; by an unusual sensibility to cold; by a tendency to coldness and lividity in the nose and fingers; by the absence of any affection of the head, or of fluttering or palpitation of the heart; by the continued good appetite and the copious, light-coloured, and fætid stools; perhaps by a fixed tenderness, and perceptible tumor, in some part of the abdomen, which is not tumid generally;—the tumor in the abdomen is not, however, always perceptible, nor are the alvine evacuatious at all times of the pale and fætid kind above described. There are a peculiar appearance of the countenance, a peculiar mode of walking, and a peculiar attitude and manner in general,—the whole denoting great disease, and a serious state of emaciation and debility.-This affection is also characterized by the circumstances detailed § 71, contrasted with §§ 2, 24, and by resisting the usual remedies of the Mimosis Acuta.

75. I think the Strumous Affection of the Mesentery in Adults by no means a rare affection. I have, within the last year and a half, seen it in the persons whose cases have been detailed*, in a young woman

^{*} First Edition, pp. 36-41.

aged 21, in a young man aged 22, in a soldier aged 30, and in a lady aged 50. Some of the symptoms of Phthisis Pulmonalis are often superadded to those of Diseased Mesentery; and this disease is often conjoined, in a greater or less degree, with Phthisis.—I intend to republish the cases which were inserted in the first edition of this work, together with some others which I have witnessed more recently, in another place; they will be contrasted with some cases of Insidious Inflammation of the Peritonæum.

76. There is an Effect of Drunkenness which resembles in many symptoms the severer form of the Mimosis Acuta; and as it admits less readily of cure, it is important to distinguish the two affections.-In this Effect of Drunkenness there are great weakness, tremor, and emaciation. The case is characterized by a peculiar state of countenance;—there is emaciation; the expression is much lost; and the eyes assume a peculiar streaked, yellow colour. The tongue and mouth are loaded, and perhaps dry, and there is a peculiar fector of the breath. The manner of the patient is dull, and often slovenly. There is a considerable loss of flesh. There is often a total loss of appetite. The bowels are generally loose. There are, I think, fewer of the complications noticed § 34, than in the Mimosis Acuta. But there is more frequently organic disease.-Particular inquiry should therefore be made into the Habits of a patient affected with symptoms of the Mimosis Acuta.

77. This affection may be exemplified by the following Case: - Mr. S. C. aged 27. He had drunk excessively of ale for 12 months, when he became affected six months ago with lowness, weakness, loss of flesh, fluttering at the heart, loss of appetite, with little affection of the digestion or bowels. He had also an early hoarseness of voice. These symptoms have continued and the loss of strength and flesh has progressively increased to the present time, -he formerly weighed 10st. 12lbs, and now 8st. 13lbs. The countenance is very thin; the tongue loaded at the back part; the voice hoarse; pulse 88; not much tremor or nervousness .- He has gradually gained strength and flesh since he began a course of tonic purgatives, and relinquished his pernicious habit of drinking.

II. THE COMPLICATIONS AND THEIR DIAGNOSIS.

78. IT may be remarked that the complications with the Mimosis Acuta in its severe form are, in general, continued, until removed by remedies; the complications with the less severe form of Mimosis Acuta, are recurrent in paroxysms, until the tendency to the original disorder is overcome. The former cases sooner take on organic derangement than the latter, and are more immediately serious.

1. Of Affections of the Head.

- 79. The affections of the head which occur in conjunction with the Mimosis Acuta, sometimes assume an apoplectic and sometimes an epileptic character; sometimes there is troublesome pain, or vertigo, or some affection of vision or of the hearing.
- 80. The Mimosis Acuta complicated with an Affection of the Head, is distinguished from an IDIO-PATHIC DISEASE OF THE HEAD, by uniting the characters noticed §§ 1, 2, 24, et seqq. with symptoms of affection of the encephalon, whilst the latter affection is denoted by its simple, definite, and regular character, and the absence of those affections.
- 81. There are two occurrences which ought ever to be present to the mind of the practitioner: the

first is the tendency to the transition of a secondary, into Organic Affection, which is particularly observed with regard to the Head;—the second, the possibility of the co-existence of the Mimosis Acuta, and of Idiopathic Disease of the Head.

82. The transition of the secondary affection of the head in the Mimosis Acuta into an Organic Disease, or the co-existence of the latter affection with the former, is to be apprehended on the occurrence of any of the following symptoms in a serious degree and continued form: - A sense of flushing or fulness about the head; acute pain of the head; unusual heaviness, dull headach, or vertigo; drowsiness, stupor, disturbed sleep, delirium, incubus, stertor; forgetfulness, timidity, confusion of mind, change of affections, tendency to laughter, or tears; affection of the senses, as temporary loss of sight, flashes of light, double vision, singing or loud noises in the ears, intolerance of light, or sound,-tenderness of the scalp. Many of these symptoms, however, occur in the Mimosis Acuta. It is only when they exist in an eminent degree that they denote danger. But it is best to take an early alarm. The danger of organic affection of the head is less equivocal on the occurrence of an unusual distortion, or an unmeaning expression, of the countenance; of a defect in articulation; of a temporary numbness or torpor, or of transient and partial weakness of any of the limbs; especially if one side of the body alone be affected. The occurrence of stupor, convulsion,

paralysis, or relaxation of the spincters, leaves little to doubt respecting the existence of internal disease of the head.

- 83. The following melancholy instance of the combination of the Mimosis Acuta with an apoplectic Affection of the Head, is particularly instructive:—
- 84. CASE XII. Mr. G. aged 33, had experienced symptoms of Mimosis, with a loaded state of the tongue, internal mouth, and breath, for several months; he had been observed to complain much of the head, frequently applying the hand to the forehead, or laying the head on a table, and stating that he suffered from severe pain. On visiting him he seemed to be affected with symptoms of the febris brevis, but the History agreed more with that of the Mimosis:-the countenance was moistened with a clammy or oily perspiration, the tongue and breath loaded; he complained of dulness about the head, but made light of the affection. His complaints increased notwithstanding the use of very active purgatives; on minute inquiry he had noticed a defect in vision,—the lines he was reading sometimes disappearing as if covered by the finger; he complained of tenderness of the scalp, of sensibility to the light, and he requested that his room might be darkened. Afterwards he was seized with stupor, the pupils dilated, and the pulse became slow-about 55 in the minute. By bleeding blistering, purging, &c. these symptoms disappeared,

and recognition returned. But in two days afterwards, restlessness, delirium, with picking of the bed-clothes, followed by stupor, with ruttling in the breathing apparently from phlegm in the fauces, came on during the night. The pulse was strong and full at six in the morning. But nothing was done for the patient. The pulse became gradually feeble, the countenance pale and sunk, the respiration laboured, with still greater ruttling. About half after ten, some remedies were resorted to, but too late, and in vain. The left side of the face, and the extremities of the left side became paralytic, and he expired about twenty-four hours after the second attack of stupor. An examination of the encephalon was refused.

- 85. The following cases assumed, in some degree, the *epileptic* form.
- 86. CASE XIII. Mr. E. aged 50, has been subject to derangement of the digestive function for many years,—always increased by anxiety or fatigue,—and requiring a frequent recourse to rhubarb and peppermint water. About six years ago he began to be frequently obliged to leave the desk of the counting house, from experiencing confusion of the head. Five years ago, after having suffered during several days from sickness, lassitude, and vertigo, Mr. E. was taken suddenly in church with severe vertigo, faintiness, general perspiration, sickness and some vomiting;—he asked for a glass of water, rejected it from

the stomach, and was then relieved, but remained feeble for the rest of the day. From this time Mr. E. experienced, with some of the symptoms just detailed, occasional attacks of singing in the ears and glimmering of the sight, but had no severe return of the affection until October, 1819, when he suddenly fell down in the yard, as if he had been shot, -again, in the winter, when sitting near the fire, the feet starting upwards and the head falling on the arm of the chair,-and once more in May last. These attacks were anticipated from vertigo and lassitude, and were followed by sickness and perspiration, and for some days, by an unpleasant stupor. On the whole, the affection has increased in the frequency of its returns during the last two years. Mr. E. mentions an overwhelming feeling on awaking which compels him to get up and not to venture to sleep any more. Fatigue, anxiety, and attention of the mind have appeared to re-induce the feelings of this distressing complaint .- Leeches, cupping, and venæsection have been employed, but only with temporary relief. Relief of a more decided and permanent kind has resulted from a course of active purgative medicine, by which much dark and fætid fæces have been evacuated. The tongue and complexion were as usually observed in protracted cases of Mimosis.

87. Case XIV. The following account is given in the patient's own words. 'The first time I experienced this malady was about the age of 15. It

came on during the time I was endeavouring to recollect my usual repetition which I had learnt but imperfectly; I had been standing some time, and at length sank on the floor, but recovered in a few minutes .-These attacks were repeated at remote intervals until I was about 21 or 22, when they became more frequent. -At this time I was engaged without intermission in the professional duties of an attorney, and was naturally of a sedentary disposition, even in my amusements. -I was about this time and previous to it, subject to an almost constant pain in my head and eyes, and occasionally to lassitude and want of energy .- I took but little exercise, lived very moderately, but ate heartily. About this time I applied to Dr. S. I adhered to his medicine for some time and found myself much benefited by it .- I have since that time been attacked more rarely, and during the time I pursued the remedies prescribed by you, I do not recollect having any return of the complaints. My general health has been much improved by the medicine which I have taken whenever I found the state of my bowels required it .- I had flattered myself that I should not be again subject to these attacks, but on Wednesday last I experienced another instance of them. I had been spending about a week in the country, and living rather more generously than usual, viz. by taking ham or meat with my breakfast, and four or five glasses of wine after dinner, with meat suppers &c .- On Wednesday afternoon I had been standing a considerable time exposed to a cold wind, taking sketches of the

church, and felt much chilled by it .- In the evening we had a party of friends and I sang to them for an hour or two until I was rather fatigued .- About 11 o'clock at night, after a supper of fruit pie, the attack came on in the usual way, by an involuntary train of ideas presenting themselves to my mind, which produced a sickening effect upon me, accompanied at the same time by a sort of fascination which led me almost irresistably to pursue the train of ideas, which at the same time I knew would occasion me to faint. —I say almost irresistably, because I have sometimes checked this train, and by rousing myself, prevented the actual fainting from taking place .- I never can completely recall the train of ideas, and on my endeavouring to do so, I feel a tendency to a repetition of the attack .- My first sensation on recovering sensibility is an anxious wish to pursue the dream, through curiosity to see the event of it; this is succeeded instantly by a consciousness of my situation and an anxiety to convince my friends that I have quite recovered from the effects of the attack. I feel a wish for a little cold water and my face is covered by a cold perspiration; an inclination to lie down and sleep succeeds and in a few minutes I am quite restored .- Previously to the attack on Wednesday I felt a drowsiness and an unusual propensity to gape and stretch my limbs several times.—On my recovery I had for several minutes a singing in the back of my head, like the noise of a boiling teakettle, or rather like the gradual escape of fixed air from a bottle when

the cork has been gently loosened.—I went to bed soon afterwards and took one of your powders and some peppermint water. I had a good deal of uneasiness at my stomach and bowels and a dislike to the usual pressure of my clothes upon them, which I attributed to flatulency. My feet were very cold and I was very drowsy.—The medicine operated in the morning and I was tolerably well, but my bowels have not yet recovered their healthy state.—I have no doubt that these attacks proceed from a disordered state of my stomach as they are generally preceded by costiveness and headach.—June 3rd, 1820.'

88. CASE XV. Mr. W. of S. aged 22. This young man had been, for some time, affected with confused pain of the head, vertigo, &c. One day he was taken suddenly, fell down on the ground, and lay in a state of insensibility for some time. He recovered, but remains subject to pain, confusion of the head, vertigo, and tendency to fall down. He has become weak, nervous, dispirited, inert, and incapable of business. The tongue is swollen, loaded, indented, and pasty, with deep folds on its surface. The breath exceedingly fætid. The countenance has a diffused appearance of oiliness and gentle flushing. The stomach and bowels are much disordered.—In this case it was dubious, at first, whether the affection arose merely from disorder of the digestive organs, or was an idiopathic disease of the head. The state of the tongue, breath, digestive organs, mind, &c. seemed to indicate a symptomatic affection of the Head, and such the History seems to have proved it; for the latter symptoms have increased materially since the first time I saw this patient, and the affection has assumed the form of hypochondriasis.

- 89. The following Cases illustrate some other affections of the head which arise from a state of Mimosis.
- 90. Case XVI. Mrs. M. October 5, 1818. On being called at 5 p. m. to this patient I found her affected with insensibility and delirium; pain about the umbilicus; hurry and rapidity of breathing and a feeling of want of breath with fluttering; the pulse was 120; the countenance rather pale and cool; she appeared to the bystanders to be faint; the tongue rather dry. The appearance of this affection was alarming; more than once she was supposed to be expiring. Five grains of calomel were given; a suppository of inspissated molasses was introduced into the rectum. They were long in operating but at length produced complete relief.
- 91. Case XVII. C. H. aged 21. He complains of severe recurrent pain of the head especially of the forehead, with great noise in the head. The tongue is very loaded, white, and indented. The bowels are costive. He is subject to these complaints and is a

framework-knitter.—He was relieved by taking two emetics and a course of purgative medicines.

- 92. Case XVIII. Mr. B. aged 40. June 8, 1819. He is subject to disorder of the stomach. A few days ago, on taking cold, this disorder was, as usual, aggravated, and he was affected with severe pain of the head and of the right side over the false ribs. The skin was moist, and clammy, and the tongue extremely loaded. All these complaints were removed by emetic and purgative medicines in the course of five days, and he is now free from acute complaint. There was a little eruption of herpes on the lip.
- 93. In one gentleman there was a state of mind approaching to mental derangement; in a lady there was an attack of affection of the head with a loss of memory,—apparently the effects of a state of Mimosis.
- 94. There are few diseases which it is more important and more difficult to recognize early, than a state of Idiopathic Inflammation or Organic Disease of the Brain.* The Diagnosis is partly negative. For the first question is, Is there a state of Fever, of Mimosis, or of other Disorder, which may account for the symptoms of cerebral affection? If not, there is greater

^{*} See Sir G. Blan's paper, in the Trans. of a Soc. for the Imp. of Med. and Surg. Knowledge, V. 2. p. 210; Abernethy, V. 1. pp. 202, 106; Bateman's Reports; passim; &c.

apprehension of Idiopathic Disease of the Head, in which there is usually an absence of the symptoms of Idiopathic, as distinguished from Symptomatic Fever,* and of the symptoms of Mimosis;—except in the cases of complication.

2. Of Affections of the Chest.

- 95. The affections of the chest which occur in consequence of a state of Mimosis, are, principally, oppressive dyspnæa resembling Asthma, and cough with expectoration resembling Phthisis Pulmonalis.
- 96. The complication of the Mimosis Acuta, with Paroxysms of Oppressive Dyspnæa, resembles Asthma in a very high degree. And it is not improbable that Asthma properly so called, frequently, if not generally, originates in a complication of this nature, but is afterwards continued, or repeated, either from a degree of disorganization induced in the heart or lungs, from the influence of the external causes of Asthma, or from the causes of Mimosis. The following cases sufficiently exemplify the complication alluded to:—
- 97. CASE XIX. D. M. aged 20, a tailor. August 30, 1813. He has complained of pain and other

^{*} See the treatise on Diagnosis, Part II, §§ I, 2, 4.6, 7, contrasted with §§ 3, 5, 8; and with §§ 107, 108.

affection of the chest for eight or ten days; but he has been affected, for a much longer period, with vomiting, pyrosis, a loaded state of the breath, mouth, and tongue, constipation of the bowels, &c. Three days ago he was bled, from the idea that the affection was a case of pneumonia.-I was called to him last night:-The breath was extremely fætid, the tongue loaded, the bowels constipated. He was affected with a paroxysm of urgent dyspnæa, complained of want of air, and said he was dying; the dyspnæa had begun an hour before, and had become gradually more and more urgent; there was great anxiety; the respiration was very rapid, and seemed a literal gasp for breath. He had also a cough, attended with a dull, hollow, peculiar sound, and with a copious, frothy, mucous expectoration. The pulse was about 100, without sharpness .- He was somewhat relieved by taking a few drops of æther, and effectually by the operation of a purgative of calomel and jalap.

98. Case XX. The Rev. Mr. A. aged 34. He has long been subject to a state of Mimosis Acuta in its less severe and more continued form, marked by a variable degree of sallowness of the complexion, loaded tongue, feetid breath, and foulness of the mouth in general, without any observable loss of flesh. He is easily hurried by surprise, and fluttered; his complaints are then aggravated, and accompanied by nervousness and tremor. The same effects are in-

duced by any unusual mental or bodily exertion.-This gentleman is subject, at these times, to a heavy and dull, or to a more acute pain of the occiput, for which cupping has been deemed necessary. The pain has sometimes been relieved by the horizontal posture, and sometimes it has been excited by an empty state of the stomach.-In other attacks the pain of the occiput has been absent, and the patient has been affected with dyspnæa, attended with restlessness, a sense of want of air, a necessity for being fanned, &c. The paroxysm has generally taken place about 1 a. m. It has been relieved by coffee. It has usually succeeded to an increase of the state of disorder constituting Mimosis, and has been obviated by correcting that state. It has recurred at distant intervals for several years. Formerly the dyspnæa was attended with cough, but not lately. A degree of oppression has generally been experienced during the day succeeding the paroxysm.*

99. The complication of the Mimosis Acuta with Cough and Expectoration is still more common, especially in protracted cases of this affection, both in its severer and less severe forms. This complication requires to be accurately and early distinguished from Phthisis Pulmonalis, or to be accurately traced into a similar state of affection of the general system and of the pulmonary organs.

^{*} See further the treatise on Diagnosis, Part II, pp. 206-209, for the Diagnostics of Asthma.

100. The discrimination is again to be begun by contrasting the character of the Mimosis Acuta, given §§ 2, 24, et seqq. with the general character of Phthisis Pulmonalis, which is very dissimilar, and almost opposite to that of the former affection. Phthisis Pulmonalis is simple and definite, in its character; in general little varied, but rather continually progressive, in its course, -although this observation has some exceptions; it is comparatively free from the tremor, fluttering, susceptibility to hurry, and general nervousness, observed in the Mimosis Acuta, § 24; there is an early loss of flesh, a symptom which ranks amongst the first which arrest attention; there is also an early weakness, but this is different from the weakness attending the Mimosis Acuta, being less attended by tremor, faintishness, &c. and the degree of remaining muscular strength is even remark, able in the latter stage of Phthisis Pulmonalis.*

101. The countenance in Phthisis Pulmonalis early assumes a peculiar and delicate paleness, but is liable to occasional, or spontaneous, gentle blushes. There is often an early movement of the nostrils, induced by the alternate acts of respiration, affording one of the earliest diagnostic signs of Phthisis. The hands and general surface are pale and delicate, the roots of the nails frequently rather livid. And there is a charac-

^{*} For the Diagnostics of Phthisis Pulmonalis from other diseases which resemble it, see the late work of M. Bayle, Armstrong on Scarlet Fever, &c. Bell's Surgical Reports, Vol. 1, pp. 18, 21; and the treatise on Diagnosis, Part II, pp. 169—172, 189—196.

teristic, irregular, alternation of chilliness, feverishness, and perspiration. There is an obvious labour in respiration, which is aggravated on going up stairs or using active exertion; a full inspiration, however, does not always occasion inconvenience. The cough is slight and tickling at first, perhaps without violence, and is less early attended with expectoration. The pulse is permanently frequent.

102. The complication of cough and expectoration may take place in either the more or the less severe form of the Mimosis Acuta, but I think the latter case is the more frequent.—In the severer form of Mimosis Acuta there is sometimes an evident labour in breathing which induces a movement and contraction of the nostrils, and there is a degree of cough with the expectoration of a little viscid mucus.—In the less severe form of this affection there is often, for a considerable period, a copious expectoration of transparent, frothy, and viscid, and, at length, of opaque, and tinged mucus. There is a cough, hard-sounding at first, and occurring in fits; afterwards easier and less in fits as the expectoration assumes more of the puriform appearance; and again difficult and in painful fits as the strength of the patient declines. The pulse is at first little accelerated; the respiration little affected, and the nostrils unmoved; but a degree of movement of the nostrils, of labour in the respiration, and of frequency of the pulse, is observed as the expectoration becomes puriform, and streaked with blood, and as

disease of structure is induced in the lungs. The usual symptoms of the less severe form of the Mimosis Acuta are superadded;—the skin is usually as already described, but sometimes harsh and dry; the tongue is sometimes furred and black at the back part; and emaciation takes place in a degree commensurate with the diseased affection of the lungs. The History of the case is peculiar,—the affection is perhaps cured by proper remedies, or becomes stationary and mitigated for a time, perhaps an apparent or decided amendment occurs even during several weeks, and again the disease proceeds,—so that the physician is alternately elated with hopes and depressed with fears,—circumstances scarcely observed in Phthisis Pulmonalis.

eminent example of this last remark relative to the course of the complication of the Mimosis Acuta with cough and expectoration, when the affection terminates fatally. Mr. N. aged 39, had long been affected with a loaded state of the tongue, swelling of the gums, feetor of the breath and other symptoms of the Mimosis Acuta, when he became seized with symptoms of an inflammatory affection of the chest, which were removed by bleeding, and blisters, conjoined with purgative medicines. He seemed to get better of the immediate inflammatory complaint of the chest, but the looked-for restoration of health and strength did not ensue, but seemed delayed week after week, without,

however, any retrograde change. He recovered, indeed, a little, and was removed to a short distance into the country. Here a manifest amendment in point of strength occurred and continued during two or three weeks. But this amendment did not proceed. He became stationary once more, and continued in a dubious way for a week or two longer. He then appeared distinctly to lose strength, and for the first time since his recovery was looked for, there was a sensible loss of flesh. Thus five weeks transpired from the abatement of the inflammatory symptoms to the period of the patient's going into the country, and seven weeks more passed between hopes and fears during his stay from Nottingham .- During the course of this complaint the countenance was affected with an oily perspiration and a sallow or swarthy complexion, with paleness of the prolabia. The mouth, tongue and breath were foul and loaded. The hands and nails pale. There was a most remarkable degree of tremor, which seemed particularly to impede the voluntary motions. There was at first scarcely any dyspnœa or movement of the nostrils; but both these symptoms were observed as the affection of the lungs was supposed to pass into organic derangement. There was at first a hard cough, with rather viscid, mucous expectoration, increased when the stomach was empty; afterwards the expectoration was greater in quantity, more viscid and opaque, and considerably tinged with blood. The pulse very frequent. The bowels were kept open by medicine. The appetite was tolerably good. - The patient survived nearly three months after this report, the affection of the lungs becoming more and more confirmed, the flesh being completely lost, and the strength exhausted. The cough was sometimes less severe; the expectoration variable in quantity, but free from the tinge of blood; the pulse less frequent; the bowels sometimes affected with diarrhæa.—An examination of the chest was refused.

- 104. CASE XXII. In another case there were considerable loss of flesh and strength, a frequent pulse, much cough with copious, mucous expectoration, and a loaded state of the tongue and breath; with some other symptoms of the Mimosis Acuta. The affection yielded very favourably to the operation of mild purgative medicines conjoined with small doses of opium.
- 105. CASE XXIII. In a third case there were all the symptoms of the severer form of the Mimosis Acuta, together with a slight cough, expectoration of phlegm, and a manifest difficulty in respiration, with movement of the nostrils. The affection yielded most favourably, and the health, strength, and flesh returned, by the use of purgative medicines, with a nutritious diet, gentle exercise, and the country air.*

^{*} See Abernethy, V. 1. p. 207; Dr. W. Philip, Med. Chir. Trans. V. 7. pp. 499, &c.

3. Of Affections of the Heart.

106. The affections of the heart which occur in consequence of the Mimosis Acuta, are fluttering, palpitation, irregular action, and syncope more or less perfect.

107. Fluttering and Palpitation of the Heart, are amongst the most frequent symptoms of the Mimosis Acuta; it is, therefore, scarcely necessary, especially after the detail of Cases IX and X, to offer any exemplification of this complication. I shall, however, make a few observations which may conduce to the establishment of a correct Diagnosis.-In the severer case of Mimosis Acuta, it will be necessary to wait, and to watch the effect of the remedies in removing this disorder; the palpitation of the heart, if a consequence and effect merely, will be mitigated or removed with the original affection.—In the less severe form of this complaint, the discrimination must be principally founded on a cautious observation of the effect of bodily exercise on the action of the heart when the symptom of palpitation is otherwise absent, and when the patient is least indisposed,—and on the continued History of the complaint. In DISEASE OF THE HEART the patient often emphatically observes, 'I should be well if I vegetated only; it is bodily exertion and mental agitation, which renew and recall the sense of my dreadful disease.' In the complica-

tion of Mimosis with palpitation, the patient, if not prevented by weakness, can, at the time when the hurried movement of the heart is absent, run pretty rapidly, or walk up stairs, without suffering more than is usual; and periods occur when he has passed several days, weeks, or months, perhaps, without experiencing the palpitation. In a disease of the heart, these circumstances are not, I think, observed; the uneasy sensations which accompany this disease, if absent at times, are always excited on any corporeal exertion; and, as I have elsewhere observed, disease of this organ is in general highly characterized, and distinguished from certain symptomatic disorders of its functions, by the permanency of the affection; by its invariable aggravation on muscular exertion, as well as mental emotion; and by the particular relief obtained at first from bloodletting.-Disease of the Heart, although its symptoms may be mitigated at one period and aggravated at another, is however permanent; the symptoms are never entirely absent; and they may at any time, be renewed, in an aggravated form, by muscular exertion. In dubious cases, the patient may be made to run up stairs; -the symptoms of an organic disease of the heart are invariably aggravated by this muscular exertion, the pulsation of the heart becoming violent, the pulse perhaps irregular, the respiration exceedingly difficult, &c. circumstances not equally observed in symptomatic derangements of the functions of this organ, unless when they are attended with great debility. There

is almost always, too, great but transitory relief from bloodletting,—in a degree not observed in the symptomatic affections.'*

- 108. Case XXIV. Dr. D. aged about 70. He became affected with the following symptoms:—loss of flesh, weakness, tremor, tendency to tears, a loaded state of the tongue, mouth, and breath, and disorder of the bowels with scanty stools; with these affections there were symptoms which led to the idea of Angina Pectoris, or of Dropsy of the Pericardium.—From the use of gentle rhubarb purgatives, with exercise in the open air, the patient gradually recovered a considerable degree of strength, the general health improved, and he has run up a high stair-case, with considerable activity, and without feeling any of his former uneasy sensations about the heart. This amendment has now continued at least a year and a half.
- 109. Case XXV. Capt. B. aged 50, was supposed to labour under an Organic Affection of the Heart, inducing palpitation and great irregularity of the pulse, for many years. It has been nearly forgotten for some months, and he has suffered exceedingly from attacks of vertigo. The complexion is sallow; the tongue swollen, lobulated, indented, and loaded; the breath tainted; the appetite good; the digestion

^{*} See the treatise on DIAGNOSIS, Part II, § 157.

often impaired; and the bowels constipated. His complaints are all mitigated by a course of purgative medicines in which he still perseveres.

110. CASE XXVI. Mr. D. of L. C. aged 34, a farmer and grazier. February 4, 1819. Six weeks ago, soon after lying down in bed, he became suddenly affected with an indescribable sensation as of approaching dissolution; he got out of bed, took some warm negus, and began to eructate copiously; -he had previously complained of vertigo, - once of pain at the stomach ascribed to wind, and several times he had asked for aloes from being constipated. Soon after taking the negus, palpitation of the heart took place, and increased to a considerable degree of violence. Since that time he has had three other violent attacks of palpitation;—the second five days after the first, and like it, almost five minutes after lying down in bed; the third a week after the second, at night, but before he was gone to bed. Since these periods he has also had palpitation, but in a less severe degree. About twenty days ago, he became affected with fluttering, referred to the scrobiculus cordis, and lower in the abdomen. At different times he has had the following symptoms: - pain at the stomach, vertigo, pain at the back of the head, restlessness, want of sleep, with starting, twitching, dreaming, alarm, and despondency,-and fear of immediate dissolution; some tremor, and several times wheezing breathing; he has felt overcome with sleep, which has induced pain

of the heart and palpitation, and a feeling as if the heart would burst.—A fortnight ago the skin was dry, the tongue much loaded, and swollen, the bowels constipated, with flatus in the bowels, the motions being very offensive and dark coloured, and the water high coloured and once or twice with much sediment. The appetite has been good all along. There is considerable loss of flesh.—The tongue is now clean; the flesh regained, and he is better in every respect. He sleeps still but little. A short time after this report this patient became affected with difficulty of urine. He afterwards became subject to dyspepsia for several months in its ordinary form, but at length recovered completely.

111. Case XXVII. Mr. W. aged 48, from siting in an over-heated counting-house, and subsequent exposure to cold, became affected by the following symptoms: — The countenance was rather flushed, but at the same time rather cold and clammy. The tongue was white, loaded, indented, and swollen. He experienced faintishness when upright, and could not bear to approach the fire. Little headach or vertigo. The pulse was rather frequent, and remarkably irregular and intermittent. There was no appetite. These symptoms gradually declined on taking an emetic followed by a dose of calomel, and other purgatives. The affection continued about eight days. The tongue got gradually clean from the edges to the centre; the pulse gradually regular and of natural frequency;

the appetite better; and the patient could in a few days bear the upright posture and being near the fire. January 15, 1819.

- 112. The cases detailed, §§ 86, 87, seem to have been attended by fainting as well as the singular affection of the head described. Indeed all these cases are complicated in an eminent degree, and all appear to me to establish the propriety and the importance, in a practical point of view, of the denomination I have appropriated to them. The following case is however still more particularly a case of fainting; it is one which I have carefully watched; I give it in the patient's own words:—
- have long been subject to a sort of fainting fit, which attacks me very suddenly and affects me in an extraordinary manner. I suddenly appear to myself to be at the point of death, and to have lost every faculty. I have sometimes completely fainted for a few minutes, and have occasionally vomited; I then experience a sense as of a stick pressing violently upon the fundament. On recovering, a copious perspiration bedews my forehead, and I am extremely pale, and feeble. About seven years ago Dr. H. suggested to me that the whole of these complaints arose from a disordered state of my bowels, and that, by a continual recourse to mild purgatives, they might be prevented. This opinion I have found to be true, having experienced

complete relief by following that prescription." To this account of the patient's sensations I add the following one of the appearances in an attack which I witnessed. He was taken on the evening of the 9th of January, 1819, with a sense of hunger succeeded by severe pain of the stomach; when I saw him, the countenance was pale, cold and clammy; the feet cold; the pulse feeble and intermittent; the pain much increased. I gave him a dose of ol ricini, and a glass of hot brandy and water. He was soon relieved, and perfectly so on the operation of the medicine.*

4. Of Affections of the Stomach and Bowels.

114. The affections of the Stomach and Bowels in the Mimosis Acuta, besides those comprised in the more general disorder, are numerous. They are chiefly the following:—hiccough, hæmatemesis, melæna, and pain in different parts of the tract of the alimentary canal, assuming the different characters about to be described.

115. The complication of Mimosis with Hiccough, § 34, has been sufficiently noticed in Case X. But as this symptom is sometimes extremely severe, and as vomiting of food is sometimes superadded in an extraordinary degree, I think it proper to detail a case or two of this affection.

^{*} See Bell's Anatomy, V. II, p. 240; Abernethy, V. 1. p. 212; Farre's Path. Researches, Ess. I, p. 46, &c.

116. Case XXIX. The first case is an instance of violent Hiccough. Mr. C. aged about 50, of Sleaford, Lincolnshire, after feeling chilly and weak, as he supposes, from cold, first experienced Hiccough on December the 18th, 1817, at 5 p. m. It was removed about 11 p. m. by a little gin and water. It returned on the 19th at 5 a.m.; it was removed for half an hour by a little rum, but recurred and continued with little interruption through the day. On the 20th, the Hiccough was arrested for a little time by laudanum and musk, but it again returned. On the 21st, a blister was applied over the stomach, but the Hiccough continued, and was accompanied on this day by a singular sort of crowing noise during inspiration. On the 22nd, snuff was recommended, which arrested the Hiccough for a short time. The affection, however, continued in a violent form through the night. On the 23d, an emetic was given, and six leeches and a blister were applied; the Hiccough ceased for a time, but recurred with the crowing and continued during the night, preventing the patient from lying in bed. The life of the patient was thought to be in imminent danger. The affection still continued on the 28th, when I recommended purgative doses of calomel and infus. sennæ, from the first operation of which the Hiccough entirely ceased with the exception of a slight return once more on the 29th. The purgative was continued for some time, by means of which a loaded state of the tongue and of the bowels was removed. I should have also stated that a little opium and carbonate of ammonia were given; but to these remedies I did not ascribe much effect.

117. Case XXX. Mr. G. C. aged 26. September 6th, 1817. This patient had an attack similar to that about to be described, eight years ago; it continued several weeks. The present attack commenced in March last, and has continued in a more or less severe form, since that period. After eating he becomes affected with chilliness, pain at the stomach, nausea, and tumidity of the abdomen; he is then suddenly affected with vomiting, and the rejection of much mucus, or there are eructation, retching, and vomiting. He remains poorly for about an hour, during which period there is some pain. He then pursues his usual occupation. His complaint again recurs on eating. He is feeble and has lost some degree of flesh. The pulse is about 80, and rather feeble. The tongue is white and loaded. The bowels constipated. The hands are apt to become cold .-The affection yielded to the remedies to be noticed hereafter.

118. Case XXXI. Mr. B. A. aged 24. He was affected in the following manner about six months ago; the attack was less severe than the present, and continued one month; the affection then ceased, but it returned ten days ago. At present, September 9th, 1817, he becomes affected with vomiting and retching after eating; with these symptoms hiccough is

sometimes conjoined, and this is occasionally of the most violent kind, affecting him like a spasm, drawing the body awry, and sometimes throwing him off his chair; he suffers such excruciating pain as to induce a flow of tears; and the breathing is irregular, rapid, and anxious. The tongue is loaded; the bowels costive. The pulse is nearly natural. There are weakness and some loss of flesh.—This affection was removed by the remedies to be noticed hereafter. It recurred after a time and again yielded to medicine. The patient became attacked by a spasmodic affection of the left arm, so as to deprive him of the use of the hand. This affection yielded to the same purgative remedies, and he has since remained well.

- 119. Case XXXII. Mr. J. R. aged 31.—He is naturally thin and feeble. He is at present affected with vomiting after meals, without hiccough. The pulse is natural. The tongue white. The bowels costive.—The affection yielded to the purgative remedies.
- 120. This affection must be carefully distinguished from Organic Disease of the Stomach. Its History,—its cessation for a period and subsequent recurrence,—its violence even at the beginning,—its spasmodic character, &c. are sufficiently contrasted with the slow formation and insidious progress, the continued form, the regularly progressive course, and the unceasing character of organic disease. It is fur-

ther distinguished by some of the symptoms of the Mimosis Acuta;—even the hiccough is, I think, a rare occurrence in the organic affection, which is rather denoted by pain, which gradually increases after eating until the stomach is unloaded by vomiting.*

- 121. MELÆNA, § 34, is an affection of much more frequent occurrence than is generally imagined. Its existence is, of course, only ascertained by accident. But I have found, on inquiry, that it has occurred in a great number of the cases of Mimosis Acuta which I have treated. It occurs also in the other forms of Mimosis to be described hereafter. It has been a very frequent occurrence during the present summer of 1818. With Melæna, HÆMATEMESIS is not unfrequently combined. The two diseases appear to be similar affections of different parts of the alimentary canal. The discharge of blood varies greatly in quantity and appearance. It yields in general to the employment of purgative medicines, with the pil. hydrarg. See Cases III. & XI. In several cases the patient has awoke in the morning with blood in the mouth.
- of Mimosis Acuta, Pain in the Epigastric, or in one of the Hypochondriac, or Chondiliac Regions, § 34. This pain is so common that I think few cases of the Mimosis occur without some degree of it. There are

^{*} See further the treatise on Diagnosis, Part II, §§ 187, 188.

sometimes all the symptoms of an attack of GALL STONES. Sometimes there is severe pain extending across the false ribs, leading to the suspicion of IN-FLAMMATION OF THE PLEURA, or, together with the affection of the complexion, it leads to the suspicion of Inflammation or Disease of the Li-VER. And its situation sometimes induces an opinion that the SPLEEN is affected. It is distinguished by being liable to recede and to recur, by varying its situation, frequently by being unattended by tenderness on pressure when the examination is made with proper care, and by conjoining the characters, §§ 2, 24, et seqq.*—This pain has appeared to me to have its seat in the Colon, some part of whose course it occupies, § 30, —from the hollow of one ilium round to that of the other.-Pain, tenderness, and even tumor, frequently exist in the course of this intestine from the remora of scybalæ; the case is perhaps considered to be an instance of organic enlargement of some viscus in this course; the cause being removed by the action of aperient medicines and by friction, the tumor is supposed to be discussed by the pil. and the ung. hydrarg.—Pain in some part of the course of the colon, but especially in the seat of the sigmoid flexure, often attends and denotes a loaded or contracted state of the RECTUM; the real state of this bowel must be ascertained by an examination, and by the aid of enemata.

^{*} See further the treatise on DIAGNOSIS, Part II. pp. 179-182, 254,

123. Case XXXIII. Mr. L. aged 40. November, 1818. When I first saw him he laboured from tenderness, pain, and fulness at the scrobiculus cordis. A similar pain had affected the right side,—and had been mitigated and had recurred several times.-The countenance was rather leucophlegmatic and bloated; the tongue loaded; the breath tainted; the bowels constipated and medicines had acted inefficiently; the pulse natural in frequency. He had suffered for several weeks previously to the attack, from lowness of spirits; tendency to shed tears; tremor; nervousness; vertigo; fluttering; and a loaded state of the tongue, breath, and bowels. He was much relieved by leeches and still more effectually by the copious action of draughts with ol. ricini, and pills with the pulvis jalapæ.

sulted me in August, for a severe pain with tenderness in the left hypochondriac and chondiliac regions; the tongue was loaded, swollen, indented, and with enlarged papillæ, the gums red and tumid, the breath extremely fætid, the bowels were said not to be much disordered; the countenance was pale and swarthy; there was some loss of flesh; the general health was manifestly impaired. The pain was almost immediately relieved by active purgative medicines; and the health was restored by a perseverance in milder doses of the same remedies.—The symptoms had led to the apprehension of organic disease. They had subsisted for several weeks, or even months.

- plains of pain in the left iliac region, accompanied by considerable tenderness. It was relieved by employ-ty-ing the rectum of Scybalæ by means of an enema.
 - 126. It scarcely appears necessary to do more than refer to the enumeration of Constipation, Diarrhæa, and Tenesmus, as occurring in the Mimosis Acuta, § 24; the source of these affections must be traced to the original disorder, -and it must be distinguished from a diseased state of the rectum.—The constipation is frequently attended by pain deep in the sacrum, which is apt to be aggravated by the movement of a carriage in travelling .- The Diarrhœa induces frequent, scanty, liquid, fætid stools, often with mucus, sometimes with more or less of blood; there are also in different instances, gripings, tenderness, and Tenesmus. The Diarrhea often occurs, recedes. and recurs during the course of the Mimosis; the latter affection is apt, however, to be overlooked, and the former to be treated as a primary affection and confounded with other kinds of Diarrhea.
 - 127. Case XXXVI. Mr. R. aged 46. He has suffered from the symptoms of the Mimosis Acuta during three months. The affection varied much in severity at different periods. There were unusual weakness, faintness, weariness, aching pains, loss of flesh, vertigo, and affection of the sight, a most tainted breath, swollen gums, ptyalism, a loaded and swollen

tongue, coldness of the hands and feet, &c. Seven days ago, a regular state of the bowels is said to have yielded to Diarrhæa, with frequent, scanty, mucous, and various stools, which are sometimes fœtid, sometimes with much filamentous substance, and sometimes bloody. There are great indisposition and an incapacity for the slightest exertion.—This affection yields slowly to the pil. hydrarg. with gentle purgative medicines, the stools becoming gradually, first fæculent, and then more and more natural.

- 128. ICTERUS has occurred in six persons, in connexion with symptoms of Mimosis Acuta, within the last year. It has appeared principally in the sedentary; it has seemed to depend on a loaded state of the intestines; and it has yielded to active doses of calomel and other purgative medicines. This appears to be by far the most frequent form of Icterus, and is observed in persons of all ages,—in very early youth, and in adult age.
- 129. Cases XXXVII. and XXXVIII. D. G. and S. P. aged about 28, and fellow-workmen in the same shop of lace-frames, have each been affected with Icterus supervening on symptoms of Mimosis Acuta. The Icterus was relieved in both by active doses of purgative medicines, and the primary affection has been totally removed, and any recurrence of it obviated, by perseverance in milder doses of the same remedies.

- tary habits, was observed during several months, to lose her wonted appearance of health and strength;—the countenance became somewhat paler, and thinner; the mouth, tongue, and breath, loaded and disagreeable, and the bowels somewhat constipated. At length Icterus supervened, the skin and tunica albuginea became greatly tinged with yellow, the urine deep coloured, the motions pale, and the bowels torpid. The affection was soon removed, and the health greatly improved, by efficient purgative medicines, and she is now ruddy, plump, and active.
- 131. Case XL. Mr. B. aged 55, has twice laboured under Icterus as a form of Mimosis. Once he was affected with vertigo, fluttering, nervousness, intermittent pulse, a loaded state of the breath, mouth, and bowels, and Icterus; he was incapable of exertion, and perspired profusely on attempting it; the second time there were violent pain of the head, and excruciating pain in the epigastre extending to the back; in both cases the Icterus soon yielded to purgative medicines, and in both there were the usual symptoms of the Mimosis Acuta.
- 132. Sometimes there are Symptoms of an AF-FECTION OF THE BLADDER. The Mimosis Acuta has begun with these symptoms in several instances, and it has been conjoined with them in some part of its course. In one case the symptoms of affection of

the bladder seemed to depend on a loaded and irritable state of the lower intestine, and soon yielded to the remedies of the Mimosis Acuta.* See § 110.

- appearances of an attack of Tic Douloureux with the symptoms of the Mimosis Acuta, the tongue, the breath, the bowels, and the urine being exceedingly loaded and tainted. In two cases the pain was promptly removed by emetic and purgative medicine. One of them was of the most severe and excruciating kind.
- 134. The next complications with the Mimosis Acuta to be noticed, are, Some Convulsive and Spasmodic Affections. I shall illustrate this subject simply by the brief detail of cases in exemplification:—
- denly seized in the following manner:—the face and limbs were affected with convulsive motions; the countenance was extremely distorted, the mouth, nostrils, and cheeks being violently and convulsively drawn to the right side; several persons were occupied in restraining the violent convulsions of the limbs; the tongue, breath, and bowels were loaded. The immediate attack was relieved by opening the temporal artery, and any return was prevented by the use of purgatives.

^{*} See Bell's Reports, V. II. p. 64.

- 136. Case XLII. Mr. A. of B. became affected with an irregular spasmodic distortion of the mouth, which was drawn, from time to time, into the most extraordinary forms. The tongue was loaded, the breath tainted, and the bowels disordered.—The affection was removed in a few days by mild purgatives, consisting principally of the pil. hydrarg. and rhubarb.
- 137. Case XLIII. Mr. B. is subject to a continued state of Mimosis, which is liable, from cold or other causes, to be considerably augmented. At these times, Mr. B. experiences a considerable degree of hesitation and difficulty in articulation, an affection which invariably recedes with the primary disorder.
- with violent and most painful cramps of the calf of the right leg and of the thigh. The muscles of the thigh were drawn into a perfect ball. The spasms were somewhat continued, and so violent as to flush the face, cause the respiration to be forcibly arrested, and induce great expression of pain and suffering. The tongue was extremely loaded; the breath fœtid; the digestive organs much disordered. He was ordered half a drachm of ipecacuan, and a calomel purge, and was immediately relieved.
 - 139. Case XLV. Mr. H. of N. on the Soar, had been affected during four weeks, with violent

and painful spasmodic contractions of the muscles of the right thigh. This thigh had long been diseased, a disease of the hip, in early youth, having impeded its growth, and the muscular substance about the thigh, having more recently, although not within the last six years, formed the seat of suppurations.-The spasms were momentary only, but extremely severe, and the pain excruciating; they were apt to be repeated on moving, or on retaining one position for too long a period; they were also induced by any surprise or hurry of mind,-by the fear of being touched on this part, and, once, most severely, on being startled by a stone thrown by a boy against an outer wall near his bed. The sleep was interrupted, and but a temporary respite was obtained, and this at first only, by laudanum with æther. The tongue was little affected; he was low-spirited and nervous; the bowels were loaded and disordered. His complaint was growing daily more and more severe.-It was removed in about a week, by the pil. hydrarg. given every second night, the sulphat of magnesia with infus. sennæ, every morning, and the effervescing saline medicine. He has now, July 3rd, been in a state of progressive convalescence for a fortnight, recovering gradually the flesh and strength he had lost before. This patient still remains free from complaint, September 18th.

140. CASE XLVI. Miss M. aged 15. She has been subject, during three springs, to the follow-

ing affection, which is said to have yielded spontaneously each time, in the course of about three months
after its commencement. The hands, and especially
the right hand, are apt to be affected with a rather
continued spasmodic contraction; they are not quite
closed, but any thing held in them is not easily removed from the grasp. The contraction comes on,
recedes and recurs, without any warning. The tongue
is much loaded, and, with the gums, considerably
swollen, and it is marked by the teeth; the breath is
fœtid; the bowels confined; the catamenia have not
yet appeared. The affection has been removed by a
month's perseverance in purgative medicines.*

- 141. Case XLVII. J. A. aged 10, has been subject for a considerable time, to the following affection;—the head is moved rapidly in a repeated, rotatory manner, from side to side; the arms and legs are subject to a similar rapid spasmodic action of their muscles; when he sits down, the rectus muscle of each thigh, but especially of the right, leaps continually, a movement best felt on applying the hand about their middle part.—This affection was gradually removed by calomel purgatives.
- 142. Case XLVIII. I have seen a similar movement of the head in a married woman about 40, which was increased by any mental affection, and particularly when remarked by a stranger. This patient

^{*} See a case in Bateman's Reports, p. 119.

did not take any medicine.—I have also been informed of a similar instance of spasmodic affection in a lady about 50.

- 143. Trismus; a state of locked hand; and a continued cramp of the foot, are not very uncommon in connexion with a similar state of disorder; but it does not appear necessary to particularize these affections.
- 144. Convulsive and Spasmodic Affections may arise from Disease within the head, from Local Irritation, from a state of Mimosis, &c. and it is of the utmost importance to ascertain the source and seat of the affection,—" neque enim credo, posse eum scire, quomodo morbos curare conveniat, qui, unde hi sint, ignoret."
- 145. In place of a spasmodic affection, there is sometimes Paralytic Weakness of the Muscles, especially of the lower extremities. This affection very insidiously and gradually deprives the patient of the use of the limbs. It is often attended by a feeling of stifness, increased on exposure to cold; and there are sometimes twitchings, or slight, rapid contractions of the muscles. The paralysis has in different cases affected the bladder, the sphincters, and the upper extremities even.
- 146. This species of Paralytic Affection must be cautiously distinguished from Paralysis arising from Disease of the Head or Spine. The Di-

agnosis must be established by observing the History and Symptoms, and the Effect of Remedies, and by contrasting them with the characters of local affection of the Head and of the Spine, especially the latter.* The presence of the symptoms of Mimosis, and the absence of discoverable local disease, should lead to the trial of the remedies of the former affection; and this experiment will afford another source of Diagnosis.

- 147. There is a singular affection, which arises from a disordered state of the stomach and bowels which has not I think been noticed. The face, and some parts of the surface of the body, become suddenly swollen and puffed.
- 148. In one case the face and eye-lids were greatly swollen and disfigured. The patient had been at a feast, and felt indisposed some days before. In about half an hour the swelling had begun to subside a little; and it was totally removed by an emetic and purge. This patient had had a similar affection a year and half before.
- 149. In another case the face, surface of the chest, &c. had become suddenly and enormously puffed up. The extremities were cold, the pulse almost imperceptible, and the patient appeared to be sinking. The attack was compared to an explosion. The affection was removed by an emetic and purgative medicine.

^{*} See the treatise on Diagnosis, Part II. § 206. and particularly Abernethy, V. I. p. 85.

III. THE LOCAL COMPLICATIONS.

- 150. In § 41, I have enumerated some Topical Affections as occurring in cases of Mimosis; they are principally if not exclusively observed in the protracted instances of that affection. I do not deem it necessary to give any illustrations of those affections; these will, I think, be afforded by each practitioner's own experience. I shall only add here an account of a diseased appearance of the Probabia, and of an ulcerated state of the Tongue, which I do not remember to have seen described.
- 151. The first is an affection of the Prolabium and immediately adjoining skin. It consists of a repeated, dry, splitting, and exfoliation of the cutis of these parts, and occupies a ring of about one fourth of an inch diameter, all round the mouth; it varies in severity at different times; and it is generally long continued.—I have seen this affection in a boy, in a young lady aged 15, in two married ladies, and in a servant. It sometimes occupies severely the whole of the prolabium and a portion of the adjoining skin; and it is sometimes more partial and less severe.
- 152. CASE XLIX. The case of ulceration of the tongue occurred in W. P. aged 27, framework-knitter. Upwards of three months ago a small deep ulcer formed near the point and another on one side of

the tongue. In about three weeks six other ulcers formed in different parts of the surface of the tongue. He took emetic and purgative medicines, which nearly removed the affection, the ulcers soon healing. In a week the affection again returned, in the form of a hard lump which suppurated leaving a deep ulcer. The tongue swelled and was loaded; and there was a copious flow of saliva. I first saw him four weeks The tongue was swollen, loaded and foul. There were two deep ulcers, one near the point, the other on one side. On the left side of the middle of the tongue there was a hardness about the size of a nut; this afterwards suppurated and formed a similar ulcer. The gums and health not much affected. Today, August 28, the ulcer at the point of the tongue is better, and the tongue is cleaner. The others are much the same. In a few weeks the ulcers healed under the influence of emetic and purgative medicines.

IV. NOTE ON A CACHECTICAL FORM OF MIMOSIS.

- 153. The Mimosis Acuta has appeared to degenerate in some cases into a state of Cachexia, and has been complicated not only with the diseases enumerated, § 41, but with other morbid affections, —especially of the skin, the mouth and throat, the periosteum, the absorbent glands, &c. Some of these have resembled syphilitic affections; others have exhibited very different appearances.
- 154. This subject is full of interest, and still presents a wide and extensive field for inquiry. I may perhaps be permitted to recommend to the young student a careful perusal and comparison of the works referred to below;* and of the following Case with those given by Willis, de Scorbuto, Cap. XI, by Mr. Abernethy, on Diseases resembling Syphilis, Sect. III, and in Dr. Bateman's Reports, p. 259.
- 155. Case L. E. M. aged 35, a framework-knitter, tall, stout, and healthy, was employed in
- * Willis, and the contemporary writers, on Scorbutus; the essay on the acceptation of the term Scorbutus, Edinb. Med. Journ. V. xvi, p. 204; Abernethy on the Constitutional Origin of Local Diseases, and on Pseudo-Syphilis; Pearson on the Cachexia Syphiloidea; Willan's Reports, on Asthenia; Dr. Bateman's Reports, on Asthenia, and on the Cachexia Syphiloidea; Dr. Philip's paper already referred to, p. 63; Dr. Hennen's paper in the Edinburgh Medical Journal, Vol. XIV, p. 201; Dr. Ayre on Marasmus, on ozæna; Dr. Crampton's paper in the Dublin Hospital Reports, V. I; &c. &c.

1815, in the most active and laborious manner, in hay-making; he was exposed to great heat, underwent great fatigue, perspired profusely, and drank copiously of beer and ale. He became affected with weakness, listlessness, loss of flesh, nocturnal perspiration,-headach and vertigo,-loss of appetite and Icterus, with pale-coloured stools and deep-coloured urine.—He recovered from these complaints, but in the year 1816 had the misfortune to break his leg. In consequence of this accident and the subsequent confinement, he became and remained indisposed, and gradually lost flesh, and from 14st. weighed between 12st. and 13st. only, and experienced, on taking cold, a loss of appetite and strength, with an inability to work, not known before. In November, 1817, he underwent much bodily exertion, and remained exposed to the cold and damp. He took cold, and became affected with hoarseness, sore throat, and cough, with ædema of the ankles. These symptoms ceased, except the ædema, which receded however gradually, but he remained affected with the following complaints which are copied from his own account of them: - 'loss of flesh, and of strength; a feeling of internal weakness; feverishness, a parched and dry state of the throat, and sometimes of the tongue; sensibility to cold, chilliness, tendency to perspiration,-especially in the night; headach; sleepiness; dulness of spirits; nervousness; fluttering at the heart and about the stomach; cough; dyspnœa; a clammy tongue and mouth, and fætid breath; loss of appetite, sense of load at the stomach, occasional rejection of food, constipation, and pain in the chondiliac regions.'—This patient became much better from the use of gentle purgatives of calomel, rhubarb, and Epsom salt. But in August, 1818, he became affected with severe and continued diarrhea, with a loss of flesh from 11st. 2lbs. to 10st. 6lbs. and loss of strength, and aching and weariness, and pain in the shoulders, sides, and legs .- He again recovered under similar remedies, but in November, 1818, was taken with feverishness, attended with a parched tongue and mouth, some delirium in the night, and further loss of flesh and of strength, and, at the same time, there occurred an extensive Ulceration of the back part of the Pharynx, and a considerable discharge of bloody mucus from the Nostrils. He was reduced from 10st. 9lbs. to 9st. 2lbs. This state continued. and Icterus again occurred, with the usual appearance of the tunica albuginea, skin, urine and stools.

of appearances and affections in this poor sufferer's complaint. 1. Swarthiness of complexion; 2. feverishness, with parched throat and mouth, and heat of the forehead and legs; 3. tendency to perspiration; 4. quivering of the chin and lips in speaking, similar to that observed before shedding tears; 5. tremor; 6. fluttering; 7. loss of flesh; 8. discharge of bloody mucus from the nostrils, with ulceration; 9. ulceration of the throat; 10. Icterus; 11. discharge of much blood and mucus,

from the bowels, preceded and attended by pain of the abdomen, with tenesmus and forcing; 12. the stools, otherwise, light-coloured; 13. some anasarca; 14. boils; 15. painful ulcers on the legs.

complaints abated, with the feverishness, and he recovered strength and a good appetite, On January 27th, 1819, I made the following report. The Nostrils and Throat are nearly free from ulceration. The Icterus has disappeared. The appetite is improved, the tongue clean, the mouth more agreeable, the bowels regular, the stools less pale, the urine pale and clear. He is improved in health, strength and flesh, weighing 9st. 8lbs. In March and May I have reported my patient getting gradually better and better, and at present he has indeed little complaint. He weighed on June the 28th, 1819, 11st. 3lbs. and on August the 2nd, 11st. 7lbs.

SECTION III. THE CAUSES AND TREATMENT.

I. THE TREATMENT OF THE GENERAL AFFECTION.

Acuta is Sedentariness. This affection is therefore frequently observed in persons of studious habits, or of sedentary and inactive occupations; it has occurred particularly in literary persons, in persons engaged in the sedentary employments of the lace-frame, the stocking-frame, the tambour, &c. in tailors, and in women.

159. WATCHING, FATIGUE, ANXIETY, ERRORS IN DIET, CONFINED AND IMPURE AIR, AND WANT OF CLEANLINESS, are also accessary causes of the Mimosis Acuta. This affection has sometimes appeared to have been immediately induced as an effect of cold, of a fit of intemperance, of indigestible diet just taken, or of exposure to much heat; it has been particularly frequent during the hot summers of 1818 and 1819. And it has occurred from alarm, or accident.

160. The treatment of the Mimosis Acuta in its simple form, embraces the administration of EMETIC

and Purgative Medicines, a particular attention to Diet, Air, Exercise, Bathing or Sponging, and Sleep, and great caution in avoiding the Causes of this affection just enumerated. On each of these points I shall make a few observations.

- 161. In the severer cases an Emetic of Ipecacuan has appeared highly beneficial. To insure its early operation plenty of weak tepid tea may be taken before the emetic draught.
- 162. The Purgative Medicines which I have generally employed have been calomel, the pil. hydrarg. rhubarb, aloes, senna, the sulphat of magnesia, and pure magnesia. They must be varied according to the previous state of the bowels, and to their effect on the alvine evacuation. This previous condition of the bowels is a state of Constipation, of Diarrhæa, of Tenesmus, or of Melæna, §§ 34, 126, 121.
- 163. In the case of Constipation an active dose of calomel may be given, and if its effect is such as was desired, it may be repeated at first every fifth night, and afterwards once a week, or once in ten days, always insuring its operation, if necessary, by a solution of the sulphat of magnesia in an infusion of senna, or by pills of aloes and rhubarb. This medicine should also be repeated on the intermediate days. The object I have ever had in view has been to avoid purging, but to induce A LARGE, COPIOUS, AND CONSISTENT

to effect this, I have considered it at once as a source and criterion of increased strength or health of the alimentary canal. Frequently, however, when the patient is very feeble and infirm, the medicine appears to accumulate without operation, and the effect on the bowels is apt, by a repetition of the dose, to be violent; this event may in general be obviated, either by the administration of a proper enema at the time when the medicine was expected to move the bowels, or by inserting into the rectum a suppository of tuffy or inspissated molasses, or of soap and honey, or other similar substance.

dark-coloured motions is by no means uncommon. This state of the bowels appears to me to have yielded best to the pil. hydrarg. given every third or fifth night, and a dose of rhubarb and calcined magnesia every morning or twice in a day. And, indeed, in all cases where the more active purgative medicines have been employed in the commencement, rhubarb appears to be particularly adapted, from its tonic properties, to complete the cure. In some cases, the state of diarrhœa has demanded the use of opium, which I have then generally united with the pil. hydrarg. The object should still be to induce copious and solid evacuations, and to restore them to their natural appearance, form, colour, and odour.

- 165. In the case of Tenesmus a small opiate enema or suppository affords almost immediate relief; the purgative medicine must of course be given at the same time, and in this case the oleum ricini has a decided advantage.
- 166. The state of Melæna is generally removed by the action of purgative medicines; the pil. hydrarg. and calomel have, however, appeared to be particularly efficacious.
- 167. The state of bowels above described, must be insured until the patient has decidedly recovered his flesh and strength; and it is incumbent to explain to him the indispensible necessity of attending to this function when he again returns to his usual occupations; for this complaint, like so many others, is particularly liable to yield imperfectly only, or to return from a repetition of its causes. In the more continued form of Mimosis Acuta this attention to the state of the bowels is still more necessary.
- 168. I have mentioned the approach, in form, of some cases of Mimosis Acuta, to the appearances of Cachexia, § 153; in these and similar cases, and, indeed, in some other instances, I have been induced to prescribe the sarsaparilla.
- 169. When the operation of purgative medicine induces weakness or faintness, a draught with the tinct.

opii, or tinct. camphoræ comp. and the sp. ammoniæ aromat. has afforded great benefit.

170. It appears scarcely necessary to make any observations respecting the absolute necessity of an assiduous attention to the DIET of the patient affected with the Mimosis Acuta; this subject is of acknowledged importance, the principles of dietetics are universally known, and little remains but to enforce them with instance and assiduity. The meats cannot be too simply done. Lean beef or mutton, and fowl, are the most unequivocally good; goose, duck, pork, veal, and even lamb, are the meats most cautiously to be avoided; vegetables, and especially the uncooked vegetables, are also injurious. I have generally recommended a slice of cold beef or mutton, or a little chicken, to breakfast, instead of bread or toast; -or biscuit, if meat could not be taken, with tea or coffee: beef, mutton, or chicken to dinner: tea with biscuit in the afternoon: a little meat, perhaps, at 11 a. m. and at supper. I have stated the necessity of taking a little nutriment, frequently; of avoiding to load thestomach; and of supplying the place of the evacuations daily procured from the bowels, with proper nourishment. And, indeed, an improvement in the appetite is generally an effect of cautiously acting upon the alimentary canal by purgatives .- To these articles of diet I have added, after the bowels have been evacuated several times, a little ALE, and I believe, with the greatest advantage. Indeed more than one patient, ascribe their recovery to the use of ale, by which their strength was sustained, and they were enabled to support the effect of the purgative medicines.—The importance of an attention to the diet is sufficiently obvious from the dreadful effects frequently observed from negligence or error in this particular; see Case X. Some complicated cases have even proved fatal apparently, from errors of this nature.

- 171. The effect of the Country Air, and of Gentle Exercise in riding or travelling, has also been extremely marked in many instances. I may refer to Case IX, as an exemplification of the benefit resulting from a journey alone; and a similar advantage has been much more remarkable when the country air and exercise have been conjoined with proper remedies and diet.
- 172. Bathing in summer, and Sponging the body with cold water, or with vinegar, or salt and water, in the other seasons, is particularly advantageous. After bathing or sponging, the surface is to be rubbed with a coarsish towel until a glow of warmth is felt over the skin.
- 173. I have mentioned SLEEP amongst the remedies of the Mimosis Acuta. In cases in which wakefulness and restlessness prevailed, I have recommended an anodyne draught with the tincture of opium, or the extract of hyosciamus, at bed time. But in other

cases I have endeavoured to procure sleep by enjoining the patient to use exercise in the open air, to bathe or sponge, immediately afterwards to lie down on a sofa or bed, and to foment the feet with hot water. The following plan has been recommended to those patients whose cases seemed to require it:-on awaking in the morning a little warm tea or coffee is to be taken, the body is then to be sponged, and the breakfast finished; the patient then frequently falls asleep;—on awaking he is to rise, dress, and ride in the open air, avoiding fatigue; -on returning he is to take a little nourishment and again to lie down to sleep; - on again awaking, he is to dine, and afterwards, to ride once more in the open air, and again, on returning, lie down for a time:—the body is again to be sponged in the evening on going to bed. In this manner the weakness is often soon diminished and the patient regains his flesh and general health.

174. It is at the same time, of the utmost moment to avoid the *causes* of this affection, §§ 158, 159, and especially anxiety of mind and fatigue of body.

II. THE TREATMENT OF THE LOCAL AFFECTIONS.

- 175. IN addition to the general treatment of the Mimosis Acuta, its complications with local affection require an additional and particular attention.
- 176. The Affection of the Head, § 79, often requires general and topical bleeding, and blisters, together with a more active exhibition of purgative medicines than is usually adopted in the uncomplicated affection; and of course the necessity for these remedies is commensurate with the degree and urgency of the local affection. If there is slight pain or vertigo, which has already continued some time without increasing materially, a blister applied to the back of the neck, with leeches to the temples, and active purgatives, may be sufficient; but if the accession of this complication has occurred lately, and if the affection itself be severe, with much dulness, acute pain, or an approach to stupor, defective vision, &c. the lancet must be employed freely, cold applications be made to the shaved scalp, cupping, leeches, blistering, and active purging be instantly enforced, and abstinence and the erect position, as far as possible, enjoined.
- 177. The Affection of the Chest, § 95, frequently demands the application of blisters, which appear particularly useful in this complication; together with

the addition of the extr. hyosciami, or opium, to the aperient medicines. It is in this complicated affection, often mistaken for Phthisis Pulmonalis, that the sponging recommended by Dr. Stewart* seems to be so efficacious.

- 178. The Paroxysm of Dyspnæa, § 96, is relieved by æther and other similar remedies; but it is removed and its return is obviated by the proper remedies for the Mimosis Acuta.
- 179. Fluttering and Palpitation of the Heart, § 106, and Hiccough, § 115, are each relieved a little by æther, but only effectually treated by the remedies for the original disorder.
- 180. In the Convulsive and Spasmodic Affections, § 134, it may be necessary sometimes to emply the lancet, but in general a course of purgatives, and sometimes an emetic, are the most efficient remedies.
- 181. Pain in the course of the Colon, §§ 30, 122, is usually removed by the operation of the purgative medicines. Friction with a liniment is also useful. And if necessary, enemata and suppositories, should be administered. Some attention is due to the mode of applying the friction, and to the posture of the patient after the exhibition of the enema or suppository, when the pain is accompanied with tumor

^{*} See the Edinb. Med. and Surg. Journal, Vol. 9, p. 356.

suspected to arise from the remora of hardened fæces, § 122.

- 182. The treatment of the cases of Constipation, Diarrhœa, Tenesmus, Melæna, and Icterus, has already been noticed, §§ 164—166, 128.
- 183. In the Pains of the Limbs, § 34, I have found an anodyne ammoniacal liniment to afford great relief. This affection is sufficiently illustrated by Case X. I shall only remark here that it must be distinguished from the pain of the shoulder in Affections of the Liver, and from Rheumatism, by a reference to the character given, §§ 24, &c.

med which it may pass. But year frequently the

CIRLS DISINCLINATION FOR EXCERTION, OF PAIN

great expression of despondency and lowness.

CHAP. II.

THE MIMOSIS CHRONICA.

SECTION I. THE DESCRIPTION.

184. BY the term Mimosis Chronica, it is intended to express the state of disorder which is usually denominated dyspepsia and hypochondriasis. This form of Mimosis is intimately allied to the less severe and more continued form of the Mimosis Acuta described §§ 39—41, from which it may originate, or into which it may pass. But very frequently the Mimosis Chronica begins, and pursues a longer or shorter course, with the character about to be given.

185. The Mimosis Chronica is denoted in general, by fits of despondency and gloom, of invincible disinclination for exertion, of pain about the head, sinking at the præcordia, and heat or fulness of the stomach.

186. The countenance is liable to be rather sallow, and occasionally rather pallid; and there is often a great expression of despondency and lowness.

- 187. The tongue is sometimes affected a little, as described § 26; but it is frequently characterized by a different appearance of very minute white grains or points; it is frequently a little marked by the teeth, and sometimes slightly formed into sulci.
- 188. There is, in general, no loss of flesh; or this is at the commencement only, or on any aggravation of the affection, and slight in degree. And there is little tremor or loss of strength.
- 189. The patient is liable to be affected with vertigo or headach. There is often despondency; a feeling of total incapacity for exertion in business or society; an impatience and irascibility of temper; intolerance of noise or disturbance; and an invincible and overwhelming state of inertia or listlessness. There is gaping, and sometimes drowsiness. These affections occur in paroxysms or fits, during the absence of which the patient is more himself, but manifests a constant disposition to speak of his different complaints.
- 190. There are sometimes irregularity of the pulse, and fluttering and irregularity in the beating of the heart.
- 191. There is often a most oppressive feeling of sinking at the pit of the stomach. There are, at different times, and in different instances, heartburn, a sense of heat or burning, acidity, load, distention, in-

flation, nausea; sometimes eructation of an acid, at other times, of a nidorous taste, and sometimes the rejection of fluid, or of food. The bowels are often constipated, or there are unsatisfactory evacuations, and the patient feels a sense of load about the rectum; sometimes there is considerable pain in the bowels. The appetite is in some cases moderate, in others much impaired, and, with the digestion, various at different periods and in different instances.

- 192. There are many uneasy feelings in different parts of the body, which vary exceedingly, but always engross the patient's attention in a forcible manner.
- 193. The Mimosis Chronica, like the Mimosis Acuta, is liable to be obscured by the predominance of some particular symptom. But it does not appear necessary to particularize these cases after the ample detail of similar complications given in the last chapter.
- 194. A severe form of this affection takes place in Youths from an indulgence in onanism. It is attended with the dread of impotency, and involuntary emissions of semen during the night, and on having an alvine evacuation. There are paleness, thinness, debility, nervousness, a dreadful state of despondency, indigestion, disturbed sleep, palpitation of the heart, &c.

SECTION II. THE DIAGNOSIS.

- 195. THE Mimosis Chronica requires to be distinguished from the Mimosis Acuta, and from the insidious approach of Obscure Organic Disease.
- 196. The Mimosis Chronica borders on the more continued and less severe form of the Mimosis Acuta, described § 39; but it differs materially from the more severe form of that affection characterized § 24. Notwithstanding the painful and distressing disorder of the function of digestion, NUTRITION does not appear to be materially interrupted, and the flesh and strength generally remain little impaired, or if a little flesh is lost at one period of the affection it is regained in another, or the case is stationary. The Mimosis Chronica is, indeed, rather characterized by UNEASY FEELINGS than by REAL IMPEDIMENT to the functions. The mind, temper, and spirits, and the corporal and functional sensations are affected, whilst there is in reality little organic change, or effect induced in the process of assimilation.*
- 197. The Mimosis Chronica is distinguished from OBSCURE ORGANIC DISEASE by the contrast drawn, § 71; to which may be added, in the present case, the absence of Loss of Flesh, the early and constant Diagnostic of the latter affection.*

^{*} See the treatise on DIAGNOSIS, Part II. §§ 82, 182, 183.

SECTION III. THE CAUSES AND TREATMENT.

- 198. THE Causes and Treatment of the Mimosis Chronica are the same, mutatis mutandis, as those of the Mimosis Acuta. As the affection is of longer continuance, a more persevering use of the pil. hydrarg. and of gentle purgatives of rhubarb and aloes; and a more constant, and indeed habitual attention to diet with gentle exercise are necessary.
- 199. In this disorder the patient must submit to minister to himself in respect to the particulars just mentioned; for nothing less than this will afford permanent relief; and it is too usual for patients to expect from medicine, what can only be imparted by proper diet and exercise.
- 200. Acidity and heartburn may be relieved by a recourse to the alkalis and the calcined magnesia. The sp. ammoniæ arom. is also very useful in these, and in other distressing feelings attending this complaint.
- 201. The load after eating may be much prevented by taking warm tea, coffee, or negus.
- 202. Too often, however, this affection is protracted by sedentary habits, and cares in business; or it is renewed by errors in diet and undue exertion. It has

appeared, in some cases, to cease spontaneously, or at least after medicine had been abandoned.

203. The first object ought to be constantly to take nourishment in a simple form and small bulk. For this purpose lean meat is the best adapted. The second object should be to induce a copious alvine evacuation daily, by means of the remedies above mentioned, and by adopting a regular habit in this respect. For the latter purpose it is useful to repair regularly to the water closet every morning within a few minutes after breakfast, until this habit be acquired. By these means this distressing complaint may be mitigated, shortened, and perhaps altogether removed.

sette at 'white receives there ! him here commend tool to see we

CHAP. III.

THE MIMOSIS DECOLOR.

SECTION I. DESCRIPTION WITH CASES.

204. THE form of Mimosis of which an account is now to be attempted, and to which I have appropriated the denomination of Mimosis Decolor, appears to have been very inadequately described by medical writers under the appellation of chlorosis, which is only one of the forms of this affection. I shall, therefore, bestow great pains to place before my reader the characteristics of the different varieties, simple and complicated, of this affection.

205. And, first, I take this early occasion of stating, that the Mimosis Decolor occurs principally in Female Youth; but frequently in Married Women, both young and old; and occasionally in the young and sedentary of the Male Sex, and even in men of adult age, from the influence of sedentary habits and mental anxiety.

206. There are two forms of the Mimosis Decolor, of which the first is more acute, and the second, more

chronic and protracted. But there is every intermediate shade, and frequently, the transition of one of these forms into the other. The Mimosis Acuta, also occasionally assumes or conjoins, by an insensible gradation, the form of the Mimosis Decolor.

- 207. The more acute form of the Mimosis Decolor may be described as observing three different stages,—the Incipient, the Confirmed, and the Inveterate. It may be characterized in general as uniting a morbid state of the complexion, and generally of the surface, with recurrent pain of the head, and of the side, palpitation of the heart, fluttering, and nervousness, and some tendency to loss of flesh, and to cedema.
- 208. The Incipient Stage of the more acute form of the Mimosis Decolor is denoted by paleness of the complexion, an exanguious state of the prolabia, and a slight appearance of tumidity of the countenance, and puffiness of the eye-lids, especially the upper one. There is sometimes a tinge of green, or yellow, or of lead colour, and frequently darkness of the eye-lids.
- 209. There are great paleness of the general surface, hands, fingers, and nails; an opaque, white, tumid, and flabby state of the skin; and a tendency to cedema of the calves and ankles. And there is a certain loss of flesh.

- 210. The tongue is white, and loaded; it is swollen, marked by pressure against the teeth, or variously formed into creases or folds; its papillæ are very numerous and much enlarged. The gums and the inside of the cheeks become tumid, and the latter as well as the former, are sometimes impressed by the teeth. The breath is tainted.
- 211. The patient is generally languid, listless, sedentary, indisposed for exertion, easily overcome by exercise, nervous, and low-spirited, drowsy, dizzy, fainty, or breathless. There is generally severe headach or vertigo; the memory and power of attention are apt to be impaired; and there is sometimes heaviness for sleep.
- 212. There is, in different instances, pain of one or both sides about the false ribs, or in the hypochondriac or chondiliac regions. Sometimes there is cough, difficulty in breathing, palpitation or irregular action of the heart, or imperfect syncope, and almost universally a sense of fluttering about the præcordia.
- 213. The appetite is generally impaired. There is frequently a morbid appetite for acids, or for magnesia. The bowels are constipated, a state which sometimes leads to diarrhæa. The fæces are dark-coloured, fætid, and scanty. The urine is frequently loaded.

- 214. The catamenia become irregular, are preceded and attended by much pain of the back and region of the uterus, and sometimes, but not always, become slowly defective in quantity, and pale in colour.
- 215. In the Confirmed Stage of this affection the state of the complexion and general surface is still more marked. The countenance is still more pallid, the prolabia and the gums exanguious, or the prolabia, especially the upper one, have a slight lilac hue, and the integuments are tumid.
- 216. The skin is smooth, but becomes preternaturally dry. The integuments are puffy, opaque, and pale, or yellowish, and there is a tendency to ædema of the feet. There is frequently scarcely any further loss of flesh.
- 217. The tongue becomes clean and smooth. But it is pale, with a slight but peculiar appearance of transparency, and of a pale lilac hue. And it remains a little swollen and indented.
- 218. The patient is affected with languor, lassitude, and even serious weakness, being at once reluctant and unable to undergo fatigue.
- 219. There are often attacks of severe pain of the head, or of equally severe pain of the side, and repeated bleeding, leeches, and blisters are usually em-

ployed, affording a temporary respite from these complaints. There are also, sometimes, fits of dyspnœa, of palpitation of the heart, or of fainting, with beating of the carotids.

- 220. The pulse is rather frequent, often about 100, and easily accelerated and rendered irregular by mental emotion.
- 221. The appetite is sometimes impaired, occasionally greater than natural, and very frequently depraved, inducing a longing or constant desire for some indigestible substance, as acids or pickles, magnesia, chalk, cinders, sand, coffee grounds, tea leaves, flour, grits, wheat, &c. which the patient likes to have constantly in her mouth, or to which recourse is had when she suffers from agitation of mind.* The bowels are slow and constipated, a state which sometimes alternates with diarrhœa, and induces melæna; the stools are dark, fœtid, and scanty.
- 222. The catamenia are attended with pain, and become paler, and less in quantity, often cease, and often yield to a state of leucorrhœa which is more or less constant.
- 223. In the Inveterate Stage of the Mimosis Decolor all the symptoms assume an aggravated charac-

^{*} In the West Indies an affection similar to the present seems to prevail amongst the Negroes, who are then termed Dirt-eaters.

ter. There is a very slow but progressive loss of flesh. The languor becomes a state of permanent debility. The ædema increases and takes on the aggravated form of anasarca. The pulse becomes frequent. There are less of the appearances of mere disorder, and more of the character of disease, § 71. Or those local affections which existed in a less continued manner before, now became either permanent, or are induced by the slightest causes, and the patient can scarcely bear the most ordinary occurrences of domestic life, and perhaps remains always in bed. Sometimes there is an almost permanent pain of the head, perhaps with intolerance of light or of noise. Sometimes there is pain of the chest, with tenderness, difficulty in breathing, and cough. Frequently there are pain and tenderness of the abdomen, with sickness and constipation, or with diarrhea. Different symptoms reign in different instances,-as some hysteric or spasmodic affection; a state of locked jaw, closed hand, contracted foot, or twisted limbs; palpitation of the heart; hurried, or suspended respiration; long fits of coughing; hiccough; retention of urine.

- 224. Besides the form of the Mimosis Decolor in its more acute form, just described, there are some varieties which require to be distinctly noticed:—
- 225. 1. Sometimes there is less pallor of the countenance and prolabia, but a ring of tumid darkness round the eye, and perhaps a tumid state of the upper

lip. 2. Sometimes the complexion is of a more yellow or icterode hue. 3. Sometimes the complexion is of a peculiar lead-colour. 4. Frequently there is a mixture of the state of the complexion observed in the Mimosis Acuta, with one or other of those observed in the Mimosis Decolor. 5. There is sometimes a peculiar state of coldness, cold moisture, and lividity of the hands and fingers, and a lilac hue of the nails, the tips of which often become white and opaque. 6. The state of Mimosis Decolor, consequent on hæmorrhagy, also deserves to be distinctly noticed; there are paleness and slight yellowness of the complexion, exanguious prolabia, a greater degree of loss of flesh, and great fluttering and nervousness.

226. Such are the usual symptoms of the different stages of the Mimosis Decolor. But, as in the Mimosis Acuta, some of these symptoms are liable to be much aggravated, and to assume the form of serious Local Disease. The following list of these complications possesses therefore great interest:—

- 1. PAIN OF THE HEAD.
- 2. COUGH AND DYSPNŒA.
- 3. PALPITATION OF THE HEART.
- 4. PAIN AND TENDERNESS OF THE SIDE.
- 5. PAIN AND TENDERNESS OF THE ABDOMEN.
- 6. CONSTIPATION.
- 7. DIARRHŒA.
- 8. MELÆNA.
- 9. MENORRHAGIA.
- 10. TENDENCY TO HÆMORRHAGY.

- 11. PURPURA.
- 12. LEUCORRHEA. Model boquitage a abered y released
- 13. HYSTERIC AFFECTIONS.
- 14. ŒDEMA-ANASARCA
- color must be distinguished from a different set of diseases from those noticed in the first chapter. The complications with the Mimosis Acuta require, in general, to be discriminated from acute, those with the Mimosis Decolor must be distinguished from sudden or from insidious, local affections. These distinctions will be attempted in the ensuing section. There is also great danger, in the complications with the Mimosis Decolor, as in those of the Mimosis Acuta, of the transition into organic affection.
- 228. There is in many cases a remarkable state apparently of the capillary system, giving rise to a hæmorrhagic tendency,—to epistaxis, melæna, hæmatemesis, menorrhagia, and even purpura. In cases of the Mimosis Decolor, the blood discharged from the nose, or taken from the arm, and the catamenia, frequently become almost aqueous and colourless, so that this affection presents an instance in which the vital fluid undergoes considerable change. I have seen the blood scarcely tinge the sheets, and I have seen it resolve itself almost entirely into serum with scarcely any crassamentum. So inadequate to the subject, are all the theories of this extraordinary affection!

- 229. In the more chronic form of the Mimosis Decolor, there is a continued though variable state of sallowness, of yellowness or icterode hue, of darkness, or of a nan, squalid, or sordid paleness of complexion, or a ring of darkness surrounding the eyes, and extending a little perhaps, towards the temples, and cheeks, and sometimes encircling the mouth, without tumiddity, without the pallidness of the prolabia mentioned § 208, and without much tendency to ædema.
- 230. The general surface is more or less affected. And in the severer and more continued cases, there is frequently a peculiar deformity of the nails, which fall in irregularly in their middle part, become brittle, and break off in laminæ, so that the patient is almost unable to take a pin out of her dress. There is less tendency to ædema in this, than in the more acute form of the Mimosis Decolor.
- 231. The state of the tongue is various,—generally but slightly loaded, or perfectly clean and moist, —sometimes of a light green, light lilac, or bright red colour,—smooth and clean with enlarged papillæ,—formed into creases,—or into lobules, or other forms, —somewhat swollen in general, and slightly marked by the teeth.
- 232. There is evidently a state of general weakness and indisposition. The patient suffers from pain of the head, continued, or in paroxysms, and from

pain of the right or left side in the hypochondriac, chondiliac, or iliac regions; she is liable to vertigo on stooping, to fluttering on being startled, and to palpitation of the heart; the respiration often appears short, and the patient seems out of breath; sometimes there is paleness with the appearance of faintness.

- 233. The appetite varies. The bowels are affected with alternate constipation and diarrhea. In the afternoon there is often flatulent distention of the abdomen, which often obliges the patient to unlace her stays; or long continued rumbling or borborigmus.
- 234. The catamenia are regular in their periods perhaps, but often irregular in their flow, paler and more scanty than natural, and there is often leucorrhæa.
- 235. The Mimosis Decolor, in all its varieties, is particularly characterized by being changeable, liable to exacerbations in itself and in its complications, and to the recurrency of its symptoms even during a general amendment. Repeated BLEEDING and BLISTERS have, in general, been employed for the affection of the head, or of the side, a circumstance which becomes a deplorable characteristic of this affection.
- 236. In those females who have been married and who have had a family, the complexion has frequently, but by no means always, been icterode; in one

case the countenance was pale and exanguious, in another, wan and squalid, and in a third, there was much colour, with a ring of deep darkness round the eyes. The affection differs in no essential particular from that of younger and unmarried females; nor does it at all appear to impede conception.

- 237. In the Mimosis Decolor I have observed an eruption of urticaria, in very large, elevated wheals, sometimes solitary, and sometimes in considerable number.
- 238. I shall now proceed to exemplify the description of the Mimosis Decolor, by a series of CASES taken succinctly in the presence of the patients themselves:—
- and sedentary habits. She was observed first to lose her colour and appearance of good health, several months ago. During this interval the countenance,—cheeks and lips,—the fingers, nails, and general surface, have become pale and exanguious, and there is a degree of puffiness of the integuments, and of cedematous swelling of the ankles. The tongue is white, loaded, and impressed somewhat by the teeth. There are headach, and pain of the sides alternately, vertigo, fluttering about the heart, listlessness and sense of fatigue from the slightest exertion, which has seemed to aggravate all her complaints indeed,—con-

stipation and painful menstruation.—An attack of violent pain and throbbing of the head, with intolerance of light, noise, or disturbance, and great nervousness, has been quite removed, and the other symptoms much mitigated, by purgative medicines after copious bleedding had failed of inducing relief. The amendment in general appearance is also rapidly progressive.

240. CASE LII. S. G. aged 17. She has been long employed at the lace frame. The bowels have generally been slow. The catamenia began to appear two years ago. Three months since she began to complain of acute and continued pain of the right side of the head; she was observed to lose her colour, to become weak, and to be easily overcome and rendered breathless by exercise.—At present the countenance is extremely pale, and a little tumid; the prolabia and gums are exanguious. The tongue is swollen, not much indented, but formed into folds and plaits, perfectly clean, with enlargement of its papillæ. The hands and general surface are pale. She has lost flesh considerably. She complains much of the pain of her head, which is always aggravated on walking out. There is no dyspnæa or cough. The pulse is 96. The appetite is impaired, with a great desire for acids. The bowels are not costive just now. The catamenia are regular, but have been paler than natural at the two last periods.—This patient recovered rapidly from the use of purgative medicine. The perfect cure was, however, impeded by the necessity

of still sitting much at the lace-frame and so continuing the baneful operation of the cause of her complaints, an occurrence too frequent amongst this class of patients.

241. CASE LIII. M. J. aged 20, housemaid. April 1st, 1818. She became indisposed nine months ago, from headach, sickness, and general feebleness. Her complaints increased gradually until Christmas 1817, when she became more seriously ill, and was placed under the care of an eminent surgeon, who stated her disorder to be a Disease of the Liver. She was bled, and took saline medicines; but her complaints augmented. About this period she consulted me, and I made the following statement of the symptoms:-The face in general, and the lips and gums, are extremely pale and exanguious; the eye-lids are swollen and puffy, -a circumstance most observed early in the morning, and then both in the eye-lids and lips. The hands are exceedingly pale and colourless; the skin in general is dry and rough, and ever free from perspiration even when she takes exercise; the hands and feet are apt to be cold. The ankles are affected with ædema, especially at night, but even in the morning she can scarcely put on her shoes. The tongue is exanguious, of a pale lilac colour, semi-transparent, smooth, clean and moist. She does not complain much of weakness. She has much pain and beating across the forehead, with a sense of weight in the eyes, and sometimes vertigo, -all of which sym-

ptoms are especially experienced on walking about, and from noise. She has much palpitation of the heart, accompanying the pain of the head, and accompanied by pain of the hypochondria. She is easily hurried, and her complaints are then always reproduced if previously absent, or aggravated if present. The bowels are habitually costive. The catamenia are irregular, appearing often only after an interval of nine weeks, and preceded, for some days, by drowsiness and much pain of the loins and in the hypogastric region. She has been affected for four or five months with leucorrhœa; this ceases before the appearance of the catamenia, which gradually resume the appearance of leucorrhea. - Since the day she consulted me, she has persevered in a course of purgative medicine, consisting of aloes and rhubarb, with daily exercise, light meat diet, and a little ale. She has now a natural complexion, a return of colour, a healthy appearance of the prolabia, gums, and tongue, and all her complaints have ceased with the exception of an occasional headach.—October 22. This patient continues well; but still finds her purgative pills requisite.-The sister of M. J. has lately suffered from a slighter attack of this affection.

242. Case LIV. Miss L. aged 25. May 18, 1819. She began to be indisposed three or four years ago. She at first lost her colour and some flesh; lately she has lost little flesh, but has become very pale. She has complained at different periods of pain of the

head, -and of both sides, and her complaint has been considered Inflammation of the Lungs and an Affection of the Liver. She has also suffered from fluttering and palpitation, breathlessness, nervousness, and faintishness. Her appetite has been impaired, she has experienced distention of the abdomen, obliging her to unlace, and has had much flatus. often violent hiccough. The bowels have been prone to costiveness. At present the countenance and prolabia are pale; the complexion is apt however to be yellowish, and once she was stated to have had Icterus. The tongue, gums, and inside of the lips are pale; the tongue is clean but a little creased and with enlarged papillæ. The hands, fingers and nails are palish; the skin rather opaque and dry. There have been attacks of pain of the right side of the head, generally ascribed to the application of leeches to the side. There have been more severe attacks of pain at the pit of the stomach, of the left side over the false ribs, and of the mammæ. The palpitation and breathlessness are induced on any hurry, or on walking quick. The pulse is usually 108. The hiccough is not ascribed to any particular cause. There is some degree of swelling of the calves and insteps. The urine often deposits a copious sediment of light pink. The catamenia have become irregular, pale, scanty,-and are succeeded by fluor albus.-May 1820. Miss L. has quite recovered under a plan of purgative medicine, and at present appears and feels in perfect health,

- 143. The three following cases exemplify the first and fifth variety of the Mimosis Decolor alluded to § 225:—
- 244. CASE LV. Miss E. S. of B. aged 19. This young lady has been December 8th, 1816. slightly indisposed for several years, complaining of languor and indisposition for moving about. Two days ago the ankles became affected with considerable ædema, which has excited alarm. There is occasionally an obvious sallowness and darkness of the complexion, not very observable at this moment. The tongue is white, doughy, and impressed with the teeth, and the papillæ are prominent. The hands and fingers are affected with a little tumidity, and with a cool, clammy moisture; they are rather dark coloured, and the nails are a little livid. She is rather heavy for sleep; there is a little want of mental energy. There is little or no headach, palpitation, or pain of the side, but sometimes hurry in the breathing. The bowels are habitually costive. The abdomen often tumid. The catamenia are regular.—This patient recovered slowly, but completely, from the use of purgative medicines, a journey, and continued exercise, with a strict attention to diet.
- 245. Case LVI. Miss P. of G. aged 19. December 7th, 1816. She has been indisposed for a very considerable period. The countenance is palish; there is a ring of darkness about the eyes and mouth; the

lips are high coloured; the complexion varies, becoming much sallower at times. The tongue is loaded, white, pasty, impressed with the teeth, and displays elongated papillæ. She is listless and sedentary, and easily fatigued. There is some loss of flesh. The hands are apt to be cold, clammy, and deep coloured, with a trifling blueness of the nails; the feet are also apt to be cold. There is occasionally headach. has suffered much from pain of the left side over the false ribs, for which fifteen blisters have been applied at different intervals. She has suffered from a severe cough, which has occurred in violent fits, and which has been generally relieved by inhaling the vapour of hot water. There have been irregular palpitation of the heart, and irregularity in the frequency of the pulse, so as to induce a suspicion of Organic Disease of the Heart; but she runs up stairs with rapidity and without unusual distress, and she generally walks quick. The bowels have required opening medicine. The catamenia are natural.—A nearly similar report was made on January the 9th, 1817. The pain of the side and chest had varied its seat and severity repeatedly. Exposure to the frosty air had induced a violent fit of coughing. The other symptoms were nearly as before.—This patient recovered slowly.

246. CASE LVI. Miss M. G. aged 18. November 27, 1817.—She was affected with scarlatina nearly two years ago, and thinks she has never been perfectly well since that time. Lately she has been ob-

served to change colour and to become occasionally pale and sallow; she has become nervous; and she has been subject to variable, recurrent pain of the stomach, and of each side in the chondiliac regions. At present the countenance is pale in general; the lips are of a deep red colour; the eye-lids are occupied by an extensive ring of blackness. The tongue is white, but with very few red points, and with little swelling or indentation. The breath is sometimes tainted. has lately felt much indisposed to exertion. hands are generally cold and clammy, the nails are of of a light blue or lilac hue, and their tips are white and opaque. The memory of recent events is said to be much impaired, but not that of more remote ones; she reads, however, and remembers what is read better than events. She is liable to pain of the occiput and in two points on the forehead, the latter being of a beating kind; to pain of the stomach of a wearisome nature; and to pain of the two chondiliac regions in a more acute and less continued form. These pains come and go without manifest cause, and rather suddenly. They are all, however, most apt to be induced by any thing which hurries and affects the mind. There is no cough. She is much liable to attacks of fluttering about the heart, especially on any recurrence of distress or mental affection. Her appetite is impaired: she is particularly fond of pickles, and of cold esculent vegetables uncooked from the garden; some things are disliked, as pudding. There is occasionally a difficulty of making water. The bowels

are costive. The catamenia are rather less frequent, and less in quantity; and their appearance has always been irregular, at six weeks, or at two months.-The following is my patient's own account of her present state of health:- 'At first I had almost constantly a pain at my stomach, and sometimes in my head; my hands and feet were generally cold; I had often a fluttering at my heart, and a pain in my side, sometimes in the right, and sometimes in the left; I had also very frequently a pain in each side of my back, a little below the shoulder; my appetite was bad, and I was altogether exceedingly weak and feeble. My health is now much better; I very seldom feel any of the complaints which I have mentioned, except the pain in my head; what I suffer most from now, is the Gravel. October 22nd, 1818.'-February 28th, 1819. 'In reply to your questions I beg to state that I have passed blood five times; the first time was in the month of January, 1818; the second in March, the third in December, the fourth in January 1819, and the last on the 25th and 6th of this month. Previously to passing the blood, I have always a pain on the left side of my bowels,' (the seat of the sigmoid flexure of the colon), 'but not at any other time, except on coughing. With respect to the gravel I think that I brought you the first specimen last October, and the last in the beginning of this month.'

247. CASE LVII. Before I proceed to the exemplification of the other forms of the Mimosis Decolor,

I wish to insert the following interesting case, which appears to unite, in some degree, the character of the more continued form of Mimosis Acuta with that of October the Mimosis Decolor. Miss J. aged 18. 29th, 1818. The breath has been tainted and the mouth has been foul and disagreeable, for many years. She first began to complain a good deal three or four years ago. During this interval there has been sometimes pain of the head with vertigo and intolerance of noise; sometimes pain of one, and generally but not always of the left, side, striking to the back; sometimes fluttering at the pit of the stomach, especially on lying down; sometimes faintness; often distressing fulness from wind at the stomach; and frequently severe hiccough; sometimes much gaping; aching, heaviness, and sense of numbness of the arms. nights are restless at first; and she is heavy for sleep in the morning. She is nervous, easily fluttered and startled by a sudden noise or other occurrence. pulse is generally about 96, and often irregular. The bowels become tumid in the evening; they are habitually costive. - The face is puffed and bloated in general, and sometimes to a much greater degree than at present; the complexion is sallow, with a dark yellowness of the eye-lids and about the mouth; just now there is a deepish diffused flush of the cheeks, but this is rare. The tongue is very white, loaded, and clammy, with enlargement of the papillæ; the gums exceedingly swollen; the mouth in general, very disagreeable, and the breath extremely tainted. The tongue occasionally becomes rather dry. The hands are puffed in a remarkable degree; the fingers and nails of a light, delicate blue; the tips of the nails white and opaque. The appetite is impaired. The bowels constipated and the operation of medicines scanty. The catamenia have been irregular until the three last periods, which have been regular; their appearance is natural in quantity and colour.

- 248. The following case exemplifies the form of Mimosis Decolor as it results from hæmorrhagy:—
- 249. CASE LVIII. S. B. aged 27. Dec. 5, 1818. She has been employed in 'mending', 13 years. Her health suffered about six years ago, and she became affected with headach, and vertigo, the bowels became constipated, and the flow of the catamenia was attended with much pain. Three years ago she ceased to have the catamenia regularly, and sometimes they were absent three months. About a year and a half ago she became pregnant, and subject to leucorrhœa. She lost much blood during labour, has suckled her infant, but has very little milk. At present the countenance, prolabia, and tongue, are extremely pale and exanguious,-with a slight tinge of yellow, and a degree of darkness round the eyes. The tongue is pale, of a peculiar pale, transparent, lilac hue, clean, but marked by the teeth at the edges. The feet are slightly ædematous towards evening. The skin is generally dry, but the hands become

moist when she is fluttered. She has still headach and vertigo; and a fluttering about the præcordia on the slightest hurry; and some dry cough. She has also suffered from hiccough, and eructation. Her appetite is tolerable but she experiences a load after eating. The bowels still costive,—often two or three days without a stool. No catamenia since the birth of her infant. She still continues her sedentary work.

250. The following cases are examples of the Inveterate Stage of the Mimosis Decolor:—

251. CASE LIX. Miss M. F. aged 19, sister to the patient whose case is given § 239. Her complaints began in 1815, three years ago, and have during this interval been so various, multiform, and numerous as almost to preclude description or even a full enumeration. The case affords, however, such a remarkable illustration of the inveterate stage of the Mimosis Decolor, described § 223, as to render it highly desirable that it should be recorded.—Even before the date of the commencement of this affection, given above, the general appearance of health and spirits in Miss F. was observed to have declined. She gradually lost her colour and some flesh, became nervous, listless, dispirited, and fainty. Since 1815, she has been continually an invalid; she has kept her room for many months, and her bed, for many weeks together. The countenance, prolabia, gums, and

tongue, -- the hands, fingers, and general surface, have become exanguious; the face is rather tumid; the skin opaque and puffy; the legs and ankles very ædematous. There has been a very slow but gradual loss of flesh. The pulse is frequent, often about 100, but easily accelerated to much greater frequency. The appetite is various, and when any thing is taken, it is generally something of an indigestible nature;once she took great quantities of pickles; -at other times toasted cheese, or frizzled meat, is the only thing she can eat. The bowels have always been torpid. The catamenia have been suppressed for a considerable time.—This patient was confined to her house and to her room for several years. She has recovered, however, at length. Her sufferings have sometimes appeared to be partly corporal and partly mental. They were always aggravated by surprise, agitation, or noise; they recurred much in the evening, and in various succession. They were, as well as the REMEDIES employed, various, and multiform in the greatest degree. The detail will be given under the head of the Mimosis Urgens.

252. Case LX. presents the inveterate stage of the Mimosis Decolor, under a different aspect.— E. B. aged 46, servant. April 10th, 1817. Her complaints have come on very gradually, beginning ten years ago. She had, at first, dull pain of the head, pain of the left side, nausea, and sickness; the bowels were constipated; and there was a difficulty in void-

ing urine.-At present she has lost all of a good complexion; the countenance is affected with a variable sallowness and yellowness, but the conjunctivæ are untinged; both eye-lids are affected with a puffy swelling, and with a yellow-black hue. The lips and gums are pale; the mouth clammy; the tongue is loaded at the back but less so at the fore part, indented at its sides, and sometimes affected with swelling. The skin is always free from perspiration, and morbidly dry; on the hands it is yellowish, opaque, and sometimes puffy; the nails have become brittle and break on the slightest occasion. The ankles swell in the evening. She is liable to headach. She is easily hurried and fluttered. She has sometimes fits of viotent coughing, which continue for half an hour, sometimes with retching, but without expectoration. There is no dyspnæa. There is sometimes palpitation of the heart. The pulse is generally 96. The catamenia are regular in their periods, but without colour, and scanty; at each period the colour and quantity diminish, and the flow is attended with greater nervousness. There is no dysury now, but the urine is scanty. The appetite is various; she is fond of chewing tealeaves. The bowels are costive. Purgatives with calomel, rhubarb, and aloes, were prescribed .- On May 16th the following report was made: - The complexion is greatly restored; the hands are become moist. She has gained flesh; and all her complaints are relieved. The medicines were continued .- On August the 30th the following statement was made:-

This patient's complaints have varied, being better and worse, since the last report; but on the whole they are aggravated, and especially lately. The paleness of the countenance is augmented, with less yellowness; there is much ædema or rather anasarca, passing up the thighs even; the pulse is more frequent,—about 108; the tongue is pale, white, swollen, and indented; the bowels are kept open; the catamenia appear regularly but are colourless: the urine is clear; the appetite moderate.—After this date the anasarca gradually disappeared. But the patient remains still much as she was in May 1817.

- 253. The ensuing cases present the symptoms and appearances of the chronic form of the Mimosis Decolor:—
- 254. Case LXI. R. B. aged 33. April 7th, 1817. She has been engaged in the sedentary occupation of chevening during fifteen years. She had formerly a good colour, which began to fade about three years ago. The bowels have generally been constipated. Her complaints have augmented considerably during the last year.—At present the countenance in general is pale and sallow, and the prolabia and gums are exanguious. The hands are also pale; the skin dry and often opaque; the general surface is deprived of its wonted perspiration. There has been some loss of flesh. There is some ædema of the ankles towards evening. The tongue, formerly loaded with

a disagreeable state of the mouth and tainted breath, is now moist and clean. She suffers from variable pain of the head, which is sometimes severe, sometimes absent, sometimes gradual, and sometimes sudden in its attacks. She is nervous. She experiences violent palpitation of the heart at times. No cough. Recurrent pain of each hypochondre, and of the back. The bowels are now regular; they were formerly constipated, but never affected with diarrhea. The catamenia are less in quantity, light coloured, and attended with much pain. The urine deposits a red sediment--On April the 10th, the complaints remained nearly as before, with the exception of diminished pain of the head .- On May the 13th, all the complaints are diminished; the complexion is much improved; but the skin remains dry; the headach and nervous symptoms have disappeared; but she suffers from pain of the back; she walks with far less fatigue. -The amendment in this patient was long progressive; but she gradually relaxed in taking her gentle purgative medicines and daily exercise, and as she resumed fully her sedentary occupation she became again more or less liable to the symptoms and affections of the chronic form of the Mimosis Decolor.

255. Case LXII. C. R. aged 28, housemaid. The following complaints began six or seven years ago, and made a slow and varied progress until about a year ago, since which time they have been nearly as about to be described. At first there was a loss of

complexion and a sensible loss of flesh; attacks of pain of the head and of the right hypochondre; vertigo; fluttering, and palpitation of the heart; fits of violent, dry dough, with retching; alternate constipation and diarrhœa; the catamenia attended with great pain of the back and lower part of the abdomen, and an appearance of paleness and coldness in the countenance, with blueness of the lips; the flow, too, would begin in the morning, cease, and recur when she went to bed, the colour and quantity being nearly natural .-At present the countenance is yellowish and icterode, with a little redness of the cheeks, but a general appearance of paleness. This colour varies exceedingly, and there are alternations of paleness, and yellowness, with darkness of the eyelids, at different times. The skin is often rather hot. The nails are split, sunk in the middle, and easily broken. The tongue is clean, moist, and almost natural. She suffers from attacks of pain occupying the crown of the head, and occurring particularly when she is employed near the fire, or more than usual. She experiences vertigo on stooping. The pain of the right hypochondre often occurs during the morning, without obvious cause, and is of an aching kind. There are sometimes fits of difficulty of breathing, which are compared to that of a person out of breath from active exertion. The fluttering and palpitation are induced, like the pain of the head, by working too near the fire. The pulse is nearly natural. There has been no loss of flesh during the last year, except perhaps a little in the face. There was

formerly a little ædema, but not lately. She is affected with leucorrhœa. -These complaints are already mitigated, and the complexion improved by adopting a course of purgative medicine. The leucorrhæa is quite removed by a lotion with sulphat of zinc, inserted, by means of a scroll of linen, into the vagina.-A still further amendment takes place in this patient: -the complexion improves, the mammæ from being flaccid are now tumid and much enlarged; she has during the last three weeks gained 2lb. of flesh; the strength, capability of exertion, and the general health are equally much improved. - It is now fourteen months since she applied to me, and she continues well, being fresh coloured, jolly, and free from her many painful complaints. The nails have much recovered, and the leucorrhea is removed.

- 256. The following cases exemplify the appearance of the Mimosis Decolor in married females, and illustrate the paragraph, § 236:—
- 257. Case LXIII. Mrs. C. aged 32, and mother of seven children. April, 1816. This patient refers her complaints to mental distress; they took place insidiously, about five years ago, and three years and a half ago she was induced to apply to medicine. At first, Mrs. C. was affected with recurrent pain of the head, under the sternum, of the right side, and of the bowels. She was alternately constipated and affected with diarrhæa, but habitually of a costive habit.—At

present there is a pale yellowness of the complexion; the upper and under eye-lids are puffy and dark coloured; the conjunctiva is white; the face is apt to be somewhat swollen in the morning. The lips, gums, and tongue are pale; the tongue, formerly loaded, is now clean; the breath, formerly tainted, is now inoffensive. The hands are very pale, sometimes a little puffy. The ankles are apt to be ædematous in the evening. The skin in general is yellowish and opaque, and always dry. There are great languor, listlessness, incapability for exertion; and some loss of memory and mental energy. Formerly there was acute pain of the head; lately the pain has been duller, with vertigo, tinnitus aurium, or a momentary defect of sight. There is some heaviness for sleep. She is easily hurried and fluttered. There is much despondency. The appetite is various, -sometimes fastidious, sometimes voracious; she is particularly fond of chewing grits. The alvine evacuation is very offensive. Mrs. C's youngest child but two is four years old; the youngest but one was born a year and a half ago; the youngest nine weeks ago; the catamenia had generally appeared once between each pregnancy, but this was not the case between the last but one and the last, which was not suspected until far advanced. Mrs. C. has scarcely been able to suckle her three last children.—This patient has recovered in a very great degree from her complaint, by a persevering use of calomel, and of rhubarb and aloetic pills, and she enjoys at present a very good

state of health. Her complexion is greatly improved; but it is apt to be affected whenever she is accidentally indisposed; at such times, too, she is reminded of her former painful complaints.

258. CASES LXIV, LXV, LXVI, and LXVII. Mrs. T. aged 40, mother of a family. She is affected with great paleness of the countenance, an exanguious state of the lips, gums, and tongue, and paleness of the hands, fingers, and general surface. To these appearances are superadded the symptoms of the Mimosis Decolor.—This patient recovered from the administration of gentle purgatives .- Mrs. C. aged 27, mother of two children. This is the most marked instance of the pale icterode countenance; there is considerable darkness of the eye-lids, loss of colour in general, and an exanguious state of the prolabia, with the several symptoms of the Mimosis Decolor.—Gentle purgatives were prescribed, and the complexion and general health were gradually improved.-Mrs. S. aged 50, is affected with squalid paleness of the countenance, and darkness of the eye-lids, united with other appearances, and the usual symptoms of the Mimosis Decolor.—She has begun the plan of gentle purgatives .- Mrs. H. aged 60, was affected with the state of countenance, tongue, general surface, and symptoms, observed in the Mimosis Decolor, together with an eruption of Urticaria in large wheals, without redness, sometimes solitary, but sometimes more numerous, and appearing particularly upon the legs.

- 259. CASE LXVIII. E. T. aged 35, engaged at the sedentary occupation of 'mending.' May 28, 1816. This patient's complaints began eight years ago. During this long interval she has suffered from severe pain of the head, of the left, and sometimes of the right, side; faintness; weakness; nervousness, and despondency. The bowels have always been constipated. The catamenia appear regularly, but are very scanty, and light coloured. From possessing a fresh complexion, she has become pale and yellow. The tongue, at first loaded, is become clean, and, with the lips and gums, very pale. The face, hands, and ankles are apt to be swollen; but especially the eyelids, and particularly in the morning. The skin is very dry. About two years ago the nails became, from being strong, very thin, brittle, exfoliating in layers, and sunk in the middle.-This patient recovered in great measure by a persevering use of gentle purgative medicines. The complexion, surface, catamenia, and bowels, became more natural.-She married about a year and a half after she became my patient, and is now the mother of a fine boy.
- 260. The following case exemplifies the occurrence of the Mimosis Decolor, in its most marked form in the Male Sex:—
- 261. Case LXIX. G. S. aged 37. May 5, 1819. Nearly 4 years ago he became affected with vomiting and diarrhœa. Since that time he has lost his health,

and has been subject to disorder of the stomach and bowels-constipation alternating with looseness,with recurrent pain of the stomach and of the bowels. He lost his colour, flesh, and strength; but first, and principally his colour. Six weeks ago he was taken in the night with gaping followed by vomiting, by which some blood was raised of a black colour; the bowels were costive, and he was affected with vertigo. A month and a fortnight ago he again vomited blood, and the last time he passed blood by stool. He became extremely pale and sallow, and very feeble, and lost flesh. He complained of pain, dulness, and heaviness across the temples, and vertigo; and during the last fortnight he has complained of fluttering and strong beating of the heart: of nervousness; and some pain of the stomach and bowels.-When I first saw him the countenance was extremely pale and yellow, with some darkness of the eye-lids; the prolabia exanguious; the eyes perfectly white. hands fingers and nails were pale and exanguious. He complained as stated, of the head, and of fluttering at the heart and stomach. Pulse 96 and strong. Tongue pale, white, and furred. No thirst. Stools with flakes of mucus, without odour, and with blood of a dark colour. No disease perceptible in the abdomen on examination. No cough or dyspnæa. A degree of shivering, but no feverishness, or perspiration. He has been better and worse.—He has had hæmatemesis three times, at intervals of a fortnight, having been taken with gaping, nausea, and vomiting of

blood, mucus, and a little food. During one day, he voided blood by stool with mucus; and he has long voided stools resembling tar in appearance.-This affection began three or four years ago, -just at the time he exchanged his point-net frame in which he worked with a double thread, for one in which a single thread was worked. In the latter there is great danger of the thread breaking, by which accident the sacrifice of several inches of the work, and if several perhaps an hour's labour, was obliged to be made. On these occasions, as he states, his heart would jump into his mouth, and his whole frame received a shock from which he did not recover perhaps for a considerable time. He also became extremely timid, apprehensive, and nervous, in a very extraordinary degree. This, together with sedentariness, appears to have been the cause of his complaint. May 15. Purgative medicines have been continued regularly. He has had neither hæmatemesis nor melæna. The stools for several days were flakes of mucus only; they have gradually become more fæculent. His complexion, the colour of the hands, appetite, and general health are much improved. But there is still much weakness, and some dizziness, and after a stool, a degree of fatigue with slight fluttering. He is far less nervous. He has ceased to loose flesh the last week. September 14, 1819. He has continued uniformly to improve in strength, flesh, complexion, appetite, and in every respect, having better health now than for three or four years past. He has however still occasional returns of paleness and sallowness, of dizziness, and of the feeling of sinking. The pulse is 76 and natural in every respect.

- 262. The appearances and symptoms of the Mimosis Decolor have occurred in a boy of 14, much confined in school; in a youth of 17, occupied as a tailor; and in a young man of 20, a painter.
- Mimosis Decolor are, I think, sufficiently illustrated by the cases already given; I suppress, therefore, many others formerly intended for insertion in this work. The accession of this affection is slow and insidious in the highest degree; the incipient stage sometimes requires many months for its developement, and occupies a longer period; the confirmed stage frequently subsists for one or two years; and the inveterate stage for an indefinite period,—being long stationary, leading to the more chronic form, to the slow formation of dropsies, or organic disease, or of a viscous affection, to a slowly progressive loss of flesh and of the powers of life.
- 264. The course of this affection is irregular. Sometimes purgative medicines induce immediate relief; but this does not continue; a more slow, regular, and progressive amendment is usually more permanent. There are attacks of the urgent symptoms, and the

patient is better and worse, even during the general amendment.

- 265. The more acute form admits more readily of remedies than the more chronic, except in the inveterate stage. But all our efforts are apt to be thwarted by want of perseverance on the part of the patient, by a return to a sedentary occupation or mode of life, and by a relaxation in taking the gentle, tonic purgatives to be recommended hereafter.
- 266. June 13, 1819. I have this day had the opportunity of ascertaining by a careful dissection, that, in a case of Mimosis Decolor, long considered a case of 'Liver Complaint,' there was no perceptible disease of that or any other viscus.

SECTION II. THE DIAGNOSIS.

I. THE DIAGNOSIS OF THE GENERAL AFFECTION.

267. THE Mimosis Decolor in general, requires to be particularly distinguished from the most Insidious forms of Organic Disease. Its complications require, as I have stated § 227, to be discriminated both from the Sudden, and the Insidious, Local Affections.

268. From Insidious Organic Disease, the Mimosis Decolor is distinguished by the state of the complexion, of the general surface, and of the tongue; by the variety and multiplicity of its symptoms, and of the organs or functions affected; and by its irregular, varied, and protracted course. In Organic Disease, the countenance is pale, perhaps partially flushed, or affected as described § 74, or § 101, and there is an expression of pain, uneasiness, or disease, with an early and progressive emaciation; whilst there is an absence of the peculiarity of countenance observed in the Mimosis Decolor, §§ 208, 215, 225, 229, 236. The surface in organic affection is equally destitute of the characters given §§ 209, 216, 230, but, on the contrary, often remains nearly natural, but is apt to be affected by the different states of symptomatic febrile affection. The tongue is free from the appearances described §§ 210, 217, 231, except when there is disorder of the stomach superadded to the organic disease. There is an absence of the characters of Mimosis and of the Mimosis Decolor, §§ 1, 2, 207, 235; Contrast further the description given of Insidious Organic Disease, §§ 71, 197.

269. The state of the complexion, the tinge of surface, and the seat of pain, in the Mimosis Decolor, has very frequently led to the suspicion of Chronic Disease of the Liver. The Diagnosis is made by an attention to the contrast of symptoms drawn in the last paragraph, by observing the state of the prolabia, conjunctiva, urine, and faces, by ascertaining the recurrent nature of the pain of the side, and by a careful examination of the region of the Liver.—By these means the list of Chronic Diseases of the Liver would be considerably curtailed, for I can recall, at this moment, numerous instances of this error in Diagnosis. This subject will be resumed shortly.

menden given 56 200, 210, 230, but, on the

at top a led Reguler whose seiters asilo, excusor

II. THE DIAGNOSIS OF THE COMPLICATIONS.

- 270. The patient affected with the Mimosis Decolor is liable to sudden attacks of the affections enumerated § 226; and he is liable to these affections in a recurrent or in a more or less continued form. This circumstance renders it necessary to distinguish the complications of this affection from the Sudden and the Chronic, Local Diseases.
- 271. The Mimosis Decolor with Affection of the Head, is, in various instances, mistaken for Sudden Tendencies or Congestions of Blood with regard to the Brain, and for Insidious or Chronic Inflammation of this organ.
- 272. Case LXX. Mrs. C. of B. aged 45. She is affected with the Mimosis Decolor, characterized by a ring of deep darkness round the eye-lids, and a little cast of yellow tinge, but united with a full colour of the face and prolabia. The tongue is affected; and there are the variety of symptoms denoting the Mimosis Decolor. This lady has for many years been subject to attacks of VIOLENT PAIN OF THE HEAD, accompanied with a sense of constriction about the neck, for which bleeding, blisters, and a seton, have been variously recommended by various practitioners. The affection continued to recur notwithstanding, for very many years, always inducing the fear of some attack

of an apoplectic nature. It has yielded, however, to a persevering use of efficient purgative medicines. It has recurred from mental anxiety and the fatigue of company; and there are shedding of tears, pain at the stomach, headach, &c. the complexion being affected more on each attack. It has also more lately changed its form and induced attacks of pain at the stomach.

- the importance of these distinctions, had been successively treated as Inflammation of the Brain, and as Inflammation of the Liver,—by bleeding, leeches, and blisters to an almost incredible extent! The patient was first long subject to severe PAIN OF THE HEAD, and afterwards to PAIN OF THE RIGHT SIDE. There were superadded the appearances and symptoms of the Mimosis Decolor. The patient perfectly and even promptly recovered, by pursuing an efficient course of purgative medicines.
- 274. If, in the sudden cases, the countenance, tongue, surface, and functions be distinctly affected as in the Mimosis Decolor, there is a great presumption that there are only symptomatic pain and affection of the head.—The History of the case, and of the attack, will also assist the Diagnosis;—the patient has been repeatedly liable to a similar affection, and the attack has probably been preceded or attended by unequivocal symptoms of the Mimosis Decolor.—The effects of

remedies may also suggest the Diagnosis; on this point some observations will be made in considering the treatment of the complications of the Mimosis Decolor.

- 275. With these remarks the reader may compare and contrast the following character of the Threatening of Apoplexy .- 'An attack of apoplexy, or of paralysis, may be apprehended on the occurrence of any of the following symptoms; especially if the person, by hereditary disposition, constitution, form of body, age, or habits, be predisposed to these affections: -A sense of flushing and of fulness about the head, or epistaxis; heaviness, dull headach, vertigo; unusual drowsiness, or stupor, disturbed sleep, incubus, stertor; forgetfulness, timidity, confusion of mind; change of affections, and tendency to laughter or tears from apparent imbecility; affection of the senses, as temporary loss of sight, flashes of light, double vision, singing, or loud noises in the ears; acute pain or tenderness of the scalp. But no symptoms are so much to be dreaded, as temporary numbness or torpor of any of the limbs, or transient and partial weakness; a defect or loss of voice; a distortion and an unusual and unmeaning expression of the countenance.-There are sometimes sickness, and faintishness, and generally constipation of the bowels.'*
- 276. From Insidious or Chronic Inflammation or Disease of the Brain, the complication of the Mimosis

^{*} See the treatise on Diagnosis, Part II. § 112.

Decolor with Affection of the Head is also to be distinguished by the state of the complexion, tongue, and surface; by the complexity and variety of its symptoms; by the History of its course; and by the recurrent nature, or varying degree, of the pain, perhaps with long intervals of ease.

- 277. The former affection is, on the contrary, denoted and distinguished from this and from some other cases of complication, by a nearly natural state of the skin over the countenance and general surface, and of the tongue, and by definitiveness in the History and symptoms of the affection. Insidious Disease of the Brain comes on with pain, variable perhaps in degree, sometimes of a dull and heavy, sometimes of an excruciating kind, attended or succeeded by delirium, but generally soon leading to stupor; the senses and muscular system are variously affected; the muscular strength, however, is often little impaired; the patient frequently takes food throughout the affection; and there are often distention of the bladder or involuntary discharges of urine and fæces, even early in the disease.
- 278. The complication of Affection of the Head with the Mimosis Decolor, has, like the similar complication with the Mimosis Acuta, assumed the character of Organic Disease; but this transition is less observed in the former case than in the latter.

- 279. The Cough and Dyspnæa in the Mimosis Decolor, § 226, are sometimes rather sudden, and sometimes more continued, affections. The former case occurs most frequently in the incipient, the latter in the later stages of the Mimosis Decolor.
- Dyspnœa, the respiration is hurried and the nostrils are moved, and the cough is very distressing. This case is distinguished by the character of the Mimosis Decolor, and by yielding to the exhibition of calomel purgatives.—The more continued case of Dyspnæa and Cough is far more alarming, and is often attended by organic affection within the chest. It is distinguished from Phthisis Pulmonalis by contrasting the appearances of the Mimosis Decolor with those described §§ 100, 101, as characteristic of the former disease.—The Mimosis Decolor may, however, be combined or followed by Phthisis Pulmonalis; the prognosis should therefore be very guarded.
- 281. The Palpitation of the Heart, symptomatic in the Mimosis Decolor, § 226, must be discriminated from Organic Disease of this Organ, by contrasting the symptoms of the Mimosis Decolor with those detailed § 107, as denoting the latter affection.* This complication is most frequent. It occurs in paroxysms, and often from mental causes.

^{*} See further the treatise on Diagnosis, pp. 213-223.

- 282. Pain and Tenderness of the Side in the Mimosis Decolor, § 226, may also be sudden, or more or less continued. In the former instance it may be deemed Acute, in the latter, Chronic Inflammation of the Pleura, or of the Liver; and under this impression, bleeding and blisters are too frequently the remedies employed, and the course of gentle purgatives is too much neglected. This pain is so common as to occur in most cases of the Mimosis Decolor. It is, therefore, doubly important to establish the Diagnosis.
- 283. The character, the symptoms, the multiplicity of concomitant complaints, and the History, sufficiently identify the state of Mimosis Decolor. But the questions remain,—is this affection combined with Inflammation, or is there only an aggravation of the Pain of the Side, so usual in it? The presumption, a priori, is that the pain is not inflammatory. The Diagnosis may be further assisted by observing that the pain of Inflammation is generally regularly progressive in its formation and increase; stationary in its situation, and incessant in its duration; it induces a constant expression of pain in the countenance; it is often aggravated by change of position; and it occasions a cautious and modified state of respiration, arresting the movements of the chest, of the diaphragm, or of the part affected, and checking a full inspiration, cough, or loud expression of pain. The Pain of Side, so usual in the Mimosis Decolor, when it exists in an aggravated form, will be found, on a cautious and vi-

gilant observation, to be various and irregular in its formation, site, degree, duration, and course :- its accession is either sudden or it follows the more common state of Pain of the Side in the Mimosis Decolor; its situation is varied, the side affected is sometimes changed, or it appears to move to some other part of the course of the colon, up the chest, or towards the back; it varies in degree, being sometimes less severe, and sometimes excruciating; its duration is uncertain, and it frequently ceases suddenly, perhaps to recur; a full inspiration sometimes, although by no means uniformly, increases the pain, but a second inspiration usually induces less aggravation of the pain than the first; there is also more urgent complaint, and more expression of suffering in the manner of the patient, than in Inflammation, which represses the movements of the body, respiration, and voice, implied in the outward expression of pain. In Inflammation the pulse is also accelerated, and affected in a degree not observed in the complication of the Mimosis Acuta with Pain of the Side.*

284. The Diagnosis of the complication of the more Protracted Pain of the Side in the Mimosis Decolor, from Insidious Inflammation of the Pleura, is, if possible, still more important. The character of the case of complication will be readily understood from the preceding observations. That of the latter case shall now be attempted:—

^{*} See further the treatise on Diagnosis, Part II. pp. 172, 179, 181.

285. In Insidious or Protracted Inflammation of the Pleura, there is an absence of the features of the Mimosis Decolor so often alluded to. The nostrils are rendered acute, and the alæ nasi are raised by the influence of continued pain, and sometimes moved by the respiration. In protracted cases there are thinness of the face, and partial flushing on the cheeks. The general surface remains long natural, but if the affection continues or advances, there are emaciation, and tendency to hectic heat of the hands, and perspiration more generally. The pain is constant, or constantly induced by a full inspiration. There is often a dry painful cough, suppressed by the patient. The pulse is frequent.—This affection sometimes remains stationary for a long period; in other cases there is a progressive loss of flesh and strength, with chronic symptomatic fever; and in a third case there is the gradual formation of Hydrothorax.*

286. Pain and Tenderness of the Abdomen in the Mimosis Decolor, § 226, are distinguished from Inflammation, by the same means as pain and tenderness of the side, § 283. I shall illustrate the sudden or urgent attack by the following case, and shall afterwards give the characteristics of the case of Protracted Inflammation of the Peritonæum.

287. Case LXXII. Mrs. B. of S. aged 45, and mother of five children, has been affected with some

^{*} See the treatise on Diagnosis, Part II. § 124.

part of the following symptoms for several years, and especially since the death of Mr. B. three years ago. -The countenance is pale, wan, and yellowish; the prolabia pale; the tongue slightly lobulated, and clean; the bowels irregular .- On September the 3rd, 1818, she was seized with hurry and difficulty in breathing; with pain in the bowels, much tenderness of the abdomen under pressure, and with discharges of blood. The affection was completely removed by calomel, senna, and sulphat of magnesia. A little ptyalism was induced, and the patient abandoned all her medicines. The affection again returned. A grain of ópium was prescribed for the symptoms, which induced great sickness and restlessness. Calomel was again resorted to; and a cautious course of this and other purgative medicines, has entirely removed the more urgent complaints, and greatly improved the general appearance, health, spirits, and strength.

288. Protracted Inflammation of the Peritonæum is denoted by an expression of pain, of suffering, and of disease, in the countenance, attended by loss of flesh; the skin in general is little affected, except at last, when it is modified by a state of chronic symptomatic fever; there is a slow, but progressive emaciation; the pulse becomes very frequent; there is a diffused tenderness over the abdomen; there are attacks of retching and vomiting, of obstruction of the bowels, or of diarrhæa.—The progress of the case is various; it is sometimes protracted for eight, or ten

months, or even longer, when there are complete hectic and atrophy.*—In the Protracted Pain and Tenderness of the Abdomen in the Mimosis Decolor, there is often a loaded state of the Colon, and sometimes a perceptible tumor in the right or left side or iliac region, or in both, from the remora of fæces in the course of this intestine; and there is constipation, or diarrhæa, and sometimes Melæna or discharges of blood, sometimes florid, sometimes dark, and sometimes tar-like.†

- 289. Melæna, 226, has frequently, but by no means exclusively, occurred in married persons affected with the Mimosis Decolor. I shall illustrate the subject by the following cases:—
- 290. Case LXXIII. Mrs. E. of L. aged 40, and mother of a family. I was consulted by this lady, who resides fifteen miles from Nottingham, on the following occasion:—She had been seriously indisposed for some weeks, during some part of which period she had kept her bed, and had become extremely weak, and had lost flesh; the most urgent complaints were great irregularity and intermission of the pulse, with a severe sense of fluttering in the chest, which

^{*} The subject of Insidious Inflammation of the Encephalon, of the Pleura, and of the Peritonæum, was intended to have been illustrated by Cases in a note in this place. But it is found too extensive. It is therefore reserved for a future opportunity.

had led to the suspicion of Organic Disease of the Heart together with diarrhea and Melæna .- On inquiring into the History of the affection, I found that Mrs. E. had lost her complexion, appearance of health, strength, and some flesh, four or five years previously, and that the irregularity of pulse, and the melæna, were not new complaints. The countenance was pale and yellowish; the prolabia and gums pale; the tongue formed into lobules; the hands and general surface pale and dry. There was some affection of the head and a degree of hurry in the respiration, with some cough.—This affection was soon relieved by calomel and gentle purgative medicines; and the general health and the complexion have been greatly improved by a continuance of the same plan under proper regulations, and with strict attention to diet, air, and exercise.

291. Case LXXIV. Mrs. H. aged 55, and the mother of a family, became indisposed, from grief and anxiety, about three years ago. The causes of her complaints were repeated, and her indisposition became gradually more and more serious. She lost her complexion and flesh, became subject to ædematous swelling of the ankles and legs, to vertigo, fluttering and palpitation, and to a state of diarrhæa attended with discharges of blood, with tenesmus or forcing, and with prolapsus ani. Once the ædema assumed the more serious form of anasarca, but was much diminished, whilst the general health was much restored,

by a long journey. For the affection of the bowels, Mrs. H. was obliged to have recourse to the tinctura opii. Such was nearly the state of her complaints about a year ago. The countenance was yellow and icterode; the integuments yellowish, puffy and opaque; the conjunctivæ quite white. The pulse was a little frequent. Not a day passed without the painful and weakening attack of looseness, with discharges of blood, and descent of the rectum. She was recommended small doses of the pil. hydrarg. and of rhubarb; and to diminish the quantity of laudanum as much as possible. By perseverance in this mode of treatment, the general health and strength are much improved, and all the other symptoms much mitigated; she has been enabled almost to leave off her laudanum, to walk a distance of three miles to Nottingham and back again in one day, and to superintend her domestic concerns with far more ease and comfort. The complexion is still yellowish: but there is a little return of colour in the cheeks; and her amendment continues obviously progressive.—The prolapsus ani had taken place previously to the present complaint, but was very much augmented by it. She had also suffered from several miscarriages.

292. Menorrhagia and Leucorrhœa are frequent occurrences in the Mimosis Decolor; they are frequently the concealed cause of the inefficacy of remedies for the original affection, and of a most distressing weakness and pain experienced in the loins.—

This affection also occurs from some other causes, as frequent labours, or abortions, too long lactation, or other sources of debility; it then appears to induce the pain and weakness of the back just mentioned, fluttering, nervousness, and other symptoms of Mimosis. It is removed, I think, universally, by the remedy to be noticed hereafter.

- 293. Case LXXV. E. S. aged 40. May 3, 1819. For half a year she has been affected with menorrhagia without intermission. Before this she was long affected with Leucorrhæa. Both these complaints have been removed by the zinc lotion and opening medicine, which was required by costiveness. The tongue was lobulated and she had many of the symptoms of Mimosis,—vertigo, fluttering, &c.
- 294. The subject of Hysteric Affections, § 226, is reserved for the ensuing chapter.

is to a second of a con-

al describa orom anteres di triccoli cilon

SECTION III. THE CAUSES AND TREATMENT.

- 295. NO age, sex, condition, or mode of life, enjoys a perfect immunity from the Mimosis Decolor; but it occurs most frequently in female youth, and next, in women in the decline of life.
- 296. The most frequent cause is Sedentariness. This affection is, therefore, usually observed in schools, in females of a delicate mode of life, or of a sedentary occupation or habit. The Mimosis Decolor is the prevailing affection of those females who, in this and other manufacturing towns, are doomed to sit from morning till evening at the lace-frame, or the tambour, or engaged in mending, seaming, chevening, &c.—Servants, and especially house-maids and cooks, have also appeared to be particularly liable to the Mimosis Decolor.
- 297. CONFINEMENT; TOO LONG LACTATION; FREQUENT HÆMORRHAGIES; PROTRACTED, OR, LONG CONTINUED HABITS OF MENORRHAGIA,—AND OF LEUCORRHŒA; ANXIETY; FATIGUE; AND LOSS OF REST, have also appeared to induce the Mimosis Decolor in persons more advanced in years.

- 298. It is needless to say that, in the treatment, it is absolutely necessary to avoid or obviate the causes of the affection; otherwise the complaint continues progressive, or assumes the chronic form.—The treatment itself consists in the administration of the remedies and regimen already recommended for the Mimosis Acuta and Chronica, properly adapted to the case of the Mimosis Decolor.
- 299. Five grains of calomel may be given once a week, once in ten days, or once a fortnight. On the intermediate days, a sufficient, consistent all-vine evacuation must be procured by pills of aloes and rhubarb, or by the infus. sennæ with the sulphat of magnesia.
- 300. Daily Exercise, riding or walking, morning, noon, and evening, —and a total change of Air and of Habits must be strenuously recommended.
- 301. The same attention to DIET is necessary as before. Lean mutton, beef, or fowl; biscuit; tea, or coffee; and after a time, a little ale, or negus, are the articles of diet principally to be recommended. Vegetables in general should be avoided as much as possible, especially greens, and even bread.
- 302. For other observations on the treatment of the Mimosis Decolor, I must refer my readers to the remarks made formerly, §§ 158—183, 198—203, which

will sufficiently suggest what I might add in this place.—I have also prescribed the preparations of Iron. But as they have never been trusted to alone, I have not been able to ascertain precisely the effects of this remedy.

- 303. The Local Complications in the Mimosis Decolor generally yield to the proper treatment of the general affection, especially purgative doses of calomel. And this is satisfactory, and a confirmation of the Diagnosis.
- 304. But in dubious cases, it may be advisable to employ remedies for the local affection itself. These consist of bleeding, leeches, and blisters, especially the latter.—In the case of Affection of the Head, it is prudent first to bleed and apply a blister in order to remove the immediate attack; and to follow this treatment by the proper remedies for the Mimosis Decolor, by means of which the recurrence of affection of the head, and of the necessity for a repetition of the former remedies, may in general be obviated .- The same remarks apply to the Painful Affection of the Side or Abdomen .- In the case of Diarrhœa or Melæna, the pil. hydrarg. and calomel afford the most efficient remedies; they must, however, be conjoined with rhubarb or other gentle purgative, during the intervals of giving the former medicines.
- 305. The remedy for Menorrhagia and Leucorrhæa formerly alluded to is a continued local application of

a solution of sulphat of zinc, in the proportion of from one to two drams and a half, to a pint of pure water. A similar solution has long been employed in cases of Leucorrhæa, injected by means of a syringe. This mode of application I have found altogether inefficient in protracted cases of the affection; the application being momentary only, the effect is transitory and the remedy itself proves inefficient. The mode I have adopted, is to direct the patient to make a scroll of linen, of a form and bulk nearly sufficient to fill the vagina; this scroll is then fully imbued with the solution of sulphat of zinc, inserted, after washing with cold water, and renewed every three hours.

CHAP. IV.

THE MIMOSIS URGENS.

SECTION I. THE DESCRIPTION.

- 306. I HAVE ventured to substitute this denomination for the very objectionable term, *Hysteria*. The adjective expresses that character of hurry, and of apparently imminent danger, so usual in almost all the forms of this affection, and will be understood by every reader by being associated with an English word, to which it gives origin.
- 307. The Mimosis Urgens, besides the characteristic just mentioned, is generally denoted by combining some considerable emotion of the mind, denoted by sighing, sobbing, tears, or laughter, with a sense and expression of suffocation, and with some urgent affection of the head, heart, respiration, stomach, or muscular system.
- 308. The Mimosis Urgens most frequently occurs as symptomatic of the Mimosis Decolor, or of the

more continued form of the Mimosis Acuta. But it is occasionally induced by severe mental emotions, as excessive joy or grief; and a less curable form of the affection has been occasioned by surprise, but especially by fright.

- 309. Of the Mimosis Urgens there are three forms, —the Mild, the Severe, and the Inveterate; and there are most numerous modifications.
- 310. The Mild Form of the Mimosis Urgens subsists as a tendency to alternate high and low spirits, to fits of laughter, to frequent deep sighing, and to tears. A fit of laughter, or of crying, sometimes takes on an aggravated character; the laughing, or the sobbing, becomes immoderate, convulsive, and involuntary, and there is frequently a peculiar spasmodic chucking in the throat. The countenance changes, being alternately flushed, and pale, and denoting great anxiety. There is frequently an urgent difficulty in breathing, with much rapid heaving of the chest. Sometimes a dry, spasmodic, and violent fit of coughing occurs. There is generally a sense, and appearance, and an urgent fear of impending suffocation. In different instances there is palpitation, hiccough, retching, or borborygmus. The patient is despondent, and aggravates all her sufferings.
- 311. The severe Form of the Mimosis Urgens consists in a various attack, catenation, or combination of the following symptoms:—

- 312. The commencement, course, or termination of this and indeed of every form of the Mimosis Urgens, is generally marked, and the case distinguished, by the signs of some inordinate mental emotion,—joy, grief, or other affection,—which constitute the most characteristic symptoms of this disorder, and have appeared to be literally *hysterical*.
- 313. The attack is frequently ushered in by an unusual appearance of the countenance, a rapid change of colour, rolling of the eyes, distortion or spasmodic affection of the face. The extremities are apt to become very cold.
- 314. A state of general or partial, of violent or of continued convulsion, or of fixed spasmodic contraction, takes place, and displays every possible variety in mode and form.
- 215. The severe form of the Mimosis Urgens sometimes consists chiefly in a severe, general or partial pain and throbbing of the head. Occasionally this pain is confined to one particular spot, and is so acute as to have obtained the appellation of clavus hystericus. Sometimes there is intolerance of light and noise. Sometimes a state of stupor; sometimes delirium.
- 216. The respiration is frequently much affected:

 —an oppressive and suffocative dyspnæa takes place;

or the breathing is rapid, anxious, and irregular; or variously attended with sobbing, sighing, much rapid heaving of the chest, and sometimes with a spasmodic action of the diaphragm inducing a peculiar elevation of the abdomen, or an equally peculiar succussory movement of the trunk in general; sometimes the respiration appears to be suspended altogether for some time, the pulse continuing to beat as before.

- 317. A crowing noise, or screaming, is apt to occur in this affection. There is occasionally, Hoarseness, or even an entire Loss of the Voice, continued for some time.
- 318. There is sometimes a painful, violent, dry, hoarse cough, continued, or recurrent in paroxysms.
- 319. There is occasionally acute Pain of the Chest or Abdomen.
- 320. Palpitation of the heart, and syncope are usual affections in the Mimosis Urgens. The pulse is otherwise little affected.
- 321. There is frequently an urgent sense of suffocation, accompanied with the feeling of a ball ascending into the throat; this symptom is so peculiar as to have obtained the denomination of globus hystericus, and is considered as diagnostic of this affection. Hiccough, and violent singultus; retching and vomiting;

the sense of a ball rolling within the abdomen; borborygmus; a peculiar, great and sudden tumidity of the abdomen, apparently from flatus; constipation, &c. are usual symptoms in the Mimosis Urgens, and sometimes occur in paroxysms, and sometimes assume a more continued form.

- 322. There is frequently difficulty or retention of urine, succeeded by a very copious flow of limpid urine.
- 323. The Inveterate Form of the Mimosis Urgens, —id enim vitium quibusdam feminis crebro revertens perpetuum evadit,—consists sometimes in an almost perpetual agitation of some part of the body, the limbs, the respiration, the throat, or the stomach;—and sometimes in a state of continued contraction of the hand or foot, or of some other part. In different instances too, there is a continued state of nervousness or agitation from the slightest noise or other cause, —of paralytic, epileptic, or spasmodic disease,—or of imbecility of the mind.
- 324. The attention has, I think, been too exclusively directed to the paroxysm of convulsion in this affection. Some of the other varieties in the attack of the Mimosis Urgens, are almost equally frequent. This affection is characterized, indeed, by affecting in the same, or in different instances, singly or conjointly, ALL THE SEVERAL SYSTEMS which consti-

tute the human frame;—the organs of animal and of organic life;—the different sets of muscles, voluntary, involuntary, mixed, and sphincter;—the faculties of the mind, and the emotions of the heart;—the functions of the head, the heart, the stomach, &c. It is in thus viewing the Mimosis *Urgens, that the diagnosis is often formed between its different and very various attacks, and other affections having a different origin, but of which it is the *imitator*,—nam nullos fere non *æmulatur* ex iis affectibus quibus atteruntur miseri mortales.*

325. The varieties of the Mimosis Urgens are more numerous even than those of the other forms of Mimosis, They are also more acute, urgent, and violent. The following list, it is hoped, will be found tolerably complete:—

- 1. CONVULSION.
- 2. PAIN OF THE HEAD.
- 3. DELIRIUM.
- 4. STUPOR.
- 5. PAIN OF THE CHEST.
- 6. DYSPNŒA.
- 7. VIOLENT COUGH.
- 8. SUSPENDED RESPIRATION.
- 9. A PAINFUL AFFECTION OF THE DIAPHRAGM.
- 10. IMITATION OF CROUP; AND OF
- 11. IMPENDING SUFFOCATION.
- 12. PALPITATION OF THE HEART.
- 13. SYNCOPE.
- 14. PAIN OF THE ABDOMEN.

^{*} Vide Sydenham de Affectione Hysterica.

- 15. HICCOUGH.
- 16. RETCHING AND VOMITING.
- 17. DYSURY.
- 18. RETENTION OF URINE.
- 19. APPARENT PARALYSIS.
- 20. TRISMUS.
- 21. TETANUS.
- 22. CONTRACTED HAND.
- 23. DISTORTED FOOT.
- 24. TWISTED LEGS.*
- 326. The attack of the Mimosis Urgens, under almost every modification, is acute, sudden, and hurried; the diagnosis must therefore be instituted between this disorder, and those morbid affections which occur suddenly or in paroxysms. The diagnostics will be attempted in the ensuing section.
- 327. It appears needless to exemplify the Mild Form of the Mimosis Urgens, § 310.
- 328. The Severe Form of this affection, §§ 311 —322, is exemplified in almost all its varieties, by the following Case, continued from § 251:—
- 329. For some time there were returns every morning, of violent PAIN OF THE HEAD, with into-lerance of light and sound,—of which she is, indeed, generally very susceptible. For several months, there were evening returns of DELIRIUM, with incessant

talking, singing, crying, or laughing, -usually ending in sleep. A number of times, but irregularly, there were attacks of STUPOR, from which she could not be roused, and which would continue during two hours. There has often been PAIN ABOUT THE STERNUM, spreading over the chest and to the shoulder-points, apparently requiring, and yielding to bleeding from the arm. With pain of the chest, there has often been very rapid, SPASMODIC BREATHING, demanding free exposure to the fresh air at an open window. At other times, there has been an apparent SUSPENSION OF RESPIRATION, during many minutes, the pulse still, however, beating; the breathing gradually returned with a sort of crowing noise like the hooping cough. At one period there were evening returns of violent and INCESSANT COUGHING, which once continued from 7 p. m. till 3 a. m. without intermission; the cough ceased gradually and at length seemed scarcely to be a cough; it was relieved by bleeding, laudanum, breathing the vapour of hot water, &c. There has been PALPITATION OF THE HEART, but not in violent attacks. There has not been much FAINTING affecting the pulse, except from bleeding. For many weeks there have been constant sickness, vomiting, and IRRITABILITY OF THE STO-MACH, every thing taken being immediately rejected. For a few times, there has been VIOLENT HICCOUGH; this was arrested by a preparation of opium termed the black drop. There has ever been great consti-PATION. Once there were the SYMPTOMS OF IN-

FLAMMATION IN THE ABDOMEN, with great tenderness under pressure. There has sometimes been complete opisthotonos, the body being drawn and suspended on the occiput and toes for several minutes; then it would be drawn in a variety of other different ways. There have been general and violent con-VULSIONS of the body. There has often been LOCK-ED-JAW, -once for eight weeks together, and several times for shorter periods; -the under lip is then sometimes drawn between the teeth and bitten. hands are often clenched, and the limbs drawn in a twisted manner round each other. There has been a loss of sight, of hearing, and of the use of some of the limbs. The catheter was daily used for RETENTION OF URINE for many months, but at length it became unnecessary. She has twice complained of pain of the spine about the loins, which has, like the other symptoms, disappeared after a time.—From a perusal of this deplorable list of complaints, the young student will be prepared what to meet in similar cases of Mimosis Urgens.

- 330. The Inveterate Form, § 323, is also illustrated by the same case, for it has at length assumed that form. The following case amongst numerous others, also presents an example of this stage of this singular affection:—
- 331. Case LXXVI. M. H. aged 30. September 16th, 1813. Her mother states that her com-

plaint began fifteen years ago, and that during the last five years she has been unable to rise unassisted from her bed. During the whole of this period she has been subject to hæmatemesis, retching and vomiting, and to constipation of the bowels .- At present there are great nervousness; starting from the slightest noise; agitation on the approach of a stranger; general rapid and forcible tremor, so that the pulse can scarcely be felt; and an almost similar affection of the respiration; there is the strangest alternation of spasmodic heaving of the chest and protrusion of the abdomen; the nostrils and the head are moved at each respiration; there is often retching; and frequently retention of urine. This state of agitation is permanent. The affection is, at different times, attended with pain of the head, and a sense of uneasiness about the throat, and at the scrobiculus cordis.

symptoms, disappeared after a time.—From a perusal of this deplorable list of complaints other young student will be prepared what ai inset in saintar cases of Mimosis Urgens of the complaints of the complaints

1830. The Inveterate Form, §, 323, is also illustrated by the same case, for it has at longth assumed that form. The following case amongst numerous arbors, also presents an example of this stage of this

singular ancount is a physicial a must were audited and six and water the said water the said so will so the said so will be s

of sy open as a shall be mother states that her keen-

Lypertilar to the Ministry Urgens will occur to biling

SECTION II. THE DIAGNOSIS, WITH CASES.

of mental substant a structured the appearances vormenty"

The last a transfer with the work of the last a transfer to

the time memarion but the Epsteful of Publication

332. THE cases of the Mimosis Urgens requiring to be discriminated from other morbid affections, will be pretty fully understood on a perusal of the list given § 325. The Diagnosis is founded partly on the peculiar and different appearance of the particular cases of the Mimosis Urgens themselves; and partly on the precursory, concomitant, or successive occurrence of some unequivocal symptom of the Mimosis Urgens, and especially of the appearances of mental emotion, &c. noticed § 312, and of hurry and apparent urgency of complaint in general.

333. It is, in particular, in this manner that the paroxysm of Convulsion in the Mimosis Urgens, § 314, is to be distinguished from Epileptic or Puerperal Convulsion. The case unites the appearances described, §§ 314, 316, &c. There is probably some symptom of mental emotion; or some appearance peculiar to the Mimosis Urgens, especially the hurried and heaving respiration; or some circumstance in the History of the attack, which may lead to the Diagnosis. Otherwise the physician must wait awhile and watch the course of the affection, and the succession of symptoms; in this manner some symptoms decided-

ly peculiar to the Mimosis Urgens will occur to prompt the discrimination. In the Epileptic or Puerperal Convulsion, there is an absence of these symptoms,of mental emotion, and of the appearances formerly deemed hysterical, § 312; and the patient seems to be rather a prey to some power which exerts a violent empire over the source of sense and muscular motion. The Mimosis Urgens may excite alarm; but the Epileptic or Puerperal Convulsion presents a far more dreadful aspect,-the face perhaps becoming deeply flushed and livid, - with foaming at the mouth, more shocking distortions of the countenance and of the body, and a very different, and a more serious affection of the respiration.* By these means these affections will generally be distinguished. The student will do well to study the general aspect and course of these affections, in order to attain an aptness in their Diagnosis. poculiar to the Mimosis Urgens .-

- 334. The affection of the Head in the Mimosis Urgens, § 325, is exemplified by the following cases:—
- 335. Case LXXVII. Miss F. aged 23, the patient whose case has been already noticed, § 239, became affected on September the 21st, 1818, with severe pain, and throbbing of the head, with intolerance of light and noise; she was bled several times, without relief. I saw her on the 24th. The pain of the head still appeared to be excruciating; she cried out on its

^{*} See farther the treatise on Diagnosis, Part II, §§ 290, 291.

being moved on the pillow by the nurse; she experienced great throbbing, and intolerance of light and noise,—the curtains were drawn, and the bells in the house were ordered not to be rung. She stated too that she experienced some pain of the left side. On my entering the room, there appeared a slight mental affection,—and shedding of tears. The pulse was from 110 to 120. The countenance was pale and exanguious; the hands pale, delicate, and puffed. There was a degree of anxiety or hurry. The motions had been scanty, dark coloured, fœtid, and scybalous.—This affection yielded most favourably to purgative medicines, as already stated.

- 336. The occurrence of Delirium is not very frequent. But I have witnessed it repeatedly. The case is identified by the occurrence of some symptoms peculiar to the Mimosis Urgens.—Delirium is noticed as occurring in the case detailed § 329.
- 337. The occurrence of Stupor as a form of the Mimosis Urgens is by no means unfrequent. The case seems to be of the most urgent nature.
- 338. Case LXXVIII. Some time ago, I received an urgent call to visit a poor woman said to be in an alarming state of insensibility. She was found without sense or motion, but, in other respects, unaffected with any particular symptom. The medical attendant had prepared his lancet to open a vein in the arm.

In a short time, however, the patient recovered herself, and manifested symptoms which are peculiar to the Mimosis Urgens.—In another case which occurred in an aged woman, a vein had been opened under the idea that she was affected with Apoplexy. Some symptoms of an anomalous kind occurred, and she became affected with an unequivocal attack of the Mimosis Urgens.—In similar or dubious cases it is proper to wait, and observe the change of symptoms, and particular inquiry must be made into the History, mode of attack, &c. of the affection. Perhaps the patient soon opens the eyes, sighs, is affected with dyspnæa, or bursts into tears. In general some unexpected and anomalous symptom occurs, to denote the nature of the affection.

- 339. In the Pain of the Chest in the Mimosis Urgens, enumerated § 325, the countenance is expressive of great anxiety, hurry, and agitation, and the nostrils are moved with rapidity. The patient complains much, manifests great impatience, is urgent for relief, and calls out from the pain. The pain of the chest is extremely acute, and the part affected is described as excruciatingly tender on being touched, and the hand applied to it is usually pushed rudely away.
- 340. With or without the pain of chest, there is often an Urgent Dyspnæa; the respiration is rapid, hurried, with much characteristic heaving of the chest, sometimes with great and rapid movements both of

the chest and abdomen, and often with a peculiar hissing noise.

the Mimoria Urgens, -In another case which occurred

341. The Cough occurring as a form of the Mimosis Urgens, is exemplified § 329. It occurs in continued fits of incessant coughing; it is frequent, hoarse, and hissing;—æger creberrime tussit, fere sine intermissione, nihil prorsus expectorans.

per to wait, and observe the change of symptoms, and

- 342. CASE LXXIX. A. T. aged 24, has long been engaged in a sedentary employment, and has suffered from some of the symptoms of the Mimosis Decolor,-paleness, pain of the side, dry cough, and dyspnæa, and, when visited, was affected in the following manner. There were the utmost anxiety of countenance, and great general agitation and impatience. The breathing was hurried, with a dry hissing sound in the throat, with much rapid heaving of the chest; there was a dry, hoarse, frequent, continued cough. The pulse was little affected. The skin natural. The bowels constipated .- She was soon relieved by an æther draught and liniment, and her general health was restored by purgatives. Previously to the employment of these measures, she had been bled without any good effect, and indeed her symptoms were aggravated rather than relieved.
- 343. The case of suspended Respiration is noticed § 329. It is not the least urgent of the forms of this affection.

- 344. The Painful Affection of the Diaphragm in the Mimosis Urgens is attended by the most acute pain in the epigastrie region, extending to each side along the false ribs, and to the back; it is augmented occasionally by moving, or by the action of the diaphragm in respiration, and causes the patient to cry out; the respiration is irregular, perhaps performed by the chest alone; the nostrils move, the face is sometimes flushed, and there is often shedding of tears.
- 345. These cases are distinguished from Inflammation, by the History,—the occurrence of symptoms peculiar to the Mimosis Urgens,—the mode of attack which is sudden,—and the general aspect of the case which is hurried and urgent,—contrasted with the character of Inflammation already given, § 283.
- 346. The Imitation of Croup by the Mimosis Urgens takes place in such a manner as to deceive a cursory observer. The respiration and the cough have precisely the character of these symptoms as occurring in Inflammation of the Trachea. It is by inquiry, waiting, and by cautiously observing the case, that the Diagnosis is to be instituted. On inquiry the attack will probably be found to have been marked by some symptom or character of the Mimosis Urgens; or by waiting, some such symptoms may occur to develope the mystery. The case is sometimes so urgent as apparently to demand an operation to prevent an impending suffocation.*

^{*} See also Mr. Charles Bell's Reports, Vol. 1. p. 40. A A

- 347. Case LXXX. In a young woman, aged 15, the first symptoms which arrested the attention was a stridulous sound of the respiration; and circumstances conduced to render an attack of an inflammatory nature probable. She had been conveyed through the cold air, and appeared to be livid from cold. On being seen in bed, however, the nature of the disease became obvious, from the presence then, of globus, from the History, by which it was ascertained that other symptoms of the Mimosis Urgens had occurred, and from the absence of any affection of the pulse.—
 The patient was speedily relieved by the operation of a purgative medicine.
- 348. Palpitation and Syncope in the Mimosis Urgens are distinguished on the same principles of observing the History of the case, assisted by the observations made § 107.
- 349. The Pain of the Abdomen in the Mimosis Urgens is attended with great urgency of complaint; much anxiety and suffering; an extreme tenderness, to the slightest touch, rather than under pressure; a hurried and irregular state of breathing, &c. The countenance is expressive of an urgent anxiety; the patient is restless, impatient, and irascible, and pushes the hand although gently applied to the abdomen rudely away; the general surface, and the pulse are, at the same time, little affected; there is sometimes vomiting, or a sort of retching; the bowels are generally constipated.

- 350. Case LXXXI. M. W. aged 35, and corpulent. She has several times been suddenly attacked with the following affection. She complains of great pain and tenderness over the abdomen; but she manifests a tendency to exaggerate all her complaints, which are stated with a degree of instance and impatience not observed in inflammation. She refuses to allow any examination of the abdomen, pushing the hand rudely away. There is also a degree of hurry in the breathing. The manner is altogether different from that of a patient affected with Inflammation.*—

 To these symptoms, others peculiar to the Mimosis Urgens succeeded.—The patient was effectually relieved by the operation of a purgative medicine.
- 351. The Hiccough or the Retching is sometimes of the most violent kind, and is apt to be long continued.
- 352. The Dysury or Retention of Urine is very common as a form of the Mimosis Urgens. Its duration is usually short. But it has continued occasionally for a long period; see § 329. It is distinguished by being combined with other symptoms of this affection.
- 353. A partial, but transient Paralysis of some of the limbs, § 325, has occasionally occurred in the Mimosis Urgens.

^{*} See the treatise on DIAGNOSIS, Part II. § 164.

- 354. Trismus, Tetanus, Clenched Hand, or Contracted Foot, §§ 325, 329, are not uncommon as forms of the Mimosis Urgens. These affections generally occur, cease, and recur suddenly and unexpectedly; but they are often of considerable duration, and sometimes even appear to induce a partial dislocation.
- 355. CASES LXXXII. and LXXXII. aged 20. April 10th, 1812. She has been affected with symptoms of the Mimosis Urgens, and for many weeks, with Locked Jaw .- S. W. aged 18. September 15th, 1813, became affected with the Mimosis Urgens some time ago, from fright. She describes the globus and the fit of convulsion as having occurred repeatedly.—At present the muscles of the face are drawn into forcible contraction, inducing sharpness and elevation of the nostrils. For many weeks there have been Locked Jaw, and Locked Hands; the arms are affected with continued spasm, press forcibly against the side, and induce great tenderness; the thighs are also affected in a similar manner; they are bound bown by means of the sheet,—otherwise they would spring forcibly upwards; the patella is immoveable; the toes are drawn forcibly towards the soles of the feet.
- 356. These affections are, like so many other forms of the Mimosis Urgens, illustrated by § 329. I think them very common in protracted cases of this disorder, and could add many other instances if necessary,

357. All these forms of the Mimosis Urgens are alike distinguished by the characters given §§ 1, 2,307, and 312,—by hurry and by urgency. The occurrence of these symptoms in other forms of the Mimosis Urgens, which have possibly been omitted here, may lead to the identification of the affection; for I cannot hope that the list of the various forms of the Mimosis Urgens given § 325, is absolutely complete, or even comprehends all the important cases of this multiform disorder.—Dies quidem me deficeret, si omnia quæ hos affectus gravant symptomata, enumerare velim, tam diversa atque invicem contraria specie variantia, quam nec Proteus lusit unquam, nec coloratus spectatur Chamæleon.*

^{*} Sydenham de Affectione Hysterica.

SECTION III. THE TREATMENT.

- 358. THE principal causes of the Mimosis Urgens have been enumerated, § 308. This affection has sometimes originated also from causes of weakness, as too long lactation;—and from fatigue, anxiety and watching. I have seen this affection in the most decided form, in the Male Sex.
- 359. The Treatment embraces two objects; 1. the means of affording immediate relief in the paroxysm, and 2. the mode of prevention.
- 360. The attack of the Mimosis Urgens is relieved by æther, the sp. ammoniæ arom., opium, &c.; by stimulating liniments containing the same substances; by fomentation with hot water; and, if necessary, by blood-letting.
- 361. The prophylaxis consists in avoiding the causes, in removing the original disorder, § 308, if the Mimosis Urgens depend on this cause; but especially in the due administration of Purgative Medicines, Diet, Air, and Exercise, the principles of which have been already detailed, §§ 160, 198, 298, et seqq.

POSTSCRIPT.

- In the former edition of this work I inserted a sketch of an affection which I denominated the MIMOSIS INQUIETA. I have more recently published a distinct Essay on that subject as already stated, p. 4, note.
- Of this Essay Reviews have appeared in the Medical and Physical Journal, for July, 1820, No. 257; in the Medico-Chirurgical Review, for September, 1820, No. 2 of the Analytical Series; and in the Medical Repository, for October, 1820, No. 82, to which the reader is referred, and from which the following extracts are made.
- "There is nowhere else such an accurate and lucid view of this disorder as is here displayed by Dr. Hall, in his general abstract, and in his particular illustrations, in the cases which have occurred to his own observation. We agree with him in his assertion, that "the morbid affection in question constitutes a great proportion among puerperal cases, and a great majority among the fatal ones; and, of these fatal ones, many are daily rendered so by a mistaken use of the lancet;" and we estimate the work before us very highly, as it must lessen the frequency of such occurrences in the practice of those who will peruse it, and who have pursued the injurious practice of which the author exposes the impropriety.
- "The cases which are related in this work form a highly valuable part of it. The descriptions of the malady are given in them in very strong traits, and mark a rare talent for clinical observation. The deleterious effects of blood-letting are forcibly shown, and well contrasted with the treatment the author advises. The infatuation with which some of the practitioners who attended previously to Dr. Hall had used the lancet in some of them, is lamentable; and it makes us particularly anxious to press on the attention of our readers this excellent addition to clinical medical literature."

The London Medical and Physical Journal, pp. 64, 68.

"The line which separates spasm and irritation from inflammation, is often so obscure, that the man who could lay down an accurate diagnosis on this subject, would do a very great service to the medical profession. We believe it is a maxim pretty generally acted upon in practice, to treat a disease as inflammation, when we are in doubt whether it be or be not spasm. This, upon the whole, is a safe rule; but cases occasionally present themselves, where it is little less destructive to bleed in irri-

tation, than to stimulate in inflammation. It appears to be the object of Dr. Hall, whose talent for observation and power of discrimination are well known, to draw the attention of his brethren to the above-mentioned cases, by a plain record of facts. The motto which he has perfixed to his work, taken from Dr. Denman, is very appropriate.

"We think the profession is under great obligations to Dr. Hall, for drawing their attention to those puerperal affections where irritation borders on, or even assumes the character of inflammation, and where strong depletory measures should be cautiously put in force. The only draw-back on the utility of the publication, is the danger of its embarrasing the inexperienced practitioner, where actual inflammation obtains, combined with, or under the guise of irritation. At all events, we trust that the work will tend to effect the object designed by the Author, that of exciting the minutest attention to the discrimination of diseases."

The Medico-Chirurgical Review, pp. 195, 204.

"The lancet has been called, "a minute instrument of mighty mischief;" and so we are persuaded it is in the hands of those "bold and decided" practitioners, who never look at disease but through the medium of inflammation, and imagine that every thing must be safe while enough of depletion is secured to their patients.

"It was with much interest we learnt that a formal denunciation of the inflammatory mania had been made by so able a physician as the author; and after an attentive perusal of the pamphlet ourselves, we can most conscientiously, and we do most carnestly recommend its contents to the consideration of our professional brethren. If there are still any among the number who think undoing and unloosing is all that medicine demands or will permit, to such we would be more urgent in the expression of our wishes that they would take Dr. Hall as a guide and preceptor."

The Medical Repository, No. 82, p. 299.