Domestic medicine : or, a treatise on the prevention and cure of diseases by regimen and simple medicines. With an appendix containing a dispensatory for the use of private practitioners. ... to which are added, some important observations concerning sea-bathing, and the use of the mineral waters; with many other additions / by William Buchan.

Contributors

Buchan William, 1729-1805. Royal College of Physicians of Edinburgh

Publication/Creation

London : And J. Balfour, and W. Creech, Edinburgh, 1805.

Persistent URL

https://wellcomecollection.org/works/bx6vbxfm

Provider

Royal College of Physicians Edinburgh

License and attribution

This material has been provided by This material has been provided by the Royal College of Physicians of Edinburgh. The original may be consulted at the Royal College of Physicians of Edinburgh. where the originals may be consulted.

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org



B ng. 3.24 BII BUCHAN (1805) R21120

Digitized by the Internet Archive in 2016

https://archive.org/details/b21722031



Dr. BUCHAN'S DOMESTIC MEDICINE.

3.24


THE NINETEENTH EDITION,

TO WHICH ARE ADDED, SOME IMPORTANT OBSERVATIONS CONCERNING

SEA-BATHING,

AND THE

USE OF THE MINERAL WATERS; with many other additions.



Domestic Medicine :

OR, A

TREATISE

ON THE

PREVENTION AND CURE

OF

DISEASES

BY

REGIMEN AND SIMPLE MEDICINES.

WITH

AN APPENDIX,

CONTAINING A DISPENSATORY FOR THE USE OF PRIVATE PRACTITIONERS.

> THE NINETEENTH EDITION, to which are added, SOME IMPORTANT OBSERVATIONS concerning SEA-BATHING,

> > AND THE

USE OF THE MINERAL WATERS;

WITH MANY OTHER ADDITIONS.

WILLIAM BUCHAN, M.D.

Fellow of the Royal College of Phyficians, Edinburgh.

LONDON:

PRINTED FOR A. STRAHAN; T. CADELL AND W. DAVIES, STRAND; AND J. BALFOUR, AND W. CREECH, EDINBURGH. Omnes homines artem medicam nosse oportet. Sapientiæ cognitionem medicinæ fororem ac contubernalem effe puto. HIPPOCRATES.

Domeflic Medicine:

PREVENTION AND CURE

RECIMEN AND SIMPLE MEDICINES

TIR E'A TI

Primoque medendi fcientia, fapientiæ par habebatur. — Rationalem quidem puto medicinam effe debere. CELSUS.

Quemadmodum fanitas omnium rerum pretium excedit, omnifque felicitatis fundamentum est, ita scientia vitæ ac fanitatis tuendæ omnium nobilissima, omnibusque hominibus commendatissima esse debet.

LINAM BUCHAN M

.2001

TO

Sir JOSEPH BANKS, Bart.

PRESIDENT OF THE ROYAL SOCIETY.

SIR,

THE DOMESTIC MEDICINE having been honoured, on its first appearance, with the patronage of your learned and worthy Predecessfor, the late Sir JOHN PRINGLE, I beg leave, in a more improved state, to dedicate it to you, as a small but fincere testimony of that veneration and esteem with which I have long beheld the man who, born to ease and affluence, had resolution to encounter the dangers of unknown seas and distant climes, in pursuit of useful Science; and whose constant object has been to render that Science fubfervient to the happines and civilization of Society.

I have the honour to be, with great respect,

SIR,

Your most obedient Servant,

LONDON, Nov. 10, 1783.

W. BUCHAN.

a

r JOSEPH BANKS, Bart.

PRESIDENT OF THE ROYAL SOCIETY.

SIR,

HE DOMESTIC MEDICINE having been honoured, on its first appearance, with the autonage of your learned and worthy Fredeoeffer, he late Sir JOHN PRINCLE, I beg leave, in a nore improved state, to dedicate it to you, as a mall but success the state of that veneration and sheen with which I have long beheld the man who, born to ease and affinence, had reformion to incounter the dangers of unknown feas and diffaut parlinges, in purfuit of useful Science; and whofe enstant object has been to render that Science other in to the happines and civilization of activity.

I have the honour to be, with great refrect, SIR,

Your moft obedient Servant,

W. BUCHAN.

ADVERTISEMENT

ill empire ; and fomeorciters of the full medical

Lavii STVAA

isty

my

THE EIGHTEENTH EDITION.

fpeflive countries. I feel the full force of fo firengian incitement, to redouble my chorts, and, as far as I an

cannence in Europe have ente

THE learned and elegant ARBUTHNOT feems to have thought it a matter of no fmall complaint, that an author, after having written a book, fhould be obliged to write another to explain it. Yet, in the difcuffion of fubjects which immediately concern the health and happines of the human species, it is of importance to leave nothing obscure or doubtful, but to place every truth in so ftrong and clear a light, as to prevent the possibility of the least misconception or mistake.

Befides, with whatever induftry and exactnefs a medical writer may prepare his first effays for the public eye, he will find, in the course of farther practice, many things to correct, others to illustrate, and, perhaps, fome valuable discoveries to introduce. The art or science, of which he treats, is in a perpetual progress to perfection. If his writings do not breathe the fame spirit, it is plain that he is either too proud to retract former errors, or too indolent to avail himself of new observations and experiments.

As gratitude also is supposed to be a greater spur to exertion than even *reason* itself, I have a still more powerful motive than any of the preceding, for using

sustries .

ADVERTISEMENT.

my beft endeavours to make every new edition of this work more worthy of the public favour. It has met with the moft flattering reception in every part of the Britifh empire; and fome writers of the firft medical eminence in Europe have enlarged the fphere of its popularity, by translating it into the languages of their refpective countries. I feel the full force of fo ftrong an incitement, to redouble my efforts, and, as far as I am able, to make improvement keep pace with the rapid extension of fuch liberal patronage.

But, while my warmest acknowledgments are due to my generous friends, I must not omit taking notice of fome obligations I am under to my adverfaries alfo. On my first attempt to give a familiar form to medical advice, and to explain what every body ought to know, in language which every body could underftand, envy, felfishness, and arrogance took the alarm. All the stories of malignant invention were exhaufted, to blacken the character of the author, to lower the public opinion of his integrity, as well as of his talents, and totally to deftroy his pretentions to honeft fame. I thank those gentlemen for having taken fo much pains to excite attention to my book; and hope they will not think this expression of my gratitude ironical, when I assure them, that the fale of above a hundred thousand copies of this work has not been lefs promoted by their illiberal abuse, than by the kindest testimonies of applaufe.

There is another defcription of perfons, whom I cannot treat with fo much pleafantry, as they have left no means untried to make the credit of my book fubfervient

viii

ADVERTISEMENT.

fervient to their private intereft, at the expence of both my fame and fortune. I mean the Invaders of Literary Property, who firft endeavoured to mutilate, mangle, and disfigure the work, in hopes of being able to conceal their fraud, and give the pirated jumble fome appearance of originality by a little alteration of the title-page, and a ftring of prefatory falfehoods *. Having completely failed in a great number of thefe attempts, they now have recourfe to a ftill more fhamelefs mode of proceeding, and pretend to publifh the "DOMESTIC MEDICINE" itfelf, with bold claims to confiderable *improvements* ! A fingle glance, however, is fufficient to detect the flimfy artifice.

In order to evade the law, by acting under the fanction of an old flatute for limiting the period of copyright, they have reprinted the *early copies* of my book, publifhed between thirty and forty years ago, which, to fay nothing of inaccuracies, did not contain above half the matter inferted in the later editions. Thus they have in reality fupprefied what I had been conftantly endeavouring to correct, improve, and enlarge. Forefeeing, however, that, in confequence of all thefe omiffions, the fize of the book, when compared with mine, would expofe the deception even to perfons incapable of examining the contents, they increafed the bulk of the volume by additions, which betray an ignorance, not only of medicine, but of language.

" Like Gipfies, left the ftolen brat be known,
" Defacing first, then claiming for their own."

ix

ADVERTISEMEN'T.

It is not likely that any great imprefion can be made on fuch characters by the force of ridicule, elfe I fhould compliment them on their extreme modefty in taking out of the hands of an author, whom they profeffed to admire, the right of improving his own works. They wifhed, no doubt, to relieve him from the trouble. Their generofity went ftill farther, and made them affix his name to their curious refearches, and to the novel beauties of their ftyle, which adorn and illustrate every page. He may, indeed, think them *too civil by half*, in placing their miftakes to the fame account, and inducing the public to rely on every thing fuppofed to come from him, though pregnant with the greatest danger.

I fhould also pay due praise to the legal shrewdness of those gentlemen, in laying hold of the most imperfect editions of my book, and leaving out all the subsequent remarks, rather than violate the laws of their country. As to the infringement of property, or the injury which the reputation of the Author might fustain from his being made responsible for their blunders, the old statute they confulted, being filent on these points, led them to conceive that they had a right to publish, under the fanction of his name, the injudicious compilations of any unfortunate hireling.

It is not very eafy to put me out of humour. I am much more inclined to amufe myfelf and the reader with laughable ftrictures on fuch frauds and follies. It would be no difficult matter, however, were the fubject to be ferioufly treated, to fhew how inconfiftent it is with common fenfe and common juffice, that property acquired by the

ADVERTISEMENT.

the labour of the hand fhould be more permanent or better protected, than that which is derived from the labour of the head. Yet the one is perpetual and well-fecured, while the other is confined to a very fhort period, and left almost defenceles. In vain does the man of genius endeavour to impress upon his works the ftamp of immortality: if he lives but a few years, he may fee them cut to pieces, without any means of redress, and his very name made use of to give currency to the most contemptible forgeries.

The affection of a parent for a darling child has often been used to illustrate an author's fond attachment to his writings. What then must be the emotions of the latter, when he finds a changeling fubstituted for the genuine offspring of his brain, or fees, as it were, his mental progeny stripped of their rights and honours by the spurious issue of a pretender!

When we confider this matter in another light, and make a fair effimate of the time and money expended in the cultivation of fuch talents as may qualify a man to write for the inftruction, or the rational entertainment, of fociety, we fhall be ftruck with the injuffice of robbing him of the fair fruits of his labour. Small as his emoluments may be, they are not fufficiently guarded by law. The acts relative to this fubject were made when literary property was not deemed of much value, and confequently little care taken to fecure it. They fhould therefore be revifed; for while the exact letter of the law protects the property of authors only for a fhort term of years, it is not in the power of a judge, however

upright,

xi

ADVERTISEMENT.

upright, to give ample redrefs of any wrong done them afterwards. Every author ought to retain an exclusive right in his own compositions during his natural life at least; and, perhaps, were the question to be brought before the tribunal of reason, no fatisfactory cause could be assigned, why the advantages which we derive from the gifts of heaven, ought not to descend to our posterity, as well as revenues arising from gifts of the crown.

I cannot conclude, without informing the public, that, befides the corrections and improvements, I have enlarged the prefent edition by the introduction of above a hundred pages of new matter, including a complete Chapter on Sea-bathing, and the Ufe of the Mineral Waters, which will enable every purchafer of this work eafily to diftinguifh the genuine from fpurious and imperfect copies, and probably fave the health and lives of numberlefs individuals from being made the fport of unprincipled impoftors.

W. BUCHAN.

PERCY-STREET, January 3, 1803.

[xiii]

PREFACE.

WHEN I first fignified my intention of publishing the following sheets, I was told by my friends it would draw on me the refentment of the whole Faculty. As I never could entertain such an unfavourable idea, I was resolved to make the experiment, which indeed came out pretty much as might have been expected. Many, whole learning and liberality of sentiments do honour to medicine, received the book in a manner which at once shewed their indulgence, and the falsity of the opinion that every Physician wishes to conceal his art; while the more felsish and narrow-minded, generally the most numerous in every profession, have not failed to perfecute both the book and its author.

The reception, however, which this Work has met with from the Public, merits my moft grateful acknowledgments. As the beft way of expressing these, I have endeavoured to render it more generally useful, by enlarging the *prophylaxis*, or that part which treats of preventing difeases; and by adding many articles which had been entirely omitted in the former impreffions. It is needless to enumerate these additions; I shall only fay, that I hope they will be found real improvements.

The obfervations relative to Nurfing and the Management of Children were chiefly fuggefted by an extensive practice among infants, in a large branch of the Foundling Hospital, where I had an opportunity not only of treating the difeases incident to childhood, but likewife of trying different plans of nurfing, and observing their effects. Whenever I had it in my power to place the children under the care of proper nurses, to instruct these nurses in their duty, and to be fatisfied that they performed it, very few of them died; but when, from diffance of place, and other unavoidable circumftances,

stances, the children were left to the fole care of mercenary nurfes, without any perfon to instruct or fuperintend them, fcarcely any of them lived.

This was fo apparent, as with me to amount to a proof of the following melancholy fact: That almost one half of the human species perists in infancy, by improper management or neglect. This reflection has made me often wish to be the happy inftrument of alleviating the miseries of those fuffering innocents, or of refcuing them from an untimely grave. No one, who has not had an opportunity of observing them, can imagine what absurd and ridiculous practices still prevail in the nursing and management of infants, and what numbers of lives are by that means loss to fociety. As these practices are chiefly owing to ignorance, it is to be hoped, that when nurses are better informed, their conduct will be more proper.

The application of medicine to the various occupations of life has been in general the refult of obfervation. An extensive practice for feveral years, in one of the largest manufacturing towns in England, afforded me fufficient opportunities of obferving the injuries which those useful people fuscain from their particular employments, and likewise of trying various methods of obviating fuch injuries. The fuccess which attended these trials was sufficient to encourage this attempt, which I hope will be of use to those who are under the necessity of earning their bread by such employments as are unfavourable to health.

I do not mean to intimidate men, far lefs to infinuate that even thofe arts, the practice of which is attended with fome degree of danger, fhould not be carried on; but to guard the lefs cautious and unwary against those dangers which they have in their power to avoid, and which they often, through mere ignorance, incur. As every occupation in life disposes those who follow it to fome particular diseases more than to others, it is certainly of importance to know these, in order that people may be upon their guard against them. It is always better to be warned of the approach of an enemy, than to

xiv

to be furprifed by him, especially where there is a possibility of avoiding the danger.

The obfervations concerning Diet, Air, Exercife, &c. are of a more general nature, and have not efcaped the attention of phyficians in any age. They are fubjects of too great importance, however, to be paffed over in an attempt of this kind, and can never be fufficiently recommended. The man who pays a proper attention to thefe, will feldom need the phyfician; and he who does not, will feldom enjoy health, let him employ as many phyficians as he pleafes.

Though we have endeavoured to point out the caufes of difeafes, and to put people upon their guard againft them, yet it muft be acknowledged that they are often of fuch a naure as to admit of being removed only by the diligence and activity of the public magiftrate. We are forry, indeed, to obferve, that the power of the magiftrate is feldom exerted in this country for the prefervation of health. The importance of a proper medical police is either not underftood, or little regarded. Many things highly injurious to the public health are daily practifed with impunity, while others, abfolutely neceffary for its prefervation, are entirely neglected.

Some of the public means of preferving health are mentioned in the general prophylaxis, as the infpection of provisions, widening the freets of great towns, keeping them clean, fupplying the inhabitants with wholefome water, &c.; but they are paffed over in a very curfory manner. A proper attention to these would have fwelled this volume to too large a fize; I have, therefore, referved them for the fubject of a future publication.

In the treatment of difeafes, I have been peculiarly attentive to regimen. The generality of people lay too much ftrefs upon Medicine, and truft too little to their own endeavours. It is always in the power of the patient, or of those about him, to do as much towards his recovery as can be effected by the phyfician. By not attending to this, the defigns of Medicine are often fruftrated; and the patient, by purfuing a wrong plan of regimen,

gimen, not only defeats the Doctor's endeavours, but renders them dangerous. I have often known patients killed by an error in regimen, when they were using very proper medicines. It will be faid, the physician always orders the regimen, when he prescribes a medicine. I wish it were so, both for the honour of the Faculty and the fafety of their patients; but physicians, as well as other people, are too little attentive to this matter.

Though many reckon it doubtful whether phyfic is more beneficial or hurtful to mankind, yet all allow the neceffity and importance of a proper regimen in difeafes. Indeed, the very appetites of the fick prove its propriety. No man in his fenfes ever imagined that a perfon in a fever, for example, could eat, drink, or conduct himfelf in the fame manner as one in perfect health. This part of medicine, therefore, is evidently founded in Nature, and is every way confiftent with reafon and common fenfe. Had men been more attentive to it, and lefs folicitous in hunting after fecret remedies, Medicine had never become an object of ridicule.

This feems to have been the first idea of Medicine. The ancient physicians acted chiefly in the capacity of nurfes. They went very little beyond aliment in their prefcriptions; and even this they generally administered themfelves, attending the fick for that purpose through the whole courfe of the difease; which gave them an opportunity not only of marking the changes of difeases with great accuracy, but likewise of observing the effects of their different applications, and adapting them to the fymptoms.

The learned Dr. Arbuthnot afferts, that by a proper attention to those things which are almost within the reach of every body, more good and less mischief will be done in acute difeases, than by medicines improperly and unfeasonably administered; and that great cures may be effected in chronical distempers by a proper regimen of the diet only. So entirely do the Doctor's fentiments and mine agree, that I would advise every person, ignorant of physic, to confine his practice folely to diet, and the

the other parts of regimen; by which means he may often do much good, and can feldom do any hurt.

This feems also to have been the opinion of the ingenious Dr. Huxham, who observes, that we often seek from Art, what all-bountiful Nature most readily, and as effectually, offers us, had we diligence and fagacity enough to observe and make use of them; that the *dietetic* part of medicine is not fo much studied as it ought to be; and that, though less pompous, yet it is the most natural method of curing diseafes.

To render this book more generally ufeful, however, as well as more acceptable to the intelligent part of mankind, I have in moft difeafes, befides regimen, recommended fome of the moft fimple and approved forms of medicine, and added fuch cautions and directions as feemed neceffary for their fafe administration. It would no doubt have been more acceptable to many, had the book abounded with pompous prefcriptions, and promifed great cures in confequence of their ufe; but this was not my plan : I think the administration of medicines always doubtful, and often dangerous, and would much rather teach men how to avoid the neceffity of using them, than how they should be used.

Several medicines, and those of confiderable efficacy, may be administered with great freedom and fafety. Physicians generally trifle a long time with medicines before they learn their proper use. Many peasants at prefent know better how to use fome of the most important articles in the *materia medica*, than physicians did a century ago; and doubtless the fame observation will hold with regard to others for fome time hence. Wherever I was convinced that medicine might be used with fafety, or where the cure depended chiefly upon it, I have taken care to recommend it; but where it was either highly dangerous, or not very necessary, it is omitted.

I have not troubled the reader with an ufelefs parade of quotations from different authors, but have in general adopted their obfervations where my own were either defective or totally wanting. Those to whom I am most obliged

obliged are, Ramazini, Arbuthnot, and Tiffot; the laft of which, in his Avis au Peuple, comes the nearest to my views of any author which I have feen. Had the Doctor's plan been as complete as the execution is mafterly, we fhould have had no occafion for any new treatife of this kind foon; but by confining himfelf to the acute difeafes, he has in my opinion omitted the most useful part of the subject. People in acute diseases may fometimes be their own phyficians; but in chronic cafes, the cure must ever depend chiefly upon the patient's own endeavours. The Doctor has also passed over the Prophylaxis, or preventive part of medicine, very flightly, though it is certainly of the greatest importance in fuch a work. He had no doubt his reafons for fo doing, and I am fo far from finding fault with him, that I think his performance does great honour both to his head and to his heart.

Several other foreign phyficians of eminence have written on nearly the fame plan with Tiffot, as the Baron Van Swieten, phyfician to their Imperial Majefties; M. Rofen, firft phyfician of the kingdom of Sweden, &c.; but thefe gentlemen's productions have never come to my hand. I cannot help wifhing, however, that fome of our diftinguifhed countrymen would follow their example. There ftill remains much to be done on this fubject, and it does not appear to me how any man could better employ his time or talents, than in eradicating hurtful prejudices, and diffufing ufeful knowledge among the people.

I know fome of the Faculty difapprove of every attempt of this nature, imagining that it muft totally deftroy their influence. But this notion appears to me to be as abfurd as it is illiberal. People in diffrefs will always apply for relief to men of fuperior abilities, when they have it in their power; and they will do this with greater confidence and readinefs when they believe that Medicine is a rational fcience, than when they take it to be only a matter of mere conjecture.

Though I have endeavoured to render this Treatife plain and useful, yet I found it impossible to avoid fome

XVIII

fome terms of art; but those are in general either explained, or are fuch as most people understand. In short, I have endeavoured to conform my style to the capacities of mankind in general; and, if my readers do not flatter either themselves or me, with some degree of fucces. On a medical subject, this is not so easy a matter as some may imagine. To make a shew of learning is easier than to write plain sense, especially in a science which has been kept at such a distance from common observation. It would, however, be no difficult matter to prove, that every thing valuable in the practical part of medicine is within the reach of common abilities.

I would be ungenerous not to express my warmeft acknowledgments to those Gentlemen who have endeavoured to extend the usefulness of this Performance, by translating it into the language of their respective countries. Most of them have not only given elegant translations of the Book, but have also enriched it with many useful observations; by which it is rendered more complete, and better adapted to the climate and the conflitutions of their countrymen. To the learned Dr. Duplanil of Paris, physician to the Count d'Artois, I lie under particular obligations; as this Gentleman has not only enlarged my treatife, but, by his very ingenious and useful notes, has rendered it fo popular on the Continent, as to occasion its being translated into all the languages of modern Europe.

I have only to add, that the book has not more exceeded my expectations in its fuccefs than in the effects it has produced. Some of the most pernicious practices, with regard to the treatment of the fick, have already given place to a more rational conduct; and many of the most hurtful prejudices, which feemed to be quite infurmountable, have in a great measure yielded to better information. Of this a stronger instance cannot be given than in the inoculation of the simil-pox. Few mothers, some years ago, would submit to have their children inoculated even by the hand of a Physician; yet nothing is more certain, than that of late many of them have performed this

13

ope-

ZZ

operation with their own hands; and as their fuccefs has been equal to that of the most dignified inoculators, there is little reason to doubt that the practice will become general. Whenever this shall be the case, more lives will be faved by inoculation alone, than are at present by all the endeavours of the Faculty.

INTRO-

INTRODUCTION.

THE improvements in Medicine, fince the revival of learning, have by no means kept pace with those of the other arts. The reason is obvious. Medicine has been studied by few, except those who intended to live by it as a business. Such, either from a mistaken zeal for the honour of Medicine, or to raise their own importance, have endeavoured to difguise and conceal the art. Medical authors have generally written in a foreign language; and those who are unequal to this task, have even valued themselves upon couching, at least, their prescriptions, in terms and characters unintelligible to the rest of mankind.

The contentions of the clergy, which happened foon after the reftoration of learning, engaged the attention of mankind, and paved the way for that freedom of thought and inquiry, which has fince prevailed in moft parts of Europe with regard to religious matters. Every man took a fide in those bloody disputes; and every gentleman, that he might distinguish himself on one fide or other, was instructed in Divinity. This taught people to think and reason for themselves in matters of religion, and at last totally destroyed that complete and absolute dominion which the clergy had obtained over the minds of men.

The fludy of law has likewife, in most civilized nations, been justly deemed a neceffary part of the education of a gentleman. Every gentleman ought certainly to know at least the laws of his own country: and, if he were also acquainted with those of others, it might be more than barely an ornament to him.

The different branches of Philosophy have also of late been very universally studied by all who pretended to a liberal education. The advantages of this are manifest.

It

INTRODUCTION.

xxii

It frees the mind from prejudice and fuperflition; fits it for the inveftigation of truth; induces habits of reafoning and judging properly; opens an inexhauftible fource of entertainment; paves the way to the improvement of arts and agriculture; and qualifies men for acting with propriety in the most important flations of life.

Natural Hiftory has likewife become an object of general attention; and it well deferves to be fo. It leads to difcoveries of the greateft importance. Indeed, agriculture, the most useful of all arts, is only a branch of Natural Hiftory, and can never arrive at a high degree of improvement where the study of that science is neglected.

Medicine, however, has not, as far as I know, in any country, been reckoned a neceffary part of the education of a gentleman. But furely no fufficient reafon can be affigned for this omiffion. No fcience lays open a more extensive field of ufeful knowledge, or affords more ample entertainment to an inquifitive mind. Anatomy, Botany, Chymistry, and the *Materia Medica*, are all branches of Natural History, and are fraught with fuch amufement and utility, that the man who entirely neglects them has but a forry claim either to tafte or learning. If a gentleman has a turn for observation, fays an excellent and fensible writer *, furely the natural history of his own species is a more interesting subject, and prefents a more ample field for the exertion of genius, than the natural history of spiders and cockle-stills.

We do not mean that every man fhould become a phyfician. This would be an attempt as ridiculous as it is impoffible. All we plead for is, that men of fenfe and learning fhould be fo far acquainted with the general principles of Medicine as to be in a condition to derive from it fome of those advantages with which it is fraught; and at the fame time to guard themselves against the destructive influence of Ignorance, Superstition, and Quackery.

As matters ftand at prefent, it is eafier to cheat a man out of his life than of a shilling, and almost impossible

· Observations on the Duties and Offices of a Physician.

either to detect or punish the offender. Notwithstanding this, people still shut their eyes, and take every thing upon truss that is administered by any Pretender to Medicine, without daring to ask him a reason for any part of his conduct. Implicit faith, every where else the object of ridicule, is still facred bere. Many of the faculty are no doubt worthy of all the confidence that can be reposed in them; but as this can never be the character of every individual in any profession, it would certainly be for the fafety, as well as the honour, of mankind, to have some check upon the conduct of those to whom they entrust fo valuable a treasure as health.

The veil of myftery, which ftill hangs over Medicine, renders it not only a conjectural, but even a fufpicious art. This has been long ago removed from the other fciences, which induces many to believe that Medicine is a mere trick, and that it will not bear a fair and candid examination. Medicine, however, needs only to be better known, in order to fecure the general effeem of mankind. Its precepts are fuch as every wife man would choofe to obferve, and it forbids nothing but what is incompatible with true happinefs.

Difguifing Medicine not only retards its improvement as a fcience, but exposes the profession to ridicule, and is injurious to the true interests of fociety. An art, founded on obfervation, can never arrive at any high degree of improvement, while it is confined to a few who make a trade of it. The united observations of all the ingenious and fenfible part of mankind, would do more in a few years towards the improvement of Medicine, than those of the Faculty alone in a great many. Any man can tell when a medicine gives him eafe as well as a phyfician; and if he only knows the name and dofe of the medicine, and the name of the difeafe, it is fufficient to perpetuate the fact. Yet the man who adds one fingle fact to the flock of Medical observations, does more real tervice to the art, than he who writes a volume in fupport of fome favourite hypothefis.

Very few of the valuable difcoveries in Medicine have been made by phyficians. They have in general either been the effect of chance or of neceffity, and have been

ufually

ufually oppofed by the Faculty, till every one elfe was convinced of their importance. An implicit faith in the opinions of teachers, an attachment to fystems and established forms, and the dread of reflections, will always operate upon those who follow Medicine as a trade. Few improvements are to be expected from a man who might ruin his character and family by even the smallest deviation from an established rule.

If men of letters, fays the author of the performance quoted above, were to claim their right of inquiry into a matter that fo nearly concerns them, the good effects of Medicine would foon appear. Such men would have no feparate intereft from that of the art. They would detect and expofe affuming Ignorance under the mafk of Gravity and Importance, and would be the judges and patrons of modelt merit. Not having their underftandings perverted in their youth by falle theories, unawed by authority, and unbiaffed by intereft, they would canvafs with freedom the most univerfally received principles in Medicine, and expose the uncertainty of many of those doctrines, of which a physician dares not fo much as feem to doubt.

No argument, continues he, can be brought againft laying open Medicine, which does not apply with equal if not greater force, to religion; yet experience has fhewn, that fince the laity have afferted their right of inquiry into these fubjects, Theology, confidered as a fcience, has been improved, the interests of real religion have been promoted, and the clergy have become a more learned, a more useful, and a more respectable body of men, than they ever were in the days of their greatest power and splendour.

Had other medical writers been as honeft as this gentleman, the art had been upon a very different footing at this day. Moft of them extol the merit of thofe men who brought Philofophy out of the fchools, and fubjected it to the rules of common fenfe. But they never confider that Medicine, at prefent, is in nearly the fame fituation that Philofophy was at that time, and that it might be as much improved by being treated in the fame manner. Indeed, no fcience can either be rendered

xxiv

dered rational or useful, without being fubmitted to the common fense and reason of mankind. These alone stamp a value upon science; and what will not bear the test of these ought to be rejected.

I know it will be faid, that diffusing medical knowledge among the people might induce them to tamper with Medicine, and to truft to their own skill, instead of calling a physician. The reverse of this, however, is true. Perfons who have most knowledge in these matters, are commonly most ready both to ask and follow advice, when it is neceffary. The ignorant are always most apt to tamper with Medicine, and have the least confidence in phyficians. Inftances of this are daily to be met with among the ignorant peafants, who, while they abfolutely refuse to take a medicine which has been preferibed by a phyfician, will fwallow with greedinefs any thing that is recommended to them by their credulous neighbours. Where men will act even without knowledge, it is certainly more rational to afford them all the light we can, than to leave them entirely in the dark.

It may alfo be alleged, that laying Medicine more open to mankind, would leffen their faith in it. This would indeed be the cafe with regard to fome; but it would have a quite contrary effect upon others. I know many people who have the utmost dread and horror of every thing preferibed by a physician, but who will neverthelefs very readily take a medicine which they know, and whofe qualities they are in fome measure acquainted with. Hence it is evident, that the dread arifes from the doctor, not from the drug. Nothing ever can or will infpire mankind with an abfolute confidence in phyficians, but an open, frank, and undifguifed behaviour. While the least fhadow of mystery remains in the conduct of the Faculty, doubts, jealousies, and fuspicions, will arife in the minds of men.

No doubt cafes will fometimes occur, where a prudent phyfician may find it expedient to difguife a Medicine. The whims and humours of men must be regarded by those who mean to do them fervice; but this can

never

INTRODUCTION.

never affect the general argument in favour of candour and opennels. A man might as well allege, becaufe there are knaves and fools in the world, that he ought to take every one he meets for fuch, and to treat him accordingly. A fenfible phyfician will always know where difguife is neceffary; but it ought never to appear on the face of his general conduct.

The appearance of mystery in the conduct of physicians not only renders their art fuspicious, but lays the foundations of Quackery, which is the difgrace of Medicine. No two characters can be more different than that of the honest physician and the quack; yet they have generally been very much confounded. The line between them is not fufficiently apparent; at least is too fine for the general eye. Few perfons are able to diftinguish fufficiently between the conduct of that man who administers a secret Medicine, and him who writes a prefeription in mystical characters and an unknown tongue. Thus the conduct of the honest physician, which needs no difguise, gives a fanction to that of the villain, whose fole confequence depends upon fecrecy.

No laws will ever be able to prevent quackery, while people believe that the quack is as honeft a man, and as well qualified as the phyfician. A very fmall degree of medical knowledge, however, would be fufficient to break this fpell; and nothing elfe can effectually undeceive them. It is the ignorance and credulity of the multitude, with regard to Medicine, which renders them fuch an eafy prey to every one who has the hardinefs to attack them on this quarter. Nor can the evil be remedied by any other means but by making them wifer.

The moft effectual way to deftroy quackery in any art or fcience, is to diffufe the knowledge of it among mankind. Did phyficians write their prefcriptions in the common language of the country, and explain their intentions to the patient, as far as he could underftand them, it would enable him to know when the Medicine had the defired effect; would infpire him with abfolute confidence in the phyfician; and would make him dread and

xxvi

and deteft every man who pretended to cram a fecret Medicine down his throat.

Men in the different flates of fociety, have very different views of the fame object. Some time ago it was the practice of this country for every perfon to fay his prayers in Latin, whether he knew any thing of that language or not. This conduct, though facred in the eyes of our anceftors, appears ridiculous enough to us; and doubtless fome parts of ours will feem as strange to posterity. Among these we may reckon the present mode of medical prefcription, which, we venture to affirm, will fome time hence appear to have been completely ridiculous, and a very high burlefque upon the common fenfe of mankind.

But this practice is not only ridiculous, it is likewife dangerous. However capable phyficians may be of writing Latin, I am certain apothecaries are not always in a condition to read it, and that dangerous miftakes, in confequence of this, often happen. But suppose the apothecary ever fo able to read the phyfician's prefcription, he is generally otherwife employed, and the bufinefs of making up prefcriptions is left entirely to the apprentice. By this means the greatest man in the kingdom, even when he employs a first-rate physician, in reality trufts his life in the hands of an idle boy, who has not only the chance of being very ignorant, but likewife giddy and carelefs. Miftakes will fometimes happen in fpite of the greatest care; but, where human lives are concerned, all poffible methods ought certainly to be taken to prevent them. For this realon, the preferiptions of phyficians, inflead of being couched in myflical characters and a dead language, ought, in my humble opinion, to be conceived in the most plain and obvious terms imaginable.

Diffusing medical knowledge among the people would not only tend to improve the art and to banifh quackery, but likewife to render Medicine more univerfally ufeful, by extending its benefits to fociety. However long Medicine may have been known as a fcience, we will venture to fay, that many of its most important purpoles to fociety have either been overlooked, or very little

INTRODUCTION.

little attended to. The cure of difeafes is doubtlefs a matter of great importance; but the prefervation of health is of ftill greater. This is the concern of every man, and furely what relates to it ought to be rendered as plain and obvious to all as poffible. It is not to be fuppofed that men can be fufficiently upon their guard againft difeafes, who are totally ignorant of their caufes. Neither can the Legiflature, in whofe power it is to do much more for preferving the public health than can ever be done by the Faculty, exert that power with propriety, and to the greateft advantage, without fome degree of medical knowledge.

Men of every occupation and condition in life might avail themfelves of a degree of medical knowledge; as it would teach them to avoid the dangers peculiar to their refpective flations; which is always eafier than to remove their effects. Medical knowledge, instead of being a check upon the enjoyments of life, only teaches men how to make the most of them. It has indeed been faid, that to live medically, is to live miferably : but it might with equal propriety be faid that to live rationally is to live miferably. If phyficians obtrude their own ridiculous whims upon mankind, or lay down rules inconfiftent with reafon or common fense, no doubt they will be defpifed. But this is not the fault of Medicine. It propofes no rules that I know, but fuch as are perfectly confiftent with the true enjoyment of life, and every way conducive to the real happiness of mankind.

We are forry indeed to obferve, that Medicine has hitherto hardly been confidered as a popular fcience, but as a branch of knowledge folely confined to a particular fet of men, while all the reft have been taught not only to neglect, but even to dread and defpife it. It will however appear, upon a more ftrict examination, that no fcience better deferves their attention, or is more capable of being rendered generally ufeful.

People are told, that if they dip the leaft into medical knowledge, it will render them fanciful, and make them believe they have every difeafe of which they read. This I am fatisfied will feldom be the cafe with fenfible people; and fuppofe it were, they must foon be undeceived.

XXVIII

ceived. A fhort time will fhew them their error, and a little more reading will infallibly correct it. A fingle inftance will fhew the abfurdity of this notion. A fenfible lady, rather than read a medical performance, which would inftruct her in the management of her children, generally leaves them entirely to the care and conduct of the most ignorant, credulous, and superstitious part of the human species.

No part of Medicine is of more general importance than that which relates to the nurfing and management of children. Yet few parents pay a proper attention to it. They leave the fole care of their tender offspring, at the very time when care and attention are most necessary, to hirelings, who are either too negligent to do their duty or too ignorant to know it. We will venture to affirm, that more human lives are lost by the careless and inattention of parents and nurfes, than are faved by the Faculty; and that the joint and well-conducted endeavours, both of private perfons and the public, for the prefervation of infant lives, would be of more advantage to fociety than the whole art of Medicine, upon its prefent footing.

The benefits of Medicine, as a trade, will ever be confined to those who are able to pay for them; and of courfe, the far greater part of mankind will be every where deprived of them. Phyficians, like other people, must live by their employment, and the poor must either want advice altogether, or take up with that which is worfe than none. There are not, however, any where wanting well-difpofed people, of better fenfe, who are willing to fupply the defect of medical advice to the poor, did not their fear of doing ill often suppress their inclination to do good. Such people are often deterred from the most noble and praise-worthy actions, by the foolifh alarms founded in their ears by a fet of men who, to raife their own importance, magnify the difficulties of doing good, find fault with what is truly commendable, and fleer at every attempt to relieve the fick which is not conducted by the precise rules of Medicine. These gentlemen must, however, excuse me for faying, that I have often known fuch well-difpofed perfons do much good;

INTRODUCTION.

XXX

good; and that their practice, which is generally the refult of good fenfe and observation, assisted by a little medical reading, is frequently more rational than that of the ignorant retainer to physic, who despises both reason and observation, that he may go wrong by rule; and who, while he is dosing his patient with Medicines, often neglects other things of far greater importance.

Many things are neceffary for the fick befides Medicine. Nor is the perfon who takes care to procure thefe for them, of lefs importance than a phyfician. The poor oftener perifh in difeafes for want of proper nurfing than of Medicine. They are frequently in want of even the neceffaries of life, and ftill more fo of what is proper for a fick-bed. No one can imagine, who has not been a witnefs of thefe fituations, how much good a welldifpofed perfon may do, by only taking care to have fuch wants fupplied. There certainly cannot be a more neceffary, a more noble, or a more godlike action, than to adminifter to the wants of our fellow-creatures in diftrefs. While virtue or religion are known among mankind, this conduct will be approved ; and while Heaven is juft, it muft be rewarded !

Perfons who do not choose to administer Medicine to the fick, may neverthelefs direct their regimen. An eminent medical author has faid, That by diet alone all the intentions of Medicine may be answered *. No doubt a great many of them may; but there are other things befide diet, which ought by no means to be neglected. Many hurtful and deftructive prejudices, with regard to the treatment of the fick, ftill prevail among the people, which perfons of better fense and learning alone can eradicate. To guard the poor against the influence of these prejudices, and to instil into their minds fome just ideas of the importance of proper food, fresh air, cleanlinefs, and other pieces of regimen neceffary in difeafes, would be a work of great merit, and productive of many happy confequences. A proper regimen, in most difeases, is at least equal to Medicine, and in many of them it is greatly fuperior.

Arbuthnot.

INTRODUCTION.

To affift the well-meant endeavours of the humane and benevolent in relieving diftres; to eradicate dangerous and hurtful prejudices; to guard the ignorant and credulous against the frauds and impositions of quacks and impoftors; and to fhew men what is in their own power, both with regard to the prevention and cure of difeafes, are certainly objects worthy of the phyfician's attention. Thefe were the leading views in compofing and publishing the following sheets. They were fuggested by an attention to the conduct of mankind, with regard to Medicine, in the courfe of a pretty long practice in different parts of this island, during which the author has often had occasion to wish that his patients, or those about them, had been poffeffed of fome fuch plain directory for regulating their conduct. How far he has fucceeded in his endeavours to fupply this deficiency, must be left to others to determine : but if they be found to contribute in any measure towards alleviating the calamities of mankind, he will think his labour very well bestowed.

The second second second

former edition of the

xxxi

HOITOUR ORTSH

To affift the weil nears endeavours of the human and the evolution in relieving diffrels; to endoare dancerous and hurtfel prejudices; '10 guard the ignorant and credulous against the diauds and inno tions of quacks and impollors ; and to (b a men what is in their own power, both with regard the prevention and cure of director, are certainly oblects worthy of the phylician's attendor, of their ware the leading views in compound and publishing the tollawing theets. They were inggalled by an attention to the conduct of mankind, with regard to Medicine, in the courie of a pretty long practice in different parts of this illand, during which the author has often had occation to with that his patients; or thole about them, had been polleffed of fome fich plain directory for regulating their comindi. How far he has fucceeded in his endervours to lupply this defidiency, mult be left to others to determine : but if they be found to contribute in any measure towards alleviating the calamities of mankind, he will think his labour yery

and the state of the second state of the

12-2

[xxxiii]

A LIST of the MEDICINES.

MANY who perufe the DOMESTIC MEDICINE have expressed a wish that the catalogue of Medicines contained in that book should be more extensive, and likewise that the dose of each article should be ascertained, as they are often at a loss to know how to administer even those Medicines, the names of which they meet with in almost every medical author. To obviate this objection, and furniss a greater scope to those who may wish to employ more articles than are contained in the Dispensatory annexed to the above work, the following List of Simples and Compounds, taken from the most improved Dispensatories, is now inferted.

To prevent miltakes, the English name of every Medicine is not only used, but the different articles are arranged according to the order of the English alphabet, and the smallest and largest dose placed opposite to each article. The doses indeed refer to adults, but may be adapted to different ages by attending to the rules laid down in the Introduction to the Appendix, p. 687. Short cautions are occasionally inferted under such articles as require to be used with care.

Though a greater variety of Medicines is contained in this than in any former edition of the Domeftic Medicine, yet the Author would advife those who peruse it, as far as possible, to adhere to simplicity in practice. Diseases are not cured by the multiplicity of Medicines, but by their proper application. A few simples, judiciously administered, and accompanied with a proper regimen, will do more good, than a farrago of Medicines employed at random.

(xxxiv)

A LIST of the MEDICINES commonly used in Practice, with their proper Dofes.

A

A CACIA, the expressed juice, from Acid, the acetous	i feruple to I drachm
Acid, the acctous	I feruple — I drachm
, muriatic	10 drops — 40 drops
There is a state of the state o	15 drops — 40 drops
, vitriolic, diluted	15 drops - 40 drops
Æther, vitriolic	30 drops - 2 drachms
	10 grains — 30 grains
Aloes	5 grains — 30 grains
Alum out of -ost totate s day	6 grains — 20 grains
, ourned	3 grains - 12 grains
A 1	1 drachm — 1 drachm
	5 grains — 30 grains
	$\frac{1}{2}$ oz. — 1 ounce
	$\frac{1}{2}$ drachm — $1\frac{1}{2}$ drachm
A .C .1 C 1	10 grains — I drachm
A	10 grains — 1 drachm
A LON TO LON MANY AND AND A REAL PROPERTY AND	1 fcruple — 1 drachm
and a state of the	4 grain — 2 grains
ACC 11	6 grains - half a dras
	half oz. — 1 ounce
	3 grains — 5 grains
) Brands

B

C

and the second to the second a second state of the second		
Balfam of capivi -	-	20 drops
Canadian —	-	
of Peru		
of Tolu	-	
Bark, Peruvian, powder -	-	2 fcrup.
Bear's foot, powder -	-	10 grains
Benzoin, refin of —	-	4 grains
	-	10 grains
Biftort, powder of the root	-	I fcruple
Bleffed thiftle	-	10 grains
, expreffed juice of	-	2 dra.
Bole, Armenian —	-	10 grains
	-	
Borax — —		10 grains
Broom, ashes of the tops -	-	I fcruple
Burdock, powder of the root	-	10 grains

Calomel

{ 1 gr. to 3 gr. alterative 3 do. to 12 do. purgative Camphor

- 60 drops

— 40 grains — 1 drachm — 1 drachm

2 drachms
20 grains
20 grains
20 grains
1 drachm
1 drachm
2 ounces
2 drachms

MEDICINES USED IN PRACTICE.

Camphor -		from	2	grains to	hal	lf a drachm
Canella alba, powder o		-				2 drachms
Cantharides						4 grains
Cardamoms						20 grains
Caraway feeds				grains		40 grains
Cafcarilla bark				grains		40 grains
Caffia, the pulp						1 ounce
Caftor -			-	grains		1 drachm
Catechu - house	M. ALTING			grains		30 grains
Camomile, in powder			-	grains		1 drachm
Chalk —				grains		2 feruples
Cinnamon -				grains		I drachm
Colocynth -			-	grains		1 drachm
Columbo -				grains		1 drachm
Confection, aromatic	Z			grains	_	
	_			grains	_	
				C	_	1 drachm
Crabs claws, prepared Conferve of rofes				grains		ALC: NOT THE REPORT OF THE REPORT
	anter			drachm		1 ounce
of fquills	-			grains		30 grains
of arum	-			grains		1 drachm
Contrayerva	-			grains		2 feruples
Corjander feed	- 1		15	grains		1 drachm
Cowhage, the spiculæ o	t one pod	mix-				

ed with honey or molaffes.

D

Dandelyon, expressed juice of ______ Decoction of hartshorn, half a pint repeated as often as necessary. _______ of broom, 1 oz. to a pint of water, to be taken by tea-cupfuls. _______ of Peruvian bark _______ _____ of the inner bark of the elm _______ of the inner bark of the elm _______ of farfaparilla _______ _____ of guaiacum, 3 drachms to a pint of water. A pint daily.

I oun	ce	- 3	ou	nces
				and the second se

I	ounce	-	4 ounces
4	ounces	-	10 oz. daily
4	ounces	-	16 oz. daily

Electuary of caffia -	- I drachm - I gunce	
of fcammony	— 20 grains — 1 drachm	
lenitive, or of fenna	- 30 grains - 6 drachms	
Elixir of vitriol —	- 15 drops - 50 drops	
Elecampane, powder of the root	20 grains - 1 drachm	
Extract of broom tops	$-\frac{1}{2}$ drachm $-$ I drachm	
Peruvian bark	- 10 grains - 1/2 drachm	
cafcarilla -	-10 grains $-\frac{1}{2}$ drachm	
camomile -	— 20 grains — 1 drachm	
colocynth - 1	- 5 grains - 25 grains	
	Extrad	

E

XXXX

xxxvi MEDICINES used in PRACTICE.

Extract of gentian -	from 10 grains to 1 drachm
liquorice	-1 drachm $-\frac{1}{2}$ ounce
logwood	- 10 grains - 1 drachm
black hellebore	- 3 grains - 10 grains
	- 10 grains - 20 grains
white poppies	- 10 grains - 20 grains
rue	- 1 grain - 5 grains - 10 grains - 20 grains
favin	- 10 grains - 30 grains
fenna —	- 10 grains - 30 grains

F

Fern, powder of the root	I	drachm		1 ounce
Fennel leed	20	grains	-	I drachm
Fox glove, powder of the leaves - or a drachm infufed in a pint of ?	12	grain	-	3 grains
boiling water, of which a dofe is	1	ounce		Call of
Should be administered with caution.				

G

Galbanum 10 grains - 30	araind
Galls 10 grains - 20	
Garlic, cloves of - No. 1 No.	6
Gentian IO grains - 40 g	
	drachm
Ginger 5 grains - 20 g	
Ginteng 20 grains - 30 g	
Guaiacum, gum-refin — Io grains — 30 g	
Cum arabic 15 grains - 10	
gambouge 2 grains - 12 g	

Η

Hartshorn, prepared -	-	20 grains	- I drachm
	-	10 drops	
ginous vehicle —	-}	5 drops	- 25 drops
, falt of	-	2 grains	- 12 grains
Hellebore, white -	-	I grain	- 5 grains
	-	5 grains	- 10 grains
Hemlock fhould always be begun	in in	1.01	2 The Day of the
very fmall dofes, of one grain	or		
lefs, and gradually increafed as	the		
conftitution will bear.			
Hiera picra —		10 grains	- 20 grains
Honey of fquills -	-	10 grains	- 40 grains
of roles	-	1 drachm	- 2 drachms
Hoffman's anodyne liquor	-	20 drops	- 60 drops
Hiddenstin + Carming On		- Shine	and an and a state of the state
- 1 - 2 man - 2 - 1			" - mainten color
Islan nowdan	_	to amaine	in quaine
	1.	and the second	- 40 grains
Infusion of gentian, compound	-	I ounce	- 3 ounces
			Infusion

MEDICINES USED IN PRACTICE. XXXVII

Infusion of roles	from	2	ounces	to	8	ounces
of fenna		12	ounce	-	2	ounces
Ipecacuanha —	-	IO	grains	-	30	grains
Iron, ruft of ammoniated	-		grains	-	20	grains ·
			grains			grains
falt of			grains			grains
Juniper, powder of the berries	-		grain			grains
Jumper, powder of the bernes	-	20	grains	-	I	drachm
read and the second second	K				Tim	
Kino, gum	-	10	grains	-	30	grains
Kermes, juice of —		I	drachm	-	3	drachms

L

T · · ·		
Lichen, afh-coloured, ground -		The state is a state of the
Icelandic, a ftrong decoction of	3 grains	- 40 grains
Icelandic, a itrong decoction of	T OURGO	
Lime-water	I ounce	- 4 ounces
Title water	4 ounces	- 8 ounces
Lixivium of tartar		
Tic 1 ica	IS drops	- 40 drops
Linfeed, an infusion of I ounce to a	3	To mope
quart of motor i ounce to a		
quart of water; may be used at		
pleafure.		
Prendic.		

M

M	a manufact o the birth of this work of
Madder powder	
Mace _	- ¹ / ₂ drachm - 1 drachm
	- 10 grains - 20 grains
Magnefia	- drachen a drachene
calcined _	$-\frac{1}{2}$ drachm -2 drachms
Manna	
	$-\frac{1}{2}$ ounce -2 ounces
Maftich, gum	
Mercury, crude	
calcined	- ¹ / ₂ ounce - 4 ounces
with chalk	- ¹ / ₂ grain - 2 grains
corrofive fublimate	- 10 grains - 30 grains
corrouve iublimate	· · · · · · · · · · · · · · · · · · ·
cinnabar of	
vellow emetic as flowers	10 grains — 30 grains
Mezereon, decost to a pint of	I grain — 3 grains
Mezereon, decoct. to a pint of water Millipedes	- 2 drachms
Mufk	
Muftard feed	- 5 grains - 40 grains
Myrrh, gum	- I drachm — I ounce
-	10 grains — I drachm
N	
Nitre, purified	the second se
Nutmeg	IO graine an and
a rucince and and	10 grains - 30 grains
	$-$ 6 grains $ \frac{1}{2}$ drachm
Oil of Almonds O	
on of Almonds	Terration in a second state
Linfeed	2 ounce — I ounce
Caftor -	
Olibanum	2 drachms - I ounce
. The set is a set of the set of	5 graine
C	5 grains - 30 grains
	Onion,
	and the second se

XXXVIII MEDICINES USED IN PRACTICE.

AAAVIII MILDICIIVIDO ODA		
Onion expressed inice of a7	· · ····	1019 10 06211-1 .
Onion, expressed juice of, a	from $\frac{1}{2}$ ounce	to 2 ounces
powerful diuretic,	- I crain	- 2 grains
Opium — —	2 grain	- 30 grains
Opoponax -	- 10 grans	n — 1 ounce
Oxymel of colchicum -		
of fquills	- 2 drachi	n — 2 drachms
and the second	and and a fig	
	P	
Petroleum	- 10 drops	- 30 drops,
Pills, aloctic	- 10 grains	- 30 grains
	- 10 grains	
- of the gums -	- 10 grains	- 20 grains
	- 20 grains	- 1 drachm
Pomegranate, powder of	- 3 grains	- 6 grains
Powder, antimonial	3 5141115	Nissalla Attriov
May be taken according to the		
directions for James's powder		Lincowyler
with which it nearly coincides.	Y M mening	an arains
- of Contrayerva, compound	- 15 grains	- 30 grams
of Chalk, compound	- 20 grains	- 40 grains
with opium	- 10 grains	- 40 grains
of Ipecacuanha, com-?	- ro grains	- 30 grains
pound, or Dover's powder 5		
		Section of the total
O service and a service of the servi		
NUMBER OF THE PROPERTY OF THE	- r grains	- 30 grains
Quaffia — Conto		- 30 8.
Two drachms to a pint of water		
for a decoction.		
Quince feeds, mucilage of, at plea-		
fure, to obtund acrimony.		
R	-	in mains
Rhubarb, powder	- 10 grains	- 40 grains
Refin, yellow -	- 3 grains	- 20 grains
Rue powder -	- 20 grains	- 40 grains
Kue powder		
additions -, aviet or S		
anterio turi ana studio o oran	- 20 grains	- I drachm
St. John's wort	- 5 grains	- zo grains
Saffron — —	- 5 grains	- 30 grains
Sagapenum -	- 10 grains	- 30 grains
Sal ammoniac —	- 10 grains - 2 dra.	- 1 ounce
Salt, Epfom	2 dia.	- 2 ounces
Glauber	- 4 dra.	- ¹ / ₂ ounce
Polychreft -	- 20 grains	- 30 grains
of Tartar	- 10 grains	- 40 grains
Sarfaparilla, powder of -	- 20 grains	- 10 grains
Scammony -	- 5 grains	
Seneka	- 20 grains	- 40 grains
	- 20 grains	- 40 grains
Senna -	- 20 grains	- i ounce
Soap	- 10 drops	- 30 drops Scurvy-
lecs 6		Dem al-

6

MEDICINES USED IN PRACTICE. XXXXX

from	I	ounce	to	4	ounces
-	20	grains	-	40	grains
-			-	8	ounces
-	I	drachm		I	ounce
	-				-
-			-		
	I 2	drachm	-	12	ounce
			-	1	drachm
	20	grains	-	I	drachm
-	20	grains	-	I	drachm
~	I	grain	-	4	grains
-	I	grain	-	3	grains
-					
				-	-
	-				
	I	drachm	-	T	ounce
		$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	 4 ounces 1 drachm 15 drops 15 drops 15 drops 15 drops 15 drops 20 grains 20 grains 20 grains 20 grains 1 grain 1 grain 5 grains 1 drachm 1 drachm 	 20 grains 4 ounces 1 drachm 15 drops 20 grains 20 grains 20 grains 20 grains 20 grains 1 grain 1 grain 5 grains 1 drachm 1 drachm 	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$

T

Tar water. A pint daily.				there and
Tartar, cream of	-	2 drachms	-	I ounce
regenerated	-	20 grains	-	1 drachm
——— foluble —	-	2 drachms		
emetic, alterative	-	1 grain		
as emetic -	-	I grain		
Terra japonica —		20 grains		
Tobacco, an infusion of, I drachm		Drame		to Branny
to a pint of water ; fhould be ad-				
ministered by table - spoonfuls :				
ftrongly diuretic.		Thing to & sty	44° 1	
Tin, powder of -	-	20 graine	-	I drachm
Turmeric — —				I drachm
Turpentine, fpirits of	11-5			
Tincture of aloes	-	10 drops		
		- 1 ounce		
compound	-			2 drachms
of afafœtida	-			2 drachms
of Benzoin, compound		10 drops		
of cantharides		10 drops		40 drops
of cardamoms		I drachm	-	1 ounce
of caftor				I_2^{\perp} drachm
of catechu	-			2 drachms
of Peruvian bark	-	I drachm	-	¹ / ₂ ounce
of iron, muriated		10 drops	-	60 drops
of Columbo	-	1 drachm		3 drachms
of Gentian, compound		ı drachm		3 drachms
of guaiacum volatile	-			3 drachms
	2 2			Tincture
and the second se				

MEDICINES USED IN PRACTICE.

Tincture of black hellebore	from 1 feruple to 1 drachm	
of jalap	- I drachm - I ounce	
of lavender, compound	- 20 drops - 2 drachme	
of myrrh	— 1 fcruple — 1 drachm	
of opium	In In drama	
camphorated,	or	
paregoric elixir	I drachm - 3 drachms	
of rhubarb	$-\frac{1}{2}$ ounce -2 ounces	
of fenna	- 2 drachms - 1 ounce	
` of fnake-root	- 1 drachm - 2 drachms	
of valerian	1 1	
volatile	— I drachm — 3 drachms	
Townson 1 C	1 drachim 2 drachms	
1 ormentil, powder of	- 10 grains - 1 drachm	

V

Valerian, powder of -	-	20 grains	-	2 drachms
Vinegar, diftilled	-	2 drachn	ns —	I ounce
	-	10 drops	-	50 drops
as emetic	-	¹ / ₂ ounce	-	I ounce
Verdigris, violent emetic -		I grain	-	2 grains
Vitriol, white, as a tonic		2 grains	-	5 grains
ting emetic as a quickly opera-	}	20 grains	-	1 drachm
blue, emetic -		I grain		3 grains
Uva urfi, in powder -	-	20 grains		1 drachm

W

Water crefs, expressed juice of Water creis, exprended und may ge- } I ounce 2 ounces 1 ounce nerally be given 3 or 4 oz. $\frac{1}{2}$ ounce I grain Wormwood, expressed juice 2 ounces White lead 3 grains ----1 ounce Wine, aloetic I ounce ----- antimonial 20 drops - 2 drachms ---- Ipecacuanha 1 drachm — $1\frac{1}{2}$ ounce ---- Rhubarb ¹₂ ounce 2 ounces

CON-

xl

PART I. Of Love - Page 113 Of the general Caufes of -Religious Melancholy 115 Difeases. CHAP. XI. Of the Common Evacuations CHAP. I. F Children Page I 116 Difeafed Parents 6 -Stool ib. -Clothing of Children -Urine 9 118 -Perspiration --Food of ditto 14 120 -Being affected by Changes in -Exercife of ditto -20 the Atmosphere — -Bad Effects of unwholefome ib. -Wet Clothes Air upon ditto -28 121 -Wet Feet -Nurfes 30 122 -- Night Air CHAP. II. ib. Of the Laborious, &c. -Damp Beds ib. 35 -Damp Houfes --the Sedentary 124 44 -Sudden Transitions from Heat -the Studious 50 to Cold CHAP. III. 125 Of Aliment ____ PART II. 57 CHAP. IV. Of Air Of Difeases. 69 CHAP. V. CHAP. XII. Of Exercife -Of the Knowledge and Cure of 75 Difeafes CHAP. VI. 129 Of Sleep 81 CHAP. XIII. -Clothing -Fevers in general ____ 84 135 CHAP. VII. CHAP. XIV. Of Intemperance -Of intermitting Fevers or Agues 89 CHAP. VIII. Of Cleanlinefs 141 CHAP. XV. 95 Of an acute continual Fever 151 CHAP. IX. Of Infection -100 CHAP. XVI. Of the Pleurify CHAP. X. Of the Paffions 157 -Baftard ditto 107 163 -Anger ---- Paraphrenitis 108 ib. -Fear ' CHAP. XVII. ib. Grief Inflammation of the Lungs 164 IIZ

[xli]

Of Confumptions Page	166
CHAP. XIX.	-
Of the Slow or Nervous	
ver — —	180
CHAP. XX.	ttad
Malignant, Putrid, or Spo Fever	186
CHAP, XXI,	100
Miliary Fever -	195
CHAP. XXII.	- ,,,
Remitting Fever	201
CHAP. XXIII.	
The Small-pox -	204
Inoculation -	215
CHAP. XXIV.	10
The Meafles -	229
	233
Bilious Fever -	234
CHAP. XXV.	
St. Anthony's Fire	236
CHAP. XXVI.	
Inflammation of the Brain	241
CHAP. XXVII.	-
Inflammation of the Eyes	245
CHAP. XXVIII.	
The Quinfey -	251
Malignant ditto	257
CHAP. XXIX.	262
Colds and Coughs	267
A common Cough Hooping-Cough -	271
CHAP, XXX.	-1-
	275
of the Inteftines	
Of the Colic -	281
Inflammation of the Kidneys	286
of the Bladder	288
of the Liver	209
CHAP. XXXI.	and
Of the Cholera Morbus, other exceffive Difcha	rges
from the Stomach and Bo	wels
	293
-a Diarrhœa, or Loofeness	296
-Vomiting -	298
CHAP. XXXII.	
Diforders of the Kidneys	and
Bladder -	301

Of the Diabetes, or exce	five
Discharge of Urine Page	302
-Supprellion of ditto	
-the Gravel and Stone	306
CHAP. XXXIII.	
Involuntary Difcharges	of
Blood -	311
Bleeding at the Nofe	313
Bleeding and Blind Piles Spitting of Blood -	316
Vomiting of ditto -	320
Bloody Urine -	323 324
Flux -	325
Cœliac Paffion -	331
CHAP. XXXIV.	220
Of the Head-ach -	332
Tooth-ach -	335
Ear-ach	339
Pain of the Stomach, &c.	340
CHAP. XXXV.	0
Of Worms	342
CHAP. XXXVI.	343
Of the Jaundice -	348
CHAP. XXXVII.	540
Of the Dropfy -	252
	352
CHAP. XXXVIII. Of the Gout	
	357
	364
CHAP. XXXIX. Of the Scurvy -	-60
	368
	373
for the second second	379
Of the Afthma -	382
CHAP. XLI.	304
Of the apoplexy -	386
	244
CHAP. XLII. Of Cofliveneis	100
- Want of Appetite	399
- Heartburn .	39 ² 393
CHAP. XLIII.	393
Of Nervous Difeafes	395
- Melancholy -	400
- the Palfy - automation	403
	ling
	405
- the Hiccup -	408
- Cramp of the Stomach	410
- the Night-Mare -	411

alii

· · · · · · · · · · · · · · · · · · ·			
Of Swoonings Page	412	Of Galling and Excoriation P.	507
- Flatulencies, or Wind	414	- Stoppage of the Nofe	508
- Low Spirits	417	Vomiting -	ib.
- Hysteric Affections -	418	- a Loofenels -	509
- Hypochondriac Affections	421	- Eruptions -	510
		- the Ĉroup -	513
CHAP. XLIV.		- Teething -	515
Diforders of the Senfes	426	- the Rickets .	517
of the Eye	ib.	- Convulfions	520
A Gutta Serena -	428	- Water in the Head	522
A Cataract -	ib.	CHAP. L.	-
Short-fightednefs -	ib.	Of Surgery -	523
Seeing only at too great a	Dif-		
tance	429	- Bleeding - - Inflammations and Abfo	524 effer
Squinting -	ib.	- Innaminations and Ilow	
Spots or Specks -	ib.	Bielding Property and	527
Bloodshot Eye	ib.	- Wounds -	529
Watery or Weeping Eye	ib.	- Burns -	532
Of the Ear -	430	- Drunes -	534
Tafte and Smell	432	— Ulcers -	535
	and the second second	CHAP. LI.	
Touch -	434	Of Diflocations -	538
CHAP. XLV.	106	Diflocation of the Jaw	539
Of Schirrus and Cancer	436	of the Neck	ib.
CHAP. XLVI.		of the Ribs	540
Of Poifons -	442	of the Shoulder	541
- Mineral Poisons -	443	of the Elbow	ib.
- Vegetable ditto -	445	of the Wrift and	Fin-
Bites of Poifonous Animals	446	gers	542
- of the Mad Dog	ib.	of the Thigh	ib.
- of the Viper	453	of the Knees	ib.
of Poifonous Infects			ib.
of the Rattle-fnake	454		
	455	of the Toes	ib.
CHAP. XLVII.	1-6	CHAP LII.	
Of the Venereal Difeafe	456		
Virulent Gonorrhœa		Strains -	546
- Gleets -	465	- Ruptures -	547
- Swelled Teffiele -	468	CHAP. LIII.	
- Buboes - - Chancres -	469	Of Cafualties -	549
- Chancres -	470	- Substances ftopt between	the
- a Confirmed Lues or Pox	473	Mouth and Stomach	550
General Obfervations	477	- Drowned Perfons	555
CHAP. XLVIII.		- Noxious Vapours	560
Difeafes of Women	483	Effects of extreme Cold	562
Of the Menstrual Discharge	484	Heat	563
- Pregnancy -	401	CHAP. LIV.	203
- Child-birth -	402		-afan
- Barrennefs -	501	Of Fainting Fits, and other (Ar
	5	which require immediate	
Difeafes of Children	-	fiftance -	564
Of the Macaning	-	- Intoxication -	568
Of the Meconium	504		570
- the Apthz or Thrush			Con-
- Acidities -	500	vulfion Fits -	573

xim

-

OF Cold Bathing, and drinking	Diet of the Common People Page 619
the Mineral Waters Page 576	General Obfervations on Ali-
Of Cold Bathing, with Remarks	ment 621
on the Cafes in which the	Of Bread - 622
Warm Bath is more advi-	Of Boiled Grain - 625
fable - 577	Of Butter - 622
Of Mineral Waters 589	Of Fruits and Roots 625
CHAP. LVI.	Of Broths and Soups 642
Obfervations concerning the	Conclusion 648

CONTENTS of the APPENDIX.

+ Des		TCC		1.02
INTRODUCTION Page	e 053	Infutions		677
Lift of Simples	658	Juleps		679
Medical Preparations.		Mixtures		681
Balfams	660	Ointments,	Liniments,	and
Bolufes -	661	Cerates	-	682
Cataplafms and Sinapifms	66z	Pills	Contraction of the	685
Clyfters	663	Plafters		689
Collyria, or Eye-waters	665	Poultices		691
Confections -	666	Powders		692
Conferves and Preferves	667	Syrups		695
Decoctions -	668		Elixirs, &c.	696
Draughts -	670	Vinegars	1.000	700
Electuaries -	671	Waters by	Infufion, &c.	702
Emulfions -	673		tilled Waters	704
Extracts -	674	Spirituous .	Distilled Waters	705
Fomentations -	675	Wheys	and a group of the second	ib.
Gargles -	676	Wines	- 0110	706

A GLOSSARY INDEX -

709

xliv

PART I.

20 - 1.

OFTHE

GENERAL CAUSES OF DISEASES.

CHAP. I.

OF CHILDREN.

THE better to trace difeafes from their original caufes, we shall take a view of the common treatment of mankind in the state of infancy. In this period of our lives, the foundations of a good or bad constitution are generally laid; it is therefore of importance, that parents be well acquainted with the various causes which may injure the health of their offspring.

It appears from the annual registers of the dead, that almost one half of the children born in Great Britain die under twelve years of age. To many, indeed, this may appear a natural evil; but on due examination it will be found to be one of our own creating. Were the death of infants a natural evil, other animals would be as liable to die young as man; but this we find is by no means the cafe.

It may feem strange that man, notwithstanding his fuperior reason, should fall so far short of other animals in the management of his young: But our surprise will soon cease, if we consider that brutes, guided by instinct, never err in this respect; while man, trusting folely to art, is feldom right. Were a catalogue of those infants who perish annually by art alone exhibited to public view, it would astonish most people.

If

2

If parents are above taking care of their children, others must be employed for that purpose: these will always endeavour to recommend themselves by the appearance of extraordinary skill and address. By this means such a number of unnecessary and destructive articles have been introduced into the diet, clothing, &c. of infants, that it is no wonder so many of them perish.

Nothing can be more prepofterous than a mother who thinks it below her to take care of her own child, or who is fo ignorant as not to know what is proper to be done for it. If we fearch Nature throughout, we cannot find a parallel to this. Every other animal is the nurfe of its own offspring, and they thrive accordingly. Were the brutes to bring up their young by proxy, they would fhare the fame fate with those of the human species.

We mean not, however, to impofe it as a tafk upon every mother to fuckle her own child. This, whatever fpeculative writers may allege, is in fome cafes impracticable, and would inevitably prove deftructive both to the mother and child. Women of delicate conffitutions, fubject to hyfteric fits, or other nervous affections, make very bad nurfes *: and thefe complaints are now fo common, that it is rare to find a woman of fashion free from them; fuch women, therefore, fuppofing them willing, are often unable to fuckle their own children.

Almost every mother would be in a condition to give fuck, did mankind live agreeably to Nature; but whoever confiders how far many mothers deviate from her dictates, will not be surprised to find some of them unable to perform that necessary office. Mothers who do not eat a sufficient quantity of solid food, nor enjoy the benefit of free air and exercise, can neither have wholesome juices themselves, nor afford proper nourishment to an infant. Hence children who are suckled

• I have known an hyfleric woman kill her child, by being feized with a fit in the night. by delicate women, either die young, or continue weak and fickly all their lives.

When we fay that mothers are not always in a condition to fuckle their own children, we would not be underftood as difcouraging that practice. Every mother who can, ought certainly to perform fo tender and agreeable an office *. But fuppofe it to be out of her power, fhe may, neverthelefs, be of great fervice to her child. The bufinefs of nurfing is by no means confined to giving fuck. To a woman who abounds with milk, this is the eafieft part of it. Numberlefs other offices are neceffary for a child, which the mother ought at leaft to fee done.

A mother who abandons the fruit of her womb, as foon as it is born, to the fole care of an hireling, hardly deferves that name. A child, by being brought up under the mother's eye, not only fecures her affection, but may reap all the advantages of a parent's care, though it be fuckled by another. How can a mother be better employed than in fuperintending the nurfery ? This is at once the most delightful and important office ; yet the most trivial bufiness or infipid amusements are often preferred to it ! A ftrong proof both of the bad tafte and wrong education of modern females.

It is indeed to be regretted, that more care is not beftowed in teaching the proper management of children to those whom Nature has designed for mothers. This, instead of being made the principal, is feldom

* Many advantages would arife to fociety, as well as to individuals, from mothers fuckling their own children. It would prevent the temptation which poor women are laid under of abandoning their children to fuckle those of the rich for the fake of gain; by which means fociety loss many of its most useful members, and mothers become in fome fense the murderers of their own offspring I am fure I speak within the truth when I fay, that not one in twenty of those children live, who are thus abandoned by their mothers. For this reason no mother should be allowed to suckle another's child till her own is either dead, or fit to be weaned. A regulation of this kind would fave many lives among the poorer fort, and could do no hurt to the rich, as most women who make good nurfes are able to fuckle two children in fuccession upon the fame milk.

confi-

confidered as any part of female education. Is it any wonder, when females fo educated come to be mothers, that they fhould be quite ignorant of the duties belonging to that character? However firange it may appear, it is certainly true, that many mothers, and thofe of fashion too, are as ignorant, when they have brought a child into the world, of what is to be done for it, as the infant itself. Indeed, the most ignorant of the fex are generally reckoned most knowing in the business of nursing. Hence, fensible people become the dupes of ignorance and superstition; and the nursing of children, instead of being conducted by reason, is the result of whim and caprice *.

Were the time that is generally fpent by females in the acquifition of trifling accomplifhments, employed in learning how to bring up their children; how to drefs them fo as not to hurt, cramp, or confine their motions; how to feed them with wholefome and nourifhing food; how to exercise their tender bodies, fo as beft to promote their growth and ftrength: were thefe made the objects of female inftruction, mankind would derive the greateft advantages from it. But while the education of females implies little more than what relates to drefs and public fhew, we have nothing to expect from them but ignorance even in the moft important concerns.

Did mothers reflect on their own importance, and lay it to heart, they would embrace every opportunity of informing themfelves of the duties which they owe to their infant offspring. It is their province, not only to form the body, but allo to give the mind its most early bias. They have it very much in their power to make

* Tacitus, the celebrated Roman hiftorian, complains greatly of the degeneracy of the Roman ladies in his time, with regard to the care of their offspring. He fays that, in former times, the greateft women in Rome used to account it their chief glory to keep the house and atter d their children; but that now the young infant was committed to the fole care of some poor Grecian wench, or other menial fervant.—We are afraid, wherever luxury and effeminacy prevail, there will be too much ground for this complaint.

men

men healthy or valetudinary, ufeful in life or the pelts of fociety.

But the mother is not the only perfon concerned in the management of children. The father has an equal interest in their welfare, and ought to affist in every thing that respects either the improvement of the body or mind.

It is pity that the men fhould be fo inattentive to this matter. Their negligence is one reafon why females know fo little of it. Women will ever be defirous to excel in fuch accomplishments as recommend them to the other fex. But men generally keep at fuch a diftance from even the fmallest acquaintance with the affairs of the nurfery, that many would reckon it an affront, were they fuppofed to know any thing of them. Not fo, however, with the kennel or the ftables : a gentleman of the first rank is not ashamed to give directions concerning the management of his dogs or horfes, yet would blufh were he furprifed in performing the fame office for that being who derived its existence from himself, who is the heir of his fortunes, and the future hope of his country.

Nor have phyficians themfelves been fufficiently attentive to the management of children: this has been generally confidered as the fole province of old women, while men of the first character in physic have refused to vifit infants even when fick. Such conduct in the faculty has not only caufed this branch of medicine to be neglected, but has also encouraged the other fex to affume an absolute title to prefcribe for children in the most dangerous difeafes. The confequence is, that a phyfician is feldom called till the good women have exhaulted all their skill; when his attendance can only ferve to divide the blame, and appeafe the difconfolate parents.

Nurfes should do all in their power to prevent difeafes; but when a child is taken ill, fome perfon of skill ought immediately to be confulted. The difeases of children are generally acute, and the least delay is dangerous.

Were phyficians more attentive to the difeases of infants, they would not only be better qualified to treat them

them properly when fick, but likewife to give ufeful directions for their management when well. The difeafes of children are by no means fo difficult to be underftood as many imagine. It is true, children cannot tell their complaints; but the caufes of them may be pretty certainly difcovered by obferving the fymptoms, and putting proper queftions to the nurfes. Befides, the difeafes of infants being lefs complicated, are eafier cured than those of adults *.

It is really aftonifhing, that fo little attention fhould in general be paid to the prefervation of infants. What labour and expence are daily beftowed to prop an old tottering carcafe for a few years, while thoufands of thofe who might be ufeful in life, perifh without being regarded! Mankind are too apt to value things according to their prefent, not their future, ufefulnefs. Though this is of all others the moft erroneous method of effimation; yet upon no other principle is it poffible to account for the general indifference with refpect to the death of infants.

Of Difeased Parents.

One great fource of the difeafes of children is, the UNHEALTHINESS OF PARENTS. It would be as reafonable to expect a rich crop from a barren foil, as that ftrong and healthy children fhould be born of parents whofe conflitutions have been worn out with intemperance or difeafe.

An ingenious writer † obferves, that on the conflitution of mothers depends originally that of their offspring. No one who believes this, will be furprifed, on a view of the female world, to find difeales and

* The common opinion, that the difeafes of infants are hard to difcover and difficult to cure, has deterred many phyficians from paying that attention to them which they deterve. I can, however, from experience declare, that this opinion is without foundation; and that the difeafes of infants are neither fo difficult to difcover, nor fo ill to cure, as those of adults.

+ Rouffeau.

death fo frequent among children. A delicate female, brought up within doors, an utter ftranger to exercife and open air, who lives on tea and other flops, may bring a child into the world, but it will hardly be fit to live. The first blass of difease will nip the tender plant in the bud: or should it struggle through a few years' existence, its feeble frame, shaken with convulsions from every trivial cause, will be unable to perform the common functions of life, and prove a burden to fociety.

If, to the delicacy of mothers, we add the irregular lives of fathers, we shall fee further caufe to believe that children are often hurt by the constitution of their parents. A fickly frame may be originally induced by hardships or intemperance, but chiefly by the latter. It is impossible that a course of vice should not spoil the best constitution : and, did the evil terminate here, it would be a just punishment for the folly of the sufferer; but when once a difease is contracted and rivetted in the habit, it is entailed on possible. What a dreadful inheritance is the gout, the feury, or the king's evil, to transmit to our offspring! how happy had it been for the heir of many a great estate, had he been born a beggar, rather than to inherit his father's fortunes at the expence of inheriting his difeases!

A perfon labouring under any incurable malady ought not to marry. He thereby not only flortens his own life, but transmits misery to others; but when both parties are deeply tainted with the fcrophula, the fcurvy, or the like, the effects must be still worfe. If fuch have any iffue, they must be miserable indeed. Want of attention to these things, in forming connections for life, has rooted out more families than plague, famine, or the fword; and as long as these connections are formed from mercenary views, the evil will be continued *.

* The Lacedemonians condemned their king Archidamus for having married a weak, puny woman; because, said they, instead of propagating a race of heroes, you will fill the throne with a progeny of changelings.

In

In our matrimonial contracts, it is amazing fo little regard is had to the health and form of the object. Our fportfmen know that the generous courfer cannot be bred out of the foundered jade, nor the fagacious fpaniel out of the fnarling cur. This is fettled upon immutable laws. The man who marries a woman of a fickly conftitution, and defcended of unhealthy parents, whatever his views may be, cannot be faid to act a prudent part. A difeafed woman may prove fertile; fhould this be the cafe, the family mult become an infirmary: what profpect of happiness the father of fuch a family has, we shall leave any one to judge *.

Such children as have the misfortune to be born of difeafed parents, will require to be nurfed with greater care than others. This is the only way to make amends for the defects of confliction; and it will often go a great length. A healthy nurfe, wholefome air, and fufficient exercife, will do wonders. But when thefe are neglected, little is to be expected from any other quarter. The defects of confliction cannot be fupplied by medicine.

Those who inherit any family-difease ought to be very circumspect in their manner of living. They should confider well the nature of such difease, and guard against it by a proper regimen. It is certain, that family-difeases have often, by proper care, been kept off for one generation; and there is reason to believe, that, by perfisting in the same course, such difeases might at length be wholly eradicated. This is a subject very little regarded, though of the greatest importance. Family-conflictutions are as capable of improvement as family-estates; and the libertine, who impairs the one,

* The Jews, by their laws, were, in certain cafes, forbid to have any manner of commerce with the difeafed; and indeed to this all wife legiflators ought to have a fpecial regard. In fome countries, difeated perfons have actually been forbid to marry. This is an evil of a complicated kind, a natural deformity, and political mifchief; and therefore requires a public confideration.

8

does

does greater injury to his posterity than the prodigal who fquanders the other.

Of the Clothing of Children.

The clothing of an infant is fo fimple a matter, that it is furprifing how any perfon fhould err in it; yet many children lofe their lives, and others are deformed, by inattention to this article.

Nature knows of no use of clothes to an infant, but to keep it warm. All that is neceffary for this purpose, is to wrap it in a fost loose covering. Were a mother left to the dictates of Nature alone, she would certainly pursue this course. But the business of dressing an infant has long been out of the hands of mothers, and has at last become a fecret which none but adepts pretend to understand.

From the most carly ages it has been thought neceffary, that a woman in labour fhould have fome perfon to attend her. This in time became a bufinefs; and, as in all others, those who were employed in it ftrove to outdo one another in the different branches of their profession. The dreffing of a child came of course to be confidered as the midwife's province; who no doubt imagined, that the more dexterity fhe could fhew in this article, the more her skill would be admired. Her attempts were feconded by the vanity of parents, who, too often defirous of making a fhew of the infant as foon as it was born, were ambitious to have as much finery heaped upon it as possible. Thus it came to be thought as neceffary for a midwife to excel in bracing and dreffing an infant, as for a forgeon to be expert in applying bandages to a broken limb; and the poor child, as foon as it came into the world, had as many rollers and wrappers applied to its body, as if every bone had been fractured in the birth ; while these were often fo tight, as not only to gall and wound its tender frame, but even to obstruct the motion of the heart, lungs, and other organs neceflary to life.

In most parts of Britain, the practice of rolling children with fo many bandages is now, in fome meafure, laid afide; but it would still be a difficult task to perfuade the generality of mankind, that the shape of an infant does not entirely depend on the care of the midwife. So far, however, are all her endeavours to mend the shape from being successful, that they conftantly operate the contrary way, and mankind become deformed in proportion to the means used to prevent it. How little deformity of body is to be found among uncivilized nations? So little indeed, that it is vulgarly believed they put all their deformed children to death. The truth is, they hardly know fuch a thing as a deformed child. Neither should we, if we followed their example. Savage nations never think of manacling their children. They allow them the full use of every organ, carry them abroad in the open air, wafh their bodies daily in cold water, &c. By this management their children become fo ftrong and hardy, that by the time our puny infants get out of the nurfe's arms, theirs are able to fhift for themfelves *.

Among brute animals, no art is neceffary to procure a fine fhape. Though many of them are extremely delicate when they come into the world, yet we never find them grow crooked for want of fwaddling bands. Is Nature lefs generous to the human kind? No: but we take the bufinefs out of Nature's hands.

Not only the analogy of other animals, but the very feelings of infants tell us, they ought to be kept eaty and free from all preffure. They cannot indeed tell their complaints, but they can fhew figns of pain; and this they never fail to do, by crying when hurt by their slothes. No fooner are they freed from their bracings, than they feem pleafed and happy: yet, ftrange infa-

• A friend of mine, who was feveral years on the coast of Africa, tells me, that the natives neither put any clothes upon their children, nor apply to their bodies bandages of any kind, but lay them on a pallet, and fuffer them to tumble about at pleasure; yet they are all straight, and feldom have any difease. tuation! the moment they hold their peace, they are again committed to their chains.

If we confider the body of an infant as a bundle of foft pipes, replenished with fluids in continual motion, the danger of preffure will appear in the strongest light. Nature, in order to make way for the growth of children, has formed their bodies fost and flexible; and less they should receive any injury from preffure in the womb, has surrounded the *fatus* everywhere with fluids. This shews the care which Nature takes to prevent all unequal preffure on the bodies of infants, and to defend them against every thing that might in the least cramp or confine their motions.

Even the bones of an infant are fo foft and cartilaginous, that they readily yield to the flighteft preffure, and eafily affume a bad fhape, which can never after be remedied. Hence it is, that fo many people appear with high fhoulders, crooked fpines, and flat breafts, who were as well-proportioned at their births as others, but had the misfortune to be fqueezed out of fhape by the application of ftays and bandages.

Preffure, by obftructing the circulation, likewife prevents the equal diffribution of nourifhment to the different parts of the body, by which means the growth becomes unequal. One part grows too large, while another remains too fmall; and thus in time the whole frame becomes difproportioned and misfhapen. To this we must add, that when a child is cramped in its clothes, it naturally fhrinks from the part that is hurt; and by putting its body into unnatural poftures, it becomes deformed by habit.

Deformity of body may indeed proceed from weaknefs or difeafe; but, in general, it is the effect of improper clothing. Nine-tenths, at leaft, of the deformity among mankind muft be imputed to this caufe. A deformed body is not only difagreeable to the eye, but by a bad figure both the animal and vital functions muft be impeded, and of courfe health impaired. Hence few people remarkably misfhapen are ftrong or healthy.

6

The new motions which commence at the birth, as the circulation of the whole mafs of blood through the lungs, refpiration, the periftaltic motion, & c. afford another firong argument for keeping the body of an infant free from all preffure. These organs, not having been accustomed to move, are easily flopped; but when this happens, death must enfue. Hardly any method could be devifed more effectually to stop these motions, than bracing the body too tight with rollers * and bandages. Were these to be applied in the fame manner to the body of an adult for an equal length of time, they would hardly fail to hurt the digestion and make him fick. How much more hurtful they must prove to the tender bodies of infants, we shall leave any one to judge.

Whoever confiders thefe things, will not be furprifed that fo many children die of convultions foon after the birth. Thefe fits are generally attributed to fome inward caufe; but, in fact, they oftener proceed from our own imprudent conduct. I have known a child feized with convultion-fits foon after the widwife had done fwaddling it, who, upon taking off the rollers and bandages, was immediately relieved, and never had the difeafe afterwards. Numerous examples of this might be given, were they neceffary.

It would be fafer to fasten the clothes of an infant with strings than pins, as these often gall and irritate their tender skins, and occasion diforders. Pins have been found sticking above half an inch into the body of a child, after it had died of convulsion fits, which in all probability proceeded from that cause.

Children are not only hurt by the tightness of their clothes, but also by the quantity. Every child has fome degree of fever after the birth; and if it be loaded with too many clothes, the fever must be increased. But this is not all; the child is generally laid in bed

with

[•] This is by no means inveighing against a thing that does not happen. In many parts of Britain at this day, a roller, eight or ten feet in length, is applied tightly tound the child's body as foon as it is born.

with the mother, who is often likewife feverifh: to which we may add the heat of the bed-chamber, the wines, and other heating things, too frequently given to children immediately after the birth. When all thefe are combined, which does not feldom happen, they mult increase the fever to fuch 'a degree as will endanger the life of the infant.

The danger of keeping infants too hot will further appear, if we confider that, after they have been for fome time in the fituation mentioned above, they are often fent into the country to be nurfed in a cold houfe. Is it any wonder, if a child, from fuch a transition, catches a mortal cold, or contracts fome other fatal difeafe? When an infant is kept too hot, its lungs, not being fufficiently expanded, are apt to remain weak and flaccid for life; hence proceed coughs, confumptions, and other difeafes of the breaft.

It would answer little purpose to specify the particular species of dress proper for an infant. These will always vary in different countries, according to custom and the humour of parents. The great rule to be observed is, That a child have no more clothes than are necessary to keep it warm, and that they be quite cosy for its body.

Stays are the very bane of infants. A volume would not fuffice to point out all the bad effects of this ridiculous piece of drefs both on children and adults. The madnefs in favour of ftays feems, however, to be fomewhat abated; and it is to be hoped the world will, in time, become wife enough to know, that the human fhape does not folely depend upon whale-bone and bend leather *.

* Stays made of bend leather are worn by all the women of lower flation in many parts of England.

I am forry to understand, that there are still mothers mad enough to lace their daughters very tight in order to improve their shape. As reasoning would be totally lost upon such people, I shall beg leave just to ask them, Why there are ten deformed women for one man ? and likewise to recommend to their perusal a faort moral precept, which forbids us to deform the human body.

I fhall

I fhall only add with refpect to the clothes of children, that they ought to be kept thoroughly clean. Children perfpire more than adults; and if their clothes be not frequently changed, they become very hurtful. Dirty clothes not only gall and fret the tender fkins of infants, but likewife occafion ill-fmells; and, what is worfe, tend to produce vermin and cutaneous difeafes.

Cleanlinefs is not only agreeable to the eye, but tends greatly to preferve the health of children. It promotes the perfpiration, and, by that means, frees the body from fuperfluous humours, which, if retained, could not fail to occafion difeafes. No mother or nurfe can have any excufe for allowing a child to be dirty. Poverty may oblige her to give it coarfe clothes; but if fhe does not keep them clean, it must be her own fault.

Of the Food of Children.

Nature not only points out the food proper for an infant, but actually prepares it. This, however, is not fufficient to prevent fome who think themfelves wifer than Nature, from attempting to bring up their children without her provision. Nothing can shew the disposition which mankind have to depart from Nature, more than their endeavouring to bring up children without the breast. The mother's milk, or that of a healthy nurse, is unquestionably the best food for an infant. Neither art nor Nature can afford a proper substitute for it. Children may seem to thrive for a few months without the breast; but when teething, the small-pox, and other diseases incident to childhood, come on, they generally perish.

A child, foon after the birth, fhews an inclination to fuck; and there is no reafon why it fhould not be gratified. It is true, the mother's milk does not always come immediately after the birth; but this is the way to bring it: befides, the first milk that the child can fqueeze fqueeze out of the breaft answers the purpose of cleanfing, better than all the drugs in the apothecary's shop, and at the same time prevents inflammations of the breast, fevers, and other diseases incident to mothers.

It is strange how people came to think that the first thing given to a child should be drugs. This is beginning with medicine by times, and no wonder if they generally end with it. It fometimes happens, indeed, that a child does not discharge the meconium, fo foon as could be wifhed; this has induced phyficians, in fuch cafes, to give fomething of an opening nature to cleanfe the first paffages. Midwives have improved upon this hint, and never fail to give fyrups, oils, &c. whether they be neceffary or not. Cramming an infant with fuch indigeftible ftuff as foon as it is born, can hardly fail to make it fick, and is more likely to occafion difeafes than to prevent them. Children are feldom long after the birth without having paffage both by ftool and urine; though these evacuations may be wanting for fome time without any danger. But if children must have fomething before they be allowed the breaft, let it be a little thin water-pap, to which may be added an equal quantity of new milk; or rather water alone, with the addition of a little raw fugar. If this be given without any wines or fpiceries, it will neither heat the blood, load the ftomach, nor occafion gripes.

Upon the first fight of an infant, almost every perfor is struck with the idea of its being weak, feeble, and wanting support. This naturally suggests the need of cordials. Accordingly wines are universally mixed with the first food of children. Nothing can be more fallacious than this way of reasoning, or more hurtful to infants than the conduct founded upon it. Children require very little food for some time after the birth; and what they receive should be thin, weak, light, and of a cooling quality. A very small quantity of wine is sufficient to heat and inflame the blood of an infant; but every perfor conversant in these matters must know, that most of the diseases of infants proceed from the heat of their humours.

If

If the mother or nurfe has enough of milk, the child will need little or no other food before the third or fourth month. It will then be proper to give it, once or twice a-day, a little of fome food that is eafy of digeftion, as water-pap, milk-pottage, weak broth with bread in it, and fuch like. This will eafe the mother, will accuftom the child by degrees to take food, and will render the weaning both lefs difficult and lefs dangerous. All great and fudden transitions are to be avoided in nurfing. For this purpofe, the food of children ought not only to be fimple, but to refemble, as nearly as poffible, the properties of milk. Indeed, milk itfelf fhould make a principal part of their food, not only before they are weaned, but for fome time after.

Next to milk, we would recommend good light bread. Bread may be given to a child as foon as it fhews an inclination to chew; and it may at all times be allowed as much plain bread as it will eat. The very chewing of bread will promote the cutting of the teeth, and the difcharge of faliva, while, by mixing with the nurfe's milk in the ftomach, it will afford an excellent nourishment. Children discover an early inclination to chew whatever is put into their hands. Parents observe the inclination, but generally mistake the object. Inftead of giving the child fomething which may at once exercife its gums and afford it nourifhment, they commonly put into its hands a piece of hard metal, or impenetrable coral. A cruft of bread is the beft gum-flick. It not only answers the purpose better than any thing elfe, but has the additional properties of nourifhing the child and carrying the faliva down into the flomach, which is too valuable a liquor to be loft.

Bread, befides being ufed dry, may be many ways prepared into food for children. One of the beft methods is to boil it in water, afterwards pouring the water off, and mixing with the bread a proper quantity of new milk unboiled. Milk is both more wholefome and nourifhing this way than boiled, and is lefs apt to occafion cofficients. For a child farther advanced, bread bread may be mixed in veal or chicken broth, made into puddings, or the like. Bread is a proper food for children at all times, provided it be plain, made of wholefome grain, and well fermented; but when enriched with fruits, fugars, or fuch things, it becomes very unwholefome.

It is foon enough to allow children animal food when they have got teeth to eat it. They fhould never tafte it till after they are weaned, and even then they ought to use it fparingly. Indeed, when children live wholly on vegetable food, it is apt to four on their ftomachs; but, on the other hand, too much flesh heats the body, and occasions fevers and other inflammatory difeases. This plainly points out a due mixture of animal and vegetable food as most proper for children.

Few things prove more hurtful to infants than the common method of fweetening their food. It entices them to take more than they ought to do, which makes them grow fat and bloated. It is pretty certain, if the food of children were quite plain, that they would never take more than enough. Their exceffes are entirely owing to nurfes. If a child be gorged with food at all hours, and enticed to take it, by making it fweet and agreeable to the palate, is it any wonder that fuch a child fhould in time be induced to crave more food than it ought to have?

Children may be hurt by too little as well as too much food. After a child is weaned, it ought to be fed four or five times a day; but fhould never be accuftomed to eat in the night; neither fhould it have too much at a time. Children thrive beft with fmall quantities of food frequently given. This neither overloads the flomach nor hurts the digeftion, and is certainly most agreeable to nature.

Writers on nurfing have inveighed with fuch vehemence against giving children too much food, that many parents, by endeavouring to shun that error, have run into the opposite extreme, and ruined the constitutions of their children. But the error of pinching children in their food is more hurtful than the other extreme. C Nature

17

Nature has many ways of relieving herfelf when overcharged; but a child, who is pinched with hunger, will never become a firong or a healthy man. That errors are frequently committed on both fides, we are ready to acknowledge; but where one child is hurt by the quantity of its food, ten fuffer from the quality. This is the principal evil, and claims our ftricteft attention.

Many people imagine, that the food which they themfelves love cannot be bad for their children: but this notion is very abfurd. In the more advanced periods of life we often acquire an inclination for food, which when children we could not endure. Befides, there are many things that by habit may agree very well with the ftomach of a grown perfon, which would be hurtful to a child: as high feafoned, falted, and fmoke-dried provifions, &c. It would alfo be improper to feed children with fat meat, ftrong broths, rich foups, or the like.

All ftrong liquors are hurtful to children. Some parents teach their children to guzzle ale, and other fermented liquors, at every meal. Such a practice cannot fail to do mifchief. Thefe children feldom efcape the violence of the fmall-pox, meafles, hooping cough, or fome inflammatory diforder. Milk, water, buttermilk, or whey, are the most proper for children to drink. If they have any thing ftronger, it may be fine fmall beer, or a little wine mixed with water. The ftomachs of children can digest well enough without the affistance of warm ftimulants: besides, being naturally hot, they are eafily hurt by every thing of a heating quality.

Few things are more hurtful to children than unripe fruits. They weaken the powers of digeftion, and four and relax the ftomach, by which means it becomes a proper neft for infects. Children indeed fhew a great inclination for fruit, and I am apt to believe, that if good ripe fruit were allowed them in proper quantity, it would have no bad effects. We never find a natural inclination wrong, if properly regulated. Fruits are generally of a cooling nature, and correct the heat and acri-

18

acrimony of the humours. This is what most children require; only care should be taken left they exceed. Indeed the best way to prevent children from going to excess in the use of fruit, or eating that which is bad, is to allow them a proper quantity of what is good *.

Roots which contain a crude vifcid juice fhould be fparingly given to children. They fill the body with grofs humours, and tend to produce eruptive difeafes. This caution is peculiarly neceffary for the poor; glad to obtain, at a fmall price, what will fill the bellies of their children, they fluff them two or three times a day with crude vegetables. Children had better eat a fmaller quantity of food which yields a wholefome nourifhment, than be crammed with what their digeftive powers are unable properly to affimilate.

Butter ought likewife to be fparingly given to children. It both relaxes the ftomach, and produces grofs humours. Indeed, most things that are fat or oily have this effect. Butter when falted becomes still more hurtful. Instead of butter, fo liberally given to children in most parts of Britain, we would recommend honey. Children who eat honey are feldom troubled with worms: they are also less subject to cutaneous difeases, as itch, scabbed head, &c.

Many people err in thinking that the diet of children ought to be altogether moift. When children live entirely upon flops, it relaxes their folids, renders them weak, and difpofes them to the rickets, the fcrophula, and other glandular diforders. Relaxation is one of the most general causes of the difeases of children. Every thing, therefore, which tends to unbrace their folids, ought to be carefully avoided.

* Children are always fickly in the fruit feafon, which may be thus accounted for: Two-thirds of the fruit which comes to market in this country is really unripe; and children, not being in a condition to judge for themfelves, eat whatever they can lay their hands upon, which often proves little better than a poifon to their tender bowels. Servants, and others who have the care of children, fhould be firifily forbidden to give them any fruit without the knowledge of their parents.

We

OF CHILDREN.

We would not be underftood by these observations as confining children to any particular kind of food. Their diet may be frequently varied, provided always that fufficient regard be had to fimplicity.

Of the Exercise of Children.

Of all the caufes which confpire to render the life of man fhort and miferable, none has greater influence than the want of proper EXERCISE: healthy parents, wholefome food, and proper clothing, will avail little, where exercife is neglected. Sufficient exercife will make up for feveral defects in nurfing; but nothing can fupply the want of it. It is abfolutely neceffary to the health, the growth, and the ftrength of children.

The defire of exercife is coëval with life itfelf. Were this principle attended to, many difeafes might be prevented. But, while indolence and fedentary employments prevent two-thirds of mankind from either taking fufficient exercife themfelves, or giving it to their children, what have we to expect but difeafes and deformity among their offspring? The rickets, fo deftructive to children, never appeared in Britain till manufactures began to flourifh, and people, attracted by the love of gain, left the country to follow fedentary employments in great towns. It is amongft thefe people that this difeafe chiefly prevails, and not only deforms but kills many of their offspring.

The conduct of other young animals fhews the propriety of giving exercise to children. Every other animal makes use of its organs of motion as soon as it can, and many of them, even when under no necessfity of moving in quest of food, cannot be restrained without force. This is evidently the cafe with the calf, the lamb, and most other young animals. If these creatures were not permitted to frisk about and take exercife, they would foon die or become diseafed. The fame inclination appears very early in the human species; but as they are not able to take exercise themfelves, felves, it is the business of their parents and nurses to affist them.

Children may be exercifed various ways. The beft method, while they are light, is to carry them about in the nurfe's arms *. This gives the nurfe an opportunity of talking to the child, and of pointing out every thing that may pleafe and delight its fancy. Befides, it is much fafer than fwinging an infant in a machine, or leaving it to the care of fuch as are not fit to take care of themfelves. Nothing can be more abfurd than to fet one child to keep another; this conduct has proved fatal to many infants, and has rendered others miferable for life.

When children begin to walk, the fafeft and beft method of leading them about is by the hands. The common way, of fwinging them in leading ftrings fixed to their backs, has feveral bad confequences. It makes them throw their bodies forward, and prefs with their whole weight upon their ftomach and breaft; by this means the breathing is obftructed, the breaft flattened, and the bowels compreffed; which muft hurt the digeftion, and occafion confumptions of the lungs, and other difeafes.

It is a common notion, that if children are fet upon their feet too foon, their legs will become crooked. There is reafon to believe, that the very reverfe of this is true. Every member acquires ftrength in proportion as it is exercifed. The limbs of children are weak indeed, but their bodies are proportionally light; and had they fkill to direct themfelves, they would foon be able to fupport their own weight. Who ever heard of any other animal that became crooked by ufing its legs too foon? Indeed, if a child is not permitted to make any use of its legs till a confiderable time after their birth, and be then fet upon them with its whole weight at

* The nurfe ought to be careful to keep the child in a proper polition; as deformity is often the confequence of inattention to this circumftance. Its fituation ought alfo to be frequently changed. I have known a child's legs bent all on one fide, by the nurfe carrying it conftantly on one arm.

once,

once, there may be fome danger; but this proceeds entirely from the child's not having been accuftomed to use its legs from the beginning.

Mothers of the poorer fort think they are great gainers, by making their children lie or fit while they themfelves work. In this they are greatly miltaken. By neglecting to give their children exercife, they are obliged to keep them a long time before they can do any thing for themfelves, and to fpend more on medicine than would have paid for proper care.

To take care of their children, is the most useful business in which even the poor can be employed: but, alas! it is not always in their power. Poverty often obliges them to neglect their offspring in order to procure the necessaries of life. When this is the case, it becomes the interest as well as the duty of the public to affist them. Ten thousand times more benefit would accrue to the State, by enabling the poor to bring up their own children, than from all the hospitals * that ever can be erected for that purpose.

Whoever confiders the ftructure of the human body will foon be convinced of the neceffity of exercise for the health of children. The body is composed of an infinite number of tubes, whose fluids cannot be pussed on without the action and preffure of the muscles. But, if the fluids remain inactive, obstructions must happen, and the humours will of course be vitiated, which cannot fail to occasion difeases. Nature has furnished both the veffels which carry the blood and lymph with numerous valves, in order that the action of every muscle might push forward their contents; but without ac-

• If it were made the intereft of the poor to keep their children alive, we fhould lofe very few of them. A fmall premium given annually to each poor family, for every child they have alive at the year's end, would fave more infant lives than if the whole revenue of the crown were expended on hofpitals for this purpofe. This would make the poor effeem fertility a bleffing; whereas many of them think it the greateft curfe that can befal them; and in place of wifning their children to live, fo far does poverty get the better of natural affection, that they are often very happy when they die. tion, this admirable contrivance can have no effect. This part of the animal œconomy proves to a demonfiration the neceffity of exercise for the prefervation of health.

Arguments to fhew the importance of exercife might be drawn from every part of the animal œconomy; without exercife, the circulation of the blood cannot be properly carried on, nor the different fecretions duly performed; without exercife, the fluids cannot be properly prepared, nor the folids rendered flrong or firm. The action of the heart, the motion of the lungs, and all the vital functions, are greatly affifted by exercife. But to point out the manner in which these effects are produced, would lead us farther into the œconomy of the human body, than most of those for whom this treatife is intended would be able to follow. We fhall therefore only add, that when exercise is neglected, none of the animal functions can be duly performed; and when this is the case, the whole constitution must go to wreck.

A good conffitution ought certainly to be our first object in the management of children. It lays a foundation for their being useful and happy in life: and whoever neglects it, not only fails in his duty to his offspring, but to fociety.

One very common error of parents, by which they hurt the conflitutions of their children, is the fending them too young to fchool. This is often done folely to prevent trouble. When the child is at fchool, he needs no keeper. Thus the fchool-mafter is made the nurfe; and the poor child is fixed to a feat feven or eight hours a-day, which time ought to be fpent in exercife and diverfions. Sitting fo long cannot fail to produce the worft effects upon the body; nor is the mind lefs injured. Early application weakens the faculties, and often fixes in the mind an averfion to books, which continues for life *.

But

• It is undoubtedly the duty of parents to inftruct their children, at leaft till they are of an age proper to take fome care of C 4 themBut fuppofe this were the way to make children fcholars, it certainly ought not to be done at the expence of their conflitutions. Our anceftors, who feldom went to fchool very young, were not lefs learned than we. But we imagine the boy's education will be quite marred, unlefs he be carried to fchool in his nurfe's arms. No wonder if fuch hot-bed plants feldom become either fcholars or men !

Not only the confinement of children in public fchools, but their number, often proves hurtful. Children are much injured by being kept in crowds within doors; their breathing not only renders the place unwholefome, but if any one of them happen to be difeafed, the reft catch the infection. A fingle child has been often known to communicate the bloody flux, the hooping cough, the itch, or other difeafes, to almost every individual in a numerous fchool.

But, if fashion must prevail, and infants are to be fent to school, we would recommend it to teachers, as they value the interests of society, not to confine them too long at a time, but allow them to run about and play at such active diversions as may promote their growth, and strengthen their constitutions. Were boys, instead of being whipped for stealing an hour to run, ride, fwim, or the like, encouraged to employ a proper part of their time in these manly and useful exercifes, it would have many excellent effects.

It would be of great fervice to boys, if, at a proper age, they were taught the military exercise. This would increase their strength, inspire them with courage, and when their country called for their affissance, would enable them to act in her defence, without being obliged to undergo a tedious and troublesome course of instruc-

themfelves. This would tend much to confirm the ties of parental tendernefs and filial affection, of the want of which there are at prefent fo many deplorable inftances. Though few fathers have time to inftruct their children, yet most mothers have; and furely they cannot be better employed.

tions,

tions, at a time when they are lefs fit to learn new motions, gestures, &c *

An effeminate education will infallibly fpoil the beft natural conftitution; and if boys are brought up in a more delicate manner than even girls ought to be, they will never be men.

Nor is the common education of girls lefs hurtful to the conflitution than that of boys. Mifs is fet down to her frame before fhe can put on her clothes; and is taught to believe, that to excel at the needle is the only thing that can entitle her to general effeem. It is unneceffary here to infift upon the dangerous confequences of obliging girls to fit too much. They are pretty well known, and are too often felt at a certain time of life. But fuppofing this critical period to be got over, greater dangers ftill wait them when they come to be mothers. Women who have been early accuftomed to a fedentary life, generally run great hazard in childbed; while thofe who have been ufed to romp about, and take fufficient exercife, are feldom in any danger.

One hardly meets with a girl who can at the fame time boaft of early performances by the needle, and a good conftitution. Clofe and early confinement generally occafions indigeftions, head-achs, pale complexions, pain of the ftomach, lofs of appetite, coughs, confumptions of the lungs, and deformity of body. The laft of thefe indeed is not to be wondered at, confidering the awkward poftures in which girls fit at many kinds of needle-work, and the delicate flexible ftate of their bodies in the early periods of life.

Would mothers, instead of having their daughters instructed in many trifling accomplishments, employ them in plain work and housewifery, and allow them

* I am happy to find that the mafters of academies now begin to put in practice this advice. Each of them ought to keep a drill-ferjeant for teaching the boys the military exercise. This, befides contributing to their health and vigour of body, would have many other happy effects. fufficient exercife in the open air, they would both make them more healthy mothers, and more ufeful members of fociety. I am no enemy to genteel accomplifhments, but would have them only confidered as fecondary, and always difregarded when they impair health.

Many people imagine it a great advantage for children to be early taught to earn their bread. This opinion is certainly right, provided they were fo employed as not to hurt their health or growth; but, when thefe fuffer, fociety, inftead of being benefited, is a real lofer by their labour. There are few employments, except fedentary ones, by which children can earn a livelihood; and if they be fet to thefe too foon, it ruins their conftitutions. Thus, by gaining a few years from childhood, we generally lofe twice as many in the latter period of life, and even render the perfon lefs ufeful while he does live.

In order to be fatisfied of the truth of this obfervation, we need only look into the great manufacturing towns, where we shall find a puny degenerate race of people, weak and fickly all their lives, feldom exceeding the middle period of life; or if they do, being unfit for bufinefs, they become a burden to fociety. Thus arts and manufactures, though they may increase the riches of a country, are by no means favourable to the health of its inhabitants. Good policy would therefore require, that fuch people as labour during life, fhould not be fet too early to work. Every perfon converfant in the breed of horfes, or other working animals, knows, that if they be fet to hard labour too foon, they never will turn out to advantage. This is equally true with refpect to the human fpecies.

There are, nevertheles, various ways of employing young people, without hurting their health. The easier parts of gardening, husbandry, or any business carried on without doors, are most proper. These are employments which most young people are fond of, and some parts

26

parts of them may always be adapted to their age, tafte, and ftrength *.

Such parents, however, as are under the neceffity of employing their children within doors, ought to allow them fufficient time for active diversions without. This would both encourage them to do more work, and prevent their conflictutions from being hurt.

Some imagine, that exercife within doors is fufficient; but they are greatly miltaken. One hour fpent in running, or any other exercife without doors, is worth ten within. When children cannot go abroad, they may indeed be exercifed at home. The beft method of doing this, is to make them run about in a long room, or dance. This laft kind of exercife, if not carried to excefs, is of excellent fervice to young people. It cheers the fpirits, promotes perfpiration, ftrengthens the limbs, &c. I knew an eminent phyfician who ufed to fay, that he made his children dance, inftead of giving them phyfic. It were well if more people followed his example.

The COLD BATH may be confidered as an aid to exercife. By it the body is braced and ftrengthened, the circulation and fecretions promoted, and, were it conducted with prudence, many difeafes, as rickets, fcrophula, &c. might thereby be prevented. The ancients, who took every method to render children hardy and robuft, were no ftrangers to the ufe of the cold bath ; and, if we may credit report, the practice of immerfing children daily in cold water muft have been very common among our anceftors.

The greatest objection to the use of the cold bath arises from the superstitutions prejudices of nurses. These are often so strong, that it is impossible to bring them to make a proper use of it. I have known some of them who would not dry a child's skin after bathing it, left it should destroy the effect of the water. Others

* I have been told that in China, where the police is the beft in the world, all the children are employed in the eafier part of gardening and hufbandry; as weeding, gathering ftones off the land, and fuch like. will even put clothes dipt in the water upon the child, and either put it to bed, or fuffer it to go about in that condition. Some believe, that the whole virtue of the water depends upon its being dedicated to a particular faint; while others place their confidence in a certain number of dips, as three, feven, nine, or the like; and the world could not perfuade them, if thefe do not fucceed, to try it a little longer. Thus, by the whims of nurfes, children lofe the benefit of the cold bath, and the hopes of the phyfician from that medicine are often fruftrated.

We ought not, however, entirely to fet afide the cold bath, becaufe fome nurles make a wrong ufe of it. Every child, when in health, fhould at leaft have its extremities daily wafhed in cold water. This is a partial ufe of the cold bath, and is better than none. In winter this may fuffice; but, in the warm feafon, if a child be relaxed, or feem to have a tendency to the rickets or fcrophula, its whole body ought to be frequently immerfed in cold water. Care however muft be taken not to do this when the body is hot, or the ftomach full. The child fhould be dipped only once at a time, fhould be taken out immediately, and have its fkin well rubbed with a dry cloth.

The bad Effects of unwholefome Air upon Children.

Few things prove more destructive to children than confined or unwholesome air. This is one reason why fo few of those infants, who are put into hospitals, or parish workhouses, live. These places are generally crowded with old, fickly, and infirm people; by which means the air is rendered fo extremely pernicious, that it becomes a poison to infants.

Want of wholefome air is likewife deftructive to many of the children born in great towns. There the poorer fort of inhabitants live in low, dirty, confined houfes, to which the fresh air has fearcely any access. Though grown people, who are hardy and robust, may live in such fituations, yet they generally prove fatal to their their offspring, few of whom arrive at maturity, and those who do are weak and deformed. As fuch people are not in a condition to carry their children abroad into the open air, we must lay our account with losing the greater part of them. But the rich have not this excuse. It is their business to see that their children be daily carried abroad, and that they be kept in the open air for a sufficient time. This will always succeed better if the mother goes along with them. Servants are often negligent in these matters, and allow a child to fit or lie on the damp ground, instead of leading or carrying it about. The mother furely needs air as well as her children ; and how can she be better employed than in attending them ?

A very bad cuftom prevails, of making children fleep in fmall apartments, or crowding two or three beds into one chamber. Inftead of this, the nurfery ought always to be the largest and best aired room in the house. When children are confined in fmall apartments, the air not only becomes unwholefome, but the heat relaxes their folids, renders them delicate, and difpofes them to colds and many other diforders. Nor is the cuftom of wrapping them up too close in cradles less pernicious. One would think that nurfes were afraid left children should suffer by breathing free air, as many of them actually cover the child's face while afleep, and others wrap a covering over the whole cradle, by which means the child is forced to breathe the fame air over and over all the time it fleeps. Cradles indeed are on many accounts hurtful to children, and it would be better if the use of them were totally laid afide *.

A child

* It is amazing how children escape fuffocation, confidering the manner in which they are often rolled up in flannels, &c. I lately attended an infant, whom 1 found muffled up over head and ears in many folds of flannel, though it was in the middle of June. I begged for a little free air to the poor babe; but though this indulgence was granted during my Itay, I found it always on my return in the fame fituation. Death, as might be expected, foon freed the infant from all its miferies: but it was not in my power to free the minds of its parents from those prejudices which proved fatal to their child.

I was

A child is generally laid to fleep with all its clothes on; and if a number of others are heaped above them, it must be overheated; by which means it cannot fail to catch cold on being taken out of the cradle, and exposed to the open air with only its usual clothing, which is too frequently the case.

Children who are kept within doors all day, and fleep all night in warm clofe apartments, may, with great propriety, be compared to plants, nurfed in a hot-houfe, inftead of the open air. Though fuch plants may by this means be kept alive for fome time, they will never arrive at that degree of ftrength, vigour, and magnitude, which they would have acquired in the open air, nor would they be able to bear it afterwards, fhould they be expofed to it.

Children brought up in the country, who have been accuftomed to open air, fhould not be too early fent to great towns, where it is confined and unwholefome. This is frequently done with a view to forward their education, but proves very hurtful to their health. All fchools and feminaries of learning ought, if poffible, to be fo fituated as to have fresh, dry, wholefome air, and should never be too much crowded.

Without entering into a detail of the particular advantages of wholefome air to children, or of the bad confequences which proceed from the want of it, I fhall only obferve, that of feveral thoufands of children which have been under my care, I do not remember one inftance of a fingle child who continued healthy in a clofe confined fituation; but have often known the most obstinate difeases cured by removing them from fuch a fituation to an open free air.

Of Nurses.

It is not here intended to lay down rules for the choice of nurfes. This would be wafting time. Com-

I was very lately called to fee an infant which was faid to be expiring in convultion-fits. I defired the mother to ftrip the child, and wrap it in a loofe covering. It had no more convultion-fits. mon mon fenfe will direct every one to choofe a woman who is healthy, and has plenty of milk *. If fhe be at the fame time cleanly, careful, and good-natured, fhe can hardly fail to make a proper nurfe. After all, however, the only certain proof of a good nurfe, is a healthy child upon her breaft. But, as the mifconduct of nurfes often proves fatal to children, it will be of importance to point out a few of their moft baneful errors, in order to roufe the attention of parents, and to make them look more ftrictly into the conduct of those to whom they commit the care of their infant offspring.

Though it admits of fome exceptions, yet we may lay it down as a general rule, *That every woman who nurfes for hire fhould be carefully looked after*, otherwife *fhe will not do her duty*. For this reafon parents ought always to have their children nurfed under their own eye, if poffible; and where this cannot be done, they fhould be extremely circumfpect in the choice of thofe perfons to whom they intruft them. It is folly to imagine that any woman, who abandons her own child to fuckle another for the fake of gain, fhould feel all the affections of a parent towards her nurfling: yet fo neceffary are thefe affections in a nurfe, that, but for them, the human race would foon be extinct.

One of the most common faults of those who nurfe for hire, is doing children with ftupefactives, or fuch things as lull them alleep. An indolent nurfe, who does not give a child fufficient exercise in the open air to make it fleep, and does not choose to be diffurbed by it in the night, will feldom fail to procure for it a dose of laudanum, diacodium, faffron, or what answers the fame purpose, a dose of spirits or other strong liquors. These, though they be certain possion to infants, are every day administered by many who bear the character of very good nurses \dagger .

* I have often known people fo imposed upon, as to give an infant to a nurse to be suckled who had not one drop of milk in her breast.

† If a mother on visiting her child at nurse finds it always assert, I would advise her to remove it immediately; otherwise it will soon fleep its last.

A nurfe

A nurfe who has not milk enough is apt to imagine that this defect may be fupplied by giving the child wines, cordial waters, or other ftrong liquors. This is an egregious miftake. The only thing that has any chance to fupply the place of the nurfe's milk, muft be fomewhat nearly of the fame quality, as cow's milk, afs's milk, or beef tea, with a little bread. It never can be done by the help of ftrong liquors. Thefe, inftead of nourifhing an infant, never fail to produce the contrary effect.

Children are often hurt by nurfes fuffering them to cry long and vehemently. This ftrains their tender bodies, and frequently occafions ruptures, inflammations of the throat, lungs, &c. A child never continues to cry long without fome caufe, which might always be difcovered by proper attention; and the nurfe who can hear an infant cry till it has almost fpent itfelf, without endeavouring to pleafe it, must be cruel indeed, and is unworthy to be intrusted with the care of an human creature.

Nurfes who deal much in medicine are always to be fufpected. They truft to it, and neglect their duty. I never knew a good nurfe who had her Godfrey's cordial, Daffy's elixirs, Dalby's carminative, &c. at hand. Such generally imagine, that a dofe of medicine will make up for all defects in food, air, exercife, and cleanlinefs. By errors of this kind, I will venture to fay, that one half the children who die annually in London lofe their lives.

Allowing children to continue long wet, is another very pernicious cuftom of indolent nurfes. This is not only difagreeable, but it galls and frets the infant, and, by relaxing the folids, occafions fcrophulas, rickets, and other difeafes. A dirty nurfe is always to be fufpected.

Nature often attempts to free the bodies of children from bad humours, by throwing them upon the fkin: by this means fevers and other difeafes are prevented. Nurfes are apt to miftake fuch critical eruptions for an itch, or fome other infectious diforder. Accordingly they they take every method to drive them in. In this way many children lofe their lives; and no wonder, as Nature is opposed in the very method she takes to relieve them. It ought to be a rule, which every nurfe fhould observe, never to stop any eruption without proper advice, or being well affured that it is not of a critical nature. At any rate, it is never to be done without previous evacuations.

Loofe ftools is another method by which Nature often prevents or carries off the difeafes of infants. If these proceed too far, no doubt they ought to be checked; but this is never to be done without the greatest caution. Nurfes, upon the first appearance of loofe stools, frequently fly to the use of aftringents, or fuch things as bind the body. Hence inflammatory fevers, and other fatal difeases, are occasioned. A dose of rhubarb, a gentle vomit, or fome other evacuation, fhould always precede the ufe of aftringent medicines.

One of the greatest faults of nurses is, concealing the difeafes of children from their parents. This they are extremely ready to do, efpecially when the difeafe is the effect of their own negligence. Many inftances might be given of perfons who have been rendered lame for life by a fall from their nurfe's arms, which fhe, through fear, concealed till the misfortune was past cure. Every parent who intrusts a nurse with the care of a child, ought to give her the ftricteft charge not to conceal the most triffing diforder or misfortune that may befal it.

We can fee no reafon why a nurfe, who conceals any misfortune which happens to a child under her care, till it loses its life or limbs, should not be punished. A few examples of this would fave the lives of many infants; but as there is little reafon to expect that it ever will be the cafe, we would earneftly recommend it to all parents to look carefully after their children, and not to truft fo valuable a treasure entirely in the hands of an hireling.

No perfon ought to imagine these things unworthy of his attention. On the proper management of children depend not only their health and usefulness in life, but likewife the fafety and profperity of the ftate to which they

they belong. Effeminacy ever will prove the ruin of any flate where it prevails; and, when its foundations are laid in infancy, it can never afterwards be wholly eradicated. Parents who love their offspring, and wifh well to their country, ought, therefore, in the management of their children, to avoid every thing that may have a tendency to make them weak or effeminate, and to take every method in their power to render their conflitutions flrong and hardy.

By arts like these Laconia nurs'd of old her hardy fons; And Rome's unconquer'd legions urg'd their way, Unhurt, thro' every toil in every clime *.

Few things tend more to the deftruction of children than drenching them with drugs. That medicine may be *fometimes* neceffary for children, I do not deny; but that it hurts them ten times for once it does them good, I will venture to affert. A London mother, the moment her child feems to ail any thing, runs immediately to the apothecary, who throws in his powders, pills, and potions, till the poor infant is poifoned; when the child might have been reftored to perfect health by a change of diet, air, exercife, clothing, or fome very eafy and fimple regulation.

But mifguided fondnefs is not fatisfied with drugging children from the apothecary's fhop: many of them are *fed* from the fame quarter. A ftarch from the Weft Indies, called flour of arrow-root, is the food of thofe infants whofe parents can afford to pay for it. I lately offended a mother very much by faying, it was not half fo good as oatmeal, though more than ten times the price. Of this, 'however, fhe had fufficient proof by a child in her arms, who had been fed on that root, and, though a year and a half old, could fcarcely put a foot to the ground, while her neighbour's child, only nine months old, but nurfed in the north country manner, could, by a hold of the finger, run all over the houfe.

* Armftrong.

I have

I have taken notice of this powder, to fhew the influence of fashion even in the feeding of an infant. I with it were the only instance I could give of the fatal effects of the fame cause. Ten thousand infants in this island, before they are out of the nurse's arms, fip tea twice a day; which, to be fure, is the true way to propagate heroes!

CHAP. II.

OF THE LABORIOUS, THE SEDENTARY, AND THE STUDIOUS.

THAT men are exposed to particular diseafes from the occupations which they follow, is a fact well known; but to remedy this evil is a matter of fome difficulty. Most people are under the necessity of following those employments to which they have been bred, whether they be favourable to health or not. For this reason, instead of inveighing, in a general way, as fome authors have done, against those occupations which are hurtful to health, we shall endeavour to point out the circumstances in each of them from which the danger chiefly arises, and to propose the most rational methods of preventing it.

Chymifts, founders, forgers, glafs-makers, and feveral other artifts, are hurt by the unwholefome air which they are obliged to breathe. This air is not only loaded with the noxious exhalations arifing from metals and minerals, but is fo charged with phlogifton as to be rendered unfit for expanding the lungs fufficiently, and anfwering the other important purpoles of refpiration. Hence proceed afthmas, coughs, and confumptions of the lungs, fo incident to perfons who follow thefe employments.

To

To prevent fuch confequences as far as poffible, the places where these occupations are carried on ought to be conftructed in fuch a manner as to discharge the smoke and other exhalations, and admit a free current of fresh air. Such artists ought never to continue long at work; and when they give over, they should fuffer themselves to cool gradually, and put on their clothes before they go into the open air. They ought never to drink large quantities of cold, weak, or watery liquors while their bodies are hot, nor to indulge in raw fruits, fallads, or any thing that is cold on the stomach *.

Miners, and all who work under ground, are likewife hurt by unwholefome air. The air, by its ftagnation in deep mines, not only lofes its proper fpring and other qualities neceffary for refpiration, but is often loaded with fuch noxious exhalations as to become a most deadly poifon.

The two kinds of air which prove most destructive to miners, are what they call the *fire damp*, and the *choke damp*. In both cases the air becomes a poison by its being loaded with phlogiston. The danger from the former may be obviated by making it explode before it accumulates in too great quantities; and the latter may be generally carried off by promoting a free circulation of air in the mine.

Miners are not only hurt by unwholefome air, but likewife by the particles of metal which adhere to their fkin, clothes, &c. Thefe are abforbed, or taken up into the body, and occafion palfies, vertigoes, and other nervous affections, which often prove fatal. Fallopius obferves, that thofe who work in mines of mercury feldom live above three or four years. Lead, and feveral other metals, are likewife very pernicious to the health.

Miners ought never to go to work fasting, nor to continue too long at work. Their food ought to be

* When perfons heated with labour have drunk cold liquor, they ought to continue at work for fome time after.

nourifhing,

nourifhing, and their liquor generous: nothing more certainly hurts them than living too low. They fhould by all means avoid coftivenels. This may either be done by chewing a little rhubarb, or taking a fufficient quantity of fallad oil. Oil not only opens the body, but fheathes and defends the inteftines from the ill effects of the metals. All who work in mines or metals ought to wash carefully, and to change their clothes as foon as they give over working. Nothing would tend more to preferve the health of such people than a strict, and almost religious, regard to cleanlinels.

Plumbers, painters, gilders, fmelters, makers of white lead, and many others who work in metals, are liable to the fame difeafes as miners; and ought to obferve the fame directions for avoiding them.

Tallow-chandlers, boilers of oil, and all who work in putrid animal fubftances, are likewife liable to fuffer from the unwholefome fmells or effluvia of thefe bodies. They ought to pay the fame regard to cleanlinefs as miners; and when they are affected with naufea, ficknefs, or indigeftion, we would advife them to take a vomit or a gentle purge. Such fubftances ought always to be manufactured as foon as poffible. When long kept, they not only become unwholefome to thofe who manufacture them, but likewife to people who live in the neighbourhood.

It would greatly exceed the limits of this part of our fubject, to specify the difeases peculiar to perfons of every occupation; we shall therefore confider mankind under the general classes of *Laborious*, *Sedentary*, and *Studious*.

THE LABORIOUS.

Though those who follow laborious employments are in general the most healthy of mankind, yet the nature of their occupations, and the places where they are carried on, expose them more particularly to some difeases. Husbandmen, for example, are exposed to all the viciffitudes of the weather, which, in this country,

are

are often very great and fudden, and occafion colds, coughs, quinfies, rheumatifms, fevers, and other acute diforders. They are likewife forced to work hard, and often to carry burdens above their ftrength, which, by overftraining the veffels, occafion afthmas, ruptures, pleurifies, &c.

Those who labour without doors are often afflicted with intermitting fevers or agues, occasioned by the frequent viciffitudes of heat and cold, poor living, bad water, fitting or lying on the damp ground, evening dews, night air, &c. to which they are frequently exposed.

Such as bear heavy burdens, as porters, labourers, &c. are obliged to draw in the air with much greater force, and alfo to keep their lungs diftended with more violence than is neceffary for common refpiration: by this means the tender veffels of the lungs are overftretched, and often burft, infomuch that a fpitting of blood or fever enfues. Hippocrates mentions an inftance to this purpofe, of a man, who, upon a wager, carried an afs; but was foon after feized with a fever, a vomiting of blood, and a rupture

Carrying heavy burdens is generally the effect of mere lazinefs, which prompts people to do at once what fhould be done at twice. Sometimes it proceeds from vanity or emulation. Hence it is, that the ftrongeft men are most commonly hurt by heavy burdens, hard labour, or feats of activity. It is rare to find one who boasts of his ftrength without a rupture, a spitting of blood, or some other difease, which he reaps as the fruit of his folly. One would imagine the daily inftances we have of the fatal effects of carrying great weights, running, wrestling, and the like, would be sufficient to prevent fuch practices.

There are indeed fome employments which neceffarily require a great exertion of firength; as porters, blackfmiths, carpenters, &c. None ought to follow thefe but men of firong body; and they fhould never exert their firength to the utmost, nor work too long. When the muscles are violently firained, frequent rest is neceffary,

10

³

in order that they may recover their tone; without this, the ftrength and conftitution will foon be worn out, and a premature old age be induced.

The erifipelas, or St. Anthony's fire, is a difeafe very incident to the laborious. It is occafioned by whatever gives a fudden check to the perfpiration, as drinking cold water when the body is warm, wet feet, keeping on wet clothes, fitting or lying on the damp ground, &c. It is impossible for those who labour without doors always to guard against these inconveniences; but it is known from experience, that their ill confequences might often be prevented by proper care.

The iliac paffion, the colic, and other complaints of the bowels, are often occafioned by the fame caufes as the erifipelas; but they may likewife proceed from flatulent and indigestible food. Labourers generally eat unfermented bread, made of peas, beans, rye, and other windy ingredients. They also devour great quantities of unripe fruits, baked, flewed, or raw, with various kinds of roots and herbs, upon which they drink four milk, stale small-beer, or the like. Such a mixture cannot fail to fill the bowels with wind, and occasion difeafes of those parts.

Inflammations, whitloes, and other difeafes of the extremities, are likewife common among those who labour without doors. These diseases are often attributed to venom, or fome kind of poifon: but they generally proceed either from fudden heat after cold, or the contrary. When labourers, milk-maids, &c. come from the field, cold or wet, they run to the fire, and often plunge their hands in warm water; by which means the blood and other humours in those parts are fuddenly expanded, and, the veffels not yielding fo quickly, a ftrangulation happens, and an inflammation or a mortification enfues.

When fuch perfons come home cold, they ought to keep at a diffance from the fire for fome time, to wash their hands in cold water, and to rub them well with a dry cloth. It fometimes happens, that people are

39

fo

fo benumbed with cold, as to be quite deprived of the use of their limbs. In this case the only remedy is to rub the parts affected with snow, or, where it cannot be had, with cold water. If they be held near the fire, or plunged into warm water, a mortification will generally enfue.

Labourers in the hot feafon are apt to lie down and fleep in the fun. This practice is fo dangerous, that they often awake in a burning fever. Thefe ardent fevers, which prove fo fatal about the end of fummer and beginning of autumn, are frequently occafioned by this means. When labourers leave off work, which they ought always to do during the heat of the day, they fhould go home, or at leaft get under fome cover, where they may repofe themfelves in fafety.

Many people follow their employments in the fields from morning till night, without eating any thing. This cannot fail to hurt their health. However homely their fare be, they ought to have it at regular times; and the harder they work, the more frequently they fhould eat. If the humours be not frequently replenifhed with fresh nourisfiment, they foon become putrid, and produce fevers of the very worst kind.

Many peafants are extremely carelefs with refpect to what they eat or drink, and often, through mere indolence ufe unwholefome food, when they might, for the fame expence, have that which is wholefome. In fome parts of Britain, the peafants are too carelefs even to take the trouble of dreffing their own victuals. Such people would live upon one meal a-day in indolence, rather than labour, though it were to procure them the greateft affluence.

Fevers of a very bad kind are often occafioned among labourers by poor living. When the body is not fufficiently nourifhed, the humours become vitiated, and the folids weak; from whence the most fatal confequences enfue. Poor living is likewife productive of many of those cutaneous difeases fo frequent among the lower class of people. It is remarkable that cattle, when pinched in their food, are generally affected with difeases difeafes of the skin, which feldom fail to disappear when they are put upon a good pasture. This shews how much a good state of the humours depends upon a sufficient quantity of proper nourishment.

Poverty not only occafions, but aggravates, many of the difeafes of the laborious. Few of them have much forefight; and if they had, it is feldom in their power to fave any thing. They are glad to make a fhift to live from day to day; and when any difeafe overtakes them, they are miferable indeed. Here the godlike virtue of charity ought always to exert itfelf. To relieve the industrious poor in distrefs, is furely the most exalted act of religion and humanity. They alone, who are witneffes of those scenes of calamity, can form a notion of what numbers perish in difeases, for want of proper affistance, and even for want of the necessaries of life.

Labourers are often hurt by a foolifh emulation, which prompts them to vie with one another, till they overheat themfelves to fuch a degree as to occafion a fever, or even to drop down dead. Such as wantonly throw away their lives in this manner, deferve to be looked upon in no better light than felfmurderers.

The office of a *foldier*, in time of war, may be ranked among the laborious employments. Soldiers fuffer many hardfhips from the inclemency of feafons, long marches, bad provifions, hunger, watching, unwholefome climates, bad water, &c. Thefe occafion fevers, fluxes, rheumatifms, and other fatal difeafes, which generally do greater execution than the fword, efpecially when campaigns are continued too late in the feafon. A few weeks of cold rainy weather will often prove more fatal than an engagement.

Those who have the command of armies should take care that their foldiers be well clothed and well fed. They ought also to finish their campaigns in due feason, and to provide their men with dry and well-aired winter-quarters. These rules, taking care, at the same time, to keep the fick at a proper distance from those in

1

ţ

li

h

18

in health, would tend greatly to preferve the lives of the foldiery *.

Sailors may alfo be numbered among the laborious. They undergo great hardfhips from change of climate, the violence of the weather, hard labour, bad provifions, &c. Sailors are of fo great importance both to the trade and fafety of this kingdom, that too much pains can never be beftowed in pointing out the means of preferving their lives.

One great fource of the difeafes of fea-faring people is excefs. When they get on fhore, after having been long at fea, without regard to the climate, or their own conftitutions, they plunge headlong into all manner of riot, and often perfift till a fever puts an end to their lives. Thus intemperance, and not the climate, is often the caufe why fo many of our brave failors die on foreign coafts. Such people ought not to live too low; but they will find moderation the beft defence against fevers and many other maladies.

Sailors, when on duty, cannot avoid fometimes getting wet. When this happens, they fhould change their clothes as foon as they are relieved, and take every method to reftore the perfpiration. They fhould not, in this cafe, make too free with fpirits or other ftrong

* It is indeed to be regretted, that foldiers fuffer not lefs from indolence and intemperance in time of peace, than from hardfhips in time of war. If men are idle they will be vicious. It would therefore be of great importance, could a fcheme be formed for rendering the military, in times of peace, both more healthy and more useful. These desirable objects might, in our opinion, be obtained, by employing them for fome hours every day, and advancing their pay accordingly. By this means, idlenefs, the mother of vice, might be prevented, the price of labour lowered, public works, as harbours, canals, turnpike roads, &c. might be made without hurting manufactures, and foldiers might be enabled to marry and bring up children. A fcheme of this kind might eafily be conducted, fo as not to deprefs the martial fpirit, provided the men were only to work four or five hours every day, and always to work without doors : no foldiers should be fuffered to work too long, or to follow any fedentary employment. Sedentary employments render men weak and effeminate, quite unfit for the hardfhips of war : whereas working for a few hours every day without doors, would inure them to the weather, brace their nerves, and increase their ftrength and courage.

liquors,

and the STUDIOUS.

liquors, but fhould rather drink them diluted with warm water, and go immediately to bed, where a found fleep and a gentle fweat would fet all to rights.

But the health of failors fuffers moft from unwholefome food. The conftant use of falted provisions inflames their humours, and occasions the fcurvy, and other obstinate maladies. It is no easy matter to prevent this difease in long voyages; yet we cannot help thinking, that much might be done towards effecting fo defirable an end, were due pains bestowed for that purpose. For example, various roots, greens, and fruits, might be kept a long time at fea, as onions, potatoes, cabbages, lemons, oranges, tamarinds, apples, &c. When fruits cannot be kept, the juices of them, either fresh or fermented, may. With these all the drink, and even the food, of the some of the fusion of

Stale bread and beer likewife contribute to vitiate the humours. Flour will keep for a long time on board, of which frefh bread might frequently be made. Malt too might be kept, and infufed with boiling water at any time. This liquor, when drank even in form of wort, is very wholefome, and is found to be an antidote againft the fcurvy. Small wines and cider might likewife be plentifully laid in; and fhould they turn four, they would ftill be ufeful as vinegar. Vinegar is a great antidote againft difeates, and fhould be ufed by all travellers, efpecially at fea. It may either be mixed with the water they drink, or taken in their food.

Such animals as can be kept alive, ought likewife to be carried on board, as hens, ducks, pigs, &c. Frefh broths made of portable foup, and puddings made of peas or other vegetables, ought to be used plentifully. Many other things will readily occur to people converfant in these matters, which would tend to preferve the health of that brave and useful set of men *.

* Our countryman, the celebrated Captain Cook, has fhewn how far, by proper care and attention, the difeafes formerly fo fatal to feamen may be prevented. In a voyage of three years and

We

We have reafon to believe, if due attention were paid to the diet, air, clothing, and, above all things, to the cleanlinefs of fea-faring people, they would be the most healthy fet of men in the world; but when these are neglected, the very reverse will happen.

The best medical antidote that we can recommend to failors or foldiers on foreign coafts, especially where dampness prevails, is the Peruvian bark. This will often prevent fevers, and other fatal diseases. About a drachm of it may be chewed every day; or if this should prove difagreeable, an ounce of bark, with half an ounce of orange-peel, and two drachms of fnakeroot coarfely powdered, may be infused for two or three days in an English quart of brandy, and half a wineglafs of it taken twice or thrice a-day, when the ftomach is empty. This has been found to be an excellent antidote against fluxes, putrid, intermitting, and other fevers, in unhealthy climates. It is not material in what form this medicine is taken. It may either be infused in water, wine, or spirits, as recommended above, or made into an electuary with fyrup of lemons, oranges, or the like.

THE SEDENTARY.

Though nothing can be more contrary to the nature of man than a fedentary life, yet this clafs comprehends by far the greater part of the fpecies. Almost the whole female world, and, in manufacturing countries, the major part of the males, may be reckoned fedentary *.

and eighteen days, during which he was exposed to every climate, from the 52° north to the 71° of fouth latitude, of one hundred and eighteen men, composing the ship's company, he lost only one, who died of a *phthis pulmonalis*. The principal means he used were, to preferve a strict attention to cleanlines, to procure abundance of vegetables and fresh provisions, especially good water, and to allow his people sufficient time for rest.

* The appellation of fedentary has generally been given only to the fludious; we can fee no reafon, however, for reftricting it to them alone. Many artificers may, with as much propriety, be denominated fedentary as the fludious, with this particular difadvantage, that they are often obliged to fit in very awkward poftures, which the fludious need not do, unlefs they pleafe.

Agriculture,

and the STUDIOUS.

Agriculture, the first and most healthful of all employments, is now followed by few who are able to carry on any other busines. But those who imagine that the culture of the earth is not sufficient to employ all its inhabitants, are greatly mistaken. An ancient Roman, we are told, could maintain his family from the produce of one acre of ground. So might a modern Briton, if he would be contented to live like a Roman. This shews what an immense increase of inhabitants Britain might admit of, and all of them live by the culture of the ground.

Agriculture is the great fource of domeftic riches. Where it is neglected, whatever wealth may be imported from abroad, poverty and mifery will abound at home. Such is, and ever will be, the fluctuating ftate of trade and manufactures, that thoufands of people may be in full employment to-day and in beggary to-morrow. This can never happen to those who cultivate the ground. They can eat the fruit of their labour, and can always by industry obtain, at least, the neceffaries of life.

Though fedentary employments are neceffary, yet there feems to be no reafon why any perfon fhould be confined for life to thefe alone. Were fuch employments intermixed with the more active and laborious, they would never do hurt. It is conftant confinement that ruins the health. A man may not be hurt by fitting five or fix hours a-day; but if he is obliged to fit ten or twelve, he will foon become difeafed.

But it is not want of exercise alone which hurts fedentary people; they likewise suffer from the confined air which they breathe. It is very common to see ten or a dozen tailors *, or stay-makers, for example, crowded

* A perfon of obfervation in that line of life told me, that most tailors die of confumptions; which he attributed chiefly to the unfavourable postures in which they fit, and the unwholefomeness of those places where their business is carried on. If more attention were not paid to profit than to the prefervation of human lives, this evil might be easily remedied; but while masters only mind their own interest, nothing will be done for the fafety of their fervants.

into one fmall apartment, where there is hardly room for one perfon to breathe freely. In this fituation they generally continue for many hours at a time, often with the addition of feveral candles, which tend likewife to wafte the air, and render it lefs fit for refpiration. Air that is breathed repeatedly becomes unfit for expanding the lungs. This is one caufe of the phthifical coughs, and other complaints of the breaft, fo incident to fedentary artificers.

Even the perfpiration from a great number of perfons pent up together, renders the air unwholefome. The danger from this quarter will be greatly increased, if any one of them happen to have bad lungs, or to be otherwife difeased. Those who fit near him, being forced to breathe the fame air, can hardly fail to be infected. It would be a rare thing, however, to find a dozen of sedentary people all in good health. The danger of crowding them together must therefore be evident to every one.

Many of those who follow fedentary employments are constantly in a bending posture, as shoemakers, tailors, cutlers, &c. Such a situation is extremely hurtful. A bending posture obstructs all the vital motions, and of course must destroy the health. Accordingly we find fuch artificers generally complaining of indigestions, flatulences, head-achs, pains of the breast, &c.

The aliment in fedentary people, inftead of being pufhed forwards by an erect pofture, and the action of the mufcles, is in a manner confined in the bowels. Hence indigeftions, coftivenels, wind, and other hypochondriacal affections, the conftant companions of the fedentary. Indeed none of the excretions can be duly performed where exercife is wanting; and when the matter which ought to be difcharged in this way is retained too long in the body, it must have bad effects, as it is again taken up into the mass of humours.

A bending posture is likewise hurtful to the lungs. When this organ is compressed, the air cannot have free access into all its parts, so as to expand them properly. Hence tubercles, adhesions, &c. are formed, which often

and the STUDIOUS.

often end in confumptions. Befides, the proper action of the lungs being abfolutely neceffary for making good blood, when that organ fails, the humours foon become univerfally depraved, and the whole conflictution goes to wreck. Being of a foft texture, and in continual action, their functions are eafily obftructed by preffure.

The fedentary are not only hurt by preffure on the bowels, but alfo on the inferior extremities, which obftructs the circulation in thefe parts, and renders them weak and feeble. Thus tailors, fhoemakers, &c. frequently lofe the ufe of their legs altogether : befides, the blood and humours are, by ftagnation, vitiated, and the perfpiration is obftructed ; whence proceed the fcab, ulcerous fores, foul blotches, and other cutaneous difeafes, fo common among fedentary artificers.

A bad figure of body is a very common confequence of close application to fedentary employments. The fpine, for example, by being continually bent, puts on a crooked fhape, and generally remains fo ever after. But a bad figure of body has already been obferved to be hurtful to health, as the vital functions are thereby impeded.

A fedentary life feldom fails to occafion an univerfal relaxation of the folids. This is the great fource from whence most of the difeases of fedentary people flow. The fcrophula, confumption, hysterics, and nervous difeases, now so common, were very little known in this country before fedentary artificers became so numerous; and they are very little known still among such of our people as follow active employments without doors, though in great towns at least two-thirds of the inhabitants are afflicted with them.

It is very difficult to remedy those evils, because many who have been accustomed to a fedentary life, like ricketty children, lose all inclination for exercise: we shall, however, throw out a few hints with respect to the most likely means for preferving the health of this useful fet of people, which some of them, we hope, will be wise enough to take.

It has been already observed, that sedentary artificers are

are often hurt by their bending pofture. They ought therefore to ftand or fit as erect as the nature of their employments will permit. They fhould likewife change their pofture frequently, and fhould never fit too long at a time, but leave off work, and walk, ride, run, or do any thing that will promote the vital functions.

Sedentary artificers are generally allowed too little time for exercife; yet, fhort as it is, they feldom employ it properly. A journeyman tailor or weaver, for example, inftead of walking abroad for exercife and frefh air, at his hours of leifure, chufes often to fpend them in a public-houfe, or in playing at fome fedentary game, by which he generally lofes both his time and his money.

The awkward poftures in which many fedentary artificers work, feem rather to be the effect of cuftom than neceffity. For example, a table might furely be contrived for ten or a dozen tailors to fit round, with liberty for their legs either to hang down, or reft upon a footboard, as they fhould chufe. A place might likewife be cut out for each perfon, in fuch a manner that he might fit as conveniently for working as in the prefent mode of fitting crofs-legged.

All fedentary artificers ought to pay the most religious regard to cleanlinefs. Both their fituation and occupations render this highly neceffary. Nothing would contribute more to preferve their health, than a strict attention to it: and such of them as neglect it, not only run the hazard of losing health, but of becoming a nuisance to their neighbours.

Sedentary people ought to avoid food that is windy or hard of digeftion, and fhould pay the ftricteft regard to fobriety. A perfon who works hard without doors will foon throw off a debauch; but one who fits has by no means an equal chance. Hence it often happens, that fedentary people are feized with fevers after hard drinking. When fuch perfons feel their fpirits low, inftead of running to the tavern for relief, they fhould ride or walk in the fields. This would remove the complaint more effectually than ftrong liquor, and would never hurt the conftitution.

Inftead

and the STUDIOUS. 49 -

Inftead of multiplying rules for preferving the health of the fedentary, we fhall recommend to them the following general plan, viz. That every perfon who follows a fedentary employment fhould cultivate a piece of ground with his own hands. This he might dig, plant, fow, and weed at leifure hours, fo as to make it both an exercife and amufement, while it produced many of the neceffaries of life. After working an hour in a garden, a man will return with more keennefs to his employment within doors, than if he had been all the while idle.

Labouring the ground is every way conducive to health. It not only gives exercise to every part of the body, but the very fmell of the earth and fresh herbs revives and cheers the fpirits, whilft the perpetual profpect of fomething coming to maturity, delights and entertains the mind. We are fo formed as to be always pleafed with fomewhat in profpect, however diftant or however trivial; hence the happiness that most men feel in planting, fowing, building, &c. Thefe feem to have been the chief employments of the more early ages; and, when kings and conquerors cultivated the ground, there is reafon to believe that they knew as well wherein true happiness confisted as we do.

It may feem romantic to recommend gardening to manufacturers in great towns; but observation proves that the plan is very practicable. In the town of Sheffield in Yorkshire, where the great iron-manufacture is carried on, there is hardly a journeyman cutler who does not poffefs a piece of ground, which he cultivates as a garden. This practice has many falutary effects. It not only induces these people to take exercise without doors, but alfo to eat many greens, roots, &c. of their own growth, which they would never think of purchasing. There can be no reason why manufacturers in any other town in Great Britain should not follow the fame plan. It is indeed to be regretted, that in fuch a place as London a plan of this kind is not practicable: yet, even there, fedentary artificers may find opportunities of taking air and exercise, if they chufe to embrace them.

Mechanica

Mechanics are too much inclined to crowd into great towns. The fituation may have fome advantages; but it has likewife many difadvantages. All mechanics who live in the country have it in their power to cultivate a piece of ground; which indeed most of them do. This not only gives them exercise, but enables them to live more comfortably. So far at least as my observation extends, mechanics who live in the country are far more happy than those in great towns. They enjoy better health, live in greater affluence, and feldom fail to rear a healthy and numerous offspring.

In a word, exercife without doors, in one fhape or another, is abfolutely neceffary to health. Thofe who neglect it, though they may for a while drag out life, can hardly be faid to enjoy it. Weak and effeminate, they languish for a few years, and soon drop into an untimely grave.

THE STUDIOUS.

mennets that

Intenfe thinking is fo deftructive to health, that few inftances can be produced of fludious perfons who are ftrong and healthy. Hard fludy always implies a fedentary life; and when intenfe thinking is joined to the want of exercife, the confequences must be bad. We have frequently known even a few months of clofe application to fludy, ruin an excellent conflitution, by inducing a train of nervous complaints which could never be removed. Man is evidently not formed for continual thought more than for perpetual action, and would be as foon worn out by the one as by the other.

So great is the power of the mind over the body, that, by its influence, the whole vital motions may be accelerated or retarded, to almost any degree. Thus cheerfulness and mirth quicken the circulation, and promote all the fecretions; whereas fadness and profound thought never fail to retard them. Hence it would appear, that even a degree of thoughtless is necessary to health. Indeed the perpetual thinker feldom enjoys either health or fpirits; while the perfon who can hardly be faid to think at all, generally enjoys both.

Perpetual

and the STUDIOUS.

Perpetual thinkers, as they are called, 'feldom think long. In a few years they generally become quite flupid, and exhibit a melancholy proof how readily the greateft bleffings may be abufed. Thinking, like every thing elfe, when carried to extreme, becomes a vice; nor can any thing afford a greater proof of wildom, than for a man frequently and feafonably to unbend his mind. This may generally be done by mixing in cheerful company, active diversions, or the like.

Inftead of attempting to inveftigate the nature of that connection which fubfifts between the mind and the body. or to inquire into the manner in which they mutually affect each other, we fhall only mention those difeases to which the learned are more peculiarly liable, and endeavour to point out the means of avoiding them.

Studious perfons are very fubject to the gout. This painful difeafe in a great meafure proceeds from indigeftion, and an obftructed perfpiration. It is impoffible that the man who fits from morning till night fhould either digeft his food, or have any of the fecretions in due quantity. But when that matter which fhould be thrown off by the fkin, is retained in the body, and the humours are not duly prepared, difeafes muft enfue.

The fludious are likewife very liable to the flone and gravel. Exercife greatly promotes both the fecretion and difcharge of urine; confequently a fedentary life muft have the contrary effect. Any one may be fatisfied of this by obferving, that he paffes much more urine by day than in the night, and alfo when he walks or rides, than when he fits. The difcharge of urine not only prevents the gravel and flone, but many other difeafes.

The circulation in the liver being flow, obftructions in that organ can hardly fail to be the confequence of inactivity. Hence fedentary people are frequently afflicted with fchirrous livers. But the proper fecretion and difcharge of the bile is fo neceffary a part of the animal ∞ conomy, that where these are not duly performed, the health must foon be impaired. Jaundice, indigestion, loss of appetite, and a wasting of the whole E_2 body,

body, feldom fail to be the confequences of a vitiated ftate of the liver, or obstructions of the bile.

Few difeafes prove more fatal to the fludious than confumptions of the lungs. It has already been obferved, that this organ cannot be duly expanded in those who do not take proper exercise; and where that is the case, obstructions and adhesions will ensue. Not only want of exercise, but the posture in which studious perfons generally fit, is very hurtful to the lungs. Those who read or write much are ready to contract a habit of bending forwards, and often press with their breast upon a table or bench. This posture cannot fail to hurt the lungs.

The functions of the heart may likewife by this means be injured. I remember to have feen a man opened, whofe pericardium adhered to the breaft-bone in fuch a manner as to obftruct the motion of the heart, and occasion his death. The only probable caufe that could be affigned for this fingular fymptom was, that the man, whofe bufinefs was writing, ufed constantly to fit in a bending posture, with his breast prefing upon the edge of a plain table.

No perfon can enjoy health who does not properly digeft his food. But intenfe thinking and inactivity never fail to weaken the powers of digeftion. Hence the humours become crude and vitiated, the folids weak and relaxed, and the whole conftitution goes to ruin.

Long and intenfe thinking often occasions grievous head-achs, which bring on vertigoes, apoplexies, palfies, and other fatal diforders. The best way to prevent these is, never to study too long at one time, and to keep the body regular, either by proper food, or taking frequently a little of fome opening medicine.

Those who read or write much are often afflicted with fore eyes. Studying by candle-light is peculiarly hurtful to the fight. This ought to be practised as feldom as possible. When it is unavoidable, the eyes should be shaded, and the head should not be held too low. When the eyes are weak or painful, they should be bathed every night and morning in cold water, to which a little brandy may be added. It has already been obferved, that the excretions are very defective in the fludious. The dropfy is often occasioned by the retention of those humours which ought to be carried off in this way. Any perfon may observe, that fitting makes his legs swell, and that this goes off by exercise; which clearly points out the method of prevention.

Fevers, efpecially of the nervous kind, are often the effect of ftudy. Nothing affects the nerves fo much as intenfe thought. It in a manner unhinges the whole human frame, and not only hurts the vital motions, but diforders the mind itfelf. Hence a delirium, melancholy, and even madnefs, are often the effect of clofe application to ftudy. In fine, there is no difeafe which can proceed either from a bad ftate of the humours, a defect of the ufual fecretions, or a debility of the nervous fyftem, which may not be induced by intenfe thinking.

But the most afflicting of all the difeases which attack the studious is the hypochondriac. This difease feldom fails to be the companion of deep thought. It may rather be called a complication of maladies than a single one. To what a wretched condition are the best of men often reduced by it! Their strength and appetite fail; a perpetual gloom hangs over their minds; they live in the constant dread of death, and are continually in fearch of relief from medicine, where, alas! it is not to be found. Those who labour under this diforder, though they are often made the subject of ridicule, justify claim our highest sympathy and compassion.

Hardly any thing can be more prepofterous than for a perfon to make fludy his fole bufinels. A mere fludent is feldom an ufeful member of fociety. He often neglects the most important duties of life, in order to purfue fludies of a very trifling nature. Indeed it rarely happens, that any ufeful invention is the effect of mere fludy. The farther men dive into profound refearches, they generally deviate the more from common fenfe, and too often lofe fight of it altogether. Profound fpeculations, inftead of making men wifer or better, generally render them abfolute fceptics, and overwhelm E_3 them

them with doubt and uncertainty. All that is neceffary for man to know, in order to be happy, is eafily obtained; and the reft, like the forbidden fruit, ferves only to increase his misery.

Studious perfons, in order to relieve their minds, muft not only difcontinue to read and write, but engage in fome employment or diversion that will fo far occupy the thought as to make them forget the business of the closet. A folitary ride or walk are fo far from relaxing the mind, that they rather encourage thought. Nothing can divert the mind when it gets into a train of ferious thinking, but attention to subjects of a more trivial nature. These prove a kind of play to the mind, and confequently relieve it.

Learned men often contract a contempt for what they call trifling company. They are afhamed to be feen with any but philofophers. This however is no proof of their being philofophers themfelves. No man deferves that name who is afhamed to unbend his mind, by affociating with the cheerful and gay. Even the fociety of children will relieve the mind, and expel the gloom which application to ftudy is too apt to occasion.

As fludious people are neceffarily much within doors, they fhould make choice of a large and well-aired place for fludy. This would not only prevent the bad effects which attend confined air, but would cheer the fpirits, and have a most happy influence both on the body and mind. It is faid of Euripides the tragedian, that he used to retire to a dark cave to compose his tragedies; and of Demosthenes the Grecian orator, that he chose a place for fludy where nothing could be either heard or feen. With all deference to fuch venerable names, we cannot help condemning their tafte. A man may furely think to as good purpose in an elegant apartment as in a cave; and may have as happy conceptions where the all-cheering rays of the fun render the air wholefome, as in places where they never enter.

Those who read or write much should be very attentive to their posture. They ought to fit and stand by turns, always keeping as nearly in an creft posture as possible.

and the STUDIOUS.

poffible. Those who dictate, may do it walking. It has an excellent effect frequently to read or speak aloud. This not only exercises the lungs, but almost the whole body. Hence studious people are greatly benefited by delivering discours in public. Public speakers, indeed, sometimes hurt themselves, by overacting their part; but this is their own fault. The martyr to mere vociferation merits not our sympathy.

The morning has, by all medical writers, been reckoned the beft time for fludy. It is fo. But it is alfo the most proper feason for exercise, while the stomach is empty, and the spirits refreshed with steps. Studious people should therefore sometimes spend the morning in walking, riding, or some manly diversions without doors. This would make them return to study with greater alacrity, and would be of more fervice than twice the time after their spirits are worn out with fatigue. It is not sufficient to take diversion only when we can think no longer. Every studious person should make it a part of his business, and should let nothing interrupt his hours of recreation more than those of study.

Mufic has a very happy effect in relieving the mind when fatigued with fludy. It would be well if every fludious perfor were fo far acquainted with that fcience as to amufe himfelf after fevere thought by playing fuch airs as have a tendency to raife the fpirits, and infpire cheerfulnefs and good humour.

It is a reproach to learning, that any of her votaries, to relieve the mind after fludy, fhould betake themfelves to the ufe of ftrong liquors. This indeed is a remedy; but it is a defperate one, and always proves deftructive. Would fuch perfons, when their fpirits are low, get on horfeback, and ride ten or a dozen miles, they would find it a more effectual remedy than any cordial medicine in the apothecary's fhop, or all the ftrong liquors in the world.

The following is my plan, and I cannot recommend a better to others. When my mind is fatigued with ftudy, or other ferious bufinefs, I mount my horfe, and

ride

ride ten or twelve miles into the country, where I fpend a day, and fometimes two, with a cheerful friend; after which I never fail to return to town with new vigour, and to purfue my fludies or bufinefs with fresh alacrity.

It is much to be regretted, that learned men, while in health, pay fo little regard to thefe things! There is not any thing more common than to fee a miferable object over-run with nervous difeafes, bathing, walking, riding, and, in a word, doing every thing for health after it is gone; yet, if any one had recommended thefe things to him by way of prevention, the advice would, in all probability, have been treated with contempt, or, at leaft, with neglect. Such is the weaknefs and folly of mankind, and fuch the want of forefight, even in thofe who ought to be wifer than others!

With regard to the diet of the fludious, we fee no reafon why they flould abstain from any kind of food that is wholefome, provided they use it in moderation. They ought, however, to be sparing in the use of every thing that is windy, rancid, or hard of digestion. Their suppers should always be light, or taken soon in the evening. Their drink may be water, fine malt liquor, not too strong, good cider, wine and water, or, if troubled with acidities, water mixed with a little brandy, rum, or any other genuine spirit.

We fhall only obferve, with regard to those kinds of exercise which are most proper for the studious, that they should not be too violent, nor ever carried to the degree of excessive fatigue. They ought likewise to be frequently varied so as to give action to all the different parts of the body; and should, as often as possible, be taken in the open air. In general, riding on horseback, walking, working in a garden, or playing at some active diversions, are the best.

We would likewife recommend the ufe of the cold bath to the ftudious. It will, in fome measure, fupply the place of exercife, and fhould not be neglected by perfons of a relaxed habit, efpecially in the warm feason.

and the STUDIOUS.

No perfon ought either to take violent exercife, or to fludy immediately after a full meal.

In the above remarks on the ufual difeafes of the fludious, my chief object was to warn them of the evil confequences of *painful* and *intenfe thinking*. But I fhould be forry to damp the ardour of their literary purfuits, which are injurious to health only when continued with inceffant toil, at late hours, and without due intervals of reft, refrefhment, relaxation, and exercife. It is not thought, fays the medical poet, 'tis painful thinking, that corrodes our clay. I deemed it neceffary to be more explicit on this head, in confequence of having found that my former cautions to men of genius and fcience had been underftood in too rigorous a fenfe, as difcouraging the manly exertion of real talents.

Though moderatilitis. A A H'D rule with regard to

to inform them when more is necellary.

their food. Mature teaches every creature when it has endoughed and the calls of thirft and hunger are full-clent

the quantity, yet the quality of food merits a farther confideration. T MrH MI J Avy F.O. by which pro-

villons may be rendered unwholefome. Bad fealons

UNWHOLESOME food, and irregularities of diet, occasion many difeases. There is no doubt but the whole constitution of body may be changed by diet alone. The fluids may be thereby attenuated or condensed, rendered mild or acrimonious, coagulated or diluted, to almost any degree. Nor are its effects upon the folids less confiderable. They may be braced or relaxed, have their sensibility, motions, &c. greatly increased or diminissed, by different kinds of aliment. A very small attention to these things will be sufficient to show, how much the prefervation of health depends upon a proper regimen of the diet.

Nor is an attention to diet neceffary for the prefervation of health only: it is likewife of importance in the cure of difeafes. Every intention in the cure of many difeafes, may be anfwered by diet alone. Its effects, indeed, are not always fo quick as those of medicine, but 58

but they are generally more lasting : besides, it is neither so difagreeable to the patient, nor so dangerous as medicine, and is always more easily obtained.

Our intention here is not to inquire minutely into the nature and properties of the various kinds of aliment in ufe among mankind; nor to fhew their effects upon the different conflictutions of the human body; but to mark fome of the most pernicious errors which people are apt to fall into, with respect both to the quantity and quality of their food, and to point out their influence upon health.

It is not indeed an eafy matter to alcertain the exact quantity of food proper for every age, fex, and conflitution: but a fcrupulous nicety here is by no means neceffary. The beft rule is to avoid all extremes. Mankind were never intended to weigh and meafure their food. Nature teaches every creature when it has enough; and the calls of thirft and hunger are fufficient to inform them when more is neceffary.

Though moderation is the chief rule with regard to the quantity, yet the quality of food merits a farther confideration. There are many ways by which provisions may be rendered unwholefome. Bad feasons may either prevent the ripening of grain, or damage it afterwards. These, indeed, are acts of Providence, and we must fubmit to them; but furely no punishment can be too fevere for those who fuffer provisions to spoil by hoarding them, on purpose to raise the price, or who promote their own interest by adulterating the necessaries of life *.

Animal as well as vegetable food may be rendered unwholefome, by being kept too long. All animal fubftances have a conftant tendency to putrefaction; and, when that has proceeded too far, they not only be-

* The poor, indeed, are generally the first who fuffer by unfound provisions; but the lives of the labouring poor are of great importance to the state: besides, difeases occasioned by unwholesome food often prove infectious, by which means they reach people in every station. It is therefore the interest of all to take care that no spoiled provisions of any kind be exposed to fale. come offenfive to the fenfes, but hurtful to health. Difeafed animals, and fuch as die of themfelves, ought never to be eaten. It is a common practice, however, in fome grazing countries, for fervants and poor people to eat fuch animals as die of any difeafe, or are killed by accident. Poverty, indeed, may oblige people to do this; but they had better eat a fmaller quantity of what is found and wholefome: it would both afford a better nourifhment, and be attended with lefs danger.

The injunction given to the Jews, not to eat any creature which died of itfelf, feems to have a ftrict regard to health; and ought to be obferved by Christians as well as Jews. Animals never die themfelves without fome previous difeafe; but how a difeafed animal should be wholefome food, is inconceivable: even those which die by accident must be hurtful, as their blood is mixed with the flesh, and foon turns putrid.

Animals which feed grofsly, as tame ducks, hogs, &c. are neither fo eafily digefted, nor afford fuch wholefome nourifhment as others. No animal can be wholefome which does not take fufficient exercife. Most of our stalled cattle are crammed with gross food, but not allowed exercise nor free air; by which means they indeed grow fat, but their juices, not being properly prepared or affimilated, remain crude, and occasion indigestions, gross humours, and oppression of the spirits, in those who feed upon them.

Animals are often rendered unwholefome by being over-heated. Exceflive heat caufes a fever, exalts the animal falts, and mixes the blood fo intimately with the flefh, that it cannot be feparated. For this reafon, butchers fhould be feverely punifhed who over-drive their cattle. No perfon would chufe to eat the flefh of an animal which had died in a high fever; yet that is the cafe with all over-drove cattle; and the fever is often raifed even to the degree of madnefs.

But this is not the only way by which butchers reader meat unwholefome. The abominable cuftom of filling the cellular membrane of animals with air, in order to make them appear fat, is every day practifed. This not

OF ALIMENT.

not only fpoils the meat, and renders it unfit for keeping, but is fuch a dirty trick, that the very idea of it is fufficient to difgult a perfon of any delicacy at every thing which comes from the fhambles. Who can bear the thought of eating meat which has been blown up with air from the lungs of a dirty fellow, perhaps labouring under the very worft of difeafes?

Butchers have likewife a method of filling the cellular membranes of animals with blood. This makes the meat feem fatter, and likewife weigh more, but is notwithftanding a very pernicious cuftom, as it both renders the meat unwholefome and unfit for keeping. I feldom fee a piece of meat from the fhambles, where the blood is not diffufed through the cellular texture. I fhall not fay that this is always the effect of defign; but I am certain it is not the cafe with animals that are killed for domeftic ufe, and properly blooded. Veal feems to be moft frequently fpoilt in this way. Perhaps that may in fome meafure be owing to the practice of carrying calves from a great diffance to market, by which means their tender flefh is bruifed, and many of their veffels burft.

No people in the world eat fuch quantities of falted animal food as the Englifh, which is one reafon why they are fo generally tainted with the fcurvy, and its numerous train of confequences, indigettion, low fpirits, hypochondriacifm, &c. Animal food was furely defigned for man, and, with a proper mixture of vegetables, it will be found the moft wholefome; but to gorge beef, mutton, pork, fifh, and fowl, twice or thrice a-day, is certainly too much. All who value health ought to be contented with making one meal of flefh in the twenty-four hours, and this ought to confift of one kind only.

The most obstinate fcurvy has often been cured by a vegetable diet; nay, milk alone will frequently do more in that difease than any medicine. Hence it is evident, that if vegetables and milk were more used in diet, we should have less fcurvy, and likewise fewer putrid and inflammatory fevers. Fresh vegetables, indeed, come to

to be daily more used in diet; this laudable practice we hope will continue to gain ground.

Our aliment ought neither to be too moift nor too dry. Moift aliment relaxes the folids, and renders the body feeble. Thus we fee females, who live much on tea and other watery diet, generally become weak and unable to digeft folid food: hence proceed hyfterics, and all their dreadful confequences. On the other hand, food that is too dry, renders the folids in a manner rigid, and the humours vifcid, which difpofes the body to inflammatory fevers, fcurvies, and the like.

Much has been faid on the ill effects of tea in diet. They are, no doubt, numerous; but they proceed rather from the imprudent ufe of it, than from any bad qualities in the tea itfelf. Tea is now the univerfal breakfaft in this part of the world; but the morning is furely the moft improper time of the day for drinking it. Moft delicate perfons, who, by the bye, are the greateft teadrinkers, cannot eat any thing in the morning. If fuch perfons, after fafting ten or twelve hours, drink four or five cups of green tea without eating almoft any bread, it muft hurt them. Good tea, taken in a moderate quantity, not too ftrong, nor too hot, nor drank upon an empty ftomach, will feldom do harm; but if it be bad, which is often the cafe, or fubftituted in the room of folid food, it muft have many ill effects.

The arts of cookery render many things unwholefome, which are not fo in their own nature. By jumbling together a number of different ingredients, in order to make a poignant fauce, or rich foup, the compolition proves almost a poison. All high feasoning, pickles, &c. are only incentives to luxury, and never fail to hurt the stomach. It were well for mankind, if cookery, as an art, were entirely prohibited. Plain roasting or boiling is all that the stomach requires. These alone are sufficient for people in health, and the fick have still less need of a cook.

The liquid part of our aliment likewife claims our attention. Water is not only the basis of most liquors, but also composes a great part of our solid food. Good

water

water must therefore be of the greatest importance in The best water is that which is most pure, and diet. free from any mixture of foreign bodies. Water takes up parts of most bodies with which it comes into contact; by this means it is often impregnated with metals or minerals of a hurtful or poifonous nature. Hence the inhabitants of fome hilly countries have peculiar difeafes, which in all probability proceed from the water. Thus the people who live near the Alps in Switzerland, and the inhabitants of the Peak of Derby in England, have large tumours or wens on their necks. This difeafe is generally imputed to the fnow water ; but there is more reafon to believe it is owing to the minerals in the mountains through which the waters pafs.

When water is impregnated with foreign bodies, it generally appears by its weight, colour, tafte, fmell, heat, or fome other fenfible quality. Our bufinefs, therefore, is to chufe fuch water, for common ufe, as is lighteft, and without any particular colour, tafte, or fmell. In most places of Britain the inhabitants have it in their power to make choice of their water, and few things would contribute more to health than a due attention to this article. But mere indolence often induces people to make ufe of the water that is nearest to them, without confidering its qualities.

Before water is brought into great towns, the ftricteft attention ought to be paid to its qualities, as many difeafes may be occafioned or aggravated by bad water; and when once it has been procured at a great expence, people are unwilling to give it up.

The common methods of rendering water clear by filtration, or foft, by exposing it to the fun and air, &c. are fo generally known that it is unneceffary to spend time in explaining them. We shall only, in general, advise all to avoid waters which stagnate long in small lakes, ponds, or the like, as such waters often become putrid, by the corruption of animal and vegetable bodies with which they abound. Even cattle frequently fuffer by drinking, in dry feasons, water which has ftood ftood long in fmall refervoirs, without being fupplied by fprings, or freshened with showers. All wells ought to be kept clean, and to have a free communication with the air.

As fermented liquors, notwithstanding they have been exclaimed against by many writers, still continue to be the common drink of almost every perfon who can afford them; we shall rather endeavour to affist people in the choice of these liquors, than pretend to condemn what custom has so firmly established. It is not the moderate use of found fermented liquors which hurts mankind: it is excess, and using such as are ill-prepared or vitiated.

Fermented liquors, which are too ftrong, hurt digeftion; and the body is fo far from being ftrengthened by them, that it is weakened and relaxed. Many imagine that hard labour could not be fupported without drinking ftrong liquors; this is a very erroneous notion. Men who never tafte ftrong liquors are not only able to endure more fatigue, but alfo live much longer, than those who use them daily. But, suppose ftrong liquors did enable a man to do more work, they must nevertheless waste the powers of life, and occasion premature old age. They keep up a constant fever, which exhausts the spirits, inflames the blood, and disposes the body to numberless difeases.

But fermented liquors may be too weak as well as too ftrong: when that is the cafe, they must either be drank new, or they become four and dead: when fuch liquors are drank new, the fermentation not being over, they generate air in the bowels, and occasion flatulencies; and, when kept till stale, they turn four on the stomach, and hurt digestion. For this reason all malt-liquors, cider, &c. ought to be of such strength as to keep till they be ripe, and then they should be used. When such liquors are kept too long, though they should not become four, yet they generally contract a hardness which renders them unwholesome.

い山山山

18

山川四

All families, who can, ought to prepare their own liquors. Since preparing and vending of liquors became

OF ALIMENT.

came one of the moft general branches of bufinefs, every method has been tried to adulterate them. The great object both to the makers and venders of liquor is, to render it intoxicating, and give it the appearance of age. But it is well known that this may be done by other ingredients than thofe which ought to be ufed for making it ftrong. It would be imprudent even to name thofe things which are daily made ufe of to render liquors heady. Suffice it to fay, that the practice is very common, and that all the ingredients ufed for this purpofe are of a narcotic or flupefactive quality. But as all opiates are poifonous, it is eafy to fee what muft be the confequence of their general ufe. Though they do not kill fuddenly, yet they hurt the nerves, relax and weaken the ftomach, and fpoil the digeftion.

Were fermented liquors faithfully prepared, kept to a proper age, and ufed in moderation, they would prove real bleffings to mankind. But, while they are ill-prepared, various ways adulterated, and taken to excefs, they must have many pernicious effects.

We would recommend it to families, not only to prepare their own liquors, but likewife their bread. Bread is fo neceffary a part of diet, that too much care cannot be beftowed in order to have it found and wholefome. For this purpole, it is not only neceffary that it be made of good grain, but likewife properly prepared, and kept free from all unwholefome ingredients. This, however, we have reafon to believe is not always the cafe vith bread prepared by thofe who make a trade of vending it. Their object is rather to pleafe the eye, than to confult the health. The beft bread is that which is neither too coarfe nor too fine; well fermented, and made of wheat-flower, or rather of wheat and rye mixed together.

To fpecify the different kinds of aliment, to explain their nature and properties, and to point out their effects in different conflictutions, would far exceed the limits of our defign. Inftead of a detail of this kind, which would not be generally underftood, and of courfe little attended to, we fhall only mention the

OF ALIMENT.

the following eafy rules with respect to the choice of aliment.

Perfons whofe folids are weak and relaxed, ought to avoid all vifcid food, or fuch things as are hard of digeftion. Their diet, however, ought to be nourifhing; and they fhould take fufficient exercife in the open air.

Such as abound with blood fhould be fparing in the use of every thing that is highly nourishing, as fat meat, rich wines, ftrong ale, and fuch like. Their food should confist chiefly of bread and other vegetable fubstances; and their drink ought to be water, whey, or fmall-beer.

Fat people fhould not eat freely of oily nourifhing diet. They ought frequently to use radifh, garlic, fpices, or fuch things as are heating and promote perfpiration and urine. Their drink fhould be water, coffee, tea, or the like; and they ought to take much exercife and little fleep.

Those who are too lean must follow an opposite courfe.

Such as are troubled with acidities, or whole food is apt to four on the ftomach, fhould live much on animal food; and those who are afflicted with hot alkaline eructations, ought to use a diet confisting chiefly of acid vegetables.

People who are affected with the gout, low fpirits, hypochondriac or hyfteric diforders, ought to avoid all flatulent food, every thing that is viscid, or hard of digeftion, all falted or fmoke-dried provisions, and whatever is auftere, acid, or apt to turn four on the ftomach. Their food fhould be light, fpare, cool, and of an opening nature.

The diet ought not only to be fuited to the age and conflitution, but alfo to the manner of life : a fedentary or studious perfon should live more sparingly than one who labours hard without doors. Many kinds of food will nourish a peafant very well, which would be almost indigestible to a citizen; and the latter will live upon a diet on which the former would starve.

65

Diet

Diet ought not to be too uniform. The conftant use of one kind of food might have fome bad effects. Nature teaches us this, by the great variety of aliment which she provided for man, and likewise by giving him an appetite for different kinds of food.

Those who labour under any particular difease, ought to avoid such aliments as have a tendency to increase it: for example, a gouty person should not indulge in rich wines, strong sources, or gravies, and should avoid all acids. One who is troubled with the gravel ought to shun all austere and astringent aliments; and those who are scorbutic should be sparing in the use of salted provisions, &c.

In the first period of life, our food ought to be light, but nourishing, and frequently taken. Food that is folid, with a fufficient degree of tenacity, is most proper for the state of manhood. The diet suited to the last period of life, when nature is upon the decline, approaches nearly to that of the first. It should be lighter and more succulent than that of vigorous age, and likewife more frequently taken.

It is not only neceffary for health that our diet be wholefome, but alfo that it be taken at regular periods. Some imagine long fafting will atone for excefs; but this, inftead of mending the matter, generally makes it worfe. When the ftomach and inteffines are overdiftended with food, they lofe their proper tone, and, by long fafting, they become weak, and inflated with wind. Thus, either gluttony or fafting deftroys the powers of digeftion.

The frequent repetition of aliment is not only neceffary for repairing the continual wafte of our bodies, but likewife to keep the fluids found and fweet. Our humours, even in the most healthy state, have a constant tendency to putrefaction, which can only be prevented by frequent supplies of fresh nourishment: when that is wanting too long, the putrefaction often proceeds so far as to occasion very dangerous fevers. From hence we may learn the necessity of regular meals. No perfon can enjoy a good state of health, whose vessels are either frequently

frequently overcharged, or the humours long deprived of fresh supplies of chyle.

Long failing is extremely hurtful to young people; it not only vitiates their humours, but prevents their growth. Nor is it lefs injurious to the aged. Moft perfons, in the decline of life, are afflicted with wind : this complaint is not only increased, but even rendered dangerous and often fatal, by long failing. Old people, when their ftomachs are empty, are frequently feized with giddines, head-achs, and faintness. These complaints may generally be removed by a piece of bread and a glass of wine, or taking any other folid food; which plainly points out the method of preventing them.

It is more than probable, that many of the fudden deaths, which happen in the advanced periods of life, are occafioned by fafting too long, as it exhaufts the fpirits, and fills the bowels with wind : we would therefore advife people in the decline of life, never to allow their flomachs to be too long empty. Many people take nothing but a few cups of tea and a little bread, from nine o'clock at night till two or three next afternoon. Such may be faid to faft almost three-fourths of their time. This can hardly fail to ruin the appetite, vitiate the humours, and fill the bowels with wind; all which might be prevented by a folid breakfast.

It is a very common practice to eat a light breakfaft and a heavy fupper. This cultom ought to be reverfed. When people fup late, their fupper fhould be very light; but the breakfaft ought always to be folid. If any one eats a light fupper, goes foon to bed, and rifes betimes in the morning, he will be fure to find an appetite for his breakfaft, and he may freely indulge it.

The ftrong and healthy do not indeed fuffer fo much from fafting as the weak and delicate; but they run great hazard from its opposite, viz. repletion. Many difeafes, especially fevers, are the effect of a plethora, or too great fulness of the veffels. Strong people, in high health, have generally a great quantity of blood and other humours. When these are fuddenly increased,

F 2

67

by

by an overcharge of rich and nourifhing diet, the veffels become too much diftended, and obstructions and inflammations enfue. Hence fo many people are feized with inflammatory and eruptive fevers, apoplexies, &c. after a feast or debauch.

All great and fudden changes in diet are dangerous. What the flomach has been long accuftomed to digeft, though lefs wholefome, will agree better with it than food of a more falutary nature to which it has not been ufed. When therefore a change becomes neceflary, it ought always to be made gradually; a fudden transition from a poor and low to a rich and luxurious diet, or the contrary, might fo difturb the functions of the body as to endanger health, or even to occasion death itfelf.

When we recommend regularity in diet, we would not be underflood as condemning every fmall deviation from it. It is next to impoffible for people at all times to avoid fome degree of excefs, and living too much by rule might make even the fmalleft deviation dangerous. It may therefore be prudent to vary a little, fometimes taking more, fometimes lefs, than the ufual quantity of meat and drink, provided always that a due regard be had to moderation.

The details which fome writers have entered into respecting the supposed qualities of every article of food and drink, as well as the proper quantities of each, appear to me just as triffing as the minuteness of the phyfician who inferted in his prefcription how many grains of falt fhould be eaten with an egg. Every man's experience of what he has found to agree or difagree with him, is a much more unerring guide than whimfical calculations of the difference between the mucilage of a carrot and a parfnip, or between the jelly contained in a leg and a shoulder of mutton. But while I point out the folly of extreme folicitude in fuch matters, I am far from advising people to eat and drink, without any choice or restraint, whatever falls in their way. This would be inconfiftent with the rules I have already laid down. Rational enjoyment of the gifts of nature, is the happy

OF ALIMENT.

happy medium between boundlefs indulgence and frivolous or unneceffary felf-denial.

- fiter - candle or the

CHAP. IV.

In great cities to many things sound to companying

OF AIR.

UNWHOLESOME air is a very common caufe of difeafes. Few are aware of the danger arifing from it. People generally pay fome attention to what they eat or drink, but feldom regard what goes into the lungs, though the latter proves often more fuddenly fatal than the former.

Air, as well as water, takes up parts of most bodies with which it comes in contact, and is often fo replenished with those of a noxious quality, as to occasion immediate death. But such violent effects feldom happen, as people are generally on their guard against them. The less perceptible influences of bad air prove more generally hurtful to mankind; we shall therefore endeavour to point out some of these, and to shew whence the danger chiefly arises.

Air may become noxious many ways. Whatever greatly alters its degrees of heat, cold, moifture, &c. renders it unwholefome: for example, that which is too hot diffipates the watery parts of the blood, exalts the bile, and renders the whole humours aduft and thick. Hence proceed bilious and inflammatory fevers, cholera morbus, &c. Very cold air obftructs the perfpiration, conftringes the folids, and condenfes the fluids. It occafions rheumatifms, coughs, and catarrhs, with other difeafes of the throat and breaft. Air that is too moift deftroys the elafticity or fpring of the folids, induces phlegmatic or lax conftitutions, and difpofes the body to agues, or intermitting fevers, dropfies, &c.

Wherever great numbers of people are crowded into one place, if the air has not a free circulation, it foon

F 3

becomes

becomes unwholefome. Hence it is that delicate perfons are fo apt to turn fick or faint in crowded churches, affemblies, or any place where the air is injured by breathing, fires, candles, or the like.

In great cities fo many things tend to contaminate the air, that it is no wonder it proves fo fatal to the inhabitants. The air in cities is not only breathed repeatedly over, but is likewife loaded with fulphur, fmoke, and other exhalations, befides the vapours continually arifing from innumerable putrid fubflances, as dunghills, flaughter-houfes, &c. All poffible care fhould be taken to keep the ftreets of large towns open and wide, that the air may have a free current through them. They ought likewife to be kept very clean. Nothing tends more to pollute and contaminate the air of a city than dirty ftreets.

It is very common in this country to have churchyards in the middle of populous cities. Whether this be the effect of ancient fuperfitition, or owing to the increase of fuch towns, is a matter of no confequence. Whatever gave rife to the custom, it is a bad one. It is habit alone which reconciles us to these things; by means of which the most ridiculous, nay pernicious customs often become facred. Certain it is, that thousands of putrid carcass, fo near the furface of the earth, in a place where the air is confined, cannot fail to taint it; and that fuch air, when breathed into the lungs, must occasion difeases *.

Burying within churches is a practice ftill more deteftable. The air in churches is feldom good, and the effluvia from putrid carcaffes muft render it ftill worfe. Churches are commonly old buildings with arched roofs. They are feldom open above once a week, are never ventilated by fires nor open windows, and rarely kept clean. This occafions that damp, mufty, unwholefome

* In most eastern countries it was customary to bury the dead at fome distance from any town. As this practice obtained among the Jews, the Greeks, and alfo the Romans, it is strange that the western parts of Europe should not have followed their example in a custom fo truly laudable.

fmell

fmell which one feels upon entering a church, and renders it a very unfafe place for the weak and valetudinary. Thefe inconveniencies might, in a great meafure, be obviated, by prohibiting all perfons from burying within churches, by keeping them clean, and permitting a ftream of fresh air to pass frequently through them, by opening opposite doors and windows *.

Wherever air ftagnates long, it becomes unwholefome. Hence the unhappy perfons confined in jails not only contract malignant fevers themfelves, but often communicate them to others. Nor are many of the holes, for we cannot call them houfes, poffeffed by the poor in great towns, much better than jails. Thefe low dirty habitations are the very lurking-places of bad air and contagious difeafes. Such as live in them feldom enjoy good health; and their children commonly die young. In the choice of a houfe, thofe who have it in their power ought always to pay the greateft attention to open free air.

The various methods which luxury has invented to make houfes clofe and warm, contribute not a little to render them unwholefome. No houfe can be wholefome unlefs the air has a free paffage through it. For which reafon, houfes ought daily to be ventilated, by opening oppofite windows, and admitting a current of frefh air into every room. Beds, inftead of being made up as foon as people rife out of them, ought to be turned down, and expofed to the frefh air from the open windows through the day. This would expel any noxious vapour, and could not fail to promote the health of the inhabitants.

In hofpitals, jails, fhips, &c. where that cannot be conveniently done, ventilators fhould be used. The method of expelling foul, and introducing fresh air, by means of ventilators, is a most falutary invention, and is indeed the most useful of all our modern medical improvements. It is capable of universal application, and

^{*} One cannot pass through a large church or cathedral, even in fummer, without feeling quite chilly.

is fraught with numerous advantages, both to those in health and fickness. In all places, where numbers of people are crowded together, ventilation becomes abfolutely neceffary.

Air which stagnates in mines, wells, cellars, &c. is extremely noxious. That kind of air is to be avoided as the most deadly poifon. It often kills almost as quickly as lightning. For this reason, people should be very cautious in opening cellars that have been long shut, or going down into deep wells or pits, especially if they have been kept close covered *.

Many people who have fplendid houfes, chufe to fleep in fmall apartments. This conduct is very imprudent. A bed-chamber ought always to be wellaired; as it is generally occupied in the night only, when all doors and windows are fhut. If a fire be kept in it, the danger from a fmall room becomes ftill greater. Numbers have been ftifled when afleep by a fire in a fmall apartment, which is always hurtful.

Those who are obliged, on account of business, to fpend the day in close towns, ought, if possible, to fleep in the country. Breathing free air in the night will, in fome measure, make up for the want of it through the day. This practice would have a greater effect in preferving the health of citizens than is commonly imagined.

Delicate perfons ought, as much as poffible, to avoid the air of great towns. It is peculiarly hurtful to the afthmatic and confumptive. Such perfons fhould avoid cities as they would the plague. The hypochondriac are likewife much hurt by it. I have often feen perfons fo much afflicted with this malady while in town, that it feemed impoffible for them to live, who, upon being removed to the country, were immediately relieved. The fame obfervation holds with regard to

• We have daily accounts of perfons who lofe their lives by going down into deep wells and other places where the air flagnates: all thefe accidents might be prevented by only letting down a lighted candle before them, and flopping when they perceive it go out; yet this precaution, fimple as it is, is feldom uied.

1

nervous and hyfteric women. Many people, indeed, have it not in their power to change their fituation in queft of better air. All we can fay to fuch perfons is, that they fhould go as often abroad into the open air as they can, that they fhould admit frefh air frequently into their houfes, and take care to keep them very clean.

It was neceffary in former times for fafety, to furround cities, colleges, and even fingle houfes, with high walls. Thefe, by obftructing the free current of air, never fail to render fuch places damp and unwholefome. As fuch walls are now, in moft parts of this country, become ufelefs, they ought to be pulled down, and every method taken to admit a free paffage to the air. Proper attention to AIR and CLEANLINESS would tend more to preferve the health of mankind, than all the prefcriptions of the Faculty.

Surrounding houfes too closely with planting or thick woods, likewife tends to render the air unwholefome. Wood not only obftructs the free current of the air, but fends forth great quantities of moift exhalations, which render it conftantly damp. Wood is very agreeable at a proper diftance from a houfe, but fhould never be planted too near it, efpecially in a flat country. Many of the gentlemen's feats in England are rendered very unwholefome from the great quantity of wood which furrounds them.

Houfes fituated in low marfhy countries, or near large lakes of flagnating water, are likewife unwholefome. Waters which flagnate, not only render the air damp, but load it with putrid exhalations, which produce the most dangerous and fatal difeases. Those who are obliged to inhabit marfhy countries, ought to make choice of the dryest fituations they can find, to live generously, and to pay the strictest regard to cleanlines.

If fresh air be necessary for those in health, it is still more fo for the fick, who often lose their lives for want of it. The notion that fick people must be kept very hot, is fo common, that one can hardly enter the chamber where a patient lies, without being ready to faint,

by

by reafon of the hot fuffocating fmell. How this muft affect the fick, any one may judge. No medicine is fo beneficial to the fick as frefh air. It is the moft reviving of all cordials, if it be administered with prudence. We are not, however, to throw open doors and windows at random upon the fick. Fresh air is to be let into the chamber gradually, and, if possible, by opening the windows of fome other apartment.

The air of a fick perfon's chamber may be greatly freshened, and the patient much revived, by sprinkling the floor, bed, &c. frequently with vinegar, juice of lemon, or any other strong vegetable acid.

In places where numbers of fick are crowded into the fame houfe, or, which is often the cafe, into the fame apartment, the frequent admiffion of frefh air becomes abfolutely neceffary. Infirmaries, hofpitals, &c. are often rendered fo noxious, for want of proper ventilation, that the fick run more hazard from them than from the difeafe. This is particularly the cafe when putrid fevers, dyfenteries, and other infectious difeafes prevail.

Phyficians, furgeons, and others who attend hofpitals, ought, for their own fafety, to take care that they be properly ventilated. Such perfons as are obliged to fpend moft of their time amongft the fick, run great hazard of being themfelves infected when the air is bad. All hofpitals, and places of reception for the fick, ought to have an open fituation, at fome diffance from any great town, and fuch patients as labour under any infectious difeafe ought never to be fuffered to come near the reft *.

It gives me great pleafure to obferve, that good fenfe and humanity have of late years been difplayed in the choice of proper fituations for those buildings, and in due attention to keep them well-aired. But my remarks

* A year feldom paffes that we do not hear of fome hofpital phyfician or furgeon having loft his life by an hofpital fever caught from his patients. For this they have themfelves alone to blame. Their patients are either in an improper fituation, or they are too carelefs with regard to their own conduct.

3

74

on

on other fources of widely-diffused infection have been hitherto productive of little effect. The practice of burying the dead in the centre of populous neighbourhoods is still too generally continued. Churches and church-yards are made the chief places of interment, in direct opposition to reason, and to the example of the most enlightened people of antiquity. The first words of the old Roman inferiptions on tomb-ftones, " Sifte, viator," ftop, traveller, fhew that the dead were buried by the fide of public roads, not in temples, nor in the heart of towns and cities. One of the laws of the late JOSEPH the Second relative to this point will do him immortal honour. After ftrictly prohibiting the interment of dead bodies in any church or chapel, " It is horrid," fays the emperor, " that a place of worthip, a temple " of the Supreme Being, fhould be converted into a " peft-houfe for living creatures! A perfon who, upon " his death-bed, makes it a condition of his will to be " buried in a church or chapel, acts like a madman: " he ought to fet his fellow-creatures a good example, " and not to do all in his power to deftroy their con-" flitutions, by expofing them to the effluvia arifing " from a corple in a ftate of putrefaction."

The admirable fentiment expressed by one of our own countrywomen, who died a few years fince, affords a ftriking contrast with the superflutious folly fo justly fligmatized by the emperor. This extraordinary female, whose mind was superior to the weakness of her fex, and to the prejudices of custom, being fully fensible, as she herfelf expressed it in her last will, " That the " bodies of the dead might be offensive to the living," ordered her body to be burnt, and the asses deposited in an urn in the burying-ground of St. George's, Hanover Square, where the remains of the fentimental YORICK are also interred,

CHAP.

[76]

CHAP. V.

mand served

OF EXERCISE.

MANY people look upon the neceffity man is under of earning his bread by labour, as a curfe. Be this as it may, it is evident, from the ftructure of the body, that exercife is not lefs neceffary than food for the prefervation of health : thofe whom poverty obliges to labour for daily bread, are not only the most healthy, but generally the most happy part of mankind. Industry feldom fails to place them above want, and activity ferves them instead of physic. This is peculiarly the cafe with those who live by the culture of the ground. The great increase of inhabitants in infant colonies, and the longevity of fuch as follow agriculture every where, evidently prove it to be the most healthful as well as the most useful employment.

The love of activity fhews itfelf very early in man. So ftrong is this principle, that a healthy youth cannot be reftrained from exercife, even by the fear of punifhment. Our love of motion is furely a ftrong proof of its utility. Nature implants no difposition in vain. It feems to be a catholic law throughout the whole animal creation, that no creature, without exercise, should enjoy health, or be able to find subsistence. Every creature, except man, takes as much of it as is neceffary. He alone, and such animals as are under his direction, deviate from this original law, and they fuffer accordingly.

Inactivity never fails to induce an univerfal relaxation of the folids, which difpofes the body to innumerable difeafes. When the folids are relaxed, neither the digeftion nor any of the fecretions can be duly performed. In this cafe, the worft confequences muft enfue. How can perfons who loll all day in eafy chairs, and fleep all night on beds of down, fail to be relaxed? Nor do fuch greatly mend the matter, who never ftir abroad but in

in a coach, fedan, or fuch like. These elegant pieces of luxury are become to common, that the inhabitants of great towns feem to be in fome danger of losing the use of their limbs altogether. It is now below any one to walk, who can afford to be carried. How ridiculous would it feem to a perfon unacquainted with modern luxury, to behold the young and healthy fwinging along on the shoulders of their fellow-creatures! or to fee a fat carcafe, over-run with difeases occasioned by inactivity, dragged through the streets by half a dozen horfes *!

Glandular obstructions, now so common, generally proceed from inactivity. These are the most obstinate of maladies. So long as the liver, kidneys, and other glands, duly perform their functions, health is feldom impaired; but when they fail, nothing can restore it. Exercise is almost the only cure we know for glandular obstructions: indeed, it does not always fucceed as a remedy; but there is reason to believe that it would feldom fail to prevent these complaints, were it used in due time. One thing is certain, that, amongst those who take sufficient exercise, glandular difeases are very little known; whereas the indolent and inactive are feldom free from them.

Weak nerves are the conftant companions of inactivity. Nothing but exercise and open air can brace and ftrengthen the nerves, or prevent the endless train of difeases which proceed from a relaxed state of these organs. We feldom hear the active or laborious complain of nervous difeases; these are referved for the sons of ease and affluence. Many have been completely cured of these diforders by being reduced, from a state of opulence, to labour for their daily bread. This plainly

* It is not neceffity, but falhion, which makes the use of carriages to common. There are many people who have not exercise enough to keep their humours wholefome, who yet dare not venture to make a visit to their next neighbours, but in a coach or fedan, left they should be looked down upon. Strange, that men should be such fools as to be laughed out of the use of their limbs, or to throw away their health, in order to gratify a piece of vanity, or to comply with a ridiculous fashion 1

points

points out the fources from whence nervous difeafes flow, and the means by which they may be prevented.

It is abfolutely impoffible to enjoy health where the perfpiration is not duly carried on; but that can never be the cafe where exercise is neglected. When the matter which ought to be thrown off by perfpiration is retained in the body, it vitiates the humours, and occafions the gout, fevers, rheumatism, &c. Exercise alone would prevent many of those difeases which cannot be cured, and would remove others where medicine proves ineffectual.

A late author *, in his excellent treatife on health, fays, that the weak and valetudinary ought to make exercife a part of their religion. We would recommend this, not only to the weak and valetudinary, but to all whofe bufinefs does not oblige them to take fufficient exercife, as fedentary artificers †, fhopkeepers, fludious perfons, &c. Such ought to use exercise as regularly as they take food. This might generally be done without any interruption to bufinefs, or real lofs of time.

No piece of indolence hurts the health more than the modern cuftom of lying a-bed too long in a morning. This is the general practice in great towns. The inhabitants of cities feldom rife before eight or nine o'clock; but the morning is undoubtedly the best time for

* Cheyne.

+ Sedentary occupations ought chiefly to be followed by women. They bear confinement much better than men, and are fitter for every kind of bufinefs which does not require much ftrength. It is ridiculous enough to fee a lufty fellow making pins, needles, or watch-wheels, while many of the laborious parts of hufbandry are carried on by the other fex. The fact is, we want men for laborious employments, while one half of the other fex are rendered ufelefs for want of occupations fuited to their ftrength, &c. Were girls bred to mechanical employments, we fhould not fee fuch numbers of them profitute themfelves for bread, nor find fuch a want of men for the important purpofes of navigation, agriculture, &c. An eminent filk-manufacturer told me, that he found women answer better for that bufinefs than men; and that he had lately taken a great many givls apprentices as filk-weavers. I hope his example will be followed by many others.

exercife;

exercife, while the flomach is empty, and the body refreshed with fleep. Besides, the morning-air braces and flrengthens the nerves, and, in some measure, answers the purpose of a cold bath. Let any one who has been accustomed to lie a-bed till eight or nine o'clock, rife by fix or feven, spend a couple of hours in walking, riding, or any active diversion without doors, and he will find his spirits cheerful and serene through the day, his appetite keen, and his body braced and strengthened. Custom soon renders early rising agreeable, and nothing contributes more to the prefervation of health.

The inactive are continually complaining of pains of the ftomach, flatulencies, indigeftions, &c. Thefe complaints, which pave the way to many others, are not to be removed by medicines. They can only be cured by a vigorous courfe of exercife, to which indeed they feldom fail to yield.

Exercife, if poffible, ought always to be taken in the open air. When that cannot be done, various methods may be contrived for exercifing the body within doors, as the dumb bell, dancing, fencing, &c. It is not neceffary to adhere ftrictly to any particular kind of exercife. The beft way is to take them by turns, and to ufe that longeft which is moft fuitable to the ftrength and conflitution. Thofe kinds of exercife which give action to moft of the bodily organs, are always to be preferred, as walking, running, riding, digging, rubbing furniture, and fuch like.

It is much to be regretted, that active and manly diversions are now so little practifed. Diversions make people take more exercise than they otherwise would do, and are of the greatest fervice to such as are not under the necessity of labouring for their bread. As active diversions lose ground, those of a fedentary kind feem to prevail. Sedentary diversions are of no other use but to confume time. Instead of relieving the mind, they often require more thought than either study or business. Every thing that induces people to fit still, unless it be fome necessary employment, ought to be avoided.

79

9

d

The diversions which afford the best exercise are, hunting, shooting, playing at cricket. hand-ball, golff*, &c. These exercise the limbs, promote perspiration, and the other secretions. They likewise strengthen the lungs, and give firmnels and agility to the whole body.

Such as can, ought to fpend two or three hours a-day on horfeback; thofe who cannot ride, fhould employ the fame time in walking. Exercise fhould never be continued too long. Over-fatigue prevents the benefit of exercise, and, instead of strengthening the body, tends to weaken it.

Every man should lay himself under some fort of necessity to take exercise. Indolence, like other vices, when indulged, gains ground, and at length becomes agreeable. Hence many who are fond of exercise in the early part of life, become quite averse from it afterwards. This is the case of most hypochondriac and gouty people, which renders their diseases in a great measure incurable.

In fome countries laws have been made, obliging every man, of whatever rank, to learn fome mechanical employment. Whether fuch laws were defigned for the prefervation of health, or the encouragement of manufacture, is a queftion of no importance. Certain it is, that if gentlemen were frequently to amufe and exercife themfelves in this way, it might have many good effects. They would at leaft derive as much honour from a few mafterly fpecimens of their own workmanfhip, as from the character of having ruined moft of their companions by gaming or drinking. Befides, men of leifure, by applying themfelves to the mechanical arts, might improve them, to the great benefit of fociety.

Indolence not only occasions difeases, and renders men useles to fociety, but promotes all manner of vice. To say a man is idle, is little better than to call him

* Golff is a diversion very common in North Britain. It is well-calculated for exercifing the body, and may always be taken in fuch moderation as neither to over-heat nor fatigue. It has greatly the preference over cricket, tennis, or any of those games which cannot be played without violence.

OF EXERCISE.

vicious. The mind, if not engaged in fome ufeful purfuit, is conftantly in queft of ideal pleafures, or imprefied with the apprehension of fome imaginary evil. From these fources proceed most of the miseries of mankind. Certainly man was never intended to be idle. Inactivity frustrates the very design of his creation; whereas an active life is the best guardian of virtue, and the greatest prefervative of health.

It is indeed evident, that the love of motion, as well as the love of food, fo obfervable in every living creature from the moment of its birth, are wifely defigned by nature as the means of its prefervation. The indolent man is therefore a rebel to her laws, and will certainly provoke her feverest punishment. In vain does he hope for enjoyment in the lap of floth : its chilling influence poisons the fource of every pleasure, and not only invites difease, but renders it almost incurable.

CHAP. NI.

OF SLEEP AND CLOTHING.

SLEEP, as well as diet, ought to be duly regulated. Too little fleep weakens the nerves, exhaufts the fpirits, and occafions difeafes; and too much renders the mind dull, the body grofs, and difpofes to apoplexies, lethargies, and other complaints of a fimilar nature. A medium ought therefore to be obferved; but this is not eafy to fix. Children require more fleep than grown perfons, the laborious than the idle, and fuch as eat and drink freely, than thofe who live abftemioufly. Befides, the real quantity of fleep cannot be meafured by time; as one perfon will be more refreshed by five or fix hours fleep, than another by eight or ten.

Children may always be allowed to take as much fleep as they pleafe; but for adults, fix or feven hours is certainly fufficient, and no one ought to exceed eight. G Those Those who lie a-bed more than eight hours may flumber, but they can hardly be faid to fleep; fuch generally toss and dream away the fore-part of the night, fink to reft towards morning, and dose till noon. The best way to make fleep found and refreshing, is to rise betimes. The custom of lying a-bed for nine or ten hours, not only makes the fleep less refreshing, but relaxes the folids, and greatly weakens the constitution.

Nature points out night as the proper feafon for fleep. Nothing more certainly deftroys the conflitution than night-watching. It is great pity that a practice fo deftructive to health flould be fo much in fashion. How quickly the want of rest in due feason will blass the most blooming complexion, or ruin the best constitution, is evident from the ghastly countenances of those who, as the phrase is, turn day into night, and night into day.

To make fleep refreshing, the following things are requisite: First, to take sufficient exercise in the open air; to avoid strong tea or coffee; next, to eat a light supper; and lastly, to lie down with a mind as cheerful and server as possible.

It is certain that too much exercife will prevent fleep, as well as too little. We feldom however hear the active and laborious complain of reftlefs nights. It is the indolent and flothful who generally have thefe complaints. Is it any wonder that a bed of down fhould not be refrefhing to a perfon who fits all day in an eafy chair? A great part of the pleafure of life confifts in alternate reft and motion; but they who neglect the latter can never relifh the former. The labourer enjoys more true luxury in plain food and found fleep, than is to be found in fumptuous tables and downy pillows, where exercife is wanting.

That light fuppers caufe found fleep, is true even to a proverb. Many perfons, if they exceed the leaft at that meal, are fure to have uneafy nights; and, if they fall afleep, the load and opprefilon on their ftomach and fpirits occafion frightful dreams, broken and diffurbed repose, the night-mare, &c. Were the fame

fame perfons to go to bed with a light fupper, or fit up till that meal was pretty well digefted, they would enjoy found fleep, and rife refreshed and cheerful. There are indeed fome people who cannot fleep unless they have eat fome folid food at night; but this does not imply the neceffity of a heavy fupper : befides, these are generally perfons who have accustomed themselves to this method, and who do not take a fufficient degree of exercise.

Nothing more certainly diffurbs our repofe than anxiety. When the mind is not at eafe, one feldom enjoys found fleep. This greateft of human bleffings flies the wretched, and vifits the happy, the cheerful, and the gay. This is a fufficient reafon why every man fhould endeavour to be as eafy in his mind as poffible when he goes to reft. Many, by indulging grief and anxious thought, have banifhed found fleep fo long, that they could never afterwards enjoy it.

Sleep, when taken in the fore-part of the night, is generally reckoned most refreshing. Whether this be the effect of habit or not, is hard to fay; but as most people are accustomed to go early to bed when young, it may be prefumed that fleep, at this feason, will prove most refreshing to them ever after. Whether the fore-part of the night be best for fleep or not, furely the fore-part of the day is fittest both for business and amusement. I hardly ever knew an early rifer who did not enjoy a good state of health *.

Early rifing is the natural confequence of going to bed early; and this habit implies fobriety, good order, and an exemption from many fashionable follies extremely prejudicial to health. The man, who accuftoms himself to go to bed at an early hour, can feldom join in the revels of Bacchus, or what are improperly

* Men-of every occupation, and in every fituation of life, have lived to a good old age; nay fome have enjoyed this bleffing whofe plan of living was by no means regular: but it confifts with obfervation, that all very old men have been early rifers. This is the only circumftance attending longevity to which I never knew an exception.

G 2

called

called the *amufements* of the gay world. His reft is not diffurbed by the effects of unfeafonable luxury. He knows, that temperance, moderate exercife, composure of mind, and external tranquillity, are the best opiates. His flumbers are found and refreshing. The waste of spirits on the preceding day is fully repaired. Every muscle, every fibre, every nerve has regained its proper tone. He rifes with cheerfulness and vigour to breathe the morning air, and to enter upon the duties of the day. In short, an attention to this single point of going to bed early, and of rising betimes, will be found to superfede a variety of other precepts, and may be justly called the golden rule for the attainment of health and long life.

base there paralele of Clothing. lot or encound and a

The clothing ought to be fuited to the climate. Cuftom has no doubt a very great influence in this article; but no cuftom can ever change the nature of things fo far, as to render the fame clothing fit for an inhabitant of Nova Zembla and the ifland of Jamaica. It is not indeed neceffary to obferve an exact proportion between the quantity of clothes we wear, and the degree of latitude which we inhabit; but, at the fame time, proper attention ought to be paid to it, as well as to the opennefs of the country, the frequency and violence of florms, &c.

In youth, while the blood is hot and the perfpiration free, it is lefs neceffary to cover the body with a great quantity of clothes; but in the decline of life, when the fkin becomes rigid and the humours more cool, the clothing fhould be increafed. Many difeafes in the latter period of life proceed from a defect of perfpiration : thele may, in fome meafure, be prevented by a fuitable addition to the clothing, or by wearing fuch as are better calculated for promoting the difcharge from the fkin, as clothes made of cotton, flannel, &c.

The clothing ought likewife to be fuited to the feafon of the year. Clothing may be warm enough for fummer,

mer, which is by no means fufficient for winter. The greateft caution, however, is neceffary in making thefe changes. We ought neither to put off our winter clothes too foon, nor to wear our fummer ones too long. In this country, the winter often fets in very early with great rigour, and we have frequently cold weather even after the commencement of the fummer months. It would likewife be prudent not to make the change all at once, but to do it gradually; and indeed the changes of apparel in this climate ought to be very inconfiderable, efpecially among those who have pafled the meridian of life *.

Clothes often become hurtful by their being made fubfervient to the purpofes of pride or vanity. Mankind in all ages feem to have confidered clothes in this view; accordingly their fallion and figure have been continually varying, with very little regard either to health, the climate, or conveniency: a farthingale, for example, may be very neceffary in hot fouthern climates, but furely nothing can be more ridiculous in the cold regions of the north.

Even the human fhape is often attempted to be mended by drefs, and those who know no better believe that mankind would be monsters without its affistance. All attempts of this nature are highly pernicious. The most destructive of them in this country is that of squeezing the stomach and bowels into as narrow a compass as possible, to procure, what is falfely called, a fine shape †. By this practice, the action of the sto-

* That colds kill more than plagues, is an old obfervation; and, with regard to this country, it holds firitily true. Every perfon of differnment, however, will perceive, that most of the colds which prove to destructive to the inhabitants of Britain, are owing to their imprudence in changing clothes. A few warm days in March or April induce them to throw off their winter garments, without confidering that our most penetrating colds generally happen in the tpring.

+ This madnefs feems to have pervaded the minds of mothers in every age and country. Terence, in his Comedy of the Eunuch, ridicules the Roman matrons for attempting to mend the fhape of their daughters. \$6

mach and bowels, the motion of the heart and lungs, and almost all the vital functions, are obstructed. Hence proceed indigestions, fyncopes or fainting fits, coughs, confumptions of the lungs, and other complaints fo common among females.

The feet likewife often fuffer by preffure. How a fmall foot came to be reckoned genteel, I will not pretend to fay; but certain it is, that this notion has made many perfons lame. Almost nine-tenths of mankind are troubled with corns: a difease that is feldom or never occasioned but by strait shoes. Corns are not only very troublesse, but by rendering people unable to walk, they may likewife be confidered as the remote cause of other difeases *.

The fize and figure of the floe ought certainly to be adapted to the foot. In children the feet are as well flhaped as the hands, and the motion of the toes as free and eafy as that of the fingers; yet few perfons in the advanced period of life are able to make any ufe of their toes. They are generally, by narrow floes, fqueezed all of a heap, and often laid over one another in fuch a manner as to be rendered altogether incapable of motion. Nor is the high heel lefs hurtful than the narrow toe. A lady may feem taller for walking on her tiptoes, but fle will never walk well in this manner. It ftrains her joints, difforts her limbs, makes her floop, and utterly deftroys all her eafe and gracefulnefs of motion : it is entirely owing to floes with high heels and

* We often fee perfons, who are rendered quite lame by the nails of their toes having grown into the flefh, and frequently hear of mortifications proceeding from this caufe. All thefe, and many other inconveniencies attending the feet, must be imputed folely to the use of flort and strait shoes.

Though we hear frequently of plafters, falves, ointments, &c. for *eradicating* corns, yet they are never known to produce that effect. The only rational mode of proceeding is to foften the corn a little by immerfion in warm water, and then to cut it carefully, and to renew this operation every week, till the fcarf fkin is reduced to its original or natural thinnefs, after which it must be preferved from the irritating preffure of ftrait fhoes, which had at first occasioned the painful callofity.

narrow

narrow toes, that not one female in ten can be faid to walk well.

In fixing on the clothes, due care fhould be taken to avoid all tight bandages. Garters, buckles, &c. when drawn too tight, not only prevent the free motion and ufe of the parts about which they are bound, but likewife obftruct the circulation of the blood, which prevents the equal nourifhment and growth of thefe parts, and occafions various difeafes. Tight bandages about the neck, as flocks, cravats, necklaces, &c. are extremely dangerous. They obftruct the blood in its courfe from the brain, by which means headachs, vertigoes, apoplexies, and other fatal difeafes, are often occafioned.

The perfection of drefs is to be eafy and clean. Nothing can be more ridiculous, than for any one to make himfelf a flave to fine clothes. Such a one, and many fuch there are, would rather remain as fixed as a flatue from morning till night, than difcompofe a fingle hair or alter the pofition of a pin. Were we to recommend any particular pattern for drefs, it would be that which is worn by the people called Quakers. They are always neat, clean, and often elegant, without any thing fuperfluous. What others lay out upon tawdry laces, ruffles, and ribands, they beftow upon fuperior cleanlinefs. Finery is only the affectation of drefs, and very often covers a great deal of dirt.

We fhall only add, with regard to clothing, that it ought not only to be fuited to the climate, the feafon of the year, and the period of life, but likewife to the temperature and conflitution. Robuft perfons are able to endure either cold or heat better than the delicate; confequently may be lefs attentive to their clothing. But the precife quantity of clothes neceffary for any perfon cannot be determined by reafoning. It is entirely a matter of experience, and every man is the beft judge for himfelf what quantity of clothes is neceffary to keep him warm *.

Since

* The celebrated Boerhaave used to fay, that nobody suffered by cold, fave fools and beggars; the latter not being able to pro-G 4 cure

Since the first publication of the preceding remarks, very important changes have taken place in the drefs of our fair countrywomen, which afford the strongest proofs of their good fense and taste. The strongest proofs of their good fense and taste. The strongest inner distorted, nor is growth checked and the vital functions impeded by a whalebone press. Easy, fase, and graceful motion in a flat-heeled shoe has completely abolished the awkwardness and danger of former attempts to totter about, as it were, upon stills. In a word, a becoming regard to health, simplicity, and elegance, feems now to have more influence over female fashions, than absurdity, caprice, or the defire of concealing any perfonal deformity.

I wish I could pay my own fex the fame compliment which the Ladies have fo well deferved. But an affectation of what is called military fmartnefs, feems to have converted their whole apparel into a fystem of bandages. "The hat is as tight as if it was intended for a helmet, and to defy the fury of a hurricane. Its form also being by no means fuited to the natural fhape of the head, it must be worn for a confiderable time, with very painful and unequal preffure, before it can be made to fit its new block. The neck is bolftered up and fwathed with the most unnatural stiffness. Easy motion without, and free circulation within, are alike obstructed. Blotches and eruptions in the face, head-achs, apoplexies, and fudden deaths, may be often traced to this caufe; and if we view its effects in another light, we shall not be furprifed at any inconfiftency in the language or conduct of people, who take fo much pains to luspend all intercourfe between the head and the heart.

cure clothes, and the former not having fenfe to wear them. Be this as it may, I can with the firiteff truth declare, that in many cafes where the powers of medicine had been tried in vain, I have cured the patient by recommending thick floes, a flannel waiftcoat and drawers, a pair of under-flockings, or a flannel petticoat, to be worn during the cold feafon at leaft Where warmer clothing is wanted, I would recommend the fleecy hofiery to be wornnext the fkin.

OF SLEEP AND CLOTHING.

The close preffure of the other articles of drefs is equally reprehensible Narrow fleeves are a great check upon the mulcular exercise of the arms. The waiftcoat, in its prefent fashionable form, may be very properly termed a strait one; and, no doubt, is in many inflances an indication of fome mental derangement. The wrifts and knees, but more particularly the latter, are braced with ligatures, or tight buttoning; and the legs, which require the utmost freedom of motion, are screwed into leathern cafes, as if to convey an idea that the wearer is fometimes mounted on horfeback. To complete the whole, and in order that the feet may be kept in as tight a prefs as the head, when fhoes are to be worn, the fhape of the foot and the eafy expansion of the toes are never confulted, but fashion regulates the form of the fhoe, fometimes fquare-toed, more frequently pointed, and always fure to produce cramps and corns, the keen, the fenfible announcers of every change of the weather. I have fo long employed ferious argument upon these subjects in vain, that I am now accultomed to view them with pleafantry; and when I meet with fuch figures, difguiled, and rendered truly awkward both in their motions and appearance. I. cannot help thinking with SHAKESPEAR, " that fome of Nature's journeymen had made them, and not made them well; they imitate humanity fo abominably !"

CHAP. VII.

the mereparation of the fracies, the prefer

OF INTEMPERANCE.

A MODERN author* obferves, that temperance and exercise are the two belt physicians in the world. He might have added, that if these were duly regarded, there would be little occasion for any other. Temper-

· Rouffeau.

1.1

OF INTEMPERANCE.

ance may justly be called the parent of health; yet numbers of mankind act as if they thought difeafes and death too flow in their progrefs, and by intemperance and debauch feem as it were to folicit their approach.

The danger of intemperance appears from the very conftruction of the human body. Health depends on that flate of the folids and fluids which fits them for the due performance of the vital functions; and while thefe go regularly on, we are found and well; but whatever diffurbs them neceffarily impairs health. Intemperance never fails to diforder the whole animal œconomy; it hurts the digeftion, relaxes the nerves, renders the different fecretions irregular, vitiates the humours, and occafions numberlefs difeafes.

The analogy between the nourifhment of plants and animals affords a firiking proof of the danger of intemperance. Moifture and manure greatly promote vegetation; yet an over-quantity of either will entirely deftroy it. The beft things become hurtful, nay deftructive, when carried to excefs. Hence we learn, that the higheft degree of human wifdom confifts in regulating our appetites and paffions fo as to avoid all extremes. It is that chiefly which entitles us to the character of rational beings. The flave of appetite will ever be the difgrace of human nature.

The Author of Nature hath endued us with various paffions, for the propagation of the fpecies, the prefervation of the individual, &c. Intemperance is the abufe of thefe paffions; and moderation confifts in the proper regulation of them. Men, not contented with fatisfying the fimple calls of Nature, create artificial wants, and are perpetually in fearch after fomething that may gratify them; but imaginary wants can never be gratified. Nature is content with little; but luxury knows no bounds. Hence the epicure, the drunkard, and the debauchee feldom ftop in their career till their money or their conflitution fails: then indeed they generally fee their error when too late.

OF INTEMPERANCE.

It is impossible to lay down fixed rules with regard to diet, on account of the different constitutions of mankind. The most ignorant perfon, however, certainly knows what is meant by excess: and it is in the power of every man, if he chooses, to avoid it.

The great rule of diet is to ftudy fimplicity. Nature delights in the moft plain and fimple food, and every animal, except man, follows her dictates. Man alone riots at large, and ranfacks the whole creation in queft of luxuries, to his own deftruction. An elegant writer * of the laft age fpeaks thus of intemperance in diet : "For my part, when I behold a fafhionable table " fet out in all its magnificence, I fancy that I fee gouts " and dropfies, fevers and lethargies, with other innu-" merable diftempers, lying in ambufcade among the " difhes."

Nor is intemperance in other things lefs deftructive than in diet. How quickly does the immoderate purfuit of carnal pleafures, or the abufe of intoxicating liquors, ruin the beft conflitution! Indeed thefe vices generally go hand in hand. Hence it is that we fo often behold the votaries of Bacchus and Venus, even before they have arrived at the prime of life, worn out with difeafes, and haftening with fwift pace to an untimely grave. Did men reflect on the painful difeafes and premature deaths which are daily occafioned by intemperance, it would be fufficient to make them fhrink back with horror from the indulgence even of their darling pleafures.

Intemperance does not hurt its votaries alone; the innocent too often feel the direful effects of it. How many wretched orphans are to be feen embracing dunghills, whofe parents, regardlefs of the future, fpent in riot and debauch what might have ferved to bring up their offspring in a decent manner! How often do we behold the miferable mother, with her helplefs infants, pining in want, while the cruel father is indulging his infatiate appetites!

* Addifon.

Families

Families are not only reduced to mifery, but even extirpated, by intemperance. Nothing tends fo much to prevent propagation, and to fhorten the lives of children, as the intemperance of parents. The poor man who labours all day, and at night lies down contented with his humble fare, can boaft a numerous offspring, while his pampered lord, funk in eafe and luxury, often languifhes without an heir to his ample fortunes. Even ftates and empires feel the influence of intemperance, and rife or fall as it prevails.

Inftead of mentioning the different kinds of intemperance, and pointing out their influence upon health, we fhall only, by way of example, make a few obfervations on one particular fpecies of that vice, viz. the abufe of intoxicating liquors.

Every act of intoxication puts nature to the expence of a fever, in order to difcharge the poilonous draught, When this is repeated almost every day, it is easy to forefee the confequences. That conflictution must be ftrong indeed, which is able long to hold out under a daily fever ! but fevers occasioned by drinking do not always go off in a day; they frequently end in an inflammation of the breast, liver, or brain, and produce fatal effects.

Though the drunkard fhould not fall by an acute difeafe, he feldom efcapes thole of a chronic kind. Intoxicating liquors, when ufed to excels, weaken the bowels and fpoil the digeftion; they deftroy the power of the nerves, and occafion paralytic and convulfive diforders; they likewife heat and inflame the blood, deftroy its balfamic quality, render it unfit for circulation, and the nourifhment of the body. Hence obftructions, atrophies, dropfies, and confumptions of the lungs. Thefe are the common ways in which drunkards make their exit. Difeafes of this kind, when brought on by hard drinking, feldom admit of a cure.

Many people injure their health by drinking, who feldom get drunk. The continual habit of foaking, as it is called, though its effects be not fo violent, is not lefs pernicious. When the veffels are kept conflantly full

OF INTEMPERANCE.

full and upon the firetch, the different digeftions can neither be duly performed, nor the humours properly prepared. Hence most people of this character are afflicted with the gout, the gravel, ulcerous fores in the legs, &c. If these diforders do not appear, they are feized with low spirits, hypochondriacal affections, and other symptoms of indigestion.

Confumptions are now fo common, that it is thought one-tenth of the inhabitants of great towns die of that difeafe. Hard drinking is no doubt one of the caufes to which we muft impute the increase of confumptions. The great quantities of viscid malt liquor drank by the common people of England, cannot fail to render the blood fizy and unfit for circulation; from whence proceed obstructions, and inflammations of the lungs. There are few great ale-drinkers who are not phthifical: nor is that to be wondered at, confidering the glutinous and almost indigestible nature of strong ale.

Those who drink ardent spirits or strong wines run fill greater hazard; these liquors heat and inflame the blood, and tear the tender vessels of the lungs to pieces; yet so great is the confumption of them in this country, that one would almost be induced to think that the inhabitants lived upon them *.

The habit of drinking proceeds frequently from miffortunes in life. The miferable fly to it for relief. It affords them indeed a temporary eafe. But, alas! this folace is fhort lived; and when it is over, the fpirits fink as much below their ufual tone as they had before been raifed above it. Hence a repetition of the dofe becomes neceffary, and every fresh dole makes way for

* We may form fome notion of the immense quantity of ardent spirits confumed in Great Britain from this circumstance, that in the city of Edinburgh and its environs, besides the great quantity of foreign spirits duly entered, and the still greater quantity which is supposed to be smuggled, it is computed that above two thousand private stills are constantly employed in preparing a poisonous liquor called *Molasses*. The common people have got so universally into the habit of drinking this base spirit, that when a porter or labourer is seen reeling along the streets, they fay, he has got molasses.

93

another.

another, till the unhappy wretch becomes a flave to the bottle, and at length falls a facrifice to what at first perhaps was taken only as a medicine. No man is fo dejected as the drunkard when his debauch is gone off. Hence it is, that those who have the greatest flow of fpirits while the glafs circulates freely, are of all others the most melancholy when sober, and often put an end to their own miferable existence in a fit of spleen or illhumour.

Drunkennefs not only proves deftructive to health, but likewife to the faculties of the mind. It is ftrange that creatures, who value themfelves on account of a fuperior degree of reafon to that of brutes, fhould take pleafure in finking fo far below them. Were fuch as voluntarily deprive themfelves of the ule of reafon, to continue ever after in that condition, it would feem but a just punishment. Though this be not the confequence of one act of intoxication, it feldom fails to fucceed a courfe of it. By a habit of drinking, the greateft genius is often reduced to a mere idiot *.

Intoxication is peculiarly hurtful to young perfons. It heats their blood, impairs their ftrength, and obstructs their growth; befides, the frequent use of ftrong liquors in the early part of life deftroys any benefit that might arife from them afterwards. Those who make a practice of drinking generous liquors when young, cannot expect to reap any benefit from them as a cordial in the decline of life.

* It is amazing that our improvements in arts, learning, and politenefs, have not put the barbarous cuftom of drinking to excefs out of fashion. It is indeed lefs common in South Britain than it was formerly; but it ftill prevails very much in the North, where this relic of barbarity is miltaken for hofpitality. There no man is fuppofed to entertain his guefts well, who does not make them drunk. Forcing people to drink is certainly the greateft piece of rudeness that any man can be guilty of. Manlinefs, complaifance, or mere good-nature, may induce a man to take his glafs, if urged to it, at a time when he might as well take poifon. The cuftom of drinking to excefs has long been out of fathion in France; and, as it begins to lofe ground among the politer part of the English, we hope it will foon be banished from every part of this ifland.

Drunkennefs

OF INTEMPERANCE.

Drunkennefs is not only in itfelf a most abominable vice, but is an inducement to many others. There is hardly any crime fo horrid that the drunkard will not perpetrate for the love of liquor. We have known mothers fell their children's clothes, the food that they should have eat, and afterwards even the infants themfelves, in order to purchase the accurfed draught.

It is of the utmost importance to check the first propensities to gluttony and intoxication, or they soon become uncontrolable. With respect to eating, the stomach being often put upon the full stretch, feels uneasiness from the least vacuity, and acquires by degrees a fort of unnatural craving, the gratifications of which are sure to be attended with stupor, debility, and difease.

The fame remark is applicable to drinking. After frequent indulgence in excefs, the fmalleft felf-denial caufes a faintnefs and depreffion of fpirits, which nothing can remove but the favourite dram or pretended cordial. Nay more, the repetition of the laft night's debauch is looked upon as the beft remedy for the ficknefs of the enfuing day. Mild diluting liquors are rejected as infipid, and fome hot ftimulant is required for the palate and ftomach, without confidering, that, by fuch means, the action of the heart and arteries is ftimulated alfo; that the lungs are inflamed; and the whole fyftem is relaxed and enfeebled.

CHAP. VIII.

OF CLEANLINESS.

THE want of cleanlines is a fault which admits of no excuse. Where water can be had for nothing, it is furely in the power of every perfon to be clean. The continual discharge from our bodies by perspiration, renders frequent change of apparel necessary. Changing apparel apparel greatly promotes the fecretion from the fkin, fo n ceffary for health. When that matter which ought to be carried off by perfpiration is either retained in the body, or re-abforbed from dirty clothes, it must occafion difeafes.

Difeafes of the fkin are chiefly owing to want of cleanlinefs*. They may indeed be caught by infection, or brought on by poor living, unwholefome food, &c.; but they will feldom continue long where cleanlinefs prevails. To the fame caufe muft we impute the various kinds of vermin which infeft the human body, houfes, &c. Thefe may always be banifhed by cleanlinefs alone, and wherever they abound, we have reafon to believe it is neglected.

One common caufe of putrid and malignant fevers is the want of cleanlinefs. Thefe fevers commonly begin among the inhabitants of clofe dirty houfes, who breathe unwholefome air, take little exercife, and wear dirty clothes. There the infection is generally hatched, which often fpreads far and wide, to the deftruction of many. Hence cleanlinefs may be confidered as an object of public attention. It is not fufficient that I be clean myfelf, while the want of it in my neighbour affects my health as well as his. If dirty people cannot be removed as a common nuifance, they ought at leaft to be avoided as infectious. All who regard their health fhould keep at a diffance even from their habitations.

In places where great numbers of people are collected, cleanlinefs becomes of the utmost importance. It is well known that infectious difeafes are communicated by tainted air. Every thing, therefore, which tends to

* Mr. Pott, in his furgical obfervations, mentions a difeafe which he calls the chimney fweepers's cancer, as it is almost peculiar to that unhappy fet of people. This he attributes to neglect of cleanlinefs, and with great juffice. I am convinced, that if that part of the body which is the feat of this cruel difeafe were kept clean by frequent washing, it would never happen. The climbing boys, as they are called, are certainly the most miferable wretches on the face of the earth; yet, for cleaning chimnies, no fuch perfons are neceffary.

9

OF CLEANLINESS.

pollute the air, or fpread the infection, ought with the utmost care to be guarded against. For this reason, in great towns, no filth, of any kind, should be permitted to lie upon the streets. Nothing is more apt to convey infection than the excrements of the diseafed.

In many great towns the ftreets are little better than dunghills, being frequently covered with afhes, dung, and naftinefs of every kind. Even flaughter-houfes, or killing fhambles, are often to be feen in the very centre of great towns. The putrid blood, excrements, &c. with which thefe places are generally covered, cannot fail to taint the air, and render it unwholefome. How eafily might this be prevented by active magiftrates, who have it always in their power to make proper laws relative to things of this nature, and to enforce the obfervance of them !

We are forry to fay, that the importance of general cleanlinefs does not feem to be fufficiently underftood by the magiftrates of most great towns in Britain; though health, pleafure, and delicacy, all confpire to recommend an attention to it. Nothing can be more agreeable to the fenfes, more to the honour of the inhabitants, or more conducive to their health, than a clean town; nor can any thing impress a stranger with a more difrespectful idea of any people than its opposite. Whatever pretenfions people may make to learning, politeness, or civilization, we will venture to affirm, that while they neglect cleanlinefs, they are in a state of barbarity *.

The peafants in most countries feem to hold cleanlianels in a fort of contempt. Were it not for the open

How truly great does the emperor Trajan appear, when giving directions to Pliny, his proconful, concerning the making of a common fewer for the health and convenience of a conquered city !

[•] In ancient Rome the greatest men did not think cleanliness an object unworthy of their attention. Pliny fays, the *Cloace*, or common fewers for the conveyance of filth and nastiness from the city, were the greatest of all the public works; and bestows higher encomiums upon Tarquinius, Agrippa, and others who made and improved them, than on those who atchieved the greatest conquests.

As many articles of diet come through the hands of peafants, every method fhould be taken to encourage and promote habits of cleanlinefs among them. This, for example, might be done by giving a fmall premium to the perfon who brings the cleaneft and beft article of any kind to market, as butter, cheefe, &c., and by punifhing feverely thofe who bring it dirty. The fame method fhould be taken with butchers, bakers, brewers, and all who are employed in preparing the neceffaries of life.

In camps the ftricteft regard fhould be paid to cleanlinefs. By negligence in this matter, infectious difeafes are often fpread amongft a whole army; and frequently more die of thefe than by the fword. The Jews, during their encampments in the wildernefs, received particular inftructions with refpect to cleanlinefs*. The rules enjoined them ought to be obferved by all in the like fituation. Indeed the whole fyftem of laws delivered to that people has a manifeft tendency to promote cleanlinefs. Whoever confiders the nature of their climate, the difeafes to which they were liable, and their dirty difpofition, will fee the propriety of fuch laws.

It is remarkable that, in most eastern countries, cleanlines makes a great part of their religion. The Mahometan as well as the Jewish religion enjoins various

* Thou fhalt have a place also without the camp, whither thou fhalt go forth abroad; and thou fhalt have a paddle upon thy weapon: and it fhall be when thou fhalt ease thyself abroad, thou fhalt dig therewith, and shall turn back, and cover that which cometh from thee, &c. Deuter. chap. xxii. ver. 12, 13. bathings,

bathings, washings, and purifications. No doubt these might be defigned to reprefent inward purity; but they were at the fame time calculated for the prefervation of health. However whimfical thefe washings may appear to fome, few things would tend more to prevent difeafes than a proper attention to many of them. Were every perfon, for example, after vifiting the fick, handling a dead body, or touching any thing that might convey infection, to wash before he went into company, or fat down to meat, he would run lefs hazard either of catching the infection himfelf, or of communicating it to others.

Frequent washing not only removes the filth and fordes which adhere to the fkin, but likewife promotes the perfpiration, braces the body, and enlivens the fpirits. How refreshed, how cheerful, and agreeable does one feel on being fhaved, washed, and shifted; especially when these offices have been neglected longer than ufual!

The eaftern cuftom of washing the feet, though lefs neceffary in this country, is neverthelefs a very agreeable piece of cleanlinefs, and contributes greatly to the prefervation of health. The fweat and dirt with which these parts are frequently covered, cannot fail to obstruct the perspiration. This piece of cleanliness would often prevent colds and fevers. Were people careful to bathe their feet and legs in lukewarm water at night, after being exposed to cold or wet through the day, they would feldom experience the ill effects which often proceed from these causes.

A proper attention to cleanlinefs is no where more neceffary than on fhipboard. If epidemical diftempers break out there, no one can be fafe. The best way to prevent them, is to take care that the whole company be cleanly in their clothes, bedding, &c. When infectious difeafes do break out, cleanliness is the most likely means to prevent their fpreading : it is likewife neceffary to prevent their returning afterwards, or being conveyed to other places. For this purpofe, the clothes, bedding, &c. of the fick ought to be carefully washed, and fumigated with brimstone. Infection will todge a long time in in dirty clothes, and afterwards break out in the molt terrible manner.

In places where great numbers of fick people are collected together, as gaols, hofpitals, &c. cleanlinefs ought to be moft religioufly obferved. The very fmell in fuch places is often fufficient to make one fick. It is eafy to imagine what effect that is likely to have upon the difeafed. In an hofpital or infirmary where cleanlinefs is neglected, a perfon in perfect health has a greater chance to become fick, than a fick perfon has to get well.

Few things are more unaccountable than that neglect, or rather dread of cleanlinefs, which appears among those who have the care of the fick; they think it almost criminal to fuffer any thing that is clean to come near a perfon in a fever, for example, and would rather allow him to wallow in all manner of filth, than change the leaft bit of his linen. If cleanlinefs be neceffary for perfons in health, it is certainly more fo for the fick. Many difeafes may be cured by cleanlinefs alone; most of them might be mitigated by it; and, where it is neglected, the flighteft diforders are often changed into the most malignant. The fame mistaken care which prompted people to prevent the least admission of fresh air to the fick, feems to have induced them to keep them dirty. Both these destructive prejudices will, we hope, be foon entirely eradicated.

Cleanlinefs is certainly agreeable to our nature. We cannot help approving it in others, even though we fhould not practife it ourfelves. It fooner attracts our regard than even finery itfelf, and often gains efteem where that fails. It is an ornament to the higheft as well as the loweft flation, and cannot be difpenfed with in either. Few virtues are of more importance to fociety than general cleanlinefs. It ought to be carefully cultivated every where; but in populous cities it fhould be almoft revered *.

I have

As it is impoffible to be thoroughly clean without a fufficient quantity of water, we would earneftly recommend it to the magiftrates of great towns to be particularly attentive to this article. Moit

OF CLEANLINESS.

I have often heard with concern the poor complain of the want of many things beyond their reach, while they difregarded other objects of the first importance which were in their own power; namely, pure, open air, and the comforts of cleanlinefs. I do not know whether a late author had this object in view, when he called the lower orders " the fwinish multitude;" but though I fhould be forry to apply to them fo reproachful an epithet, I must fay that uncleanliness has been very properly denominated " the worft affliction of in-" dolence and poverty." There is no excuse for dirt. Every body may be clean, even in rags, or in the meaneft abode; and the poor would find fuch decency not only the best prefervative of health, but the strongest recommendation to employment and to pity. Can any favourable opinion be entertained of the industry or activity of a floven or a flattern; and will not the dread of infection often drive charity herfelf from the flinking hut, and from the noxious atmosphere of a filthy object?

CHAP. IX.

OF INFECTION.

MANY difeafes are infectious. Every perfon ought therefore, as far as he can, to avoid all communication with the difeafed. The common practice of vifit-

Most great towns in Britain are so fituated as to be easily supplied with water; and those perfons who will not make a proper use of it after it is brought to their hand, certainly deserve to be feverely punished. The streets of great towns, where water can be had, ought to be washed every day. This is the only effectual method for keeping them thoroughly clean; and, upon trial, we are perfuaded it will be found the cheapest.

Some of the most dreadful difeafes incident to human nature might, in my opinion, be entirely eradicated by cleanlinefs.

OF INFECTION.

ing the fick, though often well meant, has many ill confequences. Far be it from me to difcourage any act of charity or benevolence, especially towards those in diffres; but I cannot help blaming such as endanger their own or their neighbours' lives, by a mistaken friendship, or an impertinent curiofity.

The houfes of the fick, especially in the country, are generally crowded from morning till night with idle visitors. It is customary, in such places, for servants and young people to wait upon the fick by turns, and even to fit up with them all night. It would be a miracle indeed should such always escape. Experience teaches us the danger of this conduct. People often catch fevers in this way, and communicate them to others, till at length they become epidemic.

It would be thought highly improper for one who had not had the fmall-pox, to wait upon a patient in that difeafe; yet many other fevers are almost as infectious as the fmall-pox, and not lefs fatal. Some imagine that fevers prove more fatal in villages than in great towns, for want of proper medical affistance. This may fometimes be the cafe; but I am inclined to think it oftener proceeds from the cause above mentioned.

Were a plan to be laid down for communicating infection, it could not be done more effectually than by the common method of vifiting the fick. Such vifitors not only endanger themfelves and their connexions, but likewife hurt the fick. By crowding the houfe, they render the air unwholefome, and by their private whifpers and difmal countenances, difturb the imagination of the patient, and deprefs his fpirits. Perfons who are ill, efpecially in fevers, ought to be kept as quiet as poffible. The fight of ftrange faces, and every thing that difturbs the mind, hurts them.

The common practice in country-places, of inviting great numbers of people to funerals, and crowding them into the fame apartment where the corpfe lies, is another way of fpreading infection. The infection does not always die with the patient. Every thing that comes into contact with his body while alive, receives the contagion, tagion, and fome of them, as clothes, blankets, &c. will retain it for a long time. Perfons who die of infectious diforders ought not to lie long unburied; and people should keep as much as possible at a distance from them.

It would tend greatly to prevent the fpreading of infectious difeafes, if those in health were kept at a proper diftance from the fick. The Jewish Legislator, among many other wife inftitutions for preferving health, has been peculiarly attentive to the means of preventing infection, or defilement, as it is called, either from a difeafed perfon or a dead body. In many cafes the difeafed were to be feparated from those in health; and it was deemed a crime even to approach their habitations. If a perfon only touched a difeafed or dead body, he was appointed to wash himself in water, and to keep for fome time at a diftance from fociety.

Infectious difeafes are often communicated by clothes. It is extremely dangerous to wear apparel which has been worn by a perfon who died of an infectious difeafe, unless it has been well washed and fumigated, as infection may lodge a long time in it, and afterwards produce very tragical effects. This fhews the danger of buying at random the clothes which have been worn by other people.

Infectious diforders are frequently imported. Commerce, together with the riches of foreign climes, bring us alfo their difeafes. Thefe do often more than counterbalance all the advantages of that trade by means of which they are introduced. It is to be regretted, that fo little care is commonly beftowed, either to prevent the introduction or fpreading of infectious maladies. Some attention indeed is generally paid to the plague; but other difeafes pafs unregarded *.

Infection

to

. Were the tenth part of the care taken to prevent the importation of difeafe, that there is to prevent fmuggling, it would be attended with many happy confequences. This might eafily be done by appointing a phyfician at every confiderable fea.port, to infpect the thip's company, paffengers, &c. before they came afhore, and, if any fever or other infectious diforder prevailed, to order the thip to perform a fhort quarantine, and to fend the fick H4

OF INFECTION.

Infection is often fpread through cities, by jails, hofpitals, &c. Thefe are frequently fituated in the very middle of populous towns; and when infectious difeafes break out in them, it is impoffible for the inhabitants to efcape. Did magisfrates pay any regard to the health of the people, this evil might be eafily remedied.

Many are the caufes which tend to diffufe infection through populous cities. The whole atmosphere of a large town is one contaminated mass, abounding with various kinds of infection, and must be pernicious to health. The best advice that we can give to such as are obliged to live in large cities, is to chuse an open fituation; to avoid narrow, dirty, crowded streets; to keep their own house and offices clean; and to be as much abroad in the open air as their time will permit.

It would tend greatly to prevent the fpreading of infectious difeafes, were proper nurfes every where employed to take care of the fick. This might often fave a family, or even a whole town, from being infected by one perfon. We do not mean that people fhould abandon their friends or relations in diffrefs, but only to put them on their guard againft being too much in company with thofe who are afflicted with difeafes of an infectious nature.

Such as wait upon the fick in infectious difeafes, run very great hazard. They fhould fluff their nofes with tobacco, or fome other flrong-fmelling herb, as rue, tanfy, or the like. They ought likewife to keep the patient very clean, to fprinkle the room where he lies with vinegar or other flrong acids, frequently to admit a flream of fresh air into it, and to avoid the fmell of his breath as much as they can. They ought never to

to fome hospital or proper place to be cured. He might likewife order all the clothes, bedding, &c. which had been used by the fick during the voyage, to be either destroyed, or thoroughly cleansed by fumigation, &c. before any of them were sent ashore. A scheme of this kind, if properly conducted, would prevent many fevers, and other insectious diseases, from being brought by fullors into sea-port towns, and by this means diffused all over the country.

h 1.

go.

go into company without having changed their clothes and washed their hands; otherwise, if the disease be infectious, they will in all probability carry the contagion along with them *.

However trifling it may appear to inconfiderate perfons, we will venture to affirm, that a due attention to those things which tend to diffuse infection would be of great importance in preventing difeases. As most difeases are in some degree infectious, no one should continue long with the fick, except the necessary attendants. I mean not, however, by this caution, to deter those whose duty or office leads them to wait upon the fick, from such a laudable and necessary employment.

Many things are in the power of the magistrate which would tend to prevent the spreading of infection; as the promoting of public cleanlines; removing jails, hospitals, burying grounds, and other places where infection may be generated, at a proper distance from great towns[†]; widening the streets; pulling down useless walls, and taking all methods to promote a free circulation of air through every part of the town, &c. Public hospitals, or proper places of reception for the sick, provided they were kept clean, well ventilated, and placed in an open situation, would likewise tend to prevent the spreading of infection. Such places of reception would prevent the poor, when sick, from being visited by their idle or

* There is reafon to believe that infection is often conveyed from one place to another by the careleffnefs of the faculty themfelves. Many phyficians affect a familiar way of fitting upon the patient's bedfide, and holding his arm for a confiderable time. If the patient has the fmall-pox, or any other infectious difeafe, there is no doubt but the doctor's hands, clothes, &c. will carry away fome of the infection; and if he goes directly to vifit another patient without wafning his hands, changing his clothes, or being exposed to the open air, which is not feldom the cafe, is it any wonder that he should carry the difease along with him? Phyficians not only endanger others, but also themselves, by this practice. And indeed they sometimes fuffer for their want of care.

+ The antients would not fuffer even the temples of their gods, where the fick reforted, to be built within the walls of a city.

9

officious

OF INFECTION.

officious neighbours. They would likewife render it unneceffary for fick fervants to be kept in their mafters' houfes. Mafters had better pay for having their fervants taken care of in an hofpital, than run the hazard of having an infectious difeafe diffufed among a numerous family. Sick fervants and poor people, when placed in hofpitals, are not only lefs apt to diffufe infection among their neighbours, but have likewife the advantage of being well attended.

We are not, however, to learn that hofpitals, inftead of preventing infection, may become the means of diffufing it. When they are placed in the middle of great towns; when numbers of patients are crowded together in fmall apartments; when there is a conftant communication kept up between the citizens and the patients; and when cleanlinefs and ventilation are neglected, they become nefts for hatching difeafes, and every one who goes into them not only runs a rifk of receiving infection himfelf, but likewife of communicating it to others. This however is not the fault of the hofpitals, but of those who have the management of them. It were to be wifhed, that they were both more numerous, and upon a more respectable footing, as that would induce people to go into them with lefs reluctance. This is the more to be defired, because most of the putrid fevers and other infectious diforders break out among the poor, and are by them communicated to the better fort. Were proper attention paid to the first appearances of fuch diforders, and the patients early conveyed to an hospital, we fhould feldom fee a putrid fever, which is almost as infectious as the plague, become epidemic.

In a converfation with the late Sir JOHN PRINGLE, for whom I had a great regard, he expressed fome apprehension that the contents of this chapter might deter people from attending their friends and relations in fevers. I told Sir JOHN, that was the very evil I meant to cure, having always found the country-people too apt to visit their friends and neighbours in fevers, even so as to crowd the house, and incommode the fick. Nor could I impute this to humanity, but to an inquisitive dispofition

OF INFECTION.

fition to learn what was likely to be the patient's fate, and to afk improper queftions of those about him, or of the doctor himself, were he weak enough to answer them. In this case, his answer would be fure to come back to the patient, and, if unfavourable, greatly magnified; fo ftrong is the inclination which some men feel, to spread terror, even at the risk of another's life.

Sir JOHN, not having practifed in the country, was not immediately ftruck with the force of my reafoning, till I told him what had happened to a family of his own name who lived near Edinburgh, and had nearly all perifhed in a fever. The family confifted of a father and mother, with nine or ten children, moft of them grown up, and in place. The mother was feized with a fever of the putrid kind, and the children came in turns to nurfe her. They all caught the fever : fome died ; and others narrowly efcaped with their lives. The evil did not end there. They carried the fever into the families where they lived, and spread the infection far and wide. This I have often known to happen in the country, and would advise masters and mistreffes never to fuffer their fervants to act as nurses or attendants on the fick, even though the latter fhould be their nearest relations. They had much better hire nurfes, than allow their fervants to act in that dangerous capacity.

CHAP. X.

OF THE PASSIONS.

THE paffions have great influence both in the caufe and cure of difeafes. How the mind affects the body, will, in all probability, ever remain a fecret. It is fufficient for us to know, that there is effablished a reciprocal influence between the mental and corporeal parts; and that whatever injures the one, diforders the other.

Of

Of Anger.

The paffion of *anger* ruffles the mind, difforts the countenance, hurries on the circulation of the blood, and diforders the whole vital and animal functions. It often occafions fevers, and other acute difeafes; and fometimes even fudden death. This paffion is peculiarly hurtful to the delicate, and those of weak nerves. I have known such perfors frequently lose their lives by a violent fit of anger, and would advise them to guard against the excels of this paffion with the utmost care.

It is not indeed always in our power to prevent being angry; but we may furely avoid harbouring refentment in our breaft. Refentment preys upon the mind, and occafions the most obstinate chronical diforders, which gradually waste the constitution. Nothing shews true greatness of mind more than to forgive injuries; it promotes the peace of fociety, and greatly conduces to our own ease, health, and felicity.

Such as value health fhould avoid violent gufts of anger, as they would the most deadly poifon. Neither ought they to indulge refentment, but to endeavour at all times to keep their minds calm and ferene. Nothing tends fo much to the health of the body as a constant tranquillity of mind,

Of Fear.

The influence of *fear*, both in occasioning and aggravating difeases, is very great. No man ought to be blamed for a decent concern about life; but too great a defire to preferve it, is often the cause of losing it. Fear and anxiety, by depressing the spirits, not only dispose us to difeases, but often render those diseases fatal which an undaunted mind would overcome.

Sudden fear has generally violent effects. Epileptic fits, and other convulfive diforders, are often occafioned by it. Hence the danger of that practice, fo common among young people, of frightening one another. Many have have loft their lives, and others have been rendered miferable, by frolics of this kind. It is dangerous to tamper with the human paffions. The mind may eafily be thrown into fuch diforder, as never again to act with regularity.

But the gradual effects of fear prove most hurtful. The constant dread of fome future evil, by dwelling upon the mind, often occasions the very evil itself. Hence it comes to pass, that fo many die of those very difeases of which they long had a dread, or which had been impressed on their minds by some accident, or foolish prediction. This, for example, is often the case with women in child-bed. Many of those who die in that fituation, are impressed with the notion of their death, a long time before it happens; and there is reason to believe that this impression is often the cause of it.

The methods taken to imprefs the minds of women with the apprehensions of the great *pain* and *peril* of child-birth, are very hurtful. Few women die in labour, though many lofe their lives after it; which may be thus accounted for. A woman after delivery, finding herfelf weak and exhausted, immediately apprehends she is in danger; but this fear feldom fails to obstruct the neceffary evacuations, upon which her recovery depends. Thus the fex often fall a facrifice to their own imaginations, when there would be no danger, did they apprehend none.

It feldom happens, that two or three women in a great town die in child-bed, but their death is followed by many others. Every woman of their acquaintance who is with child dreads the fame fate, and the difeafe becomes epidemical, by the mere force of imagination. This fhould induce pregnant women to defpife fear, and by all means to avoid those tattling goffips who are continually buzzing in their ears the misfortunes of others. Every thing that may in the least alarm a pregnant or child-bed woman, ought with the greatest care to be guarded against.

Many women have loft their lives in child-bed by the old fuperfitious cuftom, still kept up in most parts of Britain,

OF THE PASSIONS.

Britain, of tolling the parifh bell for every perfon who dies. People who think themfelves in danger, are very inquifitive; and if they come to know that the bell tolls for one who died in the fame fituation with themfelves, what must be the confequence? At any rate they are apt to fuppofe that this is the cafe, and it will often be found a very difficult matter to perfuade them of the contrary.

But this cuftom is not pernicious to child-bed women only. It is hurtful in many other cafes. When low fevers, in which it is difficult to fupport the patient's fpirits, prevail, what muft be the effect of a funeral peal founding five or fix times a-day in his ears? No doubt his imagination will fuggeft that others died of the fame difeafe under which he labours. This apprehenfion will have a greater tendency to deprefs his fpirits, than all the cordials of which medicine can boaft will have to raife them. The only town which has abolifhed this cuftom is Bath.

If this ufelefs piece of ceremony cannot be abolifhed, we ought to keep the fick as much from hearing it as poffible, and from every other thing that may tend to alarm them. So far, however, is this from being generally attended to, that many make it their bufinefs to vifit the fick, on purpofe to whifper difmal flories in their ears. Such may pafs for fympathizing friends, but they ought rather to be confidered as enemies: All who wifh well to the fick, ought to keep fuch perfons at the greateft diftance from them.

A cultom has long prevailed among phyficians, of prognofticating, as they call it, the patient's fate, or foretelling the iffue of the difeafe. Vanity, no doubt, introduced this practice, and ftill fupports it, in fpite of common fenfe, and the fafety of mankind. I have known a phyfician barbarous enough to boaft, that he pronounced more *fentences* than all his Majefty's judges. Would to God that fuch fentences were not often equally fatal ! It may indeed be alleged, that the doctor does not declare his opinion before the patient. So much the worfe. A fenfible patient had better hear what the doctor fays, than learn it from the difconfolate looks, the watery eyes, and the broken whilpers, of those about him.

him. It feldom happens, when the doctor gives an unfavourable opinion, that it can be concealed from the patient. The very embarraffment which the friends and attendants fhew in difguifing what he has faid, is generally fufficient to difcover the truth.

Kind heaven has, for the wifeft ends, concealed from mortals their fate; and we do not fee what right any man has to announce the death of another, efpecially if fuch a declaration has a chance to kill him. Mankind are indeed very fond of prying into future events, and feldom fail to folicit the phyfician for his opinion. A doubtful anfwer, however, or one that may tend rather to encourage the hopes of the fick, is furely the moft proper. This conduct could neither hurt the patient nor the phyfician. Nothing tends more to deftroy the credit of phyfic, than those bold prognofticators, who, by-the-bye, are generally the moft ignorant of the faculty. The mistakes which daily happen in this way are fo many standing proofs of human vanity, and the weakness of fcience.

We readily admit, that there are cafes where the phyfician ought to give intimation of the patient's danger to fome of his near connexions; though even this ought always to be done with the greateft caution: but it never can be neceffary in any cafe that the whole town and country fhould know, immediately after the doctor has made his first visit, that he has no hopes of his patient's recovery. Perfons whose impertinent curiosity leads them to question the physician with regard to the fate of his patient, certainly deferve no other than an evalue answer.

The vanity of foretelling the fate of the fick is not peculiar to the faculty. Others follow their example, and those who think themselves wifer than their neighbours often do much hurt in this way. Humanity furely calls upon every one to comfort the fick, and not add to their affliction by alarming their fears. A friend, or even a physician, may often do more good by a mild and fympathizing behaviour than by medicine, and should never neglect to administer that greatest of all cordials, HOPE.

Of

Of Grief.

Grief is the most destructive of all the passions. Its effects are permanent; and when it finks deep into the mind, it generally proves fatal. Anger and fear, being of a more violent nature, feldom last long; but grief often changes into a fixed melancholy, which preys upon the spirits, and wastes the constitution. This passion ought not to be indulged. It may generally be conquered at the beginning; but when it has gained strength, all attempts to remove it are vain.

No perfon can prevent misfortunes in life; but it fhews true greatnefs of mind to bear them with ferenity. Many perfons make a merit of indulging grief, and when misfortunes happen, they obftinately refufe all confolation, till the mind, overwhelmed with melancholy, finks under the load. Such conduct is not only deftructive to health, but inconfiftent with reafon, religion, and common fenfe.

Change of ideas is as neceffary for health as change of pofture. When the mind dwells long upon one fubject, efpecially of a difagreeable nature, it hurts the whole functions of the body. Hence grief indulged fpoils the digeftion and deftroys the appetite; by which means the fpirits are depreffed, the nerves relaxed, the bowels inflated with wind, and the humours, for want of fresh supplies of chyle, vitiated. Thus many an excellent constitution has been ruined by a family misfortune, or any thing that occasions excessive grief.

It is utterly impoffible that any perfon of a dejected mind fhould enjoy health. Life may indeed be dragged out for a few years; but whoever would live to a good old age, muft be good-humoured and cheerful. This indeed is not altogether in our own power; yet our temper of mind, as well as our actions, depend greatly upon ourfelves. We can either affociate with cheerful or melancholy companions, mingle in the amufements and offices of life, or fit ftill and brood over our calamities as we choofe. Thefe, and many fuch things, are certainly in our power, and from thefe the mind generally takes its caft.

OF THE PASSIONS.

The variety of fcenes which prefent themfelves to the fenfes, were certainly defigned to prevent our attention from being too long fixed upon any one object. Nature abounds with variety, and the mind, unlefs fixed down by habit, delights in contemplating new objects. This at once points out the method of relieving the mind in diffrefs. Turn the attention frequently to new objects. Examine them for fome time. When the mind begins to recoil, fhift the fcene. By this means a conftant fucceffion of new ideas may be kept up, till the difagreeable ones entirely difappear. Thus travelling, the ftudy of any art or fcience, reading, or writing on fuch fubjects as deeply engage the attention, will fooner expel grief than the moft fprightly amufements.

It has already been obferved, that the body cannot be healthy unlefs it be exercifed; neither can the mind. Indolence nourifhes grief. When the mind has nothing elfe to think of but calamities, no wonder that it dwells there. Few people who purfue bufinefs with attention are hurt by grief. Inftead therefore of abftracting ourfelves from the world or bufinefs when misfortunes happen, we ought to engage in it with more than ufual attention, to difcharge with double diligence the functions of our flation, and to mix with friends of a cheerful and focial temper.

Innocent amufements are by no means to be neglected. Thefe, by leading the mind infenfibly to the contemplation of agreeable objects, help to difpel the gloom which misfortunes caft over it. They make time feem lefs tedious, and have many other happy effects.

Some perfons, when overwhelmed with grief, betake themfelves to drinking. This is making the cure worfe than the difeafe. It feldom fails to end in the ruin of fortune, character, and conftitution.

Of Love.

Love is perhaps the ftrongest of all the passions. At least, when it becomes violent, it is less subject to the control either of the understanding or will, than any of the rest. Fear, anger, and several other passions, are I necessary

OF THE PASSIONS.

neceffary for the prefervation of the individual, but love is neceffary for the continuation of the fpecies itfelf: it was therefore proper that this paffion should be deeply rooted in the human breast.

Though love be a ftrong paffion, it is feldom fo rapid in its progrefs as feveral of the others. Few perfons fall desperately in love all at once. We would therefore advite every one, before he tampers with this paffion, to confider well the probability of his being able to obtain the object of his wiftes. When that is not likely, he fhould avoid every occasion of increasing it. He ought immediately to flee the company of the beloved object; to apply his mind attentively to bufiness or fludy; to take every kind of amusement; and above all, to endeavour, if poffible, to find another object which may engage his affections, and which it may be in his power to obtain.

There is no paffion with which people are fo apt to tamper as love, although none is more dangerous. Some men make love for amufement, others from mere vanity, or on purpofe to fhew their confequence with the fair. This is perhaps the greateft piece of cruelty which any one can be guilty of. What we eagerly with for, we eafily credit. Hence the too credulous fair are often betrayed into a fituation which is truly deplorable, before they are able to difcover that the pretended lover was only in jeft. But there is no jefting with this paffion. When love has got to a certain height, it admits of no other cure but the poffeffion of its object, which in this cafe ought always, if poffible, to be obtained *.

* The conduct of parents with regard to the difpofal of their children in marriage is often very blameable. An advantageous match is the conflunt aim of parents; while their children often fuffer a real martyrdoin betwixt their inclinations and duty. The first thing which parents ought to confust in difpoing their children in marriage, is certainly their inclinations. Were due regard always paid to thefe, there would be fewer unhappy couples, and parents would not have fo often caufe to repent the feverity of their conduct, after a ruined conflictution, a lost character, or a distracted mind, has shewn them their mistake.

Of

Of Religious Melancholy.

Many perfons of a religious turn of mind behave as if they thought it a crime to be cheerful. They imagine the whole of religion confifts in certain mortifications, or denying themfelves the fmalleft indulgence, even of the most innocent amusements. A perpetual gloom hangs over their countenances, while the deepest melancholy preys upon their minds. At length the faireft prospects vanish, every thing puts on a dismal appearance, and those very objects which ought to give delight, afford nothing but disgust. Life itself becomes a burden, and the unhappy wretch, perfuaded that no evil can equal what he feels, often puts an end to his miferable existence.

It is great pity that ever religion should be fo far perverted, as to become the cause of those very evils which it was designed to cure. Nothing can be better calculated than *True Religion*, to raise and support the mind of its votaries under every affliction that can befal them. It teaches men that even the sufferings of this life are preparatory to the happines of the next; and that all who persist in a course of virtue shall at length arrive at complete felicity.

Perfons whofe bufinefs it is to recommend religion to others, fhould beware of dwelling too much on gloomy fubjects. That peace and tranquillity of mind, which true religion is calculated to infpire, is a more powerful argument in its favour, than all the terrors that can be uttered. Terror may indeed deter men from outward acts of wickednefs, but can never infpire them with that love of God, and real goodnefs of heart, in which alone true religion confifts.

To conclude; the best way to counteract the violence of any passion, is to keep the mind closely engaged in fome useful pursuit.

I have often heard that the late Lord KAIMES, when he faw any literary friend finking under the preffure of melancholy, or fome other corroding paffion, always

IIS

gave this advice in a few emphatical words, " write a " book ;" which he believed to be an infallible remedy. I alfo knew the author of a very beautiful elegy cured of his grief for a wife, whom he had tenderly loved, by ftudying how to express the greatness of his loss, and the pungency of his forrows, in the most plaintive and affecting strains. Indeed, the earness direction of our thoughts to some important object is, as I before hinted, the furest method of subduing passions which may stubbornly resist the control of reason.

CHAP. XI.

OF THE COMMON EVACUATIONS.

T HE principal evacuations from the human body are those by *ftool*, urine, and infensible perspiration. None of these can be long obstructed without impairing the health. When that which ought to be thrown out of the body is too long retained, it not only occasions a *plethora*, or too great fulness of the vessels, but acquires qualities which are hurtful to the health, as acrimony, putrescence, &c.

Of the Evacuation by Stool.

Few things conduce more to health than keeping the body regular. When the *faces* lie too long in the bowels, they vitiate the humours; and when they are too foon difcharged, the body is not fufficiently nourifhed. A medium is therefore to be defired, which can only be obtained by regularity in diet, fleep, and exercife. Whenever the body is not regular, there is reafon to fufpect a fault in one or other of thefe.

Perfons who eat and drink at irregular hours, and who eat various kinds of food, and drink of feveral different liquors at every meal, have no reafon to expect either that their digeftion will be good, or their difcharges regular. Irregularity in eating and drinking difturbs every part of the animal occonomy, and never fails

fails to occafion difeafes. Either too much or too little food will have this effect. The former indeed generally occafions loofenefs, and the latter coffivenefs; but both have a tendency to hurt the health.

It would be difficult to afcertain the exact number of ftools which may be confiftent with health, as thefe differ in the different periods of life, in different conftitutions, and even in the fame conftitution under a different regimen of diet, exercife, &c. It is however generally allowed, that one ftool a-day is fufficient for an adult, and that more or lefs is hurtful. But this, like most general rules, admits of many exceptions. I have known perfons in perfect health who did not go to ftool above once a-week *. Such a degree of costiveness however is not fafe; though the perfon who labours under it may for fome time enjoy tolerable health, yet at length it may occasion difeates.

One method of procuring a ftool every day is to rife betimes, and go abroad in the open air. Not only the pofture in bed is unfavourable to regular ftools, but alfo the warmth. This, by promoting the perfpiration, leffens all the other difcharges.

The method recommended for this purpole by Mr. Locke is likewife very proper, viz. to folicit nature, by going regularly to ftool every morning whether one has a call or not. Habits of this kind may be acquired, which will in time become natural.

Perfons who have frequent recourfe to medicines for preventing coffiveness, feldom fail to ruin their conflitution. Purging medicines frequently repeated weaken the bowels, hurt the digestion, and every dose makes way for another, till at length they become as neceffary as daily bread. Those who are troubled with costiveness ought rather, if possible, to remove it by diet than drugs. They should likewise go thinly clothed, and avoid every thing of an astringent or of an heating nature. The diet and other regimen necessary in this case

• Some perfons have told me that they did not go to flool above once a month.

will be found under the article Costiveness, where this state of the bowels is treated as a difease.

Such perfons as are troubled with an habitual loofernefs, ought likewife to fuit their diet to the nature of their complaint. They fhould ufe food which braces and ftrengthens the bowels, and which is rather of an aftringent quality, as wheat-bread made of the fineft flour, cheefe, eggs, rice boiled in milk, &c. Their drink fhould be red port, claret, brandy and water, in which toafted bread has been boiled, and fuch like.

As an habitual loofeness is often owing to an obstructed perspiration, persons affected with it ought to keep their feet warm, to wear flannel next their skin, and take every other method to promote the perspiration. Further directions with regard to the treatment of this complaint will be found under the article Loofenes.

Of Urine.

So many things tend to change both the quantity and appearances of the urine, that it is very difficult to lay down any determined rules for judging of either *. Dr, Cheyne fays, the urine ought to be equal to three-fourths of the liquid part of our aliment. But fuppole any one were to take the trouble of meafuring both, he would find that every thing which altered the degree of per-

* It has long been an obfervation among phyficians, that the appearances of the urine are very uncertain, and very little to be depended on. No one will be furprifed at this who confiders how many ways it may be affected, and confequently have its appearance altered. The paffions, the flate of the atmosphere, the quantity and quality of the food, the exercife, the clothing, the flate of the other evacuations, and numberlefs other caufes, are fufficient to induce a change either in the quantity or appearance of the urine. Any one who attends to this will be aftonished at the impudence of those daring quacks, who pretend to find out difcales, and prefcribe to patients, from the bare infpection of their urine. These impoltures, however, are very common all over Britain, and, by the amazing credulity of the populace, many of them amais confiderable fortunes. Of all the medical prejudices which prevail in this country, that in favour of urine doctors is the ftrongest. The common people have still an unlimited faith in their skill, although it has been demonstrated that no one of them is able to diffinguish the urine of a horse, or any other animal, from that of a man.

fpiration,

fpiration, would alter this proportion, and likewife that different kinds of aliment would afford very different quantities of urine. Though for thefe, and other reafons, no rule can be given for judging of the precife quantity of urine which ought to be difcharged, yet a perfon of common fenfe will feldom be at a lofs to know when it is in either extreme.

As a free difcharge of urine not only prevents but actually cures many difeafes, it ought by all means to be promoted; and every thing that may obftruct it, fhould be carefully avoided. Both the fecretion and difcharge of urine are leffened by a fedentary life, fleeping on beds that are too foft and warm, food of a dry and heating, quality, liquors which are aftringent and heating, as red port, claret, and fuch like. Thofe who have reafon to fufpect that their urine is in too fmall quantity, or who have any fymptoms of the gravel, ought not only to avoid thefe things, but whatever elfe they find has a tendency to leffen the quantity of their urine.

When the urine is too long retained, it is not only reforbed, or taken up again into the mals of fluids, but by flagnating in the bladder it becomes thicker, the more watery parts flying off first, and the more grofs and earthy remaining behind. By the constant tendency which these have to concrete, the formation of stones and gravel in the bladder is promoted. Hence it comes to pass that indolent and sedentary people are much more liable to these diseases, than perfons of a more active life,

Many perfons have loft their lives, and others have brought on very tedious, and even incurable diforders by retaining their urine too long, from a falfe delicacy. When the bladder has been over-diftended, it often lofes its power of action altogether, or becomes paralytic, by which means it is rendered unable either to retain the urine, or expel it properly. The calls of nature ought never to be poftponed. Delicacy is doubtlefs a virtue, but that can never be reckoned true delicacy, which induces any one to rifk his health, or hazard his life.

But

But the urine may be in too great as well as too fmall a quantity. This may be occafioned by drinking large quantities of weak watery liquors, by the exceffive ufe of alkaline falts, or any thing that ftimulates the kidneys, dilutes the blood, &c. This diforder very foon weakens the body, and induces a confumption. It is difficult to cure, but may be mitigated by ftrengthening diet and aftringent medicines, fuch as are recommended under the article Diabetes, or exceffive difcharge of urine.

Of the Perspiration.

Infenfible perfpiration is generally reckoned the greateft of all the difcharges from the human body. It is of fo great importance to health, that few difeafes attack us while it goes properly on; but when it is obftructed, the whole frame is foon difordered. This difcharge, however, being lefs perceptible than any of the reft, is confequently lefs attended to Hence it is, that acute fevers, rheumatifms, agues, &c. often proceed from obftructed perfpiration, before we are aware of its having taken place.

On examining patients, we find most of them impute their difeases either to violent colds which they had caught, or to flight ones which had been neglected. For this reason, instead of a critical inquiry into the nature of the perspiration, its difference in different feafons, climates, constitutions, &c. we shall endeavour to point out the causes which most commonly obstruct it, and to shew how far they may be either avoided, or have their influence counteracted by timely care. The want of a due attention to these, costs Britain annually fome thousands of useful lives.

Changes in the Atmosphere.

One of the most common causes of obstructed perfpiration, or catching cold, in this country, is the changeableness of the weather, or state of the atmosphere. There is no place where such changes happen more frequently

quently than in Great Britain. With us the degrees of heat and cold are not only very different in the different feafons of the year, but often change almost from one extreme to another in a few days, and fometimes even in the courfe of one day. That fuch changes must affect the state of the perfpiration is obvious to every one *.

The best method of fortifying the body against the changes of the weather is, to be abroad every day. Those who keep most within doors are most liable to catch cold. Such perfons generally render themselves fo delicate, as to feel even the slightest changes in the atmosphere, and by their pains, coughs, and oppressions of the breast, &c. they become a kind of living barometers.

Wet Clothes.

Wet clothes not only by their coldnefs obftruct the perfpiration, but their moifture, by being abforbed, or taken up into the body, greatly increafes the danger. The most robust constitution is not proof against the danger arising from wet clothes; they daily occasion fevers, rheumatisms, and other fatal diforders, even in the young and healthy.

It is impoffible for people who go frequently abroad to avoid fometimes being wet. But the danger might generally be leffened, if not wholly prevented, by changing their clothes foon; when this cannot be done, they fhould keep in motion till they be dry. So far are many from taking this precaution, that they often fit or lie down in the fields with their clothes wet, and frequently fleep even whole nights in this condition. The frequent inftances which we have of the fatal effects of this conduct, ought certainly to deter all from being guilty of it.

* I never knew a more remarkable inftance of the uncertainty of the weather in this country, than happened when I was writing thefe notes. This morning, Aug. 14, 1783, the thermometer in the fhade was down at fifty-three degrees, and a very few days ago it flood above eighty. No one who reflects on fuch great and fudden changes in the atmosphere, will be furprifed to find colds, coughs, rheums, with other affections of the breaft and bowels, fo common in this country.

Wet Feet.

Even wet feet often occasion fatal difeases. The cholic, inflammations of the breast and of the bowels, the iliac passion, *cholera morbus*, &c. are often occasioned by wet feet. Habit will, no doubt, render this less dangerous; but it ought, as far as possible, to be avoided. The delicate, and those who are not accustomed to have their clothes or feet wet, should be peculiarly careful in this respect.

Night Air.

The perfpiration is often obftructed by night air; even in fummer, this ought to be avoided. The dews which fall plentifully after the hotteft day, make the night more dangerous than when the weather is cool. Hence, in warm countries, the evening dews are more hurtful than where the climate is more temperate.

It is very agreeable after a warm day to be abroad in a cool evening; but this is a pleafure to be avoided by all who value their health. The effects of evening dews are gradual indeed, and almost imperceptible; but they are not the lefs to be dreaded: we would therefore advife travellers, labourers, and all who are much heated by day, carefully to avoid them. When the perspiration has been great, these become dangerous in proportion, By not attending to this, in flat marshy countries, where the exhalations and dews are copious, labourers are often feized with intermitting fevers, quinfies, and other dangerous difeases.

Damp Beds.

Beds become damp, either from their not being ufed, ftanding in damp houfes, or in rooms without fire, or from the linen not being dry when laid on the bed. Nothing is more to be dreaded by travellers than damp beds, which are very common in all places where fuel is fcarce. When a traveller, cold and wet, arrives at an inn, he may by means of a good fire, warm diluting liquor, and a dry bed, have the perfpiration reftored; but if he be put into a cold room, and laid in a damp bed,

bed, it will be more obstructed, and the worst confequences will enfue. Travellers should avoid inns which are noted for damp beds, as they would a house infected with the plague, as no man, however robust, is proof against the danger arising from them.

But inns are not the only places where damp beds are to be met with. Beds kept in private families for the reception of ftrangers are often equally dangerous. All kinds of linen and bedding, when not frequently ufed, become damp. How then is it poffible that beds, which are not flept in above two or three times a-year, fhould be fafe? Nothing is more common than to hear people complain of having caught cold by changing their bed. The reafon is obvious: were they careful never to fleep in a bed but what was frequently ufed, they would feldom find any ill confequences from a change.

Nothing is more to be dreaded by a delicate perfon when on a vifit, than being laid in a bed which is kept on purpofe for ftrangers. That ill-judged piece of complaifance becomes a real injury. All the bad confequences from this quarter might eafily be prevented in private families, by caufing their fervants to fleep in the fpare beds, and refign them to ftrangers when they come. In inns, where the beds are ufed almost every night, nothing elfe is neceffary than to keep the rooms well-feafoned by frequent fires, and the linen dry.

That baneful cuftom, faid to be practifed in many inns, of damping fheets, and prefling them, in order to fave wafhing, and afterwards laying them on the beds, ought, when difcovered, to be punified with the utmost feverity. It is really a fpecies of murder, and will often prove as fatal as poifon or gun-fhot. Indeed no linen, efpecially if it has been walked in winter, ought to be ufed till it has been exposed for fome time to the fire; nor is this operation lefs neceffary for linen washed in fummer, provided it has lain by for any length of time. This caution is the more needful, as gentlemen are often exceedingly attentive to what they eat or drink at an inn,

inn, yet pay no regard to a circumstance of much more importance *.

Damp Houses.

Damp houfes frequently produce the like ill confequences; for this reafon thofe who build fhould be careful to chufe a dry fituation. A houfe which ftands on a damp marfhy foil or deep clay, will never be thoroughly dry. All houfes, unlefs where the ground is exceeding dry, fhould have the first floor a little raifed, Servants and others, who are obliged to live in cellars and funk ftories, feldom continue long in health : mafters ought furely to pay fome regard to the health of their fervants, as well as to their own.

Nothing is more common than for people, merely to avoid fome trifling inconveniency, to hazard their lives, by inhabiting a houfe almost as foon as the masons, plaisterers, &c. have done with it : fuch houses are not only dangerous from their dampness, but likewise from the second the fine paint, &c. The association of the tions, and other difeases of the lungs, so incident to people who work in these articles, are sufficient proofs of their being unwholes.

Rooms are often rendered damp by an unfeafonable piece of cleanlinefs; I mean the pernicious cuftom of washing them immediately before company is put into them. Most people catch cold, if they fit but a very short time in a room that has been lately washed; the delicate ought carefully to avoid such a structure, and even the robust are not always proof against its influence \dagger .

Sudden

* If a perfon fulpects that his bed is damp, the fimple precaution of taking off the fheets and lying in the blankets, with all, or most of his clothes on, will prevent all the danger. I have practifed this for many years, and never have been hurt by damp beds, though no constitution, without care, is proof against their baneful influence.

+ People imagine if a good fire is made in a room after it has been washed, that there is no danger from fitting in it; but they mult

Sudden Transitions from Heat to Cold.

The perfpiration is commonly obftructed by SUDDEN TRANSITIONS from heat to cold. Colds are feldom caught, unlefs when people have been too much heated. Heat rarifies the blood, quickens the circulation, and increafes the perfpiration; but when thefe are fuddenly checked, the confequences muft be bad. It is indeed impoffible for labourers not to be too hot upon fome occafions; but it is generally in their power to let themfelves cool gradually, to put on their clothes when they leave off work, to make choice of a dry place to reft themfelves in, and to avoid fleeping in the open fields. Thefe eafy rules, if obferved, would often prevent fevers and other fatal diforders.

It is very common for people, when hot, to drink freely of cold water, or fmall liquors. This conduct is extremely dangerous. Thirft indeed is hard to bear, and the inclination to gratify that appetite frequently gets the better of reafon, and makes us do what our judgment difapproves. Every peafant, however, knows, if his horfe be permitted to drink his bellyful of cold water after violent exercife, and be immediately put into the ftable, or fuffered to remain at reft, that it will kill him. This they take the utmoft care to prevent. It were well if they were equally attentive to their own fafety.

Thirst may be quenched many ways without fwallowing large quantities of cold liquor. The fields afford variety of acid fruits and plants, the very chewing of which would abate thirst. Water kept in the mouth for fome-time, and spit out again, if frequently repeated, will have the same effect. If a bit of bread be eaten along with a few mouthfuls of water, it will both quench thirst more effectually, and make the danger lefs. When

must give me leave to fay that this increases the danger. The evaporation excited by the fire generates cold, and renders the damp more active.

a perfon

a perfon is extremely hot, a mouthful of brandy, or other fpirits, if it can be obtained, ought to be preferred to any thing elfe. But if any one has been fo foolifh, when hot, as to drink freely of cold liquor, he ought to continue his exercife at leaft till what he drank be thotoughly warmed upon his ftomach.

It would be tedious to enumerate all the bad effects which flow from drinking cold liquors when the body is hot. Sometimes this has occafioned immediate death. Hoarfenels, quinfeys, and fevers of various kinds, are its common confequences. Neither is it fafe when warm to eat freely of raw fruits, fallads, or the like. These indeed have not fo fudden an effect on the body as cold liquors, but they are notwithstanding dangerous, and ought to be avoided.

Sitting in a warm room, and drinking hot liquors till the pores are quite open, and immediately going into the cold air, is extremely dangerous. Colds, coughs, and inflammations of the breaft, are the ufual effects of this conduct; yet nothing is more common than for people, after they have drank warm liquors for feveral hours, to walk or ride a number of miles in the coldeft night, or to ramble about in the ftreets *.

People are very apt, when a room is hot, to throw open a window, and to fit near it. This is a moft dangerous practice. Any perfon had better fit without doors than in fuch a fituation, as the current of air is directed against one particular part of the body. Inflammatory fevers, quinfeys, and confumptions have often been occasioned by fitting or standing thinly clothed near an open window. Nor is sleeping with open windows less to be dreaded. That ought never to be done, even in the hostess feason, unless the window is at a distance. I

* The tap-rooms in London and other great towns, where fuch numbers of people fpend their evenings, are highly pernicious. The breath of a number of people crowded into a low apartment, with the addition of fires, candles, the finoke of tobacco, and the fumes of hot liquor, &c. must not only render it hurtful to continue in fach places, but dangerous to go out of them into a cold and chilly atmosphere.

have

have known mechanics frequently contract fatal difeafes, by working ftripped at an open window, and would advife all of them to beware of fuch a practice.

Few things expose people more to catch cold than keeping their own houses too warm: such perfons may be faid to live in a fort of hot-houses; they can hardly fir abroad to visit a neighbour but at the hazard of their lives. Were there no other reason for keeping houses moderately cool, that alone is sufficient: but no house that is too hot can be wholesome; heat destroys the spring and elasticity of the air, and renders it less fit for expanding the lungs, and the other purposes of respiration. Hence it is that confumptions and other diseases of the lungs prove fo fatal to people who work in forges, glass-houses, and the like.

Some are even fo fool-hardy, as to plunge themfelves, when hot, in cold water. Not only fevers, but madnefs itfelf, has frequently been the effect of this conduct. Indeed it looks too like the action of a madman to deferve a ferious confideration.

The refult of all these observations is, that every one ought to avoid, with the utmost attention, all fudden transitions from heat to cold, and to keep the body in as uniform a temperature as possible; or where that cannot be done, to take care, when heated, to let it cool gradually.

People may imagine that too ftrict an attention to these things would tend to render them delicate. So far, however, is this from being my design, that the very first rule proposed for preventing colds is, to harden the body, by enuring it daily to the open air.

I fhall put an end to what relates to this part of my fubject, by giving an abstract of the justly celebrated advice of Celfus, with respect to the prefervation of health. "A man," fays he, "who is bleffed with good "health, should confine himself to no particular rules, "either with respect to regimen or medicine. He "ought frequently to diversify his manner of living; "to be fometimes in town, fometimes in the country; "to hunt, fail, indulge himself in rest, but more fre-2 "guently

" quently to ufe exercife. He ought to refufe no kind of food that is commonly ufed, but fometimes to eat more and fometimes lefs; fometimes to make one at an entertainment, and fometimes to forbear it; to make rather two meals a day than one, and always to eat heartily, provided he can digeft it. He ought neither too eagerly to purfue, nor too fcrupuloufly to avoid intercourfe with the fair fex : pleafures of this kind, rarely indulged, render the body alert and active; but when too frequently repeated, weak and languid. He fhould be careful in time of health not to deftroy, by exceffes of any kind, that vigour of conftitution which fhould fupport him under ficknefs."

This plain, yet elegant and judicious fummary of the most useful maxims of health confirms the justness of my former remark, that enlightened Medicine breathes the true spirit of liberal indulgence, laying down no rules but such as a man of sense would cheerfully follow, and forbidding nothing but what is incompatible with real happines. Here the votaries of fashion and folly may learn to correct their own mistaken ideas of enjoyment: the epicure may acquire a reliss for rational gratification; and the man of pleasure may be taught the æconomy of love.

and in render then deligness Er

this troub being any difficult the

PART H.

[129]

OF DISEASES.

CHAP. XII.

OF THE KNOWLEDGE AND CURE OF DISEASES.

THE cure of difeafes does not depend fo much upon fcientific principles as many imagine. It is chiefly the refult of experience and obfervations By attending the fick, and carefully obferving the various occurrences in difeafes, a great degree of accuracy may be acquired, both in diffinguifhing their fymptoms, and in the application of medicines. Hence fenfible nurfes, and other perfons who wait upon the fick, often forefee the patient's fate fooner than those who have been bred to phyfic. We do not, however, mean to infinuate that a medical education is of no ufe: It is doubtlefs of the greatest importance, but it never can fupply the place of observation and experience.

Every difeafe may be confidered as an affemblage of fymptoms, and mult be diftinguished by those which are most obvious and permanent. Instead, therefore, of giving a classical arrangement of difeases, according to the fystematic method, it will be more fuitable, in a performance of this nature, to give a full and accurate defoription of each particular difease as it occurs ; and, where any of the symptoms of one difease have a near refemblance to those of another, to take notice of that circumstance, and at the same time to point out the peculiar or characteristic symptoms by which it may be distinguished. By a due attention to these, the investigation of difeases will be found to be a less difficult matter than most people would at first be ready to imagine.

A proper

A proper attention to the patient's age, fex, temper of mind, conflictution, and manner of life, will likewife greatly affift, both in the investigation and treatment of difeafes.

In childhood, the fibres are lax and foft, the nerves extremely irritable, and the fluids thin; whereas in old age, the fibres are rigid, the nerves become almost infensible, and many of the vessels imperviable. These and other peculiarities render the diseases of the young and aged very different, and of course they must require a different method of treatment.

Females are liable to many difeafes which do not afflict the other fex : befides, the nervous fystem being more irritable in them than in men, their difeafes require to be treated with greater caution. They are lefs able to bear large evacuations ; and all stimulating medicines ought to be administered to them with a sparing hand.

Particular conflictutions not only difpofe perfons to peculiar difeafes, but likewife render it neceffary to treat thefe difeafes in a peculiar manner. A delicate perfon for example, with weak nerves, who lives moftly within doors, must not be treated, under any difeafe, precifely in the fame manner as one who is hardy and robust, and who is much exposed to the open air.

The temper and mind ought to be carefully attended to in difeafes. Fear, anxiety, and a fretful temper both occafion and aggravate difeafes. In vain do we apply medicines to the body to remove maladies which proceed from the mind. When it is affected, the beft medicine is to foothe the paffions, to divert the mind from anxious thought, and to keep the patient as eafy and cheerful as poffible.

Attention ought likewife to be paid to the climate, or place where the patient lives, the air he breathes, his diet, &c. Such as live in low marfhy fituations are fubject to many difeafes which are unknown to the inhabitants of high countries. Those who breathe the impure air of cities, have many maladies to which the more happy ruffics are entire strangers. Perfons who feed grofsly,

grofsly, and indulge in ftrong liquors, are liable to difeafes which do not affect the temperate and abstemious, &c.

It has already been obferved, that the different occupations and fituations in life difpofe men to peculiar difeafes. It is therefore neceffary to inquire into the patient's occupation, manner of life, &c. This will not only affilt us in finding out the difeafe, but will likewife direct us in the treatment of it. It would be very imprudent to treat the laborious and the fedentary precifely in the fame manner, even fuppofing them to labour under the fame difeafe.

It will likewife be proper to inquire, whether the difeafe be conflicational or accidental; whether it has been of long or fhort duration; whether it proceeds from any great and fudden alteration in the diet, manner of life, &c. The flate of the patient's body, and of the other evacuations, ought alfo to be inquired into; and likewife whether he can with eafe perform all the vital and animal functions, as breathing, digeftion. &c.

Lattly, it will be proper to inquire to what difeafes the patient has formerly been liable, and what medicines were most beneficial to him; if he has a ftrong aversion to any particular drug, &c.

As many of the indications of cure may be answered by diet alone, it is always the first thing to be attended to in the treatment of difeases. Those who know no better, imagine that every thing which goes by the name of a medicine possible forme wonderful power or fecret charm, and think, if the patient seallows enough of drugs, that he mult do well. This mistake has many ill confequences; it makes people trust to drugs, and neglect their own endeavours; besides, it discourages all attempts to relieve the fick where medicines cannot be obtained.

Medicines are no doubt ufeful in their places; and when administered with prudence, may do much good; but when they are put in place of every thing elfe, or administered at random, which is not feldom the cafe, they must do mischief. We would therefore with

OF THE KNOWLEDGE

to call the attention of mankind from the purfuit of fecret medicines, to fuch things as they are acquainted with. The proper regulation of these may often do much good, and there is little danger of their ever doing hurt.

Every difeafe weakens the digeftive powers. The diet ought therefore, in all difeafes, to be light and of eafy digeftion. It would be as prudent for a perfon with a broken leg to attempt to walk, as for one in a fever to eat the fame kind of food. and in the fame quantity, as when he was in perfect health. Even abflinence alone will often cure a fever, effectally when it has been occafioned by excefs in eating or drinking.

In all fevers attended with inflammation, as pleurifies, peripneumonies, &c. thin gruels, wheys, watery infufions of mucilaginous plants, roots, &c. are not only proper for the patient's food, but they are likewife the best medicines which can be administered.

In fevers of a flow, nervous, or putrid kind, where there are no fymptoms of inflammation, and where the patient must be supported with cordials, that intention can always be more effectually answered by nourishing diet and generous wines, than by any medicines yet known.

Nor is a proper attention to diet of lefs importance in chronic than in acute difeafes. Perfons afflicted with low fpirits, wind, weak nerves, and other hypochondriacal affections, generally find more benefit from the ufe of folid food and generous liquors, than from all the cordial and carminative medicines which can be adminiftered to them.

The fcurvy, that most obstinate malady, will fooner yield to a proper vegetable diet, than to all the boasted antifcorbutic remedies of the shops.

In confumptions, when the humours are vitiated, and the ftomach fo much weakened as to be unable to digeft the folid fibres of animals, or even to affimilate the juices of vegetables, a diet confifting chiefly of *milk* will not only fupport the patient, but will often cure the difeafe after every other medicine has failed. Nor

132

AND CURE OF DISEASES.

Nor is an attention to other things of lefs importance than to diet. The ftrange infatuation which has long induced people to fhut up the fick from all communication with the external air, has done great mifchief. Not only in fevers, but in many other difeafes, the patient will receive more benefit from having the frefh air prudently admitted into his chamber, than from all the medicines which can be given him.

Exercife may likewife in many cafes be confidered as a medicine: Sailing, or riding on horfeback, for example, will be of more fervice in the cure of confumptions, glandular obftructions, &c. than any medicine yet known. In difeafes which proceed from a relaxed flate of the folids, the cold bath, and other parts of the gymnaftic regimen, will be found equally beneficial.

Few things are of greater importance in the cure of difeafes than cleanlinefs. When a patient is fuffered to lie in dirty clothes, whatever perfpires from his body is again reforbed, or taken up into it, which ferves to nourifh the difeafe and increafe the danger. Many difeafes may be cured by cleanlinefs alone; most of them may be mitigated by it, and in all of them it is highly neceffary both for the patient and those who attend him.

Many other obfervations, were it neceffary, might be adduced to prove the importance of a proper regimen in difeafes. Regimen will often cure difeafes without medicine, but medicine will feldom fucceed where a proper regimen is neglected. For this reafon, in the treatment of difeafes, we have always given the firft place to regimen. Thofe who are ignorant of medicine may confine themfelves to it only. For others who have more knowledge, we have recommended fome of the moft fimple but approved forms of medicine in every difeafe. Thefe, however, are never to be adminiftered but by people of better underftanding; nor even by them without the greateft precaution.

The clearnefs and fimplicity with which I took care to express myself on these points, would, I thought, have prevented the possibility of any misrepresentation. Yet I find that a certain low class of felf-appointed

practi-

134 OF THE KNOWLEDGE, &c.

praclitioners, who call themfelves of the faculty, take no fmall pains to infinuate, that my obfervations on the prevention and cure of difeafes ferve only to encourage the fatal practice of domeftic quackery. This is equally inconfistent with candour and truth. The obvious tendency of all my remarks is to enlighten the minds of the people on a fubject of fuch immediate concern as their health, and thus to guard them against the bad effects of ignorance and rafhnels on their own part, and of impudence and deceit on the part of others. I believe every man of common understanding, who reads my book, will feel his caution increaled against the use of the most fimple medicines; instead of being tempted, upon every trifling occafion, to run the rifk of poifoning himfelf, or his family, by drugs, and dangerous compounds, from an apothecary's flipp.

I fhould rather have expected to be blamed for teaching people to place very little reliance on the efficacy of any medicine; which was, indeed, one of the objects I had in view, for the express purpose of directing general attention to the far more affured means of preserving health, namely air, cleanlines, diet, exercile, and the management of the passions. Upon these subjects I enlarged with peculiar earness, well-knowing how much easier it is to prevent diforders before-hand, than to cure them afterwards.

Even in cafes of actual infirmity and difeafe, I have intimated a wife, that the fe who are ignorant of phyfic would confine themfelves to regimen only, and leave the medical treatment of their complaints to perfons of better information. The remedies which I have prefcribed may be entruffed to fuch hands with perfect fafety; and if the directions I give do not operate as a check upon rafhnefs, I know of no words firong enough to produce that effect. I write in plain English; but I can affure any patient, that the prefcription will not be less falutary for wanting the boafted charm of barbarous Latin, and of ftill more barbarous hieroglyphics.

TA THIN [1357]VST TO

C H A P. XIII. Longingo T

CF FEVERS IN GENERAL.

AS more than one half of mankind is faid to perifh by fevers, it is of importance to be acquainted with their caufes. The most general caufes of fevers are, *infection*, errors in diet, unwholefome air, violent emotions of the mind, excefs or fuppression of usual evacuations, external or internal injuries, and extreme degrees of heat or cold. As most of these have already been treated of at confiderable length, and their effects shewn, we shall not now refume the confideration of them, but shall only recommend it to all, as they would wish to avoid fevers and other stated difeases, to pay the most punctual attention to these articles.

Fevers are not only the most frequent of all difeafes, but they are likewile the most complex In the most fimple tpecies of fever there is always a combination of feveral different fymptoms. The diffinguishing fymptoms of fever are, *increased heat*. frequency of pulse, loss of appetite, general debility, pain in the head, and a difficulty in performing some of the vital or animal functions, The other fymptoms usually attendant on fevers are, nausea, thirst, anxiety, delirium, wearines, wasting of the flesh, want of fleep, or the fleep diffurbed and not refreshing.

When the fever comes on gradually, the patient generally complains first of languor or listless, foreness of the flesh, or the bones, as the country people express ir, heaviness of the head, loss of appetite, fickness, with clamminess of the mouth; after some time come on excessive heat, violent thirst, rettless, &c.

When the fever attacks fuddenly, it always begins with an uneafy fenfation of exceffive cold, accompanied with debility and lots of appetite; trequently the cold is attended with fhivering, oppretion about the heart, and ficknefs at ftomach, or vomiting.

Fevers are divided into continual, remitting, intermitting, and fuch as are attended with cutaneous eruption

or topical inflammation, as the fmall-pox, erifipelas, &c. By a continual fever is meant that which never leaves the patient during the whole courfe of the difeafe, or which fhews no remarkable increase or abatement in the fymptoms. This kind of fever is likewife divided into acute, flow, and malignant. The fever is called *acute* when its progrefs is quick, and the fymptoms violent; but when these are more gentle, it is generally denominated *flow*. When livid or petechial spots thew a putrid state of the humours, the fever is called *malignant*, putrid, or petechial.

A remitting fever differs from a continual only in degree. It has frequent increases and decreases, or exacerbations and remiffions, but never wholly leaves the patient during the course of the difease. Intermitting fevers, or agues, are those which, during the time that the patient may be faid to be ill, have evident intervals or remiffions of the fymptoms.

As a fever is only an effort of Nature to free herfelf from an offending caufe, it is the bufinels of thole who have the care of the fick to observe with diligence which way Nature points, and to endeavour to affift her operations. Our bodies are fo framed, as to have a conftant tendency to expel or throw off whatever is injurious to health. This is generally done by urine, fweat, ftool, expectoration, vomit, or fome other evacuation.

There is realon to believe, if the efforts of Nature, at the beginning of a fever, were duly attended to and promoted, it would feldom continue long; but when her attempts are either neglected or counteracted, it is no wonder if the difeafe prove fatal. There are daily inftances of perfons who, after catching cold, have all the fymptoms of a beginning fever; but by keeping warm, drinking diluting liquors, bathing their feet in warm water, &c. the fymptoms in a few hours difappear, and the danger is prevented. When fevers of a putrid kind threaten, the belt method of obviating their effects is by repeated vomits.

Our defign is not to enter into a critical inquiry into the nature and immediate caufes of fevers, but to mark their

136

their most obvious fymptoms, and to point out the proper treatment of the patient with respect to his diet, drink, air, &c. in the different stages of the difease. In these articles the inclinations of the patient will in a great measure direct our conduct.

Almost every perfon in a fever complains of great thirst, and calls out for drink, especially of a cooling nature. This at once points out the use of water, and other cooling liquors. What is so likely to abate the heat, attenuate the humours, remove spass and obstructions, promote perspiration, increase the quantity of urine, and in short produce every falutary effect in an ardent or inflammatory fever, as drinking plentifully of water, thin gruel, or any other weak liquor, of which water is the basis? The necessfity of diluting liquors is pointed out by the dry tongue, the parched shin, and the burning heat, as well as by the unquenchable thirst of the patient.

Many cooling liquors, which are extremely grateful to patients in a fever, may be prepared from fruits, as decoctions of tamarinds, apple-tea, orange-whey, and the like. Mucilaginous liquors might alfo be prepared from marfh-mallow roots, linfeed, lime-tree buds, and other mild vegetables. These liquors, especially when acidulated, are highly agreeable to the patient, and fhould never be denied him.

At the beginning of a fever, the patient generally complains of great laffitude or wearinefs, and has no inclination to move. This evidently flews the propriety of keeping him eafy, and, if poffible, in bed. Lying in bed relaxes the Ipafms, abates the violence of the circulation, and gives Nature an opportunity of exerting all her force to overcome the difeafe. The bed alone would often remove a fever at the beginning; but when the patient ftruggles with the difeafe, inftead of driving it off, he only fixes it the deeper, and renders it more dangerous. This obfervation is too often verified in travellers, who happen when on a journey to be feized with a fever. Their anxiety to get home, induces them to travel with the fever upon them; which conduct feldom fails to render it fatal.

In fevers, the mind as well as the body fhould be kept eafy. Company is feldom agreeable to the fick. Indeed every thing that diffurbs the imagination, increafes the difeafe : for which reafon every perfon in a fever ought to be kept perfectly quiet, and neither allowed to fee nor hear any thing that may in the leaft affect or difcompose his mind.

Though the patient in a fever has the greatest inclination for drink, yet he feldom has any appetite for folid food : hence the impropriety of urging him to take victuals is evident. Much folid food in a fever is every way hurtful. It opprefies nature, and, instead of nourishing the patient, ferves only to feed the difease. What food the patient takes, should be in small quantity, light, and of easy digestion. It ought to be chiefly of the vegetable kind, as panada, roasted apples, gruels, and fuch like.

Poor people, when any of their family are taken ill, run directly to their rich neighbours for cordials, and pour wine, fpirits, &c. into the patient, who perhaps never had been accultomed to talte fuch liquors when in health. If there be any degree of fever, this conduct must increase it; and if there be none, this is the ready way to raife one. Stuffing the patient with fweetmeats, and other delicacies, is likewife very pernicious. These are always harder to digest than common food, and cannot fail to hurt.

Nothing is more defired by a patient in a fever, than fresh air. It not only removes his anxiety, but cools the blood, revives the spirits, and proves every way benessicial. Many patients are in a manner stilled to death in fevers for want of fresh air; yet such is the unaccountable infatuation of most people, that the moment they think a perfon in a fever, they imagine he should be kept in a close chamber, into which not one particle of fresh air mult be admitted. Instead of this, there ought to be a constant stream of fresh air into a fick perfon's chamber, so as to keep it moderately cool. Indeed, its degree of warmth ought never to be greater than is agreeable to one in perfect health.

Nothing

Nothing fpoils the air of a fick perfon's chamber, or hurts the patient more, than a number of people breathing in it. When the blood is inflamed, or the humours in a putrid flate, air that has been breathed repeatedly will greatly increase the difea e. Such air not only loses its fpring, and becomes unfit for the purpose of respiration, but acquires a noxious quality, which renders it in a manner poisonous to the fick.

In fevers, when the patient's fpirits are low and depreffed, he is not only to be fupported with cordials, but every method fhould be taken to cheer and comfort his mind. Many, from a miftaken zeal, when they think a perfon in danger, inftead of folacing his mind wi h the hopes and confolations of religion, frighten him with the views of hell and damnation. It would be unfuitable here to dwell upon the impropriety and dangerous confequences of this conduct; it often hurts the body, and there is reafon to believe feldom benefits the foul.

Among common people, the very name of a fever generally suggests the necessity of bleeding. This notion feems to have taken its rife from most fevers in this country having been formerly of an inflammatory nature; but true inflammatory fevers are now feldom to be met with. Sedentary occupations, and a different manner of living, have fo changed the flate of difeafes in Britain, that there is now hardly one fever in ten where the lancet is neceffary. In most low, nervous, and putrid tevers, which are now fo common, bleeding is really hurtful, as it weakens the patient, finks his fpirits, &c. We would recommend this general rule, never to bleed at the beginning of a fever, unless there be evident figns of inflammation. Bleeding is an excellent medicine when neceffary, but should never be wantonly performed.

It is likewife a common notion, that fweating is always neceffary in the beginning of a fever. When the fever proceeds from an obstructed perspiration, this notion is not ill founded. If the patient only he in bed, bathe his feet and legs in warm water, and drink plentifully

tifully of warm water-gruel, or any other weak, diluting liquor, he will feldom fail to perfpire freely. The warmth of the bed, and the diluting drink, will relax the univerfal fpafm, which generally affects the fkin at the beginning of a fever; it will open the pores, and promote the perfpiration, by means of which the fever may often be carried off. But inftead of this, the common practice is to heap clothes upon the patient, and to give him things of a hot nature, as fpirits, fpiceries, &c. which fire his blood, increafe the fpafms, and render the difeafe more dangerous.

In all fevers, a proper attention fhould be paid to the patient's longings. Thefe are the calls of Nature, and often point out what may be of real ufe. Patients are not indeed to be indulged in every thing that the fickly appetite may crave; but it is generally right to let them have a little of what they eagerly defire, though it may not feem altogether proper. What the patient longs for, his ftomach will generally digeft; and fuch things have fometimes a very happy effect.

When a patient is recovering from a fever, great care is neceffary to prevent a relaple. Many perfons, by too foon imagining themfelves well, have loft their lives, or contracted other difeafes of an obftinate nature. As the body after a fever is weak and delicate, it is neceffary to guard against catching cold. Moderate exercise in the open air will be of ufe, but great fatigue is by all means to be avoided; agreeable company will also have a good effect. The diet must be light, but nourifhing. It should be taken frequently, but in finall quantities. It is dangerous, at such a time, to eat as much as the ftomach may crave.

From the great variety of fevers that afflict the human body, it is impoffible to find any medicine adapted to them all, or, indeed, to all the fymptoms of any one of them. Notwithstanding this, the people of England have, for half a century, been fwallowing a powder faid to poffers wonderful virtues in the cure of fevers. Nor has the ufe of this powder been confined to England. It has been carried to every part of the globe; and great cures

cures have been attributed to it, with what truth I will not pretend to fay. L remember bleeding to have been as much in vogue in fevers, though now it is feldom preferibed, unlefs in local inflammations. But there is a fashion in physic, as well as in other things; and it is always herefy to talk against the doctrine of the day.

This fever powder, like other quack medicines, is faid to be good in a variety of complaints, and is used by fome people in every diforder, real or imaginary. I knew a lady, who not only administered it to all the poor of the parish when ill, but likewise gave it to her dogs and horses; and never failed to take it daily herfelf, till she destroyed her constitution. Many perfons look upon it as a *panacea*, or universal remedy, and keep it continually by them in case of emergencies. The fatal consequences of such credulity must be often irreparable. This, at least, was the fituation of an old General of my acquaintance, whom no argument could diffuade from taking the powder, till he lost the use of all his extremities.

There is not a greater folecism in language, nor a greater absurdity in real practice, than to pretend that any one medicine is of certain efficacy in fevers. The most skilful physicians that ever existed have always found it necessary to watch attentively the progress of a fever; and to adapt both the regimen and medicines to the different changes and symptoms as they occurred.

CHAP. XIV.

OF INTERMITTING FEVERS, OR AGUES.

INTERMITTING fevers afford the best opportunity both of observing the nature of a fever, and also the effects of medicine. No perfon can be at a loss to diftinguish an intermitting fever from any other, and the proper medicine for it is now almost universally known. 3

The feveral kinds of intermitting fevers take their names from the period in which the fit returns, as quotidian, tertian, quartan, &c.

CAUSES. — Agues are occafioned by effluvia from putrid ftagnating water. This is evident from their abounding in rainy feafons, and being moft frequent in countries where the foil is marfhy, as in Holland, the Fens of Cambridgefhire, the Hundreds of Effex, &cc. This difeafe may allo be occafioned by eating too much flone fruit, by a poor watery diet, damp houfes, evening dews, lying upon the damp ground, watching, fatigue, depreffing paffions, and the like. When the inhabitants of a high country remove to a low one, they are generally feized with intermitting fevers, and to fuch the difeafe is moft apt to prove fatal. In a word, whatever relaxes the folids, diminifhes the perfpiration, or obftructs the circulation in the capillary or finall veffels, difpofes the body to agues.

SYMPTOMS. — An intermitting fever generally begins with a pain of the head and loins, wearinefs of the limbs, coldnefs of the extremities, ftretching, yawning, with fometimes great ficknefs and vomiting; to which fucceed fhivering and violent fhaking. Afterwards the fkin becomes moift, and a profule fweat breaks out, which generally terminates the fit or paroxyfm. Sometimes indeed the difeafe comes on fuddenly, when the perfon thinks himfelf in perfect health; but it is more commonly preceded by liftleffnefs, lofs of appetite, and the fymptoms mentioned above.

REGIMEN. — While the fit continues, the patient ought to drink freely of water-gruel, orange-whey, weak camomile-tea; or, if his fpirits be low, fmall winewhey, fharpened with the juice of lemon. All his drink fhould be warm, as that will affift in bringing on the fweat, and confequently fhorten the paroxyfm *.

• Dr. Lind fays, that twenty or twenty-five drops of laudanum put into a cup of the patient's drink, and given about half an hour after the commencement of the hot fit, promotes the fweat, fhortens the fit, relieves the head, and tends greatly to remove the difeafe.

Between

Between the paroxyfms, the patient must be fupported with food that is nourishing, but light and easy of digestion, as veal or chicken broths, fago, gruel with a little wine, light puddings, and such like. His drink may be small negus, acidulated with the juice of lemons or oranges, and sometimes a little weak punch. He may likewife drink infusions of bitter herbs, as camomile, wormwood, or water-trefoil, and may now and then take a glass of small wine, in which gentian root, centaury, or fome other bitter, has been infused.

As the chief intentions of cure in an ague are to brace the folids, and promote perfpiration, the patient ought to take as much exercife between the fits as he can bear. If he be able to go abroad, riding on horfeback, or in a carriage, will be of great fervice. But if he cannot bear that kind of exercife, he ought to take fuch as his ftrength will permit. Nothing tends more to prolong an intermitting fever, than indulging a lazy indolent difpofition.

Intermitting fevers, under a proper regimen, will often go off without medicine : and when the difeafe is mild, in an open dry country, there is feldom any danger from allowing it to take its courfe ; but when the patient's strength seems to decline, or the paroxysms are so violent that his life is in danger, medicine ought immediately to be administered. This, however, should never be done till the difease be properly formed, that is to fay, till the patient has had feveral fits of shaking and sweating.

MEDICINE. — The firft thing to be done in the cure of an intermitting fever, is to cleanfe the flomach and bowels. This not only renders the application of other medicines more fafe, but likewife more efficacious. In this difeafe, the flomach is generally loaded with cold vifcid phlegm, and frequently great quantities of bile are difcharged by vomit ; which plainly points out the neceffity of fuch evacuations. Vomits are therefore to be adminiftered before the patient takes any other medicine. A dofe of ipecacuanha will generally anfwer this purpofe very well. A fcruple or half a dram of the powder

powder will be fufficient for an adult, and for a younger perfon the dofe must be less in proportion. After the vomit begins to operate, the patient ought to drink plentifully of weak camomile-tea. The vomit fhould be taken two or three hours before the return of the fit, and may be repeated at the diffance of two or three days. Vomits not only cleanfe the ftomach, but increafe the perfpiration, and all the other fecretions, which render them of fuch importance, that they often cure intermitting fevers without the affiltance of any other medicine.

Purging medicines are likewife uleful and often neceffary in intermitting fevers. A fmart purge has been known to cure an obflinate ague, after the Peruvian bark and other medicines had been used in vain. Vomits, however, are more fuitable in this difeafe, and render purging less necessary; but if the patient be afraid to take a vomit, he ought in this cafe to cleanfe the bowels by a dofe or two of Glauber's falt, jalap, or rhubarb.

Bleeding may fometimes be proper at the beginning of an intermitting fever, when exceflive heat, a delirium, &c. give reason to suspect an inflammation ; but as the blood is feldom in an inflammatory flate in intermitting fevers, this operation is rarely neceffary. When frequently repeated, it tends to prolong the difeafe.

After proper evacuations, the patient may fafely ufe the Peruvian bark, which may be taken in any way that is most agreeable to him. No preparation of the bark feems to answer better than the most simple form in which it can be given, viz. in powder.

Two ounces of the best Peruvian bark, finely powdered, may be divided into twenty-four doses. Thele may either be made into boluffes, as they are used, with a little fyrup of lemon, or mixed in a glafs of red wine, a cup of camomile-tea, water-gruel, or any other drink that is more agreeable to the patient *.

* It has lately been observed, that the red bark is more powerful than that which has for fome time been in common use. Its iuperior

In

In an ague which returns every day, one of the above dofes may be taken every two hours during the interval of the fits. By this method, the patient will be able to take five or fix dofes between each paroxyfm. In a tertian, or third day ague, it will be fufficient to take a dofe every third hour, during the interval, and in a quartan every fourth. If the patient cannot take fo large a dofe of the bark, he may divide each of the powders into two parts, and take one every hour, &c. For a young perfon, a fmaller quantity of this medicine will be fufficient, and the dofe muft be adapted to the age, conflitution, and violence of the fymptoms *.

The above quantity of bark will frequently cure an ague; the patient, however, ought not to leave off taking the medicine as foon as the paroxyfms are ftopped, but fhould continue to ufe it till there is reafon to believe the difeafe is entirely overcome. Most of the failures in the cure of this difeafe are owing to patients not continuing to ufe the medicine long enough. They are generally directed to take it till the fits are ftopped, then to leave it off, and begin again at fome diftance of time; by which means the difeafe gathers ftrength, and often returns with as much violence as before. A relapfe may always be prevented by the patient's continuing to take dofes of the medicine for fome time after the fymptoms difappear. This is both the most fafe and effectual method of cure.

fuperior efficacy feems to arife from its being of a more perfect growth than the quill-bark, and confequently more fully impregnated with the medical properties of the plant.

* In intermitting fevers of an obflinate nature, I have found it neceffary to throw in the bark much fafter. Indeed, the benefits arifing from this medicine depend chiefly upon a large quantity of it being administered in a short time. Several ounces of bark given in a few days, will do more than as many pounds taken in the courfe of some weeks. When this medicine is intended either to stop a mortification, or cure an obflinate ague, it ought to be thrown in as fast as the stomach can possibly bear it. Inattention to this circumstance has hurt the reputation of one of the best medicines of which we are in possible.

An

145

An ounce of gentian root, calamus aromaticus, and orange-peel, of each half an ounce, with three or four handfuls of camomile-flowers, and an handful of coriander-feed, all bruifed together in a mortar, may be ufed in form of infufion or tea. About half an handful of thefe ingredients may be put into a tea-pot, and an Englifh pint of boiling water poured on them. A cup of this infufion drank three or four times a-day, will greatly promote the cure. Such patients as cannot drink the watery infufion, may put two handfuls of the fame ingredients into a bottle of white wine, and take a glafs of it twice or thrice a-day. If patients drink freely of the above, or any other proper infufion of bitters, a fmaller quantity of bark than is generally ufed will be fufficient to cure an ague *.

Thole who cannot fwallow the bark in fubftance, may take it in decoction or infufion. An ounce of bark in powder may be infufed in a bottle of white wine for four or five days, frequently fhaking the bottle, afterwards let the powder fubfide, and pour off the clear liquor. A wine-glafs may be drank three or four times a day, or oftener, as there is occafion. If a decoction be more agreeable, an ounce of the bark, and two drams of fnake-root bruifed, with an equal quantity of falt of wormwood, may be boiled in a quart of water, to an Englifh pint. To the ftrained liquor may be added an equal quantity of red wine, and a glafs of it taken frequently.

In obstinate agues, the bark will be found much more efficacious when affisted by brandy, or other warm cordials, than if taken alone. This I have had frequently

• There is reafon to believe, that fundry of our own plants or barks, which are very bitter and aftringent, would fucceed in the cure of intermitting fevers, efpecially when affifted by aromatics. But as the Peruvian bark has been long approved in the cure of this difeafe, and is now to be obtained at a very reafonable rate, it is of lefs importance to fearch after new medicines. We cannot however omit taking notice, that the Peruvian bark is very often adulterated, and that it requires confiderable fkill to diffinguifh between the genuine and the falfe. This ought to make people very cautious of whom they purchafe it.

occafion

occafion to obferve in a country where intermittent fevers were endemical. The bark feldom fucceeded unlefs affifted by fnake-root, ginger, canella alba, or fome other warm aromatic. When the fits are very frequent and violent, in which cafe the fever often approaches towards an inflammatory nature, it will be fafer to keep out the aromatics, and to add falt of tartar in their ftead. But in an obftinate tertian or quartan, in the end of autumn or beginning of winter, warm and cordial medicines are abfolutely neceffary *.

As autumnal and winter agues generally prove much more obftinate than those which attack the patient in spring or summer, it will be necessary to continue the use of medicines longer in the former than in the latter. A perfon who is feized with an intermitting fever in the beginning of winter, ought frequently, if the seafon prove rainy, to take a little medicine, although the difease may seem to be cured, to prevent a relapse, till the return of the warm seafon. He ought likewise to take care not to be much abroad in wet weather, especially in cold easterly winds.

When agues are not properly cured, they often degenerate into obftinate chronical difeafes, as the dropfy, jaundice, &c. For this reafon all poffible care fhould be taken to have them radically cured, before the conftitution has been too much weakened.

Though nothing is more rational than the method of treating intermitting fevers, yet, by fome ftrange infatuation, more charms and whimfical remedies are daily ufed for removing this than any other difeafe. There is hardly an old woman who is not in pofferfion of a noftrum for ftopping an ague; and it is amazing with what readinefs their pretenfions are believed. Those in diffres

* In obfinate agues, when the patient is old, the habit phlegmatic, the feafon rainy, the fituation damp, or the like, it will be neceffary to mix with two ounces of the bark, half an ounce of Virginian fnake-root, and a quarter of an ounce of ginger, or fome other warm aromatic; but when the fymptoms are of an inflammatory nature, half an ounce of falt of wormwood or falt of tartar may be added to the above quantity of bark.

eagerly

eagerly grafp at any thing that promifes fudden relief; but the fhorteft way is not always the beft in the treatment of difeafes. The only method to obtain a fafe and lafting cure, is gradually to affift Nature in removing the caufe of the diforder.

Some indeed try bold, or rather fool-hardy experiments, to cure agues, as drinking great quantities of ftrong liquors, jumping into a river, taking arfenic, &c. Thefe may fometimes have the defired effect, but muft always be attended with danger *. When there is any degree of inflammation, or the leaft tendency to it, fuch experiments may prove fatal. The only patient whom I remember to have loft in an intermitting fever, evidently killed himfelf by drinking ftrong liquor, which fome perfon had perfuaded him would prove an infallible remedy.

Many dirty things are extelled for the cure of intermitting fevers, as fpiders' cobwebs, fnuffings of candles, &c. Though thefe may fometimes fucceed, yet their very naftinefs is fufficient to fet them afide, efpecially when cleanly medicines will anfwer the purpofe better. The only medicine that can be depended upon for thoroughly curing an intermittent fever, is the Peruvian bark. It may always be ufed with fafety: and I can honeftly declare, that in all my practice I never knew it fail, when combined with the medicines mentioned above, and duly perfifted in.

Where agues are endemical, even children are often afflicted with that difeafe. Such patients are very difficult to cure, as they can feldom be prevailed upon to take the bark, or any other difagreeable medicine. One method of rendering this medicine more palatable, is to make it into a mixture with diftilled waters and fyrup, and afterwards to give it an agreeable fharpnefs with the elixir or fpirit of vitriol. This both improves the medicine, and takes off the naufeous tafte. In cafes where

* Arfenic has of late been recommended as an infallible remedy in the ague; but I would advife that it fhould be used only under the eye of a physician. the bark cannot be administered, the *faline mixture* may be given with advantage to children *.

Wine-whey is a very proper drink for a child in an ague; to half an English pint of which may be put a tea-spoonful of the spirit of hartshorn. Exercise is likewise of confiderable service; and when the disease proves obstinate, the child ought, if possible, to be removed to a warm dry air. The food ought to be nourishing, and sometimes a little generous wine should be allowed.

To children, and fuch as cannot fwallow the bark, or when the ftomach will not bear it, it may be given by clyfter. Half an ounce of the extract of bark, diffolved in four ounces of warm water, with the addition of half an ounce of fweet oil, and fix or eight drops of laudanum, is the form recommended by Dr. Lind for an adult, and this to be repeated every fourth hour, or oftener, as the occafion fhall require. For children the quantity of extract and laudanum mult be proportionably leffened. Children have been cured of agues by making them wear a waiftcoat with powdered bark quilted between the folds of it : by bathing them frequently in a ftrong decoction of the bark, and by rubbing the fpine with ftrong fpirits, or with a mixture of equal parts of laudanum and the faponaceous liniment.

We have been the more full upon this difeafe, becaufe it is very common, and becaufe few patients in an ague apply to phyficians unlefs in extremities. There are, however, many cafes in which the difeafe is very irregular, being complicated with other difeafes, or attended with fymptoms which are both very dangerous and very difficult to underftand. All thefe we have purpofely paffed over, as they would only bewilder the generality of readers. When the difeafe is very irregular, or the fymptoms dangerous, the patient ought immediately to apply to a phyfician, and ftrictly to follow his advice.

To prevent agues, people must endeavour to avoid their causes. These have been already pointed out in the beginning of this section : we shall therefore only add one preventive medicine, which may be of use to such

* See Appendix, Saline Mixture.

961

149

as are obliged to live in low marshy countries, or who are liable to frequent attacks of this diseafe.

Take an ounce of the beft Peruvian bark; Virginian fnake-root, and orange peel, of each half an ounce; bruife them all together, and infufe for five or fix days in a bottle of brandy, Holland gin, or any good fpirit; afterwardspour off the clear liquor, and take a wine-glafs of it twice or thrice a-day. This indeed is recommending a dram; but the bitter ingredients in a great meafure take off the ill effects of the fpirit. Thofe who do not chufe it in brandy, may infufe it in wine; and fuch as can bring themfelves to chew the bark, will find that method fucceed very well. Gentian-root, or calamus aromaticus, may alfo be chewed by turns for the purpofe. All bitters feem to be antidotes to agues, efpecially thofe that are warm and aftringent.

In the directions I gave with regard to regimen, I forgot to observe, that change of air cannot be too strongly recommended. Without this, all the efforts of medical skill are sometimes exerted in vain.

The confidence which many people are fill weak enough to place in the most whimfical pretensions to cure agues, renders it neceffary to enforce with farther argument my former caution against fuch filly, and often very dangerous, experiments. I do not fpeak merely of the deceptions of quackery, which are practifed in these complaints, as well as in all others, but of more imposing specifics handed down from parents to their children with circumstantial records of the cures they performed, after the advice of the most eminent men of the faculty had been followed in vain. The accounts given of fuch cures by perfons wholly ignorant of phyfic, are not entitled to the least regard. I do not question their veracity, as far as their knowledge extends; but what can they fay more, than that the fits ceafed after taking the pretended remedy ? How do they know, whether that was the effect of its operation or not; and, if it was, whether, in ftopping the fits, their wonder-working noftrum may not have vitiated the humours, laid the foundation of some other difease, or totally destroyed the

the conflitution? Ought the evidence of fuch people to have any weight in medical experiments? If their affertions are not falle, most of them are palpably abfurd; and the testimony of all mankind cannot prove the truth of an abfurdity.

CHAP. XV.

OF AN ACUTE CONTINUAL FEVER.

THIS fever is denominated acute, ardent, or inflammatory. It most commonly attacks the young, or perfons about the prime and vigour of life, especially fuch as live high, abound with blood, and whose fibres are strong and elastic. It seizes people at all seafons of the year; but is most frequent in the spring and beginning of summer.

CAUSES.—An ardent fever may be occafioned by any thing that overheats the body, or produces plethora, as violent exercife, fleeping in the fun, drinking ftrong liquors, eating fpiceries, a full diet, with little exercife, &c. It may likewife be occafioned by whatever obftructs the perfpiration, as lying on the damp ground, drinking cold liquor when the body is hot, night-watching, or the like.

SYMPTOMS.——A rigour or chillinefs generally ufhers in this fever, which is foon fucceeded by great heat, a frequent and full pulfe, pain of the head, dry fkin, rednefs of the eyes, a florid countenance, pains in the back, loins, &c. To thefe fucceed difficulty of breathing, ficknefs, with an inclination to vomit. The patient complains of great thirft, has no appetite for folid food, is reftlefs, and his tongue generally appears black and rough.

A delirium, exceffive reftleffnefs, great oppreffion of the breaft, with laborious refpiration, ftarting of the tendons, hiccup, cold clammy fweats, and an involuntary difcharge of urine, are very dangerous fymptoms.

As this difeafe is always attended with danger, the L 4 best

beft medical affiftance ought to be procured as foon as poffible. A phyfician may be of ufe at the beginning, but his fkill is often of no avail afterwards. Nothing can be more unaccountable than the conduct of those who have it in their power, at the beginning of a fever, to procure the beft medical affiftance, yet put it off till things come to an extremity. When the difeafe, by delay or wrong treatment, has become incurable, and has exhausted the strength of the patient, it is vain to hope for relief from medicine. Physicians may indeed affift Nature; but their attempts must ever prove fruitlefs, when she is no longer able to co-operate with their endeavours.

REGIMEN.—From the fymptoms of this difeafe, it is evident, that the blood and other humours require to be attenuated; that the perfpiration, urine, faliva, and all the other fecretions, are in too fmall quantity; that the veffels are rigid, and the heat of the whole body too great: all these clearly point out the necessary of a regimen calculated to dilute the blood, correct the acrimony of the humours, allay the exceflive heat, remove the spafmodic fricture of the veffels, and promote the fecretions.

Thefe important purpofes may be greatly promoted by drinking plentifully of diluting liquors; as watergruel, or oatmeal-tea, clear whey, barley-water, balmtea, apple-tea, &c. Thefe may be fharpened with juice of orange, jelly of currants, rafpberries, and fuch like : orange-whey is likewife an excellent cooling drink. It is made by boiling among milk and water a bitter orange fliced, till the curd feparates. If no orange can be had, a lemon, a little cream of tartar, or a few fpoonfuls of vinegar, will have the fame effect. Two or three fpoonfuls of white wine may occafionally be added to the liquor when boiling.

If the patient be coffive, an ounce of tamarinds, with two ounces of ftoned raifins of the fun, and a couple of figs, may be boiled in three English pints of water to a quart. This makes a very pleasant drink and may be used at differentiation. The common pectoral decoction is like-

likewife a very proper drink in this difeafe. A tea-cup full of it may be taken every two hours, or oftener, if the patient's heat and thirft be very great *.

The above liquids must all be drank a little warm. They may be used in fmaller quantities at the beginning of a fever, but more freely atterwards, in order to affist in carrying off the disease by promoting the different excretions. We have mentioned a variety of drinks, that the patient may have it in his power to choose those which are most agreeable, and that, when tired of one, he may have recourse to another.

The patient's diet must be very fpare and light. All forts of flesh-meats, and even chicken-broths, are to be avoided. He may be allowed groat-gruel, panado, or light bread boiled in water; to which may be added a few grains of common falt, and a little fugar, which will render it more palatable. He may eat roasted apples with a little fugar, toasted bread with jelly of currants, boiled prunes, &c.

It will greatly relieve the patient, efpecially in an hot feafon, to have fresh air frequently let into his chamber. This, however, must always be done in such a manner as not to endanger his catching cold.

It is too common in fevers to load the patient with bed-clothes, under the pretence of making him fweat, or defending him from the cold. This cultom has many ill effects. It increases the heat of the body, fatigues the patient, and retards, instead of promoting, the perfpiration.

-

Sitting upright in bed, if the patient be able to bear it, will often have a good effect. It relieves the head, by retarding the motion of the blood to the brain. But this pofture ought never to be continued too long : and if the patient be inclined to fweat, it will be more fafe to let him lie, only raifing his head a little with pillows.

Sprinkling the chamber with vinegar, juice of lemon, or vinegar and rofe-water, with a little nitre diffolved in it, will greatly refresh the patient. This ought to be done frequently, especially if the weather be hot.

* See Appendix, Pettoral Decotion.

The patient's mouth should be often washed with a mixture of water and honey, to which a little vinegar may be added, or with a decoction of figs in barleywater. His feet and hands ought likewife frequently to be bathed in lukewarm water; especially if the head be affected.

The patient should be kept as quiet and easy as polfible. Company, noife, and every thing that diffurbs the mind, is hurtful. Even too much light, or any thing that affects the fenses, ought to be avoided. His attendants fhould be as few as poffible, and they ought not to be too often changed. His inclinations ought rather to be foothed than contradicted; even the promife of what he craves will often fatisfy him as much as its reality.

MEDICINE. ---- In this and all other fevers, attended with a hard, full, quick pulfe, bleeding is of the greatest importance. This operation ought always to be performed as foon as the fymptoms of an inflammatory fever appear. The quantity of blood to be taken away, however, must be in proportion to the strength of the patient and the violence of the difeafe. If after the first bleeding the fever fhould increase, and the pulse become more frequent and hard, there will be a necessity for repeating it a fecond, and perhaps a third, or even a fourth time, which may be done at the diftance of twelve, eighteen, or twenty-four hours from each other, as the fymptoms require. If the pulle continue foft, and the patient be tolerably eafy after the first bleeding, it ought not to be repeated.

If the heat and fever be very great, forty or fifty drops of the dulcified or fweet spirit of nitre may be made into a draught, with an ounce of rofe-water, two ounces of common water, and half an ounce of fimple fyrup, or a bit of loaf-fugar. This draught may be given to the patient every three or four hours while the fever is violent; afterwards once in five or fix hours will be fufficient.

If the patient be afflicted with reaching, or an inclination to vomit, it will be right to affift Nature's attempts, by

by giving him weak camomile-tea, or lukewarm water to drink.

If the body be bound, a clyfter of milk and water, with a little falt, and a fpoonful of fweet oil or fresh butter in it, ought daily to be administered. Should this not have the defired effect, a tea-spoonful of magness alba, or cream of tartar, may be frequently put into his drink. He may likewise eat tamarinds, boiled prunes, roasted apples, and the like.

If about the tenth, eleventh, or twelfth day, the pulfe become more foft, the tongue moifter, and the urine begins to let fall a reddifh fediment, there is reafon to expect a favourable iffue to the difeafe. But if, inftead of thefe fymptoms, the patient's fpirits grow languid, his pulfe finks, and his breathing becomes difficult; with a flupor, trembling of the nerves, flarting of the tendons, &cc. there is reafon to fear that the confequences will be fatal. In this cafe bliftering-plafters mult be applied to the head, ancles, infide of the legs or thighs, as there may be occafion; poultices of wheat-bread, muftard, and vinegar, may likewife be applied to the foles of the feet, and the patient muft be fupported with cordials, as ftrong wine-whey, negus, fago-gruel, with wine in it, and fuch like.

A proper regimen is not only neceffary during the fever, but likewife after the patient begins to recover. By neglecting this, many relapfe, or fall into other difeafes, and continue valetudinary for life. Though the body be weak after a fever, yet the diet for fome time ought to be rather light than of too nourifhing a nature. Too much food, drink, exercife, company, &c. are carefully to be avoided. The mind ought likewife to be kept eafy, and the patient fhould not attempt to purfue ftudy, or any bufinefs that requires intenfe thinking.

If the digeftion be bad, or the patient be feized at times with feverifh heats, an infusion of Peruvian bark in cold water will be of use. It will strengthen the stomach, and help to subdue the remains of the fever.

When the patient's ftrength is pretty well recovered, he ought to take fome gentle laxative. An ounce of tamarinds and a dram of fenna may be boiled for a few minutes

minutes in an English pint of water, and an ounce of manna diffolved in the decoction; afterwards it may be strained, and a tea-cupful drank every hour till it operates. This dose may be repeated twice or thrice, five or fix days intervening.

Those who follow laborious employments ought not to return too foon to their labour after a fever, but should keep easy till their strength and spirits are sufficiently recruited.

It requires very little argument to prove, that the body as well as the mind muft require indulgence after the leverity of fuch a difeafe. But I find it more difficult to prevent people from carrying this indulgence to excefs in what relates to eating and drinking. The appetite is ufually voracious upon recovering from moft fevers; and to fay, that its cravings are not to be fatisfied, is certainly an unpalatable doctrine. Yet felf-command is neceffary in fuch cafes, as there will be great danger, not only of a relapfe, but of other difagreeable confequences, fuch as boils, ulcers, and fettled fwellings in particular limbs. Thefe may be obviated by a light and principally vegetable diet, not however totally excluding animal food of eafy digeftion.

Though I have taken much pains to convince people of the propriety of getting the beft medical affiftance they can, upon the first attacks of a fever, before it becomes, by delay or wrong treatment, incurable, yet the number is aftonishing of those who are the victims of their own fatal neglect in this particular. Some, under a pretence of trufting to the efforts of nature, but in reality too conceited of their own strength, and too felfwilled to take advice, endeavour to keep upon their legs, as they term it, and to struggle with the diseafe as long as they can. Its violence is increased by this very attempt. The bed alone would in many cases stop a beginning fever, the posture contributing to relax the stop fast of the ardor of the circulation.

Others purfue a very opposite, but no less reprehenfible method. On the first alarm, they have recourse to the most pernicious means of exciting sweats by taking hot and volatile sudorific medicines; shutting out the

the air from all poffible admiffion into their chambers, and fmothering themfelves under enormous loads of bed-clothes. The heat and motion of the blood, already too violent, are thus increafed; fuel is added to the fire; and fweating is in reality prevented; for the higher the fever, the lefs copious will be the evacuations of every kind. Perfpiration is beft promoted by a proper quantity of diluting liquids, which at the fame time quench the patient's thirft, and abate the pains of the breaft and difficulty of breathing.

I can do no more than fairly flate the confequences of fuch errors. The ways followed by those two defcriptions of people, however different, terminate in the fame point, and that is the grave. Fevers make a dreadful havoc among the human race; but their ravages are confiderably increased by the misconduct and perverseness of the unfortunate fufferers themselves.

CHAP. XVI.

a violent pricking pate in one of the fides

OF THE PLEURISY.

T HE true pleurify is an inflammation of that membrane called the *pleura*, which lines the infide of the breaft. It is diffinguifhed into the moift and dry. In the former, the patient fpits freely; in the latter, little or none at all. There is likewife a fpecies of this difeafe, which is called the *fpurious* or *baftard pleurify*, in which the pain is more external, and chiefly affects the muscles between the ribs. The pleurify prevails among labouring people, especially such as work without doors, and are of a fanguine constitution. It is most frequent in the fpring feason.

CAUSES.——The pleurify may be occafioned by whatever obfiructs the perfpiration; as cold northerly winds; drinking cold liquors when the body is hot; fleeping without doors on the damp ground; wet clothes; plunging

plunging the body into cold water, or exposing it to the cold air, when covered with fweat, &c. It may likewife be occafioned by drinking ftrong liquors; by the stoppage of usual evacuations; as old ulcers, iffues, fweating of the feet or hands, &c. the fudden ftriking in of any eruption, as the itch, the measles, or the fmall-pox. Those who have been accustomed to bleed at a certain feafon of the year, are apt, if they neglect it, to be feized with a pleurify. Keeping the body too warm by means of fire, clothes, &c. renders it more liable to this difeafe. A pleurify may likewife be occafioned by violent exercife, as running, wreftling, leaping, or by fupporting great weight, blows on the breaft, &c. A bad conformation of the body renders perfons more liable to this difeafe, as a narrow cheft, a straitnefs of the arteries of the pleura, &c.

SYMPTOMS.——This, like moft other fevers, generally begins with chillinefs and fhivering, which are followed by heat, thirft, and reftleffnefs. To thefe fucceeds a violent pricking pain in one of the fides among the ribs. Sometimes the pain extends towards the back-bone, fometimes towards the forepart of the breaft, and at other times towards the fhoulder blades. The pain is generally moft violent when the patient draws his breath.

The pulfe in this difeafe is commonly quick and hard, the urine high-coloured; and, if blood be let, it is covered with a tough cruft, or buffy coat. The patient's fpittle is at first thin, but afterwards it becomes groffer, and is often streaked with blood.

REGIMEN.——Nature generally endeavours to carry off this difeafe by a critical difcharge of blood from fome part of the body, by expectoration, fweat, loofe ftools, thick urine, or the like. We ought therefore to fecond her intentions by leffening the force of the circulation, relaxing the veffels, diluting the humours, and promoting expectoration.

For these purposes, the diet, as in the former disease, ought to be cool, slender, and diluting. The patient must

158

must avoid all food that is viscid, hard of digestion, or that affords much nourishment; as flesh, butter, cheefe, eggs, milk, and also every thing that is of a heating nature. His drink may be whey, or an infusion of pectoral and balfamic vegetables *.

Barley-water, with a little honey or jelly of currants mixed with it, is likewife a very proper drink in this difeafe. It is made by boiling an ounce of pearl-barley in three Englifh pints of water to two, which muft afterwards be ftrained. The decoction of figs, raifins, and barley, recommended in the preceding difeafe, is here likewife very proper. Thefe and other diluting liquors are not to be drank in large quantities at a time; but the patient ought to keep continually fipping them, fo as to render his mouth and throat always moift. All his food and drink fhould be taken a little warm.

The patient fhould be kept quiet, cool, and every way eafy, as directed under the foregoing difeafe. His feet and hands ought daily to be bathed in lukewarm water; and he may fometimes fit up in bed for a fhort fpace, in order to relieve his head.

MEDICINE.——Almost every perfon knows, when a fever is attended with a violent pain of the fide, and a quick hard pulfe, that bleeding is neceffary. When these fymptoms come on, the sooner this operation is performed the better; and the quantity at first must be pretty large, provided the patient be able to bear it. A large quantity of blood let at once in the beginning of a pleurify, has a much better effect than repeated small bleedings. A man may lose twelve or fourteen ounces of blood as soon as it is certainly known that he is feized with a pleurify. For a younger perfon, or one of a delicate confliction, the quantity must be lefs.

If, after the first bleeding, the stitch, with the other violent symptoms, should still continue, it will be neceffary, at the distance of twelve or eighteen hours, to let eight or nine ounces more. If the symptoms do not then abate, and the blood shews a strong buffy coat, a

* See Appendix, Pettoral Infusion.

OF THE PLEURISY.

third or even a fourth bleeding may be requifite. If the pain of the fide abate, the pulfe become fofter, or the patient begin to fpit freely, bleeding ought not to be repeated. This operation is feldom receffary after the third or fourth day of the fever, and ought not then to be performed, unlefs in the most urgent circumstances.

The blood may be many ways attenuated without bleeding. There are likewife many things that may be done to eafe the pain of the fide without this operation, as fomenting, bliftering, &c. Fomentations may be made by boiling a handful of flowers of elder, camomile, and common mallows, or any other foft vegetables in a proper quantity of water. The herbs may be either put into a flannel bag, and applied warm to the fide, or flannels may be dipped in the decoction, afterwards wrung out, and applied to the part affected, with as much warmth as the patient can eafily bear. As the clothes grow cool, they must be changed, and great care taken that the patient do not catch cold. A bladder may be filled with warm milk and water, and applied to thefide, if the above method of fomenting be found inconvenient. Fomentations not only eafe the pain, but relax the veffels, and prevent the flagnation of the blood and other humours. The fide may likewife be frequently rubbed with a little of the volatile liniment *.

Topical bleeding has often a very good effect in this difeafe. It may either be performed by applying a number of leeches to the part affected, or by cupping, which is both a more certain and expeditious method than the other.

Leaves of various plants might likewife be applied to the patient's fide with advantage. I have often feen great benefit from young cabbage leaves applied warm to the fide in a pleurify. Thefe not only relax the parts, but likewife draw off a little moifture, and may prevent the neceffity of bliftering-plafters; which, however, when other things fail, muft be applied.

If the flitch continue after repeated bleedings, fomentations, &c. a bliftering-plafter must be applied over the

part

^{*} See Appendix, Volatile Liniment.

part affected, and fuffered to remain for two days. This not only procures a difcharge from the fide, but takes off the fpafm, and by that means affifts in removing the caufe of the difeafe. To prevent a ftrangury when the bliftering-plafter is on, the patient may drink freely of the Arabic emulfion *.

If the patient be coftive, a clyfter of thin water-gruel, or of barley-water, in which a handful of mallows, or any other emollient vegetable, has been boiled, may be daily administered. This will not only empty the bowels, but have the effect of a warm fomentation applied to the inferior vifcera, which will help to make a derivation from the breaft.

The expectoration may be promoted by fharp, oily, and mucilaginous medicines. For this purpole, an ounce of the oxymel, or the vinegar of fquills, may be added to fix ounces of the pectoral decoction, and two table-fpoonfuls of it taken every two hours.

Should the fquill difagree with the ftomach, the oily emulfion may be administered \dagger ; or, in place of it, two ounces of the oil of fweet almonds, or oil of olives, and two ounces of the fyrup of violets, may be mixed with as much fugar-candy powdered as will make an electuary of the confistence of honey. The patient may take a tea-fpoonful of this frequently, when the cough is troublefome. Should oily medicines prove naufeous, which is fometimes the cafe, two table-fpoonfuls of the folution of gum ammoniac in barley-water may be given three or four times a-day \ddagger .

If the patient does not perfpire, but has a burning heat upon his fkin, and paffes very little water, fome fmall dofes of purified nitre and camphire will be of ufe. Two drams of the former may be rubbed with five or fix grains of the latter in a mortar, and the whole divided into fix dofes, one of which may be taken every five or fix hours, in a little of the patient's ordinary drink.

We shall only mention one medicine more, which fome reckon almost a specific in the pleurify, viz. the de-

* See Appendix, Arabic Emulfion.

+ See Appendix, Oily Emulfion.

1 See Appendix, Solution of Gum Ammoniac.

coction

OF THE PLEURISY.

coction of the feneka rattle-fnake root *. After bleeding and other evacuations have been premifed, the patient may take two, three, or four table-fpoonfuls of this decoction, according as his ftomach will bear it, three or four times a-day. If it fhould occafion vomiting, two or three ounces of fimple cinnamon-water may be mixed with the quantity of decoction here directed; or it may be taken in fmaller dofes. As this medicine promotes perfpiration and urine, and likewife keeps the body eafy, it may be of fome fervice in a pleurify, or any other inflammation of the breaft.

No one will imagine that these medicines are all to be used at the fame time. We have mentioned different things, on purpose that people may have it in their power to chuse; and likewise, that when one cannot be obtained, they may make use of another. Different medicines are no doubt necessary in the different periods of a diforder; and where one fails of success, or difagrees with the patient, it will be proper to try another.

What is called the crifis, or height of the fever, is fometimes attended with very alarming fymptoms, as difficulty of breathing, an irregular pulfe, convulfive motions, &c. Thefe are apt to frighten the attendants, and induce them to do improper things, as bleeding the patient, giving him ftrong ftimulating medicines, or the like. But they are only the ftruggles of Nature to overcome the difeafe, in which fhe ought to be affifted by plenty of diluting drink, which is then peculiarly neceffary. If the patient's ftrength, however, be much exhaufted by the difeafe, it will be neceffary at this time to fupport him with frequent fmall draughts of winewhey, negus, or the like.

When the pain and fever are gone, it will be proper, after the patient has recovered fufficient ftrength, to give him fome gentle purges, as those directed towards the end of an acute continual fever. He ought likewise to use a light diet of easy digestion, and his drink should be butter-milk, whey, and other things of a cleansing nature.

* See Appendix, DecoSion of Seneka Root.

Of

OF THE PLEURISY.

Of the BASTARD PLEURISY.

That fpecies of pleurify which is called the *bastard* or *fpurious*, generally goes off by keeping warm for a few days, drinking plenty of diluting liquors, and observing a cooling regimen.

It is known by a dry cough, a quick pulfe, and a difficulty of lying on the affected fide; which laft does not always happen in the true pleurify. Sometimes, indeed, this difeafe proves obftinate, and requires bleeding, with cupping, and fcarifications of the part affected. Thefe, together with the ufe of nitrous and other cooling medicines, feldom fail to effect a cure. Bliftering is often ufeful in this difeafe.

Of the PARAPHRENITIS.

The *paraphrenitis*, or inflammation of the diaphragm, is fo nearly connected with the pleurify, and refembles it fo much in the manner of treatment, that it is fearcely neceffary to confider it as a feparate difeafe.

It is attended with a very acute fever, and an extreme pain in the part affected, which is generally augmented by coughing, fneezing, drawing in the breath, taking food, going to ftool, making water, &c. Hence the patient breathes quick, and draws in his bowels to prevent the motion of the diaphragm; is reftlefs, anxious, has a dry cough, a hiccup, and often a delirium. A convultive laugh, or rather a kind of involuntary grin, is no uncommon fymptom of this difeafe.

Every method must be taken to prevent a suppuration, as it is impossible to fave the patient's life when this happens. The regimen and medicine are in all respects the same as in the pleurify. We shall only add, that in this difease, emollient clysters are peculiarly useful, as they relax the bowels, and by that means make a derivation from the part affected.

CHAP.

[164]

CHAP. XVII.

OF A PERIPNEUMONY, OR INFLAMMA-TION OF THE LUNGS.

A S this difeafe affects an organ which is abfolutely neceffary to life, it must always be attended with danger. Perfons who abound with thick blood, whofe fibres are tenfe and rigid, who feed upon grofs aliment, and drink strong viscid liquors, are most liable to a peripneumony. It is generally fatal to those who have a flat breast, or narrow chest, and to such as are afflicted with an asthma, especially in the decline of life. Sometimes the inflammation reaches to one lobe of the lungs only, at other times the whole of the organ is affected; in which case the discase can hardly fail to prove fatal.

When the difeafe proceeds from a vifcid pituitous matter obstructing the veffels of the lungs, it is called a *fpurious* or *bastard peripneumony*. When it arifes from a thin acrid defluction on the lungs, it is denominated a *catarrhal peripneumony*, &c.

CAUSES.—An inflammation of the lungs is fometimes a primary difeafe, and fometimes it is the confequence of other difeafes, as a quinfey, a pleurify, &c. It proceeds from the fame caufes as the pleurify, viz. an obftructed perfpiration from cold, wet clothes, &c. or from an increafed circulation of the blood by violent exercife, the ufe of fpiceries, ardent fpirits, and fuch like. The pleurify and peripneumony are often complicated; in which cafe the difeafe is called a *pleuroperipneumony*.

SYMPTOMS. — Most of the symptoms of a pleurify likewife attend an inflammation of the lungs; only in the latter the pulse is more fost, and the pain less acute; but the difficulty of breathing, and oppression of the breast, are generally greater.

REGIMEN.——As the regimen and medicine are in all refpects the fame in the true peripneumony as in the

OF A PERIPNEUMONY.

the pleurify, we fhall not here repeat them, but refer the reader to the treatment of that difeafe. It may not, however, be improper to add, that the aliment ought to be more flender and thin in this than in any other inflammatory difeafe. The learned Dr. Arbuthnot afferts, that even common whey is fufficient to fupport the patient, and that decoctions of barley, and infufions of fennel roots in warm water with milk, are the most proper both for drink and nourifhment. He likewife recommends the steam of warm water taken in by the breath, which ferves as a kind of internal fomentation, and helps to attenuate the impacted humours. If the patient have loofe stopped, but rather promoted by the use of emollient clysters.

It has already been observed, that the *fpurious* or *baftard* peripneumony is occasioned by a viscid pituitous matter obstructing the vessels of the lungs. It commonly attacks the old, infirm, and phlegmatic, in winter and wet feasons.

The patient at the beginning is cold and hot by turns, has a fniall quick pulle, feels a fenfe of weight upon his breaft, breathes with difficulty, and fometimes complains of a pain and giddinefs of his head. His urine is ufually pale, and his colour very little changed.

The diet, in this as well as in the true peripneumony, must be very flender, as weak broths, fharpened with the juice of orange or lemon, and fuch like. His drink may be thin water-gruel fweetened with honey, or a decoction of the roots of fennel, liquorice, and quick grafs. An ounce of each of these may be boiled in three English pints of water to a quart, and sharpened with a little currant-jelly, or the like.

Bleeding and purging are generally proper at the beginning of this difeafe; but if the patient's fpittle be pretty thick, or well concocted, neither of them are neceffary. It will be fufficient to affift the expectoration by fome of the fharp medicines recommended for that purpofe in the pleurify, as the folution of gum-ammoniac with oxymel of fquills, &c. Bliftering-plafters have generally a good effect, and ought to be applied pretty early.

M 3

If

165

If the patient do not fpit, he must be bled, according as his strength will permit, and have a gentle purge administered. Afterwards his body may be kept open by clysters, and the expectoration promoted, by taking every four hours two table-spoonfuls of the solution mentioned above.

When an inflammation of the breaft does not yield to bleeding, bliftering, and other evacuations, it commonly ends in fuppuration, which is more or lefs dangerous, according to the part where it is fituated. When this happens in the pleura, it fometimes breaks outwardly, and the matter is difcharged by the wound.

When the fuppuration happens within the fubftance or body of the lungs, the matter may be difcharged by expectoration; but if the matter floats in the cavity of the breaft, between the pleura and the lungs, it can only be difcharged by an incifion made betwixt the ribs.

If the patient's ftrength do not return after the inflammation is to all appearance removed; if his pulfe continue quick though foft, his breathing difficult and oppreffed; if he have cold fhiverings at times, his cheeks flufhed, his lips dry; and if he complain of thirft, and want of appetite, there is reafon to fear a fuppuration, and that a phthifis or confumption of the lungs will enfue. We fhall therefore next proceed to confider the proper treatment of that difeafe.

CHAP. XVIII.

OF CONSUMPTIONS.

A CONSUMPTION is a wasting or decay of the whole body, from an ulcer, tubercles, or concretion of the lungs, an empyema, a nervous atrophy, or cachexy.

Dr. Arbuthnot obferves, that in his time confumptions made up above one-tenth part of the bills of mortality in and about London. There is reafon to believe they have

166

have rather increafed fince; and we know from experience, that they are not lefs fatal in fome other towns of England than in London.

Young perfons, between the age of fifteen and thirty, of a flender make, long neck, high fhoulders, and flat breafts, are most liable to this difease.

Confumptions prevail more in England than in any other part of the world, owing perhaps to the great ufe of animal food and malt liquors, the general application to fedentary employments, and the great quantity of pit-coal which is there burnt; to which we may add, the perpetual changes in the atmosphere, or variableness of the weather.

CAUSES.——It has already been obferved, that an inflammation of the breaft often ends in an impofthume : confequently whatever difpofes people to this difeafe mult likewife be confidered as a caufe of confumption.

Other difeafes, by vitiating the habit, may likewife occasion confumptions; as the fcurvy, the fcrophula, or king's-evil, the venereal difeafe, the afthma, fmallpox, measles, &c.

As this difeafe is feldom cured, we fhall endeavour the more particularly to point out its caufes, in order that people may be enabled to avoid it. Thefe are:

---- Confined or unwholefome air; when this fluid is impregnated with the fumes of metals or minerals, it proves extremely hurtful to the lungs, and often corrodes the tender veffels of that neceflary organ.

Great evacuations; as fweating, diarrhœas, diabetes, exceflive venery, the fluor albus, an over-difcharge of the menftrual flux, giving fuck too long, &c.

The fudden stoppage of customary evacuations; as the bleeding piles, sweating of the feet, bleeding at the nose, the menses, issues, ulcers, or eruptions of any kind.

faw the fymptoms of a phthifis occafioned by a fmall bone flicking in the bronchæ. It was afterwards vomited

along

along with a confiderable quantity of purulent matter, and the patient, by a proper regimen, and the use of the Peruvian bark, recovered.

-----Frequent and exceffive debaucheries. Late watching, and drinking ftrong liquors, which generally go together, can hardly fail to deftroy the lungs. Hence the *bon companion* generally falls a facrifice to this difeafe.

——Occupations in life. Those artificers who fit much, and are constantly leaning forward, or prefling upon the stomach and breast, as cutlers, taylors, shoemakers, seamstress, &c. often die of consumptions. They likewise prove fatal to singers, and all who have occasion to make frequent and violent exertions of the lungs.

Sharp, faline, and aromatic aliments, which heat and inflame the blood, are likewife frequently the caufe of confumptions.

We shall only add, that this difease is often owing to an hereditary taint, or a scrophulous habit; in which case it is generally incurable.

SYMPTOMS.—This difeafe generally begins with a dry cough, which often continues for fome months. If a difpofition to vomit after eating be excited by it, there is ftill greater reafon to fear an approaching confumption. The patient complains of a more than ufual degree of heat, a pain and oppreffion of the breaft, efpecially after motion; his fpittle is of a faltifh tafte, and fometimes mixed with blood. He is apt to be fad; his appetite is bad, and his thirft great. There is generally a quick,

OF CONSUMPTIONS.

a quick, foft, fmall pulfe; though fometimes the pulfe is pretty full, and rather hard. Thefe are the common fymptoms of a beginning confumption.

Afterwards the patient begins to fpit a greenifh, white, or bloody matter. His body is extenuated by the hectic fever and colliquative fweats, which mutually fucceed one another, viz. the one towards night, and the other in the morning. A loofenefs, and an exceffive difcharge of urine, are often troublefome fymptoms at this time, and greatly weaken the patient. There is a burning heat in the palms of the hands, and the face generally flufhes after eating; the fingers become remarkably fmall, the nails are bent inwards, and the hairs fall off.

At last the fwelling of the feet and legs, the total loss of ftrength, the finking of the eyes, the difficulty of fwallowing, and the coldness of the extremities, shew the immediate approach of death, which, however, the patient feldom believes to be fo near. Such is the usual progress of this fatal difease, which, if not early checked, commonly sets all medicine at defiance.

REGIMEN. — On the first appearance of a confumption, if the patient live in a large town, or any place where the air is confined, he ought immediately to quit it, and to make choice of a fituation in the country, where the air is pure and free. Here he must not remain inactive, but take every day as much exercise as he can bear.

The beft method of taking exercife is to ride on horfeback, as this gives the body a great deal of motion without much fatigue. Such as cannot bear this kind of exercife, muft make use of a carriage. A long journey, as it amuses the mind by a continual change of objects, is greatly preferable to riding the same ground over and over. Care, however, must be taken to avoid catching cold from wet clothes, damp beds, or the like. The patient ought always to finish his ride in the morning, or at least before dinner; otherwise it will oftener do harm than good.

It is pity those who attend the fick feldom recommend riding in this disease, till the patient is either unable to bear it, or the malady has become incurable. Patients are likewife apt to trifle with every thing that is in their own power. They cannot fee how one of the common actions of life fhould prove a remedy in an obfinate difeafe, and therefore they reject it, while they greedily hunt after relief from medicine, merely becaufe they do not understand it.

Those who have strength and courage to undertake a pretty long voyage, may expect great advantage from it. This to my knowledge has frequently cured a confumption after the patient was, to all appearance, far advanced in that difease, and where medicine had proved ineffectual. Hence it is reasonable to conclude, that if a voyage were undertaken in due time, it would feldom fail to perform a cure *.

Such as try this method of cure ought to carry as much frefh provisions along with them as will ferve for the whole time they are at fea. As milk is not eafily obtained in this fituation, they ought to live upon fruits, and the broth of chickens, or other young animals which can be kept alive on board. It is fearcely neceffary to add, that fuch voyages shall be undertaken, if possible, in the mildest feason, and that they ought to be towards a warmer climate \dagger .

Those who have not courage for a long voyage may travel into a more fouthern climate, as the fouth of France, Spain, or Portugal; and if they find the air of these countries agree with them, they should continue there at least till their health be confirmed.

Next to proper air and exercife, we would recommend a due attention to diet. The patient fhould eat

* Two things chiefly operate to prevent the benefits which would arife from failing. The one is, that phyficians feldom order it till the difeafe is too far advanced; and the other is, that they feldom order a voyage of a fufficient length. A patient may receive no benefit by croffing the channel, who, fhould he crofs the Atlantic, might be completely cured. Indeed we have reafon to believe, that a voyage of this kind, if taken in due time, would feldom fail to cure a confumption.

+ Though I do not remember to have feen one inftance of a genuine confumption of the lungs cured by medicine, yet I have known a Weft-India voyage work wonders in that dreadful diforder.

OF CONSUMPTIONS.

nothing that is either heating or hard of digeftion, and his drink muft be of a foft and cooling nature. All the diet ought to be calculated to leffen the acrimony of the humours, and to nourifh and fupport the patient. For this purpofe he muft keep chiefly to the use of vegetables and milk. Milk alone is of more value in this difease than the whole *materia medica*.

Affes' milk is commonly reckoned preferable to any other; but it cannot always be obtained; befides, it is generally taken in a very fmall quantity; whereas, to produce any effects, it ought to make a confiderable part of the patient's diet. It is hardly to be expected, that a gill or two of affes' milk, drank in the fpace of twenty-four hours, fhould be able to produce any confiderable change in the humours of an adult; and when people do not perceive its effects foon, they lofe hope, and fo leave it off. Hence it happens, that this medicine, however valuable, very feldom performs a cure. The reafon is obvious; it is commonly ufed too late, is taken in too fmall quantities, and is not duly perfifted in.

I have known very extraordinary effects from affes' milk in obftinate coughs, which threatened a confumption of the lungs; and do verily believe, if used at this period, that it would feldom fail; but if it be delayed till an ulcer is formed, which is generally the cafe, how can it be expected to fucceed ?

Affes' milk ought to be drank, if pofiible, in its natural warmth, and, by a grown perfon, in the quantity of half an Englifh pint at a time. Inftead of taking this quantity night and morning only, the patient ought to take it four times, or at leaft thrice a-day, and to eat a little light bread along with it, fo as to make it a kind of meal.

If the milk fhould happen to purge, it may be mixed with old conferve of rofes. When that cannot be obtained, the powder of crabs' claws may be used in its ftead. Affes' milk is usually ordered to be drank warm in bed; but as it generally throws the patient in a fweat when taken in this way, it would perhaps be better to give it after he rifes.

OF CONSUMPTIONS.

Some extraordinary cures in confumptive cafes have been performed by women's milk. Could this be obtained in fufficient quantity, we would recommend it in preference to any other. It is better if the patient can fuck it from the breaft, than to drink it afterwards. I knew a man who was reduced to fuch a degree of weaknefs in a confumption, as not to be able to turn himfelf in bed. His wife was at that time giving fuck, and the child happening to die, he fucked her breafts, not with a view to reap any advantage from the milk, but to make her eafy. Finding himfelf, however, greatly benefited by it, he continued to fuck her till he became perfectly well, and is at prefent a ftrong and healthy man.

Some prefer butter-milk to any other, and it is indeed a very valuable medicine, if the ftomach be able to bear it. It does not agree with every perfon at firft; and is therefore often laid afide without a fufficient trial. It fhould at firft be taken fparingly, and the quantity gradually increased, until it comes to be almost the sole food. I never knew it fucceed, unless where the patient almost lived upon it.

Cows' milk is most readily obtained of any, and though it be not fo easily digested as that of assor mares, it may be rendered lighter, by adding to it an equal quantity of barley-water, or allowing it to stand for some hours, and asterwards taking off the cream. If it should, notwithstanding, prove heavy on the stomach, a small quantity of brandy or rum, with a little sugar, may be added, which will render it both more light and nourishing.

It is not to be wondered, that milk fhould for fome time difagree with a ftomach that has not been accuftomed to digeft any thing but flefh and ftrong liquors, which is the cafe with many of those who fall into confumptions. We do not, however, advise those who have been accustomed to animal food and strong liquors, to leave them off all at once. This might be dangerous. It will be neceffary for such to eat a little once a-day of the flesh of some young animal, or rather to use the broth made of chickens, veal, lamb, or such like. They ought likewise to drink a little wine made into negus, or

or diluted with twice or thrice its quantity of water, and to make it gradually weaker till they can leave it off altogether.

These must be used only as preparatives to a diet confisting chiefly of milk and vegetables, which the fooner the patient can be brought to bear, the better. Rice and milk, or barley and milk, boiled with a little fugar, is very proper food. Ripe fruits, roafted, baked, or boiled, are likewise proper, as goose or currantberry tarts, apples roafted, or boiled in milk, &c. The jellies, conferves, and preferves, &c. of ripe subacid fruits, ought to be eaten plentifully, as the jelly of currants, conferve of roses, preferved plums, cherries, &c.

Wholefome air, proper exercife, and a diet confifting chiefly of thefe and other vegetables, with milk, is the only courfe that can be 'depended on in a beginning confumption. If the patient has ftrength and fufficient refolution to perfift in this courfe, he will feldom be difappointed of a cure.

In a populous town in England *, where confumptions are very common, I have frequently feen confumptive patients, who had been fent to the country with orders to ride and live upon milk and vegetables, return in a few months quite plump, and free from any complaint. This indeed was not always the cafe, efpecially when the difeafe was hereditary, or far advanced; but it was the only method in which fuccefs was to be expected: where it failed, I never knew medicine fucceed.

If the patient's ftrength and fpirits flag, he must be fupported by ftrong broths, jellies and fuch like. Some recommend shell-fish in this diforder, and with some reason, as they are nourishing and restorative \ddagger . All the food and drink ought, however, to be taken in small quantities, less an overcharge of fresh chyle should opprefs the lungs, and too much accelerate the circulation of the blood.

* Sheffield.

† I have often known perfons of a confumptive habit, where the fymptoms were not violent, reap great benefit from the use of oysters. They generally ate them raw, and drank the juice along with them. The patient's mind ought to be kept as eafy and cheers ful as poffible. Confumptions are often occafioned, and always aggravated, by a melancholy caft of mind; for which reafon mufic, cheerful company, and every thing that infpires mirth, are highly beneficial. The patient ought feldom to be left alone, as brooding over his calamities is fure to render him worfe.

MEDICINE. — Though the cure of this difeafe depends chiefly upon regimen and the patient's own endeavours, yet we shall mention a few things which may be of fervice in relieving fome of the more violent fymptoms.

In the first stage of a confumption, the cough may fometimes be appealed by bleeding; and the expectoration may be promoted by the following medicines: Take fresh squills, gum-ammoniac, and powdered cardamum seeds, of each a quarter of an ounce; beat them together in a mortar, and if the mass prove too hard for pills, a little of any kind of syrup may be added to it. This may be formed into pills of a moderate fize, and four or five of them taken twice or thrice a-day, according as the patient's stomach will bear them.

The *lac ammoniacum*, or milk of gum-ammoniac, as it is called, is likewife a proper medicine in this ftage of the difeafe. It may be used as directed in the pleurify.

A mixture made of equal parts of lemon-juice, fine honey, and fyrup of poppies, may likewife be ufed, Four ounces of each of thefe may be fimmered together in a fauce-pan, over a gentle fire, and a tablefpoonful of it taken at any time when the cough is troublefome.

It is common in this ftage of the difeafe to load the patient's ftomach with oily and balfamic medicines. Thefe, inftead of removing the caufe of the difeafe, tend rather to increafe it by heating the blood, while they pall the appetite, relax the folids, and prove every way hurtful to the patient. Whatever is ufed for removing the cough, befides riding and other proper regimen, ought to be medicines of a fharp and cleanfing nature; as oxymel, fyrup of lemon, &c.

Acids

Acids feem to have peculiarly good effects in this difeafe; they both tend to quench the patient's thirft and to cool the blood. The vegetable acids, as apples, oranges, lemons, &c. appear to be the most proper. I have known patients fuck the juice of feveral lemons every day with manifest advantage, and would for this reason recommend acid vegetables to be taken in as great quantity as the stomach will bear them.

For the patient's drink, we would recommend infufions of the bitter plants, as ground-ivy, the leffer centaury, camomile flowers, or water-trefoil. Thefe infufions may be drank at pleafure. They ftrengthen the ftomach, promote digettion, and at the fame time anfwer all the purpofes of dilution, and quench thirft much better than things that are lufcious or fweet. But if the patient spit blood, he ought to use, for his ordinary drink, infusions or decoctions of the vulnerary roots, plants, &c.*

There are many other mucilaginous plants and feeds, of a healing and agglutinating nature, from which decoctions or infufions may be prepared with the fame intention; as the orches, the quince-feed, coltsfoot, linfeed, farfaparilla, &c. It is not neceffary to mention the different ways in which thefe may be prepared. Simple infufion or boiling is all that is neceffary, and the dofe may be at different.

The conferve of roles is here peculiarly proper. It may either be put into the decoction above prefcribed, or eaten by itfelf. No benefit is to be expected from trifling doles of this medicine. I never knew it of any fervice, unlefs where three or four ounces at leaft were ufed daily for a confiderable time. In this way I have feen it produce very happy effects, and would recommend it wherever there is a difcharge of blood from the lungs.

When the fpitting up of großs matter, oppreffion of the breaft, and the hectic fymptoms, fhew that an impofthume is formed in the lungs, we would recommend the Peruvian bark, that being the only drug which has any

* See Appendix, Vulnerary Decodion.

OF CONSUMPTIONS.

176

chance to counteract the general tendency which the humours then have to putrefaction.

An ounce of the bark in powder may be divided into eighteen or twenty dofes, of which one may be taken every three hours through the day, in a little fyrup, or a cup of horehound tea.

If the bark fhould happen to purge, it may be made into an electuary, with the conferve of rofes, thus: Take old conferve of rofes, a quarter of a pound; Peruvian bark, a quarter of an ounce; fyrup of orange or lemon, as much as will make it of the confiftence of honey. This quantity will ferve the patient four or five days, and may be repeated as there is occafion.

Such as cannot take the bark in fubftance, may infufe it in cold water. This feems to be the beft menftruum for extracting the virtues of that drug. Half an ounce of bark in powder may be infufed for twenty-four hours in half an English pint of water. Afterwards, let it be paffed through a fine strainer, and an ordinary tea-cupful of it taken three or four times a-day.

We would not recommend the bark while there are any fymptoms of an inflammation of the breaft; but when it is certainly known that matter is collecting there, it is one of the beft medicines which can be ufed. Few patients, indeed, have refolution enough to give the bark a fair trial at this period of the difeafe, otherwife we have re fon to believe that fome benefit might be reaped from u.

When it is evident that there is an imposthume in the breast, and the matter can neither be spit up nor carried off by absorption, the patient must endeavour to make it break inwardly, by drawing in the steams of warm water or vinegar with his breath, coughing, laughing, or bawling aloud, &c. When it happens to burst within the lungs, the matter may be discharged by the mouth. Sometimes, indeed, the bursting of the vomica occasions immediate death, by suffocating the patient. When the quantity of matter is great, and the patient's strength exhausted, this is commonly the case. At any rate, the patient is ready to fall into a stoon, and should have volatile falts or spirits held to his note.

OF CONSUMPTIONS.

If the matter difcharged be thick, and the cough and breathing become eafier, there may be fome hopes of a cure. The diet at this time ought to be light, but reftorative, as chicken-broths, fago-gruel, rice-milk, &c. the drink, butter-milk or whey, fweetened with honey This is likewife a proper time for using the Peruvian bark, which may be taken as directed above.

If the vomica or impossible for the breast, between the pleura and the into the cavity of the breast, between the pleura and the lungs, there is no way of getting the matter out, but by an incision, as has already been observed. As this operation must always be performed by a furgeon, it is not neceffary here to describe it. We shall only add, that it is not so dreadful as people are apt to imagine, and that it is the only chance the patient in this case has for his life.

A NERVOUS CONSUMPTION, is a wasting or decay of the whole body, without any confiderable degree of fever, cough, or difficulty of breathing. It is attended with indigestion, weakness, want of appetite, &c.

Those who are of a fretful temper, who indulge in fpirituous liquors, or who breathe an unwholesome air, are most liable to this difease.

We would chiefly recommend, for the cure of a nervous confumption, a light and nourifhing diet, plenty of exercife in a free open air, and the ufe of fuch bitters as brace and ftrengthen the ftomach; as the Peruvian bark, gentian-root, camomile, horehound, &c. Thefe may be infufed in water or wine, and a glafs of it drank frequently.

It will greatly affift the digeftion, and promote the cure of this difeafe, to take twice a-day, twenty or thirty drops of the elixir of vitriol in a glass of wine or water. The chalybeate wine is likewife an excellent medicine in this cafe. It strengthens the folids, and powerfully affifts Nature in the preparation of good blood *.

Agreeable amusements, cheerful company, and riding

* See Appendix, Chalybeate Wine.

about,

about, are however preferable to all medicines in this difeafe. For which reafon, when the patient can afford it, we would recommend a long journey of pleafure, as the most likely means to restore his health.

What is called a *fymptomatic confumption*, cannot be cured without first removing the difease by which it is occasioned. Thus, when a confumption proceeds from the fcrophula, or king's evil, from the fcurvy, the afthma, the venereal difease, &c. a due attention must be paid to the malady from whence it arises, and the regimen and medicine directed accordingly.

When exceffive evacuations of any kind occafion a confumption, they mult not only be reftrained, but the patient's ftrength muft be reftored by gentle exercife, nourifhing diet, and generous cordials. Young and delicate mothers often fall into confumptions, by giving fuck too long. As foon as they perceive their ftrength and appetite begin to fail, they ought immediately to wean the child, or provide another nurfe, otherwife they cannot expect a cure.

Before we quit this fubject, we would earneftly recommend it to all, as they wifh to avoid confumptions, to take as much exercife without doors as they can, to avoid unwholefome air, and to ftudy fobriety. Confumptions owe their prefent increafe not a little to the fashion of fitting up late, eating hot suppers, and spending every evening over a bowl of punch or other strong liquors. These liquors, when too freely used, not only hurt the digestion, and spoil the appetite, but heat and inflame the blood, and set the whole constitution on fire.

At the beginning of this chapter I inferted an obfervation of Dr. ARBUTHNOT's, that, in his time, confumptions made up above one-tenth part of the bills of mortality in and about London. I alfo expressed my fear that the proportion was now greater, though I had not made the calculation. My reason for thinking so is, that the education of young people becomes every day more effeminate, and that an effeminate education produces a delicacy of habit, which paves the way to confumption. As all the other causes of a decline operated with as much

3

OF CONSUMPTIONS.

force about fifty years ago as they do now, the increase in the number of victims can only be ascribed to the enervating change which has taken place in the physical treatment of children. The seeds of disease are sown, as it were, in the cradle, and the sountain of life is poisoned in its source.

In tracing the various caufes of confumptions, I entered into minuter details, to put people more upon their guard, as the difeafe, when deeply feated, feldom admits of a cure. Not but there are plenty of perfons in London, who confidently undertake to perform cures in the moft hopelefs ftages of the complaint, though phyficians have not been fo happy as to find out the art. Perhaps, the only art which the others have difcovered, or which they have ever ftudied, is the art of impudence and deception.

An ignorant man advertifes a fyrup for the cure of confumptions. The people fwallow it; and the man gets a fortune, though he never cured a confumption in his life. Indeed, there is no occafion for the quack to cure any difeafe. The patient imagines he feels relief, and that anfwers the quack's purpofe as much as if he really did. Even if he feels no relief, he will fay that he does. No man will fuffer his underftanding to be impeached for having applied to a quack, when he can fo eafily get rid of the laugh by pretending to have experienced great benefit from the medicine.

But to return to my former argument: as confumptions feldom admit of a cure, the utmost care should be exerted to avoid them. The best general caution I can give, is to guard against catching cold, the fruitful mother of confumptions, and of many other diforders. How this is to be done, will be more fully explained when I come to treat of colds and coughs, the bane of this island, and the fource of numberless difeases, especially among the young, gay, and thoughtless part of the community, who have no fear of any ill until it overtakes them, when it is generally too late to prevent the fatal confequences.

CHAP.

CHAP. XIX.

add of madinal

ison your out the

OF THE SLOW OR NERVOUS FEVER.

NERVOUS fevers have increased greatly of late years in this island, owing doubtless to our different manner of living, and the increase of sedentary employments; as they commonly attack perfons of a weak relaxed habit, who neglect exercise, eat little folid food, study hard, or indulge in spirituous liquors.

CAUSES. — Nervous fevers may be occafioned by whatever depreffes the fpirits, or impoverifhes the blood; as grief, fear, anxiety, want of fleep, intenfe thought, living on poor watery diet, as unripe fruits, cucumbers, melons, mufhrooms, &c. They may likewife be occafioned by damp, confined, or unwholefome air. Hence they are very common in rainy feafons, and prove moft fatal to those who live in dirty low houses, crowded ftreets, hospitals, jails, or fuch like places.

Perfons whofe conflictutions have been broken by exceffive venery, frequent falivations, too free an ufe of purgative medicines, or any other exceffive evacuations, are most liable to this difeafe.

Keeping on wet clothes, lying on the damp ground, exceflive fatigue, and whatever obstructs the perspiration or causes a spasmodic stricture of the folids, may likewife occasion nervous fevers. We shall only add, frequent and great irregularities in diet. Too great abstinence, as well as excess, is hurtful. Nothing tends so much to preferve the body in a sound state, as a regular diet; nor can any thing contribute more to occasion fevers of the worst kind, than its opposite.

SYMPTOMS.—Low fpirits, want of appetite, weaknefs, wearinefs after motion, watchfulnefs, deep fighing, and dejection of mind, are generally the forerunners of this difeafe. Thefe are fucceeded by a quick low pulfe, a dry tongue without any confiderable thirft, chillnefs and flufhing in turns, &c.

After fome time the patient complains of a giddinels and pain of the head, has a nausea, with reachings and vomiting; vomiting; the pulfe is quick, and fometimes intermitting; the urine pale, refembling dead fmall-beer, and the breathing is difficult, with oppression of the breast, and flight alienations of mind.

If, towards the ninth, tenth, or twelfth day, thetongue becomes more moift, with a plentiful fpitting, a gentle purging, or a moifture upon the fkin; or if a fuppuration happen in one or both ears, or large puffules break out about the lips and nofe, there is reafon to hope for a favourable crifis.

But, if there be an exceffive loofenefs or walting fweats with frequent fainting fits; if the tongue, when put out, trembles exceffively, and the extremities feel cold, with a fluttering or flow creeping pulfe; if there be a flarting of the tendons, an almost total loss of fight and hearing, and an involuntary discharge by flool and urine, there is great reason to fear that death is approaching.

REGIMEN. — It is very neceffary in this difeafe to keep the patient cool and quiet. The leaft motion would fatigue him, and will be apt to occafion wearinefs, and even faintings. His mind ought not only to be kept eafy, but foothed and comforted with the hopes of a fpeedy recovery. Nothing is more hurtful in low fevers of this kind, than prefenting to the patient's imagination gloomy or frightful ideas Thefe of themfelves often occafion nervous fevers, and it is not to be doubted but they will likewife aggravate them.

The patient must not be kept too low His strength and spirits ought to be supported by nourishing diet and generous cordials. For this purpose his gruel, panado, or whatever food he takes, must be mixed with wine according as the symptoms may require. Pretty strong wine-whey, or small negus sharpened with the juice of orange or lemon, will be proper for his ordinary drink. Mustard-whey is likewise a very proper drink in this fever, and may be rendered an excellent cordial medicine by the addition of a proper quantity of white-wine *.

Wine in this difease, if it could be obtained genuine, is almost the only medicine that would be necessary.

* See Appendix, Muftard Whey.

N 3

Good

Good wine poffeffes all the virtues of the cordial medicines, while it is free from many of their bad qualities. I fay good wine; for however common this article of luxury is now become, it is rarely to be obtained genuine, especially by the poor, who are obliged to purchase it in small quantities.

I have often feen patients in low nervous fevers where the pulfe could hardly be felt, with a conftant delirium, coldnefs of the extremities, and almost every other mortal fymptom, recover by using, in whey, gruel, and negus, a bottle or two of strong wine every day. Good old found claret is the best, and may be made into negus, or given by itfelf, as circumstances require.

In a word, the great aim in this difeafe is to fupport the patient's ftrength, by giving him frequently fmall quantities of the above, or other drinks of a warm and cordial nature. He is not, however, to be overheated either with liquor or clothes; and his food ought to be light, and given in fmall quantities.

MEDICINE. — Where a naufea, load, and ficknefs at ftomach, prevail at the beginning of the fever, it will be neceffary to give the patient a gentle vomit. Fifteen or twenty grains of ipecacuanha in fine powder, or a few fpoonfuls of the vomiting julep *, will generally anfwer this purpofe very well. This may be repeated any time before the third or fourth day, if the above fymptoms continue. Vomits not only clean the ftomach, but, by the general fhock which they give, promote the perfpiration, and have many other excellent effects in flow fevers, where there are no figns of inflammation, and nature wants roufing.

Such as dare not venture upon a vomit, may clean the bowels by a fmall dofe of Turkey rhubarb, or an infufion of fenna and manna.

In all fevers, the great point is to regulate the fymptoms, fo as to prevent them from going to either extreme. Thus, in fevers of the inflammatory kind, where the force of the circulation is too great, or the blood denfe, and the fibres too rigid, bleeding and other evacuations are neceffary. But in nervous fevers, where

* See Appendix, Vomiting Julep.

nature

nature flags, where the blood is vapid and poor, and the folids relaxed, the lancet must be spared, and wine, with other cordials, plentifully administered.

It is the more neceffary to caution people against bleeding in this difease, as there is generally at the beginning an universal stricture upon the vessels, and sometimes an oppression and difficulty of breathing, which suggess the idea of a plethora, or too great a quantity of blood. I have known even some of the faculty deceived by their own feelings in this respect, so far as to infiss upon being bled, when it was evident from the confequences that the operation was improper.

Though bleeding is generally improper in this difeafe, yet bliftering is highly neceffary. Bliftering-plafters may be applied at all times of the fever with great advantage. If the patient is delirious he ought to be bliftered on the neck or head, and it will be the fafeft courfe, when the infenfibility continues, as foon as the difcharge occafioned by one bliftering-plafter abates, to apply another to fome other part of the body, and by that means keep up a continual fucceffion of them till he be out of danger.

I have been more fenfible of the advantage of bliftering in this, than in any other difeafe. Bliftering-plafters not only ftimulate the folids to action, but likewife occafion a continual difcharge, which may in fome meafure fupply the want of critical evacuations, which feldom happen in this kind of fever. They are most proper, however, either towards the beginning, or after fome degree of ftupor has come on, in which last cafe it will always be proper to blifter the head.

If the patient be coffive through the courfe of the difeafe, it will be neceffary to procure a ftool, by giving him every other day a clyfter of milk and water, with a little fugar, to which may be added a fpoonful of common falt, if the above does not operate.

Should a violent loofenefs come on, it may be checked by fmall quantities of Venice treacle, or giving the patient for his ordinary drink the white decoction *.

A miliary eruption fometimes breaks out about the nine or tenth day. As eruptions are often critical, great

* See Appendix, White Decection,

çare

care fhould be taken not to retard Nature's operation in this particular. The eruption ought neither to be checked by bleeding nor other evacuations, nor pushed out by a hot regimen; but the patient should be supported by gentle cordials, as wine-whey, small negus, fago-gruel with a little wine in it, and such like. He ought not to be kept too warm, yet a kindly breathing fweat should by no means be checked.

Though bliftering and the ufe of cordial liquors are the chief things to be depended on in this kind of fever; yet, for those who may choose to use them, we shall mention one or two of the forms of medicine which are commonly prefcribed in it *.

In defperate cafes, where the hiccup and flarting of the tendons have already come on, we have fometimes feen extraordinary effects from large dofes of mufk frequently repeated. Mufk is doubtlefs an antifpafmodic, and may be given to the quantity of a fcruple three or four times a day, or oftener if neceffary. Sometimes it may be proper to add to the mufk a few grains of camphire, and falt of hartfhorn, as thefe tend to promote perfpiration and the difcharge of urine. Thus, fifteen grains of mufk, with three grains of camphire, and fix grains of falt of hartfhorn, may be made into a bolus with a little fyrup, and given as above.

If the fever fhould happen to intermit, which it frequently does towards the decline, or if the patient's ftrength fhould be wafted with colliquative fweats, &c. it will be neceffary to give him the Peruvian bark. Half a drachm, or a whole drachm if the ftomach will bear it, of the bark in fine powder, may be given four or five times a-day in a glafs of red port or claret. Should

* When the patient is low, ten grains of Virginian fnake-root, and the fame quantity of contrayerva root, with five grains of Ruffian caftor, all in fine powder, may be made into a bolus with a little of the cordial confection or fyrup of faffron. One of thefe may be taken every four or five hours.

The following powder may be used with the fame intention: Take wild Valerian root in powder, one feruple, faffron and caftor each four grains. Mix these by rubbing them together in a mortar, and give one in a cup of wine-whey, three or four times a day.

the bark in fubftance not fit eafy on the ftomach, an ounce of it in powder may be infufed in a bottle of Lifbon or Rhenifh wine for two or three days, afterwards it may be ftrained, and a glafs of it taken frequently*.

Some give the bark in this and other fevers, where there are no fymptoms of inflammation, without any regard to the remiffion or intermiffion of the fever. How far future obfervations may tend to eftablish this practice, we will not pretend to fay; but we have reafon to believe, that the bark is a very universal febrifuge, and that it may be administered with advantage in most fevers, where bleeding is not necessary, or where there are no fymptoms of topical inflammation.

There is no fever that requires to be watched with more care and attention than this. If the actions of the fyftem are not kept up by ftimulating applications, and the patient's ftrength fupported by cordial medicines and nourifhing diet, he will fink under the difeafe; and it frequently happens, that, when the attendants think him better, he is actually dying.

I wifh to infpire not only patients in this fever, but their phyficians alfo, with unceafing, unabated hope till the very laft extremity. The changes for the better are often as fudden, and unforeseen, as those for the worfe. The laft gasp alone should induce us to give over the patient. I have left a patient twenty times and more, little expecting to see him alive next day. Yet I did not lose courage, but ordered a bottle, or perhaps two, of generous wine to be given in the course of twentyfour hours; and that patient, to my great fatisfaction, recovered, and enjoyed health for many years after.

* The bark may likewife be very properly administered, along with other cordials, in the following manner : Take an ounce of Peruvian bark, orange-peel half an ounce, Virginian fnake-root two drachms, faffron one drachm. Let all of them be powdered, and infused in an English pint of the best brandy for three or four days. Afterwards the liquor may be strained, and two tea-spoonfuls of it given three or four times a-day in a glass of small wine or negus. I now generally administer Huxham's Tincture.

[186]

CHAP. XX.

OF THE MALIGNANT, PUTRID, OR SPOTTED FEVER.

THIS may be called the *peftilential fever* of Europe, as in many of its fymptoms it bears a great refemblance to that dreadful difeafe the plague. Perfons of a lax habit, a melancholy difpolition, and those whose vigour has been wasted by long fasting, watching, hard labour, exceffive venery, frequent falivations, &c. are most liable to it.

CAUSES.—This fever is occafioned by foul air, from a number of people being confined in a narrow place, not properly ventilated; from putrid animal and vegetable effluvia, &c. Hence it prevails in camps, jails, hofpitals, and infirmaries, efpecially where fuch places are too much crowded, and cleanlinefs is neglected.

A clofe conftitution of the air, with long rainy or foggy weather, likewife occafions putrid fevers. They often fucceed great inundations in low and marfhy countries, efpecially when thefe are preceded or followed by a hot and fultry feafon.

Living too much upon animal food, without a proper mixture of vegetables, or eating fifh or flefh that has been kept too long, are likewife apt to occafion this kind of fever. Hence failors on long voyages, and the inhabitants of befieged cities, are very often vifited with putrid fevers.

Corn that has been greatly damaged by rainy feafons, or long keeping, and water which has become putrid by ftagnation, &c. may likewife occasion this fever.

Dead carcaffes tainting the air, efpecially in hot feafons, are very apt to occafion putrid difeafes. Hence this kind of fever often prevails in countries which are the fcenes of war and bloodshed. This shews the propriety of removing burying-grounds, flaughter-houses, &c. at a proper distance from great towns.

Want of cleanlinefs is a very general caufe of putrid fevers. Hence they prevail amongst the poor inhabitants of large towns, who breathe a confined unwholefome air, and neglect cleanlinefs. Such mechanics as carry on dirty employments, and are constantly confined within doors, are likewife very liable to this difease.

We fhall only add, that putrid, malignant, or fpotted fevers, are highly infectious, and are therefore often communicated by contagion. For which reafon, all perfons ought to keep at a diftance from those affected with fuch difeafes, unless their attendance is abfolutely neceffary.

SYMPTOMS.—The malignant fever is generally preceded by a remarkable weaknefs, or lofs of ftrength, without any apparent caufe. This is fometimes fo great, that the patient can fcarce walk, or even fit upright, without being in danger of fainting away. His mind too is greatly dejected ; he fighs, and is full of dreadful apprehenfions.

There is a naufea, and fometimes a vomiting of bile; a violent pain of the head, with a ftrong pulfation or throbbing of the temporal arteries; the eyes often appear red and inflamed, with a pain at the bottom of the orbit; there is a noife in the ears, the breathing is laborious, and often interrupted with a figh; the patient complains of a pain about the region of the ftomach, and in his back and loins; his tongue is at first white, but afterwards it appears black and chaped; and his teeth are covered with a black cruft. He fometimes paffes worms both upwards and downwards, is affected with tremors or fhaking, and often becomes delirious.

If blood be let, it appears diffolved, or with a very fmall degree of cohefion, and foon becomes putrid; the ftools fmell extremely foetid, and are fometimes of a greenifh, black, or reddifh caft. Spots of a pale, purple, dun, or black colour, often appear upon the fkin, and fometimes there are violent hæmorrhages or difcharges of blood from the mouth, eyes, nofe, &c.

Putrid feversmay be diftinguished from the inflamma, tory, by the smallness of the pulse, the great dejection

of mind, the diffolved flate of the blood, the petechiæ or purple fpots, and the putrid fmell of the excrements. They may likewife be diffinguished from the low or nervous fever, by the heat and thirst being greater, the urine of a higher colour, and the loss of strength, dejection of mind, and all the other symptoms, more violent.

It fometimes happens, however, that the inflammatory, nervous, and putrid fymptoms are fo blended together, as to render it very difficult to determine to which clafs the fever belongs. In this cafe, the greateft caution and fkill are requifite. Attention must be paid to those fymptoms which are most prevalent, and both the regimen and medicines adapted to them.

Inflammatory and nervous fevers may be converted into malignant and putrid, by too hot a regimen, or improper medicines.

The duration of putrid fevers is extremely uncertain; fometimes they terminate between the feventh and fourteenth day, and at other times they are prolonged for five or fix weeks. Their duration depends greatly upon the conflictution of the patient, and the manner of treating the difeafe.

The most favourable fymptoms are, a gentle loofeness after the fourth or fifth day, with a warm mild tweat. These, when continued for a confiderable time, often carry off the fever, and should never be imprudently stopped. Small miliary pusses appearing between the petechiæ or purple stops are likewise favourable, as also hot scabby eruptions about the mouth and nose. It is a good fign when the pulse rises upon the use of wine, or other cordials, and the nervous symptoms abate; deafness coming on towards the decline of the fever, is likewise often a favourable symptom*, as are absceffes in the groin, or parotid glands.

Among the unfavourable fymptoms may be reckoned an exceffive loofenefs with a hard fwelled belly; large black or livid blotches breaking out upon the fkin;

apthæ

^{*} Deafnels is not always a favourable fymptom in this difeafe. Perhaps it is only fe, when occasioned by abscelles formed within the cars.

apthæ in the mouth; cold clammy fweats; blindnefs; change of the voice; a wild ftaring of the eyes; difficulty of fwallowing; inability to put out the tongue; and a conftant inclination to uncover the breaft. When the fweat and faliva are tinged with blood, and the urine is black, or depofits a black footy fediment, the patient is in great danger. Starting of the tendons, and fœtid, ichorous, involuntary ftools, attended with coldnefs of the extremities, are generally the forerunners of death.

REGIMEN.—In the treatment of this difeafe, we ought to endeavour as far as poffible to counteract the putrid tendency of the humours; to fupport the patient's firength and fpirits; and to affift Nature in expelling the caufe of this difeafe, by gently promoting perfpiration and the other evacuations.

It has been obferved, that putrid fevers are often occafioned by unwholefome air, and of courfe they muft be aggravated by it. Care fhould therefore be taken to prevent the air from ftagnating in the patient's chamber, to keep it cool, and renew it frequently, by opening the doors or windows of fome adjacent apartment. The breath and perfpiration of perfons in perfect health foon render the air of a fmall apartment noxious; but this will fooner happen from the perfpiration and breath of a perfon whole whole mafs of humours are in a putrid ftate.

Befides the frequent admiffion of frefh air, we would recommend the ufe of vinegar, verjuice, juice of lemon, Seville orange, or any kind of vegetable acid that can be moft readily obtained. Thefe ought frequently to be fprinkled upon the floor, the bed, and every part of the room. They may alfo be evaporated with a hot iron, or by boiling, &cc. The frefh fkins of lemons or oranges ought likewife to be laid in different parts of the room, and they fhould be frequently held to the patient's nofe. The ufe of acids in this manner would not only prove very refrefhing to the patient, but would likewife tend to prevent the infection from fpreading among thofe who attend him. Strong fcented herbs, as rue, tanly, rofemary, wormwood, &cc, may likewife be laid in different parts

parts of the houfe, and fmelled to by those who go near the patient.

The patient must not only be kept cool, but likewise quiet and easy. The least noise will affect his head, and the smallest fatigue will be apt to make him faint.

Few things are of greater importance in this difeafe than acids, which ought to be mixed with all the patient's food as well as drink. Orange, lemon, or vinegar-whey, are all very proper, and may be drank by turns according to the patient's inclination. They may be rendered cordial by the addition of wine in fuch quantity as the patient's ftrength feems to require. When he is very low, he may drink negus, with only one half water, and fharpened with the juice of bitter orange or lemon. In fome cafes a glafs of wine may now and then be allowed. The moft proper wine is Rhenifh; but if the body be open, red port or claret is to be preferred.

When the body is bound, a tea-fpoonful of the cream of tartar may be put into a cup of the patient's drink, as there is occasion; or he may drink a decoction of tamarinds, which will both quench his thirst, and promote a discharge by stool.

If camomile-tea will fit upon his ftomach, it is a very proper drink in this difeafe. It may be fharpened by adding to every cup of tea, ten or fifteen drops of the elixir of vitriol.

The food must be light, as panado, or groat-gruel, to which a little wine may be added, if the patient be weak and low; and they ought all to be sharpened with the juice of orange, the jelly of currants, or the like. The patient ought likewife to eat freely of ripe fruits, as roafted apples, currant or goosberry tarts, preferved cherries, or plums, &c.

Taking a little food or drink frequently, not only fupports the fpirits, but counteracts the putrid tendency of the humours; for which reafon the patient ought frequently to be fipping fmall quantities of fome of the acid liquors mentioned above, or any that may be more agreeable to his palate, or more readily obtained.

If he be delirious, his feet and hands ought to be frequently fomented with a throng infufion of camomile flowers. This, or an infufion of the bark, to fuch as can afford it, cannot fail to have a good effect. Fomentations of this kind not only relieve the head, by relaxing the veffels in the extremities, but as their contents are abforbed, and taken into the fyftem, they may affift in preventing the putrefcency of the humours.

MEDICINE. — If a vomit be given at the beginning of this fever, it will hardly fail to have a good effect; but if the fever has gone on for fome days, and the fymptoms are violent, vomits are not quite fo fafe. The body, however, is always to be kept gently open, by clyfters, or mild laxative medicines.

Bleeding is feldom neceffary in putrid fevers. If there be figns of an inflammation, it may fometimes be permitted at the first onset; but the repetition of it generally proves hurtful.

Bliftering plafters are never to be used unless in the greatest extremities. If the petechiæ or spots should suddenly disappear, the patient's pulse fink remarkably, and a delirium, with other bad symptoms come on, bliftering may be permitted. In this case the bliftering plasters are to be applied to the head, and infide of the legs or thighs. But as they are sometimes apt to occasion a gangrene, we would rather recommend warm cataplasms, or poultices of mustard and vinegar to be applied to the feet, having recourse to blifters only in the utmost extremities.

It is common in the beginning of this fever to give the emetic tartar in fmall dofes, repeated every fecond or third hour, till it fhall either vomit, purge, or throw the patient into a fweat. This practice is very proper, provided it be not pushed fo far as to weaken the patient.

A very ridiculous notion has long prevailed, of expelling the poifonous matter of malignant difeafes by triffing dofes of cordial or alexipharmic medicines. In confequence of this notion, the contrayerva-root, the cordial confection, the mithridate, &c. have been extolled

tolled as infallible remedies. There is reafon however to believe, that these feldom do much good. Where cordials are neceffary, we know none that is superior to good wine; and therefore again recommend it both as the fafest and best. Wine, with acids and antiseptics, are the only things to be relied on in the cure of malignant fevers.

In the most dangerous species of this difease, when it is attended with purple, livid, or black spots, the Peruvian bark must be administered. I have seen it, when joined with acids, prove successful, even in cases where the petechiæ had the most threatening aspect. But, to answer this purpose, it must not only be given in large dose, but duly perfisted in.

The beft method of administering the bark is certainly in fubftance. An ounce of it in powder may be mixed with half an English pint of water, and the same quantity of red wine, and sharpened with the elixir or the spirit of vitriol, which will both make it fit easier on the ftomach, and render it more beneficial. Two or three ounces of the syrup of lemon may be added; and two table-spoonfuls of the mixture taken every two hours, or oftener, if the stomach is able to bear it.

Those who cannot take the bark in fubstance, may infuse it in wine, as recommended in the preceding difease.

If there be a violent loofenefs, the bark must be boiled in red wine with a little cinnamon, and sharpened with the elixir of vitriol, as above. Nothing can be more beneficial in this kind of loofenefs than plenty of acids, and fuch things as promote a gentle perspiration.

If the patient be troubled with vomiting, a dram of the falt of wormwood, diffolved in an ounce and a half of fresh lemon-juice, and made into a draught, with an ounce of simple cinnamon-water, and a bit of sugar, may be given and repeated as often as it is necessary.

If fwellings of the glands appear, their fuppuration is to be promoted by the application of poultices, ripening cataplasms, &c.; and as soon as there is any appearance of

of matter in them, they ought to be laid open, and the poultices continued.

I have known large ulcerous fores break out in various parts of the body, in the decline of this fever, of a livid gangrenous appearance, and a most putrid cadaverous fmell. These gradually healed, and the patient recovered, by the plentiful use of Peruvian bark and wine, sharpened with the vitriolic acid.

For preventing putrid fevers, we would recommend a firicit regard to cleanlinefs; a dry fituation; fufficient exercise in the open air; wholesome food, and a moderate use of generous liquors. Infection ought, above all things, to be avoided. No conflictution is proof against it. I have known perfons seized with a putrid fever, by only making a fingle visit to a patient in it; others have caught it by lodging for one night in a town where it prevailed; and some by attending the funerals of such as died of it*.

When a putrid fever feizes any perfon in a family, the greateft attention is neceffary to prevent the difeafe from fpreading. The fick ought to be placed in a large apartment, as remote from the reft of the family as poffible; he ought likewife to be kept extremely clean, and fhould have frefh air frequently let into his chamber; whatever comes from him fhould be immediately removed, his linen fhould be frequently changed, and those in health ought to avoid all unneceffary communication with him.

Any one who is apprehensive of having caught the infection, ought immediately to take a vomit, and to work it off by drinking plentifully of camomile tea. This may be repeated in a day or two, if the appre-

• The late Sir John Pringle expressed a concern left these cautions should prevent people from attending their friends or relations when afflicted with putrid fevers. I told him I meant only to difcourage unnecessary attendance, and mentioned a number of instances where putrid fevers had proved tatal to perfons, who were rather hurtful than beneficial to the fick. This fagacious physician agreed with me, in thinking that a good doctor and a careful nurse were the only necessary attendants; and that all others not only endangered themselves, but generally, by their folicitude and illdiracted care, hurt the fick.

henfions

henfions still continue, or any unfavourable fymptoms appear.

The perfon ought likewife to take an infusion of the bark and camomile flowers for his ordinary drink; and before he goes to bed, he may drink an English pint of pretty ftrong negus, or a few glaffes of generous wine. I have been frequently obliged to follow this courfe when malignant fevers prevailed, and have likewife recommended it to others with constant fuccefs.

People generally fly to bleeding and purging as antidotes against infection ; but these are so far from securing them, that they often, by debilitating the body, increafe the danger.

Those who wait upon the fick in putrid fevers, ought always to have a piece of fpunge or a handkerchief dipt in vinegar, or juice of lemon, to fmell to while near the patient. They ought likewife to wash their hands, and, if poffible, to change their clothes, before they go into company.

1 wifhed flrongly to characterife this difeafe by calling it the pestilential fever of Europe. It is certain that in our ifland nothing approaches fo near the plague as the putrid or fpotted fever. I have often marked its ravages in the narrow lanes of London, but ftill more frequently in fome of the unclean and fhamefully neglected feats of putrefaction in the environs. I have feen it in its most horrid form in St. George's Fields, particularly in the King's Bench prifon; and am afraid the day will come when the real peftilence will be generated in that quarter.

Four hundred thousand people living in a marsh, where there is not the leaft fall or defcent to carry off the perperually accumulating filth, must in time produce forme dreadful putrid diforder. How justly may one apply to the atmosphere of fuch a place the words of the medical poet, and fay, that it is not air the inhabitants breathe, but volatile corruption ! Yet that marsh, by means of fluices or proper drains, might be rendered as healthy a place of refidence as any in London, which is, perhaps, the most wholesome city of its fize in the world. I shudder at the idea that the ftagnant mafs of fo much contagion should be viewed with indifference, and that no _ fteps

fteps are likely to be taken to avert the danger, till a terrible fweep of mortality shall spread alarm through the whole kingdom.

To fhew how nearly our putrid fevers refemble the plague, I shall give a short sketch of one that I attended fome years ago at Kentish Town. A young lady, about eighteen years of age, who had come from the Weft Indies for her education, was feized with the common fymptoms of a nervous fever; but this continuing a very unufual length of time, putrefaction began to fhew itfelf, and boils and imposthumes burst forth, and penetrated fo deeply as to lay the bones bare, to an amazing extent, on the back, hips, and various parts of the body; nor were these healed without much time and trouble.

By the use of wine, bark, and other cardiac and antifeptic medicines, a perfect cure was at length effected ; and what may be particularly deferving of notice, the young lady, at one period, had no fuftenance for feveral days, but what fhe received from fipping a little brandy. The nature of a fupplementary remark will not permit me to enter into all the details of this cafe, however curious; but I must not omit mentioning one circumstance more, which was, that the nurse, for reasons best known to herfelf, yet eafily gueffed at, went often to the lady under whofe care my patient was, and complained that the never faw fuch a doctor, for be would not let her poor young lady die.

CHAP. XXI.

OF THE MILIARY FEVER.

THIS fever takes its name from the small pustules or bladders which appear on the fkin, refembling, in shape and fize, the feeds of millet. The pustules are either red or white, and fometimes both are mixed together.

The whole body is fometimes covered with pultules; but they are generally more numerous where the fweat is most abundant, as on the breast, the back, &c. A 02

gentle

gentle fweat, or moifture on the fkin, greatly promotes the eruption; but when the fkin is dry, the eruption is both more painful and dangerous.

Sometimes this is a primary difeafe; but it is much oftener only a fymptom of fome other malady, as the fmall-pox, meafles, ardent, putrid, or nervous fever, &c. In all these cafes it is generally the effect of too hot a regimen or medicines.

The miliary fever chiefly attacks the idle and the phlegmatic, or perfons of a relaxed habit. The young and the aged are more liable to it than those in the vigour and prime of life. It is likewise more incident to women than men, especially the delicate and the indolent, who, neglecting exercise, keep continually within doors, and live upon weak watery diet. Such females are extremely liable to be feized with this difease in childbed, and often lose their lives by it.

CAUSES.——The miliary fever is fometimes occafioned by violent paffions or affections of the mind; as exceffive grief, anxiety, thoughtfulnefs, &cc. It may likewife be occafioned by exceffive watching, great evacuations, a weak watery diet, rainy feafons, eating too freely of cold, crude, unripe fruits, as plums, cherries, cucumbers, melons, &cc. Impure waters, or provifions which have been fpoiled by rainy feafons, long keeping, &c. may likewife caufe miliary fevers. They may allo be occafioned by the ftoppage of any cuftomary evacuation, as iffues, fetons, ulcers, the bleeding piles in men, or the menftrual flux in women, &c.

This difeafe in childbed-women is fometimes the effect of great cofliveness during pregnancy; it may likewise be occasioned by their excessive use of green trash, and other unwholesome things, in which pregnant women are too apt to indulge. But its most general cause is indolence. Such women as lead a fedentary life, especially during pregnancy, and at the fame time live grossly, can hardly escape this difease in childbed. Hence it proves extremely fatal to women of fashion, and likewise to those women in manufacturing towns, who, in order to affist their husbands, fit close within doors for almost the whole of their time. But among women who are active active and laborious, who live in the country, and take fufficient exercife without doors, this difeafe is very little known.

SYMPTOMS. — When this is a primary difeafe, it makes its attack, like most other eruptive fevers, with a flight fhivering, which is fucceeded by heat, loss of ftrength, faintifhness, fighing, a low quick pulse, difficulty of breathing, with great anxiety and oppression of the breast. The patient is restless, and sometimes delirious; the tongue appears white, and the hands shake, with often a burning heat in the palms; and in childbed-women the milk generally goes away, and the other discharges stop.

The patient feels an itching or pricking pain under the fkin, after which innumerable fmall puffules of a red or white colour begin to appear. Upon this the fymptoms generally abate, the pulfe becomes more full and foft, the fkin grows moifter, and the fweat, as the difeafe advances, begins to have a peculiar foctid fmell; the great load on the breaft, and opprefilion of the fpirits, generally go off, and the cuftomary evacuations gradually return. About the fixth or feventh day from the eruption, the puffules begin to dry and fall off, which occafions a very difagreeable itching in the fkin.

It is impossible to afcertain the exact time when the pussible will either appear or go off. They generally come out on the third or fourth day, when the eruption is critical; but, when fymptomatical, they may appear at any time of the difease.

Sometimes the pultules appear and vanish by turns. When that is the cafe, there is always danger; but when they go in all of a fudden, and do not appear again, the danger is very great.

In childbed-women the puftules are commonly at first filled with clear water, afterwards they grow yellowish. Sometimes they are interspersed with pustules of a red colour. When these only appear, the disease goes by the name of a *rash*.

REGIMEN.——In all eruptive fevers, of whatever kind, the chief point is to prevent the fudden difappearing of the puftules, and to promote their maturation. For this purpofe, the patient must be kept in fuch a temperature,

as

198

as neither to pufh out the eruption too faft, nor to caufe it to retreat prematurely. The diet and drink ought therefore to be in a moderate degree nourifhing and cordial; but neither ftrong nor heating. The patient's chamber ought neither to be kept too hot nor cold; and he fhould not be too much covered with clothes. Above all, the mind is to be kept eafy and cheerful. Nothing fo certainly makes an eruption go in as fear, or the apprehenfion of danger.

The food must be weak chicken-broth, with bread, panado, fago, or groat-gruel, &c. to a gill of which may be added a spoonful or two of wine, as the patient's strength requires, with a few grains of falt and a little fugar. Good apples, roasted or boiled, with other ripe fruits of an opening cooling nature, may be eaten.

The drink may be fuited to the flate of the patient's flrength and fpirits. If these be pretty high, the drink ought to be weak; as water-gruel, balm-tea, or the decoction mentioned below *.

When the patient's fpirits are low, and the eruption does not rife fufficiently, his drink must be a little more generous; as wine-whey, or fmall negus, sharpened with the juice of orange or lemon, and made stronger or weaker as circumstances may require.

Sometimes the miliary fever approaches towards a putrid nature, in which cafe the patient's ftrength muft be fupported with generous cordials, joined with acids; and, if the degree of putrefcence be great, the Peruvian bark muft be administered. If the head be much affected, the body muft be kept open by emollient clysters \dagger .

MEDI-

* Take two ounces of the fhavings of hartfhorn, and the fame quantity of farfaparilla, boil them in two English quarts of water. To the ftrained decoction add a little white fugar, and let the patient take it for his ordinary drink.

† In the Commercium Literarium for the year 1735, we have the history of an epidemical miliary fever, which raged at Strafburgh in the months of November, December, and January; from which we learn the neceffity of a temperate regimen in this malady, and likewife that physicians are not always the first who discover the proper treatment of difeases. "This fever made terrible

MEDICINE. ———If the food and drink be properly regulated, there will be little occafion for medicine in this difeafe. Should the eruption, however, not rife, or the fpirits flag, it will not only be neceffary to fupport the patient with cordials, but likewife to apply bliftering plafters. The most proper cordial in this cafe is good wine, which may either be taken in the patient's food or drink; and if there be figns of putrescence, the bark and acids may be mixed with wine, as directed in the putrid fever.

Some recommend bliftering through the whole courfe of this difeafe; and where nature flags, and the eruption comes and goes, it may be neceffary to keep up a flimulus, by a continual fucceffion of fmall bliftering plafters; but we would not recommend above one at a time. If, however, the pulfe flould fink remarkably, the pultules flrike in, and the head be affected, it will be neceffary to apply feveral bliftering plafters to the moft fenfible parts, as the infide of the legs, thighs, &c.

Bleeding is feldom neceffary in this difeafe, and fometimes it does much hurt, as it weakens the patient and depreffes his fpirits. It is therefore never to be attempted unlefs by the advice of a phyfician. We mention this, becaufe it has been cuftomary to treat this difeafe in childbed women by plentiful bleeding, and other evacuations, as if it were highly inflammatory. But this practice is generally very unfafe. Patients in this

terrible havock even among men of robult conflications, and all medicine proved in vain. They were feized in an inftant with fhivering, yawning, firetching, and pains in the back, fucceeded by a most intense heat; at the same time there was a great loss of frength and appetite. On the feventh or ninth day the miliary eruptions appeared, or lpots like flea-bites, with great anxiety, a delirium, reftlessnefs, and tolling in bed. Bleeding was fatal. While matters where in this unhappy fituation, a midwife, of her own accord, gave to a patient, in the height of the dife le, a clyfter of rain water and butter without falt, and for his ordinary drink a quart of spring-water, half a pint of generous wine, the juice of a lemon, and fix ounces of the whitell fugar, gently boiled till a fcum arole, and this with great fuccels; for the belly was foon loofened, the grievous fymptoms vanished, and the patient was reflored to his fenfos, and fnatched from the jaws of death." This practice was imitated by others with the like happy effects.

199

04

fituation

200

fituation bear evacuations very ill. And, indeed, the difeafe feems often to be more of a putrid than of an in-flammatory nature.

Though this fever is often occafioned in childbedwomen by too hot a regimen, yet it would be dangerous to leave that off all of a fudden, and have recourfe to a very cool regimen, and large evacuations. We have reafon to believe, that fupporting the patient's fpirits, and promoting the natural evacuations, is here much fafer than to have recourfe to artificial ones, as thefe, by finking the fpirits, feldom fail to increafe the danger.

If the difeafe proves tedious, or the recovery flow, we would recommend the Peruvian bark, which may either be taken in fubftance, or infufed in wine or water, as the patient inclines.

The miliary fever, like other eruptive difeafes, requires gentle purging, which fhould not be neglected, as foon as the fever is gone off, and the patient's ftrength will permit.

To prevent this difeafe, a pure dry air, fufficient exercife, and wholefome food, are neceffary. Pregnant women fhould guard against costiveness, and take daily as much exercise as they can bear, avoiding all green trashy fruits, and other unwholefome things; and, when in childbed, they ought strictly to observe a cool regimen.

There is not any fever, in which the fymptoms ought to be more carefully watched than in this. The changes are frequent and rapid, and the fever itfelf often affumes a quite different character. It is, therefore, of the utmost importance upon fuch occasions to change the regimen and medicines, and adapt them to the new fymptoms. Death would often be the confequence of inattention or neglect in thefe cafes; and perhaps a ftronger proof cannot be given of what I have already pointed out, but cannot too often inculcate, the extreme folly of using or recommending any general fever medicine, when even the same fever may require, at different periods, very different modes of treatment. Really, it is not less ridiculous to prefcribe one medicine for all fevers, than for all difeafes. The quackery in the first instance may appear to the ignorant and thoughtlefs more plaufible; but is on that account more dangerous.

CHAP.

CHAP. XXII.

OF THE REMITTING FEVER.

THIS fever takes its name from a remiffion of the fymptoms, which happens fometimes fooner, and fometimes later, but generally before the eighth day. The remiffion is commonly preceded by a gentle fweat, after which the patient feems greatly relieved, but in a few hours the fever returns. Thefe remiffions return at very irregular periods, and are fometimes of longer, fometimes of fhorter duration: the nearer, however, that the fever approaches to a regular intermittent, the danger is the lefs.

CAUSES.———Remitting fevers prevail in low marfhy countries abounding with wood and ftagnating water; but they prove molt fatal in places where great heat and moifture are combined, as in fome parts of Africa, the province of Bengal in the Eaft Indies, &c. where remitting fevers are generally of a putrid kind, and prove very fatal. They are moft frequent in clofe calm weather, efpecially after rainy feafons, great inundations, or the like. No age, fex, or conftitution is exempted from the attack of this fever; but it chiefly feizes perfons of a relaxed habit, who live in low dirty habitations, breathe an impure ftagnating air, take little exercife, and ufe unwholefome diet.

SYMPTOMS.——The first fymptoms of this fever, are generally yawning, stretching, pain, and giddiness in the head, with alternate fits of heat and cold. Sometimes the patient is affected with a delirium at the very first attack. There is a pain, and sometimes a swelling, about the region of the stomach, the tongue is white, the eyes and skin frequently appear yellow, and the patient is often afflicted with bilious vomitings. The pulse is fometimes a little hard, but feldom full, and the blood, when let, rarely shews any signs of inflammation. Some patients are exceedingly costive, and others are afflicted with a very troubles loofenes.

202 OF THE REMITTING FEVER.

It is impoffible to defcribe all the fymptoms of this difeafe, as they vary according to the fituation, the feafon of the year, and the conftitution of the patient. They may likewife be greatly changed by the method of treatment, and by many other circumftances too tedious to mention. Sometimes the bilious fymptoms predominate, fometimes the nervous, and at other times the putrid. Nor is it at all uncommon to find a fucceffion of each of thefe, or even a complication of them at the fame time, in the fame perfon.

REGIMEN.——The regimen must be adapted to the prevailing fymptoms. When there are any figns of inflammation, the diet must be flender, and the drink weak and diluting. But when nervous or putrid fymptoms prevail, it will be neceffary to fupport the patient with food and liquors of a more generous nature, fuch as are recommended in the immediately preceding fevers. We must, however, be very cautious in the use of things of a heating quality, as this fever is frequently changed into a *continual* by an hot regimen, and improper medicines.

Whatever the fymptoms are, the patient ought to be kept cool, quiet, and clean. His apartment, if poffible, fhould be large and frequently ventilated by letting in frefh air at the doors and windows It ought likewife to be fprinkled with vinegar, juice of lemon, or the like. His linen, bed-clothes, &c. fhould be frequently changed, and all his excrements immediately removed. Though these things have been recommended before, we think it necessary to repeat them here, as they are of more importance to the fick than practitioners are apt to imagine *.

MEDI-

• The ingenious Dr. Lind, of Windfor, in his inaugural differtation concerning the putrid remitting fever of Bengal, has the following observation : "Industa, lodices, ac stragula, fapius funt mutanda, ac aëri exponenda; faces fordesque quam primum removendæ, oportet etiam ut loca quibus ægri decumbent fint falubria et aceto conspersa; denique ut ægris cura quanta maxima prospiciatur. Compertum ego habeo, medicum hæc fedula obfervantem, quique ea exequi potest, multo magis ægris profuturum, quam medicum peritiorem, hisce commodis destitutum."

OF THE REMITTING FEVER.

MEDICINE. — In order to cure this fever, we must endeavour to bring it to a regular intermistion. This intention may be promoted by bleeding, if there be any figns of inflammation; but when that is not the cafe, bleeding ought by no means to be attempted, as it will weaken the patient and prolong the difeafe. A vomit, however, will feldom be improper, and is generally of great fervice. Twenty or thirty grains of ipecacuanha will anfwer this purpofe very well; but, where it can be obtained, we would rather recommend a grain or two of tartar emetic, with five or fix grains of ipecacuanha, to be made into a draught, and given for a vomit. This may be repeated once or twice at proper intervals, if the ficknefs or naufea continues.

The body ought to be kept open either by clyfters or gentle laxatives, as weak infufions of fenna and manna, fmall dofes of the lenitive electuary, cream of tartar, tamarinds, ftewed prunes, or the like; but all ftrong or draftic purgatives are to be carefully avoided.

By this courfe the fever in a few days may generally be brought to a pretty regular or diffinct intermiffion, in which cafe the Peruvian bark may be administered, and it will feldom fail to perfect the cure. It is needlefs here to repeat the methods of giving the bark, as we have already had occasion frequently to mention them.

The most likely way to avoid this fever is to use a wholefome or nourifhing diet, to pay the most for upulous attention to cleanlines, to keep the body warm, to take fufficient exercise, and in hot countries to avoid damp fituations, night air, evening dews, and the like. In countries where it is endemical, the best preventive medicine which we can recommend is the Peruvian bark, which may either be chewed, or infused in brandy or

"The patient's fhirt, bed-clothes, and bedding, ought frequently to be changed, and exposed to the air, and all his excrements immediately removed; the bed-chamber should be well ventilated, and frequently sprinkled with vinegar; in short every attention should be paid to the patient. I can affirm, that a physician who puts these in practice will much oftener succeed, than one who is even more skilful, but has not opportunity of using these means."

204 ON THE REMITTING FEVER.

wine, &c. Some recommend fmoking tobacco as very beneficial in marfhy countries, both for prevention of this and intermitting fevers.

As diforders of this kind are more to be dreaded in a camp than the approach of an enemy, it is the duty of fuperior officers very earneftly to concur with their medical attendants in enforcing the proper means of prevention. The fpirit of our foldiers betrays them into a contempt of difeafe, as well as of danger ; and they are too apt to forget, that no hardihood can of itself refift the warm, fickly moisture of autumn, and the damp air of the night to which they are often unavoidably exposed. Those brave, but thoughtless men, should, therefore, be obliged to pay more attention to the fimple prefervatives from fevers above pointed out. I have too high an opinion of the talents of many eminent phyficians and furgeons now in the army, to think any farther remarks on this fubject neceffary. I am perfuaded that a hint will be fufficient to call forth the fulleft exercise of their skill, their humanity, and their zeal alfo for the honour and fecurity of their country, in faving the lives, and promoting the health and vigour of its gallant defenders.

CHAP. XXIII.

OF THE SMALL-POX.

THIS difeafe, which originally came from Arabia, is now become fo general, that very few efcape it at one time of life or another. It is a most contagious malady; and has for many years proved the fcourge of Europe.

The fmall-pox generally appear towards the fpring. They are very frequent in fummer, lefs fo in autumn, and leaft of all in winter. Children are most liable to this difease; and those whose food is unwholesome, who want proper exercise, and abound with gross humours, run the greatest hazard from it.

The difease is diffinguished into the diffinct and confluent kind; the latter of which is always attended with danger.

danger. There are likewife other diffinctions of the fmall-pox; as the chrystalline, the bloody, &c.

CAUSES. — The fmall-pox is commonly caught by infection. Since the difeafe was first brought into Europe, the infection has never been wholly extinguished, nor have any proper methods, as far as 1 know, been taken for that purpose; fo that now it has become in a manner constitutional. Children who have over-heated themselves by running, wrestling, &c. or adults after a debauch, are most apt to be feized with the small-pox.

SYMPTOMS. This difeafe is fo generally known, that a minute description of it is unneceffary. Children commonly look a little dull, feem liftlefs and drowfy for a few days before the more violent fymptoms of the fmall-pox appear. They are likewife more inclined to drink than ufual, have little appetite for folid food, complain of wearinefs, and, upon taking exercife, are apt to fweat. Thefe fymptoms are fucceeded by flight fits of cold and heat in turns, which, as the time of the eruption approaches, become more violent, and are accompanied with pains of the head and loins, vomiting, &c. The pulfe is quick, with a great heat of the fkin, and reftlefinefs. When the patient drops afleep, he wakes in a kind of horror, with a fudden ftart, which is a very common fymptom of the approaching eruption ; as are alfo convultion-fits in very young children.

About the third or fourth day from the time of fickening, the fmall-pox generally begin to appear; fometimes, indeed, they appear fooner, but that is no favourable fymptom. At first they very nearly refemble fleabites, and are foonest difference on the face, arms, and breast.

The most favourable fymptoms are a flow eruption, and an abatement of the fever as foon as the pultules appear. In a mild diffinct kind of finall-pox the pultules feldom appear before the fourth day from the time of fickening, and they generally keep coming out gradually for feveral days after. Puftules which are diftinct, with a florid red bafis, and which fill with thick purulent purulent matter, first of a whitish, and afterwards of a yellowish colour, are the best.

A livid brown colour of the puftules is an unfavourable fymptom; as also when they are small and flat, with black specks in the middle. Pustules which contain a thin watery ichor are very bad. A great number of pox on the face is always attended with danger. It is likewise a bad fign when they run into one another.

It is a most unfavourable fymptom when petechiæ or purple, brown, or black spots are interspersed among the pultules. These are figns of a putrid diffolution of the blood, and fhew the danger to be very great. Bloody stools or urine, with a fwelled belly, are bad fymptoms; as is also a continual strangury. Pale urine and a violent throbbing of the arteries of the neck are figns of an approaching delirium or of convultion-fits. When the face does not fwell, or falls before the pox come to maturity, it is very unfavourable. If the face begins to fall about the eleventh or twelfth day, and at the fame time the hands and feet begin to fwell, the patient generally does well; but when these do not fucceed each other, there is reafon to apprehend danger. When the tongue is covered with a brown cruft, it is an unfavourable fymptom. Cold fhivering fits coming on at the height of the difease, are likewise unfavourable. Grinding of the teeth, when it proceeds from an affection of the nervous system, is a bad sign; but sometimes it is occasioned by worms, or a difordered stomach.

REGIMEN. — When the first fymptoms of the fmall-pox appear, people are ready to be alarmed, and often fly to the use of medicine, to the great danger of the patient's life. I have known children, to appeale the anxiety of their parents, bled, bliftered, and purged, during the fever which preceded the eruption of the small-pox, to such a degree, that Nature was not only disturbed in her operation, but rendered unable to support the puscular they were out; fo that the patient, exhausted by mere evacuations, funk under the difease.

206

When

When convultions appear, they give a dreadful alarm. Immediately fome noftrum is applied, as if this were a primary difeafe; whereas it is only a fymptom, and far from being an unfavourable one, of the approaching eruption. As the fits generally go off before the actual appearance of the fmall-pox, it is attributed to the medicine, which by this means acquires a reputation without any merit^{*}.

All that is, generally fpeaking, neceffary during the eruptive fever, is to keep the patient cool and eafy, allowing him to drink freely of fome weak diluting liquors; as balm-tea, barley-water, clear whey, gruels, &c. He fhould not be confined to bed, but fhould fit up as much as he is able, and fhould have his feet and legs frequently bathed in lukewarm water. His food ought to be very light; and he fhould be as little difturbed with company as poffible.

Much mifchief is done at this period by confining the patient too foon to his bed, and plying him with warm cordials or fudorific medicines. Every thing that heats and inflames the blood increafes the fever, and pufhes out the puftules prematurely. This has numberlefs ill effects. It not only increafes the number of puftules, but likewife tends to make them run into one another; and when they have been pufhed out with too great violence, they generally fall in before they come to maturity.

The good women, as foon as they fee the fmall-pox begin to appear, commonly ply their tender charge with cordials, faffron, and marigold-teas, wine, punch, and even brandy itfelf. All theie are given with a view, as they term it, to throw out the eruption from the heart. This, like most other popular mistakes, is the abuse of a very just observation, that when there is a moisture on

* Convultion fits are no doubt very alarming, but their effects are often falutary. They feem to be one of the means made use of by Nature for breaking the force of a fever. I have always obferved a fever abated, and fometimes quite removed, after one or more convultion-fits. This readily accounts for convultions being a favourable fymptom in the fever which precedes the eruption of the fmall-pox, as every thing that mitigates this fever lef. fens the eruption.

the fkin, the pox rife better, and the patient is eafter, than when it continues dry and parched. But that is no reafon for forcing the patient into a fweat. Sweating never relieves unlefs where it comes fpontaneoufly, or is the effect of drinking weak diluting liquors. The patient ought to have no more covering in bed than is neceffary to prevent his catching cold, and fhould be frequently taken up, to keep him cool, and prevent too great a flux of blood towards the head.

Children are often fo peevifh, that they will not lie a-bed without a nurfe conflantly by them. Indulging them in this, we have reafon to believe, has many bad effects both upon the nurfe and the child. Even the natural heat of the nurfe cannot fail to augment the fever of the child; but if fhe too proves feverifh, which is often the cafe, the danger must be increased *.

Laying feveral children who have the fmall-pox in the fame bed, has many ill confequences. They ought, if poffible never to be in the fame chamber, as the perfpiration, the heat, fmell, &c. all tend to augment the fever, and to heighten the difeafe. It is common among the poor to fee two or three children lying in the fame bed, with fuch a load of puffules that even their fkins flick together. One can hardly view a fcene of this kind without being fickened by the fight; but how muft the effluvia affect the poor patients, many of whom perifh by this ufage [†].

A very

* I have known a nurfe, who had the fmall-pox before, fo infected by lying conftantly a-bed with a child in a bad kind of fmall-pox, that fhe had not only a great number of pullules which broke cut all over her body, but afterwards a malignant fever which terminated in a number of impofthumes or boils, and from which the narrowly efcaped with her life. We mention this to put others upon their guard against the danger of this virulent infection.

† This obfervation is likewife applicable to hofpitals, workhoufes, &c. where numbers of children happen to have the fmallpox at the fame time. I have feen above forty children cooped up in one apartment, all the while they had this difeafe, without any of them being admitted to breathe the frefh air. No one can be at a lofs to fee the impropriety of fuch conduct. It ought to be a rule not only in hofpitals for the fmall-pox, but likewite for other difeafes, that no patient fhould be within fight or hearing of another

A very dirty cuftom prevails among the lower clafs of people, of allowing children in the fmall-pox to keep on the fame linen, during the whole period of that loathfome difeafe. This is done left they fhould catch cold; but it has many ill confequences. The linen becomes hard by the moifture which it abforbs, and frets the tender fkin. It likewife occafions a bad fmell, which is very pernicious both to the patient and thofe about him; befides, the filth and fordes which adhere to the linen being reforbed, or taken up again into the body, greatly augment the difeafe.

A patient fhould not be fuffered to be dirty in an internal difeafe, far lefs in the fmall-pox. Cutaneous diforders are often occafioned by naftinefs alone, and are always increafed by it. Were the patient's linen to be changed every day, it would greatly refresh him. Care indeed is to be taken that the linen be thoroughly dry. It ought likewife to be put on when the patient is most cool.

So ftrong is the vulgar prejudice in this country, notwithstanding all that has been faid against the hot regimen in the fmall-pox, that numbers ftill fall a facrifice to that error. I have feen poor women travelling in the depth of winter, and carrying their children along with them in the fmall-pox, and have frequently obferved others begging by the way-fide, with infants in their arms covered with the puffules; yet I could never ·learn that one of these children died by this fort of treatment. This is certainly a fufficient proof of the fafety, at least, of exposing patients in the small-pox to the open air. There can be no reafon, however, for exposing them to public view. It is now very common in the environs of great towns to meet patients in the fmallpox on the public walks. This practice, however well it may fuit the purpofes of boafting inoculators, is dan-

another. This is a matter to which too little regard is paid. In most hospitals and infirmaries, the fick, the dying, and the dead, are often to be feen in the fame apartment. gerous to the citizens, and contrary to the laws of hutmanity and found policy.

The food in this difeafe ought to be very light, and of a cooling nature, as panado, or bread boiled with equal quantities of milk and water, good apples roafted or boiled with milk, and fweetened with a little fugar or fuch like.

The drink may be equal parts of milk and water, clear fweet whey, barley-water, or thin gruel, &c. After the pox are full, butter-milk, being of an opening and cleanfing nature, is a very proper drink.

MEDICINE.——This difease is generally divided into four different periods, viz. the fever which precedes the eruption, the eruption itself, the suppuration or maturation of the puscules, and the secondary fever.

It has already been obferved, that little more is neceffary during the primary fever than to keep the patient cool and quiet, allowing him to drink diluting liquors, and bathing his feet frequently in warm water. Though this be generally the fafeft courfe that can be taken with infants, yet adults of a ftrong conflitution and plethoric habit fometimes require bleeding. When a full pulfe, a dry fkin, and other fymptoms of inflammation, render this operation neceffary, it ought to be performed; but unlefs thefe fymptoms are urgent, it is fafer to let it alone; if the body is bound, emollient clyfters may be thrown in.

If there is a great naufea or inclination to vomit, weak camomile tea or lukewarm water may be drank, in order to cleanfe the flomach. At the beginning of a fever, Nature generally attempts a difcharge, either upwards or downwards, which, if promoted by gentle means, would tend greatly to abate the violence of the difeafe.

Though every method is to be taken during the primary fever, by a cool regimen, &c. to prevent too great an eruption; yet, after the puftules have made their appearance, our bufines is to promote the suppuration, by diluting drink, light food, and, if Nature seems to flag, by generous cordials. When a low creeping pulse, faintifhnes, and great loss of strength, render cordials cordials neceffary, we would recommend good wine, which may be made into negus, with an equal quantity of water, and fharpened with the juice of orange, the jelly of currants, or the like. Wine-whey, fharpened as above, is likewife a proper drink in this cafe; great care, however, must be taken not to overheat the patient by any of these things. This, instead of promoting, would retard the eruption.

The rifing of the fmall-pox is often prevented by the violence of the fever; in this cafe the cool regimen is ftrictly to be obferved. The patient's chamber must not only be kept cool, but he ought likewife frequently to be taken out of bed, and to be lightly covered with clothes while in it.

Exceffive reftlefinefs often prevents the rifing and filling of the fmall-pox. When this happens, gentle opiates are neceffary. Thefe, however, ought always to be administered with a sparing hand. To an infant, a tea-spoonful of the syrup of poppies may be given every five or fix hours till it has the defired effect. An adult will require a table-spoonful in order to answer the same purpose.

If the patient be troubled with a ftrangury, or fupprefion of urine, which often bappens in the fmall-pox, he fhould be frequently taken out of bed, and, if he be able, fhould walk acrofs the room with his feet bare. When he cannot do this, he may be frequently fet on his knees in bed, and fhould endeavour to pais his urine as often as he can. When thefe do not fucceed, a teafpoonful of the fweet fpirits of nitre may be occafionally mixed with his drink. Nothing more certainly relieves the patient, or is more beneficial in the fmall-pox, than a plentiful difcharge of urine.

If the mouth be foul, and the tongue dry and chapped, it ought frequently to be washed, and the throat gargled with water and honey, sharpened with a little vinegar or currant-jelly.

During the rifing of the finall-pox, it frequently happens that the patient is eight or ten days without a ftool. This not only tends to heat and inflame the blood, but the forces, by lodging fo long in the body, P_2 become become acrid, and even putrid; from whence bad confequences must enfue. It will therefore be proper, when the body is bound, to throw in an emollient clyster every fecond or third day, through the whole course of the difease. This will greatly cool and relieve the patient.

When petechiæ, purple, black, or livid fpots appear among the fmall-pox, the Peruvian bark muft immediately be administered in as large dofes at the patient's ftomach can bear. For a child, two drachms of the bark in powder may be mixed in three ounces of common water, one ounce of fimple cinnamon-water, and two ounces of the fyrup of orange or lemon. This may be fharpened with the fpirits of vitriol, and a tablefpoonful of it given every hour. If it be given to an adult in the fame form, he may take at least three or four fpoonfuls every hour. This medicine ought not to be trifled with, but must be administered as frequently as the ftomach can bear it; in which cafe it will often produce very happy effects. I have frequently feen the petechiæ difappear, and the fmall-pox, which had a very threatening afpect, rife and fill with laudable matter, by the use of the bark and acids.

The patient's drink ought likewife in this cafe to be generous, as wine or ftrong negus acidulated with fpirits of vitriol, vinegar, the juice of lemon, jelly of currants, or fuch like. His food must confist of apples, roasted or boiled, preferved cherries, plums, and other fruits of an acid nature.

The bark and acids are not only neceffary when the petechiæ or putrid fymptoms appear, but likewife in the lymphatic or cryftalline fmall-pox, where the matter is thin, and not duly prepared. The Peruvian bark feems to poffefs a fingular power of affifting Nature in preparing laudable pus, or what is called good matter; confequently it must be beneficial both in this and other difeases, where the crifis depends on a fuppuration. I have often observed where the fmall-pox were flat, and the matter contained in them quite clear and transparent, and where at first they had the appearance of running into one another, that the Peruvian bark, acidulated as above,

above, changed the colour and confiftence of the matter, and produced the most happy effects.

When the eruption fubfides fuddenly, or, as the good women term it, when the fmall-pox *ftrike in*, before they have arrived at maturity, the danger is very great. In this cafe bliftering-plafters must be immediately applied to the wrifts and ancles, and the patient's fpirits fupported with cordials.

Sometimes bleeding has a furprifing effect in raifing the puftules after they have fubfided; but it requires fkill to know when this is proper, or to what length the patient can bear it. Sharp cataplafms, however, may be applied to the feet and hands, as they tend to promote the fwelling of thefe parts, and by that means to draw the humours towards the extremities.

The most dangerous period of this difease is what we call the secondary fever. This generally comes on when the small-pox begin to blacken, or turn on the face; and most of those who die of the small-pox are carried off by this fever.

Nature generally attempts, at the turn of the fmallpox, to relieve the patient by loofe ftools. Her endeavours this way are by no means to be counteracted, but promoted, and the patient at the fame time fupported by food and drink of a nourifhing and cordial nature.

If, at the approach of the fecondary fever, the pulfe be very quick, hard, and ftrong, the heat intenfe, and the breathing laborious, with other fymptoms of an inflammation of the breaft, the patient must immediately be bled. The quantity of blood to be let, must be regulated by the patient's ftrength, age, and the urgency of the fymptoms.

But in the fecondary fever, if the patient be faintifh, the puftules become fuddenly pale, and if there be great coldnefs of the extremities, bliftering-plafters must be applied, and the patient must be fupported with generous cordials. Wine, and even fpirits, have fometimes been given in fuch cafes with amazing fuccefs.

As the fecondary fever is in great measure, if not wholly, owing to the abforption of the matter, it would feem highly confonant to reason, that the pustules, as soon

213

as

as they come to maturity, fhould be opened. This is every day practifed in other phlegmons which tend to fuppuration; and there feems to be no caufe why it fhould be lefs proper here. On the contrary, we have reafon to believe that by this means the fecondary fever might always be leffened, and often wholly prevented.

The puffules fhould be opened when they begin to turn of a yellow colour. Very little art is neceffary for this operation. They may either be opened with a lancet or a needle, and the matter abforbed by a little dry lint. As the puffules are generally first ripe on the face, it will be proper to begin with opening these, and the others in course as they become ripe. The puffules generally fill again, a fecond, or even a third time; for which cause the operation must be repeated, or rather continued as long as there is any confiderable appearance of matter in the pushules.

We have reafon to believe that this operation, rational as it is, has been neglected from a piece of miltaken tendernefs in parents. They believe that it must give great pain to the poor child; and, therefore, would rather fee it die than have it thus tortured. This notion, however, is entirely without foundation. I have frequently opened the pustules when the patient did not fee me, without his being in the least fensible of it; but suppose it were attended with a little pain, that is nothing in comparison to the advantages which may arise from it.

Opening the puftules not only prevents the reforption of the matter into the blood, but likewife takes off the tenfion of the fkin, and by that means greatly relieves the patient. It likewife tends to prevent the pitting, which is a matter of no fmall importance. Actid matter, by lodging long in the puftules, cannot fail to corrode the tender fkin; by which many a handfome face becomes fo deformed as hardly to bear a refemblance to the human figure *.

• Though this operation can never do harm, yet it is only neceffary when the patient has a great load of fmall-pox, or when the matter which they contain is of fo thin and acrid a nature, that there is reafon to apprehend bad confequences from its being too quickly reforbed, or taken up again into the mafs of circulating humours.

It is generally neceffary, after the fmall-pox are gone off, to purge the patient. If, however, the body has been open through the whole course of the difease, or if butter-milk and other things of an opening nature have been drank freely after the height of the fmall-pox, purging becomes lefs neceffary; but it ought never wholly to be neglected.

For very young children, an infusion of fenna and prunes, with a little rhubarb, may be fweetened with coarfe fugar, and given in fmall quantities till it operates. Those who are farther advanced must take medicines of a fharper nature. For example, a child of five or fix years of age may take eight or ten grains of fine rhubarb in powder overnight, and the fame quantity of jalap in powder next morning. This may be wrought off with fresh broth or water-gruel, and may be repeated three or four times, five or fix days intervening between each dofe. For children further advanced, and adults, the dofe must be increased in proportion to the age and constitution *.

When imposthumes happen after the small-pox, which is not feldom the cafe, they must be brought to fuppuration as foon as possible, by means of ripening poultices; and when they have been opened, or have broke of their own accord, the patient must be purged. The Peruvian bark and a milk diet will likewife be ufeful in this cafe.

When a cough, a difficulty of breathing, or other fymptoms of a confumption, fucceed to the fmall-pox, the patient must be fent to a place where the air is good, and put upon a courfe of affes' milk, with fuch exercife as he can bear. For further directions in this cafe, fee the article Confumptions.

OF INOCULATION.

Though no difease, after it is formed, baffles the power of medicine more effectually than the fmall-pox,

* I have of late been accustomed, after the fmall-pox, to give one, two, three, four, or five grains of calomel, according to the age of the patient, over night, and to work it off next morning with a fuitable dofe of jalap, Or the jalap and calomel may be mixed together, and given in the morning.

216

yet more may be done before-hand to render this difeafe favourable than any one we know, as almost all the danger from it may be prevented by inoculation. This falutary invention has been known in Europe above half a century; but, like most uleful other difcoveries, it has till of late made but flow progrefs. It must, however, be acknowledged, to the honour of this country, that inoculation has met with a more favourable reception here, than among any of our neighbours. It is still, however, far from being general, which we have reason to fear will be the cafe, as long as the practice continues in the hands of the faculty.

No difcovery can be of general utility, while the practice of it is kept in the hands of a few. Had the inoculation of the fmall-pox been introduced as a faffiion, and not as a medical difcovery, or had it been practifed by the fame kind of operators here, as it is in those countries from whence we learned it, it had long ago been universal. Fears, jealoufies, prejudices, and opposite interests are, and ever will be, the most effectual obstacles to the progress of any falutary discovery. Hence it is that the practice of inoculation never became in any measure general, even in England, till taken up by men not bred to phyfic. These have not only rendered the practice more extensive, but likewife more fafe, and by acting under lefs reftvaint than the regu-Iar practitioners, have taught them that the patient's greatest danger arose, not from the want of care, but from the excess of it.

They know very little of the matter, who impute the fuccefs of modern inoculators to any fuperior fkill, either in preparing the patient or communicating the difeafe. Some of them, indeed, from a fordid defire of engroffing the whole practice to themfelves, pretend to have extraordinary fecrets or noftrums for preparing perfons for inoculation, which never fail of fuccefs. But this is only a pretence calculated to blind the ignorant and inattentive. Common fenfe and prudence alone are fufficient, both in the choice of the fubject and management of the operation. Whoever is poffeffed of thefe may perform this this office for his children whenever he finds it convenient, provided they be in a good flate of health.

This fentiment is not the refult of theory, but of obfervation. Though few phyficians have had more opportunities of trying inoculation in all its different forms, fo little appears to me to depend on thefe, generally reckoned important circumftances, of preparing the body, communicating the infection by this or the other method, &c. that for feveral years paft I have perfuaded the parents or nurfes to perform the whole themfelves, and have found that method followed with equal fuccefs, while it is free from many inconveniences that attend the other *.

The fmall-pox may be communicated in a great variety of ways with nearly the fame degree of fafety and fuccefs. In Turkey, from whence we learned the practice, the women communicate the difeafe to children, by opening a bit of the fkin with a needle, and putting into the wound a little matter taken from a ripe puffule. On the coaft of Barbary, they pafs a thread wet with the matter through the fkin between the thumb and forefinger; and in fome of the ftates of Barbary, inoculation is performed by rubbing in the variolous matter between

* A critical fituation, too often to be met with, first put me upon trying this method. A gentleman who had loft all his children except one fon by the natural fmall-pox, was determined to have him inoculated. He told me his intention, and defired I would perfuade the mother and grandmother, &c. of its propriety. But that was impoffible. They were not to be perfuaded, and either could not get the better of their fears, or were determined against conviction. It was always a point with me not to perform the operation without the confent of the parties concerned. I therefore advised the father, after giving, his fon a dofe or two of rhubarb, to go to a patient who had the fmall-pox of a good kind, to open two or three of the pultules, taking up the matter with a little cotton, and as foon as he came home to take his fon apart, and give his arm a flight fcratch with a pin, afterwards to rub the place well with the cotton, and take no farther notice of it. All this he punctually performed : and at the ufual period the fmall-pox made their appearance, which were of an exceeding good kind, and fo mild as not to confine the boy an hour to his bed. None of the other relations knew but the difeafe had come in the natural way, till the boy was well.

the thumb and fore-finger, or on other parts of the body. The practice of communicating the fmall-pox, by rubbing the variolous matter upon the fkin has been long known in many parts of Afia and Europe as well as in Barbary, and has generally gone by the name of *buying the fmall-pox*. The fame cuftom is faid to have obtained in Wales long ago.

The prefent method of inoculating in Britain is to make two or three flanting incifions in the arm, fo fuperficial as not to pierce quite through the fkin, with a lancet wet with frefh matter taken from a ripe puffule; afterwards the wounds are clofed up, and left without any dreffing. Some make ufe of a lancet covered with the dry matter: but this is lefs certain, and ought never to be ufed unlefs where frefh matter cannot be obtained: when this is the cafe, the matter ought be moiftened by holding the lancet for fome time in the fteam of warm water*.

Indeed, if fresh matter be applied long enough to the skin, there is no occasion for any wound at all. Let a bit of thread, about half an inch long, wet with the matter, be immediately applied to the arm, midway between the shoulder and the elbow, and covered with a piece of the common flicking plaster, and kept on for eight or ten days. This will feldom fail to communicate the difease. We mention this method, because many people are afraid of a wound; and doubtless the more easily the operation can be performed, it has the greater chance to become general. Some people imagine, that the difcharge from a wound less the eruption; but there is no great stress to be laid upon this notion : besides, deep wounds often ulcerate, and become troubles forme.

We do not find that inoculation is at all confidered as a medical operation in those countries from whence we learned it. In Turkey it is performed by the women, and in the East Indies by the Brachmins or priests. In this country the custom is still in its infancy; we make no doubt, however, but it will foon become fo familiar,

* Mr. TRONCHIN communicates this difeafe by a little bit of thread dipt in the matter, which he covers with a fmall blifteringplaster. This method may no doubt be used with advantage in those cases where the patient is very much alarmed at the fight of any cutting instrument.

218

that

that parents will think no more of inoculating their children, than at prefent they do of giving them a purge.

No fet of men have it fo much in their power to render the practice of inoculation general as the clergy, the greatest opposition to it still arising from some scruples of confcience, which they alone can remove. I would recommend it to them not only to endeavour to remove the religious objections which weak minds may have to this falutary practice, but to enjoin it as a duty, and to point out the danger of neglecting to make ufe of a mean which Providence has put in our power, for faving the lives of our offspring. Surely fuch parents as wilfully neglect the means of faving their children's lives, are as guilty as those who put them to death. I wifh this matter were duly weighed. No one is more ready to make allowance for human weakness and religious prejudices, yet I cannot help recommending it, in the warmest manner, to parents to confider how great an injury they do their children, by neglecting to give them this difeafe in the early period of life.

The numerous advantages arifing from the inoculation of the fmall-pox have been pretty fully pointed out by the learned Dr. M'Kenzie, in his Hiftory of Health*. To

" " Many and great," fays this humane author, " are the dangers attending the natural infection, from all which, the inoculation is quite fecure. The natural infection may invade weak or diftempered bodies, by no means disposed for its kindly reception. It may attack them at a feafon of the year either violently hot or intenfely cold. It may be communicated from a fort of fmall-por impregnated with the utmost virulence. It may lay hold upon people unexpectedly, when a dangerous fort is imprudently imported into a maritime place. It may furprife us foon after exceffes committed in luxury, intemperance, or lewdnefs. It may likewife feize on the innocent after indifpenfable watchings, hard labour, or neceffary journeys. And is it a trivial advantage, that all these unhappy circumstances can be prevented by inoculation? By inoculation numbers are faved from deformity as well as from death. In the natural fmall-pox, how often are the fineft features, and themoft beautiful complexions, miferably disfigured? Whereas inoculation rarely leaves any ugly marks or fcars, even where the number of pufules on the face has been very confiderable, and the fymptoms by no means favourable. And many other grievous complaints that are frequently fubfequent to the natural fort, feldom

To those mentioned by the Doctor we shall only add, that fuch as have not had the fmall-pox in the early period of life are not only rendered unhappy, but likewife in a great measure unfit for fultaining many of the most useful and important offices. Few people would chufe even to hire a fervant who had not had the fmallpox, far lefs to purchase a flave, who had the chance of dying of this difeafe. How could a phyfician or a furgeon, who had never had the fmall-pox himfelf, attend others under that malady? How deplorable is the fituation of females, who arrive at mature age without having had the fmall-pox! A woman with child feldom furvives this difease : and if an infant happen to be feized with the fmall-pox upon the mother's breaft, who has not had the difeafe herfelf, the fcene must be diffreffing ! if the continue to fuckle the child, it is at the peril of her own life; and if the wean it, in all probability it will perifh. How often is the affectionate mother forced to leave her houfe, and abandon her children, at the very time when her care is most necessary? Yet, fhould parental affection get the better of her fears, the confequences would often prove fatal. I have known the tender mother and her fucking infant laid in the fame

feldom follow the artificial. Does not inoculation alfo prevent those inexpressible terrors that perpetually harafs perfons who never had this difeafe, infomuch that when the fmall-pox is epidemical, entire villages are depopulated, markets ruined, and the face of diffrefs fpread over the whole country? From this terror it arifes, that justice is frequently postponed, or discouraged, at feffions or affizes where the fmall-pox rages. Witneffes and juries dare not appear; and by reafon of the necessary absence of Iome gentlemen, our honourable and ufeful judges are not attended with that reverence and fplendour due to their office and merit. Does not inoculation, in like manner, prevent our brave failors from being feized with this diffemper on fhipboard, where they must quickly spread the infection among such of the crew who never had it before, and where they have fcarce any chance to efcape, being half stifled with the closeness of their cabins, and but very indifferently nurfed ? Laftly, with regard to the foldiery, the miferies attending these poor creatures, when attacked by the finall-pox on a march, are inconceivable, without attendance, without lodgings, without any accommodation; fo that one of three commonly perifhes."

220.

grave,

grave, both untimely victims to this dreadful malady. But these are scenes too shocking even to mention. Let parents who run away with their children to avoid the small-pox, or who refuse to inoculate them in infancy, confider to what deplorable fituations they may be reduced by this mistaken tenderness!

As the fmall-pox is now become an epidemical difeafe in most parts of the known world, no other choice remains but to render the malady as mild as poffible. This is the only manner of extirpation now left in our power; and though it may feem paradoxical, the artificial method of communicating the difeafe, could it be rendered univerfal, would amount to nearly the fame thing as rooting it out. It is a matter of fmall confequence, whether a difeafe be entirely extirpated, or rendered fo mild as neither to deftroy life nor hurt the conftitution; but that this may be done by inoculation, does not now admit of a doubt. The numbers who die under inoculation hardly deferve to be named. In the natural way, one in four or five generally dies; but by inoculation not one in a thousand. Nay, fome can boast of having inoculated ten thousand without the loss of a fingle patient.

I have often wished to see fome plan established for rendering this falutary practice universal; but am afraid I shall never be so happy. The difficulties indeed are many; yet the thing is by no means impracticable. The aim is great: no less than faving the lives of one-fourth part of mankind. What ought not to be attempted in order to accomplish fo defirable an end?

The first step towards rendering the practice universal, must be to remove the religious prejudices against it. This, as already observed, can only be done by the clergy. They must not only recommend it as a duty to others, but likewise practise it on their own children. Example will ever have more influence than precept.

The next thing requifite, is to put it in the power of all. For this purpofe, we would recommend it to the Faculty to inoculate the children of the poor gratis. It is hard that fo useful a part of mankind should, by their poverty, be excluded from such a benefit.

Should

Should this fail, it is furely in the power of any State to render the practice general, at leaft as far as their dominion extends. We do not mean that it ought to be enforced by a law. The beft way to promote it would be to employ a fufficient number of operators at the public expence to inoculate the children of the poor. This would only be neceffary till the practice became general; afterwards cuftom, the ftrongeft of all laws, would oblige every individual to inoculate his children to prevent reflections.

It may be objected to this fcheme, that the poor would refufe to employ the inoculators: this difficulty is eafily removed. A fmall premium to enable mothers to attend their children while under the difeafe, would be a fufficient inducement; befides, the fuccefs attending the operation would foon banifh all objections to it. Even confiderations of profit would induce the poor to embrace this plan. They often bring up their children to the age of ten or twelve, and when they come to be ufeful, they are fnatched away by this malady, to the great lofs of their parents, and detriment of the public.

The British legislature has of late years shewn great attention to the prefervation of infant-lives, by supporting the Foundling hospital, &c. But we will venture to fay, if one-tenth part of the fums laid out in supporting that institution, had been bestowed towards promoting the practice of inoculation of the small-pox among the poor, that not only more useful lives had been faved, but the practice, ere now, rendered quite universal in this island. It is not to be imagined what effect example and a little money will have upon the poor; yet, if left to themsfelves, they would go on for ever in the old way, without thinking of any improvement. We only mean this as a hint to the humane and public-fpirited. Should fuch a fcheme be approved, a proper plan might easily be laid down for the execution of it.

But as public plans are very difficult to bring about, and often, by the felfish views and misconduct of those entrusted with the execution of them, fail of answering the noble purposes for which they were defigned; we shall, therefore, point out some other method by which

which the benefits of inoculation may be extended to the poor.

There is no doubt but inoculators will daily become more numerous. We would, therefore, have every parifh in Britain to allow one of them a fmall annual falary for inoculating all the children of the parifh at a proper age. This might be done at a very trifling expence, and it would enable every one to enjoy the benefit of this falutary invention.

Two things chiefly operate to prevent the progrefs of inoculation. The one is a wifh to put the evil day as far off as poflible. This is a principle in our nature; and as inoculation feems rather to be anticipating a future evil, it is no wonder mankind are fo averfe to it. But this objection is fufficiently anfwered by the fuccefs. Who in his fenfes would not prefer a leffer evil to-day to a greater to-morrow, provided they were equally certain?

The other obftacle is the fear of reflections. This has a very great weight with the bulk of mankind. Should the child die, they think the world would blame them. This they cannot bear. Here lies the difficulty; and, till that be removed, inoculation will make but fmall progrefs. Nothing however can remove it but cuftom. Make the practice fashionable, and all objections will foon vanish. It is fashion alone that has led the multitude fince the beginning of the world, and will lead them to the end. We must, therefore, call upon the more enlightened part of mankind to fet a pattern to the reft. Their example, though it may for fome time meet with opposition, will at length prevail.

I am aware of an objection to this practice from the expence with which it may be attended : this is eafily obviated. We do not mean that every parifh ought to employ a Sutton or a Dimfdale as inoculators. Thefe have by their fuccefs already recommended themfelves to crowned heads, and are beyond the vulgar reach; but have not others an equal chance to fucceed? They certainly have. Let them make the fame trial, and the difficulties will foon vanifh. There is not a parifh, and hardly a village in Britain, deftitute of fome perfon who

can

can bleed. But this is a far more difficult operation, and requires both more fkill and dexterity than inoculation.

The perfons to whom we would chiefly recommend the performance of this operation are the clergy. Moft of them know fomething of medicine. Almost all of them bleed, and can order a purge, which are all the qualifications neceffary for the practice of inoculation. The priefts among the lefs enlightened Indians perform this office, and why fhould a Christian teacher think himfelf above it? Surely the bodies of men, as well as their fouls, merit a part of the paftor's care; at least the greatest Teacher who ever appeared among men, feems to have thought fo.

Should all other methods fail, we would recommend it to parents to perform the operation themfelves. Let them take any method of communicating the difeafe they pleafe; provided the fubjects be healthy, and of a proper age, they will feldom fail to fucceed to their wifh. I have known many inftances even of mothers performing the operation, and never fo much as heard of one bad confequence. A planter in one of the Weft India ifland is faid to have inoculated with his own hand, in one year, three hundred of his flaves, who, notwithflanding the warmth of the climate, and other unfavourable circumftances, all did well. Common mechanics have often, to my knowledge, performed the operation with as good fuccefs as phyficians. We do not, however, mean to difcourage those who have it in their power, from employing people of skill to inoculate their children, and attend them while under the difeafe; but only to fhew, that where fuch cannot be had, the operation ought not upon that account to be neglected.

Inftead of multiplying arguments to recommend this practice, I fhall juft beg leave to mention the method which I took with my own fon, then an only child. After giving him two gentle purges, I ordered the nurfe to take a bit of thread which had been previoufly wet with fresh matter from a pock, and to lay it upon his arm, covering it with a piece of sticking plaster. This remained on fix or feven days, till it was rubbed off by accident. At the usual time the small-pox made their appearance,

appearance, and were exceedingly favourable. Surely this, which is all that is generally neceffary, may be done without any fkill in medicine.

We have been the more full on this fubject becaufe the benefits of inoculation cannot be extended to fociety by any other means than making the practice general. While it is confined to a few, it must prove burtful to the whole. By means of it the contagion is fpread, and is communicated to many who might otherwife never have had the difeafe. Accordingly it is found that nearly the fame number die of the fmall-pox now as before inoculation was introduced; and this important difcovery, by which alone more lives might be faved than by all the endeavours of the Faculty, is in a great meafure lost by its benefits not being extended to the whole community *.

The fpring and autumn have been ufually reckoned the moft proper feafons for inoculation, on account of the weather being then moft temperate; but it ought to be confidered that thefe are generally the moft unhealthy feafons of the whole year. Undoubtedly the beft preparation for the difeafe is a previous good flate of health. I have always obferved that children in particular are more fickly towards the end of fpring and autumn than at any other time of the year. On this account, as well as for the advantage of cool air, I would propofe winter as the moft proper feafon for inoculation; though on every other confideration, the fpring would feem to be preferable.

The most proper age for inoculation is between three and five. Many approve of inoculating on the breast, and where no circumstances forbid this practice, I have no objection to it. Children, however, are more liable to convulsions at this time than afterwards; besides, the anxiety of the mother or nurse, should the child be in danger, would not fail to heighten it by spoiling the milk.

Children who have constitutional diseafes, must never-

• By a well-laid plan for extending inoculation, more lives might be faved at a fmall expence, than are at prefent preferved by all the hofpitals in England, which coft the public fuch an amazing fum.

thelefs

theless be inoculated. It will often mend the habit of body; but ought to be performed at a time when they are most healthy. Accidental diseases should always be removed before inoculation.

It is generally thought neceffary to regulate the diet for fome time before the difeafe be communicated. In children, however, great alteration in diet is feldom neceffary, their food being commonly of the most fimple and wholefome kind, as milk, water-pap, weak broths, bread, light pudding, mild roots, and white meats.

But children who have been accustomed to a richer diet, who are of a gross habit, or abound with bad humours, ought to be put upon a spare diet before they are inoculated. Their food should be of a light cooling nature, and their drink whey, butter-milk, and such like.

We would recommend no other medicinal preparation but two or three mild purges, which ought to be fuited to the age and ftrength of the patient. The fuccefs of inoculators does not depend on the preparation of their patients, but on their management of them while under the difeafe. Their conftant care is to keep them cool, and their bodies gently open, by which means the fever is kept low, and the eruption greatly leffened. The danger is feldom great when the puffules are few ; and their number is generally in proportion to the fever which precedes and attends the eruption. Hence the chief fecret of inoculation confifts in regulating the eruptive fever, which generally may be kept fufficiently low by the methods mentioned above.

The regimen during the difeafe is in all refpects the fame as under the natural finall-pox. The patient muft be kept cool, his diet fhould be light, and his drink weak and diluting, &c. Should any bad fymptoms appear, which is feldom the cafe, they muft be treated in the fame way as directed in the natural finall-pox. 1 urging is not lefs neceffary after the finall-pox by inoculation, than in the natural way, and ought by no means to be neglected.

I have already hinted how great a misfortune it was, that inoculation was first introduced into this country as a medical operation. Had Lady WORTLEY MONTAGUE brought

brought it in as a fashion, her own noble example and that of her friends would foon have rendered it popular; but while it remains in the hands of the Faculty, it cannot be generally beneficial to mankind. Though the practice lays claim to the greatest antiquity, it is nowhere confined to medical men, in the ftrict fenfe of the word, but in Europe. Mr. HOLWELL fays, that, in India, it is next to a miracle to hear that one in a million fails of receiving the infection, or fuffers any injury from it, although the business of inoculation is there performed by the bramins or priefts.

Though their practice, as defcribed by this very candid and fenfible writer, is blended with much fuperftition, and the parade of ufeless formalities, yet their fuccefs proves it to be fubftantially good, and that their conduct is in that respect highly laudable. Did the clergy of Europe follow their conduct in the effential part, inoculation would foon become general, and millions of lives would be annually faved. What a shame that christian pastors, whose master went about curing diseases, should suffer themselves to be so far outdone by the difciples of CONFUCIUS! I feel a pleafure in doing justice to a few of our clergy, who have taken the lead in this career of humane exertion; and I hope the influence of their example will fpread itfelf among the whole order of their clerical brethren.

It is evident, from the fuccels of the bramins, that medical skill is by no means necessary for the inoculation of the fmall-pox. They administer no medicine either before or after the operation, and only enjoin abstinence from certain articles of diet, which they think might prove injurious to the patient. They lay the whole ftrefs upon what they confider as a proper regimen, . both previous to the eruption, and during its continuance; and though some, even of their few restrictions, appear to me unneceffary, yet I will venture to affert, that a proper regimen is all that is requifite for the fuccelsful inoculation of the fmall-pox.

I am old enough to remember the time when the fuccels of inoculation was supposed to be entirely owing to the preparation of the body, as it was called; but I am convinced convinced that fuch preparation always has done, and ftill does more harm than good. The body cannot be better prepared to meet a difeafe, than by being in good health. Medicine may cure a difeafe; but it cannot mend good health. When a perfon enjoys this bleffing, he ought never to meddle with medicine on any account whatever.

Should the clergy decline the benevolent and chriftianlike office propoled, I have already expressed my opinion that the flate ought to undertake it; and the operators I would recommend, are mothers. A fmall premium to a mother for every child she either inoculated herfelf, or caused to be inoculated, would soon render the practice general; and then no premium would be necessary. I will venture to fay, that ten shillings per head would be a sufficient inducement to mothers in an humble sphere to inoculate their children. I take it for granted that perfons in a higher rank do so already.

Let me now afk, whether the trifling premium propofed can be put in competition with the immenfe advantages that would refult from it; with the number of lives that would be faved, and with the improvement it muft make, not only in the looks, but in the conflitutions of the great body of the people? Numbers of conflitutions are ruined by the fmall-pox, even when it does not prove immediately fatal; and how many more do we meet with, who are left by its ravages mere fpectacles of deformity? It was the defire of preventing the latter which gave rife to inoculation, and I believe no man of common fenfe will deny, that whatever can improve the human face, as well as the human form, is deferving of the greateft encouragement and attention.

The new method of vaccine inoculation has my warmelt wifhes for its ultimate fuccefs. The eafe, fafety, and fimplicity of the procefs are ftrong circumftances in its favour; but time alone can eftablish its efficacy. Should it prove a certain preventive against catching the infection in the natural way, the promulgators of fo valuable a discovery will be justly entitled to a very high rank among the benefactors of the human race.

2312 [1229]

day from the time

CHAP. XXIV.

OF THE MEASLES.

THE measles appeared in Europe about the fame time with the Imall-pox, and have a great affinity to that difeafe. They both came from the fame quarter of the world, are both infectious, and feldom attack the fame perfon more than once. The meafles are most common in the fpring feafon, and generally difappear in fummer. The difeafe itfelf, when properly managed, feldom proves fatal; but its confequences are often very troublesome.

CAUSE. This difeafe, like the fmall-pox, proceeds from infection, and is more or lefs dangerous according to the conflitution of the patient, the featon of the year, the climate, &c.

SYMPIOMS .---- The meafles, like other fevers, are preceded by alternate fits of heat and cold, with ficknefs, and lofs of appetite. The tongue is white, but generally moift. There is a fhort cough, a heavinefs of the head and eyes, drowfinefs, and a running at the nofe. Sometimes, indeed, the cough does not come before the eruption has appeared. There is an inflammation and heat in the eyes, accompanied with a defluxion of fharp rheum, and great acutenels of fensation, fo that they cannot bear the light without pain. The eye-lids frequently fwell fo as to occafion blindnefs. The patient generally complains of his throat; and a vomiting or loofenefs often precedes the eruption. The ftools in children are commonly greenifh; they complain of an itching of the fkin, and are remarkably peevifh. Bleeding at the nofe is common, both before and in the progress of the difeafe.

About the fourth day, fmall fpots, refembling fleabites, appear, first upon the face, then upon the breast. and afterwards on the extremities: these may be diftinguished from the small-pox by their scarcely rising above the fkin. The fever, cough, and difficulty of breathing, instead of being removed by the eruption, as in the fmall-pox, are rather increased; but the vomitting generally ceafes.

About

About the fixth or feventh day from the time of fickening, the meafles begin to turn pale on the face, and afterwards upon the body; fo that by the ninth day they entirely difappear. The fever, however, and difficulty of breathing, often continue, efpecially if the patient has been kept upon too hot a regimen. Petechiæ, or purple fpots, may likewife be occafioned by this error.

A violent loofeness fometimes fucceeds the measles; in which case the patient's life is in imminent danger.

Such as die of the measles, generally expire about the ninth day from the invasion, and are commonly carried off by a peripneumony, or inflammation of the lungs.

The most favourable fymptoms are a moderate loofenefs, a moist skin, and a plentiful discharge of urine.

When the eruption fuddenly falls in, and the patient is feized with a delirium, he is in the greateft danger. If the meafles turn too foon of a pale colour, it is an unfavourable fymptom, as are alfo great weaknefs, vomiting, reftleffnefs, and difficulty of fwallowing. Purple or black fpots appearing among the meafles, are very unfavourable. When a continual cough, with hoarfenefs, fucceeds the difeafe, there is reafon to fulpect an approaching confumption of the lungs.

Our bufinefs in this difeafe is to affift Nature, by proper cordials, in throwing out the eruption, if her efforts be too languid; but when they are too violent, they muft be reftrained by evacuations, and cool diluting liquors, &c. We ought likewife to endeavour to appeale the most urgent symptoms, as the cough, reftleffnefs, and difficulty of breathing.

REGIMEN. — The cool regimen is neceffary here as well as in the fmall-pox. The food too muft be light, and the drink diluting. Acids, however, do not anfwer fo well in the meafles as in the fmall-pox, as they tend to exafperate the cough. Small-beer likewife, though a good drink in the Imall-pox, is here improper. The moft fuitable liquors are decoctions of liquorice with marfh-mallow roots and farfaparilla, infufions of linfeed or of the flowers of elder, balm-tea, clarified whey, barley-water, and fuch like. Thefe, if the patient be coftive, may be fweetened with honey; or, if that fhould

OF THE MEASLES.

231

fhould difagree with the ftomach, a little manna may occafionally be added to them.

MEDICINE. — The meafles being an inflammatory difeafe, without any critical difcharge of matter, as in the fmall-pox, bleeding is commonly neceffary, efpecially when the fever runs high, with difficulty of breathing, and great oppreffion of the breaft. But if the difeafe be of a mild kind, bleeding may be omitted *.

Bathing the feet and legs frequently in lukewarm water both tends to abate the violence of the fever, and to promote the eruption.

The patient is often greatly relieved by vomiting. When there is a tendency this way, it ought to be promoted by drinking lukewarm water, or weak camomile-tea.

When the cough is very troublefome, with drynefs of the throat, and difficulty of breathing, the patient may hold his head over the fteam of warm water, and draw the fteam into his lungs.

He may likewife lick a little fpermaceti and fugar-candy pounded together; or take now and then a fpoonful of the oil of fweet almonds, with fugar-candy diffolved in it. Thefe will foften the throat, and relieve the tickling cough.

If at the turn of the difeafe the fever affumes new vigour, and there appears great danger of fuffocation, the patient muft be bled according to his ftrength, and bliftering-plafters applied, with a view to prevent the load from being thrown on the lungs, where, if an inflammation fhould fix itfelf, the patient's life will be in imminent danger.

In cafe the meafles fhould fuddenly difappear, it will be neceffary to purfue the fame method which we have recommended when the fmall-pox recede. The patient must be fupported with wine and cordials. Bliftering-plasters must be applied to the legs and arms, and the body rubbed all over with warm flannels. Warm poultices may likewife be applied to the feet and palms of the hands.

When purple or black fpots appear, the patient's drink fhould be fharpened with fpirits of vitriol; and

^{*} I do not know any difeafe wherein bleeding is more neceffary than in the meafles, efpecially when the fever runs high : in this cafe I have always found it relieve the patient.

if the putrid fymptoms increase, the Peruvian bark must be administered in the same manner as directed in the fmall-pox.

Opiates are fometimes neceffary, but fhould never be given except in cafes of extreme reftleffnefs, a violent loofenefs, or when the cough is very troublefome. For children, the fyrup of poppies is fufficient. A tea-fpoonful or two may be occafionally given, according to the patient's age, or the violence of the fymptoms.

After the measles are gone off, the patient ought to be purged. This may be conducted in the same manner as directed in the small-pox.

If a violent loofenels fucceed the meafles, it may be checked by taking for fome days a gentle dole of thubarb in the morning, and an opiate over night; but if these do not remove it, bleeding will feldom fail to have that effect.

Patients recovering after the meafles fhould be careful what they eat or drink. Their food for fome time ought to be light, and in fmall quantities, and their drink diluting, and rather of an opening nature, as butter-milk, whey, and fuch like. They ought alfo to beware of exposing themselves too foon to the cold air, left a fuffocating catarrh, an afthma, or a confumption of the lungs, should ensue.

Should a cough, with difficulty of breathing, and other fymptoms of a confumption, remain after the meafles, fmall quantities of blood may be frequently let at proper intervals, as the patient's ftrength and conflication will permit. He ought likewife to drink affes' milk, to remove to a free air, if in a large town, and to ride daily on horfeback. He must keep close to a diet confisting of milk and vegetables; and lastly, if these do not fucceed, let him remove to a warmer climate *.

* Attempts have been made to communicate the meafles. as well as the fmall-pox, by inoculation, and we make no doubt but in time the practice may fucceed. Dr. Home of Edinburgh fays, he communicated the difeafe by the blood. Others have tried this method, and have not found it fucceed. Some think the difeafe would be more certainly communicated by rubbing the fkin of a patient who has the meafles with cotton, and afterwards applying the cotton to a wound as in the fmall-pox; while others recommend

[233]

OF THE SCARLET FEVER.

The fcarlet fever is fo called from the colour of the patient's fkin, which appears as if it were tinged with red wine. It happens at any feafon of the year, but is most common towards the end of fummer: at which time it often feizes whole families; children and young perfons are most fubject to it.

It begins, like other fevers, with coldness and shivering, without any violent fickness. Afterwards the skin is covered with red spots, which are broader, more florid, and less uniform than the meass. They continue two or three days, and then disappear; after which the cuticle, or scarf-skin, falls off.

There is feldom any occasion for medicine in this difeafe. The patient ought, however, to keep within doors, to abstain from flesh, strong liquors, and cordials, and to drink freely of cool diluting liquors. If the fever runs high, the body must be kept gently open by emollient clysters, or small doses of nitre and rhubarb. A fcruple of the former, with five grains of the latter, may be taken thrice a-day, or oftener, if neceffary.

Children and young perfons are fometimes feized at the beginning of this difeafe with a kind of flupor and epileptic fits. In this cafe, the feet and legs flould be bathed in warm water, a large bliftering-plaster applied to the neck, and a dofe of the fyrup of poppies given every night till the patient recovers *.

The fcarlet fever, however, is not always of fo mild a nature. It is fometimes attended with putrid or malig-

* Sydenham.

mend a bit of flannel which had been applied to the patient's fkin all the time of the difeafe, to be afterwards laid upon the arm or leg of the perfon to whom the intection is to be communicated. There is no doubt but this difeafe, as well as the fmall-pox, may be communicated various ways; the molt probable, however, is either from cotton rubbed upon the fkin, as mentioned above, or by introducing a little of the tharp humour which diftils from the eyes of the patient into the blood. It is agreed on all hands, that fuch patients as have been inoculated had the difeafe very mildly; we therefore wifh the practice were more general, as the meafles have of late become very fatal.

OF THE SCARLET FEVER.

234

nant fymptoms, in which cafe it is always dangerous. In the malignant fcarlet fever, the patient is not only affected with coldnefs and fhivering, but with languor, ficknefs, and great opprefilion; to thefe fucceed exceffive heat, naufea, and vomiting, with a forenefs of the throat; the pulfe is extremely quick, but fmall and deprefied; the breathing frequent and laborious; the fkin hot, but not quite dry; the tongue moift, and covered with a whitifh mucus; the tonfils inflamed and ulcerated. When the eruption appears, it brings no relief; on the contrary, the fymptoms generally grow worfe, and frefh ones come on, as purging, delirium, &c.

When this difeafe is miltaken for a fimple inflammation, and treated with repeated bleedings, purging and cooling medicines, it generally proves fatal. The only medicines that can be depended on in this cafe, are cordials and antifeptics, as the Peruvian bark, wine, fnakeroot, and the like. The treatment must be in general fimilar to that of the putrid fever, or of the malignant ulcerous fore throat *.

OF THE BILIOUS FEVER.

When a continual, remitting, or intermitting fever is accompanied with a frequent or copious evacuation of bile, either by vomit or flool, the fever is denominated bilious. In Britain the bilious fever generally makes its appearance about the end of fummer, and ceafes towards the approach of winter. It is most frequent and fatal in warm countries, especially where the foil is marshy, and when great rains are succeeded by fultry heats. Perfons who work without doors, lie in camps, or who are exposed to the night-air, are most liable to this kind of fever.

* In the year 1774, during winter, a very bad fpecies of this fever prevailed in Edinburgh It raged chiefly among young people. The eruption was generally accompanied with a quinfey, and the inflammatory fymptoms were fo blended with others of a putrid nature, as to render the treatment of the difeafe very difficult. Many of the patients, towards the decline of the fever, were afflicted with large fwellings of the fubmaxillary glands, and not a few had a fuppuration in one or both ears.

OF THE BILIOUS FEVER.

If there are fymptoms of inflammation at the beginning of this fever, it will be neceffary to bleed, and to put the patient upon the cool diluting regimen recommended in the inflammatory fever. The faline draught may likewife be frequently administered, and the patient's body kept open by clysters or mild purgatives. But if the fever should remit or intermit, bleeding will feldom be neceffary. In this cafe a vomit may be administered, and, if the body be bound, a gentle purge; after which the Peruvian bark will generally complete the cure.

In cafe of a violent loofenefs, the patient must be fupported with chicken broths, jellies of hartshorn, and the like; and he may use the *white decoction* for his ordinary drink.*. If a bloody flux should accompany this fever, it must be treated in the manner recommended under the article *Dysentery*.

When there is a burning heat, and the patient does not fweat, that evacuation may be promoted by giving him, three or four times a-day, a table-fpoonful of Mindererus's fpirit † mixed in a cup of his ordinary drink.

If the bilious fever be attended with the nervous, malignant, or putrid fymptoms, which is fometimes the cafe, the patient must be treated in the fame manner as directed under these diseases.

After this fever, proper care is neceffary to prevent a relapfe. For this purpole the patient, effectially towards the end of autumn, ought to continue the ule of the Peruvian bark for fome time after he is well. He fhould likewife abstain from all trashy fruits, new liquors, and every kind of flatulent aliment.

Though few fevers bear bleeding better than that which accompanies the meafles, yet the lancet is not to be used at random, and without a strict attention to the progress of the difease. If the symptoms run high, with a full, hard pulse, and other figns of inflammation, bleeding will be proper, but not otherwise.

I have looked at fevers, as well as at other diforders, for many years: yet, were any one to afk me, what was good

* See Appendix, White Decolion.

+ See Appendix, Spirit of Mindererus.

OF THE BILIOUS FEVER.

236

for a fever, I could not tell him, without knowing the particulars of the patient's cafe. There cannot be a groffer error than that of prefcribing to the general name of a difeafe, though thousands of people in this country fwallow drugs every day on no better ground.

Nor are the inhabitants of Britain the only dupes to this notion. I had a patient very lately, a young man from a neighbouring kingdom, who, after confulting me for his own complaints, which were chiefly imaginary, requefted that I would prefcribe for his father and brother, neither of whom I had ever feen. When I told him the abfurdity of doing it, he went away feemingly much difappointed, and, I dare fay, with a far lower opinion of my abilities than he had conceived from report.

CHAP. XXV.

OF THE ERYSIPELAS, OR ST. ANTHONY'S FIRE.

THIS difeafe, which in fome parts of Britain is called the rofe, attacks perfons at any period of life, but is most common between the age of thirty and forty. Perfons of a fanguine or plethoric habit are most liable to it. It often attacks young people, and pregnant women; and fuch as have once been afflicted with it are very liable to have it again. Sometimes it is a primary difeafe, and at other times only a fymptom of fome other malady. Every part of the body is liable to be attacked by an eryfipelas, but it most frequently feizes the legs or face, efpecially the latter. It is most common in autumn, or when hot weather is fucceeded by cold and wet.

CAUSES. — The eryfipelas may be occafioned by violent paffions or affections of the mind; as fear, anger, &c. When the body has been heated to a great degree, and is immediately exposed to the cold air, fo that the perfpiration is fuddenly checked, an eryfipelas will often enfue *. It may also be occafioned by drinking to excefs,

* The country people in many parts of Britain call this difeau a blass, and imagine it proceeds from foul air, or ill wind, as they

OF THE ERYSIPELAS.

cefs, by continuing too long in a warm bath, or by any thing that overheats the blood. If any of the natural evacuations be obstructed, or in too finall quantity, it may cause an erysipelas. The same effect will follow from the stoppage of artificial evacuations; as issues, fetons, or the like.

SYMPTOMS. — The eryfipelas attacks with a fhivering, thirft, lofs of ftrength, pain in the head and back, heat, reftleffmels, and a quick pulle; to which may be added vomiting, and fometimes a delirium. On the fecond, third, or fourth day, the part fwells, becomes red, and forall pultules appear; at which time the fever generally abates.

When the eryfipelas feizes the foot, the parts contiguous fwell, the fkin fhines; and, if the pain be violent, it will afcend to the leg, and will not bear to be touched.

When it attacks the face, it fwells, appears red, and the fkin is covered with fmall puftules filled with clear water. One or both eyes are generally clofed with a fwelling; and there is a difficulty of breathing. If the mouth and noftrils be very dry, and the patient drowfy, there is reafon to fulpect an inflammation of the brain.

If the eryfipelas affects the breaft, it fwells and becomes exceedingly hard, with great pain, and is apt to fuppurate. There is a violent pain in the arm-pit on the fide affected, where an abfcefs is often formed.

If in a day or two the fwelling fubfides, the heat and pain abate, the colour of the part turns yellow, and the cuticle breaks and falls off in fcales, the danger is over.

The event of this difeafe depends greatly upon the conflitution of the patient. It is feldom dangerous; but when the conflitution is bad, the legs will fometimes fwell to a prodigious fize, and the cure prove extremely difficult. It has often proved fatal to people in the decline of life, who were of a fcorbutic habit, or whofe

they term it. The truth is, they often lie down to reft them, when warm and fatigued, upon the damp ground, where they fall afleep, and lie fo long as to catch cold, which occasions the eryfipelas. This difeafe may indeed proceed from other caufes, but we may venture to fay, that nine times out of ten it is occasioned by cold caught after the body has been greatly heated or fatigued.

persitor, april

100.57

humours were vitiated by irregular living, or unwholefome diet.

When the eryfipelas is large, deep, and affects a very fenfible part of the body, the danger is great. If the red colour changes into a livid or black, it will end in a mortification. Sometimes the inflammation cannot be difcuffed, but comes to a fuppuration; in which cafe fiftulas, a gangrene, or mortification, often enfue.

Such as die of this difeafe are commonly carried off by the fever, which is attended with difficulty of breathing, and fometimes with a delirium and great drowfinefs. They generally die about the feventh or eighth day.

REGIMEN. — In the eryfipelas the patient muft neither be kept too hot nor too cold, as either of thefe extremes will tend to make it retreat, which is always to be guarded againft. When the difeafe is mild, it will be fufficient to keep the patient within doors, without confining him to his bed, and to promote the perfpiration by diluting liquors, &c.

The diet ought to be flender, and of a moderately cooling and moiftening quality, as groat-gruel, panado, chicken or barley-broth, with cooling herbs and fruits, &c. avoiding flefh, fifh, ftrong drink, fpices, pickles, and all other things that may heat and inflame the blood; the drink may be barley-water, an infufion of elderflowers, common whey, and fuch like.

But if the pulfe be low, and the fpirits funk, the patient must be supported with negus, and other things of a cordial nature. His food may be fago-gruel, with a little wine, and nourishing broths, taken in small quantities, and often repeated. Great care, however, must be taken not to overheat him.

repel the matter too quickly. The eryfipelas in many refpects refembles the gout, and is to be treated with the greatest caution. Fine wool, or very fost flannel, are the fafest applications to the part. These not only defend it from the external air, but likewise promote the perspiration, which has a great tendency to carry off the difease. In Scotland the common people generally apply a mealy cloth to the parts affected, which is far from being improper.

It is common to bleed in the eryfipelas; but this likewife requires caution. If, however, the fever be high, the pulfe hard and ftrong, and the patient vigorous, it will be proper to bleed; but the quantity must be regulated by thefe circumstances, and the operation repeated as the fymptoms may require. If the patient has been accustomed to strong liquors, and the difease attacks his head, bleeding is absolutely necessary.

Bathing the feet and legs frequently in lukewarm water, when the difeafe attacks the face or brain, has an excellent effect. It tends to make a derivation from the head, and feldom fails to relieve the patient. When bathing proves ineffectual, poultices, or fharp finapifms, may be applied to the foles of the feet, for the fame purpofe.

In cafes where bleeding is requifite, it is likewife neceffary to keep the body open. This may be effected by emollient clyfters, or fmall dofes of nitre and rhubarb. Some indeed recommend very large dofes of nitre in the eryfipelas; but nitre feldom fits eafy on the ftomach when taken in large dofes. It is, however, one of the beft medicines when the fever and inflammation run high. Half a drachm of it, with four or five grains of rhubarb, may be taken in the patient's ordinary drink, three or four times a-day.

When the eryfipelas leaves the extremities, and feizes the head, fo as to occafion a delirium or ftupor, it is abfolutely neceffary to open the body. If clyfters and mild purgatives fail to have this effect, ftronger ones must be given. Bliftering-plafters must likewife be applied to the neck, or behind the ears, and fharp cataplafms laid to the foles of the feet.

When the inflammation cannot be difcuffed, and the part has a tendency to ulcerate, it will then be proper

to

to promote fuppuration, which may be done by the application of ripening poultices, with faffron, warm fomentations, and fuch like.

When the black, livid, or blue colour of the part fhews a tendency to mortification, the Peruvian bark muft be administered. It may be taken along with acids, as recommended in the fmall-pox, or in any other form more agreeable to the patient. It must not, however, be trifled with, as the patient's life is at stake. A drachm may be given every two hours, if the fymptoms be threatening, and cloths dipped in warm camphorated spirits of wine, or the tincture of myrrh and aloes, may be applied to the part, and frequently renewed. It may likewife be proper in this case to apply poultices of the bark, or to foment the part affected with a strong decoction of it.

In what is commonly called the *fcorbutic eryfipelas*, which continues for a confiderable time, it will only be neceffary to give gentle laxatives, and fuch things as purify the blood and promote the perfpiration. Thus, after the inflammation has been checked by opening medicines, the decoction of woods * may be drank, after which a courfe of bitters will be proper.

Such as are liable to frequent attacks of the eryfipelas ought carefully to guard against all violent passions; to abstain from strong liquors, and all fat, viscid, and highly-nouriss food. They should likewise take fufficient exercise, carefully avoiding the extremes of heat or cold. Their food should consist chiefly or milk, and such fruits, herbs, and roots, as are of a cooling quality; and their drink ought to be small-beer, whey, butter-milk, and such like. They should never suffer themsfelves to be long costive. If that cannot be prevented by fuitable diet, it will be proper to take frequently a gentle dose of rhubarb, cream of tartar, the lenitive electuary, or some other mild purgative.

Though I have fo ftrictly forbidden moift applications in the eryfipelas, yet I cannot prevail on people to leave them off. Whenever they fee inflammation, they think of fome poultice, embrocation, or fomentation, all of

* See Appendix, Decotion of Woods.

which

which do injury twenty times for once they are of the least fervice; and ought never to be used unless where suppuration is inevitable.

An abforbent, which takes up the moifture and cools the fkin, anfwers the purpofe much better. What I generally use is hair-powder, fpread upon a fost rag, and laid over the parts affected. This may be renewed twice or thrice a-day; and it is not to be imagined what eafe and comfort it gives to the patient every time.

As the eryfipelas refembles the gout in many refpects, it ought not to be rafhly tampered with. Should it be driven from the part affected, it may fix upon a more dangerous one. The alarm is generally greateft, when it removes to, or attacks the face. I have, however, known it feize upon the knee, and, after laying the bones bare, prove fatal.

CHAP. XXVI.

OF THE PHRENITIS, OR INFLAMMATION. OF THE BRAIN.

THIS is fometimes a primary difeafe, but oftener only a fymptom of fome other malady, as the inflammatory, eruptive, or fpotted fever, &c. It is very common, however, as a primary difeafe in warm climates, and is most incident to perfons about the prime or vigour of life. The passionate, the studious, and those whose nervous fystem is irritable in a high degree, are most liable to it.

CAUSES.——This difeafe is often occafioned by night-watching, efpecially when joined with hard ftudy: it may likewife proceed from hard drinking, anger, grief, or anxiety. It is often occafioned by the ftoppage of ufual evacuations; as the bleeding piles in men, the cuftomary difcharges of women, &c. Such as imprudently expose themselves to the heat of the fun, efpecially by fleeping without doors in a hot feason, with their R heads heads uncovered, are often fuddenly feized with an inflammation of the brain, fo as to awake quite delirious. When repellents are imprudently ufed in an eryfipelas, an inflammation of the brain is fometimes the confequence. It may likewife be occafioned by external injuries, as blows or bruifes upon the head, &c.

SYMPTOMS. ———— The fymptoms which ufually precede a true inflammation of the brain are, pain of the head, rednefs of the eyes, a violent flufhing of the face, diffurbed fleep, or a total want of it, great drynefs of the fkin, coftivenefs, a retention of urine, a fmall dropping of blood from the nofe, finging of the ears, and extreme fenfibility of the nervous fyftem.

When the inflammation is formed, the fymptoms in general are fimilar to those of the inflammatory fever. The pulfe indeed is often weak, irregular, and trembling; but fometimes it is hard and contracted. When the brain itfelf is inflamed, the pulfe is always foft and low; but when the inflammation only affects the integuments of the brain, viz. the dura and pia mater, it is hard. A remarkable quickness of hearing is a common fymptom of this difease; but that feldom continues long. Another ufual fymptom is a great throbbing or pulfation in the arteries of the neck and temples. Though the tongue is often black and dry, yet the patient feldom complains of thirst, and even refuses drink. The mind chiefly runs upon fuch objects as have before made a deep impreflion on it; and fometimes, from a fullen filence, the patient becomes all of a fudden quite outrageous.

A conftant trembling and ftarting of the tendons is an unfavourable fymptom, as are alfo a fupprefion of urine; a total want of fleep; a conftant fpitting; a grinding of the teetb; which laft may be confidered as a kind of convultion. When a phrenitis fucceeds an inflammation of the lungs, of the inteftines, or of the throat, &c. it is owing to a translation of the difeafe from these parts to the brain, and generally proves fatal. This fhews the neceffity of proper evacuations, and the danger of repellents in all inflammatory difeafes.

The favourable fymptoms are, a free perfpiration, a copious difcharge of blood from the nofe, the bleeding piles,

piles, a plentiful difcharge of urine, which lets fall a copious fediment. Sometimes the difeafe is carried off by a loofenefs, and in women by an exceflive flow of the menfes.

As this difeafe often proves fatal in a few days, it requires the most speedy applications. When it is prolonged, or improperly treated, it sometimes ends in madnefs, or a kind of stupidity, which continues for life.

In the cure, two things are chiefly to be attended to, viz. to leffen the quantity of blood in the brain, and to retard the circulation towards the head.

REGIMEN.——The patient ought to be kept very quiet. Company, noife, and every thing that affects the fenfes, or diffurbs the imagination, increafes the difeafe. Even too much light is hurtful; for which reafon the patient's chamber ought to be a little darkened, and he fhould neither be kept too hot nor cold. It is not, however, neceffary to exclude the company of an agreeable friend, as this has a tendency to foothe and quiet the mind. Neither ought the patient to be kept too much in the dark, left it fhould occafion a gloomy melancholy, which is too often the confequence of this difeafe.

The patient must, as far as possible, be foothed and humoured in every thing. Contradiction will ruffle his mind, and increase his malady. Even when he calls for things which are not to be obtained, or which might prove hurtful, he is not to be politively denied them, but rather put off with the promife of having them as foon as they can be obtained, or by fome other excufe. A little of any thing that the mind is fet upon, though not quite proper, will hurt the patient less than a positive refusal. In a word, whatever he was fond of, or used to be delighted with, when in health, may here be tried; as pleafing ftories, foft mufic, or whatever has a tendency to foothe the paffions, and compose the mind. Boerhaave propofes feveral mechanical experiments for this purpofe; as the foft noife of water diftilling by drops into a bafon, and the patient trying to reckon them, &c. Any uniform found, if low and continued, has a tendency to procure fleep, and confequently may be of fervice.

The aliment ought to be light, confifting chiefly of farinaceous fubftances; as panado, and water-gruel, fharpened with jelly of currants, or juice of lemons, ripe fruits roafted or boiled, jellies, preferves, &c. The drink fmall, diluting, and cooling; as whey, barley-water, or decoctions of barley and tamarinds, which latter not only render the liquor more palatable, but likewife more beneficial, as they are of an opening nature.

MEDICINE.——In an inflammation of the brain, nothing more certainly relieves the patient than a free difcharge of blood from the nofe. When this comes of its own accord, it is by no means to be flopped, but rather promoted, by applying cloths dipped in warm water to the part. When bleeding at the nofe does not happen fpontaneoufly, it may be provoked, by putting a ftraw, or any other fharp body up the noftril.

Bleeding in the temporal arteries greatly relieves the head; but as this operation cannot always be performed, we would recommend in its flead bleeding in the jugular veins. When the patient's pulfe and fpirits are fo low, that he cannot bear bleeding with the lancet, leeches may be applied to the temples. Thefe not only draw off the blood more gradually, but, by being applied nearer to the part affected, generally give more immediate relief.

A difcharge of blood from the hæmorrhoidal veins is likewife of great fervice, and ought by all means to be promoted. If the patient has been fubject to the bleeding piles, and that difcharge has been ftopped, every method must be tried to reftore it; as the application of leeches to the parts, fitting over the fteams of warm water, fharp clysters, or fuppofitories made of honey, aloes, and rock-falt.

If the inflammation of the brain be occafioned by the ftoppage of evacuations either natural or artificial, as the menfes, iffues, fetons, or fuch like, all means muft be ufed to reftore them as foon as poffible, cr to fubfitute others in their ftead.

The patient's body must be kept open by stimulating clysters or smart purges; and small quantities of nitre ought frequently to be mixed with his drink. Two or three

INFLAMMATION OF THE BRAIN. 245

three drachms, or more, if the cafe be dangerous, may be used in the space of twenty-four hours.

The head fhould be fhaved, and frequently rubbed with vinegar and rofe-water. Cloths dipped in this mixture may likewife be applied to the temples. The feet ought frequently to be bathed in lukewarm water, and foft poultices of bread and milk may be kept conftantly applied to them.

If the difeafe proves obftinate, and does not yield to thefe medicines, it will be neceffary to apply a blifteringplafter to the whole head.

I must farther observe, that, though this species of inflammation ought to be treated nearly as other inflammatory diforders are, yet more than ordinary care fhould be used to keep the patient in a state of as much ease, composure, and tranquillity as possible. A strict attention to my former hints on this head will often do more good than the beft medicines. It fhould be confidered that a fore will not bear the touch of a feather, and that the nerves of an inflamed brain are still more unfit to endure the least irritation, without torture, and without danger. Even light, found, or whatever may make a ftrong impression on the senses, is carried with such rapidity and force to the brain, as to increase the inflammation, and diforder the very organs by which it was conveyed. A docile and fenfible nurfe is, in fuch cafes, of as much fervice, as the most enlightened physician.

CHAP. XXVII.

OF THE OPHTHALMIA, OR INFLAMMATION OF THE EYES.

THIS difeafe may be occafioned by external injuries; as blows, burns, bruifes, and the like. It may likewife proceed from duft, quick-lime, or other fubftances, getting into the eyes. It is often caufed by the ftoppage of cultomary evacuations; as the healing of old fores, drying up of iffues, the fuppreffing of gentle morning R 3 fweats.

OF THE OPHTHALMIA, OR

246

fweats, or of the fweating of the feet, &c. Long expofure to the night air, efpecially in cold northerly winds, or whatever fuddenly checks the perfpiration, efpecially after the body has been much heated, is very apt to caufe an inflammation of the eyes. Viewing fnow or other white bodies for a long time, or looking ftedfaftly at the fun, a clear fire, or any bright object, will likewife occafion this malady. A fudden transition from darknefs to very bright light will often have the fame effect.

Nothing more certainly occasions an inflammation of the eyes than night-watching, efpecially reading or writing by candle-light. Drinking fpirituous liquors, and excels of venery, are likewife very hurtful to the eyes. The acrid fumes of metals, and of feveral kinds of fuel, are also pernicious. Sometimes an inflammation of the eyes proceeds from a venereal taint, and often from a fcrophulous or gouty habit. It may likewife be occafioned by hairs in the eye-lids turning inwards, and hurting the eyes. Sometimes the difeafe is epidemic, efpecially after wet feafons; and I have frequently known it prove infectious, particularly to those who lived in the fame houle with the patient. It may be occafioned by moift air, or living in low damp houfes, efpecially in perfons who are not accustomed to fuch fituations. In children it often proceeds from imprudently drying up of fcabbed heads, a running behind the ears, or any other discharge of that kind. Inflammations of the eyes often fucceed the fmall-pox or meafles, especially in children of a scrophulous habit.

SYMPTOMS.——An inflammation of the eyes is attended with acute pain, heat, rednefs, and fwelling. The patient is not able to bear the light, and fometimes he feels a pricking pain, as if his eyes were pierced with a thorn. Sometimes he imagines his eyes are full of motes, or thinks he fees flies dancing before him. The eyes are filled with a fcalding rheum, which rufhes forth in great quantities, whenever the patient attempts to look up. The pulfe is generally quick and hard, with fome degree of fever. When the difeafe is violent, the neighbouring parts fwell, and there is a throbbing or pullation in the temporal arteries, &c.

A flight

INFLAMMATION OF THE EYES. 247

A flight inflammation of the eyes, efpecially from an external caufe, is eafily cured; but when the difeafe is violent, and continues long, it often leaves fpecks upon the eyes, or dimnefs of fight, and fometimes total blindnefs.

If the patient be feized with a loofenefs, it has a good effect; and when the inflammation paffes from one eye to another, as it were by infection, it is no unfavourable fymptom. But when the difeafe is accompanied with a violent pain in the head, and continues long, the patient is in danger of lofing his fight.

REGIMEN.——The diet, unlefs in fcrophulous cafes, can hardly be too fpare, efpecially at the beginning. The patient must abstain from every thing of a heating nature. His food should confiss chiefly of mild vegetables, weak broths, and gruels. His drink may be barley-water, balm-tea, common whey, and such like.

The patient's chamber must be darkened, or his eyes fhaded by a cover, fo as to exclude the light, but not to prefs upon the eyes. He should not look at a candle, the fire, or any luminous object; and ought to avoid all smoke, as the sumes of tobacco, or any thing that may cause coughing, sneezing, or vomiting. He should be kept quiet, avoiding all violent efforts, either of body or mind, and encouraging fleep as much as possible.

MEDICINE. —— This is one of those difeases wherein great hurt is often done by external applications. Almost every perfon pretends to be possible of a remedy for the cure of fore eyes. These remedies generally confist of eye-waters and ointments, with other external applications, which do mischief twenty times for once they do good. People ought therefore to be very cautious how they use such things, as even the preffure upon the eyes often increases the malady.

Bleeding, in a violent inflammation of the eyes, is always neceffary. This fhould be performed as near the part affected as possible. An adult may lose ten or twelve ounces of blood from the jugular vein, and the operation may be repeated according to the urgency of

the

248 OF THE OPHTHALMIA, OR

the fymptoms. If it fhould not be convenient to bleed in the neck, the fame quantity may be let from the arm, or any other part of the body.

Leeches are often applied to the temples, or under the eyes, with good effect. The wounds must be fuffered to bleed for fome hours, and if the bleeding stop foon, it may be promoted by the application of cloths dipt in warm water. In obstinate cases, it will be necessary to repeat this operation feveral times.

Opening and diluting medicines are by no means to be neglected. The patient may take a fmall dofe of Glauber's falts, and cream of tartar, every fecond or third day, or a decoction of tamarinds with fenna. If thefe be not agreeable, gentle dofes of rhubarb and nitre, a little of the lenitive electuary, or any other mild purgative, will anfwer the fame end. The patient at the fame time muft drink freely of water-gruel, tea, whey, or any other weak diluting liquor. He ought likewife to take, at bed-time, a large draught of very weak wine-whey, in order to promote perfpiration. His feet and legs muft frequently be bathed in lukewarm water, and his head fhaved twice or thrice a week, and afterwards wafhed in cold water. This has often a remarkably good effect.

If the inflammation does not yield to thefe evacuations, bliftering-plafters muft be applied to the temples, behind the ears, or upon the neck, and kept open for fome time by the mild bliftering-ointment. I have feldom known thefe, if long enough kept open, fail to remove the most obstinate inflammation of the eyes; but for this purpose it is often necessary to continue the difcharge for feveral weeks.

When the difeafe has been of long ftanding, I have feen very extraordinary effects from a feton in the neck, or between the fhoulders, efpecially the latter. It fhould be put upwards and downwards, or in the direction of the fpine, and in the middle between the fhoulder blades. It may be dreffed twice a day with yellow bafilicon. I have known patients, who had been blind for a confiderable time, recover fight by means of a feton placed as above. When the feton is put acrofs the neck,

INFLAMMATION OF THE EYES. 249

neck, it foon wears out, and is both more painful and troublefome than between the fhoulders; befides, it leaves a difagreeable mark, and does not difcharge fo freely.

When the heat and pain of the eyes are very great, a poultice of bread and milk, foftened with fweet oil or fresh butter, may be applied to them, at least all night; and they may be bathed with lukewarm milk and water in the morning.

If the patient cannot fleep, which is fometimes the cafe, he may take twenty or thirty drops of laudanum, or two fpoonfuls of the fyrup of poppies, over night, more or lefs according to his age, or the violence of the fymptoms.

After the inflammation is gone off, if the eyes ftill remain weak and tender, they may be bathed every night and morning with cold water and a little brandy, fix parts of the former to one of the latter. A method fhould be contrived by which the eye can be quite immerfed in the brandy and water, where it fhould be kept for fome time. I have generally found this, or cold water and vinegar as good a ftrengthener of the eyes as any of the most celebrated collyriums.

When an inflammation of the eyes proceeds from a fcrophulous habit, it generally proves very obstinate. In this cafe the patient's diet must not be too low, and he may be allowed to drink small negus, or now and then a glass of wine. The most proper medicine is the Peruvian bark, which may either be given in substance, or prepared in the following manner:

Take an ounce of the bark in powder, with two drachms of Winter's bark, and boil them in an English quart of water to a pint: when it has boiled nearly long enough, add half an ounce of liquorice-root fliced. Let the liquor be strained. Two, three, or four table spoonfuls, according to the age of the patient, may be taken three or four times a-day. It is impossible to fay how long this medicine should be continued, as the cure is sooner performed in some than in others; but, in general, it requires a considerable time to produce any lasting effects.

Dr.

250

Dr. Cheyne fays, ' that Æthiop's mineral never fails in obstinate inflammations of the eyes, even scrophulous ones, if given in a sufficient dose, and duly persisted in.' There is no doubt but this and other preparations of mercury may be of fingular service in ophthalmias of long continuance, but they ought always to be administered with the greatest caution, or by persons of skill in physic.

It will be proper frequently to look into the eye, to fee if any hairs be turned inwards, or prefling upon them *. These ought to be removed by plucking them out with a pair of small pincers.

Those who are liable to frequent returns of this difease, ought constantly to have an issue in one or both arms. Bleeding or purging in the spring and autumn will be very beneficial to such perfons. They ought likewise to live with the greatest regularity, avoiding strong liquor, and every thing of a heating quality. Above all, let them avoid the night-air and late studies +.

Although inflammation of the eyes proceeds from a great variety of caufes, yet I find, that moft of our pretended eye-doctors treat them all nearly in the fame way. Whether the inflammation is occafioned by a blow, a fcrophulous habit, or a fpecific infection, ftill they keep fcarifying the inner coats of the eyelids, and cutting, and fcratching, till they totally extinguifh the fight. About twelve years ago, a patient of mine, troubled with an inflammation of the eyes which had arifen from a venereal taint, was not only weak enough to go to one of thofe *furgeon oculifts*, but to fay, that I was giving him calomel. The other immediately observed, that he never gave his patients mercury; and did not forget to throw out very broad hints of the impropriety of my treat-

* Any foreign body lodged in the eye may be expeditioufly removed by patting a finall hair pencil between the eye-lid and the ball of the eye. In fome places the peafants do this very effectually, by ufing their tongue in the fame manner.

† As most people are fond of using eye-waters and ointments in this and other dileases of the eyes, we have inferted some of the most approved forms of these medicines in the Appendix. See Appendix, Eye-water and Eye-falve.

ment.

INFLAMMATION OF THE EYES. 251

ment. I afterwards faw the patient, and perceiving his credulity wrought upon, I could only bid him look to the end of it. About two years after, he had occasion to go to France, when he confulted one of the most eminent medical men in Paris, who, foon difcovering one of his eyes to be irrecoverably loft, took his fee, and left the following laconic remark, inftead of a prefcription, "Your English furgeon has put your eye out ;" and, to this day, the patient wears an artificial eye.

CHAP. XXVIII.

OF THE QUINSEY, OR INFLAMMATION OF THE THROAT.

"HIS difeafe is very common in Britain, and is frequently attended with great danger. It prevails in the winter and fpring, and is most fatal to young people of a fanguine temperament.

CAUSES. — In general it proceeds from the fame caufes as other inflammatory diforders, viz. an obstructed perspiration, or whatever heats or inflames the blood. An inflammation of the throat is often occafioned by omitting fome part of the covering ufually worn about the neck, by drinking cold liquor when the body is warm, by riding or walking against a cold northerly wind, or any thing that greatly cools the throat, and parts adjacent. It may likewile proceed from the neglect of bleeding, purging, or any cultomary evacuation.

Singing, fpeaking loud and long, or whatever ftrains the throat, may likewife caufe an inflammation of that organ. I have often known the quinfey prove fatal to jovial companions, who, after fitting long in a warm room, drinking hot liquors, and finging with vehemence, were fo imprudent as to go abroad in the cold night-air. Sitting with wet feet, or keeping on wet clothes, are very apt to occafion this malady. It is likewife frequently occasioned by continuing long in a moist place, fitting near

† I

near an open window, fleeping in a damp bed, fitting in a room that has been newly plaftered, &c. I know people who never fail to have a fore throat if they fit even but a flort time in a room that has been lately washed.

Acrid or irritating food may likewife inflame the throat, and occafion a quinfey. It may alfo proceed from bones, pins, or other fharp fubftances flicking in the throat, or from the cauftic fumes of metals or minerals, as arfenic, antimony, &c. taken in by the breath. This difeafe is fometimes epidemical and infectious.

SYMPTOMS. ---- The inflammation of the throat is evident from infpection, the parts appearing red and fwelled; befides, the patient complains of pain in fwallowing. His pulfe is quick and hard, with other fymptoms of a fever. If blood be let, it is generally covered with a tough coat of a whitish colour, and the patient fpits a tough phlegm. As the fwelling and inflammation increafe, the breathing and fwallowing become more difficult; the pain affects the ears; the eyes generally appear red; and the face fwells. The patient is often obliged to keep himfelf in an erect posture, being in danger of fuffocation; there is a conftant naufea, or inclination to vomit; and the drink, inftead of paffing into the ftomach, is often returned by the nofe. The patient is fometimes starved at last, merely from an inability to fwallow any kind of food.

When the breathing is laborious with ftraitnefs of the breaft, and anxiety, the danger is great. Though the pain in fwallowing be very great, yet while the patient breathes eafy, there is not fo much danger. An external fwelling is no unfavourable fymptom : but if it fuddenly falls, and the difeafe affects the breaft, the danger is very great. When a quinfey is the confequence of fome other difeafe, which has already weakened the patient, his fituation is dangerous. A frothing at the mouth, with a fwelled tongue, a pale, ghaftly countenance, and coldnefs of the extremities, are fatal fymptoms.

REGI-

INFLAMMATION OF THE THROAT. 253

REGIMEN.——The regimen in this difeafe is in all refpects the fame as in the pleurify, or peripneumony. The food must be light, and in fmall quantity, and the drink plentiful, weak, and diluting, mixed with acids.

It is highly neceffary that the patient be kept eafy and quiet. Violent affections of the mind, or great efforts of the body may prove fatal. He fhould not even attempt to fpeak but in a low voice. Such a degree of warmth as to promote a conftant, gentle fweat, is proper. When the patient is in bed, his head ought to be raifed a little higher than ufual.

It is peculiarly neceffary that the neck be kept warm; for which purpole feveral folds of foft flannel may be wrapt round it. That alone will often remove a flight complaint of the throat, especially if applied in due time. We cannot here omit observing the propriety of a cuftom which prevails among the peafants in Scotland. When they feel any uneafiness of the throat, they wrap a flocking about it all night. So effectual is this remedy, that in many places it passes for a charm, and the flocking is applied with particular ceremonies: the cuftom, however, is undoubtedly a good one, and should never be neglected. When the throat has been thus wrapped up all night, it must not be exposed to the cold air through the day, but a handkerchief, or a piece of flannel, kept about it till the inflammation be removed.

The jelly of black currants is a medicine very much in efteem for complaints of the throat; and indeed it is of fome ufe. It fhould be almost constantly kept in the mouth, and fwallowed down leifurely. It may likewife be mixed in the patient's drink, or taken any other way. When it cannot be obtained, the jelly of red currants, or of mulberries, may be used in its stead.

Gargles for the throat are very beneficial. They may be made of fage-tea, with a little vinegar and honey, or by adding to half an English pint of the pectoral decoction two or three spoonfuls of honey, and the same quantity of currant-jelly. This may be used three or four times aday; and if the patient be troubled with tough viscid phlegm, the gargle may be rendered more sharp and cleansing, cleanfing, by adding to it a tea-fpoonful of the fpirit of *fal ammoniac*. Some recommend gargles made of a decoction of the leaves or bark of the black currantbush; but where the jelly can be had, these are unneceffary.

There is no difeafe wherein the benefit of bathing the feet and legs in lukewarm water is more apparent : that practice ought therefore never to be neglected. If people were careful to keep warm, to wrap up their throats with flannel, to bathe their feet and legs in warm water, and to ufe a fpare diet, with diluting liquors, at the beginning of this difeafe, it would feldom proceed to a great height, or be attended with any danger; but when thefe precautions are neglected, and the difeafe becomes violent, more powerful medicines are neceffary.

MEDICINE. — An inflammation of the throat being a most acute and dangerous diftemper, which fometimes takes off the patient very fuddenly, it will be proper, as foon as the fymptoms appear, to bleed in the arm, or rather in the jugular vein, and to repeat the operation if circumstances require.

The body fhould likewife be kept gently open. This may either be done, by giving the patient for his ordinary drink a decoction of figs and tamarinds, or fmall dofes of rhubarb and nitre, as recommended in the eryfipelas. Thefe may be increafed according to the age of the patient, and repeated till they have the defired effect.

I have often known very good effects from a bit of *fal prunel*, or purified nitre, held in the mouth, and fwallowed down as it melted. This promotes the difcharge of *faliva*, by which means it anfwers the end of a gargle, while at the fame time it abates the fever, by promoting the difcharge of urine, &c.

The throat ought likewife to be rubbed twice or thrice a day with a little of the volatile liniment. This feldom fails to produce fome good effects. At the fame time the neck ought to be carefully covered with wool or flannel, to prevent the cold from penetrating the fkin, as this application renders it very tender. Many other external applications are recommended in this difeafe,

INFLAMMATION OF THE THROAT. 255

as a fwallow's neft, poultices made of the fungus called Jew's ears, album Græcum, &c. But as we do not look upon any of thefe to be preferable to a common poultice of bread and milk, we fhall take no farther notice of them.

Some recommend the gum-guaiacum as a fpecific in this difeafe. Half a drachm of the gum in powder may be made into an electuary with the rob of elderberries, or the jelly of currants for a dofe, and repeated occafionally *.

Bliftering upon the neck, or behind the ears, in violent inflammations of the throat, is very beneficial; and in bad cafes, it will be neceffary to lay a bliftering-plafter quite acrofs the throat, fo as to reach from ear to ear. After the plafters are taken off, the parts ought to be kept running by the application of iffue ointment, till the inflammation is gone; otherwife, upon their drying up, the patient will be in danger of a relapfe.

When the patient has been treated as above, a fuppuration feldom happens. This, however, is fometimes the cafe, in fpite of all endeavours to prevent it. When the inflammation and fwelling continue, and it is evident that a fuppuration will enfue, it ought to be promoted by drawing the fteam of warm water into the throat through a tunnel, or the like. Soft poultices ought likewife to be applied outwardly, and the patient may keep a roafted fig conftantly in his mouth.

It fometimes happens, before the tumour breaks, that the fwelling is fo great, as entirely to prevent any thing from getting down into the ftomach. In this cafe, the patient muft inevitably perifh, unlefs he can be fupported in fome other way. This can only be done by nourifhing clyfters of broth, or gruel with milk, &c. Patients have often been fupported by thefe for feveral days, till the tumour has broke; and afterwards they have recovered.

Not only the fwallowing, but the breathing is often prevented by the tumour. In this cafe nothing can fave the patient's life, but opening the *trachea* or wind-pipe. As that has been often done with fuccefs, no perfon, in

* Dr. Home.

fuch desperate circumstances, ought to hefitate a mos ment about the operation ; but as it can only be performed by a furgeon, it is not neceffary here to give any directions about it.

When a difficulty of fwallowing is not attended with an acute pain or inflammation, it is generally owing to an obstruction of the glands about the throat, and only requires that the part be kept warm, and the throat frequently gargled with fomething that may gently flimulate the glands, as a decoction of figs with vinegar and honey; to which may be added a little mustard, or a fmall quantity of fpirits. But this gargle is never to be ufed where there are figns of an inflammation. This fpecies of angina has various names among the common people, as the pap of the throat, the falling down of the almonds of the ears, &c. Accordingly, to remove it, they lift the patient up by the hair of the head, and thrust their fingers under his jaws, &c. all which practices are at best useles, and often hurtful.

Those who are subject to inflammations of the throat, in order to avoid that difeafe, ought to live temperate. Such as do not choose to observe this rule, must have frequent recourse to purging and other evacuations, to discharge the superfluous humours. They ought likewife to beware of catching cold, and fhould abstain from aliment and medicines of an aftringent or ftimulating nature.

Violent exercife, by increasing the motion and force of the blood, is apt to occafion an inflammation of the throat, efpecially if cold liquor be drank immediately after it, or the body fuffered fuddenly to cool. Those who would avoid this difeafe ought therefore, after Speaking aloud, finging, running, drinking warm liquor, or doing any thing that may firain the throat, or increase the circulation of the blood towards it, to take care to cool gradually, and to wrap fome additional covering about their necks.

I have often known perfons who had been fubject to fore throats, entirely freed from that complaint by only wearing a ribband, or a bit of flannel, conftantly about their necks, or by wearing thicker fhoes, a flannel waiftcoats

INFLAMMATION OF THE THROAT. 257

coat, or the like. Thefe may feem trifling, but they have great effect. There is danger indeed in leaving them off after perfons have been accuftomed to them; but furely the inconveniency of using fuch things for life, is not to be compared with the danger which may attend the neglect of them.

Sometimes, after an inflammation, the glands of the throat continue fwelled, and become hard and callous. This complaint is not eafily removed, and is often rendered dangerous by the too frequent application of ftrong ftimulating and ftyptic medicines. The beft method is to keep it warm, and to gargle it twice a-day with a decoction of figs, fharpened a little with the elixir or fpirit of vitriol.

OF THE MALIGNANT QUINSEY, OR PU-TRID ULCEROUS SORE THROAT.

This kind of quinfey is but little known in the northern parts of Britain, though for fome time paft it has been fatal in the more fouthern counties. Children are more liable to it than adults, females than males, and the delicate than those who are hardy and robust. It prevails chiefly in autumn, and is most frequent after a long courfe of damp or fultry weather.

CAUSES.——This is evidently a contagious diftemper, and is generally communicated by infection. Whole families, and even entire villages, often receive the infection from one perfon. This ought to put people upon their guard against going near fuch patients as labour under the diforder; as by that means they endanger not only their own lives, but likewife those of their friends and connexions. Whatever tends to produce putrid or malignant fevers, may likewife occasion the putrid ulcerous fore throat, as unwholefome air, damaged provisions, neglect of cleanlines, &c.

SYMPTOMS. — It begins with alternate fits of fhivering and heat. The pulfe is quick, but low and unequal, and generally continues fo through the whole courfe of the difeafe. The patient complains greatly of weaknefs and oppreffion of the breaft; his fpirits are

S

low,

258 OF THE MALIGNANT QUINSEY, OR

low, and he is apt to faint away when fet upright; he is troubled with a nausea, and often with a vomiting or purging. The two latter are most common in children. The eyes appear red and watery, and the face fwells. The urine is at first pale and crude; but, as the difeafe advances, it turns more of a yellowish colour. The tongue is white, and generally moift, which diffinguishes this from an inflammatory difease. Upon looking into the throat, it appears fwelled, and of a florid red colour. Pale or afh-coloured fpots, however, are here and there interspersed, and fometimes one broad patch or spot, of an irregular figure, and pale white colour, furrounded with florid red, only appears. Thefe whitish fpots or floughs cover fo many ulcers.

An efflorescence, or eruption upon the neck, arms, breaft, and fingers, about the fecond or third day, is a common fymptom of this difeafe. When it appears, the purging and vomiting generally ceafe.

There is often a flight degree of delirium, and the face frequently appears bloated, and the infide of the nostrils red and inflamed. The patient complains of a difagreeable putrid fmell, and his breath is very offenfive.

The putrid, ulcerous fore throat may be diftinguished from the inflammatory, by the vomiting and loofenefs with which it is generally ufhered in; the foul ulcers in the throat covered with a white or livid coat; and by the exceffive weakness of the patient ; with other fymptoms of a putrid fever.

Unfavourable fymptoms are, an obstinate purging, extreme weaknefs, dimnefs of the fight, a livid or black colour of the fpots, and frequent fhiverings, with a weak, fluttering pulfe. If the eruption upon the fkin fuddenly difappears, or becomes of a livid colour, with a difcharge of blood from the nofe or mouth, the danger is very great.

If a gentle sweat break out about the third, or fourth day, and continue with a flow, firm, and equal pulfe; if the floughs caft off in a kindly manner, and appear clean and florid at the bottom; and if the breathing is foft and

PUTRID ULCEROUS SORE THROAT. 259

and free, with a lively colour of the eyes, there is reafon to hope for a falutary crifis.

REGIMEN. — The patient must be kept quiet, and for the most part in bed, as he will be apt to faint when taken out of it. His food must be nourishing and restorative; as fago-gruel with red wine, jellies, strong broths, &c. His drink ought to be generous, and of an antifeptic quality; as red-wine negus, white-wine whey, and such like.

MEDICINE. — — — The medicine in this kind of quinfey is entirely different from that which is proper in the inflammatory. All evacuations, as bleeding, purging, &c. which weaken the patient, muft be avoided. Cooling medicines, as nitre and cream of tartar, are likewife hurtful. Strengthening cordials alone can be ufed with fafety; and thefe ought never to be neglected.

If at the beginning there is a great nausea, or inclination to vomit, the patient must drink an infusion of green tea, camomile flowers, or *carduus benedictus*, in order to cleanse the stomach. If these are not sufficient, he may take a few grains of the powder of ipecacuanha, or any other gentle vomit.

If the difeafe be mild, the throat may be gargled with an infufion of fage and rofe leaves, to a gill of which may be added a fpoonful or two of honey, and as much vinegar as will make it agreeably acid; but when the fymptoms are urgent, the floughs large and thick, and the breath very offenfive, the following gargle may be ufed:

To fix or feven ounces of the pectoral decoction, when boiling, add half an ounce of contrayerva-root; let it boil for fome time, and afterwards ftrain the liquor; to which add two ounces of white-wine vinegar, an ounce of fine honey, and an ounce of the tincture of myrrh. This ought not only to be ufed as a gargle, but a little of it fhould frequently be injected with a fyringe to clean the throat, before the patient takes any meat or drink. This method is peculiarly neceffary for children, who cannot ufe a gargle.

260 OF THE MALIGNANT QUINSEY, or

It will be of great benefit if the patient frequently receives into his mouth, through an inverted funnel, the fleams of warm vinegar, myrrh, and honey.

But when the putrid fymptoms run high, and the difeafe is attended with danger, the only medicine that can be depended upon is the Peruvian bark. It may be taken in substance, if the patient's ftomach will bear it. If not, an onnce of bark großsly powdered, with two drachms of Virginian fnake-root, may be boiled in an English pint and a half of water to half a pint; to which a tea-fpoonful of the elixir of vitriol may be added, and an ordinary tea-cupful of it taken every three or four hours. Bliftering-plafters are very beneficial in this difeafe, especially when the patient's pulse and spirits are low. They may be applied to the throat, behind the ears, or upon the back part of the neck.

Should the vomiting prove troublefome, it will be proper to give the patient two table-spoonfuls of the faline julep every hour. Tea made of mint and a little cinnamon will be very proper for his ordinary drink, especially if an equal quantity of red-wine be mixed with it.

In cafe of a violent loofenefs, the fize of a nutmeg of diascordium, or the japonic confection, may be taken two or three times a-day, or oftener, if neceffary.

If a difcharge of blood from the nofe happens, the fteams of warm vinegar may be received up the noftrils frequently; and the drink must be sharpened with spirits of vitriol, or tincture of rofes.

In cafe of a strangury, the belly must be fomented with warm water, and emollient clyfters given three or four times a day. at la manual novel to zil ol

After the violence of the difeafe is over, the body should still be kept open with mild purgatives; as manna, fenna, rhubarb, or the like. To soonso own bbc doids

If great weakness and dejection of spirits, or nightfweats, with other fymptoms of a confumption, fhould enfue; we would advife the patient to continue the ufe of the Peruvian bark, with the elixir of vitriol, and to take frequently a glass of generous wine. Thefe, together with a nourifhing diet, and riding on horfe-back, are the most likely means for recovering his strength. The

PUTRID ULCEROUS SORE THROAT. 261

The quinfey, being a local difeafe, is generally caught by exposing the throat to a draught of cold air. I know many people, who are fure to be troubled with this complaint if they ftand or fit near an open window, or continue for any length of time in a room lately washed. There is not a readier or a more certain way to catch a' quinfey, than fitting near an open window in a carriage, especially during the night, or when the weather is cold or damp.

The inflammatory fore throat, though it fometimes comes to a fuppuration, generally yields to the method of treatment recommended in this chapter. Cafes, however, occur where the power of fwallowing is loft, and the patient perifhes from the mere want of fultenance. I lately faw a very ingenious invention of a young furgeon, by which a man's life was faved in a cafe of this kind. He fastened a tunnel to the skin of an eel, open at both ends; and, by means of a flexible probe, pulhed one end down the gullet, till it entered the ftomach. Afterwards, milk, broth, or whatever was deemed proper for nourifhing the patient, was put into the tunnel, and conveyed to the ftomach. Though I mention this chiefly with a view of directing others in the like alarming fituations, vet it may also ferve to confirm an opinion, often avowed by the late JOHN HUNTER, and well illustrated in his own practice, that prefence of mind, and a readinefs or fertility of mechanical contrivances, may fometimes prove more ferviceable in a critical moment, than all the refources of science.

But the most dangerous kind of quinfey, as I before observed, is that attended with a putrid fever, commonly called the malignant quinfey, or putrid ulcerous fore throat. Wherever the fymptoms of this appear, I cannot too urgently advise the patient's friends to lose no time in procuring for him the best medical affistance they can obtain. The delay of an hour may be attended with irreparable injury.

S 3

CHAP.

[262]

CHAP. XXIX.

OF COLDS AND COUGHS.

IT has already been obferved, that colds are the effect of an obftructed perfpiration; the common caufes of which we have likewife endeavoured to point out, and fhall not here repeat them. Neither fhall we fpend time in enumerating all the various fymptoms of colds, as they are pretty generally known. It may not, however, be amifs to obferve, that almost every cold is a kind of fever, which only differs in degree from fome of those that have already been treated of.

No age, fex, or conflictution, is exempted from this difeafe; neither is it in the power of any medicine or regimen to prevent it. The inhabitants of every climate are liable to catch cold, nor can even the greateft circumfpection defend them at all times from its attacks. Indeed, if the human body could be kept conftantly in an uniform degree of warmth, fuch a thing as catching cold would be impoffible; but as that cannot be effected by any means, the perfpiration muft be liable to many changes. Such changes, however, when fmall, do not affect the health; but, when great, they muft prove hurtful.

When opprefion of the breaft, a fluffing of the nofe, unufual wearinefs, pain of the head, &c. give ground to believe that the perfpiration is obftructed, or, in other words, that the perfon has caught cold, he ought immediately to leffen his diet, at leaft the ufual quantity of his folid food, and to abftain from all ftrong liquors. Inftead of flefh, fifh, eggs, milk, and other nourifhing diet, he may eat light bread-pudding, veal or chicken broth, panado, gruels, and fuch like. His drink may be watergruel fweetened with a little honey; an infufion of balm, or linfeed fharpened with the juice of bitter orange or lemon; a decoction of barley and liquorice with tamarinds, or any other cool, diluting, acid liquor.

Above

Above all, his fupper fhould be light; as fmall poffet, or water-gruel fweetened with honey, and a little toafted bread in it. If honey fhould difagree with the ftomach, the gruel may be fweetened with treacle or coarfe fugar, and fharpened with the jelly of currants. Thofe who have been accuftomed to generous liquors may take wine-whey inftead of gruel, which may be fweetened as above.

. The patient ought to lie longer than usual a-bed, and to encourage a gentle fweat, which is eafily brought on towards morning by drinking tea, or any kind of warm diluting liquor. I have often known this practice carry off a cold in one day, which, in all probability, had it been neglected, would have coft the patient his life, or have confined him for fome months. Would people facrifice a little time to eafe and warmth, and practife a moderate degree of abstinence when the first fymptoms of a cold appear, we have reafon to believe that most of the bad effects which flow from an obstructed perspiration might be prevented. But, after the difease has gathered strength by delay, all attempts to remove it often prove vain. A pleurify, a peripheumony, or a fatal confumption of the lungs, are the common effects of colds which have either been totally neglected, or treated improperly.

Many attempt to cure a cold, by getting drunk ; but this, to fay no worfe of it, is a very hazardous experiment. No doubt it may fometimes fucceed, by fuddenly reftoring the perfpiration ; but when there is any degree of inflammation, which is frequently the cafe, ftrong liquors, inflead of removing the malady, will increase it. By this means a common cold may be converted into an inflammatory fever.

When those who labour for their daily bread have the misfortune to catch cold, they cannot afford to lose a day or two, in order to keep themselves warm, and take a little medicine; by which means the diforder is often fo aggravated as to confine them for a long time, or even to render them ever after unable to fultain hard labour. But even such of the labouring poor as can afford to take care of themselves, are often too hardy to do it; they affect to despise colds, and as long as they can crawl

54

about,

about, fcorn to be confined by what they call a common cold. Hence it is that colds deftroy fuch numbers of mankind. Like an enemy defpifed, they gather ftrength from delay, till at length they become invincible. We often fee this verified in travellers, who, rather than lofe a day in the profecution of their bufinefs, throw away their lives by purfuing their journey, even in the fevereft weather, with this difeafe upon them.

It is certain, however, that colds may be too much indulged. When a perfon, for every flight cold, fhuts himfelf up in a warm room, and drinks great quantities of warm liquor, it may occafion fuch a general relaxation of the folids as will not be eafily removed. It will, therefore, be proper, when the difeafe will permit, and the weather is mild, to join to the regimen mentioned above, gentle exercife; as walking, riding on horfeback, &c. An obftinate cold, which no medicine can remove, will yield to gentle exercife and a proper regimen of the diet.

Bathing the feet and legs in warm water has a great tendency to reftore the perfpiration. But care muft be taken that the water be not too warm, otherwife it will do hurt. It fhould never be much warmer than the blood, and the patient fhould go immediately to bed after ufing it. Bathing the feet in warm water, lying in bed, and drinking warm water-gruel, or other weak liquors, will fooner take off a fpafm, and reftore the perfpiration, than all the hot fudorific medicines in the world. This is all that is neceffary for removing a common cold; and if this courfe be taken at the beginning, it will feldom fail.

But when the fymptoms do not yield to abfinence, warmth, and diluting liquors, there is reafon to fear the approach of fome other difeafe, as an inflammation of the breaft, an ardent fever, or the like. If the pulfe, therefore, be hard and frequent, the fkin hot and dry, and the patient complains of his head or breaft, it will be neceffary to bleed, and to give the cooling powders recommended in the fcarlet fever, every three or four hours, till they give a ftool.

It will likewife be proper to put a bliftering-plafter on the back, to give two table-fpoonfuls of the faline mixture every two hours, and in fhort to treat the patient in all refpects as for a flight fever. I have often feen this courfe, when obferved at the beginning, remove the complaint in two or three days, when the patient had all the fymptoms of an approaching ardent fever, or an inflammation of the breaft.

The chief fecret of preventing colds lies in avoiding, as far as poffible, all extremes either of heat or cold, and in taking care, when the body is heated, to let it cool gradually. Thefe and other circumftances relating to this important fubject are fo fully treated of under the article *Obstructed Perspiration*, that it is needlefs here to refume the confideration of them.

I believe I need not affure the reader of what he will perceive in almost every page of this book, that I am far from being an enemy to agreeable and innocent recreations. But fome of our amufements are attended with fo much injury to the health of thousands, that it would be carrying indulgence too far to pass them over unnoticed.

Public gardens, those places of general refort in the fummer feason, are as dangerous as they are inviting. Their agreeable coolnefs at the clofe of a hot day, the gaiety of the company, the charms of the mufic, and the variety of the other decorations, concur to render fuch fcenes peculiarly delightful, and to make age as well as youth forget the baneful effects of evening dews, and of the night air. In the mean time, perspiration is checked, and difeafe is inhaled at almost every breath. The dampnefs is in proportion to the heat that preceded it, and is farther increased by exhalations from the water, which always forms one of the ornaments or boundaries of the beautiful landscape. Ah! fly from the bewitching spot at an early hour; and think, that even the verdure you tread upon, fo foft to the foot, and the plants and flowers, fo pleafant to the fight and fmell, begin, foon after funfet, to emit a fort of volatile poifon, and to contaminate the air around you. A little brandy, or brandy and water, while there, is the best prefervative; but nothing

can fave you, if you ftay long. Again, then, let me urge you to withdraw foon; and, after returning home, fome warm and mild liquid, at going to bed, will tend to reftore infenfible perfpiration, and to prevent the attacks of a cold, or of a fever.

But, how am I to addrefs myfelf in a ftrain of admonition to the frequenters of our theatres, which are often the fources of rational pleafure, and where, even at my own advanced period of life, I can hardly refrain from going, when the favourites of the tragic or of the comic mufe exert their fafcinating powers? It would be writing contrary to the impulfes of my own heart, were I to ufe any diffuafives against fuch exquisite gratifications. Yet I hope I may suggest fome useful hints to guard against diffugreeable confequences.

In the relaxing atmosphere of a theatre, heated by the number of lights, and by the breath and effluvia of fo many perfons as are crowded there together, cold, weak drink, however grateful to the tafte, is extremely dangerous. Thirst may be allayed by fucking an orange or lemon, and other bad effects may be obviated by a little brandy or rum, though these fpirits, if used too freely, would rather invite than avert the apprehended evil.

After the entertainment, the greateft danger attends the fudden transition from heat to cold. Additional covering fhould always be prepared to put on, at going out of a theatre; a handkerchief or muff fhould be applied to the nofe and mouth; and the fame precautions ufed, after getting home, as I have directed in the former cafe.

Still greater care is neceffary in affemblies, where all the other caufes of heat are increafed by the exercise of dancing. The dances should end with minuets, or the flowest forms of graceful motion; and time should always be allowed to cool gradually before the breaking up of those gay meetings. On the subject of refreshments or drink, I need not add any thing to what I have already suggested. A melancholy instance has lately occurred to me of the fatal confequences of the want of due caution on such occasions. About three years ago, I was fortunate enough to contribute to the recovery of a lady in

118.3

in the early ftage of a confumption. I then gave her fome very earneft advice to guard againft the ufual caufes of fuch a complaint. But the uninterrupted enjoyment of good health for three years, blotted out of her memory my warnings of danger. In the beginning of laft winter, fhe was tempted to go to the Westminster affembly. She caught cold there; and what is very fignificantly called a *galloping* confumption, carried her in a few weeks to the grave.

As to the votaries of filly fashion, who rush to Italian operas, and the like unnatural puppet-shew performances, they fearcely appear worthy of either notice or advice. It is not likely that any argument would make much impression upon people who absurdly facrifice health and life to the allurements of false taste; —who affect to admire the most extravagant nonsense; —who, in the words of the elegant ARMSTRONG,

---- " With loudeft peals,

- " Applaud the fool that higheft lifts his heels;
- " And with infipid fhew of rapture die
- " Of idiot notes impertinently long."

OF A COMMON COUGH.

A cough is generally the effect of a cold, which has either been improperly treated, or entirely neglected. When it proves obstinate, there is always reason to fear the confequences, as this shews a weak state of the lungs, and is often the forerunner of confumption.

If the cough be violent, and the patient young and ftrong, with a hard quick pulfe, bleeding will be proper; but in weak and relaxed habits, bleeding rather prolongs the difeafe. When the patient fpits freely, bleeding is unneceffary, and fometimes hurtful, as it tends to leffen that difcharge.

When the cough is not attended with any degree of fever, and the fpittle is viscid and tough, sharp pectoral medicines are to be administered; as gum-ammoniac, fquills, &c. Two table-spoonfuls of the solution of gumammoniac may be taken three or four times a-day, more or lefs, according to the age and constitution of the patient.

OF A COMMON COUGH.

268

tient. Squills may be given various ways: two ounces of the vinegar, the oxymel, or the fyrup, may be mixed with the fame quantity of fimple cinnamon-water, to which may be added an ounce of common water and an ounce of balfamic fyrup. Two table-fpoonfuls of this mixture may be taken three or four times a-day.

A fyrup made of equal parts of lemon-juice, honey, and fugar-candy, is likewife very proper in this kind of cough. A table-fpoonful of it may be taken at pleafure.

But when the defluxion is fharp and thin, these medicines rather do hurt. In this case, gentle opiates, oils, and mucilages, are more proper. A cup of an infusion of wild poppy-leaves, and marsh-mallow roots, or the flowers of colts-foot, may be taken frequently; or a tea-spoonful of the paregoric elixir may be put into the patient's drink twice a-day. Fuller's Spanish infusion is also a very proper medicine in this case, and may be taken in the quantity of a tea-cupful three or four times a-day *.

When a cough is occafioned by acrid humours tickling the throat and *fauces*, the patient thould keep fome foft pectoral lozenges almost constantly in his mouth; as the Pontefract liquorice-cakes, barley-fugar, the common balfamic lozenges, Spanish juice, &c. These blunt the acrimony of the humours, and, by taking off their ftimulating quality, help to appeale the cough \dagger .

In obftinate coughs, proceeding from a flux of humours upon the lungs, it will often be neceffary, befides expectorating medicines, to have recourfe to iffues, fetons, or fome other drain. In this cafe I have often obferved the most happy effects from a Burgundy-

* See Appendix, Spanish Infusion.

† In a former edition of this book I recommended, for an oblinate tickling cough, an oily emulfion, made with the paregoric elixir of the Edinburgh Difpenfatory, inflead of the common alkaline fpirit. I have fince been told by feveral practitioners, that they found it to be an excellent medicine in this diforder, and every way deferving of the character which I had given it. Where this elixir is not kept, its place may be fupplied by adding to the common oily emulfion, an adequate proportion of the *Thebaic Tinclure*, or liquid laudanum.

OF A COMMON COUGH.

pitch plaster applied between the shoulders. I have ordered this simple remedy in the most obstinate coughs, in a great number of cases, and in many different constitutions, without ever knowing it fail to give relief, unless where there were evident signs of an ulcer in the lungs.

About the bulk of a nutmeg of Burgundy-pitch may be foread thin upon a piece of foft leather, about the fize of the hand, and laid between the fhoulderblades. It may be taken off and wiped every three or four days, and ought to be renewed once a fortnight or three weeks. This is indeed a cheap and fimple medicine, and confequently apt to be defpifed; but we will venture to affirm, that the whole *materia medica* does not afford an application more efficacious in almost every kind of cough. It has not indeed always an immediate effect; but, if kept on for fome time, it will fucceed where most other medicines fail.

The only inconveniency attending this plafter is the itching which it occasions; but furely this may be difpenfed with, confidering the advantage which the patient may expect to reap from the application; befides, when the itching becomes very uneafy, the plafter may be taken off, and the part rubbed with a dry cloth, or washed with a little warm milk and water. Some caution indeed is necessary in discontinuing the use of such a plaster; this, however, may be fafely done by making it so degrees, and at length quitting it altogether in a warm feason *.

But coughs proceed from many other caufes befides defluxions upon the lungs. In these cases the cure is not to be attempted by pectoral medicines. Thus, in a cough proceeding from a foulness and debility of the stomach, fyrups, oils, mucilages, and all kinds of balfamic medicines do hurt. The *flomach cough* may be known from one that is owing to a fault in the lungs by

* Some complain that the pitch plafter adheres too faft, while others find difficulty in keeping it on. This proceeds from the different kinds of pitch made use of, and likewise from the manner of making it. I generally find it answer best when mixed with a little bees-wax, and spread as cool as possible. The clear, hard, transparent pitch answers the purpose best.

OF A COMMON COUGH.

270

this, that in the latter the patient coughs whenever he infpires, or draws in his breath fully; but in the former that does not happen.

The cure of this cough depends chiefly upon cleanfing and ftrengthening the ftomach; for which purpofe gentle vomits and bitter purgatives are most proper. Thus, after a vomit or two, the facred tincture, as it is called, may be taken for a confiderable time in the dole of one or two table-fpoonfuls twice a-day, or as often as it is found neceffary, to keep the body gently open. People may make this tincture themfelves, by infufing an ounce of *hiera picra** in an English pint of white-wine, letting it ftand a few days, and then straining it off for use.

In coughs which proceed from a debility of the ftomach, the Peruvian bark is likewife of confiderable fervice. It may either be chewed, taken in powder, or made into a tincture along with other ftomachic bitters.

A nervous cough can only be removed by change of air, and proper exercife; to which may be added the ufe of gentle opiates. Inftead of the faponaceous pill, the paregoric elixir, &c. which are only opium difguifed, ten, fifteen, twenty, or twenty-five drops of liquid laudanum, more or lefs, as circumftances require, may be taken at bed-time, or when the cough is most troublefome. Immerfing the feet and hands in warm water will often appeafe the violence of a nervous cough.

When a cough is only the fymptom of fome other malady, it is in vain to attempt to remove it without first curing the difeafe from which it proceeds. Thus, when a cough is occafioned by *teething*, keeping the body open, fcarifying the gums, or whatever facilitates the cutting of the teeth, likewife appeafes the cough. In like manner, when *worms* occafion a cough, fuch medicines as remove thefe vermin will generally cure the cough; as bitter purgatives, oily clyfters, and fuch like.

Women, during the last months of pregnancy, are often greatly afflicted with a cough, which is generally

· See Appendix, Hiera Picra.

relieved

relieved by bleeding, and keeping the body gently open. They ought to avoid all flatulent food, and to wear a loofe eafy drefs.

A cough is not only a fymptom, but is often likewife the forerunner of difeafes. Thus, the gout is frequently ufhered in by a very troublefome cough, which affects the patient for fome days before the coming on of the fit. This cough is generally removed by a paroxyfm of the gout, which fhould therefore be promoted, by keeping the extremities warm, drinking warm liquors, and bathing the feet and legs frequently in lukewarm water.

OF THE HOOPING COUGH, OR CHIN-COUGH.

This cough feldom affects adults, but proves often fatal to children. Such children as live upon thin watery diet, who breathe unwholefome air, and have too little exercife, are most liable to this difease, and generally fuffer most from it.

The chin-cough is fo well known, even to nurfes, that a defcription of it is unneceffary. Whatever hurts the digestion, obstructs the perspiration, or relaxes the folids, disposes to this disease; confequently its cure must depend upon cleansing and strengthening the stomach, bracing the folids, and at the same time promoting perspiration and the different fecretions.

The diet must be light, and of eafy digestion; for children, good bread made into pap or pudding, chickenbroth, with other light spoon meats, are proper; but those who are farther advanced may be allowed sagogruel, and if the fever be not high, a little boiled chicken, or other white meats. The drink may be hyssop, or penny-royal tea, sweetened with honey or sugar-candy, small wine-whey; or, if the patient be weak, he may fometimes be allowed a little negus.

One of the most effectual remedies in the chincough is change of air. This often removes the malady, even when the change feems to be from a purer to a lefs wholefome air. This may in fome measure depend on the patient's being removed from the place where

272 OF THE HOOPING-COUGH,

where the infection prevails. Moft of the difeafes of children are infectious; nor is it at all uncommon to find the chin-cough prevailing in one town or village, when another at a very fmall diftance is quite free from it. But whatever be the caufe, we are fure of the fact. No time ought therefore to be loft in removing the patient at fome diftance from the place where he caught the difeafe, and, if poffible, into a more pure and warm air *.

When the difeafe proves violent, and the patient is in danger of being fuffocated by the cough, he ought to be bled, efpecially if there be a fever with a hard full pulfe. But as the chief intention of bleeding is to prevent an inflammation of the lungs, and to render it more fafe to give vomits, it will feldom be neceffary to repeat the operation; yet if there are fymptoms of an inflammation of the lungs, a fecond or even a third bleeding may be requifite.

It is generally reckoned a favourable fymptom when a fit of coughing makes the patient vomit. This cleanfes the ftomach, and greatly relieves the cough \ddagger . It will therefore be proper to promote this difcharge, either by fmall dofes of ipecacuanha, or the vomiting julep recommended in the Appendix \ddagger .

It is very difficult to make children drink after a vomit. I have often feen them happily deceived, by infufing a fcruple or half a drachm of the powder of ipecacuanha in a tea-pot, with half an English pint of boiling water. If this be difguised with a few drops of milk and a little fugar, they will imagine it tea, and drink it very greedily. A small tea-cupful of this may be given every quarter of an hour, or rather every ten minutes, till it operates. When the child begins to

* Some think the air ought not to be changed till the difeafe is on the decline; but there feems to be no fufficient reafon for this opinion, as patients have been known to reap benefit from a change of air at all periods of the difeafe. It is not fufficient to take the patient out daily in a carriage. This feldom anfwers any good purpofe; but often does hurt, by giving him cold.

+ As the hooping-cough is evidently a fpalmodic diforder, I have often thought that tonics might prove the most proper medicines. This, time must determine.

1 See Appendix, Vomiting Julep.

on the patient's

puke, there will be no occasion for drinking any more, as the water already on the ftomach will be fufficient.

Vomits not only cleanfe the ftomach, which in this difeafe is generally loaded with vifcid phlegm, but they likewife promote the perfpiration and other fecretions, and ought therefore to be repeated according to the obftinacy of the difeafe. They fhould not however be ftrong; gentle vomits frequently repeated are both lefs dangerous, and more beneficial than ftrong ones.

The body ought to be kept gently open. The beft medicines for this purpofe are rhubarb and its preparations, as the fyrup, tincture, &c. Of thefe a tea-fpoonful or two may be given to an infant twice or thrice a-day, as there is occafion. To fuch as are farther advanced, the dofe muft be proportionally increafed, and repeated till it has the defired effect. Thofe who cannot be brought to take the bitter tincture, may have an infufion of fenna and prunes, fweetened with manna, coarfe fugar, or honey; or a few grains of rhubarb mixed with a tea-fpoonful or two of fyrup, or currantjelly, fo as to difguife the tafte. Moft children are fond of fyrups and jellies, and feldom refufe even a difagreeable medicine when mixed with them.

Many people believe that oily, pectoral, and balfamic medicines poffefs wonderful virtues for the cure of the chin-cough, and accordingly exhibit them plentifully to patients of every age and conflitution, without confidering that every thing of this nature must load the ftomach, hurt the digestion, and of courfe aggravate the diforder *.

The millepedes, or wood-lice, are greatly recommended for the cure of a chin-cough. Those who choose to make use of these infects may infuse two ounces of them bruised in a English pint of small white-wine for

[•] Dr. DUPLANIL fays, he has feen many good effects from the kermes mineral in this complaint, the cough being frequently alleviated even by the first dose. The dose for a child of one year old is a quarter of a grain diffolved in a cup of any liquid, repeated two or three times a-day. For a child of two years, the dose is half a grain; and the quantity must be thus increased in proportion to the age of the patient.

OF THE HOOPING-COUCH,

274

one night. Afterwards the liquor may be ftrained through a cloth, and a table-fpoonful of it given to the patient three or four times a-day.

Opiates are fometimes neceffary to allay the violence of the cough. For this purpole, a little of the fyrup of poppies, or five, fix, or feven drops of laudanum, according to the age of the patient, may be taken in a cup of hyffop or penny-royal tea, and repeated occafionally *.

The garlic ointment is a well-known remedy in North Britain for the chin-cough. It is made by beating in a mortar garlic with an equal quantity of hog's lard. With this the foles of the feet may be rubbed twice or thrice a-day; but the beft method is to fpread it upon a rag, and apply it in the form of a plafter. It fhould be renewed every night and morning at leaft, as the garlick foon lofes its virtue. This is an exceeding good medicine both in the chin-cough †, and in most other coughs of an obstinate nature. It ought not, however, to be used when the patient is very hot or feverish, left it should increase these fymptoms.

The feet fhould be bathed once every two or three days in lukewarm water; and a Burgundy-pitch plafter kept conftantly between the fhoulders. But when the difeafe proves very violent, it will be neceffary, inftead of it, to apply a bliftering-plafter, and to keep the part open for fome time with iffue-ointment.

When the difeafe is prolonged, and the patient is free from a fever, the Peruvian bark, and other bitters, are the most proper medicines. The bark may either be taken in fubstance, or in a decoction or infusion, as is most agreeable. For a child, ten, fifteen, or twenty grains, according to the age of the patient, may be given three or four times a-day. For an adult half a drachm or two fcruples will be proper. Some give the

* Some recommend the extract of hemlock as an extraordinary remedy in the hooping-cough; but, fo far as I have been able to obferve, it is no way fuperior to opium, which, when properly administered, will often relieve fome of the most troublefome fymptoms of this diforder.

+ As this difeafe is evidently fpafmodic, I am inclined to think that tonic medicines will in time be found the most proper for its cure.

extract

OR CHIN-COUGH.

extract of the bark with cantharides; but to manage this requires a confiderable attention. It is more fafe to give a few grains of caftor along with the bark. A child offix or feven years of age may take feven or eight grains of caftor, with fifteen grains of powdered bark, for a dofe. This may be made into a mixture, with two or three ounces of any fimple diftilled water, and a little fyrup, and taken three or four times a-day.

CHAP. XXX.

INFLAMMATION OF THE STOMACH, AND OTHER VISCERA.

A LL inflammations of the bowels are dangerous, and require the most speedy assistance; as they frequently end in a suppuration, and sometimes in a mortification, which is certain death.

CAUSES.——An inflammation of the ftomach may proceed from any of the caufes which produce an inflammatory fever; as cold liquor drank while the body is warm, obftructed perfpiration, or the fudden ftriking in of any eruption. It may likewife proceed from the acrimony of the bile, or from acrid and ftimulating fubftances taken into the ftomach; as ftrong vomits or purges, corrofive poifons, and fuch like. When the gout has been repelled from the extremities, either by cold or improper applications, it often occafions an inflammation of the ftomach. Hard or indigeftible fubftances taken into the ftomach, as bones, the ftones of fruits, &c. may likewife have that effect.

SYMPTOMS.——It is attended with a fixed pain and burning heat in the ftomach; great reftleffnefs and anxiety; a fmall, quick, and hard pulfe; vomiting, or at leaft a naufea and ficknefs; exceffive thirft; coldnefs of the extremities; difficulty of breathing; cold clammy fweats; and fometimes convulfions and fainting fits. The ftomach is fwelled, and often feels hard to the touch. One of the most certain figns of this difeafe is the fenfe of pain, which the patient feels upon taking

any

276 INFLAMMATION OF THE STOMACH,

any kind of food or drink, especially if it be either too hot or too cold.

When the patient vomits every thing he eats or drinks, is extremely reftlefs, has a hiccup, with an intermitting pulfe, and frequent fainting fits, the danger is very great.

REGIMEN.——All acrimonious, heating, and irritating food and drink are carefully to be avoided. The weaknefs of the patient may deceive the by-ftanders, and induce them to give him wines, fpirits, or other cordials; but thefe never fail to increafe the difeafe, and often occafion fudden death. The inclination to vomit may likewife impofe on the attendants, and make them think a vomit neceffary; but that too is almost certain death.

The food must be light, thin, cool, and eafy of digeftion. It must be given in small quantities, and should neither be quite cold, nor too hot. Thin gruel made of barley or oatmeal, light toasted bread diffolved in boiling water, or very weak chicken-broth are the most proper. The drink should be clear whey, barley-water, water in which toasted bread has been boiled, or decoctions of emollient vegetables, as liquorice, and marsh-mallow roots, farfaparilla, or the like.

MEDICINE. — Bleeding in this difeafe is abfolutely neceffary, and is almost the only thing that can be depended on. When the difease proves obstinate, it will often be proper to repeat this operation several times; nor must the low state of the pulse deter us from doing fo. The pulse indeed generally rifes upon bleeding, and as long as that is the case, the operation is fase.

Frequent fomentations with lukewarm water, or a decoction of emollient vegetables, are likewife beneficial. Flannel cloths dipped in thefe must be applied to the region of the stomach, and removed as they grow cool. They must neither be applied too warm, nor be suffered to continue till they become quite cold, as either of these extremes would aggravate the difease.

The feet and legs ought likewife to be frequently bathed in lukewarm water, and warm bricks or poultices may be applied to the foles of the feet. The warm bath, if it can be conveniently ufed, will be of great fervice. In this, and all other inflammations of the bowels, an epifpastic, or bliftering-plaster, applied over the part affected, is one of the best remedies I know. I have often used it, and do not recollect one instance wherein it did not give relief to the patient.

The only internal medicines which we fhall venture to recommend in this difeafe, are mild clyfters. Thefe may be made of warm water, or thin water-gruel; and if the patient be coftive, a little fweet oil, honey, or manna, may be added. Clyfters anfwer the purpofe of an internal fomentation, while they keep the body open, and at the fame time nourifh the patient, who is often in this difeafe unable to retain any food upon his ftomach. For thefe reafons they muft not be neglected, as the patient's life may depend on them.

INFLAMMATION OF THE INTESTINES.

THIS is one of the most painful and dangerous difeafes to which mankind are liable. It generally proceeds from the fame caufes as the inflammation of the ftomach; to which may be added coftiveness, worms, eating unripe fruits, or great quantities of nuts, drinking hard windy malt liquors, as stale bottled beer or ale, four wine, cyder, &c. It may likewife be occasioned by a rupture, by fchirrous tumours of the intestines, or by their opposite fides growing together.

The inflammation of the inteffines is denominated Iliac paffion, Enteritis, Sc. according to the name of the parts affected. 'The treatment, however, is nearly the fame whatever part of the inteffinal canal be the feat of the difeafe; we shall therefore omit these distinctions, left they should perplex the reader.

The fymptoms here are nearly the fame as in the foregoing difeafe; only the pain, if poffible, is more acute, and is fituated lower. The vomiting is likewife more violent, and fometimes even the excrements, together with the clyfters, are difcharged by the mouth. The patient is continually belching up wind, and has often an obstruction of his urine.

While

278 INFLAMMATION OF THE INTESTINES.

While the pain fhifts, and the vomiting only returns at certain intervals, and while the clyfters pafs downwards, there is ground for hope; but when the clyfters and *faces* are vomited, and the patient is exceeding weak, with a low fluttering pulfe, a pale countenance, and a difagreeable or flinking breath, there is great reafon to fear that the confequences will prove fatal. Clammy fweats, black fœtid flools, with a fmall intermitting pulfe, and a total ceffation of pain, are the figns of a mortification already begun, and of approaching death.

REGIMEN.——The regimen in this difeafe is in general the fame as in an inflammation of the flomach. The patient must be kept quiet, avoiding cold, and all violent passions of the mind. His food ought to be very light, and given in small quantities; his drink weak and diluting; as clear whey, barley-water, and such like.

MEDICINE. ——Bleeding in this, as well as in the inflammation of the ftomach, is of the greateft importance. It fhould be performed as foon as the fymptoms appear, and must be repeated according to the ftrength of the patient and the violence of the difeafe.

A bliftering-plafter is here likewife to be applied immediately over the part where the most violent pain is.

This not only relieves the pain of the bowels, but even clyfters and purgative medicines, which before had no effect, will operate when the blifter begins to rife.

Fomentations and laxative clyfters are by no means to be omitted. The patient's feet and legs fhould frequently be bathed in warm water; and cloths dipped in it applied to his belly. Bladders filled with warm water may likewife be applied to the region of the navel, and warm bricks, or bottles filled with warm water, to the foles of the feet. The clyfters may be made of barley-water, or thin gruel with falt, and foftened with fweet oil or fresh butter. These may be administered every two or three hours, or oftener, if the patient continues coflive.

If the difeafe does not yield to clyfters and fomentations, recourfe must be had to pretty strong purgatives; but as thefe, by irritating the bowels, often increase their contraction,

INFLAMMATION OF THE INTESTINES. 279

contraction, and by that means frustrate their own intention, it will be neceffary to join them with opiates, which by allaying the pain, and relaxing the spafmodic contractions of the guts, greatly affist the operation of purgatives in this cafe.

What answers the purpole of opening the body very well, is a folution of the bitter purging falts. Two ounces of these may be diffolved in an English pint of warm water, or thin gruel, and a tea-cupful of it taken every half hour till it operates. At the fame time fifteen, twenty, or twenty-five drops of laudanum may be given in a glass of peppermint or fimple cinnamon-water, to appease the irritation, and prevent the vomiting, &c.

Acids have often a very happy effect in flaying the vomiting, and appealing the other violent fymptoms of this difeafe. It will therefore be of use to fharpen the patient's drink with cream of tartar, juice of lemon; or, when these cannot be obtained, with vinegar.

But it often happens that no liquid whatever will ftay on the ftomach. In this cafe the patient must take purging pills. I have generally found the following answer very well: Take jalap in powder, and vitriolated tartar, of each half a drachm, opium one grain, Castile soap as much as will make the mass fit for pills. These must be taken at one dose, and if they do not operate in a few hours, the dose may be repeated.

If a ftool cannot be procured by any of the above means, it will be neceffary to immerfe the patient in warm water up to the breaft. I have often feen this fucceed when other means had been tried in vain. The patient must continue in the water as long as he can eafily bear it without fainting, and if one immerfion has not the defired effect, it may be repeated as foon as the patient's ftrength and spirits are recruited. It is more fafe for him to go frequently into the bath, than to continue too long at a time; and it is often neceffary to repeat it feveral times before it has the defired effect.

It has fometimes happened, after all other means of procuring a stool had been tried to no purpose, that this

T 4

was

289 INFLAMMATION OF THE INTESTINES.

was brought about by immerfing the patient's lower extremities in cold water, or making him walk upon a wet pavement, and dafhing his legs and thighs with the cold water. This method, when others fail, at leaft merits a trial. It is indeed attended with fome danger; but a doubtful remedy is better than none.

In defperate cafes it is common to give quick-filver. This may be given to the quantity of feveral ounces, or even a pound, but fhould not exceed that *. When there is reafon to fulpect a mortification of the guts, this medicine ought not to be tried. In that cafe it cannot cure the patient, and will only haften his death. But when the obftruction is occafioned by any caufe that can be removed by force, quickfilver is not only a proper medicine, but the beft that can be adminiftered, as it is the fitteft body we know for making its way through the inteffinal canal.

If the difeafe proceeds from a rupture, the patient must be laid with his head very low, and the intestines returned by gentle pressure with the hand. If this, with fomentations and clysters, should not fucceed, recourse must be had to a furgical operation, which may give the patient relief.

Such as would avoid this excruciating and dangerous difeafe, muft take care never to be too long without a ftool. Some who have died of it, have had feveral pounds of hard dry *faces* taken out of their guts. They fhould likewife beware of eating too freely of four or unripe fruits, or drinking ftale windy liquors, &c. I have known it brought on by living too much on baked fruits, which are feldom good. It likewife proceeds frequently from cold caught by wet clothes, &c. but efpecially from wet feet.

* When quickfilver is given in too large quantities it defeats its own intention, as it drags down the bottom of the ftomach, which prevents its getting over the Pylorus. In this cafe the patient fhould be hung up by the heels, in order that the quickfilver may be difcharged by his mouth.

OF THE COLIC.

THE colic has a great refemblance to the two precedering difeafes, both in its fymptoms and method of cure. It is generally attended with coffiveness and acute pain of the bowels; and requires diluting diet, evacuations, fomentations, &c.

Colics are varioufly denominated according to their caufes, as the *flatulent*, the *bilious*, the *hyfteric*, the *nervous*, &c. As each of thefe requires a particular method of treatment, we fhall point out their most general fymptoms, and the means to be used for their relief.

The *flatulent*, or wind-colic, is generally occafioned by an indifcreet use of unripe fruits, meats of hard digestion, windy vegetables, fermenting liquors, and such like. It may likewise proceed from an obstructed perspiration, or catching cold. Delicate people, whose digestive powers are weak, are most liable to this kind of cholic.

The flatulent colic may either affect the flomach or inteffines. It is attended with a painful flretching of the affected part. The patient feels a rumbling in his guts, and is generally relieved by a difcharge of wind, either upwards or downwards. The pain is feldom confined to any particular part, as the vapour wanders from one division of the bowels to another, till it finds a yent.

When the difeafe proceeds from windy liquor, green fruits, four herbs, or the like, the best medicine on the first appearance of the fymptoms is a dram of brandy, gin, or any good spirits. The patient should likewife fit with his feet upon a warm hearth-stone, or apply warm bricks to them; and warm cloths may be applied to his stomach and bowels.

This is the only colic wherein ardent fpirits, fpiceries, or any thing of a hot nature, may be ventured upon. Nor indeed are they to be used here unless at the very beginning, before any symptoms of inflammation appear. We have reason to believe, that the colic occasioned

OF THE COLIC.

cafioned by wind or flatulent food might always be cured by fpirits and warm liquors, if they were taken immediately upon perceiving the first uneafines; but when the pain has continued for a confiderable time, and there is reason to fear an inflammation of the bowels is already begun, all hot things are to be avoided as poison, and the patient is to be treated in the fame manner as for the inflammation of the intestines.

Several kinds of food, as honey, eggs, &c. occasion colics in fome particular conflictutions. I have generally found the best method of cure for these was to drink plentifully of small diluting liquors, as water-gruel, small posset, toast and water, &c.

Colics which proceed from excels and indigeftion generally cure themfelves by occasioning vomiting or purging. These discharges are by no means to be stopped, but promoted by drinking plentifully of warm water, or weak posset. When their violence is over, the patient may take a dose of rhubarb, or any other gentle purge, to carry off the dregs of his debauch.

Colics which are occafioned by wet feet, or catching cold, may generally be removed at the beginning by bathing the feet and legs in warm water, and drinking fuch warm diluting liquors as will promote the perfpiration, as weak wine-whey, or water-gruel with a fmall quantity of fpirits in it.

Those flatulent colics, which prevail fo much among country people, might generally be prevented were they careful to change their clothes when they get wet. They ought likewife to take a dram, or to drink fome warm liquor after eating any kind of green trafh. We do not mean to recommend the practice of dram-drinking, but in this cafe ardent spirits prove a real medicine, and indeed the best that can be administered. A glass of good peppermint-water will have nearly the fame effect as a glass of brandy, and in fome cafes is rather to be preferred.

The bilious colic is attended with very acute pains about the region of the navel. The patient complains of great thirst, and is generally costive. He vomits a hot, bitter, yellow-coloured bile, which being discharged, feems feems to afford fome relief, but is quickly followed by the fame violent pain as before. As the diffemper advances, the propenfity to vomit fometimes increafes fo as to become almost continual, and the proper motion of the intestines is fo far perverted, that there are all the fymptoms of an impending iliac passion.

If the patient be young and ftrong, and the pulfe full and frequent, it will be proper to bleed, after which clyfters may be administered. Clear whey or gruel, fharpened with the juice of lemon, or cream of tattar, must be drank freely. Small chicken-broth, with a little manna diffolved in it, or a flight decoction of tamarinds, is likewife very proper, or any other thin, acid, opening liquor.

Befides bleeding and plentiful dilution, it will be neceffary to foment the belly with cloths dipped in warm water, and if this fhould not fucceed, the patient mult be immerfed up to the breaft in warm water.

In the bilious colic the vomiting is often very difficult to reftrain. When this happens, the patient may drink a decoction of toafted bread, or an infufion of gardenmint in boiling water. Should thele not have the defired effect, the faline draught, with a few drops of laudanum in it, may be given, and repeated according to the urgency of the fymptoms. A fmall quantity of Venice treacle may be fpread in form of a cataplalm, and applied to the pit of the ftomach. Clyfters, with a proper quantity of Venice treacle or liquid laudanum in them, may likewife be frequently adminiftered.

The *byfteric* colic bears a great refemblance to the bilious. It is attended with acute pains about the region of the ftomach, vomiting, &c. What the patient vomits in this cafe is commonly of a greenifh colour. There is a great finking of the fpirits, with dejection of mind and difficulty of breathing, which are the characteristic fymptoms of this diforder. Sometimes it is accompanied with the jaundice, but this generally goes off of its own accord in a few days.

In this colic all evacuations, as bleeding, purging, vomiting, &c. do hurt. Every thing that weakens the patient, or finks the fpirits, is to be avoided. If, how-

ever,

OF THE COLIC.

ever, the vomiting should prove violent, lukewarm water, or small posset, may be drank to cleanse the stomach. Afterwards the patient may take sisten, twenty, or twenty-five drops of liquid laudanum in a glass of cinnamon-water. This may be repeated every ten or twelve hours, till the symptoms abate.

The patient may likewife take four or five of the fætid pills every fix hours, and drink a cup of penny. royal tea after them. If afafætida fhould prove difagreeable, which is fometimes the cafe, a tea-fpoonful of the tincture of caftor in a cup of pennyroyal tea, or thirty or forty drops of the balfam of Peru dropped upon a bit of loaf-fugar, may be taken in its flead. The anti-hyfteric plafter may alfo be uled, which has often a good effect*.

The nervous colic prevails among miners, fmelters of lead, plumbers, the manufacturers of white lead, &c. It is very common in the cyder countries of England, and is fuppofed to be occafioned by the leaden veffels ufed in preparing that liquor. It is likewife a frequent difeafe in the Weft Indies, where it is termed the dry belly-ache.

No difeafe of the bowels is attended with more excruciating pain than this. Nor is it foon at an end, I have known it continue eight or ten days with very little intermiffion, the body all the while continuing bound in fpite of medicine, yet at length yield, and the patient recover \dagger . It generally, however, leaves the patient weak, and often ends in a palfy.

The general treatment of this difeafe is fo nearly the fame with that of the iliae paffion, or inflammation of the guts, that we fhall not infift upon it. The body is to be opened by mild purgatives given in fmall dofes, and frequently repeated, and their operation must be affifted by foft oily clyfters, fomentations, &c. The caftor oil is reckoned peculiarly proper in this difeafe. It

* See Appendix, Anti-hyfteric Plaster.

† As the imoke of tobaceo thrown into the bowels will often procure a flool when all other means have failed, an apparatus for this purpofe ought to be kept by every furgeon. It may be purchafed at a fmall expence, and will be of fervice in feveral other cafes, as the recovery of drowned perfons, &c. may both be mixed with the clyfters and given by the mouth *.

The Barbadoes tar is faid to be an efficacious medicine in this complaint. It may be taken to the quantity of two drachms three times a day, or oftener if the ftomach will bear it. This tar, mixed with an equal quantity of ftrong rum, is likewife proper for rubbing the fpine, in cafe any tingling, or other fymptoms of the palfy, are felt. When the tar cannot be obtained, the back may be rubbed with ftrong fpirits, or a little oil of nutmegs, or of rofemary.

If the patient remain weak and languid after this difeafe, he must take exercise on horseback, and use an infusion of the Peruvian bark in wine. When the difease ends in a palfy, the Bath waters are found to be extremely proper.

To avoid this kind of colic, people muft fhun all four fruits, acid and auftere liquors, &c. Thofe who work in lead ought never to go to their bufinefs fafting, and their food fhould be oily or fat. They may take a glafs of fallad oil, with a little brandy or rum, every morning, but fhould never take fpirits alone. Liquid aliment is beft for them; as fat broths, &c. but low living is bad. They fhould frequently go a little out of the tainted air; and fhould never fuffer themfelves to be coffive. In the Weft Indies, and on the coaft of Guinea, it has been found of great ufe, for preventing this colic, to wear a piece of flannel round the waift, and to drink an infufion of ginger by way of tea.

Sundry other kinds of this difeafe might be mentioned, but too many diffinctions would tend only to perplex the reader. Thofe already mentioned are the moft material, and fhould indeed be attended to, as their treatment is very different. But even perfons who are not in a condition to diffinguifh very accurately in thefe matters, may neverthelefs be of great fervice to patients in colics of every kind, by only obferving the following general rules, viz. To bathe the feet and legs in warm

* The dofe is from one table-fpoonful to two or three, if neceffary to open the body.

OF THE COLIC.

water; to apply bladders filled with warm water, or cloths wrung out of it, to the ftomach and bowels; to make the patient drink freely of diluting mucilaginous liquors; and to give him an emollient clyfter every two or three hours. Should thefe not fucceed, the patient ought to be immerfed in warm water.

INFLAMMATION OF THE KIDNEYS.

CAUSES. — This difeafe may proceed from any of those causes which produce an inflammatory fever. It may likewise be occasioned by wounds or bruiles of the kidneys; small stones or gravel lodging within them; by strong diuretic medicines; as spirits of turpentine, tincture of cantharides, &c. Violent motion, as hard riding or walking, especially in hot weather, or whatever drives the blood too forcibly into the kidneys, may occasion this malady. It may likewise proceed from lying too fost, too much on the back, involuntary contractions, or spafms, in the urinary vessels, &c.

SYMPTOMS. — There is a fharp pain about the region of the kidneys, with fome degree of fever, and a flupor or dull pain in the thigh of the affected fide. The urine is at first clear, and afterwards of a reddifh colour; but in the worft kind of the difeafe it generally continues pale, is paffed with difficulty, and commonly in fmall quantities at a time. The patient feels great uneafinefs when he endeavours to walk or fit upright. He lies with most ease on the affected fide, and has generally a naufea or vomiting, refembling that which happens in the colic.

This difeafe, however, may be diffinguished from the colic by the pain being feated farther back, and by the difficulty of passing urine, with which it is constantly attended.

INFLAMMATION OF THE KIDNEYS. 287

barley and liquorice, &c. The patient, notwithstanding the vomiting, must constantly keep fipping fmall quantities of these or other diluting liquors. Nothing fo fafely and certainly abates the inflammation, and expels the obstructing cause, as copious dilution. The patient must be kept easy, quiet, and free from cold, as long as any symptoms of inflammation remain.

MEDICINE. — Bleeding is generally neceffary, efpecially at the beginning. Ten or twelve ounces may be let from the arm or foot with a lancet; and if the pain and inflammation continue, the operation may be repeated in twenty-four hours, efpecially if the patient be of a full habit. Leeches may likewife be applied to the hæmorrhoidal veins, as a difcharge from thefe will greatly relieve the patient.

Cloths dipped in warm water, or bladders filled with it, must be applied as near as possible to the part affected, and renewed as they grow cool. If the bladders be filled with a decoction of mallows and camomile flowers, to which a little faffron is added, and mixed with about a third part of new milk, it will be still more beneficial.

Emollient clyfters frequently to be administered; and if these do not open the body, a little salt and honey or, manna may be added to them.

The fame courfe is to be followed where gravel or ftone is lodged in the kidney, but when the gravel or ftone is feparated from the kidney, and lodges in the Ureter *, it will be proper, befides the fomentations, to rub the fmall of the back with fweet oil, and to give gentle diuretics; as juniper-water fweetened with the fyrup of marfh-mallows: a tea-fpoonful of the fweet fpirits of nitre, with a few drops of laudanum, may now and then be put in a cup of the patient's drink. He ought likewife to take exercife on horfeback, or in a carriage, if he be able to bear it.

When the difeafe is protracted beyond the feventh or eighth day, and the patient complains of a flupor and

heavinefs

^{*} The Ureters are two long and fmall canals, one on each fide which carry the urine from the bafon of the kidneys to the bladder. They are fometimes obstructed by fmall pieces of gravel falling down from the kidneys, and lodging in them.

288 INFLAMMATION OF THE KIDNEYS.

heavinefs of the part, has frequent returns of chillnefs fhivering, &c. there is reafon to fufpect that matter is forming in the kidney, and that an abfcefs will enfue.

When matter in the urine flews that an ulcer is already formed in the kidney, the patient muff be careful to abftain from all actid, four, and falted provisions; and to live chiefly upon mild mucilaginous herbs and fruits, together with the broth of young animals, made with barley and common pot-herbs, &c. His drink may be whey, and butter-milk that is not four. The latter is by fome reckoned a fpecific remedy in ulcers of the kidneys. To anfwer this character, however, it muft be drank for a confiderable time. Chalybeate waters have likewife been found beneficial in this difeafe. This medicine is eafily obtained, as it is found in every part of Great Britain. It muft likewife be ufed for a confiderable time, in order to produce any falutary effects.

Those who are liable to frequent returns of inflammation, or obstructions of the kidneys, must abstain from wines, especially such as abound with tartar; and their food ought to be light and of easy digestion. They should use moderate exercise, not lie too hot, nor too much on their back, and avoid costiveness.

INFLAMMATION OF THE BLADDER.

THE inflammation of the bladder proceeds, in a great measure, from the same causes as that of the kidneys. It is known by an acute pain towards the bottom of the belly, and difficulty of passing urine, with some degree of fever, a constant inclination to go to stool, and a perpetual defire to make water.

This difeafe must be treated on the fame principles as the one immediately preceding. The diet must be light and thin, and the drink of a cooling nature. Bleeding is very proper at the beginning, and in robust conflitutions it will often be necessary to repeat it. The lower part of the belly should be fomented with warm water, or a decoction of mild vegetables; and emollient clysters ought frequently to be administered, &c.

The

INFLAMMATION OF THE BLADDER. 289

The patient fhould abstain from every thing that is of a hot, acrid, and stimulating quality; and should live entirely upon small broths, gruels, or mild vegetables.

A ftoppage of urine may proceed from other caufes befides an inflammation of the bladder; as a fwelling of the hæmorrhoidal veins; hard fæces lodged in the rectum; a ftone in the bladder; excrefcences in the urinary paffages, a palfy of the bladder, hysteric affections, &c. Each of these requires a particular treatment, which does not fall under our present confideration. We shall only obferve, that in all of them mild and gentle applications are the fafeft, as ftrong diuretic medicines, or things of an irritating nature, generally increase the danger. I have known fome perfons kill themfelves by introducing probes into the urinary paffages, to remove, as they thought, fomewhat that obstructed the discharge of urine, and others bring on a violent inflammation of the bladder, by using strong diuretics, as oil of turpentine, &c. for that purpole.

INFLAMMATION OF THE LIVER.

The liver is lefs fubject to inflammation than most of the other vifcera, as in it the circulation is flower; but when an inflammation does happen, it is with difficulty removed, and often ends in a fuppuration or fchirrus.

CAUSES.— Befide the common caufes of inflammation, we may here reckon the following, viz. exceflive fatnefs, a fchirrus of the liver itfelf, violent fhocks from ftrong vomits when the liver was before unfound, an adult or atrabiliarian ftate of the blood, any thing that fuddenly cools the liver after it has been greatly heated, ftones obftructing the courfe of the bile, drinking ftrong wines and fpirituous liquors, ufing hot fpicy aliment, obftinate hypochondriacal affections, &c.

SYMPTOMS.——This difeafe is known by a painful tenfion of the right fide under the falfe ribs, attended with fome degree of fever, a fenfe of weight, or fulnefs of the part, difficulty of breathing, loathing of food, great thirst, with a pale or yellowish colour of the skin and eyes.

The

290 INFLAMMATION OF THE LIVER.

The fymptoms here are various, according to the degree of inflammation, and likewife according to the particular part of the liver where the inflammation happens. Sometimes the pain is fo inconfiderable, that an inflammation is not fo much as fufpected; but when it happens in the upper or convex part of the liver, the pain is more acute, the pulfe quicker, and the patient is often troubled with a dry cough, a hiccup, and a pain extending to the fhoulder, with difficulty of lying on the left fide, &c.

This difeafe may be diftinguished from the pleurify, by the pain being lefs violent, feated under the false ribs, the pulse not so hard, and by the difficulty of lying on the left fide. It may be diffinguished from the hysteric and hypochondriac diforders by the degree of fever with which it is always attended.

This difeafe, if properly treated, is feldom mortal. A conftant hiccuping, violent fever, and exceflive thirft, are bad fymptoms. If it ends in a fuppuration, and the matter cannot be difcharged outwardly, the danger is great. When the fchirrus of the liver enfues, the patient, if he obferves a proper regimen, may neverthelefs live a number of years tolerably eafy; but if he indulge in animal food and ftrong liquors, or take medicines of an acrid or irritating nature, the fchirrus will be converted into a cancer, which muft infallibly prove fatal.

REGIMEN. — The fame regimen is to be obferved in this as in other inflammatory diforders. All hot things are to be carefully avoided, and cool diluting liquors, as whey, barley-water, &c. drank freely. The food must be light and thin, and the body, as well as the mind, kept eafy and quiet.

MEDICINE. — Bleeding is proper at the beginning of this difeafe, and it will often be neceffary, even though the pulfe fhould not feel hard, to repeat it. All violent purgatives are to be avoided; the body, however, must be kept gently open. A decoction of tamarinds, with a little honey or manna, will answer this purpofe very well. The fide affected must be fomented in the manner directed in the foregoing difeases. Mild laxative clyfters should be frequently administered; and, if the pain should notwithstanding continue violent, a blistering-

INFLAMMATION OF THE LIVER. 291

bliftering-plafter may be applied over the part affected; or rather a plafter made of gum ammoniac and vinegar of fquills.

Medicines which promote the fecretion of urine have a very good effect here. For this purpofe, half a drachm of purified nitre, or a tea-fpoonful of the fweet fpirits of nitre, may be taken in a cup of the patient's drink three or four times a-day.

When there is an inclination to fweat, it ought to be promoted, but not by warm fudorifics. The only thing to be ufed for that purpofe is plenty of diluting liquor drank about the warmth of the human blood. Indeed the patient in this cafe, as well as in all other topical inflammations, ought to drink nothing that is colder than the blood.

If the ftools fhould be loofe, and even ftreaked with blood, no means must be used to ftop them, unless they be fo frequent as to weaken the patient. Loose ftools often prove critical, and carry off the difease.

If an abfeefs or impofthume is formed in the liver, all methods fhould be tried to make it break and difcharge itfelf outwardly, as fomentations, the application of poultices, ripening cataplass, &c. Sometimes indeed the matter of an abfeefs comes away in the urine, and fometimes it is discharged by stool; but these are efforts of Nature which no means can promote. When the abfeefs bursts into the cavity of the *abdomen* at large, death must ensue; nor will the event be more favourable when the abfeefs is opened by an incision, unless in cafes where the liver adheres to the *peritonzeum*, fo as to form a bag for the matter, and prevent it from falling into the cavity of the *abdomen*; in which cafe opening the abfeefs by a fufficiently large incision will probably fave the patient's life *.

If the diforder, in fpite of all endeavours to the contrary, fhould end in a fchirrus, the patient must be careful to regulate his diet, &c. in fuch a manner as not to aggravate the difease. He must not indulge in flesh, fish,

• I know a gentleman who has had feveral abfceffes of the liver opened, and is now a firong and healthy man, though above eighty years of age.

ftrong

292 INFLAMMATION OF THE LIVER.

ftrong liquors, or any highly feafoned or falted provifions; but fhould, for the most part, live on mild vegetables, as fruits and roots, taking gentle exercise, and drinking whey, barley-water, or butter-milk. If he takes any thing ftronger, it fhould be fine mild ale, which is lefs heating than wines or spirits.

We fhall take no notice of inflammations of the other vifcera. They must in general be treated upon the fame principles as those already mentioned. The chief rule with respect to all of them is, to let blood, to avoid every thing that is strong, or of a heating nature, to apply warm fomentations to the part affected, and to cause the patient to drink a sufficient quantity of warm diluting liquors.

The difeafes mentioned in this chapter are generally relieved by warm fomentations, externally applied, and duly perfifted in. Thefe are made in a variety of ways; but the *Anodyne Fomentation*, recommended in the Appendix, to which a handful of camomile flowers may be occafionally added, will anfwer as well as any.

If the fomentations do not remove or abate the pain, recourfe muft be had to the warm bath, in which the patient is to continue as long as his ftrength will permit. The want of a proper warm bath may be fupplied by fome of the portable baths, filled with warm water. The most convenient of these contrivances, which are to be had at the tin-fhops, is commonly called the *flipper* bath, from its refembling a flipper in form. A cafk, or a common tub, may be used for the purpose upon an emergency, though not fo commodious.

Inflammations of the ftomach and bowels are ufually attended with obftinate coffiveness, for the removal of which no small skill and perfeverance are often necesfary. Sometimes a very mild medicine will operate, where a powerful one has had no effect. I have known a few spoonfuls of castor oil procure a stool, after the failure of strong drassic purges. The means, therefore, schoold be varied, not has had informed. Where one thing does not fucceed, another may be happily employed; and instances are not wanting of the efficacy even of external applications, when the best internal remedies have proved unfuccessful.

293

CHAP. XXXI.

OF THE CHOLERA MORBUS, AND OTHER EXCESSIVE DISCHARGES FROM THE STOMACH AND BOWELS.

THE cholera morbus is a violent purging and vomiting, attended with gripes, ficknefs, and a conftant defire to go to ftool. It comes on fuddenly, and is moft common in autumn. There is hardly any difeafe that kills more quickly than this, when proper means are not ufed in due time for removing it.

CAUSES.——It is occafioned by a redundancy and putrid acrimony of the bile; cold; food that eafily turns rancid or four on the ftomach; as butter, bacon, fweetmeats, cucumbers, melons, cherries, and other cold fruits*. It is fometimes the effect of ftrong acrid purges or vomits; or of poifonous fubftances taken into the ftomach. It may likewife proceed from violent paffions or affections of the mind; as fear, anger, &c.

SYMPTOMS.——It is generally preceded by a cardialgia, or heart-burn, four belchings, and flatulencies, with pain of the flomach and inteffines. To thefe fucceed exceffive vomiting, and purging of green, yellow, or blackifh coloured bile, with a diffention of the flomach, and violent griping pains. There is likewife a great thirft, with a very quick unequal pulfe, and often a fixed acute pain about the region of the navel. As the difeafe advances, the pulfe often finks fo low as to become quite imperceptible, the extremities grow cold, or cramped, and are often covered with a clammy fweat, the urine is obftructed, and there is a palpitation of the heart. Violent hiccuping, fainting, and convulfions, are the figns of approaching death.

MEDICINE. ——At the beginning of this difeafe, the efforts of Nature to expel the offending caufe fhould be affifted, by promoting the purging and voiniting.

* I have been twice brought to the gates of death by this difeafe, and both times it was occalioned by eating rancid bacon.

294 OF THE CHOLERA MORBUS, &c.

For this purpofe, the patient must drink freely of diluting liquors; as whey, butter-milk, warm water, thin water-gruel, fmall poffet, or, what is perhaps preferable to any of them, very weak chicken broth. This fhould not only be drank plentifully to promote the vomiting, but a clyster of it given every hour in order to promote the purging.

After these evacuations have been continued for some time, a decoction of toasted oat-bread may be drank to ftop the vomiting. The bread should be toasted till it is of a brown colour, and afterwards boiled in spring water. If oat-bread cannot be had, wheat-bread, or oat-meal well toasted, may be used in its stead. If this does not put a stop to the vomiting, two table-spoonfuls of the saline julep, with ten drops of laudanum, may be taken every hour till it ceases.

The vomiting and purging, however, ought never to be ftopped too foon. As long as these discharges do not weaken the patient, they are falutary, and may be allowed to go on, or rather ought to be promoted. But when the patient is weakened by the evacuations, which may be known from the finking of his pulfe, &c. recourfe must immediately be had to opiates, as recommended above; to which may be added ftrong wines, with fpirituous cinnamon-waters, and other generous cordials. Warm negus, or ftrong wine-whey, will likewife be neceffary to fupport the patient's fpirits, and promote the perspiration. His legs fhould be bathed in warm water, and afterwards rubbed with flannel cloths, or wrapped in warm blankets, and warm bricks applied to the foles of his feet. Flannels wrung out of warm fpirituous fomentations should likewife be applied to the region of the ftomach.

When the violence of the difeafe is over, to prevent a relapfe, it will be neceffary for fome time to continue the ufe of fmall dofes of laudanum. Ten or twelve drops may be taken in a glafs of wine, at leaft twice a-day, for eight or ten days. The patient's food ought to be nourifhing, but taken in fmall quantities, and he fhould ufe moderate exercife. As the ftomach and inteftines are generally much weakened, an infufion of the bark, or

OF THE CHOLERA MORBUS, &c. 295

or other bitters, in fmall wine, fharpened with the elixir of vitriol, may be drank for fome time.

Though phyficians are feldom called in due time in this difeafe, they ought not to defpair of relieving the patient even in the most defperate circumstances. Of this I lately faw a very striking proof in an old man and his fon, who had been both feized with it about the middle of the night. I did not fee them till next morning, when they had much more the appearance of dead than of living men. No pulse could be felt; the extremities were cold and rigid, the countenance was ghastly, and the strength almost quite exhausted. Yet from this deplorable condition they were both recovered by the use of opiates and cordial medicines.

I have frequently had occasion to fee this difeafe, and have fometimes felt it. Yet I never met with an instance, in my own practice, where it proved fatal, though we are told this often happens. Whether fo lamentable an iffue be owing to improper treatment, or to the extreme weaknefs of the patient's bowels, I cannot pretend to fay, without an exact knowledge of each particular cafe; but I am inclined to think, that when death is the confequence, the antidote, which is opium, has been too long delayed. No time fhould be loft in adminiftering it, upon the first ferious alarm, and before the powers of Nature are exhausted. What I generally prefcribe is laudanum, to be taken in cinnamon or fome other cordial water. Ten drops of laudanum may be added to two ounces of fimple cinnamon-water, and the draught repeated every two hours, or oftener, if neceffary.

I have found opiates no lefs fuccefsful in diarrhœa, or loofenefs. Ten grains of the powder of bole compounded with opium, given in a glafs of cordial water four or five times a-day, will feldom fail to check a recent diarrhœa, and, if judicioufly perfifted in, will often cure the most obstinate. I would therefore advise in fuch cafes, a full reliance on its final efficacy, rather than a rash impatience to try other medicines far more uncertain, and perhaps dangerous. But as a loofenefs may arife from a great variety of causes, how to adapt the mode of medical treatment to each will be explained in the next fection.

OF A DIARRHCEA, or LOOSENESS.

A loofenefs, in many cafes, is not to be confidered as a difeafe, but rather as a falutary evacuation. It ought, therefore, never to be ftopped, unlefs when it continues too long, or evidently weakens the patient. As this, however, fometimes happens, we fhall point out the most common caufes of a loofenefs, with the proper method of treatment.

When a loofenefs is occafioned by catching cold, or an obftructed perfpiration, the patient ought to keep warm, to drink freely of weak diluting liquors, to bathe his feet and legs frequently in lukewarm water, to wear flannel next his fkin, and to take every other method to reftore the perfpiration.

In a loofenels which proceeds from excels or repletion, a vomit is the proper medicine. Vomits not only cleanfe the ftomach, but promote all the fecretions, which renders them of great importance in carrying off a debauch. Half a drachm of ipecacuanha in powder will anfwer this purpole very well. A day or two after the vomit, the fame quantity of rhubarb may be taken, and repeated two or three times, if the loofenels continues. The patient ought to live upon light vegetable food of eafy digeftion, and to drink whey, thin gruel, or barley-water.

A loofenefs occafioned by the obftruction of any cuftomary evacuation, generally requires bleeding. If that does not fucceed, other evacuations may be fubftituted in the room of those which are obstructed. At the fame time, every method is to be taken to restore the usual discharges, as not only the cure of the discafe, but the patient's life, may depend on this.

A periodical loofeness ought never to be ftopped. It is always an effort of Nature to carry of fome offending matter, which, if retained in the body, might have fatal effects. Children are very liable to this kind of loofeness, especially while teething. It is, however, fo far from being hurtful to them, that fuch children generally get their teeth with least trouble. If these loofe ftools fhould

OF A DIARRHCEA, or LOOSENESS. 297

fhould at any time prove four or griping, a tea-fpoonful of magnefia alba, with four or five grains of rhubarb, may be given to the child in a little panado, or any other food. This, if repeated three or four times, will generally correct the acidity, and carry off the griping ftools. The potio cretacea, or chalk julep, may be adminiftered in dofes of two or three fpoonfuls after each evacuation; or a tea-fpoonful of fine powdered chalk may be mixed in a tea-cupful of water-gruel, and given occafionally.

A diarrhœa, or loofenefs, which proceeds from violent paffions or affections of the mind, must be treated with the greatest caution. Vomits in this cafe are highly improper. Nor are purges fafe, unlefs they be very mild, and given in fmall quantities. Opiates, and other antifpasmodic medicines, are most proper. Ten or twelve drops of liquid laudanum may be taken in a cup of valerian or penny-royal tea every eight or ten hours, till the fymptoms abate. Ease, cheerfulnes, and tranquillity of mind are here of the greatest importance.

When a loofenefs proceeds from acrid or poifonous fubftances taken into the ftomach, the patient muft drink large quantities of diluting liquors, with oil or fat broths, to promote vomiting and purging. Afterwards, if there be reafon to fufpect that the bowels are inflamed, bleeding will be neceffary. Small dofes of laudanum may likewife be taken to remove their irritation.

When the gout, repelled from the extremities occafions a loofenels, it ought to be promoted by gentle doles of rhubarb, or other mild purgatives. The gouty matter is likewife to be folicited towards the extremities by warm fomentations, cataplalms, &c. The perfpiration ought at the fame time to be promoted by warm diluting liquors; as wine-whey with fpirits of hartfhorn, or a few drops of liquid laudanum in it.

When a loolenefs proceeds from worms, which may be known from the fliminefs of the ftools, mixed with pieces of decayed worms, &c. medicines must be given to kill and carry off thefe vermin, as the powder of tin with purges of rhubarb and calomel. Afterwards limewater, either alone, or with a fmall quantity of rhubarb infufed,

298 OF A DIARRHEA, or LOOSENESS.

infufed, will be proper to ftrengthen the bowels, and prevent the new generation of worms.

A loofenefs is often occafioned by drinking bad water. When this is the cafe, the difeafe generally proves epidemical. When there is reafon to believe that this or any other difeafe proceeds from the ufe of unwholefome water, it ought immediately to be changed, or, if that cannot be done, it may be corrected by mixing with it quicklime, chalk, or the like.

In people whole ftomachs are weak, violent exercife immediately after eating will occafion a loofenefs. Though the cure of this is obvious, yet it will be proper, befides avoiding violent exercife, to use fuch medicines as tend to brace and ftrengthen the ftomach, as infufions of the bark, with other bitter and aftringent ingredients, in white-wine. Such perfons ought likewife to take frequently a glafs or two of old red port, or good claret.

From whatever caufe a loofenefs proceeds, when it is found neceffary to check it, the diet ought to confift of rice boiled with milk, and flavoured with cinnamon; rice-jelly, fago with red port; and the lighter forts of flefh-meat roafted. The drink may be thin water-gruel, rice-water, or weak broth made from lean veal, or with a fheep's head, as being more gelatinous than mutton, beef, or chicken-broth.

Perfons who, from a peculiar weaknefs, or too great an irritability of the bowels, are liable to frequent returns of this difeafe, fhould live temperately, avoiding crude fummer-fruits, all unwholefome foods, and meats of hard digeftion. They ought likewife to beware of cold, moifture, or whatever may obftruct the perfpiration, and fhould wear flannel next the fkin. All violent paffions, as fear, anger, &c. are likewife carefully to be guarded againft.

OF VOMITING.

Vomiting may proceed from various caufes; as excels in eating and drinking; foulnels of the ftomach; the acrimony of the aliments; a translation of the morbific matter of ulcers, of the gout, the eryfipelas, or other difeafes, difeafes, to the ftomach. It may likewife proceed from a loofenefs having been too fuddenly ftopped; from the ftoppage of any cuftomary evacuation, as the bleeding piles, the *menfes*, &c. from a weaknefs of the ftomach, the colic, the iliac paffion, a rupture, a fit of the gravel, worms; or from any kind of poifon taken into the ftomach. It is an ufual fymptom of injuries done to the brain; as contufions, comprefions, &c. It is likewife a fymptom of wounds or inflammations of the diaphragm, inteftines, fpleen, liver, kidneys, &c.

Vomiting may be occafioned by unufual motions, as failing, being drawn backwards in a carriage, &c. It may likewife be excited by violent paffions, or by the idea of naufeous or difagreeable objects, efpecially of fuch things as have formerly produced vomiting. Sometimes it proceeds from a regurgitation of the bile into the ftomach: in this cafe, what the patient vomits is generally of a yellow or greenifh colour, and has a bitter tafte. Perfons who are fubject to nervous affections are often fuddenly feized with violent fits of vomiting. Laftly, vomiting is a common fymptom of pregnancy. In this cafe it generally comes on about two weeks after the ftoppage of the *menfes*, and continues during the firft three or four months.

When vomiting proceeds from a foul ftomach or indigeftion, it is not to be confidered as a difeafe, but as the cure of a difeafe. It ought therefore to be promoted, by drinking lukewarm water, or thin gruel. If this does not put a ftop to the vomiting, a dofe of ipecacuanha may be taken, and worked off with weak camomile-tea.

When the retroceffion of the gout, or the obstruction of customary evacuations, occasion vomiting, all means must be used to restore these discharges; or, if that cannot be effected, their place must be supplied by others, as bleeding, purging, bathing the extremities in warm water, opening iffues, fetons, perpetual blisters, &c.

When vomiting is the effect of pregnancy, it may generally be mitigated by bleeding, and keeping the body gently open. The bleeding, however, ought to be in fmall quantities at'a time, and the purgatives fhould

be

OF VOMITING.

be of the mildeft kind, as figs, flewed prunes, manna, or fenna. Pregnant women are moft apt to vomit in the morning immediately after getting out of bed, which is owing partly to the change of pollure, but more to the emptinefs of the flomach. It may generally be prevented, by taking a difh of coffee, tea, or fome light breakfaft in bed. Pregnant women who are afflicted with vomiting, ought to be kept eafy both in body and mind. They fhould neither allow their flomachs to be quite empty, nor fhould they eat much at once. Cold water is a very proper drink in this cafe; if the flomach be weak, a little brandy may be added to it. If the fpirits be low, and the perfon apt to faint, a fpoonful of cinnamon-water, with a little marmalade of quinces or oranges, may be taken.

If vomiting proceeds from weaknefs of the ftomach, bitters will be of fervice. Peruvian bark infufed in wine or brandy, with as much rhubarb as will keep the body gently open, is an excellent medicine in this cafe. The elixir of vitriol is alfo a good medicine. It may be taken in the dofe of fifteen or twenty drops, twice or thrice a-day, in a glafs of wine or water. Habitual vomitings are fometimes alleviated by making oyfters a principal part of diet.

A vomiting which proceeds from acidities in the ftomach, is relieved by alkaline purges. The beft medicine of this kind is the magnefia alba, a tea-fpoonful of which may be taken in a difh of tea, or a little milk, three or four times a-day, or oftener if neceffary, to keep the body open.

When vomiting proceeds from violent paffions or affections of the mind, all evacuants muft be carefully avoided, efpecially vomits. Thefe are exceedingly dangerous. The patient in this cafe ought to be kept perfectly eafy and quiet, to have the mind foothed, and to take fome gentle cordial, as negus, or a little brandy and water, to which a few drops of laudanum may occafionally be added.

When vomiting proceeds from fpafmodic affections of the ftomach, mufk, caftor, and other antifpafmodic medicines, are of ufe. Warm and aromatic plafters

300

OF VOMITING.

ters have likewife a good effect. The ftomach-plafter of the London or Edinburgh difpenfatory may be applied to the pit of the ftomach, or a plafter of *theriaca*, which will anfwer rather better. Aromatic medicines may likewife be taken inwardly, as cinnamon or minttea, wine with fpiceries boiled in it, &c. The region of the ftomach may be rubbed with æther, or if that cannot be had, with ftrong brandy, or other fpirits. The belly fhould be fomented with warm water, or the patient immerfed up to the breaft in a warm bath.

I have always found the faline draughts taken in the act of effervescence, of fingular use in stopping a vomiting, from whatever cause it proceeded. These may be prepared by dissolving a drachm of the falt of tartar, in an ounce and a half of fresh lemon-juice, and adding to it an ounce of peppermint-water, the same quantity of simple cinnamon-water, and a little white sugar. This draught must be swallowed before the effervescence is quite over, and may be repeated every two hours, or oftener, if the vomiting be violent. A violent vomiting has sometimes been stopped by cupping on the region of the stomach after all other means had failed.

As the leaft motion will often bring on the vomiting again, even after it has been ftopped, the patient must avoid all manner of action. The diet must be fo regulated as to fit eafy upon the ftomach, and nothing fhould be taken that is hard of digestion. We do not, however, mean that the patient should live entirely upon flops. Solid food, in this cafe, often fits easier on the stomach than liquids.

CHAP. XXXII.

OF THE DIABETES, AND OTHER DISOR-DERS OF THE KIDNEYS AND BLADDER.

THE diabetes is a frequent and exceffive difcharge of urine. It is feldom to be met with among young people; but often attacks perfons in the decline of

OF THE DIABETES, &c.

of life, especially those who follow the more violent employments, or have been hard drinkers in their youth.

CAUSES. ----- A diabetes is often the confequence of acute difeafes, as fevers, fluxes, &c. where the patient has fuffered by exceffive evacuations; it may alfo be occafioned by great fatigue, as riding long journeys upon a hard-trotting horfe, carrying heavy burdens, running, &c. It may be brought on by hard drinking, or the use of strong stimulating diuretic medicines, as tincture of cantharides, fpirits of turpentine, and fuch like. It is often the effect of drinking too great quantities, of mineral waters. Many imagine that thefe will do them no fervice unless they be drank in great quantities, by which miftake it often happens that they occafion worfe difeafes than those they were intended to cure. In a word, this difeafe may either proceed from too great a laxity of the organs which fecrete the urine, from fomething that ftimulates the kidneys too much, or from a thin diffolved state of the blood, which makes too great a quantity of it run off by the urinary paffages.

SYMPTOMS.——In a diabetes, the urine generally exceeds in quantity all the liquid food which the patient takes. It is thin and pale, of a fweetifh tafte, and an agreeable fmell. The patient has a continual thirft, with fome degree of fever; his mouth is dry, and he fpits frequently a frothy ipittle. The ftrength fails, the appetite decays, and the flefh waftes away till the patient is reduced to fkin and bone. There is a heat of the bowels; and frequently the loins, tefficles, and feet are fwelled.

This difeafe may generally be cured at the beginning; but after it has continued long, the cure becomes very difficult. In drunkards, and very old people, a perfect cure is not to be expected.

REGIMEN. — Every thing that ftimulates the urinary paffages, or tends to relax the habit, must be avoided. For this reason the patient should live chiefly on solid food. His thirst may be quenched with acids; as forrel, juice of lemon, or vinegar. The mucilaginous vegetables, as rice, fago, and salop, with milk, are the most

302

OF THE DIABETES, &c.

most proper food. Of animal substances, shell-fish are to be preferred; as oysters, crabs, &c.

The drink may be Briftol-water. When that cannot be obtained, lime-water, in which a due proportion of oak-bark has been macerated, may be used. The white decoction *, with ifinglass diffolved in it, is likewise a very proper drink.

The patient ought daily to take exercife, but it fhould be fo gentle as not to fatigue him. He fhould lie upon a hard bed or matrafs. Nothing hurts the kidneys more than lying too foft. A warm dry air, the ufe of the flefhbrufh, and every thing that promotes perfpiration, is of fervice. For this reafon the patient ought to wear flannel next his fkin. A large ftrengthening-plafter may be applied to the back; or, what will anfwer better, a great part of the body may be wrapped in plafter.

MEDICINE. — Gentle purges, if the patient be not too much weakened by the difeafe, have a good effect. They may confift of rhubarb, with cardamum-feeds, or any other fpiceries, infufed in wine, and may be taken in fuch quantities as to keep the body gently open.

The patient must next have recourse to altringents and corroborants. Half a drachm of powder made of equal parts of allum and the infpissed juice, commonly called *Terra Japonica*, may be taken four times a day, or oftener, if the stomach will bear it. The allum must first be melted in a crucible; afterwards they may both be pounded together. Along with every dose of this powder the patient may take a tea-cupful of the tincture of roses †.

If the patient's ftomach cannot bear the allum in fubftance, whey may be made of it, and taken in the dofe of a tea-cupful three or four times a-day. The allum-whey is prepared by boiling two English quarts of milk over a flow fire, with three drachms of allum, till the curd feparates.

Opiates are of fervice in this difease, even though the patient refts well. They take off spafm and irrrita-

> * See Appendix, White Decotion. † See Appendix, Tinclure of Rofes.

E029 .

OF THE DIABETES, &c.

tion, and at the fame time leffen the force of the circulation. Ten or twelve drops of liquid laudanum may be taken in a cup of the patient's drink three or four times a-day.

The beft corroborants which we know, are the Peruvian bark and wine. A drachm of bark may be taken in a glafs of red port or claret three times a-day. The medicine will be both more efficacious and lefs difagreeable, if fifteen or twenty drops of the acid elixir of vitriol be added to each dofe. Such as cannot take the bark in fubftance, may ufe the decoction, mixed with an equal quantity of red wine, and fharpened as above.

There is a difeafe incident to labouring people in the decline of life, called an INCONTINENCY of Urine. But this is very different from a diabetes, as the water paffes off involuntarily by drops, and does not exceed the ufual quantity. This difeafe is rather troublefome than dangerous. It is owing to a relaxation of the fphincter of the bladder, and is often the effect of a palfy. Sometimes it proceeds from hurts, or injuries occafioned by blows, bruifes, preternatural labours, &c. Sometimes it is the effect of a fever. It may likewife be occafioned by a long ufe of ftrong diuretics, or of ftimulating medicines injected into the bladder.

This difease may be mitigated by the use of astringent and corroborating medicines, such as have been mentioned above; but we do not remember ever to have seen it cured.

In an incontinency of urine, from whatever caufe, a piece of fponge ought to be worn, or a bladder applied in fuch a manner as to prevent the urine from galling and excoriating the parts *.

OF A SUPPRESSION OF URINE.

It has already been obferved, that a fuppreffion of urine may proceed from various caufes; as an inflammation of the kidneys, or bladder; fmall ftones or gra-

• A bottle made of the India rubber, and properly applied, anfwers this purpose best. vel

OF A SUPPRESSION OF URINE.

vel lodging in the urinary paffages, hard faces lying in the rectum, pregnancy, a fpafm or contraction of the neck of the bladder, clotted blood in the bladder itfelf, a fwelling of the hæmorrhoidal veins, &c.

Some of these cases require the catheter, both to remove the obstructing matter, and to draw off the urine; but as this instrument can only be managed with fastery by perfors skilled in furgery, we shall say nothing further of its use. A bougee may be used by any cautious hand, and will often succeed better than the catheter.

We would chiefly recommend, in all obfructions of urine, fomentations and evacuants. Bleeding, as far as the patient's ftrength will permit, is neceffary, especially where there are fymptoms of topical inflammation. Bleeding in this cafe not only abates the fever, by leffening the force of the circulation, but, by relaxing the folids, it takes off the fpafm or ftricture upon the veffels, which occasioned the obftruction.

After bleeding, fomentations must be used. These may either confist of warm water alone, or of decoctions of mild vegetables; as mallows, camomile flowers, &c. Cloths dipped in these may either be applied to the part affected, or a large bladder filled with the decoction may be kept continually upon it. Some put the herbs themselves into a flannel bag, and apply them to the part, which is far from being a bad method. These continue longer warm than cloths dipped in the decoction, and at the fame time keep the part equally moift.

In all obstructions of urine, the body ought to be kept open. This is not, however, to be attempted by strong purgatives, but by emollient clysters, or gentle infusions of senna and manna. Clysters in this case not only open the body, but answer the purpose of an internal somentation, and greatly affist in removing the spass of the bladder and parts adjacent.

The food must be light, and taken in fmall quantities. The drink may be weak broth, or decoctions and infufions of mucilaginous vegetables, as marsh-mallow roots, lime-tree buds, &c. A tea-spoonful of the sweet spirits of nitre, or a drachm of Castile soap, may be frequently

X

put

305

306 OF A SUPPRESSION OF URINE.

put into the patient's drink; and, if there be no inflammation, he may drink fmall gin-punch.

Perfons fubject to a fuppreflion of urine ought to live very temperate. Their diet fhould be light, and their liquor diluting. They fhould avoid all acid and auftere wines, fhould take fufficient exercise, lie hard, and avoid ftudy and fedentary occupations.

OF THE GRAVEL AND STONE.

When fmall ftones are lodged in the kidneys, or difcharged along with the urine, the patient is faid to be afflicted with the gravel. If one of these ftones happen to make a lodgement in the bladder for some time, it accumulates fresh matter, and at length becomes too large to pass off with the urine. In this case the patient is faid to have the stone.

CAUSES.——The ftone and gravel may be occafioned by high living; the ufe of ftrong aftringent wines; a fedentary life; lying too hot, foft, or too much on the back; the conftant ufe of water impregnated with earthy or ftony particles; aliments of an aftringent or windy nature, &c. It may likewife proceed from an hereditary difposition. Perfons in the decline of life, and those who have been much afflicted with the gout or rheumatism, are most liable to it.

SYMPTOMS. — Small ftones or gravel in the kidneys occafion pain in the loins, ficknefs, vomiting, and fometimes bloody urine. When the ftone defcends into the *ureter*, and is too large to pafs along with eafe, all the above fymptoms are increafed; the pain extends towards the bladder; the thigh and leg of the affected fide are benumbed; the tefticles are drawn upwards, and the urine is obftructed.

A ftone in the bladder is known from a pain at the time, as well as before and after making water; from the urine coming away by drops, or ftopping fuddenly when it was running in a full ftream; by a violent pain in the neck of the bladder upon motion, efpecially on horfeback, or in a carriage on a rough road; or from a white, thick, copious ftinking mucous fediment in the urine;

urine; from an itching in the top of the *penis*; from bloody urine; from an inclination to go to ftool during the difcharge of urine; from the patient's paffing his urine more cafily when lying than in an erect pofture; from a kind of convultive motion occasioned by the fharp pain in difcharging the last drops of the urine; and lastly, from founding or fearching with the catheter.

REGIMEN. — Perfons afflicted with the gravel or ftone fhould avoid aliments of a windy or heating nature, as falt meats, four fruits, &c. Their diet ought chiefly to confift of fuch things as tend to promote the fecretion of urine, and to keep the body open. Artichokes, afparagus, fpinnage, lettuce, parfley, fuccory, purflane, turnips, potatoes, carrots, and radifhes, may be fafely eaten. Onions, leeks, and cellery are, in this cafe, reckoned medicinal. The moft proper drinks are whey, butter-milk, milk and water, barley-water ; decoctions or infufions of the roots of marfh-mallows, parfley, liquorice, or of other mild mucilaginous vegetables, as linfeed, lime-tree buds or leaves, &c. If the patient has been accuftomed to generous liquors, he may drink gin and water not too ftrong.

Gentle exercife is proper; but violent motion is apt to occafion bloody urine. We would therefore advife that it fhould be taken in moderation. Perfons afflicted with the gravel often pafs a great number of ftones after riding on horfeback, or in a carriage; but thofe who have a flone in the bladder are feldom able to bear thefe kinds of exercife. Where there is a hereditary tendency to this difeafe, a fedentary life ought never to be indulged. Were people careful, upon the first fymptoms of gravel, to obferve a proper regimen of diet, and to take fufficient exercife, it might often be carried off, or at least prevented from increasing; but if the fame courfe which occasioned the difeafe is perfifted in, it must be aggravated.

MEDICINE. In what is called a fit of the gravel, which is commonly occafioned by a ftone flicking in the *ureter*, or fome part of the urinary paffages, the patient must be bled; warm fomentations should likewife be applied to the part affected, emollient clysters X = X = X = X

administered, and diluting mucilaginous liquors drank, &c. The treatment of this cafe has been fully pointed out under the articles *inflammation of the kidneys and bladder*, to which we refer.

Dr. Whyte advifes patients who are fubject to frequent fits of gravel in the kidneys, but have no ftone in the bladder, to drink every morning, two or three hours before breakfaft, an Englifh pint of oyfter or cockle-fhell lime-water. The doctor very juftly obferves, that though this quantity might be too fmall to have any fenfible effect in diffolving a ftone in the bladder, yet it may very probably prevent its growth.

When a ftone is formed in the bladder, the Doctor recommends Spanish foap, and oyster or cockle-shell lime-water *, to be taken in the following manner: The patient must swallow every day, in any form that is least difagreeable, an ounce of the internal part of Alicant foap, and drink three or four English pints of oyster or cockle-shell lime-water: the foap is to be divided into three doses; the largest to be taken fasting in the morning early, the fecond at noon, and the third at feven in the evening; drinking above each dose a large draught of the lime-water; the remainder of which he may take any time betwixt dinner and supper, instead of other liquors.

The patient fhould begin with a fmaller quantity of the lime-water and foap than that mentioned above; at firft an Englifh pint of the former, and three drachms of the latter, may be taken daily. This quantity, however, he may increase by degrees, and ought to perfevere in the use of these medicines, especially if he finds any abatement of his complaints, for several months; nay, if the stone be very large, for years. It may likewise be proper for the patient, if he be severely pained, not only to begin with the soap and lime-water in stand of the first. However, after he has been for some time accustomed to these medicines, he may not only take the first water, but if he finds he can easily bear it, heighten its

· See Appendix, Lime-water.

diffolving

diffolving power still more by pouring it a fecond time on fresh calcined shells.

The cauftic alkali, or foap-lees, is the medicine chiefly in vogue at prefent for the ftone. It is of a very acrid nature, and ought therefore to be given in fome gelatinous or mucilaginous liquor; as veal-broth, new milk, linfeed-tea, a folution of gum-arabic, or a decoction of marfh-mallow-roots. The patient must begin with fmall dofes of the lees, as thirty or forty drops, and increase by degrees, as far as the ftomach can bear it*.

Though the foap-lees and lime-water are the molt powerful medicines which have hitherto been difcovered for the ftone, yet there are fome things of a more fimple nature, which in certain cafes are found to be beneficial, and therefore deferve a trial. An infufion of the feeds of *daucus fylve/tris*, or wild carrot, fweetened with honey, has been found to give confiderable eafe in cafes where the ftomach could not bear any thing of an acrid nature. A decoction of raw coffee-berries taken morning and evening, to the quantity of eight or ten ounces, with ten drops of fweet fpirit of nitre, has likewife been found very efficacious in bringing away large quantities of earthy matter in flakes. Honey is likewife found to be of confiderable fervice, and may be taken in gruel, or in any other form that is more agreeable.

The only other medicine which we fhall mention is the *uva urfi* It has been greatly extolled of late both for the gravel and ftone. It feems, however, to be in all refpects inferior to the foap and lime-water; but it is lefs difagreeable, and has frequently, to my knowledge, relieved gravelly complaints. It is generally taken in powder from half a drachm to a whole drachm, two or three times a-day. It may, however, be taken to the quantity of feven or eight drachms a-day, with great fafety and good effect.

* The cauftic alkali may be prepared by mixing two parts of quick-lime with one of pot-afhes, and fuffering them to fland till the lixivium be formed, which must be carefully filtrated before it be used. If the folution does not happen readily, a small quantity of water may be added to the mixture.

Of

309

Of all diforders in the urinary paffages the most tormenting is the stone in the bladder. The means of diffolving it, and bringing it away, though the frequent boast of quacks, have hitherto bassed medical inquiry. The British parliament, indeed, once paid five thousand pounds for a pretended solvent for the stone, which has long been forgotten. The fact, however, stands upon record as a signal proof of the extent to which credulity may be carried on the one hand, and successful imposture on the other.

The confiftence of fuch ftones varies fo much, that there is reafon to fear no medicine will ever be found fufficiently ftrong to diffolve the hardeft of them, without deftroying the bladder. Yet experiments on this fubject ought not to be difcontinued, as the object is great, and fome hard fubftances are known to be foluble in feemingly mild ones.

I have known feveral inftances where ftones, after getting into the urethra, were brought away by means of a bent probe; but how to get them there, is the difficulty. It can only happen while they are fmall, though I have feen flattifh ftones brought away in this manner, which meafured two inches round. I have fometimes thought that riding on a hard-trotting horfe, or in a carriage on a rough road, might tend to bring down a fmall ftone.

Most people troubled with the flone are guilty of one great error. They put off the operation too long. When it is certainly known that there is a flone in the bladder, and that it is too large to get along the urethra, no time ought to be lost in having it cut out, before the patient's habit becomes too irritable, or the flone is fo far increased in fize, that it cannot be extracted without a laceration of the parts.

CHAP.

[311]

CHAP. XXXIII.

OF INVOLUNTARY DISCHARGES OF BLOOD.

SPONTANEOUS or involuntary difcharges of blood often happen from various parts of the body. Thefe, however, are to far from being always dangerous, that they often prove falutary. When fuch difcharges are critical, which is frequently the cafe in fevers, they ought not to be ftopped. Nor, indeed, is it proper at any time to ftop them, unlefs they be fo great as to endanger the patient's life. Moft people, afraid of the fmalleft difcharge of blood from any part of the body, fly immediately to the ufe of ftyptic and aftringent medicines, by which means an inflammation of the brain, or fome other fatal difeafe, is occafioned, which, had the difcharge been allowed to go on, might have been prevented.

Periodical difcharges of blood, from whatever part of the body they proceed, must not be stopped. They are always the efforts of Nature to relieve herfelf; and fatal difeases have often been the confequence of obstructing them. It may, indeed, be sometimes necessary to check the violence of such discharges; but even this requires the greatest caution. Instances might be given where the stopping of a small periodical flux of blood from one of the stopping, has proved fatal to the health.

In the early period of life, bleeding at the nofe is very common. Those who are farther advanced in years are more liable to hæmoptoe, or discharge of blood from the lungs. After the middle period of life, hæmorrhoidal fluxes are most common; and in the decline of life, discharges of blood from the urinary passages.

Involuntary fluxes of blood may proceed from very different, and often from quite opposite causes. Sometimes they are owing to a particular construction of the body, as a sanguine temperament, a laxity of the vessels, a plethoric habit, &c. At other times they proceed from a determination of the blood towards one particular part,

as

INVOLUNTARY DISCHARGES 312

as the head, the hæmorrhoidal veins, &c. They may likewife proceed from an inflammatory disposition of the blood, in which cafe there is generally fome degree of fever : this likewife happens when the flux is occafioned by an obstructed perspiration, or a stricture upon the fkin, the bowels, or any particular part of the fystem.

But a diffolved state of the blood will likewife occafion hæmorrhages. Thus, in putrid fevers, the dyfentery, the fcurvy, the malignant fmall-pox, &c. there are often very great discharges of blood from different parts of the body. They may likewife be brought on by too liberal an use of medicines which tend to diffolve the blood, as cantharides, the volatile alkaline falts, &c. Food of an acrid or irritating quality may likewife occafion hæmorrhages; as alfo ftrong purges and vomits, or any thing that greatly ftimulates the bowels.

Violent paffions or agitations of the mind will likewife have this effect. These often cause bleeding at the nofe, and I have known them fometimes occafion an hæmorrhage in the brain. Violent efforts of the body, by overstraining or hurting the veffels, may have the fame effect, efpecially when the body is long kept in an unnatural pofture, as hanging the head very low, &c.

The cure of an hæmorrhage must be adapted to its caufe. When it proceeds from too much blood, or a tendency to inflammation, bleeding, with gentle purges and other evacuations, will be neceffary. It will, likewife, be proper for the patient in this cafe to live chiefly upon a vegetable diet, to avoid all ftrong liquors, and food that is of an acrid, hot, or flimulating quality. The body fhould be kept cool, and the mind eafy.

When an hæmorrhage is owing to a putrid or diffolved state of the blood; the patient ought to live chiefly upon acid fruits with milk, and vegetables of a nourifhing nature, as fago, falop, &c. His drink may be wine diluted with water, and fharpened with the juice of lemon, vinegar, or fpirits of vitriol. The best medicine in this cafe is the Peruvian bark, which may be taken according to the urgency of the fymptoms.

When a flux of blood is the effect of acrid food, or of ftrong ftimulating medicines, the cure is to be effected by

OF BLOOD.

by foft and mucilaginous diet. The patient may likewife take frequently about the bulk of a nutmeg of Locatelli's balfam, or the fame quantity of fpermaceti.

When an obstructed perspiration, or a stricture upon any part of the system, is the cause of an hæmorrhage, it may be removed by drinking warm diluting liquors, lying a-bed, bathing the extremities in warm water, &c.

OF BLEEDING AT THE NOSE.

Bleeding at the nofe is commonly preceded by fome degree of quickness of the pulse, flushing in the face, pulsation of the temporal arteries, heaviness in the head, dimness of the fight, heat and itching of the nostrils, &c.

To perfons who abound with blood, this difcharge is very falutary. It often cures a vertigo, the head-ach, a phrenzy, and even an epilepfy. In fevers, where there is a great determination of blood towards the head, it is of the utmost fervice. It is likewife beneficial in inflammations of the liver and spleen, and often in the gout and rheumatifm. In all difeases where bleeding is neceffary, a spontaneous difcharge of blood from the nose is of much more fervice than the fame quantity let with a lancet.

In a difcharge of blood from the nole, the great point is to determine whether it ought to be flopped or not. It is a common practice to thop the bleeding, without confidering whether it be a difease, or the cure of a difease. This conduct proceeds from fear; but it has often bad, and sometimes fatal confequences.

When a difcharge of blood from the nofe happens in an inflammatory dileafe, there is always realon to believe that it may prove falutary; and therefore it fhould be fuffered to go on, at leaft as long as the patient is not weakened by it.

When it happens to perfons in perfect health, who are full of blood, it ought not to be fuddenly itopped, effecially if the fymptoms of plethora, mentioned above, have preceded it. In this cafe it cannot be ftopped without rifking the patient's life.

314 OF BLEEDING AT THE NOSE.

In fine, whenever bleeding at the nofe relieves any bad fymptom, and does not proceed fo far as to endanger the patient's life, it ought not to be ftopped. But when it returns frequently, or continues till the pulfe becomes low, the extremities begin to grow cold, the lips pale, or the patient complains of being fick or faint, it must immediately be ftopped.

For this purpofe the patient fhould be fet nearly upright, with his head reclining a little, and his legs immerfed in water about the warmth of new milk. His hands ought likewife to be put in lukewarm water, and his garters may be tied a little tighter than ufual. Ligatures may be applied to the arms, about the place where they are ufually made for bleeding, and with nearly the fame degree of tightnefs. Thefe must be gradually flackened as the blood begins to ftop, and removed entirely as foon as it gives over.

Sometimes dry lint put up the noftrils will ftop the bleeding. When this does not fucceed, doffils of lint dipped in ftrong fpirits of wine may be put up the noftrils, or if that cannot be had, they may be dipped in brandy. Blue vitriol diffolved in water may likewife be ufed for this purpofe, or a tent dipped in the white of an egg well beat up, may be rolled in a powder made of equal parts of white fugar, burnt allum, and white vitriol, and put up the noftril from whence the blood iffues.

Internal medicines can hardly be of ufe here, as they have feldom time to operate. It may not, however, be amifs to give the patient half an ounce of Glauber's falt, and the fame quantity of manna, diffolved in four or five ounces of barley-water. This may be taken at a draught, and repeated if it does not operate in a few hours. Ten or twelve grains of nitre may be taken in a glafs of cold water and vinegar every hour, or oftener if the ftomach will bear it. If a ftronger medicine be neceffary, a tea-cupful of the tincture of rofes, with twenty or thirty drops of the weak fpirit of vitriol, may be taken every hour. When thefe things cannot be had,

OF BLEEDING AT THE NOSE.

had, the patient may drink water, with a little common falt in it, or equal parts of water and vinegar *.

If the genitals be immerfed for fome time in cold water, it will generally ftop a bleeding at the nofe. I have not known this fail.

Sometimes, when the bleeding is ftopped outwardly, it continues inwardly. This is very troublefome, and requires particular attention, as the patient is apt to be fuffocated with the blood, especially if he falls asleep, which he is very ready to do after losing a great quantity of blood.

When the patient is in danger of fuffocation from the blood getting into his throat, the paffages may be ftopped by drawing threads up the noftrils, and bringing them out at the mouth, then fastening pieces of sponge, or small rolls of linen cloth to their extremities; afterwards drawing them back, and tying them on the outfide with a sufficient degree of tightness.

After the bleeding is ftopped, the patient ought to be kept as eafy and quiet as poffible. He fhould not pick his nofe, nor take away the tents or clotted blood till they fall off of their own accord, and fhould not lie with his head low.

Those who are affected with frequent bleeding at the nose, ought to bathe their feet often in warm water, and keep them warm and dry. They ought to wear nothing tight about their necks, to keep their body as much in an erect posture as possible, and never to view any object obliquely. If they have too much blood, a vegetable diet, with now and then a cooling purge, is the fafest way to leften it.

But when the difeafe proceeds from a thin diffolved ftate of the blood, the diet fhould be rich and nourifhing; as ftrong broths and jellies, fago-gruel with wine and fugar, &c. Infufions of the Peruvian bark in wine ought likewife to be taken, and perfifted in for a confiderable time.

* From ten to twenty drops of the oil of turpentine in a little water given frequently, feldom fails to ftop a bleeding at the nofe, or from any other part.

OF THE BLEEDING AND BLIND PILES.

A difcharge of blood from the hæmorrhoidal veffels is called the *bleeding piles*. When the veffels only fwell, and difcharge no blood, but are exceeding painful, the difeafe is called the *blind piles*.

Perfons of a loofe fpongy fibre, of a bulky fize, who live high, and lead a fedentary, inactive life, are most fubject to this difeafe. It is often owing to an hereditary difposition. Where this is the cafe, it attacks perfons more early in life than when it is accidental. Men are more liable to it than women, especially those of a fanguine, plethoric, or a fcorbutic habit, or of a melancholy difposition.

The piles may be occafioned by an excefs of blood, by ftrong aloetic purges, high-feafoned food, drinking great quantities of fweet wines, the neglect of bleeding, or other cuftomary evacuations, much riding, great coftivenefs, or any thing that occafions hard or difficult ftools. Anger, grief, or other violent paffions, will likewife occafion the piles. I have often known them brought on by fitting on the damp ground. A pair of thin breeches will excite the diforder in a perfon who is fubject to it, and fometimes even in thofe who never had it before. Pregnant women are often afflicted with the piles.

A flux of blood from the *anus* is not always to be treated as a difeafe. It is even more falutary than bleeding at the nofe, and often prevents or carries off difeafes. It is peculiarly beneficial in the gout, rheumatifm, afthma, and hypochondriacal complaints, and often proves critical in colics, and inflammatory fevers.

In the management of the patient, regard must be had to his habit of body, his age, ftrength, and manner of living. A discharge which might be excessive and prove hurtful to one, may be very moderate, and even falutary to another. That only is to be esteemed dangerous, which continues too long, and is in such quantity as to waste the patient's strength, hurt the digestion, nutrition, and other functions necessary to life.

BLEEDING AND BLIND PILES. 317

When this is the cafe, the difcharge must be checked by a proper regimen, and astringent medicines. The DIET must be cool but nourifhing, confisting chiefly of bread, milk, cooling vegetables, and broths. The DRINK may be chalybeate water, orange-whey, decoctions or infusions of the astringent and mucilaginous plants, as the tormentil root, biftort, the marsh-mallow-roots, &c.

Old conferve of red rofes is a very good medicine in this cafe. It may be mixed with new milk, and taken in the quantity of an ounce three or four times a-day. This medicine is in no great repute, owing to its being feldom taken in fuch quantity as to produce any effects; but when taken as here directed, and duly perfifted in, I have known it perform very extraordinary cures in violent hæmorrhages, efpecially when affifted by the tincture of rofes; a tea-cupful of which may be taken about an hour after every dofe of the conferve.

The Peruvian bark is likewife proper in this cafe, both as a ftrengthener and aftringent. Half a drachm of it may be taken in a glafs of red wine, fharpened with a few drops of the elixir of vitriol, three or four times a-day.

The bleeding piles are fometimes periodical, and return regularly once a month, or once in three weeks. In this cafe they are always to be confidered as a falutary difcharge, and by no means to be ftopped. Some have entirely ruined their health by ftopping a periodical difcharge of blood from the hæmorrhoidal veins.

In the *blind piles*, bleeding is generally of ufe. The diet muft be light and thin, and the drink cool and diluting. It is likewife neceffary that the body be kept gently open. This may be done by fmall dofes of the flower of brimftone and cream of tartar. Thefe may be mixed in equal quantities, and a tea-fpoonful taken two or three times a-day, or oftener if neceffary. Or an ounce of the flower of brimftone and half an ounce of purified nitre may be mixed with three or four ounces of the lenitive electuary, and a tea-fpoonful of it taken three or four times a-day.

Emollient clysters are here likewife beneficial; but there is fometimes fuch an aftriction of the anus, that

they

318 BLEEDING AND BLIND PILES.

they cannot be thrown up. In this cafe I have known a vomit have a very good effect.

When the piles are exceeding painful and fwelled, but difcharge nothing, the patient muft fit over the fleams of warm water. He may likewife apply a linen cloth dipped in warm fpirits of wine to the part, or poultices made of bread and milk, or of leeks fried with butter. If thefe do not produce a difcharge, and the piles appear large, leeches muft be applied as near them as poflible, or if they will fix upon the piles themfelves, fo much the better. When leeches will not fix, the piles may be opened with a lancet. The operation is very eafy, and is attended with no danger.

Various ointments, and other external applications, are recommended in the piles; but I do not remember to have feen any effects from these worth mentioning. Their principal use is to keep the part moift, which may be done as well by a fost poultice, or an emollient cataplasm. When the pain, however, is very great, a liniment made of two ounces of emollient ointment, and half an ounce of liquid laudanum, beat up with the yolk of an egg, may be applied.

Perhaps no other caufe of the piles is fo frequent as the ufe of aloetic purgatives. It is to be observed that aloes forms a confiderable share of almost all advertised purging pills. A costive habit is more effectually, and much more fafely removed by a spoonful of castor oil taken occasionally in an evening.

When the piles are very painful, the beft external application is a weak folution of fugar of lead with a little laudanum. An ointment made of one third finely powdered galls, and two thirds hog's lard, is very ufeful. When the piles are feated high, relief may frequently be obtained from injections of lime-water, or of an infufion of galls.

The pain of the piles is very often removed by an emetic, or by taking twice a day thirty drops of balfam of copaiva on a little moift fugar. When a pile has a narrow neck, it is beft extirpated by the knife. If the pile be large, or has a broad bafis, a double ligature may be paffed through it, and tied on each fide.

When .

BLEEDING AND BLIND PILES.

When piles are neglected, they are very apt to produce a *fiftula*. This complaint is difcovered by a ftain of matter on the linen, which, on examination, will be found to proceed from a fmall orifice in the neighbourhood of the anus. Various local remedies are recommended for this complaint. The utility of all of them depends on their power of ftimulating the fides of the ulcer into more active inflammation, fo as to make them cohere together. On this principle, I think I have feen advantage from taking a wine-glafs of fea-water every night for a month or fix weeks. Irritating injections have been ufed with fimilar intentions, and have fometimes, when duly perfifted in, fucceeded.

The only certain radical cure for a fiftula is a furgical operation, the object of which is to reduce the ulcer to the ftate of a fimple wound, and as fuch to heal it. This fhould never be too long neglected. The difeafe gradually diffufes itfelf in various directions through the cellular fubftance furrounding the rectum; and new openings are formed, which render the complaint more difficult to be removed.

There are two ways of performing the operation. One is by paffing a filk thread, or piece of flexible gold wire, in at the external orifice of the fiftula, and bringing it out at the anus, and then twifting the ends together, which is daily repeated till it cuts its way out. By fome timid people this mode of cure is preferred to the knife : and, though kept a fecret by fome pretenders to medical knowledge, is as old as the hiftory of furgery. The incifion, however, is the more certain and effectual mode of eradicating the difeafe; and if fuffering is to be effimated by duration, the lefs painful alfo.

SPITTING OF BLOOD.

We mean here to treat of that difcharge of blood from the lungs only which is called an *hæmoptoe*, or *fpitting* of blood. Perfons of a flender make, and a lax fibre, who have long necks and ftrait breafts, are most liable to this difeafe. It is most common in the fpring, and generally attacks people before they arrive at the prime or middle period of life. It is a common observation, that those who have have been fubject to bleeding at the nofe when young, are afterwards most liable to an hæmoptoe.

This difeafe may likewife proceed from wounds of the These may either be received from without, or lungs. they may be occafioned by hard bodies getting into the wind-pipe, and fo falling down upon the lungs, and hurting that tender organ. The obstruction of any customary evacuation may occafion a fpitting of blood; as neglect of bleeding or purging at the ufual feafons, the ftoppage of the bleeding piles in men, or the menfes in women, &c. It may likewife proceed from a polypus, fchirrous concretions, or any thing that obstructs the circulation of the blood in the lungs. It is often the effect of a long and violent cough; in which cafe it is generally the forerunner of a confumption. A violent degree of cold fuddenly applied to the external parts of the body will occasion an hæmoptoe. It may likewife be occafioned by breathing air which is too much rarified to be able properly to expand the lungs. This is often the cafe with those who work in hot places, as furnaces, glafs-houfes, or the like. It is likewife faid to happen to fuch, as afcend to the top of very high mountains, as the Peak of Teneriff, &c.

Spitting of blood is not always to be confidered as a primary difeafe. It is often only a fymptom, and in fome difeafes not an unfavourable one. This is the cafe in pleurifies, peripneumonies, and fundry other fevers. In a dropfy, fcurvy, or confumption, it is a bad fymptom, and fhews that the lungs are ulcerated.

SYMPTOMS.——Spitting of blood is generally preceded by a fenfe of weight, and opprefiion of the breaft, a dry tickling cough, hoarfenefs, and a difficulty of breathing. Sometimes it is ufhered in with fhivering, coldnefs

nefs of the extremities, coftivenefs, great laffitude, flatulence, pain of the back and loins, &c. As thefe flew a general ftricture upon the veffels, and a tendency of the blood to inflammation, they are commonly the forerunners of a very copious difcharge. The above fymptoms do not attend a difcharge of blood from the gums or fauces, by which means they may always be diftinguifhed from an hæmoptoe. Sometimes the blood that is fpit up is thin, and of a florid red colour; and at other times it is thick, and of a dark or blackifh colour; nothing, however, can be inferred from this circumflance, but that the blood has lain a longer or fhorter time in the breaft before it was difcharged.

Spitting of blood, in a ftrong healthy perfon, of a found conflitution, is not very dangerous; but when it attacks the tender and delicate, or perfons of a weak lax fibre, it is with difficulty removed. When it proceeds from a fchirrus or polypus of the lungs, it is bad. The danger is greater when the difcharge proceeds from the rupture of a large veffel, than of a fmall one. When the extravafated blood is not fpit up, but lodges in the breaft, it corrupts, and greatly increafes the danger. When the blood proceeds from an ulcer in the lungs, it is generally fatal.

REGIMEN.——The patient ought to be kept cool and eafy. Every thing that heats the blood, or quickens the circulation, increases the danger. The mind ought likewife to be foothed, and every occasion of exciting the passions avoided. The diet should be fost, cooling, and shender; as rice boiled with milk, fmall broths, barleygruels, panado, &c. The diet, in this case, can scarces be too low. Even water-gruel is sufficient to support the patient for some days. All strong liquors must be avoided. The patient may drink milk and water, barley-water, whey, butter-milk, and fuch like. Every thing, however, should be drank cold, and in small quantities at a time. He should observe the strictes filence, or at least speak with a very low voice.

MEDICINE. ——— This, like the other involuntary difcharges of blood, ought not to be fuddenly ftopped by aftringent medicines. More mifchief is often

Y

done

done by thefe, than if it were fuffered to go on. It may however, proceed fo far as to weaken the patient, and even endanger his life; in which cafe proper means must be used for reftraining it.

The body fhould be kept gently open by laxative diet; as roafted apples, flewed prunes, and fuch like. If these should not have the defired effect, a tea-spoonful of the lenitive electuary may be taken twice or thrice aday, as is found neceffary. If the bleeding proves violent, ligatures may be applied to the extremities, as directed for a bleeding at the nose.

If the patient be hot or feverifh, bleeding and fmall dofes of nitre will be of ufe; a fcruple or half a drachm of nitre may be taken in a cup of his ordinary drink twice or thrice a-day. His drink may likewife be fharpened with acids, as juice of lemon, or a few drops of the fpirit of vitriol; or he may take frequently a cup of the tincture of rofes.

Bathing the feet and legs in lukewarm water, has likewife a very good effect in this difeafe. Opiates too are fometimes beneficial; but thefe must be administered with caution. Ten or twelve drops of laudanum may be given in a cup of barley-water twice a-day, and continued for fome time, provided they be found beneficial.

The conferve of rofes is likewife a very good medicine in this cafe, provided it be taken in fufficient quantity, and long enough perfifted in. It may be taken to the quantity of three or four ounces a-day; and, if the patient be troubled with a cough, it fhould be made into an electuary with balfamic fyrup, and a little of the fyrup of poppies.

If stronger astringents be necessary, fifteen or twenty drops of the elixir of vitriol may be given in a glass of water three or four times a-day.

Those who are subject to frequent returns of this difease should avoid all excess. Their diet should be light and cool, confisting chiefly of milk and vegetables. Above all, let them beware of vigorous efforts of the body, and violent agitations of the mind.

322

VOMIT-

VOMITING OF BLOOD.

This is not fo common as the other difcharges of blood which have already been mentioned; but it is very dangerous, and requires particular attention.

Vomiting of blood is generally preceded by pain of the fromach, ficknefs, and naufea; and is accompanied with great anxiety, and frequent fainting fits.

This difeafe is fometimes periodical; in which cafe it is lefs dangerous. It often proceeds from an obftruction of the menfes in women; and fometimes from the ftopping of the hæmorrhoidal flux in men. It may be occafioned by any thing that greatly ftimulates or wounds the ftomach, as ftrong vomits or purges, acrid poifons, fharp or hard fubftances taken into the ftomach, &c. It is often the effect of obftructions in the liver, the fpleen, or fome of the other vifcera. It may likewife proceed from external violence, as blows or bruifes, or from any of the caufes which produce inflammation. In hyfteric women, vomiting of blood is a very common, but by no means a dangerous fymptom.

A great part of the danger in this difease arises from the extravafated blood lodging in the bowels, and becoming putrid, by which means a dyfentery or putrid fever may be occafioned. The best way of preventing this, is to keep the body gently open, by frequently exhibiting emollient clyfters. Purges must not be given till the discharge is stopt, otherwise they will irritate the ftomach, and increase the diforder. All the food and drink must be of a mild cooling nature, and taken in small quantities. Even drinking cold water has fometimes proved a remedy, but it will fucceed better when fharpened with the weak fpirits of vitriol. When there are figns of an inflammation, bleeding may be neceffary; but the patient's weaknefs will feldom permit it. Opiates may be of use; but they must be given in very small doles, as four or five drops of liquid laudanum twice or thrice a-day.

After the difcharge is over, as the patient is generally troubled with gripes, occasioned by the acrimony of the

Y 2

blood

VOMITING OF BLOOD.

blood lodged in the inteffines, gentle purges will be neceffary.

OF BLOODY URINE.

This is a difcharge of blood from the veffels of the kidneys or bladder, occafioned by their being either enlarged, broken, or eroded. It is more or lefs dangerous according to the different circumstances which attend it.

When pure blood is voided fuddenly, without interruption and without pain, it proceeds from the kidneys; but if the blood be in fmall quantity, of a dark colour, and emitted with heat and pain about the bottom of the belly, it proceeds from the bladder. When bloody urine is occasioned by a rough stone descending from the kidneys to the bladder, which wounds the ureter, it is attended with a fharp pain in the back, and difficulty of making water. If the coats of the bladder are hurt by a ftone, and the bloody urine follows, it is attended with the most acute pain, and a previous stoppage of urine.

Bloody urine may, likewife, be occafioned by falls, blows, the lifting or carrying of heavy burdens, hard riding, or any violent motion. It may also proceed from ulcers of the bladder, from a ftone lodged in the kidneys, or from violent purges, or fharp diuretic medicines, efpecially cantharides.

Bloody utine is always attended with fome degree of danger : but it is peculiarly fo when mixed with purulent matter, as this fhews an ulcer fomewhere in the urinary paffages. Sometimes this difcharge proceeds from excefs of blood, in which cafe it is rather to be confidered as a falutary evacuation than a difeafe. If the difcharge, however, be very great, it may walle the patient's ftrength, and occafion an ill habit of body, a dropfy, or a confumption.

The treatment of this diforder must be varied according to the different caufes from which it proceeds.

When it is owing to a ftone in the bladder, the cure: depends upon an operation, a description of which would be foreign to our purpofe. If

OF BLOODY URINE.

If it be attended with a plethora, and fymptoms of inflammation, bleeding will be neceffary. The body muft likewife be kept open by emollient clyfters, or cooling purgative medicines; as cream of tartar, rhubarb, manna, or fmall dofes of lenitive electuary.

When bloody urine proceeds from a diffolved state of the blood, it is commonly the fymptom of fome malignant difease; as the small-pox, a putrid fever, or the like. In this case the patient's life depends on the liberal use of the Peruvian bark, wine and acids, as has already been shewn.

When there is reafon to fufpect an ulcer in the kidneys or bladder, the patient's diet must be cool, and his drink of a foft, healing, balfamic quality, as decoctions of marsh-mallow roots with liquorice, folutions of gumarabic, &c. Three ounces of marsh-mallow roots, and half an ounce of liquorice, may be boiled in two English quarts of water to one; two ounces of gum-arabic, and half an ounce of purified nitre, may be diffolved in the ftrained liquor, and a tea-cupful of it taken four or five times a-day.

The early use of aftringents in this difease has often bad confequences. When the flux is stopped too foon, the grumous blood, by being confined in the vessels, may produce inflammations, abscess, and ulcers. If, however, the case be urgent, or the patient seems to fuffer from the loss of blood, gentle astringents may be necessary. In this case the patient may take three or four ounces of lime-water, with half an ounce of the tincture of Peruvian bark, three times a-day; or he may take an ounce or two of the conferve of roses three or four times a-day, drinking a tea-cupful of the tincture of aster it; or if stronger styptics be necessary, a dram of Armenian bole in a cup of whey may be taken three or four times a-day.

OF THE DYSENTERY, OR BLOODY FLUX.

This difeafe prevails in the fpring and autumn. It is most common in marshy countries, where, after hot and dry summers, it is apt to become epidemic. Perfons are most liable to it who are much exposed to the night-air, or who live in places where the air is confined and un-

¥ 3

wholefome,

wholesome. Hence it often proves fatal in camps, on fhipboard, in jails, hofpitals, and fuch like places.

CAUSES. ---- The dyfentery may be occafioned by any thing that obstructs the perspiration, or renders the humours putrid; as damp beds, wet clothes, unwholefome diet, bad air, &c. But it is most frequently communicated by infection. This ought to make people extremely cautious in going near fuch perfons as labour under the difeafe. Even the fmell of the patient's excrements has been known to communicate the infection.

SYMPTOMS.----It is known by a flux of the belly, attended by violent pain of the bowels, a conftant inclination to go to ftool, and generally more or lefs blood in the ftools. It begins, like other fevers, with chilnefs, lofs of strength, a quick pulle, great thirst, and an inclination to vomit. The ftools are at first greafy and frothy, afterwards they are ftreaked with blood, and at laft have frequently the appearance of pure blood, mixed with fmall filaments refembling bits of fkin. Worms are fometimes paffed both upwards and downwards through the whole courfe of the difeafe. When the patient goes to ftool, he feels a bearing down, as if the whole bowels were falling out, and fometimes a part of the inteftine is actually protruded, which proves exceeding troublefome, especially in children. Flatulency is likewife a troublefome fymptom, especially towards the end of the difeafe.

This difease may be diftinguished from a diarrhœa, or loofenefs, by the acute pain of the bowels, and the blood which generally appears in the ftools. It may be diftinguished from the cholera morbus by its not being attended with fuch violent and frequent fits of vomiting, &c.

When the dyfentery attacks the old, the delicate, or fuch as have been wafted by the gout, the fcurvy, or other lingering difeafes, it generally proves fatal. Vomiting and hiccuping are bad figns, as they thew an inflammation of the ftomach. When the ftools are green, black, or have an exceeding difagreeable cadaverous fmell, the danger is very great, as it fhews the difeafe to be of the putrid kind. It is an unfavourable fymptom when clysters are immediately returned; but still more for when

6

when the paffage is fo obffinately flut, that they cannot be injected. A feeble pulfe, coldnefs of the extremities, with difficulty of fwallowing, and convultions, are figns of approaching death.

RÉGIMEN.——Nothing is of more importance in this difeafe than cleanlinefs. It contributes greatly to the recovery of the patient, and no lefs to the fafety of fuch as attend him. In all contagious difeafes the danger is increafed, and the infection fpread, by the neglect of cleanlinefs; but in no one more than in this. Every thing about the patient fhould be frequently changed. The excrement fhould never be fuffered to continue in his chamber, but removed immediately and buried under ground. A conftant ftream of frefh air fhould be admitted into the chamber; and it ought frequently to be fprinkled with vinegar, juice of lemon, or fome other ftrong acids.

The patient must not be difcouraged, but his fpirits kept up in hopes of a cure. Nothing tends more render any putrid difeafe mortal, than the fears and apprehensions of the fick. All difeases of this nature have a tendency to fink and depress the spirits, and when that is increased by fears and alarms from those whom the patient believes to be perfons of skill, it cannot fail to have the worst effects.

A flannel waiftcoat worn next the fkin has often a very good effect in the dyfentery. This promotes the perfpiration without over-heating the body. Great caution, however, is neceffary in leaving it off. I have often known a dyfentery brought on by imprudently throwing off a flannel waiftcoat before the feafon was fufficiently warm. For whatever purpofe this piece of drefs is worn, it flould never be left off but in a warm feafon.

In this difease the greatest attention must be paid to the patient's diet. Flesh, fish, and every thing that has a tendency to turn putrid or rancid on the stomach, must be abstained from. Apples boiled in milk, water-pap, and plain light pudding, with broth made of the gelatinous parts of animals, may constitute the principal part of the patient's food. Gelatinous broth not only answers the purpose of food, but likewise a medicine. I have often known dysenteries, which were not of a putrid nature,

cured

Y4

cured by it, after pompous medicines had proved ineffectual *.

Another kind of food very proper in the dyfentery, which may be used by fuch as cannot take the broth mentioned above, is made by boiling a few handfuls of fine flour, tied in a cloth, for fix or feven hours, till it becomes as hard as flarch. Two or three table-fpoonfuls of this may be grated down, and boiled in fuch a quantity of new milk and water as to be of the thickness of pap. This may be fweetened to the patient's tafte, and taken for his ordinary food \dagger .

In a *putrid dyfentery* the patient may be allowed to eat freely of most kinds of good ripe fruit; as apples, grapes, gooseberries, currant-berries, strawberries, &c. These may either be eaten raw or boiled, with or without milk, as the patient chuses. The prejudice against fruit in this difease is so great, that many believe it to be the

* The manner of making this broth is, to take a fheep's head and feet, with the fkin upon them and to burn the wool off with a hot iron; afterwards to boil them till the broth is quite a jelly. A little cinnamon or mace may be added, to give the broth an agreeable flavour, and the patient may take a little of it warm with toafted bread three or four times a-day. A clyfter of it may likewife be given twice a-day. Such as cannot ufe the broth made in this way, may have the head and feet fkinned; but we have reafon to believe that this injures the medicine. It is not our bufinefs here to reafon upon the nature and qualities of medicine, otherwife this might be fhewn to poffefs virtues every way fuited to the cure of a dyfentery which does not proceed from a putrid flate of the humours. One thing we know, which is preferable to all reafoning, that whole families have often been cured by it, after they had used many other medicines in vain. It will, however, be proper that the patient take a vomit, and a dofe or two of rhubarb, before he begins to use the broth. It will likewise be neceffary to continue the use of it for a confiderable time, and to make it the principal food.

† The learned and humane Dr. Rutherford, late profeffor of medicine in the univerfity of Edinburgh, ufed to mention this food in his public lectures with great encomiums. He directed it to be made by tying a pound or two of the fineft flour, as tight as poffible, in a linen rag, afterwards to dip it frequently in water, and to dridge the outfide with flour till a cake or cruft was formed around it, which prevents the water from foaking into it while boiling. It is then to be boiled till it becomes a hard dry mafs, as directed above. This, when mixed with milk and water, will not only anfwer the purpofe of food, but may likewife be given in clyfters.

common caufe of dyfenteries. This, however, is an egregious miftake. Both reafon and experience fhew, that good fruit is one of the beft medicines, both for the prevention and cure of the dyfentery. Good fruit is in every refpect calculated to counteract that tendency to putrefaction, from whence the most dangerous kind of dyfentery proceeds. The patient in fuch a cafe ought therefore to be allowed to eat as much fruit as he pleafes, provided it be ripe *.

The most proper drink in this diforder is whey. The dyfentery has often been cured by the ufe of clear whey alone. It may be taken both for drink, and in form of clyster. When whey cannot be had, barley water sharpened with cream of tartar may be drank, or a decoction of barley and tamarinds; two ounces of the former and one of the latter may be boiled in two Engliss quarts of water to one. Warm water, water-gruel, or water wherein hot iron has been frequently quenched, are all very proper, and may be drank in turns. Camomile-tea, if the stomach will bear it, is an exceeding proper drink. It both strengthens the stomach, and by its antifeptic quality tends to prevent a mortification of the bowels.

MEDICINE.——At the beginning of this difeafe it is always neceffary to cleanfe the first passages. For this purpose a vomit of ipecacuanha must be given, and wrought off with weak camomile tea. Strong vomits are feldom neceffary here. A scruple, or at most half a

* I lately faw a young man who had been feized with a dyfentery in North America. Many things had been tried there for his relief, but to no purpofe. At length, tired out with difappointments from medicine, and reduced to fkin and bone, he came over to Britain, rather with a view to die among his relations, than with any hopes of a cure. After taking fundry medicines here with no better fuccefs than abroad, I advifed him to leave off the ufe of drugs, and to truft entirely to a diet of milk and fruits, with gentle exercife. Strawberries was the only fruit he could procure at that feafon. Thefe he ate with milk twice and fometimes thrice a-day. The confequence was, that in a fhort time his ftools were reduced from upwards of twenty in a-day, to three or four, and fometimes not fo many. He ufed the other fruits as they came in, and was in a few weeks fo well as to leave that part of the country where I was, with a view to return to America.

329

drachm,

OF THE DYSENTERY,

330.

drachm, of ipecacuanha, is generally fufficient for an adult, and fometimes a very few grains will fuffice. The day after the vomit, half a drachm, or two fcruples of rhubarb, muft be taken; or, what will anfwer the purpofe rather better, an ounce, or an ounce and a half, of Epfom falts. This dofe may be repeated every other day for two or three times. Afterwards fmall dofes of ipecacuanha may be taken for fome time. Two or three grains of the powder may be mixed in a table-fpoonful of the fyrup of poppies, and taken three times a-day.

These evacuations, and the regimen prescribed above, will often be sufficient to effect a cure. Should it, however, happen otherwise, the following astringent medicines may be used :

A clyfter of ftarch or fat mutton broth, with thirty or forty drops of liquid laudanum in it, may be administered twice a-day. At the fame time an ounce of gum arabic, and half an ounce of gum-tragacanth, may be diffolved in an English pint of barley water, over a flow fire, and a table-spoonful of it taken every hour.

If these have not the defired effect, the patient may take, four times a-day, about the bulk of a nutmeg of the *Japonic confection*, drinking after it a tea-cupful of the decoction of logwood *.

Perfons who have been cured of this difeafe are very liable to relapfe; to prevent which, great circumfpection with refpect to diet is neceffary. The patient mult abftain from all fermented liquors, except now and then a glafs of good wine; but he mult drink no kind of malt liquor. He fhould likewife abftain from animal food, as fifh and flefh, and live principally on milk and vegetables.

Gentle exercife and wholefome air are likewife of importance. The patient fhould go to the country as foon as his ftrength will permit, and fhould take exercife daily on horfeback, or in a carriage. He may likewife ufe bitters infufed in wine or brandy, and may drink twice a day a gill of lime-water mixed with an equal quantity of new milk.

* See Appendix, Decodion of Logwood.

When

OR BLOODY FLUX.

When dyfenteries prevail, we would recommend a ftrict attention to cleanlinefs, a fpare use of animal food, and the free use of found ripe fruits, and other vegetables. The night-air is to be carefully avoided, and all communication with the fick. Bad fmells are likewife to be shunned, especially those which arise from putrid animal substances. The necessaries where the fick go are carefully to be avoided.

When the first fymptoms of the dysentery appear, the patient ought immediately to take a vomit, to go to bed, and drink plentifully of weak warm liquor, to promote a fweat. This, with a dose or two of rhubarb at the beginning, would often carry off the difease. In countries where dysenteries prevail, we would advise fuch as are liable to them, to take either a vomit or a purge every spring or autumn, as a preventive.

There are fundry other fluxes of the belly, as the LIENTERY and CŒLIAC PASSION, which, though lets dangerous than the dyfentery, yet merit confideration. These difeafes generally proceed from a relaxed ftate of the ftomach and inteffines, which is fometimes to great, that the food paffes through them with hardly any fenfible alteration; and the patient dies merely from the want of nourifhment.

When the lientery or cœliac paffion fucceeds to a dyfentery, the cafe is bad. They are always dangerous in old age, efpecially when the conflictution has been broken by excefs or acute difeafes. If the ftools be very frequent and quite crude, the thirft great, with little urine, the mouth ulcerated, and the face marked with fpots of different colours, the danger is very great.

The treatment of the patient is in general the fame as in the dyfentery. In all obfinate fluxes of the belly, the cure mult be attempted, by first cleanfing the ftomach and bowels with gentle vomits and purges; afterwards fuch a diet as has a tendency to heal and strengthen the bowels, with opiates and astringent medicines, will generally complete the cure.

The fame observation holds with respect to a TE-NESMUS, or frequent defire of going to stool. This difease

OF THE DYSENTERY, &c.

332

difeafe refembles the dyfentery fo much, both in its fymptoms and method of cure, that we think it needlefs to infift upon it.

CHAP. XXXIV.

OF THE HEAD-ACH,

A CHES and pains proceed from very different caufes, and may affect any part of the body; but we shall point out those only which occur most frequently, and are attended with the greatest danger.

When the head-ach is flight, and affects a particular part of the head only, it is called *cephalalgia*; when the whole head is affected, *cephalæa*; and when one fide only, *hemicrania*. A fixed pain in the forehead, which may be covered with the end of the thumb, is called the *clavis by/tericus*.

There are also other diffinctions. Sometimes the pain is internal, fometimes external; fometimes it is an original difease, and at other times only fymptomatic. When the head-ach proceeds from a hot bilious habit, the pain is very acute and throbbing, with a confiderable heat of the part affected. When from a cold phlegmatic habit, the patient complains of a dull heavy pain, and has a sense of coldness in the part. This kind of head-ach is fometimes attended with a degree of flupidity or folly.

Whatever obftructs the free circulation of the blood through the veffels of the head, may occafion a headach. In perfons of a full habit, who abound with blood, the head-ach often proceeds from the fuppreffion of cuftomary evacuations; as bleeding at the nofe, fweating of the feet, &c. It may likewife proceed from any caufe that determines a great flux of blood towards the head; as coldnefs of the extremities, or hanging down the head for a long time. Whatever prevents the return of the blood/from the head, will likewife occafion a head-ach; as looking long obliquely at any object, wearing

OF THE HEAD-ACH.

wearing any thing tight about the neck, a new hat, or the like.

When a head-ach proceeds from the ftoppage of a running at the nofe, there is a heavy, obtufe, preffing pain in the fore-part of the head, in which there feems to be fuch a weight, that the patient can fcarcely hold it up. When it is occafioned by the cauftic matter of the venereal difeafe, it generally affects the fkull, and often produces a *caries* of the bones.

Sometimes the head-ach proceeds from the repulfion or retroceffion of the gout, the eryfipelas, the fmall-pox, meafles, itch, or other eruptive difeafes. What is called a *hemicrania* generally proceeds from crudities or indigeftion. Inanition, or emptinefs, will also occasion head-achs. I have often feen inflances of this in nurfes who gave fuck too long, or who did not take a fufficient quantity of folid food.

There is likewife a most violent, fixed, constant, and almost intolerable head-ach, which occasions great debility both of body and mind, prevents fleep, destroys the appetite, causes a *vertigo*, dimness of fight, a noise in the ears, convulsions, epileptic fits, and sometimes vomiting, costiveness, coldness of the extremities, &c.

The head-ach is often fymptomatic in continual and intermitting fevers, efpecially quartans. It is likewife a very common fymptom in hysteric and hypochondriac complaints.

When a head-ach attends an acute fever, with pale urine, it is an unfavourable fymptom. In exceffive headachs, coldness of the extremities is a bad fign.

When the difease continues long, and is very violent, it often terminates in blindness, an apoplexy, deafness, a *wertigo*, the palfy, or the epilepfy.

In this difeafe the cool regimen in general is to be obferved. The diet ought to confift of fuch emollient fubftances as will correct the acrimony of the humours, and keep the body open; as apples boiled in milk, fpinnage, turnips, and fuch like. The drink ought to be diluting; as barley-water, infufions of mild mucilaginous vegetables, decoctions of the fudorific woods, &c. The feet and legs ought to be kept warm, and frequently bathed bathed in lukewarm water; the head fhould be fhaved, and bathed with water and vinegar. The patient ought, as much as poffible, to keep in an erect pofture, and not to lie with his head too low.

When the head-ach is owing to excels of blood, or an hot bilious conftitution, bleeding is neceffary. The patient may be bled in the jugular vein, and the operation repeated if there be occafion. Cupping allo, or the application of leeches to the temples, and behind the ears, will be of fervice; afterwards a bliftering plafter may be applied to the neck, behind the ears, or to any part of the head that is most affected. In fome cafes it will be proper to blifter the whole head. In perfons of a grofs habit, iffues, or perpetual blifters, will be of fervice. The body ought likewife to be kept open by gentle laxatives.

But when the head-ach proceeds from a copious vitiated *ferum*, ftagnating in the membranes, either within or without the fkull, with a dull, heavy, continual pain, which will neither yield to bleeding nor gentle laxatives, then more powerful purgatives are neceffary, as pills made of aloes, refin of jalap, or the like. It will alfo be neceffary in this cafe to blifter the whole head, and to keep the back-part of the neck open for a confiderable time by a perpetual blifter.

When the head-ach is occafioned by the ftoppage of a running at the nofe, the patient fhould frequently fmell to a bottle of volatile falts; he may likewife take fnuff, or any thing that will irritate the nofe, fo as to promote a difcharge from it; as the herb maftich, ground-ivy, &c.

A *hemicrania* efpecially a periodical one, is generally owing to a foulnefs of the ftomach, for which gentle vomits must be administered, as also purges of rhubarb. After the bowels have been sufficiently cleared, chalybeate waters, and such bitters as ftrengthen the ftomach, will be neceffary. A periodical head-ach has been cured by wearing a piece of flannel over the forehead during the night.

When the head-ach arifes from a vitiated flate of the humours, as in the fcurvy and venereal difease, the patient, after proper evacuations, must drink freely of the decoction of woods, or the decoction of farfaparilla, with raifins

OF THE HEAD-ACH.

fins and liquorice*. Thefe, if duly perfifted in, will produce very happy effects. When a collection of matter is felt under the fkin, it must be difcharged by an incifion, otherwife it will render the bone carious.

When the head-ach is fo intolerable as to endanger the patient's life, or is attended with continual watching and delirium, recourfe muft be had to opiates. Thefe, after proper evacuations by clyfters, or mild purgatives, may be applied both externally and internally. The affected part may be rubbed with Bate's anodyne balfam, or a cloth dipped in it may be applied to the part. The patient may, at the fame time, take twenty drops of laudanum, in a cup of valerian or penny-royal tea, twice or thrice a-day. This is only to be done in cafe of extreme pain. Proper evacuations ought always to accompany and follow the ufe of opiates \ddagger .

When the patient cannot bear the lofs of blood, his feet ought frequently to be bathed in lukewarm water, and well rubbed with a coarfe cloth. Cataplaims with muftard or horfe-radifh ought likewife to be applied to them. This courfe is peculiarly neceffary when the pain proceeds from a gouty humour affecting the head.

When the head-ach is occafioned by great heat, hard labour, or violent exercise of any kind, it may be allayed by cooling medicines; as the faline draughts with nitre, and the like.

A little æther, dropt into the palm of the hand, and applied to the forehead, will fometimes remove a violent head-ach.

OF THE TOOTH-ACH.

This difeafe is fo well known, that it needs no defcription. It has great affinity with the rheumatifm, and often fucceeds pains of the fhoulders and other parts of the body.

* See Appendix, Decotion of Sarfaparilla.

+ When the pain is very violent, and does not yield to fmall dofes of laudanum, the quantity may be increased. I have known a patient in extreme pain take three hundred drops in twentyfour hours; but fuch dofes ought only to be administered by a perfon of skill.

It may proceed from obstructed perspiration, or any of the other caufes of inflammation. I have often known the tooth-ach occafioned by neglecting fome part of the ufual coverings of the head, by fitting with the head bare near an open window, or exposing it to a draught of cold air. Food or drink taken either too hot or too cold, is very hurtful to the teeth. Great quantities of fugar, or other fweet-meats, are likewife hurtful. Nothing is more deftructive to the teeth than cracking nuts, or chewing any kind of hard fubstances. Picking the teeth with pins, needles, or any thing that may hurt the enamel with which they are covered, does great mischief, as the tooth is fure to be fpoiled whenever the air gets into it. Breeding women are very fubject to the toothach, especially during the first three or four months of pregnancy. The tooth-ach often proceeds from fcorbutic humours affecting the gums. In this cafe the teeth are fometimes wafted, and fall out without any confiderable degree of pain. The more immediate caufe of the tooth-ach is a rotten or *carious* tooth.

In order to relieve the tooth-ach, we muſt first endeavour to leffen the flux of humours to the part affected. This may be done by mild purgatives, fcarifying the gums, or applying leeches to them, and bathing the feet frequently with warm water. The perspiration ought likewife to be promoted, by drinking freely of weak wine-whey, or other diluting liquors, with small doses of nitre. Vomits too have often an exceeding good effect in the tooth-ach. It is feldom fafe to administer opiates, or any kind of heating medicines, or even to draw a tooth, till proper evacuations have been premifed; and thefe alone will often effect the cure.

If this fail, and the pain and inflammation ftill increase, a fuppuration may be expected, to promote which a toafted fig fhould be held between the gum and the cheek; bags filled with boiled camomile-flowers, flowers of elder, or the like, may be applied near the part affected, with as great a degree of warmth as the patient can bear, and renewed as they grow cool: the patient may likewife receive the steams of warm water into his mouth, through an inverted funnel, or by holding his head

OF THE TOOTH-ACH.

head over the mouth of a porringer filled with warm water. Gargles are likewife of ufe to promote a difcharge. Rob of elder diffolved in fmall beer makes a very proper gargle, or an infufion of fage or mulberry leaves.

Such things as promote the difcharge of faliva, or caufe the patient to fpit, are generally of fervice. For this purpofe, bitter, hot, or pungent vegetables may be chewed; as gentian, calamus aromaticus, or pellitory of Spain. Allen recommends the root of *yellow water flower-de-luce* in this cafe. This root may either be rubbed upon the tooth, or a little of it chewed. Brookes fays, he hardly ever knew it fail to cure the tooth-ach. It ought, however, to be ufed with caution.

Many other herbs, roots, and feeds, are recommended for curing the tooth-ach; as the leaves or roots of millefoil or yarrow chewed, tobacco fmoked or chewed, ftaves. acre, or the feeds of muftard chewed, &c. Thefe bitter, hot, and pungent things, by occasioning a greater flow of *faliva*, frequently give eafe in the tooth-ach.

Opiates often relieve the tooth-ach. For this purpofe, a little cotton wet with laudanum may be held between the teeth; or a piece of flicking-plafter, about the bignefs of a fhilling, with a bit of opium in the middle of it, of a fize not to prevent the flicking of the other, may be laid on the temporal artery, where the pulfation is most fenfible. De la Motte affirms, that there are few cafes wherein this will not give relief. If there be a hollow tooth, a fmall pill made of equal parts of camphire and opium, put into the hollow, is often beneficial. When this cannot be had, the hollow tooth may be filled with gum mastich, wax, lead, cork, or any fubstance that will flick in it, and keep out the external air.

Few applications give more relief in the tooth-ach than bliftering-plafters. These may be applied between the shoulders; but they have the best effect when put behind the ears, and made so large as to cover a great part of the lower jaw. Burning the nerve within the affected tooth with a hot iron, has frequently given ease; but this operation ought to be performed with care.

• After all, when a tooth is carious, it is often impoffible to remove the pain without extracting it; and as a fpoiled

Z

tooth

tooth never becomes found again, it is prudent to draw it foon, left it fhould affect the reft. Tooth-drawing, like bleeding, is very much practifed by mechanics, as well as perfons of the medical profeffion. The operation, however, is not without danger, and ought always to be performed with care. A perfon unacquainted with the ftructure of the parts, will be in danger of hurting the jaw-bone, or of drawing a found tooth inftead of a rotten one *. When a found tooth has been drawn, if it be replaced immediately, it will grow in again; and it is now a practice to draw a rotten tooth and put a found one from another perfon in its room.

When the tooth-ach returns periodically, and the pain chiefly affects the gums, it may be cured by the bark.

Some pretend to have found great benefit in the toothach, from the application of an artificial magnet to the affected tooth. We fhall not attempt to account for its mode of operation; but, if it be found to anfwer, though only in particular cafes, it certainly deferves a trial, as it is attended with no expence, and cannot do any harm. Electricity has likewife been recommended, and particular inftruments have been invented for fending a flock through the affected tooth.

Perfons who have returns of the tooth-ach at certain feafons, as fpring and autumn, might often prevent it by taking a purge at these times.

Keeping the teeth clean has no doubt a tendency to prevent the tooth-ach. The best method of doing this is to wash them daily with falt and water, a decoction of the bark, or with cold water alone. All brushing and fcraping of the teeth is dangerous, and, unless it be performed with great care, does mischief.

Of all the aches and pains incident to the human body, I do not know of one more diffreffing than the toothach. It is not, indeed, dangerous; but it fo much diffurbs and deranges every function, as to render the greateft man incapable of thinking or acting with propriety. I know of many things that will eafe the raging pain for

* This may always be prevented by the operator firiking upon the teeth with any piece of metal, as this never fails to excite the pain in the carious tooth.

OF THE TOOTH-ACH.

the moment, but none that will cure it, where the tooth is carious. In this cafe, I would always recommend the extraction of the unfound tooth, rather than to endure the pain for years, and be obliged to fubmit to the operation at laft. It is to be remembered that a rotten tooth can never get better, or become found; and that, by being left in, it only tends to vitiate the reft, and to do mifchief. Great care and expertnefs, however, are neceffary in extracting it.

OF THE EAR-ACH.

This diforder chiefly affects the membrane which lines the inner cavity of the ear called the *meatus auditorius*. It is often fo violent as to occafion great reftlefinefs, anxiety, and even delirium. Sometimes epileptic fits, and other convulfive diforders, have been brought on by extreme pain in the ear.

The ear-ach may proceed from any of the caufes which produce inflammation. It often proceeds from a fudden fupprefion of perfpiration, or from the head being expofed to cold when covered with fweat. It may alfo be occafioned by worms, or other infects getting into the ear, or being bred there; or from any hard body flicking in the ear. Sometimes it proceeds from the tranflation of morbific matter to the ear. This often happens in the decline of malignant fevers, and occafions deafnefs, which is generally reckoned a favourable fymptom.

When the ear-ach proceeds from infects, or any hard body flicking in the ear, every method muft be taken to remove them as foon as possible. The membranes may be relaxed by dropping into the ear oil of fweet almonds, or olive oil. Afterwards the patient should be made to fneeze, by taking souff, or some strong sternutatory. If this should not force out the body, it must be extracted by art. I have seen infects, which had got into the ear, come out of their own accord upon pouring in oil.

When the pain of the ear proceeds from inflammation, it must be treated like other topical inflammations, by a cooling regimen, and opening medicines. Bleeding at the beginning, either in the arm or jugular vein, or cupping in the neck, will be proper. The ear may likewife Z 2 be

OF THE EAR-ACH.

be fomented with fteams of warm water; or flannel bags filled with boiled mallows and camomile-flowers may be applied to it warm; or bladders filled with warm milk and water. An exceeding good method of fomenting the ear is to apply it close to the mouth of a jug filled with warm water, or a ftrong decoction of camomile-flowers.

The patient's feet fhould be frequently bathed in lukewarm water, and he ought to take fmall dofes of nitre and rhubarb, viz. a fcruple of the former, and ten grains of the latter, three times a-day. His drink may be whey, or decoction of barley and liquorice, with figs or raifins. The parts behind the ear ought frequently to be rubbed with camphorated oil, or a little of the volatile liniment, and a few drops of the camphorated fpirit of wine may be put into the ear with wool or cotton. A blifter behind the ear, if applied early, will fometimes remove this complaint.

When the inflammation cannot be difcuffed, a poultice of bread and milk, or roafted onions, may be applied to the ear, and frequently renewed, till the abfcefs breaks, or can be opened. Afterwards the humours may be diverted from the part by gentle laxatives, blifters, or iffues; but the difcharge must not be fuddenly dried up by any external application.

PAIN OF THE STOMACH, &c.

This may proceed from various caufes, as indigeftion; wind; the acrimony of the bile; fharp, acrid, or poifonous fubftances taken into the ftomach, &c. It may likewife be occafioned by worms; the ftoppage of cuftomary evacuations; or from a translation of gouty matter to the ftomach, the bowels, &c.

Women in the decline of life are very liable to pains of the ftomach and bowels, efpecially fuch as are afflicted with hyfteric complaints. It is likewife very common to hypochondriac men of a fedentary and luxurious life. In fuch perfons it often proves fo extremely obftinate as to baffle all the powers of medicine.

When the pain of the ftomach is most violent after eating, there is reason to fuspect that it proceeds from fome fault either in the digestion or the food. In this case,

PAIN OF THE STOMACH, &c.

341

ftomach

cafe, the patient ought to change his diet, till he finds what kind of food agrees beft with his ftomach, and fhould continue chiefly to ufe it. If a change of diet does not remove the complaint, the patient may take a gentle vomit, and afterwards a dofe or two of rhubarb. He ought likewife to take an infufion of camomile-flowers, or fome other ftomachic bitter, either in wine or water. I have often known exercife remove this complaint, efpecially failing, or a long journey on horfeback, or in a carriage.

When a pain of the flomach proceeds from flatulency, the patient is conftantly belching up wind, and feels an uneafy diffention of the flomach after meals. This is a most deplorable difease, and is feldom thoroughly cured. In general, the patient ought to avoid all windy diet, and every thing that sours on the flomach, as greens, roots, &c. This rule, however, admits of some exceptions. There are many inflances of perfons very much troubled with wind, who have received great benefit from eating parched pease, though that grain is generally supposed to be of a windy nature *.

This complaint may likewife be greatly relieved by labour, efpecially digging, reaping, mowing, or any kind of active employment by which the bowels are alternately comprefied and dilated. The most obstinate cafe of this kind I ever met with was in a perfon of a fedentary occupation, whom I advised, after he had tried every kind of medicine in vain, to turn gardener; which he did, and has ever fince enjoyed good health.

When a pain of the flomach is occafioned by the fwallowing of acrid or poifonous fubftances, they must be difcharged by vomit : this may be excited by butter, oils, or other foft things, which fheath and defend the flomach from the acrimony of its contents.

When a pain of the ftomach proceeds from a translation of gouty matter, warm cordials are neceffary, as generous wines, French brandy, &c. Some have drank a whole bottle of brandy or rum in this cafe in a few hours, without being in the least intoxicated, or even feeling the

* These are prepared by sceping or foaking pease in water, and afterwards drying them in a pot or kiln till they burst. They may be used at pleasure.

342 PAIN OF THE STOMACH, &c.

ftomach warmed by it. It is impoffible to afcertain the quantity neceffary upon these occasions. This must be left to the feelings and discretion of the patient. The fafer way however is, not to go too far. When there is an inclination to vomit, it may be promoted by drinking an infusion of camomile-flowers, or *carduus benedictus*.

If a pain of the ftomach proceed from the ftoppage of cuftomary evacuations, bleeding will be neceffary, especially in fanguine and very full habits. It will likewife be of use to keep the body gently open by mild purgatives; as rhubarb or fenna. When this difease affects women in the decline of life, after the ftoppage of the *menses*, making an iffue in the leg or arm will be of peculiar fervice.

When the difeafe is occafioned by worms, they must be deftroyed, or expelled by fuch means as are recommended in the following fection.

When the ftomach is greatly relaxed, and the digeftion bad, which often occafion flatulencies, the elixir of vitriol will be of fingular fervice. Fifteen or twenty drops of it may be taken in a glass of wine or water twice or thrice a-day.

Perfons afflicted with flatulency are generally unhappy unlefs they be taking fome purgative medicines; thefe, though they may give immediate eafe, tend to weaken and relax the flomach and bowels, and confequently increafe the diforder. Their beft method is to mix purgatives and flomachics together. Equal parts of Peruvian bark and rhubarb may be infufed in brandy or wine, and taken in fuch quantity as to keep the body gently open.

4 1

Pain of the ftomach proceeds from fuch a variety of caufes, that it is difficult to prefcribe a medicine for it. The treatment must of course be fuited to the nature of the complaint. But I have for fome years very generally recommended a plaster, which feldom fails to give relief. Its basis may be any kind of adhesive plaster spread upon leather, to which, while warm, a drachm and a half, or two drachms, of powdered opium may be added. It should be large enough to cover nearly the whole region of the stomach, and should be suffered to remain on as long as it will stok.

CHAP.

[343]

CHAP. XXXV.

OF WORMS.

THESE are chiefly of three kinds, viz. the tania, or tape-worm; the teres, or round and long worm; and the afcarides, or round and fhort worm. There are many other kinds of worms found in the human body; but as they proceed, in a great measure, from fimilar causes, have nearly the same symptoms, and require almost the same method of treatment as these already mentioned, we shall not spend time in enumerating them.

The tape-worm is white, very long, and full of joints. It is generally bred either in the ftomach or fmall inteftines. The round and long worm is likewife bred in the fmall guts, and fometimes in the ftomach. The round and fhort worms commonly lodge in the *rectum*, or what is called the end gut, and occasion a difagreeable itching about the feat.

The long round worms occafion fqueamifhnefs, vomiting, a difagreeable breath, gripes, loofenefs, fwelling of the belly, fwoonings, loathing of food, and at other times a voracious appetite, a dry cough, convulfions, epileptic fits, and fometimes a privation of fpeech. Thefe worms have been known to perforate the inteffines, and get into the cavity of the belly. The effects of the tapeworm are nearly the fame with those of the long and round, but rather more violent.

Andry fays, the following fymptoms particularly attend the *folium*, which is a fpecies of the tape-worm, viz. fwoonings, privation of fpeech, and a voracious appetite. The round worms called *afcarides*, befides an itching of the *anus*, caufe fwoonings, and tenefmus, or an inclination to go to ftool.

CAUSES. — Worms may proceed for various caufes; but they are feldom found except in weak and relaxed flomachs, where the digeftion is bad. Sedentary perfons are more liable to them than the active and laborious. Those who eat great quantities of unripe fruit, or who live much on raw herbs and roots, are ge-Z 4 nerally nerally fubject to worms. There feems to be an hereditary difposition in fome perfons to this difease. I have often seen all the children of a family subject to worms of a particular kind. They seem likewise frequently to be owing to the nurse. Children of the same family nursed by one woman have often worms, when those nursed by another have none.

SYMPTOMS. ---- The common fymptoms of worms are, palenels of the countenance, and, at other times, an universal flushing of the face; itching of the nose; (this, however, is doubtful, as children pick their nofes in all difeafes;) flarting, and grinding of the teeth in fleep; fwelling of the upper lip; the appetite fometimes bad, at other times quite voracious; loofeness; a four or flinking breath; a hard fwelled belly; great thirft; the urine frothy, and sometimes of a whitish colour; griping, or colic pains; an involuntary difcharge of faliva, especially when afleep; frequent pains of the fide, with a dry cough, and unequal pulle; palpitations of the heart; fwoonings; drowfinefs; cold fweats; palfy; epileptic fits, with many other unaccountable nervous fymptoms, which were formerly attributed to witchcraft, or the influence of evil fpirits. Small bodies in the excrements refembling melon or cucumber feeds are fymptoms of the tapeworm. There is no certain fymptom of worms but paffing them.

I lately faw fome very furprifing effects of worms in a girl about five years of age, who ufed to lie for whole hours as if dead. She at laft expired, and, upon opening her body, a number of the *teres*, or long round worms, were found in her guts, which were confiderably inflamed; and what anatomitts call an *intus fufceptio*, or involving of one part of the gut within another, had taken place in no lefs than four different parts of the inteftinal canal *. MEDI-

• That worms exift in the human body there can be no doubt, and that they mult fometimes be confidered as a difeafe, is equally certain; but this is not the cafe fo often as people imagine. The idea that worms occafion many difeafes, gives an opportunity to the profeffed worm-doctors of impofing on the credulity of mankind, and doing much mifchief. They find worms in every cafe, and liberally throw in their antidotes, which generally confift of ftrong draftie

MEDICINE. — Though numberlefs medicines are extolled for expelling and killing worms *, yet no difeafe more frequently baffles the phyfician's fkill. In general, the most proper medicines for their expulsion are strong purgatives; and to prevent their breeding, stomachic bitters, with now and then a glass of good wine.

The beft purge for an adult, is jalap and calomel. Five-and-twenty or thirty grains of the former, with fix or feven of the latter, mixed in fyrup, may be taken early in the morning for a dofe. It will be proper that the patient keep the houfe all day, and drink nothing cold. The dofe may be repeated once or twice a week, for a fortnight or three weeks. On the intermediate days the patient may take a drachm of the powder of tin, twice or thrice a-day, mixed with fyrup, honey, or treacle.

Those who do not chuse to take calomel, may make use of the bitter purgatives; as aloes, hiera picra, tincture of senna and rhubarb, &c.

Oily medicines are fometimes found beneficial for expelling worms. An ounce of falad oil and a table-fpoonful of common falt may be taken in a glafs of red port wine thrice a-day, or oftener if the ftomach will bear it. But the more common form of ufing oil is in clyfters. Oily clyfters, fweetened with fugar or honey, are very efficacious in bringing away the fhort round worms called *afcarides*, and likewife the *teres*.

The Harrowgate water is an excellent medicine for expelling worms, especially the *afcarides*. As this water is impregnated with fulphur, we may hence infer, that fulphur alone must be a good medicine in this cafe, which is found to be a fact. Many practitioners give flour of fulphur in very large doses, and with great fucces. It flould be made into an electuary with honey or treacle, and taken in fuch quantity as to purge the patient.

draftic purges. I have known these given in delicate conflictutions to the destruction of the patient, where there was not the least fymptom of worms.

* A medical writer of the prefent age has enumerated upwards of fifty British plants, all celebrated for killing and expelling worms.

Where

Where Harrowgate water cannot be obtained, fea-water may be used, which is far from being a contemptible medicine in this case. If fea-water cannot be had, common falt diffolved in water may be drank. I have often feen this used by country nurses with very good effect. Some flour of fulphur may be taken over-night, and the falt-water in the morning.

But worms, though expelled, will foon breed again, if the flomach remain weak and relaxed; to prevent which, we would recommend the Peruvian bark. Half a drachm of bark in powder may be taken in a glafs of red port wine three or four times a-day, after the above medicines have been ufed. Lime-water is likewife good for this purpofe, or a table-fpoonful of the chalybeate wine taken twice or thrice a-day. Infufions or decoctions of bitter herbs may likewife be drank; as the infufion of tanfy, water trefoil, camomile-flowers, tops of wormwood, the leffer centaury, &c.

The above directions are intended for adults; but for children the medicines muft be more agreeable, and in finaller dofes. For a child of four or five years old, fix grains of rhubarb, five of jalap, and two of calomel, may be mixed in a fpoonful of fyrup or honey, and given in the morning. The child fhould keep the houfe all day, and take nothing cold. This dofe may be repeated twice a-week for three or four weeks. On the intermediate days, the child may take a fcruple of powdered tin, and ten grains of æthiops mineral, in a fpoonful of treacle, twice a-day. This dofe muft be increafed or diminifhed according to the age of the patient.

Biffet fays, the great baftard black hellebore, or *bear's* foot, is a most powerful vermifuge for the long round worms. He orders the decoction of about a drachm of the green leaves, or about fifteen grains of the dried leaves in powder, for a dose to a child between four and feven years of age. This dose is to be repeated two or three times. He adds, that the green leaves made into a fyrup with coarfe fugar, is almost the only medicine he has used for round worms for three years past. Before preffing out the juice, he moss the bruised leaves with vinegar, which corrects the medicine. The dose is a tea-spoonful at bed-time, and one or two next morning. I have I have frequently known those big bellies, which in children are commonly reckoned a fign of worms, quite removed by giving them white foap in their pottage, or other food. Tanfy, garlic, and rue, are all good against worms, and may be used various ways. We might here mention many other plants, both for external and internal use, as the cabbage bark, &c. but think the powder of tin with æthiops mineral, and the purges of rhubarb and calomel, are more to be depended on.

Ball's purging vermifuge powder is a very powerful medicine. It is made of equal parts of rhubarb, fcammony, and calomel, with as much double-refined fugar as is equal to the weight of all the other ingredients. Thefe must be well mixed together, and reduced to a fine powder. The dose for a child is from ten grains to twenty, once or twice a-week. An adult may take a drachm for a dose *.

Parents who would preferve their children from worms, ought to allow them plenty of exercife in the open air; to take care that their food be wholefome and fufficiently folid; and, as far as poffible, to prevent their eating raw herbs, roots, or green trafhy fruits. It will not be amifs to allow a child who is fubject to worms, a glafs of red wine after meals; as every thing that braces and ftrengthens the ftomach, is good both for preventing and expelling thefe vermin $\frac{1}{7}$.

In order to prevent any miltake of what I have here faid in favour of *folid* food, it may be proper to observe,

* A powder for the tape-worm refembling this, was long kept a fecret on the Continent; it was lately purchased by the French King, and will be found under the article *Powder*, in the Appendix.

[†] We think it neceffary here to warn people of their danger who buy cakes, powders, and other worm medicines, at random from quacks, and give them to their children without proper care. The principal ingredients in most of these medicines is mercury, which is never to be trifled with. I lately faw a shocking instance of the danger of this conduct. A girl who had taken a dose of wormpowder, bought of a travelling quack, went out, and perhaps was so imprudent as to drink cold water during its operation : she immediately swelled, and died on the following day, with all the fymptoms of having been poisoned.

OF WORMS.

that I only made use of that word in opposition to flops of every kind; not to advise parents to cram their children with meat, two or three times a-day. This fhould only be allowed at dinner and in moderate quantities, or it would create, inftead of preventing worms; for there is no fubftance in nature, which generates fo many worms, as the flesh of animals when in a state of putrefaction. Meat, therefore, at the principal meal, fhould be always accompanied with plenty of good bread, and young, tender, and well-boiled vegetables, efpecially in the fpring, when thefe are poured forth from the bofom of the earth in fuch profusion. They promote the end in view, by keeping the body moderately open, without the aid of artificial phyfic. The ripe fruits of autumn produce the fame effect; and, from their cooling, anti-putrefcent qualities, are as wholefome as the unripe are pernicious.

I also very earnessly conjure parents not to take the alarm at every imaginary fymptom of worms, and directly run for drugs to the quack, or apothecary. They should first try the good effects of proper diet and regimen, and never have recours to medicines till after unequivocal proofs of the nature of the complaint. The danger of advertifed nostrums is fufficiently pointed out and exemplified in the preceding note.

CHAP. XXXVI.

OF THE JAUNDICE.

THIS difeafe is first observable in the white of the eye, which appears yellow. Afterwards the whole skin puts on a yellow appearance. The urine too is of a faffron hue, and dyes a white cloth, if put into it, of the fame colour. There is likewife a species of this difease called the Black Jaundice.

CAUSES.——The immediate caufe of the jaundice is an obftruction of the bile. The remote or occafional caufes are, the bites of poifonous animals, as the viper, mad

OF THE JAUNDICE.

mad dog, &c. the bilious or hyfteric colic; violent paffions, as grief, anger, &c. Strong purges or vomits will likewife occafion the jaundice. Sometimes it proceeds from obftinate agues, or from that difeafe being prematurely ftopped by aftringent medicines. In infants, it is often occafioned by the *meconium* not being fufficiently purged off. Pregnant women are very fubject to it. It is likewife a fymptom in feveral kinds of fever. Catching cold, or the ftoppage of cuftomary evacuations, as the *menfes*, the bleeding piles, iffues, &c. will occafion the jaundice.

SYMPTOMS.——The patient at first complains of excessive wearines, and has great aversion to every kind of motion. His skin is dry, and he generally feels a kind of itching or pricking pain over the whole body. The stools are of a whitish or clay colour, and the urine, as was observed above, is yellow. The breathing is difficult, and the patient complains of an unufual load or oppression on his breast. There is a heat in the nostrils, a bitter taste in the mouth, loathing of food, sickness of the stomach, vomiting, flatulency, and other symptoms of indigestion.

If the patient be young, and the difeafe complicated with no other malady, it is feldom dangerous; but in old people, where it continues long, returns frequently, or is complicated with the dropfy or hypochondriac fymptoms, it generally proves fatal. The black jaundice is more dangerous than the yellow.

REGIMEN.——The diet fhould be cool, light, and diluting, confifting chiefly of ripe fruits and mild vegetables; as apples boiled or roafted, flewed prunes, preferved plums, boiled fpinnage, &c. Veal or chickenbroth, with light bread, are likewife very proper. Many have been cured by living almost wholly for fome days on raw eggs. The drink should be butter-milk, whey fweetened with honey, or decoctions of cool opening vegetables; as marsh-mallow roots, with liquorice, &c.

The patient fhould take as much exercife as he can bear, either on horfeback, or in a carriage; walking, running, and even jumping, are likewife proper, provided vided he can bear them without pain, and there be no fymptoms of inflammation. Patients have been often cured of this difeafe by a long journey, after medicines had proved ineffectual.

Amufements are likewife of great use in the jaundice. The difease is often occasioned by a fedentary life, joined to a dull melancholy disposition. Whatever therefore tends to promote the circulation, and to cheer the spirits, must have a good effect; as dancing, laughing, singing, &c.

MEDICINE. ——If the patient be young, of a full fanguine habit, and complains of pain in the right fide, about the region of the liver, bleeding will be neceffary. After this, a vomit muft be adminifiered; and if the difeafe proves obflinate, it may be repeated once or twice. No medicines are more beneficial in the jaundice than vomits, efpecially where it is not attended with inflammation. Half a drachm of ipecacuanha in powder will be a fufficient dofe for an adult. It may be wrought off with weak camomile-tea, or luke-warm water. The body muft likewife be kept open by taking a fufficient quantity of Caftile foap, or the pills for the jaundice recommended in the Appendix.

Fomenting the parts about the region of the ftomach and liver, and rubbing them with a warm hand or flefhbrufh, are likewife beneficial; but it is ftill more fo for the patient to fit in a bath of warm water up to the breaft. He ought to do this frequently, and fhould continue in it as long as his ftrength will permit.

Many dirty things are recommended for the cure of the jaundice; as lice, millepedes, &c. But thefe do more harm than good, as people truft to them, and neglect more valuable medicines; befides, they are feldom taken in fufficient quantity to produce any effects. People always expect that fuch *things* fhould act as charms, and confequently feldom perfift in the ufe of them. Vomits, purges, fomentations, and exercife, will feldom fail to cure the jaundice when it is a fimple diteafe; and when complicated with the dropfy, a fchirrous liver, or other chronic complaints, it is hardly to be cured by any means.

Numberlefs Britifh herbs are extolled for the cure of this difeafe. The author of the Medicina Britannica mentions near a hundred, all famous for curing the jaundice. The fact is, the difeafe often goes off of its own accord; in which cafe, the laft medicine is always faid to have performed the cure. I have fometimes, however, feen confiderable benefit, in a very obflinate jaundice, from a decoction of hempfeed. Four ounces of the feed may be boiled in two Englifh quarts of ale, and fweetened with coarfe fugar. The dofe is half an Englifh pint every morning. It may be continued for eight or nine days.

I have likewife known Harrowgate fulphur water cure jaundice of very long ftanding. It fhould be used for fome weeks, and the patient must both drink and bathe.

The foluble tartar is a very proper medicine in the jaundice. A drachm of it may be taken every night and morning in a cup of tea or water gruel. If it does not open the body, the dofe may be increased.

Perfons fubject to the jaundice ought to take as much exercise as possible, and to avoid all heating and aftringent aliments.

The two laft directions are of far greater importance than fome people may imagine. In fact, taking exercife, and keeping the body open, are the only affured and rational means of removing a complaint which generally arifes from an obftruction of the biliary ducts. I knew a celebrated phyfician who was fubject to this difeafe, and who, whenever it attacked him, mounted his horfe, fet out on a journey, and never returned till he was well. For my own part, I fhould place more reliance on the efficacy of fuch a method, than on the whole catalogue of near a hundred fpecifics mentioned by the late Doctor SHORT of Sheffield, though I have known inftances, where one of them, the decoction of hempfeed, as already intimated, was found very beneficial.

[352]

CHAP. XXXVII.

OF THE DROPSY.

THE dropfy is a preternatural fwelling of the whole body, or fome part of it, occafioned by a collection of watery humour. It is diffinguished by different names, according to the part affected, as the *anafarca*, or a collection of water under the skin; the *afcites*, or a collection of water in the belly; the *bydrops pectoris*, or dropfy of the breaft; the *bydrocephalus*, or dropfy of the brain, &c.

CAUSES. — The dropfy is often owing to an hereditary difpofition. It may likewife proceed from drinking ardent fpirits, or other flrong liquors. It is true, almost to a proverb, that great drinkers die of a dropfy. The want of exercise is also a very common caufe of the dropfy. Hence it is justly reckoned among the difeafes of the fedentary. It often proceeds from exceflive evacuations, as frequent and copious bleedings, flrong purges often repeated, frequent falivations, &c. The fudden floppage of customary or neceflary evacuations, as the *menfes*, the hæmorrhoids, fluxes of the belly, &c. may likewife caufe a dropfy.

I have known the dropfy occafioned by drinking large quantities of cold, weak, watery liquor, when the body was heated by violent exercife. A low, damp, or marfhy fituation is likewife a frequent caufe of it. Hence it is a common difeafe in moift, flat, fenny countries. It may alfo be brought on by a long ufe of poor watery diet, or of vifcous aliment that is hard of digeftion. It is often the effect of other difeafes, as the jaundice, a fchirrus of the liver, a violent ague of long continuance, a diarrhœa, a dyfentery, an empyema, or a confumption of the lungs. In fhort, whatever obftructs the perfpiration, or prevents the blood from being duly prepared, may occafion a dropfy.

SYMPTOMS.——The anafarca generally begins with a fwelling of the feet and ancles towards night, 6 which

OF THE DROPSY.

which for fome time difappears in the morning. In the evening the parts, if preffed with the finger, will pit. The fwelling gradually afcends, and occupies the trunk of the body, the arms, and the head. Afterwards the breathing becomes difficult, the urine is in fmall quantity, and the thirft great; the body is bound, and the perfpiration is greatly obftructed. To thefe fucceed torpor, heavinefs, a flow wafting fever, and a troublefome cough. This laft is generally a fatal fymptom, as it fhews that the lungs are affected.

In an *afcites*, befides the above fymptoms, there is a fwelling of the belly, and often a fluctuation, which may be perceived by ftriking the belly on one fide, and laying the palm of the hand on the oppofite. This may be diftinguished from a *tympany* by the weight of the fwelling, as well as by the fluctuation. When the *anafarca* and *afcites* are combined, the cafe is very dangerous. Even a fimple *afcites* feldom admits of a radical cure. Almost all that can be done is, to let off the water by tapping, which feldom affords more than a temporary relief.

When the difeafe comes fuddenly on, and the patient is young and ftrong, there is reafon to hope for a cure, efpecially if medicine be given early. But if the patient be old, has led an irregular or a fedentary life, or if there be reafon to fulpect that the liver, lungs, or any of the vifcera are unfound, there is great reafon to fear that the confequences will prove fatal.

REGIMEN.——The patient must abstain, as much as possible, from all drink, especially weak and watery liquors, and must quench his thirst with mustard-whey, or acids, as juice of lemons, oranges, forrel, or fuch like. His aliment ought to be dry, of a stimulating and diuretic quality, as toasted bread, the flesh of birds, or other wild animals, roasted; pungent and aromatic vegetables, as garlic, mustard, onions, cresses, horfe-radish, rocambole, shalot, &c. He may also eat fea-biscuit dipped in wine or a little brandy. This is not only nourishing, but tends to quench thirst. Some have been actually cured of a dropsy by a total abstinence from all liquids, and living entirely upon fuch things as are mentioned above.

Aa

If

If the patient must have drink, the Spa-water, or Rhenish wine, with diuretic medicines infused in it, are the beft.

Exercise is of the greatest importance in a dropsy. If the patient be able to walk, dig, or the like, he ought to continue these exercises as long as he can. If he be not able to walk or labour, he must ride on horseback, or in a carriage, and the more violent the motion fo much the better, provided he can bear it. His bed ought to be hard, and the air of his apartments warm and dry. If he live in a damp country, he ought to be removed into a dry one, and, if possible, into a warmer climate. In a word, every method fhould be taken to promote the perspiration, and to brace the folids. For this purpose it will likewife be proper to rub the patient's body, two or three times a-day with a hard cloth, or the flefhbrush; and he ought constantly to wear flannel next his fkin.

MEDICINE .---- If the patient be young, his conflitution good, and the difeafe has come on fuddenly, it may generally be removed by ftrong vomits, brick purges, and fuch medicines as promote a difcharge by fweat and urine. For an adult, half a drachm of ipecacuanha in powder, and half an ounce of oxymel of fquills, will be a proper vomit. This may be repeated as often as is found neceffary, three or four days intervening between the dofes. The patient must not drink much after taking the vomit, otherwife he deftroys its effect. A cup or two of camomile-tea will be fufficient to work it off.

Between each vomit, on one of the intermediate days, the patient may take the following purge: Jalap in powder half a drachm, cream of tartar two drachms, calomel fix grains. These may be made into a bolus with a little fyrup of pale roles, and taken early in the morning. The lefs the patient drinks after it the better. If he be much griped, he may now and then take a cup of chicken-broth.

The patient may likewife take every night at bed-time the following bolus: To four or five grains of camphor, add one grain of opium, and as much fyrup of orangepeel

EA

peel as is fufficient to make them into a bolus. This will generally promote a gentle fweat, which fhould be encouraged by drinking now and then a fmall cup of wine-whey, with a tea-spoonful of the spirits of hartshorn in it. A tea-cupful of the following diuretic infufion may likewife be taken every four or five hours through the day : Take juniper-berries, mustard-feed, and horferadish, of each half an ounce, ashes of broom half a pound; infuse them in a quart of Rhenish wine or ftrong ale for a few days, and afterwards ftrain off the liquor. Such as cannot take this infusion, may use the decoction of feneka-root, which is both diuretic and sedorific. I have known an obstinate anafarca cured by an infusion of the ashes of broom in wine.

The above courfe will often cure an incidental dropfy, if the conftitution be good; but when the difeafe proceeds from a bad habit, or an unfound state of the vifcera, ftrong purges and vomits are not to be ventured upon. In this cafe, the fafer courfe is to palliate the fymptoms by the use of such medicines as promote the fecretions, and to support the patient's strength by warm and nourifhing cordials.

The fecretion of urine may be greatly promoted by nitre. Brookes fays, he knew a young woman who was cured of a dropfy by taking a drachm of nitre every morning in a draught of ale, after she had been given over as incurable. The powder of fquills is likewife a good diuretic. Six or eight grains of it, with a fcruple of nitre, may be given twice a day in a glafs of ftrong cinnamon-water. Ball fays, a large spoonful of unbruifed multard-feed taken every night and morning, and drinking half an English pint of the decoction of the tops of green broom after it, has performed a cure after other powerful medicines had proved ineffectual.

I have fometimes feen good effects from cream of tartar in this difeafe. It promotes the difcharges by ftool and urine, and will at least palliate, if it does not perform a cure. The patient may begin by taking an ounce every fecond or third day, and may increase the quantity to two or even to three ounces, if the ftomach will bear it. Aa2

This

This quantity is not, however, to be taken at once, but divided into three or four doles.

To promote perfpiration, the patient may use the decoction of feneka root, as directed above; or he may take two table-fpoonfuls of Mindererus's spirit in a cup of wine-whey three or four times a-day. To promote a discharge of urine, the following infusion of the London hospitals will likewise be beneficial:

Take of zedoary-root two drachms; dried fquills, rhubarb, and juniper-berries bruifed, of each a drachm; cinnamon in powder, three drachms; falt of wormwood, a drachm and a half; infufe in an English pint and a half of old hock wine, and, when fit for use, filter the liquor. A wine glass of it may be taken three or four times a-day.

In the *anafarca* it is ufual to fcarify the feet and legs. By this means the water is often difcharged; but the operator muft be cautious not to make the incifions too deep; they ought barely to pierce through the fkin, and efpecial care muft be taken, by fpirituous fomentations and proper digeftives, to prevent a gangrene.

In an *afcites*, when the difeafe does not evidently and fpeedily give way to purgative and diuretic medicines, the water ought to be let off by tapping. This is a very fimple and fafe operation, and would often fucceed, if it were performed in due time; but if it be delayed till the humours are vitiated, or the bowels fpoiled, by long foaking in water, it can hardly be expected that any permanent relief will be procured *.

After the evacuation of the water, the patient is to be put on a courfe of strengthening medicines; as the Peruvian bark; the elixir of vitriol; warm aromatics, with a due proportion of rhubarb, infused in wine, and such like. His diet ought to be dry and nourishing, such as is recommended in the beginning of the Chapter; and he should take as much exercise as he can bear without

• The very name of an operation is dreadful to most people, and they wish to try every thing before they have recourse to it. This is the reason why tapping to feldom fucceeds to our wish. I had a patient who was regularly tapped once a month for feveral years, and who used to eat her dinner as well after the operation as if nothing had happened. She died at last rather worn out by age than by the difease. fatigue.

OF THE DROPSY.

fatigue. He should wear flannel, or rather fleecy hosiery, next his skin, and make daily use of the flesh-brush.

We have claffical authority of two thousand years ftanding to fay, that the dropfy is an obstinate disease; not that the *incidental* dropfy is incurable, for I have often been so happy as to succeed in the treatment of it; but when the dropfy is accompanied with a schirrous liver, or a worn-out constitution, very little is to be expected from medicine.

I had lately a fingular inftance of the efficacy of nitre in a cafe of dropfy. A young man, a cornet of dragoons, was dropfical all over, even his face not excepted. After feveral things had been tried without fuccefs, I fhewed him my quotation from Dr. BROOKES, in this Chapter. He was defirous of making an experiment with nitre; and took a drachm of it in a cup of warm ale, for fome time once, and afterwards twice a-day, till he was cured.

CHAP. XXXVIII.

OF THE GOUT.

THERE is no difease which shews the imperfection of medicine, or sets the advantages of temperance and exercise in a stronger light, than the gout. Excess and idleness are the true sources from whence it originally sprung, and all who would avoid it must be active and temperate.

Though idleness and intemperance are the principal causes of the gout, yet many other things may contribute to bring on the diforder in those who are not, and to induce a paroxysm in those who are subject to it; as intense study; excess of venery; too free an use of acidulated liquors; night-watching; grief or uneasiness of mind; an obstruction or defect of any of the customary discharges, as the menses, sweating of the feet, perspiration, &c.

Aa 3

ipirits,

fpirits, and has often a pain in the limbs, with a fenfation as if wind or cold water were paffing down the thigh. The appetite is often remarkably keen a day or two before the fit, and there is a flight pain in paffing urine, and frequently an involuntary fhedding of tears. Sometimes thefe fymptoms are much more violent, efpecially upon the near approach of the fit; and it has been observed, that as is the fever which ufhers in the gout, fo will the fit be; if the fever be fhort and fharp, the fit will be fo likewife; if it be feeble, long, and lingering, the fit will be fuch alfo. But this observation can only hold with respect to very regular fits of the gout.

The regular gout generally makes its attack in the fpring or beginning of winter, in the following manner: About two or three in the morning, the patient is feized with a pain in his great toe, fometimes in the heel, and at other times in the ancle or calf of the leg. This pain is accompanied with a fenfation as if cold water were poured upon the part, which is fucceeded by a fhivering, with fome degree of fever. Afterwards the pain increases, and fixing among the fmall bones of the foot, the patient feels all the different kinds of torture, as if the part were ftretched, burnt, fqueezed, gnawed, or torn in pieces. The part at length becomes to exquisitely fensible, that the patient cannot bear to have it touched, or even fuffer any perfon to walk acrofs the room.

The patient is generally in exquifite torture for twentyfour hours, from the time of the coming on of the fit : he then becomes eafier, the part begins to fwell, appears red, and is covered with a little moifture. Towards morning he drops afleep, and generally falls into a gentle breathing fweat. This terminates the first paroxyfm, a number of which constitutes a fit of the gout; which is longer or shorter according to the patient's age, strength, the feason of the year, and the disposition of the body to this difease.

The patient is always worfe towards night, and eafier in the morning. The paroxyfms, however, generally grow milder every day, till at length the difeafe is carried off by perfpiration, urine, and the other evacuations. In fome patients this happens in a few days; in others, it requires

requires weeks, and in fome, months to finish the fit. Those whom age and frequent fits of the gout have greatly debilitated, feldom get free from it before the approach of fummer, and sometimes not till it be pretty far advanced.

REGIMEN.——As there are no medicines yet known that will cure the gout, we fhall confine our obfervations chiefly to regimen, both in and out of the fit.

In the fit, if the patient be young and ftrong, his diet ought to be thin and cooling. and his drink of a diluting nature; but where the conffitution is weak, and the patient has been accuftomed to live high, this is not a proper time to retrench. In this cafe he muft keep nearly to his ufual diet, and fhould take frequently a cup of ftrong negus, or a glafs of generous wine. Wine-whey is a very proper drink in this cafe, as it promotes the perfpiration without greatly heating the patient. It will anfwer this purpofe better if a tea-fpoonful of *fal volatile oleofum*, or fpirits of hartfhorn, be put into a cup of it twice a-day. It will likewife be proper to give at bedtime a tea-fpoonful of the volatile tincture of *guaiacum*, in a large draught of warm wine-whey. This will greatly promote perfpiration through the night.

As the most fafe and efficacious method of discharging the gouty matter is by perfpiration, this ought to be kept up by all means, especially in the affected part. For this purpose the leg and foot should be wrapt in foft flannel, fur, or wool. The last is most readily obtained, and feems to answer the purpose better than any thing elfe. The people of Lancashire look upon wool as a kind-of fpecific in the gout. They wrap a great quantity of it about the leg and foot affected, and cover it with a fkin of foft dreffed leather. This they fuffer to continue for eight or ten days, and fometimes for a fortnight or three weeks, or longer, if the pain does not ceafe. I never knew any external application answer fo well in the gout. I have often feen it applied when the fwelling and inflammation were very great, with violent pain, and have found all thefe fymptoms relieved by it in a few days. The wool which they use is generally greafed, and carded or combed. They choose the fosteft Aa4 which

which can be had, and feldom or never remove it till the fit be entirely gone off.

The patient ought likewife to be kept quiet and eafy during the fit. Every thing that affects the mind difturbs the paroxyim, and tends to throw the gout upon the nobler parts. All external applications that repel the matter are to be avoided as death. They do not cure the difease, but remove it from a safer to a more dangerous part of the body, where it often proves fatal, A fit of the gout is to be confidered as Nature's method of removing fomething that might prove deftructive to the body, and all that we can do with fafety, is to promote her intentions, and to affift her in expelling the enemy in her own way. Evacuations by bleeding, ftool, &c. are likewife to be used with caution, they do not remove the caule of the difeafe, and fometimes by weakening the patient prolong the fit; but where the conftitution is able to bear it, it will be of use to keep the body gently open by diet, or very mild laxative medicines.

Many things will indeed fhorten a fit of the gout, and fome will drive it off altogether; but nothing has yet been found which will do this with fafety to the patient. In pain we eagerly grafp at any thing that promifes immediate eafe, and even hazard life itlelf for a temporary relief. This is the true reafon why fo many infallible remedies have been propoled for the gout, and why fuch numbers have loft their lives by the use of them. Notwithftanding the acknowledged and frequently experienced danger of tampering with the gout, fuch is the effect of intenfe pain, that I never met with more than two patients who could bear their fufferings with rational composure, or, what is the fame thing, without frantic attempts to alleviate them. When the feat of the complaint is in torture, a promife to afford relief, though made by the greatest impostor upon earth, is listened to; and prefent eafe is fought for, at the rifk of any future confequences. It is not many years fince fome perfons of the first rank in the kingdom fell victims to the deceptions of a foreign quack, who foothed their impatience of pain, amufed them with the charm of fancied recovery, and rendered momentary eafe the fatal prelude to

to inevitable death. It would be as prudent to ftop the fmall-pox from rifing, and to drive them into the blood, as to attempt to repel the gouty matter after it has been thrown upon the extremities. The latter is as much an effort of Nature to free herfelf from an offending caufe as the former, and ought equally to be promoted.

When the pain, however, is very great, and the patient is reftlefs, thirty or forty drops of laudanum, more or lefs, according to the violence of the fymptoms, may be taken at bed-time. This will eafe the pain, procure reft, promote perfpiration, and forward the crifis of the difeafe.

After the fit is over, the patient ought to take a gentle dofe or two of the bitter tincture of rhubarb, or fome other warm flomachic purge. He flould alfo drink a weak infufion of flomachic bitters in fmall wine or ale, as the Peruvian bark, with cinnamon, Virginian fnakeroot, and orange-peel. The diet at this time flould be light, but nourifhing, and gentle exercise ought to be taken on horfeback, or in a carriage,

Out of the fit, it is in the patient's power to do many things towards preventing a return of the diforder, or rendering the fit, if it fhould return, lefs fevere. This, however, is not to be attempted by medicine. I have frequently known the gout kept off for feveral years by the Peruvian bark and other aftringent medicines; but in all the cafes where I had occafion to fee this tried, the perfons died fuddenly, and to all appearance for want of a regular fit of the gout. One would be apt, from hence, to conclude, that a fit of the gout, to fome conflitutions, in the decline of life, is rather falutary than hurtful,

Though it may be dangerous to ftop a fit of the gout by medicine, yet if the conftitution can be fo changed by diet and exercife, as to leffen or totally prevent its return, there certainly can be no danger in following fuch a courfe. It is well known that the whole habit may be fo altered by a proper regimen, as quite to eradicate this difeafe; and those only who have fufficient resolution to perfift in fuch a courfe have reason to expect a cure. The courfe which we would recommend for preventing the gout, is as follows: In the firft place, univerfal temperance. In the next place, fufficient exercise*. By this we do not mean fauntering about in an indolent manner, but labour, fweat, and toil. Thefe only can render the humours wholefome, and keep them fo. Going early to bed, and rifing betimes, are alfo of great importance. It is likewife proper to avoid night fludies, and intenfe thinking. The fupper fhould be light and taken early. The ufe of milk, gradually increased, till it becomes the principal part of diet, is particularly recommended. All ftrong liquors, especially generous wines and four punch, are to be avoided.

We would likewife recommend fome dofes of magnefia alba and rhubarb to be taken every fpring and autumn; and afterwards a courfe of flomachic bitters, as tanfey or water-trefoil tea, an infufion of gentian and camomile flowers, or a decoction of burdock-root, &c. Any of thefe, or an infufion of any wholefome bitter that is more agreeable to the patient, may be drank for two or three weeks in March and October twice a-day. An iffue or perpetual blifter has a great tendency to prevent the gout. If thefe were more generally ufed in the decline of life, they would not only often prevent the gout, but alfo other chronic maladies. Such as can afford to go to Bath, will find great benefit from bathing and drinking the water. It both promotes digeftion, and invigorates the habit.

Though there is little room for medicine during a regular fit of the gout, yet when it leaves the extremities, and falls on fome of the internal parts, proper applications to recal and fix it become abfolutely neceffary. When the gout affects the head, the pain of the joints ceafes, and the fwelling difappears, while either fevere head-ach, drowfinefs, trembling, giddinefs, convulfions, or delirium come on. When it feizes the lungs, great oppreffion, with cough and difficulty of breathing, enfue. If it attacks the ftomach, extreme ficknefs, vomiting, anxi-

* Some make a fecret of curing the gout by muscular exercise. This fecret, however, is as old as Celfus, who flrongly recommends that mode of cure; and whoever will submit to it in the fullest extent, may expect to reap folid and permanent advantages. ety, pain in the epigaftric region, and total loss of ftrength, will fucceed.

When the gout attacks the head or lungs, every method must be taken to fix it in the feet. They must be frequently bathed in warm water, and acrid cataplasms applied to the foles. Blistering-plasters ought likewife to be applied to the ancles or calves of the legs. Bleeding in the feet or ancles is also necessary, and warm stomachic purges. The patient ought to keep in bed for the most part, if there be any figns of inflammation, and should be very careful not to catch cold.

If it attack the ftomach, with a fenfe of cold, the moft warm cordials are neceffary; as ftrong wine boiled up with cinnamon or other fpices; cinnamon-water; peppermint-water; and even brandy or rum *. The patient fhould keep his bed, and endeavour to promote a fweat by drinking warm liquors; and if he fhould be troubled with a naufea, or inclination to vomit, he may drink camomile-tea, or any thing that will make him vomit freely.

When the gout attacks the kidneys, and imitates gravel-pains, the patient ought to drink freely of a decoction of marfh-mallows, and to have the parts fomented with warm water. An emollient clyfter ought likewife to be given, and afterwards an opiate. If the pain be very violent, twenty or thirty drops of laudanum may be taken in a cup of the decoction.

Perfons who have had the gout fhould be very attentive to any complaints that may happen to them about the time when they have reafon to expect a return of the fit. The gout imitates many other diforders, and by being miltaken for them, and treated improperly, is often diverted from its regular course, to the great danger of the patient's life.

Those who never had the gout, but who, from their conftitution or manner of living, have reason to expect it, ought likewise to be very circumspect with regard to its first approach. If the disease, by wrong conduct or improper medicines, be diverted from its proper course,

* Æther is found to be an efficacious remedy in this cafe.

OF THE GOUT.

the miferable patient has a chance to be ever after tormented with head-achs, coughs, pains of the ftomach and inteftines; and to fall at last a victim to its attack upon some of the more noble parts.

OF THE RHEUMATISM.

This difeafe has often a refemblance to the gout. It generally attacks the joints with exquisite pain, and is fometimes attended with inflammation and swelling. It is most common in the spring, and towards the end of autumn. It is usually distinguished into acute and chronic; or the rheumatism with and without a fever.

CAUSES.——The caufes of a rheumatifm are frequently the fame as those of an inflammatory fever, viz. an obstructed perspiration, the immoderate use of strong liquors, and the like. Sudden changes of the weather, and all quick transitions from heat to cold, are very apt to occasion the rheumatism. The most extraordinary case of a rheumatism that I ever faw, where almost every joint of the body was distorted, was a man who used to work one part of the day by the fire, and the other part of it in the water. Very obstinate rheumatisms have likewise been brought on by persons not accustomed to it, allowing their feet to continue long wet. The same effects are often produced by wet clothes, damp beds, fitting or lying on the damp ground, travelling in the night, &c.

The rheumatism may likewise be occasioned by exceffive evacuations, or the stoppage of customary difcharges. It is often the effect of chronic diseases, which vitiate the humours; as the scurvy, the *lues venerea*, obstinate autumnal agues, &c.

The rheumatism prevails in cold, damp, marshy countries. It is most common among the poorer fort of peasants, who are ill-clothed, live in low damp houses, and eat coarse and unwholes food, which contains but little nourishment, and is not easily digested.

SYMPTOMS. — The acute rheumatifm commonly begins with wearinefs, fhivering, a quick pulfe, reftleffnefs, thirft, and other fymptoms of fever. Afterwards the patient complains of flying pains, which are increased by

by the least motion. These at length fix in the joints, which are often affected with swelling and inflammation. If blood be let in this difease, it has generally the same appearance as in the pleurify.

In this kind of rheumatifm the treatment of the patient is nearly the fame as in an acute or inflammatory fever. If he be young and ftrong, bleeding is neceffary, which may be repeated according to the exigencies of the cafe. The body ought likewife to be kept open by emollient clyfters, or cool opening liquors; as decoctions of tamarinds, cream of tartar, whey, fenna-tea, and the like. The diet should be light, and in small quantity, confifting chiefly of roafted apples, groat-gruel, or weak chicken-broth. After the feverilh fymptoms have abated, if the pain still continues, the patient must keep his bed, and take fuch things as promote perspiration; as wine-whey, with spiritus Mindereri, &c. He may likewife take, for a few nights, at bed-time, in a cup of wine-whey, a drachm of the cream of tartar, and half a drachm of gum guaiacum in powder.

Warm bathing, after proper evacuations, has often an exceeding good effect. The patient may either be put into a bath of warm water, or have cloths wrung out of it applied to the parts affected. Great care must be taken that he do not catch cold after bathing.

The chronic rheumatism is feldom attended with any confiderable degree of fever, and is generally confined to fome particular part of the body, as the shoulders, the back, or the loins. There is feldom any inflammation or swelling in this case. Perfons in the decline of life are most subject to the chronic rheumatism. In such patients it often proves extremely obstinate and sometimes incurable.

In this kind of rheumatifm the regimen fhould be nearly the fame as in the acute. Cool and diluting diet, confifting chiefly of vegetable fubltances, as flewed prunes, coddled apples, currants or goofe-berries boiled in milk, is most proper. Arbuthnot fays, "If there be a fpecific in aliment for the rheumatifm, it is certainly whey;" and adds, "That he knew a perfon fubject to this difeafe, who could never be cured by any other method but a diet

OF THE RHEUMATISM.

a diet of whey and bread." He likewife fays, "That cream of tartar in water-gruel, taken for feveral days, will eafe rheumatic pains confiderably." This I have often expetienced, but found it always more efficacious when joined with gum guaiacum, as already directed. In this cafe the patient may take the dofe formerly mentioned twice a-day, and likewife a tea-fpoonful of the volatile tincture of gum guaiacum at bed-time in wine-whey.

This courfe may be continued for a week, or longer, if the cafe proves obstinate, and the patient's strength will permit. It ought then to be omitted for a few days, and repeated again. At the fame time leeches or a bliftering-plafter may be applied to the part affected. What I have generally found answer better than either of thefe, in obstinate fixed rheumatic pains, is the warm plaster *. I have likewife known a plaster of Burgundy pitch worn for fome time on the part affected give great relief in rheumatic pains. My ingenious friend, Dr. Alexander of Edinburgh, fays, he has frequently cured very obffinate rheumatic pains by rubbing the parts affected with tincture of cantharides. When the common tincture did not fucceed, he ufed it of a double or treble ftrength. Cupping upon the part affected is likewife often very beneficial, and fo is the application of leeches.

Though this difeafe may not feem to yield to medicines for fome time, yet they ought ftill to be perfifted in. Perfons who are fubject to frequent returns of the rheumatikm, will often find their account in ufing medicines, whether they be immediately affected with the difeafe or not. The chronic rheumatifm is fimilar to the gout in this refpect, that the most proper time for ufing medicines to extirpate it, is when the patient is most free from the diforder.

To those who can afford the expence, I would recommend the warm baths of Buxton or Matlock in Derbyfhire. These have often, to my knowledge, cured very obstinate rheumatisms, and are always fase either in or out of the fit. When the rheumatism is complicated with scorbutic complaints, which is not feldom the case,

* See Appendix, Warm Plaster.

366

13/5 5.

OF THE RHEUMATISM.

361

One

the Harrowgate waters, and those of Moffat, are proper. They should both be drank and used as a warm bath.

There are feveral of our own domeftic plants which may be used with advantage in the rheumatifm. One of the best is the white mustard. A table-spoonful of the feed of this plant may be taken twice or thrice a-day, in a glafs of water or fmall wine. The water trefoil is likewife of great use in this complaint. It may be infused in wine or ale, or drank in form of tea. The groundivy, camomile, and feveral other bitters, are also beneficial, and may be used in the fame manner. No benefit, however, is to be expected from thefe, unlefs they be taken for a confiderable time. Excellent medicines are often despised in this difease, because they do not perform an immediate cure; whereas nothing would be more certain than their effect were they duly perfifted in. Want of perfeverance in the ufe of medicines is one reafon why chronic difeafes are fo feldom cured.

Cold bathing, especially in falt water, often cures the rheumatism. We would also recommend exercise and wearing flannel next the skin. Issues are likewise very proper, especially in chronic cases. If the pain affects the shoulders, an issue may be made in the arm; but if it affects the loins, it should be put into the leg or thigh.

Perfons afflicted with the feurvy are very fubject to rheumatic complaints. The beft medicines in this cafe are bitters and mild purgatives. Thefe may either be taken feparately or together, as the patient inclines. An ounce of Peruvian bark, and half an ounce of rhubarb in powder, may be infufed in a bottle of wine; and one, two, or three wine glaffes of it taken daily, as fhall be found neceffary for keeping the body gently open. In cafes where the bark itfelf proves fufficiently purgative, the rhubarb may be omitted.

Such as are fubject to frequent attacks of the rheumatilm ought to make choice of a dry, warm fituation, to avoid the night-air, wet clothes, and wet feet, as much as poffible. Their clothing fhould be warm, and they fhould wear flaunel next their fkin, and make frequent ule of the flefh-brufh.

OF THE RHEUMATISM.

One of the best articles of drefs, not only for the prevention of the rheumatifm, but for powerful co-operation in its cure, is fleecy hofiery. A medical friend of mine, of long experience and much practice in the ifle of Ely, affured me, that the introduction of that manufacture had prevented more rheumatifms, colds, and agues, than all the medicines which had ever been used there. Such of the inhabitants of marshy countries as are in easy circumftances could not, perhaps, direct their charity and humanity to a better object, than to the fupplying of their poor neighbours with fo cheap and fimple a prefervative. I have even myfelf experienced the good effects of fuch warm covering in the rheumatifm, to which I was very fubject about thirty years ago ; but have never experienced any attack of it fince I took to warm clothing, and particularly fleecy hofiery worn next the fkin.

CHAP. XXXIX.

OF THE SCURVY.

THIS difeafe prevails chiefly in cold northern countries, efpecially in low damp fituations, near large marfhes, or great quantities of ftagnating water. Sedentary people, of a dull melancholy difpofition, are most fubject to it. It proves often fatal to failors on long voyages, particularly in ships that are not properly ventilated, have many people on board, or where cleanliness is neglected.

It is not neceffary to mention the different fpecies into which this difeafe has been divided, as they differ from one another chiefly in degree. What is called the *land fcurvy*, however, is feldom attended with those highly putrid fymptoms which appear in patients who have been long at fea, and which, we prefume, are rather owing to confined air, want of exercise, and the unwholefome food eaten by failors on long voyages, than to any specific difference in the difease.

CAUSES.

CAUSES. ——The fcurvy is occafioned by cold moift air; by the long use of falted or fmoke-dried provisions, or any kind of food that is hard of digestion, and affords little nourishment. It may also proceed from the suppression of customary evacuations; as the *menses*, the hæmorrhoidal flux, &c. It is fometimes owing to an hereditary taint, in which case a very small cause will excite the latent diforder. Grief, fear, and other depresfing passions, have a great tendency both to excite and aggravate this difease. The same observation holds with regard to neglect of cleanlines; bad clothing; the want of proper exercise; confined air; unwholesome food; or any difease which greatly weakens the body, or vitiates the humours.

SYMPTOMS .---- This difease may be known by unufual wearinefs, heavinefs, and difficulty of breathing, efpecially after motion; rottennels of the gums, which are apt to bleed on the flighteft touch; a flinking breath, frequent bleeding at the nofe; crackling of the joints; difficulty of walking; fometimes a fwelling and fometimes a falling away of the legs, on which there are livid, yellow, or violet-coloured fpots; the face is generally of a pale or leaden colour. As the difeafe advances, other fymptoms come on ; as rottennels of the teeth, hæmorrhages, or difcharges of blood from different parts of the body, foul obstinate ulcers, pains in various parts, efpecially about the breaft, dry fealy eruptions all over the body, &c. At last a wasting or hectic fever comes on, and the miferable patient is often carried off by a dylentery, a diarrhœa, a dropfy, the palfy, fainting fits, or a mortification of fome of the bowels.

CURE. — We know no way of curing this difeafe but by purfuing a plan directly opposite to that which brings it on. It proceeds from a vitiated state of the humours, occasioned by errors in diet, air, or exercise; and this cannot be removed but by a proper attention to these important articles.

If the patient has been obliged to breathe a cold, damp, or confined air, he fhould be removed, as foon as possible, to a dry, open, and moderately warm one. If there is reafon to believe that the difease proceeds from a fedentary

life,

life, or depreffing paffions, as grief, fear, &c. the patient must take daily as much exercise in the open air as he can bear, and his mind should be diverted by cheerful company and other amusements. Nothing has a greater tendency either to prevent or remove this difease, than constant cheerfulness and good humour. But this, alas! is feldom the lot of perfons afflicted with the fcurvy; they are generally furly, peevish, and morofe.

When the fcurvy has been brought on by a long ufe of falted provisions, the proper medicine is a diet confifting chiefly of frefh vegetables; as oranges, apples, lemons, limes, tamarinds, water-creffes, fcurvy-grafs, brook-lime, &c. The ufe of thefe, with milk, pot-herbs, new bread, and frefh beer or cyder, will feldom fail to remove a fcurvy of this kind, if taken before it be too far advanced; but to have this effect, they muft be perfifted in for a confiderable time. When frefh vegetables cannot be obtained, pickled or preferved ones may be ufed; and where thefe are wanting, recourfe muft be had to the chymical acids. All the patient's food and drink fhould in this cafe be fharpened with cream of tartar, elixir of vitriol, vinegar, or the fpirit of fea-falt.

Thefe things, however, will more certainly prevent than cure the fcurvy, for which reafon fea-faring people, especially in long voyages, ought to lay in plenty of them. Cabbages, onions, goofe-berries, and many other vegetables, may be kept a long time by pickling, preferving, &c. and when thefe fail, the chymical acids recommended above, which will keep for any length of time, may be used. We have reafon to believe, if ships were well ventilated, had got ftore of fruits, greens, cyder, &c. laid in, and if proper regard were paid to cleanlinefs and warmth, that failors would be the most healthy people in the world, and would feldom fuffer either from the fcurvy or putrid fevers, which are fo fatal to that useful class of men : but it is too much the temper of fuch people to defpife all precaution; they will not think of any calamity till it overtakes them, when it is too late to ward off the blow.

It must indeed be owned, that many of them have it not in their power to make the provision we are speaking

OF THE SCURVY.

ing of; but in this cafe it is the duty of their employers to make it for them; and no man ought to engage in a long voyage without having thefe articles fecured *.

I have often feen very extraordinary effects in the landfcurvy from a milk-diet. This preparation of Nature is a mixture of animal and vegetable properties, which of all others is the most fit for restoring a decayed constitution, and removing that particular acrimony of the humours, which seems to constitute the very effence of the fcurvy, and many other diseases. But people despise this wholesome and nouriss field and fermented liquors, while milk is only deemed fit for their hogs.

The most proper drink in the scurvy is whey or buttermilk. When these cannot be had, found cyder, perry, or spruce beer, may be used. Wort has likewise been found to be a proper drink in the scurvy, and may be used at sea, as malt will keep during the longest voyage. A decostion of the tops of the spruce for is likewise proper. It may be drank in the quantity of an English pint twice a-day. Tar-water may be used for the same purpose, or decostions of any of the mild mucilaginous vegetables; as farfaparilla, marshmallow roots, &c. Infusions of the bitter plants, as ground-ivy, the lesser centaury, marshtrefoil, &c. are likewise beneficial. I have seen the pea-

* Every body knows how much eafier it is to prevent than to cure any difeafe; but this is particularly true with refpect to the fcurvy. I have therefore recommended the most affured means of preferving our feamen from its formidable attacks. Vegetable and mineral acids are certainly the belt correctives of the acrimony occationed by the long ufe of falted provisions. These are one of the chief caufes of the feurvy not only at fea, but on land alfo, where falted and fmoke-dried meats are a favourite and very cuftomary article of food. It appears from the reports of modern travellers, that the feurvy is almost unknown to the natives of Canada, though they live entirely on animal food, but not falted; while the use of the latter never fails to produce the fourvy. Would it not then be worth while to make various experiments for preferving meat at fea without falt? Surely the refources of human invention are not exhausted. The absorbent and anti-putrescent properties of fugar are well known; but it might be deemed too expensive a feasoning. I must leave trials of this fort to perfons of more leifure. It is enough for me to fuggest the hint. I with it may lead to a difcovery of fo much importance.

fants

fants in fome parts of Britain express the juice of the last-mentioned plant, and drink it with good effect in those foul fcorbutic eruptions with which they are often troubled in the spring feason.

Harrowgate-water is certainly an excellent medicine in the land-fcurvy. I have often feen patients who had been reduced to the most deplorable condition by this difeafe, greatly relieved by drinking the fulphur-water, and bathing in it. The chalybeate-water may also be used with advantage, especially with a view to brace the stormach after drinking the fulphur-water, which, though it fharpens the appetite, never fails to weaken the powers of digestion.

A flight degree of fcurvy may be carried off by frequently fucking a little of the juice of a bitter orange or a lemon. When the difeafe affects the gums only, this practice, if continued for fome time, will generally carry it off. We would, however, recommend the bitter orange as greatly preferable to lemon; it feems to be as good a medicine, and is not near fo hurtful to the ftomach. Perhaps our own forrel may be little inferior to either of them.

All kinds of falad are good in the fcurvy, and ought to be eaten very plentifully, as fpinage, lettuce, parfley, celery, endive, radifh, dandelion, &c. It is amazing to fee how foon frefh vegetables in the fpring cure the brute animals of any fcab or foulnefs which is upon their fkins. It is reafonable to fuppofe that their effects would be as great upon the human fpecies, were they ufed in proper quantity for a fufficient length of time.

I have fometimes feen good effects in fcorbutic complaints of very long ftanding, from the ufe of a decoction of the roots of water-dock. It is ufually made by boiling a pound of the frefh root in fix Englifh pints of water, till about one-third of it be confumed. The dofe is from half a pint to a whole pint of the decoction every day. But in all the cafes where I have feen it prove beneficial, it was made much ftronger, and drank in larger quantities. The fafeft way, however, is for the patient to begin with fmall dofes, and increafe them both in ftrength and quantity as he finds his ftomach will bear it. It must be ufed

OF THE SCURVY.

ufed for a confiderable time. I have known fome take it for many months, and have been told of others who hadufed it for feveral years, before they were fenfible of any benefit, but who neverthelefs were cured by it at length.

The leprofy, which was fo common in this country long ago, feems to have been near a-kin to the fcurvy. Perhaps its appearing fo feldom now, may be owing to the inhabitants of Britain eating more vegetable food than formerly, living more upon tea and other diluting diet, ufing lefs falted meat, being more cleanly, better lodged and clothed, &c.—For the cure of this difeafe we would recommend the fame courfe of diet and medicine as in the 'curvy.

I have met with very few cafes of real leprofy in the courfe of my practice. The dry, fcaly eruptions all over the body, which are often the effects of the fcurvy, are very liable to be confidered as leprous fymptoms, and certainly refemble them very much. But no evil can arife even from miltake in this particular, as the fame alterative plan, which is advifable in the fcurvy, will be generally found efficacious in the leprofy. Perhaps in the latter complaint, we ought to lay a greater ftrefs, if poffible, on the benefit of good air, and of frequent changes of the linen worn next the fkin. What has been peculiarly called the difeafe of uncleannefs, can only be remedied by the practice of the opposite virtue. I have alfo found, that, after proper means for correcting internal impurities had been used for fome time, the complete difappearance of the leper's fores was often fafely and effectually promoted by the ointment for difeases of the *fkin* mentioned in the Appendix.

The medicinal virtues of the Bath waters, as well as those of Harrowgate, in the cure of the leprofy, are very highly, and I believe very justly, esteemed.

OF THE SCROPHULA, OR KING'S EVIL.

This difeafe chiefly affects the glands, efpecially those of the neck. Children, and young perions of a fedentary life, are very fubject to it. It is one of those difeafes B b 3 which

OF THE SCROPHULA,

which may be removed by proper regimen, but feldom yields to medicine. The inhabitants of cold, damp, marfhy countries are most liable to the fcrophula.

CAUSES .----- This difeafe may proceed from an hereditary taint, infection, a scrophulous nurse, &c. Children who have the misfortune to be born of fickly parents, whole conflitutions have been greatly injured by the pox, or other chronic difeafes, are apt to be affected by the fcrophula. It may likewife proceed from fuch difeafes as weaken the habit or vitiate the humours, as the fmall-pox, meafles, &c. External injuries, as blows, bruifes, and the like, fometimes produce fcrophulous ulcers; but we have reafon to believe, when this happens, that there has been a predifposition in the habit to this difeafe. In fhort, whatever tends to vitiate the humours or relax the folids, paves the way to the fcrophula; as the want of proper exercise, too much heat or cold, confined air, unwholefome food, bad water, the long ule of poor, weak, watery aliments, the neglect of cleanlinels, &c. Nothing tends more to induce this difeafe in children than allowing them to continue long wet *.

SYMPTOMS — At first, small knots appear under the chin, or behind the ears, which gradually increase in number and fize, till they form one large hard tumour. This often continues for a long time without breaking, and when it does break, it only discharges a thin *fanies*, or watery humour. Other parts of the body are likewise liable to its attack, as the arm-pits, groins, feet, hands, eyes, breasts, &c. Nor are the internal parts exempt from it. It often affects the lungs, liver, or spleen; and I have frequently seen the glands of the mysentery greatly enlarged by it.

Those obstinate ulcers which break out upon the feet and hands with swelling, and little or no redness, are of the fcrophulous kind. They feldom discharge good matter, and are exceedingly difficult to cure. The *white fwellings* of the joints seem likewise to be of this kind. They are with difficulty brought to a suppuration, and when open-

^{*} The scrophula, as well as the rickets, is found to prevail in large manufacturing towns, where people live gross, and lead fedentary lives.

ed they only difcharge a thin ichor. There is not a more general fymptom of the fcrophula than a fwelling of the upper lip and nofe. It fometimes begins in a toe or finger, which continues long fwelled, with no great degree of pain, till the bone becomes carious.

REGIMEN. — As this difeafe proceeds, in a great meafure, from relaxation, the diet ought to be generous and nourifhing, but at the fame time light and of eafy digeftion; as well-fermented bread made of found grain, the flefh and broth of young animals, with now and then a glafs of generous wine, or good ale. The air ought to be open, dry, and not too cold, and the patient fhould take as much exercife as he can bear. This is of the utmost importance. Children who have fufficient exercife are feldom troubled with the fcrophula.

MEDICINE. _____ The vulgar are remarkably credulous with regard to the cure of the fcrophula; many of them believing in the virtue of the royal touch, that of the feventh fon, &c. The truth is, we know but little either of the nature or cure of this difeafe, and where reafon or medicines fail, superstition always comes in their place. Hence it is, that in difeafes which are the most difficult to understand, we generally hear of the greatest number of miraculous cures being performed. Here, however, the deception is eafily accounted for. The fcrophula, at a certain period of life, often cures of itfelf; and, if the patient happens to be touched about this time, the cure is imputed to the touch, and not to Nature, who is really the phyfician. In the fame way, the infignificant noftrums of quacks and old women often gain applaufe when they deferve none.

There is nothing more pernicious than the cuftom of plying children in the fcrophula with ftrong purgative medicines. People imagine it proceeds from humours which muft be purged off, without confidering that thefe purgatives increase the debility and aggravate the difease. It has indeed been found, that keeping the body gently open for fome time, especially with sea-water, has a good effect; but this should only be given in gross habits, and in such quantity as to procure one, or at most two stools every day.

Bb 4.

Bathing

376

Bathing in the falt-water has likewife a very good effect, efpecially in the warm feafon. I have often known a courfe of bathing in falt-water, and drinking it in fuch quantities as to keep the body gently open, cure a fcrophula, after many other medicines had been tried in vain. When falt-water cannot be obtained, the patient may be bathed in fresh water, and his body kept open by stand quantities of falt and water, or some other mild purgative.

Next to cold bathing, and drinking the falt-water, we would recommend the Peruvian bark. The cold bath may be used in fummer, and the bark in winter. To an adult half a drachm of the bark in powder may be given in a glass of red wine four or five times a-day. Children, and such as cannot take it in substance, may use the decoction made in the following manner:

Boil an ounce of Peruvian bark and a drachm of Winter's bark, both grofsly powdered, in an Englifh quart of water to a pint: towards the end, half an ounce of fliced liquorice-root and a handful of raifins may be added, which will both render the decoction lefs difagreeable, and make it take up more of the bark. The liquor must be strained, and two, three, or four tablespoonfuls, according to the age of the patient, given three times a-day; but, in place of this, I now use the compound tincture of bark.

The Moffat and Harrowgate waters, efpecially the latter, are likewife very proper medicines in the fcrophula. They ought not, however, to be drank in large quantities, but should be taken so as to keep the body gently open, and must be used for a confiderable time.

The hemlock may fometimes be used with advantage in the fcrophula. Some lay it down as a general rule, that the fea-water is most proper before there are any fuppuration or fymptoms of *tabes*; the Peruvian bark, when there are running fores, and a degree of hectic fever; and the hemlock in old inveterate cases, approaching to the fchirrous or cancerous state. Either the extract or the fresh juice of this plant may be used. The dose may be small at first, and increased gradually as far as the stomach is able to bear it.

4

External

External applications are of little ufe. Before the tumour breaks, nothing ought to be applied to it, unlefs a piece of flannel, or fomething to keep it warm. After it breaks, the fore may be dreffed with fome digeflive ointment. What I have always found to anfwer beft, was the yellow bafilicon mixed with about a fixth or eighth part of its weight of red precipitate of mercury. The fore may be dreffed with this twice a-day; and if it be very fungous, and does not digeft well, a larger proportion of the precipitate may be added.

Medicines which mitigate this difeafe, though they do not cure it, are not to be defpifed. If the patient can be kept alive by any means till he arrives at the age of puberty, he has a great chance to get well; but if he does not recover at this time, in all probability he never will.

There is no malady which parents are fo apt to communicate to their offspring as the fcrophula, for which reafon people ought to beware of marrying into families affected with this difeafe.

For the means of preventing the fcrophula, we must refer the reader to the observations on nursing at the beginning of the book.

I am now induced to bring into one point of view fuch of those remarks as immediately relate to the fubject of prevention, in order to impress them more strongly upon the minds of those who have the care of children.

One of the most effectual means of guarding against the fcrophula, is a constant attention to keep the child dry and clean, by the immediate removal of all impurities, which not only taint the air and relax the skin, but vitiate the humours of the body, in consequence of the abforption of their most noxious particles through the pores.

Washing children frequently, forms a necessary part of this plan. At first, luke-warm water is proper, as being best fuited to the new-born infant, on account of the warm temperature to which he had been accustomed in the womb, and on account of the delicacy of habit which he may have inherited from his parents. But the warmth of the water should be gradually diminished as the infant gains strength, till it can be used quite cold with great fafety and benefit. The cold bath, so effential to the cure

of

of the fcrophula, operates with ftill greater certainty as a preventive. It braces and invigorates the frame, and thus directly counteracts one of the principal caufes of the evil, which is relaxation. The whole body ought to be wafhed every morning, and the lower half every night, after which the child is to be inftantly wiped dry, and wrapped up in a warm blanket, to guard againft the danger of fudden cold, and to fecure all the advantages of fo falutary an operation.

My former arguments, in favour of light and loofe clothing for children in general, acquire double force when there is the leaft reafon to dread the fcrophula. It is little fhort of murder to keep an infant of a delicate habit fmothered in clothes, and panting in a fort of vapour-bath caufed by the noxious fleams of its own body. The covering both by day and by night fhould be as light as is confiftent with due warmth. The linen next the fkin, which is always imbibing perfpirable matter, must be changed often; and the fame drefs ought never to be kept on for more than twelve hours together.

Wholefome unconfined air, and frequent exercife, are grand prefervatives from all difeafes, but efpecially from the fcrophula. It is not enough to felect the moft fpacious and lofty apartment in the houfe for the nurfery; children fhould be taken out into the fields every day, particularly about noon, unlefs the heat be intenfe, as the moft falutary exhalations from the earth then abound, and the air is impregnated with the balmy effence of the fweeteft plants and flowers. Cold and wet weather being deemed one of the exciting caufes of the fcrophula, any wanton expofure to it would be improper; though, even in this refpect, lefs caution is neceffary, if the ufe of the cold bath is continued every morning. This will brace the thinneft, fineft fkin, and harden it againft the imprefions of a damp, chilly atmofphere.

Exercife, befides ftrengthening the whole habit, and powerfully affifting all the vital functions, has a direct tendency to prevent obstructions of every kind, and those of the glands in particular, which constitute the earliest fymptom of the difease in question.

OR KING'S EVIL.

On the fubject of diet, fome little deviations must be made from my general plan, in rearing the child of fcrophulous parents, or one that is marked with what may be called a pre-difposition to this difease, a thin skin, and a general weaknefs and flaccidity of the habit. Extraordinary care fhould be taken to fecure a very healthy nurfe for fuch a child; and, after it is weaned, the use of animal food, but light and eafy of digeftion, should be gradually introduced, and freely allowed at dinner every day. In cafe of any just apprehensions of the scrophula, we must not trust to a mild regimen, to milk and vegetables, though in general fo wholefome and nutritious. They cannot give that tone to the itomach, and that energy to the whole fystem, which they now stand in need of. A grofs, full diet will certainly occasion humours and eruptions; but these are very different from the fcrophula, and far more eafily cured. A poverty of the blood and a relaxation of the fibres, those fure attendants, if not the principal caufes of the evil, require the most strengthening articles both of food and drink.

But I must reprobate, above all things, butter in every form, and other oily fubstances, which are fo apt to turn rancid on the stomach, loading it with phlegm, relaxing and impeding its action, inducing a debility of the folids, and occasioning a great number of complaints, as well as glandular obstructions. One of the worst compositions, of which butter or fat always forms a part, is pastry. I really shudder, whenever I fee a delicate woman, or a weak child, greedily devouring those palatable poisons. Let it be understood, that I include in this censure gingerbread, plumcakes, and all trass of the like kind. Indeed, a child of a strophulous habit should never eat any preparation of flour, except plain, well-made, and well-baked bread.

OF THE ITCH.

Though this difeafe is commonly communicated by infection, yet it feldom prevails where due regard is paid to cleanlinefs, fresh air, and wholefome diet. It generally

OF THE ITCH.

rally appears in form of fmall watery puffules, first about the wrifts, or between the fingers; afterwards it affects the arms, legs, thighs, &c. These puffules are attended with an intolerable itching, especially when the patient is warm in bed, or fits by the fire Sometimes, indeed, the skin is covered with large blotches or scabs, and at other times with a white fcurf, or scaly eruption. This last is called the Dry Itch, and is the most difficult to cure.

The itch is feldom a dangerous difeafe, unlefs when it is rendered fo by neglect or improper treatment. If it be fuffered to continue too long, it may vitiate the whole mafs of humours; and, if it be fuddenly drove in, without proper evacuations, it may occafion fevers, inflammations of the vifcera, or other internal diforders.

The beft medicine yet known for the itch is fulphur, which ought to be ufed both externally and internally. The parts moft affected may be rubbed with an ointment made of the flower of fulphur, two ounces; crude fal ammoniac finely powdered, two drachms; hog's lard, or butter, four ounces. If a fcruple or half a drachm of the effence of lemon be added, it will entirely take away the difagreeable fmell. About the bulk of a nutmeg of this may be rubbed upon the extremities at bed-time twice or thrice a-week. It is feldom neceffary to rub the whole body; but when it is, it ought not to be done all at once, but by turns, as it is dangerous to ftop too many pores at the fame time.

Before the patient begins to use the ointment, he ought, if he be of a full habit, to bleed or take a purge or two. It will likewise be proper, during the use of it, to take every night and morning as much of the flower of brimstone and cream of tartar, in a little treacle or new milk, as will keep the body gently open. He should beware of catching cold, should wear more clothes than usual, and take every thing warm. The same clothes, the linen excepted, ought to be worn all the time of using the ointment; and such clothes as have been worn while the patient was under the difease, are not to be used again, unless they have been fumigated with brimstone, and

OF THE ITCH.

and thoroughly cleanfed, otherwife they will communicate the infection anew *:

I never knew brimftone, when ufed as directed above, fail to cure the itch; and I have reafon to believe, that, if duly perfifted in, it never will fail; but if it be only ufed once or twice, and cleanlinefs neglected, it is no wonder if the diforder returns. The quantity of ointment mentioned above will generally be tufficient for the cure of one perfon; but, if any fymptoms of the difeafe fhould appear again, the medicine muft be repeated. It is both more fafe and efficacious when perfifted in for a confiderable time, than when a large quantity is applied at once. As most people diflike the fmell of fulphur, they may ufe in its place the powder of white hellebore root made up into an ointment, in the fame manner, which will feldom fail to cure the itch.

People ought to be extremely cautious left they take other eruptions for the itch; as the floppage of thefe may be attended with fatal confequences. Many of the eruptive diforders to which children are liable, have a near refemblance; and I have often known infants killed by being rubbed with greafy ointments that make thefe eruptions flrike fuddenly in, which nature had thrown out to preferve the patient's life, or prevent fome other malady.

Much mifchief is likewife done by the ufe of mercury in this difeafe. Some perfons are fo fool-hardy as to wafh the parts affected with a ftrong folution of the corrofive fublimate. Others ufe the mercurial ointment, without taking the leaft care either to avoid cold, keep the body open, or obferve a proper regimen. The confequences of fuch conduct may be eafily gueffed. I have known even the mercurial girdles produce bad effects, and would advife every perfon, as he values his health, to beware how he ufes them. Mercury ought never to

* Sir John Pringle obferves, that though this difeafe may feem triffing, there is no one in the army that is more troublefome to cure, as the infection often lurks in clothes, &c. and breaks out a second, or even a third time. The fame inconveniency occurs in private families, unlefs particular regard be paid to the changing or cleaning of their clothes, which laft is by no means an easy operation. be used as a medicine without the greatest care. Ignorant people look upon these girdles as a kind of charm, without confidering that the mercury enters the body.

It is not to be told what mifchief is done by using mercurial ointment for curing the itch and killing vermin; yet is is unneceffary for either : the former may be always more certainly cured by fulphur, and the latter will never be found where due regard is paid to cleanlinefs.

Those who would avoid this detestable difease, ought to beware of infected perfons, to use wholesome food, and to study universal cleanliness *.

CHAP. XL.

OF THE ASTHMA.

THE affhma is a difeafe of the lungs, which feldom admits of a cure. Perfons in the decline of life are most liable to it. It is diffinguished into the moist and dry, or humoural and nervous The former is attended with expectoration or spitting; but in the latter the patient feldom spits, unless sometimes a little tough phlegm, by the mere force of coughing.

CAUSES. —— The afthma is fometimes hereditary. It may likewife proceed from a bad formation of the breaft; the fumes of metals or minerals taken into the lungs; violent exercife, efpecially running; the obftruction of cultomary evacuations, as the menfes, hæmorthoids, &c. the ludden retroceffion of the gout, or ftrik-

* The itch is now by cleanlinefs banifhed from every genteel family in Britain. It ftill, however, prevails among the poorer forts of peafants in Scotland, and among the manufacturers in England. These are not only fufficient to keep the feeds of the difeate alive, but to fpread the infection among others. It were to be wifhed that fome effectual method could be devifed for extirpating it altogether. Several country clergymen have told me, that by getting fuch as were infected cured, and ftrongly recommending an attention to cleanlinefs, they have banifhed the itch entirely out of their parishes. Why might not others do the fame ?

ing in of eruptions, as the fmall-pox, meafles, &c. violent paffions of the mind, as fudden fear or furprife. In a word, the difeafe may proceed from any caufe that either impedes the circulation of the blood through the lungs, or prevents their being duly expanded by the air.

SYMPTOMS.——An afthma is known by a quick laborious breathing, which is generally performed with a kind of wheezing noife. Sometimes the difficulty of breathing is fo great, that the patient is obliged to keep in an erect pofture, otherwife he is in danger of being fuffocated. A fit or paroxyfm of the afthma generally happens after a perfon has been exposed to cold eafterly winds, or has been abroad in thick foggy weather, or has got wet, or continued long in a damp place under ground, or has taken fome food which the ftomach could not digeft, as paftries, toafted cheefe, or the like.

The paroxyfm is commonly ufhered in with liftneffnefs, want of fleep, hoarfenefs, a cough, belching of wind, a fenfe of heavinefs about the breaft, and difficulty of breathing. To thefe fucceed heat, fever, pain of the head, ficknefs and naufea, great oppreffion of the breaft, palpitation of the heart, a weak and fometimes intermitting pulfe, an involuntary flow of tears, bilious vomitings, &c. All the fymptoms grow worfe towards night; the patient is eafier when up than in bed, and is very defirous of cool air.

REGIMEN.——The food ought to be light and of eafy digeftion. Boiled meats are to be preferred to roafted, and the flefh of young animals to that of old. All windy food, and whatever is apt to fwell in the ftomach, is to be avoided. Light puddings, white broths, and ripe fruits baked, boiled, or roafted, are proper. Strong liquors of all kinds, especially malt-liquor, are hurtful. The patient should eat a very light supper, or rather none at all, and should never fuffer simfelf to be long costive. His clothing should be warm, especially in the winter feason. As all diforders of the breast are much relieved by keeping the feet warm, and promoting the perspiration, a flannel shirt or waistcoat, and thick shoes, will be of fingular fervice.

But nothing is of fo great importance in the afthma as pure and moderately warm air. Afthmatic people can feldom bear either the close heavy air of a large town, or the sharp keen atmosphere of a bleak hilly country : a medium, therefore, between these is to be chosen. The air near a large town is often better than at a diftance, provided the patient be removed fo far as not to be affected by the fmoke. Some afthmatic patients indeed breathe eafier in town than in the country; but this is feldom the cafe, especially in towns where much coal is burnt. Afthmatic perfons who are obliged to be in town all day, ought at least to fleep out of it. Even this will often prove of great fervice. Those who can afford it ought to travel into a warmer climate. Many afthmatic perfons who cannot live in Britain, enjoy very good health in the fouth of France, Portugal, Spain, or Italy.

Exercife is likewife of very great importance in the afthma, as it promotes the digeftion, and greatly affifts in the preparation of the blood. The blood of afthmatic perfons is feldom duly prepared, owing to the proper action of the lungs being impeded. For this reafon fuch people ought daily to take as much exercife, either on foot, horfeback, or in a carriage, as they can bear.

medicine in this difeafe, is to relieve the patient when feized with a violent fit. This indeed requires the greateft expedition, as the difeafe often proves fuddenly fatal. In a paroxyfin or fit the body is generally bound, a purging clyfter, with a folution of afafætida, ought therefore to be administered, and if there be occasion, it may be repeated two or three times. The patient's feet and legs ought to be immerfed in warm water, and afterwards rubbed with a warm hand, or dry cloth. Bleeding, unlefs extreme weaknefs or old age fhould forbid it, is highly proper. If there be a violent fpafin about the breaft or ftomach, warm fomentations or bladders filled with warm milk and water, may be applied to the part affected, and warm cataplaims to the foles of the feet. The patient must drink freely of diluting liquors, and may take a tea-spoonful of the tincture of caftor and of faffron mixed together in a cup of valeriantea,

tea, twice or thrice a day. Sometimes a vomit has a very good effect, and fnatches the patient, as it were, from the jaws of death. This, however, will be more fafe after other evacuations have been premifed. A very ftrong infufion of roafted coffee is faid to give eafe in an afthmatic paroxyfm.

In the moift afthma, fuch things as promote expectoration or fpitting, ought to be ufed; as the fyrup of fquills, gum-ammoniac, and fuch like. A common fpoonful of the fyrup or oxymel of fquills, mixed with an equal quantity of cinnamon-water, may be taken three or four times through the day, and four or five pills made of equal parts of afafœtida and gum-ammoniac at bedtime *.

For the convultive or nervous afthma, antifpafmodics and bracers are the most proper medicines. The patient may take a tea-spoonful of the paregoric elixir twice aday. The Peruvian bark is sometimes found to be of use in this case. It may be taken in substance, or infused in wine. In short, every thing that braces the nerves, or takes off spasm, may be of use in a nervous assesses or takes off spasm, may be of use in a nervous assesses. It is often relieved by the use of affes' milk : I have likewise known cow's milk drank warm in the morning have a very good effect in this case.

In every fpecies of afthma, fetons and iffues have a good effect; they may either be fet in the back or fide, and fhould never be allowed to dry up. We fhall here, once for all, obferve, that not only in the afthma, but in most chronic difeases, iffues are extremely proper. They are both a fase and efficacious remedy; and though they do not always cure the difease, yet they will often prolong the patient's life.

This difeafe, though fo common with us, is little known in mild climates; and, on that account, it is always advifable to try the effect of a change of climate, which has

* After copious evacuations, large dofes of æther have been found very efficacious in removing a fit of the afthma. I have likewife known the following mixture produce very happy effects: To four or five ounces of the folution or milk of gum-ammoniac add two ounces of fimple cinnamon-water, the fame quantity of balfamic fyrup, and half an ounce of paregoric elixir. Of this two table-fpoonfuls may be taken every three hours.

generally

generally been attended with great benefit. I have already intimated what little confidence I had in the power of any medicine to perform a radical cure of the affhma; but there are many things that will give the patient eafe, and, of courfe, tend to prolong his life. Much alfo may be done by regimen, when drugs are of little fervice; and I would therefore advife affhmatic patients to procure and keep by them rules for their management both in and out of the fit, adapted to their particular cafes. By a proper attention to fuch rules, a man may live many years, and enjoy tolerably good health.

I had a patient fome time ago, who was often carried home to his wife in an apparently dying flate. She felt little alarm, well-knowing what was neceffary to be done; and fhe always brought him about. This good woman did no more than may be done by any woman of common fenfe, if the doctor will deign to inftruct her. General rules will not do. They muft, as before obferved, be fuited to the patient's cafe and confliction. For want of fome fuch inftructions, which a phyfician fhould take the earlieft opportunity to give, a patient may lofe his life, before the doctor can be fent for, or any other medical advice or affiftance procured.

CHAP. XLI.

OF THE APOPLEXY.

THE apoplexy is a fudden lofs of fenfe and motion, during which the patient is to all appearance dead; the heart and lungs, however, ftill continue to move. Though this difeafe proves often fatal, yet it may fometimes be removed by proper care. It chiefly attacks fedentary perfons of a grofs habit, who ufe a rich and plentiful diet, and indulge in ftrong liquors. People in the decline of life are most fubject to the apoplexy. It prevails most in winter, especially in rainy feasons, and very low states of the barometer.

CAUSES.——The immediate caufe of an apoplexy is a compression of the brain, occasioned by an excess of blood,

4110132333

blood, or a collection of watery humours. The former is called a fanguine, and the latter a ferous apoplexy. It may be occafioned by any thing that increases the circulation towards the brain, or prevents the return of the blood from the head: as intenfe fludy; violent paffions*; viewing objects for a long time obliquely; wearing any thing too tight about the neck; a rich and luxurious diet; suppression of urine; fuffering the body to cool fuddenly after having been greatly heated; continuing long in a warm or cold bath; the exceffive use of spiceries, or high-feafoned food; excefs of venery; the fudden striking in of any eruption; fuffering iffues, fetons, &c. fuddenly to dry up, or the ftoppage of any cuftomary evacution; a mercurial falivation pushed too far, or fuddenly checked by cold; wounds or bruifes on the head ; long expolure to exceffive cold ; poifonous exhalations, &c.

SYMPTOMS, and method of cure. ——— The ufual forerunners of an apoplexy are giddinefs, pain and fwimming of the head; lofs of memory; drowfinefs; noife in the ears; the night-mare; a fpontaneous flux of tears, and laborious refpiration. When perfons of an apoplectic make, obferve thefe fymptoms, they have reafon to fear the approach of a fit, and fhould endeavour to prevent it by bleeding, a flender diet, and opening medicines.

In the fanguine apoplexy, if the patient does not die fuddenly, the countenance appears florid, the face is fwelled or puffed up, and the blood veffels, efpecially about the neck and temples, are turgid; the pulfe beats ftrong; the eyes are prominent and fixed, and the breathing is difficult, and performed with a fnorting noife. The excrements and urine are often voided fpontaneoufly, and the patient is fometimes feized with vomiting.

* I knew a woman who in a violent fit of anger was feized with a fanguine apoplexy. She at first complained of extreme pain, as if daggers had been thrust through her head, as she expressed it. Afterwards she became comatore, her pulse funk very low, and was exceeding flow. By bleeding, blistering, and other evacuations, she was kept alive for about a fortnight. When her head was opened, a large quantity of extravasfated blood was found in the left ventricle of the brain.

Cc2

In

In this species of apoplexy every method must be taken to leffen the force of the circulation towards the head. The patient should be kept perfectly eafy and cool. His head should be raifed pretty high, and his feet fuffered to hang down. His clothes ought to be loofened, especially about the neck, and freshair admitted into his chamber. His garters fhould be tied pretty tight, by which means the motion of the blood from the lower extremities will be retarded. As foon as the patient is placed in a proper posture, he should be bled freely in the neck or arm, and, if there be occafion, the operation may be repeated in two or three hours. A laxative clyfter, with plenty of fweet oil, or fresh butter, and a fpoonful or two of common falt in it, may be administered every two hours; and blistering-plasters applied between the fhoulders, and to the calves of the legs.

As foon as the fymptoms are a little abated, and the patient is able to fwallow, he ought to drink freely of fome diluting opening liquor; as a decoction of tamarinds and liquotice, cream-tartar whey, or common whey with cream of tartar diffolved in it. Or he may take any cooling purge, as Glauber's falts, manna diffolved in an infufion of fenna, or the like. All fpirits and other ftrong liquors are to be avoided. Even volatile falts held to the nofe do mifchief. Vomits, for the fame reafon, ought not to be given, or any thing that may increafe the motion of the blood towards the head.

In the ferous apoplexy, the fymptoms are nearly the fame, only the pulfe is not fo ftrong, the countenance is lefs florid, and the breathing lefs difficult. Bleeding is not fo neceffary here as in the former cafe. It may, however, generally be performed once with fafety and advantage, but fhould not be repeated. The patient fhould be placed in the fame pofture as directed above, and fhould have bliftering-plafters applied, and receive opening clyfters in the fame manner. Purges are here likewife neceffary, and the patient may drink ftrong balmtea. If he be inclined to fweat, it ought to be promoted by drinking fmall wine whey, or an infufion of carduus benedictus. A plentiful fweat kept up for a confiderable time has often carried off a ferous apoplexy.

When

OF THE APOPLEXY.

When apoplectic fymptoms proceed from opium, or other narcotic fubftances taken into the ftomach, vomits are neceffary. The patient is generally relieved as foon as he has difcharged the poifon in this way.

Perfons of an apoplectic make, or those who have been attacked by it, ought to use a very spare and flender diet, avoiding all ftrong liquors, fpiceries, and high-featoned food. They ought likewife to guard against all violent paffions, and to avoid the extremes of heat and cold. The head fhould be fhaved, and daily washed with cold water. The feet ought to be kept warm, and never fuffered to continue long wet. The body must be kept open either by food or medicine, and a little blood may be let every fpring and fall. Exercise should by no means be neglected; but it ought to be taken in moderation. Nothing has a more happy effect in preventing an apoplexy than perpetual iffues or fetons; great care, however, must be taken not to fuffer them to dry up. without opening others in their flead. Apoplectic perfons ought never to go to reft with a full ftomach, or to lie with their heads low, or wear any thing too tight about their necks.

These last cautions are of far greater importance than. fuch perfons may be aware of. The circulation, which is flower during fleep than when awake, is farther clogged by a fulnefs of the ftomach. The low pofture of the head not only favours, but feems to invite stagnation; and tight ligatures round the neck impede the return of the blood from the veffels of the brain, fo that an apoplexy, not only very naturally, but almost inevitably follows. Instead of being aftonished at the number of those who go to bed in apparent health, and are found dead in the morning, we fhould confider it as a matter of much more furprife for a perfon of a plethoric habit, after unchecked indulgence in the pleafures of the table, to go to reft without any regard to the inclination of his head or the tightness of his collar, and ever to rife again.

Cc3

CHAP_

CHAP. XLII.

OF COSTIVENESS, AND OTHER AFFEC, TIONS OF THE STOMACH AND BOWELS.

WE do not here mean to treat of those aftrictions of the bowels which are the fymptoms of difeases, as of the colic, the iliac passion, &c. but only to take notice of that infrequency of stools which sometimes happens, and which in some particular constitutions may occasion difeases.

Coftivenels may proceed from drinking rough red wines, or other aftringent liquors; too much exercife, efpecially on horfeback. It may likewife proceed from a long use of cold infipid food, which does not fufficiently ftimulate the intestines. Sometimes it is owing to the bile not descending to the intestines, as in the jaundice; and at other times it proceeds from diseases of the intesttines themselves, as a palfy, spafms, torpor, tumours, a cold dry state of the intestines, &c.

Excellive collivenels is apt to occalion pains of the head, vomiting, colics, and other complaints of the bowels. It is peculiarly hurtful to hypochondriac and hyfteric perfons, as it generates wind and other grievous fymptoms. Some people however can bear coftivenels to a great degree. I know perfons who enjoy pretty good health, yet do not go to ftool above once a-week, and others not above once a-fortnight. Indeed I have heard of fome who do not go above once amonth.

Perfons who are generally coftive fhould live upon a moiftening and laxative diet; as roafted or boiled apples, pears, flewed prunes, raifins, gruels with currants, butter, honey, fugar, and fuch like. Broths with fpinage, leeks, and other foft pot-herbs, are likewife proper. Ryebread, or that which is made of a mixture of wheat and rye together, ought to be eaten. No perfon troubled with coftiveness should eat white bread alone, especially that that which is made of fine flour. The beft bread for keeping the body foluble is what in fome parts of England they call *meflin*. It is made of a mixture of wheat and rye, and is very agreeable to those who are accustomed to it.

Coffivenels is increased by keeping the body too warm, and by every thing that promotes the perspiration; as wearing flannel, lying too long in bed, &c. Intense thought and a fedentary life are likewise hurtful. All the fecretions and excretions are promoted by moderate exercise without doors, and by a gay, cheerful, sprightly temper of mind.

The drink fhould be of an opening quality. All ardent fpirits, auftere and aftringent wines, as port, claret, &c. ought to be avoided. Malt-liquor that is fine, and of a moderate ftrength, is very proper. Butter-milk, whey, and other watery liquors, are likewife proper, and may be drank in turns, as the patient's inclination directs.

Those who are troubled with coffiveness, ought, if possible, to remedy it by diet, as the constant use of medicines for that purpose is attended with many inconveniencies, and often with bad consequences *. I never knew any one get into a habit of taking medicine for keeping the body open, who could leave it off. In time

* The learned Dr. Arbuthnot advises those who are troubled with coftiveness to use animal oils, as fresh butter, cream, marrow, fat broths, especially those made of the internal parts of animals, as the liver, heart, midriff, &c. He likewise recommends the expressed oils of mild vegetables, as olives, almonds, pattaches, and the fruits themselves; all oily and mild fruits, as figs; decoctions of mealy vegetables; these lubricate the inteftines; fome faponaceous substances which stimulate gently, as honey, hydromel, or boiled honey and water, unrefined fugar, &c.

The Doctor obferves, that fuch lenitive fubftances are proper for perfons of dry atrabilarian conflictutions, who are fubject to aftriction of the belly, and the piles, and will operate when ftronger medicinal fubftances are fometimes ineffectual; but that fuch lenitive diet hurts those whose bowels are weak and lax. He likewife observes, that all watery fubftances are lenitive, and that even common water, whey, four milk, and butter-milk, have that effect :—That new milk, especially affes milk, ftimulates ftill more when it fours on the ftomach; and that whey turned four will purge ftrongly. the cuftom becomes neceffary, and generally ends in a total relaxation of the bowels, indigeftion, lofs of appetite, wafting of the ftrength, and death.

When the body cannot be kept open without medicine, we would recommend gentle dofes of rhubarb to be taken twice or thrice a-week. This is not near fo injurious to the flomach as aloes, jalap, or the other draftic purgatives fo much in ufe. Infufions of fenna and manna may likewife be taken, or half an ounce of foluble tartar diffolved in water-gruel. About the fize of a nutmeg of lenitive electuary taken twice or thrice a-day generally anfwers the purpofe very well.

WANT OF APPETITE.

This may proceed from a foul ftomach; indigeftion; the want of free air and exercife; grief, fear, anxiety, or any of the deprefling paffions; exceffive heat; the ufe of ftrong broths, fat meats, or any thing that palls the appetite, or is hard of digeftion; the immoderate ufe of ftrong liquors, tea, tobacco, opium, &c.

The patient ought, if poffible, to make choice of an open dry air; to take exercise daily on horseback or in a carriage; to rise betimes; and to avoid all intense thought. He should use a diet of easy digestion; and should avoid excessive heat and great fatigue.

If want of appetite proceed from errors in diet, or any other part of the patient's regimen, it ought to be changed. If naufea and reachings fhew that the ftomach is loaded with crudities, a vomit will be of fervice. After this a gentle purge or two of rhubarb, or any of the bitter purging falts, may be taken. The patient ought next to use fome of the ftomachic bitters infused in wine. Though gentle evacuations be necessary, yet ftrong purges and vomits are to be avoided, as they weaken the ftomach, and hurt digestion. After proper evacuations, bitter elixirs and tinctures with aromatics may be used.

Elixir of vitriol is an excellent medicine in most cales of indigestion, weakness of the stomach, or want of appetite. From twenty to thirty drops of it may be taken twice or thrice a-day in a glass of wine or water. It may likewise be mixed with the tincture of the bark, one drachm

WANT OF APPETITE.

393

drachm of the former to an ounce of the latter, and two tea-spoonfuls of it taken in wine and water, as above.

The chalybeate waters, if drank in moderation, are generally of confiderable fervice in this cafe. The faltwater has likewife good effects; but it muft not be ufed too freely. The waters of Harrowgate, Scarborough, Moffat, and moft other fpas in Britain, may be ufed with advantage. We would advife all who are afflicted with indigeftion and want of appetite, to repair to these places of public rendezvous. The very change of air, and the cheerful company, will be of fervice; not to mention the exercise, diffipation, amufements, &c.

OF THE HEART-BURN.

What is commonly called the *heart-burn*, is not a difeafe of that organ, but an uneafy fensation of heat or acrimony about the pit of the flomach, which is fometimes attended with anxiety, nausea, and vomiting.

It may proceed from debility of the ftomach, indigeftion, bile, the abounding of an acid in the ftomach, &c. Perfons who are liable to this complaint ought to avoid ftale liquors, acids, windy or greafy aliments, and fhould never ufe violent exercise foon after a plentiful meal. I know many perfons who never fail to have the heart-burn if they ride foon after dinner, provided they have drank ale, wine, or any fermented liquor; but are never troubled with it when they have drank rum or brandy and water without any fugar or acid.

When the heart-burn proceeds from debility of the ftomach, or indigeftion, the patient ought to take a dofe or two of rhubarb; afterwards he may use infusions of the Peruvian bark, or any other of the ftomachic bitters, in wine or brandy. Drinking a cup of camomile-tea, with fifteen or twenty drops of elixir of vitriol in it, twice or thrice a-day, will ftrengthen the ftomach and promote digestion. Exercise in the open air will likewise be of use.

When bilious humours occafion the heart-burn, a teafpoonful of the fweet fpirit of nitre in a glafs of water, or a cup of tea, will generally give eafe. If it proceeds from the use of greafy aliments, a dram of brandy or rum may be taken.

If acidity or fournefs of the ftomach occasions the heart-burn, absorbents are the proper medicines. In this cafe an ounce of powdered chalk, half an ounce of fine fugar, and a quarter of an ounce of gum-arabic, may be mixed in an English quart of water, and a tea-cupful of it taken as often as is neceffary. Such as do not chufe chalk, may take a tea-fpoonful of prepared oyfter-fhells, or of the powder called crabs-eyes, in a glafs of cinnamon or peppermint-water. But the fafeft and beft abforbent is magnefia alba. This not only acts as an abforbent, but likewife as a purgative; whereas chalk, and other abforbents of that kind, are apt to lie in the inteffines, and occasion obstructions. This powder is not difagreeable, and may be taken in a cup of tea, or a glafs of mint-water. A large tea-fpoonful is the ufual dofe; but it may be taken in a much greater quantity when there is occafion. These things are now generally made up into lozenges for the conveniency of being carried in the pocket, and taken at pleafure.

If wind be the caufe of this complaint, the moft proper medicines are those called carminatives; as anifeeds, juniper-berries, ginger, canella alba, cardamom feeds, &c. These may either be chewed, or infused in wine, brandy, or other spirits; but these ought never to be used, unless they are absolutely necessary, as they are only drams in a dry form, and very pernicious to the stomach. One of the fafest medicines of this kind is the tincture made by infusing an ounce of rhubarb, and a quarter of an ounce of the less cardamom feeds, in an English pint of brandy. After this has digested for two or three days, it ought to be strained, and four ounces of white sugar-candy added to it. It must stand to digest a fecond time till the sugar be dissolved. A tablestrained of it may be taken occasionally for a dose.

I have frequently known the heart-burn cured, particularly in pregnant women, by chewing green tea. Two table-fpoonfuls of what is called the milk of gum-ammoniac, taken once or twice a-day, will fometimes cure the heart-burn.

As pregnant women are very fubject to this unealy fenfation, they fhould first confider, whether it proceeds from

OF THE HEART-BURN.

from any of the caufes already explained; in which cafe the medicines prefcribed under each head will probably remove it. But if the internal fenfe of heat be owing to the ftate of pregnancy itfelf; if it arifes from the confent between the ftomach and the womb, and is not accompanied with much fpitting or any acid eructations, the white of an egg, mixed with a little fugar and water, will often afford the only relief that can be expected for fome time.

CHAP. XLIII. OF NERVOUS DISEASES.

OF all difeafes incident to mankind, those of the nervous kind are the molt complicated and difficult to cure. A volume would not be fufficient to point out their various appearances. They imitate almost every difeafe; and are feldom alike in two different perfons, or even the fame perfon at different times. Proteus-like, they are continually changing fhape; and upon every fresh attack, the patient thinks he feels fymptoms which he never experienced before. Nor do they only affect the body; the mind likewife fuffers, and is often thereby rendered extremely weak and peevifh. The low fpirits, timoroufnefs, melancholy, and ficklenefs of temper, which generally attend nervous diforders, induce many to believe that they are entirely difeafes of the mind; but this change of temper is rather a confequence, than the caule of nervous difeafes.

CAUSES.——Every thing that tends to relax or weaken the body, difpofes it to nervous difeafes, as indolence, excellive venery, drinking too much tea, or other weak watery liquors warm, frequent bleeding, purging, vomiting, &c. Whatever hurts the digeftion, or prevents the proper affimilation of the food, has likewife this effect; as long fafting, excefs in eating or drinking, the ufe of windy, crude, or unwholefome aliments, an unfavourable pofture of the body, &c.

Nervous diforders often proceed from intenfe application to ftudy. Indeed few ftudious perfons are entirely free from them. Nor is this at all to be wondered at; intenfe thinking not only preys upon the fpirits, but prevents the perfon from taking proper exercife, by which

395

means

OF NERVOUS DISEASES. 395

means the digeftion is impaired, the nourifhment prevented, the folids relaxed, and the whole mafs of humours vitiated. Grief and difappointment likewife produce the fame effects. I have known more nervous patients who dated the commencement of their diforders from the lofs of a hufband, a favourite child, or from some disappointment in life, than from any other caufe. In a word, whatever weakens the body, or depresses the spirits, may occasion nervous diforders; as unwholesome air, want of fleep, great fatigue, difagreeable apprehenfions, anxiety, vexation, &c.

SYMPTOMS.-----We fhall only mention fome of the most general symptoms of these diforders, as it would be both an useless and impracticable task to enumerate the whole. They generally begin with windy inflations or diffentions of the ftomach and inteffines; the appetite and digeftion are ufually bad; yet fometimes there is an uncommon craving for food, and a quick digeftion. The food often turns four on the ftomach; and the patient is troubled with vomiting of clear water, tough phlegm, or a blackifh-coloured liquor refembling the grounds of coffee. Excruciating pains are often felt about the navel, attended with a rumbling or murmuring noife in the bowels. The body is fometimes loofe, but more commonly bound, which occasions a retention of wind and great uneafinefs.

The urine is fometimes in fmall quantity, at other times very copious and quite clear. There is a great ftraitnels of the breaft, with difficulty of breathing; violent palpitations of the heart; fudden flushings of heat in various parts of the body; at other times a fenfe of cold, as if water were poured on them; flying pains in the arms and limbs, pains in the back and belly, refembling those occasioned by gravel; the pulle very variable, fometimes uncommonly flow, and at other times very quick; yawning, the hiccup, frequent fighing, and a fense of fuffocation, as if from a ball or lump in the throat; alternate fits of crying and convulfive laughing; the fleep is unfound, and feldom refreshing; and the patient is often troubled with the night-mare.

As the difeafe increafes, the patient is molefted with head-aches, cramps, and fixed pains in various parts of the body;

body; the eyes are clouded, and often affected with pain and drynefs; there is a noife in the ears, and often a dulnefs of hearing; in fhort, the whole animal functions are impaired. The mind is difturbed on the moft trivial occafions, and is hurried into the moft perverfe commotions, inquietudes, terror, fadnefs, anger, diffidence, &c. The patient is apt to entertain wild imaginations, and extravagant fancies; the memory becomes weak, and the judgment fails.

Nothing is more characteriftic of this difeafe than a conftant dread of death. This renders those unhappy perfons who labour under it peevish, fickle, impatient, and apt to run from one physician to another; which is one reason why they feldom reap any benefit from medicine, as they have not sufficient resolution to perfiss in any one course till it has time to produce its proper effects. They are likewise apt to imagine that they labour under difeases from which they are quite free; and are very angry if any one attempts to fet them right, or laugh them out of their ridiculous notions.

REGIMEN .--- Perfons afflicted with nervous difeafes ought never to fast long. Their food should be folid and nourifhing, but of eafy digeftion. Fat meats and heavy fauces are hurtful. All excess should be carefully avoided. They ought never to eat more at a time than they can eafily digeft; and heavy fuppers are to be avoided. If they feel themfelves weak and faint between meals, they ought to eat a bit of bread, and drink a glafs of wine. Though wine in excels enfeebles the body, and impairs the faculties of the mind, yet taken in moderation, it ftrengthens the ftomach, and promotes digestion. Wine and water is a very proper drink at meals: but if wine fours on the flomach, or the patient is much troubled with wind, brandy and water will answer better. Every thing that is windy or hard of digeftion must be avoided. All weak and warm liquors are hurtful; as tea, coffee, punch, &c. People may find a temporary relief in the use of these, but they always increase the malady, as they weaken the ftomach, and hurt digeftion. Above all things, drams are to be avoided. Whatever immediate ease the patient may feel from the use of ardent fpirits, they are fure to aggravate the malady, and prove certain 398

certain poifons at laft. These cautions are the more neceffary, as most nervous people are peculiarly fond of tea and ardent spirits; to the use of which many of them fall victims.

Exercife in nervous diforders is fuperior to all medicines. Riding on horfeback is generally effeemed the beft, as it gives motion to the whole body, without fatiguing it. I have known fome patients, however, with whom walking agreed better, and others who were moft benefited by riding in a carriage. Every one ought to ufe that which he finds molt beneficial. Long feavoyages have an excellent effect; and to those who have fufficient resolution, we would by all means recommend this courfe. Even change of place, and the fight of new objects, by diverting the mind, have a great tendency to remove these complaints. For this reason a long journey, or a voyage, is of much more advantage than riding fhort journeys near home.

A cool and dry air is proper, as it braces and invigorates the whole body. Few things tend more to relax and enervate than hot air, especially that which is rendered fo by great fires, or floves in fmall apartments. But when the ftomach or bowels are weak, the body ought to be well-guarded against cold, especially in winter, by wearing a thin flannel waiftcoat next the fkin. This will keep up an equal perspiration, and defend the alimentary canal from many impreffions to which it would otherwife be fubject, upon every fudden change from warm to cold weather. Rubbing the body frequently with a flefh-brufh, or a coarfe linen-cloth, is likewife beneficial; as it promotes the circulation, perfpiration, &c. Perfons who have weak nerves ought to rife early, and take exercife before breakfaft, as lying too long a bed cannot fail to relax the folids. They ought likewife to be diverted, and to be kept as eafy and cheerful as poffible. There is not any thing which hurts the nervous fystem, or weakens the digestive powers, more than fear, grief, or anxiety.

MEDICINES. ———— Though nervous difeafes are feldom radically cured, yet their fymptoms may fometimes be alleviated, and the patient's life rendered at leaft more comfortable by proper medicines.

When

OF NERVOUS DISEASES.

When the patient is coftive, he ought to take a little rhubarb, or fome other mild purgative, and fhould never fuffer his body to be long bound. All ftrong and violent purgatives are, however, to be avoided; as aloes, jalap, &c. I have generally feen an infufion of fenna and rhubarb in brandy anfwer very well. This may be made of any ftrength, and taken in fuch quantity as the patient finds neceffary. When the digeftion is bad, or the ftomach relaxed and weak, the following infufion of Peruvian bark and other bitters may be used with advantage :

Take of Peruvian bark an ounce, gentian-root, orangepeel, and coriander-feed, of each half an ounce; let thefe ingredients be all bruifed in a mortar, and infufed in a bottle of brandy or rum, for the fpace of five or fix days. A table-fpoonful of the ftrained liquor may be taken in half a glafs of water, an hour before breakfaft, dinner, and fupper.

Few things tend more to ftrengthen the nervous fyftem than cold bathing. This practice, if duly perfifted in, will produce very extraordinary effects; but when the liver or other *vifcera* are obftructed, or otherwife unfound, the cold bath is improper. It is therefore to be ufed with very great caution. The most proper feafons for it are fummer and autumn. It will be fufficient, especially for perfons of a spare habit, to go into the cold bath three or four times a-week. If the patient be weakened by it, or feels chilly for a long time after coming out, it is improper.

In patients afflicted with wind, I have always obferved the greateft benefit from the elixir of vitriol. It may be taken in the quantity of fifteen, twenty, or thirty drops, twice or thrice a-day, in a glafs of water. This both expels wind, ftrengthens the ftomach, and promotes digeftion.

Opiates are generally extolled in thefe maladies; but as they only palliate the fymptoms, and generally afterwards increase the difease, we would advise people to be extremely sparing in the use of them, left habit should render them at last absolutely necessary *.

* Few days have paffed for a confiderable time, that I have not had occafion to recommend the following tincture to fome of my nervous patients, and I have feldom been difappointed with regard to its effects: Take of compound tincture of the bark and volatile

It

400

It would be an eafy matter to enumerate many medicines which have been extolled for relieving nervous diforders; but whoever wifhes for a thorough cure, must expect it from regimen alone: we shall therefore omit mentioning more medicines, and again recommend the strictest attention to DIET, AIR, EXERCISE, and AMUSE-MENTS.

OF MELANCHOLY.

Melancholy is that flate of alienation or weakness of mind which renders people incapable of enjoying the pleafures, or performing the duties of life. It is a degree of infanity, and often terminates in abfolute madnefs. CAUSES.——It may proceed from an hereditary disposition; intense thinking, especially where the mind is long occupied about one object; violent paffions or affections of the mind, as love, fear, joy, grief, pride, and fuch like. It may also be occasioned by excessive venery; narcotic or stupefactive poisons; a sedentary life; solitude; the fuppreffion of cuftomary evacuations; acute fevers; or other difeafes. Violent anger will change melancholy into madnefs; and exceffive cold, efpecially of the lower extremities, will force the blood into the brain, and produce all the fymptoms of madnefs. It may likewife proceed from the ufe of aliment that is hard of digeftion, or which cannot be eafily affimilated; from a callous state of the integuments of the brain, or a drynefs of the brain itfelf. To all which we may add gloomy and miftaken notions of religion.

SYMPTOMS. — When perfons begin to be melancholy, they are dull; dejected; timorous; watchful; fond of folitude; fretful; fickle; captious and inquifitive; folicitous about trifles; fometimes niggardly, and other times prodigal. The body is generally bound; the urine thin, and in fmall quantity; the ftomach and bowels inflated with wind; the complexion pale; the pulfe flow and weak. The functions of the mind are alfo greatly perverted, infomuch that the patient often imagines himfelf dead, or changed into fome other animal. Some have imagined their bodies were made of glafs, or other brittle fub-

tincture of valerian each an ounce; mix them; take a tea-fpoonful in a glafs of wine or water three or four times a-day.

OF MELANCHOLY.

stances, and were afraid to move, lest they should be broken to pieces. The unhappy patient, in this cafe, unless carefully watched, is apt to put an end to his own miserable life.

When the difeafe is owing to any obftruction of cultomary evacuations, or any bodily diforder, it is eafier cured than when it proceeds from affections of the mind, or an hereditary taint. A difcharge of blood from the nofe, loofenefs, fcabby eruptions, the bleeding piles, or the *menfes*, fometimes carry off this difeafe.

REGIMEN. — The diet fhould confift chiefly of vegetables of a cooling and opening quality. Animal food, efpecially falted or fmoke-dried fifh or flefh, ought to be avoided. All kinds of fhell-fifh are bad. Aliments prepared with onions, garlic, or any thing that generates thick blood are likewife improper. All kinds of fruits that are wholefome may be eaten with advantage. Boerhaave gives an inflance of a patient who, by a long ufe of whey, water, and garden-fruit, recovered, after having evacuated a great quantity of black-coloured matter.

Strong liquors of every kind ought to be avoided as poifon. The most proper drink is water, whey, or very fmall beer. Tea and coffee are improper. If honey agrees with the patient, it may be eaten freely, or his drink may be fweetened with it. Infusions of balm-leaves, pennyroyal, the roots of wild valerian, or the flowers of the lime-tree, may be drank freely, either by themselves, or fweetened with honey, as the patient shall choofe.

The patient ought to take as much exercise in the open air as he can bear. This helps to diffolve the viscid humours, it removes obstructions, promotes the perspiration, and all the other fecretions. Every kind of madness is attended with a diminished perspiration; all means ought therefore to be used to promote that necessary and falutary discharge. Nothing can have a more direct tendency to increase the disease, than confining the patient to a close apartment. Were he forced to ride or walk a certain number of miles every day, it would tend greatly to alleviate his diforder; but it would have still a better effect, if he were obliged to labour a piece of ground. By digging, hoeing, planting, fowing, &c. both the body D d

40F

and mind would be exercifed. A long journey, or a voyage, efpecially towards a warmer climate, with agreeable companions, has often very happy effects. A plan of this kind, with a ftrict attention to diet, is a much more rational method of cure, than confining the patient within doors, and plying him with medicines.

MEDICINE. --- In the cure of this difeafe, particular attention must be paid to the mind. When the patient is in a low state, his mind ought to be foothed and diverted with variety of amufements, as entertaining ftories, pastimes, mufic, &c. This feems to have been the method of curing melancholy among the Jews, as we learn from the flory of King Saul; and indeed it is a very rational one. Nothing can remove difeafes of the mind fo effectually as applications to the mind itfelf, the molt efficacious of which is mufic. The patient's company ought likewife to confift of fuch perfons as are agreeable to him. People in this flate are apt to conceive unaccountable averfions against particular perfons; and the very fight of fuch perfons is fufficient to diffract their minds, and throw them into the utmost perturbation. In all kinds of madnefs, it is better to foothe and calm the mind, than to ruffle it by contradiction.

When the patient is high, evacuations are neceffary. In this cafe he muft be bled, and have his body kept open by purging medicines, as manna, rhubarb, cream of tartar, or the foluble tartar. I have feen the laft have very happy effects. It may be taken in the dofe of half an ounce, diffolved in water-gruel, every day, for feveral weeks, or even for months, if neceffary. More or lefs may be given according as it operates. Vomits have likewife a good effect; but they muft be pretty ftrong, otherwife they will not operate.

Whatever increases the evacuation of urine or promotes perfpiration, has a tendency to remove this difease. Both these fecretions may be promoted by the use of nitre and vinegar. Half a drachm of purified nitre may be given three or four times a-day in any manner that is most agreeable to the patient; and an ounce and a half of diffilled vinegar may be daily mixed with his drink. Dr. Locker seems to think vinegar the best medicine that can be given in this difease.

Cam-

Camphire and musk have likewife been used in this cafe with advantage. Ten or twelve grains of camphire may be rubbed in a mortar with half a drachm of nitre, and taken twice a-day, or oftener, if the ftomach will bear it. If it will not fit upon the ftomach in this form, it may be made into pills with gum afafætida and Ruffian caftor, and taken in the quantity above directed. musk is to be administered, a scruple or twenty-five grains of it may be made into a bolus with a little honey or common fyrup, and taken twice or thrice a-day. The antimonial wine is by fome extolled for the cure of madnefs; it may be taken in a dofe of forty or fifty drops twice or thrice a-day in a cup of tea. We do not mean that all these medicines should be administered at once; but which-ever of them is given, muft be duly perfifted in, and where one fails another may be tried.

As it is very difficult to induce patients in this difeafe to take medicines, we shall mention a few outward applications which fometimes do good; the principal of these are iffues, fetons, and warm bathing. Iffues may be made in any part of the body, but they generally have the best effect near the spine. The discharge from these may be greatly promoted by dressing them with the mild bliftering ointment, and keeping what are commonly called the orrice-peas in them. The most proper place for a feton is between the shoulder-blades; and it ought to be placed upwards and downwards, or in the direction of the spine.

Madnefs or delirium, which proceeds from mere weaknefs, requires a different treatment. This must be removed by nourifhing diet, exercise proportioned to the patient's strength, and cordial medicines. All evacuations are carefully to be avoided. The patient may take frequently a glass of good wine in which a little Peruvian bark has been infused.

OF THE PALSY.

The palfy is a lofs or diminution of fenfe or motion, or of both, in one or more parts of the body. Of all the affections called nervous, this is the most fuddenly fatal. It is more or lefs dangerous, according to the importance

Dd 2

of

of the part affected. A palfy of the heart, lungs, or any part neceffary to life, is mortal. When it affects the ftomach, the inteffines, or the bladder, it is highly dangerous. If the face be affected, the cafe is bad, as it fhews that the difeafe proceeds from the brain. When the part affected feels cold, is infenfible, or waftes away, or when the judgment and memory begin to fail, there is fmall hope of a cure.

CAUSES.——The immediate caufe of palfy is any thing that prevents the regular exertion of the nervous power upon any particular mufcle or part of the body. The occafional and predifpofing caufes are various, as drunkennefs; wounds of the brain, or fpinal marrow; preffure upon the brain, or nerves; very cold or damp air; the fuppreffion of cuftomaty evacuations; fudden fear; want of exercife; or whatever greatly relaxes the fyftem, as drinking much tea *, or coffee. The palfy may likewife proceed from wounds of the nerves themfelves, from the poifonous fumes of metals or minerals, as mercury, lead, arfenic, &c.

In young perfons of a full habit, the palfy muft be treated in the fame manner as the fanguine apoplexy. The patient muft be bled, bliftered, and have his body opened by fharp clyfters or purgative medicines. But in old age, or when the difeafe proceeds from relaxation or debility, which is generally the cafe, a quite contrary courfe muft be purfued. The diet muft be warm and invigorating, feafoned with fpicy and aromatic vegetables, as muftard, horfe-radifh, &c. The drink may be generous wine, muftard whey, or brandy and water. Friction with the flefh-brufh, or a warm hand, is extremely proper, efpecially on the parts affected. Bliftering-plafters may likewife be applied to the affected parts with advantage. When this cannot be done, they

• Many people imagine, that tea has no tendency to hurt the nerves, and that drinking the fame quantity of warm water would be equally pernicious. This however feems to be a miltake. Many perfons drink three or four cups of warm milk and water daily, without feeling any bad confequences; yet the fame quantity of tea will make their hands fhake for twenty-four hours. That tea affects the nerves, is likewife evident from its preventing fleep, occafioning giddinefs, dimnefs of the fight, ficknefs, &c.

4 -

OF THE PALSY.

may be rubbed with the volatile liniment, or the nerve ointment of the Edinburgh dispensatory. One of the best external applications is electricity. The shocks, or rather vibrations, should be received on the part affected; and they ought daily to be repeated for several weeks.

Vomits are very beneficial in this kind of palfy, and ought frequently to be administered. Cephalic fnuff, or any thing that makes the patient fneeze, is likewife of ufe. Some pretend to have found great benefit from rubbing the parts affected with nettles; but this does not feem to be any way preferable to bliftering. If the tongue be affected, the patient may gargle his mouth frequently with brandy and multard; or he may hold a bit of fugar in his mouth, wet with the palfy-drops, or compound fpirits of lavender. The wild valerian-root is a very proper medicine in this cafe. It may either be taken in an infusion with fage-leaves, or half a drachm of it in powder may be given in a glafs of wine three or four times a-day. If the patient cannot use the valerian, he may take of fal volatile oleofum, compound fpirits of lavender, and tincture of caftor, each half an ounce; mix these together, and take forty or fifty drops in a glass of wine three or four times a-day. A table-fpoonful of muftard-feed taken frequently is a very good medicine. The patient ought likewife to chew cinnamon-bark, ginger, or other warm spiceries.

Exercife is of the utmost importance in the palfy; but the patient must beware of cold, damp, and moist air. He ought to wear flannel next his skin; and, if possible, should remove into a warmer climate.

OF THE EPILEPSY, OR FALLING SICKNESS.

The epilepfy is a fudden deprivation of all the fenfes, wherein the patient falls fuddenly down, and is affected with violent convulfive motions. Children, efpecially thole who are delicately brought up, are most fubject to it. It more frequently attacks men than women, and is very difficult to cure. When the epilepfy attacks children, there is reason to hope it may go off abut the time of puberty. When it attacks any perfon after Dd 3 twenty twenty years of age, the cure is difficult; but when after forty, a cure is hardly to be expected If the fit continues only for a fhort fpace, and returns feldom, there is reafon to hope; but if it continues long, and returns frequently, the profpect is bad. It is a very unfavourable fymptom when the patient is feized with the fits in his fleep.

CAUSES. — The epilepfy is fometimes hereditary. It may likewife proceed from blows, bruifes, or wounds on the head; a collection of water, blood, or ferous humours in the brain; a polypus; tumours or concretions within the fkull; exceffive drinking; intenfe fludy; excefs of venery; worms; teething; fuppreffion of cuftomary evacuations; too great emptinefs or repletion; violent paffions or affections of the mind, as fear, joy, &c.; hyfteric affections; contagion received into the body, as the infection of the fmall-pox, meafles, &c.

SYMPTOMS. — An epileptic fit is generally preceded by unufual wearinefs; pain of the head; dulnefs; giddinefs; noife in the ears; dimnefs of the fight; palpitation of the heart; difturbed fleep; difficult breathing; the bowels are inflated with wind; the urine is in great quantity, but thin; the complexion is pale; the extremities are cold; and the patient often feels, as it were, a flream of cold air afcending towards his head,

In the fit, the patient generally makes an unufual noife; his thumbs are drawn in towards the palms of the hands; his eyes are difforted; he flarts, and foams at the mouth; his extremities are bent or twifted various ways; he often difcharges his feed, urine, and fœces involuntarily; and is quite deltitute of all fenfe and reafon. After the fit is over, his fenfes gradually return, and he complains of a kind of ftupor, wearinefs, and pain of his head; but has no remembrance of what happened to him during the fit.

The fits are fometimes excited by violent affections of the mind, a debauch of liquor, exceflive heat, cold, or the like.

This difeale, from the difficulty of investigating its causes, and its strange symptoms, was formerly attributed to the wrath of the gods, or the agency of evil spirits. In modern times, it has often, by the vulgar, been imputed to witchcraft and fascination. It depends, however,

ever, as much upon natural oaufes as any other malady; and its cure may often be effected by perfifting in the ufe of proper means.

REGIMEN.—Epileptic patients ought, if poffible, to breathe a pure and free air. Their diet fhould be light but nourifhing. They ought to drink nothing ftrong, to avoid fwine's flefh, water-fowl, and likewife all windy and oily vegetables, as cabbage, nuts, &c. They ought to keep themfelves cheerful, carefully guarding against all violent passions, as anger, fear, exceffive joy, and the like.

Exercife is likewife of great ufe; but the patient must be careful to avoid all extremes either of heat or cold, all dangerous fituations, as standing upon precipices, riding, deep waters, and such like.

MEDICINE. ---- The intentions of cure must vary according to the caufe of the difeafe. If the patient be of a fanguine temperament, and there be reafon to fear an obstruction in the brain, bleeding and other evacuations will be neceffary. When the difease is occasioned by the stoppage of customary evacuations, these, if poffible, must be restored; if this cannot be done, others may be fubfituted in their place. Iffues or fetons in this cafe have often a very good effect. When there is reafon to believe that the difease proceeds from worms, proper medicines must be used to kill, or carry off these vermin. When the difease proceeds from teething, the body fhould be kept open by emollient clyfters, the feet frequently bathed in warm water, and, if the fits prove obstinate, a blistering-plaster may be put between the fhoulders. The fame method is to be followed, when epileptic fits precede the eruption of the fmall-pox, or meafles, &c.

When the difeafe is hereditary, or proceeds from a wrong formation of the brain, a cure is not to be expected. When it is owing to a debility, or too great an irritability of the nervous fyftem, fuch medicines as tend to brace and ftrengthen the nerves may be used, as the Peruvian bark, and steel; or the *anti-epileptic* electuaries, recommended by Fuller and Mead *.

* See Appendix, Electuary for the Epilepfy.

The

The flowers of zinc have of late been highly extolled for the cure of the epilepfy. Though this medicine will not be found to answer the expectations which have been raifed concerning it, yet in obstinate epileptic cafes it deferves a trial. The dose is from one to three or four grains, which may be taken either in pills or a bolus, as the patient inclines. The best method is to begin with a single grain four or five times a-day, and gradually to increase the dose as far as the patient can bear it. I have often known this medicine, when duly perfisted in, prove beneficial.

Mufk has fometimes been found to fucceed in the epilepfy. Ten or twelve grains of it, with the fame quantity of factitious cinnabar, may be made up into a bolus, and taken every night and morning.

Sometimes the epilepfy has been cured by electricity.

Convultion-fits proceed from the fame caufes, and must be treated in the fame manner as the epilepfy.

There is one particular species of convulsion-fits which commonly goes by the name of St. Vitus's dance, wherein the patient is agitated with strange motions and gefticulations, which by the common people are generally believed to be the effects of witchcraft. This difease may be cured by repeated bleedings and purges; and afterwards using the medicines prefcribed above for the epileps, viz. the Peruvian bark, and snake-root, &c. Chalybeate waters are found to be beneficial in this cafe. The cold bath is likewise of fingular fervice, and ought never to be neglected when the patient can bear it.

OF THE HICCUP.

The hiccup is a fpafmodic or convultive affection of the ftomach and midriff, arifing from any caufe that irritates their nervous fibres.

It may proceed from excels in eating or drinking; from a hurt in the ftomach; poifons; wind; inflammations or fchirrous tumours of the ftomach, inteftines, bladder, midriff, or the reft of the *vifcera*. In gangrenes, acute and malignant fevers, a hiccup is often the forerunner of death.

When the hiccup proceeds from the use of aliment that is flatulent, or hard of digestion, a draught of generous rous wine, or a dram of any fpirituous liquor, will generally remove it. If poifon be the caufe, plenty of milk and oil muft be drank, as has been formerly recommended. When it proceeds from an inflammation of the ftomach, &c. it is very dangerous. In this cafe the cooling regimen ought to be ftrictly obferved. The patient muft be bled, and take frequently a few drops of the fweet fpirits of nitre in a cup of wine-whey. His ftomach fhould likewife be fomented with cloths dipped in warm water, or have bladders filled with warm milk and water applied to it.

When the hiccup proceeds from a gangrene or mortification, the Peruvian bark, with other antifeptics, are the only medicines which have a chance to fucceed. When it is a primary difeafe, and proceeds from a foul ftomach, loaded either with a pituitous or a bilious humour, a gentle vomit and purge, if the patient be able to bear them, will be of fervice. If it arifes from flatulencies, the carminative medicines directed for the heart-burn muft be ufed.

When the hiccup proves very obftinate, recourfe muft be had to the moft powerful aromatic and antifpafmodic medicines. The principal of thefe is mufk; fifteen or twenty grains of which may be made into a bolus, and repeated occafionally. Opiates are likewife of fervice; but they muft be ufed with caution. A bit of fugar dipped in compound fpirits of lavender, or the volatile aromatic tincture, may be taken frequently. External applications are fometimes alfo beneficial; as the ftomach plafter, or a cataplafm of the Venice treacle of the Edinburgh or London difpenfatory, applied to the region of the ftomach.

I lately attended a patient who had almost a constant hiccup for above nine weeks. It was frequently stopped by the use of musk, opium, wine, and other cordial and antispass medicines, but always returned. Nothing, however, gave the patient fo much ease as brisk smallbeer. By drinking freely of this, the hiccup was often kept off for several days, which was more than could be done by the most powerful medicines. The patient was at length seized with a vomiting of blood, which soon put an end to his life. Upon opening the body, a large state of the patient of the body of the body of the body of the body. fchirrous tumour was found near the pylorus, or right orifice of the stomach.

The hiccup may be removed by taking vinegar; or by a few drops of the oil of vitriol taken in water.

CRAMP OF THE STOMACH.

This difeafe often feizes people fuddenly, is very dangerous, and requires immediate affiftance. It is most incident to perfons in the decline of life, especially the nervous, gouty, hysteric, and hypochondriac.

If the patient has any inclination to vomit, he ought to take fome draughts of warm water, or weak camomile-tea, to cleanfe his ftomach. After this, if he has been coftive, a laxative clyfter may be given. He ought then to take laudanum The beft way of adminiftering it is in a clyfter. Sixty or feventy drops of liquid laudanum may be given in a clyfter of warm water. This is much more certain than laudanum given by the mouth, which is often vomited, and in fome cafes increafes the pain and fpafms in the ftomach.

If the pain and cramps return with great violence, after the effects of the anodyne clyfter are over, another, with an equal or larger quantity of opium, may be given ; and every four or five hours a bolus, with ten or twelve grains of mufk, and half a drachm of the Venice treacle. In the mean time, the ftomach ought to be fomented with cloths dipped in warm water, or bladders filled with warm milk and water fhould be conftantly applied to it. I have often feen thefe produce the moft happy effects. The anodyne balfam may alfo be rubbed on the part affected; and an anti-hyfteric plafter worn upon it for fome time after the cramps are removed, to prevent their return.

In very violent and lasting pains of the stomach, fome blood ought to be let, unless the weakness of the patient forbids it. When the pain or cramps proceed from a suppression of the *menses*, bleeding is of use. If they be owing to the gout, recours must be had to spirits, or fome of the warm cordial waters. Blistering-plasters ought likewise in this case to be applied to the ancles. I have

CRAMP OF THE STOMACH.

have often feen violent cramps and pains of the ftomach removed by covering it with a large plafter of treacle of the London Difpenfatory.

OF THE NIGHT-MARE.

In this difeafe the patient, in time of fleep, imagines he feels an uncommon opprefion of weight about his breaft or ftomach, which he can by no means fhake off. He groans, and fometimes cries out, though oftener he attempts to fpeak in vain. Sometimes he imagines himfelf engaged with an enemy, and in danger of being killed, attempts to run away, but finds he cannot. Sometimes he fancies himfelf in a houfe that is on fire, or that he is in danger of being drowned in a river. He often thinks he is falling over a precipice, and the dread of being dafhed to pieces fuddenly awakes him.

This diforder has been fuppofed to proceed from too much blood; from a ftagnation of blood in the brain, lungs, &c. But it is rather a nervous affection, and arifes chiefly from indigeftion. Hence we find that perfons of weak nerves, who lead a fedentary life, and live full, are most commonly afflicted with the night-mare. Nothing tends more to produce it than heavy fuppers, especially when eaten late, or the patient goes to bed foon after. Wind is likewife a very frequent cause of this difease; for which reason those who are afflicted with it ought to avoid all flatulent food. Deep thought, anxiety, or any thing that oppress the mind, ought also to be avoided.

As perfons afflicted with the night-mare generally moan, or make fome noife in the fit, they fhould be waked, or fpoken to by fuch as hear them, as the uneafinefs generally goes off as foon as the patient is awake. Dr. Whyte fays, he generally found a dram of brandy, taken at bed-time, prevent this difeafe. That, however, is a bad cuftom, and in time lofes its effect. We would rather have the patient depend upon the ufe of food of eafy digettion, cheerfulnefs, exercife through the day, and a light fupper taken early, than to accuitom himfelf to drams. A glafs of peppermint-water will often promote

OF THE NIGHT MARE.

mote digeftion as much as a glass of brandy, and is much fafer. After a person of weak digestion, however, has eaten flatulent food, a dram may be neceffary; in this cafe we would recommend it as the most proper medicine.

Perfons who are young, and full of blood, if troubled with the night-mare, ought to take a purge frequently, and use a fpare diet.

of off SWOONINGS.

break or stomach, which he can by no means fluite off.

People of weak nerves or delicate conftitutions are liable to fwoonings or fainting fits. Thefe indeed are feldom dangerous when duly attended to; but when wholly neglected, or improperly treated, they often prove hurtful, and fometimes fatal.

The general caufes of fwoonings are, fudden tranfition from cold to heat; breathing air that is deprived of its proper fpring or elasticity; great fatigue; exceffive weaknefs; lofs of blood; long fasting; fear, grief, and other violent passions or affections of the mind.

It is well known, that perfons who have been long expoled to cold, often faint or fall into a fwoon, upon coming into the houfe, especially if they drink hot liquor, or fit near a large fire. This might eafily be prevented by people taking care not to go into a warm room immediately after they have been exposed to the cold air, to approach the fire gradually, and not to eat or drink any thing hot, till the body has been gradually brought into a warm temperature.

When any one, in confequence of neglecting thefe precautions, falls into a fwoon, he ought immediately to be removed to a cooler apartment, to have ligatures applied above his knee and elbows, and to have his hands and face fprinkled with vinegar or cold water. He fhould likewife be made to fmell to vinegar, and fhould have a fpoonful or two of water, if he can fwallow, with about a third part of vinegar mixed with it, poured into his mouth. If thefe fhould not remove the complaint, it may be neceffary to bleed the patient, and afterwards to give him a clyfter.

OF SWOONINGS.

As air that is breathed frequently lofes its elafticity or fpring, it is no wonder if perfons who refpire in it often fall into a fwoon or fainting fit. They are in this cafe deprived of the very principle of life. Hence it is that fainting fits are fo frequent in all crowded affemblies, efpecially in hot feafons. Such fits, however, must be confidered as a kind of temporary death; and, to the weak and delicate, they fometimes prove fatal. They ought therefore with the utmost care to be guarded against. The method of doing this is obvious. Let affembly-rooms, and all other places of public refort, be, large and well ventilated; and let the weak and delicate avoid fuch places, particularly in warm feafons.

A perfon who faints in fuch a fituation, ought immediately to be carried into the open air; his temples fhould be rubbed with flrong vinegar or brandy, and volatile fpirits or falts held to his nofe. He fhould be laid upon his back with his head low, and have a little wine or fome other cordial as foon as he is able to fwallow it, poured into his mouth. If the perfon has been fubject to hyfteric fits, caftor or afafœtida fhould be applied to the nofe, or burnt feathers, horn, or leather, &c.

When fainting fits proceed from mere weaknefs or exhauftion, which is often the cafe after great fatigue, long falting, lofs of blood, or the like, the patient must be fupported with generous cordials, as jellies, wines, fpirituous liquors, and fuch like. Thefe, however, muft be given at first in very small quantities, and increased gradually as the patient is able to bear them. He ought to be allowed to lie quite ftill and eafy upon his back, with his head low, and fhould have fresh air admitted into his chamber. His food fhould confift of nourifhing broths, fago-gruel with wine, new milk, and other things of a light and cordial nature. These things are to be given out of the fit. All that can be done in the fit is, to let him fmell to a bottle of Hungary-water, eau de luce, or spirits of hartshorn, and to rub his temples with warm brandy, or to lay a compress dipped in it to the pit of the ftomach.

In fainting fits that proceed from fear, grief, or other violent passions or affections of the mind, the patient must

be very cautioufly managed. He fhould be fuffered to remain at reft, and only made to fmell to fome vinegar. After he is come to himfelf, he may drink freely of warm lemonade, or balm-tea, with fome orange or lemon-peel in it. It will likewife be proper, if the fainting fits have been long and fevere, to clean the bowels by throwing in an emollient clyfter.

It is common in fainting fits, from whatever caufe they proceed, to bleed the patient. This practice may be very proper in ftrong perfons of a full habit; but in those who are weak and delicate, or fubject to nervous diforders, it is dangerous. The proper method with fuch people is, to expose them to the free air, and to use cordial and ftimulating medicines, as volatile falts, Hungary-water, spirits of lavender, tincture of caftor, and the like.

OF FLATULENCIES, OR WIND.

All nervous patients, without exception, are afflicted with wind or flatulencies in the flomach and bowels, which arife chiefly from the want of tone or vigour in thefe organs. Crude flatulent aliment, as green peas, beans, coleworts, cabbages, and fuch like, may increafe this complaint; but flrong and healthy people are feldom troubled with wind, unleis they either overload their flomachs, or drink liquors that are in a fermenting flate, and confequently full of elaftic air. While therefore the matter of flatulence proceeds from our aliments, the caufe which makes air feparate from them in fuch quantity as to occafion complaints, is almost always a fault of the bowels themfelves, which are too weak either to prevent the production of elaftic air, or to expel it after it is produced.

To relieve this complaint, fuch medicines ought to be used as have a tendency to expel wind, and by strengthening the alimentary canal, to prevent its being produced there *.

The

 Many nervous people find great benefit from eating a dry bifcuit, effectially when the ftomach is empty. I look upon this as

OF FLATULENCIES, OR WIND. 415

The lift of medicines for expelling wind is very numerous; they often however difappoint the expectations of both the phyfician and his patient. The most celebrated among the class of carminatives are juniper berries; the roots of ginger and zedoary; the feeds of anife, carraway, and coriander; gum afafœtida and opium; the warm waters, tinctures, and fpirits, as the aromatic water, the tincture of woodfoot, the volatile aromatic fpirit, æther, &c.

Dr. Whyte fays, he found no medicines more efficacious in expelling wind than æther and laudanum. He generally gave the laudanum in a mixture with peppermintwater and tincture of caftor, or fweet fpirits of nitre. Sometimes, in place of this, he gave opium in pills with afafœtida. He obferves that the good effects of opiates are equally confpicuous, whether the flatulence be contained in the ftomach or inteffines; whereas those warm medicines, commonly called *carminatives*, do not often give immediate relief, except when the wind is in the ftomach.

With regard to æther, the Doctor fays, he has often feen very good effects from it in flatulent complaints, where other medicines failed. The dofe is a tea-fpoonful mixed with two table-fpoonfuls of water *. In gouty cafes, he observes, that æther, a glass of French brandy, or of the aromatic water, or ginger, either taken in fubftance, or infused in boiling water, are among the best medicines for expelling wind.

When the cafe of flatulent patients is fuch as makes it improper to give them warm medicines inwardly, the Doctor recommends external applications, which are fometimes of advantage. Equal parts of the anti-hyfteric and ftomach plafter may be fpread upon a piece of foft leather, of fuch fize as to cover the greater part of the belly. This fhould be kept on for a confiderable time, provided the patient be able to bear it; if it fhould give

as one of the beft carminative medicines; and would recommend it in all complaints of the ftomach, arifing from flatulence, indigeftion, &c.

* Though the patient may begin with this quantity, it will be neceffary to increase the dose gradually as the stomach can bear it. Æther is now given in confiderably greater doses than it was in Dr. Whyte's time.

416 OF FLATULENCIES, OR WIND.

great uneafinefs, it may be taken off, and the following liniment used in its stead:

Take of Bates's anodyne balfam an ounce; of the expressed oil of mace half an ounce; oil of mint two drachms. Let these ingredients be mixed together, and about a table-spoonful well rubbed on the parts at bedtime.

For firengthening the flomach and bowels, and confequently for leffening the production of flatulence, the Doctor recommends the Peruvian bark, bitters, chalybeates, and exercife. In flatulent cafes, he thinks fome nutmeg or ginger flould be added to the tincture of the bark and bitters, and that the aromatic powder flould be joined with the filings of iron.

When windy complaints are attended with coffiveness, which is often the cafe, few things will be found to anfwer better than four or five of the following pills taken every night at bed-time :

Take of afafœtida two drachms; fuccotrine aloes, falt of iron, and powdered ginger, of each one drachm; as much of the *elixir proprietatis* as will be fufficient to form them into pills.

On the other hand, when the body is too open, twelve or fifteen grains of rhubarb, with half a drachm, or two fcruples of the Japonic confection, given every other evening, will have very good effects.

In those flatulent complaints which come on about the time the *menses* cease, repeated small bleedings often give more relief than any other remedy.

With regard to diet, the Doctor obferves, that tea, and likewife all flatulent aliments, are to be avoided; and that for drink, water with a little brandy or rum is not only preferable to malt liquor, but in most cafes also to wine.

As Dr. Whyte has paid great attention to this fubject, and as his fentiments upon it in a great measure agree with mine, I have taken the liberty to adopt them; and shall only add to his observations, that exercise is in my opinion superior to all medicine, both for preventing the production and likewise for expelling of flatulencies. These effects, however, are not to be expected from fauntering about, or lolling in a carriage; but from labour,

OF FLATULENCIES, OR WIND. 417

bour, or fuch active amusements as give exercise to every part of the body.

OF LOW SPIRITS.

All who have weak nerves are fubject to low fpirits in a greater or lefs degree. Generous diet, the cold bath, exercife, and amufements, are the most likely means to remove this complaint. It is greatly increased by folitude and indulging gloomy ideas, but may often be relieved by cheerful company and fprightly amufements. When low fpirits are owing to a weak relaxed ftate of the ftomach and bowels, an infusion of the Peruvian

bark with cinnamon or nutmeg will be proper. Steel joined with aromatics may likewife in this cafe be used with advantage; but riding, and a proper diet, are most to be depended on.

When they arife from a foulnefs of the flomach and inteflines, or obftruction in the hypochondriac vifcera, aloetic purges will be proper. I have fometimes known the Harrowgate or Tunbridge water of fervice in this cafe.

When low fpirits proceed from a fupprefion of the menftrual or of the hæmorrhoidal flux, thefe evacuations may either be reftored, or fome other fubftituted in their place, as iffues, fetons, or the like. Dr. Whyte obferves, that nothing has fuch fudden good effects in this cafe as bleeding.

When low fpirits have been brought on by long-continued grief, anxiety, or other diftrefs of mind, agreeable company, variety of amufements, and change of place, especially travelling into foreign countries, will afford the most certain relief.

Perfons afflicted with low fpirits fhould avoid all kinds of excefs, efpecially of venery and ftrong liquors. The moderate use of wine and other ftrong liquors is by no means hurtful; but when taken to excets they weaken the ftomach, vitiate the humours, and deprefs the fpirits. This caution is the more neceffary, as the unfortunate and melancholy often fly to ftrong liquors for relief, by which means they never fail to precipitate their own deftruction.

1 2 10

はある

1.

427

OF HYSTERIC AFFECTIONS.

bour, or fach uttive amplements as give exercite to

These likewise belong to the numerous tribe of nervous diseases, which may be justly reckoned the reproach of medicine. Women of a delicate habit, whose stomach and intestines are relaxed, and whose nervous system is extremely sensible, are most subject to hysteric complaints. In such persons an hysteric fit, as it is called, may be brought on by an irritation of the nerves of the stomach or intestines, by wind, acrid humour, or the like. A sudden suppression of the menses often gives rise to hysteric fits. They may likewise be excited by violent passions or affections of the mind, as fear, grief, anger, or great disappointments.

- Sometimes the hyfteric fit refembles a fwoon or fainting fit, during which the patient lies as in a fleep, only the breathing is fo low as fcarcely to be perceived. At other times the patient is affected with catchings and ftrong convultions. The fymptoms which precede hysteric fits are likewife various in different perfons. Sometimes the fits come on with coldness of the extremities, yawning and ftretching, lownels of spirits, oppression and anxiety. At other times the approach of the fit is foretold by a feeling, as if there were a ball at the lower part of the belly, which gradually rifes towards the ftomach, where it occasions inflation, fickness, and fometimes vomiting; afterwards it rifes into the throat, and occafions a degree of fuffocation, to which quick breathing, palpitation of the heart, giddiness of the head, dimness of the fight, lofs of hearing, with convulfive motions of the extremities and other parts of the body, fucceed. The hysteric paroxysm is often introduced by an immoderate fit of laughter, and fometimes it goes off by crying. Indeed there is not much difference between the laughing and crying of an hyfteric lady.

Our aim in the treatment of this difeafe must be to fhorten the fit or paroxyfm when prefent, and to prevent its return. The longer the fits continue, and the more frequently they return, the difeafe becomes the more obftinate. Their strength is increased by habit, and they induce duce fo great a relaxation of the fystem, that it is with difficulty removed.

It is cuftomary, during the hysteric fit or paroxyfm, to bleed the patient. In ftrong perfons of a plethoric habit, and where the pulfe is full, this may be proper; but in weak and delicate conftitutions, or where the difeafe has been of long ftanding, or arifes from inanition, it is not fafe. The best course in fuch cafes is to rouse the patient by ftrong fmells, as burnt feathers, afafœtida, or fpirits of hartfhorn, held to the nofe. Hot bricks may alfo be applied to the foles of the feet, and the legs, arms, and belly may be ftrongly rubbed with a warm cloth. But the best application is to put the feet and legs into warm water. This is peculiarly proper when the fits precede the flow of the menfes. In cafe of coftivenefs, a laxative clyfter with afafætida will be proper; and as foon as the patient can fwallow, two tablefpoonfuls of a folution of afafætida, or of fome cordial julep, may be given frequently *.

The radical cure of this diforder will be beft attempted at a time when the patient is moft free from the fits. It will be greatly promoted by a proper attention to diet. A milk and vegetable diet, when duly perfifted in, will often perform a cure. If, however, the patient has been accuftomed to a more generous diet, it will not be fafe to leave it off all at once, but by degrees. The moft proper drink is water with a fmall quantity of fpirits. A cool dry air is the beft. Cold bathing and every thing that braces the nerves, and invigorates the fyftem, is beneficial; but lying too long in bed, or whatever relaxes the body, is hurtful. It is of the greateft importance to have the mind kept conftantly eafy and

* When hyfteric fits are occafioned by fympathy, they may be cured by exciting an oppofite paffion. This is faid to have been the cafe of a whole fchool of young ladies in Holland, who were all cured by being told, that the firft who was feized lhould be burnt to death. But this method of cure, to my knowledge, will not always fucceed. I would therefore advife, that young ladies who are fubject to hyfteric fits, fhould not be fent to boarding fchools, as the difeafe may be caught by imitation. I have known madnefs itfelf brought on by fympathy.

Ee 2

cheerful, and, if poffible, to have it always engaged in fome agreeable and interefling purfuit.

The proper medicines are those which ftrengthen the alimentary canal and the whole nervous fystem, as the preparations of iron, the Peruvian bark, and other bitters. Twenty drops of the elixir of vitriol, in a cup of the infusion of the bark, may be taken twice or thrice a-day. The bark and iron may likewife be taken in substance, provided the stomach can bear them; but they are generally given in too stall doses to have any effect. The chalybeate waters generally prove beneficial in this diforder.

If the flomach is loaded with phlegm, vomits will be of ufe; but they fhould not be too flrong, nor frequently repeated, as they tend to relax and weaken the flomach. If there is a tendency to coffiveness, it must be removed either by diet, or by taking an opening pill as often as it shall be found necessfary.

To leffen the irritability of the fyftem, antifpafmodic medicines will be of ufe. The beft antifpafmodic medicines are mufk, opium, and caftor. When opium difagrees with the ftomach, it may either be applied externally, or given in clyfters. It is often fuccefsful in removing those periodical head-aches to which hyfteric and hypochondriac patients are fubject. Caftor has in fome cafes been found to procure fleep where opium failed; for which reafon Dr. Whyte advifes, that they fhould be joined together. He likewife recommends the antihyfteric plafter to be applied to the *abdomen**.

Hysteric women are often afflicted with cramps in various parts of the body, which are most apt to feize them in bed, or when asleep. The most efficacious medicines in this cafe are opium, blistering-plasters, and warm bathing or fomentations. When the cramp or spass is very violent, opium is the remedy most to be depended on. In milder cafes, immersing the feet and legs in warm water, or applying a blistering-plaster to the part affected,

* Though antifpafmodies and anodynes are univerfally recommended in this difeafe, yet all the extraordinary cures that I ever knew in hyfteric cafes, were performed by means of tonic and correborating medicines.

OF HYSTERIC AFFECTIONS.

will often be fufficient to remove the complaint. In patients whofe nerves are uncommonly delicate and fenfible, it will be better to omit the bliftering-plafter, and to attempt the cure by opiates, mufk, camphire, and the warm bath.

Cramps are often prevented or cured by comprefion. Thus cramps in the legs are prevented, and fometimes removed, by tight bandages; and when convultions arife from a flatulent diffention of the inteffines, or from fpafms beginning in them, they may be often leffened or cured by making a pretty ftrong comprefion upon the *abdomen* by means of a broad belt. A roll of brimftone held in the hand is frequently ufed as a remedy for cramps. Though this feems to owe its effect chiefly to imagination, yet, as it fometimes fucceeds, it merits a trial *. When fpafms or convultive motions arife from fharp humours in the ftomach or inteffines, no lafting relief can be procured till thefe are either corrected or expelled. The Peruvian bark has fometimes cured periodic convulfions after other medicines had failed.

OF HYPOCHONDRIAC AFFECTIONS.

This difeafe generally attacks the indolent, the luxurious, the unfortunate, and the fludious. It becomes daily more common in this country, owing, no doubt, to the increafe of luxury and fedentary employments. It has fo near a refemblance to the immediately preceding, that many authors confider them as the fame difeafe, and treat them accordingly. They require, however, a very different regimen; and the fymptoms of the latter, though lefs violent, are more permanent than those of the former.

Men of a melancholy temperament, whofe minds are capable of great attention, and whofe paffions are not eafily moved, are, in the advanced periods of life, most liable to this difease. It is usually brought on by long and ferious attention to abstruct subjects, grief, the sup-

* Some perfons afflicted with cramps pretend to reap great benefit from fmall bundles of rofemary tied all night about their feet, ancles, and knees.

preflion

421

preffion of cuftomary evacuations, excefs of venery, the repulfion of cutaneous eruptions, long continued evacuations, obstructions in fome of the viscera, as the liver, spleen, &c.

Hypochondriac perfons ought never to fast long, and their food should be folid and nourishing. All ascessent and windy vegetables are to be avoided. Flesh meats agree best with them, and their drink should be old claret or good madeira. Should these disagree with the stomach, water with a little brandy or rum in it may be drank.

Cheerfulnefs and ferenity of mind are by all means to be cultivated. Exercise of every kind is useful. The cold bath is likewise beneficial; and, where it does not agree with the patient, frictions with the flesh-brush or a coarse cloth may be tried. If the patient has it in his power, he ought to travel either by sea or land. A voyage or a long journey, especially towards a warmer climate, will be of more fervice than any medicine.

The general intentions of cure in this difeafe, are to firengthen the alimentary canal, and to promote the fecretions. These intentions will be best answered by the different preparations of iron and the Peruvian bark, which, after proper evacuations, may be taken in the fame manner as directed in the preceding difease.

If the patient be coffive, it will be neceffary to make use of fome gentle opening medicine, as pills composed of equal parts of aloes, rhubarb, and afafœtida, with as much of the elixir proprietatis as is neceffary to form the ingredients into pills. Two, three, or four of these may be taken as often as it shall be found needful to keep the body gently open. Such as cannot bear the afafœtida may substitute Spanish foap in its place.

Though a cheerful glafs may have good effects in this difeafe, yet all manner of excefs is hurtful. Intenfe ftudy, and every thing that depreffes the fpirits, are likewife pernicious.

Though the general fymptoms and treatment of nervous diforders were pointed out in the beginning of this chapter; yet, for the benefit of the unhappy perfons afflicted with those obstinate and complicated maladies, I have

I have treated feveral of their capital fymptoms under diftinct or feparate heads. Thefe, however, are not to be confidered as different difeafes, but as various modifications of the fame difeafe. They all arife from the fame general caufes, and require nearly the fame method of treatment. There are many other fymptoms that merit particular attention, which the nature of my plan will not permit me to treat of at full length. I fhall therefore omit them altogether, and conclude this chapter with a few general remarks on the molt obvious means of preventing or avoiding nervous diforders.

In all perfons afflicted with nervous diforders, there is a great delicacy and fenfibility of the whole fyftem, and an uncommon degree of weaknefs of the organs of digeftion. Thefe may be either natural or acquired. When owing to a defect in the conftitution, they are hardly to be removed; but may be mitigated by proper care. When induced by difeafes, as long or repeated fevers, profuse hæmorrhages, or the like, they prove alfo very obstinate, and will yield only to a courfe of regimen calculated to reftore and invigorate the habit.

But nervous affections arife more frequently from caufes, which it is in a great meafure in our own power to avoid, than from difeafes, or an original fault in the conflitution, &c. Exceflive grief, intenfe fludy, improper diet, and neglect of exercife, are the great fources of this extensive class of difeafes.

It has been already obferved, that grief indulged deftroys the appetite and digeftion, deprefies the fpirits; and induces an univerfal relaxation and debility of the whole fyftem. Inftances of this are daily to be feen. The lofs of a near relation, or any other misfortune in life, is often fufficient to occafion the most complicated feries of nervous fymptoms. Such misfortunes indeed are not to be avoided, but furely their effects, by a vigorous and proper exertion of the mind, might be rendered lefs hurtful. For directions in this matter we must refer the reader to the article GRIEF, in the chapter on the Paffions.

The effects of intense study are pretty similar to those occasioned by grief. It preys upon the animal spirits,

Ee4

and

and deftroys the appetite and digeftion. To prevent these effects, studious perfons ought, according to the Poet, to toy with their books*. They should never study too long at a time; nor attend long to one particular subject, especially if it be of a serious nature. They ought likewise to be attentive to their posture, and should take care frequently to unbend their minds by music, diversions, or going into agreeable company.

With regard to diet, I shall only observe, that nervous diseases may be induced either by excess or inanition. Both of these extremes hurt the digestion, and vitiate the humours. When Nature is oppressed with fresh loads of food, before she has had time to digest and affimilate the former meal, her powers are weakened, and the vessels are filled with crude humours. On the other hand, when the food is not fufficiently nouriss, or is taken too feldom, the bowels are inflated with wind, and the humours, for want of regular fresh supplies of wholesome chyle, are vitiated. These extremes are therefore with equal care to be avoided. They both tend to induce a relaxation and debility of the nervous system, with all its dreadful train of confequences.

But the most general cause of nervous diforders is indolence. The active and laborious are feldom troubled with them. They are referved for the children of ease and affluence, who generally feel their keenest force. All we shall fay to such performs is, that the means of prevention and cure are both in their own power. If the constitution of human nature be such, that man must either labour or suffer difeases, furely no individual has any right to expect an exemption from the general rule.

Those, however, who are willing to take exercise, but whose occupations confine them to the house, and perhaps to an unfavourable posture, really deferve our pity. We have in a former part of the book endeavoured to lay down rules for their conduct; and shall

* Armftrong on Health.

only add, that where these cannot be complied with, their place may, in some measure, be supplied by the use of bracing and strengthening medicines, as the Peruvian bark, with other bitters; the preparations of steel; the elixir of vitriol, and such like.

Among many remarkable cafes of the nervous kind, which I have often met with, one very lately attracted my notice in a peculiar manner. It was written by the patient himfelf, a gentleman of fortune and of liberal education; and it might be justly called a picture from nature, drawn with uncommon fentibility and force. The whole account being too long for infertion, the following extract may ferve as a fpecimen of the writer's fufferings and defcriptive talents. " It is in vain," he fays, " that I attempt to impress the Faculty with the real state of my fufferings. The fymptoms of the diforders are not to be defcribed, from their unufual preffure upon the mind; nor can they be conceived, I believe, by any but those who have fuffered under them. They may be faid to conflitute a phenomenon in the science of difeases. Since I know of no terms to express them in, or language to describe them by, I am obliged to content myfelf with denominating the diforder and its effects together a mental agony, whole influence creates a real tedium vitæ. It attacks me fometimes when fitting, fometimes when walking; and if I were not to throw myfelf on a bed during the violence of the paroxyfm, I fhould certainly dafh myfelf to pieces. This is accompanied with a laffitude, reftleffnefs, and total incapacity of attending to any concerns in life."

The fame fpirit animated every part of the affecting defcription; and the cafe was accompanied with a lift of eleven eminent phyficians, whom the patient had confulted at different times, but whofe names I fupprefs, as their prefcriptions did him no good, and did them no honour. When the primary feat of the difeafe is in the mind, it is ftooping to the low tricks of quack-ry to amufe a patient with falfe hopes of the efficacy of any medicine. The difappointment that follows aggravates every

every painful fymptom, and makes the unhappy fufferer look forward to death as the only refource. All I prefcribe for him, is travelling.

I fhould also have willingly inferted here an account of fome other nervous affections of an extraordinary nature, had not their length exceeded the limits I prefcribed to myself in these supplementary observations.

my notice in a peculiar manner. It was written

CHAP. XLIV.

adation; and it might be juffly called a picture from

DISORDERS OF THE SENSES.

WE do not mean to treat of the nature of our fenfations, or to give a minute defcription of the various organs by which they are performed; but to point out fome of the difeafes to which these organs are most liable, and to shew how they may be prevented or remedied.

OF THE EYE.

No organ of the body is fubject to more difeafes than the eye; nor is there any one of which the difeafes are more difficult to cure. Though more ignorant perfons pretend to cure these than any other class of difeafes, yet a very fuperficial acquaintance with the ftructure of the eye, and the nature of vision, will be fufficient to convince any one of the danger of truffing to them. These difeases often exceed the skill of the most learned phyfician; hence we may eafily infer the danger of truffing them to ignorant quacks, who, without all peradventure, put out more eyes than they cure. But, though the difeafes of the eye can feldom be cured, they might often, by due care, be prevented; and, even where the fight is totally loft, many things might be done, which are generally neglected, to render the unhappy perfon both more ufeful to himfelf and to fociety *.

* It is a pity those who have the misfortune to be born blind, or who lose their fight when young, should be suffered to remain in ignorance The eyes are hurt by viewing bright or luminous objects; keeping the head too long in a hanging pofture; violent head-achs, exceflive venery; the long ufe of bitters; the effluvia from acrid or volatile fubftances; various difeafes; as the fmall-pox, meafles, &c.; but, above all, from night-watching, and candle-light ftudies. Long fafting is likewife hurtful to the eyes, and frequent heats and colds are no lefs pernicious. The eyes are often hurt by the ftoppage of cuftomary evacuations; as morning fweats; fweating of the feet; the menfes in women; and the bleeding piles in men. All kinds of excefs are likewife hurtful to the fight, particularly the immoderate ufe of ardent fpirits, and other ftrong liquors.

In all difeafes of the eyes, efpecially those attended with inflammation, the cold regimen ought to be obferved. The patient muft abstain from all spirituous liquors. The smoke of tobacco, smoky rooms, the vapours of onions and garlic, and all vivid lights and glaring colours, are carefully to be avoided. The drink may be water, whey, or small beer; and the aliment must be light and of eafy digestion.

For preventing diforders of the eyes, iffues and fetons are of prime ufe. Every perfon, whofe eyes are tender, ought to have one or more of thefe in fome part of the body. It will likewife be of ufe to keep the body gently open, and either to bleed or purge every fpring and fall. All excefs and night-ftudies are to be avoided. Such as do not choofe a feton or an iffue, will reap benefit from wearing a fmall Burgundy-pitch plafter between their fhoulders.

ignorance or to beg. This is both cruelty and want of economy. There are many employments of which blind perfons are very capable, as knitting, carding, turning a wheel, teaching languages, &c. Nor are initances wanting of perfons who have arrived at the higheft pitch of learning, without having the leaft idea of light. Witness the late famous Nicholas Sanderfon of Cambridge, and my worthy friend Dr. Thomas Blacklock of Edinburgh. The former was one of the first mathematicians of his age, and the latter, besides being a good poet and philosopher, was master of all the learned languages, and a very confiderable adept in the liberal arts.

7

A gutta,

A gutta serena, or amaurofis, is an abolition of the fight, without any apparent caufe or fault in the eyes. When it is owing to a decay or wasting of the optic nerve, it does not admit of a cure; but when it proceeds from a compression of the nerves, by redundant humours, thefe may in fome meafure be drained off, and the patient relieved. For this purpose, the body must be kept open with the laxative mercurial pills. If the patient be young, and of a fanguine habit, he may be bled. Cupping, with fcarifications on the back part of the head, will likewife be of ufe. A running at the nofe may be promoted by volatile falts, ftimulating powders, &c. But the most likely means for relieving the patient are iffues or blifters kept open for a long time on the back part of the head, behind the ears, or on the neck. I have known these restore fight, even after it had been for a confiderable time loft.

Should thefe fail, recourfe must be had to a mercurial falivation; or, what will perhaps answer the purpose better, twelve grains of the corrosive sublimate of mercury may be diffolved in an English pint and a half of brandy, and a table-spoonful of it taken twice a-day, drinking half a pint of the decoction of sarfaparilla after it.

A cataract is an obftruction of the pupil, by the interpofition of fome opaque fubftance which either diminifhes or totally extinguifhes the fight. It is generally an opacity of the cryftalline humour. In a recent or beginning cataract, the fame medicines are to be ufed as in the gutta ferena; and they will fometimes fucceed. But when this does not happen, and the cataract becomes firm, it must be couched, or rather extracted. I have refolved a recent cataract by giving the patient frequent purges with calomel, keeping a poultice of fresh hemlock constantly upon the eye, and a perpetual blifter on the neck *.

The myopia, or *fbort fightednefs*, and the *prefbyopia*, or *feeing only at too great a distance*, are diforders which depend on the original structure or figure of the eye, there-

* In both these cases electricity merits a trial.

fore,

OF THE EYE.

fore, admit of no cure. The inconveniences arifing from them may however be in fome measure remedied by the help of proper glaffes. The former requires the aid of a concave, and the latter of a convex glafs.

A *ftrabifmus*, or *fquinting*, depends upon an irregular contraction of the mufcles of the eye from a fpafm, palfy, epilepfy, or an ill habit. Children often contract this diforder by having their eyes unequally exposed to the light. They may likewife acquire it by imitation from a fquinting nurfe, or play-fellow, &c. As this diforder can hardly be cured, parents ought to be careful to prevent it. Almost the only thing which can be done for it is to contrive a mask for the child to wear, which will only permit him to fee in a straight direction.

Spots or fpecks on the eyes are generally the effect of inflammation, and often appear after the fmall-pox; the meafles, or violent opthalmias. They are very difficult to cure, and often occafion total blindnefs. If the fpecks are foft and thin, they may fometimes be taken off by gentle cauftics and difcutients; as vitriol, the juice of celandine, &c. When thefe do not fucceed, a furgical operation may be tried: the fuccefs of this, however, is always very doubtful.

The blood floot eye may be occasioned by a stroke, a fall, retching, vomiting, violent coughing, &c. I have frequently known it happen to children in the hoopingcough. It appears at first like a bit of scarlet, and is afterwards of a livid or blackish colour. This disorder generally goes off without medicine. Should it prove obstinate, the patient may be bled, and have his eyes fomented with a decostion of comphry roots and elder flowers. A fost poultice may be applied to the eyes; and the body should be kept open by gentle purgatives.

The watery or weeping eye is generally occafioned by a relaxation or weaknels of the glandular parts of that organ. Thefe may be braced and ftrengthened by bathing the eye with brandy and water, Hungary-water, rofe-water, with white vitriol diffolved in it, &c. Medicines which make a revulfion are likewife proper; as mild purgatives, perpetual blifters on the neck, bathing the feet frequently in lukewarm water, &c.

When

OF THE EYE.

When this difease proceeds from an obstruction of the lachrymal duct, or natural passage of the tears, it is called a *fistula lachrymalis*, and can only be cured by a surgical operation *.

OF THE EAR.

The functions of the ear may be injured by wounds, ulcers, or any thing that hurts its fabric. The hearing may likewife be hurt by exceffive noife, violent colds in the head; fevers; hard wax, or other fubflances flicking in the cavity of the ear; too great a degree of moifture or drynefs of the ear. Deafnefs is very often the effect of old age, and is incident to most people in the decline of life. Sometimes it is owing to an original fault in the flructure or formation of the ear itself. When this is the cafe, it admits of no cure; and the unhappy perfon not only continues deaf, but generally likewife dumb for life t.

When

A weeping or watery eye is often the mark of a fcrophulous habit.

+ Though those who have the misfortune to be born deaf are generally fuffered to continue dumb, and confequently are in a great measure loft to fociety, yet nothing is more certain than that fuch perfons may be taught not only to read and write, but alfo to fpeak, and to understand what others fay to them. Teaching the dumb to fpeak, will appear parodoxical to those who do not confider that the formation of founds is merely mechanical, and may be taught without the affiftance of the ear. This is not only capable of demonstration, but is actually reduced to practice by the ingenious Mr. Thomas Braidwood of Edinburgh. This gentleman has, by the mere force of genius and application, brought the teaching of dumb perfons to fuch a degree of perfection, that his fcholars are generally more forward in their education, than those of the same age who enjoy all their faculties. They not only read and write with the utmost readiness, but likewife speak, and are capable of holding conversation with any perfon in the light. What a pity any of the human fpecies fhould remain in a state of idiotism, who are capable of being rendered as useful and intelligent as others! We mention this not only from humanity to those who have the misfortune to be born deaf, but also in juffice to Mr. Braidwood, whofe fuccels has far exceeded all former attempts this way; and indeed it exceeds imagination itfelf fo far, that no perfon who has not feen and examined his pupils, can believe what they are capable of .- As this gentleman,

When deafnefs is the effect of wounds or ulcers of the ear, or of old age, it is not eafily removed. When it proceeds from cold of the head, the patient muft be careful to keep his head warm, efpecially in the night; he fhould likewife take fome gentle purges, and keep his feet warm, and bathe them frequently in lukewarm water at bed-time. When deafnefs is the effect of a fever it generally goes off after the patient recovers. If it proceeds from dry wax flicking in the ears, it may be foftened by dropping oil into them; afterwards they muft be fyringed with warm milk and water.

If deafnels proceed from drynels of the ears, which may be known by looking into them, half an ounce of the oil of fweet almonds, and the fame quantity of liquid opodeldoch, or tincture of afafeetida, may be mixed together, and a few drops of it put into the ear every night at bed-time, ftopping them afterwards with a little wool or cotton. Some, inftead of oil, put a fmall flice of the fat of bacon into each ear, which is faid to anfwer the purpofe very well. When the ears abound with moifture, it may be drained off by an iffue or feton, which fhould be made as near the affected parts as poffible.

Some, for the cure of deafnels, recommend the gall of an eel mixed with fpirit of wine, to be dropped into the ear; others, equal parts of Hungary-water and fpirit of lavender. Etmuller extols amber and mulk; and Brookes fays, he has often known hardnels of hearing cured by putting a grain or two of mulk into the ear with cotton-wool. But thele and other applications mult be varied according to the caule of the diforder *.

man, however willing, is only able to teach a few, and as the far greater part of those who are born deaf cannot afford to attend him, it would be an act of great humanity, as well as public utility, to creft an academy for their benefit:

• A gentleman, on whole veracity I can depend, told me, that after using many things to no purpole for an oblinate deafnels, he was at lalt advised to put a few drops of his own urine warm into his ears every night and morning, from which he received great benefit. It is probable that a folution of *fal ammoniac*, in water, would produce the fame effect.

Though

Though fuch applications may fometimes be of fervice, yet they much oftener fail, and frequently they do hurt. Neither the eyes nor ears ought to be tampered with; they are tender organs, and require a very delicate touch. For this reafon, what we would chiefly recommend in deafnefs, is, to keep the head warm. From whatever caufe the diforder proceeds, this is always proper; and I have known more benefit from it alone, in the most obstinate cafes of deafnefs, than from all the medicines I ever ufed *.

OF THE TASTE AND SMELL.

Though these series are not of forgreat importance to man in a flate of fociety, as the fight and hearing, yet as the loss of them is attended with fome inconveniency, they deferve our notice. They are feldom to be reflored when lost; which ought to make us very attentive to their prefervation, by carefully avoiding whatever may in the least prove injurious to them. As there is a very great affinity between the organs of tasting and fmelling, whatever hurts the one generally affects the other.

Luxury is highly injurious to these organs. When the nose and palate are frequently stimulated by fragrant and poignant distes, they soon lose the power of distinguishing tastes and odours with any degree of nicety. Man, in a state of nature, may perhaps have these faculties as acute as any other animal.

The fenfe of fmelling may be diminifhed or deftroyed by difeafes; as the moifture, drynefs, inflammation, or fuppuration of that membrane, which lines the infide of the nofe, commonly called the olfactory membrane; the compression of the nerves which fupply this membrane, or fome fault in the brain itself at their origin. A defect, or too great a degree of folidity, of the small fpungy bones of the upper jaw, the caverns of the forehead, &c. may likewife impair the fense of smalling. It may also be injured by a collection of foetid matter in those caverns, which keeps constantly exhaling from

An obftinate deafnefs has been cured by electricity.

3

OF THE TASTE AND SMELL.

them. Few things are more hurtful to the fense of fmelling than taking great quantities of fnuff.

When the nofe abounds with moifture, after gentle evacuations, fuch things as tend to take off irritation, and coagulate the thin fharp ferum, may be applied; as the oil of annifeed mixed with fine flour; camphire diffolved in oil of almonds, &c. The vapours of amber, frankincenfe, gum-maftic, and benjamin, may likewife be received into the nofe and mouth.

For moiftening the mucus, when it is too dry, fome recommend fnuff made of the leaves of marjoram, mixed with the oil of amber, marjoram, and annifeed; or a fternutatory of calcined white vitriol; twelve grains of which may be mixed with two ounces of marjoramwater, and filtrated. The fteam or vapour of vinegar upon hot iron received up the noftrils is likewife of ufe for foftening the mucus, opening obftructions, &c.

If there is an ulcer in the nofe, it ought to be dreffed with fome emollient ointment, to which, if the pain be very great, a little laudanum may be added. If it be a venereal ulcer, it is not to be cured without mercury. In that cafe, the folution of the corrofive fublimate in brandy may be taken, as directed in the gutta ferena. The ulcer ought likewife to be wafhed with it; and the fumes of cinnabar may be received up the noftrils.

If there be reafon to fufpect that the nerves, which fupply the organs of fmelling are inert, or want ftimulating, volatile falts, ftrong fnuffs, and other things which occasion fneezing, may be applied to the nofe. The forehead may likewife be anointed with balfam of Peru, to which may be added a little of the oil of amber.

The *tafte* may be diminished by crufts, filth, mucus, apthæ, pellicles, warts, &c. covering the tongue. It may be depraved by a fault of the faliva, which, being discharged into the mouth, gives the fame fensations as if the food which the person takes had really a bad taste; or it may be entirely destroyed by injuries done to the nerves of the tongue and palate. Few things prove more hurtful either to the fense of tasting or fmelling,

than

433

434 OF THE TASTE AND SMELL.

than obstinate colds, especially those which affect the head.

When the tafte is diminifhed by filth, mucus, &c. the tongue ought to be fcraped, and frequently wafhed with a mixture of water, vinegar, and honey, or fome other detergent. When the faliva is vitiated, which feldom happens, unlefs in fevers or other difeafes, the curing of the diforder is the cure of this fymptom. To relieve it, however, in the mean time, the following things may be of ufe: If there be a bitter tafte, it may be taken away by vomits, purges, and other things which evacuate bile. What is called a nidorous tafte, arifing from putrid humours, is corrected by the juice of citrons, oranges, and other acids. A falt tafte is cured by a plentiful dilution with watery liquors. An acid tafte is deftroyed by abforbents, and alkaline falts, as powder of oyfter-fhells, falt of wormwood, &c.

When the fenfibility of the nerves, which fupply the organs of tafte, is diminished, the chewing of horferadish, or other stimulating substances, will help to recover it.

OF THE TOUCH.

The fenfe of touching may be hurt by any thing that obftructs the nervous influence, or prevents its being regularly conveyed to the organs of touching; as preffure, extreme cold, &c. It may likewife be hurt by too great a degree of fenfibility, when the nerve is not fufficiently covered by the cuticle or fcarf-fkin, or where there is too great a tenfion of it, or it is too delicate. Whatever diforders the functions of the brain and nerves, hurts the fenfe of touching. Hence it appears to proceed from the fame general caufes as palfy and apoplexy, and requires nearly the fame mode of treatment.

In a *ftupor*, or defect of touching, which arifes from an obstruction of the cutaneous nerves, the patient must first be purged; afterwards such medicines as excite the action of the nerves, or stimulate the system, may be used.

OF THE TOUCH.

ufed. For this purpofe, the fpirit of hartfhorn, fal volatile oleofum, horfe-radifh, &c. may be taken inwardly; the difordered parts, at the fame time, may be frequently rubbed with frefh nettles, or fpirit of fal ammoniac. Bliftering-plafters and finapifms applied to the parts will likewife be of ufe, as also warm bathing, especially in the natural hot baths.

In a work like this, which is wholly defigned for popular inftruction, it would have been an ufelefs difplay of anatomical fkill to mention fuch diforders of the fenfes as admit of no remedy, becaufe they are owing to a defect in the organization or ftructure of the brain, whence the nerves, those fine organs of fenfation, take their rife. But it may be proper to make a few remarks on one or two general caufes of nervous weaknefs, and of confequent debility or imperfection of the fenfes, which proceed wholly from our own mifconduct.

Nothing fo much relaxes the nervous fyftem, fo much blunts the acuteness of every fense, and deftroys its energy, as intemperance. To fay of a man when drunk, that he has lost his fenses, is literally true in the most comprehensive meaning of the word. He can neither fee, hear, taste, fmell, nor feel, with exactness; and though he may flatter himfelf, that, with the return of fobriety, he recovers his fenfes alfo, yet they become more and more impaired by every debauch, till frequent repetitions of the frantic indulgence confign him to blindnefs, to deafnefs, and to the grave. Excefs in eating produces fimilar effects, and, like the touch of the torpedo, benumbs every faculty. It particularly vitiates the tafte and fmell, and thus defeats the chief purpoles for which these sere given, to inform us of the wholefome or noxious properties of every thing we eat and drink.

Uncleannefs is alfo highly injurious to the organs of fenfation. Perhaps the benignity of Nature is not difplayed in any thing more ftrongly than in the warnings fhe gives of this evil, and in her own endeavours to avert it. She has left us fo little to do, that we delerve no pity for the feverest punishment of our neglect. See how kindly she has guarded the extremities of the nerves

435

all over the body, the interior parts of the nose, the mouth, the ear, the eye, against external annoyance! Observe with what efforts, entirely independent of our will, the ftrives to relieve those delicate organs from all impurities! The uneafinels we feel upon fuch occafions ought to roufe our immediate attention. Shall we fuffer dirt to gather upon the skin, to dull the sense of feeling, to obstruct the pores, and to drive back into the fystem the noxious particles which Nature endeavours to throw off, when the use of a little foap and water would prevent every inconvenience? Is it too much trouble to wash the ears; to dip the face with the eyes open in a bason of clean water four or five times every morning; to rinle the nofe and mouth; and to keep the tongue clean, not by fcraping it, but by attending to the flate of the flomach, of which the tongue is an index? Some people feem to be as much afraid of water as if they had been bitten by a mad dog; and if they remain obstinate in that antipathy, I can only fay, they deferve a far worfe end than that of fuch unfortunate incurables, to be fuffocated in their own filth.

CHAP. XLV.

recencts his factor allowyet

In mich march have

they fact hears to Re. Intell

OF A SCHIRRUS AND CANCER.

A SCHIRRUS is a hard indolent tumour, ufually feated in fome of the glands; as the breafts, the arm-pits, &c. If the tumour becomes large, unequal, of a livid, blackifh, or leaden colour, and is attended with violent pain, it gets the name of an occult cancer. When the fkin is broken, and a *fanies* or ichorous matter of an abominable foetid fmell is difcharged from the fore, it is called an open or ulcerated cancer. Perfons after the age of forty-five, particularly women, and thofe who lead an indolent fedentary life, are most fubject to this difeafe.

CAUSES.

436

CAUSES .---- This difeafe is often owing to fuppreffed evacuations; hence it proves fo frequently fatal to women of a grofs habit, particularly old maids and widows, about the time when the menstrual flux ceases. It may likewife be occafioned by excellive fear, grief, anger, religious melancholy, or any of the depreffing paffions. Hence the unfortunate, the choleric, and those perfons who devote themfelves to a religious life in convents or monasteries, are often afflicted with it. It may also be occasioned by the long-continued use of food that is too hard of digeftion, or of an acrid nature; by barrennefs, celibacy, indolence, cold, blows, friction, preffure, or the like. Women often fuffer from the last of these by means of their stays, which fqueeze and compress their breast fo as to occasion great mischief. Sometimes the difease is owing to an hereditary disposition.

SYMPTOMS. ——— This diforder feems often very trifling at the beginning. A hard tumour, about the fize of an hazel-nut, or perhaps finaller, is generally the first fymptom. This will often continue for a long time without feeming to increase, or giving the patient great uneafinefs; but if the conftitution be hurt, or the tumour irritated by preffure or improper treatment of any kind, it begins to extend itfelf towards the neighbouring parts, by puffing out a kind of roots or limbs. It then gets the name of *cancer*, from a fancied refemblance between these limbs and the claws of a crab. The colour of the fkin begins to change, which is first red, afterwards purple, then bluish, livid, and at last black. The patient complains of heat, with a burning, gnawing, fhooting pain. The tumour is very hard, rough, and unequal, with a protuberance, or rifing, in the middle; its fize increases daily, and the neighbouring veins become, thick, knotty, and of a blackifh colour.

The fkin at length gives way, and a thin fharp ichor begins to flow, which corrodes the neighbouring parts till it forms a large unfightly ulcer. More occult cancers arife, and communicate with the neighbouring glands. The pain and ftench become intolerable; the appetite fails; the ftrength is exhausted by a continual Ff_3 hectic

hectic fever; at last, a violent hæmorrhage, or discharge of blood, from some part of the body, with faintings or convulsion fits, generally put an end to the miserable patient's life.

REGIMEN. — The diet ought to be light, but nourifhing. All ftrong liquors, and high-feafoned or falted provifions, are to be avoided. The patient may take as much exercife as he can eafily bear; and fhould ufe every method to divert thought, and amufe his fancy. All kinds of external injury are carefully to be guarded againft, particularly of the affected part, which ought to be defended from all preffure, and even from the external air, by covering it with fur or foft flannel.

When a fchirrus tumour is first difcovered, the patient ought to obferve a proper regimen, and to take twice or thrice a-week a dofe of the common purging mercurial pill. Some blood may alfo be let, and the part affected may be gently rubbed twice a-day, with a little of the mercurial ointment, and kept warm with fur or flannel. The food must be light, and an English pint of the decoction of woods or farfaparilla may be drank daily. I have fometimes difcuffed hard tumours, which had the appearance of beginning cancers, by a courfe of this kind.

Should the tumour, however, not yield to this treatment, but, on the contrary, become larger and harder, it will be proper to extirpate it, either by the knife or cauftic. Indeed, whenever this can be done with fafety, the fooner it is done the better. It can anfwer no purpofe to extirpate a cancer after the conflictution is ruined,

10

or the whole mass of humours corrupted by it. This, however, is the common way, which makes the operation fo feldom fucceed. Few people will submit to the extirpation till death states them in the face; whereas, if it were done early, the patient's life would not be endangered by the operation, and it would generally prove a radical cure.

When the cancer is fo fituated that it cannot be cut off, or if the patient will not fubmit to the operation, fuch medicines as will mitigate or relieve the most urgent fymptoms may be used. Dr. Home fays, that half a grain of the corrosive fublimate of mercury, diffolved in a proper quantity of brandy, and taken night and morning, will often be of service in cancers of the face and nofe. He likewise recommends an infusion of the *fola*num, or night-fhade, in cancers of the breafts.

But the medicine most in repute at prefent for this difeafe is hemlock. Dr. Stock, physician at Vienna, has of late recommended the extract of this plant as very efficacious in cancers of every kind. The Doctor fays, he has given fome hundred weights of it without ever hurting any body, and often with manifest advantage. He advifes the patient, however, to begin with very fmall dofes, as two or three grains, and to increase the dose gradually till fome good effect be perceived, and there to reft without further increase. From two or three grains at first, the Doctor fays he has increased the dose to two, three, or four drachms a-day, and finds that fuch doses may be continued for feveral weeks without any bad confequences.

The regimen which the Doctor recommends during the ufe of the medicine, is to avoid farinaceous fubftances not fermented, and too acrid aromatics. He fays, good wine will not be hurtful to those who are accustomed to it, nor a moderate use of acids; and adds, that the patient should live in a pure free air, and keep his mind as quiet and cheerful as possible.

The Doctor does not pretend to fix the time in which a cancer may be refolved by the ufe of hemlock, but fays he has given it for above two years in large dofes without any apparent benefit; neverthelefs the patient has been Ff 4 cured

cured by perfifting in the ufe of it for half a year longer. This is at leaft encouragement to give it a fair trial. Though we are far from thinking the hemlock merits those extravagant encomiums which the Doctor has beflowed upon it, yet, in a difease which has so long baffled the boasted powers of medicine, we think it ought always to be tried.

The powder of hemlock is by fome preferred to the extract. They are both made of the fresh leaves, and may be used nearly in the same manner. Dr. Nicholson of Berwick says, he gradually increased the dose of the powder from a few grains to half a drachm, and gave near four drachms of it in the day with remarkably good effects. The hemlock may also be used externally either as a poultice or fomentation. The fore may likewise be kept clean by injecting daily a strong decosion of the tops and leaves into it.

Few things contribute more to the healing of foul fordid ulcers of any kind than keeping them thoroughly clean. This ought never to be neglected. The beft application for this purpofe feems to be the carrot-poultice. The root of the common carrot may be grated, and moiftened with as much water as will bring it to the confiftence of a poultice or cataplafm. This must be applied to the fore, and renewed twice a-day. It generally cleans the fore, eafes the pain, and takes away the difagreeable fmell, which are objects of no fmall importance in fuch a dreadful diforder *.

Wort, or an infusion of malt, has been recommended not only as a proper drink, but as a powerful medicine in this difease. It must be frequently made fresh, and the patient may take it at pleasure. Two, three, or even four English pints of it may be drank every day for a confiderable time. No benefit can be expected from any medicine in this difease, unless it be persisted in for a long time. It is of too obstinate a nature to be foon removed; and, when it admits of a cure at all, it must be brought about by inducing an almost total change of the habit, which must always be a work of time. Setons or iffues

* London Medical Effays.

in the neighbourhood of the cancer have fometimes good effects *.

When all other medicines fail, recourfe must be had to opium, as a kind of folace. This will not indeed cure the difease, but it will ease the patient's agony, and render life more tolerable while it continues.

To avoid this dreadful diforder, people ought to use wholefome food; to take fufficient exercise in the open air; to be as easy and cheerful as possible; and carefully to guard against all blows, bruises, and every kind of preffure upon the breasts, or other glandular parts †.

In the long catalogue of human afflictions there is fcarcely one to be more dreaded than the cancer. It is no lefs painful than loathfome : it kills by inches; is feldom cured except by the knife; and even that does not always fucceed. I have frequently feen fmall tumors in the breaft, which might perhaps have ended in cancers, yield to the camphorated mercurial ointment, applied twice a-day; but after the fchirrus had broke and become a cancer, I do not remember having ever feen it cured; nor do I believe that the whole *materia medica* can afford a remedy for it.

Yet there are plenty of people in London who cure cancers; and no one, who has a fufficient fhare of faith, can be at a lofs for a cancer doctor. One may fee even

• In a cancer which had fet all medicines, and even furgery, at defiance, I lately faw remarkable effects from an obfinate perfeverance in a courfe of antifeptics. I ordered the deep ulcers to be wafhed to the bottom by means of a fyringe, twice or thrice a day, either with an infufion of the bark, or a decoction of carrot, and that the patient fhould take four or five times a-day, a glafs of good wine, with half a drachm of the beft powdered bark in it. The fores, after being wathed, were likewife fprinkled with the fame powder. When the patient began this courfe, her death was daily expected. She continued it for above two years, with manifeft advantage; but being told by an eminent furgeon, that the bark would not cure a cancer, and that the fores ought not to be wafhed, fhe difcontinued the practice, and died in a few weeks. This courfe was not expected to cure the cancer, but to prolong the patient's life, which it evidently did almoft to a miracle.

⁺ As hemlock is the principal medicine recommended in this difeafe, we would have given fome directions for the gathering and preparing of that plant; but as its different preparations are now kept in the fhops, we think it much fafer for people to get them there, with proper directions for using them.

the fronts of houses inscribed with the words, "Cancers cured here," in large characters. I lately had a patient, who once fancied that her breast was a little cancerous, and, under that impression, was kept for two years in the hands of a female cancer-curer, though the lady in reality had not the least fymptom of a cancer about her.

But credulity is a difeafe of the mind ftill more incurable than cancers. I had occafion, a few years ago, to make feveral vifits at the houfe of one of the richeft merchants in London, whofe fifter was afflicted at the time with a cancer; and though fhe lived in the fame houfe, I was never defired to look at her.' Blind credulity prevailed over reafon. Her cure was entrufted to an American quack, who knew juft as much as my lady's lapdog, of the nature and proper treatment of cancers. He only helped to kill her, which the difeafe might ultimately have done : yet furely fhe ought to have had better advice.

CHAP. XLVI.

OF POISONS.

E VERY perfon ought, in fome measure, to be acquainted with the nature and cure of poisons. They are generally taken unawares, and their effects are often fo fudden and violent, as not to admit of delay, or allow time to procure the affistance of physicians. Happily, indeed, no great degree of medical knowledge is here necessary; the remedies for most poisons being generally at hand, are easily obtained, and nothing but common prudence needful in the application of them.

The vulgar notion that every poifon is cured by fome counter-poifon, as a fpecific, has done much hurt. People believe they can do nothing for the patient, unlefs they know the particular antidote to that kind of poifon which he has taken. Whereas the cure of all poifons taken into the ftomach, without exception, depends chiefly on difcharging them as foon as poffible.

There

There is no cafe wherein the indications of cure are more obvious. Poifon is feldom long in the ftomach before it occafions ficknefs, with an inclination to vomit. This fhews plainly what ought to be done. Indeed, common fenfe dictates to every one, that, if any thing has been taken into the ftomach which endangers life, it ought immediately to be difcharged. Were this duly regarded, the danger arifing from poifons might generally be avoided. The method of prevention is obvious, and the means are in the hands of every one.

We fhall not take up the reader's time with a detail of the ridiculous notions which have prevailed among ignorant people in different ages with regard to poifons; neither fhall we mention the boafted antidotes, which have been recommended either for preventing or obviating their effects; but fhall content ourfelves with pointing out the poifons molt common in this country, and the means of avoiding their dangerous confequences.

Poifons either belong to the mineral, the vegetable, or the animal kingdom.

Mineral poifons are commonly of an acrid or corrofive quality; as arfenic, cobalt, the corrofive fublimate of mercury, &c.

Those of the vegetable kind are generally of a narcotic or flupefactive quality; as poppy, hemlock, henbane, berries of the deadly night-fhade, &c.

Poifonous animals communicate their infection either by the bite or fting. This poifon is very different from the former, and only produces its effects when received into the body by a wound.

When a perfon has taken arfenic, he foon perceives a burning heat, and a violent pricking pain in his ftomach and bowels, with an intolerable thirft, and an inclination to vomit. The tongue and throat feel rough and dry; and, if proper means be not foon administered, the patient is feized with great anxiety, hiccuping, faintings, and coldnefs

OF POISONS.

coldness of the extremities. To these fucceed black vomits, fœtid stools, with a mortification of the stomach and intestines, which are the immediate forerunners of death.

On the first appearance of these fymptoms the patient fhould drink large quantities of new milk and fallad-oil till he vomits; or he may drink warm water mixed with oil. Fat broths are likewise proper, provided they can be got ready in time. Where no oil is to be had, fresh butter may be melted and mixed with the milk or water. These things are to be drank as long as the inclination to vomit continues. Some have drank eight or ten English quarts before the vomiting ceased; and it is never fase to leave off drinking while one particle of the poison remains in the ftomach.

These oily or fat fubstances not only provoke vomiting, but likewise blunt the acrimony of the poison, and prevent its wounding the bowels; but if they should not make the person vomit, half a drachm or two scruples of the powder of ipecacuanha must be given, or a few spoonfuls of the oxymel or vinegar of squills may be mixed with the water which he drinks. Vomiting may likewise be excited by tickling the infide of the throat with a feather. Should these methods however fail, half a drachm of white vitriol, or five or fix grains of emetic tartar, must be administered.

If tormenting pains are felt in the lower belly, and there is reafon to fear that the poifon has got down to the inteftines, clyfters of milk and oil muft be very frequently thrown up; and the patient muft drink emollient decoctions of barley, oatmeal, marfh-mallows, and fuch like. He muft likewife take an infufion of fenna and manna, a folution of Glauber's falts, or fome other purgative.

After the poifon has been evacuated, the patient ought, for fome time, to live upon fuch things as are of a healing and cooling quality; to abstain from flesh and all strong liquors, and to live upon milk, broth, gruel, light puddings, and other spoon meats of easy digestion. His drink should be barley-water, linsed-tea, or infusions of any of the mild mucilaginous vegetables.

OF POISONS.

VEGETABLE POISONS, befides heat and pain of the ftomach, commonly occasion fome degree of giddinefs, and often a kind of ftupidity or folly. Perfons who have taken these poisons must be treated in the fame manner as for the mineral or corrofive.

Though the vegetable poifons, when allowed to remain in the flomach, often prove fatal; yet the danger is generally over as foon as they are difcharged. Not being of fuch a cauftic or corrofive nature, they are lefs apt to wound or inflame the bowels, than mineral fubflances: no time, however, ought to be loft in having them difcharged.

Opium, being frequently taken by mistake, merits particular attention. It is used as a medicine both in a folid and liquid form, which latter commonly goes by the name of laudanum. It is, indeed, a valuable medicine when taken in proper quantity; but, as an overdose proves a strong poison, we shall point out its common effects, together with the method of cure.

An over-dofe of opium generally occafions great drowfinefs, with flupor and other apoplectic fymptoms. Sometimes the perfon has fo great an inclination to fleep, that it is almost impossible to keep him awake. Every method must, however, be tried for this purpose. He should be toffed, shaked, and moved about. Sharp blisteringplasters should be applied to his legs or arms, and stimulating medicines, as falts of hartshorn, &c. held under his nose. It will also be proper to let blood. At the fame time every method must be taken to make him discharge the poison. This may be done in the manner directed above, viz. by the use of strong vomits, drinking plenty of warm water, with oil, &c.

Mead, befides vomits, in this cafe, recommends acid medicines with lixivial falts. He fays, that he has often given falt of wormwood mixed with juice of lemon in repeated dofes with great fuccefs.

If the body fhould remain weak and languid after the poifon has been difcharged, nourifhing diet and cordials will be proper; but when there is reafon to fear that the ftomach or bowels are inflamed, the greateft circumfpection is neceffary both with regard to food and medicine.

OF

[446]

OF THE BITES OF POISONOUS ANIMALS.

We shall begin with the bite of a mad dog, as it is both the most common and dangerous animal-poison in this country.

The creatures naturally liable to contract this difeafe are, as far as we yet know, all of the dog-kind, viz. foxes, dogs, and wolves. Hence it is called the *rabies canina*, or dog-madnefs. Of the laft we have none in this ifland; and it fo feldom happens that any perfon is bit by the first, that they fcarcely deferve to be taken notice of. If fuch a thing should happen, the method of treatment is precifely the fame as for the bite of a mad dog.

The fymptoms of madnefs in a dog are as follows: At firft he looks dull, fhews an averfion to food and company: he does not bark as ufual, but feems to murmur, is peevifh, and apt to bite ftrangers: his ears and tail droop more than ufual, and he appears drowfy: afterwards he begins to loll out his tongue, and froth at the mouth, his eye feeming heavy and watery: he now, if not confined, takes off, runs panting along with a kind of dejected air, and endeavours to bite every one he meets. Other dogs are faid to fly from him. Some think this a certain fign of madnefs, fuppofing that they know him by the fmell; but it is not to be depended on. If he efcapes being killed, he feldom runs above two or three days, till he dies exhaufted with heat, hunger, and fatigue.

This difeafe is most frequent after long, dry, hot feafons; and fuch dogs as live upon putrid stinking carrion, without having enough of fresh-water, are most liable to it.

When any perfon has been bit by a dog, the ftricteft inquiry ought to be made whether the animal was really mad. Many difagreeable confequences arife from neglecting to afcertain this point. Some people have lived in continual anxiety for many years, becaufe they had been bit by a dog which they believed to be mad; but, as he had been killed on the fpot, it was impoffible to afcertain the fact. This fhould induce us, inftead of killing a dog the moment he has bit any perfon, to do all

all in our power to keep him alive, at leaft till we can be certain whether he be mad or not.

Many circumftances may contribute to make people imagine a dog mad. He lofes his mafter, runs about in queft of him, is fet upon by other dogs, and perhaps by men. The creature, thus frightened, beat, and abufed, looks wild, and lolls out his tongue as he runs along. Immediately a crowd is after him; while he, finding himfelf clofely purfued, and taking every one he meets for an enemy, naturally attempts to bite him in felfdefence. He foon gets knocked on the head, and it paffes currently that he was mad, as it is then impoffible to prove the contrary.

This being the true hiftory of, by far, the greater part of those dogs which pass for mad, is it any wonder that numberless whimfical medicines have been extolled for preventing the effects of their bite? This readily accounts for the great variety of infallible remedies for the bite of a mad dog, which are to be met with in almost every family. Though not one in a thoufand has any claim to merit, yet they are all supported by numberless vouchers. No wonder that imaginary difeafes fhould be cured by imaginary remedies. In this way, credulous people first impose upon themselves, and then deceive others. The fame medicine which was fuppofed to prevent the effects of the bite, when the dog was not mad, is recommended to a perfon who has had the milfortune to be bit by a dog that was really mad. He takes it, trufts to it, and is undone.

To thefe miftakes we must impute the frequent ill fuccels of the medicines used for preventing the effects of the bite of a mad dog. It is not owing fo much to a defect in medicine, as to wrong applications. I am perfuaded, if proper medicines were administered immediately after the bite is received, and continued for a fufficient length of time, we should not lose one in a thoufand of those who have the misfortune to be bit by a mad dog.

This poifon is generally communicated by a wound, which neverthelefs heals as foon as a common wound : but afterwards it begins to feel painful, and as the pain fpreads

fpreads towards the neighbouring parts, the perfon becomes heavy and liftlefs. His fleep is unquiet with frightful dreams; he fighs, looks dull, and loves folitude. Thefe are the forerunners, or rather the firft fymptoms, of that dreadful difeafe occafioned by the bite of a mad dog. But as we do not propofe to treat fully on the difeafe itfelf, but to point out the method of preventing it, we fhall not take up time in flewing its progrefs from the firft invafion to its commonly fatal end.

The common notion, that this poifon may lie in the body for many years, and afterwards prove fatal, is both hurtful and ridiculous. It muft render fuch perfons as have had the misfortune to be bit very unhappy, and can have no good effects. If the perfon takes proper medicines for forty days after the time of his being bit, and feels no fymptoms of the difeafe, there is reafon to believe him out of danger. Some have indeed gone mad twelve months after being bit, but feldom later.

The medicines recommended for preventing the effects of the bite of a mad dog, are chiefly fuch as promote the different fecretions, and antifpafmodics.

Dr. Mead recommends a preventive medicine, which he fays he never knew fail, though in the fpace of thirty years he had used it a thousand times.

The Doctor's prefcription is as follows :

"Take afh-coloured ground liver-wort, cleaned, dried, and powdered, half an ounce; of black pepper powdered, a quarter of an ounce. Mix thefe well together, and divide the powder into four dofes; one of which must be taken every morning fasting, for four mornings fucceffively, in half an English pint of cow's milk warm.

"After these four doses are taken, the patient must go into the cold bath, or a cold spring or river, every morning fasting, for a month; he must be dipped all over, but not stay in (with his head above water) longer than half a minute, if the water be very cold. After this he must go in three times a-week for a fortnight longer.

"The perfon must be bled before he begins to use the medicine *."

* Though we give this prefcription on the credit of Dr. Mead, yet we would not advife any perfon, who has reafon to believe that

We

We fhall next mention the famous East-India specific, as it is called. This medicine is composed of cinnabar and muscle. It is esteemed a great antispasmodic; and, by many, extolled as an infallible remedy for preventing the effects of the bite of a mad dog.

"Take native and factitious cinnabar, of each twenty-four grains, mufk fixteen grains. Let thefe be made into a fine powder, and taken in a glafs of arrack or brandy."

This fingle dole is faid to fecure the perfon for thirty days, at the end of which it must be repeated; but if he has any fymptoms of the difease, it must be repeated in three hours.

The following is likewife reckoned a good antifpafmodic medicine :

" Take of Virginian fnake-root in powder half a drachm, gum afafœtida twelve grains, gum camphire feven grains; make thefe into a bolus with a little fyrup of faffron."

Camphire may also be given in the following manner:

"Take purified nitre half an ounce, Virginian fnakeroot in powder two drachms, camphire one drachm; rub them together in a mortar, and divide the whole into ten dofes."

Mercury is likewife recommended as of great efficacy, both in the prevention and cure of this kind of madnefs. When ufed as a preventive, it will be fufficient to rub daily a drachm of the ointment into the parts about the wound.

Vinegar is likewife of confiderable fervice, and fhould be taken freely, either in the patient's food or drink.

These are the principal medicines recommended for preventing the effects of the bite of a mad dog. We would not, however, advise people to trust to any one of them; but from a proper combination of their different powers, there is the greatest reason to hope for fuccess.

that he has been bit by a dog which was really mad, to truft to it alone. Mead was an able phyfician; but he feems to have been no great philosopher, and was sometimes the dupe of his own credulity.

The great error in the use of these medicines lies in not taking them for a sufficient length of time. They are used more like charms, than medicines intended to produce any change in the body. To this, and not to the infufficiency of the medicines, we must impute their frequent want of success.

Dr. Mead fays, that the virtue of his medicine confifts in promoting urine. But how a poifon fhould be expelled by urine, with only three or four dofes of any medicine, however powerful, it is not eafy to conceive. More time is certainly neceffary, even though the medicine were more powerful than that which the Doctor prefcribes.

The East-India specific is still more exceptionable on this account.

As these and most other medicines, taken fingly, have frequently been found to fail, we shall recommend the following course:

If a perfon be bit in a flefhy part, where there is no hazard of hurting any large blood-veffel, the parts adjacent to the wound may be cut away. But if this be not done foon after the bite has been received, it will be better to omit it.

The wound may be dreffed with falt and water, or a pickle made of vinegar and falt, and afterwards dreffed twice a-day with yellow bafilicon mixed with red precipitate of mercury.

The patient fhould begin to use either Dr. Mead's medicine, or some of the others mentioned above. If he takes Mead's medicine, he may use it, as the Doctor directs, for sour days successively. Let him then omit it for two or three days, and again repeat the same number of doses as before.

During this courfe, he must rub into the parts about the wound, daily, one drachm of the mercurial ointment. This may be done for ten or twelve days at least.

When this courfe is over, he may take a purge or two, and wait a few days till the effect of the mercury be gone off. He must then begin to use the cold bath, into which he may go every morning for five or fix weeks. If he should feel cold and chilly for a long time after coming

coming out of the cold bath, it will be better to use a tepid one, or to have the water a little warmed.

In the mean time, we would advife him not to leave off all internal medicines, but to take either one of the bolufes of fnake-root, afafœtida, and camphire; or one of the powders of nitre, camphire, and fnake-root, twice a-day. Thefe may be ufed the whole time he is bathing.

During the use of the mercurial ointment, the patient must keep within doors, and take nothing cold.

A proper regimen must be observed throughout the whole course. The patient should abstain from flesh, and all falted and high-feasoned provisions. He must avoid strong liquors, and live mostly upon a light and rather spare diet. His mind should be kept as easy and cheerful as possible, and all excessive heat and violent passions avoided with the utmost care.

I have never feen this courfe of medicine, with proper regimen, fail to prevent the hydrophobia; and cannot help again obferving, that the want of fuccefs must generally be owing either to the application of improper medicines, or not using proper ones for a fufficient length of time.

Mankind are extremely fond of every thing that promifes a fudden or miraculous cure. By trufting to thefe they often lofe their lives, when a regular courfe of medicine would have rendered them abfolutely fafe. This holds remarkably in the prefent cafe. Numbers of people, for example, believe, if they or their cattle were once dipped in the fea, it is fufficient ; as if the falt water were a charm against the effects of the bite. This, and fuch like whims, have proved fatal to many.

It is a common notion, if a perfon be bit by a dog which is not mad, that, if he fhould go mad afterwards, the perfon would be affected with the diforder at the fame time; but this notion is too ridiculous to deferve a ferious confideration. It is a good rule, however, to avoid dogs as much as poffible, as the difeafe is often upon them for fome time before its violent fymptoms appear. The hydrophobia has been occafioned by the bite of a dog G g 2 which

which shewed no other fymptoms of the difease, but listlesines and a fullen disposition.

Though we do not mean to treat fully of the cure of the hydrophobia, yet we are far from reckoning it incurable. The notion that this difeafe could not be cured, has been productive of the moft horrid confequences. It was ufual either to abandon the unhappy perfons, as foon as they were feized with the difeafe, to their fate, to bleed them to death, or to fuffocate them between mattraffes or feather beds, &c. This conduct certainly deferved the fevereft punifhment ! We hope, for the honour of human nature, it will never again be heard of.

I have never had an opportunity of treating this difeafe, and therefore can fay nothing of it from my own experience; but the learned Dr. Tiffot fays, it may be cured in the following manner:

1. The patient must be bled to a confiderable quantity; and this may be repeated twice, or thrice, or even a fourth time, if circumstances require it.

2. The patient fhould be put, if poffible, into a warm bath; and this fhould be used twice a-day.

3. He should every day receive two, or even three emollient clysters.

4. The wound, and the parts adjoining to it, fhould be rubbed with the mercurial ointment twice a-day.

5 The whole limb which contains the wound fhould be rubbed with oil, and be wrapped up in an oily flannel.

6. Every three hours a dofe of Cob's powder fhould be taken in a cup of the infufion of lime-tree and elder flowers. This powder is made by rubbing together in a mortar, to a very fine powder, of native and factitious cinnabar, each twenty-four grains; of mufk, fixteen grains *.

* The Ormfkirk medicine, as it is called, feems to me to confift chiefly of cinnabar. Though it is faid to be infallible as a preventive, yet I would not advife any one to truft to it alone. Indeed, it is ordered to be taken in a manner which gives it more the appearance of a charm than of a medicine. Surely, if a medicine is to produce any change in the body, it must be taken for fome confiderable time, and in fufficient quantity. 7. The

7. The following bolus is to be given every night, and to be repeated in the morning, if the patient be not eafy, washing it down with the infusion mentioned above: Take one drachm of Virginian snake-root in powder; of camphire and asfactida, ten grains each; of opium, one grain; and with a sufficient quantity of conferve, or rob of elder, make a bolus.

8. If there be a great nausea at the ftomach, with a bitterness in the mouth, thirty-five or forty grains of ipecacuanha, in powder, may be taken for a vomit.

9. The patient's food, if he takes any, must be light; as panado, foups made of farinaceous or mealy vegetables, &c.

10. If the patient fhould long continue weak, and fubject to terrors, he may take half a drachm of the Peruvian bark thrice a-day.

The next poifonous animal which we fhall mention is the VIPER. The greafe of this animal, rubbed into the wound, is faid to cure the bite. Though that is all the viper-catchers generally do when bit, we fhould not think it fufficient for the bite of an enraged viper. It would furely be more fafe to have the wound well fucked *, and afterwards rubbed with warm falad-oil. A poultice of bread and milk, foftened with falad-oil, fhould likewife be applied to the wound; and the patient ought to drink freely of vinegar-whey, or watergruel with vinegar in it, to make him fweat. Vinegar is one of the beft medicines which can be ufed in any kind of poifon, and ought to be taken very liberally. If the patient be fick, he may take a vomit. This courfe

* The practice of fucking out poilons is very ancient; and indeed nothing can be more rational. Where the bite cannot be cut out, this is the moft likely way for extracting the poifon. There can be no danger in performing this office, as the poifon does no harm, unlefs it be taken into the body by a wound. The perfon who fucks the wound ought, however, to walk his month frequently with falad-oil, which will fecure him from even the lealt inconveniency. The P/ylli in Africa, and the *Marfi* in Italy, were famed for curing the bites of poifonous animals, by fucking the wound; and we are told, that the Indians in North America practife the fame at this day.

will be fufficient to cure the bite of any of the poifonous animals of this country.

With regard to poifonous infects, as the bee, the wafp, the hornet, &c. their flings are feldom attended with danger, unlefs when a perfon happens to be flung by a great number of them at the fame time; in which cafe fomething fhould be done to abate the inflammation and fwelling. Some, for this purpose, apply honey; others lay pounded parfley to the part. A mixture of vinegar and Venice treacle is likewife recommended ; but I have found rubbing the part with warm falad-oil, or frequently repeated applications of pledgits dipped in laudanum, fucceed very well. Indeed, when the flings are fo numerous as to endanger the patient's life, which is fometimes the cafe, he must not only have oily poultices, or pledgits moiftened with laudanum applied to the part, but fhould likewife be bled, and take fome cooling medicines, as nitre, or cream of tartar, and fhould drink plentifully of diluting liquors.

It is the happine's of this ifland to have very few poifonous animals, and thofe which we have are by no means of the most virulent kind. Nine-tenths of the effects attributed to poifon or venom in this country, are really other difeases, and proceed from quite different causes.

We cannot, however, make the fame obfervation with regard to poifonous vegetables. These abound every where, and prove often fatal to the ignorant and unwary. This indeed is chiefly owing to careleffness. Children ought early to be cautioned against eating any kind of fruit, roots, or berries, which they do not know; and all poifonous plants to which they can have access, ought, as far as possible, to be deftroyed. This would not be fo difficult a task as fome people imagine.

Poifonous plants have no doubt their ufe, and they ought to be propagated in proper places; but, as they prove often deftructive to cattle, they fhould be rooted out of all pafture-grounds. They ought likewife, for the fafety of the human fpecies, to be deftroyed in the neighbourhood of all towns and villages; which, by the bye, are the places where they most commonly abound. I have

I have feen the poifonous hemlock, henbane, wolfsbane, and deadly night-fhade, all growing within the *environs* of a fmall town, where, though feveral perfons, within the memory of those living in it, had lost their lives by one or other of these plants, yet no method, that I could hear of, had ever been taken to root them out; though this might be done at a very trifling expence.

Seldom a year paffes, but we have accounts of feveral perfons poifoned by eating hemlock-roots, inftead of parfnips, or fome kinds of fungus which they had gathered for mufhrooms. Thefe examples ought to put people upon their guard with refpect to the former, and to put the latter entirely out of ufe. Mufhrooms may be a delicate difh; but they are a dangerous one, as they are generally gathered by perfons who do not know one kind of fungus from another, and take every thing for a mufhroom which has that appearance.

We might here mention many other plants and animals of a poifonous nature which are found in foreign countries; but, as our observations are chiefly intended for this island, we shall pass these over. It may not, however, be amifs to obferve, for the benefit of fuch of our countrymen as go to America, that an effectual remedy is now faid to be found for the bite of the rattlefnake .- The prefcription is as follows : Take of the roots of plantain and horehound, in fummer, roots and branches together, a fufficient quantity; bruife them in a mortar, and squeeze out the juice, of which give, as foon as poffible, one large spoonful; if the patient be fwelled, you must force it down his throat. This generally will cure; but, if he finds no relief in an hour after, you may give another spoonful, which never fails. -If the roots are dried, they must be moistened with a little water. To the wound may be applied a leaf of good tobacco moistened with rum.

We give this upon the faith of Dr. Brookes, who fays it was the invention of a negro; for the difcovery of which he had his freedom purchased, and a hundred pounds *per annum* settled upon him during life, by the General Assembly of Carolina.

It is poffible there may be in nature specific remedies G g 4 for

456 BITES OF POISONOUS ANIMALS.

for every kind of poifon; but as we have very little faith in any of those which have yet been pretended to be discovered, we shall beg leave again to recommend the most strict attention to the following rules, viz. That when any poisonous substance has been taken into the ftomach, it ought, as soon as possible, to be discharged by vomits, clysters, and purges; and, when poison has been received into the body by a wound, that it be expelled by medicines which promote the different fecretions, especially those of sweat, urine, and infensible perspiration; to which may be joined antispass or such medicines as take off tension and irritation; the chief of which are opium, musc, camphire, and associate.

CHAP. XLVII.

OF THE VENEREAL DISEASE.

IN the first edition of this book the venereal difeafe was omitted. The reafons, however, which at that time induced me to leave it out, have, upon more mature confideration, vanished. Bad confequences, no doubt, may arife from ignorant perfons tampering with medicine in this diforder; but the danger from that quarter feems to be more than balanced by the great and folid advantages which must arife to the patient from an early knowledge of his cafe, and an attention to a plan of regimen, which, if it does not cure the difeafe, will be fure to render it more mild, and lefs hurtful to the conflitution.

It is peculiarly unfortunate for the unhappy perfons who contract this difeafe, that it lies under a fort of difgrace. This renders difguife neceffary, and makes the patient either conceal his diforder altogether, or apply to those who promise a fudden and fecret cure; but who, in fact, only remove the fymptoms for a time, while they fix the difease deeper in the habit. By this means a flight infection, which might have been eafily removed,

OF THE VENEREAL DISEASE.

removed, is often converted into an obstinate, and fometimes incurable malady.

Another unfavourable circumftance attending this difeafe is, that it affumes a variety of different fhapes, and may with more propriety be called an affemblage of difeafes, than a fingle one. No two difeafes can require a more different method of treatment than this does in its different ftages. Hence the folly and danger of trufting to any particular noftrum for the cure of it. Such noftrums are, however, generally administered in the fame manner to all who apply for them, without the least regard to the ftate of the difeafe, the conflictuation of the patient, the degree of infection, and a thoufand other circumftances of the utmost importance.

Though the venereal difeafe is generally the fruit of unlawful embraces, yet it may be communicated to the innocent as well as the guilty. Infants, nurfes, midwives, and married women whofe hufbands lead diffolute lives, are often affected with it, and frequently lofe their lives by not being aware of their danger in due time. The unhappy condition of fuch perfons will certainly plead our excufe, if any excufe be neceffary, for endeavouring to point out the fymptoms and cure of this too common difeafe.

To enumerate all its different fymptoms, however, and to trace the difeafe minutely through its various ftages, would require a much larger fpace than falls to this part of my fubject; I fhall therefore confine my obfervations chiefly to circumftances of importance, omitting fuch as are either trifling, or which occur but feldom. I fhall likewife pais over the hiftory of the difeafe, with the different methods of treatment which it has undergone fince it was firft introduced into Europe, and many other circumftances of a fimilar nature; all of which, though they might tend to amufe the reader, yet could afford him little or no ufeful knowledge.

OF THE VIRULENT GONORRHCEA.

The virulent Gonorrhœa is an involuntary discharge of infectious mucus from the parts of generation in either fex.

fex. It generally makes its appearance within eight or ten days after the infection has been received; sometimes indeed it appears in two or three days, and at other times not before the end of four or five weeks. Previous to the difcharge, the patient feels an itching, with a fmall degree of pain in the genitals. Afterwards a thin glary matter begins to diftil from the urinary paffage, which stains the linen, and occasions a small degree of titillation, particularly at the time of making water; this, gradually increasing, arifes at length to a degree of heat and pain, which are chiefly perceived about the extremity of the urinary paffage, where a flight degree of rednefs and inflammation likewife begins to appear.

As the diforder advances, the pain, heat of urine, and running, increase, while fresh symptoms daily ensue. In men, the erections become painful and involuntary, and are more frequent and lafting than when natural. This fymptom is most troublesome when the patient is warm in bed. The pain which was at first only perceived towards the extremity, now begins to reach all up the urinary paffage, and is most intense just after the patient has done making water. The running gradually recedes from the colour of feed, grows yellow, and at length puts on the appearance of mucus.

When the diforder has arrived at its height, all the fymptoms are more intenfe; the heat of urine is fo great, that the patient dreads the making water; and though he feels a conftant inclination this way, yet it is rendered with the greatest difficulty, and often only by drops: the involuntary erections now become extremely painful and frequent; there is also a pain, heat, and fenfe of fulnefs about the feat, and the running is plentiful and tharp, of a brown, greenith, and fometimes of a bloody colour.

By a proper treatment, the violence of the fymptoms gradually abates; the heat of urine goes off; the involuntary and painful erections, and the heat and pain about the feat, become eafier; the running alfo gradually decreafes, grows whiter and thicker, till at laft it entirely difappears. By

I

By attending to thefe fymptoms, the gonorrhœa may be generally diftinguished from any other diseafe. There are, however, some few disorders for which it may be mistaken, as an ulcer in the kidnies or bladder, the *fluor albus*, or whites, in women, &c. But in the former of these, the matter comes away only with the urine, or when the sphincter of the bladder is open; whereas in a gonorrhœa the discharge is constant. The latter is more difficult to distinguish, and must be known chiefly from its effects; as pain, communicating the infection, &c.

REGIMEN .---- When a perfon has reafon to fufpect that he has caught the venereal infection, he ought most strictly to obferve a cooling regimen, to avoid every thing of a heating nature, as wines, fpirituous liquors, rich fauces, fpices, falted, high-feafoned and fmokedried provisions, &c. as also all aromatic and flimulating vegetables, as onions, garlic, fhalot, nutmeg, muftard, cinnamon, mace, ginger, and fuch-like. His food ought chiefly to confift of mild vegetables, milk, broths, light puddings, panado, gruels, &c. His drink may be barley-water, milk and water, decoctions of marsh-mallows and liquorice, linfeed-tea, or clear whey. Of thefe he ought to drink plentifully. Violent exercise of all kinds, efpecially riding on horfeback, and venereal pleafures, are to be avoided. The patient must beware of cold, and when the inflammation is violent, he ought to keep his bed.

Sometimes, indeed, a flight infection may be carried off in a few days, by bathing the parts in warm milk and water, and injecting frequently up the urethra a little fweet oil or linfeed-tea about the warmth of new milk. Should thefe not fucceed in carrying off the infection,

infection, they will at least have a tendency to lessen its virulence.

To effect a cure, however, aftringent injections will generally be found neceffary. These may be various ways prepared, but I think those made with the white vitriol are both most fase and efficacious. They can be made stronger or weaker as circumstances may require; but it is best to begin with the more gentle, and increase their power if neceffary. I generally order a drachm of white vitriol to be dissolved in eight or nine ounces of common or rose water, and an ordinary fyringe full of it to be thrown up three or four times a-day. If this quantity does not perform a cure, it may be repeated, and the dose increased *.

Whether injections be used or not, cooling purges are always proper in the gonorrhœa. They ought not, however, to be of the ftrong or draftic kind. Whatever raises a violent commotion in the body, increases the danger, and tends to drive the difease deeper into the habit. Procuring two or three ftools every fecond or third day for the first fortnight, and the same number every fourth or fifth day for the second, will generally be fufficient to remove the inflammatory symptoms, to diminish the running, and to change its colour and confistence. It gradually becomes more white and ropy as the virulence abates †.

* Although it is now very common to cure the gonorrhœa by aftringent injections, there are ftill many practitioners who do not approve this mode of practice. I can, however, from much experience, affert, that it is both the moft eafy, elegant, and efficacious method of cure; and that any bad confequences arifing from it muft be owing to the ignorance or mifconduct of the practitioner himfelf, and not to the remedy. Many, for example, ufe ftrong preparations of lead, all of which are dangerous when applied to the internal furfaces of the body; others ufe efcharotics, which inflame and injure the parts. I have known a gonorrhœa actually cured by an injection made of green tea, and would always recommend gentle methods where they will fucceed.

† If the patient can fwallow a folution of falts and manna, he may take fix drachms; or, if his conflictution requires it, an ounce of the former, with half an ounce of the latter. Thefe may be diffolved in an English pint of boiling water, whey, or thin watergruel, and taken early in the morning.

When the inflammatory fymptoms run high, bleeding is always neceffary at the beginning. This operation, as in other topical inflammations, must be repeated according to the strength and constitution of the patient, and the vehemence and urgency of the symptoms.

Medicines which promote the fecretion of urine are likewife proper in this ftage of the diforder. For this purpofe, an ounce of nitre and two ounces of gum-arabic, pounded together, may be divided into twenty-four dofes, one of which may be taken frequently in a cup of the patient's drink. If thefe fhould make him pafs his urine fo often as to become troublefome to him, he may either take them lefs frequently, or leave out the nitre altogether, and take equal parts of gum-arabic and cream of tartar. Thefe may be pounded together, and a teafpoonful taken in a cup of the patient's drink four or five times a-day. I have generally found this anfwer extremely well, both as a diuretic, and for keeping the body gently open.

When the pain and inflammation are feated high, towards the neck of the bladder, it will be proper frequently to throw up an emollient clyfter, which, befides the benefit of procuring ftools, will ferve as a fomentation to the inflamed parts.

If an infufion of fenna and tamarinds be more agreeable, two drachms of the former, and an ounce of the latter, may be infufed all night in an English pint of boiling water. 'The infufion may be strained next morning, and half an ounce of Glauber's falts diffolved in it. A tea-cup full of this infusion may be taken every half hour till it operates.

Should the patient prefer an electuary, the following will be found to anfwer very well. Take of the lenitive electuary four ounces, cream of tartar two ounces, jalap in powder two drachms, rhubarb one drachm, and as much of the fyrup of pale rofes as will ferve to make up the whole into a foft electuary. Two or three tea-fpoonfuls of this may be taken over-night, and about the fame quantity next morning, every day that the patient choofes to take a purge.

The dofes of the above medicines may be increased or diminished, according as the patient finds it necessary. We have ordered the falts to be diffolved in a large quantity of water, because it renders their operation more mild.

Soft poultices, when they can conveniently be applied to the parts, are of great fervice. They may be made of the flour of linfeed, or of wheat-bread and milk, foftened with frefh butter or fweet oil. When poultices cannot be conveniently ufed, cloths wrung out of warm water, or bladders filled with warm milk and water, may be applied. I have often known the most excruciating pains, during the inflammatory state of the gonorrhœa, relieved by one or other of these applications.

Few things tend more to keep off inflammation in the fpermatic veffels, than a proper fulpenfory for the fcrotum. It ought to be fo contrived as to fupport the tefticles, and fhould be worn from the first appearance of the difease, till it has ceased fome weeks.

The above treatment will fometimes remove the gonorrhæa fo quickly, that the perfon will be in doubt whether he really laboured under that difeafe. This, however, is too favourable a turn to be often expected. It more frequently happens, that we are able only to procure an abatement or remiffion of the inflammatory fymptoms, fo far as to make it fafe to have recourfe to the great antidote *mercury*.

Many people, on the first appearance of a gonorrhœa, fly to the ufe of mercury. This is a bad plan. Mercury is often not at all neceffary in a gonorrhœa; and, when taken too early, it does milchief. It may be neceffary to complete the cure, but can never be proper at the commencement of it.

When bleeding, purging, fomentations, and the other things recommended as above, have eafed the pain, foftened the pulfe, relieved the heat of urine, and rendered the involuntary erections lefs frequent, the patient may begin to use mercury in any form that is least difagreeable to him.

If he takes the common mercurial pill, two at night and one in the morning will be a fufficient dofe at first. Should they affect the mouth too much, the dofe must be leffened; if not at all, it may be gradually increased to five or fix pills in the day. If calomel be thought preferable, two or three grains of it, formed into a bolus, with a little of the conferve of hips, may be taken at bedtime, and the dose gradually increased to eight or ten grains.

grains. One of the moft common preparations of mercury now in use is the corrosive sublimate. This may be taken in the manner afterwards recommended under the confirmed lues or pox. I have always found it one of the most fafe and efficacious medicines when properly used.

The above medicines may either be taken every day, or every other day, as the patient is able to bear them. They ought never to be taken in fuch quantity as to raife a falivation, unlefs in a very flight degree. The difeafe may be more fafely, and as certainly, cured without a falivation as with it. When the mercury runs off by the mouth, it is not fo fuccefsful in carrying off the difeafe, as when it continues longer in the body, and is difcharged gradually.

Should the patient be purged or griped in the night by the mercury, he must take an infusion of fenna, or fome other purgative, and drink freely of water-gruel, to prevent bloody stools, which are very apt to happen should the patient catch cold, or if the mercury has not been duly prepared. When the bowels are weak and the mercury is apt to gripe or purge, these difagreeable confequences may be prevented by taking, with the above pills or bolus, half a drachm or two scruples of diafcordium, or of the Japonic confection.

To prevent the difagreeable circumftance of the mercury's affecting the mouth too much, or bringing on a falivation, it may be combined with purgatives. With this view the laxative mercurial pill has been contrived, the ufual dofe of which is half a drachm, or three pills, night and morning, to be repeated every other day; but the fafer way is for the patient to begin with two, or even with one pill, gradually increasing the dofe.

To fuch perfons as can neither fwallow a bolus nor a pill, mercury may be given in a liquid form, as it can be fufpended even in a watery vehicle, by means of gumarabic; which not only ferves this purpofe, but likewife prevents the mercury from affecting the mouth, and renders it in many refpects a better medicine *.

In

* Take quickfilver one drachm, gum-arabic reduced to a mucilage two drachms; let the quickfilver be rubbed with the mucilage,

It happens very fortunately for those who cannot be brought to take mercury inwardly, and likewife for perfons whole bowels are too tender to bear it, that an external application of it will answer equally well, and in fome respects better. It must be acknowledged, that mercury, taken inwardly for any length of time, greatly weakens and diforders the bowels; for which reafon, when a plentiful use of it becomes necessary, we would prefer rubbing to the mercurial pills. The common mercurial or blue ointment will answer very well. Of that which is made by rubbing together equal quantities of hog's-lard and quickfilver, about a drachm may be ufed at a time. The best time for rubbing it on is at night, and the most proper place the inner fide of the thighs. The patient fhould ftand before the fire when he rubs, and fhould wear flannel drawers next his fkin at the time he is using the ointment. If ointment of a weaker or ftronger kind be ufed, the quantity must be increased or diminished in proportion.

If, during the ufe of the ointment, the inflammation of the genital parts, together with the heat and feverifhnefs, fhould return, or if the mouth fhould grow fore, the gums tender, and the breath become offenfive, a dofe or two of Glauber's falts, or fome other cooling purge, may be taken, and the rubbing intermitted for a few days. As foon, however, as the figns of fpitting are gone off, if the virulency be not quite corrected, the ointment muft be repeated, but in fmaller quantities, and at longer intervals than before. Whatever way mercury is adminiftered, its ufe muft be perfifted in as long as any virulency is fulpected to remain.

During this, which may be called the fecond ftage of the diforder, though fo ftrict a regimen is not neceffary as in the first or inflammatory state, yet intemperance

lage, in a marble mortar, until the globules of mercury entirely difappear; afterwards add gradually, ftill continuing the trituration, half an ounce of baliamic fyrup, and eight ounces of fimple cinnamon-water. Two table-fpoonfuls of this folution may be taken night and morning. Some reckon this the best form in which quickfilver can be exhibited for the cure of a gonorrhœa.

of

of every kind must be avoided. The food must be light, plain, and of easy digestion; and the greatest indulgence that may be allowed with respect to drink, is a little wine diluted with a sufficient quantity of water. Spirituous liquors are to be avoided in every shape. I have often known the inflammatory symptoms renewed and heightened, the running increased, and the cure rendered extremely difficult and tedious, by one fit of excessive drinking.

When the above treatment has removed the heat of urine, and forenels of the genital parts; when the quantity of running is confiderably leffened, without any pain or fwelling in the groin or telficle fupervening; when the patient is free from involuntary erections; and, laftly, when the running becomes pale, whitifh, thick, void of ill fmell, and tenaceous or ropy; when all or most of these fymptoms appear, the gonorrhœa is arrived at its last stage, and we may gradually proceed to treat it as a gleet with aftringent and agglutinating medicines.

OF GLEETS.

A gonorrhœa frequently repeated, or improperly treated, often ends in a gleet, which may either proceed from a relaxation, or from fome remains of the difeafe. It is, however, of the greateft importance in the cure of the gleet, to know from which of thefe caufes it proceeds. When the difcharge proves very obftinate, and receives little or no check from aftringent remedies, there is ground to fulpect that it is owing to the latter; but if the drain is conftant, and is chiefly obfervable when the patient is ftimulated by lafcivious ideas, or upon ftraining to go to ftool, we may reafonably conclude that it is chiefly owing to the former.

In the cure of a gleet proceeding from relaxation, the principal defign is to brace, and reftore a proper degree of tenfion to the debilitated and relaxed veffels. For this purpofe, befides the medicines recommended in the gonorrhœa, the patient may have recourfe to ftronger and more powerful aftringents, as the Peruvian bark *, alum,

* The Peruvian bark may be combined with other aftringents, and prepared in the following manner :

Take

OF GLEETS.

alum, vitriol, galls, tormentil, bistort, ballustines, tincture of gum kino, &c. The injections may be rendered more astringent by the addition of a few grains of alum, or increasing the quantity of vitriol as far as the parts are able to bear it.

The last remedy which we shall mention in this cafe, is the cold bath, than which there is not perhaps a more powerful bracer in the whole compass of medicine. It ought never to be omitted in this fpecies of gleet, unless there be fomething in the conflitution of the patient which renders the use of it unsafe. The chief objections to the use of the cold bath are a full habit, and an unfound state of the viscera. The danger from the former may always be leffened, if not removed, by purging and bleeding; but the latter is an infurmountable obstacle, as the preffure of the water, and the fudden contraction of the external veffels, by throwing the blood with too much force upon the internal parts, are apt to occasion ruptures of the vefiels, or a flux of humours upon the difeafed organs. But where no objection of this kind prevails, the patient ought to plunge over head in water every morning fasting, for three or four weeks together. He fhould not, however, ftay long in the water, and fhould take care to have his fkin dried as foon as he comes out.

The regimen proper in this cafe is the fame as was mentioned in the laft flage of the gonorrhœa: the diet must be drying and astringent, and the drink Spa, Pyrmont, or Bristol waters, with which a little claret or red wine may fometimes be mixed. Any perfon may now afford to drink these waters, as they can be every where prepared at almost no expence, by a mixture of common chalk and oil of vitriol.

Take of Peruvian bark bruifed fix drachms, of fresh galls bruifed two drachms; boil them in a pound and a half of water to a pound: to the strained liquor add three ounces of the simple tincture of the bark. A small tea-cup full of this may be taken three times a-day, adding to each cup fifteen or twenty drops of the acid elixir of vitriol.

466

40. 9

When

When the gleet does not in the fmalleft degree yield to these medicines, there is reason to suspect that it proceeds from ulcers. In this case recours must be had to mercury, and such medicines as tend to correct any predominant acrimony with which the juices may be affected, as the decoction of China, farsaparilla, faffafras, or the like.

Mr. Fordyce fays, he has feen many obftinate gleets, of two, three, or four years ftanding, effectually cured by a mercurial inunction, when almost every other medicine has been tried in vain. Dr. Chapman feems to be of the fame opinion; but fays, he has always found the mercury fucceed best in this cafe when joined with terebinthinate and other agglutinating medicines. For which reason the Doctor recommends pills made of calomel and Venice turpentine *; and defires that their use may be accompanied with a decoction of guaiacum or farfaparilla.

The last kind of remedy which we shall mention for the cure of ulcers in the urinary paffage, are the fuppurating candles or bougies. As thefe are prepared various ways, and are generally to be bought ready made, it is needlefs to spend time in enumerating the different ingredients of which they are compoled, or teaching the manner of preparing them. Before a bougie be introduced into the urethra, however, it fhould be fmeared all over with fweet oil, to prevent it from ftimulating too fuddenly. It may be fuffered to continue in from one to feven or eight hours, according as the patient can bear it. Obstinate ulcers are not only often healed, but tumours and excrefcences in the urinary paffages taken away, and an obstruction of urine removed by means of bougies. Obstinate gleets may be removed by the ufe of bougies.

* Take Venice turpentine, boiled to a fufficient degree of hardnefs, half an ounce, calomel half a drachm. Let thefe be mixed and formed into fixty pills, of which five or fix may be taken night and morning. If, during the ufe of thefe pills, the mouth fhould grow fore, or the breath become offenfive, they must be difcontinued till thefe fymptoms difappear.

OF

OF THE SWELLED TESTICLE.

The fwelled tefficle may either proceed from infection lately contracted, or from the venereal poifon lurking in the body; the latter indeed is not very common, but the former frequently happens both in the first and second stages of a gonorrhœa; particularly when the running is unfeasonably checked, by cold, hard drinking, strong draftic purges, violent exercise, the too early use of astringent medicines, or the like.

In the inflammatory ftage, bleeding is neceffary, which must be repeated according to the urgency of the fymptoms *. The food must be light, and the drink diluting. High-feasoned food, flesh, wines, and every thing of a heating nature, are to be avoided. Fomentations are of fingular fervice. Poultices of bread and milk, fostened with fresh butter or oil, are likewise very proper, and ought constantly to be applied when the patient is in bed; when he is up, the testicles should be kept warm, and supported by a suffective, which may easily be contrived in such a manner as to prevent the weight of the testicle from having any effect.

If it fhould be found impracticable to clear the tefficle by the cooling regimen now pointed out, and extended according to circumftances, it will be neceffary to lead the patient through fuch a complete antivenereal courfe as fhall enfure him againft any future uneafinefs. For this purpofe, befides rubbing the mercurial ointment on the thighs as directed in the gonorrhœa, the patient mult be confined to bed, if neceffary, for five or fix weeks, fufpending the tefficle, all the while, with a bag or trufs, and plying him inwardly with ftrong decoctions of farfaparilla.

When these means do not succeed, and there is reafon to suspect a scrophulous or cancerous habit, either of which may support a schirrous induration, after the venereal poison is corrected, the parts should be somented

^{*} I have been accustomed for some time past to apply leeches to inflamed testicles, which practice has always been followed with the most happy effects.

OF THE SWELLED TESTICLE.

daily with a decoction of hemlock, the bruifed leaves of which may likewife be added to the poultice, and the extract at the fame time taken inwardly *. This practice is ftrongly recommended by Dr. Storck in fchirrous and cancerous cafes; and Mr. Fordyce affures us, that by this method he has cured difeafed tefticles of two or three years ftanding, even when ulcerated, and when the fchirrus had begun to be affected with pricking and lancing pains.

OF BUBOES.

Venereal buboes are hard tumours feated in the groin, occafioned by the venereal poifon lodged in this part. They are of two kinds; viz. fuch as proceed from a recent infection, and fuch as accompany a confirmed lues.

The cure of recent buboes, that is, fuch as appear foon after impure coition, may be first attempted by *difperfion*; and, if that should not succeed, by *suppuration*. To promote the dispersion of a bubo, the same regimen must be observed as was directed in the first stage of a gonorrhœa. The patient must likewise be bled, and take fome cooling purges, as the decoclion of tamarinds and fenna, Glauber's falts, and the like. If by this course the swelling and other inflammatory symptoms abate, we may fafely proceed to the use of mercury, which must be continued till the venereal virus is quite subdued \dagger .

But if the bubo fhould, from the beginning, be attended with great heat, pain, and pulfation, it will be proper to promote its fuppuration. For this purpofe the patient may be allowed to ufe his ordinary diet, and to take now and then a glafs of wine. Emollient cataplafms, confifting of bread and milk foftened with oil or frefh butter, may be applied to the part; and, in cold conflitutions, where the tumour advances flowly, white lilyroots boiled, or fliced onions raw, and a fufficient quantity of yellow bafilicon, may be added to the poultice.

3

^{*} The extract of hemlock may be made into pills, and taken in the manner directed under the article Cancer.

⁺ For the difpersion of a bubo, a number of leeches applied to the part affected will be found equally efficacious as in the inflamed testicle.

When the tumour is ripe, which may be known by its conical figure, the foftnefs of the fkin, and a fluctuation of matter plainly to be felt under the finger, it may be opened either by a cauflic or a lancet, and afterwards dreffed with digeftive ointment.

It fometimes however happens, that buboes can neither be difperfed nor brought to a fuppuration, but remain hard indolent tumours. In this cafe the indurated glands muft be confumed by cauftic; if they fhould become fchirrous, they muft be diffolved by the application of hemlock, both externally and internally, as directed in the fchirrous tefficle.

OF CHANCRES.

Chancres are fuperficial, callous, eating ulcers; which may happen either with or without a gonorrhœa. They are commonly feated about the glands, and make their appearance in the following manner: First a little red pimple arifes, which foon becomes pointed at top, and is filled with a whitish matter inclining to yellow. This pimple is hot, and itches generally before it breaks: afterwards it degenerates into an obstinate ulcer, the bottom of which is usually covered with a viscid mucus, and whose edges gradually become hard and callous. Sometimes the first appearance refembles a fimple excoriation of the cuticle; which, however, if the cause be venereal, foon becomes a true chancre.

A chancre is fometimes a primary affection, but it is much oftener fymptomatic, and is the mark of a confirmed lues. Primary chancres difcover themfelves foon after impure coition, and are generally feated in parts covered with a thin cuticle, as the lips, the nipples of women, the glans penis of men, &c. *

* When venereal ulcers are feated in the lips, the infection may be communicated by kiffing. I have feen very obfinate venereal ulcers in the lips, which I have all the reafon in the world to believe were communicated in this manner.

Nuries ought to beware of fuckling infected children, or having their breafts drawn by perfons tainted with the venereal difeafe. This caution is peculiarly neceffary for nurles who refide in the neighbourhood of great towns.

Massi R Mark to

When

When a chancre appears foon after impure coition, its treatment is nearly fimilar to that of the virulent gonorrhœa. The patient muft obferve the cooling regimen, lofe a little blood, and take fome gentle dofes of falts and manna. The parts affected ought frequently to be bathed, or rather foaked, in warm milk and water, and, if the inflammation be great, an emollient poultice or cataplafm may be applied to them. This courfe will, in moft cafes, be fufficient to abate the inflammation, and prepare the patient for the ufe of mercury.

Symptomatic chancres are commonly accompanied with ulcers in the throat, nocturnal pains, fcurvy eruptions about the roots of the hair, and other fymptoms of a confirmed lues. Though they may be feated in any of the parts mentioned above, they commonly appear upon the private parts, or the infide of the thigh. They are alfo lefs painful, but frequently much larger and harder than primary chancres. As their cure muft depend upon that of the pox, of which they are only a fymptom, we fhall take no further notice of them, till we come to treat of a confirmed lues *.

Thus we have related most of the fymptoms which accompany or fucceed a violent gonorrhœa, and have alfo given a short view of their proper treatment; there are, however, feveral others which sometimes attend this difease, as a *strangury* or obstruction of urine, a *phymofis paraphymofis*, &c.

A ftrangury may be occafioned either by a fpafmodic confriction, or an inflammation of the urethra and parts about the neck of the bladder. In the former cafe the patient begins to void his urine with tolerable eafe; but, as foon as it touches the galled or inflamed urethra, a fudden conftriction takes place, and the urine is voided by fpurts, fometimes by drops only. When the ftrangury is owing to an inflammation about the neck of the bladder, there is a conftant heat and uneafinels of the

* I have found it anfwer extremely well to fprinkle chancres twice a-day with calomel. This will often perform a cure without any other application whatever. If the chancres are upon the glans, they may be wafhed with milk and water a little warm, and afterwards the calomel may be applied as above.

Hh4

part, a perpetual defire to make water, while the patient can only render a few drops, and a troublefome *tenefmus*, or conftant inclination to go to ftool.

When the ftrangury is owing to fpafm, fuch medicines as tend to dilute and blunt the falts of the urine will be proper. For this purpole, befides the common diluting liquors, foft and cooling emulfions, fweetened with the fyrup of poppies, may be ufed. Should thefe not have the defired effect, bleeding and emollient fomentations will be neceffary.

When the complaint is evidently owing to an inflammation about the neck of the bladder, bleeding must be more liberally performed, and repeated according to the urgency of the fymptoms. After bleeding, if the ftrangury still continues, fost clysters, with a proper quantity of laudanum in them, may be administered, and emollient fomentations applied to the region of the bladder. At the fame time, the patient may take every four hours a tea-cupful of barley-water, to an English pint of which fix ounces of the fyrup of marsh-mallows, four ounces of the oil of fweet-almonds, and half an ounce of nitre, may be added. If these remedies should not relieve the complaint, and a total suppression of urine should come on, bleeding must be repeated, and the patient fet in a warm bath up to the middle. It will be proper in this cafe to difcontinue the diuretics, and to draw off the water with a catheter; but as the patient is feldom able to bear its being introduced, we would rather recommend the use of mild bougies. These often lubricate the paffage, and greatly facilitate the difcharge of urine. Whenever they begin to ftimulate or give any uneafinefs, they may be withdrawn.

The *phymofis* is fuch a conftriction of the prepuce over the glans, as hinders it from being drawn backwards; the *paraphymofis*, on the contrary, is fuch a conftriction of the prepuce behind the glans, as hinders it from being brought forward.

The treatment of these symptoms is so nearly the fame with that of the virulent gonorrhoa, that we have no occasion to enlarge upon it. In general, bleeding, purging, poultices, and emollient fomentations are sufficient. Should

Should thefe, however, fail of removing the ftricture, and the parts be threatened with a mortification, twenty or thirty grains of ipecacuanha, and one grain of emetic tartar may be given for a vomit, and may be worked off with warm water or thin gruel.

It fometimes happens, that, in fpite of all endeavours to the contrary, the inflammation goes on, and lymptoms of a beginning mortification appear. When this is the cafe, the prepuce muft be fcarified with a lancet, and, if neceffary, divided, in order to prevent a ftrangulation, and fet the imprifoned glands at liberty. We fhall not defcribe the manner of performing this operation, as it ought always to be done by a furgeon. When a mortification has actually taken place, it will be neceffary, befides performing the above operations, to foment the parts frequently with cloths wrung out of a ftrong decoction of camomile flowers and bark, and to give the patient a drachm of the bark in powder every two or three hours.

With regard to the *priapifm*, *chordee*, and other diffortions of the *penis*, their treatment is no way different from that of the gonorrhœa. When they prove very troublefome, the patient may take a few drops of laudanum at night, efpecially after the operation of a purgative through the day.

OF A CONFIRMED LUES.

We have hitherto treated of those affections in which the venereal poilon is supposed to be confined chiefly to the particular part by which it was received, and shall next take a view of the lues in its confirmed state; that is, when the poilon is actually received into the blood, and, circulating with it through every part of the body, mixes with the feveral fecretions, and renders the whole habit tainted.

The fymptoms of a confirmed lues are, buboes in the groin, pains of the head and joints, which are peculiarly troublefome in the night, or when the patient is warm in bed; fcabs and fcurfs in various parts of the body, efpecially on the head, of a yellowifh colour, refembling. a honeycomb; corroding ulcers in various parts of the body, which generally begin about the throat, from whence

whence they creep gradually, by the palate, towards the cartilage of the nofe, which they deftroy; excrefcences or exoftofes arife in the middle of the bones, and their fpungy ends become brittle, and break upon the leaft accident; at other times they are foft, and bend like wax; the conglobate glands become hard and callous, and form in the neck, armpits, groin, and mefentery, hard moveable tumours, like the king's evil; tumours of different kinds are likewife formed in the lymphatic veffels, tendons, ligaments, and nerves, as the gummata, ganglia, nodes, tophs, &c.; the eyes are affected with itching, pain, rednefs, and fometimes with total blindnefs, and the ears with a finging noife, pain, and deafnefs, whilft their internal fubstance is exulcerated and rendered carious; at length all the animal, vital, and natural functions are depraved; the face becomes pale and livid; the body emaciated and unfit for motion, and the miferable patient falls into an atrophy or walting confumption.

Women have fymptoms peculiar to the fex; as cancers of the breaft; a fuppreffion or overflowing of the menfes; the whites; hyfteric affections; an inflammation, abfcefs, fchirrus, gangrene, cancer, or ulcer of the womb; they are generally either barren or fubject to abortion; or, if they bring children into the world, they have an univerfal eryfipelas, are half rotten, and covered with ulcers.

Such is the catalogue of fymptoms attending this dreadful difeafe in its confirmed ftate. Indeed, they are feldom all to be met with in the fame perfon, or at the fame time; fo many of them, however, are generally prefent as are fufficient to alarm the patient; and if he has reafon to fufpect the infection is lurking in his body, he ought immediately to fet about the expulsion of it, otherwife the most tragical confequences will enfue.

The only certain remedy hitherto known in Europe, for the cure of this difeafe, is mercury, which may be ufed in a great variety of forms, with nearly the fame fuccefs*. Some time ago it was reckoned impoffible to cure a confirmed lues without a falivation. This method is now, however, pretty generally laid afide, and mercury

* The preparations which I now chiefly use, in the confirmed lues, are calomel and calcined mercury.

474

59113.194

is found to be as efficacious, or rather more fo, in expelling the venereal poifon, when administered in such a manner as not to run off by the falivary glands.

Though many are of opinion, that the mercurial ointment is as efficacious as any other preparation of that mineral; yet experience has taught me to think otherwife. I have often feen the most obstinate venereal cafes, where great quantities of mercurial ointment had been used in vain, yield to the faline preparations of mercury. Nor am I fingular in this opinion. Mr. Clare, an eminent furgeon of this city, affures me, that for fome time paft he has employed in venereal cafes a faline preparation of mercury with most happy fuccess. This preparation, rubbed with a sufficient quantity of any mild powder, he applies, in fmall portions, to the tongue, where, with a gentle degree of friction, it is immediately abforbed, and produces its full effect upon the fystem, without doing the least injury to the stomach or bowels; a matter of the greatest importance in the application of this most active and powerful remedy.

It is impoffible to afcertain either the exact quantity of medicines that muft be taken, or the time they ought to be continued, in order to perform a cure. Thefe will ever vary according to the conflictution of the patient, the feafon of the year, the degree of infection, the time it has lodged in the body, &c. But though it is difficult, as Aftruc obferves, to determine à priori, what quantity of mercury will, in the whole, be neceffary to cure this diftemper completely; yet it may be judged of à pofferiori, from the abatement and ceafing of the fymptoms. The fame author adds, that commonly not lefs than two ounces of the ftrong mercurial ointment is fufficient, and not more than three or four ounces neceffary.

The only chemical preparation of mercury which we fhall take notice of, is the corrofive fublimate. This was fome time ago brought into use for the venereal difease in Germany, by the illustrious Baron Van Swieten; and was soon after introduced into Britain by the learned Sir John Pringle, at that time physician to the army. The method of giving it is as follows: One grain of corrofive sublimate is diffolved in two ounces of French brandy

brandy or malt fpirits; and of this folution, an ordinary table-fpoonful, or the quantity of half an ounce, is to be taken twice a-day, and to be continued as long as any fymptoms of the diforder remain. To those whose ftomach cannot bear the folution, the fublimate may be given in form of a pill *.

Several roots, woods, and barks, have been recommended for curing the venereal difeafe; but none of them have been found, upon experience, to answer the high encomiums which had been bestowed upon them. Though no one of these is to be depended upon alone, yet, when joined with mercury, some of them are found to be very beneficial in promoting a cure. One of the best we know yet is farfaparilla, which may be prepared and taken according to the directions in the Appendix \dagger .

The mezereon-root is likewife found to be a powerful affiftant to the fublimate, or any other mercurial. It may either be ufed along with the farfaparilla, as directed in the Appendix, or by itfelf. Thofe who choofe to ufe the mezereon by itfelf, may boil an ounce of the fresh bark, taken from the root, in twelve English pints of water to eight, adding towards the end an ounce of liquorice. The dose of this is the fame as of the decoction of farfaparilla.

We have been told that the natives of America cure the venereal difeafe, in every ftage, by a decoction of the root of a plant called the Lobelia. It is used either fresh or dried; but we have no certain accounts with regard to the proportion. Sometimes they mix other roots with it, as those of the ranunculus, the ceanothus, &c.; but whether these are designed to difguise or affiss it, is doubtful. The patient takes a large draught of the de-

* The fublimate may be given in diftilled water, or any other liquid that the patient chooses. I commonly order ten grains to be diffolved in an ounce of the spirit of wine, for the conveniency of carriage, and let the patient take twenty or thirty drops of it night and morning, in half a glass of brandy or other spirits. Mr. Debra, an ingenious chymift of this place, informs me, that he prepares a falt of mercury much more mild and gentle in its operation than the sublimate, though equally efficacious.

+ See Appendix, Decoclion of Sarfaparilla.

coction

soction early in the morning, and continues to use it for his ordinary drink through the day *.

Many other roots and woods might be mentioned which have been extolled for curing the venereal difeafe, as the china-root, the roots of foap-wort, burdock, &c. as alfo the wood of guaiacum and faffafras; but as none of these have been found to possible virtues fuperior to those already mentioned, we shall, for the fake of brevity, pass them over, and shall conclude our observations on this difease, with a few general remarks concerning the proper management of the patient, and the nature of the infection.

GENERAL OBSERVATIONS.

The condition of the patient ought always to be confidered previous to his entering upon a courfe of mercury in any form. It would be equally rafh and dangerous to administer mercury to a perfon labouring under any violent acute difease, as a putrid fever, pleurify, peripneumony, or the like. It would likewise be dangerous in some chronic cases; as a flow hectic fever, or the last stage of a confumption. Sometimes, however, these difeases proceed from a confirmed lues; in which case it will be necessary to give mercury. In chronic difeases of a less dangerous nature, as the asthma, the gravel, and such like, mercury, if necessary, may be safely ad-

* Though we are ftill very much in the dark with regard to the method of curing this difeafe among the natives of America, yet it is generally affirmed that they do cure it with speed, fafety, and fuccels, and that without the leaft knowledge of mercury. Hence it becomes an object of confiderable importance to difcover their method of cure. This might furely be done by making trials of the various plants which are found in those parts, and particularly of fuch as the natives are known to make use of. All people in a rude ftate take their medicines chiefly from the vegetable kingdom, and are often poffeffed of valuable fecrets with regard to the virtues of plants, of which more enlightened nations are ignorant. Indeed, we make no doubt but fome plants of our own growth, were proper pains taken to discover them, would be found as efficacions in curing the venereal difeafe, as those of America. It must, however, be remembered, that what will cure the venereal difeafe in one country, will not always be found to have equal fuccels in another.

477

ministered.

478

ministered. If the patient's strength has been greatly exhausted by sickness, labour, abstinence, or any other cause, the use of mercury must be postponed, till by time, rest, and a nourishing diet, it can be sufficiently restored.

Mercury ought not to be administered to women during the menstrual flux, or when the period is near at hand. Neither should it be given in the last stage of pregnancy. If, however, the woman be not near the time of her delivery, and circumstances render it neceffary, mercury may be given, but in smaller doses, and at greater intervals than usual; with these precautions, both the mother and child may be cured at the same time; if not, the disorder will at least be kept from growing worfe, till the woman be brought to bed, and sufficiently recovered, when a more effectual method may be purfued, which, if the fuckles her child, will in all probability be sufficient for the cure of both.

Mercury ought always to be administered to infants with the greatest caution. Their tender condition unfits them for supporting a falivation, and makes it necessary to administer even the mildest preparations of mercury to them with a sparing hand. A similar conduct is recommended in the treatment of old perfons, who have the misfortune to labour under a confirmed lues. No doubt the infirmities of age must render people less able to undergo the fatigues of a falivation; but this, as was formerly observed, is never necessary; besides, we have generally found that mercury had much less effect upon very old perfons, than on those who were younger.

Hyfteric and hypochondriac perfons, and fuch as are fubject to an habitual diarrhœa or dyfentery, or to frequent and violent attacks of the epilepfy, or who are afflicted with the fcrophula or the fcurvy, ought to be cautious in the ufe of mercury. Where any one of thefe diforders prevails, it ought either, if poffible, to be cured, or at leaft palliated, before the patient enters upon a courfe of mercury. When this cannot be done, the mercury muft be adminiftered in fmaller dofes, and at longer intervals than ufual.

The

479

gated.

The most proper feasons for entering upon a course of mercury, are the spring and autumn, when the air is of a moderate warmth. If the circumstances of the case, however, will not admit of delay, we must not defer the cure on account of the season, but must administer the mercury; taking care, at the same time, to keep the patient's chamber warmer or cooler, according as the season of the year requires.

The next thing to be confidered is the preparation neceffary to be obferved before we proceed to adminifter a courfe of mercury. Some lay great ftrefs upon this circumftance, obferving, that by previoufly relaxing the veffels, and correcting any diforder which may happen to prevail in the blood, not only the mercury will be difpofed to act more kindly, but many other inconveniencies will be prevented.

We have already recommended bleeding and gentle purges, previous to the administration of mercury, and shall only now add, that these are always to be repeated according to the age, strength, constitution, and other circumstances of the patient. Afterwards, if it can be conveniently done, the patient ought to bathe once or twice a-day, for a few days, in luke-warm water. His diet, in the mean time, must be light, moist, and cooling. Wine, and all heating liquors, also violent bodily exercise, and all great exertions of the mind, are carefully to be avoided.

A proper regimen is likewife to be obferved by fuch as are under a courfe of mercury. Inattention to this not only endangers the patient's life, but often alfo difappoints him of a cure. A much fmaller quantity of mercury will be fufficient for the cure of a perfon who lives low, keeps warm, and avoids all manner of excefs, than of one who cannot endure to put the fmalleft reftraint upon his appetites : indeed, it but rarely happens that fuch are thoroughly cured.

There is hardly any thing of more importance, either for preventing or removing venereal infection, than cleanlinefs. By an early attention to this, the infection might often be prevented from entering the body; and, where it has already taken place, its effects may be greatly miti-

gated. The moment any perfon has reafon to fufpect that he has received the infection, he ought to wafh the parts with water and fpirits, fweet oil, or milk and water; a fmall quantity of the laft may likewife be injected up the urethra, if it can be conveniently done. Whether this difeafe at first took its rife from dirtinefs, is hard to fay; but wherever that prevails, the infection is found in its greatest degree of virulence, which gives ground to believe that a strict attention to cleanlines would go far towards extirpating it altogether *.

When the venereal difeafe has been neglected, or improperly treated, it often becomes a diforder of the habit. In this cafe, the cure muft be attempted by reftoratives, as a milk diet, the decoction of farfaparilla, and fuch like, to which mercury may be occafionally added. It is a common practice in North Britain to fend fuch patients to drink goat-whey. This is a very proper plan, provided the infection has been totally eradicated beforehand; but when that is not the cafe, and the patient

* I have not only often feen a recent infection carried off in a few days by means of cleanlinefs, viz. bathing, fomentations, injections, &c. but have likewife found it of the greatest advantage in the more advanced ftages of the difeafe. Of this I had lately a very remarkable inftance, in a man whofe penis was almost wholly confumed by venergal ulcers; the matter had been allowed to continue on the fores, without any care having been taken to clean them, till, notwithstanding the use of mercury and other medicines, it had produced the effects above mentioned. I ordered warm milk and water to be injected three or four times a-day into all the finuous ulcers, in order to wash out the matter; after which they were fluffed with dry lint to abforb the fresh matter as it was generated. The patient at the fame time took every day half a grain of the corrofive fublimate of mercury, diffolved in an ounce of brandy, and drank an English quart of the decoction of farfaparilla. By this treatment, in about fix weeks, he was perfectly cured; and, what was very remarkable, a part of the penis was actually regenerated.

Doctor Gilchrift has given an account of a fpecies of the *luss* venerea which prevails in the weft of Scotland, to which the natives give the name of Sibbins or Sivvins. The Doctor obferves, that the fpreading of this difeafe is chiefly owing to the neglect of cleanlinefs, and feems to think, that by due attention to this virtue, it might be extirpated. The treatment of this difeafe is fimilar to that of a confirmed lues or pox. The yaws, a difeafe which is now very common both in America and the Weft India iflands, may alfo be cured in the fame manner.

trufts

trusts to the whey for finishing his cure, he will often be disappointed. I have frequently known the disease return with all its virulence after a course of goat-whey, even when that course had been thought quite fufficient for completing the cure.

One of the most unfortunate circumstances attending patients in this difeafe, is the neceffity they are often laid under of hurrying the cure. This induces them to take medicine too fast, and to leave it off too foon. A few grains more of medicine, or a few days longer confinement, would often be fufficient to perfect the cure; whereas, by neglect of these, a small degree of virulence is still left in the system, which gradually vitiates, and at length contaminates the whole mass. To avoid this, we would advise, that the patient should never leave off taking medicine immediately upon the disappearing of the symptoms, but continue it for some time after, gradually lessening the quantity, till there is fufficient ground to believe that the disease is entirely eradicated.

It is not only difficult, but abfolutely impoffible, to afcertain the exact degree of virulence that may attend the difeafe; for which reafon it will always be a much fafer rule to continue the ufe of medicine too long, than to leave it off too foon. This feems to be the leading maxim of a modern practitioner of fome note for the venereal difeafe, who always orders his patient to perform a quarantine of at leaft forty days, during which time he takes forty bottles of, I fuppofe, a ftrong decoction of farfaparilla, or fome other anti-venereal fimple. Whoever takes this method, and adds a fufficient quantity of corrofive fublimate, or fome other active preparation of mercury to the decoction, will feldom fail to cure a confirmed lues.

It is peculiarly unfortunate for the cure of this difeafe, that not one in ten of those who contract it, are either able or willing to fubmit to a proper plan of regimen. The patient is willing to take medicine; but he must follow his business, and, to prevent fuspicions, must eat and drink like the rest of the family. This is the true fource of nine-tenths of all the mischief arising from the venereal difease. I never knew the cure attended with any great difficulty or danger where the patient strictly I i followed the phyfician's advice; but a volume would not be fufficient to point out the dreadful confequences which proceed from an oppofite conduct. Schirrous tefficles, ulcerous fore throats, madnefs, confumptions, carious bones, and a rotten progeny, are a few of the bleffings derived from this fource.

There is a species of falle reasoning, with regard to this difease, which proves fatal to many. A person of a found constitution contracts a slight degree of the diforder. He gets well without taking any great care, or using much medicine, and hence concludes that this will always be the cafe. The next time the difeafe occurs. though ten times more virulent, he pursues the fame courfe, and his conftitution is ruined. Indeed, the different degrees of virulence in the fmall-pox are not greater than in this difeafe, though, as the learned Sydenham observes, in some cases the most skilful physicians cannot cure, and in others the most ignorant old woman cannot kill the patient in that diforder. Though a good conftitution is always in favour of the patient, yet too great ftress may be laid upon it. It does not appear from obfervation, that the most robust constitution is able to overcome the virulence of the venereal contagion, after it has got into the habit. In this cafe, a proper courfe of medicine is always indifpenfably neceffary.

Although it is impoffible, on account of the different degrees of virulence, &c. to lay down fixed and certain rules for the cure of this difeafe, yet the following general plan will always be found fafe, and often fuccefsful, *viz.* to bleed and administer gentle purges with diuretics during the inflammatory state, and, as soon as the fymptoms of inflammation are abated, to administer mercury, in any form that may be most agreeable to the patient. The fame medicine, affisted by the decoction of farfaparilla, and a proper regimen, will not only fecure the constitution against the further progress of a confirmed pox, but will generally perform a complete cure.

As I did not deem the venereal difease a proper subject of discussion in a book designed chiefly for regular families and the nursery, I left it out in the early editions of this work; but was afterwards prompted to introduce it, at the request of the publisher, and from a consideration

deration that a little feafonable advice might be of mate* rial fervice in a complaint, which is often too long concealed, through motives of fhame and delicacy. I have fince published a separate treatife on this complaint, and must refer to it for minuter information. Here I shall only add one temark, that, though every apothecary's boy pretends to cure the venereal difeafe, there is not any one malady that requires more diferimination, nor any medicine that is more frequently mifapplied than mercury. It is looked upon as a certain cure in every fpecies of the difeafe, whereas it is only proper in one; and, even in this, unlefs it is given with great care, much mischief may be done. Mercury is too active a medicine to be administered at random, which, God knows, is too often the cafe, when entrusted to the hands of ignorance and felf-fufficiency. But I must at the fame time add, that when mercury is properly administered, it will never hurt the conftitution, nor fail to cure the fues venerea.

CHAP. XLVIII.

DISEASES OF WOMEN.

7 OMEN, in all civilized nations, have the management of domeftic affairs, and it is very proper they fhould, as Nature has made them lefs fit for the more active, and laborious employments. This indulgence, however, is generally carried too far; and females, inftead of being benefited by it, are greatly injured, from the want of exercise and free air. To be fatisfied of this, one need only compare the fresh and ruddy looks of a milk-maid, with the pale complexion of those females whose whole time is spent within doors. Though Nature has made an evident diffinction between the male and female, with regard to bodily strength and vigour, yet she certainly never meant, either that the one should be always without, or the other always within doors.

The confinement of females, befides hurting their figure and complexion, relaxes their folids, weakens liz

their

their minds, and diforders all the functions of the body. Hence proceed obstructions, indigestion, flatulence, abortions, and the whole train of nervous diforders. These not only unfit women for being mothers and nurses, but often render them whimstical and ridiculous. A found mind depends so much upon a healthy body, that where the latter is wanting, the former is rarely to be found.

I have always obferved, that women who were chiefly employed without doors, in the different branches of hufbandry, gardening, and the like, were almost as hardy as their hufbands, and that their children were likewife ftrong and healthy. But as the bad effects of confinement and inactivity upon both fexes have been already shewn, we shall proceed to point out those circumstances in the structure and design of females, which subject them to peculiar difeases; the chief of which are, their monthly evacuations, pregnancy, and child-bearing. These, indeed, cannot properly be called difeases; but, from the delicacy of the fex, and their being often improperly managed in fuch structures.

OF THE MENSTRUAL DISCHARGE.

Females generally begin to menftruate about the age of fifteen, and leave it off about fifty, which renders thefe two periods the most critical of their lives. About the first appearance of this discharge, the constitution undergoes a very considerable change, generally indeed for the better, though sometimes for the worse. The greatest care is now necessary, as the future health and happiness of the female depends in a great measure upon her conduct at this period *.

• It is the duty of mothers, and those who are intrusted with the education of girls, to instruct them early in the conduct and management of themselves at this critical period of their lives. False modesty, inattention, and ignorance of what is beneficial or hurtful at this time, are the sources of many difeases and misfortunes in life, which a few sensible lessons from an experienced matron might have prevented. Nor is care less necessary in the subfequent returns of this discharge. Taking improper food, violent affections of the mind, or catching cold at this period, is often source further to ruin the health, or to render the female ever after incapable of procreation.

If a girl about this time of life be confined to the houfe, kept conftantly fitting, and neither allowed to romp about, nor employed in any active bufinefs, which gives exercife to the whole body, fhe becomes weak, relaxed, and puny; her blood not being duly prepared, fhe looks pale and wan; her health, fpirits, and vigour decline, and the finks into a valetudinarian for life. Such is the fate of numbers of those unhappy females, who, either from too much indulgence, or their own, narrow circumstances, are, at this critical period, denied the benefit of exercife and free air.

A lazy, indolent disposition proves likewife very hurtful to girls at this period. One feldom meets with complaints from obstructions amongst the more active and industrious part of the fex; whereas the indolent and lazy are feldom free from them. These are in a manner eaten up by the chlorofis, or green-ficknefs, and other difeafes of this nature. We would therefore recommend it to all who wish to escape these calamities, to avoid indolence and inactivity, as their greatest enemies, and to be as much abroad in the open air as poffible.

Another thing which proves very hurtful to girls about this period of life, is unwholefome food. Fond of all manner of trash, they often indulge in it, till their whole humours are quite vitiated. Hence enfue indigeftions, want of appetite, and a numerous train of evils. If the fluids be not duly prepared, it is utterly impoffible that the fecretions fhould go properly on. Accordingly we find, that fuch girls as lead an indolent life, and eat great quantities of trash, are not only subject to obstructions of the menfes, but likewife to glandular obstructions; as the fcrophula, or king's evil, &c.

A dull disposition is also very hurtful to girls at this period. It is a rare thing to fee a fprightly girl who does not enjoy good health, while the grave, moping, melancholy creature proves the very prey of vapours and hysterics. Youth is the feafon for mirth and cheerfulnels: let it therefore be indulged; it is an abfolute duty. To lay in a flock of health in time of youth, is as neceffary a piece of prudence, as to make provision against the decays of old age. While, therefore, wife Nature prompts the happy youth to join in fprightly 113 amule-

amusements, let not the fevere dictates of hoary age forbid the useful impulse, nor damp with ferious gloom the seafon defined to mirth and innocent festivity.

Another thing very hurtful to females about this period of life, is ftrait clothes. They are fond of a fine fhape, and foolifhly imagine that this can be acquired by lacing themfelves tight. Hence, by fqueezing the ftomach and bowels, they hurt the digeftion, and occafion many incurable maladies. This error is not indeed fo common as it has been, but, as fafhions change, it may come about again; we therefore think it not improper to mention it. I know many females who, to this day, feel the dreadful effects of that wretched cuftom which prevailed fome years ago, of fqueezing every girl into as fmall a fize in the middle as poffible. Human invention could not poffibly have devifed a practice more deftructive to health.

After a female has arrived at that period of life when the menfes ufually begin to flow, and they do not appear, but, on the contrary, her health and fpirits begin to decline, we would advife, inftead of fhutting the poor girl up in the houfe, and dofing her with fteel, afafœtida, and other naufeous drugs, to place her in a fituation where fhe can enjoy the benefit of free air and agreeable company. There let her eat wholefome food, take fufficient exercife, and amufe herfelf in the moft agreeable manner; and we have little reafon to fear, but Nature, thus affifted, will do her proper work. Indeed fhe feldom fails, unlefs where the fault is on our fide,

This difcharge in the beginning is feldom fo inftantaneous as to furprife females unawares. It is generally preceded by fymptoms which foretel its approach; as a fenfe of heat, weight, and dull pain in the loins; diftention and hardnefs of the breafts; head-ach; lofs of appetite; laffitude; palenefs of the countenance; and fometimes a flight degree of fever. When thefe fymptoms appear about the age at which the menftrual flux ufually begins, every thing fhould be carefully avoided which may obftruct that neceffary and falutary evacuation; and all means ufed to promote it; as fitting frequently over the fteams of warm water, drinking warm diluting liquors, &cc.

After

After the *menfes* have once begun to flow, the greateft care fhould be taken to avoid every thing that may tend to obftruct them. Females ought to be exceedingly cautious of what they eat or drink at the time they are out of order. Every thing that is cold, or apt to four on the ftomach, ought to be avoided; as fruit, buttermilk, and fuch like. Fifh, and all kinds of food that are hard of digeftion, are alfo to be avoided. As it is impoffible to mention every thing that may difagree with individuals at this time, we would recommend it to every female to be very attentive to what difagrees with herfelf, and carefully to avoid it.

Cold is extremely hurtful at this particular period. More of the fex date their difeafes from colds caught while they are out of order, than from all other caufes. This ought furely to put them upon their guard, and to make them very circumfpect in their conduct at fuch times. A degree of cold that will not in the leaft hurt them at another time, will at this period be fufficient entirely to ruin their health and conftitution.

The greatest attention ought likewife to be paid to the mind, which should be kept as easy and cheerful as poffible. Every part of the animal economy is influenced by the passions, but none more fo than this. Anger, fear, grief, and other affections of the mind, often occafion obstructions of the menstrual flux, which prove absolutely incurable.

As far as my observation goes, there are no women in the world fo inattentive to this discharge as the Engliss ; and they fuffer accordingly, as a very great number of them are obstructed, and many prove barren in consequence.

From whatever caufe this flux is obstructed, except in the state of pregnancy, proper means should be used to restore it. For this purpose we would recommend sufficient exercise, in a dry, open, and rather cool air; wholesome diet, and, if the body be weak and languid, generous liquors; also cheerful company and all manner of amusements. If these fail, recours must be had to medicine.

When obstructions proceed from a weak relaxed state of the folids, fuch medicines as tend to promote digestion, to brace the folids, and affift the body in preparing good blood, ought to be used. For this purpose, I have of late made use of pills composed of prepared steel, powdered myrrh, and the best aloes, equal parts. These must be formed into pills of the common fize, and two or three of them given every night at bed-time. The principal medicines, however, are iron, and the Peruvian bark, with other bitters and aftringents. Filings of iron may be infufed in wine or ale, two or three ounces to an English quart, and after it has stood for two or three weeks, it may be filtered, and about half a wineglafs of it taken twice a-day : or prepared fteel may be taken in the dofe of half a drachm, mixed with a little honey or treacle, three or four times a-day. The bark and other bitters may either be taken in fubftance or infusion, as is most agreeable to the patient.

When obstructions proceed from a viscid state of the blood; or for women of a gross or full habit, evacuations, and such medicines as attenuate the humours, are neceffary. The patient in this case ought to be bled, to bathe her feet frequently in warm water, to take now and then a cooling purge, and to live upon a spare thin diet. Her drink should be whey, water, or small-beer, and she ought to take sufficient exercise. A tea-spoonful of the tincture of black hellebore may also be taken twice a-day in a cup of warm water.

When obstructions proceed from affections of the mind, as grief, fear, anger, &c. every method should be taken to amule and divert the patient. And that the may the more readily forget the caufe of her affliction, the ought, if possible, to be removed from the place where it happened. A change of place, by prefenting the mind with a variety of new objects, has often a very happy influence in relieving it from the deepest distres. A foothing, kind, and affable behaviour to females in this fituation is also of the last importance.

An obstruction of the menses is often the effect of other maladies. When this is the case, instead of giving medicines to force that discharge, which might be dangerous,

gerous, we ought by all means to endeavour to reftore the patient's health and ftrength. When that is effected, the other will return of courfe.

But the menftrual flux may be too great as well as too fmall. When this happens, the patient becomes weak, the colour pale, the appetite and digeftion are bad, to which cedematous fwellings of the feet, dropfies, and confumptions often enfue. This frequently happens to women about the age of forty-five or fifty, and is very difficult to cure. It may proceed from a fedentary life; a full diet, confifting chiefly of falted, high-feafoned, or acrid food; the use of fpirituous liquors; exceffive fatigue; relaxation; a diffolved state of the blood; violent paffions of the mind, &c.

The treatment of this difeafe must be varied according to its cause. When it is occasioned by any error in the patient's regimen, an opposite course to that which induced the diforder must be pursued, and such medicines taken as have a tendency to restrain the flux, and counteract the morbid affections of the system whence it proceeds.

To reftrain the flux, the patient fhould be kept quiet and eafy both in body and mind. If it be very violent, fhe ought to lie in bed with her head low; to live upon a cool and flender diet, as veal or chicken broths with bread; and to drink decoctions of nettle-roots, or the greater comfrey. If thefe be not fufficient to ftop the flux, ftronger aftringents may be ufed, as Japan earth, alum, elixir of vitriol, the Peruvian bark, &c. *

The uterine flux may offend in quality as well as in quantity. What is ufually called the *fluor albus*, or whites, is a very common difeafe, and proves extremely hurtful to delicate women. This difcharge, however, is not always white, but fometimes pale, yellow, green, or

* Two drachms of alum and one of Japan earth may be pounded together, and divided into eight or nine dofes, one of which may be taken three times a day.

Perfons whofe ftomachs cannot bear the alum, may take two table-fpoonfuls of the tincture of rofes three or four times a-day, to each dofe of which ten drops of laudanum may be added.

If these should fail, half a drachm of the Peruvian bark, in powder, with ten drops of the elixir of vitriol, may be taken, in a glass of red wine four times a-day.

of a blackifh colour; fometimes it is fharp and corrofive, fometimes foul and fœtid, &c. It is attended with a pale complexion, pain in the back, lofs of appetite, fwelling of the feet, and other figns of debility. It generally proceeds from a relaxed ftate of the body, arifing from indolence, the exceflive use of tea, coffee, or other weak and watery diet.

To remove this difeafe, the patient muft take as much exercife as fhe can bear without fatigue. Her food fhould be folid and nourifhing, but of eafy digeftion; and her drink rather generous, as red port or claret, mixed with Pyrmont, Briftol, or lime-water. Tea and coffee are to be avoided. I have often known ftrong broths have an exceeding good effect, and fometimes a milk diet alone will perform a cure. The patient ought not to lie too long a-bed. When medicine is neceffary, we know none preferable to the Peruvian bark, which in this cafe ought always to be taken in fubftance. In warm weather, the cold bath will be of confiderable fervice.

That period of life at which the *menfes* ceafe to flow, is likewife very critical to the fex. The ftoppage of any cuftomary evacuation, however fmall, is fufficient to diforder the whole frame, and often to deftroy life itfelf. Hence it comes to pafs, that fo many women either fall into chronic diforders, or die about this time. Such of them, however, as furvive it, without contracting any chronic difeafe, often become more healthy and hardy than they were before, and enjoy ftrength and vigour to a very great age.

If the *menfes* ceafe all of a fudden in women of a full habit, they ought to abate fomewhat of their ufual quantity of food, especially of the more nourishing kind, as flesh, eggs, &c. They ought likewise to take sufficient exercise, and to keep the body open. This may be done by taking, once or twice a-week, a little rhubarb, or an infusion of hiera picra in wine or brandy.

It often happens that women of a groß habit, at this period of life, have ulcerous fores break out about their ancles, or in other parts of the body. Such ulcers ought to be confidered as critical, and fhould either be fuffered to continue open, or have artificial drains fubfituted in their ftead.

ftead. Women who will have fuch fores dried up, are often foon after carried off by acute difeafes, or fall into those of a chronic nature.

OF PREGNANCY.

Though pregnancy is not a difeafe, yet that flate is often attended with a variety of complaints which merit attention, and which fometimes require the affiftance of medicine. Some women indeed are more healthy during their pregnancy, than at any other time; but this is by no means the general cafe; most of them *breed in* forrow, and are frequently indifposed during the whole time of pregnancy. Few fatal difeases, however, happen during that period; and hardly any, except abortion, that can be called dangerous.

Pregnant women are often afflicted with the heartburn. The method of treating this complaint has been already pointed out. They are likewife, in the more early periods of pregnancy, often haraffed with ficknefs and vomiting, especially in the morning. The method of relieving these complaints has also been shewn. Both the head-ach and tooth-ach are very troublefome fymptoms of pregnancy. The former may generally be removed by keeping the body gently open, by the ufe of prunes, figs, roafted apples, and fuch like. When the pain is very violent, bleeding may be neceffary. For the treatment of the latter, we must refer to that article. Several other complaints incident to pregnant women might be mentioned, as a cough and difficulty of breathing, suppression and incontinency of urine, &c; but as all of these have been taken notice of before, it is needlefs to repeat them.

Every pregnant woman is more or lefs in danger of abortion. This should be guarded against with the greatest care, as it not only weakens the constitution, but renders the woman liable to the fame misfortune afterwards*. Abortion may happen at any period of pregnancy,

* Every mother who procures an abortion does it at the hazard of her life; yet there are not a few who run this rifk merely to prevent nancy, but it is most common in the fecond or third month. Sometimes, however, it happens in the fourth or fifth. If it happens within the first month, it is usually called a false conception; if after the feventh month, the child may often be kept alive by proper care.

The common caufes of abortion are, the death of the child; weaknefs or relaxation of the mother; great evacuations; violent exercife; raifing great weights; reaching too high; jumping, or flepping from an eminence; vomiting; coughing; convultion-fits; blows on the belly; falls; fevers; difagreeable fmells; excefs of blood; indolence; high living, or the contrary; violent paffions or affections of the mind, as fear, grief, &c.

The figns of approaching abortion are, pain in the loins, or about the bottom of the belly; a dull heavy pain in the infide of the thighs; a flight degree of coldnefs, or fhivering; ficknefs, palpitation of the heart; the breafts become flat and foft; the belly falls; and there is a difcharge of blood or watery humours from the womb.

To prevent abortion, we would advife women of a weak or relaxed habit to use folid food, avoiding great quantities of tea, and other weak and watery liquors; to rife early and go foon to bed; to shun damp houses; to take frequent exercise in the open air, but to avoid fatigue; and never to go abroad in damp foggy weather, if they can shun it.

Women of a full habit ought to ufe a fpare diet, avoiding ftrong liquors, and every thing that may tend to heat the body, or increase the quantity of blood. Their diet fhould be of an opening nature, confisting principally of vegetable fubstances. Every woman with child ought to be kept cheerful and eafy in her mind. Her appetites, even though depraved, ought to be indulged as far as prudence will permit.

vent the trouble of bearing and bringing up children. It is furely a most unnatural crime, and cannot, even in the most abandoned, be viewed without horror; but in the decent matron, it is still more unpardonable.—Those wretches who daily advertise their affistance to women in this bufines, deferve, in my opinion, the most fevere of all human punishments.

OF PREGNANCY.

When any figns of abortion appear, the woman ought to be laid in bed on a mattrafs, with her head low. She fhould be kept quiet, and her mind foothed and comforted. She ought not to be kept too hot, nor to take any thing of a heating nature. Her food fhould confift of broths, rice and milk, jellies, gruels made of oatmeal, and the like, all of which ought to be taken cold.

If the be able to bear it, the thould lofe at leaft half a pound of blood from the arm. Her drink ought to be barley-water tharpened with juice of lemon; or the may take half a drachm of powdered nitre, in a cup of watergruel, every five or fix hours. If the woman be feized with a violent loofenefs, the ought to drink the decoction of calcined hartthorn prepared. If the be affected with vomiting, let ber take frequently two table-fpoonfuls of the faline mixture. In general, opiates are of fervice; but they thould always be given with caution.

Sanguine robust women, who are liable to miscarry at a certain time of pregnancy, ought always to be bled a few days before that period arrives. By this means, and observing the regimen above prescribed, they might often escape that missfortune.

Though we recommend due care for preventing abortion, we would not be underftood as reftraining pregnant women from their ufual exercifes. This would generally operate the quite contrary way. Want of exercife not only relaxes the body, but induces a plethora, or too great a fulnefs of the veffels, which are the two principal caufes of abortion. There are, however, fome women of fo delicate a texture, that it is neceffary for them to avoid almost every kind of exercise during the whole period of pregnancy.

OF CHILD-BIRTH.

Many difeafes proceed from the want of due care in child-bed; and the more hardy part of the fex are moft apt to defpife the neceffary precautions in this ftate. This is peculiarly the cafe with young wives. They think, when the labour pains are ended, the danger is over; but in truth it may only then be faid to be begun. Nature, if left to herfelf, will feldom fail to expel the *fætus*; but proper

OF CHILD-BIRTH.

proper care and management are certainly neceffary for the recovery of the mother No doubt, mifchief may be done by too much as well as by too little care. Hence females who have the greatest number of attendants in child-bed, generally recover worst. But this is not peculiar to the state of child-bed. Excessive care always defeats its own intention, and is generally more dangerous than none at all *.

During actual labour, nothing of a heating nature ought to be given. The woman may now and then take a little panado, and her drink ought to be toaft and water, or thin groat-gruel. Spirits, wines, cordial-waters, and other things which are given with a view to ftrengthen the mother, and promote the birth, for the moft part tend only to increase the fever, inflame the womb, and retard the labour. Befides they endanger the woman afterwards, as they often occasion violent and mortal hæmorrhages, or dispose her to eruptive and other fevers.

When the labour proves tedious and difficult, to prevent inflammations, it will be proper to bleed. An emollient clyfter ought likewife frequently to be adminiftered, and the patient fhould fit over the fteams of warm water. The paffage ought to be gently rubbed with a little foft *pomatum* or frefh butter, and cloths wrung out of warm water applied over the belly. If nature feems to fink, and the woman is greatly exhausted with fatigue,

* Though the management of women in child-bed has been practifed as an employment fince the earlieft accounts of time, yet it is still in most countries on a very bad footing. Few women think of following this employment till they are reduced to the neceffity of doing it for bread. Hence not one in a hundred of them have any education, or proper knowledge of their bufinels. It is true, that Nature, if left to herfelf, will generally expel the fatus; but it is equally true, that most women, in child-bed, require to be managed with skill and attention, and that they are often hurt by the fuperflitious prejudices of ignorant and officious midwives. The mifchief done in this way is much greater than is generally imagined; most of which might be prevented by allowing no women to practife midwifery but fuch as are properly qualified. Were due attention paid to this, it would not only be the means of faving many lives, but would prevent the neceffity of employing men in this indelicate and difagreeable branch of medicine, which is, on many accounts, more proper for the other fex.

494

a draught

OF CHILD-BIRTH.

a draught of generous wine, or fome other cordial, may be given, but not otherwife. These directions are sufficient in natural labours; and in all preternatural cases, a skilful surgeon, or man-midwife, ought to be called as foon as possible.

After delivery, the woman ought to be kept as quiet and eafy as poffible *. Her food fhould be light and thin, as gruel, panado, &c. and her drink weak and diluting. To this rule, however, there are many exceptions. I have known feveral women, whofe fpirits could not be fupported in child-bed without folid food and generous liquors; to fuch, a glafs of wine and a bit of chicken must be allowed.

Sometimes an exceffive hæmorrhage or flooding happens after delivery. In this cafe the patient flould be laid with her head low, kept cool, and be in all refpects treated as for an exceffive flux of the *menfes*. If the flooding proves violent, linen cloths, which have been wrung out of a mixture of equal parts of vinegar and water, or red wine, flould be applied to the belly, the loins, and the thighs: thefe must be changed as they grow dry; and may be difcontinued as foon as the flooding abates \dagger .

If there be violent pains after delivery, the patient ought to drink plentifully of warm diluting liquors, as groat-gruel, or tea with a little faffron in it; and to take fmall broths, with carraway-feeds, or a bit of orange-peel in them; an ounce of the oil of fweet almonds may likewife be frequently taken in a cup of any of the above liquors; and if the patient be reillefs, a fpoonful of the fyrup of poppies may now and then be mixed with a cup of her drink. If the be hot or feverifh, one of the fol-

* We cannot help taking notice of that ridiculous cuftom which ftill prevails in fome parts of the country, of collecting a number of women together upon fuch occafions. Thefe, inftead of being ufeful, ferve only to crowd the houfe, and obftruct the neceffary aitendants. Befides, they hurt the patient with their noife; and often, by their untimely and impertinent advice, do much mifchief.

† In a violent flooding after delivery, I have feen very good effects from the following mixture: Take of penny-royal water, fimple cinnamon water, and fyrup of poppies, each two ounces, elixir of vitriol a drachm. Mix, and take two table-fpoonfuls every two hours, or oftener, if neceffary.

lowing powders may be taken in a cup of her ufual drink every five or fix hours *.

An inflammation of the womb is a dangerous and not unfrequent difeafe after delivery. It is known by pains in the lower part of the belly, which are greatly increafed upon touching; by the tenfion or tightnefs of the parts; great weakuefs; change of countenance; a conftant fever, with a weak and hard pulfe; a flight *delirium*, or raving; fometimes inceffant vomiting; a hiccup; a difcharge of reddifh, flinking, fharp water from the womb; an inclination to go frequently to flool; a heat, and fometimes total fuppreflion of urine.

This must be treated like other inflammatory diforders, by bleeding and plentiful dilution. The drink may be thing gruel or barley-water; in a cup of which half a drachm of nitre may be diffolved, and taken three or four times a-day. Clysters of warm milk and water must be frequently administered: and the belly should be fomented by cloths wrung out of warm water, or by applying bladders filled with warm milk and water to it.

A fupprefion of the *lochia* or ufual difcharges after delivery, and the milk-fever, muft be treated nearly in the fame manner as an inflammation of the womb. In all thefe cafes, the fafeft courfe is plentiful dilution, gentle evacuations, and fomentations of the parts affected. In the milk-fever, the breafts may be embrocated with a little warm linfeed-oil, or the leaves of red cabbage may be applied to them. The child fhould be often put to the breaft, or it fhould be drawn by fome other perfon.

Nothing would tend more to prevent the milk-fever than putting the child early to the breaft. The cuftom of not allowing children to fuck for the first two or three days, is contrary to Nature and common fense, and is very hurtful both to the mother and child.

Every mother who has milk in her breafts, ought either to fuckle her own child, or to have her breafts fre-

* Take of crabs claws prepared half an ounce, purified nitre two drachms, faffron powdered half a drachm; rub them together in a mortar, and divide the whole into eight or nine dofes.

When the patient is low fpirited, or troubled with hysterical complaints, the ought to take frequently twelve or fifteen drops of the tincture of afafectida in a cup of penny royal tea.

quently

quently drawn, at least for the first month. This would prevent many of the difeases which prove fatal to women in child-bed.

When an inflammation happens in the breaft, attended with rednefs, hardnefs, and other fymptoms of fuppuration, the fafeft application is a poultice of bread and milk, foftened with oil or frefh butter. This may be renewed twice a-day, till the tumour be either difcuffed or brought to fuppuration. The ufe of repellents, in this cafe, is very dangerous; they often occasion fevers, and fometimes cancers; whereas a fuppuration is feldom attended with any danger, and has often the moft falutary effects.

When the nipples are fretted or chapped, they may be anointed with a mixture of oil and bees-wax, or a little powdered gum-arabic may be fprinkled on them. I have feen Hungary water applied to the nipples, have a very good effect. Should the complaint prove obftinate, a cooling purge may be given, which generally removes it.

The milliary fever is a difeafe incident to women in child-bed; but as it has been treated of already, we shall take no farther notice of it. The celebrated Hoffman obferves, That this fever of child-bed women might generally be prevented, if they, during their pregnancy, were regular in their diet, ufed moderate exercise, took now and then a gentle laxative of manna, rhubarb, or cream of tartar; not forgetting to bleed in the first months, and avoid all sharp air. When the labour is coming on, it is not to be hastened with forcing medicines, which inflame the blood and humours, or put them into unnatural commotions. Care should be taken, after the birth, that the natural excretions proceed regularly; and if the pulfe be quick, a little nitrous powder, or fome other cooling medicines, should be administered.

The most fatal diforder confequent upon delivery is the *puerperal*, or child-bed, fever. It generally makes its attack upon the fecond or third day after delivery. Sometimes indeed it comes on fooner, and at other times, though rarely, it does not appear before the fifth or fixth day.

It begins, like most other fevers, with a cold or shivering fit, which is fucceeded by restlessness, pain of the K k head,

OF CHILD-BIRTH.

head, great fickness at the ftomach, and bilious vomiring. The pulfe is generally quick, the tongue dry, and there is a remarkable depression of spirits and loss of ftrength. A great pain is ufually felt in the back, hips. and region of the womb; a fudden change in the quantity or quality of the lochia alfo takes place; and the patient is frequently troubled with a tenefinus, or constant inclination to go to ftool. The urine, which is very high coloured, is discharged in small quantity, and generally with pain. The belly fometimes fwells to a confiderable bulk, and becomes fusceptible of pain from the flighteft touch. When the fever has continued for a few days, the fymptoms of inflammation ufually fubfide, and the difeafe acquires a more putrid form. At this period, if not fooner, a bilious or putrid loofenefs, of an obstinate and dangerous nature, comes on, and accompanies the difeafe through all its future progrefs.

There is not any difeafe that requires to be treated with more fkill and attention than this; confequently the beft affiftance ought always to be obtained as foon as poffible. In women of plethoric conftitutions, bleeding will generally be proper at the beginning; it ought, however, to be ufed with caution, and not to be repeated, unlefs where the figns of inflammation rife high; in which cafe it will allo be neceffary to apply a bliftering plafter to the region of the womb.

During the rigour, or cold fit, proper means fhould be ufed to abate its violence and fhorten its duration. For this purpofe, the patient may drink freely of warm diluting liquors, and, if low, may take now and then a cup of wine-whey; warm applications to the extremities, as heated bricks, bottles or bladders filled with warm water, and fuch like, may alfo be ufed with advantage.

Emollient clyfters of milk and water, or of chicken water, ought to be frequently administered through the course of the difease. These prove beneficial, by promoting a discharge from the intestines, and also by acting as a kindly fomentation to the womb and parts adjacent. Great care, however, is requisite in giving them, on

OF CHILD-BIRTH.

on account of the tendernels of the parts in the pelvis at this time.

To evacuate the offending bile from the flomach, a vomit is generally given. But as this is apt to increase the irritability of the flomach, already too great, it will be fafer to omit it, and to give in its flead a gentle laxative, which will both tend to cool the body, and to procure a free difcharge of the bile *.

The medicine which I have always found to fucceed beft in this difeafe, is the faline draught. This, if frequently repeated, will often put a ftop to the vomiting, and at the fame time leffen the violence of the fever. If it runs off by ftool, or if the patient be reftlefs, a few drops of laudanum, or fome fyrup of poppies, may occafionally be added.

If the ftools fhould prove fo frequent, as to weaken and exhauft the patient, a ftarch clyfter, with thirty or forty drops of laudanum in it, may be administered as occasion fhall require; and the drink may be rice-water, in every English pint of which half an ounce of gumarabic has been diffolved. Should these fail, recourse must be had to Columbo-root, or the powder of bole combined with opium.

Though in general the food ought to be light, and the drink diluting, yet when the difease has been long protracted, and the patient is greatly spent by evacuations, it will be necessary to support her with nourishing diet, and generous cordials.

It was obferved, that this fever, after continuing for fome time, often acquires a putrid form. In this cafe the Peruvian bark muft be given, either by itfelf, or joined with cordials, as circumftances may require. As the bark in fubftance will be apt to purge, it may be given in decoction or infufion mixed with the tincture of rofes, or other gentle aftringents; or a fcruple of the extract of bark with half an ounce of fpirituous cinnamon-water, two ounces of common water, and ten

* Midwives ought to be very cautions in administering vomits or purges to women in child-bed. I have known a woman who was recovering extremely well, thrown into the most imminent danger, by a strong purge which was given her by an officious midwife. drops of laudanum, may be made into a draught, and given every fecond, third, or fourth hour, as shall be found necessary.

When the ftomach will not bear any kind of nourifhment, the patient may be fupported for fome time by clyfters of beef-tea or chicken-broth.

To avoid this fever, every woman in child-bed ought to be kept perfectly eafy; her food fhould be light and fimple, and her bed-chamber cool and properly ventilated. There is not any thing more hurtful to a woman in this fituation than being kept too warm. She ought not to have her body bound too tight, nor to rife too foon from bed, after delivery; catching cold is alfo to be avoided; and a proper attention fhould be paid to cleanlinefs.

To prevent the milk-fever, the breafts ought to be frequently drawn; and if they are filled previous to the onfet of a fever, they fhould, upon its first appearance, be drawn, to prevent the milk from becoming acrid, and its being absorbed in this state. Costiveness is likewife to be avoided. This will be best effected by the use of mild clysters and a laxative diet.

We fhall conclude our obfervations on child-bed women, by recommending it to them, above all things, to beware of cold. Poor women, whofe circumftances oblige them to quit their bed too foon, often contract difeafes from cold, of which they never recover. It is a pity the poor are not better taken care of in this fituation.

But the better fort of women run the greatest hazard from too much heat. They are generally kept in a fort of bagnio for the first eight or ten days, and then dreffed out to see company. The danger of this conduct must be obvious to every one.

The fuperfittious cuftom of obliging women to keep the houfe till they go to church, is likewife a very common caufe of catching cold. All churches are damp, and moft of them cold; confequently they are the very worft places to which a woman can go to make her first vifit, after having been confined in a warm room for a month.

OF BARRENNESS.

Barrennefs may be very properly reckoned among the difeafes of females, as few married women, who have not children, enjoy a good ftate of health. It may proceed from various caufes, as high living, grief, relaxation, &c.; but it is chiefly owing to an obftruction or irregularity of the menftrual flux.

It is very certain that high living vitiates the humours, and prevents fecundity. We feldom find a barren woman among the labouring poor, while nothing is more common among the rich and affluent. The inhabitants of every country are prolific in proportion to their poverty; and it would be an eafy matter to adduce many inftances of women, who, by being reduced to live entirely upon a milk and vegetable diet, have conceived and brought forth children, though they never had any before. Would the rich ufe the fame fort of food and exercife as the better fort of peafants, they would feldom have caufe to envy their poor vaffals and dependants the bleffing of a numerous and healthy offspring, while they pine in forrow for the want of even a fingle heir to their extenfive domains.

Affluence begets indolence, which not only vitiates the humours, but induces a general relaxation of the folids; a ftate highly unfavourable to procreation. To remove this, we would recommend the following courfe: Firft, fufficient exercife in the open air; fecondly, a diet confifting chiefly of milk and vegetables *; thirdly, the ufe of aftringent medicines, as fteel, alum, dragon's blood, elixir of vitriol, the Spa or Tunbridge waters, Peruvian bark, &c.; and laftly, above all, the cold bath.

* Dr. Cheyne avers, that want of children is oftener the fault of the male than of the female, in this the Doctor and I do not agree, and firongly recommends a milk and vegetable diet to the former as well as the latter ; adding, that his triend Dr. Taylor, whom he calls the Milk doctor of Croydon, had brought fundry opulent families in his neighbourhood, who had continued fome years after marriage without progeny, to have feveral fine children, by keeping both parents, for a confiderable time, to a milk and vegetable diet.

Kk3

Barrennef

OF BARRENNESS.

Barrennels is often the confequence of grief, fudden fear, anxiety, or any of the paffions which tend to obftruct the menstrual flux. When barrennels is fuspected to proceed from affections of the mind, the perfon ought to be kept as easy and cheerful as possible; all difagreeable objects are to be avoided, and every method taken to amuse and entertain the fancy.

I believe I have never written, and I hope I never shall write, any thing offensive to real modelty. Yet I have not suppressed, from motives of false delicacy, what I thought might be of importance on a fubject fo clofely connected with conjugal happinefs. The remarks here made, though few and apparently fimple, contain all that is really known, in medical practice, of the caufes and remedies of barrennefs. I therefore with to put married ladies on their guard against the fallacy of private fuggestions, and of public advertisements on this head. The things commonly advifed by filly nurfes, as well as the grand restoratives of quack impostors, are all of them dangerous stimulants, that provoke desire, but enfeeble, inftead of ftrengthening, the powers of nature, and render a weaknefs, which proper treatment might have removed, abfolutely incurable.

CHAP. XLIX.

DISEASES OF CHILDREN.

MISERABLE indeed is the lot of man in the ftate of infancy! He comes into the world more helplefs than any other animal, and ftands much longer in need of the protection and care of his parents : but, alas! this care is not always beftowed upon him ; and when it is, he often fuffers as much from improper management as he would have done from neglect. Hence the officious care of parents, nurfes, and midwives, becomes one of the most fruitful fources of the diforders of infants *.

• Of the officious and ill-judged care of midwives, we shall adduce only one instance, viz. the common practice of torturing infants,

It must be obvious to every attentive perfon, that the first difeases of infants arise chiefly from their bowels. Nor is this in the least to be wondered at, as they are in a manner poisoned with indigestible drugs and improper diet as foon as they come into the world. Every thing that the stomach cannot digest may be confidered as a poison; and unless it can be thrown up, or voided by stool, it must occasion sickness, gripes, spasmodic affections of the bowels, or what the good women call inwards fits, and at last convulsions and death.

As these fymptoms evidently arise from somewhat that irritates the intestines, doubtless the proper method of cure must be to expel it as foon as possible. The most fafe and effectual method of doing this is by gentle vomits. Five or fix grains of the powder of ipecacuanha may be mixed in two table-spoonfuls of water, and sweetened with a little fugar. A tea-spoonful of this may be given to the infant every quarter of an hour till it operates; or, what will more certainly answer the purpole, a grain of emetic tartar may be diffolved in three ounces of water, fweetened with a little fyrup, and given as above. Those who are willing to use the emetic tartar, may give fix or feven drops of the antimonial wine, in a tea-spoonful of water or thin gruel. Small doses of the ipecacuanha wine will be found more gentle than any of the above, and ought to be preferred.

These medicines will not only cleanse the stomach, but will generally likewise open the body. Should this, however, not happen, and if the child be costive, some gentle purge will be necessary: for this purpose, some

fants, by fqueezing their breafts, to draw off the milk, as they call it. Though a fmall quantity of moifture is generally found in the breafts of infants, yet, as they are certainly not intended to give fuck, this ought never to be drawn off. I have feen this cruel operation bring on hardnefs, inflammation, and fuppuration of the breafts; but never knew any ill confequences from its being omitted. When the breafts are hard, the only application that we would recommend is a foft poultice, or a little of the diachylon plafter, fpread thin upon a bit of foit leather, about the fize of half a crown, and applied over each nipple. Thefe may be fuffered to continue till the hardnefs difappears.

manna

DISEASES OF CHILDREN.

504

manna and pulp of caffia may be diffolved in boiling water, and given in fmall quantities till it operates; or, what will anfwer rather better, a few grains of magnefia alba may be mixed in any kind of food that is given to the child, and continued till it has the defired effect. If thefe medicines be properly administered, and the child's belly and limbs frequently rubbed with a warm hand before the fire, they will feldom fail to relieve those affections of the stomach and bowels from which infants fuffer fo much.

Thefe general directions include most of what can be done for relieving the internal diforders of infants. They will likewife go a confiderable way in alleviating those which appear externally, as the rash, gum, or fellon, &c. These, as was formerly observed, are principally owing to too hot a regimen, and confequently will be most effectually relieved by gentle evacuations. Indeed, evacuations of one kind or other constitute a principal part of the medicine of infants, and will feldom, if administered with prudence, in any of their diseafes, fail to give relief.

OF THE MECONIUM,

The ftomach and bowels of a new-born infant are filled with a blackifh-coloured matter of the confiftence of fyrup, commonly called the *meconium*. This is generally paffed foon after the birth, by the mere effort of Nature; in which cafe it is not neceffary to give the infant any kind of medicine. But if it fhould be retained, or not fufficiently carried off, a little manna, or *magnefia alba*, may be given, as mentioned above; or, if thefe fhould not be at hand, a common fpoonful of whey, fweetened with a little honey, or raw fugar, will anfwer the purpofe.

The most proper medicine for expelling the meconium is the mother's milk, which is always at first of a purgative quality. Were children allowed to suck as soon as they shew an inclination for the breast, they would feldom have occasion for medicines to discharge the meconium; but even where this is not allowed, they ought never to have daubs of syrup, oils, and other indigestible stuff, crammed down their throats.

THE APHTHÆ, OR THRUSH.

The aphthæ are little whitish ulcers affecting the whole infide of the mouth, tongue, throat, and stomach of infants. Sometimes they reach through the whole inteftinal canal; in which case they are very dangerous, and often put an end to the infant's life.

If the aphthæ are of a pale colour, pellucid, few in number, foft, fuperficial, and fall eafily off, they are not dangerous; but if opake, yellow, brown, black, thick, or running together, they ought to be dreaded.

It is generally thought that the aphthæ owe their origin to acid humours; we have reafon, however, to believe, they are more frequently owing to too hot a regimen both of the mother and child. It is a rare thing to find a child who is not dofed with wine, punch, cinnamon-waters, or fome other hot and inflaming liquors, almost as foon as it is born. It is well known that thefe will occasion inflammatory diforders even in adults; is it any wonder then that they should heat and inflame the tender bodies of infants, and fet, as it were, the whole constitution on a blaze?

The moft proper medicines for the aphthæ are vomits, fuch as have been already recommended, and gentle laxatives. Five grains of rhubarb, and half a drachm of *magnefia alba*, may be rubbed together, and divided into fix dofes, one of which may be given to the infant every four or five hours till they operate. These powders may either be given in the child's food, or a little of the fyrup of pale roses, and may be repeated as often as is found neceffary to keep the body open. It is common in this cafe to administer calomel; but as that medicine fometimes occasions gripes, it ought always to be given to infants with caution.

Many things have been recommended for gargling the mouth and throat in this difeafe: but it is not ealy to apply thefe in very young children; we would therefore recommend it to the nurfe to rub the child's mouth frequently with a little borax and honey; or with the following mixture: Take fine honey an ounce, borax a drachm,

506 THE APHTHÆ, OR THRUSH.

drachm, burnt alum half a drachm, rofe-water two drachms; mix them together. A very proper application in this cafe, is a folution of ten or twelve grains of white vitriol in eight ounces of barley-water. Thefe may be applied with the finger, or by means of a bit of foft rag tied to the end of a probe.

OF ACIDITIES.

The food of children being for the moft part of an acefcent nature, it readily turns four upon the ftomach, efpecially if the body be any way difordered. Hence moft difeafes of children are accompanied with evident figns of acidity, as green ftools, gripes, &c. Thefe appearances have induced many to believe, that all the difeafes of children were owing to an acid abounding in the ftomach and bowels; but whoever confiders the matter attentively, will find that thefe fymptoms of acidity are oftener the effect than the caufe of their difeafes.

Nature evidently intended, that the food of children fhould be acefcent; and unlefs the body be difordered, or the digeftion hurt, from fome other caufe, we will venture to fay, that the acefcent quality of their food is feldom injurious to them. Acidity, however, is often a fymptom of diforders in children; and, as it is fometimes a troublefome one, we fhall point out the method of relieving it.

When green ftools, gripes, purgings, four fmells, &c. fhew that the bowels abound with an acid, the child fhould have a little fmall broth, with light white bread in it; and fhould have fufficient exercife, in order to promote the digeftion. It has been cuftomary in this cafe to give the pearl-julep, chalk, crabs' eyes, and other teftaceous powders. Thefe, indeed, by their abforbent quality, may correct the acidity; but they are attended with this inconvenience, that they are apt to lodge in the bowels, and occafion coffivenefs, which may prove very hurtful to the infant. For this reafon they fhould never be given unlefs mixed with purgative medicines; as rhubarb, manna, or fuch like.

The

The beft medicine which we know, in all cafes of acidity, is that fine infipid powder, called *magnefia alba*. It purges, and at the fame time corrects the acidity: by which means it not only removes the difeafe, but carries off its caufe. It may be given in any kind of food, or in a mixture, as recommended in the Appendix *.

When an infant is troubled with gripes, it ought not at firft to be dofed with brandy, fpiceries, and other hot things; but fhould have its body opened with an emollient clyfter, or the medicine mentioned above; and, at the fame time, a little brandy may be rubbed on its belly with a warm hand before the fire. I have feldom feen this fail to eafe the gripes of infants. If it fhould happen, however, not to fucceed, a little brandy or other fpirits may be mixed with thrice the quantity of warm water, and a tea-fpoonful of it given frequently, till the infant be eafier. Sometimes a little peppermint-water will anfwer this purpofe as well.

GALLING AND EXCORIATION.

These are very troublesome to children. They happen chiefly about the groin and wrinkles of the neck, under the arms, behind the ears, and in other parts that are moistened by the sweat or urine.

As these complaints are, in a great measure, owing to want of cleanliness, the most effectual means of preventing them are, to wash the parts frequently with cold water, to change the linen often, and, in a word, to keep the child in all respects thoroughly clean. When this is not fufficient, the excoriated parts may be sprinkled with absorbent or drying powders; as burnt hartshorn, tutty, chalk, crabs-claws' prepared, and the like. When the parts affected are very fore, and tend to a real ulceration, it will be proper to add a little sugar of lead to the powders; or to anoint the place with the camphorated ointment. If the parts be washed with spring-water, in which a little white vitriol has been diffolved, it will dry and heal them very powerfully. One of the best

* See Appendix, Laxative absorbent Mixture.

508 GALLING AND EXCORIATION.

applications for this purpofe is to diffolve fome fuller's. earth in a fufficient quantity of hot water; and after it has flood till it is cold, to rub it gently upon the galled parts once or twice a-day.

STOPPAGE OF THE NOSE.

The noftrils of infants are often plugged up with a grofs *mucus*, which prevents their breathing freely, and likewife renders it difficult for them to fuck or fwallow.

Some in this cafe order, after a fuitable purge, two or three grains of white vitriol diffolved in half an ounce of marjoram-water, and filtered, to be applied now and then to the noftrils with a linen rag. Wedellus fays, if two grains of white vitriol, and the fame quantity of *elaterium*, be diffolved in half an ounce of marjoram-water, and applied to the nofe, as above directed, that it brings away the *mucus* without fneezing.

In obflinate cafes these medicines may be tried; but I have never found any thing necessary, besides rubbing the nose at bed-time with a little sweet oil, or fresh butter. This resolves the filth, and renders the breathing more free *.

OF VOMITING.

From the delicate ftate of children, and the great fenfibility of their organs, a vomiting or loofenefs may be induced by any thing that irritates the nerves of the ftomach or inteftines. Hence thefe diforders are much more common in childhood than in the more advanced periods of life. They are feldom, however, dangerous, and ought never to be confidered as difeafes, unlefs when they are violent, or continue fo long as to exhauft the ftrength of the patient.

Vomiting may be excited by an over-quantity of food; by food that is of fuch a nature as to irritate the nerves of the flomach too much; or by the fen-

* Some nurfes remove this complaint by fucking the child's nofe. This is by no means a cleanly operation; but when nurfes have the refolution to do it, I am far from difeouraging the practice.

fibilit

fibility of the nerves being fo much increased as to render them unable to bear the ftimulus of even the mildest element.

When vomiting is occafioned by too much food, it ought to be promoted, as the cure will depend upon cleanfing the ftomach. This may be done either by a few grains of ipecacuanha, or a weak folution of emetic tartar, as mentioned before. When it is owing to food of an actid or irritating quality, the diet ought to be changed, and aliment of a milder nature fubfituted in its ftead.

When vomiting proceeds from an increafed degree of fenfibility, or too great an irritability of the nerves of the ftomach, fuch medicines as have a tendency to brace and ftrengthen that organ, and to abate its fenfibility, muft be ufed. The first of these intentions may be anfwered by a flight infusion of the Peruvian bark, with the addition of a little rhubarb and orange-peel; and the fecond by the faline draughts, to which a few drops of liquid laudanum may occasionally be added.

In obstinate vomitings the operation of internal medicines may be affisted by aromatic fomentations made with wine, applied warm to the pit of the stomach; or the use of the stomach-plaster, with the addition of a little *Theriaca*.

OF A LOOSENESS.

A loofenefs may generally be reckoned falutary, when the ftools are four, flimy, green, or curdled. It is not the difcharge, but the production of fuch ftools, which ought to be remedied. Even where the purging is thin and watery, it ought not to be checked too fuddenly, as it often proves critical, efpecially when the child has caught cold, or an eruption on the fkin has difappeared. Sometimes an evacuation of this kind fucceeds a humid ftate of the atmosphere, in which cafe it may also prove of advantage, by carrying off a quantity of watery humours, which would otherwise tend to relax the habit.

As the principal intention of the cure of a loofenels is to evacuate the offending matter, it is cultomary to

give

give the patient a gentle vomit of ipecacuanha, and afterwards to exhibit fmall and frequent dofes of rhubarb; interpoling abforbent medicines to mitigate the acrimony of the humours. The best purge, however, in this case, is magnefia alba. It is at the fame time abforbent and laxative, and operates without exciting gripes.

The antimonial wine, which acts both as an emetic and purge, is alfo an excellent medicine in this cafe. By being diluted with water, it may be proportioned to the weakeft conflitution; and, not being difagreeable to the palate, it may be repeated as often as occafion requires. Even one dofe will frequently mitigate the difeafe, and pave the way for the ufe of abforbents. If, however, the patient's ftrength will permit, the medicine ought to be repeated every fix or eight hours, till the ftools begin to affume a more natural appearance; afterwards a longer fpace may be allowed to intervene between the dofes. When it is neceffary to repeat the medicine frequently, the dofe ought always to be a little increafed, as its efficacy is generally diminifhed by ufe.

Some, upon the first appearance of a loofenefs, fly immediately to the use of absorbent medicines and aftringents. If these be administered before the offending humours are discharged, though the disease may appear to be mitigated for a little time, it soon afterwards breaks forth with greater violence, and often proves statal. After proper evacuations, however, these medicines may be administered with considerable advantage.

Should any gripings or reftleffnefs remain after the ftomach and bowels have been cleanfed, a tea-fpoonful of the fyrup of poppies may be given in a little fimple cinnamon-water three or four times a-day till thefe fymptoms have ceafed.

OF ERUPTIONS.

Children, while on the breaft, are feldom free from eruptions of one kind or other. Thefe, however, are not often dangerous, and ought never to be dried up but with the greatest caution. They tend to free the bodies of

OF ERUPTIONS.

of infants from hurtful humours, which, if retained, might produce fatal diforders.

The eruptions of children are chiefly owing to improper food and neglect of cleanlinefs. If a child be fluffed at all hours with food, that its flomach is not able to digeft, fuch food, not being properly affimilated, inftead of nourifhing the body, fills it with grofs humours. Thefe mult either break out in form of eruptions upon the fkin, or remain in the body, and occafion fevers and other internal diforders. That neglect of cleanlinefs is a very general caufe of eruptive diforders, muft be obvious to every one. The children of the poor, and of all who defpife cleanlinefs, are almost conftantly found to fwarm with vermin, and are generally covered with the fcab, itch, and other eruptions.

When eruptions are the effect of improper food, or want of cleanlinefs, a proper attention to thefe alone will generally be fufficient to remove them. If this fhould not be the cafe, fome drying medicines will be neceffary. When they are applied, the body ought at the fame time to be kept open, and cold is carefully to be avoided. We know no medicine that is more fafe for drying up cutaneous eruptions than fulphur, provided it be prudently ufed. A little of the flowers of fulphur may be mixed with frefh butter, oil, or hog's lard, and the parts affected frequently touched with it.

The most obstinate of all the eruptions incident to children are, the *tinea capitis*, or fcabbed head, and chilblains. The fcabbed head is often exceedingly difficult to cure, and fometimes indeed the cure proves worse than the difease. I have frequently known children feized with internal diforders, of which they died foon after, their fcabbed heads had been healed by the application of drying medicines *. The cure ought always first to be

* I fome time ago faw a very firiking inftance of the danger of fubfituting drying medicines in the place of cleanlinefs and wholefome food, in the Foundling Hofpital at Ackworth, where the children were grevioufly afflicted with fcabbed heads, and other cutaneous diforders. Upon inquiry it was found, that very little at ention

OF ERUPTIONS.

be attempted by keeping the head very clean, cutting off the hair, combing and brushing away the feabs, &c. If this is not fufficient, let the head be fhaved once a week, washed daily with foap fuds, and gently anointed with a liniment made of train oil eight ounces, red precipitate, in fine powder, one drachm. And if there be proud flefh, it should be touched with a bit of blue vitriol, or fprinkled with a little burnt alum. While these things are doing, the patient must be confined to a regular light diet, the body fhould be kept gently open; and cold, as far as possible, ought to be avoided. To prevent any bad confequences from flopping this difcharge, it will be proper, efpecially in children of a grofs habit, to make an iffue in the neck or arm, which may be kept open till the patient becomes more ftrong, and the conflitution be fomewhat mended.

Chilblains commonly attack children in cold weather. They are generally occafioned by the feet or hands being kept long wet or cold, and afterwards fuddenly heated. When children are cold, inftead of taking exercife to warm themfelves gradually, they run to the fire. This occafions a fudden rarefaction of the humours, and an infraction of the veffels; which, being often repeated, the veffels are at laft over-diftended, and forced to give way.

To prevent it, violent cold and fudden heat must be equally avoided. When the parts begin to look red and fwell, the patient ought to be purged, and to have the affected parts frequently rubbed with mustard and

tion was paid either to the propriety or foundnefs of their provifions, and that cleanlinefs was totally neglected; accordingly it was advifed that they fhould have more wholefome food, and be kept thoroughly clean. This advice, however, was not followed. It was too troublefome to the fervants, fuperintendants, &c. The bufinefs was to be done by medicine; which was accordingly attempted, but had nearly proved fatal to the whole houfe. Fevers, and other internal diforders, immediately appeared, and at length a putrid dyfentery, which proved fo infectious, that it carried off a great many of the children, and fpread over a confiderable part of the neighbouring country.

brandy,

OF ERUPTIONS.

brandy, or fomething of a warming nature. They ought likewife to be covered with flannel, and kept warm and dry. Some apply warm afhes between cloths to the fwelled parts, which frequently help to reduce them. When there is a fore, it must be dreffed with Turner's cerate, the ointment of tutty, the plaster of cerus, or fome other drying ointment. These fores are indeed troublefome, but feldom dangerous. They generally heal as foon as the warm weather fets in.

OF THE CROUP.

Children are often feized very fuddenly with this difeafe, which, if not quickly relieved, proves mortal. It is known by various names in different parts of Britain. On the eaft coaft of Scotland, it is called the *croup*. On the weft, they call it the *chock* or *fluffing*. In fome parts of England, where I have obferved it, the good women call it the *rifing of the lights*. It feems to be a fpecies of *afthma*, attended with very acute and violent catarrhal fymptoms.

This difeafe generally prevails in cold and wet feafons. It is most common upon the fea-coast, and in low marshy countries. Children of a gross and lax habit are most liable to it. I have fometimes known it hereditary. It generally attacks children in the night, after having been much exposed to damp cold easterly winds through the day. Damp houses, wet feet, thin shoes, wet clothes, or any thing that obstructs the perspiration, may occafion the croup.

It is attended with a frequent pulfe, quick and laborious breathing, which is performed with a peculiar kind of croaking noife, that may be heard at a confiderable diftance. The voice is fharp and fhrill, and the face is generally much flufhed, though fometimes it is of a livid colour.

When a child is feized with the above fymptoms, his feet fhould immediately be put into warm water. He ought likewife to be bled *, and to have a laxative clyfter

admi-

^{*} In this difease bleeding is not always proper; but in very full habits it must certainly be of use.

OF THE CROUP.

administered as soon as possible. He should be made to breathe over the steams of warm water and vinegar; or an emollient decoction, and emollient cataplasms or fomentations, may be applied round his neck. If the symptoms do not abate, a bliftering-plaster must be applied round the neck, or between the shoulders, and the child may take frequently a table-spoonful of the following julep: Take penny-royal water three ounces, syrup of althea and of poppies, each one ounce; mix them together.

Afafœtida is found to have a good effect in this cafe. It may be both given in form of clyfter, and taken by the mouth. Two drachms of afafœtida may be diffolved in one ounce of Mindererus's fpirit, and three ounces of penny-royal water. A table-fpoonful of this mixture may be given every hour, or oftener, if the patient's ftomach be able to bear it. If the child cannot be brought to take this medicine, two drachms of the afafœtida may be diffolved in a common clyfter, and adminiftered every fix or eight hours, till the violence of the difeafe abates *.

To prevent a return of the diforder, all those things which occasion it must be carefully avoided; as wet feet, cold, damp, easterly winds, &c. Children who have had frequent returns of this difease, or whose constitutions feem to dispose them to it, ought to have their diet properly regulated; all food that is viscid or hard of di-

* I was lately favoured with a letter from Dr. William Turnbull in London, a phyfician of great experience, and who, from his former fituation on the north-east coast of England, had many opportunities of observing the symptoms and progress of this dangerous difease. The Doctor's fentiments differ very little from my own; he observes, that he never found bliftering of any fervice; and recommends cataplaims of garlic, camphor, and Venice treacle to be applied both to the throat and foles of the feet. He likewife recommends boluffes of camphor, caftor, valerian-root, falt of hartfhorn, and musk, adapted to the age, ftrength, &c. of the patient; after which he advifes two fpoonfuls of the following decoction :- Take of garlic and diffilled vinegar each an ounce, hyfop-water eight ounces; beat up the ingredients together, gradually mixing the water, and adding three ounces of honey. Let the whole be fimmered over a gentle fire, and afterwards ftrained for use.

514

geftion,

OF THE CROUP.

geftion, and all crude, raw, trafhy fruits, are to be avoided. They ought likewife to have a drain conftantly kept open in fome part of their body, by means of a feton or iffue. I have fometimes known a Burgundypitch plafter, worn continually between the fhoulders for feveral years, have a very happy effect in preventing the return of this dreadful diforder.

OF TEETHING.

Dr. Arbuthnot obferves, that above a tenth part of infants die in teething, by fymptoms proceeding from the irritation of the tender nervous parts of the jaws, occafioning inflammations, fevers, convulfions, gangrenes, &c. These fymptoms are in a great measure owing to the great delicacy and exquisite fensibility of the nervous fystem at this time of life, which is too often increased by an effeminate education. Hence it comes to pass, that children who are delicately brought up, always fuffer most in teething, and often fall by convulfive diforders.

About the fixth or feventh month the teeth generally begin to make their appearance; first, the *incifores*, or fore-teeth; next, the *canini*, or dog-teeth; and, lastly, the *molares*, or grinders. About the feventh year, there comes a new set; and about the twentieth, the two inner grinders, called *dentes fapientiæ*, the teeth of wisdom.

Children about the time of cutting their teeth, flaver much, and have generally a loofenefs. When the teething is difficult, efpecially when the dog-teeth begin to make their way through the gums, the child has flartings in his fleep, tumours of the gums, watchings, gripes, green flools, the thrufh, fever, difficult breathing, and convulfions.

Difficult teething requires nearly the fame treatment as an inflammatory difeafe. If the body be bound, it must be opened either by emollient clysters or gentle purgatives; as manna, magnesia alba, rhubarb, fenna, or the like. The food should be light, and in small quantity; the drink plentiful, but weak and diluting, as infusions of balm, or of the lime-tree flowers; to which about a third or fourth part of milk may be added.

OF TEETHING.

If the fever be high, bleeding will be neceffary; but this in very young children ought always to be fparingly performed. It is an evacuation which they bear the worft of any. Purging, vomiting, or fweating agree much better with them, and are generally more beneficial. Harris, however, obferves, that when an inflammation appears, the phyfician will labour in vain, if the *cure* be not begun with applying a leech under each ear. If the child be feized with convultion fits, a blifteringplafter may be applied between the fhoulders, or one behind each ear.

Sydenham fays, that in fevers occafioned by teething, he never found any remedy fo effectual as two, three, or four drops of fpirits of hartfhorn in a fpoonful of fimple water, or other convenient vehicle, given every four hours. The number of dofes may be four, five, or fix. I have often prefcribed this medicine with fuccefs, but always found a larger dofe neceffary. It may be given from five drops to fifteen or twenty, according to the age of the child, and when coffivenefs does not forbid it, three or four drops of laudanum may be added to each dofe.

In Scotland, it is very common, when children are cutting their teeth, to put a fmall Burgundy-pitch plafter between their fhoulders. This generally eafes the tickling cough which attends teething, and is by no means an ufelefs application. When the teeth are cut with difficulty, it ought to be kept on during the whole time of teething. It may be enlarged as occafion requires, and ought to be renewed at leaft once a fortnight.

Several things have been recommended for rubbing the gums, as oils, mucilages, &c.; but from thefe much is not to be expected. If any thing of this kind is to be ufed, we would recommend a little fine honey, which may be rubbed on with the finger three or four times a-day. Children are generally at this time difpofed to chew whatever they get into their hands. For this reafon they ought never to be without fomewhat that will yield a little to the preffure of their gums, as a cruft of bread, a wax candle, a bit of liquorice-root, or fuch like.

With regard to cutting the gums, we have feldom known it of any great benefit. In obflinate cafes, however,

ever, it ought to be tried. It may be performed by the finger-nail, the edge of a fixpenny piece that is worn thin, or any fharp body which can be with fafety introduced into the mouth; but the lancet, in a fkilful hand, is certainly the most proper.

In order to render the teething lefs difficult, parents ought to take care that their children's food be light and wholefome, and that their nerves be braced by fufficient exercife without doors, the ufe of the cold bath, &c. Were thefe things duly regarded, they would have a much better effect than *teething necklaces*, or other nonfenfical amulets, worn for that purpofe.

OF THE RICKETS.

This difeafe generally attacks children between the age of nine months and two years. It appeared first in England about the time when manufactures began to flourish, and still prevails most in towns where the inhabitants follow sedentary employments, by which means they neglect either to take proper exercise themselves, or to give it to their children.

CAUSES.——One caufe of the rickets is difeafed parents. Mothers of a weak relaxed habit, who neglect exercife, and live upon weak watery diet, can neither be expected to bring forth ftrong and healthy children, or to be able to nurfe them, after they are brought forth. Accordingly we find, that the children of fuch women generally die of the rickets, the fcrophula, confumptions, or fuch-like difeafes. Children begotten by men in the decline of life, who are fubject to the gout, the gravel, or other chronic difeafes, or who have been often afflicted with the venereal difeafe in their youth, are likewife very liable to the rickets.

Any diforder that weakens the conflicution or relaxes the habit of children, as the fmall-pox, meafles, teething, the hooping-cough, &c. difpofes them to this difeafe. It may likewife be occafioned by improper diet, as food that is either too weak and watery, or fo vifcid that the ftomach cannot digeft it.

Bad nurfing is the chief caule of this difeafe. When the nurfe is either difeafed, or has not enough of milk to L 1 3 nourifh

OF THE RICKETS.

nourish the child, it cannot thrive. But children suffer oftener by want of care in nurses, than want of food. Allowing an infant to lie or fit too much, or not keeping it thoroughly clean in its clothes, has the most pernicious effects.

The want of free air is likewife very hurtful to children in this refpect. When a nurfe lives in a clofe fmall houfe, where the air is damp and confined, and is too indolent to carry her child abroad into the open air, it will hardly escape this difease. A healthy child should always be in motion, unless when assess; if it be fuffered to lie or fit, instead of being toffed and dandled about, it will not thrive.

SYMPTOMS. ——At the beginning of this difeafe the child's flefh grows foft and flabby; its ftrength is diminifhed; it lofes its wonted cheerfulnefs, looks more grave and composed than is natural for its age, and does not choofe to be moved. The head and belly become too large in proportion to the other parts; the face appears full, and the complexion florid. Afterwards the bones begin to be affected, especially in the more foft and spungy parts. Hence the wrifts and ancles become thicker than ufual; the fpine or back-bone puts on an unnatural fhape; the breaft is likewife often deformed; and the bones of the arms and legs grow crooked. All thefe fymptoms vary according to the violence of the difeafe. The pulfe is generally quick, but feeble; the appetite and digeftion for the most part bad; the teeth come flowly and with difficulty, and they often rot and fall out afterwards. Rickety children generally have great acutenefs of mind, and an understanding above their years. Whether this is owing to their being more in the company of adults than other children, or to the preternatural enlargement of the brain, is not material.

REGIMEN. — As this difeafe is always attended with evident figns of weaknefs and relaxation, our chief aim in the cure muft be to brace and ftrengthen the folids, and to promote digeftion and the due preparation of the fluids. Thefe important ends will be belt anfwered by wholefome nourifhing diet, fuited to the age and ftrength of the patient, open dry air, and fufficient exercife. If the child has a bad nurfe, who either neglects her

her duty, or does not understand it, she should be changed. If the feason be cold, the child ought to be kept warm; and when the weather is hot, it ought to be kept cool; as sweating is apt to weaken it, and too great a degree of cold has the same effect. The limbs should be rubbed frequently with a warm hand, and the child kept as cheerful as possible.

The diet ought to be dry and nourifhing, as good bread, roafted flefh, &c. Bifcuit is generally reckoned the beft bread; and pigeons, pullets, veal, rabbits, or mutton roafted or minced, are the moft proper flefh. If the child be too young for flefh meats, he may have rice, millet, or pearl-barley, boiled with raifins, to which may be added a little wine and fpice. His drink may be good claret, mixed with an equal quantity of water. Thofe who cannot afford claret, may give the child now and then a wine-glafs of mild ale, or good porter.

MEDICINE. — Medicines are here of little avail. The difeafe may often be cured by the nurfe, but feldom by the phyfician. In children of a groß habit, gentle vomits and repeated purges of rhubarb may fometimes be of ufe, but they will feldom carry off the difeafe; that muft depend chiefly upon fuch things as brace and ftrengthen the fyftem : for which purpofe, befides the regimen mentioned above, we would recommend the cold bath, efpecially in the warm feafon. It muft however be ufed with prudence, as fome rickety children cannot bear it. The beft time for ufing the cold bath is in the morning, and the child fhould be well rubbed with a dry cloth immediately after he comes out of it. If the child fhould be weakened by the cold bath, it muft be difcontinued.

Sometimes iffues have been found beneficial in this difeafe. They are peculiarly neceffary for children who abound with grofs humours. An infufion of the Peruvian bark in wine or ale would be of fervice, were it poffible to bring children to take it. We might here mention many other medicines which have been recommended for the rickets; but as there is far more danger in trufting to thele than in neglecting them altogether, we choofe rather to pafs them over, and to recommend a proper regimen as the thing chiefly to be depended on.

OF CONVULSIONS.

Though more children are faid to die of convultions than of any other difeafe, yet they are for the molt part only a fymptom of fome other malady. Whatever greatly irritates or flimulates the nerves, may occafion convultions. Hence infants whofe nerves are eafily affected, are often thrown into convultions by any thing that irritates the alimentary canal : likewife by teething, flrait clothes, the approach of the fmall-pox, meafles, or other eruptive difeafes.

When convultions proceed from an irritation of the ftomach or bowels, whatever clears them of their acrid contents, or renders thefe mild and inoffenfive, will generally perform a cure; wherefore, if the child be coftive, the beft way will be to begin with a clyfter, and afterwards to give a gentle vomit, which may be repeated occafionally, and the body in the mean time kept open by gentle dofes of *magnefia alba*, or fmall quantities of rhubarb mixed with the powder of crabs' claws.

Convultions which precede the eruption of the fmallpox or meafles, generally go off upon thefe making their appearance. The principal danger in this cafe arifes from the fears and apprehentions of thofe who have the care of the patient. Convultions are very alarming, and fomething must be done to appeale the affrighted parents, nurfes, &c. Hence the unhappy infant often undergoes bleeding, bliftering, and feveral other operations, to the great danger of its life, when a little time, bathing the feet in warm water, and throwing in a mild clyfter, would have fet all to rights.

When convultion fits arife from the cutting of teeth, befides gentle evacuations, we would recommend bliftering, and the ufe of antifpafmodic medicines, as the tincture of foot, afafœtida, or caftor. A few drops of any of thefe may be mixed in a cup of white-wine whey, and given occafionally.

When convultions proceed from any external caufe, as the preffure occationed by ftrait clothes or bandages, &c. thefe ought immediately to be removed, though in this cafe

OF CONVULSIONS.

cafe taking away the caufe will not always remove the effect, yet it ought to be done. It is not likely that the patient will recover, as long as the caufe which first gave rife to the diforder continues to act.

When a child is feized with convultions, without having any complaint in the bowels, or fymptoms of teething; or any rash or other discharge which has been fuddenly dried up; we have reafon to conclude that it is a primary difeafe, and proceeds immediately from the brain. Cafes of this kind, however, happen but feldom; which is very fortunate, as little can be done to relieve the unhappy patient. When a difease proceeds from an original fault in the formation or ftructure of the brain itfelf, we cannot expect that it fhould yield to medicine. But as this is not always the caufe, even of convultions which proceed immediately from the brain, fome attempts fhould be made to remove them. The chief intention to be purfued for this purpofe, is to make fome derivation from the head, by bliftering, purging, and the like. Should thefe fail, iffues or fetons may be put in the neck, or between the fhoulders.

OF WATER IN THE HEAD.

Though water in the head, or a dropfy of the brain, may affect adults as well as children, yet, as the latter are more peculiarly liable to it, we thought it would be most proper to place it among the difeases of infants.

CAUSES.—A dropfy of the brain may proceed from injuries done to the brain itfelf by falls, blows, or the like; it may likewife proceed from an original laxity or weaknefs of the brain; from fchirrous tumours or excrefcences within the fkull; a thin watery flate of the blood; a diminifhed fecretion of urine; a fudden check of the perfpiration; and laftly, from tedious and lingering difeafes, which wafte and confume the patient.

SYMPTOMS.——This difeafe has at first the appearance of a flow fever; the patient complains of a pain in the crown of his head, or over his eyes; he shuns the light; is fick, and sometimes vomits; his pulse is irregular and generally low; though he seems heavy and dull, yet he does not seep; he is sometimes delirious, and

522 OF WATER IN THE HEAD.

and frequently fees objects double; towards the end of this commonly fatal difeafe, the pulfe becomes more frequent, the pupils are generally dilated, the cheeks flufhed, the patient becomes comatofe, and convultions enfue*.

MEDICINE. — No medicine has hitherto been found fufficient to carry off a dropfy of the brain. It is laudable, however, to make fome attempts, as time or chance may bring many things to light, of which at prefent we have no idea. The medicines generally ufed are, purges of rhubarb or jalap, with calomel, and bliftering-plafters applied to the neck or back-part of the head. To which we would beg leave to add diuretics, or medicines which promote the fecretion of urine, fuch as are recommended in the common dropfy. A difcharge from the nofe ought likewife to be promoted by caufing the patient to fnuff the powder of afarum, white hellebore, or the like.

Some practitioners have of late pretended to cure this difeafe by the ufe of mercury. I have not been fo happy as to fee any inftances of a cure being performed in a confirmed dropfy of the brain; but in fo defperate a malady every thing deferves a trial \dagger .

To look over the long catalogue of infantile difeafes in fome medical books, one would be inclined to think, that the real defign of the authors, though concealed under the flow of precifion, was to fpread alarm through every family. I have had a very different object in view, to quiet the fears of parents, to direct their attention to the proper treatment of their children, and thus to render the use of any medicines almost unneceffary. I have shewn the folly of having recourse to physic to

* I very lately loft a patient in this difeafe, where a curious *metaftafis* feemed to take place. The water at first appeared to be in the abdomen, afterwards in the breast, and last of all it mounted up to the brain, where it soon proved fatal.

+ One reafon why this difeafe is feldom or never cured, may be, that it is feldom known till too far advanced to admit of remedy. Did parents watch the firft fymptoms, and call a phyfician in due time, I am inclined to think that fomething might be done. But thefe fymptoms are not yet fufficiently known, and are often mistaken even by phyficians themfelves. Of this I lately faw a striking instance in a patient, attended by an eminent practitioner of this city, who had all along mistaken the difease for teething. 8* bring

OF WATER IN THE HEAD.

bring away the black, vifcid, fyrup-like fubstance contained in the inteffines of a new-born infant, when the purgative quality of his mother's milk is fo admirably fuited to that very purpofe. The new milk is thin and waterifh, but acquires every day greater confiftence, and thus affords a more folid aliment to the child, as he becomes more capable of digefting it. If the mother does not vitiate by her own improper diet the pure fountains of nourifhment and health which nature has kindly given her, the child will neither be troubled with coffiveness nor gripes. He will escape those complaints of the ftomach which are occafioned by fwallowing crude, inflammatory trafh, or still more pernicious drugs. The daily use of the cold bath, and frequent exercise in the open air, will not only preferve him from colds and defluxions, but from all the diforders which are the confequences of relaxation and of nervous irritability. A child brought up in the manner I have recommended, will have little to fear even from external contagion. The firm texture of his fkin, like a fhield, will almost refift its approach, and the purity of his habit will correct its malignity. The fmallpox is the only infectious difeafe for which I would have him prepared by any particular process, because that procels is fo eafy and certain, befides affording a perfect command both of time and circumstances.

CHAP. L.

OF SURGERY.

TO defcribe all the operations of furgery, and to point out the different difeafes in which these operations are neceffary, would extend this article far beyond the limits allotted to it : we must therefore confine our obfervations to fuch cases as most generally occur, and in which proper affistance is either not asked, or not always to be obtained.

Though an acquaintance with the structure of the human body is indifpenfably necessary to qualify a man for

OF SURGERY.

for being an expert furgeon; yet many things may be done to fave the lives of their fellow-men in emergencies by those who are no adepts in anatomy. It is amazing with what facility the peasants daily perform operations upon brute animals, which are not of a less difficult nature than many of those performed on the human species: yet they feldom fail of fucces.

Indeed every man is in fome measure a furgeon, whether he will or not. He feels an inclination to affift his fellow-men in diffrefs, and accidents happen every hour which give occasion to exercise this feeling. The feelings of the heart, however, when not directed by the judgment, are apt to mislead. Thus one, by a rash attempt to fave his friend, may fometimes destroy him; while another, for fear of doing amiss, stands still and sees his bosom-friend expire without so much as attempting to relieve him, even when the means are in his power. As every good man would wish to steer a course different from either of these, it will no doubt be agreeable to him to know what ought to be done upon fuch emergencies.

OF BLEEDING.

No operation of furgery is fo frequently neceffary as bleeding; it ought therefore to be very generally underftood. But though practifed by midwives, gardeners, blackfmiths, &cc. we have reafon to believe that very few know when it is proper. Even phyficians themfelves have been fo much the dupes of theory in this article, as to render it the fubject of ridicule. It is, however, an operation of great importance, and must, when feafonably and properly performed, be of fingular fervice to those in diffres.

Bleeding is proper at the beginning of all inflammatory fevers, as pleurifies, peripneumonies, &c. It is likewife proper in all topical inflammations, as those of the inteftines, womb, bladder, ftomach, kidneys, throat, eyes, &c. as also in the afthma, fciatic pains, coughs, head-achs, rheumatifms, the apoplexy, epilepfy, and bloody flux. After falls, blows, bruifes, or any violent hurt received either externally or internally, bleeding is neceffary. It is likewife neceffary for perfons who have had

had the misfortune to be ftrangled, drowned, fuffocated with foul air, the fumes of metal, or the like. In a word, whenever the vital motions have been fuddenly ftopped from any caufe whatever, except in fwoonings occafioned by mere weaknefs or hyfteric affections, it is proper to open a vein. But in all diforders proceeding from a relaxation of the folids, and an impoverifhed ftate of the blood, as dropfies, cacochymies, &c. bleeding is improper.

Bleeding for topical inflammations ought always to be performed as near the part affected as poffible. When this can be done with a lancet, it is to be preferred to any other method; but where a vein cannot be found, recourfe must be had to leeches or cupping.

The quantity of blood to be let must always be regulated by the strength, age, constitution, manner of life, and other circumstances relating to the patient. It would be ridiculous to suppose that a child could bear to lose as much blood as a grown person, or that a delicate lady should be bled to the same extent as a robust man.

From whatever part of the body blood is to be let, a bandage must be applied between that part and the heart. As it is often necessary, in order to raife the vein, to make the bandage pretty tight, it will be proper in fuch cafes, as foon as the blood begins to flow, to flacken it a little. The bandage ought to be applied at least an inch, or an inch and a half from the place where the wound is intended to be made.

Perfons not skilled in anatomy ought never to bleed in a vein that lies over an artery or a tendon, if they can avoid it. The former may easily be known from its pulfation or beating, and the latter from its feeling hard or tight like a whipcord under the finger.

It was formerly a rule, even among those who had the character of being regular practitioners, to bleed their patients in certain difeases till they fainted. Surely a more ridiculous rule could not be proposed. One perfon will faint at the very fight of a lancet, while another will lose almost the whole blood of his body before he faints. Swooning

OF BLEEDING.

Swooning depends more upon the flate of the mind than of the body: befides, it may often be occafioned or prevented by the manner in which the operation is performed.

Children are generally bled with leeches. This, though fometimes neceffary, is a very troublefome and uncertain practice. It is impoffible to know what quantity of blood is taken away by leeches; befides, the bleeding is often very difficult to ftop, and the wounds are not eafily healed. Would those who practife bleeding take a little more pains, and accuftom themselves to bleed children, they would not find it fuch a difficult operation as they imagine.

Certain hurtful prejudices with regard to bleeding ftill prevail among the country people. They talk, for instance, of head-veins, heart-veins, breast-veins, &c. and believe that bleeding in thefe will certainly cure all difeafes of the parts from whence they are fuppoled to come, without confidering that all the blood veffels arife from the heart, and return to it again; for which reafon, unlefs in topical inflammations, it fignifies very little from what part of the body blood is taken. But this, though a foolifh prejudice, is not near fo hurtful as the vulgar notion that the first bleeding will perform wonders. This belief makes them often postpone the operation when neceffary, in order to referve it for fome more important occafion, and, when they think themfelves in extreme danger, they fly to it for relief, whether it be proper or not. Bleeding at certain stated periods or feasons has likewife bad effects.

It is a common notion that bleeding in the feet draws the humours downwards, and confequently cures difeafes of the head and other fuperior parts; but we have already obferved that, in all topical affections, the blood ought to be drawn as near the part as possible. When it is neceffary, however, to bleed in the foot or hand, as the veins are small, and the bleeding is apt to stop too foon, the part ought to be immerfed in warm water, and kept there till a fufficient quantity of blood be let.

We

We fhall not fpend time in defcribing the manner of performing this operation : that will be better learned by example than precept. Twenty pages of defcription would not convey fo juft an idea of the operation as feeing it once performed by an expert hand. Neither is it neceffary to point out the different parts of the body from whence blood may be taken, as the arm, foot, forehead, temples, neck, &cc. Thefe will readily occur to every intelligent perfon, and the foregoing obfervations will be fufficient for determining which of them is most proper upon any particular occasion. In all cafes where the intention is merely to leffen the general mass of blood, the arm is the most commodious part of the body in which the operation can be performed.

OF INFLAMMATIONS AND ABSCESSES.

From whatever caufe an inflammation proceeds, it must terminate either by difperfion, fuppuration, or gangrene. Though it is impossible to foretel with certainty in which of these ways any particular inflammation will terminate, yet a probable conjecture may be formed with regard to the event, from a knowledge of the patient's age and constitution. Inflammations happening in a flight degree upon colds, and without any previous indifposition, will most probably be dispersed; those which follow close upon a fever, or happen to persons of a gross habit of body, will generally suppurate; and those which attack very old people, or persons of a dropfical habit, will have a strong tendency to gangrene.

If the inflammation be flight, and the conftitution found, the difperfion ought always to be attempted. This will be beft promoted by a flender diluting diet, plentiful bleeding, and repeated purges. The part itfelf must be fomented, and, if the fkin be very tenfe, it may be embrocated with a mixture of three-fourths of fweet oil, and one-fourth of vinegar, and afterwards covered with a piece of wax-plaster.

If, notwithstanding these applications, the symptomatic fever increases, and the tumour becomes larger, with violent

A.

528 OF INFLAMMATIONS AND ABCESSES.

violent pain and pulfation, it will be proper to promote the fuppuration. The beft application for this purpofe is a foft poultice, which may be renewed twice a-day. If the fuppuration proceeds but flowly, a raw onion cut fmall or bruifed may be fpread upon the poultice. When the abfcefs is ripe or fit for opening, which may eafily be known from the thinnefs of the fkin in the most prominent part of it, fluctuation of matter, which may be felt under the finger, and, generally speaking, an abatement of the pain, it may be opened either with a lancet, or by means of caustic.

The laft way in which an inflammation terminates, is in a gangrene or mortification, the approach of which may be known by the following fymptoms: the inflammation lofes its rednefs, and becomes dufkifh or livid; the tenfion of the fkin goes off, and it feels flabby; little bladders filled with ichor of different colours fpread all over it; the tumour fubfides, and from a dufkifh complexion becomes black; a quick low pulfe, with cold clammy fweats, are the immediate forerunners of death.

When these fymptoms first appear, the part ought to be dreffed with London treacle, or a cataplasm made of lixivium and bran. Should the fymptoms become worse, the part must be fearified, and afterwards dreffed with bafilicum softened with oil of turpentine. All the dreffings must be applied warm. With regard to internal medicines, the patient must be supported with generous cordials, and the Peruvian bark exhibited in as large doses as the stomach will bear it. If the mortified parts should feparate, the wound will become a common ulcer, and must be treated accordingly.

This article includes the treatment of all those difeases, which, in different parts of the country, go by the names of *biles*, *impostbumes*, *whitloes*, &c. They are all abfeesses in confequence of a previous inflammation, which, if possible, ought to be discussed in the second not be done, the suppuration should be promoted, and the matter discharged by an incision, if necessary; afterwards the fore may be dreffed with yellow basilicum, or some other digestive ointment.

OF

No part of medicine has been more miftaken than the treatment or cure of wounds. Mankind in general believe that certain herbs, ointments, and plafters, are poffeffed of wonderful healing powers, and imagine that no wound can be cured without the application of them. It is, however, a fact, that no external application whatever contributes towards the cure of a wound, any other way than by keeping the parts foft, clean, and defending them from the external air, which may be as effectually done by dry lint, as by the moft pompous applications, while it is exempt from many of the bad confequences attending them.

The fame obfervation holds with refpect to internal applications. Thefe only promote the cure of wounds as far as they tend to prevent a fever, or to remove any caufe that might obftruct or impede the operations of Nature. It is Nature alone that cures wounds. All that art can do is to remove obftacles, and to put the parts in fuch a condition as is the moft favourable to Nature's efforts.

With this fimple view we fhall confider the treatment of wounds, and endeavour to point out fuch steps as ought to be taken to facilitate their cure.

The first thing to be done, when a perfon has received a wound, is to examine whether any foreign body be lodged in it, as wood, stone, iron, lead, glass, dirt, bits of cloth, or the like. These, if possible, ought to be extracted, and the wound cleaned before any dressings be applied. When that cannot be effected with fastery, on account of the patient's weakness or loss of blood, they must be suffered to remain in the wound, and afterwards extracted when he is more able to bear it.

When a wound penetrates into any of the cavities of the body, as the breaft, the bowels, &c. or where any confiderable blood-veffel is cut, a fkilful furgeon ought immediately to be called, otherwife the patient may lofe his life. But fometimes the difcharge of blood is fo great, that if it be not ftopped, the patient may die, even be-M m

fore a furgeon, though at no great diffance, can arrive. In this cafe, fomething muft be done by thofe who are prefent. If the wound be in any of the limbs, the bleeding may generally be ftopped by applying a tight ligature or bandage round the member a little above the wound. The beft method of doing this is to put a ftrong broad garter round the part, but fo flack as eafily to admit a fmall piece of flick to be put under it, which muft be twifted, in the fame manner as a countryman does a cart-rope to fecure his loading, till the bleeding ftops. Whenever this is the cafe, he muft take care to twift it no longer, as ftraining it too much might occafion an inflammation of the parts, and endanger a gangrene.

In parts where this bandage cannot be applied, various other methods may be tried to ftop the bleeding, as the application of ftyptics, aftringents, &c. Cloths dipped in a folution of blue vitriol in water, or the *ftyptic water* of the difpenfatories, may be applied to the wound. When these cannot be obtained, ftrong fpirits of wine may be used. Some recommend the *agaric* * of the oak as preferable to any of the other ftyptics; and indeed it deferves confiderable encomiums. It is eafily obtained, and ought to be kept in every family, in case of accidents. A piece of it must be laid upon the wound, and covered with a good deal of lint, above

* Dr. Tiffot, in his Advice to the People, gives the following directions for gathering, preparing, and applying the agaric .--"Gather in autumn," fays he, " while the fine weather lafts, the agaric of the oak, which is a kind of fungus or excrefcence iffuing from the wood of that tree. It confilts at first of four parts, which prefent themfelves fucceffively: 1 The outward rind or fkin, which may be thrown away. 2. The part immediately under this rind, which is the beft of all. This is to be beat well with a hammer, till it becomes foft and very pliable. This is the only preparation it requires, and a flice of it of a proper fize is to be applied directly over the burfting open blood-veffels. It conftringes and brings them clofe together, ftops the bleeding, and generally falls off at the end of two days. 3. The third part adhering to the fecond, may ferve to ftop the bleeding from the fmaller veffels; and the fourth and laft part may be reduced to powder, as conducing to the fame purpose."-Where the agaric cannot be had, fponge may be ufed in its ftead. It must be applied in the fame manner, and has nearly the fame effects.

which a bandage may be applied fo tight as to keep it firmly on.

Though fpirits, tinctures, and hot balfams may be ufed, in order to ftop the bleeding when it is exceffive, they are improper at other times. They do not promote but retard the cure, and often change a fimple wound into an ulcer. People imagine, becaufe hot balfams congeal the blood, and feem, as it were, to folder up the wound, that they therefore heal it; but this is only a deception. They may indeed ftop the flowing blood, by fearing the mouths of the veffels; but, by rendering the parts callous, they obftruct the cure.

In flight wounds, which do not penetrate much deeper than the fkin, the beft application is a bit of the common black flicking plafter. This keeps the fides of the wound together, and prevents the air from hurting it, which is all that is neceffary. When a wound penetrates deep, it is not fafe to keep its lips quite clofe: this keeps in the matter, and is apt to make the wound fefter. In this cafe the beft way is to fill the wound with foft lint, commonly called *caddis*. It, however, muft not be fluffed in too hard, otherwife it will do hurt. The lint may be covered with a cloth dipped in oil, or fpread with the common wax plafter *; and the whole muft be kept on by a proper bandage.

We shall not spend time in describing the different bandages that may be proper for wounds in different parts of the body; common sense will generally suggest the most commodious method of applying a bandage; besides, descriptions of this kind are not easily understood or remembered.

The first dreffing ought to continue on for at least two days; after which it may be removed, and fresh lint applied as before. If any part of the first dreffing flicks so close as not to be removed with ease or fastery to the patient, it may be allowed to continue, and fresh lint dipped in fweet oil laid over it. This will soften it, so as to make it come off easily at next dreffing. Afterwards the wound may be dreffed twice a-day in the same

> • See Appendix, Wax Plaster. M m 2

1:25

manner

532

TOCOTORE

manner till it be quite healed. Those who are fond of falves or ointments, may, after the wound is become very superficial, dress it with the yellow *basilicum**; and if fungous, or what is called *proud fless*, should rife in the wound, it may be checked, by mixing with the ointment a little burnt alum or red precipitate of mercury.

When a wound is greatly inflamed, the most proper application is a poultice of bread and milk, fostened with a little fweet oil or fresh butter. This must be applied instead of a plaster, and should be changed twice a-day.

If the wound be large, and there is reafon to fear an inflammation, the patient flould be kept on a very low diet. He must abstain from flesh, strong liquors, and every thing that is of a heating nature. If he be of a full habit, and has lost but little blood from the wound, he must be bled; and, if the symptoms be urgent, the operation may be repeated. But when the patient has been greatly weakened by loss of blood from the wound, it will be dangerous to bleed him, even though a fever should ensue. Nature should never be too far exhausted. It is always more fase to allow her to struggle with the difease in her own way, than to fink the patient's strength by excession.

Wounded perfons ought to be kept perfectly quiet and eafy. Every thing that ruffles the mind, or moves the paffions, as love, anger, fear, exceflive joy, &c. are very hurtful. They ought, above all things, to abftain from venery. The body fhould be kept gently open, either by laxative clyfters, or by a cool vegetable diet, as roafted apples, flewed prunes, boiled fpinage, and fuch like.

OF BURNS.

In flight burns, which do not break the fkin, it is cuftomary to hold the part near the fire for a competent time, to rub it with falt, or to lay a compress upon it, dipped in fpirits of wine or brandy. But when the burn

· See Appendix, Tellow Bafilicum.

has

OF BURNS.

has penetrated fo deep, as to blifter or break the fkin, it must be dreffed with fome of the liniments for burns mentioned in the Appendix, or with the emollient and gently-drying ointment, commonly called *Turner's ce*rate*. This may be mixed with an equal quantity of fresh olive oil, and spread upon a fost rag, and applied to the part affected. When this ointment cannot be had, an egg may be beat up with about an equal quantity of the sweetest falled oil. This will ferve very well, till a proper ointment can be prepared. When the burning is very deep, after the first two or three days, it should be dreffed with equal parts of yellow basilicum and Turner's cerate, mixed together.

When the burn is violent, or has occafioned a high degree of inflammation, and there is reafon to fear a gangrene or mortification, the fame means mult be ufed to prevent, as are recommended in other violent inflammations. The patient, in this cafe, must live low, and drink freely of weak diluting liquors. He must likewife be bled, and have his body kept open. But if the burnt parts should become livid or black, with other symptoms of mortification, it will be necessary to bathe them frequently with warm camphorated spirits of wine, tincture of myrrh, or other antifeptics, mixed with a decoction of the bark. In this cafe the bark must likewife be taken internally, and the patient's diet must be more generous, with wine, &c.

As example teaches better than precept, I shall relate the treatment of the most dreadful cafe of this kind that has occurred in my practice. A middle-aged man, of a good constitution, fell into a large vessel full of boiling water, and miserably scalded about one half of his body. As his clothes were on; the burning in some parts was very deep before they could be got off. For the first two days the scalded parts had been frequently anointed with a mixture of lime-water and oil, which is a very proper application for recent burnings. On the third day, when I first faw him, his fever was high, and his body costive, for which he was bled, and had an emol-

* See Appendix, Turner's Cerate.

lient

OF BURNS.

lient clyfter administered. Poultices of bread and milk, fostened with fresh butter, were likewise applied to the affected parts, to abate the heat and inflammation. His fever still continuing high, he was bled a second time, was kept strictly on the cooling regimen, took the faline mixture with small doses of nitre, and had an emollient clyfter administered once a-day. When the inflammation began to abate, the parts were dressed with a digestive composed of brown cerate and yellow basilicum. Where any black spots appeared, they were flightly scarified, and touched with the tincture of myrrh; and, to prevent their spreading, the Peruvian bark was administered. By this course, the man was so well in three weeks as to be able to attend his business.

OF BRUISES.

Bruifes are generally productive of worfe confequences than wounds. The danger from them does not appear immediately, by which means it often happens that they are neglected. It is needlefs to give any definition of a difeafe fo univerfally known; we fhall therefore proceed to point out the method of treating it.

In flight bruifes it will be fufficient to bathe the part with warm vinegar, to which a little brandy or rum may occafionally be added, and to keep cloths wet with this mixture conftantly applied to it. This is more proper than rubbing it with brandy, fpirits of wine, or other ardent fpirits, which are commonly ufed in fuch cafes.

In fome parts of the country the peafants apply to a recent bruife a cataplasm of fresh cow-dung. I have often seen this cataplasm applied to violent contustions, occasioned by blows, falls, bruises, and such like, and never knew it fail to have a good effect.

When a bruife is very violent, the patient ought immediately to be bled, and put upon a proper regimen. His food fhould be light and cool, and his drink weak, and of an opening nature; as whey fweetened with honey, decoctions of tamarinds, barley, cream-tartar-whey, and fuch like. The bruifed part must be bathed with vinegar

OF BRUISES.

vinegar and water, as directed above; and a poultice made by boiling crumb of bread, elder-flowers, and camomile flowers, in equal quantities of vinegar and water, applied to it. This poultice is peculiarly proper when a wound is joined to the bruife. It may be renewed two or three times a-day.

As the ftructure of the veffels is totally deftroyed by a violent bruife, there often enfues a great lofs of fubftance, which produces an ulcerous fore very difficult to cure. If the bone be affected, the fore will not heal before an exfoliation takes place; that is, before the difeafed part of the bone feparates, and comes out through the wound. This is often a very flow operation, and may even require feveral years to be completed. Hence it happens, that thefe fores are frequently miftaken for the king's evil, and treated as fuch, though in fact they proceed folely from the injury which the folid parts received from the blow.

Patients in this fituation are peftered with different advices. Every one who fees them propofes a new remedy, till the fore is fo much irritated with various and oppofite applications, that it is often at length rendered abfolutely incurable. The beft method of managing fuch fores is, to take care that the patient's conflictution does not fuffer by confinement or improper medicine, and to apply nothing to them befides fimple ointment foread upon foft lint, over which a poultice of bread and milk, with boiled camomile-flowers, or the like, may be put, to nourifh the part, and keep it foft and warm. Nature, thus affifted, will generally in time operate a cure, by throwing off the difeafed parts of the bone, after which the fore foon heals.

OF ULCERS.

Ulcers may be the confequence of wounds, bruifes, or impofthumes improperly treated; they may likewife proceed from an ill state of the humours, or what may be called a bad habit of body.

In the latter cafe, they ought not to be haftily dried up, otherwife it may prove fatal to the patient. Ulcers happen most commonly in the decline of life; and perfons who neglect exercise, and live grossly, are most M m 4 liable

OF ULCERS.

liable to them. They might often be prevented by retrenching fome part of the folid food, or by opening artificial drains, as iffues, fetons, or the like.

An ulcer may be diffinguished from a wound by its discharging a thin watery humour, which is often so acrid as to inflame and corrode the skin; by the hardness and perpendicular situation of its sides or edges; by the time of its duration, &c.

It requires confiderable skill to be able to judge whether-or not an ulcer ought to be dried up. In general, all ulcers which proceed from a bad habit of body, should be fuffered to continue open, at least till the conflitution has been fo far changed by proper regimen, or the use of medicine, that they feem disposed to heal of their own accord. Ulcers which are the effect of malignant fevers, or other acute difeafes, may generally be healed with fafety after the health has been reftored for fome time. The cure ought not, however, to be attempted too foon, nor at any time without the use of purging medicines and a proper regimen. When wounds or bruifes have, by wrong treatment, degenerated into ulcers, if the conflitution be good, they may generally be healed with fafety. When ulcers either accompany chronical difeafes, or come in their ftead, they must be cautioufly healed. If an ulcer conduces to the patient's health, from whatever caufe it proceeds, it ought not to be healed; but if, on the contrary, it waftes the ftrength, and confumes the patient by a flow fever, it fhould be healed as foon as poffible.

We would earneftly recommend a first attention to thefe particulars to all who have the misfortune to labour under this diforder, particularly perfons in the decline of life; as we have frequently known people throw away their lives by the want of it, while they were extolling and generoufly rewarding those whom they ought to have looked upon as their executioners.

The most proper regimen for promoting the cure of ulcers, is to avoid all spices, falted and high-seasoned food, all strong liquors, and to lessen the usual quantity of steff meat. The body ought to be kept gently open by a diet confissing chiefly of cooling laxative vegetables, and by drinking butter-milk, whey sweetened with honey, or the like.

OF ULCERS.

like. The patient ought to be kept cheerful, and fhould take as much exercife as he can eafily bear.

When the bottom and fides of an ulcer feem hard and callous, they may be fprinkled twice a-day with a little red precipitate of mercury, and afterwards dreffed with the yellow *bafilicum* ointment. Sometimes it will be neceffary to have the edges of the ulcer fcarified with the lancet.

Lime-water has frequently been known to have very happy effects in the cure of obftinate ulcers. It may be ufed in the fame manner as directed for the ftone and gravel.

My late learned and ingenious friend Dr. White ftrongly recommends the ufe of the folution of corrofive fublimate of mercury in brandy, for the cure of obftinate illconditioned ulcers. I have frequently found this medicine, when given according to the Doctor's directions, prove very fuccefsful. The dofe is a table-fpoonful night and morning; at the fame time wafhing the fore twice or thrice a-day with it. In a letter which I had from the Doctor a little before his death, he informed me, "That he obferved wafhing the fore thrice a-day with the folution of a triple ftrength was very beneficial *."

A fiftulous ulcer can feldom be cured without an operation. It must either be laid open fo as to have its callous parts destroyed by fome corrosive application, or they must be entirely cut away by the knife ; but as this operation requires the hand of an expert furgeon, there is no occasion to describe it. Ulcers about the anus are most apt to become fistulous, and are very difficult to cure. Some indeed pretend to have found Ward's fiftula paste very successful in this complaint. It is not a dangerous medicine, and being eafily procured, it may deferve a trial; but as these ulcers generally proceed from an ill habit of body, they will feldom yield to any thing except a long courfe of regimen, affifted by medicines, which are calculated to correct that particular habit, and to induce an almost total change in the conftitution.

* In ulcers of the lower limbs great benefit is often received from tight rollers, or wearing a laced flocking, as this prevents the flux of humours to the fores, and difpofes them to heal.

[538]

CHAP. LI.

OF DISLOCATIONS.

TATHEN a bone is moved out of its place or articulation fo as to impede its proper functions, it is faid to be luxated or diflocated. As this often happens to perfons in fituations where no medical affiftance can be obtained, by which means limbs, and even lives, are frequently loft, we shall endeavour to point out the method of reducing the most common luxations, and those which require immediate affiftance. Any perfon of common fenfe and refolution, who is prefent when a diflocation happens, may often be of more fervice to the patient, than the most expert furgeon can after the fwelling and inflammation have come on. When these are present, it is difficult to know the flate of the joint, and dangerous to attempt a reduction; and by waiting till they are gone off, the muscles become fo relaxed, and the cavity filled up, that the bone can never afterwards be retained in its place.

A recent diflocation may generally be reduced by extenfion alone, which muft always be greater or lefs according to the ftrength of the mufcles which move the joint, the age, robuftnefs, and other circumftances of the patient. When the bone has been out of its place for any confiderable time, and a fwelling or inflammation has come on, it will be neceffary to bleed the patient, and, after fomenting the part, to apply foft poultices with vinegar to it for fome time before the reduction is attempted.

All that is neceffary after the reduction, is to apply cloths dipt in vinegar or camphorated fpirits of wine to the part, and to keep it perfectly eafy. Many bad confequences proceed from the neglect of this rule. A diflocation feldom happens without the tendons and ligaments of the joint being ftretched and fometimes torn. When thefe are kept ealy till they recover their ftrength and tone, all goes on very well; but if the injury be increafed

OF DISLOCATIONS.

creafed by too frequent an exertion of the parts, no wonder if they be found weak and difeafed ever after.

DISLOCATION OF THE JAW.

The lower jaw may be luxated by yawning, blows, falls, chewing hard fubftances, or the like. It is eafily known from the patient's being unable to fhut his mouth, or to eat any thing, as the teeth of the under jaw do not correspond with those of the upper; befides, the chin either hangs down or is thrown toward one fide, and the patient is neither able to speak distinctly, nor to swallow without confiderable difficulty.

The ufual method of reducing a diflocated jaw, is to fet the patient upon a low ftool, fo as an affiftant may hold the head firm by preffing it againft his breaft. The operator is then to thruft his two thumbs, being firft wrapt up with linen cloths that they may not flip, as far back into the patient's mouth as he can, while his fingers are applied to the jaw externally. After he has got firm hold of the jaw, he is to prefs it ftrongly downwards and backwards, by which means the elapfed heads of the jaw may be eafily pufhed into their former cavities.

The peafants in fome parts of the country have a peculiar way of performing this operation. One of them puts a handkerchief under the patient's chin, then turning his back to that of the patient, pulls him up by the chin fo as to fuspend him from the ground. This method often fucceeds, but we think it a dangerous one, and therefore recommend the former.

DISLOCATION OF THE NECK.

The neck may be diflocated by falls, violent blows, or the like. In this cafe, if the patient receives no affiftance, he foon dies, which makes people imagine the neck was broken; it is, however, for the most part, only partially diflocated, and may be reduced by almost any perfon who has refolution enough to attempt it. A complete diflocation of the neck is inftantaneous death. When

540 DISLOCATION OF THE NECK.

When the neck is diflocated, the patient is immediately deprived of all fenfe and motion; his neck fwells, his countenance appears bloated; his chin lies upon his breaft, and his face is generally turned towards one fide.

To reduce this diflocation, the unhappy perfon fhould immediately be laid upon his back on the ground, and the operator muft place himfelf behind him fo as to be able to lay hold of his head with both hands, while he makes a refiftance by placing his knees against the patient's fhoulders. In this posture he muft pull the head with confiderable force, gently twifting it at the fame time, if the face be turned to one fide, till he perceives that the joint is replaced, which may be known from the noife which the bones generally make when going in, the patient's beginning to breathe, and the head continuing in its natural posture.

This is one of those operations which it is more easy to perform than describe. I have known instances of its being happily performed even by women, and often by men of no medical education. After the neck is reduced, the patient ought to be bled, and should be fuffered to rest for fome days, till the parts recover their proper tone.

DISLOCATION OF THE RIBS.

As the articulation of the ribs with the backbone is very firong, they are not often diflocated. It does, however, fometimes happen, which is a fufficient reafon for our taking notice of it. When a rib is diflocated either upwards or downwards, in order to replace it, the patient fhould be laid upon his belly on a table, and the operator muft endeavour to pufh the head of the bone into its proper place. Should this method not fucceed, the arm of the difordered fide may be fufpended over a gate or ladder, and while the ribs are thus ftretched afunder, the heads of fuch as are out of place may be thruft into their former fituation.

Those diflocations wherein the heads of the ribs are forced inwards, are both more dangerous and the most difficult to reduce, as neither the hand nor any instrument can be applied internally to direct the luxated heads of the

DISLOCATION OF THE RIBS. 541

the ribs. Almost the only thing that can be done is, to lay the patient upon his belly over a cask, or fome gibbous body, and to move the fore-part of the rib inward towards the back, fometimes shaking it; by this means the heads of the luxated ribs may flip into their former place.

DISLOCATION OF THE SHOULDER.

The humerus or upper-bone of the arm may be diflocated in various directions : it happens, however, moft frequently downwards, but very feldom directly upwards. From the nature of its articulation, as well as from its expolure to external injuries, this bone is the moft fubject to diflocation of any in the body. A diflocation of the humerus may be known by a depreffion or cavity on the top of the fhoulder, and an inability to move the arm. When the diflocation is downward or forward, the arm is elongated, and a ball or lump is perceived under the arm-pit ; but when it is backward, there appears a protuberance behind the fhoulder, and the arm is thrown forwards towards the breaft.

The ufual method of reducing diflocations of the fhoulder is to feat the patient upon a low ftool, and to caufe an affiftant to hold his body fo that it may not give way to the extension, while another lays hold of the arm a little above the elbow, and gradually extends it. The operator then puts a napkin under the patient's arm, and caufes it to be tied behind his own neck : by this, while a fufficient extension is made, he lifts up the head of the bone, and with his hands directs it into its proper place. There are various machines invented for facilitating this operation, but the hand of an expert furgeon is always more fafe. In young and delicate patients, I have generally found it a very eafy matter to reduce the fhoulder, by extending the arm with one hand, and thrufting in the head of the bone with the other. In making the extension, the arm ought always to be a little bent.

DISLOCATION OF THE ELBOW.

The bones of the fore-arm may be diflocated in any direction. When this is the cafe, a protuberance may be observed on that fide of the arm towards which the

bone

542 DISLOCATION OF THE ELBOW.

bone is pushed, from which, and the patient's inability to bend his arm, a diflocation of this joint may easily be known.

Two affiftants are generally neceffary for reducing a diflocation of the elbow; one of them muft lay hold of the arm above, and the other below the joint, and make a pretty ftrong extension, while the operator returns the bones into their proper place. Afterwards the arm muft be bent, and fuspended for fome time with a fling about the neck.

Luxations of the wrift and fingers are to be reduced in the fame manner as those of the elbow, viz. by making an extension in different directions, and thrusting the head of the bone into its place.

DISLOCATION OF THE THIGH.

When the thigh-bone is diflocated forward and downward, the knee and foot are turned out, and the leg is longer than the other; but when it is difplaced backward, it is ufually pufhed upward at the fame time, by which means the limb is fhortened, and the foot is turned inwards.

When the thigh-bone is difplaced forward and downward, the patient, in order to have it reduced, muft be laid upon his back, and made faft by bandages, or held by affiftants, while by others an extension is made by means of flings fixed about the bottom of the thigh a little above the knee. While the extension is made, the operator must push the head of the bone outward, till it gets into the focket. If the diflocation be outward, the patient must be laid upon his face, and, during the extension, the head of the bone must be pushed inward.

Diflocations of the *knees*, *ancles*, and *tces*, are reduced much in the fame manner as those of the upper extremities, *viz*. by making an extension in opposite directions, while the operator replaces the bones. In many cafes, however, the extension alone is fufficient, and the bone will flip into its place merely by pulling the limb with fufficient force. It is not hereby meant, that force alone is fufficient for the reduction of diflocations. Skill and address will often fucceed better than force. I have known

DISLOCATION OF THE THIGH. 543

known a diflocation of the thigh reduced by one man, after all the force that could be used by fix had proved ineffectual.

CHAP. LII.

OF BROKEN BONES, &c.

THERE is, in most country villages, fome perfon who pretends to the art of reducing fractures. Though in general fuch perfons are very ignorant, yet fome of them are very fuccefsful; which evidently proves, that a fmall degree of learning, with a fufficient share of common fense and a mechanical head, will enable a man to be useful in this way. We would, however, advise people never to employ such operators, when an expert and skilful furgeon can be had: but when that is impracticable, they must be employed: we shall therefore recommend the following hints to their confideration:

When a large bone is broken, the patient's diet ought in all refpects to be the fame as in an inflammatory fever. He fhould likewife be kept quiet and cool, and his body open by emollient clyfters; or, if thefe cannot be conveniently administered, by food that is of an opening quality; as stewed prunes, apples boiled in milk, boiled spinage, and the like. It ought, however, to be here remarked, that perfons who have been accustomed to live high, are not all of a fudden to be reduced to a very low diet. This might have fatal effects. There is often a neceffity for indulging even bad habits, in fome meafure, where the nature of the difease might require a different treatment.

It will generally be neceffary to bleed the patient immediately after a fracture, especially if he be young, of a full habit, or has at the same time received any bruife or contusion. This operation should not only be performed soon after the accident happens, but if the patient be very feverish, it may be repeated next day. When several of the ribs are broken, bleeding is peculiarly neceffary.

If

If any of the large bones which fupport the body are broken, the patient must keep his bed for feveral weeks. It is by no means neceffary, however, that he fhould lie all that time, as is customary, upon his back. This fituation finks the spirits, galls and frets the patient's skin, and renders him very uneasly. After the second week he may be gently raised up, and may fit several hours, supported by a bed-chair, or the like, which will greatly relieve him. Great care, however, must be taken in raising him up and laying him down, that he make no exertions himself, otherwise the action of the muscles may pull the bone out of its place *.

It is of great importance to keep the patient dry and clean while in this fituation. By neglecting this, he is often fo galled and excoriated, that he is forced to keep fhifting places for eafe. I have known a fractured thighbone, after it had been kept flraight for above a fortnight, difplaced by this means, and continue bent for life, in fpite of all that could be done.

It has been cuftomary when a bone was broken, to keep the limb for five or fix weeks continually upon the ftretch. But this is a bad pofture. It is both uneafy to the patient, and unfavourable to the cure. The beft fituation is to keep the joint a little bent. This is the pofture into which every animal puts its limbs when it goes to reft, and in which feweft mufcles are upon the ftretch. It is eafily effected, by either laying the patient upon his fide, or making the bed fo as to favour this pofition of the limb.

Bone-fetters ought carefully to examine whether the bone be not fhattered or broken into feveral pieces. In this cafe it will fometimes be neceffary to have the limb

* Various pieces of machinery have been contrived for counteracting the force of the mufcles, and retaining the fragments of broken bones; but as defcriptions of thefe without drawings would be of little ufe, I shall refer the reader to a cheap and ufeful performance On the Nature and Cure of Fradures, lately published by my ingenious triend Mr. Aitkin, surgeon, in Edinburgh; wherein that gentleman has not only given an account of the machines recommended in fractures by former authors, but has likewife added feveral improvements of his own, which are 'peculiarly useful in compound fractures, and in cases where patients with broken bones are obliged to be transported from one place to another.

OF BROKEN BONES, &c. 545

OF BROKEN BONES.

no freeuently immediately taken off, otherwife a gangrene or mortification may enfue. The horror which attends the very idea of an amputation, often occafions its being delayed in fuch cafes till too late. I have known this principle operate fo ftrongly, that a limb, where the bones were fhattered into more than twenty pieces, was not amputated before the third day after the accident, when the gangrene had proceeded fo far as to render the operation useles.

When a fracture is accompanied with a wound, it must be dreffed in all respects as a common wound.

All that art can do towards the cure of a broken bone, is to lay it perfectly straight, and to keep it quite eafy. All tight bandages do hurt. They had much better be wanting altogether. A great many of the bad confequences which fucceed to fractured bones, are owing to tight bandages. This is one of the ways in which the excels of art, or rather the abuse of it, does more mischief than would be occafioned by the want of it. Some of the most fudden cures of broken bones which were ever known, happened where no bandages were applied at all. Some method, however, must be taken to keep the member fleady; but this may be done many ways, without bracing it with a tight bandage.

The beft method of retention is by two or more fplints made of leather or pasteboard. These, if moistened before they are applied, foon affume the fhape of the included member, and are fufficient, by the affiftance of a very flight bandage, for all the purposes of retention. The bandage which we would recommend is that made with twelve or eighteen tails. It is much eafier applied and taken off than rollers, and anfwers all the purpofes of retention equally well. The fplints fhould always be as long as the limb, with holes cut for the ancles when the fracture is in the leg.

In fractures of the ribs, where a bandage cannot be properly used, an adhefive plaster may be applied over the part. The patient in this cafe ought to keep himfelf quite eafy, avoiding every thing that may occafion fneezing, laughing, coughing, or the like. He ought to keep his body in a straight posture, and should take care that his

Nn

OF BROKEN BONES, &c.

546

his ftomach be conftantly diftended, by taking frequently fome light food, and drinking freely of weak watery liquors.

The most proper external application for a fracture is oxycrate, or a mixture of vinegar and water. The bandages should be wet with this at every dreffing.

OF STRAINS.

Strains are often attended with worfe confequences than broken bones. The reafon is obvious; they are generally neglected. When a bone is broken, the patient is obliged to keep the member eafy, becaufe he cannot make ufe of it; but when a joint is only ftrained, the perfon, finding he can ftill make a fhift to move it, is forry to lofe his time for fo trifling an ailment. In this way he deceives himfelf, and converts into an incurable malady what might have been removed by only keeping the part eafy for a few days.

Country people generally immerfe a ftrained limb in cold water. This is very proper, provided it be done immediately, and not kept in too long. But the cuftom of keeping the part immerfed in cold water for a long time is certainly dangerous. It relaxes inftead of bracing the part, is more likely to produce a difeafe than remove one.

Wrapping a garter, or fome other bandage, pretty tight about the ftrained part, is likewife of ufe. It helps to reftore the proper tone of the veffels, and prevents the action of the parts from increafing the difeafe. It fhould not, however, be applied too tight. I have frequently known bleeding near the affected part have a very good effect: but what we would recommend above all, is *eafe*. It is more to be depended on than any medicine, and feldom fails to remove the complaint *.

* A great many external applications are recommended for ftrains, fome of which do good, and others hurt. The following are fuch as may be used with the greatest fafety, viz. poultices made of stale beer or vinegar and oatmeal, camphorated spirits of wine, Mindererus's spirit, volatile liniment, volatile aromatic spirit diluted with a double quantity of water, and the common fomentation, with the addition of brandy or spirit of wine.

OF

न मागणाम मा

OF RUPTURES.

Children and old people are most liable to this difease. In the former it is generally occasioned by excessive crying, coughing, vomiting, or the like. In the latter, it is commonly the effect of blows or violent exertions of the ftrength, as leaping, carrying great weights, &c. In both a relaxed habit, indolence, and an oily or very moist diet, dispose the body to this difease.

A rupture fometimes proves fatal before it is difcovered. Whenever fickness, vomiting, and obstinate coffivenels give reafon to fulpect an obstruction of the bowels, all those places where ruptures usually happen ought carefully to be examined. The protution of a very fmall part of the gut will occafion all thefe fymptoms; and, if not returned in due time, will prove fatal. On the first appearance of a rupture in an infant, it ought to be laid upon its back, with its head very low. While in this posture, if the gut does not return of itself, it may eafily be put up by gentle preffure. After it is returned, a piece of flicking-plaster may be applied over the part, and a proper trufs or bandage must be constantly worn for a confiderable time. The method of making and applying these rupture-bandages for children is pretty well known. The child must, as far as possible, be kept from crying, and from all violent exertions, till the rupture is quite healed.

In adults, when the gut has been forced down with great violence, or happens from any caufe to be inflamed, there is often great difficulty in returning it, and fometimes the thing is quite impracticable without an operation; a defcription of which is foreign to our purpofe. As I have been fortunate enough, however, always to fucceed in my attempts to return the gut, without having recourfe to any other means than what are in the power of every man, I shall briefly mention the method which I generally purfue.

After the patient has been bled, he must be laid upon his back, with his head very low, and his breech raifed high with pillows. In this fituation flannel-cloths wrung

Nn 2

ont

OF RUPTURES.

out of a decoction of mallows and camomile-flowers, or, if these are not at hand, of warm water, must be applied for a confiderable time. A clyfter made of this decoction, with a large fpoonful of butter and an ounce or two of falt, may be afterwards thrown up. If these should not prove fuccelsful, recourse must be had to preffure. If the tumour be very hard, confiderable force will be neceffary ; but it is not force alone which fucceeds here. The operator, at the fame time that he makes a preffure with the palms of his hands, must with his fingers artfully conduct the gut in by the fame aperture through which it came out. The manner of doing this can be much eafier conceived than defcribed. Should thefe endeavours prove ineffectual, clyfters of the fmoke of tobacco may be tried. Thefe have been often known to fucceed where every other method failed.

There is reafon to believe that, by perfifting in the ufe of thefe, and fuch other means as the circumftances of the cafe may fuggeft, moft *hernias* might be reduced without an operation. Cutting for the *hernia* is a nice and difficult matter. I would therefore advife furgeons to try every method of returning the gut before they have recourfe to the knife. I have once and again fucceeded by perfevering in my endeavours, after eminent furgeons had declared the reduction of the gut impracticable without an operation *.

An adult, after the gut has been returned, muft wear a fteel bandage. It is needlefs to defcribe this, as it may always be had ready-made from the artifts. Such bandages are generally uneafy to the wearer for fome time, but by cuftom they become quite eafy. No perfon who has had a rupture after he arrived at man's eftate, fhould ever be without one of thefe bandages.

Perfons who have a rupture ought carefully to avoid all violent exercife, carrying great weights, leaping, run-

• I would here beg leave to recommend it to every practitioner, when his patient complains of pain in the belly with obftinate coffivenefs, to examine the groins and every place where a rupture may happen, in order that it may be immediately reduced. By neglecting this, many perifh who were not fufpected to have had ruptures till after they were dead. I have known this happen where half a dozen of the faculty were in attendance. ning,

OF RUPTURES.

(hora)

549

ning, and the like. They fhould likewife avoid windy aliment and strong liquors; and should carefully guard against catching cold.

CHAP. LIII.

OF CASUALTIES.

T is certain that life, when to all appearance loft, may often, by due care, be reftored. Accidents frequently prove fatal, merely becaufe proper means are not ufed to counteract their effects. No perfon ought to be looked upon as killed by any accident, unless where the flructure of the heart, brain, or some organ neceffary to life, is evidently deftroyed. The action of thefe organs may be fo far impaired, as even to be for fome time imperceptible, when life is by no means gone. In this cafe, however, if the fluids be fuffered to grow cold, it will be impoffible to put them again in motion, even though the folids fhould recover their power of acting. Thus, when the motion of the lungs has been ftopped. by unwholefome vapour, the action of the heart by a ftroke on the breaft, or the functions of the brain by a blow on the head, if the perfon be fuffered to grow cold, he will in all probability continue fo; but, if the body be kept warm, as foon as the injured part has recovered its power of acting, the fluids will again begin to move, and all the vital functions will be reftored.

It is a horrid cuftom, immediately to confign over to death every perfon who has the misfortune, by a fall, a blow, or the like, to be deprived of the appearance of life. The unhappy perfon, inftead of being carried into a warm house, and laid by the fire, or put to a warm bed, is generally hurried away to church, or a barn, or fome other cold damp house, where, after a fruitless attempt has been made to bleed him, perhaps by one who knew nothing of the matter, he is given over for dead, and

OF CASUALTIES.

and no farther notice taken of him. This conduct feems to be the refult of ignorance, fupported by an ancient fuperflitious notion, which forbids the body of any perfon killed by accident to be laid in an houfe that is inhabited. What the ground of this fuperflition may be, we fhall not pretend to inquire; but furely the conduct founded upon it is contrary to all the principles of reafon, humanity, and common fenfe.

When a perfon feems to be fuddenly deprived of life, our firft bufinefs is to inquire into the caufe. We ought carefully to obferve whether any fubftance be lodged in the windpipe or gullet; and, if that is the cafe, attempts muft be made to remove it. When unwholefome air is the caufe, the patient ought immediately to be removed out of it. If the circulation be fuddenly ftopped, from any caufe whatever, except mere weaknefs, the patient fhould be bled. If the blood does not flow, he may be immerfed in warm water, or rubbed with warm cloths, &zc to promote the circulation. When the caufe cannot be fuddenly removed, our great aim muft be to keep up the vital warmth, by rubbing the patient with hot cloths, or falt, and covering his body with warm fand, afhes, or the like.

I fhould now proceed to treat more fully of those accidents, which, without immediate affistance, would often prove fatal, and to point out the most likely means for relieving the unhappy fufferers; but as I have been happily anticipated in this part of my fubject by the learned and humane Dr, Tiffot, I shall content myfelf with collecting fuch of his observations as feem to be the most important, and adding fuch of my own as have occurred in the course of practice.

OF SUBSTANCES STOPT BETWEEN THE MOUTH AND STOMACH.

Though accidents of this kind are very common, and extremely dangerous, yet they are generally the effect of careleffnefs. Children fhould be taught to chew their food well, and to put nothing into their mouths which it would

SUBSTANCES STOPT BETWEEN, &c. 551

would be dangerous for them to fwallow. But children are not the only perfons guilty of this piece of imprudence. I know many adults who put pins, nails, and other fharp-pointed fubftances in their mouths upon every occafion, and fome who even fleep with the former there all night. This conduct is exceedingly injudicious, as a fit of coughing, or twenty other accidents, may force over the fubftance before the perfon is aware *.

When any fubftance is detained in the gullet, there are two ways of removing it, viz. either by extracting it, or pufhing it down. The fafeft and most certain way is to extract it; but this is not always the easieft : it may therefore be more eligible fometimes to thrust it down, especially when the obstructing body is of fuch a nature, that there is no danger from its reception into the stomach. The substances which may be pushed down without danger are, all common nouriss ones, as bread, stork, wood, bones, pieces of metal, and such like, ought, if possible, to be extracted, especially if these bodies be sharp-pointed, as pins, needles, fish-bones, bits of glass, &c.

When fuch fubftances have not paffed in too deep, we fhould endeavour to extract them with our fingers; which method often fucceeds. When they are lower, we must make use of nippers, or a small pair of forceps, fuch as furgeons use. But this attempt to extract rarely fucceeds, if the fubstance be of a flexible nature, and has defcended far into the gullet.

If the fingers and nippers fail, or cannot be duly applied, crotchets, a kind of hooks, muft be employed. Thefe may be made at once, by bending a piece of pretty ftrong iron wire at one end. It muft be introduced in the flat way; and, for the better conducting it, there fhould likewife be a curve or bending at the end it is held by, to ferve as a kind of handle to it; which has

* A woman in one of the hofpitals of this city lately difcharged a great number of pins, whi h fhe had fwallowed in the course of her bufiness, through an ulcer in her fide.

Nn4

this

552 SUBSTANCES STOPT BETWEEN

this farther ufe, that it may be fecured by a firing tied to it; a circumftance not to be omitted in any inftrument employed on fuch occafions, to avoid fuch ill accidents as have fometimes enfued from thefe inftruments flipping out of the operator's hand. After the crotchet has paffed below the fubftance that obftructs the paffage, it is drawn up again, and hooks up the body along with it. The crotchet is alfo very convenient, when a fubftance fomewhat flexible, as a pin or fifh-bone, flicks acrofs the gullet, the hook, in fuch cates, feizing them about their middle part, crooks and thus difengages them; or, if they are very brittle fubftances, ferves to break them.

When the obstructing bodies are small, and only stop up a part of the paffage, and which may either eafily elude the hook, or straiten it by their refistance, a kind of rings, made either of wire, wool, or filk, may be ufed. A piece of fine wire of a proper length may be bent into a circle, about the middle, of about an inch diameter, and the long unbent fides brought parallel, and near each other : these are to be held in the hand, and the circular part or ring introduced into the gullet, in order to be conducted about the obstructing body, and fo to extract it. More flexible rings may be made of wool, thread, filk, or fmall pack-thread, which may be waxed for their greater ftrength and confiftence. One of these is to be tied fast to a handle of iron wire, whalebone, or any kind of flexible wood, and by this means introduced, in order to furround the obstructing fubftance, and to draw it out. Several of these rings paffed through one another may be used, the more certainly to lay hold of the obstructing body which may be involved by one, if another fhould mils it. These rings have one advantage, which is, that when the fubstance to be extracted is once laid hold of, it may then, by turning the handle, be retained fo ftrongly in the ring thus twifted, as to be moved every way, which must in many cafes be a confiderable advantage.

Another material employed on these unhappy occafions is the sponge. Its property of swelling confiderably on being wet is the principal foundation of its usefulness

THE MOUTH AND STOMACH.

fulnels here. If any fubstance is stopped in the gullet, but without filling up the whole paffage, a bit of fponge may be introduced into that part which is unftopped, and beyond the fubstance. The fponge foon dilates, and grows larger in this moift fituation ; and indeed the enlargement of it may be forwarded by making the patient fwallow a few drops of water. Afterwards it is to be drawn back by the handle to which it is fastened; and as it is now too large to return through the fmall cavity by which it was conveyed in, it draws out the obstructing body along with it.

The compreffibility of sponge is another foundation of its usefulness in such cases. A pretty large piece of fponge may be compressed or fqueezed into a small fize, by winding a ftring of tape clofely about it, which may be eafily unwound and withdrawn, after the fponge has been introduced. A bit of fponge may likewife be compreffed by a piece of whalebone split at one end; but this can hardly be introduced in fuch a manner as not to hurt the patient.

I have often known pins and other fharp bodies, which had fluck in the throat, brought up by caufing the perfon to fwallow a bit of tough meat tied to a thread. and drawing it quickly up again. This is fafer than fwallowing fponge, and will often anfwer the purpofe equally well.

When all these methods prove unfuccessful, there remains one more, which is, to make the patient vomit : but this can fcarcely be of any fervice, unlefs when fuch obstructing bodies are fimply engaged in, and not hooked or fluck into the fides of the gullet, as in this cafe vomiting might fometimes occasion farther mischief. If the patient can fwallow, vomiting may be excited by taking half a drachm or two fcruples of ipecacuanha in powder made into a draught. If he is not able to fwallow, an attempt may be made to excite vomiting, by tickling his throat with a feather; and, if that fhould not fucceed, a clyfter of tobacco may be administered. It is made by boiling an ounce of tobacco in a fufficient quantity of water. This has often been found to fucceed,

554 SUBSTANCES STOPT BETWEEN

fucceed, when other attempts to excite vomiting had failed.

When the obstructing body is of fuch a nature that it may with fafety be pushed downwards, this may be attempted by means of a wax-candle oiled, and a little heated, so as to make it flexible; or a piece of whalebone, wire, or flexible wood, with a sponge fastened to one end.

Should it be impoffible to extract even those bodies which it is dangerous to admit into the stomach, we must then prefer the least of two evils, and rather run the hazard of pushing them down, than suffer the patient to perish in a few minutes; and we ought to scruple this resolution the less, as a great many instances have happened, where the stallowing of such hurtful and indigestible substances have been followed by no diforder.

Whenever it is manifest that all endeavours either to extract or push down the substance must prove ineffectual, they should be discontinued; because the inflammation occasioned by persisting in them, might be as dangerous as the obstruction itself. Some have died in confequence of the inflammation, even after the body which caused the obstruction had been entirely removed.

While the means recommended above are making ufe of, the patient fhould often fwallow, or, if he cannot, he fhould frequently receive by injection, through a crooked tube or pipe that may reach down to the gullet, fome emollient liquor, as warm milk and water, barley-water, or a decoction of mallows. Injections of this kind not only foften and foothe the irritated parts, but, when thrown in with force, are often more fuccefsful in loofening the obftruction, than all attempts with inftruments.

When, after all our endeavours, we are obliged to leave the obftructing body in the part, the patient muft be treated as if he had an inflammatory difeafe. He fhould be bled, kept upon a low diet, and have his whole neck furrounded with emollient poultices. The like treatment muft also be ufed, if there be any reason to fuspect

THE MOUTH AND STOMACH. 555

fuspect an inflammation of the passages, though the obstructing body be removed.

A proper degree of agitation has fometimes loofened the inhering body more effectually than inftruments. Thus a blow on the back has often forced up a fubftance which fluck in the gullet; but this is ftill more proper and efficacious when the fubftance gets into the windpipe. In this cafe, vomiting and fneezing are likewife to be excited. Pins, which fluck in the gullet, have been frequently difcharged by riding on horfeback, or in a carriage.

When any indigestible fubstance has been forced down into the stomach, the patient should use a very mild and smooth diet, consisting chiefly of fruits and farinaceous substances, as puddings, pottage, and soups. He should avoid all heating and irritating things, as wine, punch, pepper, and such like; and his drink should be milk and water, barley-water, or whey.

When the gullet is fo ftrongly and fully clofed, that the patient can receive no food by the mouth, he must be nourished by clysters of foup, jelly, and the like.

When the patient is in danger of being immediately fuffocated, and all hope of freeing the paffage is vanifhed, fo that death feems at hand, if refpiration be not reftored; the operation of *bronchatomy*, or opening of the wind-pipe, must be directly performed. As this operation is neither difficult to an expert furgeon, nor very painful to the patient, and is often the only method which can be taken to preferve life in thefe emergencies, we thought proper to mention it, though it fhould only be attempted by performs fkilled in furgery.

OF DROWNED PERSONS.

When a perfon has remained above a quarter of an hour under water, there can be no confiderable hopes of his recovery. But as feveral circumftances may happen to have continued life, in fuch an unfortunate fituation, beyond the ordinary term, we fhould never too foon refign the unhappy object to his fate, but try every method for his relief, as there are many well-attefted proofs

of

556 OF DROWNED PERSONS.

of the recovery of perfons to life and health who had been taken out of the water apparently dead, and who remained a confiderable time without exhibiting any figns of life.

The first thing to be done, after the body is taken out of the water, is to convey it as foon as possible to fome convenient place where the necessary operations for its recovery may be performed. In doing this, care must be taken not to bruife or injure the body by carrying it in any unnatural posture with the head downwards, or the like. If an adult body, it ought to be laid on a bed, or on ftraw, with the head a little raised, and carried on a cart or on men's shoulders, and kept in as natural and easy a position as possible. A small body may be carried in the arms.

In attempting to recover perfons apparently drowned, the principal intention to be purfued is, to reftore the natural warmth, upon which all the vital functions depend; and to excite these functions by the application of ftimulants, not only to the skin, but likewise to the lungs, intestines, \mathfrak{Sc} .

Though cold was by no means the caufe of the perfon's death, yet it will prove an effectual obftacle to his recovery. For this reafon, after ftripping him of his wet clothes, his body muft be ftrongly rubbed for a confider, able time with coarfe linen cloths, as warm as they can be made; and, as foon as a well-heated bed can be got ready, he may be laid in it, and the rubbing fhould be continued. Warm cloths ought likewife to be frequently applied to the ftomach and bowels, and hot bricks, or bottles of warm water to the foles of his feet, and to the palms of his hands.

Strong volatile fpirits fhould be frequently applied to the nofe; and the fpine of the back and pit of the ftomach may be rubbed with warm brandy or fpirit of wine. The temples ought alfo to be chafed with volatile fpirits; and ftimulating powders, as that of tobacco or marjoram, may be blown up the noftrils.

To renew the breathing, a ftrong perfon may blow his own breath into the patient's mouth with all the force he can, holding his noftrils at the fame time. When it can

OF DROWNED PERSONS.

can be perceived by the rifing of the cheft or belly that the lungs are filled with air, the perfon ought to defift from blowing, and fhould prefs the breaft and belly fo as to expel the air again; and this operation may be repeated for fome time, alternately inflating and depreffing the lungs fo as to imitate natural refpiration.

If the lungs cannot be inflated in this manner, it may be attempted by blowing through one of the noftrils, and at the fame time keeping the other clofe. Dr. Monro, for this purpofe, recommends a wooden pipe fitted at one end for filling the noftril, and at the other for being blown into by a perfon's mouth, or for receiving the pipe of a pair of bellows, to be employed for the fame purpofe, if neceffary.

When air cannot be forced into the cheft by the mouth or nofe, it may be neceffary to make an opening into the wind pipe for this purpofe. It is needlefs, however, to fpend time in defcribing this operation, as it fhould not be attempted unlefs by perfons fkilled in furgery.

To ftimulate the inteffines the fume of tobacco may be thrown up in the form of a clyfter. There are various pieces of apparatus contrived for this purpole, which may be used when at hand; but where these cannot be obtained, the business may be done by a common tobacco-pipe. The bowl of the pipe mult be filled with tobacco well kindled, and, after the fmall tube has been introduced into the fundament, the Imoke may be forced up by blowing through a piece of paper full of holes, wrapped round the mouth of the pipe, or by blowing through an empty pipe, the mouth of which is applied close to that of the other. This may also be done in the following manner : A common clyfter-pipe, with a bag mounted upon it, may be introduced into the fundament, and the mouth of the bag may be applied round the fmall end of a tobacco-pipe, in the bowl of which tobacco is to be kindled, and the fmoke blown up as directed above. Should it be found impracticable to throw up the fmoke of tobacco, clyfters of warm warter, with the addition of a little falt, and fome wine or fpirits, may be frequently administered. This may be done by a common clysterbag

bag and pipe; but, as it ought to be thrown well up, a pretty large fyringe will answer the purpose better.

While these things are doing, some of the attendants ought to be preparing a warm bath, into which the perfon fhould be put, if the above endeavours prove ineffectual. Where there are no conveniences for using the warm bath, the body may be covered with warm falt, fand, ashes, grains, or fuch like. Tiffot mentions an inftance of a girl who was reftored to life, after the had been taken out of the water, fwelled, bloated, and to all appearance dead, by laying her naked body upon hot afhes, covering her with others equally hot, putting a bonnet round her head, and a flocking round her neck ftuffed with the fame, and heaping coverings over all. After fhe had remained half an hour in this fituation, her pulfe returned, fhe recovered fpeech, and cried out, I freeze, I freeze ; a little cherry brandy was given her, and fhe remained buried, as it were, under the afhes for eight hours: afterwards fhe was taken out, without any other complaint, except that of laffitude or wearinels, which went off in a few days. The doctor mentions likewife an inftance of a man who was reftored to life, after he had remained fix hours under water, by the heat of a dunghill.

Till the patient fhews fome figns of life, and is able to fwallow, it would be ufelefs and even dangerous to pour liquors into his mouth. His lips, however, and tongue may be frequently wet with a feather, dipped in warm brandy or other ftrong fpirits; and, as foon as he has recovered the power of fwallowing, a little warm wine, or fome other cordial, ought every now and then to be adminiftered.

Some recommend a vomit after the patient is a little re-animated; but if he can be made to puke without the fickening draught, it will be more fafe: this may generally be done by tickling the throat and fauces with an oiled feather, or fome other foft fubftance, which will not injure the parts. Tiffot, in this cafe, recommends the oxymel of fquills, a table fpoonful of which, diluted with water, may be given every quarter of an hour, till the patient has taken five or fix dofes. Where that medicine is

OF DROWNED PERSONS.

is not at hand, a ftrong infusion of fage, camomile flowers, or *carduus benedictus*, fweetened with honey, or fome warm water, with the addition of a little falt, may, he fays, fupply its place. The Doctor does not intend that any of these things should be given in fuch quantity as to occasion vomiting. He thinks emetics in this fituation are not expedient.

We are by no means to difcontinue our affiftance as foon as the patients difcover fome tokens of life, fince they fometimes expire after thefe firft appearances of recovering. The warm and ftimulating applications are ftill to be continued, and fmall quantities of fome cordial liquor ought frequently to be administered. Laftly, though the perfon should be manifestly re-animated, there fometimes remain an oppression, a cough, and feverishnes, which effectually constitute a difease. In this case, it will be necessary to bleed the patient in the arm, and to cause him to drink plentifully of barley-water, elder-flower tea, or any other fost pectoral infusion.

Such perfons as have the misfortune to be deprived of the appearances of life, by a fall, a blow, fuffocation, or the like, must be treated nearly in the fame manner as those who have been for some time under water. I once attended a patient who was fo flunned by a fall from a horfe, that for above fix hours he fcarcely exhibited any figns of life; yet this man, by being bled, and proper methods taken to keep up the vital warmth, recovered, and in a few days was perfectly well. Dr. Alexander gives an inftance to the fame purpofe, in the Edinburgh Phyfical and Literary Effays, of a man who was to all appearance killed by a blow on the breaft, but recovered upon being immerfed for fome time in warm water. Thefe, and other inftances of a fimilar nature, which might be adduced, amount to a full proof of this fact, that many of those unhappy perfons who lose their lives by falls, blows, and other accidents, might be faved by the use of proper means duly persisted in.

OF NOXIOUS VAPOURS.

Air may be many ways rendered noxious, or even destructive to animals. This may either happen from its vivifying principle being destroyed, or from subtle exhalations with which it is impregnated. Thus air that has paffed through burning fuel is neither capable of fupporting fire nor the life of animals. Hence the danger of fleeping in clofe chambers with coal fires. Some indeed fuppofe the danger here proceeds from the fulphureous oil contained in the coal, which is fet at liberty and diffused all over the chamber; while others imagine it is owing to the air of the room being charged with phlogiston. Be this as it may, it is a fituation carefully to be avoided. Indeed, it is dangerous to fleep in a fmall apartment with a fire of any kind. I lately faw four perfons who had been fuffocated by fleeping in an apartment where a fmall fire of coal had been left burning.

The vapour which exhales from wine, cyder, beer, or other liquors, in the ftate of fermentation, contains fomething poifonous, which kills in the fame manner as the vapour of coal. Hence there is always danger in going into cellars where a large quantity of these liquors is in a ftate of fermentation, especially if they have been close shut up for some time. There have been many inflances of perfons struck dead on entering fuch places, and of others who have with difficulty escaped.

When fubterraneous caves, that have been very long fhut, are opened, or when deep wells are cleaned, which have not been emptied for feveral years, the vapours arifing from them produce the fame effects as those mentioned above. For this reason, no perfon ought to venture into a well, pit, cellar, or any place that is damp, and has been long thut up, till the air has been fufficiently purified, by burning gunpowder in it. It is eafy to know, as has been observed in a former part of this work, work, when the air of fuch places is unwholefome, by letting down a lighted candle, throwing in burning fuel, or the like. If these continue to burn, people may fafely venture in; but where they are fuddenly extinguished, no one ought to enter till the air has been first purified by fire.

The offenfive fmell of lamps and of candles, efpecially when their flames are extinguished, operate like other vapours, though with lefs violence, and lefs fuddenly. There have, however, been inftances of people killed by the fumes of lamps which had been extinguished in a close chamber; and perfons of weak, delicate breafts generally find themfelves quickly opprefied in apartments illuminated with many candles.

Such as are fenfible of their danger in these fituations, and retreat feafonably from it, are generally relieved as foon as they get into the open air, or, if they have any remaining uneafiness, a little water and vinegar, or lemonade, drank hot, affords them relief. But when they are fo far poisoned, as to have lost their feeling, and understanding, the following means must be used for their recovery:

The patient fhould be exposed to a very pure, fresh, and open air; and volatile falts, or other stimulating fubstances, held to his nose. He should next be bled in the arm, or, if that does not succeed, in the neck. His legs ought to be put into warm water, and well rubbed. As soon as he can swallow, some lemonade, or water and vinegar, with the addition of a little nitre, may be given him.

Nor are fharp clyfters by any means to be neglected; thefe may be made, by adding to the common clyfter, fyrup of buckthorn and tincture of fenna, of each two ounces; or, in their flead, half an ounce of Venice turpentine diffolved in the yolk of an egg. Should thefe things not be at hand, two or three large fpoonfuls of common falt may be put into the clyfter. The fame means, if neceffary, which were recommended in the former part of this chapter, may be used to reftore the circulation, warmth, &c.

Mr.

562

Mr. Toffach, furgeon at Alloa, relates the cafe of a man fuffocated by the fteam of burning coal, whom he recovered by blowing his breath into the patient's mouth, bleeding him in the arm, and caufing him to be well rubbed and toffed about. And Dr. Frewen, of Suffex, mentions the cafe of a young man who was ftupified by the fmoke of fea-coal, but was recovered by being plunged into cold water, and afterwards laid in a warm bed.

The practice of plunging perfons fuffocated by noxious vapours in cold water, would feem to be fupported by the common experiment of fuffocating dogs in the grotto del cani, and afterwards recovering them by throwing them into the neighbouring lake.

EFFECTS OF EXTREME COLD.

When cold is extremely fevere, and a perfon is expofed to it for a long time, it proves mortal, in confequence of its ftopping the circulation in the extremities, and forcing too great a proportion of blood towards the brain; fo that the patient dies of a kind of apoplexy, preceded by great fleepinefs. The traveller, in this fituation, who finds himfelf begin to grow drowfy, fhould redouble his efforts to extricate himfelf from the imminent danger he is expofed to. This fleep, which he might confider as fome alleviation of his fufferings, would, if indulged, prove his laft.

Such violent effects of cold are happily not very common in this country; it frequently happens, however, that the hands or feet of travellers are fo benumbed or frozen, as to be in danger of a mortification, if proper means are not ufed to prevent it. The chief danger in this fituation arifes from the fudden application of heat. It is very common, when the hands or feet are pinched with cold, to hold them to the fire; yet reafon and obfervation fhew that this is a most dangerous and imprudent practice.

Every peafant knows, if frozen meat, fruits, or roots of any kind, be brought near the fire, or put into warm water, they will be deftroyed by rottennefs, or a kind

of

EFFECTS OF EXTREME COLD.

of mortification; and that the only way to recover them, it to immerfe them for fome time in very cold water. The fame obfervation holds with regard to animals in this condition.

When the hands or feet are greatly benumbed with cold, they ought either to be immerfed in cold water, or rubbed with fnow, till they recover their natural warmth and fenfibility: after which, the perfon may be removed into an apartment a little warmer, and may drink fome cups of tea, or an infufion of elder-flowers fweetened with honey. Every perfon muft have obferved, when his hands were even but flightly affected with cold, that the beft way to warm them was by wafhing them in cold water, and continuing to rub them well for fome time.

When a perfon has been fo long exposed to the cold, that all appearances of life are gone, it will be neceffary to rub him all over with fnow or cold water; or, what will answer better, if it can be obtained, to immerse him in a bath of the very coldest water. There is the greatest encouragement to persist in the use of these means, as we are assure that perfons who had remained in the fnow, or had been exposed to the freezing air during five or fix fucceffive days, and who had discovered no marks of life for several hours, have nevertheles been revived.

I have always thought, that the whitloes, kibes, chilblains, and other inflammations of the extremities, which are fo common among the peafants in the cold feafon, were chiefly occafioned by their fudden transitions from cold to heat. After they have been exposed to an extreme degree of cold, they immediately apply their hands and feet to the fire, or, if they have occafion, plunge them into warm water, by which means, if a mortification does not happen, an inflammation feldom fails to enfue. Most of the ill confequences from this quarter might be eafily avoided, by only observing the precautions mentioned above.

EFFECTS OF EXTREME HEAT.

The effects of extreme heat, though not fo common in this country, are no lefs fatal, and much more fudden O 0 2 than

564 EFFECTS OF EXTREME HEAT.

than those of cold. In hot countries people frequently drop down dead in the ftreets, exhausted with heat and fatigue. In this case, if any warm cordial can be poured into the mouth, it ought to be done. If this cannot be effected, they may be thrown up in form of a clyster. Volatile spirits, and other things of a stimulating nature, may be applied to the skin, which should be well rubbed with coarse cloths, whipped with nettles, or other stimulating things. Some of the antient physicians are said to have restored to life perfons apparently dead, by beating them with rods.

Head-aches are often occafioned by expolure to intenfe heat; and in warm climates, where people are very liable to what they call *coups de foleil*, or ftrokes of the fun, it is a common cuftom to lay linen cloths, feveral times doubled, on the head, and to keep them moistened with very cold water for half an hour, or till the ftupor is diminished. This they term *drawing the fire out of the head*.

CHAP. LIV.

OF FAINTING FITS, AND OTHER CASES WHICH REQUIRE IMMEDIATE ASSISTANCE.

STRONG and healthy perfons, who abound with blood, are often feized with fudden fainting fits, after violent exercife, drinking freely of warm or ftrong liquors, expofure to great heat, intenfe application to ftudy, or the like.

In fuch cafes the patient fhould be made to fmell to fome vinegar. His temples, forehead, and wrifts, ought at the fame time to be bathed with vinegar mixed with an equal quantity of warm water; and two or three fpoonfuls of vinegar, with four or five times as much water, may, if he can fwallow, be poured into his mouth.

If the fainting proves obflinate, or degenerates into a *lyncope*, that is, an abolition of feeling and underftanding, the patient must be bled. After the bleeding, a clyster will be proper, and then he should be kept easy and quiet, only

only giving him every half-hour a cup or two of an infufion of any mild vegetable, with the addition of a little fugar and vinegar.

When fwoonings, which arife from this caufe, occur frequently in the fame perfon, he fhould, in order to efcape them, confine himfelf to a light diet, confifting chiefly of bread, fruits, and other vegetables. His drink ought to be water or fmall beer, and he fhould fleep but moderately, and take much exercife.

But fainting fits proceed much oftener from a defect than an excels of blood. Hence they are very ready to happen after great evacuations of any kind, obstinate watching, want of appetite, or fuch like. In these, an almost directly opposite course to that mentioned above must be pursued.

The patient fhould be laid in bed, with his head low, and being covered, fhould have his legs, thighs, arms, and his whole body rubbed ftrongly with hot flannels. Hungary water, volatile falts, or ftrong finelling herbs, as rue, mint, or rofemary, may be held to his nofe. His mouth may be wet with a little rum or brandy; and, if he can fwallow, fome hot wine, mixed with fugar and cinnamon, which is an excellent cordial, may be poured into his mouth. A comprets of flannel dipt in hot wine or brandy muft be applied to the pit of his ftomach, and warm bricks, or bottles filled with hot water, laid to his feet.

As foon as the patient is recovered a little, he fhould take fome ftrong foup or broth, or a little bread or bifcuit foaked in hot-fpiced wine. To prevent the return of the fits, he ought to take often, but in fmall quantities, fome light yet ftrengthening nourifhment, as panado made with foup inflead of water, new-laid eggs lightly poached, chocolate, light roaft meats, jellies, and fuch like.

Thole fainting fits, which are the effect of bleeding or of the violent operation of purges, belong to this clafs. Such as happen after artificial bleeding are feldom dangerous, generally terminating as foon as the patient is laid upon the bed; indeed, perfons fubject to this kind fhould always be bled lying, in order to prevent it. O o 3 Should

OF FAINTING FITS, &c.

Should the fainting, however, continue longer than ufual, volatile fpirits may be held to the nofe, and rubbed on the temples, &c.

When fainting is the effect of too ftrong or acrid purges or vomits, the patient muft be treated in all respects as if he had taken poison. He should be made to drink plentifully of milk, warm water, and oil, barley-water, or such like; emollient clysters will likewise be proper, and the patient's strength should afterwards be recruited, by giving him generous cordials, and anodyne medicines.

Faintings are often occafioned by indigeftion. This may either proceed from the quantity or quality of the food. When the former of thefe is the caufe, the cure will be beft performed by vomiting, which may be promoted by caufing the patient to drink a weak infufion of camomile-flowers, *carduus benedictus*, or the like. When the diforder proceeds from the nature of the food, the patient, as in the cafe of weaknefs, must be revived by ftrong fmells, &c.; after which he should be made to fwallow a large quantity of light warm fluid, which may ferve to drown, as it were, the offending matter, to fosten its acrimony, and either to effect a discharge of it by vomiting, or force it down into the intestines.

Even difagreeable fmells will fometimes occafion fwoonings, efpecially in people of weak nerves. When this happens, the patient fhould be carried into the open air, have flimulating things held to his nofe, and thofe fubflances which are difagreeable to him ought immediately to be removed. But we have already taken notice of fwoonings which arife from nervous diforders, and fhall therefore fay no more upon that head.

Fainting fits often happen in the progress of difeases. In the beginning of putrid difeases, they generally denote an oppression at the stomach, or a mass of corrupted humours, and they cease after evacuations either by vomit or stool. When they occur at the beginning of malignant fevers, they indicate great danger. In each of these cases, vinegar used both externally and internally is the best remedy during the paroxism, and plenty of lemonjuice and water after it. Swoonings which happen in difeases accompanied with great evacuations, must be treated like

like those which are owing to weakness, and the evacuations ought to be restrained. When they happen towards the end of a violent fit of an intermitting fever, or at that of each exacerbation of a continual fever, the patient must be supported by small draughts of wine and water.

Delicate and hysteric women are very liable to fwooning or fainting fits after delivery. These might be often prevented by generous cordials, and the admittion of fresh air. When they are occasioned by excessive flooding, it ought by all means to be reftrained. They are generally the effect of mere weakness or exhaustion. Dr. Engleman relates the cafe of a woman " in child-" bed, who, after being happily delivered, fuddenly " fainted, and lay upwards of a quarter of an hour ap-" parently dead. A phyfician was fent for; her own " maid, in the mean while, being out of patience at his " delay, attempted to affift her herfelf, and extending " herfelf upon her mistrefs, applied her mouth to her's, " blew in as much breath as fhe poffibly could, and in " a very fhort time the exhausted woman awaked as out " of a profound fleep; when proper things being given " her, the foon recovered.

"The maid being afked how fhe came to think of this expedient, faid, fhe had feen it practifed at Altenburgh, by midwives, upon children with the happieft effect."

We mention this cafe chiefly that other midwives may be induced to follow fo laudable an example. Many children are born without any figns of life, and others expire foon after the birth, who might, without all doubt, by proper care, be reftored to life.

From whatever caule fainting fits proceed, frefh air is always of the greatest importance to the patient. By not attending to this circumstance, people often kill their friends while they are endeavouring to fave them. Alarmed at the patient's fituation, they call in a crowd of people to his affistance, or perhaps to witness his exit, whose breathing exhausts the air, and increases the danger. There is not the least doubt but this practice, which is very common among the lower fort of people, often $Q \circ 4$ proves proves fatal, efpecially to the delicate, and fuch perfons as fall into fainting fits from mere exhaustion, or the violence of fome dilease. No more perfons ought ever to be admitted into the room where a patient lies in a fwoon, than are absolutely neceffary for his affistance, and the windows of the apartment should always be opened, at least as far as to admit a stream of fresh air.

Perfons fubject to frequent fwoonings or fainting fits, fhould neglect no means to remove the caufe of them, as their confequences are always injurious to the conftitution. Every fainting fit leaves the perfon in dejection and weaknefs; the fecretions are thereby fulpended, the humours difpofed to ftagnation, coagulations and obftructions are formed, and, if the motion of the blood be totally intercepted, or very confiderably checked, *polypufes* are fometimes formed in the heart or larger veffels. The only kind of fwoonings not to be dreaded, are thofe which fometimes mark the *crifis* in fevers; yet even thefe ought, as foon as poffible, to be removed.

I have before remarked, but I deem it of importance to repeat the obfervation, that it is only when the fainting fit evidently arifes from a fulnefs of the habit, and is accompanied with a total abolition of feeling and underftanding, that bleeding is advifeable. The ufe of the lancet might otherwife have the moft deadly effect. Many perfons, even of robuft conflictions, are very apt to faint upon having a vein opened and lofing a little blood. How dangerous then muft the operation be, when a patient has already fainted, and moft probably from extreme weaknefs and a defect of blood ! I have no doubt but many a murder has been rafhly committed in fuch cafes.

OF INTOXICATION.

The effects of intoxication are often fatal. No kind of poifon kills more certainly than an overdole of ardent fpirits. Sometimes, by deflroying the nervous energy, they put an end to life at once; but in general their effects are more flow, and in many respects fimilar to those of opium. Other kinds of intoxicating liquors may prove

prove fatal when taken to excefs, as well as ardent fpirits; but they may generally be discharged by vomiting, which ought always to be excited when the stomach is overcharged with liquor.

More of those unhappy perfons, who die intoxicated, lose their lives from an inability to conduct themselves, than from the destructive quality of the liquor. Unable to walk, they tumble down, and lie in fome awkward pofture, which obstructs the circulation or breathing, and often continue in this fituation till they die. No perfon, when drunk, should be left by himself, till his clothes have been loofened, and his body laid in such a posture as is most favourable for continuing the vital motions, difcharging the contents of the store, &c. The best posture for discharging the contents of the store have be laid on his fide, with his head a little raised, and particular care must be taken that his neck be no way bent, twisted, or have any thing too tight about it.

The exceffive degree of thirft occafioned by drinking flrong liquors, often induces people to quench it by taking what is burtful. I have known fatal confequences even from drinking freely of milk after a debauch of wine or four punch; thefe acid liquors, together with the heat of the flomach, having coagulated the milk in fuch a manner that it could never be digefted. The fafeft drink after a debauch is water with a toaft, tea, infufions of balm, fage, barley-water, and fuch like. If the perfon wants to vomit, he may drink a weak infufion of camomile-flowers, or luke-warm water and oil; but in this condition, vomiting may generally be excited by only tickling the throat with the finger or a feather.

Inftead of giving a detail of all the different fymptoms of intoxication which indicate danger, and propofing a general plan of treatment for perfons in this fituation, I fhall briefly relate the hiftory of a cafe which lately fell under my own observation, wherein most of those fymptoms usually reckoned dangerous concurred, and where the treatment was fuccesful.

A young man, about fifteen years of age, had, for a hire, drank ten glaffes of flrong brandy. He foon after

4

fell

OF INTOXICATION.

fell fast asleep, and continued in that fituation for feveral hours, till at length his uneasy manner of breathing, the coldness of the extremities, and other threatening symptoms, alarmed his friends, and made them fend for me. I found him still strengther, his countenance ghastly, and his skin covered with a cold clammy sweat. Almost the only signs of life remaining were, a deep laborious breathing, and a convulsive motion or agitation of his bowels.

I tried to roufe him, but in vain, by pinching, fhaking, applying volatile fpirits, and other ftimulating things to his nofe, &c. A few ounces of blood were likewife taken from his arm, and a mixture of vinegar and water was poured into his mouth; but as he could not fwallow, very little of this got into the flomach. None of thefe things having the least effect, and the danger feeming to increase, I ordered his legs to be put into warm water and a fharp clyfter to be immediately administered. This gave him a ftool, and was the first thing that relieved him. It was afterwards repeated with the fame happy effect, and feemed to be the chief caufe of his recovery. He then began to flow fome figns of life, took drink when it was offered him, and came gradually to his fenfes. He continued, however, for feveral days weak and feverish, and complained much of a foreness in his bowels, which gradually went off, by means of a flender diet, and cool mucilaginous liquors.

This young man would probably have been fuffered to die, without any affiftance being called, had not a neighbour, a few days before, who had been advifed to drink a bottle of fpirits to cure him of an ague, expired under very fimilar circumftances.

OF SUFFOCATION AND STRANGLING.

These may fometimes proceed from an infraction of the lungs, produced by viscid clammy humours, or a spafmodic affection of the nerves of that organ. Perfons who feed grossly, and abound in rich blood, are very liable to suffocating fits from the former of these causes. Such ought, as soon as they are attacked, to be bled, to receive an emollient clyster, and to take frequently a cup of

SUFFOCATION AND STRANGLING. 571

of diluting liquor with a little nitre in it. They fhould likewife receive the fteams of hot vinegar into their lungs by breathing.

Nervous and afthmatic perfons are most fubject to fpafmodic affections of the lungs. In this cafe the patient's legs should be immersed in warm water, and the steams of vinegar applied as above. Warm diluting liquors should likewise be drank; to a cup of which a tea-spoonful of the paregoric elixir may occasionally be added. Burnt paper, feathers, or leather, may be held to the patient's nose, and fresh air should be freely admitted to him.

Infants are often fuffocated by the carelefinefs or inattention of their nurfes*. An infant when in bed fhould always be laid fo, that it cannot tumble down with its head under the bed-clothes; and when in a cradle, its face ought never to be covered. A finall degree of attention to thefe two fimple rules would fave the lives of many infants, and prevent others from being rendered weak and fickly all their days by the injuries done to their lungs.

Inftead of laying down a plan for the recovery of infants who are fuffocated, or overlaid, as it is termed by their nurfes, I fhall give the hiftory of a cafe related by Monfieur Janin, of the Royal College of furgery at Paris, as it was attended with fuccefs, and contains almost every thing that can be done on fuch occasions.

A nurfe having had the misfortune to overlay a child, he was called in, and found the infant without any figns of life; no pullation in the arteries, no refpiration, the face livid, the eyes open, dull, and tarnished, the nose full of fnivel, the mouth gaping, in short it was almost cold. Whilst fome linen-clothes and a parcel of ashes were warming, he had the boy unswathed, and laid him in a warm bed, and on the right fide. He then was rubbed all

• Thefe accidents are not always the effects of careleffnefs. I have known an infant overlaid by its mother being feized in the night with an hyfteric fit. This ought to ferve as a caution againft employing hyfteric women as nurles; and fhould likewife teach fuch women never to lay an infant in the fame bed with themfelves, but in a fmall adjacent one.

572 SUFFOCATION AND STRANGLING.

over with fine linen, for fear of fretting his tender and delicate fkin. As foon as the afhes had received their due degree of heat, Mr. Janin buried him in them, except the face, placed him on the fide opposite to that on which he had been at first laid, and covered him with a blanket. He had a bottle of eau de luce in his pocket, which he prefented to his nofe from time to time; and between whiles fome puffs of tobacco were blown up his noftrils : to these fucceeded the blowing into his mouth, and fqueezing tight his nofe. Animal heat began thus to be excited gradually; the pulfations of the temporal artery were foon felt, the breathing became more frequent and free, and the eyes closed and opened alternately. At length the child fetched fome cries expressive of his want of the breaft, which being applied to his mouth, he catched at it with avidity, and fucked as if nothing had happened to him. Though the pulfations of the arteries were by this time very well re-eftablished, and it was hot weather, yet Mr. Janin thought it advifeable to leave his little patient three quarters of an hour longer under the afhes. He was afterwards taken out, cleaned and dreffed as ufual; to which a gentle fleep fucceeded, and he continued perfectly well.

Mr. Janin mentions likewife an example of a young man who had hanged himfelf through defpair, to whom he administered help as effectually as in the preceding cafe.

Mr. Glover, furgeon in Doctors Commons, London, relates the cafe of a perfon who was reftored to life after twenty-nine minutes hanging, and continued in good health for many years after.

The principal means used to reftore this man to life were, opening the temporal artery and the external jugular; rubbing the back, mouth, and neck, with a quantity of volatile spirits and oil; administering the tobacco clyster by means of lighted pipes, and strong frictions of the legs and arms. This course had been continued for about four hours, when an incision was made into the windpipe, and air blown strongly through a canula into the lungs. About twenty minutes after this, the blood at the artery began to run down the sce, and a flow pulse was

SUFFOCATION AND STRANGLING. 573

was just perceptible at the wrift. The frictions were continued for fome time longer; his pulfe became more frequent, and his mouth and nofe being irritated with fpirit of fal ammoniac, he opened his eyes. Warm cordials were then administered to him, and in two days he was fo well as to be able to walk eight miles.

These cases are fufficient to shew what may be done for the recovery of those unhappy perfons who strangle themselves in a fit of despair.

OF PERSONS WHO EXPIRE IN CONVUL-SION FITS.

Convultion fits often conflitute the laft fcene of acute or chronic diforders. When this is the cafe, there can remain but fmall hopes of the patient's recovery after expiring in a fit. But when a perfon who appears to be in perfect health, is fuddenly feized with a convultion fit, and feems to expire, fome attempts ought always to be made to reftore him to life. Infants are most liable to convultions, and are often carried off very fuddenly by one or more fits about the time of teething. There are many well authenticated accounts of infants having been reftored to life, after they had to all appearance expired in convultions; but we fhall only relate the following inftance mentioned by Dr. Johnfon in his pamphlet on the practicability of recovering perfons wifibly dead.

In the parifh of St. Clements in Colchefter, a child of fix months old, lying upon its mother's lap, having had the breaft, was feized with a ftrong convultion fit, which lafted fo long, and ended with fo total a privation of motion in the body, lungs, and pulfe, that it was deemed abfolutely dead. It was accordingly ftripped, laid out, the paffing-bell ordered to be tolled, and a coffin to be made; but a neighbouring gentlewoman who used to admire the child, hearing of its fudden death, haftened to the house, and upon examining the child, found it not cold, its joints limber, and fancied that a glafs the held to its mouth and nofe was a little damped with the breath; upon which the took the child in her lap, fat down before the firerubbed it, and kept it in gentle agitation. In a quarter

to

of an hour fhe felt the heart begin to beat faintly; fhe then put a little of the mother's milk into its mouth, continued to rub its palms and foles, found the child begin to move, and the milk was fwallowed; and in another quarter of an hour fhe had the fatisfaction of reftoring to its difconfolate mother the babe quite recovered, eager to lay hold of the breaft, and able to fuck again. The child throve, had no more fits, is grown up, and at prefent alive.

These means, which are certainly in the power of every perfon, were fufficient to reftore to life an infant to all appearance dead, and who, in all probability, but for the use of these simple endeavours, would have remained fo. There are, however, many other things which might be done in case the above should not succeed; as rubbing the body with strong spirits, covering it with warm assors or falt, blowing air into the lungs, throwing up warm stimulating clysters or the stroke of tobacco into the intestines, and such like.

When children are dead born, or expire foon after the birth, the fame means ought to be used for their recovery, as if they had expired in circumstances fimilar to those mentioned above.

These directions may likewise be extended to adults, attention being always paid to the age and other circumftances of the patient.

The foregoing cafes and obfervations afford fufficient proof of the fuccels which may attend the endeavours of perfons totally ignorant of medicine, in affifting those who are fuddenly deprived of life by any accident or difease. Many facts of a fimilar nature might be adduced, were it neceffary; but these, it is hoped, will be fufficient to call up the attention of the public, and to excite the humane and benevolent to exert their utmost endeavours for the prefervation of their fellow-men.

The fociety for the recovery of drowned perfons, inflituted at Amiterdam in the year 1767, had the fatisfaction to find that no fewer than 150 perfons, in the fpace of four years, had been faved by the means pointed out by them, many of whom owed their prefervation to peafants and people of no medical knowledge. But the means

IN CONVULSION FITS.

means used with fo much efficacy in recovering drowned perfons are, with equal fuccefs, applicable to a number of cafes where the powers of life feem in reality to be only fuspended, and to remain capable of renewing all their functions, on being put into motion again. It is shocking to reflect, that, for want of this confideration, many perfons have been committed to the grave in whom the principles of life might have been revived.

The cafes wherein fuch endeavours are most likely to be attended with fuccess, are all those called fudden deaths from an invisible cause, as apoplexies, hysterics, faintings, and many other diforders wherein perfons in a moment fink down and expire. The various cafualties in which they may be tried are, fuffocations from the fulphureous damps of mines, coal-pits, &c. the unwholefome air of long unopened wells or caverns; the noxious vapours arising from fermenting liquors; the stars of burning charcoal; fulphureous mineral acids; arfenical effluvia, &c.

The various accidents of drowning, ftrangling, and apparent deaths, by blows, falls, hunger, cold, &c., likewife furnish opportunities of trying fuch endeavours. Those, perhaps, who, to appearance, are killed by lightning, or by any violent agitation of the passions, as fear, joy, furprise, and fuch like, might also be frequently recovered by the use of proper means, as blowing strongly into their lungs, &c.

The means to be used for the recovery of perfons fuddenly deprived of life, are nearly the fame in all cafes; they are practicable by every one who happens to be prefent at the accident, and require no great expence, and lefs fkill. The great aim is to reftore the warmth and vital motions. This may in general be attempted by means of heat, frictions, bleeding, blowing air into the lungs, administering clysters and generous cordials. Thefe mult be varied according to circumflances. Common fenfe, and the fituation of the patient, will fuggest the proper manner of conducting them. Above all, we would recommend *perfeverance*. People ought never to defpair on account of difcouraging circumflances, or to leave off 7

576 PERSONS WHO EXPIRE, &c.

their endeavours as long as there is the leaft hope of fuccefs. Where much good and no hurt can be done, no one ought to grudge his labour.

It were greatly to be wifhed, that an inftitution, fimilar to that of Amfterdam, was eftablifhed, upon a more extensive plan, in Great Britain; and that a reward was allowed to every one who fhould be inftrumental in reftoring to life a perfon feemingly dead *. Men will do much for fame, but ftill more for money. Should no profit, however, be annexed to those benevolent offices, the heartfelt pleasure which a good man must enjoy, on reflecting that he has been the happy inftrument of faving one of his fellow-creatures from an untimely grave, is itself a fufficient reward.

CHAP. LV.

OF COLD BATHING, AND DRINKING THE MINERAL WATERS.

IN a note fubjoined to the "Cautions," given in former editions of this work, concerning the use of the cold bath and of mineral waters, I pledged myself to treat the subject at more length on a future occasion; and I now mean to fulfil that promise. The discussion will be naturally divided into two parts. In the first, I shall endeavour to illustrate the extensive utility of the

• The Author is happy to obferve, that, fince the first publication of this work, feveral focieties have been instituted in Britain with the fame benevolent intention as that of Amsterdam, and that their endeavours have proved no lefs fuccefsful. He is likewife happy to obferve, that premiums have been awarded to those who have been active in their endeavours to reftore to life perfons who had been drowned, or fuddenly deprived of life by any accident. How much is this fuperior to the fuperstitious institution, which allows any man a premium who brings a dead perfon out of the water, fo that he may receive Christian burial; but allows nothing to the perfon who brings him out alive, or who recovers him after he has been to all appearance dead ?

cold

cold bath, in preferving as well as reftoring health; and fhall point out the cafes where it may be hurtful. I propofe, in the fecond part, to give a particular account of the qualities of the moft celebrated mineral waters in our own ifland, and on the continent;—to fpecify the difeafes wherein they are refpectively indicated ;—and to lay down the belt practical rules for employing them with fafety and benefit. I need not enlarge on the importance of fuch details. Every body knows that the beft things are liable to the greateft abufe; and as nature does not afford more active or more powerful medicines than those in queftion, any error in the application may be, and often is, attended with confiderable danger.

OF COLD BATHING, WITH REMARKS ON THE CASES IN WHICH THE WARM BATH IS MORE ADVISABLE.

Immersion in cold water is a custom which lays claim to the most remote antiquity. Indeed, it must have been coëval with man himfelf. The neceffity of water for the purpose of cleanlines, and the pleasure arising from its application to the body in hot countries, muft very early have recommended it to the human fpecies. Even the example of other animals was fufficient to give the hint to man. By inftinct, many of them are led to apply cold water in this manner; and fome, when deprived of its use, have been known to languish, and even to die. But whether the practice of cold-bathing arole from neceflity, reafoning, or imitation, is an inquiry of little confequence : our business is to point out the advantages which may be derived from it, when judicioufly reforted to, and the danger attending its improper ule.

People are apt to imagine that the fimple element of water can do no hurt, and that they may plunge into it at any time with impunity. In this, however, they are much miftaken. I have known apoplexies occafioned by going into the cold bath,—fevers excited by ftaying Pp too

too long in it,—and other maladies fo much aggravated by its continued ufe as to become abfolutely incurable. Without a proper diferimination with regard to the difeafe and the conflitution of the patient, the moft powerful medicine is more likely to do harm than good. The phyfician, who cured Augustus by cold bathing, killed his heir by the very fame prefeription. This induced the Roman fenate to make laws for regulating the baths, and preventing the numerous evils which arofe from an imprudent and promiscuous use of those elegant and fafhionable pieces of luxury. But as no fuch laws exist in this country, every one does that which is right in his own eyes, and of courfe many must do wrong. I hope, however, that, when better informed, they will learn to correct errors of fo fatal a tendency.

Abfurd prejudices against cold-bathing are not lefs blameable on the other hand. Though it fhould never be prefcribed for the cure of difeafes, without well confidering the nature of each cafe, it cannot be too earnestly or too generally recommended as a prefervative of health. I am therefore forry to fee fome modern writers attempting to revive the whimfical and long-exploded doctrine of GALEN, who faid, that immerfion in cold water was fit only for the young of lions and bears; and that warm-bathing was conducive to the growth and strength of infants. How egregiously do the greatest men err, whenever they lofe fight of facts, and fubftitute fallies of wit or fpecious arguments in physic for obfervation and experience! By these the superior excellence of the cold bath is placed beyond the poffibility of a doubt. Its tonic powers are found to be peculiarly proper for the lax fibres of young people, rendering them firm and elastic, and enabling the vital organs to perform their respective functions with eafe and regularity.

In other parts of this work I had occafion to defcribe, with greater minutenefs than is now neceffary, the many good effects of washing children; and I gave a few directions as to the manner of employing this very falutary operation, from the moment of their birth. I shewed

fhewed how the use of the cold bath might be gradually brought about with the utmost fafety; and I am perfuaded that those who give it a fair trial will readily comply with my farther advice to continue it ever after, except in fuch cafes of indifpolition or infirmity as I shall prefently notice. Nothing contributes more to the growth, vigour, and firmnels of youth, or to the activity and permanent health of manhood, than daily immerfion in cold water. It fteels the frame against changes of weather, against the impressions of cold or moisture, and many other external injuries. It is of course the best preventive of all those difeases which arise from a relaxed Ikin, obstructed or profuse perspiration, and nervous weaknefs.

When the cold bath is used merely as a means of preferving health, in which point of view I am now confidering it, a fingle plunge or dip of the whole body will be fufficient, though active fwimmers may continue their favourite amusement for five or fix minutes without injury. Any longer flay might prove dangerous, by not only occasioning an excessive flux of humours towards the head, but chilling the blood, cramping the muscles, relaxing the nerves, and wholly defeating the intention of bathing. For want of a due regard to thefe circumftances, young men have often endangered, and fometimes loft their lives. In all cafes, it is highly neceffary to be rubbed dry at the inftant of coming out of the water, and to take exercise for at least half an hour after. A little exercife is alfo advifable before bathing, fo as to excite a gentle glow or temperate degree of warmth, and thus guard against the bad confequences of a shock. when the body is either chilly or over-heated.

The like caution fhould be given against plunging into cold water after dinner, or after much fatigue. For thefe and many other reafons, the morning is very properly recommended to perfons in health as the best time for bathing. It is the least likely to interfere with their other pursuits or concerns : it washes away any particles of the peripirable matter that may have remained on the furface of the fkin, before they can be re-abforbed : it affords fresh supplies of vigour and alacrity, to enter

Pp2

upon

upon the duties of the day; and, as I have already hinted, it fortifies the body against any changes of weather, to which it may be afterwards exposed in a far lighter element.

In a ftate of perfect health, it may be farther obferved, that people need not give themfelves much trouble to enjoy the advantages of fea-bathing in preference to river-water, as the grand effect of both is nearly the fame, though fome confiderations of lefs moment may concur to render the former more inviting. Among thefe we must reckon the usual refort of gay company to different parts of the coaft in fummer, the refreshing coolnefs of the fea-air in that feafon, and the agreeable ftimulus which many perfons experience from the action of faline particles, not only in the water, but when they are floating in the atmosphere. It fhould also be confidered, that the temperature of the fea is more uniform than that of rivers, never rifing fo high, or finking fo low, in any change of weather. But fuch points of difference are chiefly interesting to valetudinarians.

What I have faid of the cold bath, when used as the means of preventing difease, will throw some light on the propriety of occasionally reforting to it as an important remedy. In cases of peculiar delicacy and danger, it is an inftrument which can only be entrusted to the most skilful hands; but in many other less critical fituations, a few plain rules may be of confiderable fervice.

The first object to be attended to in the use of the cold bath, as a remedy, is, whether the patient is not too much enfeebled to bear the shock. This cannot always be determined by appearances; but a single experiment will remove all doubt. If the immersion be followed by a pleasant glow, and a sense of increasing alacrity, it is the best proof of its agreeing with the constitution, and of its being likely to have a happy influence on the whole frame. Hence the cold bath is found to be an excellent bracer and restorative in cases of languor, of habitual lassing from much confinement, a sedentary life, intense fludy, or any of the usual causes of relaxation. But it is always understood, that, in every instance of this fort, a sufficient

noda

a fufficient strength of original stamina still remains to produce a proper re-action of the heart and arteries, upon which all the falutary effects of bathing depend.

The great efficacy of the cold bath, and particularly of fea-bathing, has often been experienced in fcrophulous complaints, which are always attended with a relaxation of the fibres, and a ftrong disposition to languor and indolence. In fuch cafes, fea-bathing is not only recommended as a tonic, or bracer, but as a powerful detergent and purifier allo, especially if the fea-water be ufed internally at the fame time. No difference of opinion prevails on this head, as far as relates to the fcrophula, but it has been alleged, that fea-bathing, though a good preventive of the scrophula, could not remove the local effects of the difease when once formed. My own practice in the treatment of fcrophulous affections has not been extensive enough to enable me to speak to this point with a tone of confidence; but the contrary doctrine appears to me supported by the fairest reasoning, and, what is more, by indifputable facts.

In the first place, a weak flaccid habit, and a thin kin, very fusceptible of impressions from cold most air, are the principal, if not the only predisposing causes of the scrophula. Now the cold bath is the best remedy for both, as it renders the texture of the skin firm, and invigorates the system. By being therefore so well adapted to remove causes, it must, according to one of the sures of medical practice, be very fit to remove effects.

The justness of fuch an inference has been placed beyond a doubt, by the reports of men of professional eminence and veracity, under whose direction, and immediate inspection also, fea-bathing has been known to resolve swellings of the glands, as well as to correct the discharge of scrophulous ulcers, and dispose them to heal. I am therefore very willing to believe, that a regular course of fea-bathing, and the internal use of scawater, with the aid of good air, proper exercise, and a light, yet nourishing diet, are the best means as yet discovered of checking the progress of the evil, or counteracting its morbid effects.

Pp3

But,

But, in order to prevent any poffible milconception of my meaning, it may be neceffary to add, that my opinion of the efficacy of fea-water in fcrophulous complaints, is confined to its probable removal of the outward fymptoms of the malady, before these have arrived at a certain pitch, or have reduced the patient to a state of extreme debility; in which case, as well as in all internal affections of the fcrophula, when it has once fastened upon the lungs, or any other vital part, bathing in the fea, or drinking its waters, would be not only useles, but extremely injurious.

It would also imply too great a confidence in the falutary virtues of fea-bathing, to prefcribe it as a remedy for cutaneous diforders in general. To many of them the warm bath is much better adapted; and the proper choice of the one or the other can only be determined by a skilful physician, after a due confideration of the patient's cafe. Some irruptions, if imprudently repelled by the action of cold on the fkin, may carry back into the habit the feeds of difeafe, to be deposited, perhaps, on some vital part, in spite of Nature's kind efforts to throw them off. But a medical man will not prefcribe fea-bathing in any cafe where pimples or blotches appear on the furface, without recommending the internal ufe of the fea-water at the fame time, to determine regularly and moderately to the bowels, fo as to carry off all impurities, without the least injury to the general health, fpirits, or appetite. I fhall have occasion to repeat this caution, when I come to fpeak of fome mineral waters, which are frequently reforted to for the cure of fimilar complaints.

Though, as I before obferved, there may be very little difference between the effects of fea-water and of river-water of the fame temperature, when applied to a found fkin and healthy body, yet the gently ftimulant, detergent, and healing properties of the faline impregnation of the former must give it a decifive fuperiority in many difeafes of the furface and habit. It cleanfes fores, and forwards the process of granulation. It often difperfes tumours that have refisted the most powerful difcutient medicines. Even deeply-feated ulcers, though beyond

beyond the reach of other applications, fometimes yield to the penetrating action of fea-water. We mult not forget, however, that its internal ufe is a neceffary auxiliary in all thefe cafes, and others of a fimilar nature. About half a pint of it, which contains fomewhat more than a quarter of an ounce of falts, taken in the morning, immediately on coming out of the fea, and the like dofe in half an hour after, will commonly anfwer the purpofe of a mild purgative. The quantity may be augmented, or the dofe repeated, if requifite, with perfect fafety, and little inconvenience. It excites thirft, but feldom naufea, unlefs the ftomach is very irritable, or the patient very fqueamifh.

In chronic difeases, where a cure cannot be expected but from the long-continued use of any remedy, it is a great recommendation of the fea-water, that it may be perfevered in for a confiderable time, without weakening the ftomach, the inteftines, or the conftitution in general. Inftances frequently occur of perfons who keep the body moderately open by its daily use for months together, and yet enjoy during the whole time a good appetite, and excellent powers of digeftion, with increafed vigour both of body and mind. It is always most advifable to make use of the fea-water externally and internally, in the manner here directed, only twice or three times a-week, till the patient is encouraged by degrees to employ the falutary process every day. It fhould also be gradually difcontinued in the fame manner, after the defired end is obtained.

There are feveral diforders, befides those already mentioned, particularly ardent fevers, and various cafes of local inflammation and muscular rigidity, in which the external application of cold water may produce good effects. But many of them require great accuracy of diffinction, as well as the utmost judgment and caution in the use of a remedy, which a small mistake, or a small change of circumstances, may render hazardous. In a work like this, designed for popular instruction, it would be improper to encourage rash experiments, by pointing out such niceties in medical practice as are fase only when under the guidance of medical skill. I do not know

PP4

any

584

any thing in its own nature fo falutary, and yet fo liable to be abufed, as the cold bath. I fhall therefore proceed to touch upon the cafes, where the inconfiderate or improper application of fuch a remedy may prove injurious, and fometimes fatal.

It is not merely in the critical cafes just alluded to, but in many flighter indispositions, that injudicious immerfion in cold water may be attended with very ferious confequences. Fevers are much oftener produced than cured by cold-bathing, if rashly reforted to. Diforders of the intellectual functions, palfies, apoplexies, and death, may be, and are frequently, occasioned by a fingle dip, in cafes either of extreme nervous debility, or of extreme fulness. When I reflect on the frantic precipitancy with which I have feen many perfons of very weak, and others of very plethoric habits, after a rapid journey from London to fome watering-place, plunge inftantly into the fea, without the least preparation, fo far from being furprized at the numbers who fuffer, I am rather aftonifhed that any fhould efcape. In order to prevent the ignorant and the thoughtless from falling victims to their indifcretion, and to guard perfons afflicted with particular complaints against the use of an improper medicine, I shall point out the principal indispositions, in which the cold bath would be likely to aggravate the fymptoms, and even to endanger the life of the patient.

When cold-bathing occafions chillnefs, lofs of appetite, liftleffnefs, pain of the breaft or bowels, a proftration of ftrength, or violent head-aches, it ought to be difcontinued. These unpleasant fensations are the furest proofs, that the actual state of the patient's habit is unfit to bear the state of the patient's habit is unfit to bear the state is too weak to overcome the cold preffure on the furface, or that the determination to the head, or 'to fome other vital part, is too rapidly increased. Every body's feelings, after immersion in cold water, are the best criterion by which we can decide on the probability of its good or its bad effects. We might otherwise be deceived by appearances, and be induced to recommend the cold bath in all cafes that might might feem to require a tonic and ftimulant plan of cure.

But it may fometimes be dangerous, or at leaft very detrimental, to make even a fingle experiment. In particular affections of the fromach and bowels, as well as in difeafes of the lungs or of the brain, and all obstinate obstructions, the effect may be fatal. The late Dr. SMOLLET, indeed, faid, that if he were perfuaded he had an ulcer in the lungs, he would jump into the cold bath. In doing fo, however, the Doctor would certainly fhew more courage than difcretion; and that he was more a man of wit than a phyfician, every one will allow. A nervous afthma, or an atrophy, may be mistaken for a pulmonary confumption : yet, in the two former, the cold bath proves often beneficial, though I never knew it fo in the latter. Indeed, all the phthifical patients I ever faw, who had tried the cold bath, were evidently hurt by it.

Perfons of very full habits, as I have already hinted, run a great rifk of burfting a blood-veffel, or of caufing an inflammation of fome important organ, by rufhing into the cold bath, without due preparation. People of this defcription ought by no means to bathe, unlefs the body has been previoufly prepared by fuitable evacuations. They will then derive the utmost benefit from what might be otherwife attended with irreparable injury to many of them.

Though I recommended the cold bath in cafes of nervous weaknefs, yet *the degree* of that weaknefs fhould be confidered, left the fhock might prove too powerful for extreme debility. Not only women of very weakly and delicate habits, but men alto in the fame predicament, as well as puny children, fhould begin with the warm bath, at the fame degree nearly as that of animal heat, about 96° of Fahrenheit's thermometer; and reduce it gradually in proportion to the increase of the patient's itrength and internal powers of re-action. The cold bath is often very neceffary to complete a cure, though not always advisable to begin with. This requires particular illustration.

In hysteric and hypochondriac cases, cold-bathing at first has done the greatest mischief, though it may be finally reforted to with good effect, after a preparatory and long-continued use of the tepid or lukewarm bath. Its warmth must be diminished very flowly, and almost imperceptibly. Nature revolts against all great transtions; and those who do violence to her dictates, have often cause to repent of their temerity.

The like gradual diminution of the temperature of the water is no lefs proper in rheumatic complaints, and in those muscular contractions and convulsive motions which are called *St. Vitus's Dance*.

Indeed, it may be laid down as a pretty general rule in that branch of nervous diforders which includes fpafms, convultions, epilepfies, and fimilar confequences of the debility or irritability of the fyftem, that we fhould always begin with the warm bath, and proceed to the cold by the most pleafing and gentle degrees.

The chief exceptions to this rule occur in the treatment of fpafmodic affections of the inteffines, hoopingcoughs, and convulfive afthmas, in which, though claffed under the general head of fpafms, the cold bath would at any time be extremely improper. But this prohibition is alfo implied in my remark on complaints of the bowels and cheft in general, the latter including coughs of every defcription. When thefe are the mere confequences of flight irritation or cold, bathing the lower extremities in warm water affords great relief; but immerfing the whole body in either the warm or the cold bath, would only aggravate the fymptoms, when the breathing is difficult.

As palfies are often occafioned by the inconfiderate ufe of the cold bath, it cannot be too ftrictly prohibited, where any paralytic fymptoms are difcoverable. There is no complaint that bears and requires a greater degree of external heat than the palfy, and there is none in which the fhock of cold water is more directly oppofite to every curative indication. The hot baths, therefore, whether natural or artificial, and particularly if impregnated with falt which increafes their ftimulus, are employed

ployed as a fovereign remedy for paralytic affections. Friction, which fbould never be neglected after bathing, is in these cases of eminent service.

In affections of the nervous coat of the ftomach, and in cafes of indigeftion, especially when occasioned by intemperance, cold-bathing is as improper as in complaints of the bowels, before taken notice of. But it is the excefs of folly, after immoderate drinking, to use the cold bath with a view of alleviating its painful effects next day. It must increase the diforder of the stomach, the violence of the head-ach, and the derangement of the circulation. It may be productive of still worfe confequences. The cooling operation may prove far more powerful and more lafting than was expected, and may extinguish for ever the remains of animal heat; or, should nature, by extraordinary efforts, be able to refift the flock, it will probably be attended with fymptoms of fever, or with very troublefome eruptions. Many painful affections of the head, as well as those which arife from intoxication, are, indeed often relieved by what is called the *shower-bath*, or by the effusion of cold water on the part affected, but never by the rafh experiment of fwimming, or of total immerfion.

I must take this opportunity to add, that the *fhower-bath* is in many other refpects a valuable contrivance. It may be easily procured: its action can be regulated at pleafure; and as the water defcends like rain, it gently impels the blood towards the lower extremities, and prevents the danger which would arise from its fudden or too rapid determination to the lungs and head in fome of the cafes already mentioned.

In uterine hæmorrhages, and other fluxes of blood, when fo confiderable as to endanger the patient's life or conftitution, cold water may be applied with good effect. It alfo forms a part of the tonic plan to be purfued in an immoderate flow of the *menses*; nor is any thing more likely to prevent the return of this complaint than cold bathing and drinking chalybeate waters in the intervals of menstruation. But when the discharge of blood is critical, as in fome affections of the brain, lungs, &c. or is become

become habitual, as in the piles, to check fo falutary an evacuation by the ufe of the cold bath would be the height of madnefs. This is no lefs true of many critical inflammations, those of the gout for inflance, in which cold water or any other repellent would evidently counteract the purpoles of nature, and very probably throw the diforder upon fome vital part. What is called the *retrocedent* gout frequently arifes from fome mismanagement of this fort, as well as from fome particular weaknefs or atony of the fystem. Cold bathing is a very hazardous experiment to be made by perfons fubject to the gout, except in the abfence of the fymptoms, when no indilposition is felt in either the head or ftomach, when the extremities are not threatened with pain ; and then only in concurrence with the beft medical advice.

Bathing the lower extremities in warm water is generally and very properly recommended both in the retention and fuppreflion of the *menfes*, to excite the action of the uterine veffels, and, in the latter cafe, to remove any ftricture of those veffels which may be induced by cold or fear. A skilful physician, however, will sometimes meet with cafes of a retention of the *menfes* after the usual age, in which the cold bath, if feasonably used at the beginning of the difease, may contribute to restore the tone of the system.

The delicacy and general irritability of the habit in a flate of pregnancy, as well as the danger of too great a determination of the blood to the womb, clearly forbid the ufe of the cold bath, unlefs it fhould be rendered advifable by fome circumflances of a peculiar nature, of which a medical man of fkill and experience is the only proper judge.

It is a great and often a fatal miftake to rely on the tonic powers of the cold bath as the beft means of repairing the injury done to the conflictution by the relaxing influence of hot climates. People, on their return to England after having refided in the Eaft or Weft Indies, would find the warm bath not only fafer, but far more conducive to the recovery of their former ftrength. I would not have them venture into a bath of a temperature

ture under 90° for a confiderable time, after which they may gradually diminish its warmth, as before recommended in cases of *extreme debility*.

I might here go into farther details, and fhew how much more falutary the warm bath is than the cold in difeafes of the liver and kidneys, and in numberlefs other cafes of internal weaknefs, irritation, or derangement; but the principles, which I have laid down, may be eafily extended to them all; and I hope that the cautions I have given will operate as fome check on the abufe of the moft powerful means of preferving and reftoring health, with which we are acquainted.

Some years ago a foreign quack made a great deal of noife in this country with his medicated baths; but, like other follies of the day, they are now almost funk into oblivion. A few writers have also been very lavish of their panegyrics on the wonderful effects of vapour baths as used in Ruffia: but I do not think that the inhabitants of these milder regions will ever have occasion to envy the rigid fibres of the north the enjoyment of fuch fanciful luxuries. The ftrength of steam is, perhaps, better known and more ufefully employed in England than in any quarter of the globe; but we meet with very few cales, where its intenfe action on the furface of the human body can be deemed effentially neceffary either for the prevention or the cure of difeafes. Surely the fkin of an Englishman may be rendered perspirable by a much gentler ftimulus, and without the aid of fo troublefome. and fuffocating a procefs.

OF MINERAL WATERS.

It is fcarcely poffible to read without a finile the numberlefs books, effays, and pamphlets which have been written on this fubject. It feems to be the favourite region for the exercife of fiction and fancy. The traditionary tales of ancient miracles, faid to be wrought by holy wells and confecrated fprings, do not much furpafs in extravagance the modern cures afcribed to those favourite haunts of valetudinarians by fraud, ignorance, and credulity. In printed Guides, as they are called, or *Pocket*

OF MINERAL WATERS.

Pocket Companions to any of those fountains of health, it may be excusable to amuse us with a little romance; but we are forry to find the same spirit pervading many medical treatises which should be distinguished by the most facred regard for truth, and a just contempt of puerile embellishments.

Some allowance indeed thould be made for the influence of local prejudices, and for the ftrong bias of intereft and ambition on the mind of a professional man, who, reliding near one of those springs, has, perhaps, no other means of acquiring popularity and reputation, but by an encomium on its virtues, and a description, quite in the poetical flyle, of the beauties of the furrounding fcenery. The latter is, indeed, a harmlefs puff : it feldom deceives any body; but is viewed in the fame light as an auctioneer's advertisement of an eftate, where frightful chafms are often described as curious grottos, a few furze bufhes as a fhrubbery laid out by the finger of nature, and a gallows as a hanging wood. There is another particular, in which those highly coloured landscapes that form the introductory part of almost every treatife on a medicinal fpring, may defeat the propofed end, and that is by exciting a reafonable fufpicion that the waters are chiefly indebted for the high reputation of their efficacy to local circumftances, to the pleafant walks and rides, or the delightful profpects round them.

A difplay of all the attractions of the favourite fpot is commonly followed by very minute details of chemical analyfis, which are just of as little use to the generality of readers as the defcription of the fcenery. A knowledge of the contents of any mineral water by no means implies a knowledge of its medical properties. These are to be learned by repeated experiments. Were it not for the evidence of facts, the late difcoveries in chemistry, as far as they refpect the analyfis of mineral waters, would only tend to leffen their credit, by fhewing the little difference between them and any common water of the fame purity and temperature. For instance, if we confined ourselves to mere fpeculation on the fubject, how could we fuppofe that a quarter of a grain of the oxyd of iron fuspended by a little fixed air in a whole quart of Tunbridge water, the

OF MINERAL WATERS.

the largeft quantity ufually taken in the courfe of a day, could produce any remarkable or peculiar effects? The fame thing might be faid of the most celebrated springs in the kingdom; and many physicians of great profeffional eminence, arguing from this principle, have not hesitated to affert, that the cures performed by those springs were not owing to the ingredients with which they were impregnated, but to the fimple elementary part, or what may be called the pure water. It would not be easy therefore to determine, whether chemical analysis has furnished more arguments in favour of, or against the boasted superiority of mineral waters.

Dr. FALCONER, the author of one of the very few books, relative to this fubject, that contain fomething more than the gratification of idle curiofity, candidly confeffes, that "chymical analyfis, as far as it has been hitherto profecuted, feems to give us a very imperfect view of the methods by which thefe effects (*i. e.* the medicinal effects of the Bath waters) have been produced; and this circumftance has induced feveral perfons to deny the truth of the facts altogether, or to reprefent them as highly exaggerated, and that fuch advantages (if any) as might be in truth received, were owing to collateral circumftances of uncertain and indeterminate efficacy, as change of air, diet, manner of life, and the like."

It is not therefore to the landscape painter, or to the chemift, that we must look for any useful information on those points, but to the modest and judicious practitioner, who, like the author now quoted, watches with care, and reports with fidelity, the bad as well as the good effects of the waters he defcribes, the inftances of their failure, as well as of their fuccels, in various diforders. I am forry to add, that the fund of fuch truly valuable materials is as yet very fcanty, and that I must confine myself to general remarks on the most frequented of our medical springs, fo as to direct invalids to the fountain, from the use of which they may form fome reafonable hopes of relief. As more particular inftructions will often be neceffary when they get to the fpot, I feel it my duty to caution them against choosing for their medical guide any man, however 3

however high his reputation may be, who has diftinguifhed himfelf as the loudeft or most eloquent trumpeter in the indifcriminate praise of the waters near which he refides.

The like caution may prove ftill more ferviceable to fuch of our countrymen as refort to foreign fprings for medicinal purpofes. A popular advocate for the ufe of any remedy is feldom to be relied on as a good phyfician ; and we have always ftrong reafon to fulpect the fkill or the integrity of a man, who fpeaks in a tone of confidence of the infallible efficacy of the waters which he prefcribes. Some of those foreign waters being alfo frequently imported into this kingdom, and ufed here medicinally, I fhall give a fhort account of a few of them, which are found to poffels virtues fuperior to any of our own in the cure of certain diforders.

Mineral waters are ufually claffed according to their fenfible qualities, as perceived by the touch, fight, tafte, and fmell, or according to fome well-known ingredient, which may predominate in this or that particular fpring. The most obvious division is into cold and hot fountains; but both these, being too comprehensive, are again subdivided into *chalybeate*, *faline*, *fulphureous*, and *calcareous*, from their being impregnated with *iron*, *falts*, *fulphur*, or *lime*. There are many still minuter distinctions, where two or more of those ingredients may be found united in the fame spring, or combined with different forts of *air*, which must have a very powerful effect in the internal use of the waters.

The first class of mineral waters, which I shall notice, are those called *chalybeates* from a Greek word that fignifies *iron*, the taste of which is very perceptible in them when fresh from the spring, though they lose it on being exposed for some time to the atmosphere. The reason is, that the small quantity of iron which they contain, being kept in solution by *fixed air*, when this evaporates, the iron sinks to the bottom, forming the sine ochre that lines the channel or water-course. As iron abounds in almost every part of the earth, it is no wonder that fo many springs should be impregnated with it, in a greater or

OF MINERAL WATERS.

or lefs degree according to the quantity of fixed air they contain, by which the iron is held in a flate of folution. Some of those waters have, in conjunction with the iron and fixed air, a pretty ftrong mixture of purgative falt, and are very different from the others in their effects as well as their tafte. In order to diffinguish each by a specific name, the former may be called *fimple chalybeates*, and the latter *faline* or *purging chalybeates*. Tunbridge Wells standing in point of reputation, or of fashionable refort, at the head of the one, and Cheltenham at the head of the other; they may be very properly chosen as examples or illustrations of the various medicinal effects of this numerous class of waters.

It has been already intimated, that the water of Tunbridge Wells is found, upon being analized to differ from that of common springs only by containing in every gallon a grain of iron fuspended by about three tablefpoonfuls in bulk of fixed air. Yet its medicinal effects are very confiderable. It gives a gentle ftimulus to the relaxed nerves, and contributes to reftore their proper tone. It affords great relief in many complaints of the ftomach, in flatulencies, bilious vomitings, irregular or imperfect digeftion, and other confequences of either debility or intemperance. It promotes the circulation of the blood, and the various fecretions; but more particularly that of urine: and this latter circumstance is one of the best proofs of its agreeing with the habit of the patient. In fhort, its natural tendency in the cafes to which its stimulant and tonic powers are adapted, is to raife the fpirits, and increase the general vigour of all the functions.

The Tunbridge waters are eminently ferviceable in what may be called the fexual diforders of females, arifing from a great weaknefs or derangement of the uterine fyftem; fuch as an immoderate flow of the *menfes*, green ficknefs, *fluor albus*, and other fimilar indifpofitions, which are not only relaxing and painful in themfelves, but are often the caufes of abortion, or of fterility. If the profuse flow of the *menfes* fhould be accompanied, as it often is, by feverish fymptoms, by pain in the back, and local irritation, the ftimulus of the waters might then

Qq

prove

OF MINERAL WATERS.

594

prove injurious. Indeed they are improper in all inflammatory cafes, except the feverifh irritation which attends the green ficknefs, and which is more frequently abated than increafed by the ufe of chalybeates.

In all obstructions of the urinary passages, and other complaints in that region, those waters are found of fingular efficacy; for though, as before observed, they in fome degree promote every secretion, yet their chief and most regular determination is to the kidneys; and experience has fully proved the good effects of their gently stimulant and diuretic properties.

We may proceed fiill farther in our recommendation of the Tunbridge waters, and preferibe them with great probability of fuccefs in fuch chronic diforders as arife from flow beginnings, and are attended with great laxity and weaknefs of the folids, but without much organic difeafe. It is neceffary to attend to this material exception; becaufe a general weaknefs may be often brought on by morbid affections of the mefentery, of the lungs, or of fome other important organ, to the cure or relief of which they would be very inadequate. Even in complaints where they have commonly proved efficacious, cafes muft often occur that require the exercife of the niceft judgment and diferimination.

But it is not enough to confider well the propriety of having recourfe to those waters in any particular instance: fome caution is also neceffary in using them. Perfons of full habits fhould not begin to drink them, without lofing a little blood, or without fome other evacuation. When the ftomach is foul, a purgative is commonly preferable to emetics. What the immediate effect of the waters may be, can only be known by trial. They often purge very brifkly at first, but this effect foon ceafes; and as their continued use has fome tendency to occasion coftivenefs, gentle opening medicines from time to time are neceffary. The water itfelf may be eafily converted into a purging chalybeate, by the addition of a little magnefia or Glauber's falts, when neceffary. In the removal of other obstructions, those more especially to which females are subject, the occasional use of the warm bath will be found an excellent auxiliary.

In

In entering upon a courfe of the Tunbridge waters, it is always beft to begin with a fmall dofe, not exceeding a quarter of a pint, about half an hour before breakfast, to be repeated, at regular intervals, once or twice in the forenoon, according to the pleafantnefs of the fenfations it excites. The quantity may be foon augmented to half a pint or even more at each dofe, if agreeable, or neceffary, as the waters lofe much of their medicinal effect by continued use, the ftimulus wearing off in about fix or eight weeks, and making no peculiar imprefiion on the ftomach or habit. The ufual hours for the three dofes are eight o'clock, ten o'clock, and twelve. A light breakfast at nine will not impede the proper action of the waters; and as one of their effects is to improve the appetite, its indulgence fhould be under the control of moderation. But I have elfewhere enlarged fo fully on the advantages of temperance, as well as of early hours and exercife, that I need not ftop here to point out their importance in promoting the falutary operation of any courfe of medicines either natural or artificial.

As fome perfons may be too foon prejudiced against the use of chalybeates by any unpleasant or unexpected fenfations at first, it is proper to inform them, that giddinefs and fometimes a heavinefs of the head, naufea, vomiting, a flight pain about the heart, and a fense of fulnels over the whole body, though by no means uncommon fymptoms on beginning a courfe of these waters, will difappear after a little use. It is only when they stubbornly continue, that they should be regarded as a proof that the waters are not fuited to the nature of the complaint, or to the patient's conftitution. The naulea or fickness being often occasioned by the coldness of the fresh-drawn water, acting on an empty or a very weak and irritable stomach, it is advisable in the first instance not to drink the waters fafting, till the ftomach becomes gradually reconciled to them; and, in cafe of extreme irritability, it is a common and judicious practice at Tunbridge to immerfe in hot water a bottle filled with the chalybeate, and well corked, that the chillneis may be diminished, with as little evaporation of the fixed air as poffible. Where this abounds, as in the waters of the Ger-

Qq2

man

OF MINERAL WATERS.

man Spa, or ftill more in those of Pyrmont, it need not be preferved with fo much caution; but a fufficient quantity of boiling water may be added to that which is taken from the fpring, to bring the whole to a moderate temperature.

There are many fprings of fimple chalybeate water in different parts of Great Britain, and even in the neighbourhood of London, which probably are little inferior in medicinal virtue or intrinfic ftrength to Tunbridge Wells, though these have acquired a higher reputation. Pure air, temperate living, regular and early hours, active diverfions, agreeable company, and a total exemption from all concern, except a rational defire to promote health, are the only circumstances that cause any real difference in the use of waters fo fimilar in quality. This difference, however, is confiderable; and the want of fome of those co-operating circumftances will always prevent the fprings of Islington or Hampstead from being fet in competition with those of Tunbridge, for the certainty of their effects. The remarks, which I have made on the proper method of drinking the latter, as well as on the diforders and particular habits of body to which they are adapted, are equally applicable to all waters of the fame defcription in our ifland. But there are a few chalybeates on the continent, particularly those of the German Spa and of Pyrmont, which, being more active and powerful in their operation, require to be used with greater caution and delicacy.

On drinking a glass of the Spa water, the taste immediately perceives a flrong infusion of iron blended with an agreeable acidity. The latter is owing to the abundance of fixed air, which amounts to near half the bulk of the water, and holds in folution more than four times as much iron as we find in the like quantity of Tunbridge water. The effects are not less perceptible than the taste. A full draught of the Spa water, especially in hot weather, or upon an empty stomach, will produce a swimming in the head, and a fort of intoxication, which fometimes continues for half an hour, and is very like that which arises from spirituous liquors, though it does not leave the fame debility after going off. A patient, therefore, fhould

fhould not drink more than a gill at first, about an hour after breakfast, and another gill after an interval of two hours more. In three or four days, the quantity may be increased, and the dose more frequently repeated; but in these particulars every body's experience will be the best guide. I before suggested the propriety of mixing as much boiling water with the chalybeate as will bring it to a degree of tepid warmth, and thereby prevent any unpleafant senfations, which its natural coldness might excite in a very weak, or a very irritable ftomach.

If fome evacuation was recommended to perfons of a plethoric habit before drinking the Tunbridge water, how much more neceffary must it be to prepare in the same manner for the use of a far stronger stimulant? This very active property of the Spa waters must also render them still more improper than those of Tunbridge in inflammatory complaints, attended with determination to the head, flufhing of the face, or any other ftrong indications of feverish heat. But if the febrile fymptoms are very flight, the Spa waters, ufed cautioufly and moderately, may prove beneficial. They quench thirst more than common water; and they have been frequently known to afford relief in ulcerated fore-throats.

What has been faid of the efficacy of our own fimple chalybeates in cafes of nervous relaxation, or of general weaknefs, in many affections of the ftomach and bowels, in diforders of the kidneys and bladder, or obstructions in the urinary paffages, and in various complaints peculiarly incident to females, may be afferted with still greater confidence of the waters of the German Spa, becaufe of their fuperior strength. Yet they do not equal the activity of the Pyrmont waters, which contain as much iron and twice as much fixed air as those of the Spa, befides a larger proportion of earthy falts. The directions already given, will apply to them both, under fimilar circumstances. Where the stimulus is evidently stronger, more caution is at first necessary; but every patient must learn from experience what quantity of either of those chalybeates the flomach or general habit will conveniently bear; and this quantity, when afcertained, whatever it may be, whether one, two, or three pints in the Qq3

OF MINERAL WATERS.

course of a day, must be continued without farther increase.

It may be thought worthy of remark, that the waters of Pyrmont and Spa, though in their own nature fomewhat intoxicating, afford the most refreshing and wholefome draught, to relieve in the morning a weakness of the stomach occasioned by the too free use of the bowl or the bottle the night before.

The only peculiarity in the virtues or medicinal effects of the Pyrmont waters is, that, when diluted with new milk, they are found very ferviceable in gouty cafes, and may therefore be fafely prefcribed in this form, during the intervals of the fits and the abfence of inflammatory fymptoms.

It is fortunate for a patient who cannot go to drink either of those waters at the fountain head, that they may be conveyed to any distance, and will retain their medicinal properties, with little or no diminution, for two years, if they are inclosed in bottles well-corked and covered with cement. They contain fo much fixed air, that it is usual, after filling the bottles, to leave them uncorked for a while, in order to let the excess of the fixed air escape, as its expansion might burst the bottles, if they were to be instantly corked and removed to a warm place.

The other species of this class of waters, which come next to be noticed, are the faline or purging chalybeates; fo called, partly from their contents and partly from their mode of operation. Befides iron and fixed air, which they have in common with the *fimple chalybeates*, they hold in folution fuch a quantity of purgative falt as gives them a regular and firongly marked determination to the bowels. Any chalybeate, whether of the milder fort, as the Tunbridge water, or of the rougher, as those of the German Spa and of Pyrmont, will often purge brifkly at first; but this is not their certain or constant effect. It feems to depend on the previous habit, or the actual state of the patient's stomach and bowels. It also goes off very foon, and is followed by a tendency to coffivenefs. But the purging chalybeates commonly operate in the fame regular, conitant, and uniform manner, as long as

OF MINERAL WATERS.

as they are continued. The chief fprings of this defcription in England are those of Cheltenham and Scarborough, upon the medicinal properties of which I shall now make a few remarks.

The Cheltenham waters require no preparation; but the quantity fufficient to produce the defired effect on the bowels, can only be known by experience. Half a pint is as much as any patient can well drink at a time; and this may be repeated three or four times, at proper intervals, in the course of the day, according to its operation, or to the intention with which it is taken. Four dofes of half a pint each contain about a quarter of an ounce of purging falts, with fomewhat more than a grain of iron held in folution by four ounces in bulk of fixed air. As purgatives act very differently in different habits, the quantity here mentioned will operate brickly upon fome patients, but fo weakly upon others as to render the occafional addition of the chryftallized falts neceffary, where a powerful and fpeedy effect is defired. On the other hand, the quantity of the water, or the number of dofes, may be leffened, when it is used merely as an alterative. But even perfons of very delicate habits may diveft themfelves of their ufual prejudices against cathartics, as those faline chalybeates do not occafion griping or languor. They only unpleafant fenfation, which they fometimes excite on first drinking them, is a fort of giddinefs or a flight head-ach, which foon goes off; but their agreeable and falutary effects are not fo transient. The improve the appetite, ftrengthen the ftomach, and promote general alertnefs, while they correct and carry off the impurities of the whole fystem. Hence it is that they may be perfevered in for a confiderable length of time, and the body kept moderately open by their uninterrupted ufe, without the least debility or inconvenience. These remarks, however, admit of fome exceptions, particularly when the conflictution is either naturally very weak, or much enfeebled by difeale, without any marks of obstruction, or of acrimony in the A conftant operation on the bowels would in fluids. fuch cafes prove very injurious.

599

Q94

Cheltenham

Cheltenham water has often been found of the greateft fervice in glandular and vifceral obftructions, in a variety of bilious and fcrophulous complaints, in the firft fymptoms of a dropfical difpofition, and in many of the moft diffreffing fcorbutic eruptions and ulcerations on the fkin. But the obftinacy of fome of thefe can only be overcome by perfeverance. In our endeavours to obtain a cure of any difeafe of the chronic kind, we fhould not forget, that complaints, flow in their progrefs, go off alfo very flowly; and that, according to the judicious remark of CELSUS, time is neceffary to remove the deeprooted evils which time has occafioned.

The proper feafon for drinking the waters of Cheltenham is the fummer; and as the warm bath may alfo be advifable in fome of the cafes to which the waters are fuited, the town is well fupplied with accommodations for that purpofe.

. The faline chalybeate at Scarborough does not contain above a third part of the purging falts which are found in a like quantity of the Cheltenham water. Of courfe, it cannot be supposed to operate with the fame force and activity, unlefs larger dofes are taken than moft ftomachs will bear, or unless the powers of the water are increased by the addition of some opening falts of the like kind as those which it already holds in folution. In its natural flate, it is chiefly employed as an alterative; but may, by the artificial means just mentioned, be adapted to all the complaints in which the Cheltenham water is found efficacious. Neither of them will keep well even in clofe bottles, or bear to be carried to any great diftance, as the iron is deposited in a few days, though the faline impregnation continues. By evaporating the water, the purgative falts are procured, for the purpole of being again diffolved in fome more water from the fame fpring, to increase its operation on the bowels.

The inferiority, in point of strength, of the Scarborough water, is more than counterbalanced by other advantages which Cheltenham does not poffefs. In the first place, near the *faline chalybeate* spring there is another

other fpring of fimple chalybeate water, like that of Tunbridge Wells, which is very convenient for patients in cafes where a determination to the kidneys may be defirable. But the fituation of Scarborough on the coaft is an object of still greater importance, as it affords an opportunity of bathing in the fea, fo conducive to the cure of many of the diforders for which those fprings are reforted to. The elevation of the ground, and the uncommon purity of the year, deferve allo peculiar notice in a medical furvey of the local advantages of Scarborough.

It is remarkable, that all the faline chalybeates in our ifland are cold, while many of those on the continent are hot, and are used as baths, as well as internally, in a number of difeafes very difficult of cure. I shall therefore referve my account of one or two of the most celebrated of them, till I come to defcribe our own hot baths, though the foreign ones, on account of their principal ingredients, may be faid more ftrictly to belong to the clafs of waters which I have just been confidering.

The details before entered into respecting the external and internal use of the fea-water, and its admirable effects, not only as a grand prefervative of health, but as an efficacious reftorative of that bleffing in various complaints, preclude the necessity of faying much of the fecond clafs of mineral waters, denominated the fimply faline, and differing from common water only in being impregnated with lome purging falts. We have feveral of these springs near the metropolis; but they are little ufed, as it is fo eafy to procure the fea-water, which is much fuperior to all others of this defcription, in the ftrength of its faline ingredients, and the certainty of its effects. Epfom water, though one of the first of the falt fprings that was brought into use, is now feldom or never prefcribed. Bagnigge Wells, those at the Dog and Duck in St. George's Fields, as well as at Kilburne and Acton, have also had their day of medicinal reputation; but even recommendatory effays and pamphlets can no longer prop up their fame. The portion of falts which they contain is not fufficient to act with certainty on the bowels, unless they are taken in fuch large quantities

602

titite as delicate ftomachs cannot bear, and as very few people can fwallow without difficulty and difgust. Two or three pints must be taken one after the other, in a fhort space of time, to ensure the full purgative effect. On this account, when they were in vogue, it was a common practice to administer more convenient doses, fmaller in quantity, but strengthened by an additional folution of fome of the fame falt as that which they already contained. But this differs fo little from an artificial dofe of phyfic, as almost to destroy the idea of a remedy prepared by nature. Befides, the facility which our infular fituation affords, of procuring falt-water of the greatest efficacy from its grand refervoir, must lessen the value of those fubstitutes. Local convenience has certainly contributed its fhare to the high repute of a fpring of the like kind at Sedlitz, a village in Bohemia, which, being much more ftrongly impregnated with bitter purging falt than the Epfom water, can be more relied upon for its medicinal effects, and is therefore very juftly effeemed in a part of the continent far removed from the fea.

But there is another faline fpring in Germany, a more particular account of which must be interesting to the English reader, not only from the peculiarity of its nature and virtues, but becaufe large quantities of it are imported into this country for medicinal purpofes. Few mineral waters have acquired a greater degree of celebrity than those which are brought from Seltzer, and which may be faid to form a peculiar fpecies, being faline and flightly alkaline, with a ftrong impregnation of fixed air. This quickly evaporates on being exposed to the atmosphere, fo that the water defigned for exportation must be instantly bottled, and kept closely corked, with the mouths of the bottles well covered with cement, or it will foon become not only vapid but putrescent. If well preserved, when poured into a glass, it is perfectly clear and fparkling, and has a gentle faline, and fomewhat pungent or acidulous tafte : but if the fixed air be fuffered to escape, through the leaft neglect, the water appears turbid, is offenfive to the fmell, and entirely lofes its pungency. The ftone bottles

tles, in which it is brought to England, contain about three pints each, a fufficient quantity for a day, to be taken in half-pint glaffes at covenient intervals. Its natural flavour is rather agreeable, and its effects on the fpirits are in general exhilarating. Perfons of very irritable flomachs may dilute it with milk; in which mixed flate it is particularly recommended in cafes of hectic fever with expectoration. It corrects and diminifhes the difcharge from the lungs, checks the violence of the fweats, and contributes very much to the patient's repofe.

The Seltzer water is alfo ufed with confiderable benefit in nervous affections of the ftomach and bowels, in cafes of indigeftion, foulnefs, bilious vomiting, acidity, heart-burn, fpafmodic pains in the alimentary canal, and various difeafes of the urinary organs. Its rapid determination to the kidneys, the action of its ftimulus, and perhaps fome confiderable power as a folvent, may concur to produce the very agreeable effects that are experienced from it in the latter complaints effectially. In any of the former it is advifable to take fome gently opening medicine every two or three days; and the eafieft method of doing this is to add to the dofe of Seltzer water fuch a quantity of vitriolated magnefia as will keep the bowels in a regular ftate.

Though this water cannot be prefcribed with fo much confidence, it may be very fafely tried in miliary puftules, and the like fudden eruptions on the fkin, attended with general irritation I confider it merely as a good diet drink in thefe diforders; for the relief of which, greater reliance is juftly placed on regimen than on medicine.

In a few of the waters already defcribed, we can difcover a flight impregnation of fulphur; but where this principle abounds or predominates, the waters are diftinguifhed by the name of *fulphureous*, from their chief ingredient. The fprings of Harrowgate take the lead in this clafs, and are certainly deferving of the reputation they have acquired, though they are alfo very frequently ufed with great indiference; and as the fame error is very common in drinking the other firong purgative

gative mineral waters, I shall take this opportunity of enlarging upon it.

A very hurtful prejudice prevails in this country, that all difeafes muft be cured by medicines taken into the ftomach, and that the more violently thefe medicines operate, they are the more likely to have the defired effect. This opinion has proved fatal to thoufands, and will, in all probability, deftroy many more, before it can be wholly eradicated. Purging is often ufeful in acute difeafes, and in chronical cafes may pave the way for the operation of other medicines; but it will feldom perform a cure; and, by exhausting the strength of the patient, will often leave him in a worse condition than it found him. That this is frequently the cafe with regard to the more active mineral waters, every perfon conversant in these matters will readily allow.

Strong ftimulants applied to the ftomach and bowels for a length of time, muft tend to weaken and deftroy their energy; and what ftimulants are more active than falt and fulphur, efpecially when thefe fubftances are intimately combined, and carried through the fyftem by the penetrating medium of water? Those bowels muft be ftrong indeed, which can withftand the daily operation of fuch active principles for months together, and not be injured. This, however, is the plan too generally purfued by those who drink the purging mineral waters, and whose circumftances permit them to continue long enough at Harrowgate, and the like places of fafhionable refort.

Many people imagine that every thing depends on the quantity of water taken, and that the more they drink, they will the fooner get well. This is an egregious error; for, while the unhappy patient thinks he is by this means eradicating his diforder, he is often, in fact, undermining the powers of life, and ruining his conftitution. Indeed, nothing can do this fo effectually as weakening the powers of digeftion by the improper application of ftrong ftimulants. The very effence of health depends on the digeftive organs performing their due functions, and the most tedious maladies are all connected with indigeftion.

604

Drinking

Drinking the water in too great quantity not only injures the bowels, and occafions indigeftion, but generally defeats the intention for which it is taken. The difeafes, for the cure of which mineral waters are chiefly celebrated, are moftly of the chronic kind; and it is well known that fuch difeafes can only be cured by the flow operation of alteratives, or fuch medicines as act by inducing a gradual change in the habit. This requires length of time, and never can be effected by medicines which run off by ftool, and operately chiefly on the firft paffages.

Those who wish for the cure of any obstinate malady from the Harrowgate waters, or others of the fulphureous or faline clafs, ought to take them in fuch a manner as hardly to produce any effect whatever on the bowels. With this view, a half-pint glafs may be drank at bed-time *, and the fame quantity an hour before breakfast, dinner, and supper. The dole, however, must vary, according to circumstances. Even the quantity mentioned above will purge fome perfons, while others will drink twice as much without being in the least moved by it. Its operation on the bowels is the only standard for using the water as an alterative. No more ought to be taken than barely to move the body; nor is it always neceffary to carry it even this length, provided the water goes off by the other emunctories, and does not occasion a chillnefs or flatulency in the flomach or bowels. When the water is intended to purge, in cafes where the nature of the patient's complaint requires a ftrong determination to the bowels, it may be neceffary to drink a pint or two before breakfaft.

I would not only caution patients who drink those waters over night, to avoid hearty suppers, but also against

• When I fpeak of drinking a glafs of the water over night, I muft beg leave to caution thole who follow this plan againft eating hearty suppers. The late Dr. DAULTRY of York, who was the first that brought the Harrowgate waters into repute, used to adwife his patients to drink a glass before they went to bed; the confequence of which was, that having eat a flesh supper, and the water operating in the night, they were often tormented with gripes, and obliged to call for medical allistance. 606

eating heavy meals at any time. The ftimulus of water, impregnated with fulphur and falts, feems to create a falfe appetite. I have feen a delicate person, after drinking the Harrowgate waters of a morning, eat a breakfast fufficient to have served two ploughmen, devour a plentiful dinner of flefh and fifh, and, to crown all, eat fuch a supper as might have fatisfied a hungry porter. All this, indeed, the ftomach feemed to crave; but this craving had better remain not quite fatisfied, than that the ftomach should be loaded with what exceeds its powers. To ftarve patients was never my plan; but I am clearly of opinion, that, in the ufe of all the ftrongly purging mineral waters, a light and rather diluting diet is the most proper; and that no perfon, during such a course, ought to eat to the full extent of what his appetite craves.

Exercife is not lefs conducive to the falutary end in view than temperance. It promotes the operation of the waters, and carries them through the fystem. It may be taken in any manner that is most agreeable to the patient; but he ought never to carry it to excels. I fcarcely need repeat a remark often made in other parts of this work, that the beft kinds of exercise are those connected with amufement. Every thing that tends to exhilarate the fpirits, not only increases the efficacy of the waters, but acts as a medicine. All those who repair to the fountains of health, ought therefore to leave every care behind, to mix with the company, and to make themfelves as cheerful and happy as poffible. From this conduct, affifted by the free and wholefome air of those fashionable places of refort, and also the regular and early hours which are ufually kept, the patient often receives more benefit than from using the waters.

During my refidence at Harrowgate, I met with many inftances of the most mischievous effects produced by drinking the waters in cases where they were absolutely improper, and adverse to the nature of the difease. When people hear of a wonderful cure having been performed by some mineral water, they immediately conclude that it will cure every thing, and accordingly swallow it down, when

when they might as well take poifon. Before patients begin to drink the more active kinds of mineral waters, they ought to be well informed of the propriety of the courfe, and fhould never perfift in using them, when they are found to aggravate the diforder.

On the other hand, I often witneffed the happy iffue of experiments made with judgment and caution at Harrowgate, when the greatest benefit was derived from the proper use of the waters in various eruptions on the fkin, of the most distreffing nature; in rheumatism complicated with fcorbutic complaints; in obstructions of the glandular and lymphatic fystem; and in difeases of the first passages, accompanied with, or proceeding from, inactivity of the ftomach and bowels, acidity, indigeftion, vitiated bile, worms, putrid fordes, the piles, and jaundice. They answer two very important purposes; first, when taken in small quantities, acting as an alterative, and inducing, by their mild operation, a gradual change in the habit; and, fecondly, when employed in larger dofes where purging is indicated, fulfilling that intention in the most defirable manner, without irritating the nerves, or weakening the patient fo much as other purgatives. After a little ufe, almost every body can drink them without any great difgust, though they are at first no less nauseous to the taste, than offensive to the fmell.

I fhall only add, that the external use of the Harrowgate waters being justly deemed a very powerful auxiliary in many of the diforders for which they are reforted to, particularly those of the cutaneous class, there are proper baths for this purpose, to the supply of which three springs out of four are devoted, that which is referved for drinking being more strongly impregnated with solution and supplies that the rest.

The fulphureous and faline waters of Moffat in North Britain are almost as much reforted to as those of Harrowgate. The impregnation of the former, indeed, is not fo strong as that of the latter, and their effects are of course fomewhat different. Two or three quarts of the Moffat water may be drank in a morning, without any fensible effect but that of increasing the flow of urine. It now

607

and

608

and then purges; but this is fo far from being its conftant or regular mode of operation, that opening medicines are almost always neceffary during a course of it. Its evident determination to the kidneys renders it of effential fervice to perfons afflicted with the ftone and gravel, particularly the latter. It has likewife afforded great relief in many bilious complaints, and in the early fymptoms of a scrophulous habit. But its chief point of celebrity, and that in which it may be faid to rival the fprings of Harrowgate, is the cure of cutaneous eruptions of every kind. In these cases, the external application of the water, warmed to a confiderable temperature, is very judiciously made a material part of the plan of treatment. One difadvantage, however, attends this procefs both at Moffat and Harrowgate. The waters, while heating, unavoidably lofe in vapour fome of their fulphureous impregnation, on which part of their efficacy, even when externally applied, muft depend. So far, therefore, a preference is justly due to the naturally hot fulphureous fprings of Aix-la-Chapelle and Bareges, of which I shall take some farther notice, after I describe the thermal fprings in our own ifland, which conflitute the next and laft clafs in my arrangement of mineral waters.

In the introductory part of this fection, the waters which now remain to be confidered are fpecifically diftinguished by the title of calcareous, because they contain more lime or calcareous falts than they do of any other folid fubstance. I adopted this title, however, in compliance with cuftom, and merely as a nominal diftinction, rather than from confidering it as an important characteristic, or by any means expressive of the grand caufe of the efficacy of fuch waters. It is not to chymical analyfis, but to experience that we are indebted for a knowledge of their virtues; and, inftead of giving them a name taken from a part of their contents, which conveys no uleful information, we had better fimply call them hot fprings, as every body will then have a clear idea of the molt perceptible difference between them and all the other mineral waters in our ifland.

but that of more ne the flow of writer then

In this part of our defcription, the hot fprings Bath have the most indisputable claims to preceden The fame of their medical virtues is more widely diffused and more firmly established than that of any other fprings in the known world. It is no wonder, therefore, that the pen of industry, of genius, or of ambition, fhould have been often exercifed upon fo popular a topic. Amidit fuch a multitude of books, Dr. FAL-CONER's " Practical Differtation on the Medicinal Effects of the Bath Waters," is acknowledged to be the beft account which has yet appeared of what he juftly calls a "powerful, but (in many inftances) nice remedy." His fituation as phyfician to the Bath Hofpital must have afforded him the beft opportunities of obferving the effects of those waters; and he has given the most fatisfactory proofs of his being well qualified to profit by those opportunities. One of his remarks is really of more confequence than a whole volume of chymical inveftigations, which have thrown just as little light upon the efficacy of the waters as the fables concerning BLA-DUD, or the frequently difcovered fragments of Roman antiquities. Chymistry, indeed, makes us acquainted with the component parts of those waters, and tells us they contain a good deal of calcareous falts, but little, if any, neutral alkaline falts; and that they are impregnated with about a fixtieth part in bulk of fixed air, which holds in folution fo very fmall a quantity of iron as to be fearcely appreciable, though it gives a flight chalybeate tafte to the water when hot from the fpring. But what inference could we deduce from these and the like amufing details, in the treatment of any particular difease? It is then to the enlightened practitioner, as I faid before, that we must look for useful information; and, in this view of the fubject, we cannot with for more respectable authority than that of Dr. FALCONER.

According to this writer, the Bath water, when drank fresh from the spring, has in most perfons the effect of raising and rather accelerating the pulse, increasing the heat, and exciting the secretions. The action on the nervous system is felt at the same time, so that the sti-

mulating

mulating properties of the water must be very diffusive; and, what is farther remarkable, though these fymptoms come on fuddenly, yet they often continue much longer than we can suppose them to be excited by the actual prefence of the water in the body.

The Bath waters not only promote urine and perfpiration, but also increase the falivary difcharge; and quench thirst better than any other fluid, in cases where there is no tendency to fever. If any disposition of this kind should appear, a hot stimulant would be evidently improper.

It has been generally imagined, that those waters were fomewhat aftringent, on account of the coffiveness which frequently accompanies the use of them. But this effect is more reasonably aferibed to their heating qualities, and to their power of exciting the other fecretions. As a proof of this, when perspiration is checked in cold weather, they fometimes prove flightly purgative.

The relief which the Bath waters afford in the cholic, in convultive retchings which often attend the gout in the ftomach, and in many other fimilar affections, is a fufficient evidence of their anti-fpafmodic powers.

This account of the primary and immediate effects of the water ufed internally will enable any patient to judge from his own feelings, whether it agrees with his conftitution or not. If it excites, on being first taken, a pleafing glow in the stomach, followed by an increase of spirits, and of appetite, particularly for breakfast, and, above all, a rapid determination to the kidneys, there is the greatest probability of its proving ferviceable. But if it occasions head-ach, thirst, and dryness of the tongue;—if it fits heavy on the stomach, or produces stickness, and does not pass off by urine or perspiration;—it may be fairly concluded, that its continuance would do injury, unless these symptoms can be removed.

If we come next to confider the external application of those waters, we shall find, in the first place, that they unite all the medicinal advantages of warm baths, from

from about a hundred and fix degrees of heat to any inferior degree that may be defired. The extent also of the baths, which affords room to move about in them freely, and the permanence and uniformity of their warmth, are no unimportant recommendations. But Dr. FALCONER is of opinion, that the Bath waters polfels some farther powers or specific qualities superior to those of common water of the fame temperature. He thinks that their action on the nervous fystem is more ftimulating than a common warm bath ;---that they raife the pulfe and heat of the body to a higher degree, yet are much less apt to produce a violent perspiration ;that they remarkably increase the urinary discharge ;--and that, fo far from caufing any relaxation or weaknefs, the bathers are observed to be in general more alert and vigorous, and to have a better appetite on the days of bathing than in the intervals. As far as my own opportunities of obfervation extended during a few fhort vifits at Bath, they are in perfect concurrence with the Doctor's opinion.

The difeafes, in which this eminent practitioner very accurately defcribes the good effects of the Bath waters, are the green ficknefs, particularly before any confiderable affection of the ftomach takes place, or any feverifh fymptoms appear; vifceral obstructions, when the confequences of intermittent fever, or of long refidence in hot climates, if the diforder in these cases has not advanced too far; the palfy, from a great variety of caufes; the gout, in that Itage of the complaint, when the inflammatory fymptoms, if any have preceded, have in a good meafure abated, and a degree of weaknefs and want of tone in the fystem begins to take place: the chronic rheumatism, and the acute also, provided the feverifh disposition be previously allayed by proper evacuations; white fwellings on the knee; hip cafes; weakness of the organs of digestion ; the cholic, accompanied with hysteric fymptoms, or produced by the poifon of lead; the jaundice, when arifing from fimple obstruction of the biliary ducts; hypochondriac and hyfterical complaints; St. Vitus's Dance; spafmdoic affections of the womb and painful menstruation; and, lastly,

Rr 2

ly, in many cutaneous, but more especially leprous eruptions.

I have confined myfelf here to a bare outline, which may be fufficient for the general direction of valetudinarians, who muft avail themfelves of more particular advice at the fountain head. Almost every cafe will require a peculiar mode of treatment; and great caution will be found neceffary to prevent fatal mistakes. I cannot too often repeat, that the more powerful any remedy is, the more liable it is to abufe; and though the efficacy of the Bath waters has been fully established in a variety of the most stubborn and afflicting diforders, yet their mistapplication has also been often attended with very ferious confequences.

Even when the use of the Bath waters may be deemed fafe or proper, confideration must be had, as Dr. FAL-CONER justly observes, to the quantity taken in, when they are drank; and to the time of stay in the bath, heat of the water, &c. when they are externally used.

The fafeft method is to begin with drinking a glafs, containing about a quarter of a pint, before breakfast; and to judge from its effects how often it may be repeated, or the quantity enlarged. If it feels eafy, warm, and cordial, to the ftomach and fpirits, and excites no pain or fense of fulness in the head or eyes, a fecond glafs of the fame fize may be taken the fame day at noon, and this quantity gradually increased to a pint in the day, taken at two or three times, as convenience may fuit. This is deemed a proper medium for the generality of people, though in fome cafes, where the habit is not very irritable, the daily allowance may be augmented to a pint and a half, with fafety and benefit. The Crofs Bath water, the temperature of which, when fresh from the spring, is 112°, is usually recommended at the commencement of a courfe, efpecially when there is any affection of the head, or when any tendency to plethora appears in the fystem. After some time, the more powerful water of the King's, or the Hot Bath, which is four degrees higher in temperature, may be used; and this change will prevent, in some measure, the

612

the difgust which is apt to come on after a long trial of the bath waters.

No lefs regard fhould be paid to the proper use of the bath. A fhort ftay of five or fix minutes is most advifeable at first; and if this trial produces no fymptoms that are difagreeable, but, on the contrary, feems to improve the health, fpirits, and ftrength, a longer ftay may be gradually indulged, till it comes to half an hour, but never to cause lassitude, faintness, or difgust. The choice of the baths is often a matter of importance, as there is a still greater variation in the temperature of the waters when in the baths than at the pumps, the waters in the King's, or the Hot Bath, being from eight to twelve degrees warmer than those in the Cross Bath. There are also private baths of any temperature to fuit the peculiarities of every patient's habit or complaint. The beft time in general for bathing is the morning; and it may be repeated twice, or at most thrice a week. As the public baths are emptied foon after nine o'clock, a much earlier hour is ufually chofen for bathing; but the private baths may be prepared at any hour of the day. Where a greater ftimulus than mere immerfion is thought neceffary, or where a partial application of the water is deemed preferable to a general one, the part may be pumped upon, fo as to increase the forcible action of the water on that part, and yet prove lefs heating to the fystem at large.

There are no fprings in our island befides those at Bath, which can be strictly called hot, though a few others have got the name, and are referred to this class of thermal waters, from being, invariably in every feafon, and independently of the state of the atmosphere, warmer than the general average of the heat of common springs. The temperature of Buxton waters is only 82°, yet approaches nearess to that of the hot springs at Bath, which, in their cooless state, are never lower than 90°. In other fensible properties the Buxton water differs very little from common pump water.

The

The medicinal effects of the Buxton waters, however, are far from being inconfiderable. Taken internally in fmall dofes at different intervals, amounting to near a pint before breakfaft, and the like quantity before dinner, they afford very great relief in heart-burn, flatulency, ficknefs at the ftomach, and other diffreffing fymptoms of defective digeftion and derangement of the alimentary organs, the fure attendants of indolence and luxury. In fuch cafes, the body muft be kept regularly open by means of medicine during the ufe of the waters, if thefe do not, as they fometimes will, produce a laxative effect, which is always falutary. They are no lefs ferviceable in alleviating painful affections of the kidneys and bladder; and here alfo the external ufe of the waters is often found to promote their internal operation.

But the employment of the Buxton waters as a bath is not confined to the relief of gravelly complaints. They contribute far more effectually to reftore the healthy action of particular limbs, that may have loft their moving powers in confequence of long or violent inflammation, occafioned either by external injury, or preceding acute rheumatifm. But if the rigidity, weaknefs, or impaired functions of the parts affected be the confequence of a paralytic ftroke or of gouty inflammation, more benefit will be experienced from the warmer temperature of the Bath waters.

In general, the Buxton bath is well fuited to fuch a languid, enfeebled, or very irritable flate of the habit, as cannot bear abfolutely cold water, but may be excited to eafy and falutary re-action, by a milder flimulus. The flight flock, which people feel at the inflant of immerfion in the Buxton waters, is almost immediately followed by a pleafant glow all over the body; and this is precifely the effect aimed at. But I have expressed myself fo fully on the nature of the particular infirmities to which the lukewarm bath is adapted, as to render any farther obfervations on this head unneceffary.

The tepid fprings at Matlock approach ftill nearer to cold water than those at Buxton, being fixteen degrees lower in point of temperature. The flock, which they give on immersion, is consequently stronger, and requires greater

greater powers of re-action in the habit ; but they form a good intermediate bath between Buxton and the fea. They are therefore very properly employed in preparing invalids for the latter, when this is neceffary to complete a cure, as is often the cafe in the chronic rheumatifm. They are used internally as a pure diluting drink, but are not known to poffefs any other remarkable medicinal properties.

The Hotwell, as it is improperly called, near Briftol, is not, indeed, fo cold as the fprings at Matlock, but does not rife to the temperature of those at Buxton, being only 74°, or twenty-two degrees below the ordinary warmth of the blood in a ftate of health. The Hotwell water is not made use of to bathe in; but, taken internally, it has acquired very high reputation for curing the incipient fymptoms of confumption, and affording confiderable relief in the more advanced ftages of this difeafe. As the effects it produces are very gradual, its agreeablenefs to the palate is a fortunate circumstance for a class of patients who can only hope by long and fleady perfeverance to check the progress of one of the most infidious deftroyers of the human species. There is no doubt but they are also indebted for a part of the relief they experience to the mild, fheltered, yet fufficiently ventilated fituation of the Hotwells, and to the judicious plan of diet, exercife, and amufements purfued there.

The other complaints, in which the purity and temperature of the Hotwell water render it very grateful and of no fmall efficacy, are relaxations of the ftomach and bowels, brought on by long refidence in hot climates; bilious diarrhœa; flight dyfentery; and a diforder ftill more difficult of cure than any of these, I mean the diabetes. I do not know any natural remedy better adapted to relieve its various fymptoms; fuch as the conftant thirft, the impeding perfpiration, the drynefs and frequent defquamation of the skin, the feverilh quickness of the pulfe, and the preternatural difcharge of urine. A regular courfe of the Hotwell water has been found to moderate this almost unquenchable thirst; to keep the fkin moift and perfpirable; to allay the fever; and to Rr4 render

render the urinary organs fitter to receive benefit from the medicines ufually prefcribed to remove their debility and that of the fyftem in general. It may be farther neceffary to obferve, that, as this water contains but a very fmall quantity of purgative falts, which are alfo counteracted by a larger folution of calcareous falts, it must have a stronger determination to the kidneys than to the bowels, so that the use of a gentle aperient medicine becomes in feveral cafes adviseable.

Particular habits and complaints require variations in the quantity to be taken of this, as well as of any other mineral water. The full dofe is a half pint, to be drank early in the morning and repeated before breakfaft, at the interval of at least half an hour fpent in gentle exercife. Two more dofes of the like quantity, and with the like interpolition of active amusement, are to be taken between breakfast and dinner, at the longest diltance from each of those meals. As this water is also used at table and for domestic purposes, every invalid unavoidably takes more than the above quantity every day; but in confequence of its altered temperature, and of the lofs of the fixed air it contained, it cannot be in fo high a ftate of medicinal perfection as when drank fresh from the fpring. The fame remark must of course apply to the immense quantities that are bottled for exportation, though the almost proverbial purity and foftnefs of the water, as well as its excellent property of keeping untainted for a great length of time in hot climates, must render it a very valuable water for long voyages.

I fhall fubjoin to these observations on the medicinal virtues of our own thermal waters, a short account of the most celebrated hot springs on the continent.

There is none of the foreign watering places more reforted to than the German Spa, of which I already took notice among the cold chalybeates, and Aix-la-Chapelle, about twenty-four miles diftant from the former, and equally famous for waters of a very different quality and temperature. Their celebrity is traced back even to the days of CHARLEMAGNE, who refided for a long time at Aix, and took fo much delight in the use of the waters,

8

as

as frequently to hold his levee in the bath, with all his attendants.

The exact degree of the heat of thefe waters is given differently by different observers; but, taking the average of their accounts, it may be reckoned in the well of the hotteft bath at 140°, and at the fountain where it is drank, about 120°. It requires to ftand feveral hours in the large baths, before it is fufficiently cooled for tepid bathing, without the addition of cold water. It contains an uncommon quantity of fulphur, and emits a fmell like that of Harrowgate water, but far more penetrating. From its heat, and its ftronger impregnation of the fulphureous principle, it is also more powerful in all the difeafes of the fkin, for which Harrowgate is reforted to. The baths at Aix-la-Chapelle are looked upon as a certain cure for almost every cutaneous eruption; but the water fhould be used internally at the fame time, to carry off impurities, and keep up the full perfpiration promoted by the bathing. These baths are equally ferviceable in stiffness of the joints and ligaments, which is left by the inflammation of gout and rheumatifm, and in the debility of palfy, where the highest degree of heat which the fkin can bear is required. In obstinate cafes, the vapour bath, formed by the steam of those very hot waters, is recommended by the German phyficians.

Numberlefs inftances ferve to eftablifh the efficacy of the waters of Aix as an internal medicine in painful affections of the kidneys and bladder, as well as in diforders of the ftomach and biliary organs occafioned by luxurious indulgence and intemperance. The common dofe is half a pint, to be repeated more or lefs often, according to its fenfible effects, and to the intention with which it is prefcribed, either as a purgative, or a diuretic. It is a ftriking proof of the power of habit, that the palate and ftomach are foon reconciled to the ufe of fuch waters, though at first extremely difgusting and naufeous.

The hot fulphureous fprings at Bareges, two little hamlets on the French fide of the Pyrenean mountains, are, indeed, inferior in the degrees of heat and ftrength of impregnation to the waters of Aix-la-Chapelle, yet are found very beneficial in the like complaints. Their highly

highly detergent powers, owing perhaps to a finall portion of oily or bituminous matter, added to the other medicinal properties which they poffefs in common with thole of Aix, render them peculiarly efficacious in deepfeated ulcers, and thole morbid affections of the womb, which French ladies call, though not with medical correctnefs, *Depôts de lait*. In fuch cafes, the waters are injected. They are also very frequently used in the form of *douche*, or by way of pumping on the part affected, as well as for general bathing.

In the interior parts of France, particularly in the central provinces of Auvergne and the Bourbonnois, now included in the Department of the Loire, there are feveral hot fprings, but of the faline chalybeate clafs, the efficacy of which, as an internal medicine, is greatly increased by their higher degree of temperature, in the diforders for which Cheltenham is reforted to in our own country. Those French springs derive another very important advantage from their heat, that of being uled as a bath in all cafes which may require that falutary flimulus on the furface. In many of these, the internal and external use of the waters co-operate with wonderful effect; and particularly in the fexual complaints of women, arifing from any defect or irregularity in the functions of the uterine organs. CATHERINE DE MEDICIS, the mother of feveral French princes, is faid to have been much indebted for her fertility to the waters of Bourbon Lancy, not far from the town of Moulins, a place configned to immortal fame by STERNE's affecting ftory of MARIA.

There is a village on the confines of Bohemia, where the waters may be faid, in the flricteft fenfe of the word, to boil up with vehemence from the fpring, and are often ufed for fealding hogs and fowls, to loofen the hair and feathers, their heat being quite fufficient for thefe purpofes. The temperature of the *Prudel*, or *furious* fountain, as it first iffues forth, is as high as 165°, and keeps invariably to the fame point. Of courfe it requires to be very much cooled before it can be ufed as a bath, or even drank. Thofe waters are faid to have been reforted to, and first brought into confiderable notice, by the emperor CHARLES IV. in 1370; to which circumstance the village owes its name of *Carlfbad*, or Charles's bath,

The natural hiftory of its waters would afford a great variety of curious and interefting particulars; but the limits of my prefent plan confine me to a fhort medical notice of their extraordinary virtues in all the difeafes for the cure of which *faline chalybeates* are internally or externally employed.

CHAP. LVI.

OBSERVATIONS CONCERNING THE DIET OF THE COMMON PEOPLE.

E XPERIENCE proves that not a few of the difeafes incident to the inhabitants of this country, are owing to their mode of living. The vegetable productions they confume, fall confiderably fhort of the proportion which they ought to bear to the animal part of their food. The conftant use of bread and animal fubftances excites an unnatural thirft, and leads to the immoderate use of beer and other flimulating liquors, which generate difease and reduce the lower orders of the people to a ftate of indigence. To teach the poor man how to live cheaper and better, is the defign of the following pages.

Though the common people of this country live at a greater expence than any where elfe, it does not follow that they live better. They are firong indeed, but by no means healthy; and it is found that, from an attachment to a particular mode of living, they are more liable to difeafe and death in foreign climes, than the inhabitants of any other country.

It is certainly proper that the poor man fhould be inftructed in every thing that can make his little earnings go as far as poffible, or which can add to the comfort of himfelf and family. Nor can œconomy in living be deemed a trivial virtue, in a country where the riches depend on the cheapnefs of labour.

620 DIET OF COMMON PEOPLE.

It is alleged that the English are for much attached to their own modes of living, that no argument will induce them to make the smalless change. Habits are indeed obstinate things, especially those which relate to diet; but there are proofs that the English are not inflexible even in this matter. The mode of living among the lower orders has been greatly changed in my time, and I am forry to fay, not for the better.

The people of England have too much good fenfe not to liften to reafon, provided due care were taken to inftruct them. But here the people may be truly faid " to perifb for want of knowledge." No means have been ufed to give them proper inftruction. Hurtful cuftoms have been fuffered to prevail, till they have ftruck fuch deep roots that it will not be an eafy matter to eradicate them. The difficulty, however, is not unfurmountable. A few experiments of reform would have the effect to render it as agreeable as it is falutary.

Adults have many old prejudices to overcome, but the cafe is different in regard to children. They may be taught to ufe any kind of food, and what they ufe when young they will love when old. If I can introduce a different method of feeding children, my purpofe will be anfwered. This alone will, in time, effect a total change in the general mode of living.

The late diffrefs of the poor has called forth many publications intended for their relief. Moft of them, however, were adapted only for the particular occafion, and not calculated to prevent the return of like evils. The following obfervations, it is hoped, will have a more permanent effect. They are intended to recommend a plan of living, which will render the people lefs dependent on bread and animal food for their fubfiftence, and confequently not fo liable to fuffer from a fcarcity or dearth of either of thefe articles in future.

Particular attention has been paid to the fubfitutes for bread, as the fcarcity of this article proves peculiarly diffreffing to the poor. It will appear from the following pages, that bread is by no means fo much a neceffary of life as is generally imagined, and that its place may, in many

DIET OF COMMON PEOPLE. 621

many inftances, be fupplied by a variety of other farinaceous fubftances.

GENERAL OBSERVATIONS ON ALIMENT.

No creature eats fuch a variety of food as man. Intended for an inhabitant of every climate, he devours the productions of them all; and if they do not fuit his palate or agree with his flomach, he calls in the aid of cookery, an art peculiar to himfelf; by which many things that, in a crude flate, would prove hurtful, or even poifonous, are rendered wholefome and falutary.

The obvious division of food is into animal and vegetable. To fay that man was intended by nature for using either the one or the other alone, would be absurd. His structure and appetite prove that he was formed for both. Judgment, however, is requisite, in adjusting the due proportions of each, so as to avoid the inconveniences arising from an extreme on either hand.

Though animal food is more nourifhing than vegetable, it is not fafe to live on that alone. Experience has fhewn that a diet, confifting folely of animal food, excites thirft and naufea, occasions putrefcence in the ftomach and bowels, and finally brings on violent griping pains, with cholera and dyfentery.

Animal food is lefs adapted to the fedentary than the laborious, and leaft of all to the fludious, whole diet ought to confift chiefly of vegetables. Indulging in animal food renders men dull, and unfit for the purfuits of fcience, especially when it is accompanied with the free use of strong liquors.

The plethoric, or perfons of a full habit, fhould eat fparingly of animal food. It yields far more blood than vegetables taken in the fame quantity, and of courfe may induce inflammatory diforders. It acts as a ftimulus to the whole fystem, by which means the circulation of the blood is greatly accelerated.

I am inclined to think that confumptions, fo common in England, are in part owing to the great use of animal food. Though the *Phibifis Pulmonalis* is not, properly

622 OBSERVATIONS ON ALIMENT.

properly fpeaking, an inflammatory difeafe, yet it generally begins with fymptoms of inflammation, and is often accompanied with them through its whole progrefs.

But the difeafe most common to this country is the fcurvy. One finds a dash of it in almost every family, and in some the taint is very deep. A difease so general must have a general cause, and there is none so obvious as the great quantity of salted animal food devoured by the natives. As a proof that source arises from this cause, we are in possession of no remedy for that difease equal to the free use of vegetables.

By the uninterrupted use of animal food, a putrid diathesis is induced in the fystem, which predisposes to a variety of diforders. I am fully convinced, that many of those obstinate complaints for which we are at a loss to account, and find it still more difficult to cure, are the effects of a scorbutic taint lurking in the habit.

Improper diet affects the mind as well as the body. The choleric difposition of the English is almost proverbial. Were I to affign a cause, it would be, their living fo much on animal food. There is no doubt but this induces a ferocity of temper unknown to men whose food is chiefly taken from the vegetable kingdom.

Though thefe and fimilar confequences may arife from the excels of animal diet, we are far from difcouraging its ufe in moderation. In all cold countries it is certainly neceffary; but the major part of the aliment ought neverthelefs to confift of vegetable fubftances. There is a continual tendency in animal food, as well as in the human body itfelf, to putrefaction, which can only be counteracted by the free ufe of vegetables.

With regard to the proportion of vegetable food to that of animal, great nicety is by no means required. It must vary according to circumstances, as the heat of the weather, the warmth of the climate, and the like. The vegetable part, however, where nothing forbids, ought certainly to preponderate, and I think in the proportion at least of two to one.

OBSERVATIONS ON ALIMENT. 623

The exceffive confumption of animal food is one great caule of the fcarcity of grain. The food that a bullock affords bears but a fmall proportion to the quantity of vegetable matter he confumes.

I am no enemy to good fruit, as an article of diet; but the greater part of what is ufed in this country, by the lower orders of the people, is mere trafh. Fruit should be eaten in the early part of the day, when the stomach is not loaded with food, and it never ought to be eaten taw till it is thoroughly ripe.

OF BREAD.

Bread, or fomething refembling it, makes a part of the diet of all nations. Hence it is emphatically denominated the ftaff of life. It may, however, be used too freely. The late Dr. Fothergill was of opinion, and I perfectly agree with him, that most people eat more bread than is conducive to their health. I do not mean to infinuate that bread is unwholefome, but that the best things may prove hurtful when taken to excess. A furfeit of bread is more dangerous than of any other food. Omnis repletio mala, repletio panis pession. The French confume vast quantities of bread; but its bad effects are prevented by their copious use of sources and fruits, which have little or no source in the diet of the common people of England.

One important use of bread is to form a mass fit for filling up the alimentary canal, and carrying the nutricious juices along that passage in such a state, as to render them fit to be acted upon by the lacteal absorbents, which take up the nourishment and convey it to the blood. In this light, bread may be confidered as a foil from whence the nourishment is drawn. I do not fay that bread contains no nourishment, but that its use, as an article of diet, does not solely depend on the quantity of nutriment it contains, but in some measure on its fitness as a vehicle for conveying the nutricious particles through the intessinal tubes. Hence it follows that the finess of nutrition. The richeft food will not nourifh an animal, unless the alimentary canal is fufficiently diftended. A dog has been fed on the richeft broth, yet could not be kept alive; while another, which had only the meat boiled to a chip and water, throve very well. This fhews the folly of attempting to nourifh men on alimentary powders and other concentrated food.

The great art, therefore, of preparing food, is to blend the nutritive part of the aliment with a fufficient quantity of fome light farinaceous fubftance, in order to fill up the canal, without overcharging it with more nutricious particles than are neceffary for the fupport of the animal. This may be done either by bread, or other farinaceous fubftances, of which there is a great variety, as will appear from the fequel.

Bread is one of the most expensive modes of using grain, and not adapted to the narrow circumstances of the lower orders of the people, as it is burthened with two heavy additional charges, in passing through the hands of both the miller and the baker. Besides, the former often grinds down extraneous matter with the wheat, and the latter as frequently bakes it up with the addition of lime, chalk, alum, and other pernicious substances. Since the articles of diet have become branches of manufacture, the public neither know what they eat, nor what they drink.

People imagine, as the fineft flour contains the greatest quantity of nourifhment, that it must therefore be the most proper for making into bread; but this by no means follows. The finest flour comes the nearest to starch, which, though it may occasionally prove a good medicine, makes bad bread. Household bread, which is made by grinding down the whole grain, and only feparating the coarser bran, is without doubt the most wholesome.

The beft household bread I ever remember to have ate, was in the county of York. It was what they call *mession bread*, and confisted of wheat and rye ground together. I am not quite certain as to the proportion, but I think there might be two parts of the former to one of the latter. This bread, when well fermented, eats light, is

is of a pleafant tafte, and foluble to the bowels. After using it for some years, I found that bread made entirely of flour was neither so agreeable to the palate, nor so conducive to health.

Bread is often fpoiled to pleafe the eye. The artificially whitened, drying, ftuffing bread, though made of the heart of the wheat, is in reality the worft of any; yet this is the bread which most people prefer, and the poorer fort will eat no other.

All the different kinds of grain are occafionally made into bread, fome giving the preference to one and fome to another, according to early cuftom and prejudice. The people of South Britain generally prefer bread made of the fineft wheat flour, while those of the northern counties eat a mixture of flour and oatmeal, or rye-meal, and many give the preference to bread made of oatmeal alone. The common people of Scotland alfo eat a mixed bread, but more frequently bread of oatmeal only. In Germany the common bread is made of rye, and the American labourer thinks no bread fo ftrengthening as that which is made of Indian corn ; nor do I much doubt but the Laplander thinks his bread made of the bones of fishes is the beft of any.

Bread made of different kinds of grain is more wholefome than what is made of one only, as their qualities ferve to correct one another. For example, wheatflour, especially the finer kind, being of a starchy nature, is apt to occasion constipation. Bread made of rye-meal, on the other hand, proves often too stippery for the bowels. A due proportion of these makes the best bread.

For the more active and laborious I would recommend a mixture of rye with the ftronger grains, as peas, beans, barley, oats, Indian corn, and the like. Thefe may be blended in many different ways: they make a hearty bread for a labouring man, and, to use his own language, they lie longer on his ftomach than bread made of wheat-flour only. Barley bread paffes too quickly through the alimentary canal to afford time for conveying the proper nourifhment; but bread made of barley mixed with peas is very nourifhing,

When

OF BREAD.

When potatoes, or boiled grain, are used, bread ceases to be a necessary article of diet. During the late fcarcity of bread, I made it a rule not to eat above one half the quantity I used to do, and I found no inconveniency whatever from the change. Nay, fome told me, that for a confiderable time they had left off the use of bread altogether, without experiencing any change in the ftate of their health.

A great part of the bread confumed in this country is by children. It is always ready, and when the child calls for food, a piece of bread is put into its hand, to fave the trouble of dreffing any other kind of victuals. Of many children this is the principal food, but it is far from being the most proper. Children are often troubled with acidities of the stomach and bowels; and it is well known that bread mixed with water, and kept in a degree of heat equal to that of the human stomach, foon turns four.

During the late fcarcity, many of the labouring men, and even artificers, could not earn as much money as was fufficient to keep their families in the article of bread only. It is certain, however, that on a different plan, fuch families might have lived very comfortably. Many of the articles of diet are cheaper than bread, and equally wholefome. Above one half of the expence of living might be faved, by a due felection of the articles of diet.

The English labourer lives chiefly on bread, which being accompanied with other dry, and often falt food, fires his blood, and excites an unquenchable thirst, fo that his perpetual cry is for drink.

But the greateft confumption of bread is occafioned by tea. It is faid that the fubjects of Great Britain confume a greater quantity of that herb, than the whole inhabitants of all the other nations of this quarter of the globe. The loweft woman in England must have her tea, and the children generally share it with her. As tea contains no nourifhment, either for young or old, there must of courfe be bread and butter to eat along with it. The quartern loaf will not go far among a family of hungry children, and if we add the cost of tea, fugar, $\frac{1}{2}$

butter, and milk, the expence of one meal will be more than would be fufficient to fill their bellies with wholefome food three times a-day.

There is reafon to believe that one half the bread confumed in England is ufed to tea, without one hearty meal ever being made of it. The higher ranks ufe tea as a luxury, while the lower orders make a diet of it. I had lately occafion to fee a ftriking inftance of this in a family, that was reprefented to me as in diffrefs for want of bread. I fent them a little money, and was informed that they ran with it directly to the tea-fhop.

To a heavy, fluggifh, phlegmatic man, a moderate ufe of tea may not prove pernicious; but where there is a debilitated ftomach and an irritability of fibre, it never fails to do much hurt. With many it has the effect to prevent fleep.

Tea will induce a total change of conflictution in the people of this country. Indeed, it has gone a great way towards effecting that evil already. A debility, and confequent irritability of fibre, are become fo common, that not only women, but even men, are affected with them. That clafs of difeafes, which, for want of a better name, we call nervous, has made almost a complete conquest of the one fex, and is making hafty ftrides towards vanquifhing the other.

Did women know the train of difeafes induced by debility, and how difagreeable thefe difeafes render them to the other fex, they would fhun tea as the most deadly poifon. No man can love a woman eaten up with vapours, or washed down with difeases arising from relaxation.

It is not tea taken as a beverage after a full meal, or in a crowded affembly, that I fo much condemn, though I think fomething as elegant and lefs pernicious might be fubfituted in its place. The mifchief occafioned by tea arifes chiefly from its being fubfituted for folid food. This is fo much the cafe at prefent, that, had I time to fpare, I think it could not be better employed than in writing against this deftructive drug.

Ss 2

OF BOILED GRAIN.

Though fatinaceous fubftances, of one kind or another, make a neceffary part of the food of man, yet there can be no reafon why fuch fubftances fhould always affume the name and form of bread. Many of them are more wholefome, and not lefs agreeable, in other forms. Bread is often ufed merely to fave the trouble of cookery; and, being portable, is the most convenient article of diet for carrying abroad.

It does not, however, admit of a doubt, that more grain is eaten boiled, though not in this country, than is made into bread; and that this mode of cookery is the most wholefome. Simple boiling precludes all adulteration, and is an operation much less laborious and artificial than baking.

The most general article of diet among mankind is rice. This may be made into a variety of diffies; but fimple boiling is all that is required, to render it a proper fubstitute for bread. It may either be eaten alone, or with milk. In the east, it is used with meat, in the fame manner as we do bread. The people of this country believe that rice proves injurious to the eyes; but this feems to be without foundation, as it has no fuch effect on those who make it the principal part of their food.

Many other kinds of grain will, when boiled, make good fubfitutes for bread. Even those which make a harsh and unpleasant fort of bread, are often rendered very palatable by boiling. This is the case with all the leguminous class of plants, as peas, beans, &c. Even oats and barley are more agreeable, as well as more wholesome, when boiled, than made into bread.

All allow that peas and beans boiled, when young, are a great luxury; but when old, they are equally wholefome, and, when properly cooked, by no means unpleafant. There are few who do not relifh peas-pudding, and even prefer it to bread. Beans are not fo fit for this purpofe; but they make an excellent ingredient in the poor poor man's broth, and whoever eats this broth will find little occasion for bread.

Peas and beans contain an equal quantity of fugar with wheat, oats, or barley, and at the fame time a greater proportion of oil, confequently are more nourifhing. This fact is confirmed by daily experience.

On those farms where peas and beans are raifed in great abundance, the labourers are much fed on that fort of grain; but when removed to farms where they are fed with other kinds of grain, they foon complain of a diminution of strength, and request a supply of peasmeal as formerly.

Nature feems to have pointed out the propriety of the extensive use of peas and beans; it being a fact, that when crops of that kind are duly alternated with crops of wheat, barley, or oats, the fertility of the foil may be maintained, without reft or manure, for many years together: whereas, if the latter be raifed on the fame foil for feveral years fucceffively, they render it barren, fo that, without reft or manure, its fertility cannot be preferved.

The people in England are but little accuftomed to the use of boiled grain, though in many countries it is eaten as a luxury. Boiled barley is a great favourite with the Dutch, and is eaten with milk, butter, or molass. It is the principal food of the Dutch failors, who, in general, are both healthy and robust.

Barley is one of the beft ingredients in foup. Count Rumford fays, it poffeffes the quality of lithing, or thickening foups, in a fuperior degree to any other grain. We have reafon, however, to believe, that grits, or coarfe oatmeal, will anfwer that purpofe ftill better.

Oatmeal is frequently made into bread; but it is a much more wholefome, as well as agreeable food, when made into hafty pudding, and eaten with milk. The peafants in many parts of Britain make two meals a-day of it, while their children almost wholly fubfiss on it; and it is well known that both old and young who are thus fed, are healthy and robust.

The opinion of oatmeal being heating, and occafioning skin difeases, is wholly without foundation. Bread

SS 3

made

made of oatmeal, when not leavened, will fometimes occafion the heart-burn; but this is no proof of its heating quality. Unleavened bread, of wheat, or any other grain, produces the fame effect on a debilitated ftomach. Oatmeal thoroughly boiled feldom gives the heart-burn.

Perfons who are fed on oatmeal bread, or hafty pudding, are not more fubject to difeafes of the fkin, than thole who live on wheat-meal. Cutaneous diforders proceed more from the want of cleanlinefs, than from any particular aliment. The French, fo far from thinking that oatmeal is heating, fpeak of it as poffeffed of a cooling quality; and even the Englifh give oatmeal, or grit gruel, to lying-in women, and fick people of every defeription, which fhews that they are inconfiftent with themfelves, in alleging that the blood is fired by the ufe of oatmeal.

A lieutenant of the army, refiding at a country village with n a few miles of Edinburgh, with a wife and ten children, having no other income than his half-pay, fed the whole of his children with hafty-pudding and butter-milk only, from a conviction that it was the moft wholefome and full diet that fell within the reach of his narrow circumftances. They grew apace; and it was the univerfal remark of the neighbourhood, that they were as fprightly, healthy, and robuft, as other children, and at the fame time perfectly free from all fkin difeafes.

Children are feldom well, unlefs when their bodies are gently open. But this is more likely to be the cafe when fed on oatmeal and milk, than when their bellies are crammed with a ftarchy fubftance made of the fineft flour; yet this in England is the common food of children. I have feen an infant ftuffed four or five times a-day with this kind of food. There needs no conjurer to tell the confequence.

A late Author, a man of learning, but the dupe of prejudice, has, by a ridiculous definition, endeavoured to reprefent oats as proper food for horfes only. I with the horfes in England devoured a fmaller quantity of that grain, and the people more. Few things would have a greater

greater tendency to leffen the expence of living. The oats in North Britain are of a fuperior quality, and I hope the people will long have the fenfe to use them as an article of diet.

Indian corn is likewife faid to make the best food when boiled. Count Rumford observes, that of all things it makes the beft pudding, and that he has made a hearty meal of it, fauce included, for five farthings. What makes good puddings will make good dumplings, and thefe will, at any time, fupply the place of bread. The Count alfo remarks, that the negroes in America perfer Indian corn to rice; and that the Bavarian peafants prefer it to wheat; that it might be imported from North America at about four or five fhillings per bushel; that, when made into flour, it would coft only one penny farthing per pound; and that it is highly nutritious, and the cheapest food known. During the late scarcity, a large quantity of this grain was imported; but fuch is the averfion of the common people of this country to every fort of food to which they are not accustomed, that they refused to purchase it, and the merchants were very great lofers by the importation. On the fame principle the Germans, till within these few years, could not be induced to eat potatoes, though now they are become extremely fond of them.

The American, the Italian, and the German, all cook Indian corn, in the fame way as the North Brton does his oatmeal, by making it into hafty pudding. It may be eaten in a variety of ways. Some eat it with a fauce composed of butter and brown fugar, or butter and molaffes. Others eat it with milk only. In either way it makes a good, cheap, and wholefome diet, by no means difagreeable to those who are accustomed to it.

The only other grain we fhall mention, as best when boiled, is buck-wheat: It is of a very mucilaginous nature, and of courfe highly nutritious. In feveral parts of Europe, it constitutes a principal part of the food of the lower people. In former times it was eaten in Russia, not by the lower classes only: even the nobility made use of it. Boiled, and then buttered, it was such a S s 4 favourite 632

favourite of the great Czar Peter, that he is faid feldom to have fupped on any thing elfe.

OF BUTTER.

It has been faid, that the English have a thousand religions, and but one fauce. It must be allowed that they use butter with almost every kind of food. Butter, though a good article of diet, may be used too freely, and in this country, I am convinced, that is the case. To weak stomaches it is hurtful, even in small quantities, and, when used freely, it proves prejudicial to the strongest.

Butter, like other things of an oily nature, has a conflant tendency to turn rancid. This procefs, by the heat of the ftomach, is greatly accelerated, infomuch that many people, foon after eating butter, complain of its rifing in their ftomachs, in a flate highly difagreeable. Oils of every kind are with difficulty mixed with watery fluids. This is the reafon why butter floats on the ftomach, and rifes in fuch an unpleafant manner.

Perfons afflicted with bile should use butter very sparingly. Some sceptical authors doubt whether or not aliment of any kind has an effect on the bile. One thing, however, is certain, that many patients, afflicted with complaints which were supposed to be occasioned by bile, have been completely cured by a total abstinence from butter.

The most violent bilious complaints that I ever met with, were evidently occasioned by food that became rancid on the flourach, as the cholera morbus, and the like. Nor can fuch complaints be cured, till the rancid matter is totally evacuated by vomiting and purging.

But fuppofing butter did not poffefs the quality of becoming rancid on the ftomach, it may, neverthelefs, prove hurtful to digeftion. Oils of all kinds are of a relaxing quality, and tend to impede the action of digeftion. Hence the cuftom of giving rich broths and fat meats to perfons who have a voracious appetite. The free use of butter, and other oily substances, not only tends to relax the stomach, and impede its action, but to induce a debility of the solids, which paves the way to many maladies. In a country where two-thirds of the inhabitants lead sedentary lives, a debility of fibre most predominate. Whatever increases that debility, ought to be avoided.

Children, without exception, are disposed to difeases arising from relaxation. Butter, of course, ought to be given to them with a sparing hand. But is this the case? By no means. Bread and butter conflitute a great part of the food of children, and I am convinced that the gross humours with which they are frequently troubled, are partly owing to this food. As children abound with moisture, bread alone is, generally speaking, better for them than bread and butter.

I have been aftonished to fee the quantities of butter eaten by großs women who lead fedentary lives. Their tea-bread is generally contrived fo as to fuck up butter like a fpunge. What quantities of crumpets and muffins they will devour in a morning, foaked with this oil; and afterwards complain of indigestion, when they have eaten what would overload the stomach of a ploughman. Dr. Fothergill is of opinion, that butter produces the nervous or sick head-ach, so common among the women of this country. As a proof of this, it is often cured by an emetic.

Oils, in certain quantities, excite nausea, and even vomiting. They must of course prove unfriendly to digestion. A Dutch failor, we are told, can digest train oil. So may an English failor; but it would be very improper food for a London lady.

To fome of the leaner farinaceous fubftances, as the potatoe, and the like, butter makes a very proper addition; but eating it to flefh and fifh, of almost every defcription, is certainly wrong. The flefh eaten in this country is generally fat enough without the addition of butter; and the more oily kinds of fifh, as falmon or herrings, are lighter on the stomach, and easier digested, when eaten without it.

Butter

OF BUTTER.

Butter is rather a groß food, and fitter for the athletic and laborious, than the fedentary and delicate. It is lefs hurtful when eaten fresh than falted. Salt butter certainly tends to induce skin diseases, and I am inclined to think, the free use of it at sea may have some share in bringing on that dreadful malady, so destructive to our brave failors, the sea scurvy.

There is a method of rendering falt butter lefs hurtful, but it feems not to be known in England. What I mean is, to mix it with an equal quantity of honey, and keep it for ufe. In this way it may be given to children with greater freedom. In North Britain, this method of mixing butter with honey is well known; and, from a common proverb, I take the cuftom to be very ancient.

Butter, in itfelf, is not near fo hurtful, as when combined with certain other things. For example: bread made with butter is almost indigestible, and pastries of every kind are little better; yet many people almost live upon pastry, and it is universally given to children. It is little better, however, than poison, and never fails to diforder their stomachs. The fond mother cannot pass a pastry-shop, without treating her darling boy with stome of the dainties, and then wonders how he got the cough, or cholic.

I have known a man feemingly in perfect health, who, by eating a penny-worth of paftry, as he paffed along the ftreet, was feized with fuch an afthmatic fit, that he was obliged to be carried home, and had nearly loft his life. This occurred whenever he inadvertently ate any thing baked with butter.

Every thing that proves very injurious to health, ought, as far as poffible, to be prohibited, by laying a high duty upon it. A duty on paftry would be ferving the public in more refpects than one. It would fave many lives, and leffen fome tax on neceffaries.

Cheefe, as a diet, is likewife injurious to health. It fhould never be eaten but as a defert. It occasions conflipation, fires the blood, and excites a constant craving for drink. It is very improper for the fedentary, and hardly to be digested even by the athletic.

OF BUTTER.

If men will live on dry bread, poor cheefe, falt butter, broiled bacon, and fuch like parching food, they will find their way to the ale-houle, the bane of the lower orders, and the fource of half the beggary in the nation.

OF FRUITS AND ROOTS.

Fruits and roots form a large clafs of the fubflitutes for bread. The latter, being produced under ground, are lefs liable to fuffer from the inclemency of the feafons than grain. Men who wifh to inflame the minds of the multitude may inveigh against the fubflitutes for bread; but reason and found fense fay, the more fubflitutes for bread, the better. When one fails, recourfe can be had to another.

In warm climates the inhabitants have many fubilitutes for bread; and as their feafons are more uniform than ours, they can generally depend on the plant, or whatever it is, proving productive. The plantain tree, commonly called the Indian fig, which has from time immemorial been cultivated in South America, bears fruit of a fweetifh tafte, which will diffolve in the mouth without chewing. It is eaten either raw, fried, or roafted. When intended to fupply the place of bread, it is gathered before it is ripe, and eaten either boiled or roafted. The banana is nearly of the fame nature, but its fruit is greatly fuperior both in tafte and flavour.

The inhabitants of the South Sea, or Ladrone iflands, are fupplied with bread from a tree, which has been lately imported into our Weft India iflands, and will, it is hoped, be found to anfwer the fame purpofe there. It has a flight degree of fweetnefs, but not much flavour. It refembles new bread, and requires to be roafted before it is eaten. Those who have tafted it fay, that it is in no refpect fuperior to the potatoe.

In fome of the West India islands the inhabitants fupply the place of grain by making bread from the root of a shrub, called the cassada, or cassava. Though, to my talte,

tafte, this bread is very infipid, yet the natives are fond of it, to fuch a degree, that I have known fome of them eat it, during their refidence in England, in preference to the fineft London bread.

But the most general fublitutes for bread in the West Indies are the yams. There are three different species of this plant, the roots of which are promiscuously used for bread. They are faid to be very nutritious, of easy digestion, and, when properly dreffed, are by some preferred to the best wheaten bread. The taste is somewhat like the potatoe, but more lussions. The negroes generally eat them boiled, and beaten into a mash. The white people have them ground into flour, and make bread and puddings of them. They can be preferved for feveral feasons, without losing any of their primitive goodness.

Of all the fubflitutes for bread in Europe, the potatoe is the moft extensively useful. This plant is a native of Peru, and has been in Europe about two hundred years. Like moft other important difcoveries, it made but a flow progrefs, and is still far from being fo generally cultivated as it deferves to be. It is indeed known in most parts of Europe, but its culture is best understood in Ireland and the northern parts of England. At Harwich, however, the preference is given to the Dutch potatoes, brought over by the packets between that place and Helvoet Sluys. There is a light fandy foil in Holland, very favourable to the culture of that inestimable root.

As this plant thrives in every foil, and feldom fuffers from the inclemency of feafons, we must blame ourfelves if we fuffer a famine to exist. Indeed, no fuch thing ever can be, where due attention is paid to the culture of potatoes. A far greater quantity of farinaceous food can be raifed on an acre of ground planted with potatoes, than fown with any kind of grain. It is not uncommon to have a return of forty for one. They are not fo hearty a food as corn, but no man will ever perish for hunger why can have potatoes.

Potatoes

Potatoes abound with an infipid juice, which induces fome to think that they are not very nutritious. Facts, however, are against this opinion. Some of the stoutest men we know, are brought up on milk and potatoes. Dr. Peatfon, who has bestowed fome pains in analizing this root, fays, that potatoes and water alone, with common falt, can nourish men completely. They differ in colour and confistence, but not materially with regard to their nutritive qualities.

Some think the firm kind are the most nutritious; but the Irish, who must be good judges, give the preference to the mealy. The difference, however, depends much on the mode of cooking them.

More than half the fubftance of potatoes confifts of water, and experience fhews, that the mode of cooking, which most diminishes their moisture, is to be preferred. In London, they are drenched in water and washed before they are brought to market, which accounts in a great measure, for the bad quality of the London potatoes.

They are dreffed in a variety of ways, but fimple boiling or roafting feems to be all the cooking they require, to render them a proper fubfitute for bread. Some are fond of making bread of them. This, in my opinion, is maring both. Why manufacture any thing into bread, which requires only the aid of fire to make it fuch? Nobody thinks of making dough of the bread fruit; but the potatoe might with as great propriety be called the bread root, as it is made into bread by the fame procefs.

Stewed mutton and potatoes make not only a nourifhing but a very palatable difh. The excefs of fat of the mutton which, when otherwife cooked, fuftains great lofs, is thus preferved, by being abforbed by the potatoes. It is, however, to be obferved, that, when potatoes are ufed in broth or flews, they ought previoufly to be boiled, and the water thrown away, as it contains fomething deletereous. Simple boiling or roafting is fufficient to prepare potatoes to fupply the place of bread, but when they are intended to ferve as a meal, they require fomething of a foftening nature, as milk, butter, or both. What a treafure

treafure is a milch cow and a potatoe-garden, to a poor man with a large family, who lives in the country! Yet, with a little attention from landlords and farmers, almost every man might be fo accommodated. What a fource of real wealth and population! Men would multiply, and poverty, unlefs among the profligate, be unknown. Horfes are fometimes fed with potatoes, and become very fond of them. With the addition of a fmall quantity of hay, they are found to be fufficiently nourifhing.

I would beg leave to recommend, both to landlords and farmers, a careful perufal of Earl Winchelfea's excellent letter to Sir John Sinclair, on the advantages of cottagers renting lands. This humane Nobleman takes up the matter in a truly patriotic light, and fhews that farmers, inftead of leffening the number of poor, do every thing they can to multiply them; and I am forry to fay that, fo far as my obfervation goes, it agrees entirely with his lordfhip's.

Another letter relating to this fubject has lately fallen into my hands, a copy of which I shall take the liberty to infert, as it contains in a few paragraphs the best practical illustration of the truths I have long been endeavouring to enforce. It was written by Sir JOHN METHUEN POORE to Sir WILLIAM PULTENEY; and is dated "Rushall, 4th April, 1801."

" Sir,

" I can prove, not by theory, but practice, the benefit of planting potatoes on fallows. In the parifu where I refide, the whole of which, except five acres, is my property, there are thirty cottages, containing one hundred and thirty-one poor people. I have, for five or fix years paft, allotted, free from rent, four acres of land, intended to be fown with wheat the following autumn, for the cottagers to plant with potatoes, by which means each raifes from ten to fifteen facks, equal to two hundred and forty pounds per fack, yearly, in proportion to the number of their children : each has not only fufficient for his family, but is enabled alfo to fat a pig. They declare, were I to give among them a hundred pounds, it would not be of fo much benefit to them; and it is not one

one shilling out of my pocket, for I have as good, if not a better crop of wheat from this land, as I have from the other part of the field.

" The method I take is this: the latter end of November I plough the land; the frost during the winter mellows it : the beginning of March following, I plough it again, and harrow it; at both which times I have little to do with my horfes: I then divide it into lots; a man with a large family has a larger lot than a fingle perfon, or one who has only two or three children, allowing about five perches (of $16\frac{1}{2}$ feet fquare) to each in a family: they then plant it, and put over their potatoes what manure they have collected the year preceding, (for every cottager has more manure than neceffary for this, from their fires, and a variety of other things,) and during the fummer, after their day's labour is done, they and their wives hoe them; and as every man works more cheerfully for himfelf than for another, they do not fuffer a weed to grow. In October they dig them up; and it is the most pleafant thing imaginable to fee the men, their wives and children, gathering the produce of their little farms, which is to ferve them the enfuing winter .--Were this plan generally adopted, the labourers would confume but little corn ; which would fupply the manufacturing towns, and we fhould have no occasion to import. As four acres are fufficient for thirty families, it would take but a fmall quantity of land from every farm in the kingdom. The way practifed here is to plant the potatoes in furrows, eighteen inches apart, and a foot apart in the rows. The land about me is of different qualities; on the hills rather light; in the vale, near the parish, inclining to clay; but all fit for turnips: the potatoes are planted in the low land, being nearer home. The poor at prefent will not live entirely without bread, as many do in Ireland, though potatoes daily get into use more and more; and I am perfuaded, were my plan generally adopted, in two or three years the labourers in the country would confume but little or no corn. Thirty years ago, the poor in this part of the country would not eat potatoes, if they could get other roots or vegetables."

It

640

It is unneceffary to make any comments on this letter, every line of which is dictated by good fenfe and humanity, as well as by liberal and enlightened policy. I shall only add an earnest with, that the example of the truly patriotic and benevolent writer may be followed by every man of landed property, and by every confiderable farmer in the kingdom !

Some think that the potatoe, unlefs it be made into bread, will not keep. An accident taught me the contrary. Many years ago a friend of mine fent me a potatoe, after it had been roafted in an oven, on account of its fingular figure. I laid it on a shelf among some other things of the like kind, and was furprifed, on removing them many years after, to find the potatoe quite fresh, though as dry as a bone. On grating it down, it was perfectly fweet; and as fit for making foup, as the day it was roafted. I apprehend that nothing made into bread would have kept fo long.

Posterity will hardly believe that a fcarcity of bread could be felt in Britain, at a time when it was known that a fufficient quantity of farinaceous food could be raifed in one county for the inhabitants of the whole ifland. Let proper encouragement be given to the culture of potatoes, and let famine at defiance.

Many other domeflic roots, fprouts, &c. are very wholefome, and may occafionally fupply the place of bread. Of these Mr. Bryant of Norwich reckons above forty ; but we shall only take notice, by way of specimen, of the most useful and productive. It is worthy of remark, that no nation can be very populous, which does not draw a great part of its food from under ground.

The Jerufalem artichoke is a native of Brazil, but, having been long cultivated in this country, it is too well known to need any defcription. From its tafte, which is like that of artichoke bottoms, it would feem to be nutritious, and is far from being unpleasant to the palate. Some reckon it windy, but this may be corrected in the cooking, by warm spices; and as the plant is very productive, we would recommend it to be used in the fame manner as potatoes, and the other farinaceous roots. Of Of the efculent roots in this country, the parfnip is reckoned the moft nourifhing. It is likewife of eafy digeftion, and is agreeable to moft palates. Some indeed diflike it on account of its fweetnefs; but that is a proof of its nutritive quality, fugar being the moft nourifhing thing in nature. We are told that, in the north of Ireland, the poor people make beer from this root.

There is not any plant that affords a more ftriking proof of the benefits of culture than the turnip. In its wild state it is good for little or nothing ; but, when properly cultivated, it not only affords wholefome nourifhment for man, but furnishes the principal winter food for cattle. There is a fpecies of this plant which grows in North Britain, called the yellow turnip, which is fweet, and of a fuperior quality to those produced in the fouth, particularly about London, which are bitter and ftringy. The yellow turnip is the most nourishing, and also the most hardy in fustaining the winter. It is eaten with milk to cure the confumption and fcurvy. Margraaf fays, he could extract no fugar from the turnip, which affords ground to conclude, that it is not fo nutritive as certain other roots. Not only the root of the turnip, but the tops, when young, make very pleafant greens. The fprouts, if gathered when very tender, make an excellent fallad.

The carrot, like the turnip, is good for little in its natural ftate, being fmall, tough, and ftringy. Manured, it grows large, fucculent, and of a pleafant flavour. It ought, however, to be eaten young, otherwife it lies on the ftomach, and is hard of digeftion. It is an ingredient in feveral foups, and, being folid, may in fome meafure fupply the place of bread.

Salfafy, skirrets, and the several kinds of beets, are all pleasant and nourishing. They are likewise of easy digestion, and may be dressed in a variety of ways. Margraaf has by experiments discovered, that both skirrets and beets contain a confiderable quantity of sugar. Though the extracting a faccharine salt from these plants may be no object while we posses the West India islands, yet it ferves to shew that they posses a quantity of nutri-

Tt

tious

tious matter, fufficient to give them a rank among the articles calculated to fupply the place of bread.

The onion, we are told, was a great favourite in Egypt four thousand years ago, and Dr. Heffelquest fays, it is not to be wondered at, for whoever has tasted the onions of Egypt, must allow that none can be better in any part of the globe. There, he fays, they are fweet, though in many countries they are strong and nauseous. There they are fost, whereas in northern countries they are hard, and their coats so compact, that they are difficult to digest. This very quality may, however, recommend them in countries where food is fcarce. The Doctor observes, that the Turks eat them roasted with their meat as we do bread, and are so fond of them that they wish to be indulged with this dist in Paradise.

From the Doctor's account one would be induced to believe that the onion ufed in Egypt was of a different fpecies from ours; but I am rather inclined to think it may depend on the mode of culture, as well as on the warmth of the climate and the difference of foil, as we find in the fouthern parts of Europe they are milder than in the more northerly. In Spain they are very mild, and a root weighing two pounds will grow from a fingle feed.

Onions are dreffed in a variety of ways, but, in regard to wholefomenefs, there is no method better than fimple boiling. By this method of cooking, they are rendered mild, of eafy digeftion, and go off without leaving any difagreeable heat on the ftomach or bowels. Many fhun them on account of the ftrong difagreeable fmell they communicate to the breath. Mr. Bryant fays, this may be remedied, by eating a few raw parfley leaves immediately after, which will effectually overcome the fcent of the onions, and likewife caufe them to fit more eafy on the ftomach.

The leek is generally reckoned among pot-herbs; but as the root is the part chiefly ufed, the confideration of it comes under the prefent head of difcuffion. Indeed, it is as properly a root as the onion, which growschiefly above ground. The leek, as well as the onion, is faid to be a conftant difh at the tables of the Egyptians, who chop them fmall, and eat them with their meat. The leek is used as a pot-herb in most parts of Britain, especially in Wales, where the natives are faid to be fond of it. In Scotland a full grown fowl and small piece of falt beef, stewed with a large quantity of leeks, is a very favourite dist. In my opinion the leek is not so generally used any where as it deferves to be. There is no ingredient goes into soup that is more wholesome, or that gives it a better flavour, than leeks. They are in many respects medicinal, and, to my taste, as an ingredient in soups, they are greatly superior to the onion or any other pot-herb whatever.

It is a fact worthy of obfervation, that the boiling of vegetable fubftances thoroughly, a thing feldom done in England, extricates a confiderable quantity of air, and makes them lefs liable to produce flatulency.

I could mention a great many more efculent plants which might occafionally fupply the place of bread, but the above fpecimen is fufficient to fhew how liberal nature is in fupplying man with food, provided he will take the trouble of cultivating and cooking it. Mr. Bryant, in his hiftory of efculent plants, enumerates above four hundred and fifty, each of which affords a wholefome nourifhment, and may occafionally be ufed in place of bread.

OF BROTHS AND SOUPS.

Thefe may likewife be confidered as fubfitutes for bread. If properly made, they will ferve both for bread and drink. Though broth is a difh of the greateft antiquity, and may be confidered as extremely delicious, yet it is not a favourite in this country. Here the people are fond of what they call folids; yet thofe very folids they make into broth, by fwallowing as much drink after them as they can get. The only difference is, the foreigner makes his broth in a pot, and the Englifhman makes his in the ftomach.

A very fenfible anonymous writer obferves, that in England a pound of meat makes fimply a pound of food; whereas in any other country in Europe, that quantity of animal food, when flewed down with vegetables and Scotch barley, will produce an ample meal for half a Tt 2 dozen

dozen people. Hence he juftly infers that, among the variety of schemes which may have been devised by the humane for relieving the diffreffes of the poor, a better and more extensive charity cannot be devised than that of inftructing them in a new mode of cookery.

The fame author adds, that the refult of his experiments on this fubject had exceeded his most fanguine expectations, and that each day gave him fresh proofs of the excellency of his plan for teaching the poor and needy to find themfelves in a wholefome and palatable diet, at the cheapest rate, in which little or no bread was required. He concludes by afferting, that there is fcarcely a place in this kingdom, where twenty perfons may not have a wholefome, hearty, and palatable meal, for three fhillings.

This anonymous letter is followed by one from Colonel Poynter, two from Dr. Johnston, of the royal hofpital at Haflar, addreffed to Admiral Waldegrave, and one from the Admiral himfelf, written for infertion in a public paper. They contain a variety of receipts for making cheap, wholefome, and nourifhing difhes for the poor. These diffies confist chiefly of broths, soups, and ftews, or what they call pottage, and are calculated to make a hearty and plentiful meal without bread or drink.

I am inclined to pay the more attention to thefe letters, as they feem all to have been written by gentlemen of observation. The pamphlet is fold by Longman and Debrett, for a charitable purpofe, at the fmall price of three-pence.

The writer who has paid most attention to the improvement of cookery, for the benefit of the poor, is Count Rumford. In his economical and philosophical effays, he has given fuch a variety of forms for making wholefome, cheap, and nourifhing foups, ftews, and other dishes for common use, that little more feems neceffary to be faid on the fubject. I shall only observe, that the mode of living on broths, foups, hafty-pudding, and fuch like, fo warmly and juftly recommended by the Count, has been practifed in the northern parts of this kingdom from time immemorial. There the food of the

71

the common people is hafty-pudding, with milk, for breakfast and supper, and broth, with vegetables and meat, for dinner. The poorer fort often make broth without meat; but they all use vegetables in great abundance, and fometimes they fupply the place of meat with butter. As the hafty-pudding and milk make a complete meal, no bread is neceffary either at fupper or breakfast; nor is much required at dinner, as the broth is made thick with barley, cabbage, and a variety of other vegetables or pot-herbs. Cabbage is a favourite ingredient in a Scotchman's broth. It is feldom made without this article, which is not eaten fo early as in England. It is there fuffered to grow to maturity, and, when that is the cafe, there is no plant more productive. This the Germans know well, and make it into four crout, one of the best antidotes against the fcurvy with which we are acquainted.

This kind of diet not only faves bread but drink. The labourer who lives on hafty-pudding and foups, feldom has occafion for drink; while he who is burnt up with dry bread and cheefe, or falt meat broiled, has a continual thirft, and fpends the greater part of his earnings in liquor. This, by acting as a powerful ftimulus, may make him do more work for fome time, but it generally cuts him off in the middle of his days. The Englifh labourer, who works hard and drinks hard, feldom lives long, and is an old man when he fhould be in his prime.

The roafting of meat is a wafteful mode of cookery, which ought to be avoided by the poorer fort of people, as much of the fubftance, and the most nutritive parts, are lost by fcorching, and what flies off by evaporation.

I know it will be faid, that I recommend flops in place of folid food. They are fuch flops, however, as the greatest heroes of antiquity lived upon; and though I have visited most parts of the island, I know of no better men than those who live in the manner described above, nor are the people any where more healthy, or longer lived.

Tt3

Broth

Broth is not only a difh of great antiquity, but one that can be made in a great variety of ways. It receives into its composition animal and vegetable fubftances of every kind that are used in diet, and it may be feasoned fo as to fuit every palate. Indeed, people early accuftomed to eat broths properly made, are generally fond of them for their whole lives.

It would be difficult to affign a reafon why the inhabitants of South Britain fhould diflike a difh fo much relifhed by other nations. Cuftom, no doubt, fettles all thefe things; but how cuftoms arife, is not fo clear a matter. If an alteration in diet is to be introduced with effect, it must begin with children. Whatever men are accustomed to eat when young, they generally prefer for the rest of their lives. Were the children in South Britain taught to eat hasty-pudding, with milk, for breakfast and supper, and broth, with vegetables and meat boiled in it, for dinner, they would reliss the diffues as long as they lived, would find little occasion for bread, and still less for drink; and would thrive better than on their prefent food.

What parents love themfelves, they generally give to their children, without any regard to its being proper for them or not. I have feen a father, who was fond of ftrong beer, make his fon, an infant, guzzle it at every meal; and the mother who delights in tea, does not fail to give it to her daughter whenever fhe takes it herfelf. By this conduct, the fon becomes a tippler, and the daughter fips tea in place of folid food, until fhe is eaten up with vapours and other nervous diforders.

Count Rumford fays, brown foup is the common breakfast of the Bavarian peasants, to which they occafionally add bread. This he avers is infinitely preferable in all respects to that pernicious wash, tea, with which the lower classes of the inhabitants of this island drench their stomachs, and ruin their constitutions. He adds, that a simple infusion of this drug, drank boiling hot, as the poor generally drink it, is certainly a poison, which, though it be sometimes slow in its operation, never fails

¹³

to produce fatal effects, even in the ftrongest constitution, where the free use of it is continued for a confiderable length of time.

The German on his *polenta*, the American on his *mufb*, and the North Britain on his *hafty-pudding*, can make a hearty breakfaft for a tenth part of what a tea-breakfaft would coft, while it is infinitely more wholefome. It has likewife the advantage that no bread is neceffary.

I have been often told, when recommending foups to the poor, that they had not time to make them, and that they could not afford fuel on account of its price, as it is dear in great towns. They can, however, find fuel twice a-day to boil a tea-kettle, and time to make the tea, which is a more tedious operation, by far, than making a mefs of hafty-pudding. For a great part of the year even the pooreft perfon must have a little fire; and it would require no more to make a comfortable mefs of foup, which is always beft when made with a flow fire.

The mode of living that I would recommend to the lower orders of the people, with a view to fave expence and improve their health, is to fubfitute occafionally other farinaceous fubftances in the place of bread, as potatoes, &c. to give up in a great measure the ufe of roafted, baked, and broiled meats, and to fupply their place with broths, foups, flews, and fuch like, made with a little meat and plenty of vegetables; to give to children, and to grown people who will eat it, for breakfaft, milk-porridge, or hafty-pudding with milk, fmall beer, or melaffes. This will be found a more wholefome breakfaft than tea, while it is much cheaper and requires no bread*.

CON-

647

• The celebrated Dr. Huffland, in his Art of prolonging Life, fays, the moderate use of soups is certainly not hurtful; and it is fingular that people should imagine it tends too much to relax the ftomach. Does not all our drink, even though cold, become in a few minutes a kind of warm soup in the stomach; and does not the stomach retain the same temperature during the whole day? Be careful only not to use it hot, in too great quantity at one time,

Tt4

CONCLUSION.

Although the place of bread may be occafionally fupplied by farinaceous roots and other vegetables, yet we would by no means wifh to difcourage the culture of grain. The culture of grain is the culture of men. While the hufbandman is raifing food for his fellowcreatures, he is laying the foundation of health and longevity to himfelf and his offspring. Innumerable benefits are connected with the culture of grain. While the artificer is fitting in fome aukward pofture, breathing confined, and perhaps contaminated air, the cultivator of the foil rifes with the fun, eats his wholefome meal of milk and farinaceous food, hies him to the field, where he fpends the day in ufeful labour, inhales the frefh breezes, and at eve returns home with a keen appetite, to enjoy his fimple repaft and found repofe.

It has been faid, as artificers can earn more money than thole who cultivate the ground, that arts ought to be encouraged, and grain, if neceffary, imported. No manufacture is equal to the manufacture of grain. It fupplies food for man and beaft, while the furplus, by being exported, enriches the nation. Nor is it fubject to the un-

or too watery. It is attended even with great advantages. It fupplies the place of drink, particularly to men of letters, women, and all those who do not drink, or drink very little except at table, and who, when they give over foup, receive into their blood too little moisture. And it is here to be remarked, that fluids ufed in the form of foups unite much better and fooner with our juices than when drunk cold and raw. On this account foup is a great preventive of dryneis and rigidity in the body, and therefore the best nourishment for old people, and those who are of an arid temperament. It even supplies the place of medicine. After catching cold, in nervous head-aches, cholics, and different kinds of cramp in the ftomach, warm foup is of excellent fervice. It may ferve as a proof of the utility, or at leaft harmleffnefs of foup, when I remark that our forefathers, who certainly had more ftrength than we have, used foup; and that it is used by ruftics, who are flill ftronger than those in refined life; and that all the old people with whom I ever was acquainted were great friends to it.

certainty

certainty of other manufactures. They often depend on fashion and caprice, but the necessaries of life will always find their value fomewhere. Though I am convinced that fome regulations are wanting for the encouragement of agriculture, I do not confider it as my province to dictate to the wisdom of the legislature. They know their duty, and I have reason to believe that they are inclined to pay it all due attention.

I will venture, however, to affert, that if proper encouragement were given to agriculture, Britain would at all times not only have a fufficiency of grain for her own confumption, but a furplus for exportation. This would contribute more to her real wealth, the happinefs of her people, and the ftability of her government, than either the increase of her trade, the flourishing of her manufactures, or the extension of her territory.

It is matter of real regret and wonder that Britain, at a time when agriculture is cultivated as a fcience, fhould not be able to raife grain for the fupply of her own inhabitants, but become every year more dependent on foreign ftates for even the neceffaries of life. Until an adequate remedy can be found for this growing evil, the free use of the various substitutes for bread cannot fail to alleviate the calamities of the poor, and to reduce the price of labour.

The great confumption of animal food, and the immenfe number of horses kept in this country, are to be reckoned among the caufes of the fcarcity of grain. Mr. Mackie computes the number of horfes in this country to be about two millions, and that every horfe, on an average, confumes the produce of three fertile acres, confequently the produce of fix millions of fertile acres; is annually confumed by horfes. These would produce a quantity of grain more than fufficient to maintain half the inhabitants of Great Britain. Two hundred and fixty thousand of these animals are kept for pleasure. I shall be told that they contribute to health. That I deny. Did our ladies of fashion and fine gentlemen make use of their limbs, instead of being dragged about in carriages, they would both benefit themfelves, and the public.

CONCLUSION.

lic. I shall conclude these remarks with the advice of the humane and benevolent Thomson :

"Ye gen'rous Britons! venerate the plough,

- " And o'er your hills and long withdrawing vales
- " Let Autumn fpread her treasures to the fun
- " Luxuriant and unbounded. As the fea
- " Far through his azure turbulent domain
- " Your empire owns, and from a thousand shores
- " Wafts all the pomp of life into your ports;
- " So with fuperior boon may your rich foil
- " Exub'rant Nature's better bleffings pour
- "O'er ev'ry land, the naked nations clothe,
- " And be the exhauftlefs gran'ry of a world."

APPENDIX:


CONTAINING

A Lift of Simples and of fuch Medicinal Preparations as ought to be kept in Readiness for private Practice :

The Method of preparing and compounding fuch Medicines as are recommended in the former Part of the Book, with the Addition of feveral others of a fimilar Nature :

Remarks on the Dofes, Uses, and Manner of applying the different Preparations.

Medicamentorum varietas ignerantiæ filia est. BACON.



APPENDIX.

being ad as , doub as adapta ge

IGNORANCE and fuperflition have attributed extraordinary medical virtues to almost every production of nature. That fuch virtues were often imaginary, time and experience have fufficiently shewn. Physicians, however, from a veneration for antiquity, still retain in their lists of medicine many things which owe their reputation entirely to the superflition and credulity of our ancestors.

The inftruments of medicine will always be multiplied, in proportion to men's ignorance of the nature and caufe of difeafes: when these are fufficiently understood, the method of cure will be fimple and obvious.

Ignorance of the real nature and permanent properties of those fubftances employed in the cure of difeases, is another reason why they have been to greatly multiplied. Physicians thought they could effect by a number of ingredients, what could not be done by any one of them. Hence arose those amazing farragos which have to long difgraced the medical art, and which were effeemed powerful in proportion to the number of fimples that entered their composition.

The great variety of forms into which almost every article of medicine has been manufactured, affords another proof of the imperfection of the medical art. A drug which is perhaps most efficacious in the fimplest form in which it can be administered, has been nevertheless ferved up in so many different shapes, that one would be induced to think the whole art of physic lay in exhibiting medicine under as many different modes as possible.

Different forms of medicine, no douot, have their ufe; but they ought never to be wantonly increased. They are by no means to neceffary as is generally imagined. A few grains of powdered rhubarb, jalap, or ipecacuanha, will actually perform all that can be done by the different preparations of these roots, and may also be exhibited in as fase and agreeable a manner. The same observation holds with regard to the Peruvian bark, and many other simples of which the preparations are very numerous.

Multiplying the ingredients of a medicine, not only renders it more expendive, but also less certain, both in its dose and operation.

ote fais, emerciants, no agreethie, ne che w

APPENDIX.

ration. Nor is this all. The compound, when kept, is apt to fpoil, or acquire qualities of a different nature. When a medicine is rendered more fafe, efficacious, or agreeable, by the addition of another, they ought, no doubt, to be joined; in all other cafes, they are better kept afunder. The combination of medicines embarraffes the phyfician, and retards the progress of medical knowledge. It is impossible to afcertain the precise effect of any one medicine, as long as it is combined with others, either of a fimilar or diffimilar nature.

In the exhibition of medicine, regard fhould not only be had to fimplicity, but likewife to elegance. Patients feldom reap much benefit from things that are highly difagreeable to their fenfes. To tafte or fmell like a drug, is become a proverb; and to fay truth, there is too much ground for it. Indeed no art can take away the difagreeable tafte or flavour of fome drugs, without entirely deftroying their efficacy; it is poffible, however, to render many medicines lefs difguffful, and others even agreeable; an object highly deferving the attention of all who adminifter medicine.

The defign of the following pages is, to exhibit fuch a lift of drugs and medicines as may be neceffary for private practice. They are confiderably more numerous indeed than those recommended in the former part of the Book, but are ftill greatly within the number contained in the most reformed dispensatories. The same medicine is feldom exhibited under different forms; and where different medicines answer nearly the same intention, there is commonly no more than one of them retained. Multiplying forms of medicine for the same intention tends rather to be wilder than affist the young practitioner, and the experienced plysician can never be at at a loss to vary his prescriptions as occasion requires.

The chemical and other difficult preparations are for the most part omitted. All of them that are used by any private practitioner are not worth preparing. He will buy them much cheaper than he can make them. Great care, however, is neceffary to obtain them genuine. They are often adulterated, and ought never to be purchased unless from persons of known veracity. Such of them as are in common use are inferted in the list of drugs and medicines. Their proper doses and manner of application are mentioned in the practical part of the book, whereever they are prescribed.

Such articles of medicine as are to be found in the house or garden of almost every peasant, as barley, eggs, onions, &c. are likewise for the most part omitted. It is needless to swell a list of medicines with such things as can be obtained whenever they are wanted, and which spoil by being kept.

654

The

The preparations made and fold by diffillers and confectioners are also generally left out. These people, by operating upon a larger plan, generally make things better, while it is in their power to afford them much cheaper than they can be prepared by any private hand.

The quantity ordered of every medicine is as fmall as could well be prepared, both to prevent unneceffary expence, and that the medicine might not (poil by keeping. Almost every medicine fuffers by being kept, and should be used as soon after it has been prepared as possible. Even simple drugs are apt to spoil, and should therefore be laid in in small quantities; they either rot, are confumed by infects, or evaporate so as to lose their peculiar tafte or flavour, and often become quite infignificant.

In the preparation of medicines, I have generally followed the most improved dispensatories; but have taken the liberty to differ from them wherever my own observations, or those of other practical writers on whose judgment I could depend, suggested an improvement.

In feveral compositions, the ingredient on which the efficacy of the medicine principally depends is increased, while the auxiliaries, which are generally ordered in such trifling quantities as to be of no importance, are lest out, or only such of them retained as are necessary to give the medicine a proper confistence, or the like.

The colouring ingredients are likewife for the moft part omitted. They increase the bulk and price of the medicine; without adding any thing to its value. It would be well if they were never used at all. Medicines are often adulterated for the fake of a colour. Acrid and even poisonous substances are, for this purpose, sometimes introduced into those medicines which ought to be most bland and emollient. Ointment of elder, for example, is often mixed with verdigrife to give it a fine green colour, which entirely frustrates the intention of that mild ointment. Those who wish to obtain genuine medicines should pay no regard to their colour.

Some regard is likewise paid to expence. Such ingredients as greatly increase the price of any composition, without adding confiderably to its virtue, are generally either omitted, or somewhat less expensive substituted in their place. Medicines are by no means powerful in proportion to their price. The cheapest are often the best; besides, they are the least apt to be adulterated, and are always most readily obtained.

With regard to the method of compounding medicines, I have generally followed that which feemed to be the most fimple and natural, mentioning the different steps of the process in the fame order in which they ought to be taken, without paying an implicit regard to the method of other difpenfatories.

For many of the remarks concerning the preparation, &c. of medicines, I have been obliged to the author of the New Difpenfatory. The other observations are either such as have occurred to myself in practice, or have been suggested in the course of reading, by authors whose names I am not able diffinctly to recollect.

I have followed the alphabetical order, both with regard to the fimples and preparations. A more fcientific method would have been agreeable to fome perfons, but lefs ufeful to the generality of readers. The different claffes of medicine have no great dependence upon one another, and, where they have, it is hard to fay which fhould ftand firft or laft; no doubt the fimple preparations ought to precede the more compound. But all the advantages arifing from this method of arrangement do not appear equal to that fingle one, of being able, on the firft opening of the book, to find out any article, which, by the alphabetical order, is rendered quite eafy.

The dole of every medicine is mentioned whenever it appeared neceflary. When this is omitted, it is to be underftood that the medicine may be uled at difcretion. The dole mentioned is always for an adult, unlels when the contrary is expressed. It is not an easy matter to proportion the doles of medicine exactly to the different ages, conflictutions, &c. of patients; but, happily for mankind, mathematical exactness here is by no means neceffary.

Several attempts have been made to afcertain the proportional dofes for the different ages and conffitutions of patients; but, after all that can be faid upon this fubject, a great deal must be left to the judgment and fkill of the perfon who administers the medicine. The following general proportions may be observed; but they are by no means intended for exact rules. A patient between twenty and fourteen may take two thirds of the dose ordered for an adult; from fourteen to nine, one half; from nine to fix, one third; from fix to four, one-fourth; from four to two, one-fixth; from two to one, a tenth; and below one, a twelfth.

Difpenfatories are ufually written in the Latin language. Even authors who write in Englifh, generally give their prefcriptions in Latin; and fome of them fhew fo great an attachment to that language, as first to write their recipes in it, and afterwards translate them; while others, to compromife the matter, write the one half in Latin and the other in English. What peculiar charm a medical prefcription when written in Latin may have, I shall not pretend to fay; but have ventured to make ufe

APPENDIX.

use of the plainest English I could, and hope my prescriptions will succeed no worse for it.

N. B. The Apothecary's weights, and the English wine measures, are used throughout the whole book, the different denominations of which will appear from the following Table: A pound contains twelve ounces.

n pound ci	onta	18 I WEIVE OI	uces.
An ounce			
A drachm	-	- three for	uples.
A scruple			
A gallon c			
A pint -			
An ounce	-	- eight dra	chms.
fpoonful is th			
The second			

Com, alocs

siders ----

munepung

Firefliors, calcined

minasausa -----

Lead, Lichards

Magnetia alba

Mercury, crude

Authorite mineral

vhite dino

- corronve fublimate

Harbs, lefter centuary

A

A LIST

walla's oranges

sohin -----

vate Frad

ce.

Ballam of Capivi

damaganio -----

Mezenon

Earthy Puller's

.[658]

ufe of the plaine? I agtif Leonid, and hope my preferiptions will

M. B. The Apothecarp's weights, and the English wine

lucceed no worid for n

A LIST of SIMPLES, and of fuch MEDICINAL PRE-PARATIONS, as ought to be kept in readiness for private Practice. miosib idgin - - sonuo nA

A drachan - - three foruples.

GARIC Antimony, crude _____ Jamaica pepper ______ juniper berries ______ nutmegs fulphur of list to stuland od tamarinds Balfam of Capivi ----- of Peru ----- of Tola Batk, cafcarilla ---- cinnamon ---- Mezerion ---- Peruvian ----- Winter's, or canella alba Borax Calamine ftone, levigated Caftor, Ruffian Cauftic, common ---- lunar Earth, Fuller's ----- Japan Armenian bole ----- French ditto Extracts of gentian ----- of guaiacum ----- of hellebore, black ----- of hemlock of jalap ----- of liquorice ----- of Peruvian bark ----- of poppies ----- of wormwood Flowers of camomile ----- colt's foot ----- elder ----- rofemary ----- damaik rofes ----- red ditto Fruits, almonds bitter apple caffia fiftularis ____ Curaffao oranges figs, dried

657

twenty grains. staig man Fruits, French prunes Gum, aloes ----- ammoniac, in tears, ---- arabic ----- alafœtida ----- camphor galbanum ----- gamboge ----- guaiacum ----- kino ----- myrrh ----- opium Hartshorn, calcined ----- fhavings of Herbs, lesser centaury ----- peppermint ----- fpearmint _____ penny-royal ----- trefoil ----- uva urfi ----- wormwood Lead, Litharge ---- white --- fugar of Lemon-peel Mace Magnefia alba Manna Mercury, crude ----- calcinated _____ Æthiops mineral _____ calomel _____ corrofive fublimate ----- red precipitate ----- white ditto Mufk Oil, effential, of amber

LIST OF SIMPLES, &c.

Oil, effential, of anife of cinnamon of juniper of lemon-peel _____ of peppermint ---- expressed, of almonds ----- of linfeed Oil of Olives, or Florence oil of palms ---- of turpentine Orange-peel Oyster-shells prepared Poppy-heads Refins benzoin ------ flowers of ----- Burgundy pitch ----- dragon's blood ----- frankincenfe liquid ftorax white, or rofin [cammony Roots, birthwort calamus aromaticus ____ contrayerva garlic gentian gentian ginger. hellebore, black, white _____ jalap ----- ipecacuanha ----- lily, white ----- liquorice ----- marfhmallow mezerion ----- rhubarb ----- farfaparilla ----- funeka ----- fquills ----- termentil turmeric ----- Virginian Inake wild valerian _____ zedoary Saffron Sal ammoniac, crude ----- volatile Salt, Epfom ---- of Glauber ---- of hartfhorn ---- nitre, purified, or prunel ---- Polychreft

Salt, Rochel ----- of tartar Seeds, anife ----- carrawsy ----- cardamom ----- coriander cummin ----- muftard ----- fweet fennel wild carrot Senna Dilaup Similar Spanish flies Spermaceti Spirits, æthereal, or æther ----- of hartfhorn ----- of lavender compound of nitre ----- ditto dulcified ----- of fal ammoniac ----- of fea falt ----- of vinegar - of vitriol ----- of wine rectified ----- volatile aromatic Steel, filings of, --- ruft of, prepared ---- foluble falt of Salphur vivum ----- balfam of ----- flowers of Tar ---- Barbadoes Tartar, cream of ----- emetic ----- foluble ---- vitriolated Tin prepared Tutty levigated Turpentine, Venice Verdegrife Vitriol, green ---- blue white Wax, white ----- yellow Woods, guaiacum ---- logwood ----- faffafras faunders red Zinc, flowers of

Uu2

MEDI-

MEDICINAL PREPARATIONS.

Salt, Mochel

5K3 TR5 10 -

BALSAMS.

THE subject of this section is not the natural balsams, but certain compositions, which, from their being supposed to posses balsamic qualities, generally go by that name.

This class of medicines was formerly very numerous, and held in great efteem, Modern practice, however, has justly reduced it to a very narrow compass.

Anodyne Balfam.

Take of white Spanish soap, one ounce; opium, unprepared, two drachms; rectified spirit of wine, nine ounces. Digest them together in a gentle heat for three days; then strain off the liquor, and add to it three drachms of camphor.

This balfam, as its title expresses, is intended to ease pain. It is of fervice in violent ftrains and rheumatic complaints, when not attended with inflammation. It must be rubbed with a warm hand on the part affected; or a linen rag moistened with it may be applied to the part, and renewed every third or fourth hour, till the pain abates. If the opium is lest out, this will be the Saponacious Balfam.

Locatelli's Balfam.

Take of olive oil, one pint; Strafburg turpentine and yellow wax, of each half a pound; red faunders, fix drachms. Melt the wax, with fome part of the oil over a gentle fire; then adding the remaining part of the oil and the turpentine; afterwards mix in the faunders, previoufly reduced to a powder, and keep them flirring together till the balfam is cold.

This ballam is recommended in erofions of the inteffines, the dyfentery, hæmorrhages, internal bruifes, and in fome complaints of the breaft. Outwardly it is ufed for healing and cleanfing wounds and ulcers. The dofe, when taken internally, is from two fcruples to two drachms.

The vulnerary Balfam.

Take of benzoin, powdered, three onnces; balfam of Peru, two ounces; hepatic aloes, in powder, half an ounce; rectified fpirit of wine, two pints. Digeft them in a gentle heat for three days, and then ftrain the balfam.

85711O 10 JiC

This balfam, or rather tincture, is applied externally to heal recent wounds and bruifes. It is likewife employed internally to remove coughs, afthmas, and other complaints of the breaft. It is faid to eafe the colic, cleanfe the kidnies, and to heal internal ulcers, &c.

The dole is from twenty to fixty drops.

This, though a medicine of fome value, does not deferve the extravagant encomiums which have been bestowed on it. It has been celebrated under the different names of The Commander's Balfam, Perfian Balfam, Balfam of Berne, Wade's Balfam, Frier's Balfam, Jefuit's Drops, Turlington's Drops, &c.

BOLUSES.

A S bolufes are intended for immediate ufe, volatile falts, and other ingredients improper for being kept, are admitted into their composition. They are generally composed of powders, with a proper quantity of fyrup, conferve, or mucilage. The lighter powders are commonly made up with fyrup, and the more ponderous, as mercury, &c. with conferve; but those of the lighter kind would be more conveniently made up with mucilage, as it increases their bulk less than the other additions, and likewife occasions the medicine to pass down more easily.

Astringent Bolus.

Take of allum, in powder, fifteen grains; gum kino, five grains; fyrup, a sufficient quantity to make a bolus.

In an exceffive flow of the menses, and other violent discharges of blood, proceeding from relaxation, this bolus may be given every four or five hours, till the discharge abates.

Diaphoretic Bolus.

Take of gum guaiacum, in powder, ten grains; flowers of fulphur and cream of tartar, of each one fcruple; fimple fyrup, a fufficient quantity.

In rheumatic complaints, and diforders of the skin, this bolus may be taken twice a-day. It will also be of service in the inflammatory quintey.

Mercurial Bolus.

Take of calomel, fix grains; conferve of rofes, half a drachm. Make a bolus.

Where mercury is neceffary, this bolus may be taken twice or thrice a week. It may be taken over night; and if it does not operate, a few grains of jalap will be proper next day to carry it off.

BOLUSES.

Bolus of Rhubarb and Mercury.

Take of the best rhubarb, in powder, from a scruple to half a drachm ; of calomel, from four to fix grains ; fimple fyrup, a fufficient quantity to make a bolus.

This is a proper purge in hypochondriac conflictutions; but its principal intention is to expel worms. Where a ftronger purge is neceffary, jalap may be used instead of the rhubarb.

Pectoral Bolus.

Take of spermaceti, a scruple ; gum ammoniae, ten grains ; falt of hartfhorn, fix grains ; fimple fyrup, as much as will make them into a bolus.

This bolus is given in colds and coughs of long flanding, afthmas, and beginning confumptions of the lungs. It is generally proper to bleed the patient before he begins to use it,

Purging Bolus.

Take of jalap, in powder, a scruple; cream of tartar, two fcruples. Let them be rubbed together, and formed into a bolus, with fimple fyrup.

Where a mild purge is wanted, this will answer the purpose very well. If a ftronger dole is neceffary, the jalap may be increafed to half a drachm or upwards.

CATAPLASMS AND SINAPISMS.

CATAPLASMS posses few or no virtues superior to a poultice, which may be fo made, as, in most cases, to supply their place. They are chiefly intended either to act as difcutients, or to promote suppuration; and as they may be of fervice in lome cales, we shall give a specimen of each kind.

Discutient Cataplasm.

Take of ba ley-meal, fix ounces; fresh hemlock leaves, bruifed, two ounces; vinegar, a fufficient quantity. Boil the meal and hemlock in the vinegar for a little time, and then add two drachms of the fugar of lead.

Ripening Cataplasm.

Take of white lily root, four ounces; fat figs and raw onions, bruifed, of each one ounce; yellow bafilicum ointment, two ounces; gum galbanum, half an ounce; linfeed meal, as much as neceffary. Boil the rocts along with the figs in a fufficient quantity of water; then bruife and add to them the other ingredients,

2

face of villant

y 18613301 5390

dients, fo as to form the whole into a fost cataplasm. The galbanum must be previously diffolved with the yolk of an egg.

Where it is neceffary to promote fuppuration, this cataplafm may he ufed by those who choose to be at the trouble and expence of making it. For my part, I have never found any application more proper for this purpose than a poultice of bread and milk, with a sufficient quantity of either boiled or raw onion in it, and softened with oil or fresh butter.

Sinapi/ms.

Sinapifms are employed to recal the blood and fpirits to a weak part, as in the palfy and atrophy. They are alfo of fervice in deep-feated pains, as the fciatica, &c. When the gout feizes the head or the ffomach, they are applied to the feet to bring the diforder to these parts. They are likewise applied to the patient's foles in the low frate of fevers. They should not be suffered to lie on, however, till they have raised blifters, but till the parts become red, and will continue fo when prefied with the finger.

The finapism is only a poultice made with vinegar instead of milk, and rendered warm and stimulating by the addition of mustard, horse-raddish, or garlic.

The common finapifm is made by taking crumb of bread and muftard-feed in powder, of each equal quantities; ftrong vinegar, as much as is fufficient, and mixing them fo as to make a poultice.

When finapisms of a more flimulating nature are wanted, a little bruised garlic may be added to the above.

CLYSTERS.

T HIS class of medicines is of more importance than is generally imagined. Clyfters ferve not only to evacuate the contents of the belly, but alfo to convey very active medicines into the fyftem. Opium, for example, may be administered in this way when it will not fit upon the flomach, and alfo in larger doses than at any time it can be taken by the mouth. The Peruvian bark may likewise be, with good effect, administered in form of clyfter to perfons who cannot take it by the mouth.

A fimple clyfter can feldom do hurt, and there are many cafes where it may do much good. A clyfter even of warm water, by ferving as a fomentation to the parts, may be of confiderable fervice in inflammations of the bladder, and the lower inteffines, &c.

Some

Some fubftances, as the fmoke of tobacco, may be thrown into the bowels in this way, which cannot be done by any other means whatever. This may be eafily effected by means of a pair of hand-bellows, with an apparatus fitted to them for that purpose.

Nor is the use of clyfters confined to medicines. Aliment may also be conveyed in this way. Perfons unable to fwallow, have been, for a confiderable time, supported by clyfters.

Emollient Clyfter.

Take of linfeed-tea and new milk, each fix ounces. Mix them.

If fifty or fixty drops of laudanum be added to this, it will fupply the place of the Anodyne Clyfter.

Laxative Clyfter.

Take of milk and water, each fix ounces; fweet oil or fresh butter, and brown fugar, of each two ounces. Mix them.

If an ounce of Glauber's falt, or two table-spoonfuls of common falt, be added to this, it will be the Purging Clyster.

Carminative Cly/ler.

Take of camomile flowers, an ounce; anife feeds, half an ounce. Boil in a pint and a half of water to one pint.

In hyfteric and hypochondriac complaints, this may be adminiftered inftead of the Fætid Clyfter, the fmell of which is fo difagreeable to most patients.

Oily Clyfter.

To four ounces of the infusion of camomile flowers, add an equal quantity of Florence oil.

This clyfter is beneficial in bringing off the fmall worms lodged in the lower parts of the alimentary canal. When given to children, the quantity must be proportionably leffened.

Starch Clyfter.

Take jelly of flarch, four ounces; linseed oil, half an ounce. Liquefy the jelly over a gentle fire, and then mix in the oil.

In the dytentery or bloody flux, this clyfter may be adminiftered after every loofe flool, to heal the ulcerated inteffines, and blunt the fharpnefs of corroding humours. Forty or fifty drops of laudanum may be occasionally added; in which cafe, it will generally fupply the place of the Aftringent Clyfter.

Turpentine

Turpentine Clyfter.

Take of common decociion, ten ounces; Venice turpentine, diffolved with the yolk of an egg, half an ounce; Florence oil, one ounce. Mix them.

This diuretic clyfter is proper in obstructions of the urinary passages, and in colicky complaints, proceeding from gravel.

Vinegar Clyfter.

This clyfter is made by mixing three ounces of vinegar with five of water-gruel.

It answers all the purposes of a common clyster, with the peculiar advantage of being proper either in inflammatory or putrid diforders, especially in the latter.

We think it unneceffary to give more examples of this clafs of medicines, as ingredients adapted to any particular intention may be occasionally added to one or other of the above forms.

COLLYRIA, or EYE-WATERS.

E YE-WATERS have been multiplied without number, almost every perfon pretending to be possefield of some fecret preparation for the cure of sore-eyes. I have examined many of them, and find that they are pretty much alike, the basis of most of them being either alum, vitriol, or lead. Their effects evidently are to brace and reftore the tone of the parts: hence they are principally of service in flight inflammations; and in that relaxed state of the parts which is induced by obstinate ones.

Camphor is commonly added to these compositions; but as it feldom incorporates properly with the water, it can be of little use. Boles and other earthy substances, as they do not diffolve in water, are likewise unfit for this purpose.

Collyrium of Alum.

Take of alum half a drachm : agitate it well together with the white of one egg.

This is the Collyrium of Riverius. It is used in inflammation of the eyes, to allay heat, and reftrain the flux of humours. It must be spread upon linen, and applied to the eyes; but should not be kept on above three or four hours at a time.

Vitriolic Collyrium.

Take of white vitriol, half a drachm; rose-water, fix ounces. Diffolve the vitriol in the water, and filter the liquor.

This,

666

This, though fimple, is perhaps equal in virtue to most of the celebrated collyria. It is an useful application in weak, watery, and inflamed eves. Though the flighter inflammations will generally yield to it, yet in those of a more obstinate nature the affistance of bleeding and bliftering will often be neceffary.

When a firong aftringent is judged proper, a double or triple quantity of the vitriol may be used. I have seen a solution of four times the firength of the above used with manifest advantage.

Collyrium of Lead.

Take fugar of lead, and crude fal ammoniac, of each four grains. Diffolve them in eight ounces of common water.

Forty or fifty drops of laudanum may be occafionally added to this collyrium.

Those who choose may substitute, instead of this, the collyrium of lead, recommended by Goulard; which is made by putting twenty-five drops of his *Extrast of Lead* to eight ounces of water, and adding a tea-spoonful of brandy.

Indeed, common water and brandy, without any other addition, will in many cafes answer very well as a collyrium. An ounce of the latter may be added to five or fix ounces of the former; and the eyes, if weak, bathed with it night and morning.

I have of late been troubled with a rheum in my eyes, for which I have found great benefit, by washing them frequently with rofe-water. Many experience the like good effect from anointing the eye lids with Smelhom's falve.

CONFECTIONS.

CONFECTIONS, containing above fixty ingredients, are ftill to be found in fome of the most reformed dispensatories. As most of their intentions, however, may be more certainly, and as effectually answered by a few glasses of wine or grains of opium, we shall pass over this class of medicines very flightly.

Japonic Confection.

Take of Japan earth, three ounces; tormentil root, nutmeg, olibanum, of each two ounces; opium diffolved in a fufficient quantity of Lisbon wine, a drachm and a half; fimple fyrup and conferve of rofes, of each fourteen ounces. Mix and make them into an electuary.

This supplies the place of the Diafcordium.

The dose of this electuary is from a fcruple to a drachm.

CON-

300

CONSERVES AND PRESERVES.

EVERY Apothecary's thop was formerly to full of these preparations, that it might have passed for a confectioner's warehouse. They posses very few medicinal properties, and may rather be classed among sweetmeats than medicines. They are fometimes, however, of use, for reducing into boluses or pills fome of the more ponderous powders, as the preparations of iron, mercury, and tin.

Conferves are compositions of fresh vegetables and fugar, beaten together into an uniform mass. In making these preparations, the leaves of vegetables must be freed from their stalks, the flowers from their cups, and the yellow part of orange-peel taken off with a rasp. They are then to be pounded in a marble mortar, with a wooden pessel, into a smooth mass; after which, thrice their weight of fine sugar is commonly added by degrees, and the beating continued till they are uniformly mixed; but the conferve will be better if only twice its weight of sugar be added.

Those who prepare large quantities of conferve generally reduce the vegetables to a pulp by the means of a mill, and afterwards beat them up with the sugar.

Conferve of Red Rofes.

Take a pound of red rofe buds, cleared of their heels; beat them well in a mortar, and adding by degrees two pounds of double-refined fugar, in powder, make a conferve.

After the fame manner are prepared the conferves of orangepeel, rofemary flowers, fea-wormwood, the leaves of woodforrel, &c.

The conferve of roles is one of the most agreeable and useful preparations belonging to this class. A drachm or two of it, diffolved in warm milk, is ordered to be given as a gentle reftringent in weakness of the ftomach, and likewise in phthifical coughs, and spitting of blood. To have any confiderable effects, however, it must be taken in larger quantities.

Conferve of Sloes.

This may be made by boiling the floes gently in water, being careful to take them out before they burft; afterwards expreffing the juice, and beating it up with three times its weight of fine fugar.

In relaxations of the uvula and glands of the throat, this makes an excellent gargle, and may be used at diferention.

Preferves are made by fteeping or boiling fresh vegetables first in water, and afterwards in fyrup, or a folution of sugar. The subject is either preferved moist in the syrup, or taken out and dried,

CONSERVES AND PRESERVES.

668

dried, that the fugar may candy upon it. The laft is the moff uleful method.

Candied Orange Peel.

Soak Seville orange-peel in feveral waters, till it lofes its bitternefs; then boil it in a folution of double-refined fugar in water, till it becomes tender and transparent.

Candied lemon peel is prepared in the fame manner.

It is needless to add more of these preparations, as they belong rather to the art of the confectioner than that of the apothecary.

DECOCTIONS.

WATER readily extracts the gummy and faline parts of vegetables; and though its action is chiefly confined to thefe, yet the refinous and oily being intimately blended with the gummy and faline, are in great part taken up along with them. Hence watery decoctions and infufions of vegetables conflitute a large, and not unufeful, clafs of medicines. Although moft vegetables yield their virtues to water, as well by infution as decoction, yet the latter is often neceffary, as it faves time, and does in a few minutes what the other would require hours, and fometimes days, to effect.

The medicines of this clafs are all intended for immediate ufe.

Decostion of Althea.

Take of the roots of marshmallows, moderately dried, three ounces; raisins of the fun, one ounce; water, three pints.

Boil the ingredients in the water till one third of it is confumed; afterwards ftrain the decoction, and let it ftand for fome time to fettle. If the roots be thoroughly dried, they must be boiled till one half of the water be confumed.

In coughs, and fharp defluxions upon the lungs, this decoction may be used for ordinary drink.

The Common Decostion.

Take of camomile flowers, one ounce; elder flowers, and fweet fennel feeds, of each half an ounce; water, two quarts. Boil them for a little while, and then ftrain the decoction.

A medicine equally good may be prepared by infufing the ingredients for fome hours in boiling water.

This decoction is chiefly intended as the bafis of clyfters, to which other ingredients may be occasionally added. It will likewife ferve as a common fomentation, spirit of wine or other things being added in such quantity as the case may require.

DECOCTIONS.

Decotion of Logwood.

Boil three ounces of the fhavings, or chips of logwood, in four pints of water, till one half the liquor is wafted. Two or three ounces of fimple cinnamon-water may be added to this decoction.

In fluxes of the belly, where the ftronger aftringents are improper, a tea-cupful of this decoction may be taken with advantage three or four times a-day.

Decostion of the Bark.

Boil an ounce of the Peruvian bark großly powdered, in a pint and a half of water to one pint; then firain the decoction. If a tea-fpoonful of the weak fpirit of vitriol be added to this medicine, it will render it both more agreeable and efficacious.

Compound Decostion of the Bark.

Take of Peruvian bark, and Virginian inake-root, grossly powdered, each three drachms. Boil them in a pint of water to one half. To the fitrained liquor add an ounce and a half of aromatic water.

Sir John Pringle recommends this as a proper medicine towards the decline of malignant fevers, when the pulfe is low, the voice weak, and the head affected with a flupor but with little delirium.

The dofe is four spoonfuls every fourth or fixth hour.

Decoction of Sarfaparilla.

Take of fresh farsaparilla root, fliced and bruised, two ounces; shavings of guaiacum wood, one ounce. Boil over a flow fire, in three quarts of water, to one; adding towards the end, half an ounce of fassafas wood, and three drachms of liquorice. Strain the decosition.

This may either be employed as an affiftant to a courfe of mercurial alteratives, or taken after the mercury has been used for fome time. It strengthens the stomach, and restores shelf and vigour to habits emaciated by the venereal difease. It may also be taken in the rheumatism, and cutaneous diforders proceeding from foulness of the blood and juices. For all these intentions it is greatly preferable to the *Decostion of Wood*.

This decoclion may be taken, from a pint and a half to two quarts in the day.

The following decoclion is faid to be fimilar to that used by *Kennedy*, in the cure of the venereal difease, and may supply the place of Lifbon diet drink :

Take of farfapatilla, three ounces; liquorice and mezerion root, of each half an ounce; fhavings of guaiacum and faffafras wood, wood, of each one ounce; crude antimony, powdered, an ounce and a half. Infufe these ingredients in eight pints of boiling water for twenty-four hours, then boil them till one half of the water is confumed; afterwards strain the decoction.

This decoction may be used in the same manner as the preceding.

Decoction of Seneka.

Take of seneka rattle-snake root, one ounce; water, a pint and a half. Boil to one pint, and strain.

This decoction is recommended in the pleurify, dropfy, rheumatism, and some obstinate diforders of the skin. The dose is two ounces, three or sour times a-day or oftener, if the stomach will bear it.

White Decostion.

Take of the puteft chalk, in powder, two ounces; gum arabic, half an ounce; water, three pints. Boil to one quart, and strain the decoction.

This is a proper drink in acute difeafes, attended with, or inclining to, a loofenefs, and where acidities abound in the flomach or bowels. It is peculiarly proper for children when afflicted with fournefs of the flomach, and for perfors who are fubject to the heartburn. It may be fweetened with fugar, as it is ufed, and two or three ounces of fimple cinnamon-water added to it.

An ounce of powdered chalk, mixed with two pints of water, will occafionally fupply the place of this decoction, and also of the chalk julep.

DRAUGHTS.

THIS is a proper form for exhibiting fuch medicines as are intended to operate immediately, and which do not need to be frequently repeated, as purges, vomits, and a few others, which are to be taken at one dofe. Where a medicine requires to be used for any length of time, it is better to make up a larger quantity of it at once, which faves both trouble and expence.

Anodyne Draught.

Take of liquid laudanum, twenty-five drops; fimple einnamon-water, an ounce; common fyrup, two drachms. Mix them.

In exceffive pain, where bleeding is not neceffary, and in great reftleffnefs, this composing draught may be taken and repeated occasionally.

Diuretic Draught.

Take of the diuretic falt, two fcruples; fyrup of poppies, two drachms; fimple cinnamon water, and common water, of each an ounce. This

DRAUGHTS.

This draught is of fervice in an obstruction or deficiency of urine.

Purging Draughts.

Take of manna, an ounce; foluble tartar, or Rochel falt, from three to four drachms. Diffolve in three ounces of boiling water; to which add Jamaica pepper-water, half an ounce.

As manna fometimes will not fit upon the ftomach, an ounce or ten drachms of the bitter purging falts, diffolved in four ounces of water, may be taken inflead of the above.

Those who cannot take falts, may use the following draught:

Take of jalap in powder, a fcruple; common water, an ounce; aromatic tincture, fix drachms. Rub the jalap with twice its weight of fugar, and add to it the other ingredients.

Sweating Draughts.

Take spirit of Mindererus, two ounces; salt of hartshorn, five grains; simple cinnamon water, and syrup of poppies, of each half an ounce. Make them into a draught.

In recent colds and rheumatic complaints, this draught is of fervice. To promote its effects, however, the patient ought to drink freely of warm water gruel, or of fome other weak diluting liquor.

Vomiting Draughts.

Take of ipecacuanha, in powder, a scruple; water, an ounce; fimple syrup, a drachm. Mix them.

Perfons who require a ftronger vomit may add to the above half a grain, or a grain, of emetic tartar.

Those who do not chuse the powder, may take ten drachms of the ipecacuanha wine; or half an ounce of the wine, and an equal quantity of the fyrup of fquills.

ELECTUARIES.

ELECTUARIES are generally composed of the lighter powders, mixed with fyrup, honey, conferve, or mucilage, into fuch a confistence, that the powders may neither feparate by keeping, nor the mass prove too stiff for fwallowing. They receive chiefly the milder alterative medicines, and such as are not ungrateful to the palate.

Aftringent electuaries, and fuch as have pulps of fruit in them, fhould be prepared only in fmall quantities; as aftringent medicines lofe their virtues by being kept in this form, and the pulps of fruits are apt to ferment.

For the extraction of pulps it will be neceffary to boil unripe fruits, and ripe ones, if they are dried, in a fmail quantity of water till they become foft. The pulp is then to be preffed out through a ftrong

ELECTUARIES.

a ftrong hair fieve, or thin cloth, and afterwards boiled to a due confiftence, in an earthen veffel, over a gentle fire, taking care to prevent the matter from burning by continually flirring it. The pulps of fruit that are both ripe and fresh may be pressed out without any previous boiling.

Lenitive Electuary.

Take of fenna, in fine powder, eight ounces; coriander feed, alfo in powder, four ounces; pulp of tamarinds and of French prunes, each a pound. Mix the pulps and powders together, and with a fufficient quantity of fimple fyrup, reduce the whole into an electuary.

A tea-fpoonful of this electuary, taken two or three times aday, generally proves an agreeable laxative. It likewife ferves as a convenient vehicle for exhibiting more active medicines, as jalap fcammony, and fuch like.

This may supply the place of the electuary of Cassia.

to a identify and Electuary for the Dyfentery.

Take of the Japonic confection, two ounces; Locatelli's balfam, one ounce; rhubarb in powder, half an ounce; fyrup of marfhmallows, enough to make an electuary.

It is often dangerous in dyfenteries to give opiates and aftringents, without interposing purgatives. The purgative is here joined with these ingredients, which renders this a very fafe and useful medicine for the purposes expressed in the title.

About the bulk of a nutmeg fhould be taken twice or thrice a day, as the fymptoms and conflitution may require.

Electuary for the Epilepsy.

Take of Peruvian bark, in powder, an ounce; of powdered tin, and wild valerian root, each half an ounce; fimple fyrup enough to make an electuary.

Dr. Mead directs a drachm of an electuary fimilar to this to be taken evening and morning, in the epileply, for the fpace of three months. It will be proper, however, to difcontinue the use of it for a few days every now and then. I have added the powdered tin, because the epileply often proceeds from worms.

Electuary for the Gonorrhæa.

Take of lenitive electuary, three ounces; jalap and rhubarb, in powder, of each two drachms; nitre, half an ounce; fimple fyrup, enough to make an electuary.

During the inflammation and tenfion of the urinary paffages, which accompany a virulent gonorrhœa, this cooling laxative may be used with advantage.

The

south a

672

The dofe is a drachm, or about the bulk of a nutmeg, two or three times a day; more or lefs, as may be neceffary to keep the body gently open.

An electuary made of cream of tartar and fimple fyrup will occafionally fupply the place of this.

After the inflammation is gone off, the following electuary may be used :

Take of lenitive electuary, two ounces; balfam of capivi, one ounce; gum guaiacum and rhubarb, in powder, of each two drachms; fimple fyrup, enough to make an electuary. The dofe is the fame as of the preceding.

Electuary of the Bark.

Take of Peruvian bark, in powder, three ounces; cafcarilla, half an ounce; fyrup of ginger, enough to make an electuary.

In the cure of obstinate intermitting fevers, the bark is affisted by the cascarilla. In hectic habits, however, it will be better to leave out the cascarilla, and put three drachms of crude sal ammoniac in its flead.

Electuary for the Piles.

Take flowers of fulphur, one ounce; cream of tartar, half an ounce; treacle, a fufficient quantity to form an electuary. A tea-fpoonful of this may be taken three or four times a-day.

Electuary for the Palfy.

Take of powdered muftard-feed, and conferve of roles, each an ounce; fyrup of ginger, enough to make an electuary. A tea-fpoonful of this may be taken three or four times a-day.

Electuary for the Rheumatism.

Take of conferve of roles, two ounces; cinnabar of antimony, levigated, an ounce and a half; gum guaiacum, in powder, an ounce; fyrup of ginger, a fufficient quantity to make an electuary.

In obstinate rheumatisms, which are not accompanied with a fever, a tea-spoonful of this electuary may be taken twice a-day with confiderable advantage.

EMULSIONS.

E MULSIONS, befide their use as medicines, are also proper vehicles for certain substances, which could not otherwise be conveniently taken in a liquid form. Thus camphor, triturated with almonds, readily unites with water into an emulfion. Pure oils, balfams, refins, and other fimilar substances, are likewise rendered miscible with water by the intervention of mucilages.

Xx

Common

EMULSIONS.

Common Emulfion.

Take of fweet almonds, an ounce; bitter almonds, a drachin ; water, two pinte.

Let the almonds be blanched, and beat up in a marble mortar; adding the water by little and little, fo as to make an emullion; afterwards let it be ftrained.

Arabic Emulsion.

This is made in the fame manner as the above, adding to the almonds, while beating, two ounces and a half of the mucilage of gum arabic.

Where fost cooling liquors are neceffary, these emulsions may be used as ordinary drink.

Camphorated Emulfion.

Take of camphor, half a drachm; fweet almonds, half a dozen; white fugar, half an ounce; mint water, eight ounces. Grind the camphor and almonds well together in a ftone mortar, and add by degrees the mint water; then ftrain the liquor, and diffolve in it the fugar.

In fevers, and other diforders which require the use of camphor, a table-spoonful of this emultion may be taken every two or three hours.

Emulsion of Gum Ammoniac.

Take of gum ammoniac, two drachms ; water, eight ounces. Grind the gum with the water poured upon it by little and little, till it is difiolved.

This emultion is used for attenuating tough, viscid phlegm, and promoting expectoration. In obflinate coughs, two ounces of the fyrup of poppies may be added to it. The dose is two table-spoonfuls three or four times a day.

Oily Emulfion.

Take of foft water, fix ounces ; volatile aromatic fpirit, two drachms ; Florence oil, an ounce ; fhake them well together, and add, of fimple fyrup, half an ounce.

In recent colds and coughs, this emulfion is generally of fervice; but if the cough proves obstinate, it will fucceed better when made with the paregoric elixir of the Edinburgh Dispensatory, instead of the volatile aromatic spirit. A table-spoonful of it may be taken every two or three hours.

EXTRACTS.

EXTRACTS are prepared by boiling the fubject in water, and evaporating the firained decoclion to a due confiftence. By this process fome of the more active parts of plants are freed from from the ufeles, indiffoluble earthy matter, which makes the larger thare of their bulk. Water, however, is not the only menftruum used in the preparation of extracts; fometimes it is joined with spirits, and at other times rectified spirit alone is employed for that purpose.

Extra s are prepared from a variety of different drugs, as the bark, gentian, jalap, &c.; but, as they require a troublefome and tedious operation, it will be more convenient for a private practitioner to purchase what he needs of them from a professed druggist, than to prepare them himself. Such of them as are generally used are inserted in our list of such of them as and cines as are to be kept for private practice.

FOMENTATIONS.

F OMENTATIONS are generally intended either to eafe pain, by taking off tenfion and fpaim; or to brace and reftore the tone and vigour of those parts to which they are applied. The first of these intentions may generally be answered by warm water, and the second by cold. Certain substances, however, are usually added to water with a view to heighten its effects, as anodynes, aromatics, astringents, &c. We shall therefore subjoin a few of the most useful medicated fomentations, that people may have it in their power to make use of them as they chuse.

Anodyne Fomentation.

Take of white poppy heads, two ounces; elder flowers, half an ounce; water, three pints. Boil till one pint is evaporated; and strain out the liquor.

This fomentation, as its title expresses, is used for relieving acute pain.

Aromatic Fomentation.

Take of Jamaica pepper, half an ounce; red wine, a pint. Boil them for a little, and then ftrain the liquor.

This is intended, not only as a topical application for external complaints, but also for relieving the internal parts. Pains of the bowels, which accompany dyfenteries and diarrhœas, flatulent colics, uneafiness of the stomach, and reachings to vomit, are frequently abated by tomenting the abdomen and region of the stomach with the warm liquor.

Common Fomentation.

Take tops of wormwood and camomile flowers, dried, of each two ounces; water, two quarts. After a flight boiling, pour off the liquor.

Brandy or ipirit of wine may be added to this fomentation, in fuch quantity as the particular circumftances of the cafe thall require; but these are not always necessary.

Xx2

Emollient

675

FOMENTATIONS.

Emollient Fomentation.

This is the fame as the common decoction.

Strengthening Fomentation.

Take of oak bark, one ounce; granate peel, half an ounce; allum, two drachms, fmith's forge water, three pints. Boil the water with the bark and peel to the confumption of one-third; then ftrain the remaining decoction, and diffolve in it the allum.

This affringent liquor is employed as an external fomentation to weak parts; it may also be used internally.

GARGLES.

HOWEVER trifling this clafs of medicines may appear, they are by no means without their ufe. They feldom, indeed, cure difeates, but they often alleviate very difagreeable fymptoms; as parchednefs of the mouth, foulnefs of the tongue and fauces, &c. they are peculiarly ufeful in fevers and fore throats. In the latter, a gargle will fometimes remove the diforder; and in the former few things are more refrefhing or agreeable to the patient, than to have his mouth frequently walked with fome foft detergent gargle.

One advantage of these medicines is, that they are easily prepared. A little barley-water and honey may be had any where; and if to these be added as much vinegar as will give them an agreeable sharpness, they will make a very useful gargle for fostening and cleansing the mouth.

Gargles have the best effect when injected with a fyringe.

Attenuating Gargle.

Take of water, fix ounces; honey, one ounce; nitre, a drachm and a half. Mix them.

This cooling gargle may be used either in the inflammatory quinfey, or in fevers, for cleaning the tongue and fauces.

Common Gargle

Take of rofe-water, fix ounces; tyrup of clove July flowers, half an ounce; fpirit of vitriol, a fufficient quantity to give it an agreeable fharpnefs. Mix them.

This gargle, belides cleanling the tongue and fauces, acts as a gentle repellent, and will fometimes remove a flight quinfey.

Detergent Gargle.

Take of the emollient gargle a pint; tincture of myrrh, an ounce; honey, two ounces. Mix them.

When exulcerations require to be cleanfed, or the excretion of tough viscid faliva promoted, this gargle will be of fervice.

Emolliene

GARGLES.

Emollient Gargle.

Take an ounce of marshmallow roots, and two or three figs: boil them in a quart of water till near one half of it be confumed; then strain out the liquor.

If an ounce of honey, and half an ounce of fpirit of fal-ammoniac, be added to the above, it will then be an exceeding good attenuating gargle.

This gargle is beneficial in fevers, where the tongue and fauces are rough and parched, to foften these parts, and promote the discharge of faliva.

The learned and accurate Sir John Pringle obferves, that in the inflammatory quinfey, or ftrangulation of the fauces, little benefit arifes from the common gargles; that fuch as are of an acid nature do more harm than good, by contracting the emunctories of the faliva and mucus, and thickening those humours; that a decoction of figs in milk and water has a contrary effect, especially if some fal-ammoniac be added; by which the faliva is made thinner, and the glands are brought to fecrete more freely; a circumftance always conducive to the cure.

INFUSIONS.

VEGETABLES yield nearly the fame properties to water by infufion as by decoclion; and though they may require a longer time to give out their virtues in this way, yet it has feveral advantages over the other; fince boiling is found to diffipate the finer parts of many bitter and aromatic fubftances, without more fully extracting their medicinal principles.

The author of the New Difpenfatory obferves, that even from those vegetables which are weak in virtue, rich infusions may be obtained, by returning the liquor upon fresh quantities of the subject, the water loading itself more and more with the active parts; and that these loaded infusions are applicable to valuable purposes in medicine, as they contain in a small compass the finer, more subtile, and active principles of vegetables, in a sorm readily miscible with the fluids of the human body.

Bitter Infusion.

Take tops of the leffer centaury and camomile flowers, of each half an ounce; yellow rind of lemon and orange-peel, carefully freed from the inner white part, of each two drachms. Cut them in fmall pieces, and infuse them in a quart of boiling water.

For indigeftion, weakness of the ftomach, or want of appetite, a tea-cupful of this infusion may be taken twice or thrice a-day.

Infusion

INFUSIONS.

Infusion of the Bark.

To an ounce of the bark, in powder, add four or five tables spoonfuls of brandy, and a pint of boiling water. Let them infule for two or three days.

This is one of the best preparations of the bark for weak ftomachs. In diforders where the corroborating virtues of that medicine are required, a tea- cup full of it may be taken two or three times a-day.

Infusion of Carduus.

Infuse an ounce of the dried leaves of carduus benedictus, or bleffed thiftle, in a pint of common water, for fix hours, without heat; then filter the liquor through paper.

This light infufion may be given, with great benefit, in weaknels of the ftomach, where the common bitters do not agree. It may be flavoured at pleafure with cinnamon, or other aromatic materials.

Infusion of Linseed.

Take of linfeed, two spoonfuls; liquorice root, fliced, half an ounce; boiling water, three pints. Let them stand to infuse by the fire for some hours, and then strain off the liquor.

If an ounce of the leaves of colt's-foot be added to these ingredients, it will then be the *Pectoral Infusion*. Both these are emollient mucilaginous liquors, and may be taken with advantage as ordinary drink in difficulty of making water; and in coughs and other complaints of the breaft.

Infusion of Roses.

Take of red roles, dried, half an ounce; boiling water, a quart; vitriolic acid, commonly called oil of vitriol, half a drachm; loaf fugar, an ounce.

Infuse the roses in the water for four hours, in an unglazed earthen veffel; afterwards pour in the acid, and, having strained the liquor, add to it the sugar.

In an exceflive flow of the menses, vomiting of blood, and other hæmorrhages, a tea-cup full of this gently aftringent infusion may be taken every three or four hours. It likewise makes an exceeding good gargle.

As the quantity of roles used here can have little or no effect, an equally valuable medicine may be prepared by mixing the acid and water without infusion.

Infusion of Tamarinds and Senna.

Take of tamarinds one ounce; fenna, and crystals of tartar, each two drachms. Let these ingredients be infused four or five

INFUSIONS.

five hours in a pint of boiling water; afterwards let the liquor be ftrained, and an ounce or two of the aromatic tincture added to it. Perfons who are eafily purged may leave out either the tamarinds or the cryftals of tartar.

This is an agreeable cooling purge. A tea-cup full may be given every half hour till it operates.

This supplies the place of the Decostion of Tamarinds and Senna.

Spanish Infusion.

Take of Spanish juice, cut into small pieces, an ounce; falt of tartar, three drachms. Infuse in a quart of boiling water for a night. To the strained liquor add an ounce and an half of the syrup of poppies.

In recent colds, coughs, and obstructions of the breast, a teacup full of this infusion may be taken with advantage three or four times a-day.

Infusion for the Palfy.

Take of horfe radifh root thaved, muftard feed bruifed, each four ounces; outer rind of orange-peel, one ounce. Infufe them in two quarts of boiling water, in a close veffel, for twenty-four hours.

In paralytic complaints, a tea-cup full of this warm ftimulating medicine may be taken three or four times a-day. It excites the action of the folids, proves diuretic, and, if the patient be kept warm, promotes perfpiration.

If two or three ounces of the dried leaves of marsh-trefoil be used instead of the mustard, it will make the Antifcorbutic Infusion.

JULEPS.

THE basis of Juleps is generally common water, or some fimple diffilled water, with one-third or one-fourth its quantity of diffilled spirituous water, and as much sugar or syrup as is sufficient to render the mixture agreeable. This is tharpened with vegetable or mineral acids, or impregnated with other medicines suitable to the intention.

Campborated Julep.

Take of camphor, one drachm; reclified spirit of wine, ten drops; double-refined sugar half an ounce; boiling distilled water, one pint. Rub the camphor first with the spirit of wine, then with the sugar; lastly, add the water by degrees, and strain the liquor.

In hysterical and other complaints, where camphor is proper, this julep may be taken in the dose of a spoonful or two as often as the stomach will bear it.

Xx4

Cordial

JULEPS.

Cordial Julep.

Take of fimple cinnamon-water, four ounces; Jamaica pepper-water, two ounces; volatile aromatic fpirit, and compound fpirit of lavender, of each two drachms; fyrup of orange-peel, an ounce. Mix them.

This is given in the dofe of two spoonfuls three or four times a-day, in diforders accompanied with great weakness and depression of spirits.

Expectorating Julep.

Take of the emultion of gum ammoniac, fix ounces; fyrup of fquills, two ounces. Mix them.

In coughs, afthmas, and obstructions of the breast, two table-spoonfuls of this julep may be taken every three or four hours.

Musk Julep.

Rub half a drachm of mosk well together with half an ounce of fugar, and add to it, gradually, of fimple cinnamon and peppermint-water, each two ounces; of the volatile aromatic spirit, two drachms.

In the low flate of nervous fevers, hiccuping, convultions, and other fpafmodic affections, two table-fpoonfuls of this julep may be taken every two or three hours.

Saline Julep.

Diffolve two drachms of falt of tartar in three ounces of fresh lemon juice, strained; when the effervescence is over, add, of mint-water, and common water, each two ounces; of simple syrup, one ounce.

This removes fickness at the flomach, relieves vomiting, promotes perspiration, and may be of some fervice in fevers, especially of the inflammatory kind.

Vomiting Julep.

Diffolve four grains of emetic tartar in eight ounces of water, and add to it half an ounce of the fyrup of clove July-flowers.

In the beginning of fevers, where there is no topical inflammation, this julep may be given in the dole of one table-fpoonful every quarter of an hour till it operates. Antimonial vomits ferve not only to evacuate the contents of the ftomach, but likewife to promote the different excretions. Hence they are found in fevers to have nearly the fame effects as Dr. James's Powder.

MIXTURES.

A MIXTURE differs from a julep in this respect, that it receives into its composition not only falts, extracts, and other substances diffoluble in water, but also earths, powders, and such substances as cannot be diffolved. A mixture is seldom either an elegant or agreeable medicine. It is nevertheles neceffary. Many persons can take a mixture, who are not able to swallow a bolus or an electuary : besides, there are medicines which act better in this than in any other form,

Astringent Mixture.

Take fimple cinnamon-water, and common water, of each three ounces; fpirituous cinnamon-water, an ounce and a half; Japonic confection, half an ounce. Mix them.

In dyfenteries which are not of long flanding, after the neceffary evacuations, a fpoonful or two of this mixture may be taken every four hours, interpofing every fecond or third day a dofe of rhubarb.

The Astringent Mixture, which I have lately made use of with great fucces, is prepared thus:

Take powder of bole with opium, two drachms; cinnamonwater and penny-royal water, of each three ounces; fpirituous cinnamon-water, fix drachms; fimple fyrup, one ounce. Mix them, and take a table-fpoonful four or five times a-day.

Diuretic Mixture.

Take of mint-water, five ounces; vinegar of fquills, fix drachms; fweet fpirit of nitre, half an ounce; fyrup of ginger, an ounce and a half. Mix them.

In obstructions of the urinary passages, two spoonfuls of this mixture may be taken twice or thrice a-day.

Laxative Abforbent Mixture.

Rub one drachm of magnefia alba in a mortar with ten or twelve grains of the best Turkey rhubarb, and add to them three ounces of common water; fimple cinnamon-water, and fyrup of fugar, of each one ounce.

As molt difeafes of infants are accompanied with acidities, this mixture may either be given with a view to correct thefe, or to open the body. A table-fpoonful may be taken for a dofe, and repeated three times a-day. To a very young child half a fpoonful will be fufficient.

When the mixture is intended to purge, the dole may either be increased, or the quantity of rhubarb doubled. This is one of the most generally useful medicines for children with which I am acquainted.

Saline Mixture.

Diffolve a drachm of the falt of tartar infour ounces of boiling water; and, when cold, drop into it fpirit of vitriol till the effervescence ceases; then add, of peppermint-water, two ounces, simple syrup, one ounce.

Where fresh lemons cannot be had, this mixture may occafionally supply the place of the faline jules.

Squill Mixture.

Take of fimple cinnamon-water, five ounces; vinegar of fquills, one ounce; fyrup of marshmallows, an ounce and a half. Mix them.

This mixture, by promoting expectoration, and the fecretion of urine, proves ferviceable in althmatic and dropfical habits. A table-spoonful of it may be taken frequently.

OINTMENTS, LINIMENTS, and CERATES.

NOTWITHSTANDING the extravagant encomiums which have been beflowed on different preparations of this kind, with regard to their efficacy in the cure of wounds, fores, &c. it is beyond a doubt, that the most proper application to a green wound is dry lint. But though ointments do not heal wounds and fores, yet they ferve to defend them from the external air, and to retain such substances as may be necessary for drying, deterging, destroying proud flesh, and such like. For these purposes, however, it will be sufficient to infert only a few of the most simple forms, as ingredients of a more active nature can occasionally be added to them.

Yellow Bafilicum Ointment.

Take of yellow wax, white refin, and frankincenfe, each a quarter of a pound; melt them together over a gentle fire; then add, of hogs' lard prepared, one pound. Strain the ointment while warm.

This ointment is employed for cleanfing and healing wounds and ulcers.

Ointment of Calamine.

Take of olive oil, a pint and a half; white wax, and calamine from levigated, of each half a pound. Let the calamine frome, reduced into a fine powder, be rubbed with fome part of the oil, and afterwards added to the reft of the oil and wax previoufly melted together, continually friring them till quite cold.

This

This ointment, which is commonly known by the name of Turner's Cerate, is an exceeding good application in burns and excoriations from whatever caufe,

Emollient Ointment.

Take of palm oil, two pounds; olive oil, a pint and a half; yellow wax, half a pound; Venice turpentine, a quarter of a pound. Melt the wax in the oils over a gentle fire; then mix in the turpentine, and firain the ointment.

This supplies the place of Althaa Ointment. It may be used for anointing inflamed parts, &c.

Eye Ointment.

Take of hogs' lard prepared, four ounces; white wax, two drachms; tutty prepared, one ounce; melt the wax with the lard over a gentle fire, and then fprukle in the tutty, continually firring them till the ointment is cold.

This outment will be more efficacious, and of a better confiftence, if two or three drachms of camphor be rubbed up with a little oil, and intimately mixed with it.

Another.

Take of camphor, and calamine from levigated, each fix drachms; verdegrife, well prepared, two drachms; hogs' lard, and mutton loet, prepared, of each two ounces. Rub the camphor well with the powder; after wards mix in the lard and fuet, continuing the tritule till they be perfectly united.

This ointment has been long in effeem for difeafes of the eyes. It ought, however, to be used with caution, when the eyes are much inflamed or very tender.

Issue Ointment.

Mix half an ounce of Spanish flies, finely powdered, in fix ounces of yellow basilicum ointment.

This ointment is chiefly intended for dreffing blifters, in order to keep them open during pleafure.

Ointment of Lead.

Take of olive oil, halt a pint; white wax, two ounces; fugar of lead, three drachms. Let the fugar of lead, reduced into a fine powder, be rubbed up with fome part of the oil, and afterwards added to the other ingredients, previoufly melted together, continually flurring them till quite cold.

I his cooling and gently aftringent ointment may be used in all cases where the intention is to dry and skin over the part, as in scalding, &c.

Mercurial Ointment.

Take of quickfilver, two ounces; hogs' lard, three ounces; mutton fuet, one ounce. Rub the quickfilver with an ounce of the hogs' lard in a warm mortar, till the globules be perfectly extinguished; then rub it up with the reft of the lard and fuet, previoufly melted together.

The principal intention of this ointment is to convey mercury into the body by being rubbed upon the fkin.

Ointment of Sulphur.

Take of hogs' lard prepared, four ounces; flowers of fulphur, an ounce and a half; crude fal ammoniac, two drachms; effence of lemon, ten or twelve drops. Make them into an ointment.

This ointment, rubbed upon the parts affected, will generally cure the itch. It is both the fafeft and beft application for that purpofe, and, when made in this way, has no difagreeable fmell.

Ointment for Diseases of the Skin.

Take of the ointment, commonly called unguentum citrinum, a drachm and a half; flour of brimftone and powder of hellebore, of each an ounce; hogs' lard, three ounces; effence of lemon, or oil of thyme, from twenty to thirty drops, to correct the offenfiveness of the smell. Make them into an ointment.

I have not only known many ordinary affections of the fkin cured by this ointment, but even fome of a very malignant nature, and approaching to leprofy.

White Ointment.

Take of olive oil, one pint; white wax and fpermaceti, of each three ounces. Melt them with a gentle heat, and keep them conftantly and brickly ftirring together, till quite cold.

If two drachms of camphor, previoufly rubbed with a fmall quantity of oil, be added to the above, it will make the White camphorated Ointment.

Liniment for Burns.

Take equal parts of Florence oil, or of fresh drawn linfeed oil, and lime water; shake them well together in a wide mouthed bottle, so as to form a liniment.

This is found to be an exceeding proper application for recent fcalds or burns. It may either be fpread upon a cloth, or the parts affected may be anointed with it twice or thrice a-day.

- White

White Liniment.

This is made in the fame manner as the white ointment, twothirds of the wax being left out.

This liniment may be applied in cafes of excoriation, where, on account of the largeness of the surface, the ointments with lead or calamine might be improper.

Liniment for the Piles.

Take of emollient ointment, two ounces; liquid laudanum, half an ounce. Mix these ingredients with the yolk of an egg, and work them well together.

Volatile Liniment.

Take of Florence oil, an ounce; fpirit of hartfhorn, half an , ounce. Shake them together.

This liniment, made with equal parts of the fpirit and oil, will be more efficacious, where the patient's fkin is able to bear it.

Sir John Pringle observes, that in the inflammatory quinfey, a piece of flannel, moistened with this liniment, and applied to the throat, to be renewed every four or five hours, is one of the most efficacious remedies; and that it feldom fails, after bleeding, either to leffen or carry off the complaint. The truth of this observation I have often experienced.

Camphoraied Oil.

Rub an ounce of camphor, with two ounces of Florence oil, in a mortar, till the camphor be entirely diffolved.

This antifpafmodic liniment may be used in obffinate rheumatifms, and in some other cases accompanied with extreme pain and tension of the parts.

FILLS.

MEDICINES which operate in a fmall dofe, and whofe difagreeable tafte or fmell makes it neceffary that they fhould be concealed from the palate, are most commodiously exhibited in this form. No medicine, however, that is intended to operate quickly, ought to be made into pills, as they often lie for a confiderable time on the ftomach before they are diffolved, fo as to produce any effect.

As the ingredients which enter the composition of pills are generally to contrived, that one pill of an ordinary fize may contain about five grains of the compound, in mentioning the dole dose we shall only specify the number of pills to be taken: as one, two, three, &c.

Composing Pill.

Take of purified opium, ten grains; Caffile foap, half a drachm. Beat them together, and form the whole into twenty pills.

When a quieting draught will not fit upon the ftomach, one, two, or three of these pills may be taken, as occasion requires.

Deobstruent Pill.

Take falt of fleel; fuccotrine aloes; myrrh in powder; of each a drachm. Make into forty pills, of which two are to be taken evening and morning.

I have found these pills of excellent service in obstructions of the menses. The late Dr. WATKINSON made it his dying request, that I would infert this prescription in the "Domestic Medicine," which he said would be immortal, and that "his soul panted for immortality."

Fætid Pill.

Take of alafœtida, half an ounce; fimple lyrup, as much as is neceffary to form it into pills.

In hyfteric complaints, four or five pills, of an ordinary fize, may be taken twice or thrice a day. They may likewife be of fervice to perfons afflicted with the affhma.

When it is neceffary to keep the body open, a proper quantity of rhubarb, aloes, or jalap, may occasionally be added to the above mais.

Hemlock Pill.

Take any quantity of the extract of hemlock, and adding to it about a fifth part its weight of the powder of the dried leaves, form it into pills of the ordinary fize.

The extract of hemlock may be taken from one grain to feveral drachms in the day. The beft method, however, of using these pills, is to begin with one or two, and to increase the dose gradually, as far as the patient can bear them, without any remarkable degree of stupor or giddines.

Mercurial Pill.

Take of purified quickfilver and honey, each half an ounce. Rub them together in a mortar, till the globules of mercury are perfectly extinguished; then add, of Castile foap, two drachms; powdered liquorice, or crumb of bread, a sufficient quantity to give the mass a proper confistence for pills.

When

When ftronger mercurial pills are wanted, the quantity of quickfilver may be doubled.

The dole of these pills is different, according to the intention with which they are given. As an alterant, two or three may be taken daily. To raise a falivation, four or five will be neceffary.

Equal parts of the above pill and powdered rhubarb made into a mais, with a fufficient quantity of fimple fyrup, will make a Mercurial Purging Pill.

Mercurial Sublimate Pill.

Diffolve fifteen grains of the corrofive fublimate of mercury in two drachms of the faturated folution of crude fal ammoniac, and make it into a pafte, in a glafs mortar, with a fufficient quantity of the crumb of bread. This mafs must be formed into one hundred and twenty pills.

This pill, which is the most agreeable form of exhibiting the fublimate, has been found efficacious, not only in curing the venereal difease, but also in killing and expelling worms, after other powerful medicines had failed*.

For the venereal difease, four of these pills may be taken twice a-day, as an alterant three, and for worms two.

Plummer's Pill.

Take of calomel, or fweet mercury, and precipitated fulphur of antimony, each three drachms; extract of liquorice, two drachms. Rub the fulphur and mercury well together : afterwards add the extract, and, with a fufficient quantity of the mucilage of gum-arabic, make them into pills.

This pill has been found a powerful, yet fafe, alterative in obflinate cutaneous diforders; and has completed a cure after falivation had failed. In venereal cafes it has likewife produced excellent effects. Two or three pills of an ordinary fize may be taken night and morning, the patient keeping moderately warm, and crinking after each dofe a draught of decoction of the woods, or of farfaparilla.

Purging Pills.

Take of fuccotrine aloes, and Caffile foap, each two drachms; of fimple fyrup, a fufficient quantity to make them into pills.

Four or five of these pills will generally prove a sufficient purge. For keeping the body gently open, one may be taken might and morning. They are reckoned both deobstruent and

* See a paper on this fubject in the Edinburgh Phyfical and Literary Effays, by the ingenious Dr. John Gardener,

ftomachic,

ftomachic, and will be found to answer all the purposes of Dr. Anderson's pills, the principal ingredient of which is aloes.

Where aloetic purges are improper, the following pills may be used :

Take extract of jalap, and vitriolated tartar, of each two drachms; fyrup of ginger, as much as will make them of a proper confiftence for pills.

These pills may be taken in the same quantity as the above.

Pill for the Bile.

Take gum pill and colocynth pill, each a drachm. Beat them together, and make the mafs into thirty pills.

In bilious and nervous patients, where it was neceffary to keep the body gently open, I have found thefe pills answer the purpose extremely well. I generally give one over night, and another next morning, once or twice a-week. But the dose must be regulated by the effect.

Pill for the Jaundice.

Take of Castile foap, succotrine aloes, and rhubarb, of each one drachm. Make them into pills, with a sufficient quantity of fyrup or mucilage.

These pills, as their title expresses, are chiefly intended for the jaundice, which, with the affistance of proper diet, they will often cure. Five or fix of them may be taken twice a-day, more or less, as is necessary to keep the body open. It will be proper, however, during their use, to interpose now and then a vomit of ipecacuanha or tartar emetic.

Stomachic Pill.

Take extract of gentian, two drachms; powdered rhubarb and vitriolated tartar, of each one drachm; oil of mint, thirty drops; fimple fyrup, a fufficient quantity.

Three or four of these pills may be taken twice a-day, for inrigorating the stomach, and keeping the body gently open.

Squill Pills.

Take powder of dried fquills, a drachm and a half; gum ammoniac, and cardamom feeds, in powder, of each three drachms; fimple fyrup, a fufficient quantity.

In dropfical and afthmatic complaints, two or three of these pills may be taken twice a-day, or oftener, if the flomach will bear them.

Strengthening Pill.

Take foft extract of the bark, and falt of fteel, each a drachm. Make into pills.

683

In diforders arifing from exceflive debility, or relaxation of the folids, as the *cholorofis*, or green fickness, two of these pills may be taken three times a-day.

PLASTERS.

PLASTERS ought to be of a different confiftence, according to the purposes for which they are intended. Such as are to be applied to the breafts or flomach, ought to be foft and yielding; while those defigned for the limbs should be firm and adhesive.

It has been fuppofed, that plafters might be impregnated with the virtues of different vegetables, by boiling the recent vegetable with the oil employed for the composition of the plaster; but this treatment does not communicate to the oils any valuable qualities.

The calces of lead boiled with oils unite with them into a plafter of a proper confiftence, which make the bafis of feveral other plafters. In boiling these compositions, a quantity of hot water must be added from time to time to prevent the plaster from burning or growing black. This, however, should be done with care, left it cause the matter to explode.

Common Plaster.

Take of common olive oil, fix pints ; litharge, reduced to a fine powder, two pounds and a half. Boil the litharge and oil together over a gentle fire, continually flirring them, and keeping always about half a gallon of water in the veffel : after they have boiled about three hours, a little of the plafter may be taken out and put into cold water, to try if it be of a proper confiftence : when that is the cafe, the whole may be fuffered to cool, and the water well preffed out of it with the hands.

This plafter is generally applied in flight wounds and excoriations of the fkin. It keeps the part foft and warm, and defends it from the air, which is all that is neceffary in fuch cafes. Its principal ufe, however, is to ferve as a bafis for other plafters.

Adhesive Plaster.

Take of common plaster, half a pound; of Burgundy pitch, a quarter of a pound. Melt them together.

This plaster is principally used for keeping on other dreffings.

Anodyne Plaster.

Melt an ounce of adhefive plafter, and, when it is cooling, mix with it a drachm of powdered optum, and the fame quantity of camphor, previoufly rubbed up with a little oil.

Yy

PLASTERS.

This plaster generally gives ease in acute pains, especially of the nervous kind.

Bliftering Plafter.

Take of Venice turpentine, fix ounces; yellow wax, two ounces; Spanish flies in fine powder, three ounces; powdered mustard, one ounce. Melt the wax, and while it is warm, add to it the turpentine, taking care not to evaporate it by too much heat. After the turpentine and wax are fufficiently incorporated, sprinkle in the powders, continually flirring the mass till it be cold.

Though this plaster is made in a variety of ways, one feldom meets with it of a proper confistence. When compounded with oils and other greafy substances, its effects are blunted, and it is apt to run; while pitch and refin render it too hard and very inconvenient.

When the bliffering plaffer is not at hand, its place may be fupplied by mixing with any fost ointment a fufficient quantity of powdered flies; or by forming them into a passe with flour and vinegar.

Bl ffering plaffers prove highly difagreeable to many people, by occasioning frangury. I have therefore of late used a plaffer, in which a small quantity of bliffering falve has been mixed with the Burgundy pitch plaffer. I lay it over the part affected, and suffer it to remain as long as it will flick. The bliffering plaffer loses its effect in a tew hours, whereas this will act for many days, or even weeks, and feldom fails to remove pain, or flight obffructions.

Gum Plaster.

Take of the common plaster, tour pounds; gum ammoniac and galbanum, strained, of each half a pound. Melt them together, and add, of Venice turpentine, fix ounces.

This plaffer is used as a digestive, and likewise for discussing indolent tumours.

Mercurial Plaster.

Take of common plafter, one pound; of gum ammoniac, ftrained, half a pound. Melt them together, and, when cooling, add eight ounces of quickfilver, previoufly extinguished by triture, with three ounces of hog's lard.

This plaster is recommended in pains of the limbs arifing from a venereal cause. Indurations of the glands, and other violent tumours, are likewise found sometimes to yield to it.

Stomach Plaster.

Take of gum plaster, half a pound; camphorated oil, an ounce and a half; black pepper, or capficum, where it can be had,

PLASTERS.

had, one ounce. Melt the plafter, and mix with it the oil; then fprinkle in the pepper, previoufly reduced to a fine powder.

An ounce or two of this plafter, fpread upon foft leather, and applied to the region of the ftomach, will be of fervice in flatulencies arising from hyfteric and hypochondriac affections. A little of the expressed oil of mace, or a few drops of the effectial oil of mint, may be rubbed upon it before it is applied.

This may supply the place of the Antihysteric Plaster.

Warm Plaster.

Take of gum plaster, one ounce; bliftering plaster, two drachms. Melt them together over a gentle fire.

This plafter is useful in the friatica and other fixed pains of the rheumatic kind : it ought, however, to be worn for fome time, and to be renewed at least once a-week. If this is found to blifter the part, which is fometimes the case, it must be made with a smaller proportion of the bliftering plaster.

Wax Plaster.

Take of yellow wax, one pound; white refin, half a pound; mutton fuet, three quarters of a pound. Melt them together.

This is generally used instead of the Meillot Plaster. It is a proper application after blifters, and in other cases where a gentle digestive is necessary.

POULTICES.

THROUGH fome overfight, this article was omitted in the earlier editions, though it relates to a clafs of medicines by no means unimportant. Poultices are often beneficial, even in the most fimple form; but more fo, when employed to retain more active medicines,—to keep them in contact with the fk n, —and to fit it for their abforption. Every nurse knows how to make a poultice.

A poor woman who had received a very dangerous wound in the tendons of her thumb from a rufty nail, called upon me fome little time fince. As her cafe properly belonged to the department of furgery, I advifed her to apply to the holpital; but the official hirelings there refufed to take her in, though I always underftood that they were obliged to take in accidents. It feems, however, that fome very confined meaning was annexed to this word by the furgeon on duty, and that he did not think the danger of a locked jaw, to be an accident as deferving of his pity and immediate affiltance, as a broken arm, or diflocated ancle.

The poor woman came back to me; and, as her fituation became every moment more and more alarming, the pain and

Xy2

inflam-

inflammation having reached as high as the arm-pit, I advifed her to apply to the whole hand and arm a large poultice, with an ounce of laudanum fprinkled over it, and to renew the poultice twice a-day. This fhe did with fo much fuccefs, that the thumb is now quite well, though the accident did not happen above three weeks ago.

Alarming as the cafe was, I had fome reafon to rely on the efficacy of the poultice, from a former trial fomewhat fimilar. One of those girls, who are employed by bookbinders in flitching the fheets, having wounded her finger with the three-edged needle used on such occasions, foon felt the pain shoot upwards with deadly tendency. I ordered her to apply the same fort of poultice with laudanum, which had the same happy effect.

Both these patients made use of the Common Poultice; but I prefer one made of linseed flour, which is more easily prepared, and keeps moist longer than any other.

POWDERS.

T HIS is one of the most fimple forms in which medicine can be administered. Many medicinal substances, however, cannot be reduced into powder, and others are too difagreeable to be taken in this form.

The lighter powders may be mixed in any agreeable thin liquor, as tea or water-gruel. The more ponderous will require a more confiftent vehicle, as fyrup, conferve, jelly, or honey.

Gums, and other fubftances, which are difficult to powder, fhould be pounded along with the drier ones; but those which are too dry, especially aromatics, ought to be sprinkled during their pulverization with a few drops of any proper water.

Aromatic powders are to be prepared only in fmall quantities at a time, and kept in glafs veffels clofely ftopped. Indeed, no powders ought to be exposed to the air, or kept too long, otherwife their virtues will be in great measure destroyed.

Aftringent Powder.

Take of alum and Japan earth, each two drachms. Pound them together, and divide the whole into ten or twelve dofes.

In an immoderate flow of the *menfes*, and other hæmorrhages, one of these powders may be taken every hour, or every halfhour, if the discharge be violent.

Powder of Bole.

Take of Bole armenic, or French bole, two ounces; cinnamon, one ounce; tormentil root and gum arabic, of each fix drachms; drachms; long pepper, one drachm. Let all these ingredients be reduced into a powder.

This warm, glutinous aftringent powder, is given in fluxes, and other diforders where medicines of that class are neceffary, in the dofe of a fcruple, or half a drachm.

If a drachm of opium be added, it will make the Powder of Bole with Opium, which is a medicine of confiderable efficacy. It may be taken in the fame quantity as the former, but not above twice or thrice a day.

Carminative Powder.

Take of coriander feed, half an ounce; ginger, one drachm; nutmegs, half a drachm; fine fugar, a drachm and a half. Reduce them into powder for twelve dofes.

This powder is employed for expelling flatulencies arifing from indigeftion, particularly those to which hyfteric and hypochondriac perfons are fo liable. It may likewife be given in fmall quantities to children in their food, when troubled with gripes.

Diuretic Powder.

Take of gum arabic, four ounces; purified nitre, one ounce. Pound them together, and divide the whole into twenty-four dofes.

During the first stage of the venereal disease, one of these cooling powders may be taken three times a-day, with confiderable advantage.

Aromatic Opening Powder.

Take of the best Turkey rhubarb, cinnamon, and fine sugar, each two drachms. Let the ingredients be pounded, and atterwards mixed well together.

When flatulency is accompanied with coffivenels, a teafpoonful of this powder may be taken once or twice a-day, according to circumflances.

Saline Laxative Powder.

Take of soluble tartar, and cream of tartar, each one drachm; purified nitre, half a drachm. Make them into a powder.

In fevers, and other inflammatory diforders, where it is neceffary to keep the body gently open, one of these cooling laxative powders may be taken in a little gruel, and repeated occasionally.

Steel Powder.

Take filings of steel, and loaf-fugar, of each two ounces; ginger, two drachms. Pound them together.

Yy3

POWDERS.

In obstructions of the menses, and other cases where steel is proper, a tea-spoonful of this powder may be taken twice a-day, and washed down with a little wine or water.

Sudorific Powder.

Take purified nitre and vitriolated tartar, of each half an ounce; opium and ipecacuanha, of each one drachm. Mix the ingredients, and reduce them to a fine powder.

This is generally known by the name of *Dover's Powder*. It is a powerful fudorific. In obfinate rheumatifms, and other cafes where it is neceffary to excite a copious fweat, this powder may be administered in the dose of a foruple or half a drachm. Some patients will require two foruples. It ought to be accompanied with the plentiful use of fome warm diluting liquor.

This powder is emp

Take of tin reduced into a fine powder, an ounce; Æthiops mineral, two drachms. Mix them well together, and divide the whole into fix dofes.

One of these powders may be taken in a little fyrup, honey, or treacle, twice a day. After they have been all used, the following anthelmintic purge may be proper.

Purging Worm Powder.

Take of powdered rhubarb, a fcruple; fcammony and calomel, of each five grains. Rub them together in a mortar for one dofe.

For children the above dofes must be leffened according to their age.

If the powder of tin be given alone, its dole may be confiderably increased. The late Dr. Alfton gave it to the amount of two ounces in three days, and fays, when thus administered, that it proved an egregious anthelminuc. He purged his patients both before they took the powder and afterwards.

Powder for the Tape Worm.

Early in the morning the patient is to take in any liquid two or three drachms, according to his age and confficution, of the root of the male fenn reduced into a fine powder. About two hours afterwards, he is to take of calomel and refin of fcammony, each ten grains; gum gamboge, fix grains. Thefe ingredients muft be finely powdered and given in a little fyrup, honey, treacle, or any thing that is noft agreeable to the patient. He is then to walk gently about, now and then drinking a difh of weak green tea, till the worm is pafled. If the powder

POWDERS.

der of the fern produces nausea, or fickness, it may be removed by fucking the juice of an orange or lemon.

This medicine, which hid been long kept a fecret abroad for the cure of the tape-worm, was fome time ago purchafed by the French King, and made public for the benefit of mankind. Not having had an opportunity of trying it, 1 can fay nothing from experience concerning its efficacy. It feems, however, from its ingredients, to be an active medicine, and ought to be taken with care. The dofe here preferibed is fufficient for the ftrongeft patient; it must, therefore, be reduced according to the age and conflitution.

SYRUPS.

SYRUPS were fome time ago looked upon as medicines of confiderable value. They are at prefent, however, regarded chiefly as vehicles for medicines of greater efficacy, and are ufed for fweetening draughts, juleps, or mixtures; and for reducing the lighter powders into bolufes, pills, and electuaries. As all thefe purpotes may be answered by the finple fyrup alone, there is little occafion for any other; especially as they are feldom found but in a flate of fermentation; and as he dofe of any medicine given in this form is very uncertain. Perfons who ferve the public muft keep whatever their cufformers call for; but, to the private practitioner, nine-tenths of the fyrups ulually kept in the thops are unneceffary.

Simple Syrup

Is made by diffolving in water, either with or without heat, about double its weight of fine fugar.

If twenty-five drops of laudanum be added to an ounce of the fimple fyrup, it will supply the place of diacodium, or the fyrup of poppies, and will be tound a more fafe and certain medicine.

The lubricating virtues of the fyrup of marfhmallows may likewife be fupplied, by adding to the common tyrup a tufficient quantity of mucilage of gum arabic.

Those who choose to preferve the juice of lemons in form of fyrup, may diffolve in it, by the heat of a warm bath, nearly double its weight of fine fugar. The juice ought to be previoufly ftrained, and suffered to ftand till it fettles.

The fyrup of ginger is fometimes of ute as a warm vehicle for giving medicine to perfons afflicted with flatulency. It may be made by intufing two ounces of bruiled groger in two pints of boiling water for twenty-four hours. After the liquor has been flrained, and has flood to tettle for fome time, it may be poured off, and a little more than double its weight of fine powdered fugar diffolved in it.

Xy4

TINC

TINCTURES, ELIXIRS, &c.

R ECTIFIED fpirit is the direct menftruum of the refins and effential oils of vegetables, and totally extracts these active principles from fundry fubftances, which yields them to water, either not at all, or only in part.

It diffolves likewife those parts of animal subfrances in which their peculiar smells and tastes refide. Hence the tinctures prepared with rectified spirits form an useful and elegant class of medicines, possessing many of the most effential virtues of simples, without being clogged with their inert or useless parts.

Water, however, being the proper menstruum of the gummy, faline, and faccharine parts of medicinal substances, it will be neceffary, in the preparation of several tinctures, to make use of a weak spirit, or a composition of rectified spirit and water.

Aromatic Tincture.

Infuse two ounces of Jamaica pepper in two pints of brandy, without heat, for a few days; then strain off the tincture.

This fimple tinclure will fufficiently anfwer all the intentions of the more coftly preparations of this kind. It is rather too hot to be taken by itfelf; but is very proper for mixing with fuch medicines as might otherwise prove too cold for the ftomach.

Compound Tinsture of the Bark.

Take of Peruvian bark, two ounces; Seville orange-peel and cinnamon, of each half an ounce. Let the bark be powdered, and the other ingredients bruifed: then infuse the whole in a pint and a half of brandy, for five or fix days, in a close veffel; afterwards ftrain off the tincture.

This tincture is not only beneficial in intermitting fevers, but alfo in the flow, nervous, and putrid kinds, especially towards their decline.

The dofe is from one drachm to three or four, every fifth or fixth hour. It may be given in any fuitable liquor, and occafionally fharpened with a few drops of the fpirit of vitriol.

Volatile Fætid Tinsture.

Infuse two ounces of asafætida in one pint of volatile aromatic spirit, for eight days, in a close bottle, frequently fhaking it; then strain the tincture. This medicine is beneficial in hyfteric diforders, especially when attended with lowness of spirits, and faintings. A teaspoonful of it may be taken in a glass of wine, or a cup of penny-royal tea.

Volatile Tincture of Gum Guaiacum.

Take of gum guaiacum, four ounces; volatile aromatic spirit, a pint. Insuse without heat, in a vessel well stopped, for a few days; then strain off the tincture *.

In rheumatic complaints, a tea-spoonful of this tincure may be taken in a cup of the infusion of water trefoil, twice or thrice a-day.

Tincture of Black Hellebore.

Infuse two ounces of the roots of black hellebore, bruised, in a pint of proof spirit, for seven or eight days; then filter the tincture through paper. A scruple of cochineal may be infused along with the roots, to give the tincture a colour.

In obstructions of the *menses*, a tea-spoonful of this tincture may be taken in a cup of camomile or penny-royal tea twice a-day.

Aftringent Tinsture.

Digeft two ounces of gum kino, in a pint and a half of brandy, for eight days; afterwards ftrain it for use.

This tincture, though not generally known, is a good affringent medicine. With this view, an ounce, or more, of it may be taken three or four times a-day.

Tincture of Myrrh and Aloes.

Take of gum myrrh, an ounce and a half; hepatic aloes, one ounce. Let them be reduced to a powder, and infueed in two pints of rectified spirits, for fix days, in a gentle heat; then ftrain the tincture.

This is principally used by furgeons for cleanfing foul ulcers, and reftraining the progress of gangrenes. It is also, by some, recommended as a proper application to green wounds.

Tincture of Opium, or Liquid Laudanum.

Take of crude opium, two ounces; fpirituous aromatic water, and mountain wine, of each ten ounces. D folve the opium, fliced, in the wine, with a gentle heat, frequently flirring it; afterwards add the fpirit, and itrain off the tincture.

A very good tincture of guaiacum, for domeflic use, may be made by infusing two or three ounces of the gum in a bottle of rum or brandy.

TINCTURES, &c.

As twenty-five drops of this tinclure contain about a grain of opium, the common dole may be from twenty to thirty drops.

Sacred Tinsture, or Tinsture of Hiera Picra.

Take of fuccotorine aloes in powder, one ounce; Virginian fnake-root and ginger, of each two drachms. Infute in a pint of mountain wine, and half a pint of brandy, for a week, frequently fhaking the bottle, then firain off the tincture.

This is a fafe and uleful purge for perfons of a languid and phlegmatic habit: but is thought to have better effects, taken in fmall dofes as a laxative.

The dole, as a purge, is from one to two ounces.

Compound Tineture of Senna.

Take of fenna, one ounce; jalap, coriander feeds, and cream of tartar, of each half an ounce. Infule them in a pint and a half of French brandy for a week; then firain the tincture, and add to it four ounces of fine fugar.

This is an agreeable purge, and anfwers all the purpofes of the Elixir Salutis, and of Daffy's Elixir.

The dole is from one to two or three ounces.

to that a bak and

Tincture of Spanish Flies.

Take of Spanish flies, reduced to a fine powder, two ounces; spirit of wine, one pint. Infuse for two or three days; then ftrain off the tincture.

This is intended as an acrid flimulant for external use. Parts affected with the palfy or chronic rheumatism may be frequently rubbed with it.

TinEture of the Balfam of Tolu.

Take of the ballam of Tolu, an ounce and a balf; reclified fpirit of wine, a pint. Infuse in a gentle heat until the ballam is diffolved; then thrain the tincture.

This tincture possesses all the virtues of the balfam. In coughs, and other complaints of the breass, a tea-spoonful or two of it may be taken in a bit of loaf sugar. But the best way of using it is in syrup. An ounce of the tincture properly mixed with two pounds of simple syrup, will make what is commonly called the *Balfamic Syrup*.

Tindure of Rhubarb.

Take of rhubarb, two ounces and a half; leffer cardamon seeds, half an ounce; brandy, two pints. Digest for a week, and strain the tincture.

Thole

Those who choose to have a vinous tincture of rhubarb may infuse the above ingredients in a bottle of Lisbon wine, adding to it about two ounces of proof spirits.

It half an ounce of gentian root, and a drachm of Virginian fnake-root be added to the above ingredients, it will make the bitter tincture of thubarb.

All these tinctures are defigned as stomachies and corroborants as well as purgatives. In weakness of the stomach, indigestion, laxity of the intestines, sluxes, cholicky and such like complaints, they are frequently of great service. The dole is from half a spoonful to three or four spoonfuls or more, according to the circumstances of the patient, and the purposes it is intended to answer.

The Tonic Tinsture.

affect to two is

Mix two ounces of the compound tincture of Peruvian bark with the like quantity of the volatile tincture of Valerian; and of this mixture a tea (poonful in a glass of wine or water is to be taken three or four times a-day.

I have long made use of this tincture for the relief of those peculiar affections of the stomach and bowels, such as indigestion, &c. which generally accompany nervous difeases. I do not fay that the tincture will cure those complaints, nor do I know of any medicine that will; but where a complete cure cannot be rationally expected, relief is certainly a very defirable object.

Paregoric Elixir.

Take of flowers of benzoin, half an ounce; opium, two drachms. Intuie in one pound of the volatile aromatic spirit, for four or five days, frequently shaking the bottle; afterwards strain the elixir.

This is an agreeable and fafe way of administering opium. It eases pain, allays tickling coughs, relieves difficult breathing, and is useful in many diforders of children, particularly the hooping cough.

The dote to an adult is from fifty to a hundred drops.

Sacred Elixir.

Take of rhubarb cut fmall, ten drachms; fuccotorine aloes, in powder, fix drachms; leffer cardamon feeds, half an ounce; French brandy, two pints. Infute for two or three days, and then itrain the elixir.

I his uteful ftomachic purge may be taken from one ounce to an ounce and a half.

Stomachic

Stomachic Elixir.

Take of gentian root, two ounces; Curaffao oranges, one ounce; Virginian inake-root, half an ounce. Let the ingredients be bruifed, and infused for three or four days in two pints of French brandy; afterwards ftrain out the elixir.

This is an excellent ftomach bitter. In flatulencies, indigestion, want of appetite, and such like complaints, a small glass of it may be taken twice a-day. It likewise relieves the gout in the stomach, when taken in a large dose.

Acid Elixir of Vitriol.

Take of the aromatic tincture, one pint; oil of vitriol, three ounces. Mix them gradually, and after the fæces have fubfided, filter the elixir through paper, in a glafs funnel.

This is one of the beft medicines which I know for hyfteric and hypochondriac patients, afflicted with flatulencies arifing from relaxation or debility of the flomach and inteffines. It will fucceed where the most celebrated flomachic bitters have no effect. The dose is from ten to forty drops, in a glass of wine or water, or a cup of any bitter infusion, twice or thrice a-day. It should be taken when the flomach is most empty.

Camphorated Spirit of Wine.

Diffolve an ounce of camphor in a pint of rectified fpirits. This folution is chiefly employed as an embrocation in bruifes, palfies, the chronic rheumatifm, and for preventing gangrenes.

The above quantity of camphor, diffolved in half a pound of the volatile aromatic spirit, makes Ward's Essence.

Spirit of Mindererus.

Take of volatile fal ammoniac, any quantity. Pour on it gradually diffilled vinegar, till the effervetcence ceafes.

This medicine is uleful in promoting a difcharge both by the fkin and urinary paffage. It is also a good external application in ftrains and bruifes.

When intended to raife a fweat, half an ounce of it in a cup of warm gruel may be given to the patient in bed every hour till it has the defired effect.

VINEGARS.

VINEGAR is an acid produced from vinous liquors by a fecond fermentation. It is an useful medicine both in inflammatory and putrid diforders. Its effects are, to cool the blood, quench thirft, counteract a tendency to putrefaction, and allay inorinordinate motions of the fystem. It likewise promotes the natural secretions, and in some cases excites a copious sweat, where the warm medicines, called alexipharmic, tend rather to prevent that falutary evacuation.

Weaknefs, faintings, vomitings, and other hyfteric affections, are often relieved by vinegar applied to the mouth and nofe, or received into the flomach. It is of excellent use also in correcting many poisonous subflances, when taken into the flomach; and in promoting their expulsion, by the different emunctories, when received into the blood.

Vinegar is not only an useful medicine, but ferves likewife to extract, in tolerable perfection, the virtues of feveral other medicinal substances. Most of the odoriferous flowers impart to it their fragrance, together with a beautiful purplish or red colour. It also affists or coincides with the intention of squills, garlic, gum ammoniac, and several other valuable medicines.

These effects, however, are not to be expected from every thing that is fold under the name of vinegar, but from such as is found and well prepared.

The beft vinegars are those prepared from French wines.

It is neceffary for fome purpofes that the vinegar be diffilled; but as this operation requires a particular chemical apparatus, we fhall not infert it.

Vinegar of Litharge.

Take of litharge, half a pound; firong vinegar, two pints. Infuse them together in a moderate heat for three days, frequently shaking the vessel; then filter the liquor for use.

This medicine is little used, from a general notion of its being dangerous. There is reason, however, to believe, that the preparations of lead with vinegar are possessed of some valuable properties, and that they may be used in many cases with fastery and success.

A preparation of a fimilar nature with the above has of late been extelled by Goulard, a French furgeon, as a fafe and extenfively uteful medicine, which he calls the *Extract of Saturn*, and orders to be made in the following manner:

Take of litharge, one pound ; vinegar made of French wine, two pints. Put them together into a glazed earthen pipkin, and let them boil, or rather fimmer, for an hour, or an hour and a quarter, taking care to flir them all the while with a wooden (patula. After the whole has flood to fettle, pour off the liquor which is upon the top into bottles for ufe.

With this extract Goulard makes his vegeto-mineral water*,

which

VINEGARS.

which he recommends in a great variety of external diforders, as inflammations, burns, bruifes, fprains, ulcers, &c.

He likewise prepares with it a number of other forms of medicine, as poultices, plasters, ointments, powders, &c.

Vinegar of Roses.

Take of red roles, half a pound ; ftrong vinegar, half a gallon. Intule in a clole veffel for leveral weeks, in a gentle heat; and then ftrain off the liquor.

This is principally used as an embrocation for headachs, &c.

Vinegar of Squills.

Take of dried squills, two ounces; distilled vinegar, two pints. Infuse for ten days or a fortnight in a gentle degree of heat, afterwards strain off the liquor, and add to it about a twelfth part its quantity of proof spring.

This medicine has good effects in diforders of the breaft, occafioned by a load of vifcid phlegm. It is also of use in hydropic cafes for promoting a difcharge of urine.

The dole is from two drachms to two ounces, according to the intention for which it is given. When intended to act as a vomit, the dole ought to be large. In other cales, it must not only be exhibited in fmall doles, but also mixed with cinnamon water, or fome other agreeable aromatic liquor, to prevent the naulea it might other wife occasion.

WATERS BY INFUSION, &c.

Lime Water.

POUR two gallons of water gradually upon a pound of fresh burnt quicklime; and when the ebullition ceases, flir them well together; then suffer the whole to stand at rest, that the lime may settle, and afterwards filter the liquor through paper, which is to be kept in vessels closely stopt.

The lime-water from calcined oyster-shells, is prepared in the same manner.

Lime-water is principally used for the gravel; in which cafe, from a pint or two or more of it may be drank daily. Externally it is used for washing foul ulcers, and removing the itch, and other diseases of the skin.

Compound Lime Water,

Take fhavings of guaiacum wood, half a pound; liquorice root,

702

root, ore ounce; faffafras bark, half an ounce; coriander feeds, three drachms; fimple lime-water, fix pints.

Infuse without heat for two days, and then firain off the liquor.

In the fame manner may lime-water be impregnated with the virtues of the other vegetable fubftances. Such impregnation not only renders the water more agreeable to the palate, but alfo a more efficacious medicine, effectially in cutaneous diforders and foulnefs of the blood and juices.

It may be taken in the fame quantity as the fimple water.

Sublimate Water.

Diffolve eight grains of the corrofive fublimate in a pint of cinnamon water.

If a stronger solution be wanted, a double or triple quantity of sublimate may be used.

The principal intention of this is to cleanfe foul ulcers, and confume proud flefh.

Styptic Water.

Take of blue vitriol and alum, each an ounce and a half; water, one pint. Boil them until the falts are diffolved; then filter the liquor, and add to it a drachm of the oil of vitriol.

This water is used for stopping a bleeding at the nose, and other hæmorrhages; for which purpose cloths or dossils dipt in it must be applied to the part.

Tar-Water.

Pour a gallon of water on two pounds of Norway tar, and flir them ftrongly together with a wooden rod : after they have flood to fettle for two days, pour off the water for use.

Though tar-water falls greatly fhort of the character which has been given of it, yet it posseful for medicinal virtues. It fensibly raises the pulse, increases the secretions, and sometimes opens the body, or occasions vomiting.

A pint of it may be drank daily, or more, if the ftomach can bear it. It is generally ordered to be taken on an empty ftomach, viz. four ounces morning and evening, and the fame quantity about two hours after breakfast and dinner.

SIMPLE DISTILLED WATERS.

A GREAT number of diffilled waters were formerly kept in the fhops, and are flill retained in fome Difpenfatories. But we confider them chiefly in the light of grateful diluents, fuitable vehicles for medicines of greater efficacy, or for rendering difguftful ones more agreeable to the palate and flomach. We fhall therefore infert only a few of those which are best adapted to these intentions.

The management of a ftill being now generally underftood, it is needless to spend time in giving directions for that purpose.

Cinnamon Water.

Steep one pound of cinnamon bark, bruifed, in a gallon and a half of water, and one pint of brandy, for two days; and then diffil off one gallon.

This is an agreeable aromatic water, poffeffing in a high degree the fragrance and cordial-virtues of the fpice.

Pennyroyal Water.

Take of pennyroyal leaves, dried, a pound and a half; water, from a gallon and a half to two gallons. Draw off by diffillation one gallon.

This water poffesses, in a confiderable degree, the smell, taste, and virtues of the plant. It is given in mixtures and juleps to hysteric patients.

An infusion of the herb in boiling water answers nearly the fame purposes.

Peppermint Water.

This is made in the fame manner as the preceding.

Spearmint Water.

This may also be prepared in the fame way as the pennyroyal water.

Both these are useful flomachic waters, and will sometimes relieve vomiting, especially when it proceeds from indigestion, or cold viscid phlegm. They are likewise useful in some cholicky complaints, the gout in the stomach, &c. particularly the peppermint water.

An infusion of the fresh plant is frequently found to have the fame effects as the distilled water.

Rofe Water.

Take of roles, fresh gathered, fix pounds; water, two gallons. Distil off one gallon.

This water is principally valued on account of its fine flavour.

Jamaica Pepper Water.

Take of Jamaica pepper, half a pound ; water, a gallon and a half. Diftil off one gallon. W00 10

This is a very elegant diffilled water, and may in most cafes fupply the place of the more coftly fpice waters.

SPIRITUOUS DISTILLED WATERS.

Spirituous Cinnamon Water.

TAKE of cinnamon bark, one pound; proof fpirit, and common water, of each one gallon. Steep the cinnamon in the liquor for two days; then diftil off one gallon.

Spirituous Jamaica Pepper Water.

Take of Jamaica pepper, half a pound ; proof spirit, three gallons; water, two gallons. Diftil off three gallons.

This is a fufficiently agreeable cordial, and may fupply the place of the Aromatic Water.

WHEYS.

Alum Whey.

BOIL two drachms of powdered alum in a pint of milk till it is curdled; then ftrain out the whey.

This whey is beneficial in an immoderate flow of the menfes, and in a diabetes, or exceffive discharge of urine.

The dole is two, three, or four ounces, according as the ftomach will bear it, three times a day. If it fhould occafion vomiting, it may be diluted.

Muftard Whey.

Take milk and water, of each a pint; bruised mustard-feed, an ounce and a half. Boil them together till the cord is perfecily separated; afterwards strain the whey through a cloth.

This is the moft elegant, and by no means the leaft efficacious method of exhibiting muffard. It warms and invigorates the babit, and promotes the different fecretions. Hence, in the Zz

705

Budvien to stell wer

Serie Children

low

WHEYS.

low ftate of nervous fevers, it will often fupply the place of wine. It is also of use in the chronic rheumatism, palsy, dropsy, &c. The addition of a little sugar will render it more agreeable.

The dole is an ordinary tea-cup-full four or five times aday.

Scorbutic Whey.

This whey is made by boiling half a pint of the fcorbutic juices in a quart of cow's milk. More benefit, however, is to be expected from eating the plants, than from their expressed juices.

The fcorbutic plants are, bitter oranges, brooklime, garden fcurvy-grafs, and water-creffes.

A number of other wheys may be prepared nearly in the fame manner, as orange-whey, cream-of-tartar-whey, &c. Thefe are cooling pleafant drinks in fevers, and may be rendered cordial, when neceffary, by the addition of wine.

WINES.

THE effects of wine are, to raife the pulfe, promote perfpiration, warm the habit, and exhilarate the fpirits. The red wines, befides these effects, have an aftringent quality, by which they ftrengthen the tone of the flomach and intestines, and by this means prove serviceable in restraining immoderate fecretions.

The thin fharp wines have a different tendency. They pass off freely by the different emunctories, and gently open the body. The effects of the full-bodied wines are, however, much more durable than those of the thinner.

All fweet wines contain a glutinous fubftance, and do not pafs off freely. Hence they will heat the body more than an equal quantity of any other wine, though it fhould contain fully as much fpirit.

From the obvious qualities of wine, it must appear to be an excellent cordial medicine. Indeed, to fay the truth, it is worth all the reft put together.

But to answer this character, it must be sound and good. No benefit is to be expected from the common trash that is often fold by the name of wine, without possessing one drop of the juice of the grape. Perhaps no medicine is more rarely obtained genuine than wine.

Wine is not only used as a medicine, but is also employed as a menstruum for extracting the virtues of other medicinal subftances; for which it is not ill adapted, being a compound of water, inflammable spirit, and acid; by which means it is enabled

700

WINES.

abled to act upon vegetable and animal fubftances, and alfo to diffolve fome bodies of the metallic kind, fo as to impregnate itfelf with their virtues, as fteel, antimony, &c.

Anthelmintic Wine.

Take of rhubarb, half an ounce; worm-feed, an ounce. Bruife them, and infuse without heat in two pints of red port wine for a few days, then strain off the wine.

As the ftomachs of perfons afflicted with worms are always debilitated, red wine alone will often prove ferviceable. It muft, however, have ftill better effects when joined with bitter and purgative ingredients, as in the above form.

A glass of this wine may be taken twice or thrice a-day.

Antimonial Wine.

Take glass of antimony, reduced to a fine powder, half an ounce; Lisbon wine, eight ounces. Digest, without heat, for three or four days, now and then shaking the bottle; afterwards filter the wine through paper.

The dofe of this wine varies according to the intention. As an alterative and diaphoretic, it may be taken from ten to fifty or fixty drops. In a large dofe it generally proves cathartic, or excites vomiting.

Bitter Wine.

Take of gentian root, yellow rind of lemon-peel, fresh, each one ounce; long pepper, two drachms; mountain wine, two pints. Infuse without heat for a week, and strain out the wine for use.

In complaints arifing from weaknefs of the ftomach, or indigestion, a glass of this wine may be taken an hour before dinner and supper.

Ipecacuanha Wine.

Take of ipecacuanha, in powder, one ounce; mountain wine, a pint. Infuse for three or four days; then filter the tincture.

This is a fafe vomit, and answers extremely well for such perfons as cannot swallow the powder, or whose stomachs are too irritable to bear it.

The dole is from one ounce to an ounce and a half.

Chalybeate, or Steel Wine.

Take filings of iron, two ounces; cinnamon and mace, of each two drachms; Rhenith wine, two pints. Inluse for three or four weeks, trequently fnaking the bottle; then pass the wine through a filter.

In

In obstructions of the menses, this preparation of iron may be taken, in the dose of half a wine-glass twice or thrice a-day.

The medicine would probably be as good if made with Lifbon wine, fharpened with half an ounce of the cream of tartar, or a fmall quantity of the vitriolic acid.

Stomach Wine.

Take of Peruvian bark, großly powdered, an ounce; cardamom feeds, and orange-peel, bruifed, of each two drachms. Infuse in a bottle of white port or Lisbon wine for five or fix days; then strain off the wine.

This wine is not only of fervice in debility of the flomach and inteffines, but may alfo be taken as a preventive, by perfons liable to the intermittent fever, or who refide in places where this difeafe prevails. It will be of use likewife to those who recover flowly after fevers of any kind, as it affifts digeftion, and helps to reftore the tone and vigour of the fyftem.

an discretive and the sharely, it gas be taken Irous that to they

CONTRACTOR DATES

Land & Sammer

A glafs of it may be taken two or three times a-day,

is a three wount, and and so is a the will be furth the

tiges as consider in allow the angeoing an whole if anar is day too

The state of a state of the state of the state of the

et meinlange GLOSSARY.

Cople. A miley Guid Separated Vrom the aliment in the intef-

tines, and conveyed by the abforhents into the blood to

The motion of the blood, which is driven by the

Lavis. A spitennels of the A ... Vill

Circulation.

all A thaid which is fearered by the liver into the

an it blodder, and from the new pairs into the intellines,

A LTHOUGH terms of art have been feduloufly avoided in the composition of this treatife, it is impossible entirely to banish technical phrases when writing on medicine, a science that has been less generally attended to by mankind, and continues therefore to be more infected with the jargon of the schools, than perhaps any other. Several perfons having expressed their opinion that a glosfary would make this work more generally intelligible, the following concise explanation of the few terms of art that occur, has been added in compliance with their fentiments, and to fulfil the original intention of this treatife, by rendering it intelligible and useful to all ranks and classes of mankind.

Abdoman. The belly.

Abforbents. Vessels that convey the nourishment from the intestines, and the secreted fluids from the various cavities into the mass of blood.

by lame surines denominated critical days, heraus

compliants have need of crived to take a docifive chang

Acrimony. Corrofive fharpnefs.

Acute. A difeafe, the fymptoms of which are violent, and tend to a speedy termination, is called acute.

Adult. Of mature age,

Adust. Dry, warm.

Antispasmodic. Whatever tends to prevent or remove spalm. Aphthæ. Small whitish ulcers appearing in the mouth. Astriation. A tightening or lettening.

Atrabilarian. An epithet commonly applied to people of a certain temperament, marked by a dark complexion, black hair, fpare habit, &c. which the ancients fupposed to arife from the atra bilis, or the black bile.

ZZ3

Biles

Bile, or Gall. A fluid which is fecreted by the liver into the gall-bladder, and from thence paffes into the inteffines, in order to promote digeftion.

Cacochymie. An unhealthy ftate of the body.

Caries. A rottenness of the bone.

Chyle. A milky fluid feparated from the aliment in the inteftines, and conveyed by the abforbents into the blood to fupply the wafte of the animal body.

Chronic. A difease whose progress is flow, in opposition to acute.

Circulation. The motion of the blood, which is driven by the heart through the arteries, and returns by the veins.

Comatofe. Sleepy.

Conglobate Gland. A fimple gland.

Conglomerate. A compound gland.

Contagion. Infectious matter.

Cutis. The fkin.

Cutaneous. Of or belonging to the fkin.

Criss. A certain period in the progress of a disease, from whence a decided alteration either for the better or the worse takes place.

Critical. Decifive or important.

Critical Days. The fourth, fifth, feventh, ninth, eleventh, thirteenth, fourteenth, feventeenth, and twenty-firft, are by fome authors denominated critical days, because febrile complaints have been observed to take a decifive change at these periods.

D

Debility. Weaknefs.

Delirium. A temporary diforder of the mental faculties. Diaphragm. A membrane feparating the cavity of the cheft from that of the belly.

Diuretic. A medicine that promotes the fecretion of urine. Draftic. Is applied to fuch purgative medicines as are violent or harfh in their operation.

Empyema. A collection of purulent matter in the cavity of the break.

Endemic.

Endemic. A difease peculiar to a certain district. Epidemic. A difease generally infectious. Exacerbation. The increase of any disease.

F

Faces. Excrements.
Flatulent. Producing wind.
Factid. Emitting an offenfive fmell.
Factus. The child before birth, or when born before the proper period, is thus termed.
Fungus. Proud flefh.

Portanal! Madicines adapted D

Gangrene. Mortification. Ganglia. Gummata. Venereal excrescences. Gymnaflic. Exercise taken with a view to preferve or reftore health.—The ancient physicians reckoned this an important branch of medicine.

Hawi to un of vapour or fweH

Hæmorrhage. Difcharge of blood.

Hæmorrhoids. The piles.

Hettic Fever. A flow confuming fever, generally attending a bad habit of body, or fome incurable and deep-rooted difeafe.

Hypochondriacifm. Low spirits.

Hypochondriac viscera. The liver, spleen, &c. so termed from their situation in the hypochondriac or upper and lateral parts of the belly.

I

 Ichor. Thin bad matter.
 Impositume. A collection of purulent matter.
 Inflammation. A furcharge of blood, and an increased action of the vessels, in any particular part of the body.

this bad matter, d.I harged from in ill cond

Ligature. Bandage. Lixivium. Ley.

M

Miliary Eruption. Eruption of fmall puffules refembling the feeds of millet.

Morbific.

to the back bone.

Morbific. Caufing disease, or diseased.

Mucus. The matter discharged from the nose, lungs, &c. Mysentery. A double membrane which connects the intestines to the back bone.

Leas. Excrements.

L'angas. Proud Rella.

L'atalant. Producing wind.

their fituation in the

Nervous. Irritable.

Naufea. An inclination to vomit. Nodes. Enlargements of the bones produced by the venereal difeafe.

P

Pectoral. Medicines adapted to cure difeases of the breast. Pelvis. The bones situated at the lower part of the trunk;

thus named from their refembling in fome measure a bafon.

Pericardium. Membrane containing the heart.

Peritonaum. A membrane lining the cavity of the belly and covering the inteffines.

Perspiration. The matter discharged from the pores of the skin in form of vapour or sweat.

Phlegmatic. Watery, relaxed.

Phlogiston. Is here used to fignify somewhat rendering the air unfit for the purposes of respiration.

Plethoric. Replete with blood.

Polypus. A difeafed excrescence, or a substance formed of coagulable lymph, frequently found in the large blood vessels.

Pus. Matter contained in a boil.

R

Regimen. Regulation of diet.

Rectum. The straight gut, in which the fæces are contained. Respiration. The act of breathing.

S

Saliva. The fluid fecreted by the glands of the mouth. Sanies. A thin bad matter, difcharged from an ill conditioned fore.

Schirrous. A ftate of difeafed hardnefs.

Slough. A part separated and thrown off by suppuration.

Spasm. A diseased contraction.

Spine. The back bone.

la noile b

Styptic.

Styptic. A medicine for flopping the difcharge of blood. Syncope. A fainting fit attended with a complete abolition of fenfation and thought.

Tabes. A species of confumption.

Temperament. A peculiar habit of body, of which there are generally reckoned four, viz. the fanguine, the bilious, the melancholic, and the phlegmatic.

U


Vertigo. Giddinefs.

Ulcer. An ill-conditioned fore.

Ureters. Two long and fmall canals which convey the urine from the kidneys to the bladder.

Urethra. The canal which conveys the urine from the bladder.





INDEX.

A

ABLUTIONS, Jewish and Mahometan, well calculated for the prefervation of health, 99.

Abortion, caufes and fymptoms of, 491. Means of prevention, 492. Proper treatment in the cafe of, 493,

Abscesses, how to be treated, 291. 527.

Accidents. See Cafualties.

Acids, of peculiar fervice in confumptions, 175. In putrid fevers, 190. Not fuitable to the measles, 230.

Acidities in the bowels of infants, the origin of, 506. Method of cure, 507.

Ackwarth, foundling hofpital at, caufe of the children there being afflicted with (cabbed heads, and fatal confequences of their ill treatment, 511, note.

Addison, his remark on the luxury of the table, gt.

Æther, very ferviceable in removing fits of the althma, 385. Is excellent for flatulencies, 415.

Æthiops mineral, strongly recommended by Dr. Cheyne in inflammations of the eyes, 250.

Africans, their treatment of children, 10, note.

Agaric of the oak, its merit as a ftyptic, 530. Method of gathering, preparing, and applying it, *ibid. note.*

Agriculture, a healthful, conftant, and profitable employment, 45. Is too much neglected in favour of manufactures, *ibid*. Gardening the most wholefome amusement for fedentary persons, 49.

Ague, a species of fever no perfon can mistake, and the proper medicine for, generally known, 141. Causes of, 142. Symptoms, *ibid.* Regimen for, *ibid.* Under a proper regimen will often go off, without medicine, 144. Medical treatment of, 145. Often degenerates into obstinate chronical diseases, is not radically cured, 147. Peruvian bark the only medicine to be relied on in, 148. Children how to be treated in, *ibid.* Preventive medicine for those who live in marshy countries, 149.

Air, confined, poifonous to children, 28. A free open air will fometimes cure the most obstinate diforders in children, 30. Occupations which injure the health by unwholefome air, 35. 46. The qualities of, act more fensibly on the body than is generally imagined, 69. The feveral noxious qualities of, specified, *ibid*. In large cities, polluted by various circumstances, 70. The air in churches, how rendered unwholefome, *ibid*. Houses ought to be ventilated daily, 72. The danger attending small apartments, *ibid*. Perfons whose business confines them to town ought to sleep in the country, *ibid*. High walls obstruct the free current of air.

air, 73. Trees should not be planted too near to houses, ibid. Fresh air peculiarly necessary for the lick, ibid. The lick in hofpitals in more danger from the want of fresh air, than from their diforders, 74. Wholefomenels of the morning air, 78. The changeableness of, one great cause of catching cold, 120. Those who keep most within doors the most fensible of these changes, 121. Of the night, to be carefully guarded againft, 122. Fresh air often of more efficacy in difeases than medicine, 133. Its importance in fevers, 138. 153. States of, liable to produce putrid fevers, 186. Must be kept cool and fresh in fick chambers under this diforder, 190. Change of, one of the most effectual remedies for the hooping cough, 272. The qualities of, a material confideration for afthmatic patients, 384. The various ways by which it may be rendered noxious, 560. Confined, how to try and purify, ibid. Method of recovering perfons poifoned by foul air, 561. Fresh, of the greatest importance in fainting fits, 567.

Aitkin, Mr. his treatife on the nature and cure of fractures, recommended, 544, note.

Alcali, cauftic, recommended in the flone, 309. How to prepare it, ibid. note.

Aliment is capable of changing the whole conflitution of the body, 57. Will in many cases answer every intention in the cure of difeafes, ibid. The calls of hunger and thirft, fufficient for regulating the due quantity of, 58. The quality of, how injured, . ibid. A due mixture of vegetables necessary with our animal food, 60. To what the bad effects of tea are principally owing, - ibid. Water, good and bad, diftinguished, 61. Inquiry into the qualities of fermented liquors, with instructions for the due making of them, 63. The qualities of good bread, and why adulterated by bakers, 64. General rules for the choice of food, ibid. Ought not to be too uniform, 65. Meals ought to be taken at regular times, 66. Long fasting injurious both to old and young, 67. Breakfasts and suppers, ibid. Changes of diet ought to be gradually made, 68. General observations on, 621.

- Amaurofis. See Gutta ferena.

American Indians, their method of curing the venereal disease, 476. Amusements, sedentary, improper for sedentary persons, 48. Ought always to be of an active kind, 79.

Anafarca. See Dropfy.

Anger, violent fits of, injurious to the conflicution, 108. Tranquillity of mind effential to health, ibid.

Animal food, cautions in giving it to children, 17. Animals that die of themselves, unfit for food, 58. Over-driven cattle rendered unwholefome, 59. The artifices of butchers exposed, ibid. Too great a quantity of, generates the fcurvy, 60. Injurious to many habits, 621.

Animals and plants, the analogy in the nourifhment of, stated, 90. Anthony's fire, St. See Erysipelas.

6

Aphthe. See Thrush.

Apoplexy, who most liable to this disorder, 386. Causes, ibid. Symptoms and method of cure, 387. Gautions to perfons of an apoplectic make, 389. Apo-

Apothecaries' weights, a table of, 657.

Apparel. See Clothing.

Appetite, want of, caules and remedies for, 392,

Arbuthnot, Dr. his advice in the inflammation of the lungs, 16g. His advice for perfons troubled with coffivenels, 391, note.

Arsenic, the effects of, on the ftomach, 443. Medical treatment when the symptoms appear, 444,

Arts. See Manufactures.

Afcites. See Dropfy.

Affes milk, why it feldom does any good, 171. Instructions for taking it, ibid.

Aftbma, the different kinds of this diforder diffinguished, with Its causes, 382. Symptoms, 383. Regimen, *ibid*. Medical treatment, 384. Remedies proper in the moist afthma, 385. Atmosphere. See Air.

B

Ball's purging vermifuge powder, preparation of, 347.

Balfams, how to prepare. Anodyne balfam, 660. Locatelli's balfam, ibid. Vulnerary balfam, ibid.

Bandages, tight, produce most of the bad confequences attending fractured bones, 545.

Bark, Pernvian, the best antidote for failors against diforders on a foreign coaft, 44. How to be administered in the ague, 145. Diffinction between the red bark and quill bark, ibid. note. A decoction or infusion of, may be taken by those who cannot swallow it in substance, 146. Is often adulterated, ibid. note. Is the only medicine to be depended on in agues, 148. How it may be rendered more palatable, ibid. May be administered by clyfter, 149. Cold water the best menstruum for extracting the virtues of this drug, 176. How to be administered in the putrid fever, 192; and in the eryfipelas, 240. In an inflammation of the eyes, 249. Its efficacy in a malignant quinfey, 260. In the hooping-cough, 274. A good medicine in vomiting, when it proceeds from weaknels of the ftomach, 300. Its efficacy in a diabetes, and how to take it, 304. Is good against the piles, 317; and worms, 346. Its use dangerous for preventing a fit of the gout, 361. A good remedy in the King's evil, 376; and in the fluor albus, 489.

Barley water, how made, 159.

Barrenness in women, the general causes of, 501. Course of relief, ibid. Dr. Cheyne's observations on, ibid. note.

Bath waters, good in the gout, 362.

Bath, cold, the good effects of, on children, 27. Recommended to the fludious, 56. Is peculiarly excellent for ftrengthening the nervous fystem, 399. Therefore should never be omitted in gleets, 466. Is good for rickety children, 519. Apoplexies frequently occasioned by the improper use of, 578. Absurd prejudices against, *ibid*. Nothing contributes more to permanent health, 579. A fingle plunge or dip fufficient, continuing too long in it dangerous, *ibid*. Morning the properest time for bathing, *ibid*. Sea-water

preferable to river-water, 580. Rules to be observed in the use of, *ibid.* 581. Swellings of the glands resolved by sea-bathing, 581. Is peculiarly falutary in ferophulous complaints, 582. Warm bath better adapted for cutaneous diforders, *ibid.* Cautions on the improper use of, 583. 586. Good effects of the shower-bath, 587. Cold bath dangerous in the gout, 588. Warm bath more ialutary in discases of the liver and kidneys, 589. Observations on the vapour-bath, *ibid.*

Bath, warm, of great service in an inflammation of the flomach, 276. Bathing, a religious duty under the Judaic and Mahometan laws, 99. Is conducive to health, *ibid*.

Bears foot, recommended as a powerful remedy against worms, 346. Bear, instead of being made up again as soon as persons rife from

them, ought to be turned down and exposed to the air, 71. Bad effects of too great indulgence in bed, 78. Damp, the danger of, 122. Soft, are injurious to the kidneys, 303. 306.

Beer, the ill confequences of making it too weak, 63. Pernicious artifices of the dealers in, 64.

Bells, parish, the tolling them for the dead, a dangerous custom, 110. Biles, 528.

Bilious colic, fymptoms and treatment of, 282.

----- fever. See Fever.

Bite of a mad dog. See Dog.

Bitters, warm and altringent, antidotes to agues, 146. Are ferviceable in vomiting when it proceeds from weakness in the stomach, 300. Bladder, inflammation of, its general causes, 289. Medical treatment of, *ibid*.

-----, ftone in. See Stone.

Blaft. See Eryfipelas.

Bleeding, cautions for the operation of, in fevers, 139. In the ague, 144. Its importance in the acute continual fever, 154. In the pleurify, 159. When necessary in an inflammation of the lungs, 166. Caution against, in a nervous fever, 183. In the putrid fever, 191. In the miliary fever, 199. When necessary in the fmall-pox, 210. When useful in the measles, 231. When neceffary in the bilious fever, 235. Under what circumstances proper in the eryfipelas, 239. Mode of, proper in an inflammation of the brain, 244. Is always necessary in an inflammation of the eyes, 247. When proper and improper, in a cough, 267. When proper in the hooping-cough, 272. Is almost the only thing to be depended on in an inflammation of the ftomach, 276. And in an inflammation of the inteffines, 278. Is necessary in an inflammation of the kidneys, 287. Its ufe in a suppression of urine, 305. Is proper in an afthma, 384. Is dangerous in fainting fits, without due caution, 414. Cautions proper in the puerperal fever, 498. Is an operation generally performed by perfons who do not understand when it is proper, 524. In what cafes it ought to be had recourfe to, ibid. The quantity taken away, how to be regulated, 525. General rules for the operation, ibid. Objections to bleeding by leeches, 526. Prevailing prejudices relating to bleeding, ibid. The arm the most commodious part to take blood from; 527.

8

Bleeding

- Bleeding at the nofe, fpontaneous, is of more fervice, where bleeding is neceffary, than the operation with the lancet, 313. Ought not to be ftopped without due confideration, *ibid*. How to ftop it when neceffary, 314. Cautions to prevent frequent returns of, 315.
- Blind persons, when born so, might be educated to employments fuited to their capacity, 426, note.
- Blifters, peculiarly advantageous in the nervous fever, 183. When only to be applied in the putrid fever, 191. When proper in the miliary fever, 199. Seldom fail to remove the most obstinate inflammation of the eyes, 248. A good remedy in the quinfey, 255. Proper for a violent hooping-cough, 274. Is one of the best remedies for an inflammation of the stomach, 277. Are efficacious in the tooth-ach, 337.
- Blood, involuntary discharges of, often falutary, and ought not to be rashly stopped, 311. The several kinds of these discharges, with their usual causes, *ibid*. Methods of cure, 312.
- _____, fpitting of, who most subject to. and at what seasons, 320. Its causes, ibid. Symptoms, ibid. Proper regimen in, 321. Medical treatment, ibid. Cautions for persons subject to it, 322.

Blood-shot eye, how to cure, 429.

Bloody-flux. See Dyfentery.

Boerbaawe, his observations on dress, 88, note. His mechanical expedients to relieve an inflammation of the brain, 243.

- Bolufes, general rules for the preparing of, 661. The affringent bolus, *ibid*. Diaphoretic bolus, *ibid*. Mercurial bolus, *ibid*. Bolus of rhubarb and mercury, 662. Pectoral bolus, *ibid*. Purging bolus, *ibid*.
- Bones, the exfoliation of, a very flow operation, 535. Bones, broken, often fuccefsfully undertaken by ignorant operators, 543. Regimen to be adopted after the accident, *ibid*. Hints of conduct if the patient is confined to his bed, 544. Cleanlinefs to be regarded during this confinement, *ibid*. The limb not to be kept continually on the firetch, *ibid*. Cautions to be obferved in fetting a bone, *ibid*. Tight bandages condemned, 545. How to keep the limb fleady by an eafy method, *ibid*. Fractures of the ribs, *ibid*.

Bowels, inflammation of. See Stomach.

Boys, the military exercise proper for them, 24.

Braidwood, Mr. his skill in teaching the dumb to speak, 430, note.

Brain, inflammation of, who most liable to it, with its caufes and fymptoms, 241. Regimen, 243. Medical treatment, 244.

Bread, proper food for children, as foon as they can chew it, 16. A cruft of, the beft gum flick, *ibid*. The beft modes of preparing it in food for children, *ibid*. Good, the qualities of, and for what purpose adulterated by the bakers, 64. Toassed, a decoction of, good to check the vomiting in a *cholera morbus*, 294. A furfeit of, more dangerous than any other food, 623. The finess not always the best adapted for nutrition, 624. Hoasfehold bread the most wholesome, *ibid*. Wheat flour apt to occassion conflipation, 625. Different kinds of grain make the best bread, *ibid*. ConfumpConfemption of bread increased by drinking tea, 626. Indian corn the best food when boiled, 631. Buck wheat highly nutritious, *ibid*. Various substitutes for bread, 642.

Brimflone. See Sulphur.

Broth, gelatinous, recommended in the dyfentery, how to make, 327.

Broths and Soups, various ingredients in the composition of 641, 642. General observations.on, 643. The most wholesome diet for the poor, *ibid*. Warmly recommended by Count Rumford, *ibid*. Brown soup recommended for breakfast in place of tea, 646.

Bruifes, why of worfe confequence than wounds, 534. Proper treatment of, *ibid*. The exfoliation of injured bones a very flow operation, 535. How to cure fores occasioned by, *ibid*.

Buboes, two kinds of, diffinguished, with their proper treatment, 469.

Burdens, heavy, injurious to the lungs, 38.

Burgundy pitch, a plaster of, between the moulders, an excéllent remedy in a cough, 260. In a hooping-cough, 274. And for children in teething, 516.

Burials, the dangers attending their being allowed in the midst of populous towns, 70.

Burns, flight, how to cure, 532. Treatment of, when violent, 533. Extraordinary cafe of, *ibid*. Liniment for, 684.

Butchers, their professional artifices explained and condemned, 59. Butter ought to be very sparingly given to children, 19. Used in large quantities very injurious to the stomach, having a constant tendency to turn rancid, 632. Bread made with butter not easily digested, 634. Passies of every kind improper for children, *ibid*.

C

Cabbage leaves, topical application of, in a pleurify, 160. Campbor, why of little use in eye-waters, 665.

Camphorated oil, preparation of, 718.

Camphorated spirit of wine, 733.

Camps, the greatest necessity of confulting cleanlines in, 98.

Cancer, its different stages described, with the producing causes, 436. Symptoms, 437. Regimen, and medical treatment, 438. Dr. Storck's method of treating this diforder, 439. Cautions for avoiding it, 441. Credulity more incurable than cancers, 442.

Canter feroti, a diforder peculiar to chimney-fweepers, owing to want of cleanlinefs, 96, note,

Carriages, the indulgence of, a facrifice of health to vanity, 77. Carrot, wild, recommended in the flone, 309.

---- poultice for cancers, how to prepare, 440.

fubstances flopped in the gullet, 550.

_____ drowning, 556.

---- noxious vapours, 56c.

extremity of cold, 562.

- Cafualies,

Cafualties, extreme heat, 563.

Cataplasms, their general intention, 662. Preparation of the difcutient cataplasm, ibid. Ripening cataplasm, ibid.

Cataraci, the diforder and its proper treatment described, 428.

Cattle, stall fed, are unwholesome sood, 59. Over driven, are killed in a high fever, *ibid*. The artifices of butchers exposed, *ibid*. 60.

Cellars, long fhut, ought to be cautioufly opened, 72. And funk flories of houses, unhealthy to live in, 124. Of liquors in a flate of fermentation, dangerous to enter, 560. How to purify the air in, 561.

Celfus, his rules for the prefervation of health, 127.

Chancres, defcribed, 470. Primary, how to treat, 471. Symptomatic, ibid.

Charcoal fire, the danger of fleeping in the fume of, 560.

Charity, the proper exercise of, 41.

Cheefe, as a diet, injurious to health, 634.

Cheyne, Dr. his perfuafive to the use of exercise, 78. His judgment of the due quantity of urine, not to be relied on, 118. Strongly recommends Æthiops mineral in inflammations of the eyes, 250. His observations on barrennels, 501, note.

Chilblains, caufe of, 512. How to cure, 513.

Child bed women, how to be treated under a miliary fever, 200. Child-bed fever. See Fever.

- Child-birsh, the feafon of, requires due care after the labour pains are over, 493. Medical advice to women in labour, 494. Ill effects of collecting a number of women on fuch occasions, 495, note. How to guard against the miliary fever, 497. Symptoms of the puerperal fever, *ibid.* Proper treatment of this fever, 498. General cautions for women at this feafon, 500, 501.
- Children, their difeafes generally acute, and delay dangerous, 5-Their diforders lefs complicated, and eafier cured, than those of adults, 6. Are often the heirs of the difeafes of their parents, ibid. Those born of diseased parents, require peculiar care in the nurfing, 8. Are often killed or deformed by injudicious clothing, 9. How treated in Africa, 10, note. The usual causes of deformity in, explained, ibid. Their clothes ought to be fastened on with ftrings, 12. General rule for clothing them, 13. Cleanlinels, an important article in their drefs, 14. The milk of the mother the most natural food for, ibid. Absurdity of giving them drugs as their first food, 15. The best method of expelling the meconium, ibid. How they ought to be weaned from the breaft, 16. A cruit of bread the beft gum-flick for them, ibid. How to prepare bread in their food, ibid. Cautions as to giving them animal food, 17. Cautions as to the quantity of their food, ibid. Errors in the quality of their food more frequent than in the quantity, 18. The food of adults improper for children, ibid. Strong liquors expole them to inflammatory dilorders, ibid. Ill effects of unripe fruit, ibid. Butter, 19. Honey, a wholefome article of food for them, ibid. The importance of exercise to promote their growth and ftrength, 20. Rules for their exercife, 21. Poverty of parents occasions their neglect of children, 22. The utility of exercise demonstrated from the organical fructure of children,

ibid.

ibid. Philofophical arguments fhewing the neceffity of exercife, 23. Ought not to be fent to fchool too foon, *ibid.* Nor be put too foom to labour, 26. Dancing an excellent exercife for them, 27. The cold bath, *ibid.* Want of wholefome air deftructive to children, 28. To wrap them up clofe in cradles pernicious, 29. Are treated like plants in a hot-houfe, 30. The uforal faulty conduct of nurfes pointed out, 31. Are crammed with cordials by indolent nurfes, 32. Eruptions ignorantly treated by nurfes, *ibid.* Loofe ftools, the proper treatment of, 33. Every method ought to be taken to make them ftrong and hardy, 34. Indications of the fmall pox in, 204.

Chin cough. See Cough.

Cholera morbus, the diforder defined, with its caules and fymptoms, 293. Medical treatment, ibid.

Churches, the feveral circumstances that render the air in, unwholefome, 70.

Churching of women after lying in, a dangerous custom, 500.

- Church-yards, the bad confequences of having them in large towns, 70.
- Cities, large, the air in, contaminated by various means, 70. The bad effects of burying the dead in, *ibid*. Houles ought to be ventilated daily, 71. The danger attending fmall apartments, 72. All who can ought to fleep in the country, *ibid*. Diforders that large towns are peculiarly hurtful to, *ibid*. Cleanlinefs not fufficiently attended to in, 97. Should be fupplied with plenty of water, 100, note. The best means to guard against infection in, 104.
- Clare, Mr. his method of applying faline preparations of mercury in venereal cafes, 475.
- Cleanlinefs, an important article of attention in the drefs of children, 14; and to fedentary artifts, 48. Finery in drefs often covers dirt, 87. Is neceffary to health, 95. Diforders originating from the want of, 96. Is not fufficiently attended to in large towns, 97. Nor by country peafants, *ibid*. Great attention paid to, by the ancient Romans, *ibid*. note. Neceffity of confulting cleanlinefs in camps. 98. Was the principal object of the whole fyftem of the Jewifh laws, 99. Is a great part of the religion of the Eaftern countries, *ibid*. Bathing and wafhing greatly conducive to health, *ibid*. Cleanlinefs peculiarly neceffary on board of fhips, *ibid*. and to the fick, *ibid*. General remarks on, 100. Many diforders may be cured by cleanlinefs alone, 133. The want of, a very general caufe of putrid fevers, 187. Is a great prefervative againft venereal infection, 479; and againft galling in infants, 507. Cutaneous diforders proceed from a want of, 630.
- Clergy, exhorted to remove popular prejudices against inoculation, 219. Might do great good by undertaking the practice of it themselves, 223.
- Clothing, the only natural use of, 9. That of children, has become a secret art, 10. Ought to be fastened on infants with strings instead of pins, 12. Pernicious consequences of stays, 13. Importance of cleanlines to children, 14. The due quantity of, distated by the climate, 84. Should be increased in the decline of life, *ibid.* and adapted to the seafons, 85. Is often hurtful

by

by being made fubfervient to the purpoles of vanity, *ibid*. Pernicious confequences of attempting to mend the fhape by drefs, *ibid*. Stays, *ibid*. Shoes, 86. Garters, buckles, and other bandages, 87. The perfection of, to be eafy and clean, *ibid*. General remarks on, 88. Wet, the danger of, and how to guard against it, 121.

Clysters, a proper form of, for an inflammation of the flomach, 277. And for an inflammation of the inteflines, 278. Of tobacco fmoke, its efficacy in procuring a flool, 284, note. Of chicken broth, falutary in the cholera morbus, 294. Their use in suppreffion of urine, 305. Ought to be frequently administered in the puerperal fever, 498. Of tobacco, to excite a vomit, 553. Of tobacco sumes, to stimulate the intestines, 557. The general intention of, 663. Preparation of the emollient clyster, 664. Laxative clyster, *ibid*. Carminative clyster, *ibid*. Oily clyster, *ibid*. Starch clyster, *ibid*. Turpentine clyster, 665. Vinegar clyster, *ibid*.

Caliac paffion, proper treatment for, 331.

Coffee berries, recommended in the stone, 309.

Cold, extreme, its effects on the human frame, 562. The fudden application of heat dangerous in fuch cafes, *ibid*. How to recover frozen or benumbed limbs, 563.

Cold Bath. See Bath.

- Colds, frequently occafioned by imprudent changes of clothes at the first approaches of summer, 85, note. Various causes of, specified, 120. Their general causes, 262. Proper regimen on the symptoms of, appearing, *ibid*. Danger of neglecting the diforder, 263. The chief secret for avoiding, 265. Public gardens at the close of a hot day as dangerous as they are inviting, on account of, *ibid*. Hints to the frequenters of theatres and affemblies, to guard against, 266. Melancholy case of a young lady, who died of a cold contracted at an affembly, *ibid*.
- Colic, different species of, 281. Medical treatment of, according to their species and causes, 282. Bilious colic, *ibid*. Hysteric colic, 283. Nervous colic, 284. Cautions necessary to guard against the nervous colic, 285. General advice in colics, *ibid*.

Collyria. See Eye-waters.

- Commerce often imports infectious diforders, 103. Means suggested to guard against this danger, *ibid. note.*
- Confections often very needlessly compounded, 666. Preparation of the Japonic confection, ibid.
- Conferves and Preferves, general remarks on, and their composition, 677. Of red roles, ibid. Of floes, ibid. Candied orange-peel, 663.
- Constitution, good or bad, the foundation of, generally laid during infancy, 1.
- Confumptions, the increase of this diforder may be attributed to hard drinking, 93. Who most liable to, and its causes, 167. Symptoms, 168. Regimen, 169. Riding, *ibid*. Much benefit to be expected from going a long voyage, 170. Travelling, *ibid*. Diet, *ibid*. Great efficacy of milk in, 171. Medical treatment, 174.

Confump-

724

Confumptions, nervous, defined, and the perfons most liable to, 177. Proper treatment of, ibid.

Convulsions, why new-born infants are fo liable to, 11. Those preceding the eruption in the small-pox favourable symptoms, 207. The general causes of, 520. Proper treatment of, *ibid*. Extraordinary recovery of an infant seemingly killed by, 573. Farther, instructions in like cases, 574.

Cook, Captain, the circumnavigator, his means of preferving the health of his men, 43, note.

Cookery, the arts of, render many things unwholefome, that are not naturally fo, 61. Roafting meat, a walteful mode of, 644.

Cordials, ought not to be given to infants, 15. Are the common refuge of nurfes who neglect their duty to children, 32. Are often fatal in an inflammation of the flomach, 276. When good in the colic, 282. Ought not to be given to a pregnant woman during labour, 404.

Corn, damaged, will produce the putrid fever, 186.

Corns in the feet are occasioned by wearing tight shoes, 86. Cortex. See Bark.

Costiweness, a frequent recourse to medicines for the prevention of, injurious to the constitution, 117. Is rather to be removed by diet than by drugs, 118. Its general causes and ill effects, 390. Regimen, *ibid*. Remedies for, 392.

Cough, the proper remedies for, 267. A plaster of Burgandy pitch laid between the shoulders an excellent remedy for, 268. The stomach cough, and cough of the lungs, distinguished, 270. Treatment for the nervous cough, *ibid*.

Cough, hooping, who most liable to, with its disposing causes, 272. Remedies, 272. Is infectious, ibid. Vomits, their use, and how

to administer them to children, 272. Garlic ointment a good remedy for, 274.

Cough, phthifical, incident to federary artificers, from their breathing bonfined air, 46.

Cradles, on many accounts hurtful to children, 29.

Gramp, proper remedies for, 421.

Cramp of the flomach, who molt subject to, 410. Medical treatment of, ibid.

Crotchets, how to use for extracting fubstances detained in the gullet, 551.

Croup in children, defcribed, 513. Its fymptoms and proper treatment, ibid.

Cyder, the ill confequences of making it too weak, 63.

Dancing, an excellent kind of exercise for young persons, 27. Daucus sylvestris. See Carrot.

Deafnefs, when a favourable fymptom in the putrid fever, 188, nots. Methods of cure, according to its caufes, 431.

Methods of cure, according to its causes, 431. Death, the evidences of, sometimes fallacious, and ought not to be too soon credited, 549. 555. 559. 576. The means to be used used for the recovery of persons from, nearly the same in all cases, 575.

- Decoctions, general remarks on, 658. Preparation of the decoction of althæa, ibid. Common decoction, ibid. Of logwood, 669. Of bark, ibid. Compound decoction of bark, ibid. Of farfaparilla, ibid. Of feneka, 670. White decoction, ibid.
- Deformity, often occafioned by the injudicious manner of draffing children, 9. Is feldom found among favage nations, 10. The ufual caufes of, explained, *ibid*.

Dews, night, dangerous to health, 122.

- Diabetes, who most liable to this diforder, 301. Its caufes and fymptoms, 302. Regimen, and medical treatment, 303. Diftinguished from incontinency of urine, 304.
- Diarrhæa. See Loofenefs.
- Diet, will often answer all the indications of cure in diseases, 132. Il-Justrations, *ibid.* General observations on, 619. Improper diet affects the mind as well as the body, 622. Vegetable ought to preponderate over animal food, *ibid.* See Aliment.
- Digestion, the powers of, equally impaired by repletion or inanition, 66.
- Difeafes, hereditary, cautions to perfons afflicted with, 8. Peculiar diforders attending particular occupations, 36. Many of them infectious, 100. The knowledge of, depends more upon experience and observation than upon scientifical principles, 129. Are to be diffinguished by the most obvious and permanent symptoms, *ibid.* The differences of fex, age, and constitution to be considered, 130. Of the mind, to be diffinguished from those of the body, *ibid.* Climate, fituation, and occupation, to be attended to, *ibid.* Other collateral circumstances, *ibid.* Many indications of cure, to be answered by diet alone, 131. Cures often effected by fresh air, by exercise, or by cleanlinets, 133. Nervous difeases, of a complicated nature, and difficult to cure, 395.
- Diflocations, fhould be reduced before the fwelling and inflammation come on, and how, 538. Of the jaw, 539. Of the neck, *ibid*. Of the ribs, 540. Of the fhoulder, 541. Of the elbow, *ibid*. Of the thigh, 542. Of the knees, ancles, and toes, *ibid*.
- Diuretic infusion for the Dropsy, how to prepare, 354, 355.
- Dog, fymptoms of madnefs in, 446. Ought to be carefully preferved after biting any perfon, to afcertain whether he is mad or not, *ibid.* Is often reputed mad when he is not fo, 447. Symptoms of the bite of a mad dog, *ibid.* The poifon cannot lie many years dormant in the body, as is fuppofed, 448. Dr. Mead's receipt for the bite, *ibid.* The famous East Indian specific for, 449. Other recipes, *ibid.* Vinegar of confiderable fervice in this diforder, *ibid.* Medical courfe of treatment, recommended, *ibid.* Regimen, 450. Dipping in the fea not to be relied on, 451. Dr. Tiffot's medical courfe for the cure of the hydrophobia, 452. Remarks on the Ormskirk medicine, *ibid. note.*
- Doses of medicines, the relative proportions of, for different ages, 687.
- Drams ought to be avoided by perfons afflicted with nervous diforders, 397.

Draught,

Draught, is the proper form for fuch medicines as are intended for immediate operation, 670. How to prepare the anodyne draught, *ibid.* Diuretic draught, *ibid.* Purging draught, 671. Sweating draught, *ibid.* Vomiting draught, *ibid.*

Drefs. See Clothing.

- Drinking, perfons who are feldom intoxicated may neverthelefs injure their conflictions by, 92. The habit of drinking frequently originates from misfortunes, 93. Frequently deftroys the powers of the mind, 94. Perfons often forced to it by miftaken holpitality, *ibid. note.* Leads to other vices, 95.
- Dropfy, the feveral diffinctions of, with its caufes, 352. Symptoms, ibid. Regimen, 353. Medical treatment, 354. Tapping a fafe and fimple operation, 356. Extraordinary efficacy of nitre in the cure of, 357.

Dropfy of the brain. See Water in the head.

Drowned perfons, ought not to be rafhly given up for dead, 555. Proper trials for the recovery of, *ibid*. Endeavours ought not to be suspended upon the first returns of life, 559. Success of the Amsterdam society for the recovery of, 574.

Drunkennefs. See Intoxication.

- Dumb persons may be taught to read, write, and discourse, 430, note.
- Dyfentery, where and when most prevalent, 325. Its causes and fymptoms, 326. Regimen, 327. Fruit, one of the best remedies for, 328. Proper drink for, 329. Medical treatment, *ibid*. Cautions to prevent a relapse, 330.

E

Ear, the feveral injuries it is liable to, 430. Deafnefs, medical treatment of, according to its caufes, 431. Ought not to be tampered with, 432.

- Ear-ach, its caufes, and proper treatment for, 339. How to drive infects out of, *ibid*.
- Education of children, should be begun at home by the parents, 23. note. That of girls hurtful to their constitution, 25.

Effluvia, putrid, will occasion the spotted fever, 186.

Electricity, beneficial in the pally, 404.

Electuaries, general rules for making, 671. Preparation of lenitive electuary, 672. Electuary for the dyfentery, *ibid*. For the epilepfy, *ibid*. For the gonorrhœa, *ibid*. Of the bark, 673. For the piles, *ibid*. For the palfy, *ibid*. For the rheumatifm, *ibid*.

Elixir, paregoric, how to prepare, 699. Sacred elixir, ibid. Stomachic elixir, 700. Acid elixir of vitriol, ibid.

Emulfions, their use, 673. Preparation of the common emulfion, 674. Atabic emulfion, *ibid*. Camphorated emulfion, *ibid*. Emulfion of gum ammoniac, *ibid*. Oily emulfion, *ibid*.

Engleman, Dr. his account of the German method of recovering perions from fainting fits, 567.

Entrails. See Intestines.

- Epilepfy, the diforder defined, 405. Its causes and symptoms, 406. Due regimen, 407. Medical treatment, ibid.
- Eruptions in children often free them from bad humours, but are miftaken and ill-treated by nurfes, 32. Ought never to be ftopped without proper advice, 33. In fevers, how to be treated, 183. 191. 197. 199. In the imall-pox, 206. 210, In children, the caufes of, 508. How to cure, 511.
- Erysipelas, a diforder incident to the laborious, 39. Its caufes explained, and who most subject to it, 236. Its symptoms, 237. Regimen, 238. Medical treatment, 240. The scorbutic erysipelas, *ibid*. Instructions for those who are subject to this diforder, *ibid*.
- Evacuations of the human body, the principal, specified, 116. By stool, ibid. Urine, 118. Perspiration, 120.
- Exercise, the importance of, to promote the growth and ftrength of children, 20. All young animals exert their organs of motion as foon as they are able, 21. The utility of, proved from anatomical confiderations, 22. And from philosophical deductions, 23. Military exercise recommended for boys. 24. Benefits of dancing, 27. Is better for fedentary perfons under lownefs of spirits than the tavern, 48. Gardening the best exercise for the fedentary, 49. Violent, ought not to be taken immediately after a full meal, 57. Is as neceffary as food for the prefervation of health, 76. Our love of activity an evidence of its utility, ibid, Indolence relaxes the folids, ibid. The indulgence of carriages as absurd as pernicious, 77. Is almost the only cure for glandular obstructions, ibid. Will prevent and remove those diforders that medicine cannot cure, 78. Is the best cure for complaints in the ftomach, 79. How to be taken within doors, when not to be done in the open air, ibid. Active fports better than fedentary amusements, 80. The golf a better exercise than cricket, ibid. note. Exercise should not be extended to fatigue, ibid. Is as neceffary for the mind as for the body, 113. Is often of more effi-cacy than any medicine whatever, 132. The best mode of taking it in a confumption, 169. Is of the greatest importance in a dropfy, 354. Muscular, for the gout, 361. Is neceffary for the afthmatic, 384. Is fuperior to all medicine in nervous diforders, 398. And in the palfy, 405. Is proper for pregnant women, unless they are of a very delicate texture, 493. Want of, the occafion of rickets in children, 518.
- Extracts, general rules for making; but are more conveniently purchafed ready made, 675.
- Eyes, inflammation of, its general caufes, 245. Symptoms, 246. Medical treatment, 247. How to be treated when it proceeds from a (crophulous habit, 249. Advice to those who are subject to this complaint, 250. Are subject to many difeases which are difficult to cure, 426. The means by which they are frequently injured, 427. General means of prevention, *ibid*. The several diforders of, with their medical treatment, 428.
- Eye-waters, general remarks on, and their principal intentions, (65. Collyrium of alum, *ibid*. Vitriolic collyrium, *ibid*. Collyrium of lead, 666.

Fainting-

Fainting-fits, how to cure, 412. 560. Cautions to persons subject to them, 568. Bleeding dangerous in, ibid.

Falling sickness. See Epilepsy.

- Fafting, long, injurious to those who labour hard, 40. Is hurtful both to old and young, 67.
- Fathers, culpably inattentive to the management of their children, 5-Their irregular lives often injure the conflictution of their children, 7.
- Fear, the influence of, very great, in occasioning and aggravating difeases, 108. Its various operations, 109.
- Feet, injured by wearing tight thoes, 86. The washing of, an agreeable article of cleanlines, 59. Wet, the danger of, 122. Bathing them in warm water, a good remedy in a cold, 264. And in the hooping-cough, 274.

Fermentation, the vapour of liquors in a state of, noxious, 560.

- Fevers, of a bad kind, often occafioned among labourers by poor living, 40. Frequently attack fedentary perfons after hard drinking, 48. Nervous, often the confequence of intenfe fludy, 53. Putrid and malignant, often occafioned by want of cleanlinefs, 96. The moft general caufes of, enumerated, 135. The diffinguifhing fymptoms of, *ibid*. The feveral fpecies of, *ibid*. Is an effort of nature, which ought to be affilted, 136. How this is to be done, 137. Cordials and fweetmeats improper in, 138. Frefh air of great importance in, *ibid*. The mind of the patient ought not to be alarmed with religious terrors, 139. Cautions as to bleeding and fweating in, *ibid*. Longings, the calls of nature, and deferve attention, *ibid*. Cautions to prevent a relapie, 140.
- Fever, acute continual, who most liable to, 151. Causes, ibid. Symptoms, ibid. Regimen, 152. Medical treatment, 154. Symptoms favourable and unfavourable, ibid. Regimen to be obferved during recovery, 155.
- Fever, bilious, general time of its appearance, 234. Proper treatment of, according to its fymptoms, 235.

Fever, intermitting. See Ague.

- Fever, miliary, from what the name derived, and its general appearances, 195. Who most liable to it, *ibid*. Causes, *ibid*. Symptoms, 197. Regimen, *ibid*. Account of a miliary fever at Strasburgh, 198, note. Proper medical treatment, 199. Cautions for avoiding this diforder, 200. How to prevent in child-bedwomen, 497.
- Fever, milk, how occasioned, 496. How to prevent, 500.
- Fever, nervous, why more common now than formerly, and who most liable to it, 180. Its caufes, *ibid*. Symptoms and proper regimen, *ibid*. Medical treatment, 182.
- Fever, puerperal, or childbed, the time of its attack, and fymptoms, 497. Medical treatment of, 498. Cautions for the prevention of this fever, 500.
- Feyer, putrid, is of a pestilential nature, and who most liable to it, 186. Its general causes, *ibid*. Symptoms of, 187. Other fevers may be converted to this, by improper treatment, *ibid*. Favourable

vourable and unfavourable symptoms of, 188. Regimen, 189. Medical treatment, 191. Cautions for the prevention of this diforder, 193. Very prevalent in the narrow lanes of London, or in the marshy grounds in its environs, 194. Extraordinary case of a young lady who was cured of this fever, 195.

Fever, remitting, derivation of its name, 201. Its caufes and fymptoms, ibid. Proper regimen, 202. Medical treatment, 203. Cautions for avoiding this fever, ibid.

Fever, scarlet, why so named, and its usual season of attack, 233, Proper treatment of, ibid. Is fometimes attended with putrid and malignant fymptoms, ibid. Medicines adapted to this flage of the malady, 234. Hiftory of a fever of this kind at Edinburgh. ibid. note.

Fever, fecondary, in the fmall-pox, proper treatment of, 213.

Flatulencies in the flomach, remedies against, 341. The feveral caufes of, 414. Medical treatment of, 415.

Flatulent colic, its caufes, and feat of the diforder, 281. Remedies for, ibid.

Floaver-de-luce, the yellow water, the root of, recommended for the tooth-ach, 337.

Fluor albus described, with its proper treatment, 489.

Fomentations, how to make and apply, 675. General intentions of, ibid. Anodyne fomentation, ibid. Aromatic fomentation, ibid. Common fomentation, ibid. Emollient fomentation, 676. Strengthening fomentation, ibid.

Food. See Aliment.

Forgiveness of injuries, ought to be practifed from a regard to our own health, 108.

Fractures. See Bones, broken.

Frozen limbs, how to recover, 563.

Fruit, unripe, very hurtful to children, 18. One of the best medicines, both for the prevention and cure of a dysentery, 328. Fruits occasional substitutes sor bread, 635. The plantain tree used for that purpose in warm climates, ibid. Yams, a proper substitute for bread in the Weft Indies, 636.

Funerals, the great number of vifitors attending them, dangerous to health, 101.

G

Galling, in infants, the caule and cure of, 507. Gangrene, proper treatment of, 528.

Gardening, a wholefome amufement for fedentary perfons, 49.

Gargles for the throat, how to make, 253. 256. 259. General intentions of, 676. Method of making the attenuating gargle, ibid, Common gargle, ibid. Detergent gargle, ibid. Emollient gargle, 677. For the mouth of infants in the thrush, 505.

Garlic ointment, a North British remedy for the hooping-cough, how to apply it, 274.

Generals of armies, how they ought to confult the health of the men they command, 41.

Gilders. See Miners.

Ginger, fyrup of, how to prepare, 695.

Girls,

Girls, the common mode of education prejudicial to their conflicution, 25. Means of rectifying it recommended, *ibid*.

Gleet, how occafioned, and its fymptoms, 465. Method of cure, ibid. Regimen, 466. Obstinate gleets cured by mercurial inunctions, 467. How to apply bougies, ibid.

- Glover, Mr. his course of treatment for the recovery of a hanged man, 572.
- Gonorrbæa, virulent, the nature of, and its fymptoms, 457. Regimen, 458. Medical treatment, 459. Is often cured by affringent injections, 460. Cooling purges always proper in, *ibid*. Mercury feldom neceffary in a gonorrhœa, 462. How to administer it when needful, 463.
- Goulard, Mr. preparation of his celebrated extract of Saturn, 734-His various applications of it, *ibid*.
- Gout, the general caufes of, 51. How to treat a loofenels occafioned by repelling it from the extremities, 207. The fources of this diforder, and its fymptoms, 357, 358. Regimen for, 359. Wool the best external application in, *ibid*. Why there are fo many noftrums for, 360. Proper medicines after the fit, 361. Proper regimen in the intervals between fits to keep off their returns, *ibid*. How to remove it from the nobler parts to the extremities, 363. General cautions to prevent danger by mittaking it for other diforders, *ibid*.
- Grain, various kinds of, when boiled, a good fubflitute for bread, 628. The culture of, the culture of man, as it promotes health and longevity, 648. No manufacture equal to the manufacture of grain, *ibid*. The fcarcity of, occasioned by the number of horfes, 649.

Gravel, how formed in the bladder, 119. How diffinguished from the flone, 306. Causes and symptoms, *ibid.* Regimen, 307. Medical treatment, *ibid.*

Green-ficknefs originates in indolence, 485.

Grief, its effects permanent, and often fatal, 112. Danger of the mind dwelling long upon one fubject, efpecially if of a difagreeable nature, *ibid*. The mind requires exercife as well as the body. 113. Innocent amufements not to be neglected, *ibid*. Is productive of nervous difeafes, 396.

Gripes in infants, proper treatment of, 507.

Guaiacum, gum, a good remedy for the quinfy, 255; and rheumatism, 366.

Gullet, how to remove fubftances detained in, 551. Cautions for the use of crotchets with this intention, *ibid*. Other mechanical expedients, 552. Treatment if the obfiruction cannot be removed, 554.

Gums of children, applications to, during teething, and how to cut them, 516.

Gutta ferena, proper treatment of, 428.

H

Hæmoptoe, spitting of blood. See Blood. Hæmorrhages. See Blood.

Harrowgate

Harrowgate water, an excellent medicine for expelling worms, 346. And for the jaundice, 351. In the fourvy, 367.

Heach-ach, the species of, distinguished, 332. Causes of, ibid. Regimen, 333. Medical treatment, 334.

Health of the people in general, a proper object of attention for the magiltrates, xv. Ought to be attended to in matrimonial contracts, 8. Is often laboured for after it is deflroyed, 51. Rules given by Celfus for the prefervation of, 127.

Heart-burn, the nature of this diforder, with its caules, and remedies for, 393.

Heat, extreme, how to recover perfons overcome by, 563.

Hemlock, a good remedy in the king's evil, 376. Is recommended by Dr. Storck, for the cure of cancers, 439.

Hemp feed, a decoction of, good in the jaundice, and how to prepare it, 351.

Hiccup, its causes and method of treatment, 408.

Hoffman, his rules for guarding child-bed women against the miliary fever, 467.

Honey, a wholefome article of food for children, 19. Is recommended in the flone, 310.

Hooping cough. See Cough.

Horfe-radifb, the chewing of, will reftore fenfibility to the organs of tafte when injured, 434.

- Hospitals, the want of fresh air in, more dangerous to the patients than their diforders, 75. Cleanliness peculiarly necessary in, 100. Often spread infection by being situated in the middle of populous towns, 104. How they might be rendered proper receptacles for the sick, 105. Particularly in infectious diforders, 106. The sick in, ought not to be crowded together, 208, note.
- Houses, inflead of contrivances to make them close and warm, ought to be regularly ventilated, 72. In marshy fituations unwholefome, 73. Ought to be built in a dry fituation, 124. Danger of inhabiting new-built houses before thoroughly dry, *ibid*. Are often rendered damp by unseafonable cleanlines, *ibid*. Are dangerous when kept too close and hot, 127.
- Husbandmen, the peculiar diforders they are exposed to, from the viciflitudes of the weather, 37.

Huxbam, Dr. recommends the study of the dietetic part of medicine, xiii.

Hydrocephalus. See Dropfy.

Hydrophobia, Dr. Tiffot's method of curing, 452.

Hydrops pectoris. See Dropfy.

Hypochondriac affections, frequently produced by intense study, 53. Their causes, and who most subject to them, 422. The general intentions of cure, *ibid*. Regimen, 423.

Hysterics, a diforder produced by the habitual use of tea, 61. General causes of, 418. Symptoms, *ibid.* Proper treatment of, 419. Regimen, *ibid.* Medicines adapted 10, 420.

Hysteric colic, symptoms and treatment of, 283.

I

Jails, why malignant fevers are often generated in them, 71. Often fpread fpread an infection by being fituated in the middle of populous towns. 104. Ought to be removed, 105.

Janin, M. his relation of the recovery of an overlaid infant, 571. And of a man who had hanged himfelf, 572.

Jaundice, the different flages of its appearance, with the caufes of this diforder, 348, 349. Symptom's and regimen, 349. Medical treatment, 350. Jejuits Bark. See Bark.

Jerus, the whole system of their laws tending to promote cleanlines, 98. 102.

Iliac passion, a particular kind of inflammation in the intestines, 277:

Imposibume in the breas, in confunctions, how to make it break inwardly, when not to be discharged by other means, 176. Imposthumes after the small-pox, proper treatment of, 215.

- Incontinency of urine, dillinguished from a diabetes, 304. Expedient for relief, ibid.
- Indigestion, is one consequence of intense study, 52. General causes, and remedies for, 392.
- Indolence, its bad effects on the conflicution, 76. Occasions glandular obstructions, 77. Ill confequences of too much indulgence in bed, 78. Is the parent of vice, 80. Is the general caufe of most mervous diforders, 424.
- Infancy, the foundation of a good or bad conflicution, generally laid in this feason of life, 1.
- Infants, nearly one half of those born in Great Britain die under twelve years of age, 1. Perifh mostly by art, 2. Ought not to be fuckled by delicate women, 3. Importance of their being nursed by their mothers, ibid. Otten lose their lives, or become deformed, by errors in clothing them, q. How the art of bandaging them became the province of the midwife, ibid. How treated in Africa, 10, note. Philosophical observations on their organical fructure, and on the caufes of deformity, 11. Why they to frequently die of convultions, 12. Why exposed to fevers, ibid. And colds, 13. Rules for their drefs, ibid. Their food, 14. Reflections on the many evils they are expoled to, 502. Why their first diforders are in their bowels, 503. How to cleanse their bowels, ibid. The meconium, 501. Thrush, 505. Acidities, 506. Gripes, ibid. Galling and exceriations, 507. Stoppage of the nofe, 508. Vomiting, ibid. Loofenefs, 509. Eruptions, 510. Scabbed heads, 511. Chilblains, 512. The croup, 513. Teething, 515. Rickets, 517. Convultions, 520. Water in the head, 521. How to recover infants feemingly dead, 567. Ought never to fleep in the fame beds with their mothers or nurfes, 571, note. Cafe of the recovery of an overlaid infant, 571. Cafe of an infant being feemingly killed by a ftrong convultion fit, and recovered, 573. See Children.
- Infection, the danger of, incurred by injudicious or unneceffary attendance on the fick, 100. And on funerals, 101. Is often communicated by clothes. 102. Is frequently imported, 103. Is fpread by hofpitals and jails being fituated in the middle of populous towns, 104. How to prevent infection in fick chambers, *ibid.* Phylicians liable to fpread infection, 103, note. In what refpects

INDEX.

telpects the fpreading of infection might be checked by the magiftrate, 105. Bleeding and purging increase the danger of, by debilitating the body, 194. Small-pox, 204. Of the imall-pox may be received again, 208, note.

Inflammations, how the laborious part of mankind expole themfelves to, 39. Proper treatment of, 527.

Inflammation of the bladder. See Bladder.

of the brain. See Brain.

of the inteffines. See Inteffines.

of the kidneys. See K dneys.

_____ of the liver. See Liver. _____ of the lungs. See Peripneumony.

of the ftomach. See Stomach.

_____ of the throat. See Quinfy.

of the womb. See Womb.

Infusions, advantages of, over decoctions, 677. How to obtain rich infutions from weak vegetables, ibid. Preparation of the bitter infusion, ibid. Infusion of the bark, 678. Infusion of carduus, ibid. Of linfeed, ibid. Of roles, ibid. Of tamarinds, and fenna, ibid. Spanish infusion, 679. For the palfy, ibid.

Inns, the great danger of meeting with damp beds in them, 122. The fheets in, how treated to fave washing, ibid.

Inoculation of the fmall-pox, more favourably received here than in neighbouring countries, 216. Cannot prove of general utility while kept in the hands of a few, ibid. No mystery in the procefs, ibid. May fafely be performed by parents or nurfes, 217. Various methods of doing it, ibid. The clergy exhorted to remove the prejudices against the operation, 219. Arguments cited from Dr. Mackenzie in favour of inoculation, ibid. note. Ought to be rendered universal, 221. Means of extending the practice of, ibid. Two obstacles to the progress of, stated, 223. Might be performed by clergymen, or by parents themfelves, 224. The proper seafons and age for performing it, 225. Will often mend the habit of body, 226. Necessary preparation and regimen for, ibid. Performed by the bramins or priefts of India, 227. Their example recommended to the clergy of Europe, ibid. Medical skill not necessary for the inoculation of the small-pox, ibid. Previous preparation of the body for inoculation not always neceffary, 228. A fmall premium from the flate to every mother who inoculated her own child, warmly recommended, ibid.

In/eds, when they creep into the ear, how to force them out, 339. Poifonous, the bites of, how to be treated, 454.

Intemperance, one great caufe of the difeafes of feamen, 42. The danger of, argued from the construction of the human body, 90. The analogy in the nourifhment of plants and animals, ibid. Is the abuse of natural passions, ibid. In diet, 91. In liquor and carnal pleafures, ibid. The bad confequence of, involve whole families, 92. Effects of drunkennefs on the conflictution, ibid. Perfons who feldom get drunk, may neverthelefs injure their conftitutions by drink, ibid. The habit of drinking frequently acquired under misfortunes, 93. Is peculiarly hurtful to young perfons, 94. Leads to all other vices, ibid.

Intermitting fever. See Ague.

Intestines, inflammation of, general causes from whence it proceeds, 277. The fymptoms, regimen, and medical treatment, 278. Cautions to guard against it, 280.

- Intoxication produces a fever, 92. Fatal confequences of a daily repetition of this vice, *ibid.* Perfons who feldom get drunk, may neverthelefs injure their conflictation by drinking, *ibid.* Getting drunk a hazardous remedy for a cold, 263. Often produces fatal effects, 568. Proper cautions for treating perfons in liquor, 569. The fatest drink after a debauch, *ibid.* Remarkable cafe, 570.
- Jobnson, Dr. extraordinary cure of an infant seemingly killed by a firong convulsion fit, related by, 573.
- Ifues, how to make them take the best effect, 483.
- Itch, the nature and fymptoms of this difeafe defcribed, 379, 380. Sulphur, the best remedy against, 380. Great danger of the injudicious use of mercurial preparations for, 381. Cleanliness the best prefervative against, 382, note.
- Juleps, the form of, explained, 679. Preparation of the camphorated julep, *ibid*. Cordial julep, 680. Expectorating julep, *ibid*. Musk julep, *ibid*. Saline julep, *ibid*. Vomiting julep, *ibid*.

K

Kermes Mineral, recommended by Dr. Duplanil for the hoopingcough, 273, note.

Kidneys, inflammation of, its general causes, 286. Its symptoms and proper regimen, *ibid*. Medical treatment, 287. Where it proceeds from the stone and gravel, *ibid*. Cautions for those subject to this disorder, *ibid*. See Gravel. King's Evil. See Scropbula.

L

Laborious employments, the peculiar diforders incident to, 37. The folly of men emulating each other in trials of ftrength, 38. Difadvantages attending their diet, 39. How they expose themfelves to inflammations, *ibid*. Danger of fleeping in the fun, 40. Long fasting hurtful to them, *ibid*. Injuries arising from poor living, *ibid*. Many of the difeases of labourers, not only occafioned, but aggravated, by poverty, 41.

Labour should not be imposed too early on children, 26.

- Labour in child-bed, medical advice for, 494. Inconveniences of collecting a number of women at, 495, note.
- Laudanum, its efficacy in fits of an ague, 142, note. How to be administered in a cholera morbus, 294. In a loofenefs, 297. In a diabetes, 304. When proper for the head-ach, 335. How to apply, for the tooth-ach, 337. Will ease pain in the gout, 361. How to administer for the cramp in the stomach, 410. Is good for flatulencies, 415. Effects of an over-dose of, 470. Medical treatment in this case, 445.

Leading-firings injurious to young children, 21. Leaches, may be successfully applied to inflamed testicles, 468, note. And

And to disperse buboes, 469, note. Are proper to apply to children where inflammations appear in teething, 516. Objections to bleeding with them, 526.

Leeks, a proper ingredient in soups, 642.

Lemons. See Oranges.

Leprofy, why lefs frequent in this country now than formerly, 373. Requires the fame treatment as the feuryy, ibid.

Lientery, proper treatment for, 331.

- Life may frequently be reflored, when the appearances of it are fulpended by fudden cafualties, 549. 555. 559. 575.
- Lightning, perfons apparently killed by, might poffibly be recovered by the use of proper means, 575.
- Lime water, recommended to prevent gravel in the kidneys from degenerating to the stone in the bladder, 308. Is a good remedy for worms, 346. Happy effects of, in the cure of obstinate ulcers, 536.
- Lind, Dr. his prescription to abate fits of an ague, 142. note. His directions for the treatment of patients under putrid remitting fevers, 202, note.
- Liniment for burns, preparation of, 684. White liniment, 685. For the piles, ibid. Volatile liniment, ibid.
- Liquors, firong, expose children to inflammatory diforders, 18. Fermented, the qualities of, examined, 63. The bad confequences of making them too weak, *ibid*. Why all families ought to prepare their own liquors, *ibid*. Cold, the danger of drinking, when a perfon is hor, 125.

Liver, schirrous, produced by fedentary employments, 51.

- Liver, inflammation of, its causes and symptoms, 289, 290. Regimen and medical treatment, 290. Abscels in, how to be treated, 291. Cautions in the event of a schirrous being formed, *ibid.*
- Lobelia, an American plant used by the natives in the venereal difeafe, 476.

Lochia, a suppression of, how to be treated, 496.

- Longings, in diseases, are the calls of nature, and often point out what may be of real use, 140.
- Loofeness, habitual, general directions for perfons subject to, 117. Its general causes, 296. A periodical looseness ought never to be stopped, *ibid*. Medical treatment of, according to its various causes, 297. Means of checking it when necessary, 298. In children, proper treatment of, 509.
- Love, why perhaps the ftrongest of all the passions, 113. Is not rapid in its progress, and may therefore be guarded against at its commencement, 114. To pretend to it for amusement, cruelty to the object, *ibid*. Children often real martyrs between inclination and duty, *ibid*. note.
- Lues, confirmed, fymptoms of, 473. Mercury the only certain remedy known in Europe for this difeafe, 474. Saline preparations of mercury more efficacious than the ointment, 475. How to administer corrofive sublimate, *ibid*. American method of curing, 476. Mercury properly administered never fails to cure this difeafe, 483.

INDEX.

Lungs, injured by artifts working in bending poftures, 46. Studious perfons liable to confumptions of, 52.

Luxury, highly injurious to the organs of talke and fmell, 432.

Mackenzie, Dr. his arguments in favour of inoculating in the fmallpox, 219, note.

Mad dog. See Dog.

736

Magnessa alba, a remedy for the heart-burn, 394. Is the best medicine in all cafes of acidity, 507.

Magnets, artificial, their reputed virtue in the tooth-ach, 338. Malt liquots, hurtful in the afthma, 383. See Beer.

Man, why inferior to brutes in the management of his young, 1. Was never intended to be idle, 81.

Manufactures, the growth of, produced the rickets in children, 19. More favourable to riches than to health, 26. Some, injurious to health by confining artifts in unwholefome air, 35. Cautions to the workmen, 36. Compared with agriculture, 45. Are injujurious to health from artifts being crowded together, 46. And from their working in confined pollures, *ibid*. Cautions offered to fedentary artifts, 47. Sedentary arts better fuited to women than to men, 78, note.

Matrimony ought not to be contracted without a due attention to health and form, 7.

Mead, Dr. his famous recipe for the bite of a mad dog, 448. His character as a physician, *ibid. note.*

Meals ought to be taken at regular times, 66. Reafons for this uniformity, ibid.

Meastes, have great affinity with the small-pox, 229. Caufe and symptoms, *ibid*. Proper regimen and medicine, 230, 231. Inoculation of, might prove very falutary, 232, note.

Mechanics ought to employ their leifure hours in gardening, 50.

Meconium, the best mode of expelling it, 15. 504.

Medicine, the origin of the art of, xvi. The operation of, doubtful at beft, xvii. 1s made a mystery of, by its professors, xxiii. The fludy of, neglected by gentlemen, xxiv. This ignorance lays men open to pretenders, *ibid*. Ought to be generally understood, xxv. A diffusion of the knowledge of, would deftroy quackery, xxvii. Objections to the cultivation of medical knowledge answered, xxviii. The theory of, can never supply the want of experience and observation, 129.

Medicines have more virtue attributed to them than they deferve, 131. Ought not to be administered by the ignorant, nor without caution, 133. Want of perfeverance in the use of, one reason why chronic diseases are so feldom cured, 367. Many retained, which owe their reputation to credulity, 653. Are multiplied and compounded in proportion to ignorance of the causes and nature of diseases, *ibid*. Disadvantages of compounded medicires, *ibid*. Are often adulterated for the sake of colour, *ibid*. The relative proportions of doses of, for different ages, 656. A lift of such medical preparations as ought to be kept for private practice, 658.

I

Melancholy.

Melancholy, religious, its effects, 115. Leads to fuicide, *ibid*. Defined, with its caufes, 400. Symptoms and regimen, *ibid*. 401. Medicinal treatment, 402.

Menstrual discharge in women, the commencement and decline of, the most critical period of their lives, 484. Confinement injurious to growing young women, 485; and tight lacing for a fine shape, 486. Symptoms of the first appearance of this discharge, 487. Objects of attention in regimen at this time, *ibid*. Ought to be reflored whenever unnaturally obstructed, and how, *ibid*. When an obstruction proceeds from another malady, the first cause is to be removed, 488. Treatment under a redundancy of the discharge, 489. Regimen and medicine proper at the final decline of the menses, 490.

Mercury may be given in desperate cases of an inflammation of the intellines, 230. Cautions for administering it, *ibid. note.* Great caution neceffary in using mercurial preparations for the itch, 38t. Is feldom neceffary in a gonorrhœa, 46z. How to administer it when needful in that disorder, *ibid.* Solution of mercury, how to make, 463, note. Is the only certain remedy known in Europe for the cure of a confirmed lues, 474. Saline preparations of, more efficacious than the mercurial ointment, 475. How to administer corrofive sublimate in venereal cases, *ibid.* Neceffary cautions in the use of mercury, 477. Proper seasons for entering on a course of, 478. Preparations for, 479. Regimen under a course of, *ibid.*

Mezereon root, a powerful affiltant in venereal cafes, 476.

Midwifery, ought not to be allowed to be practifed by any woman not properly qualified, 494, note.

Midwives, historical view of the profession, 9. How they became intrusted with the care of bandaging infants, with the ill effects of their attempts at dexterity in this office, 10. Instances of their rashness and officious ignorance, 499, note. 502, note.

Miliary Fever. See Fewer.

Military exercife recommended for boys to practife, 24.

Milk, that of the mother, the most natural food for an infant, 14. Cow's milk, better unboiled than boiled, 16. Is a good antidote against the feurvy, 60. Of more value in confumptions than the whole Materia Medica, 171. Its great efficacy in the feurvy, 371. A milk diet proper both for men and women, in cafes of barrennefs, 501.

Milk fever. See Fever.

Millipedes, how to administer for the hooping-cough, 270.

Mind, difeases of, to be diffinguished from those of the body, 130. See Paffions.

Miners, exposed to injuries from unwholesome air, and mineral particles, 36. Cautions to, ibid.

Mineral waters, the danger of drinking them in too large quantities, 302. Are of confiderable fervice in weakneffes of the ftomach, 393. Cautions to be observed in the use of, 589-619. Effays and pamphlets on watering-places, uncertain guides, 589. A knowledge of the contents of any mineral water, no criterion for its medical properties, 590. Dr. Falconer's opinion on this subject, 591. Cautions in the use of foreign springs, 592. Mineral

3 B

waters

238

waters classed according to their fenfible qualities, and their medical properties defined, *ibid.* 1. Simple Chalybeates, Tunbridge Wells, 593. 595. Islington and Hampstead, 596; German Spa and Pyrmont, *ibid.* 598. 2. Saline, or purging Chalybeates, Cheltenham, 599, 600; Scarborough, 600; Epsom, Bagnigge Wells, Dog and Duck St. George's Fields, Kilburne, Acton, 601; Sedlitz, in Bohemia, 602; Seltzer, *ibid.* 603; Harrowgate, *ibid.* 607; Moffat, *ibid.* 608. 3. Calcareous; Bath, 609. 613; Buxton, 613, 614; Briftol, 615; Aix-la-Chapelle, 608. 6.6; Bareges, 608. 617: Auvergne and the Bourbonnois, 618; Bourbon Laney, *ibid.* Carlfbad, *ibid.* 619.

Mixtures, general remarks on this form of medicines, 681. Composition of the aftringent mixture, *ibid*. Aftringent mixture, *ibid*. Diuretic mixture, *ibid*. Laxative absorbent mixture, *ibid*. Saline mixture, 682. Squill mixture, *ibid*.

Molasses, an intoxicating spirit much used by the common people at Euinburgh, 93, note.

Mothers, preputierous, when they think it below them to nurfe their own children, 2. Under what circumftances they may be really unit to perform this tafk, *ibid*. Importance of their fuckling their cwn children, 3, note. Delicate mothers produce unhealthy flort-lived children, 7. Their milk, the beft food for children, 14. Ought to give their children proper exercife, 22; and air, 28.

Mouth, cautions against putting pins or other dangerous articles into, 551.

Muscular exercise, a cure for the gout, 362.

Mushrooms, a dangerous article of food, as other fungules are often gathered instead of them, 455.

Music, the performance of, recommended as a proper amusement for itudious persons, 55.

Muse, extraordinary effects produced by, under particular circumftances, in the nervous fever, 184. Is a good remedy in the epilepfy, 408. And for the hiccup, 4c9. Deafnefs cured by, 431. Mustard, white, a good remedy in the rheumatism, 367.

proper bull for men and woured, in cafes of barron-

Natural history, the study of, necessary to the improvement of agriculture, xxii.

Nervous difeales, the most complicated and difficult to cure of all others, 395. General caules of, *ibid*. Symptoms of, 395. Regimen, 397. Medical treatment, 399. A cure only to be expected from regimen, 400. The feveral fymptoms of, though differently named, all modifications of the fame difease, 422. Remarkable cafe in, 425.

Nervous colic, its caufes and fymptoms, 284. Medical treatment of, ibid.

Nervous tever. See Fever.

\$1073.W

Night-mare, nits causes and symptoms described, 411. Proper treat-

Night-shade, an infusion of, recommended in a cancer, 439.

Nitre,

Nitre, purified, its good effects in a quinfey, 254. Is an efficacious remedy for the dropfy, 355. Promotes urine and perfpiration, 402.

Nose, ulcer in, how to cure, 433. Stoppage of, in children, how to cure, 508. See Bleeding at.

Nurfes, their fuperfittious prejudices in bathing of children, 27. The only certain evidence of a good one, 31. Their ufual faults pointed out, *ibid*. Administer cordials to remedy their neglect of duty towards children, 32. Their mistaken treatment of eruptions, *ibid*. And loofe stools, 33. Are apt to conceal the diforders of children that arife from their own negligence, *ibid*. Ought to be punished for the missfortunes they thus occasion, *ibid*. Sensible, often able to discover difeases sooner than perfons hred to physic, 129. Are liable to catch the small-pox again from those they nurfe in that diforder, 208, note.

Nurfery ought to be the largest and best aired room in a house, 29.

0

Oats and barley, more wholefome when boiled, 628.

Oatmeal, frequently made into bread, and peafants fed on it are healthy and robuft, 629. Erroneoufly supposed to occasion skin difeases, 630. Oatmeal and milk a proper food for children, *ibid.*

Oil, an antidote to the injuries arifing from working in mines or metals, 37. Salad oil, the best application to the bite of a wasp or bee, 453. Camphorated, how to prepare, 685.

Oils, effential, of vegetables, the proper menthruum for, 696.

Ointment for the itch, 380. Use of ointments when applied to wounds and fores, 632. Preparation of yellow bafilicum, *ibid*. Emollient ointment, 683. Eye ointments, *ibid*. Iffue ointment, *ibid*. Ointment of lead, *ibid*. Mercurial ointment, 684. Ointment of fulphur, *ibid*. Ointment for discases of the skin, *ibid*. White ointment, *ibid*. See Liniment.

Onions roasted, used by the Turks as bread, 642. Opthalmia. See Eye.

Opiates, efficacious in a cholera morbus, 294. In a diabetes, 304. When proper for the head-ach, 335. Recommended for the tooth-ach, *ibid*. Are delufive remedies in nervous difeafes, 399. See Laudanum.

Orange and lemon peel, how to candy, 668. How to preferve orange and lemon juice in the form of fyrup, 695.

Ormskirk medicine for the bite of a mad dog, remarks on, 452, note.

Oxycrate, the most proper external application in a fracture, 546. Oysters of great service in confumptions, 173, note.

Painters. Se Miners.

Palfy, the nature of this diforder explained, with its causes, 403. Medical treatment, 404.

Paraphrenitis, its symptoms and treatment, 163. ,

3 B 2

Parents,

Parents, their interested views in the disposal of their children in marriage, often a source of bitter repentance, 114, note. See Fathers and Mothers.

Parsnips, a nourishing food, 641.

740

Passions, intemperance the abuse of, 90. Have great influence both in the cause and cure of diseases, 107. Anger, 108. Fear, *ibid*. Grief, 112. Love, 113. Religious melancholy, 115. The best method of counteracting the violence of any of the passions, *ibid*.

Peas, parched, good in cafes of flatulency, 341.

Peruvian bark. See Bark.

Penis, ulcerated, cured, and partly regenerated, by a careful attention to cleanlinefs, 480, note.

Peripneumony, who most subject to, 164. Its causes, symptoms, and proper regimen, ibid. 165.

Perspiration, infenfible, the obstruction of, diforders the whole frame, 120. The various causes of its being checked, *ibid*. Changes in the atmosphere, *ibid*. Wet clothes, 121. Wet feet, 122. Night air, *ibid*. Damp beds, *ibid*. Damp houses, 124. Sudden tranfitions from heat to cold, 125.

Philosophy, advantages refulting from the fludy of, xxiv.

Phrenitis. See Brain.

Phymofis described, and how to treat, 472.

Physicians, ill confequences of their inattention to the management of children, 5. Are liable to spread infection, 105, note. Their custom of prognosticating the fate of their patients, a bad practice, 110. Are feldom called in until medicine can give no relief, 152. Fears, prejudices, and opposite interests, are obstacles to every falutary discovery, 216.

Pickles, provocatives injurious to the flomach, 61.

- Piles, bleeding and blind, the diffinction between, 316. Who moft fubject to the diforder, *ibid*. General caufes, *ibid*. Medical treatment of, *ibid*. Periodical difcharges of, ought not to be ftopped, 317. Proper treatment of the blind piles, *ibid*. External ointments of little ufe, 318.
- Pills, purging, proper form of, for an inflammation of the inteftines, 279. The general intention of this class of medicines, 685.
 Preparation of the composing pill, 686. Deobstruent pill, *ibid*.
 Fœtid pill, *ibid*. Hemlock pill, *ibid*. Mercurial pill, *ibid*.
 Mercurial fublimate pill, 687. Plummer's pill, *ibid*. Purging pills, *ibid*. Pill for the bile, 688. Pill for the jaundice, *ibid*.
 Stomachic pills, *ibid*. Squill pills, *ibid*. Strengthening pills, *ibid*.

Pins ought never to be used in the dreffing of children, 12. Swallowed, discharged from an ulcer in the fide, 551, note.

Plasters, the general intentions of, and their usual basis, 689. Preparation of the common plaster, *ibid*. Adhesive plaster, *ibid*. Anodyne plaster, *ibid*. Blistering plaster, 690. Gum plaster, *ibid*. Mercurial plaster, *ibid*. Stomach plaster, *ibid*. Warm plaster, 691. Wax plaster, *ibid*.

Pleurify, the nature of the diforder explained, with its causes, 157. Symptoms, 158. Regimen, *ibid*. Medical treatment, 159. A 6+ decoction

decection of seneka reckoned a specific in, 162. Bastard pleueffammerion of the inveltorify, 163. proper in a guadrinesa,

Plumbers. See Miners.

Poisons, the nature and cure of, a general concern, and eafily acquired, 442. Mineral poifons, 443. Vegetable poifons, 445. Bites of poilonous animals, 446. Bite of a mad dog, 447. Bite of a viper, 453. The practice of fucking the poilon out of wounds recommendea, ibid. note. Poisonous plants ought to be deftroyed in the neighbourhood of towns, 454. Negro remedy to cure the bite of a rattle-fnake, 455. General rules for fecurity against poisons, 456.

Poor living, the dangers of, 40.

Porters subject to diforders of the lungs, 38.

- Postures, confined, injurious to the health of fedentary artists, 46.
- Potator, the best fubstitute for bread, 636. The mealy kind deemed most nutritious, 637. Simply boiling or roasting, the best mode of cooking them, ibid. Stewed mutton and potatoes a nourifhing and palatable difh, ibid. The benefit of planting potatoes on fallows, in a letter from Sir John Methuen Poore to Sir William Pulteney, 638. His benevolent plan for promoting the culture of this root, among the poor cottagers in his parish, ibid. 639.
- Powerty, occasions parents to neglect giving their children proper exercife, 22. Deftroys parental affection, ibid. note. Not only occasions, but aggravates, many of the difeases of labourers, 41. The poor great fufferers by the fale of bad provisions, 58. And by bad air in large cities, 71.

Poultices proper for inflamed wounds, 532. General intentions of, 692. Extraordinary cafe of the good effects of, ibid.

- Powders, general instructions for making and administering, 602. Aftringent powder, preparation and doles of, ibid. Powder of bole, ibid. Carminative powder, 693. Diuretic powder, ibid. Aromatic purging powder, ibid. Saline laxative powder, ibid. Steel powder, ibid. Sudorific powder, 694. Worm powder, ibid. Purging worm powder, ibid. Powder for the tape-worm, ibid.
- Pox, fmall, who most liable to, and at what seasons, 204. Its causes and fymptoms, ibid. 205. Favourable and unfavourable fymptoms in, ibid. 206. Regimen, 206. How the patient ought to be treated during the eruptive fever, 207. Children in this diforder ought not to lie together in the fame bed, 208. Should be allowed clean linen, 209. Patients under this diforder ought not to appear in public view, ibid. Medical treatment, 210. The fecondary fever, 213. When and how to open the puffules, 214. Of inoculation, 215.
- Pregnancy, how to treat vomiting when the effect of, 299. Rules of conduct for women under the disorders incident to, 491. Causes and fymptoms of abortion, 492. How to guard against abortion, ibid. Treatment in cafes of abortion, 493. Childbirth, ibid.

Prescriptions, medical, patients exposed to danger by their being written in Latin, xxvii.

Provisions, unfound, the fale of, a public injury, 58.

Puerperal fever. See Fever.

Purges, the frequent taking of them renders the habitual use of them necessary, 117. Their efficacy in agues, 144. Proper form of, for

741

an inflammation of the intestines. 279. Cooling purges always proper in a gonorrhœa, 460. Midwives too rash in the giving of purges, 499, note. Form of a gentle purge for infants disordered in the bowels, 503. For the thrush, 505.

Puftules, in the fmall-pox, favourable and unfavourable appearances of, 20%, 20%. The suppuration of, to be promoted, 210. When and how to open, 214. slq sooroise 9 sten hills cobrammoust Putrid fever. See Fever.

bite of a ratio-funite, 455, others, tales for leounity

Quacks put out more eyes than they cure, 426. Quackery, how to deftroy, xxvi.

Quakers, their mode of dreffing recommended, 87.

Quinfey, a common and dangerous diforder, and to whom most fatal, 251. Its causes, ibid. Symptoms, 252. Regimen, 253. Medi-- cal applications, 254. How to promote suppuration, 255. How to nourish the patient when he cannot swallow, 256. Advice to perfons fubject to this diforder, ibid.

_____, malignant, who most fubject to, and its causes, 257. Its fymptoms, ibid. Regimen and medical treatment, 259. Ingenious invention in the cure of, 261.

R

Rattle-Inake, Negro remedy for the core of its bite, 455. and vd

Regimen ought to co-operate with medicine to accomplish the cure of difeases, xvi. Will often cure diseases without medicine, 133. See Aliment.

Religion, true, calculated to support the mind under every affliction, 115. The inftructions in, ought not to dwell too much on gloomy fubjects, ibid.

Remitting fever. See Fever.

Repletion impairs the digettive power, 66. Difeafes occasioned by, 6-. How to treat a loolenels produced by, 296.

Refentment, the indulgence of, injurious to the conflictution, 108.

Refins, and effential oils, the proper menstruum for, 696.

Respiration, how to reftore in a drowned perfon, 556.

Rheumarism, acute and chronic, diffinguished, 304. Caufes, ibid. Symptoms, ibid. Medical treatment, 365. Cautions to perfons fubject to this diforder, ibid.

Rice, a general article of diet, 628. Simple boiling of, renders it a proper substitute for bread, ibid.

Rickets, the appearance of, in Britains dated from the growth of manufactores and fedentary employments, z1. The caufes of, c17. Symptoms, 518. Regimen, and medical treatment, ibid. one

Rollers, pernicious tendency of applying them round the bodies of infants, 11.

Romans, ancient, their great attention to the cleanlinefs of their towns, 97, note.

Roles, conferve of, its great virtue against hamorrhages, 317, 322. Rofemary, the external application of, a popular remedy for the forame, 421, noter bal souge a gemen , 144 ston , 21 .

Ruptures,

Ruptures are chiefly incident to children and very old perfons, 547. The caufes respectively, *ibid.* Method of treatment, *ibid.* Cutting should be avoided, if possible, 548. Cautions for perfons afflicted with a rupture, *ibid.* Often prove faral before discovered, *ibid. nove.*

Rutherford, Dr. his preparation for the cure of a dyfentery, 328, note.

contequances relating from

Sailors, their health injured by change of climate, hard weather, and bad provisions, 42, 43. Many of their diseafes fpring from intemperance, *ibid*. Ought to guard against wet clothes, *ibid*. How the ill effects of falt provisions might be corrected, *ibid*. Peruvian bark the best antidote to failors on a foreign coast, 44. Cleanli-

nefs greatly conducive to their health, 98. offertal and boot

Sal-prunella, its good effects in a quinfey, 254.

Saline draughts, of good use for stopping a vomiting, 301. Preparation of, for this purpose, *ibid*. Peculiarly good in the puerperal fever, 499.

Salivation not necessary in the cure of the venereal difease, 474. Sarfaparilla, a powerful assistant in venereal cases, 476.

Scabbea head in children, difficult to cure, 511. Medical treatment, 512. Scarlet fever. See Fever.

- School, fending children there too young, its bad confequences, 23. Ought to be feated in a dry air, and not to be too much crowded, 30.
- Schirrus in the liver, proper regimen in the cafe of, 291. See Cancer.
- Scrophula, nature of this difeafe, and its caufes, 373. Symptoms, and regimen, 374. Medical treatment, 375. The molt effectual means againft, by keeping children dry and clean, 377; bathing frequently, *ibid.*; dreffing in light and loofe clothing, 378; breathing wholefome unconfined air, and using moderate exercise, *ibid.*; butter and other oily substances, and also gingerbread, plum-cakes, &c. highly injurious in this difease, 379.
- Scurvy, why prevalent among the English, 60. Where most prevalent, and the two diffinctions of, 368. Causes of, 369. Symptoms and cure, *ibid*. Instructions to fea-faring men, 370. Extraordinary effects of milk, 371. Proper liquors, *ibid*.

Sea-bathing. Sec Bathing.

Sea water, a good remedy in the king's evil, 3/5.

Sedentary life, includes the greater part of the humin species, 44. Few perfons follow agriculture who are capable of other business, 45. Sedentary and active employments ought to be intermixed, for the take of health, *ibid*. Artists fuffer from unwholesome air, by being crowded together, *ibid*. The postures artists are confined to, injurious to health, 46. Diforders produced by, *ibid*. Cautions offered to the sedentary, *ibid*. Sedentary amulements improper for sedentary persons, 47. Hints relating to improper food, 48. Exercise, a sure relief for low spirits than drinking, *ibid*. Gardening a wholesome amusement for the sedentary, 49. Diforders occasioned by intense study, 51. Dietetical advice to the sedentary, 65. Sedentary occupations better adapted to women than men, 78, note.

Senfes,

Senfes, diforders of, 426. Intemperance deftroys the energy of, 435. Excefs of eating produces the same effect, *ibid*. Uncleanness highly injurious to the organs of sensation, *ibid*.

Seton, fometimes has extraordinary effects in an inflammation of the eyes, 248. Is of fervice for preventing apoplexies, 389. The best method of making it, 403.

Shoes, tight, the bad confequences refulting from, 86. The high heels of women's fhoes, ibid.

Sibbins, a venercal diforder fo termed in the weft of Scotland, how to cure, 480, note.

Sick, the mutual danger incurred by unneceffary vifitors to, 101. Perfons in health to be kept at a diffance from the fick, 102. Proper nurfes ought to be employed about them, 104. Inftructions for avoiding infection, *ibid*. Phyficians too unguarded in their vifits to them, 105, note. Tolling of bells for the dead very dangerous to, 110. Their fears ought not to be alarmed, 111. Sight injured by fludying by candle-light, 52.

Simples, a lift of those proper to be kept for private practice, 657.

Sinapifms, the general intentions of, 660. Directions for making of, ibid.

Sleep, the due proportion of, not easy to fix, 81. How to make it refreshing, 82. Complaints of the want of, chiefly made by the indolent, *ibid*. Heavy suppers cause uneasy nights, *ibid*. Anxiety destructive of sleep, 83. That in the fore part of the night most refreshing, *ibid*. Early rifers the longest livers, 85, note.

Sleeping in the fun, the danger of; 40.

Small-pox. See Pox.

Smell, injuries to which the fense of, is liable, with the remedies applicable to, 432.

Soap, Alicant, recommended in the flone, 303. Soap-lees, how to take, 309.

Solanum. See Night-shade.

Soldiers, exposed to many diforders from the hardships they undergo, 41. Ought to be employed in moderate labour in times of peace, 42, note.

Soups. See Broths.

Spine, often bent by artifts working in unfavourable postures, 47.

Spirit, rectified, the direct mensfruum for refins and effential oils of vegetables, 696. Of wine, camphorated, how to prepare, 700. Spirit of Mindererus, *ibid*.

Spirits, lownefs of, the general forerunner of a nervous fever, 180. The proper remedies for, 417. Cautions to perfons under this complaint, *ibid*.

Spirituous liquors, when good in the colic, 281. And for gouty complaints in the flomach, 341. Should be avoided by all perfons afflicted with nervous diforders, 397. The use of, often fatal, 568. Spitting of blood. See Blood.

Sponge, may be used to supply the want of agaric as a flyptic, 530, note. Its use in extracting substances stopped in the gullet, 553.

Sports, active, far more wholesome than sedentary amusements, 80. Golf, a better exercise than cricket, *ibid. note.*

Spots in the eye, how to treat, 429.

Squinting, how to correct the habit of, 429.

Stayss

744

Stays, a ridiculous and pernicious article of female drefs, 13, 85. The wearing of, tends to produce cancers in the breafts, 437.

Sternutatories, preparations of, recommended for reftoring lost smell, 433.

Sticking-plaster is the best application for flight wounds, 531.

- Stomach, exercife the best cure for diforders of, 78. Inflammation of, a diforder that calls for speedy affiltance, 275. Its caufes, *ibid*. Symptoms, *ibid*. Regimen, 276. Medical treatment, *ibid*. Pain in, its caufes, 340. Remedies for, 341. Influctions for perfons subject to, 342.
- Stone, the formation of, in the bladder, explained, 119. This diforder how diffinguished from the gravel, 306. Caufes and fymptoms, *ibid* Regimen, 307. Medical treatment, *ibid*. Alicant foap and lime-water, how to take for this diforder, 308. The uva urfi, a remedy in prefent request for, 309.
- Stool, loofe, the benefit of, to children, 33. Proper treatment of,
 when exceflive, *ibid*. The difcharge by, cannot be regular, if the mode of living be irregular, 116. One in a day generally fufficient for an adult, 117. How to produce a regularity of, *ibid*. Frequent recourfe to medicines for coffivenels, injurious to the confliction, *ibid*.
- Storck, Dr. his method of treating cancers, 439.
- Strabifmus. See Squinting.
- Strains, proper method of treating, 546. The fafeft external applications, ibid. note.
- Stranguary, from a bliftering-plafter, how to guard againft, 161. In the fmall-pox, how to relieve, 211. From a venereal caufe, defcribed, with its proper treatment, 471.
- Strangulation, course of treatment for the recovery of perfons from, 572.

Strafburgh, fuccessful treatment of a miliary fever there, 193, note. Strength, the folly of trials of, from emulation, 38, 41.

Study, intenfe, injurious to health, 50. The diforders occafioned by, 51. Character of a mere fludent, 53. Hints of advice to fludious perfons, 54. Danger of their having recourse to cordials, 55. Health often neglected while in possefilion, and laboured for after it is deftroyed, 56. No perfon ought to fludy immediately after a full meal, 57. Dietetical advice to the flucious, 65. The general effects of, on the confliction, 305.

Sublimate, corrofive, how to administer in venereal cafes, 475.

Suffocation, by the fumes of charcoal, liable to happen in clofe chambers, 559. General caufes of fuffocation, 570. Overlaying of infants, 571.

Sugar, an improper article in the food of children, 17.

- Sulphur, a good remedy for expelling worms, 346. And for the itch, 380.
- Suppers, ought not to deftroy the appetite for breakfast, 67. Heavy suppers, sure to occasion uneasy nights, 82.
- Surgery, many of the operations of, fuccefsfully performed by perfons unfkilled in anatomy, 524. Humanity induces every one more or lefs to be a furgeon, *ibid*.

Sweating, generally excited in an improper manner, in fevers, 139, 153.

Swoonings,

Savoonings, the feveral caufes of, described, 412. Proper treatment of this diforder, *ibid.* 564. Cautions to perfons subject to them, 567.

Sydenham, Dr. his method of treating fevers in children from teething, 516.

Symptoms, difeafes better diffinguished by, than by the systematical arrangement of, 129. The differences of sex, age, and constitution, to be confidered, 130. Diseases of the mind to be diffinguished from those of the body, 131.

Syncope, proper treatment in, 564.

Syrups, the general intention of, 695. How to make fimple fyrup, and to modify it for particular purposes, *ibid*.

T

Tacitus, his remark on the degeneracy of the Roman ladies, 4, note. Tallow chandlers, and others working on putrid animal fubstances, cautions to, 37.

Tapping for the dropfy, a fafe and fimple operation, 356.

Tar, Barbadoes, its efficacy in the nervous colic, 285.

Tartar, foluble, a good remedy for the jaundice, 351. Cream of, a good remedy in a dropfy, 355. And rheumatilm, 366.

Take, how to reftore the fense of, when injured, 434.

I ailors are exposed to injuries from breathing confined air, 46. Are fubject to confumptions, *ibid. note*. Often lose the use of their legs, 47. Hints of influction offered to them in regard to their health, 48.

Tea, the cuftomary use of, injurious to female conflictions, 7. Deftroys their digettive powers, and produces hyfterics, 61. The bad qualities of, principally owing to imprudence in the use of it, *ibid*. Green, chewing of, a remedy for the heart-burn, 394. Has # powerful effect upon the nerves, 404, note. Is bad for perfons troubled with flatulencies, 416. Occasions an unnecessary confumption of bread, 626. Bad effects of an immoderate use of, 627.

Teetbing, the diforders attending, 513. Regimen, and medical treatment in, *ibid*. Applications to the gums, and how to cut them, 516.

Temperance, the parent of health, go.

Testicles, swelled, the cause of, 468. Regimen and medicine in, ibid. Treatment under a cancerous or scrophulous habit, ibid.

Thirst, how it may be quenched when a perfon is hot, without danger, 125.

Thought, intense, deftructive of health, 50.

Thrush in infants, the diforder and its causes described, 505. Medical treatment of, ibid.

Tindures and elixirs, the proper medicines to exhibit in the form of, 696. Preparations of the aromatic tincture, *ibid*. Compound tincture of the bark, *ibid*. Volatile fœtid tincture, *ibid*. Volatile rincture of gum guaiscum, 697. Tincture of black hellebore, *ibid*. Aftringent tincture, *ibid*. Tincture of myrrh and aloes, *ibid*. Tincture of opium, or liquid laudanum, *ibid*. Tincture of hiera picra,

746

picra, 698. Compound tincture of fenna, ibid. Tincture of Spanish flies, ibid. Tincture of the balfam of Tolu, ibid. Tincture of rhubarb, ibid. Tonic tincture, ibid.

Tiffot, Dr. character of his Avis au Peuple, xii. His medical course for the cure of the hydrophobia, 452. His directions for gathering, preparing, and applying agaric of the oak as a flyptic, 530, note. Initances from, of the recovery of drowned perions, 557-

Tobacco, a clyfter of a decociion of, uleful to excite a vomit, 533. A clyfter of the fumes of, will ftimulate the inteflines, and produce a ftool, 284, 557.

Toes, the free motion of, deftroyed by wearing tight thoes, 86.

Tooth ach, the general causes of, 335. Medical treatment of, 336.

When recourse must be had to extraction, 337. Directions for cleaning the teeth, 338.

Touch, injuries to which the fense of, is liable, with the remedies applicable to, 434.

Towns, great, the air of, deftructive to the children of the poor, 23. Children bred in the country, ought not to be fent too early into towns, 30. Cleanlinefs not fufficiently attended to in, 97. Ought to be fupplied with plenty of water for washing the freets, 100. note. The best means to guard against infection in, 104.

Trades, fome injurious to health by making artifts breathe unwholefome air, 35, 46.

Transitions, ludden, from heat to cold, the ill effects of, to the conftitution, 125.

Travellers, the use of vinegar recommended to, 43. O 1ght to be very careful not to fleep in damp beds, 122. Fevers why often fatal to, 137.

Trees should not be planted too near to houses, 73.

Trefoil water, a good remedy in the rheumatilm, 367.

Tumours, proper treatment of, 527.

Turnips, properly cultivated, afford wholefome nourifhment, 641. Turnbull, Dr. his method of treating the croup in children, 514, note. Turner's cerate, preparation of, 683. mide of its general

Vinentiant lipition to pois

indiana to insurant

Vapour of fermenting liquors, noxious nature of, 560. Vegetables, wholefome corrections of the bad qualities of animal food. 59, 60. Their extraordinary effects in the feurvy, 372.

Venereal difease, why omitted in the first edition of this work, 456. Unfavourable circumstances attending this diforder, 457. The virulent gonorrhoza, ibid. Gleets, 465. Swelled tefticles, 468. Bubces, 469. Chancres, 470. Stranguary, 471. Phymofis, 472. A confirmed lues, 473. American method of curing this difeafe, 476. General observations, 477. Cleanliness a great preservative againft, 479. The ule of medicines ought not to be haftily dropped, 481. Is often too much disregarded, 482.

Ventilators, the most useful of all modern medical improvements, 71. Vertigo often produced by intense fludy, 52.

Vinegar, a great antidote against difeases, and ought to be used by all travellers, 43. Should be sprinkled in lick chambers, 104, 153, 190.

190. Is of confiderable fervice in the bite of a mad dog, 449; and in any kind of poifon, 453. Its medical properties, 700. Is of use to extract the virtues of several medicinal substances, 701. How to prepare vinegar of litharge, *ibid*. Vinegar of roles, 702. Vinegar of squills, *ibid*.

Viper, the bite of, the fufficiency of the greafe for the cure of, doubted, 453. Method of treatment recommended, *ibid*.

Vitriol, elixir of, an excellent medicine in weakneffes of the flomach, 392. And for windy complaints, 399.

Vitus, St. his dance, method of cure of, 408.

Ulcers, proper treatment of, according to their different natures, 535. Lime-water a good remedy, 537. Dr. White's method of treating them, *ibid*. Fifulous ulcers, *ibid*.

Vomits, their use in agues, 143; and in the nervous fever, 182. Cautions for administering in the putrid fever, 191. Ought by no means to be administered in an inflammation of the stomach, 276. Are useful in cases of repletion, 296. Are powerful remedies in the jaundice, 350. Are the first objects to be pursued when poison has been received into the stomach, 442. Their use in the hooping-cough, and how to administer them to children, 273. Midwives too rash in the use of, 499, note. Form of a gentle one for infants difordered in the bowels, 503.

Vomiting, the feveral caufes of, 298. Medical treatment of, 299. Saline draught for ftopping of, 300. Caufes of, in children, 508, How to be treated, 509. Of blood. See Blood.

Voyage, a long one frequently cures a confumption, 170. Voyages have an excellent effect on perfons afflicted with nervous diforders, 398.

Ureters, and their use, described, 287, note.

Urine, the appearances and quality of, too uncertain to form any determined judgment from, 118. Dr. Cheyne's judgment as to the due quantity of, not to be relied on, *ibid*. The fecretion and difcharge of, how obftructed, 119. Bad confequences of retaining it too long, *ibid*. Too great a quantity of, tends to a confumption, 120. Stoppage of, its general caufes, 289. Caution as to the treatment of, *ibid*. Diabetes, 301. Incontinency of, 304. Supprefilion of, medical treatment in, *ibid*. Cautions to perform fubject to this diforder, 306. Bloody, caufes of, *ibid*. Medical treatment of, 325. An obftinate deafnefs cured by warm urine, 431, note.

Urine doctors, their impudence, and great fuccefs from the credulity of the populace, 118, note.

Uva urst, a remedy in present request for the stone, 309.

Walking, more conducive to health than riding, 649.

Walls, high, unwholesome, by obstructing the frequent current of air,

Ward's effence, preparation of, 700. His fiftula pafte, a popular remedy that may deferve trial, 537.

Wars occasion putrid fevers, by tainting the air with the effluvia of dead carcafes, 186.

Walps,

Wafps, hornets, or bees, how the bite of, ought to be treated, 454. Water, frequently unwholefome by mineral impregnations, 62. Cautions for the choice of, *ibid*. Cold, the danger of drinking when a perfon is hot, 125.

Water in the head, is a diforder chiefly incident to children, 521. Its caufes, fymptoms, and proper treatment, *ibid*. The nature of the diforder feldom difcovered in due time for cure, 522. note.

Waters by infufion, how to prepare; Lime-water, 702, Compound lime-water, *ibid*. Sublimate water, 703. Styptic water, *ibid*. Tar-water, *ibid*.

Waters, fimple difiilled, their medical ufes, 704. Preparation of cinnamon water, *ibid*. Pennyroyal water, *ibid*. Peppermint water, *ibid*. Spearmint water, *ibid*. Role water, 705. Jamaica pepper water, *ibid*.

Waters, spirituous distilled, how to prepare. Spirituous cinnamon water, 705. Spirituous Jamaica pepper water, ibid.

Watery eye, how to cure, 429.

Watkinson, Dr. his dying request, 686.

Weaning of children from the breaft, the proper mode of, 16, 17.

- Weather, states of, which produce the putrid fever, 186.

Wells, caution to perfons going down into them, 72, note. Deep ought not to be entered till the air in, is purified, 560.

Whey, an excellent drink in a dyfentery, 329; and in the rheumatifm,

365. Orange whey, how to make 152. Alum whey, 303, 705.

Muftard whey, ibid. Scorbutic whey, 706.

Whitlee, 528.

Whyte, Dr. his remedies for flatulencies, 415.

Wind. See Flatulencies.

- Windows, the danger of throwing them open on account of heat, and fitting near them, r26.
- Wine, good, almost the only medicine necessary in a nervous fever, 181.
- Wines, the medical properties of, 706. Their use in extracting the virtues of medicinal substances, *ibid*. Preparation of anthelmin-thic wine, 707. Antimonial wine, *ibid*. Bitter wine, *ibid*, Ipe-cacuanha wine, *ibid*. Chalybeate or steel wine, *ibid*. Stomachic wine, 708.
- Womb, inflammation of, its fymptoms, 496. Medical treatment of, 708.
- Women, errors in their education pointed out, 4. Why fubject to hyfterics, 61. Are better adapted to follow fedentary occupations than men, 78, note. In child-bed, often die from their apprehenfions of death, 109. Their diforders rendered epidemical by the force of imagination, *ibid*. Every thing that can alarm them to be carefully guarded againft, 110. Evil tendency of tolling bells for the dead, *ibid*. How exposed to a miliary fever during pregnancy, 196. Their conflictutions injured by living too much within doors, 483. Those who work in the open air almost as hardy as men, 484. Advice to, with reference to the menstrual discharge, *ibid*. At the commencement, 485. Fluor albus, with its proper treatment, described, 489. Advice to, at the ceasing of the menfes, 490. Rules of conduct during pregnancy, 491. Causes and iymptoms of abortion, 492. How to guard against abortion, *ibid*. Treat-

Treatment in cafes of abortion, ibid. Inftructions at the time of child birth, 493. Caufe of the milk fever, 496. How to guard against the miliary fever, 497. The puerperal fever, ibid. General cautions for women in child-bed, 500. Caufes of barrennefs, ibid.

Wool, the best external application in the gout, 359.

Workhoufes poilonous to infants, 28.

then open on account of heat, and

Worms, how to treat a loofenefs produced by, 297. Three principal kinds of, diffinguished, 343. Symptoms of, ibid. Caules, 344. Medical treatment for, 345. Remedy for children, 346. General cautions for preferving children from them, 347. Danger of taking quack medicines for them, ibid. note. Parents advifed not to feed them with meat, two or three times a-day, as the flefh of animals in a flate of putrefaction often generates worms, 348.

Wort, recommended for the fourvy, and proper to drink at fea, 371. Is a powerful remedy in cancerous cafes, 440.

Wounds, are not cured by external applications, 529. Are cured by nature, ibid. Proper courfe of treatment, ibid. How to ftop the bleeding, 530. Caution against improper flyptics, ibid. Method of dreffing them, 531. Poultices for inflamed wounds, 532. Regimen, ibid.

Writing, hints of advice to those who are much employed in, 52, 54.

Grange whey, now toy make 1 ct. - Aluth whey and to c. toc.

Yaws; general hint for the cure of, 480, note. Young animals, all exert their organs of motion as foon as they are able, 20.

7.

Zinc, the flowers of, a popular remedy for the epilepfy, 407.

this wine, yor. Antimonial star, Midd. Bitter whee, Midt. Iper caccania wine, Mid. Chalybeate or fleel wine, Mrs. Stomachic

and errors in their education pointed out, 4. Why fabred to hylosies for Are better adapted to follow lefentary occupations

fore of death, 100. Their offorders rendered opidemical by the

Carolally guarded against, 1162 Evil tendency of telling bells for the dead, which How expected to a military favor during prognancy.

196. Their conditiutions infured by invite 150 much within

Ar the commencer, at the Phan when when its ne

THE END. and Machical treatment of,

750

Strahan and Prefton, Printers-Street.

Lately publified, By T. CADELL and W. DAVIES, Strand,

ADVICE TO MOTHERS

ON THE

SUBJECT OF THEIR OWN HEALTH,

Review. 15, to Reer 4

ON THE MEANS OF PROMOTING THE HEALTH, STRENGTH, AND BEAUTY OF THEIR OFFSPRING.

By WILLIAM BUCHAN, M.D.

FELLOW OF THE ROYAL COLLEGE OF PHYSICIANS, EDINBURGH, AND AUTHOR OF "DOMESTIC MEDICINE."

One Volume, Octavo. Price Six Shillings.

"This Publication difplays much good Senfe and Benevolence. The Author ftrenuoufly maintains the Propriety of a careful Attention to the Dictates of Nature, in the various Circumftances of Mothers and Children; and he contends against those injurious, though too common, Prejudices, which still exist in the Management of Children and the Subject of Female Health. We shall give a general View of the Plan of this Work, which we may with great Safety recommend to the careful Perusal of all judicious Parents."

MONTHLY REVIEW, for September, 1804.

II.

PRACTICAL OBSERVATIONS concerning SEA BATHING. To which are added Remarks on the Use of the WARM BATH. By A. P. BUCHAN, M. D. of the Royal College of Physicians, London. One Volume, Price 5s.

NEW REVIEW OF BOOKS.

Printed for T. CADELL and W. DAVIES, Strand.

THE IMPERIAL REVIEW; or London and Dublin Literary Journal, is published regularly on the last Day of every Month.

One of the professed Objects of this new Review, is, to steer a middle courfe, between, on the one hand, that Severity which, delighting to discover and dwell on Imperfection, has its Effect in blighting or discouraging the Efforts of Genius; and on the other hand, that partial Report or indiferiminate Praife, which in an equal Degree defeats the first Purpose of a Literary Review. In consequence of its Detail of Criticism being wholly confined to the more important Works in each Department of Writing, (inferior Publications being merely registered in the Monthly Lift, with brief Notices proportionate to their Claims,) the Editors are enabled to bring forward an Account of every interefling Work very foon after its Appearance; an Advantage exclusively their own. And they purpole availing themselves of every Opportunity to introduce Notices of other Works of the fame Defcription, which have been published during the last Three Years, fo that their Journal may be confidered as virtually commencing with the Nineteenth Century.

general View of the Plan of this World, which we may with graat

To which are added Remarks on the Ute of the Wassar Bat Hat, Br.

