History of the cholera controversy with directions for the treatment of the disease / by Sir George Johnson.

Contributors

Johnson, George, 1818-1896. Royal College of Physicians of Edinburgh

Publication/Creation

London: J. & A. Churchill, 1896.

Persistent URL

https://wellcomecollection.org/works/cmepsud9

Provider

Royal College of Physicians Edinburgh

License and attribution

This material has been provided by This material has been provided by the Royal College of Physicians of Edinburgh. The original may be consulted at the Royal College of Physicians of Edinburgh. where the originals may be consulted.

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org HISTORY OF THE CHOLERA CONTROVERSY

SIR GEORGE FORMSON M.D.



24.39

HISTORY OF THE CHOLERA CONTROVERSY

By the same Author and Publisher.

MEDICAL LECTURES AND ESSAYS. 25s.
AN ESSAY ON ASPHYXIA (APNŒA). 3s.

HISTORY OF THE CHOLERA CONTROVERSY

WITH DIRECTIONS FOR THE TREATMENT OF THE DISEASE

BY

SIR GEORGE JOHNSON, M.D.LOND., F.R.C.P., F.R.S.

FELLOW OF KING'S COLLEGE, LONDON; EMERITUS PROFESSOR OF CLINICAL MEDICINE; CONSULTING PHYSICIAN TO KING'S COLLEGE HOSPITAL PHYSICIAN EXTRAORDINARY TO HER MAJESTY THE QUEEN





J. & A. CHURCHILL
11 NEW BURLINGTON STREET
1896

CONTENTS.

		PAGE.
I.	MY FIRST EXPERIENCE OF CHOLERA IN 1849	1
II.	MY SUCCESSFUL TRIAL OF THE ELIMINATIVE TREATMENT	
	IN 1854, AND THE OPPOSITION IT EXCITED	2
III.	RESULTS OF MY STUDY OF CHOLERA LITERATURE	6
IV.	PUBLICATION OF "EPIDEMIC DIARRHŒA AND CHOLERA,"	
	WITH MY THEORY OF COLLAPSE, IN 1855	7
v.	REVIEWS OF MY FIRST BOOK ON CHOLERA	12
VI.	PUBLICATION OF "NOTES ON CHOLERA" IN 1866 AND	
	ITS FAVOURABLE RECEPTION BY SIR THOMAS WATSON	
	AND OTHER EMINENT MEN	15
VII.	MEDICAL REVIEWS OF "NOTES ON CHOLERA"	21
VIII.	CONTRASTED RESULTS OF EVACUANT AND ASTRINGENT	
	TREATMENT OF CHOLERA	29
IX.	DR. SUTTON'S REPORT ON THE POST-MORTEM APPEARANCES	
	IN THE STAGE OF COLLAPSE	37
X.	MR. MACNAMARA'S THEORIES AND PRACTICE	39
XI.	THE CONTRASTED HISTORY OF OPPOSITE METHODS OF	
	TREATMENT	52
XII.	THE ACTION OF VARIOUS REMEDIES	53
XIII.	DR. BURNEY YEO AND DR. CLEMOW ON THE GENERAL	
	ADOPTION OF THE ELIMINATIVE TREATMENT	55
XIV.	DIRECTIONS FOR THE TREATMENT OF DIARRHŒA AND	
	CHOLERA	50

PREFACE.

An excellent authority has recently declared that "It is now admitted nearly on all sides" that the rational, scientific, and successful treatment of cholera has been proved, especially during the recent epidemic in Germany, Russia, and Italy, "to be mainly eliminative and antiseptic." (See p. 55.)

The subject is one of high scientific interest, and of deep practical importance, and I have undertaken to give a brief history of the controversies relating to the pathology and treatment of this terrible disease which have been carried on during the last forty years and more. I have abundant materials for such a history; for since the commencement of the controversy, during the epidemic of 1854, I have collected and arranged in a series of scrap-books the most important of the published papers, reviews and correspondence, together with many private letters, which this prolific discussion has called forth. Some of the most interesting and instructive of these documents I have reproduced in the following pages.

It will, I trust, be admitted by all unprejudiced readers of this history that the theory which attributed choleraic collapse mainly to the loss of water by the stomach and bowels, and suggested the repressive treatment by opium—this theory, plausible as it appeared, is an erroneous theory, based on an incomplete observation of the symptoms of the disease, and constructed by men who had but an imperfect knowledge of the physiology of the circulation of the blood.

In fact, when this theory was first promulgated, more than sixty years ago, the most accomplished physiologists of that day had no knowledge of the structure and functions of the muscular-walled arterioles. It was not until some years later that the important discovery was made that the resisting force of the contracting arterioles is more than equal to the propelling power of the heart; so that by the conjoint action of these Lilliputian stop-cocks the circulation may be speedily and completely arrested. This great physiological discovery is the key to the pathology of the collapse stage of cholera.

The most melancholy part of this sad story is that the men who accepted the erroneous theory and carried it into practice failed to see the disastrous results of the astringent treatment which it suggested. This inability to recognise the calamitous results of a popular but erroneous theory has been referred to by H. T. Buckle, in his "History of Civilisation in Europe." He says: "There is no well-attested case on record of any theory having been abandoned because it produced injurious results. As long as a theory is believed, men will ascribe its evil consequences to any cause except the right one. And a theory that is once believed will always be believed until there is some change of knowledge which shakes its foundation. Every practical change may, by careful analysis, be shown to depend, in the first instance, on some change of speculative opinions." I have good reason to believe that at the present time few, if any, men of influence accept the exploded "dehydration" theory of collapse, or adopt the injurious practice which it dictated.

As it would be impossible to estimate correctly the number of victims that have been sacrificed to this "idol" of the human mind, so it would be impossible to over-estimate the responsibility of any one who propounds a theory which is calculated to greatly influence medical practice. All through this prolonged and, too often, acrimonious controversy, I have had a deep and almost painful consciousness of my own responsibility; but I have, from time to time, been assured by men most competent to form a sound judgment that, in their opinion, the theory of choleraic collapse which, after a most careful inquiry, I first published in 1855, is in accordance

with all the known facts of the disease, and with the established principles of physiology.

Sir Thomas Watson, while President of the Royal College of Physicians in 1866, referred to it as "a sound as well as a most ingenious and important theory. In truth, it derives strong confirmation from the fact that it unlocks, like the true key, the whole of the pathological intricacies of the disease." This is strong testimony from a great and wise man, but that which, to my mind, is still more satisfactory is the evidence which has been gradually accumulating of the beneficial results of the evacuant treatment of cholera, especially in its early stages. What can be more conclusive than the fact, referred to in detail hereafter (p. 29), that in Liverpool* during the epidemic of 1866, out of several thousand cases of choleraic diarrhoea treated by laxatives not one passed into collapse or required removal to the "In every case relief was afforded hospital? 'quickly, safely, and pleasantly.'"

No such evidence as this ever has been or can be given in support of the astringent treatment of the disease.

Although this essay is mainly a history of past controversies, and not a complete treatise on the pathology and treatment of cholera, it will, I think,

^{* &}quot;On the Treatment of Cholera and Epidemic Diarrhea," by Dr. M'Cloy and Dr. Robertson: Med. Chir. Trans., vol. 1.

be found to give a clear sketch of the entire subject; but I must refer those who wish to read a full and complete discussion of the pathology of the disease to the chapter on "Epidemic Cholera" in my Medical Lectures and Essays.

In the concluding section of this treatise I have given detailed directions for the prevention and treatment of diarrhœa and cholera. It will be seen that the method of treatment, which ample experience has proved to be highly successful, is entirely consistent with what is now acknowledged to be the true pathology of the disease.

11 SAVILE ROW, Nov. 29th, 1895.

HISTORY OF THE CHOLERA CONTROVERSY

I. My First Experience of Cholera in 1849.

I PROPOSE in this treatise to give a brief history of the various theories with regard to the pathology of Cholera, and the different modes of practice, suggested by those theories, which have been adopted by the profession during the last forty years.

During my student days I was taught that the worst symptoms of cholera were due to the drain of water from the blood and tissues, and that the main object of treatment in the early stage of the disease was to arrest the gastro-intestinal discharges by opium.

I first saw cholera during the epidemic of 1849. I was then resident medical tutor in King's College, and I was asked to prescribe for the wife of one of the porters of the College. She was about forty years of age, and had been in good health until she was seized with vomiting, purging and cramps.

She had a warm skin and a good pulse, and I looked upon the case as a mild one. In accordance with the teaching that I had received, I gave her five grains of Dover's powder every hour until three doses had been taken. When I saw her again, in about three hours from the time of my first visit, I found to my dismay that although the vomiting, purging and cramps had ceased, she was in full collapse, and she died in a few hours.

I saw a similar result follow the like treatment by other physicians at the hospital. I was horrified to see that in case after case the speedy arrest of diarrheea by opium was directly followed, not by recovery, as in accordance with the then generally accepted theory it ought to be, but by rapidly fatal collapse. These facts convinced me that the theory was wrong, and the treatment based upon it injurious.

II. My Successful Trial of the Eliminative Treatment in 1854, and the opposition it excited.

In 1854 came another epidemic of cholera. I was then the junior assistant physician of King's College Hospital, and during almost the entire period of the epidemic I was left in entire charge of the medical wards.

It was an eventful time, and I entered upon the work with fear and trembling. I remembered that

in 1849 about three-fourths of the cases of cholera in our hospital treated by brandy and opium were fatal, while the same proportion of about an equal number treated by copious draughts of salt and water recovered.

So far as I could learn from reading, two of the most successful methods of treatment, judging by the results apart from theories, were that of Dr. Ayre, by frequent small doses of calomel, and that of Dr. Stevens, by repeated doses of salines. That which was common to these two methods was that they tended to encourage and not to repress the discharges from the bowels. Then, acting upon a provisional hypothesis, that the discharges are the means by which the cholera poison is ejected from the system, I determined to try the effect of the quickly acting evacuant castor oil, and the results were surprisingly favourable and encouraging.

The first seven cases of choleraic collapse that came under my care were treated by more frequent doses of castor oil than I afterwards found to be necessary or desirable; but they all recovered, the collapse passing off while the discharges continued. The inference was obvious: the discharges from the stomach and intestines are the means by which the cholera poison is ejected and the patient is restored to health.

On the 6th of September 1854 I addressed to the editor of the *Medical Times and Gazette* a letter, in which I stated that I had treated fifteen cases of choleraic collapse by castor oil with only three deaths,

one of which was fatal within half an hour after admission. I further stated that hundreds of cases of choleraic diarrhœa, including several nurses and one pupil in attendance in the cholera wards, had been quickly cured by one or two doses of castor oil, and that not one had passed into collapse.

The publication of this letter led to such an outbreak of censure and opposition as probably no other member of the profession has ever been assailed with. The most atrocious attack upon me by any medical journal was contained in a leading article in the Association Medical Journal. The editor was Dr. (afterwards Sir John) Rose Cormack, and if he did not write the article, he was responsible for its publication.

The writer, after commenting on what he considered the unreasonableness of my theory and the danger of my practice, proceeded as follows: "With Dr. Johnson's opinions we have now done. It is evident that the author has gone astray from the right paths, and has subjected himself to the imputation of belonging to a class by whom (to use the language of Bishop Butler) arguments are often wanted for some accidental purpose, but proof as such is what they never want for their own satisfaction of mind or conduct in life."

This sapient writer was an open enemy, who might have been prosecuted for a malicious libel, but most of my best friends believed my theory and practice to be wrong. Dr. Robert Ferguson said to Mr. Charles Hawkins, as I heard many years afterwards in a conversation with Mr. Hawkins at the Athenæum Club, "What a pity it is that Johnson has taken up this notion about cholera. It will be absolute ruin to him—absolute ruin."

Dr. Budd (who was then senior physician of the hospital, and whom I afterwards succeeded as Professor of Medicine) came back from his holiday when the epidemic was over, and gave a clinical lecture, in which he made no mention of me or the work which had been going on in his absence, but told his class that the proper treatment of choleraic diarrhea was to arrest the discharges by opium. Some of his pupils, who had watched my practice in the cholera wards, reported to me his words, but felt that they knew more of the subject than the Professor.

Dr. Todd, who was one of my kindest friends and benefactors, said to me when the epidemic was over: "I believe that in the main you are right about cholera, but you will not convince the profession that you are, and in attempting to convince them you will do yourself great harm. I advise you to publish no more on the subject."

This was kind and friendly advice, but I told Dr. Todd that I could not act upon it, and that I had determined to publish the notes of all my cases, together with such information relating to the pathology of cholera as I might be able to obtain.

III. Results of my Study of Cholera Literature.

For a period of about six months I devoted almost the whole of my time to the careful reading and analysis of books and reports on cholera, most of which were published by men who had had a large experience of the disease in India. At that time, while I was under a cloud of misrepresentation, with the nickname of "Castor Oil Johnson," my reading was but little interrupted by practice, and if I had not gained a certain reputation in connection with kidney disease, my friend Dr. Ferguson's gloomy prophecy might have been fulfilled. It was a time of great interest to me, for I was daily increasing my knowledge of the disease, and from time to time lighting upon facts which were obviously inconsistent with the theory which assumes that the chief cause of collapse is the drain of water from the blood. The majority of the Indian practitioners were agreed that, while the most deadly cases of the disease were those in which the patient falls rapidly into collapse, with little or no discharges, a complete arrest of the discharges during the stage of collapse is a sign of fatal import. Death may take place without discharge; recovery never.

Then it is very remarkable that no Indian practitioner, writing before cholera became epidemic in Europe in 1831-2, appears to have given opium for the purpose of arresting the discharges. Opium was

often given to relieve the painful cramps, but at the same time purgatives were employed to remove the offensive morbid secretions.

Dr. Edmund Parkes* was the first to publish the fact that when the inspection of the body is made soon after death during the stage of collapse, the right side of the heart, the pulmonary artery, and the large systemic veins are distended with blood, while the left side of the heart and the systemic arteries are nearly or quite empty. These facts indicate, as Dr. Parkes said, that during the collapse stage there is an impeded flow of blood through the lungs. Dr. Parkes was unable to explain this impediment, for the simple reason that his book was published four years before the earliest of M. Claude Bernard's researches on the vaso-motor nervous system were made known to the world.

IV. Publication of "Epidemic Diarrhœa and Cholera," with my Theory of Collapse, in 1855.

When, in 1855, I published my book "On Epidemic Diarrhœa and Cholera," the function of the vaso-motor nerves and of the arterioles had been ascertained, and I was enabled to suggest that the impeded circulation through the lungs during choleraic collapse is a result of contraction of the

^{* &}quot;Researches on the Pathology and Treatment of the Asiatic or Algide Cholera." 1847.

muscular arterioles excited by the poisoned blood. This action of the vaso-constrictors of the lungs explains all the most striking phenomena of choleraic collapse. It explains the emptiness of the systemic arteries, with extinction of the pulse at the wrist, the cadaverous shrinking of the features and sinking of the eye-balls, also the lividity of the surface, consequent on the venous fulness. The stream of blood through the pulmonary capillaries being much decreased, the supply of oxygen is proportionally reduced in quantity. Hence during the stage of collapse there is defective oxygenation of the blood and the tissues, with resulting lowering of the surface-temperature, diminished exhalation of carbonic acid, and almost complete suppression of bile and urine; carbonic acid and the chief constituents of bile and urine being joint products of oxidation. That this is the true explanation of the suppression of bile and urine is rendered the more probable from the curious fact that when a nursing mother becomes the subject of choleraic collapse the secretion of milk, though poor in quality, is but little diminished in quantity. Now the chief constituents of milk-casein, sugar, fat and water-are not oxidised products.

The impeded circulation through the lungs is the cause and not a result of the blood thickening in the collapse stage of cholera. So long as the circulation is free, diarrhea, no matter how copious, does not cause thickening of the blood. The loss

of water by the bowels is replaced by absorption of water from the muscles and other soft tissues, which contain water in the proportion of four-fifths of their weight; but when the vaso-constrictors of the lungs impede the onward passage of the blood, the entire systemic venous system is engorged, and water escapes from the blood into the partially dehydrated soft tissues, while venous absorption from the stomach and bowels is impeded. Defective oxidation, consequent on the scanty stream of oxygen-carrying blood through the lungs, contributes to the blood thickening by lessening the disintegration of the solid constituents, which consequently accumulate in the blood. Dr. Dundas Thompson found the blood of a patient suffering from an affection of the bronchial mucous membrane thicker than any cholera blood that he had ever examined. (Med. Chir. Trans. vol. xxxiii.)

The theory that choleraic collapse is the result of extreme constriction of the pulmonary arterioles explains, in the worst class of cases, the sudden onset of collapse, and in other cases the equally rapid recovery. Twining, an Indian practitioner, says: "In these instances recovery seems almost as rapid as in cases of patients who are resuscitated after suspension of animation from submersion in water." No such cases of rapid recovery are possible after extreme exhaustion by profuse discharges from the blood and bowels.

The marvellous temporary benefit that results

from the injection of a warm saline solution into the veins, which has been generally assumed to lend support to the theory that dehydration of the blood and the tissues is the main cause of collapse, receives a more probable explanation, from the relaxing effect of the warm liquid upon the constricted pulmonary arterioles. Dr. Mackintosh of Edinburgh injected the veins of 156 patients in 1832.* The temperature of the fluid injected ranged from 100 to 120 degrees Fahr., but he states that "the good effects of the injection were rapid in proportion to the heat of the solution."

Several of the Indian practitioners describe the great relief which has often been afforded by vene-section during the stage of collapse. The following striking case is recorded by Sir Ranald Martin:† "On visiting my hospital in the morning the European farrier-major was reported to be dying of cholera. I found that during the night he had been drained of all the fluid portion of his blood. His appearance was surprisingly altered; his respiration was oppressed, the countenance sunk and livid, the circulation flagging in the extremities. I opened a vein in each arm, but it was long before I could obtain anything but trickling of dark treacly matter. At length the blood flowed, and by degrees the

6th ed. p. 349.

^{* &}quot;Practice of Physic," 1st edition, vol. i. p. 363. + "The Influence of Tropical Climates on European Constitutions,"

darkness was exchanged for more of the hue of nature. The farrier was not of robust health, but I bled him largely; when he, whom but a moment before I thought a dying man, stood up and said, 'Sir, you have made a new man of me.' He is now alive and well."

The explanation of this striking result of a practice inconsistent with the operator's theory, is that there was a paralysing over-distension of the right side of the heart, consequent on the impeded circulation through the lungs. Venesection relieved the over-distension of the right cavities, and so increased the contractile power of their walls. The experiments of Dr. John Reid and others have proved that vene-section often restores the contraction of the over-distended right cavities in asphyxiated animals.

The distension of the right side of the heart after death from choleraic collapse is precisely the same as is found after death from asphyxia, and extreme contraction of the pulmonary arterioles is the only possible explanation of the arrested circulation in both classes of cases.

The doctrine, that the essential cause of choleraic collapse is the greatly impeded circulation through the lungs, receives remarkable confirmation from the fact that several cases are recorded in which a fibrinous plug in the pulmonary artery has caused symptoms strikingly resembling cholera.*

^{*} See the chapter on Epidemic Cholera in my "Medical Lectures and Essays."

The most remarkable case of this kind is one published by Dr. Alfred Carpenter (*The Lancet*, Sept. 23, 1871). Dr. Carpenter remarks that "the only symptoms wanting to make it apparently a case of cholera were alvine discharges and cramps in the limbs." Great thirst and almost complete suppression of urine were prominent symptoms.

The preceding is a brief abstract of facts and reasonings, most of which are fully related in my book "On Epidemic Diarrhœa and Cholera," which was published in 1855.

That work was the result of much labour and careful thought, and I naturally looked with interest to see how it would be dealt with by the journals which are supposed to lead professional opinion.

V. Reviews of my First Book on Cholera.

The Lancet (June 2, 1855) began a notice occupying about a column with the following sentence: "The essential point in connection with cholera, with which the name of Dr. George Johnson was rather prominently obtruded upon the public, is by far too well known and easily defined to be lost sight of in the physiological adornments with which the author attempts to dress the question and dazzle the reader in the 300 pages now before us."

The writer probably had not read the book, or, having read it, he had not the physiological knowledge which would have enabled him to give a fair review of the work. He says: "This work is nicely written, his physiology may appear unexceptionable, and his pathology may be as sound as the school he is attached to is capable of," but "the Committee of the Medical Council of the General Board of Health has shown in its late Report on the results of the different modes of treatment, that the procedure in question [the eliminative plan] is about the worst that has ever been promulgated." I shall hereafter show how entirely untrustworthy is the Report to which my critic referred. (See p. 27.)

The Medical Times and Gazette (June 23rd, 1855) devoted four columns to a full, fair and courteous review, which showed that the writer had carefully read the book, of which he gave a clear, correct and intelligible analysis. "The book," he says "is the production of one of the most acute and able physicians of the modern school; one whose labours in pathology entitle him to a full and fair hearing." . . . "Our own sentiments on the merits of the book are that the author seems always too eager to prove his case. His very profound knowledge of modern physiology and pathology seems to betray him into too elaborate and plausible chains of argument. His dexterity in explaining away whatever opposes his views is remarkable, no less

than his eagerness at seizing on any incident, however trivial, that supports them." This expression of opinion is fair enough; but I might ask whether my eagerness to prove my case had been as great as that of my opponents to disprove it, and whether my dexterity in explaining away whatever opposed my views had been equal to the blindness of those who failed to see the significance of collapse following directly upon the arrest of the discharges by opium.

Finally the reviewer says, "We have frankly stated our objections to this very ingenious and able work, which we advise our readers to peruse carefully for themselves, with the assurance that they will find it abundantly instructive; and even if they are not converts to the eliminative plan of treatment (for we must observe that Dr. Johnson nowhere asserts that castor oil is the specific), yet they will respect the author for the courage and zeal with which he enunciates doctrines which he knows to be unpopular, no less than for the extensive information which supplies him with arguments from every department of physiology and pathology."

The editor of the Association Medical Journal did not notice the book. Perhaps he would have found it difficult to give a fair review of the work after his statement that the author had "deviated from the right paths," &c. See ante, p. 4.

VI. Publication of "Notes on Cholera" in 1866 and its Favourable Reception by Sir Thomas Watson and other Eminent Men.

When my book was published in 1855, the epidemic had passed away, and with it departed the alarm and the interest it had excited; my work, therefore, attracted but little attention, and probably had very few readers.

After an interval of ten years cholera again appeared to be approaching our shores, and, in anticipation of a probable return of the disease, I published a series of papers in the British Medical Journal. These papers I afterwards collected, revised and published at the beginning of 1866, in a small volume entitled "Notes on Cholera." This volume excited much interest, and, perhaps on account of its smaller size, had many more readers than my previous work on the same subject, but in my judgment the most important result of its publication was that it made a convert of Sir Thomas Watson, who was at that time President of the Royal College of Physicians. I had sent him a copy of the book, and on January 24th, 1866, I received from him a letter, from which the following is an extract. "I am bound to say that I find no flaw in your reasonings. Your position is admirably well expounded and sustained, and the pathology that you adopt is rational and plausible in the best sense of that word. It seems also to be justified by the results of your peculiar treatment of the disease, and by treatment that resembles yours." I need not say that to receive this testimony as to the reasonableness of my views from one of the wisest, most cautious and conscientious of men was very gratifying and encouraging, after the adverse criticism and censure which my advocacy of the eliminative treatment had called forth.

So great an interest did Sir Thomas Watson take in the subject that, notwithstanding the many calls upon his time, he wrote a review of the book, which was published in the *Saturday Review*, June 2, 1866.

After giving, in his own inimitable style, a lucid summary of my explanation of choleraic collapse, the reviewer went on to say: "Surely this seems a reasonable theory. It is founded on a true analogy, it is consistent with the symptoms noticed during life and with the conditions discovered after death; we may therefore legitimately regard it, until fairly refuted, as a sound, as well as a most ingenious and important theory. In truth, it derives strong confirmation from the fact that it unlocks, like the true key, the whole of the pathological intricacies of the disease."

Sir Thomas Watson then concluded his article in the following terms, which, by the favour of the author, I was permitted to copy from his manuscript: "We need not abstain from expressing our belief that the true pathology of this awful disease, the sequence and relation of its various phenomena, have now for the first time been determined and made known; and we congratulate the eminent physician who has done the world this great service, on the firmness of purpose with which he has withstood the force of ridicule in high places, and upon the perseverance and sagacity which have enabled him to solve so neatly and thoroughly a very complex and deeply interesting problem. The more his facts and reasonings are scrutinised, the more widely and entirely will they, in our judgment, compel assent, and the greater will be the consequent salvage of human life."

The article, not being signed, this decided expression of opinion would not be known to have the weighty authority of the President of the Royal College of Physicians; the editor of the Review, therefore, substituted for it the following sentence: "On this point, as on others, Dr. Johnson's facts and reasonings are well deserving of careful and impartial consideration."

In 1871 Sir Thomas Watson brought out a new edition of his Lectures on "The Principles and Practice of Physic," and in the lecture on "Epidemic Cholera" he set forth at length the facts and reasonings which had induced him to change his opinion with regard to the pathology and treatment of the disease. This change of opinion had before been published, and there appeared in the Lancet,

Aug. 19, 1871, an article treating on "The Therapeutics of Cholera," from which the following is an extract: "The irrepressible castor oil controversy is threatening to assert itself again. To say nothing of the physician who is chiefly associated with the eliminative theory of treatment, a distinguished medical Baronet, not given to commit errors of judgment or of taste, does not hesitate, every now and again, by somewhat unusual courses, to indicate his adoption of the castor oil plan of treatment, or allow it to be indicated. If this treatment is to have further trial on the strength of the authority of Sir Thomas Watson, we can only hope that Sir Thomas's authority will continue to be entirely dissociated from any personal experience of the treatment, as it is now admittedly unsupported by any personal observation of cases. Sir Thomas Watson's mere authority will undoubtedly cause this practice to be tried more extensively than otherwise it would be, and there is great responsibility in such a use of authority. But the days are gone for enjoining a practice on the strength of mere authority, apart from personal or clinical experience; and we cannot doubt that the profession will agree with us in thinking that Sir Thomas Watson would have done better to withhold views on this momentous subject till he was able to support them by some facts either within his own observation or which it was in his own power to certify."

I cannot doubt that the unprejudiced members of

the profession will be of opinion that the error of judgment or of taste has been shown, not by Sir Thomas Watson, but by his critic.

When that article was published there must have been on the staff of the Lancet a man too old to learn or too proud to acknowledge that his persistent advocacy of the repressive treatment of choleraic symptoms was the result of accepting a theory of the disease which had been proved to be erroneous, and the holding of which blinds men to the mischievous results of the practice which it suggests.

The writer of the article in question speaks of the great responsibility in such a use of authority. Is there no responsibility in the attempt of an anonymous writer in an influential medical journal to discredit such an authority as Sir Thomas Watson?

We know how eloquently in his introductory lecture the great physician expressed his deep sense of the grave responsibility which attaches to a public teacher. He says: "Doctrines and maxims flow abroad from a public teacher as from a fountain, and his faulty lessons may become the indirect source of unspeakable mischief and suffering to hundreds who have never even heard his name. These thoughts fill my mind with an almost painful sense of the obligation imposed upon me by my present office of closely sifting the facts, and meditating carefully the precepts which I offer for your instruction and guidance."

This is the wise and cautious teacher whom the

anonymous leading article writer would subject to a repressive process similar to that which he blindly advocated for the treatment of cholera. One result of such vituperative writing as I have referred to, is to deter cautious and timid men from publishing opinions which they believe to be true, but which they know to be unpopular.

After the publication of my "Notes on Cholera," I received many letters from eminent members of the profession expressing entire agreement with my pathology and practice; but when I suggested that they should give public expression to their approval the reply generally was that they had nothing new to say; my facts and arguments were unanswerable, and required no further confirmation. The fact is that, as the most revered head of the profession had not escaped unseemly censure for having the courage to admit that he had abandoned opinions and a practice which he had discovered to be erroneous and mischievous, they were unwilling to expose themselves to the like unpleasant ordeal. I here reproduce one of the letters above referred to. The writer is physician to one of the largest London hospitals, a Fellow of the Royal Society, and has a great reputation for his original researches in a special department of pathology.

"I have read your book. It is the most interesting medical book I have yet read. Praise from me to you is out of place. But I must say that whatever your theory of the particular disease,

cholera, may lead to, your study of cholera as a piece of pathology is a specimen of work of the highest kind. As a student of medicine I am delighted to see such good work."

This expression of opinion by a very eminent pathologist and a most experienced physician, was very gratifying, and if he had sent a somewhat similar communication to one of the medical journals it would have made the most reckless reviewers cautious in the exercise of their responsible critical functions.

VII. Medical Reviews of "Notes on Cholera."

I now propose to refer briefly to the manner in which the principal medical journals dealt with my "Notes on Cholera."

The Lancet did me the honour to devote two leading articles to the analysis and review of the book.* The writer says: "We need not say that the book is well written; and although there is apparent now and again a too great readiness to accept whatever makes for favourite opinions, and to treat lightly an opponent, yet, admitting these faults, Dr. Johnson writes like a serious and a scientific man, and as such we shall try to treat him." The writer, although manifestly hampered by his own views as to the nature and proper treat-

^{*} The Lancet, April 18th and June 2nd, 1866.

ment of cholera, gives, on the whole, a fair analysis of the book, but with some remarkable inaccuracies. For instance, in the first article he says: "It is important that we should represent how strongly Dr. Johnson urges that the only bearing of the evacuations on the symptoms is a beneficial one;" yet in the second article it is stated that "Dr. Johnson admits that occasionally the very abundance of the discharges may be a cause of death by exhaustion." This latter statement of my opinion is true enough, but it is quite inconsistent with the previous untrue statement, unless the writer supposed me to look upon death by exhaustion as "beneficial." The reviewer demurs to my theory of collapse; but it is obvious that he has not sufficient knowledge of the forces concerned in the circulation of the blood as taught by all modern physiologists, to see clearly that the action of the pulmonary vaso-constrictors is the only possible explanation of the condition of the heart and lungs which is found soon after death in the collapse stage of cholera. With regard to the treatment of the disease the reviewer says: "Our great wisdom in the early treatment of cholera is to keep intact the composition of the blood by restraining the drain from it which almost invariably precedes, and seems largely to cause, the symptoms of collapse."

The attempt to "keep intact the composition of the blood" by preventing the expulsion of the virulent poison which causes the drain of liquid has not the appearance of "great wisdom," and its great folly has been proved by ample experience, as I shall presently show.

This controversy with the Lancet is now a matter of ancient history. It is probable that the men who from the year 1854 were so persistently opposed to me with regard to the pathology and treatment of cholera, if they are still on the staff of that journal, have changed their opinions.

The writer of a very able review of my "Medical Lectures and Essays," which appeared in the Lancet (March 24 and 31, 1888), could not have been one of my former opponents, for after a complete analysis of the chapter on cholera, he says: "This chapter alone should win for him an honoured place in the history of medicine."

The British Medical Journal (January 27, 1866), in a leading article which occupies nearly three pages of the journal, gave a very able and lucid epitome of my "Notes on Cholera." In the course of the article the writer, referring to the theory of collapse, says: "We may truly say this of it, that there is perhaps no instance on record in which the theory of a disease founded on physiological and pathological facts more strikingly explains the symptoms and dictates the course of treatment. We therefore think that the profession is greatly indebted to Dr. Johnson for thus enforcing upon their attention at this moment so vital a question.

He offers us a pathological theory which suggests a truly scientific treatment of the disease."

The writer concludes his very careful and complete analysis of the book with the following expression of opinion: "Such, we believe, is a correct exposition of what we must venture to consider the most valuable and original work which this publishing season has added to medical literature."

The Medical Times and Gazette (May 5, 1866) gives a fairly complete and accurate analysis of my "Notes on Cholera." The reviewer remarks at the beginning of his notice, that it is "a short but exceedingly elaborate and closely reasoned work," written in vindication of my practice in 1854. From time to time he interposes comments which show that if he is a convert to my pathology he is unwilling to abandon his former creed. He truly remarks upon one point which many of my opponents fail to see. "His aim (the author's) is not to increase the drain from the blood, but to empty the intestinal canal of what is really outside the system proper. His plan is not eliminative in the strict sense then, but only in the subordinate way of removing materials from the bowels which have already been removed from the system." But surely the bowels are a part of the "system," and to facilitate and quicken the expulsion of their poisonous contents is as much a process of elimination as is excretion from the blood. I have seen several cases in which the muscular coat of the bowels has been so paralysed by opium and the accumulation of their contents that even croton oil has failed to excite evacuations. Then, referring to my theory, the reviewer says: "Let us remark again that good as this may seem—and we have rarely seen so brilliant a piece of pathology—we must judge the practice founded on it separately."

That is what I did with regard to the theory to which my reviewer is inclined. It was not until I saw the deadly result of arresting diarrhœa by opium that I looked about for a theory more in accordance with facts than that which suggested so fatal a practice. Dr. Aitken's well-known textbook is quoted to show that "the facts which Dr. Johnson adduces from the morbid anatomy of cholera, in explanation of the pathology of the disease, are generally admitted." The reviewer admits the "ingenuity" and reasonableness of the theory that the blood is arrested by contraction of the pulmonary arterioles. The general conclusion is that it may be best "to do nothing in the stage of collapse, except nurse the patient carefully and soothe him as much as possible. Little as we can do in blue cholera, the experience of mankind has shown that a great deal can be done in diarrhœa."

Yes, undoubtedly a great deal may be done for diarrhea, but the vital question is whether that which is done is beneficial or the reverse. In a series of leading articles published later than the

review, the editor claims for the astringent and repressive treatment of diarrhœa the greatest amount of success.

The Medical Press and Circular (April 11, 1866) gave a brief but commendatory review of my book. The author says: "Dr. Johnson's chief object is to disprove the theory that the collapse of cholera is due to the loss of fluids and consequent thickening of the blood, and to point out the faults of practice resulting from this (as he believes) erroneous theory. We think, however, Dr. Johnson considers the views which he combats, and the lines of treatment he opposes, are much more generally received than they really are at the present day. Dr. Johnson deduces very fair and convincing arguments from facts noted by various writers on cholera in support of his views." The reviewer then describes briefly but clearly my theory and treatment of the disease, and says: "In conclusion we have to recommend Dr. Johnson's carefully written and neatly got-up little volume, so full of interesting and useful information to all members of the profession."

It would appear, however, that the editor or some one connected with the journal was of opinion that the reviewer had been too much in favour of my views, for soon afterwards, on May 23, there appeared a leading article, in which the writer says: "Without entering at all into the merits of Dr. Johnson's pathological theory, which is ingenious and may be true, we at once demur to his practical

corollary. In 1855 the Treatment Committee of the Medical Council addressed a report on the results of the different modes of treatment pursued in epidemic cholera to the President of the General Board of Health."

The general percentage of deaths from each plan of treatment was stated to be as follows:

The writer of the article thinks these figures conclusive, and he says: "From a fair experience we can confidently recommend full doses of opium, as an almost unfailing specific against the premonitory diarrhæa, while we know of no cure for true cholera save in careful nursing."

Yet the Treatment Committee reported that 79'7 per cent of cholera cases were cured by "astringents, chalk and opium."

I have not before made any public comment on this notorious report, which was published in all the papers, both general and medical, and must have had great influence in deterring fair-minded men from giving a trial to the evacuant treatment.

The obvious, if not the avowed, object of the Committee was to put an extinguisher upon me, my theory and my practice. It may be safely assumed that the Committee and those who professed to try

the eliminative treatment, were as convinced that it was wrong in theory and injurious in practice as was the chairman of the Committee, Sir George Burrows, who, when Sir Thomas Watson publicly announced that he had become a convert to my views, wrote to rebuke him for having "abandoned so many of his early principles." Sir Thomas Watson's reply to his censor was, as he told me: "Although I am advanced in years, I hope I am not too old to learn."

Now, men conscientiously believing that the eliminative treatment was injurious, could not give it a fair trial; for they would not dare to apply it to any but the most hopeless cases. It would not be difficult to go into a cholera ward, and select cases in which, under my treatment, the mortality would be not less than 100 per cent. No particulars of the cases were published, but it must have been in cases of extreme and desperate collapse that the mortality was as high as 71.7 per cent., and on the other hand the statement that with a treatment by "astringents, chalk and opium," choleraic collapse -which all the cases were represented to be-was fatal to the extent of only 20'3 per cent., is sufficient to discredit the entire report. I shall presently refer to some much more trustworthy statistics than these. But before doing so I am anxious to show that the high mortality of cholera during the stage of collapse affords significant evidence that the natural cure of the disease is by a process of elimination. No drug, not even opium, has so direct and powerful an influence in arresting or checking elimination as the greatly impeded circulation through the lungs, of which extreme contraction of the pulmonary arterioles is the cause, and the condition known as collapse the result. The supply of blood to the left side of the heart and the systemic arteries is greatly reduced, and, as an obvious consequence, the process of elimination by the gastrointestinal mucous membrane is in a corresponding degree lessened. The discharges are always greatly diminished during collapse, and too often their complete suppression is inevitably fatal. It is surprising that any one who has seen much of cholera should doubt that a continuance of the discharges during collapse is an essential condition of recovery from that perilous state.

VIII. Contrasted Results of Evacuant and Astringent Treatment of Cholera.

The most valuable record of the results of various modes of treatment that I know of is contained in a very able and most interesting paper by Drs. M'Cloy and Robertson, published in the 50th volume of the "Transactions of the Royal Medical and Chirurgical Society." The authors state that 375 cases of cholera with collapse were admitted into the Liverpool parish infirmary during the

epidemic of 1866. Of these cases 91 were treated with astringents and stimulants, camphor and iced water, applications of ice and hypodermic injections, with a mortality of 71.42 per cent. 87 cases were treated with castor oil, and a liberal use of food and alcohol, with a mortality of 42.37 per cent. The authors express their belief that food and alcohol given during collapse were injurious, and to some extent counteracted the good effects of the evacuant treatment. 197 cases were treated with castor oil, receiving little or no food of any kind or alcoholic stimulants until reaction was fairly established; of these only 30.45 per cent. died.

It is stated that the cases which were treated by astringents and stimulants occurred for the most part at the very outbreak of the epidemic, and amongst an unfavourable class of patients, but at another institution, under the same mode of treatment, the mortality continued almost as high during the whole period of the epidemic. Again they say "the *sudden* decrease in the death rate which followed the change of treatment pointed to the latter as the cause of the former."

Some interesting tables and details of typical cases are given. For instance, a boy aged ten recovered from collapse of such gravity and with so extreme an arrest of the circulation that he had gangrene of both hands, which necessitated amputation, followed by complete recovery. With regard to diarrhœa, the authors say, "Our experience was

very extensive. Several thousand cases came under our observation in the different dispensaries connected with the West Derby Union, and in the Liverpool parish infirmary. Among these were many who doubtless would have recovered under any mode of treatment, or by the vis medicatrix naturæ alone. There were many, too, of a most severe choleraic type. The treatment adopted was generally evacuant in its nature, and consisted in the administration of castor oil, calomel, rhubarb, or magnesia. In every case relief was effected quickly, pleasantly, and safely. It was but seldom that more than two or three doses of castor oil were required. In one of the public dispensaries (Bootle) many cases of diarrhoea were treated with evacuants, and the testimony of the medical officers is in accordance with our own. 'We certainly had less trouble with the evacuant mode of treatment.' We never saw a diarrhoea patient treated from the commencement of his attack, require subsequent removal to the hospital. In a large proportion of our cases (of cholera) there was premonitory diarrhœa, which had been treated, often for four or five days, with astringents. Diarrhœa patients undoubtedly recover when treated with astringents, but the recovery is not consequent on the arrest of the discharges, as these are invariably restored before the patient feels well." The authors state emphatically that "whatever the treatment adopted the result was the same-recovery never occurred without the continuance of the intestinal discharges or their restoration, if previously arrested."

I have no doubt that this important paper—although, so far as I know, Sir Thomas Watson is the only subsequent writer, except myself, who has quoted it and recognised its value—has silently exercised great influence upon the mind of the profession.

In the Lancet of August 18, 1866, there is a report by Dr. Ferris of 201 cases of diarrhœa successfully treated at the Bloomsbury dispensary, mostly by evacuants, which Dr. Pidduck, the physician to the charity, had found successful in previous epidemics. A pill of half a grain of calomel was followed in two hours by half an ounce of castor oil. If next day the purging continued, a slightly astringent mixture was given, "but this was necessary in only a small number of cases." All the patients recovered, and the report states that a considerable number of them had been previously treated at other institutions unsuccessfully by astringents. "One man had been ill for a week, taking astringents the whole time; after the above treatment he went to work the next day quite well." Twenty-two of the cases were treated by coloured mint water, of whom only two returned for further treatment.

I challenge the advocates of the repressive treatment of choleraic diarrhea by opium and other astringents to produce evidence in support of their theory and practice as conclusive as that which has here been adduced in favour of evacuants. What is often said is that by the use of astringents some cases of diarrhœa appear to have been prevented from passing into collapse, but there is no recognition of the fact that collapse has often followed directly upon the arrest of the discharges by opium.

I have, during the last forty years, treated hundreds of cases of diarrhoea by evacuants, and not one so treated has passed into collapse. Contrast the result of the several thousand cases in Liverpool and the 201 cases at the Bloomsbury dispensary with the disastrous results of the opposite treatment by two eminent French physicians, Drs. Briguet and Mignot.* They treated in the hospital 200 cases of diarrhœa, at the very commencement of the attack, by from 15 to 30 drops of laudanum, the dose to be repeated in an hour, and followed by opiate enemata if necessary. The result was that 26—that is 13 per cent.—passed into collapse. It is probable that most of them would have passed into collapse were it not for the fact that the outward flux from the blood impedes the absorption of the drug, and the natural curative efforts eject the morbid poison, together with the narcotic by which the attempt is made to arrest elimination.

Dr. Koch's experiments on guinea-pigs afford an instructive illustration of the influence of opium in

^{* &}quot;Traité pratique et analytique du Cholera Morbus, 1850."

preventing the escape of the cholera poison.* After introducing the cholera bacillus into the stomach of the animal, he injected into the peritoneal cavity a narcotic dose of tineture of opium—his object being to render "it possible for the comma bacillus to remain longer, and gain a footing in the intestine." In this manner he produced fatal cholera in 30 out of 35 guinea-pigs experimented on.

Koch's experiments have been repeated with similar results by Dr. Neil Macleod. † He induced cholera in 38 out of 54 guinea-pigs by injecting cholera bacilli into the stomach and opium into the peritoneum, the object being to check peristalsis and to allow time for the multiplication of the organisms. It must be obvious to any unprejudiced observer that the patients of those who endeavour to arrest the choleraic discharges by opium are subjected to the same kind of treatment as the before-mentioned guinea-pigs. There is, however, this important difference, all the opium injected into the peritoneum, or beneath the skin, is absorbed, but much of that taken by the mouth for the arrest of diarrhœa is ejected without entering the circulation; the latter method, therefore, is less deadly in its results than the former.

The following case affords an instructive illustration of the effect of two opposite modes of treatment on the same patient. In August 1866, I saw, with

^{*} British Medical Journal, January 1886, pp. 63-4.

⁺ Lancet, March 9, 1889.

Dr. Halse and his partner, a youth aged 15, who had been seized with diarrhoea after drinking the water from a pump in the Temple, water which was believed to have been the cause of other cases of cholera.

The diarrhea had been stopped by two doses of opium. Then after a few hours' cessation of the diarrhea he was again seized with purging, which, being stopped by opium, he rapidly passed into collapse.

When I first saw him at II A.M. on August 6, his eyes were sunk, his skin and tongue were icy cold, his pulse was too feeble to be counted, and cramps were very severe. There had been no purging for several hours, but on palpation and percussion the intestines were found to contain a considerable amount of fluid. There was a state of general torpor, and no natural effort to evacuate the bowels. He was ordered to take half an ounce of castor oil every two hours.

At 8 P.M. he was reported to have vomited once and to have passed three fœtid rice-water stools. The intestines were less distended, and he was free from cramps, but in other respects the condition was unchanged. He was to have one or two doses of oil during the night.

On the seventh, at 9.30 A.M., he had rallied completely, his countenance being natural and his skin warm. There had been five or six bile-tinged stools during the night, and the intestines were now every-

where resonant on percussion. After this he was drowsy for a few hours, and on the eighth a rash of roseola appeared on the trunk and limbs, but he made steady progress towards convalescence.

This case is more instructive than any number of mere statistics. The opium acted upon the boy as it did upon Koch's guinea-pigs. The diarrhea having been arrested a second time, collapse directly followed, and would have been fatal but for the evacuant action of the castor oil. I have never seen a case in which the life-saving action of a purgative was more manifest. If the patient had been simply "nursed and soothed," which some writers declare to be the only treatment for collapse, he would have gone to his grave with distended bowels, as many a collapsed patient has gone.

Dr. Halse had a senior partner who was much too old to learn. He could not be made to see that his opiate treatment had placed the patient's life in danger; and he looked on in a pitiful state of fear and trepidation, from a dread that castor oil would cause fatal exhaustion. This old gentleman was a type of a class of practitioners at that time very numerous, but now, it is to be hoped, nearly, if not quite extinct.

IX. Dr. Sutton's Report on the Post-mortem Appearances in the Stage of Collapse.

An important paper on the condition of the heart and lungs after death, in the collapse stage of cholera, was contributed to the fourth volume of the "London Hospital Reports," by the late Dr. Sutton. Two physicians of Guy's Hospital, who have since died, questioned the accuracy of my report of the postmortem appearances. They both denied that there was a great accumulation of blood in the right cavities of the heart and the pulmonary artery, and that the minute tissue of the lungs was remarkably anæmic after death in collapse.

One of them, who was notorious for treating grave subjects with levity, said of me (Lancet, January 19, 1867) "We believe him to be as honest as he is mistaken, and we can scarcely pay a higher compliment to his uprightness."

As I was not content to be considered honestly mistaken, it was satisfactory to find that Dr. Sutton's report of the post-mortem appearance in fifty cases of death in collapse, during the epidemic of 1866 was in exact agreement with the description which I had published, and the accuracy of which was denied by my opponents. Dr. Sutton says of the heart: "The right side was seen to be very much distended. This was particularly noticeable in the greatly distended condition of the right auricle and the auricular appendage. . . . The left ventricle

was usually contracted, often firmly so; and on cutting off the apex of the ventricle and squeezing it, the cavity was seen to be empty, or contained very little blood indeed."

"The lungs weighed very much less than normal. In some cases they were very pale in the anterior two-thirds, and of a darker colour in the posterior third. The pale portions on exposure to the air rapidly became of a scarlet colour. In other cases the lungs throughout were of a dark red colour, but both the pale and the dark-looking lungs were dry, and on pressure gave but very little blood. Dark thick blood was seen flowing out of the branches of the pulmonary artery."

This condition of the heart and lungs is the result of the stream of blood during the collapse stage of cholera having been arrested before it had reached the pulmonary capillaries; and extreme contraction of the arterioles is the only possible explanation of the arrest.

It is important to bear in mind that if the large veins of the neck are wounded in opening the chest, the distended right ventricle rapidly empties itself. Dr. Sutton mentions one case (loc. cit. p. 493) in which the right ventricle was thus emptied in two or three minutes. This result of wounding a vein gives the explanation of the marvellous relief afforded by venesection during choleraic collapse, as reported by several of the most trustworthy practitioners in India (see p. 10).

If the inspection of the body is delayed for many hours the distension of the right cavities is found to be less than when the examination is made soon after death. The obvious explanation is that the vital contraction of the pulmonary arterioles which arrests the circulation, ceases after death, and thus allows the elastic resiliency of the distended pulmonary artery and the right side of the heart to drive the blood onwards into the pulmonary capillaries and veins, and to the left side of the heart. The dark colour of the lungs which are light in weight and contain but little blood, is explained by the backward engorgement of the bronchial veins and capillaries, in common with the whole systemic venous system.

I gladly acknowledge that by this report Dr. Sutton, with whom, respecting another subject I had a sharp controversy, has made a valuable contribution to the morbid anatomy of cholera, which quite proves the accuracy of Dr. Parkes' original description of the appearances found after death in collapse, and has thus silenced further controversy on this subject.

X. Mr. Macnamara's Theories and Practice.

One of the most conspicuous and strenuous advocates of the astringent treatment of choleraic diarrhœa has been my friend Mr. N. C. Macnamara. He has written three books on cholera, and he

is the author of the article "Cholera" in the two editions of Sir Richard Quain's "Dictionary of Medicine."

It is interesting to compare and contrast the treatment recommended in his article with that advised by Dr. George Oliver in the article "Diarrhea" in the same dictionary. Mr. Macnamara says: "In the first stage of Asiatic cholera we should endeavour to stop the purging, and without doubt opium is the drug upon which we may with the greatest confidence rely for effecting this purpose." He goes on to say that he was in the habit of giving at once a pill containing a grain of opium and four grains of acetate of lead, and if the diarrhea continued, a second and even a third pill was given.

Dr. Oliver, on the other hand, says: "In choleraic diarrhœa the best results are obtained from castor oil guarded by a small dose of laudanum at the commencement, and repeated if the disease is severe, while astringents and opiates alone are withheld until the bowels are relieved of offensive materials."

Mr. Macnamara, with his fear of exciting vomiting, directs that "the patient should be prohibited from swallowing water or any other liquid than that which he gets from the ice," which he is allowed to take ad libitum. Dr. Oliver, on the contrary, says, "Vomiting should be encouraged by copious draughts of warm water, and if need be by emetics of mustard or ipecacuanha."

The two methods of treatment are as wide as the poles asunder. One endeavours to suppress the natural efforts to expel a poison, the other has for its object to aid those efforts; and the editor must surely have experienced some difficulty in allowing such irreconcilable articles to appear in his dictionary. He, doubtless, saw that they were antagonistic, and apparently left it to his readers to choose between them. Surely common-sense, apart from any profound theory, would pronounce in favour of Dr. Oliver's treatment.

Mr. Macnamara's voluminous writings, in the course of which he makes repeated reference to his large experience of cholera in India, have probably had considerable influence upon many of his readers. I propose, therefore, to set forth and to criticise some of the extraordinary pathological theories for which he has made himself responsible.

His first book, "A Treatise on Asiatic Cholera," was published in 1870, and is dedicated to the Duke of Argyle, at that time Secretary of State for India.

In this work his theory of cholera is in substance as follows. A poison, or as he says, a specific organic matter from a previous sufferer being swallowed, does not enter the blood, but by a local action destroys the epithelium of the intestinal canal. "The epithelium being thus destroyed the serum of the blood is allowed to drain away from the capillary arteries (sic) of the intestines, and at the same time venous absorption is prevented from taking place

through the intestines. Hence the symptoms of cholera. After a time, in favourable cases, the blood, by deprivation of its water having been greatly thickened, coagulates in the intestinal capillaries. This stops the further drain of liquid. Thus the epithelium is reproduced, and so absorption, which had been rendered impossible during the absence of the epithelium, recommences, the water is restored to the blood, and the patient recovers."

This strange, mechanical theory was based upon the observation of an eminent microscopist, that after death from cholera the intestinal villi were found denuded of epithelium.

This denudation was soon shown to be the result of maceration of the mucous membrane in the morbid contents of the bowels after death, so that the removal of the epithelium is a result, and not the cause of the escape of fluid into the intestines. But even if the denuding process had occurred during life, it is inconceivable that the copious outpouring of liquid from the blood could be so caused. The author has since admitted that the epithelium of the villi is not destroyed during life.

If the remarkable theory that the poison of cholera has a purely local action on the intestines without entering the blood, had a foundation in fact the practical inference would be that the noxious stuff should be expelled by a purgative, and not retained there by an opiate until it has destroyed the intestinal epithelium.

It is a peculiarity of Mr. Macnamara's method that every fact which appears to be inconsistent with his "dehydration," theory of collapse is denied or is questioned. Thus he says, that "if the body is examined soon after death from collapse, the left side of the heart contains as much blood as the right." He maintains that the distension of the right side of the heart depends on "the water of the tissues draining into the venous capillaries and right side of the heart after death in collapse, the whole venous system being full of blood from this cause." This is a good illustration of Mr. Macnamara's unphysiological method. Although he believes that during life the tissues are "wellnigh dehydrated" he imagines that after death they pour into the veins such an abundance of water as to distend the entire venous system.

The fact is that the sooner after death the inspection is made, the greater is the distension, not only of the right side of the heart and the whole systemic venous system, but of the pulmonary artery to its minutest subdivisions. The explanation of this I have before given. (See p. 39.)

In Quain's "Dictionary of Medicine," Mr. Macnamara gives his version of the condition of the heart's cavities, but without the astonishing explanation which I have just now quoted, and to this account of the appearances, the editor, in the second edition, appends the following foot-note:

"On this subject see the chapter on epidemic

cholera in Dr. George Johnson's Medical Lectures and Essays."

It would appear, therefore, that the editor doubted the accuracy of his contributor's discription of the post-mortem appearance in cholera. It is probable, too, that this editorial note made an impression on Mr. Macnamara, for in his history of Asiatic Cholera, which was published in 1892, probably after the article on "Cholera" was in print, the author says quite correctly: "The heart, as a rule, is found distended with dark blood on the right side, its left side being empty. The jugular veins, and the vena cava, together with the coronary veins, are full of blood, but the aorta and other arteries are empty, with the exception of the pulmonary artery." Then as to the lungs, he says they are "below their normal weight, and in the greater number of cases are collapsed, and found lying back against the spine. On section they appear dry, containing but little blood, and that is confined to the pulmonary arteries, and their branches, the capillaries and veins of the lungs are empty." It would have been more satisfactory if this correct description of the heart and lungs immediately after death in the collapse stage of cholera had been given in the article "Cholera" in Quain's "Dictionary of Medicine," the second edition of which was published in 1895. It is obvious that the appearances described are inconsistent with the theory that the collapse of cholera is the result of dehydration of the blood and tissues.

The sudden onset of collapse in the most malignant choleraic attacks is inexplicable by the most rapid dehydration process; therefore, Mr. Macnamara doubts the occurrence of such cases. In my "Notes on Cholera," I quote from a report on cholera in the Black Sea fleet in 1854, the statement that amongst robust English sailors "the attacks in many instances were so sudden that men fell as if they had drunk the concentrated poison of the upas tree." This he says is "high-flown language." But I find in Mr. Macnamara's treatise on "Asiatic Cholera" language even more sensational.

Thus at p. 17, in an extract from the "Proceedings of the Bengal Medical Board" for 1817 the disease is spoken of as "running its course generally in a few hours, and sometimes in a few minutes."

At p. 19 Dr. Corbyn reports that in 1818 "some of the Governor-General's servants dropped down dead behind his chair." And at p. 24 there is an extract from Jameson's report to the effect that at Bunderpoor in 1818 "the patients were described as having been knocked down dead as if struck by lightning." Surely lightning beats the upas poison in speed. Such cases must have been of exceptional malignancy; but the sudden onset is explicable by arterial contraction, as every one can understand who has studied the effect of breathing an azotic gas, such as nitrous oxide or nitrogen.

The result is, that in less than two minutes the circulation is completely arrested by the same nervo-vascular mechanism as that which stops the flow of blood in cholera. Again, in the Appendix to Mr. Macnamara's book, which he refers to as "a valuable record," "Dr. Bruce says: "It is still a favourite opinion with many, that the amount of discharge and collapse stand in the direct relation of cause and effect, but experience is opposed to such a conclusion; for who has not seen marked cases in every epidemic where there was neither purging nor vomiting sufficient to cause any degree of collapse, and yet these cases have sunk at least as rapidly as others where the discharges were profuse." He then refers to the case of a man who "stated that he had one copious fluid motion, after which, feeling unwell, he came to the hospital immediately. He vomited for the first time in my presence as he was taken out of the dooly. I found him in a state of collapse, and he was dead in six hours from that time, having been purged again only once: this last was a pure, flocculent congee motion, but it was only his second; and in the postmortem examination an unusually small amount of congee fluid was found in his intestines." Dr. Bruce also refers to some cases in which, "although the purging and vomiting were checked without difficulty, the collapse steadily increased." Yes, the collapse increased because the purging and vomiting were checked.

How long a time must elapse before it is universally known that no patient comes out of collapse without a continuance of the discharges in a greater or less degree?

If there are any facts which are inconsistent with Mr. Macnamara's theory, it is so much the worse for the facts. They are simply ignored. The continued secretion of milk by nursing mothers during collapse has been recorded by numerous observers, from Magendie in 1832 to the medical officers of the London Hospital in 1866. (See p. 8.)

Our author says: "With regard to the supposed secretion of milk in cholera, I can only say from my own experience that the mother's milk ceases as collapse comes on. If, as some state, it increases I cannot understand where its watery element comes from." I am not aware that any one has stated that it increases, but only that it is not suppressed, for reasons which I have given; but Mr. Macnamara, believing that little water is left in the body, has persuaded himself that the continued secretion of milk is impossible, and therefore doubts a repeatedly demonstrated fact.

But Mr. Macnamara's statement of the influence which loss of water has upon the buoyancy of the body is the most surprising of his many inaccuracies. In his last published work ("History of Asiatic Cholera," at p. 62) he says: "The blood and tissues of the body are well-nigh dehydrated, and if the patient is placed in a large bath it is not

easy to keep his body under the water; it floats like a cork." Here it is evident that the writer fails to distinguish the actual weight of the body from its specific gravity. Every schoolboy would know that in proportion to the decreased weight of the body by loss of water, its specific gravity must be increased; yet Mr. Macnamara writes as if he had actually seen the body of a choleraic patient floating like a cork. Since he misunderstands so simple a physical principle as this, we can appreciate his incapacity to interpret the more complex physiological phenomena of cholera.

In this same book there are indications that the author is, after all, losing faith in his own theory of collapse. He says (p. 59): "With reference to the poison which we believe the cholera bacillus produces, several theories have been advanced to explain its action. Among these theories that advocated by Sir George Johnson takes a prominent position, and by many pathologists is held to be best capable of accounting for the symptoms and the *post-mortem* changes found after death from cholera" (sic).

He then gives a brief but sufficiently accurate account of my theory, but follows it up by some futile objections, which are no more worthy of notice than his statement that the body of a cholera patient "floats like a cork." I have felt it to be a duty to deal thus seriously and at length with Mr. Macnamara's fanciful opinions; for as Sir

Thomas Watson says: "The subjects with which we have to deal are not matters of mere speculative curiosity or intellectual amusement—to be taken up to-day and dismissed perhaps with unconcern to-morrow; but they involve questions of life and death." No man with more than an elementary knowledge of physiology would have put forth such wild theories as those to which I have referred; but Mr. Macnamara's conviction, that the opiate treatment of cholera is the only successful method is so strong that he is blind to all facts and arguments which are opposed to that opinion. He quotes Dr. Koch's experiments on guinea-pigs, the object of the operator being, by means of an opiate injection, "to retain the cholera bacillus as long as possible within the intestinal canal." But he fails to see that by his opiate treatment he is repeating on the human subject an experiment, in principle, identical with that of Koch. It might be well if in future the repressive treatment of choleraic diarrhœa by opium were known by the distinctive name of Koch's guinea-pig treatment.

Mr. Macnamara in his "History of Asiatic Cholera" (p. 63), under the head of "Treatment" says: "Having gone through a severe attack of cholera I can testify to the unspeakable relief which ice affords to a person passing through an attack of this terrible disease. In my own case the onset of the disease was so sudden, and the vomiting so incessant, that everything I took was immediately rejected, and

consequently it was useless taking medicine; but I chewed and swallowed pounds of ice to my infinite comfort."

The author makes no mention of having had more than one attack of cholera, yet it is certain that the preceding account could not refer to the attack which he had when he was assisting me in the cholera wards in 1854, of which he gave me the following notes for publication a few days after his recovery:

"The following is the report of a severe attack of diarrhea I was seized with on September 4. I was in my usual state of health up to the evening of the 4th inst. and having taken a particular interest in the cholera patients in our hospital, I had for three or four days previous to the evening on which I was myself taken ill, almost lived in the cholera wards.

"On Monday night at about eleven o'clock I awoke with violent pain and spasm in my stomach, and a most distressing sense of oppression, with an inclination to vomit. These symptoms were quickly followed by cramps in my hands and feet. I at once got up and took an emetic, and as soon as the vomiting which it excited had ceased I followed it up with an ounce dose of castor oil. In about half an hour's time I was violently purged, and so much relief did I experience from it that I repeated the dose of oil during the following hour. From this time until eight o'clock in the morning I was purged

and vomited so freely that I became quite exhausted; yet I suffered little pain, and had lost that dreadful sense of oppression to which I before alluded. I then had two hours' sleep, and awoke much refreshed.

Throughout the day my bowels were opened several times, and I vomited once or twice after taking food; but in the evening I went down to Blackheath, passed a most comfortable night, and the next day returned to town ready to resume my duties, although much pulled down by my severe though short illness.

"C. N. M., M.R.C.S.

"KING'S COLLEGE HOSPITAL, "Sept. 1854."

"The dreadful sense of oppression" was probably due to the commencing embarrassment of the pulmonary circulation by the cholera poison. From this he was speedily relieved by the prompt recourse to evacuant treatment. If instead of the emetic and castor oil, of which, however, he took twice as much as there was need for, he had taken opium, the result would have been very different.

It is remarkable that Mr. Macnamara should for a number of years have been so strenuous, and, as I think I have shown, so unreasonable an opponent of a principle of treatment which was the means of saving his own life.

XI. The contrasted History of opposite Methods of Treatment.

The contrast between the history of the repressive and the evacuant treatment of cholera may be briefly stated.

The theory that loss of water is the chief cause of collapse was the result of an imperfect observation of facts. This theory suggested the employment of opium, and has blinded those who hold it to the evil results of the treatment. On the contrary, the evacuant treatment was successfully resorted to after the disastrous effects of opium had been witnessed.

Then, subsequently, a careful investigation of all the facts of the disease led to the conclusion that the main cause of collapse is a greatly impeded flow of blood through the lungs. When once it is clearly seen that this impeded circulation is a result of contraction of the pulmonary arterioles excited by a poison in the blood it is manifest that the attempt to prevent the escape of the poison by opium is irrational, and must be injurious. The different results of the two opposite modes of treatment—the astringent and the evacuant—are completely explained by the true theory of collapse.

I have never supposed or stated that evacuants of any kind are essential for the cure of choleraic diarrhea. The vis medicatrix natura suffices for the

recovery of most cases, if only the patients do not continue to drink the poisonous water which in many cases had excited the diarrhea. As many patients recover in spite of astringents, so many more would recover if they were treated by coloured water.

It has sometimes been asserted that an attack of cholera has been caused by a rhubarb pill or a dose of castor oil. I can give a parallel to this post hoc propter hoc argument. Many years ago, when I was seeing out-patients at the hospital, a woman brought a feverish child, for whom I prescribed a saline mixture. Two days afterwards the mother returned, and said in a complaining tone, "Your medicine has brought out the small-pox." Truly the small-pox had come out, but assuredly it was in the blood before it appeared in the skin. So when symptoms of cholera follow the action of an aperient the morbid poison was before in the blood, and caused the derangement of health for which the dose was taken. It is about as probable that small-pox could result from a croton-oil liniment, as that a specific disease like cholera could be caused by a purgative.

XII. The Action of various Remedies.

Some practitioners look upon diluted sulphuric acid as a valuable remedy for diarrhœa, and call it an astringent. If astringent at all, it is so in a different manner and degree from opium, and when

given in full doses it often excites griping and diarrhea. On the other hand, a chemically opposite remedy—namely, carbonate of soda, in mint tea—was reported to be very efficacious by the late Mr. Wakefield, who stated in a letter to the *Times* that during the epidemic of 1854 he had thus treated a large number of cases of diarrhea amongst the prisoners in Cold Bath Fields, and not one had passed into collapse.

Some practitioners are of opinion that the undoubted efficacy of calomel is due to its direct action on the liver, while others suppose that it has an antiseptic influence; but it would seem that even the largest doses of this insoluble compound that could, with safety, be given would have but an infinitesimal antiseptic influence on the copious intestinal discharges. Calomel in powder is an excellent substitute for castor oil when frequent vomiting prevents the retention of the latter. In such cases calomel appears to arrest the vomiting by reversing the action of the stomach. A sagacious friend of mine said to me many years ago something to this effect: "Many men would have more faith in castor oil if you could speak of it as being not merely an evacuant, but as having a specific castoroilative influence." I can understand this craving for a specific curative agent, but I do not share the feeling.

XIII. Dr. Burney Yeo and Dr. Clemow on the General Adoption of the Eliminative Treatment.

I have the satisfaction of finding that, after forty years of controversy, the evacuant or eliminative treatment is now being generally looked upon not only as rational and scientific, but in practice most successful.

Dr. Burney Yeo in his "Manual of Medical Treatment" (4th ed., vol. ii. p. 696), after referring briefly to the pathological theory, says: "It will be seen from these indications that the rational and scientific treatment of cholera is now admitted nearly on all sides to be, as Sir George Johnson insisted many years ago, mainly eliminative and antiseptic. 'Purgation and antisepsis are to some extent interchangeable terms.' When cases of cholera are seen in the first stage, when nature is making an effort to throw off the poison by the bowels, we must aid the natural effort by the administration of an unirritating purgative; and if we can at the same time combine with it an intestinal antiseptic, we shall be fulfilling very completely the first indication. Castor oil is one of the best purgatives for this purpose. Calomel is also valuable, and it is an antiseptic as well as a purgative. Gregory's powder, together with a few grains of calomel, is suggested by Sir George Johnson if, owing to vomiting, castor oil cannot be taken."

"Treatment on these lines in the early stage has been found to be the most successful in the recent epidemics in Germany, Russia, and Italy. One Russian physician began the treatment with twenty grains of calomel with an ounce of castor oil, and he reports that the mild cases recovered quickly, and so did many in the algid stage. Another started with a purgative dose of calomel combined with naphthalin, and continued with smaller doses of each. Hamburg mild threatening cases were stopped by an initial dose of castor oil, and in more advanced stages calomel in 12 grain doses, or in repeated small doses of 1/8 to 1/6 grain, was found to answer well. In severe cases, those of well-marked general intoxication, we are informed, 'calomel was the only drug that held its own.'

"In the reports of recent epidemics there is a general condemnation of the use of opium to arrest the diarrhœa in the early stage; cases so treated did worse than under any other treatment. The early use of opium increases the risk of the absorption of the cholera poison, and favours its retention within the body."

This, then, according to Dr. Burney Yeo, is the principle of treatment which is now nearly on all sides admitted to be not only rational and scientific, but successful. I confess that I read this statement with much satisfaction, for I know that Dr. Yeo has had ample opportunities to form a sound judgment as to the general opinion of the profession on this

subject; and his excellent practical work has had a very large sale, having gone through four editions within two years; it must now therefore be in the hands of several thousands of practitioners.

Dr. Frank Clemow, an English physician practising in St. Petersburgh, has published an interesting work on "The Cholera Epidemic in 1892 in the Russian Empire." In this work he says (p. 90): "The diarrhœa being looked upon as a safety-valve whereby the system got rid of large quantities of the poison that would otherwise be absorbed, was within certain limits encouraged. No attempt was, as a rule, made to check it by opium or astringents." And in a private letter with which I have been favoured by the author he says: "The St. Petersburgh physicians were in entire accord with your own teaching as to the necessity for purgatives as opposed to astringents in the treatment of cholera."

If the general body of Continental practitioners have been more ready than the English to adopt the evacuant treatment of cholera, this may perhaps be due to the fact that the dehydration theory of collapse, which suggests a repressive treatment, has never been so generally accepted abroad as at home, where, for a time, it did an incalculable amount of mischief.

The final result, then, of the cholera controversy is that the dehydration theory of collapse has been replaced by one more in accordance with facts and with physiology, and the injurious astringent treatment has been superseded by the beneficent principle of elimination. In other words, it has at length come to be generally acknowledged that it is a more rational and a more successful practice to assist than to impede the ejection of a morbid poison.

In 1892 the Local Government Board requested the Royal College of Physicians to give directions for the treatment of diarrhee and cholera. Detailed instructions were drawn up by the President, Sir Andrew Clark, and accepted, with little modification, by a large meeting of the Fellows. Amongst other directions is the following: "As soon as possible after looseness of the bowels has begun, take in capsules, or in hot milk, or in any other manner preferred, two teaspoonfuls of castor-oil."

This was a step timidly taken in the right direction, which a few years before would have been disapproved, and it was followed by detailed instructions which at the present time—so rapid has been the change of opinion and practice—would be considered unnecessary and even undesirable.

XIV.

DIRECTIONS FOR THE TREATMENT OF DIARRHŒA AND CHOLERA.

The leading principle that should guide the treatment of cholera has been clearly indicated in the preceding pages, but I have thought it desirable to give the following detailed directions for the treatment of the disease in all its stages.

Diarrhæa, during an epidemic season, is in many instances, though not in all, an early stage or a mild form of cholera; and in the great majority of cases of actual cholera, an attack of bilious diarrhæa marks the onset of the disease. A diarrhæa, when it is not the actual beginning of cholera, will weaken the patient, and so may predispose him to suffer from the more serious form of disease. Diarrhæa, therefore, during an epidemic season, ought not to be neglected even for an hour. That plan of treatment for diarrhæa is obviously the best which most speedily and completely puts a stop to the disease, without subsequent ill effects.

It may be stated as a general proposition, that

the immediate cause of diarrhea or looseness of the bowels is the presence of offending materials in the alimentary canal. These offending materials are of various kinds in different classes of cases. In one case, unwholesome and undigested food is the exciting cause of the purging; in another case, a large and unnatural accumulation of the fæculent contents of the bowel; while, in another class of cases, noxious secretions are poured from the blood into the bowel, in consequence of the action of a morbid poison upon some of the ingredients of the blood. To this last class of cases belongs what is called cholerare diarrhea.

The most reasonable theory of choleraic diarrhoea is that a morbid poison enters the blood either with the air through the lungs, or with the food and drink through the alimentary canal; and that this poison excites certain changes in the blood, in consequence of which some blood-materials are spoiled, and thus rendered not only useless, but noxious. These morbidly changed blood-materials are then discharged from the blood-vessels through the mucous membrane of the stomach and bowels, and are ultimately ejected by vomiting and purging.

Various as are the remote and primary causes of diarrhoa, this one condition is common to all classes of cases—viz., that the contents of the bowel are unnatural and offensive. These offending materials are the immediate cause of the purging; and they

must be expelled from the bowel before the diarrhœa can come to an end.*

From the above considerations we deduce one important and guiding rule of treatment, which is this: not to attempt by opiates, or by other directly repressive means, to arrest a diarrhæa while there is reason to believe that the bowel contains a considerable amount of morbid and offensive material. It is certain that these offending materials must be cast out from the bowel before the diarrhoea can permanently cease. The effect of an opiate at this stage, if, as frequently happens, its absorption be not prevented by the rapid outward flux of liquid from the blood into the stomach and bowels, is to prolong the disease, and to increase the risk of mischief, from the retention and re-absorption of the morbid contents of the bowel. If the opiate have the effect of retaining within the blood-vessels some of the morbidly changed blood-constituents, this astringent action will probably be more injurious and even deadly than the retention of morbid secretions within the bowel. The abrupt arrest of the discharges by opiates is not infrequently followed by fatal collapse.

The purging is the natural way of getting rid of the irritant cause. We may favour the recovery by directing the patient to drink copiously any simple diluent liquid—water cold or tepid, toast-

^{*} We need not here take into consideration those cases of diarrhœa which result from ulceration or other local disease of the bowel itself.

water, barley-water, or weak tea; and we may often accelerate the recovery by sweeping out the alimentary canal by some safe purgative, and then, if necessary, soothing it by an opiate. Castor oil, notwithstanding its unpleasant taste, is, on the whole, the safest and best purgative for this purpose. It has the advantage of being very mild and unirritating, yet withal very quick in its action. A tablespoonful of the oil may be taken, floating on cold water or in hot milk or any other simple liquid which may be preferred by the patient. A mixture of orange-juice or of lemon-juice with water forms an agreeable vehicle for the oil. Some prefer to take it with weak brandy and water, and there is no serious objection to this mixture, but it is very likely to excite vomiting. Another mode of giving the oil is in the form of an emulsion, with mucilage and cinnamon water, which very completely covers and disguises the taste of the oil. If the dose be vomited, it should be repeated immediately; and the patient should lie still, and take no more liquid for half an hour, by which time the oil will have passed from the stomach into the bowels. Within an hour or two, the oil will usually have acted freely. Then a tablespoonful of brandy may be taken in some thin arrowroot or gruel; and if there be much feeling of irritation, with a sense of sinking, from five to ten drops of laudanum may be given in cold water. These means will suffice for the speedy arrest of most cases of choleraic diarrhœa.

If the patient have an insuperable objection to castor oil, or if the oil cannot be retained on the stomach, ten or fifteen grains of powdered rhubarb, or a tablespoonful of the tincture of rhubarb, or a teaspoonful of Gregory's powder, may be substituted for the oil.

When vomiting is very frequent three grains of calomel in powder, washed down with cold water, may be the best form of purgative.

If the diarrhea has continued for some hours, the stools having been copious and liquid; if there is no griping pain in the bowels, no feeling or appearance of distension of the intestines—the abdomen being flaccid and empty, and the tongue clean-we may conclude that the morbid agent has already purged itself away. There will, therefore, be no need for the castor oil or other laxative, and we may immediately give the brandy in arrowroot, and the laudanum, as before directed. The rule in all cases is, not to give the opiate until the morbid poison and its products have for the most part escaped; not to close the door until "the enemy" has been expelled. While there are some cases in which the evacuant dose is not required even at the commencement of the attack, there are many more in which the opiate is unnecessary in the later stage. In some cases of severe and prolonged diarrhœa it may be necessary to repeat the oil and the laudanum alternately more than once, at intervals of three or four hours. Practical skill and

tact are required to discriminate these cases. It must be borne in mind that when the choleraic secretions are being actively poured out from the blood-vessels, the bowel, though it may have been completely evacuated by a dose of oil, may quickly again become filled with morbid secretions, and hence the need for an occasional repetition of the evacuant dose.

With reference to the use of opiates in the treatment of diarrhea, the principle to be kept constantly in view is this: opiates are useful to soothe irritation after the evacuation of the bowel; they are useless and even dangerous when the blood is poisoned or the bowel filled with morbid secretions. Opiates, in the early stage of diarrhoea would be more frequently and decidedly injurious were it not for the fact that, their absorption being prevented by the active eliminative efforts, they are quickly expelled with the morbid secretions, and they are therefore powerless to arrest the discharges. There is one symptom for which opium has often been given, but which especially forbids its use, and that symptom is cramp of the muscles. This cramp shows that the poison which excites it is still in the blood. The painful spasms can be effectually relieved only by the escape of the poison. An opiate dose might hinder the exit of this poison from the blood, and thereby induce that perilous, though painless, spasm of the small arteries of the lungs which is the essential cause of collapse.

It has been a common practice to give calomel and opium combined. The effect of this combination will depend upon the proportions of the ingredients. Equal weights of opium and calomel will have, for the most part, a narcotic and astringent action, but in proportion to the preponderance of the calomel the operation will be evacuant and purgative.

If the diarrhea be associated with vomiting, this should be encouraged and assisted by copious draughts of tepid water. The vomiting affords relief partly by the stimulus which it gives to the circulation, but mainly by the speedy ejection of morbid secretions.

If there be nausea without vomiting, and more especially if the stomach be supposed to contain undigested or unwholesome food or morbid secretions, an emetic may be given—either a teaspoonful of powdered mustard or a tablespoonful of common salt, or twenty grains of ipecacuanha powder in warm water.

When vomiting is excessive in violence or in frequency, it may sometimes be checked by small draughts of iced water at short intervals, or by lumps of ice taken and allowed to melt in the mouth. Another plan is to place three grains of calomel on the tongue and wash it down with cold water. The calomel probably acts by reversing the action of the stomach and bowels, and thus causing evacuation downwards. When given as a powder it

is less likely to be rejected by the act of vomiting than when made into a pill.

Thirst may be allayed by drinking cold water, which may be acidulated by the addition of lemonjuice or a few drops of aromatic sulphuric acid. Care should be taken that the water for drinking is pure. Organic impurities, such as result from the admixture of sewage, are especially to be dreaded. If the water be of doubtful purity, it should be carefully filtered through sand and charcoal, and then boiled. Impure water is a common exciting cause of cholera.

While the diarrhea continues, the diet should consist mainly of milk, rice, arrowroot, gruel, broth, or beef-tea.

In all cases of severe diarrhœa the patient should remain in bed.

If, while the purging continues, the stools become colourless and watery (the purging being of the kind commonly called rice-water purging), or if, without a continuance of the purging, the surface of the body become cold and blue, the disease is now passing, or has actually passed, into the stage of collapse.

In the worst forms of collapse, as cholera presents itself often at the outbreak of an epidemic, the disease is so deadly that no treatment is of any avail; and a sure way to bring discredit upon any method of treatment is to apply it in this desperate class of cases.

Choleraic collapse, as I have before explained, results from a peculiar arrest of the flow of blood through the lungs, occasioned by a morbid poison. It is not a condition of mere exhaustion. It is not relieved by the remedies for exhaustion; and it is made worse by opiates and by spirituous stimulants, which must therefore be avoided. The injurious influence of opium and alcohol in the collapse stage is now almost universally admitted. Their noxious effects are intelligible, if we bear in mind that during the collapse the oxidation of the blood and tissues is greatly diminished, and that both opium and alcohol still further impede oxidation and thus add to the peril, while opium also retards the escape of the poison.

The patient should be strictly kept in the recumbent position; he must not be raised even to go to stool. He should be abundantly supplied with fresh air, and should be allowed to drink pure water freely. The water may usually be taken cold, but it should not be iced, except occasionally when given to check excessive vomiting. Large quantities of iced water may do harm, by chilling the patient and checking the natural eliminative efforts. In cases of extreme collapse I would persuade the patient to drink hot water, for the purpose of warming the blood and quickening the circulation. Some care is required not to over-distend the stomach by liquid. Unless vomiting occurs from time to time, the drinking of large quantities of liquid may so distend the stomach

as to impede the breathing, and thus cause much distress. The circulation has sometimes been improved by the injection of large quantities of hot water into the rectum.

Hot flannels, or bottles, or bags of sand, should be applied to the feet and legs.

Cramps in the muscles are, as a rule, more frequent and severe during the diarrhea stage than during the stage of collapse. Whenever they occur, they may be relieved by rubbing the affected parts with the warm hand. There can be no objection to the use of anodyne or stimulating embrocations, but they are of doubtful advantage. Cramps in the legs may often be relieved by enveloping the limbs in flannel wrung out of hot water, with a covering of macintosh or oiled silk.

Hot baths, whether of water or of air, although they relieve the painful cramps and often effect a temporary improvement of the circulation, have been found to be, on the whole, more distressing and exhausting than beneficial.

Five grains of sesquicarbonate of ammonia, or a teaspoonful of spirit of sal volatile, may be given in an ounce of camphor mixture every two or three hours as a diffusible stimulant.

The discharges from the bowels, and the condition of the abdomen, should be carefully observed. In favourable cases the discharges always continue, more or less, during the stage of collapse and until reaction has set in. One of the earliest and surest signs of reaction is the reappearance of bile in the vomited matters and in the stools. When vomiting and purging entirely cease during the stage of collapse, the disease is nearly always fatal.

One of the main objects of treatment during this stage is to facilitate the escape of the morbid secretions from the alimentary canal. This may be done partly by the copious use of diluent drinks, and partly by an occasional dose of castor oil. If we carefully observe the condition of a patient in collapse, we shall often find that the intestines are more or less distended with liquid, and this, too, while perhaps there is general torpor and little or no effort at expulsion, sometimes the result of opium, but it may be due to the defective oxygenation during collapse. Again, it has often been found that, although there has been copious watery purging during life, the small intestines contain after death a large amount of a peculiar viscid dirty white material, having a very offensive odour. An occasional dose of castor oil may be useful in removing both these conditions: namely, over-distension of the bowel by liquid, and accumulation and retention of offensive viscid semi-solid secretions.

I have no doubt that by this treatment I have rescued patients who would have died if they had been left to the unaided efforts of nature. I can lay down no rule as to the number of doses of oil that may be required. It should be borne in mind that the object and the effect of the evacuant or

cleansing treatment are not to increase the amount of liquid which is poured from the blood into the stomach and bowels, but simply to assist and to quicken the expulsion of the morbid secretions from the alimentary canal.

It may be confidently maintained that, inasmuch as the peculiar choleraic discharges are the result of specific blood-changes, induced by a morbid poison, no ordinary purgative can *increase* those discharges, although it may facilitate and quicken their expulsion.

The volume of liquid which is discharged from the stomach and bowel, is, in the diarrhæa stage, a measure of the dose or the virulence of the poison, just as, in the case of diabetes, the flux of urine is equivalent to the amount of sugar that has to be discharged through the kidneys. During the stage of collapse the liquid discharges cease to be an index of the severity of the disease, for the reason that the partial arrest of the circulation impedes, more or less, the excretion of the poison and its products through the alimentary canal, and in the same degree increases the risk of a fatal result.

Bleeding from the bowel sometimes occurs. It is a very unfavourable and usually a fatal symptom. The best treatment consists in giving twenty drops of oil of turpentine in mucilage every two hours, and iced water may be taken freely. In such a case no castor oil should be given.

After reaction has occurred, an occasional laxa-

tive dose is required—about once in the twenty-four hours during the first two or three days.

It is worse than useless to attempt to feed a patient during collapse. The secretions of the stomach are utterly deranged; and the power of digestion is suspended. The mildest nourishment administered at this time only adds to the feeling of oppression and general distress, from which the act of vomiting often gives immediate relief.

After reaction has occurred, and when the normal secretions are restored, the mildest nourishment should be given frequently, but in small quantities—such as milk, gruel, or rice, or arrowroot with a small quantity of brandy, soup or beef-tea, or chicken-broth. After an attack of cholera the stomach is sometimes long in recovering its tone and the power to digest solid food. When this is the case, a grain of quinine, with ten or fifteen drops of dilute hydrochloric or sulphuric acid and an equal quantity of spirit of chloroform, may be taken with each meal.

Venesection has often afforded great relief during the stage of collapse. The case recorded by Sir Ranald Martin affords a striking illustration of this (see p. 10). Many similar cases are recorded in the writings of practitioners in India. The symptoms which appear especially to call for venesection are rapid breathing, and a feeling of impending suffocation. When, with these symptoms, there is a cessation of vomiting and purging, which is probably a result of the almost entire arrest of the circulation through the lungs, venesection may prove a life-saving remedy. It is difficult to obtain a stream of blood in these cases; not, as many suppose, because the blood is too thick to flow, but because, in consequence of the block in the lungs, the blood in the veins is nearly stagnant. The bleeding appears to be beneficial chiefly by lessening the distension of the right cavities of the heart, and so increasing their contractile power. Repeated doses of ammonia may help to quicken the circulation, and so favour the flow of blood.

Hot saline injections into the veins have unquestionably saved life in some apparently desperate cases. This is an operation the performance of which requires great skill and care. I have already explained its mode of action (p. 10).

Consecutive Fever.—Reaction from collapse is frequently followed by a febrile condition—a hot skin, quick pulse, coated tongue, hurried breathing, often a scanty secretion or even a complete suppression of urine, with drowsiness tending to pass into coma. These unfavourable symptoms are more common when, during the earlier stages of the disease, opium and alcoholic stimulants have been freely given; but they may occur when no such means have been employed.

The best treatment consists in a scanty liquid diet without alcohol, copious diluent drinks, with saline effervescing draughts, an occasional aperient, castor

oil, or sulphate of magnesia or soda or a seidlitz powder, counter-irritation over the lungs and kidneys, and sometimes local bleeding by leeches or cupping to relieve congestion of those organs.

When the urine is suppressed or scanty, one of the safest and most efficacious diuretics is the "Imperial Drink," made as follows: A quarter of an ounce of cream of tartar, the juice of one lemon, the outer peel of half a lemon, a pint of boiling water and sugar ad libitum. One or two pints of this may be taken cold in the course of the twenty-four hours. Alcoholic stimulants should be withheld until the function of the kidneys has been completely restored.

It will be seen that the treatment which I recommend, and which has been practised by myself and others with great success, is not a system of mere drug-giving, but a definite plan based upon a careful study of the natural history of the disease. I am confident that any one who, having an intelligent appreciation of the pathological conditions to be dealt with, will carry out this method of treatment with the same determination to arrive at a definite result as that with which he would treat a case of poisoning by opium, will soon have better reasons for continuing the practice than any with which I can furnish him. For the full success of the treatment it is necessary that the practitioner himself should have that confidence in the method without which it can never be fairly tried, and that he should

inspire his assistants, and above all his patients, with hope and confidence.

Preventive Measures.—The choleraic discharges from the bowels should be looked on as highly poisonous, and they should be disinfected and got rid of as soon as possible. Every vessel and article of clothing or bedding soiled by the discharges should be carefully cleansed and disinfected. The attendants on the sick should be warned of the necessity for thorough ventilation and for extreme personal cleanliness. The hands should be frequently cleansed with the aid of disinfectants, and always immediately before taking food.

If these simple measures are adopted, nurses and other attendants on the sick run little risk from infection,

The chief disinfectants are—chloride of lime, Burnett's liquid, Condy's fluid, and Calvert's solution of carbolic acid. These disinfectants are sold with full printed directions for the use of each. Diluted Condy's fluid is well adapted for cleansing the mouth and hands before taking food; and carbolic acid for cleansing bedding and clothing, which would be damaged by mineral disinfectants. Any one of the above may be used for disinfecting the stools.

A concentrated solution of sulphate of iron is also a cheap and efficient disinfectant for the discharges.

The meals should be taken at regular and not

too great intervals. Long fasting and over-fatigue should be carefully avoided.

Great moderation both in food and in drink is essential for safety during an epidemic of cholera. A single act of indiscretion has been followed by a severe attack. Intemperance at such a time is fraught with extreme danger.

Unwholesome articles of food, more especially tainted meat and fish and decayed vegetables, are to be carefully avoided. It is a mistake to suppose that entire abstinence from fruit and vegetables is necessary or wholesome during a cholera season. Ripe fruit and fresh vegetables may be taken in moderation with safety and advantage.

Especial attention should be paid to ensure the cleanliness and thorough ventilation of dwelling-houses. All vegetable and animal refuse should be removed as speedily as possible. Care should be taken to prevent the escape of sewer poison into the interior of dwellings.

The purity of the water employed for drinking and cooking should be most carefully provided for. A few drops of Condy's fluid may be used as a test for the purity of water. Organic impurities in the course of an hour or two decolorise the fluid; which is not only a test, but also a purifying agent by oxidising the organic impurities. It is certain that a diarrhœa is often perpetuated by the daily drinking of impure water; and it is equally certain that to arrest a diarrhœa having such an origin by

opiates and astringents is a most dangerous practice. I have seen instances in which a diarrhoea, probably caused by drinking foul water, having been arrested by opium, was quickly followed by an attack of cholera of the most formidable character.

No unnecessary medicines of any kind should be taken. When opening medicine is required, the mildest should be selected, such as castor oil or rhubarb. Saline purgatives, such as Glauber's salts and Epsom salts, are objectionable, on account of their tendency to cause profuse watery purging. On the other hand, the common belief that prolonged costiveness should not be interfered with during the prevalence of cholera is an error. An accumulation of offensive materials within the bowels may be itself a source of irritation and of danger.

Some practitioners believe and assert that cholera collapse may be caused by a dose of purgative medicine. This I entirely disbelieve. It must have happened frequently that an individual, experiencing the *malaise* which results from the cholera poison in the system, takes a dose of opening medicine, the symptoms of cholera develop themselves, and the disease may be supposed to have been caused by the medicine; yet such a conclusion would be as inconsistent with facts and with analogy as to attribute the outcoming of the rash of scarlatina to a mustard poultice applied to the

neck for the relief of the sore throat which commonly marks the commencement of that disease. Cholera and scarlatina are both specific contagious diseases; one could no more be caused by a purgative than the other by a mustard poultice.

It is important to know that cholera may be imported into a district by a person coming from an infected locality while he is suffering from diarrhea. The diarrhea may be a mild form of cholera, and the discharges will then contain the specific poison of the disease. If these discharges are permitted to contaminate the soil, the water and the air, the disease may spread with destructive rapidity. To prevent an outbreak of cholera, the diarrhea patient should, if possible, be kept in quarantine, the discharges should be disinfected, and then buried, and all soiled clothing and bedding should be carefully disinfected and cleansed or burnt.

Public latrines in a district where cholera prevails are highly dangerous and a fruitful source of infection. The use of latrines by British troops in India has often led to a great mortality, while the native troops, who never resort to latrines, have either escaped entirely or have suffered in a much less degree, at the same time and place.

The early burial of those who have died from cholera is imperative as a sanitary measure, more especially in a tropical climate, where decomposition is extremely rapid. Of course care must be taken to ascertain that life is really extinct before burial takes place. And here it may be well to mention that there are two post-mortem phenomena which the inexperienced might mistake for signs of vitality. The internal heat of the body, which even during collapse is usually above normal, comes to the surface after death, so that for an hour or two the temperature of the skin may be higher than it was for some hours before death. Again, it not infrequently happens that for half an hour or so after death some muscles, probably stimulated by the poisoned blood which excited the painful cramps during life, are affected by irregular contractions, which may be sufficiently powerful and concerted to move a finger and even a limb. These purely physical post-mortem movements have often excited a vague terror, and a suspicion that life remains in the moving corpse.

A SELECTION

FROM

J. & A. CHURCHILL'S CATALOGUE,

COMPRISING

MOST OF THE RECENT WORKS PUBLISHED BY THEM.

N.B.-J. & A. Churchill's larger Catalogue, which contains over 600 works, with a Complete Index to their Subjects, will be sent on application.

Human Anatomy:

A Treatise by various Authors. Edited by HENRY MORRIS, M.A., M.B. Lond., F.R.C.S., Surgeon to, and Lecturer on Surgery at, the Middlesex Hospital. Roy. 8vo, with 791 Illustrations, nearly all original, and many of them in several colours, 40s. (In one vol. or in three parts.)

Heath's Practical Anatomy:

A Manual of Dissections. Eighth Edition. Edited by WILLIAM ANDERSON, F.R.C.S., Surgeon and Lecturer on Anatomy at St. Thomas's Hospital, Examiner in Anatomy for R.C.P. and S. Crown 8vo, with 329 Engravings, 15s.

Wilson's Anatomist's Vade-Mecum. Eleventh Edition. By HENRY E. CLARK, M.R.C.S. Eng., F.F.P.S. Glasg., Examiner in Anatomy, F.P.S., and Professor of Surgery in St. Mungo's College, Glasgow. Crown 8vo, with 492 Engravings and 26 Coloured Plates, 18s.

An Atlas of Human Anatomy.

By RICKMAN J. GODLEE, M.S.,
F.R.C.S., Surgeon and late Demonstrator of Anatomy, University College
Hospital. With 48 Imp. 4to Plates (112
figures), and a volume of Explanatory
Text. 8vo, £4 14s. 6d.

Human Osteology.

By LUTHER HOLDEN, Consulting Surgeon to St. Bartholomew's Hospital. Seventh Edition, edited by CHARLES STEWART, Conservator of the Museum R.C.S., and ROBERT W. REID, M.D., F.R.C.S., Professor of Anatomy in the University of Aberdeen. 8vo, with 59 Lithographic Plates and 75 Engravings. 16s.

Also.

Landmarks, Medical and Surgical. Fourth Edition. 8vo, 3s. 6d.

The Student's Guide to Surgical
Anatomy. By EDWARD BELLAMY,
F.R.C.S. and Member of the Board of
Examiners. Third Edition. Fcap. 8vo,

with 81 Engravings. 7s. 6d.

Diagrams of the Nerves of the Human Body, exhibiting their Origin, Divisions, and Connections, with their Distribution to the Various Regions of the Cutaneous Surface, and to all the Muscles. By Sir W. H. FLOWER, K.C.B., F.R.S., F.R.C.S. Third Edition, with 6 Plates. Royal 4to, 12s,

Pathological Anatomy of Diseases. Arranged according to the nomenclature of the R.C.P. Lond. (Student's Guide Series). By NORMAN MOORE, M.D., F.R.C.P., Assistant Physician and Lecturer on Pathological Anatomy to St. Bartholomew's Hospital. Fcap. 8vo, with 111 Engravings, 8s. 6d.

A Manual of Clinical and Practical Pathology. By W. E. WYNTER, M.D., M.R.C.P., F.R.C.S., Medical Registrar to Middlesex Hospital, and F. J. WETHERED, M.D., M.R.C.P., Assistant Physician to Victoria Park Hospital. With 4 Coloured Plates and 67 Engravings. 8vo, 12s. 6d.

Lectures on Pathology:

Delivered at the London Hospital. By the late Henry Gawen Sutton, M.B., F.R.C.P., Physician to, and Lecturer on Pathology at, the London Hospital. Edited by Maurice E. Paul, M.D., and Revised by Samuel Wilks, M.D., LL.D., F.R.S. 8vo, 15s.

General Pathology:

An Introduction to. By John Bland Sutton, F.R.C.S., Sir E. Wilson Lecturer on Pathology, R.C.S.; Assistant Surgeon to, and Lecturer on Anatomy at, Middlesex Hospital. 8vo, with 149 Engravings, 14s.

Atlas of Pathological Anatomy. By Dr. LANCEREAUX. Translated by W. S. GREENFIELD, M.D., Professor of Pathology in the University of Edinburgh. Imp. 8vo, with 70 Coloured

Plates, £5 5s.

Pathologicus, for the Registrations of the Lesions recorded in Pathological Records or Case-books of Hospitals and Asylums. By James C. Howden, M.D., Superintendent of the Royal Lunatic Asylum, Montrose. Fcap. folio, 6s.

Atlas of the Central Nervous System. From the larger work of Hirschfeld and Léveillé. Edited by HOWARD H. TOOTH, M.D., F.R.C.P., Assistant Physician to the National Hospital for the Paralysed and Epileptic. With 37 Plates carefully coloured by Hand. Large Imp. 8vo, 4os.

The Human Brain:

Histological and Coarse Methods of Research. A Manual for Students and Asylum Medical Officers. By W. BEVAN LEWIS, L.R.C.P. Lond., Medical Superintendent, West Riding Lunatic Asylum. 8vo, with Wood Engravings and Photographs, 8s.

Elements of Human Physiology. (Student's Guide Series.) By ERNEST H. STARLING, M.D., M.R.C.P., Joint Lecturer on Physiology at Guy's Hospital. Fcap. 8vo, with 94 Engravings,

6s. 6d.

Manual of Physiology:

For the use of Junior Students of Medicine. By GERALD F. YEO, M.D., F.R.S., Emeritus Professor of Physiology in King's College, London. Third Edition. Crown 8vo, with 254 Engravings (many figures), and Plate of Spectra, 14s.

Principles of Human Physiology. By W. B. CARPENTER, C.B., M.D., F.R.S. Ninth Edition. By By HENRY POWER, M.B., F.R.C.S. 8vo, with 3 Steel Plates and 377 Wood Engravings, 31s. 6d.

Practical Lessons in Elementary Biology, for Junior Students. By PEYTON T. B. BEALE, F.R.C.S., Lecturer on Elementary Biology and Demonstrator in Physiology in King's College, London. Crown 8vo, 3s. 6d.

Medical Jurisprudence:

Its Principles and Practice. By ALFRED S. TAYLOR, M.D., F.R.C.P., F.R.S. Fourth Edition, by Thomas Stevenson, M.D., F.R.C.P., Lecturer on Medical Turisprudence at Guy's Hospital. 2 vols. 8vo, with 189 Engravings, 31s. 6d.

By the same Authors.

A Manual of Medical Jurisprudence. Twelfth Edition. Crown 8vo, with 55 Engravings, 14s.

The Student's Guide to Medical Jurisprudence. By John Aber-CROMBIE, M.D., F.R.C.P., Physician to Charing Cross Hospital. Fcap. 8vo, 7s. 6d.

Sanitary Examinations

Of Water, Air, and Food. A Vade-Mecum for the Medical Officer of Health. By Cornelius B. Fox, M.D., F.R.C.P. Second Edition. Crown 8vo, with 110 Engravings, 12s. 6d.

Microscopical Examination of Drinking Water and of Air. By J. D. MACDONALD, M.D., F.R.S., Ex-Professor of Naval Hygiene in the Army Medical School. Second Edition. 8vo,

with 25 Plates, 7s. 6d. Hygiene and Public Health.

A Treatise by various Authors. Edited by Thomas Stevenson, M.D., F.R.C.P., Lecturer on Chemistry and Medical Juris-prudence at Guy's Hospital; Official Analyst to the Home Office; and SHIRLEY F. MURPHY, Medical Officer of Health of the County of London. In 3 vols., royal 8vo, fully Illustrated. Vol. I., 28s.;

Vol. II., 32s.; Vol. III., 20s. A Manual of Practical Hygiene. By the late E. A. PARKES, M.D., F.R.S. Eighth Edition, by J. LANE NOTTER, A.M., M.D., F.R.S., Professor of Military Hygiene in the Army Medical School. 8vo, with 10 Plates and 103 Engravings, 18s.

A Handbook of Hygiene and Sanitary Science. By GEO. WILSON, M.A., M.D., LL.D., F.R.S.E., D.P.H. Camb., Medical Officer of Health for Mid-Warwickshire. Seventh Edition. Crown 8vo, with Engravings, 12s. 6d.

Elements of Health: an Introduction to the Study of Hygiene. By Louis C. Parkes, M.D., D.P.H. Lond., Lecturer on Public Health at St. George's Hospital. Post 8vo, with

27 Engravings, 3s. od.

The Prevention of Epidemics and the Construction and Management of Isolation Hospitals. By ROGER MCNEILL, M.D. Edin., D.P.H. Camb., Medical Officer of Health for the County of Argyll. 8vo. With several Hospital Plans, 10s. 6d.

Hospitals and Asylums of the World; their Origin, History, Construction, Administration, Management, and Legislation. By HENRY C. BURDETT. In 4 vols. Super Royal 8vo and Portfolio. Complete, 168s. Vols. I. and II.—Asylums: 90s. Vols. III. and IV.-Hospitals, &c., with Portfolio of Plans, 120s.

Mental Diseases:

Clinical Lectures. By T. S. CLOUSTON, M.D., F.R.C.P. Edin., Lecturer on Mental Diseases in the University of Edinburgh. Third Edition. Crown 8vo, with 13 Plates, 14s.

- The Insane and the Law: a Plain Guide for Medical Men, Solicitors, and Others as to the Detention and Treatment, Maintenance, Responsibility, and Capacity either to give evidence or make a will of Persons Mentally Afflicted. With Hints to Medical Witnesses and to Cross-Examining Counsel. By G. PITT-LEWIS, Q.C., R. PERCY SMITH, M.D., F.R.C.P., Resident Physician, Bethlem Hospital, and J. A. HAWKE, B.A., Barrister-at-Law. 8vo, 14s.
- Illustrations of the Influence of the Mind upon the Body in Health and Disease: Designed to elucidate the Action of the Imagination. By D. HACK TUKE, M.D., F.R.C.P., LL.D. Second Edition. 2 vols. crown 8vo, 15s.
- A Dictionary of Psychological Medicine, giving the Definition, Etymology, and Synonyms of the Terms used in Medical Psychology; with the Symptoms, Treatment, and Pathology of Insanity; and The Law of Lunacy in Great Britain and Ireland. Edited by D. Hack Tuke, M.D., LL.D., assisted by nearly 130 Contributors, British, Continental and American. 2 vols., 1,500 pages, royal 8vo, Illustrated. 42s.
- By CHARLES MERCIER, M.B., Lecturer on Neurology and Insanity to the Westminster Hospital Medical School, and to the Medical School for Women. Crown 8vo, 5s.
- The Journal of Mental Science.

 Published Quarterly, by Authority of the Medico-Psychological Association. 8vo, 3s. 6d.
- Mental Affections of Childhood and Youth (Lettsomian Lectures for 1887, &c.). By J. Langdon-Down, M.D., F.R.C.P., Consulting Physician to the London Hospital. 8vo, 6s.

Manual of Midwifery:

Including all that is likely to be required by Students and Practitioners. By ALFRED L. GALABIN, M.A., M.D., F.R.C.P., Obstetric Physician to, and Lecturer on, Midwifery, &c., at Guy's Hospital. Third Edition. Crown 8vo,

with 261 Engravings, 15s.

The Student's Guide to the Practice of Midwifery. By D. LLOYD ROBERTS, M.D., F.R.C.P., Lecturer on Clinical Midwifery and Diseases of Women at the Owens College; Obstetric Physician to the Manchester Royal Infirmary. Fourth Edition. Fcap. 8vo, with Coloured Plates and Engravings.

[Preparing.

Manual of the Diseases peculiar to Women. By James Oliver, M.D., F.R.S. Edin., M.R.C.P. Lond., Physician to the Hospital for Diseases of Women, London. Fcap. 8vo, 3s. 6d.

By the same Author.

Abdominal Tumours and Abdominal Dropsy in Women.

Crown 8vo, 7s. 6d.

Obstetric Aphorisms:

For the Use of Students commencing Midwifery Practice. By JOSEPH G. SWAYNE, M.D. Tenth Edition. Fcap.

8vo, with 20 Engravings, 3s. 6d.

Female Pelvic Organs (The Surgery, Surgical Pathology, and Surgical Anatomy of): in a Series of Plates taken from Nature. With Commentaries, Notes, and Cases. By Henry Savage, M.D., Consulting Physician to the Samaritan Hospital for Women and Children. Fifth Edition. 4to, Uncoloured, 15s.

Lectures on Obstetric Operations: Including the Treatment of Hæmorrhage, and forming a Guide to the Management of Difficult Labour. By ROBERT BARNES, M.D., F.R.C.P., Consulting Obstetric Physician to St. George's Hospital. Fourth Edition. 8vo, with 121 Engravings, 12s. 6d.

By the same Author.

A Clinical History of Medical and Surgical Diseases of Women. Second Edition. 8vo, with

181 Engravings, 28s.

Clinical Lectures on Diseases of Women: Delivered in St. Bartholomew's Hospital, by J. Matthews Duncan, M.D., LL.D., F.R.C.P., F.R.Ss. L. & E., late Obstetric Physician to St. Bartholomew's Hospital. Fourth Edition. Svo, 16s.

Gynæcological Operations:

(Handbook of). By Alban H. G. Doran, F.R.C.S., Surgeon to the Samaritan Hospital. 8vo, with 167 Engravings, 15s.

The Student's Guide to the Diseases of Women. By ALFRED L. GALABIN, M.A., M.D., F.R.C.P., Obstetric Physician to Guy's Hospital. Fifth Edition. Fcap. 8vo, with 142 Engravings, 8s. 6d.

A Practical Treatise on the Diseases of Women. By T. Gail-Lard Thomas, M.D. Sixth Edition, by Paul F. Mundé, M.D., Professor of Gynæcology at the New York Polyclinic and at Dartmouth College. Roy. 8vo, with 347 Engravings, 25s.

Notes on Diseases of Women:
Specially designed to assist the Student in preparing for Examination. By JAMES J. REYNOLDS, L.R.C.P., M.R.C.S. Fourth Edition, Fcap. 8vo., 3s. 6d.

Abdominal Surgery.

By J. Greig Smith, M.A., F.R.S.E., Surgeon to the Bristol Royal Infirmary, and Lecturer on Surgery in the Bristol Medical School. Fifth Edition. 8vo, with Engravings.

[In the press.

The Physiology of Death from Traumatic Fever; A Study in Abdominal Surgery. By JOHN D. MAL-COLM, M.B., C.M., F.R.C.S.E., Surgeon to the Samaritan Free Hospital. 8vo, 3s. 6d.

Notes on Gynæcological Nursing. By John Benjamin Hellier, M.D., M.R.C.S., Lecturer on the Diseases of Women and Children in the Yorkshire College, and Surgeon to the Hospital for Women, &c., Leeds. Crown 8vo, 1s. 6d.

A Manual for Hospital Nurses, and others engaged in Attending on the Sick, with a Glossary. By EDWARD J. DOMVILLE, Surgeon to the Exeter Lyingin Charity. Seventh Edition. Crown 8vo, 2s. 6d.

A Manual of Nursing, Medical and Surgical. By Charles J. Cul-LINGWORTH, M.D., F.R.C.P., Obstetric Physician to St. Thomas's Hospital. Third Edition. Fcap. 8vo, with Engravings, 2s. 6d.

By the same Author.

A Short Manual for Monthly Nurses. Third Edition. Fcap. 8vo, Is. 6d.

Diseases of Children.

For Practitioners and Students. By W. H. DAY, M.D., Physician to the Samaritan Hospital. Second Edition. Crown 8vo, 12s. 6d.

The Diseases of Children (Student's Guide Series). By Jas. F. Goodhart, M.D., F.R.C.P., Physician to Guy's Hospital. Fifth Edition. Fcap. 8vo, 10s. 6d.

A Practical Treatise on Disease in Children. By EUSTACE SMITH, M.D., F.R.C.P., Physician to the King of the Belgians, and to the East London Hospital for Children, &c. Second Edition. 8vo, 22s.

By the same Author.

Clinical Studies of Disease in Children. Second Edition. Post 8vo, 7s. 6d.

Also.

The Wasting Diseases of Infants and Children. Fifth Edition. Post 8vo, 8s. 6d.

A Practical Manual of the Diseases of Children. With a Formulary. By EDWARD ELLIS, M.D. Fifth Edition. Crown 8vo, 10s.

Materia Medica:

A Manual for the use of Students. By ISAMBARD OWEN, M.D., F.R.C.P., Lecturer on Materia Medica, &c., to St. George's Hospital. Second Edition. Crown 8vo, 6s. 6d.

Materia Medica,

Pharmacy, Pharmacology, and Therapeutics. By W. HALE WHITE, M.D., F.R.C.P., Physician to, and Lecturer on Materia Medica and Therapeutics at, Guy's Hospital; Examiner in Materia Medica on the Conjoint Board of the Royal Colleges of Physicians and Surgeons. Fcap. 8vo, 7s. 6d.

Materia Medica

And Therapeutics. By Charles D. F. Phillips, M.D., F.R.S. Edin.

Vegetable Kingdom — Organic Compounds—Animal Kingdom. 8vo, 25s Inorganic Substances. Second Edition 8vo, 21s.

Recent Materia Medica.

Notes on their Origin and Therapeutics. By F. Harwood Lescher, F.C.S., Pereira Medallist. Fourth Edition. 8vo, 2s. 6d

Galenic Pharmacy:

A Practical Handbook to the Processes of the British Pharmacopeia. By R. A. CRIPPS, M.P.S. 8vo, with 76 Engravings, 8s. 6d.

The Galenical Preparations of the British Pharmacopæia: A Handbook for Medical and Pharmaceutical Students. By CHARLES O. HAWTHORNE, M.B., C.M., Lecturer on Materia Medica and Therapeutics, Queen Margaret College, University of Glasgow. 8vo, 4s. 6d.

Practical Pharmacy.

By Barnard S. Proctor, formerly Lecturer on Pharmacy at the College of Medicine, Newcastle-on-Tyne. Third Edition. 8vo, with 44 Wood Engravings and 32 Lithograph Fac-Simile Prescriptions, 14s.

Selecta è Prescriptis:

Containing Terms, Phrases, Contractions and Abbreviations used in Prescriptions, with Explanatory Notes, &c. Also, a Series of Abbreviated Prescriptions with Translations and Key. By J. Pereira, M.D., F.R.S. Eighteenth Edition, by JOSEPH INCE, F.C.S., F.L.S. 24mo, 5s.

A Companion to the British Pharmacopœia. By Peter Squire, Revised by his Sons, P. W. and A. H. Squire. Sixteenth Edition. 8vo, 12s. 6d.

By the same Authors.

The Pharmacopæias of the London Hospitals, arranged in Groups for Easy Reference and Comparison. Sixth Edition. 18mo. 6s. The National Dispensatory:

Containing the Natural History, Chemistry, Pharmacy, Actions and Uses of Medicines, &c. By Alfred Stille, M.D., Ll.D., John M. Maisch, Phar. D., Chas. Caspari, Jun., Ph.G., and Henry C. C. Maisch, Ph.G., Ph.D. Fifth Edition, with 320 Engravings. Imp. 8vo, 36s.

Pocket Formulary

And Synopsis of the British and Foreign Pharmacopæias. By Henry Beasley. Eleventh Edition. 18mo, 6s. 6d.

By the same Author.

Druggist's General Receipt-Book. Tenth Edition. 18mo, 6s. 6d.

Book of Prescriptions:

Containing upwards of 3,000 Prescriptions from the Practice of the most eminent Physicians and Surgeons, English and Foreign. Seventh Edition. 18mo, 6s. 6d.

The Prescriber's Pharmacopæia:

The Medicines arianged in Classes according to their Action, with their Composition and Doses. By NESTOR J. C. TIRARD, M.D., F.R.C.P., Professor of Materia Medica and Therapeutics in King's College, London. Sixth Edition. 32mo, bound in leather, 3s.

Year-Book of Pharmacy:

Containing the Transactions of the British Pharmaceutical Conference. Annually. 8vo, 10s.

Royle's Manual of Materia
Medica and Therapeutics. Sixth
Edition, including additions and alterations in the B.P. 1885. By John
Harley, M.D., Physician to St.
Thomas's Hospital. Crown Svo, with
139 Engravings, 15s.

Manual of Botany. Vol. I.

Anatomy and Morphology. By
J. REYNOLDS GREEN, Sc.D., M.A.,
F.R.S., Professor of Botany to the
Pharmaceutical Society. Crown 8vo,
with about 700 Engravings.

[Nearly ready.

The Student's Guide to Systematic Botany, including the Classification of Plants and Descriptive Botany. By ROBERT BENTLEY, late Emeritus Professor of Botany in King's College and to the Pharmaceutical Society. Fcap. 8vo, with 350 Engravings, 3s. 6d.

Medicinal Plants:

Being descriptions, with original figures, of the Principal Plants employed in Medicine, and an account of their Properties and Uses. By Prof. BENTLEY and Dr. H. TRIMEN, F.R.S. In 4 vols., large 8vo, with 306 Coloured Plates, bound in Half Morocco, Gilt Edges, £11 11s.

Climate and Fevers of India, with a series of Cases (Croonian Lectures, 1882). By Sir Joseph Fayrer, K.C.S.I., M.D. 8vo, with 17 Temperature Charts, 12s.

By the same Author.

- The Natural History and Epidemiology of Cholera: Being the Annual Oration of the Medical Society of London, 1888. 8vo, 3s. 6d.
- A Manual of Family Medicine and Hygiene for India. Published under the Authority of the Government of India. By Sir WILLIAM J. MOORE, K.C.I.E., M.D., late Surgeon-General with the Government of Bombay. Sixth Edition. Post 8vo, with 71 Engravings, 12s.

By the same Author.

A Manual of the Diseases of India: With a Compendium of Diseases generally. Second Edition. Post 8vc, 10s.

Also,

- The Constitutional Requirements for Tropical Climates, &c. Crown 8vo, 4s.
- The Prevention of Disease in Tropical and Sub-Tropical Campaigns. (Parkes Memorial Prize for 1886.) By ANDREW DUNCAN, M.D., B.S. Lond., F.R.C.S., Surgeon-Major, Bengal Army. 8vo, 12s. 6d.

Practical Therapeutics:

A Manual. By EDWARD J. WARING, C.I.E., M.D., F.R.C.P., and DUDLEY W. BUXTON, M.D., B.S. Lond. Fourth Edition. Crown 8vo, 14s.

By the same Author.

Bazaar Medicines of India, And Common Medical Plants: With Full Index of Diseases, indicating their Treat-

ment by these and other Agents procurable throughout India, &c. Fourth Edition Fcap. 8vo, 5s.

Edition, Fcap. 8vo, 5s.

A Commentary on the Diseases of India. By Norman Chevers, C.I.E., M.D., F.R.C.S., Deputy Surgeon-General H.M. Indian Army. 8vo, 24s.

Hooper's Physicians' Vade-Mecum. A Manual of the Principles and Practice of Physic. Tenth Edition. By W. A. Guy, F.R.C.P., F.R.S., and J. Harley, M.D., F.R.C.P. With 118 Engravings. Fcap. 8vo, 12s. 6d. The Principles and Practice of Medicine. (Text-book.) By the late C. Hilton Fagge, M.D., and P. H. Pye-Smith, M.D., F.R.S., F.R.C.P., Physician to, and Lecturer on Medicine in, Guy's Hospital. Third Edition. 2 vols. 8vo, cloth, 40s.; Half Leather, 46s.

Manual of the Practice of Medicine. By FREDERICK TAYLOR, M.D., F.R.C.P., Physician to, and Lecturer on Medicine at, Guy's Hospital. Third Edition. Cr. 8vo, with Engravings, 15s.

The Practice of Medicine (Student's Guide Series). By M. Charteris, M.D., Professor of Therapeutics and Materia Medica in the University of Glasgow. Seventh Edition. Fcap. 8vo, with Engravings on Copper and Wood, Ios.

A Dictionary of Practical Medicine. By various writers. Edited by Jas. Kingston Fowler, M.A., M.D., F.R.C.P., Physician to Middlesex Hospital and the Hospital for Consumption. 8vo, cloth, 21s.; half calf, 25s.

How to Examine the Chest:

A Practical Guide for the use of Students. By SAMUEL WEST, M.D., F.R.C.P., Assistant Physician to St. Bartholomew's Hospital. Second Edition. With Engravings. Fcap. 8vo, 5s.

The Bronchi and Pulmonary
Blood-vessels: their Anatomy and
Nomenclature. By WILLIAM EWART,
M.D., F.R.C.P., Physician to St. George's
Hospital. 4to, with 20 Illustrations, 21s.

An Atlas of the Pathological Anatomy of the Lungs. By the late WILSON FOX, M.D., F.R.S., F.R.C.P., Physician to H.M. the Queen. With 45 Plates (mostly Coloured) and Engravings. 4to, half-bound in Calf, 70s.

By the same Author.

A Treatise on Diseases of the Lungs and Pleura. Edited by SIDNEY COUPLAND, M.D., F.R.C.P., Physician to Middlesex Hospital. Roy. 8vo, with Engravings; also Portrait and Memoir of the Author, 36s.

The Student's Guide to Diseases of the Chest. By VINCENT D. HARRIS, M.D. Lond., F.R.C.P., Physician to the City of London Hospital for Diseases of the Chest, Victoria Park. Fcap. 8vo, with 55 Illustrations (some Coloured), 7s. 6d.

Uric Acid

as a Factor in the Causation of Disease. By Alexander Haig, M.D., F.R.C.P., Physician to the Metropolitan Hospital and the Royal Hospital for Children and Women. Second Edition. With Illustrations, 8vo, 10s. 6d.

Medical Diagnosis (Student's Guide Series). By SAMUEL FENWICK, M.D., F.R.C.P., Physician to the London Hospital. Seventh Edition. Fcap. 8vo, with 117 Engravings, 7s.

By the same Author.

Outlines of Medical Treatment.

Fourth Edition. Crown Svo, with 35
Engravings, 10s.

Also.

Clinical Lectures on Some Obscure Diseases of the Abdomen.
Delivered at the London Hospital. 8vo, with Engravings, 7s. 6d.

Also.

The Saliva as a Test for Functional Diseases of the Liver. Crown 8vo, 2s.

The Microscope in Medicine.

By Lionel S. Beale, M.B., F.R.S.,
Physician to King's College Hospital.
Fourth Edition. 8vo, with 86 Plates, 21s.

By the same Author.

The Liver.

With 24 Plates (85 Figures). 8vo, 5s.

Also.

On Slight Ailments:

And on Treating Disease. Third Edition. 8vo, 5s.

The Physiology of the Carbohydrates; their Application as Food and Relation to Diabetes. By F. W. Pavy, M.D., LL.D., F.R.S., F.R.C.P., Consulting Physician to Guy's Hospital. Royal 8vo, with Plates and Engravings, 10s. 6d.

Medical Lectures and Essays.

By Sir G. Johnson, M.D., F.R.C.P.,
F.R.S., Consulting Physician to King's
College Hospital. 8vo, with 46 Engravings, 25s.

By the same Author.

An Essay on Asphyxia (Apnœa).

Bronchial Affections:

Pneumonia and Fibroid Pneumonia (their Pathological Histology). An Original Investigation. By A. G. Auld, M.D., Assistant Physician to the Glasgow Royal Infirmary. 8vo, with Illustrations, 7s. 6d.

Bronchial Asthma:

Its Pathology and Treatment. By J. B. BERKART, M.D., late Physician to the City of London Hospital for Diseases of the Chest. Second Edition, with 7 Plates (35 Figures). 8vo, 10s. 6d.

Treatment of Some of the Forms of Valvular Disease of the Heart. By A. E. SANSOM, M.D., F.R.C.P., Physician to the London Hospital. Second Edition. Fcap. 8vo, with 26 Engravings, 4s. 6d.

The Schott Methods of the Treatment of Chronic Diseases of the Heart, with an account of the Nauheim Baths and of the Therapeutic Exercises. By W. BEZLY THORNE, M.D., M.R.C.P. 8vo, with Illustrations, 5s.

Guy's Hospital Reports.

By the Medical and Surgical Staff. Vol.

XXXV. Third Series. 8vo, 10s. 6d.

St. Thomas's Hospital Reports.

By the Medical and Surgical Staff. Vol.

XXII. New Series. 8vo, 8s. 6d.

Westminster Hospital Reports.

By the Medical and Surgical Staff. Vol.

IX. 8vo, 6s.

The Climate of Rome

and the Roman Malaria. By Professor Tommasi-Crudeli. Translated by Charles Cramond Dick. Crown 8vo, 5s.

Vaccinia and Variola:

A Study of their Life History. By JOHN B. BUIST, M.D., F.R.S.E., Teacher of Vaccination for the Local Government Board. Crown 8vo, with 24 Coloured Plates, 7s. 6d.

Medical Ophthalmoscopy:

A Manual and Atlas. By W. R. Gowers, M.D., F.R.C.P., F.R.S., Physician to the National Hospital for the Paralysed and Epileptic. Third Edition. Edited with the assistance of Marcus Gunn, M.B., F.R.C.S., Surgeon to the Royal London Ophthalmic Hospital. With Coloured Plates and Woodcuts. 8vo, 16s.

By the same Author.

A Manual of Diseases of the Nervous System.

Vol. I. Diseases of the Nerves and Spinal Cord. Second Edition. Roy. 8vo, with

179 Engravings, 15s.

Vol. II. Diseases of the Brain and Cranial Nerves: General and Functional Diseases of the Nervous System. Second Edition. Roy. 8vo, with 182 Engravings, 20s.

Also.

Diagnosis of Diseases of the Brain. Second Edition. 8vo, with Engravings, 7s. 6d.

Also.

Syphilis and the Nervous System. Being a Revised Reprint of the Lettsomian Lectures for 1890. Delivered before the Medical Society of London. 8vo, 4s.

The Nervous System,

Diseases of. By J. A. Ormerod, M.D., F.R.C.P., Physician to the National Hospital for the Paralysed and Epileptic. With 66 Illustrations. Fcap. 8vo, 8s. 6d.

Handbook of the Diseases of the Nervous System. By James Ross, M.D., F.R.C.P., Professor of Medicine in the Victoria University, and Physician to the Royal Infirmary, Manchester. Roy. 8vo, with 184 Engravings, 18s.

Also.

Aphasia:

Being a Contribution to the Subject of the Dissolution of Speech from Cerebral Disease. 8vo, with Engravings, 4s. 6d.

Diseases of the Nervous System.

Lectures delivered at Guy's Hospital. By
SAMUEL WILKS, M.D., F.R.S. Second
Edition. 8vo, 18s.

Stammering:

Its Causes, Treatment, and Cure. By A. G. BERNARD, M.R.C.S., L.R.C.P. Crown 8vo, 2s.

Secondary Degenerations of the Spinal Cord (Gulstonian Lectures, 1889). By HOWARD H. TOOTH, M.D., F.R.C.P., Assistant Physician to the National Hospital for the Paralysed and Epileptic. With Plates and Engravings. 8vo, 3s. 6d.

Diseases of the Nervous System.

Clinical Lectures. By Thomas Buzzard,
M.D., F.R.C.P., Physician to the National
Hospital for the Paralysed and Epileptic.
With Engravings, 8vo. 15s.

By the same Author.

Some Forms of Paralysis from Peripheral Neuritis: of Gouty, Alcoholic, Diphtheritic, and other origin. Crown 8vo, 5s.

Also.

On the Simulation of Hysteria by Organic Disease of the Nervous System. Crown 8vo, 4s. 6d.

Gout in its Clinical Aspects.

By J. Mortimer Granville, M.D.

Crown 8vo, 6s.

Diseases of the Liver:

With and without Jaundice By GEORGE HARLEY, M.D., F.R.C.P., F.R.S. 8vo, with 2 Plates and 36 Engravings, 21s.

Rheumatic Diseases,

(Differentiation in). By HUGH LANE, Surgeon to the Royal Mineral Water Hospital, Bath, and Hon. Medical Officer to the Royal United Hospital, Bath. Second Edition, much Enlarged, with 8 Plates. Crown 8vo, 3s. 6d.

Diseases of the Abdomen,

Comprising those of the Stomach and other parts of the Alimentary Canal, Œsophagus, Cæcum, Intestines, and Peritoneum. By S. O. HABERSHON, M.D., F.R.C.P. Fourth Edition. 8vo, with 5 Plates, 21s.

On the Relief of Excessive and
Dangerous Tympanites by
Puncture of the Abdomen.
By John W. Ogle, M.A., M.D.,
F.R.C.P., Consulting Physician to St.
George's Hospital. 8vo, 5s. 6d.

Headaches:

Their Nature, Causes, and Treatment. By W. H. DAY, M.D., Physician to the Samaritan Hospital. Fourth Edition. Crown 8vo, with Engravings, 7s. 6d.

Health Resorts at Home and Abroad. By M. CHARTERIS, M.D., Professor of Therapeutics and Materia Medica in Glasgow University. Second Edition. Crown 8vo, with Map, 5s. 6d.

The Mineral Waters of France
And its Wintering Stations (Medical
Guide to). With a Special Map. By
A. VINTRAS, M.D., Physician to the
French Embassy, and to the French
Hospital, London. Second Edition.
Crown 8vo, 8s.

Canary Islands

Health Resorts, in their Climatological and Medical Aspects. By J. CLEASBY TAYLOR, M.D., M.R.C.S., Las Palmas. 8vo, with Maps, 3s. 6d.

Homburg-Spa.

An Introduction to its Waters and their use. By Dr. ARNOLD SCHETELIG. Crown 8vo, with Synoptical Table, 2s. 6d.

Illustrated Ambulance Lectures:
To which is added a Nursing Lecture.
By John M. H. Martin, M.D., F.R.C.S.,
Honorary Surgeon to the Blackburn Infirmary. Fourth Edition. Crown 8vo,
with 60 Engravings, 2s. [Nearly ready.

Surgery: its Theory and Practice. By WILLIAM J. WALSHAM, F.R.C.S., Senior Assistant Surgeon to, Lecturer on and Anatomy at, St. Bartholomew's Hospital. Fifth Edition. Crown 8vo, with 380 Engravings, 12s. 6d.

Surgical Emergencies:

Together with the Emergencies attendant on Parturition and the Treatment of Poisoning. ByW.PAULSWAIN, F.R.C.S., Surgeon to the South Devon and East Cornwall Hospital. Fourth Edition. Crown 8vo, with 120 Engravings, 5s.

Operations on the Brain (A Guide to). By ALEC FRASER, Professor of Anatomy, Royal College of Surgeons in Ireland. Illustrated by 42 life-size Plates in Autotype, and 2 Woodcuts in the text. Folio, 63s.

A Course of Operative Surgery.

By Christopher Heath, Surgeon to
University College Hospital. Second
Edition. With 20 coloured Plates (180
figures) from Nature, by M. Léveillé,
and several Woodcuts. Large 8vo, 30s.

By the same Author.

The Student's Guide to Surgical Diagnosis. Second Edition. Fcap. 8vo, 6s. 6d.

Also.

Manual of Minor Surgery and Bandaging. For the use of House-Surgeons, Dressers, and Junior Practitioners. Tenth Edition. Fcap. 8vo, with 158 Engravings, 6s.

Also.

Injuries and Diseases of the
Jaws. Fourth Edition. By HENRY
PERCY DEAN, M.S., F.R.C.S., Assistant
Surgeon to the London Hospital. 8vo,
with 187 Wood Engravings, 14s.

Also.

Lectures on Certain Diseases of the Jaws. Delivered at the R.C.S., Eng., 1887. 8vo, with 64 Engravings, 2s. 6d.

Also.

Clinical Lectures on Surgical Subjects. Delivered in University College Hospital. Fcap. 8vo, with 23 Engravings, 6s.

Surgery.

By C. W. Mansell Moullin, M.A., M.D., Oxon., F.R.C.S., Surgeon and Lecturer on Physiology to the London Hospital. Large 8vo, with 497 Engravings, 34s.

The Practice of Surgery:

A Manual. By THOMAS BRYANT, Consulting Surgeon to Guy's Hospital. Fourth Edition. 2 vols. crown 8vo, with 750 Engravings (many being coloured), and including 6 chromo plates, 32s.

By the same Author.

On Tension: Inflammation of Bone, and Head Injuries. Hunterian Lectures, 1888. 8vo, 6s.

The Surgeon's Vade-Mecum:

A Manual of Modern Surgery. By R.
DRUITT, F.R.C.S. Twelfth Edition.
By STANLEY BOYD, M.B., F.R.C.S.
Assistant Surgeon and Pathologist to
Charing Cross Hospital. Crown 8vo,
with 373 Engravings, 16s.

By CHARLES MACNAMARA, F.R.C.S., Surgeon to, and Lecturer on Surgery at, the Westminster Hospital. 8vo, with Plates and Engravings, 12s. The Operations of Surgery:

Intended for Use on the Dead and Living Subject alike. By W. H. A. JACOBSON, M.A., M.B., M.Ch. Oxon., F.R.C.S., Assistant Surgeon to, and Lecturer on Anatomy at, Guy's Hospital. Third Edition. 8vo, with many Illustrations.

[In the press.

On Anchylosis.

By BERNARD E. BRODHURST, F.R.C.S., Surgeon to the Royal Orthopædic Hospital. Fourth Edition. 8vo, with Engravings, 5s.

By the same Author.

Curvatures and Disease of the Spine. Fourth Edition. 8vo, with Engravings, 7s. 6d.

Also.

Talipes Equino-Varus, or Clubfoot. 8vo, with Engravings, 3s. 6d.

Surgical Pathology and Morbid Anatomy. By Anthony A. Bowlby, F.R.C.S., Assistant Surgeon to St. Bartholomew's Hospital. Third Edition. Crown 8vo, with 183 Engravings, 10s. 6d.

By the same Author.

Injuries and Diseases of Nerves and their Surgical Treatment. 8vo, with 20 Plates, 14s.

Illustrations of Clinical Surgery.

By Jonathan Hutchinson, F.R.S.,
Senior Surgeon to the London Hospital.
In fasciculi. 6s. 6d. each. Fasc. I. to
X. bound, with Appendix and Index,
£3 ios. Fasc. XI. to XXIII. bound,
with Index, £4 ios.

Clubfoot:

Its Causes, Pathology, and Treatment. By WM. Adams, F.R.C.S., Consulting Surgeon to the Great Northern and other Hospitals. Second Edition. Svo, with 106 Engravings and 6 Lithographic Plates, 15s.

By the same Author.

Lateral and other Forms of Curvature of the Spine: Their Pathology and Treatment. Second Edition. 8vo, with 5 Lithographic Plates and 72 Wood Engravings, 10s. 6d.

Also.

Contraction of the Fingers:

(Dupuytren's and Congenital Contractions), their Treatment by Subcutaneous Divisions of the Fascia, and Immediate Extension. Also on Hammer Toe; its Curability by Subcutaneous Division. And on The Obliteration of Depressed Cicatrices by a Subcutaneous Operation. 8vo, with 8 Plates and 31 Engravings, 6s. 6d.

Short Manual of Orthopædy.

By Heather Bigg, F.R.C.S. Ed

Part I. Deformities and Deficiencies of the Head and Neck. 8vo. 2s. 6d.

Face and Foot Deformities.

By Frederick Churchill, C.M. 8vo, with Plates and Illustrations, 10s. 6d.

The Human Foot:

Its Form and Structure, Functions and Clothing. By THOMAS S. ELLIS, Consulting Surgeon to the Gloucester Infirmary. With 7 Plates and Engravings (50 Figures). 8vo, 7s. 6d.

Royal London Ophthalmic Hospital Reports. By the Medical and Surgical Staff. Vol. XIII., Part 4. 8vo, 5s.

Ophthalmological Society

of the United Kingdom. Transactions, Vol. XIV. 8vo, 12s. 6d.

The Diseases of the Eye

(Student's Guide Series). By EDWARD NETTLESHIP, F.R.C.S., Ophthalmic Surgeon to St. Thomas's Hospital. Fifth Edition. Fcap. 8vo, with 164 Engravings and a Coloured Plate illustrating Colour-Blindness, 7s. 6d.

Diseases and Refraction of the Eye. By N. C. Macnamara, F.R.C.S., Surgeon to Westminster Hospital, and Gustavus Hartridge, F.R.C.S., Surgeon to the Royal Westminster Ophthalmic Hospital. Fifth Edition. Crown 8vo, with Plate, 156 Engravings, also Testtypes, 10s. 6d.

Diseases of the Eye: a Practical
Handbook for General Practitioners and Students. By Cecil
Edward Shaw, M.D., M.Ch., Ophthalmic Surgeon to the Ulster Hospital for Children and Women, Belfast. With a Test-Card for Colour - Blindness. Crown 8vo, 3s. 6d.

On Diseases and Injuries of the Eye: A Course of Systematic and Clinical Lectures to Students and Medical Practitioners. By J. R. Wolfe, M.D., F.R.C.S.E., Lecturer on Ophthalmic Medicine and Surgery in Anderson's College, Glasgow. With 10 Coloured Plates and 157 Wood Engravings. 8vo, £1 is.

Normal and Pathological Histology of the Human Eye and Eyelids. By C. Fred. Pollock, M.D., F.R.C.S. and F.R.S.E., Surgeon for Diseases of the Eye to Anderson's College Dispensary, Glasgow. Crown 8vo, with 100 Plates (230 drawings), 15s.

Refraction of the Eye:

A Manual for Students. By Gustavus Hartridge, F.R.C.S., Surgeon to the Royal Westminster Ophthalmic Hospital. Seventh Edition. Crown 8vo, with 98 Illustrations, also Test-types, &c., 6s.

By the same Author.

The Ophthalmoscope. A Manual for Students. Second Edition. Crown 8vo, with 67 Illustrations and 4 Plates. 4s. 6d.

Methods of Operating for Cataract and Secondary Impairments of Vision, with the results of 500 cases. By G. H. Fink, Surgeon-Captain in H.M. Indian Medical Service. Crown 8vo, with 15 Engravings, 5s.

Atlas of Ophthalmoscopy.

Composed of 12 Chromo-lithographic Plates (59 Figures drawn from nature) and Explanatory Text. By RICHARD LIEBREICH, M.R.C.S. Translated by H. ROSBOROUGH SWANZY, M.B. Third Edition, 4to, 40s.

Glaucoma:

Its Pathology and Treatment. By PRIESTLEY SMITH, Ophthalmic Surgeon to, and Clinical Lecturer on Ophthalmology at, the Queen's Hospital, Birmingham. 8vo, with 64 Engravings and 12 Zinco-photographs, 7s. 6d.

Eyestrain

(commonly called Asthenopia). By ERNEST CLARKE, M.D., B.S. Lond., Surgeon to the Central London Ophthalmic Hospital, Surgeon and Ophthalmic Surgeon to the Miller Hospital. 8vo, with 22 Illustrations, 5s.

Diseases of the Eye:

A Handbook of Ophthalmic Practice for Students and Practitioners. By G. E. DE SCHWEINITZ, M.D., Professor of Diseases of the Eye in the Philadelphia Polyclinic. With 216 Illustrations, and 2 Chromo-Lithographic Plates. 8vo, 18s.

Diseases and Injuries of the Ear. By Sir WILLIAM B. DALBY, F.R.C.S., M.B., Consulting Aural Surgeon to St. George's Hospital. Fourth Edition. Crown 8vo, with 8 Coloured Plates and 38 Wood Engravings. 10s. 6d. By the same Author.

Short Contributions to Aural Surgery, between 1875 and 1889. Second Edition. 8vo, with Engravings, 3s. 6d.

Diseases of the Ear,

Including the Anatomy and Physiology of the Organ, together with the Treatment of the Affections of the Nose and Pharynx which conduce to Aural Disease (a Treatise). By T. MARK HOVELL, F.R.C.S.E., M.R.C.S., Aural Surgeon to the London Hospital, and Lecturer on Diseases of the Throat in the College, &c. 8vo, with 122 Engravings, 18s.

- Hintson Ophthalmic Out-Patient
 Practice. By CHARLES HIGGENS,
 Ophthalmic Surgeon to Guy's Hospital.
 Third Edition. Fcap. 8vo, 3s.
- A System of Dental Surgery.

 By Sir John Tomes, F.R.S., and C. S.
 Tomes, M.A., F.R.S. Third Edition.

 Crown 8vo, with 292 Engravings, 15s.
- Dental Anatomy, Human and Comparative: A Manual. By CHARLES S. TOMES, M.A., F.R.S. Fourth Edition. Crown 8vo, with 235 Engravings, 12s. 6d.
- A Manual of Nitrous Oxide Anæsthesia, for the use of Students and General Practitioners. By J. Frederick W. Silk, M.D. Lond., M.R.C.S., Anæsthetist to the Royal Free Hospital, Dental School of Guy's Hospital, and National Epileptic Hospital. 8vo, with 26 Engravings, 5s.
- A Practical Treatise on Mechanical Dentistry. By Joseph Richardson, M.D., D.D.S. Sixth Edition revised and Edited by George W. Warren, D.D.S. Roy. 8vo, with 600 Engravings, 21s.

Notes on Dental Practice.

By Henry C. Quinby, L.D.S.I., President-Elect of the British Dental Association. Second Edition. 8vo, with 92 Illustrations, 8s.

Elements of Dental Materia
Medica and Therapeutics, with
Pharmacopœia. By James Stocken,
L.D.S.R.C.S., Pereira Prizeman for
Materia Medica, and Thomas Gaddes,
L.D.S. Eng. and Edin. Third Edition.
Fcap. 8vo, 7s. 6d.

Papers on Dermatology.

By E. D. MAPOTHER, M.D., Ex-Pres. R.C.S.I. 8vo, 3s. 6d.

Atlas of Skin Diseases.

By TILBURY FOX, M.D., F.R.C.P. With 72 Coloured Plates. Royal 4to, half morocco, £6 6s.

Diseases of the Skin:

A Practical Treatise for the Use of Students and Practitioners. By J. N. HYDE, A.M., M.D., Professor of Skin and Venereal Diseases, Rush Medical College, Chicago. Second Edition. 8vo, with 2 Coloured Plates and 96 Engravings, 20s.

Leprosy in British Guiana.

By John D. Hillis, F.R.C.S., M.R.I.A., Medical Superintendent of the Leper Asylum, British Guiana. Imp. 8vo, with 22 Lithographic Coloured Plates and Wood Engravings, £1 11s. 6d.

Diseases of the Skin

(Introduction to the Study of). By P. H. Pye-Smith, M. D., F. R. S., F.R.C.P., Physician to, and Lecturer on Medicine in, Guy's Hospital. Crown 8vo, with 26 Engravings. 7s. 6d.

Sarcoma and Carcinoma:

Their Pathology, Diagnosis, and Treatment. By Henry T. Butlin, F.R.C.S., Assistant Surgeon to St. Bartholomew's Hospital. 8vo, with 4 Plates, 8s.

By the same Author.

Malignant Disease of the Larynx (Sarcoma and Carcinoma). 8vo, with 5 Engravings, 5s.

Also.

Operative Surgery of Malignant Disease. 8vo, 14s.

On Cancer:

Its Allies, and other Tumours; their Medical and Surgical Treatment. By F. A. PURCELL, M.D., M.C., Surgeon to the Cancer Hospital, Brompton. 8vo, with 21 Engravings, 10s. 6d.

Cancers and the Cancer Process: a Treatise, Practical and Theoretic. By HERBERT L. SNOW, M.D., Surgeon to the Cancer Hospital, Brompton. 8vo, with 15 Lithographic Plates. 15s.

By the same Author.

The Re-appearance (Recurrence) of Cancer after apparent Extirpation. 8vo, 5s. 6d.

Also.

The Palliative Treatment of Incurable Cancer. Crown 8vo, 2s. 6d.

Cancerous Affections of the Skin.

(Epithelioma and Rodent Ulcer.) By
GEORGE THIN, M.D. Post 8vo, with
8 Engravings, 5s.

By the same Author.

Pathology and Treatment of Ringworm. 8vo, with 21 Engravings, 5s.

Diagnosis and Treatment of Syphilis. By Tom Robinson, M.D., Physician to St. John's Hospital for Diseases of the Skin. Crown 8vo, 3s. 6d.

By the same Author.

Eczema: its Etiology, Pathology, and Treatment. Crown 8vo, 3s. 6d.

Also.

Illustrations of Diseases of the Skin and Syphilis, with Remarks. Fasc. I with 3 Plates. Imp. 4to, 5s. BY SIR HENRY THOMPSON, F.R.C.S.

Diseases of the Urinary Organs.

Clinical Lectures. Eighth Edition.

8vo, with 121 Engravings, 10s. 6d.

Diseases of the Prostate:

Their Pathology and Treatment. Sixth
Edition. 8vo, with 39 Engravings, 6s.

Surgery of the Urinary Organs.

Some Important Points connected therewith. Lectures delivered in the R.C.S.

8vo, with 44 Engravings. Student's Edition, 2s. 6d.

Practical Lithotomy and Lithotrity; or, An Inquiryinto the Best Modes of Removing Stone from the Bladder. Third Edition. 8vo, with 87 Engravings,

The Preventive Treatment of Calculous Disease, and the Use of Solvent Remedies. Third Edition. Crown 8vo, 2s. 6d.

Tumours of the Bladder:

Their Nature, Symptoms, and Surgical Treatment. 8vo, with numerous Illustrations, 5s.

Stricture of the Urethra, and Urinary Fistulæ: their Pathology and Treatment. Fourth Edition. 8vo, with 74 Engravings, 6s.

The Suprapubic Operation of Opening the Bladder for the Stone and for Tumours. 8vo, with 14 Engravings, 3s. 6d.

Introduction to the Catalogue;
being Notes of a Thousand Cases of
Calculi of the Bladder removed by the
Author and now in the Hunterian
Museum of the Royal College of Surgeons.
8vo, 2s. 6d.

Electric Illumination of the Bladder and Urethra, as a Means of Diagnosis of Obscure Vesico-Urethral Diseases. By E. Hurry Fenwick, F.R.C.S., Surgeon to London Hospital and St. Peter's Hospital for Stone. Second Edition. 8vo, with 54 Engravings, 6s. 6d.

By the same Author.

The Cardinal Symptoms of Urinary Diseases: their Diagnostic Significance and Treatment. 8vo, with 36 Illustrations. 8s. 6d.

Atlas of Electric Cystoscopy.

By Dr. EMIL BURCKHARDT, late of the Surgical Clinique of the University of Bâle, and E. HURRY FENWICK, F.R.C.S.,

Surgeon to the London Hospital and St. Peter's Hospital for Stone. Royal 8vo, with 34 Coloured Plates, embracing 83 Figures. 21s.

Lectures on the Surgical Disorders of the Urinary Organs. By REGINALD HARRISON, F.R.C.S., Surgeon to St. Peter's Hospital. Fourth Edition. 8vo, with 156 Engravings, 16s.

Clinical Chemistry of Urine (Outlines of the). By C. A. Mac-Munn, M.A., M.D. 8vo, with 64 Engravings and Plate of Spectra, 9s.

Urinary and Renal Derangements and Calculous Disorders. By LIONEL S. BEALE, F.R.C.P., F.R.S., Physician to King's College Hospital. 8vo, 5s.

Male Organs of Generation (Diseases of). By W. H. A. JACOBSON, M.Ch. Oxon., F.R.C.S., Assistant Surgeon to Guy's Hospital. 8vo, with 88 Engravings. 22s.

The Surgical Diseases of the Genito - Urinary Organs, including Syphilis. By E. L. Keyes, M.D., Professor in Bellevue Hospital Medical College, New York (a revision of Van Buren and Keyes' Text-book). Roy. 8vo, with 114 Engravings, 21s.

Diseases of the Rectum and Anus. By Alfred Cooper, F.R.C.S., Senior Surgeon to the St. Mark's Hospital for Fistula; and F. Swinford Edwards, F.R.C.S., Senior Assistant Surgeon to St. Mark's Hospital. Second Edition, with Illustrations. 8vo, 12s.

Diseases of the Rectum and Anus. By Harrison Cripps, F.R.C.S., Assistant Surgeon to St. Bartholomew's Hospital, &c. Second Edition. 8vo, with 13 Lithographic Plates and numerous Wood Engravings, 12s. 6d.

By the same Author.

Cancer of the Rectum.

Especially considered with regard to its Surgical Treatment. Jacksonian Prize Essay. 8vo, with 13 Plates and several Wood Engravings, 6s.

The Diagnosis and Treatment of Diseases of the Rectum. By WILLIAM ALLINGHAM, F.R.C.S., Surgeon to St. Mark's Hospital for Fistula. Fifth Edition. By HERBERT WM. ALLINGHAM, F.R.C.S., Surgeon to the Great Northern Central Hospital, Demonstrator of Anatomy at St. George's Hospital. 8vo, with 53 Engravings. 10s. 6d.

Syphilis.

By Alfred Cooper, F.R.C.S., Senior Surgeon to St. Mark's Hospital for Fistula. Second Edition. Edited by Edward Cotterell, F.R.C.S., Surgeon (out-patients) to the London Lock Hospital. 8vo, with 24 Full-page Plates (12 coloured), 18s.

A Medical Vocabulary:

An Explanation of all Terms and Phrases used in the various Departments of Medical Science and Practice, their Derivation, Meaning, Application, and Pronunciation. By R. G. MAYNE, M.D., LL.D. Sixth Edition by W. W. WAGSTAFFE, B.A., F.R.C.S. Crown 8vo, 10s. 6d.

A Short Dictionary of Medical Terms. Being an Abridgment of Mayne's Vocabulary. 64mo, 2s. 6d.

Dunglison's Dictionary of Medical Science: Containing a full Explanation of its various Subjects and Terms, with their Pronunciation, Accentuation, and Derivation. Twenty-first Edition. By RICHARD J. DUNGLISON, A.M., M.D. Royal 8vo, 30s.

Terminologia Medica Polyglotta: a Concise International Dictionary of Medical Terms (French, Latin, English, German, Italian, Spanish, and Russian). By Theodore Maxwell, M.D., B.Sc., F.R.C.S. Edin. Royal 8vo, 16s.

A German-English Dictionary of Medical Terms. By FREDERICK TREVES, F.R.C.S., Surgeon to the London Hospital; and Hugo Lang, B.A. Crown 8vo, half-Persian calf, 12s.

Chemistry,

Inorganic and Organic. With Experiments. By Charles L. Bloxam. Eighth Edition, by John Millar Thomson, Professor of Chemistry in King's College, London, and Arthur G. Bloxam, Head of the Chemistry Department, The Goldsmiths' Institute, New Cross. 8vo, with nearly 300 Illustrations.

By the same Author.

Laboratory Teaching;

Or, Progressive Exercises in Practical Chemistry. Sixth Edition. By ARTHUR G. BLOXAM. Crown 8vo, with 80 Engravings, 6s. 6d.

Ouantitative Analysis.

By Frank Clowes, D.Sc. Lond., Professor of Chemistry in the University College, Nottingham, and J. Bernard Coleman, Assoc. R. C. Sci. Dublin; Senior Demonstrator of Chemistry in the University College, Nottingham. Second Edition. Post 8vo, with 94 Engravings, 8s. 6d.

By the same Authors.

Elementary Qualitative Analysis. With 40 Engravings. Post 8vo, 2s. 6d.

Practical Chemistry

And Qualitative Analysis. By Frank CLOWES, D.Sc. Lond., Professor of Chemistry in the University College, Nottingham. Sixth Edition. Post 8vo, with 84 Engravings and Frontispiece, 8s. 6d.

Watts' Manual of Chemistry,

Theoretical and Practical. By WILLIAM A. TILDEN, D.Sc., F.R.S., Professor of Chemistry in the Normal School of Science, South Kensington.

PHYSICAL AND INORGANIC CHE-MISTRY. Second Edition. Crown 8vo, with Coloured Plate of Spectra, and 122 Wood Engravings, 8s. 6d.

CHEMISTRY OF CARBON COMPOUNDS; or, ORGANIC CHEMISTRY. Second Edition. Crown 8vo, with Engravings, 10s.

Qualitative Analysis.

By R. Fresenius. Translated by Charles E. Groves, F.R.S. Tenth Edition. 8vo, with Coloured Plate of Spectra and 46 Engravings, 15s.

By the same Author.

Quantitative Analysis.

Seventh Edition.

Vol. I., Translated by A. VACHER. 8vo, with 106 Engravings, 15s. Vol. II., Parts 1 to 3, Translated by C. E. GROVES, F.R.S. 8vo, with

Engravings, 2s. 6d. each.

Practical Chemistry,

Including Analysis. By John E. Bow-MAN and CHARLES L. BLOXAM. Fcap. 8vo. Eighth Edition, with 90 Engravings, 5s. 6d.

Inorganic Chemistry.

By EDWARD FRANKLAND, Ph.D., D.C.L., LL.D., F.R.S., Professor of Chemistry in the Normal School of Science, and FRANCIS R. JAPP, M.A., Ph.D. F.I.C., F.R.S., Professor of Chemistry in the University of Aberdeen. 8vo, with numerous Illustrations on Stone and Wood, 24s.

Inorganic Chemistry

(A System of). By WILLIAM RAMSEY, Ph.D., F.R.S., Professor of Chemistry in University College, London. 8vo, with Engravings, 15s.

By the same Author.

Elementary Systematic Chemistry for the Use of Schools and Colleges. With Engravings. Crown 8vo, 4s. 6d.; Interleaved, 5s. 6d.

Valentin's Qualitative Chemical Analysis. Eighth Edition. By W.

R. Hodgkinson, Ph.D., F.R.S.E., Professor of Chemistry and Physics in the Royal Military Academy, and Artillery College, Woolwich. 8vo, with Engravings and Map of Spectra, 8s. 6d.

Analytical Chemistry.

Notes for Students in Medicine. By ALBERT J. BERNAYS, Ph.D., F.C.S., F.I.C., late Professor of Chemistry, &c., at St. Thomas's Hospital Medical School. Third Edition. Crown 8vo, 4s. 6d.

Volumetric Analysis:

(A Systematic Handbook of); or the Quantitative Estimation of Chemical Substances by Measure, applied to Liquids, Solids, and Gases. By Francis Sutton, F.C.S., F.I.C., Public Analyst for the County of Norfolk. Sixth Edition. 8vo, with 102 Engravings, 17s. 6d.

Commercial Organic Analysis:

A Treatise on the Properties, Modes of Assaying, Proximate Analytical Examination, &c., of the various Organic Chemicals and Products employed in the Arts, Manufactures, Medicine, &c. By Alfred H. Allen, F.I.C., F.C.S., Public Analyst for the West Riding of Yorkshire, the Northern Division of Derbyshire, &c.

Vol. I.—Alcohols, Neutral Alcoholic Derivatives, Sugars, Starch and its Isomers, Vegetable Acids, &c. With Illustrations. Third Edition. 8vo. [Preparing.

Vol. II.—Fixed Oils and Fats, Hydrocarbons, Phenols, &c. With Illustrations. Third Edition. 8vo.

[Preparing.

Vol. III.—Part I. Aromatic Acids, Tannins, Dyes, and Colouring Matters. Second Edition, 8vo, 14s.

Part II. Amines and Ammonium Bases, Hydrazines, Bases from Tar, Vegetable Alkaloids. Second Edition. 8vo, 18s.

Cooley's Cyclopædia

of Practical Receipts, and Collateral Information in the Arts, Manufactures, Professions, and Trades: Including Medicine, Pharmacy, Hygiene and Domestic Economy. Seventh Edition, by W. NORTH, M.A. Camb., F.C.S. 2 Vols., Roy. 8vo, with 371 Engravings, 42s.

Chemical Technology:

A Manual. By RUDOLF VON WAGNER. Translated and Edited by WILLIAM CROOKES, F.R.S., from the Thirteenth Enlarged German Edition as remodelled by Dr. FERDINAND FISCHER. 8vo, with 596 Engravings, 32s.

Chemical Technology;

Or, Chemistry in its Applications to Arts and Manufactures. Edited by CHARLES E. GROVES, F.R.S., and WILLIAM THORP, B.Sc.

Vol. I .- FUEL AND ITS APPLICA-F.R.S., and F. J. ROWAN, C.E. Royal 8vo, with 606 Engravings, 30s. Vol. II.—LIGHTING BY CANDLES AND OIL. By W. Y. DENT, J. MCARTHUR, L. FIELD and F. A. FIELD, BOVERTON REDWOOD, and D. A. Louis. Royal 8vo, with 358 Engravings and Map, 20s.

Vol. III.—GAS AND ELECTRICITY. In the press.

Technological Handbooks.

EDITED BY JOHN GARDNER, F.I.C., F.C.S., and JAMES CAMERON, F.I.C. BREWING, DISTILLING, AND WINE

MANUFACTURE. Crown 8vo, with Engravings, 6s. 6d.

BLEACHING, DYEING, AND CALICO PRINTING. With Formulæ. Crown

8vo, with Engravings, 5s.
OILS, RESINS, AND VARNISHES. Crown 8vo, with Engravings. 7s. 6d. SOAPS AND CANDLES. Crown 8vo, with 54 Engravings, 7s.

- The Microscope and its Revelations. By the late WILLIAM B. CAR-PENTER, C.B., M.D., LL.D., F.R.S. Seventh Edition, by the Rev. W. H. DALLINGER, LL.D., F.R.S. With 21 Plates and 800 Wood Engravings. 8vo, 26s. Half Calf, 30s.
- The Quarterly Journal of Microscopical Science. Edited by E. RAY LANKESTER, M.A., LL.D., F.R.S.; with the co-operation of ADAM SEDGWICK, M.A., F.R.S., and W. F. R. WELDON, M.A., F.R.S. Each Number, 10s.

Methods and Formulæ

Used in the Preparation of Animal and Vegetable Tissues for Microscopical Examination, including the Staining of Bacteria. By Peter Wyatt Squire, F.L.S. Crown 8vo, 3s. 6d.

The Microtomist's Vade-Mecum:

A Handbook of the Methods of Microscopic Anatomy. By ARTHUR BOLLES LEE, Assistant in the Russian Laboratory of Zoology at Villefranche-sur-mer (Nice). Third Edition. 8vo, 14s.

Photo-Micrography

(Guide to the Science of). By EDWARD C. BOUSFIELD, L.R.C.P. Lond. 8vo, with 34 Engravings and Frontispiece, 6s.

- Introduction to Physical Measurements, with Appendices on Absolute Electrical Measurements, &c. By Dr. F. Kohlrausch, Professor at the University of Strassburg. Third Edition, translated from the Seventh German Edition, by Thomas Hutchinson Waller, B.A., B.Sc., and Henry Richardson Procter, F.I.C., E.C.S. Svo. with of Illustrations, 12s. 6d. F.C.S. 8vo, with 91 Illustrations, 12s. 6d.
- Tuson's Veterinary Pharmacopœia, including the Outlines of Materia Medica and Therapeutics. Fifth Edition. Edited by JAMES BAYNE, F.C.S., Professor of Chemistry and Toxicology in the Royal Veterinary College. Crown 8vo, 7s. 6d.
- The Principles and Practice of Veterinary Medicine. By WILLIAM WILLIAMS, F.R.C.V.S., F.R.S.E., Principal, and Professor of Veterinary Medicine and Surgery at the New Veterinary College, Edinburgh. Seventh Edition. 8vo, with several Coloured Plates and Woodcuts, 30s.

By the same Author.

- The Principles and Practice of Veterinary Surgery. Eighth Edition. 8vo, with 9 Plates and 147 Woodcuts, 30s.
- The Veterinarian's Pocket Remembrancer: being Concise Directions for the Treatment of Urgent or Rare Cases, embracing Semeiology, Diagnosis, Prognosis, Surgery, Therapeutics, Toxicology, Detection of Poisons by their Appropriate Tests, Hygiene, &c. By George Armatage, M.R.C.V.S. Second Edition. Post 8vo, 3s.
- Chauveau's Comparative Anatomy of the Domesticated Animals. Revised and Enlarged, with the Co-operation of S. ARLOING, Director of the Lyons Veterinary School, and Edited by George Fleming, C.B., LL.D., F.R.C.V.S., late Principal Veterinary Surgeon of the British Army. Second English Edition. 8vo, with 585 Engravings, 31s. 6d.

INDEX TO J. & A. CHURCHILL'S LIST.

Abercrombie's Medical Jurisprudence, 2 Adams (W.) on Clubfoot, 9 - on Contractions of the Fingers, &c., 9 on Contractions of the Fingers, &c., 9
on Curvature of the Spine, 9
Allen's Commercial Organic Analysis, 13
Allingham (W.) on Diseases of the Rectum, 12
Armatage's Veterinary Pocket Remembrancer, 14
Auld's Bronchial Affections, 6
Barnes (R.) on Obstetric Operations, 3
on Diseases of Women, 3

Reale (L. S.) on Liver, 6 Beale (L. S.) on Liver, 6
Microscope in Medicine, 6 Slight Ailments, 6 Beale (P. T. B.) on Elementary Biology, 2
Beasley's Book of Prescriptions, 5

Druggists' General Receipt Book, 5 Pocket Formulary, 5
Bellamy's Surgical Anatomy, 1
Bentley and Trimen's Medicinal Plants, 5
Bentley's Systematic Botany, 5
Berkart's Bronchial Asthma, 6
Bernard on Stammering, 5 Bernard on Stammering, 7 Bernay's Notes on Analytical Chemistry, 1 Bigg's Short Manual of Orthopædy, 9 Bloxam's Chemistry, 12

Laboratory Teaching, 12

Bousfield's Photo-Micrography, 14

Bowlby's Injuries and Diseases of Nerves, 9

Surgical Pathology and Morbid Anatomy, 9 Bowman and Bloxam's Practical Chemistry, 13
Brodhurst's Anchylosis, 9
— Curvatures, &c., of the Spine, 9
— Talipes Equino-Varus, 9
Bryant's Practice of Surgery, 8
— Tension, Inflammation of Bone, Injuries, &c., 8
Buist's Various and Various Buist's Vaccinia and Variola, 7 Burckhardt's (E.) and Fenwick's (E. H.) Atlas of Burckhardt's (E.) and Fenwick's (E. H.) Atlas of Cystoscopy, 11

Burdett's Hospitals and Asylums of the World, 2

Butlin's Malignant Disease of the Larynx, 11

Operative Surgery of Malignant Disease, 11

Sarcoma and Carcinoma, 11

Buzzard's Diseases of the Nervous System, 7

Peripheral Neuritis, 7

Simulation of Hysteria, 7

Cameron's Oils, Resins, and Varnishes, 14

Soaps and Candles, 14

Carpenter and Dallinger on the Microscope, 14

Carpenter's Human Physiology, 2

Charteris on Health Resorts, 8

Practice of Medicine, 6 Practice of Medicine, 6 Chauveau's Comparative Anatomy, 14 Chevers' Diseases of India, 5 Churchill's Face and Foot Deformities, 9 Clarke's Eyestrain, 10 Clouston's Lectures on Mental Diseases, 2 Clowes and Coleman's Quantitative Analysis, 12 Elementary Analysis, 12 Dalby's Diseases and Injuries of the Ear, 10 Short Contributions, 10 Day on Diseases of Children, 4 — on Headaches. 8

Domville's Manual for Nurses, 4

Doran's Gynæcological Operations, 3

Druitt's Surgeon's Vade-Mecum, 8

Duncan (A.), on Prevention of Disease in Tropics, 5

Duncan (J. M.), on Diseases of Women, 3

Dunglison's Dictionary of Medical Science, 12

Ellis's (E.) Diseases of Children, 4

Ellis's (T. S.) Human Foot, 9

Ewart's Bronchi and Pulmonary Blood Vessels, 6

Fagge's Principles and Practice of Medicine, 6

Fayrer's Climate and Fevers of India, 5

Natural History, &c., of Cholera, 5 on Headaches. 8 Fenwick (E. H.), Electric Illumination of Bladder, 11
Symptoms of Urinary Diseases, 11

Fenwick's (S.) Medical Diagnosis, 6

Obscure Diseases of the Abdomen, 6 Outlines of Medical Treatment, 6 The Saliva as a Test, 6 Fink's Operating for Cataract, 10 Fink's Operating for Catalact, 10
Flower's Diagrams of the Nerves, 1
Fowler's Dictionary of Practical Medicine, 6
Fox's (C. B.) Examinations of Water, Air, and Food, 2
Fox's (T.) Atlas of Skin Diseases, 10
Fox (Wilson), Atlas of Pathological Anatomy of Lungs, 6 Treatise on Diseases of the Lungs, 6
Frankland and Japp's Inorganic Chemistry, 13
Fraser's Operations on the Brain, 8
Fresenius' Qualitative Analysis, 13
— Quantitative Analysis, 13
Galabin's Diseases of Women, 3
— Manual of Midwifery, 3
Gardner's Bleaching, Dyeing, and Calico Printing, 14
— Brewing, Distilling, and Wine Manuf. 14
Godlee's Atlas of Human Anatomy, 1
Goodhart's Diseases of Children, 4
Gowers' Diseases of the Brain, 7
— Manual of Diseases of Nervous System, 7 Treatise on Diseases of the Lungs, 6 Manual of Diseases of Nervous System, 7 Medical Ophthalmoscopy, 7 Syphilis and the Nervous System, 7 Granville on Gout, 7
Green's Manual of Botany, 5
Groves' and Thorp's Chemical Technology, 14
Guy's Hospital Reports, 7
Habershon's Diseases of the Abdomen, 7
Haig's Uric Acid, 6
Harley on Diseases of the Liver, 7
Harris's (V. D.) Diseases of Chest, 6
Harrison's Urinary Organs, 12
Hartridge's Refraction of the Eye, 10 Harrison's Orinary Organs, 12

Hartridge's Refraction of the Eye, 10

— Ophthalmoscope, 10

Hawthorne's Galenical Preparations of B P., 4

Heath's Certain Diseases of the Jaws, 8

— Clinical Lectures on Surgical Subjects, 8

— Injuries and Diseases of the Jaws, 8 Minor Surgery and Bandaging, 8 Operative Surgery, 8 Practical Anatomy, 1 Surgical Diagnosis, 8 Hellier's Notes on Gynæcological Nursing, 4 Higgens' Ophthalmic Out-patient Practice, 10 Hillis' Leprosy in British Guiana, 10 Hirschfeld's Atlas of Central Nervous System 2 Holden's Human Osteology, 1 Landmarks, 1
Hooper's Physicians' Vade-Mecum, 5
Hovel's Diseases of the Ear, 10
Howdon's Index Political Politics Howden's Index Pathologicus, 2 Hutchinson's Clinical Surgery, 9 Hyde's Diseases of the Skin, 10 Jacobson's Male Organs of Generation, 12 Operations of Surgery, 9 Johnson's Asphyxia, 6
— Medical Lectures and Essays, 6
Journal of Mental Science, 3
Keyes' Genito-Urinary Organs and Syphilis, 12
Kohlrausch's Physical Measurements, 14 Lancereaux's Atlas of Pathological Anatomy, 2 Lane's Rheumatic Diseases, 7 Langdon-Down's Mental Affections of Childhood, 3 Lee's Microtomists' Vade Mecum, 14 Lee's Microtomists' Vade Mecum, 14
Lescher's Recent Materia Medica, 4
Lewis (Bevan) on the Human Brain, 2
Liebreich's Atlas of Ophthalmoscopy, 10
Macdonald's (J. D.) Examination of Water and Air, 2
MacMunn's Clinical Chemistry of Urine, 12
Macnamara's Diseases and Refraction of the Eye,

of Bones and Joints, 8

M. N. J. Bridamics and Isolation Hospitals, 2 McNeill's Epidemics and Isolation Hospitals, 2
Malcolm's Physiology of Death, 4
Mapother's Papers on Dermatology, 10
Martin's Ambulance Lectures, 8
Maxwell's Terminologia Medica Polyglotta, 12
Mayne's Medical Vocabulary, 12
Mayrian's Lungory Law, 2 Mercier's Lunacy Law, Mercier's Lunacy Law, 3
Microscopical Journal, 14
Mills and Rowan's Fuel and its Applications, 14
Moore's (N.) Pathological Anatomy of Diseases, 1
Moore's (Sir W. J.) Family Medicine for India, 5

Manual of the Diseases of India, 5 Tropical Climates, 5 [Continued on the next page.

Morris's Human Anatomy, 1 Moullin's (Mansell) Surgery, 8 Moullin's (Mansell) Surgery, 8
Nettleship's Diseases of the Eye, 9
Ogle on Puncturing the Abdomen, 8
Oliver's Abdominal Tumours, 3
— Diseases of Women, 3
Ophthalmic (Royal London) Hospital Reports, 9
Ophthalmic (Royal London) Hospital Reports, 9
Ophthalmological Society's Transactions, 9
Ormerod's Diseases of the Nervous System, 7
Owen's Materia Medica, 4
Parkes' (E.A.) Practical Hygiene, 2
Parkes' (L.C.) Elements of Health, 2
Pavy's Carbohydrates, 6
Pereira's Selecta è Prescriptis. 4
Phillips' Materia Medica and Therapeutics, 4
Pitt-Lewis's Insane and the Law, 3
Pollock's Histology of the Eye and Eyelids, 9
Proctor's Practical Pharmacy, 4
Purcell on Cancer, 11 Richardson's Mechanical Dentistry, 10 Roberts' (D. Lloyd) Practice of Midwifery, 3 Robinson's (Tom) Eczema, 11 - Illustrations of Skin Diseases, 11 - Syphilis, 11 Ross's Aphasia, 7

Diseases of the Nervous System, 7

Royle and Harley's Materia Medica, 5

St. Thomas's Hospital Reports, 7

Sansom's Valvular Disease of the Heart, 6

Savage's Female Pelvic Organs, 3

Schetelig's Homburg-Spa, 8

Schweinitz's (G. E. de) Diseases of Eye, 10

Shaw's Diseases of the Eye, 9 Shaw's Diseases of the Eye, 9
Short Dictionary of Medical Terms, 12
Silk's Manual of Nitrous Oxide, 10
Smith's (E.) Clinical Studies, 4 - Diseases in Children, 4 - Wasting Diseases of Infants and Children, 4 Smith's (J. Greig) Abdominal Surgery, 4 Smith's (Priestley) Glaucoma, 10 Snow's Cancer and the Cancer Process, 11 - Palliative Treatment of Cancer, 11 — Reappearance of Cancer, 11
Squire's (P.) Companion to the Pharmacopœia, 4
— London Hospitals Pharmacopœias, 4
— Methods and Formulæ, 14
Starling's Elements of Human Physiology, 2

Stevenson and Murphy's Hygiene, 2 Stillé and Maisch's National Dispensatory, 5 Stocken's Dental Materia Medica and Therapeutics, 10 Stocken's Dental Materia Medica and Therapeutics Sutton's (H. G.), Lectures on Pathology, 1
Sutton's (J. B.), General Pathology, 1
Sutton's (F.) Volumetric Analysis, 13
Swaine's Surgical Emergencies, 8
Swayne's Obstetric Aphorisms, 3
Taylor's (A. S.) Medical Jurisprudence, 2
Taylor's (F.) Practice of Medicine, 6
Taylor's (J. C.), Canary Islands, 8
Thin's Cancerous Affections of the Skin, 11
— Pathology and Treatment of Ringworm, 11
Thomas's Diseases of Women, 3
Thompson's (Sir H.) Calculous Disease, 11
— Diseases of the Prostate, 11 Diseases of the Prostate, 11 Diseases of the Urinary Organs, 11 Introduction to Catalogue, 11 Lithotomy and Lithotrity, 11 Stricture of the Urethra, 11 Suprapubic Operation, 12 Surgery of the Urinary Organs, 11 Tumours of the Bladder, 11 Thorne's Diseases of the Heart, 7 Tirard's Prescriber's Pharmacopoeia, 5 Tomes' (C. S.) Dental Anatomy, 10 Tomes' (J. and C. S.) Dental Surgery, 10 Tommasi-Crudeli's Climate of Rome, 7 Tommasi-Crudeli's Climate of Rome, 7
Tooth's Spinal Cord, 7
Treves and Lang's German-English Dictionary, 12
Tuke's Dictionary of Psychological Medicine, 3
— Influence of the Mind upon the Body, 3
Tuson's Veterinary Pharmacopeia, 14
Valentin and Hodgkinson's Qualitative Analysis, 13
Vintras on the Mineral Waters, &c., of France, 8
Wagner's Chemical Technology, 13
Walsham's Surgery: its Theory and Practice, 8
Waring's Indian Bazaar Medicines, 5
— Practical Therapeutics, 5
Watts' Manual of Chemistry, 13 Practical Therapeutics, 5
Watts' Manual of Chemistry, 13
West's (S.) How to Examine the Chest, 6
Westminster Hospital Report, 7
White's (Hale) Materia Medica, Pharmacy, &c., 4
Wilks' Diseases of the Nervous System, 7
Williams' Veterinary Medicine, 14
Surgery, 14 Wilson's (Sir E.) Anatomists' Vade-Mecum, 1 Wilson's (G.) Handbook of Hygiene, 2 Wolfe's Diseases and Injuries of the Eye, 9 Wynter and Wethered's Practical Pathology, 1 Year-Book of Pharmacy, 5 Yeo's (G. F.) Manual of Physiology, 2

N.B.—J. & A. Churchill's larger Catalogue of about 600 works on Anatomy, Physiology, Hygiene, Midwifery, Materia Medica, Medicine, Surgery, Chemistry, Botany, &c. &c., with a complete Index to their Subjects, for easy reference, will be forwarded post free on application.

AMERICA.—J. & A. Churchill being in constant communication with various publishing houses in America are able to conduct negotiations favourable to English Authors.





