Hygiene and public health / by B. Arthur Whitelegge and George Newman.

Contributors

Newman, George, Sir, 1870-1948. Whitelegge, Arthur, Sir, 1852-1933. Royal College of Physicians of Edinburgh

Publication/Creation

London: Cassell, 1908.

Persistent URL

https://wellcomecollection.org/works/eankdax3

Provider

Royal College of Physicians Edinburgh

License and attribution

This material has been provided by This material has been provided by the Royal College of Physicians of Edinburgh. The original may be consulted at the Royal College of Physicians of Edinburgh. where the originals may be consulted.

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Hygiene and Public Health

Whitelegge and Newman

Fa.1.48.

By order of the College, this Book is not to be taken out of the Library (except after 6 P.M. until 10 A.M.) for one month from this date. PHYSICIANS' HALL, 19th June 1908

HYGIENE

4

HYGIENE AND PUBLIC HEALTH

BY

B. ARTHUR WHITELEGGE, C.B.

M.D., B.Sc. LOND., F.R.C.P., D.P.H.

Late County Medical Officer of Health for the West Riding of Yorkshire, Chief Sanitary Officer of the West Riding of Yorkshire Rivers Board, Marcal F Officer of Health for Nottingham, and Lecturer on Public Health, Charing Cross Hospital Medicall School

GEORGE NEW

M.D., D.P.H., F.R.S.E.

Lecturer on Public Health at St. Bartholomew's Hospital (University of London); late Medical Officer of Health of the Metropolitan Borough of Finsbury, Consulting Medical Officer of Health of the Bedfordshire County Council, and Demonstrator of Bacteriology and Infective Diseases in King's College, London

ILLUSTRATED

NEW AND REVISED EDITION

CASSELL AND COMPANY, LIMITED LONDON, PARIS, NEW YORK, TORONTO AND MELBOURNE 1908

HYGIENE, AND PUBLIC HEALTH

the Proposition of Company of the Commission of

PREFACE TO THE NEW EDITION

In order to retain in the present volume the advantages secured by a Manual of this size and nature, the authors have thought it desirable to expand the enlarged edition of 1905 as little as possible. Even so, it has been necessary, in order to bring the book down to date, to include in this Eleventh Edition a large number of additional facts and records. Every year now brings its own substantial contribution of new work, and particularly of new legislation, to the common stock of the science and practice of Preventive Medicine.

This Manual still seeks, as it originally sought, to provide a concise summary of the current knowledge concerning Public Health for the purposes of the Health Officer in every branch of the public service, as well as an elementary text-book for the student.

B. A. W.

G. N.

PHERACE TO THE NEW EDITION

request of the file of the size of the siz

considered Temperature Medicine

the provide a country was in the distribute the country and the provided to th

A 10

2001 MARKE

CONTENTS

CHAPTER										P	AGE
In	TRODUCTION	V						1			1
I.—	AIR							•			3
II.—N	METEOROLOG	Y .					100	100			18
III.—	WATER.			1911							37
IV.—I	FOOD		. 07								90
V	Soil .	angs.		W.			. 7				159
VI]	BUILDINGS .				. 74						172
VII.—	SCHOOLS AND	Scно	DL H	YGII	ENE						203
VIII.—I	HOSPITALS										222
IX.—I	REMOVAL OF	REFUS	SE								235
X	DISPOSAL OF	THE D	EAD								277
XI.—	ANIMAL PAI	RASITES									284
XII.—	INFECTION			,							294
XIII.—	DISINFECTIO	N .									318
XIV.—	SPECIFIC DIS	SEASES									334
XV	TROPICAL D	ISEASES							. 11		431
XVI	PREVENTION	of In	FECT	ious	DI	SEAS	ES				457
XVII	MEDICAL O		OF	HE	CALT	н А	ND	SANI	TAR	Y	
	INSPECTO	ORS .								100	473
XVIII	SANITARY L	AW .									490
XIX.—	FACTORIES A	AND WO	RKS	нор	S			4			557
XX	BYELAWS AN	ND REG	ULAT	TIONS	3			12000		-	578
XXI	VITAL STAT	ISTICS		MI		10			111		605

APPENDICES:—	PAGE
I.—URBAN AND RURAL DEATH RATES, 1891-1900 (Eng-	
LAND AND WALES)	631
II.—DEATH RATES AT AGES IN ENGLAND AND WALES	
(Decennia), 1860-1900	631
III.—Population of England and Wales, 1861-1901 .	632
IV.—Age and Sex Distribution of Population in	
ENGLAND AND WALES, 1901	632
VBIRTH RATES AND DEATH RATES IN ENGLAND AND	
Wales (Decennia), 1851-1900	632
VI.—DEATH RATES PER MILLION, ENGLAND AND WALES,	
1875-1900. Causes and Quinquennia	633
VII.—DEATH RATES PER MILLION, ENGLAND AND WALES,	
1891-1900. CAUSES AND AGES	633
VIIIDEATH RATES PER MILLION, ENGLAND AND WALES,	
1891-1900. CAUSES AND SEX	634
IX -LIFE TABLE, 1891-1900	634
INDEX	10222

HYGIENE

AND

PUBLIC HEALTH

INTRODUCTION

Public Health administration rests upon a wide range of sciences, almost all of which are in a progressive stage. The object of this elementary Manual is to summarise the more important practical applications of these in preventive medicine, and especially in the work of Medical Officers of Health, under present conditions; and where competent authorities differ in opinion or practice, to give prominence to official views generally accepted rather than to those of individuals. Readers who desire fuller technical detail will turn to special books of medicine, epidemiology, bacteriology, chemistry, physics, meteorology, geology, engineering and statistics; to the official publications of the Local Government Board, the Registrar-General, and other Government medical services at home and abroad; to the Reports of Royal Commissions and Departmental Committees; to the Reports of Medical Officers of Health and

School Medical Officers; and to the technical Journals and the Transactions of the many learned societies concerned with Public Health.

Any review of such a field must, it is obvious, be tentative and provisional in many points. Improved methods and appliances, fuller and later statistics and research, and new conceptions, are brought forward year by year. These, together with further legislation, are recorded in the current literature of the subject. Finally, whilst care has been taken to furnish a reliable and comprehensive statement in the present volume, it cannot be too clearly understood that practical work in administration, in the laboratory, and in the hospital, is essential to a thorough grasp of many of the principles discussed in the following pages.

the the said bears owned to each the control of the last the control of the last the control of the control of

CHAPTER I

AIR

In 100 volumes of pure dry air there are, approximately, 21 of oxygen, 78 of nitrogen, and 0.9 of argon. There are also other gases in much smaller proportion. In practice, however, air is never quite pure or dry, and it is therefore necessary to take account of watery vapour, dust, and gaseous impurities of local origin.

Oxygen ranges from 20.99 per cent. by volume in sea or mountain air to 20.7 in crowded rooms, 18.3 in ill-ventilated mines, and 16.5 in expired air. It is lessened by respiration, by combustion, by oxidisable organic effluvia, and by fog; but it is increased by vegetation and by rain. The average in the open air of towns may be taken as 20.96

volumes per cent.

Ozone, a more active form of oxygen, with the molecular formula O_3 , instead of O_2 , is an irrespirable gas readily reduced by oxidisable matter. It is only an occasional component of air, and is never present in more than minute traces, difficult of exact measurement. The proportion is greatest after thunderstorms, always higher on the coast than inland, and usually higher with S.W. rather than N.E. winds.

Nitrogen and argon are inert, and, from the point of view of public health, may be looked upon as

diluents only.

Carbonic acid gas (CO₂), said to be absent from desert air, varies from 2.5 per 10,000 in purest mountain or sea air to 20.0 or even 30.0 parts in 10,000 in crowded rooms. In fresh open air it is stated to be, as a rule, 4.0 per 10,000, but this figure, based upon Pettenkofer's process, appears to be too

high, and more exact measurement points to 3.0 per 10,000 as the correct standard. The proportion is greatly increased by expiration, for in expired air there is not infrequently as much as 450 parts per 10,000 (4.5 per cent.). The proportion of CO₂ in air is also increased by combustion, putrefaction, and fog, and it is diminished by vegetation, rain, high winds, and, in rooms, by ventilation. At great elevation the proportion is increased, owing possibly to the complete oxidation of organic matter (Angus Smith). Both carbonic acid and organic matter are greater in the air of towns than in the country, and in day air than night air. In railway tunnels Clowes found an average of 14 parts of CO₂ per 10,000, and on one occasion as much as 28.8 parts.

Ammonia salts in air, like carbonic acid gas, are due mainly to animal exhalations and putrefactive processes. Fresh air contains from 0.03 to 0.1 mgr. of carbonate or nitrate of ammonia per cubic metre, but in crowded rooms, or, for example, over middens, there may be 0.50 mgr., or even more. The variations are generally parallel with those of carbonic acid, and

are similarly affected by weather.

Marsh gas or methane (CH₄) is met with in the air of coal mines and marshes. In the former case the proportion may be sufficient to destroy life by the exclusion of oxygen, but as much as 300 parts per 1,000 can be breathed with comparative impunity. If present in certain proportions it forms an explosive mixture with air.

Carbonic oxide (CO) is present in minute quantity in the air of towns, but in larger degree in, say, that of a railway tunnel. Coal gas contains from 4 to 12 per cent., but "water gas" and "producer

^{*} Under various names gases of this nature are now largely made and supplied to the public, both separately and as an auxiliary to an ordinary coal gas supply. They have, naturally, little lighting power, but can be

gas" as much as 40 per cent. Carbonic oxide is found as a product of imperfect combustion, particularly of charcoal, and cast iron stoves or burners may give it off in considerable quantity. Air containing 0.05 parts per 1,000 causes unpleasant, and eventually poisonous, symptoms in persons inhaling it. Less than a gramme of CO may kill a man (Oliver). The toxic properties of carbon monoxide are due to the fact that the gas enters into direct combination with the hæmoglobin of the blood, forming with it an extremely stable compound, making the blood useless for respiratory purposes. Persons thus poisoned die, therefore, simply from want of oxyhæmoglobin.

Sulphurous acid (SO₂) is another product of combustion, and traces of it are found in town air, least in clear breezy weather, most during fog and under anticyclonic conditions. Coal stoves and gas burning are the chief sources of SO₂ in the air of rooms.

Sulphuretted hydrogen (H₂S) may occur in the air of marshes, in and near excavations, in collections of refuse or decaying vegetable matter, in sewer gas, and in chemical works and their waste heaps.

The inhalation of H₂S is followed by more poisonous results than is commonly supposed. An atmosphere containing 0.7 to 0.8 of H₂S in 1,000 litres of air is dangerous to human life, while air containing 1 to 2 per 1,000 parts destroys in a few minutes. When only minute quantities are present, vertigo, headache, and malaise are produced. Two types of poisoning occur, namely, that resulting in sudden death due to the action of H₂S on the respiratory nerve centres, and that resulting in slow

charged with oil-spray so as to be equal to ordinary gas, and for certain trade processes luminosity is not required. Their use calls for great care in gas fittings, and as a further precaution the gas should be "odorised," that is, charged with some strong-smelling substance. In certain Acts authorising works of supply for "Mond gas," a limit of 14 per cent. of CO is imposed.

poisoning and death by asphyxia in addition to

nervous phenomena.

It may here be stated that the various constituent parts of air are not chemically combined, but only mechanically mixed. The proofs of this are (a), that the relative proportions of oxygen and nitrogen present in air are not those of their combining weights, or of any simple multiple of those weights; (b), if air be artificially made by mixing oxygen and nitrogen together in proper proportions, no change of volume or temperature takes place as would be found had chemical combination occurred; and (c), if air and water be shaken together the water dissolves some of the air, which can be subsequently expelled by boiling, when the air which is expelled will be found to contain 14 per cent. more oxygen, which signifies that the atmosphere is not a stable chemical compound.

Physical properties. -(1) Weight. At a temperature of 32°F., the barometer being at 29.9 inches, 100 cubic inches of air weigh about 32.5 grains, and a cubic foot, 573 grains. (2) Compression and expansion. The stratum of atmosphere near the earth's surface is more dense because more subjected to pressure than the stratum remote from the earth's surface. The air is subject to Boyle and Marriotte's law: the temperature remaining the same, the volume of a given weight of air is inversely as the pressure which it bears. Under extreme pressure, with very low temperature, we now know that it may be liquefied (Dewar). The converse of compression is also true, and air is capable, in common with other gases, of expansion. This, too, follows a law, that of Charles: the volume of a given weight of air at a constant pressure is proportional to its absolute temperature, or, in other words, if the volume remains unaltered, the pressure varies as the temperature, and if the pressure remains constant the volume varies as the temperature. The increase of the unit of volume of a gas for one degree of temperature, that is, from zero to one degree, is termed the co-efficient of expansion, and is practically the same for all gases $(\frac{1}{273})$ of their volume). (3) Diffusion. If two different gases in receivers be brought into communication with each other, they will diffuse the one into the other, and similar diffusion occurs in the atmosphere. According to "Graham's Law" the diffusibility of two gases varies in inverse ratio of the square roots of their densities.

Bacteria in air vary in number and species according to certain external circumstances, of which the following are the chief, namely, the pollution of the air, the dampness of surrounding surfaces, gravity, and various seasonal and meteorological conditions.

(a) The pollution of air. It was Tyndall who in 1878 first pointed out that dust in air might carry micro-organisms ("the raft theory"), and that, other things being equal, dusty air contained more bacteria than dust-free air. Since that time Haldane and others have confirmed this by examination of the air of workshops, schoolrooms, etc. In open fields, free from habitations, few bacteria are present, and the same is true of mountain and sea air; in towns or crowded rooms the reverse is the case. Miquel found an average of 455 bacteria per cubic metre in the park of Mont Souris, but an average of 3,910 in the Rue de Rivoli, and 79,000 in a hospital ward. Haldane found 8 per litre in a dressmaker's workroom, but 850 per litre in a dusty rope-maker's room; Graham Smith found 32 per litre in the dining room of the House of Commons, and 1 per litre in the open air of the Terrace; and Andrewes, investigating the air of the Central London Railway (underground tube), found 12 per litre on the lift and 21 per litre on the platform. Carnelly, Haldane, and Anderson found 6 bacteria per 10 litres in open air, 85 in houses of four or more rooms, 430 in houses of two rooms, and 580 per 10 litres in houses of one room. A polluted or dusty atmosphere generally contains many bacteria and much carbonic acid, but there seems to be no direct relation between them. An atmosphere may be offensive and yet comparatively free from bacteria, as in sewer gas, certain offensive trades, railway

tunnels, etc.

(b) The dampness of surrounding surfaces. Air over sandy soils contains more bacteria as a rule than that over damp clay soils. Rain and snow also diminish the number of organisms in air. It has also been proved that air contained within a moist perimeter is almost germ-free. Hence the comparative absence of bacteria in expired air in ordinary quiet respiration, though in the act of coughing, sneezing, or shouting organisms may be present in the expired air (Flügge). Haldane, Andrewes, and others have shown that the same principle applies in a sewer, the air of which frequently contains fewer organisms than that of outside air, the species found belonging not to the sewage but to outside air. This is also the explanation of the retention of bacteria in sputum (the tubercle-bacillus), or excreta (the typhoid bacillus), so long as these materials remain moist.

(c) The influence of gravity operates upon microscopic cells in the same way, of course, as upon other particulate matter. Hence, fewer bacteria are found at high altitudes or on the tops of lofty buildings. Binot found from 4 to 11 microbes per cubic metre of air collected at the summit of Mont Blanc. Miquel found 28 bacteria per cubic metre on top of the Pantheon, and 750 at the street level; and Graham Smith found that at the top of the Clock Tower of the Houses of Parliament in London there was only about one-third of the number of bacteria

found at ground level.

(d) Seasonal and meteorological conditions also exert an influence upon the number of bacteria found in air. The seasonal maximum in the open air seems to occur about midsummer, and the minimum about the middle of winter, but in hospital wards and houses the reverse occurs, owing no doubt to ventilation. Air currents and wind, of course, exert a marked influence, as also do rain and snow. Direct sunlight possesses considerable bactericidal power, and thus reduces the number of air organisms.

Unlike the organic matter and carbonic acid, the number of microbes in the air of crowded rooms is dependent rather upon habitual ventilation and cleanliness than upon the conditions at the time of observation. They are not due, as we have seen, to respiration, or as a rule to want of cleanliness of persons or clothing, but are derived from the walls, ceiling, and floor of the room itself, especially if these are porous and absorbent, and if made to vibrate and thus create dust. As to the nature of the microbes found in the air, little that is final can be said. Their variety is considerable, and for the most part they are saprophytes and harmless to man, but they may include putrefactive, suppurative, and intestinal organisms, and the specific germs of disease. Under exceptional conditions bacilli of tubercle, typhoid fever and other diseases have been isolated from air, and Gordon has detected an "airstreptococcus" and a "skin staphytococcus," indicative of filth and human pollution.

Spores of moulds, though more bulky than most other air-borne organisms, are less readily deposited. Other low forms of organisms and pollen may also be found, and may be the source of certain forms of

disease (hay-fever).

Dust.—Aitken found the number of dust particles per cubic inch of air to range from 2,000 in the

open country to 3,000,000 in towns, and 30,000,000 in rooms. Apart from living organisms, dust consists chiefly of mineral particles, and of formed or unformed organic matter of vegetable or animal origin, e.g. epithelium scales, pollen, or particles of flax, wool, or cotton.

Indoor dust contains more organic components than that of the open air, although open fires add largely to the mineral part. The heavier particles subside readily in still air, the lighter more slowly, but practically deposition is always taking place, even in closed rooms and spaces, since air carrying dust is constantly passing in or out through crevices with variations of temperature and pressure. Hence the only dust-proof spaces are those which are kept hermetically sealed or in which there is provision for free entry of air through a dust-filter, as, for example, a closed bookcase, the back of which consists of baize or similar material.

Respiration. — Respiration abstracts oxygen from air and adds carbonic acid gas, water, organic matter, and a little ammonia. An average adult gives off about 0.6 cubic foot * of carbonic acid per hour, or 17 litres, of which perhaps about one-fiftieth part is yielded by the skin (Regnault). The amount is less in women, children, and aged persons, but more in animals, such as oxen or horses, which give off about three times as much CO₂ as does man.

Expired air, as we have seen, contains about 4.5 per cent. of carbonic acid, and only about 16 per cent. of oxygen. As a rough average it may be stated that 9-10 oz. of watery vapour are given off by the human lungs in 24 hours, and 20-25 oz. by the skin. In health, expired air contains few, if any, microbes, but much organic matter. It is

^{* 0:56} during sleep; 0.78 when awake, (Pettenkofer,)

nitrogenous, oxidisable, molecular, and probably exists in combination with water, since it is readily absorbed by hygroscopic objects. It is precipitated from solution by silver nitrate, blackens on ignition, reduces potassium permanganate, yields ammonia on distillation with alkaline permanganate, and has a fætid smell. Unlike carbonic acid, organic matter diffuses sluggishly through the air of a room, and is destroyed slowly by fresh air. It promotes the growth of micro-organisms, in which respect it appears to differ from CO_2 ; and milk, meat, or other foods in contact with it readily become tainted. The production per head of both carbonic acid and organic matter is increased after physical exercise.

Combustion. — Coal when burnt gives off — (1) carbonic acid, together with carbonic oxide unless the combustion is complete (it is estimated that each ton of coal burnt represents nearly 3 tons of CO₂ added to the atmosphere); (2) sulphurous and sulphuric acids (1–5 per cent. sulphur exists in coal); carbon bisulphide, ammonium sulphide, and sometimes sulphuretted hydrogen; (3) water; (4) fine

particles of carbon and tarry matter.

Coal gas yields similar products, but there is little free carbon or organic matter. The sulphur in gas should not exceed 20 grains per 100 cubic feet. The Alkali Works Regulation Act (1906) provides that not more of the acid gases of sulphur and nitrogen than the equivalent of 4 grains of sulphuric anhydride per cubic foot of air, gas, or smoke, must escape into the atmosphere.

Wood yields little of the sulphur compounds. Oil and candles give scarcely any, but they increase

the organic matter in the air.

The gaseous products, especially carbonic acid, are rapidly diffused by air-currents; but the suspended particles diffuse slowly, and are not as a rule found

at a greater elevation than 600 feet. Impurity from the use of illuminants also pollutes the air, and at the same time exhausts it. An ordinary bat's-wing burner (16 candle power), consuming $5\frac{1}{2}$ cub. ft. of gas in one hour, has been estimated to produce 2.8 cub. ft. of CO_2 and 7.3 cub. ft. of moisture, thus vitiating the atmosphere to the extent which five adults would do in respiration. There is also abstraction of oxygen.

Impurities of air include infective matter, organic effluvia, fumes, gases, and dust (poisonous or otherwise). Their relation to disease is complex,

and will be discussed later on.

Examination of air.—1. Particulate matter and bacteria suspended in the air may be collected for microscopic examination by aspirating the air through water; or by allowing a jet of air to pass into an exhausted receiver through a fine aperture, so as to impinge upon glass smeared with glycerine, which arrests the particles (Pouchet's aëroscope); or by aspirating air through a filter of powdered sugar, and by dissolving away the sugar in water, obtaining the intercepted particles available for examination. By measuring the volume of air aspirated, and counting the particles in an aliquot part of the intercepting material, these methods can be made quantitative.

Micro-organisms will, for the most part, escape detection by this method, and, indeed, no means exist of determining their number with accuracy. A rough approximation may be obtained by making a plate-cultivation from the whole (or an aliquot part) of the material obtained in the manner described above, and counting the colonies; but for obvious reasons the result gives only the number of organisms which can grow under one set of conditions. Hesse aspirates air slowly through a wide horizontal glass tube smeared on the inside with nutrient gelatine, the air entering through a pin-hole at such speed as to allow

suspended particles to fall upon the gelatine. Sedgwick, Frankland, Petri, Pasteur, and others have adopted methods of filtration through sugar, sand, etc., which is subsequently cultured in favourable media. The amount of air examined is measured by a syringe or aspirator of known capacity. Koch relied on exposure of plates of gelatine to the air to be examined. Gordon prefers using "broth-plates" in the same way, with anærobic cultivation of the broth at 37 C. for 48 hours with subsequent subculture. Search should be made for streptococci (as indices of saliva), B. coli and B. enteritidis sporogenes (as index of recent and non-recent fæcal pollution), and B. mycoides (as index of soil contamination).

2. Organic matter in air may be measured by determining the volume of air required to decolorise a known volume of a standard solution of permanganate of potash, such as 0.395 grm. in 1 litre of distilled water. The air may be aspirated through the solution, or successive volumes washed with the permanganate, by shaking in a bottle, until the solution is no longer

pink.

Another plan is to aspirate a known large volume of air through distilled water, and then examine the latter by the albuminoid ammonia process, described subsequently. In such estimations it is desirable in each case to make control experiments.

Carbonic acid.—1. Pettenkofer's method.—The solutions required are: (a) Pure lime-water, saturated or nearly so; (b) Standard solution of crystallised oxalic acid of such strength (2.25 grammes per litre) that 1 c.c. exactly neutralises 1 mgr. of lime, CaO. From the equation CaO + CO₂ = CaCO₃

56 mgr. 44 mgr. 100 mgr.

it is evident that 1 mgr. of CaO takes $\frac{44}{56}$ mgr. of CO₂. The volume of 44 mgr. of CO₂ at O·C and 760 mm. pressure, is 22.4 c.c., so that $\frac{44}{56}$ mgr. = 0.4 c.c. Hence, each c.c. of oxalic

solution corresponds to 0.4 c.c. of carbonic acid, either being capable of neutralising exactly 1 mgr. of lime.

A wide-mouthed glass bottle, provided with a tight-fitting glass or rubber stopper, is also necessary. Its capacity should

be about five litres, and must be determined accurately.

The bottle, having been cleaned and dried, is filled with sample air, either by means of a bellows, or other dry method. Moisture is held to invalidate the result. Sixty c.c. of clear lime-water are put into the bottle, the stopper is inserted and the air thoroughly washed with lime-water by means of vigorous shaking. The carbonic acid combines with lime and forms a milky precipitate of carbonate of lime; the loss of strength of the lime-water therefore measures the amount of carbonic acid present. After the bottle has stood at rest for a few hours, 30 c.c. of the lime-water are removed for analysis. It is not possible to collect the whole of the 60 c.c., nor would

there be any advantage in doing so.

It only remains to determine by means of the oxalic solution the amount of lime present in the lime-water before and after the process, and from the loss to calculate the proportion of carbonic acid in the known volume of sample air. The determination of the "causticity" of the lime-water (that is, the amount of caustic lime it contains) is made by a simple process of alkalimetry. Oxalic solution is dropped from a graduated burette into a measured quantity (30 c.c.) of lime-water, until the exact point of neutralisation is reached. Several "indicators" are available for the recognition of this point. If a drop of the lime-water is removed from time to time and put in upon turmeric paper, it will give a brown stain as long as any caustic lime remains unneutralised, but none when the neutralisation is complete. A still simpler indication is obtained by adding to the lime-water a few drops of a solution of "phenol-phthalein," which is decolorised as soon as all the lime is neutralised.

Example.—The capacity of the bottle is 4,840 c.c., and 30 c.c. of lime-water take respectively 38 and 31 c.c. of oxalic

solution before and after the process.

The volume of air is 4,840 c.c. -60 c.c. =4,780 c.c. Since 30 c.c. lost 7 mgr. of lime, the 60 c.c. employed lost 14 mgr., and from data already given this corresponds to $14 \times 0.4 = 5.6$ c.c. of carbonic acid. Hence, the sample air contained (by volume) 5.6 parts of carbonic acid in 4,780 of air, or 1.17 per thousand.

Corrections must be made if the temperature deviates materially from 32° Fahr., or the barometric pressure from 30 inches of mercury. One per cent. should be added to the

result for every 5° above 32° Fahr. If the barometric reading is B, the result must be multiplied by $\frac{30}{B}$. Thus, if in the above example the temperature were 72° Fahr., and the pressure 29.2 inches, the corrections would be as follows:-1.17 x $\frac{30}{100}$ × $\frac{30}{29.2}$ = 1.3 part of carbonic acid per thousand.

2. Haldane's method. - A method has been devised by Haldane for determining CO, without using collecting jars, etc. He adapts a gas analysis apparatus consisting of a burette with a wide ungraduated portion and a narrow portion graduated in divisions of 10000 part of the capacity of the burette. This is enclosed in a water jacket, and a control tube is used to correct for variations of temperature. The pressures under which the readings are made are found to be the same, before and after absorption, by measuring the level of the potash solution in the narrow tube. The graduated burette allows of 25 c.c. of air to be admitted for examination. Having collected the amount of air the CO2 is absorbed in the caustic potash solution. After complete abstraction of CO, the volume of the air is again read off at the same temperature and pressure. The difference between the two readings gives the result in volumes of CO2 per 10,000 of air.

3. Angus Smith's Minimetric method and "Household" test. -A certain amount of carbonic acid is needed in order to produce a cloud in a given volume of lime-water. The volume of lime-water being constant, the volume of air necessary to give a visible precipitate affords a simple inverse measure of the carbonic acid it contains. Half an ounce of clear limewater is placed in each of a series of stoppered bottles of different sizes, and well shaken. The smallest bottle which shows any cloudiness indicates the proportion of carbonic acid present in the sample air, according to the following empirical scale :-

A ppt. in bottle of	a } 443 {	e.c. capacity presence	indicates the	3 0 4 7	parts CO ₂ per 1,000
"	356	"	"	0.5	,,
"	299	HARL PARTY.	1,	0.6	"
11	259	,,	- 11	0.7	"
,,,	228	- "	"	0.8	,,
11	204	The stage of	mar, position	0.9	,,
"	185	33	,,	1.0	12
,,	157	11	"	1.2	"
"	128	,,	"	1.5	5)
"	100	.,	"	2.0	11

Adopting 0.6 per 1,000 as the permissible limit of carbonic acid, there should be no precipitate when half an ounce of limewater is shaken in a bottle of 300 c.c. or $10\frac{1}{2}$ oz. capacity. The method is not very satisfactory or accurate.

Carbonic oxide, CO, may be determined quantitatively by eudiometry, or by exposing a measured volume of air to a solution of cuprous chloride, which absorbs this gas. The loss in volume shows the amount of carbonic oxide present. On passing CO through palladium chloride solution a black precipitate of carbon is formed. Its presence may also be determined by Vogel's test with the spectroscope after using ammonium sulphide. A caged mouse is affected in $1\frac{1}{2}$ minutes by air containing 0.4 per cent. of CO, and this may be used as a sensitive test (Haldane).

Oxygen is determined quantitatively by exposing a measured volume of air to a solution of alkaline potassium pyrogallate, which absorbs the oxygen. The analysis may be conveniently made in a graduated glass tube inverted over a mercurial trough; the tube having been filled with air, the pyrogallate solution is passed up into it through the mercury. All the readings must be taken at the same pressure. Oxygen may also be determined by a Hempel burette and pipette, using phosphorus as the absorbent.

Ozone is detected by its action upon potassium iodide. Strips of porous paper are steeped in a solution containing starch and potassium iodide, and then dried. These are exposed for a definite period to the air, care being taken to exclude sun and rain. Ozone causes a blue tint, the intensity of which is taken as indicating the amount of ozone, according to a standard scale of tints;* the ozone liberates iodine from the potassium iodide, and the iodine strikes a blue

^{*} For several reasons this test is held to be inexact. The chief are (1) that nitrous acid and peroxide of hydrogen, as well as ozone, give the reaction; (2) that the conditions of exposure are not uniform—light, wind, humidity, and temperature all vary, and all affect the reaction; and (3) that some of the liberated iodine is volatilised, and some reacts again upon the potash to form inert iodide and iodate.

colour with the starch. Neutral litmus and pure potassium iodide may also be used. This paper is coloured blue at once by air containing 1/260,000 of its weight of ozone, and slightly blue by from 0002 to '0003 milligrams of ozone. Wurster's reagent (tetramethyl-paraphenylene-diamine) may be similarly

Ammonia may be collected by aspirating a known volume of air through distilled water, and then determined (quantitatively as well as qualitatively) by Nesslerising (page 78).

Nitrogen is determined with sufficient accuracy as the residue left when oxygen is removed by pyrogallate solution. The proportions of other gases are so small that their retention does not materially affect the determination of nitrogen.

Water, or rather aqueous vapour in air, may be measured by the hygrometer, by the spectroscope, or volumetrically by its absorption by sulphuric acid, etc.

It may be necessary occasionally to test for sulphuretted hydrogen, or for mineral acids due to chemical manufacturing processes. The former is detected by exposing strips of porous paper moistened with a solution of lead acetate, and noting the black colour due to formation of lead sulphide. Free mineral acids will redden moist blue litmus paper; and may be obtained for chemical analysis by aspirating the air through distilled water or solution of caustic potash.

the oil to annihilation of the the said

CHAPTER II

METEOROLOGY

The principal climatic phenomena requiring systematic record are: (1) Temperature of the air; (2) Pressure of the air; (3) Movement of the air; (4) Amount or degree of moisture in the air; (5) Ozone; (6) Rainfall; (7) Sunshine; (8) Cloud, mist, fog.

Temperature varies constantly, so that the reading at any one time has only a limited significance, and maximum and minimum registering thermometers are

required for systematic record.

The maximum thermometer has a mercurial column, a detached portion of which serves as an index, being either permanently separated from the rest by a minute bubble of air (Phillips), or else detached afresh each time by means of a constriction near the bulb (Negretti), which allows the expanding mercury to pass, but renders the cohesion of the metal insufficient to draw it back upon cooling. The thermometer is kept horizontal, and the index is shaken back after each reading, ready for a new observation.

The minimum thermometer (Rutherford's) contains coloured alcohol. The index is a little metal rod included in the column, and is drawn back by capillary attraction as the column contracts, but allows the alcohol to flow past it when expanding. To set the instrument, it is only necessary partially to invert it, and let the index fall to the top of the spirit column. It is

then replaced in the horizontal position.

If the instruments are read once daily, say at 9.0 a.m., they indicate respectively the highest and lowest temperatures attained during the previous 24 hours, irrespective of the time of occurrence.

Two sets of instruments are used, one for "shade" temperatures, i.e. the temperature of the air

itself, the other for "radiation" temperatures with

free exposure to the sun.*

Thus are obtained for each day—shade maximum, shade minimum, sun maximum, grass minimum; and from these the daily, weekly, monthly, or yearly ranges, both in shade and with free radiation.

The mean temperature of the day is calculated in

several ways :-

(a) By taking the mean of the maximum and minimum recorded temperatures. This is correct only in winter; at midsummer it is often more than 2° too high.

(b) By taking the mean of two readings at twelve-

hour intervals: e.g. 9.0 a.m. and 9.0 p.m.

(c) By taking a single reading at 9.0 p.m.

(d) By taking hourly readings, or a continuous record, by means of photography. The mean of these

is the true mean temperature of the day.

The weekly, monthly, and annual means are the averages of the daily means. Taking the average of a number of years, the daily minimum occurs at about 4.0 a.m., the maximum at 2.0 p.m. The annual

extremes are in July and January respectively.

Proximity to the sea lessens the range of temperature and moderates its extremes, water having (as compared with land) a high capacity for heat, but being slow to radiate or absorb it. Elevation lowers the temperature, by facilitating radiation. The influence of latitude, of winds and ocean currents, and of aspect and exposure, is obvious. Some soils are "hot"—dry sands and hard rocks, for example—while moist clayey soils are "cold." A surface covered with vegetation radiates much more

The former are placed in a louvred wooden box, four feet above the ground. The "sun maximum" thermometer also is placed four feet above the ground; it has a blackened bulb, which is enclosed in a second glass bulb exhausted of air. The corresponding minimum instrument is supported upon a little wooden tripod, close to the ground.

rapidly than bare earth; at night a grass field is colder than a road. The radiation thermometers are affected principally by the degree of sunshine and the humidity

of the atmosphere.*

Earth temperature, that is the temperature of the soil at fixed depths, is more uniform and less diurnally variable than atmospheric temperature. According to Ballard, when the temperature at 4 feet reaches 56° Fahr., infantile diarrhœa may be expected to become epidemic. Earth temperature, which is measured by Symonds' earth thermometer, varies but little, proportionately, from surface temperature, except that it is much more sluggish. At 24 feet of depth, however, it is almost the reverse of surface temperature, being highest in winter and lowest in summer (Forbes). Below 40 feet there is little annual variation in the temperature of the soil.

Humidity involves rather complex considerations. Water evaporates into dry air, as into a vacuum, but the volume of the resulting moist air is greater than that of the original dry air. For example, a cubic foot of dry air at 60° Fahr. weighs 536·3 grains, and is capable of taking up 5·77 grains of water; the product, however, is not 1 cubic foot of moist air, weighing 542·1 grains, but 1·0176 cubic foot, and a cubic foot of the same moist air weighs only 532·7

^{*}The duration and intensity of sunshine are recorded by Campbell's or Jordan's apparatus. The former consists essentially of a glass sphere, which acts as a lens, and brings the rays to a focus at successive points upon a curved sheet of paper placed at the right distance. A burnt track is thus described for such time as bright sunshine continues. Jordan's sunshine recorder is a small flat circular box. The sunshine is admitted through a slit, and impinges upon sensitised paper, leaving a varying photographic record of its duration and intensity. In either apparatus the papers must be renewed daily, and in the latter the sensitised paper must be "fixed" by immersing it in water.

grains. At 32° Fahr, a cubic foot of dry air can take up only 2·13 grains of water; at 80° Fahr, it can take up 10·98 grains. Further, the capacity for moisture increases with the temperature, but much

more rapidly.

Indeed, it may almost be said that, comparing temperatures which increase in arithmetic progression, the capacity of dry air for moisture at those temperatures increases in geometric progression. Taking 0.0° F. as a starting-point, the capacity for moisture (in other words, the weight of a cubic foot of aqueous vapour) doubles with successive increments of temperature, not departing greatly from an arithmetic series.

Temperature	Weight of a cubic foot
(Fahrenheit).	of vapour.
0°	0.55 grains
16°	1.09 ,,
33°	1.21 ,,
52°	4.39 ,,
73°	8.32 ,,
96°	17.68 ,,

As a rule, the water present is only about 70 or 75 per cent. of the amount required for saturation, but if the air is cooled, the same quantity of aqueous vapour may suffice to saturate it; and if the cooling is carried further some of the water will be precipitated as mist or dew. It will be understood, therefore, that, however dry the air may be, it always contains a certain proportion of water, and if cooled to a certain temperature will be "saturated" by that moisture; this temperature is called the dew-point. "Super-saturated" air deposits its moisture only upon solid surfaces, and this is the cause of mists and fogs.*

^{*} Fog is formed by the condensation of watery vapour on dust particles suspended in the air, which afford "free surfaces" on which condensation can occur. Such atmospheric dust probably consists of fine salt dust from the sea, meteoric dust, condensed gases, combustion dust, and particulate organic and inorganic matter. Aitken holds that the

Humidity is measured by hygrometers of various kinds.

Daniell's, now rarely used, consists of two dependent glass bulbs, connected by a tube bent twice at right angles. One bulb is of black glass, and contains a thermometer, the stem of which is visible in the tube above; the other bulb is covered with linen. Both bulbs contain ether. A little ether is dropped on the linen coating, and by its evaporation lowers the temperature of that bulb, so that ether distils over from the blackened bulb. This causes the temperature of the latter to fall; as soon as it sinks to the dew-point the black surface is dulled by deposit of atmospheric vapour, and this temperature is instantly read off by means of the contained thermometer.

Regnault's is somewhat similar, but the black bulb is replaced by a more sensitive bright silver cup, and the distillation is brought about more conveniently and rapidly by means

of an aspirator.

Dines' consists of a plate of black glass, covering a small chamber containing the bulb of a thermometer. Cold water is passed through this chamber until the dew-point is reached and the glass becomes dull; the current is then stopped, the temperature gradually rises, and the thermometer is read at the

moment when the cloud disappears.

Saussure's, which has the advantage of giving the relative humidity directly, consists of a human hair, fixed at one end and stretched by a light weight attached to the other. It passes round a movable axis which carries a finger pointing to a graduated arc. The hair, which must be uninjured and free from oil, elongates with moisture and contracts as it becomes dry; and in so doing rotates the axis and the indicator. The instrument has to be graduated empirically by exposing it first to perfectly dry air, and then to air saturated with moisture, and making the respective positions of the index the 0 and 100 points upon the graduated scale.

The wet and dry bulb hygrometer is usually employed. Two ordinary thermometers are fixed side by side, one of them having its bulb covered with muslin,

earth's atmosphere is greatly polluted with dust, produced by human agency, rising by heated strata of the air, and forming the "free surface" for condensation, even though the air be not saturated. The higher the humidity and the higher the vapour tension, when the air is not saturated, the greater is the amount of moisture held by the dust particles. Town fogs and smoke fogs may be dry, and are largely constituted of particles of unconsumed carbon.

kept wet with distilled water; a tail of the muslin dips into a vessel of water so as to make good the loss by evaporation. If the air is saturated with moisture, no evaporation takes place, and the two thermometers give the same reading. If it is not saturated, evaporation lowers the temperature of the wet bulb.

From the wet and dry bulb readings the dewpoint can be calculated, and from the dew-point the
weight of water present in the air. The dew-point
is determined by calculation,* or, more conveniently,
by reference to Glaisher's Tables,† in which the results
are given for all ordinary values of both readings.
The dew-point being known, the next step is to find the
relative humidity of the air, which is always expressed as the percentage of total saturation. For this
purpose reference is made once more to hygrometric
tables, giving the weight of a cubic foot of vapour at
each temperature; and the relative humidity =

weight of cubic foot of vapour at the dew-point weight of cubic foot of vapour at dry-bulb temperature × 100,

since the numerator is the weight actually present, and the denominator the weight which is required for complete saturation at the dry-bulb temperature.

In practice none of these calculations are necessary, since Glaisher's Tables give the relative humidity corresponding to all ordinary readings of the wet and dry bulb thermometers. This is by far the most important hygrometric datum, the absolute weight

^{*} By the use of Apjohn's Formula, $f'' = f' - \frac{t-t'}{87}$, where t and t' are the dry and wet bulb readings respectively, f' the elastic tension of vapour corresponding to t' (as ascertained from a table of tensions), and f'' the tension of vapour at the dew-point. Having thus found the value of f'', the temperature corresponding to it in a table of tensions is the dew-point. † Based upon the "Greenwich Factors," or "Glaisher's Factors," a different factor being employed for each temperature. If D is the factor corresponding to the dry bulb temperature, the dew-point is t - D (t - t'),

or tension of vapour present, and even the dew-point, being of minor interest. The relative humidity is an inverse measure of the drying effect of air. The capacity for taking up moisture increases with the temperature, but far more rapidly, so that the mere difference between the dew-point and the air tempera-

ture is no criterion of the humidity.

Aqueous vapour intercepts some of the sun's luminous rays, and very greatly impedes the radiation of non-luminous rays from the surface of the earth into space. Hence the difference between the sunmaximum and grass-minimum temperatures is chiefly dependent upon the humidity. The intensity of radiation at high altitudes is largely due to the absence of aqueous vapour. In Arctic regions the unimpeded solar radiation has been known to boil the pitch in the seams of ships which were surrounded by ice.

Humidity comes into question also in connection with ventilation, the aim being to keep the air of rooms as near as practicable to the accepted optimum, namely, 70% of saturation. Hence in heating by convection it is desirable to moisten the air, and for certain trade processes in which air is steamed limits

of relative humidity are imposed by law,

Evaporation from moist surfaces is affected by the temperature of the water, the temperature of the air, the humidity of the air, and the wind. It is more rapid from moist soil than from water; and deeprooted crops, such as wheat, dry the soil to a greater depth than grass. No satisfactory instrument (atmometer) has been devised for measuring evaporation, owing to the difficulty of excluding rainfall while allowing free exposure. A rough estimate may be obtained by exposing a measured volume of water in an open dish of known area, and deducting from the final measurement the ascertained rainfall. Another plan is to shelter the atmometer from rain by a cover, but this necessarily lessens the evaporation. A third is to use an index evaporation gauge (Negretti). Symons found that at Lea Bridge the mean annual evaporation per square inch of water-surface was 21 inches, the rainfall being 25 inches. In exceptionally

dry years it exceeds the rainfall.

Atmospheric pressure is measured by barometers. Mercurial barometers consist essentially of a glass tube some 36 inches long filled with mercury, and then inverted over a mercurial trough. The tube being vertical, the mercury sinks until the column is of just sufficient height to balance the atmospheric pressure acting upon the free surface of the mercury in the trough. The vertical distance between the two mercurial surfaces exactly measures the pressure of the atmosphere in terms of mercury. The level of the top of the column is read off against a fixed scale, but as the level of the mercury in the cistern also changes slightly with every change of pressure, it is necessary to take this into account.

In Fortin's barometer the cistern has a leather bottom, which can be raised or lowered by means of a fine screw, until the surface just touches a fixed ivory point. After this adjustment of the cistern to a constant standard level, the upper level is read upon a scale marked upon the casing of the tube.

In the Kew barometer the necessity for the adjustment is obviated by graduating the scale in nominal inches, which are shorter than true inches, and exactly correspond to the displacement caused by a change of pressure to the extent of one inch of mercury. A single reading therefore gives the true vertical height of the column.

In the siphon barometer the cistern is dispensed with, and the tube is of **U** shape, one arm being short and open at the end. Both levels are read upon a scale. The movement in either arm is little more than half that occurring in a cistern

barometer.

The ordinary wheel-barometer is a syphon barometer, the movements being transmitted by a string from a float upon the mercury in the open tube to an axis which carries a long finger like that of a clock.

Greater accuracy in the reading is obtained by the

use of the "vernier," a small movable scale which slides upon the principal scale (Fig. 1). If the barometer is graduated to 1 inch divisions the vernier can be so graduated that 25 of its divisions correspond to 24 of those upon the other scale, and each division is therefore only $\frac{24}{25}$ of the length of the standard divisions. With sub-graduation the vernier will show differences of $\frac{1}{25}$ of $\frac{1}{20}$, i.e. $\cdot 002$, or $\frac{1}{500}$ inch. When a reading is to be taken, the lowest mark on the vernier is accurately adjusted against the top of the mercurial column. If this level exactly corresponds with one of the divisions of the principal scale, the vernier correction is not needed; its lowest mark will coincide with the mark on the scale. If, however, the level of the mercury does not exactly coincide with a scale division, the vernier helps us to measure more accurately than is otherwise possible the fractional excess over the next division below. For this purpose we must follow the vernier scale upwards until we come to a mark upon it




Fig. 1.—Barometer Scale and Vernier, showing a reading of 29.85 + 18 × .002 = 29.886 inches.

which corresponds more closely than any other to one on the fixed scale; calling this the xth mark, the correct reading is $.002 \times x$ inch above the scale mark next below the mercury level. For example, the mercury stands above 29.85, but below 29.90, as read on the fixed scale, and upon inspection it is

found that the sixth vernier division corresponds with a division of the fixed scale. This shows that the fractional excess above 29.85 is $6 \times .002$, or .012 inch, and the true reading is therefore 29.862 inches. Had the 20th division of the vernier been the one which coincided with a fixed scale mark, the fraction would have a greater value, $20 \times .002 = .04$ inch, and the barometric reading would be 29.89 inches. A little consideration will make it clear that as compared to scale divisions we lose .002 inch for every vernier division as we follow the marks upwards, until we have lost all the fractional excess over the .29.85 mark; and then the marks on the two scales will correspond.

The reading having been accurately taken with the aid of the vernier, it remains to apply certain corrections:—

- (1) For "index error"—i.e. inaccuracies in the scale.
- (2) For "capacity"—i.e. for the change in the cistern level, unless this is allowed for as already described. Practically, this correction is never called for in good modern barometers.
- (3) For "capillarity," which tends to depress the column slightly.

These three corrections are special to each instrument, and Kew certificates give for each half inch a correction comprising them all.

- (4) For temperature, which affects the mercury, and also the brass or other scale. All readings are reduced to 32° Fahr. by means of a table giving corrections for each temperature.
- (5) For elevation. It is customary to reduce all readings to sea-level—i.e. the mean half-tide level at Liverpool. For this purpose it is necessary to know the exact height of the station and to refer to a table of corrections. Roughly speaking, the correction is about 1 inch for every 1,000 feet.

Owing to the high specific gravity of mercury, 13.5, a column of about 30 inches in height balances the atmospheric pressure, and there is the further advantage that the vapour-tension in the Torricellian vacuum

above the mercurial column is trifling. The density of mercury, however, renders the range of movement under varying atmospheric pressures small, and other liquids have been employed in order to secure greater sensitiveness of indication. Jordan's glycerine barometer is the most important of these; glycerine has a specific gravity of 1.26, and a column of 27 feet would about balance a mercurial column of 30 inches. A fall of 1 inch of mercury is represented by a fall of 10.7 inches of glycerine, and the latter is therefore far more sensitive. Water barometers have also been constructed, a column of 34 feet being required to balance 30 inches of mercury. These instruments are, of course, still more sensitive, and magnify the mercurial reading 13.5 times; but in addition to the disadvantage of their height, the vapour tension in the vacuum is a source of error that varies with the temperature.

Aneroid barometers dispense entirely with liquids, and measure the barometric pressure by means of the elasticity of metal. A small air-tight metallic box is (nearly) exhausted of air, and is so constructed that the top is slightly forced in when the atmospheric pressure rises, and (aided by a strong spring) comes out when the pressure falls. These movements are conveyed by levers, etc., to a finger moving upon a dial; the dial is graduated empirically by comparison

with a standard mercurial barometer.

There are periodic and non-periodic variations of pressure, but the former are so completely masked by the latter in the latitude of England as rarely to be perceptible except in averages. There is a tendency to a daily curve, with a range of about 0.02 inch. There are two maxima, about 9 a.m. and 9 p.m.; and two minima, about 3 a.m. and 3 p.m. This slight curve is only noticeable in the rare absence of non-periodic changes; but in the

Tropics the daily range exceeds 0.11 inch, while in Arctic regions it vanishes. The annual curve in England is somewhat irregular, but has a maximum at the end of May, and a minimum at the end of October. It is very different in other countries.

Rainfall is measured by a rain-gauge, consisting of a copper funnel leading to a bottle or other receiver. The funnel has a sharp, circular rim, usually 5 inches or 8 inches in diameter, and its area in square inches is accurately known. The water collected in the receiver is measured in a graduated glass vessel, the divisions of which correspond to the fractions of an inch of rainfall. It is found that the amount collected is greatest at the ground level. Care must be taken to place the rim of the gauge perfectly level and to select an open site. "No object ought to subtend a greater angle with the horizon than 20° in any direction from the gauge" (Scott). Snow and hail may be melted for measurement by adding a known volume of warm water, which must be deducted from the total.

As regards the daily curve of average rainfall there are indications of three maxima, the principal of which occurs about 2 or 3 p.m. The annual curve varies with locality. In London and on the east coast of England the principal maximum is in October, and the minimum in February or March; but on the west coast January is the wettest month.

The average annual rainfall varies from about 20 inches on the east coast of England to 60 or 80 inches on the west coast of Ireland and Scotland; and at Seathwaite, in Cumberland, it averaged 154 inches during six years. The average for England is about 25 inches.

The rainfall does not often exceed an inch in one day in Great Britain, but enormous quantities are recorded occasionally. At Camberwell, in 1846, 3 inches fell in $2\frac{1}{4}$ hours, and in Monmouthshire, in

1875, 5.36 inches fell in one day.

It is greatest on the westward slope of English mountains, owing to the humid westerly winds depositing moisture as soon as they are forced to ascend into colder strata. Haughton calculates that one gallon of rainfall, by its condensation from vapour into the liquid form, gives out latent heat sufficient to melt 75 lbs. of ice, or 45 lbs. of cast iron; and that on the west coast of Ireland the heat derived from the rainfall is equivalent to half that derived from the sun.

Wind may be measured, by anemometers, as regards either its pressure or its velocity.

Hooke's anemometer is a thin rectangular plate of iron, suspended from its upper edge, and set at right angles to the wind. The force of the wind is measured by the angular displacement of the plate. A pencil connected with the plate records upon a chart moved by clockwork the displacement. In Osler's anemometer the principle is the same, but the weights are replaced by a spring.

Pressure anemometers have the advantage of recording gusts and sudden changes in wind-pressure. So far as velocity and pressure are comparable, it may be said that the pressure varies as the square of the velocity. If v = the velocity in miles per hour, and P = the pressure in pounds per square foot, then

according to James,

 $V^2 = 200 \text{ P}.$

Robinson's is the only one in common use. It consists of four light arms rotating in a horizontal plane around an axis, through which their movements are transmitted to a series of recording dials. Each arm has at its extremity a hemispherical cup, facing horizontally at right angles to the arm. In whatever direction the wind blows, the vanes revolve, the resultant pressure upon the concave surfaces being greater than that upon the convex. It

was formerly believed that the cups moved with one-third the velocity of the wind, and the index dials were graduated accordingly. It appears, however, that the proportion is greater than one-third, and has a direct relation to the size of the cups and the velocity of the wind. Negretti now supplies a recording and an electric anemometer, by which Robinson's apparatus may be made self-recording.

Apart from the uncertainty as to the true value of the readings of each instrument, the records of different stations are not strictly comparable, since inequalities in elevation and in shelter from the wind are inevitable, and must materially affect the results. The position of the anemometer should be such as to secure the fullest possible exposure to wind from all quarters.


The average velocity of wind in Great Britain shows a daily maximum and minimum closely corresponding to those of temperature, about 2 p.m. and 4 a.m. The annual curve of average wind-velocity has a maximum in July and a minimum in December or January.

In Beaufort's scale the force of the wind is stated by arbitrary numbers, ranging from 0 (calm) to 12 (hurricane).

Force.	Beaufort Scale. Miles per ho	Miles per hour.	
0	Calm Up to 3.		
1	Light air 8		
2	Light breeze		
3	Gentle breeze		
4	Moderate breeze 23		
5	Fresh breeze		
6	Strong breeze		
7	Moderate gale 40		
8	Fresh gale		
9	Strong gale		
10 .	Whole cole		
11	Storm		
12	Hurricane		

In England the velocity is, upon an average, about 8 miles per hour, and rarely exceeds 40. Local winds

may be caused by geographical configuration—e.g. upon coasts and mountains. In normal conditions a wind blows at noon from sea to land, from plains to hills, but at sunset the directions are reversed. Winds increase in force with elevation.






Fig. 2.-Wind-rose.

proportions of wind-observations from each point of the compass. The Greenwich wind-rose for 1861-70 was

N	NE	E	SE	S	SW	W	NW
10	12	7	7	8	33	15	7

If the readings of the barometer at any given moment at several stations dotted over a wide area, such as Europe, are recorded upon a map, and lines are drawn connecting together all the points where the same pressure prevails, we obtain a "synoptic chart," and these lines, or **isobars**, are found to assume very commonly certain typical forms (Fig. 3).

1. Cyclones, formed by concentric isobars, the

lowest pressure being at the centre.

2. Anticyclones, the isobars being roughly con-

centric, with the highest pressure at the centre.

3. Secondary cyclones, formed by looped concentric isobars (the circle being incomplete, and thus failing to form a true cyclone) with lowest pressure in the centre.

4. V-shaped depressions, with lowest pressure in the interior, forming, for example, the angular intervals between adjoining anticyclones. 5. Wedges of high pressure—highest in the interior—inserted between two adjacent cyclones, and usually pointing to the north.

6. Cols or necks of comparatively low pressure between two adjacent anticyclones, like the pass or

"col" between two Alpine peaks.

7. Straight isobars.

Cyclones, secondary cyclones, V-depressions, and




Fig. 3.—Distribution of Pressure over the North Atlantic, and parts of the United States and Europe, Feb. 27th, 1865.

(After Abercrombie.)

wedges, usually travel eastward at the rate of about twenty miles per hour, but anticyclones often remain

stationary for days, weeks, or even months.

If the direction of the wind at each station is marked upon the synoptic chart, it will invariably be found that its direction is roughly parallel to the isobars, and that in anticyclonic areas (in the northern hemisphere) it describes a circle in the same way as the hand of a watch, while in cyclonic areas this course is reversed.* The direction of the wind in these and in all other arrangements of isobars conforms to Buys-Ballot's law—that an observer standing with his back to the wind always has lower pressure to his left, and higher pressure to his right.

The closeness of the isobars, or, in other words, the rapidity of change of atmospheric pressure, forms the "barometric gradient," to which the velocity of the

wind is directly proportionate.

Not only the force and velocity of the wind can be inferred from the arrangement of the isobars, but also the kind of weather prevailing at each station. The front part of an advancing cyclone is associated with rain, stratiform clouds, and moist, heavy atmosphere, while in its rear the opposite conditions prevail, namely, sunshine, clear "fresh" air, and cumulus clouds. Anticyclones are less intense, more stationary, and cover a wider area than cyclones. In front of them and at the centre are found sunshine, blue sky, haze, little wind, keen dry air, and free radiation, with great daily range of temperature. The weather changes which accompany secondary cyclones or V-depressions are the same as those of cyclones; wedges, like anticyclones, have fine weather in front and bad weather in their rear.

All these types are liable to break up or merge into new forms at any time, but it is often possible to "forecast" the weather in any given locality by learning from the synoptic charts the direction and velocity of cyclones approaching from the west.

Cyclones are often diverted from their course by meeting a coast line or mountain chain, or even an anticyclone; their course is not necessarily straight,

^{*} The course of the wind is not exactly parallel to the isobars. The wind crosses the isobars obliquely, and may be described as blowing spirally into a cyclone and spirally out of an anticyclone.

nor their velocity by any means uniform. Lastly, their intensity—as measured by the steepness of the barometric gradient and consequent wind-

velocity-may also change.

Cyclones are much more numerous than anticyclones. The centre of depression usually passes to the north of Britain, and hence it is more common for the wind to "veer" (or "go with the sun") than to "back." It is only necessary to imagine a watch (with its hand moving in the reversed direction) passing over a given point upon a map to understand the successive changes in the direction of the wind at that point, during the passage of a

cyclone.

Climate is the condition of a country in relation to certain meteorological elements, viewed as to their effects upon animal or vegetable life. These elements are (a) distance from the equator, or latitude; (b) relative distribution of land and water; (c) altitude; (d) presence of ocean currents; (e) rainfall; and (f) wind. Proximity of mountain ranges and their influence upon the shelter from wind and on the rainfall, soil and its permeability to moisture, and vegetation, also exert an influence on climate. The climate of the British Islands is typically marine, being greatly influenced by its insular position and the Gulf. Stream, which latter causes the "isothermal lines" (i.e. lines of equal temperature) of the North Atlantic Ocean to run from S.W. to N.E. Such a climate is essentially a temperate one, free from extremes of heat and cold, and yet marked by a high degree of atmospheric moisture. Rheumatism, lung affections, and other diseases directly or indirectly due to such climates therefore abound. It is customary to divide. climates into Warm, Temperate, Cold, Marine, and Mountain climates, and in each of these disease bears some relation to the climate. Particularly is this true

of tropical or sub-tropical climates, in which malaria, dysentery, liver abscess, cholera, yellow fever, and a variety of tropical parasitical diseases find favourable conditions. At the same time much can be done by judicious living to enable the individual accustomed to one climate to live healthily in another.

CHAPTER III

WATER

Amount.—It is generally estimated that from 10 to 15 gallons of water per head per day are required for personal and domestic use, 5 to 10 gallons for municipal purposes and a similar quantity for trade processes, but all these items are liable to wide variation. About 20 to 30 gallons per head are supplied in most towns in Great Britain. The average daily amount taken in food is about half a gallon, but at least a pint of this is contained in solid food. Half a gallon more is used in cooking. Parkes suggested as a standard in a middle-class household 6 gallons per head for domestic washing, 5 for ablutions, including a sponge-bath, 6 for water-closets, 4 for general baths, and 3 for unavoidable waste.

derived from the rainfall, which in its turn is due to distillation under the influence of the sun's rays from all humid portions of the earth's surface. Part of the rainfall is again evaporated from the surface upon which it falls, part flows along the surface to form streams and lakes. A third part sinks into the soil, descending through fissures or pores, until it reaches an impervious formation, and it then either finds its way laterally to the surface in the form of springs, or accumulates in the porous strata overlying the impervious layer, where it may be reached from the surface by wells. It has been found that upon sand or gravel surfaces as much as 90 per cent. or more of the rainfall sinks

into the ground, as compared with 40 per cent. on the chalk, and 20 per cent. on limestone, while upon clay scarcely any infiltration occurs. For obvious reasons the proportion is less in hilly districts, which offer greater facilities for surface-flow, and less in summer than in winter owing to excessive evaporation. Experiments upon a gravelly loam indicated a penetration of only 2 per cent. in summer and nearly 100 per cent. in winter.

In a memorandum dated December, 1897, addressed to District Councils, the L.G.B. observes:—

"The Council . . . should by careful inquiry make themselves acquainted with the sources, nature, and quality of the water supplies in all parts of their district, and, in every case in which the result of their inquiries is unsatisfactory, should take all such steps as may be within their powers, with the view of supplementing or improving the supplies. . . . Accurate information should be procured, if not already available, in such matters as the following:—

1. Where water is derived from gathering ground or from springs. Whether drainage from human habitations, farmyards, and the like finds its way directly or indirectly into the reservoir or to any part of the water service; and whether risk of access to the water of human excreta and similar refuse is likely

to arise.

2. Where water is derived from deep wells. Whether surface or other water liable to be contaminated by drains, sewers, cesspools, and the like, reaches, or is liable to reach the wells. The existence and direction of fissures in the strata deserve a special consideration in this respect.

3. Where the water is derived from shallow wells. Whether the wells are so circumstanced that they run risk of contamination by reason of drains, privies, cesspools, or middens, or by deposit of manure—

whether derived from human excreta or not in or on the ground in the neighbourhood of the wells."

Rain-water is soft and well aërated, but its purity and fitness for drinking or even for washing depend upon the purity of the atmosphere through which it falls. Near the coast it often contains traces of chlorides and sulphates. In inland districts there is a marked increase in the sulphuric acid, ammonia, and organic matter, owing to putrefactive processes and the combustion of coal. Near towns rain may become acid in reaction, from excess of sulphurous and sulphuric acids, and is liable to carry down considerable quantities of tarry and carbon aceous matter from the smoky air. It acquires from the air not only oxygen, nitrogen, carbonic acid and ammonia, with a minute amount of nitric acid, but also any organic or other impurities that may be present in the air. Ammonia is found in largest proportion during the early part of a shower. A litre of rain contains about 25 c.c. of gases, namely, 8 c.c. of oxygen, 16.5 c.c. of nitrogen, and 0.5 c.c. of carbonic acid. Dew and snowwater have practically the same characters as rainwater, except that the dissolved gases are in less amount in snow-water. The average of 20 years, less one-third, gives the amount of rain in the driest year, and plus one-third the amount in the wettest year. (Hawksley.)

Springs and wells.—The portion of the rainfall that sinks into the ground becomes heavily charged with carbonic acid from the air in the interstices of the soil, and aided by this and by the increasing pressure it dissolves out lime and various mineral salts from the strata through which it passes. From the superficial strata it also receives organic matter, but the tendency of filtration through the soil is to remove this, or (in the superficial layers, where oxygen

is present) oxidise it with formation of nitrates. The nature and amount of the mineral matters in solution are determined by the soluble constituents of the various strata through which the water has passed, and the organic impurities are regulated by the facilities for pollution in the superficial soil and the completeness of purification by filtration. Hence the water in shallow wells, especially if situated near dwellings or manured lands, is liable to contain much organic matter washed from the soil, with or without partial oxidation into nitrates or nitrites; and also chlorides, which usually accompany organic impurities of sewage origin. In peaty districts the water acquires a brown tint due to vegetable matter.

According to the depth of the strata from which it is derived, the water may be warm, or of the mean temperature of the locality, or varying in temperature according to the season.

River-water is derived partly from springs, but chiefly from that portion of the rainfall which runs off the surface. Hence it contains a variety of mineral salts in solution, though in less amount than in spring or well water.

All streams are inevitably polluted to a certain extent by organic matter. Moorland streams contain peaty matter, sometimes in sufficient quantity to produce diarrhea. Even in rural districts, drainage from manured land and farmsteads finds its way to the river, and in times of heavy rain much impurity of animal and vegetable origin is washed into the watercourses. Such slight degrees of pollution are not regarded as rendering water unfit for drinking purposes, especially after filtration. Some of the organic matter is oxidised by agitation and aëration, assisted by nitrifying organisms, or removed by aquatic plants, and the bacterial

contents of river water may be much reduced by natural processes. For example, it has been shown by experimental investigation that the movement of river water, and its pressure, exert such an influence. Oxidation, dilution, sedimentation, the bactericidal influence of light, and the antagonism of the fauna and flora of a river, also operate in the same way. When, however, the sewage of hamlets, villages, and towns is poured into the river, the pollution becomes more serious, and the self-purifying agencies are soon overpowered. Even if the sewage added at any one point be insufficient to make a perceptible difference in the whole volume of water, it will contribute to it, and the risk of specific pollution, discernible neither by optical nor chemical tests, must always render such a stream a questionable source of supply. Trade effluents—for example, refuse from dyeworks, paper-mills, and bleachworks-cause even more discoloration and turbidity than sewage proper. From these combined causes the rivers in the manufacturing districts become foul, discoloured, and offensive. Matters are made worse by the refuse from slaughterhouses, solid household refuse, and even contents of privy middens being cast into the river. When the pollution reaches such a degree as this, the purification by oxidation is insignificant. Putrefaction goes on, however, and the suspended matters are slowly deposited in the bed of the stream, so that there is still a tendency to self-purification. The foulness of rivers has rendered it necessary for the towns on their banks to impound pure water which formerly passed into the natural watercourses and helped to dilute the impurities cast into them. Lastly, the flow is in many rivers to a great extent intermittent.

Owing to trade exigencies, volumes of water are in many places impounded, and are only passed into the stream during work hours. In seasons of drought this may for a time almost C *

entirely arrest the flow of pure water, so that sewage or other impurities that gain access to the stream remain undiluted. This evil is greatest on Saturdays and Sundays, but occurs daily, and continues until such time in the day as the reserved flow reaches the point in question. When water is abstracted from the head of a stream for the supply of a town, it is customary to provide a "compensation reservoir" in which the storm waters are impounded, and whence they can be delivered to supplement the remaining flow in the river in dry weather. As a means of regulating the flow these reservoirs are very valuable, but as for trade purposes the discharge is often made intermittent, the lower part of the stream may suffer from delay in the delivery of the "compensation" water, in addition to the loss of the water permanently abstracted from the channel. An intermittent flow may have some beneficial influence in flushing the bed of the stream, but where there is great impurity any interruption of flow causes increased nuisance by exposure of a foul foreshore.

Mode of supply.—In thinly populated districts water is obtained from streams, springs, or wells, supple-

mented by storage of rain-water.

The depth to which it is necessary to sink a well depends upon the surface conditions as well as the arrangement of the strata beneath. Even shallow wells may yield good water unless exposed to pollution, as for instance by the proximity of leaking cesspools, ashpits, drains, or even dwellings. Water will usually be found, collected as in a basin, above the first impervious stratum of clay, and as the surface of this subterranean reservoir will vary according to season, the well must be deep enough to reach its level in times of drought. Such a well drains a wide area, and is liable to be polluted by any impurity of the soil within a considerable radius of its mouth, and as the supply is at best subsoil water, due regard must be had to the possibility of pollution at comparatively distant points. It should be lined with brick and puddled, and the brick lining should be brought a foot or two above the ground so as to exclude surface washings. If the soil is polluted, or if the

water found is impure or insufficient in quantity, the boring is carried through the first impervious stratum to other water-bearing strata deeper down. Sometimes the water in the deeper strata is under sufficient pressure to make it rise to the surface and overflow, forming an "artesian well" or artificial spring. The flow in dense strata being slow, each well exhausts a considerable area, and the yield of water cannot be greatly increased by multiple borings. Borings in sandstone or limestone give large and constant supplies,




Fig. 4.—Diagrammatic Section across a Valley. a a, Level of water in deeper permeable strata.

owing to the enormous accumulations of water they contain, but wells in superficial sand or gravel beds, or

chalk, often fail in dry seasons (Fig. 4).

Tube wells ("Abyssinian") are convenient for small or temporary supplies. A jointed iron tube is driven into the ground, section by section, the lowest being pointed and perforated at the end. When water is reached, it enters the tube through the perforations and is pumped up, if the pressure is insufficient to bring it to the surface. Where the water does not rise to within 25 feet of the surface a deep well pump is necessary. "Abyssinian" tube wells are most suitable for gravel, coarse sands, chalk, and other porous water-bearing strata, and may be driven to any depth up to 150 feet. They are not to

be recommended for clays, marls, and fine sands, and,

of course, they cannot penetrate rock.

In the country, and in certain towns where the air is free from smoke, the rain falling upon the roofs is collected and stored in cisterns for domestic use. It may, however, be collected from any other natural or artificial impervious surface. Its softness renders it valuable for washing, especially if the alternative supply is hard, but owing to sooty and other matters acquired from the air and from the roofs themselves, it is seldom obtained in a state of sufficient purity for drinking purposes. It has been estimated that on the average only about two gallons per head per diem could be collected in this way.*

If rain is relied upon for more than an auxiliary supply, the collecting surface and storage capacity must be regulated by the minimum annual rainfall

and the longest dry season.

Collection and storage.—Water is supplied to towns from reservoirs placed at a sufficient height to allow of its distribution by gravitation through open or closed aqueducts, leading to the street mains and through them to the service-pipes. Pumping stations may be requisite in order to gain this elevation, and, if the source of water is of insufficient purity, settling tanks and filter beds may also be required. When the yield is liable to intermission, large storage capacity in the reservoirs becomes necessary. Supplementary service reservoirs, fed by the main reservoirs, are often constructed in elevated parts of the town, partly for convenience of distribution to certain districts, partly in order to supplement the supply during the hours of maximum demand, and so

^{*} The annual yield is found by multiplying the inches of annual rainfall by the square inches of sectional area of the building (not the slant surface of the roof). This gives the cubic inches of water per annum, and multiplied by 0.0036, the number of gallons.

to avoid the necessity of regulating the calibre of the principal aqueduct by the greatest hourly consumption, which is double the average hourly consumption.

If water with a head of H feet flows through L feet of pipe D inches in diameter, the discharge W in cubic feet per minute will be

$$W = 4.72 \frac{\sqrt{H \times D^5}}{\sqrt{L}}$$
. Hence $D = .538 \sqrt[5]{\frac{L \times W^2}{H}}$.

The source of supply may be deep wells, springs, lakes, rivers, smaller streams, or artificial adits from gathering grounds. Some of these sources are immediately dependent upon the rainfall for their continuance, so that in planning a water supply it is necessary to take into account the area of the gathering ground, the minimum annual rainfall, the longest rainless period, the proportion of rainfall available for collection, the storage capacity of the reservoirs, and the amount of water required by the district to be served.* Water of the greatest purity is obtained from deep wells, or from barren uplands, where the risks of animal or vegetable pollution are slight.

Reservoirs are constructed either by excavation or embanking, the simplest plan being to carry an embankment across a valley. A lining of concrete or clay puddle may be needed to render them watertight. The embankments, necessarily of great strength, have a core of clay puddle, and are protected from disintegration by a covering of grass on the outer side

and dressed stone on the inner.

Means are provided of diverting the tributary

^{*} In such cases the reservoirs should be large enough to hold at least 150 days' supply, and more in dry localities. If R= mean annual rainfall, E= evaporation (page 24), and A= acreage of gathering ground, then the mean daily yield in gallons is 62 A (R-E). In the driest year the rainfall will be $\frac{1}{2}$ less, in the wettest $\frac{1}{2}$ more, than the average. (Symons.)

streams along a by-wash when they become foul in flood times. The reservoir has an overflow weir, and can be emptied for cleansing purposes by means of a pipe, controlled by a sluice, leading from its lowest point.

From the reservoirs the water passes to the aqueducts by an outlet pipe, which is bent upwards at its commencement so as to take the purest water, free alike from sediment and floating matters. The aqueduct may be an open channel, but usually consists of iron pipes buried two or three feet in the ground as a protection against frost. The pipe aqueduct may be lined with pitch or other anti-corrosive, and being watertight, its course is not necessarily downward at all points. Means of access are provided at short intervals for cleansing purposes, and if its course is undulating, sluices are needed at the lowest points for scouring out débris, and air vents at the summit.

Distributing conduits and mains convey the water to all parts of the district, and pass beneath the streets. They are similar in construction to the main aqueducts, and have "scouring valves" at all dead ends in order to wash out sediment. They should be kept as far as possible from the sewers and gas mains.* Hydrants are provided at short intervals for use in case of fire.

Service pipes convey water from the street main to the house, and are controlled by means of stop-cocks. Lead piping is generally used, owing to the many bends and joints needed in distributing water within the house. If, as is sometimes the case, the water s of such quality as to act upon lead, other materials must be employed for domestic supplies.

^{*} There is danger of in-suction if a perforation exists in a descending pipe, and especially at a point of constriction, even if the pipe is constantly full.

Constant and intermittent service.—In almost all provincial towns, and in most parts of London, the water-supply is maintained constantly, save in times of exceptional drought. In some districts, however, the supply is only afforded for a certain number of hours in the day; and it becomes necessary for each household to store a sufficient supply for twenty-four hours, in elevated cisterns. The intermittent system, now becoming obsolete, is attended by many risks and disadvantages, and is only defended on the grounds that it incurs somewhat less waste, and that the substitution of constant service entails better fittings. On the other hand the cisterns, which must be large enough to meet the maximum (not the average) daily consumption, are costly, and are liable to become foul; the water stored in them is stagnant, and absorbs impurities from the air; coal-gas, sewer gases, or liquid filth are liable to be drawn into the mains and service-pipes when empty; and, apart from this, the alternate contact with air and water tends to corrode the pipes and promotes the absorption of lead. In the event of fire, a house or district under the intermittent system is at great disadvantage.

A cistern, if any is needed, should be placed at the top of the house, and fed by a supply-pipe controlled by a ball-cock. It must be large enough to contain at least a day's supply and must not directly supply the water-closet. It may be made of lead, lined with pitch or other protective coating, or of galvanised or Barff iron, or of slate slabs set in cement. It should be effectually covered, but ventilated, and provided with an overflow discharging into the open air away from any source of effluvia. Frequent inspection and

cleansing are necessary.

Purification of water may be necessary on account of excessive hardness, excess of saline

constituents, suspended matters, organic matter in solution, or, lastly, on account of liability to specific

pollution.

Distillation is the most universally applicable mode, and even sea-water can be utilised in this way. Distilled water has an unpleasant taste, and is said to be indigestible from its want of aëration. This can be remedied by exposure to the air in finely divided currents, as, for instance, by letting it fall from a sieve, or by charging it with CO₂ under pressure.

Boiling removes temporary hardness and destroys microbes, and therefore specific contagia. Some of the organic matter is carried down with the calcium carbonate. Boiled water tastes flat, and, like distilled water, needs aëration. Lime is thrown down more

completely if sodium carbonate be added.

Chemical processes are for the most part intended to remove excess of lime. Hard water may be softened, and at the same time deprived of all suspended matter, organic or mineral, by adding about six grains of alum per gallon. Calcium sulphate and a bulky precipitate of aluminium hydrate are formed, and carry down suspended matters. If no calcium carbonate is present, calcium chloride and sodium carbonate must be added previously. Perchloride of iron (2½ grains per gallon) acts similarly.

Clark's process.—Hard waters may be softened upon the large scale in reservoirs by adding one ounce of quicklime per 700 gallons, for every degree of temporary hardness per 100,000. The lime and calcium bicarbonate react to form insoluble calcium carbonate, and this carries down with it suspended matters. A similar process may be employed on the domestic scale, using lime or washing-soda, or both.

The Porter-Clark process is a modification of Clark's. Instead of waiting for slow subsidence, which takes ten or twelve hours, the precipitated

calcium carbonate is removed rapidly by filtration

through cloth, under pressure.

Filtration aims at straining off suspended matters, and oxidising dissolved organic substances. Mineral salts in solution, such as sodic chloride or calcic sulphate, are at most only partially removed by filtration. As a rule, however, hardness is lessened, together with the nitrites and ammonia. Nitrates are increased.

For domestic filtration a sand filter capable of removing suspended matters and some of the organic impurity may be constructed like a miniature filterbed, in an ordinary flower-pot, with or without the addition of a layer of animal charcoal (6-12 inches of fine, sharp sand, superimposed on an inch or two of small gravel). Sponge-filters have deservedly fallen into disrepute; they remove suspended matters, but become foul, and harbour organisms. Carbon is the commonest of all filtering media; and animal charcoal is believed to have a far greater purifying effect than vegetable charcoal. Many efficient patterns of filters are made essentially of animal charcoal; others depend upon silicated carbon, manganous carbon, magnetic carbide of iron, or "carferal" (charcoal, iron, and clay). Maignen's "filtre rapide" has no carbon block, but the water is made to pass through a mixture of powdered charcoal and lime (carbo calcis), supported upon an asbestos cloth; like other animal charcoal filters, it seems to have the power of removing lead. The medium can be renewed without difficulty. Bischof's spongy-iron filter, besides removing the organic matter, lessens the hardness, and often reduces nitrates to ammonia.

Ordinary domestic filters do not completely remove microbes, and, indeed, some charcoal filters appear to increase the number of bacteria in water passed through them. This is owing to the fact that charcoal may add nitrogen and phosphates to water, which

are both nutritive substances on which bacteria flourish, and may absorb impurities from the air, and thus favour the growth of organisms in its substance. To remove all bacteria, filters of a different class are required, with excessively fine pores through which water can be forced under pressure, without permitting microbes to pass. Pasteur-Chamberland's, composed of porcelain formed by a mixture of kaolin and other clays, is the best example of this kind (Woodhead, Horrocks); the Berkefeld filter, somewhat similar, is made of infusorial earth. If much organic pabulum be present in the water, or the filter be used intermittently, or the water forced through it, even filters of this kind are not proof against microbes growing through; and periodic sterilisation by steam or boiling is therefore necessary.*

Filtration upon a large scale is necessary when a public water-supply is turbid or impure. The "filterbeds" constructed for this purpose are essentially shallow reservoirs, one or two feet of water, under-drained by perforated or loosely-jointed pipes, to reach which the water passes downwards through successive layers of fine sand, coarse sand, oyster shells, fine gravel, coarse gravel, and pebbles, the total thickness being several feet. Vents are carried from the deeper layers to above the surface of the water, to allow of the escape of the air displaced by the water as it first descends. From time to time the beds are run dry, for the purpose of aëration; and occasionally they are cleansed by scraping off the sediment and a little of the superficial fine sand, fresh sand being added when necessary. The suspended impurities are strained off by the upper

^{*}A filter (other than Bischof's) should be emptied from time to time to promote aëration, and requires frequent cleansing and occasional renewal of the carbon or other medium. Cleansing may be effected by scrubbing the carbon block, if removable, then running an acid solution of potassium permanganate through it, followed by copious washing with very weak hydrochloric acid, and finally several gallons of water.

portion of the filter, and a certain degree of oxidation of organic matter is effected by the aid of nitrifying organisms. Hardness is diminished somewhat, and iron is removed. It has been shown that filtration through sand removes 95 to 99 per cent. of the microbes present in water (80 per cent. of the bacteria removed were found in the upper inch of sand, and 55 per cent. in the upper quarter inch). If the gelatinous film which forms at the surface of the filter is not disturbed, and if the speed of filtration does not exceed 4 inches per hour, practically all the microbes are arrested. Those in the effluent come from the filter itself, and should not exceed 100 per c.c. (Koch.) Frankland holds that the efficiency of sand filtration depends upon (a) the storage capacity of the unfiltered water, for purposes of subsidence by sedimentation; (b) the thickness of fine sand through which the filtration is carried on; (c) rate of filtration; and (d) the renewal of the filter bed. The gelatinous film on the surface of the filter is the vital layer where the process of nitrification and arrest of bacteria goes on, and it is there that the filter exerts its chief effect. It may therefore be said that sand filtration depends upon four processes: first, there is subsidence of the grosser particles of impurity in the settling tank; secondly, there is mechanical obstruction to impurities in the interstices of the filter; thirdly, there is oxidation of organic matter in the pores of the filterbed; and fourthly, micro-organisms in the vital layer on the surface of the filter carry on the process of nitrification. The efficacy of the filtration can only be measured by periodical bacteriological andchemical examinations. The former should include an estimation of the number of bacteria present per c.c.

Other materials may be employed in the construction of filter-beds, such as magnetic carbide of iron covered by a layer of sand to intercept the suspended

impurities. Such a filter-bed must be worked slowly and intermittently, so as to renew the aëration. Spongy iron can be used on the large scale, or "batteries" of Chamberland's filters. At Antwerp "revolving purifiers" are used-cylinders containing small loose pieces of spongy iron, and having many projections from the inner surface. Their long axis, upon which they rotate, is horizontal or inclined, and the surfaces of the iron masses inside are kept constantly bright and clean by mutual friction. Water is passed slowly through the cylinders as they revolve, and is exposed to the action of the constantly fresh iron surfaces. Subsequently sand filtration is used to remove the iron. Metal cylinders containing filtering material (known as "scrubbers"), polarite, sand and polarite mixed, etc., have also been used for filtration of water.

If the source of supply is a river, storm- or floodwater should be excluded. A preliminary purification by subsidence tanks may be beneficial, and floating matters may be excluded by admitting the water

from the river through a submerged sluice.

EXAMINATION OF WATER.

Collection of samples.—The amount required depends upon the mode of analysis to be adopted. Half a gallon, or $2\frac{1}{2}$ litres, will suffice for almost all purposes, and this quantity is held by an ordinary stoppered "Winchester Quart." The bottle is rinsed out with a little hydrochloric acid, and then with water until the rinsings are no longer acid. At the time of collection it is again rinsed with sample water, and then filled up to the neck. The stopper should be tied in.

In order to obtain a fair sample, it is well, in the case of rivers or ponds, to plunge the mouth of the bottle under the surface of the water, at a distance from the margin, so as to exclude scum and débris, care

being taken not to stir up any sediment. Tap water should be allowed to run for some time before sampling, unless it is desired to ascertain the maximum impurity.

Accompanying the sample should be a more or less full statement of its source. For, in addition to a chemical and bacteriological report of a water, there should also be made a careful examination of the gathering ground. The physical configuration of the gathering ground, its subsoil (the occurrence or otherwise of peat), its geology, its rainfall, its relation to the slopes which it drains, the nature of its surface, the course of its feeding streams, and the absence or presence of vegetation and cultivated areas, of roads, of houses, of farms, of human traffic, of cattle and sheep—all these points should be noted, and their influence, direct or indirect, upon the water carefully borne in mind.

When the sample has been duly collected, sealed, and a label affixed bearing the date, time, and conditions of collection and full address, it should be transmitted (preferably in an ice-box) with the least possible delay to the laboratory, in order that the

examination may be made immediately.

Physical characters.—Water should be clear and bright, free from turbidity or smell, and have a fresh agreeable taste, without perceptible saline or other accessory character. The colour, as seen by looking down upon a white surface through a column of the water two feet in depth, should be clear, or slightly green, or blue; a yellow or brown tint raises suspicion of organic pollution.

NATURAL WATERS may contain-

In suspension—Particles of animal, vegetable, and mineral origin.

Microbes, and other living organisms, animal and vegetable.

In solution-Gases.

Mineral salts.

Soluble organic matter of animal and vegetable origin.

Suspended matters.—The water should be allowed to stand for a day in a tall glass, and the sediment then examined under the microscope.* Particles of sand have a sharp angular outline. Fragments of chalk and clay are amorphous, but a drop of acid introduced beneath the cover-glass will dissolve chalk, leaving clay unaffected. Shreds of cotton or linen are easily recognised under the microscope, and so are fragments of leaves, woody fibre, and other vegetable tissues, if not so decomposed as to be apparently structureless. Hair, wool, fragments of insects, and even epithelial scales may be found; brown globular bodies attributed to sewage contamination are sometimes seen in polluted water.

For the most part suspended matters can scarcely be deemed injurious; but some of them, cotton fibres for example, may supply important evidence of pollu-

tion by household waste, or even sewage.

Living organisms.—Rhizopoda, Infusoria, Hydrozoa, Rotifera, Scolecida, Entomostraca, and Insecta are found in water, together with Fungi, Algæ, Diatomaceæ, and many other organisms. For the most part these are believed to be harmless in themselves, though Infusoria (e.g. Paramecium, Vorticella) and Fungi indicate the presence of impurities, and a relatively large number or amount of vegetable or animal forms of life indicate the presence of organic pabulum for their sustenance. Among the parasites that are believed to be conveyed by water are

^{*} One form of sediment tube is shaped like a pipette, the point of which fits accurately into a little glass cup wherein the sediment collects. Finally a long rod is passed down the tube so as to stopper it from the inside at the point. The cup, containing the sediment and about 1 c.c. of water, can then be removed for examination.

certain tapeworms, the Guinea-worm, Dochmius duodenalis, Bilharzia, and leeches. They may be found

in the adult or embryonic form, or as ova.

Bacteria in water.—In all surface waters bacteria abound, and they are often present in small numbers in deep-well or spring waters. Water bacteria include micrococci, spirilla and bacilli, the last being by far the most common group. They multiply in water with great rapidity. If a sample of ordinary water be allowed to stand, it will be found that in 24 hours there has been enormous multiplication, which continues for three or four days, until there may be hundreds of thousands or perhaps millions of bacteria in a cubic centimetre; then, the pabulum and volume of water remaining the same, there will be a diminution, until in, say, two or three months' time there will be in all probability few or none. In a general way it may be said that foul waters rich in putrescible animal matter show a very rapid increase of bacteria; surface waters, such as river water, show a slow and persistent multiplication of organisms; whereas deep-well waters and spring water show comparatively little increase in contained bacteria. The three chief conditions affecting multiplication of bacteria, with few exceptions, are (a) temperature, (b) period of time of standing, and (c) pabulum in the water.

Number of bacteria in water. - Deepwell water generally contains but few bacteria (say 1-10) per c.c. River water varies in its bacterial content according to season (i.e. temperature) and according to degree of pollution of the river (i.e. organic pabulum). Crookes and Dewar found unfiltered Thames water in June, 1904, contained 883 bacteria per c.c. (filtered, 20), but in December, 5,875 per c.c. (filtered, 29). In 1903, a year of exceptionally heavy rainfall, and therefore of much surface pollution of the Thames, the number of bacteria in June, when the average would under ordinary circumstances have been low, was 10,337 per c.c.; in October of the same wet year the number reached 16,671, and in December 27,216. Yet after this water had passed through the filters the number was reduced to 42,66, and 37 bacteria per c.c. respectively. Prausnitz has shown that waters differ in bacterial content, as would be expected, according to the locality in the river at which examination is made. His investigations were made from the Isar before and after it receives the drainage of Munich, with the following results:—

In this country many similar investigations have been made. In 1901 Boyce found in the Severn:

```
Two miles above Shrewsbury 7,000 bacteria per c.c., including 13 B. coli. Opposite Shrewsbury .. 13,000 ,, ,, ,, 46 ,, One mile and a half lower down 23,000 ,, ,, ,, ,, 321 ,, Two miles and a half ,, 19,000 ,, ,, ,, 600 ,, Nine miles ,, 13,000 ,, ,, ,, ,, 48 ,, Sixteen miles ,, 5,000 ,, ,, ,, ,, 36 ,,
```

From these and many similar records it is evident that, in the result, the number of bacteria in river water depends upon a variety of circumstances, amongst which the most important direct conditions are (1) local pollution; (2) natural purification; (3) season and rainfall, and (4) sedimentation and filtration. Behind these direct conditions we also know that time, temperature, light, exposure to air, etc., exert more or less influence.

There is no hard and fast rule as to how many bacteria a potable water should contain, as the kind

of bacteria and the relative abundance of each species are more important data than actual numbers. Koch, however, has suggested 100 bacteria per c.c. as a maximum number for a properly filtered water. Macé proposed a standard of 0-50 per c.c. as "very pure," and 50-500 per c.c. as "good water." Estimations of water based only upon counting "colonies" of bacteria in plate cultures are of little value, but other things being equal and constant, a small number of organisms tends to indicate a small degree of organic pollution and conditions unfavourable to the multiplication of bacteria. Broadly, it is true that a water containing a large degree of organic matter, the pabulum of bacteria, will contain a higher number of bacteria than a water containing a low degree, and this, of course, is the scientific reason for quantitative estimations.

Species of bacteria in water.—There are three chief groups of water bacteria, as follows:—

(i.) Ordinary water bacteria, including the fluorescent bacilli, liquefying and non-liquefying, many of the common chromogenic organisms, and organisms of air and soil. They are of little importance unless present in large numbers, and they possess no

pathogenic power.

(ii.) Sewage bacteria, including B. coli communis and its allies, the Proteus family, B. enteritidis sporogenes of Klein, and certain streptococci and staphylococci. The most important member of this group is B. coli. This is a motile, non-spore-bearing bacillus, possessing a limited number of flagella, capable of fermenting glucose and lactose, of curdling milk with the production of acid, of forming indol in broth culture, reducing neutral red broth with the production of a green fluorescence, producing gas-bubbles in nutrient gelatine, forming a more or less brownish growth on steamed potato, and producing on the surface of

gelatine a dry, translucent growth which does not liquefy the medium. The bacilli, under the microscope, appear as cylindrical rods, showing more or less motility, having a few short flagella, and they do

not stain by the method of Gram.

The important point is the differential diagnosis of B. coli, and the following characters are now commonly relied upon :- 1. The B. coli group is nonsporing and non-liquefying. 2. The members of the group rarely stain by Gram's method. 3. They produce acid and gas with both glucose and lactose. · 4. They produce acid in milk, and they usually also coagulate it. 5. They produce acid and gas in bilesalt-glucose-broth. 6. They grow well at a temperature of 24°C. Other fairly reliable features are motility, a small number of flagella, a fairly typical growth on potato, and more rapid development on all media than the typhoid bacillus. It is also desirable to remember the characteristics of the three groups of bacilli most readily confused with the coli group. These with their characteristics are as follows :-

(a) The Proteus group, the members of which are motile, liquefy gelatine, produce gas in glucose and saccharose but not in lactose, curdle and acidulate milk very slowly, and usually produce indol; (b) the group including B. lactis aërogenes, non-motile bacilli, which do not liquefy gelatine but which curdle and acidulate milk and ferment sugars other than glucose; and (c) the enteritidis group, containing bacilli which are motile, which only ferment glucose, and which do not liquefy gelatine or curdle milk, which is ultimately rendered alkaline. This group includes B. enteritidis of Gaertner, the paracolon and the paratyphoid bacilli.

Streptococci in water usually denote recent and dangerous pollution of water with sewage matter.

They are absent from even large quantities of pure

water and from virgin soils.

(iii.) Pathogenic bacteria. — The two chief types are the typhoid bacillus and the cholera bacillus. As a rule pathogenic germs occur in water scantily and intermittently. They gain access at the source or gathering ground, in the course or from drainage and sewers, or, thirdly, in the home of the consumer.

Bacteriological Examination of Water.— Strict cleanliness must be observed in collecting samples for bacterioscopic examination. Only sterilised vessels or flasks must be used (by heating to 150° C. for 2-3 hours), and great care must be taken in manipulation at time of collecting the sample in order to avoid the smallest degree of external pollution. The object of bacteriological examination of water is fourfold. (a) The average number of microbes per c.c. on a medium incubated at room temperature (18-22° C.); (b) search for B. coli and its identification in 0.1, 1.0, and 10 c.c.; (c) enumeration of bacteria capable of growing at blood-heat (36-38° C.); (d) search for and enumeration of special microbes. Small quantities of the water to be examined (0.1, 0.2 or 0.3 c.c.) are added to tubes of sterile liquefied gelatine, which are then poured into sterilised Petri dishes. The gelatine solidifies, and the plates are incubated at 20°-22° C. and examined day by day. Soon after the lapse of 24 hours, colonies, arising from original individual bacteria, appear. These are examined by naked eye and microscope, and subcultures made in test-tubes containing suitable media - gelatine, broth, milk, agar, bile and sugar media, etc. Thus the number of organisms per c.c. can be estimated, and the species isolated and separately studied in "pure culture" under favourable conditions. Culture plates should also be made for incubation at blood heat. Slide preparations, stained or unstained, should be made for microscopical examination.

Some bacteriologists prefer to take a sterilised Berkefeld filter and pump or aspirate through it 1,000-2,000 c c. of the water under examination, and with a sterilised brush transfer the particulate matter which has collected on the candle of the filter into 10 c.c. of water or broth. This is now a concentration or emulsion of the organismal content of the water, and may be plated out as above or examined for special organisms (B. enteritidis sporogenes, B. coli, B. typhosus, strep-

tococci, or sewage organisms).

Other bacteriologists, aiming mainly at the detection of organisms of intestinal type, examine for (a) average number of microbes per c.c. on gelatin at 20-22° C. on third day; (b) B. enteritidis sporogenes in 0·1, 1·0 and 10 c.c. respectively; and (c) B. coli in 0·01, 0·1, 1·0, 10, and 100 c.c. respectively. Dr. Houston, Director of Water Examination, Metropolitan Water Board, adopts the above method and tests for B. coli as follows:—Inoculate bile-salt-glucose peptone with 0·1-100 c.c. of water; after forty-eight hours' incubation at blood heat the presence of gas necessitates secondary cultures to determine (a) fermentation of glucose or lactose or both; (b) the production or not of acid; (c) indol in peptone water; and (d) negative result in saccharose (cane sugar).

Gases in water. A litre of water can dissolve 25 c.c. of oxygen, 46 c.c. of nitrogen, and 1,000 c.c. of carbonic acid at ordinary temperature and pressure. These gases are usually present in natural waters, but the proportions are very variable, being largely dependent upon the source of the water, the degree of pollution and of exposure to air, and also upon vegetable growth; chalk and limestone waters contain excess of carbonic acid. The gases in a pure water often contain 30 per cent. of their volume of oxygen. They may be extracted for the purpose of quantitative analysis by means of a Sprengel pump, or by boiling the water for an hour, and collected over a mercurial trough. Potash will then absorb the carbonic acid, and potassium pyrogallate or sodium hyposulphite will remove the oxygen, leaving a residue of nitrogen alone.

For the determination of oxygen alone, Thresh adds measured quantities of sulphuric acid, sodic nitrite, and potassic iodide, to a measured volume of water (about 250 c.c.), in a vessel which must be full to the stopper, so as to exclude atmospheric oxygen. The nitrous acid acts as a "carrier" until all the dissolved oxygen is exhausted, liberating a proportionate amount of iodine. The water is then run into a

bottle, through which a current of coal gas is being passed, and standard hyposulphite is added from a burette until the yellow colour due to iodine disappears, the final point being accurately determined by adding a little starch, which gives a blue colour as long as any trace of free iodine remains. The hyposulphite solution is of such strength (7.75 grammes per litre) that 1 c.c. corresponds to 4 mgr. of oxygen. A correction is required for the oxygen contained in the reagents used, a blank test being made for the purpose. The amount of dissolved oxygen per litre of water is then readily calculated.

Sulphuretted hydrogen or marsh gas may be found, the former due to mineral sulphides such as pyrites, or to decomposition of sulphates by organic matter, the latter to fermentation of vegetable matter in stagnant pools, or to pollution by coal-gas. Traces of ammonia

are usually present.

Well aërated waters are bright, lustrous, and agreeable to the palate. This is especially the case with water from wells or springs, of medium depth, in calcium carbonate formations—the low temperature and greater pressure combining to increase the charge of carbonic acid. The absence of dissolved gases in distilled water renders it flat, dull, and unpleasant to taste.

Mineral salts are derived from the strata with which the water has been in contact. They may include chlorides, sulphates, carbonates, silicates, nitrates, nitrites, and phosphates, the bases being lime, magnesia, soda, potash, alumina, or iron, and more

rarely lead, zinc, copper, manganese, or arsenic.

The proportion of salts present ranges from practically nil in rain-water, or pure moorland water, to 300, or more, parts per 100,000 in "mineral" waters or brackish water from the coast. In water for drinking it should not exceed 60 parts per 100,000, and a good water will not contain more than about 40 parts. An excess of mineral salts in general leads to digestive disturbances, in the form of dyspepsia, diarrhea, or constipation; but this effect is largely dependent upon

the nature of the salts present.

Chlorides, estimated as chlorine, may reach the proportion of 10 parts per 100,000 in organically pure water from the New Red Sandstone, Greensand, and other saliferous formations. Brackish water and sea water contain excess of chlorine. In general, however, good waters contain not more than about 2 or 3 parts per 100,000, and rain-water and peaty waters little or none. Water from surfaces, shallow wells, or other sources open to contamination should be regarded with suspicion if it contains more than 10 parts of chlorine, especially if there is also excess of organic matter or oxidised nitrogen, all these being significant of sewage or animal pollution.

Tests.—I.—Qualitative.—A solution of silver nitrate gives a turbidity or a precipitate according to the amount of

chlorine present.

II.—Quantitative.—A standard solution of silver nitrate is employed, containing 4.788 grms. per litre, so that each c.c. corresponds to 1 mgr. of chlorine, according to the equation $AgNO_3 + NaCl = NaNO_3 + AgCl$. A hundred c.c. of sample water are taken in a white porcelain dish, a few drops of a solution of potassium chromate are added as an "indicator," and the silver solution is then dropped in from a burette, with constant stirring, until a faint permanent red tinge appears, showing that the chlorine is exhausted, and silver chromate is beginning to be formed in place of chloride. If the 100 c.c. of sample water take x c.c. of silver solution before the red tint appears, there are x mgr. of chlorine in the 100 c.c., and, therefore, $x \times 10$ mgrs. per litre.*

Sulphates are usually estimated as sulphuric anhydride, SO₃. The most important are calcium and magnesium sulphate, which are liable to cause dyspepsia and diarrhœa in persons unaccustomed to the use

^{*1} mgr. per litre = 0.1 part per 100,000, and 1 grain per gallon = 1 part per 70,000. Hence x grains per gallon = $\frac{190}{x}$ mgrs. per litre, and y mgrs. per litre = $\frac{7}{100}$ y grs. per gallon.

of water containing these salts. Hence limestone and dolomite waters, which may hold 5 to 20 parts per 100,000 of calcium sulphate, are less wholesome than chalk water, which contains none. Dolomite water differs from limestone water in containing magnesium sulphate, but less calcium sulphate. Some waters contain sodium sulphate. The acidity of certain peaty moorland waters that act upon lead has been attributed to the formation of acid sulphates by the oxidation of sulphides (pyrites) or of sulphurous acid absorbed from smoky air.

Tests.—I.—Qualitative.—A solution of barium chloride, with a few drops of hydrochloric acid, will give a white

precipitate.

II.—Quantitative.—200 c.c. of sample water are boiled, treated with a little hydrochloric acid, and slight excess of barium chloride solution, and then boiled again and filtered. The precipitate collected on the filter is washed, ignited, and weighed as barium sulphate, BaSO₄. Given the weight of BaSO₄ from 200 c.c., the corresponding proportion of SO₃ per litre will be readily calculated.

Carbonates. - Soluble calcium bicarbonate, CaH₂(CO₃)₂, is found in almost all well and springwaters. It occurs in excess in chalk, limestone, and dolomite water, but it is wanting in rain and peaty water. It is precipitated by boiling, in the form of normal calcium carbonate, CaCO₃. Magnesium bicarbonate, MgH₂(CO₃)₂, found in dolomite water, behaves similarly, but partially redissolves on cooling. Sodium carbonate is occasionally found in natural waters, and may give them an alkaline reaction. Free carbonic acid exists in all waters, and if in excess renders them sparkling and palatable. Apart from this, the importance of the carbonates in water depends entirely upon the bases they contain. The determination of temporary hardness will show the amount of calcium and magnesium carbonates.

Silicates of soda or alumina are occasionally found in water.

Test.—Half a litre of sample water is evaporated to dryness; the residue is acted upon by strong hydrochloric acid, and washed with boiling distilled water; then it is once more dried, ignited, treated with acid, and washed with boiling water. The final residue is silica, and may be weighed as such after drying.

Nitrates and nitrites, like chlorides, are harmless in themselves, but if found in a water exposed to risk of pollution may suffice to condemn it for use. Nitrates may be found in pure water from deep wells in the chalk, but as a rule they are due to oxidation of nitrogenous organic matter of animal origin. Even if accompanied by only a small proportion of organic matter, nitrates in water from a source open to suspicion must be regarded as oxidised filth, which may at any time be followed by unoxidised filth.

Frankland emphasised this view by recording all the inorganic nitrogen present, whether in the form of nitrates or nitrites, or ammonia, as due to previous sewage contamination, irrespective of its probable origin, recent or remote. He makes a deduction, however, of 0.032 part per 100,000, as the average amount of nitrogen in the rain-water which must have been the ultimate source of supply. Average London sewage is estimated to contain about 10 parts of nitrogen in 100,000 of sewage, so that 1 part of nitrogen represents "previous sewage contamination" to the extent of 10,000 parts.

An example will make the mode of calculation clear. A sample of water is found to contain 3·1322 parts of oxidised nitrogen (i.e. nitrogen as nitrites and nitrates) per 100,000 parts of the water; and also 0·0034 part of ammonia, which is obviously equivalent to 0·0028 part nitrogen. The "inorganic nitrogen" is therefore 3·1322 + 0·0028, or 3·1350 parts per 100,000. From this must be deducted 0·032, as a correction

for the nitrogen in average rain-water, leaving 3·1030 inorganic nitrogen, and this multiplied by 10,000 gives 31,030 (per 100,000) of "previous sewage contamination." In other words, if 31,030 parts of average London sewage had been diluted with 68,970 parts of average rain-water, and all the nitrogen then converted into nitrates, nitrites, or ammonia, the result would be a water of the composition of the sample, so far as the proportion of inorganic nitrogen is concerned.

The process of nitrification, by which nitrogenous organic matter in water or in the soil is oxidised, with formation of nitrates, is due in great measure to the action of microbes. Under other conditions, which are not clearly understood, nitrates may be reduced by microbes or by contact with oxidisable matter, yielding nitrogen or even ammonia.

Nitrites, as a rule, indicate more recent and therefore more dangerous pollution than can be inferred from the presence of nitrates. They readily

pass into nitrates.

A variety of standard terms has been proposed for the quantitative statements of nitrates and nitrites. The expression "oxidised nitrogen" is adopted here, and is precisely equivalent to "nitrogen as nitrates and nitrites" and to "nitric nitrogen." "Previous sewage contamination" has already been explained. Wanklyn gives his results in terms of "nitric acid," HNO₃.

Water from springs and deep wells contains upon an average about 5 parts of oxidised nitrogen per

million, shallow-well waters from 2 to 200 parts.

Tests for Oxidised Nitrogen.*

I. QUALITATIVE.

a. Diphenylamine test.—A 2 per cent. solution of diphenylamine in strong sulphuric acid is spread out in a thin layer upon a white porcelain plate. A drop of the sample water is allowed to fall in the centre, and any trace of oxidised nitrogen is revealed in a few seconds by a blue tint throughout, or at the line of contact.

^{*} That is, for nitrates or nitrites indifferently.

- β. Horsley's test.—2 c.c. of pure sulphuric acid are added to 1 c.c. of the water, and then a drop of pyrogallic acid solution. A pink or blue tint appears, changing to brown; it disappears momentarily upon shaking, but soon returns.
- γ. Brucine test.—A drop of pure sulphuric acid and a crystal of brucine added to the dry residue of 2 c.c. (or more) of the water will give a pink or yellow tint even with minute traces of oxidised nitrogen.

II.—QUANTITATIVE.

- δ. Aluminium process.—2 grammes of sodium are dissolved in 100 c.c. distilled water so as to obtain a pure solution of caustic soda free from nitrates; 100 c.c. of sample water are added and a piece of aluminium foil. In the course of a few hours all the oxidised nitrogen is converted into ammonia, which is distilled off and Nesslerised. Each mgr. of ammonia, after deducting the free ammonia originally present in the sample, corresponds to 0.82 mgr. of oxidised nitrogen in the 100 c.c. of sample.
- e Zinc-copper process.—Zinc foil is coated with copper by exposure to a solution of copper sulphate until it becomes black. It is then put into a bottle with 250 c.c. of the sample water and 0.5 grm. oxalic acid, and left for twenty-four hours. As in the aluminium process, all the oxidised nitrogen is converted into ammonia and measured as such by Nesslerising.

All the preceding tests are for nitrates and nitrites alike. The following apply to nitrites only:—

- 1. Ilosvay's solution, containing sulphanilic and acetic acids and napthylamine, is an extremely delicate test. A few drops added to water give a rose tint if any trace of nitrate is present.
- 2. One drop each of phenol and sulphanilic acid solution, and then ammonia, will give a yellow colour.
- 3. The addition of starch solution, potassium iodide solution, and dilute sulphuric acid causes a blue tint, due to liberation of iodine by nitrous acid and consequent formation of iodide of starch.
- 4. Meta-phenylene-diamine solution with dilute sulphuric acid (1 c.c. of each to 100 c.c. of water) will give an orange-red colour after standing a few minutes. (*Griess.*)

Phosphates are not of common occurrence except in minute traces. They are, for the most part, indicative of remote animal pollution, but their presence need not condemn a water if other indications are satisfactory.

Test.—The water is concentrated to $\frac{1}{50}$ th, 250 c.c. being boiled down to 5 c.c. Nitric acid is added, and then solution of ammonium molybdate. If phosphates are present, a yellow colour will appear, and, on standing, a precipitate of ammonium phospho-molybdate.

Sulphides are said to result from the reduction of sulphates by *Beggiatoa*, and perhaps from contact with iron pyrites. Carbonic acid liberates sulphuretted hydrogen from sulphides. They are readily detected by smell, and by the following:—

Tests.—A solution of lead acetate (or, still better, a solution of lead hydrate in caustic soda) will give a black precipitate in presence of sulphides.

A solution of sodium nitro-prusside will give a purple colour with sulphides, but not with free sulphuretted hydrogen. It

is therefore safer first to add caustic alkali.

Lime is the most important of the mineral constituents of drinking water. It occurs mainly as bicarbonate and sulphate; a good water should not, as a rule, contain more than 20 parts per 100,000 of the former, or 5 of the latter. In excess it causes constipation when used for drinking, and is wasteful for washing purposes, since it renders the water extremely hard unless previously boiled. On the other hand, absence of lime salts is held to be disadvantageous, because lime is required in the system, and also for the more practical reason that soft waters are, as a rule, less palatable than those of moderate hardness, and often dissolve lead from household service pipes. Horses are said to become rough in the coat if supplied with very hard water. Calcium sulphate is injurious (as already stated), and

renders water permanently hard. Calcium nitrate and butyrate have been found in waters that have caused diarrhœa and other symptoms of intestinal irritation.

Test.—Ammonium oxalate solution will give a turbidity or a precipitate according to the amount of lime added. This precipitate from a measured quantity of sample may be collected on a filter, washed, dried, ignited, and then weighed as calcium oxide.

Magnesia occurs chiefly as carbonate and sulphate, in water derived from magnesian limestone. It renders water hard, and although the carbonate is thrown down by boiling, it re-dissolves on standing. Magnesium salts cause dyspepsia and diarrhea.

Tests.—All the lime is first removed by adding powdered

oxalate of ammonia, and filtering.

(a) The filtrate is tested for hardness in the usual way. Any hardness beyond that of pure water is due to magnesia, and each "degree" corresponds to 0.56 grain of magnesium

carbonate per gallon.

(b) To the filtrate are added sodium phosphate, ammonium chloride, and excess of pure ammonia. In twenty-four hours the precipitate of triple phosphate, NH₄MgPO₄, is collected, washed on a filter with strong ammonia,* ignited, and weighed as magnesium pyrophosphate (MgO)₂P₂O₅.

Soda salts are almost omnipresent, and few waters are free from sodium chloride in small amount. The sulphate occurs in excess in some mineral waters, and the carbonate also is common, sometimes in sufficient

quantity to render the water alkaline.

to which it gives a characteristic inky (chalybeate) taste. It is also frequently derived from iron tanks or pipes through which the water has passed. Water is unsuitable for drinking if it contains more than one grain of iron per gallon—that is, 1.5 parts per 100,000.

^{*} Triple phosphate is soluble in water. It is necessary to allow 1 mgr. of magnesium phosphate as lost for every 50 c.c. of wash.

Ferruginous waters are said to be liable to cause dyspepsia, headache, and malaise in those unaccustomed to their use. Iron adds to the hardness of water.

Tests.—I. Qualitative.—Ammonium sulphide gives a dark colour (lost on adding acid), potassium ferrocyanide a blue, and potassium sulphocyanide (after acidulating with nitric acid) a red tint.

II. Quantitative.—A standard solution containing 1 mgr. of iron in 1 c.c. is made by dissolving 4.96 grm. of crystallised ferrous sulphate in a litre of water; if greater precision is desired, the solution may be diluted to $\frac{1}{10}$, so that 1 c.c. = $\frac{1}{10}$ mgr. of iron. 50 c.c of sample water and 50 c.c. of distilled water are placed in Nessler glasses (page 78) side by side, and to both a trace of ammonium sulphide is added. The black tint produced in the former is exactly matched in the latter by adding successive c.c. of iron solution until the right shade is reached. The amount added (to the 50 c.c.) being known, the proportion per litre (i.e., per 1,000 c.c.) will be just twentyfold, and will give, of course, the milligrammes per litre, or parts per million, of iron in the sample water.

Lead, rarely present in natural waters, is sometimes absorbed from leaden pipes or cisterns. The action is probably twofold. First, the metal is acted upon by the oxygen dissolved in the water, aided possibly by oxidised nitrogen serving as a "carrier" of oxygen. Next, the lead oxide thus formed may be dissolved, more readily, of course, if the water be acid, or may be washed away by the current in a state of fine powder. If, however, the water contains calcium carbonate there is formed on the surface of the lead a protective coating, insoluble and not readily dislodged. Hence it is found that plumbo-solvent action is comparatively rarely associated with well or spring or river waters, but quite commonly with rain-water or soft moorland waters, especially if they have an acid reaction. Moorland gathering grounds are usually rich in peat which has an acid reaction, and thus water draining from it is acid and

affects lead. Nitrates, chlorides, or excess of carbonic acid have been found to increase the corrosion, while silicates and many other substances exert some protective action.

The acid reaction and consequent intense plumbosolvent effect of certain moorland waters has been attributed to acid sulphates (due to the oxidation of iron pyrites, $FeS_2 + \hat{H}_2O + \hat{7}O = FeSO_4 + H_2SO_4$, to sulphurous or other acid in rain, or to the indefinite acid compounds found in soil, and sometimes termed humic and ulmic acid. In some cases it is caused by excess of carbonic acid; but in the main the acidity is that of peat, due, according to Houston, to acid producing bacteria in the peat. He has shown that a mixture of micro-organisms obtained from peat, when introduced into a sterile decoction made solely from peat, not uncommonly renders the peat decoction acid and plumbo-solvent in character. Further, Houston has isolated two non-motile, non-liquefying bacteria from peat soil which possess separately the properties as regards acidity and plumbo-solvency observed in the mixture of peat microbes. Acidity in water is more marked in the first flow when rain follows drought, owing no doubt to the washing out of the acid accumulated in the peat; and least marked in dry weather, when the contribution from springs is relatively largest. Where peat is abundant, and springs are relatively few and small, acidity and plumbosolvent ability may be anticipated. But in the absence of carbonate of lime, the power of acquiring lead is shared in considerable degree by many neutral and even slightly alkaline moorland waters, including those in which the acid peat washings have been neutralised by admixture with spring water. Tidy and Odling attributed the lead-dissolving power to the absence of silica, and brought forward evidence to show that this inverse relation holds good in a number of

water supplies. There are, however, many exceptions. Even distilled water, or solution of sodium carbonate in distilled water, will attack lead.

In 1887 Sinclair White called attention to the prevalence of plumbism among the consumers of the high-level water at Sheffield, and disastrous consequences have resulted in other Yorkshire towns from the distribution of plumbo-solvent water through lead

pipes.

Various remedial measures have been tried. One plan is to expose the water to fragments of limestone, and conduits have been lined with the same material, but it soon becomes inert, owing to incrustation, and frequent renewal of the surface is necessary. Measured quantities of lime have also been added (2 to 3 grs. to the gallon). A more effective method is to add powdered chalk or whitening $(1\frac{1}{2})$ grs. to the gallon), while in some instances carbonate of soda is used (500 gallons of soda solution, 1 lb. to the gallon, are added to one million gallons of acid water). In some towns reliance is placed upon filtration through calcareous sand, which, however, becomes exhausted in time and needs renewal. In pursuance of Tidy's hypothesis, the water has been charged with traces of silica by filtration through powdered granite or flint. It is at least possible that the mischief is not always due to the same cause, and that no one line of treatment is suitable in all cases. Houston recommends ordinary sand-filtration, with the addition of some neutralising material to the filter (e.g., a thin coating of lime on the surface of the sand with limestone underneath the sand) and the subsequent addition of a trace of sodium carbonate to the neutral filtered water.

Many waters, fortunately, are sufficiently protected by the salts they contain. Calcium sulphate is less protective than the carbonate, and magnesium

less than calcium salts. A crust is formed upon the interior of lead pipes, composed of carbonate and sulphate of lead, lime, and magnesia (or such of them as are present in the water), and chloride of lead. Carbonate of lead is difficult of solution except in water charged with excess of carbonic acid under pressure. Pipes which are new, which are bent against the grain, or which are alternately full and empty, are more attacked than others. The maximum amount of lead is found in water that has stood for some time in the pipes, over-night for example. Lead is now chemically purer than was usual in former years, owing to the profitable removal of silver and other metals with which it is found associated, and it would appear that the purer lead is more readily acted upon by water.

Many substitutes for ordinary lead pipes are now available, and where there is any tendency to solvent action one or other of these should be adopted. Compound pipes have long been in use, the interior of which is block-tin and the exterior lead; the two metals are solidly united throughout, and the pipe can be bent without fracturing the tin. A less reliable protective lining to lead pipes may be found in tar or bitumen. Iron pipes are more satisfactory in every respect, except their rigidity, which renders them difficult to adapt to the bends and turns that are needed in houses. Plain iron pipes may be employed, with some risk of trouble from rusting. Tin-lined, glass-lined, or galvanised iron pipes may be used, and among other protective coatings may be mentioned Angus Smith's varnish and collodion. Theoretically, the rustless Barff iron offers the greatest advantages for this purpose; iron pipes of ordinary construction are exposed while hot to superheated steam, and thereby acquire a thin coating of magnetic oxide, Fe₃O₄, which renders them free from liability to rust as long as the coating is unbroken. Unfortunately the film is easily dislodged in the process of fitting. If it is not possible to change the supply of water, or to prevent it by chemical means from acting upon lead, or to keep it from coming into contact with lead, recourse must be had to palliative measures. The water must always be run off for a few minutes in the morning, and again before drawing it for drinking purposes. All that is used for drinking or for cooking should be filtered through some medium containing animal charcoal. Such filters possess the power of removing lead from solution, owing probably to the phosphates which they contain forming an insoluble lead salt.

No water should be used for drinking that contains any trace of lead, for however minute, its presence indicates danger; moreover, the sample may not represent either the seasonal or daily maximum.

Among the symptoms caused by lead-poisoning are anemia, depression, dyspepsia, constipation, colic, and paralysis of the extensor muscles of the hand, leading to wrist-drop. A characteristic blue line appears along the edges of the gums, due to the formation of sulphide of lead in the tissue. Optic neuritis may be found. It has been observed that abortion is common in districts where lead-poisoning prevails, and sometimes to such a degree as materially to lower the birth-rate. The relation between lead-poisoning and gout and kidney disease is referred to elsewhere.

Tests.*—I. Qualitative.—(a) H₂S gives a brown tint if a trace of lead is present, and a black precipitate if much; neither is affected by acid. (b) A solution of potassium bichromate (1 grain per oz.) dropped into a tall glass of water, falls slowly, causing an opaque yellow cloud if there is any lead in solution.

II. Quantitative. - A standard solution containing 0.1 mgr.

^{*} The sample should be the first pint, or less, drawn from the pipe after standing for an hour or other stated period if it is to measure the maximum impurity.

of lead in 1 c.c. is made by dissolving 0.1831 grm. of crystallised lead acetate in a litre of distilled water. The procedure is the same as in testing for iron, but the water should be made acid.

Acidity of water and plumbo-solvency should also be tested for. The former may be estimated by using lacmoid as an indicator. The solution is best made by dissolving 0.2 grms. of lacmoid in 100 c.c. of dilute alcohol (10 per cent.). A few drops added to 10 c.c. of the sample of water at once reveals the presence of the smallest trace of acid. The amount of acid may be determined by titrating a definite quantity of the water with a decinormal solution of sodium carbonate (5.3 grms. Na₂CO₃ per litre).

Plumbo-solvency may be determined in the following way: 50 c.c. (by volume) of washed lead shot are placed in a burette $(\frac{3}{4}" \times 24")$, occupying 18 in. of its length; the sample water is allowed to filter through this lead, and is then collected in successive quantities of 50 c.c., and tested with sulphuretted hydrogen or with ammonium sulphide for the presence of

lead.

Zinc, copper, and arsenic, although rarely occurring in natural waters, may occasionally gain access by means of trade pollution, or by the solvent action of water (especially water containing organic matter, nitrites, and nitrates) upon the pipes and vessels with which it comes into contact. Any trace of these substances renders a water unfit for drinking purposes.

Tests.—Copper, like lead, strikes a black colour with ammonium sulphide, not discharged by acid. It may be determined quantitatively by a colorimetric process similar to those for iron and lead, the standard solution containing 1 mgr. of copper in each c.c. being made by dissolving 3.93 grms. of crystallised copper sulphate in a litre of distilled water.

Arsenic would be detected in the residue after evaporation,

by the Marsh-Berzelius or other test.

Hardness of water is caused by the presence of salts that decompose soap, preventing lathering until sufficient soap is added to exhaust them. The chief of these are lime salts, magnesium salts, and carbonic acid, but iron and alumina have a similar effect. By

boiling the water, the free carbonic acid is driven off, and calcium bicarbonate is reduced to the insoluble carbonate and precipitated; magnesium bicarbonate is precipitated as carbonate in like manner, though it partially redissolves on cooling, and most of the iron is also thrown down. After removal of this temporary hardness, there remain in solution calcium and magnesium sulphates and chlorides, and a little iron-assuming them to have been originally present. These, with the re-dissolved magnesium carbonate, constitute the permanent hardness. Soap consists of alkaline oleates stearates, and palmitates, which form a viscid solution in water. The phenomena of hardness depend upon the formation of insoluble calcium and magnesium oleates, etc.

Hardness is measured by a standard soap-solution, 1 c.c. of which exactly precipitates I mgr. of calcium carbonate. It is found that 1 c.c. suffices to give a lather when shaken with 70 c.c. of distilled water, and anything beyond this is due to the hardness, which is stated in terms of calcium carbonate.

70 c.c. of the water is taken in a bottle of about 250 c.c. capacity, and the soap-solution is added by degrees from a burette, with vigorous shaking after each addition, until a lather completely covers the surface and remains intact for five The number of c.c. added, deducting 1 c.c. for lathering, will give the hardness in mgrs. of calcium carbonate per 70 c.c., that is, in grains per gallon.* The permanent hardness is measured in exactly the same way, in 70 c.c. of the water, after boiling for half an hour and adding distilled water to make good the loss by evaporation. If more than 16 c.c. of soap-solution are consumed by 70 c.c. of water, it is better to dilute it by adding 70 c.c. of distilled water, in which event a deduction of 2 c.c. must be made instead of 1 c.c. for lathering A "degree" of hardness upon Clark's scale corresponds to a grain of calcium carbonate per gallon.

The disadvantages attendant upon the use of hard water have already been referred to in speaking of

^{*} Hardness, unlike other analytical data, is still commonly expressed in grains per gallon. The conversion into the standard terms is readily effected (see note, page 62), for instance, from grains per gallon to parts per 100,000 by dividing by 0.7.

lime and magnesia in water. Temporary hardness is of less moment, both for dietetic and washing purposes, than permanent hardness; it depends upon the comparatively harmless calcium carbonate, which is removed by boiling. Calcium and magnesium salts have been held responsible for goitre, which in certain districts, notably in India, seems to be limited to magnesian limestone formations. It is, however, by no means always prevalent in magnesian limestone districts, even when the water contains abundance of magnesium salts; nor is it always absent from districts free from limestone formations. Hence many authorities are inclined to attribute it to iron pyrites and other salts of iron and copper often found in limestone regions; while others consider that the concurrent agency of some organic material, or a specific organism, is necessary. Goitre appears to have been developed, under the use of certain drinking waters, in as few as eight or ten days. Urinary calculi are common in Norfolk and other chalk districts where the water is hard; but the dependence of the calculi upon the hardness of the water is far from being established. A good drinking water should not exceed 4 degrees of permanent, or 20 of total, hardness. It is estimated that each degree of hardness involves a waste of upwards of a pound of soap for every 1,000 gallons used in washing.

Organic matter in solution consists of animal and vegetable matters that have gained access to the water, and of the products of their decomposition. Little is known of its chemical composition, the variety and complexity of the components

being so great as to defy analysis.

The presence of organic matter (or rather, of oxidisable matter) may be shown by adding a few drops of a solution of potassium permanganate, which is immediately decolorised; or by slightly acidulating the sample water and boiling it for 20 minutes with a few drops of solution of chloride of gold, which will give, according to the amount of organic matter present, a rose, violet, or olive colour, or a violet or black precipitate.

A rough quantitative estimate is obtained by determining the combustible matter in the "total solid residue." 100 c.c. of sample water are measured into a platinum dish of known weight A. They are then evaporated to dryness over a water bath (after which the drying should be continued in a water oven at 100° C. for an hour more), and the weight B of the dish and its now dry contents is ascertained. B—A is the weight of the total solid residue, that is, of the organic matter and mineral salts. The dish is next heated to dull redness, to burn off the organic matter, and weighed again (C) after cooling. The loss on ignition, that is B—C, will give the amount of organic matter in the volume of water taken.* The proportion of organic matter per litre, or per gallon, is readily calculated from these data. Thus,

100 c.c. : 1,000 c.c. :: B-C : x volume taken. a litre. mgrs. lost on ignition.

x being the proportion of organic matter expressed as parts per million, or milligrammes per litre. During ignition the organic matter turns brown, or even black, before disappearing, and if of animal origin it gives off pungent fumes.

This method is not sufficiently exact for most quantitative purposes, and recourse is had to other means, of which the

chief are :-

Wanklyn's ammonia process,

Frankland's combustion process, and

Forchammer's oxygen process.

None of these professes to give the total amount of organic matter in water, but each fixes upon some one or more element, compound, or property, which is determined with precision and taken as a measure or index of the total organic matter, to which it is assumed to bear a constant ratio.

Wanklyn's Ammonia Process.—Most nitrogenous organic matter, when heated with permanganate of potash in presence of excess of potash, gives off a certain proportion of its nitrogen in the form of ammonia. Most, if not all, of the injurious organic impurities occurring in water may reasonably be assumed to be nitrogenous, and as they yield a definite and fairly constant proportion of their nitrogen

^{*} This is not strictly accurate, since water of crystallisation is driven off from the saline matters, and some small portion of the salts may be volatilised. Carbonates lose CO₂, nitrates are reduced, and chlorine volatilises.

as ammonia we have a very convenient and delicate means of measuring them. It still remains, of course, in this as in all other indirect processes, to establish empirical standards of purity, in terms of the ammonia index.

Certain standard solutions are required.

A. Standard solution of ammonium chloride, containing

0.01 mgr. of ammonia in 1 c.c.

B. Nessler's solution is a saturated solution of mercuric iodide in potassium iodide. It gives a yellowish tinge, with the faintest trace of ammonia, and a yellow-brown precipitate

(NHg₂I,H₂O) if more is present.

"Nesslerising" is a process by which the free or saline ammonia in a sample of water can be readily determined with great accuracy. Two tall cylindrical, flat-bottomed glasses, marked at 50 c.c., are placed side by side upon a white surface. In one of them are placed 50 c.c. of the sample water, and in the other 50 c.c. of distilled water free from any trace of ammonia. Two c.c. of Nessler solution are added to each glass, and the tint struck by the sample is exactly imitated by adding to the second glass successive measured quantities of the standard ammonia solution (A) from a burette, until the right tint is reached. Each c.c. of ammonia solution used corresponds to 0.01 mgr. of ammonia present in the 50 c.c. of the sample, from which data the amount per litre is readily calculated.

C. Alkaline permanganate solution. Eight grms. of potassium permanganate and 200 grms. of solid caustic potash are dissolved in a litre of distilled water.

The analysis is conducted as follows:-

Half a litre of the water to be examined, made alkaline if necessary, is put into a large flask, connected with a Liebig's condenser by means of a bent glass tube passing through a perforated indiarubber stopper. The flask is heated by a Bunsen's burner, and the first 50 c.c. of distillate are collected and Nesslerised, and the amount of ammonia they contain is recorded. This ammonia is part of the free ammonia contained in the water as such, and it is found by experience that the first 50 c.c. contain three-quarters of the free ammonia. Hence, adding \(\frac{1}{3}\) to the amount recorded in the first 50 c.c., we get the whole amount contained in the half litre of water under analysis, and, doubling this, the mgrs. of free ammonia per litre, i.e. parts per million.

Thus, if the first 50 c.c. strike a tint with Nessler, which is matched in the second glass by the addition of 1.2 c.c. standard ammonia solution, we must allow \(\frac{1}{2}\) of 1.2, that is, 0.4, making

a theoretical total of 1.6 c.c. This corresponds to 0.016 mgr. of ammonia in the half litre taken for analysis, or 0.032 part per million. Meanwhile 150 c.c. more are distilled over, to exhaust the free ammonia, but as already explained it is

unnecessary to Nesslerise these.

There are then 300 c.c. left in the flask, and to this are now added 50 c.c. of the alkaline permanganate, and the distillation is resumed. The first, second, and third 50 c.c. are collected separately and Nesslerised. Should the third 50 c.c. be found to contain noteworthy quantities of ammonia, a fourth 50 c.c. is distilled over.

The ammonia coming over at this second stage is called albuminoid ammonia, being derived from the albuminoid and other nitrogenous organic matter in the water. The second 50 c.c. contain much less than the first, and the third less than the second. Adding together the amount of ammonia found in the three lots, we have the total for the half litre of water; and twice this will give the proportion in parts per million.

> 1st 50 = 2.5 c.c. standard ammonia solution. 2nd 50 = 1.0 c c. ,, ,, ,, ,, 3rd 50 = 0.3 c c. ,, ,, ,,

Total = 3.8 e.c. = 0.038 mgr. ammonia per half litre. ** Albuminoid ammonia = 0.076 part per million.

Note. -Absence of any trace of albuminoid ammonia is suggestive of organic purity, even if free ammonia and chlorides are high. Albuminoid ammonia less than '05 part per million is consistent with extreme purity; and if free ammonia is absent or scanty, albuminoid ammonia under 0.10 is no evidence of impurity. If, however, there is much free ammonia, 0.05 albuminoid ammonia is suspicious, and 0.10 is strong evidence of pollution; 0.15 or more should condemn a water absolutely.

In good waters the free ammonia usually ranges from 0.00

or 0.01 to 0.03 part per million, and rarely exceeds 0.05.

Combustion processes involve costly and delicate apparatus, and a considerable degree of skill. The following

is an outline of Frankland's process:—

To a litre of the water are added 20 c.c. of saturated solution of sulphurous acid, to reduce the oxidised nitrogen and liberate all carbonic acid. The water is evaporated to dryness in special apparatus. The dry residue is mixed with oxide of copper and heated in vacuo in a combustion tube for an hour. The gases are collected over a mercurial trough; they contain all the organic carbon as carbonic acid, and the nitrogen as such.

They are measured volumetrically, and after the absorption of the carbonic acid by caustic potash the residue is read as nitrogen. The nitrogen present in the water as ammonia must be separately determined, and deducted from the total.

The significant points are the absolute and relative amounts of organic carbon and organic nitrogen. The lower the proportion of nitrogen to carbon, and the less the amount of each, the more favourable is the verdict, other things being equal. A low proportion of nitrogen to carbon, 1 to 8, indicates vegetable organic matter; a high proportion such as 1 to 3 is pretty certain proof of animal pollution. A river or surface water should not contain more than 0.2 organic carbon or 0.03 organic nitrogen in 100,000 parts, and spring or well water is open to suspicion if the carbon exceeds 0.1, or the nitrogen 0.03, per 100,000. In forming conclusions, due weight must be given to the other analytic data, especially the determination of chlorides and oxidised nitrogen, and to the other evidence obtainable, such as the position and surroundings of the source of supply, the facilities for pollution, and the depth and geological characters of the strata from which it is derived.

The following table, taken from the Report of the Rivers Pollution Commissioners, gives the average results of a large number of samples upon Frankland's method, stated as parts

per 100,000 :-

tion continues and him to the same of the	Rain- water.	Upland Surface Water.	Deep-Well Water.	Spring Water.
Total Solid Impurity	2.95	9.67	43.78	28.20
Organic Carbon	.070	*322	.061	.056
Organic Nitrogen	*015	.032	*018	*013
Ammonia	*029	*002	.012	.001
Nitrogen as Nitrates and			-	
Nitrites	.003	*009	*495	.383
Total Combined Nitrogen	.042	.042	*522	*396
Chlorine	0.82	1.13	5.11	2.49
Hardness:-		1000	100000000000000000000000000000000000000	
Temporary	0.4	1.5	15.8	11.0
Permanent	0.5	4.3	9.2	7.5
Total	0.9	5.8	25.0	18.5
Number of Samples Analysu	39	195	157	198

Kjeldahl's process, applied by Blair to water analysis, is based upon the conversion of the nitrogen of organic substances into ammonium sulphate if charred by heating with strong

sulphuric acid. 250 c.c. of water are distilled down to 50, with addition of sodium carbonate if acid; the first 100 c.c. of distillate contains all the free ammonia. Then 10 c.c. of pure sulphuric acid and a little arsenious acid are added, and the distillation is continued. Presently the acid begins to fume, and becomes darkened by the charring of organic matter. The heat is continued for two hours after this, or until the colour disappears again. After cooling, a slight excess of pure solution of soda is added: and the ammonia thus liberated is distilled over, and measured as in Nessler's process.

This method, like Wanklyn's, gives us a record of the "free ammonia" and "organic ammonia" at two stages; but the yield of the latter is more constant and more complete than that obtained with alkaline permanganate. The results are converted into terms of "organic nitrogen" by multiplying by 0.824. Blair finds, by this process, less than 0.06 part of organic nitrogen per million in very pure water, and 0.06 to 0.12 in ordinary drinking water; he regards as suspicious any water containing more than this, unless peaty, and condemns

any over 0.32.

Oxygen process.—There are many modifications of the Forchammer method, of which Tidy's is the best known. For the sake of uniformity the metric system is substituted for "septems" in the following description:-

The standard solutions required are—

A. Potassium permanganate, 0.395 grm. in a litre of distilled water. Every 10 c.c. yields 1 mgr. of oxygen to oxidisable matter.

B. Sodium hyposulphite, 1 grm. per litre. c. Potassium iodide, 1 part in 10 of water.

D. Starch, 1 part boiled with 20 parts of water, and filtered.

E. Sulphuric acid, 1 part to 3 parts of water.

The object is to determine the amount of oxygen absorbed in 1 hour and 3 hours respectively. As the hyposulphite solution is liable to change, it is necessary to perform a blank

test with pure distilled water every time.

Four flasks are employed, two of which contain 250 c.c. of sample water, while the other two contain 250 c.c. of pure water for the control experiments. To each flask are added 10 c.c. of permanganate solution (A), and 10 c.c. of dilute sulphuric acid (E). If the water is very bad, the pink colour due to the permanganate may soon entirely disappear, and in this event a second and even a third 10 c.c. of permanganate must be added, in order always to have a pink tinge throughout the experiment. At the end of an hour one of the samples and one of the control flasks are tested to ascertain how much undecomposed permanganate remains to be deducted from the total used. For this purpose 2 c.c. of iodide solution (c) are added, and the remaining permanganate immediately acts upon it, liberating a proportionate amount of free iodine. The iodine is measured with precision by dropping in the standard hyposulphite solution until no more free iodine remains, the exact point of disappearance of the last trace of iodine being ascertained by adding 2 c.c. of starch solution (d) near the end of the process, and noting the disappearance of the blue iodide of starch.

In the blank test the 10 c.c. of permanganate should remain unaltered at the end of the hour, so that if x c.c. of hyposulphite are used, these x c.c. correspond to 10 c.c. of the permanganate solution, that is, to 1 mgr. of oxygen. If, therefore, the sample flask, which has, of course, lost some of its permanganate, takes only y c.c. of hyposulphite, it is clear that the oxygen consumed by oxidisable matter must

have been $\frac{x-y}{x}$ mgr. If a second 10 c.c. of permanganate had been employed, the formula would obviously have become $\frac{2x-y}{x}$ mgr.

As a quarter of a litre of the sample was taken for each analysis, four times this result will give the "oxygen consumed" in milligrammes per litre—that is, in parts per million.

The determination of oxygen consumed in three hours is conducted in exactly the same way, at the end of that time.

Interpretation of results of the oxygen process.—Not only organic matter, but also nitrites, ferrous salts, or sulphuretted hydrogen, will reduce permanganates, and these latter, if present, must be removed or allowed for. No distinction is made between nitrogenous and non-nitrogenous organic matter. In London waters the organic matter is said to be about eight times the "oxygen consumed." It seems that putrescent matters, notably urine and other animal matter, are readily oxidised by permanganate, so that while the three hours' experiment gives information as to the total amount of organic matter, the one-hour reaction is important as indicating the proportion of putrescent and, therefore, presumably dangerous impurities. Sometimes four hours are

allowed instead of three, and a quarter or half an hour instead

of one hour. Peaty waters consume much oxygen.

As in all other modes of estimating the organic matter in water, the results of the oxygen process must be considered together with the other analytic data and the source of the water. In a general way, Tidy classifies waters as of great organic purity if the oxygen consumed does not exceed 0.5 part per million, of medium purity if not exceeding 1.5 part, of doubtful purity up to 2.0 parts, and as impure if the oxygen consumed exceeds 2.0 parts per million. Upland surface waters, however (the organic matters being less likely to be harmful), are judged by a more lenient standard, the degrees of which are double those given above, namely, 1.0, 3.0, and 4.0, in place of 0.5, 1.5, and 2.0.

Blair's oxygen process at 100° C. gives very satisfactory results. The reagents are similar to Tidy's, but the following

differ in strength:-

Potassium permanganate, 1.975 gramme per litre, so that 10 c.c. vield 5 mgrs. oxygen.

Sodium hyposulphite, 3.875 grammes per litre. Sulphuric acid, 100 c.c. to 900 c.c. distilled water.

Half a litre of water is taken for analysis, and after the addition of 10 c.c. of permanganate and 10 c.c. of acid, is boiled gently for two hours, further charges of permanganate being added if necessary. After cooling, the procedure is the same as in Tidy's process. Iodide is added, and the iodine set free by the unconsumed permanganate is measured by hyposulphite solution with the aid of starch.

The method of calculation also is substantially the same. Since the 10 c.c. of permanganate yields 5 mgr. of oxygen

instead of 1 mgr., the formula becomes $5 \times \frac{x-y}{x}$, or if n

charges of permanganate have been added, $5 \times \frac{x \times n - y}{x}$

This gives the mgrs. of oxygen absorbed per half litre, and doubling the result we have mgrs. per litre, or parts per million. Blair's process has obvious advantages over Tidy's, since the conditions as to temperature are constant, and the oxidation of organic matter practically complete. The empirical standards of purity necessarily range higher. Exceptionally pure waters take less than 2 parts per million, waters of medium purity 2 to 4 parts, and suspicious waters 4 to 6 parts. Even a peaty water should not be recommended if it takes much more than 6 parts.

Interpretation of the results of water examination.—From what has been said, it will be seen that there are various methods of water analysis or examination by which we may obtain a knowledge of a water sufficient to guide us as to its suitability for drinking purposes. In the past too much emphasis has been given to the results of chemical analysis and too little to the facts obtained by bacterioscopic and microscopic examination, and by inspection of the gathering ground. It cannot be too clearly stated that all the facts obtainable respecting a water are necessary to a reliable conclusion. Nothing is more likely to lead to an incorrect opinion than reliance upon isolated findings in respect of a water, whether bacteriological or chemical. The two must be taken together and in conjunction with a careful inspection of the sources, and even then the matter must be judged broadly, and the "life history" of the water, if such a term is permissible, taken into consideration.

The chief interpretations of chemical analysis have been already referred to in the text, the importance of gases, mineral salts, and soluble organic matter having been dealt with. It is therefore unnecessary to recapitulate. Here it is only necessary to add that these general rules are subject to modification according to the history and surroundings of the water. Vegetable is much less dangerous than animal contamination, and may often be recognised by the slowness with which the albuminoid ammonia comes over,* by the absence or small amount of chlorides and of free ammonia, microscopic by and examination of the sediment or knowledge of the source of the supply. Animal pollution is indicated if the albuminoid ammonia comes over rapidly, or if the chlorides or oxidised nitrogen are abundant. Pollution by absorption of effluvia, or by the

^{*} The second 50 c.c. containing little less than the first, and the third than the second.

presence of putrefying masses in the water, would, however, be unaccompanied by any excess of chlorides. It is important to take into consideration also the character of unpolluted water in the same district or geological field, since chlorides, oxidised nitrogen, and even a certain amount of free and albuminoid ammonia, may be found in waters that are free from recent and therefore dangerous pollution. It is quite possible, too, for an organically pure water to be unfit for use on account of its mineral constituents. Water from a shallow well, open to suspicion of pollution, should be condemned if the chlorides and oxidised nitrogen are high, even if albuminoid ammonia be scanty. A water that is pure at the time of sampling may still be liable to intermittent contamination. It must be remembered that Wanklyn's process tells us nothing as to the exact amount or specific nature of the pollution, or whether it is injurious or harmless. The solid matters of enteric or choleraic excreta probably do not greatly differ as regards their yield of albuminoid ammonia from those of healthy excreta, and minute proportions of either one or the other intentionally added to water in experiments by Cory and Dupré caused, as was to be expected, only a proportionately trifling addition to the albuminoid ammonia already present. Such minute but deadly pollutions, by enteric virus or any other poison, rarely occur in nature, but when they do, chemical analysis will give us no intimation of them. Hence it is true that chemical analysis "can tell us of impurity and hazard, but not of purity and safety" (Buchanan); but to this it should be added that the circumstances are rare indeed in which a careful consideration of all the analytical data, together with a study of the source of supply and its surroundings, will fail to give warning of any danger that may exist.

The summary on page 87 is slightly modified from

a table given by Parkes and De Chaumont. It shows in a general way the value to be attached to each analytical datum *per se*, but it will be evident that the interpretation is subject to material modification

according to the other information available.

Bacteriological examination is the most delicate and direct test of the safety of a water for drinking purposes, for by it we obtain information as to its potentiality to cause infective disease. Further, Klein and others have shown that by bacteriological methods it is possible to detect smaller degrees of sewage pollution than by chemistry. On the other hand, it is, of course, idle to expect to learn the exact chemical constitution of the water by bacteriological methods. Like chemistry, bacteriology must be judged by what it can tell us and not by what it cannot. There are three important groups of facts obtainable by a thorough bacteriological examination of water. These concern (a) the number of bacteria per c.c.; (b) the presence of any organisms of contamination; and (c) the presence of any specific organisms of disease. The number of bacteria per c.c. is, as a rule, only valuable when used in a series of examinations of the same water supply, or when used in the examination of water before and after filtration. The number of bacteria per c.c. is therefore a fact of only relative importance, because there is no standard of how many organisms should be present in each c.c. of a potable water, and there is no means by which the number can be accurately measured. For purposes of comparison it may be said that the metropolitan water supply, as consumed, usually contains less than 20 bacteria per c.c.

The organisms which indicate contamination of a water are certain liquefying bacilli, B. coli, B. enteritidis sporogenes, and streptococci. The chief of these as an indicator is B. coli. It is now held that if this

-	Impure Water.	Over 700. Over 70. (Much blackening, with nitrous fumes, or smell of burnt horn.)	Over 100. Marked. Marked. Over 6° Clark. Over 0.10. Over 0.15. Over 3.0.	Over 2.0. Present.	Presence of other met- als, especially lead.	Turbid. Much.	Any marked taste.	Abundant animal and vegetable forms. Fungi. Beggiatoa. Epithelial scales. Evidences of sewage. Ova of parasites.
and Dan Walens.	Suspicious Water.	400 to 700. Over 40. (Much blackening, or nitrous fumes given off.)	Over 50. Present. Marked. Over 4° Clark. Over 0.05. Over 0.10. Over 2.0. Over 0.3.	Over 1·5. nil.	Trace of iron.	Turbid. Much.	Yellowish tinge. Any marked taste.	Colourless vegetable forms. Organic débris. Fibres of cloth, or other evidence of house-refuse.
	Usable Water.	Under 400. Under 40. (Solids may blacken slightly on ignition, but no fumes should be given of	Under 50. Present. Under 4° Clark. Under 0.05. Under 2.0. Under 9.3.	Under 1.5. nil.	Trace of iron.	Clear, bright. (nil, or separable by subsidence or subside	faint blue or green tinge only. Palatable. nil.	Mineral matter. Vegetable forms with endochrome. Large animal forms. No organic débris.
	Pure Water.	Under 100. Under 15. (Solids should scarcely blacken on ignition.)	Under 15. nil, or trace only. 2° Clark. Under 0.02. Under 1.0. Under 1.0.	Under 1·0. nil.	nil.	Clear, bright.	nil. Palatable. nil.	nil, or as in next column.
	DATA.	M. Chemical (in parts per million): 1. Total dissolved Solids . 2. Loss in do. on ignition .	s ss., Permai ia, Free ia, Albumi Carbon Nitrogen	11. Oxygen taken from { Acid Permanganate } 12. Sulphides (Forchammer)	13. Metals		3. Colour	6. Microscopic Characters of Sediment

bacillus can be detected in 1 c.c. or 2 c.c. of a water, the organism has gained access recently from sewage. Its relative abundance is a matter of importance. Houston considers the presence of streptococci in water as indication of recent and dangerous pollution. The presence of any pathogenic organisms (B. typhosus, the cholera bacillus, etc.), in however few numbers, is, of course, sufficient for the condemnation of a water. Such organisms are rarely isolated from water, even though for external reasons it be evident that the water is the channel of infection, owing to the long incubation period of typhoid fever, the fact that the bacillus only lives in water for a few days, and the impossibility as a rule of examining the water

which actually caused the outbreak.

By way of summary, it may be said that a water should be condemned, as a rule, (a) if it contains a very large number of bacteria of whatever kind; (b) if it contains B. coli communis, B. enteritidis sporogenes, or streptococci, in one cubic centimetre, or any pathogenic organism; (c) if it gives the "enteritidis change" in milk cultures, or ferments glucose or lactose media. In addition to the chemical and bacteriological findings, the careful microscopic examination of the sediment and the thorough inspection of the gathering grounds, the source and course of the water, are necessary. In many ways the latter is the most important part of an examination of a water, and when in doubt as to chemical and bacteriological characteristics, topography may not unfrequently throw light upon the subject.

Drinking Water and Disease.—The relation of drinking water to typhoid and cholera will be considered subsequently. Here it is only necessary

to add a general note.

Vegetable matter in suspension and even in solution may cause diarrhea. In small quantity, however, dis-

solved peaty matter is harmless, although it gives a brownish tinge to the water.

Animal matter in suspension or solution, whether due to percolation from cesspools, leaking drains, farmyards, or manured fields, or to direct pollution of streams by drains or sewers, or to presence of decomposing animal tissues in the water, or to absorption of effluvia,

is always dangerous to health.

The products of simple decomposition of animal matter are in themselves capable of producing diarrhea and other acute alimentary disturbances, or of insidiously undermining the general health, and so possibly preparing the way for other diseases not directly dependent upon water-borne poison. In tropical countries polluted water is a frequent cause of dysentery, sporadic or epidemic. Sudden localised outbreaks of diarrhœa in Great Britain have been proved to originate in contamination of drinking water by effluvia from sewage, even without the entrance of liquid or solid filth. Apart from all this, the presence of organic filth in water shows that the way is open for the access of more specific poisons, such as those of cholera, enteric fever, dysentery, and parasitical diseases. The tendency of any such pollution is to increase in amount, while percolation through soil loses in course of time whatever purifying effect it may possess at first.

There are also certain disease conditions produced by water owing in part to its chemical constitution, as, for example, goitre, plumbism, urinary calculi, etc.

CHAPTER IV.

FOOD.

The useful constituents of all food substances, animal, vegetable, and mineral, are classified as—

1. Nitrogenous, including the animal and vegetable

albuminoids (proteids) and gelatin.

2. Fatty, including animal and vegetable fats and oils.

3. Carbohydrates, including starch and the sugars.

4. Salts, organic and inorganic.

5. Water.

Nitrogenous food.—Proteids are converted by the action of the gastric and pancreatic juices into soluble peptones, which are absorbed and used for the nutrition and repair of the nitrogenous tissues, and in some degree for the production of heat and the functional activity of the system. During life the nitrogen is eventually eliminated in the urine (as urea or urates), and in the fæces. Fat is another product of the splitting-up of albumen, and its hydrogen and carbon ultimately go to form carbonic acid and water. Muscular exertion does not increase the demand for nitrogen or the excretion of urea, these being fairly constant in a given healthy individual under suitable diet, whether working or not. Assimilation of excess of nitrogen, in the absence of exercise, is apt to lead to imperfect oxidation, with formation of urates and uric acid rather than urea. Other consequences are engorgement of the liver and general plethora, pyrexia, tendency to diarrhea, and ultimately to a gouty condition.

Deficiency of nitrogen leads to diminution of circulating albumen, anæmia, loss of weight, and impairment of energy and stamina. Ultimately the tissue albumen, and especially that of the muscles, is reduced.

Nitrogenous food is therefore essential for the growth, maintenance, repair, and functional activity of the tissues, and contributes in some measure to

the production of heat and force.*

Fats and carbohydrates supply the oxidisable material for the production of heat and force, and are ultimately eliminated as carbonic acid and water. They aid in the nutrition of the tissues and in the excretion of waste material. An excess of either lessens oxidation of nitrogenous matter and leads to storage of fat in the tissues; deficiency causes increased metabolism of circulating albumen, loss of weight, and impairment of nutrition. Sugars, starches, and certain other substances are termed carbohydrates, because their molecule contains (besides carbon) twice as many atoms of hydrogen as of oxygen, and may therefore be represented empirically as C_x(H₂O)_n, or carbon plus water. Glucose is C₆H₁₂O₆, cane sugar C₁₂H₂₂O₁₁, and starch C₆H₁₀O₅. Fats and oils (animal or vegetable) contain much less oxygen in proportion to their carbon and hydrogen, stearin, for example, being C₂₁H₄₀O₆. Carbohydrates are more readily assimilated than fats, but have a lower nutritive value by at least 40 per cent. They are converted into grape-sugar by the digestive processes. As a rule, nutrition ultimately suffers if either fats or carbohydrates are withheld; but to a certain extent they are interchangeable, and the diet of the Eskimo contains little or no carbohydrate.

Fat as a tissue is useful for its mechanical properties—namely, non-conduction of heat, lubrication, and soft elastic pressure; in excess it becomes a source of inconvenience or danger, as in extreme obesity and in

fatty infiltration of the heart.

^{*} Gelatin appears to be capable of replacing the circulating albumen, but not of forming tissue albumen or repairing its waste. Its nutritive value is said to be about one-fourth of that of albumen. Digestion converts it into a sort of peptone, which does not gelatinise.

Muscular contraction is the source of animal heat as well as of force, and the heat generated is four times the heat-equivalent of the work done. Although the hydrocarbonous matter supplies the "fuel" for these purposes, it is necessary that a certain amount of nitrogenous matter shall be associated with it. Fat or carbohydrate alone is not sufficient.

Salts.—Lime, potash, and soda are essential—lime for bone-formation and cell-growth, potash for nutrition of formed tissues, soda for fluid tissues.* Iron in small quantity is required for formation of the colouring

matter of the blood (hæmoglobin).

Chlorides and phosphates are indispensable. Salts of organic acids (lactates, tartrates, citrates, malates, acetates), which are oxidised into carbonates in the system, are important as maintaining the alkalinity of the blood.

Water constitutes two-thirds of the whole weight of the body, and forms part of every tissue. By its agency food is dissolved and carried to the tissues, waste products are removed and excreted, nutrition and chemical changes are rendered possible, and the temperature of the body is regulated by means of evaporation.

Stimulants and condiments are not essential consti-

tuents of a complete dietary.

Dietaries.—Hutchison distinguishes four criteria of the value of a dietary, (a) chemical, turning upon nutritive constituents as above; (b) physical, which measures the value of a food by the amount of heat produced by complete combustion; (c) physiological, i.e. assimilability; and (d) economic, nutritive value as compared with cost. Only the second of these calls for further comment here. The unit of heat production is the kilo-calorie, or as it is generally termed the calorie, that is the amount of heat re-

^{*} A vegetable diet calls for additional sodium salts, to make up for the increased elimination of these due to the excess of potassium salts supplied.

quired to raise 1 kilo (or 1 litre) of water 1° C. (or 1 lb. of water 4° F.). Rubner has shown that the heat value of 1 gramme (= 15½ grains) of proteid and 1 gramme of carbohydrate is 4·1 calories each, and of 1 gramme of fat is 9·3 calories. Multiplying the percentage of proteid and carbohydrate which a food contains by 4·1 and the fat by 9·3 gives the total calories per 100 parts of the food in question. For example, if 100 grammes of milk contain 2 of proteid, 4 of fat, and 6 of carbohydrate, the calories would be of proteid 8·2, of fat 37·2, and of carbohydrate 24·6, or a total of 70·0 per 100 grammes.

There are two chief requirements in a nutritious daily diet for a healthy man doing a moderate amount of work. First, it must contain food which will provide sufficient energy (average 3,500 calories); and secondly, it must provide for the upkeep and repair of the tissues of the body (average, 125 grammes of proteid, 500 grammes of carbohydrates, and 50 grammes of fat). Such a standard, according to Hutchison, yields carbon, nitrogen, and calories as follows:—

Car Child House	Carbon. (in grms.)	Nitrogen. (in grms.)	Calories.
125 grms. Proteid	62	20	512.5
500 ,, Carbohydrates 50 ,, Fat	200 38	=	2050 465
Total	300	20	3027:5

These quantities may be expressed in English ounces as $4\frac{2}{5}$ oz. of proteid, $2\frac{1}{2}$ oz. of fat, and 18 oz. of carbohydrates, making $24\frac{9}{10}$ oz. of dry food.

Rubner has suggested as a standard that whilst 2,500 calories are necessary for a person at rest, 3,121 calories are required for a person doing moderate muscular work, and 5,213 calories in hard labour.

Even more than this is actually consumed sometimes. Chittenden considers that the current estimations of the amount of proteid and total fuel-value necessary for hard work are excessive. For hard work all the chief constituents of food may be increased, but there is evidence to show that the most valuable source of muscular energy is the carbohydrates. For mental work the digestibility of food is of greater concern to the brain worker than chemical composition (Hutchison). Reduction on a muscular diet should, however, affect fats and carbohydrates more than proteids. As already remarked, age and sex exert a marked influence upon dietetic requirements. The same must be said in respect of climate, season, and personal peculiarity.

From the chemical standpoint authorities suggest that a daily dietary for a male adult should be composed somewhat as follows:—

	Nitroge- nous	Fat.	Carbohy-	Salts.	Such a diet would contain		
	Food.		drates.		Nitrogen. Carbon		
Rest Moderate \ \ work . \ \ Hard work .	3 oz. 4½,,,	1½ oz. 3 ,, 4½ ,,	12 oz. 15 ,, 18 ,,	1 oz. 1 ¹ / ₄ ,, 1 ¹ / ₂ ,,	200 grs. 300 ,, 400 ,,	4000 grs. 5000 ,, 6000 ,,	

These amounts are only approximate. The necessity for so great an increase of nitrogen during hard work is open to question, and on the other hand the hard work standard of carbon in the above table is low. More food, especially fat, is needed in cold climates, less in hot.

All the ingredients are supposed to be water-free, but would, in practice, be combined with at least their own weight of water, raising the total daily weight of solid food to 40 or 50 oz. In addition to this 50 to 80

oz. of water are taken as drink. A man consumes daily about $\frac{1}{100}$ of his weight of dry solid food and $\frac{3}{100}$ of water. Women are said to require about 10

CHEMICAL COMPOSITION (APPROXIMATE) OF CERTAIN FOOD SUBSTANCES. (Parkes.)

	Per Cent. of					Grains per lb.	
A STATE OF THE STA	Water.	Nitroge- nous.	Fatty.	Carbo- hydrates.	Salts.	Nitro- gen.	Carbon.
Good meat,* beef or mutton, with little fat (uncooked)	75	20	3.5		1.5	190	1900
Average meat* } (uncooked)	75	15	8.5		1.5		
Roast meat	55	28	15		3		
Very fat meat*	63	14	19		4		30,001
Fat pork	40	10	50		2	100	4000
Salt pork†	45	25	7		25	290	1360
Salt beef†	50	30	0.2		20	325	1115
White fish	78	18	3		1	200	875
Eggs	74	14	11.5		1		à
New milk	87	4	3.5	5	0.5	45	600
Skim milk	90	4	2	5	0.8	45	450
Cheese	37	33	24		5.5	300	3300
Butter	6	0.3	90		2.5		6500
Bread	40	8	1.5	50	1.5	90	2000
Flour	15	11	2	70	1.7	120	2700
Oatmeal	15	13	6	65	3	140	2800
Indian meal	14	10	7	65	1.5	120	3000
Peas (dry)	15	22	2	55	2.5	250	2700
Green vegetables .	90	0.2	0.5	6	0.7	14	420
Carrots	85	0.6	0.5	8.5	0.7	14	500
Potatoes	75	1.5	0.1	23	1	22	770
Rice	10	5	0.8	83	0.5	70	2700
Sugar	3			96	0.5		3100‡

^{* 20} per cent. should be allowed for bone, and 20 to 30 per cent. more is lost in cooking.

[†] Brine dissolves out myosin and other important constituents. The nutritive value of salt meat is not more than $\frac{2}{3}$ that of fresh meat. \ddagger Milk-sugar yields only 2,800 grains of carbon per lb.

per cent. less than men, both of carbon and nitrogen. Infants require nitrogenous and fatty foods with only a small proportion of carbohydrates, which is supplied by milk-sugar. A child of ten years of age needs only half as much, but at fourteen years quite as much, as a woman. The dietetic requirements of old age are just the reverse of those of childhood.

The table on page 95 supplies the means of determining how far a given dietary complies with the requirements stated above, and conversely of constructing a dietary that will satisfy these conditions.

Thus, if the question should arise whether a daily allowance of 2 lbs. of bread and ½ lb. cheese is a sufficient diet for a man in easy work, the calculation may be made on a chemical basis as follows:—

100 oz. bread contain

Nitro. Fat. Carb.-hyd.
8 oz. 1½ oz. 50 oz.

100 oz. cheese contain
33 oz. 24 oz. 0 oz.

Whereas the theoretical amounts required for moderate work are

therefore 32 oz. contain
Nitro. Fat. Carb.-hyd.
2·5 oz. 0·5 oz. 16 oz.

Nitro. Fat. Carb.-hyd.
2·5 oz. 0·5 oz. 16 oz.

Nitro. Fat. Carb.-hyd.
2·6 oz. 1·9 oz. 0·5 oz. 16 oz.

4·5 oz. 3 oz. 15 oz.

The conclusion to be drawn is that such a diet is sufficient as regards the nitrogenous matter and carbohydrates, but wanting in fat.

A still readier method is to calculate the amount of carbon and nitrogen, thus:—

1 lb. bread contains	therefore 2 lbs. contain
Nitrogen. Carbon.	Nitrogen. Carbon.
90 grs. 2000 grs.	180 grs. 4000 grs.
1 lb. cheese contains	therefore ½ lb. contains
300 grs. 3300 grs.	150 grs. 1650 grs.
	330 5650
The theoretical amounts of car nitrogen required for moder	
being	300 5000

Hence, if we leave out of consideration the differences between fats and carbohydrates the diet more

than satisfies the theoretical requirements.

The problem may also be presented the other way round. What quantity x of bread and y of cheese would be sufficient for an ordinary diet? The requirements are 300 grains of nitrogen, and 5,000 of carbon; and x lbs. of bread contain $90 \times x$ grains of nitrogen and $2,000 \times x$ grains of carbon, while y lbs. of cheese yield $300 \times y$ grains of nitrogen and $3,300 \times y$ grains of carbon. Hence we have a simultaneous equation—

$$\begin{array}{r} 90x + 300y = 300 \\ 2000x + 3300y = 5000 \end{array}$$

giving as approximate values x = 1.7 lb., and

 $y = \frac{1}{2}$ lb.

It has been found experimentally that the energy which is developed by oxidation is for one ounce of lean meat about 50 foot-tons,* carbohydrates 150 foottons, and fats 300 foot-tons. In this relation it should not be forgotten that the oxidation of food in the system is by no means complete, even as regards that portion which is assimilated.

In calculating the mechanical work done, that of respiration, circulation, and locomotion must be taken into account, as well as the load carried or lifted. According to Haughton and Parkes the force exerted daily in respiration, circulation, and other "internal" work is about 260 foot-tons; and in addition to this a strong man can do external work equivalent to 300 to 500 foot-tons more per day. Walking along a level road at three miles per hour is equivalent to climbing vertically 1/20th of the distance traversed; at 4 miles 17th. Hence, if a man weighing W lbs. carries a load of X lbs. on a horizontal track for D feet, at

3 miles per hour, he exerts energy equivalent to $\frac{(W+X)D}{20 \times 2240}$ foot-tons in so doing. A march of 20

miles along the level involves work equivalent to 350 foot-tons, taking the weight of the body as 150 lbs. If a load of 60 lbs. is added, the work is 500 foot-tons. The whole work done, estimated in this way, represents only $\frac{1}{7}$ th of the force theoretically obtainable from the food by combustion. The energy expended in a fair day's work is about one-fifth mechanical labour to

four-fifths heat (Martin).

Animal is generally held to possess certain advantages over vegetable food, of which the most certain are the ready supply of blood-pigment, the larger percentage of proteids, the greater digestibility of animal fats, and the smaller bulk required. A vegetable dietary, unless carefully selected, is apt to contain insufficient nitrogen, but it is rich in carbohydrates. It is, however, bulky, less digestible in the stomach, and less completely absorbed. The supposed inferior nutritive value of vegetable albuminoids is not so clear, although they are probably less rapidly digested; and a well-fed vegetable eater may display for a time as perfect health and energy as a meat-eater.* On the other hand, the argument from analogy with the herbivora, some of which are types of activity, loses weight from the inability of man to digest cellulose. The consistent vegetarian must either live on a diet deficient in proteid or consume an excessive bulk of food. The adoption of the former course tends to diminish energy and tissue resistance, and the latter is likely to lead to derangement of the digestive organs.

Morbid conditions dependent upon diet.— An excess of food, due to too large or too frequent meals,

^{*} Vegetarianism as ordinarily practised does not exclude animal fats or even albuminoids; milk, butter and cheese being used freely.

may accumulate in the intestine, causing fermentation and also dyspepsia, with constipation or ineffective diarrhea. Gout, obesity, gall-stones and other conditions may also arise from excess of food. Absorption of the products of putrefaction may give rise to a septic condition marked by pyrexia, furred tongue, fætid breath, heaviness, and possibly jaundice. Diseases of the blood may also arise from retention of waste products in the intestine, as for instance when chlorosis follows constipation. The effects of excessive or deficient assimilation of nitrogenous and carbonaceous food have already been considered.

Protracted insufficiency of diet is followed by wasting of the tissues (inanition). Adipose tissue is naturally the first to suffer, and may be almost completely absorbed, the other tissues following mainly in the inverse order of their importance to life. The urine still contains urea and urates, from oxidation of tissue, first of the circulating albumen, afterwards of tissue albumen. Physical and mental weakness ensue, followed by anæmia and an adynamic condition that powerfully predisposes to certain diseases, notably relapsing fever, phthisis, and pneumonia, and perhaps to all infectious diseases. Diarrhea is apt to occur, adding still further to the general emaciation and prostration. Ophthalmia, stomatitis, ulcers, and skin diseases of various kinds, are common; and any disease that may have obtained a hold upon the system is aggravated by the impairment of nutrition. Death ensues when the loss reaches about 40 per cent. of the normal weight of the body.

Apart from parasitic poison and infection, there are two well-marked morbid conditions that are almost always attributable to the absence of essential elements

of diet. These two are rickets and scurvy.

Rickets occurs in children fed upon a diet in which there is undue admixture of starchy food, and

a deficiency in fat and proteids. Starch cannot be digested by young infants, and its administration is not only useless but liable to cause diarrhea, which interferes with the digestion of other food. Prolonged indigestion has much to do with the cause of rickets. Even apart from this, an excess of starch or other carbohydrate has an injurious effect upon nutrition.

Scurvy does not occur when the diet includes plenty of fresh vegetables and fruits or their juices, or even preserved vegetables and fruits. The blood or raw flesh of recently killed animals is credited with considerable anti-scorbutic power. Citrate, tartrate, and malate of potash are also preventives, and, to a less degree, lactate and acetate. Carbonate of potash is inert. The disease affects under-fed men more readily, but cannot be checked by increase in the supply of nitrogenous food, carbohydrates, or fat. The fault probably rests with the supply of salts, and as scurvy occurs when there is no lack of phosphates or sodium chloride, potash and organic acids alone of the ordinary salines are missing. The salts of organic acids differ from those of mineral acids in one important particular—namely, their oxidation in the blood to form carbonates with alkaline reaction, although they may have been neutral or acid originally. Thus the citric acid and citrate of potash, which are the principal constituents of lime-juice, are oxidised into carbonic acid, which is removed by the lungs, and (alkaline) potassium carbonate. Potatoes are efficient anti-scorbutics, and contain a large proportion of organic acid, salts of potash, soda, and lime. The absence of land scurvy at the present time may not unreasonably be attributed in great measure to the universal use of potatoes, and other fresh vegetables and fruits, as articles of food.

Meat varies considerably in its nutritive qualities according to the age of the animal, the state of its

nutrition at the time of slaughter, and the proportion of fat. Well-fed prime meat contains more albuminoids and less connective tissue than that which is taken from animals that are badly fed, diseased, or too old or too young. The proportion of fat, roughly speaking, ranges from 50 per cent. in fat pigs or sheep to 33 per cent. in fat oxen or lambs, and 16 per cent. in calves. Absence of fat is largely compensated for by presence of additional water, so that the nitrogen remains fairly constant. As a rule, "white meat," such as fowl and rabbit, is less nitrogenous, more tender, and more digestible than "red meat," such as beef, mutton, and game. The latter become more tender if kept until rigor mortis gives place to incipient

putrefactive changes.

Relation to disease.—The flesh of animals suffering from any inflammatory disease is watery and innutritious, discoloured, possesses a stale odour, and decomposes rapidly. The use of such meat is liable to cause alimentary disturbance. The same applies, more or less, to animals emaciated from any cause. In some persons acute dyspepsia and diarrhea are caused by apparently normal meat of certain kinds, usually pork or mutton. As a rule, meat in any stage of putrefaction (the signs of which are smell, discoloration, and loss of elasticity) is liable to cause acute gastro-intestinal irritation, and other manifestations of septic poisoning, the prominent symptoms being vomiting, diarrhea, cramps, prostration, pyrexia, with weak and irregular pulse. Meat in an early stage of decomposition is frequently more toxic than when more advanced. It is not, however, easy to account for the impunity with which certain kinds of food, venison and game for instance, are habitually consumed in a state of early decomposition. It is probable that the toxic products are not present in anything like the same degree as in ordinary meat,

and it is possible that such products, and even the causal organisms, are different. Certain parasites, to which further reference is made in a later section, are clearly transmissible to man, and therefore suffice to render the whole carcase unfit for food; the chief being the *Tæniæ* in beef, mutton, and pork, and *Trichinæ* in "measley" pork. Sheep are liable to *Strongylus filaria* in the lungs, *Distoma hepaticum* in the liver, and *Cænurus cerebralis* in the brain; but these are held to require only the removal of the affected organs.

Accident, parturition, or purely local diseases, such as apoplexy or intestinal obstruction, do not necessarily render a carcase unfit for food provided that the animal has been slaughtered before any inflammatory or other constitutional mischief has occurred. An animal that has died with the blood in it is presumably diseased, and it is customary to condemn such a carcase on that account, and because decomposition rapidly sets in. The flesh of animals affected with any acute specific disease should be rejected, not only upon the grounds already stated with reference to inflammatory diseases, but also on account of the risk of imparting the specific malady. Anthrax has been thought to be so transmitted, and there is strong suspicion that actinomycosis has been similarly conveyed. Generally, it may be said that the direct transmission of disease by infected meat is best illustrated in two different classes of diseased meat represented by the Welbeck Disease and tuberculosis.

Welbeck Disease and allied conditions.—At Welbeck, Notts, in June, 1880, a large number of persons were taken ill after eating some ham at a public luncheon. Ballard found that this was the only condition common to all the sufferers. The onset occurred in some cases suddenly, for the most part between twelve and thirty-six hours later. The

symptoms included violent diarrhea, vomiting, and abdominal pain, usually accompanied or followed by pyrexia, thirst, headache, cramps in the limbs, cold sweats, and great prostration. Particulars were obtained of 72 cases, of which 4 were fatal. The ham was observed to have a disagreeable taste, but was not suspected to be tainted. It had been exposed to sewer air. Klein found spores and bacilli in the muscular fibres of the suspected ham, and similar bacilli in the kidney of a man who died of the disease. Dogs, cats, rabbits, guinea-pigs, and white mice were fed upon the ham, and nearly all developed pneumonia and other symptoms; inoculation of white mice and guinea-pigs gave the same result. Cultures were made, and guinea-pigs and white mice which were fed or inoculated therewith presented the same symptoms.

A second series of cases occurred in Nottingham, in February, 1881, among persons who had partaken of some hot baked pork. Fifteen were attacked, and one died. The time and mode of onset, and the general character of the symptoms, were similar to those noted in the Welbeck outbreak. the persons who ate of it were taken ill. Klein examined the tissues of the fatal case referred to above, and found the Welbeck bacilli in the blood, pericardial fluid, lungs, spleen, kidney, and elsewhere. Cultures inoculated into mice and guinea-pigs caused pneumonia and other symptoms in all; and the bacilli were found in the blood and exudations.

These two outbreaks drew attention to the whole question of meat poisoning, and since 1880, more than 50 such epidemics have been investigated. Some of the better known may be tabulated as on page 104.

Each outbreak has shown more or less clearly the characters common to these epidemics-1. Simultaneous attacks. 2. Similarity of symptoms and post-mortem signs. 3. A history of infection in

common. The usual symptoms include rigors, faintness, vomiting, diarrhea, abdominal pain, and occasionally skin eruptions. As a rule, certain nervous conditions have supervened, such as giddiness, headache, paralysis, mental depression, etc., and occasionally these symptoms have been predominant.

Date of Occurrence.	Place of Occurrence.	No. of Cases.	Period of Incubation in Hours.	Probable Source of Infection.
1880	Welbeck	72	12-48	Cold boiled hams
1881		15	12-34	Pork
1882	Nottingham Oldham	9	4	American tinned
1004	Olullani	9	4	pig's tongue
1882	Bishop Stortford	6	24	Beef
1882	Whitchurch	20	1-5	Brawn
1886	Carlisle	20	6-40	Ham & game pie
1886	Ironbridge	12	6-12	Veal pies
1887	Retford	80	8-36	Pork brawn
1888	Middlesbrough	114	0-00	American bacon
1889	Carlisle	25	24	Pork pies
1891	Portsmouth	13	14-17	Cold meat pie
1896	Mansfield	265	18-24	Potted meat
1898	Oldham and	200	10.21	2 octor men
120	Chadderton	54	48	Veal pies
1899	Nuneaton	42	12-48	Pork
1899	Sheffield	24	2-3	Corned beef
1902	Derby	221	4-24	Pork pie
1903	Chadderton	40	100000000000000000000000000000000000000	Ox tongue
1906	Bedfordshire	29	8-24	Pork pie
1967	Wigan	168	36	Potted tongue

Meat poisoning appears to depend not upon the number of bacteria present in the meat, but upon the particular species (that is, bacterial infection), and their products (ptomaines, toxins, that is, ptomainepoisoning). A long incubation period generally indicates poisoning by bacteria (an infection), and a short incubation period poisoning by products (an intoxication). In 1888, Gaertner of Jena investigated an outbreak of disease affecting 58 persons who had eaten uncooked meat. One victim died, and from his body, as well as from the meat, Gaertner isolated the

B. enteritidis, an organism allied to the coli group. Since that period, the B. botulinus of Ermengem, B. enteritidis sporogenes of Klein, and certain putrefactive bacteria, particularly of the coli group (p. 57), have been held responsible for causing such illnesses.

B. enteritidis (Gaertner). Characters similar to B. typhosus (morphology, motility, and staining properties), but grows more rapidly in gelatine; fewer flagella; ferments lactose and sometimes dextrose; does not produce indol or coagulate milk; positive neutral-red reaction; in litmus whey or litmus broth, acid is first produced, and then the medium becomes distinctly alkaline. Virulent to rodents and small animals (gastrointestinal symptoms, hæmorrhagic enteritis and swelling of lymph follicles). Forms no spores, and therefore cannot stand very high temperatures. (Killed at 70° C. in one minute.) It produces agglutinating properties in the blood of the patient.

B. botulinus (Ermengem). Van Ermengem describes, under the name of botulism, a state brought about by the ingestion of various articles of food, such as ham, tinned or preserved foods, oysters, mussels, etc., and which is characterised by comparatively slow onset (twelve to twenty-four hours after infection), secretory troubles, paralysis of certain muscles, particularly of tongue and pharynx, dilatation of pupil, aphonia, dysphagia, constipation, retention of urine, absence of unconsciousness and of fever, etc. Botulism differs considerably from the more common form of food-poisoning with which we are acquainted in England, and which is characterised by practically the same symptoms as those of epidemic diarrhoea. B. botulinus is 4-9 μ long and 9-12 μ broad; round, slowly motile, 4-9 flagella. Polar spores, killed in thirty minutes at 80° C. Liquefies gelatine; does not coagulate milk; anaërobic; in cultures often produces gas and a sour, rancid odour. Pathogenic for guinea-pigs, rabbits, and other small animals.

B. enteritidis sporogenes of Klein (see Epidemic Diarrhoea).

The B. enteritidis Derbiensis is a member of the Gaertner group, isolated by Delépine from pork pies the consumption of which caused the Derby outbreak in 1902. He considered the presence of the bacillus in the pork pies was due to contamination of the meat before it was cooked, and that the central parts of the pies were not thoroughly cooked. It frequently happens in these cases that some constituent part (such as jelly) of the manufactured article or prepared dish is really the polluted portion. Experience of meat poisoning outbreaks leads to the conclusion that the meat has contracted its poisonous properties in one or all of the following ways, viz., putrefaction or unsoundness in the meat itself, unclean manipulation or storage in insanitary places, and insufficient cooking. The methods of prevention are therefore obvious. Occasionally tinned goods cause poisoning, owing to metallic absorption, and this must be differentiated from bacterial poisoning.

Tuberculosis.—This disease is set up in animals by the tubercle bacillus, which is either identical or

nearly allied to the B. tuberculosis of Koch.

Tuberculosis is common among cattle, especially if stall-fed, and is increased by the practice of housing them with inadequate air-space and ventilation. Out of 23,000 cattle slaughtered at Leipzig in 1895, tuberculosis was found in 33 per cent. The tuberculin test has revealed the presence of tuberculosis in 54 per cent. of the cows of certain breeds in certain dairies in Great Britain.

Investigations carried out in 1902 in all the chief breeds in this country have shown that 14.7 per cent. are affected with tuberculosis (Geddes). Probably the general average is about 15-20 per cent. In London and other large cities, where a high degree of inspection is maintained, and where there is no in-breeding, a very small and negligible number of the cows are tubercular.

It has long been customary to condemn as unfit for food all parts of a carcase in which tubercle is found, and the whole carcase if the disease be far advanced and the animal emaciated. Following the conclusions of the Paris Congress of 1888 many authorities have adopted a more stringent standard, and condemn the whole carcase if there be any tubercle in any part, no matter how slight or localised.

The Royal Commission on Tuberculosis reported

in 1898 as follows:-

"The following principles should be observed in the inspection of tuberculous carcases of cattle:—

(a) When there is miliary tuberculosis of both lungs .

(b) When tuberculous lesions are present on the pleura and peritoneum . . .

(c) When tuberculous lesions are present in the muscular system, or in the lymphatic glands embedded in or between the muscles

(d) When tuberculous lesions exist in any part of an emaciated carcase.

Generalised tuberculosis is present, and the entire carcase and all the organs may be seized.

(a) When the lesions are confined to the lungs and the thoracic lymphatic glands | Localised tuberculosis is present, and the

(b) When the lesions are confined to the liver . .

(c) When the lesions are confined to the pharyngeal lymphatic glands . .

(d) When the lesions are confined to any combination of the foregoing, but are collectively small in extent)

culosis is present, and the carcase, if otherwise healthy, shall not be condemned, but every part of it containing tuberculous lesions shall be seized.

"In view of the greater tendency to generalisation of tuberculosis in the pig, we consider that the presence of tubercular deposit in any degree should involve seizure of the whole carcase and of the organs.

"In respect of foreign dead meat, seizure shall ensue in every case where the pleuræ have been

'stripped.'"

The tubercle bacillus is most easily found in the glands, and tuberculous deposits in the muscles are very rare. The bacillus is scarce in the caseating nodules. In the pig it is difficult to detect the bacilli as a rule, though the disease is very common in that animal, affecting chiefly the abdominal organs and the glands of the throat. It is rare in the sheep. The Royal Commission reported that "in tissues which go to form the butcher's joint, the material of tubercle is not often found even where the organs (lungs, liver, spleen, membranes, etc.) exhibit very advanced or generalised tuberculosis; indeed, in muscle and muscle juice it is very seldom that tubercle bacilli are to be met with; perhaps they are somewhat more often to be discovered in bone, or in some small lymphatic gland embedded in intermuscular fat." The chief way in which such meat substance becomes infected with tubercle appears to be through carelessness in the butcher, who perchance smears the meat substance with a knife which has been used in cutting the organs, and so has become contaminated with infected material. "Ordinary cooking, such as boiling and more especially roasting, though quite sufficient to sterilise the surface, and even the substance for a short distance from the surface of a joint, cannot be relied upon to sterilise tubercular material included in the centre of rolls of meat, especially when these are more than three pounds or four pounds weight. The least reliable method of cooking for this purpose is roasting before a fire; next comes roasting in an oven, and then boiling." (Sims Woodhead.)

It should be added that whilst it is evident that tuberculosis is not greatly spread by the consumption of tuberculous meat, the Royal Commission of 1895 definitely concluded that "we must believe that any person who takes tuberculous matter into the body as food incurs some risk of acquiring tuberculous disease." In spite of Koch's views to the contrary, which are discussed subsequently, that conclusion still repre-

sents, broadly, the views of most authorities.

Characters of good meat .- It should be firm and elastic, but not tough, and it should not "pit" on pressure; a thin red meat juice (acid) will often exude in small quantity on standing; the meat substance should be red throughout, not pale, purple, or green; marbled by layers of fat; free from purulent or gelatinous fluid in the septa between the muscular bands; fresh and pleasant in smell, as tested by a clean skewer, which has been plunged into the interior. The reaction should be slightly acid. The meat should "set" within twenty-four hours.* The flesh is pale in pork, in young animals, and in those suffering from exhausting diseases; dark in old animals, or those which have died with the blood in them, or in inflamed parts; wet in dropsy, and often in inflammatory diseases. With commencing putrefaction the colour becomes pale, and the smell disagreeable; later the meat softens in parts and turns green.

The liver, lungs, heart, and flesh should be examined for parasites; the head, stomach, and intestines for evidences of specific disease. The microscope would detect Cysticerci (usually visible to the naked eye) and Trichinæ in muscle, and Stephanurus dentatus in the brain. To demonstrate Trichinæ, a thin section should be put into liquor potassæ for a few minutes only until the muscle becomes translucent. The coiled

embryo will be seen inside the capsule.

^{*} Horse flesh may be differentiated from beef by the following characters: It does not set so rapidly or so well. It is more tendinous and fibrous, contains less fat, which is oily and yellow, and possesses a horsey smell. The colour is also more brown-red and less bright than in sound beef.;

The fat should be firm, white, and free from hæmorrhages; not yellow or gelatinous.

Fat consists of olein, palmitin, and stearin, which are respectively oleic acid, palmitic acid, and stearic acid, combined with glycerine as a base. Stearin, $(C_{18}H_{35}O)_3 \ C_3H_5$ O₃, melts at about 65° C.; palmitin, $(C_{16}H_{31}O)_3 \ C_3H_5$ O₃, at a variable point between 35° C. and 60° C.; olein, $(C_{18}H_{33}O)_3 \ C_3H_5$ O₃, at 5° C. Fats with a high melting point, such as mutton fat, consist mainly of stearin; those with a low melting point, such as bacon fat, principally of olein.

Sausages are liable to convey the same diseases as the meat from which they are made, but there are no longer any macroscopic signs to guide us. In a fatal case at Chester, investigated by Ballard, violent symptoms of gastro-intestinal irritation came on within half an hour of eating sausage, which was found to contain an organic chemical poison. There have since been similar cases. Decomposition may be detected by the smell, which is made more obvious by adding hot water and lime-water.

Preservation of meat may be effected for a

time:

(1) By exclusion of air. The meat is dipped into boiling water so as to form an impervious layer of coagulated albumen on the surface; or coated with paraffin, or simply with fat.

(2) By injection of preservative solutions. The blood-vessels are injected first with water, then with a solution of alum and aluminium chloride—or even

sodium chloride.

(3) By application of preservatives to the surface. The meat may be covered with salt, sugar, boracic acid, boroglyceride, powdered charcoal, weak carbolic acid, or other fixed antiseptic. Or it may be kept in a closed vessel containing volatile oils or antiseptics.

- (4) By pickling. Common salt, with a little potassium nitrate, is rubbed into the meat, or the latter is immersed in strong brine. Water is abstracted, and the salt acts as a preservative.
- (5) By drying. Meat is exposed in somewhat thin layers to dry air, or, still better, to the smoke from a wood fire.
- (6) By continuous exposure to cold (freezing). Meat can be kept in a cold-air chamber for an indefinite period, but decomposes rapidly upon thawing if rigor mortis had set in before freezing. It is better to keep the temperature a little above freezing-point (refrigerated or chilled meat), as actual freezing has certain disadvantages.
- (7) By hermetically sealing in tin cases or other vessels in vacuo, or in sterilised air. Various devices are adopted to sterilise the contents of the tins. Before sealing, air may be drawn off and replaced by nitrogen and sulphurous acid, or by air which has been heated to 500° F., or by steam. Other processes aim at complete exclusion of air, or exclusion of part and removal of oxygen from the remainder by means of sodium sulphite, or at exclusion of all germ life by boiling.

The dangers attending the use of preserved meat are those depending upon the original character of the meat, already referred to, together with risk of putrefaction if the process is imperfectly carried out. Tinned meat sometimes gives rise within twelve hours to symptoms of acute gastro-intestinal irritation, viz. vomiting, purging, cramps, some degree of pyrexia, irregular pulse, and prostration. These ill effects may result from decomposition due to imperfections in the process; and in this case the tin will show signs of pressure in place of the usual vacuum ("blown" tins), and the meat may have an offensive smell and

taste. Not unfrequently, however, no such changes are noticed, but salts of tin, zinc, or lead are found in the meat and jelly, due, no doubt, to the action of sodium chloride or organic acids upon the tin or

solder, perhaps aided by galvanic action.

Cooking brings about coagulation of the myosin and other albuminoids, renders the meat soft and tender by converting the connective tissue into gelatin, and reduces the amount of water and extractives. Boiling in the ordinary way causes a loss of 25 per cent. or more, but most of the salts and soluble substances may be retained by plunging the meat for five minutes into boiling water, and then continuing the cooking at a low temperature. The loss is greater in roasting than in baking, and least in boiling. The various albuminoids coagulate at temperatures ranging from 85° to 170° F., hæmoglobin at 160°. If, therefore, the temperature at any part does not reach 160° the meat is underdone; and if it exceeds 170° the tissues shrink and become hard and indigestible. The best temperature is about 160°. Roasting also entails a loss of about 25 per cent., but mainly of water, with some fat and a little gelatin. It is advisable first to expose meat to an intense heat and afterwards cook it slowly. Ordinary cooking or pickling affords littleprotection in meat infected with specific virus.

Meat inspection.—The effectual inspection of meat really requires (a) veterinary inspection of animals before slaughter, and (b) systematic inspection in some form of clearing-house of all market meat. In the absence of these two requirements, meat inspection must necessarily be incomplete. In some countries there is State inspection, and all marketable meat bears an official stamp. In this country, however, meat inspection rests with the Local Authorities, the larger of which employ meat inspectors specially qualified for the work.

Qualifications of Meat Inspectors.—The L.G.B., in circular letters addressed to local authorities in 1899 and 1901, strongly urged that Meat Inspectors should be directed to act in accordance with the principles laid down by the Royal Commission with regard to the condemnation and seizure of meat affected with tuberculosis (page 107), and, further, that they should possess proper qualifications for the office, including acquaintance with-

(a) The law of meat inspection.

(b) The names and situations of the organs of the body. (c) Signs of health and disease in animals destined for

food, both when alive and after slaughter.

(d) The appearance and character of fresh meat, organs, fat and blood, and the conditions rendering them, or preparations from them, fit or unfit for human food.

Any Inspector of Nuisances or Sanitary Inspector to whomeither of these orders applies is required in any case of doubt in connection with the inspection and seizure of meat to report the matter to the Medical Officer of Health, with the view of obtaining his advice thereon.

Methods of Inspection.—Cursory inspection will frequently mislead, and it is desirable for the meat inspector to perform his duties in a systematic manner. At the slaughter-house or in the wholesale market meat may be examined as follows (written notes being made) :-

- (a) Inspection of the Carcase.—Note in particular the species of animal, the general character of the carcase, size, weight, emaciation, the condition of the bones, the pleura (and whether "stripped" or not), the peritoneum, the kidneys, the superficial and deep glands, the colour and condition of the flesh (to be judged by the freshly cut surface), staining of the peritoneum owing to decomposition in the intestine, signs of dropsy or inflammation, signs of drugging, etc.
- (b) Inspection of the Organs (constituting the

- "offal").—The head; the heart, lungs and liver ("plucks"); spleen, udder, stomach, and intestines, should each be examined if available.
- (c) The Mode of Dressing the Carcase.—Artificial inflation of emaciated tissues by air, the smearing of fat over the surface of the carcase, stripping of the membranes, skilful dressing of the carcase, and evident preparation of it for market, should all be noted. Sometimes methods are adopted to disguise the real condition of meat on the one hand, and, on the other, meat for human consumption is dressed to a marketable standard.
- (d) Miscellaneous.—In addition to the condition of the meat itself, there are other matters not to be neglected by the meat inspector, such as the price marked upon the meat, whether it was exposed for sale, or deposited for the purpose of sale, or deposited for preparation for sale, the hour at which it was inspected (or subsequently seized), the condition and temperature of the weather, etc. All these may become of importance in any subsequent proceedings.

Mode of Proceeding.—An inspector who comes across unwholesome meat should (a) "seize" it, unless it be voluntarily surrendered by the owner taking the initiative, and (b) inform the M.O.H. If the M.O.H. considers the meat unfit for food, (c) it should at once be submitted to a magistrate for condemnation and destruction, the owner being permitted to have the meat seen by experts on his behalf if he so wishes. (d) The case should be fully reported to the S.A., and instructions may be taken as to proceedings. The portions of evidence which may be required in court are as follows:—
(i.) A copy of the summons issued by the S.A. against the owner of the meat; (ii.) the order signed by the magistrate to the effect that, having examined the meat, he had ordered its destruction; (iii.) the resolution of the S.A. or Committee,

signed by the chairman; (iv.) copies of subpoenas; (v.) a clear and concise statement of the inspector's evidence of seizure and the reasons for seizure; (vi.) a statement by the M.O.H. as to the pathological condition of the meat in question and its unfitness for human food; (vii.) collateral evidence of medical or veterinary experts, if such are to be called, and of other persons, salesmen, porters, butchers, slaughterers, etc.

Classification of Unwholesome Meat. — Before forming an opinion as to unwholesomeness of meat, it is necessary to decide as to its quality and condition; that is, whether it is fresh home-grown meat or frozen, chilled, salted, or pickled meat. Having assured himself of the conditions under which the meat is being examined, the next step for the meat inspector is to diagnose the disease with which the meat is affected. Broadly, and for practical purposes of meat inspection, these may be divided into three groups-(a) General pathological conditions; (b) Specific disease; (c) Decomposition. Among the first group are conditions of ordinary degeneration of the tissues and inflammation: fatty infiltration, calcification, and infiltration by pigment, as in melanosis and jaundice; cloudy swelling, fatty degeneration, amyloid disease, gangrene, etc.; fevered flesh and the meat of unbled animals that have died or have been slaughtered when in a moribund condition. The last named is blood-stained, sticky, pits on pressure, and possesses a uræmic smell. It rapidly decomposes, and is heavily charged with ptomaines. There are a variety of inflammatory diseases, some of which are doubtless due to specific causes, and others of which the origin is obscure.

Secondly, there are the specific diseases of animals which cause meat to be unfit for food. These diseases are not necessarily transmissible to man, but they produce in the tissues of the animal suffering from them certain poisonous toxins. It should be remem-

bered that in many cases such products occur early in the disease, and not only when it has reached an advanced stage. Among specific diseases for which meat should be seized as unfit for food are tuberculosis, actinomycosis, glanders, anthrax, swine fever, contagious pleuro-pneumonia, septicæmia, cattle plague, and a large number of parasitical diseases.

Decomposition of meat is, of course, a post-mortem change. The rapidity with which the decomposition sets in depends upon many circumstances external to the meat, such as temperature of the air, place of storage, etc. Colour, consistence, and odour are the three chief signs upon which reliance may be

placed.

Fish are, as a rule, nutritious and digestible but deficient in fat, though eels contain it in abundance, and salmon and herrings also. Fish should be eaten only in season, unbruised, fresh, firm, and free from offensive smell and disease. Unsound fish is dull in appearance, discoloured, and offensive. The least decomposition renders fish unfit for food, and liable to cause gastro-intestinal irritation if eaten. Similar symptoms occasionally follow the use of apparently sound fish. Consumption of the halibut, and probably of other fish also, has been known to cause diarrhœa and vomiting, and within a few hours a vivid red rash may appear over the upper part of the body, followed by profuse desquamation lasting several days. Fish are largely preserved by the means already referred to, and especially by drying, curing, salting, or canning.

Oysters and shell-fish.—Oysters eaten raw are readily digested, but other shell fish are as a rule indigestible and not unfrequently cause dyspepsia and urticaria. Mussels are especially prone to be poisonous, the symptoms including not only dyspepsia and urticaria, but also swelling of the tongue and

fauces, numbness of the limbs, and weak, irregular pulse. Severe and often fatal attacks have been attributed to eating mussels gathered upon shores polluted by sewage. Many of the cases of mussel-poisoning are believed to be due to simple acute dyspepsia, but others have been traced to an alkaloidal poison (mytilotoxine) contained in the liver of the mussel.

Cockles and periwinkles are usually cooked, but mussels are often eaten raw, and if taken from beds to which sewage has access appear occasionally to have

given rise to enteric fever among consumers.

Oysters, under like conditions, have been found to convey enteric fever, and the evidence clearly points to a certain ability to transmit the infection of cholera also. In the spring of 1894 Newsholme attributed a number of cases of enteric fever at Brighton to the consumption of oysters, and in October and November, 1894, a sudden and brief outbreak of enteric fever occurred among the students at Wesleyan University, Connecticut, and was investigated by Conn. In all, 23 of the students were attacked, 13 of the cases being of severe character, and 4 fatal. The other 10 were of very mild type. A consideration of the dates of attack led to the conclusion that infection must have taken place at the time of a series of initiation suppers, and, by exclusion of other possible factors, the one condition common to all sufferers was found to be the eating of raw oysters taken from a certain ground within 100 yards of sewers receiving the drainage of houses in one of which were two persons suffering from enteric fever. Apart from the students, there were a number of guests, two of whom had well-marked enteric fever, and several had less definite diarrheal attacks. Chantemesse has recorded that in June, 1896, fourteen persons living in six houses at Saint-André-de-Sangonis ate some oysters which had been stored in sewage-contaminated water. All of these persons, but no other members of their households, fell ill. Six had severe attacks of enteric fever, one ending fatally, but in the other eight the symptoms were slight. Among other microbes, the bacillus coli was found in oysters grown under like conditions; and Eberth's bacillus, as well as the colon bacillus, was found in the liquor and in the body of the oysters which had been placed in sea-water purposely contaminated with enteric discharges.

In 1896 Bulstrode and Klein reported to the L.G.B. upon Oyster Culture in Relation to Disease. Experiments by Klein gave results which are thus summarised by Thorne In the introduction to this Report:

1. The cholera vibrio, and still more, the typhoid bacillus, are difficult of demonstration in sewage known to have received them.

2. Both these organisms may persist in sea-water tanks for two or more weeks, the typhoid bacillus retaining its characteristics unimpaired, while the cholera vibrio tends to lose them.

3. Oysters from sources which appeared to be free from risk of sewage contamination exhibited none of the bacteria, specific or otherwise, which are commonly regarded as

being concerned with sewage.

4. Oysters from a few out of numerous batches derived from sources where they did appear to be exposed to risk of sewage contamination were found to exhibit colon bacilli—a circumstance which, notwithstanding the comparative universality of this intestinal organism, may be regarded as having some significance by reason of the absence of this bacillus from oysters which appeared to have been exposed to no such risk.

Since that time there have been several considerable outbreaks of enteric fever and gastro-enteritis due to eating infected oysters, including two at Winchester and Southampton in 1902, which Bulstrode traced to eating oysters grown on beds polluted with sewage. He summarised the evidence as follows; Two mayoral banquets occurred on the same

day in separate towns several miles apart. In connection with each banquet there occurred illness of analogous nature, attacking, approximately, the same percentage of guests (62 persons or 46.3 per cent. at Winchester, 55 persons or 41.3 per cent. at Southampton), and at corresponding intervals. At both banquets not every guest partook of oysters, but all those guests who suffered enteric fever, and approximately all those who suffered other illness, did partake of oysters. Oysters derived directly from the same source constituted the only article of food which was common to the guests attacked, and oysters from this source were at the same time and in other places proving themselves competent causes of enteric fever. Klein found B. coli, B. enteritidis sporogenes, and other organisms allied to the Gaertner group of bacilli in oysters from the same layings, and many other bacteriologists have demonstrated the presence of these organisms in oysters derived from polluted beds.

The results of the investigations carried out by Houston for the Royal Commission on Sewage tend to prove that the contamination of oysters by B. coli is widespread and not altogether dependent on sewage contamination. He examined over 1,000 oysters, and nearly all, from whatever laying they were taken, contained B. coli or coliform organisms. This did not hold good as regards deep-sea oysters, which were free from this organism (and spores of B. enteritidis sporogenes), as was also deep-sea water. Houston found the number of B. coli in an oyster varied from 10 to 10,000 (in 10-15 c.c.), and the contents of the stomach of the oyster contained more B. coli than the liquor in the shell. Fewer B. coli were found, as a rule, in oysters stored in pure waters, but instances occurred where the numbers of such bacilli were as great as in oysters from contaminated sources. It has been shown by various workers, most recently by Klein in 1905, that oysters, cockles and mussels have the power of clearing themselves of infective germs, partly by passing them out, and partly by "directly devitalising the microbe."

Such is some of the bacteriological evidence down to recent date, and whilst it may appear to be of a conflicting nature, there are certain conclusions which may be drawn. First, the number as well as the presence of B. coli in oysters must be considered. Secondly, topographical evidence as to pollution must be taken in conjunction with bacterial evidence, and, at present, the former is more reliable than the latter. Thirdly, oysters ordinarily grown on contaminated oyster-beds may, and do on occasion, contain the virulent specific bacillus of typhoid, which can live both in sea-water and within the shell of the oyster. This being so, the risk of typhoid infection from oysters is a real one, although in actual occurrence many conditions have to be fulfilled.

Preventive Measures.—Present legislation does not give to local authorities any power of inspection or control over oyster-layings or shell-fish beds, nor any powers to prevent the sale in their district of oysters and other molluses derived from sewage-contaminated sources, nor have they powers to control the importation of foreign oysters from whatever source. Oysters fattened on contaminated beds should be deposited for a time in pure sea-water before consumption. They should on no account be stored in unclean water.

Other shell-fish than oysters from time to time cause epidemics or individual cases of gastro-intestinal irritation, and probably contain germs acquired from their food, which is frequently filth. In a report to the **L.G.B.** on "Cockles as Agents of Infectious Diseases," Klein recorded finding *B. coli* in 3 out of 8 cockles which had been taken from a

foreshore polluted with the discharge from a sewer outfall, and also B. enteritidis sporogenes in 4 of them. No typhoid bacilli were detected. In 8 raw cockles in their shells bought from a street hawker, he found no typhoid organisms, but B. coli was found in 5 out of the 8 cockles and B. enteritidis sporogenes in 4 out of the 8. In subsequent experiments Klein came to the conclusion that a mussel immersed for 22 hours in cholera-infected water retained the bacilli of cholera for 48 hours after immersion in clean sea-water, and the same may be said in respect of typhoid infection. Indeed, evidence was obtained showing that the typhoid bacillus could multiply in cockles. Merely pouring boiling water over a heap of shell-fish, as is customary, does not necessarily destroy either cholera or typhoid infection contained in them. Steam sterilisation or boiling is necessary. Since the time of Klein's report, a number of outbreaks of disease, including enteric fever, have been traced to the consumption of mussels and cockles, and it has been shown that the cooking which these shell-fish generally undergo is not sufficient to free them from risk.

Milk contains nitrogenous substances, fat, carbohydrates, salts, and water, and may be regarded, in a sense, as a complete diet in itself. The chemical composition varies somewhat according to the breed of animal, its age, the period since calving, and the diet

and physical conditions.

the sale translate the sales of	Milk of Average Country- fed Cows.	Milk of Average Town-fed Cows.	Alderney Milk.	Fore Milk.
Water	88 %	86 %	87 % 3·2	90 %
Proteid (Casein, etc.) Milk Sugar	4 4.5	5 4·3	4·5 4·1	} 9.7
Salts (Ash)	0.7	0.7	0.7	0.7

Or the average milk constitution may be expressed thus:—

Fat .				4·1 p	er cent.
Solids no	t fat	14.5 18	100	8.8	"
To	otal solids	1		12.9	,,
Water				87.1	,,

Milks vary in standard, and the above figures can only be taken as fair averages.

Milk-sugar, or Lactose (C₁₂H₂₄O₁₂), is a constant constituent of milk, forms the chief substance in solution in whey or serum, and is a member of the cane-sugar group. Is found in varying quantities in the milk of mammals. About 5 per cent. is present in human milk, and somewhat less in that of the cow. It is very resistant to fermentation by yeast, and therefore undergoes alcoholic fermentation very slowly. It is not acted upon by rennet, pepsin, or trypsin. But of all the

sugars it is most readily acted upon by micro-organisms.

Fat occurs in milk as suspended globules of varying size. It forms the cream, and by churning is, of course, made into butter, though both cream and butter contain other constituents besides fat. Lloyd has shown that it is the large globules that form the cream, and he has also made observations upon the size of fat globules in relation to breed of cattle. The decomposition and breaking down of milk-fat by fermentation is the chief cause of gross abnormalities of cream and the rancidity of butter. The proportion of cream averages about 8 per cent., but ranges from 6 per cent. in very poor milk to 12 per cent. or 15 per cent. or more; in Alderney cows it may reach 40 per cent. One part of cream is said to correspond roughly to 0.2 part of fat. More complete separation of cream is effected by centrifugalisation in a "separator." Skimmed milk retains as a rule about 1 per cent. or more of fat; separated milk much less. Both contain the casein, sugar, and salts of the milk from which they were made, and have therefore considerable nutritive value. "Fore-milk," which is the first part of the vield of milk from the udder, contains but little fat; the "strippings," on the other hand, at the end of milking, are rich in fat.

The Proteids of Milk include casein, lactalbumin, and lactoglobulin. Casein is the most abundant and important. When milk separates naturally into its constituent parts, the fat rises and the casein falls, leaving a clear fluid, the milk plasma or

serum, between the two substances.

Mineral Matter.-The ash of milk, obtained by careful ignition of the solids, contains calcium, magnesium, potassium, sodium, phosphoric acid, sulphuric acid, chlorine, and ironphosphoric acid and lime being present in the largest amounts.

Human milk contains less fat, less proteid, and more sugar than cow's milk (e.g., an analysis of average human milk would result in total solids, 12.5; fat, 3.7; proteids, 2.3; lactose, 6.2; and ash, 0.3), and therefore to modify cow's milk so as to bring it approximately to the standard of human milk, it is necessary to dilute it and add milk-sugar. Much debate has arisen as to what should be accepted as the minimum standard for the various constituents of milk. Broadly it is now usual to fix the fat standard at 2.75-3 per cent., and the "solids not fat" at 8.5 per cent.

The following are the recommendations (1901) of

the majority of a Departmental Committee:

1.—That Regulations under s. 4 of the Food and Drugs Act, 1899, be made by the Board of Agriculture with respect

to milk (including condensed milk) and cream.

2.—(a) That in the case of any milk (other than skimmed, separated, or condensed milk) the total milk solids in which, on being dried at 100° C., do not amount to 12 per cent., a presumption shall be raised, until the contrary is proved, that the milk is deficient in the normal constituents of genuine milk.

(b) That any milk (other than skimmed, separated, or condensed milk) the total milk solids in which are less than 12 per cent., and in which the amount of milk fat is less than 3.25 per cent., shall be deemed to be so deficient in milk fat as to raise a presumption that it has been mixed with separated milk or water, or that some portion of its normal content of milk fat has been removed. In calculating the percentage amount of deficiency of fat the analyst shall have regard to the above-named limit of 3.25 per cent. of milk fat.

(c) That any milk (other than skimmed, separated, or condensed milk) the total milk solids in which are less than 12 per cent., and in which the non-fatty milk solids are less than 8.5 per cent., shall be deemed to be so deficient in normal constituents as to raise a presumption that it has been mixed 3.—That the artificial thickening of cream by any addition of gelatine or other substance shall raise a presumption that

the cream is not genuine.

4.—That any skimmed or separated milk in which the total milk solids are less than 9 per cent. shall be deemed to be so deficient in normal constituents as to raise a presumption that it has been mixed with water.

5.—That any condensed milk (other than that labelled "machine-skimmed milk" or "skimmed milk," in conformity with s. 11 of the Food and Drugs Act, 1899) in which either the amount of milk fat is less than 10 per cent. or the amount of non-fatty milk solids is less than 25 per cent. shall be deemed to be so deficient in some of the normal constituents of milk as to raise a presumption that it is not genuine.

The following expressions of opinion are added: -

(a) That it is desirable to call the attention of those engaged in the administration of the Food and Drugs Acts to the necessity of adopting effective measures to prevent any addition of water, separated or condensed milk, or other extraneous matter, for the purpose of reducing the quality of genuine milk to any limits fixed by regulation of the Board of Agriculture.

(b) That it is desirable that steps should be taken with the view of identifying or "ear-marking" separated milk by the addition of some suitable and innocuous substance, and by the adoption of procedure similar to that provided by s. 7 of

the Food and Drugs Act, 1899, in regard to margarine.

(c) That it is desirable that, so far as it may be found practicable, the procedure adopted in collecting, forwarding, and retaining, pending examination, samples of milk (including condensed milk) and cream under the Food and Drugs Acts should be uniform.

(d) That it is desirable that, so far as may be found practicable, the methods of analysis used in the examination of samples of milk (including condensed milk) or cream taken

under the Food and Drugs Acts should be uniform.

(e) That it is desirable in the case of condensed milk (other than that labelled "machine-skimmed milk" or "skimmed milk," in conformity with s. 11 of the Food and Drugs Act, 1899) that the label should state the amount of dilution required to make the proportion of milk fat equal to that

found in uncondensed milk containing not less than 3.25 per cent. of milk fat.

(f) That it is desirable in the case of condensed whole milk to limit, and in the case of machine-skimmed milk to exclude, the use of sugar.

(q) That official standardising of the measuring vessels

commercially used in the testing of milk is desirable.

The Sale of Milk Regulations of 1901 which followed this report lay down that (a) where a sample of milk contains less than 3 per cent. of milk-fat, it shall be presumed until the contrary is proved that the milk is not genuine by reason of the abstraction therefrom of milk fat or the addition thereto of water; (b) where a sample of milk contains less than 8.5 per cent. of "solids not fat," it shall be presumed until the contrary is proved that the milk is not genuine by reason of the abstraction therefrom of "solids not fat," or the addition thereto of water; and (c) when a sample of skimmed or separated milk (not being condensed milk) contains less than 9 per cent. of solids (including fat), it shall be presumed that the milk is not genuine. These regulations are in force throughout Great Britain.

Adulterations.—Addition of water and abstraction of cream are by far the most frequent frauds. Sodium carbonate, sodium chloride, starch, borax (1-2,000), formalin (1-50,000), or salicylic acid are occasionally added, either as preservatives or in order to mislead the analyst. Skimmed milk or separated milk is frequently fortified by the addition of condensed milk, and whole milk is not unfrequently diluted with separated milk. A good deal of "manipulation" of milk goes on, mixing of poor with rich, and of separated and condensed milks with water and with whole milk. Annatto and turmeric are often added to impart a yellow colour to milk.

Addition of water lowers the specific gravity, the

percentage of fat, "solids not fat," and salts; it also makes the cream rise more readily to the surface. Abstraction of fat, by skimming or by the use of the separator, increases the specific gravity and the percentage of salts and "solids not fat," but of course reduces the fat and therefore the yield of cream. The addition of condensed milk to skimmed milk may give an appearance of extreme richness, but the salts and "solids not fat" will be in excess.

Owing to the social changes by which an increasing percentage of the population lives in towns, the practice of adding preservatives to milk, with a view to "keeping" it over the necessary transit and delay in reaching the consumer, is increasing.

In 1901 another Departmental Committee appointed by the Board of Agriculture recommended:

1. That the use of formaldehyde in food and drink be absolutely prohibited, and that salicylic acid be not used in greater proportion than one grain per pint or pound respectively for liquid or solid food, its presence in all cases to be declared.

2. That the use of any preservatives or colouring matter, in milk be made an offence under the Sale of Food and Drugs

3. That boric acid preservatives only be allowed in cream the amount not to exceed 0.25 per cent, and be notified on a

4. That boric acid preservatives only be allowed in butter,

the amount not to exceed 0.5 per cent.

5. That chemical preservatives be prohibited in all dietetic preparations for the use of children and invalids.

6. That the use of copper salts for "greening" be pro-

hibited.

7. That a Court of Reference be established to supervise the use of preservatives and colouring matters in food.

Bacteria in milk.—Milk as it leaves the healthy udder is a sterile fluid, but at the time of milking, during transit to the dairy, at the milkshop, and in the home of the consumer, it becomes more or less contaminated. Being a favourable medium for the growth

of bacteria, and being frequently kept for some time at a favourable temperature for the multiplication of germs, milk becomes, as a rule, heavily impregnated with bacteria. Disease of the cow's udder when present would add a further quota. Thus it comes about that milk is often found to contain millions of bacteria per cubic centimetre, varying in number according to cleanliness or otherwise, treatment, temperature, and storage. The species of bacteria found in milk may be roughly classified as: (a) ordinary bacteria of soil and water; (b) bacteria of sewage, manure, or intestinal origin; (c) bacteria of fermentation, and (d) pathogenic organisms associated with tuberculosis, enteric fever, scarlet fever, diphtheria, sore throat and epidemic diarrhea. Here it is only necessary to consider the last two groups of bacteria.

Milk-sugar undergoes fermentation like other sugars upon the introduction of a suitable fermenting agent. There are five different kinds of such fermentation which affect milk, namely, lactic acid and butyric acid fermentations, coagulation fermentation without acid-production, alcoholic fermentation, and, lastly, a group of anomalous changes of various kinds. Lactic acid fermentation is the ordinary souring of milk. The coagulated casein is precipitated, the serum rises and carries up lumps of fat, and the whole becomes acid. This change is due to a group of bacilli (B. acidi lactici), the first member of which was discovered by Pasteur and isolated by Lister. They gain access to milk from external sources, particular kinds being found in the vicinity of particular dairies, soils, geographical localities, etc., thus giving to milk and butter characteristic flavours.

The lactic fermentation bacteria are short rods, do not liquefy gelatine, nor do they form spores. They grow readily on gelatine at room temperature, forming as a rule small circular colonies, white or grey in

colour, with sometimes a tinge of yellow, and the surface of the colony is smooth and glistening. The lactic acid organisms produce appreciable amounts of lactic acid only at somewhat elevated temperatures. If the amount of acid rises much above 2 per cent., the growth of the lactic acid bacteria is inhibited. Many other substances are produced in addition to lactic acid (e.g. acetic and formic acids, alcohol, methane, CO2, etc.). Only provisional classifications are possible at present, as, owing to variations in biology and terminology, it is probable that certain lactic organisms are described under several different terms. Generally, it may be said that some grow well in the presence of oxygen, and others do not. The most common members of the group are the facultative anærobes which sour milk best in deep vessels, and produce a right-handed lactic acid. They are widely distributed in nature, and may form 90 per cent. of the total bacteria in milk. Some produce gas, others liquefy gelatine, and some produce spores.

Butyric acid fermentation, due to the B. butyricus, is a bitter fermentation in which hydrogen and carbonic acid gas are produced. It has little of the importance of lactic fermentation or even of alcoholic fermentation by yeasts, by which "koumiss" is prepared from mare's milk and "kephir" from the milk of goats, sheep, and cows. Other fermentations of milk are represented in (a) bitter fermentations (due to Weigmann's bacillus or Conn's micrococcus); (b) slimy fermentations which produce the "ropiness" used in the manufacture of Edam cheese and teetmelk (due to Micrococcus viscosus and Streptococcus hollandicus, etc.); (c) soapy milk caused by B. lactis saponacei, and (d) a number of chromogenic fermentations.*

Pathogenic organisms gain access to milk both

^{*} Red milk (B. prodigiosus and B. lactis erythrogenes); blue milk (B. cyanogenes); yellow milk (B. synxanthus); etc.

from the cow and from sources external to milk. The tubercle bacillus is the chief example of the former channel. Probably not less than 10 to 15 per cent, of the milch cows of this country are affected with tuberculosis, and about 2 per cent. with tuberculous disease of the udder.* The milk is most infective when the udder is diseased. From 1896 to 1906 it was found that 8.0 per cent. of the country milk brought into Liverpool was tuberculous, and percentages varying from 5 to 16 have been frequently reported in market milk in this country. Whilst tuberculosis is not commonly spread among human beings by means of an infected milk supply, there is evidence, derived from pathological and clinical experience, proving that tuberculous milk can and does, on occasion, set up some form of tuberculosis, human or bovine, in the bodies of man and other animals consuming the milk. Phthisical milkmen may introduce the human tubercle bacillus to milk. All danger may be obviated by boiling the milk. The typhoid bacillus has occasionally found its way from sewage, water, or infective dust into milk, which is a favourable medium for its multiplication. The same may be said of the diphtheria bacillus, which, in four cases on record, has been actually isolated from milk suspected of conveying the disease (Bowhill, Eyre, Klein, Dean and Todd). Some authorities now hold that diphtheria is a disease of the cow and that the bacillus sometimes gains access to milk from the udder. Various forms of streptococcus have also been isolated from milk. Their exact pathological significance is unknown.

Examination of milk.

1. Specific gravity of milk is measured by a lactometer (which is a delicate hydrometer) or by a Westphal balance.

^{*} There were 4,102,000 milch cows in the United Kingdom in 1901. A cow yields on an average about 400 gallons of milk per annum,

The specific gravity averages about 1030-1032. It is lowered

by watering, and raised by abstraction of fat.

2. Total solids, which range from 12 to 13 per cent., are estimated by evaporating a small quantity of the milk in a platinum dish, weighing before and after drying (weight of residue × 100 divided by weight of milk = percentage total solids). By deducting the amount of fat, the solids not fat are obtained. The ash of milk, usually about 0.7 per cent., is of

little value as an indication of quality.

3. Fat may be determined (a) by mechanical methods such as centrifugalisation, or (b) by extraction of the fat by ether, with or without preliminary destruction of the caseine by acid. In Adams' method five c.c. of the milk are spotted on to a porous paper, which is thoroughly dried, and then rolled up and extracted with ether in a Soxhlet apparatus. The ether is syphoned off and evaporated, and when a constant weight is obtained the quantity of fat is determined. Werner-Schmidt's method is extraction of fat with ether, after destruction of casein with strong hydrochloric acid.

The fat in unadulterated milk is never less than 3.0 per cent., unless the "fore milk" alone is taken, so that when skimming as well as watering is suspected, the ratio 3.0: 8.5 may be taken as the minimum proportion of fat to "solids not fat," and if the "solids not fat" are x, the fat should be at least $\frac{3\cdot 0}{8\cdot 5}x$, and if the ascertained amount y of fat falls short of this,

it is concluded that at least $100 \times \left(\frac{3.0}{8.5}x - y\right) \div \frac{3.0}{8.5}x$ per

cent. of fat has been abstracted.

4. Water.—The results of many thousands of analyses show that, although the fat varies greatly, the percentage of "solids not fat" in unadulterated milk falls below 8.5 very rarely. Hence 8.5 per cent. is adopted as a standard. If a given milk contains x per cent. "solids not fat," and x is less than 8.5, we can affirm that however poor the milk might have been originally, it must contain added

water to the extent of at least $\left(\frac{8.5-x}{8.5}\times 100\right)$ per cent. of

the sample; or, in other words, it is $\left(\frac{8.5-x}{8.5}\times 100\right)$ per cent.

worse than genuine milk of poorest quality. If it was originally milk of fair quality, the adulteration must have been much greater.

5. Preservatives in milk. - Boracic acid is detected by moistening the ash with alcohol and H₂SO₄, and applying a light to the mixture when so little as 1 part in 10,000 will give a green flame; or the ash may be treated with dilute HCl, and turmeric paper will be turned brown, and if touched with sodium carbonate, bluish-black. Salicylic acid is detected by curdling the milk with mercuric nitrate, and then, after shaking it up with ether, evaporating, and moistening the residue with ferric chloride. A blue coloration will denote the presence of salicylic acid. Formalin in milk gives a purple colour if warmed with hydrochloric acid containing a trace of iron.

6. Bacteriological examination is somewhat similar to that of water. Plate cultivations are made, etc., but it is usually necessary to dilute the milk with sterile water. Special methods of examination have to be adopted in searching for pathogenic organisms. Generally it may be said that by the centrifuge or sedimentation flask the particulate matter is obtained and examined, and 2 c.c. are inoculated into a susceptible animal. For reliable examination for the tubercle bacillus it is necessary to inoculate animals with the milk, as many bacilli simulate the tubercle bacillus in almost every particular, except its pathogenic effect. Special staining and culture tests are necessary for B. diphtheriæ, streptococcus, B. Coli, and B. enteritidis sporogenes. The number and kind of bacteria present should be determined, the degree of acidity, and the presence or absence of pus or blood cells, yeast cells, dirt, etc.

Milk in relation to disease.—From the point of view of preventive medicine the chief milk-borne diseases are enteric fever, diphtheria, scarlet fever, sore-throat illnesses, and epidemic Probably tuberculosis is not commonly spread by milk, and only one or two outbreaks of milkborne cholera are on record. Whether other infectious diseases, such as measles, whooping cough and smallpox, are ever conveyed by milk remains to be proved.

An inflammatory affection of the udder, known as garget, has been suspected of causing outbreaks of sore throat, etc., in those using the milk. Milk from cows suffering from foot-and-mouth disease is capable of producing illness with aphthous stomatitis, swelling of the tongue, and fœtor of the breath, especially in children; in other cases a severe form of sore throat has been the chief feature. Similar symptoms may attend the consumption of milk from cows affected with inflammation or abscess of the udder. The evidence is less clear in regard to other specific diseases to which cows are subject, and, indeed, in most of them the secretion is suppressed, but the milk should never be used for food. In the Western States of America cows are subject to a disease called the "trembles," supposed to be due to their eating Rhus toxicodendron, and their milk appears to cause in children vomiting, constipation, swelling of the tongue, and prostration, with abnormally low temperature. The milk of goats that had fed on Euphorbium or Meadow Saffron has been known to induce severe diarrhœa. Even in the human subject bitter and purgative drugs, not to mention morphia and other alkaloids, if taken by the mother, act upon the infant through the milk. Colostrum, the milk yielded by cows immediately after calving, has also been shown to exert a deleterious effect on persons consuming it. Lastly, disease has been caused, it is believed, by yeast cells in milk. Such an outbreak occurred at Lincoln in 1902 (199 cases). The symptoms of the disease simulated scarlet fever. There was much swelling of the tonsils and cervical glands with pyrexia, gastritis, and "rheumatism." Klein isolated what he believes to be a pathogenic yeast possibly bearing a relationship to the spores of rusts, smuts, and mushroom fungi consumed by the cows yielding the milk.

Enteric Fever .- In 1857 Michael Taylor, of Penrith, first established the fact that milk may convey enteric fever. Since that date more than 150 epidemics of this disease have been traced to an infected milk supply. Schuder states that 17 per cent. of all typhoid epidemics are thus caused. The channels of infection

by which the bacillus obtains access to milk have been shown by a study of these outbreaks to be from direct contact with typhoid patients, from washing milk vessels with infected water, from the addition of such water to the milk, from the air by dried typhoid excreta, from contaminated cloths and clothes, and

from cooling milk in infected water.

Diphtheria.—The first epidemic of diphtheria which was shown to be milk-borne was one in North London in 1878, elucidated by Power. More than 30 similarly caused outbreaks of the disease have since been recorded. How the diphtheria bacillus gains access to milk is not known, but in all probability it is generally by direct contact. Persons suffering from the disease may remain for a long period infectious by reason of the retention for weeks after convalescence of the bacillus in the throat. Materials may also retain the infection for long periods. Water and sewage do not appear to convey the diphtheria bacillus. Klein holds that the Klebs-Löffler bacillus is inoculable in cows, producing a specific eruption on the udder, and within five days the bacillus is excreted in the cow's milk. Several outbreaks of diphtheria have suggested that bovine diphtheria is transmissible to man by milk.

Scarlet Fever. - More than 70 milk-borne attacks of scarlet fever are on record, and yet comparatively little is known either of the bacteriology of the disease or of the mode of infection of milk. Direct contact of scarlet fever patients with the milk trade is the most obvious channel of infection. In 1882 Power found reason to suspect that cows suffered from a disease which could impart to milk the virus of scarlet fever, and in 1885 in the "Hendon disease" outbreak further evidence of this was met with.

In that year a sudden and extensive outbreak of scarlet fever occurred in Marylebone, and was found to be associated with a particular milk supply derived from a farm at Hendon. The milk was also distributed in St. Pancras, Hampstead, Hendon, and St. John's Wood; in each of these districts, except the last, scarlet fever suddenly became prevalent early in December. On the 15th the milk sent to Marylebone was returned to the farmer, and some of this was given away to persons at Hendon on the 15th and 16th; a few days later, from December 20th onwards, a number of cases of scarlet fever occurred among those who had drunk the milk, and at the same time there was a sudden decrease in the number of attacks in Marylebone. The disease was conveyed by the milk, and after close investigation Power and Klein were led to the important conclusion that the cow itself was the source of infection.

There had been apparently no case of scarlet fever among the employes that could be proved to have infected the milk. Attention was next directed to the cows, and many of them were found to be suffering, or to have recently suffered, from vesicles or ulcers upon the teats and udder. These were clearly infectious, and had been first seen upon a cow that was bought on November 15th. The dates of outbreak of scarlet fever in each district being known, it was found that each outbreak was preceded by a few days by the introduction of this affection into the cowsheds from which the milk supply of the district was drawn. The early exemption of St. John's Wood was explained by the fact that the disease had not appeared in the small shed from which alone its supply was drawn; but during the inquiry this shed became affected at last, and an outbreak in St. John's Wood immediately followed. All the cows showing any signs of the disease were then isolated, and no further cases of scarlet fever occurred among the consumers of the milk. The symptoms noticed in the cow were chiefly local, but there were bald patches of skin, especially about the tail and back, the epidermis in these patches being scaly and the cutis thickened. There was no pyrexia. The vesicles, which were small, were confined to the teats and udder. They extended, and in two days formed flat irregular ulcers covered with brown scabs. Inoculated upon calves, the matter from these ulcers caused local tenderness and swelling in three days, a scabbed ulcer with vesicular margin in six days, and a further extension during the next few days, followed by healing. By cultivation a streptococcus was obtained, supposed to be identical with a similar organism found by Klein in connection with scarlet fever, and having the property of solidifying milk if kept for two days at 35° C. It was not found in the milk from unaffected teats. Inoculation of calves with pure cultivations of the streptococcus produced a constitutional disease that had some points of analogy with scarlet fever, the condition of the kidneys especially differing but little from acute scarlatinal nephritis.

Sore-throat illnesses. Somewhat allied to scarlet fever, and possibly in some instances an aberrant form of that disease, is sore-throat infection conveyed by milk. There are fully a score of wellmarked epidemics of this nature on record. One of the first to be investigated occurred in South Kensington in 1875. About 150 persons were affected after consuming cream. A similar outbreak occurred in Edinburgh in 1888. More recent outbreaks of milk-borne sore-throat occurred at Bedford in 1902, Woking 1903, Finchley 1904, and Colchester 1905. Cream was again the particular medium at Bedford, where the incidence was chiefly upon young adults. At Woking 98 families were affected, and the milk was traced to cows suffering from suppurative mammitis. The Finchley outbreak involved 500 cases. The incubation period, as in many of these sore-throat conditions, was short—24 to 48 hours. There was enlargement of the submaxillary glands, sore-throat, fever, general malaise, and in a few cases a measleslike eruption on the lower limbs. On some occasions gastro-intestinal irritation has also been present. It is possible that these sore-throat illnesses are of streptococcal origin.

In many milk-borne outbreaks of scarlet fever there occur a number of sore-throat cases simulating, but possibly not identical with, scarlet fever, in addition to those which are milder forms of scarlet fever

itself.

Epidemic diarrhea is another infective disease which appears to be conveyed by milk. Certain attacks of the disease give support to this view, and there is also the autumnal rise of infantile diarrhoea, which is

attributed by many observers, in part, to contaminated milk. It is held that the high atmospheric temperature of the third quarter of the year exerts a prejudicial effect on milk, causing it to be injurious, particularly to infants and children. It is certain that breast-fed infants suffer much less than artificially fed infants from such disease. Actual epidemics of milk-borne diarrhea have been, it appears, comparatively few. In 1894 such an outbreak occurred in Manchester, affecting 160 persons, and due probably to milk polluted with excreta; and in 1895 and 1898 three outbreaks occurred at St. Bartholomew's Hospital, traced to the consumption of milk or milkpudding from which Klein isolated the B. enteritidis sporogenes, which he regards as the vera causa of epidemic diarrhœa.

Characteristics of milk-borne epidemics.—There are certain general characteristics of outbreaks of disease transmitted by milk which it may be well briefly to

state here.

1. The outbreak is usually sudden in onset; and the cessation often almost equally so, if allowance is made for secondary cases in infected households, which have probably been infected from the earlier cases and not by the milk.

2. A large proportion of the attacks are simultaneous or nearly so. The outbreak reaches its maximum too rapidly to admit of satisfactory explanation by the hypothesis of infection from the first cases; thus, in an outbreak of scarlet fever at Wimbledon in 1886–7, the daily number of attacks was as follows:—

Date . Dec. 25 26 27 28 29 30 31 Jan. 1 2 3 Attacks . 1 11 39 53 75 118 65 86 25 15

3. It will often happen that two or more persons in the same household are taken ill at the same

time. Such multiple cases may also occur apart from milk or water infection, but it is very exceptional, especially as regards the first invasion of the household.

4. The average number of cases per infected household is usually greater than occurs under ordinary conditions, but this is of course dependent upon the number of persons who consume the milk. An average of two attacks per household may be considered high.

5. A large proportion of the households attacked have a common milk supply, which, however, may

not be distributed by the same retailer.

6. Conversely, if (a) the number of households supplied by the suspected dairy be ascertained, and also (b) the number of such households attacked, it will be found that the proportion of (b) to (a) is much greater than the proportion which the total number of infected households bears to the total number of inhabited houses in the district. Although in some cases 50 per cent. or more of the houses supplied with the implicated milk are attacked, in other instances the proportion is far lower.

7. If the households supplied by the dairy in question are classified according to the quantity of milk bought daily, the invasions are found to be proportionately more numerous among households

taking a larger supply.

8. A similar calculation, substituting individuals for households, will give similar results. Classifying the consumers according to the average consumption per head in the households to which they belong, a heavier incidence upon the larger consumers will be manifest. Hence the wealthier consumers, upon the average, suffer more than the poorer class in milk-borne outbreaks, and a classification according to rental will usually confirm this.

The Wimbledon outbreak, already referred to, illustrates these relations (Cooper):-

Incidence on Consumers, according to Supply per Head.		Incidence Houses, acco Supply per	rding to	Incidence on Houses, according to Rental.			
Pints.	Percentage of Persons supplied attacked.	Pints.	Percentage of Houses supplied invaded.	Rateable Value of Houses supplied.	Percentage of Houses supplied invaded.		
·05 to ·1	14	.25	40	£10 to £12	21		
·1 to ·2	23	.5	40	£13 to £15	39		
·2 to ·3	41	.75	50	£16 to £20	50		
·3 to ·4	39	1.0	49	£21 to £25	64		
·4 to ·5	46	1.5	58	£26 to £30	66		
·5 to ·6	57	2.0	70	£31 to £35	57		
·6 to ·7	40	3.0	67	£36 to £40	76		
·7 to ·8	46	3.5	74	£41 to £50	73		
Over ·8	49	Over 3.5	78	£51 to £60	74		
OF THE PARTY	11/11/11	STATE OF THE PARTY NAMED IN	120000000000000000000000000000000000000	£61 upwards	80		

- 9. Among persons who drink little or no milk, or only take it in tea or coffee, or always have it boiled, attacks will be rare, except after exposure to infection from earlier cases.
- 10. As a corollary to this, there is often a much heavier incidence among children and women than among men. This would be especially significant in regard to enteric fever, which under ordinary conditions chiefly affects adults. On the other hand, Littlejohn has pointed out that in milk-borne scarlet fever an exceptionally high proportion of adults may suffer.
 - 11. The type of disease in milk-borne epidemics has

been noticed to be, as a rule, mild, and attended with

low mortality.

12. The incubation period is shortened, and there is some reason to suppose that the degree of infectiousness by contact is lower than when such diseases are not milk-borne.

Ice-cream or iced-custards which have as their basis milk or cream mixed with other ingredients, may be briefly considered here in their relation to infection. A number of outbreaks of typhoid or diarrhoeal disease have been attributed to the consumption of ice-cream. Such outbreaks occurred at Deptford in 1891, Liverpool 1897, City of London and Finsbury 1902, and Paisley 1904. The ice-cream may become infected owing to storage in infected premises, between the process of "boiling" and that of "freezing," or owing to the fact that the substance has been made by a person suffering from recognised or unrecognised disease. It is on this account that, since 1895, local regulations have been in force with the view of controlling the manufacture of ice-cream. Some of the most recent of such regulations are contained in the London County Council (General Powers) Act, 1902, under which ice-cream must be made and stored in sanitary premises, not living rooms, precautions must be taken as to protection from contamination, and the name and address of the maker must appear on the street-barrows. Similar powers are conferred by local Acts in Liverpool, Glasgow, etc.

Control of milk supplies.—The requirement is a pure milk supply, or in other words a clean, whole milk, unsophisticated and unadulterated, (a) derived from healthy cows, guaranteed free from tuberculosis by the tuberculin test, and living under clean and sanitary conditions; (b) obtained by clean methods of milking, strained, and protected from contamination and from infection; (c) kept cool by means of refrigeration; and (d) not exposed to dust or uncleanliness in any way from the vessel in which it is placed, or from the persons by whom it is handled.

Low temperatures do not readily destroy germ life,

but they have a most beneficial effect upon the keeping quality of milk. If the temperature be lowered sufficiently, the contained bacteria become inactive and torpid, and eventually are unable to multiply or produce their characteristic fermentations. At about 50° F. (10° C.) the activity ceases, and at a temperature of about 40° F. organisms are practically deprived of their injurious powers. If the milk is to be conveyed long distances, then even a lower temperature is desirable. The most important point with regard to the cooling of milk is that it should take place immediately. Various kinds of apparatus are effective in accomplishing this. Perhaps those best known are Lawrence's cooler and Pfeiffer's cooler, the advantage of the latter being that during the process the milk is not exposed to the air. It must not be forgotten that cooling processes are not sterilising processes, and do not necessarily kill bacteria; they only inhibit activity, and under favourable circumstances torpid pathogenic bacteria may again acquire their injurious faculties. Hence, during the cooling of milk even greater care must be taken to prevent aërial contamination than is necessary during the process of sterilising milk. No cooling whatever should be attempted in the cowshed. Climate makes little or no difference to the practical desirability of cooling milk, yet it is obvious that less cooling will be required in the cold season. The further treatment of milk has in practice comprised the addition of preservatives, filtration, and sterilisation, but with first-rate clean milk, consumed or bottled immediately after yield, these are not necessary.

Preservation of milk may be effected by boiling, by exclusion of air, or by the addition of sugar, borax (1-2,000), formalin (1-50,000), or other antiseptic. Such preservatives are frequently added to milk, especially in towns. They prevent rapid

multiplication of germs, and so delay or prevent fermentation. On the other hand, they disguise the true condition of the milk, and by cumulative process may cause poisoning in the consumers, and therefore should be condemned. The L.G.B. (1906) advise taking action if boron preservatives are present in amount exceeding 40 grains per gallon. Condensed milk is prepared by evaporating down in vacuo at a low temperature to about one-third or one-quarter of the original volume, and most manufacturers then add cane sugar as a preservative. The added sugar renders it too carbonaceous and "fattening," though lacking in fat, but apart from this the nutritive value of condensed milk seems to be inferior to that of fresh milk, especially in regard to bone-formation and stamina. Condensed milk is, as a rule, fairly free from suspicion of ordinary adulteration or pollution, but it is by no means always sterile.

Filtration of milk is practised in Denmark, but is

little adopted in this country.

Sterilisation means the use of heat at or above boiling-point, or boiling under pressure. This may be applied in one spell of one to two hours at 212-250° F., or at stated intervals at a lower temperature. The milk is sterilised—that is to say, contains no living germs—is altered in chemical composition, and possesses a flavour which to many people is unpalatable.

Such a radical alteration is not necessary in order to secure non-infective milk. The bacteria causing the diseases conveyable by milk succumb at much lower temperatures than the boiling-point. Advantage is taken of this in the process known as pasteurisation, in which milk is heated to 167-185° F. (75-85° C.). Such a temperature kills harmful microbes, because 75° C. is above their average thermal death-point and yet the physical changes in the milk are practically

nil, because 85° C. does not relatively approach the boiling-point. There is no fixed standard for pasteurisation, except that it must be above the thermal death-point, and below the boiling-point. As a matter of fact, 158° F. (70° C.) will kill lactic acid bacteria and most disease-producing organisms found in milk. If milk is kept at that temperature for ten or fifteen minutes it has been "pasteurised." If it has been boiled, with or without pressure, for half an hour, it has been "sterilised." The only practical difference in the result is that sterilised milks have a better keeping quality than pasteurised, for the simple reason that in the latter some living germs have been unaffected.

Sterilisation may, of course, be carried out in a variety of modifications of the two chief ways above named. When the process is to be completed in one event an autoclave is used, in order to obtain increased pressure and a higher temperature. Milk so treated is physically changed in greater degree than in the slower process. The slow or intermittent method is, of course, based on Tyndall's discovery, that actively growing bacteria are more easily killed than their spores. The first sterilisation kills the bacteria, but leaves their spores. By the time of the second application the spores have developed into bacteria, which in turn are killed before they can sporulate.

The application of pasteurisation to milk is now very widely adopted by corporations, dairy companies,

hospitals, milk-dispensaries, etc.

Butter.—When cream is violently agitated, the fat globules are ruptured and their contents clot together as butter. The water is expelled by pressure, and salt is added. The flavour of butter is due to the butyrin, caproin, and caprylin, which together constitute about 7.8 per cent. of the fat, the rest being composed of olein, stearin, and palmitin.

Besides fat, which constitutes about 85 per cent., butter contains water, casein, and salts. The casein averages 2.5 per cent., but may reach 7 per cent. if the butter is carelessly made; such a butter keeps badly. The water ranges from 8 to 12 per cent. in good butter, and should never exceed 16 per cent. "Milk-blended" butters have been found to contain nearly double this amount of water. The ash should not exceed 8 per cent.; in pure fresh butter it is usually about 2 or 3 per cent. or less, and includes calcium phosphate and sodium chloride.

Adulterations and Analysis.— The addition of annatto or other harmless colouring matter is not regarded as adulteration. Starch is sometimes added, and may be recognised by its blue reaction with iodine. The chief and only important adulterations practised are the substitution of foreign fat and the addition of water or preservatives, and, therefore, the analysis of butter turns chiefly upon the proportion and composition of the fat and the proportion

of water.

1. The fat should exceed 80 per cent. of the whole. (If the butter is melted in a tall glass, the water and curd will collect at the bottom, leaving the pure fat above.)

2. The specific gravity of pure butter fat should range from 911 to 913. Margarine examined under

similar conditions would range from 901 to 906.

3. The melting-point of the fat should be about 35.8° C., but it is liable to variation, not only with variation in the relative proportions of normal constituent fats, but also with repeated meltings. Adulteration with dripping, lard, or other animal fat raises the melting-point, while vegetable fats lower it.

4. The soluble fatty acids (butyric, caproic, caprylic) should not be less than 5 per cent. of the fat, and the insoluble fatty acids should not exceed

89.5 per cent. Butter fat differs from all others in containing fatty acids that are soluble in water, and their partial or complete absence denotes substitution of foreign fat.

5. Pure butter fat is readily soluble in ether, whereas other fats dissolve slowly and leave a residue.

6. Fat crystals may be visible under the microscope, and serve to show that the butter has been melted, presumably for the purpose of adulteration.

7. Water is estimated by heating a weighed quantity of butter over a water-bath until it ceases to

lose weight.

The Regulations for the sale of butter, made in 1902 by the Board of Agriculture under section 4 of the Sale of Food and Drugs Act, 1899, provide that, "Where the proportion of water in a sample of butter exceeds 16 per cent. it shall be presumed for the purposes of the Sale of Food and Drugs Acts, 1875 to 1899, until the contrary is proved, that the butter is not genuine by reason of the excessive amount of water therein."

Nothing has been made out very clearly as to the convection of specific disease by means of butter. An outbreak of diarrhea in a large hospital was shown by Shirley Murphy to be in all probability due to the consumption of a particular kind of butter. The symptoms resembled those of tyro-toxicon poisoning, to be mentioned presently.

Margarine has been the subject of special legislation (the Margarine Act, 1887, and the Sale of Food and Drugs Act, 1899), which will be referred to in another section. The term is applied generically to fats, for the most part of animal origin, prepared by churning melted and clarified beef or mutton fat with skim milk, and so treating it as to resemble butter in appearance and flavour as far as possible. For this purpose a certain proportion of genuine

butter is frequently mixed with it. It may possess nutritive value little inferior to butter, especially if manufactured from pure animal fats. Chemically it differs from butter in containing only traces of "soluble" fatty acids. It consists mainly of olein, stearin, and palmitin. The absence of butyrin, a chief constituent of butter fat, is the most valuable means of distinguishing between butter and margarine.

Cheese is formed by the action of rennet upon milk, and consists of coagulated casein with a variable proportion of fat and salts. The best, such as Cheddar, Cheshire, double Gloucester, and most American cheeses, are made from entire milk, the poorer from skim milk, or a mixture of skim and entire milk. Stilton cheese is made from entire milk to which cream has been added. Cheeses vary widely as to the amount of fat and water which they contain. A good cheese, if kept under suitable conditions of temperature, undergoes a change, known as "ripening," which improves its flavour. Oxygen is absorbed, ammonia and carbonic acid are given off. The casein undergoes fatty metamorphosis, and the calcium phosphate reacts upon the fats and the casein, giving rise to lime salts of fatty acids, and a soluble compound of casein with phosphoric acid. Lactic acid, valerianic acid, and leucin are also formed. Both in cheese and butter-making this ripening process is due to bacteria and their products, and it is on account of this fact that much has been done in Denmark and America in the way of flavouring butter and cheese by introducing favourable organisms as "starters" of the desired fermentation. Both natural and artificial ripening occasionally go wrong, and as a result we get abnormalities, such as rancid butter, "gassy" cheese, coloured cheese, or bitter cheese. Occasionally also poisonous bodies, such as tyro-toxicon, result.

Tyro-toxicon (diazo-benzene butyrate) is a poison discovered by Vaughan in cheese that has undergone a peculiar fermentation. Such cheese is not necessarily altered in appearance or taste, but has a much stronger acid reaction than usual. Tyro-toxicon can be detected chemically by neutralising with sodium carbonate, exhausting with ether, evaporating the ethereal solution, and dissolving the residue in water. The aqueous solution strikes an orange, red, or purple colour when treated with a mixture of equal parts of carbolic and sulphuric acids. It is unstable, and present generally in only very small quantities, which are difficult of detection. It is intensely poisonous, causing dryness of the mouth and fauces, a sense of constriction in the throat, nausea, vomiting, diarrhœa, and great nervous prostration. The symptoms usually pass off in a few hours, but may end in death from collapse. Butter and cream, as well as cheese, have given rise to tyro-toxicon poisoning. In 1883-84 there occurred in Michigan, U.S.A., an outbreak of cheese poisoning, which affected 300 persons, due to tyro-toxicon (Vaughan), and in 1901 seventeen cases of tyro-toxicon poisoning occurred in Finsbury. In both outbreaks the incubation period was short (only a few hours), and the symptoms were as stated above.

Adulterations of cheese are unimportant. Starch is said to be added sometimes. The use of skimmed milk or colouring matter is not regarded as adulteration. Cotton-seed oil or other foreign fat is sometimes added in place of the abstracted cream (margarine

cheese).

Analysis.—The proportion of fat to casein should not be less than 1 to 2 at the very lowest; sometimes the fat exceeds the casein. The fat may be extracted with ether, or mechanically by the Leffmann-Beam process of melting and centrifuging. The fat should be examined by the Reichert method. Water ranges

from 20 to 35 per cent., fat from 35 per cent. down to 22 per cent. in good cheese, and to 1 per cent. in skimmed milk cheese. Casein varies between 25 and 50 per cent., and salts between 3 and 6 per cent.

Lard, which is the internal fat of the abdomen of the pig, should be absolutely free from water and from foreign fat. The addition of water, which is not now widely practised, is detected by melting the lard in a tall glass and allowing time for the water to collect at the bottom. Cotton-seed oil is the other important adulterant. It is detected by its low melting-point, high power of absorbing iodine, and reduction of nitrate of silver.

Eggs.—The average weight of a hen's egg is about 2 oz., of which 10 per cent. is shell, 60 per cent. white, and 30 per cent. yolk. The white consists of albumen, with 86 per cent. water, and the yolk of albumen and fat with 52 per cent. water. The percentage composition may also be stated thus:—Shell 10 per cent., albumen and fat 23 per cent., water 67 per cent.

Preservation.—Air may be excluded by coating the shell with butter, lard, oil, wax, gum, etc., or by boiling for half a minute so as to coagulate the surface albumen. Antiseptics are also employed, e.g. the eggs are immersed in lime water, salt water, or solution of

boro-glyceride.

Tests are only necessary for the detection of stale or rotten eggs. Two ounces of salt in a pint of water forms a solution in which good eggs sink and stale ones float. Fresh eggs are transparent at the centre, stale eggs at the top.

VEGETABLE FOODS.

Wheat-flour is rich in albumen and carbohydrates, poor in fat and salts of organic acids. It contains 8 to 12 per cent. of gluten, and can therefore be made into bread. The salts are chiefly phosphates of potash and magnesia.

Examination of Flour.—Flour should be free from acidity or mouldy smell. The colour should be white, a yellow colour indicates age, fermentation, or conversion of starch into dextrin and sugar.

The microscope may reveal any of the following:

(1) Foreign starches, especially potato and rice, the characters of which will be described presently. (2) Fungi, the most common of which are sporangia of Puccinia. One species of Puccinia produces "caries" or "smut" of wheat; smut gives the flour a disagreeable smell, and the bread a blue colour. It is said to cause diarrhea. (3) Vibriones and other organisms in fermenting flour. (4) Acarus farinæ, common in damp, inferior flour which is beginning to change. Weevils (Calandra granaria) can also be seen with the naked eye.

Adulterations and impurities.—The chief are the addition of

foreign starches, mineral matters, and other cereals.

Analysis.—Water should not exceed 16 per cent. The more the water the greater the liability to change. Gluten should not be less than 8 per cent., and may reach 12 per cent. It is separated by making the flour into a paste and working it with the fingers in a stream of water until all the starch is washed away. Ash should not be less than 0.8 per cent., or more than 2 per cent. If excessive, it should be tested for iron (indicating addition of clay), calcium sulphate, and magnesium carbonate. A simple test is to shake the flour with chloroform; the flour floats, but mineral matters sink. Alum is sometimes added during grinding.

Ergotised flour is detected by making it into a paste with dilute alkali, adding slight excess of dilute nitric acid, and then neutralising; ergot gives a reddish violet colour, changing to red in presence of nitric acid, and to violet in presence of alkali. Potash gives a herring-like smell with ergotised flour.

Foreign starches will be recognised under the microscope, but a few other tests are available. Bean meal gives off a characteristic smell when drenched with hot water, and may also be detected by the successive addition of nitric acid and ammonia, which strikes a deep red colour. Potato starch grains swell up on the addition of a trace of potash in solution, which scarcely affects wheat starch grains. Pure wheat flour, made into a dilute solution in water, is turned pink on addition of iodine; but if any potato starch is present, a dark purple colour will appear.

The flour should also be put to the practical test of bread-making.

Bread is made by mixing flour with water, and kneading it, so as to form dough by the cohesion of the moistened gluten. The dough is charged with carbonic acid, which occupies innumerable lacunæ and renders the mass porous. It is then baked and becomes bread. The carbonic acid is supplied by the use of yeast ("leaven") or baking-powders (dry alkaline carbonates mixed with tartaric or other acid), or by kneading the dough with water charged with carbonic acid gas under pressure ("aërated" bread). 100 lbs. of flour make about 130 to 150 lbs. of bread.

During baking a certain proportion of dextrin and sugar is formed from starch, and there are traces of lactic and butyric acid. Alum checks these changes, which are especially prone to occur if bad flour is used. Mechanically aërated bread is the whitest and most free from dextrin and sugar. About 1 per cent. of boiled potato is frequently added to the dough.

Examination.—Bread should be white. A yellow or dark tint may be due to old or inferior flour, bad yeast, admixture of rye, or merely to presence of bran. Acidity may be caused by old or inferior flour. Bread is heavy and sodden if from any cause the dough has not risen sufficiently; such causes are bad yeast, bad flour, too much or too little heat, or defects in

manipulation.

Water should not exceed 45 per cent. An excess lowers the nutritive value, and favours the growth of fungi. The ash ranges from 1.3 to 2.0 per cent. Alum is detected by pouring upon a slice of bread, moistened by water, a freshly made decoction of logwood chips, and then a solution of ammonium carbonate. If alum is present it acts as a mordant, and after the lapse of half-an-hour, or less, a strong blue colour is manifest, which is quite distinctive for practical purposes, so far as bread analysis is concerned, although magnesium carbonate would, if present, give the same reaction. Pure bread is merely stained pink by the reagents, and on drying this changes to a dirty brown. The blue given by alumina salts is permanent on drying—much more so than in the magnesium carbonate reaction.

For legal purposes it is necessary to determine the alumina quantitatively, and then a deduction must be made for the maximum amount of phosphate of alumina which may be present in pure bread. It is usual to deduct 16 grains (of alum) per 4-lb. loaf, anything in excess of this being regarded as adulteration. In exceptional cases pure flours may contain more than this, and many analysts now allow 8 or 10 grains per 4 lbs. Alum used for purposes of adulteration usually amounts to from 20 to 40 grains per 4-lb. loaf.

Foreign starches, especially rice and potato starch, may sometimes be detected under the microscope even after baking, but as a rule they are beyond recognition in bread. Potato makes the bread damp; the percentage of water is high, and the ash is alkaline instead of being neutral. Rice has in itself a low ash (0.85) per cent. Excessive moisture and low ash should raise suspicion of foreign flour. High ash, on the other

hand, is suggestive of mineral adulteration.

The seeds of Lolium temulentum (darnel grass) may occasionally be ground with wheat flour, and cause vertigo, vomiting, convulsions, and other symptoms of poisoning. Various fermentations also occur in bread, producing chromo-

genic changes, bitter, slimy, or mouldy bread.

Barley is in the main similar to wheat in chemical composition, but contains more nitrogen, iron, and phosphates. It is liable to cause diarrhoea. Actinomycosis is now regarded as due to eating raw barley or other cereals, upon which the actinomyces is epiphytic. Oats contain more fat and more nitrogen than wheat, and are more readily cooked, but gluten is wanting, and hence bread cannot be made of oatmeal.

Rye contains less gluten than wheat, and makes a dark sour bread, which is apt to cause diarrhoea in those unaccustomed to its use. Ergotism results from the use of rye attacked by Claviceps purpurea. Maize (Indian meal) contains much fat, and its albumen to be made digestible requires careful cooking. Pellagra has been attributed to a fungus (verdet) attacking maize. Rice is poor in nitrogen, fat, and salts, but is very digestible.

Leguminosæ contain much nitrogen, chiefly in the form of legumin. Peas and beans are less digestible than cereals, and require prolonged gentle boiling.

Watercress grown or washed in polluted water was found to be responsible for an outbreak of enteric fever in Hackney in 1903, which shows that inspection of watercress beds is necessary. Intestinal parasites may also be derived from the same source.

Potatoes are antiscorbutic and very digestible. Good potatoes sink in a saline solution of specific

gravity 1,100; bad ones float.

Starch is not a simple chemical substance, but a mixture of three bodies—granulose, erythro-granulose, and cellulose, which are respectively turned blue, red, and yellow by iodine. Granulose is the most important of these. Boiled in water, or with oxalic or other organic acid, it is converted into "soluble starch" or amidulin, a soluble dextro-rotatory substance chemically indistinguishable from granulose, and giving the blue iodine reaction. If instead of organic acid a 2 per cent, solution of sulphuric acid is boiled with the granulose or starch, the soluble starch first formed is converted into dextrin or erythro-dextrin, a sort of gum, which is dextro-rotatory, and is reddened by iodine; it is precipitated by alcohol and acetic acid, but not by tannic acid. Still further heating with sulphuric acid produces achroödextrin, which gives no colour reaction with iodine, and glucose.

Erythro-granulose is present in small proportion in wheat starch, and almost absent in potato. Its affinity for iodine is greater than that of granulose, but the red tint is completely hidden by the deep blue of the iodine-granulose. If a minimal proportion of iodine is employed, the red colour appears. Both the red and blue disappear on heating, owing to dissociation of

iodine and starch, but reappear on cooling.

Starch occurs in the various farinacea in microscopic granules, which in many cases have a sufficiently characteristic appearance to indicate their origin. The granules vary greatly in size. Some are roundish, others oval, pyriform, truncated, polyhedral; some have distinctive markings, especially faint concentric rings or a hilum.

Microscopic examination of starches.—A little of the starch, in a state of fine powder, should be mixed with a drop of water,

preferably containing 30 per cent. of glycerine, and put under the microscope. The principal varieties may be arranged in

five classes (Muter) :-

1. Large oval or pyriform grains with hilum and concentric rings.—This group includes potato starch, and several varieties of arrowroot. Tous les mois, or Canna arrowroot, has the largest grains of all; Maranta (Jamaica or St. Vincent) arrowroot is characterised by a linear hilum at the larger end.

Potato starch grains vary greatly in size and shape; the hilum is minute, but distinct, and is placed at the smaller end.*

2. Irregular, round, or oval grains with stellate hilum, but no rings.—Maize starch grains are round, or polygonal with rounded angles. Beans and pea starch grains are oval and very much alike, but the former are far more uniform in the size of the grains.

3. Round or oval grains without hilum or rings.—Wheat starch grains differ greatly in size, and there is a characteristic absence of medium sizes; they are round and flattened. Polar-

ised light shows a cross.

Barley starch is much more uniform. Rye starch granules,

if old and dry, often present a large stellate hilum.

4. Grains truncated at one end.—Sago starch has a round hilum at the convex end, and faint rings; the starch grains of prepared sago have a large depression. Tapioca starch grains are smaller than sago, and some of them show signs of altera-

tion by heat.

5. Angular grains.—Oat starch is often massed into little balls or clusters of grains fairly constant in size. Rice starch is smaller than oat, and pepper starch the smallest of all. Under very high power both rice and pepper show a hilum. Tahiti or Tacca arrowroot is larger than oat starch, and less uniform in size; it resembles maize, but has sharp angles.

Alcoholic beverages.—Alcohol, so far as it is oxidised, is in a sense a food, but by dilating the surface blood vessels, it may cause more heat to be lost than it is itself capable of producing, and in large doses the general paralysis of cellular activity produced by alcohol is so great, that heat production is diminished, and the final result is a lowering of body temperature.

* Potato starch may also be distinguished from arrowroot by certain chemical tests. When mixed with half its weight of hydrochloric acid, the former gives a transparent jelly, the latter a white paste. Potato starch gives an offensive odour when heated with sulphuric acid. Potato starch jelly made with water turns thin and sour in twelve hours, maranta not for three or four days.

Still less is it a useful producer of energy on account of its paralysing effect on nerve cells and the muscular system. Hence, though a food in an academical sense, and though having at first an apparently stimulating effect on the body, it does not give staying power. It may, of course, be associated with other substances (as in malt liquors) which have a definite value as food. Broadly speaking, it may be said that the effects of large doses of alcohol are the reverse of those produced by small doses (or amounts which may be taken without any elimination by the kidneys); that is to say, alcohol in large doses acts as a depressant and produces chronic irritation, which manifests itself in the stomach (inflammatory conditions, atrophy), in the liver (cirrhosis), in the brain (degeneration), etc. Such conditions are, of course, modified by individual habits and idiosyncrasies, age and sex, exercise, diet, occupation, and disease. The three chief forms of alcoholic beverages are (a) malt liquors, such as beer, ale, porter, and stout; (b) wines, such as the sparkling wines of the Rhine and the Moselle, champagne, hock, claret, etc.; (c) spirits, as whisky, gin, rum, and brandy.

Spirits contain about 40 per cent. of alcohol, rather less in gin, and rather more in rum. Port and sherry contain about 17 or 18 per cent., and ordinary light wines 6 to 14 per cent. English beer contains from about 3 to 7 per cent. of alcohol, German beer 2 to 5 per cent. Rectified spirit contains 90 per cent. by volume of absolute alcohol; proof spirit is a mixture of 57.3 per cent. by volume of absolute alcohol, and 42.7 per cent. of water. The ratio of alcohol to proof spirit being as 1: 1.76, the amount of alcohol in any liquid being given the amount of

proof spirit can readily be calculated.

Malt beers are made by inducing yeast fermentation of sugar, derived from the starch of barley. During germination, the insoluble starch is converted into dextrine, maltose and glucose, the process being completed by "mashing" with water at 160° F. The wort is then boiled, hops are added, and after cooling, the liquid is discharged with yeast into the fermenting tun. Non-malt beers are made by fermenting glucose and allied substances obtained by the action of sulphuric acid upon starch. Wines are produced by fermentation of the juice of the grape, complete fermentation resulting in a dry wine, incomplete in an effervescing or sparkling wine. Spirits are prepared by the distillation of some previously fermented liquor; brandy from wine, whisky from malted barley, rum from molasses. But there are many varieties in distillery methods, and much "brandy" is made from potato spirit.

From the point of view of the medical officer of health alcoholic beverages assume importance in three ways: (1) disease caused by excessive consumption; (2) adulteration, and (3) fraudulent imitations sold

to the prejudice or injury of the purchaser.

The diseases commonly attributed to excessive consumption of alcohol are chiefly alimentary or nervous, producing cirrhosis of the liver and various forms of gastric disease, or alcoholic neuritis, delirium tremens, etc. There is, of course, in addition, a large amount of disease, directly or indirectly, due to alcoholism, which does not appear in death certification.

Arsenic in Beer.—Of adulterants introduced accidentally or otherwise into alcoholic beverages, arsenic in beer is perhaps the best example. In 1900 there was wide prevalence of illness in South Lancashire and elsewhere, which was, after a time, identified by Reynolds as a form of arsenical poisoning, and traced to the consumption of beer from certain breweries in which highly arsenical brewing sugars had been used. These were found to be derived from a common

supply, in the manufacture of which, from starch, arsenical sulphuric acid had been used. The arsenic (As₀O₃) found in the sulphuric acid was 1.9 per cent., in the artificial sugars 0.008 to 0.131 per cent. (0.56 to 9.17 grs. per lb.), in beer often 1/4 to 1/2 grain per gallon, and in one sample 3 grains per gallon. This danger vanishes if care is taken to use acid free from arsenic, but a more persistent source of arsenic in beer was found in the malt itself. The usual process of drying the malt in kilns with exposure to fumes of gas-coke or anthracite entails risk of contamination by arsenic, in such proportions as 1 th of a grain per lb. The most effective remedy here would seem to be drying out of contact with the fumes; but much can be done by selection of non-arsenical fuel, by careful construction and cleanliness of the kiln, so as to avoid arsenical deposits on ledges falling into the malt, and lastly, by "brushing" the malt, whereby the fibrils (which readily become charged with arsenic) and surface dust are removed. At Halifax, in 1902, the use of arsenical malt caused a local outbreak of poisoning among beer-drinkers.

It was estimated that in the outbreak of 1900 at least 3,000 persons were attacked in Manchester and Salford alone, and 6,000 in all. Many attacks were fatal. For some time the nature of the epidemic was obscured by the unusual character of some of the prominent symptoms, more especially the peripheral neuritis and pigmentary changes. In the main, the attacks and their severity bore relation to the consumption of beer, but some of those attacked had apparently consumed but little, while some heavy drinkers escaped lightly. Similar variations in toleration and susceptibility are met with in the therapeutic use of arsenic. A high degree of insusceptibility is recorded in the case of the Styrian peasants, who take large doses without apparent ill

effect, and it may be assumed in the case of some arsenic workers. It is important to note, as an aid to diagnosis, that arsenic can be found not only in the urine, but in the hair.

The Reports of the Royal Commission on Arsenical Poisoning, dealing primarily with the outbreak of 1900, call attention also to some risk of presence of arsenic in glycerine, phosphates, boric acid, colouring matters, and other food ingredients. The Commission recommended a penal limit of $\frac{1}{100}$ grain of arsenic per pound, or per gallon, in any food substance.

In the Marsh-Berzelius test, the suspected material is first treated with hydrochloric acid in a suitable vessel. It is then placed in the Marsh-Berzelius apparatus, which consists of a flask for the generation of hydrogen from which passes a drying tube and a glass tube drawn out in the middle and at the end. In this apparatus hydrogen is obtained from pure zinc and pure sulphuric acid, and the substance under examination is placed in the flask. Hydrogen and arseniuretted hydrogen escape through the drying tube, containing calcium chloride, and thence pass into the fine glass tube, which is heated over a Bunsen burner. The arseniuretted hydrogen is decomposed, and the arsenic is deposited on the wall of the narrow part of the tube, and so forms a "mirror." Finally, this mirror is compared with standard mirrors prepared from known quantities of arsenic (e.g. $\frac{1}{50}$, $\frac{1}{66}$, $\frac{1}{88}$, and $\frac{1}{100}$ of a grain per gallon). The Reinsch test is less reliable.

The question of adulteration of alcoholic beverages is too large for consideration here, but, as an illustration, mention may be made of recent work in connection with brandy. In 1876, owing to an outbreak of phylloxera, a parasite affecting and destroying vines, the usual output of genuine brandy from the Cognac district was curtailed, and artificial preparations were substituted for the genuine article. In Islington Borough Council v. Hillyer (1904), a sample of "fine old pale brandy" proved, on analysis, to contain at least 60 per cent. of a spirit not derived from the juice of the grape. In the absence of any

official or legal standard brandy may be defined as an alcoholic liquor the spirit of which is obtained by distillation, in a pot-still, from the juice of the grape, and which should contain not less than 80 parts by weight of esters (calculated as ethyl acetate) per 100,000 parts by volume of absolute alcohol. In the above sample there were approximately 40 parts of esters, the volume being made up with plain spirit. The court held that this was not genuine brandy, and convicted.

Whisky consists of spirit derived from malted barley, or a mixture of malted or unmalted barley or wheat, distilled in a pot-still. A spirit thus produced contains at least 380 parts of secondary bodies (higher alcohols, esters, aldehydes, etc.) per 100,000 of absolute alcohol.

Summary as to relation of food and health.—From what has been said it will be seen that there are many ways in which health may become affected by articles of food.

1. Certain essential constituents of diet may be

deficient or in excess.

2. Poisonous substances may be derived from the vessels in which the food has been kept. Thus tinned provisions (fruit, fish, or meat) may become poisonous, and cider kept in lead-glazed vessels may cause lead-poisoning.

3. Injurious substances may be added or substituted accidentally, or by way of adulteration, or by improper processes of manufacture, or by the drug-

ging of animals before death.

4. Certain kinds of food are liable to be occasionally poisonous even in the fresh state (mussel-poisoning), or when grown under unfavourable conditions (oysters, watercress, etc.).

5. Putrefactive changes may have commenced.

6. Poisonous substances may be developed, either as a result of parasitic growth (fermentation) or from, at present, unknown causes (tyro-toxicon).

7. The flesh or milk of an animal suffering from certain specific or parasitic diseases may impart the disease (e.g. tuberculosis, trichinosis, hydatids). Actinomycosis affords an instance of somewhat parallel danger from vegetable, as well as from animal, food.

8. Food, and especially milk, may become infected by virus of human origin (diphtheria, enteric fever,

scarlet fever).

It should not be forgotten that disease in the individual, or more rarely idiosyncrasy apart from disease, may render certain kinds of food injurious which to ordinary persons are wholesome. Lastly, there are certain accessories of diet which may be injurious if used injudiciously (alcohol, tea).

In the systematic investigation of suspected cases of poisoning by food, help may be obtained by the use of a tabular form, a separate line being devoted to each item of food consumed in the household, with separate columns for each day and for each inmate,

whether ill or not.

isonati susti	M. 36. Ill.		F. 34.		M. 12. Not ill.			F. 5. Not ill.				
dual sensor	Mon.	Tu.	Wed.	Mon.	Tu.	Wed.	Mon.	Tu.	Wed.	Mon.	Tu.	Wed.
Beef	×		×	×		×	×		×	×		×
Pork		×		1 5	×	18.9	1330	1		1		1
Fish		×		1 6000				×		1	×	
Potatoes	×	×	×	×	×	×	×	×	×	×	×	×
Cabbage	X	×		×	×		×	×	1	×	×	
Bread	×	×	×	×	×	×	×	×	×	×	×	×
Butter (fresh) .	×	×	×	×	×	×						
,, (salt) .							×	×	×	×	×	×
Cheese	×	×	×				×	×	×			
Milk (cooked) .	×	×	×	×	×	×	×	×	×	×	×	×
etc. etc.	×	×	×	×	×	×	-	×	×	×	×	×

The list should be as complete and detailed as it can be made by the most careful inquiries. When the table is filled up, it will be seen at once what articles of diet (if any) were consumed by all the sick and by none of those who escaped attack.

CHAPTER V

SOIL

The chemical and physical constitution of the soil and the configuration of its surface have an important bearing upon many hygienic questions, particularly in relation to water-supply, burial-grounds, sewage land treatment, sites for buildings, and the prevalence of certain diseases.

Temperature.—The daily variation of temperature ceases to be perceptible at about three or four feet from the surface. The annual variation is unimportant below six or eight feet, very small below 24 feet, and vanishes at 40 feet (Galton) or 50 to 100 feet (Parkes). The maximum is later as the distance from the surface increases, and at Edinburgh it was found that in trap-rock at a depth of 24 feet the maximum was in January and the minimum in July. The absorption of heat, assuming that the exposure is equal, is determined mainly by the nature of the soil and the presence or absence of vegetation. The absorption by sand being taken as 100, that of clay is about 70, chalk 60, and humus or mould 50. Radiation varies in like manner, but is generally greater than absorption. Trees and shrubs intercept the sun's rays, and on the other hand check evaporation from the surface of the soil, the net result being to render the ground cool and moist in winter, and cool and dry in summer when the leaves are out. The evaporation from leaves is very great, and tends to moisten and cool the air, and abstract water from the soil. Pettenkofer calculated that an oak tree with 750,000 leaves had in the summer months an evaporation equal to eight times the rainfall, and in

Algeria the Eucalyptus globulus has been found to absorb and evaporate twelve times the rainfall.* Herbage lessens absorption of heat by the soil, and also true radiation, but increases evaporation and convection owing to the vast surface of the blades of grass. Grass, therefore, like trees, renders the soil cooler and drier, and more equable in temperature.

Earth temperatures are usually determined by thermometers mounted in paraffin or wax, suspended by a chain at the required depth in the soil (1 to 4 feet or more), and withdrawn at daily or other intervals for reading. An iron tube driven vertically into the ground, protected by open wooden covers, affords a passage for the thermometer. Systematic records are usually kept of the daily readings of two earth-thermometers (at 1 foot and 4 feet). At greater depths the internal heat of the earth produces an appreciable effect, causing a rise of about 1° Fahr. for every 55 feet.

Moisture.—At a certain point below the surface in permeable soil, all the interstices are full of water, to the complete exclusion of air. This subsoil water or ground-water extends downwards to the next impermeable stratum. In marshy ground the surface of the subsoil water may coincide with that of the ground itself, and under ordinary conditions it may be from two or three feet to a hundred feet or more below the ground level. The surface of the ground-water is not necessarily horizontal, or parallel with that of the ground. It flows (usually towards the nearest watercourse or the sea) with a velocity that varies with the permeability of the soil, the steepness of the gradient, and the absence of obstruction by roots of trees.† The level is affected by that portion of the rainfall which percolates through the soil; also by the height of water in adjoining streams or seas, and other conditions modifying the

^{*} As regards the air, forests render both the daily and annual temperatures more equable, and increase the humidity (Galton).

+ In Munich Pettenkofer estimated the velocity of the ground-water to

be 15 feet per day.

facility of outflow—artificial drainage, etc. Sometimes the range of rise and fall does not exceed a few

inches, but it is more usually several feet.

Above the level of the ground-water the soil, though aërated, is still kept moist by capillary attraction, by evaporation from below, by rainfall, and by the movements of the ground-water. The soil constantly loses moisture by evaporation from the surface, both directly and through vegetation.

The moisture of a sample of soil is estimated by weighing it before (B) and after (A) drying it on a water-bath. Then $\frac{B-A}{B} \times 100 = \text{percentage}$ weight of water in sample. The level of the subsoil water is necessarily that of the water in surface wells, and may be gauged by lowering into the well a cord or rod bearing little cups at short intervals, or more simply by means of an index connected by a cord with a float.

When the ground-water rises, it forces air out of the soil, and at the same time may pollute wells by bringing into them the washings of impure soil. As the ground-water falls again it leaves the soil moist and aërated, conditions favourable for fermentative and putrefactive processes in organically polluted soil.

Damp ground is rendered dry, and the level of its ground-water permanently lowered, by subsoil drainage, surface drainage, and opening the outflow. For subsoil drainage "agricultural" pipes, *i.e.* unglazed porous pipes, are laid loosely end to end at the bottom of deep trenches, and covered in. The interval to be allowed between the parallel drains will depend upon the openness of the soil.

Air ("ground-air") is present in all rocks except the very hardest, above the ground-water level. It fills all the space unoccupied by water or solid particles. The proportion by volume varies greatly with the porosity and other conditions of the soil; loose sand may contain 50 per cent., and humus several times its own volume of air. Ground-air contains moisture, organic matter of animal or vegetable origin, and sometimes NH3, H2S, and CH4. Carbonic acid is in excess, and increases with the depth; it also varies greatly according to the nature of the soil and the intensity of the local chemical changes that are going on in it. Oxygen is in relatively small proportion, and decreases with the depth. At 13 feet Fodor found 14 per cent. of CO₂ and 7.5 per cent. of oxygen. Many causes combine to keep the ground-air in constant movement, among them being wind, percolation of rain, variations in temperature and barometric pressure, and rise and fall of ground-water.

The percentage volume of air in a sample of rock of specific gravity a may be determined by finding its weight when dry (A), and when saturated with water (B). Then $\frac{B-A}{A}$ × 100 = percentage of air. For loose soils Pettenkofer used two graduated burettes connected together at the bottom by a clamped tube. One burette was filled with water, and the other with dried and crushed sample soil. Water was then allowed to pass upwards through the soil, displacing all the air until it just reached the surface.

No. of c.c. of water used × 100 = percentage of air. No. of c.c. of dry soil

All rocks are more or less permeable, and therefore capable of containing air or water in their interstices. A cubic yard of granite or marble holds about a pint of water, sandstone 25 gallons, and sand 50 gallons,

Chemical composition.—The mineral constituents of soil are of the utmost variety, but only the organic matters will be considered here. The soil undergoes constant pollution by dead animal and vegetable matter and excreta, which are as constantly

being removed by putrefaction, nitrification, and the influence of vegetation. Apart from such obvious examples as graveyards, manured fields, sewage-farms, and the like, unintentional pollution of the soil goes on upon an enormous scale, especially around dwellings. Nitrates are commonly found in abundance in the soil of any long-inhabited place. Carbonic acid and other simple compounds are formed concurrently; offensive gases are given off during putrefaction, but are deodorised and oxidised by passing through aërated soil. Vegetable matters are more persistent. They may undergo putrefaction and slow oxidation, but it is not certain that nitrification takes place.

Sometimes the natural level of the ground is artificially raised, constituting what is known as "made soil." This may be ordinary earth, but in the neighbourhood of towns it is a common practice to fill up hollows, and to elevate low-lying sites by trade refuse, ashes, household refuse, and even excremental refuse. From such soil the impurities fairly rapidly disappear, owing to oxidation and washing by rain. Sanderson, Parkes, and Savage found that in about three years most of the vegetable matters in cinder refuse had disappeared; wood, straw, and cloth were rotten, but still distinguishable. Free exposure to air and rain and free outlet for drainage will accelerate the process of purification.

Bacterial content.—Soil, particularly within a few inches of the surface, contains many bacteria. These have been divided into groups according to function. The decomposition or denitrification organisms abound particularly in soils possessing a high degree of organic matter, and their rôle is to break down compounds of ammonia and nitrogen, yielding free nitrogen, CO₂, water, ammonia, and sometimes nitrites. These latter bodies are oxidised by the nitrifying organisms first discovered by Schlösing

and Muntz in 1877. Nitrification is performed in two stages; first, an oxidation from ammonia bodies to nitrites, carried out by the nitrous organism, and secondly, the oxidation of the nitrites into nitrates by the agency of the nitric organism. For the performance of their function these organisms require pabulum, oxygen, a base with which nitric acid when formed can combine, and a favourable temperature. Both organisms are widely distributed in superficial soils (within 12 inches of the surface), and act together and in conjunction for the common purpose of nitrification. In organic solutions, nitrous organisms thrive and nitric organisms do not, the reverse being true in inorganic solutions containing nitrites, in which nitrous organisms are incapable of the oxidation process. The presence of peaty or humous matter appears to preserve the vitality of nitric organisms during the fermentation of ammonia. A third group of allied organisms are the nitrogenfixing bacteria found in soil and in the roots of leguminosæ; their function appears to be, directly or indirectly, the fixation of free nitrogen (symbiosis). This may be performed under the symbiotic condition by the leaves, by distribution in the soil of the root nodules, or by the nodules themselves. It is thus, in part, that clover, vetches, peas, beans, and other leguminous crops increase the amount of fixed nitrogen in the soil, particularly where it is especially small in quantity. Associated with these groups of bacteria are a number of saprophytic organisms, the function of which is not known. Houston has described certain acid-forming microbes in peat.

It would be difficult to over-estimate the value in the economy of nature of the three groups of organisms to which reference has been made: they remove organic matter and waste animal products, they carry on the oxidation processes which are applied in the bacterial treatment of sewage, and they fix free nitrogen. Some at least of the pathogenic organisms are also known to live and multiply in soil, and at least three find in such a medium a favourable nidus. The bacillus of tetanus (B. tetani) is common in the soil and dust of certain districts, either in a bacillary form or in the resting stage of spores. The presence of this bacillus in jute factories is possibly due to its attachment to the roots of the Corchorus in the soil of Bengal. The organisms of quarter-evil (symptomatic anthrax) and malignant ædema are two other examples of pathogenic bacteria in soil.

Recent research has shown that the typhoid bacillus bears a relation to soil. In 1888 Grancher and Deschamps showed that it was able to survive in soil for more than twenty weeks. In 1894 Dempster published the results of his work on the same subject, in which he obtained the bacillus from sand after 23 days, from garden soil after 42 days, and from peat not later than 24 hours. Four years later came Robertson's researches into the growth of the typhoid bacillus in the soil of an ordinary field, from which he was able to isolate the living bacillus twelve months after. Sidney Martin found the organism still alive after 404 days in dry sterilised garden soil; in wet soils it lived but a short time, as also in uncultivated soils which had not been sterilised, and in which ordinary putrefactive germs abounded. Under such conditions the typhoid bacillus appeared to die out rapidly. Lastly, we have the results of the investigations of Firth and Horrocks (1903), who conclude that the typhoid bacillus is able to assume a vegetative existence in ordinary soils and in sewage polluted soils for as long as 74 days. They further maintain that the controlling factor is an excess or deficiency of moisture in the soil rather than organic

nutritive material. From dry fine sand the bacillus was recovered after 25 days; from moist fine sand after 12 days; from damp (rain-water) ordinary soil after 67 days; from damp (sewage) ordinary soil after 53 days; and from ordinary soil dried to the state of dust, after 24 days. In peat the bacillus lives apparently only a few days. Firth and Horrocks, therefore, arrive at a different conclusion from Martin, namely, that the typhoid bacillus is able to assume a vegetative or saprophytic existence for considerable periods outside the body, that it can survive in ordinary earth for over two months, whether the soil be virgin or polluted with sewage, or frozen hard, and that, therefore, it follows that outbreaks of enteric fever may be due to the dissemination (for example by wind or flies) of infective soil dust. Pfuhl of Berlin (1904) has arrived at results confirmatory of these experiments. From moist garden earth he recovered the typhoid bacillus 88 days after inoculation, from dry sand after 28 days, and from moist peat 21 days.

Sewage added to ordinary garden soil leads to a temporary increase only of sewage microbes at the expense of soil microbes, the ordinary soil bacteria eventually ousting the sewage microbes in the struggle for existence. Sandy soil so treated does not recover for some months. The chief kinds of bacteria resulting from such soil treatment are indol-producing bacteria, gas-producing bacteria, spores of B. enteritidis sporogenes, B. coli and its allies, and streptococci (Houston). Further, it should not be forgotten that soil has the power of destroying some pathogenic

germs as well as sewage organisms.

Soil and disease.—Among important specific diseases, anthrax, tetanus, malaria, enteric fever, cholera, diarrhœa, and yellow fever are closely concerned with telluric conditions. Dampness of soil is favourable to phthisis and diphtheria, and, according

to some authorities, to rickets also. Goitre is often credited with a more indirect relation to the soil

(page 76).

Enteric fever and cholera exhibit in certain localities a somewhat definite inverse relation to the fluctuations of the level of the ground-water. In Munich, Leipsic, and elsewhere, it has been found that a fall in the subsoil water, and especially a rapid fall after an unusually high level, is followed by an outbreak of enteric fever. These observations, extending over several years, led Pettenkofer to the conclusion that the conditions requisite for an enteric outbreak are (1) a rapid fall (after a rise) in the ground-water, (2) pollution of the soil with animal impurities, (3) a certain earth temperature, and (4) the presence of a specific organism in the soil. These conditions imply a warm, moist, and well-aërated soil, containing pabulum suitable for the specific microbe. On the other hand, the association of falling ground-water with enteric outbreaks is not observed in other localities. It has not been traced in England, as a rule, and, indeed, outbreaks have occurred with rising groundwater. Although the existence of some direct or indirect relation between the two phenomena in Munich cannot be doubted, Pettenkofer's hypothesis is not the only possible explanation. Buchanan suggested that the connection between them may be found in pollution of wells, but Pettenkofer obtained no analytical evidence of this.

The same conditions, according to Pettenkofer, are necessary for the appearance of cholera in an epidemic

form.

Malaria, including under that term the various forms of intermittent and remittent fevers, is also, in part, a "telluric" disease. The specific protozoa have at one stage of their development their normal habitat in the soil or in surface water, and require

air, a certain degree of warmth, and decomposing vegetable organic matter. A rise of ground-water due to heavy rainfall, or impeded or insufficient outflow, is a common antecedent of severe outbreaks in malarious districts, and many such districts in England and elsewhere appear to have been permanently freed from malaria by improved drainage. Land that has gone out of cultivation may become malarious if the other conditions are favourable. Marshy land, especially if covered with rank vegetation, is the most common locus, and estuarial marshes are favourable breeding grounds. Either permanent flooding or drainage will, as a rule, render them non-malarious.

Enteric fever, cholera, and yellow fever are often endemic, a fact that in itself is strongly suggestive of telluric relations. They are also found to flourish best where the soil is polluted with excremental or other animal matter. The same may be said of diarrhæa, and Ballard has shown that outbreaks of summer diarrhœa have a close relation with the earth temperature at a depth of four feet. Soil may become infected by the spores of anthrax or tetanus, and retain the infection for an almost indefinite time.

Buchanan believed not only that phthisis mortality bears a direct relation to dampness of subsoil, i.e. to height of ground-water, but also that when the ground-water is lowered and the soil dried by artificial drainage, the recorded phthisis mortality falls, but this view must be accepted with reserve. He concluded that there is less phthisis among populations living upon pervious soils, which were higher lying and sloping, than on impervious, low lying, and flat soils. The term phthisis is, of course, to be taken in a broad sense. It appears, also, that dampness of soil is favourable to all affections of the respiratory system, including bronchitis and pneumonia. In the

absence of definite evidence the popular theory that damp and cold soils are conducive to rheumatism and all manner of catarrhs may be provisionally accepted. A high level of ground-water (say within five feet of the surface) is, therefore, objectionable, and it is preferable that it should be fifteen feet or more from the surface; but frequent and extensive fluctuations of level are probably worst of all. The presence of clay, like high ground-water, renders a soil "cold."

Building sites .- Porous soils, like sand or gravel, with low ground-water, are the warmest, and in the main the most healthy; but they are also the most liable to organic pollution, and the ground-air moves freely in their interstices. Clay, and especially a clay slope, affords a far more healthy site than gravel or sand containing organic matter, the impure groundair from which is constantly forced out of the soil, and is also liable to be drawn into the interior of houses through the basement. The most unhealthy site of all would be a hollow with impure gravelly soil and high and fluctuating level of ground-water; the most healthy, the summit of a slope with dry, pure, sandy or gravelly soil, and low and stationary level of subsoil water. The foot of a slope often receives the drainage from the higher ground, and there is the same liability to damp at the junction of an upper porous and a lower impervious bed, especially if the strata dip towards the point in question. In tropical regions it is necessary to take into account the tendency of winds to pass up valleys and ravines, from lower to higher levels, during the day, and in the reverse direction at night. The proximity of marshy or "malarious" ground may be dangerous, even if the site itself is otherwise free from objection. Herbage is beneficial, for the reasons already stated, and so too are trees, unless they are

allowed to interfere with light and circulation of air. Manured lands have not been proved to be hurtful: but irrigated lands, and especially rice-fields, which give off organic matter and much moisture, are injurious; and in Italy rice-grounds are not allowed within 14 kilometres of cities, or 1 kilometre of small towns (Parkes). There seems to be no good evidence against well-managed English sewage farms. "Made soil" is usually impure and unfit for building sites. The objection is greatest when the deposit is recent (within 3 years) and contains excremental or other organic refuse, and is so placed in hollows that drainage and aëration are impeded.

Geological formations.—The following is a summary of Parkes's conclusions in respect to the healthiness of certain geological formations, but, as he points out, they are of less importance than the

immediately local conditions:-

Granitic, Metamorphic, and Trap rocks, Millstone Grit and Clay Slate are impervious and generally healthy. The slope is great, vegetation not excessive, air dry, and water pure. Marshes are infrequent. These rocks, when weathered and disintegrated, are alleged to be unhealthy.

Limestone, Oolite, and Magnesian Limestone (Dolomite) have also much slope, and water passes off readily. Marshes may occur, even at great heights. Magnesian limestone is the worst and oolite the best.

Goitre and calculus may be met with.

Chalk is healthy if permeable and free from clay. The air and water are pure. Goitre and calculus are less common than in limestone districts. Marly chalks are impermeable, and therefore damp and cold, and may be "malarious."

Sandstone, if permeable, is healthy. The air and soil are dry, but the water is often impure. An admixture, or substratum, of clay may render the site damp. Sand is healthy if pure and dry. Many sandy soils are unhealthy, e.g. when the subsoil water is high and organic impurities are present. The water is liable to contain salts of the alkalies, lime, magnesia, and iron, and is often unfit for drinking purposes.

Gravel is healthy, except in hollows where the ground-water is high. Gravel hillocks are the healthi-

est of all sites. The water is usually pure.

Alluvial soils, including Clay and dense Marl, are usually unhealthy. Water neither runs off nor through them, marshes are common, the air is moist, and the water often impure with lime and soda salts. The deltas of great rivers present these alluvial characters in the highest degree, and should not be

adopted for sites without thorough drainage.

Much may be done to improve a site not naturally healthy. If damp it may be drained, or trenches may be cut around it so as to intercept the subsoil water on its way from higher ground. By such means the level of the ground-water will be lowered and its fluctuations lessened. Damp places near the site may be drained or filled in, and the site itself may be artificially elevated if necessary. Trees are often desirable for shelter or ornament, but should not be allowed to interfere with light or circulation of air. Rank or superabundant vegetation must be cleared away, but short grass is beneficial.

signal boards still with the ball we have been supported by the

CHAPTER VI

BUILDINGS.

Materials and general construction.— Most of the ordinary building materials are pervious to air and water to a greater or less extent. Galton states that with a temperature of 40° F. outside, and 72° F. within, the volumes of air passing through a square yard of wall vary according to the material, as follows:—

Sandstone		1.00	4.7	cubic feet	per hour.
Quarried Li	imestone		6.5	,,	**
Brick .	ACCEPTOR!	100	7.9	The said	1,,
Limestone	SVIII JUST	1	10.1	Total	,,
Mud .	Mary Mary		14.4	,,	0)

The thickness of the wall is supposed to be the same in all cases. Mortar, cement, and concrete* are also permeable, and the caustic lime they contain absorbs carbonic acid until it becomes saturated. Wood in its natural condition absorbs water, and the inevitable joints and crevices admit of the passage of air and water. Bricks are usually 9 in. \times $4\frac{1}{2}$ in. \times $2\frac{3}{4}$ in. They are porous and absorbent, but if of good quality make excellent building material. Stone varies greatly in consistence and durability. Portland stone is much used in the metropolis.

By painting, varnishing, and other means the inner surfaces of walls, floors, and ceilings can be rendered impervious, and walls lined with glazed bricks or the finer cements (Parian) are scarcely permeable.

^{*} Mortar consists of sharp sand and slaked lime in the proportion of 3 to 1. Frequently inferior material is worked up into mortar. Portland Cement is made by grinding chalk and mixing it with blue clay and river mud, then burning it in a kiln and grinding it to a fine powder. "Compo" is a mixture of Portland cement with sand for external plastering in two coats, a rough coat composed of 1 part cement to 5 parts compo, and a fine coat of 2 parts fine sand and 1 part cement. With this material walls are "rendered." Concrete is composed of 3 parts broken ballast, 2 parts sand, and 1 part lime or cement.

The disadvantages of porous materials are obvious, and the air admitted in this way though small in amount is at least doubtful in purity. Pervious walls are liable to be damp and cold. Air passing through them in either direction is filtered, and the wall becomes charged with organic impurities. This is specially the case with the inner surfaces, and in hospitals it is now customary to render the walls and floors impervious as far as possible. The exclusion of damp and impure ground-air, by means of impervious basements, is absolutely necessary in all inhabited buildings. This may most conveniently be effected by covering the basement in every part with a layer of concrete six inches thick, as required by the Model Bye-laws, but other materials may be employed.

It has been proposed to cut off the communication with the soil still more thoroughly by elevating the house upon pillars and arches, thus allowing free play of fresh air beneath it. If there are no cellars, the same result may be attained in some degree by constructing the lowest floor a foot or two above




Fig. 5.—Damp-course.

the ground level, and freely ventilating the space beneath by means of large air-grates. In houses built

without precautions of this kind the entrance of ground-air through the basement is inevitable whenever the air in the interior of the house is warm, and therefore lighter than that outside.

Measures must also be taken to prevent damp from




Fig. 6.—Double damp-course. Soil abutting upon hollow wall.

rising in the walls by capillary attraction. For this purpose a "damp-course" must be provided, that is, a continuous horizontal course of glazed earthenware, sheet lead, two layers of ordinary roofing slate set in cement, 3in. of asphalt, stone= ware slabs, two courses of Staffordshire bricks, or other impervious material, of the full thickness of the wall, above the highest point at which the wall is in contact with the earth, and below the lowest timbers or

floor supports (Fig. 5). It is undesirable to allow the soil to be in contact with the wall of any room or cellar, and this can usually be obviated by excavating it on the outside to below the level of the floor so as to form a "dry area," but if this is impracticable the device shown in the Model Bye-laws may be employed; the wall is made hollow up to a point above the ground level, and two damp-courses are inserted, one at the level of the bottom of the cavity, and of course beneath the floor level, the other at the top of the cavity, and, therefore, above the level of the ground outside (Fig. 6). In this way the inner wall is

completely isolated from the soil.

The necessity for free ventilation extends to all closed spaces, including those below floors. There is a tendency among builders to use air-grates so constructed and arranged as to allow but little air to pass. If the ventilation is insufficient, the air becomes damp and musty, and the fungous growth known as "dry-rot" (Merulius lacrimans) is liable to set in. As regards the upper storeys, however, the ventilation through the boards and ceiling is usually sufficient to prevent dry-rot, unless the floor is covered with oilcloth or otherwise rendered impervious. Fireproof

floors are now becoming common.

The thickness of walls is often regulated solely by considerations of stability, and the choice of materials by their cheapness and convenience, but regard should also be had to warmth and exclusion of damp. It is sometimes advantageous to make the outside wall hollow or double, or even to fill in the hollow of a double wall with pitch or cement. The outer surface may also be made impervious by paint, tar, cement, or slate. Damp walls may be due to ascending moisture where there is no damp-proof course, or to overflow from roof-gutters and rainwater pipes, etc. The walls separating one house from another should be carried up to or above the roof in every part. Any communication beneath the roof is objectionable for many reasons, including danger in case of fire or infectious disease in one of the houses. The Model Bye-laws require for the external walls or party-walls of houses a minimum thickness of nine inches, increasing according to a prescribed scale when the height exceeds 25 feet or the length 30 feet. Walls should be properly bonded by Flemish bond (bricks placed transversely and longitudinally by turn) or English bond (a course of longitudinal

alternating with a course of transverse).

The roof itself affords very little protection from the extremes of temperature, and the rooms immediately beneath it, if intended for occupation, should have a "false roof" or ceiling over them, so as to

leave an intervening air-space below the slates.

The size of the rooms will, of course, be regulated by convenience and by the purpose for which they are intended. As a rule, the height should not be less than ten feet, but extreme height is not desirable unless means are provided for carrying away the hot and impure air from the upper part before it condenses and falls again. A window opening top and bottom and placed in an external wall is essential in every inhabited room, and as far as possible a due northern, and therefore sunless, aspect is to be avoided. Besides the window, every room should have an open chimney, or at least some permanent and effectual means of ventilation not entirely under the control of the occupant. Rooms beneath the ground level are liable to damp, stagnation of air, and deficiency of sunlight, and should never be used as bedrooms. central hall or staircase of the full height of the house, with ventilating windows at the top, is useful in promoting free circulation of air throughout the building.

For the sake of securing a share of sunlight in every room, it is preferable that rows of houses should run north and south, and that square buildings should

have angles in those directions.

Every house should have both in front and rear an open space at least equal in length to the height of the building, in order to allow sufficient light and ventilation for the rooms on the lowest floor. The arrangement of houses in courts or small

closed squares is objectionable, since the air cannot circulate freely. Cul de sac courts and courts entered under archways or by high-walled alleys are on this account to be deprecated. Nor should buildings be so placed in relation to each other as to prevent free access of sunlight and air. The space in front of houses should be of a minimum width of 36 feet for a street of more than 100 feet in length which itself must be at least 24 feet wide (in London 40 feet). Model Bye-laws provide for at least 150 superficial feet at the rear of houses. The London Building Act crystallizes the principle that the amount of space at the rear shall be proportionate to the height of the building, and it contains a number of regulations as to building in new and old streets. Streets should not be of less width than the height of the houses in them, and a line drawn from the ridge of the roof to the foot of the wall opposite, front and rear, should not make an angle of more than 45° with the ground. The practice of building over backyards which prevailed in Central London last century is wholly pernicious, for such a system leads to backto-back houses.

Back-to-back houses are dwellings in which through ventilation cannot be obtained, and the absence of curtilage at the back is in itself a serious inconvenience. They have usually dark and ill-ventilated sculleries, pantries, storerooms, or living rooms, and it is often difficult to secure the satisfactory disconnection of sink wastes. The closets have to be built in blocks, often at an inconvenient distance, and in a public position. Tatham has shown that in such houses mortality rates are higher, by comparing the death rates in Salford (26.1 per 1,000) where there were no back-to-back houses with the death rates where 18 per cent. of the houses were of this character (29.1 per 1,000), and where

50 per cent. were of this character (37.3 per 1,000). Many similar returns have been made, and the **L.G.B.** do not now sanction bye-laws which permit of the erection of back-to-back houses.

The interior of a house will of course be arranged according to the taste and circumstances of the owner. Walls may suitably be rendered in plaster (lime mortar, or cement mortar) and finished by lime-washing, as in outbuildings, whitewashing with "whiting" (chalk, size, and alum), distempering, painting in oils, or papering. Painting of wood, iron-work, etc., has many advantages, and tends to cleanliness and preservation. It may frequently be washed. New papers should not be pasted over old ones, as this leads to filth, and, in tenement houses, to verminous conditions. The wall should be "stripped" and cleansed, and, if necessary, rendered with size before the new

paper is put on.

New houses are constructed under the supervision of the S.A., and in accordance with bye-laws prescribing all details necessary for sanitary purposes. Old houses are very often wanting in many points now regarded as essential for the health of the inmates, and it becomes a matter of considerable difficulty to decide how far it is practicable to enforce modern requirements in such houses. If the defects are such as to be prejudicial to the health of the inhabitants, it is the duty of the S.A. to require the necessary repairs or alterations to be carried out, and it is incumbent upon the M.O.H. and Inspector of Nuisances to bring such cases to the notice of the Authority for that purpose. Very often it is not possible to effect such alterations as would be considered satisfactory in new premises, and a certain degree of latitude has to be allowed. No hard and fast line can be drawn between that which is essential and that which is desirable, and a similar difficulty

frequently arises in deciding whether a house is or is not fit for habitation. Sometimes the question is only one of degree, in matters regarding which fixed standards are impossible. Several causes may combine to render a house unhealthy.

Water-supply to houses .- The Public Health Act, 1875, and the Public Health (Water) Act, 1878, give powers to S.A. to enforce supply of water to indi-

vidual houses, groups of houses and villages.

Under the 1875 Act, the S.A. can only insist on a supply to a house if the cost does not exceed the water rate authorised by any local Act in force in the district, or 2d. per week, or such other cost as the L.G.B. may, on the application of the S.A., determine to be reasonable.

The Public Health (Water) Act, 1878, makes it the duty of a Rural S.A. to see that every occupied dwelling-house in their district "has within reasonable distance an available supply of wholesome water," but, unfortunately, the interpretation of "reasonable distance," and "an available supply," has in the past often been so liberal that the Act has been largely inoperative.

The Metropolis Water Act, 1871, gave power to the Board of Trade to make regulations controlling the supply to houses by water companies. The Public Health (London) Act, 1891 (s. 48), states that a dwelling-house shall be deemed unfit for human habitation if it is without a proper and sufficient supply of water. New houses in London have to be

certified in this respect.

Ventilation.—One of the most important matters in connection with buildings, particularly dwelling-houses, is ventilation. Taking the average frequency of respiration as 16 per minute, and the average volume of each breath as 30 cubic inches, the volume of air expired is about 17 cubic feet per hour,

and if it were possible to carry this immediately away (as is done in the case of divers) a fresh-air supply of 17 cubic feet per hour would suffice. In practice, the expired air cannot be prevented from mixing with the rest, and we have to determine how much fresh air is needed to dilute down the respiratory impurities to a permissible point, assuming that the whole of the air in the room is equally contaminated.

When the carbonic acid in the air of a room exceeds 0.6 volume per 1,000, the air begins to be perceptibly "close" to a person entering from the open air, and it is usual to adopt this as a standard of purity, and to aim at introducing a sufficient supply of fresh air to keep the carbonic acid at or below this point.

A thousand cubic feet of fresh air, containing 0.4 cubic foot of carbonic acid (see page 3), can therefore take up 0.2 cubic foot more without exceeding the limit of 0.6. As already stated (p. 10), an average man gives off 0.6 cubic foot of carbonic acid per hour,

so that $\frac{0.6}{0.2} \times 1,000$, or 3,000 cubic feet of fresh air per hour, are needed to maintain the standard of purity. If other values be adopted for any of the preliminary data, the volume of fresh air needed will be proportionately modified. In general terms, if

A = no. of c.f. of CO_2 in 1,000 c.f. of fresh air = 0.4, C = proposed max. limit of CO_2 in the air of the room,

R = no. of c.f. of CO₂ given off per head per hour = 0.6, F = volume of fresh air required per head per hour to maintain the standard C.

Then $F = \frac{R \times 1,000}{C - A}$, and for *n* persons the required volume becomes $\frac{n \times R \times 1,000}{C - A}$, or $\frac{n \times 600}{C - 0.4}$

Example I.—What hourly supply of fresh air is needed for a hall containing 100 persons, in order that the air of the

hall may not contain more than 0.7 part of carbonic acid per 1,000 volumes of air?

Answer:
$$\frac{100 \times 600}{0.7 - 0.4} = 200,000$$
 cubic feet.

Example II.—If 12,000 cubic feet of fresh air per hour are supplied to a room containing 10 persons, what proportion of carbonic acid will be found in the air of the room?

The formula $F = \frac{n \times 600}{C - A}$ becomes

$$12,000 = \frac{10 \times 600}{C - 0.4}$$

Hence C = 0.9 per 1,000 cubic feet.

It is of course assumed that an equal volume of impure air is removed to make way for the fresh air.

Carnelley, Haldane, and Anderson found that the determination of carbonic acid alone is not a satisfactory measure of the impurity of the air in other respects. There is no definite connection between the number of microbes and the amount of CO_2 , but in general a high proportion of CO_2 is accompanied by a high proportion of organic matter. Nor does it of course include other impurities, such as sulphur.*

As gas forms in burning about half its volume of CO_2 , the products of combustion (on a standard of 20 grains of sulphur per 100 cubic feet) would contain less than 0.5 grains of sulphur to 500 litres of CO_2 . If the sulphur were oxidised to SO_2 this would correspond to about one volume of SO_2 to 1,500 volumes of CO_2 , which would not produce any perceptible effect in the atmosphere of a room. The sulphur contained in crude unpurified lighting gas is present chiefly as H_2S , about one-tenth being present in other forms (as CS_2). Gas which is purified from carbon bisulphide is greatly superior from a hygienic

^{*} The English gas companies are usually legally required to supply gas containing less than 20 grains per 100 cubic feet. In London hitherto the maximum has been 17 grains in summer and 22 in winter. In some provincial towns as much as 30 is allowed.

standpoint to gas which is only purified from sul-

phuretted hydrogen.

As regards the more specific diseases, the injurious effects of deficient ventilation are obvious. Crowding in a common atmosphere has long been known to be disastrous in typhus and erysipelas, and more recent experience has established this in respect of diphtheria, influenza and other maladies, both as regards transmission of infection and intensification of type, or retardation of recovery. Ventilation has greatly lessened phthisis in barracks and other institutions, and with cleanliness it has banished typhus from prisons, and hospital gangrene and erysipelas from hospitals. The success of the open-air treatment in camp hospitals and in sanatoria for phthisis has the same significance. It is evident that all diseases in which the breath is infectious must necessarily be more readily transmissible when the expired air is rebreathed in a concentrated form.

Apart from this there is abundant evidence of the gain in comfort, general health and longevity under conditions of adequate fresh air supply, and conversely of higher death rates (general, and from bronchitis and other non-specific causes), impaired health, anæmia, headache, and depression among those whose lives are largely spent in ill-ventilated rooms. Some of these effects are felt at once by susceptible persons on entering a room in which others work without discomfort, but the precise relation between them and the several physical conditions which impurity of air connotes, is not clear. Experiments have been made, with conflicting results, by removing one or other of these conditions while leaving the rest to increase in a degree far beyond that met with in practice. Thus, in experiments by Haldane and Lorrain Smith carbonic acid alone did not appear to disturb respiration until it had risen to about 300 parts per

10,000; nor did diminution of oxygen do so until the proportion had fallen from 21 to 12 per cent. of the air. But if CO₂ be removed, the accumulation of organic matter, which is usually regarded as the most harmful factor, seems to produce little effect. Humidity alone cannot be held responsible, for it may approach saturation even in the open air. Haldane is inclined to attach some importance to the depressing influence of the smell which accompanies want of ventilation.

There is, however, no doubt as to their combined effect, and it must be remembered that experimental conditions do not attempt to imitate those of daily life, in duration or in attendant circumstances such as exercise. The one is rather a question of acute

poisoning, the other of chronic poisoning.

Air space. With ordinary appliances, and under the average atmospheric conditions of the climate of England, the air of a room cannot be changed more than about three times per hour without causing an inconvenient amount of draught. Hence, in order to supply 3,000 cubic feet of fresh air per hour, we should have an air-space for each person of at least 1,000 cubic feet. In calculating this, only the available air-space should be taken into account, deduction being made for furniture and other solid objects. It must be remembered, too, that air stagnates in nooks and corners, and the useful part of the air-space is that in which the air moves freely. For these and other reasons it is more difficult to provide for the efficient ventilation of a small room than of a large one, the air-space per head being the same in each case; but on the other hand the transpiration through walls, ceilings, and floors is relatively greater in small rooms, owing to their higher ratio of surface to air contents. extent and practical importance of this transpiration is proved not only by CO, determinations but by the

experience gained in sulphur fumigation where the leakage of the gas is rapid and demonstrable in spite of every precaution to prevent its escape through

openings.

A further reservation is necessary. Above a certain height air-space ceases to be useful, since it does not assist ventilation; and in such circumstances increase in height cannot compensate for inadequacy in length and breadth, that is, floor-space. There may be overcrowding and even suffocation in the open air. It is a common rule, in rooms of moderate size, to disregard height above 15 feet in calculating air-space, but the limit varies with the floor area. In the special rules for certain dangerous processes under the Factory Act, the requirement of at least 400 cubic feet of air-space per head is coupled with a condition that height above 14 feet is not to be taken into account. In hospitals and schools there is a standard of floor-space per bed, apart from mere cubic capacity.

AVERAGE RESULTS OF ANALYSES OF AIR IN SLEEPING ROOMS BETWEEN 12.30 AND 4.30 A.M. (Carnelley, Haldane, and Anderson.)

Cubic Feet per Head.	Tempera- ture (Fahrenheit.)	Carbonic Acid per 1,000 vols.	Organic Matter (vols. of Oxygen required per million vols. of air).	Microbes per litre.
100 100	550	1.15	15.1	80
100–180	55°	1.15		
180-260	54°	1.07	15.1	49
260-340	53°	1.03	11.8	32
340-500	57°	0.92	8.4	42
500-1,000	54°	0.86	5.6	6
1,000-2,500.	53°	0.67	3.9	9
2,500-4,000	57°	0.79	-5.0	13

The experiments summarised above were made in Dundee. They tend to show that about 1,000 cubic

feet per head are sufficient, and, indeed, the impurities were somewhat greater when very large space was provided, owing probably to imperfect ventilation.

The same observers compared houses of one, two, three, and four rooms, in regard to impurity of air and mortality from different causes among the inmates. They found that the smaller the tenement the greater was the impurity of the air, as shown by increase of carbonic acid, microbes, and organic matter. The death-rate increased in like manner, and especially among children. Comparing one-roomed with four-roomed houses, the general death-rate was doubled, and the death-rate at ages below five years quadrupled. The increase was most marked in diarrhea, measles, whooping cough, bronchitis, and pneumonia. The same general conclusions are well illustrated in the following table:—

FINSBURY. DEATH RATES (per 1,000 living), 1905-6.

Tenements P la	Census Popu- lation,	All Causes.		Principal Zymotic Diseases.		Phthisis.		Respiratory disease (excluding Phthisis).	
	1901.	1905.	1906.	1905.	1906.	1905.	1906.	1905.	1906
One Two Three Four or more	14,516 31,482 21,280 33,185	32·7 19·5 12·3 6·6	39·0 22·5 14·8 6·4	3·4 2·8 1·8 0·7	6:4 5:5 2:6 0:8	3·5 2·1 1·3 0·8	3·4 2·3 1·4 0·9	6·4 5·2 2·8 1·4	8·3 4·8 2·9 1·2
The Borough	101,463	18.9	20.7	2.0	3.7	2.1	2.3	4.4	4.0

The actual air-space obtained in practice frequently falls very far short of 1,000 cubic feet per head, even in large private houses. In registered common lodging-houses the minimum is usually 300 cubic feet per head, but varies according to the regulations of the different local authorities, and in working-clas

dwellings it is frequently much less still. The London Education Committee require 120 cubic feet per head as a minimum in primary schools (96 cubic feet in infant schools). The lowest standard expressly sanctioned by law is that for canal-boats, namely, 60 cubic feet per head for adults, and 40 for children. It is important to bear in mind that air-space is chiefly valuable as making ventilation possible without "draught," and not as a store of fresh air. The foul air of an average occupied bedroom in the morning is a proof that proper ventilation has not been provided, whatever the magnitude of the cubic space. No bedroom, however large, ever contains a sufficient supply of fresh air for the night, but the larger the room the more readily can efficient ventilation be contrived. A thousand cubic feet of space per person is ample, if the air is changed three times per hour, but in itself a thousand cubic feet of air is a sufficient supply for one person for twenty minutes only, if the CO, is to be kept down to 0.6 per 1,000.

The objection to draught is largely dependent upon its chilling effect, so that in hot weather, or when the fresh air is warmed artificially, the air of a room may be changed more rapidly than three times an hour, and in that case a somewhat smaller cubic space may

be sufficient.

Although the carbonic acid of expired air diffuses readily, the organic matter is less volatile, and hangs about in invisible clouds unless dissipated by local currents. Hence the composition of air in an occupied room is not uniform, even at the same level, though for practical purposes it may usually be assumed to be so. In a large room occupied by many persons both the local currents and the foci of pollution are numerous, so that the composition of the air is much more uniform than in a small room

occupied by one person. If from any cause the air passing out of the room is not the most impure, the remaining air will be found to exceed the theoretical degree of impurity. The outlet should be at or near the highest point, since expired air, being warm and moist,* is lighter than ordinary air, and ascends at first to the upper part of the room.

The fresh-air inlets should be arranged as far as possible so as to avoid draughts impinging upon the occupants, to secure diffusion of the current, and to counteract the tendency of the fresh air (if cool) to sink to the lowest level. This may be effected by giving it an upward direction on entering, so that before descending it mixes with the air of the room.

In measuring a room for the purpose of calculating the air-space, recesses and projections may be taken separately, and then added to or deducted from the total. A similar procedure will obviate any difficulty in cases where the ceiling is not horizontal throughout. †

The ventilation of a room or building may be "natural" or "artificial," the latter term being applied to mechanical means for extraction or propulsion of air, and the former to all other forms of ventilation.

Natural ventilation is dependent upon wind,

* The vapour of water is lighter than air. The volumes occupied by 18 parts by weight of water vapour, 32 of oxygen, and 281of nitrogen, are equal; these proportions being, of course, those of their respective molecular weights.

† Very often only right angles are met with, so that the floor area = length × breadth, and the cubic capacity = area × height. The following rules cover most of the other conditions likely to occur in

Area of triangle = Base \times Height $\times \frac{1}{2}$. (Euclid I. 41.) Area of an irregular four-sided figure. Divide into two triangles, by joining two opposite angles. Find area of each triangle, and add together. Area of a circle = (Diameter) $^2 \times 0.7854$.

Area of a semicircle = (Diameter)² \times 0.3927.

Area of segment of circle = Chord \times Height $\times \frac{2}{3}$

Area of an ellipse = Long Diameter × Short Diameter × 0.7854. Capacity of a dome = Area of base \times Height $\times \frac{2}{3}$. Capacity of a cone or pyramid = Area of Base \times Height $\times \frac{1}{3}$.

upon local currents due to differences between internal and external temperatures, and still more upon local currents determined by artificial heat, as in chimneys or ventilating gas-lights, although these last may perhaps more properly be considered as artificial means of ventilation.

Air finds its way into a room by the doors and windows, however close-fitting, and to a less extent through brick or stone walls, ceilings and floors. These leakages are very small if a proper fresh-air inlet has been provided. Such inlets may be window openings, or ventilating fire-places or stoves, or special openings (ventilators). Windows should be made to open at top and bottom, but it is desirable to have in addition some means of admitting air at all seasons without draught. This may be done by having double windows, or one or more double panes, and providing an opening at the bottom of the outer and at the top of the inner one, so as to admit fresh air in an upward current. Louvred panes and several modifications of "hit-and-miss" ventilators are also available. A simple and effectual device (Hinckes-Bird's) is to place a board beneath the lower sash, propping it up and filling in the space below while providing a way for an upward indraught between the two sashes. Another plan is for the upper part of the window to be hinged below, so as to fall forward and admit air into the room as by a Sheringham valve.

Ventilating fire-places have an air-chamber at the back of the grate, heated by the fire. Fresh air brought by a pipe from outside is warmed in the hot chamber, and then passes into the room. There are many varieties, among which Galton's is the earliest and one of the best. Ventilating stoves are in use, the fresh air entering through a coiled pipe exposed

to the heat of the stove.

Special ventilators (Fig. 7) are of varied types, from simple openings through the wall to the most complex arrangements. Perforated bricks cause little draught, the holes being conical with the wider ends inside, so that the air stream slackens as it passes through the wall, and is diffused upon entering the room. Sheringham's ventilator is a small vertical flap door in the wall near the ceiling, balanced by a counterpoise and hinged below so as to fall forward towards the room; it is cased in at the sides and front so that the current can only pass upwards. Tobin's ventilator consists of a large upright tube rising five feet or more from the floor; fresh air from outside passes up this tube and into the room. Ventilating cornices are made,

consisting of a double channel of perforated metal, cold fresh air being brought into the room by the lower channel, and vitiated air being received into the upper channel and carried to the chimney or other outlet; another plan is to carry a perforated inlet tube along the cornice on three sides of the room, and a similarly perforated outlet tube along the fourth side. A hollow perforated metal beam, interrupted by a diaphragm




Fig. 7.—Ventilators.

A A, Tobin's; BB, McKinnell's; CC, simple ventilating shaft, with cap; DD, Sheringham's (acting as inlet); BE, Sheringham's (acting as outlet).

at the centre, may be taken across the room, one-half serving as an inlet and the other as an outlet, according to the direction of the wind. McKinnell's ventilator consists of a larger outer and a smaller inner tube, carried upwards through the ceiling; fresh air passes in between the tubes, and is dispersed on entering the room by a horizontal flange upon the inner tube; the inner tube, which is equal in sectional area to the inlet, and projects beyond the other both above and below, carries off impure air. Another device is a vertical shaft divided longitudinally into two or four separate channels, along which

Currents in opposite directions will always establish themselves. Workrooms may be ventilated by means of a perforated ceiling of zinc or paper, over which is a chamber open to the outer air on every side. An open-tiled roof, or one not underdrawn, allows much ventilation.

Some of the preceding are intended to act as inlets, but all are liable to become outlets. Other kinds of ventilators are, like chimneys, designed as outlets only. An opening may be made from the room into the chimney near the ceiling, reflux of smoke being prevented by a number of little valves of mica (Boyle's valves) which, in the absence of up-current, close by their own weight, owing to the forward tilt of the latticed framework to which they are attached. Ventilating gas-lights are made, in which the products of combustion are collected by means of a bell cover or glass globe, and conveyed away by a tube; this tube can be surrounded by a larger one, and the heated space between the two made to act as a further extraction shaft. A similar use may be made of chimneys, the smoke being contained in a central metal tube, leaving a heated airspace all around it, which serves for extraction. Or outlet pipes from the higher part of the room may be carried upwards for some distance in or close to the chimney so as to utilise its heat in promoting an up-current. Extraction tubes may of course be carried from any part of the room, with or without taking advantage of these various convenient sources of heat, and they may be provided with cowls.

The size of openings required for the efficient ventilation of an occupied room under average conditions is about 48 square inches per head, namely, 24 as inlet and 24 as outlet, but, according to Parkes, no single inlet should exceed 60 and no

outlet 144 square inches.

Supplies of fresh air should be taken from suitable points free from risk of contamination, and, if necessary, the air may be screened through metal gratings, canvas, cotton wool, &c. It may be washed by passing it through spray or a film of water, or by being made to impinge upon a tray containing water, and it may be warmed by means of ventilating fire-places or stoves, or by coils of hot-water or steam tubes at the inlet. If warmed, there is advantage in introducing fresh air at or near the floor level; but if cold, it is necessary to avoid draughts which might impinge upon any

person in the room, and therefore the air is delivered at a higher level and with an upward direction.

Outlets should be as near the chimney as possible; the air in that part is the warmest, as well as the most impure, so that the worst air is removed with the maximum velocity.

Any barometric pressure P may be regarded as produced by the weight of a column of air of uniform density and of height H. "Montgolfier's Rule" affirms that air under pressure will rush into a vacuum with a velocity equal to that acquired by a body falling from the corresponding height H, which velocity may be determined by the formula $V_2 = 2gH$, g being the acceleration due to gravity (32.18 feet per second), and V the velocity required. Thus under standard conditions of barometer and thermometer H is about 5 miles, or 26,400 feet, so that V₂=2 \times 32.18 \times 26,400, and V = about 1,300 feet per second.

If, however, the air under pressure P passes not into a vacuum but into space containing air under a low pressure p, the formula requires modification, the velocity being dependent upon the difference between P and p. It becomes necessary to calculate the height h of a uniform column of air (of the same standard of density as was adopted in determining H), which

would give the pressure p.

The formula then becomes $V_2 = 2g$ (H-h), and in this form

it is applicable to problems of ventilation.

For example, it may be required to determine the theoretical velocity of updraught in a vertical* flue x feet long, leading from a room containing air at a temperature to into the open air of lower temperature To. An inlet for cold air is presupposed. The column of heated air in the flue may be regarded as balanced against an equally high column of cold air outside in a sort of U-tube; and as yielding to the greater weight of the colder and less rarefied column, and the (unequal) weights of these two columns correspond to P and p. The barometric pressure at the level of the top of the flue may be neglected, since it affects both columns alike. Then taking the density of the outside air as the standard, the height x may be put for H in the formula, and it only remains to determine h, the height of the heated column when reduced to the same. standard density. Air expands with heat to the extent of '002'

^{*} The vertical height only must be taken, even if the course of the flue is oblique or bent.

of its volume for each degree Fahr.,* and hence 1 volume at T° becomes, at t°, 1 + .002 (t-T), so that the column of hot air of height x at t° corresponds to a height $x \times \frac{1}{1 + .002}$ (t - T) at the standard temperature T. This, then, is the value of h, and

$$V_2 = 2g \left(x - x \times \frac{1}{1 + \cdot 002 (t - T)} \right).$$

Since $\sqrt{2g}$ is approximately 8.0, the formula may conveniently be written.

$$V = 8 \sqrt{x - \frac{x}{1 + .002 (t - T)}}$$

If, as an example, the external and internal temperatures are respectively 40° F. and 60° F., and the height of the flue 20 feet,

$$V = 8 \sqrt{20 - \frac{20}{1 + \cdot 002 (60 - 40)}}$$

Hence the theoretical velocity of draught is 7.1 feet per second, and this multiplied by the sectional area of the flue gives the

number of cubic feet discharged per second.

The theoretical velocity is subject to correction for friction, which varies directly as the length of the tube L and the square of the velocity, and inversely as the diameter D, that is V_2L

as $\frac{V_2L}{D}$. In practice an allowance of $\frac{1}{4}$, or even $\frac{1}{2}$, has to be

made for friction, which is also increased by angles in the course of the flue.

The following formula by De Chaumont is convenient for ascertaining the relation between the size of the opening and the hourly delivery of air:—

 $D = 200 \times \Phi \times \sqrt{.002 \times x \times (t - T)},$

where D = delivery of air, in cubic feet per hour,

 Φ = the sectional area of the tube, or of the inlet or outlet, in square inches,

x = height of heated column of air, in feet,

t and T = the internal and external temperatures respectively.

Thus, if the heated column be 20 feet in height, the internal and external temperatures 60° F. and 40° F., and the required

* Strictly speaking, this is accurate only when T = 32 F., but the error is immaterial for the present purpose.

delivery 6,000 feet per hour, the formula will give the sectional area necessary :-

> $6000 = 200 \phi \sqrt{.002 \times 20 \times (60 - 40)}$ $\phi = 33.56$ square inches.

The linear velocity with which air passes through an opening may also be determined experimentally by an anemometer, which should be placed, not at the centre, but about 2 of the diameter from the side of the opening, so as to obtain the mean velocity. Very delicate anemometers are made, consisting of four light metal vanes mounted obliquely upon a common axis, the revolutions of which are indicated automatically upon dials. The registering apparatus can be thrown in or out of gear by a spring catch while the vanes are revolving, so that the time can be determined accurately. The linear movement observed during one minute, or less, is readily converted into velocity per hour or second, and this, multiplied by the sectional area of the opening, gives the cubic discharge.

Artificial ventilation involves either propulsion or extraction of air, the establishment of a plenum or vacuum in the room to be ventilated. Fresh air may be driven into a room or building by revolving fans or other motors; vitiated air may be extracted by means of fans, steam jets, or a heated outlet shaft. The fans used for the purpose have vanes set obliquely upon a rapidly revolving axis, and work in a closed box or tube so as to propel a constant current of air of any required volume and velocity, somewhat after the manner of an Archimedean screw. Steam allowed to escape in a powerful jet into a flue is capable of carrying with it 200 times its own volume of air. Gas burnt at the foot of an extraction shaft may be made to carry off a thousand times its own volume of air.

Heat is by far the most generally employed motive Ordinary fire-places, chimneys, and ventilating gas-lights act in this way. Special extraction shafts, with gas jets or furnaces at the bottom, are common. Mines are ventilated by means of a furnace at the foot of the upcast shaft, air being drawn down another shaft and made to traverse the whole of the

workings on its way to the upcast. A similar plan is adopted in steamers, the upcast being a space around the funnel and boilers, a contrivance which has been applied also to house chimneys by confining the smoke, etc., to an iron tube passing up in the centre of the chimney shaft. Extraction is necessary where dust and fumes have to be drawn away at the point, of origin, but is attended with some disadvantage from the difficulty of controlling the sources from which the supplies of fresh air are drawn, since it rushes in through all available openings. Provision may, however, be made at suitable points for free admission of pure air, which may be warmed, cooled, moistened, washed, or screened as it enters. When propulsion is adopted, the air is taken from one source only, and, being more under control, is more readily subjected to such heating or other treatment as may be needed.

The advantages of artificial ventilation are its constancy under all conditions and the facilities which it affords for regulating the source and amount of fresh air, and the preparatory treatment as to temperature, moisture, and purification. Natural ventilation is, of course, less costly, but is inconveniently subject to atmospheric conditions, and is inadequate for theatres, factories, schools, and other large buildings. The following averages are based upon a number of observations (18 to 39) in schools by

Carnelley, Haldane, and Anderson.

eritoria bestellativo chi proprie il	Natural Ventilation.	Mechanical Ventilation.
not bush to thing point out travel to	THE HOLD STATE	SCHOOL BILLIAN
Per cent. of windows open	22	3
Cubic feet of air space, per head .	168	164
Temperature	55.6° F.	62° F.
Carbonic acid (per 1,000 vols.) Organic matter (vols. of oxygen re-	18.6	12.3
quired per million vols. of air)	16.2	10.1
Microbes (per litre)	152	16.6

The difference is still more striking if only the excess over the impurities present in outside air is taken into consideration. It was found that mechanical ventilation kept the composition of the air fairly constant in all parts of the room, while natural ventilation often allowed of local stagnation.

In an ordinary room the chimney is the most important outlet when a fire is burning, air passing up with an average velocity of 4 or 5 feet per second. In the absence of a fire the chimney still acts in some degree as an upcast, and ought on no account to be closed, unless a downward current establishes itself temporarily, and causes annoyance by its sooty smell.

Warming is closely connected with ventilation. The principal methods in use may be classed as open fires, closed fires and stoves, hot air, and pipes containing hot water or steam. Some of these heat the room by radiation, warming the surfaces while leaving the air in it cool; some by convection, warming the air of the room and thus making it drier; some in both ways. Again, some (e.g. open fires) carry off large volumes of air besides their own products of combustion; some (e.g. gas or other stoves) only little; some (hot pipes) none at all; while others, and especially fender stoves, do not ventilate but add to the impurity of the air.

Ordinary **open fires** are extravagant in regard to fuel, and it is estimated that only about one-eighth part of the potential heat is utilised in warming the room, the rest being lost in unconsumed smoke and cinder, and in the hot gases passing up the chimney. The heating effect is very unequal in different parts of the room, being chiefly confined to the vicinity of the grate, and cold currents pass along the floor. They are, however, most effective ventilators, and present a more cheerful appearance than stoves or hot pipes.

The principal practical points in regard to the

construction of open fire grates are summarised by Teale as follows:—(1) use as much firebrick and as little iron as possible; (2) the back and sides should be made of firebrick; (3) the back of the fireplace should lean over the fire, the "throat" of the chimney should be contracted; (4) the bottom of the fire should be deep from before back; (5) the slits in the bottom grating should be narrow; (6) the bars in front should be narrow; (7) the space beneath the fire should be closed in front by a close-fitting iron shield, or "economiser," thus excluding cold air almost completely from the bottom of the fire, so that the fuel as it sinks is entirely burnt up, leaving only a fine ash, which drops into the space beneath. Another means of preventing waste of fuel is to make the bottom of the grate of solid fire-clay, as in the various "slow combustion" grates; and the same principle may be applied to ordinary grates by laying an iron plate upon the bars at the bottom of the grate.

Coke is the residue, consisting almost entirely of carbon and mineral ash, left in the retorts when all the combustible gases have been driven off by heat in the process of gas-making. It is smokeless and less costly than coal, but it is difficult to light, and makes a dull fire. The prevention of smoke, and the complete utilisation of the gases and other products obtainable from coal by distillation, are important

public advantages.

Gas-fires have much of the bright and attractive appearance of open fires. The gas is mingled with a considerable volume of air before ignition, as in a Bunsen burner, and the intensely hot but non-luminous flame impinges upon pieces of asbestos or iron, which become red or white-hot, giving out light and heat. Simpler forms are also made with ordinary luminous burners, the light and heat from which are thrown forward by

metal reflectors; oil-stoves are made on the same principle. In any case an adequate flue to carry off the fumes is essential.* A gas-fire burns from ten to twenty, or more, cubic feet per hour. It is economical and cleanly in use, creates no smoke, and can be lighted, extinguished, or regulated in a moment. Ventilating gas-stoves are made, but the ordinary gas-fire, unless it be composed of incandescent fragments of asbestos in an open grate, is less effectual than an open fire in promoting ventilation. If the flue is too small, as is very often the case, the air becomes dry and oppressive, and there is risk of carbonic

oxide escaping.

Stoves for coal and coke, as used in England, are commonly made of cast iron, the smoke and products of combustion being conducted by an iron flue to the chimney, or to the outside air. The fire may be more or less open, or hidden from sight, but in any case the waste of heat is less than in ordinary fire-places, the sides of the stove and also the flue contributing to the heating power. As a means of extraction of vitiated air a stove is inferior to the ordinary open fire, although there is no difficulty in utilising the hot flue for the purpose of producing an up-current in a chimney. There are also many forms of ventilating stoves, fresh air being brought in through special pipes, which in their course are exposed to the heat of the stove; in the "Calorigen" a coil of the fresh air inlet pipe is placed inside the stove. Stoves tend to make the air hot and dry, an objection which may be met by placing vessels of water upon them. A more serious difficulty is a burnt smell which is apt to result from their use, and a tendency to give off carbonic oxide,

The consumption of gas is relatively large and not limited to hours of darkness. Besides CO₂, sulphur acids and traces of carbonic oxide are given off. The condensation of fumes, which is claimed for some gas stoves, is at best only partial.

the source of which is a matter of some doubt. It has been attributed to oxidation of the organic matter in the air by contact with the heated iron, this being also assigned as the cause of the burnt smell. Oxidation of the carbon in the cast iron is another suggested explanation, and it has been found experimentally that red-hot cast iron is pervious to carbonic oxide, and even reduces carbonic acid to carbonic oxide, so that the gas found in the room may originate in the fire itself. Wrought iron offers far greater resistance. The joints, both of the stove and the flue, should be made strong and air-tight, and in order to minimise the risk of giving off carbonic oxide, it is desirable to line the stove with fire-clay, to coat it with silicate, or to use earthenware or wrought iron in place of cast iron. Stoves can only be approved from a sanitary point of view if they (a) are provided with proper chimneys; (b) are of such a pattern that they cannot become red-hot, or otherwise contaminate the air; and (c) are supplied with fresh air directly from the outside.

Hot air is advocated by some authorities as the best mode of warming houses, as well as public buildings. This may be done upon the large scale by warming the fresh air in the basement, and conveying it thence to all parts of the house along the passages or by special channels. The same end is attained in some degree by placing a stove or fire-place in the hall. Warm air may be supplied to single rooms by the ventilating grates or stoves already referred to, or by allowing the fresh air to enter through a coil of hot pipes. The air can be moistened and, if need be, purified by filtration or washing. Warming can also be effected by means of hot water or steam circulating in a system of closed pipes. The pipes may be carried round one or more sides of a room, near the floor, either open or concealed behind a skirting-board. Several turns or coils may be used at points where more heating power is required, and it is convenient to warm the fresh air at the inlets by the same means.

Hot water is usually employed for heating purposes at a temperature below 212° Fahr. The circulation is dependent upon the water being hotter, and therefore lighter, in the tube through which it leaves the boiler than in that which brings it back. Hence the boiler is placed at the lowest point, and the centrifugal and centripetal pipes are respectively connected with the top and bottom of the boiler. A feeding cistern, and vents for the escape of air and steam, are provided at the highest point. About twelve feet of four-inch iron pipe are allowed for every thousand cubic feet of air-space to be warmed. The system should be so planned with valves and connecting pipes as to allow of "short-circuiting," so that the heat may be turned on or cut off from any part at will.

Hot water may also be used under pressure at temperatures considerably exceeding 212° Fahr., on Perkins' system. Strong half-inch iron pipes are used, and, instead of a boiler, part of the pipe is carried through the fire. Steam may be used at high or low

pressure in like manner.

Inspection of houses .- The following schedule includes the principal points that may require attention. The headings are numerous, but are meant to be chiefly suggestive, so that no important detail may be overlooked. A very few notes will often suffice in practice. As in all other investigations, it is wise to make explicit written notes on the spot, and to insert negative as well as positive results of inspection.

Address. Occupier.

Owner .- Name and address.

of each

room.*

Site.—Elevation, aspect, slope. Proximity to hills, valleys, watercourses. Nature of soil. Dryness.

Surroundings .- House detached, semi-detached, in a row, back to back. Access of light and air. Obstruction by trees, high grounds, or other buildings. Open space at front, back, sides [stating area]. Proximity [stating distance in feet] of stables, cow-sheds, pig-styes, manure heaps, foul ditches, stagnant water, offensive accumulation, offensive trades, or other sources of effluvia.

Yard and outbuildings .- Condition as to cleanliness, paving,

drainage, area, closets, workshops.

Foundations.—Damp-proof course, dry area, banking of soil against walls. Dryness of basement. Exclusion of ground air. Ventilation of space beneath ground floor.

Walls.-Materials, thickness. Party walls carried up to roof? Dilapidations. Evidence of damp at any part; probable source of such damp.

Roof.—Construction. Soundness.

Floors, staircases, windows, doors, ceilings.—Soundness. Rooms.—Number on each storey, including basement.

Length, breadth, and height.

Windows. Total window-space. Space made to open. Opening top and bottom?

Other means of ventilation. Maintenance of ventilation. Chimney

Heating arrangements,—Grates, stoves, flues. Hot water or steam system. Boilers, bath-geysers.

Drainage-

Construction and course of waste-Sink wastes Bath wastes . pipes.

Lavatory wastes Description and efficiency of traps.

Disconnection. Floor gullies

Soil-pipes. Construction, dimensions, position, course, ventilation.

Gullies. Construction and trapping. Gutters around eaves. Efficiency.

Rain-pipes. Course, trapping, disconnection, destination. Leakage?

House drains. Course, construction, dimensions, grad-

^{*} As regards sleeping-rooms in basement, or "underground rooms," certain further details are necessary, namely, depth of floor below level of adjoining street; height of room; height of ceiling above footpath or nearest ground adjoining; width, depth, and lateral extent of area in front; depth of drain below floor; position of window in respect to steps (if any) bridging across open area; the conditions as to letting and occupancy.

ients, ventilation, means of access (inspection chamber), disconnection from sewer (interceptor). Flushing, cleanliness. Soundness, as tested by smoke, chemical test, or water. Drains under basement.

Cesspool (if any).—Position, construction, dimensions,

ventilation, overflow, watertight.

Sanitary Conveniences.—Number, position, cleanliness, lighting, ventilation. If outdoor, note distance from nearest door or window. If more than one house using same closet, note the number. State also number of occupants.

W.C., trough-closet, slop-closet, hand-flushed closet. Kind of apparatus, efficiency, source of water-supply,

sufficiency of flush.

Pail-, pan-, tub-, or box-closet. Size of receptacle, material, floor-level, means of access, with or without

ashes or other admixture.

Privy-midden, privy with cesspit, earth-closet. Dimensions of receptacle, relation of floor to ground-level, materials, watertight, covered, arrangement for applying ashes or earth to excreta. Connected with sewer?

Household refuse.—Means of storage. Position.

Scavenging. —Mode, frequency, and efficiency of removal of excreta and household refuse. By whom performed.

Water supply.—Public or private, from mains (by tap or stand-pipe), spring, stream, canal, dip-trough, well, rain-water cistern; distance from house, constancy, sufficiency, purity. Note any obvious risk of pollution. If from well, note depth, construction, lining, cover; distance in feet of nearest possible source of pollution, or filth accumulation of any kind. If from cistern, note position, construction, cover, cleanliness, connection with W.C., discharge of overflow pipe.

Cleanliness of premises.—Light. Ventilation. Cleanliness.

Animals kept.—Description. Number. Where kept. Nuis-

ance resulting.

Inmates.—Number of residents. Age, sex, occupation of each. Number of families; distinguishing tenement occupied by each.

Number sleeping in each room, if there is suspicion of

overcrowding. Number sleeping in basement.

Houses vary as to need for inspection according to class and kind. *Private houses* occupied by one family presumably belong to persons of such social

position and habits that the house requires but little enforcement of sanitation. Houses let in lodgings, or tenement houses, on the contrary, require a good deal of supervision. Houses of this nature are such as are let out to separate families, living each in a separate occupancy of one or more rooms. There is no common kitchen or common dormitory. Houses let in lodgings are usually placed under special bye-laws under which they are registered, and controlled in respect of occupancy, overcrowding, sanitation, periodical cleansing, etc. Not unfrequently these houses were built for use as private dwellings, and have become sublet without structural alteration. Under the Customs and Inland Revenue Act of 1890 the landlord of tenement dwellings let at rents not exceeding 7s. 6d. per week may be exempted from inhabited house duty if the medical officer of health of the district certifies the house as sanitarily fit for the purposes to which it is put. By that Act, and by bye-laws for houses let in lodgings, a certain amount of control is exercised.

Common lodging houses are lodging houses pure and simple, with living-room, kitchen and dormitory, open to use by all the lodgers, who are not members of the same family, but are isolated persons, finding it convenient to lodge together. Artisans' dwellings, such as those under the Guinness Trust, the Peabody Trust, etc., are blocks of "model" dwellings, not different in principle from, but better in equipment than houses let in lodgings. Stable dwellings are tenements over stables in "mews." They are objectionable, owing to the occurrence of effluvia arising from stables, manure, etc., and they generally have other defects. Underground dwellings are tenements limited to basement rooms, the ceiling of which is not higher than the level of the street.

CHAPTER VII

SCHOOLS AND SCHOOL HYGIENE

THE general principles as to site, surroundings, construction, and drainage are, of course, the same for

schools as for most other buildings.

Dimensions.—The Board of Education recommend that no school should ordinarily be built to accommodate more than 1,000 to 1,200 children, in three departments, and no single department to accommodate more than 400 children. For departments of this size the most suitable plan is that of a central hall ($3\frac{1}{2}$ –4 ft. floor space per scholar) with (about 7) class-rooms grouped round it. Smaller departments may have class-rooms opening from a corridor. For small schools a large room with one or more class-rooms will be sufficient. There should always be at least one class-room, in addition to the central hall, except in special cases.

Where the site is sufficiently large, open and fairly level, the most economical plan is that of a school on the ground floor, and this is preferable for educational reasons. In any case a school building should, preferably, not be on more than two floors.

For older scholars the Board of Education prescribe a minimum of 10 sq. ft. floor space per scholar. For infants 9 sq. feet, unless there is no corridor or central hall, in which case more than 9 sq. ft. is required. The Board has fixed a standard for higher elementary schools of 15 sq. ft. per scholar if single desks be used, and 12 sq. ft. for dual desks, and for secondary and "special" schools 18 sq. ft.

Lighting.—Every part and corner of a school should be well lighted, and, as far as possible, from

the left side of the scholars; all other windows in class-rooms should be regarded as supplementary or for ventilation. Where left light is impossible, right light is next best; direct front light is in no circum-

stances approved.

Windows should never be provided for the sake merely of external effect. All kinds of glazing which diminish the light and are troublesome to keep clean and in repair must be avoided. A large portion of each window should be made to open for ventilation and for cleaning. The sills of the main lighting windows should be not more than 4 feet above the floor; the tops of the windows should as a rule reach nearly to the ceiling; the upper portion may be made to swing. The ordinary rules respecting hospitals should here be remembered. Large spaces between the window heads and ceiling collect foul air.

Skylights are objectionable in schoolrooms or class-rooms. They may be allowed in central halls having ridge or apex ventilation. The colouring of the walls, ceilings, and fittings, and the size and position of the windows, are especially important in

their bearing on the eyesight of the children.

Natural ventilation is rarely sufficient to maintain the purity of the air in schools, and artificial means, coupled, when necessary, with filtration, warming, and moistening, are therefore desirable though not always attainable in practice. Kerr suggests a limit of 10 parts of CO₂ per 10,000. About 2,000 cubic feet of fresh air per head per hour is required, and it should enter the room at a rate not exceeding 5 feet per second. This is best obtained, as pointed out by Carnelley and Haldane, by using large low-pressure fans, running at low speeds. It is also desirable to have exhaust as well as plenum fans, so that the air pressure in the rooms may be kept low and short circuiting avoided. Whatever method of ventilation

is used, the design should be such as to allow large volumes of air to pass through the buildings from side to side when the windows are open. The window area should be not less than 100 of the floor area and may with advantage be $\frac{1}{5}$. When the windows are suitably designed, carried to the ceiling, easily opened at the top, and intelligently used in connection with two or more warm outlet flues or separate air chimneys from the ceiling and floor level to above the roof, and there are also large open ventilating fire-grates, stoves, and radiators, by which warmed fresh air is admitted, class-rooms can be both warmed and ventilated adequately and economically.

It is well that windows should have both the top and bottom panes arranged to open inwards as hoppers. Gratings in floors should not be provided. Besides continuous ventilation, rooms should be

flushed with fresh air as often as possible.

Warming. - The temperature of class-rooms should be about 56 to 60° F. Where schools are wholly warmed by hot water, the principle of direct radiation is recommended. Stoves are only approved by the Board of Education when provided with proper chimneys, supplied with fresh air direct from outside by flues, and not of such size as will occupy too much floor space, or of such material as will get red hot. Open fire-places have the great advantage of assisting ventilation as well as providing heat. They should be protected by fireguards. A thermometer should hang in each room.

Desks.—The edge of the desk should be vertically, or nearly so, over the front edge of the seat, and about the height of the scholar's elbow. The seat should be at least 9 inches wide, and about the height of the knee. The desk should slope (15°), and the seat be provided with a back: it is desirable to provide for a little horizontal adjustment of either the seat or the

desk, so that they may be slightly separated for writing, and brought together for reading. If a bench is used, 18 inches should be allowed to each pupil.

A simple arrangement contriving the advantages of dual desks without the disadvantages is a continuous desk of half a dozen places with single pedestal chairs (the "Sheffield System"). Many schools in the north of England are now fitted with this kind of desk. It is supported on two plain pedestals screwed to the floor. Each seat has a back, and stands on a single pedestal. The advantages are that the seats are isolated, each child is accessible, and can stand by its seat in its place for drill, etc., and schoolrooms cannot be overcrowded as with dual seats or forms. Each seat occupies about 1,000 square inches of space. Cleaning is much simplified with these seats and desks.

Playground.—The surfaces of the playground should be smooth without being slippery, and formed of materials such as asphalt or granolithic cement, which do not form dust. They should be durable and well drained. Some authorities recommend tar-paving. Gravel, cinder-ashes, shingle, and loose materials are unsuitable. A certain portion should be covered-in for use in wet weather. The whole playground should be open to the sun, and the school site should be of a size giving at least a quarter of an acre to 250 children, and the playground at least 30 square feet per child.

Cloak-rooms other than passages should be provided where damp garments can be dried. Cloak hooks should be fixed 12 inches apart, and gangways between rows of hooks should be not less than 4 feet wide.

Cloak-rooms should be amply lighted and ventilated, with separate means of ingress and egress, and should, where practicable, contain lavatory basins.

Handicraft classes, etc. — Where provision is made for the teaching of cookery, laundry work,

handicraft or similar special subjects, the Board of Education has prescribed conditions. A cookery or laundry school should accommodate from 12 to 18 pupils, each having 25 sq. ft. of floor space. Special arrangements must be made for ventilation, sinks, ranges, stoves, etc. The temperature should not exceed 70° F. Class-rooms for handicraft and science work should have 30 to 35 sq. ft. of floor space for each scholar. Such schools should have their own lavatory and cloak-room accommodation.

Water supply.—All schools should be provided with an adequate supply of wholesome drinking water. If such supply cannot be obtained from the public mains, care must be taken to ascertain that the supply proposed is adequate, is of suitable character, and is

not liable to pollution.

The pipes should be so laid as to be protected from frost, and so that in the event of their becoming unsound the water will not be liable to become fouled, or to escape without observation. There should be no direct communication between any pipe or cistern from which water is drawn for domestic purposes, and any water-closet or urinal.

All water-closets and urinals should be provided with proper service cisterns, affording adequate flush. Any cistern should be watertight, covered and ventilated, and so placed that the interior may be readily

inspected and cleansed.

Closets.—Separate closets are required for boys and girls, and there should be at least one for every 15 to 25 girls and 25 to 40 boys. They should be disconnected from the school proper by a well-ventilated lobby or covered way. Flushclosets are best adapted for the purpose in some ways, and trough-closets are used in many schools. But, probably, the time has come for schools to be supplied with a sufficient number of some simple form

of wash-down closet with separate water-waste preventer cisterns or, if necessary, flushing apparatus for half a dozen at one time. Dry earth systems in schools require careful supervision. It is, of course, possible and necessary to make pails, earth-closets, or even privies serve the purpose in rural districts, but only by constant supervision can they be prevented from becoming a serious nuisance in schools. They should be situated at least 20 feet from the school premises. Urinals, in proportion of 10 feet for 100 boys, should be properly constructed of smooth non-absorbent material, and should have a sufficient water-supply.

Soil drains from schools should be made of glazed stoneware pipes (4 to 6 in. diam.), having a fall of 1 in 30 for 4 in. and 1 in 40 for 6 in., and they must be provided with traps, ventilators, and inspection chambers.

Schools and infectious diseases.—The following is the Memorandum issued by the L.G.B. (1908) concerning this matter:—

- 2. In the Code of Regulations issued by the Board of Education in 1907, the following Article (Art. 57) prescribes, as one of the general conditions required to be fulfilled by a Public Elementary School in order to obtain an annual Parliamentary grant, that—
- "If the S.A. of the District in which the school is situated, or any two members thereof, acting on the advice of the M.O.H., require either the closure of the school or the exclusion of certain children for a specified time, with a view to preventing the spread of disease or any danger to health likely to arise from the condition of the school, such requirement must at once be complied with, but after compliance appeal may be made to the Board (of Education) if the requirement is considered unreasonable."

Article 45 (b) prescribes that if there has been a closure under medical authority, or for any other unavoidable cause, the number of meetings required [400 a year] is proportionately decreased.

(The "medical authority" referred to in this article is not necessarily that of the M.O.H.)

3. The diseases for the prevention of which school closure, or the exclusion of particular children, will be required are principally those which spread by infection directly from person to person, such as diphtheria,* scarlet fever, measles, whooping-cough, epidemic influenza, small-pox,† and rötheln. More rarely, the same question arises in connection with enteric fever and diarrhoeal diseases, which spread not so much by direct infection from person to person as indirectly through the agency of local conditions, such as infected school privies.

4. It will be seen that Article 57 quoted above confers upon sanitary authorities an alternative power with respect to public elementary schools. (A) To cause particular scholars to be for a specified time excluded from attendance, or (B) to require the

school to be closed for a specified time.

5.—A. First, as to exclusion from school of particular scholars.—Here it will be convenient to consider the circumstances under which the requirements of the public health will be satisfied by the less severe measure of the exclusion from

school of particular children.

(a) It may be laid down as a universal principle that all children suffering from any dangerous infectious disorder (i.e. of a nature dangerous to some of the persons attacked by it, however mild in other cases) should be excluded from school until there is reason to believe that they have ceased to be in an infectious condition (see s. 126 P.H. Act, 1875, also s. 57 P.H.

Acts Amd. Act, 1907).

(b) Furthermore, as it is rarely possible to provide effectual separation of the sick from the healthy within the homes of children of the class attending public elementary schools, it must commonly be necessary that all children of an infected household should be excluded from school; first, because otherwise such children might attend school while suffering from the disease in a latent form, or at an unrecognised stage, and, secondly, because it is known that infection may attach itself to, and be conveyed by, the clothes of a person living in an infected atmosphere, even though the person himself remain unaffected. The same considerations will sometimes make it desirable to prohibit the attendance at school of children who are known to have been in contact with a source of infection;

* There are special considerations that have to be borne in mind in regard to diphtheria. Conditions of school aggregation alone seem occasionally to intensify the virulence of the type of this disease. (See chapter xiv.)

† Small-pox has only a slight incidence upon children of school age, under ordinary conditions of infant vaccination, but in districts where vaccination is now neglected it has occasionally been found necessary to

close schools on account of prevalence of small-pox.

of children of certain ages or classes; or of children from a particular street or hamlet.

In the case of infectious diseases involving little or no danger to life, such as mumps or skin diseases, school interests may be more particularly considered. In such case, however, it will usually be well for the M.O.H. to advise the managers to prohibit the attendance of every child while in an infectious state.

6.—B. Secondly, as to the closing of schools.—This, by more seriously interfering with the educational work of a district, is a much more grave step for a S.A. to take than to direct the exclusion of particular scholars. It is a measure that seldom ought to be enforced, except under circumstances involving imminent risk of an epidemic, nor even then as a matter of routine, nor unless there be a clear prospect of preventing the propagation of disease such as could not be looked for from less comprehensive action.

Even where school closure is deemed necessary it need not always extend to the whole school or department, but may on suitable occasions be limited to particular classes or standards. Thus in the case of outbreaks of measles or whooping cough it is often unnecessary to close the senior departments of a school; since the majority of the elder children are commonly protected against those diseases by having already passed through an attack in previous years.

7. By Article 18 (6) of the Board's Order of 23rd March, 1891, the M.O.H. on the occasion of an outbreak of dangerous infectious disease is to advise the persons competent to act as to the measures to be taken to prevent the extension of the disease. If, therefore, he finds that the children of infected households are attending school, he should send notice of the fact to the schoolmaster, and give such advice as appears to him to be necessary with regard to the exclusion of the children from school, and as to the time for which such exclusion should continue.*

Where the number of children to be excluded is small, and the schoolmaster acts on the advice of the M.O.H., it may not be necessary to take formal action under Article 57 of the code; but where the number of children whom it is desirable to exclude from school is such as is likely seriously to diminish the average attendance, or where the advice of the M.O.H. is not

^{*} It is well always to send a second notice when the infection is at an end and the quarantine is terminated. Copies of these notices may with advantage be sent to the school authorities, in order that the school attendance officer may not in the meantime press the parents to send the children to school.

followed, and there is danger of the disease spreading by means of the school, notice for the exclusion of the children in question should be made in accordance with the requirements of Article 57.

The attention of school attendance officers and of schoolmasters should also be drawn to the following considerations. Frequently they themselves will obtain the earliest information of the occurrence of infectious disease among scholars, and it is most desirable that such officer or master should without delay communicate the facts to the M.O.H. Absence of any child from school on the plea that it is suffering under one of the before-mentioned diseases, and absence of several children of one family from school at the same time, no matter what name be given to the complaint that keeps them at home, should be reported to the health officer. In practice it has been found that this notification of absentees has materially aided the local health officer in taking measures for the suppression of infectious disease, to the advantage alike of the district and of the school. Furthermore, schoolmasters may properly be asked to take note, especially when an epidemic threatens or is present, of symptoms occurring in any of their scholars that may indicate the commencement of disease, febrile in nature. heat of skin, such symptoms are shivering, headache, and languor, especially if commencing suddenly, vomiting, rashes on the skin, and sore throat. When scarlet fever or diphtheria threatens, every trace of sore throat should be looked upon as suspicious. In any case where such symptoms are observed, the safest course will be to exclude the child from school until assurance can be had that it may attend school without harm to itself or danger to other scholars.

8. As regards duration of exclusion from school of particular children, the time to be specified will vary in different diseases and different cases, and in this matter the sanitary authority will doubtless be guided by the advice of their M.O.H.

9. In deciding whether an outbreak of infectious disease among children of school age may be best combated by closing the school, or whether it will suffice to exclude the children of infected households, the two most important points to be considered are:—

(a) The completeness and promptness of the information received by the officers of the S.A. respecting the occurrence of infectious cases.

(b) The opportunities which exist for intercourse between the children of different households elsewhere than at school.

10. (a) The more prompt and full the knowledge of cases of infectious disease that the S.A. are able to obtain, the better will be the prospect of checking such disease by keeping away from school the children of infected households, and the less will be the necessity for closing schools.* If the cases be few in number, and their origin known, the exclusion from school of the children of infected households will probably suffice, but this measure will fail where there are many undiscovered or unrecognised cases, or where the known centres of infection are very numerous.

Commonly, the failure of carefully considered measures of exclusion to stay the spread of an epidemic which shows a special incidence upon school children, may be regarded as pointing to the continued attendance at school of children with the prevalent disease in a mild or unrecognised form, and a strong case will appear for the closing of 'schools. But it must be remembered that the closing of the school will deprive the Medical Officer of Health of the information respecting attacks in the early stage or of doubtful nature which he might

otherwise be able to obtain from the school teachers.

11. (b) The second material consideration, in deciding as to the desirability of closing schools during the prevalence of infectious disease, is the amount of opportunity for intercommunication between the members of different households elsewhere than at school. In sparsely populated rural districts. where the children of different households, or of separate hamlets, rarely meet except at, or on their way to, the village school, the closing of the school is likely to be effectual in checking the spread of disease. It is less likely to be useful in a town or compact village (particularly where houses are sub-let and yards are in common), where the children of different households when not at school spend their time in playing together, and often run in and out of each others' houses. But it must be remembered that children when at play out of doors are brought into much less close association with each other than when at school.

In rural districts, where epidemic diseases are less frequently prevalent, school closing may be required as an exceptional measure to meet an exceptional state of things. As regards more populous places, it must not be forgotten that if schools were to be closed whenever an infectious disease was prevalent, there are many places where schools would hardly

ever be open.

^{*} The provisions of the Infectious Diseases Notification Act, and of s. 58 of the Public Health Acts Amendment Act, 1907 (where that has been put in force), may be referred to in this connection.

12. On occasions when by reason of the absence or exclusion of a considerable number of children the attendance at a school is greatly reduced, the managers often prefer to close it altogether, as by so doing they are enabled to obtain the advantage of Article 45 (b) of the Education Code, before quoted. This, for reasons to be mentioned, is especially apt to occur in the case of epidemics of measles and whooping cough.

But it must be borne in mind that under Article 57 of the Education Code the Sanitary Authority is only empowered to order the closure of a school with a view to preventing the spread of disease, and not in the financial interests of the schools; and that when a large proportion of the susceptible children have already contracted the disease or been exposed to infection, school closing can do little to prevent its spread.

In such cases, however, if the school managers desire to close the school, it is open to the Medical Officer of Health, if he thinks fit, to give a certificate for the purpose of Article 45 (b) of the Code, though he is not under any obligation to do so.

13. The question of school closing in outbreaks of measles merits special consideration. School closing probably more frequently takes place on account of epidemics of measles than for any other disease, but as closing is commonly deferred until a large proportion of the children are already absent, it is, in populous districts at least, useless for the purpose of preventing the spread of the disease. The following facts respecting measles have to be taken into account:—

The disease is very infectious, and almost all young children who have not already had it are susceptible to it, the susceptibility being probably greatest in the second to the fifth years of life. Persons, however, comparatively rarely contract it a second time, so that, in populous districts where the epidemics commonly recur every two or three years, most of the older children are protected against it by

having passed through a previous attack.

The incubation period of the disease from infection to the commencement of illness is usually from 12 to 14 days, and the disease is very infectious during the first three days before the characteristic rash appears. Hence, if measles is introduced into a school, the first crop of secondary cases will occur about 12 days after the original case, and in 12 days more there will be a second crop comprising the majority of the unprotected children. The rapidity of spread depends in large measure upon the number of unprotected children in the class or school; and hence the likelihood of an introduction of measles into a school being followed by an epidemic is in proportion to the length of time which has elapsed since the last previous epidemic.

In populous towns measles is constantly more or less present, and is on frequent occasions introduced into schools, hence it is not possible to prevent epidemics, but at most only to postpone them. But since the mortality from measles chiefly occurs among children under five years of age, in so far as the recurrence of an epidemic can be postponed, the mortality from measles will be reduced, since a larger number of children will have passed the age at which the risk of death from an attack is greatest.

In view of the failure of school closure, when deferred until a considerable number of children have been attacked, to prevent the spread

of measles a class closure of short duration after the occurrence of a single case of measles has been suggested. If this be done the class should be closed on the ninth day after the sickening of the first child, for a period of five days only, after which time only those who have sickened should be excluded, with those in the same households who have not had measles.

The infection of measles is probably conveyed chiefly by the nasal mucus, but is less persistent than that of small-pox, scarlet fever, or diphtheria, and is not commonly conveyed by healthy persons; hence it is unnecessary to exclude from school the children of infected households if they have themselves had measles.

The foregoing remarks apply especially to measles as it occurs in populous town districts; in rural districts, for the reasons given in paragraph 11, there is a better prospect of checking the spread of the disease by prompt measures of exclusion and school closure, especially of the infant classes.

Similar considerations apply in some degree to whooping cough, which, like measles, attacks especially young children, and is infectious in the early stage, before the characteristic whoop has appeared.

Upon any threatened outbreak of measles or whooping cough it is advisable that all children below the age of obligatory school attendance, i.e. under five years old, who may be attending school should be excluded,

unless they are known to have previously had the disease.

14. Scarlet fever in schools is spread chiefly by the agency, first, of mild unrecognised cases, presenting perhaps only a slight sore throat, the nature of which is only recognised when peeling of the skin subsequently occurs; and second, of convalescents still in an infectious condition, especially with nasal discharges. It is desirable that children who have had scarlet fever should not be re-admitted to school for a week or two after recovery or discharge from hospital, and in no case until free from

infectious sequelæ.

 Diphtheria in schools, though spreading less rapidly than measles, is often very persistent and sometimes causes serious mortality. An outbreak is, often preceded by cases of sore throat not obviously of diphtheritic nature, or not so recognised, but in which the diphtheria bacillus can often be identified in secretions taken from the throat by swabs. The specific bacillus may often be found in the throats of persons who have been in contact with cases of diphtheria, but are themselves apparently healthy. When diphtheria breaks out in a school it is desirable that the other children in the class affected should be examined with a view to the detection of such cases of sore throat; and in many cases bacteriological examinations may be necessary. It is desirable also that bacteriological examinations should be made of the throats of convalescents from diphtheria, and that they should not be allowed to return to school until the absence of the diphtheria bacillus is satisfactorily shown.

16. School closure is occasionally necessary on account of infectious sickness in the master's family, which would involve risk to the scholars. It is also sometimes necessary to close a school for a day or two in order that it may be disinfected and cleansed after cases of infectious illness have been in attendance, or to allow of the rectification of sanitary defects

of a nature to produce or spread disease.

It must be remembered that proper ventilation of school-rooms, avoidance of overcrowding, and cleanliness and, where needed, disinfection of schoolrooms and school appliances, do much to prevent the spread of infectious diseases; and if these matters were duly attended to and careful supervision kept over children slightly ailing, school closing would be comparatively rarely required,

17. In places where there are several public elementary schools, if an outbreak of infectious disease be confined to the scholars of one particular school, it may be sufficient to close that school only. But where different schools have all appeared to aid in the spread of disease (though perhaps to an unequal extent) the Sanitary Authority may consider it advisable that all should be closed lest children in an infectious state who previously attended the schools that are closed, should be sent to others that might remain open.

The playgrounds should not remain open when the schools are closed, as they may form a meeting place for the children whom it is the object of

the closure to keep apart.

It must be remembered that Sanitary Authorities have no power in respect of Sunday schools, or other private schools, except in so far as these may contravene s. 91 (5), s. 126, or other provision of the Public Health Act, 1875, or s. 57 Public Health Acts Amendment Act, 1907, where it has been put in force; but it will often be expedient to invite the co-operation of managers of such schools in efforts for securing the public health. Experience shows that they are usually ready to defer to the representations of the authority responsible for the public health of the district.

18. The M.O.H. has not power to order the closing of a school; his function in this respect is advisory only. Reports of the M.O.H. to sanitary authorities, advising the closure of a school or schools in any district, are to be treated as "special" reports within the meaning of the General Order of the L.G.B. of March 23rd, 1891, and copies of them are required by Art. 18 (15) and (16) of that Order to be sent to the Board, and to the County Council. These reports should state the grounds upon which the M.O.H. advocates the closure of the school or schools in preference to the exclusion of particular scholars.

19. All notices of the sanitary authority for the closing of Public Elementary Schools should be addressed in writing to the Managers, and should state the grounds on which the

closing is deemed necessary.

All such notices shall specify a definite time during which the school is to remain closed; this should be as short a period as can be regarded as sufficing on sanitary grounds, since a second notice may be given before the expiration of the first, if it should be found necessary to postpone the re-opening of a school. The Managers of schools, after complying with the requirements of the S.A., have the right of appeal to the Board of Education, if they consider any notice to be unreasonable.

20. Where the Medical Officer of Health is not himself the school medical officer appointed under the Education (Administrative Provisions) Act, 1907, it will be very desirable that in matters relating to the health of school children, he should co-operate with the officer appointed under that Act and should consult with him in any case of doubt or difficulty,

Administration as to infectious diseases in elementary schools. - The following is the practice now adopted in London by the Education Committee of the County Council:

(a) For notifiable diseases. On receipt of the notification from the M.O.H. the following action is taken at the schools:-(1) The head master sends the notice to the head teachers of the other departments, and all affected children (whether suffering or coming from the home in which the disease exists) are excluded from school. (2) Head teachers of all departments forward on the same day the names and addresses of all children affected by the notice, to the divisional superintendent and the medical officer (education), on a form specially provided for the purpose. The notification is then filed at the school until the end of the school year. (3) Children coming from houses in which erysipelas or enteric fever exists, but who are not themselves suffering from the disease, should not be excluded from school. (4) Children excluded on account of a notifiable infectious disease occurring either in themselves or others in the houses in which they live, must not be allowed to return to school unless a certificate has been received from the M.O.H., stating that the premises are free from infection. The certificate of the M.O.H. does not certify that the children are in a condition to resume attendance at school, for it may be that, though the premises are free from infection, the children coming from such premises may be sickening for an infectious disease. It is necessary, therefore, for a further period of seven days to elapse before the return of such children to school, unless the M.O.H. shall specially certify that a longer period of absence is necessary. (5) No child who has been in an infectious disease hospital should return to school for at least a fortnight after discharge from the hospital. (6) The London County Council have passed the following resolution in respect to diphtheria:-

"That the Education Committee be authorised to refuse, during the presence of diphtheria in any district, readmission to school of children excluded on account of diphtheria or sore throat until such children shall have obtained a medical certificate of freedom from infection, based on a bacteriological examination."

- (b) For non-notifiable diseases. The school management code in London is now as follows :-
- (1) Children suffering from measles must be excluded from school for at least one month; mumps, for one month; chickenpox, for at least two weeks, or until every scab has fallen off

the scalp or body; whooping cough, for as long as the cough continues, and not less than five weeks from the commencement of the whooping.

- (2) Children coming from houses in which measles, mumps, chicken-pox, or whooping-cough exists must be dealt with as follows: (a) Children in schools other than infants' schools who have not had the disease, and all children in infants' schools, must be excluded. 1(b) Children in schools other than infants' schools who have had the disease need not be excluded.
- (3) Children living in infected houses and excluded from school under section (2) (a) must absent themselves for the undermentioned periods: Measles: (1) Children attending other than infants' schools, until the Monday following the expiration of 14 days from the occurrence of the first case; (2) Children attending infants' schools, until the Monday following the expiration of 14 days from the occurrence of the last case. Mumps: For such time as the medical attendant dealing with the case deems necessary, or, in cases where there is no medical attendant, for three weeks. Chicken-pox: For two weeks. Whooping-cough: For two weeks.

It is understood in all cases that thorough disinfection is carried out. As a further safeguard it is desirable to insist upon a medical certificate of freedom from infection when a child returns to school after an infectious illness,

Typhus and enteric fever are not included in the list. They, of course, rarely require consideration. In either, three weeks may be adopted as the period of quarantine; and four weeks as the earliest date of return to school after an attack, if all symptoms have disappeared and convalescence is complete.

The duration of ringworm, ophthalmia, impetigo, and scabies is indefinite; and children suffering from any of these should be rigidly excluded until all trace

of the disease has disappeared.

It is important that teachers should be acquainted with the early signs of infectious diseases, and should watch for them, especially if such diseases are prevalent in the neighbourhood, or if any of the children

attending the school have been attacked. Slight attacks of disease may escape recognition until late (e.g. until the desquamation stage of scarlet fever), and multiple instances of sore throat should always be regarded with suspicion. It is important, too, that the school authorities should co-operate with the S.A. in such matters, and should report cases of non-notifiable disease. Under the Education Act, 1902, County Councils and S.A. are the bodies appointing the education committees, so that unified action is practicable.

Elementary Education (Blind and Deaf Children) Act, 1893, places upon the parent and school authority the responsibility of education of a child who is blind or deaf. The authority must obtain suitable education for such child in a school certified by the Board of Education, either established by the authority or receiving contributions from such authority with or without representation. Two or more authorities may combine for this purpose. The period of compulsory education must extend to 16 years.

Elementary Education (Defective and Epileptic Children) Act, 1899, gives power to a school authority to determine what children in its district are defective and epileptic. "Defective" children are those who "by reason of mental or physical defect are incapable of receiving proper benefit from instruction in the ordinary public elementary schools," and the defect must be certified by a medical practitioner approved by the Board of Education. Authorities may make provision for education of such children by providing or contributing to special classes or schools for the purpose, with or without "boarding out" accommodation, and periodically shall examine medically such children to ascertain whether they are fit for ordinary school. The parent is obliged to send his defective child

to such special school, which shall be entitled to grants-in-aid and the receipt of payments from parents or poor law. Compulsory education extends under

this Act also to 16 years.

The London County Council (General Powers) Act, 1907 (ss. 36 and 38), provides for examination and cleansing of verminous school children, and the Public Health Acts Amendment Act, 1907, requires for the sanitary authority lists of school children suffering from infectious disease and their exclusion from school. The Education (Provision of Meals) Act, 1906, provides for feeding school children under certain conditions with or without payment by parents.

Medical inspection of school children. Under the Education (Administrative Provisions) Act, 1907 (s. 13), local education authorities are required to provide (a) for vacation schools and playcentres, and (b) for medical inspection of children. The latter duty is "to provide for the medical inspection of children immediately before, or at the time of, or as soon as possible after their admission to a public elementary school, and on such other occasions as the Board of Education direct, and the power to make such arrangements as may be sanctioned by the Board of Education for attending to the health and physical condition of the children." This is a new power commencing from January 1, 1908. The Board have issued an official memorandum appointing at least three medical inspections of each child during school life. The work is placed in the hands of the local education authorities (which are public health authorities), and the minimum medical inspection is to include general conditions (height, weight, cleanliness, clothing, nutrition), medical history, mental capacity, an examination of nose, throat, eyes, ears, teeth, and the detection of physical defects, deformities, diseases and paralyses.

More elaborate inspection and anthropometric surveys may be undertaken where practicable. Each education authority is required to appoint school medical officers. No medical treatment as ordinarily understood is to be carried out by the education authorities except by sanction of the Board. Much, however, may be done in the way of adapting the means of education to the physical capacities and mental requirements of the children. This enactment will bring England and Wales into line with Germany and other countries which have had medical inspection for some years.

Inspection of schools.—The following schedule includes the principal points that require attention in

the sanitary inspection of day schools :-

Name of school.—Character (elementary, secondary, technical)
Local education authority.

1. Position and surroundings of school .- Nature of soil and

subsoil; site, aspect, etc.

2. Playground. - Size; covering of paving; drainage;

buildings or fixtures in; cleanliness.

3. School buildings [plans].—Structure, entrances, corridors, staircases, central hall, school-rooms, class-rooms.

Number of storeys. Infant schools. Special departments for cookery, laundry work, handicraft, science, drawing.

(a) Structure.—Condition of roofs, walls, floors, ceilings.

Purpose of rooms.

(b) Cubic capacities and accommodation; superficial area and cubic space per child. Overcrowding in class-rooms.

(c) Heating.—Mode, radiation, convection; fire-places, stoves, hot pipes, or other methods; temperature in summer and winter.

(d) Lighting. - Mode; degree; windows, size and position;

means of artificial lighting.

(e) Ventilation.—Methods; natural or artificial; measurements of Co_2 . Standard may be $\frac{10}{10,000}$. Air supply.

(f) School furniture and fittings. Desks and seats. Gymnasium. Teaching appliances — blackboards, slates, pencils, sponges.

(g) Cleanliness. — Arrangements for; scrapers. Dry

cleaning. Periodical disinfection.

4. General sanitation. — Refuse removal; receptacles; ashpits, structure and condition of; sanitary conveniences (water or earth); separation of sexes; type and character; number; condition; arrangement; position; urinals, accommodation, flushing; drainage; surface and soil drains; rain water; sink wastes; cesspits (cover, structure, cleansing); relation to well and land drainage; disconnection and ventilation; trapping, testing, etc.

5. Lavatory, bath-, and cloak-room accommodation.—Position; structure; number of basins; water supply; towels; clothes hooks; drying arrangements; size, heating, and

ventilation of cloak-room. Baths,

6. Water supply.—What provision; if well, position and covering; surroundings of well; quantity and adequacy of supply. Playground tap. Water-filters. Storage and distribution.

7. Master's house.—Position, size, accommodation; general sanitation. Teachers' rooms and accommodation.

8. General.—Number of teachers to scholars. Number on roll and average attendance. System of medical inspection in vogue. General result of records as to personal hygiene of children. Infectious diseases. Hours of school and playtime. Physical exercises. Attitude of children in school.

Much has been done in school hygiene in recent years under the above Acts, the Education Code, and by the enterprise of certain local education authorities (Bradford, Manchester, London, the County of Surrey, etc.). This work has included not only sanitation of schools and the prevention of the spread of infectious diseases, but medical inspection particularly of defective and under-nourished children. In London a cleansing scheme has been in practice, vision has been tested, a favus-school has been established, school nurses have been employed, and physically defective children have been supervised. In Croydon ringworm has been treated by X-rays and other methods. In Liverpool school nurses have been employed, and in many centres "special" schools have now been established.

CHAPTER VIII

HOSPITALS

General considerations.—All that has been said in respect of the site, surroundings, and construction of houses applies with still greater force to hospitals. Pure air and light are of especial importance in sick wards, that are occupied day and night. The site should be as dry and open as possible, not exposed to smoke or effluvium borne by wind, and not unduly sheltered by trees, buildings, or hills.

Wards.—Long wards should face east and west so as to admit sun on each side alternately, and if any deviation is necessary it should preferably be such as to give a south-east and north-west aspect (Thorne-Thorne). Single-storey hospitals are preferable, as affording the freest circulation of air. Each ward is then completely detached from the rest, and the absence of stairs is advantageous; but from motives of economy and convenience of concentration two or more storeys are usually built.

Small wards are inevitable in cottage hospitals, and, indeed, in all hospitals, since provision has to be made for the isolation of severe or infectious cases, and for the classification of patients according to sex, and sometimes according to disease. As a general proposition, however, large wards are more readily ventilated, warmed, and lighted, more economical in construction and management, and more convenient

as regards nursing.

For long wards the width should be 24 to 30 feet, the height 13 to 14 feet, and the length such as to allow a cubic space per patient of at least 1,200 feet. Not less than 2,000 feet is considered necessary in a fever or lying-in hospital, or

for severe surgical cases. It is not permissible to obtain this space by increasing the height. As already explained, extreme height is of little utility as regards ventilation, and in hospitals it is essential to have plenty of floor space around each bed. For nursing purposes at least 90 square feet per bed is necessary, and more will be required if the ward is largely used for clinical teaching. Wherever there is any possibility of septic or other infection, as for example in surgical wards, lying-in wards, and wards for infectious diseases, wide separation of beds is important. In such cases 12 feet of wall and 130 to 140 feet of floor space should be allowed. It is found that for convenience in nursing the number of beds in one ward should not exceed 32, and usually 24 or 28 would be preferable.

The beds are arranged with their heads to the wall, facing into the ward. Each bed should be placed between two adjoining windows, the space between the windows being not less than a foot wider than the bed. At most two beds may be allowed in one space, the interval between them being at least three feet. The intervals between the windows of a ward should be such as to allow of this arrangement. The windows

should reach to within a foot of the ceiling.

All the surfaces should be impervious, smooth, and washable, and all ledges, crevices, and stagnant corners in which dust may lodge must be avoided. The floors may be paraffined or waxed, the walls lined with tiles or glazed brick or Parian cement, or in the absence of

such materials simply painted.

Heating and ventilation.—The heating may be effected by open fire-places or ventilating stoves, or both, and it is well to have in addition coils of steam or hot-water pipes with radiators in corridors. Ventilation must be provided for to the extent of at least 3,000 feet per head per hour, and it may be taken as a safe rule that the whole air of the ward ought to be changed thrice hourly, so that the hourly supply of fresh air should be three times the cubic capacity of the ward. Extraction is provided for by fire-places, stoves, windows opening at the top, Sheringham valves near the ceiling, chimney shafts, warm outlet flues, and in other ways. Special extraction

shafts, with inlets at or near the ceiling, are desirable. Such shafts may be carried up alongside or surrounding the chimney or stove flue, so as to utilise the waste heat as a motive power; or, independently of these, with a gas jet in the inlet to create an upward current. Some hospitals for infectious diseases have an extraction shaft upon a larger scale, the foul germ-laden air being made to pass through a gas furnace. Most, if not all, of the germs are no doubt destroyed in this way, and the rest are prevented from diffusing horizontally. This precaution was suggested by the Hospitals Commission, and is especially important in regard to small-pox hospitals, which have been shown to be capable, under certain conditions, of acting as foci from which infection spreads through the air.

It is desirable to make arrangements for warming the air passing into the wards. This may be done by means of ventilating fire-grates, ventilating stoves, or coils of hot pipes at the inlets. Windows, Sheringhams, and Tobins may suffice in warm weather. It is a common practice to place the inlets under beds, with the intention of avoiding draughts, and supplying the

purest air close to the patients.

In the most approved plans a cross-ventilated corridor or ante-room leads from one corner of the ward to the water-closets and slop sinks, and the bathroom and lavatory are similarly placed at the adjoining corner. At the opposite end is a nurse's room

with a window overlooking the ward.

Hospitals for infectious diseases.—
Reference has already been made to the need for greater cubic space and ventilation in such hospitals, and to the importance of guarding against the tendency to aërial spread of infection in certain diseases. The communication with the outside has also to be kept under strict control. It is desirable that

"Isolation Hospitals" should themselves be isolated from other buildings, and even from thoroughfares, by reserving as wide an open space as possible. The boundary walls should be 61 feet high, Each disease must be isolated separately, and if possible in separate blocks, not less than 40 feet apart, and the same distance from the boundary.* There should be no communication whatever between the different blocks, and the intercourse with the common administrative department should be as little as possible. As regards structure and arrangement of wards the standards named in respect of general hospitals apply. A special block containing at least two or three small wards is necessary for doubtful cases, or cases not received into the principal wards. The site should be high and airy, on dry gravelly soil, removed from buildings but not too exposed. It should not in any case be less than 2 acres in extent, and the number of patients per acre should not much exceed 20.

The essential parts of a hospital for infectious diseases are-

1. An administrative block, including rooms for the medical officers, matron, nurses, and servants; dispensary, kitchen, scullery, larder, and pantry; laundry, linen-room, store-room, coal-cellar, bathrooms, closets, etc. etc.

2. Hospital blocks, namely, at least one for each disease to be isolated, and in each block at least two principal wards, one for each sex. Additional small wards for one or two beds are desirable. An "iso-

^{*} The L.G.B. require (1902) that a small-pox hospital should not be so placed as to have within a quarter of a mile a hospital (whether for infectious diseases or not), a workhouse, or any similar establishment, or a population of 200 persons; or within half a mile a population of 600 persons, whether in one or more institutions or in dwelling-houses; even where the above conditions are fulfilled a hospital must not be used at one and the same time for the reception of cases of small-pox and of any other kind of disease.

lation block," consisting entirely of small wards, is most useful. Besides the wards, closets, sinks, and bathrooms, each block should contain a nurse's room, overlooking the ward, and so arranged as to serve as a scullery and ward-kitchen for minor cookery; linen-room, store-room, and coal-place. The nurse's sleeping-room must be away from the wards; it may be in the same block, but is more usually placed in the administrative block. Arrangements must be made for the patients from each block to take exercise in the grounds without coming into contact with each other. Small-pox hospitals should be entirely independent of those for other diseases.

3. Mortuary and post-mortem room; disinfecting room; stable and ambulance shed; porter's lodge, etc. etc. The drainage, water supply, and arrangements

for heating need no special mention.

Cost.—The expenses to be incurred in respect of isolation hospitals are classified under the Isolation Hospitals Act as structural, establishment, and patients' expenses. We may first consider the capital or structural outlay. The great cost of permanent hospitals—rarely less than £200 and sometimes £400 per bed-has led to the substitution in many cases of temporary structures, more especially in the presence of an epidemic, when rapidity of construction is the first consideration (£20-£50 per bed). These may be tents or buildings of wood and galvanised or corrugated iron. Wooden hospitals are common, and can be run up in two or three weeks. The walls must be made double, and it has been found advisable to fill up the hollow with sawdust. The floors must be raised above the surface of the ground. The roof may be slated or tar-felted. The foundation required is generally a four-course brickwork with air-bricks and dampproof course. The dimensions and arrangements can be made the same as in permanent buildings.

The same may be said of galvanised iron huts, which are often used for hospital purposes. Tents can be erected still more rapidly, upon a wooden floor raised above the level of the ground. canvas should be double. Recently other forms of temporary hospitals have come into use, including those of Dæcker and Ducker types. Both of them are convenient in use, and can be erected at a few hours' notice. They consist of a stout waterproof material stretched upon wooden frames, so shaped and numbered that they can be speedily put together to form a complete and weather-proof hut, ready for the reception of patients. Willesden waterproof paper is also used for the construction of huts. Sometimes a composite construction has been preferred, with an administrative block of stone or brick, and pavilion wards of corrugated iron and wood costing about £120 a bed. It should be added that the L.G.B. does not as a rule sanction loans for the construction of temporary hospitals.

Their comparative cheapness and the rapidity with which they can be erected are important points in favour of temporary hospitals, but nevertheless their proper function is to supplement, and not to supersede, permanent buildings of brick or stone. They are less attractive in appearance than the latter, less comfortable, and more difficult to ventilate efficiently, and to keep warm or cool. True disinfection, such as to fit them for the reception of cases of different diseases, if not impossible, is at least less easy and less certain than in buildings with hard non-absorbent surfaces, free from pores and crevices. Their durability is far less than that of a permanent building, and an annually increasing outlay is needed in order to keep them in habitable repair. It is often claimed on behalf of temporary hospitals that no disinfection is necessary, since they can be burnt at the end of the epidemic, and renewed; but this plan, which would very soon prove more costly than the erection of permanent hospitals, has rarely, if ever, been put into practice. The economy in first cost is great, but is usually exaggerated. It must be remembered that the cost of erecting a permanent hospital, as

ordinarily stated, includes the preparation and laying out of the site, fencing, drainage, water and gas supply, erection of accessory buildings, such as mortuary, disinfector, lodge, etc., and the whole cost of the administrative block. These are not usually included in the stated cost of "temporary" hospitals, although just as essential in the one case as the other. Furthermore, the cost per bed largely depends on the number of beds that are allowed in a given cubic space, and it will be found that, if the standard adopted in permanent hospitals is applied to all, the nominal accommodation in temporary hospitals will often be reduced by one half, or more. If the comparison is made upon equal terms, the saving in first cost is insufficient to compensate for the want of durability, and for every purpose of use permanent hospitals have great advantages. The liability to fire must not be forgotten. The only real advantage of temporary hospitals is, therefore, the rapidity of erection, and (except as a supplementary provision) this can only be important when the S.A. has failed in its obvious duty of providing a hospital without waiting for an epidemic. It is then frequently too late, however speedily the hospital is got ready. Means of isolation are needed at all times, in every large community, and it is even more important to endeavour to delay and prevent outbreaks by secluding a few "sporadic" cases than to grapple with epidemics already in existence.

In addition to the structural expenses there are the establishment costs, which include all expenses of maintenance, repairs, salaries, etc., working out often at about £20 to £30 per bed per annum. Then there remain the expenses of removing, feeding and providing for the patients, including medicine, appliances, disinfection, etc. Many hospitals would average say £10 or £12 per patient per annum.

Number of beds.—The number of beds which should be provided has been stated as one for each thousand of population, but such an estimate can only be of the roughest kind. Much depends upon the character of the population, upon the number of diseases which it is proposed to isolate, upon the thoroughness or the reverse with which the policy of isolation is to be carried out,

and lastly upon the question whether the hospital is intended to be a final and inelastic establishment, or merely a permanent provision sufficient for nonepidemic times but capable of rapid extension at a few. days' notice in the event of serious epidemic. It is obvious also that averages are of little use as indicating the maximum possible demand for hospital accommodation at any given date, and it is not to be assumed that only one disease will be epidemic at one

By general consent small-pox, scarlet-fever, diphtheria, enteric fever, plague, and typhus are diseases which, for the public safety, ought to be treated in hospital whenever possible. Experience has shown that with compulsory notification and free hospitals it is possible to get into hospital practically all the small-pox, a great majority of the scarlet-fever, diphtheria, and enteric cases, and all the plague and typhus cases, without recourse to compulsion. For reasons stated elsewhere it appears to be very doubtful whether the time has yet come for the isolation of measles or whooping-cough upon a sufficient scale to be of any practical use; if, however, cases of special urgency occurred, the small wards would afford means of isolating them.

To sum up, therefore, the provision of one bed per thousand population is not likely to be found excessive. Indeed, it will not suffice in the event of epidemics, and if such diseases as measles and whooping cough are to be isolated, large additions will be necessary. It is, therefore, advisable to reserve a liberal area for future extensions, temporary or otherwise, and to construct the administrative portions upon such a scale as to suffice in case of such extension of the isolation blocks. The wards should be so constructed as to admit of disinfection, so that the greatest number of beds may always be




FIG. 8.—ISOLATION HOSPITALS.—PLAN OF A WARD PAVILION FOR TWELVE BEDS. (L.G.B. MODEL.)




FIG. 9.—ISOLATION HOSPITALS.—PLAN OF AN "ISOLATION" BLOCK FOR EIGHT BEDS. (L.G.B. MODEL.)

available for whatever disease is prevalent at the time. This can be done with safety and without difficulty if all the internal surfaces are impermeable. The wards, whatever the number of beds, must be sufficiently numerous to allow of the separation of the sexes wherever necessary, as well as the isolation of the several diseases one from another.

It is of the utmost importance, if isolation is attempted upon the large scale, to make the hospitals free, and this is only reasonable, since the public gain at least as much as the patient by his seclusion. The Sanitary Authority must not regard removal to hospital merely as a privilege sought by the patient and his friends, but as a concession on their part to the public safety. Any charges, however small and readily remitted, are deterrent in the very cases in which isolation is most needed. The revenue from patients' fees is at best a small fraction of the cost of maintenance, and the wholesale remission of them --without which the isolation of cases among the working classes is impracticable—has an appearance of charity which is naturally resented. An exception may be made in respect of the wealthy, who are willing to pay suitable fees for the use of private wards and special nurses.

tion for isolation is based on the broad ground that certain diseases are infectious; that is, are readily transmissible from the sick to the healthy. Infected persons act therefore as foci of the disease, and where such persons are left in the midst of densely populated communities there is a greater tendency for the disease to spread than if such persons are isolated away from others. Although it may be possible to isolate almost any case of infectious disease in a large house, where all the requisite precautions can be rigidly maintained during the whole period of in-

fectiousness, it may be stated broadly that "home isolation" in smaller houses cannot be carried out in its entirety. In houses of the working class, for example, not one case of scarlet-fever in a hundred is, or can possibly be, properly isolated, during the whole period of still infectious convalescence, from the rest of the household, or even from the public. Medical supervision is usually withdrawn at the end of the acute stage of illness. Where satisfactory isolation can be provided at home, hospital isolation may be unnecessary, and if a hospital is badly administered, or there is overcrowding, such home isolation may even be preferable. There can be no doubt that the separation of the sick from the healthy has been invaluable in preventing the spread of disease, although instances are not rare in which hospital isolation has failed to give satisfactory results. This is not surprising when it is remembered that the isolation has often been partial, and the aggregation of infectious persons, more especially in times of epidemic pressure, carried too far. Many of the factors which determine epidemic and endemic prevalence of disease are still obscure, and it is only within the last 30 years that we have begun to appreciate the germ theory and what it means in actual practice. The hospital cases include many of the more virulent kind, and many from homes where means of isolation are most inadequate, and the removal of these cannot have failed to limit infection.

It has been contended, on the basis of certain local statistics of notification, isolation, and deaths, that isolation has failed materially to reduce the prevalence and fatality of scarlet-fever, and that the fatality has fallen more in towns which have not practised hospital isolation than in those which have; and this has been attributed to a "hospitalised strain" of scarlet-fever in the latter (Millard).

Hospital isolation has also been criticised on the ground of "return cases" of scarlet-fever; that is, secondary cases of the disease occurring in the same household after the return home of the primary patient from hospital. These have been found to be about 1.3 per cent. of the number discharged. Simpson has shown that 80 per cent. of such return cases are due to discharges from mucous membranes, such discharges being in all probability the carriers and not the causal agents of infection. Local and antiseptic treatment combined with transference to less infective wards before leaving are likely to be more effective in preventing return cases than long detention (Simpson).

In dealing statistically with this question it has to be remembered that we know nothing of what the figures would have been in the absence of hospital isolation, under the same conditions of increasing density of population. It is certainly true that scarlet-fever has not been eradicated, and the milder type of the disease now prevalent (as in Sydenham's time) adds to the risk of undetected cases, in which no question of isolation of any kind arises. The causal agent and the conditions which increase or lessen its virulence are also at present unknown.

Broadly, it may be said that any limitation of the spread of infection by personal contact must reduce the prevalence of the disease, and that therefore, if rightly managed, under proper conditions, hospital isolation is of direct and marked value in the reduction of scarlet-fever. The only alternative, the so-called home isolation, affords no prospect of safety. Further, it must not be forgotten that the isolation of cases of this disease in hospital prevents the infection of milk, houses, clothes, materials, etc., with the virus of the disease.

CHAPTER IX.

REMOVAL OF REFUSE.

THE waste matter from households comprises (1) liquid refuse, including water used for washing, and slopwater; (2) ashes, and vegetable and animal refuse;

and (3) excreta.

Drainage is necessary for dealing with the first, and some system of dry removal for the second. The excreta may be disposed of in conjunction with the former ("water-carriage"), or with the latter (privy-middens, ash-closets), or independently (earth-closets, privies, or pail-closets without admixture of ashes).

Drainage.—A drain means any pipe used for the drainage of one building or premises. A sewer is a drain receiving the drainage of two or more premises. The former is vested in the owner, the

latter in the S.A.

Drains are intended to remove from the house the liquid and much of the solid filth that is produced within it, and they should be so constructed that in no circumstances can their contents, liquid, solid, or gaseous, escape from them at any but the points specially provided for this purpose. This result is attained by the choice of proper materials for their construction and by sound workmanship, by trapping all inlets so as to prevent the egress of draingases, and, lastly, by efficient ventilation at suitable points.

A trap in its simplest form is a mere bend in a pipe which retains water and thus prevents air from passing through it. For this purpose it is only necessary that the bend shall be sufficient to place some

portion of the upper surface of the pipe below the water level, the seal of the trap being the difference between the surface of the water and the depression at this point. Thus, in the diagram (Fig. 10), the seal is the water between a and b.

A water-seal alone will not always prevent the passage of gases and particulate matter through it.




Fig. 10.—Siphon Trap (Strap) with access screw-plug for cleansing.

a, b, the seal.

Gases in the drain can be absorbed by the trapping water and given off from its upper surface. Ammonia has been found to become perceptible through an unbroken trap in fifteen minutes. Moreover, when the water remains stagnant, bubbles of putrefactive gases may be formed, and, by their bursting, particulate matter may be discharged. Occasional changing of the trapping water is necessary, and the drain must be ventilated

to prevent its air contents from becoming too foul, or subjected to so much pressure that the water-seal is forced. A trap to fulfil all these requirements must be self-cleansing and must maintain an effectual seal between the drain and the outer air. For the former purpose it must be free from all angles and corners that can retain filth, and for the latter it must have a water-seal not less than 1½ inch deep.

The seal is liable to be broken by a number of circumstances. First, pressure within the drain may force its air contents through the trap. This can be prevented by ventilation of the drain. Secondly, if the momentum of the water discharge be great, the water in the trap may be driven out. This can be obviated by shaping the trap so that the water-holding portion is contracted, and the descending arm is larger

and square at the outlet. Thirdly, if the descending arm is long and runs full it may empty the trap by siphon action. This can be avoided by ventilating the top of the trap beyond the seal. Fourthly, the same result will sometimes follow when two or more pipes are connected together, the discharge from one sucking out the water from the trap of the other. This can be prevented in like manner. Lastly, the seal may disappear by evaporation, if the trap is little used.

Drainage of the subsoil of a house is not necessary in all cases, but when the site is damp some provision of the kind should be made. The drain laid for this purpose should consist of suitable earthenware pipes with the joints left open, and laid to a proper outfall, preferably in the open air, but if into a drain connected with the sewer, the connection should never be direct. The subsoil drain should always discharge into a trap, and should be ventilated at or near the trap by a shaft or upright pipe carried above the surface.

The drain must be laid with a sufficient fall for the purpose of enabling it to clear itself of its solid and fluid contents, and it is generally accepted that this should be one in forty, giving a velocity of flow in a 4-inch drain of about 3 feet per second.* Further, in discharging into a sewer, the drain should be connected with the lower half diameter of the sewer.

The drain should be constructed of glazed stoneware pipes with cemented watertight joints, or preferably, if the drain be beneath a building, of iron pipes (of which the interior has been coated with Angus Smith's solution) jointed with lead. In all circumstances the drain should be laid on a bed of concrete, and if it be made of any other material than iron the whole drain should be embedded in concrete. It is customary to use drains having a diameter of 6 inches, but a four-inch diameter is

^{*} Maguire's "decimal" rule is convenient but not invariably practicable; 1 in 40 is a proper fall for a 4-inch pipe, 1 in 60 for a 6-inch pipe, 1 in 90 for a 9-inch pipe.

frequently sufficient for houses of moderate size. It should never be less than this.

As a general rule, the drain should not be laid beneath a building, but in the case of terrace houses it is often impossible to adopt any other plan. In this situation the drain must be laid in a direct line beneath the building and be embedded and covered in concrete not less than 6 inches thick; the top of the drain at its highest point must not be less than a distance equal to the full diameter of the drain below the surface of the ground under the building. Means

of access should be provided outside the house.

An important point in connection with drain construction is the prevention of air from the sewer passing into the drain, or from the drain into the house. The former of these risks is guarded against by the insertion of a trap in the drain (interceptor) as near as possible to the point of its connection with the sewer. Such an interceptor is usually provided with a "raking arm" into the drain. The drain should be ventilated immediately on the house side of the trap, by a shaft carried up above the ground level; this will serve a double purpose, by admitting fresh air into the drain, and also affording a free outlet in case the trap is forced by pressure of sewer air. The opinion appears to be gaining ground that if drains are properly laid in concrete, and the ventilating pipe is carried up above the roof, interceptors are not necessary; that if the ventilating pipe of every house drain (whether soil pipe or not) acted as a ventilator to the sewer, the amount of foul air escaping would be so thoroughly and widely distributed as to create no nuisance; and that the interceptor may cause stoppage of the drain, especially if the gradient be slight. It should be added that, as a rule, bye-laws require an interceptor, with access thereto. A more complete and necessarily a more costly

arrangement is that of an inspection chamber at this point (Fig. 11); the drain is continued through this chamber in earthenware channels, and the arm in the trap, which is intended to allow the drain to be cleared between the trap and the sewer, is extended into the chamber. Branch drains are connected with the main channel by branch channels in the floor of the inspection chamber.

In connecting drains with one another, no right-




Fig. 11.—Disconnection (Inspection) Chamber.
A, "raking-arm," closed by a cap.

angle joints must be formed, either vertical or horizontal. They must be made to join obliquely in the direction of the flow.

The second risk, that of air from the drain entering the house, is guarded against (1) by ventilation of the drain; (2) by trapping all inlets to the drain, except those required for its ventilation; (3) by trapping and disconnecting all indoor drain inlets; (4) by proper jointings and efficient plumbing.

The proper ventilation of the drain is secured by continuing the soil pipe in its full diameter above the level of the roof. By this means air entering by the opening near the trap in the main drain can pass along

the whole drain, and issue from the soil-pipe. The openings referred to must be covered with suitable gratings to keep out leaves and dirt, and to prevent birds from building nests in them, the open spaces in each grating being at least equal to the section of the pipe thus covered. The sectional area of the ventilating pipe should not be less than that of the drain




Fig. 12.-Yard Gully.

which it ventilates.

Soil pipes should be situated on the outer walls of the house, and their junction with the drain, often a weak point in construction, will then be outside the house. If through any accident the

junction of the soil pipe with the drain should be opened, the escape of gas will be into the open air.

Whether the soil pipe is used for the purposes of ventilation or not, bends in its course should be avoided as far as possible. As a general rule, the diameter should be 4 inches. Iron and lead are both used; lead is more expensive but more durable, and free from joints that may become loose. Rolled (i.e. seamless) lead is necessary if the soil pipe is inside the house.

The old practice of trapping the foot of the soil pipe is inadmissible, since it prevents ventilation and

impedes the escape of sewage.

Waste-pipes.—The overflow of a cistern should simply be carried through an external wall to discharge in the open, so that no drain gases can enter the cistern, and waste of water from faulty apparatus is at once seen. But the foul-water wastes of sinks, lavatories, and baths must be differently dealt with. Slop-sinks used for disposal of urine must be treated

as water-closets, but all other wastes should be made to discharge over trapped gullies in the open air. Such disconnection renders it impossible for the air from the drain to enter the house.

A suitable form of gully-trap is shown in Fig. 12. Here the waste pipe discharges through a side opening beneath the grating, and above the water in the trap. The grating must be large, with a free opening kept constantly clear, or else the primary




Fig. 13.—
"Anti-D" Trap.

object of disconnection is lost. More commonly the wastepipe pours over the grating, with more effective disconnection but some risk of splashing. Neither of these methods complies with the requirements of the Model Bye-laws that the waste-pipe shall discharge over a channel leading to a trapped gully grating at least 18 inches away.

Waste-pipes should always be trapped, and the trap situated as near as possible to the sink or bath. The interior of the pipe becomes coated with decomposing soap or fat, which is itself offensive; the higher temperature of the interior of the house leads to an offensive in-draught through an untrapped pipe, still more marked after the pipe has been heated by

discharge of hot water.

A convenient form of trap is one shaped like an S (Fig.




Fig. 14.—Bell Trap.

10), and fitted with a screw at its lowest point, by the removal of which it is possible to cleanse the trap. Where the momentum of the water-discharge is great, a useful modification of the trap is that shown in Fig. 13, by which the unsealing of the trap is prevented.

From the trap the waste should be carried as rapidly as

possible through the wall, and then descend to the gully. When the waste-pipe is long, or when two or more waste-pipes are connected, ventilation of the

trap is needed to prevent siphonage, and is effected by an air-pipe passing from the top of the trap, beyond the seal, to the open air. The common "bell trap"




Fig. 15.—Antill's Trap.

shown in Fig. 14 is open to more than one objection; the depth of the seal is inadequate (about 3 inch); it readily becomes choked with grease, being far from selfcleansing, and as a result the grating is often removed to let the water pass away, and on each such occasion the waste-pipe is untrapped.

The D-trap shown in Fig. 22 is

the worst of all, as it readily becomes filthy, and always leads to the retention of a large amount of offensive matter.

In large households the gully into which the scullery sink waste discharges is liable to become choked by sand and grease (solidifying on cooling). In such cases frequent cleansing is essential. "grease-trap" (i.e. a gully, so constructed that the outflow of liquid is not prevented by solid accumula-

tion at the surface or bottom), such as is shown in Fig.16, is sometimes adopted.

Wastes from sinks and baths in upper storeys are




Fig. 16. - Grease Trap.

sometimes discharged into the head of a large pipe, which may or may not also receive rain-water. This receives disconnection, but is liable to cause nuisance, by fouling of the large pipe with soapsuds and grease.

A better arrangement is to carry the waste directly to the gully, due precautions being taken against siphonage. A waste-pipe of 11 inch diameter is sufficient even for baths, and if two or more be joined, they may discharge into a somewhat larger pipe; a ventilating pipe being carried from the upper end of the common pipe to some convenient position above the level of the windows.

The rain-water pipe should not be used as a soil

pipe, or as a ventilating pipe to the Thorough drain. ventilation of the drain is most needed at times of heavy rain, when the rainwater pipe is least able to act in this way; and as the rain-pipe cannot be carried above the eaves of the roof, it often discharges its air contents too near to a window. For the latter reason this




Fig. 17.—Flush-tank.

pipe should not be connected with the drainage system, but should be made to discharge over a trapped gully or a channel leading thereto. It is well that a waste pipe in frequent use should discharge into the same trap, thus avoiding risk of unsealing by evaporation in dry weather.

In small houses the ordinary discharge of liquid is usually sufficient to scour the drain and make it self cleansing, but where necessary this can be supplemented by a flush-tank, periodically discharging its contents into the drain (Fig. 17). This is a cistern containing a

large outer tube closed at the upper end and inverted over a small inner tube A B, thus forming a siphon. A B is open at the top, and passes vertically down through the tank, to dip slightly under the surface of water in a chamber below the tank, the water in this lower part being kept at the required level by means of a weir, practically forming a D-trap. As the water rises in the tank from the discharge into it of a tap or a drain, some of it eventually runs over the top of the tube AB; carrying with it air which is forced through the water at B, and cannot return; siphon action is started, and the whole contents of the tank are emptied into the drain.

Testing of drains and soil pipes can be most effectually done by means of (a) the water test. For the purpose of testing the horizontal drain, a plug is inserted at that portion of the drain which is near the intercepting trap intervening between the drain and the sewer; the drain is then charged with water, care being taken to see that there is no leakage through or at the side of the plug, and the water is allowed to fill the drain to the top of one of the surface traps connected with it. At the end of from half an hour to an hour the maintenance or subsidence of the water level in this trap will show whether there is any leakage. In testing a large system of drainage it is necessary to take each section separately. In testing soil pipes with water, the plug must be inserted at the bottom, and if several water-closets are connected with the same soil pipe the trap of each must be plugged.

(b) Smoke pumped into the drains previously plugged in the same way serves also as a useful test. The process must be continued until the smoke is seen issuing from the soil pipe or ventilating pipe, and if several of these are connected with the same drainage system it is necessary that each must be plugged at the top as smoke issues from it until the whole system is charged, care being taken to prevent the smoke from being forced through the traps; smoke in these circumstances, when there is leakage, soon finds its way into the interior of the

(c) Peppermint was formerly used in the same way; two or more ounces of the oil was either poured down a soil pipe with

a pailful of boiling water, or down some trap situated outside the house connected with the soil drain; the scent of the oil of peppermint found its way into the house in the event of defective pipes. Chemical "testers" such as Kemp's or Kingzett's are now commonly used. These are composed of some strongly




Fig. 18.—Valve Closet.

A, basin: flushing-rim above; B, trapped overflow; CC, valve, closed and open; D, valve-box; EE, safe; F, trap, ventilated at G.




Fig. 19.—Long Hopper Closet.




Fig. 20. - Wash-out Closet.




Fig. 21.—Wash-down or Short Hopper Closet.

smelling chemical, e.g. calcium phosphide, which, when moist, evolves phosphoretted hydrogen, and when the tester is washed into the drain over the trap, the water soaks the paper fastener, and the spring being liberated the contents escape. If the drain be defective the odour escapes into the house in about fifteen minutes.

The surfaces of the water-closet must be non-absorbent, and in its shape and capacity the basin must be able to receive the water that is discharged into it, and to allow all matter it receives to fall directly into water without soiling the sides. In both the valve closet (Fig. 18) and the wash-out pattern (Fig. 20), water is held in the basin independently of that in the trap below. In the one case it is held by a movable valve, in the other by a fixed dam, over which the contents are carried by the flush. This




Fig. 22.—Pan Closet, with D-trap.

last form is not wholly satisfactory. In the plug closet the valve or dam is replaced by a solid plug, which works vertically in a side-chamber similar to that shown in Fig. 18. Hopper closets are of two kinds; the basin of the old long hopper (Fig. 19) is from its shape and construction liable to become filthy, and it has usually a bad flush; the short hopper or washdown (Fig. 21) is free from objection, and possesses the advantage of a large water surface owing to the form of

the trap being such as to allow the water to stand high in the basin. Added to this there is a deep water seal, and the cleansing of the trap is greatly facilitated. There is at present a wholesome tendency to dispense with all unnecessary woodwork and fittings, and to reduce the closet to its simplest form.

The old container closet, or pan-closet (Fig. 22), is especially objectionable, and is prohibited by the Model Bye-laws. The basin discharges into a large receptacle, which becomes foul and, every time the valve is opened, allows polluted air to escape.

The apparatus, of whatever kind, should be provided with a flushing rim which directs the water

over the whole surface, and maintains it in a state of cleanliness. It is essential that the flush should be of sufficient volume (not less than two gallons), delivered with sufficient force (and for this purpose the pipe must not be less than 1½ inch in diameter), and properly directed so as to wash the basin effectually. The flush must on no account be taken from a service pipe or a cistern that supplies water for household purposes, but from a special cistern. Gases and even filth have been found to make their way from the closet basin up the (usually empty) deliverypipe.

It is unnecessary to discuss at any length the traps used in connection with water-closets. The Strap is here again found to be a useful form, but liable occasionally to become unsealed when the momentum of the water-discharge is great; for this reason, therefore, the form of trap (Fig. 13) recommended in connection with waste-pipes to obviate this difficulty is preferable. The D-trap is altogether

inadmissible.

Water-closet traps are liable to become unsealed inthe same manner as those of waste-pipes (page 236). Wherever two or more discharge into the same soil pipe, ventilation of the traps is necessary; the ventilating pipe from the trap of the lowest water-closet may receive the ventilating pipe of each above, and be finally connected with the soil pipe above the highest water-closet, but while this arrangement is permissible it is frequently preferable to carry the antisiphonage or ventilating pipe independently above the roof.

A closet or bath on an upper floor is usually provided with a safe to receive any overflow or splashings. The waste-pipe from the safe may be treated in the same way as an overflow from a cistern, i.e. simply carried through an external wall,

and made to discharge a few inches from the brick work.

Trough closets are sometimes adopted in schools or factories. A long trough filled with water passes beneath the seats of a number of closets placed side by side. At stated intervals the trough is flushed, automatically or by an attendant, and the contents are carried away to the sewer through a trap at the outlet end. The automatic flush may be effected by a tilting receiver placed at the upper end of the trough, and so arranged as to capsize and discharge its contents as soon as it becomes filled by a regulated stream of water; or still better by a flush tank fed by a stream of water and discharging itself into the trough by siphon action as soon as it becomes full. In either case the frequency of the flush can be regulated by the supply of water.

In waste-water closets the flush is effected by the household waste-water, the object being of course to avoid the expense of a separate water supply. The drainage from a row of houses is conducted to a flush tank placed at the end of the row farthest from the sewer. From the tank a large pipe drain passes to the sewer, and separate closets are placed upon its course, the excreta simply falling through large vertical drain pipes into the drain below, and remaining there until the next flush sweeps them away into the sewer. The drain has a siphon trap after passing the lowest closet, and a manhole or access chamber at the same point for use in the event of obstruction.

In like manner the waste-water of single houses has been utilised for flushing single out-door closets. For closets of this type, which are known as slop closets, the household waste-water, supplemented if necessary by the rainfall, is collected in a "revolver," or "tumbler," an earthenware vessel working on

pivots, and so constructed as to overbalance when full, discharging its contents forcibly along the drain, and then swinging back to its former position. On the drain, beyond the tumbler, is a large siphon trap, and the closet consists of a wide vertical shaft over this, so that the excreta fall into the water of the trap, and are carried away by the next flush.

The use of waste-water for flushing is attended with practical difficulties from the over-concentration of sewage, and the liability of automatic appliances

to get out of order.

Intermediate between the water-carriage and dry systems come certain Continental and other methods which may be described briefly. The fosse permanente is a large water-tight underground pit, to which pipes from the closets conduct the excreta; the pit is emptied at long intervals by suction through hose into an air-tight cart. The fosse mobile is a closed movable tub placed outside the house, and connected with a fixed pipe that communicates with all the closets, etc., in the house; the tub is simply removed and replaced by another when full. In Liernur's system a number of closets are connected by means of iron pipes with an air-tight tank, which is at intervals exhausted of air, so that the excreta are drawn into it.

Dry systems of excreta removal, otherwise known as "conservancy systems," include privies, pail-closets (with or without ashes), ash-closets, and dry earth-closets. In some localities privies are called earth closets, but this use of the term is misleading.

The privy and ashpit, or midden system, which is still employed in certain towns, and in most rural districts, aims at the deodorisation and drying of the excreta by admixture with ashes. The best construction is that of a small watertight pit, not drained but roofed over to exclude rain, and so arranged that the excreta and ashes become thoroughly mixed. For this purpose either the ashes must be thrown in through the closet seat, which may be hinged so as to be lifted en masse, or else a "shoot"

or sloping slab must conduct one or the other to a common point. The floor should be smooth, and raised a few inches above the level of the adjoining ground. Very few middens fulfil these conditions, however.

The contents ought to be removed at fixed short intervals by the S.A., and the work should be done at night or early in the morning so as to minimise the nuisance. Unfortunately it is still common to leave the emptying of ashpits to contractors, or even to the owners and occupiers, and the interval is frequently several months instead of a week, as prescribed by the Model Byelaws.

Privies alone, without admixture of ashes or other deodorants, are also met with. The receptacle should be small, and its contents frequently removed, but at best the system is more offensive than when ashes or

earth are added.

Pail- or tub-closets are simply miniature privies, in which the pit is represented by a movable pail placed beneath the closet seat. The full pails are removed and replaced by clean empty ones at regular intervals not exceeding a week. Close-fitting covers should be used during removal of the full pails. There are several modifications of this system. On the "Rochdale system" the pails contain excreta alone, while in some towns the ashes and other dry household refuse are added; or the ashes are thrown upon a sifter, and only the fine ash falls into the pails; or the Goux system is used, the cleansed pails being lined with a dry, absorbent packing of compressed peat or like substance.

In Morell's cinder-sifting ash-closet the ashes are thrown in by a separate opening, and by a simple automatic arrangement are sifted, so that only the fine ash falls upon the excreta, the cinders finding their way to a separate chamber for further use as fuel. The receptacle is permanent and in this respect resembles a privy rather than a pail-closet, but it is small and so arranged as to be readily emptied by a special opening. Taylor's closet is somewhat similar in principle, but the urine is collected separately.

Dry earth-closets have a small receptacle, fixed or movable, beneath the closet seat. Clean, dry earth (about 1½ lb.) is thrown upon the excreta, either by hand or by automatic delivery from a hopper, every time the closet is used. A constant supply of fine dry earth is needed, ordinary mould being the best, owing to its richness in nitrifying organisms. Sand is less efficacious.

Destination.—Water-carried excreta pass into the ordinary sewers, and their ultimate disposal will be considered in another section.

Pails are emptied and cleansed at a central depôt, and are then ready for further use. Their contents are sent into rural districts as manure, either in the crude state or after conversion into "poudrette" by drying and other treatment. For the manufacture of poudrette the excreta (free from ashes) are heated in revolving cylinders, so as to drive off the moisture. Sulphuric acid is often added to fix the ammonia.

Contents of ashpits are commonly sent away at once into the country. The refuse of ash-closets or pail-closets is also available as manure.

The earth from earth-closets can be dried and used again and again, and finally restored to the garden or field. The organic matter disappears by nitrification.

Comparative advantages of water-carriage and dry systems:-

1. Hygienic.—Water-carriage is by general consent the most satisfactory in houses of the better class, where cleanliness and freedom from effluvia are the main considerations, and due care in use can be relied upon. The excreta are at once carried right away from the premises. Water-closets may

safely be constructed practically within the house, a matter of great convenience. The objection that the sewers receive the specific poison of enteric and other fevers may be met by the use of disinfectants, and has not yet been proved to be weighty. It has been urged that indoor water-closets afford facilities for the entrance of sewer air, especially in the case of obstruction of the sewers by floods or other causes. Indeed, it has been observed in more than one enteric outbreak that the inmates of large water-closet houses upon high ground suffered more than the poorer inhabitants of lower districts with outdoor closets. This danger, however, can scarcely occur unless there is grave defect in construction or management, and it extends equally to indoor sinks. As applied to poorer dwellings water-closets often fail. Owing to carelessness or misuse, the mechanism becomes deranged, or flushing is neglected, and the drains become choked by bulky objects improperly introduced into them. Automatic flush closets, trough closets, and even slop-closets, if kept under supervision, are to a great extent free from this objection, since they are not liable to become deranged by carelessness on the part of those using them. Water-closets of ordinary construction are made serviceable, even in houses of the poorest class, by careful supervision on the part of the S.A. Neglect of hand flushing may be met by automatic mechanism connected with the door or the seat. The water-supply of outdoor closets is liable to be stopped by frost, but this rarely happens if the cistern and pipes are properly cased.

Privies and ashpits are being abandoned in most towns, although still almost inevitable in country districts where there is neither efficient drainage nor organised scavenging. The retention of large quantities of excreta and other organic matter near dwellings is in itself objectionable, and the nuisance becomes great at the time of clearing out. It is probable that such an arrangement favours the spread of diseases that have relation to filth, and especially those in which the poison is contained in the excreta. The case becomes worse when, as almost invariably happens, the pit leaks and allows its contents to pollute the soil and the

subsoil water.

Pail-closets have the advantage of extreme simplicity, and cannot be deranged by carelessness or anything short of wilful destruction. Although not removed instantly, as in water-closets, the excreta remain near the dwelling for a few days only, instead of putrefying for weeks or months as in middens. The weekly or semi-weekly visits of the scavengers ensure a certain degree of cleanliness, apart from the mere substitution

of cleansed pails. It is claimed that in the event of such diseases as enteric fever occurring, the pails and their contents can be disinfected; unfortunately not all the infective matter is thrown into the closet, and at best its disinfection in any true sense would be difficult; but it is quite possible to adopt the simple precaution of supplying specially marked pails to infected households, and cremating the contents of such pails when collected. Pail-closets that receive excreta only are inevitably offensive, however frequently changed. The addition of ashes lessens the effluvia, and the average amount of ash produced by an artisan household is found to be sufficient for the purpose.

Dry-earth closets are convenient and free from effluvia. The difficulty of ensuring a constant supply of suitable dry earth, and of removing the products, has hitherto prevented their adoption in towns. In rural districts, where these difficulties do not exist, they offer great advantages, but can only succeed when used with care and kept under supervision.

2. Economic.—Water-carriage involves some cost, either to the individual or to the public, in the extra consumption of water, although this objection does not apply to those systems in which the flushing is effected by the household waste water. Drains and sewers have to be provided in any case, so their cost cannot be charged to the account of water-carriage systems. The increased volume of sewage in water-closet towns or districts is far less material than is generally assumed, and forms at most only a small fraction of the enormous volume derived from rainfall, soil drainage, household waste water, trade effluents, and other sources. It has been found that the addition of water-closet sewage scarcely alters the chemical composition of the average sewage of a town. Hence the necessity of treating the sewage by irrigation or other means is quite independent of the admission of water-borne excreta to the sewers.

Water-carriage obviates some part of the expense of the scavenging required by the pail-closet system, but there must still be a costly organisation for collecting the dry household refuse, which, without the addition of excreta, is an unsaleable commodity, and further expense is therefore incurred in disposing of it after collection.

A similar objection is often urged from the theoretical side, namely, that the enormous dilution of excreta by water renders it difficult or impossible to recover them again for manurial purposes, and hence the soil is steadily impoverished by depriving it of what should naturally be returned to it in exchange for the cross that it yields. Vivian Poore's dictum

was that the proper destiny of organic refuse is immediate burial just below the surface of the soil, and on this principle he objected to water-carriage. This argument has some weight as bearing upon the still common practice of pouring the sewage of towns into rivers or into the sea.

From the purely financial point of view the objectionable

midden system probably costs less than any other.

The pail system involves a heavy initial outlay in buildings and plant, and an expensive staff to manage it efficiently. On the other hand, some part of the expense is recouped by the sale of manure, and a scavenging staff must in any circumstances be maintained for the removal of dry refuse. The market value of the excreta is greater if unmixed with ashes, but it is questionable if the gain is sufficient to compensate for the cost of a separate collection of dry refuse, which, as already explained, is unsaleable. The dry systems ensure the return of most of the solid excreta to the soil, but a great part of the liquid excreta, which are more important as fertilisers, goes with the household drainage in any case.

Reviewing the whole question, the privy system must be regarded as contrary to sound principles of hygiene wherever facilities exist for any other arrangement. Water-closets are most suitable in town, and water-closets or earth-closets in the

country.

Removal of ashes.—Where water-closets, or pail-closets for excreta alone, are in use, the ashes and kitchen refuse have to be removed separately. Small covered tubs or boxes (preferably of metal) should be placed in a yard or outhouse, and emptied at short and regular intervals by the scavengers. Frequently, however, large "bins" or fixed "ashpits" are employed, the contents of which are allowed to accumulate for long periods.

Household refuse ought to consist of little more than ashes, the animal and vegetable refuse from the

kitchen being easily burnt.

Trade refuse, from slaughter-houses, markets, fish and fruit shops, and many other sources, requires prompt removal. This duty is often left to the proprietors, who make private arrangements with farmers, etc., to remove the refuse for use as manure,

or convey it at their own cost to some suitable place. It is difficult to ensure proper system and regularity unless the **S.A.** itself undertakes the task, and the same difficulty may occur in urban districts in regard to manure from stables and cowsheds. Under the Public Health (London) Act, 1891, s. 36², the local authority may by formal resolution prescribe frequent removal.

Road sweepings and the contents of street gulleys have also to be dealt with.

Destructors.—Any kind of refuse which contains organic matter, but which cannot be at once disposed of as manure, should be burnt. The refuse is thrown in at the top of a furnace, and becomes dry as it sinks down; all organic matter is burnt off, and at intervals the mineral matter, or "clinker" (forming about 30 per cent. of the refuse burnt), is raked out below, to be used for road-making, filling up hollows, etc. In the best modern destructors the clinker is withdrawn in large slabs and freely slaked with water to prevent flaming when tipped. Silica and ferric oxide are the chief constituents of clinker, which is now much used for bacteria beds, mortar-making, etc. After the furnace is started, the organic matter in the refuse is usually sufficient to maintain the fire without the addition of other fuel.

Numerous forms of destructor are now in use. They consist of furnaces of a varying number of cells and are charged in different ways ("shovel fed," "top fed," mechanical charging, etc.). The furnaces are not unfrequently multitubular, or fitted with Babcock and Wilcox boilers, and the necessary draught is obtained naturally or by means of fans or steam blowers. They may be divided into two chief groups, the "single cell system" (e.g. Horsfall's, Fryer's, Warner's, Baker's) and the double cell or continuous grate system (e.g. Meldrum's, Heenan's,

Sterling's); there are, of course, various modifications. By the single-cell system is meant a destructor with a boiler between every two cells, so that there is no intermingling of the gases from two or more cells. It is desirable to have complete combustion in the cell itself, but if necessary the fumes can be passed through a second furnace.

Sewers may be required to convey :-

(i.) Household effluents, including waste water from baths, lavatories, sinks, and all liquid refuse and urine. Where there are water-closets, the volume of sewage is somewhat increased, but its composition is little altered. Latrines, etc., attached to works, schools, or public resorts may be included under this head.

(ii.) Trade effluents of all degrees of impurity, from the comparatively pure water from condensers to the foul effluents from dyeworks.

(iii.) Water used for public purposes, e.g. watering

streets, fountains, public urinals, baths, etc.

(iv.) Rainfall, of which the first portions (especially road-washings) are extremely impure.

(v.) Subsoil water, and even small natural streams.

Modern sewers, like house drains, are impervious tubes, oval or round in section. Their size varies according to requirements, say, from 9 inches to 12 feet in diameter. If the diameter is not greater than 1½ or 2 feet, glazed earthenware pipes set in cement are usually employed, laid in the manner described in speaking of house drains. Sewers of larger diameter than 18 inches are generally constructed of brick set in cement upon a bed of concrete, and are oval or ovate in section, so as to secure a better scour and less friction when little sewage is passing than if the section were circular. Sewers should be laid as far as possible in straight lines, with means of inspection and access at every change of direction. All junctions

should be obliquely in the direction of flow, and the tributary sewer or drain should have a fall into the sewer at least equal to the difference in their respective diameters. It is desirable that the rate of flow should be not less than 2 feet per second, and 3 feet is better. Hence the minimum gradient should range from 1 in 250 to 1 in 750, according to the size of the sewer. The gradient should preferably be uniform, except at angles and junctions, where it is desirable to allow extra fall.

Manholes should be provided at short distances, and at all important angles, junctions, and changes of gradient, to allow of access for inspection and flushing. Flush-tanks may with advantage be placed at the head of each sewer. Flushing is chiefly required in sewers with insufficient gradients, at "dead ends," and in hot dry seasons, when the flow is smallest and decomposition most rapid. It may be effected more or less automatically by means of flushtanks, or by temporarily damming back the stream and then allowing it to escape with a rush, or by suddenly discharging a large volume of water into the sewer through a manhole. Little benefit is likely to result from the perfunctory "flushing" of sewers that is commonly practised, consisting as it does of simply pouring a small and inadequate volume of water into the sewer through a hose, at intervals of several weeks. There is little to be gained by the addition of "disinfectants" to the water used for flushing.

The air of sewers ("sewer gas") has no constant composition, and if the sewer be properly constructed, well ventilated, and sufficiently flushed, may differ but little from outside air. Much depends upon the sewage being removed quickly, or, on the other hand, being allowed to stagnate and undergo decomposition. In the latter case the air of the sewer

becomes foul; oxygen is lessened, carbonic acid increased, and there is much organic matter, together with variable quantities of marsh gas, sulphuretted hydrogen, and ammonium sulphide. The exact composition of the organic matter varies, but its properties are similar to those of the organic matter in respired air. Micro-organisms adhere to moist surfaces, and hence the air of well-constructed sewers is, on the whole, remarkably free from them, except near fresh air inlets and at junctions, where splashing occurs. Neither bacteria nor other solid particulate matters are, under ordinary circumstances, given off from quiescent liquid surfaces; but if putrefaction be allowed to occur the bursting of bubbles may recharge the air with them. Haldane found in the air of a well-constructed but unventilated sewer in Bristol 20 volumes of carbonic acid per 10,000, but only two micro-organisms per litre, half of which were moulds; and Lawes and Andrewes demonstrated that the air of well-constructed London sewers contained fewer bacteria than the air of the street above, and the species found were not of sewage origin. It thus becomes probable that sewer gas can rarely, if ever, convey pathogenic organisms.

Air contaminated by sewage emanations may, however, be a cause of diarrhea and other gastrointestinal disturbances, and of certain forms of sore throat. Anæmia, depression, and general ill-health may result from protracted exposure to such an atmosphere. Cholera, enteric fever, pneumonia, erysipelas, puerperal fever, and diphtheria have a much heavier incidence, both in numbers and severity, upon persons exposed to these conditions. It is not necessary to assume an origin de novo in such cases, or even in the case of diarrhea and sore throat, the evidence being consistent with the supposition either that the specific poison is sometimes carried by such

emanations, or that their effect is merely to predispose to the disease. There is no evidence of any specific relation between sewer gas and small-pox,

measles, or whooping-cough.

Men working in well-ventilated sewers are not found to suffer any ill effects, but if ventilation is wanting they are liable to ophthalmia and to syncopal or apoplectic attacks. It is said that venereal disease is greatly aggravated by this employment. Meat, milk, and other food substances readily decompose if exposed

to sewage emanation.

Ventilation of sewers.—Foul gases are given off by sewers, more especially if, from insufficient gradient or obstructed outfall, they become stagnant or "sewers of deposit." Hence it becomes necessary to provide means of vent or ventilation in order to avoid the risk of the pent-up gases forcing their way into houses through traps.* For this purpose openings are provided at intervals which should not exceed 100 yards. Some of these openings act as inlets, others as outlets, and the latter only are, of course, liable to be offensive. The conditions that determine the direction of the currents of air are complex, including force and direction of wind, internal and external temperature, volume and velocity of sewage, etc., so that the result cannot be predicted, and is not even constant in the same sewer. Speaking generally, the tendency is for the air current to pass in the reverse direction to the sewage; that is, from a lower to a higher level, so that the highest openings are most liable to be outlets, and the lowest to be inlets. Nevertheless, every opening may at different times act both parts.

The plan usually adopted is to provide grated openings in the roadway, leading directly to the sewer beneath, a cage being suspended immediately beneath

^{*} In Bristol the sewers are not ventilated. The house drains are, however, provided with interceptors, and all street gullies are trapped.

the opening, to catch any solid débris that may fall through. It was formerly customary to place trays of charcoal in the openings in order to deodorise the sewer gases, but the charcoal obstructed the opening and soon became inert. To obviate annoyance from effluvia at outlets, ventilating shafts (in substitution for these openings) are often attached to houses or even to trees; these shafts should be, but rarely are, of sufficient calibre efficiently to ventilate the sewer. Sometimes factory chimneys are utilised for this purpose by carrying the vent into the stokehole or chimney. Another device (Keeling's) is to ventilate the sewers through hollow lamp posts with gas jets inside serving to create a strong upward current and (it is claimed) to destroy all effluvia.

If carried out on a large scale these special arrangements are costly. It is contended by the advocates of simple openings at the street level that the true remedy for offensive outlets lies in proper construction and flushing of sewers, and increasing the number of openings—that is, in approximating to the condition of an open sewer. Some authorities rely upon ventilating the sewer by means of the vents of each house

drain connected (see p. 238).

Rain and subsoil water.—If the sewers are water-tight, as they should be, the subsoil water cannot gain access to them, and separate provision must be made for draining the soil, although a certain quantity of water flows away along the comparatively easy track outside the sewer. Sometimes provision is made for this by leaving a space beneath the sewer for the ground-water to pass along.

It is becoming customary to exclude also the rainfall from the sewers, and thus avoid dilution of the sewage and sudden changes in volume, which add to the difficulty of treating it by precipitation or filtration. The first washings of the streets are very

foul, and should be admitted to the sewers. When the "separate" system is adopted, the rain water is taken away by another system of drains, which may with advantage be permeable, and so available for draining the subsoil.

Capacity of sewers.—In towns only a comparatively small surface of ground is pervious, and almost all the rainfall rapidly makes its way to the sewers. Hence in planning the size and gradients of sewers it is necessary to provide for the immediate removal of the maximum hourly rainfall, say, 1 inch per hour plus the maximum volume of sewage from other sources.

Parkes gives the following formula for calculating the discharge from a sewer:—

$$V = 55 \text{ A} \times \sqrt{2DF}$$
.

V being the discharge in cubic feet per minute,

A, the section area of the current,

F, the fall, in feet per mile,

D, the hydraulic mean depth, in feet. This is \(\frac{1}{4}\) of the diameter of the pipe, if running full or half-full; in all cases it is equal to the section area of the current divided by the "wetted perimeter"—that is, by the length of the arc formed by that part of the section of the circumference which is in contact with the stream.

When a sufficient fall cannot be obtained, Shone's pneumatic system offers great advantages. The sewage is conducted to one or many central collecting tanks by gravitation through drains and sewers of ordinary construction, and is forced through pipes, leading from the tanks to the outfall, by means of compressed air. Thus difficulties arising from inequalities or from low level of districts to be drained can be overcome.

The composition of sewage varies from day to day, and from hour to hour, since both the filth and the diluting water are inconstant in amount. There are variations according to season and weather, and trade effluents may be important factors. The admission or exclusion of rain-water makes a very great difference, but the presence or absence of water-carriage of excreta does not seem to affect the composition of sewage very materially. The Rivers Pollution Com-

missioners found the average composition of sewage to be somewhat as follows:-

Constituent Parts.					PARTS PER MILLION.	
					Water-closet Towns.	Midden Towns
Total Solid Residu	e .				1369	1215
	(Organic				205	213
Suspended Solids					242	178
THE RESERVE OF THE PARTY OF	(Total				447	391
Dissolved Solids					722	824
Organic Carbon					47	42
Organic Nitrogen					22	20
Ammonia .					67	54
Total combined Ni	trogen			100	77	65
Chlorine .				10	107	115

From other sources it appears that the phosphoric acid amounts to about 25 and the potash about 15 parts per million. These proportions are, of course,

liable to very wide variation.

It is calculated that in an ordinary population 21 ounces of solid excreta and 40 ounces of urine, together containing 150 grains of nitrogen,* are yielded per person per day, taking an average of all ages. This corresponds to about 10 lbs. of ammonia per annum, and a theoretical manurial value of 6s. 8d., to which the constituents of the urine contribute six-sevenths, and the fæces one-seventh only. The manurial value of sewage is dependent upon the combined nitrogen, potash, and phosphoric acid. The annual amount of sewage per head of population may be taken as 100 tons, the theoretical money value as manure being about 17s., to which the dissolved matters contribute 15s., and the suspended matters only 2s. Sewage also contains variable quantities of household waste, rain and storm water, grit, gravel, sand, and, in some localities, manufacturing waste products.

^{*} This is Parkes' estimate. Frankland's estimate, for adult males, is 250 to 300 grains.

Bacteria in sewage are chiefly denitrifying and decomposition organisms. Houston found in London crude sewage at the Barking outfall an average of nearly 4,000,000 micro-organisms per c.c., and in one sample of domestic crude sewage 28,000,000. All kinds of species are present (bacilli preponderating) and there are many spores. The most common are B. coli (600,000 per c.c.), the Proteus family represented by the "sewage proteus" (100,000 per c.c.), B. enteritidis sporogenes, sewage streptococci, and liquefying bacilli. Pathogenic organisms also occur, but, owing to the struggle for existence, rapidly disappear. All the organisms required for bacterial assimilation are present in the sewage itself. The artificial bacterial treatment of sewage is an application of the principles of nitrification as stated by Schlesing and Müntz.

Disposal of sewage. - In rural districts the household and other liquid refuse, if not turned into a ditch or watercourse, is usually conducted to a cesspool or collecting tank, from which it is periodically removed and used as manure. The former plan is bad in every respect. The latter is permissible only if the cesspool is made watertight and properly covered and ventilated, regularly emptied at suitable intervals, and situated at a sufficient distance from any house, road, or water supply. The annotated Model Bye-laws require a minimum distance of 50 feet from any dwelling, and 80 feet from any water supply. Sometimes the sewage is led to a pit, or "dumb-well," in porous soil, the solids being removed at intervals and the liquids escaping into the ground. This practice is dangerous if the pit is near to dwellings or water supplies, and is not free from risk of polluting water in any case. Simple irrigation is free from objection if well managed, even on the small scale. "Subsoil irrigation" may

work well when the conditions as to soil and fall are suitable; the drainage is conveyed into a branching system of loosely laid agricultural pipes, which allow it to escape into the subsoil. The pipes are laid about a foot or more beneath the surface. It is often necessary to relay them at intervals, owing to deposit of grease and other obstructive matter, but this may to some extent be prevented by the use of flush-tanks and grease-traps at the house-end of the drain.

The chief point is to expose the sewage to the influence of growing vegetation, not so much for the sake of utilising it as of rendering it innocuous. Sewage applied to the surface is partly absorbed by vegetation, partly oxidised (nitrified) in the superficial layers of earth. The sewage passed into leaking cesspools or dumb-wells escapes both forms of purification, and simply pollutes the soil and the subsoil water.

The disposal of town sewage may be effected in a variety of ways. The oldest and simplest is to pass it without purification into natural watercourses, a plan still common. The sea affords a ready means of disposing of the sewage of coast towns. It is allowed to escape only with the ebb tide, and reflux is prevented by valves. If it is conducted by pipes sufficiently far out to sea, it may not be washed back by currents or by the returning tide to foul the shore. Such a degree of success is far from being the rule, and there are the further objections that the removal of sewage by this means is only intermittent, and that the sewage is (at best) wasted. To a certain extent organic impurities in streams are got rid of by oxidation and by the growth of plants and other organisms, but this possibility of subsequent partial purification cannot justify the gross pollution of a stream at any given point. The purification can rarely be complete, and in no case can there be any

certainty of the destruction of specific pollution, a consideration that is especially important when the stream furnishes the water supply of districts lower down.

The alternative is to purify the sewage with the object of oxidising or retaining its dissolved and suspended impurities, and allowing only the water (with, at most, harmless dissolved matter) to pass into watercourses or the sea. This may be done by precipitation, filtration, irrigation, or bacterial treatment.

Precipitation processes aim at throwing down the organic matter chemically or mechanically, by the addition of a reagent to the sewage. For the most part their success in this respect is only partial, and they leave the ammonia and chlorides in the effluent. The reagents, dissolved or suspended in water,* are added to the raw sewage either in the sewers (Scott's and Conder's processes) or more usually in a conduit at the works. In either case the flow of the reagent is regulated as required, and it becomes thoroughly mixed with the sewage before entering the tanks. The settling tanks are large and are usually worked in series, the overflow from each into the next being over a broad shallow sill, so that only the comparatively pure upper water may escape. The current being very slight in the tanks, suspended matters are gradually deposited, accumulating at the bottom in the form of sludge, which is periodically cleared out, dried, and sold as manure, or dug into the ground, or (in Scott's process) burnt into cement. A similar subsidence takes place if no precipitant is used, but the dissolved organic matter remains in solution, and the settlement of the suspended matter is much slower and less complete. A good precipitation process will rapidly clarify the sewage by removing

^{*} In the alumino-ferric process, cakes of the reagent are laid in the sewage current, and dissolve slowly.

all, or nearly all, the suspended impurities, and will, moreover, carry down a greater or less part of the dissolved organic matter. The effluent will be fairly clear, but will contain the chlorides and certain other salts, ammonia, and more or less of the dissolved organic matter. It is therefore necessary to subject the effluent to further treatment, that is, to filtration, in order to oxidise the remaining organic matter.

Among artificial filtering media, employed after precipitation, are various forms of carbon, ferruginous materials, and burnt "ballast." All of these, like land filters, serve mainly as a nidus for the aërobic nitrifying organisms, and their efficiency depends largely upon proper management, and especially due aëration. The flow should be intermittent and not too rapid. Coke is widely used. Garfield finds that fine coal filters give excellent results, and possibly the non-porosity of the particles is an advantage. Ferruginous materials include "polarite," "magnetite," and "magnetic carbide"; they should be covered with a layer of sand, renewed at intervals.

Precipitation affords no guarantee of the removal of microbes, pathogenic or otherwise.

Many of the precipitation processes described in text-books have been abandoned on account of cost or want of efficiency, and have now only a historic interest. Lime is by far the most generally employed, either alone or in conjunction with other reagents. It has been used together with clay (Scott), calcium phosphate (Whitthread), magnesium chloride and tar (Hillé). The "Amines" process consists in adding lime together with a little herring brine. The latter, by virtue of the trimethylamine and other ingredients (aminol), acts as an antiseptic. The effluent does not decompose, and yields no microbes on cultivation. The precipitation is rapid. Lime with ferrous sulphate, ferric chloride, or aluminium sulphate gives a bulky precipitate of the respective hydrate, which entangles and carries down organic matter with it. Crude aluminium sulphate, made by acting upon clay or shale with sulphuric acid, is Anderson's (and also Bird's) precipitant; and sulphuric acid with mineral phosphate of alumina is the basis of Forbes and Price's process. Spence's "alumino-ferric" contains a very

small proportion of iron salts. Sillar's A B C process consists

in adding alum, blood, clay, and charcoal.

Another precipitant is "ferrozone," made by acting with sulphuric acid upon "polarite," described as a specially prepared ore of iron. The effluent is filtered through a bed of polarite. Another iron process, differing from the rest in being applied at the head of the sewer or house drain, is Conder's, in which a regulated quantity of ferrous sulphate is added constantly and automatically by means of a stream (of water or sewage) passing through a vessel ("ferrometer") containing crystals of ferrous sulphate. It seems to be successful in preventing decomposition and effluvia, and in keeping the drains free from deposit. A saturated oxy-chloride of iron is used as a precipitant, in the ordinary way, in the "Clarine" process. Use is sometimes made of chalybeate waters for precipitation

Webster's electrolytic process adds iron indirectly. The sewage is made to flow through a series of tanks in which it is exposed to a powerful electric current passing between electrodes consisting of iron plates. Ferrous hydrate is continuously formed, and reacts at once in the nascent condition upon the sewage, causing a rapid and fairly complete precipitation of solids. A clear and slightly greenish effluent results, which is remarkably free from microbes and has little tendency to secondary putrefaction. In the Hermite process

electrolysed sea-water is employed.

In the black-ash process, a waste product from alkali- and soap-works is used as a precipitant. This refuse contains calcium sulphide, which by exposure to air is partially oxidised into calcium sulphite and hyposulphite. The prepared black-

ash is used together with lime.

Lime processes in general yield an alkaline effluent, which has a tendency to "secondary decomposition." A lime effluent that is bright and clear on leaving the works will often be found to become turbid and offensive, if its run can be traced for a few hundred yards, without excessive dilution by other water.

Filtration of sewage may be adopted either alone or as a supplement to a precipitation process.* The filtering media employed are land, certain ferru-

^{*} E.g.: - Closed septic tank and contact beds. Open septic tank and contact beds. Chemical treatment, subsidence tanks, and contact beds. Subsidence tanks, and contact beds. Contact beds alone. Open septic tank. Chemical treatment, and subsidence tanks. Subsidence tanks.

ginous materials, and coke or other matter containing carbon. By filtration it is sought to remove the suspended matters, and to oxidise the organic matters and ammonia. The effluent should contain only chlorides and other dissolved mineral salts and nitrates, but some free and albuminoid ammonia is always found upon analysis.

Intermittent downward filtration may be defined as "the concentration of sewage at short intervals, on an area of specially-chosen porous ground as small as will absorb and cleanse it, not excluding vegetation, but making the produce of secondary importance," the flow of sewage being suspended from time to time in order to allow the filter to become charged afresh with

oxygen from the air.

Land filtration requires a porous soil, underdrained by porous pipes at a depth of four to six feet. The sewage is distributed over the surface by means of branching carriers or trenches, controlled by sluices, so that each portion of the ground in turn receives the sewage for a few hours, and each has intervals of rest. The soil is raised in ridges, upon which vegetables are grown, the sewage flowing along the furrows between the ridges. The utilisation of the sewage is a minor consideration; but vegetation is useful in absorbing and assimilating the organic matter. Land-filters planted with osiers (osier-beds) are sometimes employed for the purification of village sewage, or for the occasional treatment of storm waters that cannot be dealt with otherwise. The osiers absorb water and sewage matter freely; but constant. care is needed to prevent choking of the surface-soil, or of the subdrainage.

Broad irrigation "means the distribution of sewage over a large surface of ordinary agricultural ground, having in view a maximum growth of vegetation (consistently with due purification) for the amount of sewage supplied." For broad irrigation, as for simple filtration, the soil should be porous, but not necessarily underdrained. The sewage is distributed over each portion of the ground intermittently, by means of branching carriers, which pass along ridges of soil twenty or thirty feet apart, or along contours of slopes, and are controlled by sluices. At intervals the sewage is turned into each carrier, and overflows down the slope. The sewage is screened before distribution, unless it has been subjected to preliminary filtration or precipitation. The effect is similar to that of simple intermittent downward filtration, of which, indeed, this is only a modification. The nitrates may, however, be reduced by vegetation. The crops grown upon "sewage-farms" are very heavy. Italian rye-grass is well adapted for the purpose, since it grows rapidly and absorbs much sewage, but many other forms of vegetation can be substituted. If the land is limited in area, provision may be made for temporary excess of sewage after heavy rainfall by setting aside a portion of the ground as a land-filter.

It is usual, in cases where the approval of the L.G.B. is required, to provide at least an acre of land for each 300 of population if broad irrigation is adopted; or an acre for each 1,000 if there be either broad irrigation coupled with precipitation or intermittent downward filtration through underdrained land; or an acre for each 2,000 if precipitation is followed by land filtration. It is considered necessary to secure land, to the above extent, even if artificial filters are provided.

Apart from the action of vegetation, the purifying effect of different soils shows considerable variation. The Rivers Pollution Commissioners found that a cubic yard of chalk or sand effectually purified 5.6 gallons of sewage per diem applied intermittently; while a sample of loam purified 9.9 gallons under the same conditions. Peat had slight purifying power at first, but improved with repeated use, owing, perhaps, to increase in the nitrifying organisms. Intermittent filtration through suitable soil removed 70 per cent. of the organic nitrogen, and upwards of 80 per cent. of the organic carbon. On the whole it may be said that peat and stiff clay lands are generally unsuitable for the purification of sewage, that their use for this purpose is always attended with difficulty, and that where the depth of top soil is very small, say six inches or less, the area of such lands which would be required for efficient purification would in certain cases be so great as to render land treatment impracticable.

The first Sewage Commission (1857-1865) reported that "the right way to dispose of town sewage is to apply it continuously to land, and it is only by such application that the

pollution of rivers can be avoided."

A further Commission appointed to inquire into the best means of preventing the pollution of rivers (1868-1874) reported with regard to chemical precipitation, intermittent filtration, and broad irrigation: (1) All these are to a great extent successful in removing polluting organic matter in suspension, but intermittent filtration is best, broad irrigation ranks next, and the chemical precipitation processes are less efficient. (2) For removing organic matters in solution the processes of downward intermittent filtration and broad irrigation are greatly superior to upward filtration and chemical processes.

Another Commission (1882-1884) appointed to inquire into the system under which London sewage was discharged into the Thames, whether any evil effects resulted therefrom, and if so, what remedial or preventive measures could be applied, found that evils did exist "imperatively demanding a prompt remedy," and that by chemical precipitation a certain part of the organic matter of the sewage would be removed. They reported, however, "that the liquid so separated would not be sufficiently free from noxious matters to allow of its being discharged at the present outfalls as a permanent measure. It would require further purification, and this, according to the present state of knowledge, can only be done effectually by its application to the land."

Experience has shown that with careful management a properly constructed sewage-farm can be carried on without nuisance or injury to the health of the surrounding population; but if the land becomes waterlogged from accident, defect, or inattention, grave nuisance may readily result. It was at first feared that parasitic and other diseases in animals and man would be promoted by the use of the produce of sewage-farms as food, but no such result has been observed.

Bacterial treatment. - Biological methods of sewage purification have of late been more thoroughly studied, with important results. The oxidation effected by the aërobic organisms has been increased (1) by supplying oxygen more freely by artificial means on a contact bed; that is, a specially prepared filter charged intermittently with sewage which is allowed to rest for a time upon it, and (2) by previously subjecting the sewage to anaërobic fermentation, the effect of which is to bring about more complete solution of the organic matter and render it readily oxidisable. For the second purpose the "septic tank" has become widely used. Lowcock blows air into the interstices of the filter by means of perforated pipes. Scott-Moncrieff passes sewage, after anaërobic treatment, through a vertical series of shallow filter-boxes, with intervening air-spaces; and finds that there is a differentiation of the microbes in the successive stages, so that if the sequence of the filters is changed the purification is for a time arrested, and the effluent becomes foul. Ducat's tank is filled with porous material, such as burnt ballast — finer below and coarser above; it is ventilated by field-pipes passing through at various levels, and the walls of the tank are for the most part built of open drain-pipes, placed transversely so as to admit air freely; it is claimed that the tank can be used continuously without rest, and that no further treatment is necessary. Dibdin employs a tank filled with coarse burnt ballast; the sewage is run in and allowed to stand for two hours, after which it is slowly run off—to filters, or to a second tank with finer materials—and the tank is allowed several hours' rest for aëration.

Bacterial purification depends upon two main groups of organisms, namely, those that are able to break down and liquefy solid organic matter, and those that deal with it when in solution; but no strict line of demarcation can be drawn between them. Different systems depend upon an anaërobic action (Cameron, Scott-Moncrieff, etc.) or aërobic (Dibdin, Fowler, etc.), or a mixture of the two. Whatever system is used, the two great agencies of breaking down and oxidation must be allowed ample opportunity. Two examples are furnished in Cameron's septic tank (anaërobic) at Exeter and many other places, and in the multiple contact bacteria-beds (aërobic) at the Davyhulme Works, Manchester.

For contact beds burnt clay, coke, clinker, cinders, or various forms of gravel may all be efficient media, provided there is ample aëration and porosity. The necessary organisms exist in the sewage, but require abundant oxygen. To assist in maintaining aëration the surface should be raked over from time to time. Obstruction by frost rarely occurs, the temperature of the sewage being from December to April about 55.5° F. Frequent and prolonged periods of rest are necessary; without them the beds become clogged, and eventually inactive because lacking in aërobic bacteria. The sewage applied to the bed should be as far as possible uniform in consistence and freed from suspended matters by sedimentation. Any suspended matter not so removed should be retained as far as possible on the surface of the bed.

Fowler concludes that the bacterial process is best conducted in three stages: -- (a) settlement and screening out of the grosser solids; (b) anaërobic decomposition in the septic tank; and (c) oxidation on bacteria beds. He summarises the conditions of the successful working of contact beds as the result of the Manchester experience as follows:-

(1) The bed should be worked very slowly at first, in order to allow it to settle down and the bacterial growths to form. In this way there will be less danger of suspended matter finding its way into the body of the bed, while the material is still loose and open. (2) The burden should not be increased till analysis reveals the presence of surplus oxygen, either dissolved or in the form of nitrates in the effluent. (3) Analyses of the air in the bed may usefully be made from time to time during resting periods. (4) The variations in capacity should be carefully recorded. If the capacity is found to be rapidly decreasing, a period of rest should be allowed. (5) Long periods of rest should be avoided during winter, as when deprived of the heat of the sewage the activity of the organisms decreases. If necessary, the burden on the bed should then be decreased by reducing the number of fillings per day, rather than by giving a long rest at one time. (6) The insoluble suspended matter should be retained on the surface by covering the latter with a layer of finer material not more than three inches in depth. The suspended matter thus arrested should not be raked into the bed, but when its amount becomes excessive it should be scraped off. This should be done if possible in dry, warm weather, after the bed has rested some days. By placing the inlet and outlet penstocks as close together as possible, the suspended matter will tend to concentrate in their vicinity, and its removal will be facilitated.

It should be noted that if sewage contains pathogenic bacteria, and is then treated by bacterial methods, the effluent cannot certainly be assumed to be safer in this respect than the raw sewage diluted, and, therefore, effluents should not be passed into drinking-water streams, but on to the land.

Selection of method.—The choice among the various methods of purification of sewage must be

determined largely by local considerations. We have already quoted the findings of the Sewage Commission of 1868 (see page 270). If the sewage is free from manufacturing effluents of injurious character, and if a sufficient area of suitable land can be obtained in a convenient position at a reasonable cost, broad irrigation is entitled to preference as the most rational and economical method of treatment. If there is serious difficulty in obtaining suitable land, simple intermittent downward filtration, or the bacterial treatment, may be preferable; and precipitation processes and artificial filters for still smaller areas may suffice.

Proper construction in the first instance, and careful management afterwards, are essential in every process, and without them nuisance cannot fail to result. It is desirable, for the sake of economy and efficiency in management and supervision, that sewage should be dealt with at as few points as possible. The process of purification is greatly facilitated by the "separate" system of drainage, the volume of sewage being thereby reduced and rendered more constant.

In manufacturing districts the principal difficulty arises in connection with trade effluents. Many kinds of manufacturing refuse if admitted into the sewers interfere with the efficacy of land-filtration, either by clogging the soil or chemically checking the process of oxidation. The latter condition also operates prejudicially in bacterial methods. In such cases a preliminary purification by precipitation is necessary, and it is probable that the effluents of different trades will be found to require different chemical treatment. If they are poured into streams without purification, they cause pollution exceeding in intensity that caused by domestic sewage. For each manufacturer in such a district to deal with his own refuse would entail a very heavy aggregate expense;

and the multiplication of such works is in itself undesirable, since it must increase the risks of mismanagement and failure. Still, where trade effluents cannot be received into the sewers, the responsibility for their purification rests with those who produce them. In some cases the difficulty may be overcome by combination among manufacturers, and in others by the provision of a special intercepting sewer. A satisfactory compromise has been reached in some towns by the manufacturers partially purifying the effluents before turning them into the sewers.

The Rivers Pollution Commissioners in 1868 proposed that impurities exceeding any of the following standards should be held to render an

effluent unfit for discharge into a river :-

Impurity.	Parts per 100,000 in excess of	Other standards.
In suspension— Organic matter. Mineral matter in suspension. Oily matter in suspension In solution—	1 3 0.05	Perfect rest in subsidence tanks for at least six hours also required. Or any film of oil on the surface.
Organic matter:— Organic carbon Organic nitrogen Metals (other than K, Na, Ca, Mg.) Arsenic Chlorine (free, after addition of H ₂ SO ₄)	2 0·3 2 0·05	
Sulphur as sulphides . Acidity (as HCl)	1 2 1	Any distinct colour in depth of 1 inch when examined by daylight in a white vessel.

The following have been suggested as general conditions which an effluent ought to fulfil: (a) It must contain practically no solids in suspension; (b) it must not contain in solution a quantity of organic matter sufficient to absorb much oxygen from the stream into which it is discharged; (c) it must not be liable to putrefaction or secondary decomposition; (d) it must contain nothing inimical to microbial growth and activity, therefore it must not be treated with strong antiseptics; and (e) it must not contain pathogenic organisms.

CHAPTER X

DISPOSAL OF THE DEAD

The daily average of deaths in England and Wales is about 1,500. The disposal of this enormous number of dead (many of whom have died of infectious diseases) in such manner as to avoid danger to the living is a sanitary problem which, like many others, has only received due attention within comparatively recent years. The horrible overcrowding of graveyards which prevailed in the early part of last century is now rarely met with, and is only possible in the older grounds which are not subject to modern regulations, and which have not been formally closed by Order in Council. Seymour Haden has shown that if carcases are covered by a foot of suitable earth, the perishable parts disappear inoffensively within twelve months, and speaking broadly it may be said that for every foot of depth below the surface of the soil one year or thereabouts is necessary for resolution. The present practice, however, and even the law, are inconsistent with burial sufficiently shallow to permit of the due action of the nitrifying organisms found only in the upper layers of the soil. The objects of earth-burial are further frustrated by the still prevalent use of metallic or heavy wooden coffins.

Burial-grounds.—The objects to be aimed at are rapid resolution and complete oxidation or absorption of the products. Hence the soil should be light, open, porous, and either naturally or artificially drained to a depth of not less than eight feet, so that

air and moisture may pass freely but in a finely divided state. Loam or sandy mould is about the best soil; clay is difficult to drain, retards decomposition by excluding air and moisture, and either retains the products of decomposition or allows them to escape through fissures. A loose stony soil allows gases to escape too freely. In cases of difficulty, as regards soil or drainage, the level may be raised artificially. It is desirable that burialgrounds should not closely adjoin dwellings.* While convenient of access, they should, therefore, be placed outside the limits of present or probable future dense building, and a margin of twenty feet or more in width should be reserved for walks and planting. The surface should be grassed. Cemeteries should not be placed on elevated ground whence the natural drainage may find its way to dwellings below, or may contaminate any water supply. For obvious reasons, lands liable to floods or encroachment by streams or by the sea are unsuitable. For Model Bye-laws see p. 599. The ordinary grave space being nine feet by four, it is usually estimated that an acre of ground is a minimum allowance for a population of 1,000 persons for fifty years. The four chief sanitary considerations therefore to be held in view in the provision of burial grounds are (a) suitable soil and proper elevation of site, (b) a suitable position, especially with respect to houses and sources of water supply, (c) sufficient space, and (d) proper regulation and management. (L.G.B. Memorandum, 1906.)

There can be no question of the injurious effects of overcrowded burial-grounds upon the health of the

^{*}The Public Health (Interments) Act, 1879, prohibited the construction of a cemetery within 200 yards (reduced to 100 by the Burials Act, 1906) of any dwelling without the consent of the owner and occupier. There is no restriction if such consent is obtained, nor any prohibition of future building nearer to the cemetery.

immediately surrounding population, but there is no clear evidence of harm accruing under the conditions enforced in modern well-regulated cemeteries. The dangers to be apprehended are contamination of air and water, to which, perhaps, ought to be added the possible retention or even multiplication of the germs of specific disease, many varieties of which can live in the soil. Effluvia may rise to the surface, or pass laterally for unknown distances, possibly to the basements of houses; but this is not likely to happen if the conditions already specified are observed. The drainage, artificial or natural, must be so arranged that it cannot find access to any supply of drinking-water, be it stream, well, or other source. Investigations by Buchanan Young show that organic matter and bacteria are increased in the deeper layers of soil, but not in the upper reaches. In vaults and over graves with an insufficient covering of earth fætid organic matter, ammonia, ammonium sulphide, and sulphuretted hydrogen have been found.

Persons taking part in exhumations have in many instances suffered from febrile attacks of varying character and intensity; but as a rule nothing more than diarrhea, and occasionally vomiting, results from such exposure, and that only in a minority of cases. Much depends upon the completeness of the putrefactive changes. Grave-diggers do not appear

to be an unhealthy or short-lived class.

Brick graves, vaults, catacombs, heavy oak coffins, and (worst of all) lead coffins, are all objectionable, and represent a futile attempt to prevent decomposition. They only retard the process, and render it far more likely to injure the living than if the pent-up gases and fluids were exposed to the deodorising and oxidising influence of aërated earth. Perishable coffins of wicker, light wood, or papier mâché, are most suitable of all.

"The degree to which the purity of neighbouring wells is endangered by a cemetery, and the distance to which contamination may extend, obviously depend in each particular case upon the relative elevation of the respective sites of cemetery and well, and upon the nature and dip of the intervening strata, so that it would seem impossible to lay down a general rule for all cases. Fissured rocks might allow foul matters to traverse considerable distances, while the interposition of a bed of clay or a water-tight fault would shut them off, or the passage through an aerated stratum of finely divided earth would oxidise and destroy them on their way. A dangerous state of things occurs when graves and wells are sunk near together in a shallow superficial water-bearing stratum of loosely porous nature resting on impervious clay. From experiments made at Dresden by Professor Fleck would seem that the degree to which wells so situated are liable to pollution is greater when the surface of the subsequent clay bed is horizontal than when it is sloping, even though the slope be towards the well. In the latter case the ground-water is on the move; in the former it is stagnant, and hence the foul matters are concentrated in a smaller volume of water; just as the water of a stagnant pool is more liable to become foul than that of a running stream. It does not appear, however, that the risk to which wells are exposed from the proximity of a properly managed cemetery is in ordinary cases great. A leaky cesspool is a far greater source of danger than a grave. solid and liquid excretions voided by a human being in the course of a year amount to several times the weight of his body.

"The State Board of Health of Massachusetts . . . give a series of analyses of water from wells in the neighbourhood of cemeteries. Of seven wells in sandy and gravelly soil, varying in depth from 4 to 17 feet, situated at distances respectively of 60, 50, 10, 100, 200, 75, and 100 feet from the nearest grave, and having no other sources of contamination at hand, one only showed undoubted evidence of contamination; this was 10 feet from the nearest grave, the most recent interment, made five and a half months before, being 35 feet distant; the three purest wells were those at 60, 50, and 75 feet distance. The chemical characters by which it may be inferred that the contamination of a particular water is derived from decomposing bodies rather than from sewage, are a high proportion of nitrogenous organic matter and ammonia, or if oxidation have proceeded further, of nitrates and nitrites, relatively to the amount of chlorides present, and also the presence in notable quantity of phosphates." (L.G.B. Memorandum, 1880.)

The chief precautions to be taken to avoid pollution of wells and springs by burial grounds are (a) "the intervention of a sufficient space between the burial ground and the water supply, (b) proper drainage where necessary in order that the subsoil water of the burial ground shall be conveyed away, and (c) proper management of the burial ground, so that the amount of organic matter in one place shall not be more than the soil can dispose of." (L.G.B. Memorandum, 1906.)

Both urban and rural authorities are enabled by the P.H. (Interments) Act of 1879 to provide cemeteries for their districts, and must do so if required by the L.G.B. The cemetery need not be within the district of the S.A. The L.G.B. point out, in a memorandum issued in 1879, that it is incumbent upon the S.A. to take action-

1. Where in any burial-ground that remains in use there is not proper space for burial, and no other

suitable burial-ground has been provided.

2. Where the continuance in use of any burialground (notwithstanding there may be such space) is, by reason of its situation in relation to the water supply of the locality, or by reason of any circumstances whatsoever, injurious to the public health.

3. Where, for the protection of the public health, it is expedient to discontinue burials in a particular

town, village, or place, or within certain limits.

The necessity may also arise from unsuitability of the site or of the subsoil, or from inconvenience of

access from populous parts of the district.

If it is desirable, upon the above or other grounds, to close any existing burial-place, a representation must be made to the Home Secretary for the purpose of obtaining an Order in Council to that effect, under the provisions of the Burial Act of 1853.

In excavations for new buildings in cities not infrequently skeletons and coffins are met with, which it is necessary to remove. For removal from one consecrated ground to another a faculty from the

Bishop of the diocese may suffice, but otherwise application must be made to the Home Secretary, from whom "a licence for the removal of human remains," under the Burial Act, 1857 (s. 25), may be obtained. The remains are then suitably re-coffined in boxes and re-interred in a burial ground where burials may legally take place.

Interments underneath or within the walls of any church built after 1848 are forbidden by the P.H. Act of 1848 (s. 83; re-enacted by the P.H. Act, 1875). No buildings must be erected upon any disused burial-ground, except for the purpose of enlarging a place of worship (Disused Burial Grounds Act, 1884).

Cremation.-In a crematorium of modern construction a body of average weight is reduced to about 3 lbs. of inorganic ash within two hours. The usual fuel is coal and coke, but Fradet recommends gas. In either case, a ventilating shaft with pilot fire at its base is necessary. The chief objections to cremation are that the soil is deprived of the organic matter that would otherwise be returned to it, that it involves an unnecessary waste of the world's limited stock of combined nitrogen, and that the impossibility of exhumation would increase the facilities for concealing homicide. The first objection has no great weight at present, since little attempt is made to utilise burialgrounds by cultivation. The last is, however, more serious, and cannot be regarded as satisfactorily met by the proposition of minute and detailed autopsy in every case. Half a million examinations annually, each sufficiently detailed to cover every known poison and every possible wound, would be impracticable, especially as the result must, in the vast majority of cases, be negative. The discovery of organic disease would not by any means necessarily exclude the possibility of foul play. Some few poisons, such as copper, might be detected in the ashes, but organic and volatile mineral poisons would be dissipated by cremation. It is true that under present conditions occasion for exhumation rarely arises, but the possibility of it may to some extent act as a check upon crime. The Cremation Act, 1902, empowers Burial Authorities (Burial Boards or other local authorities maintaining cemeteries) to provide and maintain crematoria. The plans and site must be approved by the L.G.B., and the completion must be notified to the Home Secretary before any cremation takes place. No crematorium may be built within 50 yards of any public highway, within 200 yards of a dwelling-house (except by consent of owner), or within the consecrated part of a burial-ground.

Home Office Regulations of 1903 impose conditions as to maintenance, inspection (by Home Office and L.G.B.), disposal of ashes, and the further points stated below. Cremation is not allowed if the deceased has left written directions to the contrary, or if the body has not been identified. There must be (a) medical certificates as to the cause of death from two registered practitioners (one being the medical attendant and the other a practitioner of at least five years' standing who (1) is specially appointed by the Authority for the purpose, or (2) is M.O.H., Certifying Surgeon under the Factory Act, or Medical Referee under the Workmen's Compensation Act, or (3) is physician or surgeon to a public general hospital with at least 50 beds), or (b) a certificate given after post-mortem examination by a skilled pathologist, or (c) a certificate given by a coroner after inquest. In certain circumstances (b) and (c) become essential.

A Medical Referee must be appointed by the Authority, and a deputy to act for him in his absence or in cases where he is the medical attendant. His certificate that all the conditions have been fulfilled is necessary before any cremation can take place, but certain relaxation can be allowed by him in exceptional cases (persons dying of plague, yellow fever, or cholera, on ship or in an isolation hospital; still-born children; bodies exhumed after burial for more than a year). Other modifications can be made temporarily by Order of the Home Secretary on application of the S.A., during an epidemic or for other sufficient reason. In 1906 the number of cremations

in the United Kingdom was 743.

CHAPTER XI

ANIMAL PARASITES

In addition to the pathogenic bacteria, belonging rather to the vegetable than the animal kingdom, the human body is liable to invasion by a number of parasitic animal organisms, among which the following

are the most important.

Protozoa are unicellular animals, little more highly developed than some of the bacteria. They are usually divided into four classes: Rhizopods, Sporozoa, Flagellates, and Infusoria. The last named group, which includes paramecium, need not be further considered. It is only recently that the relation between these protozoa and human disease

has been demonstrated with any clearness.

The Rhizopods are the lowest group, and comprise the amæbæ which move by pseudopodia and reproduce themselves by simple division. Many kinds of amæba occur in the human intestine, some being genuine rhizopods in vegetative stage, others being amæboid stages of higher forms. Two of the forms (Gymnamæbæ) are human parasites, namely, the amæba coli (or as it is termed by Schaudin, entamæba) and the amæba dysenteriæ (entamæba histolytica). The former appears in the excreta of 20 –50 per cent. of healthy persons, though it is not easy of detection. The latter is possibly the cause, wholly or partly, of dysentery, as commonly occurring in the tropics.

The Sporozoa are so called because at one stage of their life history the individual breaks up into a number

285

of minute reproductive bodies resembling the spores of fungi, often encysted. This group includes the gregarines which inhabit the digestive tract of crustaceans, the coccidia which are intestinal parasites in birds and in animals, and the hæmoprotozoa, or parasites of blood corpuscles, classified by Sambon into four groups. The life history comprises alternation of generations and change of host. Once the sexually-produced reproductive body or spore finds its way into a suitable host it attains its full size and then splits up into individuals which carry on the cycle.

Chap. XI.]

Coccidium oviforme produces coccidiosis or psorospermosis in rabbits and other animals. Pyrosoma bigeminum, the cause of Texas fever in cattle, which is transmitted by a tick, is a closely allied organism.

The Flagellates include the *Trypanosoma*, and the *Trypanoplasma*. The latter occur in fishes only. The former cause various diseases of horses, bovines, rodents, etc. They have now been shown to be the cause of certain diseases of man, such as sleeping sickness, and some are conveyed by a fly, *Glossina palpalis*. A trypanosome has a long spermatozoon-like body fringed on one side with an undulating membrane prolonged at the anterior end into a flagellum. The body possesses a large nucleus. The organism passes through various stages in its life cycle, and a connection has recently been established between the malaria parasite and the trypanosome. The majority of trypanosomes and other hæmoprotozoa have a life cycle consisting of alternating gut parasitism and blood parasitism.

The next group of animal parasites infesting man are the Insecta and the Arachnida. Among the former are fleas, bugs, lice, the *Pulicidæ*, and the larvæ of certain flies; among the latter are the mites, ticks, and *Pentastomata*. More important are the parasitic

worms, which may be divided broadly into two groups as follows:—

- 1. The flat worms (Platyhelminthes), which are usually sub-divided into the Cestoda or tapeworms and the Trematoda or flukes.
- 2. The round or thread worms (Nemathelminthes) or Nematoda. This group includes the common thread worm (Oxyuris vermicularis), the common round worm (Ascaris lumbricoides), and the various forms of filariæ and ankylostoma.

Tæniada.—It has been said that as many as a dozen forms of tapeworm occur in man. This species of Cestoda passes through two distinct phases, in two different hosts. In one, the head, or scolex, together with a cystic expansion, is embedded in muscle and other solid tissues; in the other, the strobilus or tapeworm occupies the alimentary canal of some other animal. If the flesh containing tænia cysts (cysticerci) is eaten by another animal, the scolex reaches the alimentary canal of the new host, loses its cyst, and attaches itself to the wall of the intestine. It then develops from the caudal end joint after joint of proglottides, square or oblong flat segments, each of which is provided with double sexual organs. The chain of proglottides may attain a length of many feet. Ova are produced in the segments, and escape with the excreta of the host if the segments rupture or become detached, or if the whole of the tapeworm is expelled. Some of the ova are swallowed by a herbivorous host, and the embryo then bursts its shell, makes its way from the intestine to the solid tissues of the host, and after developing anew into a cysticercus, remains passive until it is devoured by a carnivorous animal, or perishes.

Man is subject to tapeworms, and more rarely to their cysticerci. *Tænia solium*, one of the commonest of human tapeworms, attains a length of seven to ten

287

feet or more, consisting sometimes of 700 to 800 proglottides. The head is about the size of the head of a pin, and bears four suctorial discs, and a double circle of hooklets surrounding a prominence or rostellum. It attains its full development in three or four months, but may remain in the intestine for years. The ovum is spherical, $\frac{1}{700}$ inch in diameter, and has a thick brown shell; the embryo has six hooks. The cysticercus or bladder worm of the Tænia solium is termed Cysticercus cellulosæ. It affects the pig, and more rarely man; in the latter it lodges in the muscles, connective tissues, brain, eye, and serous membranes, and attains its full development in about 2½ months. The cyst reaches the size of a pea, or even a marble; the head or scolex has a double crown of hooklets. In pigs it constitutes the affection known as "measles," and the consumption of uncooked "measly pork" is the chief source of Tania solium in man.

Tænia mediocanellata is more common than Tænia solium, and somewhat resembles it, but its length is greater (20 ft.). The head is about $\frac{1}{1.5}$ inch in diameter, and has four suckers, but no rostellum or hooklets. The eggs are oval, the short diameter being about $\frac{1}{850}$ inch. The Cysticercus tæniæ mediocanellatæ, or Cysticercus bovis, is an oval vesicle smaller than the Cysticercus cellulosæ. It consists of a cyst, and a scolex identical with the head of the tænia itself. It occurs in the flesh of cattle and produces "beef measles."

Bothriocephalus latus.—This tænia attains a length of 24 feet or more. The head is ovoid, $\frac{1}{10}$ inch long, and has two longitudinal grooves or suckers, but no hooklets. The body is ribbon-like and may have as many as 3,000 segments. The eggs are oval, about $\frac{1}{370}$ inch shorter diameter, and open by a lid at one end. The cysticercus is found in pike, turbot, perch, and

other fish. It may, perhaps, gain access to the alimentary canal of the fish by means of water, as the embryo is ciliated, and is found in river water. The tænia is met with principally in the Baltic countries

(Russia, Poland, Sweden) and in Switzerland.

Tania echinococcus affects the dog and wolf only. The head resembles that of Tania solium in presenting a rostellum surrounded with a double row of hooklets and four suckers, but it is only $\frac{1}{100}$ inch in width. There are four segments only, and the whole adult tapeworm is about 2 inches long. The eggs are spherical and contained only in the terminal or reproductive segment. Its cysticercus is a dangerous human parasite affecting the liver and various other parts of the body under the name of hydatid, which occurs more frequently in Iceland and Australia than in Great Britain. Unlike other cysticerci it increases indefinitely in size, becoming sometimes a large cyst; it also forms within itself secondary cysts, some of which ("broodcapsules") contain one or more scolices (echinococci) and remain minute, while others, containing no scolex, enlarge and form "daughter cysts," which may in turn produce new cysts by gemmation. After death the scolex or echinococcus resembles the head of the tænia, but during life it is retracted into a depression, and thereby turned inside out. Hydatids may continue to grow for an indefinite number of years, and are spread among men and animals by affected dogs.

Trematoda or flukes furnish two human parasites, the Bilharzia hæmatobia (now termed Schistosomum hæmatobium) and the Distomum hepaticum.

Bilharzia hæmatobia is believed to be the cause of a form of intermittent hæmaturia endemic in Egypt, South Africa, and elsewhere. The parasite is about \(\frac{1}{4}\) inch long, and infests the veins of the large intestine,

bladder, ureter, and pelvis of the kidney. It causes small patches of inflammation along these tracts, and the urine frequently contains blood and the ova of Bilharzia. The ureter may be obstructed. Dysenteric symptoms occasionally occur owing to the presence of the parasite in the veins of the intestine. The ova are about 1 inch long, and have a sharp terminal spine, not quite in the longitudinal axis. The parasite probably gains access to the human body through water, either by drinking or by direct infection of the urinary and alimentary tract during bathing. The larva becomes encysted in the body of aquatic crustaceans. Other varieties of Schistosomes

The common liver-fluke or Distomum hepaticum, which produces "liver-rot" in sheep, is shaped like a diminutive sole and is about an inch in length. It may infect persons consuming affected livers, and may be found in the tissues, and particularly in the

portal system.

Nematoda.—The common round worm, or Ascaris lumbricoides, affects man and other animals. Its habitat in man is chiefly the small intestine. It is pinkish in colour, and tapers to each end. The male is about 6 inches long, the female 12. The ova, enormous numbers of which are discharged, are oval and nodulated, and measure about $\frac{1}{450}$ inch in least diameter. It is believed that the ova develop only in an intermediary host. The threadworm, Oxyuris vermicularis, is much smaller, the female measuring about half an inch in length, and the male about a quarter of an inch. They occur in enormous numbers, chiefly in the rectum. The ova are unsymmetrical but oval, their short diameter being 1 and long diameter 1 inch. The origin of threadworms is uncertain, but it is probable that they are occasionally disseminated by means of drinking-water.

The adult Ankylostoma, which is about half an inch long, inhabits the upper part of the jejunum in man, A. duodenale being found in Europe and the tropics, and A. americanum in the United States.* It possesses no intermediate host, and there is no sexual generation outside the host. The infection of each host (man, sheep, etc.) is specific for each parasite. The adult alone is infective, the eggs and young larva being non-infective. The fresh ova are characterised by a thin, clear, colourless shell of oval form, encasing greyish contents divided by segmentation, and eventually becoming a morula before the emergence of an active vermiform embryo (in 18 hours). From the fresh egg the encapsuled larva appears in 4 or 5 days or longer. The second period of growth within the host from the encapsuled larva to the adult occupies about 5 weeks. The encapsuled larva is capable of prolonged saprophytic existence under favourable conditions. The eggs are killed in about 24 hours at 40° C. or at zero, and below 13° C, they will not hatch. Moisture and free oxygen are necessary for development of ova after expulsion from the body.

Ankylostoma obtains access to man's intestine by the mouth and through the unbroken skin through the hair follicles as encapsuled larva (Loos). From the skin it is carried by the blood stream to the lungs, up the trachea and down the esophagus to the alimentary tract, in which the parasite attaches itself to the villi of the jejunum, abstracting blood and producing hemorrhages. The typical result of infection is anæmia, sometimes associated with dyspeptic conditions. Many persons harbouring the worm, however, suffer from no symptoms of disease at all. Dobson found eggs in the excreta of 75 per cent. of

^{*} This nematode has also been termed Sclerostoma duodenale, Dochmius duodenalis, and Strongylus duodenalis.

the natives of Assam; and Boycott found them in 100 per cent. of workers in certain Cornish mines, although not more than 10 per cent. of the men were actually ill. The healthy worm-carrier is probably one of the chief factors in dissemination, and may continue to discharge ova for years. The disease has long been known under such names as "Egyptian chlorosis," tropical anæmia, or miners' anæmia, but only recently fully described by Haldane as "Dolcoath anæmia." It is now termed ankylostomiasis, and is recognised as an endemic disease of warm climates, readily introduced into temperate climates by dissemination of the worm. The effect of temperature operates, however, in cold climates. For instance, Tenholt has shown that in Westphalian mines where the temperature is below 17° C. the cases of ankylostomiasis (anæmia) per 1,000 men employed underground was only 0.6, whereas where the temperature was 20-22° C. it was 2.5, when it was 22-25° C. 11.7, and when above 25° C. it was 39.9 per 1,000. Treatment consists in purgation (calomel), followed by three successive doses of 30 grains of thymol. In Westphalia ethereal extract of male fern is used. Preventive measures are based upon personal precautions against ingestion or skin infection, and of separation of the diseased from healthy persons.

Trichina spiralis attacks rodents, pigs, and many other animals besides man, causing the disease known as trichinosis. It is found in the tissues, and especially in the muscles, in the form of ovoid cysts about $\frac{1}{70}$ inch in length, just visible to the naked eye. Within each cyst is coiled an immature trichina, about $\frac{1}{35}$ inch long. The encapsuled trichinæ may retain their vitality for many years, or may perish and become calcified. If the tissues in which they lodge are eaten, the capsule is dissolved by the gastric juice, and the liberated trichinæ rapidly develop, the male

attaining in one or two days a length of $\frac{1}{18}$ inch, the female $\frac{1}{8}$ inch. Ova are formed and impregnated, and are hatched within the uterus in about a week. The embryos escape, burrow into the walls of the intestine and find their way into the tissues of all parts of the body, where they become encapsuled and form the cysts already described, within a month from the ingestion of the trichinous food.

Trichinosis in man is generally due to the consumption of the imperfectly cooked flesh of a pig suffering from the disease. In Germany, where ham, pork, and sausage are often eaten almost or quite raw, trichinosis is far more common than it is in England. The symptoms commence within a week, with nausea, abdominal pain, irregularity of the bowels, prostration, rapid pulse, and elevation of temperature. The malady usually increases in severity for one or two weeks, and then gradually subsides, but fatal enteritis, peritonitis, or pneumonia may supervene, or the patient may die of exhaustion at any time from the end of the first to the end of the sixth week. Characteristic symptoms attend the invasion of the tissues by the parasites. The muscles become painful, tender, swollen, and stiff, so that the limbs are flexed and motionless. The voice is often hoarse or aphonic, from implication of the larynx. The pain in the limbs differs from that of rheumatism in not affecting the joints, but only the muscles. Œdema occurs early, first in the eyelids and face, next in the hands and feet, and may become general and involve the serous cavities. The other symptoms include increasing prostration and copious perspirations; the temperature may be little above the normal, and if high is irregular and subject to morning remissions. The mortality is sometimes very slight, sometimes 20 or 25 per cent.

Various forms of Filaria may infest man, par-

ticularly in the tropics, setting up a variety of pathological conditions, such as abscess, lymphangitis, chyluria, elephantiasis, etc. The species is rarely seen in the adult form, is a hair-like parasite, and may attain a length of three or four inches. An immature form, met with in cases of chyluria, is about $\frac{1}{7.5}$ inch long. The ova are about $\frac{1}{1.000}$ inch long, and having no distinct shell their shape is somewhat irregular. The parasite is found chiefly in the East and West Indies, but occasionally in England, even among persons who have never left the country. It abounds in the blood of affected persons during the night (Filaria nocturna), but disappears from the circulation by day; by changing the habits of the patient this periodicity can be reversed. Its presence is commonly associated with one of two diseases-chyluria and elephantiasis—due to the obstruction of the lymphatics, etc., by the parasite. The former is characterised by periodic attacks, in which the urine becomes milky and upon standing coagulates, but the coagulum soon breaks down and decomposition sets in rapidly. These phenomena have been traced to admixture of lymph with urine, and immature filariæ are visible under the microscope. Elephantiasis is attended with enormous enlargement of the greater or smaller part of the trunk or limbs, and especially the legs and generative organs (E. arabum).

Filaria medinensis vel dracunculus, or Guinea worm, is about $\frac{1}{10}$ inch in diameter, and usually one to three feet long. It is endemic in certain tropical regions. The embryos affect minute aquatic crustacea (Cyclops), in which they undergo larval development. Later they gain access to the human tissues, but whether by means of drinking-water or infection through the skin during bathing is not clear. They burrow in the tissues, especially of the legs, causing large boils and sores.

CHAPTER XII.

INFECTION.

CERTAIN diseases, the number of which is steadily increasing owing to improved methods of observation, are termed specific or infective, and are regarded as due to the invasion of the system from without by a definite "ferment" or poison, which grows and multiplies in the body. In some of them the poison is given off again, and these diseases are therefore infectious, or transmissible from person to person.

Modes of infection.—It cannot be said that the possible means by which infection is given off have been exhaustively determined in regard to any disease, but certain well-established modes may be

mentioned, namely :-

- (1) By the *breath*, in small-pox, measles, whooping cough, pneumonia, mumps, typhus, scarlet-fever, and probably the great majority of infectious fevers. Little is known of the distance to which infection can be carried through the air; long ranges seem to be possible in small-pox, and may be suspected in measles and whooping cough; typhus and plague only infect at short distances, and this is probably true of scarlet-fever.
- (2) By exhalations, from the skin in typhus and possibly other diseases, from wounds in pyæmia, erysipelas, and other septic diseases.

(3) By desquamated particles of epidermis in scarlet-fever, and pustules in small-pox. These may

be carried by currents of air.

(4) By secretions and excretions. Mucus from the mouth, throat, etc., is infective in diphtheria and

scarlet-fever; sputa in tubercular phthisis; saliva in rabies in dogs if not in man. Milk is liable to carry infection (Chapter IV.). The bowel excreta are infective in enteric fever and cholera. The urine is also a vehicle of infection in some acute specific fevers, as, for instance, in enteric fever. Syphilis and gonorrhea are transmitted by means of the

specific discharges.

As regards the recipient the mode of infection is also varied. Many diseases are conveyed through the air, and the virus is doubtless inhaled, lodging either on the fauces or in the lungs. Water is known to be able to convey the poison of enteric fever and cholera, and milk that of scarlet fever, enteric fever, and diphtheria, the poison in such cases being swallowed. It is probable that the specific organisms multiply in these media. Experiments seem to show that the acid gastric juice kills microbes that are not in the spore stage, but since the reaction is not acid in the absence of solid food, this safeguard is incomplete, especially as regards water-borne virus. Infective particles carried in fomites, i.e. in clothing, etc., or conveyed by the hand to the mouth, may be inhaled or swallowed. Mosquitoes carry the malarial poison and that of yellow fever, and flies convey infection, e.g. of anthrax, plague, enteric fever, and cholera, to a raw surface, or to milk or other food. An attempt is sometimes made to distinguish as "contagious" those diseases which are not usually transmitted without direct personal contact; but this condition is not absolute, and is favourable to the transmission of any infectious malady. Puerperal fever may be regarded as an example of infection by contact or inoculation. A few diseases, of which rabies and vaccinia are typical examples, can only be conveyed by inoculation, that is, by lodgment of the virus in an abrasion of the skin; this is the ordinary mode of infection by anthrax and glanders, although inhalation is also possible. Small-pox, tuberculosis, and certain other diseases are also inoculable. Inoculation may occasionally take place through unbroken skin. This has been shown to be possible as regards virulent cultures of the glanders bacillus if rubbed in. Ankylostoma is believed to gain access through the hair follicles (page 290).

In leprosy the mode of infection is still unknown, although the microbe is now familiar. Even less has been made out in regard to rheumatic fever, the specific nature of which, though probable, still remains

open to doubt.

De novo hypothesis. - Some authorities believe that certain specific diseases occasionally arise de novo, independently of infection from any previous case. This has been alleged more especially as regards relapsing fever, typhus, enteric fever, diphtheria, erysipelas, puerperal fever, hospital gangrene, and septicæmia. The closest examination often fails to reveal any exposure to infection, and occasionally, under favourable conditions, these diseases spring up in localities that have for years been entirely free from them. The tendency of modern research, however, is to find an adequate explanation in infection conveyed by fomites, by food, by insects, by air-borne micro-organisms, by lower animals suffering from the same disease, possibly with widely different manifestations, or by aberrant forms or unobserved attacks in man. The supposed logical necessity for a "previous case," that is, of an immediately antecedent human case, has lost some of its significance since we have learned to recognise the actual living materies morbi, to detect it in air, water, and soil, to cultivate it in artificial media for indefinite periods, to transmit it to lower animals, and to increase or lessen its virulence at will. It is noteworthy that most of the infective diseases suspected of origin de novo are filth-diseases,

and due to microbes that can be cultivated outside the living body. Further, it has been found that there are non-pathogenic microbes of wide distribution, which in every point except virulence closely resemble the specific microbes of enteric fever and diphtheria; and non-pathogenic bacilli, morphologically similar to anthrax bacilli, have been found to be protective against anthrax. Some authorities regard these harmless microbes as being probably generically identical with the virulent microbes that they resemble, and therefore capable of assuming a virulent character if cultivated under suitable conditions, a view that reintroduces the de novo hypothesis in a new form.

Endemics, epidemics, pandemics.-Many of the specific diseases, and especially those that have relation to telluric conditions, attach themselves more or less permanently to certain localities, and are termed endemic; thus cholera is endemic in the delta of the Ganges, leprosy in parts of Norway small-pox in the Soudan and elsewhere, and in a less marked degree diphtheria, enteric, and scarlet-fever may be said to be endemic in certain districts in England. From time to time most diseases of the specific class become widely prevalent over a larger or smaller area, and are said to be epidemic. Thus cholera in an epidemic form occasionally spreads westward over Europe, disappearing almost entirely in the intervals. Small-pox, scarlet-fever, measles, and other diseases, which are always present in England, assume an epidemic form every few years, locally or in widespread outbreaks. Occasionally a disease diffuses itself so generally over a great part of the globe as to constitute a pandemic.

The causes of such outbursts are still imperfectly known. They have been attributed vaguely to "epidemic constitution" of the air, to "pandemic waves" of unknown nature, and, more intelligibly, to climatic

and meteorological conditions, accumulation of susceptible persons, facilities for convection or transmission, and (as regards cholera and other "filth-diseases") imperfect sanitation. Scarlet-fever, in temperate climates where it has established itself permanently, tends to become epidemic at intervals of about five years; measles at intervals of about two years. Whooping cough is more irregular than either, but on an average becomes prevalent every second year. Diphtheria shows no very marked periodicity, apart from its dependence upon season, and enteric fever none.

It will be seen presently that the liability to each kind of infection varies in man according to age and sex, as well as locality, climate, season, and surroundings. Similar variation occurs among the different races of men, and still more conspicuously among the different genera and species constituting the animal

kingdom.

The more important specific diseases that are common to man and certain lower animals are vaccinia, tuberculosis, anthrax, rabies, tetanus, glanders, actinomycosis, and septicæmia. Influenza ought probably to be added to the list. There is some reason to believe that diphtheria and scarlet-fever affect the cow, and the same may be said of enteric fever. Variolous diseases attack sheep and other animals, but are probably not identical with small-pox.

period intervenes between infection and the appearance of the first symptoms of the disease. It is fairly constant for each disease; but shorter and more uniform when infection is due to inoculation or ingestion, and in rabies is dependent further upon the

position of the bite.

Little is accurately known of the changes that occur during incubation, beyond the fact that the

299

poison is multiplying in some part of the system.* The onset of symptoms is very gradual in enteric fever and a few others, but sudden in the majority. The temperature invariably rises, and the characteristic phenomena of each malady follow with more or less intensity. A rash makes its appearance upon the skin in certain of the infectious diseases, hence termed exanthemata; the date of its appearance is constant for each disease within narrow limits, and its distribution and other characteristics are also more or less uniform. The chief exanthemata affecting man are small-pox, varicella, scarlet-fever, measles, rötheln, enteric fever, and typhus. In exceptional cases, the rash, like any other given symptom, may be wanting.

Course of disease .- As a rule, the specific diseases have a short and definite course, ending in death, or in recovery with or without permanent changes in tissues and organs. Some, however, like syphilis and leprosy, run a life-long course, while malarial diseases tend to recur at regular intervals, or after years of apparently complete recovery. Rabies and some other diseases are almost invariably rapidly fatal, while rötheln, mumps, and varicella are attended with such slight constitutional disturbance that deaths are very rare. The tendency to death may be dependent upon constitutional symptoms such as hyperpyrexia, on local lesions as in enteric fever and diphtheria, on complications in the course of the disease (often really part of the disease itself), or, lastly, on sequelæ affecting important organs.

The intensity of any given disease is subject to

^{*} Concurrence of two distinct zymotic diseases is occasionally observed. Vaccinia and small-pox, measles and whooping cough, measles and scarletfever, measles and mumps, whooping cough and chicken-pox, are among the least rare examples. According to Squire, diseases with similar incubation (i.e. both long or both short) may run their course independently without interfering with each other, but if two diseases with dissimilar incubation are acquired at the same time, one or the other is delayed.

wide variations. Thus small-pox may be so virulent as to kill the patient before any rash appears, or so mild that only one or two spots, sometimes none, may appear. Scarlet-fever sometimes is malignant and rapidly fatal, in other cases so slight as to escape recognition. As a rule, the type, whether severe or the reverse, holds good for the majority of attacks in a given outbreak, or at all events in the same period of the outbreak; but it is commonly more intense in the earlier than in the later part of an epidemic. As regards purely seasonal variations, however, there is reason to believe that in scarlet-fever, and probably in other diseases also, increased prevalence is associated with diminished average severity, as measured by the proportion of deaths to attacks. Age and sex have an important influence upon severity of attack as well as upon susceptibility, varying with different diseases; thus scarlet-fever, whooping cough, and many others become less and less dangerous to life from infancy upwards, while in enteric fever the reverse is the case. Both scarlet-fever and smallpox are less liable to terminate fatally in females than in males, and the same seems to be true also of influenza.

characteristic of specific diseases is their tendency in the event of recovery to protect the patient against a subsequent attack of the same disease. This statement must be taken in a modified sense as regards syphilis, leprosy, and the malarial diseases. The protection is perhaps most complete in small-pox, but in this, as in all others of the class, second and even third attacks are occasionally met with. In enteric fever, diphtheria, and influenza the protection is much less marked, and in erysipelas, phthisis, and septicæmia it can scarcely be said to exist. In all cases it diminishes with the lapse of time, its

duration being dependent upon the nature of the disease, upon the amount of lesion (in vaccination at all events), and probably upon idiosyncrasy. Klein has shown that protection against anthrax, though real, is very brief in guinea-pigs, and much more lasting in cattle. Fatigue greatly increases the susceptibility of rats to anthrax. The presence of sugar in the tissues (due either to direct injection of sugar, or administration of phloridzin) lessens the resistance to glanders and to suppuration, but not to anthrax or tubercle. There are also examples of the viru-Hence of a pathogenic microbe being intensified by the presence of an otherwise harmless germ: thus attenuated cultures of the specific microbes of pneumonia or erysipelas, too feeble to give rise to any symptoms, can be made to cause an intense attack if along with them is injected a culture (sterilised (or not) of a harmless saprophyte, proteus vulgaris. It has been thought that an attack of enteric fever lessens or annuls the protective influence of vaccination. Little is known as to any protective power of one specific disease against another, unless vaccinia be regarded as absolutely distinct from small-pox. Erysipelas has been found experimentally to be protective against anthrax. Animals are inoculated with the "attenuated virus" of anthrax, fowl cholera, and other diseases, the object being to obtain protection by means of a mild form of the disease. Inoculation for small-pox, and the Pasteurian prophylaxis against rabies, are examples of similar procedure applied

The germ theory, broadly stated, affirms that certain specific diseases are invariably associated with the growth and multiplication in the system of corresponding specific microbes, and that these microbes are the actual contagia or causes of the disease. The evidence is partly direct, partly indirect.

1. In regard to a few diseases the proof is complete, since—(a) a certain microbe is invariably discovered in the blood or tissues of animals suffering from the disease in question; (b) this microbe can be cultivated in artificial media, for any required number of successive generations; (c) the same disease is reproduced by inoculation of a susceptible animal with the last cultivation; and (d) in every such inoculated animal the specific microbe is found, with the same distribution as in animals infected in the ordinary way.*

The third stage (c) being as a rule inadmissible as regards man, the complete proof is practically limited to diseases that attack some lower animal. Of those affecting man (as well as lower animals), anthrax and tuberculosis have been fully proved to be associated with specific microbes, and similar complete evidence is forthcoming in respect of many diseases

attacking lower animals only.

2. A characteristic microbe is constantly found in the blood or tissues of persons suffering from certain other diseases—e.g. relapsing fevers, enteric fever, diphtheria, leprosy, pneumonia, cholera, etc.—although the proof of causative relation is incomplete. In several diseases, however, which from analogy are nevertheless believed to be microbial, no specific or characteristic organism has been isolated with certainty. Measles and whooping cough may serve as examples.

3. There is a close analogy between the natural history of infection and that of organised living ferments such as yeast, in respect of the almost infinite multiplication of an almost infinitesimal charge planted in a suitable soil under suitable conditions,

^{*} To these four postulates, formulated by Koch, Martin has added a fifth, namely, that the secondary infective agent (toxin) separable from the tissues in the natural disease should have similar chemical and physiological action to the products obtained from a pure cultivation of the microbe.

and also in respect of the effect of reagents in checking

or permanently arresting their increase.

The germ theory is now generally accepted, even in respect of those specific diseases, such as rabies, in which no characteristic microbe has yet been isolated. The chief alternative view which has been advocated is that infection is due to an un-organised chemical ferment, comparable to diastase, the microbes being regarded as (at most) effects or collateral phenomena,

without causative influence as regards disease.

What the precise relation between the germs and the production of the varied phenomena of disease may be is less clear. Under certain conditions the enormous multiplication of the microbes may lead to obstruction of vessels—a "mechanical mycosis"—but the ordinary course of symptoms cannot be accounted for by any physical properties of such minute particles. They act mainly by modifying the physiological processes of the tissues-e.g. by depriving the red corpuscles of oxygen, or by forming some chemical pro duct (or toxin), as a result of their growth and multipli cation, which in its turn diffuses and acts as a poison.

Five principal hypotheses, in some degree overlapping each other, have been advanced to account for the protection conferred by an attack of such a

disease :-

1. The exhaustion or pabulum theory is that the microbes during the first attack remove some chemical substance necessary for their growth, the loss of this substance rendering further attack impossible. This view involves the improbable assumption that the human body contains a special and separate pabulum suited to each specific disease to which it is liable, since one disease is only protective against itself.

2. The antidote or retention theory assumes that the first attack leaves in the system some direct or indirect product of the growth of the microbes which inhibits any further multiplication. There is experimental evidence in support of this view, as will be seen presently, and sterilised or filtered cultivations of certain pathogenic microbes have been found (if injected into the system) to have a protective action, which, curiously enough, does not necessarily include any power of checking the growth of the specific microbes outside the living body, in artificial media. The effect therefore is not due to direct action on the microbes, but to increase of resistance.

3. According to the acclimatisation theory, the cells and tissues are in some way modified during an attack so as to be able to resist future invasions of the same microbe. The tissues, in short, have acquired a tolerance of the poison of a certain disease, and subsequently that infection does not produce the former

symptoms of disease.

4. The phagocyte theory, somewhat allied to the theory of acclimatisation, is based on Metchnikoff's observation that when virulent anthrax bacilli are inoculated into an insusceptible animal (such as a, frog), the white (amœboid) blood corpuscles absorb the bacilli bodily, and presumably destroy them. The same happens when attenuated virus is inoculated, even into susceptible animals; the bacilli may be seen partially or wholly embedded in the amæboid corpuscles. When a susceptible animal is inoculated with virulent anthrax, however, few if any bacilli can be seen inside the corpuscles. The inference to be drawn from Metchnikoff's observations is that the amœboid corpuscles act as "phagocytes," or destroyers of pathogenic microbes; and that "protection" is largely, if not entirely, dependent upon this action. The same hypotheses may serve to account for the termination of an attack. It may be assumed that all the pathogenic microbes perish or are eliminated at the end of an attack of most of the infectious diseases,

but pass into a quiescent stage in the intervals of malarial and relapsing fever. In tuberculosis, leprosy, and syphilis, their development may be progressive.

In 1903 Wright and Douglas found that no phagocytosis occurred if the leucocytes were separated from the blood plasma. They concluded, therefore, that the leucocyte by itself was impotent, and that the phagocytosis was dependent upon some substance in the blood plasma. Subsequently they showed that this substance did not act directly upon the leucocytes, but by combining with the micro-organisms prepared them, so to speak, for destruction by the leucocytes, hence this substance in the plasma is termed opsonin (opsono, I cater for). It follows that the amount of phagocytosis is an index of the quantity of opsonin, and does not directly represent the vital activity of the leucocyte. Opsonins are distinct from agglutinins and antitoxins, and possess apparently a high degree of specificity. In normal serum they are almost completely destroyed by heating for ten minutes at 60° C. The opsonic content of any blood is arrived at by comparing the amount of phagocytosis observed as a result of the activity of its serum with the amount of phagocytosis observed in the case of a normal serum used as a control. The figure so obtained is termed the "opsonic index" of the patient's blood; e.g. an opsonic index of 0.5 means that the blood contains one half the normal quantity of that specific opsonin.

5. Ehrlich's side chain theory.—Ehrlich looks upon a molecule of normal protoplasm as composed of a central atom cell with a large number of "sidechains" of atom groups. The central cell is the mother cell, the side chains are receptors, that is, cells having combining affinity with food-stuffs by which nutriment is brought to the mother cell. These receptors are of two kinds, those having power of combining with molecules of simple constitution, and those having power of breaking up compound bodies by ferment action for purposes of assimilation. Toxins are provisionally regarded by Ehrlich as possessing two kinds of elements, haptophorous (i.e. the atomic group responsible for union with side chains of cells) and toxophorous (i.e. the toxic atomic group).

When introduced or occurring in the system toxins are fixed to the receptors by the former, while the latter remain free, and if in sufficient number or quantity produce toxic changes. If the dose of toxin is small the mother cell throws off the receptor plus the toxin (R + T), which thus becomes free in the blood. The central atom group then produces new receptors, which in their turn are set free. Ultimately the toxin being exhausted there is over-regeneration of receptors and the excess of unfixed receptors become free in the blood, constituting antitoxin molecules. Thus, when forming part of the mother cell the receptors anchor the toxin, which sets up toxic effects, but when the receptors are free in the blood (R + T) we have an inert compound without toxic effect. This ingenious theory assumes that antitoxins are normally present in the body in varying degrees, but when stimulated by introduction of toxin become increased.

Bacteria and disease.—The terms microbe, micro-organism, bacterium, germ, are applied almost indifferently to an increasing number of microscopic organisms which are destitute of chlorophyll and consist of protoplasm enclosed in a cellulose membrane. They multiply by fission, and are therefore termed Schizomycetes, or fission fungi. The common basis of primary classification is founded upon morphology as—

Micrococci.—Minute spherical microbes, without cilia. They multiply by elongating and then dividing transversely, forming diplococci (i.e. pairs), streptococci (chains), staphylococci (clusters), sarcinæ (square groups of four cocci, or multiples of these), or zooglæa (irregular masses embedded in a gelatinous matrix).

Bacilli.—Rod-shaped, with rounded or square-cut ends; longer than they are broad. Some have flagella, others have not and are immotile. They elongate, and may form rods of any length, or divide transversely into separate short rods or chains of rods or zoogleea. Some bacilli form spores under

certain conditions, among which may be mentioned moisture, suitable temperature, and presence of oxygen. The spores are round or oval glistening bodies, which appear in the substance of the bacillus, and grow at the expense of the protoplasm until the sheath bursts and liberates them. They are extremely resistant to heat and cold, but in certain artificial media spores readily germinate, a projection appearing at one point and growing into a bacillus.

Spirilla, Spirochætæ, and Vibriones .-- Long, wavy, and motile (ciliated) filaments. Some of these are non-septate, others are sub-divided and may break into bacillary elements, as, for example, Koch's spirillum of cholera, which is known in

a bacillary form as the comma bacillus of cholera.

The higher bacteria include more highly organised members of the Schizomycetes which branch and reproduce themselves by terminal fructification in conidia. To this group belong leptothrix, cladothrix, and streptothrix. The last named seems likely to become the most important member of the group from a pathogenic point of view. The exact biology of these higher bacteria has not been finally worked out, and it is possible that further research may show that sometimes they are but forms in a life-history of an organism appearing at some stage in bacillary form.

Micrococci are of constant occurrence, and pathogenic in septic processes, acute suppuration, acute infective osteo-myelitis, and erysipelas. There is strong but not absolutely complete evidence to the same effect in respect of pneumonia, gonorrhœa, ulcerative endocarditis, scarlet fever, and puerperal fever; micrococci are generally present in active lymph of vaccine and small-pox, and the virulence is largely or wholly lost when they are removed by filtration; but beyond this the proof of their being the actual materies morbi is incomplete. Ammoniacal fermentation of urine is determined by M. ureæ, which converts urea into ammonium carbonate. Certain diseases of animals are believed to be due to specific micrococci. Micrococci are also concerned in decomposition.

Bacilli are known to be pathogenic in anthrax, tuberculosis, tetanus, glanders, and leprosy, and are believed to be so in enteric fever, diphtheria, septicæmia, influenza, syphilis, and various forms of food intoxication. Many diseases affecting lower animals are due to specific bacilli, and among them malignant ædema, swine fever, and Rauschbrand ("symptomatic anthrax" or "black leg"). The fermentation of cheese and butter, and the souring of milk and cream, are likewise due to certain bacilli. Each fermentation and most forms of decomposition possess a specific ferment of this nature.

Spirillum obermeyeri is found in the blood during acute relapsing fever; the spirillum of cholera, or Vibrio cholera asiatica, is regarded as the cause of cholera; Spirochata pallida is associated with syphilis; and other forms occur in certain tropical

diseases (yaws, etc.).

It should be mentioned that other micro-organisms, not classed with bacteria, have relation to disease and fermentative processes. Thus, thrush is due to Oidium albicans, a mycelial form of a torula (Saccharomyces mycoderma). Mycelial growths, with spores or conidia, are associated with ringworm (Tricophyton tonsurans), favus (Achorion schönleinii) and pityriasis versicolor

(Microsporon furfur).

Actinomycosis is characterised by the presence of a mycelial growth, with masses of club-shaped corpuscles radiating from the centre—the streptothrix actinomyces (or ray fungus). Somewhat similar organisms are the streptothrix madura (occurring in Madura foot), and the pathogenic streptothrices recently described. These all belong, provisionally, to the higher bacteria. Minute amæboid bodies termed plasmodia (hæmatozoa) are pathogenic in malaria, and parasitic amæbæ, twelve times the size of white blood corpuscles, have been found in the intestines and fæces in cases of dysentery, and in pus from tropical abscesses of the liver associated with dysentery.

Organisms destitute of chlorophyll, having their

habitat in living hosts, animal or vegetable, are termed parasites, while those living upon dead organic material are known as saprophytes. Many microbes are capable of fulfilling both parts, though not necessarily with equal facility. Parasites that have not been found to grow under any known conditions as saprophytes are distinguished as "obligate," others as "facultative." Bacillus anthracis is an example of a facultative parasite, and the unknown microbes of rabies and many other infective or inoculable diseases may be provisionally classed as obligate. Cultivation in artificial media, that is, a form of saprophytism, is essential for the complete proof of the pathogenic nature of a given microbe. Bacteria may therefore be classified as parasitic or other, or pathogenic or other.

Other classifications may depend upon the characteristic growth of bacteria in artificial media in pure culture, and their effect upon such media, and whether they grow under aërobic or anaërobic conditions.

The action of microbes is essentially one of oxidation. As a rule carbonic acid gas is given off, but many known and unknown bodies are formed coincidently, which vary greatly with the pabulum as well as with the microbe. Among these are the poisonous alkaloids (toxins), which, so far as certain pathogenic microbes are concerned, are usually to be regarded as the immediate agents in the production of diseasesymptoms. They are soluble, and even when freed from their respective microbes have in certain cases been found to be capable of producing a slight and transient appearance of characteristic symptoms. They may also be protective against disease, and it has been shown, for example, that the injection of filtered fluid in which anthrax bacilli have been cultivated protects against subsequent or even coincident inoculation with virulent anthrax. Hankin isolated from cultures of the anthrax bacillus an albumose, the injection of which into mice renders them insusceptible to anthrax. Brieger and Fränkel have isolated pathogenic albumoses from cultures of the specific microbes of diphtheria, enteric fever, cholera, and tetanus. Roux and Yersin believe the soluble poison to be a ferment, and Martin has shown that in anthrax and diphtheria the microbe produces such a ferment or enzyme, which reacts on the tissues to form albumoses, and that these break up into simpler bodies, which possess the specific toxic virulence and protective power.

Cultivation.—Microbes derive oxygen either from the air or from compounds containing oxygen; in the former case they are termed aërobic, in the latter anaërobic. Yeast (Saccharomyces cerevisiæ) is anaërobic, and acquires oxygen at the expense of the sugar, which is reduced to alcohol; the "vinegar plant," Mycoderma aceti, is aërobic, and oxidises alcohol into acetic acid. Aërobic microbes grow best at the surface of a liquid, anaërobic organisms deeper down away from contact with air or aërated layers of fluid. The distinction is not absolute, and many microbes are able to exist under either condition (facultative anaërobes).

Besides oxygen, microbes require nitrogen, which some are able to take from salts, such as ammonium tartrate, while others can only assimilate it from albumen or gelatin. Carbon is necessary, and also inorganic salts, including sodium, potassium, and phosphoric acid. Cultivation media should be

neutral or faintly alkaline.

Both liquid and solid media are used. Among fluid media are broths made from various kinds of meat, peptones and meat extracts, Pasteur's solution (water containing cane sugar, ammonium tartrate, and yeast-ash), Cohn's solution (water with ammonium tartrate, potassium phosphate, tricalcic phosphate, and magnesium sulphate), and bile salt and sugar solutions. Some of the more common solid media are boiled potato, solid egg-albumen, blood serum, agar-agar, nutrient gelatin (gelatin with admixture of meat extract), etc. The medium must be adapted to the organism.

After thorough sterilisation by heat, media are kept in sterilised test-tubes, guarded, by means of sterilised plugs of cotton wool, against infection by air-borne germs. Each tube may then be "inoculated" by means of a previously sterilised platinum wire charged with the microbes to be cultivated; the

wool plug is removed momentarily for this purpose, and then replaced. Potato cultures are made by inoculating the freshly cut surface of a boiled potato cut in strips and placed in sterilised test tubes. For plate-cultivation, a small portion of the liquid under examination is diffused through melted nutrient gelatin or other suitable medium, and then poured into a Petri plate (which is a flat, shallow glass dish), and allowed to solidify by cooling; protection from aërial germs is afforded by a glass cover. The temperature must be kept constantly at the point which experience has shown to be most favourable for the microbe in question. For this purpose an "incubator" is used, in which the temperature is regulated (Room temperature, or cool, incubators are automatically. regulated at 18 to 22° C., and blood heat incubators at 37 to 38° C.

The temperature is regulated by thermo-regulators.)

Liquid media which solidify offer great advantages. more than one variety of microbe is present, the resulting colonies tend to remain distinct, and it is comparatively easy to establish a pure cultivation of each by inoculating afresh from it. Plate-cultivations are especially useful in this way, since by regulating the proportions the distribution of the colonies may be made as sparse as is necessary. Accidental contamination by aërial organisms is readily prevented, Another important advantage is that pure cultivations upon solid media frequently exhibit naked-eye peculiarities of outline and characters of growth that are important as means of identification. Another method of obtaining a pure cultivation is adopted by cultivating under conditions inhibitory to some microbes, but not to certain others: this is effected by adding phenol (Parietti), tauro-cholate of soda (McConkey), crystal violet (Drigalski and Conradi), etc., or by maintaining the temperature at 45° C.

Among the points that serve for the differentiation of the large number of varieties of bacteria are the following: - Microscopic appearances, as to form, size, structure, movement; effect of certain staining and decolorising agents; media in which growth occurs, and temperature required; rate of growth; macroscopic (and if necessary microscopic) appearance of cultivation in solid media, including form, colour, distribution (superficial or deep), liquefaction of gelatin, production of gas, etc.; results of inoculation upon animals.

Attenuation of virus has been effected in

several ways.

1. Subculturing from old cultures.—Pasteur found that an organism retains its virulence unimpaired if kept in sealed tubes for a short time only, or if cultivated with a minimum interval between the successive cultures. If, however, the cultivation is allowed to remain for weeks or months without subculturing, the virus is attenuated, and when inoculated causes a mild but protective form of the disease.

- 2. Cultivation at high temperatures.—Pasteur prepared a "vaccine" (i.e. an attenuated virus) of anthrax by cultivating the bacilli for three weeks in air at 42° to 43° C. There is no visible change in the mode of growth during this period except that spores are rarely if ever formed, but spore formation is immediately resumed if the bacilli are transferred to a fresh culture at ordinary temperatures. The virus becomes more and more attenuated if the temperature is maintained, and in about a month the bacilli all perish. The attenuation is brought about more rapidly at higher temperatures; a few days suffice at 45° C., a few hours at 47° C., a few minutes at 50° to 53° C. The original virulence is regained speedily upon transferring the attenuated bacilli to fresh media at ordinary temperatures, if the attenuation has been rapid. If, however, it has been slow, it is difficult to restore the virulence.
- 3. Cultivation in presence of inhibitory reagents.— Klein cultivated anthrax bacilli in gelatin containing 0 0025 per cent. of mercuric chloride, and obtained an attenuated virus which for a time protected guineapigs. Cultivations from the attenuated virus were virulent. Other reagents, including potassium bichromate, phenol, and terchloride of iodine, have been employed in like manner; or the organism may be

grown on a medium which is either slightly too acid or too alkaline.

Anthrax bacilli from sheep or cattle cause fatal anthrax if again inoculated upon susceptible sheep or cattle, or upon white mice; but bacilli taken from white mice produce only a transient illness in sheep, which, however, is protective. Assuming Jenner's view to be correct, vaccinia is an example of attenuation of virus, that of small-pox being so modified by transmission through the (insusceptible) cow as to produce a mild but protective form of the disease.

It must not be forgotten that the opposite of attenuation may also be obtained. Exaltation of virus results from early sub-culture on favourable media or the passage of the virus through susceptible animals.

These changes brought about in the laboratory by artificial means find a close parallel in the varying virulence of epidemic diseases. Sometimes the type, whether of small-pox, scarlet fever, diphtheria, or measles, will be malignant and fatal, and at other times the same disease will be so mild as to cause scarcely any deaths.

Toxins and their effect.—When a pathogenic organism grows in the body, it produces as a result of its metabolism certain poisonous substances termed toxins. These may occur as a direct result of the life of the bacillus, or as a result of a ferment produced by the bacillus. Toxins are of various kinds, and by their effect upon the blood and body tissues they cause the symptoms of disease. Fever is produced by the action of albumoses (bodies allied to the albumins) upon the heat-regulating centres in the brain. Albumoses also cause a number of other symptoms and poisonous effects. Toxins act, broadly speaking, in two ways. They have a local effect (as in the formation of an

abscess owing to necrosis of proliferating cells), and a general effect. The latter is caused when toxins are absorbed and distributed generally throughout the body. When this occurs they produce degenerative changes in muscles, in organs, and in the blood itself. Of such a change diphtheria is an example. The bacillus occurs in a false membrane in the throat, and occasionally other parts. It first causes the inflammatory condition giving rise to the membrane, and then it breaks it down. In the body of the membrane the bacillus appears to secrete a ferment which by its action and interaction with the body cells and proteids, chiefly those of the spleen, produces albumoses and an organic acid (Martin), which are the toxins. They are absorbed, and as a result we get the frequent pulse and high temperature of fever: the toxins irritate the mucous membrane of the intestine, and cause various fermentative changes in the contents of the bowel, and thus we get symptoms of diarrhea: they penetrate the liver, spleen, and kidney, setting up fatty degeneration and its results in these organs: they finally affect many of the motor and sensory nerves, breaking up their axis-cylinders, and producing the characteristic paralysis.

Toxins have been divided by Martin into extracellular (ferments, albumoses and alkaloids) and intra-cellular, or poisons present in the body of the bacillus and not "available" if the bodies of the bacilli

be filtered out.

The virus of anthrax produces albumoses and an alkaloidal substance (Martin), the former producing fever, the latter ædema, congestion, and local irritation. Hankin holds that the bacillus first produces a ferment and then elaborates albumoses. In tetanus the bacillus produces a secretion of non-proteid toxin which causes convulsions. The albumoses present in this disease are probaby due to the secretory toxin. Ehrlich has

isolated a spasm-producing toxin (tetanospasmin), and a crude poison capable of destroying red blood cells (tetanolysin). The nature of the tetanus toxin is not determined, but it is known that it is a most powerful poison, probably less than 1 grain being poisonous to man. In diphtheria, too, we have a secretory poison in the membrane and in the tissues, and an albumose which is possibly the result of the secretion. In typhoid fever intra-cellular bacillary poisons exist, and a toxalbumin has been obtained which has pathogenic effects of an indefinite character. The toxins of the typhoid bacillus appear to have little

digestive effect.

The action of bacteria as disease producers depends then (1) upon the effects of the presence of the bacteria themselves, (2) upon their power of forming, directly or indirectly, certain chemical organic products known as toxins, and (3) upon the individual attacked. effects of bacteria, though very diverse, may be classified generally as of a necrotic or a separative character, leading to increased functional activity at first (such as phagocytosis), and subsequently to increased formative activity (such as cell growth and subdivision). In most diseases the lesion has a special site (as in typhoid fever) and the body generally is only affected indirectly. This localisation may be due to specific action, or to point of entrance of the bacillus (as in malignant pustule). Secondarily to tissue changes, the body metabolism is affected owing to the absorption and distribution of toxins, and it is to this cause, as a rule, that the chief symptoms of disease are due. Finally much depends upon the soil upon which the bacteria and their toxins are implanted. The individual is an extremely important factor in the production and character of disease.

Antitoxins are believed by some authorities to. be a kind of ultra-toxin, substances of which an early

form was a toxin; others hold that, as the toxins are products of the bacteria invading the tissues, the antitoxins are of the nature of ferments produced by the resisting tissues. A third view is that possibly antitoxins may be the result of an increased formation of molecules normally present in the tissues. Thus antitoxins came to be looked upon as protective substances produced in the body cells as a result of toxic action, and held in solution in the blood, and there and elsewhere exerting their neutralising influence by combining with the toxins. These antitoxic bodies gradually increase in the blood and tissues, and their actions fall into two groups, (a) antitoxic, which counteract the effects of the poison itself; and (b) antimicrobic, which counteract the effects of the bacillus itself. "In one and the same animal the blood may contain a substance or substances which are both antitoxic and antimicrobic, such, for example, as occurs in the process of the formation of the diphtheria and tetanus antitoxic serums" (Martin). Antitoxin must, therefore, be looked upon as a normal constituent of the living cells which is produced in increased quantity. Of the chemical nature of toxins and antitoxins very little is known. Toxins are probably of the nature of albumoses, and antitoxins probably have a molecule of greater size, and may be allied to the globulins. Antitoxin has been shown to appear in the various secretions of the body as well as in the blood, though in a less concentrated state. The relation of the antitoxin to the toxin, and its mode of antagonism, is probably analogous to chemical union.

Immunity depends upon some property in the living blood serum which opposes or annuls, partly or wholly, the products and action of the infecting organism. Buchner designated these protective bodies, held in solution in the blood, alexines, and

regarded them as belonging to the albuminous bodies of the lymph and plasma. Alexines are naturally produced antitoxins; ordinary antitoxins are acquired alexines, corresponding to "natural" and "acquired" immunity. The term natural immunity is used to denote natural resistance to some particular specific disease. It may be due to species of animal, or age, or individual idiosyncrasies. For example, cholera and typhus fever which affect man do not affect the lower animals. Swine plague does not affect man. The white rat is immune to anthrax, which readily attacks cattle. Acquired immunity is a protection not belonging to the tissues of individuals naturally and as part of their constitution, but acquired during life in one, or both, of two ways. Either it may be an involuntarily acquired immunity, or a voluntarily acquired immunity—a natural attack of disease, or an artificial attack such as that due to inoculation. Small-pox, and a number of other diseases, rarely attack the same individual twice, because each of these diseases leaves behind it, so to speak, its antitoxic influence. Hence the individual has involuntarily acquired immunity against these diseases. An example of voluntarily acquired immunity occurs in the old method of preventive inoculation for smallpox, or variolation. This was an inoculation setting up an artificial and mild attack of small-pox, by which the protective antitoxins of that disease were produced. It was a voluntarily acquired immunity. This form of artificial production of protection is artificial immunity. When artificial immunity is produced by direct inoculation of attenuated bacteria in toxins (as in vaccination or Pasteur's treatment of rabies) the condition is known as active immunity; where antitoxins are inoculated (as in the antitoxin treatment of diphtheria) the condition is that of passive immunity.

CHAPTER XIII.

DISINFECTION.

Disinfection is the destruction of the specific virus upon which infection depends, and disinfectant is equivalent to bactericide. It is, however, often loosely applied also to antiseptics, that is, to substances that arrest or impede the growth of microbes without destroying their vitality, and even to deodorants or reagents that destroy or mask the effluvia which are the frequent by-products of

bacterial growth.

It is probable that protracted exposure to air and light will ultimately destroy most, if not all, pathogenic microbes, although under favourable conditions the virus of tuberculosis and other transmissible diseases has been known to retain its vitality for many years. It would seem also that mere diffusion in the atmosphere may render a virus inert. Typhus attacks a very large proportion of those who come into close contact with typhus patients, but rarely spreads under other conditions, even if the isolation is imperfect. Small-pox, on the other hand, is believed to be carried by air currents for long distances under favourable meteorological conditions. Spores of pathogenic bacilli would doubtless be among the most resistant, but mechanical portability (influenced by the form of the microbe and the size and weight of the epithelial scales or other particles to which it may be attached), the dryness (or the reverse) of the air, and the chemical effect of the atmospheric oxygen, are possible factors, and it is at all events conceivable that a certain minimum "dose" of the virus is

necessary for infection, so that mere dilution beyond a given point would render the virus harmless, even without destroying its vitality. According to Koch, mere drying (desiccation) kills the cholera bacillus.

The complete disappearance, often for long periods, of measles and other infectious diseases from a district after a widespread epidemic is in itself a sufficient proof that an enormous amount of contagium does in some way or other speedily become inert, apart from all attempts at disinfection and the exhaustion of

available susceptible material.

Mechanical removal of infection from rooms or garments is often an important adjunct to disinfection proper. Walls are stripped of paper, or scraped, or rubbed with dough, or washed; floors are washed or swept; a strong current of air is sent through the room; garments are washed or brushed, or beaten, and hung out in the open air. As a rule, such measures, though effectual as far as they go, are in themselves incomplete safeguards, since some portion of the virus may escape dislodgment, and there is no certainty as to the future harmlessness of the rest.

True disinfection, that is destruction of germs,

may be effected by heat or by chemical methods.

Disinfection by heat is the simplest and most thorough of all methods. With articles of small value. the safest plan is of course to burn them, but when this radical remedy is inapplicable, disinfection may usually be effected by exposure to moist or dry heat.

The experiments of Koch, Parsons, and others, have shown that bacteria and even spores of some bacilli are destroyed by exposure to steam at a temperature of 212° F. for five minutes, or to hot air at a temperature of 220° F. for four hours. Less resistant organisms, for example micrococci or bacilli free from spores, are destroyed by hot air at 220° F. in an hour.

Water at a temperature of 212° F.—that is, boiling water—is at least as efficacious as steam.

The mode of experiment was to steep threads in cultures of the respective organisms, and after exposing them to known temperatures for measured periods, to test their vitality by inoculation or further cultivation; "control experiments" were, of course, made in each instance, to prove that threads treated in exactly the same way, except as regards exposure to heat, gave positive results in cultivation or inocu-Anthrax spores and anthrax bacilli serve lation. admirably as test objects, since they are themselves pathogenic, and can be readily cultivated or inoculated, with characteristic and unmistakable results. Little is known with certainty as to the specific microbes of the ordinary infectious diseases attacking man, and in regard to them inoculation experiments are inadmissible, so that for the present we can only place reliance upon such processes as are found experimentally to destroy the most resistant of other organisms, namely, the spores of bacilli.

In practice the problem of disinfection is almost always complicated by the fact that the virus is not exposed freely, but enclosed in garments, pillows, or even beds; that is, in more or less bulky articles made of materials that have been selected for use as being the worst conductors of heat.* It is found

A brief note may be introduced here respecting the microscopic and chemical characters of these materials. Under the microscope, wool fibres, like hair, are straight, unbranched, and exhibit faint cross-markings and marginal indentations, due to the imbrication of the scales. Silk filaments have no such markings, and are structureless except for occasional nodes. Linen fibres have many nodes, with ragged branching filaments. Cotton

^{*} Wool and silk are the warmest materials used for clothing. Not only are they bad conductors of heat, but they absorb a considerable amount of moisture without becoming damp. The watery vapour of perspiration is condensed, and its latent heat rendered sensible, while the tenacity with which the moisture is held prevents re-evaporation and chill. Linen and cotton are much better conductors of heat, and absorb little water. Perspiration readily makes them damp, and they are then chilly owing to evaporation.

that steam rapidly penetrates into the interior of such objects. The first portions condense, parting with their latent heat in so doing, and create a partial vacuum, so that successive supplies of steam follow continuously until a temperature of at least 212° Fahr. is attained at the centre. Dry hot air, on the contrary, being dependent upon conduction, very slightly aided by convection, has no such power, and it is practically impossible to raise the temperature at the centre of a bed or similar bulky object to 212° Fahr. by dry heat within any reasonable number of hours. Hot air moistened by steam is superior to dry hot air in penetration, but not in germicidal power, and is far inferior to steam in both respects.*

fibres are flat and twisted, without nodes or branches. The chemical reactions are shown in the following table:--

Reagent.	Wool.	Silk.	Cotton.	Linen.
Potash or Soda (hot strong solution). Sulphuric Acid Nitric (or Picric) Acid Zinc Chloride (hot strong solution). AmmoniumCuprate (solution of CuO)	Dissolves. Nil. Turns yellow. Nil. Swells.	Dissolves. Dissolves slowly. Turns yellow. Dissolves.	Nil. Gelatinises. Nil. — Dissolves.	Nil. Gelatinises Nil. — Dissolves

^{*} Under ordinary atmospheric pressure of 14.7 lbs. to the square inch, pure water boils at 212° F.; if the pressure be doubled, at 249°; if trebled, at 273°; these being the temperatures at which the vapour tension of steam is equal to the given pressure. Further heating of water beyond the boiling point at constant pressure (e.g. in the open air where the steam can escape freely) merely produces steam ("current steam"), the temperature of both water and steam remaining at the boiling point corresponding to that pressure. Steam thus in contact with water, whether under pressure or not, is "wet" or "saturated," and if cooled or if the pressure be increased it begins at once to condense to water again. If, however, it is further heated out of contact with water, it becomes "dry" or "superheated" and behaves as a permanent gas until such time as cooling or increase of pressure brings it to the condensing point.

Steam, in condensing to water, shrinks to $\frac{1}{1200}$ of its bulk, and gives off latent heat sufficient to raise its temperature 300° F. There is Another important consideration is the effect upon the colour and texture of fabrics exposed to heat. Articles composed in part of fusible substances, such as glue or sealing-wax, are, of course, ruined by heat in any form. Steam is inadmissible for leather objects, since it shrivels them up and renders them worthless; hot air merely makes them temporarily dry and brittle. With these exceptions, steam is less injurious than hot air in almost all respects. New woollen goods, such as blankets and flannels, lose some of their whiteness and fleeciness by either process, but not more than in one or two ordinary washings. Silk and cotton are not injured by steam, nor by hot air if the temperature is carefully regulated. Dyes are surprisingly little affected by either steam or hot air.

The chief difficulty in steam apparatus is to prevent loss of heat, and condensation. This is overcome by surrounding the steam chamber with a "steam-jacket," that is, by making the wall of the apparatus double, and admitting steam into the space between the inner and outer casings. A door is provided at each end, one for the reception of infected goods, and the other for removal of the goods after disinfection. The doors are steam-tight, and are fastened by strong screw-clamps. The articles to be disinfected are placed in

trays, or suspended from sliding racks.

Washington Lyon's apparatus is oval in section, and is usually worked with a pressure of 10 lbs. per square inch in the jacket and 5 lbs. in the chamber, so that the steam in the latter is superheated, a further precaution against condensation. The articles having been introduced, and the doors closed and secured, steam is first sent into the jacket so as to heat the contents of the chamber. Steam is next admitted into the chamber itself, and soon reaches the full pressure required. It is found that penetration is more rapid if the pressure is intermitted once or twice, which is readily effected by turning a

corresponding expansion and absorption of latent heat in conversion of boiling water into steam. At any given pressure the boiling point of water containing salts in solution is slightly raised.

cock. Ten to twenty minutes suffice for the penetration of even bulky objects, and at the end of that time the steam is allowed to escape from the chamber, the door is opened, and the articles are removed. Further drying may be effected if necessary by leaving the door ajar for a few minutes, and

exposing the articles to heat from the jacket.

In Goddard, Massey, and Warner's apparatus the lower part of the jacket serves as the boiler, and the pressure in the chamber is the same as that in the jacket, usually about 20 lbs., so that the steam is not superheated. After the articles are put in and the door closed, hot air is drawn through for two or three minutes, and this is repeated after the ten or twenty minutes' exposure to steam.

The Equifex disinfector, upon the model of Geneste and Herscher, largely used on the Continent, has no steam jacket, but is lagged, and hot steam-pipes within the chamber serve to lessen condensation and dry the disinfected articles. The usual pressure is about 10 lbs., and the steam is not super-

heated.

Low-pressure steam—that is, steam under pressure only slightly, if at all, exceeding the atmospheric pressure—was found by Koch to be not inferior in germicidal power to steam, at higher pressure. Van Overbeek de Meyer devised an apparatus in which the steam, generated at atmospheric pressure in a jacket-boiler surrounding the chamber, passed in through an opening at the top. The temperature did not exceed 212°, but the results were practically as good as those attainable with high pressure. In Thresh's Current Steam Disinfector a somewhat higher temperature is obtained without pressure, by the use of a saline solution, in place of plain water in the jacket-boiler. Reck's is still simpler, the jacket being dispensed with.

All of these fulfil the essential conditions—namely, rapidity of penetration, destruction of all organic life in the interior of bundles of moderate bulk without injury to fabrics, and lastly, convenience in working. High pressure, intermitted, has perhaps some advantage in rapidity of disinfection of exceptionally bulky articles; but for ordinary purposes the simpler and less costly low-pressure apparatus is almost equally good. Esmarch has cast doubt upon the germicidal efficiency of superheated steam.

Hot-air-disinfection apparatus labours under a threefold disadvantage as compared with steam. The available temperature and duration of exposure are limited by the tendency to scorch the articles exposed; the penetration of heat is so slow that it is practically impossible thoroughly to disinfect objects of moderate thickness, such as pillows; and lastly, the germicidal effect of a given temperature is far less with hot air than with steam. The only advantage that can be claimed for hot air is that leather and bound books are not spoiled by it as they are by steam.

The best apparatus of this kind is that devised many years ago by Ransom. It consists of a rectangular chamber of wood and iron, lined with non-conducting material. Hot air mixed with the products of combustion passes into the chamber after being heated by a number of gas jets; the temperature of the hot blast is kept constant by an automatic mercurial regulator, which controls the gas supply and can be set for any required temperature. An outlet with a chimney is provided at the top The best working temperature is about of the chamber. 255° F., this being the highest that can be used for long periods without danger of singeing cotton goods. Thermometers are placed at the inlet and outlet. If the contents of the stove should take fire owing to overheating or to lucifer matches being left in the pockets, the gas is automatically turned off and the entrance of air arrested, by the falling of a weighted lever, which, until such accident occurs, is held suspended by a chain attached to a fusible metal bar in the interior of the chamber. It is customary to expose articles to hot air at about 255° F. for two, four, or eight hours or more, according to their bulk, but for reasons already stated the penetration of heat is very limited.

The following figures, which give the mean values obtained in a series of 120 experiments, serve to show how slowly the temperature rises in the interior of woollen objects exposed to hot air. Registering maximum thermometers were placed beneath layers of blanket. The inlet temperature was about 255°

F. throughout the series, and the final outlet temperature 245° to 250° F. :-

Duration of Exposure.	layers.	layers.	6 layers.	layers.	18 layers.
4 hours	220° F.	206° F.	190° F.	162° F.	139° F.
	226° ,,	214° ,,	208° ,,	174° ,,	153° ,,
	230° ,,	221° ,,	215° ,,	196° ,,	182° ,,

In striking contrast to the above were the results of another series of experiments with steam apparatus, the same blankets being employed. An electric thermometer, ringing at 212° F., was placed beneath sixteen or more layers of blanket, and served to indicate the exact interval between the first exposure to steam and reaching 212° F. Different forms of steam apparatus were tested, with low pressure as well as high, and the maximum interval noted was 17 minutes.

It is essential that any apparatus for disinfection by heat should have doors at opposite ends, opening into entirely separate rooms provided with separate entrances. One of these rooms should be strictly reserved for infected and the other for disinfected goods, and no articles should on any account be allowed to enter the latter room except through the stove, the object being of course to guard against the danger of

reinfection of the purified articles.

Chemical disinfection has been shown by the accurate experiments of Koch and others to be a matter of considerable difficulty, and comparatively few of the so-called disinfectants in common use prove to be really germicidal under the conditions of actual practice. A convenient but severe test is to expose to the "disinfectant," for a definite period, threads which have been soaked in a cultivation containing anthrax spores and then dried. After the "disinfection," cultivation or inoculation experiments show whether the spores survive or not, control experiments being made at the same time. Among the very few substances which killed the spores within a day (and for practical disinfection this is far more than can often be allowed) were mercuric chloride (1 per cent.), carbolic acid and potassium permanganate (5 per cent.), and chlorine and bromine water. A 4 per cent. solution of carbolic acid took three days, and 1 per cent. permanganate had no effect in two days. Among those which failed were 5 per cent. solutions of chloride of lime, zinc salts, copper sulphate, ferrous sulphate, boracic acid, and sulphurous acid, and 5 per cent. carbolic oil. Several of these, however, are able to kill less resistant forms, such as sporeless anthrax bacilli; or, serving as antiseptics, to prevent multiplication. A substance, to be a satisfactory disinfectant, should possess five characters: (a) it should be germicidal within a reasonable time limit; (b) it should not possess chemical properties which unfit it for ordinary use; (c) it should be soluble in water, or capable of giving rise to soluble products in contact with the material to be disinfected; (d) it should not produce injurious effects on the human tissues; and (e) it should not be too costly in proportion to its germicidal value. The substances which fulfil all these requirements are not numerous. Many valueless experiments have been made respecting such chemical bodies, because the conditions necessary to valid and comparable results have been ignored. These conditions are chiefly three, namely, using disinfectants which give regular and consistent results, using standardised bacterial cultures as to age and source, and working with the same organism. Out of the confusion of contradictory findings it is possible, however, to name a number of reliable disinfectants. The liquid reagents most in use are mercuric chloride

(1 to 500), potassium permanganate (5 per cent.), formalin (40 per cent. solution), carbolic acid (5 per cent.), zinc chloride (2.5 per cent.), chloride of lime (1 to 2 per cent.), creosol, cyllin, lysol, creolin, and many compounds of the aromatic series (e.g. izal and Jeyes' fluid, which contain the higher phenols, and form emulsions with water).

A most important consideration in regard to the more potent reagents forming the first series is their "working strength." If, for example, it is proposed to disinfect a putrescent liquid by means of permanganate of potash, it is absolutely useless to add a little of a 5 per cent. solution of the salt. We must add either the solid permanganate or a highly concentrated solution, until the permanganate is present as such to the extent of 5 per cent. of the whole weight of liquid, this 5 per cent. being of course in addition to the amount required to oxidise the organic matter. These essential conditions are rarely if ever observed in practice, and "disinfection" by permanganate consists really of deodorisation with partial oxidation of organic matter. A similar consideration applies to mercuric chloride, which, if added to liquids containing organic matter, forms a precipitate that carries down part of the mercury in an inert form, and if sulphuretted hydrogen is present, the equally inert sulphide of mercury is thrown down. So, too, with carbolic acid, which must form not less than 5 per cent. of the whole weight of liquid—not merely of the stock solution—if it is to destroy anthrax spores.

Mercuric chloride is one of the chief reagents that can be conveniently employed in solution under such conditions as to destroy the most resistant microbes. One part of mercuric chloride in 1,000 of water destroys anthrax spores, according to Koch, but other observers have found this strength inadequate. The great drawback to this reagent is its extremely poisonous nature, but it may be kept in fluted poison bottles properly labelled, and the solution may be artificially coloured with indigo, and "odorised" with thymol, as further safeguards. The following proportions are suggested in a memorandum of the **L.G.B.** (1905):—½ oz. mercuric chloride, 1 oz. hydrochloric acid, and 5 grains of commercial aniline blue, in three gallons of water. This ought not to cost more than one penny per gallon and should not be further diluted. Non-metallic vessels (wooden or earthenware) should be used. Articles soaked in the mercurial solution should be steeped in water for some hours before washing. Whether the contagia of human diseases are amenable to less potent reagents is still an open question, but for the present it is not safe to assume this.

Many fluid disinfectants are now applied direct to infected surfaces by means of a sprayer (Equifex, Mackenzie, etc.), such as chloride of lime (1 lb. to 10 gallons of water), carbolic acid (5 per cent.), formalin

(40 per cent. solution, 4 ozs. to the gallon).

Fumigation.—For the disinfection of rooms and other closed spaces the principal reagents in use are sulphurous acid, chlorine, bromine, nitrous acid, and formic aldehyde. Sulphurous acid is usually generated by setting fire to fragments of roll sulphur in an iron vessel, with the addition of a little spirit to facilitate lighting. Sulphur candles are convenient substitutes, or bisulphide of carbon may be burnt in a lamp. The prescribed proportions are 1 to 3 lbs. of sulphur to every 1,000 cubic feet of air space, yielding theoretically 1.1 to 3.3 per cent. of sulphurous acid in the air of the room.

Chlorine is most conveniently produced by adding crude hydrochloric acid to chloride of lime. The strength of the latter varies, but as a rule 1½ to 2 pints of the acid should be allowed for each pound of chloride of lime. No heat is required; indeed the reaction is so rapid that it is necessary to secure time for the operator to escape, by letting the acid drip through a small hole upon a little cup which, when full, overflows into the dish containing the lime. The

quantities required, according to German experiments, are very large—about 15 lbs. of chloride of lime and 22 lbs. of hydrochloric acid * for every 1,000 cubic feet. These quantities, being somewhat prohibitive, are commonly reduced to one-tenth or less in practice. The reagents should be divided into a convenient number of portions and placed in several parts of the room in positions as near the ceiling as practicable.

Nitrous fumes are generated by adding copper

filings or other reducing agents to nitric acid.

In any case the room must be rendered as nearly air-tight as possible before fumigation. Paper should be pasted over the fireplaces and ventilators, and around the window-sashes and doors, leaving, of course, one door to be pasted up on the outside after the operator quits the room. It is desirable to render the air of the room moist by means of steam or water before fumigating by sulphurous acid or chlorine, to an extent sufficient to moisten the walls and other surfaces.

In ordinary rooms it appears that the unavoidable leakage is considerable, so that the theoretical proportion of the gas is only present for a very short time, if at all. The slightest protection shields even sensitive microbes from the action of the fumigant; very few are killed which lie in crevices, or in the pocket of a coat, or are wrapped in filter paper. Koch's experiments in an almost air-tight box showed that sulphurous acid, in the proportion of 1.0 per cent. of the cubic space, killed anthrax bacilli and other sensitive organisms in half an hour, but that anthrax spores and other spores of bacilli resisted 6.0 per cent. for days. The addition of moisture greatly accelerated the action upon bacilli. and so far increased the effect upon spores of bacilli

^{*} Strong sulphuric acid may be substituted for crude hydrochloric acid. About a third of the quantity would suffice.

that some of them were killed by exposure to 5 per

cent. sulphurous acid for twenty-four hours.

Formic aldehyde vapour can be obtained by various devices in the way of lamps (paraform lamp burning 30 tabloids per 1,000 cubic feet); more effectively by Trillat's autoclave, a somewhat costly apparatus, in which its 40 per cent. aqueous solution, known as formalin, is treated at high pressure in presence of calcium chloride, the gas being passed through the keyhole by means of a tube; or by Lingner's apparatus, in which glyco-formal (formalin 30 per cent. and glycerine 10 per cent.) is vaporised and ejected as a fine spray. If the formalin is simply heated much of the aldehyde changes into a solid polymeride paraform. The amount of formalin required for each thousand cubic feet of air-space in a room is variously stated as 150 to 500 c.c. The vapour, in the proportion of 1 per cent. in air, or solution of 0.5 per cent. strength, is found to destroy most organisms. purposes of fumigation formic aldehyde has advantages over sulphurous acid and chlorine in its lower density, which gives it greater diffusibility and some power of penetration, and in its more rapid action. It acts best if the air is dry, and is harmless to most colours and most surfaces except iron. The vapour is irritating, but readily cleared away by ventilation.

Klein, Houston, and Gordon, in 1902, found that formic aldehyde gas (30 tabloids of 034 oz. each to 1,000 cubic feet space) killed the cholera and typhoid bacilli, and the diphtheria bacillus, as well as staphylococcus pyogenes aureus, in five hours, and devitalised tubercular sputum on linen and paper. The same may be said of sulphurous acid gas (3 lbs. to 1,000 cubic feet) in 24 hours, except that the sputum remained unaffected. Anthrax spores were not killed

uniformly under any of these conditions.

It would seem, therefore, that if precautions are

by sulphurous acid, chlorine, or formic aldehyde, may be able to destroy most, if not all, of those microbes that are not very resistant, and that

happen to be freely exposed upon the surface.

Fumigation is to be regarded as—at most—a disinfection of air and of surface. Even this is an important step in the direction of safety, for the surface is precisely that portion of an infected article which is most readily and most largely charged with infective matter, and from which, moreover, infection is most readily given off to persons handling or coming into contact with the *fomites*. As a preliminary process it affords some degree of security to those engaged in carrying out later and more thorough measures of disinfection. It can be supplemented, as regards surfaces, by washing with soap and water, or with a disinfectant solution; wall-papers can be stripped off, or lime-wash renewed.

Disinfection of the air of a room is practically ensured by the free ventilation which in any case must be effected, but if infected surfaces or articles are allowed to remain, the air may at any time again become charged with infective matter accidentally

dislodged from its resting-place.

Besides the germicidal power of the reagent there are three other conditions essential to disinfection—namely, sufficient strength or intensity, maintained for a sufficient time, with sufficient contact or exposure.

In few departments of hygiene have absurd generalisations been so prevalent as in reference to disinfection. "Disinfectants," or substances labelled as such, are still widely employed under conditions which a moment's reflection would show to be utterly incompatible with any real efficacy. Passing over such trivialities as "fumigation" by burning brown paper or green sticks, we have the wholesale employment of deodorants and antiseptics under the name of

disinfection, unsupported by any scientific evidence of their utility. The few true disinfectants in general use are commonly employed under such conditions as to quantity, concentration, and duration of exposure, that little benefit can result from them. Earth is popularly believed to be a universal disinfectant, and yet earth abounds in microbes, some of which are pathogenic. Indeed, the very nitrification upon which the purifying action of earth is so largely dependent is in great part due to the life and growth of microbes. It is often asserted that the danger of infection by diseased meat is exaggerated, because the process of cooking must destroy any microbes that may be present. Even as regards microbes, the temperature in the interior of joints obviously may fall far short of that required for disinfection, and the ptomaines and unorganised ferments (toxins) are in all probability uninjured by ordinary cooking. The gastric juice also is credited with a disinfecting power much greater than the evidence warrants. The "comma" bacillus and many others are destroyed by it, but spores may escape. Sarcinæ and other microbes may be found in abundance in the stomach itself, fully exposed to the alleged germicidal gastric juice; and in the intestine millions of microbes are always present. Moreover, it is now established beyond dispute that tuberculosis and other diseases are communicable to animals by means of the alimentary canal.

Attempts are often made to "disinfect" sewers and drains by flushing them with solutions of chloride of lime, carbolic acid, or ferrous sulphate. A little consideration will render it clear that any real disinfection of this kind is wholly impracticable, even if true disinfectants such as strong solutions of mercuric chloride are used. The reagent is quickly diluted by the contents of the drain, and rapidly passes away without coming even momentarily into effectual contact with the whole of the material and surfaces it is supposed to disinfect. If the drain or sewer is properly constructed, thorough flushing by water will be more effectual

than the use of disinfectants.

Antiseptics are reagents which prevent, arrest, or impede the growth and multiplication of microbes, without necessarily destroying them. All true chemical disinfectants are still antiseptic when diluted, provided that the dilution is not carried too far (mercuric chloride has some trace of antiseptic power even at the enormous dilution of 1 in 300,000). Among

other important antiseptics are:—Boracic acid, borax, sulphurous acid; essential oils, including those of mustard, thyme, turpentine, peppermint, cloves, and

eucalyptus; quinine and other alkaloids.

Antiseptics are employed mainly to prevent or arrest decomposition of organic substances. Apart from surgical purposes, they are used for the preservation of food, to prevent or restrain decomposition in organic refuse, as a precaution against rotting of

timber, and in many other ways.

Among aërial deodorants, nitrous acid is one of the most powerful. Chlorine and the fumes given off by moist chloride of lime are also potent, and act by oxidation of organic matter; they decompose sulphuretted hydrogen, which is an important constituent of the gases of putrefaction. Hydrochloric acid fumes, like chlorine, neutralise the free ammonia and ammonium carbonate. Sulphurous acid may, perhaps, act in some degree as a reducing agent, and also as an antiseptic, but its chief effect is to overpower the effluvia and necessitate free ventilation. Fumes of wood, tar, or paper are quite useless except for the same reason.

Of the solid or liquid deodorants, ferrous sulphate and cupric sulphate act mainly by removing the sulphuretted hydrogen as a precipitate; potassium permanganate simply oxidises; carbolic acid and the essential oils exert an antiseptic effect and so check further decomposition, while at the same time their powerful odour masks all others. The ozone or peroxide of hydrogen which is believed to be associated with the essential oils may effect some small amount of oxidation.

CHAPTER XIV

SPECIFIC DISEASES

Small-pox has been described more or less clearly in Eastern countries for nearly two thousand years, and more definitely in Europe since the sixth century. No part of the world is exempt from epidemic visitation; but in India and the Soudan it is so constantly prevalent that these may be regarded as endemic foci. Pandemic extension occurs from time to time, characterised by the enormous areas attacked, and by the malignant type of the disease. The last of these was that of 1871-2, which overran Europe and America, and caused in England alone 42,000 deaths during the

two years.

The whole epidemic character of small-pox has completely changed since the introduction of vaccination. The "bills of mortality" show that upon the average 7 to 9 per cent. of the persons buried in London during the seventeenth and eighteenth centuries had died of small-pox, and in epidemic years the proportion often rose to 13, 15, or even 18 per cent. The only approach to this state of things since the introduction of vaccination was in the pandemic year 1871, when 9.8 per cent. of the deaths in London were due to small-pox. The general mortality upon which this percentage is calculated is, of course, far smaller in proportion to population than in the last During the greater part of last century small-pox became somewhat prevalent in London every four or five years, but in the decade 1891-1900 it remained at a very low ebb, although there have been signs of reviving activity since 1900. same remarks apply more or less to the other English

towns, all of which suffered heavily in 1871-2, and with few exceptions have been unprecedentedly free from the disease in recent years. A severe epidemic occurred in Sheffield in 1887-8, however, in the manufacturing districts of Yorkshire and Lancashire, and at Gloucester in 1896, and throughout the country in 1902-3.

Season.—The mortality curve for small-pox in England is above the mean from January to June, and below it from July to December. The New York curve does not differ essentially from this, but rises to a more definite maximum in May. In India the maximum occurs in the cold season, namely, in March or April.

Soil is not found to have any influence upon the

prevalence of small-pox.

Air.—In 1884-5 it was found by Power that cases of small-pox in the surrounding districts followed the admission of acute cases into Fulham Hospital. He showed that, if the district were divided into zones, by means of circles drawn upon the map from the hospital as a centre, with radii of $\frac{1}{4}$, $\frac{3}{4}$, and 1 mile respectively, and an enumeration made of all the houses in each belt, and also of all houses invaded by small-pox, the proportion of invaded houses diminished as the distance from the hospital increased, and this relation held good in each "quadrant" of each zone (Fig. 23). Within the quarter-mile zone there was only one approach to the hospital, and this was in the N.W. quadrant. The distribution of cases in the several quadrants was not such as to suggest any relation to lines of traffic or ambulance routes. Power concluded that diffusion only occurred when acute cases were aggregated, and perhaps only under certain atmospheric conditions that cannot as yet be defined.

Barry found a similar radiation of infection from

hospital upon a larger scale in the Sheffield epidemic of 1887-8, but the possibility of personal convection could not be entirely excluded. Similar observations with similar results, namely, a graduated intensity of small-pox incidence as distance between small-pox hospital and a populous neighbourhood became less,




Fig. 23.—Incidence of Small-pox upon District around Fulham Hospital,
May 25th, 1884, to September 26th, 1885.

The figures show the percentage of houses invaded in each quadrant and
quarter-mile zone.

have been made at Oldham in 1893; Warrington 1892-3; Bradford 1893; Glasgow 1900-01, and the Orsett Union, adjoining the site of the small-pox hospital ships in the Thames, from 1884 to 1902. These and other careful investigations leave little room for doubt that under favourable conditions small-pox can be conveyed through the air, and for considerable distances. An observation at Nottingham, 1887-8, tends to show that

the same radiation may occur upon a small scale from a single acute case. The town had been absolutely free from small-pox for a year, when a case was imported and concealed in a crowded house in a dense quarter. On the south side this house was closely adjoined by courts and alleys that were only accesssible (from the north) by a long and circuitous route; streets and houses on all the other sides of the house in question were freely accessible. Three houses ssituated in the isolated courts on the south were invaded, but none in the directions in which traffic The dates of attack in each case correcould pass. sponded with the hypothesis of aërial convection from the concealed case during the acute stage, and the wind was northerly at the time of probable infection in each. Confirmatory evidence as to such diffusion of small-pox from centres outwards has been repeatedly recorded.

Neither water nor milk has been shown to convey

the infection of small-pox.

Age and sex.—The mortality from small-pox is greater among males, not only at all ages taken together, but also, with few exceptions, at each age period. The exceptions are the second and third year of life, and the tenth to the fifteenth year, at all of which, from some unexplained cause, the female

mortality slightly exceeds the male.

The relation to age has to be considered with reference to vaccination. In pre-vaccination times, as McVail has shown from the Kilmarnock registers, 90 per cent. of the deaths were at ages below five years, the actual maximum being in the second year. Under present conditions, the deaths under five years, being practically limited to unvaccinated children, constitute only 30 per cent. of the total small-pox leaths, and in this age period the greatest mortality occurs in the first year, being, indeed, highest in the

first three months of life. It diminishes steadily from this point until about the fifteenth year, rises to a second maximum about the twenty-fifth year, and thenceforward diminishes again.

Race.—Coloured races, and especially negroes, have a peculiar susceptibility to small-pox. They take it more readily, and suffer a heavier case mortality, than white races, apart from all question of vaccination.

The incubation period is usually twelve days, or counting to the appearance of the rash, fourteen days, so that the rash is seen on the fifteenth day, on the same day of the week that the infection was acquired. There are, however, exceptions to this rule. Inoculation of small-pox is followed by constitutional symptoms on the seventh or eighth day, and the general rash appears on the tenth or eleventh. Even after infection by more ordinary means it is stated that the latent period may range from seven to eighteen days; and the rash is often delayed until the fourth day (i.e. the third after onset) instead of appearing on the third.

Infection is given off, presumably by the breath, from the earliest onset. The exhalations, the vesicles, pustules, and scabs are also highly infectious. The virus may be carried by air currents, as already mentioned. It readily lodges in clothing, and retains its vitality tenaciously; hence outbreaks are liable to occur in paper-works and other manufactures in which rags are employed. Isolation should be maintained for at least three weeks in the mildest case, and always until every scab has disappeared—a process which, in confluent cases, will take six weeks, and often considerably longer. After exposure to infection, a quarantine of seventeen days—to allow a margin over the fortnight—is sufficient. Second attacks are rare, except after several years' interval. Third attacks are not unknown.

Diagnosis.—The medical officer of health is not unfrequently consulted as to the diagnosis of variola, and it may be desirable therefore to add here the main points to be borne in mind as

characteristic of the disease :-

Sudden attacks of fever, headache, vomiting, backache (most marked in the lumbar and sacral regions) generally begin from 12 to 14 days after infection. Rigors are often present. Vomiting and retching are sometimes severe. In children (and even in adults occasionally) these initial symptoms may be obscured by drowsiness.

These initial symptoms last two or three days before the characteristic rash appears. But in some cases accompanying these symptoms are various forms of mild eruptions, which may simulate measles or scarlatina. Such eruptions are fre-

quently localised to different parts of the body.

The characteristic eruption of small-pox appears about two days after the onset of the backache, fever, etc. It consists at first of a number of small red pimples or spots, which gradually enlarge and become hard and "shotty" to the touch. After they have been out for about 48 hours they begin to become small bladders or vesicles, containing at first clear, and, later on, purulent matter (pustules). They are about the size of a ssplit pea, are surrounded by a thin red zone (or areola), are definitely raised above the level of the skin, often assume the colour of dull pearl, and in many cases show a central depresssion (umbilication). About the eighth or ninth day the pustules begin to dry up, so that from the first appearance of the rash, the successive stages of pimple, vesicle, and pustule occupy about ten or eleven days. Simultaneously with its appearance on the skin, the eruption appears on the mucous membrane of the mouth and palate.

In persons who have been vaccinated the rash is somewhat modified (varioloid) and comes out more rapidly; the vesicles are smaller and less abundant, not so much depressed in the

centre, and may not become pustular.

An important point is the distribution of the rash. The first spots generally come out on the face, particularly on the forehead and around the lips. Almost immediately afterwards they appear on the limbs, being particularly marked on the wrists and ankles. Afterwards the rash appears on the back, whilst the skin on the front of the chest is less affected and the skin of the abdomen least. A further point is that, as a rule, when the rash comes out the temperature falls and the patient feels better. In rare instances the rash is almost entirely suppressed, and therefore in times of epidemic the appearance of the premonitory symptoms alone should arouse suspicion.

The three most important aids to correct diagnosis arethe premonitory symptoms, the character of the rash, and the distribution of the rash.

The precautionary measures to be taken in case of an outbreak of small-pox include immediate and rigid isolation, which can only be satisfactorily carried out in hospital, especially when the danger of aërial spread of infection is considered; thorough disinfection; vaccination, or re-vaccination of all persons who have been exposed to risk of infection, however remotely, those only being excepted who have recently (say within four years) undergone successful vaccination, or who bear marks of recent small-pox. No reliance can be placed upon verbal assurances as to previous attacks, or even as to vaccination, and after a time the protection of previous small-pox, as well as of vaccination, may vanish.

The pathogenic microbe of small-pox and vaccinia may possibly be a bacillus (Klein and Copeman). It is found most readily in early lymph, about the fifth day, and can be cutivated upon the hen's egg,

though not upon ordinary media.

Inoculation of small-pox (variolation), already referred to above, was generally practised in England in the eighteenth century, having been introduced from the East in 1721 by Lady Mary Wortley Montagu. It seems to have been customary to inoculate from the primary vesicle, not from the secondary rash. Mild cases were selected, and the lymph was taken before suppuration set in. The fatality (2 or 3 per cent.), though far lower than that of "natural" small-pox, was by no means triffing, and the readiness with which the risk was incurred is a strong illustration of the magnitude of the danger to which it was the only known alternative. It was superseded by vaccination, and was finally forbidden by law in 1840.

Vaccination, introduced by Jenner in 1796, gradually superseded inoculation during the earlier part of the last century. It was provided gratuitously by the first Vaccination Act of 1840, made compulsory in

1854, and systematically enforced by paid vaccination officers from the time of the pandemic in 1871. Under the Act of 1898 more latitude has been allowed to the conscientious objector.

DEATH-RATES FROM SMALL-POX, PER 1,000 LIVING AT EACH AGE-PERIOD.*

	All Ages.	0-5.	5–10.	10-15.	15-25.	25-45.	Over 45.
1847-1853. Vaccination gratuitous but op- tional	0.302	1.617	0.337	0.094	0.109	0.066	0.022
Vaccination compulsory but not efficiently enforced .	0.223	0.817	0.243	0.088	0.163	0.131	0.052
1872-1891. Vaccination enforced .	0.089	0.177	0.095	0.054	0.097	0.086	0.038

This table shows that each extension of vaccination was attended with a reduction in the small-pox mortality, in which, however, the different ages by no means shared equally. The reduction was greatest at the ages of highest mortality—that is, among children under 10 years of age—while at ages over 15 the change was comparatively slight. In the third period the latter no longer included, as in former generations, a large proportion protected by non-fatal attacks of small-pox undergone in early life, and on the other hand the protection due to infantile vaccination had had time to fade. Moreover, even unvaccinated persons had, under the new conditions, a better chance of

^{*} Reg.-Gen. 43rd Ann. Rep.; and Foster, British Medical Journal, i., 1893, p. 1069. The ages were not abstracted before 1847.

escaping small-pox until adult life. The enormous reduction in mortality among young children whose vaccination was recent speaks for itself, and it must not be forgotten that the mortality at ages under 5 years occurs among the unvaccinated residuum, the vaccinated being practically exempt, as will be seen presently.

Another illustration of the great change in the age-incidence of small-pox has already been given. At Kilmarnock, from 1728 to 1764, there were 622 deaths from small-pox, of which 563, or 90 per cent., were at ages below 5 years. The corresponding proportion in England from 1871 to 1880 was 30 per cent. Buchanan* gave other instances in the following table, with an analysis of the age-incidence of small-pox mortality in London in 1884 upon the vaccinated and unvaccinated, separately, from which it is manifest that the old conditions remained unchanged among the unvaccinated, except that the protection which the rest of the community had acquired lessened the probability of exposure to infection:—

CONTRIBUTION OF VARIOUS AGES TO 100 SMALL-POX DEATHS AT ALL AGES.

	1580-1760.	, ý,	51.	-51.	London, 1884.			
	Geneva, 1580-	Kilmarnock, 1728–1764.	Paris, 1842-	London, 1848-51	Un- vaccinated.	Vaccinated.	Whole Community.	
0-10 years 10-20 ,, 20-30 ,, 30-40 ,, Over 40 ,,	96·1 2·6 1·0,	98·8 0·5 0·7 —	39·7 13·3 32·9 11·0 3·1	81.5 5.9 8.3 3.2 1.1	61·2 14·6 10·8 7·2 6·2	8.6 17.3 31.9 22.1 20.1	34·3 17·0 21·3 14·2 13·2	
	100.0	100.0	100.0	100.0	100.0	100.0	100.0	

^{*} Report of the Medical Officer to the L.G.B. for 1884,

An analysis of the returns of the 5,683 small-pox deaths in London during the ten years 1879-88, as given by the Registrar-General, yields the following results:—

Ages.	Of 1,406 stated to be Vaccinated.	Of 2,265 stated to be Un- vaccinated.	Of 2,012 Vaccination not stated.	Of the total 5,683, irrespective of Vaccination.
0 to 10 years 10 to 20 years Over 20 years	Per cent. 8.4 16.0 75.6	Per cent. 56.0 17.2 26.8	Per cent. 35.0 14.7 50.3	Per cent. 36.8 16.0 47.2

Confirmatory evidence, if any be needed, is afforded by Barry in reference to the Sheffield epidemic of 1887-8:—

SHEFFIELD, 1887-8.—SMALL-POX ATTACKS AND DEATHS (PER 1,000 Persons of Each Class stated).

	Attack-rate.		Death-rate.		
	Vaccin- ated.	Unvac- cinated.	Vaccin- ated.	Un vac-	
0-10 years	5	101	0.1	44	
Ditto, living in invaded	78	869	1	381	
houses	19	94	1	51	
Ditto, living in invaded houses	281	686	14	371	
All ages	15.5	97	0.7	. 48	
Ditto, living in invaded \\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	230	750	11	372	

From these statistics it appears that the vaccinated part of the population had, as compared with the unvaccinated, at ages below 10 years, a 20-fold immunity from attack, and 480-fold security against death from small-pox; at ages above 10 years, a 5-fold immunity from attack, and 51-fold security against death from

small-pox; at "all ages," a 6-fold immunity from attack, and a 64-fold security against death from small-pox.

Sheffield Hospitals, 1887-8. Proportion of mild ("varioloid" and "discrete") and severe ("coherent" and "confluent") cases per 100 attacks among persons of each class stated. (BARRY.)

			Vacci	inated.	Unvaccinated.		
			Mild.	Severe.	Mild.	Severe.	
0-10 years All ages .			91·0 82·8	9.0 17.2	21·8 18·5	78·2 81·5	

As regards London, the proportion of small-pox deaths to deaths from all causes is available as a measure of the prevalence of the disease before and after the introduction of vaccination, which came gradually into operation at the beginning of the 19th century. From the "bills of mortality" it appears that during the thirty-one years 1770-1800 small-pox caused 59,253 out of the 626,530 deaths in London, or 9.4 per cent. In the thirty-one years 1801-31 the proportion was 4.6 per cent. From 1832 to 1837 no returns are obtainable; but for the thirty-one years 1838 to 1868

Period	Period		Mean Annual Small- Pox Mortality per 1,000.		
before Vaccination.	after Vaccination.	Locality.	Before Vaccina- tion.	After Vaccina- tion.	
1777-1806 1777-1806 1781-1805 1774-1801 1751-1800	1807-1850 1807-1850 { 1810-1850 1810-1850 1801-1850	Lower Austria Upper Austria and Salzburg Berlin Sweden Copenhagen	2·48 } 1·42 3·44 2·05 3·13	0·34 0·50 0·18 0·16 0·29	

the proportion was 1.4 per cent. Even in 1871, when the pandemic wave reached London, the deaths from small-pox were only 9.8 per cent. of the total deaths, proportion very little in excess of the average which prevailed between 1770 and 1800.

The figures in the table at the foot of page 338

are quoted from Seaton.

The greater susceptibility of the unvaccinated to the small-pox virus is manifested not only in a greater liability to attack, but in a higher case mortality.

The quality of vaccination also—that is, the numper, area, and character of the cicatrices—has an important bearing upon the degree and permanence of the protection afforded. Both of these points come but clearly in all large hospital statistics, of which he following two examples will suffice:-

Pase Mortality in relation to quality of vaccination and to age, based upon an analysis of 10,403 cases in Metropolitan Small-Pox Hospitals. (GAYTON.)

Ages.		ccinated. od Marks.		In	ccinate aperfec Marks.		Va	ecinat Mark	ed.	Unv	accina	ted.
	Cases.	Deaths	Case Mort- ality,		Deaths.	Case Mort- ality.	Cases.	Deaths	Case Mort- ality,		Deaths.	Case Mort- ality,
0-5	51	0	0.0		21	11.2	128	47	36.7	677	883	56.6
0-10	267	2	0.7	714	48	6.7	325	87	26.8	1,187	568	47.4
10-20	1,045	17	16	1,976	98	5.0	419	81	19.5	521	160	30.7
20-40	725	37	5.1	1,898	258	13.6	420	140	33.3	382	181	47.4
over 40	48	6	12.5	266	51	19.2	131	44	33.8	79	84	43.0
All ages	2,085	62	3	4,854	455	9	1,295	352	27	2,169	938	43

Wase Mortality in relation to number of vaccine cicatrices. (MARSON.) Case Mortality

Unvaccinated .				47. 26		. (per cent.
tated to be vaccinated,	but	wit	hout	cicatr	ices		213
Having one cicatrix							71/2
Having two cicatrices			14				$4\frac{5}{8}$
Having three cicatrices							1 3
laving four or more cid	catri	ces					3 4

Re-vaccination renews in all respects the immunity given by primary vaccination. As regards entire populations, the results of general, if not universal, re-vaccination are shown in the following figures. taken from the report of the German Vaccination Commission of 1884. After the pandemic of 1870-74, re-vaccination was made compulsory in Germany only, and small-pox has never become seriously epidemic during the years that have since elapsed; while in other countries the incidence continued much as in the period before 1870.

SMALL-POX DEATH RATES.

I mily	Austria.	Prussia.*	Berlin.*	Hamburg. *	London.	Paris.	Vienna.
1870	0.30	0.175	0.22	0.25	0.30	5.46	0.46
1871	0.39	2.432	6.32	10.75	2.42	?	0.74
1872	1.90	2.624	1.38	0.95	0.53	0.06	5.36
1873	3.23	0.337	0.11	0.008	0.04	0.01	2.28
1874	1.78	0.092	0.024	0.002	0.02	0.05	1.35
1875	0.58	0.036	0.051	0.000	0.01	0.13	1.13
1876	0.39	0.031	0.018	0.018	0.20	0.50	1.67
1877	0.53	0.003	0.004	0.012	0.70	0.07	0.84
1878	0.61	0.007	0.007	0.002	0.38	0.04	0.75
1879	0.51	0.013	0.007	0.000	0.12	0.45	0.46
1880	0.64	0.026 (0.008	0.000	0.12	1.09	0.73
1881	0.83	0.036	0.047	0.022	0.61	0.49	1.23

The German Vaccination Law was passed in 1874 compelling vaccination and re-vaccination. Infants must be vaccinated before the end of the calendar year following the year of birth, and all children must be re-vaccinated in their twelfth year. Obedience to this law is enforced by fine or imprisonment. completely is this system worked, that there is less need for the strict isolation of small-pox than in England. For instance, in Germany, the rare cases

^{*} Italic figures are inserted for the years in which compulsory revaccination was in force.

of small-pox are placed in pavilions of general hospitals, not in separate small-pox hospitals. The population of Germany is 56 millions, and from 1891 to 1902 nelusive, there were 607 deaths recorded as due to small-pox. The small-pox deaths during the same period in England and Wales (population 32 millions) were 6,761. If the ratio had been the same in England as in Germany, there would have been only 350 deaths.

Barry shows that in the Sheffield epidemic the revaccinated had a great advantage over the rest. Of 3,198 persons re-vaccinated prior to the epidemic, 25 were attacked and 1 died, the attack-rate being therefore 3.0 per 1,000, and the death-rate 0.1. Among 56,233 persons re-vaccinated during 1887-8, 2 were loubtfully attacked and none died.

It may be convenient to summarise the evidence as follows:—

A. As regards vaccinated and unvaccinated populations respectively.—(1) In pre-vaccination times the heaviest incidence of small-pox was upon childhood. (2) In the unvaccinated part of the community it is so still. (3) Among vaccinated persons childhood is exempt, and small-pox is only fatal in later years, when the protective influence of vaccination has faded. (4) In a mixed community (vaccinated and unvaccinated) both these effects are apparent, the heavier incidence upon the unvaccinated compensating for their scanty numbers. (5) The immunity at early ages among the vaccinated may be extended indefinitely by re-vaccination. (6) In every epidemic, in every community, and in every year-if the numbers are sufficiently large to preclude fallacy—the unvaccinated suffer more attacks in proportion to their numbers than the vaccinated, and (7) a larger percentage of the attacks are fatal. (8) Among the vaccinated who suffer from small-pox in later years, the proportion of fatal to non-fatal attacks diminishes as the number or area of the cicatrices increases.

B. As regards entire populations.—(9) The introduction of vaccination has in all cases been followed by a lowered average incidence of small-pox. (10) Each extension of compulsory vaccination has led to a further reduction. (11) Compulsory re-vaccination has almost entirely suppressed small-pox.

The vaccine virus is that of cow-pox or vaccinia, a disease formerly prevalent among cattle, but now rarely met with. It occurs in the form of vesicles upon the teats and udder in cows. The virus is inoculable upon human beings as well as cows, and this inoculation constitutes vaccination. Goats also are susceptible. No person is insusceptible except as the result of previous vaccination, or an attack of small-pox.

Primary vaccination in the human subject is followed on the third day by a papule, which on the fifth or sixth day is vesicular. On the eighth day the vesicle is mature; it is greyish, elevated, tense, loculated, and contains a clear viscid lymph which exudes when the vesicle is pricked. On the ninth day inflammatory changes set in; the lymph begins to contain pus, and a reddish "areola" surrounds the vesicle. Slight constitutional symptoms are common at this stage. The vesicle becomes a pustule, begins to dry up about the tenth day, and forms a dark scab about the fourteenth. The scab falls off in the fourth week, leaving a permanent scar which should be depressed and pitted.

Vaccination may be carried on from arm to arm for an indefinite and probably unlimited number of years without having recourse to the cow again. The lymph should always be taken on the eighth day (that is, exactly a week after the vaccination) from normal and unbroken vesicles. No lymph should be collected after a re-vaccination, or from an unhealthy subject, or from vesicles showing any sign of inflammatory action. By far the surest plan is to vaccinate directly from the vesicle, but the lymph may be sealed in capillary tubes or dried upon ivory points, and retains

much of its efficacy for a considerable time.

Vaccination by calf lymph is now in use upon a large scale, owing partly to a popular suspicion that the current arm-to-arm vaccination is losing its

ower, but chiefly to the fear of the virus of syphilis r other malady being inoculated with the lymph from iseased human subjects. There appears to be no ifference either in the course of the symptoms or in esults between ordinary arm-to-arm vaccination and accination from the calf.

Vaccination is protective against itself as well as gainst small-pox. Any number of "insertions" may e made at the time of vaccination. Whether one resicle is produced or a dozen, the protection is abpolute for the time being, but all experience goes to how that the duration of the protection is limited, and is directly proportionate to the number and area ff cicatrices produced. Hence it is desirable to vacciate in at least four places, and the total area of the icatrices should not be less than half a square inch. The return of the Metropolitan Asylums Board for the small-pox epidemic in London, 1901-2, respecting his point, is given here, and a study of it will amply prove the claim made.

	Admissions.	Deaths.	Mortality per cent.
accinated cases—		Secretary S	THE STATE OF THE S
accinated cases—	- Indianal		LEtys Italy
Half square inch and up-	1 0,8900		LOSSING THE
wards	5,163	379	7.34
One third but less than half	3000000	7.01	75.00
square inch Less than one third square	835	131	15.69
inch	860	162	16.87
Not recorded	87	33	37.93
Totals of Vaccinated Class	6,945	705	10:15
nknown and Doubtful Class	436	171	39.22
nvaccinated Class	2,277	752	33.06
Grand Totals	9,658	1,628	16.87

As the protective influence of the primary vacination fades, a time arrives when re-vaccination

becomes possible. Comparatively few persons are insusceptible to re-vaccination after the lapse of ten or twelve years; many "take" readily within five years, although the primary cicatrices may be good. The course of re-vaccination in the majority of persons is different from that of the primary operation, being more rapid, and often failing to exhibit some of the typical stages. If, however, the former protection has entirely disappeared, the course of re-vaccination may be identical with that of a primary vaccination.

Vaccine lymph contains, among other microbes, bacilli which Copeman regards as pathogenic. He finds, too, that it is possible to kill out the extraneous organisms by suitable admixture with glycerine, leaving the lymph still active. Filtration renders it inert. The virus is somewhat readily destroyed by chemical disinfectants, but the lymph can be kept in the dry state, or in sealed tubes, for months or even years, although it begins to lose some of its activity from the very first.*

Jenner believed that vaccinia was small-pox modified by transmission through the cow, a form of attenuation very similar to that which we now know to be possible as regards many other diseases. Cow-pox, formerly common, has become rare among cattle coincidently with the reduction in human small-pox due to vaccination. The seat of the eruption was the teats and udders of cows, a limitation that suggests contact with the hands of milkers as a means of inoculation. It is

^{*} Glycerinated calf lymph is now prepared as follows:—A calf, aged three to six months, is kept in quarantine for a week. If then found upon examination to be quite healthy, it is removed to the vaccinating station, the lower part of its abdomen antiseptically cleaned. The animal is now vaccinated upon this sterilised area with glycerinated calf lymph. After five days the part is again thoroughly washed, and the contents of the vesicles, which have of course appeared in the interval, are removed with a sterilised sharp spoon, and transferred to a sterilised bottle. This is now removed to the laboratory, and transferred to a sterilised bottle. This is now removed to the laboratory, and the exact weight of the material ascertained. A calf thus vaccinated will yield from 18 to 24 grams of vaccine material. This is now thoroughly triturated and mixed with six times its weight of a sterilised solution of 50 per cent, chemically pure glycerine in distilled water. The resulting emulsion is aseptically stored in sealed tubes in a cool place. At intervals during four weeks it is carefully examined bacteriologically until by agar plates it is demonstrably free from extraneous organisms, when it is ready for distribution. free from extraneous organisms, when it is ready for distribution.

difficult to inoculate cows with small-pox virus, but if effected, raccinia seems to result. Hime has done so, in the calf. Ceely ucceeded twice in a large number of experiments, Badcock in only seven per cent. of his attempts, and other experimenters wave failed entirely. Voigt claims to have succeeded by the evice of vaccinating a calf in one part of its body, and at the name time inoculating with small-pox in another, true vaccine resicles appearing in both places. Chauveau failed to produce raccinia by inoculating calves with small-pox, but papules ppeared at the points of insertion, and the contents of these apules inoculated upon susceptible human subjects gave rise, ot to vaccinia but to small-pox. It is possible that Chauveau's esults correspond to Ceely's and Badcock's failure, not to their ew successes, since they, too, sometimes obtained slight local ffects which they did not regard as vaccinia; and the reproluction of human small-pox by inoculation from the papules hay have been due simply to the unaltered small-pox virus which he had inserted.

Bouley, and also Ceely, having produced such papules by moculating small-pox matter, afterwards inoculated the same

nimals with vaccine matter, and with success.

The incubation of vaccinia being shorter than that of small-pox, it is possible to modify or even entirely prevent an attack of small-pox by vaccination erformed some days after infection. This is especially the case with re-vaccination, the incubation of which

ss often shorter than in primary vaccination.

Successful vaccination within three days after excosure to infection will prevent the appearance of symptoms, and it is probable that the attack will either be arrested or modified by vaccination as late as the fifth or even sixth day. The proof of this rests apon the observation that attacks of small-pox may, and do, occur within six days of vaccination in persons who have been many days previously exposed to infection, but the few attacks that occur between six and nine days after successful vaccination are mild, and tractically none commence later. The usual incubation of small-pox, it will be remembered, is twelve days.

The Royal Commission on Vaccination, 1896,

concluded that the protection vaccination affords

against small-pox may be stated as follows:-

"(1) That it diminishes the liability to be attacked by the disease. (2) That it modifies the character of the disease and renders it less fatal and of a less severe type. (3) That the protection it affords against attacks of the disease is greatest during the years immediately succeeding the operation of vaccination. It is impossible to fix with precision the length of this period of highest protection. Though not in all cases the same, if a period is to be fixed, it might, we think, fairly be said to cover in general a period of nine or ten years. (4) That after the lapse of the period of highest protective potency, the efficacy of vaccination to protect against attack rapidly diminishes, but that it is still considerable in the next quinquennium, and possibly never altogether ceases. (5) That its power to modify the character of the disease is also greatest in the period in which its power to protect from attack is greatest, but that its power thus to modify the disease does not diminish as rapidly as its protective influence against attacks, and its efficacy during the later periods of life to modify the disease is still very considerable. (6) That revaccination restores the protection which lapse of time has diminished, but the evidence shows that this protection again diminishes, and that to ensure the highest degree of protection which vaccination can give the operation should be at intervals repeated. (7) That the beneficial effects of vaccination are most experienced by those in whose case it has been most thorough. We think it may be fairly concluded that where the vaccine matter is inserted in three or four places it is more effectual than when introduced into one or two places only, and that if the vaccination marks are of an area of half a square inch they indicate a better state of protection than if their area be at all considerably below this."

The Vaccination Acts of 1867, 1871, and 1874 have been modified by that of 1898. In general, a child must be vaccinated within six months of birth. The parent is relieved from obligation if within four months after the child is born he satisfies a magistrate that he conscientiously believes that vaccination would be prejudicial to its health.* And even in the absence of such an exemption, an order under the 1867 Act directing that a child be vaccinated cannot be made upon any person who has previously been convicted of non-compliance with a similar order relating to the same child. It is the duty of the public vaccinator, if the parent so requires, to attend the child at its home. If a child is not vaccinated within four months of its birth, the public vaccinator is to visit the home, after giving at least twenty-four hours' notice, and offer to vaccinate with glycerinated calf-lymph or such other lymph as the L.G.B. may provide. In either case, if he finds that owing to the condition of the house, or recent prevalence of infectious disease in the district, the operation cannot safely be performed, he is not to vaccinate, but must give a certificate of postponement, and inform the M.O.H.+

Public vaccinators are to use only glycerinated calf-lymph, or other lymph supplied by the L.G.B., and must keep registers from which the origin of the llymph used in each operation can be identified. The Instructions require the greatest care as to cleanliness

^{*} The Vaccination Act, 1907, provides for this claim to be made as a statutory declaration by the parent to the vaccination officer.

† The L.G.B. direct that, "except so far as any immediate danger of small-pox may require, the public vaccinator must vaccinate only subjects who are in good health. As regards infants, he must ascertain that there is not any febrile state, nor any irritation of the bowels, nor any unhealthy state of the skin, especially no chafing or eczema behind the ears or in the groin, or elsewhere in folds of skin. He must not, except of necessity, vaccinate in cases where there has been recent exposure to the infection of diseases such as measles scarlating or diphtheria, nor where infection of diseases such as measles, scarlatina, or diphtheria, nor where erysipelas is prevailing in or about the place of residence."

and sterilisation of instruments used, and aseptic precautions at all stages. "The public vaccinator must aim at producing four separate good-sized vesicles or groups of vesicles, not less than half an inch from one another. The total area of vesiculation resulting from the vaccination should not be less than

half a square inch."

The L.G.B. may, in the event of serious risk of outbreak of small-pox or other exceptional circumstances, require the Guardians to provide vaccination stations, and may temporarily and locally modify the obligation upon the public vaccinator to visit otherwise than by request. Any S.A. maintaining a hospital for small-pox must keep a register of patients, showing name, age, address, and condition as to vaccination. The entries are to be made at the time of admission, and the register is to be accessible to the public.

A child must be vaccinated within six months of its birth unless (a) the parent obtains a magistrate's certificate of exemption, or (b) the child dies, or (c) is attacked by small-pox within that period, or (d) three or more unsuccessful attempts at vaccination have been made,* or (e) a medical certificate of postponement (for not more than two months at a time) is given on the ground of ill-health, or of recent prevalence of infectious disease in the district, or of the condition of the house in which the child resides. Certificates of successful vaccination, or attack of small-pox, or of insusceptibility to vaccination, or postponement of vaccination, can be given by any registered practitioner.

Re-vaccination is entirely optional, but persons over twelve years of age are re-vaccinated gratuitously

^{*} In this event insusceptibility is inferred; but out of 38,000 primary vaccinations Cory failed in only one instance, and in that one he had no opportunity of making a third attempt.

by the public vaccinator, and if there is immediate danger of small-pox, the age-limit is reduced to ten years.

The administration of the Vaccination Acts, subject to the control of the **L.G.B.**, is entrusted to the Poor Law Guardians, not to the **S.A.** Instructions and blank forms of certificate are issued by the sub-registrars to the parents at the time of registration of birth.

Varicella or chicken-pox was only clearly distinguished from small-pox in the last century, but beyond a certain superficial resemblance there is nothing in common between the two diseases. Varicella occurs as an epidemic, which not unfrequently coincides with small-pox outbreaks, and adds to the difficulty of diagnosis of mild cases of the latter. Persons of all ages are liable to attack, but children more than adults. The mortality is practically nil, a fatal result being unknown except in cases almost moribund from other causes, or in cases of varicella gangrenosa. Nothing is known of the conditions that determine the appearance of this rare and malignant type of varicella in individual cases during an outbreak of ordinary tharacter.

The latent interval is very commonly a fortnight. The tharacteristic vesicular eruption, which may be very profuse and widely distributed, is thickest upon the back and shoulders. The vesicles are not usually umbilicated, although this is not reliable point for diagnostic purposes, and among them are some that are aptly compared to drops of scalding water upon the skin, there being no induration around them. At other times varicella spots simulate small-pox more closely, but there are almost always a want of uniformity in size, appearance, and tage of development. The premonitory symptoms (vomiting, ackache, anorexia, etc.) that accompany the onset of small-pox only, the distribution and appearance of the rash, the celation to vaccination, and the history of exposure to one or ther kind of infection will seldom fail to make the diagnosis llear.

The infection of varicella is active from the first onset, and readily imparted by contact or by means of fomites. The mode in which it is given off is unknown, but the breath is the most probable. Attempts have been made to inoculate from the vesicle, but without success. Second attacks are rare.

A quarantine of eighteen days is sufficient after exposure to

infection; after an attack, isolation should be kept up until all scabs have disappeared. In times of small-pox epidemic the L.G.B. or the county council or S.A. may make chicken-pox temporarily notifiable in order thereby to bring to light cases of small-pox simulating varicella.

Measles became clearly differentiated from smallpox and scarlet-fever in the seventeenth century. No country or race appears to be exempt from it, and it has no demonstrated relation to climate. Some dependence upon meteorological conditions may be inferred from the fact that the colder months, even in the tropics, are favourable to the rise and spread of measles epidemics. In London the seasonal curve of mortality shows two maxima, a principal one in December and a lesser one in June, the minima occurring in February and September.* The influence of age as regards liability to attack has not been worked out, but the mortality from measles is greatest in the second year of life, and rapidly falls in every succeeding year. Sixty per cent. of the deaths occur during the first two years of age, 75 per cent. in the first three, and upwards of 90 per cent. under five years. The mortality is greater among females at all ages above five years, but greater among males up to that point, so that upon the whole the female mortality is lower than the male.

Measles epidemics are occasionally so malignant in type as to have a case mortality of 30 per cent. or more, but in such instances neglect, improper treatment, or insanitary conditions have been largely responsible for the result. The epidemics now met with in Great Britain, though variable in severity of type, cause a case mortality that rarely exceeds five per cent., and is often not more than one or two per cent.

^{*} The winter maximum vanishes in many parts of Great Britain and the Continent, particularly where the measles mortality is low, leaving a simple curve with a maximum in June and a minimum in September or October. (Reg. Gen. Ann. Rept. for 1884. See also "Measles Epidemics, Major and Minor," Transactions of the Epidemiological Society, 1892-3.)

The fatal cases are as a rule due to pulmonary complications or diarrhea rather than to the primary disease itself, the tendency to the one or the other complication being often fairly constant throughout an epidemic, but apparently independent of season (Hirsch). Unlike scarlet-fever, measles often disappears almost entirely from a district in the interval between epidemics. These intervals, however, are usually short, and few towns in England escape an epidemic for much longer than two or at most three years at a time. Insular communities such as Iceland and Faröe, with little communication with the outside world, remain free for an indefinite number of years after the subsidence of an epidemic, until infection is introduced afresh. Occasionally measles, like small-pox, becomes "pandemic" and diffuses itself for a brief period over a large part of the globe.

Measles is said to have been inoculated by nasal mucus (Bristowe) with an incubation of seven or eight days. It has no known relation to soil. It does not appear to be transmitted by drinking-water, milk, or other food, or to have any connection with diseases of lower animals. "Insanitary conditions"—notably overcrowding, exposure, and want of food-diminish greatly the chance of recovery, as was found in the butbreak among French troops in the siege of Paris, when 86 out of 215 died. Whether such conditions increase the liability to attack cannot be stated. The virus appears to be readily conveyed by the air, probably for considerable distances, but of this exact proof is wanting. Measles has not yet been isolated upon the large scale, so that the question of the possiility of aërial spread of infection from hospitals has not arisen.

From analogy it may be assumed that a specific neasles microbe exists, but none has yet been satisactorily demonstrated.

Infection is probably always acquired by inhalation. The incubation period is usually about twelve days, and is followed by the stage of invasion marked by pyrexia and catarrhal symptoms affecting chiefly the respiratory tract and the conjunctiva. The rash appears on the fourth day of illness, first on the forehead, then becoming general. It consists of dusky pink, slightly elevated spots, which increase for two or three days, and then slowly fade. Slight scurfy desquamation follows. Infection is given off by the breath and mucus from the onset of the first symptoms, and very possibly even during the stage of incubation. It is not clear how long this liability continues, nor whether the desquamated cuticle is infectious, but the usual rule is to allow a quarantine of not less than three weeks. An attack is usually protective throughout life, and this is probably the main reason for the comparative immunity of adults, but well-authenticated instances of repeated attacks in the same person are far from infrequent.

Comparing measles with scarlet-fever, it is at once apparent that preventive measures are less likely to be successful against the former. Infection begins at least three days before the characteristic rash appears, and hence isolation is not adopted at a sufficiently early date to prevent the spread of the disease, so far as the household is concerned. The long incubation

adds to the difficulty of quarantine.

Measles is still allowed to run its epidemic course practically unchecked by any serious attempt at isolation or disinfection, and it is, therefore, not surprising to find that it shows no signs of sharing in the reduction of mortality which is so conspicuous in respect of scarlet-fever, enteric fever, and small-pox. It is customary to close schools during measles epidemics, and there may be some advantage in doing so in rural districts. (See Chap. VII.)

Whooping cough, or pertussis, has been recognised in Europe for 300 years, and has practically a world-wide distribution, although it appears to be on the whole less prevalent and less severe in hot countries. In London its average prevalence is greatest in March and April, least in September and October, but the seasonal curve of mortality varies greatly in different parts of Europe, and the New York curve has a second and greater maximum in

September.

By far the greatest mortality occurs in the first year of life. Of the whole number of deaths from pertussis, over 40 per cent. occur in the first year, mearly 75 per cent. under two years of age, and 96 per cent. under five years. At every age the mortality among females is greater than among males. There are no statistics sufficient to show the actual or relative incidence of attacks at different ages, but the vast majority occur in children, although age confers no complete exemption. Concurrent epidemics of measles and whooping cough are frequent. Like measles, whooping cough is a typically epidemic disease, and recurs at short but irregular intervals (usually two or three years) in outbreaks that affect a very large proportion of susceptible persons. The case mortality is usually small, and probably seldom exceeds 5 per The deaths are almost always due to bronchitis, pneumonia, convulsions, or other complications. Measles and whooping cough together are, as a rule, rresponsible for more deaths of children than all the other infectious diseases put together.

Whooping cough has not been shown to be transmissible by water, milk, or other food, or to affect Mower animals, or to have any relation with telluric conditions. It is readily conveyed by fomites. Faulty hygienic conditions - especially overcrowding and exposure-increase the danger to life, but there is no

proof that they increase susceptibility. The specific microbe has not been isolated, nor is the disease known to be inoculable.

Whooping cough is, in all probability, always acquired by inhalation of the virus. The incubation period appears to be about a fortnight, but can rarely be fixed definitely owing to the very gradual onset of characteristic symptoms. Squire believed it to be less than a week. After a week or more of apparently ordinary nasal and bronchial catarrh, the characteristic paroxysmal cough sets in, and continues for a very variable period, which is rarely less than two weeks and may exceed two months.

Whooping cough is certainly infectious in the preliminary catarrhal stage, and possibly even earlier than this. Isolation ought to be maintained until at least a week after the spasmodic cough has entirely disappeared. The infection is given off in the breath

and sputa. Second attacks are very rare.

What has been said as to the prevention of measles applies in all respects to whooping cough, and with even greater force. Owing to the absence of rash, and the insidious onset of symptoms, the disease is rarely detected until many days after it has reached the infectious stage.

Mumps (parotitis) occurs in all parts of the world in brief but often intense epidemics which cause little or no mortality. Cold and wet weather is the most favourable, and outbreaks usually occur in the winter and spring, often in concurrence with epidemics of measles, or sometimes scarlet-fever. Mumps usually disappears entirely from a district at the end of the outbreak, and there is no appearance of periodicity in its return, which may be delayed for many years.

Nothing is known of its etiology beyond its extreme infectiousness by means of the breath. Second attacks are very rare. The characteristic symptoms, including swelling of the parotid and sub-maxillary glands, set in after an incubation of about three weeks (often 19 days), and the disease runs its course in about a fortnight more. Three weeks is sufficient for

has been exposed to infection.

Rötheln, epidemic roseola, or German measles, occasionally becomes epidemic among children, but causes very slight mortality. An outbreak in the New Forest in 1880 was, however, said to have been atttended with a case mortality of 3 per cent. After a latent period, often about 21 weeks in length, slight general symptoms manifest themselves and are followed immediately, or in a day or two, by an ceruption of red, slightly elevated spots, variable in ssize and often confluent. They appear first on the face and fore-arms, increase and become more or less general in one or two days, and disappear before the end of a week. The symptoms have some resemblance tto those of scarlet-fever and measles, especially the llatter. It is, however, an entirely distinct and independent disease, and not a "hybrid," as is popularly ssupposed. The disease is infectious, but never very sactively so, and isolation for a fortnight is sufficient. Infection is probably given off by the breath, and acquired by inhalation. Second attacks do not sappear to be common. Quarantine should extend to 123 days.

Scarlet-fever, or scarlatina, has been recognised in Europe for at least 550 years, and in England upwards of 200 years. It was long confounded with measles, and only in the nineteenth century was it definitely distinguished from diphtheria. It is most widely diffused in Northern and Western Europe, and in North America, but has failed to establish itself firmly in Africa or any part of Asia, except Asia Minor. Temperate and somewhat humid climates seem to be most favourable to it, but the evidence upon this point is conflicting. In England it is more prevalent in urban than in rural areas, mining districts and several of the large manufacturing towns being most

affected of all. It is probable that the explanation of this is to be found in the greater facilities for receiving and transmitting infection from person to person. Scarlet-fever has a well-marked relation to season. In England the mortality is at its minimum in March and April, and rises to a maximum in October, subject to irregularities if only small numbers or short periods are taken. In New York, however, the curve is almost reversed, the minimum being in September and the maximum in April. The number of attacks varies according to season in the same way as the mortality, but the seasonal range is greater in the attack curve; apparently the attacks in spring are, on the average, more severe, though fewer, than those in autumn.

As to the meteorological conditions favourable to the spread of scarlet-fever, there is wide divergence of opinion. The rise and fall of scarlet-fever epidemics often cover a period of years, and the mortality in any particular week or month must not be regarded as determined solely, or even principally, by the meteorology of that brief season. It seems probable that the tendency to spread is increased by atmospheric humidity and absence of wind, but checked by heavy rainfall.

Age.*—Children under one year of age, and especially under three months, are comparatively rarely attacked, but the incidence rapidly increases and reaches its maximum in the fifth year of life. After that period it declines steadily year by year. About 45 per cent. of the attacks occur at ages under five years, 40 per cent. between five and ten years, and 11 per cent. between ten and fifteen years. The

^{*} The number of deaths from scarlet-fever increases from birth to a maximum in the third year of life, and steadily declines afterwards. 90 per cent. of the deaths occur at ages under ten years, and 65 per cent. under five years. Further details will be found in a paper on "Age, Sex, and Season in relation to Scarlet Fever' in the Transactions of the Epidemiological Society for 1887-8,

verage severity of attack, measured by the proportion of fatal cases, is greatest in infancy or in the second wear, and diminishes as age increases, but there is an apparent increase in the severity of the (very few) cases that occur in adults after the 15th or 20th wear. The lessened liability with increasing age is greater than can be accounted for either by protection afforded by previous attacks, or by diminished exposure to infection.

There is, therefore, a double gain in shielding a child from infection during the first few years of his life. Every year of escape after the fifth leaves him less and less susceptible, until finally he becomes almost immune; and secondly, if he should be attacked, every year that the attack is delayed reduces the danger to life, and also the risk of formidable complications that may become chronic and disabling even iff not immediately fatal.

Females are upon the whole more liable than males, but the attacks are somewhat less fatal, and hence up to ten years of age the mortality is higher among males than among females. At ages above ten the female deaths exceed the male, owing, no doubt, to the greater frequency of attack among women who

have the charge of children.

Scarlet-fever is for the most part an epidemic disease. Sometimes it will disappear entirely for years, or "sporadic" asses may occur without any serious general prevalence. In certain manufacturing towns, on the contrary, it may be said to be endemic, although subject to frequent epidemic outbursts.

Graves noted that scarlet-fever was prevalent and malignant in Dublin from 1801 to 1804; during the next 27 years epidemics recurred at intervals, but the type was always mild and the mortality slight; in 1834 the disease reverted once more to the old malignant form, but in the last four decennia it has been mild with low mortality. Several instances are on record in which the type has suddenly changed during an epidemic from mild to severe, or vice versa. In some epidemics the case mortality is as low as 3 per cent., or even less, while

in others it may reach 30 per cent. As a rule the cases are more severe during the early part of an epidemic, and among children more fatal in town than in country, and more fatal to

boys than girls.

No adequate explanation has been offered of the causes that determine the rise and fall of scarlet-fever epidemics, and still less of the causes of the variations in type. Seasonal changes, especially meteorological, doubtless have an influence, but this is superadded upon the true epidemic wave. During the interepidemic intervals the proportion of susceptible persons increases, but as regards ages above five years the delay lessens the susceptibility, and it does not appear that the intensity of the outbreak bears any relation to the length of the previous interval.

The pathogenic microbe of scarlet-fever, according to Klein and Gordon, is a streptococcus (streptococcus scarlatinæ or conglomeratus), which they have isolated from the blood and nasal and tonsillar discharge of persons suffering from the disease. These observers have grown this streptococcus in broth, gelatine, agar, milk, and blood serum, in several of which media it gives a characteristic culture. It is pathogenic for mice and rabbits. Gordon believes that both S. pyogenes and S. scarlatinæ play a part in the causation of scarlet-fever, but that S. scarlating is the more important of the two, and occupies a position in the bacteriological kingdom between S. pyogenes and B. diphtheriæ. Baginsky, Sommerfeld, and Class have also isolated streptococci which they believe to be the cause of scarlet-fever. They are probably forms of S. scarlating. Klein further believes that the same organism was found in cows suffering from the Hendon disease, also in a sample of condensed milk that apparently caused an outbreak of scarletfever, and in the blood of a monkey that, during an epidemic of scarlet-fever at Wimbledon, presented symptoms of the disease. It is found in the throat secretions of patients in the acute stage, but not in the desquamated cuticle or the urine, or in the

hap. XIV.]

ar discharge which occasionally affects convalescents. t has, however, been discovered in the nasal distharges of advanced convalescents, and in the throat ecretions long after apparently complete recovery. This persistence may afford an explanation of the bbscure "return cases" of scarlet-fever. It should not, nowever, be forgotten that a streptococcus occurs in the saliva of healthy persons (Gordon).

The period of incubation is usually three or four Mays, but it would appear that in exceptional cases tt may be less than twenty-four hours, or as long as eeven days. Some authorities believe the usual incupation period to be much longer, but Tonge-Smith's exact observations upon cases which, owing to error in diagnosis, were exposed to infection during transit to hospital in an ambulance, afford strong proof of

The accuracy of the estimate given above.

The onset in well-marked cases is sudden, with womiting, rigors, headache, pyrexia, and sore throat as prominent symptoms. The red rash follows in about 24 hours. It is at first punctiform, but becomes diffuse. It is usually seen first about the meck and chest, but may spread over the whole boody. The face usually remains clear of rash though it may be flushed, leaving a pale area around the mouth. The rash occurs most markedly on points of pressure and at flexures. It increases for one or two days and begins to decline in two or hree days more. Desquamation begins as soon as the rash disappears, the chest being the first and the nands and feet the last parts to clear. True relapses occur in one or two per cent. of the cases, usually about a month after the onset.

Infection is given off from the first onset and for a period that is seldom less than five weeks, and not unfrequently exceeds two months. It has been the practice to isolate until every trace of desquamation has disappeared from the feet, and still longer isolation may be necessary if any symptoms remain, such as discharges from the nose or ears, or albuminuria. But it is now not unusual to discharge cases from hospital after six weeks' detention if there are no nasal or aural discharges, regardless of desquamation, the later degrees of which are probably non-infectious.

The virus of scarlet-fever retains its vitality for long periods, possibly years, under favourable conditions, and can be lodged in crevices, or carried in

garments, etc.

By far the commonest mode of infection is from a previous case, either by contact, by proximity, or by means of infected articles. Whether the virus is usually inhaled or swallowed is less certain. Aërial convection for short distances—across a room, for example—may be assumed, but there is no proof of radiation of infection from scarlet-fever hospitals. Shirley Murphy has shown that at school ages notifications of scarlet-fever, as of diphtheria, are lessened by holidays. In Nottingham it was found that far fewer attacks began on Wednesday than on any other day of the week, owing presumably to lessened opportunity of infection on Saturday and Sunday.

Nothing is definitely known as to any relation between soil and scarlet-fever. There is no evidence that drinking-water can carry it, but milk certainly can, if infected from a human case of scarlet-fever, and epidemics such as the Hendon outbreak suggest that the cow itself may not improbably be a source

of infection.

It is at least doubtful whether "insanitary conditions" as to drainage, etc., have even a predisposing influence as regards this disease, but it seems probable that they may seriously affect the chances of recovery and promote septic complication.

It is often held that puerperal women are liable to acquire scarlet-fever in a dangerous form, and that under such conditions its course may be modified and constitute a variety of puerperal fever. In the event of infection under such conditions scarlet-fever may run a normal, and even mild, course.

Infection is given off by the breath and secretions from the nose, mouth and throat; and most freely in the earlier and acute stages. How far the desquamated cuticle is infective is a matter of some

doubt, direct experimental proof being wanting.

The preventive measures available are isolation and disinfection. Considerable difficulty arises from the long duration of the infectious stage, covering, as it almost invariably does, a period of at least four or five weeks after the patient has ceased to be ill. It is practically impossible, in a working-class household, to maintain the requisite seclusion, and the comparative freedom and better hygiene in hospital are conducive to the patient's welfare. Another source of difficulty is the occurrence of slight cases. There may be no noticeable symptoms whatever beyond sore throat, or slight desquamation, and yet any case, however slight, is capable of spreading infection. On the other hand, the short, and apparently noninfectious, incubation period, and the early appearance and distinctive characters of the rash and other symptoms, render this disease much more easy to deal with than measles or whooping cough, if proper facilities are afforded. Schools are often instrumental in spreading infection, and it may be necessary to close them when epidemic prevalence is threatened. Scholars from infected households should be excluded for at least six weeks, and only re-admitted upon a medical certificate, but a quarantine of a week is sufficient if the persons attacked have been removed to hospital and the premises disinfected. Special care is needed in guarding milk supplies from risk of direct or indirect infection from a case of scarlet-fever, and there should be no hesitation in stopping the business if necessary. At the same time it seems that boiling the milk removes all danger.

Diphtheria.—Although the term "diphtheria" was only introduced by Bretonneau in 1826, epidemics of malignant sore throat destroying life by suffocation, attacking children rather than adults, and sometimes leaving paralytic sequelæ, have been described from the earliest times. According to Hirsch, there is a tendency to cyclical epidemic or pandemic waves, lasting often for many years, and followed by long intervals of comparative quiescence. Although no climate gives immunity, the tropics suffer less than cold and temperate climates. During last century France was one of the principal centres for diphtheria. In England localised epidemics occurred in 1815-25, after which the country was almost free from the disease until 1857, when it appeared as part of a general prevalence affecting almost the whole of Europe. The mortality increased during the next two years, reaching 0.52 per thousand living, in 1859. Since that time the disease has held its ground in England, and at the beginning of the last decade of last century showed a maximum prevalence and fatality, for in London in 1893 the death-rate was 0.75 per 1,000, though in the decennium 1891-1900 it was 0.50. Its tendency, therefore, recently has been that of a steady decline since 1893. In England and Wales in 1906 the death-rate from diphtheria was 0.17 and in rural districts 0.16 per 1,000.

Season.—The curve of seasonal prevalence of diphtheria shows a relation to cold, the maximum mortality being reached in November or December, and the minimum in the summer. In New York the curve is similar. Records of epidemic prevalence show the same preference for the cold months (winter and spring), and many are associated with cold and wet weather. There are, however, many exceptions, in some of which heat has been regarded by local observers as inducing, and cold as arresting, the epidemic. In Great Britain atmospheric humidity is considered most favourable to the prevalence of diphtheria, but

malignant epidemics have occurred in excessively dry weather in California, and Newsholme finds that the relation between diphtheria prevalence and years or localities of heavy rainfall is inverse rather than direct.

Diphtheria occurs upon high and low levels, and upon dry and moist soils; but Kelly observed a greater average prevalence upon damp soils, and this relation seems to hold good in many other parts of the country. A special incidence of diphtheria

in damp houses has often been noted.

Locality.—Diphtheria was formerly, upon the average, more prevalent in rural than in urban districts; but in recent years it has shown a marked tendency to establish itself in towns, particularly affecting young children. Longstaff compared the statistics (1855-1880) of the various registration counties in respect of their average density of population and average mortality from diphtheria, with the following result:—

Mean Diph-	Dense districts	Medium districts (1 to 2 acres per person).	Sparse districts			
theria	(less than 1 acre		(over 2 acres per			
Mortality.	per person).		person).			
1855-60	0·123 per 1,000	0.182 per 1,000	0.248 per 1,000			
1861-70	0·163 ,,	0.164 ,,	0.223 ",			
1871-80	0·114 ,,	0.125 ,,	0.132 ",			

The peculiarity of the distribution of diphtheria in regard to population suggests that its causes "should not be sought for primarily in any high development of civilisation, such as sewers, but rather in some condition associated with a more primitive form of life" (Longstaff). A disease closely resembling diphtheria in many respects has often been observed to attack cats, pigeons, fowls, and horses during the prevalence

of human diphtheria. This coincidence suggests that some part of the incidence upon rural districts may be due to infection from lower animals. Some of the symptoms of diphtheria have been produced in animals by inoculating them with diphtheritic material.

Diphtheria frequently occurs in a purely sporadic form, no clue to any source of infection being discoverable, but all cases, whatever their origin, are highly infectious. It may be regarded as endemic in certain localities, sporadic cases bridging over the intervals between more or less frequent epidemics.

Air.—Several otherwise inexplicable outbreaks in secluded positions have been attributed to transmission of infection by wind for a distance of some miles. There is no à priori improbability in this explanation, but it seems clear that no such wind-convection can occur in those outbreaks that remain limited to certain streets or even houses in urban districts.

Water has been suspected of conveying infection, but no complete demonstration has been made out.

Milk, on the contrary, has been abundantly proved to convey the diphtheria poison (see p. 133). Many extensive outbreaks have been traced to this agency, and careful inquiry has brought to light a close relation in such cases between the quantity of milk consumed and the danger of attack. The milk may become infected by some person suffering from a slight attack of diphtheria, but we have now to bear in mind the possible occurrence of bovine diphtheria, and of infection from the cow itself.

Drainage effluvia.—Diphtheria is more severe and more prevalent among persons exposed to foul gases from sewers or drains. The relation is, however, by no means direct. Houses with the worst forms of

drainage defects may remain free from diphtheria for indefinite periods, and many cases of diphtheria occur in premises in which no defect can be found. The same may be said of other insanitary conditions, notably dampness of house, accumulations of manure and other filth in proximity to houses. All these ought probably to be regarded as predisposing causes

only.

Age and sex.—The mortality is greater among females than among males at all ages between three and forty-five years, and hence at all ages taken together. In both sexes the mortality increases from infancy to a maximum in the fourth year of life, and declines steadily thereafter, but there is a somewhat greater mortality (in proportion to the comparatively scanty surviving population) after the fiftieth year. Of the whole number of deaths ascribed to diphtheria, 53 per cent. occur at ages under five years, and 82

per cent. under ten years.

Incubation.—"The most common period of incubation in diphtheria is from two to three days: less frequently the period is diminished to 24 hours or extended to four days; and the limit between an almost immediate operation of the infection and a delay of six or seven days practically covers the period of incubation for preventive purposes" (Thorne Thorne). The early symptoms are often insidious, but as a rule the membrane is visible within a day or two of the onset, if not at first. In non-fatal cases the disease usually runs its course in two or three weeks. The case mortality varies greatly in different epidemics, and there is little constancy even in the same epidemic. It is usually high in well-marked outbreaks: in pre-antitoxin days it was frequently 30 per cent., or more. It is now generally between 10 and 20. The protection conferred by an attack of diphtheria does not appear to be very great.

Gresswell raised an important question in regard to those persons who have suffered from diphtheria in former years, and in whom acute inflammation of the tonsils is set up by change of weather or trifling exposure. He observed that among those associated with them at such times, diphtheria is apt to occur without discoverable cause; and suggested that the chronic morbid condition may really be a latent form of diphtheria, the occasional recrudescence of which becomes the

starting-point for the spread of infection.

Slight cases, in which the symptoms are scarcely recognisable, are extremely difficult of diagnosis. During prevalence of diphtheria any sore throat should be regarded with suspicion, even if no membrane is visible and the constitutional symptoms are wanting. There is no necessary relation between the severity of attack in two cases of which one is infected from the other; the slightest case may impart fatal infection. In doubtful cases, where no membrane can be found, some assistance may be given by the knowledge of the existence of other cases of diphtheria, and paralytic sequelæ may ultimately clear up an obscure diagnosis. The breath and sputa are infectious from the first. Clothing, furniture, and even walls and floors, may retain the virus tenaciously.

Schools afford, as Power observed in 1876, a favourable ground for the spread of diphtheria. There children at the most susceptible ages are brought together for some hours daily in close contact, often in a moist and impure atmosphere, with scanty ventilation; not unfrequently such children use the same articles, towels, slates, pencils, etc., and are brought into intimate contact with each other. Shirley Murphy has pointed out that a marked increase in the number of cases of diphtheria in children of school age followed on compulsory elementary education, and that corresponding with school holidays there is a temporary reduction in the number of notified diphtheria attacks at school ages. Nor is the danger limited to the mere spread of infection, for, as Thorne Thorne showed, a progressive development of virulence and infectiousness is brought about by the aggregation of slight (possibly

unrecognisable) cases among susceptible children at school. Many instances are recorded, especially in districts where throat affections are more or less endemic, commencing with the appearance of cases of sore throat, followed presently by cases of diphtheria, which increase in numbers and severity. The epidemic is usually arrested by closure of the school, often only to break out again, with renewed virulence, as soon as the

school is reopened.

Among preventive measures, the first are isolation and disinfection. Doubtful cases should be secluded and no children from infected households should be allowed to attend school. Supervision of "contacts" should be kept for a fortnight, and if necessary prophylactic inoculation of antitoxin should be undertaken (500 units). Closure of schools is advisable if many cases have occurred among the scholars, or if diphtheria is prevalent in the district, and especially if the school is the means of bringing children from infected villages into contact with others. Milk supplies that are found to be implicated in the spread of infection must be stopped at once, and most rigid precautions should be taken in regard to households connected with the milk trade. Lastly, insanitary conditions should be sought for and remedied.

Bacteriology. — It is now generally held that diphtheria is caused by a bacillus discovered by Klebs in 1883 and isolated by Löffler in the following year. It is a slender rod, straight or slightly curved, and beaded; there are also irregular and club-shaped forms. It differs in size according to its culture medium, but is generally 3 or 4μ in length. Cobbett and Graham-Smith recognise five morphological types of diphtheria bacilli on young serum cultures :-(1) Oval bacilli, with an unstained septum; (2) long, faintly-stained, irregularly-beaded bacilli; (3) long, regularly-beaded bacilli—"streptococcal" forms;

(4) segmented bacilli; and (5) uniformly-stained bacilli. All these forms are longer than the pseudodiphtheria bacilli, more curved, and generally with clubbed ends. In the membrane which is its strictly local habitat in the body-indeed, with the exception of the secretions of the pharynx and larynx, and occasionally in lymph-glands and the spleen, the bacillus is found practically nowhere else in the body -it sometimes shows parallel grouping, lies on the surface of the exudation, and is separated from the mucous membrane by the fibrin. The bacillus possesses at least four negative characters: it has no spores, and no power of motility; it does not liquefy gelatine, nor does it produce gas in cultures. It is stained with Löffler's methylene blue, and shows metachromatic granules and polar staining. Its favourable temperature is blood-heat, though it will grow at room temperature. It is aërobic. Löffler's medium is made by mixing three parts of ox-blood serum with one part of broth containing 1 per cent. of glucose, 1 per cent. of peptone, and 1 per cent. of common salt; the whole is coagulated. Upon this medium the Klebs-Löffler bacillus grows rapidly in 18-20 hours, producing scattered round, white colonies, becoming yellowish. The bacillus grows well in broth, but without producing either a pellicle or turbidity; it can grow on the ordinary media, though its growth on potato is not readily visible; on the white of egg it flourishes extremely well. In a moist condition, apart from the body, the bacillus has a low degree of resistance, but when dry can survive for long periods. The bacillus is secured from the throat by examination of a piece of the membrane or by a swab. Three examinations should be made before concluding that the throat is clear. About 20 per cent. of all cases of diphtheria offer no bacteriological evidence of infection. The bacillus may

exist in the throat of infected persons for long periods, even several months, causing such persons, though apparently healthy, to be infectious. It may also be present in the throats of persons who have not suffered from the disease—"carriers"—and this fact Houbtless furnishes the explanation of some sporadic cases and obscure outbreaks.

A pseudo-diphtheria bacillus described by Hofmann has somewhat similar characters to the true Klebs-Löffler bacillus, and is frequently found in healthy throats, especially in children of the poorer classes. It differs from the true diphtheria bacillus in the following ways: (a) it is thicker in the middle than at the poles; (b) forms no acid in glucose culture media; (c) does not give Neisser's reaction; (d) differs in artificial culture; (e) does not agglutinate like IB. diphtheriæ, and (f) possesses no pathogenic action. It is probably innocuous, but in practice it is the safest course to look with suspicion upon any throat marbouring pseudo-diphtheria bacilli.*

Antitoxin treatment of diphtheria.—Diphtheria untitoxin is prepared by taking a month-old culture of the Klebs-Löffler bacillus. This is filtered and the toxin inoculated into the horse in gradually increasing doses until a large degree of acquired

^{*} There are a number of other bacilli from which the true diphtheria acillus must be differentiated. These include the B. coryzæ segmentosus, B. zerosis, and a number of "diphtheroid" bacilli, and organisms from asal and aural discharge. There are, as summarised by Gordon, five thief characters by which the true diphtheria bacillus may be known:— (a) The macroscopic and microscopic appearance of the growth on blood The macroscopic and microscopic appearance of the growth on blood erum; (b) the behaviour of the bacillus io Löffler's blue, Gram's stain, and Neisser's stain for granules; (c) the reaction to litmus of a culture in likaline broth, containing 2 per cent. of dextrose after 48 hours at 37° C.; (l) the pathogenic te-t—1 c.c. of broth culture, 48 hours' growth at 37° L, injected subcutan ously into 200-300 gramme guinea-pig, produces eath generally in 48 hours, whilst post-mortem hæmorrhagic necrosis and edema are found locally, the internal organs are congested, the pleural, ericardial, and peritoneal fluids are increased, and the suprarenal apsules are enlarged and engorged with bood; (e) the virulence of the reganism or its toxin is completely neutralised by a simultaneous dose of lightheria antitoxin. For purposes of rapid diagnosis, (a) and (b) are smally accepted. sually accepted.

tolerance has been produced. After injecting, say, 500 c.c. without ill-effect, and the horse having rested for a few days, blood is drawn from the jugular vein into sterilised flasks, in which it coagulates. The separated serum, which averages about half the total blood drawn, is filtered, and tested as to strength of antitoxin value, or as it is termed, "standardised," by observation of its power of immunising a guineapig of certain weight against a minimum lethal dose of toxin. It is now ready for therapeutic use, and the patient is injected subcutaneously between the crest of the ilium and the last rib. Strict asepsis must be observed, and the dose should be as large as practicable (2,000-5,000 units). Antitoxin is now very widely used in the treatment of diphtheria, and is increasingly used as a prophylactic in "contact" persons (300-500 units).

The following summary of the Antitoxin treatment of all forms of diphtheria at the hospitals of the Metropolitan Asylums Board, 1895-1906, compared with the results obtained before the adoption of that treatment, affords striking evidence of the efficacy of diphtheria antitoxin:—

This is discred	Cases.	Deaths.	Mortality, per cent.
1890-3 (before antitoxin)	7111	2161	30:39
1894 (antitoxin occasionally used)	3042	902	29.65
1895 ,	2182	615	28.1
1896	2764	717	25.9
1897	4381	896	20.4
1898	5186	906	17.5
1899	7038	1082	15.4
1900	7271	936	12.9
1901	6499	817	12.6
1902	6015	714	11.8
1903	4839	493	10.1
1904	4070	444	10.9
1905	3734	335	9.0
1906	4149	432	10.4

Its value is particularly noticeable among children. Amongst cases in the first year of life the rate has fallen from 1.8 to 37.8; in the second year from 63.1 to 35.4; in the third year from 55.1 to 26.4; in the fourth year from 48.3 to 22.9; and in the fifth year from 39.6 to 20.7.

At the Brook Hospital, MacCombie has kept records showing the results of the antitoxin treatment, with special reference to the day of the disease on which treatment began in order to

Illustrate the effect of early administration :-

MORTALITY PER CENT, IN CASES TREATED.										
bine our one baid	1897	1898	1899	1900	1901	1902	1903	1904	1905	1906
Cases treated on 1st day ,, ,, 2nd ,, ,, 3rd ,, ,, 4th ,, ,, 5th day and after.	5·4 11·5 19·0	5.0 14.3 18.1	3·8 12·2 20·0	3·6 6·7 14·9	4·1 14·9 12·4	4:6 10:5 19:8	4·2 17·6 16·0	10.6	3·4 6·3 11·0	3·9 8·7 18·5

It is now customary to discharge diphtheria atients from hospital only after two consecutive acteriological examinations of the fauces have roved negative. In the hospitals of the Metrocolitan Asylums Board this results in an average ength of residence in hospital of 50 days.

Croup is credited with an annual death-rate of between 1 and 0.5 per 1,000 living. There can be no doubt that many eaths from diphtheria and laryngitis have been attributed to roup, and if, as some authorities still maintain, there is a true croup "—that is, a non-infectious membranous laryngitis due ainly to wet and cold, and not followed by paralysis—it is of as frequent occurrence than the death returns would imply. The death-rate from croup is declining, and the decline is proably due to improved diagnosis. Outbreaks of diphtheria are commonly attended with cases of so-called "croup," which con investigation prove to be diphtheritic.

The seasonal curve of deaths attributed rightly or wrongly croup has its maximum in the cold months from December to arch. It is much later than that of diphtheria, and approxiates more closely to the laryngitis curve. In the distribution

as regards age and sex a further difference from diphtheria is noticeable. Croup is more fatal to males than females upon the whole, the mortality among females being lower up to fifteen years of age. The highest mortality in both sexes occurs in the second year of life. In all these respects "croup" is allied to laryngitis rather than diphtheria.

Erysipelas occurs in all parts of the world, but more in temperate regions than elsewhere (Hirsch). In North America it has repeatedly taken the form of widespread and malignant epidemics, and in England and elsewhere virulent outbreaks often originated in former years in hospitals, especially in surgical wards, where cleanliness, ventilation, airspace, drainage, and other sanitary essentials were neglected. At the present time outbreaks of this kind are rare, and erysipelas is chiefly met with in sporadic form. In its seasonal curve, and also in its local attachments and periods of recurrence, it is closely allied with scarlet-fever and diphtheria, and still more closely with puerperal fever. It attacks males more than females, and kills more in early infancy than at any other period of life, although the mortality increases again after twenty-five years of age. After inoculation the incubation is usually from one to three days. It is not certain whether the disease is ever acquired except through some breach of continuity of skin or mucous membrane, but the occurrence of so-called "idiopathic" (non-traumatic) erysipelas points to this. The pathogenic microbe is a streptococcus, readily cultivated, and reproducing the disease upon subsequent inoculation.

Relapsing fever, "famine fever," or "bilious typhoid," was first recognised clearly in Ireland in the eighteenth century, and still has its principal focus there, although epidemics are not unfrequent in Scotland, and have occurred in England, Northern Europe, the Levant, India, and several eastern countries. It occurs often in conjunction with typhus and is apparently closely related to it in etiology. Epidemics of the two diseases frequently coincide, or one follows the other closely, or sporadic cases of the one are observed during prevalence of the other. Neither climate nor weather has been found to influence it in any marked degree, nor has it any apparent relation to soil. Overcrowding and want of food are powerful

bredisposing causes, especially the latter, which Murhison thought was capable of originating the disease,
but many epidemics have occurred without noteworthy
brivation. The wealthier classes are almost exempt,
and females are less susceptible than males. Under
avourable conditions it appears in districts that have
long been free from it, and in many such cases no
history of imported infection can be made out. It
has been suggested that the specific agent may be

conveyed by parasites, such as pediculi.

The disease is highly infectious, but appears to be capable of transport through the air for short distances only, though it readily attaches itself to fomites. A specific spirillum was discovered by Obermeier in the blood of patients during the febrile stage only, and inoculation with blood containing spirilla has been found to reproduce the disease in men and monkeys. It is possible that there is some relation between Obermeier's spirillum and the trypanosomes recently found in sleeping sickness. The protection afforded by an attack is slight. Only about 5 per cent. of the cases terminate fatally, but convalescence is slow.

Typhus has only comparatively recently been distinguished from enteric fever, but the two diseases are widely different in their etiology and epidemiology.

Typhus epidemics have been recorded in considerable numbers for several centuries past, more especially in times of war and famine, and were common in the seventeenth century and up to 1815. The only great epidemic in Europe since that time was in 1846-7. Ireland has been from earliest times one of the chief centres of the disease. Russia being another. Although for the most part limited to cold and temperate climates, typhus is by no means unknown in many hot countries, but usually at considerable elevation. The different habits of life, and especially the abundant ventilation, which prevail in warm countries may help to account for their comparative immunity, but as the effect of season remains marked, there is probably some further influence at work. In former times, epidemics of typhus frequently reached England, appar-

ently by invasion from Ireland; but for many years past typhus has almost entirely disappeared from the southern and midland towns of England, although still occurring in the large towns of the north and of Scotland, especially those containing a large Irish population. The last general epidemic in Ireland was in 1862-4.

Cases of typhus are most numerous in winter and spring, and in warm countries epidemics seldom occur at any other season. In countries in which typhus is common, however, the origin and course of epidemics are not checked by either extreme of temperature.

Telluric conditions are not known to have any influence.

The great predisposing causes of typhus are overcrowding, want of ventilation, filth, debility, and privation. Wars and famines in former centuries were frequently followed by outbreaks of fever, among which typhus was prominent. "Gaol fever" (typhus) was common in English prisons up to the last century, owing to their filthy and overcrowded state. Better ventilation of vessels has rendered "ship fever" almost unknown, but accidental repetition of the old conditions, even in recent years, has led to fresh outbreaks of typhus. Typhus broke out among the British troops in the Crimea in 1854-5, and among the besieged in Metz in 1870. In Ireland the years of famine and distress have been attended with typhus.

It is not clear that any race or class is especially susceptible, or the reverse, apart from the conditions under which they live; but for the reasons already stated, the poorest classes suffer by far the most

from typhus.

Age and sex.—The mortality from typhus increases from childhood to about fifty years of age, and then declines somewhat. It is greater among males, except, perhaps, in childhood. Handford finds that the susceptibility is greatest in the ten to fifteen years' ageperiod, but that the case mortality, i.e. the proportion of deaths to attacks, is at its minimum in this period,

and rapidly increases at later ages. Spear has called attention to the mildness of the symptoms in children, owing to which the initial attacks in an outbreak are

liable to escape diagnosis.

The incubation period seems to be very variable, and, according to Murchison, may be only a few hours or as long as three weeks or more. It has been observed to be short in certain cases after exposure to the poison in a concentrated form. Perhaps twelve days may be taken as the most usual duration. The rash appears within a week after the onset, and, as a rule, on the fourth or fifth day. The disease generally terminates in death or convalescence about the end of the second week, and isolation may be terminated in ten days or a fortnight more. The mortality averages about 20 per cent., but varies with the age, condition, and surroundings of the patient, and with the epidemic.

Infection is given off by the breath, probably by the exhalations from the skin and mucous memberanes, and possibly by the excreta. Some have suggested that the disease is spread by vermin, especially fleas. According to Murchison, the infectivity is more active after the first week. The air immediately surrounding the patient appears to be infectious, but its virulence is rapidly lost by diffusion—a result that may be attributed to dilution, or desiccation. The poison clings readily and strongly to clothing and other fomites, but has not been shown to be conveyed by water, or by milk or other food. Doctors and nurses in typhus wards rarely escape for long unless protected by a

Typhus is probably acquired by inhalation of the poison, whether from contact with the patient, breathing the air in his immediate vicinity, or by the intervention of fomites. The powerful predisposing influence

previous attack. Second attacks are rare.

of overcrowding and want of ventilation may be explained by the facilities which they give for infection and for concentration of the poison, as well as their effect in rendering the individual more susceptible.

No pathogenic microbe is yet known, but it may safely be inferred from analogy that, without the presence of a specific poison, no intensity of the favouring conditions can originate an outbreak of typhus. It is, nevertheless, true that the disease often makes its appearance suddenly among overcrowded and filthy communities—on shipboard, for example—in instances where no previous case can be traced.

Typhus is essentially a preventible disease. Isolation of the patient, disinfection, and quarantine of persons exposed to infection will rarely fail to arrest a localised outbreak. Cleanliness and ventilation are

auxiliaries of the first importance.

Enteric or typhoid fever was differentiated from typhus in 1850 by Jenner. It is a disease of practically world-wide distribution, but is less prevalent in tropical regions than elsewhere. In England it is decreasing in prevalence, and now contributes not more than 0·1 to the average general death-rate. For instance, the death-rate from this disease in 1906 in England and Wales was 0·09, which shows a marked decline on previous years. In 1871-80 it was 0·32, in the decade following 0·19, and in 1891-1900, 0·17. In London in 1906 it was 0·06.

Season.—In England the greatest prevalence occurs in October, and the minimum in May or June. The autumnal maximum is very general, but the exact position of the summit of the curve varies in different countries; in New York it occurs in September.

Weather.—High or low temperature of the air has in itself no clear relation to enteric prevalence, nor

has the rainfall, but instances are recorded in which hot and dry summers were followed by high mortality from the disease, upon the occurrence of rain, and the reverse after cold and wet summers. It is probable, in view of the frequent relations between telluric conditions and enteric fever, that meteorological con-Hitions act mainly by modifying the temperature and moisture of the soil, and that rainfall may either increase or diminish the chances of an outbreak according to the previous condition of the ground.

Soil .- Pollution of the earth by animal matter, especially if excrementitious in origin, is conducive to the endemic or epidemic prevalence of enteric fever. Pettenkofer and Buhl have traced a connection beween the movements of the ground water and the occurrence of enteric fever sickness and mortality in Munich, which has been confirmed by further obervation in Berlin and other parts of Germany, and Isewhere. "The total of the cases of sickness and leath from typhoid falls with the rise of the subsoil water, and rises with the fall of it; . . . the level eached by the disease is not in proportion, however, the then level of the subsoil water, but only to the variation in it on each occasion; or, in other words, is not the high or low level of the subsoil water that decisive, but only the range of fluctuation" (Hirsch). This relation is so constant as to leave no room for coubt as to its reality and significance in the localities n question, but it does not appear to hold good England, and Fodor has found exactly the reverse n Pesth. In many places where enteric fever ccurs, the subsoil water is so deep and its movement b) trifling that there is little probability of its proucing any material effect. In Munich the soil is prous, the ground-water high, and, until recently, aking cesspools are said to have been universal. As uchanan has pointed out, the purity of water in

wells is liable to be affected by changes in the level of the ground-water, and the readiness with which enteric fever is spread by means of specifically polluted water must not be lost sight of.

Air does not appear to carry the poison of enteric fever under ordinary conditions even for the shortest distances, and the chances of infection by mere proximity to a patient are slight, though occasionally instances suggestive of this are met with. Infection from the sick to the healthy by direct or indirect contact is, of course, one of the commonest means of spreading the disease; water and milk being others.

Water.—While water polluted even with excremental filth has often been drunk for years by numbers of people with perfect impunity so far as the appearance of enteric fever is concerned, contamination with the excreta of a case of enteric fever has over and over again been found to result in widespread outbreaks of the disease, the incidence of which in such instances exactly follows the distribution of the water, and affects only those persons who drink it. Such outbreaks rise suddenly to considerable proportions, but decline gradually. Corfield collected particulars of more than 50 waterborne epidemics of typhoid between 1864 and 1902. Some have arisen from specific pollution of shallow wells, as at Guildford in 1867, in the Uxbridge Rural District in 1882, at Hitchin in 1883, at Beverley in 1884, etc.; others from the pollution of deep wells. Thus an outbreak of enteric fever at Redhill and Caterham in 1879 was traced by Thorne Thorne to pollution of one of the well-adits at the Caterham Waterworks by the excreta of a workman suffering from a mild attack of enteric fever, the organic impurity of the water, as shown by analysis, being extremely small. Another case of infection of a deep well was at Houghton-le-Spring, in

Durham, in 1889. The large outbreak at Worthing in 1893 (1,315 cases) was due to the consumption of. water from a new heading, which was contaminated by sewage. The water-borne poison is not depen-Hent upon the organic matter in the water, but upon the specific pollution, which need not necessarily be

large in quantity.

Innumerable other instances are on record in which direct and obvious excretal contamination of wells, springs, streams, cisterns, reservoirs or watermains has been followed by enteric outbreaks upon a Marge or small scale, according to the number of persons drinking the water. As illustrations may be mentioned the outbreaks at Caius College, Cambridge, in 1873, at King's Lynn and Maidstone in 1897, and at Lincoln in 1905. Buchanan found that in the case of Caius College, the water in the main at Tree Court, which was chiefly affected, was contaminated by suction into it, during the periods of intermission of supply, of foul air from a soil pipe, through a "weeping pipe" which supplied the trap on the waste-pipe of the safe-tray of a water-closet, the trap being liable to siphonage. The King's Lynn outbreaks (1892 and 1897) were traced to pollution of the sources of the water supply, namely, the river Gaywood, with typhoid excreta, and the same applies to the great outbreak at Maidstone (1,847 cases) in 11897, which was believed to have been caused by pollution of the springs by excremental matter from colony of hop-pickers. In both these cases the filtration of the water as delivered was inefficient. The outbreak at Lincoln in 1905 included some 900 cases, and commenced towards the end of January. The river Witham and its tributaries, from which the supply was taken, were polluted "by drainage from land highly cultivated, and occasionally nanured with night soil." The effluent from a

sewage farm discharged into the river above the intake. In the autumn of 1904, and in January, 1905, there were a number of cases of enteric fever in Lincoln, and, therefore, opportunity for the typhoid virus to gain access to the water supply, which, owing to increased acceleration due to heavy demands, was not effectually filtered in the filter-beds. The result was a water-borne outbreak of enteric fever, chiefly affecting households which did not boil their drinking water. Many confirmatory observations have been made of places in which enteric fever, once endemic, has become rare coincidently with the substitution of a pure for an impure water supply. Thus at Millbank Prison, artesian well water was substituted for Thames water in 1854, and the previous endemic prevalence of enteric fever disappeared.

In several of the cases mentioned above, the proof of specific pollution was incomplete, and it has been held that sewage-polluted water may acquire the property of causing enteric fever in those who drink it, without the access of enteric excreta. Even in country districts, where the movements, identity, and health of every person coming into the locality are easily traced, it has often been found impossible to obtain evidence of specific pollution of water that was nevertheless instrumental in spreading enteric fever. On the other hand, it is clear that something beyond simple excremental pollution is necessary to account for the sudden acquisition of morbific properties in a water that has long been polluted; and, moreover, attacks of enteric fever may be so mild as to escape recognition, nor is it safe to disregard the possibility that the specific poison may gain access in ways as yet unsuspected.

Shirley Murphy has found distinct increase in notified enteric attacks in London following floods

above the water-intakes, and he has further pointed out that disease carried by water or milk may affect so few persons, or be in so mild a form, as to escape

recognition.

It is necessary to remember that the urine of the patient as well as the bowel discharges may contain the infective bacillus. Horton Smith has shown that the urine is infective in one in four cases, and most frequently towards the end of the disease and during convalescence. In one recorded case the typhoid bacillus was found in the urine eight months after convalescence. Other observers have found bacilluria of this kind present in between 10 and 30 per cent. of the cases examined.

Food.—Milk was first recognised as a medium for the dissemination of enteric fever in 1857. Many instances have since been recorded, in England and other countries, the infection being sometimes traced to the use of polluted water for washing the cans or diluting the milk, sometimes to the milk Ibeing infected more or less directly by a person suffering from enteric fever (see page 132). Oysters lhave on many occasions recently been suspected of conveying the infection of enteric fever. In 1894 Newsholme reported a number of such cases, and in the same year an outbreak of the disease occurred at a college in Connecticut, traced to coysters. In this country, between 1894 and 1902, increasing evidence was forthcoming in confirmation of this channel of infection, and in the latter year occurred the well-known outbreaks at Winchester and Southampton, traced to the consumption of polluted coysters at mayoral banquets (Bulstrode). It is now well understood that oysters fattened on sewage contaminated beds may cause disease in persons eating them (see page 120). Fried fish (Hamer), ice-cream, watercress, and other foods have also carried enteric infection. When the disease is thus contracted it is characterised by multiple cases in houses, and an

exceptional age-incidence.

Drain effluvia, sewer gases, and emanations from manure heaps and other filth accumulations, are often regarded as directly causing enteric fever. That they are powerful predisposing causes is beyond doubt, and it may be supposed that persons exposed to their influence become more susceptible to the enteric poison. Alessi has shown experimentally that this is true of rats, rabbits and guinea-pigs, after exposure to the effluvia from cesspools or excreta. The incidence of attack is heaviest upon households or districts in which drainage defects, or insanitary privies and earth closets, exist, although it is often difficult to dissociate their influence from that of impurity of soil and water. Buchanan has shown that in the large English towns improved drainage has greatly lowered the death-rate from enteric fever. In towns that have been newly sewered, without due attention to the disconnection and trapping of house drains, enteric fever has sometimes become far more prevalent in large houses upon high ground, tenanted by the wealthier classes, than in the poorer districts down below, in which very commonly there were no drain-inlets within the houses. Here, however, the presence of enteric excreta in the sewage may be suspected. Precisely the same conditions are recognised, and with equal justice, to be conducive to outbreaks of diphtheria, a totally distinct disease, and in either case the presence of a specific poison is doubtless essential to the manifestation of the specific disease, be its mode of access what it may. Grossly insanitary conditions may exist for years without any outbreak of either disease. Men working in sewers do not appear to be specially liable to enteric fever, nor do those employed in scavenging in pail-closet towns, although

they come into close and frequent contact with enteric excreta.

Locality. — Enteric fever is more prevalent and more fatal, at all ages, in towns than in the country, and often fixes persistently upon one district, where the predisposing conditions mentioned elsewhere will usually be found to exist. In such districts newcomers are especially liable to attack; this has been observed on a large scale as regards soldiers newly carrived in India. It seems probable that some sort of acclimatisation is at work, relative immunity being conferred by repeated minimal infection. As a rule, but by no means always, the areas in which enteric fever is endemic are occupied by the poorest class, among whom insanitary conditions abound, and llittle care is taken for preventive measures.

Age.—According to Handford, "the susceptibility tto enteric fever decreases with age from the earliest years onwards, though the decrease is slight between fifteen and twenty-five, whereas the risk of a fatal ttermination steadily increases with age." The Registtrar-General's figures, however, tend to show that the fatal incidence is less during the first five years of life than between five and ten years. The registered mortality is low in infancy, high in the second to the fifteenth year, and then rises further to a maximum lbetween 15 and 25 years, after which it falls permanently. It is probable that true enteric fever is rare among infants and young children, and that many of the cases recorded at those ages are due to faulty diagnosis.

Sex.—More males than females die from enteric fever, but the mortality is greater among females from the fourth to the twentieth year of age. The susceptibility, according to the Registrar-General, is greater among males from about the fifth to about the twenty-fifth year, and greater among females at

all other ages. The average severity of attack is somewhat greater among females; 18.8 per cent. of female cases end fatally, and 17.1 of male cases.

The most frequent mode of infection is by contact or by the mouth, the virus either being contained in water, milk, or other food, or conveyed more directly by the unwashed hands after contact with infected matter. That the excreta and urine are infective is certain, and it is possible that infection is sometimes carried by the breath. Enteric patients are sometimes treated in hospitals side by side with other cases; but with scrupulous cleanliness, especially as regards the nurse's hands, and any clothing that may have become soiled in however minute degree with the patient's discharges, it seems that, although nurses are not unfrequently attacked, the infection has little tendency to spread to other patients. It is otherwise in crowded homes, and there can be no question that direct personal contagion has been the cause of not a few of the smaller outbreaks of the disease, and of many secondary cases in the larger epidemics (e.g. in the Lincoln outbreak of 1905).

The incubation period is variable, from two or three days to three or even four weeks, perhaps most usually twelve to fourteen days. The insidious mode of onset renders exact determinations difficult. The incubation is often short when the poison is introduced by water or milk. In some of Budd's cases it was only two or three days, at Guildford ten or eleven days, in the Caterham outbreak fourteen days. Jancken records an instance in which a body of soldiers, not otherwise exposed to infection, marched through two villages in which enteric fever was rife, and drank freely of water. Two days later 3 were attacked by enteric fever; then 7, 6, 0, 4, 5 in successive days, and 7 more within the fortnight—32 in all. The onset was sudden, but the course mild. Another section of the

ttroops passed through the same villages without

drinking, and escaped illness.

The duration of illness is usually about three to four weeks, but may be protracted by relapses. Second attacks are not uncommon.

The greatest care is needed to prevent all risk of contamination of milk supplies or water. The disposal of the excreta is a matter of great importance. They should be disinfected and disposed of at once. In the country it is customary to bury them away from Thouses or sources of water supply; in towns they are usually dealt with like other excreta, but where pail-closets are used enteric excreta may be separately collected and cremated. The urine, as we have seen, often contains the bacilli (in about 20 per cent. of

cases).

Bacteriology.—In 1880-81 Eberth announced the discovery of the typhoid bacillus in cases of clinical enteric fever. In 1884 it was first cultivated outside the body by Gaffky. Since then other organisms have been held responsible for the causation of the disease. In 1885 the B. coli communis was recognised, and it was a matter of debate among bacteriologists as to how far these two organisms were the same species, and interchangeable. Bacteriologists now generally regard the Eberth-Gaffky bacillus as the specific cause of typhoid fever, though complete proof is still wanting. About 3 per cent. of typhoid cases are due to paratyphoid organisms (Boycott). Typhoid bacilli appear as rods, 2-4 μ long, 5 μ broad, having round ends. Sometimes threads are observable, being 10 μ in length. The bacilli differ in length from each other, but are approximately of the same thickness. Round and oval cells constantly occur even in pure culture, and many of these shorter forms appear to be identical in morphology with some of the many forms of B. coli. There are no spores. Motility is marked, due to some five to twenty flagella of varying length. The flagella are both terminal and lateral,

and are elastic and wavy.

The organism may be isolated from the ulcerated Peyer's patches in the intestine, from the spleen, the mesenteric glands, and the urine. Owing to the mixture of bacteria found elsewhere, it is generally most readily isolated from the spleen. For this purpose the whole spleen is removed, and a portion of its capsule seared with a hot iron to destroy superficial organisms. With a sterilised knife a small cut is made into the substance of the organ, and by means of a sterilised platinum wire a little of the pulp is removed and traced over the surface of agar, on which growth occurs in 24 hours at 37° C. The bacillus grows on all the ordinary media. It is decolorised by Gram's method. The typhoid bacillus is capable of saprophytic existence in soil, dust, water, and milk, and may persist in the excreta for months. It can survive in ordinary earth for two months, on sterilised linen for 60 days, on woollen cloth for 80 days, in sterilised water 196 days, in certain soils 400 days. Wright has prepared a vaccine from a virulent 24 hours' culture in broth. It is killed by heating at 60° C. for five minutes, and is injected hypodermically in the flank.

The typhoid bacillus agglutinates with serum from a typhoid patient, and this has been used as a bacteriological aid in diagnosis. It is known as Widal's application of Gruber's reaction, and is carried out in the following way:—A few drops of blood are drawn from the lobe of the ear or finger of the patient; the serum, which exudes as the blood coagulates, is then diluted with about 9 parts of neutral bouillon. A drop of this dilution is mixed with an equal quantity of a typhoid broth cultivation of 18 to 24 hours' growth made from a virulent culture. Under the microscope it may be seen whether the reaction is positive or negative. If in the mixture of serum and of culture the bacilli become immotile and also become agglutinated, or grouped together in colonies, the

reaction is positive. If, however, they remain, as they do in nealthy blood serum, motile and scattered, the reaction is negative. The Widal reaction is obtainable in about 70 to 80 per cent. of the cases between the 4th and 14th day of the ase and in 90 per cent. of the cases during the third and ourth weeks.

Simple continued fever is still assigned as the ause of a considerable number of deaths, but is steadily losing ground year by year with improving diagnosis and certification. Many of the deaths so returned are probably due to enteric ever, others to septic conditions, pneumonia, etc. Longstaff has shown that the seasonal mortality curve for "simple coninued fever" is very different from that of enteric fever, and resembles the typhus curve rather closely, having its maximum in winter and spring.

Epidemic diarrhœa* is returned as the cause of a death-rate that steadily averaged 0.9 per 1,000 rom 1850 to 1880, and in 1906 was, for England and Wales, 0.87 per 1,000. In towns it is much higher 11.0 per 1,000) than in rural districts (0.52). In London, in 1906, it was 0.94, and 0.81 for ten prereding years. In many cases diarrhea is merely ymptomatic, but the existence of an epidemic disease f which diarrhea is the prominent manifestation is now ecognised. Much light has been thrown on the subject by Ballard, Newsholme, and other observers.

Density of population.—Epidemic diarrhea is sssentially a disease of towns, or, in more general erms, of crowded areas. The diarrhea death-rate in 891-1900 among children 0-5 years of age was 5.4; in urban counties it was 6.7 and in rural districts 2.6. The disease is ordinarily twice as fatal in urban as in tural counties and more dangerous to boys than girls.

^{*} Under this term are included all forms of zymotic diarrhea and bidemic enteritis. The Royal College of Physicians (1900) laid down that gastro-enteritis," "muco-enteritis," "gastric catarrh," and such monyms of epidemic diarrhoea in medical certificates of death should be scarded. The College regarded epidemic diarrhœa as a general disease a specific character in the same sense as enteric or other fevers, and herefore authorised the use of the terms "epidemic enteritis" or, if pre-prred, "zymotic enteritis" or "epidemic diarrhea," to designate the sease in its various forms and degrees.

In counties such as Dorset and Wilts, the mortality (among children 0-5 years) in 1903 was below 2 per 1,000, whereas in Warwickshire it was 8.8, in the East Riding 8.9, and in Lancashire 9.13.

An analysis of the diarrhea mortality in the various counties according to their average density of population gives a similar result. Thus, in 1871-80 the mean annual diarrhea mortality per 1,000 at ages under five years was

In 6	counties	with m						per	person	2.6
,, 9	"	,		,, :				,,		3.6
,, 12	"	,	,				3 ,,	"	"	4.2
,, 11	counties	-:41 1	,	,,,	1,	, 2	,,	"	"	4.8
,, 6	England	with I	less	than	1	1	acre	"	,,	7.1
111 1111	England an ave	erage of	ares,	WITH	1	1	·5 acre	,,	"	5.6

In 1898-1902 the rate was 11.5 in urban districts and 5.0 in rural. Urban conditions are favourable to the disease, chiefly because of some or all of the following:—

Density of buildings upon an area increases the tendency to diarrhea mortality, in addition to the density of population that usually, though not always, accompanies it. It is chiefly a disease of urban life.

Elevation of site tends to reduce diarrhæa mortality, but only in the same degree that it affects infant

mortality from all causes.

Lack of ventilation and light is conducive to diarrhea mortality. Among the common conditions that are harmful in this way are narrow dark courts and streets, obstructive walls or buildings, back-to-back houses, overcrowding, and neglect of ventilation of rooms.

Lack of cleanliness has a similar effect, and is usually found in association with the above, particularly in houses. But further, towns which have adopted the water-carriage system of sewerage have less

diarrhea than those which retain other methods of removal of excrement. Towns also which have the most effectual scavenging arrangements, including the methods of removal of house refuse, have the least epidemic diarrhea.

Foul air from sewers, cesspools, and filth accumulations of any kind promotes diarrhea mortality. Smoke and mere chemical effluvia seem to be inoperative.

Drinking-water may cause outbreaks of diarrhoa, independently of season, but is not responsible for

ordinary epidemics of summer diarrhea.

Soil.—Diarrhea mortality is low in places built upon solid rock, high where the soil is loose and poorous. Sand and deep mould are the worst, clay is fiar better. Organic pollution of the soil, whether wegetable or animal in origin, is a most potent fiactor, and occurs especially upon such sites as "made ground," pervious soils, town refuse, market gardens, or in soil polluted by leakage from drains or cesspools. Dampness of soil, accompanied by aëration, is the most favourable condition for diarrhea, dryness or saturation being alike preventive.

Temperature.—High temperature of the air has long been observed to be associated with high liarrheal mortality, and the reverse with low air temperature, but Ballard shows that the relation is madirect. The maximum mortality by no means necessarily coincides with the highest readings of the hir-thermometer. The temperature of the soil, being less sensitive indicator and more steady in its record,

ss on the whole a safer guide.

Ballard concluded that-

[&]quot;(a) The summer rise of diarrhoeal mortality does not comnience until the mean temperature recorded by the 4-foot earth-thermometer has attained somewhere about 56° F., no matter what may have been the temperature previously attained by the atmosphere or recorded by the 1-foot earth-thermometer.

"(b) The maximum diarrhoeal mortality of the year is usually attained in the week in which the temperature recorded by the 4-foot earth-thermometer attains its mean weekly maximum.

"(c) The decline of the diarrhoeal mortality . . . coincides with the decline of the temperature recorded by the 4-foot earth-thermometer, which temperature declines much more slowly than the atmospheric temperature or than that recorded

by the 1-foot earth-thermometer.

"(d) The influence of the atmospheric temperature, and of the temperature of the more superficial layers of the earth, . . . is little, if at all, apparent until the temperature of the 4-foot earth-thermometer has risen as stated above; then their influence is apparent, but it is a subsidiary one."

Tomkins found that epidemic diarrhea did not occur in Leicester until the temperature of the 1-foot thermometer reached 60° F.

Rainfall is operative by its cleansing effect upon the atmosphere, by laying dust, and by its reducing effect upon the temperature of the soil and the atmosphere. Diarrheal mortality is greater in dry and less in wet seasons. Deficiency of rainfall plus high atmospheric temperature are probably the two chief external conditions favourable to epidemic diarrhea. Wind tends to reduce diarrhea mortality, calm in the diarrheal season promotes it.

It was upon broad facts such as these that Ballard,

in 1887, based his "provisional hypothesis":—

That the essential cause of diarrhea resides ordinarily in the superficial layers of the earth, where it is intimately associated with the life-processes of some micro-organism not yet isolated.

That the vital manifestations of such organism are dependent, amongst other things, perhaps, principally upon conditions of season and on the presence of

dead organic matter, which is its pabulum.

That on occasion such micro-organism is capable of getting abroad from its primary habitat, the earth, and having become air-borne obtains opportunity for

astening on non-living organic material (especially ood whether inside or outside the body), and of using nch organic material both as nidus and pabulum in ndergoing various phases in its life-history.

That from food and from organic matter in certain bils it can manufacture a virulent chemical poison, which is the material cause of epidemic diarrhea.

Social position.—By far the greatest incidence is pon the poorest class, the artisan and lower labour-

ng classes.

Food, and particularly milk, may "ferment" or ecome contaminated by exposure to dust or filth manations, and thus cause diarrhea. Breast-fed hildren are remarkably exempt, and those partially reast-fed come next; the mortality is much higher mong children artificially fed, and especially if fed y the bottle. On the whole there is evidence to show nat infantile diarrheea is usually a form of food coisoning.

Maternal neglect.—The mortality among illegitimate children is higher than that of legitimate mildren from all causes, but the excess is greatest in egard to diarrhœa. The difference is most marked n years of low epidemicity, and tends to disappear n years of high diarrhea mortality, but in the latter me average age of attack is earlier among illegitimate children. Occupation of mothers from home contributes to neglect and improper, irregular, and

rtificial feeding of infants.

Age.—Diarrhœa is fatal at both extremes of life. The first year of life, especially from the third to the inth month, has by far the greatest mortality. In 871-80, 63 per cent. of the deaths attributed to iarrhœa were at ages under one year, 80 per cent. nder two years. It was by far the most fatal of the nfective diseases in infancy, and caused a mortality f over 25 per thousand births. In 1881-90 the

mean mortality rate showed further decline, and in 1891-1900 73 per cent. of the deaths from this disease were under one year and 86 under two. From infancy the mortality diminishes until about the 20th year, after which it again increases until the end of life. The liability to attack seems to be greater in the second year than the first, and at all events is far greater in the first two years than in the third or later years. It is comparatively small in the first three months, and probably increases up to the end of the first or beginning of the second year. No age is exempt from attack, but only severe and acute attacks kill at ages between infancy and old age. Tomkins states that "infants and young children form only a small proportion of those attacked, although they furnish nearly the whole of the deaths."

Sex.—The mortality is greater among females from the third to about the 45th year, but greater among males in infancy and old age. The liability to attack, however, is greater among males at all

ages.

Season.—Fatal diarrhœa occurs at all seasons, but always increases greatly in summer. In London the mortality curve shows a slight rise throughout June, rapidly increasing in July, and reaching its maximum in the first week of August, after which it again falls steadily throughout September and October. During the rest of the year there is little variation. An outburst of epidemic diarrhœa occurs every year, but the date and intensity of the epidemic vary considerably from year to year, and according to locality. The relation to temperature and other atmospheric and telluric conditions have already been discussed. There is very frequently an alternation of years of high and low diarrheal mortality, but a simple explanation of this is usually to be found in the meteorological records of the summer months, there

being coincident alternation of hot, dry summers with thers which are cool and wet.

Locality.—Diarrhœa is endemic in many large English towns, notably Leicester and Preston, and auses heavy mortality annually. Others, such as Bristol, Derby, and Halifax, are affected much less, hough all, as in the Metropolis, show the summer

iise every year.

Previous health affects the chances of recovery more than the liability to attack. The incubation is apparently very short, from a few hours to a day, or at most two days. Half the fatal cases terminate within a week, and the course is more rapid in the ater periods of an epidemic (Ballard). Characteristic oathological changes are found in the kidneys as well s in the intestines, a fact that goes far to prove the specific character of epidemic diarrhea. Pneumonia is common and fatty degeneration of the liver almost invariable.

In many instances diarrhea has appeared to be nighly infectious by means of the excreta, but this is

not always the case.

Outbreaks of diarrhea occur from time to time, especially in hospitals and other public institutions, which cannot be referred to climatic conditions, but upon investigation are traced to articles of food, and especially to water and milk. Water may acquire the power of causing diarrhea in many ways, including the presence of suspended mineral matter, such as clay or mica (Parkes); excess of dissolved mineral matter; suspended or dissolved sewage matter or other unimal impurity, especially if undergoing putrefaction; suspended vegetable matter; absorption of fætid gases, including sulphuretted hydrogen and "sewer gas." Milk is a frequent cause of diarrhea, owing either to fermentative changes in itself, or to contamination by specific poison or effluvia. An outbreak at Moorfields Hospital was traced to butter. Putrid food of any kind tends to cause diarrhea, and fish is especially prone to be injurious if not fresh. Lastly, specific diseases must be mentioned in this connection, as for example the Welbeck cases described in an earlier

chapter.

Bacteriology.—In 1885 Escherich published his researches on Bacillus coli communis, and since that date many authorities have maintained that this organism is the causal agent of epidemic diarrhea. Booker, on the other hand, as the result of investigation, concluded that many different varieties of bacteria are concerned in the etiology of the summer diarrhœas of children. He considered "streptococcus enteritidis" and Proteus vulgaris the most important. In 1895 Klein isolated an organism from the evacuations of patients suffering from an epidemic of diarrhœa at St. Bartholomew's Hospital, which he named B. enteritidis sporogenes,* and which he believes to be the cause of epidemic diarrhea. . B. enteritidis sporogenes is widely distributed, and occurs in normal and typhoid excreta, in sewage, manure, soil, dust, and milk. It has been frequently found in the dejecta of patients suffering from epidemic diarrhæa. It fulfils in a somewhat exceptional degree the requirements suggested by Ballard. But the same is true of B. coli, and the question is therefore sub judice.

Influenza stands almost alone in its purely epidemic or pandemic occurrence, with little trace

^{*} B. enteritidis sporogenes (Klein) is an anaërobic bacillus: $1^{\circ}6-4^{\circ}8$ μ long, and $0^{\circ}8$ μ broad; stains by Gram's method and ordinary stains. Motile; spore formation present; large oval spores often situated near one end of bacillus; grows well on gelatine and agar. In the former gas is produced and the gelatine liquefies. It grows well in milk. After thirty-six hours of anaërobic incubation at 37° C. the surface is covered with stringy, pinkish-white masses of coagulated casein enclosing a number of gas-bubbles. The main portion of the tube of milk contains a colourless thin, watery whey, with a few casein lumps here and there adhering to the sides of the tube. The whey has a smell of butyric acid, and is acid in reaction. It contains many bacilli.

of endemic localisation. Much confusion has arisen rom the prevalent misapplication of the term nfluenza to severe catarrhs, infectious or otherwise. Many epidemics of true influenza have been observed during the last few centuries, spreading over a greater or less portion of the globe. Although there is clear indication of preference for lines of traffic, the progress of an epidemic is highly irregular. It appears simultaneously at widely separated points, often ceaping over or avoiding entire countries, perhaps to appear in them later. It may be limited to particular towns or villages, sometimes even to a small section of the population in one part of a town. The advance is frequently rapid, but not more rapid than human traffic, and at other times its progress is llow and halting. The pandemic of 1889 was first neard of in Siberia, in May, and had spread to almost every part of Europe before the end of the year. eached London late in December, and in the course of the next three months diffused itself over the whole of England.

The epidemic quickly reaches its maximum limits in a given centre, owing probably to its infectiousness and short incubation, and the susceptibility of the greater part of the population. It seldom remains conger than a few weeks in any locality, but may return in the course of the same pandemic. The interval between epidemics is irregular. Parsons states that major epidemics occurred in England in 1803, 1833, 837-8, 1847-8, and minor epidemics about every three ears. From 1860 onwards these practically ceased, ntil 1889, when a new series of epidemics began, ach covering almost the whole country, at intervals f about a year, and which in ten years destroyed no ewer than 87,767 lives in England and Wales. In 891-1900 the influenza mortality was 361 per million. During the decennium there were two epidemics, in 1895 when the death rate rose to 423 per million, and

in 1898-1900 when it reached 504 per million.

In the same district the type of the epidemic remains fairly constant as regards intensity, severity, and perhaps the tendency to one or other group of local symptoms or complications. There is, however, considerable variety in different centres during the progress of an epidemic, and the type may change in a given locality.

Climate.—Epidemics are more common in hot countries, but no country, however cold, is exempt

from invasion.

Season and weather.—Neither cold nor heat, nor any known meteorological condition, has been shown to have any effect in checking or promoting the spread of the disease. Intense outbreaks occur in the hottest and coldest seasons, in good, bad, and changeable weather, apparently indifferently. Each type of weather in turn has been alleged to cause epidemic prevalence, upon the strength of narrow observation confined frequently to one locality and one outbreak. Nothing is definitely known of any relation to atmospheric electricity, or the presence of ozone. It is, however, conceivable that conditions of climate, season, and weather may affect the severity of the epidemic, and especially the mortality from the respiratory complications that usually attend it.

Telluric conditions are apparently without influence. The disease attacks persons of all ages and both sexes, sometimes to the extent of quarter or even half the entire population. In general both incidence and severity of attack are less at ages below 20 years, and less among women than among men; though in some few localities and instances children, and especially school children, suffered more than adults. It would appear that influenza, like diphtheria, may become intensified by the aggregation of susceptible persons in a confined atmosphere. Whether it can spread through the air for long distances is doubtful. Many instances are on record in which

was apparently conveyed in that way to ships and ght-houses, but Parsons has shown that in some of mese the isolation was not complete. On the whole, secluded ersons and populations suffered least, the incidence being exceptionally heavy upon postal and railway officials, and there whose work brought them most into contact with the public.

Dogs and cats apparently suffer from influenza, and horses are liable to a severe and often fatal disease (termed "pink-ye" from one of its prominent symptoms), which has been egarded as a form of influenza. On several occasions in England and elsewhere epidemics of influenza have been preceded by outbreaks of "pink-eye." But in a Report to the L. G.B. (1893) Klein gives reasons for doubting the transmissibility of influenza to lower animals.

The period of incubation appears to be short, from one to three days. The onset is sudden, with chills, elevation of temperature, and severe pain in the eyes, mead, and back, and tenderness of muscles, especially of the legs and back. Three principal forms of the elisease are described, with affection respectively of the respiratory or alimentary tracts, or of the nervous system alone. Intense prostration and depression and rapid loss of weight are almost invariable. The symptoms usually abate in three or four days, but complications and sequelæ may protract the duration of illness indefinitely. The breath is in all probability infectious from the first.

The mortality from influenza is usually slight, except among persons already weakened by disease or predisposed to bronchitis or pneumonia. The increased death-rate which follows the appearance of influenza in a district is largely due to deaths attributed to respiratory diseases. The protection, if any, conferred by an attack is slight and evanescent, repeated

attacks being common.

In 1892 Pfeiffer discovered a small bacillus in the bronchial mucus of patients suffering from influenza. It is aërobic, non-motile and non-resistant. It does

not stain by Gram's method, and grows feebly on artificial media.

The ordinary preventive measures—notification, isolation, and disinfection—have not yet been em-

ployed on a large scale against influenza.

Cerebro-spinal fever.—Cerebro-spinal fever, known also as epidemic cerebro-spinal meningitis or spotted fever, is not rare in this country, and has recently caused serious mortality in New York and on the continent of Europe. In 1866-7 a severe epidemic in Ireland was marked by high mortality

and hæmorrhagic rashes.

The disease may be described as "an acute, epidemic disease, characterised by profound disturbance of the central nervous system, indicated at the onset chiefly by shivering, intense headache or vertigo, or both, and persistent vomiting; subsequently by delirium, often violent, alternating with somnolence or a state of apathy or stupor; an acutely painful condition with spasm—sometimes tetanoid—of certain groups of muscles, especially the posterior muscles of the neck, occasioning retraction of the head; and an increased sensitiveness of the surface of the body" (Radcliffe). Throughout the disease there is marked depression of the vital powers, and not unfrequently collapse. In its course an eruption of vesicles, petechial or purpuric spots, or mottling of the skin, sometimes not unlike the rash of typhus, is apt to occur. If the disease tend to recovery, the symptoms gradually subside without any critical phenomena, and convalescence is protracted; if to a fatal termination, death is almost invariably preceded by coma. After death the membranes of the brain and spinal cord are found engorged with blood, and an effusion of sero-purulent matter is often present in the meshes of the pia mater and beneath the arachnoid.

The clinical manifestations include the presence

of Kernig's sign* and of tache cérébrale.†

There are also milder or anomalous forms which render identification difficult, and which lead to its being mistaken for other ailments of more common occurrence. In certain localised outbreaks in the eastern counties in 1890 this disease was generally mistaken for sunstroke or enteric fever; in Northamptonshire in 1890-91 it was diagnosed as pneumonia or sore-throat; and in Irthlingborough in 1905 many of the persons attacked were regarded as suffering from influenza. When death ensues rapidly the disease has been mistaken for typhus, tetanus, or malignant measles.

Cerebro-spinal fluid, withdrawn from the lower part of the spinal canal by lumbar puncture, should be examined for the presence of the "Diplococcus meningitidis intracellularis" of Weichselbaum, now generally regarded as the specific cause of the

disease.

Whether cerebro-spinal fever is spread by direct infection from person to person is matter of uncertainty, but the isolation of the sick from the healthy should be practised and disinfection carried out.

Pneumonia occasionally occurs in an epidemic form, and in many instances has shown distinctly infectious properties without attaining epidemic proportions. Epidemic pneumonia is, with few exceptions, of the "croupous" or "fibrinous" form, but it is not certain that all cases of sporadic croupous pneumonia are due to infection, or that infectious pneumonia is

^{*} To obtain this sign the patient is placed so that the hip-joint is semi-flexed, and while the hip-joint is retained in this position, the knee-joint is extended passively. A contraction of the hamstrings then sets in, which causes resistance to the extension.

[†] A congested streak produced by drawing the finger-nail gently over the patient's thigh, abdomen, or face. In cases of meningitis this appears earlier (within 30 seconds), lasts longer (8, 10, or 15 minutes), and is broader and of deeper colour than in healthy persons.

always caused by the same virus. Infectious pneumonia is very often limited to the upper lobe, and often accompanied by pleurisy or empyema. Gastric symptoms, diarrhea, and jaundice are common, and prostration and cerebral symptoms are frequently intense, out of proportion to the physical signs of pneumonia. The latter may be delayed for days after the onset. The mortality is often high, and usually so in outbreaks limited to a few persons. Epidemics have been described in considerable numbers in England and various other parts of Europe during the last two centuries. Although more frequent in temperate climates, many are recorded in the West Indies, Mexico, Peru, and India. Nearly all occur in winter or spring, and the seasonal curve of epidemic prevalence coincides pretty closely with that of pneumonia mortality, which has its maximum in December, and is high from November to April. Hence the prevalence of pneumonia-epidemic or otherwise -- is associated with the colder months, and a closer analysis shows that in each climate the greatest prevalence of pneumonia occurs at the season of most rapid and sudden changes of temperature, be it winter or spring (Hirsch), and there is evidence tending to show that the prevalence varies in some measure with the intensity of the changes of temperature. Epidemics occurring at unusual seasons have often been associated with unusual meteorological conditions of the same kind.

The popular belief in "chill" as an exciting cause of pneumonia cannot be entertained as regards infectious pneumonia, in spite of the strength of the evidence which connects pneumonia outbreaks with the weather conditions in which "chill" is believed to be most liable to occur. Exposure to sudden changes of temperature or extreme cold may increase the activity of infection or the susceptibility of the individual, but nothing more. If, however, there is a non-infectious and non-specific croupous pneumonia, the question of "chill" may be

rargued upon different lines. It is said that croupous pneumonia has been produced by injecting irritants into the lungs of dogs, by mechanical injury to lung tissue, by experimentally injuring parts of the nervous system, and by various other mechanical means, which may be admitted as establishing some sort of prima facie possibility of like effect from sudden change of temperature, especially as the whole mortality attributed to pneumonia (and not merely that part which is already recognised as due to an infectious form of the disease) reaches its maximum at times of greatest liability to "chill." On the other hand, attempts to produce pneumonia by exposing animals to sudden alternations of temperature have uniformly ended in failure, and even traumatic pneumonia may conceivably be dependent upon the co-operation of specific microbes accidentally gaining access to the injured tissue.

There are now on record many instances of outlbreaks of pneumonia which remain limited to a single lhousehold or a small circle, but in which nevertheless the evidence points strongly to infection.

Nothing conclusive has been established as regards the influence of rainfall or telluric conditions upon pneumonia, although it has been asserted that absence of rain (dry cold) and low level of subsoil water are favourable conditions.

Males are far more liable to pneumonia than females, but the attacks are usually less severe and fatal. In the Middlesbrough epidemic of 1888 Ballard found that the female case mortality only exceeded the male at ages above sixty-five years. Both the liability to attack and the average case mortality increase greatly as age advances. In child-hood and old age the mortality is highest, and urban death-rates are usually in excess of rural.

All depressing conditions predispose to attack during epidemic prevalence, among them fatigue, anxiety, poverty, and debility from any cause. Insanitary conditions, especially filth, overcrowding, and want of ventilation, act apparently as powerful but not indispensable predisposing causes. Effluvia from

graveyards have also been held responsible for outbreaks. Repeated outbreaks have sometimes been observed in the same buildings, especially barracks and prisons. In many extensive epidemics it has been found that those streets or houses suffered most which were in the worst sanitary condition, and outbreaks in barracks, prisons, and ships have usually been attributed to the same causes. Such conditions must not be regarded as more than predisposing causes in any case, and in many instances—notably in several of the small outbreaks of infectious pneumonia limited to one household—they are conspicuously absent. Negroes are especially susceptible to pneumonia, even in their native climate.

Friedländer's oval, capsulated pneumo-bacillus is found in large numbers in the affected lungs, and in the blood and sputa in croupous pneumonia (but not in acute catarrhal pneumonia or septic pneumonias). It was found in abundance in the substance of the floors and ceilings of rooms in a prison at Amberg where repeated outbreaks of pneumonia had occurred for many years (Emmerich). Pneumonia has been produced in rabbits by the injection of cultivations; on the other hand these results have been disputed, and the true pathogenic microbe asserted to be Fränkel's small, oval, capsulated diplococcus, which is non-motile, non-liquefying and aërobic. It is difficult to cultivate, and rapidly loses virulence. It is stained by Gram's method. It is by far the most frequently present organism in croupous pneumonia, and may be present in other diseases, suppuration, otitis media, ulcerative endocarditis, and meningitis.

The widespread and fatal epidemic of pneumonia that occurred at Middlesbrough in 1888 was investigated by Ballard. Out of 1,633 cases in a population of 97,000, 369 ended fatally, the case mortality being

409

11 per cent. The poorer classes suffered more than ne wealthy, and cases were exceptionally numerous and severe in the workhouse, where the drainage was ery faulty. The workhouse children suffered six mes, but adults only one and a half times, as much ss the corresponding class outside. Exposure and atigue seem to have acted as predisposing causes, and many apparent instances were recorded of direct inection from contact with a sick person. The progress ff the epidemic seemed to be arrested by heavy rains, and to be most rapid during rainless periods. Klein bund neither Friedländer's nor Fränkel's organisms, tut large numbers of short bacilli, which he samed Bacillus pneumoniæ. Inoculation of human ung-juice or of cultivations of the bacillus into mice caused an acute disease, the chief and constant lesion of which was pneumonia; further moculations from such mice imparted the same lisease to other mice. Samples of bacon were purhased in the infected districts, and it was found that ff mice fed upon this bacon a large proportion became III, with the same symptoms as those mentioned above. The Bacillus pneumonia was recoverable by cultivation from their tissues, and by inoculation the disease could be transferred to other mice. Whether the acon had or had not become infected by human cases ff pneumonia is not clear, but it may be suspected hat the disease was capable of being spread by means if infected food.

The incubation appears to be short, frequently bout five to seven days. The onset is sudden, and sually marked by rigors and severe constitutional ymptoms, the signs of pneumonia being often delayed or three or four days. The fifth day is often fatal; and, on the other hand, crisis often occurs on or bout the fifth, seventh, or ninth day, but sometimes he course, even in non-fatal cases, is severe and protracted, the symptoms being of a "typhoid" character.

Both the breath and the sputa may be assumed to be infective.

The mortality varies considerably, but is usually high, especially in elderly persons and in the outbreaks limited to a narrow circle. In some epidemics,

however, it has been as low as 5 per cent.

Epidemics of pneumonia often occur coincidently with outbreaks of other diseases, and especially enteric fever. Pneumonia is a common complication or sequela of enteric fever, and it has been suggested that an invasion of the system by the enteric poison may have its sole manifestation in pneumonia, the

usual intestinal lesions being slight or absent ..

Tuberculosis is a specific and infective disease common to man and the lower animals. The principal form in which it affects man is pulmonary tuberculosis or phthisis, but all parts of the body are liable to be invaded. The mortality due to tuberculosis is enormous, but does not admit of exact statement, owing to imperfections of diagnosis, nomenclature, and classification. Pulmonary phthisis may be accepted as a fairly well-defined division, although it undoubtedly includes a certain proportion of non-tubercular destructive diseases of the lungs. Tubercular meningitis is not sufficiently distinguished in death-returns from other diseases that are attended with somewhat similar symptoms, and "tabes mesenterica," which is classed as a tubercular disease, includes a large annual number of deaths among children from wasting diseases, of which no exact diagnosis has been made.

Phthisis, "consumption," or pulmonary tuberculosis is largely dependent upon known and remediable conditions, and improved hygiene has led to a steady decline in mortality from this tal diseases, especially during adult life, and as cently as the decennial period 1891-1900 it was no recorded cause of one-twelfth of the whole cortality in England, and of about thirty per cent. If the deaths at ages between twenty and fortywe.

Its geographical distribution is almost world-wide. Weither hot nor cold climates are exempt, but huhidity, especially if the daily range of temperature is igh, is frequently associated with prevalence of hthisis. Cold, and especially Arctic countries, suffer omparatively little as a rule, and the exceptions re mostly explicable by social conditions incolving overcrowding and want of ventilation. Other things being equal, elevated and mountainous egions are less affected than lowlands, owing, it is elieved, to the dryness and purity of the air and oil, and the fuller and deeper respiratory movements. A general relation between dampness of soil and prevalence of phthisis has been affirmed by Buchanan* in England and Bowditch in America. Donsiderable and early reduction in phthisis mortality has, in many instances, followed artificial drainage of damp localities. In Ely the reduction was 47 poer cent., and in Salisbury 49 per cent., but at Ashby-de-la-Zouch, Alnwick, and Carlisle, Buchanan found an increase coincident with drainage of the towns, and the same was observed at Danzig and

The mortality statistics of the decennium 1851-60, for Surrey, Kent, and Sussex, showed that there was less phthisis among populations living on pervious soil than among those living on impervious soils; less on high-lying pervious soils than on low-lying pervious soils; less on sloping impervious soils than on flat impervious soils. Buchanan also found a general agreement in phthisis mortality between districts that have common geological and topographical features of a nature to affect the water-holding quality of the soil—an agreement not observed in districts differently circumstanced in such respects; and further, a general parallelism between the degree of wetness of soil and the degree of phthisis incidence.

elsewhere. In several of these cases, however, the local conditions were such that the drainage works did not materially dry the soil. It is stated by many authorities that phthisis does not occur in marshy regions, and Kelly found that the phthisis mortality is

greater upon the drier soil.

Towns suffer more than rural districts, and there is a close relation between density of population and phthisis mortality, due to the co-operation of several causes, chief among which are overcrowding, poverty, and stagnation and impurity of air. The heaviest incidence is upon the poorer classes, and especially those living in narrow streets, alleys and courts, and in back-to-back houses. Conditions inside the houses such as overcrowding and want of ventilation are even more potent, and repeated experience in barracks, workhouses, prisons, and other public institutions, as well as in ships, has shown that with improved ventilation and ampler air-space the mortality from phthisis is greatly reduced. Among nomadic races the disease is rare.

Many trades have a tendency to induce phthisis, and especially those industries which are carried on in overcrowded, hot, moist workrooms, or which charge the air with mineral or organic dust of an irritating kind. Exposure to rapid alternations of temperature is very injurious. Greenhow attached much importance to a stooping or cramped posture

at work, as predisposing to phthisis.

Among other predisposing conditions, intemperance and debilitating causes of all kinds are important. Brouardel considers that "alcoholism is the most potent factor in propagating tuberculosis," and he quotes Baudran as showing that the deaths from tuberculosis per 10,000 rise in proportion to the annual consumption of litres of alcohol per head. In this country the Registrar-General has shown that more

rounts in London result from phthisis; more than tee-third of the deaths of public-house servants, free, and potmen in London are due to phthisis; death the mortality from this disease among London ankeepers and their servants is more than twice as that among all occupied males, and the cortality amongst the servants alone is more than tree times as great. No doubt some of this mortality

due to inhalation of tuberculous dust from ried sputum. Then there is the lowering effect of revious disease. Attacks of pleurisy, bronchitis, or neumonia increase the liability to phthisis, and so, to, do many forms of specific fevers (notably measles, thooping cough, and enteric fever), and other diseases, are example diabetes and insanity. The low phthisis cortality among fishermen, agricultural labourers, and others following essentially out-door employments, seems to indicate that exposure to weather as been overrated as a predisposing cause, for this pondition in a great measure implies abundance of tesh air.

Rightly or wrongly, a large but decreasing number if deaths of young children under five years of age is tetributed to phthisis. The annual mortality per million living at ages under five years was 973 in 8861-70, 770 in 1871-80, 535 in 1881-90, and 413 in 1891-1900. From five to ten years the mortality at a minimum, and then rises steadily to its maximum at ages between 35 and 45, after which it teclines again as age advances.

Generally speaking, the decline since 1861 has been due to improved sanitation and nutrition, combined with increased knowledge as to infection and the influence of institutional and sanatorium segregation.

The influence of sex is very marked, but is to be explained for the most part by differences of surround-

ings rather than by variation in natural susceptibility. In proportion to their numbers males suffer a higher phthisis mortality than females at all ages except between 5 and 25 years. While the mortality is decreasing rapidly in both sexes, the decline among females is greater than that among males.

PHTHISIS MORTALITY IN ENGLAND AND WALES, 1851-1900 (PER 1,000 LIVING).

Decennia.				Males.	Females.	Total.	
1851-60	101	BUI	4	2.6	2.8	2.7	
1861-70			To.	2.5	2.5	2.5	
1871-80				2.2	2.0	2.1	
1881-90			THE STATE OF	1.8	1.6	1.7	
1891-1900	1 400			1.5	1.2	1.3	

In London also there has been a steady decline. In 1906 the phthisis death-rate was 1.44 per 1,000, having fallen in the four preceding years. In 1906, when the phthisis death-rate was 1.44 for the whole of London, it was as low as 0.67 in Hampstead and as high as 2.01 in Shoreditch, 2.06 in Southwark, 2.18 in Finsbury, and 2.37 in Holborn.

No race is exempt, but among Jews phthisis appears to be rare, and this comparative immunity has been partly attributed to the care that is taken in the selection of animals slaughtered for food. Natives of hot countries migrating to cold, damp

climates are very prone to phthisis.

The tendency to phthisis may be inherited, in the form of constitutional weakness, or of a narrow, contracted chest. Whether heredity can do more than strongly predispose to the disease is very doubtful. Tubercle has in rare instances been found in children within a few days of birth, and even in the fœtus. Various observers have obtained evidence of parental phthisis in from 15 to 30 per cent. of the cases coming

inder their notice, but this proportion is scarcely afficient to establish the proof of heredity. A family istory of phthisis is obtainable in 50 per cent. of

ospital cases of this nature.

Deaths from phthisis are most frequent in spring March and April), and least so in autumn (September and October). The seasonal curve of mortality is merefore later than in the ordinary respiratory isseases, but it serves to indicate seasonal conditions occelerating death rather than those primarily induc-

ng a disease of long and uncertain course.

Bacteriology.—The pathogenic microbe of tuberulosis, discovered by Koch in 1882, is a motionless, Mender, beaded bacillus which, under suitable conitions, forms spores. The bacilli are very slow in growth and multiplication, taking one, two, or even Three weeks to form a colony on artificial media, and They require for that purpose a somewhat high temperature, which must further be maintained within narrow limits, about 37° to 39° C. A peculiarity that listinguishes them from all other bacilli except those of leprosy and syphilis is that after staining with fuchsin they are not decolorised by nitric acid (Ziehl-Neelsen method). They are therefore termed acid-fast. There are in addition to the tubercle bacillus a group of acid-fast bacilli allied to the streptothrix family. They possess, however, characters which differentiate them from the tubercle-bacillus. These latter bacilli are found in all tuberculous lesions, both in man and the lower animals. They have been found also in the blood, and in certain secretions (milk), and they abound in phthisical sputa. Cultivations if inoculated or injected into susceptible animals reproduce the disease. The bacilli are readily destroyed by many chemical or thermic means, and by prolonged exposure to sunlight, but the spores are tenacious of vitality.

Tuberculosis can be acquired by inoculation, inhalation, or swallowing. Inoculation is comparatively rare, and usually causes a local lesion only. Inhalation would seem to be by far the most common source of The bacilli, or rather their spores, are found in air-borne dust, especially in rooms inhabited by phthisical persons. The daily sputa of a single patient may contain many millions of bacilli, and drying for months will not destroy their virulence. The prevalence of the pulmonary form of tuberculosis, and its close relation to air-conditions, are not without significance, as pointing to air-borne infection. A further analogy to the ordinary infectious diseases is to be found in the strong evidence which has been adduced by Ransome and others to show that tuberculosis attaches itself to particular small localities ("Tuberculous Infective Areas"), houses, and even rooms. The bacilli have been found not only in the air and dust, but in the walls of rooms occupied by phthisical persons. Coates investigated this point in houses in Manchester, in 1900, with the following results:-

(1) Dirty houses in which a consumptive patient is living who takes no precautions to dispose of his expectoration, but spits freely upon the floor and into his pocket handkerchief: In 66.6 per cent. of these houses virulent tubercle bacilli were found, showing the large amount of dangerous infective material

present in an infected house.

(2) Clean houses in which a patient is living who is not sufficiently careful as to the disposal of his sputa: In 50 per cent. of these instances the bacillus was found. It is evident that ordinary household cleanliness alone is insufficient to prevent the accumulation of infective material in rooms occupied by a consumptive.

(3) Very dirty houses in which there had been no case of consumption for some years: In this class of

ouse no tubercle bacilli were present, showing that irulent dust found in classes 1 and 2 must have been

ue to the presence of the consumptive patient.

Niven examined the records of 5,000 deaths from. his cause in Oldham, and found that whereas the mathematical probability was that 68 houses would e invaded twice, and 7.6 three times, the actual tumbers were 274 and 24 respectively. Many ther workers have confirmed this. The endemic rreas occur chiefly in the worst parts of towns, when the predisposing conditions already referred to exist in maximum intensity. Whenever tuberculous sputum ss allowed to dry the risks are therefore great that the Hust so produced may be inhaled in a virulent form, and lodging at one or more points may set up varying Hegrees of tuberculosis. This broad fact is based upon experimental and clinical evidence. Tubercucosis has been produced experimentally in animals in this way, and there is clinically the overwhelming requency of tuberculosis of the lungs among men exposed to this form of infection. But Koch, Flügge, and others have shown that not only is sputum a source of infection when dried and pulverised, but also when disseminated by coughing, shouting, etc., in the form of minute moist particles of spray.* Koch exposed rabbits, guinea-pigs, rats, and mice to an infected spray for half an hour on three successive days, and produced tuberculosis in every nimal. Heymann found that such spray particles from human beings inoculated into guinea-pigs profluced tuberculosis. Most of the droplets are large and settle rapidly, but some may remain suspended in the air for more than an hour, retaining, of course.

^{*} Mervyn Gordon in this country has confirmed these results by applying a streptococcus test for saliva (in which secretion 10 million streptococci per c.c. may commonly occur). He finds that during loud speaking bacteria may be projected by means of salivary spray a distance of 40 ft.

their virulent properties. Heymann found the duration of life of the bacilli in these droplets was eighteen days in the dark, and three days when exposed to light. Under ordinary circumstances and an absence of draughts, the zone of danger from a coughing consumptive extends to a distance of about three feet. It must be remembered that the tubercle bacilli in the moist particles of "cough-spray" are probably of higher virulence than those in dried sputum dust, and therefore it seems reasonable to suppose that the cough-spray is one of the most ready modes of infection. The degree of infectivity of phthisis is not, however, a very high one. It is a true infective

disease, but only a sub-infectious one.

There is still much doubt as to the frequency of direct infection from person to person, but none as to its occasional occurrence, or as to the fact that almost all persons must frequently inhale and swallow living tubercle-bacilli. Fortunately the microbe is very slow in its development, and exacting in its requirements as to temperature and surroundings, so that the vast majority that gain entrance to the respiratory tract are expelled or perish. The long duration of the disease, the wide and general diffusion of the virus, the paramount importance of predisposing conditions, and the difficulty of infection in their absence, all combine to render obscure the time and source of infection. Statistics are inconclusive even in regard to transmission of phthisis between husband and wife.

We have previously referred to the conveyance of tuberculosis by milk and meat (see page 106). The risk of contracting the disease by consuming tuberculous milk and meat is undoubted, but the disease does not appear to be widely spread by these means. As regards milk, all danger of infection may be obviated by boiling, and prevented by using milk

om tubercle-free herds only (tested by tuberculin, hich is a glycerine extract of dried residue of young altures of the tubercle bacillus). The cooking to hich meat is subjected is frequently insufficient to estroy bacilli that may be present in the deeper arts.

The heavy mortality among children from "tabes esenterica" (tubercular peritonitis) and tubercular eningitis is urged as a confirmation of the extreme revalence of tubercular disease, but it is probable at both of these terms are employed in the loosest ay in making out certificates, and it would not be fe to place much reliance upon the tubercular ature of the majority of the cases so certified. he former disease (tubercular peritonitis) has een attributed to the consumption by children f tuberculous milk. Still, however, has shown that ne commonest channel of infection in tuberculosis f children is through the lung, and that infection hough the intestine, as would occur if it were milkorne, is less common in infancy than in later childood. There is, however, substantial evidence nat tuberculous milk can and does set up some form f tuberculosis (bovine or human) in the bodies of rersons consuming it.

Tuberculosis often ends in recovery. The records f long series of autopsies of persons who have died f other diseases show that traces of cured phthisis re found in a proportion variously estimated as 5 per cent. and upwards. It is not in any degree

rotective against a further attack.

Bovine and human tuberculosis. — Since the iscovery by Koch in 1882 of the tubercle acillus, it has generally been held that tuberulosis in man and animal is one and the same isease. Villemin (1865) was the first to maintain his identity on the results of inoculation of bovine

and human tubercular matter into small animals. Chauveau (1868) carried out similar experiments upon cattle. Both workers were successful in transmitting the disease, which produced similar effects in the inoculated animals. Many other workers have obtained like results, which were more or less uniformly in support of the identity of bovine and human tuberculosis. The Royal Commission on Tuberculosis, 1895, said: "We regard the disease as being the same disease in man and the food animals, no matter though there are differences in the one and the other in their manifestations of the disease; and we consider the bacilli of tubercle to form an integral part of the disease in each, and (whatever be its origin) to be transmissible from man to animals and from animals to animals. Of such transmission there exists a quantity of evidence, altogether conclusive, derived from experiment."

It was, however, conceded on all hands that tuberculosis was a more virulent disease in animals than in man, and that the bacillus in the two species differed in morphological, biological, and pathological properties. In 1901, at the London Congress on Tuberculosis, however, Koch expressed the opinion that, "human tuberculosis differs from bovine, and cannot be transmitted to cattle," and that bovine tuberculosis was scarcely, if at all, transmissible to man. On the same occasion counter-evidence was produced by MacFadyean, Ravenel, Crookshank, and many others.

Koch based his opinion upon the negative results which followed the feeding of six cows with human tubercular sputum, making 4 others inhale human tubercle bacilli and otherwise infecting 9 more. None of the 19 cattle suffered, and no tuberculosis was found in their body tissues post-mortem. When, however, such cattle were infected with bovine disease they contracted it. Swine were also fed in a similar way, with like results.

In 1902 Koch again emphasised the comparative rarity of primary intestinal tuberculosis in the human being, and the local, as distinguished from the general, infective nature of accidental bovine inoculation of man (tuberculosis verrucosa cutis). In isolated cases the nearest lymph glands might become affected, but the disease remained nevertheless a local one. Koch further expressed the view that if bovine tubercu-Mosis was transmissible to man by means of the milk of cows with tuberculous udders, it would be reasonable to suppose that "groups of illnesses" would occur, in a manner analogous to other infective diseases, though the circumstances would differ owing to the different length of the incubation periods. To carry conviction as to milkborne tuberculosis, Koch maintains that the following conditions must be fulfilled:—(i.) Certain proof of tubercle in the person affected; (ii.) exclusion of other sources of infection; (iii.) the condition of all the consumers of the suspected milk; (iv.) the exact source of the ssuspected milk, particularly in respect to the disease of the udder of the cow yielding the milk. Finally, he concludes that all that can be said at present is that the injurious effects of milk infected with bovine tuberculosis and its products are not proven.

As a result of the wide differences of opinion revealed by the pronouncement of Koch's views, special Commissions of Inquiry were instituted in Germany, Great Britain, and other countries, in addition to the individual research work carried out

by many workers.

Kossel, Weber, and Heuss, who carried out a comparative research upon tubercle bacilli of different origins for the German Imperial Health Office, made a number of experiments on calves by injecting some forty different strains of human bacilli and fifteen strains from cows, fowls, and swine. They concluded (1904) that in a preponderating number of cases of human tuberculosis bacilli occur which were found distinguishable from the bovine bacilli of perlsucht morphologically, culturally, and in pathogenic properties, but that exceptionally in man tubercle bacilli occur which cannot be distinguished. They hold that

the possibility of infection in man under certain circumstances by milk from tuberculous udders is proved. They found that generalized tuberculosis was produced in animals by injecting strains of tubercle bacilli obtained from tuberculous diseases in children.

A Royal Commission was also appointed in this country, and in an interim report (1907) concluded that bovine and human tuberculosis was generically one and the same disease, though wide variation in degree of virulence may occur; that the human body can be infected by the bovine tubercle bacillus and the bovine body by the human tubercle bacillus; that in a certain number of cases human tuberculosis, especially in children, is the direct result of the introduction of the bovine bacillus, introduced in the majority of cases through cow's milk; that "cow's milk containing bovine tubercle bacilli is clearly a cause of tuberculosis, and of fatal tuberculosis in man;" that a very large proportion of tuberculosis contracted by ingestion is due to tubercle bacilli of bovine source; that a very considerable amount of disease and loss of life, especially among the young, must be attributed to the consumption of cow's milk containing tubercle bacilli; and that on this account the findings of the Commission point clearly to the necessity of measures more stringent than those at present enforced being taken to prevent the sale or consumption of such milk. (Second Report Royal Commission on Tuberculosis, 1907, pp. 20-37.)

Rabies (hydrophobia).—There are two chief forms, "furious" and "paralytic." The former is the commoner in dogs. The disease is comparatively rare in man, but is one of interest and importance from the point of view of preventive medicine. There were 761 deaths from hydrophobia in England and Wales between 1866 and 1885, Lancashire, London, the West Riding, and Cheshire being the principal

foci," whence the rest of the country was subjected repeated inroads. There were no deaths due to rabies in this country during the six years 1899-1904, except two in 1902. In the decennium ended 1890 there were 29 deaths on an average yearly.

Hydrophobia causes most deaths at ages between five and fifteen years; more among males than females; and more in late summer and autumn than at other

sseasons (Longstaff).

The disease is imparted to man by the bite of rabid dogs; or more rarely of rabid cats, foxes, or wolves. The ordinary incubation period is six weeks. It is stated to have been as short as one week in children bitten about the face. The disease almost invariably ends in death within a few lays. It is not known to have been transmitted from one human being to another. Bites about the face are more deadly than others, and in such cases the incubation is short. The danger is less when the part bitten is protected by clothing. Taking only cases of bites by animals proved beyond doubt to be rabid (the proof being the occurrence of a genuine case of rabies in some person or animal bitten by them or inoculated from them), hydrophobia manifests itself in about 15 per cent. of the persons bitten. By means to be described presently, this mortality may be greatly reduced, and by other preventive measures rabies can be, and has been, stamped out altogether.

Pasteur discovered that the virus, though absent from the blood, is present in the spinal cord of rabid animals as well as in the saliva, and that a portion of the spinal cord of a rabid animal (or human being dead from hydrophobia), inoculated beneath the dura mater of a rabbit, imparts the disease with certainty. This alone would be important as a certain means of diagnosis; but Pasteur further demonstrated that the virulent cord gradually loses its

virulence if kept in perfectly dry air, and that this attenuation afforded a means of curative or protective inoculation. He inoculated at short intervals with successively more and more virulent material, commencing with very attenuated virus (14 days old), until at last fresh cord was employed. It is surmised that the protection is due to a chemical substance, and not to any real attenuation of the microbe. However this may be, the result of the course of inoculations—so far from causing rabies, as a single inoculation with fresh rabic cord would do—is to render dogs insusceptible to rabies. The same treatment is now extensively practised in regard to persons bitten by rabid animals; and the mortality among such, under early treatment, is no longer 15, but a fraction of 1 per cent. In the decennium 1893-1902 an average of 1,450 persons were treated annually at the Pasteur Institute in Paris and the rate of mortality was 0.32 per 1,000.

The microbe has not been isolated with any certainty, although both micrococci and bacilli have been

found in the rabic cord.

Rabies can be stamped out by muzzling all dogs for a sufficient length of time. This was done in Sweden many years ago, and the country remains free from rabies. Less isolated countries are subject to constant new importations of the disease across the frontiers, but the success of repressive measures has been very great. Thus hydrophobia, though formerly prevalent, has been unknown in Berlin since 1874, and the Prussian provinces are practically free except on the eastern (Russian) frontier.

In England several local attempts were made in the same direction, but none upon a large scale until 1890, when it was made compulsory for a time throughout Lancashire, Cheshire, the West Riding, and London, and was followed by the cessation of peedily put an end to the prevalence of rabies. The Rabies Order of 1892, issued by the Board of Agriculture, enables county authorities to make regulations for muzzling and other preventive measures.

Glanders or farcy, as affecting man, was only described during the last century. Although in isolated instances t has been transmitted from one human being to another, it is practically always acquired from the horse. The virus does not appear to be capable of aërial transmission, except, perhaps, for very short ranges, and nothing is established as regards convection by water or milk. Inoculation is the almost invariable mode of infection so far as man is concerned; but it s probable that this may sometimes occur without abrasion of kin or mucous membrane. Among animals the disease spreads rapidly. It is very rare in man, but attended with an extremely high mortality, which is stated to be not less than 50 per cent. in chronic cases, and not far from 100 per cent. in the acute variety. The usual incubation ranges from three to eight lays, and is shortest in acute cases. For obvious reasons, men are much more liable to glanders than women or children. The disease affects the nasal and respiratory mucous membranes, and also the lymphatic glands. When the latter are attacked first, the disease is termed farcy.

Anthrax affects man in two forms, external and internal. External anthrax, or "malignant pustule," has its usual seat about the face or neck, and is no doubt due to inoculation. The first local manifestation is the appearance of a papule or vesicle, which develops in the course of a few days into an inflamed indurated mass, with a central black slough. The border may be fringed with vesicles or inflamed. The surrounding tissues and the lymphatic glands are swollen and indurated. The disease may remain localised, and end in resolution, or at most in suppuration. Usually, however, constitutional symptoms attend its course, and general infection may follow. Occasionally, malignant pustule supervenes upon internal anthrax. Internal anthrax appears to be due to inhalation or

swallowing of the poison. After a very variable incubation period, ranging perhaps from two to twelve days, the early symptoms are chills, weariness, depression, restlessness, and a feeling of constriction in the chest. This prodromal stage may last only a few hours, but more usually two to six days, and then graver symptoms set in suddenly. The prostration becomes extreme; pulse and respiration are rapid. The temperature is somewhat elevated, but always liable to sudden remissions, accompanied by perspiration. The patient may die from heart-failure, or brain symptoms may become prominent, or pneumonia or diarrhœa may supervene. Remissions may occur. The mortality is high, but recovery may happen even in serious cases. The protection derived from an attack is very slight, if any.

Anthrax received the name of "wool-sorters' disease" from its prevalence among men employed in sorting infected foreign wool and hair. Tanners, butchers, and persons engaged in handling raw hides

or horse-hair are liable to malignant pustule.

Anthrax attacks sheep, goats, pigs, cattle, and horses, and may be communicated to mice, guineapigs, and many other animals. As in man, the disease may be either localised or constitutional, and in the latter variety enlargement of the spleen is so prominent a characteristic that anthrax is known as "splenic fever." A field may become infected with anthrax, and healthy animals turned into it after the lapse of months, or even years, may acquire the The infection is probably imparted to the superficial layers of the soil by the blood or secretions of affected animals, in which the resistant spores form. Pasteur suggested that the spores from buried carcases are brought to the surface by earth-worms, but Klein has found that spores are not formed under such conditions, and that within a week all bacilli and all

infectivity have been destroyed by putrefaction, if the marcase is buried intact. Other sources of infection of animals are the use of infected dust or other trade effuse as manure, and drinking from streams receiving infected trade effluents.

Although exact evidence is wanting, it may be ssumed that anthrax can be acquired by eating the lesh of diseased animals. The usual modes of infection in man are by inoculation and inhalation, and it has been suggested that the poison may be carried by lies and other insects.

Bacteriology.—The pathogenic organism is the Bacillus anthracis, which forms spores when exposed oo air. The bacillus, which grows readily on ordinary media, is square-ended, 1 \mu broad and 4-5 long, someimes in filaments; aërobic, non-motile, liquefying; thiefly occurring in spleen and blood stream, and readily destroyed by heat or other disinecting agencies, though the spores are extremely esistant. It has been found that animals can be infected by inhaling or swallowing the spores, but not by the bacilli unless there is some abrasion such as to Allow practically of inoculation. Bacilli are destroyed by the gastric juice, spores are not. A further marked lifference between bacilli and spores is shown by the results of inoculation (Klein). The former cause a slight and localised form of the disease, the latter a severe constitutional malady, which is usually fatal. Thus, the fresh blood of a mouse dead of anthrax contains bacilli only, but upon exposure to air, or artificial cultivation, spores are readily formed. Inoculation of the fresh blood into sheep is followed by slight local anthrax, but if blood containing spores s used a fatal attack ensues. Moreover, if two animals of the same species are inoculated, one with apores of a broth culture and the other with bacilli of a gelatine culture, the blood bacilli of the latter will

be more attenuated than the former. But apart from this, anthrax virus varies in intensity according to the animal from which it is taken. Comparing the results of inoculation with virus from mice, guineapigs, sheep, and cattle, it is found that virulence increases in the order stated, mouse anthrax being the

least and cattle anthrax the most potent.

Martin has shown that the Bacillus anthracis produces albumoses, which in turn yield toxins much more virulent than themselves. Sclave of Siena has prepared an anti-anthrax serum from which excellent results have been obtained, reducing case mortality from 24 to 6 per cent. The animal used for immunisation is the ass, which is inoculated with slowly increasing doses of virulent cultures of the anthrax bacillus. Blood is drawn from the left jugular vein, and after clotting the serum is decanted and ether added to the extent of 3 per cent. of the whole bulk. The serum is then bottled with antiseptic precautions in 10 c.c. tubes.

In external anthrax early excision is of the greatest importance, and usually results in recovery.

Cases of anthrax in horses, cattle, sheep, pigs, etc., are dealt with under the Anthrax Order of 1899, issued by the Board of Agriculture. The owner reports the outbreak to the inspector specially appointed for the purpose by the local authority. It is the inspector's duty to inform the M.O.H. and to direct precautionary measures as to isolation, disinfection, and destruction of the carcase, as prescribed in the Order.

Tetanus.—Tetanus occurs in man and horses most commonly, though it may affect other animals. There is usually a wound, often an insignificant one, which may occur in any part of the body, but is nearly always on one or other of the limbs. It is not the locality of the wound nor its size that affects the

lisease. Wounds which are jagged, and occurring in absorptive tissues, are those most fitted to allow the entrance of the bacillus. The wound forms a local manufactory, so to speak, of the bacillus and its secreted poisons; the bacillus remains in the wound, out the toxins may pass throughout the body, and are especially absorbed by the cells of the central nervous system. Suppuration generally occurs in the wound. After a few days or, it may be, as much as a fortnight, when the primary wound may be almost forgotten, general symptoms occur. Their appearance is often the first sign of the disease. Stiffness of the neck and ffacial muscles, including the muscles of the jaw, is the most prominent sign. This is rapidly followed by spasms and local convulsions, which, when affecting the respiratory or alimentary tract, may cause a fatal result. Fever and increased rate of pulse and respiration are further signs of the disease becoming general. After death, which results in the majority of cases, there is very little to show the cause of fatality. The wound is observable, and patches of congestion may be found on different parts of the nervous system, particularly the medulla (grey matter), pons, and even cerebellum. The experience of the Pasteur Institute ttends to support the theory that tetanus is a "nervous" disease, more or less allied to rabies, and is best treated by intra-cerebral injection of antitoxin, which then has an opportunity of opposing the toxins at their favourite site. The toxins diffuse throughout the tissues of the body, but particularly affect the spinal cord. The long incubation period indicates that the toxins are probably produced by a ferment which is a most powerful poison.

Tetanus bacilli spores have been found in considerable quantities in the dust of dry jute fibre (Andrewes), and various cases are on record where the disease has been contracted in jute mills. Legge

attributes the presence of the bacilli to the soil in

which the jute is grown in Bengal.

In the wound the bacillus is present in large numbers, but mixed up with a great variety of suppurative bacteria and extraneous organisms. It is a straight short rod with rounded ends, occurring singly or in pairs or threads, and slightly motile. Flagella may be demonstrated (lateral and terminal). Branching also has been described. Indeed, it would appear that, like the bacillus of tubercle, this organism has various polymorphic forms. A large round spore occurs at one or other pole of the bacillus ("drumstick" appearance). The spores are very resistant. bacillus is an anaërobe, and grows at blood heat. In some soils it is common, and infection has frequently been derived from dust in jute factories, etc. disease is more common in hot countries than in temperate climates, and can frequently be induced in animals experimentally, by inoculation of a little garden earth, manure, refuse, or soil.

Puerperal fever.—In 1891-1900 the mortality from puerperal sepsis and childbirth averaged 5.09 per 1,000 births, which was higher than in any of the three preceding decennia. This result is due in large measure to differences of certification and collection

of statistics.

CHAPTER XV

TROPICAL DISEASES

In this chapter will be considered briefly some of the chief tropical diseases. Many diseases, such as pneumonia, diarrhœa, relapsing fever, typhus and typhoid, occur of course in sub-tropical and tropical countries; but such diseases as cholera, yellow fever, plague, malaria, and leprosy may rightly be considered as more correctly "tropical" diseases.

Asiatic cholera has its endemic area in certain parts of India, including the delta of the Ganges. The first great epidemic in India occurred in 1817. At irregular intervals it spreads in epidemic or pandemic form over a great part of the world. follows the lines of traffic by land or water, but no reason has been found for the apparently capricious selection of some routes and omission of others. The invasion of each new country along its line of march is almost invariably traceable to infection through some point of communication with a country already attacked. In temperate climates the outbreak often subsides or disappears in winter, but frequently reappears with the warm weather in the late spring or early summer; and may even recur again in the third year, apparently without fresh introduction.

Cholera gained a firm footing in England on four occasions, viz. 1831-2, 1848-9, 1853-4, and 1865-6, the epidemic extension from India having occupied respectively five, two, one, and two years in transit.

In 1831 infection was carried to Sunderland and Newcastle, from Baltic ports, in October, and in the course of the next year caused great mortality throughout the kingdom. The system of registration of deaths was not then in operation, but

according to Hirsch, the total mortality in England was 30,924. In 1848 Hull was infected from Hamburg early in October, and the disease at once spread. Only 988 deaths were recorded in England from October to March, but the epidemic made rapid progress in the summer, and before its final disappearance in December, 1849, it had caused 53,293 deaths, besides a heavy diarrhoea mortality, part of which may be supposed to have been due to cholera. In 1853-4 the invasion was again from Germany, London being attacked in August. outbreak continued through the winter, especially in the north; became more active in July, 1854, and overran the country, causing during the year 20,097 deaths in England and Wales. The 1865-6 epidemic was milder, but still resulted in 14,378 deaths.

On several other occasions the disease has invaded Europe, but failed to reach England, notably in 1871-4 and 1884-7. In 1892 it again spread to many parts of Western Europe, notably Hamburg, but apart from imported cases none appeared in England until September, 1893, when a few occurred at Grimsby, Hull, and elsewhere. Since then only small isolated outbreaks have occurred in this country.

Climate, season, and temperature.—Heat is a predisposing condition of great importance, but cold does not necessarily arrest the disease. Severe epidemics have occurred in Sweden and Russia in the depth of winter.

Even in India the seasonal curve of cholera prevalence is by no means always parallel with that of temperature. In Bombay the maximum of cholera deaths is in April; in the N.W. Provinces and the Deccan in August; in Madras there are two maxima, in February and September; and in Calcutta a chief maximum in April and a smaller one in November. In all these regions, except one, the highest mean temperature is reached in May; in the N.W. it is in June. In Madras cholera mortality is at its minimum in June, when the mean temperature is at its highest. In Europe, as a rule, epidemics attain their maximum in August, but sometimes in September or July.

Rainfall also has a marked influence upon the prevalence of cholera, and supplies a clue to many of the discrepancies mentioned above. In Calcutta the

conset of the rainy season in April and May is attended with lowered cholera prevalence, although the temperature is still rising; and cholera increases again in November, when the rains terminate, although the temperature is then falling. In the N. W. Provinces the maximum cholera mortality coincides with the maximum rainfall in August, two months later than the highest mean temperature. Outbreaks during the dry season in the N. W. are usually associated with sudden rains. Both temperature and rainfall are indirect or predisposing causes only, and the maximum of either may actually coincide with small prevalence of cholera. This is readily explained if we assume, with Pettenkofer, that moisture of the soil (as distinguished from dryness or saturation), heat, aëration, the presence of organic matter, and the specific poison itself, are the essential telluric conditions for the propagation of cholera. Rainfall sufficient to saturate the soil ((i.e. to raise the level of the ground water) will tend tto arrest cholera, however high the temperature may be; but so long as it merely moistens a previously dry soil, the other conditions being present, rain is apt to induce an outbreak. Given a moist soil, prolonged heat and drought may be the conditions most conducive to cholera.

Soil.—A direct relation between earth temperature, at a depth of 3 to 6 feet, and the prevalence of cholera during an epidemic, has been affirmed by Pfeiffer as regards England and Prussia. Pollution, and especially animal pollution, of the soil is doubtless an important factor. The subsoil water, according to Pettenkofer, by its movement affects very materially the development of cholera as well as enteric fever. In a dry, porous soil its influence may be trifling, but in general its fall—especially after great height—leaves the soil above it moist and aërated, and is

therefore favourable to cholera if the other conditions are present. Its rise to a high level has the reverse effect. The level of the ground water in Calcutta is highest in September, lowest in May, and therefore accords closely with the inverse relation to cholera affirmed by Pettenkofer.

Epidemics are especially apt to follow the banks of rivers and lakes, whether navigable or not. Hilly regions seem to escape frequently, although outbreaks may occur on high ground. The influence of elevation is often recognisable even in the distribution of

cholera in a single town.

Particular quarters, streets, and even houses are often selected for attack in a town invaded by cholera, and the selection can usually be explained by the existence of some filth condition in or about the houses, some pollution of water supply, or other insanitary condition.

Drinking-water has been closely associated with several outbreaks of cholera in England and India.

A few classical instances may be quoted here. The 1849 epidemic was very fatal in a part of London supplied mainly by two water companies—the Southwark and the Lambeth both of which distributed polluted Thames water after imperfect filtration. The incidence of cholera was substantially the same upon the customers of both companies, whose mains supplied the same districts and even houses in the same streets. In 1854, however, the population supplied by the Southwark Company suffered in 14 weeks a mortality from cholera of 13.0 per 1,000, and the customers of the Lambeth Company only 3.7 per 1,000. The disease singled out the houses supplied by the Southwark Company, whose intake remained as in 1849; while the Lambeth Company had in the meantime obtained a purer supply from a point many miles higher up the river.

The 1866 epidemic in London chiefly affected Poplar, and Netten Radcliffe found a similarly close coincidence between the area affected and the distribution of polluted water by the East London Company. In 1854 a sudden and intense outbreak occurred among the inhabitants of a small area near Golden Square, upwards of 400 persons being attacked within who had drunk the water from a pump in Golden Square—including persons residing at a distance who had sent for the water—and upon investigation the well was found to be polluted with sewage. A few fatal cases had occurred in the vicinity during the month previous to the outbreak, and there was strong probability of specific pollution of the well.

More recently, the experience of Hamburg, Altona and other places, has afforded further proof of the possibility of

econvection of cholera on a large scale by drinking-water.

Air has been regarded as the usual means of convection, but there is no clear evidence of this, and

many facts are opposed to it.

Race.—The incidence and severity are greater among negroes than Europeans, but apparently less among natives of hot countries in which cholera occurs than among European settlers.

Insanitary conditions of air, water, food (including milk), soil, drainage, and housing determine the points of attack. Hence the disease attacks more especially the poorest quarters of towns. Excre-

mental filth is most dangerous of all.

The poison doubtless gains access to the system by inhalation or swallowing. The incubation period is stated to be usually a few hours to three days, but may reach ten or even twenty days (Parkes). The mortality is, as a rule, not far short of 50 per cent. of the cases, and may be greater. Infection is given off in the discharges from the bowels, and probably in the vomit also. These may infect water, soil, or fomites.

Bacteriology.—Koch isolated from cholera discharges a microbe which from its shape received the name of "comma" bacillus. It is a curved rod, often about half the length of the tubercle bacillus, and motile. Strictly speaking, the "comma" bacilli are segments of a spirillum or vibrio, and not true bacilli. They are present in choleraic evacuations, generally, if not always, in spirillum form. They

occur in the intestinal wall and in the detached flakes of mucous membrane, in which they often lie in parallel arrangement, creating the "fish in stream appearance" under the microscope. They do not occur in the blood, and but little in the body generally. They have been found in wells polluted by choleraic excreta. They can be readily cultivated, and have a characteristic mode of growth; they flourish in sewage, and have been found to survive for 81 days in Marseilles harbour water, but disappear in a few days from pure water (Frankland).

The bacillus liquefies gelatine and grows in involution forms, but does not produce spores. It is stained by ordinary aniline dyes but not by Gram's method. In cultures it grows aërobically, and produces nitrates, but can be acclimatised to anaërobic conditions, and is then more virulent but less resistant. The organism produces characteristic growth on the ordinary media and produces indol, which, with the nitrates, gives a red reaction on addition of sulphuric acid (nitroso-indol). This test is known as "the cholera-red reaction." The bacilli are killed by acid liquids, so that they cannot live in presence of the acid gastric juice, but it has been shown that the contents of the stomach may remain alkaline if only liquids are taken. Koch neutralised the gastric juice of fasting guinea-pigs by alkali, checked the peristalsis by opium, and then introduced cultivations of "comma" bacilli into the stomach. The animals died in a day or two, the intestines being filled with fluid containing numbers of "comma" bacilli. Infected water and milk, and in one case an artificial culture, have been found to convey infection. It is possible also infection may occasionally occur through the respiratory tract, but this is certainly not the ordinary mode.

The "comma" bacillus is readily destroyed by heat

or other disinfectants, or by mere drying. The organism can, however, resist cold. Spores, if ever formed, would doubtless be more resistant. Emmerich has found in cholera evacuations another still more minute bacillus, straight and immobile, which

he regards as pathogenic in cholera.

That the comma bacillus is pathogenic in Asiatic cholera seems to be proved by the unintentional infection of a worker in Koch's laboratory from a culture of comma bacilli, under conditions that excluded other sources of infection. This bacillus has been "intensified" in various ways, and is then pathogenic to lower animals, including dogs and rabbits. It has also been attenuated, and a protective "vaccine" has been obtained. Haffkine prepares two vaccines, the weaker of which is made from cultures of the bacillus attenuated and the virus exalté from similar cultures the virulence of which has been increased by passage through animals. The first vaccine is inoculated and followed, three or four days afterwards, by the second.

Broadly, the reasons for believing Koch's comma bacillus to be the cause of cholera are (a) its constant presence in cases of the disease, (b) the results of accidental infection, (c) the agglutinative and protective properties of the serums of cholera patients, and (d) the result of Haffkine's pre-

ventive inoculation with his two vaccines.

Cholera nostras is the name given to a form of diarrhea endemic in Egypt and elsewhere. It does not spread in epidemic form, nor does it cause high mortality. Finkler and Prior have detected in the evacuations a "comma" bacillus, which, however, differs in size and mode of growth from Koch's.

Yellow fever has its endemic areas in the West Indies and certain parts of the Mexican and West African coasts, and frequent and intense epidemics occur between 33° N. and 23° S.—that is, from Charleston to Rio Janeiro. It is, therefore, essentially a tropical disease, but has occasionally occurred in epidemic form as far north as 43° N., at Leghorn and at Portsmouth in New Hampshire, and a few isolated cases occurred at Swansea (51° N.) in 1864 among persons who came into direct contact with infected vessels. Among European countries only Spain and Portugal come practically within its epidemic range. Infected ships reach European ports, but without causing any spread of the disease.

In the endemic area the greatest prevalence is between April and September, but in higher latitudes the epidemics occur in summer and autumn, beginning usually in July and August and never lasting beyond December. It is only endemic in regions where the mean temperature reaches 20° C. (68° F.), and epidemics do not as a rule occur in any latitude unless the temperature is above that point. A fall below 20° C., however, will not necessarily arrest the course of an epidemic when once begun, though it will usually cause a remission, but frost always arrests it more or less completely. According to Hirsch, atmospheric humidity and rainfall are as a rule favourable to yellow fever, unless accompanied by great lowering of temperature. With few exceptions the incidence of the disease is limited to the sea coast and the shores of great navigable rivers; only in intense epidemics does it affect elevated points. Even in the West Indies it rarely extends as high as 700 feet above the sea.

Yellow fever is a disease of towns, and has a strong tendency to localise itself in certain quarters, streets, or blocks of houses. Successive epidemics frequently single out the same local foci. As a rule these are in the filthiest and most crowded parts, where insanitary conditions of every kind abound, and more especially excremental pollution of the air, soil, and water. Outbreaks frequently occur on board ship, and if virulent are usually associated with similar filth conditions.

Telluric conditions have probably some influence, but it is not known to be conveyed by drinking-

water. Some authorities believe that wind can carry the infection for short distances.

Negroes, and perhaps also Chinese, have a remarkable immunity from yellow fever. Strangers of other races, and especially those from temperate climates, are very liable to attack. They may acquire a considerable and increasing degree of immunity tby "acclimatisation," but for this it appears to the essential that they reside in a yellow-fever disttrict for some years, including an epidemic period. It is possible that natives gain their immunity in the same way during childhood. Those who have undergone an attack are safest of all. It seems, however, that negroes as well as other persons may lose their immunity by a few years' residence in temperate climates, and even by migration from one yellow-fever district to another. Both susceptibility and severity of attack are found to increase in inverse proportion to the temperature of the native country. In Guiana the case mortality among West Indians was 7 per cent., Italians and French 17 per cent., English 19 per cent., Germans and Dutch 20 per cent., Scandinavians and Russians 28 per cent. Young and vigorous persons are said to be especially liable to attack, and men more so than women.

Yellow fever follows almost exclusively the lines of maritime traffic, and is not carried far by land. It is certain that ships may become infected and retain the infection with much tenacity, and baggage and other fomites have been known to convey infection without the intervention of infected persons. That a person whether infected or not may carry the infection with him appears to be undoubted, but the most intimate contact with the sick, wearing their unwashed clothes, occupying the same beds, making autopsies, and even inoculating the vomit, may, so it is alleged, fail to impart the disease.

The incubation is stated to vary greatly, from two or three to fourteen or sixteen days. The attack is sudden, with high fever and intense headache and lumbar and spinal pains. Vomiting sets in early, becomes incessant, and in three or four days blood is seen in the vomited matter, and often in the fæces. The urine is albuminous, and scanty or suppressed. Jaundice follows, and the patient passes into a typhoid state. A large proportion of the cases end fatally, the rest in lysis and convalescence in two or three weeks.

Bacteriology.—The etiology of yellow fever is obscure, but two theories share at the present time the chief support; the first is that the disease is due to Sanarelli's Bacillus icteroides, the second (and more probable) that it is due to a protozoan parasite, conveyed by a species of mosquito.

Sanarelli's bacillus is an anaërobic, motile, nonliquefying, resistant organism, obtained from the blood and tissues (not the alimentary canal) of persons suffering from yellow fever. It grows on ordinary media and ferments sugar. It is decolorised by Gram's staining method. Sanarelli holds that moulds are the "natural protectors" of the bacillus.

In 1901 the United States Army Commission reported that whilst B. icteroides was not always present in yellow fever, the blood appeared to contain the virus, and retained it after being passed through a Berkefeld filter. The commission further reported that the disease was not communicable by direct contact, but probably by mosquitoes, as in malaria. The species of mosquito suspected was the Stegomyia fasciata. Preventive measures were adopted in Havana on the mosquito hypothesis, with the remarkable result that the disease was stamped out. Guitéras of Havana has carried out further experiments which confirm many of the Commission's

ndings, and, in particular, the transmission of the isease by mosquitoes. A United States Army Expedition reinvestigated the subject in 1902-3, and reported that: (1) Bacteriological examination of the blood during life, and of organs and blood immediately tfter death, is negative. (2) Stegomyia fasciata, when allowed to suck the blood of a yellow fever poatient 41 hours after the onset, can on biting a non-immune person 22 days later produce the disease. (3) Stegomyia fasciata, contaminated by sucking the blood of a yellow-fever patient, and then killed, cut into sections and appropriately stained, presents with regularity a protozoan parasite, Myxo-coccidium stegomyiæ, which can be traced through a cycle of developments from the gamete to the sporozite. (4) Stegomyia fasciata, fed on the blood of a person with malarial fever, on normal blood, or artificially, does not harbour the myxo-coccidium.

The preventive measures to be adopted against yellow fever are the avoidance of districts in which it is known to prevail, the selection of inland, thinly-populated, and especially elevated points for residence, avoidance of the *Stegomyia* mosquitoes, and strict attention to sanitary conditions. Fatigue, exposure to sun, drinking, and improper food are said to predispose to

infection.

Dengue is now known to have occurred in epidemic form in the last century, but was first recognised during outbreaks in Farther India in 1824, and in the West Indies, Mexico, and the Southern States in 1827-8. Dengue resembles influenza in the intensity of its epidemics, which often affect almost every person in a town. It spreads mainly along lines of traffic. Even in tropical countries its prevalence is greatest in the hot season, and elsewhere it occurs only in summer and autumn, and in years of high temperatures. It affects maritime towns, and rarely spreads inland. Low-lying, filthy and overcrowded quarters are the first to be attacked. After a short incubation of two to four days the onset is sudden, with rigors, pyrexia, and acute pains and tenderness in the small

joints of the hand and feet. Very soon the large joints become affected, then the bones and muscles, and the head and back. A transient erythema appears at this stage in half the cases. In one or two days a remission or intermission of all the symptoms occurs after profuse sweats having a pungent odour, but after a short interval the symptoms return in a modified degree, often accompanied by glandular swellings, and a rash which as a rule consists of irregular red elevated spots. The second febrile stage passes off in three or four days at longest, and slow convalescence follows. Dengue is fatal only in debilitated persons, or in the very young or old. In some epidemics the disease has apparently attacked cattle, dogs, and cats.

Plague, first introduced from the East, and possibly identical with the Black Death of the Middle Ages (1348), was frequently epidemic in England up to the time of the Great Plague in 1665. In the following century it retreated eastward, and disappeared from Europe in 1841 and from Egypt in 1844. In Asia it has lingered on endemically in certain temperate regions, and of late has shown renewed activity, especially in India. Grave epidemics have occurred (from 1896-1907 it caused five and a half million deaths in India) and infection has been brought in recent years by ship to Great Britain. Plague is an infectious filth disease, fostered by destitution, uncleanliness, overcrowding, and lack of ventilation.

There are four chief varieties of plague: the bubonic form, the most common and typical, the lymph glands being chiefly affected in the groin, the axilla, or the neck; the septicæmic, in which the bacillus reaches the blood; the pneumonic, in which the lungs are mainly affected; and pestis minor, in which the affection of the glands stops short of the septicæmic stage, and even the local symptoms are slight. There are certain symptoms common to all forms of plague, when at all severe. An ordinary attack of plague usually begins three to five days after infection. Such attack may develop gradually,

but, generally, there is sudden onset with much fever, as indicated by a high temperature, rapid pulse, headtche, backache, hot skin, and thirst. The eyes are injected as if inflamed; the expression, at first haggard, anxious and frightened, becomes subsequently wacant, listless, and dull; the utterance is thick, and the gait unsteady as in one under the influence of salcohol. Mental aberration develops quickly. There iis frequently a marked tendency to faint. The ttongue is at first covered with a moist white fur except at the edges, which are red, but later on it becomes dry and of a mahogany colour. Vomiting and nausea are present from the onset. Sleeplessness is a characteristic symptom. The most distinctive sign, however, is the presence of swellings, or "buboes" in the groin, armpit, or neck, which appear on the second or third day of the disease, as large, smooth, tense swellings, painful and tender on pressure, and in size varying from that of an almond to that of an orange.

Cases occur in which these manifestations of the disease are greatly delayed or even absent, as, for instance, in "pneumonic," "gastric," and "septicæmic" plague; forms of the malady which may be mistaken respectively for inflammation of the lungs, typhoid fever, or acute blood poisoning. Plague in these forms is always grave; not only because of the fatality of the cases but for the reason that they are highly infectious to other persons. It is important, therefore, that in localities where plague is present or is threatened, cases of anomalous illness of the above sorts should be, without loss of time, brought under medical supervision.

There is also the so-called "ambulant form." In plague of this description the affected person is hardly ill at all, presenting no definite symptoms perhaps beyond indolent, though painful, swellings in groin or

armpit. Such plague cases may nevertheless be in-

strumental in spreading the disease.

Bacteriology.—The causal organism, Bacillus pestis, was discovered in 1894 by Kitasato in the blood of patients suffering from the disease. It is a small, round-ended, oval cell, with marked polar staining, hence presenting an appearance not unlike a diplococcus. In the middle there is a clear interspace, and the whole is surrounded with a thick capsule, stained only with difficulty. The organisms are nonmotile and often linked together in pairs or even chains (especially in fluid cultures), and exhibit polymorphic forms. In culture the bacillus may be coccal or bacillary in form.

It grows readily on the ordinary media at bloodheat, producing smooth, shining circular creamcoloured colonies, with a wavy outline, which eventually coalesce to form a greyish film. If melted butter (or ghee) or oil be added to bouillon, the bacillus grows in "stalactite" form. The following negative characters assist in differentiating the bacillus: There is no growth on potato; milk is not coagulated; gelatine is not liquefied; Gram's method does not stain the bacillus; and there are no spores.

over sulphuric acid.

In the body the bacillus chiefly frequents the spleen and lymph glands, the latter becoming inflamed,

bacillus is readily killed by heat or by desiccation

especially the inguinal and axillary.

Haffkine has prepared a vaccine which diminishes the incidence and fatality of the disease in inoculated populations, although the degree of protection is not The vaccine is made from a six-weeks' old broth culture of B. pestis, sterilised at 65-70° C. for an hour, and preserved in bottles containing 0.5 per cent. carbolic acid. The dose is 5-10 c.c.

Plague is, in a general sense, a disease of locality

irect agent is more or less subordinate. Probably he essential cause of epidemics is the plague stricken at, and the intermediate agent between rat and man, and rat and rat, is the common rat flea (P. cheopis) which acts as a carrier of the bacillus. The blood of an infected rat generally contains enormous numbers of bacilli, and insects sucking such blood readily become infected. Rats, which usually suffer from blague before epidemics occur among men, thus convey infection from place to place and from port to port.

Preventive measures, therefore, include vigorous disinfection of infected houses; inspection and segregation; inoculation; and the extermination of rrats. The latter may be accomplished by various rrat traps and poisons, and on board ship also by

ccarbonic acid gas.

Malarial diseases may conveniently be considered as a single group, although they include many varieties that have received specific names. All have a characteristic tendency to periodicity in the course which the symptoms follow, and this peculiarity is made the basis of a broad subdivision into intermittent and remittent types of fever, the former being the less severe.

Intermittent fever or ague is further subdivided into quotidian, tertian, and quartan varieties, according to the length of interval between the attacks, which may occur every two, three or four days. Each attack is marked by a cold stage, a hot stage, and a sweating stage; but the temperature in the intervals always becomes normal. Of these sub-varieties, quotidian ague, with daily attacks, is the most severe, has the longest hot stage and the shortest cold stage, and approaches the most closely to the remittent type.

In remittent fevers there are one or two exacerbations every day, but the temperature during the intervals does not subside to the normal level, nor do the constitutional symptoms entirely disappear. The cold stage is very short and the hot stage protracted. Remittent fevers are met with principally in tropical regions, but are endemic in many marshy districts in Southern Europe, and occasionally spread as far north as the Baltic.

General conditions .- As the intensity of the endemic or epidemic prevalence diminishes, the individual cases tend to assume more and more the intermittent type. Broadly speaking, the tendency to malaria increases with the temperature. especially if the diurnal range of temperature is high. Atmospheric humidity is also favourable to malaria. Moisture of the upper part of the soil is an almost invariable condition of malarious districts, and the effect has been said to be heightened by a slight superficial drying. Clay soils, which retain moisture, are among the worst, but chalk, or sand, or other porous soil, may become highly malarious if from any cause the subsoil water is high. Malarious soils almost always contain vegetable matter in greater or less quantity. In such districts heavy rainfall, especially after long heat and drought, is generally followed by outbreaks of malaria, and irrigation has a similar tendency. They cease to be malarious when the ground is either completely flooded or dried-temporarily or permanently—by drainage and extensive planting (e.g., with Eucalyptus globulus). As regards the surface, the prevalence of malaria decreases with the elevation, other things being equal. High ground is the most healthy, but exception must be made in respect of spurs of hills and depressions even at high levels, and more particularly of ravines. Among other causes conducive to malaria are abandonment of cultivation, or, on the other hand, the breaking up of new ground, excavation, or cutting down trees. Volcanic disturbances and earthquakes have, it would appear, sometimes been followed by malarial outbreaks. Fodor concludes from observations at Pesth that only the superficial layers of the soil are concerned in producing conditions favourable to malaria.

Such being the general characteristics of malarious districts as regards climate and soil, it must next be pointed out that they are in themselves inadequate to produce malaria, and also

447

calities. Malaria is unknown, for example, in many parts of cland, in the Bermudas, and the Pampas of South America, there the physical conditions would à priori appear to be aghly favourable. On the other hand, some of the malarious stricts in the Deccan and Mysore are steep mountain slopes, and many others in India and elsewhere are equally free from coisture of the soil. Instances are common in which districts, reviously healthy, become temporarily or permanently malarious, without apparent change in their physical conditions. Malaria being a specific disease, the presence of a poecific organism is necessary, and the conditions above esscribed are to be regarded merely as more or less favourable its growth and dissemination.

ap. XV.]

The etiology of malaria.—In 1880 Laveran first liscovered and described parasites in the blood cells of malarial patients, and on further investigation it was soon found that these assumed many different forms. These differences depend upon the kind of lever and the stage of fever. The reasons for believing that Laveran's bodies—though they have not yet been cultivated outside the human body—are the specific cause of malaria are that the parasites found in the blood of malarial patients of all countries are the same, and such parasites are not found in healthy persons; that their development fully accounts for the production of the melanæmia and malarial pigmentations of viscera owing to the melanin-forming property of the parasite; that the phases in the development of such parasites correspond with the clinical course of malaria (Golgi); that quinine, which cures malaria, kills the parasite; and that malaria can be conveyed by the introduction of this parasite into the blood of man, and the parasite reappears in the blood of the individual so inoculated. Laveran's bodies have been variously classified as knowledge of them has grown. It is now agreed that these parasites belong to the Sporozoa, to the order of Hæmocytozoa, and to the genus of Hæmamæba. They gain access

to the blood corpuscles in which they develop, producing sporocytes which, on the rupture of the corpuscle, become free in the blood and attack other corpuscles passing through the same changes (the cycle of Golgi), and resulting in a fresh generation of spores. The exact changes, however, depend partly upon the form of disease (quartan, or benign tertian, or malignant quotidian), but much more so upon the immediate needs of the parasite, that is to say, whether the development is to be in the human body only and asexual, or in the mosquito and sexual. (Manson, Ross, and others.) If it is to be the latter the amœbula does not sporulate in the corpuscle, but becomes a crescentshaped body absorbing the corpuscle. These crescents, or gametocytes, are of two forms, male and female, and are the first stage in the extra-corporeal stage of the life-cycle, that is to say in the mosquito. The male gamete (microgametocyte) develops flagella, which are of the nature of spermatozoa, and break off from the cell and become free in the blood ingested by the mosquito. The female crescent, or gamete (macrogametocyte) starts in the same way as the male form, but becomes spherical and extrudes two polar bodies, receiving at the site of extrusion an impregnation by the free flagellum of the male cell. The result is a zygote, a travelling vermicule, which passes into the stomach wall of the mosquito, and becomes encysted there, and eventually reproduces spores or sporozoits. About the twelfth day it bursts, discharging the reproductive elements into the body cavity of the mosquito, which, passing from thence, become accumulated in the salivary gland, and are thus in position to be injected into the blood of the next human being bitten by the mosquito. In the human blood again they attack the corpuscles, and recommence the asexual life cycle. Just as the sporocytes are the means of propagating the parasite in the blood, so the flagellated body and

sexual cycle is the means of propagation in the

mosquito.

The species of mosquito which has been shown to be the host of the malaria parasite is the Anopheles, the body of which is slim, the proboscis long and thick, the palpi long, and the wing "dappled" with dark spots on its anterior margin. It is a "puddle-Ibreeding" insect. The common mosquito, or Culex, lhas short palpi, thick body, thin proboscis and unspotted wings. It is a "pot-breeding" mosquito. Certain conditions are necessary for the multiplication of malarial mosquitoes, namely, high atmospheric temperature, 75-104° F., collections of water, fresh or brackish, and the presence in the breeding pools of llow forms of animal and vegetable life. Many of the physical conditions hitherto cited as favourable to the disease are now seen to be favourable to the specific mosquito, and their relationship to malaria is thus, in part, explained. The prevention of malaria depends upon the fulfilment of four chief requirements: (1) The prevention of mosquito breeding by landdrainage and removal of collections of water; (2) the destruction of mosquitoes (Anopheles) and of their eggs, larvæ, and nymphæ; (3) avoidance of mosquito bites, and (4) the therapeutic and prophylactic use of quinine.

Leprosy is a disease of great antiquity in Egypt, India, China, and Japan. The Biblical term "leprosy" doubtless included many forms of skin disease, and the same ambiguity attends most, if not all, of the other records of the disease in former ages.

Leprosy has been especially confounded with syphilis and elephantiasis arabum, true leprosy being sometimes termed elephantiasis gracorum. It probably made its appearance in Europe about the beginning of the Christian era, and became widely diffused during the next eight or nine centuries. A further increase—attributed by some to the Crusades—took

place in the twelfth and thirteenth centuries, but in the fourteenth and fifteenth centuries it began to disappear, and at the present time has vanished from Europe, with the exception of certain limited districts, more especially in Norway, Sweden, Finland, Spain, Portugal, and some parts of Italy, Greece, Turkey, South Russia, and some of the Mediterranean islands, including Sicily and Cyprus. It is prevalent in the West Indies, the west coast of South America, India, China, and the greater part of Asia; most of the islands in the Pacific, notably Hawaii, and in many parts of Africa. It is common in Mexico, and minor foci exist in Louisiana, New Brunswick, and British Columbia. Special attention has recently been directed to its increase at the Cape. Districts upon or near the coast are by far the most frequent seats of endemic leprosy.

In England, special legislation was directed against leprosy in the tenth century, and leper hospitals were established in the eleventh, twelfth and thirteenth centuries in many of the larger towns. The last record of any extensive prevalence in England was at the end of the seventeenth century, and it is

now unknown except as a disease acquired abroad.

Leprosy is diminishing in its few remaining European centres, but probably slowly increasing in the West Indies, Demerara, South Africa, and some other countries. Chinese immigration is said to have introduced it into North Australia and British Columbia.

General conditions.—Climate would appear to have little influence, since the geographical distribution is or has been almost world-wide, nor has anything been made out definitely in regard to more local meteorological conditions. The effect of season, if any, cannot be traced in a disease which begins insidiously and runs a protracted course. The greater incidence upon coast regions seems to give a clue to the etiology, but the disease prevails in elevated regions far removed from the sea. The use of fish, and especially of decomposing fish, as a principal diet is supposed to contribute largely to the causation of leprosy (Hutchinson). This view is supported mainly by the prevalence of the disease in littoral districts, but

also to some extent by its frequent appearance in mew regions among Chinese settlers, whose skill in cookery enables them to make use of putrid fish, and llastly by its observed disappearance from certain districts where fish diet ceased to be general. On the other lhand, there are many inland leprous regions in which fish is rare, or, as in Central China, limited to the ruse of the wealthier classes. The disease has declined in many localities without any material change in diet, and, further, the vegetarian Brahmins are by no means exempt from leprosy. It would seem, therefore, that a fish dietary, if it has any etiological significance at all, is not operative in all cases. Another hypothesis is that deficiency of common salt in food is conducive to leprosy. Sanitary improvements in the habits and surroundings of the population have, in all probability, been chiefly instrumental in suppressing leprosy in England and elsewhere. Although never confined to the poorer classes, its incidence has been greater upon those living in filthy and unhygienic conditions.

Nothing is known as to any relation with telluric conditions, or transmission by water or air. Negroes, Hottentots, and Chinese are more affected than white races living in the same localities. Males suffer in greater numbers than females, and among the very old and the very young the disease is of rare occurrence.

Authorities are at variance as to the hereditary transmission of leprosy, and very little is known of the way in which the disease is ordinarily acquired. It is communicable by direct or indirect means, from the sick to the healthy, and there is no certain evidence of its arising or spreading in any other way. Hansen states that no sign of leprosy had appeared among the descendants of 160 Norwegian lepers who settled in the United States. Arning has proved it to be inoculable, having successfully inoculated a condemned convict.

Clinical characters.—The incubation period in Arning's case was two years. It is believed to be very variable, extending over months or years, with or without premonitory symptoms. Three chief types are described—(a) the anæsthetic (or maculo-anæsthetic), (b) the tuberculated, and (c) the mixed. The anæsthetic form causes a destruction of nerve fibres, the tuberculated chiefly affects the skin, resulting in a nodular growth or diffuse infiltration. It produces marked disfigurement. The average duration is given as eight years in the former and sixteen in the latter, but with wide variations. When the disease begins in early life, its duration is comparatively short. The result is almost always fatal, but it is stated that in Norway thirty-eight cases were cured in the five years 1881-5, and Kaurin believes that if the disease is localised, early amputation may sometimes permanently arrest it. According to Sand, the disease may exhaust itself in thirty or forty years if the constitution is exceptionally strong.

Bacteriology.—The B. lepræ was discovered by Hansen in 1874. It is present in enormous numbers in the skin and tissues, and has a form very similar to B. tuberculosis. Spores have not been seen, but Neisser holds that this is the form in which the bacillus gains entrance to the body. The arrangement in clumps and colonies inside the round cells is useful for diagnostic purposes. The bacilli are also found free in the lymphatics, inside connective tissue cells, in the walls of blood vessels, in the lymphatic glands, and in the internal organs. Bordoni-Uffreduzzi, Carrasquilla (1899), and Campana claim to have isolated the bacillus and grown it on artificial media, the two former aërobically on peptoneglycerine-blood-serum, at 37° C., the last-named anaërobically. Cultivated bacteria from the organs of lepers, described rather later by Babés, and still

more recently by Czaplewski, differ from the genuine oacillus of leprosy in their incomplete resistance to cids. There is little doubt that the disease is prolluced by Hansen's bacillus. Bordoni-Uffreduzzi holds that the bacillus has a saprophytic stage in its life

Hansen, Ehlers, and others believe that the disease ss spread by contagion, Virchow held the reverse opinion. Besnier and Sticker consider the infection s conveyed in nasal secretion. The Leprosy Com-

mission in India (1890-91) reported that—

(1.) Leprosy is a disease sui generis; it is not a form of syphilis or tuberculosis, but has striking etiological analogies with the latter. (2.) It is not inherited, and has a natural sendency to die out. (3.) Though contagious and inoculable, the extent to which it is propagated by these means is exceedingly small. (4.) It is not directly originated by the use of any particular article of food, nor by any climatic or telluric conditions, nor by insanitary surroundings, neither does it peculiarly affect any race or caste. (5.) It is indirectly influenced by insanitary surroundings, such as poverty, bad food, or deficient drainage or ventilation, for these increase the susceptibility.

Preventive measures.—The practical suggestions of the Commission for preventive treatment included voluntary isolation of the sick; prohibition of the sale of articles of food by lepers; leper farms; orphanages; and "improved sanitation and good dietetic conditions" generally. Serum-therapy has been attempted, but without success. Up to the present time no curative agent has been found.

In Norway, leper asylums were established in 1856, and the number of lepers is believed to have steadily declined. The isolation in these asylums is not rigid, but in Hawaii an island is set apart for the leper settlement, and the seclusion is practically life-long. It seems clear that the infection is not acquired readily, if we take the ordinary infectious diseases as

a standard, and there is some probability that even a partial segregation of lepers may have considerable influence in arresting the disease, assuming its spread to be due mainly to personal intercourse. The difficulties of obtaining a tolerably complete isolation or even registration of lepers are of course very great.

Malta fever (Mediterranean fever) is common along the coast of the Mediterranean and in its islands. It is a disease of long duration and variable symptoms, including remittent fever. Perspiration, pains, swelling of joints, enlargement of the spleen, etc., are among the common signs. In 1887 Bruce cultivated from the spleen the Micrococcus melitensis, a small round or slightly oval coccus, occurring singly, in pairs, or in chains. It can be cultivated from the spleen, on agar at 37° C.; colonies appear about the third day as small, round, slightly raised growths; old cultures assume a buff tint. The organism is present in the peripheral blood in all cases during the early stages of the disease, and in severe pyrexial relapses. It has been isolated from the urine. The disease is transmissible to man by the medium of milk from infected goats.

Sleeping sickness is probably caused by the entrance into the blood, and thence into the cerebrospinal fluid, of a species of trypanosome (probably the Trypanosoma gambiense), which is transmitted from the sick to the healthy by a tsetse fly (Glossina palpalis), and, apparently, by it alone. It has been demonstrated by experiment that not only were these flies, fed on sleeping sickness cases, capable of conveying the disease to healthy monkeys, but that freshly caught flies from an infected area, without any artificial feeding, were also capable of conveying the disease. This fly, like its congener the tsetse fly of South Africa, is confined to well-defined areas, which correspond absolutely with the distribution of sleeping

sickness; whereas, in regions where no Glossina palpalis is found, although other biting flies abound, there is no sleeping sickness. While a large percentage of the inhabitants of the sleeping sickness areas have the trypanosome in their blood, in not a single inhabitant of non-sleeping sickness areas was this parasite found. Monkeys inoculated with cerebro-spinal fluid from sleeping sickness patients, or with blood ffrom natives not as yet showing symptoms of sleeping ssickness, but containing a similar parasite, sickened and died with all the symptoms of sleeping sickness. As in the case of ankylostoma, individuals may have this trypanosome in their blood, and yet reveal no ssymptoms of the disease. A Portuguese Commission llaid emphasis upon a diplo-streptococcus which a number of observers have found to be present.

The disease has spread in the Congo region, Uganda, and northwards along lines of communication, that is rivers, and has become a scourge in certain riverine tribes. Clinically the early stage of sleeping sickness may be recognised by enlargement of the cervical glands (polyadenitis), enlargement of the spleen, edema, and certain pathological conditions of the blood, including lymphocytosis. Sleeping sickness is probably the last stage of a disease and

seems to be invariably fatal.

Dysentery.—Endemic or tropical dysentery is possibly due to Amæba coli vel dysenteriæ. Epidemic dysentery is more probably due to Bacillus dysenteriæ, and sporadic and parasitic dysentery to various parasites, such as Balantidium coli and the Bilharzia. Mott holds that "asylum dysentery" is identical with tropical dysentery, and that both conditions are in all probability of bacillary origin.

The disease is not confined to tropical climates. At one time it was "one of the most prevalent and fatal diseases of London" (Baly). It is believed to

be influenced by season and soil, autumn and polluted subsoil favouring its occurrence, though personal predisposing conditions doubtless play an important part. From a sanitary point of view water supply, dietary, and uncleanliness are the conditions into which inquiry should be made. It is probable that the excreta serve as one of the chief media of infection, and they should therefore be disinfected. Apart from institutional outbreaks (e.g. asylum dysentery) dysentery is

a comparatively rare disease in this country.

Beri-beri is endemic in Western India, in the India Archipelago, China and Japan. It is mostly confined to the working classes, and is said by some to be due to an organism infecting stale, white rice; others consider it a "place disease," and Durham holds that it is more a "gang" or "institutional" disease. It is a kind of scurvy of the tropics, affecting the underfed, and is characterised by weakness, dropsical distension of the abdomen, limbs, and face, and many of the features of alcoholic neuritis. Probably the disease is communicated from person to person directly or through fomites, though Braddon holds that it is caused by a non-volatile alkaloid.

CHAPTER XVI

PREVENTION OF INFECTIOUS DISEASES

IIn the preceding chapters reference has been made to the various means by which specific and other diseases are disseminated or checked. It remains to consider the powers and duties of the S.A., and of the M.O.H., in relation to preventible diseases, and especially infectious diseases. All infectious diseases may be said to be preventible, but not all are amenable to the preventive measures which at present can be enforced by the most active sanitary administration. Small-pox and rabies may be taken as types of specific diseases which can be and ought to be suppressed, and are now, as a matter of fact, limited to such persons and such populations as do not choose tto adopt the measures which reiterated experience has shown to afford safety. Measles and whooping cough, on the other hand, although strictly speaking preventible diseases, always dependent (so far as we know) upon direct or indirect infection from other cases, have not yet been brought under control. No prophylaxis has been discovered, and as infants are especially liable, the susceptibility of the population is speedily renewed, even after an outbreak. Isolation and disinfection are theoretically capable of arresting the spread of these diseases if applied in every case and at the commencement of the infectious stage, but in practice this can scarcely be accomplished at present, for reasons that will be stated later.

Scarlet-fever, diphtheria, and enteric fever are examples of a different class. They are not entirely dependent upon infection from person to person, but are affected by a variety of conditions of which every

year brings us fuller knowledge. Some of these conditions, including personal infection, infection from lower animals (by consumption of milk or flesh, or in other ways), and infection by drinking-water, are more or less under our control. So, too, are the various "insanitary conditions" which conduce to the spread and malignancy of specific diseases, and telluric conditions also are to a great extent remediable. Experience has shown that if carried out thoroughly, isolation, disinfection, and quarantine can very greatly reduce the incidence of these diseases, but there is always danger from unrecognised cases, from the early stages of the disease before the diagnosis is made, from importation of infection, and from accidental or wilful neglect of precautions. It must also be remembered that there may be modes of infection in these diseases at present unknown to us, and this is especially probable in respect of diphtheria.

Compulsory notification of certain infectious diseases rests in London upon the Public Health (London) Act, 1891, and elsewhere upon the Infectious Diseases (Notification) Act of 1889, as amended by the Infectious Diseases (Notification) Extension Act of 1899, which also enables any S.A. with the sanction of the L.G.B. to add temporarily or otherwise other infectious diseases to those scheduled in the Act, which are-smallpox, cholera, diphtheria, membranous croup, erysipelas, scarlet-fever, typhus, enteric fever, relapsing fever, continued fever, puerperal fever. The inclusion of membranous croup in the list tends to make the reporting of diphtheria more complete, and "continued fever "covers obscure cases of enteric fever, or possibly typhus. A large number of trivial cases are reported as erysipelas, but the disease has practical importance in connection with vaccination and puerperal fever.

Doubt having arisen as to the meaning to be attached to "puerperal fever," the Royal College of Physicians have expressed the opinion that it should be understood tto include "septicæmia, pyæmia, septic peritonitis, sseptic metritis, and other septic inflammations in the pelvis occurring as the direct result of childbirth."

Certain minor diseases such as chicken-pox may the made notifiable temporarily or otherwise in order tto ensure detection of more serious complaints such

as smallpox.

Forms of certificate are supplied to every practitioner practising in the district, and a fee of 2s. 6d. is paid to him for each certificate regarding a private patient and 11s. for each case in public practice. "Every medical practtitioner attending on, or called in to visit, the patient shall forthwith, on becoming aware that the patient is ssuffering from an infectious disease to which this Act sapplies, send to the M.O.H. for the district a certificate stating the name of the patient, the ssituation of the building, and the infectious disease ffrom which, in the opinion of such medical practitioner, tthe patient is suffering." The penalty for default is a fine not exceeding 40s. Under the same penalty the lhouseholder is compelled to notify, but in a less formal way, and without receiving any fee. The Act applies tto "every ship, vessel, boat, tent, van, shed, or similar structure used for human habitation," but not to any "hospital in which persons suffering from an infectious disease are received." It gives no power of compulsory removal of patients to hospital, nor even power of entering upon the premises for the purpose of making inquiries, but it is very rarely that any difficulty is met with in the latter respect.

Among the advantages to be derived by the M.O.H. from compulsory notification are the following:— 1. Early and complete knowledge of all the cases of notifiable disease, and thus of the whole prevalence and distribution in the district. Death-returns give only a small fraction of

the cases, and after an interval of many days or weeks.

2. Power to exercise such supervision as may be necessary over every case during its whole course, and to enforce due observance of the provisions of the Public Health Act as regards isolation and disinfection. This has special importance in connection with outbreaks of scarlet-fever, diphtheria, or enteric fever in dairies, schools, etc.

3. Opportunity of offering, and in some cases compelling, removal to hospital in every suitable case, unless the S.A. has failed to provide hospital accommodation. Vaccination can be offered, and is rarely refused in households invaded by

small-pox.

4. Opportunity of investigating the sanitary conditions of all households in which cases of enteric fever, or diphtheria, or other notified disease may occur.

5. Power to control the spread of infection through schools or other centres, by excluding members of infected households.

6. Means of detecting at once any suspicious grouping of cases around schools, milk supplies, water supplies, or other common foci. A comparison of the data obtained by the routine inquiries into each notified case will at once reveal this, although there may be nothing in any single case to arouse the suspicions of the friends or of the medical attendant.

In a number of towns it has been found possible, with the aid of notification, to isolate in hospital all the cases of small-pox and three-quarters, or more, of the scarlet-fever, and to establish a cordon of vaccination around each new small-pox centre. Very few towns have made adequate provision for the hospital isolation of enteric fever, nor is there as yet any absolutely reliable prophylactic such as vaccination to be offered, and, moreover, the danger of spread of infection from person to person by ordinary proximity or contact is less marked than in small-pox or scarletfever; hence notification brings to bear comparatively slight reinforcements of preventive measures, but it has great value in bringing to light any grouping of cases around a water or milk supply or other common cause. Diphtheria is more readily communicated

rom person to person, but in other respects the same considerations apply. The notification of diphtheria an scarcely be complete or satisfactory, owing to the number of slight and almost undistinguishable cases, and hospital isolation upon an adequate scale has not ret been tried.

Epidemic prevalence of disease is determined by many causes, of which some are still unknown to us, and others (including climatic conditions) are beyond numan control. Against such influences, unless they should prove to be operative merely by increasing the hacilities for the spread of infection in ordinary ways from person to person, notification and the precautions which it renders possible are of no avail, and epidemics will continue to occur in spite of it. Nevertheless the known and preventible causes, such as the transmission of infection from person to person, or by means of milk, are factors of the utmost importance, and it is safe to affirm that, however great or however small may be the total number of victims, the suppression of known and preventible favouring conditions renders the danger less than it would otherwise have been. Undividuals may be saved, and widespread outbreaks Idelayed, restricted, or averted.

The question whether measles ought to be added to the list of notified diseases is an important one. It is more than doubtful if any legislation could at present bring about a complete notification of measles, and, if such could be obtained, whether it would be worth while, for the following reasons. (i.) Medical advice is frequently not taken in mild forms of measles. (ii.) It is a purely epidemic disease, which has no relation to water or milk supplies, or to defects of drainage. (iii.) There is no vaccination. Isolation and disinfection are, therefore, the sole preventive measures which could be brought to bear against measles, even if it were possible to

obtain complete notification. Isolation in the homes of the poor is impracticable, and isolation in hospitals would entail a formidable addition to their present size. The results of isolation also are far less likely to be satisfactory in measles than in scarlet-fever, and as a very large proportion of the sufferers are infants, there would be great reluctance on the part of parents to consent to removal. (iv.) Infection begins generally at least three days before the rash appears. (v.) There is a long incubation period which adds to the difficulty. (vi.) Lastly, measles is often so widespread that few susceptible children among the working classes escape, and measures of notification and isolation would be on too vast a scale to make them worth while.

That valuable information would be derived from notification of measles cannot be doubted, and from this point of view its inclusion in the schedule of notified diseases is desirable. It is urged, also, that even an incomplete notification would be invaluable in rendering possible early control over school-infection; that hospital isolation of measles would no longer appear impracticable if epidemics were not allowed to proceed unchecked; and that notification of measles would bring to light many cases of scarlet-fever which, in the absence of any medical inspection, are regarded by the parents as "only measles." This last argument points to the notification of rötheln and chicken-pox.

The same considerations apply to whooping cough also. About 2 per cent. of the sanitary authorities of Great Britain have voluntarily made measles a notifiable disease for a longer or shorter period. On the whole, however, the system has not been of permanent value. In London in 1904 the disease was scheduled under the Public Health (London) Act as a "dangerous infectious disease" to which sections dealing with disinfection, exposure of persons, etc.,

sshould apply. Voluntary notification has also been applied to phthisis in a number of sanitary districts, including Manchester Brighton, and many of the metropolitan boroughs. In Sheffield, by a local Act of 1903, compulsory notification of phthisis has been established for an experimental period of seven years.

The regulations of the L.G.B. require that upon receiving information of any outbreak of dangerous infectious disease the M.O.H. shall at once make full inquiries, and adopt such precautionary measures as he may consider requisite. The necessary inquiries and investigations are simplified and made more systematic by the use of printed forms with spaces for the insertion of the details for each notifiable disease. Those used for scarlet-fever, enteric fever, and diphtheria, should give prominence to the details of milk supply and other channels of infection.

The following inquiries should be made in all cases:

- 1. Patient.—Address, name, sex, age, date of onset, date of rash, present and past isolation, probable source of infection, recent contact with infected persons or things.
- 2. Household (including patient).—Sex and age of each inmate, susceptibility (as shown by history as to previous attack), date of previous attack (if any), occupation, place of work or school, and date of last attendance thereat.
 - 3. Work or business carried on in the house.
 - 4. Water supply and milk supply.
 - 5. Sanitary condition of premises and surroundings.
- 6. Previous cases of the disease, or of conditions allied to it or simulating it in the home or in the vicinity, or at the school or work-place.
- 7. Possible sources of infection:—(1) Visits; (2) Visitors; (3) Place of work; (4) Food, milk, ice cream, shell fish, watercress, etc. -what, when, where purchased, condition; (5) Water; (6) Previous cases of sick headache, sore throat, or diarrhoea in the house or neighbourhood; (7) School, including Sundays; (8) Drain nuisances, at or near house; (9) Tailoring, washing, mangling.

In dealing with an outbreak of small-pox, further inquiry must be made regarding the condition of each individual as to

vaccination and revaccination, noting the dates.

In enteric and diphtheria cases note should be taken of the previous state of health of the patient, the duration of his residence in the house, and the occurrence of any sickness among domestic animals.

The precautionary measures to be enjoined, and if necessary enforced, will vary with the nature of the disease and the

circumstances of each individual case.

Registers of sickness.—From the report sheets more important details can be entered in registers, a process which involves very little cost of trouble and time, and gives a comprehensive view of the facts that cannot be attained in any other way. Each disease should have its separate register, the columns of which may be headed as follows:—

No.
District.
Address.
Name of patient.
Occupation.
Sex.
Age.
Date of onset.
,, ,, notification.
,, ,, receipt of ditto.
,, ,, removal to hospital.
,, ,, disinfection.

Result, and date of recovery

or death.

Place of work or school.

Date of last attendance at ditto.

Milk supply.

Water supply.

Sanitary notes.

Reference to other cases.

Other cases in household.

Medical attendant.

General notes, as to precautions, history, etc.

(Vaccination: date and condition.)

If such registers are kept systematically, any suspicious grouping of cases in any given locality, or in connection with any particular water supply, milk supply, school, or workplace, cannot fail to attract attention at once.

Disease maps.—Maps showing the distribution of disease according to locality can be readily prepared from the information given in the report sheets or

registers. A large-scale map is needed for crowded listricts. Each disease being represented by a particular colour, a spot of this colour or a coloured pin is placed on every house invaded by the disease in question. Each map may serve for one or more diseases, and for a whole year, or such shorter time as may be determined.

Isolation of the patient.—Hospital isolation should be urged in all cases where practicable, and can be enforced under the conditions specified in the Public Health Act, 1875, s. 124, and the Public Health (London) Act, 1891, ss. 66 and 67. Much may be done by persuasion, by pointing out the benefit to be derived by the patient as regards nursing, food, and greater liberty during convalescence, and by explaining the protracted course of the disease, the risk of infection, and the inconvenience or impossibility of maintaining due isolation at home. There will rarely be much difficulty in removing small-pox cases (see pp. 224-234).

Failing removal to hospital, the arrangements for **home isolation** must be made as complete as possible, but in an ordinary household of the working class due isolation is impracticable, except perhaps at first. In large households it may be possible to obtain complete isolation and to maintain it through-

out the illness upon the lines indicated below.

(i.) Arrangement of the sick room.—The patient should be kept in one or two rooms, preferably at the top of the house, or in a detached wing, exclusively set apart for his use. A fire should be kept burning for the sake of ventilation and destroying waste material, even when not required for heating purposes. It tends to secure an indraught at the door, but provision must be made for the constant entry of fresh air by the window or other ventilator. The door must be kept shut. A sheet is often hung

outside the door, and is useful in some ways; but little is gained by wetting it with "disinfectant" solutions, since any infected air which might escape would pass under or round it rather than through it. No one except the persons in actual charge of the patient should be allowed to enter the room. No one should leave the room without washing hands and changing outer garments, and an adjoining room may be used for this purpose. No food should be taken out of the sick room, nor any clothing or utensils, without previous disinfection. Any effluvia in the room should be prevented or removed by free ventilation; and, if necessary, the air may be "sweetened" by the moderate use of sanitas, thymol, chloride of lime, or other deodorant, in the form of spray; but deodorants should not be regarded as disinfectants, and in other parts of the house they serve no useful purpose whatever.

(ii.) Disinfection during the illness.—A plentiful supply of 1 per 1,000 mercuric chloride solution (or 10 per cent. carbolic solution) should be provided. Disinfection of excreta and secretions may be attempted by adding an equal bulk of the solution, which may with advantage be placed in the vessel before receiving the excreta. Old linen should be used in place of pocket-handkerchiefs, and after use thrown into the fire. Linen, etc., requiring washing should be placed in a large vessel filled with the disinfectant solution; such articles should subsequently be rinsed in water before washing, to remove the disinfectant. All utensils should be similarly disinfected before leaving the room.

(iii.) Final disinfection at the end of the case. (a) All linen, cotton, and silk articles which can be removed should be boiled for ten minutes, but exception must be made in respect of certain dyed fabrics and other special articles; (b) all movable textile materials which cannot be boiled—including blankets and other

woollen materials, beds, pillows, mattresses (not, of course, spring mattresses), curtains, hangings, carpets, rugs, etc., must be removed and disinfected by steam; (c) the remaining articles must be laid as open as possible, and the room fumigated by sulphurous acid, from aldehyde, or chlorine; (d) all furniture and other movable articles should be taken out into the open air, and brushed, washed, or beaten; (e) the walls, ceiling, floor, and all surfaces must be cleansed, sprayed, or washed, especial attention being given to crevices and corners. It is an additional safeguard to strip off the paper, and to limewash the ceiling and walls.

None of these precautions can safely be omitted when a patient suffering from small-pox, scarlet-fever, diphtheria, typhus, measles, whooping-cough, mumps, rötheln, or varicella, is kept in a household that contains susceptible persons. In enteric fever, however, the infection appears to be limited to the excretions, with little tendency to aërial diffusion, and a lless rigid isolation may suffice if strict attention is paid to the disinfection of everything that has been in contact with the patient, especially bedding, clothing, and other articles that almost inevitably become infected.

In enteric fever and cholera the evacuations should be regarded as capable of communicating an infectious quality to any nightsoil with which they are mingled in privies, drains, or cesspools; and after such disinfection of them as is practicable, they should be disposed of without delay and under the safest conditions that local circumstances permit. They should not be thrown into any fixed privy receptacle, and above all, they must never be cast where they can run or soak into sources of drinking water.

In the event of death taking place from an

infectious disease, the body should as soon as possible be placed in a coffin with chlorinated lime or other suitable disinfectant, and should be buried (or cremated) with no longer delay than is necessary to allow the fact of death to be verified. Holding of "wakes," large funeral assemblages, and exposure of the corpse to visitors, are especially to be avoided, as is also borrowing of mourning dress for the occasion of the funeral. (See also L.G.B. Memorandum, 1905.)

Disinfection should always be carried out under the direction of the sanitary staff at the end of the case, but disinfectants may be supplied gratuitously by the S.A. for use during the illness. Clothing and all movable articles should either be boiled at home, or removed to a disinfecting station and treated by steam or hot air; rooms, and articles that cannot be removed, should be disinfected in the manner described.

When the patient is removed to hospital the premises may be disinfected at once, but the occurrence of any later case will, of course, necessitate a

repetition of the process.

Home cases offer more difficulty. The final disinfection cannot usefully take place until after the end of the infectious stage, as determined by the written certificate of the medical attendant or the judgment of the **M.O.H.** It is desirable, in order to prevent misunderstanding, to fix a minimum interval between onset and final disinfection, which in scarlet-fever should not be less than six weeks, and to explain this and the reasons for it upon the printed notices left at the house.

Sanitary defects must be remedied, and any clue or even suspicion as to milk or water infection must be followed up at once. It may be necessary to communicate with employers or other persons, if there is reason to anticipate neglect of due precautions against

conveying infection.

In outbreaks of small-pox it is necessary to provide for immediate vaccination or revaccination, and to urge it not only upon the inmates of the infected house, but also upon all who have been in contact with the patient, and upon his neighbours and fellow workmen.

Quarantine.—If the patient is removed to hospital and disinfection properly carried out, it is only necessary to keep the household under observation for the maximum period of incubation of the disease in question. If the case is treated at home, the household must be kept under a certain degree of quarantine until the last case has ceased to be infectious

and the final disinfection has been completed.

During the quarantine period no children from the house should be allowed to attend school, but this restriction may be relaxed if the case is one of enteric fever. The same exclusion should be insisted upon as regards certain occupations, notably milk traffic, nursing, washing, and businesses involving the handling of food; and this is especially important when the work in question is carried on in the house. Tailoring and all forms of seamstress's homework should be forbidden. Strict injunctions should be given against admitting persons, especially children, into the infected house, but there is no power to enforce this byelaw, or to inflict punishment in case of default. Exposure of infected persons or articles in any public place is, however, dealt with under s. 126 of the Public Health Act, 1875, and s. 68 of the London Act, 1891. As regards small-pox, the quarantine should be rigidly enforced, and no member of an infected household should be allowed to follow his business until the period of danger is over.

In some towns quarantine stations are provided; and are usually attached to the isolation hospitals. The sick person having been removed to hospital, free lodging is offered to the rest of the household during the quarantine period, and meanwhile the house and its contents are thoroughly disinfected. This plan has been adopted in many towns in cases of small-pox, and its utility is obvious.

In all these matters, and especially in the removal of the patient to hospital, due consideration should be given to the opinion and advice of the medical

attendant by whom the case has been notified.

Cautionary and advisory notices may be left at the house, explaining in simple language the precautions that are necessary, and the penalties attending neglect of them. Such leaflets should include directions as to isolation and disinfection, and instruction to report immediately any further cases. It is convenient also to leave a post-card addressed to the M.O.H., to be signed by the medical attendant when in his judgment the period of infection is at an end, and final disinfection of the premises is needed.

Quarantine, in the original sense, is not observed in this country. The special precautions directed against the importation of infection from abroad are those set forth in the new **L.G.B.** Order of September 9th, 1907, summarised below.

Regulations as to cholera, yellow fever, and plague.—The Customs Officer, who is the first to board the ship, must as far as possible ascertain whether it is "infected" or "suspected," and if he has reason to believe that it is, or that it has come from an infected port, he must obtain from the master (or from the surgeon if there be one) a written statement in prescribed form declaring the occurrence or non-occurrence of cases or suspected cases during the voyage. If he finds the ship to be infected he must order the master to anchor, and must give notice to the S.A. of the port at which the ship is about to call.

If from such warning, or from other information, the M.O.H. has reason to believe that any ship within the jurisdiction of his S.A. is "infected" or "suspected," he must forthwith visit and examine it; and may do so if it comes from an infected

port. If he finds that there is or has been a case (of one of the hree diseases in question) on board, he must certify accordingly to the master, who is thereupon bound to moor in the blace appointed. The M.O.H. must then examine every person on board, none being allowed to leave the ship until the examination is made. All who are found to be suffering from such disease are to be removed to the hospital or place provided by the S.A., if their condition admit of it, and must not leave such place until the M.O.H. certifies that they are free from the disease. If they cannot be removed, the ship remains subject to the control of the M.O.H., without whose written consent the infected persons cannot leave the ship. Persons certified by the M.O.H. to be suffering from an Illness which he suspects may prove to be one of the three, may be detained either on the ship or in some place provided by the S.A., for not more than two days, in order that it may be ascertained whether their illness is or is not of the kind suspected. No person, not certified as above, is to be permitted to land, unless he satisfies the M.O.H. as to his name, place of destination, and address at such place; and the M.O.H. must give such names and addresses to the clerk of the S.A., who must transmit them to the S.A. of the districts in question. The M.O.H. must give directions and take such steps as may appear to him to be necessary for preventing the spread of infection, and the master of the ship must carry out such directions as are given to him. In the event of a case ending fatally on board, the master must, at the direction of the S.A., either bury the body at sea, properly weighted, or deliver it to the S.A. for interment. must disinfect, and if necessary destroy, the clothing, bedding, and other articles of personal use likely to retain infection, which have been used by infected persons; and disinfect the ship, and disinfect or destroy all articles therein probably infected, according to the directions of the M.O.H. If he believes the ship to be infected, or to come from an infected port, the M.O.H. may order the bilge water and water ballast to be pumped out before the ship enters dock; or may cause the water-ballast tanks to be sealed, if emptying them would endanger the ship. He may order all casks or tanks containing drinking water to be emptied and cleansed, on the S.A. providing a proper supply.

Where a vessel is not infected, but has passengers on board who are in a filthy or otherwise unwholesome condition, the **M.O.H.** may certify to the master that in his opinion it is desirable, with a view to checking the introduction or spread of cholera, yellow fever, or plague, that no persons should be

allowed to land until they have satisfied him as to their names and places of destination, and addresses at such place. Thereupon the same measures are to be adopted as in case of persons

permitted to leave an infected ship.

The master of a ship "infected" with plague or "suspected" of it shall, under the direction of the M.O.H., employ suitable means for the effectual destruction of rats. The same applies to ships coming from plague-infected ports, or those upon which there has been an unusual mortality among rats. In ships "infected" by yellow fever, or "suspected" of it, or which have come from parts infected with that disease, the master shall effectually destroy mosquitoes and their larvæ to the satisfaction of the M.O.H., who shall certify the action taken.

The master of an "infected" or "suspected" ship must hoist at the mast-head a large yellow and black flag, borne quarterly, when the ship is within three miles of the coast of England and Wales, and at night, at a height of not less than 20 feet above the hull of the ship, a night signal of three lights in triangular form, six feet apart, the top light white and the

two lights at the base red in colour.

The expression "infected ship" means a ship which at the time of its arrival from a foreign port has on board a case of cholera, yellow fever, or plague, or which has had on board a case of cholera or plague within a period of seven days, or of yellow fever within a period of eighteen days, prior to that time. The expression "suspected ship" means a ship which has had on board during the voyage, or during the stay of the ship in the port of departure, or in a port in the course of the voyage, a case of cholera, yellow fever, or plague, but which has not had on board a fresh case of cholera or plague within a period of seven days or of yellow fever within a period of eighteen days prior to the time of its arrival from a foreign port.

The L.G.B. have from time to time issued and revoked Orders of a similar nature to the above, both for ships coming foreign and for outward bound and coasting ships, under powers conferred on the Board by the Public Health Acts 1875, 1891, 1896, and 1904.

CHAPTER XVII

MEDICAL OFFICERS OF HEALTH AND SANITARY INSPECTORS

MVERY Sanitary Authority is required (Public Health Let, 1875, secs. 189, 190) to appoint one or more medical officers of health and an inspector of nuisances. I'wo or more S.A. may appoint the same M.O.H.; and, apart from this, the L.G.B. is empowered so 286) compulsorily to unite districts for the purpose of appointing a M.O.H. County Councils are authorised, by s. 17 of the Local Government Act, 1888, so appoint (county) medical officers of health, who are forbidden to hold other appointments or engage an private practice without the written consent of the Council.

If any part of the salary of the M.O.H. of a cocal authority is repaid, the L.G.B. has the same cowers in regard to qualification, appointment, duties, alary, and tenure of office as it has in the case of a coor Law medical officer (Public Health Act, s. 191).

Qualifications.— The Orders issued by the L.G.B. require that the M.O.H. shall be registered under the Medical Act of 1858, and be qualified by law o practise both medicine and surgery in England and Wales. The powers of the Board in this respect are practically superseded by the Local Government Act, 888, s. 18 of which requires that (except when the L.G.B., for reasons brought to their notice, may see fit in particular cases especially to allow) every M.O.H. appointed after the passing of the Act shall be legally qualified for the practice of medicine, surgery, and midwifery; and further, by the same section, he must, if appointed after the 1st of January, 1892, to a district having at the last

census 50,000 inhabitants or more, appear in the Medical Register as the holder of a diploma in Sanitary Science, Public Health, or State Medicine under s. 21 of the Medical Act, 1886; or have been, during some three consecutive years prior to 1892, a medical officer of a district with a population (according to the last census) of not less than 20,000; or have been for not less than three years a medical

officer or inspector of the L.G.B.

Tenure of office. The L.G.B. have no control over the tenure of office unless a portion of the salary is repaid to the S.A.* If the M.O.H. be appointed under the latter condition, he may continue to hold office for such period as the S.A. may, with the approval of the L.G.B., determine, or until he die or resign, or be removed by such Authority with the consent of the L.G.B., or by the L.G.B. The S.A. may suspend him, but must forthwith report their action, together with the cause, to the L.G.B., the latter Board having the power to remove the suspension. If a S.A. desire to change the duties or salary of an officer (or any change be made in the extent of the district), and he decline to acquiesce therein, they may give him six months' notice to determine his appointment, but only with the consent of the L.G.B.

Salary .- This is only controlled by the L.G.B. in cases as above; the salary has then to be approved by the L.G.B., and with the consent of this Board a reasonable compensation may be paid by the Authority on account of extraordinary services or other unforeseen or special circumstances in connection with the duties of the M.O.H. or the necessities of the district.

^{*} These contributions, amounting to half the salary, are paid by County Councils on the certificate of the L.G.B., if claimed by the S.A., and if copies of the reports have been duly received by the Board and by the County Council.

Duties. - The duties of medical officers of health ss prescribed by the L.G.B. are the same whether a ontribution is made to the salary or not, except that n the latter case the officer is obliged to report his ppointment within seven days to the L.G.B.

The following duties are prescribed by the Board's Order of 1891 (Art. 18), for every medical officer of mealth appointed or reappointed after that date:-

(1) He shall inform himself as far as practicable respecting Ill influences affecting, or threatening to affect, injuriously the bublic health within the district.

(2) He shall inquire into and ascertain by such means as are t his disposal the causes, origin, and distribution of diseases within the district, and ascertain to what extent the same have epended on conditions capable of removal or mitigation.

(3) He shall, by inspection of the district, both systematially at certain periods, and at intervals as occasion may equire, keep himself informed of the conditions injurious to

nealth existing therein.

(4) He shall be prepared to advise the S.A. on all matters affecting the health of the district, and on all sanitary points involved in the action of the S.A.; and in cases requiring it ne shall certify for the guidance of the S.A. or of the justices as to any matter in respect of which the certificate of a M.O.H. or a medical practitioner is required as the basis or n aid of sanitary action.

(5) He shall advise the S.A. on any question relating to nealth involved in the framing and subsequent working of such byelaws and regulations as they may have power to make, and as to the adoption by the S.A. of the Infectious Disease (Prevention) Act, 1890, or of any section or sections of such Act.

- (6) On receiving information of the outbreak of any contagious, infectious, or epidemic disease of a dangerous character within the district, he shall visit the spot without delay, and inquire into the causes and circumstances of such outbreak, and in case he is not satisfied that all due precautions are being taken, he shall advise the persons competent to act as to the measures which may appear to him to be required to prevent the extension of the disease, and take such measures for the prevention of the disease as he is legally authorised to take under any statute in force in the district, or by any resolution of the S.A.
 - (7) Subject to the instructions of the S.A., he shall direct

or superintend the work of the inspector of nuisances in the way and to the extent that the S.A. shall approve, and on receiving information from the inspector of nuisances that his intervention is required in consequence of the existence of any nuisance injurious to health, or of any overcrowding in a house, he shall, as early as practicable, take such steps as he is legally authorised to take under any statute in force in the district, or by any resolution of the S.A., as the circumstances of the case may justify and require.

(8) In any case in which it may appear to him to be necessary or advisable, or in which he shall be so directed by the S.A. he shall himself inspect and examine any animal, carcase, meat, poultry, game, flesh, fish, fruit, vegetables, corn, bread, flour, or milk, and any other article to which the provisions of the Public Health Act, 1875, in this behalf shall apply, exposed for sale, or deposited for the purpose of sale, or of preparation for sale, and intended for the food of man, which is deemed to be diseased, or unsound, or unwholesome, or unfit for the food of man; and if he finds that such animal or article is diseased, or unsound, or unwholesome, or unfit for the food of man, he shall give such directions as may be necessary for causing the same to be dealt with by a justice according to the provisions of the statutes applicable to the case.

(9) He shall perform all the duties imposed upon him by any byelaws and regulations of the S.A., duly confirmed, where confirmation is legally required, in respect of any matter affecting the public health, and touching which they

are authorised to frame byelaws and regulations.

(10) He shall inquire into any offensive process of trade carried on within the district, and report on the appropriate means for the prevention of any nuisance or injury to health therefrom.

(11) He shall attend at the office of the S.A., or at some other appointed place, at such stated times as they may direct.

- (12) He shall from time to time report in writing to the S.A. his proceedings, and the measures which may require to be adopted for the improvement or protection of the public health in the district. He shall in like manner report with respect to the sickness and mortality within the district, so far as he has been enabled to ascertain the same.
- (13) He shall keep a book or books, to be provided by the S.A., in which he shall make an entry of his visits, and notes of his observations and instructions thereon, and also the date and nature of applications made to him, the date and result of the action taken thereon, and of any action taken on previous

reports; and shall produce such book or books, whenever

required, to the S.A.

- (14) He shall also make an annual report to the S.A. up to the 31st of December in each year, comprising a summary of • the action taken, or which he has advised the S.A. to take, during the year for preventing the spread of disease, and an account of the sanitary state of his district generally at the end of the year. The report shall also contain an account of the inquiries which he has made as to conditions injurious to health existing in his district, and of the proceedings in which hie has taken part or advised under any statute, so far as such proceedings relate to those conditions; and also an account of the supervision exercised by him, or on his advice, for sanitary pourposes, over places and houses that the S.A. have power to regulate, with the nature and results of any proceedings which may have been so required and taken in respect of the same luring the year. It shall also record the action taken by him, or on his advice, during the year, in regard to offensive trades, to dairies, cowsheds, and milkshops, and to factories and workshops. The report shall also contain tabular statements on forms to be supplied by the L.G.B. or to the like effect) of the sickness and mortality within the district, classified according to diseases, ages, and localities. Provided that if the M.O.H. shall cease to hold office before December 31st in any year, he shall make the like report for so much of the year as shall have expired when he ceases to hold office.
- (15) He shall give immediate information to the L.G.B. of any outbreak of dangerous epidemic disease within the district, and shall transmit to the Board a copy of each annual report and of any special report. He shall make a special report to the L.G.B. of the grounds of any advice he may give to the S.A. with a view to the closure of any school or schools, in pursuance of the Code of Regulations approved by the Education Department, and for the time being in force.

(16) At the same time that he gives information to the L.G.B. of the outbreak of infectious disease, or transmits to them a copy of his annual or any special report, he must give the like information, or transmit a copy of such report, to the County Council of the county in which his district is situated.

- (17) In matters not specifically provided for in this Order he shall observe and execute any instructions issued by the L.G.B., and the lawful orders and directions of the S.A. applicable to his office.
- (18) Whenever the L.G.B. shall make regulations for all or any of the purposes specified in s. 134 of the Public

Health Act, 1875, and shall declare the regulations so made to be in force within any area comprising the whole or any part of the district, he shall observe such regulations, so far as the same relate to or concern his office.

The duties of the medical officer of health to a Port S.A. are defined by the L.G.B. in terms which are closely parallel to those given above, omitting the references to regulated trades and inspection of food, and substituting "ships" for "houses," and "shipping within the district" for "district."

"He shall inform himself as far as practicable respecting all conditions affecting or threatening to affect injuriously the health of crews and other persons on ship-board within the district He shall inquire into and ascertain by such means as are at his disposal, the causes, origin, and distribution of diseases in the ships and other vessels within the district, and ascertain to what extent the same have depended on conditions capable of removal or mitigation He shall, by inspection of the shipping in the district, keep himself informed of the condition injurious to health existing therein On receiving information of the arrival within the district of any ship having any infectious or epidemic disease of a dangerous character on board, or of the outbreak of any such disease on board any ship within the district, he shall visit the vessel without delay, and inquire into the causes and circumstances of such outbreak, and advise the persons competent to act as to the measures which may appear to him to be required to prevent the extension of the disease, and so far as he may be lawfully authorised to assist in the execution of the same On receiving information from the Inspector of Nuisances that his intervention is required in consequence of the existence of any nuisance injurious to health, or of any overcrowding in a ship, he shall, as early as practicable, take such steps authorised by the Public Health Act, 1875, on that behalf, as the circumstances of the case may justify and require When any vessel within his district has had dangerous infectious disease on board, he shall give notice thereof to the M.O.H. of any port within the United Kingdom whither such vessel is about to sail."

The substance of Clauses 4, 5 (omitting the reference to the Infectious Disease Prevention Act), 9, 11,

2, 13, 14 (omitting the reference to trades, etc.), 15 omitting the reference to schools), 17, and 18, of the nuties of M.O.H., is applicable to Port M.O.H., utatis mutandis.

The statutory duties of county medical officers of

ealth have not yet been authoritatively defined.

Annual reports .- The following instructions ave been issued by the L.G.B., 1907:—

Every M.O.H., appointed under Order of the L.G.B., required to make an annual report with regard to each anitary district, or division of a district, which is under his superintendence. This report is to be for the year ending the 11st of December, or, if the officer at that date has not been in office for a whole year, then for so much of the year as has lapsed since his appointment. The report is to be made to he Council by whom he is appointed, and the M.O.H. nimself should send a copy of it to the L.G.B. and to the County Council or County Councils of the county or counties within which his district may be situated. It should be made as soon as practicable after the expiration of the year to which t relates. The M.O.H. ought not, in general, to have any difficulty in doing this within a month or six weeks; but if from any special circumstances the report cannot be completed within six weeks, it should be understood that the delay must not be indefinite, and that the report should be in the hands of his Council, and of the Board, within, at most, three months from the end of the year. Any special circumstances preventing the delivery of the Board's copy within two months from the end of the year should at once be reported to the Board. The Board's copy of the report should be forwarded to them when the original is sent to the Council, except where the report is likely to be printed by order of the Council. In such cases the Board need only be supplied with a printed copy. It is very desirable that the annual report should be printed, for the sake of facility of reference and in order that a supply of copies may be available for distribution among the Town or District Councillors and other persons interested.

Article 18 (s. 14) of the Board's Order of March, 1891, respecting the duties of the M.O.H., specifies the infor-

mation to be contained in the annual report.

The report should be chiefly concerned with the conditions affecting health in the district and with the means for improving those conditions. It should contain an account, brought up to the end of the year under review, of the sanitary circumstances of the district, and of any improvement or deterioration in these circumstances which may have occurred during the year. Care should be taken to report fully and explicitly on the influences affecting or threatening to affect injuriously the public health in the district, and on the action which has been taken, or which may still be needed, with a view to combat those influences. It is of especial importance that the M.O.H. should record what action has been taken to remedy unhealthy conditions which have been reported by him in previous annual reports, or in special reports presented during the year under review, and that attention should be called afresh, year by year, to such as remain unremedied.

As subjects concerning which the Board desire to obtain, through annual reports of the M.O.H., not only definite general information, but record also of particular changes of condition that may have occurred incidentally or by action of the local authority, the following deserve to be especially

borne in mind:-*

General and physical features of the district and conditions of its population.

The chief occupations of the inhabitants and the influence

of any particular occupation on public health.

House accommodation, especially for the working classes: its adequacy and fitness for habitation. Sufficiency of open space about houses, and cleanliness of surroundings. Supervision over erection of new houses. Action under the Housing of the Working Classes Act.

Sewerage and drainage: its sufficiency in all parts of the district. Condition of sewers and house drains. Method or methods of disposal of sewage. Localities where improvements

are needed.

Pollution of rivers and streams in the district; the sources and nature of such pollution and any action taken to check it.

Excrement disposal: system in vogue; defects, if any. Removal and disposal of house refuse—whether by public

scavenger or occupiers: frequency and method.

Water supply of the district or its several parts: its source (from public service or otherwise), nature (river water, well water, upland water, etc.), sufficiency, wholesomeness, and

^{*} This list is, of course, not exhaustive. Additional details are:—Common lodging-houses, tenement-houses, back-to-back houses; House drainage and connections; Burial grounds; Action taken under the above, and under the Sale of Food and Drugs Acts; Canal Boats Act; Meteorological data; Prevalence of epizootic or epiphytic disease; Vaccination (latest available returns); Legal proceedings.

reedom (by special treatment or otherwise) from risks of

poollution.

Places over which the Council have supervision, e.g., hodging-houses, slaughterhouses, dairies, cowsheds, and milk-shops, bakehouses, factories and workshops, and offensive trades.

Schools, especially public elementary schools; sanitary condition of, including water supply; action taken in relation to the health of the scholars, and for preventing the spread of infectious disease.

Byelaws: steps taken for their enforcement, any need of amendment, or of further byelaws.

Nuisances: proceedings for their abatement-any remain-

ing unabated.

Methods of dealing with infectious diseases: notification; isolation hospital accommodation and its sufficiency; disinfection.

With regard to such points it should be remembered that these reports are for the information of the Board and of the County Council as well as of the Council of the district, and that a statement of the local circumstances and a history of Mocal sanitary questions which may seem superfluous for the

Matter may often be needed by the former bodies.

In reference to schools it is desired to draw particular sattention to the medical inspection of school children, under the Education (Administrative Provisions) Act, 1907. This Act does not confer powers in supersession of those heretofore exercised generally in a public health sense by the S.A. runder previous enactments; rather it is supplementary to existing Public Health Law in that it requires supervision of the health of the individual child. The arrangements to this end tto be made by Local Education Authorities will no doubt vary with the diverse circumstances of different districts, but it is anticipated that in most districts the work will, in accordance with the advice of the Board of Education, be carried out in conjunction with that of the M.O.H., or at least under his Where the M.O.H. is also School Medical supervision. Officer it may be convenient that the annual report which he is required to make in the latter capacity should be issued together with his annual report on the health of his district. It is the desire alike of the L.G.B. and of the Board of Education that the relations of the local S.A. and the Local Education Authority, whether or not these bodies or their officers happen to be identical, should be intimate and cordial in order that the administrative procedures of both bodies should be reciprocally helpful.

Section 132 of the Factory and Workshop Act, 1901,

requires that-

"The M.O.H. of every District Council shall, in his annual report to them, report specifically on the administration of this Act in workshops and workplaces, and he shall send a copy of his annual report, or so much of it as deals with this subject, to the Secretary of State."

The copy should be addressed to the Secretary of State, Home Office, Whitehall, and the foregoing remarks with respect to the transmission of the report to the L.G.B. apply

also to its transmission to the Home Office.

The M.O.H. is required to report on the administration of the Factory and Workshop Act, 1901, only in so far as this administration is in the hands of the District Council and is concerned with matters in his department. In reporting on the sanitary administration of workshops and workplaces he should include an account of the action with respect to factories, workshops, and workplaces taken under the Public Health Acts as well as under the Factory and Workshop Act.

In reporting on the sanitary condition of workshops and workplaces the M.O.H. should state what action has been taken to remedy any defective conditions met with under each heading. He should state whether s. 22 of the Public Health Acts (Amendment) Act, 1890, is in force in the district, and what standard of "sufficiency and suitability" of sanitary accommodation is adopted locally.

The M.O.H., in reporting his proceedings and advice, should put on record whether he has made systematic inspections of his district. By "systematic inspections" are meant inspections independent of such inquiries as the M.O.H. may have to make into particular outbreaks of disease, or into unwholesome conditions to which his attention has been specially called by complaints or otherwise; and such inspections will include the house-to-house inspections which may be necessary in particular localities.

In making systematic inspections, as in much of his other action, the M.O.H. will usually have required the assistance of the Inspector of Nuisances; and the Medical Officer should include in his report an account of the action which, at his instance, the Inspector may have taken for the removal of nuisances injurious to health.

The report should deal with the extent, distribution, and causes of disease, especially of epidemic and notifiable diseases, within the district; and should give an account of any noteworthy outbreaks of such diseases during the year under

review, stating the result of his investigations into their origin and propagation, and the steps taken by him, or on his advice,

with a view to check their spread.

The tabular statements of sickness and mortality in the district during the year, to be made on the forms supplied for the purpose, should be the subject of comment in the text of the report, in so far as deductions from them may assist the Board and the Councils concerned to an appreciation of the lines of action needful in the future.

Sources of information.—Returns of registered deaths should be obtained weekly from each of the registration sub-districts within the area.* These returns should give, as regards every death registered during the week, the name, age, sex, residence, and occupation of the deceased, the nature and duration (so far as stated on the certificate of cause of death) of each primary and secondary cause of death, and the name of the certifying practitioner. "Immediate notice should be given of all deaths from infectious disease in fresh localities, and all groups of deaths from such disease, or from diarrhea, in any localities" (L.G.B. Memorandum, dated June, 1882). A fee of twopence for each return and twopence for each entry is payable by the Authority. Returns should also be obtained from the registrars, at weekly, monthly, or quarterly intervals, of the births registered in such intervals; these returns should distinguish between males and females, and between legitimate and illegitimate births. Deaths of non-residents in the locality must, of course, be distinguished from the rest, and the necessary indications for this purpose

^{* &}quot;Every registrar, when and as required by a S.A. shall transmit by post or otherwise a return, certified under the hand of such registrar to be a true return, of such of the particulars registered by him concerning any death as may be specified in the requisition of the S.A. The S.A. may supply a form of the prescribed character, for the purpose of the return, and in that case the return shall be made in the form so supplied. The registrar making such returns shall be entitled to a fee of twopence, and to a further fee of twopence for every death entered in such return, which fee shall be paid by the Authority requiring the return." (Births and Deaths Registration Act, 1874, s. 28.)

should be given in the returns. Arrangements must be made separately for information of the deaths in distant public institutions of residents in the locality.

Notification returns.—Compulsory notification of certain infectious diseases, voluntary notification of others, notification from schools, etc., all furnish the Medical Officer with information respecting the occurrence of such disease.

Returns of pauper sickness.—An order of the L.G.B., dated February 12th, 1879, requires all district and workhouse medical officers appointed after February 28th, 1879, to furnish the M.O.H. with returns of pauper sickness and deaths, and to notify the outbreak of any dangerous infectious disease. The same is required (by an order dated June 14th, 1879) of medical officers of district schools appointed after June 24th, 1879. The Guardians should be asked to instruct their clerk to copy from the district medical officer's relief lists the new cases which are reported at each meeting of the Guardians, and to forward the same promptly and regularly to the M.O.H., and to instruct the Poor Law medical officers to give to the M.O.H. the earliest possible information of cases of dangerous infectious disease under their charge.

Cases of dangerous infectious diseases occurring on canal boats or in common lodging-houses must be reported to the M.O.H., and the S.A. have power to make byelaws requiring notification of such diseases in "tenement" lodging-houses, and in

vans or tents.

Maps.—A good map is essential. It should show the boundaries of the district, and of each of its subdivisions (whether parishes, townships, registration sub-districts, special drainage districts, or other) which have a bearing upon sanitary administration, and also the main sewers and water-mains. Maps showing the geology of the district, the contour lines, and the different watersheds, should be obtained, if possible. Maps showing the local distribution of disease and

mortality have already been referred to.

Periodical reports, etc.—The annual and quarterly reports of the Registrar-General are indispensable. The annual summaries and weekly returns deal only with the statistics of the large towns. Reports of the inspectors of the L.G.B. affecting any part of the district should, of course, be obtained.

Acts of Parliament. - In addition to any local Acts in force in the district, and all the chief statutes dealing with public health, the M.O.H. should be familiar with any other Acts which the special circumstances of his district may render important, e.g. the

Alkali Act.*

Byelaws and regulations.—The byelaws and regullations in force in the district can readily be obtained,

and are not less important than the preceding.

Census data.—The M.O.H. should be in possession of all the information regarding his district given by ceach census taken since its formation. Arrangements should, if possible, be made at the time when the ccensus is taken to obtain a record of the particulars regarding each "enumeration district" for the use of tthe S.A.

It is desirable to keep registers showing for each village or hamlet the dates and particulars of every death, and of every notified case of infectious disease. In towns this would take the form of a "street and house register." If cases of disease as well as deaths are entered, they may be distinguished by red ink. Registers of notified diseases have already been referred to. Registers of all deaths occurring among certain trades will greatly facilitate any future in-

^{*} Lumley's or Glen's Public Health Acts, Glen's L.G.B. Sanitary Orders, and the Model Byelaws of the L.G.B. are valuable for reference.

vestigation which the M.O.H. may have to make into the influence of those trades upon mortality.

The seemingly formidable records mentioned above, if well arranged in the first instance, and systematically kept, involve much less labour than would appear at first sight. They are invaluable in inquiries regarding the incidence of mortality and disease upon any given locality or occupation, and materially simplify the preparation of the annual report.

Auxiliary agencies.—In addition to the routine work of the M.O.H. there are many auxiliary agencies which may be established in his district or by his S.A. which call for more or less of his attention; such, for example, are voluntary health societies, crêches, infant milk depôts, bacteriological work, chemical analysis, meteorological apparatus, reception houses for infectious "contacts," baths for cleansing of verminous persons, etc.

Infants' Milk Depôts .- Infantile mortality, as we have seen, bears a definite relationship to (a) the feeding of infants, (b) personal care of infants by parents, (c) urban life, and (d) certain meteorological conditions affecting temperature and the dissemination of dust. Other elements enter into the problem, but, so far as municipal action is concerned, those are the four main elements. The problem, in practice, is largely one of polluted milk. The sources of the pollution are not only in unsatisfactory methods of milking, and in storing and conveying the milk supplied, but also in dirty domestic conditions, and particularly in carelessness in the use of feeding bottles. The ideal is breast-feeding, and in respect of the artificial feeding of infants a pure modified milk supply which needs no sterilisation; and towards that end our efforts should be directed. The best possible under existing conditions involves sterilisation or pasteurisation, modification, and protection from home contamination. Such a milk supply should be under medical supervision and adopted only in suitable cases.

It would appear that there are five possible means of supplying a suitable and pure milk for infants—(1) By means of Municipal Milk Depôts similar to institutions at Nancy and Fécamp; (2) By one or more dairymen undertaking to furnish

By obtaining such a supply from some central source; (4) By means of medical "milk commissions," as in the United States; or (5) By means of a voluntary society supplying such milk

funder necessary supervision and control.

The French depôts are, broadly speaking, of two types, the "consultation de nourrissons," as at Nancy, and the "goutte de lait," as at Fécamp. The latter is the prototype of the infants' milk depôts established in this country at Liverpool, Finsbury, and elsewhere, to succour those infants of the poor for whom breast-feeding is practically impossible. The milk is modified, pasteurised, or sterilised, and served in "one-feed" bottles, the amount and kind of modification depending, of course, upon the age of the infant. It is too early to speak of the effect such milk depôts exert upon infant mortality, but there can be no question of their value if properly controlled. It is, of course, essential that the milk should be derived from healthy clean cows, and all the dairying carried out on hygienic principles.

Bacteriological Laboratories.—The great value of early and accurate diagnosis in such diseases as tuberculosis, diphtheria, anthrax, enteric fever, and the bacteriological investigation of water, milk, cheese, of sewage and its effluents, of air, etc., has now become generally recognised, with the result that a Public Health Department is not considered fully equipped unless it is provided with facilities for bacteriological research, either by establishing a municipal laboratory or by utilising a private laboratory such as those in connection with hospitals and medical schools. Facilities should be provided for the free examination of any materials from infectious cases submitted

by medical men.

Duties of sanitary inspectors.—The work of the Sanitary Inspector (or "Inspector of Nuisances") is closely connected with that of the M.O.H., but the broad lines separating them will be apparent from the following definition of the inspector's duties, formulated by the L.G.B.

"(1) He shall perform, either under the special directions of the S.A., or (so far as authorised by the S.A.) under the directions of the M.O.H., or, in cases where no such directions are required, without such directions, all the duties specially imposed upon an Inspector of Nuisances by the Public Health Act, 1875, or by any other statute or statutes, or

by the Orders of the L.G.B., so far as the same apply to his office.

"(2) He shall attend all meetings of the S.A. when so

required.

"(3) He shall, by inspection of the district, both systematically at certain periods, and at intervals as occasion may require, keep himself informed in respect of the nuisances

existing therein that require abatement.

"(4) On receiving notice of the existence of any nuisance within the district, or of the breach of any byelaws or regulations made by the S.A. for the suppression of nuisances, he shall, as early as practicable, visit the spot, and inquire into such alleged nuisance or breach of byelaws or regulations.

"(5) He shall report to the S.A. any noxious or offensive businesses, trades, or manufactories established within the district, and the breach or non-observance of any byelaws or

regulations made in respect of the same.

"(6) He shall report to the S.A. any damage done to any works of water supply, or other works belonging to them, and also any case of wilful or negligent waste of water supplied by them, or any fouling, by gas, filth, or otherwise, of water used

for domestic purposes.

- "(7) He shall from time to time, and forthwith upon complaint, visit and inspect the shops and places kept or used for the preparation or sale of butcher's meat, poultry, fish, fruit, vegetables, corn, bread, flour, milk, or any other article to which the provisions of the Public Health Act, 1875, in this behalf shall apply, and examine any animal, carcase, meat, poultry, game, flesh, fish, fruit, vegetables, corn, bread, flour, milk, or other article as aforesaid, which may be therein; and in case any such article appear to him to be intended for the food of man, and to be unfit for such food, he shall cause the same to be seized, and take such other proceedings as may be necessary in order to have the same dealt with by a Justice: provided that in any case of doubt arising under this clause, he shall report the matter to the M.O.H., with the view of obtaining his advice thereon.
- "(8) He shall, when and as directed by the S.A., procure and submit samples of food, drink, or drugs, suspected to be adulterated, to be analysed by the analyst appointed under the Sale of Food and Drugs Act, 1875, and upon receiving a certificate stating that the articles of food, drink, or drugs are adulterated, cause a complaint to be made, and take other proceedings prescribed by that Act.

"(9) He shall give immediate notice to the M.O.H. of the occurrence within the district of any contagious, infectious, or epidemic disease; and whenever it appears to him that the intervention of such officer is necessary, in consequence of the existence of any nuisance injurious to health, or of any overcrowding in a house, he shall forthwith inform the M.O.H. thereof.

"(10) He shall, subject to the directions of the S.A., attend to the instructions of the M.O H. with respect to any measures which can be lawfully taken by an Inspector of Nuisances under the Public Health Act, 1875, or under any statute or statutes, for preventing the spread of contagious,

infectious, or epidemic disease of a dangerous character.

"(11) He shall enter from day to day, in a book provided by the S.A., particulars of his inspections, and of the action taken by him in the execution of his duties. He shall also keep a book or books, to be provided by the S.A., so arranged as to form as far as possible a continuous record of the sanitary condition of each of the premises in respect of which any action has been taken under the Public Health Act, 1875, or under any other statute or statutes, and shall keep any other systematic records that the S.A. may require.

"(12) He shall, at all reasonable times, when applied to by the M.O.H., produce to him his books, or any of them, and render to him such information as he may be able to furnish with respect to any matter to which the duties of Inspector of

Nuisances relate.

"(13) He shall, if directed by the S.A. to do so, superintend and see to the due execution of all works which may be undertaken under their direction for the suppression or removal of nuisances within the district.

"(14) He shall, if directed by the S.A. to do so, act as Officer of the said Authority as Local Authority under the Contagious Diseases (Animals) Act, 1886, and any Orders or

Regulations made thereunder.

"(15) In matters not specially provided for in this Order, he shall observe and execute all the lawful orders and directions of the S.A., and the Orders of the L.G.B., which may be hereafter issued, applicable to his office."

The above statement of duties is general, and the details must be sought in the Acts, Orders, Bye-laws, and Regulations, and in the instructions of the S.A. It is competent for the S.A. to specialise the work of the Sanitary Inspectors (e.g. inspection of meat, inspection of workshops), or to appoint women in that capacity.

Q*

CHAPTER XVIII

SANITARY LAW

Under this heading it is proposed to give a summary of the principal Acts of Parliament bearing upon public health in general, reserving for consideration in later chapters industrial legislation and local byelaws.

As elsewhere in this volume, S.A. stands for Sanitary Authority, L.G.B. for Local Government Board, and M.O.H. for Medical Officer of Health.

Public Health Act, 1875.—The whole of England and Wales, except London, consists of Urban and Rural Sanitary Districts, under the jurisdiction respectively of Urban and Rural S.A.'s (s. 5), which, in pursuance of the Local Government Act of 1894, are now termed Urban and Rural District Councils. That portion of the area of a Union which is not included in any Urban District constitutes a Rural Sanitary District, and the guardians representing that part of the union are the Rural S.A. (s. 9).

Sewers, except certain private sewers, are vested in the S.A. of the district (s. 13). The S.A. may purchase (s. 14) or construct (s. 15) sewers. They must provide such sewers as are necessary for effectually draining their district (s. 15). Powers are given by s. 27 for the treatment and disposal of sewage. The sewers must be so constructed, covered, ventilated, and kept as not to be a nuisance or injurious to health, and must be properly cleansed (s. 19).

House drainage.—If any house is without a sufficient drain the S.A. must require the owner or occupier to provide a drain, and may prescribe the materials, size, and levels. This drain must lead to

the public sewer if there be any within 100 feet of the site of the house; if not, to a covered cesspool in such position (not under a house) as the S.A. may direct. Failing compliance, the Authority may carry tout the works. No house in an urban district must be built or occupied until proper covered drains have been constructed to the satisfaction of the S.A. The S.A. may make regulations as to the mode in which connections between drains and sewers are to be made. Upon giving due notice, and complying with such regulations, the owner of any premises is entitled to carry drains into the public sewers (sections 21–25).

Excrement and refuse disposal.—It is unlawful tto erect any house without a sufficient water-closet, earth-closet, or privy, and an ashpit with proper doors and coverings, and the same must be provided for any eexisting house on the order of the S.A., who may require a separate closet for each house (sections 35-136). The S.A. may order the provision of separate closets for each sex in factories and workshops where both sexes are employed (s. 38); this applies also to above-ground parts of coal-mines if women and girls are eemployed (Coal Mines Regulation Act, 1872, s. 74)the corresponding parts (pit-banks) of metalliferous mines are factories or workshops. Urban S.A.'s may provide public closets, urinals, or receptacles for refuse (sections 39, 45). Every S.A. must see that all drains, closets, ashpits, and cesspools are properly constructed and kept (s. 40).

On written complaint that any drain, closet, ashpit, or cesspool is a nuisance, the S.A. may authorise their officer to enter the premises (after giving twenty-four hours' notice, except in case of emergency) and open the ground; if any defect is found, the S.A. must serve notice upon the owner or occupier; but if there be no defect, the S.A. must close the ground and make good any damage.

Scavenging and cleansing.—The S.A. may, and when required by order of the L.G.B. shall, themselves undertake or contract for the removal of house refuse, the cleansing of earth-closets, privies, ashpits, and cesspools, and cleansing of streets (s. 42). In that case they are liable to penalty for any delay exceeding seven days after written notice from any householder.

Cleansing of Houses.—If the M.O.H. or two medical practitioners certify that any house or part thereof is so filthy as to endanger health, or that the whitewashing, cleansing, or purifying thereof would tend to prevent infectious disease, the S.A. may require the owner or occupier to cleanse, etc., and in his default may themselves do what is necessary (s. 46).

It is unlawful in an urban district to keep swine in any dwelling-house, or so as to be a nuisance, or to suffer any waste water to remain within any dwelling-house for twenty-four hours after written notice from the **S.A.**, or to allow the contents of any water-closet, privy, or cesspool to overflow or soak therefrom. In case of default, the **S.A.** must abate

such nuisance (s. 47).

An inspector of nuisances in an urban district may give notice to the owner of any accumulation of offensive matter, or to the occupier of the premises whereon it exists, and if not removed within twenty-four hours, it becomes vested in the **S.A.** to be disposed of by them (s. 49). An urban **S.A.** may give public notice requiring the periodical removal of manure or other refuse matter from stables or other premises and enforce the same under penalty (s. 50).

Water supply.—Any S.A. may * construct works for supplying any part of their district with water;

^{*} The wording here is permissive; but see pages 506-507.

or may hire or purchase such works, or contract for a supply (s. 51); and may require houses to be supplied with water when they are without a proper supply, if it can be furnished at a cost not exceeding the water-rate authorised by any local Act, or two-pence a week, or such other cost as the **L.G.B.** may, upon application, determine to be reasonable in the circumstances (s. 62).

Heavy penalties are incurred by persons polluting any stream, pond, etc., with gas washings (s. 68), and a local authority may proceed against anyone polluting any watercourse with sewage (s. 69). A Justice's order may be obtained for the closing of a well or cistern the water of which is used, or likely to be used, for drinking or domestic purposes, when the water is so polluted as to be injurious to health, and the Court may cause the water to be analysed at the cost of the S.A. (s. 70).

Cellar dwellings.—No cellar built or rebuilt since 11848 (the date of a previous Act) can be separately occupied as a dwelling; nor any cellar whatever, unless it comply with the following requirements:

(a) The height must in every part be at least seven feet, three feet of which must be above the level of the adjoining street. (b) An open area at least $2\frac{1}{2}$ feet wide in every part, and 6 in. below the level of the floor, must extend along the whole frontage. It may be crossed by steps, but not opposite the window. (c) The cellar must be drained by a drain at least 1 foot below the floor. (d) There must be proper closet and ashpit accommodation. (e) There must be a fireplace and chimney and (f) a window at least 9 square feet in area, made to open. The window of a back cellar let or occupied along with a front cellar need only be 4 square feet in area (sections 71—73).

Anyone passing the night in a cellar is deemed to

occupy it (s. 74).

Common lodging-houses.*—The S.A. are required to keep a register of common lodging-houses, and to make byelaws in respect of them (sections 76-80). It is unlawful to keep a common lodging-house unless it be registered (s. 77), and this can only be done after it has been inspected and approved (s. 78). The keeper when required is to affix notice of registration to the house.

When the lodging-house is without a proper supply of water, and this can be furnished at a reasonable rate, the S.A. may enforce it (s. 81). The keeper is required to limewash the walls and ceilings in the first week of April and October of every year (s. 82). The S.A. may require the keeper of a house in which beggars or vagrants are received, to make returns of persons who have slept there the previous night (s. 83), and the keeper must always give notice to the M.O.H. and to the Relieving Officer of any case of infectious disease (s. 84).

Houses let in lodgings .- Any S.A. may make byelaws respecting houses let in lodgings or occupied by members of more than one family (s. 90, modified by s. 8 of the Housing of the Working

Classes Act, 1885).

Nuisances.—(s. 91.) The following are nuisances. (1) Any premises † in such a state as to be a nuisance or injurious to health. (2) Any pool, ditch, gutter, watercourse, privy, urinal, cesspool, drain, t or ashpit, so foul or in such a state as to be

† Including buildings or lands.

† "'Drain' means any drain of and used for the drainage of one building only, or premises within the same curtilage" A sewer is a drain receiving the drainage of two or more buildings.

^{*} Lodging-houses in which persons of the poorer class are received for short periods, and, though strangers to each other, are allowed to inhabit one common room. The term does not cover inns, or lodgings let to the middle or upper classes. In London there are Common Lodging Houses Acts, 1851-3, in force, incorporating similar powers.

a nuisance or injurious to health. (3) Any animal so kept as to be a nuisance or injurious to health. (4) Any accumulation or deposit which is a nuisance or injurious to health.* (5) Any house,† or part of a house, so overcrowded as to be dangerous or injurious to the health of the inmates, whether or not members of the same family. ‡ (6) Any workshop, workplace, or domestic factory, not kept in a cleanly state, or not ventilated in such a manner as to render harmless as far as practicable any gases, vapours, dust, or other impurities generated in the course of the work carried on therein that are a nuisance or injurious to health, or so overcrowded as to be dangerous or injurious to the health of those employed therein. § (7) Any fireplace or furnace which does not, as far as practicable, consume the smoke arising from the combustible used therein, and which is used for working engines by steam, or in any manufacturing or trade process whatever; and any chimney (not being the chimney of a private dwelling-house) sending forth black smoke in such quantity as to be a nuisance.

^{*} There is no penalty if the accumulation or deposit is necessary for, and has not been kept longer than is necessary for, the carrying on of any business or manufacture, and if the best available means have been taken for preventing injury to the public health.

[†] Including schools, and also workshops and other buildings in which persons are employed. Domestic factories also are included.

[‡] It is customary to adopt the standards of the Model Byelaws, viz. 300 cubic feet per head, counting two children as one adult; but for workshops and workplaces, see p. 561. In the event of a second conviction for overcrowding within three months, the Court may order the closing of the premises for any period (section 109).

[§] The provisions of this sub-section apply to all buildings, including schools, factories, and workshops, except such as are subject to the special provisions, relating to cleanliness, ventilation, or overcrowding, of the Factory and Workshop Act.

^{||} There is no penalty if the Court is satisfied that such fireplace or furnace is constructed in such manner as to consume as far as practicable, having regard to the nature of the manufacture or trade, all smoke arising therefrom, and that such fireplace or furnace has been carefully attended to by the person having the charge thereof.

Later Acts have extended this definition. Thus workshops or workplaces not kept free from effluvia arising from any drain, closet, or other nuisance (Factory Act 1901, s. 2); workshops in any room of which sufficient ventilation is not provided and maintained, or in which the processes carried on render the floor wet, and the floor is not adequately drained (Factory Act, s. 78-see also p. 559); tents or vans so filthy or overcrowded as to be a nuisance or injurious to health (Housing Working Classes Act, 1885, s. 9), and unfenced quarries (Quarry Fencing Act, 1887, s. 3) or unfenced shafts of disused metalliferous mines (Metalliferous Mines Regulation Act, 1872, s. 13) within 50 yards of a highway, are nuisances under s. 91, 1875.

Procedure in respect of nuisances.—It is (1) the duty of the S.A. to cause to be made from time to time an inspection of their district, to ascertain what nuisances exist, but (2) the S.A. may be put in motion by any person aggrieved, or by any two householders or by any officer of the S.A., or by the Relieving Officer, or by any officer of the police. (3) If satisfied as to the existence of a nuisance, the S.A. must serve a notice on the person responsible, or if he cannot be found, on the owner or occupier, requiring him to abate it, and to execute such works and do such things as may be necessary. Where the nuisance arises from the want or defect of any structural convenience, or where there is no occupier, notice must be served on the owner. If the person causing the nuisance cannot be found, and the owner or occupier is not responsible, the local authority may abate the nuisance. (4) If the person on whom the notice is served fails to comply, or if the nuisance, although abated, is likely to recur, the S.A. may apply to a Justice, who shall summon the person responsible, and may make an order requiring compliance or prohibiting the recurrence of the nuisance, and directing the execution of any works necessary. The Court may further impose a penalty (sections 92-96). Where the nuisance is such as to render the house

nfit for habitation, the Court may order the house be closed, and may rescind this by a further order when satisfied that the house has been made tt for habitation (s. 97). Any person not obeying he order of the Court, or failing to use due diligence, ; liable to penalty; and the S.A. may carry out he order and charge him with the expenses (s. 98). Where the person responsible for the nuisance cannot ee found, the order of the Court may be executed by he S.A. Any matter or thing removed by the S.A. n abating any nuisance may be sold. Where a ruisance within a district is caused by some act or efault beyond its limits, the S.A. may, nevertheless, nstitute proceedings (s. 108).

Ships.—Any ship or vessel lying in any water within the district of any S.A. is subject to their urisdiction, as if it were a house.* If in any other vater, it is deemed to be within such district as may be prescribed by the L.G.B., and in the absence if such prescription, then within the nearest district ss. 110).

Right of entry upon private premises. — The 33.A. and their officers have rights of entry between a.m. and 6 p.m., and in the case of a nuisance rrising in respect of any business, at any hour when wach business is in progress. If admission be refused Justice's order may be obtained (sections 102 and 103).

Offensive trades.—It is illegal to establish an offenlive trade within the district of an urban S.A. without their consent. The offensive trades are: bloodooiler, bone-boiler, fellmonger, soap-boiler, tallow-nelter, tripe-boiler, any other noxious or offensive rade, business, or manufacture (s. 112). An urban

^{*} This section originally brought ships under those provisions of the act that relate to nuisances only; but the Public Health (Ships) Act, 885, extended its scope to include also infectious diseases and hospitals sections 120, 121, 124, 125, 126, 128, 131, 132, and 133).

S.A. may make byelaws with respect to such trades as have been established with their consent,* to prevent or diminish nuisance arising therefrom (s. 113). If the M.O.H., or two medical practitioners, or ten inhabitants, certify that any of the premises named below are a nuisance, or injurious to health, the S.A. must take proceedings against the offender, who is liable to penalty, unless he can show that he has used the best practical means for abating such nuisance, or preventing or counteracting such effluvia. The premises in question are: candle-house, melting-house, melting-place, soap-house, slaughterhouse, any building or place for boiling offal or blood, or for boiling, burning, or crushing bones, or any manufactory, building, or place used for any trade, business, process, or manufacture, causing effluvia (s. 114). The same powers are applicable in the case of nuisances affecting the inhabitants of a district, but resulting from premises situated beyond the limits of the district.

[As regards the prohibition (s. 112) of the establishment of any given trade without the consent of the S.A., it is incumbent upon the prosecuting authority to show that the trade is either one of those specifically mentioned above, or ejusdem generis with them; that is, that it is necessarily an offensive trade apart from neglect or mismanagement. The higher Courts have held that this is the case with rag and bone stores for example, but not with brick-making, manure works, or fish-frying. Nevertheless, a very large number of trade processes to which the above restriction may be considered inapplicable will come within the scope of s. 114, if it can be shown that they are carried on so as to give rise to effluvium nuisances.]

^{*} As this consent only became necessary in 1848, the byelaws can only apply to trades established since that date.

Unsound food.—The M.O.H. or Inspector of Nuisances may, at all reasonable times, examine any unimal,* carcase, meat, poultry, game, flesh, fish, fruit, wegetables, corn, bread, flour, or milk, exposed for sale, or deposited for the purpose of sale, or of preparation for sale, and intended for the food of man; and may seize the same if diseased, unsound, or unwholesome, and take it to a magistrate, who may order it to be destroyed, and inflict a penalty upon the offender. The proof that it was not intended for the food of man rests with the person charged (sections 116, 117). Any person obstructing is subject to penalty (s. 118). On complaint made on oath a Justice may grant a search warrant to the M.O.H. or Inspector.

[Eggs, butter, cheese, and other important articles of food, are not mentioned specifically, and are, therefore, not included in the scope of the Act. No proceedings can be taken in regard to articles already sold. Both these defects are avoided in the Public Health Amendment Act of 1890, and in the Public

Health (London) Act, 1891.

Infectious diseases. — Upon the certificate of the M.O.H. or other medical practitioner that the cleansing and disinfecting of any house or part thereof, and of any articles therein, would tend to prevent infectious diseases, it is incumbent on the S.A. to serve notice upon either the owner or occupier, requiring him to cleanse and disinfect. A penalty is incurred by default, and the S.A. may do what is necessary and recover the costs, or may undertake this duty in the first instance, with the consent of the occupier, at their own cost (s. 120). The S.A. may destroy infected bedding, clothing, or other articles, and give compensation (s. 121); may provide a disinfecting apparatus and disinfect free of charge

^{*} Living animals may be seized under this section.

(s. 122); may provide an ambulance and pay expenses of conveyance of infectious persons to hospital (s. 123).

Where a hospital is provided within convenient distance, a Justice may, on the certificate of a medical practitioner, order the removal of any person who is suffering from any dangerous infectious disorder, and is without proper lodging or accommodation, or lodged in a room occupied by more than one family,* or is on board any ship or vessel (s. 124). The S.A. may make regulations for removing to any available hospital, and for keeping there as long as necessary, any persons brought within their district by vessel who are infected with a dangerous infectious disorder (s. 125).

It is unlawful for any person so suffering to expose himself wilfully, without proper precautions against spreading the disorder, in any street, public place, shop, inn, or public conveyance, or to enter any public conveyance without previously notifying to the owner, conductor, or driver thereof that he is so suffering; or, being in charge of any person so suffering, to expose such sufferer, or to give, lend, sell, transmit, or expose without previous disinfection any bedding, clothing, rags, or other things which have been exposed to infection from any such disorder; but this does not apply to the transmission with proper precautions of articles for the purpose of having them disinfected (s. 126). The owner or driver of a public conveyance so used is required under penalty to have the same immediately disinfected, but he need not convey any person so suffering until he has been paid a sum sufficient to cover any loss or expense incurred by him (s. 127). Any person who knowingly lets for hire any house, or room, in

^{*} This detail is omitted as superfluous in the London Act of 1891; but lodgment in a tent or van is made per se a reason for removal.

thich any person has suffered from such disorder, ithout having it and the contents disinfected to the atisfaction of a medical practitioner as testified by certificate signed by him, is liable to penalty so 128). Any person letting or offering for hire my house, or part of a house, who on being questioned so to the fact of there being, or within six weeks reviously having been therein, any person suffering from any dangerous infectious disorder, knowingly makes a false answer to such question, becomes liable to penalty or imprisonment (s. 129).

Special Regulations of L.G.B.—The L.G.B. may make regulations for the treatment of persons affected with cholera, or any other epidemic, endemic, or intectious diseases, and for preventing the spread of such liseases as well on the seas, rivers, and waters of the United Kingdom, and on the high seas within three miles of the coast thereof, as on land, and may declare by what Sanitary Authorities such regulations shall be enforced and executed (s. 130). Cholera Regulations have been issued under this section (p. 470).

Furthermore, whenever any part of England apmears to be threatened with, or is affected by, any formidable epidemic, endemic, or infectious disease, he L.G.B. may make, and from time to time alter or revoke, regulations for any of the following purposes, viz. for the speedy interment of the dead, for nouse-to-house visitation, for the provision of medical wid and accommodation, for the promotion of cleansing, ventilation, and disinfection, and for guarding against the spread of disease; and may by order leclare all or any of the regulations so made to be in force within the whole or any part of the district of any S.A., and to apply to any vessels whether on inland waters or on parts of the sea within the jurisdiction of the Lord High Admiral of the United Kingdom (s. 134). The S.A.'s are required

to do everything that is necessary to carry out these

regulations.

Hospitals.—Any S.A. may build or contract for the use of hospitals for their district. Two or more S.A.'s may combine for this purpose (s. 131). The S.A. may recover from a patient, who is not a pauper, the cost of his maintenance (s. 132). The S.A. may with the sanction of the L.G.B. themselves provide or contract for a temporary supply of medicine and medical assistance for the poor of their district (s. 133). (See also Isolation Hospitals Acts.)

Mortuaries.—Every S.A. may, and, if required by the L.G.B., must, provide a mortuary, and may make byelaws with respect to its management and charges for its use. They may also provide for the decent and economical interment, at charges to be fixed by such byelaws, of any dead body which may be received into a mortuary (s. 141). A Justice may, on a certificate signed by a practitioner, order to be removed to a mortuary, at the cost of the S.A., (1) the body of anyone who has died of any infectious disease, and which is retained in a room in which persons live or sleep, or (2) any dead body which is in such a state as to endanger the health of the inmates of the house or room in which it is retained. He may direct the same to be buried within a specified time. If the friends fail to comply, it is the duty of the Relieving Officer to bury, and the expenses may be recovered from the proper person (s. 142). The S.A. may provide a place for the postmortem examination ordered by a coroner.

New streets and buildings.—In urban districts all public streets (i.e. those reparable by the inhabitants at large) are vested in the **S.A.**, who shall cause them to be levelled, paved, and repaired, as occasion may require (s. 149). The owners of property abutting on any private street or part of a street may

be required by the (urban) **S.A.** to level, pave, sewer, light, or make good such street or part of a street; and in case of default the **S.A.** may carry out the work and recover the expenses from the owners according to the frontage of their respective premises (s. 150). An urban **S.A.** may make byelaws with regard to new streets and buildings. For the purposes of the Act the re-erection of any building pulled down to or below the ground floor, or the conversion into a dwelling-house of any building not originally constructed for human habitation, or the conversion into more than one dwelling-house of a building originally constructed as one dwelling-house only, shall be considered the erection of a new building (s. 159).

Dangerous structures.*—The surveyor of an urban SS.A., if satisfied that any building or wall is in a ruinous state so as to be dangerous to passengers or to the inmates of neighbouring houses, shall cause at fence to be put up, and shall order the owner forthwith to secure or pull down such building; and in diefault thereof the surveyor may obtain a Justice's order to carry out the necessary works, and may recover the expenses. (Towns Improvement Clauses Act, sections 75–78, incorporated with the Public Health Act by s. 160 of the latter.)

Rain-pipes and eaves-gutters.—An urban S.A. may compel the owner of any building adjoining or near to a street to provide within seven days efficient eavesgutters and rain-pipes. (Towns Improvement Clauses Act, s. 74; Public Health Act, s. 160.)

Slaughter-houses.—An urban S.A. may provide battoirs or slaughter-houses, and if they do so must make byelaws with respect to management and harges (s. 169). They may also license slaughter-

^{*} These and some of the following provisions are not to be found in the ext of the Public Health Act, but in the incorporated portion of the owns Improvement Clauses Act.

houses and knackers' yards, and without their licence no place shall be used for such purposes which was not so used at the time of the passing of the Act, i.e. in 1875. Every place used as a slaughterhouse or knackers' yard shall be registered by the owner or occupier in a book kept by the S.A. An urban S.A. must make byelaws in regard to slaughter-houses and knackers' yards. If any person is convicted of killing or dressing any cattle contrary to the provisions of the Public Health Act, or of the non-observance of any of the byelaws or regulations made under this Act, the Justices before whom he is convicted may suspend the licence for two months or less, and in the event of a second offence may revoke the licence. (Sections 125 to 130 Towns Improvement Clauses Act, incorporated in the Public Health Act by virtue of s. 169.)

A legible notice bearing the words Licensed Slaughter-House or Registered Slaughter-House must be attached and maintained in some conspicuous place on every registered slaughter-house by the owner or

occupier (s. 170).

Combination of sanitary authorities for appointment of Medical Officer of Health .- The L.G.B. may (compulsorily or otherwise) unite two or more districts situated wholly or partially in the same county, for the purpose of appointing a Medical Officer of Health; but no urban district with more than 25,000 inhabitants shall be included without the consent of the S.A., and in the event of any S.A. objecting to being included, the L.G.B. can only include their district by means of Provisional Order subject to confirmation by Parliament (s. 286).

A S.A. may delegate all or any of its powers to a committee of its own members (sections 200 and 201); and a rural authority may form parochial committees consisting wholly or partially of its own members, and

may give to such committees certain limited sanitary functions (s. 202). (See under Loc. Govt. Act, p. 520.)

Urban powers for rural districts.—The L.G.B. may give to a rural S.A., as regards the whole or part of their area, any powers and responsibilities conferred

by this Act upon urban S.A.'s (s. 276).

Special drainage districts.—A rural S.A. may, with the consent of the L.G.B., constitute any portion of their area a special drainage district for the purpose of charging thereon exclusively the expenses of works of sewerage, water supply, or other works which may be declared by the L.G.B. to be special expenses, and thereupon such area shall become a separate contributory place (s. 277). The S.A. have also power to Heclare expenses which they have incurred to be private improvement expenses, and to charge a special private improvement rate in respect thereof upon the estate benefited (sections 213 and 232).

Union of districts for sanitary purposes.—Joint boards may be formed, by order of the **L.G.B.**, for the purpose of water-supply or sewerage, or for other burposes of the Public Health Act, as applied to a district comprising the whole or parts of two or more sanitary districts (s. 279). Under this section joint

Hospital Boards may be formed.

Default of S.A.—If satisfied, after due inquiry, hat any S.A. have made default in providing or mainaining sewers, or in providing and maintaining a upply of pure water,* or in enforcing any provisions of the Public Health Act which it is their duty to unforce, the L.G.B. shall make an order limiting a time for the performance of their duty in the matter. If such duty is not performed in the time allowed, the order may either be enforced by writ of mandamus,

^{* &}quot;In cases where danger arises to the health of the inhabitants from neinsufficiency or unwholesomeness of the existing supply of water, and proper supply can be got at a reasonable cost."

or the L.G.B. may appoint some person to perform the work, and charge all expenses to the authority in default (s. 299).

The 7th section of the Housing of the Working

Classes Act, 1885, has a wider scope.]

Port Sanitary Authority.—The L.G.B. may, by order, constitute any S.A. whose district abuts upon any port in England the S.A. for the whole or for any part of such port. The Board may also combine two or more riparian authorities for the purpose; or may constitute one Port S.A. for any two or more ports. The Authority may be constituted permanently or temporarily. The order constituting the Authority may assign to it any of the powers or duties of an urban S.A. so far as applicable to a Port S.A., and to vessels, waters, or persons within its jurisdiction namely, under sections 91 to 111 (nuisances), 120 to 133 (infectious diseases and hospitals), 134 to 138 (prevention of epidemic diseases), 141 and 142 (mortuaries), 182 to 186 and 188 (byelaws), 189 (appointment of M.O.H. and inspector of nuisances), and other sections.*

The Public Health (Water) Act, 1878, applies to every rural S.A., and also to such urban S.A.'s as the L.G.B. may order (sections 3 and 11). Every rural S.A. shall see that every occupied dwellinghouse within their district has, within reasonable distance, an available and sufficient supply of wholesome water. If the M.O.H. reports that an occupied house is without such supply, and the S.A. are of opinion that such a supply can be provided at a reasonable cost (the interest on which, at 5 per cent.,

^{*} The powers and duties of a Port S.A. and its officers consist primarily of those conferred by such of the specified portions of the Public Health Act as the particular Order may decide. They are extended by the Cholera Regulations (page 470), and the Order prescribing the duties of the Port M.O.H, A ship comes within the definition of "house" for many purposes (page 497).

shall not exceed twopence or threepence per week, as the L.G.B. may, on the application of the S.A., Retermine to be reasonable in the circumstances), the S.A. may require the owner (subject to appeal to the **L.G.B.**) to provide such supply within a specified time, and, in case of default, may themselves carry but the necessary works at his expense. It remains, however, the duty of the S.A. to provide any part of their district with a supply of water, in cases where llanger arises to the health of the inhabitants from the insufficiency or unwholesomeness of the existing supply, and a general scheme of supply is required, and can be got at reasonable cost (s. 3).

The owner of any dwelling-house in a rural district that may be erected or rebuilt from the ground-floor after the date of this Act shall not permit such house to be occupied without a certificate from the S.A. that it is provided with a sufficient and available supply of wholesome water, such certificate to be based on the report of the M.O.H. or inspector

(s. 6). If the S.A. provide a stand-pipe for waterssupply, they may levy water charges upon every dwelling within 200 feet, as if the supply were given on the premises; not, however, upon such houses as have a good supply within reasonable distance, from another source, unless water from the stand-pipe is used by

the inmates (s. 9).

Every rural S.A. must, through their officers or otherwise, provide for the periodical inspection of the water-supply of their district. The same powers of entry upon premises are given as are conferred by sections 102 and 103 of the Public Health Act, 1875, in respect of nuisances (s. 7).

The Public Health Act, 1896, repeals the old Quarantine Act of 1825, and extends the powers of the L.G.B. to make Orders under sections 130 and 134 of the 1875 Act, dealing with epidemic, endemic, or infectious diseases. Accordingly (see p. 470), regulations as to cholera, yellow fever, and plague were issued in an Order dated 1896 and renewed September 9th, 1907.

The Public Health (Ports) Act, 1896, enables the L.G.B. to issue orders making all or any of the provisions of the Infectious Diseases Prevention Act, 1890, applicable to any Port

Sanitary Authority.

The Public Health Acts Amendment Act, 1890, is an "adoptive" Act, of which Part III. is concerned with sanitary matters. Urban S.A.'s may by resolution adopt any part of the Act; rural Authorities can adopt those powers which are not expressly limited to urban districts, and as regards the rest, can apply to the L.G.B. for full urban powers in the same manner as under the 1875 Act.

The following are the chief provisions of Part III. Sewers.—It is unlawful to pass into any drain or sewer (a) any substance which may injure it or impede the flow of its contents; or (b) any chemical refuse, or waste steam, or water or other liquid heated above 110° F., which either alone or in combination with sewage causes a nuisance or is dangerous to health.

Closets, etc.—Where any sanitary convenience is used in common by the occupants of two or more houses, any person fouling or injuring it is liable to penalty; and if nuisance arises from want of cleanliness of any part of it or of the approaches, each of the persons having the right of use is liable to penalty, in the absence of proof as to which of them is in default. Sufficient and suitable sanitary conveniences must be provided in all factories and workshops; with separate accommodation for each sex, where persons of both sexes are employed (s. 22).

Byelaws.—Urban S.A.'s may make byelaws respecting public sanitary conveniences provided by

aem. Section 157 of the 1875 Act is extended so as enable any urban S.A. to make further byelaws oncerning new buildings, upon the following points: adequate water supply to closets; (b) construction floors, hearths and staircases; (c) height of rooms ttended for habitation; (d) paving of yards and open paces in connection with houses; (e) provision of condary approaches to houses, for the purpose of reoving refuse. It is further provided that byelaws specting closets and drainage may be made applible to old as well as new houses. Similar power framing byelaws respecting buildings (under s. 57 as amended) is given to rural S.A.'s, but with ertain exceptions. Apart from this the L.G.B. nn, as already stated, grant full urban powers to ural Authorities. Any S.A. may make byelaws to revent buildings erected in accordance with byelaws om being altered in such a way that if at first so onstructed they would have contravened the byelaws.

An urban **S.A.** may make byelaws respecting the criage through the streets of offensive matter or quid, prescribing (a) certain hours for such removal, proper construction and covering of the receptacle sed for the removal, and (c) the cleaning of any

ace fouled by matters dropped or spilled.

Any S.A., themselves undertaking or contracting or the removal of house-refuse, may make byelaws imposing upon the occupier duties in connection with uch removal.

Polluted sites.—No new building may be erected pon ground impregnated with animal or vegetable latter, or upon which such matter has been deposited, nless such matter has been properly removed or has ecome innocuous. If (in an urban district) any porton of a room is immediately over any privy (not eing a water-closet or earth-closet), cesspool, midden, r ash-pit, it is illegal to occupy it, or suffer it to be

occupied, as a dwelling-place, sleeping-place, work-

room, or place of business.

Cleansing of common passages.—If in an urban district any private court or passage, leading to the back of several buildings separately occupied, is not regularly and effectually swept and kept clean, the **S.A.** may cause it to be swept and cleaned, and recover the expenses from the occupiers.

Pollution of streams by solid refuse.—It is forbidden "to throw or place in any watercourse within any district in which this part of this Act is adopted, any cinders, ashes, bricks, stone, rubbish, dust, filth, or other matter which is likely to

cause annoyance."

Food.—Sections 116—119 of the 1875 Act are extended so as to apply to "all articles intended for the food of man, sold or exposed for sale, or deposited in any place for the purpose of sale, or of preparation for sale." A Justice may condemn any such article and order it to be destroyed, if satisfied that it is diseased, unsound, unwholesome, or unfit for the food of man, even if it has not been formally seized under s. 116.

Slaughter - houses.—Licences granted after the adoption of this Act are to remain in force only for such period, not being less than a year, as the S.A. shall specify in the licence. Every change of occupation of a licensed slaughter-house must (under penalty) be notified in writing to the inspector of nuisances; and notice of this requirement must be endorsed on all new licences. If the occupier of any licensed slaughter-house is convicted under sections 116—119 of the 1875 Act, the Court may revoke the licence.

Common lodging-houses.— A penalty is imposed upon the keeper if he fails to give notice of any case of infectious disease, as required by s. 84 of the 1875

Act (p. 494).

The Act does not apply to London.

The Public Health Acts Amendment Act, 9907, is also an "adoptive" Act in the sense that .A.'s may apply to the L.G.B. to make an Order its enforcement as regards Parts I.—VI. The ct does not apply to London. Parts VII. (Police), III. (Fire Brigade), and IX. (Sky Signs) are encorceable by the Home Secretary. Part I. deals with me general application of the Act. Part II. concerns preets and buildings, and provides for deposit of tans, means of access, repairs, enclosing dangerous aces, fencing lands, re-naming, temporary buildings, to. The only strictly sanitary clause in this Part is are the effectual paving and drainage of subsoil or urface of yards of dwelling-houses.

Part III. contains the chief sanitary provisions. 35 adds to the nuisances enumerated in the Public Kealth Act, 1875, any cistern used for the supply of ater for domestic purposes so placed, constructed, or cept as to render the water therein liable to conmination likely to cause risk to health; any gutter, rain, shoot, stack-pipe so defective as to cause ampness to a building; and any deposit of material n or on any building or land which shall cause a impness dangerous or injurious to health. Rainater pipes must not be used as soil pipes (s. 36), nor ust water or stack pipes be used as ventilating hafts (s. 37). S. 39 deals with the provision of afficient and proper water closet and slop-closet ccommodation in new and old buildings, and power given to S.A.'s to require the provision of rinals to public-houses and similar establishments. in report of defective or suspicious drainage, tests may be applied (except the water test) subject to onsent of owner or occupier, or by order of a court f summary jurisdiction, and provision is made for lling up old cesspools. The S.A. may provide and maintain public sanitary conveniences in or under

streets. By s. 48 the Authority must remove trade refuse if required by owners or occupiers so to do.

Various new powers are given in Part IV. as to infectious diseases. If any person knows that he is suffering from an infectious disease he may not engage in any industrial occupation unless he can do so without risk of spreading infection. If the M.O.H. certifies to the S.A. that milk is suspected of conveying infection, the S.A. may require the dairyman to furnish a list of sources of his milk supply, for which he shall be paid sixpence for every 25 names. Every dairyman shall notify to the S.A. the occurrence of all cases of infectious disease among his servants. Infected clothes may not be sent, unless disinfected, to a public laundry. S. 57 provides that no child suffering from infectious disease or who has been exposed to infection shall, after notice from the M.O.H., be sent to school without a certificate from the M.O H., and if the S.A. require it the head teacher of a school shall furnish (for sixpence per 25 names) a complete list of the names and addresses of the scholars in a school in which any scholar is suffering from an infectious disease. Persons suffering from infectious diseases shall not use any book from a public library or enter any public vehicle, in the latter case both patient and driver being held responsible. If any public vehicle be so used it must be disinfected. Extended powers for cleansing and disinfecting premises are also given; for the management of common lodging-houses; for the provision of recreation grounds; for policing streets; for fire brigades; and sky signs.

The Infectious Diseases Prevention Act, 1890, is another adoptive or optional Act; and any extra-metropolitan S.A. may by resolution adopt all or any of its sections, and may by further resolution rescind them. For London its leading provisions are

ontained in the Public Health (London) Act. The infectious diseases to which this Act applies are those pecified in (or added to) the Infectious Diseases Notication Act. The principal provisions are as follows:—

Milk supplies .- If the M.O.H. has reason to believe that the consumption of milk from any dairy* within or without his district) has caused or is likely o cause infectious disease to any person residing in he district, he may (if authorised by a Justice having urisdiction in the place where the dairy is situate) inspect the dairy. He may further, if accompanied by a veterinary surgeon, inspect the animals therein. If after inspection he is of opinion that infectious disase is caused by consumption of the milk, he must eport to the S.A., forwarding also any report furhished to him by the veterinary surgeon. The S.A. may then give not less than 24 hours' notice to the hairyman to appear before them, and show cause why The supply of the milk in their district should not be prohibited. If in their opinion he fails to show such ause, they may order accordingly, and must give motice of the facts to the S.A. and the County Council of the district in which the dairy is situate, and to the G.G.B. The order must be forthwith withdrawn on he S.A. or the M.O.H. being satisfied that the milk upply has been changed, or that the cause of infection has been removed. Penalties are provided for conravention of this section of the Act.

Disinfection.—Where the M.O.H. or any registered medical practitioner certifies that the cleansing and disinfection of any house, or part thereof, and of any articles therein, would tend to prevent infection, the S.A. may, after 24 hours' notice to the owner or occupier, proceed to carry out such disinfection or

^{*} Including any farm, farmhouse, cowshed, milk store, milk-shop, or ther place from which milk is supplied or in which it is kept for the surpose of sale.

S.A. that he will, within a period fixed in the notice, himself carry out the work to the satisfaction of the M.O.H. If he fail to do this within the specified period, it is to be done by the officers of the S.A., under the superintendence of the M.O.H., and the expenses may be recovered. Power of entry between 10 a.m. and 6 p.m. is given for the purposes of this section. Where the section is adopted, the 120th section of the Public Health Act, 1875, is repealed.

The S.A. may, by written notice, require (under penalty) any infected clothing or other articles to be delivered to their officer for disinfection. The S.A. must take away, disinfect, and return such articles free of charge, and, in the event of any unnecessary

damage, must compensate the owner.

Any person who shall cease to occupy any house or room in which any person has, within six weeks, been suffering from any infectious disease (a) must have such house or room (and all articles therein liable to retain infection) disinfected to the satisfaction of a registered medical practitioner, as testified by a certificate signed by him; (b) must give to the owner notice of the previous existence of such disease; and (c) must not knowingly make a false answer when questioned by the owner, or by any person negotiating for the hire of the house or room, as to there having, within six weeks previously, been therein any person suffering from any infectious disease. Penalties are provided in each case. The S.A. must give notice of the provisions of this section to the occupier of any house in which they are aware there is a person suffering from an infections disease.

Prompt interment.—The body of a person who has died of any infectious disease must not, without a certificate from the M.O.H. or a registered medical

where than in a mortuary, or in a room not used at the time as a dwelling-place, sleeping-place, or workroom. In such cases, and also where any corpse is retained in building so as to endanger the health of the inmates,

Justice may, upon the application of the M.O.H., order the body to be removed by the S.A. to a mortuary, and to be buried within a specified time. Unless the riends undertake to bury, and do bury within the specified time, the relieving officer must do so. The body of any person who has died from infectious disease an a hospital must not be removed except for immediate interment or to a mortuary, if the M.O.H. or registered practitioner certify that such restriction desirable for preventing infection. The body of any terson who has died from an infectious disease must not be conveyed in any public conveyance (other than hearse) without due warning to the owner or driver, who must forthwith provide for disinfection.

Detention in hospital.—Any person suffering from my infectious disease, and being an inmate of a cospital for infectious diseases, and who upon leaving would be without accommodation in which due presutions could be taken against the spread of infection, may, by order of a Justice, be detained in hospital (at me cost of the S.A.) for any specified period, and such

eriod may be extended as often as necessary.

The S.A. shall provide free temporary shelter, ith any necessary attendance, for the members of ny family in which infectious disease has appeared, the have to leave their dwellings to allow of dis-

infection by the S.A.

Infectious rubbish must not be thrown into any eceptacle for refuse without previous disinfection. In lotice of this prohibition must be served by the S.A. pon the master of any house in which there is any ase of dangerous infectious disease.

The Isolation Hospitals Act, 1893, gives to County Councils limited power to secure the provision of isolation hospitals in their various counties. It applies to England and Wales generally, but not to London or to any county borough; other boroughs also are exempt, except by Order of the L.G.B., if the population be less than 10,000 (according to the latest census), or by consent of the corporation if the population be 10,000 or more.

A Hospital District may consist of one or more local areas; a "local area" includes an urban or rural sanitary district, or any "contributory place," and is constituted by Order of the County Council.

Procedure.—The County Council may take the initiative by directing their M.O.H. to report as to the hospital requirement of any part of their county, and acting upon his report; but they may also be set in motion by a petition from any local authority, or from twenty-five ratepayers in any contributory place. The next step is for the County Council to hold a local inquiry, after which they make an Order constituting the hospital district and defining its extent. No local area can be included in a hospital district without the consent of its local authority, if it already has (in the judgment of the County Council) adequate accommodation; nor must a hospital district be formed for one local area only, or for one or more local areas within the same rural sanitary district, without the consent of the S.A., unless the County Council are satisfied that the S.A. are unable or unwilling to make suitable provision for the purpose. The Order constitutes a hospital committee, consisting of local representatives, but if a grant be made out of county funds the committee may consist wholly or in part of county councillors. The Order further gives the committee power to provide and maintain a hospital; and apart from this

they are authorised by the Act to make temporary arrangements for isolation, and to establish district hospitals in cottages or small buildings. They may also (subject to any regulations made by the County Council) undertake the training of nurses, and may charge for their attendance outside the hospital. Every hospital is to be provided with one or more ambulances, and must, if practicable, be "in connection with the system of telegraphs."

The County Council have the power of inspecting any such hospital, and of raising money by loan for

the purposes of the hospital.

Expenses.—"Structural" (e.g. site, building, furmishing, extensions, etc.) and "Establishment" (e.g. maintenance of hospital, salaries, etc.) expenses are borne by the several local rates of the constituent local areas, in proportions to be fixed by the Order. The costs of conveying, removing, feeding, medicines, disinfecting, and "all other things required for patients individually, exclusive of structural and establishment expenses," are termed patients' expenses. For ordinary mon-pauper patients they are to be paid by the local authority out of the rates of the local area from which the patient came, but the Guardians are responsible if Poor Law relief had been given at or within a fortnight of the time of admission. Patients lesiring exceptional accommodation are themselves responsible for the cost of maintenance, etc., on such terms ("special patients' expenses") as the committee may appoint.

The Isolation Hospital Act, 1901, provides that a S.A. having possession of a hospital for the reception of the sick may, with the sanction of the IL.G.B., transfer it to the County Council for appropriation to a district under the Isolation Hospitals Act, 1893, as an infectious diseases hospital, and the County Council may contribute to the expenses

of such a hospital. Minor amendments and definitions are included in the Act.

Midwives Act, 1902, is another Act directly concerned with disease-prevention, and may therefore be quoted here. The duties of the Local Supervising Authorities (the Councils of Counties or County Boroughs, or bodies to whom they may delegate their powers) are defined in Rules drawn up by the Central Midwives Board under s. 3, and approved by the Privy Council. In addition to the registration of midwives, the following duties arise under s. 3:—

(1) To exercise a general supervision over all midwives practising within their area, in accord-

ance with the above rules.

(2) To investigate charges of malpractices, negligence, or misconduct on the part of a midwife. If a primâ facie case be established, it is to be reported to the Central Board.

(3) To suspend a midwife from practice, when necessary, in accordance with the Midwives

Board rules.

The rules (1907) in question include the following:—

Whenever a midwife has been in attendance upon a patient suffering from puerperal fever, or from any other illness supposed to be infectious, she must disinfect herself, and all her instruments and other appliances, to the satisfaction of the S.A., and must have her clothing thoroughly disinfected before going to another labour. Unless otherwise directed by the Local Supervising Authority, all washable clothing should be boiled, and other clothing should be disinfected by the S.A.

Formal notification must be made by the midwife to the Local Supervising Authority when any of the following

occur :-

(a) When the mother or the child dies without having been seen by a registered medical practitioner.

(b) When the child is deemed to be stillborn.
(c) When, owing to the occurrence of various complications either of the labour or of its aftercourse, as

defined in the rules, it has been necessary for the midwife to advise that a registered medical practitioner be sent for.

The Local Supervising Authority must secure proper inpection of every midwife's case book, bag of instruments and ppliances, etc., and must, when necessary, make arrangements for inspection of her residence, and for investigation of

mer mode of practice.

The Local Supervising Authority are to suspend from praclice a midwife who contravenes the directions issued for the use If disinfectants and for the proper safeguards against the spread If infection laid down by the Central Board, or other rules of he Board. Any suspension (with the grounds thereof) must

e reported to the Central Board.

Local Government Act, 1888.—The sanitary provisions of this important Act are few. Besides the powers nn respect of pollution of streams conferred upon County Councils by s. 14, and the provisions as to the appointment and qualifications of county and other M.O.H. contained in ections 17 and 18 (page 473), only the following require mention here :-

A County Council may, subject to the approval of the L.G.B., make byelaws in relation to the whole or any specified part of their county (boroughs excepted*) for various purposes, including the prevention and suppression of nuisances not already punishable in a summary manner by virtue of any Act in force through-

out the county (s. 16).

Every M.O.H. for a district in any county shall send to the County Council a copy of every periodical report + of which a copylis, for the time being, required by the regulations of the L.G.B. to be sent to the Board; and if the M.O.H. fail to send such copy, the County Council may refuse to pay any contribution towards his salary. If it appear to the County Council from any such report that the Public Health Act, 1875, has not been properly put in force within the district to which the report relates, or that any other matter affecting the public health of the district requires to be remedied, the Council may cause a representation to be made to the L.G.B. on the matter (s. 19).

* Boroughs have the same power under s. 23 of the Municipal Corpora-

[†] The Regulations of the L.G.B. require the M.O.H. to furnish the County Council with copies of special as well as annual reports. (See page 477.)

County Councils have now certain powers under the Housing of the Working Classes Act, and they have, of course, power to make formal or informal representations to the L.G.B., or other controlling authority, upon any matter.

The Local Government Act, 1894, does not apply to county boroughs, or to London, Scotland, or Ireland.

Area and boundaries.—Boundaries are to be adjusted so as to prevent intersection of those which define counties, county districts, unions, and parishes. Small rural parishes are to be grouped into workable units for parish council purposes, but each is to have its parish meeting. In rural districts Parish Meetings and Parish Councils are constituted.

The Parish Meeting has the exclusive power of adopting certain optional Acts, including the Burials Acts and the Baths and Washhouses Act, and its consent is required before the Parish Council can incur any expense involving a loan or a rate exceeding

3d. in any year.

The Parish Council have power (s. 8) "to utilise any well, spring, or stream within their parish and provide facilities for obtaining water therefrom,* but so as not to interfere with the rights of any corporation or person; and to deal with any pond, pool, open ditch, drain, or place containing or used for the collection of any drainage, filth, stagnant water or matter likely to be prejudicial to health, by draining, cleansing, covering it, or otherwise preventing it from being prejudicial to health, but so as not to interfere with any private right or the sewage or drainage works of any Local Authority." They may execute works for these purposes, or contribute to the cost. They may complain (s. 16) to the County Council if the Rural District Council have failed to pro-

^{*} They cannot, except by agreement, acquire land for the purpose of any supply of water (s. 9).

vide or to maintain sufficient drainage or water supply ffor the parish, or to enforce any provision of the Public Health Act, and in that event the County (Council may take over to themselves the powers of the District Council for the purpose, or may make an Order for the necessary works to be carried out by the District Council or by some person appointed by the County Council. Notice must be given to the Parish Council concerned if the Rural District Council decide to undertake works of sewerage or water supply. The Parish Council may acquire or hire land for allotments, and may, if necessary, appeal to the County Council for compulsory powers of purchase or hire; the Order if granted requires confirmation by the **L.G.B.** (sections 9 and 10). They may also make official representations to the District Council under the Allotments Acts; or to the M.O.H. as to unhealthy dwellings or obstructive buildings under the Housing of the Working Classes Act (see p. 522 et seq.); or to the L.G.B. with a view to the grant of urban powers for their parish or any part of it. The District Council may delegate to them any powers which under the Public Health Acts could be delegated to a parochial committee (page 504). Apart from expenditure under adoptive Acts, the Parish Council must not incur expenses involving more than a 6d. rate in any year, nor more than a 3d. rate without the consent of the parish meeting. They may raise money on loan, but only with the approval of the parish meeting, the County Council, and the L.G.B.

Rural District Councils, notwithstanding the limited initiative possessed by the Parish Council, are the responsible executive sanitary authority, and must carry out all necessary works, giving notice to the Parish Council. The **L.G.B.** may by General Order confer upon Rural District Councils

such urban powers as they think fit; this is in addition to their right under the Public Health Act of 1875 (see page 505) to grant urban powers for particular areas, which right may under the 1894 Act be exercised by the Board on the application of the County or Parish Council concerned.

Urban District Councils are the S.A. for urban districts. Boroughs have their "corporation" or "town council," but are included in the term "county district," which comprises all urban and

rural districts.

All District Councils have the charge of highways. Among other powers given to them are the administration of the Acts relating to petroleum and infant life protection, and the licensing of knackers' yards. A District or Parish Council may delegate their powers to a committee of their number, or to a joint committee representing other councils also.

County Councils obtain under the 1894 Act certain powers of supervision and control, specified in

the foregoing summary.

Housing of the Working Classes Act, 1885.

—Every **S.A.** must exercise powers with which they are invested, to secure the proper sanitary condition of all premises within their area (s. 7).

A tent, van, shed, or similar structure, used for human habitation, which is in such a state as to be a nuisance or injurious to health, or which is so overcrowded as to be injurious to the health of the inmates, whether or not members of the same family, shall be deemed to be a nuisance within the meaning of s. 91 of the Public Health Act, 1875, and the provisions of that Act shall apply accordingly (s. 9). A S.A. may make byelaws in regard to such habitations. Power of entry between 6 a.m. and 9 p.m. is given to any person duly authorised by the S.A. or by a justice of the peace, if such person has reasonable cause to

suspect any contravention of this Act or of any byellaws made under it, or that there is in such habitation any person suffering from any dangerous infectious disease (s. 9).

Housing of the Working Classes Act, 11890.—Part I. deals with unhealthy areas, and is applicable in urban districts only. It is the duty of the M.O.H. to make an official repressentation to the S.A., whenever he sees cause to do so, that within a certain area either (a) any houses or courts are unfit for habitation, or (b) the bad arrangement or condition of the streets or houses, or the want of light, ventilation, or proper conveniences, or any other sanitary defects, are dangerous to the health of the inhabitants, and that the evils cannot be effectually remedied otherwise than by rearrangement and reconstruction of some or all of the streets or houses. The S.A. must consider this representation, and if satisfied of the truth thereof, and of the sufficiency of their resources, must declare the area to be an unhealthy area, and frame an improvement scheme. The scheme may exclude any part of the area, or include any neighbouring lands. It must provide for proper sanitary arrangements, and may provide for widening the approaches, or otherwise opening out the area. Due publicity having been given to the scheme in a specified manner, application is to be made to the "confirming authority" for a provisional Order. The Confirming Authority (the L.G.B.), if it approves the scheme, directs a local inquiry to be held respecting the correctness of the official representation and the sufficiency of the scheme. The provisional Order, if granted, is subject to confirmation by Parliament. It may modify the scheme, and must determine the extent of the provision to be made for the working classes displaced. In London, accommodation must be provided in or

near the area for the whole number displaced, unless the Order decrees otherwise; under certain conditions the Order may accept in substitution equally convenient accommodation not in or near the area, and may dispense with the obligation to any extent not exceeding one half. Outside the metropolis such provision is only compulsory if (and to the extent) prescribed by the Order. In assessing the value of property, no additional allowance for compulsory purchase is to be made in regard to any unhealthy portion of the area; and evidence may be given showing that any premises are (1) unfit for habitation, and cannot reasonably be made fit; or (2) in bad repair, or in an insanitary condition; or (3) that the rental is enhanced by reason of overcrowding or use for illegal purposes. In the first case the compensation is to be based upon the value of the land and building materials only; in the second, upon the value after allowing for the cost of necessary repairs; in the third, upon the value apart from such illegal use. After the provisional Order has been confirmed by Parliament, it becomes "the Confirming Act," and is executed by the local authority who take steps to receive and adjudicate upon claims, if necessary submitting them to arbitration.

Upon complaint by two Justices, or twelve rate-payers, the M.O.H. must inspect and report upon any area. If he fails to do so, or reports that it is not an unhealthy area, such ratepayers may appeal to the Confirming Authority, who may then appoint a medical practitioner to inspect the area. If the latter reports that it is an unhealthy area, the S.A. are required to prepare an improvement scheme. In any case, if the S.A. refuse or fail to prepare a scheme upon receipt of an official representation, they must report the facts to the Confirming Authority, who may then order a local inquiry to be held.

Part II. applies to rural as well as urban districts, and relates to unhealthy or obstructive dwellings. It is the duty of the M.O.H. to represent to the S.A. any dwelling which appears to him to be in a state dangerous to health so as to be unfit for human lhabitation; but the S.A. are also required to cause inspections of their district to be made from time to ttime for the same purpose. If satisfied that any dwelling is unfit for habitation, the S.A. must first take proceedings against the owner for the closure of the house, as directed by the Public Health Act, 1875. (See pp. 496-7.) If the closing order of the court is not terminated by a further order, the S.A. may proceed to make an order for the demolition of the premises; but before doing so must afford the owner an opportunity of stating his objections. There is an appeal to Quarter Sessions. Upon written complaint from four or more householders that any dwelling is unfit for habitation, the M.O.H. must inspect such house, and report to the S.A. If the S.A. (not being a rural S.A., or in the county of London) fail to take action within three months, the complainants may appeal to the L.G.B., who may, after local inquiry, order the S.A. to proceed.

Obstructive Buildings.—If the M.O.H. finds that any building—though not in itself unfit for habitation—by its proximity to other buildings, either (a) stops ventilation, or otherwise conduces to make them unfit for habitation, or (b) prevents the proper remedy of any nuisance or other evils in respect of such other buildings, he must make a representation to the S.A., stating that in his opinion the obstructive building should be pulled down. A similar representation may be made by any four or more inhabitant householders. In either case the S.A. must make inquiry as to the facts, and as to the cost of acquiring the land and pulling down the building. If they

decide to proceed, the S.A. can make an order for the demolition of the building, after giving the owner an opportunity of stating his objections, and subject to appeal to Quarter Sessions. The compensation to be paid by the S.A. to the owner, and also the amount recoverable by the S.A. from the owners of the houses benefited by the demolition, are settled by

arbitration in case of dispute. The S.A. may prepare a scheme for dedicating as a highway or open space, or appropriating or exchanging for the erection of working-class dwellings, the site of any building ordered to be demolished under Part II., if it appears to them that it would benefit the health of the inhabitants of the adjoining houses. They may also prepare an improvement scheme for any unhealthy area too small to be dealt with under Part I. Such schemes require confirmation by provisional Order of the L.G.B. after a local inquiry, and with or without modification. The Order may require provision to be made for the accommodation of persons of the working classes displaced. When an official representation has been made under Part II., or a closing order obtained, a rural S.A. must forward a copy to the County Council, and report from time to time all further proceedings. If it appears to the County Council that a closing order should be applied for, or an order for demolition made, and if the S.A., after due notice from the County Council, fails to adopt such measures, the Council may by resolution take over and exercise the powers of the S.A. in respect of such buildings, recovering the expenses from the S.A. Except in boroughs, a representation by the county M.O.H., if forwarded by the County Council to any S.A., has, for the purposes of Part II., the same effect as if made by the M.O.H. of the district.

In the county of London, Part I. is entrusted to the County Council, and Part II. primarily to the Borough Councils. In the event of doubt, the Home Secretary has to decide under which part of the Act a given area shall be dealt with. But the County Council may prepare schemes under Part II. If they think fit, and may apply to the Home Secretary to order a contribution from the District Authority, who, in like manner, if they proceed, may apply for a contribution from the County Council.

Part III. authorises the S.A. to provide lodginghouses for the working classes, subject to the sanction of the L.G.B. as regards urban authorities, and of the County Council in rural districts; and to make byelaws respecting such houses. For the administration of this part of the Act it is necessary for the S.A. tto adopt it.

The administration of Parts I., II., and III. in the district of the City of London is vested in the Commissioners, and the County Council have no juris-

diction in this area.

Part IV. contains supplemental provisions, of which the following is important:—"In any contract made after August 14th, 1885, for letting for habitation by persons of the working classes a house or part of a house, there shall be implied a condition that the house is at the commencement of the holding in all respects reasonably fit for human habitation."

Housing of the Working Classes Act, 1900.

This Act amends Part III. of the Act of 1890. It allows any Urban S.A. which has adopted Part III. to establish or acquire lodging-houses for the working classes outside their district; and any Rural S.A., with the consent of the County Council, to adopt Part III. either for the whole or part of their district. In giving or withholding consent the County Council shall have regard to the area proposed, to the necessity for accommodation for the housing of the working classes in that area; to the probability of such accommodation being provided without the adoption of the said Part; and to the liability which will be incurred by the rates, and to the question

whether it is, in all the circumstances, prudent to adopt the said Part.

An Urban S.A., with the consent of the L.G.B., and a Rural S.A. with the consent of the County Council, may lease any land acquired by them under and for the purposes of Part III. to any lessee for the purpose and under the condition that the lessee will carry the Act into execution by building and maintaining on the land lodging-houses within the meaning of the Act.

Section 6 defines the powers of the County Council to act in

default of a Rural Council.

Compensation questions which may arise shall in default of agreement be determined by a single arbitrator to be appointed

and removable by the L.G.B.

The Housing of the Working Classes Act, 1903, extends the maximum period for the repayment of loans raised for the purposes of these Acts to 80 years (from 60), leaving the actual period for repayment, subject to this limitation, to be determined with the sanction of the L.G.B.

The Board propose as a general rule to allow the full term of 80 years for the repayment of loans for the purchase of freehold land, and 60 years for the repayment of loans for the erection of buildings, where the circumstances are such that

this may properly be done.

The powers given to the Board to direct a local inquiry to be held, and a report made to them as to the correctness of the official representation (when the S.A. has taken no action) are supplemented by s. 4 (1) of the 1903 Act. If, on report made to them after inquiry, they are satisfied that a scheme ought to have been made for the improvement of the area to which the inquiry relates, or some part thereof, they can order the S.A. to make such a scheme (under either Part I., or Part II.)

The Order of the Board is enforceable by mandamus.

Section 4 (2) provides that any 12 or more ratepayers of the district shall have the same right of appeal, under s. 16 of the 1890 Act, as is given to the 12 or more ratepayers who have made complaint to the M.O.H. mentioned in that section.

The advertisements of an improvement scheme under Part I. may be published for three consecutive weeks at any period of the year, and the prescribed notices may be served during the 30 days next following the date of the last publication of the advertisement.

Section 8 concerns procedure for obtaining closing orders in regard to insanitary houses. - Where a S.A. desires to obtain a closing Order it was necessary under s. 32 of the 1890 Act, first give notice to the owner or occupier of the house to abate the uisance. S. 8, 1903, dispenses with this notice as regards any welling-house which, in the opinion of the S.A., is either on, or (b) is in such a state that its occupation should be inmediately discentinued.

A more speedy and efficacious way of obtaining possession if a house in respect of which a closing Order has been made than that provided by s. 32 (3) of the 1890 Act is afforded by

10, 1903. Whatever the value or rent of the house, the S.A. may proceed either under Sections 138 to 145 of the County courts Act, 1888, or under the Small Tenements Recovery act, 1838. The expenses of these proceedings can be recovered

com the owner under the Summary Jurisdiction Acts.

S. 11 (1) empowers the S.A., if they provide lodging couses under Part III., or if they supply accommodation under me Housing Acts or under any scheme made in pursuance of mese Acts, to provide and maintain, in connection therewith, my building adapted for use as a shop, any recreation grounds, or other buildings or land which, in the opinion of the Board, will serve a beneficial purpose in connection with the requirements of the persons for whom the lodging-houses or dwelling eccommodation are provided. The consent of the Board is equired to the exercise of these powers, and they may require actuatory provisions of control.

Service of notices. - The service of notices and other docu-

ents is facilitated by s. 13.

person to mix, colour, stain, or powder) any drug with any ingredient or material so as to affect injuriously the quality or potency of such drug, with intent that the same may be sold in that state; and no person shall sell any drug-so mixed, coloured, stained, or powdered . . ." (s. 4). The penalty for such injurious admixture is a fine not exceeding £50 for a first offence; subsequent offences are misdemeanours, punishable by imprisonment with hard labour for a period not exceeding six months. No liability is, however, incurred if the accused person can show that he was unaware of the admixture, and could not, with reasonable diligence, have ascertained it (s. 5).

"No person shall sell, to the prejudice of the purchaser any article of food or any drug which is not of the nature, substance, and quality of the article demanded by such purchaser, under a penalty not exceeding £20; but no offence shall be deemed to be committed under this section in the following cases:-(1) Where any matter or ingredient not injurious to health has been added to the food or drug because the same is required for the production or preparation thereof as an article of commerce in a state fit for carriage or consumption, and not fraudulently to increase the bulk, weight, or measure of the food or drug, or conceal the inferior quality thereof; (2) where the drug or food is a proprietary medicine, or is the subject of a patent in force, and is supplied in the state required by the specification of the patent; (3) where the food or drug is compounded . . . [and the provisions of the seventh and eighth sections are observed]; (4) where the food or drug is unavoidably mixed with some extraneous matter in the process of collection or preparation" (s. 6). As regards these exemptions, the onus probandi rests with the defendant (s. 24). No person shall sell any compound, drug, or article of food which is not comosed of ingredients in accordance with the demand of the purchaser, under a penalty not exceeding £20 ss. 7); but no offence under this section is committed in respect of the sale of a drug or article of food mixed with an ingredient not injurious to health, if it is labelled as "mixed" at the time of sale (s. 8).

"No person shall (with the intent that the same may be sold in its altered state without notice) betract from an article of food any part of it, so so to affect injuriously its quality, substance, or ature; and no person shall sell any article so altered without making disclosure of the alteration, under a cenalty not exceeding £20" (s. 9). In any prosecution under this Act, the defendant is to be discharged if he proves to the satisfaction of the Court without that he bought the article as being the same in ature, substance, and quality with that demanded by the purchaser, and with a written warranty to that affect; (b) that at the time of sale he had no reason to believe it to be otherwise; and (c) that he sold it the same state as when he purchased it (s. 25).

In every district a competent person may be (and, if required by the **L.G.B.**, must be) appointed as Public Analyst* (s. 10); and any person shall be intitled to have any drug or article of food purchased by him in such district analysed by the public analyst in payment of a sum not exceeding 10s. 6d., and to eceive from him a certificate of the result of the nalysis (s. 12). The **M.O.H.**, inspector, or constable, harged by the **S.A.** with the execution of the Act, may procure samples of food and drugs, and submit hem to the public analyst (s. 13). Any person purchasing an article for analysis shall, upon the

^{*} The appointment is made by the Town Council of any borough aving a separate Court of Quarter Sessions, or a separate police establishment. The County Council appoints the Public Analyst for all parts of he county not included in the above. All appointments and re-appointments are subject to the approval of the L.G.B.

completion of the purchase, forthwith notify to the seller his intention to have it analysed by the public analyst; and shall offer to divide it into three parts, to be then and there separated, and each part to be marked and sealed or fastened up, and shall, if required to do so, proceed accordingly, and shall deliver one of the parts to the seller. He shall retain one of the three parts for future comparison, and deliver the third to the public analyst (s. 14). If the seller do not accept the offer of division, the analyst must divide the article into two parts, and seal up and deliver one of them to the purchaser (s. 15). Samples may be sent to the analyst by registered parcel post, if his residence is two miles from that of the purchaser (s. 16). Any person refusing to sell to an officer of the S.A. any article of food or drug on sale by retail, the price being tendered, and the quantity demanded not being greater than is reasonably requisite, is liable to a penalty not exceeding £10 (s. 17). The certificate of the analyst must be in a prescribed form (s. 18), and is sufficient evidence of the facts therein stated, unless the defendant requires the analyst to be called as a witness (s. 21). The Justices before whom a case is heard may, at the request of either party, cause any article of food or drug to be sent to the Commissioners of Inland Revenue for analysis at the Government Laboratory (s. 22).

Sale of Food and Drugs Act Amendment Act, 1879.—This amending Act was passed in order to settle certain points upon which the 1875 Act was

found to be insufficiently explicit.

"It shall be no defence to allege that the purchaser is not prejudiced by the sale of adulterated articles, on the ground that he bought it for analysis only; or to allege that the article in question, though defective in nature, or substance, or quality, was not defective

in all three respects" (s. 2). "The M.O.H., inspector, or constable charged with the execution of the Act, may procure, at the place of delivery, a sample of milk in course of delivery to the purchaser or consignee, in pursuance of any contract; and may ssubmit the sample to the Public Analyst" (s. 3). The seller, or his representative, if he refuses to allow a sufficient sample to be taken, is liable to a penalty not exceeding £10" (s. 4). "As regards spirits not adulterated otherwise than by admixture of water, it is a good defence to prove that the admixture has not reduced the spirit more than 25 degrees under proof for brandy, whisky, or rum; or 35

dlegrees under proof for gin" (s. 6).

Sale of Food and Drugs Act, 1899, provides that articles imported into the United Kingdom, such as margarine, margarine cheese, impoverished or adulterated milk, condensed, separated, or skimmed milk, or any other article hereafter added by Order in Council, shall be conspicuously marked with name and description (s. 1); the L.G.B. and the Board of Agriculture may procure samples for analysis if it is deemed desirable in the general interest of the consumer, and the results of the analysis shall be communicated to the S.A., who shall pay the analyst's fiees (s. 2). Under s. 3 similar power is given to these Boards to act in default of the S.A. The Board of Agriculture may make regulations for determining what deficiency in any normal constituents, or adulterations, shall raise a presumption that milk, cream, butter and cheese is not genuine (s. 4). The Margarine Act, 1887, is extended to apply to "margarine cheese," which must be clearly marked on the package in black letters not less than half an inch long. Every maker of margarine and margarine cheese shall keep a register as to quantity and destination of his consignments, and he shall submit to

inspection (s. 7). The fat of margarine must not contain more than 10 per cent. of butter fat (s. 8), and condensed milk of all kinds may only be sold when clearly described on the label. A person selling milk on any highway must have conspicuously marked on the vehicle his name and address.

Penalties for wilful obstruction of officers in the course of their duties in food sampling and for offences under the Acts are, for the first offence, sums not exceeding £20, and for subsequent offences, £50 and £100; but proceedings may not be instituted after the expiration of 28 days from the time of purchase. A warranty is not available as a defence unless a copy has been sent to the purchaser within seven days of service of the summons, with a written notice stating that reliance will be placed on it, and specifying the name and address of the warrantor.

These Acts are directed not only against fraudulent dilution, abstraction and substitution, but also against injury to health, and the sampling and analysis should be guided by consideration of both points. In the case of beer the risk of lead contamination from pewter pipes has long been known, but that of arsenic was overlooked until 1900.

The Margarine Act, 1887, defines "butter" as made exclusively from milk, or cream, or both, with or without salt or other preservative, or added colouring matter. "Margarine" includes all substances, whether compounds or otherwise, prepared in imitation of butter, whether mixed with butter or not (s. 3). Every package of margarine must be so marked, in capital letters not less than $\frac{3}{4}$ inch square, and in shops, by labels with capital letters not less than $1\frac{1}{2}$ in. square). All margarine factories must be registered with the Authority appointing the Public Analyst of the district (s. 9). Officers authorised under the Sale of Food and Drugs Acts may take samples of butter (or substances purporting to be

nargarine) without going through the form of purchase required by that Act, but otherwise complying with its provisions as to dealing with the samples (s. 10). Any such substance not being marked as margarine is to be presumed to be exposed for sale as butter. The wendor is absolved if he proves that he bought the article with a written warranty, and sold it, in the same state us when bought, believing it to be butter. The penalty

for a first offence is a fine not exceeding £20.

The Butter and Margarine Act, 1907, substitutes for the definition of margarine in the Act above mentioned the following words: "Margarine shall mean any article of food, whether mixed with butter or not, which resembles butter and is not milk-blended butter." This Act also gives powers of inspection to the Board of Agriculture, the L.G.B. and the S.A. over all processes of manufacture, blending, re-working, or treatment of butter and similar substances, and a limit is laid down by which mot more than 16 per cent. of moisture is permitted in butter and margarine, and not more than 24 per cent. in milk-blended butter. Nor may such goods be imported. New powers are also given as to the proportion of milk solids in butter, as to the making of regulations by the L.G.B. for prohibiting the use of preservatives in butter, margarine, and milkblended butter, and as to labelling these several articles

Sale of Horseflesh, etc., Regulation Act, 1889.—The flesh of horses, asses, or mules must not be sold or kept for sale for human food, except in a shop or stall over upon which is placed conspicuously, in legible characters our inches long, an announcement that horseflesh is sold there. If otherwise, the M.O.H. or inspector may seize the meat and carry it before a magistrate, who may order as to its lisposal as he thinks fit; and the offender is liable to penalty. These conditions apply to such flesh cooked or uncooked, and

alone or mixed with other substances. It is illegal to supply horseflesh for human food to a purchaser asking for other meat, or for a compounded article not usually made of horseflesh.

The Rivers Pollution Prevention Act, 1876, deals separately with (a) solid matters, (b) sewage, (c) trade effluents, and (d) mining effluents.

(a) No solid refuse of any kind must be put into any stream, so as to interfere with its due flow or to

pollute its waters (s. 2).*

(b) No solid or liquid sewage matter must be passed into any stream; but as regards conditions in existence prior to 1876 no offence is committed under this Act if it is shown that the best practicable and available means are being used to render harmless the sewage entering the stream (s. 3).

(c) No poisonous, noxious, or polluting liquid from any factory or manufacturing process must be passed into any stream. The saving clause as to conditions

existent prior to 1876 is repeated here (s. 4).

(d) No solid matter from mines must be put into any stream so as prejudicially to affect its flow; nor must any poisonous, noxious, or polluting solid or liquid matter from mines be passed into any stream, other than water in the same condition in which it has been raised or drained from such mine (s. 5).

Proceedings may be instituted, in respect of pollution of streams by sewage or solid matters, by any private person or S.A. aggrieved (s. 8); but in respect of manufacturing or mining effluents S.A.'s only can take action, and subject to the approval of the L.G.B. The Board, in giving or withholding consent, shall have regard to the industrial interests involved, and the circumstances and requirements of the locality. They shall not give their consent to proceedings by the S.A. of a district which

^{*} See also Public Health Acts Amendment Act, 1890 (p. 508).

they are satisfied, after due inquiry, that means for endering harmless the effluents from such manufacturing processes are reasonably practical and available, and that no material injury to the interests of such industry will be caused by the proceedings (s.). Any person interested may call the attention of the S.A. to pollution by manufacturing or mining ffluents, and, if the S.A. refuse to move, may appeal to the L.G.B., and the Board may, after due inquiry, hirect the Authority to take proceedings (s. 6).

The S.A. must afford facilities for admitting trade iffluents into sewers, unless they would prejudicially iffect the sewers or interfere with the disposal of the ewage, or be injurious in a sanitary point of view, from their temperature or otherwise; or unless the ewers are only large enough for the ordinary require-

ments of the district (s. 7).

[S. 14 of the Local Government Act, 1888, confers pon County Councils the powers of S.A.'s under the Rivers Pollution Prevention Act; and enables the L.G.B. to form a joint committee representing all the dministrative counties through or by which a river, or any specified portion of a river, or any tributary thereof passes, and to confer upon the joint committee all or any of the powers of a S.A. under the Act.]

The Royal Commission on Sewage Disposal ecommended in 1903 the formation of River Boards broughout the country composed of joint committees of County Councils for the purpose of properly inforcing the R.P.P. Act. Such Boards have the advantage of jurisdiction over the whole of a drainage area and are sufficiently large to secure the appointment of skilled officers. Some have extended powers under special Acts. Many methods are now available for the treatment of sewage at noderate cost, in such manner as to deprive it of most,

if not all, of its impurities, leaving an effluent which may be passed into streams without material ill effect. As to trade pollutions, until recently comparatively little had been done. The difficulties, both legal and sanitary, are far greater than in respect of domestic or town sewage. Now, however, the treatment of sewage mixed with trade effluents is being practically dealt with and presents less difficulty where manufacturers adopt reasonable means for removing the solids, equalising the discharge, and, when necessary, neutralising the trade effluent.

A short explanatory Act (R.P.P. Act, 1893), removes a technical difficulty liable to be raised by **S.A.'s** seeking to evade their responsibilities under s. 3.

Canal Boats Act, 1877 .- The term "canal" includes all navigable waters, tidal or otherwise, within the body of a county, and the term "canal boat" includes every vessel, however propelled, used for conveyance of goods along such waters, but does not include a ship registered under the Merchant Shipping Act, unless the L.G.B. orders otherwise. No canal boat must be occupied as a dwelling unless it is registered under this Act, and then only by the number of persons of the age and sex for which it is registered (s. 1). The registration authority must be a S.A. whose district abuts on the canal on which the boat is intended to ply; and the boat must be registered as belonging to some place which is within the said district, and which is also within a school district (s. 7). Upon registration two certificates must be given to the owner, identifying the owner and the boat, and stating the place to which it belongs, and the number, age, and sex of persons allowed to dwell in the boat. The master of the boat must carry one of these certificates. Every registered boat must have conspicuously painted upon it the registered number, the place to which it belongs, and

he word "registered" (s. 3). If any person on canal boat is suffering from an infectious disease, the J.A. of the place where the boat is shall adopt such recautions as appear necessary, upon the certificate if the M.O.H. or other legally qualified practitioner; and may remove such sick person and exercise the ther powers conferred by the Public Health Act in his respect; and may detain the boat as long as necessary for cleansing and disinfecting (s. 4). if any person duly authorised by the S.A. has reason suspect any contravention of the Act, or that a erson on board is suffering from an infectious disease, e may enter the boat for the purpose of inspection eetween 6 a.m. and 9 p.m., and may require the master if the boat to afford him facilities for so doing, and to roduce the certificate of registry of the boat (s. 5).

The second section directs the L.G.B. to make egulations (i.) for the registration of canal boats; ii.) for lettering, marking, and numbering such coats; (iii.) for fixing the number, age, and sex of eersons who may be allowed to dwell in a canal boat, aving regard to the cubic space, ventilation, separaion of the sexes, general healthiness, and convenience ff accommodation; (iv.) for promoting cleanliness and abitable condition of such boats; and (v.) for preenting the spread of infectious disease by canal boats.

Such regulations were accordingly issued by the L.G.B. in 1878, and the following is a summary of their principal

anitary provisions :-

There must be at least one dry, clean, weatherproof cabin, in good repair. An after-cabin intended to be used as a dwelling must contain not less than 180 cubic feet of free air space, and a fore-cabin 80 cubic feet; every such cabin must have means of ventilation besides the door, and must be so constructed as to provide adequate sleeping accommodation. One cabin must contain a stove and chimney. The boat must be furnished with suitable storage for three gallons of water. If intended to be

ordinarily used for foul cargoes, the hold must be separated from any inhabited cabin by a double bulkhead with an interspace of four inches, and the bulkhead next the cargo must be water-tight. Not less than 60 cubic feet of air space must be allowed for each person over twelve years of age, and not less than 40 cubic feet for each person under that age. In "fly-boats" worked by shifts, a cabin occupied at the same time by two persons must have a capacity of 180 cubic feet. A cabin in which a married couple sleep must not be occupied at the same time by any other male above 14, or female above 12 years of age. Males above 14 and females above 12 years of age must not occupy the same sleeping cabin at the same time, but reservation is made for married couples, and also (under certain conditions) in respect of boats constructed prior to 1878. The interior of the cabin must be repainted every three years, and must be kept clean. Bilge water must be pumped out daily. The master of the boat must at once notify the occurrence of any case of infectious disease on the boat to the S.A. of the district through which the boat may be passing, and also to the S.A. of the place of destination; he must also inform the owner, who is required to notify to the S.A. of the place to which the boat belongs. If the boat is detained by the S.A. for purposes of disinfection, the S.A. must obtain a medical certificate that the boat has been cleansed and disinfected, and must give it to the master.

Act in certain details, and makes it incumbent upon every S.A. through whose districts any canal passes to enforce the provisions of the Act and Regulations, and to make an annual report upon the subject to the L.G.B.* The Board is required to make inquiries from time to time (by an inspector or inspectors specially appointed for the purpose) as to the working of the Acts and Regulations, and to report annually thereon to Parliament.

^{*} This report has to be drawn up in prescribed form and must include particulars as to registration, notification of change of master, certificate, marking of boat, overcrowding, separation of sexes, cleanliness, ventilation, painting, provision of water cask, removal of bilge water, notification of infectious disease, and admittance of inspector.

The Dairies, Cowsheds, and Milkshops Order of 1885 was made by the Privy Council under s. 34 of the Contagious Diseases (Animals) Act of 1878. An amending Act (C.D.A. Act, 1886) ransferred to the L.G.B. the powers of the Privy Council under this section, and the D.C. and M. Amending Order, 1886, substituted the Board for the Privy Council and remedied certain omissions in regard to penalties, giving the Order the force of law. The D.C. and M. Order of 1899 amended the previous Orders by including tuberculosis among the infective diseases on the occurrence of which milk should not be sold under s. 15. The effect of the three Orders is to throw upon every S.A. the duty of supervising the milk trade in their district, and of carrying out certain general regulations prescribed by the Orders. These duties are common to all districts alike, but any SS.A. may arm itself with further powers by making Regulations under s. 13, having the force of byelaws. The chief provisions of the Orders are summarised below :-

Section 6. (1) It shall not be lawful for any person to carry on . . . the trade of cowkeeper, dairyman, or purveyor of milk unless he is registered as below. ((2) Every S.A. shall keep a register of such persons, and shall from time to time revise and correct the register. (3) The S.A. shall register every such person, but registration shall not be deemed to authorise such person to occupy as a dairy or cowshed any particular building, or preclude proceedings. (4) The S.A. shall from time to time give public notice of registration being required, and of the mode of registration. (5) A person who carries on the trade of cowkeeper or dairyman for the purpose only of making and selling butter or cheese, or both, and who is not also a purveyor of milk, need not be registered. (6) A person who sells milk

of his own cows in small quantities to his workmen or neighbours for their accommodation need not, by

reason thereof, be registered.

Section 7. (1) It shall not be lawful to begin to occupy as a dairy or cowshed any building not so occupied at the commencement of this Order until provision is made, to the reasonable satisfaction of the S.A., for the lighting and ventilation, including air-space, and the cleansing, drainage, and water-supply; or (2) without first giving one month's notice in writing to the S.A.

Section 8. It shall not be lawful to occupy as a dairy or cowshed any building-whether so occupied at the commencement of this Order or not-if the lighting and ventilation, including air-space, and the cleansing, drainage, and water-supply thereof, are not such as are necessary or proper (a) for the health and good condition of the cattle therein; (b) for the cleanliness of milk-vessels used therein for containing milk for sale; and (c) for the protection of the milk

therein against infection or contamination.

Section 9. It shall not be lawful for any cowkeeper, or dairyman, or purveyor of milk, or occupier of a milkshop (a) to allow any person suffering from a dangerous infectious disorder, or having recently been in contact with a person so suffering, to milk cows or to handle vessels used for containing milk for sale, or in any way to take part or assist in the conduct of the trade so far as regards the production, distribution, or storage of milk; or (b) if himself so suffering, or having recently been in contact as aforesaid, to milk cows or handle vessels containing milk for sale, or in any way to take part in the conduct of his trade as far as regards the production, distribution, or storage of milk; until, in each case, all danger therefrom of the communication of infection to the milk or of its contamination has ceased.

Section 10. It shall not be lawful for any wkeeper, dairyman, or purveyor of milk, or cupier of a milk store or milkshop, after the receipt notice of not less than one month from the Local attention to the provisions of this rticle, to permit any water-closet, earth-closet, privy, sspool, or urinal to be within, communicate directly th, or ventilate into, any dairy or any room used as

milk store or milkshop.

Section 11. It shall not be lawful for any wkeeper, or dairyman, or purveyor of milk, or cupier of a milk store or milkshop, to use a milk ore or milkshop in his occupation, or permit the me to be used, as a sleeping apartment, or for any rpose incompatible with the proper preservation of es cleanliness of the milk store or milkshop, and the milk-vessels and milk therein, or in any unner likely to cause contamination of the milk erein.

Section 12. It shall not be lawful for any wkeeper, or dairyman, or purveyor of milk to keep y swine in any building used by him for eping cows, or in any milk store or other place used

him for keeping milk for sale.

Section 13. Any S.A. may make regulations. for the inspection of cattle in dairies; (b) for escribing and regulating the lighting, ventilan,* cleansing, drainage, and water supply, of dairies d cowsheds; (c) for securing the cleanliness milk stores, milkshops, and milk-vessels used for ntaining milk for sale; and (d) for prescribing precauons to be taken by purveyors of milk, and persons ling milk by retail, against infection or contaminon.

^{*} Where a minimum cubic space is specified for cowsheds in such sulations, the L.G.B. have suggested 800 cubic feet per head, but in ny districts a lower standard has been adopted.

Section 14. The following provisions shall apply to regulations made by any S.A. under this Order:—
(1) Every regulation shall be published by advertisement in a newspaper circulating in the district.
(2) The S.A. shall send to the L.G.B. a copy of every regulation at least a month before the date of operation. (3) The L.G.B. may at any time revoke any regulation if satisfied on inquiry that it is too restrictive or otherwise objectionable.

Section 15. The milk of a cow suffering from cattle-plague, pleuro-pneumonia, or foot-and-mouth disease (a) shall not be mixed with other milk; (b) shall not be sold or used for human food; and (c) shall not be sold or used for food of animals, unless it has been boiled. Tuberculosis is now included in

this section under the D.C. and M.O. of 1899.

The following Model Regulations under section 13 of the Order have been drawn up by the L.G.B. for Dairies, Cowsheds and Milkshops.

Interpretation.

1. The expression "cowshed" includes any dairy in which milking cows may be kept, and the expression "cowkeeper" means any person following the trade of a cowkeeper or dairyman required to be registered under the Order of 1885.

For the Inspection of Cattle in Dairies.

2. Every occupier of a dairy wherein any cattle may be kept, and which the M.O.H., or the Inspector of Nuisances, or any other officer of the Council specially authorised by them in that behalf, may visit for the purpose of inspecting a cattle, and every person for the time being having the care or control of any such dairy, or of any cattle therein, shall afford such officer all reasonable assistance that may, for the purpose of the inspection, be required by him.

FOR PRESCRIBING AND REGULATING THE LIGHTING, VENTILA-TION, CLEANSING, DRAINAGE, AND WATER SUPPLY OF COWSHEDS AND DAIRIES.

PART I.

The regulations in this Part shall apply to cowsheds the cows from which are habitually grazed on grass land during the greater

part of the year, and when not so grazed are habitually turned out during a portion of each day.

3. Lighting.—Every cowkeeper shall provide that every wshed in his occupation shall be sufficiently lighted with

indows, whether in the sides or roof thereof.

4. Ventilation.—Every cowkeeper shall cause every cowed in his occupation to be sufficiently ventilated, and for this rpose to be provided with a sufficient number of openings to the external air to keep the air in the cowshed in a wholene condition.

5. Cleansing.—(1) Every cowkeeper shall cause every part the interior of every cowshed in his occupation to be broughly cleansed from time to time as often as may be so cessary to secure that such cowshed shall be at all times

sonably clean and sweet.

(2) Such person shall cause the ceiling or interior of the of and walls of every cowshed in his occupation to be operly limewashed twice at least in every year, that is to say, e during the month of May and once during the month of tober, and at such other times as may be necessary.

Provided that this requirement shall not apply to any part such ceiling, roof, or walls that may be properly painted, or mished, or constructed of or covered with any material such to render the limewashing unsuitable or inexpedient, and

t may be otherwise properly cleansed.

(3) He shall cause the floor of every such cowshed to be roughly swept, and all dung and other offensive matter to removed from such cowshed as often as may be necessary,

I not less than once in every day.

6. Drainage.—(1) Every cowkeeper shall cause the drainof every cowshed in his occupation to be so arranged t all liquid matter which may fall or be cast upon the floor y be conveyed by a suitable open channel to a drain inlet nate in the open air at proper distance from any door or ndow of such cowshed, or to some other suitable place of posal which is so situate.

(2) He shall not cause or suffer any inlet to any drain of

h cowshed to be within such cowshed.

7. Water Supply.—(1) Every cowkeeper shall keep in, or connection with, every cowshed in his occupation a supply water suitable and sufficient for all such purposes as may m time to time be reasonably necessary.

(2) He shall cause any receptacle which may be provided such water to be emptied and thoroughly cleansed from e to time as often as may be necessary to prevent the pollution of any water that may be stored therein, and where such receptacle is used for the storage only of water he shall cause it to be properly covered and ventilated, and so placed as to be at all times readily accessible.

PART II.

- The regulations in Part I., and also the following regulation. shall apply to all cowsheds other than those the cows from which are habitually grazed on grass land during the greater part of the year, and when not so grazed are habitually turned out during a portion of each day.
- 8. A cowkeeper shall not cause or allow any cowshed in his occupation to be occupied by a larger number of cows than will leave not less than eight hundred feet of air space for each cow.

Provided as follows:

(a) In calculating the air space for the purposes of this regulation, no space shall be reckoned which is more than sixteen feet above the floor; but if the roof or ceiling is inclined, then the mean height of the same above the floor may be taken as the height thereof for the purposes of this regulation.

(b) This regulation shall not apply to any cowshed constructed and used before the date of these regulations

coming into effect, until two years after that date.

PART III.

9. In this Part the expression "dairy" means a dairy in

which cattle are not kept.

10. Lighting.—Every cowkeeper shall provide that every dairy in his occupation shall be sufficiently lighted with

windows, whether in the sides or roof thereof.

- 11. Ventilation.—Every cowkeeper shall cause every dairy in his occupation to be sufficiently ventilated, and for this purpose to be provided with a sufficient number of openings into the external air to keep the air in the dairy in a wholesome condition.
- 12. Cleansing.—(1) Every cowkeeper shall cause every part of the interior of every dairy in his occupation to be thoroughly cleansed from time to time as often as may be necessary to secure that such dairy shall be at all times reasonably clean and sweet.

(2) He shall cause the floor of every such dairy to be

thoroughly cleansed with water at least once in every day.

13. Drainage.—(1) Every cowkeeper shall cause the drainge of every dairy in his occupation to be so arranged that all quid matter which may fall or be cast upon the floor may be onveyed by a suitable opening channel to the outside of such airy, and may there be received in a suitable gully communiating with a proper and sufficient drain.

(2) He shall not cause or suffer any inlet to any drain of

such dairy to be within such dairy.

14. Water Supply.—(1) Every cowkeeper shall cause every airy in his occupation to be provided with an adequate supply If good and wholesome water for the cleansing of such dairy and of any vessels that may be used therein for containing milk, and for all other reasonable and necessary purposes in onnection with the use thereof.

(2) He shall cause every cistern or other receptacle in which any such water may be stored to be properly covered and entilated, and so placed as to be at all times readily accessible.

(3) He shall cause every such cistern or receptacle to be implied and thoroughly cleansed from time to time as often as may be necessary to prevent the pollution of any water that may be stored therein.

FOR SECURING THE CLEANLINESS OF MILK-STORES, MILK-SHOPS, AND OF MILK-VESSELS.

- 15. Cleanliness of Milk-stores and Milk-shops. Every eccupier of a milk-store or milk-shop shall cause every part of he interior of such milk-store or milk-shop to be thoroughly ceansed from time to time as often as may be necessary to maintain such milk-store or milk-shop in a thorough state of eeanliness.
- 16. Cleanliness of Milk-vessels. (1) Every cowkeeper wall from time to time, as often as may be necessary, cause very milk-vessel that may be used by him for containing alk for sale to be thoroughly cleansed with steam or clean biling water, and shall otherwise take all proper precautions or the maintenance of such milk-vessel in a constant state of eanliness.
- (2) He shall, on every occasion when any such vessel nall have been used to contain milk, or shall have been bturned to him after having been out of his possession, cause ach vessel to be forthwith so cleansed.

FOR PRESCRIBING PRECAUTIONS AGAINST INFECTION OR CONTAMINATION OF MILK.

17. (1) Every purveyor of milk or person selling milk by stail shall take all reasonable and proper precautions, in and

in connection with the storage and distribution of the milk, and otherwise, to prevent the exposure of the milk to any infection or contamination.

(2) He shall not deposit or keep any milk intended for

sale-

(a) in any room or place where it would be liable to become infected or contaminated by impure air, or by any offensive, noxious, or deleterious gas or substance, or by any noxious or injurious emanation, exhalation, or effluvium; or

(b) in any room used as a kitchen or as a living

room; or

(e) in any room or building, or part of a building, communicating directly by door, window, or otherwise with any room used as a sleeping room, or in which there may be any person suffering from any infectious or contagious disease, or which may have been used by any person suffering from any such disease and may not have been properly disinfected; or

(d) in any room or building or part of a building

in which there may be any direct inlet to any drain.

(3) He shall not keep milk for sale, or cause or suffer any such milk to be placed, in any vessel, receptacle, or utensil

which is not thoroughly clean.

(4) He shall cause every vessel, receptacle, or utensil used by him for containing milk for sale to be thoroughly cleansed with steam or clean boiling water after it shall have been used, and to be maintained in a constant state of cleanliness.

(5) He shall not cause or suffer any cow belonging to him or under his care or control to be milked for the purpose of

obtaining milk for sale-

(a) unless, at the time of milking, the udder and teats

of such cow are thoroughly clean; and

(b) unless the hands of the person milking such cow, also, are thoroughly clean and free from all infection and contamination.

PENALTIES.

18. Every person who shall offend against any of the foregoing regulations shall be liable for every such offence to a penalty not exceeding five pounds, and in the case of a continuing offence to a further penalty not exceeding forty shillings for each day after written notice of the offence from the Council.

Provided, nevertheless, that the Justices or Court before whom any complaint may be made or any proceedings may be

ken in respect of any such offence may, if they think fit, ljudge the payment as a penalty of any sum less than the all amount of the penalty imposed by this regulation.

The Public Health (London) Act, 1891, orms a sanitary code for London, and includes (with many important additions and amendments) the main rrovisions of the Public Health Acts and the Infectious Diseases Notification and Prevention Acts. A brief mmary of the principal points of divergence from

ne Acts named above is appended.

Water supply.—Absence of proper water supply, re of proper fittings, is held to render a house unfit for abitation. A new house must not be occupied until me S.A. grant a certificate that it has a proper water apply. A water company cutting off the supply of my house must give immediate notice to the S.A. for closure of polluted wells, etc., the S.A. have only satisfy the Court that the water is "so polluted, or kely to be so polluted, as to be injurious or dangerous b) health."

Closets.—A new house must have "one or more vater-closets, as circumstances may require," with roper water supply and trapped soil-pan, and other ccessories. The same applies to all houses, irrespecwe of date, under notice from the S.A. Substitution Privies or earth-closets is only permissible if the vailable sewerage or water supply is insufficient for water-closet. Penalties are prescribed for (a) wililly injuring any drain or closet or water supply in onnection therewith; (b) discontinuing any such vater supply without lawful authority; and (c) so contructing or repairing a water-closet or drain that it is nuisance or injurious to health.

Scavenging .- The S.A. must cleanse streets, footaths, cesspits, earth-closets, privies, and cesspools. hey must remove house-refuse at proper intervals; and trade-refuse also, if required to do so, on payment.

They may undertake the collection of manure and other refuse, on request; or may, by order, require periodical removal by the owner.

Cellar Dwellings.—No underground room not let or occupied separately as a dwelling before January 1st,

1892, can be so let or occupied, unless

(a) Every part is seven feet high, and at least three feet above the level of the adjoining ground; but if the outside area is six feet wide, or if its width is not less than the depth of the floor below ground-level, then only one foot of the height need be above the ground-level.

(b) Every wall has a damp-course, and if in contact with the soil, is effectually secured from damp

from the soil.

(c) There is an open area outside along the whole frontage, four feet wide in every part, and six inches below the floor-level. It may be crossed by steps, but not opposite a window.

(d) The area and the soil immediately below the

room are effectually drained.

(e) The hollow space (if any) below the floor is ventilated to the outer air.

(f) Any drain under the room is made of gas-tight pipe.

(g) The room is effectually secured against the

rising of any effluvia or exhalation.

(h) There is proper water-closet and ashpit accommodation.

(i) There is effectual ventilation.

(j) There is a fire-place, with chimney.

(k) There are one or more windows opening directly into the open air; the window-area being at least one-tenth of the floor area, at least half made to open, and in each case opening to the top.

The same conditions apply to underground rooms occupied separately as dwellings before January 1st

892; but the S.A., either by general regulations upon special application by the owner, may modify my conditions newly imposed by this Act which volve structural alteration of the building.

Keeping of animals.—Swine must not be kept ithin forty yards of a street or public place, nor be llowed to stray in any public place. A Court may cohibit the keeping of any animal in any specified

ace shown to be unfit for the purpose.

Offensive trades.—None of the following must be tablished anew in London:-Blood-boiling, bonebiling, manure manufacture, soap-boiling, tallowelting, or knacker's yard. Nor any of the following, ithout the sanction of the County Council:-Fellonger, tripe-boiling, slaughtering of cattle or horses, other business which the County Council may, ith the approval of the L.G.B., add to the list. aughter-houses and knackers' yards must obtain cences, issued by the County Council, lasting or one year only, but renewable at the discreon of the County Council. The same applies to wsheds.

Notification is compulsory. The medical cerficate must state the patient's full name, postal ldress, age, sex, whether attended in public or rivate practice; if in hospital, the date of admission, and the place from which brought. The M.O.H. oust, within twelve hours of receipt, send a copy f the certificate to the Metropolitan Asylums coard, and another to the head-teacher of the schools ttended by the patient or by other inmates of the ouse. The Asylums Board must furnish to the ounty Council, and to every M.O.H., such weekly eturns of certificates so received by them as the ounty Council may prescribe. Any S.A. may, with ne consent of the L.G.B., add other dangerous inectious diseases to the list, temporarily or permanently, and the County Council have like power for

the county as a whole.

Isolation.—A person who knows himself to be suffering from a dangerous infectious disease must not milk any animal, or pick fruit, or engage in any business in such manner as to be likely to spread the disease. He must not enter any public conveyance, nor must any other person place him, or convey him, in such.

Disinfection.—Every S.A. must provide disinfect-

ing apparatus.

Mortuaries.—Every S.A. must provide a mortuary; the addition of a room for post-mortems is optional, unless ordered by the County Council. The County Council may provide other mortuaries wherein bodies may be kept for identification, and must provide proper accommodation for inquests.

Appointment of M.O.H. must not be for a limited period only. It is not terminable by the **S.A.** without the consent of the **L.G.B.** The **M.O.H.** must live in his district, or within a mile of the boundary. He

must be qualified as stated on page 473.

Sanitary Inspectors appointed after January 1st, 1895 (if they have not held office as sanitary inspector during three consecutive years prior to that date in a London district or in an urban district outside London with a population of not less than 20,000) must hold a certificate of competence granted after examination by such body as the **L.G.B.** may from time to time approve.*

Default of S.A.—In the event of default of the S.A. with respect to the removal of any nuisance, the institution of any proceedings, or the enforcement of any byelaw, the County Council may act in their stead, and recover the expenses from the S.A. If the L.G.B. are satisfied, upon complaint from the County Council

^{*} Such examinations are now conducted and certificates issued by the Sanitary Inspectors' Examination Board.

d after due inquiry, that the S.A. have made default carrying out any part of this Act, they may order S.A. to perform the duty within a specified time; if this is not done, the order may be enforced by andamus, or the L.G.B. may appoint the County uncil to perform such duty, at the cost of the S.A. Byelaws.—The London County Council must make elaws—

(1) As to the time and manner of conveying offen-

e matters by road or water.

(2) As to the closure and filling up of cesspools d privies, removal and disposal of refuse, and duties householders in facilitating removal of house-refuse.

(3) As to water-closets, earth-closets, ashpits, cessols, and receptacles for dung, and the proper accesries thereof in connection with new or old buildings.

They may also make byelaws as to the conduct of train offensive trades, the buildings in which they carried on, and the mode of application for sanction. eese trades are fellmonger, tripe-boiler, slaughterer cattle or horses; but other businesses may be afterords added to the list.

Every London S.A. must make and enforce byews relating to nuisances and keeping of animals, uses let in lodgings, and also the following:—

(1) For the prevention of nuisances arising from ensive matter running out of any manufactory, ewery, slaughter-house, knacker's yard, butcher's or hmonger's shop, or dunghill, into any uncovered uce, whether enclosed or not.

(2) For the paving of yards and open spaces in

nnection with dwellings.

(3) For securing cleanliness, and freedom from llution, of tanks, cisterns, and other receptacles drinking-water.

(4) For maintaining the efficiency of flush for

iter-closets.

The S.A. may also make byelaws relating to shipborne infectious cases, public conveniences, mortuaries,

and inhabited tents and vans (p. 578 et seq.).

The London Building Act, 1894, is a consolidation Act amending various enactments relating to streets and buildings in the metropolis. The Act is divided into an introduction and fifteen parts, dealing with the formation and widening of streets, lines of building frontage, naming and numbering of streets, heights and construction of buildings, temporary, dangerous and neglected structures, sky-signs, and a number of similar miscellaneous matters. Some duties under the Act, such, for instance, as respect wooden structures, and in relation to the removal of obstructions in streets, are to be exercised by the Borough Councils; otherwise the Act is administered by the London County Council.

An Amending Act was passed in 1898 dealing with building extensions, fore-courts, the height of

working-class dwellings, etc.

The Infant Life Protection Act, 1897, requires that persons receiving for hire infants for the purpose of maintenance must give notice to the local authority (in London the County Council, elsewhere the Board of Guardians, except in Scotland, where it is the Parish Council), furnish particulars, and submit to inspection and supervision. The local authority may, by order, require the removal of such infant to a workhouse or "place of safety," if it be kept in premises which are so unfit or so overcrowded as to endanger its health, or if it be kept by persons unfit to care for it, or who, by negligence, ignorance or other cause, endanger its health.

The Cleansing of Persons Act, 1897, gives power to S.A. to provide cleansing facilities free of charge (not to be considered as parochial relief or charity) to persons infested with vermin.

n practice such cleansing includes a bath and disinection of clothes.

The Public Health (Regulations as to rood Act), 1907, amends the Public Health Act of 896 by giving power to the L.G.B. to make regulations authorising measures to be taken for the prevention of danger arising to public health from the importation, preparation, storage, and distribution of food r drink (other than drugs or water) intended for sale or human consumption. Such regulations have to be published and may provide for the examination

and taking of samples of any articles of food.

The Notification of Births Act, 1907, is an doptive measure passed with the view of assisting adirectly in the reduction of infant mortality. rovides, where adopted, for notification in writing to me M.O.H. within thirty-six hours of the occurrence f a birth. The duty falls either upon the father of ne child or any person attending the mother at the me of or within six hours of the birth. The S.A. is supply without fee stamped forms or notification. enalty for non-compliance is 20s. This notification in addition to and not in substitution of ordinary arth registration. A certifiable birth is defined as any child which has issued forth from its mother ter the expiration of the twenty-eighth week of regnancy whether alive or dead." The Act may be lopted by a resolution of the S.A. which must eceive the sanction of the L.G.B. and be advertised the district. Any S.A., including County Councils ccept the County of London, may adopt the Act, or ie L.G.B. may, by order, declare the Act in force in ny area they think expedient.

The London County Council (General owers Act), 1907, gives power to the L.C.C. to ke samples of milk for examination for the presence tubercle bacillus. If it be found, provision is

made for inspection of cows, and cause must be shown by the dairyman why the sale of the milk should not be prohibited. If such a prohibition order be made without due cause, the dairyman is entitled to compensation. Every dairyman selling milk within the county must notify the presence of tuberculosis in his cows. The same Act provides for the examination of school children for uncleanliness and vermin, and, if necessary, compulsory cleansing of the body and clothing. The same powers are given for cleansing inmates of common lodging-houses.

will designed beyond not since I wanted the

CHAPTER XIX

FACTORIES AND WORKSHOPS

INCIDENTAL reference has been made in Chapters XVII. and XVIII. to the powers and duties of the S.A. with regard to industrial premises. Large additions are made by the Factory Acts, and in stating these it will be convenient to refer briefly to some of the leading points in industrial hygiene, and to indicate generally the division of responsibility between the local and central authorities.

The Factory and Workshop Acts, 1901 and 1907, which apply to some 100,000 factories and 150,000 workshops, giving employment to about 5,000,000 workpeople, are administered chiefly by H.M. Inspectors of Factories, but also in important particulars by the S.A. The latter are charged with duties as to (1) sanitation of workshops (and, in certain particulars, factories also), (2) bakehouses, (3) home-work, and (4) means of escape from fire. Many of these fall to the M.O.H., upon whom are placed further duties of reporting.

Factories include (1) works in which mechanical power is used in aid of a manufacturing process, or in quarries, pit banks of metalliferous mines, dry cleaning, carpet-beating or laundries; and (2) the following, whether power be used or not:—print works, bleaching and dyeing works, earthenware works, lucifer match works, percussion-cap works, cartridge works, paper-staining works, fustian-cutting works, blast furnaces, copper mills, iron mills, foundries, metal and india-rubber works, paper mills, glass works, tobacco factories, letterpress printing works, book-binding works, flax scutch mills, electrical stations. Those in italies are technical expressions, defined in the Act.

A tenement factory is one in which mechanical power is supplied to different parts of the same building occupied by different persons for the purpose of any manufacturing process or handicraft in such manner that these parts con-

stitute in law separate factories.

Workshops include (1) the following, unless they are factories by reason of use of mechanical power:—hat works, rope works, bakehouses ("any places in which are baked bread, biscuits, or confectionery from the baking or selling of which a profit is derived"), lace-warehouses, shipbuilding yards, quarries, pit-banks of metalliferous mines, dry-cleaning works, carpetbeating works, bottle-washing works, and laundries.

(2) Premises (not being factories) in which manual labour is exercised by way of trade or for purposes of gain, in or incidental to the making, altering, repairing, ornamenting, finishing or adapting for sale of any article, if the employer has

the right of access or control.

(3) Tenement workshops: that is, workplaces in which, "with the permission of or under agreement with the owner or occupier, two or more persons carry on any work which would constitute the workplace's workshop if the persons working therein were in the employment of the owner or occupier."

There may be several distinct factories or workshops, each comprising one or more rooms in different occupation, in the same building, which may also be used for dwelling or other

purposes.

Domestic factories and domestic workshops are those carried on in private houses without use of mechanical power, where the only persons employed are members of the same family dwelling there. If any work certified by the Home Secretary to be dangerous is carried on they are treated as ordinary factories and workshops: otherwise they are, so far as sanitation is concerned, treated as ordinary workshops, except that the requirements as to general ventilation and drainage of floors do not apply.

The following, if carried on under conditions which would otherwise make the premises a domestic workshop, are

exempted.

(1) Straw-plaiting, pillow-lace making, glove-making.

(2) Work done at irregular intervals and not furnishing the

principal means of living to the family.

Men's workshops (those in which only males over 18 are employed) are exempted from several of the provisions of the Act, including those relating to general ventilation, temperature, drainage of floors, sanitary accommodation, lavatories, fans for removal of dust or fumes, and means of escape from fire.

Crown factories and workshops are excluded from the juris-

diction of the S.A.

Workplace is not defined, but has a wider meaning than workshop, and has been held to include any place where people assemble together to do work permanently: e.g. a stable and

tableyard where men were employed as cab-cleaners and norse-keepers.

In workshops and workplaces the S.A. are primarily responsible for enforcing sanitary requirements generally, and means of escape from fire; and their officers have the same powers of entry, inspection, and egal proceedings as Inspectors of Factories. Certain anitary requirements, however, more particularly those relating to temperature, fans for removal of dust and fumes, and those arising under the Factory Act as distinguished from the powers of the S.A. under the P.H. Acts) with regard to sanitary accommodation, are enforced by the Factory Inspectors. The S.A. are required to keep a register of all workshops in their district. If the M.O.H. finds any woman, young person, or child employed in a workshop in which no abstract of the Factory Act is affixed, he is required o inform the Factory Inspector. The occupier of a new factory or workshop must notify the Factory Inspector, who, in the case of a workshop, forwards he notice to the S.A.

In factories, sanitation generally is enforced by the Factory Inspector, not by the S.A., but the atter deal with means of escape from fire (in London this rests with the County Council), and saniary accommodation (under the P.H. Acts only), and have special duties in connection with bakehouses and lomestic factories.

If any act or default, punishable or remediable under the P.H. Acts but not under the Factory Act, is found by the Factory Inspector in a factory or workshop, he is to report it to the S.A., who must make enquiry, take such action as may seem proper, and inform the Factory Inspector what has been done. If proceedings are not taken by the S.A. within a month, the Factory Inspector may institute proceedings in default, and recover from the S.A. any expenses not recovered from other persons, and not incurred in unsuccessful proceedings. If the S.A. fail generally to carry out their

duties with regard to factories, workshops or workplaces, the Home Secretary may authorise the Factory Inspector to act in default for a specified period, and the expenses not recovered from other persons may be recovered from the S.A.

The M.O.H. is required in his Annual Report to report specifically upon the administration with regard to workshops and workplaces, and to send a copy to the Home Secretary. A table issued by the L.G.B. for the guidance of the M.O.H. includes the items calling for statistical record.

Outside the Factory Act (which is directed to the protection of the workers), the S.A. are concerned with certain factories and workshops on public health grounds, as sources of smoke, poisonous fumes, trade effluents; as places where offensive trades or food manufacture needing control are carried on; as new buildings needing approval of plans, or old buildings in a dangerous state; or more indirectly as places in which infection may be spread from one person to another, or (as in laundries)

infected articles may be handled.

Nor are these the only points claiming the attention of architects, builders, and S.A.'s in connection with plans for new factories and workshops. Others are strength of foundations and structure generally where heavy machinery is to be fixed; means of escape from fire; sanitary accommodation. Others, again, are the statutory requirements as to air space, ventilation and warming; the surfaces of walls and ceilings must be such as to admit of due cleansing, and the proper construction of floors (soundness, impermeability, slope, drainage) is of importance in certain processes. In certain dangerous trades additional structural requirements arise, and in some the plans have also to be approved by the Chief Inspector of Factories.

Cleanliness.—Workshops and workplaces must be kept in a cleanly state; and a workshop, or any specified part thereof (in London, a workplace or domestic factory also), must be limewashed, cleansed, or purified when so required by notice from the S.A. on certificate of the M.O.H. or Sanitary Inspector that it is necessary for the health of the workers.

Factories must be kept in a cleanly state and (subject to certain exceptions) either (a) limewashed

every 14 months, or (b) painted with oil or varnished every 7 years, and washed with soap and hot water every 14 months.

For the cleaning of floors, etc., damp processes are

preferable, as checking the production of dust.

Air-space.—Factories, workshops and work-places must not be so overcrowded as to be dangerous or injurious to the health of the workers, and there must be at least 250* c.f. of air-space for each worker. The air-space of each workroom must be stated in a notice affixed in the works.

A much larger air-space is required in certain dangerous trades, and in works where certain special exceptions are permitted—e.g. in match factories in which yellow phosphorus is used the minimum is 400 c.f., and height above 14 feet is not counted, and in engineering works exemption from routine annual limewashing is allowed if the air-space be 2,500 c.f. per head.

General ventilation must be efficient in every workroom of a factory or workshop (other than a men's workshop). Standards† of ventilation are prescribed for certain workrooms. A departmental Committee in 1902 recommended a general limit for

workrooms of 12 parts of CO, per 10,000.

Air samples are usually taken at the breathing level, at points where work is carried on, and not in extreme corners or near to inlets or outlets, or close to persons or to sources of CO₂. In breweries, aërated water factories, bakehouses, etc., where CO₂ is given off by trade processes, allowance has to be made; and similarly where artificial lighting yields CO₂. As a rule it is well to make a parallel determination in the open air.

Mechanical ventilation by fans may be either on the plenum or vacuum system. The number, position, dimensions and speed of the fans are matters requiring expert knowledge; as

* 400 during overtime of women, 500 in underground bakehouses, 400 between 9 p.m. and 6 a.m. in other bakehouses with light other than electric, 400 in workshops (not domestic) used also as sleeping rooms.

t e.g.—A limit of 9 vols. CO₂ per 10,000 in humid cotton cloth factories, and in flax, hemp and jute works, and 600 c.f. fresh air per head per hour in other humid textile factories.

a rule better results are obtained by large fans moving slowly than by small fans at a high speed. Plenum is on the whole preferable, and offers facilities for warming, screening or humidifying the air when necessary. Exhaust ventilation calls for care in guarding against indraught through closets or other unsuitable channels. Paddle fans, which churn the air without changing it, are not to be regarded as means of ventilation, though the currents they cause may promote evaporation, and thus have some cooling effect. If reliance is placed on "natural" ventilation (whether by chimneys, open windows and doors, ventilators or shafts), supervision becomes necessary, especially in cold weather. In any case care should be taken to secure ventilation of all parts of the room, to avoid short-circuiting and direct draughts upon workers, and to provide adequately for both inlets and outlets.

Local ventilation, for removal of dust, gases, vapours or other impurities generated in the course of work, which are a nuisance or injurious to health, must be such as to render them harmless as far as possible.

Usually this involves exhaust ventilation, applied at or near the point of origin; and fans (or other efficient mechanical means) for the purpose are prescribed in certain dangerous processes, and can be required by the Factory Inspector in others. Exhaust ventilation may sometimes be secured by connection with flues or other heated shafts, but in general fans are required and need careful planning. The essentials are that the dust or fumes should be intercepted as near as possible to their source, and carried away from the workers, without entering the general air of the room: that the draught should be adequate for the purpose, and guided where necessary by hoods: and that the air-ducts should be kept clean and discharge at a suitable point, preferably in a closed dust-chamber. As a rule a downward draught is best, since it carries away the dust more readily. The efficiency of exhaust can be tested by anemometer, or by smoke, light dust, or fumes of ammonium chloride. In some instances standards are fixed, in terms of sectional area of openings and velocity at given points.

Other measures are the use of closed chambers or boxes, and as regards dust the adoption of wet processes, handling the dusty materials over gratings through which the dust can fall, cleanliness of premises and floors, and the use of overalls.

As a last resource respirators may be adopted, especially

where the occupation is intermittent, but there is difficulty in enforcing the systematic use of them, and most of them entail some physical discomfort if worn for long periods or in heavy work. The essentials in a respirator are that it should be light, simple, inexpensive, the filtering material readily renewed or cleaned; that it should be as little unsightly as possible; that it should arrest all dust, filter the whole of the inspired air, and prevent the rebreathing of expired air; and that it should offer the minimum resistance to the free passage of air without respiratory effort, and without becoming obstructed in use by moisture from the breath. A fold of cambric covering the mouth and nostrils is an efficient simple respirator. Many elaborate forms fail when tried under the actual conditions of the work for which they are proposed.

Some kinds of dust are directly infective, e.g. anthrax, tetanus, small-pox (among rag-sorters in paper works and elsewhere); others poisonous (e.g. lead, mercury); and others, again, mechanically injurious in varying degree. Similarly the wide variety of gases, fumes and vapours which arise in manufacturing processes and require removal include many which are directly poisonous (e.g. CO, CS₂, H₂S, SO₂, chlorine, nitrous and other acid fumes, AsH₃; lead, zinc and mercury fumes), and some which are harmful in other ways (e.g. steam)

or offensive (organic effluvia).

Explosion may arise from ignition of inflammable dust or fumes mixed with air: e.g. ordinary gas, marsh gas (in coal

mines), naphtha, coal dust, or celluloid borings.

Temperature.—In every workroom of a factory or workshop provision must be made for securing a reasonable temperature, and without interfering with the purity of the air of the room. This requirement is enforced by the Factory Inspector. Gas jets cannot be accepted as means of warming, nor must the ventilators be closed. Flueless stoves are objectionable, owing to the risk of adding sulphur compounds,

CO, and other impurities to the air.

As a lower limit 60° F. is suitable for sedentary occupations involving little exertion, but active work can be carried on, indoors as well as in the open air, at lower temperatures. The upper limit is not easily defined, since in certain processes a high temperature is a trade necessity, while in others it is only accidental (e.g. from use of furnaces, consumption of gas, moving machinery, steam pipes), and calls for preventive measures such as free ventilation, screening off sources of heat, "lagging" of hot pipes and reducing their dimensions as far as practicable. In humid cotton cloth factories the prescribed precautions of the kind include whitewashing of the roof in

summer, and it is forbidden to raise the temperature artificially above 70° F., except so far as is necessary in humidifying.

If the air be dry enough, high temperatures can be borne, but as a matter of health as well as comfort high wet-bulb temperatures are to be avoided as far as possible, especially

where the work is hard.

Humidity is expressly regulated in humid textile factories generally,* a limit being fixed for each dry-bulb degree: 88% from 59° to 70°, decreasing thenceforward to 77.5% at 80°, 69% at 90°, and 64% at 100°. In certain textile processes, however, a difference of 2° between wet and dry bulb is accepted at all temperatures. Hygrometers must be affixed in each room, and (except in cotton spinning) records kept of two (in cotton cloth factories three) daily readings at [7 to 8 a.m.], 10 to 11 a.m., and 3 to 4 p.m. The water used for humidification must either be taken from a pure source or be effectively purified; a water which gives off offensive gases of any kind on heating, or which absorbs from acid solution of KMnO4 at 60° F. in 2 hours more than 0.5 mgr. of oxygen, may be regarded as unsuitable. In moist air the clothing becomes damp, and warmed ventilated cloak-rooms are obligatory in new humid cotton weaving factories.

Steam in excess arises incidentally in laundries, dyehouses, etc., and can be removed by exhaust ventilation or by intro-

ducing hot dry air.

Lighting is only regulated by the Factory Act as regards underground bakehouses (p. 568) and in certain dangerous trades. For artificial lighting electric light or ventilated gas burners are preferable, as they do not add to the impurity of the air or call for additional ventilation.

Floors have to be sound and impervious for the purpose of certain dangerous processes, and to be drained in wet processes generally (except in men's workshops) and especially in laundries.

Sanitary accommodation is enforced by S.A.'s in both factories and workshops, in London (Pub. Health Act, 1891, s. 38) and where s. 22 of the P.H. Acts Amendment Act of 1890 is in force (p. 508). Elsewhere the S.A. have similar powers under s. 38 of the P.H. Act, 1875; but s. 9 of the Factory Act also applies, and the administration of this (and of

^{*} Spinning and weaving factories in which the air is artificially humidified by steam jets or otherwise, in order to correct the dryness of the air, which beyond a certain limit interferes with the treatment of the fibre

the order which interprets it) rests with the Factory IInspector alone.

Factory and Workshop Act, 1901: s. 9. "(a). Every factory and workshop must be provided with sufficient and suitable accommodation in the way of sanitary conveniences, regard being had to the number of persons employed or in attendance at the factory or workshop, and also where persons of both sexes are or are intended to be employed or in attendance, with proper separate accommodation for persons of each sex. (b). The Secretary of State shall, by special order, determine what its sufficient and suitable accommodation within the meaning of this section."

Sanitary Accommodation Order of February 4th, 1903.

"In pursuance of Section 9 of the Factory and Workshop Act, 1901, I hereby determine that the accommodation in the way of sanitary conveniences provided in a factory or workshop shall be deemed to be sufficient and suitable within the meaning of the said section if the following conditions are complied with and not otherwise:—

"1. In factories or workshops where females are employed cor in attendance there shall be one sanitary convenience for

eevery 25 females.

"In factories or workshops where males are employed or in attendance there shall be one sanitary convenience for every 25

males; provided that—

(a) in factories or workshops where the number of males employed or in attendance exceeds 100, and sufficient urinal accommodation is also provided, it shall be sufficient if there is one sanitary convenience for every 25 males up to the first 100, and one for every

40 after;

(b) in factories or workshops where the number of males employed or in attendance exceeds 500, and the District Inspector of Factories certifies in writing that by means of a check system, or otherwise, proper supervision and control in regard to the use of the conveniences are exercised by officers specially appointed for that purpose, it shall be sufficient if one sanitary convenience is provided for every 60 males, in addition to sufficient urinal accommodation. Any certificate given by an Inspector shall be kept attached to the general register, and shall be liable at any time to be revoked by notice in writing from the Inspector.

"In calculating the number of conveniences required by this order, any odd number of persons less than 25, 40, or 60,

as the case may be, shall be reckoned as 25, 40, or 60.

"2. Every sanitary convenience shall be kept in a cleanly state, shall be sufficiently ventilated and lighted, and shall not communicate with any work-room except through the open air or through an intervening ventilated space; provided that in work-rooms in use prior to 1st January, 1903, and mechanically ventilated in such manner that air cannot be drawn into the work-room through the sanitary convenience, an intervening ventilated space shall not be required.

"3. Every sanitary convenience shall be under cover and so partitioned off as to secure privacy, and if for the use of

females shall have a proper door and fastenings.

"4. The sanitary conveniences in a factory or workshop shall be so arranged and maintained as to be conveniently accessible to all persons employed therein at all times during

their employment.

"5. Where persons of both sexes are employed, the conveniences for each sex shall be so placed or so screened that the interior shall not be visible, even when the door of any convenience is open, from any place where persons of the other sex have to work or pass; and, if the conveniences for one sex adjoin those for the other sex, the approaches shall be separate."

Other sanitary regulations.—Lavatories are required where any poisonous substance is used. For certain dangerous trades details are prescribed, e.g. that there shall be soap, nailbrushes, towels, and at least one basin (with water supply and waste pipe) for every 5 persons. In some, baths also are required. Cloak-rooms are obligatory in new humid cotton cloth factories, and where certain dangerous trades are carried on.

In certain rooms, e.g. where poisonous substances are so used as to give rise to dust or fumes, it is forbidden to take food or to remain during meal hours, and accommodation for meals

must be provided elsewhere on the premises.

Dangerous trades, certified as such by the Home Secretary, are further subject to regulations (of automatic application) or special rules (established in individual works) issued by the Home Office. These prescribe additional precautions, according to the nature of the danger, and are enforced by the Factory Inspectors.

The Acts allow certain special exceptions, which are often coupled with additional requirements, sanitary and other; these also are administered solely by the Factory Inspectors.

Certifying surgeons are appointed for every locality. Their duties include (1) examination of children and of young

persons under 16 for certificates of fitness for employment in factories and certain workshops, (2) investigation of accidents and poisoning, (3) periodical examination, when required by regulations or special rules, of persons employed in processes

entailing danger of poisoning.

The dangers to health against which the above provisions are directed are obvious, but there are, of course, many others of a different kind, not limited to employment under the Factory Act. Thus coal miners are liable to nystagmus from working in constrained positions, and to "beat hand" and "beat knee" (chronic inflammation due to local pressure); men working in compressed air to "caisson disease" if care be not taken to pass slowly from higher to lower pressures, through air-locks. Writer's cramp, housemaid's knee, the emphysema of performers on wind instruments, are other examples. Hernia or aneurysm may be determined by violent exertion; carrying unduly heavy weights is injurious, and is indirectly prohibited as regards girls under 18 as a condition of special exception as to hours in fruit preserving. Sight may suffer from strain in fine work in inadequate light, and cataract is common among glass workers. Habitual exposure to weather in open-air occupations, and violent alternations of temperature in connection with furnaces or otherwise, present dangers of their own.

Means of escape from fire.—Factories built after 1891 and workshops built after 1895, must, if more than 40 persons are employed, have a certificate from the S.A. as to reasonable adequacy in this respect, specifying in detail the means provided. In older works with more than 40 hands the S.A. must see that reasonable provision is made, and if not, must require the owner by notice to carry out specified details in a specified time: any dispute to be settled by arbitration. The whole of a tenement factory or tenement workshop is counted as one. The S.A. may, subject to confirmation by the L.G.B., make byelaws as to means of escape in any factories or workshops. In London these duties rest with the County Council.

Doors of workrooms must not during working hours be fastened so as not to be readily opened from inside: and in

works built after 1895, doors of rooms in which more than 10

persons are employed must not open inwards.

Adequate means of escape are all the more essential where the nature of the manufacture entails special risk of fire. Alternative paths of escape are desirable (since the main line of exit may be readily involved in the fire) especially where large numbers are employed, and these paths should be obvious and

kept clear of unnecessary obstruction.

Laundries are technically factories or workshops. In factory laundries fans are required for regulating the temperature of ironing-rooms, and for removing steam from washhouses; and in all laundries (a) stoves for heating irons must be sufficiently separated from the ironing-room or ironing-table, and gas-irons emitting noxious fumes must not be used; (b) floors must be kept in good condition and so drained as to allow the water to flow off freely.

Bakehouses are factories or workshops, but subject to certain special requirements as below. Retail bakehouses are workshop bakehouses in which the articles baked are sold by retail only in a shop occupied therewith. Underground bakehouses are those in which any room used for baking or processes incidental thereto has its floor more than 3 feet below the footway of the adjoining street or ground adjoining or nearest to the room.

If a bakehouse is on sanitary grounds unfit for that use, a Court may, on application by the Factory Inspector or S.A., impose a penalty and require the necessary alterations to be made. It must not contain or communicate directly with a closet, nor derive its water supply from a cistern supplying a w.c. A drain or pipe for carrying off fæcal or sewage matter must not have an opening in the bakehouse (thus any wastepipe must be disconnected outside). The walls and ceilings of rooms, passages and staircases must be limewashed every 6 months; or receive 3 coats of oil-paint or varnish every 7 years, and be washed with hot water and soap every 6 months. A part of the same building, on the same level with the bakehouse, must not be used as a sleeping-room unless partitioned off from floor to ceiling, and having an external glazed window 9 square feet in area, 41 of which is made to open.

An underground bakehouse must not be used as such unless (a) it was so used on August 17th, 1901, and further (b) the S.A. has certified it to be suitable as regards construction,

light, ventilation and in all other respects.

The administration of these special requirements in retail bakehouses (in London in all workshop bakehouses), and the certification of underground bakehouses, rests with the S.A.

an other cases the Factory Inspector and S.A. have concurrent owers.

A high standard of sanitation is needed in premises where ood is manufactured, prepared, cooked or stored; including akehouses, restaurant kitchens, ice-cream premises, fish-shops, deat-shops, preserved meat works, confectionery works, airies and milk-shops. The following summary, directed rimarily to bakehouses, may serve to indicate generally the nief points needing attention.

Surfaces of floors, walls and ceilings to be hard, smooth,

urable and impervious.

Floors to be laid to proper falls, and drained by proper

nannels to a gully outside.

Walls to be absolutely dry; failing this, to have an inner all at least 1-inch thick, tied to the outer wall, but with 2-inch entilated and drained intervening space.

Any drain beneath the floor to be gas and water-tight. All

pes and fittings to be efficient.

Minimum height 8 feet; minimum air-space 400 c.f. per ead and 1,500 c.f. in all, clear of ovens, machinery, and stored sbocd.

Efficient lighting: where practicable natural light only; mong artificial lights electric is preferable, or if gas is necessary, candescent burners protected from dust and ventilated to emove fumes.

Ventilation, by fan if necessary; ensuring constant and infficient change of air without draughts, and a temperature not dinarily exceeding 60° F., or 80° F. at any time except ithin half-an-hour after "drawing" an oven.

Oven furnaces to have efficient flues to carry away fumes.

An outlet for heat and steam immediately before and above ne oven door. Oven doors and outer doors of furnaces to fit osely. Proving ovens to be efficiently ventilated. If gas is sed for heating, means of removing fumes. Hoods and ducts ading to the open air, over stoves and ranges.

Strict cleanliness of premises and appliances.

Water supply, with jet and hose, for washing floors; cisterns f any) to be of proper construction, with dust-tight covers, and ept clean.

Tables, benches, dough-troughs, etc., to be freely movable

a casters.

Any necessary shelves to be fixed 2 inches away from walls, ut unnecessary woodwork on walls to be avoided.

Covered metal boxes for refuse.

Suitable provision, outside the workroom, for storage of food naterials (milk in clean covered vessels, in a cool place); storage of fuel; deposit of clothing not in use; lavatory; sanitary accommodation.

Home work of certain kinds* defined (1907) by order of the Home Secretary is under special control. The employer must keep a list of the names and addresses of the outworkers, and send copies to the S.A. of his district on or before February 1st and August 1st of each year, in a prescribed form. If any such addresses are outside the area of the S.A., particulars of them must be sent by the S.A. to the S.A. concerned. If the home premises are injurious or dangerous to the health of the workers, the S.A. may prohibit the employer from giving out work to be done there.

If an inmate of the house is suffering from a notifiable infectious disease (even if he has been removed) the S.A. may by order prohibit the employer from giving out work to be done in that house or in any specified part of it, during a specified term. This power applies only to the classes of work named in the first part of the footnote below. In emergency it may be exercised by two or more members of the S.A. on the advice of the M.O.H. The order may be subject to conditions as to disinfection or other precautions.

In case of scarlet fever or small-pox, an employer giving out work or wearing apparel is subject to

The making of electro-plate; files; iron and steel cables and chains, anchors and grapnels; cart gear, including swivels, rings, loops, gear buckles, mullin bits, hooks, and attachments of all kinds; locks, latches,

and keys.

^{*} The making, cleaning, washing, altering, ornamenting, finishing, and repairing of wearing apparel; the making, ornamenting, mending, and finishing of lace and of lace curtains and nets; cabinet and furniture making and upholstery work; fur-pulling; the making or repairing of umbrellas, sunshades, parasols, or parts thereof; the making of artificial flowers; nets other than wire nets; tents; the making or repairing of sacks; the covering of racquet or tennis balls; the making of paper bags, boxes or other receptacles or parts thereof made wholly or partially of paper, card board, chip, or similar material; the making of brushes; pea picking; feather sorting; the carding, boxing, or packeting of buttons, hooks and eyes, pins, and hair pins; the making of stuffed toys; baskets; and any processes incidental to the above.

The making of electro-plate; files; iron and steel cables and chains,

penalty, apart from any order from the S.A. as above, unless he proves that he was not aware of the illness and could not reasonably have been expected to become aware of it.

Industrial poisoning .- "Every medical practtitioner attending on or called in to visit a patient whom he believes to be suffering from lead, phosphorus, arsenical or mercurial poisoning, or anthrax, contracted in a factory or workshop, shall . . . send tto the Chief Inspector of Factories, at the Home Office, London, a notice stating the name and full postal address of the patient and the disease from which in the opinion of the medical practitioner the patient is suffering." A fee of 2s. 6d. is allowed. The employer must notify such cases to the Inspector cof Factories and to the Certifying Surgeon.

Lead poisoning arises in a number of industries, from the materials or appliances used. Owing to more stringent precautions the number of reported cases in the United Kingdom fell from 1,258 in 1899, to 632 in 1906, the eight years' totals being-

*White Lead	1,512	Ship-building	235
*Earthenware & China		Printing	149
Coach-building	527	*Tinning & Enamelling	
*Paints and Colours	398	of Metals	142
*Smelting of Metals	311	Other ,	1,464
*File Cutting	225	Manufacture and the sales	-
*Electrical Accumulators	244	Total	6,243

Those marked * are under special rules or regulations as dangerous trades. The great danger is that of inhalation or swallowing of dust, but there is also risk from carrying unwashed hands to the mouth, from eating or using tobacco while at work, and in some industries from fumes containing lead. The chief preventive measures, apart from those directed against dust and fumes (p. 562), are the substitution of materials free from lead or containing it only in an insoluble and comparatively harmless form, facilities for washing and due use of them, overalls in dusty processes, periodical medical examination of workers, and exclusion of persons found to be susceptible.

Women appear to be more susceptible than men, and the employment of children and even of young persons under 15 is forbidden in certain lead processes, as less attention to pre-

cautions can be expected from them.

The figures given above are those for industries as a whole. each including a variety of processes involving very different degrees of risk: thus in pottery processes the incidence is much greater among dippers and ware-cleaners, who handle the glaze before it is fired; and in the processes named much depends upon the nature of the materials handled, the frequency and duration of employment, the conditions of work, including observance of precautions, and upon the age and sex of the workers. These considerations are material in estimating the statistical value of attack-rates or mortality rates, in all industrial diseases and not merely as regards plumbism. A further necessary consideration is the character of attack, which varies greatly as regards the nature, site, and severity of symptoms, and shades off into constitutional conditions which can hardly be regarded as poisoning or disease, and may only be revealed by large statistics showing greater tendency to more remote affections, such as the kidney disease and gout which are found from mortality returns to be more common among lead-workers than others.

Phosphorus poisoning is now rare, stringent precautions being in force in the lucifer match factories in which yellow phosphorus is used. Its characteristic manifestation is necrosis of the jaw, arising in connection with carious teeth: it rarely occurs until the employment has been long continued, and there is some reason to suppose that it is dependent upon a constitutional change predisposing to local attack. Substitutes for yellow phosphorus have been found in red phosphorus (a harmless form), sesquisulphide of phosphorus, and other Other precautionary measures in force are the chemicals. use of closed vessels for phosphorus, separation of phosphorus processes from each other and from the rest, free general ventilation, exhaust ventilation, washing facilities and systematic use of them, overalls, and periodical examination by a dentist (with power to forbid employment), in order to secure sound-

ness of teeth.

The reported attacks in 1899-1906 were 8, 3, 4, 1, 0, 1, 3, 0. Arsenical poisoning arises chiefly in connection with paints and colours and in chemical works, or exceptionally from the evolution of arseniuretted hydrogen in metallurgical processes. In 1900-1906, 55 attacks were reported.

Mercurial poisoning is met with in industries in which mercury or a mercurial salt is used: including the making of thernometers and other scientific instruments, furriers' work (in carrotting") and chemical works. Formerly it was observed mong "water gilders" employed in "silvering" mirrors by neans of mercury. In addition to salivation, alimentary isturbances and cachexia, there are peculiar fine tremors of the ands and arms, with occasional pains in the joints. In 1899-906, 68 attacks were reported.

Anthrax was reported in 386 instances in 1899-1906.

INDUSTRY.	ATTACKS.	DEATHS.
Wool, Worsted	146	43
Horsehair, Bristles	87	23
Hides, Skins	121	30
Other	32	10

Internal anthrax is generally fatal. Among external cases he pustule is usually (85%) on the head or neck, but in 12°/ on the hand or arm.

The infection in the trades named above is due to materials imported from Eastern countries in which anthrax is rife, e.g., Wan mohair, Persian wool, Siberian and Chinese horsehair, Chinese and Persian skins. Dry and dusty materials and processes are especially dangerous, and greasy or moist goods e.g. salted hides) less so. Where disinfection is impracticable and infected materials are used, the preventive measures burn upon suppression of dust, personal cleanliness, protective garments, and immediate attention to abrasions of the skin, however slight. Early diagnosis, prompt excision of anthrax pustules, and the use of Sclavo's serum are important.

Brass-founders' ague, attributed to zinc fumes, is characterised by recurrent attacks, with cold, hot, and sweating stages, differing from those of malaria in not being periodic.

Other provisions of the Factory Act require only brief mention here, and the following summary may suffice.

Safety.—Dangerous machinery must be fenced or made safe by position. The use of dangerous ways, works, machinery or plant, or of dangerous premises, can be forbidden by a Court. Certain dangerous employments are forbidden to "children" and "young persons," and further limitations of this kind can be imposed by the Certifying Surgeon. Where any manufacture, plant, process, or description of manual labour is certified by the Home Secretary as dangerous or injurious to health, or dangerous to life or limb, regulations (of general application) or special rules (for particular works)

may be made: many such codes are in force.

Accidents which are fatal, or which cause disablement for ordinary work for seven days (or, if caused by power machinery, molten metal, hot liquid, explosion, escape of gas or steam or electricity, such disablement for one whole day) must be registered and reported to the Factory Inspector; and also (except the seven-day accidents above) to the Certifying Surgeon. These obligations arise under an Amending Act (Notice of Accidents Act, 1906) and extend to dock processes and building operations. About 100,000 accidents per year are

so reported, and of these about 1% are fatal.

Age and Sex.—Children under 12 may not be employed. Above 12 they may be employed for half-time (if they have attained a certain standard and if the local educational byelaws permit), but must attend school for the other half. In factories and certain workshops they must have a certificate of fitness (as to age and physique) from the Certifying Surgeon, which may be coupled with conditions as to the nature of the employment permitted. "Young persons" are those between 14 and 18 (or between 13 and 14, subject to conditions as to educational certificates), and are eligible for full time employment, with certain limitations. Certificates of fitness are required for those under 16 employed in factories and certain workshops. A woman must not be employed within 4 weeks after giving birth to a child.

Hours and Intervals.—The period of employment for women and young persons in textile factories is on ordinary days 12 hours (6 to 6 or 7 to 7), less 2 hours intervals, and on Saturday 6 hours (6 to 12 or 7 to 1, with $\frac{1}{2}$ hour interval, or 6 to 12.30, with an hour interval): or a weekly total of $55\frac{1}{2}$ working hours. In non-textile factories and in workshops it is 12 hours on ordinary days (6 to 6, 7 to 7, or 8 to 8), with $1\frac{1}{2}$ hours intervals, and on Saturdays 8 hours (6 to 2, 7 to 3, or 8 to 4), with $\frac{1}{2}$ hour interval; or a weekly total of 60 hours.

In laundries additional latitude is allowed. There is a further limit as to continuous spell of employment, which must not exceed $4\frac{1}{2}$ hours in textile factories and printing, bleaching and dyeing works, or 5 hours in non-textile factories and workshops, without an interval of $\frac{1}{2}$ hour. Certain holidays, amounting to six days in the year, are prescribed. Work on Sundays is prohibited. The hours, intervals and holidays must be fixed by notice. None of the above provisions

apply to men over 18, and certain exceptions are allowed as egards women, young persons and children in special cases: .g. women may be employed "overtime" for 2 hours, on not nere than 3 days in any week, nor more than 30 days in any 2 months, in certain specified branches; and in others male coung persons can be employed at night.

The Alkali, etc., Works Regulation Act, 11906, applies to a wide variety of chemical and other works, and is administered by Inspectors appointed by the L.G.B. It regulates inter alia the discharge of acid gases into the atmosphere; thus in alkali works the escaping air or smoke must not contain more than ½ grain of muriatic acid per cubic foot, and in other specified works the acid gases discharged must not exceed the equivalent of $1\frac{1}{2}$ grains of sulphuric anhydride per cubic foot. The best practicable means must be adopted for preventing the escape of noxious and offensive gases, and rendering them harmless and inoffensive. Acid drainage must not be allowed to mix with alkali waste so as to cause nuisance. The owner may require the S.A. to prowide and maintain at his expense a drain for carrying the acid waste into the sea or any watercourse into wwhich it can be taken without breach of the law. Alkali waste must not be deposited or discharged without the best practicable and available means being used to prevent nuisance.

On complaint from the S.A. or from ten inhabitants, that nuisance arises from contravention of the Act, the L.G.B. must institute such inquiry and

direct such proceedings as they think fit.

Mines are controlled by the Coal Mines Regulation Act, 1887, and the Metalliferous Mines Regulation Act, 1872 (many of the provisions of which are extended to quarries, over 20 feet in depth, by the Quarries Act, 1894), and amending Acts. These Acts are administered by H.M. Inspectors of Mines, but in one or two minor points (see pages 491 and 496) they give powers under the Public Health Acts to the S.A.

The Shop Hours Acts, 1892, 1893, and 1895, are administered by the County and Borough Councils. They contain no sanitary provisions, but regulate the hours of employment of young persons under 18, whether wholly in a shop or partly also in a factory or workshop. The total number of hours, including meal hours, must not exceed 74 in any week, and a notice of this limit must be kept affixed in the shop. "Shops" include warehouses and licensed public houses and refreshment houses of all kinds. A further Act (Seats for Shop Assistants Act, 1899) requires one seat to be provided for every three female assistants.

The Employment of Children Act, 1903, enables urban S.A.'s with a population over 20,000 (or, in the case of boroughs over 10,000, and elsewhere the County Council) to make byelaws regulating the employment of children under 14 (not under the Factory or Mines Acts) and street trading by persons under 16; it forbids carrying of heavy weights or other injurious occupation, street trading by children under 11, and (in general) work between 9 p.m. and 6 a.m.; but as regards children employed in mines, factories, and workshops, these provisions are administered by Home Office Inspectors only.

Under the Prevention of Cruelty to Children Act, 1904, the S.A. are charged with the duty of seeing to the observance of the conditions of licences granted by a Court for the employment of children over 10 in public entertainments.

The Workmen's Compensation Act, 1906, deals with certain industrial diseases as well as accidents. Where (a) the Certifying Surgeon (or other practitioner appointed by the Home Secretary for the purpose) certifies that a workman is suffering from a disease named below, and thereby disabled from earning full wages at the work on which he was employed, or (b) a workman is suspended in pursuance of Regulations or special Rules from his usual employment on account of having contracted such a disease, or (c) the death of a workman is caused by such a disease, claim to compensation arises if the disease can be shown to be due to the nature of the employment. In certain specified cases (col. 2) this industrial causation is assumed, unless the Certifying Surgeon

certifies, or the employer proves, the contrary. Additions to the Act can be made by order of the Home Secretary.

Disease. (1)
Proisoning (or the sequelæ of such poisoning) by— Lead Mercury Phosphorus Arsenic Carbon bisulphide Nitro- and amido-derivatives of benzene (e.g. anilin, di-nitro-benzol) Nitrous fumes Niekel carbonyl African boxwood (Gonioma Kamassi)
AAnthrax
CChrome ulceration, or its sequelæ
PEczematous ulceration of the skin, produced by dust or caustic or corrosive liquids Ulceration of the nose or mouth produced by dust Epithelioma of the skin or cornea due to pitch tar or tarry compounds Epithelioma of the scrotum Glanders
Compressed air illness (caisson disease) Nystagmus "Beat hand" \ subcutaneous "Beat knee" \ cellulitis Beat elbow(acute bursitis) Synovitis of wrist joint and tender

Processes.

Any process involving the use of these, or their preparations or compounds

Any process in which these gases are evolved Manufacturing articles therefrom

Handling of wool, hair, bristles, hides, skins Use of chromic acid, or bichromates, or their preparations

Handling or use of pitch tar or tarry compounds

Chimney-sweeping
Care of any equine animal suffering
from glanders; handling the
carcase of such animal
Work in compressed air

Mining

sheaths Ankylostomiasis

CHAPTER XX

BYELAWS AND REGULATIONS

Byelaws, having the force of law, may be made by Sanitary Authorities for the better government of their districts, but only in respect of certain matters, and under certain conditions, expressly stated in They are designed to the Acts of Parliament. supplement, not to summarise, vary, or supersede, the express provisions of the statute law. It is, of course, quite open to any S.A. to append to their byelaws a statement or summary of the requirements of the statute law bearing upon the same matters. some cases this course is advised by the L.G.B. Byelaws must be reasonable, and must not in any particular be repugnant to the provisions of the general law. They have (as a rule) no force until approved by the L.G.B. Any byelaw may be altered or repealed by a subsequent byelaw. In framing byelaws, a S.A. should impose reasonable penalties, not exceeding £5 for each offence; and in the event of a continuing offence, a further penalty not exceeding £2 per day.

Urban authorities are empowered to make bye-

laws as follows :--

1. Private scavenging.—For imposing upon occupiers the duty of cleansing pavements and footways, of removing house-refuse, and of cleansing earth-closets, ashpits, privies, and cesspools. Such byelaws are only to be made if the **S.A.** do not undertake or contract for these matters * (Public Health Act, 1875, s. 44).

2. Prevention of nuisances.—For the prevention of nuisances arising from snow, filth, dust, ashes, and

^{*} But see the Public Health Acts Amendment Act, 1890, p. 508.

- rubbish. For the prevention of the keeping of named and premises so as to be injurious to health Public Health Act, 1875, s. 44).
- 3. Common lodging-houses.—For regulating the number of lodgers and the separation of the sexes; for promoting cleanliness and ventilation; for giving notice and taking precautions in case of infectious disease; and for the general well-ordering of such nouses (Public Health Act, 1875, s. 80).
- 4. Houses let in lodgings.—For regulating the number of persons and separation of the sexes in a mouse or part of a house let in lodgings or occupied by members of more than one family; for the registration and inspection of such houses; for drainage, privy accommodation, cleanliness and ventilation; for cleansing and whitewashing at fixed times; for paving of yards; and for giving notice and taking precautions in case of infectious disease * (Public Health Act, 1875, s. 90). The Merchant Shipping Fishing Boats) Act of 1883 gives power to the S.A. on make byelaws for the regulation of seamen's codging-houses.
- 5. Hop-pickers.—For securing the decent lodging and accommodation of persons engaged in picking the loops (Public Health Act, s. 314).
- 6. Fruit-pickers.—For securing the decent lodging and accommodation of persons engaged in picking fruit and vegetables [Public Health (Fruit-pickers) Act 1882].
- 7. Tents and vans.—For promoting cleanliness in, and habitable condition of, tents, vans, sheds, and similar structures used for human habitation; for pre-
- The M.O.H. must on request inspect any house wholly let in enements at rents not exceeding 7s. 6d. weekly, and (if satisfied) certify that the accommodation and sanitary arrangements are suitable. The effect is to exempt from Inhabited House Duty. The S.A. may appoint another practitioner, qualified to act as M.O.H., to give such certificate Customs and Inland Revenue Act, 1890).

venting the spread of infectious disease by the occupants thereof; and generally for the prevention of nuisances in connection with the same (Housing of

the Working Classes Act, 1885).

8. New streets and buildings.—With respect to the level, width, construction, and sewerage of new streets; with respect to the structure of walls, foundations, roofs, and chimneys of new buildings, for securing stability and for the prevention of fires, and for purposes of health; with respect to ventilation, and sufficiency of air-space about buildings; with respect to drainage, closets, ashpits, and cesspools; with respect to the depositing of plans and sections of proposed new streets and buildings; with respect to the power of the Authority to remove, alter, or pull down any work begun or done in contravention of the byelaws; and with respect to the closure of buildings unfit for habitation * (Public Health Act, 1875, s. 157).

9. Mortuaries.—For the regulation of charges and

management of public mortuaries (s. 141).

10. Cemeteries.—For the regulation of charges and management [Public Health (Interments) Act, 1879].

11. Slaughter-houses.—For the licensing, registering, and inspection of slaughter-houses and knackers' yards; for preventing cruelty therein; for cleanliness and daily removal of filth; and for providing a supply of water (Public Health Act, 1875, s. 169).

12. Offensive trades.—For preventing or diminishing the noxious or injurious effects of any offensive

trades (Public Health Act, 1875, s. 113).

13. Markets and fairs.—For preventing nuisances in markets and fairs; for inspection of slaughter-houses, and daily removal of refuse therefrom; for preventing the sale or exposure for sale of unwhole-some provisions; and for many other purposes (Public Health Act, 1875, s. 167).

^{*} For further powers, see pages 509 and 587.

14. Open spaces.—For the regulation of public walks and pleasure grounds (Public Health Act, s. 1164). For the regulation of any open space, churchyard, cemetery, or burial ground over which the S.A.

has control (Open Spaces Act, 1887).

15. The Municipal Corporations Act, 1882, s. 123, gives power to the Council of a borough to make byelaws for the prevention and suppression of nuisances not already punishable in a summary manner by virtue of any Act in force throughout the borough. County Councils have similar power under the 16th section of the Local Government Act, 1888.

16. For the regulation of buildings provided under

the Housing of the Working Classes Acts.

17. By adopting certain portions of the Public Health Acts Amendment Act, the S.A. may make by elaws as to public conveniences provided by them; and also additional byelaws as to new (and old) buildings, removal of house-refuse, and prevention of nuisances (page 509).

18. For means of escape from fire in factories and workshops (Factory and Workshop Act, 1901, s. 15).

Rural Authorities have similar powers in respect of the following:-

Private Scavenging (1, 17). Common Lodging-houses (3). Houses let in Lodgings, and Seamen's Lodging-houses (4). Hop-pickers (5). Fruit-pickers (6).

Tents and Vans (7). Mortuaries (9). Means of escape from fire, in factories and workshops (18).

And also under the Housing of the Working Classes Acts (16). By adopting portions of the Public Health Acts Amendment Act, they can make certain byelaws as to new (and old) buildings (8, 17). The L.G.B. may confer upon them any other powers as to byelaws which the Public Health Acts give to Urban Authorities (Public Health Act, 1875, s. 276).

Byelaws as to common lodging-houses must be made by every S.A., and as to slaughter-houses by every urban S.A.; the rest are optional.

Any S.A. may make regulations under the Dairies

Order (see p. 544).

A S.A. may make regulations (to be approved by the L.G.B.) for the removal to hospital, and detention in hospital as long as necessary, of persons brought within their district by ship or boat, and infected with any dangerous infectious disorder (Public Health Act, s. 125). They may also make regulations for the management of places provided by them for post-mortem examinations ordered by a coroner.

Model Byelaws have been prepared by the L.G.B. in respect of certain of the matters upon which S.A. have power to frame byelaws. The following

have been issued up to the present time:-

Private Scavenging.
Prevention of Nuisances.
Common Lodging-houses.
New Streets and Buildings.
Markets.
Slaughter-houses.
Hackney-carriages.
Public Bathing.
Baths and Wash-houses.

Pleasure Grounds.
Horses, etc., let for Hire.
Pleasure-boats.
Houses let in Lodgings.
Cemeteries.
Mortuaries.
Offensive Trades.
Hop-pickers.
Means of escape from fire, in factories and workshops.

These Models* are now very generally adopted, with occasional slight modifications, by **S.A.** seeking to frame byelaws. The following summary includes only those which have a more direct bearing upon sanitary matters:—

1. Private Scavenging.†—The occupier of any premises must cleanse the footways and pavements

^{*} The semi-official Annotated Model Byelaws (Knight and Co.) contains many important notes, additions, and illustrations.

† Applicable to districts where the S.A. do not undertake these duties.

adjoining his premises daily except Sunday. He must remove the house-refuse once a week, and excreta at intervals not exceeding the following maximum limits :-

Must be cleansed at least-Earth-closets, with fixed receptacle once in three months " movable " once a week. Privies, whether the receptacles are fixed) once a week. Ashpits, whether receiving excreta or once a week. Cesspools once in three months.

2. Prevention of Nuisances.

- (a) Clearing away snow.—The occupier of any premises must clear away snow from the footways and pavements adjoining his premises, as soon as possible after it ceases to fall.
- (b) Scavenging.—The refuse from any premises shall only be removed in a suitable covered receptacle or carriage; and if from premises within 20 yards of any dwelling, public building, or place of business only between [7.0 and 9.30 a.m.] from November to February, and between [6.0 and 8.30 a.m.] from March to October. Refuse must not be deposited upon any road, and any refuse accidentally falling upon a road must be immediately gathered up and the place cleansed.
- (c) Deposit of nightsoil and other refuse.—No load of filth must be deposited for more than [24] hours within [100] yards of any street, dwelling, public building, or place of business. Nightsoil deposited for agricultural purposes upon land within [100] yards of a street, dwelling, etc., and not deodorised, must at once be dug or ploughed into the ground.
- (d) Keeping of animals. Swine must not be kept within [100] feet of any dwelling, nor cattle where they may pollute water likely to be used for drinking, domestic, or dairy purposes, or for

manufacturing drinks. The same prohibitions apply to storage of dung. Premises wherein are kept any swine, cattle, horses, etc., must be provided with proper receptacles for manure, and with efficient drainage; the receptacle must be watertight, covered, and entirely above the level of the ground, and it must be cleansed at least once a week; the drain must be properly constructed and kept in order at all times, so as to convey all liquid filth to a sewer, cesspool, or other suitable receptacle.

3. Common Lodging-houses.

(a) Number of lodgers.—The S.A. may specify by notice in writing the maximum number of lodgers to be accommodated in each room, and the keeper must not allow this number to be exceeded; notice stating the maximum number allowed must be exhibited in each room.* The S.A. may vary the number from time

to time by further notice.

(b) Separation of sexes.—In general no person above ten years of age must occupy the same sleeping-room as persons of the opposite sex, but rooms may be set apart for the sole use of married couples, to the exclusion of other persons over ten years of age, on condition that every bed is screened off. No bed must be occupied by more than one male above ten years of age.

(c) Cleanliness of premises.—The yards, etc., must be kept clean and in good order; all floors swept daily and washed once a week; all windows, painted surfaces, and fittings of wood, stone, or metal, kept clean.

(d) Closets must be kept clean and in good and

efficient order.

(e) Ashpits must be kept clean and in good order. No filth or wet refuse must be thrown into ashpits designed for dry refuse only.

^{*} It is usual to require at least 300 cubic feet of air-space per head, but to count two children as one adult.

(f) Management of sleeping-rooms.—The windows must be opened fully for an hour in the morning, and an hour in the afternoon, except in case of stress of weather or occupation of the room by a sick person, or other sufficient cause. Beds must be stripped of blothes and exposed freely to the air for an hour each day, and must not be reoccupied within eight hours after being vacated. All refuse and slops must be removed every day before 10 a.m., and all utensils cleansed daily. Every sleeping-room must be provided with sufficient bedsteads, beds, bedclothes, and utensils, for the use of the maximum number of lodgers to be received therein.

(g) Facilities for washing.—A sufficient supply of suitable basins, water, and towels must be provided for the use of lodgers; and must be kept clean and

renewed as required.

(h) Precautions in respect of infectious diseases.— IIf the keeper finds that any lodger is suffering from an infectious disease, he must at once take all necessary precautions. No person, except a relative or lattendant, must occupy the same room as the sick person. If the patient is removed to hospital by the S.A. the keeper must afford all facilities for removal and must adopt all precautions directed by the M.O.H.

He must, if required by the M.O.H. to do so, temporarily cease to receive lodgers into any infected room. At the end of the case by removal, recovery, or death, the keeper must at once give notice to the M.O.H. and must cleanse and disinfect every part of the infected rooms and their contents, and in doing so must comply with all instructions of the M.O.H. When the cleansing and disinfection are completed, he must give notice thereof to the M.O.H., and must not receive any lodger into the rooms in question until two days after such notice has been given.

(i) A copy of the byelaws in force with respect to Common Lodging-Houses, supplied by the S.A., must be exhibited in some conspicuous place in the house, and must not be concealed, altered, obliterated or injured.

As some of the most important regulations regarding common lodging-houses are contained in the Public Health Act, and are, therefore, omitted in the byelaws, it is desirable that a statement of the provisions of sections 75 to 89 of the Public Health Act should accompany the copy of byelaws supplied to each common lodging-house.

As regards the question of the suitability of any particular premises or particular rooms for registration, a memorandum, issued by the L.G.B., points out that a common lodging-house should, in the first place, possess the conditions of wholesomeness needed for houses in general, and secondly, that it should have arrangements fitting it for the special purpose of receiving a given number of lodgers. Inside walls should not be papered. Every registered room should have special means of ventilation (by chimney if possible), and a window opening freely and directly upon the outer air. There should be kitchen and day-room accommodation apart from the bedrooms. Rooms partially underground should not be registered as sleeping rooms. There should be a supply of pure water allowing at least ten gallons per head per day for the maximum number of inmates, and one closet for every twenty registered lodgers. Wherever practicable, there should be a separate place for washing, not in the bedrooms, with fixed basins and trapped and disconnected waste-pipes.

4. New Streets and Buildings.

(a) Streets.—No new street must be less than 36 feet wide, if it exceeds 100 feet in length or is intended to be a carriage road: nor less than 24

feet in any case. One end at least must be quite

open.

- (b) Sites.—No buildings must be erected upon soil polluted with animal or vegetable matter. Sites in low and damp situations, near rivers or in excavations, must be elevated artificially. The site of a new house must be entirely asphalted or covered with 6 inches of concrete.
- (c) Walls of all new buildings must be constructed of good bricks, stone, or other hard and incombustible materials, properly bonded and solidly put together with good mortar compounded of good lime and clean sharp sand or other suitable material, or with good cement, or with good cement mixed with clean sharp sand. Every wall must have a proper damp course of durable and impervious material beneath the level of the lowest timbers and at least 6 inches above the ground. If the ground is to be in contact with a wall above the level of the floor of the lowest storey, that wall must be made double, with a cavity 21 inches wide extending from the base of the wall to 6 inches above the surface of the adjoining ground; and damp courses must be inserted both at the base of the wall and at the level of the top of the cavity. Walls of new houses must be at least 9 inches thick, increasing according to a prescribed scale when the height is greater than 25 feet, or the length greater than 30 feet. Party-walls must be carried up at least 15 inches above the roof, the distance to be measured at right angles to the slope of the roof. The L.G.B. now permit certain modifications in the byelaws for buildings in rural districts. (Circular, 5th Jan. 1906).

(d) Roofs must be made of incombustible materials,

and provided with gutters leading to rain-pipes.

(e) Open space.—A new house must have along its whole frontage an open space measuring at least 24 feet to the boundary of any land or premises

immediately opposite or to the opposite side of the street. In the rear there must be an open space exclusively belonging to the house, at least 150 square feet in area, and free from any erection above the ground level except a closet and an ashpit; the open space must extend along the entire width of the house, and must measure in no case less than 10 feet from every part of the back wall of the house; if the house is 15 feet high, the distance must be 15 feet; if 25 feet, then 20 feet, and if 35 feet or more, then 25 feet at least.

(f) Ventilation beneath floors.—If the floor of the lowest storey is boarded, there must be a clear space of at least 3 inches between the boards and the impervious covering of the site, and the space must be ventilated.

(g) Windows opening directly into the external air must be provided in every habitable room. The window area must be at least one-tenth of the floor area; at least half of each window must be made to

open, and it must open at the top.

(h) Ventilation. — Every habitable room must either have a fireplace and chimney, or a special ventilating aperture or air-shaft with an unobstructed sectional area of at least 100 square inches. Every new building must be provided with adequate means of ventilation.*

(i) Drainage.—Damp sites must be drained by earthenware field pipes properly laid to a suitable outfall, but not directly communicating with any sewer or cesspool or drain containing sewage. Rainpipes must be provided to carry away all water falling on the roof without causing dampness of the walls or foundations. The level of the lowest storey must be such as to allow of the construction of a drain sufficient

^{*} The semi-official Model Byelaws under the Public Health Acts Amendment Act prescribe a minimum height of 9 feet in every part, except in attics used as bedrooms, when a minimum height of 5 feet is permitted, if in two-thirds of the area the height is not less than 9 feet.

for the drainage of the building communicating with a sewer at a point above the centre of the sewer. All drains for sewage must be made of impervious pipes 4 inches or more in internal diameter, laid with a proper fall in a bed of concrete, and with watertight joints. Every drain inlet not intended for ventilation must be trapped. No drain conveying sewage must pass under a building unless no other mode of construction is practicable; in that case it must be laid in a direct line for the whole distance beneath the house, and must be embedded in and covered with concrete 6 inches thick all round, and must be laid at a depth below the surface at least equal to its diameter, and, lastly, must be ventilated at each end of the portion beneath the building. The main drain must be trapped at a point within the curtilage, but as distant as practicable from the building. Branch drains must join other drains obliquely in the direction of flow.

(j) Ventilation of drains.—There must be at least two untrapped ventilating openings into the drains, according to one of the following alternative arrangements:—(1) One opening consists of a shaft or disconnecting chamber opening at or near the ground level, and situated as close as possible to the trap specified above, but on the house side of it; the other opening is a pipe or shaft carried from a point as far distant as possible from the said trap (i.e. as near as possible to the head of the drain) vertically upwards in such manner and to such height (in no case less than 10 feet) as to prevent any escape of foul air into any building; but (2) if more convenient, the relative positions of these openings may be reversed, the shaft being placed near the trap, and the opening at the ground level at the head of the drain. The groundlevel opening must have a grating, with apertures equal in total area to the sectional area of the drain.

The pipe or shaft at the other end of the drain (whether used as a soil pipe or not) is required to have a sectional area equal to that of the drain, and in no case less than 4 inches; and bends and angles are

to be avoided as far as practicable.

- (k) Disconnection of drains from house. No drain inlet is permitted within a building except the inlet necessary for a water-closet. Every soil pipe must be at least 4 inches in diameter, must be placed outside the building, and must be continued upwards in full diameter without bends or angles (except where unavoidable), to such a height and such a point as to afford a safe outlet for sewer air. There must be no trap between the soil pipe and the drain to which it leads, nor in any part of the soil pipe except such as may be necessary in the construction of the watercloset. The waste pipe from a slop sink must conform to the same requirements as a soil pipe. The waste pipes from any other sink, bath or lavatory, the overflow pipe from any cistern and from any "safe" under a bath or water-closet, and every pipe for conveying waste water, must be taken through an external wall and must discharge in the open air over a channel leading to a trapped gulley grating at least 18 inches distant.
- (l) Water-closets must have a window opening directly into the external air, and measuring 2 feet by 1 foot clear of the frame; and, in addition to the window, adequate means of constant ventilation by air-bricks, air-shafts, etc. Such closets, if within the building, must adjoin an external wall. The water must be supplied to a water-closet by means of a special cistern. The apparatus must be suitable for effectual flushing and cleansing of the basin; the basin must be made of non-absorbent material, and of such shape and capacity as to receive and contain a sufficient quantity of water, and to allow all filth to

fall free of the sides directly into the water. "Containers" and "D-traps" are forbidden.

(m) Earth-closets are subject to the same conditions as water-closets so far as regards position, lighting, and ventilation. Proper arrangements must be made for the supply of dry earth, and its effectual and frequent application to the excreta; also for convenience of scavenging, and for exclusion of rainfall and drainage. The receptacle for excreta, whether fixed or movable, must be so constructed as to prevent absorption or escape of the contents, and to exclude rainfall and drainage; if fixed, its capacity must not be greater than may suffice for (three months?) nor in any case greater than (forty?) cubic feet, and it must in every part be (three?) inches above the ground. (The maximum limit of size may with advantage be reduced to two cubic feet in the case of earth-closets

placed inside houses.)

(n) Privies must not be erected within 6 feet of a dwelling, public building, or place of business, nor within (fifty?) feet of any water likely to be used for drinking, or domestic purposes, or for manufacturing drinks, nor otherwise in such a position as to entail danger of the pollution of such water. Privies must be built so as to admit of convenient scavenging, without carrying the contents through any dwelling, public building, or place of business. There must be an opening for ventilation at the top; the floor must be paved, and raised 6 inches above the ground in all parts, with a fall of half an inch per foot towards the door. The receptacle may be fixed or movable. If movable, as in pail-closets, the floor of the area beneath the seat must be flagged or asphalted, and raised 3 inches above the ground level, and all the sides of the said area must be made of flag, slate, or brick, at least 9 inches thick, and rendered in cement. If the receptacle is fixed, it must be in every part

3 inches above the ground level, and its capacity must not exceed 8 cubic feet; * means must be provided for the application of ashes and dry refuse to the excreta, and for the exclusion of rainfall and drainage, and for convenient access for scavenging; the materials and construction must be such as to prevent absorption or leakage, and there must be no

connection with any drain.

(o) Ashpits must not be constructed within 6 feet of any dwelling, public building, or place of business, nor within (50 ?) feet of any water likely to be used for drinking or domestic purposes, etc., nor otherwise in such a position as to entail danger of the pollution of such water. Ashpits must be so placed and constructed as conveniently to allow of scavenging without carrying the contents through any dwelling, public building, or place of business. The capacity must not exceed 6 cubic feet or such less capacity as may suffice for a period not exceeding one week. The walls must be of flag, slate, or brick, at least 9 inches thick, and rendered inside with cement; the floor must be flagged or asphalted, and raised at least 3 inches above the ground level. The ashpit must be roofed and ventilated, and provided with a door so arranged as to allow of convenient removal of the contents, and to allow also of being closed and fastened. The ashpit must not be connected with any drain.

(p) Cesspools must not be constructed within (50?) feet of any dwelling, public building, or place of business, nor within (100?) feet of any water likely to be used for drinking, or domestic purposes, or for manufacturing drinks, or otherwise in such a position as to entail danger of pollution of such water. Cesspools must be so constructed and placed as con-

^{*} These dimensions are based upon the assumption of weekly scavenging.

veniently to admit of scavenging and cleansing without carrying the contents through any dwelling, public building, or place of business. They must not be connected with any sewer. They must be covered over by an arch or otherwise, and adequately ventilated; and must be constructed of brick in cement, rendered inside with cement, and with a backing of at least 9 inches of clay.

(q) Closure of premises unfit for occupation.—The S.A. may under certificate from the M.O.H. or Surveyor, declare any building or part of a building erected after unfit for habitation, and order it to be closed until rendered fit for habitation. Opportunity must be given to the owner to show cause why

such order should not be made.

(r) Plans and sections must be submitted, showing in detail the construction of all proposed new streets or buildings. (The Public Health Act, s. 158, requires that the S.A. shall signify their approval or disapproval of the plans within a month after receiving them.)

(s) Inspection.—Notice must be given to the Surveyor of the dates upon which work is to be commenced, and upon which any sewer, drain, or foundation is to be covered up; notice must also be given of the completion of the work. Free access for inspection must be afforded to him at all times during the progress of the work.

(t) Demolition of illicit works.—If any work to which the byelaws apply is done in contravention of such byelaws, the S.A. are empowered to remove,

alter, or pull down such work.*

^{*} The Model Byelaws contain no provision for the sewerage of new streets, owing to the wide variation in the conditions in different localities. "It may be doubted whether any powers, which under such bye-laws may be lawfully assumed by Sanitary Authorities, will, as regards extent and efficacy, compare with the powers which they derive from the express provisions of the Public Health Act."—L.G.B. Gircular to Urban S.A.

5. Slaughter-houses.

(a) Licences.—Applications for licence of existing premises, or erection of new slaughter-houses, must be made upon a specified form, and must include full particulars as to the position, form, area, cubic space, etc., of the buildings and appendages; materials and construction of walls and floors; means of water supply, drainage, lighting, and ventilation; means of access for cattle; number, position, and size of stalls or lairs, and number of animals to be accommodated therein, distinguishing oxen, calves, sheep, and swine. The boundaries must also be shown, and, in the case of old premises, particulars as to the ownership and the applicant's tenure must be given.

(b) Registration.—If the S.A. approve the application, a licence shall be issued to the applicant, and must be registered by him at the office of the S.A.

(c) Inspection.—Free access to every slaughterhouse for the purpose of inspection must be afforded at all reasonable times to the M.O.H., Inspector, Surveyor, and Committees appointed by the S.A.

(d) Water must be supplied to every animal kept

in a lair prior to slaughter.

(e) Mode of slaughter.—Cattle must be secured by the head so as to be felled with as little pain as practicable.

(f) Drainage, water supply, and ventilation must

be kept in efficient order.

(g) Cleanliness.—The walls and floor must be kept in good order and repair, and must be thoroughly cleansed within three hours after any slaughtering; the walls and ceiling must be limewashed within the first ten days of March, June, September, and December respectively.

(h) Animals not to be kept.—No dog may be kept in a slaughter-house: nor other animal, unless intended for slaughter upon the premises, and then only

or preparing it for slaughter by fasting or otherwise.

(i) Removal of refuse.—Suitable vessels made of mon-absorbent materials, and provided with close-itting covers, must be provided for the reception of blood, manure, garbage, and other refuse; all such matters must be placed in these vessels immediately after the slaughtering; the refuse must be removed within 24 hours, and the vessels forthwith cleansed. All skins, fat, and offal must be removed within 24 hours.

"The Board have considered that the statutory terms do not warrant...... regulations directly affecting the structure of the premises. But.....in the exercise of the discretionary power of licensing which has been conferred upon the S.A., the following rules as to site and structure should influence their decision upon each application for a licence:—

"1. The premises should not be within 100 feet of any dwelling-house; and the site should be such as to admit of free eventilation by direct communication with the external air on

two sides at least of the slaughter-house.

"2. Lairs for cattle in connection with the slaughter-house should not be within 100 feet of a dwelling-house.

"3. The slaughter-house should not in any part be below

the surface of the ground.

"4. The approach to the slaughter-house should not be on an incline of more than one in four, and should not be through any dwelling-house or shop.

"5. No room or loft should be constructed over the

slaughter-house.

"6. The slaughter-house should be provided with an adequate tank or other proper receptacle for water, so placed that the bottom shall not be less than six feet above the level of the floor of the slaughter-house.

"7. The slaughter-house shall be provided with means of

thorough ventilation.

"8. The slaughter-house should be well paved with asphalte or concrete, and laid with proper slope and channel towards a gulley, which should be properly trapped and covered with a grating, the bars of which should not be more than three-eighths of an inch apart. Provision for the effectual drainage of the slaughter-house should also be made.

"9. The surface of the walls in the interior of the slaughterhouse should be covered with hard, smooth, impervious

material to a sufficient height.

"10. No water-closet, privy, or cesspool should be constructed within the slaughter-house. There should be no direct communication between the slaughter-house and any stable, water-closet, privy, or cesspool.

"11. Every lair for cattle in connection with the slaughter-

house should be properly paved, drained and ventilated.

"No habitable room should be constructed over any lair."

-L.G.B., Memorandum.

6. Houses let in lodgings (or "tenements")

-(a) Definitions.-A "lodging-house" is defined as a house or part of a house let in lodgings, or occupied by members of more than one family. The "landlord" is the person letting such lodging-house, or entitled to receive the profits of such letting; and a "lodger" is a person to whom any room or rooms in such house, or part of a house, have been let for his use and occupation. (b) Limitation as to class.— (1) On a rent limit, say of 3s. to 8s. if unfurnished, or 5s. to 10s. if furnished; or (2) where the landlord resides on the premises, and the letting is to not more than one lodger. (c) Cubic space.—A room used exclusively as a sleeping-apartment shall not be so used at any one time by a greater number of persons than will admit of the provision of 300 cubic feet of air-space for each person over ten years of age, and 150 cubic feet for each person under ten years. A room not exclusively used as a sleepingroom shall not be used at any one time by a greater number of persons than will admit of the provision of 400 cubic feet of air-space for each person over ten years of age, and 200 cubic feet for each person under ten years. (d) Landlord to furnish returns .-The landlord shall, if required by the S.A., furnish the following particulars:—(1) Number of rooms in the house; (2) number of rooms let in lodg-

ings or occupied by members of more than one family; (3) the manner of use of each room; (4) the number, age, and sex of the occupants of each room used for sleeping; (5) full name of the lessee of each room; (6) the rent payable by each lessee. (e) Inspection.—Facilities for inspection shall be afforded tto the M.O.H., Inspector, or Surveyor at all times by the landlord and lodgers. (f) Closet accom-*modation.—The landlord shall provide sufficient closet saccommodation, in the proportion of at least one closet for every twelve persons of the maximum number of lodgers allowed by the byelaws, and maintain the closets and all necessary appurtenances in efficient order. All closets must be kept clean, and supplied with dry earth (if earth-closets) or water (if water-closets) as required.* (g) Ashpits must be maintained in repair by the landlord, and must be kept clean; no filth or wet refuse must be thrown into an ashpit designed for dry refuse only. (h) Cleanliness of rooms.—The floors of all rooms must be swept daily, and washed every week. All windows, all wood, stone, or metal fixtures, and all painted surfaces must be kept clean. All solid or liquid filth or refuse must be removed from every room every day, and all receptacles thereof cleansed. (i) Keeping of animals. -No animal must be kept upon the premises in such manner as to render the premises filthy. (j) Water storage.—All cisterns for storing water must be kept clean. (k) Ventilation.—The landlord must maintain in efficient order the means of ventilation of all parts of the house. (1) Annual cleansing of premises.— Every year, in the first week of the month of the landlord must cleanse every part of the premises, and whitewash the ceilings and walls throughout the

^{*} If the byelaws respecting new streets and buildings are not in force in the district, those of them which refer to the construction of closets may be introduced here.

house, except surfaces to which limewashing may be unsuitable, and these must be cleansed and, if necessary, painted. (m) Yard and open spaces. - The yard and all open spaces must be kept clean and in good order. The yard must have a hard, impervious pavement laid upon concrete, and sloped to a channel leading to a trapped gulley grating; all these must be kept in repair by the landlord. (n) Ventilation of sleeping-rooms.— The window of every sleeping-room shall be kept open for an hour in the forenoon and an hour in the afternoon, except in case of stress of weather, or occupation of the room by a sick person, or other sufficient reason. (o) Infectious diseases.—The landlord must immediately give written notice to the M.O.H. if he learns that any person in the house is ill of an infectious disease. A lodger must immediately give written notice to the M.O.H., and verbal or written notice to every lodger in the house, if he has reason to believe that any occupant of any room is ill of an infectious disease. Where a justice's order has been obtained for the removal to a hospital of a person suffering from any dangerous infectious disease, both the landlord and the lodger in whose tenement such sick person is shall take such steps as may be requisite on their respective parts for the safe and prompt removal of such person; and shall adopt all such precautions as may be most suitable, in accordance with any instructions of the M.O.H.

Where a duty imposed upon or a prohibition applied to the landlord of a lodging-house is by any byelaw expressly imposed or applied subject to the provisions of these byelaws, proceedings shall not be taken against him for an offence against the byelaw unless and until the landlord, after service upon him of a notice in writing from the Council, signed by the M.O.H., or clerk of the authority, requiring him within a reasonable time specified in the notice to

comply with the byelaw, has failed to comply with the byelaw within that time.

The landlord is made responsible for the structural repair of every part of the premises, for the efficiency of the closets and of the means of ventilation, for the annual cleansing, and for the detailed cleansing of all parts (including yards, closets, staircases, and passages) which are not let for the sole use of one tenant. The tenant is responsible for the cleanliness and management of the rooms rented by him, and of all other parts of the premises of which he has the sole use.

7. Cemeteries.—(a) Definitions.—A "grave" is defined as a burial-place formed in the ground by excavation, and without any internal wall of brickwork or stonework, or any other artificial lining. A "" vault" is an underground burial-place of any other construction. (b) Vaults.—Every vault shall be enclosed with walls of brick or stone, solidly put together with good mortar or cement. (c) Common graves.— Not more than one body shall be buried at any one time in a grave in respect of which no exclusive right of burial has been granted. (Exception is made in the case of two or more members of the same family.) Such a grave shall not be reopened for the purpose of a further burial within eight years after the burial of a person aged less than twelve years, nor within fourteen years after the burial of a person aged more than twelve years. (Exception is made in the case of members of the same family.) (d) Minimum covering of earth .- No part of a coffin shall be buried at a less depth than three feet below the ground adjoining the grave, if it contains the body of a person aged less than twelve years; nor at a less depth than four feet if the age of the deceased was over twelve years. A layer of earth, not less than one foot in thickness, shall be interposed between every coffin and the coffin nearest to it. (e) Closure of vaults.—A coffin buried

in a vault shall, within twenty-four hours after burial, be wholly and permanently embedded in and covered with good cement concrete, not less in any part than 6 inches in thickness; or wholly and permanently enclosed in a separate cell, constructed of slate or flag, not less than two inches thick, and jointed in cement, or of brick in cement, and in such manner as to prevent as far as practicable the escape of noxious gas.

8. Mortuaries.—The only provisions of sanitary importance suggested in this series are the following: A body deposited in the mortuary shall be removed therefrom for interment within......days after death; but if the deceased has died of an infectious disease, the body shall be removed for interment within......days after death.

An L.G.B. memorandum makes the following general suggestions in regard to construction and

management :-

The buildings should be isolated and unobtrusive, but substantial, structures of brick or stone. Every chamber for the reception of corpses should be on the ground floor. In addition to such chamber there should be a waiting-room, a caretaker's house, and a shed or outhouse. Every mortuary chamber should be lofty, and there should be a ceiling or a double roof, with an intervening space of eight inches, for the sake of coolness. The area should be sufficient to allow freedom of movement between the slabs. windows should be on the north side, if practicable; if otherwise, they should have external louvre blinds. Louvres, or air gratings, under the eaves will be the best means of ventilation. The pavement must be even and close, and a cement floor is preferable. The slabs should be of slate, and 21 ft. to 3 ft. from the floor. Water should be laid on within the chamber. The walls and ceiling should be whitewashed,

and the outside of the roof also whitened. There should be at least two chambers, one of which may be reserved for bodies of persons who have died of infectious disease.

There should be a resident caretaker, and bodies should be received at any hour of the day or night.

A register should be kept.

9. Offensive trades.—The offensive trades in regard to which Model Byelaws have been issued by the L.G.B. are those of—

Blood-boiler.
Bone-boiler.
Fellmonger.
Soap-boiler.
Tallow-melter.
Tripe-boiler.

Blood-drier.
Leather-dresser.
Tanner.
Fat-melter or Extractor.
Glue-maker.
Size-maker.
Gut-scraper.

Of these the first six only are specifically mentioned in s. 112 of the Public Health Act, the rest being examples of "other offensive trades."* The same general provisions appear in all, with numerous additions or variations as required by the conditions of the particular trade in question.

The following summary will show their general

character:

(i.) Storage of offensive materials.—Materials not required for immediate use or treatment shall be

^{*} A blood-boiler boils blood or treats it with steam and prepares it for manure manufacture by reducing it to powder in heated "driers." A bone-boiler extracts fat and gelatine from bones by boiling under pressure. The fellmonger prepares dry or fresh skins for leather-dressing. Fresh skins are beaten with a mallet, then soaked in water and milk, of lime, after which the hair or wool is plucked off. Fat, tallow, oil; etc., together with alkali solutions (K or Na) termed "lyes," are boiled by the soap-boiler preparatory to soap manufacture. Tallow-melting is a similar process of treating rendered fat without alkalies. The liquid is run off into barrels. A tripe-boiler cleans and boils the stomach of the ox or sheep, and prepares it for human food. Other offensive trades include such processes as involve the collection or treatment of large quantities of animal matter or substances likely to become a nuisance.

so stored as to prevent effluvium. (ii.) Offensive vapours emitted during boiling, melting, etc. The best practicable means shall be adopted for rendering such vapours innocuous. The vapour shall either (a) be discharged into the external air in such a manner and at such a height as to admit of its diffusion without noxious or injurious effects, or (b) shall be passed directly from the pan, etc., through a fire, or (c) into a condensing apparatus, or (d) through a condensing apparatus and then through a fire, in such manner as effectually to consume the vapour or deprive it of all noxious or injurious properties. (iii.) Drainage must be kept in efficient order. Boneboilers are required to cool all hot liquid refuse before passing it into any drain. (iv.) Floors must be kept in good order so as to prevent the absorption of filth. In respect of several trades it is required that at the close of every working day the floors shall be swept (e.g. tanner's, leather-dresser's) or washed (blooddrier's, etc.) or scraped, or otherwise cleansed. All refuse collected by sweeping or scraping shall be forthwith removed from the premises in covered receptacles, unless intended to be forthwith subjected to further trade processes on the premises. (v.) Walls shall be kept in good order so as to prevent the absorption of filth, and, if necessary, be scraped. Limewashing of walls and ceiling twice yearly is required in regard to certain trades-fellmonger's, for example. (vi.) Apparatus, including all implements and vessels, to be kept clean; daily cleansing is required in some cases. (vii.) Waste lime (fellmonger's, tanner's) must be removed at once, and under close cover. (viii.) Tanks used by fell-mongers for washing or soaking skins shall be emptied and cleansed as often as necessary to prevent effluvia. (ix.) Free access for the purpose of inspection shall at all reasonable times be afforded to the M.O.H.,

Inspector of Nuisances, Surveyor, or any committee

specially appointed by the S.A. on that behalf.

10. Hop-pickers.—(a) Application.—These byelaws apply to all tents, sheds, barns, or other places occupied as temporary dwellings by hop-pickers, but not to places inhabited throughout the year. (b) Habitable condition .- Such temporary habitations are required to be clean, dry, weatherproof, and to be ventilated and lighted. (c) Closet accommodation. There must be adequate privy accommodation for the separate use of each sex. (d) Water.—There must be a sufficient supply of good water for drinking, cooking, and washing. (e) Cooking.—There must be a separate cooking-place for every fifteen persons authorised to be received. (f) Area.—An average floor space of 15 square feet must be allowed for each occupant, but two children under ten years of age may be counted as one adult. (g) Bedding.—Clean straw or other suitable material must be supplied for bedding, and renewed as required. The beds must be screened off in places occupied by adults of different sexes. (h) Cleansing.—The premises and appointments must be thoroughly cleansed immediately before occupation, the internal surfaces limewashed, and all offensive accumulations cleared away. The cleansing must be repeated as required from time to time during the period of occupation.

(Similar byelaws may be made in respect of the accommodation of persons engaged in picking fruit or

vegetables.)

11. Means of escape from fire, in factories and workshops.—Buildings of more than one storey must, if any part above the ground floor is used as a factory or workshop, have adequate fixed stairs (with handrails), lighted, and giving free access from every such part to the ground floor, and where practicable to an open space outside. In new

factories or workshops, buildings, and buildings newly altered structurally for that purpose, the stairs must be of fire-resisting material, e.g. brick, stone, concrete, iron, steel, copper, hard timber (oak, teak),

or timber rendered uninflammable.

If any floor over thirty feet from the ground is used as a factory or workshop, and either more than ten persons are employed, or readily inflammable materials are stored, there must, in addition, be (1) an external fire-resisting staircase, or (2) an efficient fire escape, or (3) ready access to the roof, and where practicable to the roof of an adjoining building. The window or door leading to (1), (2), or (3) must be distinctively marked.

Every room (of a factory or workshop) in which persons are habitually employed must have at least one window (distinctively marked if there be others) opening easily at the sill level to an extent allowing

easy escape.

CHAPTER XXI

VITAL STATISTICS

Population.—A census has been taken decennially since 1801, and gives detailed information regarding each county, union, registration district and subdistrict, and urban district. The chief data are the total number of inhabitants in each area, the numbers living of each sex and at certain age-periods, and the numbers employed in certain callings.

The official estimate of population for any year is made by assuming that increase or decrease has gone on since the last census at exactly the same rate as between that and the previous census. This assumption is arbitrary, and seldom accords with the facts as

ascertained by the next census.

If a population of 100,000 in a given year becomes 101,000 in the next year, the estimate for the third

year is 102,010, not 102,000.

If p = the population in any given year, and r = the factor of annual increase (in the above example r = 1.01), then in one year the population will become

 $p \times r$, in two years $p \times r^2$, and in n years $p \times r^n$:

The census is taken, not at the middle of the year, but at the end of the first quarter. Given a census population of x in 1891, and y in 1901, the rate of decennial increase is $\frac{y}{x}$, that of annual increase $\sqrt[10]{\frac{y}{x}}$, and that of

quarterly increase
$$\sqrt[40]{\frac{y}{x}}$$
. Hence—

 $\log y - \log x = \log$ rate of decennial increase, and $\frac{\log y - \log x}{10} = \log$ rate of annual increase,

and $\frac{\log y - \log x}{10 \times 4} = \log$. rate of quarterly increase.

Thus the logarithm of the estimated mid-year population for 1901 is, $\log y + \frac{\log y - \log x}{10 \times 4}$; and for 1908,

$$\log_{10} y + \frac{\log_{10} y - \log_{10} x}{10 \times 4} + 7 \frac{\log_{10} y - \log_{10} x}{10}$$

The following calculation of the mid-year 1905 population from census data of 1891 and 1901 may serve as an example :—

(a) The census population of 1901 was 100,970; log. 100,970 =	5.0041924
(b) The census population of 1891 was 87,842; log. 87,842 =	4.9437022
(c) By subtraction, the logarithm of decennial increase was, therefore	0 0004302
is obtained	
(e) Again dividing by 4, the log. of quarterly increase is obtained =	0.0015122
(f) Adding together log. 1901 census population (a), and log. quarterly increase (b), and four times log. annual increase (d), we get the log of 1905 mid-year population.	5.0299017

And on referring to the tables, the number corresponding to this logarithm is found to be 107,128.

There are obvious limitations to the value of such an estimate. The rate of increase is not the same in the centre of a town as in the growing suburbs. New houses are springing up and becoming inhabited in the latter, apart from the "natural increase" by excess of births over deaths; while in the former there may even be an actual decrease of population, owing to the displacement of dwellings by business premises.

If the estimate is very wide of the mark, some indication of this may be given by an apparently excessively high or low birth-rate and death-rate; by the number of inhabited houses,* as ascertained from the rate-books or other sources; by returns

^{*} The average number of persons per inhabited house is fairly constant for each locality, though varying considerably in different towns, according to the proportion of "tenement houses," and houses occupied by one family.

of those who by migration have evaded vaccination; by returns of school attendance; or by inference from the condition of the local industries. In villages, where the local conditions are well known, a better estimate can sometimes be made from such data than from calculations based upon a remote census.

The actual increase is dependent upon the balance between births and immigration on the one hand, and deaths and emigration on the other. The excess of births over deaths is termed the "natural increase." As regards towns, the actual is, upon the average, greater than the natural increase, since there is a tendency to migration from rural to urban districts. In times of depression of local trade there is emigration to other districts; whereas, when trade is flourishing, labour is attracted from rural districts and from other towns.

The age-distribution of a population is ascertained by each census, and may be assumed to remain constant until the next census; that is, the rate of increase is assumed to be uniform at all ages. There are. however, very material differences among towns, and between town and country, in this respect, and as the tendency to death is much greater among the very young and very old, it becomes important to allow for these variations in drawing inferences from death-rates as to the health conditions of a community. Owing to the immigration of young adults from rural districts, town populations almost always contain a larger proportion of persons between 15 and 45 than is found in the country at large; and hence a higher birth-rate and (under equal health conditions) a lower death-rate are to be expected a priori.

Births must, according to the Births and Deaths Registration Act of 1874, be registered within 42

days.

More males than females are born, in the proportion of nearly 104 to 100, but this excess is diminishing. Owing to the higher death-rate among males, the survivors of the two sexes are nearly equal in number at the end of the first year of life, and thenceforward the females are in the majority.

Still-births, which Farr estimated at 4 per cent. of the total births, are not included in the returns.

The Birth-rate is the proportion of births in one

year per 1,000 persons living; that is-

No. of births in the year mid-year population × 1,000.

For shorter periods, such as quarters or weeks, the rate is calculated on the assumption that the like periods throughout the year would yield the same proportion: so that the birth-rate during $\frac{1}{x}$ th part of a year is—

No. of births during the period in question $x \times x \times 1,000$. mid-year population

For ordinary purposes it is sufficient to regard the year as consisting of 365 days, or 52 weeks; but if greater precision is required, the astronomical year may be used—namely, 365.24 days, or 52.177 weeks.

The English birth-rate, which was 35.4 in 1871-80, and 36.3 in 1876, declined to 27.2 in 1905. It is higher in towns, and in times of prosperity; lower in rural districts, and in times of depression of trade.

A sounder basis of population for the purpose is that of women at ages 15-45; or, still better, married women at those ages (with exclusion of illegitimate births, the proportion of which is variable and decreasing, and averages about 4 per cent. of the total). On this latter basis the rate per 1,000 was 293 in 1870-2, 286 in 1880-2, 264 in 1890-2, and 236 in 1900-2. Even this fails to take account of material differences in fertility at the several ages within the limits stated.

Deaths must be registered within five days.

Death-rates are calculated in the same way as birth-rates. The death-rate in England and Wales declined from 22.4 in 1851–70 to 18.2 in 1891–1900, and 15.2 in 1905, when the annual total of deaths was 520,031. Local death-rates require certain

corrections, as below, in order to render them com-

parable with that of the whole country.

Correction for non-residents.—The deaths of strangers who happen to die in the district should be excluded. Without this correction the true rate may be seriously overstated if the district contains large institutions admitting non-residents. Similarly, deaths of residents dying outside the district should be included.

Correction for age and sex distribution is made by a "factor" obtained as below, which is greater than unity in nearly all towns, and less than unity in rural districts. Multiplying the local death-rate by the factor, the corrected local death-rate is obtained. The comparative mortality figure for the district is—

 $\frac{\text{corrected local death-rate}}{\text{death-rate for the whole country}} \times 1,000.$

A standard death-rate for a locality is the death-rate for the whole country, re-calculated on the basis of the local age and sex distribution, and may be found as follows. The age and sex distribution per million of both local and general population are separately determined from census data as at page 632. The recorded deaths (in the whole country) are distributed in the same groups, and the number of deaths in each group corrected in the proportion of the local to the general population at those ages. If, for example, the males living at a given group of ages are x per million of the local population and y per million of that of the whole country, and the male deaths at those ages are m, the corrected number of deaths is $m \times \frac{x}{y}$. Adding together all the corrected numbers, we have a corrected total a, differing from the recorded total b (of deaths in the whole country), and representing the deaths which would have occurred had the age and sex distribution been the same as that of the locality

Hence $\frac{a \times 1000}{\text{population of whole country}}$ is the standard death-rate

for the locality. The death-rate of whole country is the standard local death-rate

factor sought, and is equal to $\frac{b}{a}$.

Urban death-rates are higher than rural, and the correction for age and sex distribution increases the inequality. The causes of this are complex, and include differences of occupation and mode of life, and the direct and indirect consequences of aggregation of population. Ogle found that in 1871-80 the death-rates of districts increased with the density of population, if exceeding 400 per square mile, but below that point the influence was not apparent. (The average density for England and Wales exceeds 500 per square mile, or 0.8 per acre.) Farr concluded that, other things being equal, urban death-rates vary as the twelfth root of the density; but this rule is of limited practical application, since many variable conditions are involved, and notably the degree of crowding of rooms. Newsholme showed that in 1889 the death-rate among the 20,000 inhabitants of Peabody Buildings (density, 750 per acre), was lower than that of London generally (average density, 49 per acre).

The death-rate is diminishing in both urban and rural districts, but more rapidly in the former than

in the latter.

The death-rate among males is higher than among females, subject, however, to exceptions at certain age-periods, notably 10-20, as Appendix II. shows. In the decennium 1891-1900, the former was 19.3 and the latter 17.1. Both are decreasing, but the saving of life is chiefly effected in the earlier age-periods, and is inconspicuous at ages over 40, owing, perhaps, partly to the increasing wear and tear of modern life, and partly to the more frequent survival of weakly persons at middle life who under former conditions would have died in youth.

In both sexes the tendency to death is high in infancy, reaches its minimum in the 10 to 15 year age-period, and afterwards increases steadily through-

out life.

Infant mortality is calculated upon the birthreturns, which are precise and free from many of the errors attendant upon estimates. The formula is:—

 $\frac{\text{deaths of children under 1 year}}{\text{registered births}} \times 1000.$

Influence of the birth-rate upon the death-rate.—Other things being equal, a high birth-rate in a given year would theoretically increase the death-rate, since the mortality among the newly born is much greater than that of the population generally. But a high birth-rate implies (a) a large proportion of persons at the reproductive ages, and in a few years (b) a new population of children and young adults, since under the most adverse conditions a large majority of the infants survive. The mortality among both classes (a and b) is low, and more than balances the higher mortality in the first year or two of life. (Farr.)

Cause of death.—The death-returns obtained from the registrars are copied from the certificates signed by medical practitioners or coroners. Very commonly two or more causes are assigned—one being regarded as "primary" and the other as "secondary." As only one cause in each case can be accepted for statistical purposes, preference should be given to (1) infective or specific diseases, and (2) primary as compared with secondary causes.

Progressive changes are noticeable, in regard to the deathrate from certain well-defined diseases, in the direction of increase or decrease. It may extend to all ages, or be more or less limited to certain periods of life. The change does not necessarily affect both sexes alike, phthisis, for example, having decreased much more rapidly among females than

among males.

Decrease is manifest in the mortality from small-pox, scarlet-fever, diarrhoea, typhus, enteric fever, simple continued fever, thrush, phthisis, convulsions, croup, diseases of the digestive system, and diseases of parturition; also from accident, negligence, homicide, and "ill-defined causes." As regards the first five, the reduction is real, and attributable in great part to improved measures of precaution and sanitation. Phthisis has been greatly lessened, but there has

probably been some transfer of cases by improved diagnosis from phthisis to other respiratory diseases. Better diagnosis and better certification are responsible for the gradual disappearance of "croup," "convulsions," and "simple continued fever," and also, of course, for the lessened total of "ill-defined" diseases; nearly all these are cases of transfer to other headings, and not true reductions in mortality.

On the other hand, there is a distinct increase in the mortality attributed to cancer, diabetes, suicide, and diseases of the nervous, circulatory, respiratory, and urinary systems. It is probable that some real increase has occurred in respect of nervous and circulatory diseases, and perhaps also urinary diseases, but improvement in diagnosis has also to be taken

into account.

The cancer death-rate in 1861-70 was 387 per million, in 1871-80 473, in 1881-90 589, in 1891-1900 758; and fairly well defined areas yielded exceptionally high rates. Age-distribution of population is important in this connection, since the mortality is chiefly at ages over 35. The increase is partly due to improved diagnosis, and to inclusion of forms of malignant disease not previously grouped with cancer. Recent researches in this country, while showing a wider difference among lower animals than had been suspected (Bashford), have not confirmed the suggestion that cancer is an infective disease due to micro-organisms. Instances are met with of repeated occurrence of the disease in "cancer houses," but little is known of dependence upon dietetic or other causes, and no special preventive measures are in force.

Zymotic death-rate is a term formerly applied to the aggregate death-rate from the "seven principal zymotic diseases" — small-pox, measles, scarlet-fever, diphtheria, whooping cough, "fever" (typhus, simple continued, and enteric), and diarrhoea. It is liable to great fluctuation, according to the epidemic prevalence of one or other of the

diseases included. (See p. 614).

Deaths due to protracted sequelæ of acute specific diseases are usually credited to the secondary and not to the true primary cause. It will be found that the mortality from bronchitis and pneumonia is increased by outbreaks of influenza, whooping cough and measles, and scarlet-fever prevalence is followed by deaths referred merely to kidney diseases.

Sex has an important bearing upon the mortality from certain diseases, owing partly to differences in constitutional predisposition, and partly to exposure to

the conditions favourable or unfavourable to the development of the diseases in question. The male mortality exceeds the female in most well defined diseases or groups of diseases, but there are important exceptions, such as diphtheria, whooping cough, erysipelas, rheumatism, cancer, and anæmia

(including chlorosis and pernicious anæmia).

Age too materially affects the average liability to certain diseases, by reason of physiological changes, acquired protection, and degree of exposure. Vaccination completely alters the age-curve in smallpox. In all protective diseases the accumulation of protected survivors must be allowed for before any lessened average susceptibility at higher ages can be inferred. The maximum mortality from small-pox, whooping cough, erysipelas, and diarrhea occurs in the first year of life, from measles in the second, from scarletfever in the third, and from diphtheria in the fourth. A second maximum follows in small-pox about the 25th year; and from about the same point diarrhea mortality again rises until the end of life. Phthisis is at its minimum from the 5th to the 10th year, increases up to the 45th, and afterwards diminishes. Cancer mortality is small in the earlier years, but increases rapidly from about the 25th year to the end of life. The single curve, with a minimum from the 10th to the 15th year, which is characteristic of the total deathrate, is common also to the mortality from nervous, respiratory, digestive, and urinary diseases, and violence; diseases of the circulatory system increase steadily in mortality from birth to the end of life. varying influence of age is seen even in infancy.

The true influence of sex and age cannot be ascertained by returns of mortality alone. The mere number of deaths at each age-period conveys little information, since the numbers living diminish as age advances. Even death-rates for each group tell only of the liability to death, but nothing as to the incidence

of sickness or the chance of recovery. The variations in incidence and in severity, according to age and sex, are not always parallel, and indeed are often inverse, as in scarletfever and enteric fever. From the number of persons living, of each sex and at each age-period, it is possible to calculate, for each sub-group, (1) the incidence of disease, that is, the proportion of attacks to population, and (2) the case-mortality, or proportion of deaths to attacks. Evidence as to incidence of disease at different ages can be obtained from notification returns, but hospital statistics are misleading, since age materially affects the chances of removal to hospital. As regards case-mortality, however, both hospital and notification data may be admitted.

Statistical evidence of the health of communities.—The usual criteria are either actual death-rates (general, "zymotic," or infant mortality) or figures representing the average longevity-namely, "expectation of life," "probable duration of life,"

"mean age at death."

The corrected death-rate affords a simple and, in the main, accurate measure of the comparative prevalence of disease. It is untrustworthy if the figures are so small as to be exposed to violent fluctuations; thus, the "weekly death rate" is only useful in large communities. Other sources of error are (i) uncertainty as to population, and (ii) severe epidemics, which may have no known relation to

impaired public health in general.

A high death-rate from enteric fever, diphtheria, or diarrhea may in general fairly be taken to imply a defective sanitary state; but may also be due to temporary and accidental causes, such as climatic conditions or accidental contamination of milk or water. Little is known of the determining causes of epidemics of small-pox, measles, whooping cough, and scarlet-fever; but their predisposing causes are all widely different, and are for the most part not affected by what are known as "sanitary conditions." The death-rate due to such a heterogeneous group

as that of the "principal zymotic diseases" (p. 612) denotes simply the presence or absence of grave epidemics, and connotes nothing as to the health

condition of the community in other respects.

Infant mortality has not shared in the decline common to other death-rates. It is one-third higher in urban districts than in rural, higher among illegitimate children, and higher in the first trimester of the first year than in other trimesters. About one-fifth of the total deaths of infants occurs in the first week of life. The three chief diseases responsible for high infant mortality are (a) immaturity in all its forms, (b) diarrhœa, and (c) lung disease. The first is due to prematurity, debility, and congenital conditions. The second causes death about the middle of the first year, and is due to food causes; and the lung disease is brought about chiefly by exposure to cold and unequable temperatures.

The phthisis death-rate, if excessive, indicates dampness of soil, unhealthy occupations, or overcrowding of tenements. The death-rate from respiratory diseases

(other than phthisis) is also important.

The mean age at death is obtained by adding up the ages and dividing by the number of deaths. It is controlled by the birth-rate, and by the age-distribution of the population. A large proportion of infant deaths must necessarily reduce the average age at death.

The probable duration of life is the age at which exactly half of any given number of children born will have died. It can only be ascertained from a life-table, and is of no great value. The probable duration of life for males is about 53, and for females 58 years.

The expectation of life, at any age, is the average number of years which a person at that age will live, as shown by a life-table. The expectation of life at age 0-i.e. at birth-is also known as the mean

duration of life.* At other ages than the time of birth it is sometimes termed mean after-lifetime, and the present age plus the mean after-lifetime is the age to which a person may expect to survive.

The expectation of life is the true measure of the vitality of a community. The expectation at birth is the most convenient for comparative purposes, but if necessary we can eliminate the influence of infant mortality by taking the expectation at a later age.

A life-table shows how many, out of a million persons supposed to be born simultaneously, will survive at the end of each year or each term of years. The data required are (1) a census population—that is, a population of which the distribution according to ages and sexes is known; (2) returns of deaths (grouped for each sex in the same age-periods as have been adopted for the census population) for one or more years among this same population. The simplest plan is to take only the deaths in the census year, when the population is known with precision; but as it is important to obtain large numbers, it is convenient to use the death returns of a series of three or more years, in which the census year is central. Thus the 1901 census population may fairly be assumed to be substantially the same as the average population in the years 1900-1-2. A still better method is to take the death returns for a whole inter-censal period, and the mean population. A separate table should be constructed for each sex.

Having decided upon the intervals to be taken (annual or quinquennial), the first step is to ascertain from the census returns the number living at each

The "mean duration of life" differs from the "probable duration of life" just as the arithmetical mean of a list of numbers differs from the middle value of the series. The fact that one term has as many terms above as below it does not render it the mean of the series.

The "mean duration of life" must also be carefully distinguished from the "mean age at death." The latter expression is not employed in reference to a life-table population.

group of ages, and, from the death returns, the mean annual number of deaths among each group. As already explained, $\frac{\text{number of deaths}}{\text{number living}} \times 1,000 = D = \text{the}$ death-rate for that group of ages per thousand living. These D deaths (among 1,000 living) are assumed to be evenly distributed over the whole age-period included in the group, so that half $(i.e. \frac{D}{2})$ will occur in the earlier half of the ages, and half in the later.* Hence the 1,000 persons are regarded as decreasing from $1,000 + \frac{D}{2}$ at the beginning of the period to $1,000 - \frac{D}{2}$ at the end of it; and the ratio of the final to the initial population is $\left(1,000 - \frac{D}{2}\right): 1,000 + \frac{D}{2}$ $\frac{2,000-D}{2,000+D}$. Having found the value of this ratio for each age or group of ages, we have all the data necessary for the construction of a life-table.

For the first year of life the formula is not required. If the infant mortality (deaths under 1 year of age per 1,000 births) be 200, it is evident that 1,000,000 persons at birth (i.e. living at the commencement of the year in question) will be reduced to 800,000 during the year. The number commencing the second year is 800,000; taking the death-rate during that year of age as 65 per thousand living, the formula becomes $\frac{2,000-65}{2,000+65}$, or $\frac{1,935}{2,065}$, and, as already explained the 800,000 living at the beginning of the second year will be reduced to $\frac{800,000}{1} \times \frac{1,935}{2,065}$, or 749,637 at its

^{*} In other words, "the number of the living in any year of their age is an arithmetical mean proportional between the numbers that annually enter upon and that annually complete that year" (Milne, quoted by Farr). An earlier hypothesis (De Moivre's) assumed that the numbers living decreased in arithmetical progression down to nothing at the age of eighty-six years. U*

close. In like manner are determined the survivors at the end of the third, fourth, and fifth years. After the fifth year it is usual to proceed by 5-yearly periods, but the method is substantially the same. Suppose, for instance, that we have already found that of the million only 700,000 survive at the end of the fifth year; and that the mean annual death-rate among persons aged five to ten years is 6.5. This death-rate is assumed to be true of each of the five years, hence the formula of reduction for each year is $\frac{2,000-6.5}{2,000+6.5}$, and for the five years $(\frac{2,000-6.5}{2,000+6.5})^5$. At the end of the tenth year, therefore, the 700,000 will have become reduced to $700,000 \times (\frac{2,000-6.5}{2,000+6.5})^5$ —which, by the aid of logarithms, is readily found to be 677.615.

The process is repeated for each quinquennium

until no more survivors are left.

If decennial be substituted for quinquennial periods, the annual death-rate D must be calculated for the ten-year age period as a whole, and the formula will be $\left(\frac{2000-D}{2000+D}\right)^{10}$.

A life-table traces the history of a hypothetical generation which lives through each period of its life under the conditions that attend each such period during the era under examination. It eliminates the disturbing influence of a variable birth-rate and migration-rate, which introduce many fallacies into any attempted estimation of longevity from mere death-rates in a changing population. A census return gives the numbers living at ages between certain fixed points; a life-table, on the other hand, shows the relative numbers surviving at certain points of time, or rather of age, and thus enables us to calculate the expectation of life at any such point.

The expectation of life at any age A is calculated by adding together the years of life lived through by the whole of the life-table population after that age, and dividing by the number of survivors at age A. Thus to find the expectation of life at fifty years of age in males according to the 1891-1900 life-table (p. 634), the first step is to add together the numbers surviving at each later age: this gives the number of complete quinquennia lived through = 1,745,421. But in addition to the quinquennia which he completes, each of the 530,888 males surviving at age 50 lives through some portion of that quinquennium in which he dies. By a further application of Milne's principle this fraction is averaged as half a quinquennium, and hence we have to add 265,444 half-quinquennia to the 1,745,421 quinquennia. We thus get a total of 2,010,865 quinquennia or 10,054,325 years of future life; and this, divided among the 530,888 who are alive at age 50, gives the expectation of life as 18.9 years.

The expectation of life is somewhat greater among females than among males, at all ages. It is greatest

in both sexes at about three years of age.

Seasonal variations. - Most diseases, and well-defined groups of diseases, have a fairly constant relation to season, and present characteristic curves of average seasonal mortality, subject, however, to variation in time and range in any given year. Many of them have an obvious relation to heat and cold; thus there is a winter maximum in deaths from apoplexy and diseases of lungs, heart, and kidneys; and a summer maximum as regards diarrhea, tabes mesenterica, and thrush. Scarlet-fever, enteric fever, diphtheria, puerperal fever, erysipelas, and rheumatism have their maximum about November; small-pox and whooping cough have a spring maximum; and so, too, have phthisis, "convulsions," gout, and laryngitis. Measles has a double seasonal curve, with two maxima and two minima.

	Mortali	TY.
Cause of Death.	Greatest.	Least.
ALL CAUSES	(1) Dec. to March. (2) July to August. January to May. (1) December. (2) June. October. Nov., December. March, April. November. January. Nov., December. November. July (4th week), Aug. (1st and 2nd weeks). November. July. April. March, April. Dec. to March. March. Nov. to January. March. January. Jan. and March. December. Nov. to April. January. Jan. and March. December. Nov. to April. January.	(1) June. (2) October. Sept. and Oct. (1) September. (2) February. March to May. May, June. Sept., Oct. June. September. May to Aug. August. June, July. Dec. to May. August. Dec. to Mar. September. September. September. July, August. September. August. June to Aug. June to Oct.

There are also characteristic seasonal curves of mortality from all causes at certain ages :—

A man	MORTALITY.								
Ages.	Greatest.	Least.							
0-1 { 1-5 { 5-20 Over 20	4th week in July to 2nd. week in Aug. (1) Feb. to April (2) July, August December December	April, May, June June, July September . July, August August							

The mortality at ages from 5 to 20 shows only a slight excess at mid-winter and a slight falling off




Fig. 24.—Seasonal Curves of Mortality (London). Mean of 40 Years, except Diphtheria (20 years), and Enteric Fever (12 years).

Each division corresponds to 10 per cent. above or below the mean annual mortality indicated by the dotted line.

at mid-summer. The position of the maximum and minimum points remains the same at all later ages, but a rapidly increasing sensitiveness to cold is manifested by a wider range of seasonal curve. Below five years of age a similar increase in sensitiveness to climatic conditions is found; but the greatest danger is from heat, not from cold. From one to five years the principal maximum occurs in the early spring, owing to seasonal prevalence of whooping cough and measles; but there is also a distinct rise in July and August, due to diarrheal diseases. Under one year of age this minor elevation becomes an enormous peak, corresponding to the diarrhea maximum.

Occupation and mortality. — In tracing, from mortality data, the influence of occupation upon health, it is necessary to take into account not only the numbers employed, but also their sex and age-distribution, and the inevitable selection of the more robust for certain kinds of employment, and

the weaker for others.

Comparing the results of the 1891 census with the mean mortality of the three years 1890–1–2, Tatham has shown that among the whole male population aged 25–65 the mean annual death-rate is 16·3, which may be alternatively stated as 1,000 deaths among 61,215 persons. The age-distribution of the 61,215 is that of the whole male population of the 25–65 age period, and it is found that there are 22,586 aged 25–35, 17,418 aged 35–45, 12,885 aged 45–55, and 8,326 aged 55–65. The rates of mortality from the several causes (such as phthisis, accident, etc.) can also be stated in figures the total of which is the 1,000 which represents the entire mortality.

In like manner the smaller statistics of persons engaged in different occupations are dealt with, and in order to arrive at comparable results the standard age- and sex-distribution has to be observed. Thus,

as regards the earthenware manufacture it is ascertained from the census returns how many male potters are living at each of the four age-decennia between 25 and 65; and from the death returns what is the annual number of deaths among male potters at each age decennium. It is then easy to calculate the annual number of deaths among 22,586 potters aged 25-35, 17,418 aged 35-45, 12,885 aged 45-55, and 8,326 aged 55-65, making up the standard population of 61,215. The sum of these deaths, which among "all males" was 1,000, in the case of the potters reaches 1,706. These are comparable

Comparative Mortality according to Occupation, among Males 25-65 Years of Age.

England and Wales. 1890-91-92.	Phthisis.	Respiratory Diseases.	Nervous Diseases.	Urinary Diseases.	Liver Diseases.	Circulatory Diseases.	Gout.	Alcoholism.	Plumbism.	Suicide.	Accident.	Comp. Mort. Figures.
ALL MALES -	192	224	102	44	29	132	2	13	1	15	56	1900
Agriculturist	106	115	51	24	17	83	1	4	-	10	36	602
Fishermen	114	123	86	28	24	128		3	_	12	148	845
Cotton Manufacture	202	338	114	42	25	152	0	10	0	18	26	1141
Wool Manufacture -	191	256	100	45	21	131	1	3	3	12	21	991
Coal Miners	97	269	67	30	17	120	0	4	0	9	141	925
Coal Heavers	215	472	98	50	37	221	-	29	-	7		1528
Tin Miners	508	377	99	45	28	95	-	4	-	1000		1409
Lead Miners	380	225	62	41	34	142	-	5	5	5		1310
Glass Manufacture -	295	445	155	63	30	157	9	24	12	17	1	1487
Earthenware	333	668	123	63	32	227	1	9	17	16		1706
Plumbers	217	224	131	84	22	142	10	13	19	16		1120
Lead Workers	148	397	232	161	80	272	-	-	211	13		1783
Cutlers	382	518	91	56	27	167	-	18	3	29		1516
File-Makers	402	423	212	124	36	204	4	4	75	31	and the control of	1810
Printers	326	214	98	52	30	133	4	10	3	17	19	1096
Tailors	271	195	98	46	31	121	1	12	0	15	22	989
Drapers	260	181	102	36	33	135	-	14	1	15 12	19	1014 664
Grocers	131	121	66	40	26	94	1	8	=	19	25	920
Bakers	185	207	69	49	39	130	1	11	1	23	39	1096
Butchers	195	209	105	48	56	157	6	35	-	17		1427
Brewers	273	315	125	78	59	195	10	41	-	32		1642
Innkeepers	259	287	160	90	201	193	12	92	0	14		1652
Costermongers	443	403	139	70	29	222	3	36	-1	22		1829
Dock Labourers	325	564	114	68	26	236	3	52	1	44	102	1020
	1			-	200	100	100		100	014	-	1

figures, and in the same way comparable data as to mortality from phthisis and other special causes can be obtained for each occupation. A few examples are given on page 620.

The following list shows the comparative mortality figure, calculated in the same way, for a

number of other occupations:-

ALL MALI	ES	1		1,000	Blacksmiths 914
Occupied !	Male	S.		953	Shoemakers 920
Unoccupie	ed Ma	ales	1	2,215	Commercial Travellers 961
No more				LALER	Medical Men 966
					Shopkeepers generally 973
Clergy				533	Law Clerks 1,070
Gardeners		1		553	Hairdressers 1,099
Teachers	1			604	Hatters 1,109
Hosiery M	Ianuf	facture	9 .	698	Musicians 1,214
Lace Wor	kers			709	General Labourers . 1,221
Tanners				756	Iron and Steel Manu-
Artists				778	facture 1,301
Lawyers				821	Chimney Sweepers . 1,311

Phthisis and respiratory diseases cause a high mortality among debilitated persons, especially those exposed to weather, to hot or impure air, and to certain forms of organic or inorganic dust. Hence the mortality from these causes is high among costermongers, cutlers, file-makers, potters, printers, and highest of all in Cornish tin-miners. The comparatively low mortality from phthisis among coalminers may be attributed to the particles of coal being comparatively free from sharp angles and hence less irritating than stone or metallic dust (Hirt). The same immunity from phthisis has been observed among coal-miners in other countries, but it must not be forgotten that the nature of the employment excludes weakly persons, and that coal-mines are well ventilated.

Nervous diseases cause excessive mortality in occupations associated with alcohol or lead; the most fatal occupations in this respect are those of innkeepers, glass-makers, file-makers, and lead-workers. The incidence of gout and of fatal kidney disease is very similar. Mortality from diseases of the liver is in the main associated with mortality from alcoholism, both being high, for example, among commercial travellers, butchers, brewers, and innkeepers. Suicide also has a fairly close relation to intemperance.

Diseases of the circulatory system are most fatal among costermongers, cabmen, brewers, publicans, potters, cutlers, file-makers, and workers in glass.

These figures tend to establish a relation between intemperance and diseases of the heart, liver, kidneys, and nervous system, and also phthisis, gout, and suicide. Lead poisoning, similarly, is seen to be associated with diseases of the kidney, heart, and nervous system, and also with gout.

Value of statistical series and averages.— The mean or average of a series of numbers is a number which lies between the greatest and least of these, and stands in a definite dependence upon the whole of the series (Radicke). There are several kinds of means. Thus, if the series consists of four numbers—namely, a, b, c, and d—we have

The arithmetic mean, or simple average,
$$\frac{a+b+c+d}{4}$$
. The geometric mean, $\sqrt[4]{a\ b\ c\ d}$.

The harmonic mean,
$$\frac{4}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d}$$
.

The quadratic mean,
$$\sqrt{\frac{a^2+b^2+c^2+d^2}{4}}$$
.

When the terms of the series are equal, all kinds of means are also equal; but if otherwise, the quadratic is the highest, then the arithmetic, geometric, and harmonic, in the order stated.

The simple average, or arithmetic mean, is by far the most generally adopted, and mainly in three classes of cases:—

- 1. As a pure average; for example, the average age of a number of persons. The terms are exact, and the "mean" has no significance as regards any individual. Thus, the ages of ten persons whose deaths were attributed to diarrhea might be 1, 1, 1, 1, 71, 71, 1, 1, 71; and those of ten who died from enteric fever 17, 18, 26, 29, 25, 20, 16, 19, 24, 26. These two series have nothing in common as regards the individuals, but the average of each is 22. The mean temperature of a day or year is a simple average of the observations which are taken into account. Similarly, if we mix together equal volumes—say, an ounce of each — of four saline solutions containing respectively 10, 20, 30, and 40 grains of salt per ounce, the whole will contain 25 grains per ounce.
- Given two results equally likely to occur, the frequency of the one becomes more and more certainly equal to that of the other, in proportion as the number of observations increases. If a coin be tossed 1,000 times, it is improbable that heads will turn up much more or much less than 500 times; but with only 10 trials, there is little certainty of approximate equality. If in a given series of measurements of a fixed quantity (an angle, for instance), the errors of experiment are exactly as likely to be in excess as in deficiency, these errors will neutralise each other if we take the mean of a sufficiently large number of observations. An error of magnitude x in a single observation will

affect the mean to the extent $\frac{x}{10}$ in a series of ten ob-

servations, but only $\frac{x}{1,000}$ in a series of a thousand terms. The more extended the series, the greater the probability of accuracy, which increases as the square root of the number of observations.

If the measurements are of equal precision, a closer but less simple approximation to the probable true value sought may be obtained by adopting the principle of least squares: that is, by finding such a mean value that the sum of the squares of the residual errors of the observations shall be a minimum.

3. As a probable value of a variable quantity determined under average conditions; for example, the average daily excretion of urea, as determined by a series of more or less exact measurements under varying conditions as to food and exercise. In vital statistics absolutely fixed points are of rare occurrence, and most of the problems belong strictly to this class.

It may be important to determine the value of a series of observations; that is, how far the mean is a trustworthy approximation to the true value sought. As already explained, the value of a series increases with the number of observations, and with their equality. Another test is that of "successive means." For this purpose we take the mean of the first two terms $\left(\frac{a+b}{2}\right)$, of the first three $\left(\frac{a+b+c}{3}\right)$, of the first four $\left(\frac{a+b+c+d}{4}\right)$, and so on. These "successive means"

may vary sharply at first; but presently the first figure becomes stationary; then the second; and finally, if the series is sufficiently extended, the later successive means cease to vary materially. If the successive means, carried to the end of the series of terms, do not attain the required degree of constancy, the series is too short.

A further criterion is the determination of the

error; that is, the divergence of the individual terms of the series from its mean. The mean error in excess and the mean error in deficiency are simply the arithmetic means of the errors of those terms which fall respectively above and below the mean of the series. The mean error of the series is the average of the mean errors in excess and deficiency. The greater the mean error, the greater is the need for an extended series in order to compensate for the uncertainty of each observation. Radicke prefers to adopt the quadratic mean error rather than the arithmetic, described above, since the former increases more rapidly with the inequality of the terms.

Multiplying the mean error by 6745 or $\frac{2}{3}$ we get the *probable error*, so called because it is probable that, if the series were prolonged indefinitely, the errors would as often exceed as fall short of this

quantity.

The maximum error, whether in excess or deficiency, is useful as showing the possible error of a limited number of observations. Another way of estimating the error is to determine the "error of mean square." In a series of n terms, q being the quadratic mean and q the arithmetic mean, the "error of mean square" is $\frac{Q-A}{(n^2-n)}$.

The value of a series increases as the proportion of the "error" (of whatever kind) to the mean diminishes. The relative values of two or more series vary inversely as the square of their "probable errors."

Statistics are often presented in bulk, and the same question of sufficiency arises. For instance, out of 6,288 reported cases of scarlet fever 2,861 are at ages under five years; are these numbers sufficiently great to warrant the conclusion that the proportion, 45.5 per cent., is approximately constant? Such data are really the sum or average of suppressed series, and

their value is proportionate to the square root of their magnitude. Poisson proposed to measure the liability to error thus:—If of μ observations, m are in one category and n in other categories, and $m + n = \mu$, then the true proportion of the m category to the total

 μ lies between $\frac{m}{\mu} + 2$ $\sqrt{\frac{2mn}{\mu^3}}$ and $\frac{m}{\mu} - 2$ $\sqrt{\frac{2mn}{\mu^3}}$; that is, within a possible range of 4 $\sqrt{\frac{2mn}{\mu^3}}$. In the instance in point the range is 4 $\sqrt{\frac{2 \times 2,861 \times 3,427}{(6,288)^3}}$,

or about $\frac{36}{1,000}$. Hence the indicated proportion, 45.5 per cent., is liable to a range of error of 3.6 per cent., from 43.7 to 47.3.

It may also be possible to divide the total μ into two or more fairly equal parts, and thus ascertain if the proportion $\frac{m}{\mu}$ is approximately constant in each.

Such constancy would tend to show that the proportion is a fixed quantity, and not a mere average.

The preceding remarks have reference mainly to series that are intended to show by their mean some true fixed quantity. Sometimes they are used for a different purpose—namely, the exhibition of progressive changes or fluctuations. Here the general considerations as to magnitude still apply, but the determinations of "error" have no significance except in regard of the individual terms of the series. It is often convenient, for the purpose of comparing the fluctuations of different series, to adopt Buchan and Mitchell's method—namely, to state each term of a series as a percentage of the arithmetic mean of the series. In this way all series, whether the terms are large or small, are reduced to the same scale, and their true differences and resemblances can be seen at a

glance. The curves in Fig. 24 may serve as an

example.

If the terms of a series are based upon scanty statistics, the accidental irregularities may be so great as to obscure the true curve. Some degree of this is evident in respect of diphtheria in Fig. 24. A clearer indication of the general outline may in such cases often be obtained by Bloxam's method—that is, by substituting for each term the mean of three adjoining terms. Thus in a series a, b, c, d, e, f, g, etc., we should substitute (a), $\frac{a+b+c}{3}$, $\frac{b+c+d}{3}$, $\frac{c+d+e}{3}$, etc. This treatment is, of course, arbitrary, and curves so constructed are not to be regarded as exact in detail.

APPENDICES.

I. DEATH RATES IN SELECTED URBAN AND RURAL COUNTIES, 1891—1900.

ENGLAND AND WALES (TATHAM).

and a given	Cause	3,	10.01	Urban Counties.	Rural.
All Causes	7			20,325	14,800
Small Pox			 	13	3
Measles			 	512	212
Scarlet Fever			 	200	99
Whooping Cou			 	433	292
Diphtheria			 	297	201
Enteric Fever			 	198	116
Diarrhœa			 	909	382
Phthisis			 14	1,493	1,292
Infant Mortality			 	167	126

II. DEATH RATES AT AGES, ENGLAND AND WALES.

AGES.	A GRAD		Males	· ·		Females.						
Ten years } ended }	1860	1870	1880	1890	1900	1860	1870	1880	1890	1900		
All Ages	21.9	22.3	21.5	19.7	19.3	20.6	20.4	19:3	17.6	17:		
0-5	72.7	73.5	68.5	61.6	62.7	63.0	63.7	58.4	52.0	52.8		
5-10	8.5	8.2	6.7	5.4	4.3	8.4	7.8	6.3	5.3	4.		
10-15	4.9	4.5	3.7	3.0	2.4	5.1	4.5	3.7	3.1	2.0		
15-20	6.7	6.2	5.3	4.3	3.8	7.4	6.7	5.5	4.4	3.7		
20-25	8.8	8.5	7.4	5.7	5.1	8.6	8.0	6.8	5.5	4:		
25-35	9.6	9.9	9.4	7.8	6.8	10.0	9.7	8.6	7.4	6.1		
35-45	12.5	13.5	13.8	12.4	11.5	12.2	12.1	11.6	10.6	9.6		
45-55	18.0	19.2	20.1	19.4	19:0	15.3	15.6	15.6	15.1	14.8		
55-65	31.0	33.1	34.9	34.7	35.0	27.1	27.9	28.7	28.5	28.4		
65-75	65.5	67.1	69.7	70.4	70.4	58.9	59.1	61.0	60.4	60.7		
75-85	146.7	147.2	150.8	146.6	146.1	134.5	134.9	135.4	130.6	130.6		
Over 85	308-2	315.0	327.4	305.8	286:8	288 9	285.1	296.4	270.8	261 .4		

III. POPULATION.—ENGLAND AND WALES.

		Rate of	Females	Density.			
Census,	Population.	Decennial Increase.	per 100 Males.	Per Square Mile.	Per Inhabited House.		
1861	20,066,224	11.9	105:3	344	5.4		
1871	22,712,266	13.2	105.4	390	5.3		
1881	25,974,439	14.4	105.5	445	5.4		
1891	29,002,525	11.7	106.4	497	5.3		
1901	32,526,075	12.2*	106.9	558	5.2		

^{*} Highest (23.1) in urban districts of 50,000 to 100,000, diminishing in each direction to 9.3 in those over 250,000 and to 4.6 in those under 3,000.

IV. AGE AND SEX DISTRIBUTION OF POPULATION PER MILLION, ENGLAND AND WALES, CENSUS 1901.

Age- Periods.	Males.	Females.	Total.	Urban.	Rural.	
0-5	57,039	57,223	114,262	114,348	113,978	
5-10	53,462	53,747	107,209	105,555	112,760	
10-15	51,370	51,365	102,735	101,147	108,397	
15-20	49,420	50,376	99,796	101,080	95,489	
20-25	45,273	50,673	95,946	100,291	81,370	
25-35	76,425	85,154	161,579	167,597	141,390	
35-45	59,394	63,455	122,849	124,282	118,039	
45-55	42,924	46,298	89,222	88,368	92,086	
55-65	27,913	31,828	59,741	56,642	70,139	
65-75	14,691	18,389	33,080	29,502	45,080	
75-85	5,080	7,010	12,090	10,102	18,758	
Over 85	552	939	1,491	1,186	2,514	
All Ages	483,543	516,457	1,000,000	1,000,000*	1,000,000	

^{*} Males, 479,444; females, 520,556. + Males, 497,293; females, 502,707.

V. BIRTH RATES AND DEATH RATES, ENGLAND AND WALES.

Decennium. Birth Rate.		Births of Males per 100 Births	I	Infant Mor-		
	of Females.	Male.	Female. Total.		tality.	
1851—1860	34.1	104.6	23.0	21.3	22.4	154
1861-1870	35.2	104.2	23.6	21.3	22.4	154
1871-1880	35.5	103.8	22.6	20.0	21.3	149
1881-1890	32.4	103.7	20.5	18.0	19.1	142
1891-1900	29.9	103.6	19.3	17.2	18.2	154

VI. DEATH RATES PER MILLION, ENGLAND AND WALES.

CAUSES AND 5-YEAR PERIODS.

CAUSES.		years {	1880	1885	1890	1895	1900	1905
All Causes			20,791	19,403	18,895	18,718	17,685	16,004
Small Pox			78	78	14	20	7	25
Measles			385	413	468	407	421	326
Scarlet Fever			680	436	241	182	135	126
Typhus			34	23	7	4	1	1
Enteric Fever			277	216	179	174	175	113
Whooping Coup	gh		527	459	444	398	359	300
Diphtheria			122	156	170	253	272	204
Diarrhœa			853	672	681	651	817	673
Pneumonia			999	1,002	1,131	1,251	1,203	1,271
Erysipelas			81	83	54	48	35	36
Phthisis			2,040	1,830	1,635	1,462	1,322	1,215
Alcoholism			42	48	56	68	86	78
Rheumatic Feve			62	98	89	88	83	77
Cancer			494	548	632	711	800	864
Diabetes			40	51	62	69	81	89
Premature Birtl		-000	473	475	507	562	574	569
Heart			2,475	2,383	2,581	2,504	2,417	2,296
Bronchitis			2,377	2,154	2,136	2,074	1,563	1,237
Cirrhosis of Liv			110	120	122	120	135	121
Bright's Disease		4	188	222	248	257	278	379

II. DEATH RATES PER MILLION, ENGLAND AND WALES, 1891-1900.

CAUSES AND AGES.

Auses. Ages {	0-5	5-10	10-15	15-20	20-25	25-35	35-45	45-55	55-65	65-75	Ove 75
all Pox	29	10		5	11	16	18		10	10	
asles	3247	221	18		4	4	3	2	1	1	(
rlet Fever	844	353	81	33	22	15	8	4	2	1	
ooping Cough	3086	96		1	0	0	0	0	1	1	
htheria	1362	679	125	36		16	14	12	15	12	1
teric Fever	82	127	162	256	272	239	190	144	112	69	2
rrhœa	5485	37	10	8	12	22	40	85	234	708	207
eumonia	4945	297	112	206	281	448	852	1271	1953	2939	397
thisis	413	206	368	1144	1730	2135	2592	2362	1881	1154	43
icer	30	16	16	29	44	139	646	1833	3662	5604	614
betes	4	7	17	29	36	49	71	136	390	460	38
nchitis	6306	110	27	31	45	106	367	1214	3850	10060	2326
ght's disease	149	64	49	67	99	171	371	690	1218	1849	199

VIII. DEATH RATES PER MILLION, ENGLAND AND WALES, 1891-1900. CAUSES AND SEX.

Causes.	Male.	Female.	Causes.	Male.	Female.
All Causes	19,316	17,142	Diarrhœa	802	678
Small Pox	16	11	Pneumonia	5,425	4,468
Measles	439	391	Phthisis	441	385
Scarlet Fever	162	153	Cancer	600	906
Whooping Cough	349	404	Diabetes	86	66
Diphtheria	264	262	Bronchitis	1,849	1,775
Enteric Fever	200	150	Bright's Disease	379	299

IX. ENGLISH LIFE TABLE, BASED UPON THE MORTALITY STATISTICS OF 1891-1900 (TATHAM).

Males.				Females.				
Age.	Survivors at each age, out of	Expectation of Life.			Survivors at each age, out of	Expectation of Life.		
100	1,000,000 born.	1891-00	1881-90	1838-54	1,000,000 born.	1891-00	1831-90	1838-54
0	1,000,000	44.1	43.7	39.9	1,000,000	47.8	47.2	41.8
1	828,136	52.2	51.0	46.6	859,342	54.5	53.2	47.3
2	784,090	54.1	53.0	48.8	816,810	56.3	55.2	49.4
3	767,754	54.3	53.3	49.6	800,357	56.5	55.5	50.2
4	757,631	54.0	53.1	49.8	789,683	56.2	55.3	50.4
5	750,281	53.5	52.7	49.7	782,144	55.8	54.9	50.3
10	734,299	49.6	49.0	47.0	765,267	52.0	51.1	47.7
15	725,373	45.2	44.5	43.2	755,499	47.6	46.5	43.9
20	711,714	41.0	40.3	39.5	741,766	43.4	42.4	40.3
25	693,894	37.0	36.3	36.1	725,386	39.4	38.5	37.0
30	673,200	-33.1	32.5	32.8	705,819	35.4	34.8	33.8
35	648,169	29.2	28.9	29.4	682,147	31.5	31.2	30.6
40	615,964	25.6	25.4	26.1	653,014	27.8	27.6	27.3
45	577,010	22.2	22.1	22.8	619,184	24.2	24.0	24.1
50	530,888	18.9	18.8	19.5	580,320	20.6	20.6	20.7
55	475,849	15.8	15.7	16.4	533,105	17.2	17.2	17.4
60	409,518	12.9	12.9	13.5	473,037	14.1	14.1	14.3
65	332,344	10.3	10.3	10.8	398, 299	11.3	11.3	11.5
70	246,630	8.0	8.0	8.4	307,168	8.8	8.8	9.0
75	158,608	6.1	6.1	6.5	210,688	6.7	6.7	6.9
80	82,298	4.6	4.5	4.9	118,068	5.0	5.0	5.3
85	31,323	3.4	3.3	3.7	49,925	3.8	3.7	4.0
90	7,724	2.6	2.4	2.8	14,330	2.9	2.7	3.0
95	1,059	1.9	1.7	2.2	2,494	2.2	2.0	2.3
100	68	1.5	1.2	1.7	241	1.8	1.5	1.8

INDEX

A B C process, 267 Absorption of heat by soil, 159 Abyssinian tube wells, 43 Acclimatisation theory of protection, 304 Acid, Carbonic, in air, 3, 13, 180 -, Sulphurous, in gas, 5 Actinomycosis, 150, 308 Acts of Parliament, 490 Adam's method, 130 Adoptive Act, 508 Adulterations of brandy, 156

— of butter, 143

— of flour, 148

— of milk, 125 Aerated waters, 61 Aerobic microbes, 310 Aeroscope, 12 Age-distribution of population, 607, 632 - as affecting mortality, 609, 613, 621, 622 and severity of attack, 300 — and susceptibility, 300 Ague, 445 Air, 3 -, Analysis of, 180, 184 —, Bacteria in, 7 —, Composition of, 3 -, Dust in, 9 -, Examination of, 12 —, Impurities of, 12 — of sewers, 257 —, Physical properties of, 6 Air space. See Cubic space. Albuminoid ammonia, 79 Albumoses, 310 Alcoholic Beverages, 152 Alexines, 316 Alkali Acts, 11, 575
— works, Fumes from, 11, 575 Alkaline permanganate solution, Alluvial soil, 171 Alumino-ferric, 265 Ambulant plague, 443 Amines process, 266

Ammonia in air, 4, 17
—, Albuminoid, 79
—, Free 78 —, Free, 78 — process of water analysis, 78 Ammoniacal fermentation of urine, 307 Amœbæ, Parasitic, 284 Anæmia due to parasites, 291 Anaërobic microbes, 310 Analyst, Public, 531
Anderson's process, 266 Anemometer, 30 Aneroid barometer, 28 Animal food, Advantages of, 98 — matter in water, 84, 89 - -, Diseases attributed — parasites, 284 Animals, Keeping of, 481, 495, 551, 583, 594, 597 Ankylostoma, 289 Ankylostomiasis, 291 Anopheles, 449 Annual variations of barometer, 29 —— of birth-rate, 608 —— of death-rate, 620 — of disease, 619, 621 — of rainfall, 29 — of thermometer, 19 Anthrax, 425, 573
—, Bacillus of, 427 Anticyclone, 32 "Anti D" trap, 241 Antidote theory of protection, 303 Antill's trap, 242 Antiscorbutics, 100 Antiseptics, 332 Antitoxins, 315 Apjohn's formula, 23 Arachnida, 285 Argon in air, 3 Arithmetic mean, 625 Arrowroot, 152 Arsenic in beer, 154 — in water, 74 Arsenical poisoning, 572 Artesian wells, 43

Artificial ventilation, 193 Artisans' dwellings, 202 Ascaris lumbricoides, 289 Ash-closets, 249 Ashes, Removal of, 235, 254, 492, 509, 578 Ashpits, Byelaws as to, 597 —, Cleansing of, 583 -, Construction of, 592 Asiatic cholera, 431 Aspect of houses, 176 Atmometer, 24 Atmospheric pressure, 25 Attenuation of virus, 312 Auxiliary Agencies, 486 Averages, 625

Bacilli, 306 -, Pathogenic, 307 Bacillus anthracis, 427 - dysenteriæ, 455 --- enteritidis sporogenes, 400 - icteroides, 440 --- lepræ, 452 --- pestis, 444 Back-to-back houses, 177 Bacteria and disease, 306 — in air, 7 — in milk, 126 —— in sewage, 57, 263 — in soil, 163 —, Pathogenic, 59 Bacterial infection, 104 - treatment of sewage, 271 Bacteriological laboratories, 437 Bakehouses, 482, 568
Ballard on diarrheal mortality, 395 Barff iron, 72 Barley, 150 Barometers, Aneroid, 28 -, Glycerine, 28 —, Mercurial, 25 —, Water, 28 Beaufort's scale, 31 Beer, 151 -, Arsenic in, 154 Beggiatoa, 67 Bell trap, 242 Beri-beri, 456 Berkefeld filter, 50 Bilharzia hæmatobia, 288 Bird's process, 266 Birth-rates, 607, 632 - as affecting death-rates, 611 Births, Legitimate and illegiti-mate, 607

Births, Male and female, 607 Births and Deaths Registration Act, 483, 607 - ___, Seasonal curve of, 608 Bischof's filter, 49 Bishopsgate, Plague on board the, Black-ash process, 267 Blair's process, 83 Bloxam's method, 630 Boiling of water, 48 Boracic acid as antiseptic, 333 — as disinfectant, 326 Borax as antiseptic, 333 Borough Councils, 576 Bothriocephalus, 287 Bovine diphtheria, 133 Bovine tuberculosis, 419 Boyle and Marriotte's law, 6 Boyle's valves, 190 Brandy, Adulteration of, 156 Brassfounder's ague, 573 Bread, 95, 149 -, Adulterations of, 149 Brick, Permeability of, 172 Bristol, Sewers at, 259 Broad irrigation, 268 Bromine, Disinfection by, 326 Bubonic plague, 442 Buchan and Mitchell's method, 629Building materials, 172 — sites, 169 Buildings, Byelaws as to, 580, 586 —, Dangerous, 503 —, New, 502, 509, 580 —, Obstructive, 525 Burial Act, 281, 282 — grounds, 277 — grounds, 277
— , Air of, 279
— , Byelaws as to, 580, 599
— , Closure of, 281
— , Disused, 281
— , L.G.B. Memo. as to, 280
— , S.A. to provide, 281 ---, Power to provide for and enforce, 515 Butter, 142 - adulteration, 143 — analysis, 95, 143 —, Composition of, 143
— and disease, 144 - and Margarine Act, 1907, 535 Butyric acid fermentation, 128 Buys-Ballot's law, 34 Byelaws, 508, 578 —, Annotated Model, 582 — for urban and rural districts, 490, 508

Byelaws, L.C.C. must make, 552 Cholera and soil, 433 -, Model, 582 — and water, 434 Calcium salts in water, 62, 67, 74 - in England, 431 Calculi, Urinary, 76 Calf vaccination, 348, 350 Chyluria, 293 Cisterns, House, 47, 493 Campbell's sunshine recorder, 20 —, Separate, for closets, 247 Canal Boats Acts, 538, 540 Clarine process, 267 Cancer mortality, 612 Clark's process, 48 Carbohydrates, 91 Claviceps purpurea, 150 Carbolic acid as deodorant, 333 Clay sites, 169 ----- as disinfectant, 327 ----- oil, 326 Cleansing of houses, Compulsory. Cleansing of Persons Act, 554 Carbon filters, 49 Carbonates in water, 63 Climate, 35 Carbonic acid in air, 3, 13, 180 Closets, Ash, 249 ----- in soil, 162 —, Byelaws as to, 583 ---- in water, 60 - common to two or more — oxide in air, 4, 16, 197 houses, 508 —, Earth, 249, 251, 583, 591 —, Flush, 248 Carferal filter, 49 Carrots, 95 — for factories, 491 Casein, 121 Caterham outbreak, 384, 390 — for schools, 207 - for tenement houses, number Causes of death, Classification of, 611, 620 required, 597 Cellar dwellings, 493, 550 - for houses, Compulsory, 491, Cement, 172 Cemeteries, 278, 580, 599
—, Air of, 279 —, Pail, 250, 252, 253, 254 —, S.A. responsible for supervision of, 491 Census, 605 -, Revolver, 248 Cerebro-spinal fever, 404 —, Trough, 248 —, Water, 245, 590 - meningitis, Epidemic. See Cerebro-spinal fever. Closure of houses unfit for habi-Cesspools, 263, 583, 592 tation, 524, 593 Chalk, 170 —— soils, 170 - of polluted wells or cisterns, —, Water from, 63, 64 493 Chamberland's filter, 52 Coccidia, 285 Cockles, 126 Charcoal filters, 49 Cœnurus cerebralis, 102 Charles's law, 6 Coffins, 279 Cheese, 96, 145 Cohn's solution, 310 Chemical composition of food substances, 95 Coke, 196 — disinfection, 325 — purification of water, 48 Cols, 33 Combination of sanitary districts, Chicken-pox, 215, 355 Children in factories, 574 Combustion, 11 - process, 79 Chittenden on dietary, 94 Comma bacillus, 435 Chloride of lime, 326 Committees, Powers of, 505 Chlorides in water, 62 Common lodging houses, 494, 510 Chlorine as deodorant, 333 -- ---, byelaws, 579, 584 - as disinfectant, 328 Comparative mortality figure, 609, Cholera, 431 -, Microbe of, 435 Compo for walls, 172 - nostras, 437 — regulations, 470 — and ground water, 168, 433 Compulsory detention in hospital, 515 — disinfection, 500, 513 — and locality, 434 — and rainfall, 432 --- notification, 458 —— removal to hospital, 500 — and season, 432

Concrete, 172
Condensed milk, 141
Conder's process, 265, 267
Conduits, 46
Constant water supply, 47
Consumption. See Phthisis.
Contagious diseases, 295
Container closet, 246
Contamination of air, 7
— of water by lead, 69
by organic matter, 40
Contributory place, 505
Cooking, Chemical effect of, 112
Copper salts in water, 74
Cornices, Ventilating, 188
Corrected death-rates, 608, 614
Cotton clothing, 320
- workers, Mortality among,
623
Cough-spray, 418
County authorities may regulate
dog-muzzling, 425
— Councils, 473, 519, 522, 526,
537, 551, 552, 559, 576
Cow-pox, 350
Cow-sheds. See Dairies.
Cream, 122
Cremation, 282
Croup, 377
Cubic space, 183
— and mortality, 185
———— in canal boats, 539
in common lodging-
houses, 584
———— in cowsheds, 543
in factories, 561
in hospitals, 222
in tenement houses, 596
Culex, 449
Cultivation of microbes, 310
Cupric sulphate as deodorant, 333
———— as disinfectant, 326
Customs and Inland Revenue Act,
1890, 579
Cyclones, 32
Cysticercus, 287
Cysticercus, 207
D twon 949
D-trap, 242
Dairies, cowsheds, and milkshops
Orders, 541
, Drainage of, 542,
547
. — , Infectious diseases
in, 547
, Lighting of, 545,
546
— — , Registration of, 541
— — , Regulations re-
specting, 544

Dairies, cowsheds, and milkshops. Swine in, 543 - ---, Ventilation of, 545, 546 Damp and disease, 167 — course, 174, 588 — soil, 161 Dangerous buildings, 503 Daniells' hygrometer, 22 Darnel, 150 Dead, Disposal of the, 277 Death, Causes of, 611, 633 —, daily average, 277 —, Registration of, 608 - - and density of population, 185, 394, 610 — at ages, 609, 613, 631, 632, 633, 634 — —, Calculation of, 605 — —, Corrected, 608, 614 - influenced by birth-rates. 611 , Male and female, 610, 612, 631, 632 - - Progressive changes in, 611, 620 — —, Urban and rural, 609,631 — —, Zymotic, 612, 614 -, Zymotic, 612, 614 Deaths at all ages, Statistical table, 631 - from all causes, Statistical table, 633 Default of S.A., Procedure in case of, 505, 525, 552 Dengue, 441 De novo hypothesis, 296 Density of population as affecting mortality, 185, 394, 610 Deodorants, 333 Destructors, 255 Dew-point of temperature, 21, 23 Diarrhœa, Age and, 397 — and density of population, 393 -, Bacillus of, 400 —, Earth-temperatures and, 396 — Epidemic, 393, 396 —, Locality and, 399 -, Milk and, 135, 398 -, Outbreaks in public institutions, 399 —, Season and, 398 ---, Sex and, 398 -, Soil and, 395 —, Temperature and, 395 —, Water and, 395, 399 Dibden's treatment of bacteria, 272 Diet, Animal and vegetable, 98 ---, Disease and, 99

Diet, Essentials of, 90 Dietaries, 92 -, Calculation of, 94 Dines' hygrometer, 22 Diphtheria, 214, 367, 460, 631, 633 -, Air and, 370 -, Animals and, 369 -, Antitoxin treatment of, 375 -, Bacteriology of, 373 Bovine, 133 -, Locality and, 369 -, Milk and, 370 -, Season and, 368 -, Sewer-gas and, 36 -, Virus of, 372 Diplococci, 306 Disconnection chamber, 239 - from drain, 239, 590 - from sewer, 239 - of waste-pipes, 241 Disease maps, 464 Diseases common to man and lower animals, 298 -, Contagious, 295 -, Endemic, 297 -, Epidemic, 297
- for which schools must be closed, 208 -, Infectious, 294, 499 -, Inoculable, 295 -, Malarial, 445 -, Notification of, 458 -, Pandemic, 297 -, Specific, 294, 334 -, Telluric, 166 -, Tropical, 431 Disinfecting apparatus, S.A. to provide, 552 Disinfection, 318 - by heat, 319 by hot air, 321, 324
by steam, 321, 325 -, Chemical, 325 —, Compulsory, 500, 513 - in sick-rooms, 466 — of rooms, 328, 469 , Shelter for inmates during, 468 Disposal of ashes and household refuse, 235, 254, 491, 509, 516, 583, Disposal of the dead, 277 — of sewage, 263, 537 — of trade refuse, 254, 537,595, 602 Distillation of water, 48 Distoma hepaticum, 102, 289 District Councils, 521 Disused Burial Grounds Act, 282 Dochmius duodenalis, 290

Dœcker huts, 227 Dolcoath anæmia, 291 Dolomite. See Magnesian limestone. Dog-muzzling and rabies, 425 Domestic factories andworkshops, Drainage, 161, 235 —— and disease, 167, 168 ——, Byelaws as to, 509, 579, 588, 602 -, Compulsory, for every house, 490 —— districts, Special, 505 — of subsoil, 161, 237 Drains, Construction of, 235, 589 -, Definition of, 494 -, Dimensions of, 237 -, Disconnection of, 238, 590 —, Testing of, 244 —, Trapping of, 589 —, Ventilation of, 239, 589 Draught, Velocity of, 191 Dry heat, Disinfection by, 321, 324 — rot, 175 systems of excreta disposal, 249 Ducat's treatment of sewage bacteria, 271 Ducker huts, 227 Dumb wells, 263 Duration of life, Probable, 615 Dust, 9 - and disease, 7 Dysentery, Amoboid parasite in, Earth-burial, 277 —— closets, 249, 251, 583, 591 - temperatures, 20 Eaves-gutters, 503

Eberth-Gaffky bacillus, 391 Education Committee of L.C.C. and infectious diseases, 216 Education, Elementary (Blind and deaf children), Act, 1893, 218 - (Defective and epileptic children), Act, 1899, 218 Effluents, Sewage, 256 —, Trade, 41, 256, 274, 536 Eggs, 95, 147 Egyptian chlorosis, 291 Ehrlich's side chain protection theory, 305 Electrolytic process, 267 Elementary Education. See Education. Elephantiasis, 293 Employment of Children Act, 576 Endemic diseases, 297

English life table, 634
Enteric fever, 382, 463, 633, 634
— and ground-water, 167,
— and locality, 389 [384
— and milk, 387
and organia 207
———— and oysters, 387
— and season ,382
———— and sewer-gas, 388
———— and soil, 383
———— and water, 384
— at Caterham, 384, 390
——— at Lincoln, 385, 390
— at Munich, 166, 383
at Worthing, 385
———, Bacillus of, 391
Entry upon private premises, 497
Epidemic diseases, 297
—, variation in type, 300
Epidemics, Milk, 132, 513
—, Water, 89, 384, 434
Equifex disinfector, 323
Ergot in flour, 148
— in rye, 150
Error, Mean, 628
of moon governo 690
of mean square, 628
—, Probable, 628
Erysipelas, 378
Essential oils as antiseptics, 333
— as deodorants, 333
Evaporation, 24
—— from leaves, 159
Exanthemata, 299
Excreta, Amount of, 262
—, Disposal of, 245, 251
, Removal of, 251, 491
, Different systems com-
pared, 251
, Economic advantages of
different systems compared, 251
—, Theoretical value of, 262
Exhaustion-theory of protection,
303 Enhumation Effect on health 272
Exhumation, Effect on health, 273
Expectation of life, 615 Expired air, Composition of, 10
Expired air, Composition of, 10
—, Volume of, 179
Factory and Workshop Acts, 557
Factories, Fire escape from, 567
-, Sanitary accommodation of,
564
Ventilation and cleanliness
of, 495, 560, 561
Facultative parasites, 309
Fairs and markets, Byelaws, 580
Famine fever, 378
Farcy, 425
Fat as food, 91
—, Composition of, 110

Fat in butter, 143 — in milk, 122, 130 Ferozone, 267 Ferrous sulphate as deodorant, 333 - — as disinfectant, 326 Ferruginous water, 69 Fever, Cerebro-spinal, 404 -, Enteric, 382, 633, 634 ---, Famine, 378 —, Gaol, 380 —, Intermittent, 445 —, Malta, 454 ---, Puerperal, 367, 430 —, Relapsing, 378 —, Remittent, 446 —, Scarlet, 361, 462, 463, 631, 633, 634 -, Simple continued, 393 —, Typhoid. See Enteric. —, Typhus, 379 —, Yellow, 437 Filaria, 292 Filter-beds, 50 Filters, 49 Filth-conditions and disease, 89, 168, 373, 385, 394, 434, 435, 438 Filtration, Effects of, 49, 50 - of milk, 141 — of sewage, 266, 267 — of water, 49 Filtre Rapide, 49 Fire, means of escape from, in factories and workshops, 603 -Fires, Gas, 196 —, Open, 34 Fish, 116 — and disease, 116 —, Composition of, 95 —, Preservation of, 116 Flagellates, 285 Flour, 95, 147 Flush-closets, 248 — tanks, 243 Flushing of closets, 246 - of drains, 243 - of sewers, 257 Fogs, Cause of, 21 Fomites, 295 Food and disease, 98, 397 and Drugs Acts, 529, 532, 533 -, Classification of, 90 -, Composition of, 95 —, unsound, Seizure of, 114, 499, Foot and mouth disease, 131 Forbes and Price's process, 266 Forchammer's process, 81 Forecasts, Weather, 34 Formic aldehyde vapour as disinfectant, 330

Fortin's barometer, 25 Fosse mobile, 249 - permanente, 49 Foul air and disease, 182 Frankland's combustion process, - filtration process, 13 Fruit-pickers, Byelaws as to, 579 Fulham hospital and small-pox,335 Fumigation, 328 Gametocytes, 448 Gaol fever, 380 Garget, 131 Gases in air, 3, 8 - in sewers, 257 —— in soil, 162 ___ in water, 62 Gas-fires, 196 —, Lighting, 181 — washings, 493 Gelatin as food, 91 General Powers Act. See London County Council. Geological formations, 170 Geometric mean, 625 Germ theory, 301 German measles, 361 Glaisher's tables, 23 Glanders, 425 Glossina palpalis, 285 Gloucester, Small-pox at, 335 Glycerinated calf lymph, 3 Glycerine barometer, 28 Goddard, Massey, and Warner's disinfector, 323 Goitre, 76 Gordon's process, 13 Goux closets, 250 Granite, Permeability of, 162 - sites, 170 -, Water from, 170 Gravel sites, 171 -, Water from, 171 Graves, 277, 278, 279, 599 Graveyards, Air of, 279 —, Byelaws, 580, 599 —, effect upon health, 278 - -, Pollution of water by, ----, Position of, 278 Grease traps, 242 Greenwich factors, 23 Gregarines, 285 - water, 160 — and disease, 167, 168, 169

Guinea-worm, 293 Gullies, 241 Hæmaturia, Intermittent, 288 Hæmosporidia, 285 Haffkine's cholera vaccine, 437

Haldane's method, 15 Halibut, Poisoning by, 116 Handicraft classes for schools 207

Hardness of water, 74
—, Effects of, 75, 76
—, Tests for, 75

Harmonic mean, 625 Health affected by articles of

food, 157 Heat, Disinfection by, 319 Heating by fires, 195

- by hot pipes, 198 — by stoves, 197 Height of rooms, 588 Hendon disease, 133 Hermite process, 267 Hesse's process, 12

Hille's process, 266 Home isolation, 465 - work, 570

Hooke's anemometer, 30 Hopper closets, 246

Hop-pickers, Byelaws as to, 579,

Horseflesh, Sale of, Act to regulate, 535

Hospitals, Compulsory removal to, 300

-, Construction of, 222, 516

—, Cubic space in, 222 —, Detention in, 515

—, Effect of Isolation in, 232

—, Floor space in, 223
— for infectious diseases, 224, 502
—, Number of beds in, 228
—, Permanent and temporary,

—, Ventilation of, 223 —, Wards, 222

Hot air, Disinfection by, 321, 324

———, Warming by, 198 - water, Warming by, 198

House-cleansing, 492 — -drainage, 490

Household refuse, 235, 254, 491,

509, 583, 595 - test for carbonic acid, 15 Houses, Construction of, 172, 587

_, ___, Byelaws regarding, 580,

- Inspection of, 199, 593

- let in lodgings, Byelaws re garding, 494, 579 596

Houses unfit for habitation, 497,	Infective diseases, 228
524, 593	Influenza, 400
Housing of the Working Classes	— and season, 402
Acts, 522, 523, 527, 528	—, Bacillus of, 403
Houston, Dr., test of, for B. coli, 60	— epidemics, 401
Humidity of air, 20	, Incubation of, 403
— in workrooms, 564	Inhabited vans and tents, 496, 522
, Relative, 23	————, Byelaws, 579
—— of soil, 150	Inoculation of disease, 295
Hydatids, 288	— of leprosy, 451
Hydraulic mean depth, 261	of small-pox, 317, 340
Hydrochloric acid as deodorant,	— of tuberculosis, 416
333	, Protective, 312, 313
Hydrophobia, 422	Inorganic nitrogen, 64
—, Pasteur's experiments, 423	Insecta, 285
Hygrometers, 22	Inspection of houses, 199, 593
T	Inspectors, Meat, 113
Ice-cream and Infection, 139	-, Sanitary. See Sanitary In-
Immunity from specific diseases,	spector
303, 316	Interment, Power to provide for
Improvement schemes, 523	and enforce, 514
Impurities of air, 7, 12	Intermittent fever, 445
— of soil, 163	—— filtration, 268
— of water, 54, 280	hæmaturia, 288
Incubation of disease, 298	
	— water supply, 47
Industrial poisoning, 571	Iron hospitals, 227
Infant Life Protection Act, 554	salts in water, 68
— mortality, 611, 615	—, Sulphate of, as disinfectant,
Infants' Milk Depôts, 486	326, 332, 333
Infection, 294	Irrigation, Broad, 268
by air, 234, 295, 318, 335	—— of sewage, 263
by food, 98, 397	—, Subsoil, 263
— by milk, 131, 397, 418, 513	Isobars, 32
— by rats, 445	Isolation hospitals, 225, 232, 516
— by water, 89, 295, 384, 434	——————————————————————————————————————
Infectious diseases, 294	————, 1901, 517
	— —, Removal to, 465, 514
, Administration as to, in	Compulsory 500
elementary schools, 216	—, —, Compulsory, 500
— and schools, 208	
—— ——, Disease maps, 464	Joint boards for sanitary pur-
———, Disinfection, 466	poses, 505
————, Hospitals for, 224, 502	Jordan's barometer, 28
in canal boats, 538	
- in common lodging-	— sunshine recorder, 20
houses, 585	
- in inhabited tents or	Kefir, 128
vans, 533	Kemp's tester, 245
— in tenement houses, 598	Kernig's sign, 405
,Inquiries respecting, 463,	Kew barometer, 25
475	Kilmarnock, Small-pox at, 342
, Isolation of, 465, 552	Kingzett's tester, 245
— , Law regarding, 499	Kjeldahl's process, 80
— Notification Act, 458	Klebs-Löffler bacillus, 373
———, Precautions, 463	Knackers' yards, 504
——————————————————————————————————————	Koch's bovine tuberculosis, 420
, Prevention of, 457	experiments with cholera
, Register of sickness, 464	microbe, 437
, Regulations, 463	filtration process, 13
refuse, 515	Koumiss, 128

Lactic fermentation, 127 Lactose, 122 Land-filters for sewage, 268 Lard, 147 Laundries, 568 Laveran's bodies, 447 Lead in water, 69 poisoning, 73, 571Tests for, 73 Leguminosæ, 150 Leprosy, 449
and fish diet, 450 -, Bacillus of, 452 - Commission, 453 -, Heredity in, 451 —, Inoculation of, 451 —, Isolation of, 453 Liernur's system, 249 Life, Expectation of, 615 Life-tables, 616 ---, English, 634 Lighting of schools, 203 Lime-juice, 100 - salts in water, 67 — stone, 62, 63, 170, 172 -, Treatment of sewage by, 266, 267 Lincoln outbreak, 385, 390 Linen clothing, 320 Living organisms in water, 54 Local Government Act, 473, 519, — Board, Dairies Order, 541 - — Memoranda regarding cemeteries, 280 - - Memoranda regarding schools, 208
— — Memoranda regarding water, 38 - Orders relating to M.O.H., 473, 475, 477 - - Regulations as to canal boats, 538, 540 — — as to cholera, diseases, 470 — as to epidemic etc., 470 - - as to infectious diseases, 463, 501 Lodging-houses, 202 for seamen, Byelaws, 579 for the Working Class, Lodgings, Houses let in, 494, 579, Lolium temulentum, 150 London Building Act, 554 - County Council, Education Committee of, 216

London County Council (General Powers) Act, 1907, 555 - Public Health Act, 549 Lowcock's treatment of sewage bacteria, 271 Macrogametocytes, 448 Made soil, 163, 170 Magnesia salts in water, 61, 68 -, Tests for, 68 Magnesian limestone, 62, 63, 68, 170 Maignen's Filtre Rapide, 49 Mains, Water, 46 Maize, 95, 150, 152 Malaria, 167, 445 —, Etiology of, 447 Malt liquors, 153 Malta fever, 454 Manchester, Sewage treatment at, Manholes, 239, 257 Manufactures, Effluents from, 41, 256, 536 Manurial value of sewage, 262 Maps, 484 —, Disease, 464 Margarine, 144 — Act, 535 Markets and Fairs, Byelaws as to, Marsh gas, 4 Marsh-Berzelius test, 156 McKinnell's ventilator, 189 Mean after life-time, 616 - age at death, 615 — duration of life, 615 — error, 628 Means and averages, 620 —, Arithmetic, 625 —, Geometric, 625 —, Harmonic, 625 —, Quadratic, 625 -, Successive, 627 Measles, 213, 356, 619, 620, 631, 633 —, German, 361 — of swine, 287 Meat and disease, 101 —, Characters of good, 109 —, Classification of unwhole. some, 115 —, Composition of, 95, 100 —, Inspection of, 112 —, Parasites in, 102, 109 — poisoning, 102 —, Preservation of, 110 Mechanical impurities in air, 7

--- ventilation, 193

- work done by food, 97

Medical inspection of school
children, 219, 481
- Officer of Health, 473
, Appointment of,
473, 504, 552 Appointment of,
— — , Duties of, 475. (See
also chapters xviii. and xx.)
473
————, Reports of, 477, 479
, Tenure of office, 474
Mediterranean fever, 454
Meningitis, Tubercular, 419
Monehant Chinaian (Tinling
Merchant Shipping (Fishing
Boats) Act, 1883, 579
Mercurial poisoning, 572
Mercuric chloride as antiseptic,
332
as disinfectant, 327
Metchnikoff's theory of protec-
tion, 304
Meteorology, 18
Methane in air, 4
Microbes, 306
, Pathogenic, 309
- See Bacteria.
Micrococci, 306
—, Pathogenic, 307
Micrococcus melitensis, 454
Microgametocyte, 448
Middens, 249, 252, 253, 254
75:231 1 T
Middlesbrough, Pneumonia at,
409
Midwives Act, 518
Milk, 121
—— and disease, 131, 418, 513
, Adulteration of, 125
—, Bacteria in, 126
—, Composition of, 95, 122
, Condensed, 141
—, Condensed, 141 —, Control of supplies, 139
—, Condensed, 141 —, Control of supplies, 139
—, Control of supplies, 139 — epidemics, 132, 513
—, Control of supplies, 139 — epidemics, 132, 513
—, Control of supplies, 139 — epidemics, 132, 513
—, Control of supplies, 139 — epidemics, 132, 513
——, Control of supplies, 139 —— epidemics, 132, 513 ——, Examination of, 129 ——, Fermentation of, 127 ——, Preservation of, 140
——, Control of supplies, 139 —— epidemics, 132, 513 ——, Examination of, 129 ——, Fermentation of, 127 ——, Preservation of, 140 ——, Prohibition of sale of, 513, 544
——, Control of supplies, 139 —— epidemics, 132, 513 ——, Examination of, 129 ——, Fermentation of, 127 ——, Preservation of, 140 ——, Prohibition of sale of, 513, 544 ——, Regulations for sale of, 125
——, Control of supplies, 139 —— epidemics, 132, 513 ——, Examination of, 129 ——, Fermentation of, 127 ——, Preservation of, 140 ——, Prohibition of sale of, 513, 544 ——, Regulations for sale of, 125
——, Control of supplies, 139 —— epidemics, 132, 513 ——, Examination of, 129 ——, Fermentation of, 127 ——, Preservation of, 140 ——, Prohibition of sale of, 513, 544 ——, Regulations for sale of, 125 ——, Standard for, 123
—, Control of supplies, 139 — epidemics, 132, 513 —, Examination of, 129 —, Fermentation of, 127 —, Preservation of, 140 —, Prohibition of sale of, 513, 544 —, Regulations for sale of, 125 —, Standard for, 123 —, Sterilisation of, 141
—, Control of supplies, 139 — epidemics, 132, 513 —, Examination of, 129 —, Fermentation of, 127 —, Preservation of, 140 —, Prohibition of sale of, 513, 544 —, Regulations for sale of, 125 —, Standard for, 123 —, Sterilisation of, 141 — sugar, 122, 127
—, Control of supplies, 139 — epidemics, 132, 513 —, Examination of, 129 —, Fermentation of, 127 —, Preservation of, 140 —, Prohibition of sale of, 513, 544 —, Regulations for sale of, 125 —, Standard for, 123 —, Sterilisation of, 141 — sugar, 122, 127
—, Control of supplies, 139 — epidemics, 132, 513 —, Examination of, 129 —, Fermentation of, 127 —, Preservation of, 140 —, Prohibition of sale of, 513, 544 —, Regulations for sale of, 125 —, Standard for, 123 —, Sterilisation of, 141 — sugar, 122, 127 Milkshops. See Dairies.
—, Control of supplies, 139 — epidemics, 132, 513 —, Examination of, 129 —, Fermentation of, 127 —, Preservation of, 140 —, Prohibition of sale of, 513, 544 —, Regulations for sale of, 125 —, Standard for, 123 —, Sterilisation of, 141 — sugar, 122, 127 Milkshops. See Dairies. Millstone grit, 170
—, Control of supplies, 139 — epidemics, 132, 513 —, Examination of, 129 —, Fermentation of, 127 —, Preservation of, 140 —, Prohibition of sale of, 513, 544 —, Regulations for sale of, 125 —, Standard for, 123 —, Sterilisation of, 141 — sugar, 122, 127 Milkshops. See Dairies. Millstone grit, 170 Milne's hypothesis, 617
——, Control of supplies, 139 —— epidemics, 132, 513 ——, Examination of, 129 ——, Fermentation of, 127 ——, Preservation of, 140 ——, Prohibition of sale of, 513, 544 ——, Regulations for sale of, 125 ——, Standard for, 123 ——, Sterilisation of, 141 —— sugar, 122, 127 Milkshops. See Dairies. Millstone grit, 170 Milne's hypothesis, 617 Mineral salts in water, 61
——, Control of supplies, 139 —— epidemics, 132, 513 ——, Examination of, 129 ——, Fermentation of, 127 ——, Preservation of, 140 ——, Prohibition of sale of, 513, 544 ——, Regulations for sale of, 125 ——, Standard for, 123 ——, Sterilisation of, 141 —— sugar, 122, 127 Milkshops. See Dairies. Millstone grit, 170 Milne's hypothesis, 617 Mineral salts in water, 61
—, Control of supplies, 139 — epidemics, 132, 513 —, Examination of, 129 —, Fermentation of, 127 —, Preservation of, 140 —, Prohibition of sale of, 513, 544 —, Regulations for sale of, 125 —, Standard for, 123 —, Sterilisation of, 141 — sugar, 122, 127 Milkshops. See Dairies. Millstone grit, 170 Milne's hypothesis, 617 Mineral salts in water, 61 Miners, Mortality among, 623, 624
—, Control of supplies, 139 — epidemics, 132, 513 —, Examination of, 129 —, Fermentation of, 127 —, Preservation of, 140 —, Prohibition of sale of, 513, 544 —, Regulations for sale of, 125 —, Standard for, 123 —, Sterilisation of, 141 — sugar, 122, 127 Milkshops. See Dairies. Millstone grit, 170 Milne's hypothesis, 617 Mineral salts in water, 61 Miners, Mortality among, 623, 624 Mines, Control of, 575
—, Control of supplies, 139 — epidemics, 132, 513 —, Examination of, 129 —, Fermentation of, 127 —, Preservation of, 140 —, Prohibition of sale of, 513, 544 —, Regulations for sale of, 125 —, Standard for, 123 —, Sterilisation of, 141 — sugar, 122, 127 Milkshops. See Dairies. Millstone grit, 170 Milne's hypothesis, 617 Mineral salts in water, 61 Miners, Mortality among, 623, 624 Mines, Control of, 575 —, Refuse from, 536
—, Control of supplies, 139 — epidemics, 132, 513 —, Examination of, 129 —, Fermentation of, 127 —, Preservation of, 140 —, Prohibition of sale of, 513, 544 —, Regulations for sale of, 125 —, Standard for, 123 —, Sterilisation of, 141 — sugar, 122, 127 Milkshops. See Dairies. Millstone grit, 170 Milne's hypothesis, 617 Mineral salts in water, 61 Miners, Mortality among, 623, 624 Mines, Control of, 575 —, Refuse from, 536 Minimetric test, 15
—, Control of supplies, 139 — epidemics, 132, 513 —, Examination of, 129 —, Fermentation of, 127 —, Preservation of, 140 —, Prohibition of sale of, 513, 544 —, Regulations for sale of, 125 —, Standard for, 123 —, Sterilisation of, 141 — sugar, 122, 127 Milkshops. See Dairies. Millstone grit, 170 Milne's hypothesis, 617 Mineral salts in water, 61 Miners, Mortality among, 623, 624 Mines, Control of, 575 —, Refuse from, 536 Minimetric test, 15
—, Control of supplies, 139 — epidemics, 132, 513 —, Examination of, 129 —, Fermentation of, 127 —, Preservation of, 140 —, Prohibition of sale of, 513, 544 —, Regulations for sale of, 125 —, Standard for, 123 —, Sterilisation of, 141 — sugar, 122, 127 Milkshops. See Dairies. Millstone grit, 170 Milne's hypothesis, 617 Mineral salts in water, 61 Miners, Mortality among, 623, 624 Mines, Control of, 575 —, Refuse from, 536

Moisture of air, 20 of soil, 160 Moncrieff's treatment of sewage bacteria, 271 Montgolfier's rule, 191 Morell's ash closet, 250 Mortality and age, 609, 613, 621, 622 — and occupation, 622 --- and season, 620 - and sex, 610, 612 Mortar, 172 Mortuaries, 502, 552, 580, 600 Mosquitoes and yellow fever, 440 - and malaria, 449 Mumps, 215, 360 Munich, Enteric fever at, 167, 383 Municipal Corporations Act, 581 Mussels, Poisoning by, 116, 121 Muzzling dogs for rabies, 425 Mytilotoxine, 117 Natural increase of population, 607 ventilation, 187 Necks of low pressure, 33 Negretti's recording and electric anemometers, 31 Nematoda, 289 Nessler's solution, 78 New streets and buildings, Bye-laws as to, 502, 580, 586 Nightsoil and other refuse, 583 Nitrates and nitrites in soil, 163 - in water, 64 -, Tests for 65 Nitrification, 164 Nitrites in water, 64 Nitrogen in air, 3, 16 —— in water, 60 -, Inorganic, 64 -, Organic, 79 -, Oxidised, 65 Nitrogenous food, 90 Nitrous acid, 329, 333 Notification of Births Act, 1907, 555 Notification, Compulsory, of infectious diseases, 458, 552 - and canal boats, 484, 540 - and common lodging-houses, 484, 494 — and tenement houses, 484, 598 - and vans and tents, 483, 580 - of measles and whooping cough, 461, 462 Nottingnam, Welbeck disease at, Nuisances, 494

-, Byelaws regarding, 508, 578,

-, Procedure in regard to, 496

Oatmeal, 95, 150 Oats, 150 Obligate parasites, 309 Obstructive buildings, 525 Occupation and mortality, 622 Offensive accumulations, 495, 498, 509, 578, 583, 584 — materials, Storage of, 601 - trades, 497, 550, 580, 601 Open fires, 194 - space around houses, 587, 598 — Spaces Act, 581 Opsonins, 305 Optical tests for milk, 130 Organic matter in air, 13 — in soil, 164 — in water, 76 Organisms in air, 12 --- in soil, 163 - in water, 54 Overcrowding, 495 Oxidised nitrogen, 65 Oxygen in air, 3, 16 — in soil, 162 —— in water, 60 — process of water analysis, 81 Oxyuris vermicularis, 289 Oyster culture in relation to disease, 118, 387 Oysters, Poisoning by, 116 Ozone, 3, 16 Pabulum theory of protection, Pail-closets, 250, 252, 253, 254

Pan-closets, 246 Pandemics, 297 Parasites, 308 —, Amœboid, 308 -, Animal, 284 —, Facultative, 309 — in meat, 102, 109 — in water, 55 —, Malarial, 447, 449 —, Obligate, 309 —, Vegetable, 306, 308 —, Worm, 286 Parish Councils, 520 Parotitis, 360 Passages, Cleansing common, 510 Pasteur-Chamberland's filter, 50 Pasteurisation, 141 Pasteur's experiments, Rabies, 423 — solution, 310 Pauper sickness, Returns of, 483 Peas, 95, 150, 152

Pellagra, 150 Pepper starch, 152 Peppermint test, 244 Percolation of water, 37 Perforated bricks as ventilators, Periodicity of epidemics, 297 Peritonitis, Tubercular, 419 Permanganate, Alkaline, in water analysis, 78 -, Potassic, as deodorant, 333 _____, _____, as disinfectant, 327 ______, in water analysis, 76, 77, -, -, Standard solutions of, 78, 81 Persons per inhabited house, Number of, 606 Pertussis. See Whooping Cough. Pestis minor plague, 442 Petri's process, 13 Pettenkofer's method, 13 observations on cholera, 167, - — on enteric fever, 167, 383 Phagocyte theory of protection, Phosphates in water, 67 Phosphorus poisoning, 572 Phthisis, 168, 410, 463, 615, 624 — and dampness of soil, 168, 411 — and locality, 411, 414 - and occupation, 412, 623 Pipes, Rain, 243, 503
—, Waste, 240
—, Water, 72 Plague, 442 - regulations, 470 Plasmodia, 308 Plate cultivations, 310 Playground of schools, 206 Plug closets, 246 Plumbism, 71 Pneumonia, Epidemic, 405 —, Bacillus of, 408, 409 Pneumonic plague, 442 Poisson's formula, 629 Polarite, 267 Pollution of rivers, 40, 80, 264, 510, 536 — of soil, 40, 42, 163 - of water, 40, 42, 84, 89 Population, 605 -, Age distribution of, 607 —, Estimates of, 605 -, Natural increase of, 607 — per house, 606 —, Urban and rural, 607

Population of England and Wales, tion of, 632 - Sanitary Authority, 506 - S.A. and cholera, 470 Porter-Clark process, 48 Portland cement, 172 Potato, 95, 100, 151 -, Microbe cultivation on, 310 --- starch, 152 Pouchet's aëroscope, 12 Power of entry, 497 Precipitation processes, 265 Preservation of eggs, 147 — of fish, 116 — of meat, 110 — of milk, 140 Pressure anemometer, 30 Prevention of Cruelty to Children Act, 576 - of infectious diseases, 457 --- of nuisances, Byelaws as to, 508, 578, 583 Private improvement expenses, - premises, Power of entry upon, 497 - scavenging, Byelaws as to, 492, 549, 578, 582 Privies, 249, 252, 254, 491, 591 -, Byelaws as to, 583 Probable duration of life, 615 Protection in specific diseases, —, Theories of, 301 Protective inoculation, 309, 313 Protozoa, 284 Ptomaines, 104 Public Analyst, 531 —— Health Act (1875), 490 _____, (1896), 507, 508 _____ Acts Amendment Act (1890), 508 --- (1907), 511 - — (Fruit-pickers) Act, 579 - (Interments) Act, 278 --- (London) Act, 549 — (Ports) Act (1896), 508 — (Regulations as to Food) Act (1907), 555 — (Water) Act, 506 — Vaccination, 353 Puerperal fever, 367, 430 Purification of sewage, 264 - of water, 47 Pyrosoma bigeminum, 285 Quadratic mean, 625

Quarantine, Domestic, 469
—, School, 215
Quarries, 496

Rabies, 422 - Order of 1892, 425 Raft theory, 7 Rainfall, 29 Rain-gauge, 29 --- pipes, 243, 503 - water, 39 — —, Disposal of, 260 Ransom's disinfector, 324 Rats, Infection by, 445 Rauschbrand, 308 Refuse, Household, 235, 254, 491, 509, 583, 595
—, Town, 255
—, Trade, 254, 547, 595, 602
Registers of sickness, 464 Registrars of births and deaths, Registration of births, 607 —— of canal boats, 533 - of common lodging -houses, 494 - of dairies and cow-sheds, 541 —— of deaths, 608 —— of slaughter-houses, 504, 510, 580, 594 — of tenement houses, 202 Regnault's hygrometer, 22 Regulations, Power of S.A. to frame, 492, 578, 581 Relapsing fever, 378 Remittent fever, 446 Removal to hospital, 465, 500, 515 - of refuse, 235, 254, 491, 509, 595 Reservoirs, Water, 45 Respiration, 10 Retention theory of protection, Returns of births, 483 — of deaths, 483 - of pauper-sickness, 484 Revaccination, 346, 354 Revolver closets, 248 Revolving purifier, 52 Rhizopods, 284 Rice, 95, 150 - fields, 170 Rickets, 99 Right of entry, 497 Ringworm, 308 River Boards, 537 - water, 40 Rivers, 40

Diam Dellution Committee	AND THE PERSON NAMED IN COLUMN TWO IS NOT THE OWNER.
Rivers Pollution Commission,	Sanitary districts, 490
Standards suggested by, 275	- inspectors, Appointment of,
Commissioners' Report,	552
80	———, Duties of, 487
, of, 40, 80, 264, 510, 536	—— law, 490
— Prevention Act, 536	- purposes, Union of districts
Robinson's anemometer, 30	for, 505
Rochdale system, 250	Saprophytes, 309
Roofs, Requirements as to, 587	Sarcinæ, 306
Rooms, Disinfection of, 328, 433	Sausages and disease, 110
- over privies, cesspools, etc.,	
PAA PAA	Saussure's hygrometer, 22
	Scarlet-fever, 215, 361, 462, 463,
—, Ventilation of, 176, 187, 588,	631, 633, 634
Roseola, Epidemic, 361	——— and milk, 133
Rötheln, 215, 361	—— —— and season, 362
Royal Commission on bovine	— —, Epidemic type of, 363
tuberculosis, 1904, 422	——— in schools, 366
on arsenical poisoning,	———, Isolation of, 267
156	, Microbe of, 364
on sewage disposal, 537	———, Puerperal, 367
on tuberculosis, 1898, 107	, Surgical, 366
——————, 1895, 108, 420	Scavenging, 255, 492, 549, 578, 581,
—— on vaccination, 1896, 351	583
Rural District Councils, 521	
	Schizomycetes, 306
— districts, Urban powers for,	School children, Medical inspec-
505	tion of, 219, 481
- Sanitary Authorities, 490	School grants and closing during
— — Power to frame bye-	epidemics, 209
laws, 506, 581	School hygiene. See Schools.
— Districts, 490	Schools, 203
Rye, 150	—, Closets, 207
	—, Closure of, 208—215
THE RESIDENCE OF THE PARTY OF T	—, Cloak-rooms, 20;
Sago starch, 152	—, Desks, 205
Sale of Horseflesh Act, 535	Dimensions of 202
	—, Dimensions of, 203
of Food and Drugs Acts, 529,	, Handicraft classes for, 207
532, 533	, Infectious disease in, 208,216
Salt meat, 95	—, Inspection of, 220
Salts in diet, 92	——, Lighting, 203
Samples of food and drugs, 532	—, Playground, 206
— of water, 52	—, Powers of S.A. over, 208
Sanarelli's bacillus, 439	—, Ventilation of, 204
Sand filters, 49	—, Warming of, 205
, Permeability of, 162	, Water supply, 207
—, Sites upon, 170	Sclerostoma duodenale, 290
—, Water from, 170	Scott's process, 265, 266
Sandstone permeability, 162, 172	Seurvy, 100
, Sites upon, 170	Seasonal variations of barometer,
, Water from, 62, 170	28
Sanitary anthomitics 400	— of birth-rate, 603
Sanitary authorities, 490 ————————————————————————————————————	— of birth-rate, oos
— combining to appoint	
MOH 501	— of death-rate, 620
M.O.H., 504	—— of death-rate, 620 —— of disease, 363, 619
M.O.H., 504 ————, Default of, 505, 552	—— of death-rate, 620 —— of disease, 363, 619 —— of microbes in air, 9
M.O.H., 504 ————————————————————————————————————	— — of death-rate, 620 — of disease, 363, 619 — of microbes in air, 9 — of rainfall, 29
M.O.H., 504 ———————————————————————————————————	
M.O.H., 504 —, Default of, 505, 552 —, Duties of, as to burial- grounds, 281 —, Port, 506	— of death-rate, 620 — of disease, 363, 619 — of microbes in air, 9 — of rainfall, 29 — of thermometer, 19 Secondary cyclones, 32
M.O.H., 504 —, Default of, 505, 552 —, Duties of, as to burial- grounds, 281 —, Port, 506 —, Powers and duties of.	
M.O.H., 504 —, Default of, 505, 552 —, Duties of, as to burial- grounds, 281 —, Port, 506 —, Powers and duties of.	— of death-rate, 620 — of disease, 363, 619 — of microbes in air, 9 — of rainfall, 29 — of thermometer, 19 Secondary cyclones, 32 Sedgwick's process, 13
M.O.H., 504 —, Default of, 505, 552 —, Duties of, as to burial- grounds, 281 —, Port, 506	— of death-rate, 620 — of disease, 363, 619 — of microbes in air, 9 — of rainfall, 29 — of thermometer, 19 Secondary cyclones, 32

Separate contributory places, 505
— system of sewage, 264, 274
Separated milk, 125
Series, Statistical, Comparative
value of 625
value of, 625
Service pipes, Water, 45
reservoirs, Water, 44
— system, 47
Severity of attack influenced by
age, 300
influenced by epidemic
type, 300
influenced by season, 300
influenced by sex, 300
Sewage bacteria, 57, 263
—, Bacterial treatment of, 271
— Commission Reports, 270
—, Composition of, 258, 261
—, Disposal of, 263, 537
— farms, 269, 271
—, Filtration of, 267
—, Pollution of rivers by, 40, 264,
536
, Precipitation of, 265
—, Value of, 262
Sewer, Definition of, 494
—— gas, 257
— gas, 257 Sewers, 256, 490, 508 —, Air of, 257
, Air of, 257
—, Capacity of, 261
Sex and age-distribution, 609, 632
— and death rates, 610, 612, 631
— and diseases, 363
- and expectation of life, 619
—, Numbers born of each, 607
Chaffeld Small count 200 244 244
Sheffield, Small-pox at, 336,343,344
— system of desks, 206
Shell-fish, 116, 120
Sheringham's ventilator, 188
Ships infected with cholera, 470
- subject to jurisdiction of
S.A., 497
Shone's system, 261
Shop Hours Acts, 576
Sickness, Registers of, 464
Sick-room, Arrangement of, 465
Silicates in water, 64
Silk as clothing, 320
Sillar's process, 267
Simple continued fever, 393
Siphon barometer, 25
Sites, 509, 587, 588
Skim-milk, 122
Slaughter-houses, 503, 510, 580, 594
Sleeping-rooms, 585, 598
———, Analysis of air in, 184
sickness, 454
Slop-closets, 248
Statement of the last of the l

Small-pox, 215, 334, 459, 602, 633, 634 Aërial transmission of, 335 - and vaccination, 340 -, Diagnosis of, 339 -, Hospitals for, 225 - in Gloucester, 335 - in Kilmarnock, 342 - in Sheffield, 335, 343, 347 -, Inoculation of, 317, 340 Smith's, Angus, method, 15 Smoke nuisances, 495 - test, 244 Snow, Clearing away, 583 Soap, 75 Soda-salts in water, 68 Softening of water, 48 Soil, 159 —, Air in, 161 - and disease, 166 -, Bacteria in, 163 -, Composition of, 162 -, Made, 163, 170 -, Moisture of, 160 -, Organic matter in, 164 -pipes, 240 -, Testing of, 244 -, Temperature of, 159 Solids not fat, 123, 125 Sore-throat, Milk-borne, 135 Special drainage districts, 505 Specific diseases, 294, 334 - gravity of milk, 130 Spence's process, 266 Spirilla, 307 -, Pathogenic, 307 Spirits, 153 Spirochætæ, 307 Sponge filters, 49 Spongy-iron filter, 49, 52 Spontaneous origin of specific diseases, 296 Sporocytes, 448 Sporozoa, 284, 447 Spotted fever. See Cerebro-spinal fever. Springs, 39 Stable dwellings, 202 Standards of purity of air, 180, 183 - of sewage and trade effluents, 275 of water, 79, 80, 82, 87 Staphylococci, 306 Starch, 151 Statistical series, Value of, 625 —, M.O.H., 281 Steam disinfection, 321, 325 -, Heating by, 198 in sewers, 508

Sterilisation of Milk, 141 Stoves, Heating by, 197 Streams, Pollution of, 40, 510 Streets, New, 502 —, —, Byelaws as to, 580, 586 Streptococci, 59, 306 Streptococcus scarlatinæ, 364 Streptothrix, 308 Strongylus duodenalis, 290 - filaria, 102 Subsoil drainage, 161, 237 —— irrigation, 263 —— water, 160 —, Disposal of, 260 Successive means, 627 Sulphates of iron, copper, and zinc, as deodorants and disinfectants, 326, 333 - in water, 62 Sulphides in water, 67 Sulphuretted hydrogen in air, 5 --- in water, 71 Sulphurous acid in gas, 5 Sunday-schools, S.A. have no power over, 214 Sunshine recorders, 20 Superheated steam, 322 Susceptibility to disease, Variations in, 298, 300 Suspended matters in air, 3, 7, 10 ——— in water, 51 Swine, Keeping of, 543, 551, 583 Synoptic charts, 32, 34 Tabes mesenterica, 419 Tache cérébrale, 405 Tæniada, 102, 286 Tapeworms, 102, 286 Tapioca starch, 152 Taylor's closet, 251 Telluric diseases, 166 Temperature, Dew-point of, 21, 23 --- of air, 18 — of factories, 563 — of soil, 159 Tenement factory, 557 —— houses, 202 ———, Byelaws as to, 579 598 - workshop, 558 Tent hospitals, 227 Tents and vans, 496, 522, 579 Tetanus, 428 Thermometers, Air, 18

—, Earth, 160

- process, 60

Thrush, 308

Threadworms, 289

Thresh's disinfector 323

Stegomyia fasciata, 440

Tidy's process, 81 Tinned foods, 111 Tobin's ventilator, 188 Toxins, 313 Trade effluents, 41, 256, 274, 536 --- refuse, 254, 546, 595, 602 Trades, Offensive, 497, 551, 580, 601 Traps, 235, 239, 247 Trematoda, 288 Trichina spiralis, 291 Trichinosis, 292 Tropical abscess of liver, Parasites in, 308 -— diseases, 431 Trough closets, 248 Trypanosomes, 285, 454 Tub-closets, 250 Tube wells, 43 Tuberculosis, 106, 410 —— and food, 418 —— and locality, 416 — and lower animals, 417 —— and milk, 129, 418 —, Bacillus of, 415 —, Bovine, 419 — in animals, 106 Tubercular meningitis, 419 — peritonitis, 419 Tumbler closets, 249 Typhoid bacillus in soil, 165 — fever, 382 Typhus, 379 Tyro-toxicon, 146 Unhealthy areas, 523 Unsound food, Seizure of, 499 Urban and rural birth and death rates, 631 - mortality compared, - Authorities, power to make byelaws, 578 - District Councils, 522 — powers for rural districts, 505 —— Sanitary Authorities, 490 —— —— Districts, 490 Urinary calculi, 76 Vaccination, 340 —— Acts, 353 —, Calf, 348 - certificates, Magistrates', 353 ———, Postponement, 354 —, Insusceptibility to, 354 —, Protection by, 349 —, Public, 353 —, Quality of, 345 ---, Report of Royal Commission on, 351 —, Rabies, 424

Transfer (1) -1 100	W. F. M. W. 400
Vaccine, Cholera, 436	Water in soil, 160
Value of sewage, 262	— in tissues, 92
—— of statistical series, 625	—, Organisms in, 54
Valve closets, 246	, Purification of, 47
Van Overbeek de Meyer's disin-	, Physical characters of, 53
fector, 323	—— pipes, 72
Vans, Inhabited, 494, 522, 579	, Rain, 39
Varicella, 215, 355	, River, 40
Variolation, 317, 340	, Storage of, 44
Vegetable dietary, 98	— supplies, 42, 47, 179, 207, 492,
—— foods, 147	506, 539
—— matter in soil, 164	
———— in water, 88	yards, 279
Vegetation, effect on climate, 159	-, Suspended matters in, 54
Ventilation, 179	Watercress, 150
—, Artificial, 193	Webster's process, 267
— beneath floors, 173, 588	"Wedges" of high pressure, 33
—, Haldane and Lorrain Smith's	
	Welbeck disease, 102
experiments, 182	Wells, 39, 42
—, Mechanical, 193	—, Closure of, 493
—, Natural, 187, 194	—, Dumb, 263
— of drains, 239, 589 [561	Werner-Schmidt's method, 130
- of factories and workshops,	Wheat flour, 147
— of hospitals, 223	Wheel barometer, 25
— of new buildings; 588	Whisky, definition of, 157
— of rooms, 176, 187, 588, 598	Whitthread's process, 266
—— of schools, 204	Whooping cough, 215, 359, 462,
— of sewers, 259	623, 634
Ventilators, 188	Widal's application of Grüber's
Verdet, 150	reaction, 392
Vernier scale, 26	Wimbledon, Scarlet-fever at, 138
Vessels infected with cholera, 470	Winchester quart, 52
Vibriones, 307	Wind, 30
Virus, Attenuation of, 312	Windows, 588
Vital statistics, 605	Windrose, 32
V-shaped depressions, 32	Wooden hospitals, 226
, samped depression,	Woollen clothing, 320
Walls, Construction of, 172, 175,	Woolsorters' disease, 426
587	Work, Mechanical, 97
Wanklyn's ammonia process, 77	Working Classes, Housing of,
Warming, Methods of, 195	Acts, 522, 523, 527, 528
— of Schools, 205	Workmen's Compensation Act,
Wash-out closet, 246	1906, 576
Washington Lyon's disinfector, 322	Workshops. See Factories.
	Worms, Parasitic, 286
Waste-pipes, 240	Worthing, Outbreak, 385
Water, 37	Worthing, Outbreak, 500
—, Analysis of, 76	77 3 - 63-3-in- houses 500
— and disease, 88, 384, 434	Yards of lodging-houses, 598
— as food, 92	Yellow fever, 437
—, Bacteria in, 55	— , Bacteriology of, 440
barometer, 28	— Regulations, 470
closets, 245, 590	
—, Collection of, 44	Zinc salts as disinfectants, 326
—, Dissolved matter in, 53, 84	in water, 74
—, Examination of, 52, 59, 84	Zooglos 306
—, Gases in, 60	Zooglea, 306
, Good and bad, 87	Zygote, 448 Zymotic death-rate, 612
—— in air, 17	Zymoric death-rate, ora

Printed by Cassell & Company, Limited, La Belle Sauvage, London, E.C. 10.4.08



