

Landmarks and surface markings of the human body / by Louis Bathe Rawling.

Contributors

Rawling, L. Bathe
Royal College of Physicians of Edinburgh

Publication/Creation

London : H.K. Lewis, 1904.

Persistent URL

<https://wellcomecollection.org/works/jbbyyc65>

Provider

Royal College of Physicians Edinburgh

License and attribution

This material has been provided by This material has been provided by the Royal College of Physicians of Edinburgh. The original may be consulted at the Royal College of Physicians of Edinburgh. where the originals may be consulted.

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).

LANDMARKS
AND
SURFACE MARKINGS
OF THE
HUMAN BODY

R.C.P. EDINBURGH LIBRARY



R27732A0236

*y order of the College, this Book is not to be taken out
the Library (except after 6 P.M. until 10 A.M.) for one
month from this date.*

PHYSICIANS' HALL,

8th Dec 190 *4*

Sprint

T. 5. 18.

LANDMARKS AND SURFACE MARKINGS
OF THE HUMAN BODY

THE HISTORY OF THE
ROYAL NAVY

LANDMARKS AND
SURFACE MARKINGS OF
THE HUMAN BODY

BY

LOUIS BATHE RAWLING

M.B., B.C. (CANT.), F.R.C.S. (ENG.)

SENIOR DEMONSTRATOR OF ANATOMY, ST. BARTHOLOMEW'S HOSPITAL; ASSISTANT-
SURGEON TO THE GERMAN HOSPITAL, DALSTON; HUNTERIAN PROFESSOR,
ROYAL COLLEGE OF SURGEONS, ENGLAND




WITH TWENTY-NINE ILLUSTRATIONS

LONDON
H. K. LEWIS, 136, GOWER STREET

1904

LANDMARKS AND
SURFACE MARKINGS OF
THE HUMAN BODY



LONDON
H. K. LEWIS, 65 GOWER STREET
1904

PREFACE

THIS book is intended to fill a gap in the series of textbooks that are offered to the medical student. No apology is needed for the introduction of such a work, as the author, during several years of hospital and of private teaching, has fully realized the urgent want of a handy book which describes and illustrates the more important landmarks and surface markings of the human body. To the student in anatomy such a book is a necessity, as the ordinary "dryness" of his work is mitigated by the realization of the importance of the structures which are here alluded to, and to the student of medicine and of surgery the knowledge gained is of obvious practical utility.

The writer has endeavoured to include all the more important landmarks and surface markings, and the details given are derived partly from personal experience and partly from references to the best-known anatomical, medical, and surgical text-books.

Any omissions will be repaired in a future edition if, as the author trusts, this work should meet with a favourable reception. The writer desires to tender his best

thanks to his colleague, Mr. S. R. Scott, for many valuable suggestions, and to Mr. Frank Butterworth, of the Savoy Buildings, Strand, who is responsible for the reproduction of all the illustrations, and who has taken infinite pains to follow out the desires and suggestions of the author.

LOUIS BATHE RAWLING.

16, MONTAGU STREET,
PORTMAN SQUARE, W.

October, 1904.

CONTENTS

	CHAPTER I	PAGE
THE HEAD AND NECK	- - - - -	I
	CHAPTER II	
THE UPPER EXTREMITY	- - - - -	15
	CHAPTER III	
THE THORAX	- - - - -	28
	CHAPTER IV	
THE ABDOMEN	- - - - -	40
	CHAPTER V	
THE LOWER EXTREMITY	- - - - -	57
	APPENDIX	
THE LENGTH OF THE VARIOUS PASSAGES, TUBES, ETC.	- - - - -	70
THE WEIGHT OF SOME ORGANS	- - - - -	71
INDEX	- - - - -	73

CONTENTS

CHAPTER I	THE HISTORY OF THE
CHAPTER II	THE HISTORY OF THE
CHAPTER III	THE HISTORY OF THE
CHAPTER IV	THE HISTORY OF THE
CHAPTER V	THE HISTORY OF THE
CHAPTER VI	THE HISTORY OF THE
CHAPTER VII	THE HISTORY OF THE
CHAPTER VIII	THE HISTORY OF THE
CHAPTER IX	THE HISTORY OF THE
CHAPTER X	THE HISTORY OF THE
APPENDIX	THE HISTORY OF THE
INDEX	THE HISTORY OF THE

LANDMARKS AND SURFACE MARKINGS OF THE HUMAN BODY

CHAPTER I

THE HEAD AND NECK

CRANIO-CEREBRAL TOPOGRAPHY.

ONLY those surface markings will be given which are of practical value, and as far as possible each landmark will be rendered independent of any other, as by such means any given structure can be rapidly depicted on the surface, the important question of time and of space rendering the more complicated systems, in which it is necessary to map out a network of intersecting lines in order to fix the position of any single structure, of little surgical value. It is necessary, however, to recognise first certain important bony points, etc.

1. The *nasion*, situated at the base of the nose at the central point of the naso-frontal suture.

Fig. i., 1.

2. The *inion*, or external occipital protuberance.

Fig. i., 2.

A line uniting these two points over the vertex of the skull corresponds in position to the upper attached margin of the falx cerebri, to the mesial longitudinal fissure of the brain, and to the superior longi-

tudinal sinus, which, though narrow in front, broadens out rapidly as it passes backwards towards the internal occipital protuberance.

This line also corresponds in direction to the occasionally persistent metopic suture between the two halves of the frontal bone, and to the sagittal suture between the two parietal bones.

At the junction of the sagittal and coronal sutures is the *bregma*, the site of the foetal anterior fontanelle, which opening is usually closed by the end of the second year, and at the junction of the sagittal and lambdoid sutures the posterior fontanelle is situated, closed at or soon after birth. The point of junction of these two latter sutures is known as the *lambda*, and it is situated usually about $2\frac{1}{2}$ inches above the inion or external occipital protuberance.

3. The *external angular frontal process* articulates with the frontal process of the malar bone, and the articulation between the two processes is easily felt at the upper and outer border of the orbital cavity.


4. The *malar tubercle*, a small prominence to be felt along the posterior border of the frontal process of the malar bone, a short distance below the fronto-malar suture.

5. The *zygomatic process* should then be traced backwards towards the ear, and an examination of the skull will show that the zygoma in front of the ear divides into three roots, the anterior of which merges into the eminentia articularis, whilst the middle helps in the formation of the post-glenoid process. The upper root can, on deep palpation, be traced backwards above the external auditory meatus, where it becomes continuous with the *supra-mastoid crest*, a prominent ridge at the upper and anterior part of the base of the mastoid

Fig. ii., 16.

Fig. i., 12.

Fig. ii., 13.






FIG. I.




FIG. II.


To face p. 3.

FIG. I.

1. The nasion.
2. The inion.
3. The mid-point between nasion and inion.
4. The Rolandic fissure.
5. The superior temporal crest.
6. The inferior temporal crest.
7. The Sylvian point.
8. The anterior horizontal limb of the Sylvian fissure.
9. The vertical limb of the Sylvian fissure.
10. The posterior horizontal limb of the Sylvian fissure.
11. The parietal prominence.
12. The malar tubercle.
13. The lambda.
14. The first temporo-sphenoidal sulcus.
15. The external parieto-occipital sulcus.
16. The lateral sinus.
- 17, 17, 17. The level of the base of the cerebrum.
18. The mastoid process.
- 19, 19. Reid's base line.

FIG. II.

- 1, 1. Reid's base line.
- 2, 2. A line parallel to the above at the level of the supra-orbital margin.
3. The middle meningeal artery.
4. The anterior branch.
- 5, 5, 5. The three sites for trephination.
6. The posterior branch.
7. The site for trephination.
8. The point for trephining to reach the descending horn of the lateral ventricle.
9. The lateral sinus.
10. The inion.
11. The mastoid process.
12. Macewen's suprameatal triangle.
- 12*a*. The mastoid antrum.
- 12*b*. The facial nerve.
13. The supramastoid crest.
- 14, 14. The temporal crest.
15. The temporal fossa.
16. The external angular frontal process.
17. The tendo-oculi attachment.
18. The lachrymal groove.



process, which becomes continuous behind with the posterior part of the temporal crest. This supramastoid crest is of great surgical value, insomuch that it forms the upper boundary of Macewen's suprameatal triangle, and also corresponds to the level of the base of the brain in this situation.

The *lateral sinus* describes a gentle curve from the external occipital protuberance to the upper and posterior part of the base of the mastoid process, curving finally downwards towards, though not reaching to, the apex of that process. The degree of upward curve is subject to a good deal of variation, and the sinus frequently presents a lesser upward convexity than was the case in the specimen from which the illustration was taken.

The *lower limit of the cerebrum* can, with sufficient accuracy, be mapped out in the following manner: A point is taken in the median antero-posterior line about $\frac{1}{2}$ inch above the nasion, and from this point a line is drawn outwards which lies about $\frac{1}{2}$ inch above, and follows the curve of the upper border of the orbit. This line is carried backwards as far as the level of the external angular frontal process, then being directed downwards and backwards towards the midpoint of the zygoma, and running backwards parallel with the upper border of that process to the level of the external auditory meatus. At and behind the ear the brain lies flush with the supramastoid crest, and subsequently follows the curve of the lateral sinus from the base of the mastoid process to the external occipital protuberance.

The curve already mapped out as formed by the lateral sinus also represents, therefore, the lower level of the

posterior part of the cerebrum, and it further corresponds to the outer attachment of the tentorium cerebelli which assists in the formation of the lateral sinus. Below the curve the cerebellum lies.

Reid's base line is drawn backwards from the lower border of the orbit to the middle of the external auditory meatus, and when further produced the line will be found to fall just below the level of the inion, and to lie almost entirely below the level of the lateral sinus. This line is utilized by some surgeons in trephination of the skull, distances being measured along this line and points taken above or below, according to the seat of the lesion.

The *Sylvian point* represents the site of divergence of the three limbs of the Sylvian fissure. It lies $1\frac{1}{4}$ inches behind the external angular frontal process, and $1\frac{1}{2}$ inches above the upper border of the zygoma. The main posterior horizontal limb of the Sylvian fissure passes backwards and upwards from the Sylvian point to a second point situated $\frac{3}{4}$ inch below the most prominent part of the parietal bone.

The vertical limb is directed upwards for about $\frac{3}{4}$ inch, whilst the anterior horizontal limb passes forwards for about the same distance.

The Sylvian point corresponds also to the anterior pole of the island of Reil and to the middle cerebral artery, as that vessel lies deeply embedded in the anterior part of the Sylvian fissure.

To mark out the *external parieto-occipital* fissure and the first *temporo-sphenoidal* or parallel sulcus, it is necessary to find two bony points—the malar tubercle and the lambda. A line uniting these two points

corresponds in its posterior part to the external parieto-occipital sulcus, and in its middle third to the line of the first temporo-sphenoidal sulcus.

The *Rolandic Fissure*.—Take a point $\frac{1}{2}$ inch behind the centre of a line drawn across the vertex of the skull from the nasion to the inion, and from this point draw a line downwards and forwards for $3\frac{1}{2}$ to 4 inches at an angle of $67\frac{1}{2}^\circ$ (three-quarters of a right-angle) to the median antero-posterior line. In front of this sulcus is the precentral or ascending frontal convolution, an area better known as the Rolandic or motor area. The main centres here situated correspond from above downwards to the movements of the leg, arm, and face of the opposite side of the body.

The superior temporal crest cuts across the Rolandic line at the junction of its lower and middle thirds, and it therefore roughly separates the arm and face areas one from the other.

On the left side of the head that part of the brain which is included in the obtuse angle between the anterior and posterior horizontal limbs of the Sylvian fissure is known as Broca's area (the speech centre).

The *middle meningeal artery*, a branch of the internal maxillary, enters the skull through the foramen spinosum, and divides, after a short and variable course across the middle fossa of the skull, into two main trunks. The seat of bifurcation usually corresponds to a point just above the centre of the zygoma.

The anterior branch is not only the larger of the two, but it is also more liable to injury, since it is protected in the temporal region by a comparatively thin osseous barrier.

Fig. ii., 5,
5, 5. The danger zone in the course of this branch may be mapped out by taking three points :

(1) 1 inch behind the external angular frontal process, and 1 inch above the zygoma.

(2) $1\frac{1}{2}$ inches behind the external angular frontal process, and $1\frac{1}{2}$ inches above the zygoma.

(3) 2 inches behind the external angular frontal process, and 2 inches above the zygoma.

A line uniting these three points indicates, therefore, that part of the anterior division of the middle meningeal artery which is most liable to injury.

The anterior division of the vessel will be exposed by trephining over any of these three points, but it is generally preferable to choose the highest point, as by such means the outer part of the lesser wing of the sphenoid is avoided ; and, as an additional reason, it should be added that in the position of points 1 and 2 the artery frequently runs in an osseous canal. After trephining over the upper point the bone can be chipped away in a downward and forward direction, if such an exposure of the artery is deemed necessary.

The *posterior branch of the artery* passes almost horizontally backwards, parallel to the zygoma and to the supramastoid crest, and it can be exposed by trephining over a point where a vertical line drawn upwards from the posterior border of the mastoid process

Fig. ii., 6.
Fig. ii., 7. cuts another line drawn backwards from the supra-orbital margin parallel to Reid's base line.

The *Lateral Ventricles*.—The descending cornu of the lateral ventricle may be tapped by trephining
Fig. ii., 8. $1\frac{1}{4}$ inches above Reid's base line and $1\frac{1}{4}$ inches behind the external auditory meatus. The trocar should

be directed towards the summit of the opposite ear, the ventricle being reached within 2 inches from the surface (Keen).

The *Basic Fossæ*.—"There is no external sign to indicate the situation of the fossæ of the skull. In general, however, it may be said that the anterior fossa extends as far back as the anterior end of the zygoma; that the middle fossa lies between this and the mastoid process, and the posterior includes all the base behind the process" (Eisendrath).

The *mastoid antrum* may be exposed by trephining Fig. ii., 12a. in Macewen's suprameatal triangle, a space Fig. ii., 12. which is bounded above by the backward continuation of the upper root of the zygoma (the supramastoid crest), behind by a vertical line drawn upwards from the posterior border of the external auditory meatus, and below and in front by the suprameatal spine, a prominent bony process which assists in the formation of the posterior superior quadrant of the external auditory meatus. In this triangle there is usually a well-marked depression, the suprameatal fossa. The supramastoid crest not only indicates the uppermost possible limit of the mastoid antrum, but, as has already been stated, it corresponds also to the level of the base of brain in this situation. Fig. ii., 13. The crest, therefore, represents the level of the tegmen antri, and in mastoid explorations the scene of operation must be confined to an area below this crest. In the adult the antrum usually lies at a depth of $\frac{1}{2}$ to $\frac{3}{4}$ inch from the surface.

The lateral sinus lies posterior and nearer to the Fig. ii., 9. surface, whilst the facial nerve pursues its Fig. ii., 12b. course in front and on a deeper plane.

The *frontal sinuses* are very variable in extent. They

occupy the space between the inner and outer tables of the frontal bone, above the base of the nose and above the inner half of the supra-orbital margin. The duct opens into the middle meatus of the nose, under cover of the middle turbinated bone, on a level with the inner part of the palpebral fissure.


The *parotid gland* occupies the space which is bounded
 Fig. iii., 5. above by the zygomatic arch, behind by the auricle and the mastoid process, and below by a line drawn from the angle of the jaw to the apex of the mastoid process. In front the gland extends a variable distance over the anterior surface of the masseter muscle.

Stenson's duct, the duct of the parotid gland, corresponds
 Fig. iii., 7. to the middle third of a line drawn from the lower border of the tragus of the ear to a point situated halfway between the ala of the nose and the red line of the upper lip. At the anterior border of the masseter muscle the duct dips inwards, through the buccinator muscle, to open on the buccal mucous membrane, opposite the second molar tooth of the upper jaw.

The *transverse facial artery*, a branch of the superficial
 Fig. iii., 6. temporal, runs inwards parallel to and immediately below the zygoma, lying above the level of Stenson's duct.

The *facial nerve*, after emerging from the stylo-mastoid
 Fig. iii., 8. foramen, curls round the condyle of the jaw, and traverses the substance of the parotid gland, in which part of its course it divides into numerous branches. The general trans-parotid course of the nerve and the direction of its buccal branch may be indicated by a line drawn forwards parallel to and below Stenson's duct from the lobule of the ear.

The *Tendo Oculi and the Nasal Duct*.—By alternate






FIG. III.

- | | |
|----------------------------------|---|
| 1. The supra-orbital foramen. | 11. The internal carotid artery. |
| 2. The infra-orbital foramen. | 12. The superior thyroid artery. |
| 3. The mental foramen. | 13. The lingual artery. |
| 4. The zygoma. | 14. The occipital artery. |
| 5. The parotid gland. | 15. The internal jugular vein. |
| 6. The transverse facial artery. | 16. The external jugular vein. |
| 7. Stenson's duct. | 17. The upper limit of the brachial plexus. |
| 8. The facial nerve. | 18. The subclavian artery. |
| 9. The common carotid artery. | 19. The clavicle. |
| 10. The external carotid artery. | |

To face p. 9.

forcible closure and opening of the lids the internal tarsal ligament, or tendo oculi, can be felt passing to its insertion into the nasal process of the superior maxilla. Immediately below the tendon, at the junction of the inner and lower walls of the orbital cavity, is the depression for the lachrymal sac, which sac narrows below into the *nasal duct*. The duct passes downwards and slightly outwards to open into the anterior part of the inferior meatus of the nose under cover of the inferior turbinated bone. The duct is about $\frac{1}{2}$ inch long.

The *Supra-orbital, Infra-orbital and Mental Foramina*.—

At the junction of the inner and middle thirds of the supra-orbital margin the supra-orbital notch or foramen may be felt, and a line drawn downwards from this foramen through the interval between the two lower bicuspid teeth will pass through both infra-orbital and mental foramina. The former foramen lies $\frac{1}{4}$ to $\frac{1}{2}$ inch below the orbital margin, whilst the latter (in the adult) lies midway between the alveolar and inferior borders of the lower jaw.

The *facial artery* curls round the inferior border of the lower jaw at the anterior margin of the masseter muscle, about $1\frac{1}{2}$ inches in front of the angle. The artery then passes upwards and inwards in a tortuous manner towards the inner canthus of the eye, there terminating as the “angular artery.” The *facial vein* has a similar general direction. It lies, however, posterior to the artery, and it does not pursue a tortuous course.

The *inferior dental nerve* may be exposed by trephining over the ascending ramus of the lower jaw, midway between the anterior and posterior borders, and on a level with the last molar tooth. In

this manner the nerve, accompanied by the corresponding artery, is exposed as it enters the inferior dental canal.

The *tonsil* corresponds in position to a point situated
 Fig. iv., 3. just above and in front of the angle of the
 lower jaw.

The *antrum of Highmore* usually occupies the greater part of the interior of the body of the superior maxilla, and it opens into the middle meatus of the nose under cover of the middle turbinated bone.

The *occipital artery* and the *great occipital nerve* turn upwards on to the scalp about $1\frac{1}{2}$ inches external to the external occipital protuberance.

The *superficial temporal artery* and the *auriculo-temporal nerve* cross the base of the zygomatic process of the temporal bone immediately above and in front of the tragus of the ear.

The *posterior auricular artery and nerve* pass upwards on to the scalp after lying deeply placed in the hollow between the ear and the mastoid process.

The *external jugular vein* is formed just behind the angle
 Fig. iii., 16. of the jaw by the junction of the posterior division of the temporo-maxillary trunk with the posterior auricular vein. The vessel so formed passes downwards and backwards superficial to the sternomastoid muscle towards the middle of the clavicle, above which bone the vein pierces the deep fascia to join the subclavian vein.

The *carotid arteries* correspond in direction to a line
 Fig. iii., 9. from the sterno-clavicular joint to the hollow between the angle of the jaw and the mastoid process. The *common carotid* usually bifurcates at the
 Fig. iii., 10, level of the upper border of the thyroid cartilage (fourth cervical vertebra), the external
 11. carotid subsequently lying superficial to and slightly to the

inner side of the internal carotid. The *omo-hyoid* muscle (upper belly) crosses the common carotid at the level of the cricoid cartilage, and in this situation the artery may be compressed against the prominent anterior tubercle of the transverse process of the sixth cervical vertebra. At the same level the larynx and pharynx end, and the trachea and œsophagus begin.

The *superior thyroid artery* arises from the external carotid immediately above the level of the upper border of the thyroid cartilage; the *lingual* midway between the thyroid cartilage and the great cornu of the hyoid bone; whilst the *facial* arises just below the level of the great cornu of the hyoid. All these vessels are given off in the carotid triangle, the superior thyroid turning down to the apex of the lateral lobe of the thyroid gland; the lingual running upwards to gain the upper aspect of the hyoid, with which bone it runs parallel for a short distance inwards towards the middle line; whilst the facial artery is directed forwards and upwards through the submaxillary gland to a point about $1\frac{1}{2}$ inches in front of the angle of the jaw.

The *occipital artery* arises from the outer aspect of the external carotid just below the level of the great cornu of the hyoid, and it passes backwards and upwards towards the base of the mastoid process.

The *internal jugular vein* runs parallel and external to the internal and common carotid arteries, and it therefore presents a similar surface marking to that already given for those arteries.

The *vagus nerve* passes downwards in the carotid sheath behind and between the carotid arteries and the internal jugular vein.

The *cervical sympathetic trunk* also lies in the line of the

carotid arteries, being placed behind the carotid sheath. The *superior cervical ganglion* is situated in front of the transverse processes of the second and third cervical vertebræ; the *middle ganglion* overlies the corresponding process of the sixth vertebra; whilst the *inferior ganglion*, which is frequently fused with the first thoracic, lies behind the first part of the subclavian artery between the transverse process of the seventh cervical vertebra and the neck of the first rib.

The *phrenic nerve* is formed below the level of the hyoid bone, and it passes downwards and slightly inwards towards the sternal end of the clavicle. At the level of the cricoid cartilage the nerve lies midway between the anterior and posterior borders of the sterno-mastoid muscle.

The *spinal accessory nerve* crosses the transverse process of the atlas, a bony prominence to be felt immediately below and in front of the apex of the mastoid process. The nerve now enters the substance of the sterno-mastoid at the junction of the upper and second quarters along the anterior border of the muscle, emerging from the posterior border of the muscle at the junction of the upper and middle thirds. The point of emergence is, however, subject to some variation, and the nerve may enter the posterior triangle of the neck at a somewhat lower level, pursuing subsequently a downward and backward course towards the anterior border of the trapezius muscle, beneath which muscle it sinks.

The *Superficial Cervical Plexus*.—Take a point midway along the posterior border of the sterno-mastoid muscle, and from this point draw three lines :

1. Upwards towards the lobe of the ear = the great auricular nerve.





FIG. IV.

- | | |
|--|---|
| 1. The inferior maxilla. | 8. The third part of the subclavian artery. |
| 2. The inferior dental nerve. | 9. The clavicle. |
| 3. The tonsil. | 10. The clavicular head of the sterno-mastoid muscle. |
| 4. The transverse process of the atlas. | 11. The sternal head of the sterno-mastoid muscle. |
| 5, 5. The spinal accessory nerve. | |
| 6. The sterno-mastoid muscle. | |
| 7. The upper limit of the brachial plexus. | |

To face p. 12.



2. Upwards along the posterior border of the sterno-mastoid muscle = small occipital nerve.

3. Forwards towards the middle line of the neck = the transverse cervical nerve.

By producing these three lines in a downward direction, the descending branches of the plexus are roughly indicated.

Thus the great auricular produced = the supraclavicular nerve; the small occipital produced = the suprasternal nerve; the transverse cervical produced = the supra-acromial nerve.

The *subclavian artery* (cervical course) is represented by a curved line from the sterno-clavicular joint to the mid-point of the corresponding clavicle, the convexity of the line extending upwards into the supraclavicular fossa about $\frac{3}{4}$ to 1 inch above the clavicle. In marking out this vessel the shoulders should be well depressed.

The artery passes behind the scalenus anticus muscle, the second part of the artery being covered by that muscle. The outer border of the scalene muscle usually corresponds to the outer border of the sterno-mastoid muscle, and consequently the third part of the subclavian artery is represented by that part of the curve which lies between the outer border of the sterno-mastoid muscle and the mid-point of the clavicle.

The subclavian artery ends anatomically at the outer border of the first rib.

The *Brachial Plexus*.—The upper limit of the nerve-trunks which form this plexus is represented by a line drawn from the mid-point between the anterior and posterior borders of the sterno-mastoid muscle at the level of the cricoid cartilage to a second

Fig. iii., 18.
Fig. iv., 8.
Fig. v., 14.

Fig. iv., 8.

Fig. iii., 17.
Fig. iv., 7.

point situated just external to the mid-point of the clavicle. The lowest cord lies behind the third part of the sub-clavian artery.

The *rima glottidis*, which in its front part is laterally bounded by the true vocal cords, lies opposite the mid-point along the anterior border of the thyroid cartilage.

The *epiglottis*, though fixed below to the thyroid angle immediately above the point of attachment of the true vocal cords, extends upwards to above the level of the body of the hyoid bone.

A suicidal cut throat generally involves the thyro-hyoid space, and if of sufficient depth the epiglottis is severed from its thyroid attachment.

The *isthmus of the thyroid gland* in the adult crosses the trachea about $\frac{1}{2}$ to $\frac{3}{4}$ inch below the cricoid cartilage.

The structures in the *middle line of the neck* are sufficiently indicated in the diagram as to require no special enumeration.





FIG. V.

- | | |
|-------------------------------|---|
| 1. The supra-orbital foramen. | 9. The trachea. |
| 2. The infra-orbital foramen. | 10. The thyroid isthmus. |
| 3. The mental foramen. | 11. The phrenic nerve. |
| 4. The hyoid bone. | 12. The upper limit of the brachial plexus. |
| 5. The thyroid cartilage. | 13. The common carotid artery. |
| 6. The epiglottis. | 14. The subclavian artery. |
| 7. The rima glottidis. | |
| 8. The cricoid cartilage. | |

To face p. 14.



CHAPTER II

THE UPPER EXTREMITY

THE SHOULDER REGION.

AT the acromial end of the clavicle there is usually a slight elevation, and immediately external to this is the acromio-clavicular joint, the long axis of which lies in the antero-posterior direction. The acromial spine and process can both be palpated throughout their entire length. The prominence of the shoulder results from the outward displacement of the deltoid muscle by the underlying head and great tuberosity of the humerus. This normal outward displacement of the deltoid muscle is taken advantage of by Hamilton in the diagnosis of a dislocation of the shoulder-joint. The resultant inward displacement of the head of the humerus after such an injury leads to such flattening of the shoulder that a ruler placed alongside the outer part of the arm will be in contact with the acromial process and with the external condyle of the humerus at one and the same time. In the normal condition this is not possible. To measure the length of the humerus the tape measure should be carried between the same two points. About 1 inch below the outer part of the clavicle the coracoid process of the scapula may be felt lying under cover of the anterior border of the deltoid muscle. The lesser tuberosity of the humerus is not easily felt, but it can usually be distinguished lying rather below the mid-point of a line uniting the

Fig. x., 1, 2.

Fig. vi., 3.

coracoid process and the great tuberosity of the humerus, the arm being well everted and the elbow in contact with the side of the body.

Between the two tuberosities is the bicipital groove, in which runs the long tendon of the biceps muscle. The groove may be represented by a line, about 2 inches long, which runs downwards from the tip of the acromial process, parallel to the long axis of the humerus.

The upper epiphysis of the humerus includes the head and both tuberosities, and the epiphysial line runs transversely at right angles to the long axis of the humerus at the lower border of the great tuberosity.

The *anterior axillary wall* is formed by the major and minor pectoral muscles, whilst the anterior fold of this space is due to the former muscle only.

The posterior wall is formed mainly by the subscapularis muscle overlying the ventral surface of the scapula, the axillary border of which bone can be made out by deep palpation from the inferior angle below to the head above.

The rounded posterior fold of the axilla results from the fact that the latissimus dorsi curls round the teres major muscle from behind forwards in order to reach its insertion into the floor of the bicipital groove. The narrow outer boundary of the axilla corresponds to the upper part of the shaft and to the head of the humerus, and in this situation, in a well-developed

arm, two prominent longitudinal folds are seen, the anterior of which corresponds to the coracobrachialis and biceps (short head) muscles, whilst the more

posterior fold results from the projection of the neuro-vascular bundle. To examine the apex of the axilla, the elbow should be supported, and the patient instructed to relax all muscles. The head of the

Fig. vi., 8.

Fig. x., 7.

Fig. vi., 10.

Fig. vi., 11.






FIG. VI.

- | | | |
|--|---|--|
| 1. The subclavian artery. | 8. The anterior axillary fold. | 15. The level of the head of the radius. |
| 2. The clavicle. | 9. The musculo-cutaneous nerve. | 16. The radial nerve. |
| 3. The coracoid process. | 10. The coraco-brachialis prominence. | 17. The radial artery. |
| 4. The pectoralis minor. | 11. The median nerve, upper third. | 18. The ulnar artery. |
| 5. The axillary artery. | 12. The brachial artery. | 19. The median nerve in the forearm. |
| 6. The anterior border of the deltoid. | 13. The ulnar nerve. | 20. The ulnar nerve in the forearm. |
| 7. The circumflex nerve and arteries. | 14. The median nerve, lower third (middle third unrepresented). | |

To face p. 16.








FIG. VII.

1. The external condyle. 2. The internal condyle.
3. The olecranon process.

To face p. 17.

humerus and of the scapula can then be felt at the upper and posterior part of the axilla, the second rib on the inner side, and the coracoid process in front. The head of the humerus looks in the same direction as the internal condyle of the humerus. The neuro-vascular bundle can be traced upwards as far as the second rib. The inner wall of the axilla is formed by the upper part of the lateral wall of the thorax, which is here clothed by the serrations of the serratus magnus muscle.

The *Elbow Region*.—When the forearm is extended, a line joining the internal and external condyles of the humerus cuts across the tip of the olecranon process, which bony prominence lies well to the inner side of the mid-point of the intercondyloid line. When the forearm is flexed the olecranon moves downwards, and by uniting the three bony points a triangle is formed.

Immediately below the external condyle the head of the radius is felt "lying in the valley behind the supinator longus" (Holden). The humero-radial articulation is transverse, but the humero-ulnar articulation slopes obliquely downwards and inwards, and consequently, whilst the external condyle is about $\frac{3}{4}$ inch above the humero-radial joint, the internal condyle lies rather more than 1 inch above the line of the humero-ulnar articulation.

The junction of the diaphysis and lower epiphysis of the humerus corresponds to a transverse line drawn across the humerus immediately above the tips of the condyles. The bony points on the outer side of the joint are generally obscured in those cases where there is considerable effusion into the elbow-joint, the synovial membrane bulging outwards below the external condyle of the humerus and between that process and the olecranon process. Under

similar conditions there is also an outward projection of the synovial membrane between the olecranon and the internal condyle of the humerus, obscuring the deep depression that normally exists in that situation—a depression at the base of which the ulnar nerve can be rolled beneath the finger.

In front of the elbow is the *antecubital triangle*, the base of which corresponds to a line drawn across the front of the elbow between the two humeral condyles, whilst the inner and outer boundaries are formed respectively by the pronator radii teres and supinator longus muscles. This triangular space is vertically subdivided by the biceps tendon, on either side of which a depression exists, the inner and outer bicipital sulci. In the outer sulcus the musculo-spiral nerve divides into its two terminal branches, radial and posterior interosseous, whilst the median nerve and the brachial artery lie in the inner sulcus, the artery intermediate between the tendon and the nerve. The artery and nerve are, however, obscured in the lower part of the inner sulcus by the overlying bicipital fascia, which can be traced inwards to the pronator region, whilst its upper free margin presents a well-marked crescentic edge which looks upwards and inwards.

The *superficial veins* in front of the bend of the elbow are arranged in the form of a letter **M**, the radial, median, and ulnar veins being received from below, whilst two main efferent vessels, the cephalic and the basilic, carry the blood upwards. The *basilic vein* passes upwards in the superficial fascia, along the inner side of the arm, and pierces the deep fascia about half-way between the axilla and the



FIG. VIII.

- | | |
|------------------------------|--------------------------|
| 1. The intercondyloid line. | 5. The biceps. |
| 2. The supinator longus. | 6. The bicipital fascia. |
| 3. The pronator radii teres. | 7. The brachial artery. |
| 4. The musculo-spiral nerve. | 8. The median nerve. |





FIG. IX.

1. The radial vein.
2. The median vein.
3. The median-cephalic vein.
4. The cephalic vein.
5. The point at which the cephalic vein pierces the costocoracoid membrane.
6. The median-basilic vein.
7. The anterior ulnar vein.
8. The posterior ulnar vein.
9. The basilic vein.
10. The point at which the basilic vein pierces the deep fascia.
11. The epitrochlear gland.



internal condyle, and at the foramen so produced in the deep fascia the internal cutaneous nerve emerges, to become superficial. The *epitrochlear gland* lies in close connection with the median basilic or basilic veins above and in front of the internal condyle.

The *cephalic vein* can be traced upwards along the outer side of the arm as far as the groove between the deltoid and pectoralis major muscles. In the interval between these two muscles the vein lies embedded, and it eventually pierces the costo-coracoid membrane in the infraclavicular region to open into the axillary vein.

The *axillary artery* extends anatomically from the outer border of the first rib to the lower margin of the teres major muscle. When the arm is held out at right angles to the long axis of the body, and the palmar surface of the hand turned upwards, the artery corresponds in direction to a line drawn from the middle of the clavicle to the junction of the anterior and middle thirds of the outer axillary wall at the outlet of that space. At its termination the artery and the accompanying nerves, the neuro-vascular bundle, form a projection which lies behind that due to the coraco-brachialis and biceps (short head) muscles. The artery is divided into three parts by the pectoralis minor muscle, which muscle can be represented by a triangle, the base corresponding to the anterior extremities of the third, fourth and fifth ribs, whilst the apex is situated at the end of the coracoid process.

The *Brachial Artery*.—The arm and forearm being held in the position already indicated as necessary in order to map out the axillary artery, the

brachial artery corresponds to a line drawn from the outer wall of the axillary outlet at the junction of its anterior and middle thirds to the mid-point in front of the bend of

the elbow at the level of the head of the radius.
 Fig. vi., 15. At the last point the artery bifurcates into radial and ulnar arteries.

The *musculo-spiral nerve*, accompanied by the anterior branch of the superior profunda artery, pierces the external intermuscular septum of the arm from behind forwards at the junction of the upper and middle thirds of a line drawn from the insertion of the deltoid muscle to the external condyle of the humerus. Below this point the nerve passes downwards and inwards to the outer bicipital sulcus, where it bifurcates into its two terminal branches. Above the point at which the nerve pierces the external intermuscular septum the course of the nerve may be represented by a curved line drawn upwards and inwards to the junction of the upper arm with the posterior fold of the axilla.

The *musculo-cutaneous nerve* usually pierces the inner aspect of the coraco-brachialis muscle about 1 to 2 inches below the coracoid process. It then runs downwards and outwards deep to the biceps muscle towards the outer bicipital sulcus, at which level it becomes cutaneous. The course of the nerve in the arm can, therefore, be roughly indicated by a line from the coracoid process above to the outer bicipital sulcus below.

The *Median Nerve in the Arm*.—This nerve accompanies the brachial artery, and therefore presents a similar surface marking. It is necessary, however, to bear in mind that the nerve crosses the artery superficially from above downwards and from without





FIG. X.

1. The acromion spine.
2. The acromion process.
3. The circumflex nerve and arteries.
4. The insertion of the deltoid.
5. The external condyle of the humerus.

6. The musculo-spiral nerve.
7. The posterior fold of the axilla.
8. The radius.
9. The posterior interosseous nerve.
10. The radial nerve.

To face p. 20.



inwards. In the diagram the middle third of the nerve is not represented.

The *ulnar nerve* in the upper third of the arm lies along the inner side of the brachial artery. It then leaves that vessel, and, accompanied by the inferior profunda artery, passes downwards and backwards to reach the hollow between the internal condyle and the olecranon process.

The *circumflex nerve and the posterior circumflex artery* both pass backwards through the quadrilateral muscular space, and curl round the surgical neck of the humerus towards the outer and front aspect of the shoulder region. The artery anastomoses with the anterior circumflex artery, and the level of the arterial circle so formed, and of the posterior circumflex nerve, may be represented by a line drawn at right angles to the shaft of the humerus from a point just above the central point of the deltoid muscle.

THE REGION OF THE WRIST AND HAND.

Two tendons only are conspicuous at the front of the wrist—the palmaris longus in the middle line, and the flexor carpi radialis to the outer side of the palmaris longus. The flexor carpi ulnaris can, however, be distinguished by palpation along the ulnar border of the forearm, and it can be traced downwards to its insertion into the pisiform bone. Between the palmaris longus and the flexor carpi ulnaris the main mass of the flexor sublimis digitorum lies.

Two transverse creases are seen in this situation, the upper of which roughly corresponds to the level of the radio-carpal joint, whilst the lower represents

almost exactly the upper limit of the anterior annular ligament.

Just external to where the flexor carpi radialis tendon cuts across the two transverse creases there is a depression in the floor of which the lower end of the radius and the tubercle of the scaphoid bone may be felt. The radial artery crosses this space in a downward and outward direction. The trapezium lies at the lower limit of the depression, immediately below and external to the scaphoid tuberosity. The prominent pisiform bone can be distinguished by tracing downwards the tendon of the flexor carpi ulnaris muscle, and posterior to this bone both cuneiform and unciform bones are situated. A finger's breadth below and external to the pisiform bone deep palpation will verify the position of the hook of the unciform bone.

The *anterior annular ligament* is attached to four bony points, two on the radial side, the scaphoid tubercle and the ridge on the trapezium, and two on the ulnar side, the pisiform and the hook of the unciform. The upper limit of the ligament corresponds to the lower of the two transverse creases in front of the wrist, whilst the lower limit of the ligament lies about $\frac{3}{4}$ inch below.

The *Flexor Synovial Sheaths*.—The flexor longus pollicis, the flexor sublimis and the flexor profundus digitorum all pass beneath the anterior annular ligament. In this situation the flexor sublimis consists of four tendons, of which the medius and annularis lie superficial to the tendons which pass to the index and little fingers. The profundus consists of two parts only, the tendon to the index-finger being alone differentiated




FIG. XI.

1. The flexor carpi radialis.
2. The palmaris longus.
3. The ulnar artery.
4. The flexor carpi ulnaris.
5. The pisiform bone.
6. The transverse creases of the wrist.
7. The superficial branch of the ulnar artery.
8. The deep branch of the ulnar artery.
9. The superficial palmar arch.
10. The deep palmar arch.
11. The digital branches.
12. The superficialis volæ.

FIG. XII.

1. The pisiform.
2. The unciform.
3. The trapezium.
4. The scaphoid tuberosity.
5. The anterior annular ligament.
- 6, 6. The flexor longus pollicis sheath.
- 7, 7. The main flexor synovial sheath.
8. The distal flexor synovial sheaths.
9. The continuation of the main sheath along the little finger.




FIG. XII.

To face p 22.





FIG. XI.



off from the main mass. Beneath the ligament these tendons are surrounded by two synovial sheaths, one for the flexor longus pollicis and one for the remaining tendons plus the median nerve. The sheaths extend upwards about 1 inch above the upper limit of the ligament, and therefore the same distance above the lower transverse

Fig. xii.,
6, 6.

crease in front of the wrist. The flexor longus pollicis sheath is continued downwards to the insertion of the tendon into the distal phalanx of the thumb. The main sheath broadens out below the ligament, and though generally continued onwards to the end

Fig. xii., 7,
7, 9.

of the little finger, the major portion terminates at the level of the upper transverse crease of the palm. The flexor tendons to the fore, middle and ring fingers also possess more distally distinct synovial sheaths,

Fig. xii., 8,
8, 8.

which extend from the terminal phalanges of the fingers upwards to the necks of the metacarpal bones, a level corresponding roughly to the lower transverse crease of the palm.

A distance of $\frac{1}{2}$ inch separates the main synovial sheath above from the more distal segments below.

On the outer side of the wrist the most marked feature is the "anatomical snuff-box," a space bounded on the radial side by the tendons of the extensor ossis metacarpi and primi internodii pollicis muscles, and on the ulnar side by the tendon of the extensor secundi internodii pollicis.

In the floor of the space the styloid process of the radius is felt, this prominence lying fully $\frac{1}{2}$ inch below the level of the corresponding process of the ulna, and also on a slightly more anterior plane. Immediately below the radial styloid process the scaphoid bone lies, most prominent when the hand is well adducted. Below this, again,

the trapezium and the bases of the first and second metacarpals are to be felt.

On the dorsum of the hand there is a well-marked elevation, most noticeable when the wrist is fully flexed, due to the projection of the bases of the second and third metacarpal bones, the styloid process of the latter bone being especially prominent.

Immediately above this elevation there is a depression where the tendons of the extensor carpi radialis longior and brevior are felt as they pass to their insertion into the bases of the second and third metacarpal bones.

Near the middle of the posterior aspect of the lower end of the radius a tubercle can generally be distinguished, the radial tubercle, separating the extensor secundi internodii pollicis on the inner side from the tendon of the extensor carpi radialis longior, which lies more external.


The *posterior annular ligament*, about $\frac{3}{4}$ inch broad, extends from the lower part of the outer border of the radius to the styloid process of the ulna and the carpal bones below the ulna. The ligament has, therefore, a downward and inward direction, and beneath it pass the extensor tendons. These occupy distinct compartments and possess synovial sheaths as under :

1. One compartment and synovial sheath for the extensor ossis metacarpi and extensor primi internodii pollicis.

2. One for the extensor carpi radialis longior and brevior.

3. One for the extensor secundi internodii pollicis.

4. One for the extensor communis digitorum and extensor indicis.






FIG. XIII.

1. The posterior annular ligament.
4. The radial tubercle.
- 2, 3, 5-8. The compartments and synovial sheaths of the
extensor tendons (see text).

To face p. 25.

Fig. xiii., 7. 5. One for the extensor minimi digiti.
 Fig. xiii., 8. 6. One for the extensor carpi ulnaris. The extent of the synovial sheaths is indicated in the diagram.

The *radial artery* extends from the middle of the bend of the elbow at the level of the head of the radius to the radial side of the tendon of the flexor carpi radialis muscle just above the base of the thumb. The artery then crosses the "anatomical snuff-box" towards the base of the first interosseous space.

The *radial nerve*, a branch of the musculo-spiral, arises in the outer bicipital sulcus, and passes vertically down the forearm to meet and accompany the radial artery in the middle third of the course of that vessel, lying to the outer side of the artery. At the junction of the middle and lower thirds of the forearm the nerve turns round the outer border of the radius under cover of the supinator longus tendon, and is distributed to the back of the wrist and hand.

The *posterior interosseous* nerve arises from the musculo-spiral in the outer bicipital sulcus, and it curls round the neck of the radius in the substance of the supinator brevis muscle, emerging from the posterior border of that muscle 2 inches below the head of the radius. It then passes vertically down the posterior aspect of the forearm, lying about midway between the inner and outer borders. It terminates in a gangliform enlargement at the posterior aspect of the wrist.

The *median nerve* in the forearm passes vertically downwards from the inner bicipital sulcus to the front of the wrist, there lying to the ulnar side of the flexor carpi radialis and under cover of the palmaris

longus tendon. The nerve then passes under the anterior annular ligament to the palm.

The *ulnar nerve* in the forearm corresponds in direction to a line drawn from the internal condyle of the humerus to the radial side of the pisiform bone. In front of the wrist the nerve lies to the radial side of the tendon of the flexor carpi ulnaris muscle, and it subsequently passes superficial to the anterior annular ligament to its palmar distribution.

The *ulnar artery* in the lower two-thirds of its course accompanies and lies to the radial side of the ulnar nerve. The upper third of its course is represented by a line which passes obliquely upwards and outwards to the middle of the bend of the elbow at the level of the head of the radius.

The *superficial palmar arch* is formed by the anastomosis of the superficial division of the ulnar artery with the superficialis volæ, or with some other branch of the radial artery. The convexity of the arch looks towards the fingers, and lies on a level with the lower border of the outstretched thumb. Occasionally the arch extends lower down, reaching as far as the upper of the two transverse creases on the palmar aspect of the hand.

The *deep palmar arch*, formed by the anastomosis of the radial artery with the deep branch of the ulna, lies about one finger's breadth above the level of the superficial palmar arch.

The *digital branches* of the superficial palmar arch pass downwards in the intervals between the metacarpal bones to within $\frac{1}{2}$ inch of the digital clefts, where the vessels bifurcate to run along the adjacent sides of the fingers.

The *palmar fascia* is triangular in shape, the apex being attached to the anterior annular ligament between the thenar and hypothenar eminences, whilst the base corresponds to the proximal ends of the four inner fingers.

THE CREASES OF THE PALM AND FINGERS.

The upper transverse crease on the palmar aspect of the hand lies just below the normal limit of the superficial palmar arch, but at the level of the lower limit of the main flexor synovial sheath. The lower crease crosses the necks of the metacarpal bones, and corresponds to the upper limit of the distal flexor synovial sheaths.

The metacarpo-phalangeal joints lie about half-way between the two transverse creases of the palm. The middle and distal creases on the palmar aspect of the fingers correspond fairly accurately to the respective inter-phalangeal joints. The "knuckles" are formed by the heads of metacarpal bones and by the heads of the proximal and middle phalanges.

CHAPTER III

THE THORAX


THE female *mamma*, when well developed, extends upwards to the second rib, inwards to the outer border of the sternum, downwards to the sixth or seventh rib, and outwards to the mid-axillary line. The true glandular substance is, however, less regular in disposition, prolongations passing in various directions. Of these, the most important is the so-called "axillary prolongation," which is directed upwards and outwards under cover of the anterior fold of the axilla.

The *nipple* usually corresponds to the fourth interspace.

Fig. xiv. It is variable in position, and not infrequently it overlies the fourth rib. In the female the nipple lies just below, and external to, the apex or central point of the breast.

The *heart* lies opposite the middle four dorsal vertebræ, and the projection of its anterior surface upon the front of the chest wall may be indicated in the following manner.

- Fig. xiv.,
1-4.
- Take the following four points:
1. The lower border of the second left costal cartilage about 1 inch from the left border of the sternum.
 2. The upper border of the third right costal cartilage $\frac{1}{4}$ to $\frac{1}{2}$ inch from the right border of the sternum.
 3. The lower border of the sixth right costal cartilage $\frac{3}{4}$ inch from the right border of the sternum.






FIG. XIV.

To face p. 29.

FIG. XIV

N.B.—In this and in other figures in which the costal cartilages are depicted the numbered references do *not* refer to the corresponding cartilages. These are numbered for general convenience only.

- 1-4. The four points of the heart.
5. The auricular area.
6. The ventricular area.
7. The left ventricle.
8. The apex beat.
9. The pulmonary valve.
10. The pulmonary artery.
11. The aortic valve.
12. The mitral valve.
13. The tricuspid valve.
14. The ascending aorta.
15. The aortic arch.
16. The innominate artery.
17. The right and left common carotid arteries.
18. The right and left subclavian arteries.
19. The right and left subclavian veins.
20. The right and left internal jugular veins.
21. The right and left innominate veins.
22. The superior vena cava.
23. The inferior vena cava.
24. The abdominal aorta.
25. The coeliac axis.
26. The superior mesenteric artery.
27. The renal arteries.
28. The inferior mesenteric artery.
29. The common iliac arteries.
30. The internal iliac arteries.
31. The external iliac arteries.
32. The kidney.
33. The ureters.
34. The ovary.

THE UNIVERSITY OF CHICAGO
LIBRARY

4. The fifth left interspace $1\frac{1}{2}$ inches below and just internal to a line drawn vertically downwards from the nipple. If any abnormality exists with regard to the position of the nipple, this last point may be fixed with greater accuracy by taking a point in the fifth left interspace nearly 4 inches from the middle line.

The apex beat, or the point of maximum impulse of the left ventricle against the chest wall, lies in the fifth left interspace $1\frac{1}{2}$ inches below and $\frac{1}{2}$ inch internal to the nipple, or $3\frac{1}{2}$ inches from the middle line. The position of the apex beat, therefore, does not exactly represent the outermost limit of the left ventricle. The above four points should now be joined in the following manner :

Points 1 and 2 by a straight line.

Points 2 and 3 by a curved line, the heart reaching the greatest distance from the middle line, $1\frac{1}{2}$ inches, in the fourth interspace.

Points 3 and 4 by a line presenting a slight downward convexity, and cutting across the middle line in close relation to the xiphisternal junction.

Points 1 and 4 by a line presenting a fairly well marked convexity to the left. In the illustration this line is drawn rather too straight.

A line joining Points 1 and 3 divides this cardiac area into two parts, which roughly correspond to the *auricular* area above and to the right, and the *ventricular* area below and to the left. This ventricular area is occupied mainly by the right ventricle, whilst a narrow strip along the left border represents that part of the left ventricle which comes to the surface.

THE VALVES OF THE HEART.

1. The *pulmonary* valve is situated at the highest level, and lies opposite the upper border of the third left costal cartilage, close to its junction with the sternum.

2. The *aortic* valve lies just below and internal to the pulmonary valve at the lower border of the third left costal cartilage at its junction with the sternum.

3. The *mitral* or left auriculo-ventricular valve is situated behind the left half of the sternum at the level of the fourth chondro-sternal junction.

4. The *tricuspid* or right auriculo-ventricular valve lies very obliquely behind the sternum at the level of the fourth interspace and of the anterior extremities of the fifth costal cartilages, extending downwards and to the right almost as far as the sixth chondro-sternal junction.

THE AORTA.

The *ascending aorta*, 2 to 2½ inches long, arises behind the left border of the sternum at the level of the third costal cartilage, and it passes upwards and to the right towards the right border of the sternum at the level of the second costal cartilage.

The *aortic arch* is directed backwards and to the left, the upper limit lying about 1 inch below the suprasternal notch, or half-way between that notch and the angle of Ludwig. The arch becomes the descending aorta at the left side of the lower part of the body of the fourth dorsal vertebra.

The *descending thoracic aorta*, 7 to 8 inches long, passes

onwards through the posterior mediastinum, and pierces the diaphragm at the level of the twelfth dorsal vertebra. The site of diaphragmatic perforation is represented on the surface by a point situated just to the left of the middle line, two fingers' breadth above the transpyloric plane.*

The *innominate artery*, $1\frac{1}{2}$ to 2 inches long, arises from
 Fig. xiv., the aortic arch in the middle line 1 inch below
 16. the suprasternal notch, and it passes upwards and to the right to the right sterno-clavicular articulation, at which level it bifurcates into its two terminal branches.

The *left common carotid* (thoracic course) arises from
 Fig. xiv., the aortic arch on a posterior plane to, and
 17. slightly to the left of, the trunk of the innominate artery, and it passes upwards and to the left to the left sterno-clavicular articulation.

The *left subclavian artery* (thoracic course) arises from
 Fig. xiv., the aortic arch on a posterior plane to, and
 18. slightly to the left of, the thoracic part of the left common carotid artery, and it passes almost vertically upwards behind the left border of the sternum to the left sterno-clavicular joint.

The *superior mediastinum* is bounded above by the plane of the thoracic inlet and below by a plane which passes backwards from the angle of Ludwig in front to the lower border of the fourth dorsal vertebra behind (Ludwig's plane).

The *pulmonary artery* arises opposite the upper border
 Fig. xiv., of the third left costal cartilage at its junction
 10. with the sternum, and passes backwards and slightly upwards to its bifurcation, which takes place opposite the second left costal cartilage.

* Any "planes" mentioned in this chapter will be explained in the chapter on the abdomen.

The *internal mammary artery* arises from the first part of the cervical course of the subclavian artery, and it passes almost vertically downwards behind the corresponding sterno-clavicular joint. In its further thoracic course the artery lies $\frac{1}{2}$ inch external to the outer border of the sternum, bifurcating opposite the sixth costal cartilage or the sixth interspace into the *musculo-phrenic* and *superior epigastric* arteries. The former vessel curves outwards, following the line of the costal arch, whilst the latter passes onwards to enter the sheath of the rectus abdominis muscle.

The *left innominate vein*, 3 inches long, is formed opposite the left sterno-clavicular joint, and it passes to the right, slightly overlapping the upper part of the aortic arch, and occupying the greater part of the space between the summit of the arch below and the suprasternal notch above.

The *right innominate vein* is formed at the right sterno-clavicular joint, and it passes obliquely downwards and inwards to meet the corresponding vein of the opposite side at the lower border of the first right costal cartilage close to its junction with the sternum.

The *superior vena cava*, formed by the junction of the above two veins, is directed almost vertically downwards from the lower border of the first right costal cartilage close to its junction with the sternum, to open into the right auricle of the heart at the level of the upper border of the third right chondro-sternal junction.

The *inferior vena cava* enters upon its short intrathoracic course by passing through the quadrate opening of the diaphragm at the level of the eighth dorsal vertebra, opening into the right auricle of the heart

opposite the fifth right interspace and the adjoining part of the sternum.

The *vena azygos major* drains the whole thoracic wall except the first space on the right side and the upper three spaces on the left. It opens into the superior vena cava at the level of the lower part of the second right interspace, curling round the root of the right lung in order to reach its destination.

The main *aortic intercostal* vessels occupy the subcostal groove of a rib as they pass round the chest wall, lying between the corresponding vein above and the nerve below.

THE PLEURA AND LUNGS.

I. The *Pleural Sacs*.—When the shoulders are depressed, Fig. xv., 5. the two clavicles lie practically at right angles to the long axis of the body, and in this position Fig. xvi., 4. the apices of the pleural sacs extend into the supra-clavicular region, lying about $1\frac{1}{2}$ inches above the clavicle under cover of the clavicular head of the sterno-cleido-mastoid muscle.

The anterior margin of each sac sweeps downwards and inwards behind the corresponding sterno-clavicular joint, the two sacs converging towards the angle of Ludwig, at which level they meet one another just to the left of the middle line. They then pass vertically downwards parallel to one another as far as the level of the fourth chondro-sternal junction.

The right sac passes onwards in the same straight line to the sixth or seventh chondro-sternal articulation, and then sweeps round the anterior, lateral, and posterior aspects of the chest wall, cutting across—

(1) The upper part of the eighth costal cartilage in the lateral vertical line;

(2) The tenth rib in the mid-axillary line ;

(3) The eleventh rib in the line of the inferior angle of the scapula ;

(4) The twelfth rib at the outer border of the erector spinæ muscle.

The obliquity of the twelfth rib causes the pleura to fall below the level of the inner half of the rib, the pleura in this last part of its course being directed inwards towards the spine of the twelfth dorsal vertebra.


The *left pleura*, from the level of the fourth left chondrosternal articulation, sweeps obliquely outwards and downwards behind the costal cartilages of the fifth, sixth, and seventh ribs to the eighth costal cartilage in the lateral vertical plane. Beyond this point the left pleura follows the same general direction as the right sac, descending, however, to a slightly lower level.

The lowest limit reached by the two pleural sacs is situated in the mid-axillary line, the sacs there cutting across the tenth rib about 2 inches above the costal margin, which is in this situation usually represented by the tip of the eleventh rib.

Difficulty is sometimes experienced in endeavouring to verify the position of the twelfth dorsal spine, and in such cases the pleural reflection behind the mid-axillary region may be represented by a line which passes almost transversely backwards towards the median posterior line from the tenth rib in the mid-axillary line.

The two sacs are for the most part separated from one another by the width of the vertebral bodies—
 Fig. xvi., 4. about $1\frac{1}{2}$ inches. In the lower part of the posterior mediastinum, however, the right sac approaches the median line.

2. The *lungs* may be mapped out in part by lines similar






FIG. XV.

- | | |
|------------------------------|--|
| 1. The trachea. | 7, 7. The main oblique fissure of the lungs. |
| 2. The left bronchus. | 8. The small transverse fissure of the right lung. |
| 3. The eparterial bronchus. | 9. The liver |
| 4. The hyparterial bronchus. | |
| 5, 5. The pleuræ. | |
| 6, 6. The lungs. | |

to those given for the pleural sacs. The apex of the upper lobe of each lung extends into the supra-clavicular region, and the anterior margins converge towards the angle of Ludwig, cutting obliquely across the corresponding sterno-clavicular joint.

The two anterior borders do not, however, meet at the angle of Ludwig, for the right lung on reaching the middle line passes vertically downwards as far as the level of the sixth or seventh chondro-sternal junction, whilst the left lung runs down behind the left border of the sternum to the junction of the fourth costal cartilage with the sternum. The *right lung*, from the level of the sixth or seventh chondro-sternal junction, sweeps outwards, cutting across—

- (1) The sixth costal cartilage in the lateral vertical line ;
- (2) The eighth rib in the mid-axillary line ;
- (3) The tenth rib in the line of the inferior angle of the scapula, and finally passing inwards towards the tenth dorsal spine.

The *left lung*, from the outer border of the sternum, at the level of the fourth chondro-sternal articulation, passes outwards for a short distance along the lower border of the fourth costal cartilage, and then turns downwards and inwards in a curved direction to the sixth costal cartilage in the lateral vertical plane. The lung then sweeps round the chest wall, following a course similar to that already indicated as pursued by the right lung, the left lung lying, however, at a slightly lower level.


Stress should, perhaps, be laid on the fact that, as the lower border of each lung sweeps round the antero-lateral, lateral, and posterior aspect of the chest wall from the sixth costal cartilage in front to the tenth dorsal spine behind, the course pursued is practically transverse to the long axis of the body.

*The Lower Limit of the Lung, the Pleura, and the Liver
in the Right Mid-axillary Line.*

- (1) The lung corresponds to the eighth rib.
 (2) „ pleura „ „ tenth rib.
 (3) „ liver „ „ costal margin, or even
 lower.

The *Fissures of the Lungs*.—The main fissure of each lung is indicated by a curved line which starts behind Fig. xv., 7. at the level of the second dorsal spine, the apex of the main lower lobe of each lung, therefore, being situated immediately below and external to this dorsal spine. When the arms fall naturally to the side of the body, the line representing the main fissure cuts across the infraspinous surface of the scapula, and crossing then the fifth rib in the mid-axillary line, it terminates in front at the inferior border of the lung at the level of the sixth chondro-sternal junction. When the arms are extended above the head, the inferior angle of the scapula slides upwards and outwards on the chest wall. In this position the main fissure is represented by a line from the second dorsal spine which passes downwards and outwards to the inferior angle of the scapula, and then forwards to the termination of the fissure at the inferior border of the lung.

The smaller or upper fissure of the right lung separates off from the main fissure in the mid-axillary line, Fig. xv., 8. and passes almost transversely forwards along the lower border of the fourth rib and costal cartilage to the anterior border of the lung. It is thus manifest that the anterior aspect of the chest, above the level of the sixth costal cartilage, corresponds to the upper two lobes of the right lung and to the upper lobe of the left lung, whilst the posterior aspect of the chest below the level of the






FIG. XVI.

- | | |
|--|--|
| 1, 1. The spinal dural sheath. | 9. The spleen. |
| 2. The spinal cord. | 10. The left kidney in Morris's quadrilateral. |
| 3. The filum terminale. | 11. The ureter. |
| 4, 4. The pleurae. | 12. The liver. |
| 5, 5. The lungs. | 13. The descending colon. |
| 6, 6. The main oblique fissures of the lungs. | 14. The rectum. |
| 7, 7. The apex of the lower lobe of the lungs. | 15. The posterior superior iliac spine. |
| 8, 8. The roots of the lungs. | |

second dorsal spine corresponds to the right and left main lower lobes.

The *Apices of the Upper and Lower Lobes*.—It will be here in place to again lay stress on the fact that the apex of the upper main lobe lies about 1 inch above the clavicle in the supraclavicular fossa, under cover of the clavicular head of a well-developed sterno-cleido-mastoid muscle, and that
 Fig. xvi., 7. the apex of the lower lobe lies immediately below and external to the spine of the second dorsal vertebra.

The *roots of the lungs* lie opposite the spines of the fourth, fifth, and sixth dorsal vertebræ, and the bodies
 Fig. xvi., 8. of the fifth, sixth, and seventh vertebræ. They may be so represented, lying also midway between the median posterior line and the vertebral border of the scapula, the arms hanging loosely from the shoulders.

The *Areas of Deep and Superficial Cardiac Dulness*.—
 1. The area of deep cardiac dulness, quadrate in form,
 Fig. xiv., 1-4. corresponds to the complete area already mapped out as representing the projection of the heart on to the anterior aspect of the chest wall.

2. The area of superficial cardiac dulness, more or less triangular in shape, corresponds to that part of the heart
 Fig. xiv. which is not covered by the thin anterior margin of the lung. This area can, with sufficient accuracy, be defined as a triangular space, the left border being formed by a line from the fourth left chondro-sternal articulation to the apex beat of the heart in the fifth left interspace, the right border by a line which passes downwards along the middle of the sternum from the level of the anterior extremities of the fourth to the seventh costal cartilages, and the base by a line which passes out-

wards from the level of the seventh costal cartilage to the position of the apex beat.

The *Trachea and Bronchi*.—The trachea, $4\frac{1}{2}$ inches long, commences immediately below the cricoid cartilage, on a level with the sixth cervical vertebra, and passing downwards through the superior mediastinum, it bifurcates opposite the level of the lower part of the body of the fourth dorsal vertebra (Ludwig's plane).
Fig. xv., 1.

The two bronchi diverge, the left being the longer and the narrower. The tendency of foreign bodies to pass more frequently into the right bronchus is explained by the fact that the septum between the two bronchi is placed to the left of the middle line of the trachea. The right bronchus, previous to the giving-off of the eparterial bronchus, is less obliquely inclined than the left bronchus, though subsequently it follows much the same course.
Fig. xv., 2, 3, 4.

The greater obliquity of the left bronchus accounts also for the fact that the left pulmonary artery tends to lie at the higher level, whilst the right pulmonary artery lies below the level of the corresponding bronchus.

The *œsophagus*, 9 inches long, also commences at the level of the cricoid cartilage, and passing downwards through the superior and posterior mediastina, it pierces the diaphragm at the level of the tenth dorsal vertebra, entering the stomach at the level of the eleventh vertebra.
Fig. xviii., 1, 1.

The point of entrance of the *œsophagus* into the stomach may be indicated by taking a point on the seventh left costal cartilage $\frac{1}{2}$ inch away from the left side of the xiphisternal junction.

The *thoracic duct*, 15 to 18 inches long, commences

as the receptaculum chyli, a spindle-shaped sac which is situated opposite the bodies of the first and second lumbar vertebræ, and which may be represented on the surface by an oval enlargement placed just to the right of the middle line, occupying the upper two-thirds of the space between the transyloric (first lumbar) and subcostal (third lumbar) planes.

The efferent duct pierces the diaphragm through the aortic orifice opposite the twelfth dorsal vertebra, and passes almost vertically upwards through the posterior mediastinum, just to the right of the middle line, as far as the lower part of the fourth dorsal vertebra (Ludwig's plane). The duct now crosses behind the œsophagus to the left of the middle line, and then again passes vertically upwards through the superior mediastinum and into the neck as far as the level of the transverse process of the seventh cervical vertebra. Finally, the duct curls outwards and downwards to open

into the angle between the internal jugular and subclavian veins of the left side. The duct drains the whole of the lymphatic area of the body, except the right side of the head and neck, the right arm, the right side of the thorax, and the convexity of the liver.

CHAPTER IV
THE ABDOMEN

THE anterior aspect of the trunk (*i.e.*, thorax and abdomen) is divisible into right and left halves by a *median vertical plane* from the middle point at the suprasternal notch above to the symphysis pubis below. Each half is again divided by a *lateral vertical plane* which is drawn parallel to the median plane half-way between that plane and the anterior superior iliac spine. Prolonged downwards, this lateral plane crosses Poupart's ligament rather nearer to the inner than to the outer end. Prolonged upwards, it crosses the clavicle about midway between the median point at the suprasternal notch and the acromio-clavicular joint.

That part of the lateral vertical plane which traverses the mammary region is sometimes called the "mammary plane," that part which crosses the clavicle the "clavicular plane," and the downward prolongation which cuts across Poupart's ligament the "Poupart plane."

The clavicular, mammary, and Poupart planes are, however, continuous, and they together form the *lateral vertical plane*, which is chosen in preference to the mid-Poupart plane of many anatomists, since it is measured from the median plane to a fixed bony point. Two vertical planes only will be consequently retained in the subdivision of the anterior aspect of the trunk, the *median and lateral vertical planes*.

The median plane can be bisected by a horizontal plane

which is on the same level as the body of the first lumbar vertebra. This plane so constantly cuts across the pyloric end of the stomach that it is called the *transpyloric plane*, and it will be found that not only does this plane lie half-way between the suprasternal notch and the symphysis pubis, but that it also lies midway between the umbilicus and the sterno-xiphoid junction. It is, therefore, not necessary to expose the whole of the anterior aspect of the trunk in order to verify the position of the transpyloric plane, a plane of the greatest value in defining the position of several abdominal viscera.

The point at which the median, vertical, and transpyloric planes intersect has been suitably called the "central point," and the point of intersection of the lateral vertical and transpyloric planes may be tentatively called the lateral central or paracentral point. This latter point usually corresponds to some part of the ninth costal cartilage.

The distance between the central point and the top of the symphysis pubis is bisected by a horizontal plane which passes through the tubercles of the iliac crests, the *intertubercular plane*, a plane corresponding to the level of the body of the fifth lumbar vertebra. It has also been suggested that the distance between the central point and the suprasternal notch should likewise be bisected by a plane, the *thoracic plane*, which crosses the gladiolus at the level of some part of the anterior extremities of the fourth costal cartilages. This plane is, however, of little value, and it is merely mentioned as completing the symmetrical subdivision of the median vertical plane into four equal parts.

The following planes are, therefore, suggested as the

most scientific in the subdivision of the anterior aspect of the trunk :

Two vertical planes—(1) The median ; (2) the lateral.

Three transverse planes—(1) The intertubercular ; (2) the transpyloric ; (3) the thoracic.

Two important points are also named—(1) The central point ; (2) the lateral central point.

The abdominal regions mapped out by the intersection of the transpyloric and intertubercular planes with the lateral vertical planes receive the same nomenclature as in the older methods of regional subdivision of the abdomen. These regions are nine in number :

1. Right hypochondriac. 2. Epigastric. 3. Left hypochondriac. 4. Right lumbar. 5. Umbilical.
 Fig. xvii. 6. Left lumbar. 7. Right iliac. 8. Hypogastric. 9. Left iliac.

OTHER TRANSVERSE PLANES, WITH THEIR CORRESPONDING VERTEBRAL LEVELS.

(a) The *suprasternal plane*, on a level with the disc between the second and third dorsal vertebræ.
 Fig. xvii.


(b) *Ludwig's plane* (junction of manubrium and gladiolus), on a level with the lower part of the body of the fourth dorsal vertebra.

(c) The *sterno-xiphoid plane* (junction of sternum and xiphoid cartilage), on a level with the disc between the ninth and tenth dorsal vertebræ.

(d) The *subcostal plane*, on a level with the upper part of the third lumbar vertebra.

(e) The *umbilical plane*, on a level with the disc between the third and fourth lumbar vertebræ.

(f) The *spinous plane*, drawn between the two anterior



Level of hard palate = first cervical.
 Level of free margin of upper teeth = second cervical.
 Level of hyoid bone = second to third cervical.
 Level of upper part of thyroid cartilage = fourth cervical.
 Level of cricoid cartilage = sixth cervical.




FIG. XVII.

1. The central point.
- 2, 2. The lateral central or paracentral point.
- 3, 3. The median vertical plane.
- 4, 4. The lateral vertical plane.
- 5, 5. The mid-Poupart point.
- 6, 6. The pubic spines.

superior iliac spines, and usually falling below the level of the sacral promontory.

The *lineæ semilunares* correspond to the outer border of the rectus abdominis muscle, and they extend with a slight outward convexity from the pubic spine below to the tip of the ninth costal cartilage above (the lateral central point).

The *lineæ transversæ* result from the tendinous inter-sections in the rectus abdominis muscle. They are three in number, and are situated—(1) at the level of the umbilicus; (2) midway between the umbilicus and the xiphoid cartilage; (3) immediately below the xiphoid cartilage.

The *semilunar fold of Douglas*, representing the lower limit of the posterior lamella of the rectus sheath, lies about half-way between the umbilicus and the upper border of the pubic symphysis.

The *umbilicus* usually lies 1 to 1½ inches above the intertubercular plane, and corresponds to the level of the disc between the third and fourth lumbar vertebræ. The umbilicus is, however, so inconstant in position that the umbilical plane is rejected as often as possible in favour of a more definite and scientific plane.

The *Iliac Spines and Crest*.—When the body is in the dorsal recumbent position, the anterior superior iliac spine is usually visible to the eye, and no palpation is needful in order to fix its position. In the obese, however, it is generally necessary to trace forward the iliac crest to its anterior termination. By tracing the iliac crests in a backward direction the “iliac tubercles” will be found lying about 2 to 2½ inches behind the anterior superior spines, and a line uniting these two

tubercles (the intertubercular plane) corresponds to the level of the fifth lumbar vertebra. Still further backward the posterior superior iliac spines will be found at the posterior termination of the iliac crest. A line which joins the posterior superior iliac spines cuts across the spine of the second sacral vertebra in the middle line behind.

The *pubic spine* lies at the outer limit of the pubic crest. In the male it is advisable to invaginate the scrotum in order to locate the position of this spine; whilst in the female, owing to the prominence of the mons veneris, it is usually necessary to abduct the thigh, to feel for the rounded tendon of the adductor longus muscle, and to trace this tendon up to its origin from a depression on the pubic bone, which is situated immediately below and internal to the pubic spine.

The *Inguinal Canal*.—In the adult this canal is about $1\frac{1}{2}$ inches long, and it extends from the internal to the external abdominal rings. The *internal abdominal ring*, a funnel-shaped prolongation of the transversalis fascia, is situated $\frac{1}{2}$ inch above the midpoint of Poupart's ligament. The *external abdominal ring*, formed by the splitting of the aponeurosis of the external oblique muscle, is triangular in shape, the base directed downwards and inwards and opening up immediately above the pubic spine, whilst the apex is directed upwards and outwards.

Petit's Triangle.—This triangle is bounded in front by the posterior border of the external oblique, and behind by the anterior border of the latissimus dorsi muscle, whilst the base is formed by part of the iliac crest. The external oblique is inserted into the anterior half of the

iliac crest, and the base of the triangle corresponds to 1 to 2 inches of the bone behind the mid-point of the crest. The triangle is subject to great variation in size, the two bounding muscles converging rapidly above to form the apex of the triangle. The floor is formed by the internal oblique muscle.

THE ALIMENTARY CANAL.

The *Stomach*.—Capacity about 2 pints. The *cardiac orifice* lies opposite the eleventh dorsal vertebra, and it is situated about 4 inches away from the surface. It corresponds in position to a point on the seventh costal cartilage $\frac{1}{2}$ inch away from the outer border of the sterno-xiphoid junction. The seventh costal cartilage is the lowest of the series of cartilages which articulate in front with the mesial sterno-xiphoid bar, and it forms, therefore, the upper lateral boundary of the epigastric triangle. The *pyloric orifice* lies opposite the first lumbar vertebra, and it corresponds in position to a point in the transpyloric plane just to the right of the middle line.

The *lesser curvature* is represented by a curved line, convexity to the left, uniting the above two points. The *greater curvature*, in the moderately-distended condition of the stomach, ascends to the lower border of the left fifth costal cartilage and rib, lying immediately above and behind the apex of the heart. Sweeping then downwards, the greater curvature usually cuts the left costal margin at some part of the ninth costal cartilage, and finally curves upwards and inwards to the pylorus. The upper limit of the fundus of the stomach corresponds to the level of the left dome of the diaphragm.


The *Duodenum*.—Total length about 10 inches. Part 1 = 2 inches; part 2 = 3 to 4 inches; part 3 = 4 to 5 inches.

The pyloric orifice of the stomach lies opposite the first lumbar vertebra, and the first part of the duodenum is directed backwards, with a slight inclination upwards, to the right side of the body of the first lumbar vertebra.

Part 2 descends on the right side of the median vertical plane from the level of the first lumbar vertebra (transpyloric plane) to the level of the third lumbar vertebra (subcostal plane). The third part of the duodenum passes at first almost transversely across the middle line at the level of the subcostal plane, and having reached the left side of the middle line, it ascends sharply to the *duodeno-jejunal* flexure, which is placed on a level with the second lumbar vertebra just below the transpyloric plane, and 1 to 1½ inches to the left of the middle line. The duodenum is subject to great variation in position, and the description given merely represents the average situation of this loop of the small gut. The pyloric orifice of the stomach and the duodeno-jejunal flexure are both fairly constant in position, and between these two more or less fixed points the gut describes a loop which varies both in shape and in extent.

The *Pancreas*.—The head of the pancreas occupies the concavity of the duodenal loop, the body crossing the middle line at the level of the first and second lumbar vertebræ, and occupying, therefore, the upper two-thirds of the space between the transpyloric (first lumbar) and subcostal (third lumbar) planes. The tail of the pancreas extends to the left as far as the hilum of the spleen.

The *small intestine* is about 23 feet in length, the upper







FIG. XVIII.

a, a and *a', a'* = the lateral vertical planes.
b, b. The transpyloric plane.
c, c. The subcostal plane.
d, d. The intertubercular plane.

FIG. XVIII.

- 1, 1. The œsophagus.
2. The stomach.
3. The pylorus.
- 4, 4, 4. The three parts of the duodenum. 4'. The pancreas.
5. The duodeno-jejunal flexure.
6. The attachment of the mesentery of the small intestine.
7. The ileo-cæcal valve.
8. The cæcum.
9. The vermiform appendix.
10. The ascending colon.
11. The hepatic flexure.
12. The splenic flexure.
13. The descending colon.
14. The iliac colon.
15. The ileo-pelvic colon.
16. The gastro-hepatic omentum.
17. The foramen of Winslow.
18. The common bile-duct.



two-fifths being known as the jejunum, the lower three-fifths as the ileum.

The *mesenteric attachment* of the small gut extends from Fig. xviii., a point 1 to $1\frac{1}{2}$ inches to the left of the middle
6. line, and just below the transpyloric plane (duodeno-jejunal flexure), to the junction of the right lateral vertical and intertubercular planes. The line drawn between these two points should pass at first obliquely outwards and downwards to the right iliac fossa, curving finally outwards to the region of the ileo-cæcal valve.

The *ileo-cæcal valve* is placed opposite the junction of the Fig. xviii., right lateral vertical and intertubercular planes.
7. The ileum opens into the cæcum on the postero-internal aspect of that part of the large gut. The opening is almost horizontal, though in the diagram, owing to want of space, it has been represented as lying in the vertical direction.

The *cæcum* is about $2\frac{1}{2}$ inches long, and the long axis of Fig. xviii., the sac is directed downwards, forwards, and
8. inwards. This blind end of the large gut lies below the level of the intertubercular plane, and it occupies the right iliac fossa and part of the right half of the hypogastric region.

The Vermiform Appendix.—The opening of the appendix Fig. xviii., into the cæcum is situated just below and
9. internal to the junction of the right lateral vertical and intertubercular planes, at the top right-hand corner of the hypogastric region. The surface marking of the cæcal orifice of the appendix does not coincide with McBurney's point, which is situated at the junction of the outer and middle thirds of a line drawn from the right anterior superior iliac spine to the umbilicus. This point represents the usual seat of maximum pain on palpa-

tion in an attack of appendicitis. The appendix is usually 3 to 4 inches long, and, according to Testut, it is in 40 per cent. of cases directed downwards and inwards, overhanging the pelvic brim, whilst in 26 per cent. of cases only is it directed upwards and inwards (towards the spleen). Stress should be laid on the fact that the ileo-cæcal valve and the cæcal orifice of the appendix are both situated on the postero-internal aspect of the cæcum.

The *ascending colon* passes upwards from the level of the Fig. xviii.,
10. intertubercular plane to the upper part of the ninth right costal cartilage, the gut there turning on itself to form the *hepatic flexure*. In its upward course the ascending colon lies almost entirely to the right of the right lateral vertical plane.

The *transverse colon* extends from the hepatic flexure on Fig. xviii.,
11. the right to the *splenic flexure* on the left. The former flexure corresponds to the ninth costal cartilage, whilst the latter reaches upwards as high as the eighth. In between these two points the gut varies greatly in direction in different subjects. Most commonly the Fig. xviii.,
12. gut passes almost transversely from one side to the other, crossing the middle line at about the level of the second lumbar vertebra. It also crosses the second part of the duodenum, and lies, therefore, usually above the umbilical plane. In the diagram the two flexures are depicted, but the intervening portion of the gut has been intentionally omitted.

The *descending colon* passes almost vertically downwards Fig. xviii.,
13. from the region of the splenic flexure to the level of the posterior part of the iliac crest, below which level it becomes known as the *iliac colon*. The descending colon lies wholly to the left of the left lateral vertical plane.

The operation of lumbar colotomy is now seldom performed, but it is nevertheless necessary to indicate the position of the descending colon on the posterior aspect of the trunk. It corresponds in direction to a line drawn vertically upwards to the tip of the last rib, from a point situated $\frac{1}{2}$ inch behind the mid-point along the iliac crest between the anterior and posterior superior iliac spines.

The *Iliac and Ileo-Pelvic Colons*.—Between the termination of the descending colon at the level of the iliac crest, and the beginning of the rectum proper at the level of the third piece of the sacrum, the large gut describes so varied a course that no definite detailed account can be given of its surface-marking. It may, however, be briefly described as passing downwards and inwards from the level of the iliac crest, parallel to Poupart's ligament, as far as the left side of the pelvic brim (the iliac colon). The gut then forms a great loop, which sweeps over to the right side of the pelvic brim, turning on itself to become the rectum at the level of the third sacral vertebra (ileo-pelvic colon).

The *Rectum*.—A line which unites the two posterior superior iliac spines crosses the spinous process of the second sacral vertebra. The rectum begins at the level of the third sacral vertebra, and it may be indicated on the surface by drawing in the gut as starting about $\frac{1}{2}$ to $\frac{3}{4}$ inch below the above-mentioned line, and extending downwards, following the curves of the sacrum and coccyx to the anal orifice, which is placed about 2 inches below the level of the tip of the coccyx.

The dura mater enclosing the spinal cord (see "Spinal Cord") reaches downwards to the level of the third sacral vertebra. The spinal dura, there-

fore, terminates at the same level as the rectum begins, a point to be borne in mind in those operations carried out in the sacral region for the exposure of a growth involving the gut in the neighbourhood of the ileo-pelvic and rectal junction.

THE KIDNEY.

Length, $4\frac{1}{2}$ inches ; breadth, $2\frac{1}{2}$ inches ; thickness, $1\frac{1}{2}$ inches ; weight, $4\frac{1}{2}$ ounces.

(a) *Anterior Surface Marking*.—The two kidneys are obliquely placed in such a manner that the superior poles lie $1\frac{1}{2}$ to 2 inches, and the inferior poles $2\frac{1}{2}$ to 3 inches, distant from the middle line. The left kidney lies at a slightly higher level than its fellow, and the hilum is placed just below and internal to the junction of the transpyloric and left lateral vertical planes ; or, in other words, the hilum of the left kidney lies just internal to the anterior extremity of the ninth costal cartilage. The upper pole lies half-way between the sternoxiphoid and transpyloric planes, whilst the lower pole corresponds to the subcostal plane. The right kidney does not ascend to quite such a high level, and the inferior pole lies opposite the umbilical plane. The hilum of this kidney also lies just below the level of the hilum of the opposite kidney.

(b) *Posterior Surface Marking—Morris's Quadrilateral*.—Two vertical lines are drawn at a distance of 1 inch and $3\frac{1}{2}$ inches respectively from the median posterior line, and two horizontal lines are drawn outwards at the level of the spinous processes of the eleventh dorsal and third lumbar vertebræ. In the quadrilateral so marked out, the kidneys are drawn, care being taken to place the long axis of each kidney in the required oblique direction.

THE URETERS (10 INCHES LONG).

(a) *Anterior Surface Marking.*—The ureter passes nearly vertically downwards from the hilum of the kidney (just below and internal to the junction of the transpyloric and lateral vertical planes), and it dips into the true pelvis in close relation to the bifurcation of the common iliac artery. This vessel bifurcates into internal and external iliacs at the junction of the upper and middle thirds of a line drawn from a point $\frac{1}{2}$ inch below and to the left of the umbilicus (the aortic bifurcation) to a second point situated half-way between the anterior superior iliac spine and the symphysis pubis. This apparently complicated surface marking for the ureter will be rendered more easy by a reference to the diagram. The right ureter generally dips into the pelvis just below the bifurcation of the common iliac artery.

(b) *Posterior Surface Marking.*—The course of the ureter on the posterior aspect of the trunk can be represented by a line drawn vertically upwards from the posterior superior iliac spine to the level of the spinous process of the second lumbar vertebra.

The *ovary* lies opposite the middle point of a line drawn from the anterior superior iliac spine of the same side to the pubic spine of the opposite side. Anatomically, the ovary lies in close relation to the angle formed between the internal and external iliac arteries.

The *urachus* is directed upwards from the apex of the bladder at the upper border of the pubic symphysis to the umbilicus.


ABDOMINAL VESSELS.

The *Abdominal Aorta*.—The thoracic aorta enters the abdominal cavity by passing beneath the middle arcuate ligament of the diaphragm at the level of the twelfth dorsal vertebra, and it bifurcates into the two common iliac arteries on the left side of the body of the fourth lumbar vertebra. The course of the vessel may be mapped out on the surface by taking a point about two fingers' breadth above the transpyloric plane and slightly to the left of the middle line, and by drawing a line vertically downwards from that point to a second point situated $\frac{1}{2}$ inch below and to the left of the umbilicus.

The *celiac axis* arises immediately below the origin of the abdominal aorta, and it is at once followed by the *superior mesenteric*. The *renals* arise about 3 inches above the umbilicus, and the *inferior mesenteric* is given off about 1 inch above that level.

The *common iliac artery* corresponds to the upper third of a line drawn from a point $\frac{1}{2}$ inch below and to the left of the umbilicus to a second point situated half-way between the anterior superior iliac spine and the symphysis pubis. The *external iliac artery* corresponds in direction to the lower two-thirds of the line.

The *deep epigastric artery* is given off from the external iliac just as that vessel passes under Poupart's ligament half-way between the anterior superior iliac spine and the symphysis pubis. The epigastric artery then passes upwards and inwards along the inner side of the internal abdominal ring towards a point situated







FIG. XIX.

FIG. XIX.

- 1, 1. The diaphragm.
2. The liver.
3. The gall-bladder.
4. The ligamentum teres.
5. The receptaculum chyli.
6. The thoracic duct.
7. The venous termination of the duct.
8. The internal mammary artery.
9. The superior epigastric artery.
10. The musculo-phrenic artery.
11. The rectus abdominis muscle.
- 12, 12. The lineæ semilunares.
- 13, 13, 13. The lineæ transversæ.
14. The semilunar fold of Douglas.
15. The urachus.
16. Hesselbach's triangle.
17. The internal abdominal ring.
18. The external abdominal ring.
19. The inguinal canal.
20. The deep epigastric artery.



$\frac{1}{2}$ inch to 1 inch outside the umbilicus, entering the rectus sheath at the level of the semilunar fold of Douglas.

This vessel forms the outer boundary of *Hesselbach's triangle*, the inner boundary of this space being formed by the linea semilunaris of the same side, and the base by Poupart's ligament. Each triangle is vertically subdivided into two parts by the obliterated hypogastric artery, on either side of which herniæ may protrude.

The *inferior vena cava* is formed by the junction of the two common iliac veins on the right side of the body of the fifth lumbar vertebra, about 1 inch below and $\frac{1}{2}$ inch to the right of the umbilicus. The vein passes upwards to pierce the quadrate opening of the diaphragm at the level of the eighth dorsal vertebra, entering the right auricle of the heart opposite the fifth right interspace and the adjoining part of the sternum.

The *Liver*.—The anterior border can be mapped out by drawing a curved line from a point in the fifth left interspace $3\frac{1}{2}$ inches from the middle line (the position of the apex of the heart), the line cutting the left costal margin at the tip of the eighth costal cartilage and the right costal margin at the tip of the ninth costal cartilage. In between these two latter points the anterior border of the liver crosses the middle line half-way between the umbilicus and the sterno-xiphoid junction (= transpyloric plane), whilst a notch to the right of the middle line indicates the hepatic attachment of the *ligamentum teres*, which passes from that notch downwards and inwards to the umbilicus.

Beyond the tip of the ninth right costal cartilage the anterior border of the liver follows the lower limit of the

costal arch, descending sometimes even below that level, and after cutting across the twelfth rib, it ascends towards the level of the eleventh dorsal spine.

The *upper limit of the liver* is indicated by a line starting as before in the fifth left interspace $3\frac{1}{2}$ inches from the middle line, and ascending slightly as it passes to the right. This line cuts across the sixth right chondro-sternal articulation, the upper border of the right fifth costal cartilage in the right lateral vertical plane, the sixth rib in the mid-axillary line, sweeping thence just below the angle of the scapula towards the eighth dorsal spine. A posterior surface marking for the liver to the right of the middle line is purposely omitted as being unnecessary.

The *Gall-bladder*.—The fundus projects from under the anterior border of the liver in the angle between the tips of the ninth and tenth costal cartilages and the outer border of the rectus abdominis muscle.

The *Diaphragm*.—On ordinary inspiration the right dome of the diaphragm corresponds in level to the lower part of the fourth right interspace, whilst the left dome ascends to the lower part of the fifth left rib and costal cartilage.

THE COMMON BILE-DUCT, ETC.

The *gastro-hepatic omentum*, passing upwards from the lesser curvature of the stomach to the transverse fissure of the liver, presents a free edge, which looks downwards and to the right. This free edge forms the anterior boundary of the foramen of Winslow, and in the free edge three important structures run—the portal vein, the hepatic artery, and the common bile-duct.

In mapping out any of these structures, it is, therefore, advisable to first draw in the lesser curvature of the stomach, the pylorus, the three parts of the duodenum, and the pancreas. The pylorus occupies such a definite position in the transpyloric plan that all these structures are easily and quickly drawn in. The free margin of the lesser omentum should be represented as a curved line passing upwards and to the right for $1\frac{1}{2}$ to 2 inches from the duodeno-pyloric junction. The *portal vein* is formed behind the head of the pancreas by the union of the superior mesenteric and splenic veins, and it passes upwards to the transverse fissure of the liver behind the first part of the duodenum, and in the free edge of the lesser omentum.

The *hepatic artery*, a branch of the cœliac axis, passes upwards from the upper border of the first part of the duodenum, in the free edge of the lesser omentum, to the transverse fissure of the liver.

The *common bile-duct* is 3 inches long, and it is formed
 Fig. xviii.,
 18. by the union of the hepatic and cystic ducts. It passes downwards in the free edge of the gastro-hepatic omentum, behind the first part of the duodenum, behind the head of the pancreas, and it opens on the inner aspect of the second or descending part of the duodenum.

The *Spleen*.—The long axis corresponds to the tenth rib,
 Fig. xvi., 9. and the viscus extends upwards to the upper border of the ninth rib, and downwards to the lower border of the eleventh rib. The upper and inner pole lies $1\frac{1}{2}$ to 2 inches away from the tenth dorsal spine, whilst the lower or anterior pole reaches as far forwards as the mid-axillary line.

The *spinal cord* extends from the foramen magnum to

the lower border of the first lumbar vertebra (trans-
 Fig. xvi., 2. pyloric plane). The cord follows the curves of
 the vertebral column, and presents also two
 enlargements, the *intumescencia cervicalis* and *lumbalis*.
 The former swelling lies between the third cervical and
 second dorsal vertebræ, the latter between the ninth
 and twelfth dorsal vertebræ. Near its termination the
 cord tapers away as the *conus medullaris*.

The *filum terminale*, the prolongation of the cord, is
 Fig. xvi., 3. continued onwards from the lower part of the
 body of the first lumbar vertebra to near the
 tip of the coccyx, at which level it blends with the
 periosteum lining that bone.

The *theca vertebralis*, or dural sheath, extends as low
 Fig. xvi.,
 1, 1. as the third sacral vertebra, at which level it is
 pierced by the *filum terminale*.

A line uniting the two posterior superior iliac spines
 Fig. xvi.,
 15. cuts across the second sacral spine, and the
 dural sac, therefore, terminates about $\frac{1}{2}$ inch
 below the level of this interspinous line. At the third
 month of intra-uterine life the cord extends the whole
 length of the vertebral canal, whilst at birth it reaches as
 low down as the third lumbar vertebra.

CHAPTER V

THE LOWER EXTREMITY

THE pubic spine, the iliac crest, the anterior and posterior superior iliac spines, and the iliac tubercles, have all been previously alluded to and located. It has also been stated that a line uniting the two posterior superior iliac spines cuts across the spine of the second sacral vertebra. Below this line the remaining sacral spines and the coccyx are easily felt, though the coccyx itself is more readily examined by inserting the forefinger into the rectum whilst the thumb is placed over the bone externally.

The great trochanter of the femur and the ischial tuberosity must now be examined. The latter process lies under cover of the lower part of the gluteus maximus muscle, whilst the trochanter is subcutaneous in its lower part, and covered over in front and above by the insertion of the gluteus medius muscle.

The trochanter is more or less quadrilateral in shape, fading off into the shaft of the femur below, and presenting a well-marked posterior border, whilst the highest part is situated at the posterior superior angle. When the body is in the erect position there is a well-marked depression situated above and behind the great trochanter, which becomes less marked when, as the result of disease or disuse, the gluteus maximus undergoes atrophic changes. The fold of the nates does not correspond to the lower border of the gluteus maximus muscle,

as it runs transversely across the lower oblique fibres of that muscle. This fold also becomes less distinct when the glutei muscles degenerate. The head and neck of the femur form with the shaft of that bone an angle of 125 to 130 degrees.

Nélaton's Line.—“ If in the normal state you examine Fig. xx, 8. the relation of the great trochanter to the other Fig. xxii., 5. bony prominences of the pelvis, you will find that the top of the great trochanter corresponds to a line drawn from the anterior superior iliac spine of the ilium to the most prominent point of the tuberosity of the ischium. This line also runs through the centre of the acetabulum. The extent of displacement in dislocation or in fracture is marked by the projection of the trochanter behind and above this line ” (Nélaton).

Bryant's Triangle.—When the patient is in the dorsal Fig. xxii., 4. recumbent position, draw a line round the body at the level of the anterior superior iliac spine, and from this line drop a perpendicular to the top of the great trochanter. To complete the triangle, draw a line from the anterior superior iliac spine to the top of the trochanter. When the trochanter is displaced upwards the perpendicular line is diminished in length as compared with the sound side, and when it undergoes a backward displacement the spino-trochanteric line is relatively increased in length.

In order to *compare the length* of the lower extremities the limbs should be placed parallel to one another, and the tape-measure carried from the anterior superior iliac spine to the tip of the internal malleolus of the tibia of Fig. xxiii., the same side. The distance between these 16. two points may be subdivided if necessary by marking out on each leg the tranverse line which indicates







FIG. XX.

To face p. 59.

FIG. XX.

1. The iliac crest.
2. The posterior superior iliac spine.
3. The second sacral spine.
4. The ischial tuberosity.
5. The sciatic artery.
6. The great trochanter.
7. The gluteal artery.
8. Nélaton's line.
9. The great sciatic nerve.
10. The internal popliteal nerve.
11. The external popliteal nerve.
12. The biceps tendon.
13. The semimembranosus.
14. The semitendinosus.
15. The outer head of the gastrocnemius.
16. The inner head of the gastrocnemius.
17. The posterior tibial artery and nerve.
18. The external saphenous vein.



the level of the femoro-tibial articulation. The length of the femur or of the tibia may so be estimated.

The *gluteal artery* emerges from the great sacro-sciatic notch at the junction of the inner and middle thirds of a line drawn from the posterior superior iliac spine to the top of the great trochanter of the femur of the same side.

The *sciatic artery* may be ligatured at the junction of the middle and lower thirds of a line drawn from the posterior superior iliac spine to the outer part of the ischial tuberosity of the same side. This line also cuts across the *posterior inferior iliac spine* and the tip of the *ischial spine*, the latter spine and the *internal pudic artery* lying immediately internal to the seat of election for ligation of the sciatic artery.

The *great sciatic nerve* emerges from under cover of the lower border of the gluteus maximus muscle just to the inner side of the mid-point between the ischial tuberosity and the great trochanter of the femur. The nerve corresponds in direction to the upper two-thirds of a line drawn downwards from the above point to the middle of the popliteal space below.

The *small sciatic nerve* lies in the same line as the great sciatic, but extends downwards as far as the inferior angle of the popliteal space.

The *Region of the Knee*.—The biceps tendon forms the upper and outer boundary of the popliteal space, and under cover of this tendon, on its inner or popliteal aspect, a cord-like structure is felt, the external popliteal or peroneal nerve. This intimate relation of tendon and of nerve must be remembered in the operation of tenotomy of the biceps tendon. If the biceps tendon be

Fig. xx., 12.

Fig. xxii.,
10.

Fig. xx., 11.

Fig. xxii.,
11.

now traced downwards the head of the fibula is reached, this process lying below, external, and on a posterior plane to the outer tuberosity of the tibia. The styloid process of the head of the fibula projects upwards from the posterior part of the head, and in front of this the rounded long external lateral ligament of the knee-joint can be traced upwards to its femoral attachment. In front of the biceps tendon there is a depression which is bounded anteriorly by the broad ilio-tibial band. Two well-marked tendons bound the popliteal space on the upper and inner side, the semi-membranosus and semitendinosus. The latter is the more external, the more superficial, and the narrower, and the long rounded tendon can be traced some distance up into the thigh. The semi-membranosus tendon lies to the inner side of the semitendinosus and on a deeper plane. The broad tendon can be traced downwards to its insertion into the inner and posterior aspect of the internal tuberosity of the tibia. On the inner aspect of the knee the tendon of the gracilis muscle and the lower part of the sartorius muscle form a fairly well marked prominence, the individual muscles being, however, usually incapable of clear definition owing to their flattened shape. Between these tendons and the prominent vastus internus muscle a depression exists, and by deep palpation the adductor magnus tendon may be felt lying under cover of the inner margin of the vastus internus muscle. By tracing this tendon downwards to its insertion the adductor tubercle is reached. This tubercle corresponds also to the level of the lower epiphysial line of the femur.

The sartorius and gracilis muscles, though not easily




FIG. XXI.

1. The anterior superior iliac spine.
2. The pubic spine.
3. Poupart's ligament.
4. The anterior crural nerve.
5. The common femoral artery.
6. The femoral vein.
7. The femoral or crural ring.
8. The saphenous opening.
9. The profunda femoris artery.
10. The superficial femoral artery.
11. The synovial membrane of the knee-joint.
12. The patella.
13. The prepatellar bursa.
14. The tibial tubercle.
- 15, 15. The ligamentum patellæ.
16. The outer tibial tuberosity.
17. The head of the fibula.
18. The peroneal nerve.
- 19, 19. The anterior tibial nerve.
20. The musculo-cutaneous nerve.
- 21, 21. The anterior tibial artery.
22. The dorsalis pedis artery.
23. The dorsal venous arch.





FIG. XXI.

To face p. 60.






FIG. XXII.

1. The iliac crest.
2. The anterior superior iliac spine.
3. Poupart's ligament.
4. Bryant's triangle.
5. Nélaton's line.
6. The great trochanter.
7. The ilio-tibial band.
8. The outer tibial tuberosity.
9. The head of the fibula.
10. The biceps tendon.
11. The peroneal nerve.
- 12, 12. The anterior tibial nerve.
13. The musculo-cutaneous nerve.
- 14, 14. The anterior tibial artery.
15. The dorsalis pedis artery.
16. The external saphenous vein.
17. The venous arch.





FIG. XXII.

To follow Fig. xxi.



defined on the inner side of the knee, form, together with
 Fig. xxiii.,
 17. the semitendinosus muscle, a fairly well marked elevation below the internal tuberosity of the tibia, which is directed downwards, forwards, and outwards.

The ligamentum patellæ narrows off as it passes from
 Fig. xxi.
 the inferior border of the patella to the tibial tuberosity, and on each side of the ligament depressions exist in the lower part of which the inner and outer tuberosities of the tibia are readily felt.

The *prepatellar bursa* extends from the middle of the
 Fig. xxi.,
 13. patella to the tibial tubercle. Laterally, the bursa falls just short of the patellar border.

The *synovial membrane* of the knee-joint extends upwards
 Fig. xxi.,
 11. about three fingers' breadth above the upper border of the patella when the leg is in the extended position, reaching up under the vastus internus to a slightly higher level than on the outer side. Laterally, the synovial membrane extends to near the inner and outer margins of the femoral condyles, whilst the lower limit is situated just above the tubercle of the tibia. When the joint is distended with fluid the outline of the joint cavity becomes marked, and the depressions which normally exist each side of the patellar ligament become obliterated.

The *Common and Superficial Femoral Arteries*.—With
 Fig. xxiii.
 the thigh flexed, everted, and slightly abducted, these vessels correspond in direction to a line drawn from a point midway between the anterior superior iliac spine and the symphysis pubis to the adductor tubercle of the femur below.

Fig. xxiii., 5. The upper $1\frac{1}{2}$ inches of this line = the common femoral artery.

Fig. xxiii.,
5, 11. The upper third = the common and superficial femoral arteries in Scarpa's triangle.

Fig. xxiii.,
5, 11, 12. The upper two-thirds = the complete common and superficial femoral arteries.

Fig. xxiii.,
12. The middle third = the superficial femoral artery in Hunter's canal.

The *femoral ring*, through which a femoral hernia commonly escapes from the abdominal cavity, lies below the inner part of Poupart's ligament, and external to the pubic spine. A good way to define the ring with precision is that recommended by Holden: "Feel for the pulsation of the common femoral artery, allow $\frac{1}{2}$ inch on the inner side for the femoral vein, then comes the femoral ring." The crural or femoral canal is $\frac{1}{2}$ inch long.

The *anterior crural nerve* emerges from under cover of Poupart's ligament, about half-way between the anterior superior iliac spine and the pubic spine. The nerve lies nearly $\frac{1}{2}$ inch external to the common femoral artery, and the same distance external to the femoral sheath.

The *saphenous opening* is oval in shape, being 1 inch long and $\frac{1}{2}$ to $\frac{3}{4}$ inch broad, the long axis vertical. The central point of the opening is situated $1\frac{1}{2}$ inches below and $1\frac{1}{2}$ inches external to the pubic spine.

The *internal saphenous vein* arises from the inner side of the venous arch found on the dorsum of the foot. It passes upwards in front of the internal malleolus, along the inner side of the leg and knee, and its further upward course in the thigh is indicated by a line drawn from the adductor tubercle to the saphenous opening. Attention has previously been drawn to the elevation below the internal tuberosity of the tibia which




FIG. XXIII.

1. The anterior superior iliac spine.
- 2, 2. The pubic spines.
3. Poupart's ligament.
4. The anterior crural nerve.
5. The common femoral artery.
6. The common femoral vein.
7. The femoral ring.
8. The saphenous opening.
- 9, 9, 9. The internal saphenous vein.
10. The profunda femoris artery.
11. The superficial femoral in Scarpa's triangle.
12. The superficial femoral in Hunter's canal.
13. The adductor magnus tendon.
14. The adductor tubercle.
15. The lower epiphysial line of the femur.
16. The line of the knee-joint.
17. The gracilis, sartorius, and semitendinosus.
18. The internal saphenous nerve.
19. The posterior tibial artery.
20. The internal plantar artery.
21. The external plantar artery.
22. The anterior tibial nerve.





FIG. XXIII.

To face p. 62.



is formed by the sartorius, gracilis, and semitendinosus muscles, and below this prominence the saphenous vein is accompanied by the *internal saphenous nerve*, a branch of the deep division of the anterior crural. The saphenous nerve runs down the leg with the vein, in front of the internal malleolus of the tibia, and extends as far forwards as the ball of the great-toe. In the thigh the nerve crosses in front of the superficial femoral artery from without inwards, and it accompanies that artery throughout the whole length of Hunter's canal.

The *popliteal artery* enters the upper angle of the popliteal space from the inner side by passing between the femur and the adductor magnus tendon. The vessel at first passes obliquely outwards and downwards to the mid-point of the space, and then changes direction by passing vertically downwards as far as the lower border of the popliteus muscle, at which level it bifurcates into anterior and posterior tibial arteries. The point of bifurcation corresponds to the level of the tubercle of the tibia.

The *Ankle and Foot*.—The external malleolus projects about $\frac{1}{2}$ inch below the internal, and it also lies on a more posterior plane. The ankle-joint corresponds in level to a point about $\frac{1}{2}$ inch above the tip of the internal malleolus.

About 1 inch below and $\frac{1}{2}$ inch in front of the styloid process of the external malleolus is the *peroneal tubercle*, which separates the peroneus brevis above from the longus below. The two peronei tendons, when traced upwards, are found to pass behind the external malleolus. About 1 inch in front of the peroneal tubercle is the prominent styloid process of the base of the fifth metatarsal bone to which the peroneus brevis is attached. Between the peroneal tubercle and the base of the fifth metatarsal bone the

cuboid bone may be felt, grooved on its outer and under aspect by the peroneus longus tendon. The tendon crosses the plantar aspect of the foot in a forward and inward direction, to be inserted into the outer aspect of the base of the first metatarsal bone. Immediately in front of the external malleolus there is a well-marked depression, which is bounded in front by a prominence due to the fleshy mass of the extensor brevis digitorum, and above by the tendon of the peroneus tertius.

Fig. xxiv.,
3, 4.

If the floor of this depression be examined, the head of the astragalus will be felt above and to the inner side, and the head of the os calcis below and to the outer side.

Fig. xxiv., 2.

Fig. xxiv., 3.

Between the two malleoli in front of the ankle-joint four tendons can be felt. The most prominent and the innermost is the tendon of the tibialis anticus muscle. External to this follow the extensor longus hallucis, the extensor longus digitorum, and the peroneus tertius. When the foot is well extended the head of the astragalus can also be made out lying under cover of the extensor tendons.

Fig. xxv., 1.

Fig. xxiv.,
4, 10.

Immediately below the internal malleolus is the *sustentaculum tali*, grooved on its under aspect by the flexor longus hallucis tendon, and between the sustentaculum and the internal malleolus the tibialis posticus and flexor longus digitorum pass.

Fig. xxv., 6.
Fig. xxv., 7.


Fig. xxv.,
3, 5.

The tibialis posticus tendon can be traced upwards behind the internal malleolus and downwards to the scaphoid tuberosity, to which process the tendon gains its main attachment. In front of the internal malleolus there is another depression, which lies below the line of the tibialis anticus tendon, and here the head of

Fig. xxv., 3.

Fig. xxv., 4.

Fig. xxv., 1.






FIG. XXIV.

1. The external malleolus.
2. The head of the astragalus.
3. The head of the os calcis.
4. The peroneus tertius.
5. The base of the fifth metatarsal.
6. The peroneus brevis.
7. The peroneus longus.
8. The peroneal tubercle.
9. The tendo-Achilles.
10. The extensor longus digitorum.



FIG. XXV.

1. The tibialis anticus.
2. The internal malleolus.
3. The tibialis posticus.
4. The scaphoid tuberosity.
5. The flexor longus digitorum (upper part only).
6. The sustentaculum tali.
7. The flexor longus hallucis.
8. The tendo-Achilles.
9. The head of the astragalus.
10. The scapho-cuneiform joint.
11. The metatarso-cuneiform joint.
12. The metatarso-phalangeal joint.

Fig. xxv., 9. the astragalus can again be felt, especially prominent when the foot is well everted.

About 1 inch below and in front of the internal malleolus

Fig. xxv., 4. the tuberosity of the scaphoid bone forms the most prominent bony point on the inner side of the foot, and a line which joins the tip of the internal malleolus, the head of the astragalus, and the scaphoid tubercle, normally presents a slight upward convexity.

In flat-foot the head of the astragalus undergoes a downward displacement, and the line uniting the three bony points becomes straight, or even downwardly convex. A line drawn almost transversely across the foot from a point just behind the scaphoid tubercle indicates the level of the mid-tarsal joint.

Fig. xxv., 10-11. In front of the scaphoid tuberosity the internal cuneiform and the first metatarsal bone
Fig. xxv., 11-12. may be located and verified.

Behind the ankle-joint the tendo Achilles is placed, the tendon being at its narrowest at a point about
Fig. xxiv., 9. $1\frac{1}{2}$ inches above the insertion of the tendon
Fig. xxv., 8. into the posterior part of the os calcis. When distended with fluid, the synovial membrane of the ankle-joint bulges outwards, so as to obliterate the depressions that normally lie between the tendo Achilles and the two malleoli.

The *Anterior Annular Ligament of the Ankle*.—The upper
Fig. xxvi., 3. portion of this ligament, about 1 inch broad, extends transversely across the ankle from tibia to fibula. It presents two compartments only, one for the
Fig. xxvi., 6. tibialis anticus, and one for the remaining extensor tendons. The former tendon alone possesses a synovial sheath.

The lower portion of the ligament is Y-shaped, the

Fig. xxvi., 4. single limb arising from the upper and outer
 Fig. xxvi., 5. aspect of the head of the os calcis in close
 connection with the origin of the extensor brevis digitorum
 muscle. The upper limb of the divided portion becomes
 attached to the internal malleolus, whilst the lower limb
 sweeps over to the scaphoid tuberosity and to the inner side
 Fig. xxvi., of the foot. The extensor communis digitorum
 8, 8. and the peroneus tertius pass under the single
 undivided limb, and possess in this situation a common
 Fig. xxvi., synovial sheath, whilst the extensor longus
 6, 7. hallucis and the tibialis anticus pass through
 separate compartments in each limb of the divided portion
 of the ligament, and each tendon in so doing is surrounded
 by a synovial sheath, that enveloping the tibialis anticus
 tendon being continuous with the sheath already alluded
 to as enclosing the tendon under the transverse portion of
 the ligament.

The *internal annular ligament* is triangular in shape, the
 Fig. xxvii., 6. apex being attached to the internal malleolus,
 and the base to the lower margin of the os
 calcis. From the deep aspect of the ligament septa are
 given off which form separate compartments for the
 Fig. xxvii., passage of the tendons of the tibialis posticus,
 7, 8, 9. flexor longus digitorum, and flexor longus
 hallucis muscles, each tendon having its own synovial
 sheath. These three sheaths extend for about 1 inch above
 the upper limit of the annular ligament, and although the
 sheath enveloping the tibialis posticus reaches almost as
 far forwards as the scaphoid tuberosity, the other two
 sheaths usually terminate about $\frac{1}{2}$ inch below the inferior
 Fig. xxix., margin of the ligament. The flexor longus
 3, 3, 3, 3. hallucis and flexor longus digitorum have, again,
 distinct synovial sheaths just before their insertion into





FIG. XXVI.

1. The internal malleolus.]
2. The external malleolus.
3. The transverse band of the anterior annular ligament.
4. The Y-shaped band of the anterior annular ligament.
5. The head of the os calcis and the extensor brevis digitorum.
6. The tibialis anticus synovial sheath.
7. The extensor longus hallucis synovial sheath.
8. The extensor longus digitorum and peroneus tertius synovial sheath.

To face p. 66.







FIG. XXVII.

1. The transverse band.
- 2, 3. The upper and lower limbs of the Y-shaped part of the anterior annular ligament.
4. The tibialis anticus synovial sheath.
5. The internal malleolus.
6. The internal annular ligament.
7. The tibialis posticus synovial sheath.
8. The flexor longus digitorum synovial sheath.
9. The flexor longus hallucis synovial sheath.

To follow Fig. xxvi.



the distal phalanges of the toes, these sheaths, however, being very variable, and rarely extending further backwards than the heads of the metatarsal bones.

The *external annular ligament* is less definite in shape, and it can only be described as a broad band passing from the external malleolus to the lower margin of the os calcis.

Fig. xxviii.,
4, 5, 6. Beneath it two tendons pass, the peroneus longus and brevis. These two tendons possess a common synovial sheath, which extends upwards 2 to 3 inches above the tip of the external malleolus, and downwards as far as the "peroneal tubercle," where the sac divides into two, one part accompanying the peroneus brevis to near the base of the fifth metatarsal bone, the other extending forwards to the outer and under aspect of the cuboid bone. The peroneus longus is also usually enclosed in a synovial sheath in the last inch or so of its course previous to its insertion into the outer aspect of the base of the first metatarsal bone.

The *Anterior Tibial Artery*.—The course of this vessel may be indicated by a line drawn from a point just below the level of the tibial tubercle and midway between the outer tuberosity of the tibia and the head of the fibula to a second point in front of the ankle midway between the two malleoli, at which latter level the artery lies between the tendons of the extensor longus hallucis and longus digitorum muscles.


Fig. xxi.,
21, 21.
Fig. xxii.,
14, 14. The anterior tibial artery is continued onwards as the *dorsalis pedis* as far as the base of the first interosseous space.

The *posterior tibial artery* starts at the lower border of the popliteus muscle as one of the terminal branches of the popliteal artery. It can be represented by a line which starts at the inferior angle of

the popliteal space, on a level with the tubercle of the tibia, and which passes downwards and inwards to the mid-point between the posterior border of the internal malleolus and the inner border of the os calcis. At this level it lies under cover of the internal annular ligament, and it bifurcates in this situation into the internal and external plantar arteries. Behind the internal malleolus the posterior tibial artery lies between the tendons of the flexor longus hallucis and longus digitorum muscles, but on a slightly superficial plane.

The *small internal plantar artery* passes forwards to the cleft between the first and second toes, whilst the more important *external plantar artery* is first directed forwards and outwards towards the base of the fifth metatarsal bone, and then, changing direction, it passes forwards and inwards to the base of the first interosseous space, forming in this latter part of its course the *deep plantar arch*. It anastomoses with the *dorsalis pedis artery*, which dips downwards between the two heads of the first dorsal interosseous muscle.

The *internal popliteal nerve* crosses the popliteal artery superficially from without inwards; its onward continuation, the posterior tibial nerve, and the two terminal branches of the posterior tibial nerve, the internal and external plantars, all have the same surface marking as the corresponding arteries. Two points, however, need to be borne in mind: firstly, the posterior tibial nerve crosses the corresponding artery superficially from within outwards and downwards; and, secondly, the internal plantar nerve is relatively much more important than the artery of the same name.






FIG. XXVIII.

1. The transverse limb.
2. The stem of the Y-shaped part of the anterior annular ligament.
3. The extensor digitorum and peroneus tertius synovial sheath.
4. The peroneus longus and brevis synovial sheath.
5. The peroneus brevis sheath.




FIG. XXIX.

1. The external plantar artery.
2. The internal plantar artery.
- 3, 3. The distal flexor synovial sheaths.

To face p. 69.

The *external popliteal or peroneal nerve* was last seen to lie under cover of the biceps femoris tendon at the upper and outer boundary of the popliteal space. The nerve follows the tendon downwards to the head of the fibula, and curls round to the antero-external aspect of the leg about 1 inch below the head of that bone, dividing there into its two terminal branches, anterior tibial and musculo-cutaneous.

Fig. xx.,
11.
Fig. xxii.,
11.

Fig. xxii.,
12, 13.

The *anterior tibial nerve* passes downwards and inwards to join the corresponding artery, lying external to the upper third of the artery, superficial to the middle third, and external again to the lower third.

Fig. xxi.,
19, 19.

Fig. xxii.,
12, 12.

The nerve extends forwards along the outer side of the dorsalis pedis artery as far as the cleft between the first and second toes, the contiguous sides of which toes it supplies.

The *musculo-cutaneous nerve* running down in the substance of the peronei muscles becomes cutaneous below the middle of the leg. It then passes obliquely downwards and inwards across the anterior annular ligament, to be distributed to the greater part of the dorsum of the foot.

Fig. xxi.,
20.

Fig. xxii.,
13.

The *external saphenous vein* arises from the outer side of the venous arch on the dorsum of the foot, passing upwards behind the external malleolus and along the outer and back part of the leg to the middle of the popliteal space, where it pierces the deep fascia to open into the popliteal vein. It is accompanied in the greater part of its course by the *external saphenous nerve*, which extends forwards on the outer side of the foot as far as the tip of the little toe.

Fig. xx.,
18.

Fig. xxii.,
17.

APPENDIX

THE LENGTH OF VARIOUS PASSAGES TUBES, ETC.

THE spinal cord, 16 to 18 inches.

The trachea, $4\frac{1}{2}$ inches.

The right bronchus, 1 inch.

The left bronchus, $1\frac{1}{2}$ to 2 inches.

The pharynx, $4\frac{1}{2}$ inches.

The œsophagus, 9 inches.

The stomach :

Capacity, about 2 pints.

Length, 10 inches.

Width, 4 to 5 inches.

Duodenum, 8 to 10 inches.

Bile-duct, 3 inches.

Small intestine, 23 feet.

Jejunum, upper two-fifths.

Ileum, lower three-fifths.

Appendix, 3 to 4 inches.

Cæcum, $2\frac{1}{2}$ inches.

Ascending colon, 8 inches.

Transverse colon, 20 inches.

Descending colon, 4 to 6 inches.

Iliac colon, 5 to 6 inches.

- Pelvic colon, 16 to 18 inches.
 Rectum, 5 to 6 inches.
 Anal canal, 1 to 1½ inches.
 Crural canal, ½ inch.
 Inguinal canal, 1½ inches.
 Receptaculum chyli, 1 to 2 inches.
 Thoracic duct, 16 to 18 inches.
 Kidney, 4½ inches by 2½ inches by 1½ inches.
 Ureter, 10 inches.
 Male urethra, 8 to 10 inches.
 Prostatic, 1 to 1½ inches.
 Membranous anterior wall, ¾ inch.
 Membranous posterior wall, ½ inch.
 Spongy and penile, 6 to 8 inches.
 Testis, 1½ inches by 1 inch by ¾ inch.
 Seminiferous tubules, 2 to 3 feet.
 Canal of the epididymis, 19 to 20 feet.
 Vas deferens, 16 to 18 inches.
 Ovary, 1 inch by ½ inch.
 Fallopian tubes, 4 to 4½ inches.
 Uterus, 3 inches by 2 inches by 1 inch.
 Vagina, anterior wall, 3 inches.
 Vagina, posterior wall, 4 inches.
 Female urethra, 1 to 1½ inches.

THE WEIGHT OF SOME ORGANS.

- The brain : Male, 50 ounces ; female, 45 ounces.
 The lungs : Together, 42 ounces ; right, 22 ounces ;
 left, 20 ounces.
 The heart : Male, 10 to 12 ounces ; female, 8 to
 10 ounces.
 The liver, 50 to 60 ounces.

- The kidneys, $4\frac{1}{2}$ ounces.
The suprarenals, 1 to 2 drachms.
The prostate, 6 drachms.
The testis, 6 to 8 drachms.
The ovary, 1 to 2 drachms.
The spinal cord, $1\frac{1}{2}$ ounces.
The pancreas, 2 to 4 ounces.
The spleen, 7 ounces.

INDEX

- ABDOMINAL AORTA, 52
 planes, 40
 regions, 42
 Acromial spine and process, 15
 Acromio-clavicular articulation,
 15
 Adductor magnus tendon, 60
 tubercle, 60
 Anatomical 'snuff-box,' 23
 Ankle region, 63
 Antecubital space, 18
 Anterior annular ligament (ankle),
 65
 Anterior annular ligament (wrist),
 22
 Anterior crural nerve, 62
 tibial artery, 10
 nerve, 67
 Antrum of Highmore, 68
 Aortic arch, 30
 intercostals, 33
 valve, 30
 Apex beat, 29
 Apices of lung, 37
 Appendix, 47
 Ascending aorta, 30
 colon, 48
 Astragalus, 64, 65
 Auriculo-temporal nerve, 10
 Axilla, 16
 Axillary artery, 19

 Base of brain, 3
 Basic fossæ, 7
 Basilic vein, 18
 Biceps femoris, 59
 humeri, 16-18
 Bicipital fascia, 18
 groove, 16
 sulci, 18
 Bile-duct, 55
 Bony points of elbow, 17
 Brachial artery, 18, 19
 plexus, 13
 Bregma, 5

 Broca's area, 2
 Bronchi, 38
 Bryant's triangle, 58

 Cæcum, 47
 Cardiac orifice, 45
 Carotid tubercle, 11
 Central point, 41
 Cephalic vein, 18, 19
 Cerebellum, 4
 Cerebrum, 3
 Cervical sympathetic, 11
 Circumflex nerve and artery, 21
 Cœliac axis, 52
 Common carotid artery, 10, 31
 femoral artery, 61
 iliac artery, 52
 Coraco-brachialis fold, 16
 Coracoid process, 15
 Coronal suture, 2
 Creases of palm and fingers, 27
 of wrist, 21
 Cricoid cartilage, 11
 Cuboid bone, 64
 Cuneiform bone, 22

 Deep cardiac dulness, 37
 epigastric artery, 52
 palmar arch, 26
 plantar arch, 68
 Deltoid, 15
 Descending aorta, 30
 colon, 48
 Diaphragm, 54
 Digital arteries, 26
 Dorsalis pedis artery, 67, 69
 Douglas's fold, 43, 53
 Duodeno-jejunal fold, 46
 Duodenum, 46

 Elbow region, 17
 Epiglottis, 14
 Epiphysial line of humerus, 16, 17
 femur, 60
 Epitrochlear gland, 19

- Extensor tendons of foot, 64
 External abdominal ring, 44
 angular frontal process, 2
 annular ligament (foot), 65
 carotid artery, 10
 iliac artery, 52
 jugular vein, 10
 lateral ligament (knee), 60
 plantar artery and nerve, 68
 popliteal nerve, 59, 69
 saphenous nerve and vein, 69
- Facial artery, 9, 11
 nerve, 7, 8
 vein, 9
- Falx cerebri, 1
 Femoral canal and ring, 62
 Fibula, 60, 69
 Filum terminale, 56
 Fissures of lung, 36
 Flexor carpi radialis, 21
 ulnaris, 21
 sublimis, 21
 synovial sheaths, 22
 Fontanelle, anterior, 2
 posterior, 2
 Foot region, 63
 Foramen of Winslow, 54
 Frontal sinus, 7
- Gall-bladder, 54
 Gastro-hepatic omentum, 54
 Gluteal artery, 59
 Gluteus maximus, 57
 Gracilis, 60, 61
 Great auricular nerve, 12
 occipital nerve, 10
 sciatic nerve, 59
- Hamilton's test, 15
 Hand region, 21
 Heart, 28
 Hepatic artery, 55
 flexure, 48
 Hesselbach's triangle, 53
- Ileo-cæcal valve, 47
 Ileo-pelvic colon, 49
 Iliac colon, 49
 crest, 43
 spines, 43, 59
 Ilio-tibial band, 60
 Inferior dental nerve and artery, 9
 mesenteric artery, 52
 vena cava, 32, 53
 Infra-orbital foramen, 9
 Inguinal canal, 44
 Inion, 1
 Innominate artery, 31
 veins, 32
- Internal abdominal ring, 44, 52
 annular ligament, 66
 carotid artery, 10
 jugular vein, 11
 mammary artery, 32
 plantar artery and nerve, 68
 popliteal nerve, 68
 pudic artery, 59
 saphenous nerve, 63
 vein, 62
- Intertubercular plane, 41
 Ischial spine, 59
 tuberosity, 57
 Island of Reil, 4
- Kidney, 50
 Knee region, 59
 Knuckles, 27
- Lachrymal sac, 9
 Lambda, 2
 Larynx, 11
 Lateral sinus, 3, 7
 ventricles, 6
 vertical plane, 40
 Length of femur, 58
 of humerus, 15
 of tibia, 58
 Ligamentum patellæ, 61
 teres, 53
 Lineæ semilunares, 43
 transversæ, 43
 Lingual artery, 11
 Liver, 53
 Ludwig's plane, 42
 Lungs, 34
- Macewen's triangle, 7
 Malar tubercle, 2
 Mamma, 28
 Mastoid antrum, 7
 McBurney's point, 47
 Median nerve, 18, 20, 25
 Median vertical plane, 40
 Mental foramen, 9
 Mesentery of small intestine, 47
 Mesial fissure of brain, 1
 Metopic suture, 2
 Middle cerebral artery, 4
 meningeal artery, 5, 6
 Mitral valve, 30
 Morris's quadrilateral, 50
 Musculo-cutaneous nerve (arm),
 20
 Musculo-cutaneous nerve (leg), 69
 Musculo-spiral nerve, 18, 20
- Nasal duct, 8
 Nasion, 1
 Natal fold, 57

- Nélaton's line, 58
Nipple, 28
- Occipital artery, 10, 11
 nerve (great), 10, 11
 (small), 12
- Olecranon, 17
Omo-hyoid muscle, 11
Os calcis, 64
Ovary, 51
- Palmar fascia, 21
Palmaris longus, 27
Pancreas, 46
Paracentral point, 41
Parieto-occipital sulcus, 4
Parotid gland, 8
Pectoralis minor, 19
Peroneal nerve, 59, 69
 tendons, 63, 64
 tubercle, 63
Petit's triangle, 44
Pharynx, 11
Phrenic nerve, 12
Pisiform bone, 21, 22
Plantar arch, 68
Pleura, 33
Popliteal artery, 63
 space, 59
Portal vein, 55
Posterior annular ligament (wrist),
 24
 auricular nerve, 10
 interosseous nerve, 18, 25
 tibial artery, 67
 nerve, 68
Prepatellar bursa, 61
Pubic spine, 44
Pulmonary artery, 31
 valve, 30
Pylorus, 45
- Radial artery, 20, 22, 25
 nerve, 18, 25
 tubercle, 24
Radius, 17
Receptaculum chyli, 39
Rectum, 49
Reid's base line, 4
Renal artery, 52
Rima glottidis, 14
Rolandic area, 5
 fissure, 5
Roots of lung, 37
- Sagittal suture, 2
Saphenous opening, 62
Sartorius, 60, 61
Scalenus anticus, 13
Scaphoid bone (carpus), 22, 23
 (tarsus), 65
- Sciatic artery, 59
 nerve (great), 59
 (small), 59
Semilunar fold of Douglas, 43
Semimembranosus, 60
 tendinosus, 60, 61
Shoulder region, 15
Small intestine, 46
 occipital nerve, 13
 sciatic nerve, 59
Spinal accessory nerve, 12
 cord, 55
Spinous plane, 42
Spleen, 55
Splenic flexure, 48
Stenson's duct, 8
Sterno-mastoid, 13
Sterno-xiphoid plane, 42
Stomach, 45
Subclavian artery, 13, 31
Subcostal plane, 42
Superficial cardiac dulness, 37
 cervical plexus, 12, 13
 femoral artery, 61
 palmar arch, 26
 temporal artery, 10
Superior longitudinal sinus, 1
 mediastinum, 31
 mesenteric artery, 52
 thyroid artery, 11
 vena cava, 32
Suprameatal crest or spine, 2, 7
 fossa, 7
Supra-orbital foramen, 9
Suprasternal plane, 42
Sustentaculum tali, 64
Sylvian fissure, 4
 point, 4
Sympathetic chain and ganglia, 11
Synovial membrane of knee, 61
 sheaths of ankle, 65, 66
 of wrist (extensor), 24
 (flexor), 22
- Tegmen antri, 7
Temporo-sphenoidal sulcus, 4
Tendo-Achilles, 65
 oculi, 8
Tendons in front of ankle, 64
Tentorium cerebelli, 4
Theca vertebralis, 56
Thoracic aorta, 30
 duct, 38
 plane, 41
Thyroid isthmus, 14
Tibial tuberosities, 61
Tibialis anticus, 64, 65
 posticus, 64
Tonsil, 10
Trachea, 11, 38
Transpyloric plane, 41

- Transverse cervical nerve, 13
 colon, 48
 creases of palm, 27
 of wrist, 22
 facial artery, 8
 process of atlas, 12
Trapezium, 24
Tricuspid valve, 30
Trochanter of femur, 57
Tuberosities of humerus, 15
- Ulnar artery, 20, 26
 nerve, 18, 21, 26
Umbilical plane, 42
Umbilicus, 43
- Unciform bone, 22
Urachus, 51
Ureter, 51
- Vagus nerve, 11
Valves of heart, 30
Veins of elbow, 18
Vena azygos major, 33
Vermiform appendix, 47
Vocal cords, 14
- Wrist region, 21
- Zygomatic process, 2

THE END

November, 1904.

CATALOGUE OF WORKS

PUBLISHED BY

H. K. LEWIS,

136 GOWER STREET, LONDON, W.C.

ESTABLISHED 1844.

Telegrams: PUBLICAVIT, LONDON.

Telephone: 10721 CENTRAL.

A. C. ABBOTT, M.D.

Professor of Hygiene and Bacteriology, University of Pennsylvania.

THE PRINCIPLES OF BACTERIOLOGY: A Practical Manual
for Students and Physicians. Sixth Edition, with 111 Illustrations, 26 being
coloured, post 8vo, 12s. 6d. *nett*.

H. ALDERSMITH, M.B. LOND., F.R.C.S.

Medical Officer, Christ's Hospital, West Horsham.

RINGWORM AND ALOPECIA AREATA: Their Pathology,
Diagnosis, and Treatment. Fourth Edition, enlarged, with new Illustrations,
demy 8vo, 10s. 6d.

IZETT ANDERSON, M.D. EDIN.

*Extraordinary Member of the Royal Medical Society of Edinburgh Honorary Member of the Institute
of Jamaica.*

YELLOW FEVER IN THE WEST INDIES. Crown 8vo, 3s. 6d.

JAMES ANDERSON, M.D., F.R.C.P.

Late Assistant Physician to the London Hospital, &c.

NOTES ON MEDICAL NURSING from the Lectures given
to the Probationers at the London Hospital. Edited by E. F. LAMPORT, Asso-
ciate of the Sanitary Institute. With an Introductory Biographical Notice by
the late SIR ANDREW CLARK, BART. Third Edition, with Glossary, crown 8vo,
2s. 6d.

G. GRANVILLE BANTOCK, M.D., F.R.C.S. EDIN.

Surgeon to the Samaritan Free Hospital for Women and Children.

ON THE TREATMENT OF RUPTURE OF THE FEMALE
Perineum Immediate and Remote. Second Edition, with Illustrations, 8vo,
3s. 6d.

A PLEA FOR EARLY OVARIOTOMY. Demy 8vo, 2s.

FANCOURT BARNES, M.D., M.R.C.P.

Physician to the Chelsea Hospital for Women; Obstetric Physician to the Great Northern Hospital, &c.

A GERMAN-ENGLISH DICTIONARY OF WORDS AND Terms used in Medicine and its Cognate Sciences. Square 12mo, Roxburgh binding, 9s.

JAMES BARR, M.D.

Physician to the Northern Hospital, Liverpool; Medical Officer of Her Majesty's Prison, Kirkdale, &c.

THE TREATMENT OF TYPHOID FEVER, and reports of fifty-five consecutive cases with only one death. With Introduction by W. T. GAIRDNER, M.D., LL.D., Professor of Medicine in the University of Glasgow. With Illustrations, demy 8vo, 6s.

ASHLEY W. BARRETT, M.B. LOND., M.R.C.S., L.D.S.E.

Consulting Dental Surgeon to the London Hospital; Examiner in Dental Surgery to the Royal College of Surgeons, England.

DENTAL SURGERY FOR MEDICAL PRACTITIONERS AND Students of Medicine. Third Edition, with Illustrations, crown 8vo, 3s. 6d.
[LEWIS'S PRACTICAL SERIES.]

H. CHARLTON BASTIAN, M.A., M.D., F.R.S., F.R.C.P.

Emeritus Professor of the Principles and Practice of Medicine in University College, London; Consulting Physician to University College Hospital, &c.

I.
A TREATISE ON APHASIA AND OTHER SPEECH DEFECTS.
With illustrations, medium 8vo, 15s.

II.
PARALYSES: CEREBRAL, BULBAR, AND SPINAL. A Manual of Diagnosis for Students and Practitioners. With numerous Illustrations, 8vo, 12s. 6d.

III.
VARIOUS FORMS OF HYSTERICAL OR FUNCTIONAL Paralysis. Demy 8vo, 7s. 6d.

W. BATESON, M.A., F.R.S.

MENDEL'S PRINCIPLES OF HEREDITY. A Defence, with a Translation of Mendel's original Paper on Hybridisation. Crown 8vo, 4s. *nett.*

W. M. BEAUMONT.

Surgeon to the Bath Eye Infirmary.

THE SHADOW-TEST IN THE DIAGNOSIS AND ESTIMATION of Ametropia. Post 8vo, 2s. 6d.

F. E. BEDDARD, M.A., F.R.S.

[See Cambridge Biological Series, page 5.]

CHARLES E. BEEVOR, M.D. LOND., F.R.C.P.

Physician to the National Hospital for the Paralysed and Epileptic, the Great Northern Central Hospital, and the National Orthopædic Hospital.

DISEASES OF THE NERVOUS SYSTEM. A Handbook for Students and Practitioners. With Illustrations, crown 8vo, 10s. 6d.

[LEWIS'S PRACTICAL SERIES.]

A. HUGHES BENNETT, M.D., M.R.C.P.

Physician to the Hospital for Epilepsy and Paralysis, Regent's Park, &c.

A PRACTICAL TREATISE ON ELECTRO-DIAGNOSIS IN Diseases of the Nervous System. With Illustrations, 8vo, 8s. 6d.

E. H. BENNETT, M.D., F.R.C.S.I.

Professor of Surgery, University of Dublin,

AND

D. J. CUNNINGHAM, M.D., F.R.C.S.I.

Professor of Anatomy and Chirurgery, University of Dublin.

THE SECTIONAL ANATOMY OF CONGENITAL CŒCAL Hernia. With coloured plates, sm. folio, 5s. 6d.

HORATIO R. BIGELOW, M.D.

Permanent Member of the American Medical Association, &c.

PLAIN TALKS ON ELECTRICITY AND BATTERIES, WITH Therapeutic Index. With Illustrations, crown 8vo, 4s. 6d.

DRS. BOURNEVILLE AND BRICON.

MANUAL OF HYPODERMIC MEDICATION.

Translated from the Second Edition, and Edited, with Therapeutic Index of Diseases, by ANDREW S. CURRIE, M.D. Edin., &c. With Illustrations, crown 8vo, 3s. 6d.

RUBERT BOYCE, F.R.S., M.B., M.R.C.S.

Professor of Pathology in University College, Liverpool.

A TEXT-BOOK OF MORBID HISTOLOGY for Students and Practitioners. With 130 coloured Illustrations, royal 8vo, 31s. 6d.

A. BROCA, M.D.

Chirurgien des Hôpitaux de Paris, &c.

AND

F. LUBET-BARBON, M.D.

Ancien interne des Hôpitaux de Paris.

MASTOID ABSCESSSES AND THEIR TREATMENT.

Translated and edited by HENRY J. CURTIS, B.S. and M.D. (Lond.), F.R.C.S. (Eng.), formerly Assistant to the Professor of Pathology, University College, London, &c. With coloured Illustrations, cr. 8vo, 6s.

O. A. BROWNE, M.A., M.D.

Physician to the Royal Hospital for Diseases of the Chest, &c.

ANEURYSMS OF THE AORTA. 4to, 2s. 6d.

MARY BULLAR & J. F. BULLAR, M.B. CANTAB., F.R.C.S.

RECEIPTS FOR FLUID FOODS. 16mo, 1s.

CHARLES H. BURNETT, A.M., M.D.

Emeritus Professor of Otology in the Philadelphia Polyclinic; Clinical Professor of Otology in the Woman's Medical College of Pennsylvania, &c.

SYSTEM OF DISEASES OF THE EAR, NOSE AND THROAT.

By 45 Eminent American, British, Canadian and Spanish Authors. Edited by CHARLES H. BURNETT. With Illustrations, in two imperial 8vo vols., half morocco, 48s. *nett.*

DUDLEY W. BUXTON, M.D., B.S., M.R.C.P.

Administrator of Anæsthetics and Lecturer in University College Hospital; Consulting Anæsthetist to the National Hospital for Paralysis and Epilepsy, Queen's Square, &c.

ANÆSTHETICS: THEIR USES AND ADMINISTRATION.

Fourth Edition, with Illustrations, crown 8vo. [LEWIS'S PRACTICAL SERIES.]
[In the Press.]

JAMES CALVERT, B.A., B.SC., M.D. LOND.

Fellow of the Royal College of Physicians; Lecturer on Materia Medica, Pharmacology, and Therapeutics to St. Bartholomew's Hospital.

PRACTICAL PHARMACY AND PRESCRIBING FOR STUDENTS OF MEDICINE.

Being the Course in Use at St. Bartholomew's Hospital. Second Edition, crown 8vo, interleaved, 4s. 6d.

HARRY CAMPBELL, M.D., B.S. LOND., F.R.C.P.

Physician to the North-West London Hospital.

I.
THE CAUSATION OF DISEASE: An Exposition of the ultimate factors which induce it. Demy 8vo, 12s. 6d.

II.
FLUSHING AND MORBID BLUSHING: Their Pathology and Treatment. With plates and wood engravings, royal 8vo, 10s. 6d.

III.
DIFFERENCES IN THE NERVOUS ORGANISATION OF Man and Woman, Physiological and Pathological. Royal 8vo, 15s.

IV.
HEADACHE AND OTHER MORBID CEPHALIC SENSATIONS. Royal 8vo, 12s. 6d.

CAMBRIDGE BIOLOGICAL SERIES.*(General Editor: A. E. SHIPLEY, M.A., Fellow and Tutor of Christ's College).*

- THE VERTEBRATE SKELETON.** By S. H. REYNOLDS, M.A. Trinity College, Cambridge; Lecturer and Demonstrator in Geology and Zoology at University College, Bristol. With numerous Illustrations, crown 8vo, 12s. 6d.
- PRACTICAL MORBID ANATOMY.** By H. D. ROLLESTON, M.D., F.R.C.P., Fellow of St. John's College, Cambridge; Assistant Physician and Lecturer on Pathology, St. George's Hospital, London; and A. A. KANTHACK, M.D., M.R.C.P., Lecturer on Pathology, St. Bartholomew's Hospital, London. Crown 8vo, 6s.
- ZOOGEOGRAPHY.** By F. E. BEDDARD, M.A., F.R.S. With Maps, crown 8vo, 6s.
- PRACTICAL PHYSIOLOGY OF PLANTS.** By F. DARWIN, M.A., F.R.S., and E. H. ACTON, M.A. Third Edition, with Illustrations, crown 8vo, 4s. 6d.
- ELEMENTS OF BOTANY.** By F. DARWIN, M.A., F.R.S. Second Edition, with numerous Illustrations, crown 8vo, 4s. 6d.
- A MANUAL AND DICTIONARY OF THE FLOWERING PLANTS AND Ferns: Morphology, Natural History and Classification.** Alphabetically arranged by J. C. WILLIS, M.A., Director of the Royal Botanic Gardens, Ceylon, &c. Second edition, 10s. 6d. *[Just published.]*
- FOSSIL PLANTS, a Manual for Students of Botany and Geology.** By A. C. SEWARD, M.A. F.R.S., St. John's College. In two volumes, demy 8vo. Vol. I., with Illustrations, 12s. *[Vol. II. just ready.]*
- PALÆONTOLOGY—INVERTEBRATE.** By HENRY WOODS, M.A., F.G.S., University Lecturer in Palæo-zoology, Cambridge. Third Edition, revised and enlarged, with additional Illustrations, crown 8vo, 6s.
- OUTLINES OF VERTEBRATE PALÆONTOLOGY FOR STUDENTS OF Zoology.** By A. S. WOODWARD, M.A. With Illustrations, demy 8vo, 14s.
- THE SOLUBLE FERMENTS AND FERMENTATION.** By J. REYNOLDS GREEN, Sc.D., F.R.S., Professor of Botany to the Pharmaceutical Society of Great Britain, &c. Second Edition, 8vo, 12s.
- ZOOLOGY, an Elementary Textbook.** By A. E. SHIPLEY, M.A., Fellow and Tutor of Christ's College, Cambridge, and E. W. MACBRIDE, M.A. (Cantab.), D.Sc. (Lond.). With numerous Illustrations, 8vo, 10s. 6d. *net.*
- GRASSES: a Handbook for use in the Field and Laboratory.** By H. MARSHALL WARD, Sc.D., F.R.S., Crown 8vo, 6s.
- LECTURES ON THE HISTORY OF PHYSIOLOGY DURING THE SIXTEENTH, SEVENTEENTH, and EIGHTEENTH CENTURIES.** By Sir M. FOSTER, K.C.B., M.P., M.D., &c., Professor of Physiology in the University of Cambridge, &c. 8vo, 9s.
- THE NATURAL HISTORY OF SOME COMMON ANIMALS.** By O. H. LATTER, M.A., Science Master at Charterhouse School. Crown 8vo, 5s. *net.* *[Just published.]*
- THE CLASSIFICATION OF FLOWERING PLANTS.** By A. B. RENDLE, M.A. (Cantab.), D.Sc. (Lond.), Assistant in Botany, British Museum. Vol. I., Introduction, Gymnosperms, Monocotyledons. 8vo, 10s. 6d. *net.* *[Just published.]*
- TREES.** By H. M. WARD, Sc.D., F.R.S., Professor of Botany in the University of Cambridge, &c. Vol. I., Buds and Twigs. Crown 8vo, 4s. 6d. *net.* *[Just published.]*
- THE MORPHOLOGY OF PLANTS.** By J. C. WILLIS, M.A. *[In the press.]*

CAMBRIDGE GEOLOGICAL SERIES.

- HANDBOOK TO THE GEOLOGY OF CAMBRIDGESHIRE.** For the use of Students. By F. R. COWPER REED, M.A., F.G.S., Assistant to the Woodwardian Professor of Geology. Crown 8vo, 7s. 6d.
- PETROLOGY FOR STUDENTS: An Introduction to the Study of Rocks under the Microscope.** By A. HARKER, M.A., F.G.S., Fellow of St. John's College; Demonstrator in Geology (Petrology) in the University of Cambridge. Third Edition, revised, cr. 8vo, 7s. 6d.
- THE PRINCIPLES OF STRATIGRAPHICAL GEOLOGY.** By J. E. MARR, M.A., F.R.S., Fellow and Lecturer of St. John's College, Cambridge, cr. 8vo, 6s.
- A TREATISE ON CRYSTALLOGRAPHY.** By W. J. LEWIS, M.A. Professor of Mineralogy in the University of Cambridge. 8vo, 14s. *net.*

CAMBRIDGE PHYSICAL SERIES.*(General Editor: R. T. GLAZEBROOK, M.A., F.R.S., Fellow of Trinity College; Assistant Director of the Cavendish Laboratory).*

- HEAT AND LIGHT.** By R. T. GLAZEBROOK, M.A. Crown 8vo, 5s.
The two Parts are also published separately. HEAT, 3s. LIGHT, 3s.
- MECHANICS AND HYDROSTATICS.** By the same Author. Crown 8vo, 8s. 6d. Also in separate Parts. Part I.—DYNAMICS, 4s. Part II.—STATICS, 3s. Part III.—HYDROSTATICS, 3s.
- A TREATISE ON THE THEORY OF SOLUTION, INCLUDING PHENOMENA OF ELECTROLYSIS.** By W. C. D. WHETHAM, M.A., F.R.S., Fellow of Trinity College. 8vo, 10s. *net.*
- MECHANICS.** By J. COX, M.A., F.R.S.C. 8vo, 9s. *net.*
- ELECTRICITY AND MAGNETISM.** By R. T. GLAZEBROOK, M.A., F.R.S. Crown 8vo, 7s. 6d. *[Just published.]*
- CONDUCTION OF ELECTRICITY THROUGH GASES.** By J. J. THOMSON, D.Sc., LL.D., F.R.S., Fellow of Trinity College, Cambridge. 8vo, 16s. *[Just published.]*
- RADIO-ACTIVITY.** By E. RUTHERFORD, Trinity College, Cambridge, Professor of Physics at McGill University, Montreal, 8vo, 10s. 6d. *nett.* *[Just published.]*
- SOUND.** By J. W. CAPSTICK, M.A., Fellow of Trinity College. *[In preparation.]*

J. W. CAPSTICK, M.A.

[See Cambridge Physical Series, page 5.]

ALFRED H. CARTER, M.D., M.SC. LOND.

Fellow of the Royal College of Physicians; Professor of Medicine, University of Birmingham; Senior Physician to the Queen's Hospital, Birmingham, &c.

ELEMENTS OF PRACTICAL MEDICINE.

Eighth Edition, crown 8vo, 10s. 6d.

F. H. CHAMPNEYS, M.A., M.D. OXON., F.R.C.P.

Physician-Accoucheur and Lecturer on Obstetric Medicine at St. Bartholomew's Hospital; Examiner in Obstetric Medicine in the University of Oxford, &c.

I.

LECTURES ON PAINFUL MENSTRUATION. The Harveian Lectures, 1890. Roy. 8vo, 7s. 6d.

II.

EXPERIMENTAL RESEARCHES IN ARTIFICIAL RESPIRATION in Stillborn Children, and Allied Subjects. Crown 8vo, 3s. 6d.

W. BRUCE CLARKE, M.A., M.B. OXON., F.R.C.S.

Assistant Surgeon to, and Senior Demonstrator of Anatomy and Operative Surgery at St. Bartholomew's Hospital; Surgeon to the West London Hospital, &c.

THE DIAGNOSIS AND TREATMENT OF DISEASES OF THE Kidney Amenable to direct Surgical Interference. With Illustrations, demy 8vo, 7s. 6d.

ALEXANDER COLLIE, M.D. ABERD., M.R.C.P. LOND.

Secretary of the Epidemiological Society for Germany and Russia, &c.

ON FEVERS: THEIR HISTORY, ETIOLOGY, DIAGNOSIS, Prognosis, and Treatment. Illustrated with Coloured Plates, crown 8vo, 8s. 6d.

[LEWIS'S PRACTICAL SERIES.]

E. TREACHER COLLINS, F.R.C.S.

Assistant Surgeon to the Royal London Ophthalmic Hospital, Moorfields; Hunterian Professor, Royal College of Surgeons, England, 1893-94.

RESEARCHES INTO THE ANATOMY AND PATHOLOGY OF the Eye. With 10 Plates and 28 Figures in the Text, demy 8vo, 6s.

WALTER S. COLMAN, M.D., F.R.C.P. LOND.

Assistant Physician to the National Hospital for the Paralysed and Epileptic, &c.

SECTION CUTTING AND STAINING: A Practical Introduction to Histological Methods for Students and Practitioners. Second Edition, with Illustrations, crown 8vo, 3s. 6d.

W. H. CORFIELD, M.A., M.D. OXON., F.R.C.P. LOND.

Consulting Sanitary Adviser to H. M. Office of Works; Hon. Sanitary Adviser to University College Hospital; Professor of Hygiene and Public Health in University College, London; Medical Officer of Health for St. George's, Hanover Square, &c.

I.

DWELLING HOUSES: their Sanitary Construction and Arrangements. Fourth Edition, with Illustrations, crown 8vo, 3s. 6d.

II.

DISEASE AND DEFECTIVE HOUSE SANITATION: Being Two Lectures delivered before the Harveian Society of London. With Illustrations, crown 8vo, 2s.

III.

THE ETIOLOGY OF TYPHOID FEVER AND ITS PREVENTION. Being the Milroy Lectures delivered at the Royal College of Physicians, 1902. Demy 8vo, 2s. 6d.

J. COX, M.A., F.R.C.S.

See Cambridge Physical Series, page 5.

SIDNEY COUPLAND, M.D., F.R.C.P.

Physician to the Middlesex Hospital, and Lecturer on Practical Medicine in the Medical School; late Examiner in Medicine at the Examining Board for England.

NOTES ON THE CLINICAL EXAMINATION OF THE BLOOD and Excreta. Third Edition, 12mo, 1s. 6d.

CHARLES CREIGHTON, M.A., M.D.

Formerly Demonstrator of Anatomy in the University of Cambridge.

I.

A HISTORY OF EPIDEMICS IN BRITAIN. Vol. I., From A.D. 664 to the Extinction of the Plague. Demy 8vo, 18s. Vol. II., From the Extinction of the Plague to the Present Time. Demy 8vo, 20s.

II.

ILLUSTRATIONS OF UNCONSCIOUS MEMORY IN DISEASE, including a Theory of Alteratives. Post 8vo, 6s.

H. RADCLIFFE-CROCKER, M.D. LOND., F.R.C.P.

Physician for Diseases of the Skin in University College Hospital, &c.

I.

DISEASES OF THE SKIN; THEIR DESCRIPTION, PATHOLOGY, Diagnosis, and Treatment. With special Reference to the Skin Eruptions of Children, and an Analysis of Fifteen Thousand Cases of Skin Disease. Third Edition, with 4 Plates and 112 Illustrations, 2 vols., med. 8vo, 28s. *nett.*
[Now ready.]

II.

THE CONDITIONS WHICH MODIFY THE CHARACTERS of Inflammations of the Skin, and their Influence on Treatment. Being the Lettsomian Lectures at the Medical Society of London, 1903. 8vo, 2s. 6d.
[Now ready.]

F. DARWIN, M.A., F.R.S.

[See Cambridge Biological Series, page 5.]

HERBERT DAVIES, M.D., F.R.C.P.

Late Consulting Physician to the London Hospital.

THE MECHANISM OF THE CIRCULATION OF THE BLOOD
through Organically Diseased Hearts. Edited by ARTHUR TEMPLER DAVIES,
B.A. (Nat. Science Honours), M.D. Cantab., F.R.C.P., Physician to the Royal
Hospital for Diseases of the Chest. Crown 8vo, 3s. 6d.

HORACE DOBELL, M.D.

Consulting Physician to the Royal Hospital for Diseases of the Chest, &c.

ON DIET AND REGIMEN IN SICKNESS AND HEALTH,
and on the Interdependence and Prevention of Diseases and the Diminution of
their Fatality. Seventh Edition, 8vo, 5s. *nett.*

ROBERT W. DOYNE, F.R.C.S.

*Surgeon to the Oxford Eye Hospital; Ophthalmic Surgeon to St. John's Hospital, Cowley, and to the
Bourton on Water Cottage Hospital.*

NOTES ON THE MORE COMMON DISEASES OF THE EYE.
With Test Types, crown 8vo, 2s.

PROF. A. DÜHRSEN, M.D.

Professor in Midwifery and Gynæcology in the University of Berlin.

I.
A MANUAL OF GYNÆCOLOGICAL PRACTICE FOR STU-
dents and Practitioners. Second English, translated and edited from the Sixth
German edition, by JOHN W. TAYLOR, F.R.C.S., Professor of Gynæcology, the
University of Birmingham, and FREDERICK EDGE, M.D. LOND., M.R.C.P.,
F.R.C.S., Surgeon to the Wolverhampton and District Hospital for Women.
With 125 Illustrations, crown 8vo, 3s. 6d. *nett.*

II.
A MANUAL OF OBSTETRIC PRACTICE FOR STUDENTS
and Practitioners. Translated and edited from the Sixth German Edition, by
JOHN W. TAYLOR and FREDERICK EDGE. With Illustrations, crown 8vo, 3s. 6d. *nett.*

ARTHUR W. EDIS, M.D. LOND., F.R.C.P.

*Senior Physician to the Chelsea Hospital for Women; Late Obstetric Physician to the
Middlesex Hospital.*

STERILITY IN WOMEN: including its Causation and Treat-
ment. With 33 Illustrations, demy 8vo, 6s.

EDWARD J. EDWARDES, M.D. LOND.

Member of the Royal College of Physicians, London.

A CONCISE HISTORY OF SMALL-POX AND VACCINATION
in Europe. Crown 8vo, 2s. 6d. *nett.*

W. ELDER, M.D., F.R.C.P. EDIN.

Physician to Leith Hospital.

APHASIA AND THE CEREBRAL SPEECH MECHANISM.

With Illustrations, demy 8vo, 10s. 6d.

W. D'ESTE EMERY, M.D., B.SC. LOND.

Assistant Bacteriologist to the Laboratories of the Royal Colleges of Physicians and Surgeons, London; formerly Lecturer on Pathology and Bacteriology in the University of Birmingham, &c.

A HANDBOOK OF BACTERIOLOGICAL DIAGNOSIS FOR Practitioners (including Instructions for the Clinical Examination of the Blood).

With 2 Coloured Plates and 33 Illustrations, crown 8vo, 5s. 6d.

[LEWIS'S PRACTICAL SERIES.]

ALEXANDER S. FAULKNER.

Surgeon-Major, Indian Medical Service.

A GUIDE TO THE PUBLIC MEDICAL SERVICES.

Compiled from Official Sources. 8vo, 2s.

W. SOLTAU FENWICK, M.D., B.S. LOND., M.R.C.P.

Physician to Out-patients at the Evelina Hospital for Sick Children; Physician to the London Temperance Hospital.

I.

DISORDERS OF DIGESTION IN INFANCY AND CHILDHOOD. With Illustrations, demy 8vo, 10s. 6d.

II.

THE DYSPEPSIA OF PHTHISIS: Its Varieties and Treatment, including a Description of Certain Forms of Dyspepsia associated with the Tubercular Diathesis. Demy 8vo, 6s.

DR. FERBER.

MODEL DIAGRAM OF THE ORGANS IN THE THORAX and Upper Part of the Abdomen. With Letter-press Description. In 4to, coloured, 5s. *nett.*

G. H. FINK, M.R.C.S., L.S.A. LOND.

Major, Indian Medical Service (Retired).

CANCER, AND PRECANCEROUS CHANGES, their Origin and Treatment. 8vo, 5s.

W. H. RUSSELL FORSBROOK, M.D. LOND., M.R.C.S.

Consulting Medical Officer to the Government of the Cape of Good Hope; formerly Surgical Registrar to Westminster Hospital.

A DISSERTATION ON OSTEO-ARTHRITIS. Demy 8vo, 5s.

SIR M. FOSTER, K.C.B., M.P., M.D., ETC.

[See Cambridge Biological Series, page 5.]

J. MILNER FOTHERGILL, M.D., M.R.C.P.

Late Physician to the City of London Hospital for Diseases of the Chest, Victoria Park, &c.

I.

INDIGESTION AND BILIOUSNESS. Second Edition, post 8vo, 7s. 6d.

II.

GOUT IN ITS PROTEAN ASPECTS. Post 8vo, 7s. 6d.

III.

THE TOWN DWELLER: His Needs and His Wants. With an Introduction by SIR B. W. RICHARDSON, M.D., LL.D., F.R.S. Post 8vo, 3s. 6d.

R. HINGSTON FOX, M.D. BRUX., M.R.C.P. LOND.

WILLIAM HUNTER: Anatomist, Physician, Obstetrician, (1718-1783). With notices of his friends CULLEN, SMELLIE, FOTHERGILL and BAILLIE. With seven Portrait-Prints, Chronological Chart of Life and Times, and View of Hunter's Birthplace, 8vo, 4s. 6d. *nett.*

PROFESSOR DR. PAUL FÜRBRINGER.

Director of the Friedrichshain Hospital, Berlin, &c.

TEXT-BOOK OF DISEASES OF THE KIDNEYS AND Genito-Urinary Organs. Translated by W. H. GILBERT, M.D., Physician in Baden-Baden, &c. Vol. I., demy 8vo, 7s. 6d. Vol. II., demy 8vo, 10s. 6d.

SIR DOUGLAS GALTON.

Late Royal Engineers, K.C.B., Hon. D.C.L., LL.D., F.R.S., Assoc. Inst. C.E., M.I.Mech.E., F.S.A., F.G.S., F.L.S., F.C.S., F.R.G.S., &c.

HEALTHY HOSPITALS. Observations on some points connected with Hospital Construction. With Illustrations, 8vo, 10s. 6d.

JOHN HENRY GARRETT, M.D.

Licentiate in Sanitary Science and Diplomat in Public Health, Universities of Durham and Cambridge, &c.

THE ACTION OF WATER ON LEAD; being an inquiry into the Cause and Mode of the Action and its Prevention. Crown 8vo, 4s. 6d.

ALFRED W. GERRARD, F.C.S.

Examiner to the Pharmaceutical Society, &c.

ELEMENTS OF MATERIA MEDICA AND PHARMACY. With Supplement (1890). Crown 8vo, 8s. 6d.

R. T. GLAZEBROOK, M.A., F.R.S.

[See Cambridge Physical Series, page 5.]

E. W. GOODALL, M.D. LOND.

*Medical Superintendent of the Eastern Hospital of the Metropolitan Asylums Board, formerly
Medical Registrar to Guy's Hospital,*

AND

J. W. WASHBOURN, C.M.G., M.D. LOND.

*Fellow of the Royal College of Physicians; Physician to the London Fever Hospital; Assistant
Physician to Guy's Hospital, and Lecturer in the Medical School.*

A MANUAL OF INFECTIOUS DISEASES. Illustrated with Plates,
Diagrams, and Charts, demy 8vo, 15s.

JAMES F. GOODHART, M.D. ABERD., F.R.C.P.

Physician to Guy's Hospital, and Consulting Physician to the Evelina Hospital for Sick Children.

ON COMMON NEUROSES: or the Neurotic Element in Disease
and its Rational Treatment. Second Edition, crown 8vo, 3s. 6d.

JOHN GORHAM, M.R.C.S.

TOOTH EXTRACTION: a Manual on the proper mode of Ex-
tracting Teeth. Fourth Edition, fcap. 8vo, 1s. 6d.

GEORGE M. GOULD, A.M., M.D.

Ophthalmic Surgeon to the Philadelphia Hospital, &c.

I.

THE STUDENT'S MEDICAL DICTIONARY: including all the
words and phrases generally used in Medicine, with their proper pronunciation
and definitions, based on recent medical literature. Eleventh Edition, with
elaborate Tables and many Illustrations, 8vo, 14s. nett.

II.

A POCKET MEDICAL DICTIONARY, Giving the Pronuncia-
tion and Definition of 30,000 of the Principal Words used in Medicine and the
Collateral Sciences. Fourth Edition, bound limp, 32mo, 5s. nett.

SIR W. R. GOWERS, M.D., F.R.C.P., M.R.C.S.

Physician to University College Hospital, &c.

DIAGRAMS FOR THE RECORD OF PHYSICAL SIGNS.
In books of 12 sets of figures, 1s.

LANDON C. GRAY, M.D.

*Professor of Nervous and Mental Diseases in the New York Polyclinic; Visiting Physician to
St. Mary's Hospital, &c.*

A TREATISE ON NERVOUS AND MENTAL DISEASES FOR
Students and Practitioners of Medicine. With 168 illustrations, 8vo, 21s.

WILLIAM GRAY, M.D., C.M. (EDIN.).

INFLUENZA, with Special Reference to some Peculiar Sym-
ptoms. 8vo, 3s. 6d.

J. REYNOLDS GREEN, SC.D., F.R.S.

[See Cambridge Biological Series, page 5.]

A. HILL GRIFFITH, M.D.

Surgeon, Manchester Royal Eye Hospital.

THE DIAGNOSIS OF INTRA-OCULAR GROWTHS. With woodcuts, 8vo, 1s. 6d.

DR. JOSEF GRUBER.

Professor of Otology in the University of Vienna, Etc.

A TEXT-BOOK OF THE DISEASES OF THE EAR.

Translated from the Second German edition, and Edited, with additions, by EDWARD LAW, M.D., C.M. EDIN., M.R.C.S. ENG., Surgeon to the London Throat Hospital for Diseases of the Throat, Nose and Ear; and COLEMAN JEWELL, M.B. LOND., M.R.C.S. ENG., late Surgeon and Pathologist to the London Throat Hospital. Second English Edition, with 165 Illustrations, and 70 coloured figures on 2 lithographic plates, royal 8vo, 28s.

F. DE HAVILLAND HALL, M.D., F.R.C.P. LOND.

Physician to the Westminster Hospital; President of the Laryngological Society of London,

AND

HERBERT TILLEY, M.D., B.S. LOND., F.R.C.S. ENG.

Surgeon to the Throat Hospital, Golden Square; Lecturer on Diseases of the Nose and Throat, London Post-Graduate College and Polyclinic.

DISEASES OF THE NOSE AND THROAT. Second edition, thoroughly revised, with 2 coloured Plates and 80 Illustrations, crown 8vo, 10s. 6d.

[LEWIS'S PRACTICAL SERIES.]

H. J. HANSEN.

AND

W. SORENSEN.

Fellow Royal Danish Academy, Copenhagen.

ON TWO ORDERS OF ARACHNIDA OPILIONES. Especially the Sub-order, Cyphophthalmi and Ricinulei, namely the Cryptostemmatoidae. With 9 plates, 4to, 15s. *nett.* [Just published.]

A. HARKER, M.A., F.G.S.

[See Cambridge Geological Series, page 5.]

VINCENT DORMER HARRIS, M.D. LOND., F.R.C.P.

Physician to the City of London Hospital for Diseases of the Chest, Victoria Park; Examining Physician to the Royal National Hospital for Consumption and Diseases of the Chest, Ventnor, &c.,

AND

EDWIN CLIFFORD BEALE, M.A., M.B. CANTAB., F.R.C.P.

Physician to the City of London Hospital for Diseases of the Chest, Victoria Park, and to the Great Northern Central Hospital, &c.

THE TREATMENT OF PULMONARY CONSUMPTION.

A Practical Manual. Crown 8vo, 10s. 6d.

[LEWIS'S PRACTICAL SERIES.]

Drs. HARVEY and DAVIDSON'S

SYLLABUS OF MATERIA MEDICA. Revised in accordance with the "British Pharmacopœia," 1898, by WILLIAM MARTINDALE, F.L.S., F.C.S., Member of Council of Pharmaceutical Society, and late Examiner; Joint Author of "The Extra Pharmacopœia." Tenth Edition, foolscap 16mo, 1s. *nett.*

W. S. HEDLEY, M.D.

Medical Officer in charge of the Electro-Therapeutic Department of the London Hospital.

I.

THE HYDRO-ELECTRIC METHODS IN MEDICINE.

Second Edition, with Illustrations, demy 8vo, 4s. 6d.

II.

CURRENT FROM THE MAIN: The Medical Employment of Electric Lighting Currents. With Illustrations, demy 8vo, 2s. 6d.

III.

PRACTICAL MUSCLE-TESTING; AND, THE TREATMENT of Muscular Atrophies. With Illustrations, demy 8vo, 3s. 6d.

H. HELBING, F.C.S.

MODERN MATERIA MEDICA: For Medical Men, Pharmacists, and Students. Fourth Edition, 8vo, 8s. *nett*.

HERBERT T. HERRING, M.B., B.S. (DURH.), M.R.C.S.

THE STERILISATION OF URETHRAL INSTRUMENTS, and their Use in some Urinary Complaints. With Illustrations, demy 8vo, 5s.

[*Now ready.*]

C. HIGGENS, F.R.C.S.

Ophthalmic Surgeon to Guy's Hospital; Lecturer on Ophthalmology at Guy's Hospital Medical School.

A MANUAL OF OPHTHALMIC PRACTICE.

Second Edition, revised and edited by A. W. ORMOND, F.R.C.S.E., Assistant Surgeon, Royal Eye Hospital, Southwark, &c. With 66 Illustrations, crown 8vo, 7s. 6d. [LEWIS'S PRACTICAL SERIES.]

BERKELEY HILL, M.B. LOND., F.R.C.S.

Professor of Clinical Surgery in University College; Surgeon to University College Hospital and to the Lock Hospital;

AND

ARTHUR COOPER, L.R.C.P., M.R.C.S.

Surgeon to the Westminster General Dispensary.

SYPHILIS AND LOCAL CONTAGIOUS DISORDERS.

Second Edition, entirely re-written, royal 8vo, 18s.

JAMES HINSELWOOD, M.A., M.D., F.F.P.S. GLAS.

Surgeon to the Glasgow Eye Infirmary.

LETTER-, WORD- AND MIND-BLINDNESS.

Crown 8vo, 3s.

FROM HOSPITAL WARD TO CONSULTING ROOM, with Notes by the Way; a Medical Autobiography. By a Graduate of the London University. Post 8vo, 3s. 6d.

GEORGE LINDSAY JOHNSON, M.A., M.B., B.C. CANTAB.

Clinical Assistant, late House Surgeon and Chloroformist, Royal Westminster Ophthalmic Hospital, &c

A NEW METHOD OF TREATING CHRONIC GLAUCOMA, based on Recent Researches into its Pathology. With Illustrations and coloured frontispiece, demy 8vo, 3s. 6d.

H. LEWIS JONES, M.A., M.D., F.R.C.P.

Medical Officer in charge of the Electrical Department in St. Bartholomew's Hospital; President of the British Electro-Therapeutic Society, &c.

MEDICAL ELECTRICITY. A Practical Handbook for Students and Practitioners. Fourth Edition, thoroughly revised and enlarged, with 12 plates and 168 Illustrations, demy 8vo, 12s. 6d. *nett.*

[LEWIS'S PRACTICAL SERIES.]

[Just published.]

L. VERNON JONES, M.D.

GONORRHOËAL ARTHRITIS, its Pathology, Symptoms, and Treatment. With Illustrations, crown 8vo, 2s. 6d.

T. N. KELYNACK, M.D.

Pathologist to the Manchester Royal Infirmary; Demonstrator and Lecturer on Pathology in the Owens College.

A CONTRIBUTION TO THE PATHOLOGY OF THE VER-
miform Appendix. With Illustrations, large 8vo, 10s. 6d.

HENRY R. KENWOOD, M.B., D.P.H., F.C.S.

Professor of Hygiene and Public Health at University College, London, and Medical Officer of Health and Public Analyst for Stoke Newington and Finchley.

PUBLIC HEALTH LABORATORY WORK. The part on Public Health Bacteriological Work contributed by W. G. SAVAGE, M.D., B.Sc., D.P.H., Medical Officer of Health, Colchester, formerly Assistant to the Professor of Pathology, with Charge of the Bacteriological Department, University College, London, &c. Third Edition, with Illustrations, cr. 8vo, 10s. 6d.

[LEWIS'S PRACTICAL SERIES.] [Just published.]

NORMAN KERR, M.D., F.L.S.

President of the Society for the Study of Inebriety; Consulting Physician, Dalrymple Home for Inebriates, etc.

INEBRIETY OR NARCOMANIA: its Etiology, Pathology, Treatment, and Jurisprudence. Third Edition, 8vo, 7s. 6d. *nett.*

S. A. KNOPF, M.D.

Physician to the Lung Department of the New York Throat and Nose Hospital; formerly Assistant Physician to Professor Dettweiler, Falkenstein Sanatorium, Germany, &c.

PULMONARY TUBERCULOSIS: its Modern Prophylaxis and the Treatment in Special Institutions and at Home. With 76 Illustrations, royal 8vo, 14s. *nett.*

DR. PHILALETHES KUHN.

Staff Surgeon to the Imperial Troops of the South West African Protectorate.

INOCULATION AGAINST MALARIA.

Translated by H. A. NESBIT, M.A., with a Table of Curves. 8vo, 2s. *nett.*

F. CHARLES LARKIN, F.R.C.S. ENG.

Surgeon to the Stanley Hospital; late Assistant Lecturer in Physiology in University College, Liverpool,

AND

RANDLE LEIGH, M.B., B.SC. LOND.

Senior Demonstrator of Physiology in University College, Liverpool.

OUTLINES OF PRACTICAL PHYSIOLOGICAL CHEMISTRY.

Second Edition, with Illustrations, crown 8vo, paper 2s. 6d. *nett*, or cloth 3s. *nett*.

O. H. LATTER, M.A.

[See Cambridge Biological Series, page 5.]

J. WICKHAM LEGG, F.R.C.P.

Formerly Assistant Physician to Saint Bartholomew's Hospital, and Lecturer on Pathological Anatomy in the Medical School.

I.

ON THE BILE, JAUNDICE, AND BILIOUS DISEASES.

With Illustrations in chromo-lithography, roy. 8vo, 25s.

II.

A GUIDE TO THE EXAMINATION OF THE URINE.

Seventh Edition, edited and revised, by H. LEWIS JONES, M.D., M.A., F.R.C.P., Medical Officer in charge of the Electrical Department in St. Bartholomew's Hospital. With Illustrations, fcap. 8vo, 3s. 6d.

ARTHUR H. N. LEWERS, M.D. LOND., F.R.C.P. LOND.

Senior Obstetric Physician to the London Hospital, and Lecturer on Midwifery, London Hospital Medical School; Examiner in Obstetric Medicine to the University of London; Examiner in Midwifery and Diseases of Women at the Conjoint Board of the Royal College of Physicians of London, and of the Royal College of Surgeons of England, &c.

I.

CANCER OF THE UTERUS: A Clinical Monograph on its Diagnosis and Treatment. With the After Results in Seventy-Three Cases Treated by Radical Operation. With 3 coloured Plates and 51 original Illustrations, 8vo, 10s. 6d. *nett*.

II.

A PRACTICAL TEXTBOOK OF THE DISEASES OF WOMEN.

Sixth Edition, with 166 Illustrations and 4 coloured Plates, crown 8vo, 10s. 6d.

[Just published.]

[LEWIS'S PRACTICAL SERIES].

PROF. W. J. LEWIS, M.A.

[See Cambridge Geological Series, page 5.]

LEWIS'S POCKET MEDICAL VOCABULARY. Second Edition, thoroughly revised, 32mo, roan, 3s. 6d.

ROBERT LINDSAY, A.M., M.B., F.R.C.S.E.

Retired Surgeon, Army Medical Department.

AN ESSAY ON MALARIA AND ITS CONSEQUENCES.

Crown 8vo, 4s.

A. W. MACFARLANE, M.D., F.R.C.P. EDIN.

Examiner in Medical Jurisprudence in the University of Glasgow; Honorary Consulting Physician (late Physician) Kilmarnock Infirmary.

INSOMNIA AND ITS THERAPEUTICS. Medium 8vo, 12s. 6d.

WILLIAM A. M'KEOWN, M.D., M.CH.

Surgeon to the Ulster Eye, Ear, and Throat Hospital, Belfast; Member of the Senate of the Royal University of Ireland; Lecturer on Ophthalmology and Otology, Queen's College, Belfast.

A TREATISE ON "UNRIPE" CATARACT, and its Successful Treatment by Operation, with Tables comprising 151 Cases. With Illustrations, royal 8vo, 12s. 6d. *nett.*

DONALD J. MACKINTOSH, M.B., M.V.O.

Medical Superintendent, Western Infirmary, Glasgow.

SKIAGRAPHIC ATLAS OF FRACTURES AND DISLOCATIONS, with Notes on Treatment, for the use of Students. With 80 plates, demy 4to, 12s. 6d. *nett.*

J. M. H. MACLEOD, M.A., M.D., M.R.C.P.

Assistant Physician for Diseases of the Skin, Charing Cross Hospital; Physician for Diseases of the Skin, Victoria Hospital for Children; Lecturer on Skin Diseases, London School of Tropical Medicine.

PRACTICAL HANDBOOK OF THE PATHOLOGY OF THE SKIN. An Introduction to the Histology, Pathology, and Bacteriology of the Skin, with Special Reference to Technique. With 40 Plates, 8 being in Colours, from original Drawings, demy 8vo, 15s. *nett.*

JOHN MACPHERSON, M.D.

Inspector-General of Hospitals H.M. Bengal Army (Retired).

ANNALS OF CHOLERA FROM THE EARLIEST PERIODS to the Year 1817. With a map. Demy 8vo, 7s. 6d.

A. COWLEY MALLEY, B.A., M.B., B.CH., T.C.D.

PHOTO-MICROGRAPHY; including a description of the Wet Collodion and Gelatino-Bromide Processes, together with the best methods of Mounting and Preparing Microscopic Objects for Photo-Micrography. Second Edition, with Photographs and Illustrations, crown 8vo, 7s. 6d.

SIR PATRICK MANSON, K.C.M.G., M.D., LL.D.

THE FILARIA SANGUINIS HOMINIS; AND CERTAIN New Forms of Parasitic Disease in India, China, and Warm Countries. Illustrated with Plates and Charts, 8vo, 10s. 6d.

J. E. MARR, M.A.

[See Cambridge Geological Series, page 5.]

JEFFERY A. MARSTON, M.D., C.B., F.R.C.S., M.R.C.P. LOND.
Surgeon General Medical Staff (Retired).

NOTES ON TYPHOID FEVER: Tropical Life and its Sequelæ.
 Crown 8vo, 3s. 6d.

WILLIAM MARTINDALE, F.L.S., F.C.S.
Late President and Examiner of the Pharmaceutical Society,
 AND
 W. WYNN WESTCOTT, M.B. LOND., D.P.H.
H.M.'s Coroner for North-East London.

THE EXTRA PHARMACOPŒIA. Eleventh Edition, revised by W.
 HARRISON MARTINDALE, PH.D., F.C.S., and W. WYNN WESTCOTT, M.B., &c.,
 limp roan, med. 24mo, 9s. 6d. *nett.*

WILLIAM MARTINDALE, F.C.S.
Late Examiner of the Pharmaceutical Society, &c.

I.
 COCA, AND COCAINE: their History, Medical and Economic
 Uses, and Medicinal Preparations. Fourth Edition, fcap. 8vo, 2s.

II.
 ANALYSES OF TWELVE THOUSAND PRESCRIPTIONS:
 being Statistics of the Frequency of Use therein of all Official and Unofficial
 Preparations. Fcap. 4to, 2s. 6d. *nett.*

MATERIA MEDICA LABELS.

Adapted for Public and Private Collections. Compiled from the British Pharma-
 copœia of 1898 and other sources. The Labels are arranged in Two Divi-
 sions:—

Division I.—Comprises Chemical Materia Medica, including Alcohols, Alka-
 loids, Sugars, and Neutral Bodies.

Division II.—Comprises, with few exceptions, Substances of Organized Struc-
 ture, obtained from the Vegetable and Animal Kingdoms.

On gummed paper, 12s. 6d. *nett.*

. Specimens of the Labels, of which there are about 500, will be sent on application.

S. E. MAUNSELL, L.R.C.S.I.
Surgeon-Major, Medical Staff.

NOTES OF MEDICAL EXPERIENCES IN INDIA PRINCIPALLY
 with Reference to Diseases of the Eye. With Map, post 8vo, 3s. 6d.

C. W. MANSELL MOULLIN, M.D. OXON., F.R.C.S.
*Surgeon to and Lecturer on Surgery at the London Hospital; Examiner in Surgery at the University of
 Oxford, &c.*

I.
 INFLAMMATION OF THE BLADDER AND URINARY
 Fever. 8vo, 5s.

II.
 ENLARGEMENT OF THE PROSTATE: its Treatment and
 Radical Cure. Third edition, with plates, 8vo, 6s. [*Just published.*]

III.
 SPRAINS; THEIR CONSEQUENCES AND TREATMENT.
 Second Edition, crown 8vo, 4s. 6d.

GEORGE R. MURRAY, M.A., M.D. CAMB., F.R.C.P.

Heath Professor of Comparative Pathology in the University of Durham; Physician to the Royal Infirmary, Newcastle.

DISEASES OF THE THYROID GLAND. Part I., MYXŒDEMA AND
CRETINISM. With 25 Illustrations, demy 8vo, 7s. 6d.

WILLIAM MURRAY, M.D., F.R.C.P. LOND.

I.

ROUGH NOTES ON REMEDIES. Fourth Edition, with additional
Chapter on Rothbury as a Health Resort. Crown 8vo, 4s. nett.

II.

**ILLUSTRATIONS OF THE INDUCTIVE METHOD IN MEDI-
cine.** Crown 8vo, 3s. 6d.

WILLIAM MURRELL, M.D., F.R.C.P.

*Physician to, and Lecturer on Clinical Medicine and Joint Lecturer on the Principles and Practice of
Medicine at, the Westminster Hospital; Examiner in Materia Medica in the
University of Glasgow.*

WHAT TO DO IN CASES OF POISONING. Ninth Edition,
royal 32mo, 3s. 6d.

GEORGE OLIVER, M.D. LOND., F.R.C.P. LOND.

I.

BLOOD-PRESSURE AND TISSUE-LYMPH CIRCULATION:
The Oliver-Sharpey Lectures, 1904, with addenda. [Preparing.]

II.

**A CONTRIBUTION TO THE STUDY OF THE BLOOD AND
BLOOD-PRESSURE;** founded on Portions of the Croonian Lectures delivered
before the Royal College of Physicians, London, 1896, with Considerable Exten-
sions. With Illustrations, demy 8vo, 7s. 6d.

III.

**PULSE-GAUGING: A Clinical Study of Radial Measurement
and Pulse Pressure.** With Illustrations, fcap. 8vo, 3s. 6d.

IV.

HARROGATE AND ITS WATERS: Notes on the Climate of
Harrogate, and on the Chemistry of the Mineral Spring. With Map of the
Wells, crown 8vo, 2s. 6d.

V.

ON BEDSIDE URINE TESTING: a Clinical Guide to the
Observation of Urine in the course of Work. Fourth Edition, fcap. 8vo, 3s. 6d.

DR. A. ONODI.

Lecturer on Rhino-Laryngology in the University of Budapest.

**THE ANATOMY OF THE NASAL CAVITY, AND ITS ACCES-
sory Sinuses.** An Atlas for Practitioners and Students. Translated by ST CLAIR
THOMSON, M.D. LOND., F.R.C.S. ENG., M.R.C.P. LOND. With plates, small 4to,
6s. nett.

SAML. OSBORN, F.R.C.S.

Knight of Grace of the Order of St. John of Jerusalem; Lecturer and Examiner to the St. John Ambulance Association; Surgeon to the Hospital for Women, Soho Square, &c.

I.

AMBULANCE LECTURES: FIRST AID TO THE INJURED.

Fourth Edition, with Illustrations, fcap. 8vo, 2s.

II.

AMBULANCE LECTURES: HOME NURSING AND HY-

giene. Third Edition, with Illustrations, fcap. 8vo, 2s.

WILLIAM OSLER, M.D., F.R.C.P. LOND., F.R.S.

Regius Professor of Medicine, University of Oxford; President of the Association of American Physicians; Professor of Medicine, Johns Hopkins University, and Physician-in-Chief Johns Hopkins Hospital, Baltimore.

I.

ÆQUANIMITAS. With other Addresses to Medical Students, Nurses, and Practitioners of Medicine, p. 8vo. [Just published.]

II.

ON CHOREA AND CHOREIFORM AFFECTIONS. Large 8vo, 5s.

III.

THE CEREBRAL PALSIES OF CHILDREN. A Clinical Study from the Infirmary for Nervous Diseases, Philadelphia. Demy 8vo, 5s.

WILLIAM OSLER, M.D., F.R.S., F.R.C.P. LOND.

AND

THOMAS McCRAE, M.B. TOR., L.R.C.P. LOND.

Of the Johns Hopkins Hospital, Baltimore.

CANCER OF THE STOMACH; a Clinical Study. With Illustrations, med. 8vo, 6s.

KURRE W. OSTROM.

Instructor in Massage and Swedish Movements in the Philadelphia Polyclinic and College for Graduates in Medicine.

MASSAGE AND THE ORIGINAL SWEDISH MOVEMENTS; their application to various diseases of the body. Fifth Edition, with 105 Illustrations, crown 8vo, 3s. 6d. *nett.*

CHARLES A. PARKER. F.R.C.S. EDIN.

Assistant Surgeon to the Hospital for Diseases of the Throat, Golden Square, London.

POST-NASAL GROWTHS. Demy 8vo, 4s. 6d.

ROBERT W. PARKER.

Senior Surgeon to the East London Hospital for Children; Surgeon to the German Hospital.

DIPHTHERIA: ITS NATURE AND TREATMENT, WITH
Special Reference to the Operation, After-Treatment, and Complications of
Tracheotomy. Third Edition, with Illustrations, 8vo, 6s.

LOUIS PARKES, M.D., D.P.H. LOND. UNIV.

*Fellow of the Sanitary Institute; Lecturer on Public Health at St. George's Hospital Medical School;
Medical Officer of Health for Chelsea,*

AND

HENRY R. KENWOOD, M.B., D.P.H., F.C.S.

*Fellow of the Sanitary Institute and Member of the Board of Examiners; Professor of Hygiene and
Public Health at University College, London, &c.*

HYGIENE AND PUBLIC HEALTH. Second edit., with 88 Illustrations,
crown 8vo, 12s. [LEWIS'S PRACTICAL SERIES.]

LOUIS PARKES, M.D., D.P.H. LOND. UNIV.

INFECTIOUS DISEASES, NOTIFICATION AND PREVEN-
tion. Fcap. 8vo, cloth, 2s. 6d; roan, 4s. 6d.

LESLIE PHILLIPS, M.D.

Surgeon to the Birmingham and Midland Skin and Lock Hospital.

MEDICATED BATHS IN THE TREATMENT OF SKIN DIS-
eases. Crown 8vo, 4s. 6d.

HENRY G. PIFFARD, A.M., M.D.

*Clinical Professor of Dermatology, University of the City of New York; Surgeon in Charge of the
New York Dispensary for Diseases of the Skin, &c.*

A PRACTICAL TREATISE ON DISEASES OF THE SKIN.
With 50 full page Original Plates and 33 Illustrations in the Text, 4to,
£2 12s. 6d. nett.

J. E. PLATT, M.S. LOND., F.R.C.S.

Honorary Surgeon to the Hulme Dispensary; Surgical Officer to the Cancer Hospital, Manchester, &c.

A CONTRIBUTION TO THE SURGERY OF FRACTURES
and Dislocations of the Upper Extremity, based upon an analysis of about 700
Consecutive Cases observed at the Manchester Royal Infirmary. With Illus-
trations, royal 8vo, 10s.

G. V. POORE, M.D., F.R.C.P.

*Professor of Medical Jurisprudence, University College; Assistant Physician to, and Physician in
charge of the Throat Department of, University College Hospital.*

LECTURES ON THE PHYSICAL EXAMINATION OF THE
Mouth and Throat. With an Appendix of Cases. 8vo, 3s. 6d.

SIR RICHARD DOUGLAS POWELL, Bart., M.D. LOND., F.R.C.P.

*Physician Extra-ordinary to His Majesty the King; Physician to the Middlesex Hospital;
Consulting Physician to the Brompton Hospital, &c.*

THE LUMLEIAN LECTURES ON THE PRINCIPLES WHICH
Govern Treatment in Diseases and Disorders of the Heart. With coloured
Diagrams, demy 8vo, 6s.

BY THE SAME AUTHOR

DISEASES OF THE LUNGS AND PLEURÆ, INCLUDING
Consumption. Fourth Edition, with coloured plates and wood engravings,
8vo, 18s.

TABLE OF PHYSICAL EXAMINATION OF THE LUNGS:
with Note on International Nomenclature of Physical Signs (reprinted from
SIR R. D. POWELL'S "Diseases of the Lungs"). On one sheet, 6d.

D'ARCY POWER, M.A., M.B. OXON., F.R.C.S. ENG.

*Assistant Surgeon at St. Bartholomew's Hospital; Surgeon to the Victoria Hospital for Children,
Chelsea; Examiner in the University of Durham; Member of the Conjoint Examining Board
of the Royal College of Physicians (Lond.) and of Surgeons (Eng.).*

THE SURGICAL DISEASES OF CHILDREN AND THEIR
Treatment by Modern Methods. With Illustrations, crown 8vo, 10s. 6d.
[LEWIS'S PRACTICAL SERIES.]

URBAN PRITCHARD, M.D. EDIN., F.R.C.S. ENG.

*Professor of Aural Surgery at King's College, London; Aural Surgeon to King's College Hospital;
Senior Surgeon to the Royal Ear Hospital.*

HANDBOOK OF DISEASES OF THE EAR FOR THE USE
of Students and Practitioners. Fourth Edition, with Illustrations, crown 8vo.
[LEWIS'S PRACTICAL SERIES.]
[In preparation.]

DR. THEODOR PUSCHMANN.

Public Professor in Ordinary at the University of Vienna.

A HISTORY OF MEDICAL EDUCATION FROM THE MOST
Remote to the Most Recent Times. Translated and edited by EVAN H. HARE,
M.A. OXON., F.R.C.S. ENG., L.S.A. Demy 8vo, 21s.

C. H. RALFE, M.A., M.D. CANTAB., F.R.C.P. LOND.

*Assistant Physician to the London Hospital; Examiner in Medicine to the University of
Durham, &c., &c.*

A PRACTICAL TREATISE ON DISEASES OF THE KID-
neys and Urinary Derangements. With Illustrations, crown 8vo, 10s. 6d.
[LEWIS'S PRACTICAL SERIES.]

LOUIS BATHE RAWLING, M.B., B.C. (CANT.), F.R.C.S. (ENG.).
Senior Demonstrator of Anatomy, St. Bartholomew's Hospital; Assistant Surgeon to the German Hospital, Dalston; Hunterian Professor, Royal College of Surgeons, England.

LANDMARKS AND SURFACE MARKINGS OF THE HUMAN
 Body. Demy 8vo, 24 plates (3 in colours) 5s. nett. [Just published.]

F. R. COWPER REED, M.A., F.G.S.

[See Cambridge Geological Series, page 5.]

H. A. REEVES, F.R.C.S. EDIN.

*Senior Assistant Surgeon and Teacher of Practical Surgery at the London Hospital;
 Surgeon to the Royal Orthopaedic Hospital.*

BODILY DEFORMITIES AND THEIR TREATMENT: A
 Handbook of Practical Orthopaedics. With Illustrations, crown 8vo, 8s. 6d.
 [LEWIS'S PRACTICAL SERIES.]

A. B. RENDLE, M.A. (CANTAB.), B.SC. (LOND.).

[See Cambridge Biological Series, page 5.]

SIDNEY H. REYNOLDS, M.A.

[See Cambridge Biological Series, page 5.]

SAMUEL RIDEAL, D.SC. LOND., F.I.C., F.C.S.
Fellow of University College, London.

I.
PRACTICAL ORGANIC CHEMISTRY; The Detection and
 Properties of some of the more important Organic Compounds. Second edition,
 12mo, 2s. 6d.

II.
PRACTICAL CHEMISTRY FOR MEDICAL STUDENTS, re-
 quired at the First Examination of the Conjoint Examining Board in England.
 Fcap 8vo, 2s.

J. JAMES RIDGE, M.D.

Medical Officer of Health, Enfield.

ALCOHOL AND PUBLIC HEALTH. Second Edition, crown 8vo, 2s.

E. A. RIDSDALE.

Associate of the Royal School of Mines.

COSMIC EVOLUTION; being Speculations on the Origin of our
 Environment. Fcap. 8vo, 3s.

SYDNEY RINGER, M.D., F.R.S.

Holme Professor of Clinical Medicine in University College; Physician to University College Hospital,
AND

HARRINGTON SAINSBURY, M.D., F.R.C.P.

*Physician to the Royal Free Hospital and the City of London Hospital for Diseases of the Chest,
Victoria Park.*

A HANDBOOK OF THERAPEUTICS. Thirteenth Edition, thoroughly revised, 8vo, 16s.

SYDNEY RINGER, M.D., F.R.S.

ON THE TEMPERATURE OF THE BODY AS A MEANS
of Diagnosis and Prognosis in Phthisis. Second Edition, small 8vo, 2s. 6d.

R. LAWTON ROBERTS, M.D. LOND., D.P.H. CAMB., M.R.C.S. ENG.

*Honorary Life Member of, and Lecturer and Examiner to, the St. John Ambulance Association;
J.P. for County of Denbigh.*

I.

ILLUSTRATED LECTURES ON AMBULANCE WORK.

Fifth Edition, copiously Illustrated, crown 8vo, 2s. 6d.

II.

ILLUSTRATED LECTURES ON NURSING AND HYGIENE.

Third Edition, with Illustrations, crown 8vo, 2s. 6d.

FREDERICK T. ROBERTS, M.D., B.SC., F.R.C.P.

*Professor of the Principles and Practice of Medicine in University College; Physician to University
College Hospital; Consulting Physician to Brompton Consumption Hospital, &c.*

THE THEORY AND PRACTICE OF MEDICINE. Tenth Edition,
with Illustrations, large 8vo. [In the press.]

H. D. ROLLESTON, M.D., F.R.C.P.

[See Cambridge Biological Series, page 5.]

ROBSON ROOSE, M.D., LL.D., F.C.S.

Fellow of the Royal College of Physicians in Edinburgh, &c.

LEPROSY AND ITS PREVENTION: as Illustrated by Nor-
wegian Experience. Crown 8vo, 3s. 6d.

WILLIAM ROSE, M.B., B.S. LOND., F.R.C.S.

Professor of Surgery in King's College, London, and Surgeon to King's College Hospital.

HARELIP AND CLEFT PALATE. With Illustrations, demy 8vo, 6s.

BERNARD ROTH, F.R.C.S.

Orthopædic Surgeon to the Royal Alexandra Children's Hospital, Brighton, &c.

THE TREATMENT OF LATERAL CURVATURE OF THE Spine: with Appendix giving an Analysis of 1000 Consecutive Cases treated by "Posture and Exercise" exclusively (without Mechanical Support). Second edition, with Photographic and other Illustrations, royal 8vo, 10s. 6d.

PROF. E. RUTHERFORD.

[See Cambridge Physical Series, page 5.]

JOHN SAVORY.

Member of the Society of Apothecaries, London.

A COMPENDIUM OF DOMESTIC MEDICINE AND COMPANION to the Medicine Chest: Intended as a source of easy reference for Clergymen, Master Mariners, and Travellers; and for Families resident at a distance from professional assistance. Tenth Edition, sm. 8vo, 5s.

DR. B. S. SCHULTZE.

Professor of Gynecology; Director of the Lying-in Hospital, and of the Gynecological Clinic at Jena.

THE PATHOLOGY AND TREATMENT OF DISPLACEMENTS of the Uterus. Translated by J. J. MACAN, M.A., M.R.C.S., and edited by A. V. MACAN, M.B., M.Ch., Master of the Rotunda Lying-in Hospital, Dublin. With 120 Illustrations, medium 8vo, 12s. 6d.

A. C. SEWARD, M.A., F.G.S.

[See Cambridge Biological Series, page 5.]

A. E. SHIPLEY, M.A.,

AND

E. W. MACBRIDE, M.A.

[See Cambridge Biological Series, page 5.]

G. E. SHUTTLEWORTH, B.A., M.D.

Medical Examiner of Defective Children, School Board for London; late Medical Superintendent, Royal Albert Asylum for Idiots and Imbeciles of the Northern Counties, Lancaster, &c.

MENTALLY-DEFICIENT CHILDREN: THEIR TREATMENT and Training. Second edition, with Illustrations, crown 8vo, 5s. *nett.*

DAVID SIME, M.D.

RABIES: Its Place Amongst Germ Diseases, and its Origin in the Animal Kingdom. Royal 8vo, 10s. 6d. *nett.*

FRANCIS W. SMITH, M.B., B.S.

THE SALINE WATERS OF LEAMINGTON. Second Edition, with Illustrations, crown 8vo, 1s. nett.

J. LEWIS SMITH, M.D.

Physician to the New York Foundling Asylum; Clinical Professor of Diseases of Children in Bellevue Hospital Medical College.

A TREATISE ON THE MEDICAL AND SURGICAL DISEASES OF INFANCY AND CHILDHOOD. Eighth Edition, with 273 Illustrations and four plates, large 8vo, 21s.

E. HUGH SNELL, M.D., B.SC. LOND.

Diplomate in Public Health of the University of Cambridge; Medical Officer of Health to the City of Coventry; late London County Council Medical Officer to the Blackwall Tunnel.

COMPRESSED AIR ILLNESS, OR SO-CALLED CAISSON DISEASE. With Illustrations, demy 8vo, 10s. 6d.

JOHN KENT SPENDER, M.D. LOND.

Physician to the Royal Mineral Water Hospital, Bath.

THE EARLY SYMPTOMS AND THE EARLY TREATMENT OF OSTEO-ARTHRITIS, commonly called Rheumatoid Arthritis, with special reference to the Bath Thermal Waters. Sm. 8vo, 2s. 6d.

LOUIS STARR, M.D.

Physician to the Children's Hospital, Philadelphia, &c.

HYGIENE OF THE NURSERY. Including the General Regimen and Feeding of Infants and Children; Massage, and the Domestic Management of the Ordinary Emergencies of Early Life. Sixth Edition, with Illustrations, crown 8vo, 3s. 6d.

JOHN LINDSAY STEVEN, M.D.

Assistant Physician and Pathologist, Glasgow Royal Infirmary; Physician for Out-patients, Royal Hospital for Sick Children, Glasgow, &c.

THE PATHOLOGY OF MEDIASTINAL TUMOURS. With special reference to Diagnosis. With Plates, 8vo, 4s. 6d.

JUKES DE STYRAP, M.R.C.P.I., ETC.

Physician-Extraordinary, late Physician in Ordinary, to the Salop Infirmary; Consulting Physician to the South Salop and Montgomeryshire Infirmaries, etc.

I.

THE YOUNG PRACTITIONER: WITH PRACTICAL HINTS
and Instructive Suggestions, as Subsidiary Aids, for his Guidance on Entering
into Private Practice. Demy 8vo, 7s. 6d. *nett*.

II.

A CODE OF MEDICAL ETHICS: WITH GENERAL AND
Special Rules for the Guidance of the Faculty and the Public in the Complex
Relations of Professional Life. Fourth Edition, demy 8vo, 3s. 6d. *nett*.

III.

MEDICO-CHIRURGICAL TARIFFS. Fifth Edition, revised and en-
larged, fcap. 4to, 2s. *nett*.

IV.

THE YOUNG PRACTITIONER: HIS CODE AND TARIFF.
Being the above three works in one volume. Demy 8vo, 10s. 6d. *nett*.

C. W. SUCKLING, M.D. LOND., M.R.C.P.

*Professor of Materia Medica and Therapeutics at the Queen's College, Physician to the Queen's
Hospital, Birmingham, etc.*

I.

ON THE DIAGNOSIS OF DISEASES OF THE BRAIN
Spinal Cord, and Nerves. With Illustrations, crown 8vo, 8s. 6d.

II.

ON THE TREATMENT OF DISEASES OF THE NERVOUS
System. Crown 8vo, 7s. 6d.

J. BLAND-SUTTON, F.R.C.S.

*Assistant Surgeon to the Middlesex Hospital; Examiner in Anatomy for the Fellowship to the Royal
College of Surgeons, England.*

LIGAMENTS: THEIR NATURE AND MORPHOLOGY.

Third Edition, with numerous Illustrations, post 8vo, 4s. 6d. [Now ready.]

HENRY R. SWANZY, A.M., M.B., F.R.C.S.I.

*Surgeon to the Royal Victoria Eye and Ear Hospital, and Ophthalmic Surgeon to the Adelaide
Hospital, Dublin; Ex-President of the Ophthalmological Society of the United Kingdom.*

A HANDBOOK OF THE DISEASES OF THE EYE AND
their Treatment. Eighth Edition, illustrated with 168 wood-engravings, colour
tests, &c., small 8vo, 12s. 6d. [Now ready.]

ALBERT TAYLOR.

Member Sanitary Institute; Demonstrator to the Students of the Sanitary Institute; Sanitary Inspector, City of Westminster; Late Chief Sanitary Inspector to the Vestry of St. George, Hanover Square, &c.

THE SANITARY INSPECTOR'S HANDBOOK. Third Edition, with Illustrations, crown 8vo, 6s.

H. COUPLAND TAYLOR, M.D.

Fellow of the Royal Meteorological Society.

WANDERINGS IN SEARCH OF HEALTH, OR MEDICAL and Meteorological Notes on Various Foreign Health Resorts. With Illustrations, crown 8vo, 6s.

JOHN W. TAYLOR, F.R.C.S. ENG.

Professor of Gynæcology in the University of Birmingham; Senior In-patient Surgeon to the Birmingham and Midland Hospital for Women; Consulting Surgeon to the Wolverhampton Hospital for Women; Consulting Gynæcological Surgeon to the Birmingham Skin and Lock Hospital, &c.

EXTRA-UTERINE PREGNANCY. A Clinical and Operative Study. With Illustrations, demy 8vo, 7s. 6d.

J. J. THOMSON, D.SC., LL.D., F.R.S.

[See Cambridge Physical Series, page 5.]

J. C. THOROWGOOD, M.D.

Assistant Physician to the City of London Hospital for Diseases of the Chest.

THE CLIMATIC TREATMENT OF CONSUMPTION AND Chronic Lung Diseases. Third Edition, post 8vo, 3s. 6d.

HERBERT TILLEY, M.D., B.S. LOND., F.R.C.S. ENG.

Surgeon to the Throat Hospital, Golden Square; Lecturer on Diseases of the Nose and Throat, London Post-Graduate College and Polyclinic.

PURULENT NASAL DISCHARGES, their Diagnosis and Treatment. Second Edition, with Illustrations, crown 8vo, 4s. *nett.*

M. TUCHMANN, M.R.C.S. ENG., M.D. WURZBURG.

Formerly Surgeon to Out-Patients, German Hospital, Dalston.

THE EXPLORATION OF THE URETHRA AND BLADDER. With 26 Illustrations, demy 8vo, 5s.

A. DUNBAR WALKER, M.D., C.M.

THE PARENT'S MEDICAL NOTE BOOK. Oblong post 8vo, cloth, 1s. 6d.

E. W. AINLEY WALKER, M.A., D.M. (OXON.).

Fellow and Praelector of University College, Oxford; Late Goraon Lecturer in Experimental Pathology at Guy's Hospital; formerly Radcliffe Travelling Fellow in the University of Oxford, &c.

THE GENERAL PATHOLOGY OF INFLAMMATION, INfection, and Fever, being the Gordon Lectures for 1902. Crown 8vo, 4s. 6d. *nett.*
[Just published.]

A. J. WALL, M.D. LOND.

Fellow of the Royal College of Surgeons of England; of the Medical Staff of H. M. Indian Army (Retired List).

ASIATIC CHOLERA: its History, Pathology, and Modern Treatment. Demy 8vo, 6s.

H. MARSHALL WARD, SC.D., F.R.S.

[See Cambridge Biological Series, page 5.]

W. SPENCER WATSON, B.M. LOND., F.R.C.S. ENG.

Surgeon to the Throat Department of the Great Northern Hospital; Senior Surgeon to the Royal South London Ophthalmic Hospital.

I.

DISEASES OF THE NOSE AND ITS ACCESSORY CAVITIES.
Second Edition, with Illustrations, demy 8vo, 12s. 6d.

II.

THE ANATOMY AND DISEASES OF THE LACHRYMAL Passages. With Illustrations, demy 8vo, 2s. 6d.

III.

EYEBALL-TENSION: Its Effects on the Sight and its Treatment. With woodcuts, post 8vo, 2s. 6d.

W. WYNN WESTCOTT, M.B.

Deputy Coroner for Central Middlesex.

SUICIDE; its History, Literature, Jurisprudence, and Prevention. Crown 8vo, 6s.

FRANK J. WETHERED, M.D.

Medical Registrar to the Middlesex Hospital, and Demonstrator of Practical Medicine in the Middlesex Hospital Medical School; late Assistant Physician to the City of London Chest Hospital, Victoria Park.

MEDICAL MICROSCOPY. A Guide to the Use of the Microscope in Medical Practice. With Illustrations, crown 8vo, 9s.

[LEWIS'S PRACTICAL SERIES.]

W. C. D. WHETHAM, M.A.

[See Cambridge Physical Series, page 5.]

L. R. WILBERFORCE, M.A.

AND

T. C. FITZPATRICK, M.A.

Demonstrators at the Cavendish Laboratory, Cambridge.

A LABORATORY NOTE-BOOK OF ELEMENTARY PRACTICAL PHYSICS. I. MECHANICS AND HYDROSTATICS. II. HEAT AND OPTICS. III. MAGNETISM AND ELECTRICITY. Quarto, paper covers, 1s. each.

SIR JOHN WILLIAMS, BART., M.D., F.R.C.P.
*Consulting Physician to University College Hospital; Physician Accoucheur to H.R.H.
Princess Beatrice, &c.*

CANCER OF THE UTERUS: Being the Harveian Lectures for 1886. Illustrated with Lithographic Plates, royal 8vo, 10s. 6d.

W. WILLIAMS, M.A., M.D., D.P.H. (OXON.).
*Medical Officer of Health to the Glamorgan County Council; Lecturer in Public Health to the
University College of South Wales and Monmouthshire, Cardiff; Examiner in
State Medicine to the University of London, &c.*

DEATHS IN CHILDBED: A Preventable Mortality, being the Milroy Lectures for 1904. Demy 8vo, 2s. 6d. *nett.* [Just published.]

R. T. WILLIAMSON, M.D. LOND., F.R.C.P.
Medical Registrar, Royal Infirmary, and Assistant in Medicine, Owens College, Manchester.

ON THE RELATION OF DISEASES OF THE SPINAL CORD
to the Distribution and Lesions of the Spinal Blood Vessels. Royal 8vo, 2s.

J. C. WILLIS, M.A.

[See Cambridge Biological Series, page 5.]

E. T. WILSON, M.B. OXON., F.R.C.P. LOND.
*Physician to the Cheltenham General Hospital; Associate Metropolitan Association of Medical
Officers of Health.*

DISINFECTANTS AND ANTISEPTICS: HOW TO USE THEM.
40th Thousand. In Packets of one doz. price 1s., by post 1s. 1d.
[Thoroughly revised.]

BERTRAM C. A. WINDLE, F.R.S., SC.D., M.D., M.A. DUBL.
*Professor of Anatomy in the University of Birmingham; sometime Examiner in Anatomy in the
Universities of Cambridge, Aberdeen, and Durham.*

A HANDBOOK OF SURFACE ANATOMY AND LANDMARKS.
Third Edition, Illustrated with plain and coloured figures, post 8vo, 4s. *nett.*
[Now ready.]

EDWARD WOAKES, M.D. LOND.
*Senior Aural Surgeon, London Hospital; Lecturer on Diseases of the Ear, London Hospital Medical
College.*

ON DEAFNESS, GIDDINESS AND NOISES IN THE HEAD.
Fourth Edition, Part i., with Illustrations, 8vo, 10s. 6d.

HENRY WOODS, B.A., F.G.S.

[See Cambridge Biological Series, page 5.]

A. S. WOODWARD, M.A.

[See Cambridge Biological Series, page 5.]

OSWALD ZIEMSEN, M.D.

Knight of the Iron Cross, and of the Prussian Order of the Crown.

THE TREATMENT OF CONSTITUTIONAL SYPHILIS.

Post 8vo, 3s. 6d.

Lewis's Diet Charts.

Price 5s. per packet of 100 charts (assorted) post free.

A suggestive set of Diet Tables for the use of Physicians, for handing to patients after consultation, modified to suit individual requirements, for Albuminuria, Anæmia and Debility, Constipation, Diabetes, Diarrhœa, Dyspepsia, Eczema, Fevers, Gall Stones, Gout and Gravel, Heart Disease (chronic), Nervous Diseases, Obesity, Phthisis, Rheumatism (chronic), and Blank Chart for other diseases.

A special leaflet on the Diet and Management of Infants is sold separately, price 7s. 6d. per 100, or 1s. per dozen, post free.

Lewis's Handy Temperature Chart.

25s. per 1000, 14s. per 500, 3s. 6d. per 100, 2s. per 50, 1s. per 20, carriage free.

Arranged for three weeks, and specially ruled on back for recording observations on Urine.

Lewis's Four-Hour Temperature Chart.

25s. per 1000, 14s. per 500, 3s. 6d. per 100, 2s. per 50, 1s. per 20, carriage free.

This form has been drawn up to meet the requirements of a chart on which the temperature and other observations can be recorded at intervals of four hours. They will be found most convenient in hospital and private practice. Each chart will last a week.

Clinical Chart for Temperature Observations, etc.

Arranged by W. RIGDEN, M.R.C.S. 50s. per 1000, 28s. per 500, 15s. per 250, 7s. per 100, or 1s. per dozen, carriage free.

Each Chart is arranged for four weeks, and is ruled at the back for making notes of Cases they are convenient in size, and are suitable both for hospital and private practice.

Lewis's Nursing Chart.

25s. per 1000, 14s. per 500, 3s. 6d. per 100, 2s. per 50, or 1s. per 20, post free.

This Chart affords a ready method of recording the progress of the case from day to day. Printed on both sides.

* * Boards to hold either of the above Charts, price 1s.

Lewis's Clinical Chart, specially designed for use with the Visiting

List. This Temperature Chart is arranged for four weeks and measures 6 × 3 inches. 20s. per 1000, 11s. 6d. per 500, 2s. 6d. per 100, 6d. per dozen, post free.

Chart for Recording the Examination of Urine.

40s. per 1000; 25s. per 500; 15s. per 250; 7s. 6d. per 100; 1s. per 10.

This Chart is designed for the use of Medical Men, Analysts, and others making examinations of the Urine of patients and affords a very ready and convenient method of recording the results of the examination.

LEWIS'S PRACTICAL SERIES.

Under this title a Series of Monographs is published, embracing the various branches of Medicine and Surgery. The volumes are written by well known Hospital Physicians and Surgeons, recognised as authorities in the subjects of which they treat. The works are of a THOROUGHLY PRACTICAL nature, calculated to meet the requirements of the practitioner and student and to present the most recent information in a compact form.

A HANDBOOK OF BACTERIOLOGICAL DIAGNOSIS FOR PRACTITIONERS (including Instructions for the Clinical Examination of the Blood). By W. D'ESTE EMERY, M.D., B.Sc. Lond., Assistant Bacteriologist to the Laboratories of the Royal Colleges of Physicians and Surgeons, London, &c. With 2 Coloured Plates and 33 Illustrations, crown 8vo, 5s. 6d.

DISEASES OF THE NERVOUS SYSTEM: A Handbook for Students and Practitioners. By CHARLES E. BEEVOR, M.D., LOND., F.R.C.P., Physician to the National Hospital for the Paralyse and Epileptic, the Great Northern Central Hospital, and the National Orthopædic Hospital. With Illustrations, crown 8vo, 10s. 6d.

THE TREATMENT OF PULMONARY CONSUMPTION. By VINCENT D. HARRIS, M.D. LOND., F.R.C.P., Physician to the City of London Hospital for Diseases of the Chest, Victoria Park, &c., and E. CLIFFORD BEALE, M.A., M.B. CANTAB., F.R.C.P., Physician to the City of London Hospital for Diseases of the Chest, Victoria Park, &c. Crown 8vo, 10s. 6d.

THE SURGICAL DISEASES OF CHILDREN AND THEIR TREATMENT BY Modern Methods. By D'ARCY POWER, M.A., M.B. Oxon., F.R.C.S. Eng., Assistant Surgeon at St. Bartholomew's Hospital, &c. With Illustrations, crown 8vo, 10s. 6d.

DISEASES OF THE NOSE AND THROAT. By F. de HAVILLAND HALL, M.D., F.R.C.P. Lond., Physician to the Westminster Hospital, and HERBERT TILLEY, M.D., B.S. Lond., F.R.C.S. Eng., Surgeon to the Throat Hospital, Golden Square, &c. Second edition, with two coloured plates and 80 illustrations, crown 8vo, 10s. 6d.

PUBLIC HEALTH LABORATORY WORK. By HENRY R. KENWOOD, M.B., D.P.H., F.C.S., Professor of Hygiene and Public Health, University College, &c. Third Edition, Part VII. contributed by W. G. SAVAGE, M.D., &c. With Illustrations, cr. 8vo, 10s. 6d.

[Just published.]

MEDICAL MICROSCOPY: A Guide to the use of the Microscope in Medical Practice. By FRANK J. WETHERED, M.D., M.R.C.P., Demonstrator of Practical Medicine in the Middlesex Hospital Medical School, &c. With Illustrations, crown 8vo, 9s.

MEDICAL ELECTRICITY: A Practical Handbook for Students and Practitioners. By H. LEWIS JONES, M.A., M.D., F.R.C.P. Fourth edition, with 168 illustrations, demy 8vo, 12s. 6d. nett.

[Just published.]

HYGIENE AND PUBLIC HEALTH. By LOUIS PARKES, M.D., D.P.H. LOND. UNIV., Fellow of the Sanitary Institute, and HENRY R. KENWOOD, M.B., D.P.H., F.C.S., Professor of Hygiene and Public Health, University College. Second Edition, with 88 illustrations, cr. 8vo, 12s.

MANUAL OF OPHTHALMIC PRACTICE. By C. HIGGENS, F.R.C.S., Ophthalmic Surgeon to Guy's Hospital; Lecturer on Ophthalmology at Guy's Hospital Medical School. Second Edition, revised and edited by A. W. ORMOND, F.R.C.S.E., &c. With Illustrations, cr. 8vo, 7s. 6d.

[Just published.]

A PRACTICAL TEXTBOOK OF THE DISEASES OF WOMEN. By ARTHUR H. N. LEWERS, M.D. Lond., F.R.C.P. Lond., Senior Obstetric Physician to the London Hospital, etc. Sixth Edition, with Illustrations, crown 8vo, 10s. 6d.

[Just published.]

ANÆSTHETICS: Their Uses and Administration. By DUDLEY W. BUXTON, M.D., B.S., M.R.C.P., Administrator of Anæsthetics and Lecturer in University College Hospital, &c. Fourth Edition, with Illustrations, crown 8vo.

[In the press.]

ON FEVERS: Their History, Etiology, Diagnosis, Prognosis, and Treatment. By ALEXANDER COLLIE, M.D. Aberd., M.R.C.P. Illustrated with Coloured Plates, crown 8vo, 8s. 6d.

HANDBOOK OF DISEASES OF THE EAR: For the Use of Students and Practitioners. By URBAN PRITCHARD, M.D. Edin., F.R.C.S. Eng., Professor of Aural Surgery at King's College, London. Fourth Edition, with Illustrations, crown 8vo. [In preparation.]

A PRACTICAL TREATISE ON DISEASES OF THE KIDNEYS AND Urinary Derangements. By C. H. RALFE, M.A., M.D., F.R.C.P., Assistant Physician to the London Hospital, &c. With Illustrations, cr. 8vo, 10s. 6d.

DENTAL SURGERY FOR MEDICAL PRACTITIONERS AND STUDENTS of Medicine. By ASHLEY W. BARRETT, M.B. Lond., M.R.C.S., L.D.S., Consulting Dental Surgeon to the London Hospital. Third edition, with Illustrations, cr. 8vo, 3s. 6d.

PRACTICAL DEFORMITIES AND THEIR TREATMENT. A Handbook of Practical Orthopædics. By H. A. REEVES, F.R.C.S. Edin., Senior Assistant Surgeon at the London Hospital, etc. With Illustrations, cr. 8vo, 8s. 6d.

THE NEW SYDENHAM SOCIETY'S PUBLICATIONS.

President :—W. OSLER, M.D., LL.D., F.R.S.

Honorary Secretary :—JONATHAN HUTCHINSON, ESQ., F.R.S.

Treasurer :—HENRY POWER, ESQ., F.R.C.S.

ANNUAL SUBSCRIPTION, ONE GUINEA.

The Society is now issuing an important "Atlas of Clinical Medicine, Surgery, and Pathology," designed to form a pictorial guide to diagnosis. A volume of Selected Monographs, or a translation of some important Continental work, will be issued from time to time as well.

Amongst the works issued are Limbeck's "Pathology of the Blood," Helferich "On Fractures and Dislocations," Laveran's "Paludism," Pozzi's "Gynecology," Spiegelberg's "Midwifery," works by Charcot, Duchenne, Begbie, Billroth, Graves, Koch, Hebra, Guttmann, and a valuable and exhaustive "Lexicon of Medicine and the Allied Sciences," just completed.

The Annual Report, with full list of works published, prospectus of the new Atlas of Clinical Illustrations of Medicine and Pathology, and all further information will be sent on application.

PERIODICAL WORKS PUBLISHED BY H. K. LEWIS.

THE BRITISH JOURNAL OF DERMATOLOGY. Under the direction of H. G. Brooke, H. Radcliffe-Crocker, T. Colcott Fox, E. Graham Little, Sir Stephen Mackenzie, Malcolm Morris, J. J. Pringle, J. Herbert Stowers, Arthur Whitfield; Edited by James Galloway and J. M. H. MacLeod. Published monthly, 1s. Annual Subscription, 12s. post free.

THE GLASGOW MEDICAL JOURNAL. Edited by John Lindsay Steven, M.D., and T. K. Monro, M.D. Published Monthly. Annual Subscription, 20s., post free. Single numbers, 2s. each.

LIVERPOOL MEDICO-CHIRURGICAL JOURNAL, including the Proceedings of the Liverpool Medical Institution. Published twice yearly, 3s. 6d. each number.

TRANSACTIONS OF THE DERMATOLOGICAL SOCIETY OF GREAT BRITAIN AND IRELAND. Vols. I. to X., royal 8vo, 5s. each.

NORTH LONDON OR UNIVERSITY COLLEGE HOSPITAL REPORTS, SURGICAL CASES. 1901, 8vo, 3s. 6d. *nett*.

THE THERAPEUTIC GAZETTE. A Monthly Journal, devoted to the Science of Pharmacology, and to the introduction of New Therapeutic Agents. Edited by Dr. H. A. Hare and Dr. E. Martin. Annual Subscription, 10s., post free.

MIDDLESEX HOSPITAL, REPORTS OF THE MEDICAL, SURGICAL, AND PATHOLOGICAL REGISTRARS. Demy 8vo, 2s. 6d. *nett*. each volume.

* * MR. LEWIS is in constant communication with the leading publishing firms in America, and has transactions with them for the sale of his publications in that country. Advantageous arrangements are made in the interests of Authors for the publishing of their works in the United States.

Mr. LEWIS's publications can be procured of all Booksellers in any part of the world.

110





