On diet and regimen in sickness and health: and on the interdependence and prevention of diseases and the diminution of their fatality / by Horace Dobell.

#### **Contributors**

Dobell, Horace, 1828-1917. Royal College of Physicians of Edinburgh

#### **Publication/Creation**

London: H.K. Lewis, 1870.

#### **Persistent URL**

https://wellcomecollection.org/works/s8ank9sb

#### **Provider**

Royal College of Physicians Edinburgh

#### License and attribution

This material has been provided by This material has been provided by the Royal College of Physicians of Edinburgh. The original may be consulted at the Royal College of Physicians of Edinburgh. where the originals may be consulted.

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.


You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org DET AND REGIMEN
IN
BEAUTH AND DISEASE
DR. DOBELL.

3/6

bc1.51





# DIET AND REGIMEN

IN

## SICKNESS AND HEALTH

AND

ON THE INTERDEPENDENCE AND PREVENTION OF DISEASES
AND THE DIMINUTION OF THEIR FATALITY.

BY

# HORACE DOBELL, M.D.

SENIOR PHYSICIAN TO THE ROYAL HOSPITAL FOR DISEASES OF THE CHEST;

(Fourth edition, rewritten and much enlarged.)

H. K. LEWIS, 136 GOWER STREET, W.C. 1870.

LONDON: PRINTED BY H. K. LEWIS.

## PREFACE.

My "Manual of Diet and Regimen" which appeared in 1864 having run through two editions in its first year, I published a third and revised edition in January, 1865. This has long been out of print and, as the demand for the Manual did not decrease, I have been repeatedly urged to prepare a fourth edition. After many delays, from the interruption of other professional work, I have, at last, rewritten the book, adding much new matter and incorporating several contributions which I have published from time to time on subjects relating to the preservation of health. I hope it will be found that I have thus materially increased the usefulness of the work without adding inconveniently to its length. My object and endeavour has been to produce a small book founded upon accurate scientific data but essentially practical: a book from which a doctor may refresh his memory, and at the same time one which he may safely place in the hands of his patients, to enlighten them on some of those important points which ought to be understood by the non-professional, and to teach them how vast and intricate is the science and art of rational medicine. Thus, to make them more intelligent patients, and in so doing to remove

some of the doctor's difficulties in healing them when they are sick, and when they are well, to strengthen his hands in the preservation of their health.

It gives me great pleasure to be able to renew the thanks I expressed in 1864 to my friend Mr. Farrants, at that time President of the Microscopical Society, for the valuable assistance I have so frequently received from his remarkable analytical and arithmetical talents.

84 HARLEY STREET. May, 1870.

## CONTENTS.

#### CHAPTER I.

#### CHAPTER II.

Rules for Promoting and Maintaining Vigorous Health in Adults living in the Climate of the United Kingdom—Ventilation and Heating—Clothing and Protection from Cold—Sleep—London Noise and London Sleep—Exercise—Posture—Bathing—Regulation of the Bowels Rest and Change—Meals—Proper Hours for Meals

## CHAPTER III.

Food, Heat, and Motion—A foot-pound, a foot-ton, a British unit of heat—Mechanical Equivalent of a British unit of Heat—Combustion of Carbon, Consumption of Oxygen, Evolution of Heat—Mode in which the Heat Evolved from Food is disposed of—Composition of some of the Principal Articles of Food and their Heat Equivalents.

48

## CHAPTER IV.

OHAT IBIO IV.	
Essentials of a Normal Diet—Mode of Constructing Diets  —Expensive Diets and Cheap Diets for maintaining  Health—Tables and Analyses—Effects of neglecting  even one of the Essentials of a Normal Diet	
CHAPTER V.	
Fermented Liquors. — Acid and Sugar in Spirituous Liquors—Absolute Alcohol and Proof Spirit—Alcohol table showing the Analysis of Spirituous Liquors— Effects of Alcohol in Disease 6	3
CHAPTER VI.	
Fat essential to Healthy Nutrition—Importance of distinguishing between Solid and Liquid Fat—Bantingism—Getting Fat and getting Thin—Analysis of Bantingdiet, showing its Errors and Advantages	70
CHAPTER VII.	
On the Diet of Sickness—Principles upon which to arrange the Diet of Disease—Rules for carrying out these Principles—Examples of Diets for Consumption, and for Diabetes	76
CHAPTER VIII.	
The Duties of Medical Men—The Interdependence of Diseases shewing the importance of Periodical Examinations—Anæmia and Fatty Degeneration as Vestiges of Disease—Abnornal Physiological States—Conditions not recognised as Disease but which certainly are not health	91

#### CHAPTER IX.

											PAGE
Disi	nfecti	on-Med	ia by	which	h the	Ca	tching	Dise	ases a	re	
C	ommi	unicated -	- Me	ans fo	or Pr	ever	nting	Infec	tion a	nd	
C	ontag	rion-Dir	ection	s for	the S	Sick	Room	and	Atter	id-	
aı	nts—	Precautio	ns to	be	used	by	Docto	rs-	Mode	of	
C	leansi	ing Apart	ment	s afte	r Illn	ess-	-Prop	er Me	ethods	of	
us	sing	Chloride	of I	ime,	Carb	olic	Acid	and	Condy	y's	
F	luid.										134

Special Recipes, Directions, and Appliances for the Sickroom—I. Farinaceous articles—2. Beef tea—3. Leibig's Extract of meat—4. White of Egg, Eggs—5. Whey—6. Animal foods—7. Bread—8, Special Restorative—9. Invalid Soup—10. Combinations of alimentary principles in nearly exact normal proportions—11. Port Wine Jelly—12. Suet and Milk—13. Milk with Rum, Brandy, or Whiskey—14. Ardent Spirits and their peculiarities—15. Nutritive Enemata—16. Poultices—17. Fluids for Inhalations—18. Warm Baths—19. The Lights in a Sick-room—20. A Nursing Schedule. 140



# DIET, REGIMEN,

&c. &c.

#### CHAPTER I.

The Interdependence of Diseases—Narrow Specialities and Broad Departments in the Practice of Medicine—Normal Diet—Errors in Diet—The Diet of Disease—Food, Heat, and Motion—Getting fat and getting thin—Alcohol, its proper place in Diet—Regimen—The regulation of habits—Importance of sleep—Proper hours for meals—Disinfection—Pseudo-medical Dogmas, and rational Medicine.

In the practice of Rational Medicine, there are many subjects upon which the wise Physician is always glad to find that his patient has some knowledge as well as himself. Among these I know of none more important than the interdependence of Diseases and the principles of Diet and Regimen.

There are few medical questions which so test the depth and extent of a doctor's acquirements as THE INTERDEPENDENCE OF DISEASES—by which I mean, the way in which one disease leads to another, substitutes another, aggravates, or relieves another; the way in which disease appearing at one part of the body, depends upon the derange-

ment of another part, perhaps distantly separated; the way in which disease of one kind in an ancestor, leads to disease of several different kinds in the descendants; the way in which one general morbid cause may produce different effects upon different persons, according to the conditions of health in which they happen to be at the time; and numberless other similar phenomena.

These are matters of the highest importance in the right management and treatment of disease, yet they are so little understood that the pains taken by a conscientious doctor in their investigation are not appreciated by the majority of persons. Thus it is that patients so often fail to discriminate between the wisdom of the man who will not prescribe for a part of the body until he has learnt the condition of the whole, and the ad captandum ignorance of another man who will prescribe unhesitatingly for a disorder of the whole body without previously ascertaining whether it depends upon disease of a part.

The chief objection to specialities in medical practice is the danger, that the doctor who treats but one part of the body may lose sight of the interdependence of diseases. This must always be a grave objection to very narrow specialities, but it does not apply to the broad division of the practice of medicine into Departments—that is to say, such a reasonable application of the principles of the "division of labour" as shall enable each practitioner, not only to completely master the department of Medicine which he selects for special study, but to advance the general knowledge of the profession by his investigations and experience.

For example, diseases of the ear are often dependent upon, or are connected with, diseases of other parts of the body; and the doctor who confines his practice to diseases

of the ear may be in danger of losing sight of the broad facts of medicine, and in attempting to cure the ear may damage the body; but by extending his practice to all the organs of special sense, which might fairly form a department of Medicine, he would be obliged to take in a range of observation and thought which would enforce a know-

ledge of the interdependence of diseases.

Again—Diseases of the heart and arteries are intimately connected with those of the lungs and throat, diseases of the lungs with those of the heart, arteries and windpipe, diseases of the brain with those of the lungs, heart and arteries, and all of these are often inseparably connected, in the relation of cause or effect, with affections of the stomach, liver, pancreas, kidneys, intestinal glands, etc. etc. For a doctor, therefore, to limit his practice to diseases of the heart, to diseases of the throat, to diseases of the lungs, or to diseases of the windpipe, is to constitute a speciality so narrow that he is in danger of losing sight of the great principles of medicine. If, on the other hand, he selects for special study and practice such a broad department as Diseases of the Chest-that is to say, of all the respiratory and circulatory organs, and the associated, processes of digestion and assimilation—he will be obliged to take in a range of observation and thought necessitating a constant remembrance of the interdependence of diseases, and he will be competent to deal with all classes of disease as well as those peculiar to his own department. I have instanced Diseases of the Chest because it is one of the most important "lines" of practice, and the one with which I am most intimately associated, but the same remarks apply to Obstetrics, to Fevers, to the Diseases of Children, and to other large divisions of medicine. Thus departments of medical practice may be wisely selected by different medical men for special study-but

narrow specialities are dangerous and objectionable, unless those who confine themselves to them are content to act simply as the assistants of the more general physicians and surgeons, which is not likely to be the case. For those who may be interested to pursue this subject further I have introduced some extracts from my "Lectures on the germs and vestiges of disease." (See Index.)

DIET is so little understood that, very often, those are the best off who abstain from all attempts to meddle with it and are content to follow the dictates of their instinct. This, of course, ought not to be the case, for diet, properly understood, may be made a powerful agent in the restoration and maintenance of health; and errors of diet are at all times capable of becoming serious causes of disease.

But unfortunately, interference with diet, like all good things, is particularly open to abuse, for nothing is easier than to lay down a complicated code of restrictions and rules as to "what to eat and what to drink" and the patient is very apt to think that the skill of the doctor increases with the number and variety of his orders. But those who understand the principles of diet know that the reverse is much nearer to the truth, and that learning and skill in dieting a patient, are shown by the wisdom with which the doctor, instead of meddling with unimportant details, seizes upon the few essential points on which the vice or virtue of a diet will generally be found to turn. Thus, in a case of diabetes, the ignorant intermeddler may order fifty restrictions without doing his patient the slightest good; whereas, the doctor who understands the nature of the disease and the principles of diet, will speedily relieve his sufferings by telling him to take whatever he likes, so long as he touches nothing which contains starch or sugar. Thus, again, I have often seen patients suffering from acid dyspepsia who have been ordered and forbidden so many different articles of food that their lives were rendered miserable, without the slightest relief to their complaint, whereas, by forbidding them to take cheese and malt liquor, the chief factors of their malady, and allowing them to eat and drink whatever else they pleased, their sufferings have been speedily removed and their lives made enjoyable.

It is absolutely necessary that he who ventures to interfere with a person's diet should first understand the principles upon which the food of health is regulated in nature; and nothing assists the intelligent doctor more in getting his orders carried out, than for his patient, as well as himself, to be acquainted with this subject. Ignorance and stupidity are the constant obstacles in the proper regulation of these matters. To elucidate this part of the subject I have drawn up a brief statement of the "Essentials of a normal diet." This is illustrated by a set of tables, showing the method by which diets may be arranged so that they shall equally well support the human body in a state of normal health and strength, whether they consist of a complicated list of expensive articles, or, of no more than Bread, Cheese, Butter, Cresses and Water. (See Index.)

The advances of science have, of late, thrown most important light upon the connection between food and force, a subject of the greatest interest to all, whether medical or not; for, as we all take food and all desire to be strong, we are unavoidably fascinated by the enquiry what kind of food will give us the greatest amount of strength. I have not hesitated, therefore, to introduce a short paper on this subject which some may think rather dry. (See Index: "Food, Heat and Motion.")

The question of getting fat or thin, whether in health or disease, is one inseparable from the subject of "food and force," for we have discovered that heat is but a mode of

motion,\* and that those elements of food capable of liberating the largest amount of heat, may be utilized as accumulators of fat by economising the motion and heat of the body, and vice versa. Under this head I have introduced a table, shewing the composition of some of the most important articles of food, and the amount of heat to be got out of equal quantities of each, and I have also given directions how to estimate the amount of mechanical force equivalent to a given quantity of heat. (See Index). In order to elucidate the importance of fat, and the subject of getting fat or losing fat by alterations of diet, I have printed a portion of my letter on "Bantingism", written some time ago, and an article on the importance of distinguishing between solid and liquid fat. (See Index).

From what I have already said, and from the articles referred to, it must be clear that nothing can take precedence of proper food among the means for the preservation of the public health; and when health is lost, whether by unavoidable disease or by other causes, nothing can be more important than to understand the leading principles which should guide us in altering the food to meet the conditions of disease, and thus to convert diet into a powerful instrument for the restoration of health. This is a subject which could not be fully discussed in a small work like the present, but I have given a short lecture "On some Principles of Diet in Disease," which I delivered at the Royal Hospital for Diseases of the Chest, in 1865.

The subject of Alcohol which has of late years occupied so much attention, both in the scientific and in the religious world, and which has been so grievously misrepresented and misunderstood, is now placed upon a more reasonable foot-

<sup>\*</sup> See Prof. Tyndall's Lectures : " Heat a Mode of Motion."

ing, some plain facts having been unquestionably established by the dispassionate investigations of practical and scientific men. In this country we are particularly indebted to Dr. Anstie for his investigations on this subject; I have therefore introduced a leading article on alcohol, from the Lancet, which is, I believe, from his pen. (See Index.)

It is now quite certain, as I have long maintained, that alcohol is food—that is to say, that it is consumed in the body by a process of oxidation similar to that which other kinds of food undergo, that during its combustion force is liberated, and the demand for other kinds of food diminished. Alcohol, therefore, is able to save the combustion of the elements of tissue formation, by vicarious oxidation of itself.

It is quite certain that alcohol is one of the most readily oxidisable substances capable of being introduced into the animal organism. In this consists its chief value and its chief danger; for, on the one hand, it is able to replenish the lamp of life with so little tax upon the digestive functions and with so little loss of time, that nothing can take its place when these are the desiderata of most urgent import. But, on the other hand, this extraordinary facility for oxidation makes the presence of alcohol in the system a most dangerous impediment to the combustion of those less readily oxidisable matters, the full and regular combustion of which is absolutely essential to the continuance of normal health.

It is quite certain that alcohol is not an indispensable article of daily food; a normal diet may be perfectly well constructed without it, and a healthy person, under favourable circumstances, can live without it, just in fact, as the lower animals do. But it is also quite certain that alcohol is a most blessed gift of Providence, when rightly understood and wisely used. It is the endowment of man with those

higher attributes which principally distinguish him from the brutes, that constitutes the element in his nature which has made it necessary to his welfare that Providence should give him this peculiar article of food in addition to all those placed within the reach of the less intellectual animals.

Thus is it a fact, that while man is brutalized by the abuse of alcohol, its proper use is one of his distinctions from the brutes.

Alcohol may perhaps be best described as a *Medicinal Food*. It is essentially a poison, although in its proper use it is an indispensable element in the life and happiness of highly civilized society. But it ought always to be regarded as a most treacherous friend.

I have drawn up an "alcohol table," (see Index), the result of many careful analyses of each article so as to present a fair average result, from which it can be seen, at a glance, what quantity by measure of each of the ordinary fermented liquors represents one ounce by weight of absolute alcohol. The table includes a statement of the proportion of carbon and of some other matters, contained in fermented liquors, besides the alcohol; and it must always be remembered that different kinds of fermented liquors represent many other influences for good or evil, in a dietetic point of view, in addition to those possessed by the alcohol, all of which require consideration in selecting the particular beverage suitable to any given case.

It will be observed that I have omitted alcohol from all of the Tables of Normal Diet, leaving it to be added in any form and quantity which circumstances may render most advisable. According to my own experience, based upon long and careful observation, an average-sized adult man taking moderate exercise may drink with advantage enough

fermented liquors, each 24 hours, to represent from one to two ounces avoirdupois of absolute alcohol—provided it be always diluted to the extent of 10 fluid ounces (half-a-pint imperial) of water or some other unfermented liquor to each ounce avoirdupois of absolute alcohol, and that it be taken when there is food in the stomach. (See Index).

REGIMEN. The regulation of habits could not be fully discussed without involving physiological, and other details which would have far over-run the limits of this book .- I have, therefore, contented myself with laying down, as concisely as possible, such rules as my own experience, added to that of others, has taught me are most necessary and most suitable to the generality of persons. They must, of course, be modified to suit peculiar constitutions, idiosyncrasies, and other circumstances, and to meet the various conditions of disease. In connection with these Rules for Regimen, I have introduced a letter "on the Proper Hours for Meals." (See Index), and an important leading article from the Times of August 17th, 1869, suggested by a letter of mine on "London Noise and London Sleep," which had appeared on the 13th. (See Index). On the subject of exercise and training, in addition to the rules I have laid down, I advise everyone, medical or not, to read "A system of Physical education," an admirable treatise issued from the Clarendon Press in 1869, by Archibald Maclaren, of the Gymnasium, Oxford.

As a necessary part of a work like the present, having for its object the preservation of health, I have added a chapter on DISINFECTION, with rules for preventing the spread of the "Catching Diseases." And at the end of the book will be found a few special directions for the sick room.

In concluding this introductory chapter I must not

lose the opportunity of expressing my opinion upon a matter regarding which there is great misapprehension, and, I regret to say, intentional misrepresentation. I refer to the subject of PSEUDO-MEDICAL-DOGMAS, such as are involved in the terms Allopathy, Hydropathy, Homoeopathy, Kinesipathy, &c., and in such expressions as "a stimulating system," "a depleting system," "similia similibus, curantur," "contraria contraribus curantur," and the like.

I disclaim them all, and I feel sure that in this statement I express the sentiment of the majority of thinking medical men. It is utterly unworthy of the professor of a science and art so great as medicine, to "pin himself" to such narrow dogmas and rules of thought and practice as are indicated by one and all of these expressions.

The practice of BATIONAL MEDICINE—having for its subject, not only that most complex and wonderful fabric the animal organism, but that organism endowed with all those attributes which place man but "a little lower than the angels"—demands that there shall not be anything excluded from its service, and that every science and every art shall combine, at its bidding, to assuage the suffering and to save the life of Nature's highest earthly creature.

Every honest and intelligent practitioner of rational medicine knows, that there are remedies of unquestionable potency the action of which could never have been discovered by any such dogmas as "contraria contraribus curantur" or "similia similibus curantur." On the other hand, he knows that there are remedies the action of which may appear to be explained by one or other of these principles, but that the more intimately we become acquainted with the occult properties of medicines and the occult physiology of diseases, the more plainly do we see that these apparent explanations of the modus operandi of remedies are ab-

surdly superficial and incorrect. He will not, then, refuse to benefit his patient by the use of the one remedy because he cannot explain its action, or of the other because it appears to act by similarity, or of a third because he thinks it acts by contrariety. Again, he knows that the same disease may assume such different phases, in different constitutions, at different times and in different places, that in one case it may require "stimulation," in another "depletion," that at one period of the same case, "water treatment" may be advisable, at another "gymnastic," at another "mechanical," at another "climatic." And he claims these, and all other means beneficently placed within our reach which can protect health or benefit the sick, as the legitimate weapons in the armoury of rational medicine.

Every theory, every system, every dogma, must give place, at last, before the fact, once unquestionably proved, that such or such a remedy is best for such or such an occasion.

Therefore, in my opinion, all men who "pin their faith" on narrow dogmas, stamp themselves thereby as unfit for the practice of rational medicine.

#### CHAPTER II.

Rules for Promoting and Maintaining Vigorous Health in Adults living in the Climate of the United Kingdom—Ventilation and Heating—Clothing and Protection from Cold—Sleep—London Noise and London Sleep—Exercise—Posture—Bathing—Regulation of the Bowels—Rest and Change—Meals—Proper Hours for Meals.

\*\* These Rules will require to be modified by the medical man to suit special cases.

## VENTILATION AND HEATING.

- 1. No sitting or sleeping room should be left long without a fire, and every room in which persons live, either by day or by night, should have some opening by which it communicates directly with the outer air; but this should be so arranged that no draught can fall upon the persons in the room.
- 2. If several rooms are occupied by turns during the twenty-four hours, the temperature of any one should not differ greatly from that of the rest.
- 3. No draught should blow upon a bed, and during sleep the whole body should have one covering at least of woollen material; for, while it is very important to keep

the air of sleeping rooms fresh, it must be remembered that the body is more susceptible to chills during sleep than waking, and that changes in the temperature of the outer air are especially apt to occur during the night, and are, therefore, in danger of producing chills before they are observed. (See Sleep.)

4. All arrangements for ventilation must be based upon

the following facts :-

The rate of respiration in an average sized adult man is about 16 times per minute, and each such respiration vitiates about 1 cubic foot of normal atmospheric air; so that each adult man vitiates 960 cubic feet of air per hour, and consequently will require a supply of fresh air at that rate; or in round numbers 1000 cubic feet per hour.

This supply of fresh air can be provided by observing

the following conditions:-

A current of air travelling at the rate of 36 linear inches per second is not perceptible as a draught, and at that rate of movement 1½ cubic feet per minute or 75 per hour will be admitted by an aperture of 1 square inch sectional area, communicating directly with the outer air.

If, therefore, an apartment is provided with an aperture having a ratio of 1 square inch sectional area to each 75 feet of cubic capacity, the whole air of the apartment will be changed once per hour.

Supposing, then, that the cubic capacity of an apartment is in the proportion of 1000 feet to each person in it, the requisite supply of fresh air may be obtained, without draught, through an aperture of from 13 to 14 square inches sectional area to each 1000 feet of cubic capacity. But if the cubic capacity of the apartment is only in the proportion of 500 feet for each person in it, the air must be changed twice per hour in order to supply the requisite 1000 cubic feet of fresh air for each person.

When gas is burnt in an apartment the following facts must be borne in mind, in addition to the foregoing, for the purposes of ventilation:—

One cubic foot of ordinary London coal gas produces, in burning, 2-thirds of a cubic foot of carbonic acid, about the amount produced per hour by the respiration of one adult man,—and as an ordinary fish-tail gas-burner consumes about 3 cubic feet of gas per hour, it produces in that time as much carbonic acid as the respiration of three adults.

#### CLOTHING.

- 5. In winter, the body and limbs, from the root of the neck to the toes and elbows, should be covered, next the skin, with some woollen material, such as lamb's wool or flannel.
- 6. In summer, the material may be lighter, as merino, and need not cover the limbs.
- 7. The same woollen dress should not be worn both night and day, but should be changed for a woollen sleeping vest of the same material as that worn during the day. (See Sleep).
- 8. Having provided that the skin is so covered that it is protected against sudden changes of temperature, all other clothing should be limited to that which is sufficient to preserve a comfortable feeling of warmth under the different changes of the season and of the weather.
  - 9. Over-clothing, i.e., such as keeps the body perspiring

while at rest or produces perspiration under very slight exertion, should be avoided, especially over-clothing of any one part of the body by which it is kept hotter than the other parts.

10. It is of the greatest importance to keep the feet dry and as warm as the rest of the body. If the weather is damp, this can only be done by wearing goloshes when out of doors. (See Bathing).

This subject may be further elucidated by the following extract from the Author's Lectures on Winter Cough. Catarrh, Bronchitis, Emphysema, Asthma, delivered at the Royal Hospital for Diseases of the Chest in 1865.

Fresh colds were the only causes which brought on or aggravated the cough in 72 per cent of the cases, and the most frequent and potent causes of these fresh colds, as stated by the patients themselves, were:—

- 1. Sudden changes of temperature in 21 per cent.
- 2. Fogs and damp air in 19 per cent.
- 3. Draughts of cold air in 16 per cent.
- 4. Cold winds in 10 per cent.
- 5. Getting wet in 14 per cent.
- 6. Wet feet in 17 per cent. And we find the same list of causes of fresh colds recurring as the potent provokers of short breathing. This important list is no less striking for the power for evil which it is shown to possess, than for the remarkable simplicity of the evil powers themselves.

If, then, these unfortunate sufferers from Winter Cough could have been protected from sudden changes of temperature, fogs, mists, cold winds, draughts, wet feet, and wet coats and dresses, 72 per cent would have kept free from their coughs. It would seem strange if we could not find means of protection against such common-place influences. In truth, there is no deficiency of means of protect-

tion against them, and it is because of the very commonplace character of these means of protection and of the influences themselves that both are so much neglected and under-valued.

But 72 per cent of the cases of Winter Cough which I have analysed might probably have been prevented by attention to these common-place things. Let us then give a few minutes to their consideration.

1. Sudden changes of temperature. This is the most difficult to avoid of any on the list. The occupations and amusements of all classes involve such changes, and we cannot stop these occupations and amusements, even were it desirable to do so. The workshop, the counting house, the committee-room, the opera-house, the ball-room, must be warm when the outer air is cold, and changes from one to the other cannot be avoided. But very much could be done to prevent the body from feeling these changes. The first and most important is the complete envelopment of the body and limbs in wool next the skin, thus interposing a bad conductor of heat between the surface of the body and the outer air. It is surprising that even at the present day this simple and common-sense protection is neglected by so large a number of persons both of the educated and of the uneducated classes. It is not sufficient for the purpose in view that a little body-vest should be worn just big enough to cover the thorax and abdomen, leaving the extremities unprotected. It should be insisted on by medical men that the arms and legs require to be protected from sudden transitions of temperature as well as the trunk.

In fashionable life the greatest practical difficulty we have to encounter is the question of exposing the necks and shoulders of ladies in evening dress. It is useless to order body-clothing of wool to the throat, and to expect

that ladies will give up a fashion which has been followed and thought charming in all countries and all ages. The difficulty is, however, to be got over pretty well. Every lady in evening dress should carry with her, as invariably as she does her pocket handkerchief, a Shetland shawl or a mantilla of wool or fur, of a size and shape to cover all those parts not protected by woollen underclothing, and it should only be removed while actually within warm rooms and should be kept at hand to replace on passing through passages, or if the rooms become cold, or if sitting in draughts.

The main source of protection, then, against sudden changes of temperature to the surface of the body, is to be found in a complete covering of wool next the skin. But besides this, much greater attention, than is common, should be paid to putting on and taking off complete and efficient over-clothing when going from hot to cold and from cold to hot temperatures. This is particularly neglected by the working classes and by girls and boys at school. ..... What I have said with regard to sudden changes of temperature will apply equally to two other causes of fresh colds on our list, viz., draughts of cold air, and cold winds. Both are to be deprived of their sting by proper clothing of the skin and mucous orifices—of the latter I shall speak again.

Getting wet, and wet feet occupy a very serious place in our list, and there is no doubt that damp and cold applied to the general surface is a most efficient means of producing chill and vital depression, with congestion of the internal organs. It is necessary that cold be combined with moisture to produce this effect.

Even if all the clothes on the body are wet, no harm will come so long as they are kept warm, and this suggests the very great value, to all persons liable to expos-

ure to wet, of light waterproof over-alls. They may either be put on to keep the underclothing dry, or, if the underclothing has become wet, either by weather or by perspiration, they may be put on to prevent too rapid evaporation and consequent reduction of temperature, especially when the person is about to remain still after getting warm with exercise. In this variable climate, therefore, school girls, governesses, shop and factory girls, and all women whose occupations call upon them to brave the weather, ought to carry with them complete water-proof mantles made as light as possible, but extending from the neck to the ankles, which can be put on or not as required; and boys and men similarly exposed should carry waterproof over-alls. But if wet and cold to the surface of the body is a fruitful source of catarrh, wet feet-which means wet and cold feet-is a still more prolific source. There is no external influence which so surely produces congestion of the naso-pulmonary mucous membrane, as wet and cold to the soles of the feet. There is nothing so universally neglected, and yet there is nothing more easy to avoid. Warm socks, horsehair soles, goloshes, provide efficient protection against wet and cold feet. It does not seem to be half understood that, although a shoe or boot may not be wet through, if the sole is damp it will, by conduction and evaporation, most effectually carry away the heat from the sole of the foot, and, therefore, ought never to be worn after exercise is over. . . . .

We have still one item left on our list—viz., Fogs and Damp air, which were the things most inclined to make the breath short in 24 per cent of the cases, and the most potent causes of fresh colds in 19 per cent. I have particularly remarked, that although the smoke and other irritating matters constituting fog are unquestionably very injurious, it is the moisture and cold of the fog which are

the qualities most potent for mischief to the naso-pulmonary tract. There is but one means of depriving a fog or mist of its injurious properties, and that is a respirator; and the same may be said of the changes of temperature of which I spoke just now; a respirator is the only means of protecting the respiratory passages from the effects of transitions of temperature. . . . .

Although it is quite proper to cover the neck and throat lightly, I am decidedly of opinion that warm wrappers round the neck are objectionable; they produce congestion of the nasal and faucial mucous membrane and thus dispose to the very complaints they are supposed to prevent".—p. 148 to p. 153 Lectures.

I am glad to find the use of respirators, which I have so long advocated, recently demonstrated in one of Professor Tyndall's beautiful lectures, of which the following abstract appeared January 29th, 1870, in the British Medical Journal.

# "Dust and Disease, A Lecture delivered at the Royal Institution."

"Professor Tyndall, in making some experiments on vapours, wished for a current of air quite free from the particles of dust always seen in a beam of sunlight. He tried various means for this purpose; one consisted in passing air through a tube filled with sulphuric acid; another, in passing the air through a tube filled with solution of potash. In each case, particles, capable of refracting light, and rendering themselves visible, were still present. In October 1868 he hit on the plan of allowing the air to pass over the flame of a spirit-lamp. 'The floating matter no longer appeared, having been burnt up by the flame. It was, therefore, organic matter. If the air were

sent too rapidly through the flame, a fine blue cloud was noticed. This was the smoke of the organic particles.' The Professor was not prepared for the discovery that the dust of our air wss organic. He had always considered it inorganic and non-combustible. Mr. Valentin now furnished him with a small gas furnace with a platinum tube, which could be heated to redness. Air was passed through this tube when cold, and then when hot. When combustion was perfect, no particles could be detected."

"Further experiments led to still more interesting results. A beam of light was made to illumine the dust of the laboratory, and the flame of a spirit-lamp allowed to play on it. Wreaths of darkness were at once seen to mingle with the flame, just like intensely black smoke. 'When the flame was placed below the beam of light, the same dark masses steamed upwards.' They were at times blacker than the blackest smoke. A red-hot poker placed under the beam produced the same dark wreaths. A large hydrogen flame led to the same result. Smoke was therefore out of the question. What, then, was the blackness? Simply that of stellar space resulting from the absence, from the track of the beam, of all matter capable of scattering its light."

"The Professor then remarked: 'Nobody could, without repugnance in the first instance, place the mouth at the illuminated focus of the electric beam and inhale the dirt revealed there. Nor is the disgust abolished by the reflection that, although we do not see the nastiness, we are churning it in our lungs every hour and minute of our lives.' The wonder is, that so small a portion of this dirt should appear to be really deadly to man. What is this portion?"

"The lecturer then alluded to the notion, at one time prevalent, that malarious diseases were due to organic matter in a state of decay (fermentation). It was then shown that fermentation really depended on the growth of the yeast-plant. Further, Schwann, in 1837, showed that meat, in contact with air which had been heated, did not putrefy; and he affirmed that putrefaction was caused by something derived from the air, which could be destroyed by high temperature. The germ-theory of epidemic disease soon followed, and found an energetic supporter in Sir Henry Holland, the present President of the Institution. The spread of cholera and that of small-pox were adduced as instances in support of the germ-theory."

"Professor Tyndall alluded to the difficulty that must be experienced in freeing surgical instruments (a canula, for instance) from the means of carrying contagion, in the presence of an atmosphere such as ours, unless a high temperature were employed, and this is not done. Thus, not-withstanding all the surgeon's care, inflammation often sets in after the use of such an instrument. When an abscess has been tapped, the pus, which was at first sweet, becomes feetid and swarms with vibrios. Professor Lister's views were quoted."

"The Professor then went back to the dust. It could not be blown away with a pair of bellows; but, if the muzzle of the bellows were stuffed with cotton-wool, it was found that the air which escaped was free from particles. Schræder used cotton-wool as a filter in his experiments on spontaneous generation; and subsequently it was used in those of Pasteur. Since 1868 Dr. Tyndall has used it himself."

"The most interesting and important illustration of such a filtering process is furnished by the human breath. After inspiring a quantity of common air, a long expiration is made through a glass tube across the electric beam. At first the luminous track is uninterrupted. The breath im-

22 SLEEP.

presses on the floating matter a transverse motion, but the dust from the lungs makes good the particles displaced. After a time, however, an obscure disc appears upon the beam, and, at the end of expiration, the beam is, as it were, pierced by an intensely black hole, in which no particles whatever can be discerned. The air, in fact, has lodged its dirt in the lungs. A handful of cotton-wool placed over the nose and mouth during inspiration makes the dark hole in the beam of light appear from the beginning of expiration. A silk handkerchief answers nearly as well, but the filtration is not nearly so perfect as with cotton-wool."

"In conclusion, the use of cotton-wool respirators to exclude the entrance of the germs of contagious diseases by means of respiration, was strongly advocated. 'By means of the respirator, so far as the germs are concerned, the air of the highest Alps may be brought into the chamber of the invalid.'"

## SLEEP.

11. During ordinary health, the hours spent in actual sleep should not exceed eight; and if the sleep is sound, continuous, and refreshing, six will be sufficient for many

persons. (See Rest and Change).

12. During sleep, it is equally important to keep the body pleasantly warm, and to avoid keeping it over-heated; and, as serious changes in the weather may happen in the night, and the lowest temperature in the twenty-four hours naturally occurs between 2 o'clock and 6 o'clock a.m., the

ventilation and clothing must be prepared for these contingencies. (See Ventilation, and Clothing).

- 13. If sleep is taken after meals, it should not exceed half an hour in duration; it should be taken sitting back in an easy chair, with the head supported behind; not lying down, and not sitting with the chin resting on the breast; the feet should be kept warm, and the dress loose round the neck and waist.
- 14. Every precaution should be taken to secure quiet sleeping-rooms.

Letter by the Author on London Noise and London Sleep, published in the Times, August 13th, 1869.

TO THE EDITOR OF THE TIMES.

Sir,—The London season is over once more, and everyone is hurrying to get a dash of sea water or of mountain
air, before returning to London work. Brain-fag, nervous
exhaustion, and worry are the universal complaints; and
who can wonder at it? For where and when in this great
city, I should like to be told, can anyone secure six hours
of undisturbed sleep?

If insanity increases, if doctors are more busy every year with diseases of the nervous system, if men and women wear out faster and faster, who can wonder, if he will take the trouble to consider, how utterly our municipal arrangements ignore the necessity for sleep?

It is time, indeed, that we turned our attention to this vital question, upon which hangs not only the comfort and happiness of our best men and women of the day, but the actual value of their mental work, the continued sanity of their minds, and the mental calibre of their progeny. I am not the only member of my profession who has long

desired to bring this matter into public notice; far otherwise. Your columns might easily be filled day by day with the sore experiences of medical men on this point. But I see the evil increasing year by year, and I cannot longer keep silent when I see "holiday tasks" set to thinking men, to rob them of the only chance they had left of recreation and rest.

If some of us are to be doomed to spend the recess in digesting education Bills, and in speculations on the value of clipped sovereigns, at least let some others of us consider the value of a little sleep.

This is a very simple question, and the evil appears to be susceptible of a simple remedy. It is a question of police; and attention being once prominently called to its importance there can be no excuse for its being longer neglected. The police should be made to keep our streets quiet during certain hours of the night. No attempt at this is made at present. The night policemen walk tacitly up and down, while every house in a street is being roused by the most abominable noises, without making the slightest attempt at checking them. It might not unfairly be asked that people should have a chance of sleeping from 12 o'clock till 8; but in the name of all that is sane, let them have the possibility of sleeping from 2 o'clock till 8.

There is no such chance now. No one interferes to stop any amount of noise in the night and early morning. A party of cats may hold an uproarious concert in the middle of the road without even a "hiss" from the policeman to disperse them. Two "cabbies" may career down the opposite gutters, and hold a conversation across the road at the top of their voices. A train of scavengers' carts may be driven down the streets, rumbling like thunder, while the driver in the last cart holloas his jokes to the

man in the front. In some districts it is thought necessary to create the most infernal noise about 5 o'clock in the morning by setting a host of garrulous old men to scrape and stone the roads at that pleasant hour. On Sunday mornings the paper boys are allowed to bawl with all their might. At any hour of the night, a fool in love with a concertina may disturb a whole neighbourhood with the noise he pleases to think music; and no interruption is given to any number of drunken rollickers who choose to sing and holloa up and down our streets and squares. I have said nothing of the early organ-grinders, collectors of hares' and rabbits' skins, sellers of water-cresses, the inevitable dustman, the rows attendant upon balls and receptions, or a hundred other sleep preventers, too painfully familiar to those who turn into their beds between 12 and 3 o'clock in the morning, their brains fagged and excited by work-Parliamentary, scientific, judicial, professional, it matters not which, or even by those unavoidable and wearying pursuits of social life which we call "society." Of the sick I dare not even speak, for to them this everlasting noise is simply murder. But for those in ordinary health, for those who are doing the brainwork of the nation, I ask for some chance of at least six hours' quiet sleep out of the 24. Even this moderate allowance might suffice to restore the wear and tear of the day and refresh the spirits for the ever recurring fight. But such sleep is utterly impossible, as a rule, in the present state of things, and hence the yearly increasing decay of mental and nervous health.

It needs but your aid, Sir, to procure a remedy. With proper police regulations the noisiest place in London can be kept comparatively quiet, as the citizens have shown in the main thoroughfare through their new meat market in Smithfield. There a couple of policemen, properly in

structed, stop all unpleasant noise. I trust, then, that before London assembles again for work some steps may have been taken to give us quiet streets for at least six hours out of the twenty-four.

I am, Sir, yours truly, M.D. Harley-street. Aug. 11, 1869.

Leader from the "Times" August 17, 1869, referring to the Authors letter.

"Our correspondent 'M.D.' may, we think, claim the credit of having suggested, or at least drawn public attention to, a new remedy for a very old and common complaint. The complaint cannot, indeed, be altogether got rid of without revolutionizing our whole system of modern life, for it simply amounts to this-that men now-a-days work so hard, and live in such a constant whirl of excitement, that neither body nor mind gets sufficient recreation or rest. The mind is naturally the greater sufferer, for the obvious reason that it may be kept almost incessantly in harness, and may be working its very hardest, while the body, comfortably stretched on bed or sofa, is enjoying luxurious repose. It is scarcely too much to say, for instance, that men of anxious temperament engaged in commercial speculations during critical times do not, from the moment they open their eyes in the morning to the moment they close them at night, know what it is to have ten minutes altogether free from absorbing calculations how to avoid this possible piece of bad luck, or turn good luck to the best advantage. Even when they try to banish these exciting topics-an effort, however, which over-anxious minds seldom make very strenuously—their brain, nevertheless, will sometimes work in spite of them. And what happens to them happens, if in less degree, to nearly all who have to win their bread in professions and occupations where competition is keen. Hence it is, as 'M.D.' points out, that we see what are vaguely called 'nervous complaints' steadily on the increase, often leading, if they are not at once taken vigorously in hand, to confirmed ill-health, and sometimes even to insanity. 'It is time,' he declares "that we turned our attention to this vital question, upon which hang not only the comfort and happiness of our best men and women of the day, but the actual value of their mental work, the the continued sanity of their minds, and the mental calibre of their offspring."

"We are afraid that nothing short of a miracle could provide a complete remedy for the evil of which our correspondent complains. We should require a second Joshua to bid modern civilization stand still or change its course. There are, indeed, philosophers who look hopefully forward to the time when men will be content to do just enough work to supply them with the means of gratifying their proper mental and bodily wants-when they will not be for ever jostling and struggling with each other in the arduous race for power and wealth. The philosophers may, for aught we know, be right, and it at least gratifies the benevolent instincts to picture our over-worked world some day settling down by universal consent into a blessed state of repose. But certainly there are at present no signs of this coming millennium. The nineteenth century notion of 'enough' means 'always a little more.' Thus A may have plenty to eat and drink of the best, a comfortable house, a charming wife and family, a large balance at his banker's to spend in the most pleasurable

ways prompted by a cultivated taste-may, in short, have all that the philosophers consider essential to human happiness; but all this avails nothing to keep A quiet while B remains a yard ahead of him in the race of life, and no sooner is B distanced than he becomes equally solicitous about C. Nor is this always A's own fault, if fault it is to be considered, or his deliberate choice. He cannot stand still if he would, since to remain stationary where everybody else is moving on is to lose ground which cannot, perhaps, be made good. The successful merchant or professional man toils in his office all day, and then goes home late at night too tired and worn out to enjoy the home comforts and luxuries with which his wealth has enabled him to surround himself. If you tell him that he is throwing all his opportunities away, that he might as well be a poor man without these advantages as rich without time for enjoying them, and that he would be wiser to content himself with doing two-thirds of his present work for two-thirds of his present pay, he will probably reply that he gladly would do this if he could, but that he has no choice. He cannot do just as much work as he likes, and leave the rest undone, without danger of losing his position altogether. In the general rush of eager competitors from all quarters, he must go ahead with the crowd, or risk being left far behind, and perhaps trodden down by it. When a man has attained so secure a position that he can please himself about taking or rejecting work, it usually either is time that he should retire upon his laurels, or else, if his health is good, long habit has made constant work such a second nature that he cannot get on without it. We are afraid that, until the promised millennium provides some substitute for the keen spirit of competition, as understood and practised in the nineteenth century, there is no hope of escaping altogether

from the growing evils of over-work, since every new invention which diminishes space and time, or otherwise multiplies facilities for communication between man and man, only supplies fresh materials for the competitive spirit to work upon."

"However, that we cannot effectually remedy the evils of over-work, and that it is steadily on the increase, are only additional reasons for adopting what partial remedies we can find; and 'M.D.' has at least one to suggest well worth public consideration. The best cure for over-work is obviously rest, and the best kind of rest is sound sleep at night. The first question usually that the doctor asks a patient suffering from 'nervous exhaustion' 'worry' or 'brain-fag' is whether he sleeps well, and, if the answer is in the negative, every plan is tried to secure him that surest and safest of all restoratives, a sound night's rest. Yet in London, as 'M.D' bitterly complains-and the complaint is repeated by other correspondents, whose letters we publish to-day-in which, as the busiest centre of industrial and social life, the wear and tear of overwork is greatest, and the need for the best kind of rest proportionately pressing, no attention whatever is paid to procuring quiet nights. Anybody and everybody is allowed, without let or hindrance, to make night hideous by whatever discordant sounds may suit his jovial or fantastic mood, his pleasure in the performance being probably enhanced by the thought that he is disturbing all the commonplace, prosaic people who have tamely slunk off to bed instead of 'enjoying life' as he does. Our correspondent, 'M.D.,' graphically described with the indignant energy apparently of a victim, the various nocturnal noises which in London murder sleep. It is not necessary here to re-enumerate them, or to say more about them than that they nearly all-certainly all the most objectionable-might be, if not altogether stopped, at least kept within less troublesome limits by the police. The police, for instance, cannot prevent cabs from remaining in or driving about the streets all night, but they can prevent the cabmen from carrying on a conversation in Stentorian tones with each other or the watermen or any late reveller who provokes them to a wordy war; and although the reveller himself cannot be kept from wandering all night about London, he may at least be kept quiet without any unwarrantable infringement upon the liberties of the British subject, and it is to revellers that much of the nightly disturbance in town is due. Besides, the interference of the police, independently of its immediate and special effects, would generally exercise a beneficial influence by directing public attention to a question which, notwithstanding its vital importance, is now strangely neglected. Unless a law is so unpopular that people deliberately rebel against it, its natural tendency is to educate the public mind into a spontaneous appreciation of the practice which it enjoins. Many Londoners, otherwise harmless and well-disposed enough, are often noisy at night simply because the duty of keeping quiet and allowing their neighbours to sleep undisturbed has never been fairly brought home to them. But if the violation of this duty came to be recognized as an offence, not merely in theory, but in practice, rendering the offender liable to be locked up some hours in a police-cell and punished next morning by a magistrate, the duty itself would acquire its proper importance in the eyes of all respectable law-regarding people. Nor would the indirect effects of police interference cease, we think, here. In various other ways attention would be drawn to the necessity of securing quiet nights in London. Streets would be paved, houses and windows be constructed, and even bedrooms chosen with

much greater regard than at present to this point if its importance were more generally recognized—if, in fact, it became one of the standing sanitary arrangements of the metropolis. Now, although a vast number of Londoners suffer sadly from want of sleep at night, they have learnt by long habit to look upon it as a necessary evil, inseparable from London life. They bear with it, because their only notion of escaping from it is to rush off to some quiet place in the country where there are no night cabmen or drunken revellers, and by most Londoners the remedy would probably be considered worse than the disease. Yet, for all this, they do not at all relish the disease, but would gladly be rid of it, and if the police will take the matter up and endeavour gradually and judiciously, by making a few well-chosen examples, to protect the streets of London and other large towns from needless disturbance between, say, midnight and 6 or 7 o'clock in the morning, we are quite sure they will receive the thanks of the community, and may rely on public support."

#### EXERCISE.

15. During ordinary health, some part of every day ought to be spent out of doors; and in ill-health it is of great importance not to discontinue the observance of this rule without good reason; for, although in eertain states of disease it may be very important to remain in-doors, it must not be forgotten that proper clothing, goloshes, respirators, and umbrellas, may make it not only safe but advantageous

to go out of doors for exercise, when, without them, it would be very injurious. (See Posture).

16. Out-of-door exercise should be as active as the strength will allow, and should always be continued up to the point of slight—but not over—fatigue. This will be the best measure of the proper amount for both the weak and the strong.

17. Unless the air is pure, and the person strong, exercise before breakfast is more likely to do harm than good; a tumbler of milk and a biscuit, however, will be a sufficient meal to take before the walk or ride—a more substantial breakfast being taken afterwards. (See Meals).

18. Especial care is needed not to expose the body to chills when heated by exercise; and cold drinks should not be taken at that time, unless the exercise is about to be continued immediately; and even then the quantity of cold drink taken at once should be very small. (See Meals).

The following excellent remarks are from Mr. Maclaren's work "on Physical Education" referred to at p. 9.

"Such in brief is Exercise, such the ends which it accomplishes, and such the manner of their accomplishment; viz. the destruction of the tissues, the hastening of the decay and death of every part coming within its influence; but also the speedy removal of all waste, the hastening forward of fresh material for its replacement; and in doing this it attains three distinct but co-relative results.

1. It increases the size and power of the voluntary muscles employed.

EXERCISE 33

2. It increases the functional capacity of the involuntary muscles employed.

3. It promotes the health and strength of the whole body, by increasing respiration and quickening circulation." p. 15.

"It is health rather than strength that is the great requirement of modern men at modern occupations; it is not the power to travel great distances, carry great burdens, lift great weights, or overcome great material obstructions; it is simply that condition of body and that mere want of vital capacity, which shall enable each man in his place to pursue his calling, and work on in his working life, with the greatest amount of comfort to himself and usefulness to his fellow men. . . . . Let it not be inferred from this that I consider health and strength as in any manner opposed to each other, on the contrary, they are most intimately allied and are usually, by the same means and by the same manner obtained, very closely are they connected but they are not the same, and a man may possess either without the other." p. 24.... A most important principle in Exercise, and one which should ever be borne in mind, is, that it should be regulated by individual fitness, for the exercise that scarcely amounts to exertion in one person will be injurious and dangerous to another, and not only is this inequality observable among different individuals, but, the same individual may have parts of his body possessing special power or presenting special weakness. ' man may have limbs capable of transporting him at he rate of four miles an hour throughout the day, and for many days in succession, but with heart or lungs all unequal to the effort. Or he may have an organization so frail and a temperament so susceptible to stimulation or excitement, that the one is an abiding danger to the other." p. 55.

#### POSTURE.

19. It is of great importance to acquire a habit of drawing the breath deeply and slowly, so as freely to expand the lungs during ordinary breathing. This requires that the head and shoulders be thrown well back in walking, sitting, and standing, and that no clothing be worn tight round the ribs. Those engaged in sedentary or stooping occupations should especially attend to this advice.

20. Those whose occupation obliges them to maintain the erect posture for a number of hours each day, should take every opportunity of lying flat down, even if only for a few minutes at a time. They should also bear in mind that standing will not take the place of walking exercise. (See Exercise).

21. All persons whose pursuits require the long continuance or frequent assumption of any particular position or movement of the body or limbs, should take every opportunity of changing it for an opposite position or movement. (See Rest and Change, and Sleep).\*

#### BATHING.

- 22. Warm baths, Turkish baths, vapour baths, shower baths, and cold plunges, should only be used under special medical orders.
- 23. During ordinary health, the skin of the body and limbs should be smartly rubbed once in twenty-four hours,
- \* For "Gymnastics" see Mr. Maclaren's "System of Physical Education," already quoted. For Effects of unhealthy posture, see Mr. Heather Bigg's "Orthopraxy."

first with a rough towel, wet with cold water, and then with a dry one till in a glow. The bather should stand on a dry rug while using this "cold friction bath," and it should not last more than one or two minutes, including both the wet and the dry rub. Salt may be advantageously added to the water; and the bath may be used either on rising or going to bed, according to the feelings and convenience of each individual. When water cannot be borne cold, it must not be used tepid, but scalding hot.—The momentary application of very hot water to the skin, immediately followed by a brisk friction, will produce a direct-action glow nearly approaching the re-action glow following the application of cold.

- 24. If the weather is very cold or the person delicate and chilly, the upper half of the body should be uncovered and rubbed first, and then a woollen vest should be put on and the lower half uncovered and rubbed.
- 25. It is well to accustom the feet to being washed in cold water, but it must be done cautiously at first, and they should never be kept in the water more than a few seconds.
- 26. It is obvious, that there are times when cold bathing of all kinds must be temporarily discontinued.

# REGULATION OF THE BOWELS.

27. As a general rule, the bowels ought to act, at some stated time, once in the twenty-four hours; and it is best to accustom them to act in the morning, after breakfast.

28. If they do not act spontaneously, they should be

assisted by some wholesome article of diet which is found to affect them, or by some harmless aperient medicine, which must be prescribed by a medical man, to suit the particular case; as the best aperient for one person may be the worst for another.

29. Provided that an aperient medicine is suited to the case, contains no drug injurious to the general health, and is not taken oftener than every second night, there is no harm in taking it at bed-time, whenever the bowels have not acted satisfactorily during the day.

#### REST AND CHANGE.

30. Active life is essential to the health of body and mind; but both require periods of rest, in addition to the regular hours of sleep. It is much better, therefore, to work vigorously for a time and then to rest, than to keep up a monotonous round of lifeless drudgery. (See Sleep; and Exercise).

31. The "current of the thoughts" is to the mind what posture is to the body; and both require change to prevent

weariness and deformity. (See Posture).

32. Rest of body or mind may be obtained either by abstaining from all bodily or mental exercise, or by change of occupation, and as the one gives entire rest and the other only partial rest, it is best to adopt each of these measures at different times.

33. It must be remembered, that, as the mind acts by means of the brain, which is a part of the body, it cannot act

MEALS. 37

healthfully while the body is suffering under disease or exhaustion. The "gait" of the mind is affected by the condition of the brain; as much as the "gait" of the body is affected by the condition of the limbs.

#### MEALS.

34. Counting from the time of beginning one meal to that of beginning the next, food should be taken at regular intervals of from four to five hours, except the interval between dinner and a very slight tea, which may be reduced to two or three hours. In weakly persons, and when the appetite will allow only a very small meal to be taken at one time, the intervals between all the meals may be reduced to from three to four hours. In illness, the interval must be ordered day by day by the medical man.

35. The chief meal of the day—the full-meal—by whatever name it is called, should be taken at whatever hour active occupation, both bodily and mental, can be suspended for about two hours; provided always that not less than two hours elapse between the conclusion of the full meal

and bedtime.

36. Breakfast should be the second best meal of the day, and should be taken leisurely immediately after rising in the morning. (See Exercise).

37. The other meals should be taken punctually at the fixed hours, but should be only light refreshments, and

small in bulk.

38. No food should be taken in the intervals between the regular meals.

39. As a general rule, pure water may be taken at any time, if indicated by thirst, so that the body is not heated by exercise, and the quantity drunk at once does not ex-

ceed a quarter of a pint. (See Exercise).

40. Spirituous liquors should not be taken the first thing in the morning or the last thing at night, without medical orders, they should not be taken when the stomach is empty, and they should not be drunk stronger than in the proportion of one ounce avoirdupois of absolute alcohol in about ten fluid ounces of liquid. One of the principal sources of mischief in the use of alcoholic liquors—is the practice of taking them to quench thirst in the place of unfermented drinks. The sense of thirst is a call from the organism for water, not for alcohol. Let the alcohol be taken as food, as medicine, or as a luxury, but not to quench thirst in the place of those unfermented liquors which are essential to health as diluents and solvents. (See Alcohol Table, and Remarks on Alcohol, Chapter I.)

41. Alcohol Fasts.—Those who habitually take alcohol daily, should abstain from it entirely for a few days twice

or thrice a year.

The importance of "PROPER HOURS FOR MEALS," is more fully explained in the following extracts from a letter by the Author, addressed to Employers and printed for private circulation in 1852, to advance the first step towards improving the national dining-hours of the commercial and working classes.

Gentlemen,—I address you as individuals who have a certain influence over the habits of a large portion of the community. Many past events testify so much anxiety on your part to increase the domestic comfort and social

happiness of your dependants, that I feel desirous to assist you in your praiseworthy object, by calling your attention to a most fruitful source of discomfort, pain, and even misery, peculiarly rife among the objects of your solicitude; I beg, therefore, to point out to you, that it is in your power either to ameliorate these evils, or to increase them, according as you frame some of your regulations. . . . .

That animal life cannot go on in even passable health and comfort without sufficient food and effectual digestion, is a fact perfectly familiar to all; yet, unhappily, among those who live an in-door life, effectual digestion is scarcely ever known. They constitute the bulk of that enormous number of persons who suffer from what is popularly known as "indigestion."

Now, there can be no question that they are pre-disposed to this class of diseases by many circumstances, quite inseparable from their occupations. But that makes it the more desirable to avoid all those causes of disease which are not necessary to their pursuits.

I must request you to remember that I am not speaking of any particular form of indigestion, but in a general and broad sense, of all those various maladies classed under the popular term.

The choice of the hours at which clerks, shopmen, mechanics, labourers, and other business-servants of both sexes take their meals, depends almost entirely upon the decision of their superiors.

If these hours are not well chosen, indigestion, in some form, is the certain consequence sooner or later. And you will see, from the few plain facts which I shall lay before you, that the notions which have long prevailed as to the proper hours for taking food, and consequently the regulations of nearly all business establishments, in this respect, are inconsistent with the conditions which physiology

teaches us to be essential to healthy digestion—with the requirements of the various occupations by which servants gain their livelihood, and with the conventional arrangements of society in the present day. . . . . . .

I shall hope, by avoiding all technicalities and minute physiological details, and by employing the plainest popular terms, to make my meaning perfectly intelligible to the unprofessional, to whom, in this instance, it is necessary to address my remarks; and I now ask your particular attention to the following statement of the conditions necessary to healthy digestion.

1. Food of appropriate quality and bulk.

2. (In the case of a full meal), rest of body and tranquility of mind for a short time previous to taking food, and for at least an hour afterwards. (In the case of a light meal), gentle exercise, and moderate mental activity may be allowed with impunity.

3. Effectual mastication, by which the alimentary matter is completely disintegrated and saturated with saliva, and

the meal taken slowly.

4. A sufficient interval between any two meals, to allow the first to have been digested and removed from the stomach, long enough for the digestive functions to regain their full vigour, before the second is introduced. The period which should intervene will depend upon the rapidity of digestion, which may vary according to the quantity eaten, the state of the health, the nature and amount of the previous exercise, the condition of the mind, and many other circumstances.

5. Caution that the stomach is not left empty long enough for the system to become exhausted, and the di-

gestive powers thereby weakened.

6. Strict adherence to the same hours for taking food, that the stomach may acquire a habit of preparing for its reception.

7. An interval between the last meal and bed time, sufficient to allow the work of digestion to be concluded before lying down to sleep.

Many other conditions might be mentioned, but as the few main points already stated are enough for our present

purpose, I shall confine myself to them.

The circumstances under which the majority of those persons are placed, whose health we are principally considering, must now be reviewed, in order to compare them with the conditions essential to healthy digestion, and to draw the necessary conclusions as to the proper hours for meals.

From the factory artisan up to the banker's or merchant's clerk, several important circumstances are common to all.

- 1. The day is devoted to labour either of the body or of the mind, or of both.
- 2. Business must be continued unremittingly up to the moment of leaving for meals, and resumed immediately on returning from them.
- 3. The time which can be spared for meals during business hours, must necessarily be very limited.

4. Some portion of this short time must be occupied in repairing to, and returning from the refreshment place.

In many instances, especially in retail trades, the nature of the business renders it impossible to keep to fixed times

for those meals taken during working hours.

The fact of taking time for meals out of the middle of the day, necessarily makes it later in the evening before the business can be finished, and this not only to the extent of the time during which work is actually stopped, for we all know how much the progress of business is interrupted by the simple acts of discontinuing and recommencing it, more especially when books and calculations are concerned. The effects of these combined circumstances may be viewed daily by visiting the public dining rooms attended by clerks, the lodgings of labourers at their dinner hour, and the offices and factories before and after meal-time—dinner-time more particularly, and dinner being the principal meal in the day—the full meal—it is of it that I shall chiefly speak.

Twelve, one and two o'clock seem to be the national dining hours for the working classes, and sixty minutes the maximum time allowed from business for this chief meal. The hour arriving, books or tools are hastily laid aside, and the dining place is reached by a sharp walk, which adds to the bodily fatigue of the labourer, and is not long enough or sufficiently leisurely to rest the brain of the accountant. The dinner must be despatched hurriedly, or there will not be time for the artizan to smoke his pipe, or for the clerk to glance at the Times, or perhaps the reading and eating are carried on at once: mastication must be carelessly performed, the mind is kept occupied, and the stomach rapidly loaded with food before it has had time to make ready for it; and, in some cases, a larger quantity is taken than the stomach has power to dispose of, simply because it is introduced too expeditiously for the system to become acquainted, as it were, that the supply of its wants is being effected. The meal finished, and the paper glanced at, or the pipe smoked, the sharp walk must be repeated, now with a full stomach, and business resumed before digestion has had time even to commence: and at this period, when the organic energies ought to be all concentrated about the stomach, they are at once summoned to the brain or to the muscular system. Consequently, the meal remains imperfectly digested, or not digested at all, lingers in the stomach beyond its proper time, and is finally expelled

in an unnatural condition, unfitted to undergo the important changes necessary before it can be appropriated for healthy nutrition; in all probability the next meal is introduced before the former one has entirely left the stomach, and thus is the mischief increased. Comparing the circumstances, here briefly enumerated, with those essential to healthy digestion, laid down before, the antagonism, so evident between them, plainly shows that "dyspepsia," or difficult digestion, is only the natural consequence of so persistent a disregard to the laws of health.

I am aware that different businesses do, from their nature, present various obstacles in the way of carrying out such sanitary measures as are most desirable, and that one set of rules will not always be practicable in two establishments. Therefore I should advise that the few principal conditions of healthy digestion before mentioned be carefully borne in mind; and that in each establishment the hours for meals be so selected, that these conditions and the calls of business may be arranged in unison with each other.

For a considerable period of time, during which my attention has been more particularly directed to this subject, I have made extensive inquiries on points connected with it, among employers and their servants, and have also had sufficient opportunities of enforcing the better regulation of meals upon both classes, with effects so decidedly beneficial, that I can with confidence impress the importance of the alterations I propose, and am convinced of their general practicability. After this experience and a mature consideration of the subject scientifically, I beg leave to call attention to those regulations, which appear to me to afford the nearest approach to the healthy standard that can be practicably

arranged to meet the necessary obligations of business establishments in general.\*

Let me assume that a substantial breakfast has been taken soon after rising, and a space of time allowed to elapse between the meal and the commencement of active business; (in those cases where the hour for business is very early, the breakfast should be divided, the fast being broken by a light fluid meal, and a second light meal taken some hours after). But to return to the better plan. Breakfast having been taken, and the day's employment commenced, the arrangements are now under the direction of the employer. From this moment until labour, whether mental or bodily, has ceased, no full-meal should be allowed. This is to be considered the FIRST GOLDEN RULE. The SECOND is not less important—that some light refection should be taken punctually every four or five hours. By adhering to the first rule, the system will never be charged with the task of active digestion at times when it is amply engaged in other functions; the digestive organs will not be injured by being called upon to undertake what they cannot properly perform; while the servant will not suffer from that oppressive languor and inertia, so unavoidable after a full meal, and will be so much the better fitted for his duties. By the second rule the system is maintained in a state of energy, the light refreshment, being easily digested during bodily or mental activity, supports the strength much more than a full but ill-digested and unassimilated meal can possibly do, and leaves the stomach itself unwearied, in a state of healthy vigour when the hour of relaxation from business arrives. Even in weak persons, a light meal will almost invariably have quitted the stomach in four or at most five

<sup>\*</sup> This was written eighteen years ago and subsequent experience has amply proved the value of the rules laid down.

hours, and in the strong considerably sooner than this; at intervals of four or five hours, therefore, according to the strength of the individual and the rapidity of his digestion, the stomach will be preparing for fresh work, and will call for it by the return of appetite, which must not on any account be disregarded. And in this place I must again impress upon you the serious importance to health, of not neglecting these occasional refreshments during business.

The day's labour over, the objections to taking a full meal are at an end, provided the second golden rule has been observed; and the amount of refreshment, necessary to repair the wear and tear of the day's exertions, may now be taken with decided advantage. I must here remind both employers and employed, that rest of mind and of body are necessary for a short time before taking a meal.\* To those who have been mentally occupied with sedentary business, a short leisurely walk will be a very proper prelude to dinner; but to those whose physical powers have been taxed during the day, there should be a short period of perfect rest before commencing the chief meal. It is a habit too common among commercial men to return home, impatient for their dinners, and to commence eating immediately, while in a fatigued condition. I have, in many instances, induced such persons to lie down for a quarter of an hour before beginning their meal; and have seen so much benefit arise from this practice, that I can confidently recommend it to all who are actively engaged during the day. When the dinner is finished, one hour at least should be devoted to mental and physical tranquility-some leisurely amusement being preferable to sleep. When the meal has not exceeded the bounds of moderation, a sufficient quantity of gastric fluid for the

<sup>\*</sup> See p. 40 for the conditions essential to healthy digestion.

digestion of the whole will generally be secreted within this period of rest; after which, the same quantity of blood and nervous energy being no longer required by the stomach, some more active employment of the mind or body may be indulged in, and all will go on well.

The full meal of the day, then—the dinner—should, under ordinary circumstances, be taken between the hours of five and seven p.m., which will allow time for it to be entirely disposed of before the hour of sleep arrives; and as the whole night passes without refreshment, a light refection such as our national "tea," is very desirable a few hours before bed-time. These arrangements will be found perfectly consistent with the essential conditions of healthy digestion; and, with a little contrivance and modification of hours in particular instances, are practicable in the majority of large establishments. And here let the friends of "early closing" observe, that by doing away with the injurious mid-day dinner, there will be so much saving of time in the best part of the day, and consequently a better opportunity of concluding business earlier in the evening.

It remains for me briefly to point out some of the inconsistencies of the system now most popular among dyspeptics and those employed in business. The dinner being taken at one or two o'clock must necessarily be a hurried one; it is impossible in the middle of business to allow time for rest, before or after the meal, to any serviceable extent: the evil consequences of this, in a large number of instances, have been already referred to. The "tea" being taken between five and seven o'clock, can only be a light meal, for the stomach does not require more so soon after dinner, therefore there is but one alternative, to leave the system without substantial nourishment from two o'clock p.m. until breakfast next day, or to take another

meal just before bed-time: of the two the latter would be the better, if the supper could be limited to a very moderate refection; but after waiting from an early dinner until nine or ten o'clock in the evening, a person in health has too good an appetite to be contented with this; therefore the stomach is loaded at a time when it cannot empty itself properly before the hour of sleep; the disadvantages of which are too familiar to need repeating here. Any one, therefore, who is anxious to regulate his diet according to the popular ideas of what is healthy, finds himself in this dilemma. He believes it to be an essential point that he should dine at an early hour. He knows that rest before meals, eating them slowly, and rest after them, are all necessary to proper digestion. He also knows perfectly well that, in the middle of the day, his business will not allow him either the rest or the leisurely dinner. Supper he is convinced is a most unwholesome meal, yet he knows from sore experience, that to fast from his early "tea" until next morning, leaves him too exhausted to sleep comfortably, to rise with vigour, or to enjoy breakfast. At length, if he reason at all, it becomes clear to him that to do what he thinks right in one respect, he must do wrong in another-a sufficient proof that there is error in the whole system.

#### CHAPTER III.

FOOD, HEAT, AND MOTION—A FOOT-POUND A FOOT-TON A BRITISH UNIT OF HEAT—MECHANICAL EQUIVALENT OF A BRITISH UNIT OF HEAT—COMBUSTION OF CARBON, CONSUMPTION OF OXYGEN, EVOLUTION OF HEAT—MODE IN WHICH THE HEAT EVOLVED FROM FOOD IS DISPOSED OF—COMPOSITION OF SOME OF THE PRINCIPAL ARTICLES OF FOOD AND THEIR HEAT EQUIVALENTS.

## \* \* See Essentials of a Normal Diet.

The relation between food, heat, and mechanical force has been the subject of some of the most important investigations of late years, and the light which has been thrown upon it is one of the greatest steps in the progress of medical science. But so far as medical art and practice are concerned, very little application has yet been made of this advance in scientific knowledge. This is only what might be expected, considering the occult nature of the subject, and the scarcity of time among the great body of medical practitioners to master and familiarise themselves with the details of such departments of medical progress.

Nevertheless, it is high time that such facts as are at present known, and such applications of them as are at present possible, should be added to the common stock of knowledge possessed by practical medical men; for it is only in this way that the public can derive benefit from our advances in science.

For the purpose, therefore, of enabling practitioners easily to familiarise themselves with these matters, the following concise statement has been prepared:—

1. A Foot-pound is the amount of mechanical force re-

quired to raise a pound weight one foot.

2. A British Unit of Heat is the amount of heat required to raise the temperature of one pound of water one degree Fahrenheit.

- 3. This amount of heat (a British unit) may be converted into mechanical force sufficient to raise a pound weight 772 feet.
- 4. This amount of heat (a British unit) is generated by a pound weight falling through a space of 772 feet. Hence 772 foot-pounds is called the Mechanical Equivalent of a British unit of heat.
- 5. One ton weight is 2,240 lbs.;  $\frac{772}{2240} = 0.34465$ ; therefore, the heat units multiplied by 0.34465, will always give the mechanical equivalent in Foot-tons.

6. The heat of combustion of carbon and hydrogen fully

oxidised (to CO2 and H2O) is as follows:-

1 lb.  ${C \text{ evolves } 14,500 }$  British Units of Heat.

- 7. The total heat of combustion of a compound containing H and C is the sum of the quantities of heat which the C and H contained in it would produce separately by their combustion.
- 8. It has been found that the presence of Hydrogen with Oxygen in the proportion to form water does not affect the total heat of combustion. It is only the excess of Hydrogen that can be made serviceable as a source of available heat.
- 9. The mean consumption of oxygen by an adult man of average stature (weight 150 lbs.) taking ordinary exercise is about 30 ozs. avoirdupois in 24 hours, and the heat evolved by each 1 oz. of oxygen in combining with carbon, hydrogen, etc., is about 340 British units. Hence, 10,000 British units of heat will be evolved every 24 hours by the

combination of 30 ozs. of oxygen with carbon, hydrogen, etc.; therefore, the food of an adult man, under ordinary circumstances, should be such as may, in addition to other purposes, evolve at least 10,000 British units of heat.

10. Practical experience in the dieting of large numbers of men, and other means, have enabled us to establish the fact, that such an average man, as above described, requires, for the maintenance of health, a diet which shall contain about 4 ozs. of plastic materials, 3 ozs. of fat, and 10 ozs. of Carbo-hydrates; and on careful analysis of this diet, we find that it can supply the required 10,000 British units of heat—viz., 2,516 from the plastic, 3,357 from the fat, and 4150 from the carbo-hydrates; total, 10,023.

11. The mode in which these 10,000 British units of heat are disposed of, and the purposes which they serve, may be seen in the following calculation which has been made as nearly correct as possible. 8,000 British units are required as sensible heat,—to raise the temperature of the inspired air to the temperature of the body, to vaporise the pulmonary halitus, and to maintain animal heat. The mechanical equivalent of 2,000 British units (equal to 690 foot-tons) is expended in actual work, more than half of which is employed in internal vital work (the mechanical work of the heart alone is equivalent to 200 foot-tons; respiratory and other vital movements may be estimated as equivalent to nearly 200 foot-tons more), leaving about 290 foot-tons available for external work, which may be represented by the labour of walking 16 miles; but of course only so much is available for actual walking, as is not used in the other external movements of the body, which we daily perform.

In the following table the conditions laid down in No. 6, 7, 8, of the foregoing paragraphs have been observed in calculating the British units of heat given in the last column.

The composition of the articles of food in this table will not be found to agree exactly with any one published analysis. 186 separate analyses by eminent chemists have been examined and compared for the purpose of fixing, as nearly as practicable, the approximate mean composition of each article.

-	THE RESERVE OF THE PERSON NAMED IN	CONTRACTOR OF THE PARTY OF	Name and		OF REAL PROPERTY.	-	-	_				and the same	
ARTICLES OF FOOD.	Ozs. Avoirdu-	Plastic,	Fat.	Saccharine,	Acid, Gelatine,	Complement,	Mineral.	Water,	N.	Н.	0.	c.	British Units of Heat,
Apple	1.000	.002		.136	.001	.022		.839		.000	072	058	53
Bacon (dried)	1.000	.084	625	1			1		11				754
Bread	1.000	.100	.007	453		.010							259
Butter	1,000	.010	·86o						'002	1			
Cheese	1.000	.308	.256	.024					.049				
Egg	1,000	.120	.108				1000	717				1	218
Fig (dried)	1,000	.020	.009	.570		.150	.034	187	008	1 3	1		
Fish (fresh)	1,000	.166	.008						.026	1000		1	
Flour (Wheat)	1,000	142	,010	.698		.013	'012	125	.023	.056	377	'394	392
Indian Corn Meal	1.000	.081	.045	.700		1	I The same of	1	.013		100	100000000000000000000000000000000000000	A STATE OF THE PARTY OF
Meat (Butcher's, cooked)	1,000	.225	.089						.036		1000		
Milk (new)	1,000	.045	.031	.048					.007		1		81
Oatmeal	1.000	.120	.058	.532		.096			.024		1000		
Pea (dry)	1,000	.219	.012	•469					.035				354
Pork (fresh)	1.000	175	.160				Acres de la constante de la co	1	.028	1		22.2	
Potatoes	1,000	.012		.230									
Rice	1.000	.021	004	.817		.033	005	.090	.008	057	414	393	376
Suet									.003				
	1.000							.100		059	460	381	350
Vegetables (fresh)	1,000	'01	002	078		.008	007	895	'002	006	043	039	38
(III. 36 1 1 1 1 1 I	- 11		-					ا					

The Mechanical Equivalent of the British Units of Heat may be calculated according to the directions given at p. 49.

## CHAPTER IV.

ESSENTIALS OF A NORMAL DIET—MODE OF CONSTRUCTING DIETS

—EXPENSIVE DIETS AND CHEAP DIETS FOR MAINTAINING
HEALTH—TABLES AND ANALYSES—EFFECTS OF NEGLECTING
EVEN ONE OF THE ESSENTIALS OF A NORMAL DIET.

# ESSENTIALS OF A NORMAL DIET.

(See Food, Heat, Motion.)

A healthy adult man of average stature, taking moderate exercise, will require and can consume, daily, from 32 to 40 ounces avoirdupois of dry nutritious food, which should have the following characters:—

1. About  $\frac{1}{200}$  must be mineral matter.

2. From  $\frac{2}{5}$  to  $\frac{1}{2}$  may be water, leaving  $\frac{3}{5}$  or not less than  $\frac{1}{2}$  or from 15 to 20 ozs. of anhydrous solid alimentary material.

3. Three or four ounces of plastic matter must be combined with three or four times that quantity of heat-giving material.

4. The heat-giving constituents must contain a mixture of fats (hydro-carbons) with saccharine materials (carbo-hydrates), in the proportion of about 1 of the former to 3 of the latter.

5. These heat-giving constituents should supply from 6 to 10 ounces of carbon, the exact amount required varying with season, exercise, etc.

6. The Articles of Food must be sufficiently varied to meet the requirements of the taste and of the appetite, and their Mechanical and other Conditions must be suited to the digestive powers of the stomach.

In addition to these characters, every complete diet must contain some potash-vegetable or fruit; and the total amount of water taken in 24 hours, including that contained in the dry food, must not be less than 70 ozs. avoirdupois.

# DIET TABLES FOR THE MAINTENANCE OF HEALTH IN ADULTS LIVING IN THE CLIMATE OF THE UNITED KINGDOM.

Each diet table contains all the essential elements of nutrition, in forms, quantities, and proportions necessary to the maintenance of health. The tables have been made complete without alcohol, leaving this to be ordered or not, according to circumstances. But, if spirituous liquors are added to any diet table, the quantity of carbon which they contain must be borne in mind. (See Alcohol Table).

It will be observed that the totals of the corresponding columns of the analyses are nearly the same in all the diet tables, showing the important fact that all the essentials of a normal diet may be equally secured in a diet that is simple and cheap and in one that is complicated and expensive.

In all diets Salt must be used and in those which do not include Potatoes some other Potash-vegetable or Fruit or Cresses or Lime or Lemon juice must be taken.

No. I.

Liquid.	Dry.	T-16-	1			Water	Plastic	Fat.	Saccha- rine.	Carbon
fluid ozs.	ozs.	Food for	24 not	ırs.		ozs.	ozs.	ozs.	ozs.	ozs.
TOTAL	HATO	Meat, Poul	ltry, o	r Ga	me					
Total	6	(cooked)				3'97	1.32	*53	****	.42
	4	Fish .				3.10	.66	.03		'02
	10	Bread				4.50	1.00	*07	4'53	1.03
	8	Potatoes				5.81	'14	••••	1.84	.76
	2	Rice .				.18	,IO	.01	1.63	.73
	21/2	Sugar				****	••••	* ****	2.20	1.02
	21/2	Butter				••••	****	2.20		1.85
5		Milk .				4.34	*25	.12	*21	(50.00)
		Coffee				15.77	****		*22	1 2 3 3 3
16		Tea .				15.95	****		'05	
17		Water				17'				
54	35					70.41	3.20	3.31	10.08	7.1
711	26.00		TIT		I	n Plast	ic matte	r .		1.8
							То	tal		8.9

No. 2.

Liquid.	Dry.	W. or The Land	Water.	Plastic	Fat.	Saccha- rine.	Carbon.
fluid ozs.	ozs.	Food for 24 hours.	ozs.	ozs.	ozs.	ozs.	ozs.
5 16 16	8 12 12 2 2	Meat, Poultry, or Game (cooked) Bread Potatoes Butter Sugar Milk Chocolate Tea Water	5°29 5°04 8°71  4°34 15°20 15°95 16°00	1.80 1.20 .20  .25 .07	71 ·08  2·00  ·18 ·32 	2.76 2.76 2.00 21 38 05	·56 2·32 1·14 1·48 ·84 ·23 ·42 ·02
53	36		70.23	3.25	3.59	10.84	
7			In Pl	astic ma	tter		8.01

No. 3.

Liquid.	Dry.	Food for	r 24 h	ours.	Water.	Plastic.	Fat.	Saccha- rine.	Carbon.
ozs.	ozs.				ozs.	ozs.	ozs.	ozs.	ozs.
5 20 20 20	18 3½ 3 1½	Bread Cheese Bacon Sugar Milk Chocolate Tea. Water			 7.56 1.28 .76  4.33 19.00 19.93 20.00	1.80 1.07 .25  .25 .10	.12 .89 1.88  .17 .40	8·15 ·08  1·50 ·21 ·50 .07	3'47 '70 1'48 '63 '22 '50 '04
65	26				72.86	3.47	3.46	10.21	7.04
					In Pla	stic mat	ter		1.87
						1	Cotal		8.91

No. 4.

Liquid.	Dry.	Food fo	ог 24	hours.	Water.	Plastic.	Fat.	Saccha- rine.	Carbon.
ozs.	ozs.				ozs.	ozs.	ozs.	ozs.	ozs.
10 20 25	8 16 8 2 <sup>1</sup> / <sub>2</sub> 1 <sup>1</sup> / <sub>2</sub>	Fish . Bread Potatoes Butter Milk . Cocoa Sugar Water			 6·38 6·72 5·81  6·93 19·10  25·00	1.33 1.60 .14  .40 .10	°06 'II 2°50 '28 '60	 7'25 1'84  '34 '20 1'50	*05 3*09 *76 1*85 *36 *56 *63
55	36				69.94	3.57	3.22	11.13	7:30
		1777000		-70	In Plas	stic matt	ter		1.93
		The state of				Т	Total		9.23

No. 5.

Liquid.	Dry.	Food fo	or 24 h	ours.	Water.	Plastic.	Fat.	Saccha- rine.	Carbon.
fluid ozs.	ozs.	1004			ozs.	ozs.	ozs.	ozs.	ozs.
	16	Bread.			6.72	1.60	·11	7.25	3.09
307	3	Peas .			.41	.65	.04	1.40	.65
	4	Bacon			1.14	.33	2.20	****	1.08
	. 2	Cheese			'73	.61	.21	.02	.40
10		Milk .			6.94	'40	.28	'34	.36
20		Coffee			19.71			.28	.13
	I	Sugar						I.00	.42
35		Water			35.00				
63	26				70.65	3.29	3:41	10.35	7.03
					In Pla	stic ma	tter		1.04
							Total		8.97

No. 6.

Liquid.	Dry.	Food for 24 hours.	Water.	Plastic.	Fat.	Saccha- rine.	Carbon.
ozs. 40	0zs.  4 3 2½ 1 9	Milk	ozs. 34.68 . 36 . 2.15 3.78 . 30.00	2:00 :20 :45  :90	1'40 '02 '32  1'00 '06	1.68 3.26  2.50  4.08	1.80 1.46 .26 1.05 .74 1.74
70	191		70.97	3.22	2.80	11.2	7.05
100			In Pla	astic ma			8.97

No. 7.

Liquid.	Dry.	Food for 24 hours.	Water.	Plastic.	Fat.	Saccha- rine.	Carbon
ozs.	ozs.		ozs.	ozs.	ozs.	ozs.	ozs.
20 20 25	20 3 1½	Eggs (two) Butter Milk		2.00 .44  1.00 .10	'14 '32 1'50 '70 '60	9.06  .84 .20 .75	3.86 .26 1.11 .90 .56
65	251/4		70.02	3.24	3.26	10.85	7.00
			In Plas	stic matt	ter		1.01
				П	otal		8.91

No. 8.

Liquid.	Dry.	Food for 24 hours.	Water.	Plastic.	Fat.	Saccha- rine.	Carbon
ozs.	ozs.		ozs.	ozs.	ozs.	ozs.	ozs.
40 30	8 7½ 2½ 1½ 1½	Rice	72 5'40  34'68 30'00	'40 I'II  2'00	'03 '80  1'25 1'40	6·54  2·50  1·68	2.93 .65 1.05 .92 1.80
70	19	berliner jamil	70.80	3.21	3.48	10.42	7:35
		to the policy opins	In Plas	stic mat	ter		1.89
				7	Total		9.24

No. 9.

Liquid.	Dry.	Food for	r 24 h	ours.	Water.	Plastic.	Fat.	Saccha- rine.	Carbon
ozs.	ozs.				ozs.	ozs.	ozs.	ozs.	ozs.
27 42	9 1 3 3 6	Flour. Suet. Sugar Eggs (four Milk. Water	r)		 1.13  4.32 23.41 42.00	1.28  .88 1.35	.09 1'75  '64 '94	6.58 3.00 	2.82 1.38 1.26 .52 1.21
69	193				70.86	3.21	3.42	10.41	7.19
				1	In Pla	stic ma	tter		1.89
							Total		9.08

No. 10.

Liquid. fluid ozs	Dry.	Food fo	or 24 l	hours.	Water.	Plastic.	Fat.	Saccha- rine. ozs.	Carbon.
22	16 13/4 3/4	Oatmeal Milk . Butter Sugar Water		: : : :	 2°14 19°07  49°00	2.40 1.10	 1.75 	8·51 ·90  ·75	4.46 .99 1.30 .32
71	181	1			70.31	3.20	3*45	10.19	7.07
1007		Dan T		100	In Pla	astic ma	tter Total		8.96

No. II.

Liquid. fluid ozs.	Dry.	Food in 24 ho	ours.	Water.	Plastic.	Fat.	Saccha- rine. ozs.	Carbon.
60	25 3 <sup>1</sup> / <sub>2</sub>	Bread . Cheese . Butter . Water .	: :	10.20	2.20 1.00	.17 .83 2.00	.08	4·82 ·65 1·48
60	301			71.68	3.20	3.00	11.41	6.95
				In Plas	stic mat	ter		1.89
					. 0	[otal		8.84

In the following letter published during the prevalence of Typhus fever in the Metropolis, I called attention to the mischief which may accrue from neglecting even one of the essentials of a Normal diet.

# BREAD DIET AND TYPHUS FEVER.

TO THE EDITOR OF "THE DAILY TELEGRAPH."

SIR,—The folly of the man who built his house upon the sands is familiar to all the world; but there is a parable of a folly even greater than his. A people, knowing that a tempest was approaching, left out the mortar and the girders from the houses that they were building, and when the tempest came, and the winds blew upon the houses, they crumbled to pieces before the storm.

It will hardly be believed that such an act as this is being performed at the present hour in the most enlightened

city of Christendom, under the eyes of scientific and learned men, in the purest innocence and good faith, and in the name of charity. Yet such is the plain truth.

Typhus fever has made its appearance in the houses of the half-starved poor of this metropolis. It is a contagious disease, but requires a particular soil to insure its fructification. Hence, out of any number of persons exposed to its poison, those suffer from the disease whose bodies present the most appropriate soil, and it proves especially fatal to those who are most strongly predisposed to its attack. It is well known to pathologists that in those who die of typhus all the tissues of the body, and the blood itself, are found to have lost their plasticity. Loss of plasticity in the body, and general want of vital force, constitute the characteristics of the soil in which the seed of typhus is sure to fructify. It would seem, then, but the most obvious common sense, that wherever the seed of typhus is known to be lurking every possible effort should be made to secure that it shall find no appropriate soil—that no human bodies deficient in plasticity shall be at hand. Yet what is the real state of the case? Typhus fever follows in the track of want. It fixes on some household where the effects of want are rife-i. e. where the inmates are reduced to the last extreme of poverty, and as a consequence are depending upon parish relief. Such a household is almost sure to be closely associated with others of the same class-depending upon parish relief. This parish relief means bread.

The seed of typhus, then, is scattered amongst a community subsisting upon bread. Now the houses in the parable, built up without mortar or girders, are the analogues of the human bodies built up out of bread. They have lost plasticity, and when the seed of typhus falls upon them, it takes root and fructifies, and they crumble

to pieces before the storm of sickness. It is hardly possible to maintain the body in health upon bread alone. A long study of the composition of food, in relation to health, warrants me in making this assertion; and I feel bound, in the cause of humanity, to call attention to this fact at the present moment, when the winter is at hand, and the poor of this city are threatened with such a scourge as typhus fever.

No food can maintain health in an adult, unless the average allowance for twenty-four hours yields to analysis from 3 to 4 oz. of plastic material, from 2 to 3 oz. of fat, and from 8 to 10 oz. of saccharine material. (See Essentials of a Normal Diet). Compare this with the composition of bread. At least 32 oz. of bread must be not only eaten but assimilated in order to obtain 3 oz. of plastic material, the smallest amount which can be accepted as necessary to supply the daily waste of tissue. This assimilation cannot take place without a certain proportion of fat, which fat is not supplied in the necessary quantity by the bread, and no other source of supply is provided by the parish relief. It thus becomes evident that all those poor creatures who are entirely dependent upon bread for their existence are specially prepared for the reception of typhus, and not less specially prepared for its fatal issue.

Setting every other consideration aside, then, and taking simply the ground of economy, it is a frightful extravagance of the funds of a parish, not less than of its life, to issue rations of bread only to the inhabitants of districts where typhus fever is rife. Immediately a patient is stricken with typhus the medical officer must call for wine, brandy, beef-tea, and the like, to the consternation of the board of guardians; and after all this there will probably come at last the expense of burial. So that every case of typhus

been made by supplying the extra food necessary to prevent it. If then the spread of typhus is to be checked, instead of encouraged—if it is to be rendered less fatal when it occurs, instead of more fatal—rations containing the proper elements of nutrition in their proper proportions and quantities must be secured to every member of a family in which one individual is affected with typhus, and also to all those persons who are associated with the infected household.

There is considerable difficulty in selecting diets which combine cheapness and simplicity with the essential chemical composition, and a form which the stomach will tolerate. But the two following will be found to fulfil all these conditions, so far as it is possible to do so without the introduction of meat; and in one of them bread is given in the largest quantity consistent with health :-1. Allowance for twenty-four hours: Bread, 25 oz.; cheese, 2 oz.; butter, suet, or dripping, 2 oz. These yield to analysis: Plastic material, 3.1 oz.; fat, 2.3 oz.; saccharine material, 11.4 oz.-2. Allowance for twenty-four hours: Oatmeal, 16 oz.; milk, ½ pint; butter, suet, or dripping, 1 oz. These yield to analysis: Plastic material, 3.0 oz.; fat, 2.3 oz.; saccharine, 9.0 oz. To each of these diets must be added limejuice or some land vegetable, salt, and a free supply of pure water. \*-I am, Sir, yours, &c., M.D.

M.D

London, Dec. 15.

<sup>\*</sup> The attention of those who visit the poor is particularly requested to these Diets.

# CHAPTER V.

FERMENTED LIQUORS.—ACID AND SUGAR IN SPIRITUOUS LIQ-UORS—ABSOLUTE ALCOHOL AND PROOF SPIRIT—ALCOHOL TABLE SHOWING THE ANALYSIS OF SPIRITUOUS LIQUORS— EFFECTS OF ALCOHOL IN DISEASE.

# ACID AND SUGAR IN SPIRITUOUS LIQUORS.

It is important to remember that although a fluid may contain much acid and therefore be unfit for some stomachs, it will not generate acid if it is free from saccharine matter, whereas a fluid containing saccharine matter will generate acid although it may not contain any.

### ACID.

"Proceeding from the least acid wine to the most acid we have Sherry, Port, Champagne, Claret, Madeira, Burgundy, Rhine wine, Moselle. The least acid fluids examined were Geneva and Whisky; then Rum, Brandy, Ale, Porter, Stout; the wines were all more acid than the malt-liquors."

"The nature of the acid was not absolutely determined, but a volatile acid distils over from wine, which is not acetic acid; and the action of polarised light shows that tartaric acid is seldom present, hence the fixed acid is most probably racemic, and perhaps malic acid."

#### SUGAR.

"I found no Sherry, Port, Madeira, or Champagne that did not contain more or less uncrystallisable sugar; (two samples of Sherry excepted, which were free from sugar.) I met with no Claret, Burgundy, Rhine, or Moselle wine, (excepting only one sample of Sauterne,) which was not free from every kind of sugar. Usually spirits contain no sugar; but one specimen of genuine French brandy had some cane-sugar, added to it. All kinds of Ale, Porter, and Stout contain much glucose [grape-sugar]. Hard cider, I found also to be perfectly free from sugar. Sweet cider contained uncrystallisable sugar."

"The fluids examined may be arranged in the following order, commencing with those which contain no sugar,

and ending with the most saccharine:

"Geneva, Rum, Whiskey, Claret, Burgundy, Rhine, Moselle. These have no sugar. Brandy, Sherry, Madeira, Champagne, Port, Cider, Porter, Stout, Malmsey, Ale, Tokay, Samos, Paxarete, Cyprus." (From the Appendix to "The Chemistry of Wine," by G. J. Mülder. Edited by H. Bence Jones, M.D., F.R.S., pp. 381-3.)\*

It will be seen from the following Alcohol Table that 2.26 fluid ounces of Proof spirit—contain 1 ounce avoirdupois of Absolute alcohol and that the other articles in the table contain less and less alcohol as the list descends until it finishes with London Porter, of which, 35.6 fluid ounces contain only 1 ounce avoirdupois of absolute alcohol.

For directions as to the quantity of alcohol to be taken during health and the extent to which it ought to be di-

luted see p. 9.

<sup>\*</sup>On the subject of Wines the reader is advised to refer to Mr. J. Denman's little book "What to drink," published by Longmans.

### ALCOHOL TABLE.

The weight of Absolute Alcohol (spec. grav. '796 at 68° Fht), and of solid dissolved, in measured quantities of Spirituous Liquors. \*\* This column shows the Quantity per One Imperial Pint quantity by 1 oz. contains. cent. measure of avoir. of SPIRITUOUS LIQUORS. each Spiritu- Carbon ous Liquor is con-(For ACID and SUGAR, see p. 25). which con-Alcohol. Extract. Carbon. tained Alcohol. Extract tains 1 OZ. in fluid AVOIRD. of OZS. OZS. ozs. ozs. Meaabsolute Alavoird. avoird. avoird. Weight. sure. cohol. Fld. oz. Proof Spirit 2.26 8.8 56.0 4'3 4.6 ARDENT SPIRITS. Whisky 2.6 5.0 7.6 •1 48.3 .6 4'0 Brandy 2.7 5.0 7.4 .2 4.0 47'I 1'2 Rum 2.8 7°I 5'3 .2 3.8 45'0 I.I Arrack 5.8 6.6 3.0 \* I 3.4 .4 41.4 Gin 3.5 6.3 5.2 .0 3.6 39.8 4.5 Roussillon 6.4 8.6 3.I 1.7 2.3 10.6 8.0 Sherry. 6.6 10'2 3.0 I.O 2'0 19.2 4.6 Cape Maderia 6.8 II.O 2'9 1.8 18.6 .7 3.2 South African Port 6.8 10.0 2.0 I.I 18.6 2.0 5.6 Port 6.9 10'2 2.0 I.I 2.0 18.5 5.5 Bucellas . 7.3 11.5 2.7 .0 1.8 17.2 4.4 Marsala . 7.5 11.7 2.6 .0 1.7 16.0 4.4 East India Madeia WINES 7.6 11.3 2.6 1.7 1.0 16.2 4.8 Frontignac . 7.8 9.0 2.2 3.2 2.6 14'0 19.1 Champagne . 12.6 11.3 1.0 2.3 1.8 10.4 11.0 Hock . 13.4 20.4 1.2 .5 I.O 9.5 2.5 Hungarian Red Voslau 14'1 21.5 1.4 .5 .0 8.0 2.5 Burgundy 15'2 23.8 1.3 .8 .4 8.3 1.8 Moselle . 15'2 23'3 1.3 .8 .4 8.3 2.0 Claret . 16.3 23.4 1.5 .5 .8 8.0 2.5 Sauterne . 10.0 27.2 I.O .5 .7 6.6 23 Hungarian White Ne- ) 10.0 28.8 szmely. . I'O .4 .7 6.6 1.8 CIDER . . . . 64.0 40.0 .5 .8 .6 3.0 3.7 Ale, Burton, Bass, 84s. 12.5 0.0 1.6 3.4 2'2 IO.I 15.7 60s. 14.2 13.0 I'4 2.0 1.2 8.0 9.6 Pale 2.2 60s. 10.0 17.5 I.0 1.2 I.I 6.6 India (Gardner, 7.0 MALT LIQUORS. 28.0 23.0 .0 .6 X 54s.) .7 5.5 3.0 Scotch? 10.0 Bottled 13.1 (Edin.) I.O 2.4 6.6 1.2 11.4 Pale 25'0 20'5 .8 I'4 6.6 I.O "Eightpenny" 5.0 22.7 22.2 .0 I.I .0 5.7 Family 1s gallon 5.4 24'9 22'0 .8 I'2 .0 "Fourpenny". 5'4 5.7 25.4 23.0 .8 I.I .8 5.2 Stout, Dublin, (bottled). 5.3 20.8 16.7 .0 1.7 I'2 6.I 8.2 London . 21.5 18.0 1.2 .0 I.I 5'9 6.9 Porter, London 35.6 26.6 .6

3.7

5.6

.8

1.3

<sup>20</sup> Fluid oz. = 1 Imperial pint (measure). 16 oz. avoird. = 1 Pound (weight).

### ON ALCOHOL IN DISEASE.

(Leading Article from the Lancet. Jan. 1st., 1870, referred to at p. 9.)

"After a great many fluctuations, professional opinion upon the question of alcoholic stimulation in disease appears to have recently achieved some real progress towards the establishment of satisfactory principles. There are still very wide divergencies between different authorities, and we fear that there is still a good deal of extreme and irrational practice, both in the direction of excessive stimulation, and of the opposite fault of an unreasonable fear of the remedy. But it seems to us that the outlines of a greatly improved knowledge, both of the therapeutic powers, and the capacity for mischief which alcohol

possesses, may now be discerned.

In the first place, as regards acute diseases attended with febrile phenomena, three things are tolerably plain. The first is, that alcohol, when it acts well, acts as an antiphlogistic stimulant; that is to say, it lowers abnormally high temperature, it reduces the frequency of the pulse, and, while raising nervous power, it calms those disturbances of the nervous system which attend, if they are not caused by, the elevated temperature of the blood in pyrexia. The second great fact is, that there are the utmost differences between different pyrexial patients as to their capacity for receiving benefit in this way; that a large number of persons, especially among the young, and previously robust, do best without any alcohol; and that among those to whom it is beneficial, there are some for whom three ounces of wine per diem is fully the physiological and therapeutical equivalent of twenty-four ounces given to another and smaller class of patients; and that nothing but careful tentative use of the remedy can tell, in any particular case, whether it is needed, and, if so, in what quantities. In short, that all generalisations to the effect that you must, or must not, give wine by ounces, or by bottles, in fevers or in inflammations generally, are worthless and misleading. The third settled fact is this, that it is often in those cases where alcohol used (on purely theoretical grounds) to be thought most dangerous—viz., in cases with very high temperature and flushed face—that it produces its best effects; and that to pour large quantities of stimulants into a fever patient simply because he is pale, and has a small pulse, is an unwarrantable proceeding.

The above are certainties, and the following are probabilities.

It is probable that alcohol owes part of its influence in fevers to an antiseptic agency, by which it destroys the activity of certain bodies-call them organisms, or not, as you please-by means of which the contagium sets up the febrile disturbance within the blood. In the case of inflammation, it is probable that alcohol, when it acts well, does so in part because it stimulates the sympathetic and contracts the arterioles, and in part interferes with the migration of blood-corpuscles through the vascular walls, as Binz and his pupils have shown that quinine can also do. It is probable that so far as alcohol can be applied to these purposes within the organism, it is of unmixed benefit. And there is much reason to believe that the singular differences between different individuals, as to the quantity of alcohol they can bear, depend on some unexplained difference in the respective rapidity with which alcohol is oxidised in the blood in different persons. For it is now known with certainty, on the one hand, that nearly the whole, even of a poisonous

dose, is always oxidised in the body, and on the other hand, that the presence of large quantities of unchanged alcohol for any length of time in the blood, inevitably poisons the nervous system. The antiseptic action, and the influence on the migrative tendency of the corpuscles, are most likely produced immediately that the alcohol mingles with the blood; it probably depends on the subsequent progress of oxidation whether the general effect on the patient will be good or bad. One thing is certain: if signs of narcotism—i.e., paralysis of the nervous system—are produced, the alcohol is doing harm and must be immediately diminished or stopped.

As regards chronic diseases, we are sorry to observe that there is very much less of intelligent progress in medical opinion than in the case of acute diseases. It is much to be regretted that a large number, even of highlyeducated practitioners, will persist in acting on the assumption that in non-febrile diseases the amount of alcohol to be administered ought to be measured by the degree of debility, merely as such. The direct and very mischievous corollary of this is the practice, unfortunately daily increasing, of prescribing stimulants with lavish profusion in those numerous nervous affections to which weakly persons (more especially women) are prone. It is our duty, as medical journalists, to raise our voice to the utmost against this tendency. We are no bigots against alcohol; and we are heartily sick of the unthinking abuse which has been lavished on what it is the fashion to call "indiscriminate stimulation in acute disease." We declare our belief that the real mischief lies at the door of those who are indiscriminate (because they are unthinking and illogical) in their prescription of alcohol for debility, merely as such. It is no figure of speech, but the literal truth, when we say that hundreds of neuralgic,

hysteric, and epileptic patients have been driven into drunkeness or lunacy, or both, by the careless folly of advisers, who had no better reason for the prescription of large doses of alcohol than the fact that these diseases are attended with nervous weakness, as they undoubtedly are. The assumption involved—that so much ingested alcohol is necessarily so much added nervous strength—is so gross a fallacy that no one would assent to it if expressed in plain words. Yet we constantly see it acted upon. We repeat with all the energy of which we are capable, that it is a grave scandal and mischief that medical men should endanger in this serious way the powers of moral resistence of women and other weak persons while basing their practice upon ideas that are illogical and untenable; and we trust that a reform in this respect will immediately be commenced."

### CHAPTER VI.

FAT ESSENTIAL TO HEALTHY NUTRITION—IMPORTANCE OF DISTINGUISHING BETWEEN SOLID AND LIQUID FAT—BANT-INGISM—GETTING FAT AND GETTING THIN—ANALYSIS OF BANTING-DIET, SHOWING ITS ERRORS AND ADVANTAGES.

Fat is so essential to the maintenance of healthy nutrition, that the quantity contained in the daily food cannot be reduced much below the proportions given in the "Essentials of a normal diet" and in the diet tables, without great risk of damaging the health.

When it is necessary for any special object to reduce the quantity of carbon taken in the aliments, this can more safely be done by diminishing the saccharine and amylaceous matters than the fats.

The importance of fat in nutrition should be studiously borne in mind by those who construct diets for the poor, for public institutions, or for the treatment of obesity, diabetes, fatty degeneration, dyspepsia, and the like.\*

# ON THE IMPORTANCE OF DISTINGUISHING BETWEEN SOLID AND LIQUID FATS.

Extract from an Article by the Author in the "CHEMICAL NEWS, Sept. 4th, 1868."

The peculiar isomeric modifications of which stearin and palmitin are susceptible, as shown by Duffy, pointedly distinguish them from olein, which, so far as at present

<sup>\*</sup> See Index "ON BREAD DIET, AND TYPHUS FEVER," also "ON DIET IN DISEASE,"

known, has not this susceptibility; a distinction which is supported by the different behaviour of oleic acid towards chlorine and bromine, from that of stearic or margaric acids (Lefort), and by the different action of bile upon stearic acid and upon oleic acid (Marcet).

But I think we ought to be prepared to learn that solid and liquid fats differ in some important physiological properties, by the first general fact concerning the constitution of all natural fats—viz., that they are mixtures in varying proportions of at least four different bodies, of which the melting points so widely differ—stearin melting at 144° F., palmitin at 114.8° F., margarin (probably a compound of stearin and palmitin) at 116° F., while olein remains fluid at 32° F.

That the different degrees of solidity of fats depend upon the proportions in which the solid ingredients are mixed with olein, that olein has a peculiar power of dissolving the solid ingredients, and that the melting point of the mixture is thereby reduced, appear to me to be facts pointing in the same direction as the foregoing, especially when we remember that the affinity of oleic acid for oxygen is much greater than that of the other fatty acids.

The fatty bodies obtained from warm-blooded animals are generally solid at ordinary temperatures, whilst those from fish and from cold-blooded animals are liquid. And when we consider the high melting points of the solid fats as compared with the temperature of the body in warm-blooded animals, it is evident that the fat in them would be solid at the temperature of their blood, but for the mixture of olein, by which the melting point is reduced. Therefore the solidity or fluidity of the fat in living animals is determined by the proportion of olein, which is able to be mixed with the stearin, palmitin, and margarin

in each individual; and we are forced to conclude, either that it is of no importance whether the fats of the body during life are in a solid or liquid state, or that it is important in what proportion the olein, stearin, etc., shall be combined.

It has been already proved by experiments on the fattening of cattle, that the solidity or fluidity of the fat in the body varies with the food—that cattle fattened upon linseed cake, for example, accumulate, in their adipose tissue, an oily material of unusual fluidity (Draper), and that the consistence of butter is dependent upon the kind of food given to the animals from which it is produced (Fownes).

The fat in animals is particularly liable to accumulate immediately beneath the cutis, in the omentum, and around the kidneys; and the fat found in the latter situation, where it is subject to a more uniformly elevated temperature than in the integument, is well-known to be of a more suety character—that is to say, it contains a smaller proportion of olein, and has a higher melting point. These familiar facts point again to some importance in the animal economy, attaching to the melting point of the fat and the consequent degree of fluidity in which it should exist during life.

With regard to the fat of the integument—the principal deposit of adipose tissue in the body—it appears to me self-evident that the fluidity of this fat must vary with the temperature of the atmosphere in which the animal is placed; to what extent this is the case, is, in my opinion, a most important subject for enquiry; and although the experiments to determine the question are yet deficient, I hope soon to be able to supply them.

In conclusion, what I now suggest as a general propositiod, is this:—That, in all probability, the stability of the

fats of the animal body in resisting too rapid oxidation is dependent upon the degree of solidity which they possess at the temperature of the living animal at any given time; that alterations in external temperature may affect the solidity of the adipose tissue of the integument, and, consequently, its power of resisting oxidation; and that, therefore, in all probability, it is of great importance that the food of an animal shall contain a certain proportion of material capable of supplying the adipose tissue with solid fat.

# ON GETTING FAT AND GETTING THIN.

On comparing the following analysis of Mr. Banting's diet for getting thin with my tables of normal diets it will be seen that it yields less than half the normal quantity of Carbon, leaving the deficiency to be obtained from the fat already stored up in the system, by the consumption of which the obesity is removed. The fault consists in this reduction of Carbon being obtained by diminishing the Hydrocarbons (fats) of the food instead of only cutting off the Carbo-hydrates (Sugar and Starch).

Meat, Poultry, Game, and fish . Bread	9°25	3°15	ozs.	, ozs.	ozs.
Game, and fish . Bread			1.25		•08
Vegetables, (green) Fruit Tea Wine (Claret)	1.40 2.25 16.00 13.00	*20 *02 		.00 .15 .40 	·98 ·39 ·07 ·15 ·02 ·70
	43.40	3:37	1.56	1.45	2'31
	Tea	Tea	Tea	Tea	Tea : : : : : : : : : : : : : : : : : : :

Extract from the Author's Letter "ON BANTINGISM," written in 1864.

It has happened to me to have to do with a great number of persons who have tried Bantingism, and I do not hesitate to say that, up to the present time, Mr. Banting has done a great deal more good than harm. He has not brought forward a single new fact or new idea but he has had the luck, by zealously advertising a striking case of the effects of a plan of treatment long familiar to every medical man who understood physiology, to convince the public of the immense influence on the animal organism of modifications in the quality of food—an influence, in the importance of which they did not half believe when urged upon them in the form of medical advice.

Mr. Banting candidly told his readers that he was ignorant of the physiology of food, but they did not care for this while determined to try his plan, and hence it is not to be wondered at that while many have been benefitted many have done themselves harm. But the harm that has been done at present is not very great and is nothing to compare with the harm that will come, if, frightened by some striking case in which Bantingism has nearly cost a man his life, the public rush away from it as blindly as they rushed into it, and establish for a time such a popular prejudice against systems of diet, that a medical man shall not be able to institute those rational and scientific regulations of diet which are often more important than any other means of treatment in the management of disease . . . . . The plain and simple facts of the case are these. 1. A certain amount of fat in the system is one of the most essential elements of health. quantity required by different individuals to maintain

health differs. 3. The effects of a deficiency of the quantity actually required in any given organism are most disastrous, the tissues of the body and the brain and nerves being at length disintegrated to supply the elements of fat which they contain. 4. When there is a quantity of fat in the body in excess of that necessary to health, it may be lessened with great and needed advantage, provided it be done slowly and cautiously,allowing the nutritive functions of the system to accommodate themselves gradually to the altered condition, -and most peremptorily stopping the loss of fat before it has amounted to a deficiency in the quantity necessary to healthy nutrition. Nothing is easier then to make a fat man thinner by altering his diet, but to do this safely and well is by no means devoid of difficulty, and in a large number of cases requires the greatest caution consideration and judgment on the part of those who rule the modifications in the diet of the patient. But there is no denying that, with such care and judgment, it can in most cases be done safely and advantageously.....

Further information regarding fat, its importance in the organism, its relation to Disease, and the means of accumulating fat, will be found in my works "on Tubercu-Losis" and "on the True First Stage of Consumption."

### CHAPTER VII.

On the Diet of Sickness—Principles upon which to arrange the Diet of Disease—Rules for carrying out these Principles—Examples of Diets for Consumption, and for Diabetes.

### ON SOME PRINCIPLES OF DIET IN DISEASE.

A lecture delivered by the Author at the Royal Hospital for Diseases of the Chest, April, 18 5,

(Illustrated by Diet Tables.)

The subject which I have chosen for to-day's lecture is one upon which it would be very easy to write a large book, but which is very difficult to treat in a short lecture. I must content myself, therefore, with condensing what I have to say into a few statements of opinion, avoiding all attempts at discussion, and if I appear to deliver my views dogmatically, you will, I hope, understand that it is not because I feel at all inclined to dogmatise, but because I am pressed by the extent of the subject, and the narrow limits of our time.

In my "Manual of Diet and Regimen"\* I have given "the Essentials of a Normal Diet" for health. The question now is, how ought these to be modified in disease? You will observe that these "Essentials of a Normal Diet" provide for the maintenance of healthy nutrition in a "healthy adult man of average stature taking moderate

<sup>\*</sup> Now embodied in this work.

exercise," and, for simplicity sake, it will be best to make all our remarks apply to this "adult man of average stature" under altered circumstances.

The first aud leading principle of diet in disease is, to provide for the maintenance of healthy nutrition, under the peculiar alterations of circumstances attendant upon disease. In other words, the diet of disease should be as nearly that of health as the altered condition of the nutritive functions, and the altered conditions of the patient's life will allow; the grand object being to keep up healthy nutrition of the whole organism.

The true appreciation of this first principle, in its various bearings, will save us from running into either of those extremes which at different times have disfigured medical practice. We shall not be led to starve our patients upon water-gruel, when they are craving for natural food; or to stuff them with beef-steaks and porter, when every instinct of their nature rebels against it. In fact, you may take it as a very safe rule, that it is better cautiously to supply a patient with the kind and quantity of food that his stomach calls for, than to deny it to him without an unquestionably good reason for so doing.

This may seem to you very much like letting a patient eat and drink what he pleases. But that is not at all what I intend to recommend. All I mean is this, and I wish particularly to impress it, that if we intend to interfere in the subject of diet, we must take care that we thoroughly understand what we are about; and in order to do this it is necessary to keep well up in the following subjects:—

1. The physiology of healthy nutrition.

2. The composition of food, and the essentials of a normal diet.

3. The physiology of disease.

If we keep these matters well before the mind, and at

the same time keep our wits about us in watching the case under treatment, it is surprising what an immense deal of good may be done by interfering with the diet; but not otherwise.

We start, then, with this as our first principle, never to be lost sight of—that healthy nutrition is to be maintained, if possible, under all circumstances. And we assume that to do this in a healthy adult man of average stature, taking moderate exercise, the essentials of a normal diet, must be supplied.

It must be borne in mind that the proportions and quantities of the different elements of this normal diet are arranged to meet the requirements of the different functions of the organism when in a healthy state of activity; and it therefore follows, that if the activity of any of these functions is altered, the requirements will be altered; and hence, the second general principle is this:—To alter the quantities and proportions of the elements of a normal diet to correspond with any alterations in the conditions of life. Thus, when a man is overtaken by sickness, and confined to his room or bed, the adult man taking moderate exercise becomes an adult man taking no exercise; and the ingredients of his diet which were proportioned to his moderate exercise must now be proportioned to his no exercise; and other alterations must be made in like manner to correspond with other altered circumstances, in addition to any that may be specially required by the nature of his disease.

But I must remind you that, even when a man is confined to his bed, and precluded from taking any kind of exercise, he is still necessarily undergoing a considerable amount of muscular exertion, which must be provided for in his diet. For example, so long as life remains, such all-important muscles as those of respiration, and the

heart itself, continue to act, and to require that their healthy nutrition shall be provided for by a supply of plastic materials in the food.\*

We come next to the long list of alterations of function which may be involved in the term "Sickness." And the third principle is-To alter the forms, quantities, and proportions of the elements of a normal diet, to meet the altered relations in the activity and condition of organs consequent upon disease. It is evident that, in order to carry out our first principle of maintaining healthy nutrition under all circumstances, it may be necessary, under some conditions, to reduce the quantity of every element of diet; and also, under some circumstances, to alter the proportions of the different elements. This we see demonstrated in some of the lower animals by the phenomena of hybernation. When an animal gives itself up to its winter sleep, every vital function is reduced to its lowest degree of activity; and the animal is able to maintain healthy nutrition for a long period without taking any food at all; but as respiration has to be kept up more actively than the rest of the functions, a special store of carbon for this purpose is laid up beforehand in the body.

Now, supposing a man to suffer from any state of disease which should place him in the position, as regards his functions, of an animal during hybernation, it is clear that, while his whole diet must be reduced to a very low scale, the respiratory (heat-giving) elements must be supplied in quantities out of the normal proportions as compared with the rest; because no supply of carbon is stored up in preparation for his illness, as it is in the hybernating animal in preparation for its sleep.

We see conditions, in many respects similar to these

<sup>\*</sup>See " Food, Heat, and Motion."

in some stages of fevers, in which absorption, nutrition, and every vital function is at its lowest point consistent with life, respiration being the only one sufficiently active to call for any considerable supply of food. But here, of course, we must not lose sight of an element in the case not present in hybernation-viz., the existence of a poison, which by some means, natural or artificial, has to be eliminated or destroyed, and which may be keeping some functions in activity, the requirements of which must be met. The precision with which we are able to do this in any given case, will depend upon the correctness of our knowledge of the nature of the poison, and of the organs concerned in the restorative process. Here, no doubt, we are often obliged to act in the dark, and to supply many ingredients which may not be needed, in the hope of furnishing among them that which is required, but which our ignorance prevents us from identifying. And we had far better, whenever our knowledge is at fault, act in this safe manner and supply much that may be useless rather than run the risk of witholding that which may be essential to life. But, in the majority of cases, our knowledge will be sufficient for the emergency, if we keep in mind the general principles of action.

The fourth principle is this:—To obtain rest for every organ while it is suffering under active disease, by removing from the diet such elements as increase its functions. These are conditions which it is not always easy to fulfil without deviating from our first principle. For example—in the case of diseased kidney—the healthy nutrition of this organ requires a supply of albuminoid materials, while its function is increased by any surplus of these materials in the organism; and when its function is interrupted by disease, a proportion of albuminoids in the diet, necessary to the healthy nutrition of the organism

generally, will be tantamount to an excess as regards the function of the kidney, and the accumulation of retained excretory matters will press injuriously upon the affected organ. In such a case other medical aids than diet must be brought to bear; and while the albuminoids in the food are reduced as low as is consistent with healthy nutrition, some auxiliary organs which are not damaged must be stimulated for the time, to save the diseased part from undue pressure upon its functions.

A simpler, but still important principle, may be stated as the fifth, viz .: - In all alterations of diet, to avoid any unnecessary reduction in the number and variety of the forms in which food is allowed to be taken. This becomes especially necessary to be borne in mind when dieting the dyspeptic, who are often still engaged in the active avocations of business and of society while under medical treatment. To treat such cases by cutting off from the daily bill of fare first one article and then another, till the food consists only of two or three permitted forms, is to destroy the appetite and the digestive powers by monotony of diet, and to depress the spirits of the patient by a constant series of petty denials. This plan of dieting can only be regarded as the resource of ignorance; because an enlightened view of the case will discover some particular defect in the function of digestion or assimilation which will at once indicate the form or element of the food which is to be avoided; and thus it will be only necessary to cut off those articles which specially represent this element, or simply to alter the forms in which they are presented to the stomach. (See p. 5.)

The sixth principle is also of great importance, viz.:— When it is necessary to remove from the food any of the essentials of a normal diet, to aim at selecting that which will answer the desired end with least danger to the nutri-

tion of the vital organs. For example, if it is necessary for any special purpose to diminish the heat-giving elements of the diet, it is safer to remove the carbo-hydrates than the hydro-carbons, because the latter not only supply carbon for the evolution of animal heat, but are essential to the nutrition of the nervous system, and of the albuminoid tissues generally.\*

The seventh and last principle which I shall give in this Lecture is of very general application:—When it is desired to *increase* the normal nutrition of a tissue or organ, we must not only supply it freely with the special materials requisite for its development, growth, and repair, but at the same time call upon it for the performance of its normal functions—over-fed idleness insures morbid nutrition, not healthy life.

In the next place I will give you a few rules which may assist you in carrying out these general principles—

Rule 1.—When the power of appropriating any essential ingredient of a normal diet is lost to the organism, the lost function must be substituted by some artificial process, or the ingredient in question must be withdrawn from the diet till the normal function is restored. In obedience to this rule you see me at this hospital administer pancreatic emulsions of fat to patients, who have lost the power of assimilating fat without this artificial assistance, while I adopt all practicable means of restoring the normal function.†

Rule 2.—Is inseparable from the first, and it is this:— No essential of a normal diet must be withdrawn without an attempt being made either to supply to the organism in some other way the ingredients of which it is deprived, or to suspend those functions which call for a supply of this ingredient. Thus, to take a simple illustration:—

<sup>\*</sup> See Fat. + See Diets for Consumption.

Suppose the power of digesting meat to be lost through a deficient secretion of gastric juice, meat must be withdrawn from the diet till the lost function is restored, or else an artificial digestive fluid must be introduced; or if it is impossible by these means to maintain the digestion of meat, the physiological ingredients of meat must be supplied in the form of some albuminoid solution; or, finally, if this cannot be done, then those functions which principally waste the albuminoid tissues of the body must be placed as far as possible in a state of rest, muscular action must be suspended until the function is restored.

Rule 3.—If an undue waste of any elements of normal nutrition is found to be going on in the organism, and the means remain of appropriating those elements from the food, they must be supplied in the food in quantities as much in excess of those proper to the normal diet of health as will be sufficient to supply the waste until it is stopped. This also may be illustrated by a very simple example. In Bright's disease of the kidney there is no loss of the power to appropriate the albuminoids from the food, whereas a constant loss of albumen is going on through the kidneys, which must be met by proportionate increase of the albuminoids in the diet. But in following this rule, in this particular case, it will be necessary to observe the precautions which I mentioned when speaking of the fourth general principle.

Rule 4.—When through any defect in the organism, the elements of a normal diet are lost to nutrition if presented in the usual forms, those forms must be changed; but care must be taken that in the altered form all the essential elements of a normal diet are supplied in their proper quantities and proportions. Nothing can illustrate this better than the use of milk as a substitute for solid or mixed foods in diarrhœa or sickness.

Rule 5.—Has to deal with more complicated difficulties. If such a defect exists in the organism that some of the essentials of a normal diet are misappropriated, so that the organism is deprived of one or more of the normal elements of nutrition, and at the same time a disease is constituted out of the misappropriated food, then we have a double duty in interfering with the diet. First, the source of the disease must be stopped by withdrawing that part of the diet out of which it is constituted; and, secondly, the elements of nutrition thus removed must be supplied by some other means or in some other form.

Thus, in Diabetes the saccharine and amylaceous elements of the diet are misappropriated; they do not serve their normal function of supplying carbon for the evolution of heat, and by passing off through the kidneys they constitute an exhausting disease. It is necessary, therefore, to stop the source of this disease by cutting off the saccharine and amylaceous ingredients of the diet till normal nutrition is restored. But, in the meantime, as carbon must be obtained by some means, it is taken from the fat stored up in the body so long as that lasts, and when it is gone, from the albuminoid tissues themselves, till the whole organism is disintegrated; unless at the same time that we cut off the starch and sugar, we increase the quantity of fat supplied in the food as much in excess of the proportion proper to a normal diet as shall fully supply the demand.\*

The modern dietetic treatment of Diabetes may be taken as a good example of the way in which increased knowledge of the nature of disease and of the physiology of food enables us to act under what I have called the fifth principle of diet, viz., to avoid any unnecessary reduction in the number and variety of the forms in which food can

<sup>\*</sup> See Diet for Diabetes.

be taken. In former days the poor parched diabetic was forbidden to drink water lest he should increase his flow of urine; now we are able to let him quench his thirst as much as he pleases, so that he takes nothing which contains starch or sugar; and again, by preparing his articles of food in such a manner as to exclude the injurious ingredients, and by selecting those which are known to contain them in the smallest quantities, or not to contain them at all, we are able to present the diabetic with a fairly tempting and varied diet, so that he is able to keep to it for months and years with comparatively little difficulty. (See p. 4.)

In conclusion I shall venture to give you a sixth rule, which I call the "golden rule," as it relates to the carrying out of the first of our principles throughout all the difficulties involved in the rest. The "golden rule" is this:—Never let a patient become weak through a defective diet if you can prevent it by any possible contrivance; and if a patient has unavoidably become weak, never let him remain so, through any defect of diet, one hour longer than it is absolutely impossible to prevent.

Other principles and rules I must reserve for future Lectures, when treating of special cases as they come before us. But I hope that you will find, when you come to put them to the test of practice, that the few which I have given you to-day will serve both your patients and yourselves many a good turn, if you will only bear in mind what I have endeavoured to impress; that in order to carry them out satisfactorily you must keep up your knowledge of your profession generally, and especially of the various subjects which I have enumerated, as essential to a proper understanding of the alterations of diet required in disease.

# DIETS FOR CONSUMPTION AND FOR DIABETES.

In the Diets for Consumption it is assumed that no fat is assimilated except that artifically pancreatised.

- In Table I. The required amount of carbon is supplied by an excess of carbo-hydrates.
- In Table II. The required amount of carbon is supplied by an excess of albuminoids.
- In Table III. The amount of carbon is kept low, because it is only intended as a temporary diet to be used during periods of rest in a warm room. The arrowroot and some of the fat of the milk are pancreatised by mixture with the "pancreatic emulsion."

In the Diet for Diabetes. Pancreatic Emulsion is added and Pepsine ordered to assist in the digestion of the large quantity of fat and of albuminoids, necessitated by the nature of the disease.

DIETS FOR CONSUMPTION. TABLE I. CARBO-HYDRATE.

Food for 24 hours.	Oz.	Plastic.	Fat.	Saccharine.	Total Carbon.	Carbon from	Carbon from Non-
Cooked Meat	4						- Grande Grande
Bread	0 01	1.350	.534		1.152	0.732	0.450
Potatoes	000	0.136	020	1.840	2.470	0.540	1.930
Sugar	2			008.1	0.848	7/00	0.848
Liebia's Food for Infants	CJ (	000.I	0.200	0.840	1.440	0.540	006.0
Faringceous Foods	27 0	0.300	911.0	1.064	0.720	0.162	0.558
Fermented Liquots*	0	0.300	0.050	4.800	2.350	091.0	2.190
Pancreatic Finalsion			:	:	000.I		000.I
· · · · · · · · · · · · · · · · · · ·	N		1.000		0.740	:	0.740
Totals	372	4.086	2.440	14.974	11.552	2.206	9.340
Deduct Carbon from Non-pan-	:	:	:		0.945	:	:
Total available Carbon	:	1	:	:	209.01	:	:

\* Either-Half a pint (Imperial) of Port, Sherry, or Marsala; Or, One Pint of Burgundy. Claret, or other similar Wine; Or, One pint of good Ale or Stout; Or, A quarter of a pint of Rum, Whiskey or Brandy, diluted with one pint of water,

DIETS FOR CONSUMPTION. - TABLE II. - ALBUMINOID.

With this diet Hydrochloric Acid and Pepsine should be given to assist in digesting the very large quantity of plastic matter.

\* Either—Half a pint (Imperial) of Port, Sherry, or Marsala; Or, One pint of Burgundy, Claret, or other similar Wine; Or, One pint of good Ale or Stout; Or, A Quarter of a pint of Rum, Whiskey, or Brandy, diluted with one pint of water,

DIETS FOR CONSUMPTION. - TABLE III. - FLUID DIET.

THE RESERVE THE PROPERTY OF THE PERSON NAMED IN COLUMN TWO IS NOT THE PERSON NAMED IN COLUMN TWO IS NAMED IN COLUMN TWO IS NOT THE PERSON NAMED IN COLUMN TWO IS NAMED IN COLUMN TWO IS	AMERICAN COLUMNS OF THE PERSON NAMED IN	THE RESIDENCE OF THE PARTY OF T	A REPORT OF PERSONS ASSESSED.	NATIONAL PROPERTY AND PERSONS ASSESSMENT	OCCUPANT OF REAL PROPERTY.		
	Oz.	Plastic.	Fat.	Saccharine.	Total Carbon.	Carbon from Non-Nitrogenous.	Carbon from Non-
Milk 78 Fluid oz	IO	3.600	3.730	3.276	5.620	2.106	3.510
Arrowroot	9	0.300	0.000	4.300	2.350	091.0	2.190
Pancreatic Emulsion	73	:	000.1	:	0.740	:	0.740
Totals	18	4.200	4.750	8.176	8.710	2.266	6.440

This diet is to be given as follows:-

8 ozs. of Milk and 1 oz. of Arrowroot every 4 hours (6 times in 24 hours) for 24 hours.

10 ozs. of Milk and 1 oz. of Arrowroot every 4 hours for 24 hours.

12 ozs. of Milk and 1 oz. of Arrowroot every 4 hours for 24 hours.

13 ozs. of Milk and 1 oz. of Arrowroot every 4 hours for 24 hours.

The last quantity to be continued until solid diet can be borne by the stomach.

One-third of an oz. of Pancreatic Emulsion is to be mixed with a little water, or with a portion of the milk, and given

directly after each dose of Arrowroot and Milk, not mixed with the whole bulk.

DIET FOR DIABETES.

With this diet Hydrochloric Acid and Pepsine should be given to assist in digesting the very large quantity of Plastic Matter.

MARINEMANTEN				
Carbon from Saccharine por- tion of Non-ni- trogenous.	0.510 0.612			·
Carbon from Non- ogen- Nitrogen- ous.	0.5°0 000 005 0°2°0 0°510 0°510 0°102 1°480 0°750 0°260 1°485	6.472	:	:
Carbo Nitrogen- ous.	0 976 0 740 0 710 0 166 1 1928 0 0 15	4.670		:
Total Carbon.	1.536 0.830 0.745 0.366 0.366 0.117 1.000 1.480 0.555 0.260 1.620	10.947	0.612	10.335
Saccha- rine.	024 I.148 0.234	1.406		:
Fat.	0.712 0.110 0.055 0.250 0.055 0.055 0.055 0.050	6.113	:	:
Plastic.	1.800 1.380 1.310 0.308 3.556 0.030	9.074	:	:
Dry Oz.	жонн ок : ини ок	363	:	:
Food for 24 hours.	Cooked Meat and Poultry Cooked Pigeon or Game	Totals	Carbon from Saccharine to be deducted as waste	Total Carbon available

Salt to taste. Water as much as required by thirst. Tea without Sugar, with a slice of lemon peal in it. \* Green vegetables permitted—Cress, Celery, Endive, Greens, Lettuce, Mustard, Spinach, Water-cress. The following Wines may take the place of Spirits, (for equivalent quantities see Alcohol Table). Claret, Moselle (still), Rhine Wine, Manzanilla, Greek (St. Elie), very dry Amontillado Sherry.

### CHAPTER VIII.

"Amid all the dangers that threaten this Metropolis there is a sad certainty more serious than any one of them . . . . it is not disease but it is not health. It is a low state of vitality, of physical power, of mental energy, of enjoyment, and even of moral strength."—Times, August 4th, 1858.

THE DUTIES OF MEDICAL MEN—THE INTERDEPENDENCE OF DISEASES SHEWING THE IMPORTANCE OF PERIODICAL EXAMINATIONS—ANÆMIA AND FATTY DEGENERATION AS VESTIGES OF DISEASE—ABNORMAL PHYSIOLOGICAL STATES—CONDITIONS NOT RECOGNISED AS DISEASE BUT WHICH CERTAINLY ARE NOT HEALTH.

Selections from Lectures on the Germs and Vestiges of Disease and on the Prevention of the Invasion and Fatality of Disease by Periodical Examinations, delivered by the Author at the Royal Hospital for Diseases of the Chest, in 1861.\*

# (From Lecture 6.)

Gentlemen, we are all members of a practical profession. We have taken upon ourselves high and responsible duties, all culminating in action. So long as we choose to assume these duties in a profession, as yet so far from perfection, we are not justified in spending our time in scientific investigations or speculations, unless they have for their end some practical application for the good of humanity. It is the hope of attaining such an end which has led me on in the design and labour of these Lectures.

I proceed at once to lay it before you.

In the beginning of the first Lecture, I asserted, that

<sup>\*</sup> Published by Messrs Churchill, New Burlington Street.

we are justified in practising the profession of medicine only in proportion as we believe in the articles of the following creed :-

1. That man may be the instrument through whom the invasion and progress of premature destructive changes in the human organism may be prevented or arrested.

2. That man may be the instrument through whom the damaged organism may be more efficiently repaired.

3. That man may be the instrument through whom the

sufferings of the human being may be alleviated.

In what sense the organism is capable of accomplishing these ends without the instrumentality of man, I have shown you by numerous arguments and examples. I have also shown you the various modes by which it becomes deprived of this capability. The conclusion at which I now arrive is this, that man may be the instrument through whom the capability of accomplishing these ends may be preserved and restored to the organism.

The manner in which man is to exercise this instrumentality is the next point for our consideration. But I think we have almost reduced it to a necessary conclusion. For as we have plainly seen (in the previous lectures) that the organism is competent to take care of itself, provided that it possesses a normal Vital force, and is surrounded by normal conditions of life; and as we have also seen that the great causes of defect in the force, are the vestiges of disease and abnormal conditions of life; and as we have also learnt that the diseases, from which the vestiges result, are invited, by defects of the force; and that when thus invited and received into the organism, they are capable of being disposed of without leaving vestiges behind, if the Vital force is free from excessive defect; that thus these vestiges are due to defective force; and as we have learnt that the earliest invasion of defects in the force, upon which all the long and intricate succession of ills depend as a germ—as we have learnt, I say, that this state of germination exists at a period anterior to the manifestation of disease in its ordinary characters, and that it is to be found in the garb of slight impairments of the general health, the indications of which are more and more evasive and occult, the earlier the stage of germination; and, finally, as we have learnt that it is in this occult and evasive stage of germination that the defect is most easily and most efficiently to be remedied; I think you will agree with me in the practical conclusion at which I have arrived.

This conclusion is—that the manner in which man is to exercise his instrumentality for the prevention of disease, the prevention of the vestiges of disease, and the prevention of fatality in disease, is to search out these earliest evasive periods, of defect in the physiological state, and to adopt measures for their remedy. This appears to me to be the highest, the most ennobled duty of the physician, calling for the most abstruse knowledge of the science of life, the deepest experience in disease, the keenest exercise of the perceptive faculties, the calmest, most far-sighted reasoning and the wisest judgment,—a duty as much above the management of acute disease as to rule an empire is above fighting a pitched battle.

Now, Gentlemen, I am perfectly convinced, from my own observation and experience in practice, that patients never think of consulting their doctors till these conditions of impaired general health have advanced far enough to have been developed into some form of disease: that thousands and thousands of persons, believing themselves to be in health, are nevertheless undergoing these early, occult, and evasive stages of defect in the physiological state; and that such persons may be considered to

be in health, not only by themselves, but by any one accustomed to associate with them.

The only means by which to reach this evil and to obtain the good, would be for persons to submit themselves and their children to systematic periodical examinations.

Such examinations ought to include an inquiry into the family history, to learn the hereditary constitution; into the personal history, to learn all the previous diseases that have been passed through, and the habits and vicissitudes of life; into all the conditions of life surrounding the individual; into the condition of the organs and functions of the body; into the state of the secretions and fluids of the body by analyses and microscopical examinations; and so forth.

The examination should be reported in writing; and, after due consideration, such advice should be given as a careful judgment may dictate, for the future conduct, pursuits, and habits of the patient, with a view to correcting any defects or tendency to defects in the organism. Advice should also be given as to the means of removing any vestiges of disease that have been detected, or if they are not removable, advice as to the best way of overcoming their influence or of averting their increase. To this must be added precautions to be adopted in certain contingencies which, according to the judgment of the case, appear probable.

If such a plan as I have here proposed were to be faithfully and conscientiously carried out by the present and rising generation of well-educated studious medical men, immense benefit would be conferred upon the public. The next question is, then, what would be the effect upon the profession in a pecuniary and in an ethical sense. With regard to the pecuniary question, it is only necessary to observe that of course I do not expect that any man in

good practice, whose time is profitably employed, could conduct such an examination and give such advice for the usual consultation-fee. Every man who attempts to follow out the plan, should require such a fee as might enable him to give the necessary time and consideration to every case.

But this should in no way interfere with the power of the poor to participate in such a system. Every hospital and dispensary should institute a distinct department for the conduct of such examinations, and for giving the necessary advice. Every patient discharged from its wards should be submitted to this department before returning to the duties of life.\*

I have again and again referred in these Lectures to the numberless anomalous symptoms, the pains, discomforts, nervous disturbances, etc., etc., which affect persons in abnormal physiological states, and which increase in their severity, obviousness, and number, as the states of health become more and more degraded, and the occurrence of some acute disease becomes more imminent. I have pointed out to you that these are the states of health which fill our consulting-rooms and the out-patient departments of hospitals and dispensaries.

In further confirmation of my statements on this point I may remind you of some apposite observations by Sir Henry Holland on the subject of symptomatic complaints. In his Medical Notes and Reflections he says, speaking of gouty blood—"Irregular actions of the heart, hypochondriacal depression, as well as the more common symptoms of dyspepsia and disordered secretions, frequently antecede by months

<sup>\*</sup> For details of the mode of carrying out the Examinations at public Charities, see a paper by the Author read at the Guildhall, June 1862. "Transactions of the National Association for the promotion of Social Science."

the first appearance of gout in the extremities, and occa sionally give serious alarm even to those who look with medical eyes upon these ambiguous cases."—(p. 246, 3rd edition.)

"Modern observation has led us to recognise some of these relations (of gout with local or constitutional disorders) under the names of gouty headache, gouty ophthalmia, and gouty bronchitis. My own experience would lead me to add many cases of asthma to the number. I have so often seen this disorder prevalent in gouty families, affecting those who do not undergo the disease in the joints, and ceasing wholly or in part when the gout appears externally, that I cannot doubt the existence of this relation.

"The greater tendency to apoplexy in this habit is noticed by many of the older writers, and confirmed by general experience.

"Reference has already been made to hypochondriasis and hysteria; and it is probable that other disorders of the same class, still less generally viewed under this connection, will hereafter be submitted to it.

"The relation of gout to the functions and disorders of the liver is another point of much interest in pathology, clearly attested both in the active symptoms of the disease and by those which are common under other forms of the gouty temperament. This, moreover, is one of the points associating it with that group of maladies bearing the vague name of dyspepsia.

"The connection of gout with cutaneous affections is an additional topic, yet almost unexamined; though I cannot doubt, from my own observation, that certain of these disorders occur as effects of the habit in question."—(Ibid, pp. 253-54.)

The enormous quantities of medicine dispensed in the

out-patient departments of hospitals and dispensaries for the temporary relief of this class of functional derangements and local diseases-for complaints which might be prevented by the patients themselves if they were properly informed of the causes and premonitory symptoms of their maladies-these are facts which must be perfectly familiar to all my hearers. Such a system of examination and advice as I propose, if properly carried out, must strike at the root of these evils, and would at the same time reduce the miserable over-crowding of the hospital waiting-rooms, and the enormous expenses incurred for drugs. These are considerations which, however important as elements of social and political economy, are elevated far above the rank of financial questions by the fact I have endeavoured to demonstrate in these Lectures, that by these same means, and at the same time, we shall so largely promote the economy of life.

I hope, Gentlemen, that you will draw the attention of the treasurers and governors of any hospitals to which you may belong to this subject. It is necessary to the credit and honour of our profession, that improvements in these medical establishments should not come from the public to us, but should originate among ourselves, and be

urged by medical men upon the public attention.

The following sketch of a common complaint, and a common story, may illustrate the need for some such system as I propose. A lad of fifteen is admitted into a hospital ward with a first attack of rheumatic fever. He is treated with skill, and nursed with care, and in two or three weeks he is convalescent, and returns to his home without having received any damage to his heart or other organs. But he has no proper understanding of the nature of his complaint, of the conditions of life calculated to keep up the morbid influence in his organism, no clear

notions of the diet which he ought or ought not to adopt, no knowledge of the premonitory symptoms by which a fresh attack of rheumatic fever is heralded, or of the precautions necessary when such an attack is feared. He has probably a general idea that his great enemy is cold, and his great friend flannel; and that is as likely to lead him

wrong as right in the measures he adopts.

He goes back to his home and his pursuits apparently well. He keeps his skin closely cased in flannel, and his dwelling-rooms warm, but he neglects altogether to provide for efficient excretion by the skin. He chooses a business, either utterly regardless of its fitness for his constitution, or makes as great a mistake by selecting one in which he may be sure of warm, that is to say close, rooms and workshops by which he is deprived of exercise and oxygen. He drinks beer, eats cheese, and so forth, like other people in his position. After a few months he comes back as an out-patient at the hospital, with severe acid dyspepsia, and after consuming the usual amount of drugs for several weeks, gets relief, and goes back to his old habits. A few months more, and he appears again, the subject of a skin-disease; goes again through the consumption of drugs, and gets well and goes back to work. By and by he comes again with diarrhœa, and goes through the same process; at another time with gravel, and gets relief again. At length, a few years perhaps having elapsed, and after some months of depressed health, with palpitations of the heart, gloomy thoughts, irritable temper, and general malaise, he happens to be out on a damp raw day, gets a chill, and applies again at the hospital, with a fresh attack of rheumatic fever. time he suffers from endocarditis, and after some weeks returns again to his home, "discharged cured," but with disease of the valves of his heart. He has been thoroughly well treated, and is very free from rheumatic poison, and hence goes on for a considerable time without much inconvenience; but having no clear ideas of the nature of the damage he has received, or of the precautions necessary to prevent its increase or the production of secondary diseases dependent upon this damage for their cause, he gradually becomes the subject of congested lungs and liver; of attacks of bronchitis to which he was not formerly inclined; his breathing becomes short, his old dyspeptic troubles, cutaneous affections and gravel, recur again and again, and his capability of following his former occupations gets less and less.

If he does not have another attack of rheumatic fever, he comes back to the hospital some future day with chronic bronchitis, or with apoplectic symptoms, or with congested liver, or still later with albuminuria and dropsy. At length he dies, and his death is registered under the

head of apoplexy, heart-disease, or dropsy.

You will admit that this sketch is no exaggeration; that I have, in fact, omitted numerous details of minor diseases and discomforts, that are sure to have existed in the case itself. I have said nothing of the effects upon this man's children of his continued ill-health, or of the poverty and want of food brought upon his wife and family by the same cause, and acting as fresh causes of disease in them. But I have said enough to make it quite clear, that, in the course of a life prematurely ended, he must have consumed a vast amount of money in the form of drugs, and a vast amount of nervous energy, if not of brains, in the form of medical advice.

It is to the first causes, to the "wells and springs" of such a series of calamities as this, that I have directed your attention as fellow medical practitioners, and to which I now beg that you will direct the attention of

the Treasurers and Governors of Hospitals and Dispensaries, of your private patients, and of the public at large.

In my fourth lecture I gave an Etiological analysis of those states of disease assembled under the following 6 headings in the Registrar-General's Reports of the deaths in London. (Report of the deaths in London registered in the 21st week of each of the ten years 1848-57. The Report current at the time the notes for these Lectures were made.)

- 1. Typhus, typhoid and other forms of continued fever.
- 2. Apoplexy and Paralysis.
- 3. Heart diseases and Pericarditis.
- 4. Rheumatism and Gout.
- 5. Bronchitis.
- 6. Atrophy and Debility.

In this analysis I demonstrated that when we analyse the natural history of any disease we find that the principal factors of its essential cause, of its predisposing causes and of the causes of its fatality fall necessarily under one of the three headings:—1. Conditions of life. 2. Coetaneous diseases. 3. Vestiges of Disease.

I showed not only theoretically but from actual observations made by others as well as by myself, how the organism becomes damaged by these Vestiges of Disease—how the vital force becomes defective through these Vestiges, how this defective state of the vital force becomes the essential cause and the predisposing cause of disease; and how the Vestiges of one disease become the causes of fatality in whole families of other diseases.

I endeavoured to prove, by an array of facts, that the vestiges of disease become causes of fatality in other diseases principally in two ways.

1. By destroying those modes of matter and that corre-

lation of conditions upon which the existence of the vital force in its normal condition depends, thus producing excessive defect of the vital force.

2. By producing excessive defect in the condition of some part of the organism, occupying the position, at the time, of an essential *instrument* in the processes of life, and thus causing the organism to break down at this its weakest part.

I pointed out that in the large majority of deaths from disease, the fatality is due, not to the disease itself, but to the vestiges of some pre-existent disease, operating in one or other of the above ways.

In illustration of this great fact, I have set forth the course of events, by which the vestiges of disease, passing under the names of ANÆMIA and FATTY DEGENERATION, become the actual causes of a large number of the deaths registered under the following names:—

Tabes mesenterica, croup, measles, hydrocephalus, whooping-cough, dentition, convulsions, apoplexy, paralysis, delirium tremens, intemperance, angina pectoris, diseases of the heart, pneumonia, diarrhœa, mortification, influenza, peritonitis, childbirth, bronchitis, jaundice, liver disease, kidney disease, and some others.

As an indication of the *insidious way* in which the deadly influence is exerted by these states,—ANÆMIA and FATTY DEGENERATION—(which are only examples of a class), I called attention to the fact that their names do not appear in the bills of mortality. (See Anæmia, Fatty Degeneration.)

My object now is to exhibit in a conspicuous manner the Interdependence of diseases.

I have therefore restricted myself to a statement of the influence exerted by the vestiges of each of the diseases analysed in the preceding Lectures upon the rest of the

same little group, and in order that the different relationships may be seen at a glance I have arranged the facts in a tabular form.

In the following tables the first column gives a list of the vestiges of the disease at its head. The second column shows to which of the diseases already analysed each vestige may become the predisposing cause. The third column shows to which it may become the essential cause, and the fourth column shows to which it may become the cause of fatality. It must be remembered that several vestiges usually concur in producing their effects.

## TABLES SHEWING THE INTERDEPENDENCE OF DISEASES REGISTERED UNDER THE FOLLOWING NAMES: --CONTINUED FEVER-APOPLEXY AND PARALYSIS -HEART DISEASE AND PERICARDITIS-BRONCHITIS-ATROPHY AND DE-BILITY.

AND CAUSES OF FATALITY IN-	Bronchitis. Typhus. Heart-dis- ease. Subsequent attacks of rheumatism and gout,	Bronchitis. Rheumatism and gout. Typhus and heart-disease.	Bronchitis, Rheumatism and gout, Typhus, Heart-disease, Atrophy	and debility.		Paralysis. Bronchitis. Typhus. Heart-disease.	Rheumatism and gout. Typhus. Heart-disease. Atrophy and de- bility.
ESSENTIAL CAUSES OF	Apoplexy. Mechanical heart-disease			debility, de- Local paralysis.		Atrophy and debility in the subject, and in the offspring if the subject is a female.	Dilatation and degeneration in hypertrophy of heart .
BECOME PREDISPOSING CAUSES OF	Bronchitis			Atrophy and debility, degenerative heart-disease .	A repetition of an attack of gout or rheumatic fever.	Typhus; heart-diseases, (de- generative and mechanical)	Bronchitis, rheumatic and gouty attacks. Typhus .
VESTIGES OF ONE OR MORE AT- TACKS OF RHEUMATIC FEVER.	Valvular disease	Pericardial effusion. Pericardial adhesions to the heart or pleura	Pleuritic adhesions and effusions	Stiffened joints	Tendency to a repetition of the attack, and of its determination to the damaged parts	Anæmia	Debility and nervous exhaustion, especially from repeated attacks

			TALLET.	DELENDE	NCE OF	DISEAS	ES.			
AND CAUSES OF FATALITY IN-	Bronchitis. Rheumatic fever.		Bronchitis. Rheumatic fever.	Bronchitis, Heart-disease, Rheu- matic fever. Typhus.	Rheumatic fever. Typhus.	Apoplexy. Bronchitis. Heart-disease. Rheumatism and gout. Typhus.	Heart-disease. Typhus.	All forms of heart-disease. Rheumatism and gout. Bronchitis. Apoplexy and paralysis. Typhus.	Bronchitis. Heart-disease.	Heart-disease. Bronchitis. Atro- phy and debility. Typhus.
ESSENTIAL CAUSES OF	Dilatation and hypertrophy of the heart.			Apoplexy and paralysis		Aggravated hypertrophy and dilatation of the heart. Appoplexy and paralysis .	Atrophy and debility.			Apoplexy
BECOME PREDISPOSING CAUSES OF	Bronchitis, Apoplexy and Paralysis (by inclining to fibrinous clots in the circulation).	Fatty degeneration of the heart	Fatty degeneration of the heart.	Bronchitis. Apoplexy .	Bronchitis. Toxic heart-disease	Bronchitis		Debility		Apoplexy
VESTIGES OF DISEASES OF THE HEART AND PERICARDIUM.	Constriction or insufficiency of the valves. Adhesions of the pericardium	Tendency to degeneration after repeated inflammations	Hypertrophy of the heart	Venous congestion of the lungs, brain, liver, and all parts in arrear of the obstruction in the heart	Tendency to inflammations, hæmorrhages, and fluxes in the organs congested	Obstruction to the systemic arterial circulation, secondary to the venous congestion	Dilatation and defect in the force of the systemic circulation .	Tendency to death from any subsequent disease, surgical operation or accident	Angina pectoris	Diseases of the kidneys, from continued congestion. Dropsy

			INT	ERDEPE	NDE	NCE	OF DI	SEAS	ES.		
AND CAUSES OF FATALITY IN-	Bronchitis.	Bronchitis. Atrophy and debility. Heart-disease.	Bronchitis. Apoplexy.	Heart-disease. Rheumatic fever. Atrophy and debility. Bron-chitis. by exhaustion from ex	cessive discharge, or asphyxia from accumulation of secretion.		Typhus. Apoplexy. Subsequent	Bronchitis.	Bronchitis, Paralysis, Any severe disease, especially typhus and	others due to a morbid poison in the blood.	Heart disease. Rheumatism and gout. Typhus. Apoplexy.
OF				and .							
ESSENTIAL CAUSES OF				Heart-disease. Atrophy and debility							ease.
CAU				Atro					bility		t-dis
TAL		sase .		sase.					nd de		Неал
SEN		Heart disease .		eart-disc debility					hy ar		lexy.
Es		Hear		Hear					Atrophy and debility.		Apoplexy. Heart-disease.
Ð									,		
BECOME PREDISPOSING CAUSES OF		Heart-disease							n.		ism
EDIS:		eart-c							natisı		Rheumatism
PRAUSE									Rheumatism. lisease.		Rhe
CC	chitis	chitis	hitis	lexy		hitis.	exy		s. B		exy.
BEG	Bronchitis	Bronchitis.	Bron	Apop		Bronc	Apopl		Typhus. Rheum Heart-disease.		Apoplexy.
ri.			Hypertrophy of the heart, especial- Bronchitis ly dilatation .	ith.		er-	de-				
HITTE			· espec	or wi		1 (K	and				1 to
RONC			eart,	with		estion	stion d bloc	walls			secondary to
F B		•	the b	tis, (hœa)		cong	conge	hial			seco.
SES (	onchi	na	hy of tion	onchi		snou .	nous	bronc			ease,
VESTIGES OF BRONCHITIS.	Dilated bronchi	Emphysema	ypertrophy o	nonic bronchitis, out bronchorhæa		tic ve	eneral venous congestion a ficiently oxygenated blood	ened	ty		idney-disease, other vestiges
Δ	Dila	Emp	Hype ly	Chronic bronchitis, (with or with- Apoplexy out bronchorrhæa)		Hepatic venous congestion (Kier- Bronchitis.	General venous congestion and deficiently oxygenated blood	Thickened bronchial walls.	Debility		Kidney-disease, other vestiges
								(200	1		-

AND CAUSES OF FAIALLIY IN—			Bronchitis, rheumatic and Atrophy and debility. Atro- Bronchitis. Rheumatism and gout. gouty attacks, typhus phy and debility in the off- Typhus. Heart-disease. Atrophy spring if the subject is a and debility.
ESSENTIAL CAUSES OF	Heart-disease.		Atrophy and debility. Atro- phy and debility in the off- spring if the subject is a
BECOME PREDISPOSING CAUSES OF	Apoplexy	Attack of gout.	Bronchitis, rheumatic and gouty attacks, typhus .
VESTIGES OF ONE OR MORE AT- TACKS OF GOUT.	Deposits of urate of soda in and Apoplexy about joints and some other parts	Tendency to a return of the attack Attack of gout, in the parts previously affected.	Anæmia and nervous exhaustion, especially from repeated attacks

And Causes of Fatality In—	Rheumatism and gout. Apoplexy and paralysis. Bronchitis and typhus.	Paralysis. Apoplexy. Rheumatism and gout.			Paralysis. Bronchitis. Rheumatism and gout. Typhus. Heartdisease. Atrophy and debility.
ESSENTIAL CAUSES OF	Apoplexy and paralysis. Heart-disease (mechanical and degenerative). Atrophy and debility.	Apoplexy (uræmic). Heart-disease			Atrophy and debility in the subject. Atrophy and debility in the offspring if the subject is a female. Dilatation and degenerative hypertrophy of the heart
BECOME PREDISPOSING CAUSES OF	Apoplexy and paralysis. Bronchitis	Bronchitis	Degenerative heart-disease.	Attacks of gouty or rheumatic inflammations, when the skin disease is suppressd.	Degenerative and mechanical heart-disease. Bronchitis. Rheumatic and gouty attacks. Typhus
GOUT, WHEN THE ESSENTIAL CAUSE HAS REMAINED LONG IN THE ORGANISM WITHOUT PRODUCING ACUTE GOUTY OR RHEUMATIC INFLAMMATION.	Fatty and calcareous degenerations or deposits in the heart, arteries and capillaries. Degeneration of the tissues generally.	Chronic disease of the kidneys	Stiffened and contracted joints .	Cutaneous affections	Anæmia and other forms of debility.

VESTIGES OF RHEUMATISM OR

AND CAUSES OF FATALITY IN—	AND CAUSES OF FATALITY IN-	Paralysis.	Bronchitis. Atrophy and debility. Rheumatism and gout. Heart- disease. Paralysis.		(When affecting the brain), apoplexy.	Heart-disease. Apoplexy. Bron-	Any other form of continued fever.
ESSENTIAL CAUSES OF Rheumatism, gout, heart-disease and debility in the generation concerned.	ESSENTIAL CAUSES OF	Degenerative heart-disease, in cases of pre-existent hypertrophy	Atrophy and debility. Heart-disease.				Atrophy and debility.
BECOME PREDISPOSING CAUSES OF Rheumatism, gout, and heart- disease in the generation concerned	BECOME PREDISPOSING CAUSES OF	Debility. Degenerative heart-disease	Bronchitis. Attacks of rheu- matic or gouty inflamma- tion. Acute specific dis- eases.	Heart-disease and apoplexy, in those previously affected by other Predisposing Causes		Apoplexy and paralysis.	
VESTIGES OF RHEUMATISM OR GOUT ACTIVE OR LATENT.  The rheumatic or gouty constitution transmitted to a future generation	VESTIGES OF TYPHUS, TYPHOID, AND OTHER FORMS OF CON- TINUED FEVER.	Tendency to fatty degeneration .	Great defect in the vital force especially during protracted convalescence.	Increased irritability in the nervous system	Softening of parenchymatous organs and of the tissue generally.	Tendency to extravasations of blood	Typhoid deposits, and enlarged mesenteric glands, spleen, etc

INTERDEPENDENCE OF DISEASES.										109
AND CAUSES OF FATALITY IN-	Bronchitis. Heart-disease. Apo-		Typhus. Apoplexy.	AND CAUSES OF FATALITY IN		Atrophy and debility. Typhus.	Apoplexy. Typhus.	Apoplexy. Paralysis. Heart-dis-		Typhus. Atrophy and debility. Bronchitis. Rheumatism and gout.
EDISPOSING ESSENTIAL CAUSES OF	. Atrophy and debility, by producing fatty degeneration.		· Paralysis · ·	DISPOSING ESSENTIAL CAUSES OF	plexy.				. Heart-disease, (degenerative.)	
BECOME PREDISPOSING CAUSES OF		Apoplexy and paralysis.	Apoplexy .	BECOME PREDISPOSING CAUSES OF	Paralysis. Apoplexy.					
VESTIGES OF APOPLEXY,	Paralysis of sensibility or motion •	Strong tendency to a recurrence of apoplexy in a more severe form.	Softening of some part of the brain Apoplexy	VESTIGES OF PARALYSIS,	Tendency to recurrence of the attack in a more severe form, or to apoplexy	Tendency to mortification in the paralysed parts	Softening of a portion of brain or spinal cord	Diseases of the urinary tract	Degeneration of the paralysed parts, or of the organism generally	Tendency to death in an attack of acute or chronic disease.

110	INTERDEPENDENCE OF DISEASES.								
AND CAUSES OF FATALITY IN-	Bronchitis. Typhus.	Bronchitis, Typhus,	Apoplexy and paralysis.	Bronchitis, Typhus. Rheumatism. Apoplexy and paralysis. Heart-disease.	Apoplexy. Heart disease. Rheu- matism. Typhus. Atrophy and debility.	Typhus.	Bronchitis. Heart-disease.		Bronchitis. Heart-disease. Ty- phus. Rheumatism and gout.
ESSENTIAL CAUSES OF A	Atrophy and debility B	Atrophy and debility E	Debility. Heart-disease. Apo- A plexy and paralysis.	Atrophy and debility from E deficient circulation .	Apoplexy and paralysis (if A clots float into the cerebral or spinal vessels)	Atrophy and debility T		Atrophy and debility, local paralysis	Atrophy and debility . I
BECOME PREDISPOSING CAUSES OF	Typhus and acute specific diseases generally	Paralysis (from strumous disease of vertebræ)	Apoplexy and paralysis, heart-disease	Bronchitis	Apoplexy or paralysis	Atrophy and debility.	Atrophy and debility. Local paralysis		
VESTIGES OF ATROPHY AND DEBILITY,	Rickets, in children produced while the mother is suffering from atrophy and debility.	Scrofulosis or tuberculosis, in the children produced while either parent is suffering from atrophy or debility	Degeneration of tissues generally, or of those of some organ or part	Dilatation of the heart, especially in those who have made frequent ascents during atrophy and debility.	Fibrinous clots in the heart or vessels which may float off	Lardaceous or albuminoid deposits especially in the liver, spleen, lymphatic glands, and kidneys.	Deformities in the osseous skeleton	Arrest of development in certain muscles or other parts of the organism	Effusions into serous cavities, and discharges from the mucous passages .

ch disease, rheumatism, and

Tendency to become affected by Typhus, any morbid influence to which diseas the organism may be exposed gout. (general predisposition to disease)

Tendency to death from all acute and chronic diseases, occurring during the continuance of atrophy or debility

Apoplexy and paralysis, bronchitis, heart-disease, typhus, rheumatism and gout. Atrophy and debility.

Having shown the remarkable interdependence which exists between different diseases—how they act and react upon one another. I must attempt in this Lecture, to show to what extent one vestige of disease, or the several vestiges of one disease, may act as essential causes and as causes of death in other diseases. I will first take some Diseases which have no place in the Registrar's report. This will enable us to observe how many of the headings under which deaths are in that report owe their existence there to diseases which never figure at all as causes of death.

The first example I have chosen is Anæmia, a name that does not appear in the Registrar's Reports.

## ANÆMIA.

A careful examination of the subject teaches us that Anæmia is one of the most frequent vestiges of all severe diseases, when they do not terminate in death—the patient remains anæmic during convalescence, and in a large number of cases continues so long after returning to the duties of life. Again, we observe, that anæmia is continually occurring in connection with the periods of puberty and the cessation of the catamenia, and that during the intervening years, it is one of the most frequent forms of impaired health connected with derangements of menstruation, with lactation, with convalescence from the puerperal state, and during pregnancy.

Innumerable other causes may be found for this condition, which is, in fact, the type of most of those miserable states of health referred to by *The Times* in the memorable article already quoted—"There is a certainty worse than any occasional and precarious plague. We may anticipate it from our own experience—we may see the great fact with our own eyes. . . . . It is not disease,

ANÆMIA. 113

but it is not health. It is a low state of vitality, of physical power, of mental energy, of enjoyment, and even of moral strength. . . . . Shocking as it may seem, a plague once in twenty years seems but a light evil to so low a condition of humanity."

From a concatenation of circumstances, it so happens that the female portion of the population have to support the principal weight of this dire affliction. Anæmiated girls, anæmiated brides, anæmiated spinsters, anæmiated mothers, abound in the consulting room, the outpatient room, the hospital ward, the home; wherever, in fact, the physician's duties call him there they are to be found. Yet from the long list of diseases named in the mortality tables, this name, anæmia, is absent. Is it, then, a thing to be endured, but not to be feared? Is it a name for sickness only, and not a cause of death? Quite the reverse, gentlemen, the fact is that, like some individuals of great influence and importance, anæmia travels incognito. When we discover its various "nommes de guerre," we are startled to find ourselves face to face with the impersonation of long-dreaded enemies.

Tabes mesenterica		18.3	deaths	per week.
Croup		6.8	"	
Measles		25.0	"	"
Hydrocephalus .		32.5	"	"
Whooping-cough		51.1	,,	"
Pneumonia		60.3	"	"
Dentition		12.8	"	"
Diarrhœa		17.5	"	"
Convulsions .		36.5	"	"
Bronchitis		64.5	"	"
Childbirth	14.	4.3	"	"
			,,	"
	6	329.6		

These are some of the names under which anæmia travels among the sick, and takes its place in the bills of mortality. The way in which it does so is as stealthy as it is deadly. I will tell you something of how it happens.

"Among the children of the poor of London, the most widely-spread of these diseases (tuberculosis, scrofulosis, rickets,) is rickets. It is, however, by no means limited to the poor, or to London, or even to large towns." I quote from one of the most correct medical philosophers of modern times.—(Dr., now Sir W., Jenner, Lectures on Rickets, Medical Times and Gazette, March 17, 1860.)

"Rickets is essentially and purely a disease of nutrition, not of one part only, but of the whole body."

"Rickets causes, primarily or secondarily, more deaths

than any other disease of childhood."

"The great causes of death in rickets are:—1. Intensity of the general cachexia. 2. Catarrh and bronchitis. 3. Albuminoid infiltration of organs, especially of the lymphatic glands and spleen (but also of the liver). 4. Laryngismus stridulus. 5. Chronic Hydrocephalus. 6. Convulsions. 7. Diarrhœa."

You see, gentlemen, how closely this list of the causes of death in rickets corresponds with the list I have just given of what I have called the "nommes de guerre" of anæmia; and the correspondence is really closer than it

appears at first sight.

"The connection between rickets and laryngismus stridulus," continues Dr. Jenner, "is very close. . . . . I believe the reason of laryngismus stridulus being so constantly referred to teething is that the ricketty condition retards the development of the teeth, and the medical practitioner refers the laryngismus to that which, like itself, is the consequence of constitutional disease.

"If a child pass over the ninth month without teeth, you should carefully inquire into its cause. . . . . It may be, and this is infinitely the most common cause of late dentition, that the child is rickety."

Many deaths, therefore, registered under the head of DENTITION, may be referred to rickets.

"Catarrh and Bronchitis," says Dr. Jenner, "are unquestionably the most common causes of death in rickets. The softening of the ribs rendering the mechanical power by which inspiration is performed so defective, that the impediment offered to the entrance of the air by the mucus in the bronchial tubes cannot be overcome; and collapse of large portions of the lung follows.

"This want of inspiratory power, and the consequent accumulation of mucus in the bronchial tubes, affords an explanation of the extraordinary mortality of measles, whooping cough, and bronchitis in ricketty subjects."

Thus may we refer two more of the headings of the Registrar—MEASLES and WHOOPING COUGH—to rickets.

For similar reasons, and still more from the intensity of the general cachexia and the corresponding defect in the vital force, rickets will be found to be the true cause of fatality in many of the deaths registered under the name of PNEUMONIA.

Then we have the deaths from CHILDBIRTH 4.3 per week, and it is very interesting to trace out the influence of rickets in causing these miserable deaths. The mode of its operation is twofold. First, there is the large size of the head in the rickety fœtus; and, secondly, deformities of the pelvis in women who have been rickety in their childhood. On examining into the causes of death in childbirth, as carefully collated by Dr. R. Collins, of the Dublin lying-in hospital, I find that of eighty-one cases of

death during parturition, thirty-two were due to rupture of the uterus, eleven to tedious or difficult labour. If we, then, enquire into the causes of rupture of the uterus and tedious or difficult labour, to which forty-three out of eighty-one deaths were due, we find that narrowing and deformity of the pelvis, and abnormal size of the child's head, stand prominently forward in their importance; and thus we are brought back to rickets as the chief cause of these deaths.

But you will perhaps ask, how anæmia is proved to be the cause of the deaths under these different headings, by proving that they are due to rickets? That is, in fact, the important question. The answer is this, that anæmia in the mother produces rickets in the child, and anæmia in the child may lead to rickets also. This may be as familiar to you all as it is to me; but I will give you the authority of Dr. Jenner again, who has devoted great

attention to this subject :-

"The health of the mother has a decided influence in the development of rickets in the offspring. . . . Of this much, I am sure—that when the mother is in delicate health, in a state of which anæmia and general want of power form the prominent features, without being the subject of disease usually so-called, there the children are often in a very decided degree rickety, although the father is in robust health, and the hygienic conditions in which the children are placed are most favourable." "It is very common for the first two or three children to be free from any signs of rickets, and yet for every subsequent child to be rickety.

"Whatever external conditions are favourable to the formation of hydræmic blood in a child, seem to be favour-

able to the development of rickets.

"Albuminoid infiltration of the lymphatic glands,

spleen, and other organs is by no means an uncommon cause of death in rickets. The two great features, during life, of albuminoid infiltration of these organs in a young child, are emaciation and pallor; the anæmia is often most remarkable."

You will not fail, then, to see at once the intimate connection between rickets and anæmia, and between anæmia and that list of terrible names in the Registrar's Reports.

I do not wish you to suppose that I attribute all the deaths under those headings to anæmia; that would be a great mistake. But from an elaborate and careful analysis of such deaths, I have found, and you may find, that a very large, a very remarkably large, proportion of them are due to anæmia in the individual, or to anæmia in a parent and rickets in the offspring. It is important to bear in mind, that, with the exception of hydrocephalus, the diseases I have enumerated from the mortality-tables are extremely common every-day complaints-that they occur and terminate favourably thousands of times every year; that it is only the fatal cases that come into our list, and with which we now are concerned. And what I have endeavoured to show, and what I wish to impress upon you is, that we must look to anæmia as the chief cause of this fatality; that it is anæmia in the individual or anæmia in a parent that brings these remediable, every-day complaints into the bills of mortality.

When, therefore, we see anæmia establishing itself, as a vestige of the diseases from which our convalescents have lately suffered, or in any other way, we must regard it as the grim harbinger of death in a vast family of diseases.

The next example I shall give is Fatty degeneration which like the last does not appear in the Registrar's Reports.

## FATTY DEGENERATION.

In the present day, Gentlemen, we have all learnt that "something much more than a general tendency to form fat, or a general excess of fat in the blood is necessary to produce a local fatty degeneration."—(Paget's Lectures,

Vol. I., p. 112).

Every year throws new light on the physiology of the process by which this change is brought about; but of this it is not within the province of these Lectures to treat. I wish, however, to point out to you that the pathological condition termed fatty degeneration is to be regarded as a vestige of disease—a vestige of perverted assimilation, either local or general, and hence a vestige also of any diseases by which such perverted assimilation is produced. In the third series of Guy's Hospital Reports (Vol. iii. 1857), Dr. Wilks has related a number of interesting cases of fatty degeneration, of which the only appreciable and probable causes were hæmorrhage, diarrhœa, miasmata. In these cases the subjects were comparatively young, the heart was the organ in which the diseased change was most marked, and the body generally was neither fat nor wasted.

Fatty degeneration appears to be especially prone to occur in tissues which have passed from a condition of active assimilation to one of comparatively inactive or feeble nutrition, by whatever cause this change may have been brought about. Thus, the muscular tissue is

particularly subject to fatty degeneration; it is also peculiar for the vigour of its normal assimilation, and it is when this is rendered feeble that degeneration occurs. This appears to be the case whether the vigorous assimilation is hindered by arrest of function in the muscles, as in the case of paralyzed limbs, or by deficient supply of blood, as in disease of the nutrient arteries of a part, or by a depraved condition of the blood supplied to the part, as in the case of persons who, after having led active muscular lives, gradually become anæmiated by passive hæmorrhages. I have seen many instances of strong men, accustomed to vigorous country occupations, who, having become the subjects of bleeding hæmorrhoids, by which the blood lost its red globules, and nutrition became enfeebled, suffered from degeneration of their previously strong and vigorous hearts. Many other examples might be adduced of the degeneration of tissues when their assimilation is changed from a vigorous to a feeble state. Thus we find, in watching the course of diseases of the heart, that the heart which has become hypertrophied in opposing some obstruction to the circulation during the active life of the individual, becomes degenerated when, in the further progress of the case, the patient is forced to relinquish active pursuits, and thus ceases to call upon the hypertrophied organ for the full exercise of its muscular power. Thus, also, parts which have been inflamed are especially liable to degenerate, and fatty degeneration frequently takes place in organs deprived of their proper functions by disease, as in kidneys spoilt by Bright's disease. Thus, also, the products of inflammation, when they have no further functions to perform, are peculiarly subject to fatty degeneration. Without pressing the subject further, I think we shall all agree that fatty degeneration must be regarded as a vestige of disease, not

as a disease in itself, and that in thus regarding it, we must often look beyond the defect in assimilation to the causes of that defect, in order to find the disease of which the degeneration is truly a vestige.

I must now proceed to show to what a wide extent this vestige of disease acts as a cause of those deaths classed in the Registrar's reports under the heads of a variety of diseases.

"The most interesting examples (of fatty degeneration)," says Mr. Paget, "are those of primary degeneration of blood-vessels. This has long been known in the atheromatous disease, as it was called, of the larger arteries, the true nature of which, as a fatty and calcareous degeneration of the inner and, consecutively, of the middle arterial coat, was discovered by Mr. Gulliver (see Med. Chir. Transactions, Vol. XXVI.) The descriptions of this complaint by him and by Rokitansky have left nothing unsaid that is yet known; but the observations are each year becoming more numerous and more interesting of similar changes in the minutest blood-vessels. Such changes are especially observable in the minutest cerebral vessels, and their importance in relation to APOPLEXY of which they seem to be the most frequent precedent, cannot be overstated."-(Ibid., Vol. I., p. 139.)

Dr. Wilks' experience has taught him that, "In the majority of cases of sanguineous apoplexy, disease of the blood-vessels exists."—(Lectures on Pathological Anatomy, 1859.)

Dr. Kirkes has shown (Med. Chir. Transactions, Vol. XXXV.) that paralysis, consequent upon arrested circulation in some portion of brain, is frequently the result of the obstruction of healthy cerebral arteries by masses of fibrine carried into them, after being dislodged from the

valves of the left side of the heart, or from some part of the arterial system.

And Dr. Ormerod has pointed out that it is in the cachetic subjects, with feeble circulations, that such masses are likely to form in the heart, in the very persons, in fact, who are likely to be the subjects of fatty degeneration of this organ.—(Observations on the Clinical History and Pathology of one form of Fatty Degeneration of the Heart, by E. L. Ormerod, M.D. Medical Gazette, 1849.)

And although Dr. Ormerod was not prepared at that time (1849) to consider the occurrence of fibrinous clots in the subjects of fatty degeneration to be more than a frequent coincidence, we do not now doubt that the languid circulation, and the inefficient contractions of the heart in the subjects of fatty degeneration, act, together with other circumstances, in causing these deposits of fibrine from the blood.

Describing the appearances after death in fatal cases of Delirium tremens, Dr. Wilks says, "The body, as a rule, presents many degenerative changes, brought about by the intemperate habits. It is this alteration of the viscera, I think, to which death is owing. Delirium tremens is a recoverable affection until such changes have occurred in the tissues that improvement is no longer possible; and we then find in the body various morbid changes. These are mostly of the fatty kind, as all alcoholic drinks tend to this condition."—(Ibid.)

In Angina Pectoris, again, Dr. Wilks, in common with other observers, has found that "the heart is usually fatty, and the coronary arteries ossified."—(Ibid.)

"It is impossible," says Dr. Ormerod (*Ibid.*), "to read any collection of cases of angina pectoris without . . . . feeling how much further fatty degeneration goes to explain the symptoms than does any other morbid change usually found on dissection."

Mr. Paget classes the heart and arteries first among the frequent seats of fatty degeneration.—(*Ibid.*, Vol. I., p. 116).

Among the vessels of which the coats have been found degenerated are those of the lungs and placenta; and pulmonary or uterine hæmorrhages may result from this state of the vessels.

Against the extreme vital depression which accompanies such diseases as Peritonitis, influenza, and diarrhea, the subjects of fatty degeneration have no resisting power; they are among the first to succumb; and thus fatty degeneration becomes a cause of death in these and other depressing diseases when they attack adults.

Out of 164 deaths from Childbirth in Dublin Hospital, during the seven years ending November, 1826, as recorded by Dr. R. Collins, I find thirty two attributed to rupture of the uterus, and eleven to tedious or difficult labour. Recent investigations into the causes of rupture of the uterus, show that fatty degeneration of its walls is a most frequent if not a constant coincidence.

Fatty degeneration of the uterus, then, is a cause of death in childbirth by producing or disposing to rupture of the organ; but it has yet another influence on these deaths, for we may be quite sure that a degree of degeneration must often exist, not sufficient to lead to rupture, but quite sufficient to render the organ incompetent to vigorous muscular contraction, even under the influence of unusual stimuli. And thus it may become the cause of those hopelessly tedious labours which, as we have seen, assist to swell the death-rate in parturition.

Thus do we see that among the rational causes of dread that haunt the brain of the accoucheur during a tedious labour, fatty degeneration of the uterus has a right to stand foremost. In PNEUMONIA, Dr. Wilks says that, in "far the majority of fatal cases, some pre-existing and more chronic disease is found in some organ of the body."—(*Ibid.*)

Speaking of the probabilities of death in pneumonia, Dr. Walshe has made the following very well-considered observation—

"There are certain other circumstances (besides treatment) beyond the control of the physician, which exercise a most indubitable influence on the issue (of pneumonia). Among these, the pre-existence of organic disease and the state of health generally of the individual hold an important place. But of all the collateral conditions, age is the most important. While at the two extremities of life, in the new-born infant and in the octogenarian, pneumonia is almost inevitably fatal, the mortality between the ages of six and twelve years scarcely exceeds two-and-a-half per cent."—(Manual of Diseases of the Chest, 2nd edition, p. 438.)

To this I would add, that the octogenarian may be represented at any period of life by the subject of fatty degeneration.

I need hardly remind you that, in protracted Chronic Bronchitis there is no one feature more fearfully prognostic of a fatal issue, sooner or later, than the co-existence of fatty degeneration.

In speaking of Senile Gangrene, Mr. Paget says it occurs, "as its name implies, in the old, and often in those who are old in structure, rather than in years; it affects tissues already degenerate. . . . . I think that, in some cases, its beginning may be when the progressive degeneration of the part has arrived at death. But, if this do not happen, some injury or disease, even a trivial one, kills that which was nearly already dead, as a severe injury might kill any part, however actively alive."—(Ibid., Vol. I., p. 461.)

Under the heads of LIVER-DISEASE, JAUNDICE, and KID-NEY-DISEASE, I may mention some interesting cases examined by Rokitansky. (On Fatal Steatosis—Fatty Degeneration—of the Liver and Kidneys.)

"The cases referred to consisted in steatosis of the liver, accompanied by a high degree of steatosis of the kidneys. Their importance rests upon the possibility of proving them to be parallel to the cases of acute atrophy of the liver, and the analogous renal affection which co-exists with it.

"It is evident that in our cases we have not to deal with that steatosis of the liver which occurs so commonly in the course of consuming suppurative processes, but with fatty livers, as they not rarely develop themselves to a high degree, at the side of an abundant formation of fat in the areolar tissue, without the disease being always attributable to gross feeding.

"There exists thus a steatosis of the liver, occurring in individuals inclined to the formation of fat, to which sooner or later a steatosis of the kidney is added, both which diseases attain slowly and imperceptibly so high a degree that, finally, a cessation of the biliary and urinary secretion supervenes, and, after a slight degree of icterus, death rapidly sets in from anæmia and a hæmorrhagic decomposition of the blood."—(Ranking's Abstract, Vol. XXXI., p. 40.)

Gentlemen, I might prolong this subject much further—so widely spread is the influence of this seemingly spontaneous atrophy, this vestige of disease. I might give you quotations from reliable authorities to show in how many more diseases than I have yet referred to, the fatality, is determined by fatty degeneration; but I should exhaust your patience and overstep our time. I will, therefore,

conclude the list by simply enumerating the causes to which death was attributed in sixty-eight cases analysed by Dr. Quain, in all of which there was fatty degeneration.—
(Med. Chir. Transactions. 1850.)

LIST OF CAUSES TO WHICH DEATH WAS ATTRIBUTED IN SIXTY-EIGHT CASES OF FATTY DEGENERATION.

Rupture of the heart.
Exhaustion.
Coma.
Pleuropneumonia.
Syncope anginosa.
Cardiac apoplexy.
Syncope.
Cancrum oris.
Gradual decay.
Lethargie.
Hæmorrhage into the pericardium.
Cerebral hæmorrhage.
Diarrhæa.
Gangrene of the intestines.

I must briefly remind you, in the words of Mr. Paget, that "Fatty degeneration of the heart often introduces unexpected dangers into the ordinary practice of surgery. It is, I believe, not rarely the cause of sudden death after operations. It is one of the conditions in which chloroform should be administered with more than ordinary caution. They who labour under it may be fit for all the ordinary events of a calm and quiet life, but they are unable to resist the storm of sickness, an accident, or an operation."—(Ibid., Vol. 1., p. 129.)

To sum up, Gentlemen, we have seen that this vestige of disease, fatty degeneration, may claim as its victims a certain number of deaths out of each of the following headings of the Registrar's report:—

Apoplexy .			26.0	deaths per	week.
Paralysis .			23.7	"	,,
Delirium tremen	s aı	nd			
intemperance			4.5	,,	,,
Angina pectoris	s aı	nd			
other diseases	of th	he			
heart .			31.9	,,	,,
Pneumonia.			60.3	,,	,,
Diarrhœa .			17.5	,,	,,
Mortification			4.6	,,	,,
Influenza .			3.1	,,	,,
Peritonitis .			4.4	,,	,,
Childbirth .			4.3	,,	,,
Bronchitis .			64.5	,,	,,
Jaundice .			2.8	,,	,,
Liver disease			11.0	,,	,,
Kidney disease			4.9	,,	,,

263.5 deaths per week;

out of which fatty degeneration claims so large a share.

## ABNORMAL PHYSIOLOGICAL STATES.

We pass on now, Gentlemen, to another and most important branch of our subject.

I must now speak more at length of those states of health to which I referred in my second Lecture, when I said that I should impress upon you that "they are intimately related to the definitely marked diseases"—those states so puzzling to the young practitioner, because they do not fall under any of the nosological headings which have been his landmarks in the study of disease—those states which, although perhaps familiar in their aspect to most old practitioners, are, nevertheless, most inefficiently treated, or not treated at all, because their interpretation is so little understood, and because their importance is not appreciated—states which are not recognised as disease, but which certainly are not health.

A family of four children were exposed to the infection of measles at the same time, and from the same source; all of them were supposed to be in health at the time. One had the measles simply and slightly; one had a severe attack of pneumonia combined with it; one indicated a disposition to typhoid symptoms, and was completely oppressed by the morbid poison; a fourth lingered in its convalescence, and was found to have become the subject of an eruption of tubercles in the lungs.

A party of friends, all apparently in what is called health, met at a funeral; they went together into a damp unwarmed cemetery chapel, on a raw winter's day, and returned together, one and all complaining that they had taken "a severe chill." They dined together and went to their homes. One suffered an attack of rheumatic fever; one had anasarca; one jaundice; another bronchitis; a fifth pneumonia; a sixth diarrhœa; a seventh had erysipelas; and another had pleurisy. One coughed up a quantity of blood; while the rest got a restless night, and a cold in the head, and thought no more about it. These are no imaginary stories; analagous cases frequently occur within the experience of medical men in large practice. But what is their interpretation? Why did the

same cause—the chill—produce such different effects, under external conditions, apparently the same? No doubt, the first answer which suggests itself is, that the circumstances, so apparently the same, must have been really Well, Gentlemen, I will admit that, as the first step towards solving the difficulty. But I will insist that in a given case, the cause, so far as the chill is concerned, shall be the same in each individual; and the circumstances, so far as they are external, shall be the same for each individual. Nevertheless, these different effects shall be produced; and the reason we shall find to be this, that there are other causes and other circumstances, different in each case, existing within the organisms of the sufferers, with which the one cause—the chill—has to combine in producing its effect, and that the effect is the result of this combination of causes, different in each individual.

Some of you, perhaps, will say that this simply means that the different effects are explained by the different idiosyncrasies of the individual. And I must warn you against adopting a word as the explanation of a difficulty, lest in doing so, you fail to investigate the multitude of facts which that word may represent. I have no objection to the word, so long as you bear in mind that you must be able to explain what it means, if it is to be accepted as any explanation at all. Professor Bernard has spoken well on this subject: "I discovered," he says, "that section of large divisions of the sympathetic nerve was apparently unattended with the slightest inconvenience, as long as the health of these animals (rabbits) remained perfect . . . . but as soon as a general debilitation of the system arose from want of proper nourishment, acute inflammation was produced in the organs deprived on nervous influence. We had, therefore, succeeded if

artificially creating particular idiosyncrasies in these animals, and could predict with certainty that, as soon as health failed, disease would arise at a given point.

.... Morbid predispositions must, therefore, be viewed in the light of peculiar physiological conditions." And he concludes thus: "Let me advise you not to consider idiosyncrasies in the light of mysterious powers residing within the depths of our organs, nor as entirely novel functions superadded, as it were, to those which already exist. They must be viewed as mere manifestations of the ordinary laws of physiology."—(Bernard, Lecture V., Medical Times and Gazette.)

Health, Gentlemen, is the normal physiological state; and peculiarities in, or divergencies from this condition must be regarded as greater or less degradations of health, in proportion as they predispose to contingencies which increase the probabilities of death before the normal term—before the attainment of the ultimatum.

There are very few persons who pass through life in the normal physiological state. At some period of life almost every individual diverges, more or less, from this state in one direction or another; and during that divergence, although escaping an attack of what is recognised as disease, he certainly is not in health; and in almost every individual there is a tendency to diverge in some particular direction, during which divergence—i. e., during that period of degraded health—he is particularly prone to certain classes of disease. In the case I have taken as an example of the effects of chill, for instance:-The man who had rheumatic fever, was already surcharged with uric acid. He who had jaundice was suffering from defective excretion by the liver, requiring only a certain increase in the defect, or of the demands upon the secreting function, to throw the secretion back into the blood.

The patient who had anasarca was suffering either from hydræmia, or from defect in the excreting powers of the kidneys. The sufferer from bronchitis, I attended myself: he had chronic congestion of the bronchi, from repeated former attacks of bronchitis, and the circulation through his chest was defective, through a feeble degenerated heart; but he had been accustomed to pass as a man in health, competent to perform the onerous duties of a taxcollector and county-court agent. He who had an attack of diarrhœa, found in his bile ducts or in the intestinal mucous membrane a safety valve, by which he was saved from either anasarca, jaundice, or rheumatic fever. The subjects of erysipelas and pneumonia, and the man who had hæmoptysis, were already suffering from depraved states of the blood, or of the organs to which it was determined; and from which, in the last two cases, it escaped in different quantities; while the patient who was attacked with pleurisy, was surcharged with urea from defective action of his kidneys.

These several persons, therefore, were suffering, when they considered themselves in health—before the occurrence of the chill—from physiological conditions to which we can attribute the particular form of the disease which was set up, by the addition, in each case, of one and the same cause, viz., the chill.—But some escaped unhurt! Because in them, the physiological state was sufficiently normal, that a resisting and reactionary power existed, which was competent speedily to restore the functions of the organs subjected to the shock of the chill, and to make them compensate for the temporary arrest by increased activity.

I have chosen this group of somewhat crude examples on purpose, that their meaning may be the more perspicuous. They, most of them, exemplify states of health edpendent on the *fluids and excretory organs* of the body. It would be easy to bring many examples of degraded health consequent on disease attributable to the *nervous* system.

\* \* \* \* \* \* \*

I assume, then, Gentlemen, that I have sufficiently demonstrated that the vestiges of disease stand first among the causes of death.

I have shown that so long as these vestiges exist, they are causes of defect in the vital force and thus act as factors of the essential and of the predisposing causes of fresh attacks of disease.

I have shown that the diseases, from which these vestiges result, are but the manifestations of pre-existent physiological states, to which, by some means, the last condition has been supplied, necessary to complete the conditions of existence proper to the disease which then is developed in its characteristic features.

I have shown that these ABNORMAL PHYSIOLOGICAL STATES are indicated by the various conditions of impaired, general health, "conditions not recognised as disease, but which certainly are not health."

The sum of it is this :-

- 1. The majority of diseases which we see excited by the various accidents of life are but the manifestations of pre-existent physiological states, which required only this last condition (the accident of life) to complete their development into the characteristic features of disease.
- 2. Those conditions "not recognised as disease, but which certainly are not health," are the faint expressions of these morbid physiological states, while still deficient in the condition necessary to complete their development into the recognised features of disease.
- 3. The multifarious and anomalous functional derangements which puzzle the physician, and make martyrs of

the patients, depend, for the most part, upon the influence exerted by these morbid physiological states over the ordinary incidents of animal existence, which are thereby modified, coloured, and distorted.

4. During the whole of the time that the physiological conditions are disturbed, there is a greater or less defect in the vital force, and this defect, therefore, exists at a period anterior to what are usually understood as structural changes.

I have shown that these abnormal physiological states, recognisable under various forms of impaired health and attended by a legion of anomalous symptoms, may be traced back to still earlier periods in their history, when they require the greatest vigilance of the physician to detect any deviation from the standard of normal health.

Then I endeavoured to show in what direction we must look for the causes of these earliest and most occult deviations from the normal physiological state.

With this intention I pointed out that the conditions of life and the vestiges of disease have a direct influence on the vital force; that these alterations in it are transmissible to the germs of a succeeding generation; and that thus defects of force may be due to abnormal conditions of life in the individual, and to the vestiges, or vestiges of vestiges of disease in an ancestor. And I wish to draw your marked attention in this place to the fact, that it is to these defects transmitted from an ancestor, and to the conditions of life in the individual, that we must especially look for the causes of those first, insidious deviations from normal health.

I endeavoured to show that these incipient and insidious degradations of the vital force exist at a period anterior to such changes as are understood by the terms structural and organic; and I think you will now understand why I said

in my last Lecture, that there could be no fact in Medicine of greater practical importance than this. My reason was —that the very dependence upon the conditions of life which exposes the force in the individual to degradation under unfavourable conditions, must render it amenable to elevation under the influence of conditions which are favourable. And thus we learn that there is a possibility of cure in states that would otherwise be beyond the reach of treatment.

# CHAPTER IX.

DISINFECTION—MEDIA BY WHICH THE CATCHING DISEASES ARE COMMUNICATED—MEANS FOR PREVENTING INFECTION AND CONTAGION—DIRECTIONS FOR THE SICK ROOM AND ATTENDANTS—PRECAUTIONS TO BE USED BY DOCTORS—MODE OF CLEANSING APARTMENTS AFTER ILLNESS—PROPER METHODS OF USING CHLORIDE OF LIME, CARBOLIC ACID AND CONDY'S FLUID.

Infectious and Contagious (catching) diseases, Scarlet fever, Small-pox, Measles, Typhus, Relapsing fever, Puerperal fever, are communicated principally by the following media:—

- 1. The breath.
- 2. The secretions and excretions.
  - a. The saliva.
  - b. The perspiration.
  - c. The urine.
  - d. The evacuations from the bowels.
  - e. The sputa.
- 3. The skin of the body; especially the dead peeling skin and the powdered skin which adhere to clothing and are wafted about in the air.
  - 4. The body-clothes.
  - 5. The bed-clothing.
  - 6. Discharges from eruptions, abscesses, wounds, &c.
  - 7. The hair.
  - 8. The walls, floors, and furniture of apartments.

It is probable that no infectious or contagious disease would spread if all these media were promptly disinfected.

The difficulty lies, not in disinfecting any one of them when it is specially submitted to us, for that is easily done, but in keeping up such a rigorous system of disinfection around the sick person, that none of these media shall escape disinfection; for common sense tells us that if only one escapes the protection from infection may be lost. It is evident that the disinfection of the secretions and excretions must be performed as they leave the body, so that the air may not be infected during their passage. For this reason the disinfectants put into spitting pots, urinals, and bed pans should be Volatile, like Chlorine and Carbolic Acid, so that an atmosphere of the disinfectant may rise from the utensils while they are being used. I therefore, advise for these purposes Chloride of Lime and water, or Carbolic Acid, and they have the advantage of continuing to give off their disinfecting principles after the contents of the utensils are thrown down the drains.

For most other purposes Condy's fluid will suffice, it is clean, free from offensive taste or smell and not poisonous. It is not volatile but it disinfects whatever comes in contact with it. The air of a room therefore may be disinfected to a great extent by freely dispersing Condy's fluid with a spray producer. Clothes can be disinfected by being soaked in it. The skin can be disinfected by being washed with it. Sponges can be disinfected by being soaked in it, but sponges are dangerous spreaders of infection and it is much better to use pieces of cloth, which can be burnt, and in cleansing wounds to irrigate them with a syringe. Linen and other washing materials can be disinfected by boiling, but they should be previously put into Condy's fluid or diluted Carbolic Acid directly they are done with and kept in it for about two hours, otherwise they may spread infection before they are boiled. For all these purposes Carbolic Acid is most efficient but its smell is objectionable.

Whatever is not susceptible of boiling but will bear baking can be disinfected by being submitted to a temperature of from 212° to 250° in an oven, and for greater safety a little Carbolic Acid should be put into the oven at the time.

1. When a catching disease occurs in a house the first thing to be done is—to select a room for the patient as much isolated as possible from the rest of the house—a room through which no one has to pass and by which there are as few passers as possible—the top of the house is best as a rule. The room should have a window opening directly into the fresh air and an open chimney and fire-place, and it should be supplied with not less than 1000 cubic feet of fresh air every hour for each occupant whether patient or attendant.\* Whenever practicable a second room should be set apart near the patient's room for the attendants to cook and take meals in and to sleep in when off duty.

2. The second thing to be done is—to remove from the room or rooms selected every thing that cannot be washed, boiled, baked, or burnt, and then to hang over the outside of each doorway a curtain kept constantly wet

with Condy's fluid.

3. The third thing is—to decide who is to attend upon the patient. There should always be two persons and no more. They should cover their hair with washing caps, dress in washing clothes, and not associate with the rest of the household or with any other persons. They should each take at least six hours sleep out of the 24. They should each walk out in the fresh air 1 hour per day. They should wash their hands and faces with Condy's fluid and water when they leave the patient's room. They should avoid inhaling the breath or exhalations of the

\* See Ventilation and Heating.

patient and they should neither eat nor drink in the sick-room.

- 4. The fourth thing is—to place in a corner of the patient's room a large glazed pan containing water and Carbolic Acid or Condy's fluid. Into this pan everything that can be washed or boiled should be put, directly it is done with, before it leaves the sick-room. The pan should be emptied night and morning or oftener.
- 5. The patient should be sponged all over with warm water and Calvert's Carbolic Acid Soap twice a day (except during periods in which it may be thought unadvisable by the doctor) and if the skin is peeling or powdery it should be anointed after the sponging with olive oil containing a little Carbolic Acid (about 10 per cent.)
- 6. Chloride of Lime and Water or Carbolic Acid should be put into every utensil before it is used by the patient and after being used the contents should be immediately thrown down the drain together with some fresh Chloride of Lime.
- 7. The patient's teeth and mouth should be washed with Condy's fluid, or Calvert's Carbolic Acid Soap and water several times a day and when there is any discharge from the nose or other passages these should be cleansed in the same way. When the discharges from the body are feetid Savory and Moore's Carbolic Acid Vaporiser should be burnt in the room.
- 8. It must constantly be borne in mind that no amount or kind of disinfection will take the place of fresh air, and therefore the sick room must be kept freely ventilated, carefully avoiding draughts.\* Gas should not be burnt in a sick room.
- 9. The doctor should not communicate with any other members of the household after he leaves the sick-room

<sup>\*</sup> See Ventilation and Heating.

and he should always be provided with a basin of water, Calvert's Carbolic Acid Soap, a nail brush, and a clean towel with which to wash his hands on leaving.

Let it be especially remembered that the main points are the prompt disinfection of the infecting media enumerated, in the beginning of this Chapter, and the free circulation of fresh air.

After the acute specific disease has run its course, infection may last as long as there are any unhealthy discharges left by the disease, as long as there is any peeling or powdering of skin, shedding of hair and the like; and precautions are needed for disinfecting these means of spreading disease so long as they exist.

No doctor who has been in attendance upon Erysipelas, Scarlet fever, Puerperal fever, Typhus, Gangrene, Pyæmia, or who has been engaged in a post-mortem examination should enter a lying-in-room until he has subjected himself to effectual purification and disinfection; and it is a safe rule, under all circumstances, for a doctor not to touch a lying-in patient until he has washed his hands with some disinfectant.

The difficulties of carrying out this rule are very great in the hurry of practice, but every doctor engaged in Obstetric practice is bound to provide for doing so. And he should never go upon his rounds without some disinfectant with him to mix with the water in which he washes his hands in case of unexpected need.

To facilitate this important hygienic pecaution I have induced Messrs. Maw, the well-known instrument makers, of 11 Aldersgate Street, and Messrs. Calvert the celebrated manufacturers of Carbolic Acid, to provide a little metal case containing a piece of strong Carbolic Acid Soap, a nail-brush, (for it is useless to wash the hands without brushing out the nails) and a little tube of Carbolic

Acid, for medical men to carry constantly in their pockets when on their rounds. The whole is no larger than a small snuff-box. Directions for use are on the inside of the lid. It ought to be used by nurses and visitors upon the sick as well as by doctors.

When the infectious period of the illness is over and the patient removed from the sick-room—everything which has been used or worn by the patient and attendants having been either disinfected or destroyed as already directed—the room must be thoroughly disinfected in the following manner.

- 1. The doors and windows being shut, the room must be kept full of Carbolic Acid fumes for 12 hours by means of one of Savory and Moore's Vaporisers—Then the windows must be opened for 12 hours before cleaners and workpeople are admitted.
- 2. The floors, paint, and furniture, must be washed with Water and Chloride of Lime or with Calverts Carbolic Acid Soap. The ceiling and walls must be Lime-washed, after which the walls may be re-papered if required. While this work, is in progress a fire should be kept burning in the grate and the windows open.

The following particulars as to the mode and proportions in which to use the Chloride of Lime, Carbolic Acid, and Condy's fluid, mentioned above, have been kindly furnished by Messrs. Savory and Moore, and by Mr. Calvert.

For fumigation by Chlorides.—Dissolve one pound of the Chloride of Lime in 4 gallons of Water and after stirring the mixture well allow it to settle, pour off the clear liquor and place in shallow dishes or other convenient vessels in the rooms and passages of the House. For use in chamber utensils, bed-pans, spittoons, &c., and where a volatile

disinfectant is required, a wineglassful of the above solution should be added to a pint of water.

Carbolic Acid.—When required for the above purposes, that prepared "for Disinfecting" should be obtained—one ounce of this added to a quart of Water is sufficiently strong to purify drains, water-closets, &c. The mixture should be vigorously stirred or shaken (as Carbolic Acid is not readily miscible with water). Half a pint of this solution put into any utensil used in the sick-room will disinfect the contents and enable them to be preserved for the inspection of the medical men without any unpleasant effluvia arising.

Carbolic Acid Oil for anointing patients should consist of Olive Oil and about 10 per cent of Carbolic Acid.

Condy's fluid.—For use either with the spray producer to disinfect rooms, for cleaning sponges or bandages, washing the skin of the patient, the personal use of Nurses or Medical Attendants, dressing wounds, &c. Condy's fluid should be mixed with water in the proportion of a teaspoonful to a pint. Linen and Bed-clothes should be immersed in a solution made by mixing a wineglassful in a pailful of Water, diluted in this proportion it does not stain the linen.

# SPECIAL RECIPES, DIRECTIONS, AND APPLIANCES FOR THE SICK-ROOM.

As this does not profess to be a "cookery-book" only a few Medical-food Recipes will be here given, with some directions for the cooking and appliances of the sick-room which cannot easily be found elsewhere.—For the ordinary kitchen recipes the reader is referred to the cookery-books of Miss Acton, and Mrs. Beeton, and to Cre-fydd's "Family Fare," pp. 240 to 249.

- 1. Farinaceous Articles should all be submitted to a temperature of 212° Faht. (boiling water) to make them digestible.
- 2. BEEF TEA should not be boiled, and should not be strained through a fine sieve or muslin. It should be made as follows. Take of Rumpsteak, free from fat and minced, 1 lb, cold water 1 pint, a pinch of salt. Put them into a jar and tie it down. Place the jar in a saucepan of cold water, raise this water slowly to boiling and keep it slowly boiling for two hours. Remove the jar and strain its contents through a very coarse sieve so that all finely powdered sediment may run through. Then pass a piece of bread over the surface to remove any fat that may float upon it.
- 3. Liebig's Extract of Meat and other similar preparations. It is important to bear in mind that these contain very little, if any, nourishment properly so called, that is to say, they contain no plastic material, no fat, no saccharine matter. Their principal virtues belong to the class of stimulants and blood-tonics. When mixed with water, they are excellent menstrua in which to administer nutritive materials, such as eggs, bread, oatmeal, cornflour, vermicelli; but without such additions they are quite incapable of supporting life for any length of time. Baron Leibig's own writings support this statement. Unless these facts are borne in mind a patient may easily be starved unintentionally.
- 4. White of Egg differs from the yolk principally in containing no fat. On this account it is often better borne by bilious persons. Yolk of Egg contains 29.8 per cent of fat; when the stomach can bear it, therefore, it is a more complete nutriment than the white. But white of Egg beaten up in milk answers every purpose.

Eggs for the sick should be either raw, or very lightly boiled. They should never be given hard.

- 5. When may be made either with Crosse and Blackwell's Essence of Rennet, or with Treacle, or with Sherry, according to circumstances.
  - a. Rennet Whey—see the Directions on Crosse and Blackwell's bottles.
  - b. Treacle Whey—pour three tablespoonfuls of best Treacle into a pint of new milk while it is boiling; see that it boils up once after the treacle is added; set it aside to cool and then strain.
  - c. White Wine Whey is made in the same way as Treacle Whey, 3 glasses of sherry being used to a pint of milk.

Note—When a more nutritious food is desired and suitable, well beat up the curds with the whey instead of straining; see that they are not in lumps.

6. ALL ANIMAL FOODS are apt to lose digestibility by re-

peated cooking.

7. Bread should be either toasted or 24 hours old.

8. Special Restorative. I have found the following to be the most efficient restorative food, and the mixture is agreeable to most palates.

New milk (cold) 4 parts. Beef tea (cold) 2 parts. Brandy (pale) 1 part.

If no other food is taken, about 5 fluid-ounces (a quarter of a pint Imperial) should be given every 2 hours, or half that quantity every hour. It should be sucked out of a syphon Infants-feeding bottle, not drunk out of a spoon or cup. When desirable this food may be gradually thickened by the addition of boiled corn-flour or other farinaceous articles. The flavour may be varied by adding different spices.

9. INVALID Soup. The following Invalid soup has

proved extremely useful in a large number of cases, and since I first published the recipe in 1864 it has been usually kept nicely prepared by Mr. Donges, Confectioner, Gower Street, W.C.

Gravy beef 1 lb., scragg of mutton 1 lb., isinglass 2 oz., vermicelli 3 ozs., mushroom ketchup 3 tablespoonfuls, corns of allspice 24, sage a sprig, cold water 3 quarts; put the isinglass and the meat cut small into the cold water, gradually boil, skim well, and then add the other ingredients; simmer four or five hours till reduced to 1 quart; strain through a fine hair sieve, and carefully remove all fat; add salt to the taste. This may be taken cold as a jelly, or warm as a soup. Calf's-foot may be used instead of isinglass when procurable—and when allowable a little solution of cayenne pepper should be added; and the taste may be varied by the addition of a little Worcester, or other wholesome sauce.

- 10. Combinations of alimentary principles in nearly exact normal proportions.
- a. Flour 4 oz., sugar 1½ oz., suet ¾ oz., milk ¾ pint Imperial, 1 egg.—This will make a good pudding, or it may be given in any other form desired; with the addition of a little cress and salt and water it forms a complete diet, upon a sufficient quantity of which a person can live healthfully for an indefinite length of time without any other food.
- b. The same may be said of the following.—Rice 3 oz., sugar 1 oz., 2 eggs, butter  $\frac{1}{2}$  oz., milk  $\frac{3}{4}$  pint (Imperial), water as much as is sufficient to boil the rice in.
- 11. Port Wine Jelly. Take of port wine 1 pint, isinglass 1 oz., sugar 1 oz.; put the isinglass and sugar into 4 pint of water, warm till all is dissolved, then add the wine, strain through muslin and set to jelly. (An excellent way of giving port wine.)

12. SUET AND MILK. Put a tablespoonful of shredded beef-suet into ½ pint of fresh milk, warm it sufficiently to completely melt the suet, then skim it, pour it into a warm glass or cup, and drink it before it cools. If there is any difficulty in digesting the suet add 10 gr. of Pancreatine powder. (Savory and Moore's.)

13. MILK WITH RUM, BRANDY, OR WHISKEY. Put I table-spoonful of Rum, Brandy, or Whiskey, into half a pint of new milk, and mix well by pouring several times from one vessel to another. "Bilious" persons should heat the rum before adding it to the milk.

14. Ardent Spirits and their Peculiarities. Brandy makes people nervous, Gin weakens, Rum and Whiskey produce biliousness, Whiskey is the least objectionable. None should be taken neat, and when mixed with water or other fluids they should be well incorporated by pouring from one vessel to another. When Whiskey produces biliousness, Rum, which has been made hot and then cooled, may be tried instead.—(See Alcohol.)

15. NUTRITIVE ENEMATA. When nutriment is given in enemata the quantity should not exceed 2 to 4 oz., and the temperature should be about 80°.

The bowel should be first washed out with half a pint of warm water.—An elastic bottle holding the required quantity is better for nutritive enemata than the ordinary enema syringe. They should be given while the patient is lying on the back with the hips raised on a pillow.

16. Poultices. Linseed poultices should be made by filling a muslin bag with crushed linseed, (not linseed meal) then putting it into a basin or dish and pouring boiling water upon it.—When thoroughly soaked it should be squeezed between towels till no water drips from it. The same poultice may be made hot four or five times by pouring fresh boiling water upon it. Bread poultices should be

made of finely crumbled bread treated in the same way as the linseed.

- 17. Fluids for Inhalations should have a temperature of about 170° Faht.
- 18. WARM BATHS should not exceed a temperature of 98° without medical orders.
- 19. The LIGHTS IN A SICK-ROOM should always be placed behind the patient not in front. Gas should not be used at all. (See "Ventilation.") Candles are better than lamps.

20. A Nursing Schedule should be used in all Fevers, Inflammations, Surgical Cases, and other Acute Illnesses.

Those who have been called upon to bear the responsibility of the Nursing in a severe case of Fever, Inflammation, Surgical Operation, or the like, will have a vivid recollection of the difficulties they encountered in carrying out the various Medical Orders upon the punctual attendance to which, through anxious days and nights, the patient's life depended.

How to arrange the hours for Nutriment, for Medicine, for Wine, for Brandy, for dressing wounds, for external applications, for Sleep, &c., &c., without one important item clashing with another, is often a most puzzling question, requiring the details to be carefully considered and arranged in writing, before it is practicable to work them out.

Not unfrequently the orders left by the doctor are misunderstood, or found to be impracticable when they come to be put together and arranged according to time—the hours for dressing wounds or taking Medicine clashing with those for Food, the hours for Wine coming upon those for Brandy, or interfering with the necessary sleep, and so forth. When the doctor has gone and, perhaps, as in country houses, cannot possibly be consulted again for many hours, it is difficult to overstate the distress of friends and nurses on discovering that orders, which have been impressed upon them with all the weight of questions of life and death, cannot be implicitly carried out from want of a consistent arrangement of their details.

Happily for the sick, and for the doctors, nursing is now passing into the hands of educated Ladies and well trained Nurses, competent to understand and to perform with intelligence that systematic nursing to which the enlightened practical Medicine of the present day attaches such vital importance. And the introduction of the Clinical Thermometer and similar appliances into the sick room, makes it necessary in private houses (where there are no clinical assistants or house-surgeons as in hospitals) to entrust the nurse with the task of keeping a register of the temperature, the rate of pulse and respiration, &c., at hours when the doctor cannot be in attendance.

The doctor who believes in the importance of his own orders will be strict in requiring at each visit an accurate report of how they have been obeyed. The nurse who intends to obey them faithfully will not be satisfied unless she sees that they are both intelligible and practicable before the doctor leaves the house, and she should write them down directly he is gone.\*

\* To meet the necessity thus arising in the sick room of the present day and to assist, at once, the patient, the doctor and the nurse, I have arranged "The Nursing Schedule," and in order that it may be widely useful it has been brought out by Mr. H. K. Lewis, 136 Gower Street, at as low a price as the expenses of paper, printing and publishing, will permit.

Each sheet is arranged for the report of two days and two nights and contains directions for use. A doctor wishing the Nursing Schedule to be used in any particular case, has only to order it on his prescription, and the Chemist will supply it with the medicines.

# INDEX.

PAGE	PAGE
Acid, wines and other fluids	Bowels, regulation of 35
with least and most 63	Bread 142
Air, fresh, conditions for supply	Bread diet, not sufficient to
of to another aut	Diead diet, not sumcient to
of to apartments	maintain health 61
Albuminoids, increase of, in the	and typhus fever . 59
diet, in Bright's disease 83	Breakfast, the second best meal
Alcohol, table, (notice) 8	of the day 37
- table showing the one	Bronchitis, vestiges of, the causes
lysis of spirituous liquors	of other discourses
Tysis of spirituous inquors . 100	of other diseases 105
absolute, and proof	death from, in rickets 115
spirit 64	
Anstie on, (notice) . 7	
——— fasts	
fasts	0 1 11 12 2 2 4
body as food 7	Carbolic acid, mode of using for
it le - c' - c'	Carbon, heat of combustion of,
its place in diet 6	Carbon, heat of combustion of
——— in disease, effects . 66	with hydrogen
acute 66	with hydrogen
	Change and rest
"Leading article from	see Posture.
Lancet"	Childbirth, death from rickets in 115
Anomia a fraguent resting of	Chill, severe, case of different
Anæmia, a frequent vestige of	sequelæ from, in different per-
severe diseases, and names of	sons 197 190 20
diseases under which it proves	Chloride of Lime, modes of using
fatal	for disinfection
- connection with rick-	for disinfection 139
ets 116-17	Cider, analysis of 65
Anstie on alcohol, (notice) 7	Climate of United Kingdom,
Apoplexy, relation of changes in	rules for promoting and main-
blood was all to	taining vigorous health in 12-47
blood vessels to 120	Clothing and protection from
vestiges of, the causes	cold protocolon from
of other diseases 109	Cold, protection from, by cloth-
Arteries, atheromatous disease of 120	Cold, protection from, by cloth-
Atrophy, vestiges of, the causes	ing
of other diseases 110	Colds, per centage of causes of . 15
	Collins (R) deaths in childbirth
Banting's diet, (approximate ana-	from fatty degeneration of
lysis)	Uterus
Bantingism, good and harm	Condy's fluid, mode of using for
from, as a system of diet . 74	diginfaction
Bathing 24	disinfection . 140
Batha " cold friction " de il	disinfection
Bathing Baths, "cold friction" daily use	carbo-hydrate 87
	- albuminoid co
— warm	fluid diet . 89
Beef Tea 141	Contagion, see Infection.
Bernard, experiments showing	Cotton-wool, use of as a filter
morbid predispositions to be	and recovered as a miter
peculiar physiological condi-	"Cough (winter) Lectures on,"
	Cough (winter) Lectures on,"
tions	notice 15

76 66 134
66 134 3 9
66 134 3 9
134 3 9
134 3 9
134 3 9
3 9
3 9
3 9
9
9
21_0
34-9 5-19
-19
23
20
90
38
70
10
76
195
07
91
95
10
10
-
19
81
OI
-
5
5
5
32
32 141
32 141 144
32 141 144
32 141 144
32 141 144
32 141 144 92-9
32 141 144 92-9 31
32 141 144 92-9 31
32 141 144 92-9 31
32 141 144 92.9 31
32 141 144 92.9 31
32 141 144 92-9 31
32 141 144 92-9 31
32 141 144 92-9 31
32 141 144 92-9 31
32 141 144 92.9 31

### INDEX.

DACE	PAGE
PAGE	Heart and Pericardium, vestiges
Fat in the body, its character	
varied by food	of diseases of, causes of other
- importance of proper food	diseases
to supply the adipose tissue	Heat, food and motion . 5, 48
with	British unit of, mechanical
- see Pancreatic Emulsion.	equivalent of
getting, or thin 5, 73	heat-equivalents of food . 49
solid and liquid, importance	evolution of, food required
of distinguishing between . 70	for 50
	for
Fatty degeneration, as a vestige	evolved from food, mode in
of disease	which disposed of 50
supposed	Heating and ventilation 12
causes of death in cases of 125-6	Holland (Sir H.) on Symptoma-
Fermented liquors, amount	tic complaints (from "Medi-
which may be taken daily	cal Notes") 95
with advantage 8-9	cal Notes")
with advantage 8-9 Fevers, contagious, media by	giving elementain diagrages ve
which communicated 194	giving elements in diseases re-
which communicated 134	sembling 79
continued, vestiges of,	T
the causes of other diseases , 108	Indigestion common among
—— see Typhus, Typhoid.	those who live an in-door life 39
see Rheumatic Fever.	Infection, means of preventing . 135
Fluids, amount of acidity in . 63	directions for the sick-
Fogs and damp air as causes of	room and attendants to pre-
colds 18	vent
colds	vent
Food, heat and motion . 5, 48	precautions to be used
— connection with force . 5	by medical men 138
— heat equivalents of, 49	modes of preventing
mode in which heat evolved	after illness
from is disposed of 50	Inhalation, fluids for 145
- its precedence anong means	Interdependence of diseases . 1
of preserving health 6	. I
- table of the composition of	Tally part wine
some of the principal articles	Jelly, port wine
of	Jenner (Sir W.) Lecture on
of	Rickets," extracts from 114
see Diet.	
Foot-pound, its mechanical force 49	Kirkes, on fibrinous obstruction
Foot-tons, mechanical equivalent	of cerebral arteries 120
of heat-units in 49	
	Laryngismus stridulus, connec-
Gangrene, senile, in tissues al-	tion with rickets
ready degenerate 123	Liphic's outpool of
Germ theory of disease 21	tion with rickets
Goloshes, use of 15	Lights in the sick-room Liquors, see Fermented, Malt,
Bout, antecedents of, and rela-	Liquors, see Fermented, Malt,
tions with constitutional dis-	Spirituous.
orders 05 6	"London Noise and London
mostings of the same of	Sleep" Letter on by Dr. Dobell 23
orders	- Loader from the
other diseases 106-8	
aulliver, atheromatous disease	Times" on ditto 26
of arteries 120	enumeration of special
T 11/	night noises 24-5
Health more required by modern	preventible ones 30
Health more required by modern	
men at their occupations than	Maclaren "on Physical educa-
strength	tion" (notice of) 9  Malt 1: ditto, extracts from . 32
rules for promoting and	ditto, extracts from . 32
rules for promoting and maintaining in the climate of	
the United Kingdom . 12-47	
	Meals

Meals, proper hours for 9, 38	Physiological conditions not re-
modern regulations of busi-	cognised as disease, and yet
ness establishments relative	not health
to, inconsistent with healthy	not health  possibility of cure in ditto  not health  possibility of cure in ditto  not health  possibility of cure in ditto
digestion 39-40	in ditto
regulations for improving	Pneumonia, a cause of death in
the times of, connected with	rickets
the obligations of business . 44	fatality in from other
light, importance of during	Chronic disease
business hours 45	Posture Jelly
Measles, variety in attacks, from	Tosture
the same source 127	Posture
Meat, rules in loss of power of	
digesting 83	
Liebig's extract of, 141	Recipes, &c., special, for the
Medical Men, the duties of, 91-3	sick-room 140-6
Medicine, departments of, which	Regimen 9
are not specialities 2-3	Regulation of habits 9
rational medicine and	Respiration, amount of air vitiat-
pseudo-medical dogmas 10	ed by, per hour
Milk, use in diarrhœa 83	Respirator, the only protection from damp fogs, &c 19
— and Suet 144 — with Spirit	Rest and Change
Motion, heat and food . 5, 48	— see Sleep
"Mulder's Chemistry of Wine,"	Rest for diseased organs, to be
extract from 64	obtained by removing elements
	from diet which increase their
	functions
Nervous complaints, increase of	Restorative food
from modern excessive compe-	Rheumatic fever, case of, and
tition in commerce and the pro-	probable sequelæ from want of
fessions	precautions
Nursing schodule passesity for 145	causes of other diseases 103
Nursing schedule, necessity for . 145 Nutrition, healthy, necessity of	Rheumatism, vestiges of, the
preserving by diet in disease 77-8	causes of other diseases . 107-8
preserving by drev in discusse 77 o	Rickets, causes of death in . 114
Olein, its relations to solid and	
liquid fatty bodies, 71	116-17
Ormerod on fatty degeneration of	diseases in which they
the heart	are primarily or secondarily
Oxygen, mean consumption of by	the cause of death 114-17
adults 49	Rokitansky, cases of fatty degen- eration of Liver and Kidneys . 124
	eration of liver and Kidneys . 121
Paret (I) namente en fattu de	
Paget (J.) remarks on fatty degeneration, &c 120-2	Schreeder, cotton-wool as a filter 21
Pancreatic emulsion 82	Schwann, putrefaction of meat . 21
Paralysis, vestiges of, the causes	Sickness, see Disease
of other diseases 109	Sick-room, special recipes, direc-
Periodical examinations, import-	tions and appliances for, . 140-6
ance of, for the prevention of	Sleep
disease 92-9	III LIOILGON 4
"Physical education" extracts	
from Maclaren's work on . 32	Specialities, narrow, in Medicine 2 Spirits, table of analyses of . 65
Physiological States (abnormal) 126-33.	ardent, their peculiarities 144
120-00.	tertitoria, orrore February

PAGE	PAGE
Spirituous liquors, acid and sugar in	Uterus, fatty degeneration of, deaths in childbirth from . 122
taken, &c	Ventilation and Heating 12
Tables of the composition of articles of food	Walshe, fatality of pneumonia from pre-existent organic disease

# CORRIGENDA.

Page 1 line 5 for interdependance read interdependence.
— 10 — 8 dele comma between similibus and curantur.
— 64 — 21 insert heading of "Alcohol"
—— 112 —— 9 insert the word registered between are and in.
—— 127 —— 32 for analogous read analogous.
—— 128 —— 33 for on, read of.
—— 128 —— 34 for if, read in.

# OTHER WORKS BY DR. DOBELL.

Practitioners' and Students' Guide to Physical Diagnosis.

COLOURED PLATES.

# DEMONSTRATIONS OF DISEASES IN

THE CHEST, AND THEIR PHYSICAL DIAGNOSIS; A Treatise on the Acoustics of Auscultation and Percussion—Directions for Educating the Ear by Experiments on Inanimate Bodies—Instructions for Practice at the Bedside—and Coloured Plates of thirty-six Pathological Conditions. Each Plate accompanied by a Statement of the Physical Signs to be observed during Life, and by a Corollary upon its leading points.

# LECTURES ON WINTER COUGH

(CATARRH, BRONCHITIS, EMPHYSEMA, ASTHMA).

"The chapter on treatment is of great practical interest. We think this book calculated to make us more thoughtful and more successful in regard to a most important class of diseases."—The Lancet.

# REPORT ON THE EXPERIENCE OF

MEDICAL MEN WHO HAVE USED "PANCREATIC EMULSION OF FAT." 1867. Issued in reply to the following Editorial note which appeared in the Lancet, Nov. 17, 1866:—

- "It would be interesting to know to what extent the experience of those medical men who have used the 'pancreatic emulsion' of fat accords with that which Dr. Dobell has now recorded in our columns in about two hundred cases of consumption . . . The remedy has been placed fairly before the profession, the author reserving to himself no pecuniary interest or advantage whatever; and if the utility of the remedy bears any proportion to the large demand which has arisen for it, we ought to know this clearly, and to understand and appreciate our obligatioes."—Lancet.
- "The pancreatic preparations are in high favour. . . . To have introduced them is a good deed"—MED. TIMES AND GAZETTE.
- "Pancreatic Emulsion, as every one now knows, was originated by Dr. Dobell."—Dublin Quarterly Journal of Med. Science, (Nov. 1867.)

# ON THE TRUE FIRST STAGE OF CON-

SUMPTION: Its Nature Diagnosis, and Treatment by Diet, Climate, Medicine, Etc.

# ON TUBERCULOSIS: ITS NATURE, CAUSES,

AND TREATMENT. With Notes on Pancreatic Juice and Pancreatic Emulsion. Second Edition.

# REPORTS ON THE PROGRESS OF PRAC-

TICAL AND SCIENTIFIC MEDICINE IN DIFFERENT PARTS OF THE WORLD. Edited by HORACE DOBELL, M.D., Assisted by numerous and distinguished Coadjutors.

"This is truly a valuable work, and Dr. Dobell will not only receive for it the thanks of the physicians of his own country, but of those of all other countries. Every scientific fact of interest in medicine that has been brought out by the researches of the learned in the profession is here noted."—CINCINNATI MEDICAL REPERTORY, April, 1870.

"Dr. Dobell, in projecting the serial work before us, has aimed at bringing together, in the English language, original and independent Reports from all parts of the World of New Researches in connection especially with Practical Medicine. The idea is an excellent one . . . . The work is an interesting one; a few references will show this."—LANCET, April 9.

"The work is difficult of analysis, but we trust we have already sufficiently indicated its practical value to show our readers that it should be in the hands of every busy practitioner of medicine."—New York Medical Gazette, April 9.

"The volume is cheerful in its style and aspect, its perusal easy, and it includes something interesting for everyone who cares for his Profession... A sort of Medical Exhibition of All Nations, an admirable idea."—MEDICAL PRESS. May 4.

# NEW BOOKS

PUBLISHED AND SOLD BY

# H. K. LEWIS, 136 GOWER STREET. W.C.

Sir William Jenner, Bart., M.D., D.C.L., F.R.S.

THE PRACTICAL MEDICINE OF TO-DAY: Two Addresses delivered before the British Medical Association, and the Epidemiological Society. Small 8vo, 1s 6d.

### Sidney Ringer, M.D.

PROFESSOR OF THERAPEUTICS IN UNIVERSITY COLLEGE, ETC.

A HANDBOOK OF THERAPEUTICS. Sm. 8vo, nearly 500 pages, cloth, 10s. 6d.

"The author, as we have said, designed the book simply as an introduction to Therapeutics for the use of students and young practitioners. That to such it will be found most useful there is no doubt; but we are also sure that, from the mass of valuable, readable, and well-arranged information it contains, and the numberless minor practical hints scattered throughout its pages, the work will be almost equally useful to the busy practitioner; and that it will receive from the profession generally the warm welcome which it deserves."—BRITISH MED. JOURNAL.

"The work contains a large amount of most useful matter, much of which will be novel to the English student. There is moreover an honest directness of statement, and an apparent desire to balance opposing opinions with candour, which is calculated to attract confidence. We can scarcely doubt that the present edition will be quickly sold."—The Practitioner.

"We can congratulate Dr. Ringer on turning out a book for which we predict almost certain success. Professedly founded on the admirable treatise of Buchheim, it is the embodiment of the best and most recent views on therapeutics. Dr. Ringer wisely avoids collecting into an omnium gatherum everything that has been said as to the action of certain remedies. Such a mélange is precisely what is not wanted. Quality, and not quantity, is the modern requisite in therapeutics, and Dr. Ringer has done much to eliminate the certain from the uncertain. The classification is the classification of Buchheim—the most scientific now in use—and Dr. Ringer has also borrowed from the valuable, if diffuse, treatise of Clarus. The works of Parkes and Stillé have also been consulted. The remedies are grouped as far as possible in accordance with the effects of some substance common to or similar in all. A table of doses is added, and—a somewhat novel feature in a Medical work—there is a dietary for invalids. Altogether, within the small space available, a great deal of sound matter, most of it tested by careful clinical experience, has been collected."—Medical. Times and Gazette.

"Dr. Sidney Ringer's "Handbook of Therapeutics" is a very well printed, a very nicely bound, and a well cut book. The work bears evidence of careful writing."—Medical Mirror.

## Joseph Toynbee, F.R.S.

THE DISEASES OF THE EAR: Their Diagnosis and Treatment. With a Supplement by JAMES HINTON Demy 8vo, 8s. 6d.

"The mantle of the master has fallen upon a worthy successor, and in the hands of its present accomplished editor the volume will maintain its high position as the first text-book on the subject."—Medical Times and Gazette.

"Great credit is due to Mr. Hinton for the manner in which he has supplemented the labours of his deceased friend. The work in its new form constitutes a most reliable guide to the treatment of aural disease, and is without a rival in the English language."—Lances.

### Felix von Niemeyer.

A TEXT-BOOK OF PRACTICAL MEDICINE, With Particular Reference to Physiology and Pathological Anatomy. Translated from the seventh German Edition, by special permission of the Author. By GEORGE H. HUMPHREYS, M.D., and CHARLES E. HACKLEY, M.D. Two vols. roy. 8vo, 1500 pages, 36s.

"To America we owe a debt of gratitude for the work already done, and the debt is increased in no small degree by the translation of Niemeyer now before us. The publication of these volumes brings within the reach of the busy practitioner, the student and the teacher in our schools, one of the best text-books in any language. As the translators say, "Professor Niemeyer's volumes present a concise and well-digested epitome of the results of ten years of carefully rendered clinical observation by the most illustrious medical authorities of Europe," elaborated, we may add, by a most able teacher and an experienced physician. Like 'Watson's Practice of Physic' in these islands, Niemeyer is a classic among the text-books of the Continent. To translate such a work worthily is no easy task, and we congratulate Drs. Humphreys and Hackley on their success. The style throughout is clear and good, and the rendering, though often close, has little, if any, of that ruggedness which so often mars an otherwise good translation. . . . Not the least valuable part of the book will be found in the therapeutical sections, and the numerous formulæ, which are given most clearly, will prove a boon to many a hardworked practitioner. . . . We cannot omit the praise due to the publisher. The type is good and clear, the paper excellent. The publisher has in a word, done his work in the same spirit as the translators, and the result is very creditable to both."—BRIT. AND FOR. MED. CHIR. REVIEW.

# Meigs and Pepper.

A PRACTICAL TREATISE ON THE DISEASES OF CHILDREN. Fourth edition, roy. 8vo, cloth, cut edges, 920 pages, 26s.

"Good books on the diseases of children are so scarce that we cordially welcome this new edition of the once popular work of Dr. Meigs. The labour of revision was found to involve so many changes, and the addition of so much new material, that Dr. Meigs called to his assistance Dr. William Pepper, a gentleman of large experience, and well-qualified for the task. The result of their joint labours is now presented to the public, and so great have been the changes that the volume before us has really the interest of a new book."—New York Medical Record.

### Dr. Wickham Legg.

# A GUIDE TO THE EXAMINATION OF THE URINE. Second Edition, revised and enlarged, sm. 8vo, cloth, 2s. 6d.

"We are glad to welcome this little work . . . . Just what was wanted."—BRITISH MEDICAL JOURNAL.

"This little work forms a concise guide to the complete 'clinical' examination of the urine . . . . The work is a very handy one, and we recommend the student to make it his companion in the wards of the hospital."—LANCET Jan. 8th, 1870.

# Spencer Watson, F.R.C.S., Eng.

ON ABSCESS AND TUMOURS OF THE ORBIT. 8vo, cloth, 6s. or in two parts, 3s. each.

### Thomas Inman, M.D.

ON THE PRESERVATION OF HEALTH. Demy 8vo, cloth, 160 pages, 5s.

ON THE RESTORATION OF HEALTH. 8vo, cloth,

Charles Alexander Gordon, M.D., C.B.

REMARKS ON ARMY SURGEONS AND THEIR WORKS. 8vo, cloth, 5s.

# S. Scott Alison, M.D.

MORBID CONDITIONS OF THE THROAT, in their relation to Pulmonary Consumption. Post 8vo, 3s. 6d.

# Dr. Graily Hewitt.

OUTLINES FOR PICTORIAL DIAGNOSIS OF DIS-EASES OF WOMEN. 4to, 6s.

#### Weber.

CLINICAL HANDBOOK OF AUSCULTATION AND PERCUSSION. An exposition from first principles, of the method of investigating diseases of the respiratory and circulating organs. By JOHN COCKLE, A.M., M.D. With illustrations, 12mo, cloth, 1s. 6d. (published at 5s.)

### Lindley.

SYNOPSIS OF THE BRITISH FLORA. Arranged according to the natural orders. Containing vasculares or flowering plants, third edition, 12mo, 3s. (published at 10s. 6d.)

### Rev. J. Berkeley.

INTRODUCTION TO CRYPTOGAMIC BOTANY. Numerous woodcuts, 8vo, 5s. 6d. (published at £1.)

# Joseph Sampson Gamgee, F.R.C.S.

THE STARCHED APPARATUS IN THE TREAT-MENT OF FRACTURES AND DISEASES OF THE JOINTS. 8vo, cloth, 5s.

# John Henry Griesbach.

ANALYSIS OF MUSICAL SOUNDS with illustrative figures of the ratios of vibrations of musical intervals and their compounds, temperament, harmonic vibration, resultant sounds, &c., &c. Small folio, cloth, £1 5s.

#### Dr. Dobell.

THE NURSING SCHEDULE. To be used in Fevers, Inflammations, Surgical Cases, and other acute Illnesses. In packets reporting 14 days and 14 nights. 15.

#### Alexander Bruce.

OBSERVATIONS IN THE MILITARY HOSPITALS OF DRESDEN. 8vo, 1s.

# C. H. F. Routh, M.D.

ON THE CHEAPEST AND MOST NUTRITIOUS FOOD FOR CHARITABLE INSTITUTIONS AND THE POOR; Being the Result of an inquiry, made by desire on the Food supplied to the Hill Street Female Refuge. 15.

# Dr. Rigby.

TREATISE ON MIDWIFERY. Post 8vo, 3s. (published at 10s. 6d.)

### Dr. R. Hooper.

ON THE MORBID ANATOMY OF THE HUMAN BRAIN. Illustrated by coloured engravings of the most frequent and important Organic Diseases to which that Viscus is subject. 4to, 16s. (published at £2 12s. 6d.)

ON THE MORBID ANATOMY OF THE HUMAN UTERUS AND ITS APPENDAGES; With illustrations of the most important Organic Diseases to which those Viscera are Subject. Coloured plates, 4to, £1 5s. (published £3 3s.)

#### Dr. W. Abbotts Smith.

ON HUMAN ENTOZOA, AND THE DISORDERS PRODUCED BY WORMS; With numerous engraving and a Glosary of Terms, 8vo, 8s.

### Robert Collins, M.D.

PRACTICAL TREATISE ON MIDWIFERY; containing the result of 16,954 births occurring in the Dubling Lying-in Hospital, 8vo, 5s. (published at 12s. 6d.)

#### R. E. Grant, M.D.

OUTLINES OF COMPARATIVE ANATOMY. Illustrations, 8vo, cloth 5s. (published at 26s.)

#### Dr. R. Lee.

LECTURES ON MIDWIFERY. 8vo, 6s. (published at 16s).

ANATOMY OF THE NERVES OF THE UTERUS. 2 plates, folio, is. 6d. (published at 8s.)

#### R. D. Lyons.

HAND-BOOK OF HOSPITAL PRACTICE; Or an Introduction to the practical study of Medicine at the bed-side, with cases, &c. post 8vo, 2s. (published at 6s. 6d.)

TREATISE ON FEVER. 8vo, 6s. (published at 12s.)

### Dr. J. C. Prichard.

ON INSANITY IN RELATION TO JURISPRUDENCE. 12mo, cloth, 1s. 6d. (pubished at 5s.)

# John Quekett.

LECTURES ON HISTOLOGY, Delivered at the Royal College of Surgeons, woodcuts, 2 vols., 8vo, 12s. (pub. £1 8s. 6d.)

Vogel.

PATHOLOGICAL ANATOMY OF THE HUMAN BODY. Translated by Dr. Day, plates, some coloured, 8vo, cl., 8s, (published at 18s.)

#### Dr. W. H. Walshe.

NATURE AND TREATMENT OF CANCER, plates, 8vo, 3s. (published at 16s.)

### Dr. R. Williams.

ELEMENTS OF MEDICINE, (Morbid Poisons), 2 vols., 8vo, cl., 13s. (published at £1 6s.)

#### Dr. Forbes Winslow.

ANATOMY OF SUICIDE, 8vo, cl. 3s. (published at 12s.)

#### Marshall Hall.

DISEASES OF THE NERVOUS SYSTEM, plates, 8vo, 3s. 6d. (published at 15s.)

#### Dr. F. H. Hamilton.

MILITARY SURGERY AND HYGIENE, 127 Illustrations, 8vo, 12s. (published at £1 5s.)

#### Trousseau and Reveil

THE PRESCRIBER'S COMPLETE HANDBOOK. Edited by NEVINS. Square 18mo, limp roan, 2s. 6d. (published at 6s. 6d.)

#### Thomas Graham, F.R.S.

THE ELEMENTS OF CHEMISTRY, including the application of the Science in the Arts, illustrations, 2 vols., 8vo, 12s. (published at £2.)

# THE NEW SYDENHAM SOCIETY.

Annual Subscription, One Guinea.

#### WORKS FOR 1870.

I.

TROUSSEAU'S CLINICAL MEDICINE, Vol. III. (now ready).

NIEMEYER'S CLINICAL LECTURES ON PHTHISIS. Translated by Dr. BAUMLER. (ready).

STRICKER'S HISTOLOGY, Vol. I. With numerous illustrations. by Dr. POWER. (In the Press).

TENTH FASCICULUS OF THE ATLAS OF SKIN DISEASES.

Names and Subscriptions of new Members received by any of the Hon. Local Secretaries, and by the Society's agent, Mr. H. K. Lewis, 136 Gower Street.





