A treatise on the art of cupping: in which the history of that operation is traced; the various diseases in which it is useful indicated; and the most approved method of performing it described / [Thomas Mapleson].

Contributors

Mapleson, Thomas, Cupper, active 1813-1829.

Publication/Creation

London: printed for the author, 1813.

Persistent URL

https://wellcomecollection.org/works/b59h9f6q

License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.




Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org TISE ON CUPPING.

BY.

THOMAS MAPLESON

35305/A



Mittere autem sanguinem cum sit expeditissimum usum habenti; tamen ignare difficillimum.—To let blood is extremely easy to an experienced operator, but very difcult to one who is ignorant.

Idque auxilium, [sc: cucurbitulæ] ut minus vehemens, ita majus tutum; neque unquam periculosum.—And this remedy, cupping, as it is less violent, so it is more safe; nor is it ever attended with danger. Celsus, lib: ii. eap. xi.

35997

TREATISE

ON THE

ART OF CUPPING:

In which the History of that Operation is traced; the various Diseases in which it is useful indicated; and the most approved method of performing it described.

BY THOMAS MAPLESON, CUPPER

To HIS ROYAL HIGHNESS THE PRINCE REGENT,

To the Westminster Hospital,

THE ST. PANCRAS PAROCHIAL INFIRMARY.

London:

PRINTED FOR THE AUTHOR;

AND SOLD, PRICE 3s.6d. AT HIS RESIDENCE, GOLDENSQUARE; ALSO BY J. CALLOW, MEDICAL
BOOKSELLER, CROWN-COURT, SOHO.

1813.

HISTORICAL MEDICAL

G. Sidney, Printer, Northumberland-street, Strand.

PREFACE.

HAVING been frequently solicited by medical practitioners, residing in the country, to procure for them the apparatus for cupping, as well as to furnish them with directions how to perform the operation, to save trouble, I determined to have them printed. As all that I had to say concerning the subject, could occupy but a very few pages, it was suggested to me, that a slight historical sketch of the operation, which appears to be of great antiquity, together with an account of the complaints in the cure of which it is now most generally employed in the metropolis, and its vicinity, might not be unfavourably received, especially as no separate treatise on the subject, as far as I know, exists.

Not to acknowledge that, in the compilation of these remarks, I have been much indebted to the kind assistance of a professional friend, would be equally ungenerous and uncandid.

Golden Square, Nov. 10, 1813.

CONTENTS.

PART I.—Historical Sketch of the opera-	y ex
tion of Cupping	1
PART II A synopsis of the diseases, in	
the treatment of which cupping is found	
useful	30
PART III Practical instructions for per-	
forming the operation	59

Diseases in which Cupping is generally employed with advantage.

APOPLEXY.

ANGINA PECTORIS.

ASTHMA.

BLOOD, SPITTING OF

BRUISES.

COUGH.

CATARRH.

CONSUMPTION.

CONTUSION.

CONVULSIONS.

CRAMP.

DISEASES OF THE HIP

AND KNEE JOINTS.

DEAFNESS.

DELIRIUM.

DROPSY.

EPILEPSY.

ERYSIPELAS.

ERUPTIONS.

GIDDINESS.

GOUT.

HOOPING COUCH.

HYDROCEPHALUS.

HEAD ACHE.

INFLAMMATION OF THE EYES.

LUNGS.

INTOXICATION.

LETHARGY.

LUNACY.

LUMBAGO.

MEASLES.

NUMBNESS OF THE

LIMBS.

OBSTRUCTIONS.

OPHTHALMIA.

PLEURISY.

PALSY.

PERSPIRATION, DE-

FECTIVE

PERIPNEUMONY.

RHEUMATISM.

REST, TO PROCURE

SCIATICA.

SHORTNESSOF BREATH

SORE THROAT.

PAINS OF THE SIDE

AND CHEST.

HISTORY

OF THE

ART OF CUPPING.

Remedies were divided by the ancients into two general classes:—those by means of which something was abstracted from the living body,—and those by which something was added to it. Among the former the art of diminishing, by means of an artificial operation, the quantity of the vital fluid, or circulating blood, by opening a blood-vessel, may certainly be considered as one of the most efficacious and important.

The abstraction of blood, as a means of removing disease, is, indeed, mentioned in the earliest records of the science of medicine.

Blood letting is divided into general and topical. The former is well known to be effected by making a puncture in a vein, or more rarely in an artery, and allowing the blood to flow till the quantity of the whole mass be duly diminished. Topical or local bleeding is performed by making slight incisions, or scarifications, hardly extending deeper than the thickness of the skin. It must have been very early discerned, however, that the blood did not flow readily from these superficial wounds, without the aid of some

species of suction. The most natural, and probably the earliest, means, of effecting this purpose, would be, by the application of the lips, probably suggested by the custom of sucking the poison out of wounds. The utility of some intermediate vacuity must have been discovered at a very early period, as HIPPOCRATES takes notice of two kinds of instruments for the purpose of eliciting blood after the operation of scarifying. The first a small gourd, or cucurbit, furnished with two orifices, one of a sufficient size to include the carifications, the other small; by applying the mouth to the latter, the air was sucked out, and a partial vacuum formed; it was then closed by means of a bit of wax, till a proper quantity of blood was obtained. The other, of similar form and size, was constructed of brass, but with only one aperture. In this a vacuum was produced by inserting a piece of burning flax or linen previously to applying it to the skin; a method very analogous to that which is employed for making a vacuum in the cupping glass at the present day.

HIPPOCRATES mentions also the application of exhausted cucurbits over the orifices made by leeches.

It is probable, however, that the art of scarifying and abstracting blood, by the application of vessels in which the air is rarified by means of heat, as a mode of removing disease, was, like many other arts, adopted by the physicians of

Greece from the more ancient practice of the natives of Ægypt.

PROSPER ALPINUS, who resided several years in Ægypt, as physician to the Venetian consulate, about the close of the sixteenth century, has given an excellent account of the practice of medicine among the people of Ægypt, at that period; and as it is well known that although the nations of the east have frequently changed masters, they are not very prone to change manners and customs, it may with probability be conjectured, that the state of medicine was then much the same as at a period of very remote antiquity.

Among the Ægyptians, and the various nations who inhabit that peculiar

country, the abstraction of blood appears to have been considered as a remedy for almost every species of disease, as well as an important means of preserving health. The reason they give for supposing that they laboured at all times under a plethora, or redundance of blood, is not a little singular. They contend that the water of the Nile, which is their chief drink, is in the living body converted entirely into blood, and that it has been possessed of this property ever since Moses, at the command of the Lord, turned the waters into blood; " and he lift up the rod, and smote the waters that were in the river, in the sight of Pharaoh, and in the sight of his servants; and all the waters that

were in the river were turned into blood."

The complaints in which the Ægyptions used scarifying, for which he quotes the authority of Herodotus, a celebrated physician of antiquity, were the following: "Scarification with cupping possesses the power of evacuating offending matter from the head, of diminishing pain of the same part; of lessening inflammation, of discussing inflations; of restoring appetite; of strengthening a weak stomach; of removing vertigo, and tendencies to faint; of drawing deep-seated offending matter towards the surface; of drying up fluxions; checking hæmorrhages; promoting menstrualevacuations; arresting the tendency

to putrefaction in fevers; allaying rigors; accelerating and moderating the crisis of diseases; removing a propensity to somnolency; conciliating natural repose; removing heaviness;—these and many analogous maladies are relieved by the judicious application of the cucurbits, dry or bloody."

They appear, however, to have considered cupping rather as a means of producing derivation, than of diminishing the whole quantity of blood.

ALPINUS states, that children in Ægypt are peculiarly liable to inflammations of the eyes, and enlargement of the tonsils. For the cure of these complaints they scarify the lobes of the ears; from which, after being well fomented with

hotwater, a considerable quantity of blood may be obtained; so common at certain seasons of the year is this operation, that of a hundred children taken promiscuously, you will find sixty with pieces of lint adhering to the parts of the ear where scarification had been employed.

In head-aches they take blood by scarification from the inside of the nostrils with much advantage. On the first attack of the plague they bleed largely.

Cupping behind the ears they consider as peculiarly efficacious in conciliating sleep.

They are either unacquainted with the use of leeches, or have a prejudice against them, as considering their bite to be venomous. They also use scarifying with cupping successfully, to remove the swellings consequent to reiterated attacks of the gout.

But by far the most general place to take blood by scarifying and cupping is the calves of the legs. In all complaints of the head, and in suppressions of the evacuations, peculiar to the female sex, this is considered as much the most efficacious remedy that can be employed. In order to make the operation succeed properly, the legs must be carefully fomented with hot water, and a slight bandage applied below the knees, to arrest the course of the blood through the superficial veins. The practice of cupping on this part of the body is at present

neglected, but I can see no reason why it should not be revived, as it is attended with less pain than perhaps any other part of the body.

Among the Romans the utility of cupping appears to have been properly appreciated, and the practice to have been very general. The description of the operation, and the account of the diseases in which it is beneficial, given by the celebrated Celsus, are so correct and judicious, that I shall insert them at full length for the sake of the information they contain, as well as to mark the general employment of, and confidence in, this remedy at that period.

" * Of cucurbitulæthereare two kinds;

^{*} Celsus de Medecina, lib. ii. cap. xi. De sanguinis detractione per cucurbitulas.

brass and horn. Those of brass have an aperture on one side, the other is close; those of horn have a similar aperture on one part, and on the opposite a small foramen or hole. Into those of brass a piece of burning flax is thrown, the aperture is then immediately applied to the surface of the body, and pressed on till it adheres. Those of horn are simply applied to the skin, then, by applying the mouth to the small aperture, the enclosed air is sucked out, the hole is then stopped with a bit of wax, and adhesion takes place. Both kinds may be formed not only of these materials, but of any thing else that will answer the like purpose. If no. thing else is at hand, a little cup or pot, provided it be somewhat narrow towards the mouth, may be used for this purpose. When adhesion has taken place, if the skin had been previously scarified with a lancet, blood is drawn forth, if entire, air. Therefore, when the matter within is of a noxious quality, the former method is to be employed; when it is merely flatus the latter."

"Cupping is more peculiarly useful in local than in general diseases, where offending matter seems to be seated in some particular part of the body, the extraction of which is sufficient to restore health. Of this it is a sufficient proof that if it be necessary to take blood from any diseased part by scarification, the nearer that it is done to the injured part

of the member it is the more efficacious; nor does any person apply a cucurbit to a distant part of the body, unless he is desirous of making a derivation of blood from the part affected; but as nearly as possible to the seat of pain which it is intended to remove."

"Cucurbits may also be useful in chronic complaints (even although some time has already elapsed), if there be peccant matter or improper flatus. Moreover, in some acute diseases, they are requisite, where it is necessary to relieve the system, and the strength does not admit of taking blood from a vein. And this remedy, as it is less violent, so it is more safe; nor indeed is it ever dangerous, even if employed in the first attack of fever, or

while crudity still prevails. Therefore, when there is a necessity for taking away blood, and there is obvious danger in opening a vein, or when the disease is seated in some vital part, to this remedy we must have recourse; aware, however, that while it has no danger, it is at the same time less efficacious than general blood-letting; for it is impossible to arrest the progress of a violent disease, otherwise than by the assistance of a remedy equally violent."

This operation as a remedy for disease appears indeed to have spread all over the world. I have cupped various gentlemen who have resided in India, who have informed me, that they have had the same operation performed in that distant

region by natives, generally of the female sex, who continue to use the same means as the Ægyptian cuppers. They extract the air by applying their mouths to the smaller orifice of a suitable vessel, generally formed of earthen ware, having previously made a great number of incisions in the skin, by means of a razor, which they contrive to use in such a manner as to occasion little or no pain. It is indeed a remedy in common use among the natives of India, chiefly employed by them for diseases of the liver and the spleen, in the removal of which it is found eminently successful.

I have been informed, that local extraction of blood, by applying the mouth to scarifications previously made by incisions with a sharp reed, is common in the islands of the South Sea. Something of the same kind has been observed to be practised by the rude natives of New Holland.

It is, however, very singular, that the original inhabitants of America, the savages, as we are pleased to term them, who inhabited the Isthmus of Darien, when that part of America was first visited by the Buccaneers, appear to have trusted the cure of their diseases chiefly to an operation of this kind. It is minutely detailed by Lionel Wafer, Surgeon to Captain Dampier, in his very curious description of the Isthmus of America.

After giving an account of his being

taken captive by the native Indians, he proceeds to observe: "We had not been long here, before an occurrence happened, which tended much to increase the good opinion Lacenta and his people had conceived of us, and brought me into particular esteem with them.

"It so happened, that one of Lacenta's wives being indisposed, was to be let blood, which the Indians perform in this manner.—The patient is seated on a stone in the river, and one with a small bow shoots little arrows into the naked body of the patient, up and down, shooting them as fast as he can, and not missing any part. But the arrows are guarded, so that they pene-

trate no farther than we commonly thrust our lancets; and if by chance they hit a vein which is full of wind, and the blood spirts out a little, they will leap and skip about, shewing many antic gestures, by way of rejoicing and triumph.

"I was by, while this was performing on Lacenta's lady, and, perceiving their ignorance, told Lacenta, that, if he pleased, I would shew him a better way, without putting the patient to so much torment. 'Let me see,' says he; and at his command I bound up her arm with a piece of bark, and with my lancet breathed a vein. But this rash attempt had like to have cost me my life; for Lacenta, seeing the blood issue

out in a stream, which used to come only drop by drop, got hold of his lance, and swore by his tooth, that if she did otherwise than well, he would have my heart's blood. I was not moved, but desired him to be patient, and I drew off about 12 ounces, and bound up her arm, and desired she might rest till the next day; by which means the fever abated, and she had not another fit. This gained me so much reputation, that Lacenta came to me, and, before all his attendants, bowed, and kissed my hand. Then the rest came thick about me, and some kissed my hand, and others my knee, and some my foot; after which, I was taken up into a hammock, and carried on men's shoulders, Lacenta himself making a speech in my praise, and commending me as much superior to any of their own Doctors. Thus was I carried from plantation to plantation, and lived in great splendour and repute, administering both physic and phlebotomy to those that wanted."

Although honest Lionel Wafer may have introduced a considerable improvement in the medical art of his Indian friends, his plain narrative abundantly testifies, that previous to his arrival they were in possession of the art of taking away blood by superficial scarification; and that they even made use of a spring lancet, (for the small bow and guarded arrow comes to the same thing,) which

is, comparatively speaking, a recent improvement in the art of surgery.

I was lately cupping a gentleman from the West Indies, who informed me, that perceiving a fresh imported negro singularly occupied, upon more particular investigation, he found him sucking the blood from a number of small orifices in the skin of his arm, by the intervention of a gourd, which he said was to remove the effects of a bruise.

In the works of Galen there are a great many observations on the utility of scarifying and cupping, and of the distinctions to be observed between that operation, blood-letting, and the application of leeches. As Galen was the oracle of the medical profession, during what

are commonly termed the dark ages, there can be no doubt that the operation would then continue to be performed according to his directions.

One of the most important improvements in the art of cupping, is connected with the discovery of the spring box, by means of which a number of incisions are made at once, instead of being done in succession, by repeated strokes of the lancet. By the use of this instrument, the pain of the operation is considerably diminished, and the time required to perform it much curtailed.

All my endeavours have been hitherto ineffectual to ascertain the precise period of the introduction of this useful improvement. It is not noticed in the vo-

luminous works of AMBROSE PARRY, surgeon to Henry the Fourth of France, and which appear to contain cuts and descriptions of all the chirurgical instruments in use at that period. Neither is it to be found in the Armamentarium of Schultens, published in a work, professing to exhibit figures of every instrument known at that day. There is however a good figure of it in the works of Heister, the first edition of which was published about the year 1710. Indeed, I have had spring scarificators in my own possession, the workmanship of which clearly indicated them to have been made about that period.

Until little more than a century ago, scarification and cupping appear to have

been operations performed by the regular surgeon, when deemed necessary. About that period the use of warm baths was introduced into this country by a person who had resided some years in Asia, and which still continue to be designated nearly by their original appellation Haumaum, the Turkish appellation for a warm bath, corrupted in common parlance into Hummums. As these baths were copied from those of Ægypt, a country to which I have endeavoured to trace the origin of scarification and cupping, which were there generally performed in the warm bath, as when the practice of warm bathing was introduced into this country, the practice of cupping came along with it.

The following documents I think curious, in as far as they shew the manner in which these novelties were first made public.

*The Queen's Bagnio, in Long Acre, is made very convenient for both sexes to sweat and bathe privately every day, and to be cupped in the best perfection, there being the best and newest instruments for that purpose. Price 5s. for one single person; but if two or more come together 4s. each.—There is no entertainment for women after 12 o'clock at night, but all gentlemen who desire beds may have them at 2s. per night.

Persons may be cupped at their own houses; the way of cupping is the very same as was used by the late Mr. Verdier, deceased.

^{*} Tatler, Vol. ii. No. 95, p. 429, Chalmers's edit.

Wash Balls, perfumed, camphired, and plain, shall restore complexions to that degree, that a country Fox Hunter, who uses them, shall in a week's time look with a courtly and affable paleness, without using the bagnio or cupping.

*Airpumps, single and double barrelled, with apparatus for demonstrating the several properties of the air.—Small air pumps, with glasses for the new way of cupping; scarificators, one of which makes at once 10, another 13, another 16, effectual incisions.

The custom, which appears to have become prevalent of resorting to these Bagnios, or Haumaums, to be bathed and cupped, appears to have superseded the

^{*} Spectator, Vol. iv. No. 289, Note.

practice of this operation by the regular surgeons. Falling into the hands of mere hirelings, who practised without knowledge, and without any other principle than one merely mercenary, the operation appears to have fallen into contempt, to have been neglected by Physicians, because patients had recourse to it without previous advice, and disparaged by regular Surgeons, because, being performed by others, it diminished the profits of their profession. Of late years, however, the utility of this local abstraction of blood has been recognised by some of the more enlightened and eminent practitioners of both medicine and surgery. The practice has been rescued from that class of inferior practitioners into whose

hands it had fallen. Men of experience have devoted themselves to this peculiar operation; among whom I trust I may be allowed to class myself as an humble individual; an instrument, I hope, for good in the hands of Providence. Still the adroit performance of this operation, simple as it may appear, continues to be confined to a few individuals in the capital. My purpose in committing these few pages to the press, is, to render more extensive the practice of an operation now generally acknowledged to be useful, and which the remarks contained in the subsequent pages, especially if aided by a very few practical lessons, will, I trust, enable any man, possessed of common ingenuity, to perform with propriety and even elegance.

OF COMPLAINTS IN WHICH CUPPING,
WITH OR WITHOUT SCARIFICATION, HAS BEEN FOUND USEFUL.

I beg leave here to disclaim any intention of intruding on the province of the regular practitioner of medicine, by pretending to direct in what complaints cupping is to be used; but having had the honour of being employed by many of the most eminent Physicians of the present day, I trust I may, without any impropriety, be allowed to state some of the diseases in which, under their direction, this remedy has been successfully employed.

The physicians of Antiquity, by whom Cupping, both with and without scarification, appears to be very frequently employed, seem to have considered, that this operation was not merely useful by diminishing the general mass of blood, but that it also acted by producing a derivation of that fluid from the part affected. Without pretending to offer any opinion upon the theories of derivation and revulsion, which have occasioned so many disputes in the medical world, I may be permitted to state, that I have seen many instances of inflammation of the eyes, being immediately relieved by taking blood from the temples, or from the nape of the neck, and behind the ears, which had

received no benefit whatever from the application of numerous leeches in their more immediate vicinity. Indeed, several cases have occurred to me, where the redness of the eyes seemed to be augmented, and the sense of fulness to the feelings of the patient increased, after the application of leeches to the temples, when, by applying a cupping glass over the part bitten, and taking away more blood, immediate relief has been produced.

It is upon this principle of revulsion that the ancient Physicians appear to have recommended the application of cupping glasses to the legs and thighs, in suppression of the Catamenia; a practice which I know, by experience, to be eminently successful; and for a

similar reason, when that secretion exceeded in quantity, they were in the habit of applying large cups, with or without scarification, upon and in the neighbourhood of the breasts. I understand this practice is particularly recommended, and insisted upon, by Hippocrates.

In all diseases occasioned by fulness of blood in the head, Cupping is particularly useful.

Head-ache, when seated in the fore part of the head, and attended with a sense of fulness, is generally immediately relieved by cupping on the occiput. When the pain is seated in the back part of the head, which I believe is generally considered as a symptom of debi-

lity, I have not found the abstraction of blood so useful, nor would I recommend it. When the pain is felt chiefly over one eye, a complaint to which the name of clavus hystericus has been given, and which is generally connected with disorder of the digestive faculties, I have not found cupping to be of service.

In the Delirum attendant on the beginning of fever, Cupping never fails to do good; and I have reason to believe, that if a sufficient quantity of blood be drawn away, the further progress of the disease, in most cases, may, by this means, be effectually checked.

In Phrenitis, or Inflammation of the Brain, cupping is the most active and energetic means of depletion that can be employed.

In slighter cases of INSANITY, abstraction of blood always does good, and in the more confirmed states of this deplorable calamity, seldom fails to afford (at least) temporary relief. I was lately sent for, by an eminent practitioner, to a gentleman who had passed two nights totally without sleep, and was committing all manner of extravagances, such as tossing his watch and clothes out of the window, &c. I was directed to take full twenty ounces of blood from the occiput. In a short time the patient fell into a sound sleep, which continued for about twelve hours. awoke perfectly tranquil, and rational, and apparently without any recollection of his frantic conduct.

In that species of Delirium, or Insa-

nity, to which some delicate females are liable after parturition, I have witnessed many instances, where cupping has been eminently useful.

In Vertico, or giddiness of the head, and disposition to lethargy, which may perhaps all be considered as species of, or tendencies towards, APOPLEXY, it is well known cupping always affords relief, and, indeed, is generally recommended. I have heard a very eminent practitioner, of long and extensive experience observe, that if the effects of cupping were more generally known, and duly appreciated, we should not hear of so many instances of sudden death.

In Palsy, which is generally the sequel of apoplexy, the utility of cupping

must depend upon the general health of the patient. But as in this disease there is a tendency to fulness of the head, occasional abstraction of blood may, in most instances, be had recourse to with advantage, to obviate repeated attacks of the complaint.

The whizzing noise, or singing in the ears, a very troublesome complaint, is generally removed by cupping; and in certain cases of DEAFNESS, I have also known it extremely useful.

In hydrocephalus, or the watery head, cupping is now recommended, by the most eminent practitioners of the metropolis; and I have been employed to perform the operation in many hundred cases of this complaint, some of

which have terminated favourably, especially if the remedy was used at a sufficiently early period of the disease.

In EPILEPSY, taking blood from the immediate vicinity of the brain, is frequently very beneficial. In a case of epilepsy, which recently occurred at the Westminster Hospital, where the fits amounted to ten and even twelve in the course of the day, I applied cupping glasses, under the direction of Dr. Buchan; and so sensible was the patient, (a young man,) of relief, that he frequently solicited to be cupped, even when it was not ordered by his physician, and by this means, I believe, alone, he obtained a perfect cure.

DEJECTION OF SPIRITS, where that un-

pleasant feeling is produced by pressure of blood upon thebrain, is always relieved by cupping. That Low spirits frequently proceed from this cause, I have reason to believe, from the very general expressions of cheerfulness and relief from weight, which I have heard after the operation, even when this has been performed for other complaints.

Perhaps in no disease whatever is cupping more generally used than in INFLAMMATION OF THE EYES. Taking blood from the temples, the nape of the neck, or behind the ears, produces in such cases almost instantaneous relief. I can assert from experience, that abstracting blood by this means, is much more efficacious than by leeches, and, as I have before observed, has

frequently done good, where these useful animals have been applied in vain.

In cases where it is deemed expedient to open the Temporal Artery, this operation is performed with great facility, by means of the small spring scarificator; and the flow of blood may be immediately arrested by the application of an exhausted glass.

In acute inflammation of the throat, where respiration has been so much impeded as to threaten immediate suffocation, I have witnessed the most decided and instantaneous relief produced by taking blood from the neck.

In the CROUP, that disease so fatal to infant life, I have frequently been employed to cup, and have witnessed the repeated use of the operation, attended with manifest advantage.

In cases of ENLARGED TONSILS, a complaint frequently met with in delicate children, I have seen many instances where cupping has been employed with success.

In the more serious inflammatory affections of the lungs, denominated Pleurisy, and Peripheumony, topical abstraction of blood may always be considered as an useful auxiliary to general venesection. I have observed that many eminent practitioners are in the habit of ordering blistering plasters over the parts that have been recently scarified; and I presume with advantage.

A cough which, when recent, may

be considered as a slight inflammatory affection of the lungs, is in many instances immediately removed by cupping.

Slight bleedings are found to give relief in PULMONARY CONSUMPTION; but when they are topical, they certainly weaken less, than when blood is taken from the general system.

SHORTNESS OF BREATHING, and PAROXYSMS of ASTHMA, are generally relieved by cupping. In the last named distressing complaint, I believe doubts exist, respecting the propriety of taking blood from the system, but I never saw any mischief result from topical bleeding; on the contrary, it generally gives immediate and considerable relief.

Of Palpitation of the Heart, I have seen several instances where relief from cupping on the chest was not only almost immediate, but permanent.

PAIN, OR STITCH OF THE SIDE, is in general immediately relieved by topical bleeding.

In Measles I have, in many cases, been employed to cup upon the chest, with a view to obviate the troublesome cough that is too frequently the consequence of that disease; and it may not be improper to observe, that children in general complain less of cupping, than persons who had not seen them undergo the operation would imagine.

In obstinate cases of Hooping Cough,
I have also known the local abstraction
of blood to be attended with advantage.

In Obstructions of Urine, I have in many cases been directed to take blood from the loins, and vicinity of the bladder.

In recent attacks of Lumbago, the application of several glasses to the lower parts of the loins, and taking away a sufficient quantity of blood, will generally remove the pain.

In Rheumatism, cupping with scarification is much recommended by the celebrated Hoffman; he observes, that when the part where the pain is seated feels cold to the patient, less blood is obtained from the part itself, than from the neighbourhood; upon which, therefore, he recommends the glasses to be applied in preference. He quotes a va-

riety of passages from Celsus, in favour of the utility of this operation, which he considers, to use his own terms, as vicarious to blood letting.

In Inflammation, and in obstinate constipation of the Bowels, I have been in many cases directed to cup upon the abdomen, previous to the use of the warm bath.

In Sciatica cupping is always of service. A number of exhausted glasses are applied to the sacrum, and along the outside of the affected thigh, and blood taken from some of them. I have seen many severe attacks of this complaint entirely removed by this means in the cource of a few hours.

In cases of IRRITATION, and SPASM of

the BLADDER, and diseased PROSTATE GLAND, it is now the practice of some of the most eminent Surgeons, to order cupping on the perinæum; and, in many similar cases, on the lower parts of the loins, and I have always heard patients express their conviction of its utility.

In CRAMP, and NUMBNESS of the Limbs, I have seen many instances of immediate relief from the application of the dry glasses.

In some cases of weakness of the ANKLES, and KNEE JOINTS, after blistering and many other remedies had been ineffectually employed, dry cupping was advised as a kind of experiment, or last resort; but the happy result was the complete cure of the

disease.—I recollect some years since, a female relation of mine, being unable to walk for some weeks, from a contraction of the muscles of the legs, I proposed scarifying the parts slightly; she objected to the loss of blood, but permitted the application of dry cupping, frequently repeated, which, in less than a week, enabled her to walk about as usual.

In one case of DEFECTIVE PERSPIRA-TION, accompanied with a dry and scabrous state of the skin, cupping was attended with the most beneficial consequences.

I have known the operation also useful in removing many kinds of cutaneous eruptions on the face, as well as on other parts of the body.

In contusions, from falls and blows, and in cases of extravasated blood, cupping is very serviceable.

When it is thought proper to scarify the legs, to draw off the fluid in ANASARCA, the application of the springscarificator, and partially exhausted glass, answers the purpose extremely well.

Taking blood from the Feet, especicially by applying scarification and cupping to the soles, is mentioned by the celebrated HOPFMAN, as an efficacious means of obviating a return of the cour, in persons liable to that complaint, if used when the early indications of a fit are perceived.—Of the utility of this practice on the feet, I have no personal

experience, although I have been frequently ordered to take blood from gouty patients, for affections of the head, even during a severe attack of that disease.

Of the various purposes of cupping, the most extraordinary I ever witnessed, is the effect of that operation, (when performed in the vicinity of the head) in almost immediately suspending the state of intoxication, the consequence of taking too large a quantity of fermented liquor. To illustrate this subject, I shall, from many cases within my knowledge, detail the following circumstances, which occurred some months By the desire of a Physician I repaired, about 10 o'clock at night, to a

celebrated Tavern, where we found four gentlemen, one of whom was laid on a sofa; his face extremely red, his eyes suffused with tears, the pupils dilated, and his knees every half minute drawn up to his chin, by violent spasmodic convulsions.—Although the state of their companion was such as to create alarm, not one of the party could articulate sufficiently plain to give any account of what had occasioned it; but we learned from the waiter, that a great deal of wine had been drunk.

I was desired to take blood freely from the shoulders; in a short time after the operation was over, the gentleman, who was the youngest of the company, perfectly recovered his senses, and stated every circumstance that had occurred; that they had been hunting all the morning, and had hastily taken a good deal of wine upon an empty stomach, a condition, in which it is very apt to induce sudden intoxication. He soon became so perfectly sober, as to be able to see his companions safely home in a coach.

I have seen many examples of this kind, although none perhaps quite so striking; and am inclined to believe, if the efficacy of this operation in such cases, was more generally known, we should more rarely hear of instances of perhaps unpremeditated intoxication terminating in death.

It may be necessary here to attempt to obviate some prejudices respecting the consequences of cupping.

Frequent repetition of cupping has been said to injure the sight. I can with truth assert, that no instance of this kind has fallen within my knowledge; on the contrary, I have heard many persons observe, that they could discern objects more distinctly after the operation than before; and some gentlemen, whom I frequently cup for inflammatory affections, have told me that their sight continues remarkably good, and even seems to be improved since they had recourse to this remedy.

It is also said, that if a person acquires a habit of losing blood, he cannot

leave it off with impunity; this observation is unquestionably true; but it applies equally to all remedies.—If a person is in the habit of using purgatives, he cannot do without them; but that would not be a reason for omitting to take a cathartic when necessary.—Doubtless a person in perfect health requires no physic, and would not take any; but those whose constitution, either by nature or accident, deviates from this standard of perfection, must use such remedies, as are requisite, to correct the errors which they cannot otherwise remove.

On this subject I shall beg leave to offer the opinion of a sensible writer, and most respectable practitioner of medicine, which will, at least, have the merit of being less interested than my own, as the author has long since paid the debt of nature.

" Blood letting is in itself extremely innocent, in the limited degree in which it is used, and productive of no bad consequences, but imaginary ones, and those who have most experience, will tell us, that we can bleed oftener than we can purge or vomit, with safety, in many cases; and the same can inform us, that in being too saving of our vital fluid, as we fondly call it, we often cherish a snake in our bosoms; for that they know it on many occasions, to act a very troublesome and a very traiterous part in our microcosm, and never so often as when it exceeds in quantity.*

Our ancestors were accustomed to lose blood regularly twice a year, spring and fall, and I have heard medical men of sound judgment, and extensive experience, doubt whether the custom is more honoured in the breach than in the observance.

I am acquainted with a medical gentleman, who considers it as a means of preserving health to lose blood now and then. After cupping, he compares his sensations to those of a person recovering from disease.

He thinks it affords the constitution

^{*} Vide Empirical Observations on the Use of Bleeding in the Island of Jamaica, &c. By GEORGE SPENCE, 1777.

an opportunity of renovating the vital fluid occasionally. He is a man of accurate observation; and I have heard him state, that after losing eight or ten ounces of blood by cupping, he experiences an increase of appetite for about three days; at the end of which period, he supposes the loss to be compensated by a fresh supply of new blood.

To prove that cupping, or the occasional loss of blood, has no tendency to shorten the duration of life, I might cite the example of a Physician, by far the most eminent and the most respected of his time, who not only recommended the operation, very generally; but who, I have the best authority for saying, was himself cupped at least twice a

year, and who died full of age, and of honours, upwards of ninety.

The late Dr. HEBERDEN observes, "the symptom of giddiness is moderated in those who can bear this small loss of blood, by taking away six ounces by cupping glasses, more than by any other means: this has been well borne by those who could not bear the loss of blood from a vein by a lancet. I have known it experienced in several, and particularly in a woman of sixty-eight, who had such bad fits of it, as made her several times fall, and frequently threatened apoplexy. She began the cupping at that time of life, and used it constantly every six weeks, until she died, which happened at the age of eighty-five. She was in no

danger of ever forgetting it; for she felt the most evident marks of wanting this relief, whenever she deferred it beyond the usual period. During all this time, the giddiness was inconsiderable, and came but seldom. She was struck at last with a palsy, which had probably been kept off for many years by this practice of cupping."*

Did the limits of this little publication permit, I could recite the cases of a great number of persons very far advanced in life, who find the operation indispensable; one of which I lately cupped for the seventeenth time within the period of one year.

^{*} Commentaries on the History and Cure of Diseases, 1806.

OF THE OPERATION OF CUPPING.

Notwithstanding the acknowledged advantages derived from cupping with or without scarification, the proper performance of the operation is at present confined within very narrow limits. Indeed, I believe I am justified in stating, that it can hardly be said to extend beyond the boundaries of the Metropolis of these realms.

Cupping is not practised as a distinct profession, either in Scotland or Ireland; and I understand it is rarely, if at all, recommended, even in Edinburgh, that centre of medical information.

In Paris, I am told, the operation is hardly known; of course, seldom practised.

The many applications for instructions in the art of cupping, which I have received from practitioners residing in various parts of the country, suffice to convince me, that the performance of this operation, forms no part of the present system of regular medical Education; though, when gentlemen are settled in practice, they become sensible of the advantage of being able to perform it with propriety. I shall, therefore, proceed to give as plain and explicit directions for the performance of the operation as lies in my power, such as I trust may enable any man of common

sense and observation to perform it; yet ocular observation and manual practice are doubtless requisite to impart neatness and dexterity.

It has frequently occurred to me, that if every young gentleman intended for the medical department of the Navy or Army, was, in the course of his studies, required to make himself master of the mode of performing an operation so extensively useful, the public service would be greatly benefited, and perhaps many valuable lives saved. An apparatus for cupping is indeed, I believe, now furnished by the public medical boards, but as the surgeons are not instructed in the mode of using it, the expence is incurred in vain.

The first step in the operation of cupping is to produce a partial vacuum over one or more portions of the surface of the skin.—This purpose, as has been already stated, was formerly effected by sucking the air out of a hollow gourd or cucurbit, by the application of the mouth, or expelling it from a similarly shaped metalic vessel, by the introduction of a piece of inflamed flax. The invention of glass has wholly superseded these contrivances, and small bellshaped glass cups are now universally used for the purpose of exhaustion, as being neat, cleanly, light, permitting the colour and quantity of blood to be judged of, as it flows from the wounds. For the purpose of exhausting the contained air, these are occasionally fitted up with brass caps and valves, and a small syringe of the same metal. To this plan there are two objections: 1st, the exhaustion is apt to be carried too far; which, instead of promoting, tends to obstruct the flow of blood; 2nd, the operation is rendered extremely tedious and fatiguing, both to the patient and the operator; besides, the glasses are very apt to be broke by the valves losing their elasticity.

The air contained in the glass may also be rarified by steam; but the mode now, I believe, universally adopted by regular cuppers, is the momentary introduction of the flame of a spirit lamp, with a thick wick; the larger the glass,

(if properly exhausted), the less pain does the patient suffer, and the more freely does the blood flow.

When about to perform the operation, let there be provided a hand bason with warm water, a piece of fine sponge, and a lighted candle. Place as many glasses in the bason as may be judged requisite to draw the quantity of blood intended to be taken away. If sixteen or twenty ounces are ordered, four glasses of a size adapted to the surface will, in most cases, be required. Each glass is then separately to be held, for an instant, over the flame of the burning lamp, and immediately placed upon the skin of the patient. Upon the quickness with which this is done, depends the

whole neatness and efficacy of the operation .- To obviate their want of dexterity, many operators in the country throw a little bit of tow, dipped in spirits, and inflamed, into the cupping glass, the moment before it is applied, -a very clumsy expedient, adding unnecessarily to the sufferings of the patient by cauterising the skin; doing harm also by rarefying the air more than necessary within the glass, in consequence of which the edges of the cup compress the cutaneous vessels so much as to obstruct the influx of the blood.

If the glasses have been duly exhausted, the skin will be seen gradually to swell up within the cup, owing to the pressure of the air upon the parts in the vicinity,

as well as the expansion of the fluids contained in the cellular membrane. The skin becomes also of a dark purple colour, owing to the influx of blood into the smaller vessels. If dry cupping be only intended, the glasses may be allowed to remain on the skin for a few moments, and replaced five or six times, varying their position a little, to prevent bruising the skin .- If the intention be to scarify and take away blood, the glass ought not to remain more than a minute, when it is to be removed by gently introducing the nail of the fore finger under the edge, and the scarificator instantly applied and discharged upon the skin, before the tumour has had time to subside. Upon the rapidity or slowness with which the application of the scarificator succeeds the removal of the glass depends all the sufferings of the patient.—If the skin has completely subsided before the stroke of the instrument, much unnecessary pain is inflicted.

The glasses are thus to be removed and reapplied in succession. They should be a second time removed, if necessary, as soon as the blood is perceived to coagulate within them, or when they are so full as to be in danger of dropping off. For the sake of neatness, care should be taken to insert the nail under the upper part of the glass, and open them downwards, gently

wiping the wounds at the same time with a warm sponge.

The glasses previous to every application should be rinced in the warm water, but not dried. To obviate the unpleasant sensation produced by the coldness of the metal, it is adviseable to pass the instrument for a moment over the flame of the lamp before using.

To ascertain the precise quantity of blood, a circumstance generally interesting to all parties concerned, the contents of the cups should be emptied into a graduated glass measure, with which the operator should always be provided.

When the operation is finished, it is common to apply a piece of fine linen rag to the wounds, but f the patient does not object to a little smarting, either Arquebusade water, or spirits of wine, is a preferable application, as it immediately stops the oozing of the blood, promotes the healing of the wounds, and prevents the subsequent itching, which I have heard some patients complain of, as the most unpleasant part of the operation.

In some gross and inflammatory habits I have seen slight suppuration of the wounds made by the scarificator, but never any case that might not be healed by the application of a little of the common wax ointment.

It is a common error to make the incisions too deep, especially if the

object be to take away much blood, but I am convinced nothing is gained by going deeper than the cutis, or true skin, except an unnecessary increase of pain to the patient.

The quantity of blood obtained (in most cases) depends wholly upon the due exhaustion and proper application of the glasses, to attain which requires some practice and experience. In cupping upon the back or neck, the glasses should never be placed upon the spine itself, which is very painful, but on each side of it. If it can possibly be avoided, the glasses should not be applied to parts where the skin is in immediate contact with the subjacent bones; though, if care be taken to apply the scarificator, before

the tumour caused by the glass has subsided, there is no danger.

It is certainly preferable to make the incisions in the direction of the fibres of the subjacent muscles, but it is not of much importance, as in my opinion the incisions should never penetrate so deep.

In some diseases of the head, it is requisite to cup upon the posterior part of the hairy scalp, which of course must be previously shaved. As the shin is peculiarly loose and thick upon this part, the cups often bury themselves so deeply as to require the thin edge of a shilling to displace them, and the scarificator also requires to be screw-

mands more the attention of the oper

ed up so as to cut a little deeper than is necessary in other cases.

From cupping on the temples, I have seen many hundred instances of decided benefit in severe inflammation of the eyes, but I would not advise any person to attempt this operation unless he has had considerable practice upon other parts of the body.

Previously to applying the glasses it is of importance to foment the skin well with a sponge and warm water, especially the calves of the legs.—Cupping in the warm bath is rendered more efficacious, by the relaxed and softened state of the skin.

No part of the operation of cupping demands more the attention of the operator exquisitely keen, they occasion unnecessary pain, if foul or rusty they may communicate disease, or give rise to festering sores.—The lancets or cutters should be kept as sharp and in as fine order as possible, so as to make a clean incision without bruising or giving much pain.

The intention of the operation is, I believe, frequently defeated from want of attention to these apparently trivial circumstances.

An ingenious artist has lately invented an instrument which cuts from the centre outwards. I have frequently tried it, but cannot say that it has any particular advantage over that commonly employed, excepting that it is more readily taken to pieces, and by shifting pinions, and different sets of lancets, may be useful to those persons who are incompetent to put their instruments in order when they become dull.

After cupping, persons are much less liable to faint, except when the operation is performed on the temples, than after bleeding from a vein. If that accident should supervene, the application of volatiles to the nose, and an horizontal posture, will generally be found to restore animation. If the stomach be quite empty, persons are more apt to faint from losing blood, than if it be moderately distended with food. I know many practitioners of medicine, who are of opinion that abstraction of blood produces more benefit when accompanied by fainting, than if it be not.

Many persons prefer being cupped in the evening, to which there can be no objection, provided some hours have elapsed since the principal meal. If the operation is performed while the process of digestion is going on, the serum of the blood has often the appearance of whey.

Cupping glasses have been formerly applied with the view of producing a rapid vesication, which the inventor particularly extols as the most efficient mode of removing the consequences of intoxication. "The organist of a certain Convent of English Benedictines, (says BURNET), having become extremely intoxicated, fell down senseless upon the

stone floor of the tavern where he had been drinking, and bruised the back part of his head. I was sent for to visit him as one past all hope. Finding, however, that there was no fracture, though sterterous breathing and total insensibility, I did not despair, but after administering an active purgative of extract of colocynth, &c. I applied to the nape of his neck, a wide-mouthed cupping glass, exhausted with a large flame.

"After remaining a quarter of an hour I removed it, and immediately replaced it, exhausted as much as possible. In a short time, small vesicles made their appearance, almost the size of pins' heads.

"These gradually coalescing formed, in the course of half an hour, a vesicle so large, as completely to fill the cavity of the glass.

"If the cupping glass be not provided with an aperture, stopped with wax, on piercing which with a needle the vapour escapes, and the glass drops off, I generally break the glass by the stroke of a knife, for if it be too forcibly dragged away by the hand, it lacerates the blister and occasions acute pain. By this means I produce to a certainty in less than an hour, extensive vesication, while blistering plasters, even after being applied for many hours, often disappoint our expectations. This is my own invention, and, as far as I know, hitherto

unemployed by any other person, and by this means, accompanied by the usual remedies, I cured my patient of his apoplexy, as well as of a paralytic affection of his tongue which supervened."*

Of the effects of this mode of producing vesication I profess to know nothing; but it may probably answer equally well as the recently proposed mode of blistering by the application of scalding water.

The operation of cupping differs from venesection in this, that by the former arterial as well as veinous blood is abstracted. The crimson and purple colour may be clearly distinguished, as the two kinds of fluid mingle in the

^{*} Vide Burnet's Thesaurus Medic. Practicæ, 1675.

glass.—The arterial blood generally comes more abundantly on the second application of the glasses than the first. I have observed that the more abundantly the arterial blood appears, the more benefit is generally derived from the operation.

To me it appears that in most cases the state of the constitution may be judged of equally well by the appearances of the blood taken by cupping, as if drawn from a vein.—The bluish film is often very obvious, said to be indicative of the first or slightest degree of inflammatory action.

The strength of the coagulum, the appearance of the buff, often more obvious on the second than the first appli-

cation of the glasses, its different degrees of tenacity, and the tendency to become hollow or cup as it is termed, all present themselves to the practiced and discriminating eye.

Sometimes the blood flows very slowly, and does not coagulate, but retains precisely the appearance of common treacle. Experience has taught me, that this is, in general, a mortal symptom, the almost certain harbinger of death. Physiologists, perhaps, would call it an indication, that the blood was deprived of the principle of vitality.

FINIS.

G. Sidney, Printer, Northumberland street, Strand.



CORNE OF MARCH STATES The state of the same and a comment the party of the last the



