

The pharmacopoeia of the Royal College of Physicians of Edinburgh.

Contributors

Royal College of Physicians of Edinburgh.

Publication/Creation

Edinburgh : A. & C. Black, 1841.

Persistent URL

<https://wellcomecollection.org/works/f6cmafhp>


License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.

**wellcome
collection**

Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>



EDINBURGH
PHARMACOPŒIA

SECOND EDITION

Price 5s.


Alex. Mitchell,
BOOKSELLER,
20, Upperkirkgate,
ABERDEEN.

[2 ed.]

40783/A 1700 - 1850 slacc

CANCELLED

University of Aberdeen




TIMOR DOMINI.


IN SCIENTIÆ

Bought with
Compensation Grant.

~~57B~~

6. 24





PHILLIPS LIBRARY
OF PHARMACOLOGY AND
THERAPEUTICS

lex
Be
Op
A B

atom of C 8

THE
PHARMACOPŒIA
OF THE
ROYAL COLLEGE
OF
PHYSICIANS OF EDINBURGH.

EDINBURGH:
ADAM AND CHARLES BLACK,
AND BELL AND BRADFUTE;
AND LONGMAN & CO. LONDON.



303929

6th October 1841.—Published by Authority of the
College.

R. GRAHAM, President.
G. A. BORTHWICK, } Censors.
R. LEWINS,



DEDICATED
TO THE
QUEEN'S
MOST GRACIOUS MAJESTY,
BY HER MAJESTY'S
FAITHFUL SUBJECTS AND SERVANTS,
THE
ROYAL COLLEGE OF PHYSICIANS
OF
EDINBURGH.

MEDICAL

BY

GEORGE

THE

THE

ROYAL

THE

P R E F A C E.

BEING informed by our Publisher, that our first English Pharmacopœia, which appeared two years ago, has been already disposed of, we present to the Medical Profession this new, and, as we hope, improved Edition. Few additions have been called for. But we have rectified several errors, which have occurred to ourselves, or have been pointed out by various critics :—and we have made some extensive changes, particularly in the formulas for pills and tinctures, by which, although in general little alteration is made in their composition, and scarcely any in their strength, greater facilities will be given in compounding them.

THE A. O. L.

Having informed by our Librarian that
the first English Manuscript which we
found two years ago, has been already the
subject of a present to the Medical Library
of the University, and as we have improved
it. Few additional persons are
yet known to have written several other
works, and we have been
pointed out by various critics:—and we
have made some extensive changes, parti-
cularly in the headings for bills and titles,
and in which, although in general little
alteration is made in their composition, and
scarcely any in their arrangement, greater light
will be given in comparing them.

P R E F A C E

TO THE

FIRST ENGLISH EDITION.

1ST JUNE 1839.

AFTER much delay, the causes of which it would be tedious and of little moment to mention, we are at length enabled to present to the Medical Profession a new Edition of the Edinburgh Pharmacopœia. We do so with no little anxiety for its success; for, notwithstanding the time and care that have been bestowed upon it, we are conscious that the extraordinary progress, which has been made in the sciences of Therapeutics, Chemistry and Pharmacy during the last two-and-twenty years, has rendered the adaptation of a Pharmacopœia to the modern state of Physic no light task, and

constituted the work itself almost a new one rather than a corrected Edition.

In the plan of the present Pharmacopœia we have thought it advisable to deviate materially in several respects from those of former years.

That we have departed from all previous practice of Colleges in this country by publishing our Pharmacopœia in the English language is an alteration, which, as it has been sanctioned by the almost unanimous consent of the College, will also, we apprehend, meet with the general approbation of the medical public. The time is perhaps gone by when public opinion required as a test of learning that a College of Medicine should write in Latin alone; and it may even be questioned whether the practice be not open to censure as leading to risks of inaccuracy in preparing and compounding drugs. Besides, the favourable reception of unauthorized translations of former Pharmacopœias, together with the slow sale of the

last Latin Edition of 1817, seemed sufficiently to indicate the wishes of the profession on this subject, and to show that the Latin language cannot be any longer retained, without occasioning, as of late, serious delays and obstructions in the way of future improvement.

The increasing frequency and extent of the adulteration of drugs induced us to propose, a few years ago, to the Royal College of Physicians of London, in the course of certain negotiations relative to an Imperial or General Pharmacopœia for the Empire, that to the List of the *Materia Medica* there should be added a short statement of characters for ascertaining, that the leading articles are free from known sophistications, and of the due degree of purity for medical use. The suggestion has been partly adopted in the recent edition of the London Pharmacopœia; and in the present work we have endeavoured to carry more completely into effect the principles we propounded. In

judging of the attempt now made to enable practitioners and druggists to defend themselves and the public against the present notorious practices, it must be steadily kept in view, that our object has been to avoid all tedious or difficult methods of analysis ; to disregard those means which an accomplished chemist alone can turn to account ; and to adopt such characters only as may be applied with the aid of that ordinary knowledge of Practical Chemistry and Pharmacy, which, according to the rules of Universities and other medical institutions, every medical student ought now to acquire. Several of the formulæ, we are aware, are more defective than is desirable : For several articles of importance it has been found impracticable at present to furnish any simple and trustworthy characters : But nevertheless we have not hesitated to produce our inquiries as they stand, trusting their correction and extension to time, and the criticisms of competent authorities.

It has occurred to the College that, as the Chemical preparations now in use are prepared in the present day not by practitioners or druggists, but in a great measure by chemical manufacturers, who will scarcely be guided by Colleges in their processes, a considerable proportion of the Chemical formulæ might have been omitted without injury. On the whole, however, it has been thought advisable to put it in the power of every medical man to prepare his own chemical compounds if he chooses; and with that view we have taken care to select those processes which are the most simple and the most certain, though they may not be always either the cheapest or the most productive. As in former editions, so in the present Pharmacopœia, it has not been thought requisite to describe particularly the apparatus to be employed. At the same time it is right to mention here, that in one process of very frequent application, the process of distillation, complete success cannot be easily attained, especially on the small scale

without the substitution of a different apparatus for the retort and receiver hitherto commonly used in this country. In all operations, except where inorganic acids are to be distilled, it is greatly preferable to use a globular matrass, to which is fitted with a cork a tube cut obliquely at its lower end, curved above at a somewhat acute angle, and fitted at its other end to a refrigeratory. This refrigeratory consists of a long narrow cylinder slightly inclined to the horizon, and of a tube which passes along the centre of the cylinder, and is fixed at each end so that the space between them is air-tight; and by means of a funnel entering at the lower end of this interspace, and an exit tube from its upper extremity, a stream of cold water may be kept constantly running, by which refrigeration and the condensation of vapours within the inner tube are far more effectually accomplished than by any other mode that has hitherto been devised.

There is no department of our late inquiries which has given us less satisfac-

tion than that of the Nomenclature of the articles of the *Materia Medica*. We believe there are few physicians, and not many chemists, who now entertain any doubt that the Colleges committed a great error, when they were first seduced by the philosophical attractions of modern Chemical nomenclature, to abandon for the terms of scientific chemistry the trite names formerly used in Pharmacy and medical practice. The more decorous dress of science or philosophy has been dearly purchased at the cost of being compelled to follow the changing fashion of the day. We apprehend that practitioners will not submit much longer to the constant fluctuations which have been for some time forced upon them in pharmaceutic nomenclature. We have done our best to put a stop to this evil. The result has been necessarily a patchwork, of which we cannot boast, but which the public will probably receive in consideration of its convenience. A uniform nomenclature for pharmacy is now unattainable, unless, indeed, we were

to imitate the example of others by following the footsteps of chemistry through its changes and refinements.

We continue to employ the system of weights hitherto adopted by the Colleges of this country, namely, that commonly called Apothecaries' Weight. But at the same time we must confess we have never been able to see the force of those objections, which prevented the introduction of the Imperial system of weights into the practice of Medicine and Pharmacy, on the occasion of the late reform in the national weights and measures.

The following table represents the relations of the different apothecaries' weights, and the signs by which they are to be denoted in prescriptions.

one grain, gr.i

one scruple, \mathfrak{ss} = ... 20 grains.

one drachm, \mathfrak{z} i = ... \mathfrak{ss} iii = 60 grains.

one ounce, \mathfrak{z} i = ... \mathfrak{z} viii = 480 grains.

one pound, lbi = ... \mathfrak{z} xii = 5760 grains.

In former editions we directed that the

quantities of fluids, as well as solids, should be determined by weight. Measurement, however, is so much more convenient for fluids, that in practice it will always be followed; and we have therefore adopted in the present Pharmacopœia the system of measures lately introduced by the Royal College of Physicians of London, the basis of which is the Imperial Standard measure. The following table represents the denominations of this system of measures and their signs for use in prescriptions.

one minim,	m.	
one fluidrachm.	fʒ = ...	60 minims.
one fluidounce,	fʒ = ... fʒviii =	480 minims.
one pint,	O = ... fʒxx =	9600 minims.
one gallon,	C = ... O viii =	76,800 mins.

In using these measures it must be observed that the minim, fluidrachm, and fluidounce differ somewhat from those currently used till two years ago; so that most graduated vessels hitherto employed in Scotland

are not exactly available for the prescriptions and processes of this Pharmacopœia. It must likewise be remembered, that the denominations of measure now adopted bear no precise relation to the seemingly equivalent denominations of weight: That is, the minim, fluidrachm, and fluidounce, do not indicate by weight a grain, drachm, and ounce of water,—these weights being about a tenth more than the corresponding measures, or equal to $1 + \frac{17}{173}$ of the measures.

The substitution of measures instead of weights for ascertaining the quantities of fluids has rendered many alterations necessary in the apparent proportions of the ingredients in the formulas; but on due attention being paid to the densities of the liquids, and to the relation subsisting between weights and measures, it will be seen that in very few of the old formulæ has any material change been really made upon the strength of the preparations.

For ascertaining the densities of fluids,

we recommend the Hydrometer of Twaddell, or Lovi's Density Beads: and we understand the temperature to be taken at 60° of Fahrenheit's scale.

The account of the first number of the
 year of the first year of the
 year of the first year of the
 year of the first year of the

The account of the first number of the
 year of the first year of the
 year of the first year of the
 year of the first year of the

The account of the first number of the
 year of the first year of the
 year of the first year of the
 year of the first year of the

The account of the first number of the
 year of the first year of the
 year of the first year of the
 year of the first year of the

The account of the first number of the
 year of the first year of the
 year of the first year of the
 year of the first year of the

The account of the first number of the
 year of the first year of the
 year of the first year of the
 year of the first year of the

LIST
OF THE
ROYAL COLLEGE OF PHYSICIANS
OF EDINBURGH.

October 6, 1841.

RESIDENT FELLOWS
in the order of their Election.

* * * *A few Members since their election as
Resident Fellows have ceased to reside in
Edinburgh. Their present residences are
here indicated.*

Thomas Spens, *Treasurer and Councillor.*

James Home.

Thomas Charles Hope.

Alexander Monro.

Sir Alexander Morrison, (London.)

Joshua Henry Davidson.

William Fergusson, (Windsor.)

William Pulteney Alison, *Councillor.*

William Preston Lauder, (London.)

Walter Adam.
James George Playfair, (Florence.)
Robert Graham, *President*.
Robert Renton, *Councillor*.
Robert Christison, *Vice-President* and *Coun-
cillor*.
John Abercrombie.
George Augustus Borthwick, *Censor*.
Robert Carnegy, (Dundee.)
William Beilby.
John Macwhirter.
Thomas Shortt.
James Wood.
Richard Poole, (Montrose.)
Robert Grant, (London.)
John Thatcher.
William Gregory, (Aberdeen.)
John Thomson.
Robert Lewins, *Censor*.
David Boswell Reid, (London.)
John Mackenzie, (Kinellan, Dingwall.)
Montgomery Robertson, (Richmond.)
Andrew Combe.
David Craigie, *Secretary* and *Councillor*.
John Home Peebles, (Torquay.)
Peter Fairbairn.
Thomas Stewart Traill.
Jacob Dickson Hunter.
William Thomson, *Librarian*.
John Smith.
John Davie Morries Stirling, (Norway.)

James Paterson, (Downham, Norfolk.)
 Robert Spittal.
 Charles Ransford.
 Ralph Richardson, (Chester.)
 William Macdonald.
 William Seller.
 John Reid, (St Andrews.)
 James Young Simpson.
 William Reid.
 John Spens.
 James Cox.
 Charles Bell.
 John Moir.
 Martin Barry.
 George Paterson.
 William Henderson.
 James Marr.
 James Stark, *Keeper of Museum.*
 James Andrew.
 George Lund.
 Robert Bowes Malcolm.
 Alexander Wood.
 John Rose Cormack.
 Henry Lonsdale.

NON-RESIDENT FELLOWS
 in the order of their Election.

Robert Steavenson,	Newcastle.
Joseph Fox,	Falmouth.
James Macdonnell,	Belfast.
Alex. Philip Wilson Philip,	London.

George Dickson,	Berwickshire.
Andrew Keltie,	Perth.
George Browne Mill,	Bath.
Sir Alexander Wilson,	Bath.
Matthew Poole,	Waterford.
Samuel M'Dowall,	
Alexander Mackenzie,	Madras.
Sir James Macgrigor,	London.
James Proud Johnston,	Shrewsbury.
David Daniel Davies,	Sheffield.
John Bigsby,	Nottingham.
John Clark,	Dumfries-shire.
Samuel Fergusson,	
John Bowen,	
Benjamin Lara,	Portsea.
Sir George Magrath,	Plymouth.
William Beatty,	Plymouth.
George Drysdale,	
Sir William Pym,	London.
Sir Isaac Wilson,	Plymouth.
Sir D. J. Hamilton Dickson,	Plymouth.
Robert John Hume,	London.
John Ramsay,	Italy.
Stephen Macmullen,	Bridgewater.
James Gillies,	Bath.
Sir James R. Grant,	Cumberland.
John Butter,	Plymouth.
William Arnold,	Jamaica.
David Campbell,	Lancaster.
Thomas Kidd,	Jamaica.

Thomas Magrath,	Biggleswade.
Alexander Boyle,	
John Murray,	
James Mellis,	Calcutta.
James Macdonald,	London.
Peter Ramsay,	Liverpool.
John Price,	Llangollen.
Alexander George Home,	2d Drag. Guards.
Patrick Charles,	Putney, Surrey.
William Glover,	
James Burnes,	Bombay Army.
Archibald Robertson,	Northampton.
Thomas Burnford Harness,	Tavistock.
Samuel Hobart,	Cork.
John Tilstone,	Congalton, Cheshire.
Henry Atkinson,	Boulogne-sur- mer.
Edmund B. Lockyer.	
Patrick Rolland,	South America.
John M'Naught,	Jamaica.
James Lynch O'Connor,	Trinidad.
John Tomlinson Ingleby,	Birmingham.
Robert G. Holland,	Sheffield.
John Ward Dowsley,	Clonmel.
John Stevenson Bushnan,	Castle Cary, So- mersetshire.
Ralph Fawsett Ainsworth,	Manchester.
Andrew Henderson,	Royal Navy.
Thomas Radford,	Manchester.
Keith Imray,	Manchester

Evan Philip Cameron,	Berbice.
Thomas Richardson Colledge,	China.
John Willet Lavington,	Wilts.
George Hull,	Peckham.
Henry Hawkins,	Southampton.

LICENTIATES.

James Saunders.
Alexander Reid.

CONTENTS.

Preface,	Page v
Preface to the last Edition,	vii
List of the Members of the College,	xix
List of the Materia Medica,	1
Preparations and Compounds,	45
Acids,	47
Alcohol and Ethers,	54
Alkalis,	57
Alkaloids and their Salts,	61
Conserves and Electuaries,	68
Decoctions,	71
Distilled Waters,	75
Enemas,	78
Extracts,	79
Honeys,	88
Infusions,	89
Metals and their Compounds,	95
Mixtures and Emulsions,	121
Ointments, Liniments, and Cerates,	126
Oxidifiable non-metallic Elements,	134
Pills,	136
Plasters,	143
Powders,	148
Spirits,	151

Syrups,	Page 155
Tinctures,	161
Troches,	181
Volatile Oils,	184
Vinegars,	188
Wines,	190
Tests,	193
Index of changes in Nomenclature,	195
Latin Index of Preparations and Compounds,	199
English Index of Articles of the Materia Medica with their Preparations and Compounds,	206

LIST
OF THE
MATERIA MEDICA.

LIST

OF THE

MATHEMATICAL

LIST
OF THE
MATERIA MEDICA.

WITH FORMULAS FOR ASCERTAINING THE RE-
QUISITE PURITY OF THE MORE IMPORTANT
ARTICLES AND THEIR FREEDOM FROM KNOWN
ADULTERATIONS.

* * * IN the following List of the Materia Medica, the articles are arranged in alphabetical order, according to Pharmaceutic names, which have been chosen either for their convenience, or because they are familiarly employed, and without regard always to their correct designations in the Nomenclature of Chemistry or Natural History. The explanation of the names has been added, so that no one may be at a loss to understand their exact meaning. In this explanation reference is made in chemistry to the nomenclature in general use by scientific writers in Britain. In Botany reference respecting plants yielding

drugs is made to the author who first assigned the name adopted, if subsequent to Linnæus,—to the second edition of *Linnæus's* Species Plantarum, to that by *Willdenow*, to *Sprengel's* Systema Vegetabilium, and to *Decandolle's* Prodromus, as works of easy access,—and specially to any other works which it has been thought necessary or more advisable to quote.

ABSINTHIUM. Herb of *Artemisia Absinthium* (*L. W. Spr. DC*); *Wormwood*.

ACETUM BRITANNICUM. *British vinegar*.

Density 1006 to 1019. Sulphuretted hydrogen does not colour it. In four fluidounces complete precipitation takes place with 30 minims of Solution of nitrate of baryta. (See *Tests*).

ACETUM DESTILLATUM. *Distilled vinegar*.

Density 1005 : colourless : unaltered by sulphuretted hydrogen : one hundred minims neutralize 8 grains of carbonate of soda.

ACETUM GALLICUM. *French vinegar*.

Density 1014 to 1022. Ammonia in slight excess causes a purplish muddiness, and slowly a purplish precipitate. In four fluidounces complete precipitation takes place with 30 minims of solution of nitrate of baryta. (See *Tests*).

ACIDUM ACETICUM. *Acetic Acid*.

Density not above 1068.5, and increased

by 20 per cent of water : colourless : unaltered by sulphuretted hydrogen or nitrate of baryta : one hundred minims neutralize at least 216 grains of carbonate of soda.

ACIDUM BENZOICUM. *Benzoic Acid.*

Colourless : sublimed entirely by heat.

ACIDUM CITRICUM. *Citric acid.*

A solution in four parts of water is not precipitated by carbonate of potash : when incinerated with the aid of red oxide of mercury, no ash is left, or a mere trace.

ACIDUM HYDROCYANICUM. Hydrocyanic acid diluted with about thirty parts of water.

Hydrocyanic acid.

Solution of nitrate of baryta occasions no precipitate. Fifty minims diluted with one fluidounce of distilled water, agitated with 390 minims of Solution of nitrate of silver, and allowed to settle, will again give a precipitate with forty minims more of the test ; but a farther addition of the test after agitation and rest has no effect. The precipitate entirely disappears in boiling nitric acid.

ACIDUM MURIATICUM. *Hydrochloric acid of commerce.*

Density at least 1180. It is always yellow, and commonly contains a little sulphuric acid, oxide of iron, and chlorine.

ACIDUM MURIATICUM PURUM. *Hydrochloric acid.*

Density 1170 : nearly or entirely colourless :

without action on gold-leaf: If previously diluted with distilled water, it is not altered by solution of nitrate of baryta.

ACIDUM NITRICUM. *Nitric acid of commerce.*

Density 1380 to 1390; colourless or nearly so: if diluted with distilled water, it precipitates but slightly, or not at all, with solution of nitrate of baryta, or of nitrate of silver.

ACIDUM NITRICUM PURUM. *Pure nitric acid.*

Density 1500: colourless or pale-yellow: unaffected by solution of nitrate of silver, or nitrate of baryta, if previously diluted with distilled water.

ACIDUM PYROLIGNEUM. Diluted acetic acid, obtained by the destructive distillation of wood. *Pyroligneous acid.*

Density at least 1034: nearly or entirely colourless: unaffected by sulphuretted hydrogen, or solution of nitrate of baryta: one hundred minims neutralize at least 53 grains of carbonate of soda.

ACIDUM SULPHURICUM. *Sulphuric acid of commerce.*

Density 1840 or near it: colourless: when diluted with its own volume of water, only a scanty muddiness arises, and no orange fumes escape: when diluted with twelve volumes of water, sulphuretted hydrogen causes a white muddiness, but not a yellow precipitate.

ACIDUM SULPHURICUM PURUM. *Sulphuric acid.*

Density 1845 : colourless : dilution causes no muddiness : solution of sulphate of iron shows no reddening at the line of contact, when poured over it.

ACIDUM TARTARICUM. *Tartaric acid.*

When incinerated with the aid of red oxide of mercury, it leaves no residuum, or a mere trace only.

ACONITUM. Leaves of Aconitum Napellus (*L. W. DC. Spr.*); *Monkshood.*

AERUGO. Commercial diacetate of copper. *Verdigris.*

It is dissolved in a great measure by muriatic acid, not above five per cent of impurity being left.

AETHER SULPHURICUS. *Sulphuric ether.*

Density 735 or under ; when agitated in a minim measure with half its volume of concentrated solution of muriate of lime, its volume is not lessened,

ALCOHOL. *Alcohol : Absolute alcohol.*

Density 794-6 : when mixed with a little solution of nitrate of silver and exposed to bright light, it remains unchanged, or only a very scanty dark precipitate forms.

ALLIUM. Bulb of Allium sativum (*L. W. Spr.*); *Garlic.*

ALOE BARBADENSIS. Extract or inspissated juice of the leaves of one or more undeter-

- mined species of Aloe (*L. W. Spr.*) *Barbadoes aloes*.
- ALOE INDICA. From one or more undetermined species of Aloe (*L. W. Spr.*) *East Indian aloes*.
- ALOE SOCOTORINA. Probably inspissated juice of an undetermined species of Aloe (*L. W. Spr.*) *Socotorine aloes*.
In thin pieces translucent and garnet-red: almost entirely soluble in spirit of the strength of sherry. *Very rare*.
- ALTHÆÆ FOLIA. Leaves of *Althæa officinalis* (*L. W. DC. Spr.*); *Marsh-mallow*.
- ALTHÆÆ RADIX. Root of *Althæa officinalis* (*L. W. DC. Spr.*) *Marsh-mallow*.
- ALUMEN. Sulphate of alumina and potash.
Alum.
Not subject to adulteration.
- AMMONIACUM. Gummy-resinous exudation of *Dorema Ammoniacum* (*Don in Linn. Trans. xvi.*); *Ammoniac*.
- AMMONIÆ CARBONAS. Sesquicarbonate of ammonia. *Carbonate of ammonia*.
Heat sublimes it entirely: a solution in water, when treated with nitric acid in excess, does not precipitate with solution of nitrate of baryta or nitrate of silver.
- AMMONIÆ MURIAS. Hydrochlorate of ammonia. *Sal-ammoniac*.
Not liable to adulteration.
- AMMONIÆ SPIRITUS. Solution of ammonia in rectified spirit. *Spirit of ammonia*.

It has a density about 845, and a strong ammoniacal odour : it does not effervesce with diluted muriatic acid.

AMYGDALA AMARA. Kernel of *Amygdalus communis*, var. α (DC.) ; *Bitter almond*.

AMYGDALA DULCIS. Kernel of *Amygdalus communis*, var. β and γ (DC.) ; *Sweet almond*.

AMYLUM. Fecula of the seeds of *Triticum vulgare* (*Villars, Delph.—Willd. Hort. Berol.—Spr.*) ; *Starch*.

ANETHUM. Fruit of *Anethum graveolens* (L. W. DC.) ; *Dill*.

ANGELICA. Root of *Angelica Archangelica* (L. W. Spr.) ; *Angelica*.

ANISUM. Fruit of *Pimpinella Anisum* (L. W. DC.) ; *Anise*.

ANTHEMIS. Simple flowers of *Anthemis nobilis* (L. W. Spr. DC.) ; *Chamomile*.

ANTIMONII OXIDUM. *Sesquioxide of antimony*.

Entirely soluble in muriatic acid, and also in a boiling mixture of water and bitartrate of potash : snow-white : fusible at a full-red heat.

ANTIMONII SULPHURETUM. Native sesquisulphuret of antimony. *Sulphuret of antimony*. Entirely soluble in muriatic acid with the aid of heat.

ANTIMONII SULPHURETUM AUREUM. A mixture or compound of sesquisulphuret of

antimony, sesquioxide of antimony, and sulphur. *Golden sulphuret of antimony.*

Tasteless: twelve times its weight of muriatic acid aided by heat will dissolve most of it, forming a colourless solution, and leaving a little sulphur.

ANTIMONIUM TARTARIZATUM. Tartrate of potash and antimony; *Tartar-emetic.*

Entirely soluble in twenty parts of water: solution colourless, and not affected by solution of ferrocyanide of potassium: a solution in forty parts of water is not affected by its own volume of a solution of eight parts of acetate of lead in thirty-two parts of water and fifteen parts of acetic acid.

AQUA. *Spring water.*

For pharmaceutic use spring water must be so far at least free of saline matter as not to possess the quality of hardness, or contain above a 6000th of solid matter.

AQUA DESTILLATA. *Distilled water.*

Free of colour and odour: unaltered by sulphuretted hydrogen, or solution of nitrate of silver, nitrate of baryta, or oxalate of ammonia.

AQUA AMMONIÆ. Diluted aqueous solution of ammonia. *Ammonia.*

Density 960; diluted nitric acid occasions no effervescence: when saturated with nitric acid, it is not precipitated by solution of nitrate of silver.

AQUA AMMONIAE FORTIOR. Concentrated aqueous solution of ammonia. *Strong ammonia.*

Density 880: one fluidounce with $2\frac{1}{2}$ fluidounces of water makes Aqua Ammoniae, for which other characters are given above.

AQUA AMMONIAE ACETATIS. Diluted aqueous solution of acetate of ammonia.

Without action on litmus: density 1011: free of colour or odour: Solution of potash disengages an ammoniacal, sulphuric acid an acetous, odour: unaffected by solution of nitrate of silver.

AQUA POTASSAE. Diluted aqueous solution of potash. *Solution of potash.*

Density 1072: colourless: sulphuric acid does not occasion effervescence.

ARGENTUM. *Silver; Virgin or pure silver.*

Soluble entirely in diluted nitric acid: this solution, treated with an excess of muriate of soda, gives a white precipitate entirely soluble in Aqua ammoniae, and a fluid which is not affected by sulphuretted hydrogen.

ARGENTI NITRAS. *Nitrate of silver.*

Soluble in distilled water with the exception of a very scanty black powder; twenty-nine grains dissolved in one fluidounce of distilled water, acidulated with nitric acid, precipitated with a solution of nine grains of muriate of ammonia, briskly agitated for a few seconds, and then allowed to rest a little,

will yield a clear supernatant liquid, which still precipitates with more of the test.

ARMORACIA. Fresh root of *Cochlearia Armoracia* (*L. W. DC. Spr.*); *Horse-radish*.

ARSENICUM ALBUM. Sesquioxide of arsenic; *Arsenious acid*.

Entirely sublimed by heat.

ASSAFOETIDA. Gummy-resinous exudation of *Ferula Assafoetida* (*L. W. Spr. DC.*) and probably *Ferula persica* (*W. Spr. DC.*); *Assafœtida*.

AURANTII AQUA. Distilled water of the flowers of *Citrus vulgaris* (*Risso, Annales du Museum, xx. DC.*), and sometimes of *Citrus Aurantium* (*Ibid.*); *Orange-flower water*.

Nearly colourless: unaffected by sulphuretted hydrogen.

AURANTII CORTEX. Rind of the fruit of *Citrus vulgaris* (*Risso, Annales du Museum, xx. DC.*); *Bitter orange rind*.

AURANTII OLEUM. Volatile oil of the flowers of *Citrus vulgaris* (*Risso, Annales du Museum, xx. DC.*), and sometimes of *Citrus Aurantium* (*Ibid.*); *Neroli oil*.

AVENA. Seeds of *Avena sativa* (*L. W. Spr.*)

AXUNGIA. Fat of *Sus Scrofa*; *Axunge*.

BALSAMUM CANADENSE. Fluid resinous exudation of *Abies balsamea* (*Marsh, Arb. Amer.*); *Canada Balsam*.

BALSAMUM PERUVIANUM. Fluid balsamic

exudation of *Myrospermum peruiferum* (*DC.*); *Peru Balsam.*

BALSAMUM TOLUTANUM. Concrete balsamic exudation of *Myrospermum toluiferum* (*DC. Spr.*); *Tolu Balsam.*

BARYTAE CARBONAS. *Carbonate of baryta.*
One hundred grains dissolved in an excess of nitric acid are not entirely precipitated with 124 grains of sulphate of magnesia.

BARYTAE MURIAS. *Chloride of barium.*
One hundred grains in solution are not entirely precipitated by 100 grains of sulphate of magnesia.

BARYTAE SULPHAS. *Sulphate of baryta : heavy spar.*

White or flesh-red : heavy : lamellar ; brittle.

BELLADONNA. Leaves of *Atropa Belladonna* (*L. W. Spr.*); *Deadly Nightshade.*

BENZOINUM. Concrete balsamic exudation of *Styrax Benzoin* (*Dryand. in Phil. Trans. — W. Spr.*); *Benzoin.*

BERGAMOTAE OLEUM. Volatile oil of the rind of the fruit of *Citrus Limetta* (*Risso in Ann. du Museum, xx. DC.*); *Oil of Bergamot.*

BISMUTHUM. *Bismuth.*

Its powder is entirely soluble in nitric acid with the aid of heat ; and the solution is colourless or nearly so, and deposits a white powder when much diluted with cold water.

BISMUTHUM ALBUM. *Trisnitrate of bismuth.*
It forms a colourless solution with nitric

acid, and without effervescence : not subject to adulteration.

BORAX. Borate of soda. *Borax.*

A hot concentrated solution, if treated with sulphuric acid, deposits copious scaly crystals on cooling. Not subject to adulteration.

BUCKU. Leaves of various species of *Barosma* (*W. in Hort. Berol.*) ; *Bucku.*

CAJUPUTI OLEUM. Volatile oil of the leaves of *Melaleuca minor* (*Smith in Rees's Cycl. DC.*) ; *Oil of Cajuput.*

CALAMINA PRÆPARATA. Levigated impure carbonate of zinc ; *Calamine.*

CALAMUS AROMATICUS. Rhizoma of *Acorus Calamus*, var. *α*, vulgaris (*L. W.*) ; *Sweet Flag.*

CALCIS MURIAS. [CRYSTALLIZATUM.] *Hydrochlorate of lime.*

Extremely deliquescent : a solution of 76 grains in one fluidounce of distilled water, precipitated by 49 grains of oxalate of ammonia, remains precipitable by more of the test.

CALOMELAS. Chloride of mercury. *Calomel.*

Heat sublimes it without any residuum : sulphuric ether agitated with it, filtered, and then evaporated to dryness, leaves no crystalline residuum, and what residuum may be left is not turned yellow with *Aqua potassæ*.

CALUMBA. Root of *Cocculus palmatus* (*DC.*) ; *Calumba.*

Its infusion is precipitated by infusion of

galls, rendered blue by iodine, and not altered by sesquioxide of iron.

CALX. *Lime.*

It is slaked by water; muriatic acid then dissolves it entirely, without any effervescence; and the solution does not precipitate with ammonia in excess.

CALX CHLORINATA. Hypochlorite of lime.
Chloride of lime.

Pale grayish-white: dry: 50 grains are nearly all soluble in two fluidounces of water, forming a solution of the density 1027, and of which 100 measures treated with an excess of oxalic acid give off much chlorine, and if then boiled and allowed to rest 24 hours yield a precipitate which occupies nineteen measures of the liquid.

CAMBOGIA [SIAMENSIS]. Gum-resin from an unascertained plant inhabiting Siam, probably a species of Hebradendron (*Graham, ut infra*); *Siam Gamboge.*

Fracture somewhat conchoidal, smooth, and glistening: a decoction of its powder cooled is not rendered green by tincture of iodine, but merely somewhat tawny.

CAMBOGIA [ZEYLANICA.] Gummy-resinous exudation of Hebradendron cambogioides (*Graham in Comp. to Bot. Mag. ii.*); *Ceylon Gamboge.*

CAMPHORA. Camphor of Camphora officinarum (*Nees von Esenbeck, Laurineæ.*)
Camphor.

Its powder evaporates entirely when gently heated.

CANELLA. Bark of *Canella alba* (*Murr. Syst.—W. DC. Spr.*); *Canella*.

CANTHARIS. *Cantharis vesicatoria*,—the whole fly; *Cantharides*.

CAPSICUM. Fruit of *Capsicum annum* (*L. W. Spr.*) and other species; *Capsicum*; *Chillies*.

CARBO ANIMALIS. Impure animal charcoal obtained commonly from bones: *Ivory-black*.

CARBO ANIMALIS PURIFICATUS. *Animal charcoal*; *Purified ivory-black*.

When incinerated with its own volume of red oxide of mercury, it is dissipated, leaving only a scanty ash.

CARBO LIGNI. *Charcoal*.

CARDAMOMUM. Fruit of *Renealmia Cardamomum* (*Roscoe, Monandrous Plants.*); *Cardamoms*.

CARUI. Fruit of *Carum Carui* (*L. W. Spr. DC.*); *Caraway*.

CARYOPHYLLUS. Dried undeveloped flower of *Caryophyllus aromaticus* (*L. DC.*); *Clove*.

CARYOPHYLLI OLEUM. Volatile oil of the undeveloped flowers of *Caryophyllus aromaticus* (*L. DC.*); *Oil of Cloves*.

CASCARILLA. Bark probably of *Croton Eleuteria* (*Swartz, Fl. Ind. Occident.—W. Spr.*), and possibly of other species of the same genus; *Cascarilla*.

CASSIAE CORTEX. Bark of *Cinnamomum Cassia* (*Blume, Bijdrag tot de Flora van Nederl. Ind.—Hayne, Darstellung, &c. xii.*); *Cassia-bark.*

CASSIAE OLEUM. Volatile oil of the bark of *Cinnamomum Cassia* (*Blume, &c. ut supra*); *Oil of Cassia.*

CASSIAE PULPA. Pulp of the pods of *Cassia Fistula* (*L. W. Spr. DC.*); *Cassia-pulp.*

CASTOREUM. Castor: a peculiar secretion from the præputial follicles of Castor fiber.

CATECHU. Extract of the wood of *Acacia Catechu* (*W. DC. Spr.*), of the kernels of *Areca Catechu* (*L. W. Spr.*), and of the leaves of *Uncaria Gambier* (*Roxburgh, Fl. Indica. DC.*), probably too from other plants; *Catechu.*

The finest qualities yield to sulphuric ether 53, and the lowest qualities 28 per cent of tannin dried at 280°.

CENTAURIUM. The flowering heads of *Erythraea Centaurium* (*Persoon, Synopsis. Spr.*); *Common Centaury.*

CERA ALBA. Bleached Bees'-wax. *White-wax.*

CERA FLAVA. Waxy secretion of *Apis mellifica*; *Bees'-wax.*

CETACEUM. Cetine of *Physeter macrocephalus*, nearly pure; *Spermaceti.*

CETRARIA. *Cetraria islandica* (*Achar. Synopsis.*); *Iceland-moss.*

CHIRETTA. Herb and root of *Agathotes Chi-*

rayta (*Don in Lond. and Ed. Phil. Mag.* 1836.); *Chiretta*.

CHLORINEI AQUA. Concentrated watery solution of chlorine, with a little sulphate of soda; *Chlorine-water*.

CINCHONA CORONAE. Bark of Cinchona Condaminea, (*Humboldt et Bonp. Pl. Aeq. —Spr. DC.*); *Crown-Bark*.

CINCHONA CINEREA. Bark of Cinchona micrantha (*Ruiz and Pavon in Fl. Peruv. —DC.*); *Gray-Bark: Silver-Bark*.

CINCHONA FLAVA. Bark of an unascertained species of Cinchona (*L. W. Spr. DC.*) *Yellow-Bark*.

A filtered decoction of 100 grains in two fluidounces of distilled water gives, with a fluidounce of concentrated solution of carbonate of soda, a precipitate, which when heated in the fluid becomes a fused mass, weighing when cold 2 grains or more, and easily soluble in solution of oxalic acid.

CINCHONA RUBRA. Bark of an undetermined species of Cinchona (*L. W. Spr. DC.*) *Red-Bark*.

CINNABARIS. Bisulphuret of mercury. *Cinnabar*.

It is sublimed entirely by heat, and without any metallic globules being formed.

CINNAMOMUM. Bark of Cinnamomum zeylanicum (*Nees von Esenbeck, Laurineæ. Hayne's Darstellung, &c.*); *Cinnamon*.

CINNAMOMI OLEUM. Volatile oil of the bark of *Cinnamomum zeylanicum* (*Nees, ut supra*); *Oil of Cinnamon*.

Cherry-red when old, wine-yellow when recent: odour purely cinnamomic; nitric acid converts it nearly into a uniform crystalline mass.

COCCI. *Coccus cacti*; the entire insects; *Cochineal*.

COCCULUS. Fruit of *Anamirta Cocculus* (*Wight and Arnott, Flora Penins. Ind. Or.*); *Cocculus Indicus*.

The kernels should fill at least two-thirds of the fruit.

COLCHICI CORMUS. The Cormus of *Colchicum autumnale* (*L. W. Spr.*); *Colchicum-bulb*.

COLCHICI SEMINA. Seeds of *Colchicum autumnale* (*L. W. Spr.*); *Colchicum-seeds*.

COLOCYNTHIS. Pulp of the fruit of *Cucumis Colocynthis* (*L. W. Spr. DC.*); *Colocynth*.

CONIUM. Leaves of *Conium maculatum* (*L. W. Spr. DC.*); *Hemlock*.

The powder triturated with *Aqua potassæ* exhales a powerful odour of conia.

COPAIBA. Fluid resinous exudation of various species of *Copaifera* (*L. W. DC. Spr.*); *Copaiva*.

Transparent: free of turpentine odour when heated: soluble in two parts of alcohol: it dissolves a fourth of its weight of carbonate of magnesia, with the aid of a gentle heat, and continues translucent.

COPAIBAE OLEUM. Volatile oil of Copaiva.

See Copaiba.

CORIANDRUM. Fruit of *Coriandrum sativum* (*L. W. Spr. DC.*); *Coriander*.

CORNU. Horn of *Cervus Elaphus*.

CREASOTUM. *Creasote*.

Colourless, and remains so under sunshine : density 1066 : entirely and easily soluble in its own volume of acetic acid : a drop on white filtering paper heated for ten minutes about 212° leaves no translucent stain.

CRETA. Friable carbonate of lime : *Chalk*.

CRETA PRÆPARATA. Chalk, finely pulverized by levigation. *Prepared chalk*.

A solution of 25 grains in ten fluidrachms of pyroligneous acid, when neutralized by carbonate of soda, and precipitated by 32 grains of oxalate of ammonia, continues precipitable after filtration by more of the test.

CROCUS. The stigmata of *Crocus sativus* (*Allioni, Fl. Ped.—DC. Flore Franc.*); *Saffron*.

CROTONIS OLEUM. Expressed oil of the seeds of *Croton Tiglium* (*W. Spr.*); *Croton-oil*.

When agitated with its own volume of pure alcohol and gently heated, it separates on standing, without having undergone any apparent diminution.

CUBEBAE. Fruit of *Piper Cubeba* (*L. Suppl.—W. Spr.*); *Cubebs*.

CUMINUM. Fruit of *Cuminum Cyminum* (*L. W. Spr. DC.*); *Cumin*.

CUPRI SULPHAS. *Sulphate of Copper.*

Not subject to adulteration.

CUPRUM AMMONIATUM. Sulphate of copper and ammonia; *Ammoniacal sulphate of copper.*

CURCUMA. Rhizoma of *Curcuma longa* (*L. W. Spr.*); *Turmeric.*

CUSPARIA. Bark of *Galipea officinalis* (*Hancock in Trans. of Med. Botan. Soc.*); *Angustura-bark.*

Its outer surface is not turned dark-green, nor its transverse fracture red, by nitric acid.

DAUCI RADIX. Root of *Daucus Carota*, var. *sativa* (*L. W. DC.*); *Common Carrot.*

DIGITALIS. Leaves of *Digitalis purpurea* (*L. W. Spr.*); *Foxglove.*

DULCAMARA. Twigs of *Solanum Dulcamara* (*L. W. Spr.*); *Bittersweet.*

ELATERIUM. Feculence of the juice of the fruit of *Momordica Elaterium* (*L. W. Spr. DC.*); *Elaterium.*

Colour pale gray: when exhausted by rectified spirit, the solution, concentrated and poured into hot diluted *Aqua potassæ*, deposits, on cooling, minute silky, colourless crystals weighing from a seventh to a fourth of the elaterium.

ELEMI. Concrete resinous exudation from one or more unascertained plants. *Elemi.*

ERGOTA. An undetermined fungus, with degenerated seed of *Secale cereale* (*L. W. Spr.*); *Ergot of rye.*

EUPHORBIIUM. Concrete resinous juice of undetermined species of *Euphorbia* (*L. W. Spr.*); *Euphorbium*.

FARINA. Flour of the seeds of *Triticum vulgare* (*Villars, Delph.—W. in Hort. Berol.—Spr.*); *Flour*.

FERRI CARBONAS SACCHARATUM. Carbonate of protoxide of iron in an undetermined state of combination with sugar and sesquioxide of iron. *Saccharine carbonate of iron*. Colour grayish-green; easily soluble in muriatic acid, with brisk effervescence.

FERRI FILUM. *Iron-wire*.

FERRI IODIDUM. *Iodide of iron: protiodide of iron*.

Entirely soluble in water, or nearly so; forming a greenish solution.

FERRI IODIDI SYRUPUS. Solution of Iodide of iron in syrup. *Syrup of Iodide of iron*.

Colourless, or pale-green; transparent; without sediment even when exposed to air.

FERRI LIMATURA. *Iron-filings*.

FERRI MURIATIS TINCTURA. Solution of sesquichloride of iron in rectified spirit. *Tincture of iron*.

FERRI OXIDUM NIGRUM. Ferroso-ferric oxide (*Berzelius*): a compound of protoxide and sesquioxide of iron. *Black oxide of iron*.

Dark grayish-black: strongly attracted by the magnet: heat expels water from it: muriatic acid dissolves it entirely; and ammonia precipitates a black powder from this solution.

FERRI OXIDUM RUBRUM. Sesquioxide of iron: *Red oxide of iron.*

Entirely soluble in muriatic acid, aided by gentle heat.

FERRISULPHAS. Sulphate of protoxide of iron. *Sulphate of iron.*

Pale bluish-green crystals, with little or no efflorescence.

FERRI SULPHURETUM. Protosulphuret of Iron. *Sulphuret of iron.*

Soluble in a great measure in diluted sulphuric acid, with effervescence and disengagement of sulphuretted hydrogen-gas.

FERRUGO. Hydrated sesquioxide of iron. *Rust of iron.*

Entirely and very easily soluble in muriatic acid, without effervescence: if previously dried at 180°, a stronger heat drives off about 18 per cent of water: the magnet does not attract it.

FERRUM TARTARIZATUM. Tartrate of potash and sesquioxide of iron. *Tartrate of iron.*

Entirely soluble in cold water: taste feebly chalybeate: the solution is not altered by Aqua potassæ, and not precipitated by solution of ferrocyanide of potassium.

FICI. Dried fruit of Ficus Carica (*L. W. Spr.*); *Figs.*

FILIX. Rhizoma of Nephrodium Filix-mas (*Richard, Botan. Méd.*): *Male Shield Fern.*

FOENICULUM. Fruit of Fœniculum officinale (*Allioni, Fl. Ped.*); *Fennel.*

GALBANUM. Concrete gummy-resinous exudation of an imperfectly ascertained umbelliferous plant, probably a species of *Opoidia* (*Lindley, Bot. Reg.* 1839); *Galbanum*.

GALLAE. Excrescences of *Quercus infectoria* [*W. Spr.*]; formed by *Diplolepis gallæ tinctorum* (*Olivier, Voyage*); *Galls*.

GENTIANA. Root of *Gentiana lutea* (*L. W. Spr.*); *Gentian*.

GLYCYRRHIZÆ RADIX. Root of *Glycyrrhiza glabra* (*L. W. DC. Spr.*); *Liquorice-root*.

GLYCYRRHIZÆ EXTRACTUM. Extract of the root of *Glycyrrhiza glabra* (*L. W. DC. Spr.*) *Liquorice*.

GOSSYPIUM. Hairs attached to the seeds of *Gossypium herbaceum* (*L. W. DC. Spr.*) and other species of the genus: *Raw cotton*.

GRANATI RADIX. Root-bark of *Punica Granatum* (*L. W. DC. Spr.*); *Pomegranate-bark*.

GUAIACI LIGNUM, Wood of *Guaiacum officinale* (*L. W. DC. Spr.*); *Lignum-vitæ*.

GUAIAACUM. Resin obtained by heat from the wood of *Guaiacum officinale* (*L. W. DC. Spr.*); *Guaiac*.

Fresh fracture red, slowly passing to green: the tincture slowly strikes a lively blue colour on the inner surface of a thin paring of a raw potato.

GUMMI ACACIAE. Gum of various species of *Acacia* (*W. DC. Spr.*); *Gum-Arabic*.

HAEMATOTOXYLON. Wood of *Hæmatoxylon campechianum* (*L. W. DC. Spr.*) ; *Log-wood.*

HELLEBORUS. Root of *Helleborus niger* (*L. W. DC. Spr.*) ; *Black Hellebore.*

HORDEUM. Decorticated seeds of *Hordeum distichon* (*L. W. Spr.*) ; *Pearl-Barley.*

HYDRARGYRUM. *Mercury.*

Entirely sublimed by heat : a globule moved along a sheet of paper leaves no trail : pure sulphuric acid agitated with it evaporates when heated without leaving any residuum.

HYDRARGYRI BINIODIDUM. *Biniodide of mercury.*

Entirely vaporizable : soluble entirely in 40 parts of a concentrated solution of muriate of soda at 212° ; and again deposited in fine red crystals on cooling.

HYDRARGYRI OXIDUM RUBRUM. *Binoxide of mercury. Red Precipitate.*

Entirely soluble in muriatic acid : heat decomposes and sublimes it entirely in metallic globules, without any discharge of nitrous fumes.

HYDRARGYRI PRECIPITATUM ALBUM. *Chloride of mercury and ammonia. White Precipitate.*

HYOSCYAMUS. Leaves of *Hyoscyamus niger* (*L. W. Spr.*) ; *Henbane.*

IODINEUM. *Iodine.*

Entirely vaporizable : Thirty-nine grains,

with nine grains of quicklime and three ounces of water, when heated short of ebullition, slowly form a perfect solution, which is yellowish or brownish, if the iodine be pure, but colourless if there be above two per cent of water or other impurity.

IPECACUANHA. Root of *Cephaelis Ipecacuanha* (*Richard, Hist. Ipec.—DC. Spr.*); *Ipecacuan.*

JALAPA. Root of *Ipomæa Purga* (*Nees von Esenbeck, Plantæ Medic.—Hayne's Darstellung, &c.*); *Jalap.*

JUNIPERI CACUMINA. Tops of *Juniperus communis* (*L. W. Spr.*); *Juniper-tops.*

JUNIPERI FRUCTUS. Berries of *Juniperus communis* (*L. W. Spr.*); *Juniper-berries.*

JUNIPERI OLEUM. Volatile oil of the fruit of *Juniperus communis* (*L. W. Spr.*) *Oil of Juniper.*

KINO. Concrete exudation of *Pterocarpus erinaceus* (*Lam. Encyc. DC.*) and of other undetermined Genera and species; *Kino.*

KRAMERIA. Root of *Krameria triandra*. (*Ruiz and Pavon in Flor. Peruv.—DC. Spr.*); *Rhatany-root*

LACMUS. A peculiar colouring matter from *Roccella tinctoria* (*Acharius Lichenog. Univ.*); *Litmus.*

LACTUCARIUM. Inspissated juice of *Lactuca virosa* (*L. W. Spr. DC.*) and *sativa* (*Ibid.*); *Lettuce-Opium.*

- LAURO-CERASUS. Leaves of *Prunus lauro-cerasus* (*L. W. Spr. DC.*); *Cherry-laurel*.
- LAVANDULA. The flowering heads of *Lavandula vera* (*DC. Flore Fran.*); *Lavender*.
- LAVANDULAE OLEUM. Volatile oil of the flowering heads of *Lavandula vera* (*DC. Flore Franc.*); *Oil of Lavender*.
- LIMONES. Fruit of *Citrus medica* and *Citrus Limonum* (*Risso, Ann. du Mus. xx. DC.*); *Lemons and Limes*.
- LIMONUM CORTEX. Rind of the fruit of *Citrus medica* (*Risso, ut supra.—DC.*); *Lemon-peel*.
- LIMONUM OLEUM. Volatile oil of the rind of the fruit of *Citrus medica* (*Risso, ut supra, DC.*) *Oil of Lemons*.
- LINI SEMINA. Seeds of *Linum usitatissimum* (*L. W. DC. Spr.*); *Linseed*.
- LINI FARINA. Meal of the seeds of *Linum usitatissimum* (*L. W. DC. Spr.*) deprived of their fixed oil by expression. *Linseed-meal*.
- LINI OLEUM. Expressed oil of the seeds of *Linum usitatissimum* (*L. W. DC. Spr.*) *Linseed-oil*.
- LINUM CATHARTICUM. Herb of *Linum catharticum* (*L. W. DC. Spr.*); *Purging Flax*.
- LITHARGYRUM. Protoxide of lead, partially fused; *Litharge*.
Fifty grains dissolve entirely, without effervescence, in a fluidounce and a half of pyro-ligneous acid; and the solution, precipitated

by 53 grains of phosphate of soda, remains precipitable by more of the test.

LOBELIA. Herb of *Lobelia inflata* (*L. W. Spr.*); *Lobelia*.

LUPULUS. Catkin of *Humulus Lupulus* (*L. W. Spr.*); *Hops*.

MAGNESIA. *Magnesia*.

Fifty grains are entirely soluble, without effervescence, in a fluidounce of muriatic acid: an excess of ammonia occasions in the solution only a scanty precipitate of alumina: the filtered fluid is not precipitated by solution of oxalate of ammonia.

MAGNESIAE CARBONAS. *Carbonate of magnesia*.

When dissolved in an excess of muriatic acid, an excess of ammonia occasions only a scanty precipitate of alumina; and the filtered fluid is not precipitated by oxalate of ammonia.

MAGNESIAE SULPHAS. *Sulphate of magnesia*.

Ten grains dissolved in a fluidounce of water and treated with solution of carbonate of ammonia, are not entirely precipitated by 280 minims of Solution of phosphate of soda.

[See *Tests*.]

MALVA. Herb of *Malva sylvestris* (*L. W. DC. Spr.*); *Common Mallow*.

MANGANESII OXIDUM. Native impure peroxide of manganese. *Oxide of manganese*. Muriatic acid aided by heat dissolves it al-

most entirely, disengaging chlorine : heat disengages oxygen.

MANNA. Sweet concrete exudation probably from several species of *Fraxinus* (*L. W. Spr.*) and *Ornus* (*Persoon, Synopsis*); *Manna*.

MARANTA. Fecula of the tubers of *Maranta arundinacea* (*L. W. Spr.*) and *Maranta indica* (*De Tussac, Journ. Bot.—Spr.*); *Arrow-root*.

MARMOR. Massive crystalline carbonate of lime; *White marble*.

A neutral solution in diluted nitric acid, precipitated by an excess of oxalate of ammonia, and filtered, yields no white precipitate with phosphate of ammonia.

MASTICHE. Concrete resinous exudation of *Pistacia Lentiscus* (*L. W. DC. Spr.*); *Mastic*.

MEL. Saccharine secretion of *Apis mellifica*. *Honey*.

MELISSA. Herb of *Melissa officinalis* (*L. W. Spr.*); *Balm*.

MENTHA PIPERITA. Herb of *Mentha piperita* (*L. W. Spr.*); *Peppermint*.

MENTHÆ PIPERITÆ OLEUM. Volatile oil of *Mentha piperita* (*L. W. Spr.*)

MENTHA VIRIDIS. Herb of *Mentha viridis* (*L. W. Spr.*); *Spearmint*.

MENYANTHES. Leaves of *Menyanthes trifoliata* (*L. W. Spr.*); *Buckbean*.

MEZEREON. Root-bark of *Daphne Mezereon* (*L. W. Spr.*); *Mezereon*.

MORPHIAE ACETAS. *Acetate of morphia.*

One hundred measures of a solution of ten grains in half a fluidounce of water and five minims of acetic acid, heated near to 212° and decomposed by a faint excess of ammonia, yield by agitation a precipitate which in 24 hours occupies 15.5 measures of the liquid.

MORPHIAE MURIAS. *Hydrochlorate of morphia.*

Snow-white: entirely soluble: solution colourless: loss of weight at 212° not above 13 per cent: one hundred measures of a solution of 10 grains in half a fluidounce of water, heated near to 212° , and decomposed with agitation by a faint excess of ammonia, yield a precipitate which in 24 hours occupies 12.5 measures of the liquid.

MOSCHUS. Inspissated secretion in the follicle of the prepuce of *Moschus moschiferus*.
Musk.

MUCUNA. Hairs from the pod of *Mucuna pruriens* (*DC.*); *Cowitch.*

MYRISTICA. Kernel of the fruit of *Myristica officinalis* (*L. Suppl.*); *Nutmeg.*

MYRISTICAE ADEPS. Concrete expressed oil from the kernel of the fruit of *Myristica officinalis*. (*Linn. Suppl.*); *Concrete oil of nutmeg.*

MYRISTICAE OLEUM. Volatile oil from the

kernel of the fruit of *Myristica officinalis* (*L. Suppl.*); *Volatile oil of Nutmeg.*

MYRRHA. Gummy-resinous exudation of *Balsamodendron* (*Protium?*) *Myrrha* (*Nees von Esenbeck, Plantæ Medic.*); *Myrrh.*

NUX-VOMICA. Seeds of *Strychnos Nux-vomica* (*L. W. Spr.*); *Nux-vomica.*

OLIVAE OLEUM. Expressed oil of the pericarp of *Olea europea* (*L. W. Spr.*); *Olive-oil.*

When carefully mixed with a twelfth of its volume of solution of nitrate of mercury prepared as for the *Unguentum Citrinum*, it becomes in three or four hours like a firm fat, without any separation of liquid oil.

OPIUM. Concrete juice from the unripe capsules of *Papaver somniferum* (*L. W. DC. Spr.*); *Opium.*

A solution from 100 grains of fine opium macerated 24 hours in two fluidounces of water, filtered, and strongly squeezed in a cloth, if treated with a cold solution of half an ounce of carbonate of soda in two waters, yields a precipitate, which weighs when dry at least ten grains, and dissolves entirely in solution of oxalic acid.

ORIGANUM. Herb of *Origanum vulgare* (*L. W. Spr.*); *Marjoram.*

OVUM. Egg of *Phasianus gallus.*

PAPAVER. Capsules of *Papaver somniferum.* (*L. W. DC. Spr.*) not quite ripe; *Poppy-heads.*

- PAREIRA. Root of *Cissampelos Pareira*. (*L. W. DC. Spr.*); *Pareira*.
- PETROLEUM. *Petroleum: rock-oil*.
- PIMENTA. Unripe berries of *Eugenia Pimenta* (*DC.*); *Pimento*.
- PIPER LONGUM. Dried spikes of *Piper longum* (*L. W. Spr.*); *Long pepper*.
- PIPER NIGRUM. Dried unripe berries of *Piper nigrum* (*L. W. Spr.*); *Black pepper*.
- PIX ARIDA. Pitch: from various species of *Pinus* (*L. W. Spr.*) and *Abies* (*Lam. Enc. Meth.*)
- PIX BURGUNDICA. Concrete resinous exudation probably in a great measure from *Abies excelsa* (*Lam. in Enc. Method. vi. 518.*); *Burgundy Pitch*.
- PIX LIQUIDA. Tar: from various species of *Pinus* (*L. W. Spr.*) and *Abies* (*Lam. Enc. Meth.*)
- PLUMBI ACETAS. *Acetate of lead*.
Entirely soluble in distilled water acidulated with acetic acid: Forty-eight grains thus dissolved are not entirely precipitated by a solution of 30 grains of phosphate of soda.
- PLUMBI DIACETATIS AQUA. Solution of diacetate of lead. *Goulard's extract*.
A copious precipitate is gradually formed when the breath is propelled through it by means of a tube.
- PLUMBI CARBONAS. *Carbonate of lead*.
It does not lose weight at a temperature of

212°: Sixty-eight grains are entirely dissolved in 150 minims of acetic acid diluted with a fluidounce of distilled water; and the solution is not entirely precipitated by a solution of 60 grains of phosphate of soda.

PLUMBI IODIDUM. *Iodide of lead.*

Bright yellow: five grains are entirely soluble, with the aid of ebullition, in one fluidrachm of pyroligneous acid diluted with a fluidounce and a-half of distilled water; and golden crystals are abundantly deposited on cooling.

PLUMBI NITRAS. *Nitrate of lead.*

Not subject to adulteration.

PLUMBI OXIDUM RUBRUM. A compound of protoxide and peroxide of lead. *Red oxide of lead.*

Entirely soluble in highly fuming nitrous acid; partially soluble in diluted nitric acid, a brown powder being left.

POTASSA. Protoxide of potassium: *Potash.*

Boiling water commonly leaves oxide of iron undissolved, which should not exceed 1.25 per cent: the solution supersaturated with nitric acid gives a faint precipitate with solution of nitrate of baryta, and more with solution of nitrate of silver,—owing to the presence of impurities.

POTASSA CUM CALCE. A mixture of potash and lime. *Potash-and-lime caustic.*

POTASSAE ACETAS. *Acetate of potash.*

Not subject to adulteration.

POTASSAE AQUA EFFERVESCENS. Solution of bicarbonate of potash, surcharged with carbonic acid : *Kali-water*.

POTASSAE BICARBONAS. *Bicarbonate of potash*.

A solution in 40 parts of water does not give a brick-red precipitate with solution of corrosive sublimate ; and when supersaturated with nitric acid, is not affected by solution of nitrate of baryta or nitrate of silver.

POTASSAE BISULPHAS. *Bisulphate of potash*.

A solution in eight waters effervesces briskly with alkaline carbonates.

POTASSAE BITARTRAS. *Bitartrate of potash*.

Entirely soluble in 40 parts of boiling water : Forty grains in solution are neutralized with 30 grains of crystallized carbonate of soda ; and when then precipitated by 70 grains of nitrate of lead, the liquid remains precipitable by more of the test.

POTASSAE CARBONAS. Carbonate of potash not quite pure, obtained by lixiviating, evaporating, and granulating by fusion and refrigeration the potashes of commerce. *Potashes*.

One hundred grains lose not more than twenty on exposure to a red heat : and when dissolved and supersaturated by pure diluted nitric acid, the solution gives a faint haze only with solution of nitrate of baryta, and is

entirely precipitated by 100 minims of solution of nitrate of silver. (See *Tests*.)

POTASSAE CARBONAS PURUM. *Carbonate of Potash.*

It does not lose weight at a low red heat; and a solution supersaturated with pure nitric acid is precipitated either faintly, or not at all, by solution of nitrate of baryta or nitrate of silver.

POTASSAE NITRAS. *Nitrate of potash.*

Entirely soluble: its solution is not affected by solution of nitrate of baryta, and faintly or not at all, by solution of nitrate of silver.

POTASSAE SULPHAS. *Sulphate of potash.*

Not subject to adulteration.

POTASSAE SULPHAS CUM SULPHURE. Nature undetermined.

POTASSAE TARTRAS. *Tartrate of potash.*

Entirely and easily soluble in four parts of boiling water: solution neutral, and yielding a crystalline precipitate with muriatic acid: 44 grains in solution are not entirely precipitated by 55 grains of nitrate of lead.

POTASSAE ET SODAE TARTRAS. *Tartrate of potash and soda.*

Entirely and easily soluble in five parts of boiling water: muriatic acid occasions a crystalline precipitate in a strong solution: 37 grains in solution are not entirely precipitated by 43 grains of nitrate of lead.

POTASSII FERROCYANIDUM. *Ferrocyanide of potassium : ferrocyanate of potash.*

Not subject to adulteration.

POTASSII IODIDUM. *Iodide of potassium : hydriodate of potash.*

Its solution is not affected, or is merely rendered hazy, by solution of nitrate of baryta : a solution of five grains in a fluidounce of distilled water, precipitated by an excess of solution of nitrate of silver, and then agitated in a bottle with a little Aqua ammoniæ, yields quickly by subsidence a clear supernatant liquid, which is not altered by an excess of nitric acid or is rendered merely hazy.

POTASSII SULPHURETUM. A mixture of sulphate of potash with persulphuret of potassium. *Sulphuret of potash.*

PRUNA. Dried fruit of *Prunus domestica* (*L. W. DC. Spr.*) ; *Prunes.*

PTEROCARPUS. Wood of *Pterocarpus santalinus* (*L. Sup. W. DC. Spr.*) ; *Red Sandalwood.*

PULEGIUM. Herb of *Mentha Pulegium* (*L. W. Spr.*) ; *Pennyroyal.*

PULVIS ANTIMONIALIS. A mixture chiefly of antimonious acid and phosphate of lime, with some sesquioxide of antimony, and a little antimonite of lime ; *Antimonial powder.*

Distilled water, boiled with it and filtered, gives with sulphuretted hydrogen an orange precipitate : muriatic acid digested with the

residue becomes yellow, does not become turbid by dilution, but gives a copious orange precipitate with sulphuretted hydrogen.

PYRETHRUM. Root of *Anacyclus Pyrethrum* (DC.); *Pellitory of Spain*.

PYROLA. Herb of *Chimaphila umbellata* (Nuttal, *Gen. Spr.*); *Pyrola*.

QUASSIA. Wood chiefly of *Picraena excelsa* (Lindley, *Fl. Med.*) seldom of *Quassia amara* (L. *Supp. W. DC. Spr.*); *Quassia*.

QUERCUS CORTEX. Bark of *Quercus pedunculata* (W. *Spr.*); *Oakbark*.

QUINAE SULPHAS. *Sulphate of Quina*.

A solution of 10 grains in a fluidounce of distilled water, and two or three drops of sulphuric acid, if decomposed by a solution of half an ounce of carbonate of soda in two waters, and heated till the precipitate shrinks and fuses, yields on cooling a solid mass, which when dry weighs 7.4 grains, and in powder dissolves entirely in solution of oxalic acid.

RESINA. Residue of the distillation of the turpentine of various species of *Pinus* (L. *W. Spr.*) and *Abies* (Lam. *Enc. Meth.*) *Resin*.

RHAMNI BACCAE. Fruit of *Rhamnus catharticus* (L. *W. DC. Spr.*); *Buckthorn*.

RHEUM. Root of an undetermined species of *Rheum* (L. *W. Spr.*); *Rhubarb*.

RHOEAS. Petals of *Papaver Rhœas* (L. *W. DC. Sp.*); *Corn-poppy*.

- RICINI OLEUM. Expressed oil of the seeds of *Ricinus communis* (*L. W. Spr.*); *Castor-oil*. It is entirely dissolved by its own volume of alcohol.
- ROSA CENTIFOLIA. Petals of *Rosa centifolia* (*L. W. DC. Spr.*); *Damask-rose*.
- ROSA GALLICA. Petals of *Rosa gallica* (*L. W. DC. Spr.*); *Red-rose*.
- ROSAE FRUCTUS. Hip of *Rosa canina*, (*L. W. DC. Spr.*), and of several allied species, deprived of the carpels. *Hips*.
- ROSAE OLEUM. Volatile oil of the petals of *Rosa centifolia* (*L. W. DC. Spr.*); *Attar of Roses*.
- ROSMARINUS. Tops of *Rosmarinus officinalis* (*L. W. Spr.*); *Rosemary*.
- RUTA. Leaves and unripe fruit of *Ruta graveolens* (*L. W. DC. Spr.*); *Rue*.
- RUTAE OLEUM. Volatile oil of *Ruta graveolens* (*L. W. DC. Spr.*); *Oil of Rue*.
- SABADILLA. Fruit of *Veratrum Sabadilla* (*Retz. Obs.—W. Spr.*) of *Helonias officinalis* (*Don in Edin. Phil. Journ. 1832*), and probably of other *Melanthaceæ*. *Cevadilla*.
- SABINA. Tops of *Juniperus Sabina* (*L. W. Spr.*); *Savin*.
- SACCHARUM COMMUNE. Impure sugar, from *Saccharum officinarum*, (*L. W. Spr.*); *Muscovado*.
- SACCHARI FAEX. Concentrated uncrystallizable juice of *Saccharum officinarum* (*L. W. Spr.*); *Treacle*.

SACCHARUM PURUM. Pure sugar, from Saccharum officinarum (*L. W. Spr.*); *White sugar.*

SAGO. Farina from the interior of the trunk of various Palmaceæ and species of Cycas. (*L. W. Spr.*); *Sago.*

SALICIS CORTEX. Bark of Salix Caprea (*L. W. Spr.*); *Willow-bark.*

SAMBUCUS. Flowers of Sambucus nigra (*L. W. DC. Spr.*); *Elder-flowers.*

SAPO DURUS. *Spanish or Castile soap*, made with olive oil and soda.

White: it does not stain paper, is free of odour, and dissolves entirely in rectified spirit.

SAPO MOLLIS. *Soft soap*, made with olive oil and potash.

SARZA. Root of Smilax officinalis (*Humb. et Bonpl. Nov. Gen. i.—Spr.*) and probably other species; *Sarsaparilla.*

SASSAFRAS. Root of Sassafras officinale (*Nees und Ebermaier, Handb.*); *Sassafras.*

SCAMMONIUM. Gummy-resinous exudation from incisions into the root of Convolvulus Scammonia (*L. W. Spr.*); *Scammony.*

Fracture glistening, almost resinous, if the specimen be old and dry: muriatic acid does not cause effervescence on its surface: the decoction of its powder, filtered and cooled, is not rendered blue by tincture of iodine. Sulphuric ether separates at least eighty per cent of resin dried at 280°.

SCILLA. Bulb of *Squilla maritima* (*Steinheil in Ann. des Sc. Natur. 2ème Sér. vi.*); *Squill.*

SCOPARIUM. Tops of *Cytisus Scoparius* (*DC.*); *Broomtops.*

SENEGA. Root of *Polygala Senega* (*L. W. DC. Spr.*); *Seneka Snake-root.*

SENNA ALEXANDRINA. Leaves of various species of *Cassia*, probably of *Cassia lanceolata* (*Forskal, Flora Ægypt. Arab.*) *Cassia acutifolia* (*Delile, Egypte*), and *Cassia obovata* (*Colladon—DC. Spr.*); *Alexandrian Senna.*

As imported, it also contains an abundant admixture of leaves of *Cynanchum Argel* (*Delile. DC. Spr.*); which ought to be removed as far as possible by picking.

SENNA INDICA. Leaves of *Cassia elongata* (*Lemaire-Lisancourt, Journ de Pharm. vii.*); *East Indian senna, var. Tinnivelly.* Leaves for the most part large, unbroken, and free of brownness or blackening.

SERPENTARIA. Root of *Aristolochia Serpentaria* (*L. W. Spr.*); *Virginian Snakeroot.*

SEVUM. Fat of *Ovis Aries*; *Suet.*

SIMARUBA. Root-bark of *Simaruba amara* (*Aublet. Guian.*); *Simaruba-root.*

SINAPI. Flour of the seeds of *Sinapis nigra* (*L. W. DC. Spr.*), generally mixed with those of *Sinapis alba*, (*Ibid.*) and deprived of fixed oil by expression; *Mustard.*

A decoction allowed to cool is not turned blue with tincture of iodine.

SODAE AQUA EFFERVESCENS. Solution of bi-carbonate of soda surcharged with carbonic acid; *Soda-water*.

SODAE BICARBONAS. *Bi-carbonate of soda.*
A solution in 40 parts of water does not give an orange precipitate with solution of corrosive sublimate.

SODAE CARBONAS. *Carbonate of soda (crystallized.)*

A solution of 21 grains in a fluidounce of distilled water, precipitated by 19 grains of nitrate of baryta, remains precipitable by more of the test; and the precipitate is entirely soluble in nitric acid. Little subject to adulteration.

SODAE MURIAS. Impure commercial chloride of sodium. *Salt.*

SODAE MURIAS PURUM. *Chloride of sodium.*
A solution is not precipitated by solution of carbonate of ammonia followed by solution of phosphate of soda: a solution of 9 grains in distilled water is not entirely precipitated by a solution of 26 grains of nitrate of silver.

SODAE PHOSPHAS. *Phosphate of soda.*

An efflorescent salt: 45 grains dissolved in two fluidounces of boiling distilled water, and precipitated by a solution of 50 grains of carbonate of lead in a fluidounce of pyroligneous acid, will remain precipitable by solution of acetate of lead.

SODAE SULPHAS. *Sulphate of soda.*

Not subject to adulteration.

SPIGELIA. Root of *Spigelia marilandica* (L. W. Spr.); *Carolina-pink.*

SPIRITUS AETHERIS NITRICI. Nitric (Hyponitrous) ether with four volumes of rectified spirit. *Spirit of nitrous ether.*

Density 847 : it effervesces feebly, or not at all, with solution of bi-carbonate of potash : when agitated with twice its volume of concentrated solution of muriate of lime, 12 per cent of ether slowly separates.

SPIRITUS AETHERIS SULPHURICI. Sulphuric ether with alcohol. *Spirit of sulphuric ether.*

Density 809 : it does not affect litmus-paper, or render water muddy : when agitated with twice its volume of a concentrated solution of muriate of lime, 28 per cent of ether separates by rest.

SPIRITUS RECTIFICATUS. *Rectified spirit.*

Density 838 (56 over proof) : Four fluid-ounces treated with 25 minims of Solution of nitrate of silver, exposed to bright light for twenty-four hours, and then passed through a filter purified by weak nitric acid, so as to separate the black powder which forms, undergo no farther change when again exposed to light with more of the test.

SPIRITUS TENUIOR. *Proof spirit.*

Density 912 (7 over proof) : Tests otherwise as for rectified spirit.

SPONGIA. *Spongia officinalis*; *Sponge*.

STANNUM. *Tin*.

When finely granulated, 100 grains are entirely converted into a white powder by three fluidrachms of nitric acid (D. 1380); and distilled water, boiled with this powder and filtered, is colourless, and precipitates but faintly, or not at all with solution of sulphate of magnesia.

STAPHISAGRIA. Seeds of *Delphinium Staphisagria* (L. W. DC. Spr.); *Stavesacre*.

STRAMONIUM. Herb of *Datura Stramonium* (L. W. Spr.); *Thornapple*.

STRYCHNIA. *Strychnia*. Always more or less impure.

Intensely bitter: nitric acid strongly reddens it: a solution of 10 grains in 4 fluidrachms of water by means of a fluidrachm of pyroligneous acid, when decomposed by one fluidounce of concentrated solution of carbonate of soda, yields on brisk agitation a coherent mass, weighing when dry 10 grains, and entirely soluble in solution of oxalic acid.

STYRAX. Balsamic exudation of *Styrax officinale* (L. W. Spr.); *Storax*.

SUBLIMATUS CORROSIVUS. Bichloride of mercury. *Corrosive-sublimate*.

It sublimes entirely by heat; and its powder is entirely and easily soluble in sulphuric ether.

SULPHUR. *Sulphur.*

It is entirely sublimed by heat; and distilled water agitated with it does not affect litmus-paper. When nitric acid is heated with it, the solution diluted with water, neutralized with carbonate of soda, and acidulated with muriatic acid, does not give a yellow precipitate with sulphuretted-hydrogen.

TABACUM. Leaves of *Nicotiana Tabacum* (*L. W. Spr.*); *Tobacco.*

TAMARINDUS. Pulp of the pods of *Tamarindus indica* (*L. W. DC. Spr.*); *Tamarind-pulp.*

TAPIOCA. Fecula of the root of *Janipha Manihot* (*Humb. and Bonpl. Nov. Gen. et Spec. ii.—Spr.*); *Tapioca.*

TARAXACUM. Root of *Taraxacum Dens-leonis* (*Desfontaines, Fl. Atlant.—DC.*); *Dandelion.*

TEREBINTHA CHIA. Liquid resinous exudation of *Pistacia Terebinthus* (*L. W. DC.*); *Chian turpentine.*

TEREBINTHINA VENETA. Liquid resinous exudation of *Abies Larix* (*Lam. Illustr.*); *Venice turpentine.*

TEREBINTHINAE OLEUM. Volatile oil of the liquid resinous exudation of various species of *Pinus* (*L. W. Spr.*) and *Abies* (*Lam. Enc. Meth.*); *Oil of turpentine.*

TORMENTILLA. Root of *Potentilla Tormentilla* (*Sibthorpe, Flor. Oxon.—DC. Spr.*); *Tormentil.*

TRAGACANTHA. Gummy exudation from Astragalus gummifer (*Labillardière, W. DC. Spr.*) and probably Astragalus verus (*Olivier, Voyage,—DC.*) and other species; *Tragacanth.*

UVAE PASSAE. Dried fruit of *Vitis vinifera* (*L. W. DC. Spr.*); *Raisins.*

UVA-URSI. Leaves of *Arctostaphylos Uva-ursi* (*Spr.*); *Bear-berry.*

VALERIANA. Root of *Valeriana officinalis* (*L. W. Spr. DC.*); *Valerian.*

VERATRUM. Rhizoma of *Veratrum album* (*L. W. Spr.*); *White Hellebore.*

VINUM ALBUM. *Sherry.*

VIOLA. Flowers of *Viola odorata* (*L. W. DC. Spr.*) *Violet.*

ZINCI OXIDUM. *Oxide of Zinc.*

White: tasteless: entirely soluble in diluted nitric acid without effervescence: this solution is not affected by nitrate of baryta, but gives with ammonia a white precipitate entirely soluble in an excess of the test.

ZINCI SULPHAS. *Sulphate of zinc.*

When a solution in six waters is boiled with a little nitric acid, and solution of ammonia is then added till the oxide of zinc first thrown down is all redissolved, no yellow precipitate remains, or a trace only, and the solution is colourless.

ZINCUM. *Zinc.*

It dissolves in a great measure in diluted sulphuric acid, leaving only a scanty gray-

ish-black residuum : this solution presents the characters just given for the solution of sulphate of zinc.

ZINGIBER. Rhizoma of *Zingiber officinale*.
(*Roscoe in Trans. Lin. Soc. viii.—Spr.*) ;
Ginger.

OMITTED.

CANNA. Fecula of the root of an imperfectly determined species of canna (*L. W. Spr.*)
Tous-les-mois.

PREPARATIONS

AND

COMPOUNDS.

THE ALIEN

COMPANIES

PREPARATIONS

AND

COMPOUNDS.

ACIDS.

ACETUM DESTILLATUM.

Take of Vinegar (French, by preference) eight parts : distil over with a gentle heat seven parts : dilute the product, if necessary, with distilled water, till the density is 1005.

ACIDUM ACETICUM.

Take of Acetate of lead any convenient quantity : heat it gradually in a porcelain basin by means of a bath of oil or fusible metal, (8 tin, 4 lead, 3 bismuth) to 320° F. ; and stir till the fused mass concretes again : pulverize this when cold, and heat the powder again to 320°, with frequent stirring, till the particles cease to accrete. Add six ounces of the powder to nine fluidrachms and a half of Pure sulphuric acid

contained in a glass-mattrass : attach a proper tube and refrigeratory ; and distil from a fusible-metal-bath with a heat of 320° to complete dryness. Agitate the distilled liquid with a few grains of red oxide of lead to remove a little sulphurous acid, allow the vessel to rest a few minutes, pour off the clear liquor, and redistil it. The density is commonly from 1063 to 1065, but must not exceed 1068.5

ACIDUM BENZOICUM.

Take of Benzoin any convenient quantity : put it into a glass-mattrass ; and by means of a gradually increasing heat sublime as long as any thing rises : squeeze the sublimate between folds of filtering-paper to remove the oil as much as possible ; and sublime the residuum again.

ACIDUM CITRICUM.

Take of Lemon-juice, four pints ;
Prepared Chalk, four ounces and a half, or a sufficiency ;
Diluted Sulphuric acid, thirty-six fluidounces, or in the same proportion to the chalk required.

Boil the Lemon-juice, allow it to rest, pour off the clear liquor, boil this again, and add the Chalk to it while hot by degrees till there is no more effervescence, and the liquor ceases to taste acid. Collect the

precipitate, and wash it with hot water till the water passes from it colourless. Squeeze the residuum in a powerful press; mix it uniformly with two pints of distilled water; and then add the Sulphuric acid by degrees and with constant stirring. Try whether a small portion of the liquid, when filtered, gives with solution of nitrate of baryta a precipitate almost entirely soluble in nitric acid; and if the precipitate is not nearly all soluble, add a little citrate of lime to the whole liquor till it stand this test. Separate now the clear liquor by subsidence or filtration, washing the insoluble matter with cold water, and adding the washings to the liquor: concentrate with a gentle heat till crystals form on the surface: set the liquor aside to cool and crystallize; and purify the crystals by repeated solution and crystallization till they are colourless.

ACIDUM HYDROCYANICUM.

Take of Ferrocyanide of Potassium, three ounces;

Sulphuric acid, two fluidounces;

Water, sixteen fluidounces.

Dissolve the salt in eleven fluidounces of the water, and put the solution into a matrass with a little sand: add the acid previously diluted with five fluidounces of the water and allowed to cool: connect the matrass with a

proper refrigeratory : distil with a gentle heat, by means of a sand-bath or naked gas-flame, till fourteen fluidounces pass over, or till the residuum begins to froth up. Dilute the product with distilled water till it measures sixteen fluidounces.

ACIDUM MURIATICUM PURUM.

Purify Muriate of soda by dissolving it in boiling water, concentrating the solution, skimming off the crystals as they form on the surface, draining from them the adhering solution as much as possible, and subsequently washing them with cold water slightly. Take of this salt, previously well dried, of Pure sulphuric acid, and of water, equal weights. Put the salt into a glass retort, and add the acid previously diluted with a third part of the water and allowed to cool. Fit on a receiver containing the rest of the water. Distil with a gentle heat by means of a sand-bath or naked gas-flame so long as any liquid passes over, preserving the receiver constantly cool by snow or a stream of cold water.

ACIDUM MURIATICUM DILUTUM.

Take of Muriatic Acid, four fluidounces ;
Distilled water, twelve fluidounces.
Mix them together ; the density of this preparation is 1050.

ACIDUM NITRICUM PURUM.

Purify Nitrate of potash, if necessary, by two or more crystallizations till nitrate of silver does not act on its solution in distilled water. Put into a glass retort equal weights of this purified nitrate and of sulphuric acid; and distil into a cool receiver with a moderate heat from a sand-bath or naked gas-flame so long as the fused material continues to give off vapour. The pale-yellow acid thus obtained may be rendered colourless, should this be thought necessary, by heating it gently in a retort.

ACIDUM NITRICUM DILUTUM.

Mix together one fluidounce of Pure Nitric acid (D. 1500), and nine fluidounces of distilled water. If the Commercial Nitric acid of D. 1390 be used, one fluidounce and five fluidrachms and a-half are required.—The density of this diluted acid is 1077.

ACIDUM SULPHURICUM PURUM.

If commercial sulphuric acid contain nitrous acid, heat eight fluidounces of it with between ten and fifteen grains of sugar, at a temperature not quite sufficient to boil the acid, till the dark colour at first produced shall have nearly or altogether disappeared. This process removes nitrous acid. Other impurities may be removed by distillation;

which on the small scale is easily managed by boiling the acid with a few platinum chips in a glass retort by means of a sand-bath or gas-flame,—rejecting the first half ounce.

ACIDUM SULPHURICUM DILUTUM.

Mix together one fluidounce of sulphuric acid and thirteen fluidounces of water. The density of this preparation is about 1090.

ACIDUM SULPHURICUM AROMATICUM.

Take of Sulphuric Acid (commercial,) three fluidounces and a-half;

Rectified spirit, one pint and a-half;

Cinnamon, in moderately fine powder, an ounce and a-half;

Ginger, in moderately fine powder, one ounce;

Add the acid gradually to the spirit, let the mixture digest at a very gentle heat for three days in a closed vessel; mix the powders, moisten them with a little of the acid spirit, let the mass rest for twelve hours, and then put it into a percolator and transmit the rest of the acid spirit. This preparation may also be made by digesting the powders for six days in the acid spirit, and then straining the liquor.

ACIDUM TARTARICUM.

Take of Bitartrate of potash, four pounds;

Boiling distilled water, two gallons
and a-half ;

Prepared chalk, twenty-five ounces
and six drachms ;

Diluted sulphuric acid, ten pints and
seven fluidounces ;

Muriatic acid, twenty-six fluidounces
and a-half, or a sufficiency.

Boil the bitartrate with two gallons of the water, and add gradually half the chalk, constantly stirring : when the effervescence is over, add a solution obtained by dissolving the rest of the chalk in the muriatic acid diluted with four pints of the water. After the tartrate of lime has subsided, pour off the liquid, and wash the tartrate with distilled water till it is tasteless. Then pour the diluted sulphuric acid on the tartrate, and boil for fifteen minutes. Evaporate with a gentle heat to obtain crystals. Purify these by repeated solution, filtration, and crystallization.

ALCOHOL AND ETHERS.

ALCOHOL.

Take of Rectified-spirit, one pint.

Lime, eighteen ounces.

Break down the Lime into small fragments : expose the Spirit and lime together to a gentle heat in a glass matrass till the lime begins to slake : withdraw the heat till the slaking is finished, preserving the upper part of the matrass cool with damp cloths. Then attach a proper refrigeratory, and with a gradually increasing heat distil off seventeen fluidounces. The density of this alcohol should not exceed 796 : If higher, the distillation must have been begun before the slaking of the lime was finished.

AETHER SULPHURICUS.

Take of Rectified-spirit, fifty fluidounces.

Sulphuric acid, ten fluidounces.

Pour twelve fluidounces of the Spirit gently over the Acid contained in an open vessel, and then stir them together briskly and thoroughly. Transfer the mixture immediately into a glass matrass connected with a refrigeratory, and raise the heat quickly to about 280° . As soon as the ethereal fluid begins to distil over, supply fresh Spirit through a tube into the matrass in a continuous stream, and in such quantity as to equal

that of the fluid which distils over. This is best accomplished by connecting one end of the tube with a graduated vessel containing the spirit,—passing the other end through a cork fitted into the matrass,—and having a stop-cock on the tube to regulate the discharge. When forty-two ounces have distilled over and the whole spirit has been added, the process may be stopped. Agitate the impure ether with sixteen fluidounces of a saturated solution of muriate of lime, containing about half an ounce of lime recently slaked. When all odour of sulphurous acid has been thus removed, pour off the supernatant liquor, and distil it with a very gentle heat so long as the liquid which passes over has a density not above 735. More ether of the same strength is then to be obtained from the solution of muriate of lime. From the residuum of both distillations a weaker ether may be obtained in small quantity, which must be rectified by distilling it gently again.

SPIRITUS AETHERIS SULPHURICI.

Take of Sulphuric ether, a pint.

Rectified-spirit, two pints.

Mix them. The density of this preparation ought to be 809.

SPIRITUS AETHERIS NITRICI.

Take of Rectified-spirit, two pints and six fluidounces ;

Pure nitric acid (D. 1500) seven fluidounces.

Put fifteen fluidounces of the spirit, with a little clean sand, into a two-pint matrass, fitted with a cork, through which are passed a safety-tube terminating an inch above the spirit, and another tube leading to a refrigerator. The safety-tube being filled with pure nitric acid, add through it gradually three fluidounces and a-half of the acid. When the ebullition which slowly rises is nearly over, add the rest of the acid gradually, half a fluidounce at a time, waiting till the ebullition caused by each portion is nearly over before adding more, and cooling the refrigerator with a stream of water, iced in summer. The ether thus distilled over, being received in a bottle, is to be agitated first with a little milk of lime, till it ceases to redden litmus-paper, and then with half its volume of concentrated solution of muriate of lime. The pure hyponitrous ether thus obtained, which should have a density of 899, is then to be mixed with the remainder of the rectified spirit, or exactly four times its volume.

Spirit of nitric ether ought not to be kept long, as it always undergoes decomposition, and becomes at length strongly acid. Its density by this process is 847.

SPIRITUS DILUTOR. See *Spirits*.

ALKALIS.

AMMONIAE AQUA, et

AMMONIAE AQUA FORTIOR.

Take of Muriate of ammonia, thirteen ounces ;

Quicklime, thirteen ounces ;

Water, seven fluidounces and a half ;

Distilled water, twelve fluidounces.

Slake the Lime with the water, cover it up till it cool, triturate it well and quickly with the Muriate of ammonia previously in fine powder, and put the mixture into a glass retort, to which is attached a receiver with a safety-tube. Connect with the receiver a bottle also provided with a safety-tube, and containing four ounces of the Distilled water, but capable of holding twice as much. Connect this bottle with another loosely corked, and containing the remaining eight ounces of distilled water. The communicating tubes must descend to the bottom of the bottles at the further end from the retort ; and the receiver and bottles must be kept cool by snow, ice, or a running stream of very cold water. Apply to the retort a gradually-increasing heat till gas ceases to be evolved ; remove the retort, cork up the aperture in the receiver where it was connected with the re-

tort, and apply to the receiver a gentle and gradually-increasing heat, to drive over as much of the gas in the liquid contained in it, but as little of the water, as possible. Should the liquid in the last bottle not have the density of 960, reduce it with some of the Stronger Aqua ammoniæ in the first bottle, or raise it with distilled water, so as to form Aqua ammoniæ of the prescribed density.

AMMONIAE CARBONAS.

Take of Sal-ammoniac, one pound ;

Chalk, one pound and a-half.

Reduce them separately to fine powder, mix them thoroughly, and subject the mixture in a retort with a proper receiver to a gradually-increasing heat so long as any vapours sublime.

AMMONIAE CARBONATIS AQUA.

Take of Carbonate of Ammonia, four ounces

Distilled water, one pint.

Dissolve the salt in the water.

AMMONIAE ACETATIS AQUA.

Take of Distilled vinegar, (from French vinegar in preference) twenty-four fluidounces ;

Carbonate of ammonia, one ounce.

Mix them and dissolve the salt. If the solution has any bitterness, add by degrees a

little distilled vinegar till that taste be removed. The density of the distilled vinegar should be 1005, and that of the Aqua acetatis ammoniæ 1011.

POTASSA.

Take any convenient quantity of Aqua potassæ; evaporate it in a clean and covered iron vessel, increasing gradually the heat, till an oily-looking fluid remains, a drop of which, when removed on a rod, becomes hard on cooling. Then pour out the liquid upon a bright iron plate, and as soon as it solidifies, break it quickly and put it into glass bottles secured with glass stoppers.

POTASSAE AQUA.

Take of Carbonate of potash (dry) four ounces;

Lime recently burnt, two ounces;

Water, forty-five fluidounces.

Let the Lime be slaked and converted into milk of lime with seven fluidounces of the water. Dissolve the Carbonate in the remaining thirty-eight fluidounces of water; boil the solution, and add to it the milk of lime in successive portions, about an eighth at a time,—boiling briskly for a few minutes after each addition. Pour the whole into a deep narrow glass-vessel for twenty-four hours; and then withdraw with a syphon the

clear liquid, which should amount to at least thirty-five fluidounces, and ought to have a density of 1072.

POTASSA CUM CALCE.

Take any convenient quantity of Aqua potassæ; evaporate it in a clean covered iron vessel to one-third of its volume; add slaked Lime till the fluid has the consistence of firm pulp: preserve the product in carefully covered vessels.

ALKALOIDS AND THEIR SALTS.

MORPHIAE ACETAS.

Take of Muriate of morphia any convenient quantity. Dissolve it in fourteen times its weight of warm water, and when the solution is cool add Aqua ammoniæ gradually and with constant agitation until there is a permanent but faint odour of ammonia in the fluid. Collect the precipitate on a calico filter, wash it moderately with cold water, and dissolve it by means of a slight excess of Pyroligneous acid in twelve parts of warm water for every part of muriate of morphia that was used. Concentrate the solution over the vapour-bath and set it aside to crystallize. Drain and squeeze the crystals, and dry them with a gentle heat. More acetate of morphia may be obtained on concentrating the mother liquor.

MORPHIAE MURIAS.

Take of Opium, twenty ounces ;
Water, eight pints ;
Muriate of lime, one ounce, or a
slight excess.

Macerate the opium in fragments for twenty-four hours in two pints of the water ; and separate the infusion, squeezing well the re-

sidue. Repeat the maceration successively with two pints more of the water till the whole is made use of. Concentrate the whole infusions over the vapour bath to one pint, and add the Muriate of lime dissolved in four fluidounces of water. Set the whole aside to settle ; pour off the liquid ; wash the sediment with a little water, adding the washings to the liquid. Evaporate the liquid sufficiently in the vapour bath for it to solidify on cooling. Subject the cooled mass to very strong pressure in a cloth ; redissolve the cake in a sufficiency of warm distilled water ; add a little fine powder of white marble, and filter ; acidulate the filtered fluid with a very little muriatic acid ; and concentrate a second time in the vapour-bath for crystallization. Subject the crystals again to very strong pressure in a cloth. Repeat the process of solution, clarification by marble and muriatic acid, concentration, and crystallization, until a snow-white mass be obtained.

On the small scale trouble and loss are saved by decolorizing the solution of muriate of morphia by means of a little purified animal charcoal after two crystallizations. But on the large scale it is better to purify the salt by repeated crystallizations alone, and to treat all the expressed fluids, except the first, in the same way with the original solution, of impure muriate of morphia. An ad-

ditional quantity of salt may often be got from the first dark and resinous fluid obtained by expression, on merely allowing it to remain at rest for a few months, when a little muriate of morphia may be deposited in an impure condition.

The opium, which yields the largest quantity of precipitate by carbonate of soda according to the formula in p. 29, yields muriate of morphia not only in greatest proportion, but likewise with the fewest crystallizations.

MORPHIAE MURIATIS SOLUTIO.

Take of Muriate of morphia, one drachm and a half;

Rectified-spirit, five fluidounces;

Distilled water, fifteen fluidounces.

Mix the spirit and water, and dissolve the muriate of morphia in the mixture with the aid of a gentle heat.

QUINAE SULPHAS.

Take of Yellow Bark in coarse powder, one pound;

Carbonate of soda, eight ounces;

Sulphuric acid, half a fluidounce;

Purified Animal Charcoal, two drachms.

Boil the bark for an hour in four pints of water, in which half the carbonate of soda

has been dissolved ; strain and express strongly through linen or calico ; moisten the residuum with water and express again ; and repeat this twice. Boil the residuum for half an hour with four pints of water and half the sulphuric acid ; strain, express strongly, moisten with water, and express again. Boil the residuum with three pints of water and a fourth part of the acid ; strain and squeeze as before. Boil again the residuum with the same quantity of water and acid, strain and squeeze as formerly. Concentrate the whole acid liquids to about a pint ; let the product cool ; filter it ; and dissolve in it the remainder of the Carbonate of soda. Collect the impure quina on a cloth, wash it slightly, and squeeze out the liquor with the hand. Break down the moist precipitate in a pint of distilled water, add one fluidscruple of Sulphuric acid, heat it to 212° , and stir occasionally. Should any precipitate retain its gray colour, and the liquid be neutral, add sulphuric acid drop by drop, stirring constantly, till the gray colour disappears. Should the liquid red- den litmus, neutralize it with a little carbonate of soda. Should crystals form on the surface, add boiling distilled water to dissolve them. Filter through paper, preserving the funnel hot ; set the liquid aside to

crystallize ; collect, and squeeze the crystals ; dissolve them in a pint of distilled water heated to 212° ; digest the solution for fifteen minutes with the Animal charcoal ; filter, and crystallize as before. Dry the crystals with a heat not exceeding 140° .

The mother liquors of each crystallization will yield a little more salt by concentration and cooling.

STRYCHNIA.

Take of Nux-vomica, one pound ;
Quicklime, one ounce and a-half ;
Rectified spirit, a sufficiency.

Subject the Nux-vomica for two hours to the vapour of steam, chop or slice it, dry it thoroughly in the vapour-bath, or hot air-press, and immediately grind it in a coffee-mill. Macerate it for twelve hours in two pints of water and boil it ; strain through linen or calico, and squeeze the residuum ; repeat the maceration and decoction twice with a pint and a half of water. Concentrate the decoctions to the consistence of thin syrup ; add the Lime in the form of milk of lime ; dry the precipitate in the vapour-bath ; pulverize it, and boil it with successive portions of Rectified spirit till the spirit cease to acquire a bitter taste. Distil off the spirit till the residuum be sufficiently concentrated to crys-

tallize on cooling. Purify the crystals by repeated crystallizations.

VERATRIA.

Take any convenient quantity of Cevadilla : pour boiling water over it in a covered vessel, and let it macerate for twenty-four hours ; remove the cevadilla, squeeze it, and dry it thoroughly with a gentle heat. Beat it now in a mortar, and separate the seeds from the capsules by brisk agitation in a deep narrow vessel. Grind the seeds in a coffee-mill, and form them into a thick paste with Rectified spirit. Pack this firmly in a percolator, and pass rectified spirit through it till the spirit ceases to be coloured. Concentrate the spirituous solutions by distillation so long as no deposit forms ; and pour the residuum while hot into twelve times its volume of cold water. Filter through calico, and wash the residuum on the filter so long as the washings precipitate with ammonia. Unite the filtered liquid with the washings, and add an excess of ammonia. Collect the precipitate on a filter, wash it slightly with cold water, and dry it first by imbibition with filtering-paper, and then in the vapour-bath. A small additional quantity may be got by concentrating the filtered ammoniacal fluid and allowing it to cool.

Veratria thus obtained is not pure, but sufficiently so for medical use. From this coloured substance it may be obtained white, though at considerable loss, by solution in very weak muriatic acid, decolorization with animal charcoal, and re-precipitation with ammonia.

CONSERVES AND ELECTUARIES.

CONSERVA AMYGDALARUM.

Take of Sweet almonds, eight ounces ;
Powder of gum-arabic, one ounce ;
White sugar, four ounces.

Blanch the almonds by maceration and peeling ; and beat them with the gum and sugar into a uniform pulpy mass.

ELECTUARIUM AROMATICUM.

Take of Aromatic powder, one part ;
Syrup of orange peel, two parts.
Mix them and triturate them into a uniform pulp.

CONSERVA AURANTII.

Grate off the outer rind of bitter oranges, and beat it into a pulp, adding gradually thrice its weight of white sugar.

ELECTUARIUM CATECHU.

Take of Catechu, and
Kino, of each four ounces ;
Cinnamon, and
Nutmeg, of each one ounce ;
Opium, diffused in a little sherry,
one drachm and a half ;
Syrup of red roses, reduced to the
consistence of honey, one pint
and a half.

Pulverize the solids ; mix the opium and syrup, then the powders, and beat them thoroughly into a uniform mass.

ELECTUARIUM OPII.

Take of Aromatic powder, six ounces ;
Senega, in fine powder, three ounces ;
Opium diffused in a little sherry, half
an ounce ;
Syrup of ginger, a pound.
Mix them together, and beat them into an
electuary.

ELECTUARIUM PIPERIS.

Take of Black Pepper, and
Liquorice-root in powder, of each a
pound ;
Fennel, three pounds ;
Honey, and
White sugar, of each two pounds ;
Triturate the solids together into a very fine
powder ; add the honey ; and beat the whole
into a uniform mass.

CONSERVA ROSAE.

Beat the petals of the *Rosa gallica* to a pulp,
gradually adding twice their weight of white
sugar.

CONSERVA ROSAE FRUCTUS.

Take any convenient quantity of hips, care-

fully deprived of their carpels ; beat them to a fine pulp, adding gradually thrice their weight of white sugar.

ELECTUARIUM SENNAE.

Take of Senna, eight ounces ;
Coriander, four ounces ;
Liquorice-root, bruised, three ounces ;
Figs, a pound ;
Pulp of prunes, a pound ;
White sugar, two pounds and a half ;
Water, three pints and a quarter.

Powder the senna and coriander ; sift out ten ounces of the mixture ; boil the residue with the figs and liquorice, in the water down to one-half ; express and strain the liquor, and evaporate it to twenty-four fluidounces ; dissolve in this the sugar, and add the liquid by degrees to the pulp of prunes ; mix gradually the powder, and triturate the whole carefully to a uniform pulp.

DECOCTIONS.

DECOCTUM ALOES.

Take of Socotorine or Hepatic Aloes,
Powder of myrrh, and
Saffron, of each one drachm ;
Extract of liquorice, half an ounce ;
Carbonate of potash, two scruples ;
Compound tincture of cardamom,
four fluidounces ;
Water, sixteen fluidounces.

Mix the aloes, myrrh, saffron, liquorice, and carbonate of potash with the water ; boil down to twelve ounces ; filter, and add the compound tincture of cardamom.

DECOCTUM CINCHONAE.

Take of Crown, Gray, Yellow, or Red cinchona, one ounce, bruised ;
Water, twenty-four fluidounces.

Boil for ten minutes, let the decoction cool, then filter it, and evaporate to sixteen fluidounces.

DECOCTUM DULCAMARAE.

Take of Dulcamara, chopped down, one ounce ;
Water, twenty-four fluidounces.

Mix them, boil, and concentrate by evaporation to sixteen fluidounces.

DECOCTUM GUAIACI.

Take of Guaiac turnings, three ounces ;
Raisins, two ounces :
Sassafras, rasped, and
Liquorice-root, bruised, each one
ounce ;
Water, eight pints.

Boil the guaiac and raisins with the water gently down to five pints, adding the liquorice and sassafras towards the end. Strain the decoction.

DECOCTUM HAEMATOXYLI.

Take of Logwood, in chips, one ounce ;
Water, a pint ;
Cinnamon, one drachm, in powder.

Boil the logwood in the water down to ten fluidounces, adding the cinnamon towards the end ; and then strain.

DECOCTUM MEZEREI.

Take of Mezereon, in chips, two drachms ;
Liquorice-root, bruised, half an
ounce ;
Water, two pints.

Mix them and boil down with a gentle heat to a pint and a-half ; and then strain.

DECOCTUM PAPAVERIS.

Take of Poppy-heads, sliced, four ounces ;
Water, three pints ;
Boil for fifteen minutes, and then strain.

DECOCTUM QUERCUS.

Take of Oak-bark, ten drachms ;
Water, two pints ;
Boil down to one pint, and then strain.

DECOCTUM SARZAE.

Take of Sarza, in chips, five ounces ;
Boiling water, four pints.
Digest the root in the water for two hours
at a temperature somewhat below ebullition,
take out the root, bruise it, replace it, boil
down to two pints, and then squeeze out the
decoction and strain it.

DECOCTUM SARZAE COMPOSITUM.

Take of Decoction of Sarza, boiling hot,
four pints ;
Sassafras, in chips, and bruised,
Guaiac turnings, and
Fresh Liquorice-root, of each, ten
drachms ;
Mezereon, half an ounce ;
Boil them together for fifteen minutes, and
then strain.

DECOCTUM SCOPARII.

Take of Broom-tops, and
Juniper-tops, of each, half an ounce ;
Bitartrate of potash, two drachms
and a-half ;
Water, a pint and a-half.

Boil them together down to a pint; and then strain.

DECOCTUM TARAXACI.

Take of Taraxacum, herb and root, fresh,
seven ounces;

Water, two pints.

Boil together down to one pint; and then strain.

DISTILLED WATERS.

Distilled waters may be prepared from fresh, and generally also from dried, vegetables. In the latter case only half the weight of material should be used. They may also be prepared for the most part by agitating the volatile oils of the plants with water and filtering the solution. But distilled waters obtained in this way have seldom so fine a flavour as those obtained from the plants themselves.

AQUA DESTILLATA.

Take any convenient quantity of spring-water; distil it from a proper vessel, rejecting the first twentieth part, and preserving the first half of the remainder.

AQUA ANETHI.

Take of Anethum seeds, bruised, eighteen ounces;
Water, two gallons;
Rectified spirit, three fluidounces.
Mix together, and distil off one gallon.

AQUA CASSIAE.

Take of Cassia-bark bruised, eighteen ounces;
Water, two gallons;
Rectified spirit, three fluidounces.
Mix them together, and distil off one gallon

AQUA CINNAMOMI.

This distilled water is to be prepared with Cinnamon in the same way as *Aqua cassiæ*.

AQUA FOENICULI.

This distilled water is prepared with Fennel in the same way as *Aqua Anethi*.

AQUA LAUROCERASI.

Take of fresh leaves of Cherry-laurel, a pound ;
Water, two pints and a half ;
Compound spirit of Lavender, an ounce.

Chop down the leaves, mix them with the water, distil off one pint, agitate the distilled liquid well, filter it if any milkiness remain after a few seconds of rest, and then add the lavender spirit.

AQUA MENTHÆ PIPERITÆ.

This distilled water is prepared as *Aqua menthæ viridis*.

AQUA MENTHÆ VIRIDIS.

Take of Spearmint four pounds if fresh, two two pounds if dry ;
Water, two gallons ;
Rectified-spirit, three fluidounces.
Mix them ; and distil off one gallon.

AQUA PIMENTAE.

Take of Pimento bruised, one pound ;
Water, two gallons ;
Rectified-spirit, three fluidounces.
Mix them ; and distil off one gallon.

AQUA PULEGII.

This distilled water is prepared like Aqua menthæ viridis.

AQUA ROSAE.

Take of petals of Rosa centifolia, ten pounds ;
Water, two gallons ;
Rectified-spirit, three fluidounces.
Mix them and distil off one gallon. The petals should be preferred when fresh ; but it also answers well to use those which have been preserved by beating them with twice their weight of muriate of soda.

AQUA SAMBUCCI.

Take of Elder-flowers, fresh, ten pounds ;
Water, two gallons ;
Rectified-spirit, three fluidounces.
Mix them ; and distil off one gallon.

ENEMAS.

ENEMA CATHARTICUM.

Take of Olive oil, one ounce ;
Sulphate of magnesia, half an ounce ;
Sugar, one ounce ;
Senna, half an ounce ;
Boiling water, sixteen fluidounces :
Infuse the senna for an hour in the water ;
then dissolve the salt and sugar ; add the oil,
and mix them by agitation.

ENEMA FOETIDUM.

Add to the cathartic enema two drachms of
tincture of assafoetida.

ENEMA OPII vel ANODYNUM.

Take of Starch, half a drachm ;
Tincture of opium, half a fluidrachm
to one fluidrachm ;
Water, two fluidounces :
Boil the starch in the water, and when it is
cool enough for use, add the tincture of opium.

ENEMA TEREBINTHINAE.

Take of Oil of Turpentine, one fluidounce ;
Yolk of egg, a sufficiency ;
Water, nineteen fluidounces ;
Rub the oil and yolk carefully together, and
then add the water gradually.

ENEMA TABACI.

Take of Tobacco, 15 grains to half a drachm ;
Boiling water, eight fluidounces :
Infuse for half an hour, and then strain.

EXTRACTS.

Extracts are usually prepared by evaporating the expressed juices of plants, or their infusions and decoctions in water, proof-spirit, or rectified spirit, at a temperature not exceeding 212° F. by means of a vapour-bath. Most of them, however, may be obtained of greatly superior quality by the process of evaporation in vacuo. And the extracts of expressed juices cannot, perhaps, be better prepared than by spontaneous evaporation in shallow vessels, exposed to a current of air. Extracts should be evaporated to such a consistence as to form a firm pill-mass when cold.

EXTRACTUM ACONITI.

Take of the leaves of Monkshood, fresh, any convenient quantity; beat them into a pulp; express the juice; subject the residuum to percolation with rectified spirit, so long as the spirit passes materially coloured; unite the expressed juice and the spirituous infusion; filter; distil off the spirit; and evaporate the residuum in the vapour-bath, taking care to remove the vessel from the heat so soon as the due degree of consistence shall be attained.

EXTRACTUM ANTHEMIDIS.

Take of Chamomile, a pound; boil it with a

gallon of water down to four pints ; filter the liquor hot ; evaporate in the vapour-bath to the due consistence.

EXTRACTUM BELLADONNAE.

Take of Belladonna fresh, any convenient quantity ; bruise it in a marble mortar into a uniform pulp ; express the juice ; moisten the residuum with water, and express again. Unite the expressed fluids, filter them, and evaporate the filtered liquid in the vapour-bath to the consistence of firm extract, stirring constantly towards the close.

EXTRACTUM CINCHONAE.

Take of any of the varieties of Cinchona, but especially the Yellow or Red cinchona, in fine powder, four ounces ;
Proof spirit, twenty-four fluidounces ;
Percolate the Cinchona with the spirit ; distil off the greater part of the spirit ; and evaporate what remains in an open vessel over the vapour-bath to a due consistence.

EXTRACTUM COLCHICI ACETICUM.

Take of the Bulb of Colchicum, a pound ;
Pyroligneous acid, three fluidounces ;
Beat the colchicum to a pulp, gradually adding the acid ; express the liquid, and evaporate it in a porcelain vessel (not glazed with lead) over the vapour-bath to the due consistence.

EXTRACTUM COLOCYNTHIDIS.

Take of Colocynth, a pound ;

Water, two gallons.

Boil gently for six hours, replacing the evaporated water occasionally. Strain the liquor while hot ; and evaporate it in the vapour-bath to the due consistence.

EXTRACTUM CONII.

Take of Conium any convenient quantity ; beat it into a uniform pulp in a marble mortar, express the juice, and filter it. Let this juice be evaporated to the consistence of a very firm extract either in a vacuum with the aid of heat, or spontaneously in shallow vessels exposed to a strong current of air freed of dust by gauze-skreens.

This extract is of good quality only when a very strong odour of conia is disengaged by degrees on its being carefully triturated with Aqua potassæ.

EXTRACTUM DIGITALIS.

This extract is best prepared from the fresh leaves of Digitalis, by any of the processes indicated for extract of conium.

EXTRACTUM ELATERII, SEU ELATERIUM.

Take of the fruit of Momordica elaterium before it is quite ripe, any convenient quantity ; cut the fruit and express the juice gently through a fine sieve ; allow the liquid to

rest till it becomes pretty clear ; pour off the supernatant liquor, which may be thrown away; and dry the feculence with a gentle heat.

EXTRACTUM GENTIANAE.

Take of Gentian, any convenient quantity ; bruise it to a moderately fine powder ; mix it thoroughly with half its weight of distilled water ; in twelve hours put it into a proper percolator, and exhaust it by percolation with temperate distilled water ; concentrate the liquid, filter before it becomes too thick, and evaporate in the vapour-bath to a due consistence.

EXTRACTUM GLYCIRRHIZAE.

Cut Liquorice-root into small chips, dry it thoroughly with a gentle heat, reduce it to a moderately fine powder, and proceed as for extract of Gentian.

EXTRACTUM HAEMATOKYLI.

Take of Logwood, in fine chips, a pound ;
Boiling water, a gallon.

Macerate for twenty-four hours, then boil down to four pints, strain, and concentrate in the vapour-bath to the due consistence.

EXTRACTUM HYOSCYAMI.

This extract is to be prepared from the fresh leaves of Hyoscyamus by any of the processes directed for extract of Conium.

EXTRACTUM sive RESINA JALAPAE.

Take any convenient quantity of Jalap, in moderately fine powder; mix it thoroughly with enough of rectified spirit to moisten it well; put it in twelve hours into a percolator, and exhaust the powder with rectified spirit; distil off the greater part of the spirit, and concentrate the residuum over the vapour-bath to a due consistence.

EXTRACTUM KRAMERIAE.

This extract is to be prepared from Krameria-root in the same way with that of Liquorice-root.

EXTRACTUM LUPULI.

This extract is prepared from Hops in the same way with the Extract of Logwood.

EXTRACTUM NUCIS-VOMICAE.

Take of nux-vomica any convenient quantity; expose it in a proper vessel to steam till it is completely softened; slice it, dry it thoroughly, and immediately grind it in a coffee-mill; exhaust the powder either by percolating it with rectified spirit, or by boiling it with repeated portions of rectified spirit, until the spirit comes off free of bitterness. Distil off the greater part of the spirit; and evaporate what remains in the vapour-bath to a proper consistence.

EXTRACTUM OPII.

Take of Opium, one pound ;

Water, five pints ;

Cut the opium into small fragments, macerate it for twenty-four hours in a pint of water, break down the fragments with the hand, express the liquid with pretty strong pressure ; break down the residuum again in another pint of the water, let it macerate for twenty-four hours, and express the liquid ; repeat the maceration and expression in the same way till the water is all used. Filter the successive infusions as they are made, passing them through the same filter ; unite and evaporate them in the vapour-bath to the due consistence.

EXTRACTUM PAPAVERIS.

Take of Poppy-heads without the seeds, fifteen ounces ;

Boiling water, a gallon ;

Macerate for twenty-four hours ; boil down to four pints ; filter the liquor hot, and evaporate over the vapour-bath to the due consistence.

EXTRACTUM PAREIRÆ.

This extract is to be prepared from Pareira-root in the same way with the extract of Liquorice-root.

EXTRACTUM QUASSIÆ.

This extract is to be prepared from Quassia in the same way with the extract of Liquorice-root.

EXTRACTUM RHEI.

Take of Rhubarb, one pound ;

Water, five pints ;

Cut the Rhubarb into small fragments, macerate it for twenty-four hours in three pints of the water, filter the liquor through a cloth, and express it with the hands or otherwise moderately ; macerate the residuum with the rest of the water for twelve hours at least, filter the liquor with the same cloth as before, and express the residuum strongly. The liquors, filtered again if necessary, are then to be evaporated together to a proper consistence in the vapour-bath. The extract however is obtained of finer quality by evaporation in a vacuum with a gentle heat.

EXTRACTUM *sive* RESINA SCAMMONII.

Take any convenient quantity of Scammony in fine powder ; boil it in successive portions of proof-spirit till the spirit ceases to dissolve any thing ; filter ; distil the liquid till little but water passes over. Then pour away the watery solution from the resin at the bottom ; agitate the resin with succes-

sive portions of boiling water till it is well washed ; and lastly, dry it at a temperature not exceeding 240°.

EXTRACTUM SARZAE FLUIDUM.

Take of Sarza in chips, one pound ;

Boiling water, six pints ;

Digest the root for two hours in four pints of the water ; take it out, bruise it, replace it, and boil for two hours ; filter and squeeze out the liquid ; boil the residuum in the remaining two pints of water, and filter and squeeze out this liquor also ; evaporate the united liquors to the consistence of thin syrup ; add, when the product is cool, as much rectified spirit as will make in all sixteen fluidounces. Filter.

This fluid extract may be aromatized with volatile oils or warm aromatics.

EXTRACTUM STYRACIS.

Take any convenient quantity of Storax, in fine powder ; exhaust it by boiling it in successive quantities of rectified spirit ; filter the spirituous solutions ; distil off the greater part of the spirit ; evaporate the remainder over the vapour-bath to the consistence of a thin extract.

EXTRACTUM STRAMONII.

Take of seeds of Stramonium any conve-

nient quantity ; grind them well in a coffee-mill.

Rub the powder into a thick mass with Proof-spirit ; put the pulp into a percolator, and transmit proof-spirit till it passes colourless ; distil off the spirit, and evaporate what remains in the vapour-bath to a proper consistence.

EXTRACTUM TARAXACI.

Take of fresh root of Taraxacum, a pound ;
Boiling water a gallon.

Proceed as for the preparation of extract of Poppy heads.

HONEYYS.

MEL BORACIS.

Take of Borax, one drachm ;

Honey, one ounce ;

Mix them.

MEL ROSAE.

Take of the dried petals of *Rosa gallica*, four
ounces ;

Boiling water, two pints and a-half ;

Honey, five pounds ;

Infuse the petals in the water for six hours ;
strain and squeeze ; let the impurities sub-
side ; pour off the clear liquor ; mix the honey
with it ; and evaporate the whole in the
vapour-bath to the consistence of syrup, re-
moving the scum which forms.

INFUSIONS.

INFUSUM ANTHEMIDIS.

Take of Chamomile, five drachms ;

Boiling water, one pint.

Infuse for twenty minutes in a covered vessel, and then strain.

INFUSUM AURANTII.

Take of Bitter orange-peel, dried, half an ounce ;

Lemon-peel, fresh, two drachms ;

Cloves, bruised, one drachm ;

Boiling water, one pint ;

Infuse for fifteen minutes in a covered vessel, and strain through linen or calico.

INFUSUM BUCKU.

Take of Bucku, an ounce ;

Boiling water, one pint ;

Infuse for two hours in a covered vessel, and strain through linen or calico.

INFUSUM CALUMBAE.

Take of Calumba, in coarse powder, half an ounce ;

Cold water, about a pint.

Triturate the Calumba with a little of the water, so as to moisten it thoroughly ; put it into a percolator, and transmit cold water

till sixteen fluidounces of infusion be obtained.

INFUSUM CARYOPHYLLI.

Take of bruised Cloves, three drachms ;
Boiling water, one pint ;
Infuse for two hours in a covered vessel,
and strain through linen or calico.

INFUSUM CASCARILLÆ.

Take of Cascarilla, bruised, an ounce and a
half ;
Boiling water, one pint ;
Infuse for two hours, in a covered vessel,
and strain through linen or calico.

INFUSUM CATECHU.

Take of Catechu, in powder, six drachms ;
Cinnamon, in powder, one drachm ;
Syrup, three fluidounces ;
Boiling water, seventeen fluid-
ounces ;
Infuse the catechu and cinnamon with the
water for two hours, strain through linen or
calico, and add the syrup.

INFUSUM CHIRETTÆ.

Take of Chiretta, four drachms ;
Boiling water, one pint ;
Infuse for two hours, and strain through li-
nen or calico.

INFUSUM CINCHONÆ.

Take of any species of Cinchona, according to prescription, one ounce in powder ;

Boiling water, one pint ;

Infuse for four hours in a covered vessel, and then strain through linen or calico.

INFUSUM CUSPARIÆ.

Take of Cusparia, bruised, five drachms ;

Boiling water, one pint ;

Infuse for two hours in a covered vessel, and then strain through linen or calico.

INFUSUM DIGITALIS.

Take of Digitalis, dried, two drachms ;

Spirit of Cinnamon, two fluidounces ;

Boiling water, eighteen fluidounces ;

Infuse the Digitalis in the water, in a covered vessel for four hours ; strain through linen or calico ; and then add the spirit of cinnamon.

INFUSUM GENTIANÆ.

Take of Gentian, sliced, half an ounce ;

Bitter orange-peel, dried and bruised, one drachm ;

Coriander, bruised, one drachm ;

Proof-spirit, four fluidounces ;

Cold water, sixteen fluidounces ;

Pour the spirit upon the solids ; in three

hours add the water ; and in twelve hours more strain through linen or calico.

INFUSUM LINI.

Take of Linseed, six drachms ;
Liquorice-root, bruised, two drachms ;
Boiling-water, one pint ;
Digest near the fire in a covered vessel for four hours, and then strain through linen or calico.

INFUSUM PAREIRAE.

Take of Pareira, six drachms ;
Boiling water, one pint ;
Infuse for two hours in a covered vessel, and then strain through linen or calico.

INFUSUM QUASSIAE.

Take of Quassia, in chips, one drachm ;
Boiling water, one pint ;
Infuse for two hours in a covered vessel, and then strain through linen or calico.

INFUSUM RHEI.

Take of Rhubarb, bruised into coarse powder, one ounce ;
Spirit of cinnamon, two fluidounces ;
Boiling water, eighteen fluidounces ;
Infuse the rhubarb for twelve hours in the water in a covered vessel ; add the spirit, and strain through linen or calico.

INFUSUM ROSAE.

Take of Rosa gallica dried, three drachms ;
Diluted Sulphuric acid, one fluidrachm and a half ;
Pure Sugar, six drachms ;
Boiling water, one pint ;

Infuse the rose-petals in the water, in a covered vessel of glass or porcelain, not glazed with lead, for one hour ; then add the acid, strain through linen or calico, and dissolve the sugar in the liquor.

INFUSUM SENEGAE.

Take of Senega, ten drachms ;
Boiling water, one pint ;
Infuse for four hours in a covered vessel, and strain.

INFUSUM SENNAE.

Take of Senna, an ounce and a half ;
Ginger, bruised, four scruples ;
Boiling water, one pint ;
Infuse for an hour in a covered vessel ; and then strain through linen or calico.

INFUSUM SENNAE COMPOSITUM.

Take of Senna, one drachm ;
Tamarinds, one ounce ;
Coriander, bruised, one drachm ;
Muscovado, half an ounce ;
Boiling water, eight fluidounces ;

Infuse for four hours, with occasional stirring in a covered vessel, not glazed with lead, and then strain through linen or calico.

This infusion may be likewise made with twice or thrice the prescribed quantity of senna.

INFUSUM SERPENTARIAE.

Take of *Serpentaria*, half an ounce ;

Boiling water, a pint ;

Infuse for four hours in a covered vessel, and then strain through linen or calico.

INFUSUM SIMARUBAE.

Take of *Simaruba*, bruised, three drachms ;

Boiling water, a pint ;

Infuse for two hours in a covered vessel ; and then strain through linen or calico.

METALS AND THEIR COMPOUNDS.

ALUMEN EXSICCATUM.

Take any convenient quantity of alum ; fuse it over the fire in a vessel of iron or earthenware ; continue the heat till ebullition ceases and vapour is no longer discharged ; and then reduce it to powder.

ANTIMONII OXIDUM.

Take of Sulphuret of Antimony in fine powder, four ounces ;
Muriatic acid (commercial,) one pint ;
Water, five pints.

Dissolve the sulphuret in the acid with the aid of a gentle heat ; boil for half an hour ; filter ; pour the fluid into the water ; collect the precipitate on a calico filter ; wash it well with cold water, then with a weak solution of carbonate of soda, and again with cold water till the water ceases to affect reddened litmus paper. Dry the powder over the vapour bath.

PULVIS ANTIMONIALIS.

Take of Sulphuret of Antimony, in coarse powder ;
Hartshorn shavings, equal weights ;
Mix them, put them into a red-hot iron pot,

and stir constantly till they acquire an ash-gray colour and vapours no longer arise. Pulverise the product, put it into a crucible with a perforated cover, and expose this to a gradually-increasing heat till a white heat be produced, which is to be maintained for two hours. Reduce the product when cold to fine powder.

ANTIMONII SULPHURETUM AUREUM.

Take of Sulphuret of Antimony, in fine powder, one ounce ;

Solution of Potash, eleven fluid-ounces ;

Water, two pints ;

Mix the water and solution of potash, add the sulphuret, boil for an hour, filter immediately, and precipitate the liquid, while hot, with an excess of diluted sulphuric acid. Collect the precipitate on a calico filter, wash it thoroughly with water, and dry it with a gentle heat.

ANTIMONII TARTARIZATUM.

Take of Sulphuret of Antimony, in fine powder, four ounces ;

Muriatic acid (commercial,) one pint ;

Water, five pints ;

Dissolve the sulphuret in the acid with the aid of a gentle heat ; boil for half an hour ;

filter; pour the liquid into the water; collect the precipitate on a calico filter, wash it with cold water till the water ceases to redden litmus-paper; dry the precipitate over the vapour-bath.

Take of this precipitate three ounces;

Bitartrate of potash, four ounces and two drachms;

Water, twenty-seven fluidounces;

Mix the powders, add the water, boil for an hour, filter, and set the liquid aside to crystallize. The mother-liquor when concentrated yields more crystals, but not so free of colour, and therefore requiring a second crystallization.

VINUM ANTIMONIALE.

Take of Tartar-emetic, two scruples;

Sherry, one pint;

Dissolve the salt in the wine.

ARGENTI NITRAS.

Take of Pure Silver, an ounce and a half;

Pure Nitric acid, one fluidounce;

Distilled water, two fluidounces;

Mix the acid and water, add the silver, and dissolve it with the aid of a gentle heat; increase the heat gradually till a dry salt be obtained; fuse the salt in an earthen-ware or porcelain crucible, and pour the fused matter into iron moulds previously heated and

greased slightly with tallow. Preserve the product in glass vessels.

LIQUOR ARSENICALIS.

Take of White Arsenic in powder, and
Carbonate of Potash, of each, four
scruples ;
Compound Tincture of Lavender,
five fluidrachms ;
Water, one pint ;

Dissolve the oxide and carbonate together in half the water, with the aid of heat ; filter, if necessary ; add the tincture to the liquid when cold, and then dilute it with water till the whole measure one pint.

BARYTAE MURIAS.

Take of Carbonate of Baryta, in fragments,
ten ounces ;
Pure Muriatic acid, half a pint ;
Distilled water, two pints ;

Mix the acid and water ; add the carbonate by degrees ; apply a gentle heat towards the close of the effervescence ; and when the action is over, filter, concentrate, and set aside the solution to crystallize.

or

Take of Sulphate of Baryta, two pounds ;
Charcoal in fine powder, four ounces ;
Pure Muriatic acid, a sufficiency ;
Heat the sulphate to redness, reduce it to

fine powder, mix the charcoal with it thoroughly, heat the mixture in a covered crucible for three hours at a low white heat. Pulverize the product, put it gradually into five pints of boiling water; boil for a few minutes; let it rest for a little over a vapour-bath; pour off the clear liquor, and filter it if necessary, keeping it hot. Pour three pints of boiling water over the residuum, and proceed as before. Unite the two liquids; and while they are still hot, or, if cooled, after heating them again, add pure muriatic acid gradually so long as effervescence is occasioned. In this process the solutions ought to be as little exposed to the air as possible; and in the last step the disengaged gas should be discharged by a proper tube into a chimney or the ash-pit of a furnace. Strain the liquor, concentrate it, and set it aside to crystallize.

SOLUTIO BARYTAE MURIATIS.

Take of Muriate of Baryta, one drachm;
Distilled water, one fluidounce;
Dissolve the salt in the water.

BISMUTHUM ALBUM.

Take of Bismuth, in fine powder, one ounce;
Nitric acid (D.1380) one fluidounce
and a-half;
Water, three pints;

Add the metal gradually to the acid, favouring the action with a gentle heat, and adding a very little distilled water so soon as crystals or a white powder may begin to form. When the solution is complete, pour the liquid into the water. Collect the precipitate immediately on a calico filter, wash it quickly with cold water, and dry it in a dark place.

CALX.

Heat white marble broken into small fragments in a covered crucible at a full-red heat for three hours, or till the residuum when slaked and suspended in water no longer effervesces on the addition of muriatic acid.

AQUA CALCIS.

Take any convenient quantity of water ; pour a little of it over about a twentieth of its weight of lime ; when the lime is slaked, add it to the rest of the water in a bottle ; agitate well ; allow the undissolved matter to subside ; pour off the clear liquor when it is wanted, replacing it with more water, and agitating briskly as before.

CALCIS MURIAS.

Take of White marble, in fragments, ten ounces ;
Muriatic acid, (commercial) and
Water, of each one pint ;

Mix the acid and water ; add the marble by degrees, and when the effervescence is over, add a little marble in fine powder till the liquid no longer reddens litmus ; filter and concentrate to one-half ; put the remaining fluid in a cold place to crystallize ; preserve the crystals in a well-closed bottle. More crystals will be obtained by concentrating the mother-liquor.

CALCIS MURIATIS SOLUTIO.

Take of Muriate of lime, eight ounces ;

Water, twelve fluidounces :

Dissolve the salt in the water.

CRETA PRÆPARATA.

Take any convenient quantity of chalk ; triturate it well in a mortar with a little water ; then pour it into a large vessel nearly full of water, and agitate briskly ; allow it to rest for a short time, and pour the milky water into another vessel, in which the fine suspended chalk is to be left slowly to subside ; repeat this process with the coarsely powdered chalk which subsided quickly in the first vessel ; collect the fine powder in the second vessel on a filter of linen or calico, and dry it.

CUPRUM AMMONIATUM.

Take of Sulphate of Copper, two ounces ;

Carbonate of Ammonia, three ounces ;

Triturate them thoroughly together, till effervescence ceases, wrap the product in blotting-paper, and dry it first by folds of blotting-paper, afterwards by exposure to the air for a little ; and preserve it in closely-stopped bottles.

CUPRI AMMONIATI SOLUTIO.

Take of Ammoniated Copper, one drachm ;

Water, one pint ;

Dissolve the salt in the water, and filter.

FERRI CARBONAS SACCHARATUM.

Take of Sulphate of Iron, four ounces ;

Carbonate of Soda, five ounces ;

Pure Sugar, two ounces ;

Water, four pints.

Dissolve the sulphate and carbonate each in two pints of the water ; add the solutions and mix them ; collect the precipitate on a cloth filter, and immediately wash it with cold water, squeeze out as much of the water as possible, and without delay triturate the pulp which remains with the sugar previously in fine powder. Dry the mixture at a temperature not much above 120°.

FERRI IODIDI SYRUPUS.

Take of Iodine (dry) 200 grains ;

Fine Iron-wire recently cleaned, 100 grains ;

White-sugar in powder, four ounces and a-half ;

Distilled water, six fluidounces.

Boil the iodine, iron, and water together in a glass matrass, at first gently to avoid the expulsion of iodine-vapour, afterwards briskly, until about two fluidounces of liquid remain. Filter this quickly, while hot, into a matrass containing the sugar ; dissolve the sugar with a gentle heat ; and add distilled water, if necessary, to make up six fluidounces.—Twelve minims contain one grain of iodide of iron.

FERRI IODIDUM.

Take any convenient quantity of Iodine, Iron-wire and Distilled water in the proportions for making Solution of Iodide of Iron. Proceed as directed for that process ; but before filtering the solution concentrate it to one-sixth of its volume, without removing the excess of iron-wire. Put the filtered liquor quickly in an evaporating basin, along with twelve times its weight of quicklime around the basin, in some convenient apparatus in which it may be shut up accurately in a small space not communicating with the general atmosphere. Heat the whole apparatus in a hot air-press, or otherwise, until the

water be entirely evaporated; and preserve the dry iodide in small well-closed bottles.

FERRI MURIATIS TINCTURA.

Take of Red Oxide of Iron, six ounces;
Muriatic acid, (commercial) one
pint;
Rectified Spirit, three pints;
Add the oxide to the acid in a glass vessel;
digest with a gentle heat, and occasional agi-
tation, for a day, or till most of the oxide be
dissolved; then add the spirit, and filter.

FERRI OXIDUM NIGRUM.

Take of Sulphate of Iron, six ounces;
Sulphuric Acid, (commercial) two
fluidrachms and two fluidscruples;
Pure Nitric Acid, four fluidrachms
and a-half;
Stronger Aqua Ammoniaë, four fluid-
ounces and a-half;
Boiling water, three pints.
Dissolve half the sulphate in half the boil-
ing water and add the sulphuric acid; boil;
add the nitric acid by degrees, boiling the
liquid after each addition briskly for a few
minutes. Dissolve the rest of the sulphate
in the rest of the boiling water; mix tho-
roughly the two solutions; and immediately
add the ammonia in a full stream, stirring
the mixture at the same time briskly. Collect

the black powder on a calico-filter ; wash it with water till the water is scarcely precipitated by solution of nitrate of baryta ; and dry it at a temperature not exceeding 180°.

FERRI OXIDUM RUBRUM.

Take of Sulphate of Iron, four ounces ;
Carbonate of Soda, five ounces ;
Boiling water, half a pint ;
Cold water, three pints and a-half ;
Dissolve the sulphate in the boiling water, add the cold water, and then the carbonate of soda previously dissolved in about thrice its weight of water. Collect the precipitate on a calico-filter ; wash it with water till the water is but little affected with solution of nitrate of baryta ; and dry it in the hot air-press or over the vapour-bath.

FERRI SULPHAS.

If the Sulphate of iron of commerce be not in transparent green crystals, without efflorescence, dissolve it in its own weight of boiling water acidulated with a little sulphuric acid ; filter ; and set the solution aside to crystallize. Preserve the crystals in well-closed bottles.

FERRI SULPHAS EXSICCATUM.

Expose any convenient quantity of Sulphate of iron to a moderate heat in a porcelain or

earthen-ware vessel not glazed with lead, till it is converted into a dry grayish-white mass, which is to be reduced to powder.

FERRI SULPHURETUM.

The best sulphuret of Iron is made by heating an iron rod to a full-white heat in a forge, and rubbing it with a roll of sulphur over a deep vessel filled with water to receive the fused globules of sulphuret which form. An inferior sort, good enough however for pharmaceutic purposes, is obtained by heating one part of sublimed sulphur and three of iron-filings in a crucible in a common fire till the mixture begins to glow, and then removing the crucible and covering it, until the action, which at first increases considerably, shall come to an end.

FERRUGO.

Take of Sulphate of Iron, four ounces ;
Sulphuric Acid (commercial) three
fluidrachms and a-half ;
Nitric Acid (D. 1380), nine fluid-
drachms ;
Stronger Aqua Ammoniaë three
fluidounces and a-half.
Water, two pints ;

Dissolve the Sulphate in the water, add the Sulphuric acid, and boil the solution ; add then the Nitric acid in small portions, boil-

ing the liquid for a minute or two after each addition, until it acquires a yellowish-brown colour and yields a precipitate of the same colour with ammonia. Filter; allow the liquid to cool; and add in a full stream the Aqua Ammoniaë, stirring the mixture briskly. Collect the precipitate on a calico filter; wash it with water till the washings cease to precipitate with nitrate of baryta; squeeze out the water as much as possible; and dry the precipitate at a temperature not exceeding 180°.

When this preparation is kept as an antidote for poisoning with arsenic, it is preferable to preserve it in the moist state, after being simply squeezed.

FERRUM TARTARIZATUM.

Take of Sulphate of Iron, five ounces;
Bitartrate of potash, five ounces and
one drachm;
Carbonate of Ammonia in fine powder, a sufficiency.

Prepare the Rust of iron from the sulphate as directed under Ferrugo, and without drying it. Mix the pulpy mass with four pints of water; add the Bitartrate; boil till the rust of iron is dissolved; let the solution cool; pour off the clear liquid, and add to this the Carbonate of ammonia so long as it occasions effervescence. Concentrate the

liquid over the vapour bath to the consistence of a thick extract, or till the residuum becomes on cooling a firm solid; which must be preserved in well closed vessels.

HYDRARGYRI BINIODIDUM.

Take of Mercury, two ounces;
Iodine, two ounces and a-half;
Concentrated Solution of Muriate
of Soda, a gallon;

Triturate the Mercury and Iodine together, adding occasionally a little rectified spirit till a uniform red powder be obtained. Reduce the product to fine powder, and dissolve it in the solution of muriate of soda with the aid of brisk ebullition. Filter, if necessary, through calico, keeping the funnel hot; wash and dry the crystals which form on cooling.

CALOMELAS.

Take of Mercury, eight ounces;
Sulphuric acid (commercial), two
fluidounces and three fluidrachms;
Pure Nitric acid, half a fluidounce;
Muriate of Soda, three ounces;

Mix the acids, add four ounces of the mercury, and dissolve it with the aid of a moderate heat. Raise the heat so as to obtain a dry salt. Triturate this with the Muriate of soda and the rest of the Mercury till the globules entirely disappear. Heat the mix-

ture by means of a sandbath in a proper subliming apparatus. Reduce the sublimate to fine powder; wash the powder with boiling distilled water until the water ceases to precipitate with solution of Iodide of potassium; and then dry it.

CINNABARIS.

Take of Mercury, two pounds;

Sulphur, five ounces;

Melt the sulphur, add the mercury, and continue the heat till the mixture begins to swell up. Then remove the vessel, and cover it closely to prevent the mixture taking fire. When the material is cold, reduce it to powder, and sublime it.

HYDRARGYRUM CUM CRETA.

Take of Mercury, three ounces;

Prepared Chalk, five ounces;

Triturate them together till the globules entirely disappear.

HYDRARGYRI OXIDUM RUBRUM.

Take of Mercury, eight ounces;

Diluted Nitric acid (D. 1280), five fluid ounces;

Dissolve half of the mercury in the acid with the aid of a moderate heat; and continue the heat till a dry salt is formed. Triturate the rest of the mercury with the

salt till a fine uniform powder be obtained ; heat the powder in a porcelain vessel and constantly stir it, till acid fumes cease to be discharged.

HYDRARGYRI PRECIPITATUM ALBUM.

Take of Corrosive Sublimate, six ounces ;

Distilled water, six pints ;

Aqua Ammoniaë, eight fluidounces ;

Dissolve the Corrosive sublimate with the aid of heat in the Distilled water ; and when the solution is cold add the Aqua Ammoniaë ; stir the whole well ; collect the powder on a calico filter, and wash it thoroughly with cold water.

SUBLIMATUS CORROSIVUS.

Take of Mercury, four ounces ;

Sulphuric acid (commercial), two fluidounces and three fluidrachms ;

Pure Nitric Acid, half a fluidounce ;

Muriate of Soda, three ounces.

Mix the acids ; add the mercury ; dissolve it with the aid of a moderate heat ; and then raise the heat so as to obtain a dry salt. Triturate this thoroughly with the muriate of soda ; and sublime in a proper apparatus.

MAGNESIA.

Take any convenient quantity of Carbonate of Magnesia, expose it in a crucible to a full

red heat for two hours, or till the powder, when suspended in water, presents no effervescence on the addition of muriatic acid. Preserve the product in well-closed bottles.

MAGNESIAE CARBONAS.

Take of Sulphate of Magnesia, four pounds ;
Carbonate of Soda, four pounds and
eight ounces ;
Water, four gallons.

Dissolve the salts separately, each in two gallons of the water ; mix the solutions, boil the mixture, and stir briskly for fifteen or twenty minutes. Collect the precipitate on a filter of calico or linen, wash it thoroughly with boiling water, and then dry it.

PLUMBI ACETAS.

Take of Pyroligneous acid (D. 1034), two
pints ;
Distilled water, one pint :
Litharge, fourteen ounces.

Mix the acid and water, add the litharge, dissolve it with the aid of a gentle heat, filter, concentrate the solution sufficiently for crystallization on cooling.

PLUMBI DIACETATIS SOLUTIO.

Take of Acetate of lead, six ounces and six
drachms :
Litharge in fine powder, four ounces ;
Water, a pint and a half.

Boil the salt and litharge with the water for half an hour, stirring occasionally. When the solution is cold add water, if necessary, to make up a pint and a half; and then filter. Preserve the solution in well-closed bottles.

PLUMBI IODIDUM.

Take of Iodide of Potassium and
Nitrate of Lead of each an ounce;
Water, a pint and a half;

Dissolve the salts separately, each in one-half of the water; add the solutions; collect the precipitate on a filter of linen or calico, and wash it with water. Boil the powder in three gallons of water acidulated with three fluidounces of pyroligneous acid. Let any undissolved matter subside, maintaining the temperature near the boiling point; and pour off the clear liquor, from which the iodide of lead will crystallize on cooling.

PLUMBI NITRAS.

Take of Litharge, four ounces and a half;
Diluted Nitric acid, a pint;

Dissolve the litharge to saturation with the aid of a gentle heat. Filter and set the liquid aside to crystallize. Concentrate the residual liquid to obtain more crystals.

POTASSÆ ACETAS.

Take of Pyroligneous acid, a pint and a-half;

Carbonate of potash (dry) seven ounces, or a sufficiency ;

Add the carbonate gradually to the acid till complete neutralization is accomplished. Evaporate the solution over the vapour-bath till it is so concentrated as to form a concrete mass when cold. Allow it to cool and crystallize in a solid cake ; which must be broken up and immediately put into well-closed bottles.

POTASSÆ AQUA EFFERVESCENS.

Take of Bicarbonate of potash, one drachm ;
Distilled water, one pint ;

Dissolve the salt in the water, and transmit through the solution carbonic acid gas under strong pressure.

POTASSÆ BICARBONAS.

Take of Carbonate of Potash, six ounces ;
Carbonate of Ammonia, three ounces and a half ;

Triturate the Carbonate of Ammonia to a very fine powder ; mix with it the carbonate of potash ; triturate them thoroughly together, adding by degrees a very little water, till a smooth and uniform pulp be formed. Dry this gradually at a temperature not exceeding 140°, triturating occasionally towards the close ; and continue the desiccation till a fine powder be obtained, entirely free of ammoniacal odour.

POTASSAE BISULPHAS.

Take of the residuum in the preparation of
Pure nitric acid, two pounds ;
Sulphuric acid (commercial), seven
fluidounces and one fluidrachm ;
Boiling water, six pints ;
Dissolve the salt in the water, add the acid,
concentrate the solution, and set it aside to
cool and form crystals.

POTASSAE CARBONAS PURUM.

Pure Carbonate of potash may be most readily
obtained by heating crystallized Bicarbonate
of potash to redness in a crucible, but more
cheaply by dissolving Bitartrate of potash in
thirty parts of boiling water, separating and
washing the crystals which form on cooling,
heating these in a loosely-covered crucible
to redness so long as fumes are discharged,
breaking down the mass, and roasting it in an
open crucible for two hours, with occasional
stirring, lixiviating the product with dis-
tilled water, filtering the solution thus ob-
tained, evaporating the solution to dryness,
granulating the salt towards the close by
brisk agitation, and heating the granular
salt nearly to redness. The product of either
process must be kept in well-closed vessels.

POTASSAE SULPHAS.

Take of the residuum of the preparation of

Pure nitric acid, two pounds ;
Boiling water, two gallons ;
White marble in powder, a suffi-
ciency ;

Dissolve the salt in the water ; add the marble gradually till effervescence ceases, and the solution is completely neutralized ; filter the liquid, and evaporate it till a pellicle forms on its surface ; then set it aside to cool and form crystals.

POTASSAE SULPHAS CUM SULPHURE.

Take of Nitrate of Potash, and
Sulphur, equal parts ;

Mix them thoroughly ; throw the mixture in small successive portions into a red-hot crucible ; and when the deflagration is over, and the salt has cooled, reduce it to powder, and preserve it in well-closed bottles.

POTASSAE TARTRAS.

Take of Bitartrate of Potash, three pounds ;
Carbonate of Potash, sixteen ounces,
or a sufficiency ;
Boiling water, six pints ;

Dissolve the carbonate in the water, add the bitartrate till the liquor is neutralized, boil and filter. Concentrate the liquor till a pellicle form on its surface, and then set it aside to cool and crystallize. The residual liquor will yield more crystals by farther concentration and cooling.

POTASSAE ET SODAE TARTRAS.

Take of Bitartrate of Potash, sixteen ounces ;
Carbonate of soda, twelve ounces ;
Boiling water, four pints ;
Proceed for this preparation exactly as for
the tartrate of potash.

POTASSII IODIDUM.

Take of Iodine (dry), five ounces ;
Fine iron-wire, three ounces ;
Water, four pints ;
Carbonate of Potash (dry), two
ounces and six drachms.

With the water, iodine and iron-wire prepare the solution of iodide of iron as directed in p. 103. Add immediately, while it is hot, the carbonate of potash previously dissolved in a few ounces of water, stir carefully, filter the product, and wash the powder on the filter with a little water. Concentrate the liquor at a temperature short of ebullition, till a dry salt be obtained, which is to be purified from a little red oxide of iron and other impurities, by dissolving it in less than its own weight of boiling water, or still better by boiling it in twice its weight of rectified spirit, filtering the solution, and setting it aside to crystallize. More crystals will be obtained by concentrating and cooling the residual liquor.

POTASSII SULPHURETUM.

Take of Sulphur, one ounce ;

Carbonate of potash, four ounces ;
Triturate them well together, and heat them in a covered crucible till they form a uniform fused mass : which, when cold, is to be broken into fragments, and kept in well-closed vessels.

SODAE AQUA EFFERVESCENS.

Take of Bicarbonate of Soda, one drachm ;
Water, one pint ;

Dissolve the Bicarbonate in the water and saturate it with carbonic acid under strong pressure. Preserve the liquid in well-closed vessels.

SODAE BICARBONAS.

Fill with fragments of marble a glass jar, open at the bottom and tubulated at the top ; close the bottom in such way as to keep in the marble without preventing the free passage of a fluid ; connect the tubulature closely by a bent tube and corks with an empty bottle, and this in like manner with another bottle filled with one part of Carbonate of soda and two parts of Dried carbonate of soda well triturated together ; and let the tube be long enough to reach the bottom of the bottle. Before closing the last cork closely, immerse the jar to the top in diluted

muriatic acid contained in any convenient vessel; when the whole apparatus is thus filled with carbonic acid gas, secure the last cork tightly; and let the action go on till next morning, or till gas is no longer absorbed by the salt. Remove the damp salt which is formed, and dry it, either in the air without heat, or at a temperature not above 120°.

SODAE CARBONAS SICCATUM.

Heat any convenient quantity of Carbonate of Soda in a shallow vessel till it is dry, then urge it with a red heat in a crucible, and reduce it to powder when cold.

SODAE MURIAS PURUM.

Take any convenient quantity of Muriate of Soda; dissolve it in boiling water; filter the solution: and boil it down over the fire, skimming off the crystals which form; wash the crystals quickly with cold water and dry them.

SODAE PHOSPHAS.

Take of Bones burnt to whiteness, ten pounds;
Sulphuric acid, two pints, and four fluidounces:
Carbonate of Soda, a sufficiency;
Pulverize the bones and mix them with the acid; add gradually six pints of water; digest for three days, replacing the water which

evaporates ; add six pints of boiling water, and strain through strong linen : pass more boiling water through the mass on the filter till it comes away nearly tasteless. Let the impurities subside in the united liquors, pour off the clear fluid, and concentrate to six pints. Let the impurities again settle ; and to the clear liquor, which is to be poured off and heated to ebullition, add carbonate of soda, previously dissolved in boiling water, until the acid is completely neutralized. Set the solution aside to cool and crystallize. More crystals will be obtained by successively evaporating, adding a little carbonate of soda till the liquid exerts a feeble alkaline reaction on litmus-paper, and then allowing it to cool. Preserve the crystals in well-closed vessels.

SODAE SULPHAS.

Take of the Salt which remains after preparing
pure muriatic acid, two pounds ;
Boiling-water, three pints :
White marble, in powder, a suffi-
ciency ;

Dissolve the salt in the water, add the marble so long as effervescence takes place, boil the liquid, and when neutral filter it ; wash the insoluble matter with boiling-water, adding the water to the original liquid ; concentrate till a pellicle begins to form, and then let the liquid cool and crystallize.

STANNI PULVIS.

Melt tin in an iron vessel; pour it into an earthen-ware mortar heated a little above the melting point of the metal; triturate briskly as the metal cools, ceasing as soon as a considerable proportion is pulverized; sift the product, and repeat the process with what remains in the sieve.

ZINCI OXIDUM.

Take of Sulphate of Zinc, twelve ounces;
Carbonate of Ammonia, six ounces;
Dissolve each in two pints of water; mix the solutions; collect the precipitate on a filter of linen or calico; wash it thoroughly; squeeze and dry it, and expose it for two hours to a red heat.

ZINCI SULPHAS.

This salt may be prepared either by dissolving fragments of zinc in diluted sulphuric acid till a neutral liquid be obtained, filtering the solution, and concentrating sufficiently for it to crystallize on cooling,—or by repeatedly dissolving and crystallizing the impure sulphate of zinc of commerce, until the product when dissolved in water does not yield a black precipitate with tincture of galls, and corresponds with the characters laid down for sulphate of zinc in the List of the *Materia Medica*.

MIXTURES AND EMULSIONS.

MISTURA ACACIAE.

Take of Mucilage, three fluidounces ;
 Sweet Almonds, one ounce and
 two drachms ;
 Pure Sugar, five drachms ;
 Water, two pints ;

Steep the almonds in hot water and peel them ; beat them to a smooth pulp in an earthen-ware or marble mortar, first with the sugar, and then with the mucilage ; add the water gradually, stirring constantly ; then strain through linen or calico.

MISTURA AMYGDALARUM.

Take of the Conserve of Almonds, two
 ounces ;
 Water, two pints ;

Add the water gradually to the confection, triturating constantly ; and then strain through linen or calico.

Or

Take of Sweet Almonds, one ounce and two
 drachms ;
 Pure sugar, five drachms ;
 Mucilage, half a fluidounce ;
 Water, two pints ;

Steep the almonds in hot water and peel them ; and proceed as for the Mistura acaciæ.

MISTURA ALTHÆÆ.

Take of Althæa-root, dried, four ounces ;
Raisins, freed of the seeds, two
ounces ;

Boiling-water, five pints ;

Boil down to three pints ; strain through
linen or calico, and when the sediment has
subsided, pour off the clear liquor for use.

MISTURA CAMPHORÆ.

Take of Camphor, one scruple ;
Sweet Almonds, and
Pure Sugar, of each half an ounce ;
Water, one pint.

Steep the almonds in hot water and peel
them ; rub the camphor and sugar well to-
gether in a mortar ; add the almonds ; beat
the whole into a smooth pulp ; add the wa-
ter gradually, with constant stirring, and
then strain.

MISTURA CAMPHORÆ CUM MAGNESIA.

Take of Camphor, ten grains ;
Carbonate of Magnesia, twenty-five
grains ;
Water, six fluidounces.

Triturate the camphor and carbonate of
magnesia together, adding the water gra-
dually.

MISTURA CREASOTI.

Take of Creasote, and

Acetic acid, of each sixteen minims;

Compound Spirit of Juniper, and

Syrup, of each one fluidounce;

Water, fourteen fluidounces;

Mix the creasote with the acid, then gradually the water, and lastly the syrup and spirit.

MISTURA CRETAE.

Take of Prepared Chalk, ten drachms;

Pure Sugar, five drachms;

Mucilage, three fluidounces;

Spirit of Cinnamon, two ounces;

Water, two pints.

Triturate the chalk, sugar, and mucilage together; and then add gradually the water and spirit of cinnamon.

MISTURA FERRI COMPOSITA.

Take of Myrrh, bruised, two drachms;

Carbonate of Potash, one drachm;

Rose-water, eighteen fluidounces;

Sulphate of Iron in coarse powder,
two scruples and a-half;

Spirit of Nutmeg, one fluidounce;

Pure Sugar, two drachms.

Triturate the myrrh with the spirit of nutmeg and carbonate of potash, add the rose-water and sugar, with constant trituration,

and then the sulphate of iron. Preserve the mixture in well-closed bottles.

MISTURA GUAIACI.

Take of Guaiac, three drachms ;
Sugar, half an ounce ;
Mucilage, half a fluidounce ;
Cinnamon-water, nineteen fluid-
ounces and a half.

Triturate the guaiac with the sugar, then with the mucilage, and then add gradually the cinnamon water, with constant trituration.

MISTURA HORDEI.

Take of Pearl-Barley,
Figs, sliced,
Raisins, freed of the seeds, of each
two ounces and a half.
Liquorice-root, sliced and bruised,
five drachms ;
Water, five pints and a-half ;

Clean the Barley, if necessary, by washing it with cold water ; boil it with four pints and a half of the water down to two pints ; add the figs, raisins, and liquorice-root, with the remaining pint of water ; and again boil down to two pints ; then strain.

MISTURA SCAMMONII.

Take of Resin of Scammony, seven grains ;

Unskimmed Milk, three fluid-ounces ;

Triturate the resin with a little of the milk, and gradually with the rest of it till a uniform emulsion is formed.

MUCILAGO.

Take of Gum-arabic, nine ounces ;

Water (cold), one pint.

Mix them, allow the gum to dissolve without applying heat, but with occasional stirring ; then strain through linen or calico.

MUCILAGO AMYLI.

Take of Starch, half an ounce ;

Water, one pint.

Triturate the starch with a little of the water ; add the rest of the water gradually ; then boil for a few minutes.

MUCILAGO TRAGACANTHÆ.

Take of Tragacanth, two drachms ;

Boiling-water, nine fluidounces ;

Macerate for four-and-twenty hours, then triturate to dissolve the gum, and express through linen or calico.

OINTMENTS, LINIMENTS, AND
CERATES.

CERATUM CALAMINAE.

Take of Calamine, prepared in the same manner as Prepared Chalk, one part ;
Simple Cerate, five parts ;
Mix them well together.

CERATUM SABINAE.

Take of fresh Savin, two parts ;
Bees'-wax, one part ;
Axunge, four parts ;
Melt the wax and axunge together, add the savin, and boil them together, till the leaves are friable ; then strain.

CERATUM SIMPLEX.

Take of Olive-oil, six parts ;
White-wax, three parts ;
Spermaceti, one part ;
Heat the oil gently, add the wax and spermaceti, stir the whole briskly when it is fluid, and continue the agitation as it cools.

LINIMENTUM AMMONIAE.

Take of Olive-oil, two fluidounces ;
Aqua ammoniæ (D. 960), one fluidounce ;
Mix and agitate them well together.

LINIMENTUM AMMONIAE COMPOSITUM.

Take of Stronger Aqua Ammoniaë (D. 880),
five fluidounces ;

Tincture of Camphor, two fluid-
ounces ;

Spirit of Rosemary, one fluidounce ;

Mix them well together. This liniment may
be also made weaker for some purposes with
three fluidounces of tincture of camphor and
two of Spirit of rosemary.

LINIMENTUM CALCIS.

Take of Linseed-oil, and

Lime-water, of each equal measures ;

Mix and agitate them well together.

LINIMENTUM CAMPHORAE.

Take of Olive-oil, four fluidounces ;

Camphor, one ounce.

Rub them together in a mortar till the cam-
phor is dissolved.

LINIMENTUM OPII.

Take of Castile Soap, six ounces ;

Opium, an ounce and a-half ;

Camphor, three ounces ;

Oil of Rosemary, six fluidrachms ;

Rectified spirit, two pints.

Macerate the soap and opium in the spirit
for three days ; filter, add the oil and cam-
phor, and agitate briskly.

LINIMENTUM SAPONIS.

Take of Castile Soap, five ounces ;
Camphor, two ounces and a-half ;
Volatile oil of Rosemary, six fluidrachms.

Rectified Spirit, two pints ;

Digest the soap in the spirit for three days ;
add the camphor and oil, and agitate briskly.

LINIMENTUM SIMPLEX.

Take of Olive oil, four parts ;

White-wax, one part.

Dissolve the wax in the oil with a gentle heat ; and agitate well as the fused mass cools and concretes.

LINIMENTUM TEREBINTHINATUM.

Take of Resinous ointment, four ounces ;

Oil of Turpentine, five fluidounces :

Camphor, half an ounce.

Melt the ointment, and gradually mix with it the camphor and oil, till a uniform liniment be obtained.

UNGUENTUM AERUGINIS.

Take of Resinous ointment, fifteen ounces ;

Verdigris, in fine powder, one ounce :

Melt the ointment, sprinkle into it the powder of verdigris, and stir the mixture briskly as it cools and concretes.

UNGUENTUM ANTIMONIALE.

Take of Axunge, four ounces ;
Tartar-emetic, in very fine powder,
one ounce.

Triturate them carefully together into a smooth and uniform mass.

UNGUENTUM CANTHARIDIS.

Take of Resinous ointment, seven ounces ;
Cantharides, in very fine powder, one
ounce.

Melt the ointment ; sprinkle into it the cantharides powder ; and stir the mixture briskly as it concretes on cooling.

UNGUENTUM INFUSI CANTHARIDIS.

Take of Cantharides in moderately fine powder,
Resin, and
Bees'-wax, of each one ounce ;
Venice Turpentine, and
Axunge, of each two ounces ;
Boiling water, five fluidounces.

Infuse the Cantharides in the water for one night, squeeze strongly, and filter the expressed liquid. Add the axunge, and boil till the water is dispersed. Then add the wax and resin : and when these have become liquid, remove the vessel from the fire, add the Turpentine, and mix the whole thoroughly.

UNGUENTUM CITRINUM.

Take of Pure Nitric acid, eight fluidounces
and six fluidrachms ;
Mercury, four ounces ;
Axunge, fifteen ounces ;
Olive-oil, thirty-two fluidounces.

Dissolve the mercury in the acid with the aid of a gentle heat. Melt the axunge in the oil with the aid of a moderate heat in a vessel capable of holding six times the quantity ; and while the mixture is hot, add the solution of mercury, also hot, and mix them thoroughly. If the mixture do not froth up, increase the heat a little till this take place. Keep this ointment in earthen-ware vessels, or in glass-vessels secluded from the light.

UNGUENTUM COCCULI.

Take any convenient quantity of Cocculus Indicus, separate and preserve the kernels, beat them well in a mortar, first alone and then with a little axunge ; and then add axunge till it amounts altogether to five times the weight of the kernels.

UNGUENTUM CREASOTI.

Take of Axunge, three ounces ;
Creasote, one drachm.
Melt the Axunge, add the Creasote, stir

them briskly, and continue to do so as the mixture concretes on cooling.

UNGUENTUM GALLAE ET OPII.

Take of Galls in fine powder, two drachms ;
Opium in powder, one drachm ;
Axunge, one ounce.

Triturate them together into a uniform mass.

UNGUENTUM HYDRARGYRI.

Take of Mercury, two pounds ;
Axunge, twenty-three ounces ;
Suet, one ounce.

Triturate the mercury with the suet and a little of the axunge till globules are no longer visible ; then add the rest of the axunge, and mix the whole thoroughly. This ointment is not well prepared so long as metallic globules may be seen in it with a magnifier of four powers.

The mercurial ointment with the proportions here directed may be diluted at pleasure with twice or thrice its weight of axunge.

UNGUENTUM IODINEI.

Take of Iodine, one drachm ;
Iodide of potassium, two drachms ;
Axunge, four ounces.

Triturate the Iodine and Iodide together,

132 OINTMENTS, LINIMENTS,

and then add gradually the axunge, continuing the trituration till a uniform ointment be obtained.

UNGUENTUM OXIDI HYDRARGYRI.

Take of Red oxide of mercury, one ounce ;
Axunge, eight ounces.
Triturate them into a uniform mass.

UNGUENTUM PICIS LIQUIDAE.

Take of Tar, five ounces :
Bees'-wax, two ounces.
Melt the wax with a gentle heat, add the tar, and stir the mixture briskly while it concretes on cooling.

UNGUENTUM PLUMBI ACETATIS.

Take of Simple ointment, twenty ounces ;
Acetate of lead, in fine powder, one ounce.
Mix them thoroughly.

UNGUENTUM PLUMBI CARBONATIS.

Take of Simple ointment, five ounces ;
Carbonate of Lead, one ounce ;
Mix them thoroughly.

UNGUENTUM PRECIPITATI ALBI.

Take of White Precipitate, two drachms ;
Axunge, three ounces ;
Melt the axunge, add the white precipitate,

and stir the mixture briskly while it concretes on cooling.

UNGUENTUM RESINOSUM.

Take of Resin, five ounces ;

 Axunge, eight ounces ;

 Bees'-wax, two ounces ;

Melt them together with a gentle heat, and then stir the mixture briskly while it cools and concretes.

UNGUENTUM SIMPLEX.

Take of Olive-oil, five fluidounces and a half ;

 White wax, two ounces ;

Melt the wax in the oil, and stir the mixture briskly while it concretes on cooling.

UNGUENTUM SULPHURIS.

Take of Axunge, four ounces ;

 Sublimed Sulphur, one ounce ;

Mix them thoroughly together.

UNGUENTUM ZINCI.

Take of Simple Liniment six ounces ;

 Oxide of Zinc, one ounce.

Mix them thoroughly together.

OXIDIFIABLE NON-METALLIC
ELEMENTS.

CARBO ANIMALIS PURIFICATUS.

Take of Ivory-black, one pound ;
Muriatic Acid, (commercial), and
Water, of each twelve fluidounces ;
Mix the acid and water ; add gradually the
ivory-black, stirring occasionally. Digest
with a gentle heat for two days, agitating
from time to time. Then boil ; dilute with
two pints of water ; collect the undissolved
charcoal on a filter of linen or calico, and
wash it with water till what passes through
scarcely precipitates with solution of carbo-
nate of soda. Heat the charcoal first mo-
derately, and then to redness in a closely-
covered crucible.

CHLORINEI AQUA.

Take of Muriate of Soda, sixty grains ;
Sulphuric Acid (commercial), two
fluidrachms ;
Red Oxide of Lead, three hundred
and fifty grains ;
Water, eight fluidounces ;
Triturate the muriate of soda and oxide to-
gether ; put them into the water contained
in a bottle with a glass-stopper ; add the
acid ; agitate occasionally till the red oxide

becomes almost all white. Allow the insoluble matter to subside before using the liquid.

IODINEUM.

Iodine, as obtained in commerce, being almost always adulterated with variable proportions of water, and being consequently unfit for making pharmaceutic preparations of fixed and uniform strength, it must be dried by being placed in a shallow basin of earthen-ware in a small confined space of air with ten or twelve times its weight of fresh-burnt lime, till it scarcely adheres to the inside of a dry bottle.

IODINEI LIQUOR COMPOSITUS.

Take of Iodine, two drachms ;

Iodide of potassium, an ounce ;

Distilled water, sixteen fluidounces;

Dissolve the iodide and iodine in the water with gentle heat and agitation.

SULPHUR SUBLIMATUM.

Sublime sulphur in a proper vessel ; wash the powder thus obtained with boiling water in successive portions till the water ceases to have an acid taste ; then dry the sulphur with a gentle heat.

PILLS.

PILULAE ALOES.

Take of Socotorine Aloes, and
Castile Soap, equal parts ;
Conserve of Red Roses, a sufficiency;
Beat them into a proper pill mass. This
pill may be also correctly made with the finer
qualities of East-Indian aloes, as the Soco-
torine variety is very scarce ; and many, not
without reason, prefer the stronger Barba-
dos aloes.

PILULAE ALOES ET ASSAFOETIDAE.

Take of Aloes (Socotorine or East-Indian);
Assafoetida, and
Castile Soap, equal parts ;
Beat them with Conserve of Red roses into
a proper pill mass.

PILULAE ALOES ET FERRI.

Take of Sulphate of iron, three parts ;
Barbados aloes, two parts ;
Aromatic powder, six parts ;
Conserve of Red roses, eight parts ;
Pulverize the aloes and sulphate of iron se-
parately ; mix the whole ingredients ; and
beat them into a proper mass ; which is to
be divided into five-grain pills.

PILULAE ALOES ET MYRRHAE.

Take of Aloes (Socotorine or East-Indian)
 four parts ;
 Myrrh, two parts ;
 Saffron, one part ;

Beat them into a proper mass with a sufficient quantity of Conserve of Red roses.

PILULAE ASSAFOETIDAE.

Take of Assafoetida,
 Galbanum, and
 Myrrh, three parts of each ;
 Conserve of Red Roses, four parts
 or a sufficiency ;

Mix them, and beat them into a proper pill mass.

PILULAE CAMBOGIAE.

Take of Gamboge,
 East-Indian or Barbados aloes, and
 Aromatic powder, of each, one part ;
 Castile Soap, two parts ;

Pulverize the gamboge and aloes separately, mix all the powders, add the soap, and then a sufficiency of syrup ; beat the whole into a proper pill mass.

PILULAE CALOMELANOS COMPOSITAE.

Take of Calomel, and
 Golden sulphuret of antimony, of
 each one part ;

Guaiac, in fine powder, and
Treacle, of each two parts ;
Mix the solids in fine powder, then the
treacle, and beat the whole into a proper pill
mass ; to be divided into six-grain pills.

PILULAE CALOMELANOS ET OPII.

Take of Calomel, three parts ;
Opium, one part ;
Conserve of Red roses, a sufficiency ;
Beat them into a proper mass, which is to
be divided into pills, each containing two
grains of calomel.

PILULAE COLOCYNTHIDIS.

Take of Socotorine or East-Indian aloes, and
Scammony, of each eight parts ;
Colocynth, four parts ;
Sulphate of potash, and
Oil of Cloves, of each one part ;
Rectified spirit, a sufficiency ;
Pulverize the aloes, scammony, and sulphate
of potash together ; mix with them the co-
locynth previously reduced to fine powder ;
add the oil of cloves ; and with the aid of a
small quantity of rectified spirit beat the
whole into a proper pill mass ; which is to be
divided into five-grain pills.

PILULAE COLOCYNTHIDIS ET HYOSCYAMI.

Take of the Colocynth-pill mass, two parts ;

Extract of Hyoscyamus, one part ;
Beat them well together, adding a few drops
of rectified spirit, if necessary ; and divide
the mass into five-grain pills.

PILULAE CUPRI AMMONIATI.

Take of Ammoniated Copper in fine powder,
one part ;
Bread-crumbs, six parts ;
Solution of Carbonate of Ammonia,
a sufficiency ;

Beat them into a proper mass ; and divide it
into pills, containing each half a grain of
ammoniated copper.

PILULAE DIGITALIS ET SCILLAE.

Take of Digitalis, and
Squill, of each one part ;
Aromatic Electuary, two parts ;

Beat them into a proper mass with conserve
of Red roses ; and divide the mass into four-
grain pills.

PILULAE FERRI CARBONATIS.

Take of the Saccharine Carbonate of Iron,
four parts ;
Conserve of Red roses, one part ;
Beat them into a proper mass, to be divided
into five-grain pills.

PILULAE FERRI SULPHATIS.

Take of dried Sulphate of Iron, two parts ;

Extract of Taraxacum, five parts ;
Conserve of Red roses, two parts ;
Liquorice-root powder, three parts ;
Beat them together into a proper mass, which
is to be divided into five-grain pills.

PILULAE HYDRARGYRI.

Take of Mercury, two parts :
Liquorice-root, in powder, one part ;
Conserve of Red roses, three parts ;
Beat the mercury and conserve into a uni-
form mass till globules of mercury can no
longer be detected, then add the liquorice-
root, and beat the whole again into a proper
mass, which is to be divided into five-grain
pills.

PILULAE IPECACUANHAE ET OPII.

Take of the Powder of Ipecacuan and Opium,
three parts.
Conserve of Red roses, one part ;
Beat them into a proper mass, which is to be
divided into four-grain pills.

PILULAE OPII *sive* THEBAICAE.

Take of Opium, one part :
Sulphate of Potass, three parts ;
Conserve of Red roses, one part ;
Beat them into a proper mass, which is to
be divided into five-grain pills ;
It is to be observed that this pill contains

twice as much opium as the Opiate pill of the last Latin edition of this Pharmacopœia.

PILULAE PLUMBI OPIATAE.

Take of Acetate of lead, six parts ;

Opium, one part ;

Conserve of Red roses, about one part ;

Beat them into a proper mass, which is to be divided into four-grain pills.—This pill may be made also with twice the quantity of opium.

PILULAE RHEI.

Take of Rhubarb, in fine powder, nine parts ;

Acetate of potash, one part :

Conserve of Red roses, five parts ;

Beat them into a proper mass, and divide it into five-grain pills.

PILULAE RHEI COMPOSITAE.

Take of Rhubarb, in fine powder, twelve parts ;

Aloes, in fine powder, nine parts ;

Myrrh, and

Castile soap, of each six parts ;

Oil of Peppermint, one part ;

Conserve of Red roses, five parts ;

Mix them, and beat them into a proper mass, and divide it into five-grain pills.—

This pill may be also made without oil of peppermint, when so preferred.

PILULAE RHEI ET FERRI.

Take of Dried Sulphate of Iron, four parts ;
Extract of Rhubarb, ten parts ;
Conserve of Red Roses, five parts.
Beat them into a proper pill mass, and divide it into five-grain pills.

PILULAE SCILLAE.

Take of Squill, in fine powder, five parts ;
Ammoniac,
Ginger, in fine powder, and
Spanish Soap, of each, four parts ;
Conserve of Red roses, two parts ;
Mix the powders, add the other articles, beat them into a uniform mass, and divide it into five-grain pills.

PILULAE STYRACIS.

Take of Opium, and
Saffron, of each one part ;
Extract of Storax, two parts ;
Beat them into a uniform mass, which is to be divided into four-grain pills.

PLASTERS.

EMPLASTRUM AMMONIACI.

Take of Ammoniac, five ounces ;
Distilled vinegar, nine fluidounces ;
Dissolve the ammoniac in the vinegar, and
then evaporate to a proper consistence over
the vapour-bath, frequently stirring the li-
quid.

EMPLASTRUM AMMONIACI ET HYDRARGYRI.

Take of Ammoniac, one pound ;
Mercury, three ounces ;
Olive-oil, one fluidrachm ;
Sulphur, eight grains ;
Heat the oil, add the sulphur by degrees, stir
them till they unite, add the mercury, and
triturate till the globules disappear, then add
also the ammoniac previously liquefied, and
mix the whole carefully.

EMPLASTRUM ASSAFOETIDAE.

Take of Litharge plaster, and
Assafoetida, of each, two ounces ;
Galbanum, and
Bees'-wax, of each, one ounce ;
Liquefy the gum-resins together and strain
them, then add the plaster and wax also in
the fluid state, and mix them all thoroughly.

EMPLASTRUM BELLADONNAE.

Take of Resin plaster, three ounces ;
Extract of Belladonna, an ounce
and a-half ;
Liquefy the plaster with a gentle heat, add
the extract, and agitate briskly.

EMPLASTRUM CANTHARIDIS.

Take of Cantharides, in very fine powder,
Resin,
Bees'-wax, and
Suet, of each two ounces ;
Liquefy the fats, remove from the heat,
sprinkle in the cantharides, and stir briskly
as the mixture concretes on cooling.

EMPLASTRUM CANTHARIDIS COMPOSITUM.

Take of Venice Turpentine, four ounces and
a-half ;
Burgundy-Pitch, and
Cantharides, of each three ounces ;
Bees'-wax, one ounce ;
Verdigris, half an ounce ;
White-mustard seed, and
Black Pepper, of each two drachms ;
Liquefy the wax and Burgundy-pitch, add
the turpentine, and while the mixture is hot
sprinkle into it the remaining articles pre-
viously in fine powder, and mixed together.
Stir the whole briskly as it concretes in
cooling.

EMPLASTRUM FERRI.

Take of Litharge-plaster, three ounces ;
Resin, six drachms ;
Olive-oil, three fluidrachms and a
half ;
Bees'-wax, three drachms ;
Red oxide of iron, one ounce ;
Triturate the oxide of iron with the oil, and
add the mixture to the other articles pre-
viously liquefied by gentle heat. Mix the
whole thoroughly.

EMPLASTRUM GUMMOSUM.

Take of Litharge plaster, four ounces ;
Ammoniac,
Galbanum, and
Bees'-wax, of each half an ounce ;
Melt the gum-resins together and strain
them ; melt also together the plaster and
wax ; add the former to the latter mixture,
and mix the whole thoroughly.

EMPLASTRUM HYDRARGYRI.

Take of Mercury three ounces ;
Olive-oil, nine fluidrachms ;
Resin, one ounce ;
Litharge plaster, six ounces ;
Liquefy together the oil and resin, let them
cool, add the mercury, and triturate till its
globules disappear ; then add to the mixture
the plaster previously liquefied ; and mix
the whole thoroughly.

EMPLASTRUM LITHARGYRI.

Take of Litharge, in fine powder, five ounces ;
Olive oil, twelve fluidounces ;
Water, three fluidounces.

Mix them ; boil and stir constantly till the oil and litharge unite, replacing the water if it evaporate too far.

EMPLASTRUM OPII.

Take of Powder of Opium, half an ounce ;
Burgundy pitch, three ounces ;
Litharge plaster, twelve ounces ;
Liquefy the plaster and pitch, add the opium by degrees, and mix them thoroughly.

EMPLASTRUM PICIS.

Take of Burgundy pitch, one pound and a half ;
Resin, and
Bees'-wax, of each two ounces ;
Oil of mace, half an ounce ;
Olive oil, one fluidounce ;
Water, one fluidounce.

Liquefy the pitch, resin and wax with a gentle heat ; add the other articles ; mix them well together ; and boil till the mixture acquires the proper consistence.

EMPLASTRUM RESINOSUM.

Take of Litharge plaster, five ounces ;
Resin, one ounce ;

Melt them together with a moderate heat, and stir the mixture well till it concretes on cooling.

EMPLASTRUM SAPONIS.

Take of Litharge plaster, four ounces ;

Gum plaster, two ounces ;

Castile soap, in shavings, one ounce ;

Melt the plasters together with a moderate heat, add the soap, and boil for a little.

EMPLASTRUM SIMPLEX.

Take of Bees'-wax, three ounces ;

Suet, and

Resin, of each two ounces ;

Melt them together with a moderate heat, and stir the mixture briskly till it concretes on cooling.

POWDERS.

PULVIS ALUMINIS COMPOSITUS.

Take of Alum, four ounces ;
Kino, one ounce ;
Mix them and reduce them to fine powder.

PULVIS AROMATICUS.

Take of Cinnamon,
Cardamom seeds, and
Ginger, of each equal parts ;
Mix them, and reduce to a very fine powder,
which is to be kept in well closed glass vessels.

PULVIS CRETAE COMPOSITUS.

Take of Prepared chalk, four ounces ;
Cinnamon, in fine powder, one
drachm and a half ;
Nutmeg, in fine powder, a drachm ;
Triturate them well together.

PULVIS CRETAE OPIATUS.

Take of Compound chalk powder, six ounces ;
Powder of Opium, four scruples ;
Triturate them together thoroughly.

PULVERES EFFERVESCENTES.

Take of Tartaric acid, one ounce ;
Bicarbonate of Soda, one ounce and
54 grains ;

Or

Bicarbonate of Potass, one ounce
and 160 grains ;

Reduce the acid and either bicarbonate separately to fine powder, and divide each into sixteen powders ; preserve the acid and alkaline powders in separate papers of different colours.

PULVIS IPECACUANHAE COMPOSITUS.

Take of Ipecacuan, in powder and
Powder of Opium, of each, one
ounce ;
Sulphate of Potass, eight ounces ;
Triturate them together thoroughly.

PULVIS JALAPAE COMPOSITUS.

Take of Jalap, in powder, one ounce ;
Bitartrate of Potash, two ounces ;
Triturate them to a very fine powder.

PULVIS RHEI COMPOSITUS.

Take of Magnesia, one pound ;
Ginger in fine powder, two ounces ;
Rhubarb in fine powder, 4 ounces ;
Mix them thoroughly, and preserve the
powder in well-closed bottles.

PULVIS SALINUS COMPOSITUS.

Take of Pure Muriate of Soda, and
Sulphate of Magnesia, four ounces ;

Sulphate of Potash, three ounces ;
Dry the salts separately with a gentle heat,
and pulverize each, then triturate them well
together, and preserve the mixture in well-
closed vessels.

PULVIS SCAMMONII COMPOSITUS.

Take of Scammony, and
Bitartrate of Potash, of each, equal
parts.
Triturate them together to a very fine powder.

PULVIS TRAGACANTHAE COMPOSITUS.

Take of Tragacanth, bruised,
Gum-arabic, bruised, and
Starch, of each one ounce and a half ;
Pure sugar, three ounces ;
Reduce the starch and sugar together to
powder, then add the tragacanth and gum-
arabic, and pulverize the mixture thoroughly.

SPIRITS.

SPIRITUS AMMONIAE.

Take of Rectified spirit, two pints ;
Fresh-burnt Lime, twelve ounces ;
Muriate of ammonia, in very fine
powder, eight ounces ;
Water, six fluidounces and a half ;

Let the lime be slaked with the water in an iron or earthen-ware vessel, and cover the vessel till the powder be cold ; mix the lime and muriate of ammonia quickly and thoroughly in a mortar, and transfer the mixture at once into a glass retort ; adapt to the retort a tube which passes nearly to the bottom of a bottle containing the rectified spirit ; heat the retort in a sand-bath gradually, so long as any thing passes over, preserving the bottle cool. The bottle should be large enough to contain one-half more than the spirit used.

SPIRITUS AMMONIAE AROMATICUS.

Take of Spirit of ammonia, eight fluidounces ;
Volatile oil of Lemon-peel, one fluid-
drachm ;
Volatile oil of Rosemary, one fluid-
drachm and a half ;
Dissolve the oils in the spirit by agitation.

SPIRITUS AMMONIAE FOETIDUS.

Take of Spirit of ammonia, ten fluidounces
and a half ;

Assafœtida, half an ounce ;

Break the assafœtida into small fragments,
digest it in the spirit for twelve hours, and
distil over ten fluidounces and a half by means
of a vapour-bath heat.

SPIRITUS CARUI.

Take of Caraway, bruised, half a pound ;

Proof-spirit, seven pints ;

Macerate for two days in a covered vessel ;
add a pint and a half of water ; and distil off
seven pints.

SIRITUS CASSIAE.

Take of Cassia in coarse powder, one pound.

Proceed as for the Spirit of Caraway.

SPIRITUS CINNAMOMI.

Take of Cinnamon, in coarse powder, one
pound. Proceed as for the spirit of Caraway.

SPIRITUS JUNIPERI COMPOSITUS.

Take of Juniper berries, bruised, a pound ;

Fennel bruised, and

Caraway bruised, of each, an ounce
and a half ;

Proof-spirit, seven pints ;

Water, two pints ;

Macerate the fruits in the spirit for two days, add the water, and distil off seven pints.

SPIRITUS LAVANDULAE.

Take of Lavender, fresh, two pounds and a-half;

Rectified spirit, a gallon;

Mix them, and with the heat of a vapour-bath distil over seven pints.

SPIRITUS LAVANDULAE COMPOSITUS.

Take of Spirit of Lavender two pints;

Spirit of Rosemary, twelve fluid-ounces;

Cinnamon in coarse powder, one ounce;

Cloves bruised, two drachms;

Nutmeg bruised, half an ounce;

Red Sandal-wood, in shavings, three drachms.

Let the whole macerate for seven days and then strain the liquor through calico.

SPIRITUS MENTHAE.

Take of Peppermint, fresh, one pound and a-half. Proceed as for Spirit of Caraway.

SPIRITUS MYRISTICAE.

Take of Bruised Nutmeg, two ounces and a-half;

Proof-spirit, one gallon ;

Water one pint ;

Mix them together, and distil over one gallon.

SPIRITUS PIMENTAE.

Take of Pimento bruised, half a pound. Proceed as for the Spirit of Caraway.

SPIRITUS ROSMARINI.

Take of Rosemary, two pounds and a-half. Proceed as for the Spirit of Lavender.

SPIRITUS TENUIOR.

Take of Rectified-spirit, two pints ;

Distilled water, one pint ;

Mix them. The density of the product should be 912.

SYRUPS.

SYRUPUS ACETI.

Take of Vinegar, French in preference, eleven fluidounces ;

Pure sugar, fourteen ounces.

Boil them together.

SYRUPUS ALTHÆAE.

Take of Althæa-root, fresh and sliced, eight ounces.

Boiling water, four pints ;

Pure sugar, two pounds and a-half ;

Boil the althæa-root with the water down to two pints ; strain and express strongly through calico ; let the impurities subside ; and dissolve the sugar in the clear liquor with the aid of heat.

SYRUPUS AURANTII.

Take of fresh Bitter Orange-Peel, two ounces and a-half ;

Boiling water, one pint ;

Pure sugar, three pounds ;

Infuse the peel in the water for twelve hours in a covered vessel, pour off the liquor, and filter it if necessary ; add the sugar to the liquor and dissolve it with the aid of heat.

SYRUPUS CROCI.

Take of Saffron, ten drachms ;

Boiling water, one pint ;

Sugar, three pounds.

Proceed as for the Syrup of orange-peel.

SYRUPUS IPECACUANHAE.

Take of Ipecacuan in coarse powder, four ounces ;

Rectified-spirit, one pint ;

Proof-spirit, and

Water, of each fourteen fluidounces ;

Syrup, seven pints ;

Digest the ipecacuan in fifteen fluidounces of the rectified spirit at a gentle heat for twenty-four hours ; strain, squeeze the residuum and filter. Repeat this process with the residuum and proof-spirit ; and again with the water. Unite the fluids, and distil off the spirit till the residuum amount to twelve ounces ; add to the residuum five fluidounces of the rectified spirit, and then the syrup.

SYRUPUS LIMONUM.

Take of Lemon-juice freed of impurities by subsidence and filtration, a pint ;

Sugar, two pounds and a half ;

Dissolve the sugar in the lemon-juice with the aid of a gentle heat, and after twenty-four hours' rest remove the scum, and pour the clear liquor from the dregs.

SYRUPUS PAPAVERIS.

Take of Poppy-heads without the seeds, one pound and a-half ;

Boiling water, fifteen pints ;

Pure sugar, three pounds ;

Slice the Poppy heads, infuse them in the water for twelve hours, boil down to five pints, strain and express strongly through calico, boil again down to two pints and a-half ; then add the sugar and dissolve it with the aid of heat.

SYRUPUS RHAMNI.

Take of fresh juice of Buckthorn berries, four pints ;

Ginger sliced, and

Pimento bruised, of each six drachms ;

Pure sugar, four pounds ;

Let the juice rest for three days ; pour off the clear liquor and filter it. Digest the ginger and pimento in a pint of the filtered liquor at a gentle heat for four hours, and filter. Boil down the rest of the juice to one pint and a-half ; mix the two liquors, add the sugar, and dissolve it with the aid of heat.

SYRUPUS RHOEADOS.

Take of Corn-poppy petals, one pound ;

Boiling water, one pint ;

Pure sugar, two pounds and a-half ;

Heat the water over a vapour-bath, add the petals by degrees, stirring occasionally ; remove the vessel from the bath, infuse for twelve hours ; strain and express the liquor ; add to it the sugar, and dissolve this with the aid of heat.

SYRUPUS ROSAE CENTIFOLIAE.

Take of fresh Damask-rose petals, one pound ;
Boiling water, three pints ;
Pure sugar, three pounds ;
Infuse the petals in the water for twelve hours, strain the liquor, and dissolve the sugar in it with the aid of heat.

SYRUPUS ROSAE GALLICAE.

Take of Dried Red rose-petals, two ounces ;
Boiling water, one pint ;
Pure sugar, twenty ounces ;
Proceed as for the syrup of damask-rose.

SYRUPUS SARZAE.

Take of Sarza, in chips, fifteen ounces ;
Boiling water, one gallon ;
Pure sugar, fifteen ounces ;
Infuse the sarsaparilla in the water for twenty-four hours ; boil down to four pints, and strain the liquor while hot ; add the sugar, and evaporate to the consistence of syrup.

SYRUPUS SCILLAE.

Take of Vinegar of squill, three pints ;
Pure sugar in powder, seven pounds ;
Dissolve the sugar in the vinegar of squills
with the aid of a gentle heat and agitation.

SYRUPUS SENNAE.

Take of Senna, four ounces ;
Boiling water, one pint and four
fluidounces ;
Treacle, forty-eight ounces ;
Infuse the senna in the water for twelve
hours ; strain and express strongly through
calico, so as to obtain a pint and two fluid-
ounces at least of liquid. Concentrate the
treacle in the vapour-bath as far as possible,
or till a little taken out upon a rod becomes
nearly concrete on cooling ; and while the
treacle is still hot, add the infusion, stirring
carefully, and removing the vessel from the
vapour-bath as soon as the mixture is com-
plete.

If Alexandrian Senna be used for this pre-
paration, it must be carefully freed of Cyn-
anchum leaves by picking it.

SYRUPUS SIMPLEX.

Take of Pure sugar, ten pounds ;
Boiling water, three pints ;
Dissolve the sugar in the water with the aid
of a gentle heat.

SYRUPUS TOLUTANUS.

Take of Simple syrup, two pounds ;

Tincture of Tolu, an ounce ;

When the syrup has been recently prepared and has not altogether cooled, add the tincture of Tolu by degrees, agitating briskly.

SYRUPUS VIOLAE.

Take of Fresh Violets, one pound ;

Boiling water, two pints and a half ;

Pure sugar, seven pounds and a half ;

Infuse the flowers for twenty-four hours in the water, in a covered glass or earthen-ware vessel ; strain without squeezing, and dissolve the sugar in the filtered liquor.

SYRUPUS ZINGIBERIS.

Take of Ginger, two ounces and a half ;

Boiling water, one pint ;

Pure sugar, two pounds and a half ;

Bruise the ginger, infuse it for four hours in the water, and to the strained liquor add the sugar and dissolve it with the aid of heat.

TINCTURES.

TINCTURES are usually made by reducing the solid ingredients to small fragments, coarse powder, or fine powder, macerating them for seven days or upwards in proof-spirit or rectified spirit, straining the solution through linen or calico, and finally expressing the residuum strongly to obtain what fluid is still retained in the mass. A much superior method, however, has been lately introduced, which answers well for most tinctures, namely, the method of displacement by percolation. According to this process, the solid materials, usually in coarse or moderately fine powder, are moistened with a sufficiency of the solvent to form a thick pulp; in twelve hours, or frequently without any delay, the mass is put into a cylinder of glass, porcelain, or tinned iron, open at both ends, but obstructed at the lower end by a piece of calico or linen, tied tightly over it as a filter; and the pulp being packed by pressure, varying as to degree with various articles, the remainder of the solvent is poured into the upper part of the cylinder, and allowed gradually to percolate. In order to obtain the portion of the fluid which is kept in the residuum, an additional quantity of the solvent is poured into the cylinder until the

tincture which has passed through equals in amount the spirit originally prescribed; and the spirit employed for this purpose is then recovered for the most part by pouring over the residuum as much water as there is of spirit retained in it, which may be easily known by an obvious calculation in each case. The method by percolation, where applicable, will be found much more convenient and expeditious, than the mode hitherto commonly followed, and it exhausts the solid materials in general much more completely. As considerable practice, however, is required for managing the details in different cases, more especially in regard to the degree of minuteness of division of the solids, and the degree of firmness with which they are to be packed in the cylinder, we have thought it right to direct that the method by maceration may be followed as an alternative. But the method by percolation is now preferred by all who have made sufficient trial of it to apply it correctly.

TINCTURA ALOES.

Take of Aloes (Socotorine or Indian), in coarse powder, one ounce;
Extract of Liquorice, three ounces;
Rectified-spirit, twelve fluidounces;
Water, one pint and eight fluidounces;
Mix them and digest for seven days, with occasional agitation; filter the clear liquor,

separated from the sediment. This tincture cannot without difficulty and delay be prepared by percolation.

TINCTURA ALOES ET MYRRHAE.

Take of Aloes (Socotorine or Indian), in coarse powder four ounces ;

Saffron two ounces ;

Tincture of Myrrh, two pints ;

Digest for seven days ; and filter the clear superincumbent liquor. This tincture cannot be well prepared by percolation.

TINCTURA ASSAFOETIDAE.

Take of Assafoetida, in small fragments, five ounces ;

Rectified spirit, two pints ;

Digest for seven days, and filter the clear liquor. This tincture cannot be made by percolation without much delay.

TINCTURA AURANTII.

Take of Bitter Orange-peel, dried, three ounces and a-half ;

Proof-spirit, two pints ;

Digest for seven days, strain and express strongly, and filter the liquor. This tincture may be prepared by percolation, by cutting the peel into small fragments, macerating it in a little of the spirit for twelve hours, and beating the mass into a coarse pulp before putting it into the percolator.

TINCTURA BENZOINI COMPOSITA.

Take of Benzoin, in coarse powder, four ounces ;

Peru-balsam, two ounces and a half ;

East-Indian Aloes, half an ounce ;

Rectified spirit, two pints.

Digest for seven days, pour off the clear liquor, and filter it.

TINCTURA BUCKU.

Take of Bucku, five ounces ;

Proof-spirit, two pints ;

Digest for seven days, pour off the clear liquor, and filter it. This tincture may be conveniently and quickly made also by the process of percolation.

TINCTURA CALUMBAE.

Take of Calumba, in small fragments (if by percolation, in moderately fine powder), three ounces ;

Proof-spirit, two pints ;

Digest for seven days, pour off the clear liquor, express the residuum strongly, and filter the liquors. This tincture is much more conveniently prepared by the process of percolation, allowing the powder to be soaked with a little of the spirit for six hours before putting it into the percolator.

TINCTURA CAMPHORÆ.

Take of Camphor, in small fragments, two ounces and a half ;
Rectified Spirit, two pints ;
Dissolve the camphor in the spirit.

TINCTURA CANTHARIDIS.

Take of Cantharides, half an ounce ;
Proof-spirit, two pints ;
Digest for seven days, strain, and express strongly the residuum ; filter the liquor. This tincture may be obtained more conveniently and expeditiously by percolation, provided the cantharides be reduced to coarse powder, and left with a little of the spirit in the state of pulp for twelve hours before the process of percolation is commenced.

TINCTURA CAPSICI.

Take of Capsicum bruised, (or if percolation be followed, in moderately fine powder,) ten drachms ;
Proof-spirit, two pints ;
Digest for seven days, strain, squeeze the residuum, and filter the liquors. This tincture is best prepared by percolation, which may be commenced so soon as the capsicum is made into a pulp with a little of the spirit.

TINCTURA CARDAMOMI.

Take of the seeds of Cardamoms, bruised,
four ounces and a-half ;

Proof-spirit, two pints ;

Digest for seven days, strain, squeeze the residuum, and filter the liquors. This tincture may be better prepared by the process of percolation, in the same way with the tincture of capsicum, the seeds being first ground in a coffee-mill.

TINCTURA CARDAMOMI COMPOSITA.

Take of Cardamom-seeds, bruised, and
Caraway, bruised, of each two
drachms and-a half ;

Cochineal, bruised, one drachm ;

Cinnamon, bruised, five drachms ;

Raisins, five ounces ;

Proof-spirit, two pints ;

Digest for seven days, strain, express strongly the residuum, and filter the liquors. This tincture may be also prepared by the method of percolation, if the solid materials be first beat together, moistened with a little spirit, and left thus for twelve hours before being put into the percolator.

TINCTURA CASCARILLAE.

Take of Cascarilla, in moderately fine powder,
five ounces ;

Proof-spirit, two pints ;

Proceed by percolation or digestion, as afterwards directed for Tincture of Cinchona.

TINCTURA CASSIAE.

Take of Cassia, in moderately fine powder,
three ounces and a half.

Proof-spirit, two pints ;

Digest for seven days, strain, express the residuum strongly, and filter. This tincture is more conveniently made by the process of percolation, the cassia being allowed to macerate in a little of the spirit for twelve hours before being put into the percolator.

TINCTURA CASTOREI.

Take of Castor, bruised, two ounces and a half ;

Rectified spirit, two pints ;

This tincture may be prepared either by digestion or percolation, like the tincture of cassia.

TINCTURA CASTOREI AMMONIATA.

Take of Castor, bruised, two ounces and a half ;

Assafoetida in small fragments, ten drachms ;

Spirit of Ammonia, two pints ;

Digest for seven days in a well-closed vessel ; strain and express strongly the residuum ; and filter the liquor. This tincture cannot be so conveniently prepared by the method of percolation.

TINCTURA CATECHU.

Take of Catechu, in moderately fine powder,
three ounces and a half ;

Cinnamon, in fine powder, two ounces
and a half ;

Proof-spirit, two pints ;

Digest for seven days ; strain and express
strongly the residuum ; filter the liquors.
This tincture may be also prepared by the
process of percolation, the mixed powders
being put into the percolator without being
previously moistened with the spirit.

TINCTURA CINCHONAE.

Take of Yellow-Bark, in fine powder (or of
any other species of Cinchona, ac-
cording to prescription,) eight
ounces.

Proof-spirit, two pints ;

Percolate the bark with the spirit, the bark
being previously moistened with a very little
spirit, left thus for ten or twelve hours, and
then firmly packed in the cylinder. This
tincture may also be prepared, though much
less expeditiously, and with much greater
loss, by the usual process of digestion, the
bark being in that case reduced to coarse
powder only.

TINCTURA CINCHONAE COMPOSITA.

Take of Yellow-Bark in coarse powder, (fine

if percolation be followed) four ounces ;

Bitter Orange-Peel, bruised, three ounces ;

Serpentaria, in moderately fine powder, six drachms ;

Saffron, chopped, two drachms ;

Cochineal, bruised, one drachm ;

Proof-spirit, two pints ;

Digest for seven days ; strain and express strongly ; filter the liquors. This tincture may also be conveniently prepared by the method of percolation in the same way as the compound tincture of cardamom.

TINCTURA CINNAMOMI.

Take of Cinnamon, in moderately fine powder, three ounces and a-half ;

Proof-spirit, two pints ;

Proceed by percolation or digestion as directed for tincture of cassia.

TINCTURA CINNAMOMI COMPOSITA.

Take of Cinnamon in coarse powder (fine, if percolation be followed), and Cardamom Seeds, bruised, of each an ounce ;

Long Pepper, ground finely, three drachms ;

Proof-spirit, two pints ;

This tincture is best prepared by the method of percolation as directed for the compound

tincture of cardamom. But it may also be made in the ordinary way by digestion for seven days, straining and expressing the liquor, and then filtering it.

TINCTURA COLCHICI.

Take of Colchicum seeds, ground finely in a coffee-mill, five ounces ;

Proof-spirit, two pints.

This tincture is to be prepared like the tincture of cinchona ; and percolation is much more convenient and speedy than digestion.

TINCTURA CONII.

Take of fresh leaves of Conium, twelve ounces ;

Tincture of Cardamom, half a pint ;

Rectified spirit, one pint and a half.

Bruise the hemlock leaves, express the juice strongly ; bruise the residuum, pack it firmly in a percolator ; transmit first the tincture of cardamom, and then the rectified spirit, allowing the spirituous liquors to mix with the expressed juice as they pass through ; add gently water enough to the percolator for pushing through the spirit remaining in the residuum. Filter the liquor after agitation.

TINCTURA CROCI.

Take of Saffron, chopped fine, two ounces ;

Proof-spirit, two pints ;

This tincture is to be prepared like tincture of cinchona, either by percolation or by digestion, the former method being the more convenient and expeditious.

TINCTURA CUSPARIAE.

Take of Cusparia, in moderately fine powder, four ounces and a-half ;

Proof-spirit, two pints ;

This tincture is to be made like the tincture of cinchona, and most expeditiously by the process of percolation.

TINCTURA DIGITALIS.

Take of Digitalis, in moderately fine powder, four ounces ;

Proof-spirit, two pints ;

This tincture is best prepared by the process of percolation, as directed for the tincture of capsicum. If forty fluidounces of spirit be passed through, the density is 944, and the solid contents of a fluidounce amount to twenty-four grains. It may also be made by digestion.

TINCTURA GALLARUM.

Take of Galls, in fine powder, five ounces ;

Proof-spirit, two pints ;

This tincture may be prepared either by digestion or percolation as directed for tincture of capsicum.

TINCTURA GENTIANAE COMPOSITA.

Take of Gentian sliced and bruised, two ounces and a-half.

Dried Bitter-orange peel bruised, ten drachms.

Canella, in moderately fine powder, six drachms ;

Cochineal bruised, half a drachm ;

Proof-spirit, two pints ;

Digest for seven days ; strain and express strongly : and then filter the liquor. This tincture may be more conveniently prepared by percolation, as directed for the compound tincture of cardamom.

TINCTURA GUAIACI.

Take of Guaiac, in coarse powder, seven ounces ;

Rectified spirit, two pints ;

Digest for seven days, and then filter the liquor.

TINCTURA GUAIACI AMMONIATA.

Take of Guaiac, in coarse powder, seven ounces ;

Spirit of ammonia, two pints ;

Digest for seven days in a well-closed vessel, and then filter the liquor.

TINCTURA HYOSCYAMI.

Take of Hyoscyamus, dried and in mode-

moderately fine powder, five ounces ;
Proof-spirit, two pints ;

This tincture is best prepared by the process of percolation as directed for tincture of cap-sicum ; but it may also be obtained, though with greater loss, by the process of digestion.

TINCTURA IODINEI.

Take of Iodine, two ounces and a-half.

Rectified spirit, two pints.

Dissolve the iodine in the spirit with the aid of a gentle heat and agitation ; keep the tincture in well-closed bottles.

TINCTURA JALAPAE.

Take of Jalap, in moderately fine powder,
seven ounces ;

Proof-spirit, two pints ;

This tincture may be prepared either by digestion or percolation, as directed for tincture of cinchona.

TINCTURA KINO.

Take of Kino, in moderately fine powder,
three ounces and a-half.

Rectified spirit, two pints ;

Digest for seven days, and then filter. This tincture cannot be conveniently prepared by the process of percolation.

TINCTURA LACTUCARII.

Take of Lactucarium, in fine powder, four ounces ;

Proof-spirit, two pints :

This tincture is best prepared by percolation as directed for tincture of myrrh ; but may also be prepared by digestion with coarse powder of Lactucarium.

TINCTURA LOBELIAE.

Take of Lobelia, dried and in moderately fine powder, five ounces ;

Proof-spirit, two pints ;

This tincture is best prepared by the process of percolation as directed for tincture of capsicum ; but it may also be made in the usual way by digestion.

TINCTURA LOBELIAE AETHEREA.

Take of dry Lobelia, in moderately fine powder, five ounces ;

Spirit of Sulphuric Ether, two pints ;

This tincture is best prepared by percolation, as directed for tincture of capsicum ; but it may be also obtained by digestion in a well-closed vessel for seven days.

TINCTURA LUPULI.

Take any convenient quantity of hops, recently dried ; separate by friction and sifting the yellowish-brown powder attached to their scales. Then take of this powder five

ounces, and of rectified spirit, two pints ; and prepare the tincture by percolation or digestion as directed for tincture of capsicum.

TINCTURA MYRRHÆ.

Take of Myrrh in moderately fine powder,
three ounces and a-half.

Rectified spirit, two pints ;

Pack the myrrh very gently without any spirit in a percolator ; then pour on the spirit ; and when thirty-three fluidounces have passed through, agitate well to dissolve the oleo-resinous matter which first passes and which lies at the bottom. This tincture is much less conveniently obtained by the process of digestion for seven days.

TINCTURA OPII.

Take of Opium sliced, three ounces ;

Rectified spirit, one pint and seven
fluidounces ;

Water, thirteen fluidounces and a
half ;

Digest* the opium in the water at a temperature near 212° for two hours ; break down the opium with the hand ; strain and express the infusion ; macerate the residuum in the rectified spirit for about twenty hours, and then strain and express very strongly. Mix the watery and spirituous infusions, and filter.

This tincture is not easily obtained by the process of percolation ; but when the opium is of fine quality, it may be prepared thus : Slice the opium finely ; mix the spirit and water ; let the opium macerate in fourteen fluidounces of the mixture for twelve hours, and then break it down thoroughly with the hand ; pour the whole pulpy mass and fluid into a percolator, and let the fluid part pass through, add the rest of the spirit without packing the opium in the cylinder, and continue the process of percolation till two pints are obtained.

TINCTURA OPII AMMONIATA.

Take of Benzoic acid, and
Saffron, chopped, six drachms of
each ;
Opium sliced, half an ounce ;
Oil of Anise, a drachm ;
Spirit of Ammonia, two pints ;
Digest for seven days, and then filter.

TINCTURA OPII CAMPHORATA.

Take of Camphor, two scruples and a half ;
Opium sliced, and
Benzoic acid, four scruples ;
Oil of Anise, one fluidrachm ;
Proof-spirit, two pints ;
Digest for seven days, and then filter.

TINCTURA QUASSIAE.

Take of Quassia, in chips, ten drachms ;
Proof-spirit, two pints ;
Digest for seven days, and then filter.

TINCTURA QUASSIAE COMPOSITA.

Take of Cardamom-seeds bruised, and
Cochineal, bruised, of each half an
ounce ;
Cinnamon in moderately fine powder, and
Quassia, in chips, of each six
drachms ;
Raisins, seven ounces ;
Proof-spirit, two pints ;
Digest for seven days, strain the liquor, express strongly the residuum, and filter This tincture may also be obtained by percolation as directed for Compound tincture of Cardamom, provided the quassia be rasped or in powder.

TINCTURA RHEI.

Take of Rhubarb, in moderately fine powder, three ounces and a half ;
Cardamom Seeds bruised, half an ounce ;
Proof-spirit, two pints ;
Mix the rhubarb and cardamom seeds, and proceed by the process of percolation as directed for tincture of cinchona. This tincture may be also prepared by digestion.

TINCTURA RHEI ET ALOES.

Take of Rhubarb, in moderately fine powder,
an ounce and a half;

Socotorine or East-Indian Aloes,
in moderately fine powder, six
drachms;

Cardamom Seeds bruised, five
drachms;

Proof-spirit, two pints.

Mix the powders, and proceed as for the
tincture of cinchona.

TINCTURA RHEI ET GENTIANAE.

Take of Rhubarb, in moderately fine powder,
two ounces;

Gentian, finely cut or in coarse pow-
der, half an ounce;

Proof-spirit, two pints.

Mix the powders, and proceed as directed
for tincture of cinchona.

TINCTURA SCILLAE.

Take of Squill, in coarse powder, five ounces;
Proof-spirit, two pints.

Prepare this tincture by percolation, as di-
rected for tincture of cinchona, but without
packing the pulp firmly in the percolator. It
may likewise be obtained by digestion from
the sliced bulb.

TINCTURAE SENNA COMPOSITA.

Take of Sugar, two ounces and a half;

Coriander bruised, one ounce ;
Jalap in moderately fine powder, six
drachms ;
Senna, four ounces ;
Caraway bruised, and
Cardamom Seeds bruised, of each
five drachms ;
Raisins bruised, four ounces ;
Proof-spirit, two pints ;

Digest for seven days, strain the liquor, express strongly the residuum, and filter the liquids. This tincture may be more conveniently and expeditiously prepared by percolation, as directed for the compound tincture of cardamom.

If Alexandrian Senna be used for this preparation it must be freed of *Cynanchum* leaves by picking.

TINCTURA SERPENTARIAE.

Take of *Serpentaria* in moderately fine powder, three ounces and a half ;
Cochineal bruised, one drachm ;
Proof-spirit, two pints ;
Proceed by percolation or digestion as for the tincture of cinchona.

TINCTURA TOLUTANA.

Take of Tolu Balsam, in coarse powder, three ounces and a half ;
Rectified-spirit, two pints ;

Digest the balsam in the spirit with a gentle heat till it is dissolved.

TINCTURA VALERIANAE.

Take of Valerian bruised, five ounces ;
Proof-spirit, two pints ;
Proceed by percolation or digestion as for
tincture of cinchona.

TINCTURA VALERIANAE AMMONIATA.

Take of Valerian bruised, five ounces ;
Spirit of Ammonia, two pints.
Proceed by percolation, or by digestion in a
well-closed vessel, as directed for tincture of
cinchona.

TINCTURA ZINGIBERIS.

Take of Ginger, in coarse powder, two ounces and a-half ;
Rectified-spirit, two pints ;
Proceed by percolation or digestion, as directed for tincture of cinchona.

TROCHES.

TROCHISCI ACACIAE.

Take of Gum-arabic, four ounces ;

Starch, one ounce ;

Pure sugar, one pound ;

Mix and pulverize them, and make them into a proper mass with rose-water for forming lozenges.

TROCHISCI ACIDI TARTARICI.

Take of Tartaric acid, two drachms ;

Pure Sugar, eight ounces ;

Volatile oil of Lemons, ten minims.

Pulverize the sugar and acid, add the oil, mix them thoroughly, and with Mucilage beat them into a proper mass for making lozenges.

TROCHISCI CRETAE.

Take of Prepared Chalk, four ounces ;

Gum-arabic, one ounce ;

Nutmeg, one drachm ;

Pure sugar, six ounces ;

Reduce them to powder, and beat them with a little water into a proper mass for making lozenges.

TROCHISCI GLYCIRRHIZAE.

Take of Extract of Liquorice, and

Gum-arabic, of each six ounces ;
Pure Sugar, one pound ;
Dissolve them in a sufficiency of boiling water ; and then concentrate the solution over the vapour-bath to a proper consistence for making lozenges.

TROCHISCI LACTUCARII.

To be prepared with Lactucarium in the same proportion and in the same manner as the Opium Lozenge.

TROCHISCI MAGNESIAE.

Take of Carbonate of Magnesia, six ounces ;
Pure Sugar, three ounces ;
Nutmeg, one scruple ;
Pulverize them, and with mucilage of Tragacanth beat them into a proper mass for making lozenges.

TROCHISCI MORPHIAE.

Take of Muriate of Morphia, one scruple ;
Tincture of Tolu, half an ounce ;
Pure Sugar, twenty-five ounces ;
Dissolve the muriate of morphia in a little hot water ; mix it and the Tincture of Tolu with the sugar ; and with a sufficiency of Mucilage form a proper mass for making lozenges ; each of which should weigh about fifteen grains.

TROCHISCI MORPHIAE ET IPECACUANHAE.

Take of Muriate of Morphia, one scruple ;
Ipecacuan, in fine powder, one
drachm ;
Tincture of Tolu, half a fluidounce ;
Pure Sugar, twenty five ounces ;
Dissolve the muriate in a little hot-water ;
mix it with the tincture and the ipecacuan
and sugar ; and with a sufficiency of Mucilage
beat the whole into a proper mass, which
is to be divided into fifteen-grain lozenges.

TROCHISCI OPII.

Take of Opium, two drachms ;
Tincture of Tolu, half an ounce ;
Pure Sugar, in fine powder, six
ounces ;
Powder of Gum-Arabic, and
Extract of Liquorice, of each five
ounces ;
Reduce the opium to a fluid extract by for-
mula, p. 84 ; mix it intimately with the li-
quorice previously reduced to the consistence
of treacle ; add the tincture ; sprinkle the gum
and sugar into the mixture, and beat it into
a proper mass, which is to be divided into
lozenges of ten grains.

TROCHISCI SODAE BICARBONATIS.

Take of Bicarbonate of Soda, one ounce ;
Pure sugar, three ounces ;
Gum-Arabic, half an ounce ;
Pulverize them, and with mucilage beat them
into a proper mass for making lozenges.

VOLATILE OILS.

VOLATILE oils are obtained chiefly from the flowers, leaves, fruits, barks, and roots of plants, by distilling them with water, in which they have been allowed to macerate for some time. In order to obtain these oils profitably and of good quality, a great variety of conditions must be attended to, differing in regard to each, and such as it would be out of place to enumerate here in detail. Certain general principles, however, may be mentioned.

Flowers, leaves, and fruits generally yield the finest oils, and in greatest quantity, when they are used fresh. Many, however, answer equally well, if they have been preserved by beating them into a pulp with about twice their weight of muriate of soda, and keeping the mixture in well-closed vessels.

Substances yielding volatile oils must be distilled with water, the proper proportion of which varies for each article, and for the several qualities of each. In all instances the quantity must be such as to prevent any of the material from being empyreumatized before the whole oil is carried over. In operations where the material is of pulpy consistence, other contrivances must be resorted to for the same purpose. These chiefly consist of particular modes

of applying heat so as to maintain a regulated temperature not much above 212° . On the small scale heat may be thus conveniently applied by means of a bath of a strong solution of muriate of lime, or by means of an oil-bath, kept at a stationary temperature with the aid of a thermometer. On the large scale heat is often applied by means of steam under regulated pressure. In other operations it is found sufficient to hang the material within the still in a cage or bag of fine net-work; and sometimes the material is not mingled with the water at all, but is subjected to a current of steam passing through it.

The best mode of collecting the oil is by means of the refrigeratory described in the Preface; from which the water and oil drop together into a tall narrow vessel provided with a lateral tube or lip near the top, and another tube rising from the bottom to about a quarter of an inch below the level of the former. It is evident that with a receiver of this construction the water will escape by the lower tube; while the volatile oil, as it accumulates, will be discharged by the upper one, except in the very few instances where the oil is heavier than water.

By attending to the general principles now explained, Volatile oils may be readily obtained of excellent quality from the flowers of

ANTHEMIS NOBILIS,
LAVANDULA VERA, and

RUTA GRAVEOLENS ;

From the fruit of

ANETHUM GRAVEOLENS, bruised,

CARUM CARUI, bruised,

EUGENIA PIMENTO, bruised,

FÆNICULUM OFFICINALE, bruised,

JUNIPERUS COMMUNIS, bruised,

PIPER CUBEBA, ground, and

PIMPINELLA ANISUM, ground ;

From the undeveloped dried flowers of

CARYOPHYLLUS AROMATICUS ;

From the tops of

JUNIPERUS SABINA, and

ROSMARINUS OFFICINALIS ;

From the entire herb of

MENTHA PIPERITA,

MENTHA PULEGIUM,

MENTHA VIRIDIS, and

ORIGANUM MAJORANA ;

And also from the bruised root of

SASSAFRAS OFFICINALE.

OLEUM TEREBINTHINÆ PURIFICATUM.

Take of oil of Turpentine, one pint ;

Water, four pints ;

Distil as long as oil comes over with the
water.

OLEUM COPAIBÆ.

Take of Copaiva, one ounce ;

Water, one pint and a-half ;

Distil, preserving the water; when most of the water has passed over, heat it, return it into the still, and resume the distillation; repeat this process so long as a sensible quantity of oil passes over with the water.

VINEGARS.

ACIDUM ACETICUM CAMPHORATUM.

Take of Camphor, half an ounce ;

Acetic acid, six fluidounces and a half ;

Pulverize the camphor with the aid of a little rectified spirit, and dissolve it in the acid.

ACETUM CANTHARIDIS.

Take of Cantharides, in powder, three ounces ;

Acetic acid, five fluidounces ;

Pyroligneous acid, fifteen fluidounces ;

Euphorbium, in coarse powder, half an ounce ;

Mix the acids, add the powders, macerate for seven days, strain and express strongly, and filter the liquors.

ACETUM COLCHICI.

Take of Colchicum-bulb, fresh and sliced, one ounce ;

Distilled vinegar, sixteen fluidounces ;

Proof-spirit, one fluidounce ;

Macerate the Colchicum in the vinegar for three days in a covered glass vessel ; strain

and express strongly ; filter the liquors, and add the spirit.

ACETUM OPII.

Take of Opium, four ounces ;
Distilled Vinegar, sixteen fluidounces ;

Cut the Opium into small fragments, triturate it into a pulp with a little of the vinegar, add the rest of the vinegar, macerate in a closed vessel for seven days, and agitate occasionally. Then strain and express strongly, and filter the liquor.

ACETUM SCILLAE.

Take of dried Squill, in small fragments,
five ounces ;

Distilled Vinegar, two pints ;

Proof-spirit, three fluidounces ;

Macerate the Squill in the vinegar for seven days in a covered glass vessel, strain and express the liquor, add the spirit, and filter the whole.

WINES.

VINUM ALOES.

Take of Socotorine, or East Indian Aloes,
an ounce and a half ;
Cardamom-seeds, ground,
Ginger, in coarse powder, of each
a drachm and a half ;
Sherry, two pints ;
Digest for seven days, and strain through
linen or calico.

VINUM COLCHICI.

Take of Colchicum-bulb, dried and sliced,
eight ounces ;
Sherry, two pints ;
Digest for seven days, strain, express strong-
ly the residuum, and filter the liquors.

VINUM GENTIANAE.

Take of Gentian, in coarse powder, half an
ounce ;
Yellow-bark, in coarse powder, one
ounce ;
Bitter-orange peel, dried and sliced,
two drachms ;
Canella, in coarse powder, one
drachm ;
Proof-spirit, four fluidounces and
a-half ;

Sherry, one pint and sixteen fluid-ounces ;

Digest the root and barks for twenty-four hours in the spirit ; add the wine, and digest for seven days more ; strain and express the residuum strongly, and filter the liquors.

VINUM IPECACUANHAE.

Take of Ipecacuan, in moderately fine powder, two ounces and a half ;

Sherry, two pints ;

Digest for seven days, and then filter.

VINUM OPII.

Take of Opium, three ounces ;

Cinnamon, in moderately fine powder, and

Cloves, bruised, of each, two drachms and a-half ;

Sherry, two pints ;

Digest for seven days, and then filter.

VINUM RHEI.

Take of Rhubarb, in coarse powder, five ounces ;

Canella, in coarse powder, two drachms ;

Proof-spirit, five fluidounces ;

Sherry, one pint, and fifteen fluid-ounces.

Digest for seven days, strain, express strongly the residuum, and filter the liquors.

VINUM TABACI.

Take of Tobacco, three ounces and a-half,
Sherry, two pints ;

Digest for seven days, strain, express strongly the residuum, and filter the liquors.

TESTS.

AMMONIAE OXALAS.

Take of Oxalic Acid, four ounces ;
Carbonate of Ammonia, eight ounces ;
Distilled Water, four pints ;
Dissolve the carbonate in the water, add gradually the acid, boil and concentrate sufficiently for crystals to form on cooling.

BARYTAE NITRAS.

This salt is to be prepared like the muriate of baryta, p. 98, substituting pure nitric acid for the muriatic acid.

SOLUTIO ARGENTI NITRATIS.

Take of Nitrate of Silver, forty grains ;
Distilled Water, sixteen hundred grains ;
Dissolve the salt in the water, and keep the solution in well-closed bottles.

SOLUTIO ARGENTI AMMONIATI.

Take of Nitrate of Silver, forty-four grains ;
Distilled Water, one fluidounce ;
Aqua Ammoniaë, a sufficiency ;
Dissolve the salt in the water, and add the aqua ammoniaë gradually, and towards the end cautiously, till the precipitate at first

thrown down is very nearly, but not entirely, redissolved.

SOLUTIO BARYTAE NITRATIS.

Take of Nitrate of Baryta, forty grains ;
Distilled Water, eight hundred
grains ;

Dissolve the salt in the water ; and keep
the solution in well-closed bottles.

SOLUTIO SODAE PHOSPHATIS.

Take of Phosphate of Soda (free of efflorescence), one hundred and seventy-five grains ;

Distilled Water, eight fluidounces ;
Dissolve the salt in the water, and keep
the solution in well-closed bottles.

INDEX

OF CHANGES IN NOMENCLATURE.

* * * Many of the changes in Nomenclature adopted in the present edition of the Pharmacopœia are mere Abbreviations, which it is unnecessary to introduce into this Index. The following comprehend all where the departure from the names of the last Latin Edition of the Pharmacopœia is important.

<i>Old Names.</i>	<i>New Names.</i>
Acaciæ Catechu Extract.	Catechu
Acori Calami Radix	Calamus Aromaticus
Adeps Ovillus	Sevum
Adeps Suillus	Axungia
Aether Sulphuricus cum Alcohole	Spirit. Ætheris Sulphur.
Alcohol Ammoniatum	Spiritus Ammoniaë
Alcohol Dilutius	Spiritus Tenuior
Alcohol Fortius	Spiritus Rectificatus
Ammoniaretum Cupri	Cuprum Ammoniatum
Amomi Repentis Semina	Cardamomum
Anethi Fœniculi Semina	Fœniculum
Aqua Supercarb. Potassæ	Aqua Potassæ Effervescens
Aqua Supercarb. Sodæ	Aqua Sodæ Effervescens
Aspidii Filicis Maris Radix	Filix
Astragali Tragacanthæ Gummi	Tragacantha
Bonplandiæ Trifoliatæ Cortex	Cusparia

Bubonis Galbani Gummi- Resina	Galbanum
Carbonas Ferri Precipitat.	Ferri Oxidum Rubrum
Carbonas Potassæ	Potassæ Bicarbonas
Carbonas Sodæ	Sodæ Bicarbonas
Carbonas Zinci Impurus	Calamina
Ceratum Carbon. Zinc. Impurum	Ceratum Calaminæ
Cervi Elaphi Cornu	Cornu
Cinchonæ Cordifoliæ Cortex	Cinchona Flava
Cinchonæ Lancifoliæ Cortex	Cinchona Coronæ
Cinchonæ Oblongifoliæ Cortex	Cinchona Rubra
Citrus Medicæ Fructus	Limones
Colchici Autumnalis Radix	Colchici Cormus
Colombæ Radix	Calumba
Conserva Rosæ Caninæ Gallicæ	Conserva Rosæ Fructus Conserva Rosæ
Convolvulus Scammonii Gummi-resina	Scammonium
Copaiferæ Officinalis Re- sina	Copaiba
Crotonis Eleutheriæ Cortex	Cascarilla
Decoctum Smilacis Sar- saparillæ	Decoctum Sarzæ
Althææ	Mistura Althææ
Hordei Comp.	Mistura Hordei
Dolichi prurientis Pubes	Mucuna
Emplastrum Oxidi Ferri rubri	Emplastrum Ferri
Oxidi Plumbi	Emplastrum Lithargyri semivitrei
Emulsio Acaciæ Arabicæ	Mistura Acaciæ
Amygdali Com- munis	Mistura Amygdalarum

Emulsio Camphoræ	Mistura Camphoræ
Eugeniae Caryophyllatæ Flores	Caryophyllus
Gambogia	Cambogia
Guaiaci Officinalis Resina	Guaiacum
Humuli Lupuli Strobili	Lupulus
Infusum Colombæ	Infusum Calumbæ
Infusum Gentianæ comp.	Infusum Gentianæ
Lichen Islandicus	Cetraria
Melaleucæ Leucodendri Oleum Volat.	Cajuputi Oleum
Menispermi Cocculi Baccæ	Cocculus
Mucilago Acaciæ Arabicæ	Mucilago
Murias Hydrarg. Corros.	Corrosivus Sublimatus
Oleæ Europææ Oleum Fixum	Olivæ Oleum
Oleum Ammoniatum Camphoratum Lini cum Calce	Linimentum Ammoniae Camphoræ Calcis
Oxidum Antimonii cum Phosphate Calcis	Pulvis Antimonialis
Oxidum Arsenici Hydrarg. Rubri per Acid. Nit.	Arsenicum Album Oxidum Hydrargyri Rubri
Oxidum Plumbi Semivit.	Lithargyrum
Papaveris Somnif. Caps.	Papaver
Pilulæ Ammoniareti Cupri Assafœtidæ Comp. Colocynth. Comp. Gambogiæ Comp. Submuriatis Hy- drarg. Comp.	Pilulæ Cupri Ammoniatæ Assafœtidæ Colocynthidis Cambogiæ Calomelanos Comp.
Pini Balsamei Resina Oleum Volatile	Balsamum Canadense Terebinthinæ Oleum
Pistaciæ Lentisci Resin.	Mastiche
Potio Carbonatis Calcis	Mistura Cretæ
Pruni Domesticæ Fructus	Pruna
Pulvis Carb. Calcis Comp. Opiatus	Pulvis Cretæ Compositus Pulvis Cretæ Opiatus

198 INDEX CHANGES IN NOMENCLATURE.

Saccharum non purificat.	Saccharum Commune
purificatum	Saccharum Purum
Smilac. Sarsaparillæ Radix	Sarza
Spartii Scoparii Cacumina	Scoparium
Spermaceti	Cetaceum
Styracis Officinalis Bals.	Styrax
Subacetas Cupri	Ærugo
Sub-boras Sodæ	Borax
Subcarbonas Ammoniaë	Ammoniaë Carbonas
Potassæ	Potassæ Carbonas
Sodæ	Sodæ Carbonas
Submurias Hydrargyri	Calomelas
Sulphur. Antim. Precip.	Antim. Sulphur. Aureum
Hydrarg. Rubrum	Cinnabaris
Supertartras Potassæ	Potassæ Bitartras
Syrupus Citri Medicæ	Syrupus Limonum
Empyreumaticus	Sacchari Fæx
Toluif. Balsami	Syrupus Tolutanus
Tartras Antimonii	Antimonium Tartarizatum
Potassæ et Ferri	Ferrum Tartarizatum
Tinctura Amomi Repentis	Tinctura Cardamomi
Assafoetidæ Am-	Spiritus Ammoniaë Foetidus
moniaata	
Colombæ	Tinctura Calumbæ
Crotonis Eleuth.	Cascarilla
Saponis Camph.	Linimentum Saponis
Saponis et Opii	Opii
Toluiferæ Bals.	Tinctura Tolutana
Toluiferæ Balsami Bals.	Balsamum Tolutanum
Trochisci Carb. Calcis	Trochisci Cretæ
Glycir. cum Op.	Opii
Gummosi	Acaciæ
Unguentum Nit. Hydrarg.	Unguentum Citrinum
Oxidi Zinci	Zinci
Pulveris Can-	Cantharidis
tharidis vesicatorii	
Subacet. Cupri	Aeruginis
Vitis Viniferæ Fructus	Uvæ Passæ

INDEX

MEDICAMENTORUM PREPARATORUM ET COMPOSITORUM.

<p>Acetum Cantharidis 188 Colchici 188 Destillatum 47 Opii . 189 Scillæ 189 Acidum Aceticum 47 Acet. Camph. 188 Benzoicum 48 Citricum 48 Hydrocyanicum 49 Muriat. dilutum 50 Muriat. purum 50 Nitricum dilutum 51 Nitricum purum 51 Sulph. aromaticum . 52 Sulph. dilutum 52 Sulph. purum 51 Tartaricum 52 Aether Sulphuricus 54 Alcohol . 54 Alumen Exsiccatum 95 Ammoniaë Aqua 57 Aqua fortior 57 Acetatis Aqua 58 Carbonas 58 Carbonatis aqua 58</p>	<p>Ammoniaë Oxalas 193 Spiritus 146 Antimonii oxidum 95 Sulphuretum Aureum 96 Antimonium Tartarizatum 96 Aqua Acetatis Ammoniaë 58 Ammoniaë 57 Ammoniaë fortior . 57 Anethi . 75 Calcis . 100 Carbonat. Ammoniaë 58 Cassiaë . 75 Cinnamoni 76 Destillata 75 Fœniculi 76 Lauro-cerasi 76 Menthæ Piperitæ . 76 Menthæ viridis 76 Pimentæ 77 Potassæ . 59 Rosæ . 77 Sambuci 77</p>
--	---

Argenti Ammoniati		Decoctum Mezerei	72
Solutio	193	Papaveris	72
Nitras	97	Quercus	73
Nitratis Solutio	193	Sarzæ	73
Barytæ Murias	98	Sarzæ Compos.	73
Muriatis Solutio	99	Scoparii	73
Nitras	193	Taraxaci	74
Nitratis Solutio	194	Elaterium	81
Bismuthum album	99	Electuarium Aromaticum	68
Calcis Aqua	100	Catechu	68
Murias	100	Opii	69
Muriat. Solutio	101	Piperis	69
Calomelas	108	Sennæ	70
Calx	100	Emplastrum Ammoniaci	143
Carbo animalis purificatus	134	Ammon. cum Hydrargyro	143
Ceratum Calaminæ	126	Assafoetidæ	143
Sabinae	126	Belladonnæ	144
Simplex	126	Cantharidis	144
Chlorinei aqua	134	Canth. compositum	144
Cinnabaris	109	Ferri	145
Conserva Amygdalarum	68	Gummosum	145
Aurantii	68	Hydrargyri	145
Rosæ Fructus	69	Lithargyri	146
Rosæ	69	Opii	146
Creta Preparata	101	Picis	146
Cupri Ammoniati Solutio	102	Resinosum	146
Cuprum Ammoniatum	101	Saponis	147
Decoctum Aloes	71	Simplex	147
Cinchonæ	71	Enema Anodynum	78
Dulcamaræ	71	Catharticum	78
Guaiaci	72	Fœtidum	78
Hæmatoxyli	72	Opii	78
		Terebinthinæ	78

PREPARATORUM ET COMPOSITORUM. 201

Enema Tabaci .	78	Ferri Oxidum Rubrum	105
Extractum Aconiti	79	Sulphas	105
Anthemidis	79	Sulphas Exsic-	
Belladonnæ	80	catum	105
Cinchonæ	80	Sulphuretum	106
Colchici Ace-		Ferrugo .	106
ticum .	80	Ferrum Tartarizatum	107
Colocynthis	81		
Conii .	81	Hydrargyri Biniodi-	
Digitalis	81	dum .	108
Elaterii .	81	cum Creta	109
Gentianæ	82	Oxidum Rub-	
Glycirrhizæ	82	rum .	109
Hæmatoxyli	82	Precipitatum	
Hyoscyami	82	Album	110
Jalapæ .	83		
Kramerizæ	83	Infusum Anthemidis	89
Lupuli .	83	Aurantii	89
Nucis-vomicæ	83	Bucku .	89
Opii .	84	Calumbæ	89
Papaveris	84	Caryophylli	90
Pareiræ	84	Cascarillæ	90
Quassizæ	85	Catechu .	90
Rhei .	85	Chirettæ	90
Scammonii	85	Cinchonæ	91
Sarzæ Fluidum	86	Cusparizæ	91
Styracis .	86	Digitalis .	91
Stramonii	86	Gentianæ	91
Taraxaci	87	Lini .	92
		Pareiræ .	92
Ferri Carbonas Sac-		Quassizæ	92
charatum	102	Rhei .	92
Iodidi Syrupus	102	Rosæ .	93
Iodidum	103	Senegæ	93
Muriatis Tinc-		Sennæ .	93
tura .	104	Sennæ Compo-	
Oxidum Nig-		situm	93
rum .	104	Serpentariæ	94

Infusum Simarubæ	94	Oleum purificat.	186
Iodineum .	135	Ol. Volatile Anisi	186
Iodinei liquor compo- situs .	135	Anethi	186
Linimentum Ammoniaë	126	Anthemidis	185
Ammoniaë Com- positum	127	Carui .	186
Calcis .	127	Caryophylli	186
Camphoræ	127	Copaibæ	186
Opii .	127	Cubebæ	186
Saponis	128	Fœniculi	186
Simplex	128	Juniperi	186
Terebinthina- tum .	128	Lavandulæ	185
Magnesiaë .	110	Menthæ Pipe- ritæ .	186
Magnesiaë Carbonas	111	Menth. Virid.	186
Mel Boracis .	88	Origani	186
Mel Rosæ .	88	Pimentæ	185
Mistura Acaciaë	121	Pulegii .	186
Amygdalarum	121	Rosmarini	186
Althææ	122	Rutæ .	186
Camphoræ	122	Sabinaë	186
Camphoræ cum Magnesia	122	Sassafras	186
Creasoti	123	Pilulæ Aloes .	136
Cretæ .	123	Aloes et Assa- fœtidæ	136
Ferri composita	123	Aloes et Ferri	136
Guaiaci	124	Aloes et Myrrh.	137
Hordei .	124	Assafœtidæ	137
Scammonii	124	Cambogiaë	137
Morphiaë Acetas	61	Calomelanos compositæ	137
Murias	61	Calomelanos et Opii	138
Muriatis Solutio	63	Colocynthidis	138
Mucilago .	125	Colocynthidis et Hyoscyami	138
Amyli .	125	Cupri Ammon.	139
Tragacanthæ	125	Digitalis et Scillaë .	139.
Oleum Terebinthinæ			

Pilulæ Ferri Carbonat.	139	Pulvis Cretæ Opiatus	148
Ferri Sulphatis	139	Ipecac. Comp.	149
Hydrargyri	140	Jalapæ Comp.	149
Ipecacuanhæ et		Rhei Comp.	149
Opii .	140	Salinus Comp.	149
Opii .	140	Scammonii	
Plumbi Opiatæ	141	Compositus	150
Rhei .	141	Tragacanthæ	
Rhei Composit.	141	Compositus	150
Rhei et Ferri	142		
Scillæ .	142	Quinæ Sulphas .	63
Styracis .	142		
Thebaicæ .	140	Resina Jalapæ .	83
Plumbi Acetas	111	Scammonii	85
Diacetatis so-			
lutio .	111	Sodæ Aqua Efferves.	117
Iodidum .	112	Bicarbonas	117
Nitras .	112	Carbonas Sic-	
Potassa .	59	catum .	118
cum Calce .	60	Murias Purum	118
Potassæ Acetas .	112	Phosphas	118
Aqua .	59	Phosphatis so-	
Aqua Efferves.	113	lutio .	194
Bicarbonas	113	Sulphas .	119
Bisulphas .	114	Solutio Arsenicalis	98
Carb. Purum	114	Spiritus Ætheris nitrici	55
et Sodæ Tartras	116	Ætheris Sul-	
Sulphas	114	phurici .	55
Sulphas cum		Ammoniaë .	151
Sulphure	115	Ammon. Aro-	
Tartras .	115	maticus .	151
Potassii Iodidum	116	Ammon. Fœ-	
Sulphuretum	117	tidus .	152
Pulveres effervescentes	148	Carui .	152
Pulvis Aluminis Com-		Cassiaë .	152
positus	148	Cinnamomi	152
Antimonialis	95	Juniperi Com-	
Aromaticus	148	positus .	152
Cretæ Comp.	148	Lavandulæ	153

Spiritus Lavand. Com-		Tinctura Calumbæ	164
positus	153	Camphoræ	165
Menthæ .	153	Cantharidis	165
Myristicæ	153	Capsici .	165
Pimentæ .	154	Cardamomi	165
Rosmarini	154	Cardamomi	
Tenuior .	154	composita	166
Stanni Pulvis .	120	Cascarillæ	166
Strychnia .	65	Cassiae .	167
Sublimatus Corrosi-		Castorei .	167
vus . . .	110	Castorei Am-	
Sulphur Sublimatum	135	moniata	167
Syrupus Aceti .	155	Catechu .	168
Althææ .	155	Cinchonæ	168
Aurantii .	155	Cinch. Comp.	168
Croci .	155	Cinnamomi	169
Ipecacuanhæ	156	Cinnam. Com-	
Limonum .	156	posita .	169
Papaveris .	157	Colchici .	170
Rhamni .	157	Conii .	170
Rhæados	157	Croci .	170
RosæCentifoliæ	158	Cuspariæ	171
Rosæ Gallicæ	158	Digitalis .	171
Sarzæ .	158	Ferri Muriatis	104
Scillæ .	159	Gallarum .	171
Sennæ .	159	Gentianæ com-	
Simplex .	159	posita .	172
Tolutanus	160	Guaiaci .	172
Violæ .	160	Guaiaci Am-	
Zingiberis	160	moniata	172
		Hyoseyami	172
Tinctura Aloes .	162	Iodinei .	173
Aloes et Myr-		Jalapæ .	173
rhæ .	163	Kino .	173
Assafoetidæ	163	Lactucarii	173
Aurantii .	163	Lobeliæ .	174
Benzoini comp.	164	Lobeliæ Æth.	
Bucku .	164	erea .	174

Tinctura Lupuli .	174	Unguentum Antimon.	129
Myrrhæ .	175	Cantharidis	129
Opii .	175	Infusi Cantha-	
Opii Ammo-		ridis .	129
niata .	176	Citrinum .	130
Opii Campho-		Cocculi .	130
rata .	176	Creasoti .	130
Quassiæ .	177	Gallæ et Opii	131
Quassiæ Comp.	177	Hydrargyri	131
Rhei .	177	Iodinei .	131
Rhei et Aloes	178	Oxydi Hydrar-	
Rhei et Gen-		gyri .	132
tianæ .	178	Picis Liquidæ	132
Scillæ .	178	Plumbi Ace-	
Sennæ Comp.	178	tatis .	132
Serpentariæ	179	Plumbi Car-	
Tolutana .	179	bonatis	132
Valerianæ	180	Precipitati Albi	132
Valerianæ Am-		Resinosum	133
moniata	180	Simplex .	133
Zingiberis	180	Sulphuris	133
Trochisci Acaciæ	181	Zinci .	133
Acidi Tartarici	181	Veratria .	66
Cretæ .	181	Vinum Aloes .	190
Glycirrhizæ	181	Antimoniale .	97
Lactucarii	182	Colchici .	190
Magnesiæ .	182	Gentianæ .	190
Morphiæ	182	Ipecacuanhæ	191
Morph. et Ipe-		Opii .	191
cacuanhæ	183	Rhei . .	191
Opii .	183	Tabaci .	192
Sodæ Bicarb.	183	Zinci Oxidum .	120
Unguentum Aerugi-		Sulphas .	120
nis .	128		

ENGLISH INDEX

OF THE

ARTICLES OF THE MATERIA MEDICA AND
THEIR PREPARATIONS AND COMPOUNDS.

Acid, Acetic	2, 47	Almond Conserve	68
Arom. Sulph.	52	Mixture	121
Benzoic	3, 48	Aloes, Barbados	5
Camphorated		East Indian	6
Acetic	188	Socotorine	6
Citric	3, 48	Decoction	71
Hydrocyanic	3, 49	Pill	136
Muriatic	3	and Assafoetida	
Dil. Muriatic	50	pill	136
Pure Muriatic	3, 50	and Iron pill	136
Nitric	4	and Myrrh pill	137
Dilut. Nitric	51	Tincture of	162
Pure Nitric	4, 51	Tincture of	
Sulphuric	4	and Myrrh,	163
Diluted Sul-		Wine of	190
phuric	52	Alum	6
Pure Sulp.	5, 51	Dried	95
Pyroligneous	4	Compound	
Tartaric	5, 52	powder of	148
Tartaric Lo-		Ammonia, Carbonate	
zenges	181	of	6, 58
Alcohol	5, 54	Solution of	
Almonds, Bitter	7	Carbonate of	58
Sweet	7	Liniment of	126

- | | |
|-------------------------------------|------------------------------------|
| Ammonia, Compound | Assafœtida, Ammoniated |
| Liniment of 127 | spirit of . 152 |
| Muriate of 6 | Pill . 137 |
| Oxalate of 193 | Plaster . 143 |
| Solution of 8, 57 | Tincture of 163 |
| Stronger Solution of 9, 57 | Axunge . 10 |
| Solution of | Balm . . . 27 |
| Acetate of 9, 58 | Balsam, Canada . 10 |
| Spirit of . 6, 151 | Peru . 10 |
| Aromatic Spirit of . 151 | Tolu . 11 |
| Ammoniac . 6 | Bark. <i>See</i> Cinchona |
| Plaster . 143 | Barley . . . 23 |
| and Mercury plaster . 143 | Mixture . 124 |
| Ammoniated Spirit of | Baryta, carbonate of 10 |
| Assafœtida 152 | Muriate of 11, 98 |
| Angelica . 7 | Solution of |
| Angustura-bark. <i>See</i> Cusparia | Muriate of 99 |
| Anise, . . . 7 | Nitrate of . 193 |
| Oil of . 186 | Nitrate, solution of . 194 |
| Anodyne Enema . 78 | Sulphate of . 11 |
| Antimonial Ointment 129 | Bear-berry . 43 |
| Powder . 34, 95 | Bees'-wax . 15 |
| Wine . 97 | Bleached . 15 |
| Antimony, Oxide of 7, 95 | Belladonna . 11 |
| Sulphuret of . 7 | Extract of . 80 |
| Golden Sulphuret of 7, 96 | Plaster . 144 |
| Tartar emetic 8, 96 | Benzoin . . . 11 |
| Aromatic electuary 68 | Comp.tinct. of 164 |
| powder . 148 | Bergamot, oil of . 11 |
| Arrow-root . 27 | Bismuth . 11 |
| Arsenic . . . 10 | White . 11, 99 |
| Arsenical solution . 98 | Bitter-sweet. <i>See</i> Dulcamara |
| Assafœtida . 10 | Borax . . . 12 |
| | Honey of . 88 |
| | Broom-tops . 33 |

Broom-tops, Decoction	73	Caraway, Spirit of	152
Buckbean	27	Cardamom	14
Buckthorn	35	Tincture of	165
Syrup	157	Compound	
Bucku	12	tincture of	166
Infusion of	89	Carolina-pink	40
Tincture of	164	Carrot	19
Burgundy pitch	30	Cascarilla	14
Cajuput-oil	12	Infusion of	90
Calamine	12	Tincture of	166
Cerate	126	Cassia-bark	15
Calomel	12, 108	Cassia, Oil of	15
Pill, comp.	137	Pulp	15
and Opium		Spirit of	152
pill	138	Tincture of	167
Calumba	12	Water of	75
Infusion of	89	Castor	15
Tincture of	164	Tincture of	167
Camphor	13	Ammoniated	
Liniment	127	tincture of	167
Mixture	122	Castor-Oil	36
and Magnesia		Catechu	15
Mixture	122	Electuary of	68
Tincture of	165	Infusion of	90
Car. ella	14	Tincture of	168
Cantharides	14	Cathartic enema	78
Ointment	129	Cayenne-pepper. <i>See</i>	
Infusion, oint-		Capsicum	
ment of	129	Centaury	15
Plaster	144	Cerates	126
Plaster comp.	144	Cerate Calamine	126
Tincture of	165	Savin	126
Vinegar	188	Simple	126
Capsicum	14	Cevadilla	36
Tincture of	165	Chalk	18
Caraway	14	Lozenge	181
Oil of	186	Mixture	123
		Powder comp.	148

- | | | | |
|----------------------------|-----------|----------------------------|-----------|
| Chalk Powder with
opium | 148 | Cloves Infusion of | 90 |
| Prepared | 18, 101 | Oil of | . 14, 186 |
| Chamomile | . 7 | Clysters. <i>See</i> Enema | |
| Extract of | . 79 | Cocculus-indicus | . 17 |
| Infusion | . 89 | Ointment | . 130 |
| Oil of | . 185 | Cochineal | . 17 |
| Charcoal | . 14 | Colchicum-bulb | . 17 |
| Animal | . 14 | Extract of | . 80 |
| Purified ani-
mal | . 14, 134 | Seeds | . 17 |
| Cherry-laurel | . 25 | Tincture of | 170 |
| Laurel-water | 76 | Vinegar | . 188 |
| Chiretta | . 15 | Wine | . 190 |
| Infusion of | 90 | Colocynth | . 17 |
| Chlorine-water | 16, 134 | Extract of | . 80 |
| Cinchona-bark | . 16 | Pill | . 138 |
| Crown-bark | 16 | and Hyoscyamus | |
| Gray-bark | . 16 | pill | . 138 |
| Silver-bark | 16 | Conium. <i>See</i> Hemlock | |
| Red-bark | . 16 | Conserve of Almonds | 68 |
| Yellow-bark | 16 | of Hips | . 69 |
| Decoction of | 71 | of Orange | . 68 |
| Extract of | 80 | of Red Roses | 69 |
| Infusion of | 91 | Copaiva | . 17 |
| Tincture of | 168 | Oil of | 18, 186 |
| Compound | | Copper, Ammoniat-
ed | . 19, 101 |
| tincture of | 168 | Solution of | |
| Cinnabar | . 109 | ammoniated | 102 |
| Cinnamon | . 16 | Pill of ammo-
niated | . 139 |
| Oil of | . 17 | Sulphate of | 19 |
| Spirit of | . 152 | Coriander | . 18 |
| Tincture of | 169 | Corn-poppy | . 35 |
| Compound | | Syrup | . 157 |
| tincture of | 169 | Corrosive-sublimate | 41, 110 |
| Water of | . 76 | Cotton, raw | . 22 |
| Citrine ointment | . 130 | Cowitch | . 28 |
| Cloves | . 14 | Creasote | . 18 |

- | | | | |
|--------------------------------|-----|--------------------------------|--------|
| Creasote Mixture . . . | 123 | Elaterium . . . | 19, 81 |
| Ointment . . . | 130 | Elder-flowers . . . | 37 |
| Croton-oil . . . | 18 | Elder-flower-water . . . | 77 |
| Cubebs . . . | 18 | Electuaries . . . | 68 |
| Oil of . . . | 186 | Electuary, aromatic . . . | 68 |
| Cumin . . . | 18 | of Catechu . . . | 68 |
| Cusparia . . . | 19 | of Opium . . . | 69 |
| Infusion of . . . | 91 | of Pepper . . . | 69 |
| Tincture of . . . | 171 | of Senna . . . | 70 |
| Dandelion . . . | 42 | Elemi . . . | 19 |
| Decoction of . . . | 74 | Emulsions. <i>See</i> Mixtures | |
| Extract of . . . | 87 | Enemas . . . | 78 |
| Deadly-nightshade. <i>See</i> | | Enema, Anodyne . . . | 78 |
| Belladonna | | Cathartic . . . | 78 |
| Decoction of Aloes . . . | 71 | Fetid . . . | 78 |
| of Broom-tops . . . | 73 | Opiate . . . | 78 |
| of Cinchona . . . | 71 | Tobacco . . . | 78 |
| of Dandelion . . . | 74 | Turpentine . . . | 78 |
| of Dulcamara . . . | 71 | Ergot . . . | 19 |
| of Guaiac . . . | 72 | Ether, Sulphuric . . . | 5, 54 |
| of Logwood . . . | 72 | Spirit of Sul- | |
| of Mezereon . . . | 72 | phuric . . . | 40, 55 |
| of Oakbark . . . | 73 | Spirit of Ni- | |
| of Poppy-heads . . . | 72 | tric . . . | 40, 55 |
| of Sarza . . . | 73 | Euphorbium . . . | 20 |
| of Sarza, comp. . . | 73 | Extracts, preparation | |
| Digitalis. <i>See</i> Foxglove | | of . . . | 79 |
| Dill . . . | 7 | Extract of Belladonna . . . | 80 |
| Oil of . . . | 186 | of Chamomile . . . | 79 |
| Water . . . | 75 | of Cinchona . . . | 80 |
| Distilled water . . . | 75 | of Colchicum . . . | 80 |
| Distilled waters . . . | 75 | of Colocynth . . . | 81 |
| Dulcamara . . . | 19 | of Dandelion . . . | 87 |
| Decoction of . . . | 71 | of Elaterium . . . | 81 |
| Effervescing powders . . . | 148 | of Foxglove . . . | 81 |
| Egg . . . | 29 | of Gentian . . . | 82 |
| | | of Hemlock . . . | 81 |
| | | of Hops . . . | 83 |

Extract of Hyoscyamus	82	Garlick	5
of Jalap	83	Gentian	22
of Liquorice	82	Extract of	82
of Logwood	82	Compound Tinc-	
of Monkshood	79	ture of	172
of Nux-vomica	83	Infusion of	91
of Opium	84	Wine	190
of Pareira	84	Ginger	44
of Poppy-heads	84	Syrup of	160
of Quassia	85	Tincture of	180
of Rhatany	83	Guaiac	22
of Rhubarb	85	Decoction of	72
of Sarza, Fluid	86	Mixture	124
of Scammony	85	Tincture of	172
of Storax	86	Ammoniated	
of Thornapple	86	Tincture of	172
Fennel	21	Gum-arabic	22
Oil of	186	Mixture	121
Water of	76	Mucilage of	125
Fern, male shield	21	Lozenge	181
Fetid Enema	78	Gum-Plaster	145
Figs	21	Hartshorn. <i>See</i> Am-	
Flour	20	monia	
Foxglove	19	Hellebore, Black	23
Extract of	81	White	43
Infusion of	91	Hemlock	17
and Squill pill	139	Extract of	81
Tincture of	171	Tincture of	170
Galbanum	22	Henbane. <i>See</i> Hyoscy-	
Galls	22	amus	
and Opium Oint-		Hips	36
ment	131	Conserve of	69
Tincture of,	171	Honey	27
Gamboge, Ceylon	13	of Borax	88
Siam	13	of Roses	88
Pill	137	Hops	26
		Extract of	83

Hops, Tincture of	174	Ipecacuan. Powder,	
Horn	18	Compound	149
Horse-raddish	10	Syrup	156
Hyoscyamus	23	Wine	191
Extract of	82	Iron, Black Oxide	
Tincture of	172	of	20, 104
Iceland-moss	15	Filings	20
Infusion of Bucku	89	Iodide of	20, 103
Calumba	89	Syrup of iodide	
Cascarilla	90	of	20, 102
Catechu	90	Mixture, Com-	
Chamomile	89	pound	123
Chiretta	90	Pill of Carbo-	
Cinchona	91	nate of	139
Cloves	90	Plaster	145
Cusparia	91	Red Oxide	
Foxglove	91	of	21, 105
Gentian	91	Rust of	21, 106
Linseed	92	Saccharine Car-	
Orange Peel	89	bonate	20, 102
Pareira	92	Sulphate of	21, 105
Quassia	92	Dried sulphate	
Rhubarb	92	of	105
Roses	93	Pill of sulphate	
Senega	93	of	139
Senna	93	Sulphuret of	
Senna, Comp.	93	21, 106	
Serpentaria	94	Tartrate of	21, 107
Simaruba	94	Tincture of mu-	
Iodine	23	riate of,	20, 104
Compound solu-		Wire	20
tion of,	135	Jalap	24
Ointment	131	Extract of	83
Purification of	135	Powder, Com-	
Tincture of	173	pound,	149
Ipecacuan	24	Tincture of	173
and Opium pill	140	Juniper-berries	24

- | | | | |
|--|---------|-----------------------------|---------|
| Juniper, Oil of | 24, 186 | Lime, Liniment | 127 |
| Tops | 24 | Muriate of | 12, 100 |
| Compound Spi-
rit of | 152 | Solution of mu-
riate of | 101 |
| Kali-water | 32, 113 | Water | 100 |
| Kino | 24 | Limes | 25 |
| Tincture of | 173 | Liniment of Am-
monia | 126 |
| Lactucarium | 24 | of Ammonia,
compound | 127 |
| Lozenges | 182 | of Camphor | 127 |
| Tincture of | 173 | of Lime | 127 |
| Lavender | 25 | of Opium | 127 |
| Oil of | 25, 185 | Simple | 128 |
| Spirit of | 153 | Soap | 128 |
| Compound spi-
rit of | 153 | Turpentine | 128 |
| Lead, Acetate of | 30, 111 | Linseed | 25 |
| Carbonate of | 30 | Infusion | 92 |
| Iodide of | 31, 112 | Meal | 25 |
| Nitrate of | 31, 112 | Oil | 25 |
| and Opium Pill | 141 | Liquorice-root | 22 |
| Ointment of
acetate of | 132 | Extract of | 22, 82 |
| Ointment of
carbonate of | 132 | Lozenge | 181 |
| Red Oxide of | 31 | Litharge | 25 |
| Solution of di-
acetate of | 30, 111 | Plaster | 146 |
| Lemon | 25 | Litmus | 24 |
| Juice, Syrup of | 156 | Lobelia | 26 |
| Oil of | 25 | Ethereal tinc-
ture of | 174 |
| Peel | 25 | Tincture of | 174 |
| Lettuce-opium. <i>See</i>
Lactucarium | | Logwood | 23 |
| Lignum-vitæ | 22 | Decoction | 72 |
| Lime | 13, 100 | Extract of | 82 |
| Chloride of | 13 | Lozenges, or Troches | 181 |
| | | Chalk | 181 |
| | | Gum-Arabic | 181 |
| | | Lactucarium | 182 |
| | | Liquorice | 181 |

- | | | | |
|-----------------------------|---------|----------------------------|---------|
| Lozenges, Magnesia | 182 | Mercury, White precipitate | 23, 110 |
| Morphia | 182 | Mezereon | 28 |
| Morphia and Ipecacuan | 183 | Decoction | 72 |
| Opium | 183 | Mixtures | 121 |
| Soda | 183 | Gum-Arabic | 121 |
| Tartaric acid | 181 | Almond | 121 |
| Magnesia | 26, 110 | Barley | 124 |
| Carbonate of | 26, 111 | Camphor | 122 |
| Lozenge | 182 | Camphor and Magnesia | 122 |
| Sulphate of | 26 | Chalk | 123 |
| Mallow, Common | 26 | Compound Iron | 123 |
| Marsh. <i>See</i> | | Creasote | 123 |
| Marsh-mallow | | Guaiaac | 124 |
| Manganese | 26 | Marsh-mallow | 122 |
| Manna | 27 | Scammony | 124 |
| Marble | 27 | Monkshood | 5 |
| Marjoram | 29 | Extract of | 79 |
| Oil of | 186 | Morphia, acetate of | 28, 61 |
| Marsh-Mallow Leaves | 6 | Lozenge | 182 |
| Root | 6 | and Ipecacuan | |
| Mixture | 122 | Lozenge | 183 |
| Syrup | 155 | Muriate of | 28, 61 |
| Mastic | 27 | Solution of | 63 |
| Mercury | 23 | Mucilage | 125 |
| Biniodide of | 23, 108 | of Starch | 125 |
| and Chalk | 109 | of Tragacanth | 125 |
| Ointment of | 131 | Muscovado | 36 |
| Ointment of oxide of | 132 | Musk | 28 |
| Ointment of white precipit. | 132 | Mustard | 38 |
| Pill of | 140 | Myrrh | 29 |
| Plaster of | 145 | Tincture of | 175 |
| Red oxide of | 23, 109 | Neroli-oil | 10 |
| Sulphuret of | 109 | Nutmeg | 28 |
| | | Concrete oil of | 28 |

Nutmeg Spirit of	153	Opium Camphorated	
Volatile oil of	28	tincture of	176
Nux-vomica .	29	Electuary	69
Extract of	83	Enema .	78
Oak-bark .	35	Extract of	84
Decoction .	73	Liniment	127
Oats .	10	Lozenge	183
Oils. <i>See</i> Volatile Oils		Pill .	140
Ointments .	126	Plaster	146
Ointment of Acetate of		Tincture of	175
Lead	132	Vinegar	189
Antimonial	129	Wine .	191
of Carb. of lead	132	Orange-Flower Water	10
of Cantharides	129	Oil .	10
of Infusion of		Orange-Peel .	10
Cantharides	129	Conserve of	68
Citrine	130	Infusion of	89
Cocculus-indi-		Syrup of	155
cus .	130	Tincture of	163
Creasote	130	Pareira .	30
of Galls and		Extract of	84
Opium	131	Infusion of	92
Iodine .	131	Pellitory of Spain	35
Mercurial	131	Penny-royal .	34
of Oxide of		Oil .	186
Mercury	132	Water .	77
Resinous .	133	Pepper, Black	30
Simple .	133	Electuary of	69
Sulphur .	133	Long .	30
Tar .	132	Peppermint .	27
Verdigris	128	Oil	27, 186
White Precipi-		Spirit .	153
tate .	132	Water .	76
Zinc .	133	Petroleum .	30
Olive-oil .	29	Pills .	136
Opium .	29	Aloetic	136
Ammoniated		Aloes and As-	
tincture of	176	safœtida	136

Pills, Aloes and Iron	136	Plaster, Ammoniac	143
Aloes and Myrrh	137	Ammoniac and Mercury	143
Ammoniated Copper	139	Assafoetida	143
Assafoetida	137	Belladonna	144
Calomel and Opium	138	Cantharides	144
Carb. of Iron	139	Compound Cantharides	144
Colocynth	138	Gum	145
Colocynth and Hyoscyamus	138	of Iron	145
Compound Calomel	137	Litharge	146
Compound Rhubarb	141	Mercurial	145
Digitalis and Squill	139	Opium	146
Gamboge	137	Pitch	146
Ipecacuan and Opium	140	Resin	146
Lead and Opium	141	Simple	147
Mercurial	140	Soap	147
Opiate	140	Pomegranate-root-bark	22
Rhubarb	141	Poppy-Heads	29
Rhubarb and Iron	142	Decoction of	72
Squill	142	Extract of	84
Storax	142	Syrup of	157
Sulph. of Iron	139	Potash	31, 59
Pimento	30	Acetate of	31, 112
Oil	186	Bicarbonate of	32, 113
Spirit	154	Bisulphate of	32, 114
Water	77	Bitartrate of	32
Pitch	30	Carbonate of	32
Plaster	146	Carbonate of, Pure	33, 114
Plasters	143	and Lime	31, 60
		Ferrocyanide of	34
		Iodide of	34, 116
		Nitrate of	33
		Solution of	9, 59

- | | | | |
|-----------------------|---------|-------------------|---------|
| Potash, Sulphate of | 33, 114 | Rectified Spirit | 40 |
| Sulphate of, with | | Red-Sandal wood | 34 |
| Sulphur | 33, 115 | Resin | 35 |
| Sulphuret | | of Jalap | 83 |
| of | 34, 117 | of Scammony | 85 |
| Tartrate of | 33, 115 | Resinous Ointment | 133 |
| Tartrate of So- | | Plaster | 146 |
| da and | 33, 116 | Rhatany-root | 24 |
| Powders | 148 | Extract of | 83 |
| Powder, Antimonial | 95 | Rhubarb | 35 |
| Aromatic | 148 | Extract of | 85 |
| Chalk and | | Infusion of | 92 |
| Opium | 148 | and Iron Pills | 142 |
| Comp. Alum | 148 | Pill | 141 |
| Comp. Chalk | 148 | Pill, compound | 141 |
| Compound | | Powder, comp. | 149 |
| Ipecacuan | 149 | Tincture of | 177 |
| Comp, Jalap | 149 | and Aloes, Tinc- | |
| Comp. Rhubarb | 149 | ture of | 177 |
| Comp. Saline | 149 | and Gentian, | |
| Comp. Scam- | | Tincture of | 177 |
| mony | 150 | Wine | 191 |
| Comp. Traga- | | Rose, Damask | 36 |
| canth | 150 | Damask, Syrup | |
| Powders, Effervescing | 148 | of | 158 |
| Proof-spirit | 40, 154 | Oil or Attar of | 36 |
| Prunes | 34 | Red | 36 |
| Pyrola | 35 | Red, Conserve of | 69 |
| Purging-flax | 25 | Red, Honey of | 88 |
| | | Red, Infusion of | 93 |
| Quassia | 35 | Red, Syrup of | 158 |
| Extract of | 85 | Water | 77 |
| Infusion of | 92 | Rosemary | 36 |
| Tincture of | 177 | Oil | 186 |
| Compound | | Spirit of | 154 |
| Tincture of | 177 | Rue | 36 |
| Quina, Sulphate of | 35, 63 | Oil of | 36, 186 |
| Raisins, | 43 | Saffron, | 18 |


Saffron, Syrup of	155	Silver	9
Tincture of	170	Ammoniated, So-	
Sago	37	lution of	193
Saline powder, com-		Nitrate of	9, 93
pound	149	Solution of Ni-	97
Sarza	37	trate of	1
Compound De-		Simaruba-root	93
coction	73	Infusion of	38
Decoction of	73	Simple Cerate	126
Fluid Extract	86	Liniment	128
Syrup of	158	Ointment	133
Sassafras	37	Plaster	147
Oil	186	Syrup	159
Savin	36	Snake-root	38
Cerate	126	Infusion of	93
Oil	186	Soap, Castile	37
Scammony	37	Liniment	128
Compound Pow-		Plaster	147
der	150	Soft	37
Extract of	85	Soda, Bicarbonate	
Mixture	124	of	39, 117
Resin of	85	Carbonate of	39
Senega	38	Dried Carbo-	
Infusion of	93	nate of	118
Senna, Alexandrian	38	Lozenge	183
East Indian	38	Muriate of	39
Electuary	70	Pure Muriate	
Infusion of	93	of	39, 118
Compound In-		Phosphate of	39, 118
fusion of	93	Solution of	
Compound		Phosph. of	194
Tincture of	178	Sulphate of	40, 119
Syrup of	159	Water	39, 117
Tinnivelly	38	Spearmint	27
Serpentaria	38	Water of	77
Infusion of	94	Oil of	186
Tincture of	179	Spermaceti	15
Sherry	43	Spirits	151

Spirit of Ammonia	151	Suet	38
of Ammonia,		Sugar, White	37
Aromatic	151	Sulphur	42
of Assafoetida		Ointment	133
Ammoniat.	152	Sublimed	135
of Caraway	152	Sweet Flag	12
of Cassia	152	Syrups	155
of Cinnamon	152	Syrup of Buckthorn	157
of Juniper,		of Corn-poppy	157
Compound	152	of Damask-rose	158
of Lavender	153	of Ginger	160
of Lavender,		of Ipecacuan	156
Compound	153	of Lemon-juice	156
of Nitric		of Marsh-mal-	
Ether	40, 55	low	155
of Nutmeg	153	of Orange	155
of Peppermint	153	of Poppy-head	157
of Pimento	154	of Red Rose	158
Proof	40, 154	of Saffron	155
Rectified	40	of Sarza	158
of Rosemary	154	of Senna	159
of Sulphuric		Simple	159
Ether	40, 55	of Squill	159
Sponge	40	of Tolu	160
Squill	38	of Vinegar	155
Pill	142	of Violets	160
Syrup of	159	Tamarind Pulp	42
Tincture of	178	Tapioca	42
Vinegar	189	Tar	30
Starch	7	Ointment	132
Mucilage of	125	Tartar Emetic	8, 96
Stavesacre	41	Emetic Oint.	129
Storax	41	Tests	193
Extract of	86	Thebaic Pill	140
Pill	142	Thorn-Apple	41
Stramonium	41	Extract of	86
Extract of	86	Tin	41
Strychnia	41, 65		

Tin Powder .	120	Tincture of Hemlock	170
Tinctures, Preparation		Hops .	174
of .	161	Hyoscyamus	172
Tincture of Aloes	162	Iodine .	173
Aloes and		Jalap .	173
Myrrh	163	Kino .	173
Assafœtida	163	Lactucarium	173
Benzoin;		Lobelia .	174
Compound	164	Lobelia, Ethe-	
Bucku	164	real .	174
Calumba	164	Muriate of Iron	104
Camphor	165	Myrrh	175
Cantharides	165	Opium .	175
Capsicum	165	Opium, Am-	
Cardamom	165	moniated	176
Card. Comp.	166	Opium, Cam-	
Cascarilla	166	phorated	176
Cassia .	167	Orange Peel	163
Castor .	167	Quassia,	177
Castor, Am-		Quassia, Com-	
moniated	167	pound	177
Catechu .	168	Rhubarb .	177
Cinchona .	168	Rhubarb and	
Cinchona,		Aloes	178
Compound	168	Rhubarb and	
Cinnamon	169	Gentian	178
Cinnamon,		Saffron .	170
Compound	169	Senna, Com-	
Colchicum	170	pound	178
Cusparia	171	Serpentaria	179
Foxglove	171	Squill .	178
Galls .	171	Tolu .	179
Gentian, Com-		Valerian	180
pound	172	Valerian, Am-	
Ginger .	180	moniated	180
Guaiaic	172	Tobacco .	42
Guaiaic Am-		Enema	78
moniated	17	Wine .	192

Tolu-balsam	11	Volatile Oils, Extrac-	
Syrup of	160	tion of	184
Tincture of	179	Oil of Anise	186
Tormentil	42	Oil of Caraway	186
Tous-les-mois	44	Oil of Chamo-	
Tragacanth	43	mile	185
Mucilage of	125	Oil of Cloves	186
Compound pow-		Oil of Copaiva	186
der of	150	Oil of Cubebs	186
Treacle	36	Oil of Dill	186
Troches. <i>See</i> Lozenges		Oil of Fennel	186
Turmeric	19	Oil of Juniper	186
Turpentine, Chian	42	Oil of Lavender	185
Enema	78	Oil of Marjo-	
Liniment	128	ram	186
Oil	42	Oil of Penny-	
Oil, Purified	186	royal	186
Venice	42	Oil of Pepper-	
Valerian	43	mint	186
Tincture of	180	Oil of Pimento	186
Ammoniated		Oil of Rose-	
Tincture of	180	mary	186
Veratria	66	Oil of Rue	186
Verdigris	5	Oil of Sassafras	186
Ointment	128	Oil of Savin	186
Vinegar, British	2	Oil of Spear-	
Cantharides	188	mint	186
Colchicum	188	Oil of Turpen-	
Distilled	2, 47	tine	42, 186
French	2	Water	8
Opium	189	Cassia	75
Squill	189	Cherry-laurel	76
Syrup of	155	Cinnamon	76
Violet	43	Dill	75
Syrup of	160	Distilled	8, 75
Virginian Snake-root	38	Elder-flower	77
Infusion of	94	Fennel	76

Water, Pennyroyal	77	Wine, Antimonial	97
Peppermint	76	Colchicum	190
Pimento	77	Gentian	190
Rose	77	Ipecacuan	191
Spearmint	76	Opium	191
White-Hellebore	43	Rhubarb	191
White Precipitate	23, 110	Tobacco	192
Precipitate Ointment	132	Zinc	43
Willow-bark	37	Ointment	133
Wormwood	2	Oxide of	43, 120
Wine, Aloetic	190	Sulphate of	43, 120



4

5 -

6 -

7 -

8

9 -

10 -

11 -

12 -

Soz 1847

