

Chemical recreations: a series of amusing ... experiments ... To which are prefixed, first lines of chemistry, etc / [Anon].

Contributors

Griffin, John Joseph, 1802-1877

Publication/Creation

Glasgow : R. Griffin, 1824.

Persistent URL

<https://wellcomecollection.org/works/beb2nmh8>


License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.




Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>




25, 744/A


GRIFIN







CHEMICAL RECREATIONS *Pl. 1.*



272
CHEMICAL RECREATIONS:

A SERIES OF

AMUSING AND INSTRUCTIVE
EXPERIMENTS,

WHICH MAY BE PERFORMED
EASILY, SAFELY, AND AT LITTLE EXPENSE.

TO WHICH ARE PREFIXED,

FIRST LINES OF CHEMISTRY;

WHEREIN

*The principal facts of the Science, as stated by the most
Celebrated Experimentalists, are familiarly explained.*

WITH A

MINUTE DESCRIPTION

OF A

CHEAP AND SIMPLE APPARATUS;

ILLUSTRATED BY

SEVENTY ENGRAVED FIGURES

OF THE DIFFERENT PARTS OF IT.

THIRD EDITION, CORRECTED AND ENLARGED.

PRINTED FOR

RICHARD GRIFFIN & CO. GLASGOW;

E. WEST & CO. EDINBURGH; AND THOMAS TEGG, LONDON.

MDCCCXXIV.

CHEMICAL REPERATIONS:

THE FARMINGTON HOSPITAL

BY J. H. WELLS

AS PUBLISHED BY THE

EAST LITTLE EXPENSE

TO THE

FIRST LITTLE CHEMISTRY:



The following is a description of the

and is intended to be used as a

with a

MINUTE DESCRIPTION

OF A

OF THE

OF THE

OF THE

OF THE

OF THE

OF THE

OF THE

OF THE

OF THE

OF THE

OF THE

OF THE

TO THE MEMBERS
OF
THE MECHANICS' CLASS
OF THE
ANDERSONIAN INSTITUTION,
GLASGOW.

GENTLEMEN,

The following work, written by one of your number, was undertaken for the purpose of furnishing to junior students of Chemistry, a *text-book* at a moderate price.

The contents of the work, are, an introductory view of Chemistry, and a collection of interesting experiments, with such directions for the performance of them, as, I hope, will render the understanding of the processes perfectly easy. For the materials of the work, I must (like the author of every elementary work) acknowledge myself chiefly indebted to the writings of others; and more particularly to those of the eminent chemists of the present day. In the arrangement of the different subjects, I have studied simplicity; and in

language, conciseness. My readers, I trust, will neither find themselves puzzled by confusion in theory, nor wearied by circumlocutory details.

To none can I dedicate this work with more propriety than to yourselves. When I first sat on the same benches with you, and felt (what many young members of the class still feel) the want of a book of this nature, and the impossibility of obtaining it, I determined to compile such a one, as soon as my attainments in science rendered me capable of so doing. Since then, some time has elapsed, and now I have endeavoured to perform what I had projected. The result of this undertaking, I beg leave to present to you, and hope may be found not unworthy of your patronage.

I am, GENTLEMEN,

Your obedient Servant,

J. G.

GLASGOW, *September 18th*, 1823.

ADVERTISEMENT
TO
THE THIRD EDITION.

SINCE this work was last presented to the public, it has undergone a thorough revision, and now makes its appearance in a greatly improved state. Many hundred corrections have been made; the language has been rendered more explicit; the details more minute. The Index also has been enlarged; and a considerable Glossary has been incorporated in it.

The Public will please to consider my endeavours to *improve* the book, as my best mode of returning thanks for the favourable reception it met with at its first appearance.

Now that Mechanics' Institutions have been established in many parts of the kingdom, it is gratifying to me to be able to add, that this book has been recommended, by very competent judges, as being "peculiarly adapted for the use of the members of those Institutions." It was expressly for the use of Mechanics that the book was written: I have therefore obtained the object that I had in view.

London, May 31st, 1824.

ADVERTISEMENT

THE THIRD EDITION.


Since this work was last presented to the public, it has undergone a thorough revision, and now makes its appearance in a greatly improved state. Many hundred corrections have been made; the language has been rendered more explicit; the details more minute. The index also has been enlarged; and a considerable Glossary has been incorporated in it.


The Public will please to consider my endeavours to improve the book, as my last mode of retaining thanks for the favourable reception it met with at its first appearance.


Now that Mechanics' Institutions have been established in many parts of the Kingdom, it is gratifying to me to be able to add, that this book has been recommended, by very competent judges, as being "peculiarly adapted for the use of the members of those Institutions." It was expressly for the use of Mechanics that the book was written: I have therefore obtained the object that I had in view.


London, May 31st, 1834.

CHEMICAL RECREATIONS *Pl 2*













CHEMICAL RECREATIONS *Pl. 4.*











CHEMICAL RECREATIONS *Pl. 6.*



R. Gray Sc.



FIRST LINES OF CHEMISTRY;

IN WHICH

THE PRINCIPAL FACTS OF THE SCIENCE, AS STATED
BY THE MOST CELEBRATED EXPERIMENTALISTS,
ARE FAMILIARLY EXPLAINED.

Introduction.

1. "WHAT IS CHEMISTRY?—IS IT INTERESTING?—
IS IT USEFUL? HOW AM I TO PROCEED IN STUDYING IT?"

2. Chemistry is the science which makes known to us the nature and properties of all natural bodies, whether they be simple or compound—solid, liquid, or aëriform: a science which, as its objects are inexhaustible, and infinitely varied, furnishes us with continual entertainment; and which is essentially useful in many of the arts upon which depend the comforts, and even the very existence, of civilized life.

3. The great importance of the science of Chemistry is rendered evident by the following considerations.—It is useful in explaining natural phenomena. In acquiring a knowledge of the constitution of the atmosphere, in investigating the changes to which it is subject, the variations of temperature, the laws of winds, dew, rain, hail, and snow, chemistry is our principal, our only satisfactory guide. These remarkable changes—changes which, because familiar, do not produce any emotion in the mind, though in themselves truly wonderful—are chemical operations on a magnificent scale, and can only be explained by chemical laws.

4. In man's researches into the nature of the things whence he derives the means of his comfort, his happiness, his luxuries, and even his existence—in examining the various objects which compose the mineral, the vegetable and the animal kingdoms, Chemistry is essentially requisite for the successful progress of his inquiries.

5. In considering the application of Chemistry to the improvement of the arts of civilized life, a wide field of contemplation opens to our view. So extensive, indeed, are its influence and importance, that, in most of the arts, many of the processes, in some all, that are employed depend on chemical principles. The bare mention of some of these arts will suggest ample illustrations of its extensive utility.

6. In the medical art, of so great service is a knowledge of Chemistry, that it is now universally regarded as an essential branch of a medical education. In the art of extracting metals from their ores, in purifying and combining them with each other, and in forming instruments and metals—whether for useful or ornamental purposes—almost all the processes are purely chemical. The art of glass and porcelain making—of tanning, soapmaking, dyeing, and bleaching—depend entirely upon Chemistry, and all the processes in baking, brewing, and distilling, most of the culinary arts, and many others in domestic economy, are chemical operations. In short, wherever in any of the processes of nature or of art, the addition or the abstraction of heat takes place—wherever substances in combination are to be separated—wherever the union of simple substances and the formation of new compounds are effected;—the operations and their results can only be explained on chemical principles.

7. From this general view of the extensive applications of chemical science, those who have not considered the objects which it embraces will be enabled to judge of the importance of this study.

8. If we consider Chemistry purely as a science, we shall find no study better calculated to encourage that generous love of truth which confers dignity and superiority on those who successfully pursue it. No science

holds out more interesting subjects of research, and none affords more striking proofs of the wisdom and beneficence of the Creator of the universe. A machine constructed by human art, is admired in proportion to the simplicity of its contrivance, the extent of its usefulness, and the niceness of its adaptations. But the works of man sink into nothing when brought into comparison with the works of nature. When examining the former, every step of our progress is obscured with comparative clumsiness and defect: in contemplating the latter, we behold perfection rise on perfection, and more exquisite wonders still meeting our view. It is the merit of Chemistry, that by its aid we are enabled to take a minuter survey of the great system of the universe. And, so far as our limited powers can comprehend it, the whole is nicely balanced and adjusted, and all its changes tend to the most beneficial purposes. Circumstances which, on a superficial view, were seeming imperfections and defects, a closer inspection points out to be real excellencies. In all the singular and surprising changes which everywhere present themselves, the more closely we observe and examine them, the more we shall admire the simple means by which they are accomplished, and the intelligent design and perfect wisdom displayed in the beneficial ends to which they are directed.

9. Chemistry is a science that is founded entirely upon experiment; and no person can understand it unless he performs such experiments as verify its fundamental truths. The hearing of lectures, and the reading of books, will never benefit him who attends to nothing else; for Chemistry can only be studied to advantage *practically*. One *experiment*, well-conducted, and carefully observed by the student, from first to last, will afford more knowledge than the mere perusal of a whole volume. It may be added to this, that chemical operations are, in general, the most interesting that could possibly be devised—Reader! what more is requisite to induce you to

MAKE EXPERIMENTS?

10. It has been found, that all the marvellous diversity of appearance under which bodies are presented to the eye, and the changes of state to which they are incessantly sub-

jected, are occasioned by the reciprocal actions and combinations of a few unchangeable primary bodies. The properties of these bodies, and the nature of the laws which regulate their actions, are, therefore, the objects of which the chemical student is to enter into an investigation.

11. There are two methods by which chemical knowledge is acquired, *analysis* and *synthesis*;—the former signify the *separating* of the constituent parts of a compound body—the latter, the *formation* of a compound body, by the *putting-together* of its component principles. Both *analysis* and *synthesis* are effected by means of certain processes or operations, thence called *chemical operations*.

12. It is evident from what has been said that the whole subjects of Chemistry are resolved into—*First*, the art of performing the necessary chemical operations, (that is, *of making experiments*,)—and, *Secondly*, the obtaining, through experiments, a knowledge of the nature and properties of all natural bodies, simple and compound; so as to be enabled to apply these bodies to useful or ornamental purposes.

SIMPLE BODIES.

13. Formerly, fire, water, air, and earth, were regarded as—and as the only—*simple bodies*: that is to say, as the elementary bodies, or *different sorts of matter*, of which all other known bodies were composed. But water, air, and earth, have been now proved to be compound bodies, (the nature of fire is still unknown,) and there are many substances in nature which have no place at all in the list. This system, thus shown to be very erroneous, has been long given up.

14. When we say that a body is *simple*, it is to be understood, that it has not yet been *decomposed*—that is to say, has not been resolved into other different substances. It is probable that we are not yet acquainted with any one of the *elements* of matter; but yet, as long as the bodies with which we are acquainted continue *undecomposed*, they are to be regarded as *simple*, or *elementary* bodies. It is surely needless to give any account of what is meant by a *compound* body, for the term is self-explanatory.

15. The metal *gold* affords a very good idea of a *simple*

body ; for, though that substance may be melted by heat, or dissolved in a corrosive menstruum, yet it is recovered unchanged in its properties ; but, on the other hand, *marble* is, by a strong heat, converted into two other bodies—one of which is *lime*, the other, an *elastic fluid* (or *air*) that is disengaged during the process. *Marble*, therefore, is a *compound* body.

16. The number of hitherto-undecompounded bodies is *fifty-three*. Four others—light, heat, electricity, and magnetism, called the *imponderable* bodies—have, by some, been added to these ; but, as their separate identity has not been clearly ascertained, they are not generally reckoned with the others. As, however, they produce, by their various and important actions, very remarkable effects in other substances, we shall hereafter take such notice of them as may be deemed necessary. The whole of these *fifty-three* bodies may be *weighed* and *measured*, and hence (in contradistinction to the four bodies just mentioned, which cannot be weighed and measured) are called *ponderable* bodies. These, in order to facilitate the acquirement of a knowledge of their properties, have been arranged as shown in the following

17. TABLE OF SIMPLE BODIES.

CLASS FIRST.

Bodies having an immense affinity for the simple bodies of the succeeding two classes ; with which bodies they combine, and thereby form substances that are totally different in their properties from the substances of which they are composed :—

- | | | |
|-------------|--|--------------|
| 1. OXYGEN | | 3. IODINE |
| 2. CHLORINE | | 4. FLUORINE. |

CLASS SECOND.

Bodies of a non-metallic nature, but inflammable or acidifiable :—

- | | | |
|-----------------|---|------------------------------------|
| 5. HYDROGEN | } | <i>Gaseous Bodies</i> |
| 6. NITROGEN | | |
| 7. CARBON | } | <i>Fixed and Infusible Solids</i> |
| 8. BORON | | |
| 9. SULPHUR | } | <i>Fusible and Volatile Solids</i> |
| 10. PHOSPHORUS. | | |

CLASS THIRD,

Inflammable substances of a metallic nature. This is the most numerous class of simple bodies; the individual of which it is composed being in number *forty-three*—as shown in the Table of Metals, paragraph 29. These substances combine with nearly all the ten bodies named above; but the most important compounds into which they enter, are the bodies formed by their combination with oxygen.

18.—1. OXYGEN.—This is one of the most important agents in nature. Scarcely a process of any description takes place, in which it has not a share. In a simple state it is obtained only in the form of *gas*. It is an exceedingly abundant body; the air of the atmosphere contains one-fifth of its bulk of it; and water, 75 parts in every 100. It also exists in most natural products, animal, vegetable, and mineral. *Oxygen gas* is, like common air, colourless, invisible, tasteless, inodorous, and elastic. But it is heavier than common air, in the proportion $11\frac{1}{2}$ to 10. It is slightly soluble in water. It is a powerful supporter of combustion; that is to say, when any inflammable body, as a lighted taper, is put into it, it burns very vigorously—much more so than if it were put into common air; indeed, it is owing to the oxygen it contains, that common air supports combustion at all, as will be shown hereafter. Its presence is also essential for the continuance of animal life. We cannot breathe air which has been deprived of its oxygen; and it must be noticed, that an animal lives, and a combustible body burns, much longer in a definite quantity of oxygen gas, than it would in the same quantity of atmospherical air. Hence it is evident, that oxygen is the principle which supports both life and fire. Oxygen is not only found combined in natural bodies, but it can be made, by means of art, to combine with a great variety of substances, with which it forms very peculiar compounds—see 97, 133. The method of obtaining oxygen gas, and of making experiments with it, is described at 408 to 427.

NOTE.—Properly speaking, *oxygen gas* is not a simple body: since the gaseous state is not the natural state of

oxygen, but is owing to the presence of a peculiar chemical agent, which will be presently described, called *caloric*. But as we know of no substances that are *separated* from caloric, it is customary to apply the term simple to such as are combined with caloric only. *Gas* is the name given to all permanently-elastic fluids, both simple and compound; except the atmosphere, to which the term *air* is appropriated. It is necessary to distinguish between *gas* and *vapour*. The latter is elastic and fluid, but not permanently so. The vapour of water (steam) upon cooling becomes a liquid: it is, therefore, not a *gas*, for *gases* are bodies whose aëriiform state is permanent. The methods of obtaining and exhibiting the properties of oxygen, and other gases, is minutely described in the experimental part of this work.

19.—II. CHLORINE is a *gas*, possessing the mechanical properties of common air. Its colour is greenish-yellow (which its name literally signifies). Its taste is very disagreeable, and its smell exceedingly strong and suffocating. It would kill the person who presumed to breathe it; it is dangerous, even when largely diluted with common air. Though not respirable, it is an eminent supporter of combustion; some bodies indeed inflame in it spontaneously. It mixes very readily and largely with water, and then acquires the property (for it has it not in its dry gaseous state) of destroying vegetable colours. This has rendered it useful in some *bleaching* operations. The weight of chlorine gas is to that of common air, as 5 to 2. The compounds of chlorine are generally *chlorides*; it forms, however, a few *acids*. For the method of obtaining chlorine gas, and of examining its properties, see 472 to 476.

20.—III. IODINE is a solid body which has the colour and lustre of plumbago. It is capable of crystallization. It is slightly soluble in water. When exposed to heat about that of boiling water, it combines with caloric, and is converted into a beautiful violet-coloured vapour, nearly nine times heavier than common air. Iodine has an acrid taste, and is strongly poisonous. In its smell and its action on vegetable colours it resembles chlorine; but its effects are less violent than those of that substance. It is

incombustible; but, in combining with several bodies, the intensity of mutual action is such, as to produce the phenomena of combustion. It is obtained from seaweed, but it can be only got in small quantities, and its preparation requires considerable skill. *Acids* and *iodides* are the products of its combinations.

21.—IV. FLUORINE.—So strong are the tendencies of this substance to combination, that as yet no vessel has been found, that is capable of containing it in its pure state. When united to hydrogen, it produces an intensely-acid compound, called fluoric acid.

22.—V. HYDROGEN is only known in the state of *gas*, and is sometimes called *inflammable air*. It is the lightest species of ponderable matter with which we are acquainted; compared to oxygen, its density is as 1 is to 15. It is the basis of *water*, from which body only it can be procured. Hydrogen gas, when pure, is possessed of all the physical properties of common air; a slight odour, which it sometimes has, is produced by some substance that is held in solution by it. It does not support combustion, though it is itself one of the most combustible of all bodies; being that which gives the power of burning with flame to all the substances used for the economical production of heat and light. But it only burns in the presence of oxygen. It is not fit for respiration; for animals which breathe it, die almost instantaneously. If pure oxygen and hydrogen gas be mixed together, they remain unaltered; but if a lighted taper be brought into contact with the mixture, it explodes with astonishing violence; and, if the two gaseous bodies have been mixed in certain proportions, the whole is condensed into water: hence we see the origin of the term *hydrogen*, which literally signifies *the water-former*. Hydrogen gas is the substance which, on account of its rarity, is employed to inflate air-balloons. The method of obtaining and making experiments with hydrogen gas is described at 428 to 441.

23.—VI. NITROGEN, called also *azote* (or *life-destroyer*), is a gaseous body, rather lighter than common air; of which it forms 4-5th parts in bulk, the remaining 1-5th being oxygen. It is tasteless, inodorous, colourless, and

capable of being condensed and dilated. It extinguishes flame, and is fatal to animal life. It combines with oxygen in various proportions, forming compounds which differ greatly in their properties. In one proportion it forms the air of the atmosphere: in other proportions, the bodies described at 106, 107, 134, 135. With hydrogen, it forms the alkaline body ammonia (121); and with chlorine and iodine, two very formidable detonating compounds.

24.—VII. CARBON is the name given to the pure inflammable part of *charcoal*, of which substance, the diamond is only a variety in a pure crystallized state; for pure charcoal and diamond, when treated in the same manner, produce precisely the same results. Carbon is insoluble in water, and infusible by the most intense heat. Carbon has a powerful affinity for oxygen, and is thence useful in several operations, to be hereafter described: it is employed with advantage to purify various substances. Carbon combines with oxygen, and produces a gas called carbonic acid; and, when combined with hydrogen gas, forms carburetted hydrogen gas—the same that is now used to light up shops. Animal and vegetable oils, are composed almost entirely of carbon and hydrogen; the difference in their properties resulting chiefly from the variation in the proportions of these two bodies. The same may be observed of gum, sugar, and starch. All these bodies, however, contain oxygen.

25.—VIII. BORON is a solid of a dark olive colour, procured from boracic acid, which is one of the constituents of a substance called borax. It is infusible; but when heated takes fire, and burns with a red light, and brilliant scintillations. It was lately discovered, and is difficultly prepared.

26.—IX. SULPHUR, a well-known substance, distinguished commonly by the name of brimstone. It is a hard brittle body, of a yellow colour, destitute of smell, and of a weak taste. It is universally diffused in nature; but commonly combined with other bodies. It is insoluble in water, but if poured into that liquid when liquified by heat, it retains its softness; and in this state is employed

for taking impressions from seals and medals. When exposed to heat in close vessels, it is sublimed or volatilized in the form of a very fine powder, called, very absurdly, *flowers of sulphur*. At a heat of about twice that of boiling water, it takes fire, if in contact with the air, and burns with a flame of a pale blue colour. [In this process it dissolves in the oxygen of the atmosphere, and produces an elastic fluid acid. It is a substance of great importance in chemistry and the arts. Oxygen unites with it in four proportions, its compounds forming an interesting series of acids, see 100, 101, 102. The compounds of sulphur with metals are called sulphurets. With hydrogen it forms sulphuretted hydrogen gas.

27.—X. PHOSPHORUS is a semi-transparent yellow matter, of the consistence of wax. It is procured, in general, by the decomposition of bones. It is so inflammable, that it is set on fire by a heat of about one-third the of boiling water. Indeed, it has a luminous appearance arising from a slow combustion, at the common temperature of the atmosphere. During its combustion, it emits a dense white smoke, which has the smell of garlic, and the dark is luminous. When heated to five times the height necessary to inflame it (air being excluded) it boils. On account of its very combustible nature, it requires to be handled with great caution. It is a violent poison. Alcohol, oils, and ether, dissolve it in small quantities forming solutions, which are possessed of some curious properties, as is shown among the *experiments*. Phosphorus, when burnt in oxygen, forms, by combining with it in different proportions, a series of acids.

ON METALS.

28. The forty-three elementary substances which now remain to be described, are all METALS. They compose the most numerous class of undecomposed chemical bodies, and are distinguished by the following general characters:—

I. They possess a peculiar lustre.—II. They are opaque.—III. They are fusible by heat; and in fusion retain their lustre and opacity.—IV. They are excellent

conductors of electricity and heat.—V. Many of them may be extended under the hammer, and are called malleable ; or under the rolling press, and are called laminable ; or drawn into wire, and are called ductile.—VI. When exposed, highly-heated, to the action of oxygen, chlorine, or iodine, they take fire, and are converted by the combustion into oxides, chlorides, or iodides ; bodies destitute of lustre, and other metallic characteristics.—VII. They will combine, in almost any proportion, with each other, when in a state of fusion, and thus form compounds, which are termed alloys, bodies that *retain* the properties of metals.—VIII. From their brilliancy and opacity, conjointly, they reflect the greater part of the light which falls on their surface ; hence they form excellent mirrors.—IX. When combined with simple bodies of the second class, (11,) they produce bodies of very peculiar characters, some of them being gaseous, others semi-metallic, &c.—X. Many of them may by peculiar management be crystallized.—XI. They are very heavy : to this character, however, (though it was till lately considered one of their most prominent features,) there are important exceptions ; since *metals* have been obtained, (potassium and sodium, for instance,) which are lighter than water.

The relative weights, or specific gravities, of the different metals, are noted in the general table of specific gravities, at the end of the book.

The acquirement, retention, and application of knowledge, is much facilitated by a methodical distribution of its different parts. The relations of the metals, however, to the various objects of chemistry, are so complex and diversified, as to render their classification a task of peculiar difficulty. In the following “ Table,” they are arranged according to their perfectability. Those at the head of the list are scarcely at all affected by the power of oxygen. But as we progressively descend, the influence of that beautiful element progressively increases ; so that among those near the bottom, it exercises an almost despotic sway, which the magical pile of Volta, directed by the genius of Davy, can only suspend for a season. The

emancipated metal soon relapses under the dominion of oxygen.

In an annexed column, is described the nature of the products formed by their union with oxygen. These combinations will be duly considered hereafter.

29. GENERAL TABLE OF THE METALS.

<i>Names.</i>	<i>United to Oxygen, form</i>	<i>Names.</i>	<i>United to Oxygen, form</i>		
1. Platinum, ~~~~~	Ordinary Metallic Oxides, or Neutral Salifiable Bases.	24. Osmium, ~~~~~	Metallic Oxides, not much known.		
2. Gold, ~~~~~		25. Rhodium, ~~~~~			
3. Silver, ~~~~~		26. Iridium, ~~~~~			
4. Paladium, ~~~~~		27. Uranium, ~~~~~			
5. Mercury, ~~~~~		28. Titanium, ~~~~~			
6. Copper, ~~~~~		29. Cerium, ~~~~~			
7. Iron, ~~~~~		30. Wodanium, ~~~~~			
8. Tin, ~~~~~		31. Potassium, ~~~~~ 32. Sodium, ~~~~~ 33. Lithium, ~~~~~	Alkalies.		
9. Lead, ~~~~~					
10. Nickel, ~~~~~					
11. Cadmium, ~~~~~		34. Calcium, ~~~~~ 35. Barium, ~~~~~ 36. Strontium, ~~~~~ 37. Magnesium, ~~~~~ 38. Yttrium, ~~~~~ 39. Glucinum, ~~~~~ 40. Aluminum, ~~~~~ 41. Thorinum, ~~~~~ 42. Zirconium, ~~~~~ 43. Silicium, ~~~~~	Earths.		
12. Zinc, ~~~~~					
13. Bismuth, ~~~~~					
14. Antimony, ~~~~~					
15. Manganese, ~~~~~					
16. Cobalt, ~~~~~					
17. Tellurium, ~~~~~	Acids.				
18. Arsenic, ~~~~~					
19. Chromium, ~~~~~					
20. Molybdenum, ~~~~~					
21. Tungstenum, ~~~~~					
22. Columbium, ~~~~~					
23. Selenium, ~~~~~					

NOTE.—Two of the classes of bodies which the above Table points out as *compounds of metals with oxygen*, the *alkalies*, and the *earths*, were, till very lately, supposed to be *simple substances*; and were, consequently, in every system of chemistry, arranged with the other simple bodies: as, however, it has been fully proved that they are *not* such, they are now, of course, treated of with the compounds. But their great importance entitles them to claim and obtains for them, a very conspicuous place in every work on chemistry: we shall shortly treat of them; in the mean time, we proceed to give a sketch of the individual characteristics of the metals.

30.—I. PLATINUM, a metal of a dull white colour, the heaviest body in nature. It requires an intense heat for its fusion; and has so little affinity for oxygen, that it may be exposed for any length of time to the atmosphere, without being oxidised (rusted or tarnished): hence it would make very useful utensils, could it be obtained in large quantities. But it is scarce. Two red hot pieces of it may, by hammering, be joined into one. This operation is termed *welding*, and is peculiar to itself and iron. It is extremely malleable and ductile—may be hammered into thin plates, and drawn into wire $\frac{1}{3000}$ ths of an inch in diameter. It is unacted upon by all acids, except such as contain chlorine.

31.—II. GOLD is a beautiful yellow metal, and, next to platinum, the heaviest of all known bodies. It bears the same relation to acid *menstrua* that platinum does; requiring those containing chlorine to dissolve it. Of all metals, gold is the most ductile and malleable; it is also soft, very tough, and unalterable even by the strongest heat of a furnace. It does not oxidise when exposed to the air. Most metals unite to gold by fusion. Mercury is very strongly disposed to do so without fusion. Lead makes it brittle. Copper deepens its colour, and, by rendering it harder and more fusible, makes it more fit for coin and other articles used. The gold coins of Great Britain contain 11 parts of gold and 1 of copper.

32.—III. SILVER. This is the whitest of all metals, and in brilliancy only inferior to polished steel. It is softer than copper, but harder than gold. It is very ductile and malleable, though less so than gold: it may, however, be beat into leaves $\frac{1}{10000}$ ths of an inch thick, and a single grain of it has been drawn into a wire 400 feet long. It is also very tenacious: a fine wire will support a very great weight without breaking. Silver, by means of an intense heat, that produced by a galvanic battery, for instance, may, like any other combustible body, be burnt. Its inflammation is attended by a green light of great beauty and brilliancy. Silver is oxidised by ignition in an open vessel, but not by exposure to the atmosphere. It is rapidly tarnished by the vapour of sulphur. To ren-

der it harder, and more fitting for the purposes of coin than it is in its pure state, it is alloyed with copper.

33.—IV. PALLADIUM, a metal that is very scarce. Its colour is grayish-white. It is ductile, malleable, and harder than iron. It takes a fine polish.

34.—V. MERCURY, a heavy metal, of a blueish-white colour. It is also called quicksilver. Its chief characteristic is that of always being fluid at the common temperature of the atmosphere. But it becomes solid, when exposed to a sufficient degree of cold. The temperature necessary for freezing it, is 39° below 0° of Fahrenheit's thermometer. At 600° it boils, and volatilizes, so that it may be distilled from one vessel into another. By this process it is purified. It is strongly inclined to unite with metals; its metallic compounds are termed *amalgams*. Looking-glasses are silvered by an *amalgam* of tin. Among the preparations of mercury, are the bodies usually termed vermilion, calomel, and corrosive sublimate; the last is a most violent poison, but is, as well as the one preceding it, frequently employed in medicine. Mercury is easily dissolved in nitric acid.

35.—VI. COPPER is a reddish-brown metal; hard and sonorous; very malleable and ductile; and of great tenacity. It is one of the most useful and most abundant metals. It is about nine times heavier than water. It is melted by a heat seven times greater than that of boiling water. By a still greater heat, it evaporates in visible fumes. It is easily oxidised by exposure to the air. In making a plate of copper red hot, and plunging it in this state into cold water, oxide, in small scales, will separate from the metal; any quantity of oxide may be thus obtained. When copper is exposed to a very violent heat it burns with a vivid green-coloured flame. Copper combines with most other metals. With zinc, it forms the very useful compounds, called *brass*, *pinchbeck*, *princemetal*, &c. With tin it forms bell-metal, bronze, the alloys used for cannons, mirrors of telescopes, &c. Some of the compounds formed by the combinations of copper with acids are very useful.—See 203 to 206.

36.—VII. IRON is the most useful, and most abundant

dant, of all the metals we are acquainted with. Its colour is blueish white. It has the properties of hardness, tenacity, and ductility, in a very eminent degree. It is malleable also, but less so than silver. It requires an intense degree of heat for its fusion. One of its most advantageous properties, is that of *welding*. By undergoing this process, (peculiar to itself and platinum,) two pieces of this metal may be joined into one. Susceptibility of magnetism, in a very high degree, is also one of its distinguishing properties. The *loadstone* itself (in which the power of magnetism chiefly resides) is an iron ore. Iron is widely diffused: it is found in animals, vegetables, and minerals. It has a strong affinity for oxygen. When exposed to the air, its surface gradually becomes covered with a brown or red powder, commonly called *rust*. This powder is an oxide of iron. Carbon and iron united, form the exceedingly useful compound, called *steel*. Iron unites with other metals. The most useful of its alloys, is that with tin, called *tin-plate*. Iron enters into the composition of the beautiful pigment, called *prussian-blue*. Green vitriol, or copperas, is iron combined with sulphuric acid.

37.—VIII. TIN is a yellowish-white metal, possessed of great brilliancy, harder than lead, and very malleable. It may be beat into leaves, $\frac{1}{2000}$ ths of an inch thick. But it is neither very tenacious, nor very ductile. It is very flexible. It melts, at a low heat; by a great heat, it may be evaporated. Tin soon tarnishes when exposed to air, becoming slightly oxidated; and, when it is melted in an open vessel, its surface is soon covered with an oxide, in the form of a gray powder. The alloys of tin (one of which is pewter) are of considerable use; but, as well as the uses of tin in general, are too well known to need pointing out. The combinations of tin with acids, are used as chemical re-agents, and in the art of dying.

38.—IX. LEAD is a pale blueish metal, very soft, very heavy, and very malleable; but, though it may be drawn into wire, not very ductile. It melts at a low heat, and at a strong heat boils and evaporates. When rubbed on paper, it leaves a black mark. Lead unites to oxygen in

different proportions, forming differently-coloured oxides (see 90.) The oxides of lead are, by heat, converted into glass. Most of the acids combine with lead. The well-known poisonous substance called *sugar of lead*, is composed of lead and acetic acid. The proper counter-poison for a dangerous dose of sugar-of-lead, is a solution of Epsom or Glauber salts, swallowed in considerable quantity; either of which medicines instantly converts the *poisonous acetate* of lead, into the inert and *innocuous sulphate*.

39.—X. NICKEL is a fine white metal, hard, ductile, malleable, and very difficult of fusion. It is magnetical, but not so much so as iron. It is scarce, and little used. The combinations of nickel with acids, are distinguished by their fine green colour.

40.—XI. CADMIUM bears a considerable resemblance to *tin*, but it is more tenacious, and more fusible. It is rare, and not applied to any use.

41.—XII. ZINC is of a brilliant blueish-white colour, and has a crystallized appearance. It is hard, ductile, tenacious, and malleable, but only in a slight degree. It melts readily: just before melting, it is so brittle, that it may be pulverized. It is very combustible: when the fused metal has become red hot, it takes fire, continues to burn with a dazzling white flame, and is oxidised so rapidly, that it flies up in light filaments, called *flowers of zinc*, or *philosophical wool*. Zinc precipitates lead, tin, copper, silver, and some other metals, from their solutions. The substance known by the name of *white vitriol*, is a sulphate of zinc, and is, as are several of the alloys of this metal, exceedingly useful.

42.—XIII. BISMUTH is of a reddish-white colour, hard, brittle, and very fusible. When heated considerably, it evaporates. When raised to a strong red heat in contact with the air, it takes fire, and burns with a light blue flame, emitting a dense yellow smoke, which is an oxide of bismuth. When the metal, after being melted, is cooled gradually, it crystallizes. The alloys of bismuth (*pewter* is one of them) are all very fusible. The nitrate of bis-

muth affords, upon the addition of water, a white powder, which is used as a paint, under the name of *flake-white*.

43.—XIV. ANTIMONY is of a dusky-white colour, very brittle, but so soft, that, like lead, it may be cut with a knife. When heated to redness, it melts. If after this the heat be increased, the metal volatilizes in the form of white fumes, consisting of an oxide, formerly called *flowers of antimony*. A variety of useful alloys have antimony for one of their constituents: the metal for printers' types, that on which music is engraved, and pewter, belong to this class. Antimonial preparations serve also as paints and medicines.

44.—XV. MANGANESE has much the appearance of cast-iron. Its attraction for oxygen is so powerful, that it is preserved in the metallic state with great difficulty. The oxide of manganese is of great utility in the arts. The chemist, too, frequently uses that substance for the procuration of oxygen gas.

45.—XVI. COBALT is of a reddish-gray colour; brittle, rather soft, possessed of little lustre, and difficultly fusible. It is slightly susceptible of magnetism.

46.—XVII. TELLURIUM is a tin-coloured highly-lustrous metal, very brittle, and very volatile. It may be inflamed, and burns with a vivid blue light. It forms, with hydrogen, a gaseous acid, called telluretted hydrogen gas. But little is known, either of tellurium, or its combinations.

47.—XVIII. ARSENIC is a metal of a blueish-white colour, subject to become black by exposure to the air. It is the softest, most brittle, and most volatile of all the metals that are known. Its fusing-point cannot be discovered, because it evaporates without melting. It is exceedingly poisonous. It combines with two proportions of oxygen, forming *arsenious acid*, and *arsenic acid*. It combines with hydrogen, forming a very noxious gas, called *arsenuretted hydrogen gas*. The common arsenic of the shops is arsenious acid.

48.—XIX. CHROMIUM, a very rare metal. Slightly magnetical. White, brittle, lustrous. Acids act upon it with great difficulty. It unites with three different doses

of oxygen, forming *green* oxide, *brown* oxide, and with the largest proportion *chromic acid*, which is obtained in small ruby-red crystals.

49.—XX. MOLYBDENUM has only been obtained in very small quantities. It appears in the form of small blackish, lustrous, and brittle globules. It is the base of the molybdic acid.

50.—XXI. TUNGSTENUM, the base of the tungstic acid, is a metal much like steel. It is so hard that a file can scarcely make an impression on it, and in weight it is only inferior to gold. It is scarce, and not used for any thing.

51.—XXII. COLUMBIUM, which forms with oxygen columbic acid, is of an iron colour, hard, and brittle: other of its properties are not known. It is unacted upon by the strongest acids.

52.—XXIII. SELENIUM, the base of the selenic acid, is a gray metal, lustrous, brittle, soft, fusible, and volatile. It is distinguished by having the smell of *horse-raddish*.

53.—XXIV. OSMIUM has a dark-gray colour. When heated in the open air, is readily oxidised; but it resists the action of the strongest acids. It has not been melted, and is little known.

54.—XXV. RHODIUM is a whitish metal, very brittle, and as hard, and infusible, as iron. It is insoluble in acids. It unites to oxygen in several proportions, forming different-coloured oxides. It is very little known.

55.—XXVI. IRIDIUM is a heavy whitish metal, malleable, and infusible. It alloys with other metals. It has its name from the striking variety of colours it affords, while dissolving in muriatic acid. But little is known of its several combinations.

56.—XXVII. URANIUM is a hard, glittering, gray metal, obtained with great difficulty, and in very small quantities, from a mineral body resembling *pitch*.

57.—XXVIII. TITANIUM, a rare metal, of a dark copper colour, very heavy, lustrous, brittle, elastic, and highly-infusible. It is readily oxidised by heat. All the

dense acids act upon it energetically. It forms three oxides, the *blue*, the *red*, and the *white*.

58.—XXIX. CERIUM, a metal but little known, being scarce, and obtained with difficulty. It dissolves in acids, forming *sweet* salts. There are two oxides, one, the protoxide, being *white*; the other, the peroxide, *red*.

59.—XXX. WODANIUM, a metal of a bronze-yellow colour; hard and malleable. It is strongly attracted by the magnet. Its acid solutions are colourless; but its hydrated oxide, precipitated by caustic ammonia, is *indigo blue*.

60.—XXXI. POTASSIUM.—This marvellous metal was first revealed to the world by Sir Humphrey Davy, in 1807. He produced it, by means of voltaic electricity, from a substance which had been, till that time, considered simple, namely, the fixed alkali *potass*, which substance, however, he then proved to be a metallic oxide.

The properties of potassium are very extraordinary. It is lighter, not only than all other metals, but even than water. It is solid, soft, and of the colour and lustre of silver. But its metallic nature is not retained a single minute when it is exposed to the air: it instantly absorbs oxygen, and becomes covered with a crust of *potass*. This oxide absorbs water, which is rapidly decomposed, and in a short time the whole becomes a saturated solution of *potass*. Potassium is exceedingly inflammable; it fuses at 136° , and rises in vapour at a heat below redness. When heated in oxygen gas to the temperature at which it begins to evaporate, it burns with a brilliant white light, and an intense heat. When thrown upon the surface of water, it acts with great violence. The liquid is rapidly decomposed: its hydrogen is evolved, and spontaneously inflames in the air; it then communicates the combustion to the potassium, and the whole burns (while swimming about on the water) with a beautiful light of a violet-red colour: meantime, the oxygen of the water, and that drawn from the atmosphere by the combustion, unite with the potassium, and produce an oxide; which, as it is produced, dissolves in the water, and the final result is a *solution of pure potass*. Potassium likewise burns with a kind of explosion, when

placed (in a perfectly cold state) upon a piece of ice, dissolving the ice, and forming a little hole, in which it burns, till entirely consumed, forming, as before, a solution of potass. It spontaneously inflames, when put into chlorine gas, burning with great brilliancy, and producing chloride of potassium ; a substance which upon solution becomes muriate of potass.

Upon all fluids containing water, or much oxygen or chlorine, potassium acts with great rapidity ; and, in its general powers of chemical combination, that substance says its illustrious discoverer, may be compared to the *alkahest*, or *universal solvent*, so much talked of and longed for by the alchymists.

Potassium has, of all known substances, the strongest affinity for oxygen ; and it can only be preserved in the metallic state, by keeping it immersed in naphtha, a liquid of which oxygen is not a constituent.

61.—XXXII. SODIUM, the base of the alkali *soda*, was discovered by Sir Humphrey Davy, a few days after he discovered potassium. It was obtained in a similar manner, and is possessed of the following properties.

In several respects it bears a great resemblance to potassium, but is much less energetic in its actions. It is lighter than water, has the colour of silver, is solid, very soft, malleable, and a conductor of electricity ; but it is heavier than potass ; and requires a greater heat to fuse it, and a much higher still to volatilize it, than potassium does. When sodium is thrown upon water, it effervesces violently, produces a hissing noise, and swims about in a state of great agitation ; but it does not inflame. The water, however, is decomposed, and its oxygen combines with the sodium, and produces oxide of sodium, or soda. This product is dissolved by the water ; and the operation closes with the formation of a solution of pure soda. The combination of sodium with chlorine, is well known as *common table salt*, which substance, however, is, by the smallest addition of water, changed to muriate of soda. It must be observed, in explanation of this, that the water is decomposed : its oxygen unites to the sodium, forming soda ; and its hydrogen to the chlorine, forming muriatic

acid. It is evident, from this, that the aqueous compound is not chloride of sodium and water, but muriatic acid and soda, or the salt produced by their combination. When the solution of the salt is evaporated, the solid obtained is not dry muriate of soda, but chloride of sodium.

Sodium must, like potassium, and on the same account, be kept in naphtha.

62.—XXXIII. LITHIUM is a metal obtained by voltaism from a newly-discovered mineral alkali, called *lithia*. The quantity hitherto obtained has been much too small to allow of a full examination of its properties. From what is known, however, we conclude it to resemble sodium.

63.—XXXIV. CALCIUM, the metallic base of *lime*, has been obtained, by a voltaic process, from that base, but only in quantities too small to admit of an examination of its nature. It is of a bright white colour, burns brilliantly when heated in the air, and produces a white oxide, which is pure dry *lime*, a well known and very important substance.

64.—XXXV. BARIUM was procured by Sir Humphrey Davy, from its oxide *barytes*, by subjecting that earth to powerful electricity. It is of a dark-gray colour, is lustrous, heavy, and fusible. When exposed to the air, it rapidly attracts oxygen, and its surface becomes covered with a crust of barytes. It burns with a deep red light, when gently heated in the air. When thrown upon water, it effervesces violently; and, as it is heavy, it sinks into the fluid, which is decomposed with great rapidity; hydrogen gas is emitted, and the barium becomes barytes.

65.—XXXVI. STRONTIUM, the base of *strontia*,

66.—XXXVII. MAGNESIUM, the base of *magnesia*,

67.—XXXVIII. YTTRIUM, the base of *yttria*,

68.—XXXIX. GLUCINUM, the base of *glucina*,

69.—XL. ALUMINUM, the base of *alumina*,

70.—XLI. THORIUM, the base of *thorina*,

71.—XLII. ZIRCONIUM, the base of *zirconia*,

72.—XLIII. SILICIUM, the base of *silica*,

have, like barium, been procured in such minute quantities, that, as yet, their properties are but little known.

ON THE GENERAL POWERS PRODUCTIVE
OF CHEMICAL PHENOMENA.

73. Having presented to the notice of the student, the various simple or elementary bodies, we are now to perform a like office for the substances which result from their various combinations. Before, however, we commence the examination of the properties of the compound bodies, it will be proper for us to enter into an investigation of the nature of those Grand Powers, which, by their continual action upon common matter, change its form and produce arrangements fitted for the purposes of life.

74. **ATTRACTION.**—This is that unknown force, which causes bodies to approach each other. The instances of its exertion, exhibited in the phenomena around us, are exceedingly numerous, and continually present themselves to our observation.

75. Attraction exemplifies itself in a great variety of modes: *gravitation*, the *wonders of electricity and magnetism*, the *ascension of liquids in narrow tubes*, or porous substances all point out the exertion of this power. But, besides these here named, there is another kind of attraction—kind which comes immediately under the chemist's cognizance—that of *cohesion*. This operates among the *particles* of bodies, causing them to cohere, and form a whole.

76. The attraction of cohesion is differently named when spoken of with regard to different substances. If it causes the fixing-together of homogeneous particles, it is called the *attraction of aggregation*. But when it acts upon particles of different kinds, it is called the *attraction of composition*, or **CHEMICAL AFFINITY**.

77. The study of **CHEMICAL AFFINITY** is of vast importance; in fact, the whole science rests upon a knowledge of its modes of taking place. From the careful observance of the effects produced by certain experiments, chemists have deduced the following *laws*.

I.—The attraction of composition, or chemical affinity, takes place only between bodies of a different nature.

II.—It takes place only between the most minute particles of bodies.

III.—It can take place between two, three, four, or a greater number of bodies.

IV.—A change of temperature always takes place at the moment of combination.

V.—The properties which characterize bodies, when separate, are altered or destroyed by combination.

VI.—The force of chemical affinity, between the constituents of a body, is estimated by that which is required for their separation.

VII.—Every substance, though it has a certain affinity for all other substances, has different degrees of affinity for different substances.

This last law is the grand principle of all chemical operations; for it is by a proper arrangement of bodies which differ in their degrees of affinity, that all chemical decompositions and compositions are effected.

78. When a simple substance is presented or applied to another substance, compounded of two principles, and unites to one of those principles, so as to separate or exclude the other, the effect is said to be produced by *simple elective affinity*.

79. Double elective attraction, or *compound affinity*, (the terms are synonymous) takes place when two bodies, each consisting of two principles, are presented to each other, and mutually exchange a principle of each, by which means two new bodies, or compounds, are produced, of a different nature from the original compounds.

80. Upon the principle, that “bodies differ in their degrees of affinity,” chemists have constructed *tables of affinity*: these, though, like most other things, liable to objections, are so extremely useful, that the study of them cannot be too strongly recommended. An accurate “TABLE OF CHEMICAL DECOMPOSITIONS” is given in the APPENDIX. The illustrations of chemical affinity contained in *section fifth* of our *Experiments*, develope the doctrine in a very agreeable manner.

81. REPULSION—CALORIC.—The opponent power of *attraction*, is that which causes bodies to *recede* from each other, and which is known by the name of *repulsion*. That

substance, possessed of this power, to which the attention of chemists is chiefly directed, is *caloric*.

82. The appellation of *caloric* has been given, by modern chemists, to that which produces the well-known sensation called *heat*.

83. Caloric penetrates all bodies; separating their particles from each other, and thus increasing their bulk. Solids, by an increase of heat, become fluids, and fluids, gases. Thus ice is converted, by an accession of caloric, into water, and by a still farther accession, it becomes steam.

84. The subject of *heat* is pursued at considerable length in *section the first* of the experimental part of this work. The student should make this division of the science of Chemistry an early object of his attention; as, unless he comprehend the manner in which caloric acts, the rationale of scarcely a single chemical process will be intelligible to him.

85. COMBUSTION—COMBINATION.—Combustion is considered as the general result of intense chemical action. When oxygen and chlorine unite with the simple combustibles, compounds possessed of most remarkable properties are formed: the combinations take place with astonishing energy, and with the evolutions of light and heat.

86. In all common cases, *combustion* is the process of the solution of a body in oxygen; as when sulphur or phosphorus are burnt in it, or when it forms water, by combining with hydrogen,

87. The union of considerable quantities of oxygen and chlorine with combustible bodies, generally produce acids; thus sulphuric, phosphoric, and boracic acids, are formed by the union of considerable quantities of oxygen with portions of sulphur, phosphorus, and boron—and muriatic acid gas is formed by the union of chlorine and hydrogen. See the article ACIDS, (97.)

88. When these solvent substances, oxygen and chlorine, unite with combustible bodies in *smaller* quantities, they produce compounds, not acid, and, in a greater or less degree, soluble in water. A vast number of compounds are thus formed by oxygen: the earths, the fixed alkalies, and

the metallic oxides, bodies connected by analogies with each other, are all of this class. See the article OXIDES, (133).

89. ATOMIC THEORY, or *Doctrine of Definite Proportions*.—One of the latest and most important chemical discoveries, is, that the elementary bodies enter into combination in *definite* proportions; that is to say, if two simple bodies combine and form a particular compound, they always unite in the same proportions to form that compound: they, however, sometimes enter in other proportions into the composition of other compounds.

This law is well exhibited in the combinations of gaseous bodies. These are seen to unite in simple ratios of volumes. Water is composed of hydrogen and oxygen, and 1 part by weight of the former gas, unites to 7.5 of the latter. The specific gravity of hydrogen, compared with that of oxygen, is as 1 to 15; it is obvious, therefore, that one volume of hydrogen unites to half a volume of oxygen, and that the composition of water will be represented by weight and volume, thus:

1	7.5
Hydrogen.	Oxygen.

No other proportions of these gases than those here represented, can ever combine and form water: nor is it known that they form any other body, by combining in other proportions. This, however, is not the case with all bodies.

90. The metallic oxides, in general, consist of the metals, united to from one to four proportions of oxygen; and there are, in some cases, many different oxides of the same metal: thus, there are three oxides of lead—the yellow oxide, or massicot, containing two proportions of

oxygen; the red oxide, or minium, three; and the puce-coloured oxide, four proportions. Again, there are two oxides of copper, the black and the orange; the black contains two proportions of oxygen, the orange, one.

91. When a metal is oxidised in different degrees, the several compounds have a prefix to their names, to distinguish them. The smallest addition of oxygen forms a *protoxide*, the second quantity gives rise to the *deutoxide*, and the third the *tritoxide*—farther, the oxide which contains the largest quantity of oxygen, is generally called the *peroxide*.

92. As the doctrine of definite proportions must, to beginners in the science, appear somewhat intricate, we shall not, in this place, pursue it at any length; more especially as the other doctrines of Chemistry, and the other chemical properties of different kinds of matter, may be understood without it. The subject will, however, be spoken of in an article to be given in the *Appendix* to this work, to which the student is referred.

93. CRYSTALLIZATION.—When fluid substances are suffered to pass with adequate slowness to the solid state, the attractive forces frequently arrange their ultimate particles, so as to form regular geometrical solids, to which has been given the name of *crystals*.

Perfect mobility among the corpuscles is essential to crystallization. The chemist produces it either by igneous fusion, or by solution in a liquid. When the temperature is slowly lowered in the former case, or the liquid slowly extracted by evaporation in the latter, the attractive forces resume the ascendancy, and arrange the particles in symmetrical forms. Bodies, in crystallizing from their watery solution, always retain a small portion of water, which remains confined in the crystals in a state of solidity, and does not re-appear till the crystalline form of those bodies is destroyed. This is called the *water of crystallization*.

It is well worthy of remark, that every individual salt has a certain determinate figure, which it assumes upon crystallization.

The operation of crystallizing is of great utility in the purifying of various saline substances.

COMPOUND BODIES.

94. These are innumerable, and some of them are of vast importance. The list of those with which chemists are intimately acquainted, is very extensive; but as, in this small volume, nothing can be expected but the *Rudiments* of the science, and a familiar account of the most important bodies, it would be ridiculous, were we to present the student with a large catalogue of compounds, without (for we should have no room to include) descriptions of their properties. Such a course of proceeding, instead of directing him in his pursuit of the facts of chemistry, would serve only to perplex him. We shall, therefore, confine our attention to such as, by their great utility or striking peculiarities, seem especially to require it.

95. ATMOSPHERIC AIR, is the term applied to that immense mass of permanently elastic fluid, which surrounds the globe we inhabit. This substance was for a long time supposed to be simple, but recently it has been proved, by experiment, to be a compound of oxygen and nitrogen. 100 parts of common air, contain 21 parts by measure of the former body, and 79 of the latter. The proportions of these two gases, in atmospheric air, are uniform and constant. They have been found to be the same in all parts of the world, and in all seasons of the year. The density of air is $\frac{1}{8\frac{1}{2}}$ th of that of water. Its physical properties, as its transparency, its elasticity, and so on—are so obvious, as not to require description. Air which has been breathed, is found to have lost its oxygen, and is then no longer fit to support life. This principle is retained in the lungs, where it is absorbed by the blood, from which it expels *carbonic acid gas*, and which it renders capable of supporting life. The red colour of the blood is owing to the oxygen it acquires in passing through the lungs. As air which will not support combustion is not fit for respiration, we hence have recourse to the efficacy of air in supporting combustion, in order to determine its purity.—At 455 is shown how air may be *composed*, and at 451, how it may be *decomposed*.

96. WATER, a fluid of which the most useful properties are universally known, and properly appreciated. Without it, neither vegetable nor animal life could be supported. Like common air, *water*, which was formerly considered a simple body, has been found to be a compound one. Its principles being *hydrogen* and *oxygen*; consigned in the proportions of, by weight, 1 part of the former to 7.5 of the latter, or, by measure, of 2 of the former to 1 of the latter.

Pure water is perfectly transparent, tasteless, inodorous, and not liable to change. Rain water is the purest we see in nature. Spring water has always some mineral impurities. By distillation, it is freed from whatever contaminates it. The composition of water is proved at 440; and at 673, the mode of its decomposition is described.

ACIDS.

97. These form the most important class of chemical compounds. They are generally produced by the combination of particular substances with oxygen: thus, sulphuric acid is formed by the union of sulphur with oxygen—and carbonic acid, by the union of carbon (charcoal) with oxygen. But oxygen, however, though the general, is not the only acidifying principle, for hydrogen, chlorine, and iodine, are producers of acids likewise. The following are the general properties of the acids:—

I. The taste of these bodies is, as their name denotes, *sour*; and in the stronger species, it is acrid and corrosive.—II. They combine with water in every proportion, with a condensation of volume, and evolution of heat.—III. They are volatilized or decomposed at a moderate heat.—IV. They change the blue, green, and purple colours of vegetables, to a bright red.—V. They unite in definite proportions with the alkalies, earths, and metallic oxides, and form the important class of bodies termed *salts* (153). This may be reckoned their characteristic and indispensable property.

98 ACIDS, in order to give general views to students, by grouping together substances which are in some respects similar, have been classed as follows:

Division 1st. Acids from inorganic nature, or such as are procurable without having recourse to animal or vegetable products.—Division 2d. Acids elaborated by means of organization.

The first group is subdivided into three families:—

I. Oxygen acids, $\left\{ \begin{array}{l} \text{non-metallic,} \\ \text{metallic.} \end{array} \right.$

II. Hydrogen acids.

III. Acids containing neither oxygen nor hydrogen.

The number of the animal and vegetable acids (those belonging to the second division) is forty-six. The principal of which are described at 238. The total number of acids now proved to exist, is eighty-seven. As many of these are, however, but little known, are of no importance, and may be found particularized in any large work on Chemistry, we shall not here describe them; but confine our attention to such as are much employed, and which it is necessary for the young chemist to become early and fully acquainted with.

99. Previously, however, it will be necessary to explain the nomenclature of acids; for to be acquainted with the mode of *naming* these curious compounds is a thing of moment.—When an acid is formed by the combination of oxygen with another body, as sulphur, which is called the *base*, the acid takes for *its* name the name of its base, with the addition of the syllable *ic*—thus *sulphuric acid*. In the cases where oxygen forms two acids from the same base, by combining with it in different proportions, the termination *ic* is put to that acid which contains the *largest* proportion of oxygen; and to the other acid is affixed the syllable *ous*. This is fully illustrated by the Table of Acids at 102.

100. SULPHURIC ACID.—When sulphur is burnt in oxygen gas, and the gaseous combination of sulphur and oxygen is mixed with water, the liquid product is found to possess, in a very eminent degree, the distinguishing properties of acids. It was formerly called *oil of vitriol*; a foolish name, as the liquid possesses not one of the properties of an *oil*. Its modern appellation, *sulphuric acid*, at once denotes its composition and its chief properties.

It is composed of two parts of sulphur united to three of oxygen. When in the pure concentrated liquid state, it has neither smell nor colour. It is violently caustic, and is therefore poisonous. Its density is twice that of water for which body it has a strong attraction, see 625. When mixed with water, great heat is produced, see 386; but when mixed with ice, cold is produced,—see Table of Freezing Mixtures in the Appendix. Sulphuric acid is prepared in the large way, by burning sulphur mixed with a large portion of nitre, in close chambers lined with lead. The nitre furnishes oxygen to the sulphur, and the acid, as it is produced, combines with a quantity of water with which the floor of the chambers is previously covered. This liquid acid is afterwards distilled, to free it as much from water as possible. That sulphuric acid is formed of sulphur and oxygen, is proved by the experiment at 418. The use of this acid in chemistry, metallurgy, bleaching, dying, medicine, and other arts, is very extensive. It combines with the various alkalies, earths and metals, forming a class of bodies called *sulphates*, (or *salts of the sulphuric acid*,) which are described under the head “SALTS.”

101. **SULPHUROUS ACID.**—This is formed of sulphur combined with a smaller proportion of oxygen than is necessary to form sulphuric acid. Sulphurous acid, in the ordinary temperature of the atmosphere, is a gas, composed of one volume of oxygen, and one of vapour of sulphur. It has a strong suffocating smell, and possesses the property of destroying vegetable colours, (see 469,) which renders it useful in some bleaching operations. Its combinations with salifiable bases, are termed *sulphites*. These are of little use.

102. **HYPOSULPHUROUS ACID.**—This is composed of sulphur, united to a less portion of oxygen than sulphurous acid. **HYPOSULPHURIC ACID.**—This, which is also sulphur acid, contains more oxygen than the sulphurous but less than the sulphuric. These acids are not of much importance.

* * * The following table of the just-described acid com-

pounds of sulphur and oxygen, may serve to fix their compositions in the learner's memory.

ACIDS.	PARTS OF SULPHUR.	PARTS OF OXYGEN.
Sulphuric, ~~~~~	2	3
Hyposulphuric, ~~~~~	2	2.5
Sulphurous, ~~~~~	2	2
Hyposulphurous, ~~~~~	2	1

103. PHOSPHORIC ACID.—This is a compound of oxygen and phosphorus. It may be procured by burning the latter body in the former, (see 419,) but is much more cheaply obtained from calcined bones. The mineral, vegetable, and animal kingdoms abound, either with the base of this acid, or the acid itself. Pure phosphoric acid is in a solid state, but it is soluble, in all proportions, in water. It has no smell, is sour, but not corrosive. Its salts are termed *phosphates*.

104. CARBONIC ACID.—This is a compound of oxygen and carbon. It may be obtained by burning carbon in oxygen gas, (see 416); but it exists ready formed, in great abundance, in nature. It composes forty-four per cent. of the weight of all lime-stone, chalk, marble, &c.; from which it may easily be separated, as almost every other acid has a superior affinity for the base of these stones.

Carbonic acid is obtained in the gaseous state, and cannot be condensed into the liquid or solid state. It is capable, however, of being absorbed or condensed in water, and of thus forming a weak acid. Carbonic acid gas extinguishes flame, and destroys animal life. It is much denser than common air, and hence always occupies the lower part of places where it is formed. Miners call it the *choke-damp*. It is emitted in large quantities, by bodies that are *fermenting*, and is the principle that gives to bottled porter, ginger-beer, &c. the effervescing, or frothing property. It has a peculiar sharp taste. It reddens

infusion of litmus fugaciously. 100 parts of carbonic acid gas, contain 72 oxygen, 28 carbon. The salts of this acid, which are an important class, are called *carbonates*.

105. BORACIC ACID.—This is a compound of oxygen and a lately discovered body, called boron (25). A substance well known by the name of borax, is composed of this acid united to soda. Boracic acid is obtained in the form of thin crystals, (or scales,) of a silvery-white colour; having a greasy feel, no smell, but a very strange taste—being first sourish, then bitterish, cooling, and at last agreeably sweet. It is soluble in water, but only in a slight degree. It is more so in alcohol, and gives to the flame of that body, when burning, a green colour (see 589.) It is fusible, and is converted by fusion into a kind of glass, which is used in the composition of false gems. The salts of boracic acid are called *borates*.

106. NITRIC ACID.—This body, which is of a deadly poisonous nature, has for its component principles the same bodies as common air, namely, oxygen and nitrogen. They are united in it, however, in different proportions. Atmospheric air is composed of 1 volume of oxygen, and 4 of nitrogen: nitric acid gas of $2\frac{1}{2}$ volumes of oxygen and 1 of nitrogen. Nitric acid gas combines with water very eagerly, and communicates to the water its properties.—Strong pure liquid nitric acid is about one-half denser than water, and colourless. It is eminently corrosive: of this property, [its old name, *aqua fortis*, (strong water) is very expressive. The *aqua fortis* of commerce is nitric acid in a state of impurity. Nitric acid is obtained by pouring two parts of strong sulphuric acid upon one part of saltpetre, in a retort, and distilling the mixture. The process is not a dangerous one, if the red fumes be avoided. It is proper to insert a bent tube into the tublature of the receiver, one end of which is immersed in a small quantity of water. This will condense the fuming portion. Nitric acid is of considerable use in the arts: being employed for etching on copper, in dying, metallurgy, assaying, and medicinal preparations; also, in various chemical processes, on account

of the facility with which it dissolves metals, by first parting with a portion of its oxygen to them, and then attacking the oxides. But there are two metals upon which it has no effect—these are gold and platinum; even these, however, may be dissolved by a liquid, of which nitric acid forms a part.—The salts of nitric acid are of considerable use; they are termed *nitrates*.

107. NITROUS ACID.—This is said to be composed of $\frac{1}{2}$ oxygen to 1 nitrogen, but much uncertainty prevails regarding it. Indeed, its very existence is doubted. The orange-coloured volatile fluid, to which the name has been usually applied, is held by some to be nitric acid, holding a particular sort of air, called nitrous gas, in solution. Its salts (if there are such) are NITRITES.

108. MURIATIC ACID.—This, in a gaseous state, consists of chlorine gas, and hydrogen gas, united in equal volumes. Muriatic acid is obtained by pouring, by degrees, 6 parts of sulphuric acid upon 5 parts of dry sea salt, previously put into a tubelated retort, the beak of which is connected with Woolf's apparatus, containing 5 or 6 parts of water, and allowing the gas to pass through it. Muriatic acid gas is invisible, has a peculiar pungent odour, and a sour corrosive taste. It is one-fourth heavier than common air. It extinguishes flame, and is destructive of animal life. Water absorbs 480 times its bulk of this gas, forming liquid muriatic acid. This, when pure, is colourless, but owing to impurities, the muriatic acid of commerce has usually a straw colour. It is generally obtained from common salt, and hence has been called *spirit of salt*. It is an acid that is much used in the arts, and chemical laboratories. Its salts, (termed *muriates*,) when in a state of dryness, are actually *chlorides*, consisting of chlorine and the metallic base; but since moisture makes them instantly pass to the state of *muriates*, we shall describe them under that name.

109. FLUORIC ACID.—This acid has hitherto resisted every attempt to decompose it completely, and consequently its real nature is uncertain. It is commonly supposed to consist of a base called fluorine united to hydrogen; and hence is sometimes called hydrofluoric acid. The gaseous

state is that in which it is procured : it is soluble in water. It is intensely sour, corrodes and instantly dissolves glass (see 613,) and produces very painful wounds when touches the skin, so that it requires to be examined with great caution. Its salts are called *fluates*.

110. PRUSSIC ACID.—This is formed of 1 part of cyrogen and 1 part of hydrogen. It is liquid, possessed a hot, but sweetish taste, and having a very pungent smell which resembles that of bitter almonds. It is the most violent of all poisons ; causing the death of animals, when applied to the skin only. Prussic acid may be obtained readily by making a saturated solution of prussiate of potash and dropping into it as much tartaric acid as will precipitate the potash, which it does by uniting with it, and forming tartrate of potash, (cream of tartar). When the precipitate has subsided, the fluid may be decanted, and slowly distilled, with a moderate heat, into a receiver kept cool with water, or pounded ice. It combines with iron and forms the well-known pigment—*prussian blue*. Some of its salts (the *prussiates*) are valuable as *re-agents*.

111. ACETIC ACID.—The same acid, that, in a very dilute, and somewhat impure state, is called *vinegar*. Its constituent principles are hydrogen, carbon, and oxygen. When pure it is colourless, pungent, acrid, and volatile. It can be concentrated till capable of corroding the skin. Almost all dry vegetable substances, and some animals when subjected to a red heat in close vessels, yield vinegar copiously. It is the result of spontaneous fermentation likewise. A very strong vinegar, used for pickling, &c. is now obtained by distilling dry wood in iron retort. Acetic acid is of great importance, for it is not only extensively used for domestic purposes, but applied to many of the arts. Its salts are called *acetates*.

112. CITRIC ACID, is the purified juice of limes or lemons. It is obtained in the form of crystals, which have an extremely sharp taste, are fusible by heat, and very soluble in water. Citric acid is of considerable service in domestic economy, and in some of the arts. Some of its saline compounds (termed *citrates*) are also much used.

113. BENZOIC ACID, is commonly known by the name

of flowers of benjamin, a substance obtained by sublimation from gum benzoin (see 644). It appears in white needle-shaped crystals, having a bitter taste, and an aromatic odour. It very readily dissolves in alcohol, but is scarcely at all soluble in water. Its salts are termed *benzoates*.

114. GALLIC ACID is obtained from various vegetable substances, particularly from nut-galls. It is a crystallizable acid, soluble in 20 parts of water. It has an astringent acidulous taste, and, when heated, an aromatic, though not pleasant, odour. It is extensively used in dying, ink-making, and by the chemist as a re-agent. Its salts are called *gallates*.

115. OXALIC ACID is a deadly poison, which, when in crystallized state, has much the appearance of Epsom salt—in place of which compound it has been taken accidentally, and has more than once proved fatal. The mode of distinguishing oxalic acid from Epsom salt, is explained at 547. The best antidote to a dose of it is an emetic, aided by copious draughts of warm water, containing chalk, or common carbonate of magnesia. Oxalic acid has an exceedingly sharp, sour, but not unpleasant taste; it is soluble in water; and is possessed of some bleaching properties. Oxalic acid may be obtained by pouring 4 parts of strong nitric acid upon 1 part of lump sugar, in small pieces, previously put into a stoppered retort, to which a receiver is fitted, distill off a considerable portion of the acid, and set the retort aside to cool, when a great number of needle-shaped crystals will be obtained. It is obtained also from the juice of wood-sorrell, and from many other vegetables. Its salts are termed *oxalates*.

116. TARTARIC ACID.—This is that which, united to potass, forms what is called *cream of tartar*. It is obtained in crystals, which are soluble in water. It has a very acid and agreeable taste—so that it may supply the place of citric acid, than which it is cheaper. Its salts are called *tartarates*.

ALKALIES.

117. These are substances of great importance. They may be defined to be bodies which combine with acids, neutralize their powers, and produce salts. They are divided into three classes, thus:—

I. Those consisting of a metallic base united to oxygen, which are in number three—*potass*, *soda*, and *lithia*.—II. That containing no oxygen—*ammonia*.—III. Those containing oxygen, hydrogen, and carbon;—these are some recently-discovered vegetable alkalies, of which little is known.

Besides the already-mentioned properties of alkalies they possess the following also:—I. They change the purple colour of many vegetables to a green, the red to purple, the yellow to a brown. They also re-produce the purples which have been reddened by acids.—II. They continue in possession of this property, and retain the solubility in water, when saturated with carbonic acid; these properties distinguish them from the alkaline earths.—III. They have an acrid and urinous taste.—IV. They are powerful corrosives, or solvents of animal matter, with which, as well as with oils in general, they combine, so as to produce neutrality.—V. They are decomposed or volatilized by a strong heat.—VI. With water they combine in every proportion; and with alcohol, very largely.

118. POTASS, a *fixed alkali*, procured from wood-ash, and called on that account, the *vegetable alkali*. It is the oxide of the newly-discovered metal, *potassium*.

Potass, when pure, is solid, white, and extremely caustic. It changes the purple colour of vegetables to a green, and yellow to a brown. When exposed to the air, it rapidly attracts humidity from it, and becomes a liquid. In Chemistry it is employed very extensively, both in manufactures and as a re-agent in analysis. It is the basis of the common soft soaps. Potass combines with the other acids also, forming neutral salts. The substance known in commerce by the name of potash, is not pure potash.

but potass united to one proportion of carbonic acid ; it is a sub-carbonate of potass.

119. SODA, which, because supposed to be peculiar to the mineral kingdom, was called the *mineral alkali*, is, when pure, of a gray colour, and in its properties bears considerable resemblance to *potass*. It does not, however, like that substance, liquify if exposed to the air, but falls to powder. The metallic base of this alkali is *sodium*. *Soda* is the basis of common salt, of plate and crown glass, and of all hard soaps.

120. LITHIA, a new fixed alkali, whose metallic base is LITHIAM. It has a very sharp burning taste, is soluble in water, changes vegetable blues to green, and forms neutral salts with the acids. It is scarce.

121. AMMONIA, also called the *volatile alkali*, is composed of nitrogen and hydrogen ; and in its purest form, (the gaseous,) is called ammoniacal gas. Water dissolves 460 times its bulk of this gas, forming with it liquid ammonia. Ammoniacal gas has an exceedingly pungent smell, well known by the name of *spirits of hartshorn*. An animal plunged into it, speedily dies. It extinguishes combustion, has a very acrid taste, and, like *fixed* alkalies, changes the blue colour of vegetables to green. It is a valuable material in manufactures, in medicine, and in Chemistry. All the acids are capable of combining with it. For the method of procuring ammoniacal gas, see 468.

* * * The alkalies of the third class—those containing oxygen, hydrogen, and carbon,—are of so little importance, that in this place no notice need be taken of them.

EARTHS.

122. These are the substances that compose the vast crust of rocks and soils on the surface of the globe, and which, from their very diversified appearance, appear to be innumerable. Analysis, however, has shown, that all these heterogeneous stony masses may be resolved into only *ten* different bodies. The names given to these bodies, are :—

1 *Barytes*, 3 *Lime*, 5 *Alumina*, 7 *Glucina*, 9 *Silica*,
2 *Strontia*, 4 *Magnesia*, 6 *Yttria*, 8 *Zirconia*, 10 *Thorina*.

The *earths* are, in general, infusible and incombustible, have no smell, and are rather less than five times heavier than water. On account of some *particular* properties, they have been divided into two classes :

1. *Alkaline* earths.—2. *Common* earths.

The *alkaline* earths possess some properties similar to those possessed by *alkalies*, but, nevertheless, widely differ from those bodies in their *general* properties. The earths of this class, are the first four in the above list.—The *common* earths possess no alkaline properties : they are the latter six in the list.

One of the most important of modern chemical discoveries, is, that the *earths*, instead of being, as was supposed, *simple bodies*, are combinations of *oxygen* with peculiar *metallic bases*.

123. BARYTES is an earth of a grayish-white colour, possessed of strong alkaline properties, changing, like them, the vegetable blues to green. Its taste is extremely caustic, and it is a violent poison. It has no smell. When water is poured upon dry *barytes*, it slakes like quicklime, evolving, however, more heat. It is soluble in water, forms an insoluble compound with sulphuric acid, and tinges flame yellow. Under the name of *permanent white*, it is used as a paint. It is employed by the chemist as a re-agent. The metallic base of barytes is *barium*.

124. STRONTIA, in its properties, has a considerable affinity to *barytes*. It differs from it chiefly in being infusible, much less soluble, of a different form when crystallized, weaker in its affinities, and not poisonous. Its base is *strontium*. A peculiar characteristic of strontia, is the giving a blood-red colour to flame. Its *salts* are numerous, but not useful.

125. LIME is one of the most abundant substances in nature. It is the chief constituent of vast rocks and mountains, under the names of chalk, marble, limestone, calcareous spar, &c., in the whole of which it exists, combined with carbonic acid. By exposure to a strong heat, these substances are rendered pure, or form what is com-

monly called *quicklime*. The recently discovered metallic base of lime, is named *calcium*.

Lime is a soft white substance, with a caustic, astringent, and alkaline taste. It is fused with great difficulty. It is soluble in water. It renders vegetable blues, green; yellows, brown; and restores to reddened litmus, its usual purple. If a little water only be sprinkled upon dry newly burnt lime, it is rapidly absorbed, with the evolution of much heat and vapour. It has an immense affinity for carbonic acid, which, when it is exposed to the atmosphere, it rapidly absorbs.

126. *MAGNESIA* is a white soft powder, having a metallic base, called *magnesium*. It renders the infusion of red cabbage, green, and reddens turmeric. By an intense heat, and only by that, it may be fused. It has little taste, and no smell. It is nearly insoluble in water, but absorbs a quantity of that liquid with the production of heat. *Magnesia*, and its combinations, (*Epsom-salt* is its *sulphate*,) are chiefly used in medicine.

127. *ALUMINA* is an earth, which, as constituting the plastic principle of all clays and loams, was called *argil*, or the *argillaceous earth*, but now, as being obtained in greater purity from *alum*, is styled *alumina*. It has a metallic base, named *aluminum*. Pure *alumina* is soft, pulverulent, white; adheres to the tongue, forms with water a smooth paste, produces no change on vegetable colours, is insipid, inodorous, and insoluble in water, but mixes with it in all proportions. It is hardened by fire. It is the basis of some hard gems, and of earthenware. It is dissolved by most acids. *Alum* is a *sulphate* of *alumina* and *potash*.

128. *SILICA* is a compound of a peculiar combustible principle, (the metal *silicium*,) with oxygen. This earth exists nearly pure in flint and rock crystal. It is white, tasteless, infusible, and insoluble in water. It feels gritty. It unites with *alkalies*, and forms *glass*. In the state of gravel, it is, on account of its hardness, much valued for roads. It is a necessary ingredient in cements, porcelain, and glass. No acids dissolve it, except those containing fluorine.

129. GLUCINA, the oxide of the metal *glucinum*, is exceedingly rare: as yet, it has been obtained only from one or two precious stones. It is a soft white powder, light, tasteless, and adhering to the tongue. Its name is derived from its distinguishing character of forming with acids, salts which are sweet to the taste.

130. ZIRCONIA is, like glucina, procured from gems; consequently it is rare. Its metallic base is *zirconium*. It is a fine white powder, hard to the touch, without taste or smell. Insoluble in water, but uniting with all acids.

131. YTTRIA, whose base is *yttrium*, is white, tasteless, inodorous, infusible, and very heavy. Water, and pure alkalies, do not dissolve it, but carbonate of ammonia does. With borax it melts into a glass; and is soluble in most acids, forming with them sweet and coloured salts.

132. THORINA is a lately discovered earth, resembling *zirconia*. To its supposed metallic base, the name *thorium* has been applied. Carbonic acid is very readily absorbed by it, and forms with it, (as will almost any other acid,) an astringent-tasted *salt*.

133. OXIDES.—This term denotes, that the bodies to which it is applied, are combinations of various substances with oxygen, in all cases where the oxygen is not in sufficient quantity to make the compound *acid*.

Some observations respecting oxidation are to be found at 88, 90, 91. There are many bodies, which are properly oxides, that are not generally known by that name. Atmospheric air, for instance, is an oxide of azote, and water, an oxide of hydrogen. The earths, and alkalis also, (see NOTE 29,) are oxides. Now, under the names here mentioned, and at the paragraphs referred to, the most important oxides have been already described; and a few only remain to be noticed in this place.

134. NITRIC OXIDE, or *nitrous gas*, is a compound of oxygen and nitrogen, in equal volumes. It is colourless, but when suffered to mix with air, or oxygen gas, produces reddish fumes, being by its union with oxygen converted into nitrous acid gas. Nitric oxide is fatal to animal life, and extinguishes flame. There are, however, a few bodies

that can be burnt in it. Water absorbs about 1.20ths of its bulk of this gas.—See experiments, 456 to 459.

135. NITROUS OXIDE, a gas composed of two proportions of nitrogen, and one of oxygen. It supports combustion, but (though it may be respired) is not capable of supporting life. It has a sweet taste, and a faint, but agreeable odour. Water (which has been boiled) absorbs about one-half its bulk of it; but acquires thereby no other property than that of sweetness, and gives out the whole of it again, when boiled or frozen. The most extraordinary property of this gas, is its action on the human body, when respired. The sensations that are produced vary greatly in persons of different constitutions; but, in general, they are highly pleasurable, and resemble those attendant on the pleasant period of intoxication. It has been called *intoxicating gas*, *laughing gas*, and *gas of Paradise*. This characteristic of nitrous oxide was discovered by Sir Humphrey Davy. The method of obtaining and of respiring this gas, is described at 460 to 462.

136. CARBONIC OXIDE, a gaseous body, composed of half as much oxygen, united to a certain proportion of carbon, as is necessary to form carbonic acid. It is void of taste and smell, and is fatal to animal life. But it is inflammable; its combustion is attended with a blue flame.

137. CHLOROUS OXIDE, and CHLORIC OXIDE, are gaseous compounds of oxygen and chlorine. The preparation of them, and examination of their properties, is attended with much danger, for they are of an exceedingly explosive nature: consequently, they are little known, and they are not at all used.

138. CHLORIDES.—The most important of the bodies to which this name properly belongs, are described under the name of *muriates*. One of the *chlorides*, (that of *lime*,) commonly called *bleaching powder*, is much used in bleaching processes. It is made by forcing chlorine gas into a close vessel, containing dry newly-burnt lime.

139. IODIDES.—Bodies of this class are numerous, but very unimportant, being entirely useless.

140. BORURETS are compounds of metals with boron. They are but of little consequence.

141. CARBURETTED HYDROGEN is a gas composed of equal parts of carbon and hydrogen; sometimes, however, the gas is obtained with a less proportion of carbon in which case it is called sub-carburetted hydrogen. Carburetted hydrogen gas is heavy, invisible, and, when pure, void of taste and smell: it generally, however, has a sulphurous smell, owing to its holding a portion of sulphur in solution. By the combustion of carburetted hydrogen we obtain a splendid white flame, which is now much employed in domestic economy, in preference to the light of candles. The much-dreaded *fire-damp* of the coal mines is identical with the gas under consideration. The way to procure it is shown at 463.

142. PHOSPHURETTED HYDROGEN, is a gaseous compound of phosphorus and hydrogen. Its characteristic property is extreme combustibility: it inflames by merely coming into contact with the atmosphere. A very brilliant white light attends its inflammation in oxygen gas. When brought into contact with chlorine, it detonates with a brilliant green light. It has a very disagreeable smell, resembling that of putrid fish. It combines in a very slight degree with water. Those flashes of light, called by the vulgar *will-of-the-wisp*, and by some *ignes-fatui*, which are often seen in churchyards, and other places where vapours are exhaled from putrifying animal matter, are produced by the formation and inflammation of this gas. A very pretty experiment with this gas, is described at 464.

143. SULPHURETTED HYDROGEN is a gas, whose component principles are hydrogen and sulphur. It is twice the weight of common air, and is slightly absorbed by water. It burns (when lighted) with a pale blue flame, depositing sulphur. It does not support combustion. Its smell is extremely fetid, resembling that of rotten eggs. Its taste is sour. It reddens vegetable blues. Hence, by some, it has been reckoned an acid, but is not fully (though nearly) entitled to that appellation. Of all the gases, sul-

phuretted hydrogen is perhaps the most deleterious to animal life. A dog of middle size is destroyed in air, containing only 1.800ths of its bulk of it. Indeed, it has been proved, that to kill an animal, it is sufficient to make the sulphuretted hydrogen act on the surface of its body, when it is absorbed by the inhalents. Yet, to the presence of this gas, is chiefly owing the beneficial medicinal properties of Harrowgate, Aix-la-Chapelle, and some other mineral waters. Sulphuretted hydrogen is employed by the chemist as a re-agent. It unites with several bodies, forming hydro-sulphurets. The process for procuring this gas is given at 465.

144. ARSENURETTE D HYDROGEN is a gaseous compound of arsenic and hydrogen. It is inflammable, extinguishes flame, and destroys animal life instantaneously.

145. TELLURETTE D HYDROGEN, a gas composed of tellurium and hydrogen, is soluble in water, and gives a claret-coloured solution. It combines with the alkalies. It burns with a blue flame. It smells like rotten eggs.

146. POTASSURETTE D HYDROGEN.—When hydrogen and potassium are passed through a tube heated to whiteness, the gas dissolves the metal, and potassuretted hydrogen gas is formed, and inflames spontaneously.

147. CARBURET OF SULPHUR—SULPHURET OF CARBON—ALCOHOL OF SULPHUR.—(It is differently named) an interesting compound of carbon and sulphur,—prepared with difficulty,—possessed of the following properties: it is an oily liquid, eminently transparent, and perfectly colourless. It has a pungent aromatic taste, and a nauseous fetid smell. It boils briskly at about half the heat of boiling water, and at 600° inflames, burning with a blue flame. Mixed with oxygen, and having the elective spark passed through it, it detonates.

148. CARBURET OF IRON—STEEL.—This is a compound of iron and carbon. It may be made, by keeping iron mixed with charcoal in a crucible, for a number of hours, in a strong heat. It is hardened by being thrown, when red-hot, into cold water. Steel contains scarcely 1.200ths of its weight of carbon, and yet its properties differ very

greatly from those of pure iron. The uses of this compound are well known.

149. PLUMBAGO is another carburet of iron. This substance is generally called *black-lead*, which, as it contains not an atom of lead, is exceedingly absurd. The only thing in which plumbago and lead agree, and from which the name must have originated, is the making a black mark on paper. Plumbago is composed of 90 carbon + 10 iron. Large quantities of it, in a very pure state, are found at Keswick, in Cumberland.

150. PRUSSINE, or *cyanogen*, a substance formed by the combination of carbon and nitrogen. It is a gas. It has a peculiar penetrating smell, burns with a blue flame, and reddens vegetable blues. It forms, with hydrogen, prussic acid (110).

151. SULPHURETS.—Bodies of this class are very numerous, and some of them very important. All the metals appear capable of forming sulphurets. Copper pyrites, and iron pyrites, are two native compounds of metals with sulphur. Alkalies and earths, by combining with sulphur, likewise form sulphurets.

152. PHOSPHURETS.—Compounds of phosphorus with metallic, and other bases. Phosphuret of sulphur is employed to form matches for the instantaneous production of light (see 584.) Phosphuret of lime is remarkable for having the property of decomposing water, at the common temperature of the atmosphere: when a piece of it is thrown into water, the hydrogen that is liberated, unites with a portion of phosphorus, and forms phosphuretted hydrogen gas, which takes fire spontaneously (see 585).

SALTS.

153. These are compounds of acids, in definite proportions, with alkalies, earths, and metallic oxides. They form a very numerous, and very important class of chemical bodies; and the study of their composition and properties, must occupy a large share of the learner's attention. Nothing can give such clear ideas concerning the nature of chemical changes, so well explain the laws

f affinity, and exhibit to advantage the doctrine of definite proportions, as the synthesis and analysis of salts. There is another thing also which may be mentioned here, as it serves to show with what facility some parts of chemical knowledge may be acquired. The student, upon being told that the number of salts amounts to several thousands, might be terrified into the opinion, that it would be scarcely possible to remember even the *names* of so many bodies. But when he is shown how philosophically these bodies have been arranged, and how excellently named, he will perceive, that the learning and remembering—not only their names, but their constituent principles also, is a task of but little difficulty. To illustrate this.

154. Let the student be told, that every salt has a double name : one part of it intimating the acid of the compound, and the other, the particular base. Let him also be told, that the terminations of the names of the salts invariably agree with the terminations of the names of their components ; and then the nomenclature of salts will be perfectly intelligible. Thus, if an acid, whose name ends in IC, as sulphuric acid, combines with a base, as soda, and forms a salt, the part of the name of that salt which intimates its acid constituent, will end in ATE ; the compound being a *sulphATE*. But, on the contrary, if the salt be formed by an acid ending in OUS, as sulphurous acid, then the name of the salt ends in ITE, as *sulphITE of soda*.—A few more examples may render this familiar :—

ACIDS.	BASES.	SALTS PRODUCED.
Sulphuric.	Lime.	Sulphate of Lime.
Sulphurous.	Lime.	Sulphite of Lime.
Nitric.	Potass.	Nitrate of Potass.
Nitrous.	Potass.	Nitrite of Potass.

The salts which end in *ite*, like the acids they are formed from, are generally of very little value.

155. When the proportions of the constituents of a salt are so adjusted, that the resulting substance does not exhibit vegetable colours, it is called a *neutral salt*. When an alteration of colour evinces a predominance of acid, the salt is said to be acidulous, and the prefix *super*, or *bi*, is used to denote the excess of acid. If, on the contrary, the acid matter appears to be in defect, or short of the quantity necessary for neutralizing the properties of the base, the salt is then said to be with excess of base, and the prefix *sub*, is attached to its name.

156. It is to be remembered, that all salts are combinations of metals with acids, and that the bond of union between the two bodies is oxygen. If a piece of pure metal is put into an acid, it remains unacted upon, until, by some means, it acquires oxygen, and is converted into a metallic oxide: *then* its dissolution immediately commences. But whence does pure metal, when put into an acid, obtain the oxygen necessary for its conversion into an oxide?—To this question must be given the following answer: The metal, previous to its dissolution in an acid, decomposes either part of the acid itself, or else the water with which the acid is diluted. It then attracts to itself the oxygen of the decomposed liquid, and sets the other constituent of it at liberty. It will be worth while to consider the chief phenomena of metallic dissolution more closely, in order to show, that this is a right explanation of what takes place.

157. The first and most striking of these phenomena, is the effervescence which takes place: what is this occasioned by?—The effervescence itself is a disengagement of gas; but what is this gas disengaged from? Let us consider nitric acid. We may suppose that the dissolution is taking place in nitric acid. Now, we know, that the constituent principles of nitric acid (oxygen and nitrogen) can only exist when separated from each other, or when united in other proportions, in the state of gas; we also know, that nitric acid imparts its oxygen to other bodies with great readiness. Hence it is easy to infer what takes place when this acid has a metal put into it: a portion of the acid is decomposed; part of the oxygen of this decomposed portion of acid combines with the metal, another part combines

th the other element of the acid, (nitrogen,) and constitutes a body *not* soluble in water. This newly-formed body is *nitrous gas*, and this it is, which, by its sudden conversion from the liquid to the gaseous state, produces its disengagement, the noise and frothing which is termed effervescence.

158. The same decomposition, and consequent formation of gas, takes place when solutions of metals are made with other acids, whether they be diluted or concentrated. In solutions with sulphuric acid, the disengaged aëriiform body is either sulphurous acid gas, or hydrogen gas, according as the oxidation of the metal happens to be made at the expense of the sulphuric acid, or of the water.

159. The second observable phenomenon is, that when the metal has been previously oxidised, its dissolution takes place *without effervescence*—and why?—because, not needing oxygen, the metal does not decompose the acid, or water, to obtain it, and no gas, therefore, is set at liberty. The earths, and alkalies, dissolve without effervescence, for this reason.

160. Another phenomenon worthy of notice, is, that all metals which have a weaker affinity for oxygen, than the bases of certain acids, are absolutely insoluble in those acids. Hence iron, and some other metals, are, in their metallic state, insoluble in sulphuric acid; but, if those metals are put into the acid in an oxidised state, they become readily dissolved, without effervescence; or, if the acid be diluted with water, to oxidise the metals by its decomposition, they are then dissolved *with* effervescence.

161. The solubility of salts in water, is their most important general habitude. In this menstruum they are usually crystallized; and, by its agency, they are purified and separated from each other, in the inverse order of their solubility. It is not very difficult to determine the quantity of a salt, which water can dissolve. Water, at a certain temperature, must be saturated exactly with the salt whose solubility we wish to know; a certain quantity of this solution must be weighed out and evaporated, and the saline residue, upon being weighed, gives the amount. If, for example, 3 parts of the saturated solution,

upon being evaporated, leave 1 part of salt; then, the salt is said to be soluble in 2 parts of water, at the given temperature.

162. To obtain a perfectly saturated saline solution, heat the water with the salt, and then allow it to cool at the temperature required. In very particular cases, it is necessary to keep the final temperature constant for at least two hours, and to stir the solution frequently, to insure of its saturation.

163. The following pages contain descriptions of some of the most important salts. They will be found very interesting, and will repay the student for the closest attention.

164. NITRATE OF POTASS—SALTPETRE.—This well-known salt is a compound of nitric acid and potass. It crystallizes, and its crystals contain very little water of crystallization. Its component parts are 47 base, and 53 acid. Its solution requires four times its weight of cold water, but only an equal weight of boiling, water. If a red heat be applied, it is decomposed, in consequence of the destruction of its acid. By distilling in an earthen retort or in a gun barrel, oxygen gas may be obtained in great abundance, one pound of nitrate of potass yielding about 12,000 cubic inches, sufficiently pure for common experiments. This salt powerfully promotes the combustion of inflammable substances: it is the principal ingredient in the composition of gunpowder. Besides the uses already mentioned, nitre enters into the composition of fluxes, and is extensively employed in metallurgy: it serves to promote the combustion of sulphur in fuming, by eating its acid; it is used in the art of dying; it is added to common salt for preserving meat, to which it gives a red hue; it is an ingredient in some frigorific mixtures, and is frequently prescribed in medicine. It is also from this body that the nitric acid of commerce is generally obtained.

165. CHLORATE OF POTASS, a salt of a very singular nature, composed of chloric acid and potass, or, according to some, of chlorine 1 part, potassium 1 part, and oxygen 6 parts. Its properties are the following:—It crys-

lizes in shining scales. One part requires for solution 17 of cold water, or $2\frac{1}{2}$ of hot. Its taste is cooling and unpleasant. When one hundred grains of this salt are exposed to a gentle red heat, they yield 115 cubic inches of very pure oxygen gas. It is decomposed by all the strong acids. It exerts very powerful effects on inflammable bodies: The mode of preparing and performing experiments with this body, will be found described among the other experiments.

166. CARBONATE OF POTASS.—This salt has been long known. There are two varieties of it. The first is when the alkali is fully saturated with carbonic acid; the second is the potass of commerce, which is in the state of a sub-carbonate—the alkali being in excess. The sub-carbonate is composed of carbonic acid 32 + potass 68; the carbonate, of equal portions of the two bodies. The sub-carbonate has a taste far less penetrating and corrosive than pure alkali; but it still turns to green, the blue infusion of vegetables. Upon the addition of an acid, carbonic acid gas is given out with effervescence. It dissolves very readily in water, which, at the ordinary temperature, takes up more than its own bulk. When exposed to the atmosphere, it attracts so much moisture, as to pass rapidly to a liquid state. This change is termed deliquescence.—The carbonate of potass differs from the sub-carbonate in the following particulars:—In the greater mildness of its taste. Though still alkaline, it is but slightly so. It is composed of carbonic acid 64 + potass 68. It is unchanged by exposure to the air. It assumes the shape of regular crystals. It requires, for solution, four times its weight of water at 60° ; but boiling water dissolves nearly an equal weight; the heat, however, in this case, drives off a portion of carbonic acid gas.

167. PRUSSIAN OF POTASS.—This salt is composed of prussic acid and potass. It is one of the most valuable re-agents which the chemist possesses, in metallic analysis. The crystals of this salt are transparent, and of a beautiful lemon, or topaz yellow colour. It has a saline, cooling, but not unpleasant taste. In large crystals, it possesses a kind of toughness, and, in thin scales, of elas-

ticity. Water, when cold, dissolves one-third of its weight of this salt, and an equal weight at the boiling point. Its solution is not affected by alkalies ; but it is decomposed by almost all the salts of the permanent metals,—the alkali combines with the acid of the metallic salt, and the prussic acid with the precipitated metallic oxide, to which according to the nature of the metal, it communicates a peculiar colour. Hence the use of prussiate of potass as a test. The following table presents a view of the colours of some of the metallic precipitates thus obtained :—

<i>Solutions of</i>	<i>Give a</i>
Manganese,	White precipitate.
Protoxide of iron,	Copious white.
Deutoxide of iron,	Copious clear blue.
Tritoxide of iron,	Copious dark blue.
Tin,	White.
Zinc,	White.
Cobalt,	Grass green.
Bismuth,	White.
Protoxide of copper,	White.
Deutoxide of copper,	Crimson brown.
Lead,	White. [in the air
Silver,	White, passing to blue
Gold,	None.

168. OXALATE OF POTASS, sometimes called *essentielle salt of lemons*, exists in two states : one, the super-oxalate is obtained from the leaves of wood-sorrell ; the other the oxalate, may be obtained by adding more potass to the super-oxalate. Both may be formed by the direct union of potass with oxalic acid, certain cautions being observed. Oxalate of potass is a crystallizable salt ; but it does not readily crystallize, unless one of the ingredients be in excess. The crystals are nearly insoluble in cold water, but will dissolve in 10 parts of boiling water. The taste of this salt is sharp, acrid, and bitter. It is used for removing stains from linen, and for various purposes of the arts.

169. TARTRATE OF POTASS.—There are two varieties

of this salt, the tartrate, commonly called *soluble tartar*; and the super-tartrate, well known by the name of *cream of tartar*. 1. The super-tartrate of potass is composed of tartaric acid, 18 + potass 6. It is found encrusted on the bottoms and sides of casks in which wine has been kept. The crust is purified, and the salt is procured from it in crystals. This dissolves in 60 parts of cold water, or 14 of boiling. It is much used in chemical and medicinal preparations. 2. The tartrate of potass is composed of tartaric acid, 9 + potass 6. This salt has a bitter taste. It is used in medicine as a mild purgative. It is dissolved by its own weight of cold water, and by a still smaller quantity of hot water. It is decomposed by heat.

170. MURIATE OF SODA—CULINARY SALT.—This abounds in the waters of the ocean, and chiefly occasions their peculiar taste. It is likewise found, in great abundance, in various parts of the globe, in a dry state. At Northwich, in England, is a salt mine, which yields 4000 tons of rock-salt a-year; and, in Spain, there is a remarkable solid mountain of salt, 500 feet high, and a league in circuit; its depth below the surface of the earth is unknown. This mountain is composed of salt, in a state of perfect purity. In short, salt is one of the most abundant bodies in nature. And it is a very useful body. The shape of its crystals is a regular cube. Its taste is well known. It dissolves in $2\frac{1}{2}$ parts of water, whether it be cold or boiling. It is not affected by exposure to the atmosphere. When heated, it breaks with a crackling noise, which phenomenon is termed *decrepitation*. By a greater heat it may be fused, and by a still greater volatilized; but heat cannot decompose it, for, after volatilization, it remains muriate of soda. It is composed of muriatic acid, 14 + soda 16; or, if it be regarded as a compound of chlorine and sodium, of chlorine 18 + sodium 12. The muriatic acid of the shops is obtained from this salt.

171. SULPHATE OF SODA—GLAUBER'S SALT.—A compound of sulphuric acid 19 + soda 15. When crystallized, this salt is composed of 19 acid, + 15 base, +

44 water of crystallization. It is found native in various saline springs, and sometimes efflorescent on the walls of old houses. It may be formed artificially by the combination of its constituent parts. The taste of this salt is cooling and bitter. Its crystals, when in contact with the atmosphere, lose their water of crystallization, and become pulverulent. It is soluble in three parts of cold water, and less than its own weight of boiling water; a saturated hot solution, when cooled, readily shoots into beautiful crystals. The crystals, on exposure to fire, liquify in a short time; and, when an intense heat is applied, fuse. Sulphate of soda is decomposed by barytes, strontia, and potass; the first two take its acid from it entirely, the last only partially. It is generally used in medicine, as a purgative.

172. BORATE OF SODA—BORAX.—Neutral borate of soda is an artificial production, not much known; but the sub-borate of soda, or common borax, is a body that is well known, and very useful. It is brought from the East Indies in an impure state, and called *tincal*. It is purified by being boiled very strongly, and for a long time. It is white, transparent, has a greasy feel, and a styptic alkaline taste; it converts vegetable blues to green; it is soluble in 6 parts of boiling water, and 18 of cold water. When exposed to heat, it swells, boils, loses its water of crystallization, and becomes converted into a porous, white, opaque mass, termed calcined borax. With a stronger heat it melts into a transparent glass, but remains soluble in water. The component parts of sub-borate of soda, are boracic acid 36, soda 17, water 47. It is much used as a flux for metals, and is therefore of great importance in analysis by the blow-pipe.

173. PHOSPHATE OF SODA.—The taste of this salt is very similar to that of common salt. Its crystals are soluble in 2 parts of boiling water, or 4 of cold water: they effloresce in the air, and when heated, undergo the watery fusion. At a red heat they melt into a white enamel, and before the blow-pipe into a transparent globule, which, on cooling, becomes opaque. It is much used by mineralogists in the fusion of minerals.

174. CARBONATE OF SODA.—This compound of carbonic acid and soda, is known in commerce by the names of *kelp*, *barilla*, and *soda*, and is obtained either from the decomposition of common salt, or from the combustion of marine plants. This salt, like the carbonate of potass, exists in two states—1. The *carbonate*, properly so called, in which the soda is fully saturated with carbonic acid, (composition 20 acid + 14 base.) 2. The *sub-carbonate*, in which the soda is in excess, (composition, in its dry state, 10 acid, 14 base; composition, when crystallized, 10 acid + 14 base, 43 + water of crystallization.) The latter sort is the one generally found in commerce, and known by the above-mentioned names.

175. Carbonate of soda is found native, and may be formed artificially. It is not a crystallizable salt, but is obtained in the form of a white solid mass, which is so little liable to change by exposure to air, that it has been used in the construction of walls of edifices, instead of stone.

176. Subcarbonate of soda, crystallizes. Its taste is urinous, and slightly acrid, without being caustic. It changes blue vegetable colours to a green. It is soluble in twice its weight of cold water, and less than its weight of boiling: hence a saturated hot solution crystallizes on cooling; the crystals, are, however, more beautiful, when the evaporation is carried on slowly. It is one of the most efflorescent salts known, falling completely to powder in a very little time after its exposure to the atmosphere. On the application of heat, it is soon rendered fluid, is dried by the continuance of the heat, and then melted. As it is more fusible than the carbonate of potass, it is preferred to that salt in the manufacture of glass; because it promotes the fusion of the earth in a greater degree, and produces a glass of better quality.

177. NITRATE OF AMMONIA—*Inflammable Nitre*.—This is a combination of nitric acid and the volatile alkali. It is obtained in crystals, (though its components cannot be crystallized,) which differ in their appearance, according to the degree of heat employed to evaporate the water of solution. The taste is extremely cold and acrid. It dis-

solves in 2 parts of cold water, and in half its weight of boiling water. It deliquesces. The most remarkable property of this salt, is, that, when thrown on a red hot iron, it explodes, with a loud noise, accompanied by a white flame. By this it is decomposed: the oxygen of the acid forms water, by combining with the hydrogen of the alkali; and the nitrogen of both bodies is dissipated in the shape of gas. The chief use of nitrate of ammonia, is for affording nitrous oxide gas, by a certain mode of decomposition. When dry, it consists of 6.75 acid, 2.13 ammonia, 1.125 water.

178. MURIATE OF AMMONIA—*Sal Ammoniac*.—This salt was brought to this country for a long time from Egypt; but it is now economically prepared at home. This salt is purified by sublimation; hence, heat does not decompose it. When sublimed, it is in cakes, which are remarkable for possessing a certain degree of ductility, so as not to be easily pulverable. But it may be obtained from its solution, (which requires $3\frac{1}{2}$ parts of cold, or little more than 1 of boiling, water,) in regular quadrangular crystals. The compact salt is not affected by air, but the crystals deliquesce. It has a pungent, acrid, cooling taste, and the white smoke in which it sublimes has a very peculiar smell. Great cold is produced, both by its solution in water, and its mixture with pounded ice: hence, here, it is useful for frigorific mixtures, and, in warm countries, for cooling wines. It is composed of 48 acid, 16 base, 37 water. Few salts are of more extensive use than this. In chemistry, medicine, metallurgy, dying, and many other arts, it is much employed; and it is indispensable in operations whereby the surfaces of some metals are coated with others—for example, in the tinning of copper.

179. CARBONATE OF AMMONIA, vulgarly called *smelting salts*, when very pure, is in small crystals. The taste and smell of this salt are the same with those of pure ammonia, but much weaker. It is soluble in rather more than twice its weight of cold water; and in its own weight of hot water; but a boiling heat volatilizes it. It is not decomposed, however, by heat, though so readily

sublimed. It is soluble in the atmosphere: if left in a state of exposure, it diffuses its smell, and diminishes in weight. It is prepared in different ways; and as these vary, so do its component parts. The salt of the shops consists of 55 acid, 30 base, 15 water. The solid salt produced by the union of its constituents in the state of gas, is composed of 56 acid + 44 alkali.

180. SULPHATE OF BARYTES.—This salt may easily be, though it seldom is, formed by art; because it abounds in nature, especially in veins of metals. It is generally obtained in crystals, differently shaped. It is the heaviest of all salts, and is insoluble in water. It is tasteless and inodorous. It is *not poisonous*,—this property distinguishes the sulphate from all the other salts of this earth, as they are poisonous. When formed into a thin cake with flour and water, and exposed for sometime to a very strong heat, it shines afterwards in the dark. This property was discovered in Bologna, and hence the mineral has been termed *Bolognian phosphorus*. It is composed of sulphuric acid 1 + barytes 2.

181. CARBONATE OF BARYTES.—This is found native, but not so abundantly as the sulphate of barytes. It crystallizes, is very heavy, tasteless, inodorous, extremely poisonous, and nearly insoluble in water. It is not altered by air, but is decomposed by heat. Its constituents are 2 acid + 7 base.

182. NITRATE OF BARYTES.—This is obtained by processes hereafter described, either from the native carbonate, or sulphate of the earth. Its taste is hot and acrid. It is soluble in 12 parts cold, and in 4 parts boiling, water. It crystallizes, sometimes in stars, sometimes in small brilliant plates. When heated, it decrepitates: if the heat is continued, the acid is expelled, and the barytes remain pure. This salt is composed of 6.75 acid + 9.75 base. It is chiefly used as a *test* for sulphuric acid, for which purpose it is exceedingly well suited, as is shown at 535.

183. MURIATE OF BARYTES.—The observation prefixed to nitrate of barytes, respecting the mode of obtain-

ing that salt, applies to this also. Muriate of barytes crystallizes. It is soluble in 5 parts of cold water, in still less of hot water, and also in alcohol. It is not altered by air, and only partially decomposed by heat. It has a nauseous burning taste, and is poisonous; but is sometimes given medicinally in small doses. It consists of 6 acid + 19 base, in a dry state; and 5 acid, 16 base, 4 water, when crystallized. The muriate, like the nitrate of barytes, is of great use as a *test* for sulphuric acid, see 535.

184. SULPHATE OF STRONTIA bears, in many of its properties, considerable resemblance to sulphate of barytes; but its specific gravity is less, and its affinities are not so energetical. It occurs native, in great abundance in various parts of the world, usually in a crystallized form. It is insoluble. Its composition is 5 acid + 6½ base.

185. CARBONATE OF STRONTIA is found native, but as well as the preceding salt, may be formed artificially. The native substance appears in semi-transparent crystals, of a white colour tinged with green. It is insipid; insoluble; is not altered by exposure to the air; but is decomposed by heat. If thrown in powder on well kindled coals, it exhibits red sparks. It is much lighter than carbonate of barytes, (with which body it was long thought identical,) and is not poisonous. It consists of 2.75 acid + 6.5 strontia.

186. NITRATE OF STRONTIA may be obtained in the same manner as that of barytes, with which it agrees in the shape of its crystals, and most of its properties. It is much more soluble, however; requiring but an equal weight of water at 60°, and half its weight of boiling water. In dry air, it effloresces; in moist air, deliquesces. By heat it is decomposed. Applied to the wick of a candle, or added to burning alcohol, it gives a deep red, or purple colour to the flame. This is the characteristic property of strontia. It consists of 6.75 acid + 6.5 strontia.

187. MURIATE OF STRONTIA, formed artificially; tastes cold and sharp; soluble in less than its own weight of

old water ; and in any quantity of boiling water. It deliquesces in moist air. Its constituents are 24 acid + 36 base + 40 water.

188. CARBONATE OF LIME forms a considerable part of the solid strata of the globe. It is sometimes obtained pure ; but generally mixed with other earths, with metals, coal, &c. : it is occasionally obtained in transparent crystals. In these various states it is known by the names of *chalk, marble, limestone, stalactites*, &c. With the exception of muriate of soda, it is of more importance than any other of the salts. Carbonate of lime has no taste, is insoluble in water, but slightly soluble (like the other carbonates) in water impregnated with carbonic acid. But, by exposure to the air, the extra dose of carbonic acid flies off, and the salt reverts to its state of insolubility. No alteration is effected in it, on exposure to the air. When heated, it decrepitates, its water flies off, and lastly its acid ; but this requires a pretty strong heat. By this process it is burned into lime. It is composed of 14 lime + 11 carbonic acid. The specific gravity of this body varies, as does its state of aggregation : chalk and marble are essentially the same in constitution, but very different in their relative weights. Perhaps it may be stated at 2.7.

189. SULPHATE OF LIME.—This is known also by the names of *alabaster, gypsum, plaster of Paris*, and various others. It is found native in great abundance ; sometimes in a crystallized state, sometimes otherwise. It requires for its solution 500 parts of cold water, and 400 of hot. At a red heat it loses its water of crystallization, and falls into a white powder. When this powder is mixed with water, heat is disengaged, and the product is a solid mass. In this case, the water loses its heat of liquidity, by entering into a state of solidity with the calcined salt. A hasty crystallization takes place, with an enlargement of bulk. Owing to this property, calcined sulphate of lime has been much employed in making casts from anatomical and ornamental figures. It also forms one of the bases of *tucco*, is used as a cement, and for making various mouldings and other ornaments in building. Its consti-

tuents (when deprived of its water of crystallization) are 7 lime + 10 sulphuric acid.

190. FLUATE OF LIME.—This body is very abundant in the native state, and is called fluor spar, Derbyshire spar, phosphoric spar, &c. The substance found in Derbyshire is a very beautiful variety, and is much employed in the formation of various ornaments for cabinets, and on cups, candlesticks, &c. It is insoluble. When heated it decrepitates, loses its water of crystallization, and afterwards shines in the dark. At a strong heat, it melts into a glass. Several acids, when assisted by heat, decompose it, and expel the fluoric acid in white fumes. The constituents of fluate of lime are acid 16 + base 44 + water of crystallization 60.

191. NITRATE OF LIME.—This salt is formed in the operations attending the crystallizing of nitrate of potash of commerce. Its crystals resemble bundles of needles diverging from a centre. Its taste is hot and bitter. It deliquesces in the air, and is so extremely soluble, that it can only be kept in the dry state with great difficulty. It is soluble in an equal weight of alcohol. Nitrate of lime is of very little use. Its constituents are 45 acid + 2 lime + 35 water of crystallization.

192. MURIATE OF LIME.—This salt exists in nature but neither very abundantly, nor very pure. It is easily formed artificially, but very difficultly crystallized: it being one of the most deliquescent salts known. Its taste is acrid, bitter, and very disagreeable. It is soluble by heat in its own water of crystallization, and in half its weight of cold water. It dissolves in alcohol; during the solution, so much heat is generated, that the alcohol boils. Muriate of lime is chiefly used in frigorific mixtures; and with snow it produces a very great degree of cold. Its constituents, when deprived of its water of crystallization, are 11 acid, 11 lime.

193. PHOSPHATE OF LIME.—This is the basis of the bones of animals, and the principle which gives to them their hardness and strength: hence it is a body of great importance. It is not only found in bones, however, but

in considerable abundance in a native mineral state, and it may be formed artificially. It is sometimes obtained crystallized, but is generally in the form of a white powder, which is insoluble in water. By a violent heat, it may be fused into an enamel. It is used for making cupels, which are vessels employed by silversmiths in the operation of refining silver; for polishing gems and metals, to absorb grease from linen, paper, and silk, and as a medicine. It is identical with the burnt hartshorn of the shops. Its constituents are 30.5 acid, 69.5 base.

194. OXALATE OF LIME is a white powder, which is insoluble in water, not decomposable, except by fire, and turning vegetable blues green. It is composed of 14 acid + 11 lime.

195. SULPHATE OF MAGNESIA—*Epsom-Salt*.—This salt obtained the latter name, from having been obtained for a long time by evaporating the water of the saline springs at Epsom, in England. It is now generally obtained from sea water, in which it exists in large quantities, mixed with muriate of soda. It may be formed artificially. When pure, it crystallizes. Its taste is cool and bitter. It is very soluble, requiring only an equal weight of cold water, and three-fourths its weight of hot. It effloresces in the air, though but slowly. When heated, it dissolves in its water of crystallization, but is not decomposed. It is chiefly used as a purgative in medicine, and to furnish magnesia by its decomposition. It is composed of 5 acid + 2.5 magnesia + 7.9 water of crystallization.

196. CARBONATE OF MAGNESIA exists in two states: one in which the constituents are neutral, the carbonate; another, in which the acid is in defect, the sub-carbonate. This latter kind is the same as the *magnesia* of the shops; what is really simple magnesia, is commonly termed *calcined magnesia*. The methods of preparing both kinds of the salt will be described hereafter. The carbonate crystallizes, and its crystals are soluble in 48 times their weight of cold water; the sub-carbonate is a white powder, and requires for its solution ten times that quantity of water. The crystals slightly effloresce, and, upon exposure to the fire, decrepitate, lose their water and acid,

fall to powder, and are reduced to one-fourth of their original weight. The powder, upon being heated, boils; its acid flies off, it loses half its weight, and the magnesia remains quite pure.

197. SULPHATE OF ALUMINA AND POTASS—ALUM.—

This salt is sometimes found in a native state, but is generally formed from various minerals, called alum ores. It crystallizes, has a sweetish, but very astringent taste, and reddens vegetable blues, showing the acid to be in excess. It is soluble in from 15 to 20 parts of cold water, and in three-fourths its weight of hot. Its crystals effloresce, and, when heated, melt in their water of crystallization; at a strong heat, they lose 44 per cent. (chiefly water) of their weight, and are reduced to a white porous mass called *burnt alum*. This salt is a very important one: it is extensively used in medicine, bleaching, dying, calico printing, tanning, paper-making, and various other arts. Bodies which have been steeped in a solution of alum, are not apt to take fire. Its composition is stated thus: acid 10, alumina 3, potass 3, water of crystallization 16.

198. METALLIC SALTS.—This term is applied to the bodies which are formed by the combination of acids with those metals, that are better known in their metallic states than in their oxidised. It might certainly, with strict propriety, include the alkaline and earthy salts, for *there* are as much metallic salts as any others are, (as will be evident to any one who reads paragraphs 156 to 160;); but the term is, however, generally prescribed to what we have stated. The following are among the most curious and useful of the metallic salts.

199. NITRO-MURIATE OF GOLD.—No simple acid, however strong it may be, has the power of dissolving gold; but a compound, made by mixing one part of nitric acid with two of muriatic, *does* possess this property. This menstruum was formerly called *aqua regia*, but is now more properly named nitro-muriatic acid. On mixing the two acids, heat is given out, an effervescence takes place, and the mixture acquires an orange colour.

A saturated solution of nitro-muriate of gold, is of a deep yellow colour, very caustic, and of an astringent

metallic taste. It tinges the skin, and almost all animal and vegetable bodies, and even marble, of a deep and indelible purple. It crystallizes, its beautiful golden colour being preserved. The crystals are very soluble in water, and still more so in ether. Decomposition of this salt is effected by light, heat, earths, alkalies, and several metals; as in another part of this work will be proven.

200. NITRATE OF SILVER.—Nitric acid dissolves half its weight of silver. The solution is limpid, colourless, heavy, and caustic. It conveys to hair, the skin, and most animal substances, an indelible black colour. It forms brilliant transparent crystals, possessed of a very bitter metallic taste, and being very soluble in water; not deliquescent, but decomposed by light; converted by heat into a preparation, used in medicine, under the name of *lunar caustic*, which is a sub-nitrate. It enters into the composition of several detonating or fulminating compounds. It is composed of acid 1 + base 2. It is decomposed by nearly the same bodies, and in the same manner, that nitro-muriate of gold is. The method of forming nitrate of silver is shown at 494.

201. MURIATE OF SILVER is an insoluble salt, formed by adding muriatic acid, or compounds containing it, to nitrate of silver. It cannot be decomposed by heat, for, after melting, it rises in vapour; but the silver may be reduced by a process, which will be hereafter described. On account of the immense affinity which muriatic acid has for silver, these two bodies serve as excellent tests for each other. See 516. The composition of muriate of silver is acid 1 + silver 3.

202. NITRATE OF MERCURY (495) is colourless, heavy, and very caustic: it tinges the skin of an indelible black colour, and yields transparent crystals, which are soluble in hot water. It is decomposed by sulphuretted hydrogen gas, by alkalies, and by muriate of tin. It is the basis of a fulminating compound. See 612.

203. NITRATE OF COPPER is very caustic. It is easily formed, see 496. Its composition is 37 acid + 31 base + 32 water. Its crystals are of a beautiful blue colour; deliquescent, soluble, and decomposed by heat. This

salt, from the facility with which it parts with oxygen, is capable of acting on several substances with much energy, see 601. Nitrate of copper is decomposed by the alkalis, by iron, and by several other bodies.

204. SULPHATE OF COPPER is a regularly crystallized salt, easily dissolved by water. It is of a beautiful deep blue colour, and hence obtained the name of *blue vitriol*. It is a salt that is of great use in several of the arts. It is caustic, reddens vegetable blues, and effloresces, when exposed to the air. It is decomposed by heat, by pure and carbonated earths and alkalies, and by several metals. It is composed of acid 32 + copper 32 + water 36.

205. MURIATE OF COPPER is of a beautiful deep green colour, and very acrid. It may be formed by corking up, in a small phial, some copper clippings and a little muriatic acid: after a few weeks, a white powder will be seen at the bottom of the phial, and the liquid acid will have acquired the above-mentioned colour. The white powder is oxide of copper precipitated from solution; the green liquid is a solution of the salt. Its constituents are 40 black oxide of copper + 24 acid + 36 water.

206. ACETATE OF COPPER.—When plates of copper are exposed to the vapour of vinegar, a substance is formed, of a blueish green colour, used in dying and painting, and called *verdigris*. This substance, when dissolved in acetic acid, forms acetate of copper. This salt, like the sub-acetate, (*verdigris*,) is blueish-green; its crystals are soluble in 5 parts of hot water, but scarcely at all in cold. They effloresce in the air, but are decomposed by heat, yielding pure acetic acid. Acetate of copper is a violent poison, like all other bodies of which this metal forms a part. But we possess an antidote of undoubted efficacy to this poison—this is *sugar*. The chief use of this salt is in dying.

207. SULPHATE OF IRON, the body so well known by the name of *green vitriol* and *copperas*. It forms beautiful transparent green crystals. Its taste is harsh and styptic. It reddens vegetable blues, the acid being always in excess. Two parts of cold, and three-fourths of boiling water, dissolve it. It is efflorescent. A moderate heat

drives off its water of crystallization, and a strong heat its acid. Sulphate of iron is of extensive use in the arts. Its constituents are 29 acid + 28 oxide of iron + 45 water.

208. MURIATE OF IRON.—The iron in this salt is sometimes in the state of a protoxide, at others, of a peroxide; and the salts are thence called promuriate, and permuriate.—The former is of a green colour, crystallizable, and very soluble. It absorbs nitrous gas. From it, alkalies throw down a green precipitate; prussiate of potass, a white one; and gallic acid, none at all.—The permuriate has a brown colour, is uncrystallizable, very soluble, and readily decomposed by heat: the addition of sulphuric acid expels chlorine; and alkalies throw down a yellowish brown precipitate.

209. CARBONATE OF IRON is the chief ingredient in *Chalybeate* mineral waters. It has a green colour, is brittle, somewhat transparent, and soluble in water. It possesses the singular property of becoming a permanent magnet, when heated red hot. Common *rust* of iron is a compound of carbonic acid, and peroxide of iron. It is not soluble in water.

210. ACETATE OF IRON crystallizes, has a green colour, and a sweetish styptic taste. A solution of it is much used by calico-printers, under the name of *iron liquor*, in dying black.

211. GALLATE OF IRON.—This salt constitutes black dyes and ink. It is formed by the union of gallic acid with peroxide of iron. It is decomposed by the stronger acids; hence ink-spots are taken from paper or linen by their application.

212. PRUSSIATE OF IRON, the bases of the beautiful pigment called *prussian blue*. The combined iron in this substance is in the state of a peroxide.

213. NITRATE OF TIN is a salt that is not permanent (though the acid acts upon the metal with amazing violence;) the tin continues to attract oxygen from the nitric acid, until it becomes too highly oxidised to remain soluble. It is accordingly precipitated in the form of a gray powder.

214. MURIATE OF TIN is of a yellow colour, and may be crystallized. The crystals are deliquescent. This salt is a test for several metals; especially for mercury and gold.

215. CARBONATE OF LEAD, commonly known by the name of *white lead*, is formed by exposing thin plates of lead, coiled up, to the vapour of vinegar. The metal is oxidised by the decomposition of the vapour of the acetic acid; and the salt is formed by the abstraction of carbonic acid, partly from the atmosphere, and partly from the acetic acid vapour. It is used as a white paint.

216. ACETATE OF LEAD.—*Sugar of Lead*.—This salt obtained the latter name, from its sweet taste. It is a deadly poison, like all other preparations of lead. It crystallizes in needles, which have a silky appearance. These are soluble in $3\frac{1}{2}$ parts of cold water, and something less of hot. Acetate of lead is of great utility. The constituents of this salt are 27 acid + 58.7 base + 14.3 water.

217. SULPHATE OF ZINC—the *White Vitriol* of commerce—a compound of some importance. It crystallizes. Its crystals are soluble in $2\frac{1}{4}$ parts of cold water, and less of boiling. Heated, they melt, lose their water of crystallization, and then their acid. Heated with charcoal, the zinc is reduced, it then rises in vapour, and inflames. This salt is used in dyeing, to deepen several red colours. It is decomposed by alkalis. Its constituents are 50 acid + 51 oxide of zinc, in a dry state, and 50 sulphuric acid + 51 oxide of zinc + 79 water, when crystallized. For the method of forming sulphate of zinc, see 430, 501.

218. NITRATE OF BISMUTH is colourless, yields white crystals, which effloresce and detonate weakly. When the crystals are put into water, they are decomposed; the oxide of bismuth falling down in a white powder, called *magistry of bismuth*, which is used as a paint for the complexion, though it is liable to turn black, by exposure to sulphuretted hydrogen gas. The *pearl powder* of the perfumers is an oxide of bismuth precipitated by cold water, from the nitro-muriate of the metal.

MINERALS.

219. The inorganic and inanimate substances, both simple and compound, and of whatever nature, that are found native, either on the surface, or in the bowels, of the earth. The totality of these bodies has received the name of the mineral kingdom.

The following arrangement of minerals is the one generally adopted :—

Class 1. Metallic minerals, or ores.

2. Earths and stones.

3. Saline minerals, or native salts.

4. Inflammable fossils.

From bodies of the first class, we obtain, by certain processes, *pure metals*, from those of class 2, *pure earths*; from those of class 3, various *salts*, but chiefly nitrates, muriates, sulphates, and carbonates; and from bodies of the 4th class, such substances as *sulphur* and *bitumens*.

The study of what relates to mineral bodies, properly constitutes the science of *mineralogy*. The subject, though interesting, is one which we are not able to do more than just touch upon. From what is said above, and from the *analyses of ores, stones, marls, soils, salts, &c.* to be found among the experiments described hereafter, the student will obtain as much knowledge of mineralogy as will be sufficient for him, till he advances further in other divisions of chemistry.

MINERAL WATERS.

220. This is the term applied to the waters of certain springs, which, because they possess peculiar tastes, smells, colours, and other singular properties, cannot be applied to purposes of domestic economy. But, though these waters cannot be used for what water in general is used, yet they are not altogether unserviceable. Mineral springs have been resorted to in all ages for the cure of various diseases; and some of them have effected wonderful cures in several obstinate disorders.

The examination of mineral waters with a view to ascertain their ingredients, and thence their medical pro-

perties, and the means of compounding them artificially, is an object of considerable importance to society. In the analysis of mineral waters, we, in the first place, by means of certain tests, ascertain what are the *ingredients*; and then, by a more minute process, determine the *proportions*. But the quantity of mineral substance which gives to water those peculiar properties, is sometimes so very small, and the ingredients so various, that few problems in Chemistry are attended with more difficulty than the correct analysis of mineral waters. On this account, we shall be excused from going into the subject very deeply, and, therefore, instead of giving a formal account of *all* the substances that have *ever* been found in mineral waters, we shall mention merely those which are *generally* found in them; and furnish some pleasing experiments, to show the mode of detecting the presence of these substances. See SECTION 5. of the experimental part of this work.

NATURAL CHEMISTRY.

221. Having now considered the properties of the elementary substances, and such of their compounds as can be formed artificially, or are found in the mineral world, we proceed to examine the states of combination in which they occur in organic substances.

By *organic substances*, we mean those which constitute the vegetable and animal kingdoms: and which essentially differ from those composing the mineral kingdom inasmuch as, while the latter seem to be composed by the accidental combinations of their constituents, *their* constitution bears the most striking and impressive marks of design, and they are in possession of, and are governed by, the unknown, but vastly important principle called *life*. With the real *nature of life*, we are perfectly unacquainted; all we know of it regards its operations.

222. ON VEGETABLE SUBSTANCES.—The elementary bodies which enter into this class of compounds, are very few. Plants are found by analysis to consist principally of aëriform matter, and charcoal. They give out, by distillation, volatile compounds, which are composed of oxygen, hydrogen, nitrogen, and coally matter, or charcoal.

Yet, although these elements are few, the principles of life enables plants to form from them a great variety of new chemical arrangements, by means of appropriate organs. The various solid parts and juices of vegetables, thus composed, and existing ready-made in them, may, therefore, be regarded, as their immediate materials, or *proximate* constituents. The following are the most important of these substances:—

223. GUM.—Colourless, tasteless, and inodorous. Does not crystallize. Dissolves in water, forming mucilage. Insoluble in alcohol. There are several kinds of gum; the one most used in the arts is *gum arabic*. Gum exudes from growing trees.

224. SUGAR.—To be found in every vegetable. Is very abundant in, and easily obtained from, the plant called the sugar-cane. Taste sweet. Crystallizes. Soluble in water, alcohol, and nitric acid. When distilled with the latter, yields oxalic acid.

225. FECULA, or *Starch*.—Obtained from seeds and roots, particularly wheat and potatoes. An insipid white powder. Insoluble in cold water. Forms an opaque glutinous solution in hot water. Insoluble in alcohol.

226. GLUTEN.—Obtained from wheat flour. Forms a ductile elastic mass with water. Partially soluble in water, but insoluble in alcohol. It is this substance that gives to flour the property of forming good paste with water.

227. OILS are of two kinds, fixed and volatile; and both are obtained from plants. *Fixed* oils exist chiefly in seeds, and each oil bears the name of the plant from whose seed it is extracted; thus lintseed, rapeseed, almond oil. Sometimes a fixed oil is obtained from the pulp of fruit; olive oil is one that is. They are extracted by pressure and boiling, have no smell, are insoluble in water and alcohol, form soaps with alkalies, and are coagulated by salts.

228. *Volatile* oils are obtained from all parts of plants, *excepting* the seeds. Extracted by distillation and pressure. Strong peculiar smell to each individual oil. In-

soluble in water. Soluble in alcohol. Volatilized by heat.

229. CAMPHOR.—Obtained from a species of laurel tree. Analogous in its nature to volatile oils. Strong smell. Extremely volatile and inflammable. Insoluble in water and alkalies. Soluble in alcohol, oils, acids. Used in medicine.

230. RESINS.—These are volatile oils, peculiarly modified by the action of oxygen. Solid. Melt, when heated. Insoluble in water. Soluble in alcohol, ether, and alkalies. By nitric acid converted into artificial tannin. Pitch, tar, and turpentine, are the most common resins: they exude from trees. Copal, mastic, and frankincense, are also of this class of compounds.

231. GUM RESINS are precisely what their name denotes—resins combined with mucilage. Strong smell. Brittle, opaque, infusible. Form milky solutions with water; transparent with alcohol. Myrrh and assafœtida are of this class.

232. BALSAMS.—Peculiar resinous juices, combined with benzoic acid, which body sublimes, when they are heated. Strong smell.

233. CAOUTCHOUC. A milky glutinous fluid, which exudes from certain trees, and in drying turns black and acquires consistence; in this state it is called Indian-rubber. Very elastic. Insoluble in water and alcohol. Very combustible.

234. COLOURING MATTER.—The colours of vegetables are owing to certain matters, which may be extracted, and converted into dyes and pigments. The art of dying consists in extracting colours from vegetables, and fixing them on cloth.

235. TANNIN.—Taste astringent. Soluble in water. Forms a precipitate with gelatin, which is insoluble in water. This precipitate is formed when skins are tanned and made into leather. It is found abundantly in oak-bark and nut-galls, and may be formed (though not economically) artificially.

236. WAX.—Insoluble in water. Fusible. Soluble in alcohol, ether, and oils. Forms soap with alkalies.

237. WOOD.—Composed of fibres. Tasteless. Insoluble in water and alcohol. Yields much charcoal, (giving out impure acetic acid,) when distilled in a red heat.

238. VEGETABLE ACIDS.—The most important of these are the mucous, obtained from gum; the suberic, from cork; the camphoric, from camphor; the benzoic, from balsams; the gallic, from galls, &c.; the malic, from ripe fruit; the citric, from lemon and lime juice; the oxalic, from sorrell, and from bodies distilled with nitric acid; the succinic, from amber; the tartaric, from cream of tartar; the acetic, from vinegar. They are all decomposable by heat, and soluble in water. The other properties of the most useful amongst them, have been stated already (see 111 to 116.)

239. The *Decomposition of Plants* begins as soon as they are dead. The reduction of the various matters composing them to their simple elements, is a process of very considerable length; during which many new combinations are successively established, and successively destroyed. The process here spoken of is *fermentation*.

240. The *Vinous Fermentation* derives its name from its product, which is *wine*.—When saccharine vegetable matter, accompanied by some other bodies, as fecula, mucilage, &c., are exposed to water and heat, they shortly experience a very striking alteration. An internal commotion takes place; the mass grows turbid; carbonic acid gas is disengaged, in considerable quantity, and, being enclosed in viscid matter, forms, on the surface of the liquor, a stratum of *yeast*. After a time these appearances cease, and the fermented liquor becomes clear and transparent. It will now have lost its sweetness and viscosity, and have acquired the vinous taste and intoxicating quality. Thus *wine* is made from the juice of the grape, and thus *beer* (which may be chemically regarded as the wine of grain) is made from malt.

241. If the fermentation be stopped, by the exclusion of air, before the whole of the carbonic acid gas is evolved, the wine is brisk and sparkling, like Champagne, from the carbonic acid gas imprisoned in it; it is also

sweet, like cyder, from the sugar not being completely decomposed.

242. Sugar alone does not furnish wine, by undergoing fermentation ; for, when it is decomposed, its constituents are recombined into two new substances ; the one a peculiar liquid substance, called *alcohol*, or *spirit of wine* ; the other, carbonic acid gas.

243. It is to *alcohol* that fermented liquors owe their intoxicating qualities, and from those bodies it may readily be separated by distillation. Alcohol, when pure, is of a strong heating nature, very light, inflammable, volatile, and possesses a peculiar odour. It dissolves resins, volatile oils, camphor, and many other substances. It is composed of small quantities of carbon and oxygen, united to a considerable proportion of hydrogen. Of these elements (in different proportions though,) sugar too must be composed, of course. But, though sugar may be decomposed into alcohol and carbonic acid, it is not possible to compose sugar, by uniting carbonic acid and alcohol.

244. *Ether*, the lightest of all liquids, highly volatile, inflammable, and odorous, is alcohol, deprived of a portion of its carbon. The decarbonization of alcohol is effected by the action of strong acids on it ; and the ether differs in its nature, and is differently named, according to the particular acid that is made use of. Thus nitric acid with alcohol, produces *nitric ether*, and sulphuric acid, *sulphuric ether*.

245. The *Acetous Fermentation*, is that which produces *vinegar*. It is generally said, that, in this species of fermentation, the product is obtained from liquors which have already undergone the *vinous* fermentation ; indeed, *vinegar* is chiefly prepared from wines, but the *vinous* fermentation is not necessarily preliminary, for vinegar may be obtained from simple mucilage. Common vinegar may be purified and concentrated by distillation, and it is then called distilled vinegar, or acetic acid (111.)

246. The phenomena of *putrefaction* is occasioned by the last change, or final decomposition, which vegetables undergo. The putrefactive fermentation is a complete

analysis of vegetable substance, during which the constituent elements are all disengaged in the form of *gas*, with the exception of some fixed earthy products, which remain in the state of mould, and are used as a manure.

247. But the element *nitrogen* is one that is exceedingly favourable to putrefaction ; and substances which contain it, or to which it has been added, are decomposed very rapidly. Among the gaseous bodies which are evolved during this process, are those described in paragraphs 121, 141, 142, 143. These have all peculiar and disagreeable smells ; and from the mixture of these different odours proceeds the feter that accompanies putrefaction.

248. No substance can enter into the putrefactive fermentation, if unaccessible by moisture and air ; we can, therefore, by keeping bodies in a state of dryness, and where air has no access, hinder their putrifying.

249. ANIMAL KINGDOM.—The principal primary constituents of animal substances are nearly the same as those of vegetables ; but they have less carbon and hydrogen, and a far greater quantity of nitrogen and phosphorus. The most important of the immediate materials of animals, which are formed by the combinations of these substances, are *gelatin*, *albumen*, and *fibrin*. These three kinds of animal matter form the bases of all the various parts of the animal system : *viz.* solid, as the *skin*, *flesh*, *nerves*, *membranes*, *cartilages*, and *bones* ; and fluids, as *blood*, *chyle*, *milk*, the *gastric juice*, *saliva*, *tears*, &c.

250. GELATIN, or *jelly*, the chief ingredient in skin. It is obtained under the forms of glue, size, isinglass, jelly, &c. It is a viscid substance, very soluble in water, but not in alcohol ; insipid, and without smell ; when cold, it congeals into a cohesive tremulous substance, but liquifies again when heated. It forms the bases of soups, &c. The union of gelatin with tannin in a skin, constitutes leather.

251. ALBUMEN is the principal ingredient in the serum of blood, the white of eggs. It is miscible with cold water, but is coagulated by heat, which forms the best test of its presence. It is also coagulated by acids and alcohol ; but is dissolved by pure alkalies.

252. FIBRIN forms the basis of the muscular part of animals. It is an insipid and inodorous substance, having somewhat the appearance of fine white threads adhering together. It is insoluble both in water and alcohol.

253. Animal matter, although the most complicated of all natural substances, returns to its elementary state by one single spontaneous process, the *putrefactive fermentation*. By this, the substances forming its different parts are slowly reduced to the state of oxygen, hydrogen, nitrogen, and carbon; and thus the circle of changes through which these principles have passed, is finally completed. Quitting their elementary forms, they entered the vegetable system, thence passed to the animal kingdom, and from that return again to their original simplicity; soon to re-enter the sphere of organized existence.

ON
CHEMICAL OPERATIONS
AND
APPARATUS.

254. THE very prevalent notion, that "a laboratory, fitted up with furnaces, and expensive and complicated instruments, &c. &c. &c., is absolutely necessary for the performance of chemical experiments," is exceedingly erroneous. In fact, the truth of the matter is diametrically opposite to this opinion. "For general and ordinary chemical purposes, (says Dr. Henry,) and even for the prosecution of new and important inquiries, very simple means are sufficient: some of the most interesting facts of the science may be exhibited and ascertained, with the aid merely of Florence flasks, of common phials, and of wine glasses. In converting these to the purposes of apparatus, a considerable saving of expense will accrue to the experimentalist; and he will avoid the encumbrance of various instruments, the value of which consists in show, rather than real utility." It is a curious and instructive fact, that some of the most important discoveries in chemistry were made by persons who, either from choice, or motives of economy, used utensils of the very simplest character. The laboratory of the great Priestley cost a mere trifle; and it is well known how savingly Franklin went to work.

We intend, in this section of our work, to give an account of the chemical processes of most frequent recurrence, and to describe with minuteness the instruments whose real utility, simplicity, and cheapness, entitle them

to be so distinguished. The student will also be informed where he may obtain the different utensils, and how much they will cost him. We shall also notice the precautions necessary to be observed in conducting processes, or handling substances, that are in the least degree dangerous. In short, as we proceed with our instructions, we shall continually bear in mind, that the persons we are addressing both need and are inclined to receive information; that they have neither time to be lost, nor money to be thrown away,—and thus understanding the situation of our readers, we shall suitably model our discourse. We are not unaware, however, that while we proceed thus, we are laying ourselves open to the charge of puerility. We shall probably be told, that the mention of such trifling things is not befitting the pages of a philosophical treatise. But, being of opinion that elementary writers in general are by far too concise in their practical instructions,—that they presume the student knows many things which he really does not know,—we do humbly conceive, that we shall be serviceably employed, in giving precise directions for the performance of the most curious and instructive chemical experiments.

The following observations, by a writer of celebrity, on the conducting of chemical processes in general, are truly valuable and judicious. Method, order, and cleanliness, are essentially necessary in a chemical laboratory. Every vessel and utensil ought to be well cleansed as often as it is used, and put again into its place: labels ought to be put upon all glasses and boxes containing preparations. These cares, however, which seem to be trifling, are very fatiguing and tedious; but they are also very important, though frequently little observed. When a person is keenly engaged, experiments succeed each other quickly; some seem nearly to decide the matter, and others suggest new ideas: he cannot but proceed to them immediately, and he is led from one to another: he thinks he shall easily know again the products of the first experiments, and therefore he does not take time to put them in order: he prosecutes with eagerness the experiments which he has last thought of; and, in the mean time, the ves-

sels employed, the glasses and bottles, and products so accumulate, that he can no longer distinguish them; or at least, he is uncertain concerning some particular product. This evil is increased, if a new series of operations succeed, and occupy all the laboratory, or if he be obliged to quit it for some time, for then every thing goes into confusion. Thence it frequently happens, that he loses the fruits of much labour, and that he must throw away almost all the products of his experiments.

When new researches and inquiries are made, the mixtures, results, and products of all the operations ought to be kept a long time, distinctly labelled and registered; for these things, when kept some time, frequently present phenomena that were not at all suspected. Many fine discoveries in chemistry have been made in this manner; and many have certainly been lost, by throwing away too hastily, or neglecting the products.

The great agents whose aid the chemical analyst most frequently needs, are *fire* and *water*. When a process requires the first chiefly, it is said to be executed in the *dry* way; but when the latter is employed, the operation is performed in the *humid* way.

But, besides modes of proceeding which are purely chemical, and independent of the employment of chemical agents, the operator is frequently obliged to resort to processes entirely mechanical. The operations of chemistry might be naturally divided, therefore, into different classes, according to the several objects they have in view, and to the ways and means employed to accomplish those objects. All the operations which tend to change the form, without changing the nature, of a substance; those, for example, performed by the hammer, the knife, the pestle, and instruments of a like nature; and all those which determine the quantities of bodies are mechanical operations. But the operations performed by the aid of chemical powers and agents, and which separate the constituents of bodies, are purely chemical operations. We know that chemical action can only take place between bodies of different kinds, and that at least one of the bodies must be in a fluid state; we also know that chemical action is accelerated, when the

solid that is put into the fluid is divided into small particles. Now the division of a solid body into small particles is accomplished by pounding, grinding, rasping, cutting, and other operations of a like nature: hence the adoption and employment of chemical agents by the practical chemist.

It would be of little account to draw a line of distinction between these different kinds of processes: we shall therefore, instead of classifying them, proceed to give directions for conducting them.

255. WEIGHING.—The beginning and end of every exact chemical process consists in *weighing*; and the best means of ascertaining measures of weight is by means of *scales*. Accurate balances are therefore indispensable. What are termed *apothecaries' scales*, are very convenient for students who operate on small quantities of the different substances. These may be bought at the Apothecaries' Hall (and perhaps at druggists' shops). They are put up in a little box, which, besides the scales, contains a series of weights from half a grain to two drachms, in all about 20. These scales are sufficiently accurate for all common purposes. The charge made for the box complete is about 4s. 6d. The student must be careful not to use scales, or any other delicate metallic apparatus, in any place where acid vapours are flying about; for if he does, they will be seriously injured. It would be found very convenient, to have, besides these scales, others, to weigh quantities as high as a quarter of a pound.

We here annex the table by which, in chemistry, the weights of bodies are calculated. It may be mentioned that ambiguity is most easily avoided by reckoning by grains.

TROY WEIGHT.

Pound. Ounces. Drachms. Scruples. Grains.

1 — 12 — 96 — 288 — 5760.

1 — 8 — 24 — 480.

1 — 3 — 60.

1 — 20.

256. LIQUID MEASURE. In order to estimate the quantities of *fluid* bodies recourse is had to measures of

capacity. In chemistry the *wine pint* is usually employed. This measure corresponds to 28.875 cubical inches of water, at the temperature of 60°. For chemical uses, the graduated test-tube, fig. 12. is a measure of great convenience. It holds two troy ounces of water, equal to $\frac{1}{8}$ of the pint, to 960 troy grains, and to 3.6096 cubical inches. The following is a table of this measure :—

LIQUID MEASURE.

Gal. Pints. TroyOunces. Drach. Cubical Inches.

1	—	1	—	128	—	1024	—	231.
		1	—	16	—	128	—	28.875.
				1	—	8	—	1.8048.
						1	—	0.2256.

257. ON TAKING SPECIFIC GRAVITIES.—By the term specific gravity is understood the density (or quantity of matter under a certain bulk) of one body, compared to the density of another. This latter body is assumed as standard, and the standard to which bodies are generally compared is pure water, at the temperature of 60°. In other words, specific gravity is the comparative weights of different sorts of matter. Having found, by a certain process, that a given quantity of water weighs 1000, we employ the same method to ascertain the weight of the same quantity of the metal mercury; we find it to be 13000: thus we have the comparative weights of the same bulk of these two bodies, and we say that the specific gravity of mercury is, to that of water, as 13 to 1. If water at 1000 is assumed as a standard of specific gravities, which, as it has been said, it generally is, then the specific gravity of mercury (given with a reference to the standard) is 13000. The object of finding the specific gravity of bodies, is to distinguish them from each other in one of their most obvious qualities—namely, weight of matter contained in a given space. We proceed to describe the methods by which the specific gravities of different kinds of bodies are obtained; and give in the APPENDIX a general TABLE OF SPECIFIC GRAVITIES.

258. TO DETERMINE THE SPECIFIC GRAVITY OF A SOLID it is weighed, first in air, and then in water. To do this it is necessary to be provided with very accurate scales, the bottom of one of which is affixed a small hook, to which the substance is fastened by a fine thread, or hair. When the solid, after being weighed in air, is lowered into the water, it loses of its weight, a quantity precisely equal to the weight of its own bulk of water; and hence, by comparing this weight with its total weight, we find its specific gravity. The rule therefore is, divide the total weight by the *loss* of weight in water, the quotient is the specific gravity. Thus, if a mineral weighs 3 ounces in air, and 2 ounces in water, and the *total* weight be divided by the *loss* which is 1, the quotient, or specific gravity of that mineral will be 3.

259. A VERY READY WAY TO DETERMINE THE SPECIFIC GRAVITY OF SOLIDS, is to fill a phial with water, and note the weight of the whole accurately in grains. Then weigh 100 grains of the mineral or other substance to be examined, and drop it gradually into the phial of water. The difference of weight of the phial with its contents now, and when it was filled with water only, will give the specific gravity of the matter under consideration. For example, if the bottle weighs 50 grains more than it did when filled with water only, it shows that 100 grains of the mineral displace only 50 grains of water, and consequently that its specific gravity is 2000, or twice that of water. The kind of phial which may be advantageously used for experiments of this kind, is represented by fig. 41, plate 3.

260. TO DETERMINE WITH READINESS THE SPECIFIC GRAVITY OF A LIQUID.—We use for this, the *specific gravity bottle* (fig. 41). This, when filled with water, contains 500, 1000, or any even number of grains. This bottle is filled with the liquid, the specific gravity of which is required, and then weighed with its contents: the result, deducting the weight of the bottle, is the weight of the liquid under examination, compared to the weight of water. If, for instance, the bottle holds 1000 grains of water, and 1850 grains of sulphuric acid, then the specific gravity of the latter, is to that of the former, as 1850 is to 1000.

* * * In taking the specific gravities of bodies, attention should always be paid to their temperature; because the specific gravity of a body when heated, is much less than the specific gravity of the same body, in a cold state.

261. PULVERIZATION,—LEVIGATION,—TRITURATION.

—These, strictly speaking are merely mechanical operations, by which bodies are reduced into fine powder, and rendered more fitting to undergo chemical action than they were in their compact form. This kind of process never separates from each other the elementary principles of bodies; for every particle of an impalpable powder, is a small whole, perfectly similar in its composition to the original mass from which it was divided. But, on the contrary, chemical action separates the constituents of a body. The one class of processes is employed to overcome the attraction of aggregation, the other, to overcome the attraction of composition.—Brittle substances are reduced to powder by means of *the pestle and mortar*. (See 300.) Some substances are reduced to powder by a dexterous use of the pestle round the sides of the mortar; in fact, by *rubbing*, and this is that which is termed *trituration*. Reiterated blows of the pestle (which constitutes *pulverization*) are made use of, to powder hard bodies in iron mortars. Only a small quantity of the substance to be powdered should be put into the mortar at one time. *Levigation* is generally performed by rubbing a body (sometimes with the addition of water) on a flat stone, with another stone, round on one side to suit the hand, and flat on the other, which is called a *muller*. A thin flexible knife, either of iron or bone, is employed to collect the substance under operation, from the sides to the centre of the flat stone or mortar. Bodies that are not brittle are reduced to small particles by means of *files, rasps, knives, and graters*.

262. SIFTING and WASHING are performed to separate the finer particles of bodies from the coarser, which may want further pulverization. For the operation of *sifting*, the well-known instrument called a *sieve* is employed. Sieves, for nice cases, should be kept of different degrees of fineness. *Washing* is used for procuring powders of a more uniform degree of fineness, than can be done by means of

the sieve ; but it can only be used for such substances as are not acted upon by the fluid which is used. The powdered substance is mixed with water, or other convenient fluid ; the liquid is allowed to settle for a few moments, and is then decanted off ; the coarser powder remains at the bottom of the vessel, and the finer passes over with the liquid. By repeated decantations in this manner, various sediments are obtained, of different degrees of fineness ; the last, or that which remains longest suspended in the liquor, being the finest. Any of the glasses, figs. 6 7, 8, 9, 14, 16, 17, may be used for this purpose : those with lips, figs. 7, 8, 9, 14, are most convenient.

263. FILTRATION is employed to separate solids from fluids. Filters are of different materials ; for ordinary purposes, paper is used. The paper must be of a porous nature : and unsized, *blotting paper*, without colour, which may be had at the stationers, answers very well. The filtering paper is folded into a conical form, and placed in a funnel, in order that, when wetted, it may not break. Fig. 13, is a ribbed glass funnel, which is the best that can be used for this purpose. The substance to be filtered must be poured into the filter gradually. The portion of liquid that passes through first must be re-filtered, as it is seldom clear. But in a short time, the fibres of the paper swell, by accumulating moisture, and then the liquid that runs through is perfectly transparent. Acids, alkalies, and other corrosive fluids, are best filtered, by means of a glass funnel filled with pounded glass, a few large pieces being first put into the neck of the funnel, smaller pieces over these, and the finer powder at the top. The porosity of this filter retains much of the fluid, which may, however, be recovered (though at the inconvenience of dilution) by gently pouring on the filter a portion of water. The fluid will then be displaced, and pass through, and part of the water be retained in its stead. The funnel should be placed on one of the tall glasses, figs. 7, 8, 9, which may serve as a recipient for the filtered liquid.

264. DECANTATION, as well as filtration, is an expedient by which fluids are separated from solids that are diffused in them. The operation consists in allowing the

quid to settle till the solid matter have subsided, when the clear supernatant fluid is gently poured off. It is best performed with tall cylindrical glasses, furnished with spouts—such as figs. 7, 8, 9. If the sediment be so light as to mix again with the fluid when gently moved, the clear fluid must be drawn off by means of a syphon.

265. SOLUTION is when a solid put into a fluid entirely disappears in it, leaving the liquor clear. The body which thus disappears, is said to be *soluble*, and the liquid it dissolves in, is called the *solvent*, or *menstruum*. Sugar and salt are soluble bodies; for when they are put into water, they disappear entirely; but chalk is an insoluble body; for when that is put into water, it only makes the fluid *turbid*, or muddy, for a short time (being merely *diffused* through it) and then sinks to the bottom. Some bodies are capable of being dissolved in one kind of liquid, but not in another kind. Camphor, for instance, is soluble in alcohol, but insoluble in water. Metals are soluble, but their *dissolution* only takes place when they are put into acids. For some observations regarding metallic dissolution, see 156 to 160.

The operation of *solution* is more speedy in proportion as the substance to be dissolved presents a greater surface: on this principle is founded the practice of pounding, cutting, and otherwise dividing the bodies intended to be dissolved.

The solution of a body invariably produces cold; (*how* as explained at 382) and advantage has been taken of this phenomenon, to procure artificial cold, much greater than the most rigorous temperature ever observed in any climate. A *table* showing the different degrees of cold produced by different mixtures, will be found in the APPENDIX.

Solution is much accelerated by heat and agitation. But whether a cold liquid, or heat, or agitation should be employed in any particular case of solution, must be determined by the nature of the substance operated upon. In making solutions, it is necessary to use a vessel of such materials as shall not be acted upon by its contents, and of sufficient capacity to admit of any sudden expansion, or frothing, to which chemical action may give rise. The

glasses, figs. 6, 7, 8, 9, 14, 16, 17, may be employed when heat is *not* required, and those, figs. 53, 54, when heat is required. When either of the latter vessels is employed it should not be above $\frac{1}{3}$ full. It should be tied over with a piece of wet bladder, so that none of the fluid may be thrown out by the agitation of the contents. A few pinholes must be made in the bladder to admit of the escape of any gas which may be generated. Heat is applied to the vessel in the manner described at 314.

266. LIXIVIATION is used for separating such substances as are soluble in water from such as are insoluble. Suppose, for example, it is required to separate the *sarsaparilla* from a mixture of sand and salt: the compound body is placed in water; the salt is dissolved by the water; the sand is diffused through it. The mixture is filtered; the salt passes through with the water; the sand remains on the filter. The apparatus used in filtration, with the addition of a jug, is all that is required for this operation.

267. INFUSION is performed when we pour a hot liquid upon a substance that is partly soluble and partly insoluble in order to extract something from it. The making of *tea* is an instance of the performance of this operation.

268. DIGESTION.—This operation consists in soaking for a long time, a solid substance in a liquid kept constantly hot.

269. MACERATION.—The continued steeping of a solid body in a cold liquid. Ink is produced by macerating the materials of which it is composed.

270. EVAPORATION is a process employed to separate a fluid from a solid, or a more volatile fluid from another which is less volatile, by means of heat. The vessels in which it is performed, are described at 299.

As during evaporation, the fluid carried off in the state of vapour, is entirely lost, being sacrificed for the sake of the fixed substances with which it was combined, and which remain behind, this process is only employed when the fluid is of small value, as water, for instance. But when the fluid is of sufficient consequence to be preserved we have recourse to a process termed *distillation*.

271. DISTILLATION is evaporation performed in close

vessels, composed of different materials, and varying in their forms. The vessel usually employed, in the large way, is called a *still*, and is represented by fig. 56. It consists of a copper vessel, of the shape of a tea-kettle, but without its spout and handle, (*a*,) enclosed in the brick-work of a furnace. Into the opening of this vessel, instead of a common lid, a moveable head (*b*,) is affixed, which ends in a narrow open pipe. This pipe is received into what is called a *worm*, which is a tube of lead, twisted spirally, and fixed in a wooden tub, (*c*,) so that it may be surrounded by cold water. When the apparatus is to be used, the liquid intended to be distilled, is put into the body of the still, and the head is fixed in its place, the pipe, which terminates it, being received into the leaden worm. When fire is then kindled in the furnace, the door of which is seen at *f*, and the liquid is raised into vapour, which passes into the worm, is there condensed by the surrounding cold water, and flows out of the extremity of the pipe, by the cock *d*, into the vessel *e*, placed to receive it.

The common still, however, can only be employed for volatilizing substances, that do not act on the copper (or other metal) of which it is made; and it is, therefore, limited to a few operations. The vessel, fig. 26, is of glass, and is also used for distillation. It is termed an *alembic*, and consists of two parts; the body *a*, for containing the materials, and the head *b*, in which the vapour is condensed, by the application of wet cloths. The condensed fluid runs in a groove at the base of the head, and is conveyed, by the pipe *c*, into the receiver *d*. At *e*, is presented one of the rings of the lamp-furnace, (fig. 23), to which the alembic is placed, in order to be held over the flame of a lamp.

Besides the alembic and still, there is another utensil, which is employed perhaps more than either of those, in the operation of distilling. This is the *Retort*. It is a glass vessel, represented by fig. 18; *a* is the body of it, *b* the neck. In the top of the body is an opening, through which the materials to be distilled, are inserted. To this opening a glass stopple is fitted, by grinding, so as to be air-tight. Retorts are sometimes made without

the opening at the top, they are then cheaper, and are called *plain* retorts; but those with the opening, (called *tubulated* retorts,) are far more convenient. Retorts are also made of earthen-ware, and of metal.

A necessary appendage to the retort, is a *receiver*. This also is a vessel of glass. Fig. 19, represents a tubulated one; *a* represents the body of the receiver, *b* the neck which receives into it the neck *b* of the retort. Sometimes the neck of the retort is much too small to fit closely the neck of the receiver. In that case, a cork must be provided that fits the latter, and must have a hole burnt or bored through it, of a sufficient size to hold the former. The plan with the cork will not answer, however, when corrosive fluids are distilled. We must then use an instrument of glass, called an *Adopter*. This is represented in fig. 20, *b* is the adopter, *a* the neck of the retort going into the wide end of it, *c* the neck of the receiver enclosing the narrow end of it. The joinings are, as represented in the figure, secured by *lutes*.

Heat may be applied to glass retorts by means of the lamps. figs. 21, 24, properly regulated, to make a regular flame. Earthen-ware and iron retorts are exposed to the naked fire.—In order that the vapour which rises may be condensed as fast as it comes over from the retort, the body of the receiver is either placed in a tub of cold water or kept cool by the continually-renewed application of wet cloths.

272. **SUBLIMATION** is a process, by which volatile substances are raised by heat, and again condensed in the solid form. This operation is founded on the same principles as distillation, and its rules are the same, as it is nothing but a dry distillation. The apparatus for sublimation is very simple: that represented by fig. 26, and described under “distillation,” with the head *b*, wanting the neck *c*, will answer for most purposes. For example of sublimation, see 644.

273. **ON CRYSTALLIZING.**—The nature of crystallization has been already explained. The mode of obtaining crystals of certain bodies, differs according to the nature of those bodies. If it is desired to obtain crystals of a salt

that is more soluble in hot water than in cold, (there are many salts of this kind,) all that there is to do, is to put into hot water as much of that salt as it will dissolve; in short, to make a hot saturated solution—and then to allow it to cool gradually; the slower the better. As the caloric which contributed to the fluidity of the salt flies off, crystals will be deposited at the bottom of the vessel. Salts that are soluble in equal parts of cold and hot water, can only be crystallized by driving the water of solution off in vapour. But this must be done very slowly; for a rapid evaporation leaves a salt, not in a crystallized state, but in that of a solid irregular mass. By the operation of crystallizing, salts which differ in their degrees of solubility, or whose solution is unequally accelerated by heat, may be obtained separately from the same solution. Thus, if two salts be dissolved in the same liquid, and if one of them be much more soluble in hot than in cold water, and the other be equally soluble at any temperature, on evaporating the solution sufficiently, the latter salt will crystallize while the liquor is hot; on cooling, the other will shoot into crystals; and by alternate evaporation and cooling, the two may be obtained uncombined, though perhaps with a little intermixture of each other.

The only general rule that can be given to the young student, for the purpose of directing him how to crystallize bodies, is this: slowly evaporate the solution, until a pellicle (or thin skin) is formed on the surface of it; then set it in a cool place, where it will be free from dust, and can remain undisturbed.—This rule will not, by any means, apply to all salts, nor is there any other rule that will. Nothing but experience, and a knowledge of the habitudes of the various crystallizable substances, can be of much avail. For evaporating the solutions, the vessels described at 299, may be generally made use of. Metallic vessels are used in the large way.

274. PRECIPITATION.—Sometimes when two bodies (one at least of them being in a liquid state) are placed together, chemical action ensues, and a powder is formed and thrown to the bottom of the vessel. This powder is called a *precipitate*, the agent which causes it to be thrown down, the

precipitant, and the process which causes its production, *precipitation*. Figs. 6, 7, 8, 9, 14, 16, 17, represent the jars which may be used for this purpose.

275. GRANULATION ; the method of dividing metallic substances into grains or small particles, in order to fit them for different purposes. It is performed either by pouring the melted metal into water from a considerable height, or by shaking it in a box, previously well-rubbed with chalk, till the moment of congelation, at which instant it becomes converted into powder.

276. REDUCTION.—The operation by which metals are restored to their metallic state, after having been deprived of it, either by combustion, as the metallic oxides, or by the union of some heterogeneous matters which disguise them, as fulminating gold, horn silver, cinnabar, and other compounds of the same kind. Reduction is also called *revivification*. The operation is performed in crucibles, by the aid of heat, and with the addition of certain substances, which act chemically upon the body to be reduced.

277. FUSION.—The act of converting a solid into a fluid by means of heat. The operation is generally performed in vessels called *crucibles*, various kinds of which are described at 298.

278. ON FURNACES.—In order that substances to be fused may be submitted to the necessary degree of heat, instruments of different forms and sizes, termed *furnaces*, are employed ; and, in most books on chemistry, particular directions are given for the construction of them. But we do not intend to give any descriptions of furnaces in this work ; because, first, for the simple experiments here recommended, the use of furnaces is not required ; as the heat of a clear kitchen fire, urged by common bellows, will answer every object ; and, secondly, because the student would act wrong in buying or building furnaces, for which perhaps he might never have any use.

For a great number of chemical processes, in the small way, heat may be applied, in the very best possible manner, by means of a lamp.—Refer to what is said in the description of the *lamp*, fig. 21, and of the lamp-

furnace, fig. 23.—The characteristics of operations performed by means of the instruments here recommended, are simplicity, quickness, and cheapness,—qualities which few students know not how to prize.

279. ON THE BLOW-PIPE.—Every effect of the most violent heat of furnaces, may be produced by the flame of a candle or lamp, urged upon a small particle of any substance, by an instrument called the *blow-pipe*. This consists of a conical brass pipe, about 8 inches long, $\frac{1}{4}$ of an inch diameter at the top, with a curvature near the lower end, whence the point tapers off, with a very small perforation for the wind to escape. For nice purposes, the blow-pipe is provided with a bowl or enlargement, in which the vapours of the breath are condensed and retained, and also with two or three moveable nozzles, having holes of different diameters, to afford a larger or smaller flame. Fig. 59, represents a blow-pipe agreeing with the above description ;—*a* is the part of it that is taken into the mouth, *b* the bowl which condenses the humidity of breath, and *c* the point directed against the flame.

There is an artifice in the blowing through this pipe, which is more difficult to describe than to acquire. The effect intended to be produced is a continual stream of air for many minutes, if necessary, without ceasing. This is done by applying the tongue to the roof of the mouth, so as to interrupt the communication between the mouth and the passage of the nostrils ; by which means the operator is at liberty to breathe through the nostrils, at the same time that, by the muscles of the lip, he forces a continual stream of air from the anterior part of the mouth, through the blow-pipe. When the mouth begins to be empty, it is replenished by the lungs in an instant, while the tongue is withdrawn from the roof of the mouth, and replaced again in the same manner as in pronouncing the monosyllable *tut*. In this way the stream may be continued for a long time without any fatigue, if the flame be not urged too impetuously, and even in this case, no other fatigue is felt, than that of the muscles of the lips.

A candle with a thicker wick than they are generally used with, may be conveniently employed for occa-

sional experiments. It should be snuffed rather short, and the wick turned on one side towards the object, so that a part of it should lie horizontally. The stream of air must be blown along this horizontal part, as near as may be without striking the wick. If the flame be ragged and irregular, it is a proof, that the hole is not round or smooth; and if the flame have a cavity through it, the aperture of the pipe is too large. When the hole is of a proper figure and duly proportioned, the flame consists of a neat luminous blue cone, surrounded by another flame of a more faint and indistinct appearance. The strongest heat is at the point of the inner flame.

Considerable disadvantages, however, attend the use of candles. The difficulty of getting good ones, or such as have thick wicks, is frequently great. By far the best flame that can be employed, is that of a lamp, because it is large, and easily and cheaply obtained. Fig. 44, represents a blow-pipe lamp, of perhaps as simple a construction as could be devised.—Refer to the description of that figure.

The substance intended to be acted upon by the blow-pipe ought not to exceed the size of a peppercorn. It may be laid upon a piece of close-grained, well-burned charcoal; using, not the ends of the fibres, but the sides; otherwise the substance, when fused, instead of forming a round bead, is scattered about. In general, a small hole is made in the charcoal, and the body put into it. When the substance under examination is of such a nature as to sink into the pores of the charcoal, or to have its properties affected by the inflammable qualities of that substance, *then*, instead of being laid on charcoal, to be exposed to the heat, it is placed in a small metallic spoon, formed of gold, of silver, or of platinum. Some information regarding a spoon of this kind may be obtained from fig. 60, and the description of it.—The heat of the blow-pipe should be communicated gradually. The flame should not be immediately applied to the substance under examination, but directed a little above it, or else to the part of the spoon just beneath it.

Small plates of clay, prepared as directed below, are found to be a very useful addition to the blow-pipe apparatus. The colours of bodies melted with borax, &c. are shown to great advantage on them; and quantities of matter too minute to be tried on charcoal, or in the platinum spoon, may on them be readily examined, either alone, or with fluxes. *Process for forming clay plates.* Extend a white, refractory clay, by blows with the hammer, between the folds of a piece of paper, in the same manner as gold is extended between skins. Then, the clay and paper together, must be cut with scissors, into pieces about half an inch long, and a quarter of an inch wide, and afterwards hardened in the fire in a tobacco-pipe.

Many advantages may be derived from the use of this simple, and valuable instrument (the blow pipe.) Its smallness, which renders it suitable to the pocket, is no inconsiderable recommendation. The most expensive materials, and the minutest specimens of bodies, may be used in these experiments; and the whole process, instead of being carried on in an opaque vessel, is under the eye of the observer from beginning to end. It is true, that very little can be determined in this way concerning the quantities of products; but in most cases, a knowledge of the contents of any substance is a great acquisition, which is thus obtained in a very short time, and will, at all events, serve to show the best and least expensive way of conducting processes, with the same matters, in the larger way.

280. FOR PERFORMING EXPERIMENTS ON GASES, many articles of apparatus, not hitherto described, are absolutely necessary. These consist, partly of vessels for containing the materials which afford the gases, and partly of vessels adapted to contain the gases, and submit them to experiments.

Some gases are procurable by the mere mixture of the substances, which upon combining evolve them; but others cannot be obtained without submitting the materials to heat. For these different modes of proceeding, different vessels are requisite of course.

281. Fig. 29, represents a glass bottle with two necks, one of which has a glass stopple fitted to it by

grinding. This utensil is employed, when a gas is to be obtained without the aid of heat. The materials which are to afford the gas are put into the bottle. The solid part being broken into small pieces, to facilitate chemical action, and the liquid being poured in through a funnel, to avoid soiling the neck of the bottle. The stopple is then inserted in the neck to which it is fitted, and the gas, as it is formed, escapes through the neck that remains open. Now, in order to convey the gas thus evolved into a proper recipient, the instrument represented by fig. 30 is employed. This is a bent glass tube, with one end closely fitted in a cork that fits the open neck of the bottle, fig. 29. This tube is fixed in its proper place, before the materials for evolving a gas are put in the bottle; and, if it is thought necessary, the juncture is luted. The gas, therefore, when formed, escapes from the open end *b* of this tube.

282. For procuring the gases that are producible by the aid of heat, the retort, (fig. 18) is generally employed. The materials are inserted through the tubulure (or opening at the top of the body of the retort) which is then closed by a glass stopple; and the round part of the retort (which must be very thin in order that it may bear the application of heat) is then placed in one of the rings of the lamp-furnace, over a lamp giving a gentle heat. The materials will then give out the gas, which of course escapes through the neck *b* of the retort.—In many cases, a Florence flask (fig. 54) with a cork perforated by a bent glass tube, or even by a tin pipe, will serve instead of a retort and is much cheaper.

283. For receiving the gases, glass jars, of various sizes and shapes, (figs. 34 *a* 35, 36, 37, 51, 52,) are indispensable. The descriptions of these different figures will explain the peculiarities of each. Read, therefore, those descriptions; and read, also, the note 409.

284. To contain these jars, when in use, the *Pneumatic Trough* (fig. 25) is required. The description of figs. 25 and 27, gives a minute account of this apparatus, and must now be read. When the pneumatic trough is used, water must be poured into it, till it rises about one inch above the upper surface of the shelf (to the point marked *g* in

fig. 27). When the glass jars *e* and *f*, (fig. 27) or any other vessels, open only at one end, are plunged under water, and inverted after they are filled, they will remain full, notwithstanding their being raised out of the water, provided their mouths be kept immersed; for, in this case, the water in the jar is sustained by the pressure of the atmosphere, in the same manner as mercury in a barometer. It may without difficulty be imagined, that if common air, or any other fluid resembling common air in lightness and elasticity, be suffered to enter these vessels, it will rise to the upper part, and the surface of the water will subside. If a bottle, a cup, or any other vessel, in that state which is usually called empty, though in reality full—of *air*, be plunged into the water with its mouth downwards, scarcely any water will enter, because its entrance is opposed by the elasticity of the included air: but if, while the vessel is immersed, its mouth be turned upwards, the air will rise in bubbles to the surface of the water, leaving that body to occupy its place in the vessel. Suppose this operation to be performed under one of the jars which are filled with water, the air will ascend as before; but, instead of escaping, it will be detained in the upper part of the jar. In this manner, therefore, we see, that air may be emptied out of one vessel into another by an inverted pouring. Just in this manner are gases collected in vessels placed in the pneumatic trough: the jars which are to receive certain elastic fluids, are filled with water, and placed, mouths downwards, upon the shelf, over the holes; and the necks of retorts, and ends of tubes, from which gases are evolved, are directed below the holes under the jars: then the gases, as they issue forth, rise in bubbles through the holes, enter the jars, driving thence the water, and occupying its place. When, therefore, the jars are thus emptied of water, they are filled with gas.

285. When air is to be poured from a vessel with a wide mouth (such as fig. 36) into one with a narrow mouth, (such as fig. 37), the instrument (fig. 15,) called an *air funnel*, is employed: it is held under the shelf of the trough, and its neck is put through one of the holes in the shelf, so as to enter the mouth of the receiver.

286. Many kinds of gas combine with water, and therefore cannot be collected in jars placed in the water-trough described above. For retained gases of this kind, the chemist usually employs a trough containing mercury; but this mode is much too expensive for the student to proceed in. A readier and cheaper plan of filling vessels with gases of this kind, is described at 450, 468, 470. It is founded on the difference in the specific gravity of the *airs*.

287. An apparatus, almost indispensable in experiments on this class of bodies, is a *Gas-holder*, which enables the chemist to collect and preserve large quantities of gas, and to apply it to certain purposes, with great convenience. See fig. 61, and read the description of it.

288. When trial is to be made of any kind of air, whether or not it is fit for maintaining combustion, the air may be put into a long narrow glass vessel, the mouth of which, being carefully covered, may be turned upward. A bit of wax candle being then fastened to the end of a wire, which is so bent that the flame of the candle may be uppermost, is to be let down into the vessel, which must be kept covered till the instant of plunging the lighted candle into it. Fig. 46 is a representation of this mode of making experiments.

289. Where the change of dimensions which follows from the mixture of several kinds of air, is to be ascertained, an instrument, called an *eudiometer* is made use of. This is represented by fig. 34. See the description of that figure. It should be accompanied by a glass tube containing exactly either one or two cubical inches.

290. It is frequently an interesting object, to pass the electric spark through different kinds of air. But, for the performance of this kind of experiment, an electrifying machine (an instrument which many persons may not have) is necessary: as also is an eudiometer of a peculiar construction. Fig. 63, shows this eudiometer, and the description of that figure, explains the method of using it.

291. As for the purpose of experiment, it is frequently necessary to fill bladders with gases, we shall describe the

ode of doing it.—This is *represented* by figure 35—a is glass receiver, open at the bottom, and having a short neck at the top. To this neck is cemented a brass cap, on which is screwed a stop-cock. We are to suppose, in the present case, that this receiver is standing on the shelf of the pneumatic trough, and that it is filled with gas. Take a bladder, with a stop-cock fastened to its neck by means of a ferrule, one end of which has the mouth of the bladder tied securely round it. The bladder must be moistened, to render it flexible, and then pressed or wrung, to expel all the common air from it. (*This is necessary to prevent accidents.*) Shut the stop-cock, and then screw it to the stop-cock on the top of the receiver. Next, open both the stop-cocks, hold the apparatus in the manner shown by the figure, gently slide the receiver off the shelf, and press it down into the water: the gas will soon enter and fill the bladder, being forced through the opening, by the upward pressure of the water. The stop-cocks are then to be closed, the receiver replaced on the shelf, and the two vessels disunited.

292. When it is necessary, in order to make room on the shelf of the trough, to move some of the filled jars off, proceed thus: if the jar to be removed has a wide mouth, it must be gently slid off the shelf into a flat dish filled with water. See fig. 36. But if the jar has a narrow mouth, like fig. 37, all that is necessary, is to insert in its neck (while it remains under water) a ground glass stopple, or a good cork. Some gases, such as oxygen, may be kept in common wine bottles, for several months, secured merely by corks.

293. If any thing, as a gallipot, is to be supported at a considerable height within a jar, a stand of wire (represented by fig. 48) is employed. This answers better than any other kind of stand; because it takes up but little room, and is easily bent to any height or figure.

293.* The method of *weighing gases* with accuracy is very simple, but it requires the aid of an air-pump. But few students, therefore, will be able to take the specific gravities of gases, for few will possess a good air-pump. It will be as well, however, to explain the operation.—A

stop-cock is affixed to a glass globe or flask, and this apparatus is then weighed. The glass vessel is then exhausted of air, and is again weighed. The difference of the weights shows the weight of the atmospheric air withdrawn from it. The stop-cock of the flask is now affixed to a gas-holder, and the flask is filled with the gas to be tried. It is again weighed, and the result is compared with the results of the other weighings; the difference shows the specific gravity of the gas. Thus, if the apparatus filled with atmospheric air weighed 6 ounces, 10 grains; when exhausted 6 ounces; and when filled with gas, 6 ounces, 20 grains; then the specific gravity of the gas is to that of common air as 2 is to 1.

* *. Previously to undertaking experiments on other gases, it will be well for an unpractised experimentalist to accustom himself to the dexterous management of gas by performing, with common air, the processes of filling bladders, transferred from vessel to vessel, &c.

294. WOULFE'S APPARATUS FOR COMPOUND DISTILLATION.—In several cases of *distillation*, the substance raised is partly a condensable fluid, and partly a gas, which gas is incondensable by itself, but capable of being condensed by being transmitted through a liquid. The apparatus required by a process, in which this double purpose is effected, is represented by fig. 57, and is commonly termed Woulfe's apparatus. It is a series of receivers connected in a particular manner, and more or less in number as the case may require. The distilling vessel made use of, is the *retort*, (fig. 18,) into the tubulure of which, instead of a glass stopple, is inserted the safety tube, fig. 58.—See the description of that utensil. The first receiver (*a*) is joined to the retort, and has a right angled glass tube, open at both ends, fixed into its tubulure. *b*, the second receiver, is a bottle which, beside its usual neck at top, has an opening just at the place where its sides fall in to form the top: into this opening a glass tube is fixed diagonally, and the juncture is secured by the application of a lute. The lower end of this tube must be about an inch from the bottom of the receiver. *c* is in every respect the same as *b*. The openings spoken of are at *g g*, and the tubes fixed in them are shown by

The small tube *e*, which rises from the first receiver passes down the diagonal tube in the second receiver; and another small tube, likewise marked *e* rising from (and fitted in) the neck of the second receiver, passes down the diagonal tube in the third receiver. If there were more receivers, they would be like *b* and *c*, and would be connected in the same manner. The lower ends of the tubes *e e*, must project as far beyond the ends of the tubes *d d*, as they can do without touching the bottom of the bottles. The liquids by which the gas is to be absorbed are put into the second, and subsequent receivers, each being filled two-thirds full. The nature of this liquid is regulated by the nature of the gas to be absorbed. For gases that are readily absorbed by water, (such as sulphurous acid, muriatic acid, &c.) distilled water is made use of: for other gases (the carbonic acid for instance) solution of caustic potass is employed. In general, it is adviseable to put water into the second receiver, and the alkaline solution into the third: the *first* (*a*) is always left empty.

295. The materials being introduced into the retort, the arrangements completed, and the joints secured by cements, the distillation is begun. The condensable vapour collects in a liquid form in the receiver *a*, which is kept cool by being placed in water, or by having wet cloths applied to it. In the meantime, the evolved gas passes through the bent tube *e*, into the water contained in *b*, which continues to absorb it (if it is a gas absorbable by water) till it is saturated. When saturation takes place, or when no absorption ensues, the gas bubbles up through the water, passes through the second pipe *e*, and enters the receiver *c*.—And so the process continues till the liquids in all the bottles are saturated; and then, if any gas continues to be produced, it escapes through the neck *f* of the last receiver. Should it be required to preserve this overplus gas, it may be conducted into a receiver placed in the pneumatic trough, by fitting into the neck *f*, the bent tube, fig. 30.

296. It will be now proper to point out the use of the tube of safety, (fig. 58,) and of the wide diagonally-fixed tubes *d d*. Supposing the retort to be closed by a stopple

as usual; and supposing the bottles to be destitute of those tubes, and consequently, that $e e$, were luted into the receivers at $g g$, the process would be then liable to be interrupted by an accident. for if, in consequence of the irregularity of the heat, or other circumstances, a vacuum should be produced in the retort, by the re-absorption of gas, the liquids in the different receivers (being acted upon by the pressure of the atmosphere at f) would rush from one into another to supply that vacuum, and by such a mixture of products the whole experiment would be spoiled. If, on the contrary, gas were to be evolved faster than it could be absorbed by the different liquids, than it could escape at f , the apparatus might burst, with considerable danger to the operator. Should the operator close f as well as all the other places, then the apparatus would be destroyed, if either absorption or expansion took place; for, in the one case, the external air would press on the vessels till they broke, and in the other, the same effect would be produced by the elasticity of the gas confined within. Now, all these inconveniences are obviated by the employment of the different tubes. If an absorption takes place, when these tubes are fixed in the manner that has been described, the vacuum is instantly supplied by the external air, which rushes down the tubes $d d$, into the receivers, and down the tube of safety into the retort. The experiment being thus prevented from failing altogether, at the price of having a small portion of common air mixed with its products. On the other hand, no gas can escape for any pressure within is instantly followed by the formation of a high column of liquid in the tubes $d d$, which resists the egress of gas, as long as is consistent with safety.

297. We have now described all those chemical processes, which the experimentalist is in general called upon to conduct; as likewise, the apparatus he finds it necessary to employ: and in such a manner, as, we hope, will render our instructions beneficial to those who consult us. But besides the instruments used in general operations, many others are requisite for the obtainment of various particular objects. The figures on the annexed plates, with the following explanation of them, will give the student an

idea of a vast variety of the most useful and necessary articles of *the* chemical apparatus with which, in consistence with economy, he may provide himself.

298. Figs. 1, 2, 3 (plate 1.)—CRUCIBLES.—These vessels are of indispensable use in the operation of *fusion*. They are formed of the following substances: earthen-ware, black-lead, silver, platinum. The best kind of earthen-ware ones, are those called Hessian crucibles. These are of the shape represented by fig. 3. They support a very intense heat without melting; but, unfortunately, are liable to crack unless heated and cooled very gradually. The crucibles made of black-lead, (which are generally made of the shape of fig. 1,) are *not* liable to be cracked by alteration of heat; but they are unfit for the fusion of alkaline or saline matters. For these last mentioned bodies, when particular nicety is required, crucibles of silver must be used; and, for exposing bodies to a very great heat, those of platinum are necessary.

But for all ordinary purposes, the earthen-ware crucibles are the best in every respect. They are strong and cheap, and those are the grand points. You may buy at an iron-monger's, a set, (of shape fig. 3,) containing six, of different sizes, for *ninepence*. The metallic crucibles are sold by the ounce. Crucible covers (see fig. above the 2d. crucible,) cost from 2*d.* to 6*d.* each. These, in certain cases, are obliged to be luted to the crucibles to prevent the escape of any portion of the substances submitted to experiment. A tobacco-pipe with a large bowl may be used as a crucible in a vast number of processes.

299. Fig. 4.—A WEDGEWOOD'S WARE EVAPORATING BASIN.—This is an utensil of great utility. The substance of which it is formed, is so compact, that a strong heat may be suddenly applied to it, without any danger of its breaking. It also has the advantageous property of being unacted upon by acids. For the convenience of pouring liquids out of it, it is furnished with a spout. The student will find, that, for the numerous operations of evaporation and crystallization, he will require two or three of this article. Let him buy, at the shops where

Wedgewood's-ware is sold, one at 6*d.* one at 9*d.* and at 1*s.* These will measure (across the top) from $2\frac{1}{2}$ to 5 inches. Sometimes, for evaporating small portions of fluids, small capsules, resembling watch-glasses, are employed. The vessels used for evaporation should always be thin at the bottom, in order that they may bear, without breaking, the sudden application of heat.

300. Fig. 5.—A WEDGEWOOD'S-WARE PESTLE AND MORTAR.—This, like the article just described, has the properties of strength and unalteration by acid menstrea. It is so useful an utensil that the student *must* have it. He may buy, at the place where he gets the evaporating basin, a mortar measuring 4 or 5 inches across the top. This is a very convenient size : such an one will cost about 2*s.* 6*d.*

The Carron Company sell cast iron mortars, at very low prices. The writer of this bought of one of their Agents, (an Agent for the Carron Co. is to be found in most large towns,) a strong neat pestle and mortar, weighing 7 pounds, for *fourteen-pence*. This iron mortar should be got (if convenient) in *addition* to the one of Wedgewood's-ware.

301. Figs. 6, 7, 8, 9, PRECIPITATING AND DECANTING GLASSES of various shapes and capacities. The height of them is about 6 or 7 inches. Their other proportions may be calculated from this standard. The kind employed for decantation, are those with spouts. Go to a glass-house for these articles. No. 6, will cost 9*d.* ; No. 7, 1*s.* 6*d.* ; No. 8, 1*s.* 6*d.* ; No. 9, 1*s.* They should be made of thick glass, so as to be strong ; for they are frequently used : and the glass must be clear, otherwise the colours produced by the action of the precipitant, or test, will not be visible.

302. Fig. 10.—A GLASS FUNNEL *with a long neck*.—It should be about twelve inches in length. Its use is described in the experiments for the performance of which it is necessary. This, like all other glass utensils, must be got at a glass-house. It will cost 1*s.*

303. Fig. 11.—A DROPPING TUBE.—This is a glass tube, six inches in length, with a bulb about one inch in

diameter blown in the middle of it, and having its lowermost point drawn out till the opening in it is very small. It is used to convey from one vessel into another, a small quantity of water, or of any other liquid that the purpose may require. The manner of using it is this: while the lower end is immersed in the liquid, the ball is filled by the action of the mouth which is applied to its upper extremity. The latter is then closed by the finger; the water remains suspended in the tube, until it is removed to the place where it (the water) is to be put, when, by cautiously removing the finger, it is expelled in drops. Filters may be washed very neatly by means of this tube. A glass-blower will make the instrument for sixpence. When the student can manage the blow-pipe, he may easily make such things as this himself.

304. Fig. 12.—A GRADUATED JAR FOR MEASURING LIQUIDS.—This is a cylindrical glass vessel, which holds, when filled to a certain mark made near the top, two ounces of distilled water, at the temperature of 60° , in short, two ounces of clear cold water. It has other marks all down it, as low as one drachm; so that any quantity of water from one drachm (or, indeed the $\frac{1}{2}$ or $\frac{1}{3}$ of a drachm) to sixteen drachms, may be readily obtained for any required purpose.

305. But, besides the utility of this utensil for measuring water, it may be also employed, when the purpose does not require great accuracy, to measure liquids whose specific gravities are different. If, for example, we wish to obtain an ounce of sulphuric acid, we proceed thus: knowing, that the specific gravity of sulphuric acid is to the specific gravity of water, a little less than as 2 is to 1; and, consequently, that if an ounce of water occupies a certain number of divisions in this tube, then an ounce of the acid can only occupy a little more than half that number; of course, an ounce of sulphuric acid is readily obtained by pouring that liquid into the tube till it rises a little above the mark for $\frac{1}{2}$ an oz. of water. We proceed in like manner for other fluids, calculating the bulk according to the specific gravity. It is by no means intended, by the recommendation of this manner of measur-

ing liquid in general, to do away with more accurate modes: we only point out a method of proceeding in common cases, calculated to save time.

306. It may be observed here, that when, in giving directions for the performance of an experiment, we say, "an ounce *by measure*," it signifies by the measure for water. "An ounce of sulphuric acid by measure" is really almost two ounce of that liquid.—The price of this jar is about 2s. 6d.

307. Fig. 13.—A RIBBED GLASS FUNNEL used for *filtration*, and described under that head. It costs about 6d. or 8d. according to the size.

308. Figs. 14, 16, 17.—TEST GLASSES.—These are used to hold small quantities of liquids to which *tests* or *re-agents* are to be applied, in order to discover what bodies those liquids hold in solution. They are also used for the purposes to which figs. 6, 7, 8, 9, are applied. Fig. 17, is quite round and smooth at the bottom, so that any substances which may subside, can be readily taken out for the purpose of being weighed. The price of these is about 6d. each. It is very common to use pieces of thin glass tubes as test glasses.

309. Fig. 15.—A GLASS FUNNEL *used to convey Gas from a wide mouthed vessel into one having a narrow mouth.* It may measure five inches across the wide part, and be about 4 inches in depth. Get it made at a glass-house: it will cost 8d.

310. Fig. 18.—A TUBULATED GLASS RETORT, which, as well as its use, is described at 271, 282. Retorts are made of various sizes; one of which the body (*a*) will hold 8 or 10 ounces of water, is a very convenient size for ordinary purposes. The price of such an one is about 2s. 6d. NOTE.—*In selecting, at the glass-house, a retort, or any other utensil to be placed over a lamp or in a sand-bath, be careful to take one with as thin a bottom as can be got. Thin glasses are the only ones that will bear the application of heat; thick ones crack immediately upon being put to the fire.*

311. Fig. 19.—A TUBULATED GLASS RECEIVER, whose use is described under "distillation." One holding 10

ances (which is a very good size) will cost about 2*s*. Both *receivers* and *retorts* may be had without *tubulures*, and they then cost less than the others; but we recommend the student to get those that *are* tubulated, because the difference in price is more than overbalanced by the convenience of them.

312. Fig. 28.—AN ADOPTER; a tube of glass of unequal dimensions; used for lengthening the necks of retorts, and for fitting retorts to receivers. It costs about *l*. or 8*d*.

313. Fig. 21, (plate 2.)—A CHEMICAL LAMP for the portable furnace, fig. 23. This instrument affords the means of obtaining either a strong or a gentle heat, for a number of hours at once. It is simple in its construction, and, therefore, the more convenient for those who are inexpert.

a represents the body of the lamp, or that part of it which contains the oil. This (which is circular) measures in altitude $1\frac{1}{2}$ inches, in diameter 3 inches. Fig. 2, represents the top of it: *b* is the outer extremity, *a* is a tube passing through (but of exactly the same length as) the lamp, serving as a passage for a current of air which is necessary to support the combustion of the oil. This tube is supported in its place, by having its lower end soldered to the bottom of the lamp at the place where it enters it. Round its upper end, but a little below the top of it, is fitted a rim of brass, which has six cylinders of tin $\frac{1}{4}$ inch in length, and $\frac{1}{6}$ th of an inch in diameter, soldered in so many holes made at equable distances in it to receive them. This is shown by fig. 22. Only about $\frac{1}{4}$ of an inch of the length of these small pipes projects above the top of the rim, the long parts go down into the body of the lamp. These tubes are, as will readily be imagined, intended to hold the wicks. The rim is fastened in its place, by another brass rim which screws on above it. *d* figs. 21, and 22, show a small pipe, through which the oil is introduced, and which, when the lamp is filled, is closed with a brass screw. At *g*, fig. 22, is a small hole intended to receive the wire *e*, fig. 1. This wire rises two inches perpendicularly from the top of the lamp, and then turns off horizontally, terminat-

ing in a ring of the same size as, and hanging directly over the afore-mentioned rim of brass which bears the wicks. Into this ring is fitted a tin cylinder (*b*) (which must be kept *bright*) of such a length as to come to 5-8ths of an inch from the tops of the little tubes whence the cotton issues. The wire *e* fits easily into the hole that receives it, so that it may be readily moved about. The use of the cylinder is to condense the flame. Its action differs in no respect from that of the glass ones used in common, but it is to be preferred to them, on account of its being cheaper, and not liable to be broken. Its opacity is no argument against it; because the flame of this lamp is not valued for the *light* it furnishes, but for the *heat*.

From what has been said, and from an inspection of the accompanying figures, a clear idea of the construction of this instrument may readily be obtained.

By depressing three of the wicks, and lighting the remainder, a small flame is produced. By lighting all the wicks a strong flame is produced. The heat is kept up very regularly; and *no smoke arises*. Any tin smith will be able to make a lamp of this kind, and the price of it will be about 3s. 6d.

314. Fig. 23.—A TABLE LAMP-FURNACE.—This is an instrument of great utility. There may be performed by its aid a vast variety of chemical operations on a small scale, in a very easy and expeditious manner. It consists of a rod of iron *a*, two feet long, and rather less than half an inch thick, screwed into a foot of the same metal. Upon this upright slide three sockets with horizontal arms terminating in rings, (all made of iron,) which are of diameter $2\frac{1}{2}$, $3\frac{1}{2}$, and $4\frac{1}{2}$ inches; and they are fixable at any height, or may be turned round in any direction, by means of screws (see *b*) affixed to the sockets. These rings serve for supporting glass alembics, retorts, flasks, evaporating basins, &c.; for performing distillations, digestions, solutions, evaporation, analyses with the pneumatic apparatus &c. &c. *c* is an evaporating basin placed on a ring in the manner described. The lamp, (either fig. 21, or 22) which is to furnish the heat, is placed beneath the utensil to be heated at any distance which may be required.

if a strong heat is wanted, the distance must be small ; if a gentle heat, the distance may be greater.

A working smith, or ironmonger, will make a stand of this kind (not including a lamp, of course,) for 4s. 6d. Sometimes, in addition to the three rings, the stand has another arm with a board fixed to it for supporting a lamp (see the figure ;) but it is as well to lay a piece of board across one of the rings, or to give the lamp a separate support.

315. Fig. 24. A SPIRIT LAMP. This, for experiments in the small way, where but a moderate degree of heat is required, is an excellent contrivance. It is a glass vessel, of a flattish-globular shape, with a wide neck, and holding somewhat less than a common wine glass. On the neck is laid a circular concave piece of brass, with a small cylindrical pipe, an inch long, passing through the middle of it, to contain the wick (which is of common cotton.) The price charged for this lamp, by the chemical instrument makers, is 1s. 6d. Alcohol burns with a flame which is large or small, according to the size of the surface of the cotton wick. It gives no smoke, it has no disagreeable smell, nor does it spoil any thing if it happens to be overturned. But, with its many advantages, this lamp has one great disadvantage, namely, it cannot be used much, on account of the great expense attending the consumption of the spirit of wine.

316. Fig. 25.—A PNEUMATIC TROUGH.—This is a strong, square, water-tight, wooden box ; the internal dimensions of which are as follows : depth eight inches, length and breadth each sixteen inches. A shelf, sixteen inches long, and eight inches wide, having four holes cut in it, as is represented in the figure, is fixed on one side of the interior of the *trough* ; at such a height as to leave a space of two inches and a half between *its* top and the top of the box.—For an account of the uses of the pneumatic trough, see 284.

317. Fig. 27, is a section of the box : *a* is the body or open part of the trough, *b* is the shelf, *e* and *f* are two glass jars, inverted over two holes (*c* and *d*,) made in the shelf.—If the retort is used, (in the manner directed in para-

graph 282,) the neck of it is most conveniently brought under the hole *c*; but if the gas-bottle (fig. 29,) is employed the end *b* of the bent tube (fig. 30,) which is fitted to that bottle, is put through the opening *d*, in such a manner as that its point may turn up directly under a jar placed as *f* is.

Sometimes pneumatic troughs are made of tin, but wooden ones have many advantages; and the only great objection to them is, that they are not always water-tight. If, however, the carpenter who makes the box, takes care that the joinings of it are, by dove-tailing and re-batting, fitted closely together; and if a strong coat of paint be applied to it, both within and without, it will bear water without losing a drop. The shelf may be made moveable, and a moveable top may be fitted to the box, which may be also furnished with a lock and key. All these things are useful. When the box is not to be used for some time, place the shelf at the bottom of it; upon that, place the instruments formed of glass, then lock on the lid of the box, and all will be free from injury. For a box, made of strong wood, according to the directions here given, the writer of this paid—to the Carpenter, 8*s.*—to the Painter, 2*s.*

It may not be amiss to observe, that in a case of emergency, a good substitute for a pneumatic trough, is a wash hand-basin, or small wooden tub, with a board, which has a hole in it, large enough to admit the neck of a common wine bottle, laid across it.

318. Fig. 26.—A GLASS ALEMBIC for distillation and sublimation, described at 271. The head must be made to fit the body of it very exactly, or the substance that is volatilized will escape between them, and be lost. It must likewise be depressed round its lower part, so that a circular channel may be formed, to conduct the condensed fluid into the pipe *c*. An alembic, of which the body will hold a wine-pint, with a head proportionably large, will be of a sufficient size for common purposes, and will cost about 5*s.*

319. Fig. 28.—A SMALL GLASS RECEIVER.—This is sometimes fitted to the neck *c* of the *alembic*, to serve as a

ipient of the product of distillation, in lieu of the vessel, fig. 26.

320. Fig. 29. (plate 3.)—A Woulfe's BOTTLE with two necks, one of which is furnished with a glass stopple, fitted closely to it by grinding. This apparatus is much used in experiments on gases.—See 281. It may be had of any glass-house. One capable of holding twelve ounces of water is a good size, and such an one will cost 2s. 3d. 2s. 6d.

321. Fig. 30.—This is a glass tube, of the size shown in fig. 55, bent in such a manner, and of such a length as to fit it to convey a gas from the bottle, fig. 29, into a receiver placed in the pneumatic trough, fig. 25; *a* is a cork fitted to one end of the tube, and adapted to the neck of Woulfe's bottle; *b* is the end of the tube which is to be directed under a receiver.—See 281. This bent tube cannot be provided till both the trough (fig. 25) and the bottle (fig. 29) are ready for use: when the student has these, he must bend a piece of wire into the exact figure of which the tube (in order to fit the two instruments) must be a counterpart. Then, according to this pattern he must bend his tube; but if he cannot manage that himself, he must get it done at a glass-house, where he will have to pay about 8d. for it.

322. Fig. 31.—A JET-PIPE for making some experiments with hydrogen gas. It is simply the shank of a tobacco-pipe cemented into a cork, which fits one of the necks of the Woulfe's bottle, fig. 29.

323. Fig. 32.—A STOP-COCK represented on a large scale, and such an one as ought to be fitted to a bladder for performing some experiments with gases. *a* represents a screw on which is fitted the ferrule or plug that goes into the neck of the bladder.—See 291. This ferrule must be indented or roughened on the outside, in order that the string by which the bladder is bound upon it may have something to hold by. At the end of the stop-cock marked *b*, there are two screws; an internal one which fits the screw of fig. 49, and an external one adapted to the screw *b* of fig. 50, and to the screw *b* of the receiver *a*, fig. 25. *c* is a screw that opens or shuts the communication through the stop-cock. It may

be observed here, once for all, that whenever stop-cocks or other apparatus, are screwed together, great care must be taken to fit them closely. Every juncture must be rendered air-tight, by the interposition of circular pieces (or rings) of oiled leather.—These stop-cocks are made of brass, and may be bought of the tradesmen who fit up gas-pipes in shops, at 1s. to 1s. 3d. each.

324. Fig. 34.—An EUDIOMETER, or graduated glass tube used to measure the purity of air. *How the purity of air is ascertained* will be shown amongst the experiments; we have here to describe the instrument. Its length is 18 inches, and its diameter 1 inch; its bottom is shaped like a funnel, in order that it may stand firm, and its side has a scale drawn down it, graduated into equal measures of capacity—into cubical inches, or halves, or fifths, or tenths of cubical inches. One graduated half inch will be sufficiently fine for all the purposes a student will require it.

325. One method of graduating (or drawing scales on) glass jars is as follows: fix the jar to be graduated mouth upwards, in such a manner as to make it stand firmly without being held by the hands, then take a vessel capable of holding a certain quantity of water—for want of one holding a cubical inch, or half a cubical inch, use the graduated jar, fig. 12, and mark half-ounces by it; fill the measure with cold water, pour the water into the jar, and make a mark on the jar, exactly at the surface of the water with a file, or (which is much better) a glazier's diamond. Continue thus to pour in succession equal measures of water, and mark its surface at each addition.

326. Fig. 35.—A REPRESENTATION OF THE MODE OF FILLING BLADDERS WITH GASES.—See 291. *f* is part of the pneumatic trough fig. 25; *h* is the shelf; *g* shows the height of the water in the trough; *a*, a glass receiver, holding three pints, open at the bottom, and with a narrow neck at top; *c*, a brass cap, cemented on the neck of the receiver and having a screw adapted to one of the screws of stop-cock *b*; *d* is a bladder into which the gas is being driven from the receiver; and *e*, *b*, are the stop-cocks of the bladder and receiver fitted together to open a communication. A jar of this capacity may be got made at

glass-house for 3s. and the brass cap will be fitted on by furnishing iron monger for 1s. 6d. The prices of the other parts of this apparatus are mentioned elsewhere.

327. Fig. 36.—A GAS RECEIVER of a very convenient form. It is of glass, holds a quart, is open at bottom, and has a knob by which it may be held at top. It is represented as filled with gas, and standing in a dish filled with water, which forms a portable pneumatic trough.—See 2.—Such a jar costs 1s. 6d.

328. Fig. 37. A GAS BOTTLE *for experiments on the combustion of bodies in oxygen gas.* It must hold about a quart; and be *thin* round the sides, in order that it may not be cracked by the heat produced by deflagrating a body within. When this utensil is used, it is filled with gas, in the common method, over water, and its mouth is secured by the insertion of a ground glass stopple. In this position it may stand upright on a table, and is ready for an experiment. The price of a gas bottle of this kind, with stopple, at a glass-house, is about 1s. 6d.

329. Fig. 38. A COPPER DEFLAGRATING LADE *for containing substances that are to be immersed, in a state of combustion, in the oxygen gas contained in the bottle, fig. 37.* The bowl of the ladle is about the size, and the piece of metal it is made of about the thickness, of a shilling. The thin part, or shank of the ladle, passes through a cork which is adapted to the neck of fig. 37. When the cork is pushed into place, the bowl of the ladle must be about $\frac{1}{4}$ the whole height of the bottle from the bottom of it. The price of this ladle is 1s. Any Coppersmith will make it.

330. Fig. 39.—A SPIRAL IRON WIRE with a cork in its upper part that fits the neck of the gas bottle, fig. 37, and on its lower extremity, a bit of tinder. This apparatus is employed in the experiment which exhibits the formation of a metallic oxide, by the combustion of a metal in oxygen gas. The kind of wire that is used is about $\frac{1}{4}$ of an inch thick, and is called by the ironmongers to sell it, *binding-wire*. In order to bring it into the spiral, or cork-screw shape, it is coiled lightly round a stick of half an inch diameter, and then drawn off it. Afterwards, the cork is fitted on it, it is drawn out to a

proper length, ($\frac{3}{4}$ that of the bottle, say,) and has a morsel of tinder, charcoal, or thread dipped in brimstone or turpentine, fixed upon its lower end. Twopence worth of this wire is enough for a great many experiments.

331. Fig 40.—AN ECONOMICAL GALVANIC APPARATUS. Obtain a number (say thirty) of the smooth penny pieces; and the same quantity of plates of zinc of the same size. The pieces of zinc may be easily cast in a mould made of clay, for that metal is of about the same degree of fusibility as lead. Crude zinc to make them may be purchased at a brass-foundry, at tenpence a-pound. In addition to the plates of copper and zinc, it is necessary to be provided with the same number of pieces of wadded cloth, rather smaller than those plates in size. These be soaked in a solution of common salt, till they have thoroughly imbibed it; when they are to be used they must be taken out of the solution and gently squeezed in order to force out the superabundant water.

The *Voitaic Pile* (a, fig. 40) is to be formed of these materials in the following manner: lay down a circular piece of wood a plate of zinc, upon that a plate of copper, and then a piece of moistened cloth—let this arrangement be continued—zinc, copper, cloth—zinc, copper, cloth,—till all the pieces that have been provided be laid on. As the pile began with zinc, it must be concluded with copper. The number of pieces of each substance must be at least twenty, and may be advantageously extended to fifty; for the more pieces there are, the greater will be the effect produced by the pile which they form.

But a pile thus formed, is, when tall, very liable to be overthrown, and must, therefore, in order to avoid this accident, be supported. The method usually added to hold it up steadily, is to fix into the piece of wood in which the pile is formed, three rods of glass, which must touch the pieces of metal at three equidistant points. Do these rods may slide a piece of wood similar to the bottom piece, for the purpose of keeping by its pressure the different parts of the pile in close contact. Fig. 33 is intended to represent this top—the circle in the centre shows the situation of the pile—the three small circles, between

the outer and inner large circles, show the places where holes are cut for the reception of the glass rods. The upper side of the piece of wood *b*, and the under side of the other piece of wood, must be lined with tin foil. In the figure is represented the arrangement employed in decomposing water. *c* is a glass tube filled with water, and having the ends closed with corks: *d* and *e* are wires, one of them having an end fixed to the top of the pile, and the other an end fixed to the bottom of it. The other ends of these wires are passed through the corks, and in the tube *c*, approach to within a quarter of an inch of each other.

The zinc plates in the voltaic pile become oxydated after being used a certain time, and then the galvanic action ceases; but they may be cleaned by being put into diluted muriatic acid, by which the oxide is dissolved. They may also be cleared of the oxide by being filed.—It is advisable, when the voltaic apparatus is not to be used for some time, to take it down, by which means the zinc plates will be prevented from wearing out too fast.

332. A SPECIFIC-GRAVITY BOTTLE.—The use of this is described at 260. It is a little globular bottle with a flat bottom. It has a glass-ground stopple, with a small hole through it. When the bottle is filled with water, or any other liquid, and the stopple put in its place, the superfluous water escapes through the hole, and the bottle remains quite full, without any portion of air. A bottle holding 500 grains costs about 2s.—A weight to counterpoise the bottle must be obtained, made of brass or lead.

333. Fig. 42. (Plate 4.)—APPARATUS FOR EXHIBITING SOME OF THE PROPERTIES OF HEAT.—*a* is a glass tube of the thickness of fig. 55, and about ten inches long. *b* is a ball or globe blown at one end of the glass tube. A glass-blower will charge about 9d. for this instrument. *c* is a small stool, having a hole in the top, through which the lower end of the tube *a* is put, in order that it may be kept steady, and upright.

334. Fig. 43.—Dr. WOLLASTON'S CRYOPHORUS, or frost-bearer. It is a glass tube of the thickness of fig. 55, and about twelve inches long. At each end of this tube is a

ball of one inch diameter, and the tube is bent to a right angle at the distance of half an inch from each ball. One of these balls (as *a*) should be somewhat less than half full of water, and the remaining cavity should be as perfect a vacuum as can readily be obtained; which is effected by making the water boil briskly in the one ball before sealing up the capillary opening left in (*b*) the other ball. A glass blower will charge about 1s. 3d. for this instrument.

335. Fig. 44.—A LAMP to be used with the blow-pipe. *a* is a little oblong tin dish, similar to those that tarts are baked in; *c* is a tin tube soldered to the bottom of it, and is used to hold the lamp on a candlestick. Over half, or three-fourths, of the top of this vessel is fixed a broad piece of tin, and through the part remaining open, rises, in a diagonal direction, a wide tin tube (*b*) intended to hold a cotton wick. This tube is shaped at the bottom in the manner shown by the figure.—A tinsmith will make this lamp for 4d.

336. Fig. 45.—APPARATUS FOR PRODUCING MUSICAL SOUNDS BY THE COMBUSTION OF HYDROGEN GAS.—*a* the bottle whence hydrogen gas is issuing through the pipe *b*; *c* a glass tube eighteen inches long, and one inch wide, held over the flame produced by the burning gas.

337. Fig. 46.—APPARATUS FOR ASCERTAINING WHETHER A GAS WILL SUPPORT COMBUSTION.—See 288.

338. Fig. 47.—APPARATUS FOR SHOWING THE COMPOSITION OF WATER BY THE COMBUSTION OF HYDROGEN GAS.—*a* is a glass vessel held over the flame produced by the burning of the hydrogen gas.—See 336.

339. Fig. 48.—A WIRE-STAND.—See 293. The wire should be $\frac{1}{10}$ of an inch thick; the metal may be brass or iron. The foot is a solid lump of lead.

340. Fig. 49.—A JET-PIPE, made of brass. The lower end is a screw, adapted to the screw of a stop-cock, or a bladder.—See 323. The opening at the point of the pipe must be very small, like the jet of a blow-pipe. The length of this jet-pipe must be about three inches. It may be procured at the place where the stop-cocks are purchased. The price of it will be 1s. 3d.

341. Fig. 50.—APPARATUS FOR FILLING SOAP-BUBBLES.

WITH HYDROGEN GAS.—*a* is a piece of a common tobacco-pipe. *c* the end of a small brass cylinder into which the pipe is cemented. *b* the end of the brass cylinder in which there is a screw fitting a screw on the stop-cock of a bladder.—See 323. Before the end of the tobacco-pipe is put into the end of the brass tube, it should have a piece of thin oiled leather wrapped round it, to make the juncture air-tight. The piece of brass may be about an inch long; it can be made by the person who makes the jet-pipe, and will cost 1s.

342. Fig. 51.—APPARATUS FOR OBTAINING, WITHOUT THE AID OF THE PNEUMATIC TROUGH, GASES THAT ARE LIGHTER THAN COMMON AIR.—See the experiments on ammoniacal gas.

343. Fig. 52.—APPARATUS FOR OBTAINING WITHOUT THE AID OF THE PNEUMATIC TROUGH, GASES THAT ARE HEAVIER THAN COMMON AIR.—See the experiments on carbonic acid gas.

344. Fig. 53.—A MATRAS, OR BOLT-HEAD.—*a* is a glass tube six inches long, and wide enough to allow a sixpence to pass down it. *b* is a ball of rather more than two inches diameter blown at the end of the tube. The ball must be very thin at the bottom, as it is to be exposed to heat. For the use of this instrument see 265, 268. The glass blower charges about 1s. 3d. for it.

345. Fig. 54.—A FLORENCE FLASK; an instrument of great utility in performing the operations of digestion, solution, &c. Being made of very thin glass, it bears the sudden application of heat excellently; but, on account of *that* thinness, is very liable to be broken by a slight blow: it therefore requires to be handled carefully. The student should be provided with several of them. They are to be had of oilmen, who sell the empty flasks, after having disposed of the Florence oil they contained. Sometimes they are charged 4d. at other times 3d. and 6d.

346. Fig. 55.—A PIECE OF GLASS TUBE of the size generally employed to convey gases from the vessels they are formed in, to the receivers which are to contain them. *b, b*, show the thickness of the tube; *a, a*, show the width of the internal part of it. The instruments represented by

figures 30; 58; *e, e*, 57; *a*, 42; and 43; are all formed of tubes of this size.

Every chemist should be provided with an assortment of glass tubes of different lengths and thickness; for in various operations he is obliged to employ them, and whenever he wanted a bit of tube, it would be a very great waste of time to have to go and buy it. The student should go to a glass-house, and choose out about a pound weight of tubes, sorted. The thinnest about the size of a small quill, the thickest about an inch in width. The tubes marked *d, d*, fig. 57, should be $\frac{3}{4}$ of an inch in diameter.—These tubes cost from 3s. to 4s. a pound. The highest-priced ones being those which are very regular in the bore, and are used to form barometers and thermometers. For many purposes, green glass tubes will suffice, and they are much cheaper than the others. A considerable quantity may be bought at a bottle work for 1s.—Glass tubes may be easily cut to any required length by a small file, and (if thin) they may be bent (as fig. 30, and *e, e*, fig. 57, are, or to any other shape), by the aid of the blow-pipe. But, for boltheads, and other apparatus made of thick tubes, drawings must be given to a glass blower.—Occasionally, for conveying gases, metallic tubes are employed. Gases that have no action on metals (such as oxygen and hydrogen) may be economically transferred through tubes of tin, lead, or copper. Small gas-pipes are easily procured, are cheap, durable, and very flexible; they may be easily bent into any required shape.

347. Fig. 56, (*Plate 5.*)—THE COMMON STILL.—This figure is given to illustrate the ordinary method of distilling, and has been described already—see 271, article *distillation*.

348. Fig. 57.—WOULFE'S APPARATUS described at 294. The retort used with the apparatus is that represented by fig. 18. The receiver *a* is represented by fig. 19. The receivers *b, c*, may each hold 10 or 12 ounces, and will cost (without any of the tubes) about 2s. each. They are to be had at the glass houses.

349. Fig. 58.—A TUBE OF SAFETY.—This is a glass tube as thick as fig. 55, and of about sixteen inches in

length, bent as represented in the figure. One end of this is fastened into a cork, and the other made into the form of a small funnel, *a* is the cork, *b*, the funnel. In cases of distillation in which sudden absorption, or expansion may take place in the retort,—see 296,—this utensil is fitted into the tubulure, and the bended part of it (*c, c,*) is filled with water. Then, if a vacuum happens to be produced in the retort, the external air forces its way through the tube to supply that vacuum; and, on the contrary, if expansion takes place, the elastic fluid gains room by forcing the whole water between *c, c,* up into the straight part of the tube under the funnel *b*. A number of other uses to which the safety-tube is applicable, are described in particular experiments. It costs about 1s. 3d. when bought ready made, but it is easily formed out of a glass tube, which costs 9d.

350. Fig. 59. A BLOW PIPE.—For an account of the mode of using this instrument, see 279. The blow-pipe is generally formed of brass; a single plain tube without a bowl for condensing the vapour, may be bought for 1s. This is the kind that working jewellers use. The blow-pipe for very accurate experiments (young students need not have this) *must* have the enlargement, and be provided with moveable nozzles. Such an one will cost 5s. Chemists frequently employ blow-pipes made of glass, chiefly because it is more easy to give to glass than to brass a small smooth opening; but glass ones have the disadvantage of being easily broken. A glass blow-pipe resembling fig. 59, may be bought for 1s.

351. Fig. 60.—A PLATINUM BLOW-PIPE SPOON.—See 279 for an account of the use of this instrument. *a*, the bowl, should measure $\frac{1}{2}$ an inch across the top; *b* is the handle, made of wood; *c* is the shank, made of platinum, or sometimes (for cheapness) of silver. The whole length of the instrument should be about six inches. The metal platinum being almost infusible, bears uninjured a very strong heat; but it cannot be used with alkaline fluxes, because it has a strong tendency to combine with those substances. Whenever, therefore, these fluxes are employed,

charcoal, or spoons of silver, must be used. The platinum blow-pipe spoon costs 2s.

352. Fig. 6. (plate 6.)—A GAS-HOLDER.—(This apparatus is sometimes made sufficiently large to hold eight gallons; but *the student* need not get one larger than that which is described here.) *a* is the body, or reservoir, of the gas-holder; which is of a circular figure, and measures in diameter $8\frac{1}{2}$ inches, in height, 10 inches. It holds about nineteen wine pints. This reservoir is made of tin plate, as, indeed, are all the other parts of the apparatus, except where the contrary is mentioned; and the whole of it is japanned, both on the inside and the outside. *b* is a shallow cistern, fixed by four small supports over the reservoir at the height of four inches; *c* and *d* are stop-cocks, communicating with both the reservoir and the cistern; *d* just enters into the top of *a*; but, to *c*, there is a tin tube affixed, which goes to half an inch from the bottom of *a*. This tube is rather less than half an inch in diameter. The tops of the stop-cocks (or rather of the tubes in which they are inserted) are exactly level with the bottom of the cistern *b*. *e* is a funnel, the pipe of which is two feet long, and half an inch in diameter. It is made to fit the pipe of *c*; but can be readily taken away, when required. *f* is a glass tube, of a quarter of an inch bore, communicating at both ends with the reservoir. The manner in which it is properly fixed in its place is shown by fig. 62; where *ac* represents the body of the gas-holder; *b*, and *c*, two tin tubes projecting from it, and *d*, (dotted lines,) the glass tube, fastened, by glazier's putty, into the tin tubes. *g* is a short tube issuing from the bottom of the reservoir; and *h* is a stop-cock communicating with the top of the reservoir; *i* is a blow-pipe affixed to the stop-cock *h*, and serving to direct a stream of gas, upon a substance held to it on the stand *k*; *l* is one of the handles by which the machine is lifted. A corresponding handle is on the other side of the reservoir.

The following is an account of the cost of the gasholder of which the above is a description. 3 Stop-cocks, 3s. 6d.; brass screw for *g*, 1s.; glass tube *f*, 3d. To the tinsmith, for reservoir, cistern, funnel, for putting all to-

gether, and for japanning, 5s., *k* and *i* are supernumeraries. Such a blow-pipe as *i*, however, costs 1s.

353. When it is intended to fill this apparatus with gas, the funnel *e* is removed, (*that* being wanted in particular cases only), the tube *g*, and the stop-cock *h*, are closed, and the stop-cocks *c*, and *d*, are opened. Water is then poured into the cistern, and is thence conveyed into the reservoir, by the tube affixed to *c*; while the air in the reservoir is forced out through *d*. Thus, in a short time, the body of the gas-holder is filled with water and all the air is driven out of it. When this is the case, shut *c* and *d*, and open *g*, into which introduce the beak of the retort, or the tube, whence gas is issuing. The mode of introducing the tube into this opening is shown by fig. 67.—The water does not run out at the aperture *g*, when the screw is taken from it; because the stop-cocks, *c*, *d*, cut off the pressure of the atmosphere. But, as gas which issues from a tube put in *g*, rises through the water to the top of the reservoir, the displaced water, must, of course, run out at the opening *g*, where the gas-pipe is introduced. By looking in the tube *f*, we can readily see how much gas the reservoir, at any period, contains; and, in order to determine this with precision, we may affix a scale to the tube in the manner shown by the figure. When, while filling the reservoir with gas, we perceive, in the tube *f*, that it is nearly full, the beak of the retort, or other vessel, must be withdrawn, and the opening *g* must be closed by its screw. The gas-holder being now filled with gas, may be removed to any place where it may be required to make use of the gas.

354. If a jar such as fig. 37, is to be filled with the gas, the jar is first filled with water, and inverted, in the cistern *b*, (also filled with water,) over the stop-cock *d*:—then *d* and *c* are both opened; the water descends through *c*, and forces the gas out through *d*, into the jar, which, when filled, (the stop-cocks being then immediately closed,) may be removed from the cistern, as directed 292.

355. If the bladder *d*, fig. 35, is to be filled with the gas, let the stop-cock (*e*, fig. 35) affixed to that bladder, be screwed to the stop-cock *h* of the gas-holder. (The squeez-

ing of the bladder, &c., (see 291), being observed particularly.) Then pour water into the cistern, and open the stop-cocks, *e*, *h*, *c*, when the gas will be forced into the bladder.

356. When the funnel *e* is fixed into the stop-cock of the pressure of a column of water is obtained, and the gas in the reservoir is forced out through the stop-cock, which may be opened with considerable force. If a blow-pipe as *i*, is screwed on *h*, and a stream of oxygen gas driven through it, in the manner just described; and if a burning substance be exposed to this jet of gas, a heat of very great intensity is produced instantaneously. The stand *k* is employed to expose bodies to this blow-pipe: the top of it is of iron, and can be fixed at a greater or less height, by means of a screw. Perhaps there is no instrument which can enable any one to perform so many beautiful experiments as this gas-holder. A few more of the very many uses to which it may be applied, are described at 425 to 427.

357. Fig. 63.—DOCTOR URE'S EUDIOMETER.—“THE analysis of gases, by explosion with the electric spark, furnishes, when it can be applied, one of the speediest and most elegant methods of chemical research. The risk of failure to which the chemist is exposed, in operating with the simple tube, (see fig. 65.) from the ejection of the mercury, and escape or introduction of the air; or of injury from the bursting of the glass, by the forcible expansion of some gaseous mixtures, has given rise to several modifications of apparatus.

“A very simple form of the eudiometer occurred to me about two years ago; and this, which I have frequently used since that time, I can now recommend to the chemical world, as possessing every requisite advantage of convenience, cheapness, safety, and precision.

“It consists of a glass syphon, having an interior diameter of 1-4th of an inch. Its legs are each six inches long. The open extremity is slightly funnel shaped; the other is hermetically sealed; and has inserted near by the blow-pipe, two platinum wires. The outer end of the one wire is incurvated across, so as nearly to touch

edge of the aperture; (look at the top part of figure 1) that of the other is formed into a little hook, (see fig. 2) to allow a small spherical button to be attached to it, when the electrical spark is to be transmitted. The two legs of the syphon are from one-fourth to one-half of an inch asunder.

"The sealed leg is graduated by introducing successively equal weights of mercury;—137 grains of that metal occupy the space of the 1.25th of a cubic inch. (The graduations may be marked by a diamond, a file, or a piece of rock crystal). The instrument is then finished.

"To use it, we first fill the whole syphon with mercury or water, which a little practice will render easy. We then introduce into the open leg, plunged in a pneumatic trough, any convenient quantity of the gases, from a glass measure tube, containing them previously mixed in determinate proportions. Applying a finger to the orifice, we next remove it from the trough in which it stands, like a simple tube; and by a little dexterity, we transfer the gas to the sealed leg of the syphon. When we conceive enough to have been passed up, we remove the finger, and next bring the mercury to a level in both legs, either by the addition of a few drops, or by the displacement of a portion, by thrusting down into it a small cylinder of wood. We now ascertain, by careful inspection, the volume of included gas. Applying the fore-finger again to the orifice, so as also to touch the end of the platinum wire, we then approach the pendent ball or button to the electrical machine, and transmit the spark. Even when the included gas is considerable in quantity, and of a strongly explosive power, we feel at the instant, nothing but a slight shock or pressure at the tip of the finger. After explosion, when condensation of volume ensues, the finger will feel pressed down to the orifice by the superincumbent atmosphere. On gradually sliding the finger to one side, and admitting the air, the mercurial column in the sealed leg will rise more or less above that in the other. We then stir in this liquid metal, till the equilibrium be again restored, when we read off as before, without any reduction, the true resulting volume of gas.

“ As we ought always to leave two inches or more air, between the finger and the mercury, this atmospheric column serves as a perfect recoil spring, enabling us to explode very large quantities without any inconvenience or danger. The manipulation is also, after a little practice as easy as that of the single tube. But a peculiar advantage of this detachable instrument is, to enable us to keep our pneumatic trough, and electrical machine, at any distance which convenience may require; even in different chambers; which, in the case of wet-weather, or a dark apartment, may be found necessary to ensure electrical excitation. In the immediate vicinity of the water pneumatic cistern, we know how often the electric spark refuses to issue from a machine. Besides, no discharging rod or communicating wire is here required. Holding the eudiometer in the left hand, we turn the handle of the machine with the right hand, and approaching the little ball, explosion ensues. The electrician is well aware, that a spark so small as to excite no unpleasant feeling in the finger, is capable, when drawn off by a smooth ball, of exploding flaming combustible gas. Even this trifling circumstance may be obviated, by hanging on a slender wire, instead of applying the finger.”—*Extracted from a paper by Dr. Ure in the Transactions of the Royal Society of Edinburgh.*

358. Fig. 65.—VOLTA'S EUDIOMETER.—A strong glass tube $\frac{3}{4}$ of an inch in diameter, and twelve inches long. The bottom end of it is shaped like a funnel, in order that, when placed upright, it may stand firmly. In the top end, a piece of brass wire, or better, platinum wire is sealed hermetically. It must, like other eudiometers, be graduated.

When this eudiometer, containing the gaseous mixture that is to be detonated, is standing over water, introduce into it a brass wire; one end of which must terminate $\frac{1}{16}$ of an inch from the upper wire, and the other end of it project from the bottom of the tube, and be attached to the electrical apparatus, to form part of a circuit. The gas in the tube is exploded by completing the circuit externally, and thereby sending the electric spark from one wire to the other, in the tube. When the spark is taken

sed, the tube must be held firmly, either by the hand, or some other contrivance.—It will be seen that this eudiometer is much inferior to Dr. Ure's.

359. Fig. 66.—Part of an apparatus for ascertaining the quantity of carbonic acid discharged from any substance by addition of an acid. It is a piece of glass tube, of half the diameter of fig. 55, bent into a zig-zag form, (which may be easily done by the blow-pipe), with a cork fitted to one end, and the other end drawn out to a small point.

LIST OF CHEMICAL PREPARATIONS,

AND OTHER MISCELLANEOUS ARTICLES WHICH THE STUDENT SHOULD BE PROVIDED WITH, TO ENABLE HIM TO PERFORM THE EXPERIMENTS.

It is necessary, for experiments of research, where particular nicety is required, to be furnished with re-agents, &c., in the very greatest possible degree of purity; but, for ordinary purposes—for the performance of such experiments as the student finds either necessary or agreeable—

different substances may be made use of in the degree of purity at which they are sold by respectable druggists, and at Apothecaries' Hall. In the experiments in this

book, for which *pure* preparations are requisite, directions are given for purifying the preparations of *commerce*,

ACIDS.—Sulphuric, 4 oz. per oz. $\frac{1}{2}d$.—Muriatic, 4 oz. per oz. $1d$.—Nitric, 4 oz. per oz. $3d$.—Oxalic, $\frac{1}{2}$ oz., per $6d$.—Tartaric, 1 oz. $5d$.—The sulphuric, muriatic, and nitric, must be kept in bottles with glass-stopples—*note*: always wipe the neck of the bottle from which a strong acid has been poured, before replacing the stopple: this will keep you from burning your fingers or cloths.

ALKALIES.—Potass (caustic) 1 oz. $8d$. This is obtained in small sticks resembling slate pencils, which must be kept in a phial well secured from air.—Liquid Ammonia 2 oz. per oz. $2d$.—to be kept in a phial with a glass stopple.

METALS.—Iron filings, and thick polished wire.—Copper clippings—get some slips of thin copperplate from a tinsmith, which clean bright with pumice stone and

water, and clip into very small pieces with strong old scissors. Cut some of the same copper slips, into pieces $\frac{1}{2}$ an inch wide, and three inches long, and finely polish these are to be used in precipitating silver, &c. from solution.—Zinc, in lumps, may be bought of brass-foundry for 9d. a pound. If you cast plates for a galvanic pile you will want two pounds or so, of this. Zinc, granulated may be bought at chemists, at 1s. per pound. Half a pound will be needed for making hydrogen gas.—Gold leaf, a small quantity from a gold-beater.—Granulated Tin, (called by dyers *dropt tin*), 1 oz. 1d.—Tin foil, square foot, 3d.—Mercury, 1 oz. 6d. To be kept corked in a phial.—Bismuth, 1 oz. 6d.—Antimony, 1 oz. 6d.—Cobalt, 1 drachm 3d.

SALTS.—Acetate of lead, 1 oz. 4d.—Borax, 1 oz. 2d.—Carbonate of Ammonia, 3 oz. per oz. 2d.—Carbonate of Barytes (Native), and Carbonate of Strontia (Native), to be had (in small pieces) of those who deal in minerals.—Carbonate of magnesia (common magnesia) 1 oz. 3d.—Carbonate (super-carbonate of the druggists) of Potass 4 oz. per oz. 3d.—Sub-carbonate of potass (the potash of the dry-salters, of whom it may be bought) 4 oz. per oz. $\frac{1}{2}$ d. This must be purified before use, and must be kept from air, in a stone jar.—Carbonate (super-carbonate of the druggists) of soda 1 oz. 6d.—Sub-carbonate of soda (the *soda* of the dry-salters) 4 oz. per oz. $\frac{1}{2}$ d. The soda of soda, like those of potass, must be kept from air.—Fluate of Lime (fluor spar) to be had of dealers in minerals.—Muriate of Ammonia, 2 oz. per oz. 2d.—Nitrate of Potass (saltpetre) 4 oz. per oz. $\frac{1}{2}$ d.—Prussiate of Potass, 1 drachm, 3d.—Sulphate of Alumina and potass (Alum), 4 oz. per oz. $\frac{1}{2}$ d.—Sulphate of Copper (Blue Vitriol) 1 oz. 2d.—Sulphate of Iron (Green Vitriol) 4 oz. per oz. $\frac{1}{2}$ d.—Sulphate of Magnesia 2 oz. per oz. 1d.—Sulphate of Soda, 2 oz. per oz. $\frac{1}{2}$ d.—Cream of Tartar 1 oz. 1 $\frac{1}{2}$ d.

MISCELLANEOUS ARTICLES.—Alcohol, 4 oz. per oz.—Olive Oil, 1 oz. 2d.—Oil of Turpentine, 2 oz. per oz.—Phosphorus, $\frac{1}{4}$ oz. per oz. 6s.—The manner in which it is preserved, is mentioned at 700.—Roll sulphur, flow

of sulphur, chalk, pipe-clay, red-lead, of each a penny worth.—Pieces of white marble.—Sulphuric Ether, $\frac{1}{2}$ oz. 6d. To be kept in a well closed phial, in a cool place.—Turmeric, $\frac{1}{2}$ oz. $1\frac{1}{2}d.$ —Litmus, $\frac{1}{2}$ oz. 3d.—Nut-galls, $\frac{1}{4}$ oz. $1\frac{1}{2}d.$ —Oxide of Manganese, a-pound 6d. or 8d. :—This is to furnish oxygen.—Sulphuret of Antimony 1 oz. 3d.—Gum benzoin, $\frac{1}{2}$ oz. $1\frac{1}{2}d.$ —Some charcoal, newly made.

SMALL ARTICLES OF APPARATUS.

AN iron ladle.—A deep iron pan holding about a quart : to be filled with sand—this is to serve for a *sand-bath*.—A magnifier.—An artificial magnet.—Glass rods, and clean straws, for stirring mixtures in glasses.—A collection of glass tubes :—see 454.—corks of all sizes ; and a considerable number of spare phials should be constantly at hand, to receive the products of experiments.—About a dozen from 2 to 4 ounce phials, with ground glass-stopples, should be provided, to hold corrosive liquids.—Bottles with wide mouths, or jars with tin covers, such as are seen on the shelves of apothecaries' shops, should be provided, to hold salts and substances that require to be kept from the air.—A number of small wooden and pasteboard boxes will also be found convenient.

EXPERIMENTS.

~~~~~  
\* \* *It is earnestly recommended, that the student will perform no experiment without, at the same time, making himself acquainted with the chemical history of the substances employed, as detailed in the preceding pages.*  
~~~~~

SECTION I.—ON THE PROPERTIES OF HEAT.

360. HEAT AND COLD PRODUCED BY THE SAME LIQUID AT THE SAME TIME.—*Process.* Put your right hand into a basin, containing water made as hot as you can well bear it, and put your left hand into a basin containing cold water. After a few minutes, take out both hands, and instantly plunge them into water warmed moderately: what effect will be produced? The water will cool your right hand and warm your left.

361. *Explanation of this experiment.*—What we call heat, is the effect produced by the presence of the peculiar substance which chemists call *caloric*. Cold is merely a negative quality: it signifies the absence of heat, or rather, a diminution of heat. This producer of heat (*caloric*), always tends to an equilibrium; that is to say, heated bodies placed among cool ones, always part with their heat to the cool ones, till all are brought to the same temperature. Of course, by such a process, the cool bodies are heated, and the heated ones, cooled. This doctrine of the distribution of heat enables us fully to comprehend the phenomenon of the above experiment. A hot hand put into cold water, communicates a part of its heat to that water, and thus becomes cooled. Again, a cold hand put into hot water, takes a portion of heat from that water, and consequently is heated. Hence, we readily discover

how water of a medium temperature heated the hand that had been cooled by cold water, and cooled the hand that had been heated by hot water.

People very generally imagine that the sensation of heat is an accurate test of temperature, and they are thereby frequently led to misal things. They come into this room from the open air to-day, and exclaim "How warm it is!" To-morrow, they will again come into it from a still warmer room, and will cry "How cold it is!" In the first case they gain heat, and therefore call the room warm; in the latter case, they lose heat, and then they term it cold; while, in reality, the air of the room continues, during the whole time, heated precisely to the same degree of temperature.

Two men were travelling on a high mountain; one of them was ascending, the other, descending it. About the middle they met. "Bless me!" exclaimed he who was going down, "how extremely hot it is to-day!" "Hot?" cried the other, "why, I never felt so cold in all my life." These two men judged from their sensations, and truly expressed what they felt. At the top of the mountain, the air was cold; at the bottom of it, the air was warm. He who was descending, came, therefore, into warm air, and was heated; on the contrary, he who was ascending, was, by coming to the cool air, cooled. We learn from this, that our sense of feeling can never inform us respecting the true temperature of the bodies by which we are surrounded.

362. HEAT EXPANDS BODIES, as the following experiments will prove.—*Process 1.* Take a piece of iron that exactly fits a ring made to receive it. Let it be made red-hot. It will be then so much enlarged in bulk, that it will not go into the ring.

363. *Process 2.* Put into the bulb of the instrument represented by fig. 42, some cold water, or alcohol. Then, holding the part *a* of the instrument in the hand, plunge the bulb into hot water: upon this, the enclosed liquid, as it gains heat from the hot water, will rise in the tube. The expansion will be seen more clearly, if the liquid in the bulb be coloured.

364. *Process 3.* Let a bladder, partly filled with air, be tied at the neck, and held near a fire. The small quantity of air inclosed will speedily expand, and fill the bladder completely.

* * * That, in all these cases, the expansion of the different bodies is really occasioned by the caloric they acquire, is very evident, from the circumstance, that, when they become cool again, they become small again.

365. *Remarks on this phenomenon.*—Of all the properties of heat, none is more remarkable than its power of expansion. Matter of every description, bodies in every state, are, by an accession of caloric, enlarged in bulk; and, on the contrary, when bodies are deprived of heat, they are reduced in bulk.

In order to account for so curious a phenomenon as increase of size by the addition of invisible matter, philosophers have supposed caloric to be a peculiar fluid, possessed of so subtile a nature, as to be capable of forcing itself between, and consequently of driving asunder, the particles of matter of even the very densest kind. By this means it is, that it is able to overcome the attraction of cohesion, and thereby to dilate bodies.

The fact shown by *Process 1*, (362) is taken advantage of by coopers in fastening the staves of a cask with iron hoops: before the hoops are put on, they are heated, and of course enlarged; they then, on cooling, contract, and the cask is tightly bound.—The expansion of fluids (*Process 2*, 363,) has been very usefully applied in the construction of *thermometers*, by which it is made the means of measuring the variations in the quantity of caloric contained in other bodies.

366. A THERMOMETER (or, *measurer of heat*,) is a small tube of glass, with a bulb at the end of it; the bore of the tube is very small, and perfectly cylindrical; the bulb is entirely, and the tube partly, filled with the fluid by whose expansion the degrees of heat are to be measured. The tube is applied to a scale, formed on fixed principles, and the expansion of the fluid is thereby measured with accuracy. The fluid with which thermometers are generally filled is the metal mercury, *that* being found to

combine more advantages than any other fluid. The scale just mentioned, is divided into two hundred and twelve equal portions, which are called degrees; the two hundred and twelfth degree (usually written 212°) being the topmost one. The mark at the bottom of the scale, near the bulb, (0°) is called *zero*.

367. When the thermometer is immersed in melting ice, the surface of the mercury stands invariably at the 32d degree, (shortly said, at 32°), and this is on that account called the *freezing point of water*. When the thermometer is plunged into boiling water, the mercury rises to 212° , which, therefore, is called the *boiling point of water*. All bodies that are as hot as boiling water, make the mercury rise as high as boiling water does; and all bodies that are as cold as melting ice, (or freezing water) make the mercury sink to 32° . If the thermometer is immersed in a mixture of equal parts of water at 32° and 212° , the mercury stands level with 122° ; we have, in this case, an excellent illustration of what has been said (361) regarding the tendency which caloric has to an equilibrium: the hot water communicates 90° of heat to the cold, and its own temperature is thereby reduced 90° .

368. The thermometer does not tell the precise quantity of caloric contained in the bodies it is applied to; it only shows by how many degrees of heat the temperature of certain substances exceeds that of others. For we cannot deprive any body of all the caloric it contains. The mercury rises by abstracting heat, and falls by communicating it. If the mercury is sunk to 0° by being plunged into a certain mixture, still we cannot say that that mixture is deprived of heat, since it is evident, that the mercury fell upon giving out part of *its* heat to the mixture, and became stationary upon the production of an equilibrium. Though the lowest point on the thermometer scale is 0° , yet cold can be produced by many degrees more intense than that; so, also, can much greater heat than that of boiling water be produced. Thermometers for chemical purposes are sometimes marked to 600° above 0° and to

40° below it; which marks show the boiling and freezing points of mercury.

369. We must now notice a very remarkable exception to the laws of expansion upon increase of heat. When water at 32° is heated, instead of expanding as other bodies in the same situation would do, it becomes denser, continuing to do so, more and more, till it arrives at 40° after which it regularly expands! When water at 212° is cooled, it contracts till it comes to 40°, and then it gradually expands till it comes to 32°! This property of water is a very beneficial one, as we shall presently see.

370. DIFFERENT BODIES EXPAND IN DIFFERENT DEGREES. As may be proved by the following *Process*.—Fill the bulb of the instrument, fig. 42, with water, and fill the bulb of another instrument exactly like that with alcohol. The two liquids had better be coloured by some vegetable infusion. Now, take the two tubes, and plunge the bulbs together into hot water: both the liquids will expand and rise in the tubes, but the alcohol with far more rapidity than the water. The cause of the difference in the expansibility of bodies is unknown.

371. HEAT PASSES QUICKLY THROUGH SOME BODIES SLOWLY THROUGH OTHERS. All bodies through which heat passes are called *conductors* of it; those through which it passes *quickly* are termed *good* conductors, and those through which it passes *slowly*, *bad* conductors. Dense bodies are, in general, the best conductors; light porous bodies, the worst. Bodies which are the worst conductors of heat, have the greatest capacity for it; that is to say, they take a greater quantity of heat than good conductors, to raise an equal bulk to a given theometrical temperature, within a certain time. When bodies are compressed, heat is evolved; because, as, by their condensation, their conducting powers are increased, so are their capacities for heat diminished. The following experiments *prove* the conducting powers of bodies to be different.

Process 1.—Take a rod of iron a foot long, and a rod of wood just the same size. Put one end of each into a

fire, and hold the other extremities by the hands. When the one end of the iron has become red hot, the other end will be almost unbearably hot; but the wood will scarcely give any heat to the hand, though held till nearly all consumed by the fire. Hence we learn, that iron is a good conductor of heat, but wood, a bad one.

372. *Process 2.*—Prepare a number of equally-sized rods of different substances, copper, lead, tin, iron, glass, bone, and wood of various sorts. Coat one end of each, by dipping it into melted wax or tallow. When they are all ready plunge their uncoated extremities into boiling water, or hot sand,—in a short time the wax or tallow will be melted; but it will be observed, that all the coatings do not melt together, but, that of each particular rod in the order of its power of conducting heat. The metallic rods (these too in their particular order) produce the effect first, next the glass, and last of all the wood.

373. *CURIOUS MOTION PRODUCED IN LIQUIDS BY HEATING AND COOLING THEM.*—Fill a large phial with water, and put into it a small quantity of powdered amber, having previously added to the water a sufficient quantity of potash, to make a solution of the same specific gravity as amber. Immerse the phial in a glass of hot water; upon which a very singular internal motion will be immediately perceived. A current of the fluid will rise up the sides of the phial, and another descend in the centre of it. Now take the phial out of the hot water, and observe the effects of its cooling. The currents will be reversed: the external one will descend, and the internal one ascend.—The use of the powdered amber is to make the opposite currents into which the water is thrown, visible.

374. *Rationale of this Experiment.*—The manner in which a quantity of water, put in a vessel over a fire, is heated throughout, is by the constant agitation of its particles; for though heat is transmitted from one particle of a fluid to another in the same manner as in solid bodies, yet it is only so in a very small degree. As the particles of water immediately at the bottom of a vessel are expanded by heat, they become, of course, specifically lighter than the rest of the fluid; and consequently rise to the

surface, communicating, in their progress, part of their heat to the colder particles around them. Successive portions are thus heated at the bottom, and rise to the surface; and this agitation continues till all the liquid is brought to the greatest degree of heat it can acquire, namely, to its boiling point. If a liquid is heated at the surface, the heated particles being specifically lighter than those below, cannot descend, and therefore no agitation can be produced to communicate the heat to the bottom of the water. It was once asserted that heat could not descend, and in apparent proof of this, the upper surface of a vessel of water can be boiled and evaporated, while a cake of ice remains frozen at the bottom. See 377. But, it has been proved that heat is propagated downwards, (by transmission from particle to particle,) though but slowly.

In the experiment just performed, the water rises at the sides—this is because the heat is communicated through the medium of the sides of the bottle. The descending current in the centre, is occasioned by the sinking of the water which parts with a portion of its heat, to the atmosphere, at the surface. When the fluid was held out of the water, the currents were reversed, because the external particles being in contact with cool air, are cooled; they therefore descend and force up the central particles, which are lighter because warmer.

From what is said at 369 and 374, it will be seen, that water in large quantities, as, for instance, in deep lakes, when cooled by the atmosphere in winter, sinks in successive portions, till the whole bulk of it is of the temperature of 40° ; after which, as it cools to 32° , (its freezing point,) it continues unagitated, that is, its cool particles, being lighter than its warm ones, remain at the surface, and are at last converted into a sheet of ice. How manifest is the wisdom and goodness of the GREAT ARTIFICER of the world in this arrangement! If water continued to condense to 32° , the water on the surface of our rivers, would sink as it froze, another sheet of water would freeze immediately, and sink also; the ultimate consequence of which would be, that the beds of our rivers would become depositories of immense masses of

which no subsequent summer could unbind; and the world would have been shortly converted into a frozen mass!

375. TO PROVE THAT WATER EXPANDS WHEN FREEZING.—Fill a small phial with water, cork it securely, and place it in a situation where the water in it may be frozen: whenever solidification takes place the bottle will burst.

The force with which water expands when in the act of freezing, is immense. A small brass globe, which would have required a force equal to 27700 pounds to have burst, has been burst by the freezing of a little water in it.

The expansion of water during frosts, trees and rocks are often split asunder. Slate is dug from quarries in large blocks, which are placed, edge uppermost, exposed to the rain; this penetrates their fissures, and, when a frost takes place, expands, and splits the slate into thin layers.

376. TO SHOW THAT HOT WATER IS LIGHTER THAN COLD.—Pour gently hot water into a tall glass nearly filled with cold water; it will remain on the surface: but if cold water be poured upon hot water, it will sink to the bottom. This experiment may be rendered more striking by colouring that portion of water which is poured

377. TO CAUSE WATER TO BOIL ON THE SURFACE OF ICE.—To effect this, first freeze a quantity of water in the bottom of a long glass tube, closed at one end, either by exposure to cold air, or by means of a *freezing mixture* (of which we shall speak presently). Then cover the top of the tube with ice by a quantity of water, and hold the tube (without handling the part of it containing the ice) in such a manner over a lamp, that the surface of the water may be heated to the point of boiling: for this the tube requires to be placed in a diagonal direction, which is such as to allow the water at the top of it to be heated, while the bottom remains unheated below.

378. SUDDEN CONVERSION OF A LIQUID INTO AN INSOLUBLE FLUID OR VAPOUR, AND RE-CONVERSION OF IT INTO A LIQUID.—Put into the instrument, fig. 42, two or three spoonfuls of sulphuric ether, and then fill it (tube and

all) with water. The instrument should, for this experiment, be sufficiently large to hold about half-a-pint, and the water may be coloured. Let the mouth of the tube (the bulb being turned upwards) be put into a vessel of water: it may be kept in this situation, by being fixed into a hole made in a little stool; just as it is represented by the figure above referred to. Next pour boiling water gently upon the top of the ball: the sudden addition of heat will instantly change the ether into a vapour which, by its expansion, forces the coloured water out of the vessel, and occupies its place. It is proper, however, not to allow the whole of the water to be forced out; but after expansion has proceeded to a certain degree, to pour then *cold* water upon the vessel. This will immediately re-convert the vapour into a liquid, and cause the water to rush up into the bulb, to supply the vacuum.

379. *Rationale*.—This experiment is intended to show that heat has great influence on the form or state of bodies. When solids are heated, they become liquids, as is proved in every instance of fusion; and when liquids are heated they acquire the gaseous form, and become invisible elastic fluids, possessed of the mechanical properties of common air. They retain this form or state as long as their temperature remains sufficiently high, but re-assume the liquid form when cooled again. Different fluids pass into the æriform state at different temperatures: the temperature at which a liquid becomes changed into a vapour, is called its *boiling point*.

380. TO CAUSE WATER TO BOIL BY THE APPLICATION OF COLD, AND TO CEASE TO BOIL BY THE APPLICATION OF HEAT.—Provide a Florence flask, a good cork, that fits it closely, and a piece of bladder, softened by having been soaked in water. Fill the flask half full of water and place it over a lamp, by means of the stand, fig. 2, till the water boils; then remove the flask, instantly cork it securely, and tie the moistened bladder over the neck and cork, so as to prevent the least access of air. Till the boiling of the water, which continues a little while after the flask is removed from the lamp, soon ceases entirely; but it will immediately recommence, if the flask is plung-

to cold water; and cease again, when the flask is held near the fire.

381. *The explanation of the experiment is this:—*The boiling points of liquids are regulated by the pressure of the atmosphere. The greater the pressure, the greater is the degree of heat requisite for making the liquor boil; and, consequently, the degree of heat required to cause the ebullition of a liquid is small, under a small pressure.

If the pressure on the surface of the earth could be entirely removed, every substance, whatever, would begin to boil instantly.

Now, in what manner do these facts explain the curious phenomena of the above experiment? Why, when the flask is taken from the lamp, in order to be closed, the upper part of it was filled with vapour, the quantity of which was increased by that generated during the short time that the water continued to boil, after being removed from the lamp; and it was the great pressure of this body of vapour, which made the still hot water cease to boil. The boiling recommenced when the flask was cooled, because, by that cooling, the vapour was condensed into water, and a vacuum formed. This being the case, the water will boil

at a very low temperature; and the heat that it retains is sufficient for the purpose. The boiling of the water ceased again, upon the flask being heated, because the vapour formed by that means renewed the pressure on its surface.

* * * Sometimes Florence flasks are so thin, that, when the vapour is condensed, the glass is broken by the pressure of the atmosphere. The following is another mode of showing this experiment, calculated to obviate this inconvenience. Provide a six-ounce medicine phial, half fill it with water, and place it in a pan of boiling water over a fire, till the water it contains boils also. Then cork it closely. Place this phial, thus prepared, in a tea saucer; pour cold water upon it; when the water in the phial will boil—next, pour hot water upon it, when the water in the phial will cease to boil.—The *rationale* need not be repeated.

382. TO FREEZE WATER BY PUTTING IT INTO A PAN ON THE FIRE.—Prepare a FREEZING MIXTURE, as direct-

ed in the APPENDIX, and place the pan containing over the fire; then immediately plunge into it a tube thin glass, about $\frac{3}{8}$ of an inch in diameter, containing small quantity of water. In a short time, by the act of the freezing mixture, the water in the tube will be converted into ice.—Care must be taken not to let it remain too long, or the ice after being formed will be melted.

Rationale.—During the liquefaction of bodies, a quantity of heat is absorbed, which is essential to the state of liquidity, and which does not increase the sensible or thermometric temperature. The same may be observed of the conversion of liquids into vapours. Consequently, if a cold solid body, and the same body hot and in a liquid state, be mixed in known proportions, the temperature after mixture will not be the proportional mean, as would be the case if both were liquid, but will fall short of it, much of the heat of the liquid body being consumed in rendering the solid body *liquid*, before it produces any effect upon its sensible temperature.

Equal parts of *water* at 32° , and of *water* at 212° , will produce, on mixture, a mean temperature of 122° . If equal parts of *ice* at 32° , and of *water* at 212° , will only produce (after the liquefaction of the ice) a temperature of 52° , the greater portion of the heat of the water being employed in thawing the ice, before it produces any rise of temperature in the mixture. Thus the water is cooled 160 degrees, while the ice is heated only 20; consequently 140 degrees of heat have disappeared, this having been united to the solid water (ice) to produce fluidity. Heat thus rendered insensible, or combined, is termed *latent heat*.

The same phenomena are observable in all cases of liquefaction, and we produce artificial cold, often of great intensity, by the rapid solution of certain saline bodies in water. Upon this principle the action of freezing mixtures depends.—In the above Process, the water becomes solid, by giving out its *heat of fluidity* to the mixture.

The solution must be boiled some time, filtered, and evaporated till a pellicle is formed on the surface; when by slow cooling, it deposits crystals.

383. TO FREEZE WATER IN A FEW MINUTES, EVEN

THE MIDST OF SUMMER.—Take a thin glass tube, four or five inches long, and two or three eighths of an inch in diameter. This must be closed at one end, and have water poured into it, to the height of an inch. Now, by means of the dropping tube, (see 303,) let a stream of sulphuric ether fall upon that part of the tube where the water is. The ether, as soon as exposed to the atmosphere, rapidly evaporates; but, in order to be converted into vapour, it requires a considerable quantity of caloric; and it therefore immediately robs the water in the tube, of its heat of fluidity. The consequence is, that the water is changed to ice; and if a thin spiral wire (such as fig. 39) be previously put into the tube, the ice will adhere to it, and may be drawn out.

Cold is produced in all cases of evaporation. The inhabitants of warm climates cool their liquors for drinking, by wrapping the vessels containing them in wet cloths, and hanging them up in the sun. The water in the cloth evaporates quickly, and thus produces cold. A person whose clothes are wet, feels cold, even when near a fire; it is because the water, as it evaporates, robs his body of caloric.—Under certain circumstances, the cold produced by evaporation is very great, being even sufficient to freeze water. This has been proved by the above experiment; and is also most elegantly shown by that which follows.

384. DR. WOLLASTON'S CRYOPHORUS.—This instrument is described at 334. It was invented to demonstrate the relation between evaporation at low temperatures, and the production of cold. To make use of it, prepare a freezing mixture, and plunge the ball *b* into it; upon which, the water in the other ball will be frozen in a few minutes.

By referring to 334, and to 381, we see, that the instrument, as it was closed while the water in it was boiling, is filled with vapour. This vapour, when the ball *b* is plunged in the freezing mixture, is condensed by the common operation of cold; and the vacuum produced by this condensation, gives opportunity for a fresh quantity of vapour to rise from the opposite ball. Now, the small quantity of water which rises from *a* to supply this va-

cuum, takes, in order to be converted into vapour, a large quantity of heat from the remainder of the water ; and it is by the reduction of temperature thus effected, that the water is eventually changed into ice.

385. INSTANTANEOUS CRYSTALLIZATION : A CURIOUS EXAMPLE OF THE PRODUCTION OF HEAT, BY THE CONVERSION OF A LIQUID INTO A SOLID.—Into two ounces of boiling water, put as much sulphate of soda, as it will dissolve (about 3 ounces). Pour as much of this saturated solution, when boiling hot, into a phial, as will nearly, but not quite, fill it; cork the phial closely, and let it stand to cool. When cold, the solution is still fluid, but the instant you draw the cork, a very beautiful, but confused crystallization, of the whole mass, will immediately take place ; and, at the same time, so much heat is evolved, as to make the phial warm.

The explanation of the experiment is this : water will dissolve more sulphate of soda when hot than when cold ; and cold water will dissolve more in proportion as the pressure of the atmosphere is diminished. The hot water was here saturated, and, had it been suffered to cool in an open vessel, would have deposited part of the salt. But, in this case, none was deposited, for by suffering the solution to cool in a close vessel, a partial vacuum was produced at the surface of it, (the steam which occupied the top part of the phial when the cork was inserted, being, by the subsequent cold, condensed,) and the water when cold, was thus enabled to hold in solution, all the salt which, when hot, it had dissolved. As soon, however, as, by drawing the cork, you admitted the usual pressure of the atmosphere, the cold water was rendered incapable of holding so much salt in solution, and part was, therefore, instantly crystallized. The heat which was evolved was the heat of liquidity of the portion of the salt which thus became solid. If, when the salt has crystallized, you plunge the phial containing it into hot water, it will be again dissolved. You may then cork the phial, as before, and the same solution will serve for a repetition of the experiment.

386. TO PRODUCE A BOILING HOT LIQUID BY MIXING

TOGETHER TWO COLD ONES.—Take a small phial about half-full of cold water; grasp it gently in the left hand, and from another phial pour sulphuric acid very gradually into the water. The mixture will immediately become so hot, that the phial cannot be held.—If a thin glass tube, three-eighths of an inch in diameter, containing a small quantity of water, be plunged into a mixture of one part water to four parts acid, the water in the tube will *boil*.—*Rationale of this experiment*.—Whenever two liquids unite chemically, the compound has greater density than the mean density; and whenever the density of bodies is increased, they evolve heat.

SECTION II.—ON CHEMICAL AFFINITY.

387. TO SHOW WHAT CHEMICAL AFFINITY IS.—*Process*. Shake together, in a small phial, a quarter of an ounce of olive oil, and the same quantity of water; then allow the mixture to settle, upon which the oil and water will separate. Now, add to the mixture a drachm of a solution of an alkali (either potass or soda) and shake the phial again. Upon this, these three bodies will form a substance resembling thick cream.—*Rationale*.—Oil has no affinity for water, and, therefore, does not combine with it; but, it has a strong affinity for alkalies, with which it readily combines, and forms *soap*.—The cream-like appearance of the above mixture, is occasioned by the diffusion of the soap that is formed, through the water.

388. PROOF THAT A SUBSTANCE HAS DIFFERENT DEGREES OF AFFINITY FOR DIFFERENT SUBSTANCES.—To the product of the preceding experiment, add a little diluted sulphuric acid. The soap will instantly be decomposed. For the acid combines with the alkali (having a stronger affinity than the oil for it,) and forms a *salt* which is dissolved in the water: while the liberated oil rises to the surface of the mixture.

389. CURIOUS EXAMPLES OF CHEMICAL AFFINITY.

Process 1.—Take a little solution of sulphate of iron, and a little infusion of galls, both diluted till colourless,

pour them together,—the mixture will be black.—*Rationale.* The gallic acid contained in the infusion of galls has a stronger affinity for iron than sulphuric acid has. Consequently, the sulphate of iron is decomposed, and gallate of iron immediately produced.

390. *Process 2.* Into the black liquor, which is the product of the preceding experiment, pour a little diluted muriatic acid: the mixture will immediately become colourless.—*Rationale.* Muriatic acid has a still stronger affinity for iron than even gallic acid. We, therefore, in this experiment, form muriate of iron, which gives a colourless solution.

391. *Process 3.*—To the limpid liquid produced by the last experiment, add a little solution of potass, (which is colourless,) the mixture will again become black.—*Rationale.* Here the muriatic acid quits the iron to unite with the potass, and the disengaged iron is caught up by the gallic acid which remains in the solution, to re-produce the black gallate.

392. *Process 4.*—Immerse a piece of gold—a coin will do—in a glass of mercury. After a few seconds take it out: it will be apparently transformed into silver. Drop the silver-looking metal into a little diluted nitric acid: upon taking it out, after a little while, it will be found to have resumed its natural appearance.—*Rationale.* Mercury has a strong affinity for gold, so that amalgamation takes place whenever the two metals are placed in contact. The coin, in the above case, becomes cased in an amalgam of gold. The nitric acid, into which the metal is afterwards put, dissolves the mercury, but does not act upon the gold.

393. *Process 5.*—Take an ounce of mercury and an ounce of sulphur; melt them together in a crucible, stirring the mass continually—pour the mixture on a piece of marble or glass, greased and warmed. The substance thus obtained is sulphuret of mercury, which, if you have an alembic, you may sublime, (see 318,) and you will then form the beautiful pigment called vermilion.—*Rationale.* This experiment is an example of simple affinity, by which bodies combine and form a substance differing entirely from its constituents. Mercury is white, brilliant, and

luid, and sulphur is yellow. Now, the body formed by their union is a solid of a red colour.

394. *Process 6.*—Take equal parts of muriate of ammonia and lime, (newly burnt,) these bodies are inodorous: triturate them in a mortar; upon this, a very pungent smell is immediately produced. If the operation is performed quickly, and the mixture put in a bottle, well closed, it serves as a smelling bottle for a long time.—*Rationale.* The muriatic acid which is a constituent of the muriate of ammonia, has a stronger affinity for the lime, than for the ammonia. It therefore combines with the lime, and yields up the ammonia in the state of gas. It is this ammoniacal gas which has the pungent smell.

395. ORDER OF THE AFFINITIES OF SOME OF THE ACIDS FOR POTASS.—*Acetic greater than Carbonic.* Put some solution of carbonate of potass into a tumbler, and pour over it a little acetic acid. The carbonic acid will be expelled with effervescence, and acetate of potass will remain.

396. *Muriatic greater than Acetic.*—Into the newly-formed acetate of potass, pour muriatic acid as long as acetic acid (known by the smell) continues to be evolved. The new compound, muriate of potass, is a salt that may be crystallized.

397. *Nitric greater than Muriatic.*—Into the solution of muriate of potass, pour nitric acid; the muriatic acid (which has a very pungent smell) will be expelled; and nitrate of potass (saltpetre) will remain.

398. *Sulphuric greater than Nitric.*—Pour sulphuric acid into the solution of nitrate of potass obtained in the last experiment. A solution of sulphate of potass will result.

* * * If this series of experiments was performed in a retort, and the gaseous acids were discharged thence into proper receivers, and examined by tests, in the manner hereafter described, the results would be far more satisfactory.

399. COMPARATIVE AFFINITIES OF SEVERAL SUBSTANCES FOR SULPHURIC ACID.—*Ammonia greater than Iron.*—Into a solution of sulphate of iron, contained in a test glass,

drop as much liquid ammonia as will precipitate the whole of the oxide of iron. The *then* solution will be sulphate of ammonia.

400. *Magnesia greater than Ammonia*.—Having decanted the solution furnished by the last experiment into a clean tumbler, stir in it as much carbonate of magnesia (common magnesia) as can be dissolved. The magnesia will be separated from the carbonic acid, which flies off in the state of gas, as likewise does the ammonia; and there remains in the solution, sulphate of magnesia (Epsom salt.)

401. *Soda greater than Magnesia*.—Into the solution of sulphate of magnesia, pour a solution of carbonate of soda until the whole of the magnesia is precipitated. The carbonate of soda is decomposed: the carbonic acid unites to the precipitated magnesia, and the soda with the sulphuric acid form sulphate of soda, which remains in solution.

402. *Potass greater than Soda*.—Pour a solution of carbonate of potass, till the commencement of effervescence, into the solution of sulphate of soda. Here the sulphuric acid seizes the potass, and the liberated carbonic acid combines with the soda. There is no precipitate; because both the compounds are very soluble. But, that decomposition certainly takes place, may be proved by crystallizing the product of the experiment, when the salts may be distinguished from each other.

403. *Strontia greater than potass*.—Into the solution of sulphate of potass, pour a solution of pure strontia in cold water or of carbonate of strontia in hot water: sulphate of strontia will be precipitated.

404. *Barytes greater than Strontia*.—Dissolve the last mentioned precipitate in boiling water, and pour in a solution of barytes, or of muriate of barytes. The sulphuric acid will now make one more election—seizing on the barytes, and forming with it a very insoluble salt, sulphate of barytes.

* * * The foregoing experiments, showing the comparative affinities of different substances for one of the strong acids, form a very interesting lesson. We recommend the

young student to perform them with care ; and, in order to prove that the products do really contain the bodies specified, he may try by means of appropriate tests.

405. MUTUAL DECOMPOSITION OF TWO SALTS—AN EXAMPLE OF COMPOUND AFFINITY.—Prepare a solution of carbonate of soda, and a solution of muriate of barytes, (both transparent liquors),—mix them together in a test glass ; this done, a very heavy white powder will be thrown down immediately. Here, a mutual decomposition, and formation of two new salts, carbonate of barytes, and muriate of soda, take place. The former is precipitated : whilst the latter remains in solution, and may be crystallized into that salt known by the name of *table salt*.

406. THE DENSITY OF BODIES IS ALTERED BY CHEMICAL ACTION.—*Process 1.*—If two cubical inches of copper, and the same bulk of tin, are melted together, they form an alloy of the size of only three cubical inches. One-fourth of their bulk, therefore, is lost ; while, in weight, they are unaltered.

407. *Process 2.*—If one part of sulphuric acid, be mixed with three parts of water, the compound will occupy less space than the bodies did when separate.

SECTION III.—ON GASES.

408. TO PROCURE OXYGEN GAS.—Put into the retort, fig. 18, an ounce of black oxide of manganese, in powder, and pour over it as much sulphuric acid as will convert it into a thin paste. Stir the mixture with a glass rod. Proceed next as directed, 282. Give a very gentle heat. The gas will shortly be evolved ; and may be collected in jars as directed 284, 317. Or, the gas may be collected in the gas-holder, fig. 61.—See 353. There are several other methods of procuring oxygen gas ; but this is the one generally adopted, for small quantities.

409. *Note.*—The first portion of gas (of whatever kind it may be) that is evolved from the vessel in which it is formed, is always contaminated with the common air, with which the said vessel was filled in the first instance. A quantity of the first air received, rather more than the ca-

capacity of the vessel in bulk, must, therefore, in order to avoid accidents and failures, be thrown away.

410. TO PROVE THAT "OXYGEN GAS IS AN EMINENT SUPPORTER OF COMBUSTION." *Process 1.*—Plunge a lighted candle into a quantity of the gas contained in a glass in the manner directed at 288. The cover to confine the gas till it is wanted, may be a piece of glass, or paste-board.

411. *Process 2.*—If the light of a taper be blown out and the taper be let down into a glass of this gas while the snuff (which should be a thick one) remains red hot it instantly rekindles, with a slight explosion. When the taper is re-lighted, it continues to burn, as in the preceding case, with a rapidity, a brilliancy of flame, and an evolution of light, truly wonderful.

412. During *combustion in oxygen gas* the volume of the gas is decreased, and, if the combustion continues long enough, the gas wholly disappears. This is owing to a combination which takes place between the oxygen of the oxygen gas, and the body that is burnt in the oxygen gas. The result of this union is either an oxide, or an acid.

413. Sometimes the product of such combustion is a gaseous body, and sometimes a solid. Thus, sulphur produces sulphuric acid gas, and carbon, carbonic acid gas;—but, phosphorus produces phosphoric acid, which is deposited in a *solid* state, and a vacuum is produced in the vessel wherein the combination of the two elements take place.

414. When combustion takes place in *common air*, the same phenomena occur, but less rapidly, and to a less extent. By burning substances in a given portion of common air, the bulk of that portion of air is diminished one-fifth, and the remaining quantity will support neither combustion, nor animal life. The portion of air thus abstracted has been proved to be oxygen, and the air remaining is nitrogen. And by mixing nitrogen gas and oxygen gas in the above-mentioned proportions, a compound is obtained which possesses precisely the same properties as common air. Thus, therefore, the composition, and the proportions of the constituents, of atmospheric air, is proved both by analysis and synthesis.

415. The grand uses of air being to support life and combustion, and its pure part being abstracted thereby, a continual supply becomes necessary wherever those processes are carried on. This shows us how important it is to renew the fresh air of the rooms we live in, in order that breathing, and the burning of fires and candles, may be readily carried on.

416. CHARCOAL BURNS BRILLIANTLY IN OXYGEN GAS, AND PRODUCES AN ACID—THE CARBONIC. *Process 1.*—Fill the bottle, fig. 37, with oxygen gas, as directed 328. Then, put a piece of red-hot charcoal into the spoon, fig. 3, and plunge it into the gas; allowing the instrument to be sustained in its place, by the cork which is laid upon (not fastened into) the neck of the bottle. As soon as the red-hot charcoal comes into contact with the gas, it begins to burn very vividly, its combustion proceeds with great splendour, and brilliant scintillating sparks are thrown out in all directions. When the combustion is at an end, it will be found that the oxygen gas has been converted into carbonic acid gas.—This is an instance of *the formation of an acid*, by the union of a simple body (carbon) with oxygen.—The reason that the cork to which the spoon is attached must not be screwed tightly into the neck of the bottle, is, that the gas, upon being heated, expands, and would burst the bottle, were it closely fastened up.

417. *Process 2.*—The preceding experiment may be performed on a smaller scale, by employing a jar that holds less gas, and using a copper wire, with a bit of charcoal fastened to the end of it. In this case, beautiful sparks will be thrown out, as before.

418. SULPHUR BURNS BEAUTIFULLY IN OXYGEN GAS, AND PRODUCES AN ACID—THE SULPHURIC.—A piece of sulphur, the size of a pea, is to be put into the copper spoon, set fire to by a candle and blow-pipe, and plunged into the same jar, and in the same manner as directed for performing the experiment with charcoal. The sulphur will burn with a beautiful violet-coloured scintillating flame, and the jar will shortly be filled with a brown vapour, which is sulphuric acid gas. This gas, if water has

been put into the jar, will speedily combine with it, and produce sulphuric acid.

419. **SPLendid COMBUSTION OF PHOSPHORUS IN OXYGEN GAS, AND PRODUCTION OF PHOSPHORIC ACID.**—*Process 1.*—The light of phosphorus in combustion in oxygen gas, is the most splendid that can be, by any means, produced. Place the size of a small pea of phosphorus in a little hemispherical tin cap, raised, by means of the wire stand, (fig. 48), an inch or two above the surface of water contained in a broad shallow dish. Fill the receiver *a*, fig. 35, with oxygen gas, and screw on the top of it the bladder *d*, compressed—(see 291); next open the stopcocks *b*, *e*, so as to make a communication between the receiver and the bladder; and then press over the mouth of the receiver, as it stands in the pneumatic trough, a circular piece of pasteboard, rather exceeding its diameter. Now, instantly cover the phosphorus, when an assistant has set fire to it, with the vessel of oxygen gas, retaining the pasteboard in its place till the receiver is immediately over the cup. When this has been skilfully managed, a very small portion only of the gas can escape. The inflammation of the phosphorus will be so extremely brilliant that it will be found almost impossible for the eyes to bear the light. The use of the flaccid bladder is to receive the expanded gas, which is thus prevented from escaping into the room, and proving disagreeable by its suffocating smell.—The odorous compound produced being phosphoric acid, which at last settles on the sides of the receiver in white flakes.

420. *Process 2.*—The foregoing experiment may be more easily, but less agreeably, performed, by fastening a bit (of the size of half a pea) of phosphorus to a wire, putting it in the copper spoon, and then immersing it in a bottle. See 416, 417.—The student is particularly cautioned against using larger pieces of phosphorus than those directed.

421. **IRON MAY BE BURNED IN OXYGEN GAS: THE COMBUSTION IS ATTENDED BY A BRILLIANT LIGHT, AND THE PRODUCT IS A METALLIC OXIDE.**—Prepare a bottle of oxygen gas, as directed 328; and, also, prepare the

fire, fig. 39, as directed 330. Light the inflammable matter at the bottom of the wire, and plunge it into the bottle, suspending the whole by the cork. The flame will be instantly communicated to the wire, which will continue to burn with an appearance inconceivably brilliant and striking: proceeding with a meteor-like body, in a spiral form, and throwing out beautiful sparks in all directions.

422. These sparks, upon being examined when cold, are found to be very different from the iron of which they have been formed.—They are brittle, and destitute of metallic lustre. The weight of the drops, too, is greater than that of the metal made use of; so that, in burning, something must have been added to them: this something, is the oxygen, which, united to caloric, formed the oxygen gas. The term applied to this compound is *oxide of iron*, or iron and oxygen. When the drops fly off in their fire state, they are so hot, that unless the bottom of the jar be covered an inch or so with sand or water, they perhaps crack it.

423. COMBUSTION OF ZINC, AND FORMATION OF OXIDE OF ZINC.—Substitute, for the phosphorus in experiment 419, a small ball formed of turnings of zinc, in which about a grain of phosphorus is enclosed. Set fire to the phosphorus, and cover it expeditiously with the jar of oxygen. The zinc will be inflamed, and burn with a beautiful white light.

424. PROOF THAT METALS ARE INCREASED IN WEIGHT BY COMBINING WITH OXYGEN.—Coil up a drachm of very slender iron wire, (see 330), and put it into the bowl of the pipe, fig. 50, which place in a clear fire. Have ready the bladder *d* fig. 35, filled with oxygen gas. When the iron in the pipe is red hot, force from the bladder, through the pipe, a stream of oxygen gas. The iron will burn very rapidly, and, by combining with the base of the gas, be converted into oxide of iron—see 422. If the bowl of the pipe is kept free from dust, the iron, upon being weighed, will be found to have increased from 1 drachm, to 1 drachm and 20 grains, by its oxygenation.

425. READY AND ELEGANT MODE OF BURNING METALS.—*Process 1.*—Have ready a quantity of oxygen gas in the gas-holder, fig. 61. Fix the blow-pipe *i* to the stop-cock *h* of the instrument; and let the funnel *e* be placed as in the figure. The stand *k* is to be put directly under the opening of the blow-pipe. Take now a small square piece of charcoal, and set it on fire at top by means of a common blow-pipe. Lay this charcoal, when red-hot, upon the stand *k*, open the stop-cock *h*, and pour water down the funnel *e*. As the water descends, the gas will issue from *i* with great violence, and falling directly against the inflamed charcoal will produce a very intense heat. Now, drop iron filings upon this charcoal and they will exhibit a very brilliant light. The process being, in fact, only a variation of experiment 421.

426. *Process 2.* The same as the last, only, in place of iron filings, use tin filings, or tin finely granulated.

427. *Process 3.* The same as the preceding, with the exception of employing copper filings. These burn with a beautiful greenish flame.

* * We shall give no directions, though they might easily be multiplied, for oxidising other metals thus; because the student may try the effects of the gas upon various substances he may have at hand. By means of this apparatus, which is represented complete in plate 6, a series of most beautiful experiments may be performed with great ease.

428. TO PROCURE HYDROGEN GAS, OR INFLAMMABLE AIR.—*Process 1.*—Fix fig. 30 into fig. 29, closing the juncture very accurately, by the application of putty, or any other lute which may be found necessary or convenient—see 281. Into the bottle thus prepared, introduce one ounce of clean iron filings, and the same quantity of sulphuric acid, diluted with five or six times its quantity of water. Close the bottle with its stopple: the gas will be rapidly evolved, and may be collected over water, as directed 317, or in the gas-holder.—See 353.

429. *Process 2.*—Instead of the materials mentioned above, use one ounce of granulated zinc, half an ounce measure, (see 306), of sulphuric acid, and four ounces of water. This mixture yields a very large quantity of gas.

430. In these experiments, the hydrogen gas is furnished at the expense of the water, which latter, is, by the metal, assisted by the acid, decomposed : its hydrogen escapes in the form of gas ; its oxygen combines with the metal, and thus renders it fit (for it is not fit in its pure state) to be dissolved by the acid. The solution which remains behind, is, therefore, a sulphate, either of zinc or iron, according to the metal that is made use of. The student will not, of course, throw this residuum away, as useless ; but filter, evaporate, and crystallize it.

NOTE.—It will be recollected, that the first portion of gas which is collected in this, as well as in all other cases, must be rejected. See 409.

431. MINIATURE BALLOONS.—A MODE OF ILLUSTRATING THE EXTREME LIGHTNESS OF HYDROGEN GAS. *Process.* Fill a bladder with hydrogen gas, in the manner directed 291, 355. Fix to the stop-cock of the bladder the pipe fig. 50. Prepare a strong solution of soap, (a lather such as children use to blow common soap bubbles with), dip the bowl of the pipe into it, and by compressing the bladder, after having opened the stop-cock, fill soap bubbles with the hydrogen gas. These, when shaken from the pipe, instead of falling downwards, like common bubbles, will rapidly ascend to the ceiling of the room. This experiment affords not only a proof of the comparative little specific gravity of hydrogen gas, but also a good illustration of the principles of *Aërostation* : for it is with hydrogen gas that *air-balloons* are generally inflated.

AIR-BALLOONS.—A very pretty apparatus, of recent invention, is now to be had of the philosophical instrument makers. It is a little balloon, in shape resembling a bladder, and is to be had of various sizes. It is made of the maw of a Turkey, and is so extremely light, that, when filled with hydrogen gas, and left free in the atmosphere, it *ascends*.—The price of this apparatus is from 1s. 6d. to 6s., according to the size. One of 4 inches diameter will cost about 2s. 6d.

432. If one of these soap bubbles be arrested in its flight by the application of a lighted paper, the hydrogen gas will explode, and the bubble burst with a vivid flash of light.

NOTE.—Take care and not inflame the bubbles till they are detached from the pipe, or the whole of the gas in the bladder may explode.

433. DETONATING BALLOONS.—Fill a bladder with a mixture of two parts of hydrogen gas, and one part of oxygen gas. Bubbles blown with this mixture will ascend, though not so rapidly as those filled with pure hydrogen. But, upon the application of flame, they will explode with far greater violence; without, however, occasioning any accident, unless they are fired before they are away from the pipe.

434. THE PHILOSOPHICAL TAPER—AN ILLUSTRATION OF THE COMBUSTIBILITY OF HYDROGEN GAS.—Fit the jet-pipe, fig. 31, into one of the necks of the gas bottle fig. 29. Then, introduce through the other neck, materials for producing hydrogen gas (see 429.) In a short time, the gas will be heard issuing from the top of the pipe *a*. Let it escape, till you think as much has issued as served in the beginning to fill the bottle; then, apply to the top of the tube, a lighted paper; upon this the gas will be inflamed, and will burn with a blueish-coloured jet, as long as it continues to be produced.—The reason that a quantity of air must be suffered to escape is explained by *note* 409. Hydrogen gas mixed with common air, violently explodes when inflamed, so that particular care is requisite, in this experiment, to let all the common air escape.

435. *Another mode of performing this experiment, is by fixing the pipe, fig. 31, into the top of the stop-cock *d* of fig. 61. Then pouring water down *e*, and inflaming the gas, which, of course (see 354) will issue from the pipe. Or, the jet-pipe, fig. 49, may be fixed to a bladder filled with gas, (461), and the gas inflamed as before. When by pressing the bladder more or less, a very pretty jet, or stream of fire, either large or small, is easily produced. No danger need be apprehended, because the smallness of the hole in the end of the pipe prevents the flame from entering into the bladder.*

436. ARTIFICIAL FLOWERS are made by passing hydrogen gas through metallic tubes, bent into various forms and having a number of small orifices bored in them as

certain distances. These tubes are affixed to a main pipe which enters the bag or bladder containing the gas; and the gas is forced through them by pressing the bladder. The gas then issues from the small orifices into the atmosphere, and is there inflamed by the application of a lighted candle. If the machine is made partly, or in whole, moveable, a considerable increase is made to the effect.

437. We account for the luminous appearance in the sky, called the AURORA BOREALIS, or *northern lights*, by supposing that a stratum of hydrogen gas, above the atmosphere of common air, is lighted by the electrical fluid, and burns slowly where it comes in contact with common air. The levity of this gas, accounts for its ascending to so high a region; and as it can only burn when in contact with the latter, the appearances we observe in the *aurora borealis* are exactly such as we should expect from such a cause.

438. HYDROGEN GAS EXTINGUISHES FLAME.—To show this, take a phial which holds about two ounces of water, and fill it with hydrogen gas in the usual manner (284). Then, raise it from the shelf of the pneumatic trough, (holding the mouth downwards), and immerse a lighted taper in the gas: the flame will be extinguished.—*Note.* The phial should be made of strong glass, and a handkerchief should be folded round it when used, to prevent injury being received from the glass, in case of bursting.

439. HYDROGEN GAS ONLY BURNS IN THE PRESENCE OF AIR.—Fill a medicine phial with a narrow neck with hydrogen gas, and cork it while under water. Place it then on a table, withdraw the cork gently, and apply a light; the gas will burn with a pretty blue flame, just at the mouth of the phial, sinking at last into the phial, and then disappearing.

440. FORMATION OF WATER BY THE COMBUSTION OF HYDROGEN GAS. The mode of exhibiting this experiment is shown by figure 47. It consists in holding a bell glass over the flame of the *philosophical taper* (434). The hydrogen gas, as it burns, combines with the oxygen of the atmosphere, and forms water, which rises in vapour, and is

deposited on the internal surface of the glass, in a kind of fine dew.

441. MUSICAL SOUNDS PRODUCED BY THE COMBUSTION OF HYDROGEN GAS.—Take a tube, either of glass, earthenware, or metal, from 18 to 24 inches long, and from 1 to 2 inches wide, and open at both ends. Bring it down a few inches over the flame of the *philosophical taper*, (434).—see figure 45,—and very strange but pleasing sounds, somewhat resembling those of an *Æolian harp*, will be immediately produced. By raising or depressing the tube, or by using tubes of different sizes, the intensity of the musical chord may be greatly varied. The production of the sound is occasioned by the rushing of the air into the tube, to supply the vacuum repeatedly formed by the condensation of the oxygen of the air into water, as it combines with the burning hydrogen.

442. TO PROCURE CARBONIC ACID GAS.—Put into the gas-bottle, fig. 29, an ounce or two of chalk or marble, broken into pieces the size of peas, and pour over that 3 or 4 ounces of water. Next, fit the conducting tube, fig. 30, in its place, and fit into the other neck of the bottle, instead of its glass stopple, the hydrostatic funnel, fig. 58. This instrument should, for this purpose, be shorter in the bended part than is represented in the figure, and the bottom end of it should go into the water in the bottle. All things being ready, pour gradually into the funnel, either sulphuric acid, or muriatic acid. A violent effervescence will be produced, as soon as the acid descends into the water: carbonic acid gas is disengaged, and may be collected over water, in the same manner as hydrogen gas, (428). Sometimes the acid and water are mixed before they are put in the bottle, and the funnel is not used; but so violent is the action which takes place, that when this mode is adopted, a quantity of gas escapes before the apparatus can be properly secured.

443. TO SHOW THAT CARBONIC ACID GAS EXTINGUISHES FLAME, AND DESTROYS ANIMAL LIFE.—Fill a glass (such as fig. 46) with carbonic acid gas, as directed 288, and plunge a lighted candle into it: the flame will be extinguished. A person who is quite a stranger to the proper-

es of this kind of gas, will be agreeably amused by extinguishing lighted candles, or blazing chips of wood, on its surface; as the smoke readily mixes with the gas, and little or none of it escapes into the atmosphere. The smoke floats, in a very curious manner, on the surface of the gas, forming a smooth well defined-plain, which, if the vessel be agitated, is thrown into the form of waves. Insects, which it is desirable to preserve in their true form and brilliancy of colours, for cabinets, may be instantly killed by immersion in carbonic acid gas.

444. PLEASING MODE OF SHOWING THE GREAT WEIGHT OF CARBONIC ACID GAS.—Place a lighted candle in the bottom of a jar which has its open part uppermost, (the jar being filled with atmospherical air); take then a jar filled with carbonic acid gas, and invert it over the jar in which the candle is placed: the effect is very striking: the invisible fluid descends like water, and extinguishes the flame. The whole, to spectators who have no idea of substance without sensible matter, having the appearance of *magic*!

445. AN ILLUSTRATION OF THE CHARACTERISTIC POWERS OF OXYGEN GAS, CARBONIC ACID GAS, AND ATMOSPHERICAL AIR, WITH RESPECT TO COMBUSTION.—Set three jars resembling fig. 46 or 52, of equal size, mouths upwards, on a table. The first must contain common air; the second, carbonic acid gas; and the third, oxygen gas. Take a lighted candle with a pretty large wick, and lower it, by means of a wire, see fig. 46, into the *first* jar—the flame will have its usual brightness. Lower it next into the *second* jar—the flame will be extinguished. Lower it now, while the wick continues red, into the *third* jar—it will be relighted, and will burn for some time with a dazzling splendour. To ordinary spectators, this experiment will be the subject of much wonder. The whole of the jars will, by them, be deemed *empty*; and the different effects resulting from plunging the same candle into seemingly similar vessels, will be quite incomprehensible.

446. TO PROCURE LIQUID CARBONIC ACID.—Having filled a quart bottle with carbonic acid gas, pour into it half a pint of pure water, then cork it securely, and shake

it violently for a few minutes. Let it now remain for a quarter of an hour, (cork downwards), and then shake again. Continue to do this for three hours. At the expiration of which time, the water, if very cold when put into the gas, (for the colder it is the more gas it absorbs) will be very strongly impregnated.

447. TO FORM CARBONATE OF POTASS.—*Process 1.*—Fill a common phial with carbonic acid gas. Apply the thumb to its mouth, and invert it in a solution of pure potass, contained in a cup, and rather exceeding in quantity what is sufficient to fill the phial. The liquid will instantly rise in the phial, and the gas, if it be pure, will disappear entirely; being all absorbed. Pour out the alkaline liquor into the cup, fill the phial again with gas and again invert it in the solution. The absorption will take place again. Repeat the process till the alkali is saturated.

448. *Process 2.*—Take such a bottle as *c*, fig. 57. Put into it a solution of pure potass, and then connect it, by means of one of the tubes, *e*, fig. 57, with the gas-bottle fig. 29. Now, put into the latter, materials for producing carbonic acid gas, as directed 442. The gas, as it is formed, passes of course into the solution of potass, in which it is absorbed. The neck *f* of the bottle *c* is to be closed. Then, when the solution of potass is saturated with gas, it will be forced up the tube *d*, by the superfluous gas which presses its surface in the bottle.

449. TO FORM CARBONATE OF LIME.—Mix lime water with water impregnated with carbonic acid gas, (446). These liquids are individually transparent, but the mixture is opaque: a white powder gradually sinks to the bottom of the vessel.—*Rationale.* The affinity of carbonic acid for lime is very strong: here therefore, the two bodies quitting the water which held them in solution, unite and form an insoluble salt, carbonate of lime (chalk).

450. ILLUSTRATION OF THE MODE OF COLLECTING GASES THAT ARE ABSORBABLE BY WATER, WITHOUT THE AID OF A MERCURIAL TROUGH.—Prepare the apparatus when the gas is to be evolved, as described 442. But let the part *b* of the tube fig. 30, be sufficiently long to reach

the bottom of a long cylindrical jar, just as the tube *a*, fig. 52, reaches to the bottom of that jar. Then, the carbonic acid gas, as it is evolved, forces the common air, which is lighter than itself, out of the jar, and occupies its place. We discover when the jar is filled with the gas, by holding to the top of it a piece of wetted litmus paper, the blue colour of which is changed by the gas to a red.

It will be observed, perhaps, that carbonic acid gas, is *not* incapable of being collected over water. That is true: but we only give directions for obtaining *this* gas without the trough, by way of example. It will be seen, that this mode can only be efficient, when the gas to be collected is heavier than common air: another plan is to be adopted, when the gas is lighter than common air.—See 468.

451. TO PROCURE NITROGEN GAS. *Process 1.*—Introduce a lighted taper under a glass jar, which stands over water, and is filled with common air. The light will shortly be extinguished, a cloudiness will be perceived, which, however, soon subsides, and the water in the basin rises in the jar.—*Rationale.* The atmospheric air is decomposed; the oxygen is absorbed by the burning taper, and the nitrogen remains. The cloudiness proceeds from the unconsumed smoke of the taper. The water rises in the jar, because the included volume of air, is, by the absorption of its oxygen, diminished.

452. *Process 2.*—Mix equal weights of clean iron filings and sulphur into a paste with water, and place the mixture over water, in a cup supported by the stand, fig. 48: then invert over it, a jar full of common air, and let it remain thus for a day or two, when the air will be diminished in bulk one-fifth, and nitrogen gas will remain. The mixture having absorbed the oxygen of the air.

453. *Process 3.*—Wash a piece of lean beef well, and cut it into very small pieces; put these into a retort, and pour upon them nitric acid, diluted with a considerable quantity of water. Apply a gentle heat. Collect the gas over water.

454. TO PROVE THAT NITROGEN GAS DOES NOT SUPPORT COMBUSTION.—The same as process 438. Only that the

gas employed must be nitrogen. The light of the candle will be instantly put out.

455. TO COMPOSE ATMOSPHERICAL AIR.—Mix four parts of nitrogen gas, with one part of oxygen gas. Examine this mixture—by plunging a lighted taper in it, or by any other test you please—it will be found to possess all the properties of common air.

456. TO PROCURE NITROUS GAS—Put some shreds of copper into the retort, fig. 18, and add a portion of nitric acid, diluted with five times its weight of water. Apply a gentle heat, and collect the gas, which will be rapidly evolved, as directed 408.

457. TO CONVERT NITROUS GAS INTO NITROUS ACID GAS.—*Process 1.*—Open a bottle of nitrous gas (which is colourless) in the atmosphere: *red fumes* appear.

458. *Process 2.* Fill the eudiometer, fig. 34, with water and place it on the shelf of the pneumatic trough; pass up into it, three parts of nitrous gas, and two parts of oxygen gas. The gaseous mixture will become of red colour and will then disappear.

* * * In both these cases, nitrous acid gas is produced by the combination of oxygen gas and nitrous gas: in *process 1*, the oxygen is taken from the air of the atmosphere—in the latter case, the acid is absorbed by the water.

459. WHEN AN INFLAMED TAPER IS PLUNGED INTO NITROUS GAS, THE LIGHT IS INSTANTLY EXTINGUISHED. But inflamed phosphorus burns in it with great brilliancy.—*Rationale.* The burning phosphorus decomposes the nitrous gas, and absorbs its oxygen; but the heat of the inflamed taper is not sufficiently intense to do that.

460. TO PROCURE NITROUS OXIDE, OR INTOXICATING GAS.—Put a quantity of nitrate of ammonia into a glass retort, and apply the heat of a lamp, which must be gentle and well regulated. The salt will in a short time liquify and must then be kept gently simmering, avoiding violent ebullition. The gas may be collected over water, and must be allowed to stand a few hours before it is used, during which time it will deposit a kind of white vapour and will become perfectly transparent.

461. SOME OF THE PROPERTIES OF NITROUS OXIDE GAS.

-1. A candle burns in it with a brilliant greenish flame, and a crackling noise.—2. It is heavier than common air.—3. Phosphorus, charcoal, and sulphur burn in it,—as likewise does iron wire.—4. When mixed with oxygen gas, upon the application of flame, it detonates.—All these things may be proved, by proceeding as directed for performing the experiments which demonstrate the properties of oxygen gas. It must be observed, however, that bodies to be burned in nitrous oxide gas, must, when introduced into it, be in a state of complete ignition.

462. INTOXICATING POWER OF NITROUS OXIDE GAS.—

Though this gas is not fitted to support life, yet it may be respired for a short time, and the effects produced by it upon the animal frame, are its most extraordinary properties. The manner of breathing it is this: put nitrous oxide gas that has been purified by standing over water into a large bladder, or varnished silk bag, having a wide glass tube, or a stop-cock with a large bore, affixed to its neck. Then, hold the bladder by the tube (closing the mouth of the tube by applying the thumb) in the right hand; close the nostrils with the left hand; expel the air contained in the lungs by a long expiration; and instantly apply the tube of the bladder to the mouth, and breath the gas from, and into, the bladder as long as possible, which, perhaps, will be about two or three minutes. What effects will be produced? Why, it is impossible to say: for they differ greatly according to the constitutions of the persons by whom the gas is respired. In general, however, they are *highly pleasureable*, and resemble those attendant on the agreeable period of intoxication. “Exquisite sensation” of pleasure—an irresistible propensity to laughter—a rapid flow of vivid ideas—singular thrilling in the toes, fingers, and ears—a strong incitement to muscular motions—are the ordinary feelings produced by it. The celebrated Mr. Wedgwood, “after breathing the gas sometime, drew the bag from him, and kept breathing on laboriously with an open mouth, holding his nose with his fingers, without power to remove them, though aware of the ludicrousness of his situation; he had a violent inclination to jump

over the chairs and tables, and seemed so light, that I thought he was going to fly." What is exceedingly remarkable, is, that the intoxication thus produced, instead of being succeeded by the debility subsequent to intoxication by fermented liquors, does, on the contrary, generally render the person who takes it cheerful and high spirited for the remainder of the day.

463. CARBURETTED HYDROGEN GAS—THAT WITH WHICH SHOPS ARE LIGHTED—PRODUCED ON A SMALL SCALE.—Fill the bowl with the largest tobacco-pipe, that can be procured with pulverised coal of a good quality, and close the top of it by the application of pipe-clay, or, what is better, a mixture of sand and beer. When the lute is dry, place the bowl of the pipe in a clear fire: in a few minutes dense smoke will issue from the stem, which, on the application of a lighted paper, will inflame, and will continue in a state of beautiful combustion as long as any gas continues to be distilled from the coal. The body which will be found remaining in the bowl of the pipe is the substance called *coke*.

464. A FOUNTAIN OF FIRE—FORMED BY PHOSPHURETTED HYDROGEN GAS.—Put fifteen grains of finely granulated zinc, and six grains of phosphorus, cut in small pieces, (cut this under water,) in the glass, fig. 17. Mix in another glass, a drachm by measure of sulphuric acid with two drachms of water. Now, take the two glasses to a dark room, and there pour the diluted acid over the zinc and phosphorus in the other glass: in a short time, phosphuretted hydrogen gas will be produced, and beautiful jets of blueish flame will dart from all parts of the surface of the liquid, the mixture will be quite luminous, and a quantity of beautiful luminous smoke will rise in a column from the glass. A *Fountain of Fire* is a very apt name for the appearance that is produced. The experiment is very easily performed, and is a very beautiful one.

465. TO PROCURE SULPHURETTED HYDROGEN GAS.—Put into the retort, (fig. 18), one part of sulphuret of iron of commerce, or of sulphuret of iron, broken into a coarse powder, and pour upon it three or four parts

strong muriatic acid; apply a gentle heat, and receive the gas over water.

466. LIQUID SULPHURETTED HYDROGEN.—Water may be impregnated with sulphuretted hydrogen gas, in the same manner that it is impregnated with carbonic acid gas.

—See 446. This compound is chiefly used as a test.

467. TO PROCURE AMMONIACAL GAS.—*Process 1.*—Take equal parts of muriate of ammonia, and quicklime, each separately powdered, and introduce them into the flask, fig. 54, then fit to the flask the pipe, fig. 31. Apply a gentle heat, and the gas will be rapidly evolved. The rationale of the production of ammoniacal gas in this case, has been already described, (394). If this gas is conveyed into water, it is rapidly absorbed: the water acquiring the properties of what is known by the name of *liquid ammonia*. To collect it, we therefore, do not use the water trough, but proceed as follows: over the pipe whence the gas issues, we invert a glass jar, bringing it down till the top of the pipe nearly touches the upper part of the jar,—this is represented by fig. 51. This mode is founded on the difference between the specific gravity of this gas, and that of common air: for here, the gas entering the jar at the top, forces down the heavier common air, and occupies its place. We discover when the jar is full of gas, by holding at the bottom of it a feather moistened with muriatic acid; for when the fume of muriatic acid comes in contact with ammoniacal gas, a white vapour is produced. Now, muriatic acid gas (of which we shall speak presently) must be collected in the manner shown by fig. 2.—See 470. And, consequently, in order to discover when the jar is full of muriatic acid gas, we have only to hold at the mouth of it, a feather moistened with liquid ammonia. *Process 2.* The same as the preceding—only, instead of the powdered materials for producing the gas, use liquid ammonia.

468. SULPHUROUS ACID is formed when a common flintstone match is burnt. Its peculiar smell may be then perceived. Its effects on vegetable colours may be shown in a pleasing manner, by holding a red rose over the blue flame produced by the burning sulphur: wherever the

sulphurous acid vapour comes in contact with the red, the colour will be discharged so as to render it beautifully variegated, or entirely white. Afterwards, the rose, dipped in water, recovers its redness.

469. TO PROCURE MURIATIC ACID GAS.—Put a quantity of dried muriate of soda (common salt) into the bottle, fig. 29, then fit the tube, fig. 30, and the funnel, fig. 58, into the bottle, as directed at 591. Pour strong sulphuric acid down the funnel, and muriatic acid will be rapidly evolved. For the mode of discovery when the jar is full of gas, see the preceding experiment *Rationale of this experiment*. The muriate of soda is decomposed by the sulphuric acid: the muriatic acid is discharged in the state of gas, and sulphate of soda (which may be easily crystallized) remains behind.

470. TWO PUNGENT AND INVISIBLE GASES UNITE AND FORM AN INODOROUS SOLID.—*Process 1.*—Fill a jar with ammoniacal gas, as shown by fig. 51, (see 468), and fill another with muriatic gas, as shown by fig. 52.—See 470. Apply the two jars mouth to mouth, and the above-mentioned effect will instantly be produced: muriate of ammonia being formed and precipitated on the sides of the jars. *Process 2.*—Dip a clean feather into muriatic acid, and moisten with it the interior of the glass, fig. 6. Moist in like manner, the interior of the glass, fig. 16, with liquid ammonia. The glasses, in this state, still seem empty; but, if they are put mouth to mouth together, the whole included space will be filled with a dense white vapour; which in the end settles on the sides of the glasses in the form of a white powder, (solid muriate of ammonia).—*Process 3.* If two jars, one containing ammoniacal gas and the other containing carbonic acid gas, are thus put together, solid carbonate of ammonia will be formed in like manner.

471. TO PROCURE CHLORINE GAS—Grind, in a mortar, three parts of common salt, with one part of black oxide of manganese. Introduce this mixture into the retort (fig. 18,) and add two parts of sulphuric acid, diluted with the same weight of water. Instead of the above-mentioned materials, you may use one ounce of fine

powdered black oxide of manganese, and two ounces of muriatic acid. In either case, apply a gentle heat, and chlorine gas will be evolved. Chlorine gas, being absorbable by *cold* water, must be received in bottles filled with, and inverted in, *warm* water, (of a temperature made by mixing two parts of cold, with one of the boiling water). The bottles must be provided with accurately ground stoppers, which must be introduced under water, while the bottles remain full of gas, and inverted, and no water must be left in the bottle with the gas.

NOTE:—In experiments with this gas, great care should be taken that it does not escape, in any considerable quantity, into the apartment; as its action on the lungs is extremely injurious.

472. LIQUID CHLORINE.—This may be prepared by agitating chlorine gas with water, in the manner directed for such operations at 446. It possesses the property of destroying vegetable colours.

473. ILLUSTRATION OF THE ART OF BLEACHING.—Put into a phial of liquid chlorine, strips of linen cloth, dyed of different colours—the colours will be quickly discharged.

474. TO SHOW THAT CHLORINE GAS SUPPORTS COMBUSTION.—1. A candle, previously lighted, when plunged into a jar of it, burns with a red flame, and a copious emission of dense fumes. 2. Charcoal dust; 3. tinfoil; 4. copperleaf; 5. powdered antimony; 6. phosphorus; and some other bodies; when dropt in it, *inflammes spontaneously*.

475. TO PROCURE CHLORATE OF POTASS.—Pass chlorine gas, as it comes from the vessel in which it is formed, through a nearly-saturated solution of pure, or sub-carbonate of, potass. This may be done by means of Woulfe's apparatus, (fig. 57), using only the bottle into which the alkaline solution is put in addition to the receiver. But, for this purpose, the tube *e*, which passes into the solution, must be half an inch wide, in order that it may not be choaked up by crystals which form during the process. The chlorine gas as it passes into the liquid alkali will be rapidly absorbed, and, if the carbonate of potass is used,

the carbonic acid escapes with effervescence. The liquor is generally of a pinkish hue from the presence of manganese. When it is saturated, it may be put aside, in a cool dark place, for 24 hours, when it will be found to have deposited a considerable portion of crystallized chlorate, which may be taken out, drained, and purified by solution in boiling water, which, during cooling, deposits the salt in brilliant crystalline scales. Dr. Henry, in his directions for preparing this salt, says—"the solution when saturated with the gas, may be gently evaporated and the first products only of crystals are to be reserved for use; for the subsequent products consist of muriate of potass only." NOTE:—By passing the current of gas into the solution the water is decomposed: its oxygen unites to one portion of chlorine to form chloric acid; its hydrogen unites to another portion of chlorine to form muriatic acid: hence a chlorate and a muriate of the alkali are simultaneously produced, and are to be separated by one of the methods above described.

SECTION IV.—ON THE PREPARATION OF SALTS

476. CARBONATE OF POTASS may be formed by passing a current of carbonic acid gas through pure (or sub-carbonate of) potass, as directed 448. The salt may be crystallized by very slow evaporation.

477. SUB-CARBONATE OF POTASS may be obtained by dissolving the potash of the shops, and filtering and evaporating the solution. It does not crystallize.

478. NITRATE OF POTASS is obtained by adding to a solution of sub-carbonate of potass, nitric acid, as long as effervescence continues. The solution is then filtered and crystallized. To obtain cheaply, for purposes of experiment, nitrate of potass in a state of purity, dissolve common saltpetre in boiling water, nearly to the point of saturation; strain the solution while hot; and add a few drops of a solution of caustic potass. If precipitation ensues, continue to add the potass till it ceases. If precipitation takes place, or when it is at an end, filter the solution, and crystallize by slow evaporation.

479. TARTRATE OF POTASS may be obtained by adding sub-carbonate of potass to cream of tartar, or crystallizing

tartaric acid, in solution, till the cessation of effervescence, 100 parts of cream of tartar, (which is a super-tartrate) require $43\frac{1}{2}$ parts of sub-carbonate of potass to make a neutral tartrate.

480. TO OBTAIN PURE SUB-CARBONATE OF SODA—dissolve soda of commerce in a small portion of water, filter the solution; this clears it from the admixture of earthy bodies. Slowly evaporate the filtered solution by a low heat, very small crystals of muriate of soda will form on the surface. Skim them off. When these cease to form, the solution may be suffered to cool, and the purified sub-carbonate of soda will crystallize.

481. MURIATE OF SODA—(CULINARY SALT)—may be obtained by dropping into a saturated solution of sub-carbonate of soda, muriatic acid, as long as effervescence takes place. The solution is then to be filtered, and the salt crystallized by evaporation.

482. TO PURIFY MURIATE OF SODA for chemical purposes, dissolve it in boiled water; filter the solution; and add a solution of carbonate of soda as long as any milkiness ensues. Filter the solution, and evaporate till crystallization.

483. SULPHATE OF SODA is formed by adding sulphuric acid to a solution of sub-carbonate of soda, till effervescence ceases. The solution must be filtered and evaporated for a short time, and then set by to cool, when crystals will be deposited. The remaining solution must be again evaporated, and again cooled; and this must be repeated till all the water is driven off. *Note.*—If you put a ready formed crystal of the salt into the solution which you set by to cool, crystallization will take place very speedily.

484. NITRATE OF AMMONIA,—To form this; saturate nitric acid diluted with five parts of water, with carbonate of ammonia. Filter the solution. Then, evaporate by a gentle heat, and, to supply the waste of the alkali, (which is volatilized by the heat,) add, occasionally, a little more of the carbonate. When the evaporation has proceeded to a certain extent (which experience alone can point out), the solution is set aside, and the salt, upon cooling, crys-

tallizes. The chief use of this salt is to afford, by its decomposition, nitrous oxide.

486. MURIATE OF AMMONIA may be formed by any of the following processes.—1. By saturating sub-carbonate of ammonia with muriatic acid, and filtering the product.—2. By combining the constituents in the gaseous state, see 471.—3. By putting into a wine glass a small quantity of liquid ammonia, and pouring upon it liquid muriatic acid till saturation, which will have taken place when the addition of the acid ceases to cause a white vapour to ascend.

487. MURIATE OF BARYTES is formed by dissolving native carbonate of barytes, in diluted muriatic acid. The solution, when saturated, must be filtered, and gently evaporated, when the salt will crystallize. This salt may also be obtained as directed 489.

488. NITRATE OF BARYTES is formed in the same manner as *muriate* of barytes.

489. SULPHATE OF BARYTES is formed whenever the acid and base which constitute it, or bodies containing that acid and base, are placed in contact. The salt is very abundant in a native state. To convert it into the *muriate* proceed thus: reduce it to powder, mix it with charcoal powder, and keep it for some hours red-hot in a crucible: during this operation, the oxygen of the sulphuric acid continues with the carbon and heat, and forms carbonic acid gas, which flies off, and leaves sulphuret of barytes behind. This product must be dissolved in water, and have nitric acid added to it: when the sulphur will be precipitated, and nitrate of barytes will remain in solution. Filtration, evaporation, and crystallization then follow, as above directed.

490. STRONTIA SALTS are prepared precisely in the same manner as the *salts of barytes*.

401. NITRATE OF LIME.—To form this, nitric acid, diluted with five or six parts of water, must be saturated with carbonate of lime, 63 parts of which are decomposed by 90 parts of nitric acid, and give 103 parts of dry nitrate of lime. When this solution is boiled down to the

consistence of a syrup, and exposed in a cool place, the salt crystallizes.

491. MURIATE OF LIME.—This may be obtained by dissolving carbonate of lime in muriatic acid. To be crystallized, it must be boiled down to the consistence of a syrup, and then set in a cool place. *Note* :—This salt, and the preceding one, can only be preserved in the solid state by being kept in closed bottles—they are so extremely deliquescent.

492. SULPHATE OF MAGNESIA.—To form this, saturate the common magnesia of the shops with diluted sulphuric acid, filter the solution, and crystallize the salt, by concentrating the solution, and then allowing it to cool. For the purposes of experiment, dissolve, filter, and recrystallize, the sulphate of magnesia (Epsom salts) of the shops.

493. TO MAKE NITRATE OF SILVER.—Put a small quantity of pure silver into a test glass, (fig. 17,) and pour over it twice its weight of nitric acid and twice as much water as acid. The silver will be quickly dissolved, and at the same time nitrous gas will be discharged. The solution, if the metal and acid be both pure, will be transparent and colourless; but, if any copper be present, the solution will have a green colour; and if any gold be present, it will be precipitated in the form of a black powder. Should the nitric acid contain any muriatic or sulphuric acid, an insoluble white powder will be formed, which will be a muriate, or sulphate, of silver. Nitrate of silver may be crystallized.

494. NITRATE OF MERCURY is made by dissolving mercury in diluted nitric acid. The saturated solution is very ponderous and colourless; and yields, by evaporation, large transparent crystals.

495. TO MAKE NITRATE OF COPPER.—Put a quarter of an ounce of shreds of copper into a test glass, (fig. 17), and pour over them half an ounce of nitric acid diluted with its bulk of water. An effervescence instantly commences, which is accompanied by a copious discharge of nitrous gas. The liquor assumes a muddy greenish ap-

pearance ; but afterwards becomes clear and of a beautiful blue colour, which is permanent ; and a yellowish precipitate is found at the bottom of the vessel. To crystallize this salt, evaporate the saturated solution till a strong pellicle is formed on its surface, then set it by to cool. The crystals have the fine blue colour of the solution.

496. TO MAKE SULPHATE OF IRON, all that is necessary is to pour diluted sulphuric acid over iron filings. See 428, 430. The solution, by evaporation, yields crystals which have a beautiful green colour.

497. MURIATE OF TIN is formed by digesting pure tin in concentrated muriatic acid, in the proportion of one part of the former to two of the latter. The vessel, fig. 53 may be made use of. The solution, when evaporated, furnishes crystals.

498. ACETATE OF LEAD is obtained by dissolving *white-lead* in distilled vinegar. When the solution is evaporated and cooled, the salt crystallizes.

499. NITRATE OF LEAD is formed by digesting lead with nitric acid. It forms crystals, when evaporated and cooled.

500. SULPHATE OF ZINC is formed by pouring sulphuric acid diluted with six parts of water upon granulated zinc. The water is decomposed by the metal : its oxygen forms an oxide by combining with the zinc, and this is dissolved by the acid, forming a colourless solution. The hydrogen of the water, escapes meanwhile, in the gaseous state. See 429. If the solution is rapidly evaporated to dryness, the salt is obtained in a mass which bears a resemblance to loaf-sugar. But, if slowly evaporated to the due degree of density, it shoots, on being allowed to cool, into regular crystals.

501. NITRATE OF BISMUTH.—Nitric acid dissolves bismuth with great rapidity. To one part and a half of nitric acid (undiluted), add, at distant intervals, one of bismuth, broken into small pieces. The salt is crystallizable. The solution cannot be diluted, for water precipitates the metal in the state of an oxide.

502. NITRO-MURIATE OF COBALT.—Digest, in a sand-bath, for some hours, one part of cobalt or zaffre, with four parts of nitric acid, add to the solution, one part of

muriate of soda, and four times as much water as acid, and filter the solution. This is the green sympathetic ink.

SECTION V.—ON TESTS, OR RE-AGENTS.

503. CHEMICAL ANALYSIS consists of a great variety of operations, performed for the purpose of separating the component parts of bodies. It might seem an impossible task, to exhibit, in a separate state, five or six substances, the weight of some of which is, perhaps, less than the one-five-hundredth part of the whole; yet, to so wondrous a degree of perfection has the science of chemistry arrived, that we are enabled to do this with ease.

When, in an analytical pursuit, the object of inquiry is—what are the elementary parts of a certain compound?—we place that compound, under particular circumstances, to the action of certain bodies or powers, which it is expected will chemically act upon it; and which, when they do, produce changes so obvious to the senses, as to enable us to decide whether the compound does, or does not, contain the principles which it was suspected to contain. The bodies which produce these changes are called *tests*, or *re-agents*; the proper application of which constitutes the chief part of the proceeding called chemical analysis.

To illustrate the agencies of chemical re-agents, we shall subjoin some striking and beautiful experiments; but, first, it will be necessary to describe the most important tests, individually.

504. TINCTURE OF LITMUS—*Preparation.* Tie bruised litmus root in a linen rag and steep it in pure water. Strain the infusion, and, to keep it from spoiling, add to it an eighth part of alcohol. Should the tincture be more of a purple colour than of a blue, it must have a drop or two of liquid ammonia added to it.—*Use.* It is a test of most uncombined acids. Its blue colour is changed to red, by water containing 1-20,000 of its bulk of sulphuric acid. The redness produced by carbonic acid and sulphuretted hydrogen, goes away when the mixture is heated. The other acids permanently redden the test: the red colour is changed to blue by the alkalies and alkaline earths.

505. LITMUS TEST PAPERS are prepared by staining slips of paper, four inches long, and half an inch wide with the tincture. The tint should be pale: the paper for this purpose must be *unsized writing paper*; or, if sized, must be well washed with warm water. The litmus papers are more conveniently applied than the tincture: but they are not so delicate a test. They are sensible however, to acid diluted to 1-14,000.—Test papers must be kept from light and air, or they will be spoiled.

506. TINCTURE OF CABBAGE.—To prepare this, pour boiling water upon sliced red cabbage, decant the clear infusion (which will be of a fine blue colour) when cold and mix it with an eight part of alcohol. Red cabbage leaves may be preserved for a long time, by drying them in a warm place till they are crisp, having previously cut them into small pieces. It is necessary to be provided with some of these dried leaves, because the tincture spoils by keeping. The blue colour of this test is changed to red by acids, and to green by alkalies and alkaline earths.

507. TINCTURE OF TURMERIC is prepared in the same manner as tincture of litmus. It has a fine yellow colour which alkalies change to a reddish-brown. The carbonated earths have no action upon it, so that, in solution, they may be thus distinguished from alkalies. It is a delicate test, being affected by a solution containing only 1-2000 of potass. *Turmeric test-papers* are prepared as litmus test-papers are.

508. TINCTURE OF GALLS is prepared by steeping bruised nut-galls in boiling water, filtering the solution, and adding to it an eight part of alcohol. It is the test generally employed for detecting iron, with all the combinations of which it produces a black tinge. It is, likewise, owing to the *tan* contained in it, a good test for gelatin, with which it forms an insoluble precipitate.

509. SULPHURIC ACID.—1. Discovers, by a slight effervescence, the presence of carbonic acid.—2. Barytes, for which it is the best test, is precipitated instantly, in the form of a white powder, which is a sulphate of barytes.—3. It throws down lime from most of its solutions.—4. It precipitates lead, as a white powder.

510. OXALIC ACID (in solution) is a most delicate test of lime, which it separates from all its combinations. *Oxalate of Ammonia* (which may be easily formed by saturating the carbonate of ammonia with a solution of oxalic acid) is preferable to the pure acid as a re-agent, being less affected by other substances.

511. PURE ALKALIES, AND CARBONATE OF ALKALIES.—(in solution), precipitate most of the earths and all the metals: the colour, and other properties, of the precipitate, determining the nature of it.

512. LIME WATER, made by dissolving pure lime in rain water, is chiefly used as a test for carbonic acid, with which it forms a white precipitate. On the same principle *liquid carbonic acid* is a test for lime. Lime water also shows the presence of corrosive sublimate by a sediment of a brick-dust colour.

513. BARYTIC WATER is more effectual than lime water in denoting carbonic acid, and is more portable and convenient; since, from the pure earth, the solution may be at any time readily prepared. The barytic solution is also a most sensible test for sulphuric acid, which it indicates by an insoluble precipitate.

514. METALS may be used as tests of each other on the principle of elective affinity. Thus, for example, a polished iron plate, immersed in a solution of sulphate of copper, soon acquires a coat of this metal—this is occasioned by the sulphuric acid seizing on the iron, and letting fall the copper; and the same in other instances.

515. NITRATE OF SILVER (in solution) is peculiarly adapted to the discovery of muriatic acid and muriates. For the silver combines with the muriatic acid, and forms a flaky precipitate, which, at first, is white, but on exposure to the sun's light, acquires a blueish and finally a black colour. Muriatic acid, consequently, is a test for silver.

516. ACETATE OF LEAD (in solution) is a test for sulphuretted hydrogen, which occasions a precipitate of a black colour. It also forms a white precipitate with sulphuric acid, for which, however, it is not so good a test as the following.

517. MURIATE OF BARYTES (in solution) is the best test for sulphuric acid. A sensible precipitation is produced by it in water, containing only 1.910256 part of sulphuric acid. This is supposing the acid to be uncombined: if the acid in a state of combination (as in sulphate of soda &c.) the test is not so sensible.

518. PRUSSIAN OF POTASS is a very sensible test for iron, with the solutions of which in acids it produces Prussian blue precipitate. To render its effect certain, a little muriatic acid must be added previously, to saturate bodies which might interfere. It also precipitates many other metals, see 167.

519. MURIATE OF LIME (in solution) is sometimes used to discover sulphuric acid, with which it forms a white precipitate; but, it is not so good a test as muriate of barytes.

520. SOLUTION OF SOAP IN ALCOHOL is employed to ascertain the comparative hardness of waters. With distilled water it may be mixed, without any change ensuing; but, if added to a hard water, it produces a milkiness more considerable as the water is less pure; and from the degree of this milkiness, an experienced eye will derive a tolerable indication of the quality of the water. This effect is owing to the alkali quitting the oil whenever there is present in a water any substance for which the alkali has a stronger affinity than it has for oil. Thus all uncombined acids, and all salts, except those of alkalis, decompose soap, and occasion that property in water which is termed hardness.

521. CARBONIC ACID (applied either in the gaseous or liquid state) occasions a white precipitate in lime,—barytes—or strontia water, soluble, with effervescence, in muriatic acid.

522. NITRATE OF MERCURY (in solution) is the most sensible test of ammonia, one part of which, with 30, or 40 parts of water, is indicated by a slight blackish yellow tinge on adding the test.

523. LIQUID AMMONIA is an excellent test for copper, with which it strikes a fine blue colour.

524. METALLIC TIN, AND NITRO MURIATE OF TIN

(in solution), throw down *Gold* in form of a purple oxide. On the same principle, nitro-muriate of gold serves as a test for tin.

525. LIQUID SULPHURETTED HYDROGEN is a test for both lead and arsenic; with the former it gives a black, and with the latter a yellow, precipitate.

* * * *The following experiments are intended to give an idea of the great power and usefulness of these tests.*

526. A BEAUTIFUL GREEN LIQUID PRODUCED BY MIXING A BLUE ONE WITH A COLOURLESS ONE.—Nearly fill the test glass, fig. 6, with water, and add to it a spoonful of the tincture of cabbage. The mixture will be of a fine blue colour: add, next, a few drops of colourless liquid ammonia: the mixture will instantly become green. *Rationale*.—This experiment is designed to exhibit one of the distinguishing properties of alkalies, namely, that of changing vegetable colours. A solution of potash or soda produces the same effect as liquid ammonia; but, if the liquid made green by ammonia be boiled, it regains its blue colour—because that alkali, being volatile, is driven off in the gaseous state by the heat.—The following experiment shows the mode of using the common test for alkalies.

527. *Process*.—Into a test glass of water, pour a few drops of an alkaline solution, and then dip into the mixture a slip of turmeric paper. The pale yellow colour of the test-paper will be changed to a deep brown.

528. TO PRODUCE A BEAUTIFUL CRIMSON LIQUID BY MIXING A BLUE ONE AND A COLOURLESS ONE.—To a mixture of a tea-spoonful of tincture of cabbage with a glass full of water, add a few drops of sulphuric or muriatic acid; upon which, the blue colour given by the tincture will be changed to a red.—This is designed to show the influence of acids in changing vegetable colours. If a red colour is produced by dropping a little tincture of cabbage into a certain solution, we know that that solution contains an acid; and if the reddened solution upon being heated becomes blue, we learn that the acid is either carbonic acid, or sulphuretted hydrogen—as only those two acids are volatile.

529. TO CHANGE THE COLOUR OF A LIQUID FROM GREEN TO RED, BY ADDING A COLOURLESS ONE TO IT.—The green liquid is the one produced by experiment 527.—The colour is changed by pouring a little sulphuric acid into it. A portion of the acid neutralizes the alkali by combining with it, and another portion produces its ordinary reddening effect.

530. TO MAKE THE SAME LIQUID ALTERNATELY RED AND GREEN BY THE ADDITION OF TWO COLOURLESS LIQUIDS, you have only to add alternately a few drops of sulphuric acid and liquid ammonia, to diluted tincture of cabbage.

531. WATER, CONTAINED IN A TALL GLASS, MAY BE COLOURED CRIMSON AT THE BOTTOM, PURPLE IN THE MIDDLE AND GREEN AT THE TOP.—Nearly fill a tall test glass with water, and colour it blue by adding a table spoonful of tincture of cabbage—then make it green by adding a little liquid ammonia, and afterwards pour a little sulphuric acid gently down the sides of the glass: upon which the effect mentioned will be produced.

532. THREE DIFFERENT COLOURS PRODUCED FROM THE SAME VEGETABLE INFUSION, BY THE ADDITION OF THREE COLOURLESS LIQUIDS.—Into each of three test glasses put a little diluted tincture of cabbage. To one, add a solution of alum, to the second, a little solution of potassa and to the third, a few drops of muriatic acid. The product of the first mixture will have a purple colour, that of the second, a bright green, and that of the third, a beautiful crimson.

533. TEST FOR MURIATIC ACID.—Add a drop of muriatic acid to a quart of water; pour some of the mixture into a test glass, and let fall into it a single drop of nitrate of silver—the whole will instantly be pervaded by a milkiness, as the muriatic acid combines with the silver and forms muriate of silver, a salt highly insoluble. So great is the power of this test, that if a single grain of common salt is dissolved in 42,250 grains of water, the muriatic acid, though amounting to only the 1-108,333 part of the weight of the solution is detected.

534. TESTS FOR SULPHURIC ACID.—Mix a drop of sulphuric acid with a quart of water, and pour some of this into four test-glasses. Then add to the four glasses se-

parately, the four tests described 513, 516, 517, 519. In all the cases, there will follow a precipitation of a white powder: in the first case, sulphate of barytes (a salt highly insoluble) is formed; in the second case, sulphate of lead; in the third, sulphate of barytes; in the fourth, sulphate of lime.—The muriate of barytes is the best test.

535. TESTS FOR IRON.—*Process 1.* In a goblet-full of water, dissolve one grain of sulphate of iron, and add a very little tincture of galls: a purple or blackish colour will be produced immediately, showing that every drop of the water is united to a portion of iron.—*Process 2.* To a similar diluted solution of sulphate of iron, add a drop or two of a solution of prussiate of potass,—by which a beautiful Prussian blue colour will be produced.

536. TEST FOR GOLD. To a diluted colourless solution of nitro-muriate of gold, add a few drops of a solution of any salt of tin—or stir the solution of gold with a slip of metallic tin: in either case, the production of a beautiful purple or port-wine colour will be the immediate result. If the mixture is allowed to settle, it becomes colourless; a purple powder (which is an oxide of gold combined with a little tin) being precipitated. This powder is employed in the painting of china, and is called the purple precipitate of Cassius.

537. TO CONVERT IRON (APPARENTLY) INTO COPPER.—Dip a piece of polished iron (as the blade of a knife) into a solution either of nitrate or sulphate of copper: it will assume the appearance of a piece of pure copper. For an explanation of this phenomenon, see 514.

538. TEST FOR COPPER—A BEAUTIFUL BLUE LIQUID PRODUCED BY MIXING TWO COLOURLESS ONES—Add a few drops of a solution of nitrate of copper to a test glass of water—the mixture will be colourless: pour into it, a little liquid ammonia—the mixture will then assume a fine deep blue colour.—*Rationale.* The alkali precipitates the copper, and then dissolves it, forming a blue liquid ammoniuret of copper.

539. TO PRODUCE A COLOURLESS LIQUID BY MIXING A DEEP BLUE ONE WITH A COLOURLESS ONE.—Add a little nitric acid to the product of the preceding experiment: upon

which the effect described will be produced.—*Rational*
The ammoniuret of copper is decomposed, and nitrate of ammonia formed. The copper re-unites to the nitric acid, and is dissolved in the water, which, on account of its quantity, renders the blue colour of the salt insensible.

540. TEST FOR LEAD.—Dissolve a grain of nitrate of acetate of lead in a glass of water, and add a little solution of sulphate of soda—a dense white precipitate (which is sulphate of lead) will be produced immediately; allow the mixture to settle, decant the clear part, and digest the precipitate in nitric acid: to the solution obtained by this means, add liquid sulphuretted hydrogen, upon which a precipitate of a black colour will be produced. Thus, in this experiment, the lead passes through the several states of an acetate, a sulphate, a nitrate, a hydrosulphuret.

541. TEST FOR SILVER.—Let fall a drop of a solution of nitrate of silver into a glassful of water, and add to it a grain of common salt. Mutual decomposition of the salts will take place, and muriate of silver (in the form of a white powder) will be precipitated—see 515.

SECTION VI.—APPLICATION OF CHEMICAL KNOWLEDGE TO THE ARTS AND TO DOMESTIC PURPOSES.

542. PREPARATION OF GINGER-BEER POWDER.—Take 2 drachms of fine loaf sugar, 8 grains of ginger, and 1 grain of carbonate of potass, all in fine powder—mix them intimately in a Wedgewood's-ware-mortar. Take also, 27 grains of citric or tartaric acid, (the first is the pleasantest, but the last the cheapest.) The acid is to be kept separate from the mixture. The beer is prepared from the powders thus: take two tumbler glasses, each half filled with water—stir up the compound powder in one of them, and the acid powder in the other: then mix the two liquors—an effervescence takes place—the beer is prepared, and may be drank off. The effervescence is occasioned by the discharge of the carbonic acid of the carbonate of potass, which is given up because the potass has a stronger affinity for the tartaric acid. If the bl

is allowed to stand for a few minutes, it becomes *flat*: this is owing to its having lost all its carbonic acid.—The cost of these powders is 8*d.* a-dozen sets.

543. METHOD OF PREPARING SODA WATER.—Soda water is prepared (from powders) precisely in the same manner as ginger beer, (see preceding exp.) except that, instead of the two powders there mentioned, the two following are used: for one glass, 30 grains of carbonate of Soda; for the other glass, 25 grains of tartaric (or citric) acid.

544. LEMONADE.—Mix one part of citric acid with six parts of finely-pounded loaf-sugar: a very fine dry lemonade is thus prepared, which may be preserved for any length of time. The quantity of this mixture necessary to be put into a glass of water to make a pleasant drink, must be regulated by the taste of the person using it.

545. INSTANTANEOUS PRODUCTION OF CURDS AND WHEY FROM MILK.—A very ready and elegant mode of procuring curds, and also a pleasant acidulous whey, is, by adding to a glassful of milk a little solution of citric acid—taking care not to add too much; an experiment or two will readily show the quantity necessary to effect the purpose.

546. TO DISTINGUISH A SOLUTION OF EPSOM SALT FROM A SOLUTION OF OXALIC ACID—1. Taste the solution; Epsom salt is bitter; oxalic acid extremely sour.—2. Pour a little tincture of litmus into the solution: if Epsom salt be present, the blue colour will remain unchanged; if oxalic acid be present, the blue will be turned to red.—3. Tincture of cabbage, or any other vegetable infusion, or a slip of litmus test-paper, are all acted upon by the acid, (which changes their colours,) but not by the salt.—4. Oxalic acid when dropped into water makes a *crackling noise*, which *Epsom* salt does not.

547. TEST FOR THE PURITY OF WINE.—Put into a phial, sixteen grains of sulphuret of lime (prepared by exposing to a red heat, in a covered crucible, equal weights of powdered lime and sulphur,) and twenty grains of super-tartrate of potass (cream of tartar.) Fill the phial with water, cork it well, and shake it occasionally for the space of ten minutes. Separate the clear liquid by decantation,

and preserve it in a well stopped bottle for use. A portion of this liquor fresh prepared, when added to wine containing lead, produces a blackish precipitate.

548. TO DETECT COPPER IN PICKLES OR GREEN TEA.—Put a few leaves of the tea, or some of the pickles cut small, into a phial, with two or three drachms of liquid ammonia, diluted with one half the quantity of water. Shake the phial: when, if the most minute portion of copper be present, the liquid will assume a fine blue colour.

549. TEST FOR THE PURITY OF VOLATILE OILS.—Many of the volatile oils being produced in small quantity, are high priced. There is, therefore, some temptation to adulterate them with fixed oils to increase the quantity. To detect the fraud, let a single drop of the suspected oil fall on clean paper, and expose it to a gentle heat. If the oil be pure, the whole will be evaporated, and no trace will remain in the paper; but if it has been mixed with a fixed oil, a greasy spot remains behind.

550. TO DISTINGUISH IRON FROM STEEL.—Let a drop of diluted nitric acid fall on the metal; and after a few minutes wash it off with water. If the metal be steel, a black spot will be left on it; if it be iron, a whitish-grey spot will remain.—*Rationale.* The nitric acid dissolves the iron in both cases; but the charcoal that enters into the composition of the steel, remains undissolved, and constitutes the blackness.

551. TO DETERMINE WHETHER WATER BE HARD OR SOFT; that is whether or not it be fit for domestic purposes.—To a glassful of the water, add a few drops of solution of soap in alcohol. If the water be pure, it will continue limpid; if it be impure, white flakes will be formed—see 520.

552. TO DETERMINE WHETHER WHEAT FLOUR OR BREAD, BE ADULTERATED WITH CHALK.—1. Mix with the flour to be tried, a little sulphuric acid: if chalk or whiting be present, an effervescence (arising from the discharge of the carbonic acid of the chalk) will take place; but, if the flour be pure, no effervescence is produced. 2. Pour boiling water on some slices of bread, and then pour into

the water a little sulphuric acid : if there be any chalk in the bread, an effervescence ensues.

553. TO DISCOVER IF BREAD IS ADULTERATED WITH ALUM.—The bread must be soaked in water, and to the water in which it has been soaked, a little of any test for Sulphuric acid must be added. (Solution of muriate of lime will do). Upon which, if any alum be present, the liquid will be pervaded with milkiness ; but, if the bread be pure, liquid will remain limpid.—*Rationale.* Sulphuric acid has a stronger affinity for lime than for the alumina and potass with which it forms alum ; it therefore quits those bodies, to form sulphate of lime, with the lime of the test—which produces the milkiness.

554. TEST FOR JELLY.—Let a grain of isinglass, glue, or any other gelatinous matter, be dissolved in a goblet-full of water, and let a few drops of tincture of galls be added to the solution : the immediate product will be an abundant flocculent precipitate. This precipitate is a compound of the *tan* of the gall, and the pure *gelatin* of the jelly.

555. TEST FOR THE PURITY OF MAGNESIA.—The common magnesia of the shops (which is a carbonate) is frequently adulterated with chalk ; this may be detected by adding a little diluted sulphuric acid, which, with magnesia, forms a very soluble salt, but with lime, a very insoluble one. Pure magnesia (called *calcined* magnesia, in the shops) dissolves in diluted sulphuric acid entirely, and without effervescence.

556. READY METHOD OF ASCERTAINING THE PREPARATION OF ALCOHOL IN WINES, BEER, CIDER, AND OTHER SPIRITUOUS LIQUORS.—*Process.* To 100 parts in volume of the liquid to be tried, add 12 parts of solution of subacetate of lead, (prepared as directed below,) a precipitation ensues ; which, by a slight agitation, is rendered general. On filtering, a colourless liquid, containing the alcohol, is procured. By mixing with this dry and warm carbonate of potass, (calcined pearl-ash,) as long as it is dissolved, we separate the water from the alcohol. The latter is seen floating above in a well-marked stratum ; the quantity of which can be estimated at once, in a mea-

sure tube, (such as figure 12).—*Preparation of the solution of sub-acetate of lead.*—Boil 15 parts of pulverized (and calcined) litharge, with 10 of acetate of lead, in 200 of water, for 20 minutes, and concentrate the liquid by slow evaporation to one half; it must be kept in well-corked phials quite full.

557. APPLICATION OF CHEMISTRY TO AGRICULTURE.

ANALYSIS OF SOILS.—Among the substances that compose soils, the following are the principal: mixtures of the *earths* (see 122); animal and vegetable matter in decomposing state; certain saline bodies; and oxide of iron; together with water. Now, in order that a soil may answer a certain purpose, these bodies should be present in particular proportions, and united in a particular manner; and the end of analytical experiments performed with soils, is to determine whether any substance is either super-abundant or wanting.

The earths commonly found in soils are principally silica, or the earth of flints; alumina, or the pure matter of clay; lime, or calcareous earth; and magnesia; for the characters of which see articles 128, 127, 125, 126. Silica composes a considerable part of hard gravelly soils, hard sandy soils, and hard stony lands. Alumina abounds most in clayey soils, and clayey loams; but even in the smallest particles of these soils it is generally united with silica and oxide of iron. Lime always exists in soils in a state of combination, and chiefly with carbonic acid, where it is called carbonate of lime. This carbonate in its hardest state is marble; in its softest, chalk. Lime united with sulphuric acid is sulphate of lime, or gypsum; with phosphoric acid, phosphate of lime, or the earth of bones. Carbonate of lime, mixed with other substances, composes chalky soils and marles, and is found in soft sandy soils. Magnesia is rarely found in soils; when it is, it is combined with carbonic acid, or with silica and alumina. Decomposing animal matter exists in different states: it contains much carbonaceous substance, volatile alkali, inflammable aëriform products, and carbonic acid. It is found chiefly in lands lately manured. Decomposing vegetable matter usually contains still more carbonaceous

substance, and principally differs from the preceding, in not producing volatile alkali. It forms a great proportion of all peats, abounds in rich mould, and is found in larger or smaller quantities in all lands. The saline compounds are few, and in small quantity: they are chiefly muriate of soda, (common salt,) sulphate of magnesia, muriate and sulphate of potash, nitrate of lime, and the mild alkalies. Oxide of iron, which is the same with the rust produced by exposing iron to air and water, is found in all soils, but most abundantly in red and yellow clays, and red and yellow siliceous sands.

Process 1. When the general nature of the soil of a field is to be ascertained, specimens of it should be taken from different places; the total quantity submitted to analysis may be 400 grains. For weighing the various products of analysis, a pair of scales, capable of holding a quarter of a pound of common soil, and turning with a grain when loaded, must be provided. Soil, when collected, if it cannot be immediately examined, should be preserved in phials quite filled with it, and closed with ground glass stopples. It should be collected in dry weather, and exposed to the air till it feels dry. The first step to be taken, is an examination of the physical properties of the soil, because *they*, to a certain extent, denote its composition, and therefore serve as guides in directing the experiments: thus, it is of importance that the *specific gravity* of it should be known, as it affords an idea of the quantity of animal and vegetable matter it contains, these substances being always most abundant in the lighter soils; thus, also, a roughness to the touch, and a capability of scratching glass when rubbed upon it, denote siliceous soils; if a soil adheres strongly to the tongue, and emits a strong earthy smell when breathed upon, it is aluminous; and if it is soft, and possesses little adheviseness, it is calcareous.

Process 2. The next step is to free the soil as much as possible with water, without, in any other respect, affecting its composition. This is done by heating the soil for ten or twelve minutes in the Wedgewood's-ware basin, fig. 4, by means of the lamp-furnace; the proper degree

of heat being ascertained by keeping a piece of wood in contact with the bottom of the basin : the process must be stopped, as soon as the wood begins to be charred. The loss of weight in this process must be carefully noted ; it amounts to 50 grains in 400 of the soil, then clay, aluminous earth will be abundant ; if it amounts to about 10 or 20 grains in 400, the soil is to be considered as chiefly siliceous.

Process 3. Now, gently bruise the soil in the Wedgwood's-ware mortar, and separate from it, (by means of a wire sieve, coarse enough to let a pepper corn pass through) the large vegetable fibres, and loose stones ; note down separately the weights of these substances, and let the nature of the latter be ascertained. If the stones or gravel be calcareous, they will effervesce with acids ; siliceous, they will scratch glass ; if aluminous, they will be soft, easily dissolved and incapable of effervescing with acids.

Process 4. Mix the soil well with water ; let it settle for a minute or two, during which time, the coarse sand will sink to the bottom ; then, pour off the water in which the fine part of the soil remains suspended through a filter. Preserve the water ; it will contain the soluble matter of the soil. Dry and weigh the coarse sand, and also, what remains on the filter ; and note down their respective quantities.

Process 5. The coarse sand thus separated is always siliceous, or calcareous sand, or a mixture of the two ; its nature may be detected in the same way as that of the stones and gravel. (*Process 3.*) If it consist wholly of carbonate of lime, it will dissolve rapidly, with effervescence in diluted muriatic acid (the experiment may be tried in a glass tumbler ;) but if it consist partly of this and partly of siliceous matter, a residuum will be left after the acid has ceased to act upon it, the acid being added till the mixture has a sour taste, and has ceased to effervesce. This residuum is the siliceous part ; which being washed (by being placed in a filtering paper, and having rain water poured over it,) dried, and heated strongly in a crucible, shows, by the difference of its weight from that of the whole, the quantity of the calcareous sand.

Process 6. The purely divided matter of the soil (separated by *Process 4*.) sometimes contains the four usual earths of soils, as well as animal and vegetable matter; and it is the most difficult part of *the analysis* to ascertain with accuracy the proportions of these. Put the earthy matter into the evaporating basin, fig. 4, and add twice its weight (measured as directed 305) of a mixture of one part of muriatic acid with two parts of water. Stir the mixture often with a glass rod, and suffer it to remain for an hour and a-half before it is examined. During this time, the acid will have dissolved (what was present) of the carbonate of lime, or of magnesia; and, sometimes too, a little oxide of iron. Filter the mixture; wash the solid matter; dry it at a moderate heat, and weigh it; add the washings to the solution. The weight lost by the solid matter denotes the quantity of it dissolved by the muriatic acid. The solution must be made sour to the taste (if not so already) by the addition of fresh acid. Into this solution drop a solution of prussiate of potass: if a blue colour is produced, continue to drop it in, till the production of a blue precipitate ceases. Collect the precipitate on a filter, in the same manner as the other precipitates, and heat it red; the result will be oxide of iron, which must be weighed. Into the remaining solution, drop solution of carbonate of potass till effervescence entirely ceases, and till the solution has a strongly alkaline taste. Collect the precipitate by filtration, dry it at a heat below redness and weigh it—it is carbonate of lime. Boil the filtered fluid (in a Florence flask) for a quarter of an hour; when the magnesia (if any be present) will be precipitated, combined with carbonic acid: proceed with this as with the carbonate of lime.

Process 7. To ascertain the quantity of insoluble animal and vegetable matter contained in the solid matter separated in the preceding process, from the muriatic solution, heat it in a crucible, over a common fire, at a strong red heat, till the mass retains no blackness, stirring frequently meanwhile with a metallic wire. The loss of weight shows the quantity of animal and vegetable matter there was, but not the proportions of each. The

presence of animal matter is indicated by a smell of burnt feathers emitted during the process ; and a copious blue flame generally denotes a considerable proportion of vegetable matter.

Process 8. What remains now is a mixture of silica, alumina, and oxide of iron. To separate these, boil the residue for 2 or 3 hours, in sulphuric acid diluted with four times its weight of water, allowing about 6 grains of acid to every 5 grains of the residuum. At the end of this time whatever remains undissolved is silica, which may be separated, washed, dried, and weighed, in the usual manner. Carbonate of ammonia being added to the solution in quantity more than sufficient to saturate the acid, the alumina will be precipitated, and the oxide of iron, if any, may be obtained by boiling away the remaining liquid.

Process 9. Let the filtered liquid of *process 4*, be evaporated to dryness, at a heat below boiling : the nature of the residuum (if any) may be ascertained as follows : if it be of a brown colour, and inflammable, it is vegetable extract ; if, when heated, its smell be strong and fetid, animal mucilaginous, or gelatinous matter. If it be white and transparent, it is saline. Nitrate of potash or of lime in this saline matter is indicated by its sparks when thrown on burning coals : sulphate of magnesia may be detected by its bitter taste : and sulphate of potash produces no alteration in a solution of carbonate of ammonia, but gives a precipitate with a solution of muriatic barytes.

Process 10. A particular operation is required to detect sulphate of lime (or gypsum) and phosphate of lime in soil. A given weight (400 grains) of the entire soil must be mixed with one-third as much powdered charcoal, and kept at a red heat, in a crucible, for half an hour. The mixture must then be boiled a quarter of an hour, in a pint of water, and the solution, being filtered, exposed some days to the open air. A white precipitate will form in the fluid, which, being weighed, gives the proportion of sulphate of lime existing in the soil.—To separate

phosphate of lime, add to the soil, after the process for gypsum, muriatic acid, more than sufficient to saturate all the soluble earths; form a solution by digestion; then, evaporate the solution to dryness, and pour water upon the solid matter. This will dissolve the muriates, and leave the phosphate of lime untouched.

When the experimenter is become acquainted with the use of the different instruments, the properties of the reagents, and the relations between the external and chemical qualities of soils, he will seldom find it necessary to perform, in any one case, all the processes that have been described. When his soil, for instance, contains no notable proportion of calcareous matter, the action of the muriatic acid, *Process 6*, may be omitted: in examining peat soils, he will have to attend principally to the operation by fire and air, *Process 7*: and in the analysis of chalks and loams, he will often be able to omit the experiment with sulphuric acid, *Process 8*.

The examination of a soil being completed, the products are to be classed, and their quantities added together; and if they nearly equal the original quantity of soil, the analysis may be considered as accurate. When, however, products are obtained by *Process 10*, a sum equal to their weights, must be deducted from the weight of the carbonate of lime obtained by precipitation, in *Process 6*. In arranging the products, the form should be in the order of the experiments, by which they are obtained. Thus, 400 grains of a good siliceous sandy soil, may be supposed to contain,

	grains.
Of water of absorption, - - -	18.
Of loose stones and gravel, chiefly siliceous, -	42.
Of undecomposed vegetable fibres, - -	10.
Of siliceous sand, - - - -	200.
Of minutely-divided matter, consists of	
Carbonate of lime, - - -	25.
Carbonate of Magnesia, - -	4.
Matter destructible by heat, principally vegetable, - - -	10.
Carried over, - - - -	39. 270.

Brought forward, - - - -	39. 27
Silica, - - - -	40.
Alumina, - - - -	32.
Oxide of iron, - - - -	4.
Soluble matter, principally sulphate of potass, and vegetable extract, -	5.
	— 122
Of sulphate of lime, - - - -	
Of phosphate of lime, - - - -	
	—
Amount of all the products, - -	39
Loss, - - - -	
	40
	—

In this instance the loss is supposed small, but, in general, in actual experiments, it will be found much greater in consequence of the difficulty of collecting the whole quantities of the different precipitates; and when it is within 30 for 400 grains, there is no reason to suspect any want of due precision in the processes.

In the first trials that are made by persons unacquainted with chemistry, they must not expect much precision of result. Many difficulties will be met with; but in overcoming them the most useful kind of practical knowledge will be obtained; and nothing is so instructive in experimental science as the detection of mistakes. The correct analyst ought to be well grounded in general chemical information; but perhaps there is no better mode of gaining it than that of attempting original investigations. In pursuing his experiments, he will be continually obliged to learn from books the history of the substances he is employed or acting upon; and his theoretical ideas will be more valuable in being connected with practical operation and acquired for the purpose of discovery.

558. ANALYSIS OF MARLS.—The ingredients of marl on which their fitness for agricultural purposes depends is carbonate of lime. To find the composition of a marl pour a few ounces of diluted muriatic acid into a Florence flask, place the flask in a scale, and counterpoise it. The

reduce some dry marl to powder, take a few ounces of this powder, and drop it carefully and gradually into the diluted acid, until an addition ceases to cause effervescence. Let the remainder of the quantity of marl taken be weighed, by which the quantity projected will be known. Let the balance be then restored. The difference of weight between the quantity projected, and that requisite to restore the balance, will show the weight of air lost during effervescence. This air is carbonic acid gas. For every 45 grains of which that escape, reckon 100 grains of carbonate of lime as present in the projected quantity of marl. If the loss (by the escape of gas) amount to 13 *per cent* of the quantity of marl projected, or from 13 to 32 *per cent*, the marl assayed is rich in calcareous earths. Clayey and sandy marls lose only 8 or 10 *per cent* by this treatment.—The presence of much argillaceous earth may be judged by drying the marl after being washed by muriatic acid, when it will harden and form a brick. To determine, with still greater precision, the quantity of calcareous earth in a marl, let the muriatic solution be treated with carbonate of potass, as described at *Process 6* of the preceding Analysis. NOTE:—In some of the processes instituted to determine the quantity of carbonic acid discharged by another acid, errors arose, in consequence of a portion of water being thrown from the vessel along with the gas, by the violence of the effervescence. To remedy this inconvenience, the instrument described at 359 was contrived. A given quantity of any matter to be examined—as carbonate of potass or of lime—being introduced into a bottle, and weighed, and a given weight of diluted acid being added, the cork affixed to the bottom of the bended tube is *instantly* inserted in the neck of the bottle. Any water therefore that rises with the gas, is condensed in the worm, and runs down again into the vessel, nothing but gas escaping.

559. HOW TO IMPROVE SOILS.—Having given an abstract of Sir Humphry Davy's method of analyzing soils, we subjoin the rules he has laid down for improving them. In cases where a barren soil is examined with a view to its improvement, it ought if possible to be compared with an

extremely fertile soil, situated in a similar place: the difference given by their analysis will indicate the means of improvement. If, for instance, the fertile soil contain much more sand than the barren one, then sand is the material which the barren soil is in need of; and so of other kinds of matter. In the application of clay, marble, chalk &c., to lands, there are no particular chemical principles to be observed; but when quicklime is used, great care must be taken that it is not obtained from the magnesian limestone, for such lime is exceedingly injurious to lands. The magnesia limestone may be distinguished from the common limestone by the slowness of its solution in acids—the softest kind of it being much longer in dissolving than marble. When the analytical comparison indicates an excess of vegetable matter as the cause of sterility, it may be destroyed by much pulverization and exposure to air, by paring and burning, or by the agency of newly-made quicklime. And a defect of animal and vegetable matter must be supplied by suitable manure.

560. INSTANTANEOUS LIGHT-BOXES, sold by Chemists, contain a little phial filled with a liquid, and a number of small matches: you take a match and dip it into the liquid upon which it takes fire.—The *liquid* is concentrated sulphuric acid. The bottle containing it is never opened except when it is to be used; for the acid, when exposed to the air, imbibes moisture very rapidly, and is soon spoiled. The *matches* are prepared as follows: the ends of some small slips of light wood are dipped into a strong solution of gum, and afterwards into the mixture of chlorate of potass and sulphur prepared as described 591: the powder is fastened to the wood by the gum, and the matches when dry are fit for use.

561. ILLUSTRATION OF THE ART OF CALICO-PRINTING.—*Process 1.* To produce a white pattern on a black ground. Let a piece of white calico be dipped in a cold solution of sulphate of iron, and dried. Then imprint any figure upon it with a strong solution of colourless citric acid, and let this dry also. If the piece be then well washed in warm water, and afterwards boiled in a decoction of logwood, the ground will be dyed either a slate or black colour.

lour, according to the strength of the metallic solution, while the printed figures will remain beautifully white.

562. *Process 2.* TO PRODUCE A SCARLET PATTERN ON A BLACK GROUND.—Dip a piece of white calico in a strong solution of acetate of iron (which may be made by boiling sulphate of iron with acetate of lead); dry it by the fire and lay it aside for 3 or 4 days. After this, wash it well in hot water, and then dye it black by boiling it for ten minutes in a strong decoction of logwood; and, lastly, rinse it in water, and let it dry. Now, any figures printed upon it with a *colourless* solution of muriate of tin, will acquire a beautiful scarlet colour,—though the ground remains black.

563. *Process 3.*—OF DYING DIFFERENT COLOURS WITH THE SAME LIQUOR.—Dissolve indigo in sulphuric acid, and add to the solution an equal quantity of solution of carbonate of potass.—If a piece of *white* cloth be dipped in this mixture, it will be changed to *blue*: in the same way, *yellow* may be changed to *green*; and *red* to *purple*; and a piece of *blue* litmus paper will become *red*.

564. EXTEMPORANEOUS FORMATION OF A BLEACHING LIQUID.—Add a few grains of chlorate of potass to a spoonful of muriatic or sulphuric acid diluted with half a wine-glassful of water. The liquid thus formed will possess the bleaching property of that described 474.

565. MOIREE METAL, OR CRYSTALLIZED TIN.—Clean a piece of tin from dirt or grease, by well washing it with warm soapy water; and rinse it in clean water. Then, heat the tin to the temperature of bare sufferance to the hand, and pour on it, or apply with a brush or sponge, one of the following mixtures.—1. Dissolve two ounces of muriate of soda in four ounces of water, and add one ounce of nitric acid.—2. To two ounces of water, add half an ounce of muriatic acid, and one fourth of an ounce of sulphuric acid.—3. Either of the preceding in a greater or less degree of dilution.—When the mixture has been applied, the tin is to be dipped into water slightly acidulated, and afterwards well washed in clean water. This process will cause the tin to assume an appearance of great beauty: an apparent crystallization takes place, somewhat resembling mother of

pearl; and curious figures, stars, and rude leaves appear. A transparent varnish (either coloured or colourless) may be laid on now, which will give a fine polish to the work, and produce that brilliant covering so much employed for snuffer-stands, for covering pillars in shop-windows, &c.

GOLD-VARNISH to be laid on the tin which has been ornamented by undergoing the process just described. —Take of shell-lac two ounces, of arnatto and turmeric of each one ounce, of dragon's blood thirty grains: make an extract with twenty ounces of alcohol in a gentle heat.

566. TO MAKE THE ADMIRER COSMETIC CALLED PEARL POWDER. —Dissolve bismuth in nitro-muriatic acid, and gradually add to the solution cold distilled water: upon which, a beautiful white powder, which is an oxide of bismuth, will be precipitated.

567. PREPARATION OF WRITING INK. —*Process 1.* Take two ounces of gall-nuts (in coarse powder); one ounce of logwood (in thin chips); one ounce of sulphate of iron (green vitriol); three-fourths of an ounce of gum-arabic (in powder); one-fourth of an ounce of sulphate of copper (blue vitriol); and one-fourth of an ounce of sugar-candy. Boil the galls and logwood together in three pints of water, till the quantity is reduced to one-half; when the liquor must be strained through a flannel into a proper vessel, and the remainder of the ingredients be added to it. The mixture is now to be frequently stirred, till the whole is dissolved: after which, it must be left at rest for 24 hours. The ink may then be decanted from the gross sediment, and must be preserved in a glass, or stone bottle, well corked. —Writing performed with this ink has a beautiful black cast, which it retains, unaltered, for a considerable length of time. —There is an inconvenience attends the use of the sulphate of copper (which some, therefore, dispense with) that requires notice: —upon mending a pen, which is moist with the ink, a film of copper is deposited upon the edge of the penknife, and takes considerably from its keenness. NOTE: —Ink-holders ought always to be made of glass.

568. *Process 2.* —Take four ounces of the best galls; two ounces and a half of sulphate of iron calcined to white-

ness; two pints of water; and ten drachms of gum-arabic. The galls, and calcined vitriol, are to be left to macerate for 24 hours, then the gum is to be added, and the mixture is to be preserved in a stone jar, open, or merely covered with paper.

569. INDELIBLE INKS, USEFUL FOR WRITING LABELS FOR BOTTLES CONTAINING ACIDS, &c.—1. Take oil of lavender, 200 grains; gum copal, in powder, 25 grains; and lamp black, 3 grains. Dissolve the copal in the oil of lavender, in a small flask or phial, by the aid of a gentle heat; and then mix the lamp-black with the solution, by trituration in the Wedgewood's-ware mortar. After a repose of some hours, the ink, before use, requires to be shaken, or must be stirred with an iron wire. If it be found too thick it may be diluted with a little oil of lavender, or of turpentine.—2. A solution may be made, in like manner, of 120 grains of oil of lavender; 17 grains of copal, and 60 grains of vermilion.—3. Boil one ounce of finely-rasped Brazil-wood, and half an ounce of alum, in twelve ounces of water, till the liquid is reduced to eight ounces; then decant the clear part, and add to it an ounce of calcined manganese mixed with half an ounce of gum-arabic.

570. COLOURED INKS.—*Red.* Boil an ounce of Brazil-wood (in fine chips) in half a pint of water, for a quarter of an hour; and add to the decoction, 3 drachms of gum-arabic, and as much alum as it will dissolve.—*Blue.* Diffuse Prussian blue or indigo through strong gum-water.—*Yellow.* Dissolve gamboge in gum water.—*Scarlet.* Dissolve vermilion in gum-water.—Inks of other colours may be made from a strong decoction of the materials used in dying, mixed with a little alum and gum-arabic.

571. INK POWDER.—For the use of travellers, who may not find it convenient to carry liquid ink with them, the materials used for producing good ink—see 568, 569—may be finely powdered, and intimately blended together; by which means, an ink powder is made, from which, at any time, good ink may be extemporaneously prepared, by mere mixture with water.

572. TO REMOVE INK STAINS FROM BOOKS, LINEN, &c.—Apply to the spot, muriatic acid diluted with five

or six times its weight of water, and, after a minute or two wash it off; repeating the application as often as may be found necessary. Strong solutions of oxalic, citric, and tartaric acids, also effect the purpose, and, being less likely than the muriatic acid to injure the fabric they are applied to, are preferable.

573. PERMANENT INK FOR MARKING LINEN.—Dissolve a drachm of lunar caustic (fused nitrate of silver) in three times its weight of distilled (or rain) water, and add about half a drachm of gum-arabic. This forms the *ink*, with which you must write, or rather *paint*, with a clean pen, upon the linen prepared as follows: dissolve half an ounce of sulphate of soda in an ounce of distilled water, and add 20 grains of gum-arabic; this forms the *mordant*. Wet and moisten, with the mordant, the part of the linen that is to be written on, dry it well by a gentle heat, and then apply the ink in the manner above directed, to the place that has been moistened. The writing, when exposed to the sun, becomes black.

574. INK FOR PRINTING ON LINEN WITH TYPES.—Dissolve one part of asphaltum in four parts of oil of turpentine, and add lamp-black, or black-lead, in fine powder, in sufficient quantity to render the ink of a proper consistence for printing with types.

SECTION VII.—ON COMBUSTION, DETONATION AND LIGHT.

575. WRITING WHICH IS LUMINOUS IN THE DARK.—Trace letters or figures on a smooth board, or on dark coloured paper, with a stick of phosphorus, in the same manner as you would trace them with a crayon. Every line thus made will be beautifully luminous in the dark, and will continue so for some minutes.

576. *Rationale*.—The luminous appearance of the writing arises from a slow combustion of the phosphorous which adheres to the board.—In all experiments with this substance, the greatest degree of care is required, on account of its very combustible nature. A very small portion of it

(the quantity is generally mentioned in the experiments) should be operated upon at once. When it is taken in the hands, it never should be held for more than a few seconds, for the heat thus applied is sufficient to inflame it, if continued; and a burn from phosphorous is more painful than any other kind of burn. A basin of cold water ought always to be at hand, to dip the phosphorus in occasionally; and, when it is cut to pieces it must be cut in water. Phosphorus can only be preserved by keeping it in places where neither light nor heat has access. It is obtained from druggists in rolls about the thickness of a quill; these are put into a phial filled with cold water, which has been boiled to expel air from it, and the phial is inclosed in an opaque case.

577. TO SHOW THE INFLAMMABLE NATURE OF PHOSPHORUS.—1. Wrap a grain of it, dried on blotting paper, in a piece of brown paper, and rub it with some hard body; it will set fire to the paper.—2. Put into the middle of some dry cotton, a piece of phosphorus the size of a large pin's head, (previously dried, as before): strike it with a hammer and it will inflame.—3. Upon a piece of glass, lay a small piece of phosphorus, and place the glass upon the surface of hot water in a basin: the phosphorus will inflame.

578. PREPARATION OF PHOSPHORIZED ETHER.—Suffer sulphuric ether to stand, for some weeks, over a quantity of phosphorous in a well-stopped phial. The solution is aided by occasional agitation.

579. PREPARATION OF PHOSPHORIZED OIL.—Put one part of phosphorus with six parts of olive-oil into a Florence flask, and digest the mixture in a gentle sand heat for two hours. The solution must be kept in a dark place.

580. TO MAKE WAVES OF FIRE ON THE SURFACE OF WATER.—On a lump of loaf sugar, let fall a few drops of phosphORIZED ether, and place the sugar in a glass of warm water; a very beautiful appearance will be instantly exhibited, and the effect is increased, if the surface of the water is made to undulate, by blowing gently with the breath.

581. TO MAKE FACES AND HANDS LUMINOUS ; SO THAT IN THE DARK, THEY APPEAR AS IF ON FIRE.—Though the phosphorized oil and ether are luminous in the dark, yet they have not the power to burn any thing ; so that either of them may be rubbed on the face and hands without danger ; and the appearance thereby produced is most hideously frightful. All the parts of the face that have been rubbed, appear to be covered with a luminous blueish flame, and the mouth and eyes appear as black spots.—When the bottles containing phosphorized oil and ether are opened in the dark, light enough to tell the hour on a watch is evolved.

582. BRILLIANT COMBUSTION UNDER WATER.—Drop a piece of phosphorus, half the size of a small pea, into a glass of hot water. It will immediately inflame. Then force upon it, from a bladder with the jet-pipe, fig. 44 fitted to it, a stream of oxygen gas. Upon this, there will be produced a flame of great vividness.

583. PREPARATIONS FOR THE INSTANTANEOUS PRODUCTION OF LIGHT.—Put a little phosphorus, dried on blotting paper, into a small phial ; heat the phial by placing it in a ladle full of hot sand, and turn it round, so that the melted phosphorus may adhere to its sides. Cool the phial closely, and it is prepared. Another method of preparing it, consists in mixing one part of flowers of sulphur, with eight parts of phosphorus.—On putting a common sulphur match into this fire-bottle, stirring about a little, and then withdrawing it into the air, it will take fire. Sometimes, however, it is found necessary to rub the match when withdrawn from the phial on a cork before it will inflame.

584. CURIOUS PROPERTY OF PHOSPHURET OF LIME.—*Process 1.* Drop a small piece of phosphuret of lime into a wine-glass of water : in a short time, bubbles of gas will be produced, which, rising to the surface of the water, will take fire and explode. After each explosion, a beautiful column of white smoke will ascend from the glass.—*Rationale*, see 152.—If the phosphuret of lime be recently fresh made, it may be proper to warm the water it is added to. If the residue of the phosphuret of lime

taken out of the water, and, after being dried, has muriatic acid poured upon it, it will inflame.

585. *Process 2.* Into the glass, fig. 6, put one part of chlorate of potass, and two parts of phosphuret of lime, in pieces about the size of peas (not in powder). Fill the glass with water, and put into it the funnel, fig. 10, which will reach to the bottom of it. Through this, pour six or eight parts of strong sulphuric acid, which will decompose the chlorate; and the phosphuret also decomposing the water at the same time, flashes of fire dart from the surface of the fluid, and the bottom of the vessel is illuminated by a beautiful green light.

586. *METHOD OF PREPARING AND USING A FULMINATING POWDER.*—Take three parts of nitrate of potass, two parts of sub-carbonate of potass, and one part of sulphur. Powder them separately, and dry them, by placing them on a tile before the fire. Then, mix them intimately, by rubbing all together in a warm mortar; and preserve the compound in a corked phial.—Let half a drachm of this powder be spread on the bottom of a small iron ladle, and heated over a fire. It gradually blackens, and at last melts, at which instant, it explodes with a violent report: it is not, however, attended by any danger. The explosion is occasioned by a sudden expansion of the solid powder into a variety of gases.

587. *TO MAKE GUNPOWDER.*—Pulverise separately, 5 drachms of nitrate of potass, 1 of sulphur, and 1 of newly-burnt charcoal. Mix them together, with a little water, in a mortar, so as to make the compound into a dough; which must be rolled out into round pieces, the thickness of a pin, between two boards. Lay a few of these pieces together, and cut them with a knife into small grains, which are to be placed on a sheet of paper, in a warm place, to dry. During granulation, the dough must be prevented sticking to the board, by rubbing on it a little of the dry compound powder. The effects produced by the explosion of gunpowder, are so well known, that it would be useless to describe them. It may be observed, however, that the explosion takes place in consequence of the generation of a large quantity of various gases.

588. TO PRODUCE COLOURED FLAMES, *which, in dark, are very curiously reflected by the faces of the spectators.*—To do this, certain substances are mixed with burning alcohol. A beautiful *rose*-coloured flame is produced by inflaming four parts of alcohol, poured over one part of muriate of strontia, in a small iron ladle. An *orange*-coloured flame is produced, by burning spirit of wine, on muriate of lime deprived of its water of crystallization.—A flame having a fine *green* tinge, is produced by burning alcohol on boracic acid, or nitrate of copper.—A *yellow* flame is produced, by burning alcohol on muriate of soda, or nitrate of potass.—NOTE. It should be observed, that the ladle ought previously to be warmed and ought to be kept heated while the alcohol is burning. The salts remaining behind, after being dried, may be used for the same purpose again.

589. A METAL WHICH BURSTS INTO FLAME WHEN THROWN UPON COLD WATER.—Place a piece of potass of about two grains weight, upon cold water in a basin—the effect is described at 60.

590. EXHIBITION OF THE POWERFUL DETONATING PROPERTIES OF ALL MIXTURES OF INFLAMMABLE BODIES WITH CHLORATE OF POTASS.—NOTE: In the performance of experiments with chlorate of potass, proceed with caution. Never use more than the prescribed quantities. Let the hands be covered with stout gloves, keep the mixtures, and vessels containing them, as far as possible from the face. Chlorate of potass must never be kept ready mixed with inflammable bodies; for such mixtures sometimes explode spontaneously.

Process 1.—Rub two grains of chlorate of potass to a fine powder in a mortar, and add one grain of flower of sulphur. Mix them very accurately, by well triturating them in the gentlest possible manner; and then, pressing collected the mixture to one part of the mortar, press the pestle down upon it suddenly and forcibly—a detonation, accompanied by a flash of light, will instantly ensue.

591. *Process 2.*—Let the mixture of the salt and sulphur, just described, be wrapped in some strong paper.

or tin-foil, and then struck with a hammer—a still louder report will be produced.

592. *Process 3.*—Let *half* the quantity of a similar mixture be forcibly triturated in a dry mortar—several explosions, like the cracking of a whip, accompanied by flashes of purple light, take place.

593. *Process 4.*—Four grains of the salt are to be reduced to a fine powder, and then mixed with two grains of finely-pulverized charcoal. To this mixture, rather more than one grain of sulphur is to be added; and the whole must be intimately incorporated by mixture, with the least possible friction, on a piece of paper. If this mixture be triturated, it will burst into flame—but not with much noise.

594. *Process 5.*—Mix gently, but intimately, three grains of the salt, with three grains of loaf sugar, both previously reduced to fine powder. Place the mixture on a plate, and touch it with a thread which has just been dipped in strong sulphuric acid—a sudden and vehement inflammation will be immediately produced.—It will be prudent to fasten the string to the end of a stick.

595. *Process 6.*—Upon one grain of the powdered salt, in a mortar, place half a grain of phosphorus; and then, strike the phosphorus with a pestle—a violent detonation takes place instantly.—The utmost degree of caution is requisite in this experiment.

596. *Process 7.*—Put into the glass, fig. 17, one part (about 3 grains) of phosphorus, and two parts (6 grains) of the salt; nearly fill the glass with water; and then convey to the bottom of it, by means of the funnel, fig. 10, three or four parts of sulphuric acid.—A violent action ensues, and the phosphorus burns very vividly, with a curious light, under the water.

597. *Process 8.*—Oil may be inflamed on the surface of water, by repeating the preceding experiment, omitting however, the phosphorus, and substituting a little olive or lintseed oil.

598. *Process 9.*—If a slip of cotton cloth be dipped in a strong solution of the salt, and afterwards well dried,

upon being rubbed in a mortar, it will emit flashes of fire with explosions similar to the cracking of whips. If the cloth and the mortar be very dry and warm, the trituration causes the cloth to take fire.—Inflammation is also produced by pouring sulphuric acid upon a similar piece of cloth.

599. *Process 10.*—If a little of the mixture described at *process 1*, be taken on the point of a knife, and dropped into a wine-glass containing sulphuric acid, a beautiful column of flame will be produced.

600. CURIOUS INSTANCE OF COMBUSTION RESULTING FROM INTENSE CHEMICAL ACTION.—Coarsely powdered few crystals of nitrate of copper, lay them on a piece of tin-foil, add enough of water to make them into a kind of paste, and then quickly fold up the tin-foil, doubling the sides and corners well together, so as to exclude air. In a short time, nitrous gas will force its way out of the packet, and the tin-foil will be set on fire.

601. FIRE PRODUCED BY THE MIXTURE OF TWO COOL LIQUIDS.—Into a gallipot, placed upon a hearth, pour about 3 teaspoonsful of oil of turpentine; then, mix in a phial about 3 teaspoonsful of fuming nitrous acid with $\frac{1}{4}$ part of sulphuric acid, and pour this mixture suddenly upon the oil of turpentine. Instantaneous inflammation accompanied by the production of a large quantity of black smoke, is the result.—It is adviseable to fix the phial from which the acid is poured, to the end of a long stick, as the sudden combustion sometimes occasions a part of the liquid to be thrown out of the vessel.

602. LAMP WITHOUT FLAME.—Take platinum wire about 1-100th of an inch in thickness. Coil it up, and stick the coil loosely on the wick of a spirit lamp. The cotton of the lamp must be very straight, and not pressed by the wire. There should be about 16 spiral turns, of half of which should surround the wick, and the rest is above it. Having lighted the lamp for an instant, and blowing it out, the wire will become brightly ignited, and will continue to glow as long as any alcohol remains.

603. A WIRE INSTANTLY MADE RED-HOT BY BEING

PLACED IN CONTACT WITH A VAPOUR.—Let a few drops of ether be thrown into a cold glass, or a few of alcohol into a warm one; let a few coils of platinum wire, of the 1-60th or 1-70th of an inch in thickness, be heated to redness by a candle, then, when it has ceased to be red-hot, let it be held in the glass over the ether: in some parts of the glass, it will become glowing, almost white-hot, and will continue so, as long as a sufficient quantity of vapour and air remains in the glass.

604. BRILLIANT INFLAMMATION, WITHOUT THE PRESENCE OF AIR.—Mix intimately together, eight parts of copper filings, and two parts of flowers of sulphur, and expose the mixture in the glass vessel, fig. 53, to a gentle heat, by means of the lamp furnace. As soon as the sulphur is heated to a little above its melting temperature, combustion suddenly pervades the whole mass, and the two bodies unite with an explosion—the result being the formation of *sulphuret of copper*. The most curious circumstance attending this experiment is, that it succeeds equally well if the vessel is closed so as to exclude air, or even if filled with *nitrogen gas*. Here, then, is an instance of combustion taking place, without the presence of either oxygen or chlorine. So that the theory which terms those bodies the supporters, and the only supporters, of combustion, is erroneous. The modern notion of combustion, that “it is the result of intense chemical action,” appears to be the more correct one.

605. TO MELT A PIECE OF METAL IN A NUTSHELL.—Take three parts of nitre, (freed from water of crystallization by exposure to heat,) one part of sulphur, and one of very fine dry saw-dust—mingle the whole intimately together. This is called the *powder of fusion*, and is a kind of chemical flux. Let a quantity of this be well-pressed into a walnut-shell, with a thin piece of copper coiled up in the midst of it, and then set on fire: it will burn rapidly, and the metal will be fused into a round globule, while the shell is only blackened.—A combination takes place between the metal and sulphur, (which is aided by the potass,) and the result is the formation of a sulphuret.

606. TO MAKE A POWDER WHICH INFLAMES ON EXPOSURE TO AIR.—To the substance possessing this property, is given the name of *Homburg's pyrophorus*: it is prepared in the following manner. Equal parts of alum and coarse brown sugar are to be mixed together and dried over the fire in an iron ladle, being diligently stirred all the while with an iron rod. The mixture melts, becomes thick and swells up, and runs into small dry lumps. These are coarsely powdered in a mortar, and again roasted till the operator is well assured that the mass contains not the least moisture, when it looks like a blackish powder of charcoal. This, while hot, must be put into a common phial which must be previously luted on the inside, by being washed with a weak solution of borax, made as thick as cream, by the addition of pipe-clay, and which must have a narrow glass tube, six inches long, and open at both ends, luted into its neck. The phial, which, however, is to be filled three-fourths full only, is to be placed in a crucible, or deep iron pan, and covered with sand; and the crucible is to be put into a fire, and heated to redness: a thick smoke will rise from the tube for a quarter of an hour, which is succeeded by a sulphurous vapour, which must be inflamed. When this flame disappears, the operation is completed: the tube must be then closed by a plug of soft clay, and the crucible must be removed from the fire. As soon as the phial becomes cool enough to be handled, the contents of it are to be hastily transferred into a dry and warm stout glass phial, which must be secured by a ground glass stopple.—*Experiment*. Throw a little of this powder from the bottle upon some flat dry surface: in a short time it will burst into flame.—*Rationale*. The sulphuret of potass derived from the alum attracts moisture, or (perhaps) a little potassium formed in the process, attracts oxygen, from the air, and generates sufficient heat to kindle the carbonaceous matter mingled with it.

607. PHOSPHORESCENCE —1. If coarsely powdered fluo spar be sprinkled in a dark room, on a fire shovel made hot, (but not to redness,) a beautiful phosphorescent light will be emitted for some time.—2. “About 4 drachms

of the substance of a fresh herring were put into a solution of 2 drachms of sulphate of magnesia, in 2 ounces of water. On the succeeding evening, the whole of the liquor, upon shaking the phial, became beautifully luminous; and it continued luminous till the fourth night."

—3. Herrings, whittings, and many other fish, if examined in the dark, shortly after being caught, will be found to possess a considerable share of this phosphorescence. The English glow-worm, the Chinese lanthorn-fly, and the West-India fire-fly, are *living* phosphorescent bodies. The cause of the curious property thus possessed by some bodies is unknown.

608. TO MAKE CANTON'S PHOSPHORUS.—Take thick oyster shells, wash them, and keep them red-hot in an open fire for half an hour: then, select the whitest parts, and pound them in the Wedgewood's-ware mortar. Mix three parts of this powder with one part of flowers of sulphur, and ram the compound into a crucible, till it is nearly full. The crucible is then to be kept red-hot for at least an hour; and when it has afterwards become cold, the contents of it are to be turned out, and the whitest part scraped off, and preserved in a phial with a ground-glass stopple. If the bottle containing this phosphorus be exposed for a few minutes to the light of the day, and be then taken into the dark, light enough will be emitted to enable one to distinguish the figures on the dial-plate of a watch. If the bottle containing the phosphorus is heated, the luminous appearance increases; if cooled, it decreases. The phosphorescence ceases after a while; but is renewed when the phosphorus is again exposed to light.—*Rationale.* It is generally supposed, that light is capable of entering into bodies, and, after remaining in them, of being extricated unaltered.

609. TO PREPARE BALDWIN'S PHOSPHORUS, WHICH EMITS LIGHT IN THE DARK.—Put some dry nitrate of lime into a crucible, place it in a clear fire, and let it remain in a state of fusion for about ten minutes; then pour it out into a warm iron vessel and it will become solid; break it into pieces, and enclose them in a well-stopped phial. If this phial be exposed for some hours to the di-

rect rays of the sun, and be then taken into a dark place, the substance within it will emit a considerable quantity of light.

610. HOMBERG'S PHOSPHORUS possesses the same property as Baldwin's phosphorus, and is prepared in the same manner—only substituting muriate of lime for nitrate of lime.

611. TO MAKE FULMINATING QUICKSILVER.—Dissolve 100 grains of mercury in an ounce and a-half (by measure) of nitric acid, assisting the solution by heat. When the solution is cold, pour it upon two ounces (by measure) of strong alcohol, previously introduced into a small glass retort, and apply a moderate heat till effervescence is excited. (Do not forget that the mercurial solution must be poured upon the alcohol, and not the alcohol upon the solution.) A white fume then begins to undulate on the surface of the liquor, and flows through the neck of the retort; and a white powder is gradually precipitated. As soon as any precipitate ceases to form, quickly pour the contents of the retort upon a filter, well wash the powder with distilled water, and cautiously dry it by a heat not exceeding that of boiling water. The immediate washing of the powder is material, because it is liable to the re-action of the nitric acid; and while any of that acid adheres to it, it is very subject to be decomposed by the action of light. From 100 grains of mercury, about 120 or 130 grains of the fulminating powder are obtained. —*Experiments.*—1. Lay about a grain of the powder on a smooth iron, and strike it with a hammer—it will detonate with violence.—2. The third of a grain of it may be enclosed between the ends of two slips of card pasted together, and the other end of the card marked by a notch. If that end of the card which contains the powder be held over a candle, detonation takes place.—3. Half a grain of the powder may be wrapped up with a hard pea in a piece of tissue paper. This, (which is the *fulminating ball*, or *Waterloo cracker*,) when thrown on the ground forcibly, explodes.—4. The other fulminating curiosities such as *spiders*, &c., are too well known to need describing, and may be easily prepared. NOTE:—This com-

pound is less dangerous than the fulminating compounds of gold and silver, as it never explodes spontaneously ; but yet it cannot be handled with too much caution ; and no student should attempt to make it who is not tolerably expert at chemical processes in general.

SECTION VIII.—MISCELLANEOUS EXPERIMENTS.

612. TO ENGRAVE FIGURES ON GLASS.—Cover one side of a flat piece of glass, after having made it perfectly clean, with bees' wax, and trace figures upon it with a needle, taking care that every stroke cuts completely through the wax. Next, make a border of wax all round the glass, to prevent any liquid when poured on from running off. Now, take some finely-powdered fluuate of lime, (fluor spar ;) strew it evenly over the glass plate, (upon the waxed side ;) and then, gently pour upon it, so as not to displace the powder, as much sulphuric acid, diluted with thrice its weight of water, as is sufficient to cover the powdered fluor spar. Let every thing remain in this state for three hours, then remove the mixture, and clean the glass by washing it with oil of turpentine : the figures which were traced through the wax will be found engraven on the glass ; while the parts which the wax covered, will be uncorroded.—*Rationale.* The fluuate of lime is decomposed by the sulphuric acid, and sulphate of lime is formed. The fluoric acid disengaged in the gaseous state, combines with the water that diluted the sulphuric acid, and forms liquid fluoric acid, by which the glass is corroded.

613. EXHIBITION OF IODINE.—This substance is prepared with great difficulty, and is very expensive. But, at some of the shops where chemical preparations are sold small glass tubes, hermetically sealed, and containing two or three grains of it, may be purchased.—If one of these glass tubes is gently heated, by holding it over a lamp, the iodine is converted into a beautiful violet-coloured vapour.

614. THE CHAMELEON SPIRIT—A LIQUID WHICH IS

BLUE WHEN THE BOTTLE CONTAINING IT IS OPEN, BUT COLOURLESS WHEN THE BOTTLE IS CLOSED.—Put some shreds of clean copper into a small phial, fill it with liquid ammonia, and cork it air-tight. Nothing will take place. Open the bottle, and let it remain for a few hours. The liquid will become blue. Cork the phial. After some time, the liquid will be colourless. Re-open the phial,—the liquid will become blue again.—It will be found necessary, sometimes, to add to the mixture, after it has remained exposed to the air for some time, a little more copper.—*Rationale.* Metallic copper is not acted upon by ammonia; but if the copper is allowed to attract oxygen from the air, it then becomes soluble. But copper oxidised only so much as to render it soluble produces a colourless solution; and it is only when it is highly oxidised that the *blue* solution is produced. Thus, in the above case, the copper, when the phial was left open, acquired enough of oxygen from the atmosphere to produce blue solution. When, however, the phial was closed, the blue colour disappeared—why?—because the remainder of the metallic copper took, in order to render *it* soluble, the super-abundant oxygen from the portion of copper which was dissolved. The blue colour returned, when the bottle was opened, because more oxygen was attracted.

615. SPOONS WHICH MELT IN HOT WATER.—Fuse together in a crucible, eight parts of bismuth, five of lead, and three of tin: these metals will combine, and form an alloy, (of which spoons may be made possessed of the remarkable property of melting in boiling water.)

616. TO MAKE BEAUTIFUL GOLDEN FIGURES UPON SILK.—Dilute a solution of nitro muriate of gold with thrice its quantity of water, and with the solution paint flowers, or shrubs, on a piece of white silk, then expose the silk (kept moist by being wetted with pure water) to a current of hydrogen gas, which is to be forced through a pipe, from the vessel containing it, as directed 434, 435. The hydrogen gas will reduce the oxide of gold,—the colour of the paintings will change from yellow to green—and the

figures will soon shine with all the splendour of the finest gold. NOTE:—Unless the silk is kept wet, the effect will not be produced.

617. TO MAKE BEAUTIFUL SILVERY FIGURES UPON SILK.—Immerse a piece of silk in a diluted solution of nitrate of silver, and expose it while *wet* to a current of hydrogen gas (as described in the preceding experiment) the silver will quickly be reduced; various colours,—blue, purple, red, orange, yellow,—accompany the reduction, and at last, the threads of the silk appear like silver wire.

618. TO MAKE CINDERS, OR LITTLE WICKER-BASKETS, APPEAR AS IF THEY WERE CRYSTALLIZED.—Saturate water *kept boiling*, with alum; then set the solution in a cool place, suspending in it, by a hair, or fine silk thread, a cinder, a sprig of a plant, or any other trifle; as the solution cools, a beautiful crystallization of the salt takes place upon the cinder, &c. which resembles specimens of mineralogical spars.

619. TO MAKE A PIECE OF CHARCOAL APPEAR AS THOUGH IT WERE COATED WITH GOLD. Dilute a saturated solution of nitro-muriate of gold with 5 times its bulk of water, place a thin slip of fresh-burned charcoal into it, and apply heat, gradually increasing which, till the solution gently boils.—The heat will make the charcoal rob the oxide of gold in solution of its oxygen, whereby the metal is rendered insoluble, and is precipitated on the charcoal, in the form of brilliant spangles.

620. TO GIVE A PIECE OF CHARCOAL A RICH COAT OF SILVER.—Lay a crystal of nitrate of silver upon a piece of burning charcoal,—the metallic salt will deflagrate and throw out the most beautiful scintillations that can be imagined. The silver is reduced, and, in the end, produces upon the charcoal an appearance very brilliant.

621. BEAUTIFUL PRECIPITATION OF PURE METALS FROM METALLIC SALTS.—*Process 1.* Let a small piece of phosphorus be scraped to free it from oxide, and dried by immersion in alcohol. Pass a thread through it with a needle, and then suspend it in a solution of nitro-muriate of gold. In a few minutes, the phosphorus will be covered with pure gold.—*Process 2.* Suspend, in like manner, a

bit of phosphorus in a solution of sulphate of copper, for a few hours; the result will be a precipitation of pure copper.—*Process 3.* Let the same be suspended, for a day or two, in a solution of nitrate of silver—a brilliant precipitation of silver will ensue.—*Rationale.* The oxygen which enters into the composition of the metallic salts by forming soluble oxides with the metals, is abstracted by the phosphorus; and the metals, being in its absence insoluble, are precipitated.

622. TO SHOW THE FIXITY OF THE NOBLE METALS.—Expose to a current of oxygen gas, in the manner described, 425, an alloy of silver and lead: as soon as the alloy is in a state of complete fusion, the lead will begin to burn, and in a short time, will be entirely dissipated in a white smoke; the silver being left behind in a state of purity.

623. TO MAKE A BEAUTIFUL GREEN PAINT.—Add, to a hot solution of sulphate of copper, a little solution of carbonate of soda—a beautiful powder (known by the name of French Green) will be precipitated.—The powder is a sub-carbonate of copper.

624. TO SHOW THAT THE ATMOSPHERE CONTAINS WATER EVEN IN THE DRIEST WEATHER.—1. Expose to the open air, for a few days, a spoonful of dry sub-carbonate of potash, or muriate of lime, spread on a saucer: the salt will attract so much moisture from the air, that it will become liquid.—2. Put a given quantity of strong sulphuric acid into a vessel exposed to the air; at the expiration of twenty-four hours, its weight will be found to have increased one-third.

625. TO SHOW THAT THE ATMOSPHERE CONTAINS CARBONIC ACID.—Expose to the open air, in an open vessel, a quantity of transparent lime water; a white crust will soon form on its surface, which on being broken, falls to the bottom of the vessel, and is succeeded by another—this precipitate, upon being examined, proves to be carbonate of lime—therefore, carbonic acid is attracted from the atmosphere by the lime in solution.

626. TO SHOW THAT CARBONIC ACID IS CONTAINED IN AIR RESPIRED FROM THE LUNGS. *Process 1.*—Put into

test glass a little water tinged blue by tincture of cabbage ; then, blow into this water, through a tobacco-pipe, air from the lungs—the blue colour will soon be changed to red. This proves that the air blown from the lungs contains an acid—now, *query*, what acid is it? Let us see,—*Process 2.* Warm the product of *process 1*, the blue colour will be restored—hence the acid is volatile, and must be either carbonic acid or sulphuretted hydrogen. That it is *not* the latter, may be known by its want of smell ; that it is the former, may be proved thus ;—*Process 3.* Blow air, in the manner described above, into lime or barytic water. The transparent solution will be shortly rendered opaque, by the formation of carbonate of lime, or of barytes.

627. CHEMICAL MIRACLE! TWO LIMPID LIQUIDS CONVERTED BY MIXTURE INTO A SOLID MASS.—*Process 1.* If a saturated solution of muriate of lime be mixed with a saturated solution of carbonate of potass, (both transparent liquids), the result is the formation of an opaque and almost solid mass.—*Rationale.* Mutual decomposition of the salts takes place—muriate of potass, and carbonate of lime, are formed ; and the latter absorbs the whole of the water of solution—and thus a degree of solidity is produced.—*Process 2.* Drop sulphuric acid into a saturated solution of muriate of lime ; in this case also an opaque mass is produced.—*Rationale.* The muriate is decomposed, and sulphate of lime (a highly insoluble salt) is formed.—*Process 3.* Pour a saturated solution of caustic potass into a saturated solution of sulphate of magnesia (Epsom salt)—a nearly solid mass is again produced.—*Rationale.* The sulphuric acid leaves the magnesia (which is then precipitated in the form of a white powder) in order to combine with the potass, for which it has a greater affinity. * * * If a little nitric acid be added to the product of *process 1*, the solid mass will be changed to a transparent liquid : the insoluble carbonate of lime being converted into the soluble nitrate.

628. A FLUID PRODUCED BY RUBBING TOGETHER TWO SOLID METALS.—Triturate an amalgam of lead with an amalgam of bismuth—the product will be fluid, like mer-

cury. Fluids are likewise produced when any of the mixtures which follow are triturated: acetate of lead and sulphate of zinc—or, sulphate of soda and nitrate of ammonia—or, sulphate of soda and carbonate of potass.—These salts should be all fresh crystallized.

629. A GREEN COLOURED SOLID PRODUCED BY MIXING A BLUE ONE WITH A WHITE ONE.—Triturate crystallized sulphate of copper with crystallized super-acetate of lead. In this process, acetate of copper (which is of a green colour) is formed.

630. AN ALLOY WHICH MAY BE KEPT IN FUSION UPON A PAPER HELD OVER A CANDLE.—Melt together in a crucible—lead one part, zinc one part, and bismuth one part.

631. METALLIC TREES.—Some of the metals, when precipitated in a particular manner from their solutions, crystallize into the appearance of very beautiful trees or plants, which are usually known by the Latin name of *arbores*. These experiments are not only curious and entertaining, but very simple; and, therefore, we subjoin the processes for working them.

632. FORMATION OF THE SILVER TREE, OR ARBOR DIANAÆ.—Mix one part of a saturated solution of nitrate of silver with twenty parts of distilled water, and pour the mixture upon two parts of mercury in a phial. After some time (the phial being left standing quietly) the branches and a figure of a tree formed of brilliant silver will appear to grow from the mercury, in a very beautiful manner.—*Rationale.* The silver in solution is robbed of its oxygen by the metallic mercury, and is then consequently precipitated.

633. FORMATION OF THE IRON TREE, OR ARBOR MARTIALI.—Add a saturated solution of sub-carbonate of potass to a solution of nitrate of iron. The mixture swells up considerably, then sinks to rest, and metallic branches spring out in a very curious manner on the surface of the glass.

634. FORMATION OF THE LEAD TREE, OR ARBOR PLUMBÆ.—Dissolve two drachms of acetate of lead (sugar of lead) in six ounces of rain water; filter the solution; and

pour it into a clean wide phial. Then, suspend a thin roll or a granule of zinc, by a thread or wire fastened to the cork of the phial, in the middle of the solution, and place the phial where it will not be disturbed. After a few hours, the lead (which is de-oxidized by the zinc) is precipitated on the zinc in the shape of scales or leaves, which have a very brilliant appearance.

635. EASY PRODUCTION OF SILVER SHRUBBERY.—*Process 1.* Bend a piece of copper wire into the shape of a tree, (but *flat*), or any figure you choose, and lay it upon a piece of glass or slate over which you have previously spread a few drops of nitrate of silver. In a few hours, the copper will be covered by brilliant ramifications of silver.—*Process 2.* Drop a little solution of nitrate of silver upon a clean plate of copper; in a short time, a very elegant and pleasing metallic vegetation will be observed to branch out.

636. EASY METHOD OF MAKING BEAUTIFUL GOLDEN FIGURES UPON STEEL.—Add to a saturated solution of nitromuriate of gold, about a fourth part of sulphuric ether; shake the mixture, and then allow it to settle. The ether will take the gold from the acid, and will separate itself from it also and form an upper stratum in the vessel. Carefully pour this ethereal gold into another glass, and immerse in it any steel utensil that is highly polished, then take it out and instantly plunge it into water—when the surface will have acquired a coat of pure gold, the beauty of which may be increased by burnishing.—You may use a pen, and draw figures on razors, &c., and the gold will remain on them, as just described.

637. TO MAKE BRASS.—Put into the bowl of a large tobacco-pipe, a portion of copper filings, mixed with about twice its quantity of finely granulated zinc; cover it with charcoal powder, and press it well together; then, expose it to the action of a clear fire for some time:—the two metals will combine and form brass.

638. TO SILVER COPPER.—Precipitate silver from its nitric solution, by the immersion of polished plates of copper. Take of this silver 20 grains, of super-tartrate of potass 2 drachms, of common salt 2 drachms, and of alum

half a drachm ; mix the whole well together. Take then the article to be silvered, clean it well, and rub some of the mixture, previously a little moistened, upon its surface: the silvered surface may be polished with a piece of soft leather.—The dial-plates of clocks, scales of barometers &c. are all plated thus.

639. TO OBTAIN LARGE AND BEAUTIFUL ARTIFICIAL CRYSTALS.—This operation requires considerable address and much patient attention ; it is as follows : a solution of the salt to be crystallized is to be slowly evaporated to such a consistency that it shall crystallize upon cooling, which may be known by letting a drop of it fall on a plate of glass. When it is in this state, set it by ; and when it is cold, pour into a flat-bottomed vessel the liquid part of the solution off the mass of crystals, which will have formed at the bottom of it. After a few days, solitary crystals will be formed, which will gradually increase in size. Pick out the most regular of these, and put them into another flat-bottomed vessel ; and pour over them a fresh solution of the salt evaporated till it crystallizes on cooling. After this, alter the position of every crystal once a day with a glass rod, so that all the faces of it may be alternately exposed to the liquid ; for the face on which the crystal rests never receives any increment. By this process the crystals gradually increase in size. When they are so large that their forms can be easily distinguished, take the best of them, and put each into a vessel separately ; add a fresh solution of the salt as before directed ; and turn every crystal several times a-day. By this treatment we may obtain them of almost any size we wish. It is necessary to pour off the liquid from the crystals, and add fresh liquid in its place, very frequently ; for the solution, after depositing a certain portion of its salts, becomes weakened, and then attacks the crystals—rounding off their angles in the first place, as an attentive observer may perceive, and infallibly destroying them unless renewed.—The student may endeavour to form regular crystal of *alum* thus, to exercise his dexterity.

640. TO PROVE THAT “ EVERY SALT HAS A CERTAIN DETERMINATE FIGURE WHICH IT ASSUMES UPON CRYSTALL

LIZATION."—*Process 1.* Dissolve in separate portions of boiling water, equal weights of sulphate of copper and nitrate of potass; pour them together while hot, into an evaporating dish, drive off by heat a little of the water, and then suffer the mixture to cool, when the salts will shoot:—the sulphate of copper into *blue* crystals, the nitrate of potass into *white* ones—precisely similar to what they were before they were dissolved.—*Process 2.* Treat, in the same manner, sulphate of iron and muriate of soda: the former salt is separated from the solution (in *green* crystals) by alternate evaporation and cooling; the latter salt can only be obtained (in *white* crystals) by converting all the water into steam.—*Process 3.* Dissolve a quarter of an oz. of each of the following salts in a small quantity of water, in separate glasses,—sulphates of magnesia, iron, copper, soda, alumina and potass (alum),—pour the whole into the Wedgewood's-ware basin, and slowly evaporate a portion of the water. When the liquid has acquired its proper degree of density (the student will be taught by practice to know when this is the case) it must be set by to cool—when every salt will crystallize in its own peculiar manner; having the identical colour, form, taste, and other properties by which it was distinguished, previous to its solution: the mixture of the different crystallized salts having a most pleasing, singular, and beautiful appearance.

641. PLEASING EXAMPLE OF CRYSTALLIZATION BY FUSION.—Take common red litharge, pour over and stir well with it one-third of its weight of strong muriatic acid. Let the mixture become dry, then melt it in a crucible, and pour it into a metallic vessel while in a state of fusion. The mass when cool will be of a beautiful yellow colour, and when broken will exhibit the most regular crystallization that can be conceived.

642. THE MINERAL CHAMELEON.—When one part of black oxide of manganese and three parts of nitrate of potass, both reduced to powder, and mixed together, are exposed in a crucible to a strong heat for about an hour, or, as long as any gas continues to be disengaged, a compound of highly oxidised manganese and potass, possess-

ed of some very curious properties, is obtained.—*Experiment 1.*—A few grains of this compound put into a wine-glassful of water produces a *green* colour; an increase of the quantity changes the colour to a *blue*; more still to a *purple*; and a yet further increase produces a beautiful *deep purple*.—*Experiment 2.* Put equal quantities of this substance into two separate wine-glasses, and add to the one *hot*, and to the other *cold*, water. The hot solution will be of a beautiful green colour; the cold one of a deep purple. By using more glasses, and water more or less in quantity, and at different temperatures, a great variety of colours will be produced in this way from the same substance.

643. EXAMPLES OF SUBLIMATION.—*Process 1.* Put a quantity of roll sulphur, previously reduced to a powder, into the mattrass *a*, fig. 26, then fasten upon *a* the head *b*, and close by a cork the neck *c*. Apply now a gentle heat, and the sulphur will be converted into a vapour, which will be condensed in the head of the vessel in a state of tolerable purity; its purification may be completed by washing it with distilled water.—*Process 2.* Spread a small quantity of grossly-powdered gum benzoin on the bottom of a Wedgewood's-ware basin, invert over it a glass tumbler, and apply to it a gentle heat by means of the lamp furnace: the gum will melt, and dense fumes will immediately rise from it and deposit themselves in beautiful silky crystals (which are benzoic acid) on the sides of the glass.—*Process 3.* Take a large glass jar, containing at its top a sprig of rosemary or some such shrub, and invert it over a flat thick piece of heated iron on which coarse powder of gum benzoin has just been spread—then, the benzoic acid which rises, as in the preceding experiment, will be deposited on the branches of the shrub, producing a singular and beautiful representation of hoar frost.

644. TO OBTAIN PURE GOLD.—Add to a solution of nitro-muriate of gold, a solution of green sulphate of iron, as long as precipitation ensues: the precipitate is to be well-washed on a filter, (first with diluted muriatic acid,

and then with pure water), and dried—it is then pure gold.

645. TO FORM SULPHURET OF IRON,—Bring a bar of iron to a white heat, hold it over a pan of water, and touch it with a roll of sulphur. A portion of the metal will instantly combine with the sulphur, and fall, in drops, into the water.—Sulphuret of iron may also be formed, by melting equal portions of iron filings and sulphur in a crucible.

646. PURE LIME IS OBTAINED by exposing oyster-shells, marble, or chalk, to the action of a clear kitchen fire, for about an hour, the heat drives off the carbonic acid, and the substance remaining, is pure lime, which must, while warm, be put into a bottle and be well-secured from the air.

647. CURIOUS MODE OF SILVERING IVORY.—Immerse a small slip of ivory in a weak solution of nitrate of silver; and let it remain till the solution has given to it a deep yellow colour; then take it out and immerse it in a tumbler of clear water; and expose it in the water to the rays of the sun. After it has been thus exposed for about three hours, the ivory acquires a black colour; but, the black surface, on being rubbed, soon becomes changed to a brilliant silvery one.

648. SYMPATHETIC INKS.—Among the amusing experiments of the science of chemistry, the exhibition of sympathetic ink holds a distinguished place. These liquids, when written with, leave no visible traces behind: the writing only becomes visible when some known *re-agent* is applied to it.—We shall here mention a few of these “inks” out of the great number which but a slight acquaintance with chemistry will suggest to the student.—NOTE.—The sympathetic inks may be laid on paper either with a camel hair pencil or a common pen; but, whichever is used, it is necessary that the instrument be perfectly clean—the presence of the smallest conceivable quantity of any foreign body will go nigh to spoil the effect. The best thing to employ is a clean fresh-cut pen.

Process 1.—Write with weak tincture of galls—the characters will be invisible. Moisten the paper with a

feather dipped in a weak solution of sulphate of iron—the writing will become *black*.—For an explanation of this phenomenon, see 389.

649. *Process 2*.—Write with a weak solution of prussiate of potass—the letters will be invisible. Moisten the paper, as in the preceding experiment, with a weak solution of sulphate of iron—the writing will assume a fine *blue* colour.—*Rationale*. Prussiate of iron is formed, see 167.

650. *Process 3*.—Wash paper with a solution of sulphate of iron, and suffer it to dry: when written upon this paper, solution of prussiate of potass produces *blue* letters, and tincture of galls *black* ones; but upon common paper, they make colourless marks.

651. *Process 4*.—Most acids, diluted, and written with, leave marks which are invisible till the paper is heated, when they become *black*; the heat concentrating the weak acid, and enabling it to char the paper.

652. *Process 5*.—Write with a weak solution of nitrate of silver, and let the writing dry *in the dark*—it will be invisible; fold up the paper so as to keep the writing in the dark—it will continue invisible; but, expose the writing to the light of the sun—it will become *black*.—*Rationale*. The nitrate of silver has the property of being decomposed by light; a black colour being acquired by the metallic oxide.

653. *Process 6*.—Characters written with a solution of equal parts of sulphate of copper and muriate of ammonia, have a *yellow* colour when heated; but, are invisible when cold.

654. *Process 7*.—Write with a weak solution of muriate of copper—the writing is invisible when cold but *yellow* when heated.

655. *Process 8*.—Write with a weak solution of nitromuriate of gold, and dry the writing in the dark—it will be invisible. Moisten the paper, by means of a feather or bit of sponge, with a solution of muriate of tin—the writing will then assume a *purple* colour.

656. *Process 9*.—Write with a solution of nitrate of bismuth—the writing will be invisible. Immerse the paper in water—the characters will then be legible.—*Rationale*.

The water decomposes the salt, and causes the white oxide of bismuth to be precipitated.

657. *Process 10.*—Expose a paper upon which you have written with nitrate of bismuth, to the vapour of water impregnated with sulphuretted hydrogen—the writing will become *black*; It being the property of bismuth to be thus affected by sulphuretted hydrogen.

658. *Process 11.*—Let a paper upon which you have written with nitrate of bismuth be moistened with solution of prussiate of potass—the writing will assume a *beautiful yellow* colour: a prussiate of bismuth being formed.

659. *Process 12.*—Write with a solution of sulphate of copper—no writing will be visible. Wash the paper with solution of prussiate of potass—the writing will then get a *reddish-brown* colour: prussiate of copper being formed.

660. *Process 13.*—Write with a solution of super-acetate of lead—the writing will be invisible. Hold the paper over a saucer containing liquid sulphuretted hydrogen—the writing will become, first black, and then glittering like silver.—*Rationale.* The metallic salt is decomposed by the sulphuretted hydrogen, which robs it of its oxygen, and, consequently, the lead is gradually reduced to its metallic state.

661. *Process 14.*—Upon a fire-screen let there be drawn a representation of winter, with trees destitute of foliage, and ground covered with snow. Let however, every part of the picture which, if the scene represented summer, would be green, be covered with the sympathetic ink, for making which directions are given at 503. Draw, for instance, leaves on the trees, and grass on the ground. These marks will not be visible; the picture will still bear the aspect of winter. But, let the fire-screen be placed near the fire, *then* the view will exhibit all the verdure of the summer. When allowed to cool, the verdure disappears; but it may be again revived, by the same means as before, and that as often as desired—provided, that the paper be not heated beyond a certain point: for, if heated too much, the ink will assume a permanent brown colour.

662. *Process 15.*—Write with a diluted solution of acetate of cobalt (made by dissolving oxide of cobalt in acetic

acid):—the writing will be invisible till the paper is heated, when it becomes *blue*.

663. CURIOUS EFFECT OF COLOUR WITH RESPECT TO THE ABSORPTION AND REFLECTION OF HEAT.—Take 6 pieces of tin plate, each one inch square—paint (on one side only) 1 black, 1 blue, 1 green, 1 red, 1 yellow, and let the others remain bright. On the backs of these pieces of metal, lay a thin coat of cerate, composed of oil and wax, which easily melts. Place them, thus prepared, on a board painted white, and expose the coloured surfaces of the metal to the rays of the sun. The heat will melt the cerate, and (if the board rests in a proper position) it will run down—from the black, the blue, the green, the red, and the yellow, pieces—while the bright one remains nearly unaffected.—This experiment shows that light-coloured surfaces reflect heat and dark ones absorb it. Hence we learn, that, if we wish to be warmed by the sun, we should clothe ourselves in black; and that, in the summer, light coloured dresses are preferable.

664. VERY SINGULAR PROPERTY OF CAMPHOR.—If a small piece of camphor is placed on the surface of water contained in a basin, it immediately begins to move round and round with considerable rapidity. It is necessary that the water be pure, for, if dust or grease be present, the camphor will not move, or, if, while its rotatory motion is proceeding, a single drop of grease be let fall into the water, the camphor will instantly start away from it to the sides of the basin, and its motion will be put a stop to.—Nothing satisfactory has yet been advanced in explanation of this phenomenon.

665. TO MAKE PEWTER.—Melt, in a crucible, one part of lead, or bismuth, with about twenty parts of tin.

666. AN EXAMPLE OF THE MODE OF ANALYZING A METALLIC ALLOY.—Suppose the alloy to contain silver, zinc, lead, and bismuth.—*Process 1.* Let it be dissolved with the aid of heat, in an excess of nitric acid.—*2.* Evaporate the solution to dryness.—*3.* Pour water on the residuum which, of course, contains *nitrates* of the different metals—solutions of nitrate of silver, zinc, and lead, will be obtained; the nitrate of bismuth will be decomposed by the

water, and oxide of bismuth (for every 100 grains of which reckon 90 grains of metallic bismuth) will be left at the bottom.—4. Into a hot and pretty dilute solution of the remaining salts, pour muriatic acid—muriate of silver (containing 83 grains of metal in every 100 grains of the salt) will be precipitated.—5. Add to the solution, having separated the product of the last process by filtration, sulphate of soda—this will throw down sulphate of lead (which contains 69 grains *per cent.* of the metal).—6. Finally, add carbonate of potass to precipitate the zinc. NOTE:—the papers through which the mixture is filtered, are to be carefully weighed before they are used; and are to be weighed again, when perfectly dry, after the process, with the precipitates upon them. The number of grains shows the weight of the precipitates.

667. EXHIBITION OF THE EFFECTS PRODUCED BY GALVANIC ACTION.—*Process 1.* Place a piece of the metal zinc under the tongue, and a piece of copper (as a penny) upon the tongue, and bring the extremities of the metals into contact—the result will be the immediate perception of a curious *taste*, something resembling the taste of copperas. The metals have not this taste when not in contact; and, therefore, this experiment clearly shows the influence of excited galvanic electricity upon the organ of taste.

668. *Process 2.* Moisten the under lip; lay upon it a shilling, with the edge of it *beneath* the tongue; lay *upon* the tongue a piece of zinc, and bring the edges of the pieces of metal into contact—instantly, the shilling will feel so intolerably hot, that you will be obliged to let it fall from your mouth. You will then not hesitate to say—galvanic electricity has an influence over the sense of feeling.

669. *Process 3.* Place a piece of zinc under the tongue, and put the small end of a silver tea-spoon in the mouth, between the teeth and the under-lip, passing it up as high as possible, and pressing the extremity of it against the gums; then, let the lower part of the spoon touch the edge of the piece of zinc—at the moment of contact, a flash, like lightning, appears to the eye, which is renewed as often as you let the two metals touch each other. If

you shut your eyes when you perform this experiment, you still see the light; and if you look sternly at a candle, or other light, still you see it. Does not this afford ocular demonstration, that galvanic electricity affects the organ of sight?

670. *Process 4.*—A Voltaic Pile, constructed as directed, 331, affords a constant current of the galvanic fluid, for many hours, through any conducting body which may be employed to connect its two ends. If the hands be moistened, and one applied to each end of the pile, or if a rod of metal be held in each hand, and a communication be made by means of them, a shock will be received; the intensity of which is according to the number of pieces that compose the pile; thus, zinc, copper, and cloth, repeated 20 times, forms a pile that gives a shock perceptible to the wrists, and a pile five times as high, gives a tremulous but severe and continued sensation extending to the shoulders.

671. *Process 5.*—When a wire communicating with the top of a small pile is held between the teeth, so as to rest upon the tongue (the hands at the same time touching the bottom of the pile), the lips, and the tongue, will become convulsed, a flash will appear to the eye, and a very pungent taste will be perceived in the mouth.

672. *Process 6.*—Put distilled water into a small glass tube (of a diameter of about one-seventh of an inch), and connect the tube, by means of two copper wires, to the Voltaic pile, in the manner described, 331. (The pile, for this experiment, must be formed of at least 50 or 60 sets of plates.) After some time, the wire connected with the zinc or positive end of the pile, will be oxidised at the end in the tube; while from the end of the other (the negative) wire in the tube, a stream of small bubbles of gas will arise. This gas is hydrogen; as may be proved by pulling out the cork, and instantly applying an ignited body to the tube, by which the gas will be inflamed. As the gas that is formed in the tube occupies more space than the water that is decomposed, it is necessary to make room in it by allowing some of the water to escape; this is done by cutting a small slit in the cork fastened in the bottom

of the tube.—If two wires of gold or platinum (which are not oxidable) be used instead of the copper wires, a stream of gas then issues from each; and collects at the top of the tube; forming a mixture of hydrogen and oxygen gas, which explodes on the approach of an ignited body.—The gases may be obtained separately, by passing each of the gold wires down a leg of a glass syphon, making them nearly meet at the bottom, and closing the openings at top by means of corks as before. The gas that will be formed will rise up the two legs of the syphon—the hydrogen up the leg containing the negative wire—the oxygen up the leg containing the positive wire—and it will be seen that the gases are formed from the decomposition of the water precisely in the proportions that are required to compose it, namely two volumes of hydrogen to one of oxygen. In this case, the water that must be suffered to escape to make room in the tube, must pass through a small hole ground in the bended part of the syphon, near where the two wires are opposed to each other.

673. *Process 7.* Fill the tube *c*, fig. 40, with a solution of acetate of lead, and let the galvanic influence be passed through the tube by copper wires, in the manner described in the preceding process: no gas will be perceived, for the hydrogen arising from the decomposition of the water reduces the oxide of lead held in solution by the acetic acid, and uniting to its oxygen, re-forms water; the lead, in a brilliant metallic state, being precipitated on the negative wire, first in the form of beautiful needles, and then like fern.

674. *Process 8.* Lay a live flounder upon a wetted plate of copper, and lay on the top of it a piece of zinc; then, by means of a bended wire, make a communication between the pieces of metal placed on each side of the fish; instantly, the animal (though previously lying perfectly quiet) will be violently convulsed.

675. *Process 9.* If a piece of zinc and a piece of silver have each one extremity immersed in the same vessel, containing sulphuric or muriatic acid largely diluted with water, the zinc is dissolved and yields hydrogen gas by de-

composition of the water; the silver not being acted upon has no power of decomposing the water; but whenever the zinc and silver are made to touch, or any metallic communication is made between them, hydrogen gas is always formed at the surface of the silver.

676. *Process 10.*—Experiments analogous to the former and equally simple, may also be made with metallic solutions. Take a solution of sulphate of copper; immerse a piece of silver in it—it will come out unchanged; immerse a piece of polished iron—it will receive a coat of copper (see 538): bring the iron and silver into contact, and immerse them in the solution together—then, both will be coated with copper.

677. SKETCH OF THE SCIENCE OF GALVANISM—explanatory of the preceding experiments.—1. It was accidentally discovered by Galvani, an Italian, that if the cranial nerve and the muscles of the leg of a recently-killed frog be laid bare, and we place a piece of zinc upon the nerve, and a piece of copper upon the muscle, and make the two metals, while so placed, touch each other, the leg of the frog is immediately thrown into violent convulsions. 2. Volta, a friend of Galvani's, invented the *pile* we have described; and showed that shocks like those produced by electricity could be received from it.—3. Shortly after it was found that water, and many other substances, could be decomposed by the galvanic energy.—4. It was next discovered, that when compound bodies are decomposed, oxygen and acids accumulate round the positive pole, while hydrogen, alkalies, earths, and metals, accumulate round the negative pole. Hence it was deduced, that chemical affinity is identical with electrical attraction, and, that the reason that oxygen combines with hydrogen, and acid with alkalies, earths, and metals, is that they are in opposite states of electricity.—5. Galvanic electricity, when properly excited and directed, is sufficiently powerful to decompose any compound whatever. And even pure gold may be burned by being made to form a part of a galvanic circuit.

678. TO OBTAIN METALLIC ZINC FROM AN ORE. Rub together in a mortar 1 ounce of lapis calaminarius

(which is an oxy-carbonate of zinc mixed with earths, &c.), and rather more than $\frac{1}{2}$ the quantity of powdered charcoal, mixed with a little quick-lime; press a portion of this mixture into the bowl of a large tobacco-pipe, and submit it to the action of a strong clear fire; after some time, the zinc will be reduced, and will be found at the bottom of the vessel.

679. TO ANALYSE GALENA, AN ORE OF LEAD.—A very common ore of lead is *galena*, which is a compound of lead and sulphur. It is of a blue colour, has a brilliant metallic lustre, and is easily broken. Before the blow-pipe it decrepitates, melts with a sulphureous smell, and yields a globule of lead.—Let a quantity of this mineral (say 200 grains) be pounded, and put into nitric acid, diluted with an equal weight of water. Nitrate of lead will be formed, and the sulphur will remain undissolved. Filter the solution, and crystallize the salt by evaporation. Or, add to the solution, concentrated sulphuric acid: a white precipitate of sulphate of lead will be thrown down: weigh this, and from the weight deduct 70 per cent.; the remainder shows the quantity of lead.

680. FLUXES are substances employed to assist the fusion of minerals. *Crude flux* is a mixture of nitre and tartar, which is put into the crucible with the mineral intended to be fused. *White flux* is formed by projecting a mixture of equal parts of nitre and tartar, by moderate portions at a time, into an ignited crucible. It is potass in a state of tolerable purity. *Black flux* differs from white flux (both being made in the same manner) in the proportion of its ingredient. In this, the weight of the tartar is double that of the nitre. *Microcosmic Salt* is procured, by mixing equal part of phosphate of soda and phosphate of ammonia in solution, and then crystallizing. A slight excess of ammonia is useful.

681. LUTES, are substances applied to the junctures of vessels used in distillation, to prevent the escape of vapours. In fig. 20, the adopter *b* is represented as *luted* to the necks of the retort and receiver—*a* and *c*. When vapours of watery liquors, and such others as are not corrosive, are to be contained, it is sufficient to surround the joining of the receiver to the retort with slips of wet blad-

der, or of linen, or paper, covered with flower paste, mucilage of gum-arabic. When more penetrating and dissolving vapours are to be contained, a lute is to be employed of quicklime slacked in the air, and beaten into a liquid paste with white of eggs. This must be applied in strips of linen; it is very convenient, as it easily dries and becomes firm. This lute is very useful for joining broken china-ware.—For containing the vapour of acids or highly-corrosive substances, the *fat-lute* is made use of. This is formed by beating perfectly dry and finely-sifted tobacco-pipe clay, with painters' drying oil, in a mortar to such a consistence that it may be moulded by the hand. To use it, it is rolled into cylinders of a convenient size which are applied, by flattening them, to the joinings of the vessels, which must be quite dry, as the least moisture prevents the lute from adhering. The lute when applied is to be covered with slips of linen spread with the second mentioned lute; which slips are to be fastened with pack-thread.

682. TO PROCURE BORACIC ACID.—Dissolve borate of soda (borax) in hot water, and filter the solution; then add sulphuric acid by little and little, till the liquor has a sensibly acid taste. Lay it aside to cool, and a great number of small laminated crystals (scales) will be gradually formed. These are the boracic acid. They are to be purified by washing with cold water, which carries off any extraneous soluble body, but leaves the acid, which is very sparingly soluble, almost untouched. When the crystals have been washed, they are to be drained upon brown paper.

683. EASY METHOD OF BREAKING GLASS TO ANY REQUIRED FIGURE.—Make a small notch, by means of a file, on the edge of a piece of glass; then, make the end of a tobacco-pipe, or of a rod of iron of the same size red-hot in the fire: apply the hot iron to the notch, and draw it slowly along the surface of the glass in any direction you please; a crack will be made in the glass, and will follow the direction of the iron. Cylindrical glass vessels, such as flasks, may be cut in the middle, by wrapping round them a worsted thread dipped in spirit of turpentine, and setting it on fire when fastened on the glass.

APPENDIX.

TABLE OF CHEMICAL DECOMPOSITIONS.

ALKALIES.	BARYTES AND STRONTIA.	LIME.	MAGNESIA.
Sulphuric acid, Nitric, Muriatic, Oxalic, Tartaric, Carbonic.	Sulphuric acid, Oxalic, Nitric, Muriatic, Tartaric, Carbonic.	Oxalic acid, Sulphuric, Tartaric, Nitric, Muriatic, Carbonic.	Oxalic acid, Sulphuric, Nitric, Muriatic, Tartaric, Carbonic.
SULPHURIC ACID.	NITRIC ACID.	MURIATIC ACID.	CARBONIC ACID.
Barytes, Strontia, Potass, Soda, Lime, Magnesia, Ammonia.	Barytes, Potass, Soda, Strontia, Lime, Magnesia, Ammonia.	Barytes, Potass, Soda, Strontia, Lime, Ammonia, Magnesia.	Barytes, Strontia, Lime, Potass, Soda, Magnesia, Ammonia.
OXALIC AND TARTARIC ACIDS.	ACIDS WITH OXYGEN.	OXIDE OF SILVER.	OXIDE OF IRON.
Lime, Barytes, Strontia, Magnesia, Potass, Soda, Ammonia.	Zinc, Iron, Tin, Lead, Copper, Mercury, Silver,	Galic acid, Muriatic, Oxalic, Sulphuric, Nitric, Prussic, Carbonic.	Galic acid, Sulphuric, Muriatic, Nitric, Acetic, Prussic, Carbonic.

See 514.

EXPLANATION: This table shows what would result from the mutual action of two chemical agents, if placed in contact. Thus the column headed *sulphuric acid* is designed to show, that barytes has a stronger affinity than any other body for that acid, and will detach it from any of the succeeding substances in the list; and that strontia will separate it from potass; and so on of the rest. This may be verified by actual experiment.—When, therefore, any question regarding decomposition occurs, if, for instance, we wish to know, whether potass will decompose sulphate of magnesia, instead of putting the bodies together to try the experiment, we refer to the table, in order to save trouble; and there, under the head *sulphuric acid*, we find it stated, that potass has a stronger affinity for that acid than magnesia, and, therefore, that it *will* decompose the above mentioned salt.

TABLE OF FREEZING MIXTURES.

Mixtures.		Thermometer sinks,
Muriate of Ammonia,	5 parts	} From 50° to 10°
Nitrate of Potass,	- 5	
Water, - - -	16	
Nitrate of Ammonia,	1 part	} From 50° to 4°
Water, - - -	1	
Sulphate of Soda, -	5 parts	} From 58° to 30°
Diluted Sulphuric Acid,	4	
Snow, - - -	1 part	} From 32° to 0°
Common Salt, - -	1	
Muriate of Lime, -	3 parts	} From 32° to 50°
Snow, - - -	2	
Snow, - - -	2 parts	} From 10° to 50°
Diluted Sulphuric Acid,	1	
Diluted Nitric Acid,	1	
Snow, or pounded Ice,	12 parts	} From 18° to 25°
Common Salt, -	5	
Nitrate of Ammonia, -	5	
Muriate of Lime, -	3 parts	} From 40° to 73°
Snow, - - -	1	
Diluted Sulphuric Acid,	10 parts	} From 68° to 91°
Snow, - - -	8	

NOTE:—The principle upon which the action of *freezing mixtures* depends is mentioned at 382. In order to produce the effects described in the above table, several things must be attended to. The salts employed must be fresh crystallized, and should contain as much water of crystallization as possible; they should also be quite dry, and reduced to fine powder immediately before being used. The vessel in which the freezing mixture is made should be very thin; and the substance it is made of should be a good conductor of caloric. Tin vessels answer very well; but, when it is intended to put acid mixtures into them, they must be coated with wax. The materials should be mixed as quickly as possible, and must be in sufficient quantity to fill the vessel.—When great cold is to be produced, the materials for the mixture must first, separately, be reduced to the temperature marked in the table, by being placed in some of the other freezing mixtures; and they are then to be mixed together in a vessel placed in a similar freezing mixture.

Mercury freezes at -39° , so that, by the aid of some of the mixtures described in the above table, put together in the manner we have just directed, the interesting experiment of the congelation of this singular metal may be performed. The readiest method of exposing the mercury to the action of the mixture, is to place a little of it, on a watch glass, or concave piece of tin, supported in the middle of a pretty large mass of the freezing mixture.

For procuring moderate refrigeration, the most convenient mixture is the *first* in the above table; because, the water of solution being afterwards removed by evaporation, the pulverized salts are equally efficacious as at first.

TABLE OF SPECIFIC GRAVITIES.

METALS.		LIQUIDS.	
Water (the Standard),	1.00	Water, - - - -	1.00
Platinum, - - -	21.50	Sulphuric Acid, -	1.90
Gold, - - - -	19.30	Nitric Acid, - -	1.50
Standard Gold, -	17.50	Carburet of sulphur,	1.22
Mercury, - - - -	13.50	Muriatic Acid, -	1.22
Lead, - - - -	11.35	Sea Water, - - -	1.00
Silver, - - - -	10.50	Lintseed Oil, - -	0.90
Bismuth, - - - -	9.80	Olive Oil, - - -	0.90
Copper, - - - -	8.90	Ammonia, - - -	0.88
Brass, - - - -	8.50	Oil of Turpentine, -	0.88
Arsenic, - - - -	8.35	Alcohol (Com.), -	0.85
Nickel, - - - -	7.55	Alcohol (Pure), -	0.79
Cobalt, - - - -	8.00	Naphtha, - - -	0.75
Iron, - - - -	7.78	Sulphuric Ether, -	0.75
Tin, - - - -	7.30	SOLIDS.	
Zinc, - - - -	7.00	Water (the Standard),	1.00
Manganese, - - -	6.85	Barytes, - - - -	4.20
Antimony, - - -	6.70	Strontia, - - - -	3.70
Sodium, - - - -	0.97	Carbon (Diamond),	3.50
Potassium, - - -	6.86	Flint Glass, - - -	3.29
GASES.		Fluor Spar, - - -	3.18
Air (the Standard),	1.00	Carbon. of lime, as } fr. 2.70	to 1.38
Iodine (Vapour of),	8.77	marble, chalk, &c. }	
Chlorine, - - - -	2.50	Rock Crystal, - -	2.65
Nitrous Oxide, -	1.53	Silica, - - - -	2.60
Muriatic Acid, -	1.29	Potass, - - - -	2.50
Sulph. Hydrogen,	1.19	Lime, - - - -	2.30
Oxygen, - - - -	1.12	Magnesia, - - -	2.30
Nitric Oxide, - -	1.04	Alumina, - - -	2.00
Carb. Hydrogen, -	0.99	Sulphur, - - -	1.99
Carbonic Oxide, -	0.98	Phosphorus, - -	1.77
Nitrogen, - - -	0.97	Camphor, - - -	0.99
Phos. Hydrogen, -	0.89	Oak (Dry), - - -	0.92
Water (Vapour of),	0.62	Pumice-Stone, -	0.91
Ammonia, - - -	0.60	Fir (Dry), - - -	0.55
Hydrogen, - - -	0.07	Cork, - - - -	0.24

CHEMICAL EQUIVALENTS.

At paragraph 92 the student is referred to the Appendix for a continuation of the subject of *definite proportions*. All that seems necessary to be done, is to give some account of Dr. Wollaston's admirable *Scale of Chemical Equivalents*; "an instrument which," as Dr. Ure observes, "has contributed more to facilitate the general study and practice of chemistry than any other invention of man." It is of so much advantage to the practical chemist, so simple and easily used, and so cheap, (the price is 4s.) that it cannot fail to come into general use. It answers at once a great many important questions, which perpetually occur to the practical chemist, and sometimes cannot otherwise be answered than by a long arithmetical calculation.

Dr. Wollaston's scale is a list of simple chemical bodies and chemical compounds, arranged on one or the other side of a scale of numbers, in the order of their relative weights, and at such distances from each other, according to their weights, that the series of numbers placed on a sliding scale can at pleasure be moved, so that any number expressing the weight of a compound, may be brought to correspond with the place of that compound in the adjacent column. The arrangement is then such, that the weight of any ingredient in its composition, of any re-agent to be employed, or precipitate that might be obtained in its analysis, will be found opposite to the point at which its name is placed.

In order to show more clearly the use of this scale, the table annexed exhibits a portion of the scale with two different situations of the slider, or, more strictly speaking, the table shows two sets of cases *selected* from the scale.

First view of the Scale.		
	38	Carbonic acid
	46	Muriatic acid, <i>dry</i>
Lime	48	
Soda	53	
	86	Carbonate of Lime
Carbonate of Soda	91	
	94	Muriate of Lime, <i>dry</i>
Muriate of Soda	99	
Second view of the Scale.		
	24	Carbonic acid
	29	Muriatic acid, <i>dry</i>
Lime	30	
Soda	33	
	54	Carbonate of Lime
Carbonate of Soda	57	
	59	Muriate of Lime, <i>dry</i>
Muriate of Soda	62	


In the *first view* all the following questions are answered. In 94 parts of *dry* (that is *not* crystallized) muriate of lime, how many parts of lime and how many parts of muriatic acid are there?—48 lime, 46 acid. If I wish to decompose this salt by means of soda, how much of the latter will be required?—53 parts, or 91 parts of carbonate of soda, which will leave 38 parts of carbonic acid. Then how many parts of muriate of soda will result? and how many parts of lime will be precipitated? 99 parts of muriate of soda, and 48 parts of lime. Another case: if I wish to form muriate of lime from carbonate of lime (chalk) how many parts of muriatic acid, will 86 parts of carbonate of lime require for its decomposition? 46, which will give 94 parts of muriate of lime, and leave 38 parts of free carbonic acid. These are but a few of the queries capable of being solved by this small table, but it is needless to enumerate any more.


In the *second view of the scale*, questions similar to the above are answered, with the difference that the quantities

of the bodies are not the same—this is owing to the sliders being placed at a different height; the same relationship exists between the numbers as before. In Dr. Wollaston's scale, as the slider of numbers can be set at any height, an infinite number of questions can be answered respecting any quantity of every article in the list; and in this list is to be found every important known chemical body. Now, surely, the accurate and immediate solution of so many important practical problems, is an incalculable benefit conferred on the chemist. The *Young Student* will obtain from the scale what will be of great use to him, namely, a knowledge of the proportions of the constituents of most compound bodies; and he will finally be satisfied, after a few experiments, of the truth of the doctrine laid down in paragraph 89, that "bodies enter into combination in *definite* proportions."

the body of the paper. The paper is divided into two parts, the first part being devoted to a discussion of the general principles of the theory, and the second part to a discussion of the particular case of the theory. In the first part, the author discusses the general principles of the theory, and in the second part, he discusses the particular case of the theory. The author's discussion is based on the assumption that the theory is correct, and he shows that the theory is correct by showing that it is consistent with the experimental results. The author's discussion is based on the assumption that the theory is correct, and he shows that the theory is correct by showing that it is consistent with the experimental results. The author's discussion is based on the assumption that the theory is correct, and he shows that the theory is correct by showing that it is consistent with the experimental results.

Index and Glossary.



 *The Figures refer to the Paragraphs of the Work.*

-
- ACETATES, 111
Acetic acid, 111, 238, 245
Acidifiable. Capable of being made into an acid. Sulphur is an example of an acidifiable base.
Acids, 87
———, properties of, 97
———, classification of, 98
———, nomenclature of, 99
Adopter, 271, 312
Affinity, Chemical. See Chemical Affinity.
Agent, Chemical. Any substance capable of producing chemical action.
Aggregation, attraction of, 76
Agriculture, chemistry applied to, 557
Air, (atmospheric,) *Note to*, 18, 95; to compose, 455; to decompose, 452
———, contains water and carbonic acid, 624, 625
Air-Funnel, 285
Air-Holder, 287, 352
Aix-la-Chapelle mineral water, 143
Alabaster, 189
Albumen, 249, 251
Alcohol, 242, 243
———, ready method of ascertaining the quantity of, in wine, &c. 556
Alcohol of sulphur, 147
Alembic, 271, 318
Alkahest, 60
Alkali, fixed, 118
———, mineral, 119
———, vegetable, 118
———, volatile, 121
Alkalies, *Note to*, 29, 88, 117, 128
———, as tests, 511
Alloys, 28, 628, 630, 666
———, mode of analysing, 666
Alum, 127, 197
Alumina, 69, 127
——— and potass, sulphate of, 197
Aluminum, 29, 69
Amalgams, 34
Amber, 238
Ammonia, 121
———, carbonate of, 179
———, muriate of, 178, 486
———, nitrate of, 177, 484
———, oxalate of, 510
——— liquid, 523
Ammoniacal gas, 467
Analysis defined, 11
Animal kingdom, 249
Antimony, 29, 43
Apparatus, 254—359
Aqua-fortis, 106
Aqua-regia, 199
Arbor Dianæ, 632

- Arbor Martis, 633
 ——— Plumbum, 634
 Argil, 127
 Argillaceous earths, 127
 Arsenic, 29, 47
 Arsenic acid, 47
 Arsenious acid, 47
 Assafoetida, 231
 Atomic theory, 89—92
 Attraction, 74
 Aurora Borealis, 437
 Azote, 23
 BALDWIN'S Phosphorus, 609
 Balloons (Miniature), 431, 432, 433
 Balsams, 232, 238
 Barilla, 174
 Barium, 29, 64
 Barytes, 64, 123
 ———, carbonate of, 181
 ———, muriate of, 183, 487, 517
 ———, nitrate of, 182, 488
 ———, sulphate of, 180, 489
 Barytic water, 513
 Base. A term applied to denote the *oxide* which is combined with an acid, to form a salt:
 Bath (Sand). A vessel filled with sand, and placed over a fire. It is employed to give heat to matrasses, retorts, and other vessels, which it would not be prudent to expose to a naked fire.
 Bath (Water). A vessel of boiling water, in which other vessels to be heated are placed; or on the cover of which, certain precipitates, &c. are placed to dry. The boiling water bath gives a constant heat of 212°.
 Beer, 240
 Bell-metal, 35
 Benzoates, 113
 Benzoic acid, 113, 238
 Bismuth, 29, 42
 ———, nitrate of, 42, 218, 501
 ———, nitro-muriate of, 218
 Bitumen. A generic term applied to a variety of inflammable mineral substances. The following are some of the most important of this class of bodies: Naphtha, Petroleum, Asphaltum, Jet, Amber, Coal.
 Black-lead, 149
 Bleaching, 469, 473, 564
 Bleaching-Powder, 138
 Blow-pipe, 279, 350
 ——— spoon, 351
 Bolognian phosphorus, 1800
 Bolt-head, 344
 Bones, 27, 193
 Boracic acid, 105, 682
 Borates, 105
 Borax, 25, 105, 172
 Boron, 17, 25
 Borurets, 140
 Brass, 35, 637
 Brimstone, 26
 Bronze, 35
 CABBAGE, tincture of, 5066
 Cadmium, 29, 40
 Calcareous. Of a chalky nature, containing lime.
 Calcareous spar, 125
 Calcination. The act of burning certain substances, in a particular manner. Chalk is calcined when it is made into lime.

- Lead is calcined, when it is made into dross, by being kept melted, and exposed to the air. The *dross* of Lead is an *oxide* of that metal; it formerly was termed the *calx* of Lead.
- Calcium, 29, 63
- Calico-Printing, 561, 562, 563
- Calomel, 34
- Caloric, a simple body, 16
 ———, nature of, 81—84
 ———, experiments on, 360 to 386
 ———, expands bodies, 362
 ———, conductors of, 371
- Camphor, 229, 238, 664
- Camphoric acid, 238
- Canton's phosphorus, 608
- Caoutchouc, 233
- Carbon, 17, 24
- Carbonates, 104
- Carbonic acid, 24, 104, 416, 442—450, 521
- Carbonic acid contained in air respired from the lungs, 626
- Carbonic acid gas, directions for obtaining, and performing experiments with, 442—450
- Carbonic oxide, 136
- Carburet of iron, 148, 149
 ——— of sulphur, 147
- Cassius, precipitate of, 536
- Causticity. A property which some substances possess, of burning or corroding animal bodies to which they are applied.
- Cerium, 29, 58
- Chalk, 125, 188
- Chalybeate mineral waters, 209
- Chameleon (mineral), 642
 ——— spirit, 614
- Charcoal, 24
- Chemical Affinity, defined, 76; laws of, 77; simple, 78; compound, 79; experiments on, 387—407
- Chemical Decompositions, table of, see APPENDIX.
- Chemical miracle, 627
- Chemical preparations necessary for the performance of the experiments, *page* 119
- Chemistry, usefulness of, 3—8
- Chloric oxide, 137
- Chlorides, 19, 108, 138
- Chlorine, 17, 19, 471
 ——— liquid, 472
- Chlorous oxide, 137
- Choke-damp, 104
- Chromic acid, 48
- Chromium, 29, 48
- Cinders, to chrySTALLIZE, so as to resemble specimens of minerals, 618
- Citrates, 112
- Citric acid, 112, 238
- Clay, 127
- Coal gas, 463
- Cobalt, 29, 45
 ———, nitro-muriate of, 502
- Cohesion, attraction of, 75
- Coke, 463
- Cold. This is treated of along with Heat, or Caloric, which see.
- Colour, 663; see also "Liquids."
- Coloured flames, 588
- Colouring matter, 234
- Columbic acid, 51
- Columbium, 29, 51
- Combination, 85

- Combustion, 85, 599, 600, 604
 Combustion under water, 582
 Concentration. The act of increasing the specific gravity of bodies. In general, the term is applied to fluids which are rendered stronger by evaporating a portion of the water they contain.
 Compound bodies, 14, 94
 Copal, 230
 Copper, 29, 35
 ———, acetate of, 206
 ———, carbonate of, 623
 ———, muriate of, 205
 ———, nitrate of, 203, 495
 ———, sulphate of, 204
 ———, oxide of, to make, 35
 ———, to silver, 638
 Copperas, 36, 207
 Cork, 238
 Corrosive sublimate, 34
 Cream of Tartar, 116, 169, 238
 Crucibles, 277, 298
 Cryophorous, 334, 384
 Crystallization, 93
 ———, water of, 93
 ———, instantaneous, 385
 Crystallizing, process of, 273, 639, 640, 641
 Crystals, beautiful, 641
 Cupel, 192
 Culinary salt, 170
 Cyanogen, 150
 DECANTATION, 264
 Decanting glasses, 301
 Decomposition. The separation of the constituent principles of compound bodies by chemical means.
 Decrepitation, 170
 Definite proportions, doctrine of, 89
 Deflagrating ladle, 329
 Deliquescence, 167
 Derbyshire spar, 190
 Deutoxide, 91
 Diamond, 24
 Digestion, 268
 Dissolution of metals, 156, 265
 Distillation, 271
 ———, compound, 294
 Dropping tube, 303
 Ductility. A property of certain bodies, in consequence of which they may be drawn out to a certain length without fracture.
 Dying, (art of,) 234, 561
 EARTHS, *Note to*, 29, 88, 122
 Effervescence. An internal motion which takes place in certain bodies, occasioned by the sudden escape of a gaseous substance.
 Efflorescent. Saline crystals, to which this term is applied, when exposed to the air, fall into powder, in consequence of losing a part of their water of crystallization.
 Electricity, 16
 Elements. The simple constituent parts of bodies, which are incapable of decomposition; they are frequently called principles. See *Simple Bodies*.
 Elutriation. The same as *washing*, which see.
 Empyreuma. A peculiar and indescribably disagreeable smell, arising

- from the burning of animal and vegetable matter in close vessels.
- Epsom-salt, 126, 195
- Equivalents (chemical). See APPENDIX.
- Ether, 244
- Eudiometer, common, 289, 324
- , Dr. Ure's, 357
- , Volta's, 358
- Evaporating basin, 299
- Evaporation, 270
- Expansion by heat, 362
- Experiments (chemical), general observations respecting, 254, and *page* 123
- Experiments:—On the properties of Heat, 360 to 386
- Experiments:—On Chemical Affinity, 387 to 407
- Experiments:—On Gases, 408 to 475
- Experiments:—On the preparation of Salts, 476 to 502
- Experiments:—On Chemical Tests, or Re-agents, 503 to 541
- Experiments:—Application of chemical knowledge to the Arts, and to Domestic Purposes, 542 to 574
- Experiments:—On Combustion, Detonation, and Light, 575 to 611
- Experiments:—Miscellaneous, 612 to 683
- FACE, luminous, 581
- Fecula, 225
- Fermentation, 239
- , vinious, 240
- , acetous, 245
- Fibrin, 249, 252
- Filters, 263
- Filtration, 263
- Fire produced by the mixture of two cold liquids, 601
- Fire, a fountain of, 464
- Fire-damp, 141
- Fireworks, with gas, 435
- Fixity of the noble metals, to show the, 622
- Flake-white, 42
- Flask, 345
- Flint, 128
- Flowers, in chemical language, are solid dry substances reduced to a powder by sublimation.
- Flowers of antimony, 43
- of benjamin, 113, 643
- of sulphur, 26, 643
- of zinc, 41
- Fluates, 109
- Fluor spar, 190
- Fluoric acid, 109, 612
- Fluorine, 17, 21, 128
- Fluxes, 680
- Frankincense, 230
- Frigorific mixtures, table of, see APPENDIX.
- Frost bearer, 334, 384
- Fulminating mercury, 611
- mixtures, 590 to 599
- powder, 586
- Furnace (Table-lamp), 314
- Furnaces, 278, 314
- Fusible metal, 615
- Fusion, 277
- GALENA, an ore of lead, analysed, 679
- Gallates, 114
- Gallic acid, 114, 238
- Galls, tincture of, 508
- Galvanic experiments, 667 to 677
- Galvanic pile, 331

- Galvanism, sketch of, 677
 Gas differs from vapour,
Note to, 18
 Gas-bottles, 281, 320, 327,
 328
 Gas-holder, 287, 352
 Gas, intoxicating, 135
 Gas, laughing, 135
 Gas-lights illustrated, 463
 Gas of Paradise, 135
 Gas, to fill bladders with,
 291, 326
 Gas, to try if combustible,
 288
 Gases, apparatus for, 280 to
 293
 Gases, experiments on, 408
 to 475
 Gases, how collected, 284,
 286
 Gases, how weighed, 293*
 Gasometer. An instrument
 used for the same purpose
 as a Gas-holder; but the
 name here intimates, that
 the instrument serves not
 only to contain the gas,
 but to measure its quan-
 tity also.
 Gelatin, 249, 250
 Ginger beer, to make, 542
 Glass, of what made, 128,
 176
 Glass, to cut, 683
 Glass, to engrave on, 612
 Glass jars, to graduate, 325
 Glass tubes, remarks on, 346
 Glauber's salts, 171
 Glucina, 68, 129
 Glucinum, 29, 68
 Glue, 250
 Gluten, 226
 Gold, 29, 30, 644
 Gold, nitro-muriate of, 199
 Golden figures, to make,
 upon charcoal, 619; upon
 silk, 616; upon steel, 636
 Graduated measurer, 304
 Granulation, 275
 Green Paint, 623
 Gum, of what composed, 24
 Gum, properties of, 223
 Gum Benzoin, 113, 643
 Gum Resins, 231
 Gunpowder, 587
 Gypsum, 189
 HARROWGATE water, 143
 Hartshorn, 193
 Heat, see Caloric.
 Homberg's phosphorus, 610
 ——— pyrophorus, 606
 Hydrates. Those substan-
 ces that have formed so
 intimate an union with
 water, as to solidify the
 water, and render it one
 of their component parts.
 Common slaked lime is
 properly a hydrate of lime.
 Hydrogen, 17, 22
 Hydrogen gas, directions for
 obtaining, and performing
 experiments with, 428 to
 441
 Hydrogen, arsenuretted, 47,
 144
 Hydrogen, carburetted, 24,
 141, 463
 Hydrogen, phosphuretted,
 142, 152, 464
 Hydrogen, potassuretted,
 146
 Hydrogen, sulphuretted, 26,
 143, 465
 Hydrogen, sulphuretted, li-
 quid, 466, 525
 Hydrogen, telluretted, 46,
 145
 Hydro-sulphurets, 143
 Hyposulphuric acid, 102

Hyposulphurous acid, 102

IGNES fatui, 142

Imponderable bodies, 16

Incineration. The burning of vegetables, for the sake of their ashes. It is usually applied to the burning of kelp on the sea-coast, for making mineral alkali.

Indian rubber, 233

Inflammable air, 22

Inflammation, curious cases of, 589, 600, 604

Infusion, 267

Ink-powder, 571

Ink-stains, to remove, 572

Inks, coloured, 570

Inks, indelible, 569

Inks, indelible, for marking linen, 573, 574

Inks, sympathetic, 648 to 662

Inks, writing, 567, 568

Instantaneous light, 560, 583

Iodides, 20, 139

Iodine, 17, 20

Iodine, vapour of, 613

Iridium, 29, 55

Iron, 29, 36

—, acetate of, 210

—, carbonate of, 209

—, gallate of, 211

—, muriate of, 208

—, nitrate of,

—, prussiate of, 212

—, sulphate of, 207, 496

—, oxide of, 421

Iron liquor, 210

Iron tree, 633

Isinglass, 250

Ivory, to silver, 647

JELLY, 250

Jet-pipes, for gas, 322, 340

KALI. A genus of marine

plants, which is burnt for the preparation of mineral alkali. The alkali is afterwards obtained by lixiviating the ashes, and then concentrating the saline liquor by evaporation.

Kelp, 174

Kermes, mineral. A medicinal preparation, formed by the union of sulphuret of antimony and potash.

LABORATORY, observations on fitting up a, 254

Lakes. Certain colours made by combining the colouring matter of cochineal, or of certain vegetables, with pure alumina, or with oxide of tin, zinc, &c.

Lamps, 313, 315, 335, 602

Lamp-Furnace, 314

Lapis calaminaris, 678

Latent heat, 382

Lead, 29, 38

—, acetate of, 216, 499, 516

—, carbonate of, 215

—, nitrate of, 499

—, oxides of, 90

—, ore of, analysed, 679

Lead tree, 634

Leather, 235, 250

Lemon juice, 238

Lemonade, to make, 544

Levigation, 261

Light, 16

Lime, 63, 125, 646

Lime, carbonate of, 188, 449

—, chloride of, 138

—, fluuate of, 190

—, muriate of, 192, 491, 519

—, nitrate of, 191, 491

—, oxalate of, 194

- Lime, phosphate of, 193
 —, sulphate of, 189
 Limestone, 125, 188
 Limestones, analysis of, 558
 Lime-water, 512
 Liquefaction. The change of a solid to the state of a fluid, in consequence of the combination of caloric with it.
 Liquids, remarkable alterations in, resulting from chemical action, 520 to 541
 Liquids, to measure given quantities of, 304—306
 Lithia, 62, 120
 Lithium, 29, 62
 Litmus, tincture of, 504
 Litmus test-papers, 505
 Lixiviation, 266
 Loadstone, 36
 Lunar caustic, 200
 Lutes, 271, 681
 MACERATION, 269
 Magnesia, 66, 126, 196
 —, carbonate of, 196
 —, sulphate of, 126, 195, 492
 Magnesium, 29, 66
 Magnetism, 16, 36, 39, 45
 Malic acid, 238
 Malleability. That property of metals which gives them the capacity of being extended and flattened by hammering.
 Manganese, 29, 44
 Marble, 125, 188
 Marls, analysis of, 558
 Martial. An old term for chemical preparations of iron.
 Massicot, 90
 Mastic, 230
 Matrass, 344
 Menstruum, 265
 Mercury, 29, 34
 —, fulminating, 611
 —, nitrate of, 202, 494, 522
 Metallic oxides, 90
 Metallic salts, 198
 Metallic trees, 631
 Metallurgy. The art of extracting and purifying metals.
 Metals, 17, 28, 29
 — as tests, 514
 Microcosmic salt. A compound formed of phosphoric acid, soda, and ammonia.
 Milk, 545
 Mineral waters, 220
 Mineralizers. Those substances that are combined with metals in their ores; such as sulphur, arsenic, oxygen, carbonic acid, &c.
 Mineralogy, 219
 Minerals, 219
 Minium, 90
 Moiree metal, 565
 Molybdenum, 29, 49
 Molybdic acid, 49
 Mordants. Substances which have a chemical affinity for particular colours; they are employed by dyers as a bond of union between colours and the cloth intended to be dyed with them. Alum is a body of this class.
 Mortar, 261, 300
 Mucous acid, 238
 Muriates, 108, 138
 Muriatic acid, 108, 469
 —, as a test, 515
 Musical sounds produced by the combustion of gas, 441

Myrrh, 231

NATRON. One of the names for mineral alkali, or soda.

Neutral salts, 155

Neutralize. When two or more substances mutually disguise each other's properties, they are said to neutralize one another.

Nickel, 29, 39

Nitrates, 106

Nitre. The same substance as nitrate of potass, or saltpetre.

Nitre, inflammable, 177

Nitric acid, 106

Nitric ether, 244

Nitric oxide, 134, 456

Nitrites, 107

Nitrogen, 17, 23, 247, 451 to 453

Nitro-muriates. Salts formed by the combination of bases with nitro-muriatic acid.

Nitro-muriatic acid, 199

Nitrous acid, 107

Nitrous gas, 134, 456

Nitrous oxide, 135; to procure, 460; to breathe, 462

OCHRES. Various combinations of the earths with oxide or carbonate of iron.

Oil of lime. Muriate of lime in a state of deliquescence.

Oil of vitriol, 100

Oil, fixed, 227

Oil, volatile, 228

Oils, 24, 227

Ores. Metallic earths mixed with extraneous matters; such as sulphur, arsenic, &c.

Osmium, 29, 53

Oxalates, 115

Oxalic acid, 115, 224, 238

———— as a test, 510

Oxides, 29, 88, 90, 133

Oxidizement. The operation by which any substance is combined with oxygen, in a degree not sufficient to produce acidity.

Oxygen, 17, 18

Oxygen gas, directions for obtaining and performing experiments with, 408 to 427

Oxygenize. To acidify a substance by oxygen.

PALLADIUM, 29, 33

Parting. The operation of separating gold from silver.

Pearl-powder, 218

Permanent white, 123

Peroxide, 91

Pewter, 37, 42, 43, 665

Philosophical taper, 434

Philosophical wool, 41

Phlogiston. This is an old chemical name for a substance now held to be *imaginary*; supposed anciently to be a combination of fire with some other body, and to be a constituent of all inflammable bodies, and of many other substances.

Phosphates, 103

Phosphorescence, 607

Phosphoric acid, 103, 419

Phosphoric fire-bottle, 583

Phosphoric spar, 190

Phosphorized ether, 578

Phosphorized oil, 581

Phosphorus, 17, 27, 575 to 582

- Phosphorus, caution respecting the use of, 576
 Phosphuret of lime, 152, 584, 585
 Phosphuret of sulphur, 152
 Phosphurets, 152
 Pinchbeck, 35
 Pitch, 230
 Plaster of Paris, 189
 Platinum, 29, 30
 Plumbago, 149
 Pneumatic. Any thing relating to the airs and gases.
 Pneumatic trough, 284, 316, 317
 Potash, 118
 Potass, 29, 60, 118
 ——— affinities of some of the acids for, 395 to 398
 ——— carbonate of, 166, 447, 448, 477
 ——— chlorate of, 165, 475, 590 to 599
 ——— nitrate of, 164, 478
 ——— oxalate of, 168
 ——— prussiate of, 167, 518
 ——— sub-carbonate of, 118, 166, 477
 ——— super-tartrate of, 169
 ——— tartrate of, 169, 479
 Potassium, 29, 60, 518-589
 Precipitant, 274
 Precipitate, 274
 Precipitating glasses, 301
 Precipitation, 274
 Protoxides, 91
 Prussian Blue, 36, 110, 212
 Prussiates, 110
 Prussic acid, 110
 Prussine, 150
 Pulverization, 261
 Putrefaction, 246, 253
 Pyrites. A mineral found abundantly in England. Some are sulphurets of iron, and others sulphurets of copper, with a portion of alumina and silica. The former are worked for the sake of the sulphur and the latter for sulphur and copper. This mineral is sometimes called Fire-stone; see 151.
 Pyrites (Martial). That species of pyrites which has iron for its base.
 Pyrometer. An instrument for measuring high degrees of heat; that is, for degrees of heat above the range of thermometers.
 Pyrophori. Certain substances that heat of themselves, and take fire when exposed to atmospheric air; see 606.
 QUICKLIME, 125
 Quicksilver, 34
 REAGENTS. Substances that are added to mineral waters or other liquids, to discover their nature and composition. *Reagent* is synonymous with *Test*; which see.
 Receiver, 271
 Red-lead, 90
 Reduction, 276
 Refining. The process of separating the perfect metals from other metallic substances by capellation.
 Refractory. A term applied to earths or metals that are infusible, or that require an extraordinary degree of heat to change or melt them.
 Regulus, in its chemical acceptance, signifies a me-

- tal purified from all extraneous matter. The term is an antiquated one.
- Repulsion, 81
- Resins, 230
- Retort, 271, 282
- Revivification, 276
- Rhodium, 29, 54
- Roasting. A preparative operation in mineralogy, by which the sulphur, arsenic, and other volatile bodies, combined with some minerals, are dissipated.
- Rock-crystal, 128
- Rust of iron, 36
- SAL ammoniac, 178
- Sal polychrest. An old name for sulphate of potass.
- Salifiable bases. All the bodies capable of combining with acids and forming salts.
- Saline. Partaking of the properties of a salt.
- Salt, common table, 61, 170
- Salt of lemons, 168
- Saltpetre, 164
- Salts, bi-, 155
- neutral, 155
- nomenclature of, 154
- phenomena attending the formation of, 156 to 160
- sub-, 155
- super-, 155
- the solubility of, 161
- what they are, 153
- Salts, triple. Salts formed by the combination of an acid with two bases. Alum (a sulphate of alumina and potass), and Rochelle salt (a tartrate of soda and potass), are both instances of this kind of combination.
- Saturation. The act of impregnating a liquid with another substance till no more can be received or imbibed. A fluid which holds as much of any substance as it can dissolve, is said to be saturated with that substance. A liquid, however, which is fully saturated with one substance, is still capable of dissolving other substances. At 162 is explained the mode of obtaining a saturated solution of a salt.
- Scales, 255
- Selenic acid, 52
- Selenium, 29, 52
- Sifting, 262
- Silica, 72, 128
- Siliceous earths. A term used to describe a variety of natural substances which are composed chiefly of silica; as quartz, flint, sand, &c.
- Silicium, 29, 72
- Silver, 29, 32
- Silver, muriate of, 201
- Silver, nitrate of, 200, 493, 515
- Silver, sub-nitrate of, 200
- Silver shrubbery, 635
- Silver tree, 632
- Simple bodies, what, 13 to 15
- Simple bodies described, 13 to 72
- Simple bodies, table of, 17
- Smelling salts, 179
- Smelting. The operation of fusing ores for the purpose of separating the me-

- tals they contain, from the sulphur and arsenic with which they are mineralized, and also from other heterogeneous matter.
 Soap, how made, 387
 Soap as a test, 520
 Soda, 61, 119, 174
 — borate of, 172
 — carbonate of, 174, 175
 — muriate of, 170, 481, 482
 — phosphate of, 173
 — sub-carbonate of, 174, 176, 480
 — sulphate of, 171, 483
 Soda-water, to make, 543
 Sodium, 29, 61
 Sodium, chloride of, 61
 Soils, analysis of, 557
 Soils, how to improve, 559
 Soluble tartar, 169
 Soluble bodies, 265
 Solution, 265
 Specific gravities, on taking, 257
 Specific gravities, table of, see APPENDIX.
 Specific gravity bottle, 260, 332
 Specific gravity of gases, to determine the, 293*
 Specific gravity of liquids, to determine the, 260
 Specific gravity of solids, to determine the, 258, 259
 Spirit-lamp, 315
 Spirit of salt, 108
 Spirit of wine, 242
 Spirits of hartshorn, 121
 Spoons which melt in hot water, 615
 Stalactites. Certain concretions of calcareous earth found suspended like icicles in caverns. They are formed by the oozing of water charged with this kind of earth through the crevices of the rocks.
 Starch, 24, 225
 Steel, 36, 148
 Still, 271
 Stop-cock, 323
 Strontia, 65, 124
 Strontia, carbonate of, 185
 Strontia, muriate of, 187, 490
 Strontia, nitrate of, 186, 490
 Strontia, sulphate of, 184
 Strontium, 29, 65
 Sub-salts, 155
 Suberic acid, 238
 Sublimation, 272, 643
 Succinic acid, 238
 Sugar, 24, 224
 Sugar of lead, 38, 216
 Sulphates, 100
 Sulphites, 101
 Sulphur, 17, 26, 643
 Sulphuret of carbon, 147
 Sulphuret of copper, 151
 Sulphurets, 26, 151
 Sulphuric acid, 100, 418
 Sulphuric acid, comparative affinities of several substances for, 399 to 404
 Sulphuric acid as a test, 509
 Sulphuric ether, 244
 Sulphurous acid, 101, 468
 Super-salts, 155
 Sweet Salts, 129
 Synthesis defined, 11
 TANNIN, 235
 Tannin, artificial, 235
 Tar, 230
 Tartaric acid, 116, 238
 Tartrates, 116
 Tea, 267
 Telluric acid, 46
 Tellurium, 29, 46
 Test-glasses, 303
 Test-papers, 505, 507

Tests, or re-agents, experiments on, 503 to 541

TESTS, *References to, for the presence of the following substances.*—

Acids in general, 504, 505, 528

Albumen, 251

Alkalies in general, 504, 506, 507, 526, 527

Ammonia, 522

Arsenic, 525

Barytes, 509, 521

Carbonates;—effervesce on adding acids.

Carbonic acids, 509, 512, 513, 528

Copper, 514, 537, 523, 540

Corrosive sublimate, 512

Earths in general, 511

Earths, dissolved by carbonic acid, precipitate on boiling.

Gallic acid, 389, 508

Gelatine, 508, 554

Gold, 524, 536

Iron, 508, 518, 535

Lead, 509, 525, 540

Lime, 509, 510, 512, 521

Magnesia, dissolved by acids;—precipitated by pure ammonia.

Magnesia, dissolved by carbonic acid, precipitates on boiling.

Metals, 511, 514, 518

Muriates, 515

Muriatic acid, 515, 533

Silver, 515, 541

Strontia, 521

Sulphuretted hydrogen, 516, 528

Sulphates, 534

Sulphate of lime, 551

Sulphuric acid, 513, 516, 517, 519, 534

Tin, 524

To distinguish iron from steel, 550

For impurities in wine, 547

To determine whether water be hard or soft, 520, 551

To detect copper in pickles or green tea, 548

To determine whether wheat-flour or bread be adulterated with chalk, 552

To discover if bread be adulterated with alum, 553

For impurities in magnesia, 555

To distinguish a solution of oxalic acid from one of Epsom salt, 546

For the purity of volatile oils, 549

Thermometer, 366 to 368

Thorina, 70, 132

Thorium, 29, 70

Tin, 29, 37

Tin, muriate of, 214, 497

— nitrate of, 213

— as a test, 524

Tincal, 172

Tin-plate, 36

Titanium, 29, 57

Tritoxide, 91

Trituration, 261

Tube of safety, 296, 349

Tungstenum, 29, 50

Tungstic acid, 50

Turmeric, tincture of, 507

Turmeric test-papers, 507

Turpentine, 230

UNION, chemical. When a mere mixture of two or more substances is made, they are said to be *mechanically* united; but

- when each or either substance forms a permanent component part of the product, the substances have formed a *chemical union*.
- Uranium, 29, 56
- VAPOUR differs from gas, *Note to*, 18
- Vegetable acids, 238
- Vegetable substances, 222
- Verdigris, 206
- Vermilion, 34
- Vinegar; see Acetic acid.
- Vital air, one of the names applied to Oxygen gas.
- Vitriol, blue, 204
- Vitriol, green, 36, 207
- Vitriol, white, 41, 217
- Volatile alkali, 121
- Voltaic pile, 331
- WASHING, 262
- Water, the basis of, 22
- Water, composition of, 89, 440
- Water, freezing and boiling points of, 366
- Water, properties of, 96
- Water, mineral. Waters which are impregnated with mineral and other substances, are known by this appellation. These minerals are generally held in solution by carbonic, sulphuric, or muriatic acid.
- Waves of fire on water, 58
- Wax, 236
- Way, dry and humid, *page* 75
- Weighing, 255
- Welding, 30, 36
- White-lead, 215
- Will-of-the-wisp, 142
- Wine, 240
- Wine-test, 547
- Wodanium, 29, 59
- Wood, 237
- Worm, 271
- Woulfe's apparatus, 29, 348
- Woulfe's gas bottle, 320
- Writing, luminous, 575
- YEAST, 240
- Yttria, 67, 131
- Yttrium, 29, 67
- ZAFFRE. An oxide of cobalt, mixed with a portion of siliceous matter. In this state it is imported from Saxony.
- Zero, 366
- Zinc, 29, 41
- Zinc, ore of, analyzed, 677
- Zinc, oxide of, 423
- Zinc, sulphate of, 217, 500
- Zirconia, 71, 130
- Zirconium, 29, 71

FINIS.

