A calendar of flora composed during the year 1809, at Warrington, lat. 53*0 30' / [George Crosfield].

Contributors

Crosfield, George, 1785-1847.

Publication/Creation

Warrington [England]: J. Haddock for Wilkie & Robinson, London, 1810.

Persistent URL

https://wellcomecollection.org/works/ah474vsw

License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
https://wellcomecollection.org

CALENDAR

OF

FLORA,

COMPOSED DURING THE YEAR

1809,

at

Warrington,

Lat. 53°. 30'.

BY GEORGE CROSFIELD, SECRETARY

TO THE BOTANICAL SOCIETY OF WARRINGTON.

These, as they change, ALMIGHTY FATHER, these, Are but the varied God. The rolling year Is full of Thee.-----

Mysterious round! what skill, what force divine,
Deep felt, in these appear! a simple train,
Yet so delightful mix'd, with such kind art,
Such beauty and beneficence combin'd;
Shade, unperceiv'd, so softening into shade;
And all so forming an harmonious whole;
That, as they still succeed, they ravish still.

THOMSON.

WARRINGTON:
Printed by J. Haddock.

SOLD BY WILKIE AND ROBINSON, PATERNOSTER-ROW;
AND WHITE, FLEET-STREET,
LONDON.

1810.

CALENDAR

FLORA,

25336(2) 0081

Tellarrington,

Lat. 53". 30'.

BY GEORGE CROSFYELD,

O THE ROTANICAL SOCIETY OF WARRISCHOOL

And on the change 'Agustage Parties, Lond.

Despite the contract of the co

THOMESON'S

Prince by J. Pittled

SOLD BY WILKIE AND POSITION PATERNOSTEE-ROW,

tell tot too bearing by the first

advanta 810 afforded, by my sinceron a

PREFACE.

An asterist has been prefered to all those plants, which,

memirers of which I seknowledge myself under considerable

obligation for their ready authance.

DINCE the science of Botany became a branch of study, considered at once useful and instructive, it has been deemed of some importance to point out the different vegetable productions of particular districts. Some very respectable Botanists have likewise expressed a wish that the periods of inflorescence in different seasons and in different latitudes should be marked down with accuracy. To unite these two plans, at the least possible expence, is my design in the following pages, which contain a list of upwards of eight hundred British Plants, comprized in the twenty-three first Classes of the Linnean system, growing, either wild or in a cultivated state, in the neighbourhood of Warrington, and arranged according to their earliest periods of inflorescence during the year 1809. It would, I am aware, have added much to its value, to have been accompanied by an accurate meteorological journal; but such an undertaking would have occupied more time than was consistent with my commercial avocations. It may not however be amiss to say, that, with the exception of a few warm days early in the 3d Month, (March.) the Spring was delayed later than usual by constant easterly winds, vegetation being kept back until the third of the 5th Month, (May.). From this time until the beginning of the following month the progress of vegetation was very rapid. The latter end of Summer was attended with much rain.

In compiling the following Calendar of Flora, I have not only depended on my own personal observation, but have availed myself of the advantages afforded by my situation as

Secretary to a provincial Botanical Society, to some of the members of which I acknowledge myself under considerable obligation for their ready assistance.

With a few exceptions, no plant has been admitted into the following list, which is not ranked as indigenous in the very excellent *Flora Britannica* of Dr. Smith. Where the rule is departed from, the authority for it is specified.

An asterisk has been prefixed to all those plants, which, although indigenous, are only found, in this neighbourhood, in a state of cultivation. A few, observed in the more western, and northern parts of this county, are distinguished by a dagger.

The notes will not I trust, be deemed superfluous. For those marked K, I am indebted to my esteemed friend James Kendrick, M. D. F. L. S. President of the Botanical Society of Warrington.

The town of Warrington is situated at the southern extremity of the county of Lancaster, the river Mersey dividing it from that of Chester. The broad estuary of this River commences about six miles westward. In its immediate vicinity, is a large quantity of rich meadow land, formed principally of alluvial soil. A large extent of peat moss commences about two miles eastward of the town. The only high ground in its neighbourhood is a hill about two miles to the south, formed of red sand stone, covered with a stratum of sandy soil.

In every instance I have adopted the nomenclature of Dr. Smith: where he has differed from Linnæus, or where subsequent writers have differed from him their synonyms are also given.

GEORGE CROSFIELD.

WARRINGTON,
1st of 1st Month, 1810.

A CALENDAR, &c.

FIRST MONTH (January).

- 14. ULEX europæus. Common Furze. It handher
- 16. BELLIS perennis. Common Daisy. (1)

SECOND MONTH (February).

- 2. * HELLEBORUS fatidus. Stinking Hellebore. 40
- 12. SENECIO vulgaris. Groundsel.
- 14. GALANTHUS nivalis. Snowdrop.

 * CROCUS vernus. Spring Crocus.

 VINCA minor. Lesser Periwinkle.
- 26. DAPHNE Mezereum. Mezereon.

THIRD MONTH (March).

- 2. LAMIUM purpureum. Red Archangel.
- 7. \$VIOLA odorata, Sweet Violet.
- 10. PRIMULA vulgaris. Primrose. Staron.
- 11. TUSSILAGO Farfara. Colts-foot.

----- Petasites. Common Butter-bur.

Reil 40

- 11. LEONTODON Taraxacum. Common Dandelion. (2)
 STELLARIA media.
 ALSINE media. Linn. Common Chickweed.
- 12. * SCILLA bifolia. Two-leaved Squill.

 RANUNCULUS Ficaria. Pilewort. (3)

 BETULA Alnus.

 ALNUS glutinosa. Willdenow. Common Alder.

POPULUS alba. Great White Poplar.

- 13. ULMUS campestris.
 ULMUS suberosa. Willd. Common Elm-tree.
- 14. FRAGARIA sterilis. Barren Strawberry.
 SALICIS species variæ. Various kinds of Willow. (4)
- POA annua. Annual Meadow-grass.
 DRABA verna. Common Whitlow-grass.
- 19. * PULMONARIA officinalis. Common Lung-wort.
- 23. a VIOLA odorata. Sweet Violet.

 NARCISSUS Pseudo-narcissus. Common Daffodil.

 DAPHNE Laureola. Spurge-Laurel.

 CARDAMINE hirsuta. Hairy Ladies-Smock.
- 25. VERONICA verna. Vernal Speedwell.

 OXALIS Acetosella. Common Wood-Sorrel.
- (2) Whether this plant possesses the deobstruent properties attributed to it by some authors of reputation may be doubted. Its diuretic effects are in some instances considerable. Its vulgar name, in more than one language, implies a knowledge of its powers. K_{\bullet}
- (3) It may not be amiss to state that this termination so common in the names of British plants, is derived from the Saxon Wyrt, a herb or plant, and implies that the plant so denominated, has been supposed to possess some specific power of curing the disease of a particular organ, e.g. Lungwort, Liverwort, &c. &c. K.
- (4) The bark of all the species of Salix, containing a considerable portion of Tannin, is useful in the art of tanning leather. With regard to its medical properties, it may suffice to say, that it is by no means a despicable tonic, though experience and analysis prove it an improper substitute for the Bark of the Cinchona lancifolia (Peruvian Bark). K.

25. ULMUS montana. Willd. Broad-leaved Elm-tree. Rich Al

26. β VIOLA tricolor. Pansy Violet.

ADOXA Moschatellina. Tuberous Moschatel.

ANEMONE nemorosa. Wood Anemone. has a kibbliller. III

THLASPI arvense. Smooth Mithridate Mustard.

MERCURIALIS perennis. Perennial Mercury.

PRIMULA elatior.
β PRIMULA veris. Linn.
β Oxlip.

CHRYSOPLENIUM oppositifolium. Opposite-leaved
Golden-Saxifrage.

RANUNCULUS hederaceus. Ivy Crowfoot.

CALTHA palustris. Marsh Marygold.

CHEIRANTHUS fruticulosus. Wallflower.

FOURTH MONTH (April).

ALCHEMILLA arvensis.
APHANES arvensis. Linn.
ALCHEMILLA aphanes. Willd. Field Ladies Mantle.

CERASTIUM vulgatum. Broad-leaved Mouse-ear

Chickweed.
GLECHOMA hederacea. Ground-Ivy.

GLECHOMA hederacea. Ground-Ivy.
ARABIS thaliana. Common Wall-Cress.

2. VIOLA canina. Dog's Violet.

JUNCUS sylvaticus.

JUNCUS pilosus. Linn.

Wood Rush.

3. CHARA vulgaris. Common Chara, Ju Read : 54

- 3. *ORNITHOGALUM luteum. YellowStar of Bethlehem.

 *ANEMONE apennina. Blue mountain Anemone.

 *FUMARIA solida. Solid bulbous Fumitory.
 - THLASPI campestre. Common Mithridate Mustard.
 - BUXUS sempervirens. Box-tree.
 - * RUSCUS aculeatus. Butcher's-Broom.
 - * VISCUM album. Misseltoe.
- 4. * VINCA major. Greater Periwinkle.
- 5. RIBES rubrum. Common Currants.
 - * FRITILLARIA Meleagris. Common Fritillary.
 - * ANEMONE Pulsatilla. Pasque Flower. MYRICA Gale. Sweet Gale.
- 7. PRIMULA farinosa. Bird's eye Primrose. Jen powden
- 8. * HELLEBORUS viridis. Green Hellebore.
- 9. GALIUM Aparine. Goose-grass. (5)
- 10. RIBES Uva-crispa. Smooth Gooseberry.

 CARPINUS Betulus. Horn-beam.

 POPULUS canescens. Common White Poplar.
- 11. * ECHIUM vulgare. Common blue Vipers-Bugloss.
 TUSSILAGO hybrida. Longer stalked Butter-bur.
 * ARABIS stricta. Bristol Rock-cress.
- a CALLITRICHE aquatica.

 14. CALLITRICHE verna. Linn. } Water Star-wert.

 ALCHEMILLA vulgaris. Common Ladies Mantle.
 - * DENTARIA bulbifera. Bulbiferous Coral-wort. CARDAMINE pratensis. Meadow Ladies-smock.
- 2 VIOLA tricolor. Pansy Violet. PRUNUS spinosa. Sloe-tree.
- 17. ----- insititia. Bullace-tree.
 - * SCROPHULARIA vernatis. Yellow Fig-wort.

⁽⁵⁾ The powers of late attributed to this plant, as an anticarcinomatous remedy, may reasonably be doubted. It has long been a popular remedy in those complaints commonly called scorbutic. Its root, in common with that of Madder, (Rubia tinctorum Linn.) has the property of tinging the bones of animals fed upon it, of a red colour. K.

17. * COCHLEARIA danica. Danish Scurvy-grass.
ERYSIMUM Alliaria. Garlick Hedge-mustard. Rech 36

18. * NARCISSUS poeticus. Poetic Narcissus.
PRUNUS domestica. Plum-tree.

19. CHARA hispida. Prickly Chara. Red 54
RANUNCULUS auricomus. Wood Crowfoot.

23. CERASTIUM aquaticum. Water Chickweed.

PRUNUS Cerasus. Cherry-tree.

THLASPI Bursa-Pastoris. Common Shepherd's Purse.

BRASSICA Rapa. Turnep. Rect. 36

25. * IXIA Bulbocodium. Crocus-leaved Ixia. (Curtis.)

* EPIMEDIUM alpinum. Barren-wort.

* NARCISSUS biftorus. Pale Narcissus.
PYRUS communis. Iron Pear-tree. Rich 36

26. RIBES nigrum. Black Currants.

* ASARUM europæum. Asarabacca.

27. CHÆROPHYLLUM sylvestre. Smooth Cow Parsley.
STELLARIA nemorum. Wood Stitchwort.
COCHLEARIA anglica. English Scurvy-grass.
SALIX repens. Creeping dwarf Willow.

28. STELLARIA holostea. Greater Stitchwort. recy land companies BETULA alba. Common Birch. Cathele field from ment the street.

29. VERONICA arvensis. Wall Speedwell. ASPERULA odorata. Sweet Woodruff.

EUPHORBIA Cyparissias. Cypress Spurge. Rep. 16, 39

FRAGARIA vesca. Wood Strawberry.

α RANUNCULUS aquatilis. β RANUNCULUS aquatilis. β Water Crowfoot

So. a VALERIANA Locusta.

VALERIANA olitoria. Willd. Corn Sallad. Confidence of the Confidence of the

⁽⁶⁾ A year or two ago, I met with a variety of this species, having the summit of the spike surmounted by leaves; which I considered as an accidental conversion of the terminal flowers, rather than as a permanent variety.

Qu. Is this var. 3. of Withering, described as having "the summit of the stalk surrounded with leaves longer than the spike?"

30. * LEUCOJUM astivium. Summer Snow-flake. MENZIESIA polifolia. Jussieu. ANDROMEDA polifolia. Marsh Rosemary. a LYCHNIS dioica. Red Campion. IBERIS nudicaulis. Naked-stalked Candy-tuft. TURRITIS hirsuta. Hairy Tower-Mustard. SINAPIS arvensis. Charlock.

Rich F. 24 _ GERANIUM molle. Dove's-foot Crane's-bill. Richard Tape 23 ORCHIS mascula. Early purple Orchis. Rost in

FIFTH MONTH (May).

- That's Just 1. VERONICA agrestis. Procumbent Speedwell. Rich 40 ANCHUSA sempervirens. Evergreen Alkanet. ACER Pseudo-platanus. Greater Maple. Friend 228 CERASTIUM semidecandrum. Little Mouse-ear-Chickweed.
 - * TROLLIUS europæus. Globe-flower.
 - * FUMARIA lutea. Yellow Fumitory.
 - * SALIX herbacea. Least Willow.
 - 2. * GENTIANA acaulis. Dwarf Gentian. (E. Bot.)
 - 3. VACCINIUM Myrtillus. Bilberry. ACER campestre. Common Maple. HOLOSTEUM umbellatum. Umbelliferous Chickweed. CARDAMINE amara. Bitter Ladies-smock.

GERANIUM cicutarium. Linn. Hemlock Stork's-bill. GERANIUM pusillum. Small-flowered Crane's-bill. FAGUS sylvatica. Beech-tree. Forusty lan a complete cornin

5. HIPPURIS vulgaris. Common Mare's-tail. (7) * ORNITHOGALUM umbellatum. Common Star of Bethlehem. This flowers ofens its blow at 11 & about the

GERANIUM lucidum. Shining Crane's-bill. CAREX pracox. Vernal Carex.

Satifica granulata . Thecenta Hestry

(7) Was observed to emerge on the 12th of 4th Month. (April.) K. torres in blent, or the dornt of lain adjunity Teddington lock

of teal year, undersourface burkle

- 6. * ARBUTUS Uva-ursi. Red-berried trailing Arbutus. (8)
 * SAXIFRAGA umbrosa. London Pride.
- BUNIUM flexuosum. Common Earth-nut.
 * SAXIFRAGA oppositifolia. Purple Saxifrage.

MESPILUS Oxyacantha. CRATÆGUS Oxyacantha. Linn. } Hawthorn. (9) Red 5764

CAREX paludosa. Lesser common Carex.

8. VERONICA serpyllifolia. Smooth Speedwell. Parties Filary Hels

* VIOLA lutea. Yellow mountain Pansy.

SCANDIX odorata. Great Chervil.

PYRUS Malus. Apple-tree. Rich 36, 40, 44, 57

* GEUM rivale. Water Avens.

PEDICULARIS palustris. Marsh Louse-wort. Has an arillus Rich 3

----- sylvatica. Pasture Louse-wort. (10)

POLYGALA vulgaris. Milk-wort.

CAREX dioica. Common separate-headed Carex.

* SALIX reticulata. Wrinkled Willow.

9. VERONICA montana. Mountain Speedwell. The standard or

ANTHOXANTHUM odoratum. Sweet-scented Vernal-grass.

SAGINA apetala. Annual small-flowered Pearl-wort.

CHENOPODIUM Bonus Henricus. Perennial Goosefoot.

SCILLA nutans.
HYASCINTHUS nonscriptus. Linn. } Hare-bell. chista of rocie les

* SAXIFRAGA hypnoides. Mossy Saxifrage.

+ Throbanga granulata

* CERASTIUM tomentosum. Woolly Mouse-ear. (HULL.)

abortens ANTIRRHINUM Cymbalaria. Ivy-leaved Snapdragon. approprie

(8) The tonic and sedative properties of this plant are not, I apprehend, in general fully appreciated, notwithstanding what has been said on the subject by Doctors Ferriar and Bourne. In many cases I have found it not a little efficacious. K.

(10) Frequently varies to white.

the avoto have a mass of made tuberder containing and whatle of reproducing the plant without being unrounded with Mular tipue full of anylaceous Je outs as the potatore

⁽⁹⁾ Its inflorescence usually commences on the first of this month (MAY).

B. 40

9. * IBERIS amara. Bitter Candy-tuft. FUMARIA officinalis. Common Fumitory. Matrices in 2 bundle * GNAPHALIUM dioicum. Mountain Cud-weed. * SAXIFRAGA tridectylites. Rue-leaved Saxifrage. 10. RANUNCULUS sceleratus. Celery-leaved Crowfoot. ----- hirsutus. Pale hairy Crowfoot. ---- acris. Upright meadow Crowfoot. Item law GALEOBDOLON luteum. GALEOPSIS Galeobdolon. Linn, Yellow Dead-Nettle. LEONURUS Galeobdolon. Willd.) * GERANIUM phæum. Dusky Crane's-bill. * CENTAUREA Cyanus. Corn Blue-bottle. FRAXINUS excelsior. Common Ash. Wat officed done to VERONICA Chamedrys. Germander Speedwell. VALERIANA dioica. Marsh Valerian. Page of the con-SHERARDIA arvensis. Blue Sherardia. β MYOSOTIS scorpioides. Mouse-ear Scorpion-grass. MENYANTHES trifoliata. Buckbean. (11) Pools of wat * LONICERA Caprifolium. Pale perfoliate Honeysuckle. Black Salt-wort. GLAUX maritima. STATICE Armeria. Common Thrift. JUNCUS campestris. Hairy field Rush. RUMEX Acetosa. Common Sorrel. TRIGLOCHIN maritimum. Sea Arrow-grass PARIS quadrifolia. Herb Paris. Filent brolongo lega ARENARIA rubra. Purple Sandwort. PRUNUS Padus. Bird Cherry. PYRUS aucuparia. Mountain-ash. SORBUS aucuparia. POTENTILLA anserina. Silver-weed. TORMENTILLA officinalis. Common Tormentil. TORMENTILLA erecta. Linn.

---- reptans. Trailing Tormentil.

⁽¹¹⁾ Varies frequently in the number of stamens, each flower having often six, and sometimes seven. K.

CHELIDONIUM majus. Celandine. 11. * AQUILEGIA vulgaris. Columbine. RANUNCULUS bulbosus. Bulbous Crowfoot. Jone and denthe hills AJUGA reptans. Common Bugle. GERANIUM robertianum. Stinking Crane's-bill. (12) then frage GENISTA anglica. Needle Green-weed. Its and with things TRIFOLIUM procumbens. Hop Trefoil. a LOTUS corniculatus. Common Bird's-foot Trefoil. CAREX pulicaris. Flea Carex. ---- arenaria. Sea Carex. QUERCUS Robur.

QUERCUS pedunculata. Willd. } Common British Oak. 12. VALERIANA rubra. Red Valerian, try about in Chalkhets his SCIRPUS caspitosus. Scaly-stalked Club-rush. a MYOSOTIS scorpioides. Mouse-ear Scorpion-grass. POLEMONIUM caruleum. Jacob's Ladder. Jacob's Ladder. * CONVALLARIA multiflora. Common Solomon's Seal x Julian RUMEX Acetosella. Sheep's Sorrel. * RUBUS arcticus. Dwarf crimson Bramble. RANUNCULUS Flammula. Lesser Spear-wort. Rich 52 THLASPI perfoliatum. Perfoliate Shepherd's Purse. COCHLEARIA Armoracia. Horse-radish. CORONOPUS Ruellii.
COCHLEARIA Coronopus. Linn. Common Wart-cress ERYSIMUM Barbarea. Yellow Rocket. FUMARIA claviculata. White climbing Fumitory. CAREX vulpina. Great spiked Carex. URTICA dioica. Great Nettle. 13. SAGINA procumbens. Procumbent Pearl-wort. Rich 3/ -ALLIUM ursinum. Broad-leaved Garlick. * CONVALLARIA majalis. Lily of the Valley.

SCLERANTHUS annuus. Annual Knawel. RANUNCULUS repens. Creeping Crowfoot. ORNITHOPUS perpusillus. Common Bird's-foot. TRIFOLIUM pratense, Common purple Trefoil.

⁽¹²⁾ Rarely varies to white.

SCIRPUS triqueter. Triangular Club-rush. ALOPECURUS pratensis. Meadow Fox-tail-grass. SAGINA erecta. Upright Pearl-wort. GALIUM cruciatum. Crosswort. Ju Rul 724 VALANTIA cruciata. Linn. LYSIMACHIA nemorum. Wood Loosestrife. HYDROCOTYLE vulgaris. Marsh Pennywort. * STAPHYLEA pinnata. Bladder-nut-tree. BERBERIS vulgaris. Barberry. (13) * VACCINIUM Vitis Idaa. Red Whortle-berry. STELLARIA uliginosa. Bog Stitchwort. ARENARIA peploides. Sea Sandwort. * PAPAVER cambricum. Yellow Poppy. ERYSIMUM officinale. Common Hedge-mustard. SPARTIUM scoparium. Common Broom. LATHYRUS Nissolia. Crimson Lathyrus. HIERACIUM Pilosella. Mouse-ear Hawkweed. URTICA wens. Small Nettle.

ARUM maculatum. (Spadice purpureo.) Cuckowpint.

STELLARIA graminea. Lesser Stitchwort.

* DORONICUM Pardalianches. Great Leopard's-bane. CAREX acuta. Slender-spiked Carex.

----- riparia. Great common Carex.

VERONICA Beccabunga. Brooklime. 16. & MYOSOTIS scorpioides. Mouse-ear Scorpion-grass. EUPHORBIA Peplus. Petty Spurge. SISYMBRIUM Nasturtium. Water Cress.

-- terrestre. SISYMBRIUM palustre. Willd. Annual water Rocket HOLCUS avenaceus.

AVENA elatior. Linn. } Oat-like Soft-grass. POA trivialis. Roughish Meadow-grass.

GALIUM saxatile. Smooth heath Bed-straw.

⁽¹³⁾ It would be worth while to ascertain by experiment, whether the very probable conjecture of Dr. Smith respecting the fructification of this plant has any foundation in fact. K.

16. BUNIUM Bulbocastanum. Great Earth-nut.

HOTTONIA palustris. Water Violet. Leaves submered

* CONVALLARIA Polygonatum. Angular Solomon's Seal.

VICIA sativa. Common Vetch.

TRIFOLIUM minus. Lesser yellow Trefoil.

17. CORNUS sanguinea. Wild Cornel-tree.

ILEX Aquifolium. Holly-tree.

* LONICERA Xylosteum. Upright Honeysuckle.

VACCINIUM Oxycoccus. Cranberry.

POTENTILLA reptans. Common creeping Cinquefoil.

GEUM urbanum. Common Avens.

RANUNCULUS arvensis. Corn Crowfoot.

CAREX ovalis. Oval-spiked Carex.

18. VERONICA spicata. Spiked Speedwell.

----- officinalis. Male Speedwell.

LOLIUM perenne. Ray-grass.

SCANDIX Anthriscus. Rough Chervil.

ÆGOPODIUM Podagraria. Gout-weed.

RUMEX obtusifolius. Broad-leaved Dock.

GERANIUM dissectum. Jagged-leaved Crane's-bill.

TRIFOLIUM repens. White Trefoil.

ANTHEMIS Cotula. Stinking May-weed.

19. * ALLIUM Schenoprasum. Chive Garlick.

* CISTUS Helianthemum. Common dwarf Cistus.

* ARISTOLOCHIA Clematitis. Common Birth-wort. Cale

* RHODIOLA rosea. Rose-root.

20. PARIETARIA officinalis. Pellitory of the Wall. Ju P.

P. 24 SYMPHYTUM officinale. Common Comfrey.

* RHAMNUS Frangula. Alder Buckthorn.

* PYRUS Aria.

CRATÆGUS Aria. Linn.

White Beam-tree.

CRAMBE maritima. Sea Kale.

GERANIUM sanguineum. Bloody Crane's-bill.

MALVA rotundifolia. Dwarf Mallow. Comolis

* JUNIPERUS communis. Common Juniper.

21. SCIRPUS palustris. Marsh-creeping Club-rush.

JUNCUS effusus. Soft Rush.

LYCHNIS Flos-Cuculi. Meadow Lychnis.

THYMUS Serpyllum. Wild Thyme.

SENECIO aquaticus. Marsh Ragwort.

ORCHIS Morio. Meadow Orchis.

23. SCIRPUS maritimus. Salt-marsh Club-rush.

POA fluitans.
FESTUCA fluitans. Linn. } Flote Meadow-grass.

POA rigida. Hard Meadow-grass.

DACTYLIS glomerata. Rough Cock's-foot-grass.

PLANTAGO Coronopus. Buck's-horn Plantain. Purcant at particular to proper the proper to the prope

B JUNCUS acutus. Linn. S Polygonum Bistorta. Great Bistort. He State B LYCHNIS dioica. White Campion. (14)

HEDYPNOIS hispida.

LEONTODON hispidum. Linn.

APARGIA hispida. Willd.

Rough Hedypnois.

TAMUS communis. Black Bryony.

ALOPECURUS agrestis. Slender Fox-tail-grass.

----- geniculatus. Floating Fox-tail-grass.

AGROSTIS vulgaris. Fine Bent-grass. HOLCUS lanatus. Meadow Soft-grass.

⁽¹⁴⁾ LYCHNIS vespertina of Sibthorp and Symons, so called from its

24. POTAMOGETON gramineum. Grassy Pond-weed. CENANTHE crocata. Hemlock Water Dropwort. Rook at Kno ALLIUM vineale. Crow Garlick. ORCHIS maculata. Spotted palmate Orchis.

AIRA aquatica. Water Hair-grass. 25.

* BRIZA maxima. Great Quake-grass. (Hull.)

* ALCHEMILLA alpina. Alpine Ladies-mantle.

* CYNOGLOSSUM officinale. Common Hounds-tongue.

HERACLEUM Sphondylium. Common Cow-parsnep.

* SAXIFRAGA palmata. Palmate Saxifrage.

SCROPHULARIA nodosa. Knotty-rooted Fig-wort. Rech 27, 28

BRASSICA campestris. Field Cabbage. Rich 36,50

RAPHANUS Raphanistrum. Wild Radish. Ju Rich 36

TRAGOPOGON porrifolius. Purple Goat's-beard. Tappen feathery

OPHRYS ovata.

EPIPACTIS ovata. Swartz.

Common Twayblade.

26. SCIRPUS setaceus. Least Club-rush.

SANGUISORBA officinalis. Great Burnet.

LITHOSPERMUM arvense. Corn Gromwell.

VIBURNUM Opulus. Common Guelder-rose. 14ct toling to

PEPLIS Portula. Water Purslane.

TRIGLOCHIN palustre. Marsh Arrow-grass.

FUMARIA capreolata. Ramping Fumitory. (15)

HIERACIUM murorum. Wall Hawkweed.

27. AVENA fatua. Wild Oat.

* RHAMNUS catharticus. Purging Buckthorn. Perksy

HYDROCOTYLE inundata. } Floating White-rot.

SISON inundatum. Linn.

- * MESPILUS germanica. Common Medlar.
- * SPIRÆA Filipendula. Common Dropwort.
- * GERANIUM columbinum. Long-stalked Crane's-bill.
- AIRA præcox. Early Hair-grass.
 - * ASPARAGUS officinalis. Common Asparagus. (Current)
 - * SAXIFRAGA granulata. White Saxifrage. Veronica agrestes in cultivated freezes

(15) Very common in the hedges in many places in this neighbourhood.

* CISTUS marifolius. Hoary dwarf Cistus. 28. RANUNCULUS parvulus. Little upright Crowfoot. * ERODIUM moschatum. Musky Stork's-bill. GERANIUM moschatum. Linn. * VICIA lathyroides. Spring Vetch. Juds cubord

29. VERONICA Anagallis. Water Speedwell. CYNOSURUS cristatus. Crested Dog's-tail-grass. GALIUM uliginosum. Rough marsh Bed-straw. SOLANUM Dulcamara. Woody Nightshade. (16) CHÆROPHYLLUM temulentum. Rough Cow-parsley. COTYLEDON Umbilicus. BCOTYLEDON Umbilicus. Linn. Common Navel-wort. COMARUM palustre. Marsh Cinquefoil.

Tighe Stigma butkley PAPAVER Rhags. Common red Poppy. Calgy of 2 10 26,27 DIGITALIS purpurea. Purple Fox-glove. of of the a ORCHIS bifolia. Butterfly Orchis.

30. AVENA pratensis. Narrow-leaved Oat-grass. ANAGALLIS arvensis. Scarlet Pimpernel. Just hajed

* EUONYMUS europæus. Common Spindle-tree. It. DAUCUS Carota. Wild Carrot.

ACORUS Calamus. Sweet Flag.

* TOFIELDIA palustris. ANTHERICUM calyculatum. Linn. Scottish Asphodel. HELONIAS borealis. Willd. RESEDA Luteola. Dyer's-weed.

26 57 NYMPHÆA lutea. Yellow Water-lily. * HESPERIS inodora. Scentless Dame's-violet. OROBUS tuberosus. Common Orobus.

ALOPECURUS bulbosus. Bulbous Fox-tail-grass. MELICA uniflora. Wood Melic-grass. * POA bulbosa. Bulbous Meadow-grass.

Eurof: Flowers greenish while of foetra, Cupiniles purple

⁽¹⁶⁾ That the berries of this plant possess, highly poisonous qualities is well known. That the stems are not entirely destitute of narcotic properties is equally true. In one instance where a very large dose had been swallowed, paralysis of the tongue and fauces, with diminished sensibility of the retina was the consequence. Its effects were not entirely removed for several days. K.

* PINGUICULA vulgaris. Common Butterwort. 31. PLANTAGO maritima. Sea Plantain. POTAMOGETON densum. Close-leaved Pond-weed. * LITHOSPERMUM officinale. Common Gromwell. LYCOPSIS arvensis. Small Bugloss. MEUM athamanticum. ATHAMANTA Meum. Linn. > Spignel. ÆTHUSA Meum. Willd. JUNCUS acutus. Great sharp Rush. ----- conglomeratus. Common Rush. RUMEX sanguineus. Bloody-veined Dock. SILENE maritima. Sea Campion. & CUCUBALUS Behen. ARENARIA marina. Sea spurrey Sandwort. BARENARIA rubra. RUBUS idwus. Raspberry. PAPAVER Argemone. Long rough headed Poppy.

----- dubium. Long smooth headed Poppy.

NYMPHÆA alba. White Water-lily. Filements He. CARDUUS lanceolatus. Spear Thistle.

CHRYSANTHEMUM Leucanthemum. Great white Ox-eye.

ORCHIS latifolia. Marsh Orchis.

ARUM maculatum. (spadice albo) Cuckow-pint. (17)

SIXITH MONTH (June).

- 1. SONCHUS arvensis. Corn Sow-thistle.

 ACHILLEA Millefolium. Common Yarrow.
- 2. RUMEX acutus. Sharp Dock.
 POLYGONUM aviculare. Knotgrass.
 * ALYSSUM sativum.
 MYAGRUM sativum. Linn. } Gold of Pleasure.

3. BRIZA media. Common Quaking-grass.

alledota - an orcher Jones &

⁽¹⁷⁾ I have observed that the inflorescence of those plants in which the Spadix is white, uniformly takes place later than when it is purple.

3. RUMEX crispus. Curled Dock.

LATHYRUS latifolius. Broad-leaved Lathyrus.

4. CAUCALIS nodosa.
TORDYLIUM nodosum. Linn. } Knotted stone Parsley.

* SEDUM rupestre. Rock Stonecrop.

RUBUS casius. Dewberry.

----- corylifolius. Hasel-leaved Bramble.

TRAGOPOGON pratensis. Yellow Goat's-beard.

ANTHEMIS arvensis. Corn Chamomile.

* CALENDULA arvensis. Marigold. (Hull.) (18) CAREX distans. Loose Carex.

SPERGULA nodosa. Knotted Spurrey.
 SISYMBRIUM amphibium. Great water Rocket.

290 LEMNA minor. Lesser Duck-weed. Com in January

6. * SALVIA verbenaca. Wild English Clary. Compon chalk SAMBUCUS nigra. Dwarf Elder.

MALVA sylvestris. Common Mallow. Talk muttified

in ray large - Rich 48 TO I HIM paranea. Par cross the faith horsen to heavy

7. LOLIUM perenne. Ray-grass. TRITICUM repens. Couch-grass.

* SAXIFRAGA stellaris. Hairy Saxifrage.

* SILENE acaulis. Moss Campion.

8. PLANTAGO media. Hoary Plantain. Pericont a Myring.

JUNCUS uliginosus. Little bulbous Rush.

* ANTIRRHINUM majus. Great Snapdragon. Rich 31, 4

. * VERONICA saxatilis. Blue rock Speedwell. Back Safe

* ----- alpina. Alpine Speedwell. A Boy on the self-

* POA alpina. Alpine Meadow-grass.

* STIPA pennata. Downy Feather-grass.

* CAMPANULA glomerata. Clustered Bell-flower.

rulesca com in ditakent bonts place cash in

(18) It is not uncommon for the flowers of this plant to become double; its multiplication arising from a conversion of the tubular into ligulate florets. It is worthy of remark that its fertility is increased in proportion to its multiplication. It is apt to become proliferous. K.

This Hower closes it's petals at noon are from there has her called go to her at noon - It's leaves are amplexical.

6 26,27

un roy way

stanen adortive

SIXTH MONTH. or Croporate Leaves to in a whole; leave * ATROPA Belladonna. Deadly Nightshade. Lander in Twing LONICERA Periclyminum. Common Honeysuckle, * ANGELICA Archangelica. Garden Angelica. abund on hunks * IMPERATORIA Ostruthium. Masterwort. RUMEX aquaticus. Great water Dock. ALISMA Plantago. Greater water Plantain. Point of attachment of) under water, trainer above the Surface ERICA Tetralix. Cross-leaved Heath. (19) ERICA botuliformis. Salisbury. * SEDUM anglicum. English Stonecrop. * OXALIS corniculata. Yellow procumbent Wood-Sorrel. ROSA arvensis. White Dog-rose. Repl 31 RUBUS fruticosus. Common Bramble. 10. FESTUCA elatior. Tall Fescue-grass. ELYMUS europæus. Wood Lyme-grass. EPILOBIUM palustre. Round-stalked marsh Willow-herb. GENISTA tinctoria. Dyer's Green-weed. SONCHUS oleraceus. Common Sow-thistle. CREPIS tectorum. Smooth Hawk's-beard. * LAGURUS ovatus. Hare's-tail-grass. ÆTHUSA Cynapium. Fool's-Parsley. * EPILOBIUM angustifolium. Rose bay Willow-herb. MELAMPYRUM pratense. Common yellow Cow-wheat. HIERACIUM paludosum. Succory-leaved mountain Hawk-weed. VALERIANA officinalis. Great wild Valerian. Maither the 40.12. LOLIUM arvense. White Darnel.

HYOSCYAMUS niger. Common Henbane. Teresarpea piece

CONIUM maculatum. Common Hemlock.

* ORNITHOGALUM pyrenaicum. Spiked Star of Bethlehem.

* DIANTHUS deltoides. Maiden Pink.

----- cæsius. Mountain Pink. & DIANTHUS virgineus. Linn. ARENARIA serpyllifolia. Thyme-leaved Sandwort.

(19) All the species of Erica (E. Dabeoci excepted, and this I presume, will scarcely be considered as belonging to the genus) not unfrequently vary to white.

other moniocot plants it has a vingle involvere 3 divis berry notes lord new 3 external and calveal yet the six dion, form but one ums circle on the recentable - Terbers Edeble de Burnett 424

incurrella in Bosi in heit of Enclase ten Scotlants

- 12. SPIRÆA Ulmaria. Meadow-sweet.

 LINNÆA borealis. Two-flowered Linnæa.

 MEDICAGO polymorpha. Heart Medic.

 CARDUUS pratensis. Meadow Thistle.
- 14. AGROSTIS Spica-venti. Silky Bent-grass.

 FESTUCA ovina. Sheep's Fescue-grass.

 BROMUS erectus. Upright perennial Brome-grass.

 POTAMOGETON lucens. Shining Pond-weed.

 RUPPIA maritima. Sea Ruppia.

 ROSA spinosissima. Burnet Rose.

 OROBANCHE major. Greater Broom-rape. The state of the sta
- 15. * POA flexuosa. Zigzag Meadow-grass. * ----- cæsia. Sea-green Meadow-grass.
 - * ---- nemoralis. Wood Meadow-grass.

 EPILOBIUM tetragonum. Square-stalked Willow-herb.

 BUTOMUS umbellatus. Flowering Rush.

* LYCHNIS Viscaria. Red German Catchfly.

* HEDYSARUM Onobrychis. Saintfoin. book officer jointed

* HIERACIUM auranticum. Orange Hawkweed.

16. CIRCÆA lutetiana. Common Enchanter's Nightshade.

Tentimenta alpina . Bop opolnohte plan and than non

(20) Sometimes, although rarely, varies to purple.

O. Minn in Clara fields - Lee M. Mat Hist V. 1. 300

ik 34

16. PHLEUM pratense. Common Cat's-tail-grass.

* FESTUCA rubra. Creeping Fescue-grass.

* ----- loliacea. Spiked Fescue-grass.

BROMUS secalinus. Smooth rye Brome-grass.

JASIONE montana. Sheep's Scabious.

APIUM graveolens. Wild Celery.

ROSA canina. Common Dog-rose. Richstx

* POTENTILLA fruticosa. Shrubby Cinquefoil.

MENTHA arvensis. Corn Mint.

PRUNELLA vulgaris. Self-heal.

HYPERICUM quadrangulum. Square St. John's Wort.

SENECIO Jacobea. Common Ragwort.

BRYONIA dioica. Red-berried Bryouv.

17. AIRA cristata. Crested Hair-grass.

POTAMOGETON perfoliatum. Perfoliate Pond-weed.

VICIA sepium. Bush Vetch.

GNAPHALIUM gallicum. } Narrow-leaved Cud-weed.

SENECIO sylvaticus. Mountain Groundsel.

13. BROMUS mollis. Soft Brome-grass.

GALIUM Mollugo. Great hedge Bed-straw.

JUNCUS articulatus. Jointed Rush.

ALISMA ranunculoides. Small water Plantain.

SEDUM acre. Biting Stonecrop.

RESEDA lutea. Wild Mignonette.

EUPHORBIA helioscopia. Sun Spurge.

CREPIS biennis. Rough Hawk's-beard.

HYOSERIS minima. Swine's Succory.

19. ARUNDO Calamagrostis. Small Reed.

ROSA rubiginosa. Sweetbriar.

* PAPAVER somniferum. White Poppy.

SCUTELLARIA galericulata. Common Skull-cap.

* GERANIUM pratense (flore albo). Crowfoot-leaved

TRIFOLIUM officinale.

TRIFOLIUM Melilotus officinalis. Linn. Common Melilot.

---- striatum. Soft knotted Trefoil.

of Min Darinatura monostrin garlic minut hack

* GNAPHALIUM luteo-album. Jersey Cud-weed.

LIGUSTRUM vulgare. Privet. 20.

* ARUNDO colorata. PHALARIS arundinacea. Linn. Canary Reed-grass.

GALIUM palustre. White-water Bed-straw.

LINUM usitatissimum. Common Flax.

----- catharticum. Purging Flax.

* SILENE Armeria. Lobel's Catch-fly.

EUPHORBIA exigua. Dwarf Spurge.

† ----- paralia. Sea Spurge.

THALICTRUM flavum. Common Meadow-rue.

RANUNCULUS Lingua. Great Spear-wort.

SCROPHULARIA aquatica. Water Fig-wort.

ERVUM hirsutum. Hairy Tare.

HYPERICUM androsamum. Tutsan St. John's Wort.

MATRICARIA Chamomilla, Wild Chamomile,

TYPHA angustifolia. Lesser Reed-mace.

HYDROCHARIS Morsus rana. Common Frog-bit.

*VERONICA fruticulosa. Flesh-coloured shrubby Speed-21. well. Ben Guachan Bentpourt.

SCIRPUS Holoschoenus. Round Cluster-headed Club-rush.

FESTUCA duriuscula. Hard Fescue-grass.

LOLIUM temulentum. Bearded Darnel.

LYSIMACHIA Nummularia. Creeping Loosestrife.

CAMPANULA rotundifolia. Round-leaved Bell-flower.

HERNIARIA glabra. Smooth Rupture-wort.

GERANIUM pratense. Crowfoot-leaved Crane's-bill.

HYPERICUM humifusum. Trailing St. John's-Wort.

* CIRCÆA alpina. Mountain Enchanter's Nightshade. 22. † ARUNDO arenaria. Sea Reed.

SEDUM album. White Stonecrop.

CHENOPODIUM urbicum. Upright Goosefoot.

POLYGONUM Convolvolus. Climbing Buckwheat. HYPERICUM pulchrum. Upright St. John's-Wort.

LAPSANA communis. Nipple-wort.

PYRETHRUM Parthenium. MATRICARIA Parthenium. Linn. Common Feverfew.

why shot by the The vier side between Now and Kings to alix and corolla court with sid granular shots - Its

ing are rooting ake appeare repens

24. VERONICA scutellata. Narrow-leaved Marsh Speedwell
GALIUM verum. Yellow Bed-straw.
ANTIRRHINUM Linaria. Common yellow Toadflax.
* HYPERICUM hirsutum. Hairy St. John's Wort.

PLANTAGO major. Greater Plantain.

* IMPATIENS Noli me tangere. Yellow Balsam.

SOLANUM nigrum. Common Nightshade. (21) Rich 27,20

* ERYNGIUM campestre. Field Eryngo.

SIUM latifolium. Broad-leaved Water Parsnep.

AGROSTEMMA Githago. Corn Cockle. Caly longer than Carolla

* DELPHINIUM Consolida. Field Larkspur.

* ADONIS autumnalis. Corn Adonis.

† & LOTUS corniculatus. Common Bird's-foot Trefoil CHRYSANTHEMUM segetum. Yellow Ox-eye. ATRIPLEX pedunculata. Pedunculated sea Orache.

26. HORDEUM pratense.
β HORDEUM murinum. Linn. } Meadow Barley.

CICUTA virosa. Water Hemlock.

TRIFOLIUM ochroleucum. Sulphur coloured Trefoil.
α CENTAUREA nigra. Lesser Knapweed.

27. BORAGO officinalis. Common Borage.

ENANTHE fistulosa. Common Water Dropwort.

CHLORA perfoliata.
GENTIANA perfoliata. Linn. Yellow Centaury. (22)
POLYGONUM Fagopyrum. Buckwheat.
MARRUBIUM vulgare. White Horehound.
HYPERICUM perforatum. Perforated St. John's Wort.

⁽²¹⁾ Too much pains cannot be taken to extirpate this poisonous weed, the berries of which have an appearance likely to tempt the appetites of children. The situations in which it is most frequently found near Warrington, are the gardens at the western end of the town and the road side at Lower Walton. K.

⁽²²⁾ This plant may be added to the list of those which are usually denominated aquinoctial; its expansion commencing at 7 o'clock A. M. and ending at 2 o'clock P. M. K.

49

the bare,

HORDEUM murinum. Wall Barley. SAMBUCUS Ebulus. Dwarf Elder. NARTHECIUM ossifragum. ANTHERICUM ossifragum. Linn. Lancashire Asphodel am abortine VERBENA officinalis. Common Vervain. SERRATULA arvensis. Linn. Creeping Thistle. CARDUUS arvensis. EUPATORIUM cannabinum. Hemp-agrimony. TYPHA latifolia. Broad-leaved Reed-mace. It bit is ap POLYGONUM Persicaria. Spotted Persicaria. 29. SEMPERVIVUM tectorum. Common Houseleek. * ACTEA spicata. Herb Christopher. TILIA europaa. Common Lime-tree. Its bracke adheres to the GALEOPSIS versicolor. Large-flowered & GALEOPSIS Tetrahit. Linn. GALEOPSIS cannabina. Willd. \ Hemp-nettle. step and tubrai † SISYMBRIUM Sophia. Flix-weed. 30. CHENOPODIUM nubrum. Red Goose-foot, White Goose-foot. --- album. & SAMBUCUS nigra. Common Elder. + PYROLA minor. Lesser Wintergreen. (23) Vitoma Vita

SEDUM reflexum. Yellow Stonecrop.

* POTENTILLA alba. White Cinquefoil.

* SIBTHORPIA europæa. Cornish Moneywort.

GNAPHALIUM germanicum. Common Cudweed. FILAGO germanica. Linn.

B CENTAUREA nigra. Lesser Knapweed.

SPARGANIUM simplex. Unbranched upright Bur-reed.

SEVENTH MONTH (July).

VERBASCUM Thapsus. Great Mullein. Mann James * LIGUSTICUM scoticum. Scottish Lovage. STACHYS arvensis. Corn Woundwort. + y ONONIS arrensis. Rest-harrow. of has dee ONONIS repens. Linn. Luceine for a percential

⁽²³⁾ This rare plant was brought to me from a rock near the waterfall at Ambleside, Westmoreland.

2. VALERIANA dentata. Oval-fruited Corn Sallad. Roll to

† ERYNGIUM maritimum. Sea Eryngo.

PHELLANDRIUM aquaticum. Water Hemloek.

† GLAUCIUM luteum.

CHELIDONIUM Glaucium. Linn. Spoppy.

† ANTHYLLIS vulneraria. Kidney Vetch.

† VICIA sylvatica. Wood Vetch.

* ------ Cracca. Tufted Vetch.

3. FESTUCA gigantea. BROMUS giganteus. Linn. } Tallest Fescue-grass.

AGRIMONIA cupatoria. Common Agrimony. Standay have alte STACHYS palustris. Marsh Woundwort.

a ONONIS arvensis. Rest-harrow.

ANAGALLIS tenella.
LYSIMACHIA tenella. Linn. Bog Pimpernel.
CHIRONIA Centaurium.
GENTIANA Centaurium. Linn. Common Centaury.

5. LYSIMACHIA vulgaris. Yellow Loosestrife.

† GERANIUM sylvaticum. Wood Crane's-bill.

MALVA moschata. Musk Mallow.

† HYPERICUM montanum. Mountain St. John's Wort.

6. † \$CIRCÆA ulpina. | Mountain Enchanter's CIRCÆA intermedia. Ehrhart. | Nightshade. | CONVOLVOLUS arvensis. Small Bindweed. | SIUM nodiflorum. Procumbent Water Parsnep. | PIMPINELLA saxifraga. Common Burnet-Saxifrage. | * OENOTHERA biennis. Biennial Oenothera. (E. Bot.)

BALLOTA nigra. Stinking Black Horehound.

LEONURUS Cardiaca. Motherwort.

* HYPERICUM calycinum. Great-flowered St. John's Wort. (E. Bor.)

BIDENS tripartita. Trifid Bur Marygold.

CENTAUREA Scabiosa. Greater Knapweed.

7. SCABIOSA arvensis. Field Scabious.

CAMPANULA latifolia. Giant Bell-flower.

Lyhida D 2 Graig Miedhis hank

Malaxis Paleidora grows on net hearth beau bulls on mayin of leaf the Red + May Wat

above it 1. 1.

Note & top 21 Sandarea Japellefolen - leaves Macherune EPILOBIUM roseum. Pale Smooth-leaved Willow-herb. TEUCRIUM Scorodonia. Wood Germander. BONONIS arvenis. B ONONIS spinosa. CARLINA vulgaris. Common Carline. * GNAPHALIUM margaritaceum. American Cudweed. ACHILLEA Ptarmica. Sneeze-wort.

> * CAMPANULA Trachelium. Nettle-leaved Bell-flower. 8. * CHENOPODIUM olidum. CHENOPODIUM Vulvaria. Linn. Stinking Goose-foot.

--- maritimum. Sea Goose-foot.

* SWERTIA perennis. Marsh Felwort.

SELINUM palustre. Marsh Milky Parsley.

CAUCALIS Anthriscus. TORDYLIUM Anthriscus. Linn. Upright Hedge Parsley JUNCUS bufonius. Toad Rush.

EPILOBIUM hirsutum. Great hairy Willow-herb. ARENARIA tenuifolia. Fine-leaved Sand-wort. * CARDUUS marianus. Milk Thistle.

MELICA carulea. Purple Melic-grass. AIRA carulea. Linn. GALEOPSIS Tetrahit. Common Hemp-nettle. ERVUM tetraspermum. Smooth Tare. CENTAUREA Calcitrapa. Common Star-thistle. + ORCHIS conopsea. Aromatic Orchis.

10. SAMOLUS Valerandi. Brookweed. + CONVOLVOLUS Soldanella. Sea Bindweed. + SALSOLA Kali. Prickly Saltwort. LYTHRUM Salicaria. Purple Loosestrife. (24) BETONICA officinalis. Wood Betony. + BUNIAS Cakile. Sea Rocket. CAKILE maritima. Willd. CICHORIUM Intybus. Wild Succery.

3 cel Colorecan rotered an work elect; t

⁽²⁴⁾ A singular variety of this is sometimes found, which differs in the foliafion, its leaves being alternate, ovate, and acuminate. This is not accidental but permanent,

SERAPIAS latifolia. SERAPIAS Helleborine. Linn. | Broad-leaved Helleborine EPIPACTIS latifolia. Swartz. SERAPIAS palustris. Y SERAPIAS longifolia. Linn. Marsh Helleborine. EPIPACTIS palustris. Swartz. + ATRIPLEX laciniata, Frosted sea Orache. * PHALARIS canariensis. Manured Canary-grass. EPILOBIUM parviflorum. Small-flowered & EPILOBIUM hirsutum. Linn. Willow-herb. * DIANTHUS Caryophyllus. Clove Pink. * SERRATULA alpina. Alpine Saw-wort. 12. * IRIS fatidissima. Stinking Iris. SCABIOSA succisa. Devil's-bit Scabious. JUNCUS bulbosus. Round-fruited Rush. * DIANTHUS prolifer. Proliferous Pink. STACHYS sylvatica. Hedge Woundwort. ASTER Tripolium. Sea Star-wort. Thanks trend all PYRETHRUM maritimum. Sea Feverfew. MATRICARIA maritimum. Linn. LYCOPUS europæus. Water Horehound. Banky of DIPSACUS sylvestris. Wild Teasel. Re a DIPSACUS fullonum. Linn. * CAMPANULA Rapunculus. Rampion Bell-flower. VERBASCUM Blattaria. Moth Mullein. SIUM anguistifolium. Narrow-leaved Water Parsnep. PIMPINELLA magna. Great Burnet-Saxifrage. * STRATIOTES aloides. Water Soldier. GNAPHALIUM minimum. Least Cudweed. * LYSIMACHIA thyrsiflora. Tufted Loosestrife. 14. ANETHUM Faniculum. Common Fennel. STATICE Limonium. Sea Lavender. DROSERA rotundifolia. Round-leaved Sun-dew. (25) --- longifolia. Long-leaved Sun-dew.

* he a curious article on Arojera in May. Nat 4 NO 32

⁽²⁵⁾ The following accurate description of the Phænomenon exhibited by this Plant, may be considered worthy of insertion in this place.

[&]quot; Ses seuilles sont bordées de poils terminés par une petite goutte de liqueur.

" Quand un insecte se pose sur le disque d'une seuille, elle se serme comme

14. † β GALEOPSIS Ladanum. Red Hemp-nettle.

* LAVATERA arborea. Sea Tree-mallow.

HYPERICUM elodes. Marsh St. John's Wort.

HEDYPNOIS autumnalis.

LEONTODON autumnalis. Linn.

APARGIA autumnalis. Willd.

* INULA Helenium. Elecampane.

- 15. SAXIFRAGA Hirculus. Yellow Marsh Saxifrag'z. (26)
 BARTSIA viscosa. Yellow viscid Bartsia. (27)
 ARTEMISIA vulgaris. Mugwort.
 † CONYZA squarrosa. Plowman's Spilenard.
- 16. POA procumbens. Procumbent sea Meadow-grass,
 † DIPSACUS pilosus. Small Teasel.
 * NEPETA Cataria. Cat-mint.
 SONCHUS palustris. Tall marsh Sow-thistle.
 TANACETUM vulgare. Common Tansy.

une bourse à jetons dont on tire les cordons, et l'insecte meurt, tout couvert du suc visqueux qui s'echappe de l'extremité des poils. C'est un spectacle aussi bizarre qu'il est etonnant, de voir, comme j'ai eu plusieurs fois occasion de l'observer, presque toutes les feuilles de cette petite plante formant des prisons enduites de glu, où sont retenus quelques insectes, tels que des fourmis ou des petites mouches. Ainsi la rencontre d'un gazon de rossolis est aussi terrible, pour ces légions de créatures foibles et miscrables, que l'explosion d'un feu souterrain pour les peuplades humaines; mais ces desordres ont leur limites; et, tandis que les individus sont en proie à mille dangers, les espéces ne cessent pas de se propager, et ces innombrables races d'êtres ephémeres ne sont pas moins que la race humaine sous la main protectrice de la Nature."

MIRBEL Traite d'Anatomie
et de Physiologie Vegetales
Tom. 1. p. 265.

- (26) Observed upon Knutsford moor; the habitat already pointed out.
- (27) This rare plant has been found at Ormskirk, South-port, Allerton, Speke, Hale, and Ditton in Lancashire, and on Bartington heath in Cheshire, by different members of the Botanical Society of Warrington.

17. † PARNASSIA palustris. Grass of Parnassus. HUMULUS Lupulus. Common Hop.

18. SEDUM Telephium. Orpine.
* SPIRÆA salicifolia. Willow-leaved Spiræa.
GALEOPSIS villosa. Downy Hemp-nettle.
α ONONIS arvensis (flore albo). Rest-harrow. (28)
INULA dysenterica. Common Flea-bane. (29)
ATRIPLEX patula. Spreading halberd-leaved Orache.

19. GENTIANA Pneumonanthe. Marsh Gentian. (30)

ERICA vulgaris.
CALLUNA vulgaris. Salisbury. Common Heath.

BARTSIA Odontites.
EUPHRASIA Odontites. Linn. Red Eyebright.

PICRIS hieracioides. Hawkweed Ox-tongue. Pappen simple not fraction.

LACTUCA Scariola. Prickly Lettuce. Pappen simple not fraction.

ARTEMISIA maritima. Sea Wormwood.

20. BROMUS sylvaticas. Slender wood Brome-grass,

* HERNIARIA hirsuta. Hairy Rupture-wort.

* ERICA vagans.
ERICA vaga. Salisbury.

POLYGONUM lapathifolium. Pale-flowered Persicaria.

ANTHEMIS maritima. Sea Chamomile.

21. * VERBASCUM Lychnitis. White Mullein.

MENTHA sylvestris. Horse Mint.

(28) This variety, differing from a Ononis arcensis, not only in the whiteness of its petals, but in the colour of the stalk which is of a pale green instead of purple, is found near the pier, at Runcorn. K.

⁽²⁹⁾ This plant is said to have been occasionally useful in cases of Dysentery. As this disease, after an interval of more than 100 years, has of late appeared among us, it may perhaps, as a cheap remedy, be worthy of trial. It has an unpleasant smell, compared by some to that of soap; and a bitterish subacid taste. Its acrimony a little resembles that of Arum maculatum. K.

⁽³⁰⁾ A white-flowering variety of this plant, is said to have been found upon Bartington heath.

- 21. SOLIDAGO Virgaurea. Common Golden-rod. (31)
 - 22. * CORIANDRUM sativum. Common Coriander.

 *TEUCRIUM Chamædrys. Common Germander.

 CINERARIA palustris. Marsh Flea-wort.

23. POLYGONUM Hydropiper. Biting Persicaria.

24. *SALVIA pratensis. Meadow Clary. a rug rate flam

* DIPSACUS fullonum. Manured Teasel.

DROSERA anglica. Great Sun-dew.

ANDROMEDA polifolia. Marsh Rosemary. (Second inflorescence.) (32)

ARCTIUM Lappa. Burdock.

- 25. SENECIO viscosus. Stinking Groundsel.
- 26. SALICORNIA herbacea. Marsh Samphire. ERIGERON acre. Blue Flea-bane.
- 27. CAUCALIS infesta.

 SCANDIX infesta. Linn. Spreading hedge Parsley.

 SENECIO tenuifolius. Hoary Ragwort.
- 28. ANGELICA sylvestris. Wild Angelica. ANTHEMIS nobilis. Common Chamomile.
- 29. CHIRONIA Centaurium. (flore albo) Common Centaury. (33)

 MENTHA hirsuta. Hairy Mint.

 LATHYRUS palustris. Marsh Lathyrus.
- 30. MENTHA rotundifolia. Round-leaved Mint. HYPOCHÆRIS radicata. Long-rooted Cat's-ear.
- 31. GENTIANA Amarella. Autumnal Gentian. STACHYS germanica. Downy Woundwort.

(31) The powdered Root, in doses of half a dram, three times a day, has been found a powerful remedy in *Dropsy*. It has long been in use in cases of *Nephritis*, and with good effect. K.

⁽³²⁾ Repeated observation confirms the opinion that this species has two distinct periods of inflorescence in each year.

⁽³³⁾ This variety has been found in several parts of this neighbourhood. K.

31. HIERACIUM sabaudum. Shrubby broad-leaved Hawkweed.

URTICA pilulifera. Roman Nettle.

To region Canadense: Lineaus thought their plant was brught from show and lang EIGHTH MONTH (August). all the plants of Europe

- * COLCHICUM autimizate. Mendow Soffian SCUTELLARIA minor. Lesser Skull-cap. 1.
- GNAPHALIUM uliginosum. Marsh Cudweed. 2.
- * MENTHA viridis. Spear Mint. 3.
- GNAPHALIUM rectum. Upright Cudweed. 4.
- BIDENS cernua. Nodding Bur Marygold. 5.
- POLYGONUM minus. Small creeping Persicaria. 6.
- 7. CLINOPODIUM vulgare. Wild Basil. LEONTODON palustre. LEONTODON lividus. Willd. Marsh Dandelion.
- ARUNDO Phragmites. Common Reed. 8.
- MENTHA piperita. Pepper Mint.
- 10. ---- rubra. Tall red Mint.
- * ORIGANUM vulgare. Common Marjoram. 11.
- ARUNDO epijegos. Wood Reed. 12.
- 13. MENTHA odorata. Bergamot Mint.
- UTRICULARIA vulgaris. Hooded Milfoil. Itagh! distaher 14.
- * SENECIO sarracenicus. Broad-leaved Groundsel. 15. ULEX nanus. Dwarf Furze. β ULEX europæus. Linn.
- 16. MENTHA Pulegium. Penny Royal.
- SAPONARIA officinalis. Soapwort. 17.
- CHIRONIA pulchella. Dwarf branched Centaury. 19.
- CHENOPODIUM hybridum. Maple leaved Goose-foot. 20. * HIERACIUM umbellatum. Narrow-leaved Hawkweed. + SERRATULA tinctoria. Common Saw-wort.
- 24. OPHRYS spiralis. NEOTTIA spiralis. Swartz. } Spiral Ophrys.
- y BIDENS cernua. BIDENS minima. Linn. | Nodding Bur Marygold.

Unicularea internersa - near butter trest of a minor Justiche Westerbir lake Ferland

- 27. CORNUS sanguinea. Wild Cornel-tree (second inflorescence). (34)
- 28. SCLERANTHUS perennis. Perennial Knawel.

NINTH MONTH (September).

- 8. * COLCHICUM autumnale. Meadow Saffron.
- 10. * ARBUTUS Unedo. Strawberry-tree.
- 18. * CROCUS sativus. Saffron Crocus.

TENTH MONTH (October).

13. HEDERA Helir. Common Ivy. Roll 3/51

(34) The time here mentioned was evidently the second period of inflorescence, but whether it is permanently the case as in Andromeda I do not know.

12. ARUNDO spiegos. Wood Road.

UTRICULARIA sulgaria. Hooded Milfoil. ... SENECTO carracences. Broad-leaved Groundsell.

& ULLEX encopance Linn. | Dwaif Force.

MENTHA Polegium. Penny Royal Comment.

CHENOPODIUM hybridami. Maple leaved Goose-foot.

*HIERACIUM umbellatum, Narrow-leaved Hawkweed.

OPHRYS spindie.
NEOTTIA spindis. Swartz. } Spiral Ophrys.

y BIDENS cornua. Linn. | Nodding Bur Marygold.

INDEX GENERUM.

The Genera printed in Italics occur only as synonyms.

01 - 1			A DECEMBER OF THE PERSON OF TH
West of the second		Page.	Page.
54	A	ally parts	Angelica - 21, 32.
ACER	-	- 10.	Anthemis - 15, 20, 31, 32.
Achillea		19, 28.	Anthericum - 18, 26.
Acorus -		- 18.	
Actæa	- Charles	26.	Anthyllis 27.
Adonis -		- 25.	
Adoxa	- 12	1.	Apargia - 16, 30.
Ægopodium		- 15.	Aphanes - 7.
Æthusa		19, 21.	Apium 23.
Agrimonia		- 27.	Aquilegia - 13.
Agrostemma		- 25.	Arabis 7, 8.
Agrostis		16, 22.	Arbutus - 11, 34.
Aira -		17, 23, 28.	Arctium - 32.
Ajuga	-	- 13.	Arenaria 12, 14, 19, 21, 28.
Alchemilla	1/00	7, 8, 17.	Aristolochia - 15.
Alisma	14 4	21, 23.	Artemisia - 30, 31.
Allium -	4	13, 15, 17.	Arum 14, 19.
Alnus -	1	- 6.	Arundo - 23, 24, 33.
Alopecurus	-	14, 16, 18.	{ Asarum - 9.
Alsine	-	- 6.	Allman of the ball
Alyssum	-	- 19.	
Anagallis		- 18, 27.	
Anchusa	-	- 10.	
Andromeda			Atriplex - 25, 29, 31.
Anemone			{ Atropa 21.
Anethum	-	- 29.	Avena - 14, 17, 18.

B. Page.	C. Pari
Ballota - 27.	Cochlearia - 7, 9, 13.
Bartsia 30, 31.	Colchicum - 34.
Bellis - 5	Comarum 18.
Berberis - 14.	Conium - 21.
Betonica - 28	Convallaria - 13, 15.
Betula - 1 6, 9.	Convolvolus 27, 28.
Bidens 27, 33	Conyza 30.
Borago 05	Coriandrum - 32.
Brassica - 9, 17.	Cornus - 15, 34.
Briza 17, 10	Coronopus - 13.
Bromus - 22, 23, 27, 31	Corylus - 6.
Bryonia 23	Cotyledon - 18.
Bunias 28.	Crambe - 15.
Bunium - 11, 15.	Cratægus - 11, 15.
Butomus 22	Crepis - 21, 23,
Buxus - 8.	Crocus - 5, 34
C	Cucubalus - 19, 22.
Cakile 28.	Cynoglossum - 17-
Calendula - 20.	Cynosurus - 18.
Callitriche - 8, 34.	D
Caltha 7.	Dactylis 16.
Calluna - 31.	Daphne 5, 6.
Campanula 20, 24, 27, 28, 29.	Datura 25.
Cardamine - 6, 8, 10.	Daucus 18.
Carduus - 19, 22, 26, 28.	Delphinium - 25.
Carex 10,11,13,14,15,16,20.	Dentaria 8
Carlina - 28.	Dianthus 21, 29.
Carpinus - 8.	Digitalis - 18.
Caucalis - 20, 28, 32.	Dipsacus - 29, 30, 32.
Centaurea 12,25,26,27,28.	Doronicum - 14.
Cerastium - 7, 9, 10, 11.	Draba - 6.
Cherophyllum - 9, 18.	Drosera - 29, 32.
Chara 7, 9.	E - AMA
Cheiranthus 7.	Echium 8.
Chelidonium - 13, 27.	Elymus - 21.
Chenopodium 11,24,26, 28, 33.	Epilobium > 21, 22, 28, 29.
Chironia - 27, 32, 33.	Epimedium - 9.
Chlora - 25.	Epipactis 17, 29.
Chrysanthemum' - 19, 25.	Erica - 10, 21, 22, 31.
Chrysoplenium - 7. 28.	Erigeron 32.
The state of the s	Eriophorum - 9.
	Erodium 10, 18.
The state of the s	Ervum 24,28.
The state of the s	
NO. OF THE RESERVE OF THE PARTY	Erysimum - 9, 13, 14.
- 53.)	Euonymus - 18.

Page.	Page.
Eupatorium 26.	Hyoseris - 28.
Euphorbia - 9, 14, 23, 24.	Hypericum 23, 24, 25, 27, 30.
Euphrasia - 22, 31.	Hypochæris - 32.
Potamoveton - H. 17, 18,00 on	adjordenialis I- 192
Fagus - 10, 20.	Iberis - 10, 12.
Festuca 16, 21, 22, 23, 24, 27.	Ilex 115.
Filago - 23, 26.	Impatiens - 25.
Fragaria 6, 9.	Imperatoria - 21.
Fraxinus - 12.	Inula 30, 31.
Fritillaria Minamond 8.	Iris - 16, 29.
Fumaria - 8, 10, 12, 13, 17.	Ixia - 9.
G slory	J- zuzzinaki
Galanthus 5.	Jasione 23.
Galeobdolon - 12.	Juneus 7,12,16,19,20,23,28,29.
Galeopsis 12,22,26,28,30,31.	Juniperus 115.
Galium 8,14, 18, 23, 24, 25.	. Nymphaa L 18, 15.
Genista 13, 21.	Lactuca - 0 - 31.
Gentiana - 10,25,27,31,32.	Lagurus odio 21.
Geranium 10, 12, 13, 15, 17, 18,	Lamium 5, 7.
23,24,27.	Lapsana 24.
Geum 11, 15.	Lathyrus - 14, 20, 22, 32
Glaucium 27.	Lavatera 30.
Glaux 1010012.	Lemna 20.
Glechoma di7.	Leontodon - 6, 16, 30, 33.
Gnaphalium 12,23,24,26,28,	Leonurus - 12, 27.
19 02 01 41 29,33. audull	Leucojum 10.
Rumes 12 H 15, 18, 20, 21.	Ligusticum - 26.
Hedera 34.	Ligustrum 24.
Hedypnois - 16,30.	Linnæa - 22.
Hedysarum - 22.	Linum 24.
Helleborus - 5, 8.	Lithospermum - 17, 19.
Helonias - 18.	
Heracleum - 17.	Lonicera - 12, 15, 21.
Herniaria - 24, 31.	Lotus 13, 25.
Hesperis 18.	Lychnis - 10, 16, 22.
Hieracium 14, 17, 21, 22, 33.	
Hippuris - 10.	
Holcus - 14, 16.	Lysimachia - 14,24,27,29.
Holosteum - 10.	
Hordeum - 25, 26.	M allendaring
Hottonia - 15.	
Humulus - 31.	Marrubium - 25.
Hyascinthus - 11.	Matricaria - 24, 29.
Hydrocharis - 24.	Medicago 22.
Hydrocotyle - 14, 17.	Melampyrum - 21.
Hyoscyamus - 21	. Melica - 18, 28.

Page.	Page.
Montha - 23, 31, 32, 33.	Polygonum 16, 19, 22, 24, 25,
Menyanthes 9 29 -months 121	19 12 41 26, 31, 32, 33, 44
Menziesia 10.	Populus - 6, 7, 8.
Mercurialis - 7, 32.	Potamogeton 16, 17, 19, 22, 23.
Mespilus - 11, 17.	Potentilla - 12, 15, 23, 26.
Meum 19.	Prenanthes 22.
Montia 7.	Primula - 5, 7, 8.
Myagrum 19.	Prunella 231
Myosotis 12, 13, 14.	Prunus 8, 9, 12.
Myrica 8.	Pulmonaria smellini 6.
N w minimaged	Pyrethrum 01 - 24, 29.
Narcisens - 6.9.	Pyrola - 26.
Narcissus 6, 9. Narthecium - 26.	Pyrus - 9, 11, 12, 15.
Neottia 200-11.01.04.7 abou 33.	Calcobdolen Q - P
Nepeta	Quercus - 13.
Nymphæa - 18, 19.	Calings 8 A 18,23,24,25
Nymphaa O - complete	Ranunculus 6,7,9,12,13,15,
Enanthe - 17, 25.	
Oenothera - 27.	Raphanus et 1- 01 -minor 17.
Ononis - 26, 27, 28, 31.	Reseda
Ophrys 17, 33.	
Orchis 10, 16, 17, 18, 19, 22, 28.	
Origanum - 33.	Rhodiola 15.
Ornithogalum - 8, 10, 21.	Ribes 7, 8, 9.
Ornithopus - 13.	Rosa 22, 21, 22, 23,
Orobanche - 22.	Rubus - 13, 19, 20, 21.
	Rumex 12, 13, 15, 19, 20, 21.
Orobus 6, 21.	1
P - pommit	Ruscus - Fingybi S.
100 00 00 00	
THE RESIDENCE OF THE PARTY OF T	2.2.
THE RESERVE OF THE PARTY OF THE	Salicornia 32.
Larm	Salix - 6, 9, 10, 11.
· · · · · · · · · · · · · · · · · · ·	200 1
rememans	Salvia - 20, 32.
Tepan	00 00
The state of the s	20 1 00
I licitation tunt	177
I meum	10
FICTIS .	00
T IIII Piliteria	1 10 17 00 90
1 mg mcuts	Scabiosa - 27, 29,
Plantago - 9, 16, 19, 20, 25.	1 1 1 00
Poa 6, 14, 16, 18, 20, 22, 27, 30.	Scheenus - 20.
Potemomani	
Polygala - 11.	TO signormovida
THE RESERVE OF THE PARTY OF THE	The state of the s

Page. \$	Page.
	Taxus - 7.
Scirpus 13, 14, 15, 16, 17, 24.	Teucrium - 28, 32.
Scleranthus - 13, 34.	Thalictrum - 22, 24.
Scrophularia - 8, 17, 24.	Thlaspi - 7, 8, 9, 13.
Scutellaria - 23, 33.	Thymus 16.
Sedum 20, 21, 23, 24, 26, 31.	Tilia 26.
Selinum 28.	Tofieldia 18.
Sempervivum - 26.	Tollerata
Senecio - 5, 16, 23, 32, 33.	Tordylium - 20, 28. Tormentilla - 12.
Serapias - 29.	10rmentina
Serratula - 26, 29, 33.	2000
Sherardia - 12.	Trifolium 13, 15, 22, 23, 25. Triglochin - 12, 17.
Sibthorpia - 26.	00
Silene - 19, 20, 22, 24.	Triticum
Sinapis 10.	Tromus
Sison 17.	Tunpa
Sisymbrium - 14, 20, 20.	Luities
Sium - 25, 27, 29.	1 4301145
Solanum - 18, 25.	Typha - 24, 20.
Solidago 32.	1 2 00
Sonchus - 19, 21, 30.	(0101
Sorbus 12.	Ulmus 6, 7.
Sparganium - 26.	Urtica - 13, 14, 33.
Spartium 14.	Utricularia - 33.
Spergula - 16, 20.	} V
Spiræa - 17, 22, 31.	Vaccinium - 10, 14, 15.
Stachys - 26, 27, 29, 32.	Valantia - 14.
Staphylea 14.	Valeriana - 9, 12, 13, 21, 21.
Statice - 12, 29.	Verbascum - 26, 29, 31.
Stellaria - 6, 9, 14.	Verbena - 26.
Stipa 20.	Veronica 6,7,9,10,11,12,14,
Stratiotes - 29.	
Swertia 28.	
Symphytum - 15.	The second secon
T on mine	{ Vinca 5, 8.
Tamus - 16.	
Tanacetum - 30.	. } Viscum 8.

THE END.

J. Haddock, Printer, Horse-Market, Warrington.

	THE RESERVE OF THE PARTY OF THE
100	Page.
Taxus	
Tenerium - 28, 32,	lemathus - 13, 34 5
The intrum - 22, 24.	rophularia - 8, 17, 24, 9
Thlaspi 7, 8, 9, 13,	atellaria - 23, 83.
Thymus 16.	dum 20, 21, 23, 24, 26, 81.
Tilia 20.	189 281
Tofieldia 18.	mpeyvirum 20.
Lordylium - 190, 28.	archo 5,16, 23, 32, 35. [
Tormentilla - 12.	rapins 20.
Tragopogon - 17, 20.	ratula 20, 29, 33.
Trifolium 13, 15, 22, 23, 25.	écurdia - 12.
Triglochin c 12, 17.	nigrochia 20.
Triticum 20.	me - , 19, 20, 22, 24.
Trollies - 10.	tapis 10.
Tulipa.	17.7
Of amount	cymbrium - 14, 20, 26,
Tussingo 5.8.	im 25,27,39
Typha - 24, 26.	lamma - 18,25.
U	lidago 32.
Ulex = 5, 38.	nches - 19, 21, 30.
Ulmus - 6,7.	rbus 12.
Urtica" - 13, 14, 33.	arganium - 26.
Urricularia - 33.	14.
7	ergula : 16, 20,
(Vaccinium - 10, 14, 15,	iran - 17, 22, 31,
	schys 26, 27, 29, 32.
Valeriana - 0,12,13,21,27,-	
Verbaseum : 26120, 31	
Verbena - 26.	
Veronica 6,7,9,10,11,12,14,	ipa
The state of the s	rationes - 20.
Vibaruum 17,	vertia - 28.
	mighytam - 15.
Vioca 5,8.	T
Viola - 5, 6, 7; 8, 11;	.81 - 1 semi
Viscuin	the state of the s
A STATE OF THE PARTY OF THE PAR	A CONTRACTOR OF THE PARTY OF TH

THE END.

J. Haldock, Printer, Holes-Market, Washington