

On the coagulation of the blood / by L.C. Wooldridge.

Contributors

Wooldridge, Leonard Charles, 1857-1889.
Royal College of Surgeons of England

Publication/Creation

[London] : [Physiological Society], [1883]

Persistent URL

<https://wellcomecollection.org/works/kaddeynx>

Provider

Royal College of Surgeons

License and attribution

This material has been provided by This material has been provided by The Royal College of Surgeons of England. The original may be consulted at The Royal College of Surgeons of England. where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

ON THE COAGULATION OF THE BLOOD. BY L. C. WOOLDRIDGE, D.Sc., M.B., *George Henry Lewes Student*.

THE following communication is a continuation of the results already published by myself as to the influence of lecithin in producing coagulation of the blood¹.

The plasma used in those experiments was peptone plasma. In the present case I have made use of blood which has been prevented from coagulating by being, immediately after leaving the body, cooled down to a temperature of about 0°.

For experiments on cooled plasma it is best to use the blood of the horse. I was however unable to obtain any horse's blood and therefore employed dog's blood. In horse's blood the corpuscles sink rapidly and the coagulation is very tardy, it is hence easy to obtain plasma. This is not the case with the dog, and I was therefore led to adopt a particular method of experimenting.

The coagulation of the blood is brought about by a certain interaction of the white corpuscles and the plasma. By rapid cooling this interaction is to a great extent suppressed and hence the blood does not coagulate. But if one of the substances of which the white cells are made up be diffused through the cooled blood, coagulation does occur. This substance is lecithin.

The method of experimenting is as follows. The blood is taken from a large artery and flows into a thin cylindrical metal vessel of about $\frac{1}{2}$ -inch diameter. This tube stands in a large vessel filled with broken ice and a small quantity of water to fill up the interspaces.

For each experiment two such tubes are used. Each holds 40 c.c. In the one is contained 15 c.c. of .6 per cent. NaCl solution, in the other a similar quantity of normal salt solution through which finely emulsified lecithin is diffused.

The blood flows directly from the artery into the metal vessels, one

¹ "Further Observations on the Coagulation of Blood." *Journal of Physiology*, Vol. iv. Nos. 2 & 3. "Zur Gerinnung des Blutes." *Archiv für Anat. u. Physiol.* 1883.

vessel being filled immediately after the other. Absolutely no interruption to the flow must occur. The first small quantity of blood is not used.

The following examples will shew clearly the nature of the results.

I. Large Dog. Blood from Femoral.

Tube 1 contains lecithin.

Tube 2 contains simple salt solution.

By means of glass rod mixing is carried out in each tube, a separate rod being used for each tube.

Tubes filled at 4.15 P.M.

At 4h. 45m., as is ascertained by feeling with glass rod in tube 1 (lecithin), the blood is firmly coagulated.

At this time the blood in tube 2 is perfectly fluid. The temperature in the two tubes is the same, 2° at upper, 3° at lower part of tube.

On removing tube 1 from ice and inverting, a solid clot, forming a complete cast of the tube, slips out.

At 5h. 30m., blood in tube 2, on trying with glass rod is evidently completely fluid; on taking it out of ice and inverting the blood flows out; it is perfectly free from coagula; there are no traces of coagula on the walls of the tube.

II. Similar arrangements as in I. Large Dog. Blood from Carotid.

Tube 1, lecithin.

Tube 2, simple salt solution.

Tubes filled at 4h. 3m. P.M.

At 4h. 10m. tube 1 completely coagulated.

„ tube 2 completely fluid.

Temperature in the two tubes equal, 2.5° in middle of tube.

At 4h. 45m. tube 2 still quite fluid.

At 5h. imperfectly coagulated.

III. Arrangement the same. The lecithin is badly emulsified and in flocculent pieces which tend to rise to the surface of the NaCl solution.

Time of filling the two tubes 4.50 P.M.

At 5.30 P.M., tube 1 (lecithin), several loose coagula can be fished out. Tube 2 apparently quite free. Temp. tube 1, 2.5° bottom, 1.5° top.

„ tube 2, 4° bottom, 2½° top.

On emptying tube 1 it is found to contain loose clots and fluid blood, it is coagulated firmly at the top so that it can be inverted. The walls of the tube covered with clot.

At 6 P.M. tube 2 emptied. The contained blood is perfectly fluid. No trace of coagulation on the walls of the tube.

I have made many such experiments and always with the like result. That is to say, the addition of lecithin causes coagulation.

The lecithin used is prepared from lymph-glands in the manner I have described in the above-quoted papers. It has a slightly acid reaction and is not perfectly pure. It is rubbed up with a drop or two of dilute Na_2CO_3 solution to a paste and this is diffused through the salt solution. The salt solution after the addition of the lecithin is either neutral or very faintly alkaline. It should be an emulsion as opaque as milk if the coagulation is to occur quickly.

In my experiments on peptone plasma I have shewn that the small quantities of impurities contained in the lecithin preparations are of no effect in producing coagulation and also that lecithin from other sources is active, and hence I have felt justified in speaking of the alcohol-ether extract of the glands as lecithin, though it is not perfectly pure. From its mode of preparation it is evident that this extract cannot contain any paraglobulin.

In discussing the results I have obtained on peptone plasma with friends whose opinions are of worth, there has always been a tendency to suppose that the results obtained are due to presence of fibrin ferment in the lecithin preparations. In fact this is, so far as I can see, the only objection one can raise. It is however certainly not a valid objection. The lecithin is not influenced in its activity by being boiled with water. I have tried this several times both with peptone and cooled blood, and as a matter of fact it has no fermentative activity.

There are many coagulable fluids which do not coagulate in the slightest on the addition of lecithin but do so readily with fibrin ferment, although the lecithin in question acts perfectly with plasma. Such fluids are human pericardial fluid, hydrocele fluid so far as I have examined, and solution of fibrinogen prepared according to the salt method from peptone plasma.

The experiments described were carried out in the Pathological Institute in Berlin. I am much indebted to Prof. Salkowski for the use of his Laboratory.

Dec. 6, 1883.

1. The first of these is the fact that the American Medical Association is a voluntary association of physicians and surgeons. It is not a government agency, nor is it a part of the government. It is a private organization, and its members are free to join or leave it at will. This is one of the reasons why it is able to maintain its independence and to act in the best interests of the medical profession as a whole.

2. The second reason is that the American Medical Association is a national organization. It represents the interests of physicians and surgeons throughout the United States. This gives it a broad perspective and the ability to act in the best interests of the entire medical profession, rather than just a local or regional group.

3. The third reason is that the American Medical Association is a professional organization. Its members are all physicians and surgeons, and they are all committed to the highest standards of medical practice. This gives the organization a high level of credibility and respectability, and it enables it to act in the best interests of the public as well as the medical profession.

4. The fourth reason is that the American Medical Association is a non-profit organization. It does not have any financial interests, and its only purpose is to serve the interests of the medical profession and the public. This gives it a high level of integrity and trustworthiness, and it enables it to act in the best interests of the medical profession and the public without any ulterior motives.

5. The fifth reason is that the American Medical Association is a democratic organization. Its members are all equal, and they all have the right to vote in the election of the organization's leaders. This gives the organization a high level of accountability and transparency, and it enables it to act in the best interests of the medical profession and the public without any undue influence.

6. The sixth reason is that the American Medical Association is a cooperative organization. Its members all work together to achieve the organization's goals, and they all share in the organization's success. This gives the organization a high level of unity and solidarity, and it enables it to act in the best interests of the medical profession and the public with a high level of effectiveness.

7. The seventh reason is that the American Medical Association is a progressive organization. It is always looking for ways to improve the medical profession and the public's health, and it is always willing to embrace new ideas and technologies. This gives the organization a high level of dynamism and innovation, and it enables it to act in the best interests of the medical profession and the public with a high level of effectiveness.

8. The eighth reason is that the American Medical Association is a patriotic organization. It is committed to the United States and its values, and it is always willing to act in the best interests of the country. This gives the organization a high level of loyalty and dedication, and it enables it to act in the best interests of the medical profession and the public with a high level of effectiveness.

9. The ninth reason is that the American Medical Association is a humanitarian organization. It is committed to the well-being of all people, and it is always willing to act in the best interests of the most vulnerable members of society. This gives the organization a high level of compassion and empathy, and it enables it to act in the best interests of the medical profession and the public with a high level of effectiveness.

10. The tenth reason is that the American Medical Association is a responsible organization. It is always aware of its obligations to the medical profession and the public, and it is always willing to act in the best interests of both. This gives the organization a high level of responsibility and accountability, and it enables it to act in the best interests of the medical profession and the public with a high level of effectiveness.