Catalogue of North American mammals, chiefly in the museum of the Smithsonian Institution / by Spencer F. Baird.

Contributors

Baird, Spencer Fullerton, 1823-1888. Royal College of Surgeons of England

Publication/Creation

Washington : Smithsonian Institution, 1857.

Persistent URL

https://wellcomecollection.org/works/uv98yjf6

Provider

Royal College of Surgeons

License and attribution

This material has been provided by This material has been provided by The Royal College of Surgeons of England. The original may be consulted at The Royal College of Surgeons of England. where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org

CATALOGUE

OF

Miscella

NORTH AMERICAN MAMMALS,

CHIEFLY IN THE MUSEUM OF THE

SMITHSONIAN INSTITUTION.

BY

SPENCER F. BAIRD,

ASSSISTANT SECRETARY OF THE SMITHSONIAN INSTITUTION.



PRESENTED-BY

WASHINGTON: -SMITHSONIAN INSTITUTION. JULY, 1857.

Digitized by the Internet Archive in 2016

https://archive.org/details/b22297121

Order.	Page.	Family.	Page.	Sub-family.	Page.	Genus.	Page.	Sub-genus.	Page.	Species examined and identified.	Species not identified.
RAPACIA.											1
1. Sub-order.											
sectivora	2	1. Soricidae	7	Soricinae	7	1. Neosorex	* 11			1	
						2. Sorex	13			12	
		And the second second		· · · · · · · · · · · ·	41	3. Blarina				7	
		2. Talpidae	57	•••••	••••••	4. Scalops	58	Scalops		2	١.
	1.11					f. Can Johns	71	Scapanus		2	
2. Sub-order.		1.1.1				 Condylura Urotrichus 	76	•••••		1	
Carnivora	78	3. Felidae	80			7. Felis	81			5	
						8. Lynx	89			4	
		4. Canidae	102	Lupinae	103	9. Canis	104			2	
		100000000000000000000000000000000000000		Vulpinae	121	10. Vulpes	191	Vulpes	121	4	
								Urocyon	121	2	
		5. Viverridae	146			11. Bassaris	147			1	
		6. Mustelidae	148	Martinae	148	12. Mustela	149			2	1.
						13. Putorious	159	••••••		10	
				Lutrinae	183	14. Guið 15. Lutra	181			1 2	-
			1	Lutrinae	100	16. Enhydra	189			1	:
				Melinae	190	17. Mephitis	191	Thiosmus	191	1	1
		1.		*				Mephitis		4	
						18. Taxidea	201			2	
		7. Ursidae	206			19. Procyon	207			2	
						20. Ursus	216			3	
II. MARSUPIATA		8. Didelphidae	231			21. Didelphys	232			2	
III. RODENTIA	235	9. Sciuridae	240	Sciurinae	240	22. Sciurus	243	••••••		12	
						23. Pteromys	286	••••••		4	
*						24. Tamias	291 304	Otospermophilus		4 5	
						25. Spermophilus	501	Colobotis	305	9	
						26. Cynomys	329			2	
						27. Arctomys				2	
				Myoxinae	350						
				Castorinae	350	28. Aplodontia	350			1	
						29. Castor	355			1	
		Contract of the second	100	NEW REPORT	1	30. Castoroides	363	••••••		1	
		10. Saccomyidae	365	Geomyinae	366	31. Geomys	368	••••••		5	1.1
				Saccomyinae	405	32. Thomomys	388			7	
				estecomymac	100	 33. Dipodomys 34. Perognathus 	1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.	Perognathus		4	4.
						on recomments		Cricetodipus	10000	2	1
		11. Muridae	427	Dipodinae	428	35. Jacutus	429			1	
				Murinae	436	36. Mus	436			4	
						37. Reithrodon	447			4	
						38. Hesperomys	453	Hesperomys		13	
					8			Onychomys		1	
						20 Nantown	400	Oryzomys		1	1
						 Neotoma Sigmodon 				6	1
				Arvicolinas	507	41. Arvicola	501	Hypudaeus		1	1.1
								Hemiotomys		10	1
								Chilotus		1	
								Pedomys	1	3	
								Pitymys		1	
						42. Myodes	554	Synaptomys	558	1	
							in second	Myodes		2	
						43. Fiber	560				

I. TABLE OF THE HIGHER GROUPS.

...

NORTH AMERICAN MAMMALS.

Order.	Page.	Family.	Page.	Sub-family.	Page.	Genus.	Page.	Sub-genus.	Page.	Species examined and identified.	Species not identified
		12. Hystricidae	565			44. Erethizon	568			2	
		13. Leporidae	572			45. Lepus	573			13	2
and for the second						46. Lagomys	618			1	
IV. EDENTATA	621	14. Effodienta				47. Dasypus	100.55			1	
V. PACHYDERMATA.	625	15. Sciuridae	626	•••••		48. Dicotyles	a state of the second	•••••		1 1 1 1	
VI. RUMINANTIA	628	16. Cervidae	629	Cervinae	630	49. Alce	631			- E1	
	1					50. Rangifer	633	••••••			
	-	in the second	1.4.4			51. Cervus	637		•••••		
	-	17. Cavicornia	664	Aptilopinae	664	52. Antilocapra	665	•••••	•••••		
and the second						53. Aplocerus				100	
Contraction of the local sectors of the local secto				Ovinae	664	54. Ovis	673			1	
				Bovinae	664	55. Ovibos	680			1	
		1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1				56. Bos	682	••••••		1	
										220	35

TABLE OF THE HIGHER GROUPS-Continued.

II. LIST OF SPECIES.

ALIST OF EXAMINED AND IDENTIFIED SPECIES OF NORTH AMERICAN MAMMALS.	
1. NEOSOREX, Baird.	Page. 11
1Neosorex navigator, Cooper, (n. s.)-Water ShrewFort Vancouver	11
2. SOREX, L.	13
2Sorex trowbridgii, Baird, (n. s.)-Trowbridge's ShrewAstoria to Steilacoom	13
3.—Sorex vagrans, Cooper, (n. s.)—Wandering Shrew.—Cape Flattery, W. T., to Petaluma, Cal. Range in lat. 48.30 to 38.20	15
4Sorex suckleyi, Baird, (n. s.)-Suckley's ShrewSteilacoom, W. T., to Monterey, Cal. Range in lat. 47.90 to 36.30	18
5.—Sorex pachyurus, Baird, (n. s.)—Thick-Tailed Shrew.—Pembina, Minn., to Fort Ripley. Lat. 49.00 to 46.19	20
6.—Sorex forsteri, Rich.—Forster's Shrew.—Carlisle, Pa. (Hudson's Bay, Rich. New York. DeKay.)	22
7.—Sorex richardsonii, Bachm.—Richardson's Shrew.—Racine, Wisc.? (Hudson's Bay, Rich. N. W. Territory, Bach.)	24
8.—Sorex platyrhinus, Wagner.—Eared Shrew.—Hingham, Mass., to Cleveland, Ohio, and Burlington, Vt	25
9Sorex cooperi, BachCooper's ShrewLabrador to Medicine Bow Cr., Neb	27
10.—Sorex haydeni, Baird, (n. s.)—Hayden's Shrew.—Fort Pierre to Ft. Union, Neb. Range on the Missouri river, from lat. 44.20 to 48.00	29
11Sorex personatus, GeoffMasked ShrewSouthern Atlantic States. Washington	30
12.—Sorex hoyi, Baird, (n. s.)—Hoy's Shrew.—Racine, Wisc	32
13Sorex thompsoni, Baird, (n. s.)-Thompson's ShrewHalifax, N. Scotia, to Zanes- ville, Ohio	34
3. BLARINA, Gray.	36

14.—Blarina	talpoides, G	ray Mole-	Shrew]	Halifax, 1	N. 1	S.	Montreal to Columbus,	
	Ohio, an	d Western	Georgia.	Range	in 1	lat.	45.30 to 32.30; in long.	
	from 63.	30 to 81.30.						30

	Page.
15Blarina brevicauda, GrayShort-tailed ShrewIllinois to Nebraska	42
16.—Blarina carolinensis.—Carolina Shrew.—S. Carolina to Missouri	45
17Blarina angusticeps, Baird, (n. s.)-Burlington, Vermont	47
18.—Blarina cinerea.—Ash-colored Shrew.—S. Pennsylvania to Florida	45
19Blarina exilipes, Baird, (n. s.)-Short-footed ShrewTexas to Mississippi	51
20.—Blarina berlandieri, Baird, (n. s.)—Matamoras, Mexico	53

4. SCALOPS, Cuv.

58

a. Scalops.

21.—Scalops aquaticus, Cuv.—Common mole.—Middleboro', Mass., to Indian river, Fla., and west to the Lower Mississippi. Range along the Atlantic States from lat. 42. to 27.30.	60
22Scalops argentatus, Aud. & BachSilvery MoleDetroit to Fort Riley, and south to Prairie Mer Rouge? Range in lat. from 42.20 to Louisi- ana? in long. 83. to 97	63
b. Scapanus, Pomel.	58
23.—Scalops townsendii, Bach.—Oregon Mole.—Steilacoom, W. T., to San Francisco, Cal. Range on the Pacific coast from lat. 47.10 to 37.48	65
24.—Scalops breweri, Bach.—Hairy-Tailed Mole.—Connecticut and New York, to Cleve- land, Ohio	68
5. CONDYLURA, III.	71
25.—Condylura cristata, Ill.—Star-Nosed Mole.—Halifax, N. S., to Fort Ripley, and to Carlisle, Pa. Range in lat. 40.12 to 46.20; in long. 63.30 to 94.20.	71
6. UROTRICHUS, Temm.	76
26Urotrichus gibbsii, Baird, (n. s.)-Cascade Mountains, W. T	76
7. FELIS, L.	81
27.—Felis concolor, L.—Panther.—Fort Steilacoom, W. T., to southern Chile, and from the Atlantic to the Pacific. Range from 47 (or 55. according	
to Richardson) N. lat. to 54 S. lat	83
28Felis onca, LJaguar; Mexican Tiger-Brazos river, Texas, to South America	86
29.—Felis pardalis, Linn.—Tiger Cat; Ocelot.—Northern Texas?	87
30Felis eyra, DesmarestRio Grande of Texas	88

	Page.
31.—Felis yaguarundi, Desmarest.—Yaguarundi.—Rio Grande of Texas	88
8. LYNX, Raf.	89
32Lynx rufus, RafWild CatAtlantic to Pacific. Upper Missouri to the Gila river. Not on the Rio Grande?	90
33Lynx rufus var. maculatus, Aud. & BachTexas Wild CatMatamoras, Mexico, to Fort Yuma, Cal., and to Fort Tejon, Cal	93
34Lynx fasciatus, RafRed CatWashington Territory	96
35.—Lynx canadensis, Raf.—Canada Lynx.—Nova Scotia to Rocky Mountains. Penn. to 66 N. lat.?	99
9. CANIS, L.	104
36a.—Canis occidentalis var. griseo-albus.—White and Gray Wolf.—North America generally?	104
36bCanis occidentalis var. nubilusDusky WolfMissouri river to the Pacific	111
36c.—Canis occidentalis var. mexicanus.—Mexican Wolf.—Rio Grande of Texas and Sonora	113
37Canis latrans, SayCoyote; Prairie WolfFort Riley, Kansas, to the Pacific, and Upper Missouri to the Rio Grande of Texas	113
10. VULPES.	121
a. Vulpes, Baird.	121
38aVulpes fulvus var. fulvusRed FoxAtlantic States to the Missouri, and Penn- sylvania to Canada	124
38b Vulpes fulvus var. decussatus Cross Fox Northern New York to Canada	127
38cVulpes fulvus var. argentatusSilver Fox ; Black FoxWashington Territory	128
39Vulpes macrourus, BairdPrairie FoxUpper Missouri to Plains of Colum- bia, O. T	130
40Vulpes velox, Aud. & BachKit Fox; Swift FoxPlains west of Missouri to the Cascade Mountains of Oregon	133
41Vulpes lagopus, RichArctic FoxArctic America	137
b. Urocyon, Baird.	121
42Vulpus virginianus, RichGray FoxPennsylvania to the Southern States, and from the Atlantic to the Pacific; from the Gila to Puget's Sound.	

According to Aud. & Bach., from Maine to Florida...... 138

LIST OF S	PECIES.
-----------	---------

43Vulpes littoralis, Baird, (n. s.)-Island FoxIsl. of San Miguel, coast of Cal	Page. 143
11. BASSARIS, Licht.	147
44Bassaris astuta, LichtCivet CatCity of Mexico to Texas and California	. 147
12. MUSTELA, Linn.	149
45.—Mustela pennantii, Erxl.—Fisher; Black Cat.—Southern Penn. to New York, (to 65 N., Richardson,) and from the Allegheny range to Washing- ton Territory	149
46.—Mustela americana, Turton.—Pine Marten; American Sable.—Adirondac Moun- tains, N. Y., and west to the Cascade Mountains and Puget's Sound	152
13. PUTORIUS, Cuvier.	159
47.—Putorius pusillus, Aud. & Bach.—Least Weasel.—Pembina, Minn., to Puget's Sound: New York, (DeKay.)	159
48.—Putorius cicognanii.—Small Brown Weasel.—Labrador and Nova Scotia to Puget's Sound, and south to Boston, Massachusetts	161
49.—Putorius richardsonii, Bp.—Little Ermine.—Nova Scotia to Puget's Sound, and south to Boston	164
50.—Putorius noveboracensis, DeKay.—White Weasel; Ermine.—Boston and North- ern New York, and Illinois, to southern Penn. and Arkansas. Not in the far west?	166
51.—Putorius longicauda, Rich.—Long-Tailed Weasel.—Upper Missouri and Platte. (Carlton House, H. B. T., Rich.)	169
52.—Putorius kaneii, Baird, (n. s.)—Kane's Ermine.—Behring's Straits and Siberia	172
53.—Putorius frenatus, Aud. & Bach.—Bridled Weasel.—City of Mexico to the Rio Grande, Texas	173
54.—Putorius xanthogenys.—Yellow-Cheeked Weasel.—San Diego to San Francisco, Cal., on the coast	176
55.—Putorius vison, Rich.—Brown Mink.—New York and Massachusetts to Florida, and west to the Pacific and Puget's Sound	177
56.—Putorius nigrescens, Aud. & Bach.—Little Black Mink.—Massachusetts and Northern New York to Fort Leavenworth	180
14. GULO, Storr.	181
57Gulo luscus, SabineWolverineSalt Lake, and Black Hills, Neb., to Arctic America, (northern New York, Aud. & Bach.)	181

•

LIST OF SPECIES.	9
15. LUTRA, Linn.	Page. 184
58.—Lutra canadensis, Sab.—American Otter.—Northern parts of the United States to Florida, and west to the Rocky Mountains	184
59.—Lutra californica, Gray.—California Otter.—Coast region and Cascade Mountains of California and Oregon	187
16. ENHYDRA, Fleming.	189
60Enhydra marina, FlemingSea OtterPacific coast of the United States	. 189
17. MEPHITIS, Cuvier.	191
a. Thiosmus, Licht.	191
61.—Mephitis mesoleuca, Licht.—White-Backed Skunk.—Chico, Mexico, to Western Texas.	192
62.—Mephitis varians, Gray.—Texas or Long-Tailed Skunk.—Texas and Northern Mexico	193
63.—Mephitis occidentalis, Baird, (n.s.)—California Skunk.—Western coast of the United States	194
64.—Mephitis mephitica.—Common Skunk.—United States, east of the Missouri plains and north of Texas.	195
65.—Mephitis bicolor, Gray.—Little Striped Skunk.—Southern Texas and California	197
18. TAXIDEA, Waterhouse.	201
66.—Taxidea americana, Waterh.—Missouri Badger.—Iowa and Wisconsin to the Pacific coast, and from Arkansas to 49 N. lat.—(To 58 N. lat., Rich.)	202
67.—Taxidea berlandieri, Baird.—Mexican Badger.—Northern Mexico, Southern and Western Texas. (To Lower California?)	205
19. PROCYON, Storr.	207
68.—Procyon lotor, Storr.—Common Raccoon.—Massachusetts to Florida, and west to Fort Kearney. Not in Southern Texas?	209
69.—Procyon hernandezii, Wagler.—California or Black-Footed Raccoon.—Rio Grande of Texas to Pacific coast, and north along the coast to Puget's Sound	212
69a.—Procyon hernandezii, var. mexicana, Baird.—Sonora Raccoon.—Gulf of California	215
20. URSUS, L.	216
70. Ursus horibilis, Ord.—Grizzly Bear.—Plains of Upper Missouri to the Rocky Moun- tains, and along their base, thence to the Coast of California, (not of Oregon or Washington?)	219

.

LIST	OF	SPEC	CIES.
------	----	------	-------

•

71Ursus americanus, PallasBlack BearUnited States generally	Page. 225
71 aUrsus americanus var. cinnamoneus, Aud. & BachRocky Mountains and	000
72 Ursus maritimus, LinnWhite or Polar BearShores and Islands of Arctic Regions	228 229
21. DIDELPHYS, Linn.	232
73 Didelphys virginiana, Shaw.—Possum.—From Hudson River to the southern part of the eastern United States, and west to the Missouri. Not in south- ern Texas	232
74 Didelphys californica, BennetTexas PossumRio Grande of Texas to the City of Mexico, and west to Lower California	233
22. SCIURUS, Linnæus.	243
75.—Sciurus vulpinus, Gmelin.—Southern Fox Squirrel.—North Carolina to Florida, and along the Gulf States to eastern Texas	246
76.—Sciurus cinereus, Linn.—Cat Squirrel; Pennsylvania Fox Squirrel.—New Jersey to Virginia, and west to the Alleghenies	248
77.—Sciurus ludovicianus, Custis.—Western Fox Squirrel.—Mississippi valley	251
78.—Sciurus limitis, Baird, (n. s.)—Pecos Fox Squirrel.—Pecos River, Texas	256
79.—Sciurus carolinensis, Gm.—Gray Squirrel; Black Squirrel. (Black var.)—Kastern United States to the Missouri River	256
79 a.—Sciurus carolinensis, var. mexicanus?—Mexican Gray Squirrel.—Northern Mexico	263
80.—Sciurus fossor, Peale.—California Gray Squirrel.—Pacific Coast from San Diego, California, to the Columbia River, at Fort Dalles	264
81.—Sciurus castanonotus—Baird, (n. s.)—Chestnut Backed Squirrel.—Southern Rocky Mountains. (Fort Webster)	266
82Sciurus aberti, WoodhouseTuft-Eared SquirrelSan Francisco Mountains, New Mexico	267
83.—Sciurus hudsonius, Pallas.—Red Squirrel; Chickaree.—Labrador (Lat. 56°) to Mississippi; and in the United States from the Atlantic to the Missouri River	260
84.—Sciurus fremontii, Towns.—Mountain Gray Squirrel	200
85Sciurus richardsonii, BachRichardson's SquirrelFrom the Main Rocky Moun- tains to the Cascade Range.	273

LIST OF SPECIES.	11
86Sciurus douglassii, BachOregon Red SquirrelSan Francisco to Puget's Sound,	Page.
near the coast	275
AN DEPONING OF I	
23. PTEROMYS, Cuvier.	86
87.—Pteromys volucella, Cuv.(?)—Flying Squirrel.—United States, to the Missouri River	286
88.—Pteromys hudsonius, Fischer.—Northern Flying Squirrel.—Northern parts of the United States from Maine to Minnesota, and north	288
89.—Pteromys alpinus, Richardson.—Rocky Mountain Flying Squirrel.—Rocky Moun- tains?	289
90-Pteromys oregonensis, BachOregon Flying SquirrelNear mouth of Columbia River to Southern California?	290
24. TAMIAS, Illiger.	291
91Tamias striatusChipmunkMontreal to Virginia, and west to the Missouri	292
92.—Tamias quadrivittatus, Rich.—Missouri Striped Squirrel.—Upper Missouri to Rocky Mountains, and west to the Cascade Range. Along the Rocky Mountains as far south as Fort Staunton, New Mexico. Lat. 33° 30'	297
93.—Tamias dorsalis, Baird.—Gila Striped Squirrel.—Fort Webster, New Mexico, about Lat. 33.00, Long. 108.00	300
94.—Tamias townsendii, Bach.—Townsend's Striped Squirrel.—From Puget's Sound to Petaluma, California, along and west of Cascade Mountains	301
25. SPERMOPHILUS, Cuvier.	304
a. Otospermophilus, Brandt.	305
95.—Spermophilus beecheyi, Rich.—California Ground Squirrel.—Coast of California, from San Diego to Fort Reading	307
96Spermophilus douglassii, RichColumbia Ground SquirrelColumbia River, from Fort Dalles to its mouth, and south to Klamath Lake	309
97. Spermophilus grammurus, Bachman.—Line-Tailed Squirrel.—Head of Arkansas River; along the Rocky Mountains to Sonora	310
98Spermophilus couchii, Baird, (n. s.)-Black Ground SquirrelNorthern Mexico	311
99Spermophilus lateralis, RichSay's Striped SquirrelRocky Mountains to Cas- cades, and between about Lat. 38° 25' to northern line of California	312
b. Colobotis, Brandt.	306
100Spermophilus harrisii, Aud. & BachHarris's SpermophileMohave Desert	313

LIST	OF	SPEC	IES.
------	----	------	------

	Page.
101. Spermophilus franklini, Rich.—Gray Gopher.—Northern Illinois and Wisconsin, and to Minnesota and the Saskatchewan	314
102.—Spermophilus tereticauda, Baird, (n. s.)—Round-Tailed Spermophile.—Fort Yuma, Cal.	315
103.—Spermophilus tridecem-lineatus, Aud. & Bach.—Striped Gopher: Prairie Squir- rel.—Eastern Michigan to the plains of the Missouri, and south to Red River, Arkansas, and Fort Thorn, N. M	316
104.—Spermophilus mexicanus, Wagner.—Mexican Ground Squirrel.—Rio Grande of Texas and the Pecos, and south into Mexico	319
105.—Spermophilus spilosoma, Bennett.—Sonora Ground Squirrel.—Fort Staunton, N. M. and El Paso, to Chihuahua, and west to the Gulf of California	321
106.—Spermophilus parryi, Rich.—Parry's Marmot.—Shores and islands of Arctic seas; island of Arikamtchitchi, on the Asiatic side of Behring's Straits	323
107.—Spermophilus richardsonii.—Yellow Gopher.—Sault St. Marie, Mich., and west to the Rocky Mountains north of Upper Missouri	325
108.—Spermophilus townsendii, Bach.—Townsend's Spermophile.—Rocky Mountains to the north	326
AL OWNOR D. C	000
26. CYNOMYS, Raf.	329
109.—Cynomys ludovicianus.—Prairie Dog.—Milk river and Upper Missouri; west towards the Rocky Mountains; south to Red River; southwest to the Upper Rio Grande, and a short distance into Sonora?	Q
110.—Cynomys gunnisonii, Baird (n. s.)—Short-Tailed Prairie Dog.—Coochetope Pass to Lodge Pole Creek, in or near the Rocky Mountains	335
27. ARCTOMYS, Schreber.	348
111Arctomys monax, GmelinWoodchuck ; Ground HogMassachusetts to Wis-	
consin and Canada; south to Virginia. (To South Carolina, Aud. & Bach.)	
112.—Arctomys flaviventer, Bachman.—Yellow-Footed Marmot.—Black Hills, Neb	343
28. APLODONTIA, Rich.	350
113.—Aplodontia leporina, Rich.—Sewellel; Showt'l.—Puget's Sound	353
29. CASTOR, Linn.	355
114.—Castor canadensis, Kuhl.—American Beaver.—Throughout the entire area of North America	f 355

LIST OF SPECIES.	13
	Page.
30. CASTOROIDES, Foster.	363
115Castoroides ohioensis, FosterFossil BeaverNorthern Ohio, New York and Michigan	363
31. GEOMYS, Rafinesque.	368
116Geomys bursarius, RichPouched or Pocket GopherMissouri to Minnesota and Nebraska	372
117Geomys breviceps, Baird, (n. s.)-Short-Headed GopherLouisiana to Middle or Western Texas	378
118.—Geomys pinetis, Rafinesque.—Salamander.—Georgia and Florida. (Alabama, Aud. & Bach.)	380
119.—Geomys clarkii, Baird, (n. s.)—Pecos Gopher.—Pecos River, Texas	383
120.—Geomys castanops, Leconte.—Chestnut-Faced Gopher.—Bent's Fort, Upper Arkansas River	384
Norg.—The two following Mexican species are referred to— Geomys hispidus, Leconte.—Mexico	386 387
32. THOMOMYS, Maxim.	388
121.—Thomomys bulbivorus.—California Gopher.—Coast of California, from Tejon valley to Petaluma	389
122Thomomys laticeps, Baird, (n. s.)-Broad-Headed GopherHumbolt Bay ,Cal	392
123.—Thomomys douglassii, Giebel.—Oregon Gopher.—Columbia River and Puget's Sound to the Dalles	394
124. Thomomys borealis.—Upper Columbia River? Canoe Creek, Cal. (East of Cascade mountains.)	396
125.—Thomomys rufescens, Maxim.—Fort Union Gopher.—Fort Pierre to Fort Union, on the Missouri	397
126.—Thomomys umbrinus.—Sonora Gopher.—Cimarron River, N. M., west to the Rocky Mountains, and south into Sonora	399
127Thomomys fulvusSan Francisco mountains, N. M., to Fort Yuma and San Diego	402
33. DIPODOMYS, Gray.	406
128.—Dipodomys ordii, Woodhouse.—Kangaroo Rat.—Platte River, along the eastern slope of the Rocky Mountains, into Durango and Coahuila, Mexico	410
129.—Dipodomys phillipii, Gray.—Jumping Rat.—Los Angeles, Cal., to Fort Walla- Walla, O. T., on the Columbia	412

LIST OF SPECIES.	LI	ST	OF	SPE	CIES.
------------------	----	----	----	-----	-------

	Page.
130.—Dipodomys agilis, Gambel.—Fort Yuma and San Diego, Cal., to San Francisco	414
34. PEROGNATHUS, Maxim.	416
a. Perognathus, Max.	417
131.—Perognathus penicillatus, Woodhouse.—Tuft-tailed Mouse.—San Francisco Moun- tains, New Mexico, to Fort Yuma, California, along Colorado	418
132.—Perognathus fasciatus, Pr. Max.—Fort Union, Nebraska, and Fort Riley, Kansas Territory, to Chihuahua	420
133.—Perognathus hispidus, Baird, (n. s.)—Matamoras	421
134.—Perognathus monticola, Baird, (n. s.)—St. Mary's, Rocky Mountains	422
b. Cricetodipus, Peale.	418
135.—Perognathus flavus, Baird, (n. s.)—Upper Missouri, along eastern slopes of Rocky Mountains to Sonora and Chihuahua; and along the Rio Grande to Matamoras.	423
136.—Perognathus parvus, Leconte.—King's River, California. (Oregon, Peale.)	425
35. JACULUS, Wagler.	429
137.—Jaculus hudsonius.—Jumping Mouse.—Nova Scotia, (Labrador, Pennant,) to Southern Pennsylvania, and west to the Pacific Ocean	430
TRIBE MURES.	
36. MUS, Linnæus.	436
138Mus decumanus, PallasBrown RatThroughout North America. (Introduced.)	438
139Mus rattus, LBlack RatThroughout North America. (Introduced.)	439
140.—Mus tectorum, Savi.—White-Bellied Rat; Roof Rat.—Tar River North Carolina, to Georgia; Northern Mexico. (Introduced.)	441
141Mus musculus, LinnCommon MouseNorth America generally. (Introduced.)	443
TRIBE SIGMODONTES.	445
37. REITHRODON, Waterhouse.	447
142Reithrodon humilisHarvest MouseSouth Carolina and Georgia, to St. Louis?	448
143.—Reithrodon montanus, Baird, (n. s.)—Rocky Mountains. Lat. 39°	449
144.—Reithrodon megalotis, Baird, (n. s.)—Sonora	451
145. Reithrodon longicauda Baird (n. s.)-San Francisco and Petaluma California	451

LIST OF SPECIES.	15
38. HESPEROMYS, Waterhouse.	Page. 453
a. Hesperomys.	458
146Hesperomys leucopus, WagnerWhite-footed MouseNova Scotia (and Lab- rador,) to Virginia, and west to the Mississippi River	459
147Hesperomys texanus, WoodhouseTexas MouseRio Grande, Texas	464
148Hesperomys gambelii, Baird, (n. s.)-California and Oregon	464
149.—Hesperomys austerus, Baird, (n. s.)—Washington Territory	466
150.—Hesperomys nuttalli.—Red Mouse.—Southern Pennsylvania to Georgia, and west to St. Louis.	467
151.—Hesperomys gossypinus, Leconte.—Cotton Mouse.—Lowlands of Georgia and South Carolina	469
152Hesperomys cognatus, LeconteGray-Bellied MouseSouthern Atlantic and Gulf States, from North Carolina to Louisiana	469
153.—Hesperomys boylii, Baird, (n. s.)—Long-Tailed Mouse.—Western California and Oregon	471
154Hesperomys myoidesHamster MouseCanada, Vermont, and New York	472
155.—Hesperomys sonoriensis, Leconte.—Upper Missouri, and Rocky Mountains to El Paso and Sonora	474
156.—Hesperomys michiganensis, Wagner.—Prairie Mouse.—Michigan, Wisconsin, and Illinois, to St. Louis.	476
157.—Hesperomys californicus.—Great-Eared Mouse.—Santa Clara, California	478
185.—Hesperomys eremicus, Baird, (n. s.)—Desert Mouse.—Fort Yuma, California	479
b. Onychomys, Baird.	458
159Hesperomys leucogasterMissouri MousePlains along the upper Missouri	480
c. Oryzomys, Baird.	458
160.—Hesperomys palustris, Wagner.—Rice Field Mouse.—South Carolina and Georgia	482
39. NEOTOMA, Say & Ord.	486
161.—Neotoma floridana, Say & Ord.—Florida Rat; Wood Rat.—Southern Atlantic and Gulf States, and up the Missouri. Stragglers as far as New York	487
162Neotoma mexicana, Baird, (n.s.)-Mexican Bush RatPecos river to Chi- huahua, and west to Gulf of California	490

LIST (OF S	SPE	CIES.
--------	------	-----	-------

	Page.
163.—Neotoma micropus, Baird, (n. s.)—Black Wood Rat.—Lower Rio Grande of Texas and Mexico	492
164.—Neotoma fuscipes, Cooper, Mss., (n. s.)—Brown-Footed Rat.—Santa Clara to Petaluma, Cal	495
165.—Neotoma occidentalis, Cooper, (n. s.)—Hairy-Tailed Rat.—Mouth of Columbia, and Puget's Sound	496
166.—Neotoma cinerea.—Rocky Mountain Rat.—Eastern slope of northern Rocky Mountains, and upper Missouri	499
166.*-Neotoma magister, Baird, (n. s.)-Carlisle Bone Cave, Pemnsylvania. Fossil.	
40. SIGMODON, Say & Ord.	500
167.—Sigmodon hispidus, Say & Ord.—Cotton Rat.—Southern Atlantic and Gulf States	502
168.—Sigmodon berlandieri, Baird, (n. s.)—Texas Cotton Rat.—Texas	504
41. ARVICOLA, Lacep.	509
a. Hypudaeus, Keys. & Blas.	
169.—Arvicola gapperi, Vigors.—Red-Backed Mouse.—Nova Scotia, Massachusetts, Canada, and Lake Superior	518
b. Hemiotomys, Selys.	
170Arvicola riparia, OrdBank MouseNorthern and Middle States	522
171.—Arvicola breweri, Baird, (n. s.)—Gray Mouse.—Muskeget Island, Mass	525
172.—Arvicola rufidorsum, Baird, (n. s.)—Holmes' Hole, Mass. (Martha's Vineyard.)	526
173.—Arvicola townsendii, Bachman.—Washington & Oregon	527
174.—Arvicola montana, Peale.—Cascade Mountains, Oregon & California; Western Nebraska?	528
175.—Arvicola longirostris, Baird, (n. s.)—Cascade Mountains, Cal	530
176.— Arvicola edax, Leconte.—Monterey to San Diego, Cal	531
177Arvicola californica, PealeCalifornia	532
178.—Arvicola occidentalis, Peale.—Puget's Sound	534
179.—Arvicola modesta, Baird, (n. s.)—Sawatch Pass, Rocky Mountains	535
c. Chilotus, Baird.	516
180Arvicola oregoni, BachmanSteilacoom, W. T., to Tomales Bay, Cal	537
d. Pedomys, Baird.	517
181.—Arvicola austera, Leconte —Prairie Meadow Mouse —Wisconsin to Louisiana	539

LIST OF SPECIES.	17
182.—Arvicola cinnamonea, Baird, (n. s.).—Pembina, Minnesota	Page, 541
183.—Arvicola haydenii, Baird, (n. s.)—Fort Pierre, Nebraska	543
e. Pitymys, McMurtrie.	517
184.—Arvicola pinetorum, Leconte.—Upland Mouse.—Long Island, N. Y., to Florida and the Gulf States, Illinois and Ohio	5 4 4
42. MYODES, Pallas.	554
a. Synaptomys, Baird.	554
185.—Myodes cooperii, Baird, (n. s.)—United States	558
b. Myodes.	554
186Myodes torquatus, Keys. & BlasLemmingArctic Regions	558
187.—Myodes obensis, Brants.—Lemming.—Arctic Regions	559
43. FIBER, Cuv.	560
188.—Fiber zibethicus, Cuv.—Musk Rat.—North America, generally	561
44. ERETHIZON, F. Cuvier.	568
 189.—Erethizon dorsatus, F. Cuv.—White-Haired Porcupine.—Eastern United States to the Mississippi river, and from Northern Pennsylvania to Canada.— (To 67°, Rich.) 190.—Erethizon epixanthus, Brandt.—Yellow-Haired Porcupine.—Upper Missouri ; 	568
whole Pacific Coast	569
45. LEPUS.	573
191.—Lepus glacialis, Leach.—Polar Hare.—Newfoundland, north to the Polar Seas, and Greenland	577
192Lepus americanus, ErxlNorthern Hare; White RabbitMountains of Vir- ginia to Labrador. Not west of Minnesota?	579
193Lepus washingtonii, Baird, (n. s.)-Red HareWashington Territory, to 54° 40'	583
194.—Lepus campestris, Bach.—Prairie Hare.—Upper Missouri and Saskatchewan Plains to the Cascade Range of Oregon	585
195.—Lepus callotis, Wagler.—Jackass or Mule Rabbit.—Northern Mexico through southern Texas, and west through New Mexico to the Rocky Moun- tains. Fort Boisé, Oregon Territory? Southern Sonora?	
196.—Lepus californicus, Gray.—California Hare.—Lower Colorado and San Diego, to Fort Jones, Cal. Not found east of Cascades	

	Page.
197.—Lepus sylvaticus, Bach.—Gray Rabbit.—From Massachusetts throughout the United States, as far as Indianola, Texas? and west to the Missouri, as far as Fort Union, Nebraska	597
	591
198.—Lepus artemisia, Bach.—Sage Hare.—Regions west of the Missouri to the Rocky Mountains, and to the Cascade Mountains of Oregon, along the Colum-	
bia on the north, and to the City of Chihuahua on the south 199.—Lepus bachmani, Waterhouse.—Bachman's Hare.—Lower Rio Grande of Texas.	602 606
200.—Lepus audubonii, Baird, (n. s.)—Audubon's Hare.—Coast of California, from San Diego to San Francisco	
201.—Lepus trowbridgii, Baird, (n. s.)—Coast of California, from San Francisco to San Diego	610
202Lepus aquaticus, BachWater RabbitLower Mississippi River	612
203.—Lepus palustris, Bach.—Marsh Rabbit.—Lowlands of southern coast, from South Carolina to Florida	615
46. LAGOMYS, Cuv.	618
204.—Lagomys princeps, Rich.—Little Chief Hare.—South Pass of Rocky Mountains northward	619
47. DASYPUS, Linnæus.	623
205.—Dasypus novem-cinctus, Linn.—Texas Armadillo.—Lower Rio Grande of Texas to Paraguay	623
18 DICOTVIES Com	627
48. DICOTYLES, Cuv.	027
206.—Dicotyles torquatus, Cuv.—Texas Peccary; Mexican Hog.—Texas to Red River, Arkansas	627
49. ALCE, Ham. Smith.	631
207.—Alce americanus, Jardine.—American Moose.—Northern portions of the eastern United States to Labrador, whence it extends west to the Pacific Ocean	
50. RANGIFER, Ham. Smith.	633
208-Rangifer caribou, Aud. & BachWoodland CaribouMaine and New Bruns- wick, and west to north shore of Lake Superior	633
209.—Rangifer groenlandicus.—Barren Ground Caribou.—Barren Grounds of Arctic America and Greenland.	634

51.	\mathbf{CE}	PVI	ITS:	Т.
01.	ULL.	LUY	00,	11.

	Page.
210-Cervus canadensis, ErxlAmerican ElkNorthern portions of United States, to Upper Missouri, and west to the Pacific. Found in the Alleghenies	537
of Pennsylvania and Virginia. (North to 57°, Rich.)	638
211.—Cervus virginianus, Boddaert.—Virginia Deer.—United States generally, east of the Missouri, and as far north as Maine. Eastern Texas?	643
212.—Cervus leucurus, Douglass.—White-Tailed Deer.—Upper Missouri and Platte to the Columbia River and Washington Territory. Western Texas and New Mexico?	649
213.—Cervus mexicanus, Gmelin.—Sonora Deer.—Sonora and near El Paso?	658
214.—Cervus macrotis, Say.—Mule Deer.—Upper Missouri and Platte to the Cascade Range, (head of Des Chutes River.) Oregon Territory. Not extend- ing to the Pacific. Headwaters of Arkansas	656
215.—Cervus columbianus, Rich.—Columbia Black-Tailed Deer.—Pacific Coast, from San Diego to Puget's Sound	659
52. ANTILOCAPRA, Ord.	665
916 Antileconne omericane Ord Brong Harn Antelena ; Cabres Plaine west of	
216—Antilocapra americana, Ord.—Prong-Horn Antelope; Cabree.—Plains west of Missouri, from the Lower Rio Grande to the Saskatchewan, and west	
to the Cascade and Coast range of the Pacific slope	666
53. APLOCERUS, Ham. Smith.	671
217.—Aplocerus montanus, Rich.—Mountain Goat.—Northern Rocky Mountains of the United States and British America; also on the northern Cascade	
Mountains	671
54. OVIS, Linn.	673
218.—Ovis montana, Cuvier.—Bighorn; Mountain Sheep.—Broken ground on the Upper	
Missouri and Platte. Rocky Mountains, generally, as far west, at least, as Coast and Cascade Mountains of the Pacific slope	673
55. OVIBOS, Blainville.	680
219.—Ovibos moschatus, Blainville.—Musk Ox.—Barren Grounds of Arctic America	680
BOS, Linnæus.	682
220.—Bos americanus, Gmelin.—American Buffalo.—Formerly found throughout nearly the whole of North America, east of the Rocky Mountains. Now	
confined to the plains west of the Missouri and along the slopes of the Rocky Mountains.	682

II.-LIST OF SPECIES-Continued.

B .-- LIST OF DESCRIBED NORTH AMERICAN SPECIES NOT SEEN AND IDENTIFIED.

2. SOREX, Linn.

	Page.
1.—Sorex palustris, Rich.—Marsh Shrew.—Hudson's Bay region	55
2Sorex fimbripes, BachFringed ShrewAlleghenies of Pennsylvania	55
3.—Sorex parvus, Say.—Least Shrew.—Missouri River	56
4.—Sorex harlani, Duvernoy.—New Harmony, Indiana. (Doubtful)	56
4. SCALOPS, Cuv.	
5.—Scalops latimanus, Bach.—Mexico and Texas	65
9. CANIS, L.	
6 — Canis occidentalis, var. ater. — Black Wolf. — Southern States	113
6 aCanis occidentalis, var. rufusRed WolfTexas	113
13. PUTORIUS, L.	
7Putorius nigripes, Aud. & BachBlack-Footed FerretLower waters of Platte	180
17. MEPHITIS, Licht.	
8Mephitis mesomelas, LichtBlack-Backed SkunkLouisiana? (Doubtful)	199
The following Mexican species may yet be found in the United States : Mephitis leuconota, Licht.—Alvarado River, Mephitis macroura, Licht.—Temperate, Mexico. Mephitis vittata, Licht.—Oaxaca, Mexico.	Mexico.
19. PROCYON, Storr.	
9.—Procyon psora, Gray.—Short-Tailed Racoon.—Sacramento River, California. (Very doubtful)	215
22. SCIURUS, L.	
10Sciurus colliaei, RichCollie's SquirrelWest Coast of Mexico	280
11.—Sciurus mustelinus, Aud. & Bach.—Weasel Squirrel.—(California, Aud. & Bach.) Probably Mexican?	280
12.—Sciurus nigrescens, Bennett.—Dusky Squirrel.—Lower California. Not belonging to the fauna of the United States?	280
13Sciurus lanigerus, Aud. & BachWoolly Squirrel(California, Aud. & Bach.)	280
14.—Sciurus leporinus, Aud. & Bach.—Hare Squirrel.—(California, Aud. & Bach.) Doubtful	280
15.—Sciurus ferruginiventris, Aud. & Bach.—(California, Aud. & Bach.) Probably Mexican	281
For an enumeration of other supposed North American Squirrels, see pages 281-285.	

LIST OF SPECIES.	21				
	Page.				
25. SPERMOPHILUS, Cuv.	304				
16.—Spermophilus macrourus, Bennet.—West Coast of Mexico. (Not belonging to the United States.)	327				
27. ARCTOMYS, Schreb.	338				
17.—Arctomys pruinosus, Gmelin.—Hoary Marmot; Whistler.—Hudson's Bay to Northwest coast of America	34 5				
18.—Arctomys lewisii, Aud. & Bach.—Lewis' Marmot.—Columbia River?	347				
32. THOMOMYS, Max.	388				
19.—Thomomys talpoides, Giebel.—Mole Gopher.—Hudson's Bay	403				
33. DIPODOMYS, Gray.	406				
20Dipodomys wagneri, LeconteSouth Carolina(!) (Very doubtful)	415				
21.—Dipodomys heermannii, Leconte.—Sierra Nevada, California	415				
37. REITHRODON, Waterh.	447				
22.—Reithrodon carolinensis.—South Carolina	452				
38. HESPEROMYS, Waterh.	453				
23.—Hesperomys campestris, Lec.—New Jersey. (Uncertain)	485				
41. ARVICOLA, Lac.	509				
24.—Arvicola albo-rufescens, Emmons.—Massachusetts	549				
25.—Arvicola borealis, Rich.—Great Bear Lake	549				
26.—Arvicola dekayi, Aud. &. Bach.—New York					
27Arvicola drummondii, Aud. & BachBritish Rocky Mountains	550				
28.—Arvicola hirsutus, Dekay.—New York.	550				
29.—Arvicola nasuta, Aud. & Bach.—Massachusetts	550				
30.—Arvicola oneida, Dekay.—New York	551				
31.—Arvicola richardsonii, Dekay.—British Rocky Mountains	551				
32.—Arvicola rubricatus, Rich.—Behring's Straits	551				
33.—Arvicola texiana, Aud. & Bach.—Texas	552				
34.—Arvicola xanthognathus, Leach.—Hudson's Bay	552				
Note Of the above Arvicolae, those most likely to prove additional species are Nos. 25, 27, 31, 32, and 34.					

45.	LEPUS.	Lin.	573
-----	--------	------	-----

35Lepus texianus, Aud. & BachTexas RabbitTexas and Arkansas	617
36Lepus nuttalii, BachLeast HareColumbia River. (Very doubtful)	617

