

The Alpine winter cure : with notes on Davos Platz, Wiesen, St. Moritz, and the Maloja / by A.T. Tucker Wise.

Contributors

Wise, A. T. Tucker 1848-1928.
Royal College of Surgeons of England

Publication/Creation

London : Baillière, Tindall, & Cox, 1884.

Persistent URL

<https://wellcomecollection.org/works/e7cqjpbe>

Provider

Royal College of Surgeons

License and attribution

This material has been provided by This material has been provided by The Royal College of Surgeons of England. The original may be consulted at The Royal College of Surgeons of England. where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.


You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

14





14

THE ALPINE WINTER CURE:

WITH NOTES ON

DAVOS PLATZ, WIESEN, ST. MORITZ,

AND


THE MALOJA.

BY

A. T. TUCKER WISE, M.D., L.R.C.P., M.R.C.S.,


*Formerly Visiting Physician to the Infirmary for Consumption, Margaret St., Cavendish Sq. ;
Physician to the Western General Dispensary ; Honorary Medical Officer to the
Kilburn and Maida Vale Dispensary ;
House Physician, House Surgeon, and Resident Obstetric Officer, St. Mary's Hospital, London ;
Member of the Harveian Society ; Fellow of the Royal Meteorological Society, &c. ;
Author of "Davos Platz, and the Effects of High Altitude on Phthisis ;" "Wiesen as a Health
Resort in Early Phthisis ;" &c., &c.*

(NEW EDITION OF THE ALPINE CLIMATE SERIES.)



LONDON: BAILLIÈRE, TINDALL, & COX,
20 KING WILLIAM STREET, STRAND, W.C.

1884.



CONTENTS.

	PAGE
INTRODUCTION 	5
CHAPTER I.	
The Health Resorts—Davos Platz, Wiesen, and St. Moritz— Pontresina, Andermatt—Water and Soil in the Swiss Moun- tains 	9
CHAPTER II.	
Atmosphere in the High Alps—Ventilation of the Hotels—Dust— Therapeutic Effects of “Cold”—Sunlight—Barometric Pressure —Ozone—Oxygen... 	20
CHAPTER III.	
“Cold” as a Sensation and as Temperature—Characteristics of Alpine Climate and its Effects on the Vascularity of the Lungs and on Nutrition	34
CHAPTER IV.	
The “Drawbacks” of Alpine Winter Stations	42
CHAPTER V.	
Winter Clothing—Diet in the Swiss Alps—Exercise, Meals, Sleep, &c. 	51
CHAPTER VI.	
The Maloja—Necessity for Sanitary Precautions in the High Valleys of Switzerland—The Hôtel-Kursaal de la Maloja—Drainage— The Plateau : its Mountain-Screen—Rarity of the Föhn Wind— Duration of Sunlight, and Advantages of Solar Rays in Anæmia —Level Promenades and Healthy Amusements—Facility in Reaching Lower Levels from the Maloja—A Description of the Heating and Ventilating Arrangements used in the Hôtel- Kursaal	63

CONTENTS.

CHAPTER VII.

The Maloja Spa and Summer Cure—The Chalybeate Spring— Analysis of the Waters by Professor Reichhardt, and a Com- parison with the Paracelsus Spring of St. Moritz—Summer Alpine Climate for Early Phthisis—Duration of the “Summer Season,” and Length of Time Necessary for Renovation of Lost Health—Contra-Indications of High Elevations—Healthy Indoor Amusements possible in the Maloja Hotel	73
The Journey from England	80

CHAPTER VIII.

The Meteorology of the Maloja—Drying Power of the Air	83
Meteorological Observations taken in the Maloja during Winter— Tabular Comparison of the Health Resorts in Grisons—Com- parison in Sunshine of the Maloja with other Stations, and Mean Temperature.	

PREFACE.

IN this new edition of the Alpine Climate Series, I have made a *résumé* of my studies in the mountain health-stations of the Grisons. Although other matters have claimed attention and limited the time at my disposal, I have been enabled to include winter observations of the Maloja Plateau, in the Upper Engadine.

As the Belgian enterprise of building a Kursaal there is the first true attempt to construct a building in these cold climates, with an efficient and costly system of ventilation and warming, *without the aid of stoves*, a special portion of the book has been devoted to its description. It is to be hoped that the example set in sanitary matters, in this undertaking, may be soon imitated by hotel-companies and proprietors, so that the great blot of the "Alpine Cure Stations"—viz., faulty ventilation of dwellings and bad drainage—may meet with the scrutiny which such important points demand.

In addition to the steps taken for renewing the air in this new establishment by artificial means, a plan has been adopted, at my suggestion, of introducing ozone into the building. This is effected by electricity in connection with the electric lighting of the Kursaal, a fall of the river Inn being the motor-force employed for the purpose. It is also possible, by a simple contrivance, to place vola-

tilised remedial agents in the hot air of the *batteries* if necessary, and distribute the drug to any single chamber without infecting other rooms by its odour. The extreme cleanness of the internal atmosphere, however, renders such a proceeding quite superfluous, as the frequent changes in the air of the whole building, produce with the ozonised currents, a state of purity perhaps never before attained in any large habitation. Coupled with the large quantities of solar light received by the Maloja the results on the quality of the blood (especially in anæmic persons dwelling at these altitudes) can only be highly beneficial.


The comparatively slight attention which is given to the morbid condition of impoverished blood might lead one to estimate lowly its importance. Cases of "pernicious anæmia" are doubtless not very common, and a serious termination of the general disease rare; but the check this constitutional state gives to healthy development, physical force, and mental vigour, as well as to convalescence from other debilitating causes must never be forgotten. Nor must either the susceptibility to other complaints which anæmia favours, be lost sight of. Even in slight cases, the state of languor felt, discontent with ordinary existence, lack of interest and energy in worldly affairs, sometimes makes life scarcely worth living. The sudden change which the air, sunlight, and scenery of the Engadine give to this depression is little short of marvellous. Marked improvement usually setting in after a short stay; unlike the results of a sojourn in the South, where, not infrequently, the full force of "change" is not perceived until a return is made to the cooler climate at home.

The intention in the present volume is to exhibit the remarkable curative and health-giving properties of Alpine climate in their true light without exaggeration, or an omission of those minor details termed "drawbacks," so necessary to be portrayed in the consideration of any foreign or home health-resort.

A. T. W.

HAUTE ENGADINE,

May, 1884.



INTRODUCTION.

THE first impression on beholding a snow-clad landscape in a cold climate during fine weather is one of surprise and admiration. Conscious of excessive cold by external appearances rather than by disagreeable sensations of chilliness, one almost doubts the reality of the low temperature. A bright sun and blue sky overhead, a clear and quiet atmosphere, distant sounds transmitted to the ear through the still air, combine with the charms of the scenery to produce such buoyancy of spirits that a man is braced and invigorated for almost any exertion.

An Englishman inured to a damp and windy climate, and with an inherent love for outdoor sports and exercises, feels new life and energy as he views the glassy expanse of a frozen Canadian lake surrounded by dense forest, or makes his way through the rugged boulders and pines of Nova Scotia. A blazing sun above would almost persuade him it was summer, whilst he recognises the sing-song of the sleigh-bells in the distance, miles away it may be, and tastes the keen atmosphere of a bright winter's day.

There may be several degrees of frost, but there is seldom that piercing, raw cold experienced in England, which depends mainly on the quantity of moisture in the air and the force of the wind rather than on the pitch of the thermometer.

The effects on the Anglo-Saxon race of living in a cold climate may be seen in Canada and some of the Northern States of America, where the race is physically superior and more vigorous than that of the parent stock. This, no doubt,

depends to a large extent on the mode of life, quality of food, and outdoor exercise taken, in addition to an artificial selection of species in emigration. Nevertheless, such is the fact that these cold climates, with wide ranges of temperature, harden the European constitution by invigorating the whole frame, and rendering the body less susceptible to changes in temperature, privation, or disease.

What the eventual results will be on the English race by the growth of population and consequent increase of indoor employment, it is difficult to foretell. Certain it is that dwellers in large towns, *employées* in factories, workshops, warehouses, offices, &c., are not exposed to the most favourable conditions of life in regard to health and robust development; and although the "survival of the fittest" will select the strongest and most suitable being to withstand what may almost be termed the ravages of civilisation, even the selected one will scarcely be improved by the deteriorating effects of over-crowding, impure air, improper food, want of sunlight and equable exercise of all the organs of the body.

With examples before us of the health-giving properties of cold regions, it may seem an oversight that cold climates as "change" never received the attention they merited until within recent years.

One of the first to foreshadow the track which is now being pursued by many of the leaders of medicine in England, France, and Germany was Dr. Bodington, a practitioner in Warwickshire, who recommended "dry, frosty air" in the treatment of pulmonary consumption. In a small *brochure*, published by him forty-four years ago on this subject, he said: "To live in and breathe freely the open air, without being deterred by the wind or weather, is one important and essential remedy in arresting its progress—one about which there appears to have generally prevailed a groundless alarm lest the consumptive patient should take cold . . . The abode of the patient should be in an airy house in the

country—if on an eminence, the better. The neighbourhood chosen should be dry and high; the soil generally of a light loam, a sandy, or gravelly bottom. The atmosphere is in such situations comparatively free from fogs and dampness. The patient ought never to be deterred by the state of the weather from exercise in the open air; if wet and rainy, a covered vehicle should be employed, with open windows. The cold is never too severe for the consumptive patient in this climate. The cooler the air which passes into the lungs, the greater will be the benefit the patient will derive. *Sharp, frosty days in the winter season* are most favourable. The application of cold, pure air to the interior surface of the lungs is the most powerful sedative that can be applied, and does more to promote the healing and closing of cavities and ulcers of the lungs than any other means that can be employed. . . . Many persons are alarmed and deterred from taking much exercise in the open air from the circumstance of their coughing much on their first emerging from the warm room of a house, but this shows that the air of the room was too warm, not that the common atmosphere was too cold.” As adjuncts, he advocated the use of milk, wine, and generous diet. (a) In these remarkable sentences are clearly signalled the indications for residence at such health resorts as the high Swiss Valleys, and they anticipate in an unmistakable manner the treatment of pulmonary disease or weakness by what is now termed, somewhat euphoniously, “the high altitude cure.”

This English practitioner appears to have been the first, or among the first, to introduce to notice as a therapeutic agent “dry, frosty air,” which has proved itself to be a remedy of practical value and application in the treatment of disease.

In more recent years Dr. Hermann Weber, with eventual

(a) “*Essay on the Treatment and Cure of Pulmonary Consumption.*”
By George Bodington. 1840.

success, drew attention to high, cold altitudes as presenting favourable conditions during winter months for the arrest or amelioration of phthisis. To him undoubtedly belongs the credit of having brought the subject prominently before the profession in a clear and intelligible manner, with results on patients who had wintered in the mountains. In 1864 Dr. Hermann Weber wrote: "The question of the influence of Alpine climate—*i.e.*, localities in Switzerland elevated beyond 5,000 feet above the level of the sea—on the tendency to consumption, and on the very first stage of this disease, is one of such vast importance that I cannot help asking for the co-operation of the whole medical profession in its further examination and decision." It may appear surprising that so many years elapsed before the medical profession began to take notice of climates with low temperature. In England this can be readily accounted for, as probably the majority of persons would judge a low temperature from their own experience of cold in the mother country. For about six or seven years Davos Platz has attracted large numbers of *poitrinaires*, and from reports published by English physicians at home the efficiency of mountain air in some cases of phthisis is now undeniable. (a)

As may be expected, there are numerous sheltered spots in Switzerland where health may be sought with advantage, and where the benefits of living amongst a scanty population at a high altitude are associated with clean air and sunlight.

(a) Dr. Clifford Allbutt, *Lancet*, 1879; Dr. C. Theodore Williams, International Medical Congress Transactions, 1881; Dr. Burney Yeo, "Health Resorts and their Uses;" Dr. Drysdale, *British Medical Journal*, 1882; Dr. Marcet, Symes Thompson, and others.

THE ALPINE WINTER CURE.

CHAPTER I.

DAVOS PLATZ, WIESEN, AND ST. MORITZ.

Site and Elevation of Davos.—Water, and Drainage of Surface-Moisture in the Mountains.

THE district of Davos (an elevated valley of the Canton des Grisons, in Switzerland) extends to about fourteen miles in length, and contains between 3,000 and 4,000 inhabitants. The three miles of valley, with the health-resorts Davos Platz and Dörfli, has an altitude of 5,100 feet, and is surrounded by Alps ranging to 9,000 and 10,000 feet above sea-level. This portion of the valley is effectually sheltered from the north and west winds; rather less so from the south and east. The larger village, Platz, is situated on the north-western side of the valley, and consists of hotels, "pensions," and shops. Of these there are four or five large hotels (*a*), seven or eight smaller ones, and several pensions affording fair accommodation.

Some years ago the place was frequented principally by Germans and Swiss, but in recent times many English and French have wintered there, also Belgians, Russians, Spaniards, Dutch, and even Americans.

The scenery is picturesque and grand to a stranger in

(*a*) The Kurhaus is the largest and finest hotel, and has a *calorifère* in use for warming and ventilating a portion of the house.

Switzerland, although somewhat limited in extent, the breadth of the valley being from 500 to 1,000 yards across. Numerous pine trees cover the steep slopes, and a winding stream, the "Landwasser," drains the lake at the north-east end of the valley. The soil is dry and thin, excepting the central portion, where it is of a peaty character.

Davos Dörfli lies about one and a-half miles from Platz, towards the "Davoser See," is a few feet higher, and receives plenty of sun.

Nearly the whole extent of land is used for cattle-grazing and hay, and the sides of the valley are quite free from surface moisture.

The present system of drainage is undergoing alteration, none too soon, as within the last two or three years the increase in the winter population of Davos has been so large that this subject has forced itself on the notice of those most interested in attracting visitors to the place. When a small village, even in a healthy locality, with a scanty population and scattered dwellings, develops into a little town, unless sanitary matters go hand-in-hand with building and increase of population, dangers will arise which in the case of health-resorts may prove disastrous to their future prosperity. For the sake of Davos and the number of sick who frequent the place with the object of prolonging their lives, it is earnestly to be hoped that the new drainage system will soon reach completion. Before it is completed, however, let us trust that some hotel proprietors be encouraged to make a little outlay on *airing their houses*. This is, in reality, of more importance during the winter season than the present drainage question.

No one has at any time seriously made the attempt to uphold Davos as faultless; nevertheless, in the present instance, it will serve a very convenient purpose to recognise in the well-known Alpine station a suitable standard for

comparison. A great deal has been said for and against the place, both by those who understood something of the climate, and by others who based their observations on short visits, or on accounts gleaned second-hand. In this way, although some error has crept in, much truth and information has been elicited, greatly to the benefit of those who had never seen the place, and who have eventually experienced the curative properties of its climate.

Some misunderstanding still prevails on the problem of "cold" as a sensation, many believing that the low temperature must necessarily be unbearable, and the thought of being surrounded with snow for the whole winter awakens visions of peril to invalids.

The results which have been obtained at Davos on patients with pulmonary complaints have been highly satisfactory, and even remarkable. Dr. C. Theodore Williams, who has published the details of many cases, has shown that the principal features noticed in persons who have undergone the Alpine cure, are unusual expansion of the chest, gain in flesh, and improvement in sanguification.

Davos, like all the other mountain resorts in the Grisons, can be reached by diligence from Coire, where a few days may be spent *en route* with benefit. Landquart, the railway station in advance of Coire, is nearer Davos, but the hotel accommodation is not very good, and a stay there could scarcely be made with any comfort. In winter the post leaves Coire at 7 a.m., passes Lenz at noon, where a rest is made for dinner; at 1 p.m. arrives at Wiesen, and 4 p.m. Davos Platz. From Landquart the diligence leaves at 10 a.m., and, traversing the Prättigau valley, gets into Davos at 5.30 p.m., the distance being twenty-nine miles instead of thirty-four miles from Coire; but the Landwasser route from the latter place is well worth seeing, and the opportunity should not be missed of travelling over an interesting and wonderful road.

WIESEN (5,050 ft. above sea-level by the Calculations of the Swiss Meteorological Society; 4,990 ft. by Dr. Frankland; and 4,771 ft. from Dufour's trigonometrical measurements).

This small village stands about twenty-four miles from Chûr (Coire), on the picturesque Landwasser route. About 11 miles further on, after presenting several structural difficulties and passing through the Züge gorge, the road winds into Davos.

The position of Wiesen exhibits many peculiarities and advantages as an Alpine health resort in winter. Situated on the slope of the Wiesener Alp, facing south and protected on the N. and E. by mountain ranges of 8,000 ft. and 10,000 ft. (Sandhubel, Foppa, Alteingrat, Leidbachhorn), it is effectually screened from the cold winds of winter. Sheltered equally on the southern and western aspect by the Buhlenhorn and huge Stulsergrat (8,390 ft.), the Tinzenhorn (10,278 ft.), and Piz St. Michel (10,374 ft.), and containing the circle, by the Motta Palusa, Piz Toisa and Piz Curvèr (9,760 ft.), a landlocked region is formed, radiant in winter with dazzling sunbeams. A thousand feet below is the Landwasser chafing through a narrow course to join the Albula, a mile or so beyond.

Covering the slopes are innumerable pine tree. The odour from these is frequently perceived by new comers, and the antiseptic vapours exhaled presumably contribute to cure. The most noteworthy feature of the vicinity is an extensive plateau partially filling up the N.W. side of the basin, and jutting out towards the centre, forming an excellent and picturesque promenade. The view from the border of this plateau is seldom excelled on a small scale. Far down at one's feet is seen the floor of the gorge, with the rushing of the Landwasser audible. A peep at the Albula Valley is gained, and beyond rises the Curvèr. On the left lies the little village of Jenisberg; the Stulsergrat towering above.

Behind to the east is the Alteingrat, Leidbachhorn, &c., and on the right the two hotels separated from the plateau by a small vale down which the road descends to the Züge, passing the Känzeli waterfall.

Distant 11 miles from Davos, 334 ft. lower in altitude, and with a different contour of land, slight modifications in the meteorological details may be anticipated; whilst the general character of a cold, bracing, and stimulating climate is maintained. The chief points of variation from Davos are—its position on the side of a hill; sparser population; slightly higher and more equable temperature, with less wind. Wiesen may be considered to be about 2 or 3 degrees warmer than Davos, and, although an increase of mean temperature might lead to the belief that dangers would arise from thaws recurring, the periods of liability to actual snow-melting seldom take place with more frequency than in the latter valley. The intense heat of the sun sometimes renders parts of the main road damp under-foot when the snow is discoloured, but this could be greatly remedied by removing or covering all dirty patches as soon as formed. The plan of cleaning the roads in this manner is adopted at Davos, and would, no doubt, be undertaken at Wiesen if the number of visitors increased.

With a higher mean temperature, of course the winter is shorter compared with Davos: this might be two or three weeks at the commencement of cold weather, and about the same at the termination of the winter. During the worst time of snow-melting in Davos (which happens either in March or April, according to the mildness or severity of the season), Wiesen clears of snow rapidly; and there being no marshy valley below, like at Davos when the snow melts, any injurious consequences proceeding from dampness and evaporation from marshy land are avoided. The hard frosts at night during this period depend much on terrestrial radiation and gravitation of cold air; the larger sky expanse

presented to the Davos Valley favours the former, whilst the situation of the village is low. The bad part of the winter season is then prolonged, as the warmth of mid-day and afternoon causes a thaw. The evening and night change the wet snow into ice, and so delay its speedy disappearance. An early clearance of snow makes Wiesen a desirable locality for a change from Davos towards the end of the season. Or by judiciously-timed transits from one place to the other much of the bad weather can be escaped at either station in the months of October and March. The journey can be made within two hours, and if a tolerably fine day be selected, little risk is incurred, even in an open sleigh. Covered sleighs may be chosen; but with plenty of wraps, in bright weather an open vehicle is much to be preferred. Neither would it be at all a perilous test if the experiment were made of sending cases here when Davos was found to be unsuitable for them. If satisfactory progress were not induced, the patient would be spared the long drive to Chûr or Lanquart. The deplorable results which have taken place after sending people who were seriously ill from Davos on to Thusis and Ragatz in cold weather perhaps supports this suggestion. On the other hand, the advisability of making a rapid descent from a high altitude to the low-lands, even for convalescents is questionable. Many patients, when leaving Davos for the plains, lose much of the ground they have gained, in consequence, it may be, of the fatigues of travelling, exposure to windy or damp weather, neglect of ordinary precautions or care in personal management. Dr. Spengler, of Davos Platz, makes a statement which not only suggests the need of investigation into this matter, but also raises doubts as to whether the "high altitude cure" can be considered a permanent cure or not; in other words, if the advantages gained by breathing this mountain air are lasting, and if persons can return to their old occupations with impunity.

“Un médecin, résidant à Davos, a observé, durant une pratique de treize ans, non seulement que l'état sanitaire et les rapports de la mortalité sont des plus favorables parmi les habitants de cette vallée, mais encore que les cas de phtisie pulmonaire, en particulier, y sont complètement inconnus ou tout ou moins extrêmement isolés. Par contre, il a vu des natifs de Davos, tels que pâtissiers, cafetiers, etc., qui, après avoir échangé le séjour de leur vallée contre celui de grandes villes de l'Europe, sont revenus à Davos avec tous les symptômes de la phtisie pulmonaire. Mais lorsqu'ils eurent aspiré pendant quelque temps l'air natal, leur santé s'améliora et finit par se rétablir complètement. L'arrivée de plusieurs malades suivit de près la publicité de ces constatations, et si le succès qui a couronné leur séjour à Davos n'a pas été aussi rapide que pour les natifs mêmes de la vallée, il n'en a été ni moins heureux, ni moins complet.” (a)

It would appear from these observations that a prolonged existence in Davos does not carry with it absolute immunity from phthisis, even in presumably healthy persons with little or no hereditary predisposition to the disease. To what extent, then, must “cured” patients be liable to a return of pulmonary symptoms?—when are these likely to re-develop themselves?—when does the expanded chest shrink again? These are questions which will be answered by increased experience of Alpine climate.

ST. MORITZ.—Leaving the Davos Valley with its three stations—Davos Dörfli, Davos Platz, and Wiesen—three passes are available for a journey to the Engadine,—viz. the Fluela (7,890 feet), Albula (7,582 feet), and the Julier (7,504 feet). Strange as it may seem to those ignorant of the winter climate of these regions, a passage fraught with

but little risk can be undertaken in the depth of winter, and the journey to St. Moritz or the Maloja made in one day, even by persons who are not in robust health. Many change their *locale* in this way, and when tired of one place, seek the novelty and freshness of new quarters. There seems to be no objection to the practice, always provided the wanderer is in a fit state of health for the trip. St. Moritz lies in the Upper Engadine, over 6,000 feet above the level of the sea, is surrounded by lofty mountains, and well sheltered from wind; it possesses the climatic characteristics of an Alpine health resort; the air is similar in quality and effects to that of Davos, but being a little colder has less absolute moisture in suspension. Situated nearly 1,000 feet higher than Davos Platz, the barometric pressure is lessened to about 3 lbs in the square inch, instead of $2\frac{1}{2}$ lbs. at the latter place. The winter season is longer, and a larger quantity of snow generally falls. Dr. Burney Yeo represented the advantages of this cold and dry climate in 1866, but at that period the high lands in Switzerland were not appreciated as they are now, and the experiment of sending patients there failed through a misunderstanding of the requirements of such cases. Now, however, two hotels are open to receive those who winter at St. Moritz, two ice rinks are in constant use, and several tracks are kept in good order for tobogging.

PONTRESINA (5,915 feet), about three miles east of St. Moritz, has had a few visitors during winter-time, but it is mostly in the summer months that this attractive spot is frequented. The chief feature of Pontresina is its sunlight. The village being placed on the slope, and having a magnificent mountainous but comparatively open view to the east and south, receives the first rays of the sun at 8.10 a.m., and 8.30 a.m., during December and January; whilst the sunset varies in those months from 3.5 p.m., to 3.20 p.m.

The air is bracing and dry: the day-temperature about the same degree as Davos.

Another place which has been set forth as a winter health resort is Andermatt (4,738 feet), situated in the Urseren Valley. From the unusual facilities offered by the St. Gothard line, this spot deserves to be noticed, being only 29 hours from London, with a drive of less than an hour for mounting from Göschenen. The position of this locality is excellent for convenience in travelling either to England or to the Riviera, if the mountain air is found to be unsuitable, but no personal experiences can be given here extending over sufficient time to warrant an opinion on its meteorology, or of its fitness for a winter station.

Water and Soil.—The water supply of the dwellings at all the high level stations is upland surface water, and appears to be of wholesome and excellent quality. (a) It falls on the extensive summits and slopes of lofty mountains, mostly as snow, and undergoes filtration in its descent to the valleys, where it makes its appearance in small streams and springs. The rivulets selected to supply the houses are protected from cattle, &c., and the water conveyed in pine-tree tubes and iron pipes to the hotels. In winter it is constantly running to prevent the pipes freezing. The likelihood of contamination is then at a minimum, as no storage takes place in the houses, nor are there any habitations above the level or in the vicinity of the mountain streams. The water itself is without odour or flavour, and has a bright sparkling appearance from the carbonic acid taken up in the interstices of the rocky soil, through which the springs percolate. Although deprived of its oxygen in winter, by freezing when falling as snow, it becomes sufficiently aerated in its course to the lower elevations. At Weisen the water

(a) The discovery in the Maloja of a spring with medicinal properties, will be noticed further on.

is drawn from sandstone, which makes it considerably softer than if obtained from the Dolomite limestone, as is generally the case at these levels. An analysis has been made of the water at Davos Platz, by Charles P. Holland, F.C.S., of Manchester, who found a somewhat considerable proportion of magnesium carbonate (4.506 gr. to the gallon), but considered the water to be exceptionally pure.

The condition and nature of the soil in the Swiss mountains, and its bearings to health, present an interesting subject of study, but one which it is impossible to deal with, at any length, in the limits of the present work. One or two of the salient features connected with the temperature, and quality of the land must not, however, be passed over.

The soil of these mountain pastures is thin, of a rich, fertile character, and absorbs roughly 1.4 (*a*) times its weight of water, underneath is the solid rock, mostly Dolomite limestone, except in the centre and at the edges of the valleys or ravines, where lie the moraines of ancient glaciers composed of rubbish or deep earth intermixed with smooth blocks of rock and pebbles.

From the steep declivities on which moisture falls no injurious collection of water can take place. The drainage both of the surface and subsoil is rapid and effective. Being also of shallow depth, the danger arising from a heavy rainfall pressing up effluvia from the deeper strata is obviated; and having also no stagnant underground sheet of water, the evolution of organic emanations, forwarded by constant moisture, is not aided; or if these decompositions do occur, it is probable that absorption of their deleterious properties, is mainly affected by contact with comparatively dry earth. Vegetation also, as short thick grass covering the slopes, speedily absorbs the products of decomposition. But the

(*a*) 2,500 grms. of soil taken from a garden at Wiesen and dried, weighed 1,850 grms. On being wetted again and allowed to drain, weighed 2,600 grms.

condition of the land with which we are principally concerned during the winter season, takes on altogether a changed aspect, and one which until recently has attracted little attention. Uninterruptedly covered with snow for about four months during the year, the ground itself assumes the nature of "sub-soil," whilst the layer of thick snow above the actual service, modifies in a marked degree the effects of telluric influence.

CHAPTER II.

*Atmosphere in the High Alps.—Ventilation of the Hotels.—
Dust.—Therapeutic Effects of Cold, Sunlight, and
Barometric Pressure.—Oxygen.—Ozone.*

Atmosphere.—As the purity, freshness, and stimulating properties of the air are important points for consideration, attention must be directed to its particular characters of humidity, temperature, barometric pressure, sunlight, wind, &c.

The quantity of moisture in the atmosphere affecting the rate of evaporation from the lungs and skin is a point which requires much consideration; but at present observations have been so limited that we have not much data to go upon. Certainly, warm moist air, is sometimes very grateful to cases of congestion and irritability of the bronchial tubes; but a more permanent condition of moisture, of which our English winter affords an example, undoubtedly has an injurious influence on the majority of lung diseases; the watery vapour abstracting an undue amount of heat from the respiratory tract, and giving rise to catarrhs, coughs, or perhaps inflammations. It is a generally accepted fact that dampness of soil, apart from hereditary tendency, favours the development of phthisis. This dampness supplying the atmosphere with large quantities of moisture, shows what an important observation in climatic influences “humidity” becomes in its results on the mitigation or development of disease. Now, it may be expected that in a place where the surface formation is favourable for the rapid escape of surface water; where dry snow covers the ground, and the rainfall is slight, the reverse of this condition, so pernicious to the

lungs, is to be expected. Such is the case in the high altitudes of Switzerland; although an occasional thaw or fall of snow may saturate the air with moisture, the actual quantity of watery vapour held in suspension is extremely small, from the fact that air, at a low temperature can contain but little.

What the precise pathological effects are, of the incessant breathing of moist air is not yet quite clear; but it has been shown that bacteria and germs flourish in such a medium, whilst cold and dryness are antagonistic to their development or increase. (a) Excessive cooling of a part of the respiratory tract from moisture, on account of the conduction of heat being more free in a damp atmosphere is frequently experienced when breathing a cold London fog, the sensation of cold being felt sometimes as far down the trachea as the notch of the sternum, and even lower. The result on the skin is to abstract an undue amount of heat, and retard the normal evaporation of waste products, thereby throwing more work on the lungs and kidneys. Although there is a diminished death-rate from phthisis in some parts of Scotland and Ireland, where excessive dampness prevails, the people there are mostly employed in out-door pursuits, which makes up for a great deal in the matter of health; but if a phthisical patient courted extreme ventilation in a damp climate the result would not be so satisfactory as exposure to drier surroundings. Perhaps it is hardly necessary to mention that air containing less than 35 per cent. of moisture, with a

(a) During the progress of cutting the St. Gothard Tunnel the workmen employed were exposed to *high* temperature (80° to 90° F.) and an excessively damp atmosphere. Their health, in consequence, suffered severely, one of the principal complaints being accompanied by an intestinal parasite. Catarrhs with unusual dyspnoea were constant, and the colour and appearance of the men indicated imperfect aëration of the blood, and consequent anæmia. Their strength and appetite failed them, and the body temperature was raised. There was great mortality also amongst the horses used within the tunnel.

temperature of 60° Fahr. is much too dry for health, and would give rise to irritation of the air passages.

The freedom, or otherwise, from germs, mechanical irritants, noxious gases, or animal exhalations, necessitates but little discussion, the presence of these contaminations rendering the air impure in proportion to their quantity.

Ventilation.—Intimately connected with the subject of pure air is that of ventilation. Receiving, as it does, some scientific attention, many persons would be led to expect that a perfect system existed in any place laying claim to the appellation of "health resort." This is by no means the case in Switzerland. Like elsewhere, much depends on the individual as to the amount of air which he considers advantageous. In his private rooms he may exercise discretion at will; but in dining and sitting rooms, and especially in smoking or billiard rooms, a difficulty always presents itself. Deserving, as the subject is, of the greatest consideration in the Alps, it is to be regretted that so little real attention is paid to it. The purity of air in the interior of dwellings, both in health and disease, and especially in the treatment of chest affections, demands every scrutiny when we take into account the number of hours spent in-doors. On the shortest day of mid-winter the sun remains but five or six hours in the valleys, therefore delicate persons must be content to pass 18 and 19 hours in the house. (a) It is obvious that much of the benefit derived from open-air exposure will then be neutralised unless constant currents of fresh air are supplied for respiration.

Dust.—The great depth of snow buries all impurities, the noxious emanations of which are destroyed or lie dormant until the spring thaw. Frequent falls of snow during the winter keep any surface refuse, as it were, below ground, and

(a) For the sun-hours of the Maloja *vide* page 68.

assist to free the atmosphere also, from aërial germs and mechanical irritants, fixing them in a freezing medium; the former being destroyed or absorbed by the soil and vegetation, at the time of thaw; the latter being washed into, and mixed with the earth. Under these circumstances it is apparent that the air above is as pure as Nature can possibly produce it—clean, dry, calm, and laden with balsamic vapours from the pines. There is almost complete freedom from dust, with the exception of the carbon, &c., from the chimneys, which, of course, depends on the number of habitations, and the stillness of the air.

Taken as a whole, these conditions are healthy and exhilarating, and the mental effect of the sunlight on many patients who have been habitually spending much of their time indoors at home, is most cheering and beneficial. The value of exposure to fresh air in many diseases is becoming more widely known and appreciated, and its benefit to phthisical cases, provided they escape attacks of catarrh and bronchitis, &c., cannot be doubted, whilst the tonicity of mountain air in anæmia, debility, dyspepsia, some uterine complaints, &c., and convalescence from acute diseases, are too well known by Anglo-Indians to require repetition.

Therapeutical Effects of Cold.—Cold increases the appetite, probably from its exhilarating powers, and the requirements in the body of more hydro-carbons to meet a normal increase in combustion. A well-known writer on therapeutics remarks that the most vigorous health is maintained by a rapid construction and destruction of tissue within certain bounds, provided these processes are fairly balanced. Cold, when judiciously applied, is well-known to be a powerful tonic. A cold climate and cold bathing are tonic and bracing. The theory of the tonic action of cold may, perhaps, be stated thus:—During exposure to cold the loss of body-heat, as tested by the

thermometer, is by no means the measure of the quantity withdrawn. Many observers have shown that at such times increased combustion occurs, whereby much of the lost heat is compensated, and the temperature is maintained as soon as restored. This increased oxidation of tissue is demonstrated by the greatly increased quantity of carbonic acid thrown off by the lungs on exposure to cold (Ringer). Cold may also be considered a nervine tonic, as it stimulates the nervous system, abolishing that languor and want of energy that heat produces; the most striking example being the "plunge" or "cold douche" after a Turkish bath, and also the capacity for exertion that one possesses in a cold climate. With regard to sanguification, a low temperature causes the lungs to absorb more oxygen, and by thus inducing quicker change in the blood corpuscles, exerts a highly beneficial influence on anæmia.

It has been noticed that many patients have lost their night sweats and high temperatures in the Alps—sometimes after a few days' residence. It is not unlikely that the lower temperature of the surrounding medium, viz., the atmosphere, influences this to some extent, and also, in the case of perspiration, that its stimulating properties brace up the coats of the cutaneous vessels and promote a healthy action of the skin and sudoriparous glands. (See also page 19). Some analogy appears to exist between the action of cold on temperature, and the effects of quinine and salicylic acid: these agents often reduce a high temperature, but have a markedly diminished action on the normal heat of the body.

Moderate cold appears to influence the desire, as well as the capability, for movement and exercise; thereby maintaining the body in working order or training deficient organs up to a state of physiological activity by the stir which Nature urges, for the comfort, satisfaction, or warmth of the individual. It can be readily understood that the

employment of the various organs and limbs of the human frame is essential to their efficiency. From disuse man, has lost the grasping power of the foot, and the toes of the present species are atrophied, although the tendons remain. It becomes a question to be considered whether the sedentary occupations of modern civilisation will not eventually favour a pseudo-atrophy of organs connected with the respiratory and circulatory systems. How many men, dwellers in any of the large cities, having reached the term of years generally styled the "prime of life," could run a hundred yards with comfort, or mount a staircase rapidly without breathlessness or palpitation? With women the case is even worse, an additional curb being placed on the act of respiration by modern listlessness and fashionable attire.

The value of cold high climates in pulmonary complaints, is, in a great measure, the consequence of a gentle training of the respiratory function, the lungs becoming expanded without calling for the usual stimulus of muscular effort, which in many conditions of disease would be injurious or perhaps dangerous from the consequent force given to the pulmonary circulation.

Rattray has made observations on the weight and height of forty-eight naval cadets, aged from $14\frac{1}{2}$ to 17 years, during four successive changes of climate during a voyage. The results show that in the Tropics they increased in height more rapidly than in cold climates, but that they lost weight very considerably, and, in spite of their rapid growth, Rattray concludes that the heat impaired the strength, weight, and health of these lads. His figures seem conclusive on these points, and show the beneficial influence of cold on youths belonging to races long resident in temperate climates. (Parkes' "Hygiene," p. 436.)

Sunlight.—That light has an action on the blood corpuscles may be easily proved by observing the large

number of workers in mines and dark factories, shop girls, clerks, &c., who suffer from anæmia; likewise stokers on board ship and sailors employed entirely between decks or in the hold of a vessel, where the amount of light is necessarily limited, or perhaps entirely absent; these men, if contrasted with the workers on the upper deck, compare unfavourably in their healthful appearance, although as regards diet and sea-air, some of them are very nearly under the same conditions. Referring to Arctic experiences, there is every reason to expect the converse of the depressing influence exercised by the prolonged and intense darkness of the Arctic night.

Dr. W. Hammond contributed a paper in the "Medico-Chirurgical Transactions," on the influence of light, showing that the development of tadpoles may be retarded by depriving them of light; and that in an experiment with two kittens, where one was confined in a dark box and the other in a box to which light was admitted, the weight was perceptibly increased by light, while the growth of the other was retarded. Various experiments demonstrate that the action of light is of benefit in many conditions, anæmia, chlorosis, and phthisis, being among the number. Other factors enter largely into the cause of anæmia, &c., but the want of sunlight bears on it very strongly. The aspect of health which is created by the sun's rays speaks for itself, showing that light is a therapeutic agent of much value.

The chemical action in plants depends greatly on the presence of sunlight with the chlorophyl. Some gases also which do not combine in darkness, immediately do so on exposure to strong light.

Barometric Pressure.—In approaching the subject of barometric pressure, it will be interesting to first quote the experiences of Glaisher, Gay Lussac, and others, of its result on the action of the pulse and respiration.

Balloon ascents of	Feet.	Increase in pulse.
Biot and Gay Lussac	9,000	18 to 30
Glaisher.	17,000	10 to 24
	24,000	24 to 31

An ascent by Glaisher and Coxwell on the 17th July, 1862, gave these results:

Mr. Glaisher's pulse	76
Mr. Coxwell's pulse	74
At 17,000 feet, Glaisher . . .	100
„ „ Coxwell	84

	21st August at 1,000 feet.	At 11,000.
Mr. Coxwell	95	90
Mr. Inglelow	80	100
Captain Percival	90	88

The humidity of the air was found to decrease with the height in a wonderfully decreasing ratio, till at heights exceeding five miles the amount of aqueous vapour in the atmosphere was found to be very small indeed, (*a*)

The number of pulsations usually increased with elevation, as also the number of respirations. (*b*)

Armieux, in the case of eighty-six invalids removed from the plains to Barèges at a height of 4,000 feet, satisfied himself, after a residence of four months, the respirations were increased by two, and the beats of the pulse reduced by four. He had also found on careful examination that the eighty-six men had in four months gained on an average one inch in girth round the chest,

Dr. Kellett found that the invalids at Landour gained one inch, chiefly during the first two weeks. (*c*)

Jourdanet has asserted ("Du Mexique," p. 76) that the usual notion that the respirations are augmented in number

(*a*) "Lectures in Exeter Hall," by Glaisher.

(*b*) Glaisher's "Travels in the Air," 1871.

(*c*) "M. R.," vol. lviii., 1876.

in the inhabitants of high lands is "completely erroneous; that the respirations are, in fact, lessened; and that from time to time a deeper inspiration is involuntarily made as partial compensation."

But Coindet from 1,500 observations on French and Mexicans does not confirm this. The mean number of respirations was—

19·36 per minute for the French,
20·297 " " Mexicans.

(PARKES.)

From these and more recent observations, evidence is in favour of a slight increase both in the pulse and respirations in persons first dwelling at high altitudes, but the length of time these phenomena last has not been noted with much accuracy. It must not be forgotten, however, that the increase in the measurements of the chest, and the excursions forward of the sternum, after a short residence in Alpine valleys, may also in a great measure be explained by the gain in flesh and strength, for we know that on convalescence from many diseases at *low* altitudes this event is a consequence of returning health, and a token of general improvement.

Oxygen.—The difference in the amount of oxygen inhaled at an ascent of 6,000 feet is as follows:—

In a cubic foot of dry air at 32° Fahr., and 30 inches barometric pressure, we find 130·375 grains of oxygen. A man draws on an average when tranquil, 16·6 cubic feet of air into his lungs per hour, $130·375 \times 16·6 = 2164·2$ grains of oxygen (Parkes). An ascent (about 6,000 feet) where the barometer stands at 24 inches will reduce this 1-5th, or $\left(\frac{24 \times 130·4}{30}\right) = 104·32$ grains, lessening the quantity per hour by 432·4 grains.

Without allowing for a slight difference of oxygen at high altitudes, owing to the small amount of moisture in the air, about four additional respirations per minute would be

necessary to compensate for a barometric fall of six inches, but by experiments on animals it has been found that as long as the percentage of oxygen was not below 14, the same quantity was absorbed into the blood as when this gas was in normal proportion. The quantity of oxygen in the atmosphere surrounding animals appears to have very little influence on the amount of this gas absorbed by them, for the quantity consumed is not greater, even though an excess of oxygen be added to the atmosphere experimented with (Regnault and Reiset). It therefore does not seem at all probable that the lessened *weight* of oxygen taken into the lungs, when breathing rarefied air at 6,000 feet, necessitates any increase in the number of respirations. This is not unworthy of notice when we reflect that in phthisis and some forms of anæmia there is diminished respiratory function.

The explanation why breathing mountain air should increase the number of respirations and expand the lungs cannot be satisfactorily accounted for by the laws of mechanical pressure, as the diminution is everywhere the same, both internally and externally, and such an equilibrium of force being established disposes of any theory which attributes a freedom of the circulation, or increased thoracic capacity, directly to diminished barometric pressure.

The rhythm of the involuntary movements of the chest-walls and diaphragm depend entirely on nervous influence, and it would appear that the cause of extended respiratory movements depends on the excitation of the respiratory centres, influenced, amongst other causes, by certain fibres which run in the course of the pneumogastric. Rarefied air irritating these fibres would therefore account for the additional number of respirations and extended chest-movements. An increased proportion of blood in the lungs would also tend to this result.

In the cold high altitudes we may, then, attribute the change to both these causes, but whether this phenomenon

occurs in persons whose lungs are already quickened in action by disease is a matter for observation. In these cases the exciting cause of respiratory rhythm depends more on the proportion of carbonic acid and oxygen in the blood than on the density of the air breathed.

The following facts prove that this condition of the blood influences the respiratory movements:—1. The respiratory movements can be totally arrested if, either by a forced artificial respiration (by blowing air into the lungs) or by forced voluntary breathing, the blood becomes saturated with oxygen and poor in carbonic acid ("apnœa"). 2. Respiration becomes stronger, and the more accessory muscles take part in it ("dyspnœa"). The poorer in oxygen and the richer in carbonic acid the blood is, as, *e.g.*, on the entrance of air or fluid into the pleural cavities, causing a collapse of the lung, or when, by inflammation, &c., the lungs are unfit for respiration (Herman's "Physiology").

There are three causes at high altitudes which advance the combination of the carbon and hydrogen of the body with the oxygen of the air, viz., cold air, sunlight, and lessened pressure; therefore, it is conceivable that the additional weight of oxygen absorbed by the blood does not become such an overplus as would, when reaching the medulla, induce a tendency to apnœa, but may be sufficient to exert some inhibitory influence and balance irritation of the peripheral fibres of the pneumogastric in the lungs. Or, to summarise thus:

1. Rarefied air and larger proportion of blood in the lungs, increasing respiration.

2. Oxygen in blood, circulating to medulla, retarding respiration.

The balance being a little in favour of No. 1 in healthy persons; but in impaired lungs with quickened action (owing to excess of carbonic acid in the blood) the increased quantity of oxygen absorbed by the blood at high levels seems

to have a proportionately stronger inhibitory influence. This appears to be borne out by the fact of many patients breathing comparatively freely when leaving the plains who again experience dyspnoea on their return from the mountains.

From the experiments of Dr. Marcet at high altitudes in Switzerland and the Island of Teneriffe, it appears that more air in bulk, but less in weight, is breathed at high altitudes, and that a larger proportion of carbonic acid is excreted in the cold altitudes of Switzerland (showing that more oxygen has been absorbed).

At Teneriffe the carbonic acid was not increased in amount, whereas in the Swiss altitudes of 13,000 feet an increase of 15 per cent. was discovered. We can attribute this to the lower temperature of the latter country (as Dr. Marcet observes), and also to the larger amount of sunlight, and in a *cold* high climate there would be a greater volume of blood in the lungs to avail itself of the absorption of the oxygen.

One may be said to live quickly at these altitudes, and the most perfect health is maintained by a rapid waste and repair of the tissues of the body. This increased combustion does not mean shortened existence, but improved health, provided that repair and loss are equally balanced.

Theoretically, lungs which would be incapable of performing the respiratory functions completely at sea-level would, on the patient rising to higher cold levels, utilise more oxygen in proportion to the weight of the air inhaled the higher they ascend, within ordinary limits, and that the functions of the skin would also be promoted by reduced pressure, favouring the action of osmosis. (*a*)

(*a*) There is evidence that the interchange of gases between the air and the blood through the skin has an important share in keeping up the temperature of the body, and we find the temperature of the surface much elevated in cases of pneumonia, phthisis, &c., in which

Although, perhaps, not quite so much fresh air is breathed in a cold climate by those who sleep with closed windows as is breathed in the warmer health resorts by persons who sleep with windows open, the three conditions of excessive sunlight, cold atmosphere, and lessened barometric pressure, may partly compensate for the advantage in that respect. Indeed, it is difficult anywhere to persuade many patients to ventilate their rooms at night.

Before leaving this subject, it will be interesting to note the diminution of barometric pressure, in the Engadine, viz., 6,774·76 lbs. off the whole of the human body. Although the considerable reduction of three tons (*a*) taken off the fluids, and solid parts of the skeleton carries with it no remarkable phenomenon, it is not illogical to assume that such a declension in the weight of the atmosphere, must exert some peculiar powers. Whether these affect physiological actions, or influence the animal economy in any way other than favouring the interchange of gases; increasing the beats of the pulse and the number of respirations; may be elucidated by future observations.

Ozone.—This peculiar condition of oxygen is generally found in healthy localities. Sea-air and mountainous tracts, especially give indications of its presence, whilst in densely populated districts or unhealthy places, ozone is observed only in small quantities. In a work on Ozone and Autozone, Dr. Cornelius Fox states as follows: "The salubrity of a town or city may be pretty accurately estimated by the effects of its air on ozonoscopes, as the feebleness and

the lungs seem to perform their function very insufficiently (Carpenter's "Human Physiology").

(*a*) The barometer is lowered about 6 inches, $\frac{1}{5}$ th of the capacity of the mercurial column at sea-level. A cubic inch of mercury weighs 3,433·5 grains, or ·49 lb. The surface of an ordinary sized man is about 16 square feet: therefore the calculation can be easily made.

sluggishness of the re-action is a very good gauge of the amount of impurities which it contains. Ozone is a deodorising and purifying agent of the highest order, resolving and decomposing into innocuous forms. The oils of the cod's liver, the cocoa-nut, and sun-flower, *when ozonised*, have been found, by Drs. Theophilus and Symes Thompson, to be very useful in reducing the rapidity of the pulse, and exerting at the same time an invigorating influence on the heart's action in consumption. Last in order, but first in importance, ozone has been considered to be probably concerned in a work most gigantic in magnitude and of vital consequence. It has been thought to be influential in the modification of climate, to exercise a beneficial action on animal and vegetable life, and to be indispensable to the relief and cure of functional disorder and disease. It has been doubted whether life could continue to exist on this planet, according to the present constitution of terrestrial laws, if the formation of ozone should cease in nature."

CHAPTER III.

“Cold” as a Sensation and as Temperature.—Characteristics of Alpine Climate, and its Effects on the Vascularity of the Lungs and on Nutrition.

THOSE who are unacquainted with high altitudes in winter may perhaps be inclined to judge the sensations at a high cold region from an English standard of cold, thinking possible that twenty degrees of frost signifies twenty degrees of chilliness, and that any temperature below freezing point would be likely to cause discomfort to delicate persons. A brief explanation may tend to correct this view. The body can be deprived of its heat in four different ways:—

1. By conduction, or contact with colder substances, either solid, fluid, or gaseous.
2. By evaporation from the surface of the skin, and the mucous membrane of the respiratory tract.
3. By excretory matters leaving the body ; and
4. By radiation.

Now, although it is possible from the hygrometric state of the atmosphere that an additional quantity of moisture is evaporated from the skin and respiratory tract, at high levels, (a) this variation plays a very minor part in sensibly reducing the temperature of the body compared to the abstraction of heat by conduction ; or, in other words,

(a) It must not be supposed that evaporation from the body depends entirely on the percentage of humidity in the air. The conservative and balancing agencies of physiological action amongst other things constrict or dilate the cutaneous capillaries in response to cold or heat, rest or exertion, &c.

contact with cold air in movement. This latter cause is the one which principally bears on the question of sensation, inasmuch as cause No. 3 is too insignificant to be felt, and No. 4 can be guarded against to a great extent by clothing.

The physical sensibility of cold is produced by the amount of heat rather suddenly abstracted from the body (which does not always depend on the temperature in contact with it). For example, if the hand be placed on fur at 30° Fahr., it feels warm in comparison with iron at the same temperature. The former being a bad conductor—owing to the *motionless* air in its interstices—does not abstract much heat from the hand; the metal, being a good conductor of heat, appears intensely cold to the touch.

If, therefore, cold, motionless, dry air surrounds the body, heat is not abstracted nearly so readily as it would be by somewhat warmer air in movement. It must be remembered that the sensation of cold cannot be accurately gauged by reference to the thermometer. Two other conditions are intimately connected with temperature in causing impressions of cold or heat—viz., wind and moisture, for it is these that cool the body, by conduction. If their temperature is lower than ours, they appear colder than they really are, because from their conductivity heat quickly passes away from us.

In the high valleys of the Alps, although the thermometer may register some 15° or 20° of frost, this low temperature is by no means disagreeable, as the calm air and intense solar heat enable many persons to sit in the open and bask in the sun during the depth of winter without feeling the slightest sensation of chilliness. Even excessive tanning and reddening of the skin takes place with almost everyone who takes plenty of outdoor exercise: ladies, who are generally well protected by sunshades or umbrellas,

do not escape a healthful aspect. This is mostly owing to the reflection of light from the snow, which, coming in upward and parallel directions, cannot be well screened from the face. The habit and necessity of wearing smoked-glass spectacles also enables persons to face the glare, and thereby receive a much larger proportion of light than in England. (*a*) According to Dr. Cornelius Fox, ozone also causes a healthy colouration of exposed parts of the body. The speedy tanning that one undergoes when crossing a mountain pass, or driving in an open sleigh, supports this view. As more ozone is brought into contact with the skin by movement through air when driving, or on the passes where there is generally a breeze present, so a greater colouration ensues: whereas one may be exposed to the influence of wind in large cities (where there is an absence or great diminution of ozone) without wearing a rosy or tanned appearance.

The general climatic characteristics are—

1. Dryness of the air (*b*) and its comparative freedom from mechanical irritants, germs, and noxious gases, (aseptic air).

(*a*) It has been frequently noticed that dark-complexioned individuals become sunburnt more readily than “blondes.” This depends principally on the sensitiveness of the retina and the colour of the eyes. For instance, “fair” people cannot face the light with such ease and comfort as those who have plenty of pigment in their irises; for the pigment absorbing the rays of light, protects the retina, and even enables some with very “dark” eyes to gaze on the sun itself. On the other hand, a person with a grey or pale iris averts or screens the eyes from the sunlight as much as possible, and in this way escapes the effect of the rays on the face.

(*b*) The drying and preserving of meat hung in the air, has been alluded to as an illustration of the dryness of these climates. This takes place in an analogous manner to the drying of turtle in the sun in the West Indies, viz., a rapid evaporation of moisture before putrefaction takes place. In the Swiss mountains putrefaction is so long delayed by a low temperature, that thick masses can be gradually dried.

2. Profusion of sunlight, with a low temperature.
3. Diminished barometric pressure.
4. Ozoniferous atmosphere.

The results on pulmonary complaints, anæmia and allied disorders, may be stated thus :—

1. Lessened irritation of the respiratory tract from absence of dust.
2. Vaporisation of morbid secretions in the lungs, promoted by reduced barometric pressure and dryness of the atmosphere.
3. Increased oxidation of blood and tissue, from sunlight, cold air, and reduced pressure.
4. Increased quantity of blood circulating in the lungs—caused by the low temperature—the freedom of the circulation being aided by extended chest movements.
5. Increased activity in the pulmonary lymphatics (depending on circulation and expansion) and a general improvement in nutrition and glandular secretion; also an exhilarating effect on the nervous system.

Some of these results are obtainable under no other conditions than those presented at high cold regions. With regard to the increased quantity of blood circulating in the lungs (presumably influencing the nutrition of those organs), it may be contended that this is not a desirable sequence. Perhaps it is not in hæmorrhagic phthisis; but in some other forms, especially early tubercular deposits, it would not seem to be disadvantageous. What would lead one to suppose this is the rare occurrence of tubercular phthisis in persons affected with mitral disease. Even when hæmoptysis takes place, and when some of the blood presumably gravitates into the air cells, tubercular disease rarely follows; whilst, on the other hand, phthisis is not an uncommon consequence after hæmoptysis from other causes. This would appear to indicate that a general hyperæmic condition of the lungs

impedes the deposition of tubercle and restrains phthisical processes. Support is also lent to this view by tubercle generally attacking the apices of the lungs, which parts contain rather less blood, owing to gravitation.

Conversely, where the quantity of blood circulating in the lungs is lessened, as in hot climates, phthisis is frequently seen to run a very rapid course.

It is not improbable that this increased volume of blood moving in an impaired and imperfect lung at a high altitude plays an important part in the nutrition of the tissues and in augmenting the movement of lymph through the pulmonary lymphatics, so removing by absorption many of the smaller morbid cell growths. With the lessened barometric pressure accelerating the action of osmosis, compensation, to some extent, is made for the loss in pulmonary capacity, for we know that a more rapid interchange of gases takes place under reduced pressure.

On the other hand, the emphysematous signs presented by patients who may be said to be cured, after a prolonged residence at high stations, may seem to contra-indicate any theory based on this assumption. It must be conceded, however, that the expansion of the chest obtained, it is doubtful if emphysema occasions such compression of the pulmonary capillaries as to decrease the *whole* volume of blood circulating in the lungs.

That the lungs contain more blood in a cold climate is pretty clear, if we accept the evidence of Dr. Francis (Bengal Army), who found from a large number of observations, that the lungs are lighter after death in Europeans in India than the European standard. Parkes confirms this, and also Rattray, in his observations of diminished respiratory function in hot climates.

When the activity of any organ of the body is augmented, more blood is attracted to it than when at rest or during lessened exertion. By breathing rarefied air at high levels

the respiratory movements are usually quickened and extended, *especially on taking exercise*.

The liability, also, of the natives of these high valleys to pneumonia, whilst exempt from phthisis, would seem to point to some alteration in the vascular condition of the organs affected.

What result any variation in the vascularity of the lungs would have on the bacillus tuberculosis is rather premature to surmise. No bacilli have, however, up to the present time been discovered in the blood of tubercular subjects. It appears, therefore, that either they do not enter the vessels, or, if entering, are changed in character or destroyed. That the state of the blood, chemical or pathological condition, or functional activity of the tissues must be agencies governing the suitability of the soil for the reception of the bacillus, is supported by the fact that infection is very rare, although there are numberless cases in which bacilli have undoubtedly been inhaled.

There is, however, no decided criterion for the determination of the question whether—

1. A major quantity of blood in the lungs, such as may be diffused through these organs in a high cold climate, with slightly augmented freedom (from acceleration of respiratory and cardiac movements) and more complete oxidation, is or is not less disposed to bring about pulmonary hæmorrhage than—

2. A minor proportion of blood at sea-level, circulating with less facility, and not so effectually oxidised.

The problem also appears to present itself as one of “nutrition of the lung,” viz., whether slight hyperæmia (under those circumstances) is not a more desirable sequence than the inclination to slight blood stasis of No. 2.

It is well known, also, that the hygrometric and barometric states of the atmosphere modify the process of evaporation from the lungs and skin. The evaporation of

morbid secretions in the lungs was pointed out in 1881 (a) as being one of the circumstances which probably has a favourable bearing on phthisis. The process of evaporation in dry climates, acting on ulcers, cavities, or suppurating surfaces, if not analogous to the dry treatment of wounds (so successfully carried out by Mr. Gamgee, of Birmingham), brings about a less moist and watery condition of the secretions from diseased bronchial tubes or cavities of the lung; virtually lessening expectoration without the patient undergoing the effort of coughing.

The effects on the body of sunlight and reduced pressure are doubtless those facilitating the interchange of gases in the blood and tissues, whilst the cold air necessitates the requirement of a larger absorption of oxygen and assimilation of hydro-carbons to maintain the heat of the body. It may be conjectured that this contributes to the sudden and considerable push given to nutrition on arrival at a cold high altitude, when the appetite is, in most cases, at once improved in a remarkable way, and animal food that could hardly be thought of previously without disgust is eaten promptly. Where improvement begins it is difficult to say; indeed, it is only to a combination of causes that the variety of effects can be attributed. This push to nutrition is a reliable feature in the first evidences of progression, and assures a certain amount of hopefulness in the case.

The exhilarating feeling produced by the *consciousness* of moving about amid snow and ice, without taking cold or feeling pinched, is not to be despised as contributing towards cure. The contrast of this with the life of an invalid in England during winter, where every change of weather has to be guarded against, is so marked that the hope of recovery presents itself, and despondency is banished.

There is every reason to suppose that under the many favourable circumstances presented by these climates the

(a) "Davos Platz, and the Effects of Altitude on Phthisis."

treatment of suitable cases of anæmia and its allies, scrofula, consumption, and affections of the chest, and some cachetic states of the system, can be undertaken with greater confidence, and those measures which have of late years prolonged many valuable lives, are certainly more likely to be efficient and successful, when supported by the curative effects of mountain air; whilst for the somewhat minor maladies, such as debility, either from physical causes, or from mental fatigue and worry, malaria, &c., some forms of dyspepsia, chronic discharges or suppurations (that do not incapacitate the patient from taking gentle exercise), and during convalescence from many acute diseases, the renovating power of these remarkable climates are, in suitable cases, doubtless far superior in rapidity of effect, to warmer and lower latitudes.

It is not for a moment implied that climatic conditions, grateful at all times in health and disease, are not found in certain localities, where the range of the thermometer is generally from 50° to 65° , or thereabouts. This equality of temperature with dryness, &c., enables patients to be constantly in the open air, and, if not out of the house, to be able to avail themselves of a system of "hyperventilation" day and night; but, on the other hand, it is now well known that other places can be found as favourable, if not more so, for these complaints in cold climates, although the range of the temperature is not so limited; whilst the languor and depression of strength felt in warm climates is altogether escaped.

CHAPTER IV.

Including the Drawbacks of the Alpine Winter Stations.

AFTER leaving Coire (*a*) for the ascent to Churwalden (4,976 ft.) and Parpan (4,938 ft.) the beauties of the mountain journey begin. The rise in the altitude gives a sensation of lightness and elasticity to the frame. All the surroundings cheer the spirits—the tinkling music of the cow-bells, the autumnal changes in foliage, golden larches and aspens contrasting with the dark green of the firs, topped here and there by brilliant sun-beams, undulating into gorges and hollows of obscure impenetrable shades—whilst in the distance are occasionally seen the higher peaks crowned with dazzling snow of snowy whiteness. After a railway journey, perhaps from foggy London, these scenes alone stimulate and recruit the strength. It is well to be cautious at this stage, that the powers are not overtaxed by fatigue; for this sudden vigour is not yet permanent, but may permit a delicate person unconsciously to out-run the supply of force. It is, in some instances, desirable that the journey from London, for example, should be broken at various points, so that the patient shall not reach his destination in a wearied and jaded state.

The transformation scene after the winter snow has fallen intensifies all the virtues of mountain air. With the sudden covering of the earth by a non-conducting material an entirely new condition of the climate is established. Fresh breezes have died away, packs of cumulus clouds have been

(*a*) Davos Platz and Davos Dörfli can be reached from Landquart, the station before Chûr (*vide page 11*).

condensed into snow, the moisture in the atmosphere is of the smallest quantity, whilst the sun shines with the lustre and power of the South, and the sky presents an almost unclouded surface of azure blue.

There are, however, many cases of advanced disease buoyed up with false hopes, in real ignorance of their true condition, who, instead of seeking the soothing and languid influences of a Southern clime, or the comforts of home, venture to face suddenly the stimulating and force-producing powers of the keen mountain air.

To countenance the reception of unsuitable cases with the chance of recovery that occasionally presents itself in pulmonary disease, is not only reprehensible and bad, but detracts from the value of "climate" as a remedy of, even limited application. We cannot but think that this is one of the reasons which has somewhat retarded the frank reception by the medical faculty of Alpine regions, as offering a means of cure to many who can arrange to spend a winter amidst the snow and sunlight of the Swiss mountains.

The great drawback to most of these health-resorts, as with many others in the South, is defective sanitation. As long as overcrowding does not take place, the cold and snow will, to a great extent, mitigate the evils of bad drainage and impure emanations; but when hotels get filled with a large number of visitors, many of whom may be in delicate health, the air within will not (except by artificial ventilation and good house drainage) be free from the usual indoor impurities, as sewer-gas, kitchen and basement air, exhalations from the lungs and skin of organic matter, scales of epithelium, fibres of cotton, wool, wood, &c.; the products of combustion from gas, lamps, and candles; bacteria and fungi; and, what is, perhaps, more important still in the case of delicate lungs, the bacillus tuberculosis floating about in the air. Patients must, therefore, exercise their own discretion in opening windows and airing their rooms as much as possible. A fair

quantity of fresh air can always be permitted to enter the bed-rooms, according to the desire of the individual; but in dining-rooms, billiard, smoking, and conversation rooms, the means of admitting pure air and providing for the escape of foul are very imperfect. Until appliances are introduced into the hotels (as is done in the Maloja Kursaal) to clean the air, warm, moisten, and medicate it, if necessary, the high altitude stations can scarcely be designated "air-cure places," for most of the patient's time is spent indoors, where the conditions in some of the hotels are no better than at home—perhaps worse, if the house should be full of people.

By boxing-up soiled air for the sake of warmth, the pollution of the interior of dwellings by organic matter can be readily discerned by anyone who is familiar with its peculiar odour. In crowded places of entertainment, on first entering from the outside, a faint, musty smell is not uncommon. This indicates an atmosphere loaded with organic impurities, and the immediate effects of such commence with faintness and headache. The quantity of carbonic acid evolved is in no sense so dangerous as the animal matter, although it gives a good clue in estimating the quantity of the latter poison.

Where stoves are in use, attention should be given to them to see that servants do not entirely close the valve which regulates the flue. Occasionally this is done, when the fuel is consumed and the remains are glowing red, with the intention of economising heat, which it effectually does, by preventing air passing through and cooling the stove; but the most poisonous of all gases, viz., carbonic oxide, is slowly evolved into the room from the dying embers and quickly gives rise to headache, which is not rapidly shaken off even by fresh air. (a)

(a) Carbonic oxide replaces oxygen in the blood, and cannot be again replaced by oxygen, but has to be slowly converted into carbonic acid, before elimination.

It is essential too, that some special employment should be undertaken when persons remain several months abroad: languages can be studied, or the mind occupied with work of some sort. By this means time does not drag heavily, *ennui* is not experienced, and restoration to health is not retarded by dejection of spirits.

The travelling also has to be considered. Those who are in pretty fair health can make the journey in three days, either to St. Moritz, Davos, or Wiesen, but a longer time is recommended for those whose health does not admit of prolonged confinement in railway carriages. The journey may be broken at Paris or Brussels, Bâle, and Chûr.

On first arrival it occasionally happens with some people that they are unable to obtain a good night's rest for the first night or so. This depends on the sudden rise which they have made to high regions. The difficulty soon passes off; but in the case of sensitive persons who sleep badly, it may be met by remaining a few days at Chûr (1,936 ft.). The stay there will frequently help in training the system to the new conditions. It must not be forgotten that the altered form of bed may, in many instances, prevent sound and refreshing sleep. The wedge-shaped hair bolster can sometimes be removed with great advantage to the sleeper, and extra clothing be placed over the shoulders and upper part of the body, which is usually not covered by the eider-down quilt. In addition to equalising the bed temperature in this way, the head should not be raised at too great an angle from the chest and shoulders, or compression of the vessels of the neck will ensue and so impede the ready return of blood from the brain, keeping this organ active and full, when it should be in the anæmic state which usually accompanies sleep.

A little discomfort is sometimes caused by chilblains during the winter time. To prevent these, walking exercise should be taken immediately after breakfast, and the endea-

vour be made to keep the feet warm throughout the day by this means. The parquet floors, although healthy and advantageous in many respects, are, in a great measure, the cause of cold feet and the source of chilblains, therefore all the corridors and bed-rooms in hotels should certainly be carpeted in these climates during winter time.

What has been advanced as an objection to Alpine stations is the use of the German stove as a means of warming the interior of houses and hotels. The fault, however, does not lie so much with the ponderous German *Ofen* as with improper management in neglecting to provide sufficient ventilation for the rooms and moisture for the air. An improvement on these stoves is the steam reservoirs in use at the Kurhaus, Davos Platz, the admission of steam being under the control of the occupant of the room in which they are placed. But this plan also raises the temperature without raising the "dew point," or, in popular language, "dries the air," creating a necessity for a small supply of watery vapour to render the atmosphere fit for healthy and agreeable respiration; for, although dryness of the climate is one of the main features of the Swiss Alps, a limit can be reached beyond which a dessicating effect may be produced.

The introduction to the bright and calm winters of these regions is sometimes a little unsettled, but not dangerously so. Falls of snow occur and are apt to thaw again; the roads, however, soon dry, by reason of the sloping land and rapid evaporation. Acclimatisation has frequently been held up as a pre-requisite for visitors, but, as the winter is generally bright, calm, with dry cold and plenty of sun, the risk of arrival, even after the commencement of winter, is in no wise greater than during the autumn. To become acclimatised for one of the best seasons of the year, and one which is regarded as a "cure" period, requires no inurement—many persons even arriving in mid-winter and crossing a mountain pass without inconvenience or harm resulting.

The selection of an Alpine station should certainly be influenced by the opinion of physicians at home, even where the intention is entertained of returning for a second or third winter. Perhaps this may seem to be written in the interests of the medical profession, but those who have spent three or four seasons at Davos will have observed that some cases return again who would do better in a southern clime. An over-estimation of their own recuperative powers, or an enthusiastic belief in the potency of the charming Alpine climate to sweep away *all* maladies, is possibly the cause of an injudicious decision.

Children over three years of age do extremely well in the Alps, make healthy blood and muscle, gain flesh, and expand their chests. Under this age, it is doubtful if sufficient exercise can be taken by a child to ward off the cold, without being swathed to such an extent in furs and flannels as would impede the free motion of the limbs and thorax, so essential to development in childhood.

It is also a moot question whether some invalids should not quit the Alps immediately the end of the winter seems approaching, as the changeable weather, with wet roads, winds, &c., is likely to upset many of the benefits gained during the dry season. In this case, it would probably be well to go south, Lugano (982 ft.), Como (705 ft.), Sorrento, &c.; but if spring has set in, the shores of Lac Léman (1,230 ft.) offer a fairly safe change. Thusis (2,448 ft.), Ragatz (1,709 ft.), Mels (1,637 ft.) are reached easily, but a residence at these latter has seldom been attended with much improvement in health. Promontogno (2,700 ft.) is very convenient for the Engadine; its position and climate go a great way in recommending it as a halting-place. A month might be spent at any of these, previous to returning home, provided the weather is fine, but it is well to be impressed with the necessity of the great personal care needed during this migratory period, especially with regard

to clothing. Thick flannels, socks, &c., should not be dispensed with hastily, and it would be well to guard against the change from the calm regions by adopting outer garments of close texture, impervious to the chilling effects of wind.

Any remarks on the climate of Switzerland would be incomplete without reference being made to the föhn wind. (*a*)

Altitude and low temperature modify, to some extent, the noxious qualities of this wind.

The föhn is a moist southerly aerial current. Its temperature in Switzerland becomes elevated from various causes, the chief of which is probably atmospheric pressure. By this accession of heat the capability of the föhn for holding more moisture in suspension is at once augmented. The "absolute" humidity may remain the same, but by a rise in temperature the percentage of relative humidity (*i.e.*, humidity in relation to the point of saturation), is considerably reduced, for warm air can contain more watery vapour; therefore, the föhn seizes with avidity on any moisture which is present. The extreme dryness is again advanced by the stoves in use, further raising the temperature to a dessicating point. In this way many bad symptoms are aggravated and vaporisation of organic

(*a*) Dr. Wild says, the föhn as such is known only in the north-eastern valleys of Switzerland, and it is there distinguished by its great heat, and still more by its peculiar dryness before which the snow disappears both by rapid melting and also by that rapid evaporation which has obtained for it the appropriate name of the snow-eater (*Schneefresser*).

But while the föhn proper is blowing in the valleys of the north-east eating away the snow in winter, or in summer and autumn drying the hay or ripening the grapes, over the south-west of Switzerland a warm and wet wind blows, which precipitates its moisture in a heavy down-pour, and floods the country with rain and melted snow. The distinctly föhn stations named by Dr. Wild are, Glarus, Auen, Altdorf, Engelberg, Schwyz, Chûr, and Klosters.

matter takes place, which is doubtless re-inhaled by the occupants of a room badly ventilated or unprovided with sufficient water near the stoves to bedew the air.

The nervous system of most persons becomes remarkably depressed by this wind; the inclination to undergo exertion is diminished; sleep and digestion are disturbed; the animals, too, seem to suffer—effects which closely resemble those caused by the “Vent d’Espagne,” in the Pyrenees (Hermann Weber). Indisposition is usually attributed to this wind at Davos, and in several instances with justice.

After two or three days, sometimes longer, the second phase of the föhn is developed. The thermometer falls and with the cooling process the air is gradually brought below the dew-point, and what a few days before was an exceedingly dry wind, becomes converted into an atmosphere saturated with moisture, which falls as snow or rain, according to the time of the year, and temperature.

The days of invasion by the föhn wind are about equal at Davos and Wiesen. On the Maloja plateau, 1,000 feet higher, this wind is rarely felt, and its noxious qualities are greatly modified by the higher altitude, extended width of the valley, and the proximity of immense glaciers to the S. and S.W. (*vide page 66*).

The teeth are apt to cause some trouble in the Alps, if caries be present. A good dental surgeon should be consulted previous to leaving home, and the habit be acquired of breathing through the nose as much as possible, even when taking exercise. The rapid changes in the temperature of the teeth occurring to persons who eat, drink, and breathe with the mouth, give rise to a variety of derangements, especially in cold climates. Amongst the natives of Switzerland the noticeable deterioration of the teeth may probably be due to the consumption of the rough wines of the country, which doubtless contain much acetic acid, caries being very marked in the lake districts.

In conclusion, it may not be superfluous to remind those interested that the high cold regions are not yet proved to be the best for *all* cases. The varieties of chest and other affections and the individual differences of affected persons denote that one general climatic panacea is quite inadmissible, and what may be a suitable climate for one, may prove of very little value to another. Whilst some misconception prevails on these points, cold mountain air will not receive the appreciation it merits. It would be inconsistent to contend that any particular health-resort had not its drawbacks. It is misleading to make light of these, and most essential in the recommendation of foreign health-resorts that their disadvantages be clearly stated, as well as the probable benefits likely to accrue to persons visiting these places. Much disappointment is thereby averted, and many, who otherwise would expect impossibilities, do not become discontented. In spite of much misunderstanding and opposition, high altitude treatment is gaining ground with the medical profession, and will, no doubt, some day take its right place in climatic therapeutics. Time is needed for the subject to be well threshed out by advocates and opponents alike.

CHAPTER V.

Winter Clothing.

THE selection of suitable clothing for an Alpine climate will contribute greatly to health as well as to comfort.

The first advice tendered to ladies who seek restoration to health from lung disease is:—Abandon corsets absolutely, and wear the loosest *ceinture* on the waist. This injunction cannot be emphasised too strongly. One prominent sign of what may be termed “cure” of phthisis, at high altitudes, is the almost invariable expansion of the whole or portions of the chest walls. To impede such a beneficial result is to limit the advantages of breathing mountain air. No one could wittingly be guilty of such an irrational procedure as to curtail the actual physical improvement they travel many miles to secure. Yet usually, whilst the upper regions of the chest are allowed free play, the most mobile section, the diaphragm, is fettered and confined in its movements by an unyielding investment pressing on the abdominal viscera. In this way an opposite conformation is maintained against what physiology and common sense would indicate; for in many instances where damage exists in the superior thoracic regions, limitation of movement is desirable in *these* situations, rather than in the lower and sounder parts of the lung.

The substitute for corsets should be a thick flannel waistcoat or jersey. Flannel should also be worn next the skin over the trunk and extremities. Chamois leather may be substituted when over-sensitiveness of the skin renders the use of flannel unbearable; generally, however, a fine texture can be procured which is not too irritating. A good stock should be taken, as frequent changes are beneficial.

The coverings for the feet and legs require attention. In all cases woollen socks or stockings are indispensable. Flannel pants should be made long enough for socks to be drawn over them. By enveloping the whole body in flannel the patient is spared the necessity of loading the exterior with thick weighty dresses or heavy overcoats, which fatigue the wearer and confine the act of respiration. Those who are liable to suffer from cold feet, or are susceptible to chilblains, will find thick worsted socks and cloth gaiters a necessity, or thickly-lined cloth boots, to come well up the leg. As a prophylactic against chilblains, bathing the feet in salt and water, avoiding tight boots, and changing the socks whenever the feet are damp from perspiration, &c., will carry many through the winter of a cold climate, who might otherwise suffer much inconvenience and pain from these minor but troublesome ailments. Boots should not be over thick and heavy, which renders them inelastic. There is no utility in having clump soles, as leather gets saturated quickly with moisture and takes a long time to become thoroughly dry. Ordinary stout skating boots, with broad toes, low heels, and plenty of room inside them, which may be filled up with a cork sock, are suitable both for ladies and gentlemen; they should be well greased every day and always changed immediately after a long walk. Goloshes are useful for short journeys, but cannot be recommended for constant use, on account of confining the perspiration. All clothes should be light in colour, as near grey as the taste will allow. A light colour does not absorb nor radiate heat so much as black; therefore a wearer of grey will be comfortably free from the heat of solar radiation and warmer in the shade. Waistcoats should be lined in the back with flannel, as with modern garments all the protection is in front of the body. (a) If

(a) With double-breasted apparel a man may have from ten to thirteen thicknesses of woollen clothing in front of him, whilst his back is covered with three layers of wool only.

men endeavoured to dispense with the use of braces greater play would be given to the muscles of inspiration. Hats may be of straw for the sunny weather, but light-coloured thin felt answers very well. Furs should not be worn when taking exercise. The ladies' fur tippet can be discarded as superfluous at any time, from the fact of its covering the shoulders only, causing that part of the body to become overheated, and therefore liable to chill. A short list is appended of articles found to be serviceable for these climates :—

One fur or railway rug. One warm ulster, with hood or fur-lined coat for travelling or sleighing. One thin overcoat, a Shetland shawl. Two suits of clothes (waistcoats lined). Thick flannel vests and flannel shirts. Flannel night-shirts. Half-a-dozen or more worsted socks. Flannel pants. Cloth or fur gloves with gauntlets. Cork socks. A pair of dark neutral tint spectacles. A few rough bath towels and a flesh glove. One woollen muffler. Two pairs of stout boots (one pair smooth for skating, the other with a few spikes in the soles, not thickly studded with nails, or the snow will ball on them). A pair of leggings are required for "coasting," unless long boots are taken. One pair of shoes or thick slippers with heels can be worn indoors with spats over them. Boot-laces, dubbin, and skates may be added, and for ladies a sunshade and goloshes with or without cloth tops. A fur foot-warmer or large fur-lined boots are also very useful.

Diet in the Swiss Alps.—The food at St. Moritz, Davos, and Wiesen, as almost everywhere in Swiss hotels, consists largely of azotized or nitrogenous matter. Many who have visited Davos, for example, will recollect that much lean meat is eaten, whilst the proportion of animal fats is very small.

It is essential to the health and well-being of individuals that a proper proportion of nitrogenous and non-nitrogenous food should be consumed. The latter consists chiefly of

starch, fats, sugar, saline substances, and water, which in the ordinary way with meats form the mixed dietary most suitable for man; as the structure of his teeth and past experience indicate. But whilst an almost exact estimate can be formed of the elements necessary for the system to ingest, it must not be forgotten that the quantity and kind of food taken will depend very much on the condition or idiosyncrasy of the consumer. As a general rule, few delicate persons would be capable of effectually digesting the quantity of bread, pastry, milk, and root vegetables, &c., which would be requisite to constitute with the meat eaten, a fair combination of nitrogenous and non-nitrogenous matter, and as there is a marked absence of palatable fat at these places, a little consideration of the means for balancing the needful combination, will conduce to the recovery of health and avert many ill-effects arising from perverted nutrition—as dyspepsia, troubled sleep, a loaded tongue, vitiated secretions, &c.

“Many people,” Dr. Pavy remarks, “seem to look upon meat almost as though it formed the only food that really nourished and supplied what is wanted for work. The physician is constantly coming across an expression of this view.”

The greatest importance must be attached to the use of fats during winter in the Alps; for it is well-known that the inhabitants of Siberia, Greenland, &c., and all cold countries, eat enormous quantities of these heat-producing materials, without which they would be unable to resist the intense cold of their frigid climate. Sir John Ross observes—“It would be very desirable, indeed, if the men could acquire the taste for Greenland food, since all experience has shown that the large use of oil and fat meats is the true secret of life in these frozen countries.”

Dr. Cheadle also lays much stress on the value of fats in cold climates. “One effect of the cold was to give a most ravenous appetite for fat. It is the most valuable part of food

in winter, and horses and dogs will not stand work in the cold unless fat." (a)

Besides forming the chief articles of diet which are required for a calorifacient or heat-producing agent, they may almost certainly supply a pabulum for the oxidising process of fever in phthisis to act upon, thereby not only restricting the waste of tissue in that process, but perhaps in some way diverting its occurrence, as is witnessed when cod-liver oil is taken. With an increase in animal food, which can be readily eaten at high cold levels, the appetite being sharpened by exercise and low temperature, the necessity for fats is by no means abated; on the contrary, a physiological demand is created for additional food capable of undergoing the process of oxidation, which cannot be wholly supplied by the lean meat or by starchy and saccharine bodies.

"It appears from the experiments of Pettenkofer and Voit that increasing the proportion of nitrogenous matter in the food determines an increased absorption of oxygen by the lungs. Nitrogenous matter it is which starts the changes occurring in the system, and the suggestion presents itself that upon the amount of nitrogenous matter may, to some extent, depend the application of oxygen to the oxidation of fatty matter. Under this view the success of Mr. Banting's system may be due, not exclusively to the restriction of the principles that tend to produce fat, but in part, also, to an increased oxidising action promoted by the large amount of nitrogenous matter consumed." (Pavy).

Let us draw attention to the substances at these health resorts that furnish the calorifacient group of alimentary principles. Disregarding the fat produced by a complicated metamorphosis of the carbo-hydrates and a small part of the

(a) "The North West Passage by Land." Viscount Milton and Dr. Cheadle.

nitrogenous food ingested (*a*) the main articles of diet from which fat is derived, would be butter and milk. About 1 oz. of the former at breakfast, and 2 pints of milk during the twenty-four hours together furnish at the most 2·6 ozs. of fat, allowing from $\frac{1}{2}$ oz. to 1 oz. for that contained in the lean meat with gravies, &c., 3 to $3\frac{1}{2}$ ozs, are obtained. It is doubtful if this is sufficient with exercise and low temperature, nor is it to be recommended that those with poor appetites should drink more milk. Persons should eat plentifully of butter at breakfast, especially as that meal is not a substantial one.

It would also be very advantageous to continue taking cod-liver oil if it had been found to agree in England, for reasons that are plainly apparent. Should the stomach have been unable to digest it hitherto, trial should again be made in a cold climate, as it might be then more easily assimilated, its oxidation being assisted by the increased proportion of nitrogenous material ingested. A good time for taking it is immediately after lunch or dinner in a glass of marsala, or in milk half an hour after a meal, commencing with one teaspoonful, and gradually increasing the dose.

Pancreatine and pancreatine emulsion are sometimes valuable in assisting the digestion, and malt extracts also. All these substances can be looked upon as supplementary to diet, and not as medicines. If cod-liver oil is objectionable, butter should be eaten at every meal, or cream be made use of.

To further demonstrate the virtue in cod-liver oil, fat, and butter, the table on next page taken from Frankland will give a clear idea of their force-producing value.

(*a*) It is questionable, if with the low temperature in the Alps, phthisical persons should be physiologically compelled to maintain much of the body-heat by the carbo-hydrates, as the changes in their principles previous to their becoming calorifacient take place in the liver; and the function of that organ is often impaired in phthisis.

Name of Food.	Per cent. of Water present.	Force-producing value.		
		In units of heat.	In kilogrammètres of Force.	
			When burnt in oxygen.	When oxidised in the body.
Cod-liver oil'	—	9107	3857	3857
Beef fat	—	9069	3841	3841
Butter	—	7264	3077	3077
Cocoa-nibs... ..	—	6873	2911	2902
Cheese (Cheshire)	24	4647	1969	1846
Isinglass	—	4520	1914	1550
Bread-crust	—	4459	1888	—
Oatmeal	—	4004	1696	1665
Flour	—	3936	1669	1627
Pea-meal	—	3936	1667	1598
Arrowroot	—	3912	1657	1657
Ground rice	—	3813	1615	1591
Yolk of egg	47·0	3423	1449	1400
Lump sugar	—	3348	1418	1418
Grape sugar (commercial)	—	3277	1388	1388
Hard-boiled egg	62·3	2383	1009	966
Bread-crumbs	44·0	2231	945	910
Ham, lean (boiled)	54·4	1980	839	711
Mackerel	70·5	1789	758	683
Beef (lean)	70·5	1567	664	604
Veal (lean)	70·9	1314	556	496
Guinness's stout	88·4	1076	455	455
Potatoes	73·0	1013	429	422
Whiting	80·0	904	383	335
Bass's ale (alcohol reckoned)	88·4	775	328	328
White of egg	86·3	671	284	244
Milk	87·0	662	280	266
Apples	82·0	660	280	273
Carrots	86·0	527	223	220
Cabbage	88·5	434	184	178

In the event of high temperature supervening, the digestion of meats is greatly interfered with, and the usual diet stands in need of modification, as the nitrogenous matter will, if excessive, embarrass the digestive powers, and prove an encumbrance to the stomach, leading to further complications, which may be avoided by substituting food of a different nature that has not to undergo such a complicated

process of absorption and elimination. For this purpose, milk and raw eggs are to be chiefly relied on, with beef-tea, soups, jellies, light puddings, toast, biscuits, and farinaceous substances, as arrowroot, or one of the numerous "foods." By this regimen, bearing in mind that the system requires much less nourishment when the body is at rest, an ample dietary is furnished. During bad weather, also, when but little exercise can be undertaken, and confinement indoors is called for, diminished diet proves of some service, not only to the comfort of the patient, but to his general condition, and, consequently, the local state of disease.

On the cold, dry days, with outdoor exercise, the appetite can be wholly satisfied, with safety and advantage, for it is on these occasions that the "push" is given to nutrition, and any excess in nourishment is more likely to be burnt up or assimilated in the system, to maintain heat, produce force, or counterbalance waste and change.

As the breakfast is not a substantial meal, it will be perceived that the latter part of the day, between noon and 8 p.m., is the period principally occupied by the digestive process, the remaining sixteen hours, therefore, will, in the case of delicate persons, prove to be a great tax on the force and heat-producing powers unless sustained in some way or other. Although it is impossible to lay down rules to apply to every one, a short dietary table for an ordinary case of loss of flesh can be modified to suit the temperament or capacity of any individual, bearing in mind that in many instances "suitable diet" is a matter of experiment.

Regimen at the Swiss Health Resorts.

7 or 7.30 a.m.—Warm milk, $\frac{1}{2}$ litre.

8 or 8.30 a.m.—Breakfast : Tea, coffee, or chocolate, $\frac{1}{2}$ litre ; bread, butter, honey. Extras not provided *en pension*—eggs, cold meat, bacon, omelette, &c.

Noon or 1 p.m.—Lunch: Soup, meats, sweets, cheese, 2 glasses of red wine, or wine and water ($\frac{1}{2}$ pint).

4 or 4.30 p.m.—Warm milk, $\frac{1}{2}$ pint, with a biscuit, or other light refreshment, as tea, coffee, with bread and butter.

5.30 or 6 p.m.—Dinner: Soup, fish, or *entrées*, meats, sweets, cheese, red wine or wine and water ($\frac{1}{2}$ litre).

9 p.m.—Supper not provided *en pension*. Milk, $\frac{1}{2}$ pint, with biscuit, &c., or beef-tea, or some “food” prepared with milk.

A glass of milk, with a biscuit, may also be taken sometimes at 11 a.m. if it is found to agree.

If night-sweats occur, nourishment should be taken at frequent intervals, especially during the night. Stimulants are efficacious at these times—whisky, rum, or cognac; but spirits should always be mixed with milk or egg, or both combined. Their efficiency seems to be increased in this way. Neat spirits as a “*petit verre*” cannot be recommended with much benefit unless there is food in the stomach. Brand’s extract of meat and Liebig’s, are suitable also. These can be mixed together, when they form a better and more natural beef-tea, than when taken separately. The Veltliner wine drunk with meals will act as a good astringent. A frequent cause of perspiration at night is an excessive quantity of clothes on the bed. The usual eiderdown quilt should not cover the patient if night-sweats are frequent. A flannel night-shirt is of great service on these occasions, and the temperature of the bedroom ought never to rise over sixty degrees. Fifty to fifty-five degrees is generally found to be a comfortable temperature in winter.

Exercise, Meals, Sleep, etc.—It is unnecessary to point out the need for individual management in this matter. Much will depend on the state of health and capability for exertion. One rule can, however, be laid down—viz., keep in the open air as much as possible. This will of itself

entail a fair amount of movement, but if the state of the lungs do not preclude skating, tobogganing, or walking ascents—care must, of course, be taken in beginning exercise gradually. A state of breathlessness or fatigue must never be permitted, nor must the body be allowed to cool rapidly if perspiring.

The time available at these altitudes in the depth of winter is somewhat limited for delicate persons, as the sun at this time of the year remains but a short time in the valleys (five or six hours). It is therefore incumbent on those seeking health to make the most of this time, and when in-doors to breathe as much fresh air as can be admitted with comfort. Bedroom windows can be left a little open if the nights are clear, and the heat of the stoves regulated accordingly. In giving an outline of personal hygienic management a description of how the day may be passed will be of some assistance.

At 6.30 or 7 a.m. the stove should be lighted, and a half-litre of milk, warm from the cow, be brought to the bedside of the patient. After drinking this, an hour's sleep may be had. If a cold bath be prohibited, a rapid sponging of the chest and back, followed by friction, is of great service in keeping the skin in a healthy condition. A fairly vigorous patient may have a cold bath, or the chill may be taken off; but precaution must be observed in having the air of the bedroom at this time fresh and warm (not below 55 degs. F.), so that the deep inspirations caused by the shock of cold to the skin shall not take in the used-up bedroom air. A bath or sponging may be tolerated in many cases, provided it takes place *immediately on rising*, while the body is hot. If the skin be allowed to cool down by tardiness in preparing for the bath the water will feel intensely cold. Slight dumb-bell movements may be executed when dressed, throwing the shoulders back and taking deep inspirations. Soon after breakfast, which is taken at 8.30 or

9 a.m., the patient should get out for a walk, making a slight ascent if not too short of breath; and returning at 11 a.m. for a glass of warm milk. The afternoons are mostly spent in sitting out of doors, skating, coasting, walking or sleighing, the latter is not to be recommended in very cold weather. Another half-pint of milk can be drunk in the afternoon, or a little light refreshment such as tea or coffee will do no harm, the patient being generally able to determine if it can be taken with benefit. If he finds it too much fluid, a less quantity of thin cream might be constituted. Dr. Symes Thompson gives a short piece of advice free from mysticism on the subject of exercise in the Alps (*a*):—"Those in health need few restraints, but for those with active lung disease sudden exertion on arrival should be discouraged lest it lead to hæmorrhage. If there is active disease or hæmorrhagic tendency or moist sounds in the lung, the patient should sit out in the sun till dry sounds replace moist ones. He may then walk on the level, or skate, or gently stroll up and down hill, thus causing deep inspiration. Quiet skating can be indulged in by almost all. Tobogging is more severe, as patients are apt to talk and laugh when walking up hill. This is very good for the vigorous, as it expands the chest. Lawn tennis is suited only for the strongest, in whom lung disease is quiescent."

The dinner hour varies in different hotels from 5.30 to 7 o'clock; 6 or 6.30 p.m. is a very good hour. Food should be taken leisurely, and masticated well. Half an hour's rest before and after meals facilitates digestion. The evenings are spent in various ways. Davos possesses a theatre and concert-room (*b*) and amateur theatricals are got up at the hotels, with concerts, tableaux vivants, &c.

(*a*) "On the Winter Health Resorts of the Alps." E. Symes Thompson, M.D.

(*b*) This handsome *Salon* at the Kurhaus is ventilated by a calorifère.

It would be a great boon if, on the occasions of such gatherings, artificial means were provided for the purification of contaminated air similar to those used in the theatre salon of the Maloja. With the intense external cold, air cannot be plentifully admitted to dilute the effete products of respiration, &c., therefore, although entertainments in crowded rooms generally conduce to a healthy frame of mind, and tend to banish despondency, evils may originate if sanitary precautions are neglected.

With respect to the length of time for sleep, the temperament and habits of the individual will have to be considered. The old dictum of 6 hours for a man, and 7 for a woman will scarcely commend itself to most people, nor does experience teach us that any definite duration of time for mental and physical rest can be determined with exactitude. The intensity of muscular, mental, or nervous exhaustion during the day will in all cases influence the desire for repose. Cold also predisposes to sleep, as may be witnessed in hibernating animals. Some human beings also hibernate. It will not be too much to say that 8 or 9 hours slumber in winter, at these cold stations, is near the mark; remembering that nothing is to be gained by remaining in bed in a semi-state of wakefulness after this time is past.

If smoking cannot be entirely given up, the quantity of tobacco used should be cut down as low as possible. Cigarette smoking should certainly be discontinued by those whose lungs are affected, as the habit generally acquired of inhaling the smoke or passing it through the nasal passages, proves very irritating to the mucous membrane, and is more injurious to the throat and lungs than the same portion of tobacco smoked in a pipe.

CHAPTER VI.

The South West End of the Upper Engadine,

The MALOJA.—Since the development of Davos Platz into an Alpine winter resort and a rapid growth in size and population, other places have sprung up having strong claims to be considered equal, if not superior, to that well-known but rather well-filled little town. St. Moritz revives an ancient effort to attract people in winter, this time with some success, the Kulm Hôtel having more than a hundred and fifty guests living there during the past season. Wiesen the picturesque, with its ponderous flanks and million pine trees, is on a fair way to offer health to a limited number who desire quietude and the charms of country life. The milder temperature and extreme calmness give a special character to this resort, which renders it suitable as a change at the end of the winter season for those who have been living at a higher level.

It is indispensable for the reputation of the high valleys in Switzerland that more places spring up for the "winter cure," as it is notorious to those who give real attention to the subject that, although these places increase in size, population, and independence, individual and collective energy is needed to keep pace with overcrowding and to preserve the clean air of the Alpine heights free from contamination. As but few hours can be spent out of the house in winter, it is imperative that hotels which harbour a number of delicate individuals in search of health within their walls should be rendered perfect in a sanitary point of view, both as regards drainage and ventilation. Of how much therapeutic value must that air be which, in five or six

hours can, not only neutralise the ill-effects of a contaminated atmosphere for the remainder of the day and night, but can, in numerous instances, favour an almost miraculous recovery.

Heretofore, the curative properties of these climates have only been developed to a fraction of their full power; and whilst the restorative agent has lain at the closed doors and double windows of over-loaded hotels, patients have been sent home cured in spite of what would seem to be to on-lookers either medical ignorance of the true antidote or want of common sense.

These drawbacks, however, have now been met by the construction on the Maloja plateau of a large Kursaal (designed especially for winter residence by M. Jules Rau, a Belgian architect), fitted with arrangements for the propulsion and extraction of air throughout the building, as well as for the addition of ozone to the atmosphere of the corridors and some of the large salons. The latter is effected by the surplus electricity at command in use for the electric lighting of the building.

In addition to these appliances for rendering the atmosphere as healthy and pure as possible, baths supplied with mineral waters from a chalybeate spring discovered in the neighbourhood are being established for the organisation of hydropathic treatment. This, of course, is employed mostly for the summer cure, but, as the douches and sprays can be rendered tepid, with high pressure, an additional agent for the treatment of many morbid conditions will be available. A double value is lent to these restoratives by the uniform temperature throughout the building (including the baths and corridors) frustrating any liability to chill on going to and from a bath. A thermal room likewise is to be furnished with the water, where masseurs and masseuses will execute the processes of shampooing and medical rubbing. Electric baths and needle sprays complete the winter set.

The Kursaal, which may be considered unique in all sanitary arrangements, faces the lake of Sils, between which and the N.E. façade lies the *parc*, and on the S.W. side of the building a large garden (in all, 13 acres), with skating-rink, pavilions, &c. A theatre salon is semi-detached from the main corridor to guard against any transmission of sound to the interior of the house. This structure is also warmed and ventilated by propulsion and extraction, but in a slightly modified manner to that of the general plan.

An entirely new plan of drainage has been adopted, by which it is impossible for any sewer-gas to enter the dwelling. The main difficulty in these climates—viz., the freezing of the water supply to *cabinets*, has been overcome, and each *cabinet urinoir*, &c., is efficiently trapped and ventilated with the external air.

It is anticipated, and with some reason, that what has been termed the air-cure will be exhibited in its true meaning, and that results hitherto unattainable where “air-nourishment” is the essentiality will be achieved with greater rapidity and success than in many other health-resorts where but little mindfulness is bestowed on a point of such signal importance.

As much attention is devoted at the present time to all matters connected with hygiene, notably that section relating to ventilation, a description of the system in use at the Kursaal Hôtel will be given further on.

The MALOJA PLATEAU is situated at the higher extremity of the Upper Engadine, and is well protected from northerly, easterly, and southerly winds. Facing the plateau is the lake of Sils; the largest in the chain of lakes between Maloja and St. Moritz. In contrast with the glassiness of its tranquil waters, and the clear blue of the sky, are the rugged crags of the higher Alps, clothed below with red firs

and larches. The site is considered by many to be the most romantic and picturesque part of the Engadine.

On the eastern wing of the lake and plateau lie the Bernina chain: comprising the loftiest mountains of the Grisons and the whole of Switzerland, matted with glaciers and snow fields to an extent of more than 350 square miles. The slopes of the Corvatsch, Surlej, and Della Margna (their peaks elevated to ten and eleven thousand feet), abut on the lake of Sils, and eastern side of the plateau, protecting the lower ground. Opposite and rounding towards the north-east the rough broadside of Lunghino, Gravasalvas, and Materdell are in close proximity. To the south the Muretto, Dei Rossi, Del Forno, and Salecina rear their summits to heights of 10,000 feet, whilst a mile or so to the south-west is seen the prominent Piz Lizzone and the serrated crests on the eastern ridge of the Val Bregaglia.

By these shelters the plateau is screened from harsh upper currents of air. The thalwind or valley-wind, which blows in every Swiss valley, seldom exceeds a force of one degree Beaufort scale, is by no means insupportable, and dies away in force and frequency when snow covers the adjacent regions. At this season, and when the lake of Sils freezes, still greater calm prevails, and the locality partakes of that Alpine stillness and sunshine which allows the most delicate individuals to be exposed to a low temperature without feeling the sensation of cold.

Preceding and during a fall of snow in the mountains of Switzerland, storms are not uncommon, alternating with lulls and blasts of wind. Commonly these proceed from a southerly or south-westerly direction and bring with them the peculiar changes in the atmosphere, which have given rise to a special name for southerly currents of air, viz., föhn wind. This föhn, which is considered the pernicious wind for invalids is rarely perceived at the S.W. end of the Engadine, its peculiar characteristics being tempered by the high altitude, and by

passage over a portion of the Bernina snow fields and glaciers. In this way the temperature of a southerly current is greatly reduced, and the commencement of the föhn shorn of its sultriness. The rapid rises in temperature at the onset of a southerly wind were not common during the past winter, nor was there ever recorded a percentage of "relative humidity," so low as to render the atmosphere of the interior of warmed dwellings too dry for healthy respiration.

One of the most important features of Alpine climate—and one which unquestionably acts as an adjuvant in the chemical process of the natural and healthy change taking place in the blood discs—is the amount and duration of sunlight. Were it not for the transparent air and powerful sun the high Alps would be quite unsuited to delicate people. By reason of the direct chemical action of light, and the increased time afforded for out-door amusement and exercise which a longer day admits of, an additional hour's sunlight during the short days of December and January is an inconceivable advantage, both for the mind as well as the body. The inequality, in this respect, of Davos, Wiesen, and St. Moritz, with the Maloja, is due, as may be supposed, to the configuration of the surrounding mountains. If a slice could be taken off the Jacobshorn at Davos, and the Stulsergrat, at Wiesen, the sun would flood those districts before half-past ten, on the shortest day.

St. Moritz at this time loses the sun at 3 p.m., and although it rises during the middle of February at a quarter to eight, it sets at 3.50 p.m. The sunset of the Maloja on this date takes place at 4.30 p.m., and during the months of December and January the Hôtel-Kursaal has more than one hour's sunshine longer than Davos, Wiesen, or St. Moritz. In respect of duration of sunlight, Pontresina still exceeds the Maloja, but does not enjoy the *late sunset of the latter*. At the end the reader will find the sun-hours given for the various health resorts during the winter months. Sunrise

and sunset take place on the first day of January as follows :

		Sunrise.	Sunset.
Maloja	...	9.35	3.45 P.M.
Wiesen	...	10.35	3.45 „
(a) Pontresina	...	8.30	3.10 „
(a) St. Moritz	...	10.0	3.5 „
Davos Platz	...	10.3	3.0 „
(b) Andermatt	...	11.45	3.15 „

It will be seen that Wiesen and Pontresina stand next to the Maloja, the late departure of the sun being of most consequence.

The necessity for exposure to Alpine solar rays in the treatment of various complaints, is well illustrated by cases of anæmia among some of the domestics employed in hotels at these levels. Attracted by the rumours of cure in anæmia they seek employment at some health station, but being principally confined to badly ventilated kitchens and shady places their anæmia rests with but little improvement. The case with nurse-girls is different as they very soon gain a healthy tint.

Another advantageous feature of this part of the Engadine, is the vast extent of level ground which admits of exercise being taken without fatigue. After the breathing becomes freer and there is a tendency for the chest to expand, tobogging and gentle ascents can be undertaken. For the more vigorous a descent into the Bergell Valley on a toboggan is one of the finest runs in Switzerland. Skating of course, can be indulged in by those who are equal to the pastime ; a good rink is in course of preparation, whilst the lake of Sils is but 300 yards from the Hotel. It is in the choice of amusements and graduation of exercise where

(a) Taken from the houses on the main road of the village (Dr. Ludwig's "Ober-Engadin").

(b) Reported by Dr. Schmid.

judgment ought to be shown. During summer time more variety in out-door sports and games can be pursued, mountaineering, boating, fishing, lawn-tennis, &c., and consequently more care must be taken in refraining from over-exertion. The long evenings of winter can be enlivened by healthy in-door amusements, as fresh air and ozone are supplied to the theatre-salon as well as to other recreation rooms, and a clean atmosphere is breathed day and night.

The climate can be described in the general terms of all Swiss high level stations, viz., a stimulating, bracing, tonic, and cold climate; increasing the power of the heart and other muscles, expanding the chest and exciting the nervous and glandular systems into healthy activity. The Maloja (6,000 feet) shows a mean temperature two or three degrees lower than Wiesen and about the same as Davos Platz (5,100 feet).

During mid-winter a dip of over 3,200 feet can be made in an hour and a-half to Promotongno, where there is a good hotel; and the Italian climate of Chiavenna (1,090 feet) can be reached in less than three hours, if the upper level appears to disagree. This facility in being able to quit the highlands without a prolonged, and in cases of illness, a dangerous journey from any other Alpine station, cannot be over-rated. Doubtless there are instances in which any climate, either cold or temperate, does not always effect the salubrious change anticipated; it then becomes a question of the utmost importance to an invalid, how to escape without incurring risk. In the present instance the dangers of moving, even in the depth of winter, are at a minimum. A drive to the lake of Como requires less time than it takes to reach Davos from Landquart.

The soil and vegetation of the Maloja are of the usual character found at these altitudes, a thin, rich, dark incrustation of the former, rapidly absorbing moisture. The bed of the plateau (formed by an ancient recession of the lake of

Sils) consists of peaty mould, resting on a thick layer of gravel. Up to a height of 1,000 feet and 2,000 feet above the level ground, the slopes on the eastern side are covered with larches and red pines. The western side is rugged, large boulders and crags being interspersed with patches of nutritious short grass. On the plateau itself, and the little hills adjoining, are dwarf *melèzes* (red pines), heath and cranberry bushes.

The coldest part of the Engadine is near Bevers; the temperature in winter becomes a little higher as we approach the Maloja, owing to the fact that its south-west end terminates in an abrupt fall of 1,200 feet (*a*) into the extensive valley of Bergell, which after passing Promontogno terminates near Chiavenna in Italy, eighteen miles distant; the Italian frontier being twelve miles from the Maloja. To the proximity of a channel at this end for milder air to mount and replace the denser and colder strata, the deviation in temperature may be attributed.

The difference, however, with the several Alpine health resorts, lies not so much in their grades of temperature, as in their sanitary characters, especially as regards the purity of internal air. As the term "air-cure" (*luft kurort*) has rightly or wrongly grown into use, it will be expedient to include in this chapter a description of the special appliances which have been placed in this modern building with a view of furnishing a larger breathing space to its inmates.

The system admits of being adjusted to suit any temperature or degree of humidity of external atmosphere. The calculation by M. Cortella of the volume of air provided for each individual, shows the possibility of supplying 100 mètres cubes (3,500 cubic feet) per hour for each person, supposing the Kursaal to contain 500 individuals. By the double means of propulsion and extraction furnished to each

(*a*) The precipitous descent is cut by a road with sixteen zig-zags.

chamber this breathing space exceeds by 0·6 times the estimate given by one of the greatest authorities for ventilating soldiers' barracks at night, (a) and the same care has been exhibited in ventilating the basement of the house as in ventilating the bed and sitting rooms.

In the basement beneath the theatre and concert room, situated apart from the main building, are placed three *chaudières* in which steam is generated to a low pressure (1 to three atmospheres). This steam is conveyed to forty-nine *batteries*, distributed throughout the basement, where it transverses a series of pipes with flanges attached to increase the surface of heat. A *batterie* is made up of twenty pipes enclosed in a case, through which air is drawn from the outside of the building, previously passing through wire gauze. This air receives heat by passing over the steam pipes, whilst the steam, having no communication with the interior of the *batterie* is condensed, and returns as tepid water to the *chaudière* for the economy of fuel. Within these *batteries* are contrivances for moistening the dry external air to a healthful degree of humidity in harmony with the increase of temperature and hygrometric condition of the atmosphere, or to medicate the air of one or more rooms with volatilised remedial agents; as leading from each *batterie* are five tubes that control by valves the quantity of warmed air discharged through them to the five chambers which they supply.

The ascending power of the warmed air at a temperature of 50° centigrade (122° Fahr.) raises it through the conduits to each room in the house, where it enters a dormitory at the topmost storey, having an agreeable temperature, and at a sufficient rate to change the air of a bedroom once in every hour, day and night. In the theatre, smoking and billiard

(a) In barracks, 30 mètres cubes, per hour during the day; 60 by night. Workshops, 60 mètré cubes. Schools 30. Hospitals 80. (Movin).

rooms, and public salons, a necessary number of conduits are provided for renewing the atmosphere two or three times every hour.

Each delivery tube is fitted with a sliding valve which can be adjusted at pleasure according to the degree of heat desirable for comfort.

To allow of the escape of used-up air two tubes of exit are placed in a chamber, one under the bed for the escape of the heavier impure gases, another near the ceiling to liberate the lighter impurities. A certain number of these exit tubes are in connection with an iron casing surrounding the main flue of the *chaudières*, that acts as an extraction shaft for the foul air of the central portion of the building. The kitchen chimney acts in this manner for another part of the house and together with two stoves placed in the basement, aspirate the air of other rooms into the double chimneys. As the upward motion in these extraction shafts would scarcely be forcible and rapid enough during summer time to effectually withdraw the air from so many chambers, and as perchance some air might find its way back again to the upper stories, a contrivance with steam tubes is fixed in the space beneath the roof to again heat and thereby promote the ascent of foul air and so accelerate its escape externally.

Every corridor, room, and *cabinet* in the house is thus ventilated, and the atmosphere of any special chamber can be medicated at will by placing an antiseptic agent above the valve of the *batterie* in the air-tube supplying the apartment: being volatilised at a temperature of 50° centigrade, the product is rapidly carried to its destination without tainting the air of other rooms, as the escape will take place in the exit shafts, each chamber having its own separate ventilation by three channels—one inlet and two outlets (one for the escape of warm impure air, the other for the cooler air near the floor). By the simple adjustment of the valves in a room, the temperature can be regulated with nicety. Even in summer, if

the *batteries* are not heated, extraction will go on as usual ; and the windows may be opened without interfering in any way with the general ventilation of the house.

It will easily be seen that with this system various therapeutic agents of undoubted importance are placed under immediate control, viz.:—

1. Temperature.
2. Moist fresh air.
3. Dry fresh air.
4. Volatilised remedial agents.

In addition, ozone generated by a powerful apparatus placed in the vestibule, permeates the corridors, &c., and passes to the *salle à manger*, *théâtre-salon*, *salle de lecture*, &c., by means of a “blower” and tubes.

CHAPTER VII.

The Maloja Spa and Summer Cure.

THE term summer cure has been confined to those localities in Switzerland where waters having special chemical properties were available for the treatment of various complaints. The Canton of the Grisons is especially famous for the number of its spas; the most frequented up to the present being St. Moritz in the Upper Engadine, and Tarasp-Schuls in the Lower Engadine.

Although Davos has valuable sulphur springs within a few miles (Clavadel and Spinabad) scarcely anyone pays the little town a lengthened summer visit. At this season, the place, for some reason or other, is quite empty. The reverse may be said of Wiesen, eleven miles nearer Coire. People reach this secluded haven and make a stay, frequently when *en route* to the Engadine by the Albula Pass, but Wiesen holds no pretensions to a summer "cure."

The therapeutic utility of the waters of St. Moritz has been known since the time of Paracelsus, and the treatment by baths and internal administration has been followed without intermission for a number of years. Thousands have visited the village, and still do so. A variety of complaints are said to have improved, the most marked being anæmia and chlorosis, debility from physical or nervous causes, dyspepsia and some cachetic states of the system. The scrofulous diathesis appears to be greatly benefited by the treatment; also sluggish circulation, chronic gastric and pulmonary catarrh, and other cases in which a tonic stimulating course is desirable. Malnutrition with phosphatic urine and extreme low tension of arteries undergoes a salutary change by residence alone. Individuals of both sexes

who have been exposed to tropical heat and malaria, regain health rapidly in summer and winter.

It is a matter of good fortune that a mineral water has been discovered in the Maloja, having the same characteristics as that of St. Moritz,—viz., chalybeate with large quantities of free carbonic acid. Much of the crowding at the latter springs during the summer season will now be avoided and visitors be enabled to take the waters with less hurry and inconvenience, and thereby obtain their full benefit.

Dr. E. Reichhardt, Professor of Chemistry in the University of Jena, pronounces a strong probability of the Maloja spring being actually the same as that which supplies St. Moritz, eleven miles further down the Engadine.

The analysis made by him during the autumn of the past year is as follows :—

			Maloja.		St. Moritz. (Paracelsus).
1. Sulphate of lime	2·2470	...	—
2. Sulphate of soda	—	...	3·2101
3. Sulphate of potass	—	...	0·1480
4. Chlorate of potass	0·2215	...	—
5. Chlorate of sodium	0·1433	...	—
6. Bicarbonate of soda	4·8227	...	1·81518
7. Bicarbonate of potass	5·0845	...	13·01950
8. Bicarbonate of magnesia	1·4224	...	2·02188
9. Bicarbonate of iron	0·900	...	0·38648
10. Free carbonic acid	1·60 or 859·7	in 10,000 parts.	
11. Sulphuric acid	1·32	...	1·877
12. Carbonic acid	8·98	...	35·960
13. Chlorine	0·19	...	0·218
14. Lime	2·91	...	5·063
15. Magnesia	0·45	...	0·631
16. Soda	1·74	...	2·376
17. Potass	0·14	...	0·080
18. Oxyde of iron	0·90	...	0·234 (?)

From the foregoing analysis, it is noticeable that the water is especially rich in carbonic acid, soda, potass, and iron. The importance of this combination will be appre-

ciated at once, when it is remembered that the presence of carbonic acid aids the assimilation in the system of the iron by enabling the stomach to tolerate the latter by reason of the sedative effects of the gas.

In a recent work by Tichborne and Prosser James on the "Mineral Waters of Europe," it is said of chalybeate, at page 28, p. 3, "The medical use of iron is so everyday an occurrence that no wonder waters known to contain the metal should be popular. There is little doubt that iron may be assimilated if taken in these waters, in cases in which it is scarcely tolerated in the most refined pharmaceutical productions. When it occurs as carbonate it is most palatable, and often most effectual."

In the report appended to the analysis of the Maloja waters, by Professor Reichhardt he remarks, "It can be well understood that I have only been able to define and calculate the fixed parts of this mineral water. It is evidently a water having a very close resemblance to that of St. Moritz."

Much of the carbonic acid has naturally escaped during transport, nevertheless there is found a quantity sufficient to cover double the carbonates and then there still remains a portion quite "free." This never occurs in the case of any ordinary water.

Its contents in bicarbonate of soda doubly proves it to be of the same origin as the waters of St. Moritz ; at the same time iron is found. Alone, iron would not carry with it much importance, seeing that in springs, *non captivées*, it is easily found in the turbid parts, but disappears from the clear portion when transported.

The same thing happens with the waters of St. Moritz, although the contrary is reported, in fact some iron waters do not even retain their taste, which is the case here. Hausemann also distinguishes between "combined" and "free" iron, the first precipitate is no longer part of the water, The foregoing is, then, a true mineral water resembling that of St. Moritz,

if not absolutely the same spring." He then goes on to state that some analysts wrongly calculate the bicarbonate of soda in obtaining the sulphate, and suggests that this mistake has been made with the St. Moritz water.

In no sense does there appear to be a contra-indication of the summer Alpine climate for persons with chest complaints, that is with some of the early phases of phthisis, not those with hopeless advanced symptoms, or who are in a feeble irritable condition, with high fever, and incapacity for gentle exercises. The determination of suitable cases for the high altitudes, both in summer and winter, can be roughly gauged by the amount of physical power conserved, taken in conjunction with the duration and apparent rate with which consolidation or softening is advancing. The slow scrofulous forms and fibroid conditions receive undoubted benefit, but these must not be complicated with secondary lesions.

A little misapprehension exists on the duration of what is known as the "summer season." Fashion has led the custom, of visiting the Engadine in the months of July and August: the majority of persons entertaining an impression that those are the only months to see the Engadine in all its beauty, and at the same time acquire the succour of its restorative climate. This is quite a false notion formed on no foundation of reality. The months of June and September are sometimes among the best that can be spent in the Alps at these altitudes, and the air at the same time carries with it the remarkable invigorating agencies which characterise the mountain climate. The worst period of the year is the snow-melting, occurring some time during the month of April; but this can scarcely be termed a dangerous season, even to the most delicate persons, provided that ordinary precautions are taken in keeping the feet dry and not incurring thoughtless exposure to inclement weather. This applies equally to the introductory snow-falls and storms preceding the settled weather of winter. Invalids must guard

against unnecessary exposure everywhere, both on the Riviera and in the Alps, although it would perhaps be less injurious to face a mild Alpine snow-storm than remain shut up in a badly ventilated room in a southern health resort.

The length of time also, needful for a permanent recovery of lost strength or health, has ever been considered too short to carry with it a firm re-establishment of exhausted energy. A summer month spent in the Engadine—with assiduous regard to diet, exercise, sleep, and medical treatment—should, in almost every instance, be prolonged to six weeks *at least*; longer if time permitted. The lengthened stay would, in the long run, be an economy of time as far as health was concerned, and prove a greater satisfaction to doctors and patients alike. Force of a more lasting and durable nature would be accumulated, the system settle down in a more staple condition, with enlarged powers of resistance to morbid impressions, than if the treatment be suddenly relinquished to return to old occupations and former habits.

It is true that physiological changes take place under the summer treatment with great rapidity and completeness. A good indication of such being the case is the demand made by nature for more fuel, shown by the increased appetite, by the skin acting under the stimulus of the baths, the kidneys and the whole glandular system being awakened into activity, blood changes quickened and amplified, and above all, the perception that the capacity for mental and physical exertion is extended, but if a departure is made immediately improvement of the health is attained and strength begins to be felt, some disappointment may eventually be experienced, and the climate be discredited by the transient effects of too short a stay to allow the frame to be seasoned in the new conditions, or undergo the peculiar alteration known as acclimatisation. This is especially noticeable in anæmia and chlorosis, where the blood changes take place rather rapidly, and a hue of health is acquired in a few weeks leading to an erroneous

conclusion that a permanent cure is effected, whereas a further time is needed for its completeness. In chronic cases, either of catarrhs (pulmonary, gastric, or uterine), phthisis, anæmia, some abdominal affections, and even dyspepsia, a medical mind is aware of the value of prudent management after the disappearance of symptoms. Whilst it is evident that in constitutional affections, such as scrofula, rachitis, &c., patience and persistent care are a *sine quâ non*, it is doubtful if anyone would be bold enough, from the comparatively limited number of observations at our disposal, to lay down a clear indication for patients who should go south, and those who should mount the Alpine heights. At present the same can be said of the mountains as of the South—a variety of cases have improved in a most remarkable manner. It is also undoubtedly true that in the high altitudes some rapidly advancing symptoms have been arrested when scarcely any hope of a return to health could be held out by physicians.

Patients with the following diseases should, however, refrain from seeking health at high elevations :—

1. Diseases of the heart or large vessels.
2. Tendency to articular rheumatism.
3. Kidney diseases (during winter).
4. Acute inflammations of throat or larynx.
5. Some diseases of bladder or prostate.

6. Persons somewhat advanced in years, should not visit the mountains, unless the circulatory system is sound.

In every instance the body should be clothed in flannel, a thin fine texture for summer wear. No great coats or heavy mantles will then be necessary, except for driving or sitting out. Even the danger in summer evenings from a fall in temperature, or from change of wind, is by no means serious, for the dry atmosphere does not carry with it the chilliness and cold which is felt in climates where the absolute humidity is greater. Moreover, the concert room, salons, corridors, &c.,

within the Maloja hotel permit of in-door recreation, with ample breathing space, whilst in the *parc* and gardens, numerous *pavillons* afford shelter externally, the mind being enlivened by variety in amusements and magnificent scenery. There is no lack of walks or drives either in St. Moritz direction—passing the villages of Sils, Silvaplana, and Campfer, with their superb lakes, or down by the steep incline of the Maloja Kulm into the beautiful Bergell Valley.

To those who cast about for the best and surest means of correcting the unnatural conditions of life and health which so commonly present themselves in the forms specified at the commencement of this introductory study, the consideration of humidity of the air, temperature, sunlight, barometric pressure, and ozone, will assuredly not prove to be a fruitless inquiry devoid of interest.

We are from the commencement of life immersed in a fluid with varying pressures and compositions; can we yet accurately gauge the influences on health and disease, of our atmospheric medium?

THE JOURNEY FROM ENGLAND.

THE quickest route from London to the Engadine is by Dover, Calais, Boulogne, Amiens, Tergnier, Laon, Rheims, Belfort, Bâle (Basel), Chûr (Coire).

The tidal boats from Folkestone to Boulogne are more agreeable to travel by, and the luggage by this route is taken care of, and not so roughly handled as between Dover and Calais. Inquiries must be made of the South-Eastern Railway Co. for the hours of departure, as these boats depend on the tide.

Viâ Dover and Calais, a start is made at 10 a.m. from Cannon Street, Charing Cross, or Victoria. Lunch can be obtained at Calais at 2 p.m., dinner at Tergnier about 6 p.m., and Bâle reached at 6 a.m., where breakfast is taken at the railway station. From Bâle the journey is continued, after breakfast, either to Como by the St. Gothard line, and from thence to Colico by steamer, then by diligence to Chiavenna and Maloja; or leaving Bâle at 7 a.m. Chûr is reached at 3 p.m. (A stay of two hours can be made at Zürich, if desired, and Chûr reached later.)

From Chûr the diligence leaves in the early morning for Maloja, *viâ* the Julier Pass and Silvaplana, and arrives in time for dinner.

It is advisable for delicate persons to break the journey either at Bâle or Chûr. Those who are unable to obtain any sleep in a train would do well to alight at Bâle and go to bed; otherwise, a break at Zürich for two hours, partaking of dinner at one of the numerous hotels in that interesting town, will be sufficiently refreshing to allow of a continua-

tion of the journey without fatigue. A telegram might be sent to Hôtel Steinbock, requesting a room to be warmed, as the air at Chûr will feel chilly in autumn after sitting still in the train. Should the weather be fine, a few days' stay can be made with benefit, previous to mounting. A private conveyance should then be taken for a short journey to Mühlen, spending the night there. From Mühlen the diligence leaves at 2 p.m. for Silvaplana, and the Maloja is reached at 6 p.m.

CHAPTER VIII.

Meteorology of the Maloja.

THE instruments were by Casella, of Holborn, and were all corrected at Kew previous to use. A Stevenson's screen of the old pattern was placed in the shade of a large chalet, and was fitted with an arrangement admitting of its adjustment after a snow-fall to the ordinary distance of 4 ft. from the surface of snow. The nearest building was the chalet, 60 ft. distant.

In the Alpine climate a thermometer screen of whatever pattern should be placed in shade, otherwise the intense solar heat warms the woodwork, and thereby raises the temperature of the interior many degrees above that of the external air.

The treatment of the ice-covered bulb also requires great attention, or a wider difference will be noted between the two thermometers than ought actually to occur. This takes place during a rise in temperature, and if not met by moistening the muslin around the bulb, half or three-quarters of an-hour previous to observation, a false estimate of the amount of humidity will be made.

The dew-point and weight of vapour in the atmosphere were calculated from Glaisher's tables, in conjunction with Apjohn's formula :—

$$F = f \frac{d}{88} \times \frac{h}{30}$$

when the temperature of the wet bulb was above 32° ;

and $F = f \frac{d}{96} \times \frac{h}{30}$ when below 32° Fahr.

The force of the wind is noted according to the Beaufort scale, judged from the readings, in miles, of an anemometer placed on high ground in an exposed part of the plateau, 14 ft. above snow. Another anemometer was erected 6 ft

above snow, in the garden S.W. of the Hôtel-Kursaal. This instrument registered from $\frac{1}{3}$ to $\frac{1}{2}$ the number of miles of the former. The past winter in the Engadine was not so calm as the previous four winters, owing to the exceptionally small quantity of snow which fell in November and December. The depth of snow even in mid-winter being unusually shallow, many slopes and rocks were left bare, favouring absorption of solar heat by the earth, to be again radiated off into space during the night, and in this way giving rise to local movements of air.

At the end of January and during February the snow-covering of the earth was from one to three feet deep, and the calmness, blue sky, and sunshine at this time were typical of Alpine winter. Although the mean temperature for February was 26·5 Fahr. (-3° Centigrade) at noon, there was never felt any necessity to put on extra garments. No great coats were needed, and a cold bath at a temperature of 41° Fahr. could be taken with advantage, and less positive feeling of cold than in London during winter.

The coloured sunsets which were so remarkable over the whole of Europe, partook of a gorgeous display at these heights. On several occasions a complete representation of the solar spectrum was stretched across the heavens from E to W, The earth appeared to be occupying the yellow spectral tint and the other six principal colours followed in the usual order of dispersion, as when white light is decomposed by refraction.

	Indigo.	Violet.
	Blue.	S. Orange.
	Green.	S.E. S.W. Red.
Yellow.	E.	W. Yellow

The atmosphere at these times was singularly free from cloud, and contained a small quantity of watery vapour, but not less than is generally found at 6,000 feet in the Alps, during fine weather. The phenomena were preceded and followed by wind.

Mist or fog was observed five times during the winter :— In the month of November, not at all ; December, twice ; January, once ; and February, twice.


The hygrometric state of the atmosphere reached saturation on fourteen occasions only :—November, three times ; December, five ; January, one ; February, five. The evaporation from snow, ice, or even an expanse of pure water, is not in any way injurious to health : it is badly drained ground marshy land, or sodden vegetable matter which is to be avoided. The quantity of water evaporated from the latter is considerably greater than from a lake or sea, by reason of absorption of heat from the sun's rays, and the larger evaporating surface presented by wet soils, &c., in an equivalent area ; added to which, in some instances is malarial poison, or emanations from decaying vegetable substances ; but, as is perfectly well known, the altitudes under consideration are quite secure from any suspicion of malaria.

The extreme day temperatures noted were :—

1° Fahr. (17·2° Centigrade) on the 19th of February ;
45° Fahr. (7·2° Centigrade) on the 21st December ; and
143° Fahr. (61·6° Centigrade) for the black bulb in vacuo
on the 19th of February.

The “drying power of the air,” noted in the tables, is the weight of vapour which ten cubic feet of air were still capable of absorbing at the time of observation. The mean “drying power” for the winter was 5·4 grains, against an average of 12 grains of vapour per 10 cubic feet of the atmosphere. Owing to this dry air being a bad conductor of heat, it is possible to quit a heated room and remain in the open air several minutes without being able to recognise the wide difference in temperature, which sometimes attains as much as 50° Fahr. To this absence of moisture suspended in the air may be ascribed the immunity from catarrhs and the capability of animals to support the low temperature in these high regions. (a)

(a) Extract from a paper by the author read before the Royal Meteorological Society.



METEOROLOGY OF THE MALOJA.

IN the following Meteorological Observations the abbreviations c.c., cn., cum., indicate respectively cirro-cumulus, cirro-nimbus, cumulus ; sn., snow ; blue, blue sky.

The "earth temperature" was taken at one foot below the surface ; and the note of "ozone" made by Schönbien's method, with an exposure to the air of the test papers, for six hours instead of twelve.

The maximum and minimum thermometers were noted three times a day, for the purpose of determining the variation of temperature during forenoon and afternoon.

METEOROLOGICAL OBSERVATIONS

TAKEN IN THE MALOJA DURING THE WINTER OF 1883—84,

BY

A. TUCKER WISE, M.D.

*From a Paper read before the ROYAL METEOROLOGICAL
SOCIETY, on the 21st May, 1884.*

	AT 9 A.M.						AT NOON.							
	Dry Bulb.	Wet Bulb.	Maximum.	Minimum.	Cloud.		Dry Bulb.	Wet Bulb.	Maximum.	Minimum.	Cloud.		Elastic Force of Vapour.	Weight of
					Form.	Amount.					Form.	Amount.		
1883														
Nov.														
1	36.5	31.0	44.0	27.0	blue	0	40.0	36.5	43.0	35.5	blue	0	.184	
2	35.0	29.5	38.0	28.5		0	40.5	33.5	42.0	34.5	cir	1	.128	
3	33.0	28.0	39.0	26.0	cum	1	39.0	33.5	40.0	33	c.c.	5	.142	
4	33	28.5	38	27	cir	1	40	32	40.5	33	blue	0	.114	
5	32.5	32	39.5	30.5	over	10	35	34.5	36	2	cum	8	.153	
6	30	25	38.5	24.5	cum	6	31	28	31	29	cum	5	.128	
7	34	32.5	34	28	c.n.	9	35	33.5	36	33	c.n.	9	.178	
8	35	35	36	33	over	10	35.5	35.5	36	35	over	10	.208	
9	29	25	35	26.5	cum	1	29.8	25.2	30	28	blue	0	.097	
10	28.7	26	34.5	21.5	c.c.	2	31.8	28.5	34	27.5	c.n.	9	.128	
11	20	16.5	31	17.5	c.c.	6	23.8	19.8	24.5	18.5	over	10	.073	
12	23.5	20	25	11	cum	7	26.2	23.5	27	23.5	cum	6	.103	
13	26	23	32	17.5	c.n.	7	27.5	22	28	25	c.c.	4	.072	
14	21.8	18.8	23	12.5	c.c.	6	27.5	24	27.5	20.5	c.n.	7	.099	
15	15	13	26	12	blue	0	22	20	27	15	blue	0	.091	
16	23.8	21.8	24	18	cum	6	22.8	22.2	26	21.5	cum	3	.114	
17	20.5	18.5	23	12	c.c.	6	26	22.5	26	19.5	cir	4	.090	
18	30	28	30	16	c.c.	3	32	30	35	29	over	10	.150	
19	25	24	31.5	17.5	c.n.	6	34.5	29	34.5	24	cum	2	.114	
20	31.8	28	37	25	c.c.	8	31	27	32	28	c.n.	8	.113	
21	20	17.8	30.5	17	cir	1	27.2	22.5	27.5	18.5	cir	2	.082	
22	22	19	29.5	15	c.n.	8	29.2	23	29.5	20	over	10	.071	
23	17	16	27.5	13	cir	1	27.5	23.5	28	16	cir	2	.092	
24	18.5	16.8	27.5	17	blue	0	23	19.5	22.5	17.5	blue	0	.075	
25	19	16.2	23	15		0	29.5	24	29.5	17	cir	4	.083	
26	27.8	26.5	29	14	over	10	29	28	29	26	over	10	.144	
27	30.5	30.5	30.5	26.5	over	10	31	30.5	31	29	over	10	.128	
28	29.5	28.5	31	27	cum	8	31	29.2	31.5	28.5	cum	5	.146	
29	20	19	31.5	16.5	blue	0	29	25.5	29	19	blue	0	.108	
30	24.2	22.8	35.5	18.5		0	29.2	26.5	29.5	23	blue	0	.121	
	26.4		44	11		4	30.5		43	15		4		

OF THE MALOJA.

AT 3 P.M.

Time.	Drying power of air per 10 cub. ft.	Dry Bulb.	Wet Bulb.	Maximum.	Minimum.	Cloud.		Wind.		Solar Radiation.	Earth Temperature	Rain Gauge.	Snow.	Ozone six hours
						Form.	Amount.	Dir., upr. and lower Currents.	Force (1 to 10).					
	Grains.													
5	7	38.0	37.0	42.0	37.0	c. c.	1	SW	0.5	120	35.5			9
8	13	39.0	34.0	41.5	38.5	cir	1	SW	0.5	127	35.5			5
2	10	37.5	31.5	40	36	cum	4	SW	0.5	113	35			4
2	14	39.8	32.2	41	38.5	blue	0	W	0.5	126	35			6
	5	38	31	40	35.5	nim	8	NE	0	120	34.5	1.6	2	11
9	5	30.5	29	31	29.5	over	10	W	1	86	34			5
6	3	35	34.5	36.5	35	c. n.	9	W	1	104	34	12.5	rain	13
5	0	34.8	34.2	36	34	over	10	W	0.5	90	34			12
8	7	32	26.2	32	30	blue	0	—	2	130	33.5	2.3	3	—
8	6	30	29	32.5	29.5	over	10	E	1	124	33	3.1	4	6
8	6	24	20.5	27.5	22.5	cum	4	W	1	115	33			5
	5	27	23.8	28.5	25	c. w.	2	W	1	119	32.5			5
5	9	22.5	19	27.5	21.5	c. c.	3	SW	1	120	32.5			7
2	6	25	22	27.5	19.5	c. n.	7	W	1	111	32.5			5
4	3	26	23	27	20.5	blue	0	N	0.5	126	32			5
2	1	24	22	25	21.5	cum	4	NE	0.5	116	32			15
	6	26	22	28.5	24.5	cir	3	SW	0.5	122	32			5
5	3	31.5	29	35.5	29.5	c. n.	4	W	0.5	125	32.5			8
2	9	33.8	31	37.5	33.5	blue	0	NE	1	130	32			8
	7	30.5	26	32	29.5	c. n.	3	W	2	124	32			5
1	7	29.8	24.2	29.8	26.5	blue	0	NE	1	114	32			6
1	10	27.5	22.2	29.5	26.5	over	10	—	0	102	32			5
6	7	27	24	28.5	25.5	c. n.	9	W	1	119	31.5			7
3	6	23.8	19.2	25	21.5	cir	1	W	1	124	31.5	4	5	7
3	9	30.5	26	31	28	blue	0	E	0	120	31.5	1.6	2	4
5	1	29.8	29.5	29.8	26	over	10	E	0.5	62	31			16
8	5	30.2	30	31	29	over	10	W	1	51	31	2.5	36	12
9	3	31.5	30	31.5	28	blue	0	—	1	120	31.5	3.1	4	10
	6	30.2	26.5	31.5	28	blue	0	NE	1	120	31.5			8
7	5	30.5	27	33.5	28.5	blue	0	NE	0	127	31.5			7
	6	30.5		42	19.5		4	NE	0.7	113		53.2	56	7

METEOROLOGICAL OBSERVATIONS

	AT 9 A.M.						AT NOON							
	Dry Bulb.	Wet Bulb.	Maximum.	Minimum.	Cloud.		Dry Bulb.	Wet Bulb.	Maximum.	Minimum.	Cloud.		Elastic Force of Vapour.	Weight of Moisture in 10 cub. ft. of Air.
					Form.	Amount.					Form.	Amount.		
1883														
Dec.														Grains.
1	26	23.2	31.5	22.5	blue	0	31	23	31	25	blue	0	.056	7
2	26	24	29	23	over	10	27	26	29	26	cum	9	.132	16
3	19	18	21	18	blue	0	23	20	23	19	blue	0	.083	10
4	20	17	27.2	14	snow	10	24	20	29	16	snow	10	.074	9
5	16	16	24	14	snow	10	19	18	27	16	snow	10	.089	10
6	24	22	29	10	blue	0	29	27	30	22	blue	0	.130	15
7	4	3	29	0.5	blue	0	9	7.2	9.2	1.5	blue	0	.045	5
8	3.2	3	10	-2	blue	0	12.5	9	12.5	0	blue	0	.035	4
9	11	10	14	9.5	blue	0	20	15.5	20	9.5	blue	0	.050	6
10	20	18.5	20.5	12	c.c.	3	25	21.5	25	18	over	10	.085	10
11	18.2	17	26	13	c.c.	9	22	20	22.5	16	over	10	.091	11
12	23.8	21.5	26	20	c.c.	6	26	25	28	22	snow	10	.126	15
13	17.5	16.8	23	14	snow	10	18.2	17.5	19	15.5	snow	10	.090	11
14	31	29	36.5	18	c.n.	6	38.8	32.5	39.5	17.5	c.c.	5	.126	15
15	27	26	38	25	cum	3	37	32	38	25	cum	3	.139	16
16	23.2	21.5	38	18	over	10	27.5	25	27.5	22	over	10	.114	14
17	19	17.5	26	17	over	10	20	17.2	20	17	snow	10	.070	9
18	10	9	19	7	blue	0	10	8.5	11.5	6	blue	0	.050	6
19	11	10.2	14	-1	snow	10	20	17.5	22	9	over	10	.075	9
20	20.5	20	26	17	snow	0	23.5	21	24	18.5	c.c.	3	.092	11
21	21.5	21	27.5	20.5	blue	0	23.5	22.2	25	20	blue	0	.108	13
22	19.3	17	27	11	cir	3	27.2	22.8	27.5	17.5	cir	3	.085	10
23	15.5	14.8	28	11	blue	0	24.2	21.2	26	12.5	blue	0	.089	11
24	28.8	27.8	32	17	cir	3	35	28	35.5	27	blue	0	.094	11
25	30.8	29	37	22	c.n.	5	40.2	35.5	41.5	29	c.c.	2	.165	19
26	30	27	45	29	cir	1	41.5	34.5	43	29	blue	0	.135	16
27	23.5	21	40	20	blue	0	31	25.2	31.5	20	blue	0	.087	10
28	23.5	21	35	17	cir str	1	31	26.5	31	21	blue	0	.106	13
29	12	11	37	9	blue	0	17.5	15	18	10	blue	0	.065	8
30	11.8	11.5	20.5	5	c.n.	9	19	18.5	19	10	over	10	.095	12
31	12	11.5	20	12	c.n.	9	21	20	21	14	cum	6	.099	12

OF THE MALOJA.

AT 3 P.M.

Dew Point.	Drying Power of Air.	Dry Bulb.	Wet Bulb.	Maximum.	Minimum.	Cloud.		Wind.		Solar Radiation.	Earth Temperature.	Rain Gauge.	Snow.	Ozone
						Form.	Amount.	Dir. upr. and lower Currents.	Force.					
	Grains.											m m	Cent.	
5.7	13	29	23.1	29	27.3	blue	0	— NE	0	73	31.5			
24.5	1	21	20	27	20	over	10	N	3	50	31.5			
14.3	5	22	21	26	20	blue	0	— NE	1	65	31.5			
12	6	23	23	25	20	snow	10	N	2	63	31.5	2.1	2	
15.8	2	19	17	28	17	snow	10	N	3	46	31.5			
24.1	4	27	24	29	27	blue	0	— NE	5		31.5			
0.5	3	9	7	11	7	blue	0	— NE	1		30			
—4.5	6	14.5	10.8	15	10.5	blue	0	— E	2	124	30			4
3	7	21	17	25	18	blue	0	— E	1	113	30			5
14.8	6	23	22.5	25	18	snow	10	— E	0.5	46	30	1.4	1.5	7
16.4	3	23.5	22	25	20	snow	10	— E	0.5	70	30.5			7
23.5	2	23	21	29	22	snow	10	NW S	0.5	100	31	4	5	9
16	1	19.5	18.8	24	16	snow	10	SW SW	0.5	41	30.5			6
23.5	12	37	33.8	41	36	c.c.	5	W NE	0	123	31	8.2	9	7
25.6	9	38.5	31.5	40	35	cum	3	W	0	108	31.5			7
21.2	3	25	25	27	24	snow	10	SW SW	1	60	31.5			17
10.8	4	17	16	21	16.5	snow	10	— NE	1	60	31	8	9	6
3	2	10	8.5	11	7	blue	0	— E	1	84	30.5			5
12.3	4	20.2	18.2	22	18.5	over	10	NE W	0.5	83	30			7
16.5	4	25	22	26	22	c.n.	9	— W	0	110	31	2.5	3	6
20	2	26.8	24.5	29	21	cir.	1	— NE	0	120	31.5			7
14.8	7	27.5	23.8	30	25	blue	0	— N	0.5	117	31			5
15.8	4	30	26	30	22	blue	0	— NE	0	117	31			8
17	13	35.2	27.5	36	33	blue	0	— NNE	1	115	31.5			6
29.7	9	41.8	35.5	43.5	40	c.c.	2	NNE NNE	2	117	31.5			6
25	14	37.2	22.2	43	35	blue	0	— NNE	1	120	32			4
15.3	10	32	28	35	28.5	blue	0	— NE	2	108	31.5			4
19.6	7	32	27.2	33.5	28	blue	0	— E	1	118	32			4
9	3	20	20	23	16	fog	10	— NE	0	116	31.8			9
17.3	1	20	19.5	21	12.5	snow	10	SW	0	50	31.8			
18.2	1	20	20	21.5	19	fog	10	S SW	0	65	31.5			14
5.4	24.8						4.8		1	80		26.2	20.5	

METEOROLOGICAL OBSERVATIONS

	AT 9 A.M.						AT NOON.						
	Dry Bulb.	Wet Bulb.	Maximum.	Minimum.	Cloud.		Dry Bulb.	Wet Bulb.	Maximum.	Minimum.	Cloud.		Elastic Force of Vapour.
					Form.	Amount.					Form.	Amount.	
1884													
Jan.													
1	6.8	6.5	21	2.5	blue sky	0	14.5	13.8	16	2	blue sky	0	.075
2	16.8	15	17	4.5	c.c.	2	23.5	20.5	24	15	cum	7	.085
3	16.8	14.6	22	5	cum	6	22.5	19	22	14.5	cum	5	.073
4	24	23	25.5	18	cum	3	34	30	34	22	c.c.	4	.133
5	18	17	35	17	blue	0	28.8	25	29	16	cir	3	.103
6	25	23.0	29.5	12	over	10	30	26	30.5	23	c.c.	4	.107
7	16	15.2	29.5	11	c.c.	4	28	25	33	14	cum	2	.110
8	19.8	19.0	34.5	16	over	10	24	21	24	18	cum	10	.088
9	9.8	8.5	25	6.5	blue	0	16.8	13.8	16.8	7.5	blue	0	.056
10	24	22	25	13.5	blue	0	29	24	31	19	blue	0	.087
11	15	13	34	12	cir	1	33	28	39	13	blue	0	.111
12	23	22	40	12.5	cum	9	21.5	17.5	23.5	19.5	cum	5	.062
13	17.5	16	20	14	snow	10	18	17	19	15	snow	0	.085
14	25.5	23	25.5	12.5	cum	6	26	22.5	26.5	22.5	cum	4	.090
15	24	23	26	11.5	snow	10	25	24	26	22	snow	10	.120
16	29	28	29	22	snow	10	29.5	25.8	30	26.5	cum	6	.109
17	23	22	29	15.5	blue	0	25.6	23	26	21	blue	0	.101
18	23	20.8	29	13.5	blue	0	29.5	24	30	19	blue	0	.083
19	19	17.5	32	13	blue	0	29	24	29.5	17	blue	0	.087
20	15	13	34	11	blue	0	27	22	29	13	blue	0	.076
21	20	17.5	29	14	blue	0	30	25.2	31	18	blue	0	.096
22	21.8	19	23	17	blue	0	28.2	23.5	28.5	16	blue	0	.086
23	24	22	31	10.5	cn	7	30	25	30	20	cn	9	.093
24	27	24	31.5	19.5	blue	0	31.5	26.2	34.5	23.5	cum	4	.097
25	6.8	6	26	4.5	blue	0	15	12	15.5	4.5	blue	0	.049
26	2	1	24	-4	cir	2	16	12.8	16	0	blue	0	.050
27	19.5	19	22.5	11	snow	10	22	21.5	22.5	13	snow	10	.110
28	21.2	19.8	23	14.5	snow	10	24.2	22.5	26	17.5	snow	10	.105
29	18	17	20	10	snow	10	24	23.2	25	16	cum	6	.117
30	29	24	37	19.5	blue	0	28.5	26	31	20	blue	0	.120
31	19	17.5	29.5	9	cir	2	33.5	28	34	17	cir	3	.107

AT 3 P.M.

Time of day. alt. ft. of Air.	Dew Point.	Drying Power of Air.	Dry Bulb.	Wet Bulb.	Maximum.	Minimum.	Cloud.		Wind.		Solar Radiation.	Earth Temperature	Rain Gauge.	Snow.	Ozone.
							Form.	Amount.	Direction.	Force.					
9	12.2	1	16.8	16.8	19	15	fog	10	S	0	110	31.8	m.m.	ctm.	4
0	14.7	5	22	19.4	27	20.5	c.c.	3	N NE	0	101	31.5			7
9	11.7	5	23.8	21.8	25	20.5	cum	1	N	1	116	31			5
6	24.6	7	34.5	31	36	30.5	cum	8	NNE	1	120	30.5			4
2	19	6	27	24.2	31.5	25	c.c.	3	NE	1	112	31			7
3	19.8	6	27.8	26.8	33	22.5	c.c.	4	NW	0.5	115	31.5			4
3	20.4	5	34	28.5	36	26	cn	6	N	0.5	115	31			6
0	15.5	5	24.5	21.5	25	22	blue	0	NE	2	60	31			4
7	5.7	4	22	17.8	23	15	blue	0	E	2	95	30.5			7
0	15.3	8	32.5	23	35	27	blue	0	E	2	119	30.8			5
3	20.6	9	40	32.5	41	35.5	blue	0	W	0	121	30.8			4
7	7.9	7	19	16	22	17.5	blue	0	N E N E	2	99	30.5			7
0	14.7	1	18.8	17.5	20	15.5	snow	0	N E N E	5	40	31	1.2	3	7
1	16	5	24.5	22	26.5	22	cum	1	NE	3	117	30.5			4
4	22.4	2	24	23.2	26	22.5	snow	10	NE	0	45	30.8			9
3	20.2	6	28.5	25.5	30	27	cum	7	N	2	112	31	1.4	3.2	5
2	18.6	4	29	25	29	24.5	blue	0	ENE	3	121	31			8
0	14.3	9	32	27	33.5	27.5	blue	0	NE	1	123	31			4
0	15.3	9	32.5	30	36	27	blue	0	NE	3	122	31			6
9	12.5	8	30	26	36	24	blue	0	ENE	3	126	31			5
1	17.5	7	32	27	36	27.5	blue	0	NE	2	126	31			5
0	15	8	30	25.2	31.5	26.5	blue	0	NE	1	128	31			6
1	16.7	8	37	32	38	28	cum	4	NW W N NE	0.5	88	31			4
2	17.7	8	25.5	22.2	32.5	29	blue	0	N NE	1	122	31			4
6	2.5	4	23.5	17.8	23.5	13.5	blue	0	E	1	118	30.8			6
6	3	4	22.5	18	22.5	14	blue	0	NE	0	124	30.5			7
3	20.4	1	22.5	21.8	23	17	cum	6	W SWW	2	60	30.5	3.4	6	18
3	19.4	2	19.8	19.2	25	18	snow	10	W W W	0	72	30.5			17
4	21.8	1	30	28.8	32	22.5	cn	7	N W	0	103	30.5	7.3	14	15
4	22.4	4	29	26.5	36	27	cum	3	NW	0.5	129	30.5			7
3	19.8	9	31	27	34	28	cum	4	W WSW	0	113	30.5			5

METEOROLOGICAL OBSERVAT

	At 9 A.M.						At Noon						
	Dry Bulb.	Wet Bulb.	Maximum.	Minimum.	Cloud.		Dry Bulb.	Wet Bulb.	Maximum.	Minimum.	Cloud.		Elastic Force of Vapour.
					Form.	Amount.					Form.	Amount.	
1884 Feb.													
1	27.2	26.5	31.5	13.5	c.n.	6	30	24.5	31	25	c.n.	6	.086
2	27	27	29	25	sn.	10	29	28.5	29.5	25.5	c.n.	9	.151
3	11.8	11	29	7	blue	0	23	21	23	9.5	c.c.	1	.096
4	12	11	28	3	blue	0	24.5	19	24.5	9	blue	0	.057
5	15.5	14	30	13	blue	0	29	25	30	18.5	blue	0	.101
6	15	13	37	7	c.c.	2	26.8	22.2	29	13	blue	0	.080
7	17.2	16	32	6	blue	0	27	23.2	29	14	blue	0	.092
8	22	20	31.5	7	cum	9	29	24	32	20	over	10	.087
9	15.2	13	31	11	c.c.	3	31	27.5	31	13	c.c.	4	.120
10	15.5	14.5	32	9	c.c.	6	29	26	30	15	c.c.	1	.116
11	26	25	32	13.5	over	10	31	29	31.5	29	over	10	.143
12	29.8	29.5	25	31	fog snow	10	33.5	32	38	27.5	sn.	10	.168
13	18	17	35	14	blue	0	29	26	29	15	blue	0	.116
14	13.5	12	32	4	blue	0	31	26	31	11	blue	0	.099
15	24	23.5	35	12	sn.	10	21.2	21	24	19	sn.	10	.111
16	18	18	21.5	15	sn.	10	18.5	18	22	15	sn.	10	.093
17	13	12	19.5	3	c.c.	6	23	22	26	20	c.c.	4	.109
18	15	15	21	10.5	sn.	10	17	16	18	12.5	cum	7	.081
19	1	1	18	-7.5	blue	0	17	16	17	-2	blue	0	.081
20	2	1	28.5	-6.5	blue	0	23.5	20.1	23.5	-0.5	blue	0	.079
21	10.8	9	24	-2.5	blue	0	31.5	25.8	32.5	8.5	blue	0	.092
22	19	17	32	3	c.c.	2	33	28	36.5	16.5	cir	3	.111
23	30.5	27.5	34	16	c.n.	4	30.2	27.5	32	27	over	10	.127
24	28.5	28	30	29	c.c.	7	30	29	31.5	27	sn.	10	.151
25	17	15.8	30.5	9.5	c.c.	1	28	24	31.5	14	cum	4	.095
26	18	17	29	8	c.c.	2	24	20	24.5	15	cum	4	.074
27	11.8	10.2	23	-2.5	c.c.	2	22	18	24	8.5	c.c.	1	.064
28	6.5	5	23	-2.5	c.c.	3	21.8	20	22.5	4	cum	8	.093
29	23	22	23	16	cum	3	25.5	24	26	20.5	cum	3	.116
	17.3		37	-7.5		4	26.5		38	-2		4.3	

AT 3 P.M.

Dew point.	Drying power of Air.	Dry Bulb.	Wet Bulb.	Maximum.	Minimum.	Cloud.		Wind.		Solar Radiation.	Earth Temperature.	Rain Gauge.	Snow.	Ozone.
						Form.	Amount.	Direction.	Force.					
15.1	9	28.5	28	31.5	27	cum	3	SW	2	125	30.5			20
27.6	1	27.5	27.5	30	27	fog	10	SW W WSW	0.5	89	30.8	3.1	4	20
17.5	4	27	24	27.5	21	c.c.	4	NE NE	2	120	31			11
6.1	9	27	22.5	30	20.5	blue	0	— NE	2	132	31			5
18.7	7	36	30.5	38	27.5	blue	0	— NE	0.5	130	31			7
13.5	8	32.5	27	34.5	24.5	blue	0	— NE	0.5	126	31			5
16.5	6	31	26	33	26	blue	0	— NE	0.5	126	31			4
15.3	8	30	26	32	27	c.c.	5	WSW WSW	0.5	88	31			4
22.4	6	31.8	28.8	33	29	c.c.	8	SW SW	0	125	30.8			8
21.6	4	32	29	33	29	c.c.	3	— NE	0	134	31			8
26.3	3	30	27.8	32	28.5	c.c.	8	W W	0.5	66	31			7
30.1	2	32.8	32	35	31	cum	9	— SW	0	124	30.8	4.3	7	11
21.6	4	31	26.5	34	27	blue	0	— NE	1	143	30.8			5
18.2	8	34	28	34.5	29	blue	0	— NE	1	136	31			5
20.6	1	21	21	22	19	sn.	10	SW SW	1	50	31			15
16.8	1	19	19	19.5	15.5	c.c.	7	W SW	1	52	30.8	10.5	17	17
20.2	2	20	20	23	19	over	10	— NNW	0	85	31			14
13.7	1	18	17.5	19	15	c.c.	6	— SW	0	100	31	2.1	3	10
13.7	1	24	22	24	15	blue	0	— N	0.5	131	31			6
13.3	5	24	20.2	30	22	blue	0	— NE	0	137	31			7
16.5	9	32	26	34	28	blue	0	— SW	0	135	31			6
20.6	9	33	27.5	36	31	cn	3	— SW	0	130	30.8			6
23.6	4	29	27.5	31	27	c.c.	8	SW SW	0.5	75	30.5			12
27.6	1	30	29	31	27.5	c.c.	7	W W	1	60	30.5	1	2	18
17.3	7	28	24	30	25	cum	4	N NE	1	100	30.5			7
12	6	22	19	26	20	c.c.	2	N NE	2	125	30.5	0.4	1	7
8.7	6	23	19	24	20	blue	0	NW NW	1	135	30.5			6
16.7	3	20	19	22	17	cn	8	W W	1	93	30.5			5
21.6	2	25.2	23.8	27	23	cum	9	W W	1	81	30.5			9
	4.7	27.5		38	15		4.3		0.7	108		21.4	34.4	9

SUMMARY OF METEOROLOGICAL OBSERVATIONS, TAKEN IN THE

	AT 9 A.M.							AT NOON.						
	Air Temperature.		Maximum.		Minimum.		Amount of Cloud.	Air Temperature.		Maximum.		Minimum.		Amount of Cloud.
	Fahr.	Centi-grade.	Fahr.	Centi-grade.	Fahr.	Centi-grade.		Fahr.	Centi-grade.	Fahr.	Centi-grade.	Fahr.	Centi-grade.	
Nov., '83	26.4	-3.1	44	6.7	11	-11.7	4	30.5	-0.8	43	6.1	15	-9.4	4
Dec., '83	19.3	-7	45	7.2	-2	-18.3	4.1	25	-3.9	43	6.1	0	-17.8	4
Jan., '84	19.3	-7	40	4.4	-4	-20	4	25.7	-3.5	39	3.9	0	-17.8	3
Feb. '84	17.3	-8.1	37	2.8	-7.5	-22	4	26.5	-3	38	3.4	-2	-18.9	4
	20.6	-6.3	45	7.2	-7.5	-22	4	26.9	-2.8	43	6.1	-2	-18.9	4

Mean Temperature for the Winter = 25° Fahr. (-3° 9 Centigrade).

Average weight of Moisture = 12 grains per 10 cubic feet of Air.

Average drying Power of Air = 5.4 grains per 10 cubic feet of Air.

Average daily quantity of Ozone, with 6 hours exposure (scale 0° to 20°) 7°.4.

Highest Maximum Thermometer = 45° Fahr. (7°.2 Cent.), 26th December.

Lowest Minimum Thermometer = -7°.5 Fahr. (-22° Cent.), 19th February.

Highest Solar Radiation = 143° Fahr. (61°.6 Cent.), 13th February.

MALOJA DURING THE WINTER 1883-84, BY A. TUCKER WISE, M.D.

		AT 3 P.M.												
Weight of moisture in 10 cub. ft. of air.	Drying power of air per 10 cubic feet.	Air Tempe- rature		Maximum.		Minimum.		Amount of Cloud.	Force of wind.	Mean Solar Radiation.		Rain Gauge.	Snow.	Ozone, 6 hours exposure.
		Fahr.	Centi- grade.	Fahr.	Centi- grade.	Fahr.	Centi- grade.			Fahr.	Centi- grade.			
Grains.	Grains.							1 to 10.	1 to 12.			mm.	Centi- mètres.	1 to 10.
14	6	30.5	-0.8	42	5.6	19.5	-7	4	0.7	113	45	53.2	56	7
11	5.4	24.8	-4	43.5	6.4	7	-13.9	4.8	1	89	31.7	26.2	29.5	7
11	5.4	27.3	-2.4	41	5	13.5	-10.2	2.8	1.3	105	40.6	13.3	26.2	6.6
12.2	4.7	27.5	-2.5	38	3.3	15	-9.4	4.3	0.7	108	42.2	21.4	34	9
12	5.4	27.5	-2.4	43.5	6.4	7	-13.9	4	0.9	104	40	114.1	145.7	7.4

TABULAR COMPARISON OF THE FOUR

MALOJA.

Position.—Situated at the S.W. end of the Upper Engadine. Receives more sun than any other Alpine Station. No morning nor evening mist. The Kursaal is placed on the western side of the plateau, and in this situation escapes most of the valley draught (thalwind). Extremely picturesque scenery at this part of the Engadine. Mountain screen, 3,000 to 6,000 feet.

Proximity to Glaciers.—Partially surrounded by glaciers.* Bernina glaciers, W.; Murtel, 10 kilometres; Fedoz, Forno, and Albigna, 7 to 10 kilomètres distant, on the S.W. and S.

* The effects of glacial air on the respiration has been noted by Dr. Burney Yeo, who considers that its condensed state gives a sensation of freedom in breathing.—“Health Resorts and their Uses.”

Pernicious Winds.—South-west wind. Fairly-well sheltered from South wind. The sultry characters of the föhn appear to be extremely rare, probably from the influence of the surrounding glaciers.

Thalwind (valley wind).—Blows from the N.E. principally on the eastern side of the plateau, in a line with the Bergell valley, and is mostly noticed on the Kulm, about one mile distant from the hotel.

WIESEN.

Wiesen is located on the hill-side, the Landwasser being about 1,000 feet below the houses. The mountain screen ranges from 3,000 feet to 5,000 feet above. There is in fine weather no morning nor evening mist, and no smoke accumulates, as, with rare exceptions, a constant but imperceptible current of cold air travels down the declivities. Notwithstanding that this motion of the atmosphere is unfelt, it is sufficient to obviate any tendency to stagnation.

Scaletta, 13 miles to the E. (Small glacier.) Silvretta Glacier, 20 miles to N.E.

The pernicious winds are the south and south-west currents, known locally by the name of “föhn.”

This wind, which blows in every Swiss valley, is infrequent, as the village is situated far above the bed of the gorge, and consequently out of the zone of commotion of air, caused by the descending cold currents which converge and flow down the gorges and ravines.

HEALTH RESORTS IN THE GRISONS.

DAVOS.

Situated on rising ground in the valley itself. The mountain screen ranges from 3,000 feet to 5,000 feet. A perceptible mist generally covers the valley each morning; this is soon dissipated by the sun, but the usual haze which is seen over villages and small towns is plainly apparent, and remains stagnant all day, unless moved by wind.

Scaletta, 8 miles to S.E. Silvretta,
12 miles N.E. by E.

South and south-west winds, as at
Wiesen.

A gusty thalwind is often felt between
2 and 3 P.M. from the N.E.

ST. MORITZ.

The Kulm Hotel is about 300 feet above the lake. A thin mist hangs over the lake in the early morning, but below the level of the dwellings. The aspect is more open than Davos, but is sheltered from wind.

Extensive Bernina glaciers to S. and
S.W. Piz Bernina is 15 kilomètres
distant.

South and south-west wind.

No regular valley wind, blowing from a definite direction, independently of the upper current. Nevertheless, the wind rises regularly in the course of the day, and is unpleasant from either the north or south (Mr. Waters' observations). Thalwind felt on the St. Moritz Kulm.

NUMBER OF HOURS OF POSSIBLE SUNSHINE DURING WINTER.

		MALOJA.†		WIESEN.		DAVOS.		ST. MORITZ. *
1 Nov.	...	7 $\frac{1}{8}$ hours	...	7 $\frac{1}{3}$ hours	...	7 $\frac{1}{3}$ hours	...	— hours
15 „	...	6 $\frac{3}{4}$ „	...	— „	...	— „	...	6 „
1 Dec.	...	6 $\frac{1}{2}$ „	...	5 $\frac{1}{4}$ „	...	5 $\frac{1}{4}$ „	...	5 $\frac{1}{4}$ „
15 „	...	6 $\frac{1}{4}$ „	...	5 $\frac{1}{12}$ „	...	5 $\frac{1}{12}$ „	...	5 $\frac{1}{12}$ „
1 Jan.	...	6 $\frac{1}{10}$ „	...	5 $\frac{1}{6}$ „	...	5 „	...	5 $\frac{1}{12}$ „
15 „	...	6 $\frac{1}{3}$ „	...	5 $\frac{5}{6}$ „	...	5 $\frac{1}{2}$ „	...	5 $\frac{1}{3}$ „
1 Feb.	...	6 $\frac{5}{8}$ „	...	7 $\frac{1}{4}$ „	...	6 $\frac{1}{3}$ „	...	7 $\frac{1}{12}$ „
15 „	...	7 $\frac{1}{8}$ „	...	7 $\frac{2}{3}$ „	...	7 $\frac{3}{4}$ „	...	8 $\frac{1}{12}$ „

* Das Oberengadin (Dr. J. M. Ludwig). There is no meteorological station at St. Moritz.

† These observations were taken from one spot in the Maloja (The Kursaal); if the hâlets or grounds of the hotel had been included, a very much larger duration of sunshine would be recorded.

MEAN DAILY TEMPERATURE AT 7 A.M., GIVEN IN DEGREES CENTIGRADE.


		Nov.	Dec.	Jan.	Feb.	Mean for the whole winter '83-84.
Maloja	(6,000 ft.)	—3·1	—6·3	—6·5	—7·9	—5·9
Wiesen	(4,771 ft.)	—1·5	—5·3	—3·3	—3·1	—3·3
Davos	(5,105 ft.)	—3·7	—7·1	—5·9	—5·5	—5·5
Andermatt	(4,738 ft.)	—3·4	—8·1	—5·5	—5·0	—5·5

† Furnished to the author by Professor Billwiller, Director of the Meteorological Stations.

* Calculated from the observations of M. Kuoni.

HYGROMETRIC AND BAROMETRIC CURVES FOR THE MALOJA, during the Winter 1883-84, given in millimètres of pressure and grains of vapour in 10 cubic feet of air.

The calculation of vapour was made at Noon, and the barometer noted one hour after.



THE ALPINE CLIMATE SERIES.

By the same Author.

DAVOS PLATZ,

AND THE EFFECTS OF HIGH ALTITUDE ON PHTHISIS.

Fcap. 8vo.

"As far as we can see, Dr. Wise has collected all the information invalids need in a convenient form."—*Saturday Review*.

"A good deal of useful information is afforded."—*Edinburgh Med. Journal*.

"This is about the best of the many books on Davos which we have read. There can be no doubt that Davos is an invaluable resort for many young and active people with incipient phthisis. Dr. Wise's little book will be found full of useful information."—*Lancet*.

"The remarks on the therapeutical effects of cold are excellent, and in a few words show the why and the wherefore of a dry cold air being so eminently beneficial to a large class of consumptive patients. It will not take long for anyone to read Dr. Wise's book through, and we should advise all contemplating a visit to Davos to get the work and study it well."—*Medical Press and Circular*.

"Contains a great deal of information."—*British Medical Journal*.

J. & A. CHURCHILL, 11 New Burlington Street, London.

WIESEN AS A HEALTH RESORT IN EARLY PHTHISIS, WITH DIRECTIONS FOR DIET, CLOTHING, AND EXERCISE IN THE SWISS ALPS DURING WINTER.

"We can recommend Dr. Wise's Handbook for Wiesen as full of practical details, and bristling with statistics."—*Saturday Review*.

"It is the first treatise written with strict impartiality by a physician of experience, who has taken the trouble to visit the places of which he speaks, to reside many months in each of them, and to compare the results of his observations with candid frankness. Dr. Wise does not puff any one locality. He does not flatter existing institutions, as is proved by his severe remarks on hotel ventilation (pp. 6, 37). He is no one-sided enthusiast, as may be gathered from his chapter on "The Drawbacks of High Altitude Stations." His plain-spoken criticism adds weight to what he finds to say in favour of each place he treats of, and is calculated to inspire the intending visitor with confidence in the substantial soundness of the system."—*Pall Mall Gazette*.

"Those seeking information concerning new health resorts for consumption should read this guide-book."—*New York Medical Record*.

"It abounds in interesting and useful information, Physiological, Local Hygienic, Sanitary and Meteorological."—*Canadian Practitioner*.

BAILLIERE, TINDALL, & COX, 20 King William Street, Strand W.C.