Final report of the Anthropometric Committee / consisting in 1882-3 of F. Galton (chairman) [and others]; drawn up by C. Roberts and Sir Rawson W. Rawson.

Contributors

British Association for the Advancement of Science. Anthropometric Committee.

Galton, Sir Francis, 1822-1911. Roberts, Charles, 1836-1901. Rawson, Sir Rawson William, 1812-1899. Royal College of Surgeons of England

Publication/Creation

[London]: [Printed by Spottiswoode], [1883]

Persistent URL

https://wellcomecollection.org/works/hjdm8ta8

Provider

Royal College of Surgeons

License and attribution


This material has been provided by This material has been provided by The Royal College of Surgeons of England. The original may be consulted at The Royal College of Surgeons of England. Where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org

Final Reports of the Anthropometric Committee fle Butish Apociation fate Advancement of Voience 11/25



Final Report of the Anthropometric Committee, consisting in 1882-3 of Mr. F. Galton (Chairman), Dr. Beddoe, Mr. Brabrook (Secretary), Mr. Frank Fellows, Mr. James Heywood, Professor Leone Levi, Dr. F. A. Mahomed, Mr. J. E. Price, Lieut.-General Pitt-Rivers, Sir Rawson W. Rawson, and Mr. C. Roberts. Associates, Dr. T. G. Balfour, Dr. J. H. GLADSTONE, Inspector-General Lawson, Dr. W. OGLE. Drawn up by Mr. C. Roberts and Sir Rawson W. Rawson.

[PLATES IV .- X.]

1. The Committee, originally appointed in 1875, and aided by successive grants, of which it has expended 2801., has made a Report in each of

the five years 1878 to 1882, and now submits its final Report.

2. Not that the work open to the Committee is exhausted, although it has to a great extent supplied what was pointed out in its Reports of 1881 and 1882 as chiefly wanting, or that its conclusions are to its own mind complete and satisfactory. But it would require more time and larger funds than are at the disposal of the Committee to prosecute its inquiries, even with the materials now in its possession, to the end which it has had in view; and the Committee is of opinion that the most useful course will be to bring before the Association the results of its past labours, indicating at the same time the conclusions which it considers to be sufficiently established by the facts ascertained, and the deficiencies, both of data and methods, which remain to be supplemented, either by individual exertion, or by the reappointment of a similar Committee at some future period under the auspices of the Association.

3. In order to furnish a complete review of the information obtained, it will be necessary to refer to tables and data contained in previous

reports. A list of these Reports is furnished in a note.2

Objects and Operations of the Committee.

4. The Committee was appointed for the purpose of collecting observations on the systematic examination of the height, weight, and other

physical characters of the inhabitants of the British Isles.

5. Its operations in each year are described in the introduction to its Report of 1881. The description and amount of the statistics which it has collected, and the names of the persons to whom it is indebted for the collection, are detailed chiefly at the commencement of its several Reports from 1880 to 1882.

6. Among the objects early aimed at by the Committee, and prosecuted by it up to the year 1881, was the collection and comparison of photographs of the typical races of the United Kingdom; but at the meeting of that year this inquiry was assigned to a separate Committee, upon whom will devolve the duty of reporting upon this branch of the general subject.

¹ The late Dr. William Farr was a member, and Chairman of the Committee from 1875 to 1879.

² 1, Report for 1878, 5 pp. (numbered pp. 182-6 in the Annual Report of the Association). 2, Report, 1879, 35 pp.; *ibid.* pp. 175-209. 3, Report, 1880, 41 pp.; *ibid.* pp. 120-59. 4, Report, 1881, 48 pp.; *ibid.* pp. 225-72. 5, Report, 1882, 3 pp.; *ibid.* pp. 278-80. An Index to the Tables is given in Appendix C.

- 7. The points to which the Committee has addressed its inquiries
 - Stature.
 Weight.

(3) Girth of chest.

- (4) Colour of eyes (5) , hair Complexion.
- (6) Breathing capacity.(7) Strength of arm.

(8) Sight.

(9) Span of arms.

To these might have been added others, especially-

(10) Size and shape of head.

(11) Length of lower limbs as shown by the difference between the sitting and standing positions.

(12) Girth, length, and breadth of other parts of the body.

But the Committee was afraid of seeking to obtain more information than their contributors would be likely to furnish; and experience has shown that many of them have been unable to supply more than a portion of that which was requested. Few have furnished complete returns on all the subjects, but where one has failed another has succeeded, and sufficient data have been collected to give trustworthy statistical results on all the subjects of inquiry except those of breathing capacity and sight. An abstract of one of the complete returns will be given in its proper place, as exhibiting a good epitome of what the Committee has sought to obtain in all cases. (See Table XXIII.)

8. The large body of observations on stature, weight, and complexion collected by Dr. Beddoe, and those on stature, weight, and chest-girth collected by Mr. Roberts, previously to the formation of the Committee, have been made use of; and the Committee has thus had observations made on a total number of about 53,000 individuals of both sexes and of all ages, from which to construct their tables and to base their conclusions.

9. The statistics are unique in range and numbers, and have been obtained from a very large number of independent observers living in different parts of the country, without prejudice, and often in ignorance of the use which would be made of them; and they have been analysed and tabulated in a perfectly impartial manner, irrespective of all preconceived opinions. The Committee does not claim for them exemption from the liability to that amount of imperfection and probable error which must attach to all conclusions drawn from a disproportionate, and from a comparatively small number of observations. But great care has been taken in the examination and classification of all the returns to eliminate obvious errors, and to call attention in the body of the Report to any apparent discrepancies from faulty observation or deficient numbers.\(^1\)

^{&#}x27;If an exceedingly large number of measurements, weights, &c. be taken—supposing no bias, or any cause of error acting preferably in any one direction to exist—not only will the number of small errors vastly exceed that of large ones, but the results will be found to group themselves about the mean of the whole always according to one invariable law of numbers, and that the more precisely, the greater the total number of determinations. . . . Rude and unskilful measurements of any kind, accumulated in very great numbers, are competent to afford precise mean results. The only conditions are the continual animus mensurandi, the absence of

Methods.

10. The forms and instruments used have been explained in the Reports for 1878 and 1880; but practical difficulties have been found to exist in obtaining trustworthy observations with regard to breathing capacity. Experience has also led the Committee to believe that the use of Snellen's test-types for sight, Nos. 1 and 10, is more convenient, and will yield more trustworthy results, than that of the army test-dots, which were adopted in its original circulars. Since 1879, also, the Committee has introduced the use of cards for recording the observations relating to single persons, which has been extensively adopted in Germany and the United States, and recently by the Investigation Committee of the British Medical Association, and which offers great facilities in analysing and grouping the facts observed. The Committee appends copies of the forms of the cards and of the methods of measurement and observation which they have employed. (See Appendix A.)

11. The difference between the average and mean of a number of observations, and its importance in dealing with the subjects under consideration, has been pointed out and discussed by Mr. Roberts in the Report for 1881, at p. 233; ² and the special sense in which Mr. Roberts employs the term mean, being that value in an arithmetic series of observed values of which the observations are the most frequent, has been adopted by

the Committee.3

12. In connection with the question of the applicability of the exponential law of error to statistical results relating to anthropometry, Mr. Francis Galton has contributed a valuable series of tables, with remarks, on the range in height, weight, and strength, in which he introduces his method of the calculation of deciles, quartiles, and medians.⁴

bias, the correctness of the scale with which the measures are compared, and the assurance that we have the entire range of error, at least in one direction, within the record.'—Sir J. F. W. Herschel, *Edin. Rev.* vol. xcii.

1 See the Report for 1881 for a discussion of this subject by Mr. Lawson and

Mr. Roberts.

² Also in a note at p. 121 of the Report for 1880.

3 Mr. Roberts has followed Quetelet in the use of the word mean, and its difference from an average is thus explained by Sir John Herschel. Speaking of Quetelet's homme moyen he says:—'Now, this result, be it observed, is a mean as distinguished from an average. The distinction is one of much importance, and is very properly insisted on by M. Quetelet, who proposes to use the word mean only for the former, and to speak of the latter (average) as the "arithmetical mean." An average may exist of the most different objects, as of the height of houses in a town, or the size of books in a library. It may be convenient to convey a general notion of the things averaged, but involves no conception of a natural and recognised central magnitude, all differences from which ought to be regarded as deviations from a standard. The notion of a mean, on the other hand, does imply such a conception, standing distinguished from an average by this very feature, viz., the regular march of the groups, increasing to a maximum and then again diminishing. An average gives us no assurance that the future will be like the past. A mean may be reckoned on with the most implicit confidence. All the philosophical value of statistical results depends on a due appreciation of this distinction, and acceptance of its consequences.'- Edin. Rev. vol. xcii. Mr. Galton, however, desires to state that considering many statistical groups which are regular in their distribution are at the same time normally asymmetrical, he does not recognise the expressions of 'mean value 'and 'the value most likely to be observed 'as strictly equivalent.

Report for 1881, p. 245.

TABLE I.—Showing the STATURE, WEIGHT, CHEST-GIRTH, and STRENGTH Kingdom, arranged according

| | | | | | STATU | JRE | MODE. | | | | | | | |
|---|---|---|--|------------------------|---|---|--|------------------------|------------------|---|--|---|--|------------------------|
| Height without shoes Scotland | | nd | Ireland | | England | | Wales | | Total | | Weight with | | th Scotl | |
| nches | Mètres | No. of observations | No. per 1,000 | No. of observations | No. per 1,000 | No. of observations | No. per 1,000 | No. of observations | No. per 1,000 | No. of observations | No. per 1,000 | lbs. | kilos. | No. of observations |
| 77- 76- 75- 74- 73- 72- 71- 70- 68- 66- 65- 64- 63- 62- 61- 60- 58- 58- | 1'957 1'931 1'906 1'881 1'855 1'830 1'804 1'779 1'754 | 1 4 6 15 26 69 102 115 218 210 139 109 47 19 9 2 2 1 | 1 3 4 12 20 53 78 88 167 -161 107 84 36 14 7 2 1 | | - 8 29 44 72 116 179 -211 167 96 44 20 6 5 - 3 | 1 1 9 16 48 117 254 473 753 886 918 | - 2 2 8 8 19 41 76 122 143 -148 142 119 85 52 20 12 6 2 1 | | | 2 5 16 32 79 202 392 646 1063 1230 —1329—1223 990 669 394 169 83 41 14 4 2 | 1 2 3 9 24 46 75 124 143 -155 143 115 78 46 20 9 5 | 280 270 260 250 240 230 220 210 200 190 180 170 160 130 120 110 100 90 | 127'3 122'7 118'2 113'6 109'1 104'5 100'0 95'5 90'9 86'4 81'8 77'3 72'7 68'2 63'6 59'1 54'5 50'0 45'5 40'9 | |
| Total | | 1304 | 1000 | 346 | 1000 | 6194 | 1000 | 741 | 1000 | 8585 | 1000 | Total | | 1212 |
| Averag | ge inches mètres | 68:71 1'746 | = | 67·90 1·726 | = | 67·36 1·712 | = | 66.66 1.694 | = | 67:66 1'720 | = | Aver | rage lbs. | 165-3 75°1 |
| Mean | inches | 68·5 1·741 | - | 67 5 | = | 67·5 1·715 | -= | 1.690 1.690 | - | 67·5 1·715 | 1-1 | Mea. | n lbs kilos | 160-0 72'7 |
| Heigh | t÷weight es per lb. weight) | -416 | - | *441 | - | •435 | - | *421 | - | •428 | - | - Weight+hgt. (lbs. per in. of height) | | |

Note.—The factors in the bottom line give some means of ascertaining the most probable stature, weight, chest-girth, or strength of a man, when only one of these data is known. They also give modified values when the birthplace of the man is also known, whether it be in Scotland, Ireland, England, or Wales. The results so obtained are based on the supposition that the proportion between the values of these qualities is constant, which is practically true for values that do not differ widely from the mean.

The method of employing the factors is simple: thus, the first five of them are the number of inches in height divided by the number of pounds in

Adult Males (age from 23 to 50) of the Population of the United lace of Birth.

| | WE | IGHT | N TO | in fact | NI W | ted in | | 1 | CHEST- | GIRTH | | | STRI | ENGTH | |
|----|---|------------------------|----------------------|------------------------|------------------|--|------------------|--|---|---------------------------|---|-------|--------------------|------------------------|------------------|
| 18 | iles | Eng | agland Ireland Total | | chest | Empty chest-girth: Total: chiefly military English | | Strength: drawing- power, as in drawing a bow | | Total: chiefly English | | | | | |
| | No. per 1,000 | No. of observations | No. per 1,000 | No. of observations | No. per 1,000 | No. of observations | No. per 1,000 | Inches | Centimètres | No. of observations | No. per 1,000 | lbs. | kilos. | No. of observations | No. per 1,000 |
| | 1 — 1 — 3 2 2 111 9 466 1388 182 — 242 7 — 92 31 13 3 — — | | | | | 1 — 1 8 11 16 41 85 107 263 476 787 1326 —1559—1623 867 390 152 34 2 — | | | 114'3 111'7 109'2 106 6 104'1 101'6 99'0 95'5 93'9 91'4- 88'9 86'3 83'8 81'2 78'7 76'2 73'6 71'1 68'5 | | 1 2 6 6 17 22 35 63 97 130 —173 162 158 75 35 12 10 2 — — | | | | |
| I | 1000 | 5552 | 1000 | 247 | 1000 | 7749 | 1000 | Total | | 3407 | 1000 | Total | | 1497 | 1000 |
| | 11 | 155-0 70'5 | - | 154·1 70°0 | Ξ | 158·2 71·9 | = | Avera | ge ins | 36:46 92:6 | 1 | Avera | age lbs. kilos. | 79·6 35·2 | = |
| | - | 150·0 68·2 | = | 150·0 68·2 | - | 155·0 70°5 | _ | Mean | inches . | 36·50 92°7 | = | Mean | lbs kilos | 77·5 35°2 | - |
| | - | 2:301 | - | 2:270 | - | 2:323 | - | Girt Girt | h÷hgt. h÷wgt. | ·542 ·235 | - | | th.÷ht. th.÷wt. | 1·182 ·513 | = |

weight, in the five following cases, natives of Scotland, Ireland, England, and Wales, and in the British Isles generally. The factor for Scotland is 0.416, consequently a Scotchman whose weight is 150 lbs. has most probably a height of 150×0.416 inches, or 62.4 inches. Similarly, in the next group of pounds of weight divided by inches of height, the factor for Englishmen is 2.301, consequently an Englishman 66 inches in height should weigh 66×2.301 lbs., or 152 lbs. In the same way we may calculate the other elements by the remaining factors.

Summary of Information Obtained.

13. The Committee submit in this, its final Report, a review of all the information which it has collected under the different heads of inquiry, giving references to those tables and conclusions which have been published in its previous Reports, and adding such others as it has been able to draw from the several sources at its command.

14. The first object of the Committee has been to ascertain the prin-

cipal characteristics of the adult population :-

a. As to the stature, weight, chest-girth, and strength of the whole country and of each of its four provinces, shown in Table I., pages 256, 257.

b. The relative stature, weight, and strength of men and women.

Table II., page 261.

c. The stature, weight, and complexion (colour of eyes and hair) of men in different counties as indicating their racial origin, and the influence of soil, climate, occupation, and other sanitary surroundings. Tables III. and IV., and Plates V.–IX., pages 262 to 265.

d. The relative stature of men of British origin, and that of other nationalities and races as far as they have been ascertained. Tables V.

and VI., pages 268, 269.

15. The second object the Committee has had in view has been to ascertain the rate of growth and development of children of both sexes under different conditions of life (media); the period of the attainment of maturity; and the influence of advancing age on the physical condition of the body. Tables XII. to XXV.

ADULT POPULATION OF THE BRITISH ISLES.


a. Adult Males-Table I.


16. Table I. shows the stature, weight, chest-girth, and strength of adult males of the ages from twenty-three to fifty years, the number of men at each measurement, and the ratio per thousand of the male population.

17. The observations are grouped according to the place of birth in England, Wales, Scotland, and Ireland; and, with the exception of the Irish, they were chiefly derived from the division of the country under which they are entered in the table. The Irish returns are almost entirely those of men born in Ireland, but living in England, Scotland, or Wales; and the Committee regrets that it has not been able to obtain more than one return direct from Ireland. The Scotch and Welsh by birth, living in England, have been entered under their respective nationalities. The columns are arranged in the order of the superiority of the average

stature and weight.

18. The general results indicated by this table may be summarised as follows:—In height the Scotch stand first (68.71 inches; 1.746 mètres), the Irish second (67.90 inches; 1.726 mètres), the English third (67.36 inches; 1.712 mètres), and the Welsh last (66.66 inches; 1.694 mètres), the average of the whole being 67.66 inches (1.720 mètres). In weight the Scotch take the first place (165.3 lbs.; 75.1 kilos.), the Welsh the second (158.3 lbs.; 71.9 kilos.), the English the third (155.0 lbs.; 70.5 kilos.), and the Irish the fourth (154.1 lbs.; 70.0 kilos.), the average weight of the whole being 158.2 lbs. (71.9 kilos.). Thus the Scotch are the tallest and heaviest, the English take the third place in both tables, while the position of the Welsh and Irish is reversed—the





Irish, occupying the second place in stature, come last in weight, and the Welsh, though lowest in stature, stand second in weight. For each inch of stature a Scotchman weighs 2.406 lbs., a Welshman 2.375 lbs.,

an Englishman 2:301 lbs., and an Irishman 2:270 lbs.

19. The columns showing the number of individuals per thousand at each height, besides showing in a uniform manner the relative stature and weight of the different nationalities, will be useful to military surgeons for determining the minimum stature of recruits for the army. From the run of the figures it is obvious that if each country has to contribute its relative quota of soldiers, the minimum standard for Welsh recruits should be two inches lower, and for English and Irish recruits one inch lower, than for Scotch recruits. This difference in the relative stature is best shown by the black line running across the table, which marks the mean height—that is to say, the height at which the greatest number of observations occur in each nationality.

20. It is probable that too much importance has been attached to stature in selecting recruits for the army in this country, and that a high standard does not necessarily produce men best fitted for military duties. In the Report for 1879 are given two tables of the stature and weight of the English, Scotch, and Irish recruits for the years 1862–3, when the minimum standard of height was 66 inches (1.677 mètres), and in 1864–65, when it was reduced to 65 inches (1.626 mètres); and the result of this change was to lower the general average stature of English recruits by only 0.17 inch, of the Scotch by 0.21 inch, and the Irish by 0.25 inch, but in all three nationalities to increase the average weight—the English by 1.3 lbs., the Scotch by 6.7 lbs., and the Irish by 0.8 lb.

21. Although the minimum standard was the same for all the nationalities, the influence of race is indicated by the difference in the average stature of the recruits. The English and Welsh recruits (who were not distinguished from each other) were shorter in stature than the

Irish by 0.30 inch, and the Scotch by 0.44 of an inch.1

22. The measurements of the chest given in Table I. are almost entirely those of Englishmen, and must be studied in connection with the English observations of height and weight; and the same remark applies to the figures relative to strength. The chest-girths were taken by the method adopted in the British army, and the strengths by the springbalance introduced by this Committee, and described in Appendix A.

23. An examination of Table I. shows that an adult Englishman or typical proportions has a stature of 5 feet $7\frac{1}{2}$ inches; a chest-girth of $36\frac{1}{2}$ inches; a weight of 10 stones 10 lbs.; and is able to draw, as in drawing a bow, a weight of $77\frac{1}{2}$ pounds. These are the mean proportions. The averages give greater weight for height; they are:—Height, 5 feet $7\frac{1}{3}$ inches; weight, 11 stones 1 lb.; empty chest-girth, 36·46 inches; and strength, 79·6 lbs. For every variation of an inch in stature above or below the average, 2·301 lbs. weight, ·542 inch chest-girth, and 1·182 lbs. strength must be added or subtracted to keep up the typical proportions. This rule of proportion is, however, only approximately correct, as variations in the stature depend largely on the length of the lower limbs, while the other qualities depend chiefly on the size of the trunk. In ascending the scale of height, therefore, the above figures are probably a little too great, while in the opposite direction they are barely sufficient, but in either case they are sufficiently near for all practical

¹ Further tables relating to recruits are given in Appendix B to this Report.

purposes.1 A further development of this rule as applicable to both

sexes and at all ages will be found in Table XX.

24. Plate IV. shows the relative stature of the four British nationalities, traced from the columns in the table showing the number of men at each height per thousand. The curve of the English very nearly corresponds with that of the average for the whole kingdom. The Scotch curve is above the average, and from its irregularity it is evident that the observations on which it is based are not quite representative of that part of the kingdom. The Welsh curve is below the general average, and in a manner balances the excess of the Scotch, while the Irish curve is somewhat too acute, owing to the comparatively small number of observations on which it is based.

b. Adult Males and Females-Table II.

25. Table II. shows the relative stature, weight, and strength of adult males and females in England, no returns for females having been received from other parts of the kingdom. The average stature of adult males is 67.36 inches (1.712 metres), and of females, 62.65 inches (1.592 metres), showing a difference of 4.71 inches (120 mètres), or nearly 43 inches. The average weight of males is 155.0 lbs. (70.5 kilos.), and that of females 122.8 lbs. (55.8 kilos.), showing an excess of 32.2 lbs. (14.7 kilos.), or about 21 stones on the side of males, the percentage difference of weight being just threefold that of height. The ratio between the stature of men and women in England is as 1 to 0.930, or as 16 to 14.88, the difference being somewhat greater than in Belgium, where, according to Quetelet, the ratio is as 1 to 0.937, or about 16 to 15 (strictly 16 to 14.99). The observations of the strength of females were obtained from pupils in training institutions for schoolmistresses and from shop assistants, and the average is no doubt much lower than if the labouring classes were also represented. The difference of strength is 35 lbs., the females being little more than half as strong as males. In these tables, the age of the attainment of maturity is fixed at 23 years for males, and 20 years for females, the reasons for which will be explained in another part of the Report.

¹ The following measurements show the difference between the height of the body of men in the standing and recumbent positions, and the span of arms measured across the front of the chest. Also the difference between the height of the body in the standing and the sitting positions, showing the relative length of the trunk and of the lower limbs. The English figures are calculated from the American measurements of Dr. Hitchcock, taken in 1882.

| | Age | No. of obs. | | Standing height | Horizontal length | Span of arms | Sitting height | |
|------------------------------------|------|-------------|-----------------------------|--------------------|--------------------------------|--------------------------------------|------------------------------|-----------------------|
| American Amherst College | 21.5 | 327 | mètres | 1·729 68·07 | 1·748 68 82 | 1·787 70·36 | 0·907 35·71 | Length of trunk |
| English Profes- sional class | 21.5 | 364 | mètres inches | 1·746 68·70 | 1.765 69.45 | 1·804 71·01 | ·915 36·04 | and head |
| Difference { | Ame | rican dish | mètres inches inches inches | +·017 +·63 | +·019 +·75 +·019 +·75 | + ·058 + 2·29 + ·058 + 2·31 | 822 32-36 831 32-66 | Length of lower limbs |

The ratio between the total height and the sitting height is 1 to 1.906.

Table II.—Showing the Relative Stature, Weight, and Strength of Adult Males (23-50 years) and Females (20-50 years) of English Origin.

| | | Height | | | | W | eight | | Strength | | | | |
|------|-------------------------------|----------------|----------------|--------------------|------------|-----------------|-----------|--------------------|--------------|------------------|------------------------|--------------|--|
| | Heig | | | nber of vations | | ht with thes | | nber of vations | | ngth, g-power | Number of observations | | |
| | Inches | Mètres | Males | Females | lbs. | Kilos. | Males | Females | lbs. | Kilos. | Males | Females | |
| | 77- 76- | 1·957 1·931 | 1 | _ | 260 250 | 118·2 113·6 | 1 3 | _ | - | - | _ | - | |
| | 75_ | 1.906 | 9 | | 240 | 109.1 | 9 | | | - | | | |
| | 74_ | 1.881 | 16 | _ | 230 | 104.5 | 10 | | 150 | 68.2 | 4 | | |
| | 73- | 1.855 | 48 | _ | 220 | 100.0 | 33 | | 140 | 63.6 | 4 | | |
| | 72- | 1.830 | 117 | - | 210 | 95.5 | 62 | _ | 130 | 59.1 | 2 | | |
| | 71_ | 1.804 | 254 | 1 | 200 | 90.9 | 75 | 1 | 120 | 54.5 | 15 | | |
| | 70_ | 1.779 | 473 | - | 190 | 86.4 | 174 | - | 110 | 50.0 | 18 | - | |
| | 69_ | 1.754 | 753 | - | 180 | 81 8 | 304 | 1 | 100 | 45.5 | 73 | - | |
| Mean | 68_ | 1.728 | 886 | 3 | 170 | 77:3 | 492 | - | 90 | 40.9 | 226 | 1 | |
| Me | -67 66- | -1.702 | | | 160 | 72.7 | 881 | 2 | 80 | 36.4 | 296 | - | |
| | 65- | 1·677 1·653 | 881 740 | 22 24 | 150 | 68.2 | 1075 | 14 | -70 - | -31.8- | | 7 2 | |
| | 64- | 1.626 | 524 | 44 | 140 | 63.6 | 1240 | 7 20 | 60 | 27:3 | 250 | 5 | |
| | 63_ | 1.601 | 320 | 57 | 130 | 59 1 | 694 | 58 | 50 40 | 22.7 | 69 | 25 | |
| | 62_ | 1.575 | 128 | <u></u> | 120 110 | 54.5 | 338 | 101 | 30 | 18·2 13·6 | 15 3 | 101 | |
| | 61_ | 1.550 | 70 | 59 | 100 | 50·0 45·5 | 133 26 | 108 | 20 | 9.1 | 3 | 98 | |
| | 60- | 1.525 | 39 | 37 | 90 | 40 9 | 20 | 53 10 | - 20 | 31 | | 9 | |
| | 59- | 1.499 | 12 | 22 | | | | 10 | | | | - | |
| | 58- | 1.474 | 3 | 17 | | | | | _ | _ | _ | - | |
| | 57- | 1.448 | 1 | 6 | _ | | _ | _ | _ | _ | _ | - | |
| | 56- | 1.423 | - | 3 | _ | _ | _ | | _ | _ | _ | _ | |
| | 55- | 1.398 | - | 2 | - | - | - | - | - | - | - | | |
| - | Total n berofol vations | oser-} | 6194 | 379 | | | 5552 | 368 | - | - | 1497 | 241 | |
| | Aver- Si | | | 62.65 | Aver- f | lbs. | 155.0 | 122.8 | Aver- | Clhe | 79.6 | 11.5 | |
| | age 11 | mètres | 1.712 | 1.592 | age (| | 70.5 | 55.8 | | kilos | 36.2 | 44·5 20·2 | |
| | | nches | 67·50 1·715 | 62·5 1·588 | Mean | lbs. | 150.0 | 120.0 | Mean | ſlbs. | 77.5 | 40.0 | |
| 1 | | 00200 | - 110 | 1 000 | | kilos | 68.2 | 54.6 | Mean | kilos | 35.2 | 18.2 | |

c. Distribution of Adult Males according to Stature, Weight, and Complexion.

Table III., and Plates V.-IX. (Maps Nos. 1 to 5).

26. Table III. exhibits the average stature, weight, and complexion (colour of eyes and hair) of adult males born in the several counties of Great Britain and Wales and in each province of Ireland, arranged in the order of the greatest stature. The Committee is sensible that the number of observations in some of the counties is not sufficient to furnish an average which may be fully relied upon; but the results, as detailed in the remarks upon this summary, show that there is such a consistency between the data and the records of history as to justify a general trust in the conclusions to be drawn from the figures.

TABLE III -- Showing the STATURE, WEIGHT, and COMPLEXION of 8,614 Adult Males (age from 23 to 50) of the Population of the United Kingdom, arranged according to birthplace in Counties in the order of greatest Stature. Illustrated by Maps.

| na- | ass tthith or air | 1 | T | | | - | | | | - | | - | - | - | - | | - | 41. | - | - |
|---|---|----------------------|---|-----------------------|---|-------|--|-----------------|-------|---------------------|--------------------|-------------------------------------|----------|--------------------------|----------------|-------------------------------------|----------------|--------------|---------|---------|
| Other combina- | such as green, light brown eves with light or dark hair | per cent. | 2.0 | 3.1 | 8.0 | 1.6 | 0.7 | 1 | 0.4 | 0.9 | 6-0 | 17 | - 1 | 1 | 3.5 | 79 | 4.0 | 5-9 | 20 00 | 2.9 |
| | Total Dark eyes | per cent, 24-0 | 27.0 | 18.6 | 29.2 | 28.7 | 18.3 | 26.0 | 96.0 | 25.2 | 15.5 | 7-68 | 65.6 | 1 | 33.7 | 27.0 | 21.7 | 29.8 | 2.98 | 87-1 |
| r black | Red and dark red hair | per cent. | J | 3.1 | 3.5 | 1.3 | 2-0 | 1 | 1.0 | 5.2 | 1 | 17 | 1 | 1 | 9-0 | 0.8 | 0.4 | 10 | 25.0 | 9.0 |
| i, hazel, or eves, with | Fair | per cent. | 1 | 1 | 150 | 4:1 | 1 | 1.3 | P-0 | 5.2 | 6.0 | 8.1 | 8.0 | 1 | 1.6 | 6.6 | 8.0 | 1; | 1.1 | 9-0 |
| Brown, hazel, or black eyes, with | Brown, dark brown, and black hair | per cent. 22·0 | 27.0 | 15.5 | 26.0 | 23.3 | 17.6 | 24.7 | 24.6 | 17.5 | 14.6 | 6.86 | 21.4 | 1 | 31.5 | 23.8 | 20-2 | 6.83 | 96-8 | 85-9 |
| | Total Fair eyes | per cent. 75-2 | 72.8 | 78.3 | 70.0 | 71.3 | 81.0 | 74.0 | 78.6 | 72.5 | 83.6 | 66.5 | 77.8 | 1 | 68-1 | 9-99 | 74.8 | 87.8 | 0.19 | 57.5 |
| dark s, with | Golden or red hair | per cent. 4·2 | 3.3 | 1.0 | 7.1 | 5.7 | 9.9 | 8.5 | 8.5 | 7.5 | 9.7 | 2:1 | 4.8 | 1 | 3.1 | 5.6 | | | 0.60 | - |
| Light blue, blue, dark lue, and grey eyes, wit | Black or dark brown hair | per cent. 24-9 | 25.6 | 25.8 | 22.0 | 22.1 | 82.4 | 26.6 | 17.6 | 27.5 | 23.9 | 24-0 | 11:1 | 1 | 19-6 | 20.5 | 23.8 | 20.9 | 25.5 | 26.4 |
| Light blue, blue, dark blue, and grey eyes, with | Very fair, light brown, or brown hair | per cent. 46·1 | 48.4 | 51.5 | 40.9 | 58.4 | 43.0 | 44.2 | 52.8 | 37.5 | 38.0 | 40.1 | 62.4 | 1 | 40-4 | 40.5 | 43.2 | 41.4 | 41.7 | 26-4 |
| Ratio, 1bs. | t re t | 2.406 | 2.465 | 2.551 | 2.501 | 2.579 | 2.356 | 2.459 | 2.219 | 2.464 | 2.040 | 2.488 | 2.595 | 2.490 | 2.301 | 2.877 | 2.858 | 0.000 | 2.858 | 2.809 |
| age ht, ling | 80 | 75-1 | 78.6 | 81.2 | 78.5 | 78-9 | 78.4 | 75.5 | 8.89 | 76.4 | 277.8 | 75.4 | 8.02 | 8.92 | 2.02 | 74.5 | 78.8 | 1.77 | 72.7 | 21.8 |
| Average weight, including | clothes Ibs. Kii | 165-3 | 172.9 | 1/8.6 | 172.9 | 162.7 | 9-191 | 166.3 | 151-4 | 168-1 | 171-8 | 165-9 | 155.9 | 169-1 | 155.0 | 164-0 | 161.4 | 0.001 | 160-1 | 156-9 |
| ht but | s lètres | 1-746 | 1-782 | F.769 | 1-757 | 1-745 | 1-741 | 1-740 | 1.734 | 1.784 | 1-728 | 1.728 | 1.726 | 1.726 | 1.712 | 1.754 | 1.793 | 1.700 | 1.728 | 1-727 |
| Average height without | shoes Inches Mètres | 68-71 | | 09-69 | 69-13 | 68.63 | 68.29 | 68-45 | 68-21 | 68-22 | 68-04 | 68-04 | 67-92 | 67-91 | 67.36 | 00-69 | 68-93 | 68.97 | 00-89 | 67-95 |
| | Num- ber of obs. | 1369 | | 09 | 97 | 85 | 113 | 88 | 189 | 89 | 109 | 109 | 108 | 77 | 6194 | 231 | 291 | 212 | 128 | 133 |
| | Counties | SCOTLAND, Total . | Kirkcudbright, Ayrshire, and Wigton Edinburgh. Linlithcow. Hadding-1 | ton, and Berwickshire | Perth, Stirling, and Dumbarton . Sutherland, Ross, Cromarty, and Skye | | Dumfries, Koxburgh, Selkirk, and Peebles | Inverness-shire | gow) | Caithness Frankling | Islav and Colonsav | Aberdeen, Banff, Elgin, and Nairn . | Shetland | Hebrides-Harris and Uist | ENGLAND, Total | Yorkshire, North and East Ridings . | Northumberland | Lincolnahira | Norfolk | Nasex . |

| 100 | | | | The state of the s | |
|---|---|---|---|--|---|
| 3-9 1-1 6-9 | 1.8 4.8 | 62666144774 | 1.0 9.3 9.3 1.6 8.1 8.1 8.1 8.1 | 8.4 5.1 1.5 8.0 | 9.6 9.6 9.6 9.6 9.6 9.6 9.6 9.6 9.6 9.6 |
| 22.5 28.8 38.8 | 38.5 39.6 39.0 39.0 | 246.2 36.6 48.6 48.6 41.3 32.5 32.5 32.5 | 31.3 34.8 45.4 35.0 30.9 31.1 28.8 40.1 34.4 24.6 | 31·6 41·1 33·1 33·1 34·8 28·4 | 14:5 24:8 23:4 32:2 |
| 1123 | 311289 | 889 1 649 | 0.2 2.0 0.7 0.7 0.8 1.0 0.4 0.7 0.7 | 11 1 1 1 1 1 1 1 1 1 | 1-2 |
| 1 8 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 | 14188 | 2.8 9.3 1.6 1.6 1.6 | 1.5 0.5 0.8 1.0 1.0 1.4 0.8 0.8 | 8.7 8.7 6.4 6.4 | 1881 |
| 81-4 19-1 82-8 | 86.1 88.5 82.9 87.4 | 229 244.8 257.7 25.6 28.8 28.8 32.8 | 29.6 29.1 30.2 30.2 30.2 30.2 30.2 30.2 30.2 30.3 30.3 | 26-8 41-1 27-9 83-8 21-2 | 18:3 24:0 22:2 28:9 |
| 54.5 54.8 | 56.7 56.7 57.1 64.0 57.1 | 59.9 59.5 59.5 57.8 57.8 65.8 59.9 | 67.7 55.9 68.3 66.2 67.0 67.0 69.9 61.2 67.3 | 64-3 60-0 53-8 66-9 63-7 63-6 | 84·3 65·6 74·1 64·5 |
| 3.54.4 | 851185 | 1.6 1.6 1.6 1.6 1.6 1.6 1.6 1.6 1.6 1.6 | 6.4 2.8 2.8 1.0 1.0 1.0 2.2 2.2 2.0 2.0 | 8·7 11·6 5·1 13·0 1·4 8·2 | 4.8 |
| 27.0 22.8 22.8 | 16.9 16.4 16.4 15.7 | 16.3 16.3 18.5 118.5 17.7 13.7 23.0 | 19.2 17.2 111.2 21.7 18.0 19.8 22.3 26.2 15.7 14.8 17.4 | 21.2 18.9 12.8 23.0 20.3 20.3 | 24·1 24·0 24·7 11·5 |
| 44.9 27-2 18.5 | 88.6 41.9 45.8 40.2 | 24.4 24.4 24.4 25.5 25.6 25.6 25.6 25.6 25.8 | 42.1 386.3 39.9 40.8 40.8 40.8 40.8 88.9 29.0 | 29.5 35.9 30.9 42.0 39.6 | 59.0 36.8 49.4 50.3 |
| 2-366 2-323 2-366 | 2:308 2:323 2:247 2:247 2:301 | 2.284 2.821 2.821 2.821 2.824 2.825 2.839 2.889 | 2.278 2.285 2.285 2.275 2.204 2.269 2.252 2.249 2.249 2.361 | 2.875 2.481 2.840 2.891 2.839 2.839 | 2-253 2-253 2-308 2-181 |
| | | 72:5 71:0 71:6 71:8 71:8 71:8 | 69.5 70.6 69.0 68.6 66.6 70.0 71.9 67.9 67.8 | 71.9 78.1 70.9 70.6 70.6 | 70-2 69-5 71-8 67-9 |
| | | | 152.6 152.9 155.3 155.3 150.9 146.5 158.2 149.4 148.3 149.1 | 158.3 162.5 155.9 155.4 155.4 | 154-9 158-0 157-9 149-4 |
| 1.724 | | | 1.702 1.696 1.696 1.696 1.689 1.688 1.685 1.685 1.685 1.685 | 1.694 1.703 1.699 1.692 1.689 1.689 | 1.746 1.741 1.789 1.784 |
| 67.71 67.71 67.71 | | 67.29 67.26 67.26 67.26 67.12 67.12 67.07 67.08 | 66.98 66.75 66.74 66.47 66.47 66.47 66.45 66.84 66.83 66.83 66.83 | 66-66 66-85 66-58 66-58 66-58 66-47 | 68-73 68-52 68-41 68-21 |
| 164 | 228 228 243 243 166 | 136 136 147 165 165 173 173 173 173 173 173 173 173 173 173 | 458 259 122 72 270 28 141 60 836 447 160 | 735 82 82 82 389 60 1122 346 | 85 55 148 148 |
| Derbyshire Suffolk | Berkshire Kent Lancashire Hannashire Notting hamshire | land | | Wales. Total . Flint and Denbigh . Garnarvon, Anglesea, Merioneth, and Montgomery Cardigan Brecon and Radnor . Glamorgan, Caermarthen, and Pem- broke . IRELAND. Total . | |

27. To save much detailed description, the Committee has thought it desirable to illustrate Table III. by a series of shaded maps (Plates V.-IX.), which present at once to the eye the relative distribution of the stature, weight, and complexion of the adult male population in the several counties of Great Britain and in each province of Ireland.

Map No. 1 shows the distribution of the average stature (without shoes) of adult males, in degrees of half an inch each from 66 to 70 inches.

The darkest shade represents the shortest stature.

Map No. 2 shows the distribution of the average weight (including the clothes) of adult males, in degrees of five pounds from 145 pounds to 180 pounds. The darkest shade represents the lightest weight.


Map No. 3 shows the distribution of adult males with fair complexion, i.e. blue and grey eyes with fair, light-brown, brown, and light-red hair. The darkest shade represents the lowest percentage of fair complexion.

Map No. 4 shows the distribution of adult males with dark complexion, i.e. brown and black eyes, with brown, dark brown, dark red, and black hair. The darkest shade represents the highest percentage of dark complexion, or its greatest prevalence.


dark complexion, or its greatest prevalence.

Map No. 5 shows the distribution of adult males with mixed complexion, i.e. blue and grey eyes with dark brown and black hair. The darkest shade represents the highest percentage, or the greatest prevalence of this complexion.


- 28. As the observations were necessarily made on a limited number of individuals, and as doubts may exist as to whether the results can be accepted as representing the whole of the male population at the ages specified, the counties having similar statures have been grouped together, and the male population for each group ascertained from the Census returns of 1881. The average stature worked out from these figures is 67.58 inches, while that obtained from the actual observations on 8,585 individuals, given in Table I., is 67.66 inches, the difference between the two being only 0.08 of an inch. Table IV. shows the grouping of the counties, having the same stature according to the Committee's returns, and the total male population of each group at the ages from 25 to 55 years.
- ¹ These returns for England and Scotland are not yet published, and the Committee is indebted to the courtesy of the Registrars-General of those portions of the kingdom for manuscript copies of the returns. The ages of the men on whom the observations were made are not exactly the same as those obtained from the Census office, but they are sufficiently near for any practical purpose. The measurements were made on men from 23 to 51 years of age, while the Census returns are those of men from 25 to 55 years, but the four years above 51 will about compensate for the two years wanting below 25 years both in numbers and stature, in consequence of losses by death. Both periods correspond with the best portion of men's lives, at least as far as stature is concerned.



















TABLE IV.—Showing the Number of Adult Males of the Ages above 25 and under 55 years for each group of counties possessing the same Average Stature, and the ratio per 1,000. From the Census returns of 1881.

| Observed average stature without shoes in inches | Counties of the United Kingdom | Adult male population age 25-55 years | Per 1,000 |
|---|---|--|-----------|
| 69½ and upwards | Kirkcudbright, Ayr, Wigton; Edinburgh, Linlithgow, Haddington, Berwickshire. | 125,103 | 22.2 |
| 69 to 69½ | Sutherland, Ross and Cromarty, Skye, Perth, Stirling, Dumbarton, Fife, Kinross, Clackmannan; North and East Ridings of Yorkshire. | 167,914 | 30.0 |
| 68½ to 69 | Argyle, Bute, Arran, Dumfries, Rox- burgh, Selkirk, Peebles; Northum- berland; Connaught, Munster. (Caithness, Inverness, Aberdeen, Banff,) | 459,055 | 81.7 |
| 68 to 68½ | Elgin, Nairn, Forfar, Kincardine; Lanark, Renfrew; Cumberland, Westmoreland; Lincoln, Norfolk; Ulster, Leinster. | 974,177 | 173.4 |
| 67½ to 68 | Shetland, Western Hebrides; Durham, Lancashire, Derby, Stafford; Suf- folk, Essex, Kent; Berkshire; Corn- wall. | 1,326,292 | 236.0 |
| 67 to 67½ | Northampton, Bedford; Warwick, Worcester; Flint, Denbigh; Sussex, Hampshire, Dorset, Devon. | 688,465 | 122.6 |
| _ | London (66.92 inches). | 667,118 | 118.7 |
| 66½ to 67 | West Riding of Yorkshire, Chester; Carnarvon, Anglesea, Merioneth, Montgomery, Cardigan, Brecon, Radnor; Cambridge, Huntingdon; Buckinghamshire, Oxfordshire. | 636,769 | 113-3 |
| 66 to 66½ | Hertford, Middlesex (ex. metrop.); Surrey (ex. metrop.); Shropshire, Hereford, Monmouth, Gloucester, Wiltshire, Somerset; Glamorgan, Caermarthen, Pembroke. | 573,774 | 102·1 |
| | | 5,618,677 | 1000 |

Stature \times Population Total male population $= \begin{cases} 67.58 \text{ inches, average stature of adult males (25-55 years of age) of the United Kingdom.} \end{cases}$

29. Ethnology.—The variations in stature, weight, and complexion shown to exist in different districts of the British Isles by the maps, appear to be chiefly due to difference of racial origin, and this influence predominates over all others. 'We have reason to believe, from historical and antiquarian researches, that the ancient Caledonii, the Belgæ and Cimbri, and the Saxons and Frisians, as well as the Danes and Normans, were all people of great stature. On the other hand, the prehistoric (neolithic) race or races in Britain appear to have been of low or moderate stature. Accordingly the higher statures are found in the Pictish or

Cimbro-British districts of Galloway; in the Anglo-Danish ones of North and East Yorkshire, Westmoreland and Lincolnshire, and in Cumberland, whose people are ethnologically intermediate between the two. Lothian and Berwickshire are mainly Anglian, while the Perthshire Highlanders are the most clearly identified as the descendants of the Caledonii. The high position of Norfolk in the list is due to a large admixture of Danish blood on the coast. There is a fringe of moderately high stature all round the coast from Norfolk to Cornwall, while the inland people, retaining more of the ancient British blood, yield lower averages. Middlesex and Hertfordshire, which stand very low, were later and less perfectly colonised by the Anglo-Saxon than the surrounding counties, and nearly the same may be said of the counties around the Severn estuary and the Welsh border. Cornwall stands higher than the surrounding counties, and this is probably due to its having become the refuge of the military class of Southern Britain, in the main of Belgic origin. Flint and Denbigh owe their superiority to the other Welsh counties to the immigration of the Cumbrian and Strathclyde Britons.' —Dr. Beddoe.

30. According to the Committee's returns, the western provinces of Ireland possess a high stature, similar to the Scotch Highlands, with which they may have a common racial origin, while the lower stature of the eastern provinces is probably traceable to the comparatively recent Scotch and English immigrations. The Irish returns are, however, too few to be relied on (although the closeness of the averages for all the provinces would suggest the absence of any errors of observation), and any conclusions drawn from them must be received with great reserve until they are confirmed by more extended inquiries. In some of the returns the county origin and birthplace was not recorded, which accounts for the difference between the totals for the whole of Ireland and those living in each province.

31. The racial elements of the British population are best demonstrated by separating a few of the counties where there has been the least admixture of foreign blood, and comparing these together, thus:—

| Race | District | Stature | Weight |
|--|--|----------------------------------|----------------------------------|
| Early British . Saxon Anglian Scandinavian . | Cardigan, Radnor, and Brecon Sussex, Berkshire, and Oxfordshire Lothians, Northumberland, and Norfolk . Shetland, Caithness, North and East York- shire, and Lincolnshire. | 66·59 67·22 68·73 68·32 | 169·3 155·8 166·7 162·7 |

32. Geographical distribution.—The inhabitants of the more elevated districts possess a greater stature than those of alluvial plains. The counties forming the river valleys of the Severn and Wye, the Thames, the Dee and Mersey, the Clyde, the Trent, and the fen district of Cambridge and Huntingdon, show a lower stature than the surrounding counties inhabited by persons of a similar racial origin.

33. With respect to latitude and climate, the inhabitants of the northern and colder districts possess greater stature than those of the southern and warmer parts of the island; those of the north-eastern and drier regions are taller than those of the south-western and damper climates. A similar disposition of stature has been found to exist in France and Italy, the

inhabitants of both these countries being taller in the northern than in the southern provinces. The same rule applies to the whole of the countries of Europe, in their relation to each other, as will be seen in Table IV., constructed to show the position held by the inhabitants of the British Isles relative to the stature of other European countries. The Committee regrets that it has not been able to obtain any information on this subject direct from the European countries (except some referring to conscripts, which were not suitable for their purpose), and has been obliged to avail itself of the observations made in the United States of America on emigrants from European States. In reading this table it must be borne in mind that the statistics referring to the United Kingdom, collected by the Committee, and to the native-born population of the United States, refer to men of all classes; while those collected by the military authorities of 1863-4 in the United States, referring to Canada and the other American countries, and to those of all Europe, refer to emigrants, who belong almost entirely to the labouring classes. The close accord between the average stature of the United Kingdom (67.66 inches) and that of the native white population of the United States (67.67 inches) is accounted for in this way; and, on the other hand, the marked differences between the statures of the Scotch (68.71), Irish (67.90), English (67.36), and Welsh (66.66 inches), as given by the Committee and those given by the United States Government (67.07, 66.74, 66.58, and 66.42 respectively) is explained. Some American writers on the subject have overlooked this important distinction, and, studying only the statistics obtained in their own country, have concluded that the Anglo-Saxon race is of greater stature in America than in Great Britain. In the Report of the Committee for 1879 Mr. Roberts has given a paper, illustrated by a series of diagrams and statistical tables, of English and Americans, showing the close similarity which exists between the stature and weight of the two branches of our race, both in children and adults; and the more extended observations of the Committee appear to confirm

34. Occupation and sanitary surroundings.—The various industries of this country are not often so defined by the county boundaries as to show their effects on the physical development. It is probable, however, that the low stature in the West Riding of Yorkshire is due to the large manufacturing town population included in the returns, and the relatively low stature of Durham to the large mining population. Lancashire and Stafford, which contain similar industries to those of the West Riding and Durham, do not show any falling off in stature, and it is probable that a large number of returns received from Sheffield have unfairly lowered the West Riding. The very low position, lower than can be accounted for by their racial origin, taken by the home counties-Hertford, Middlesex, and Surrey-is no doubt due to their proximity to London; the more vigorous men are attracted to the town by high wages, and the more feeble overflow into the surrounding districts. The counties which fringe the sea-coast possess a higher stature than those adjoining them but lying further inland. This may be due to race, as has already been suggested; but it may also be due to the more healthy situation or the fishing occupation. The lower stature of the river valleys would seem to imply that such situations are not favourable to physical development, especially as some of them were originally settled by the Scandinavian races.

Table V.—Showing the Average Stature of Adult Males in each Division of the United Kingdom, according to the returns collected by the Anthropometric Committee, compared with that of Adult Males of American and European Origin, who were examined for admission into the United States Army in the year 1863-4; the natives of European origin being arranged in the order of their average stature, showing also the medium stature, and the proportions above and below it, with the proportions of the extremes of high and low stature. (See 'Statistics, Medical and Anthropological, U.S. Army, 1875.')

| Medical and Anthropologic | ai, U.D. 1 | rimy, | 1010. | | | | |
|---|---------------------|----------------------------|-----------------|---------------------------------|-----------------|--|-----------------|
| | vations | ature. | port | entage p ion of to number | | Extre Percer proport total no | itage ion of |
| Countries | No. of observations | Average stature. Inches | Under 65 inches | 65 to 69 inches | Above 69 inches | Under 61 inches | Above 73 inches |
| Observations of Anthropo- metric Committee:— | | | | | - | | |
| Scotland Ireland | 1,304 346 | 68·71 67·90 | 5·6 6·7 | 50·2 65·3 | 44·2 28·0 | 0·19 0·32 | 2·13 0·00 |
| England | 6,194 | 67.36 | 17.8 | 55.5 | 26.7 | 0.93 | 0.43 |
| Wales | 741 | 66.66 | 22.8 | 62.0 | 15.2 | _ | - |
| Total, United Kingdom | 8,585 | 67.66 | 16.1 | 55.7 | 28.2 | - | _ |
| Observations on Conscripts in U.S. America:— | | Printles Con lo | | | Park de | | |
| United States. White, native born . | 315,620 | 67.67 | 15.3 | 54.1 | 30-6 | 0.53 | 2.02 |
| Coloured, of all degrees | 25,828 | 66.63 | 29.6 | 51.9 | 18.5 | 1.79 | 1.00 |
| Indians, N.A. tribes . | 121 | 67.93 | 14.2 | 52.0 | 33.8 | - | 0.08 |
| Immigrants from— | | | | | | | |
| Canada (chiefly French) | . 21,645 | 67.01 | 218 | 56.3 | 21.9 | 0.74 | 1.01 |
| Mexico | . 91 | 66-11 | 25.2 | 51.7 | 13.1 | 3.29 | 1.09 |
| South America | . 79 | 65.90 | 41.7 | 40·4 56·4 | 17.9 | 2.13 | 0.34 |
| West Indies | . 580 | 66.31 | 28.9 | 90.4 | LEI | 0 00 | 0.51 |
| Europe. Norway | 2,290 | 67.47 | 16.6 | 57.0 | 26.4 | 0.74 | 1.31 |
| Scotland | 3,476 | 67.07 | 20.4 | 58.3 | 21.3 | 0.46 | 1.03 |
| Sweden | . 1,190 | 66.90 | 21.3 | 59.5 | 19.2 | 0.42 | 0.76 |
| Ireland | . 30,557 | 66.74 | 23.2 | 60.1 | 16.7 | 0.70 | 0.49 |
| Denmark | . 383 | 66.65 | 25.1 | 57.7 | 17:2 | 0.78 | 0.26 |
| Holland | . 989 | 66.64 | 26.6 | 56.3 | 17:1 | 1.31 | 0.50 |
| England | . 16,196 | 66.58 | 25.9 | 58·3 58·4 | 15.8 | 3.37 | 1.12 |
| Hungary | . 89 | 66.58 66.54 | 22·5 27·0 | 57.0 | 16.0 | 1.31 | 0.51 |
| Germany | . 54,944 | 66.42 | 29.3 | 53.6 | 17.1 | 0.82 | 0.63 |
| Russia | 1,104 | 66.39 | 29.6 | 51.0 | 16.4 | 3.28 | 0.82 |
| Switzerland | 1.302 | 66.38 | 29.5 | 55 7 | 14.8 | 1.61 | 0.44 |
| France | . 3,243 | 66.28 | 30.0 | 56.5 | 13.5 | 1.85 | 0.57 |
| Poland | . 171 | 66.21 | 32.1 | 56.7 | 11.2 | 1.75 | 1.17 |
| Italy | . 339 | 66.00 | 37.8 | 48.9 | 13.3 | 2.06 | 0.29 |
| Spain | . 148 | 65.64 | 43.3 | 49·3 56·8 | 7·4 3·7 | 3.70 | _ |
| Portugal | . 81 | 65.43 | 39.5 | 00.0 | 01 | 1 010 | |

d. British compared with other Races and Nationalities.

35. Considering the large number of different races included in the British Empire, and the political and commercial relations of its people with nearly every other country, the Committee think it will be interesting and useful to give a table showing the average stature of the different races and nationalities of the world, as far as it has been able to ascertain them from published records. The list is very imperfect, and it is probable that many of the measurements need revision by more extensive observation. No nation is so favourably situated for revising and completing the list as our own; and the Committee hope that the table will be instrumental in promoting further observations of the kind, especially by medical officers in the Navy and Army, and others practising in our numerous colonies and dependencies. It is interesting to find that, with the exception of a few imperfectly-observed South Sea Islanders, and whose actual numbers, if the measurements are correct, are very few, the English professional classes head the long list, and that the Anglo-Saxon race takes the chief place in it among the civilised communities, although it is possible it might stand second to the Scandinavian countries if a fair sample of their population were obtained.

Table VI.—Showing the Stature of Adult Males of the British Isles relative to that of other Races and Nationalities, arranged in the order of greatest Stature.

| Race or Nationality | Authority | Mètres | Ft. in. |
|--|--|---|--|
| Polynesians Samoa | Lapeyrouse Garnot, Beechey Porter, Cook, &c. Various Wilkes, Novara Lesson, Rollin | 1.762 | 5- 9.33 |
| English professional class | Anthropometric Com. | 1.757 | 5- 9.14 |
| Patagonians $\begin{cases} 1.778 \\ 1.730 \end{cases}$ | Musters | 1.754 | 5- 9 00 |
| Angamis of the Naga Hills Negroes of the Congo Scotch, all classes (recruits, 5 ft. 8·03) Amakosa Kaffirs, South Africa Iroquois Indians Todas of the Nilghiries Negroes of Calabar North American Indians Irish, all classes (recruits, 5 ft. 8·04) United States (whites, all classes) English, all classes (recruits, 5 ft. 7·71) | Woodthorp Topinard Anthropometric Com. Sir A. Smith Gold Marshall Topinard Baxter Anthropometric Com. Baxter Anthropometric Com. | 1·754 1·752 1·746 1·741 1·735 1·727 1·727 1·726 1·725 1·719 1·719 | 5- 9.00 5- 8.95 5- 8.71 5- 8.50 5- 8.28 5- 7.95 5- 7.95 5- 7.93 5- 7.67 5- 7.66 |
| Norwegians { immigrants to U.S. 1.717 Zulus | Beddoe | 1·719 1·707 1·705 | 5 - 7.66 5 - 7.19 5 - 7.08 |
| U.S. America Tajiks of Ferghana and Samarkand Swedes, immigrants to U.S. America Chipeway Indians Kabyles, large race | | 1.705 | 5- 7·01 5- 7·10 5- 6·90 5- 6·90 5- 6·85 |

TABLE VI. (continued).

| Race or Nationality | Authority | Mètres | Ft. in. |
|---|-------------------------|----------------|--------------------|
| Welsh, all classes | Anthropometric Com. | 1.695 | 5- 6.66 |
| Welsh, all classes | Baxter | 1.694 | 5- 6-65 |
| Dutch | Baxter | 1.693 | 5- 6.62 |
| Dutch ,, American negroes of all degrees of | | | |
| | Baxter | 1.693 | 5- 6.62 |
| English immigrants to U.S. America | Baxter | 1.692 | 5- 6.58 |
| | Baxter | 1.692 | 5- 6.58 |
| Hungarians , , , | Anthropometric Com. | 1.692 | 5- 6.57 |
| Germans, immigrants to U.S. America . | Baxter | 1.691 | 5- 6.54 |
| Swiss of Geneva | Baxter | 1.688 | 5- 6.43 |
| Swiss immigrants to U.S. America . | Baxter | 1.687 | 5- 6.38 |
| Russians ,, , , | Baytor | 1.687 | 5- 6.38 |
| Belgians | Quetelet | 1.687 | 5- 6.38 |
| French immigrants to U.S. America . | Baxter | 1.683 | 5- 6-23 |
| Poles " . | Baxter | 1.682 | 5- 6.20 |
| Poles ,, | De Quatrefages | 1.681 | 5- 6:14 |
| Germans | Novara | 1.680 | 5- 6.10 |
| Mexicans | Baxter | 1.680 | 5- 6.10 |
| Mexicans | Topinard | 1.680 | 5- 6.10 |
| Arabs | Various Ujfalvy | 1.679 | 5- 6.08 |
| Usbeks of Ferghana and Samarkand . | Ujfalvy | 1.679 | 5- 6.08 |
| Javanese | Novara | 1.679 | 5- 6.08 |
| Russians | Shultz | 1.678 | 5- 6.04 |
| Italians, immigrants to U.S. America . | Baxter | 1.677 | 5- 6.00 |
| South Americans ,, , , | Baxter | 1.675 | 5- 5.90 |
| Australian Aborigines | Various | 1.669 | 5- 5.68 |
| Austrian Sclaves | Novara | 1.669 | 5- 5.68 |
| Galchas, Iranian Mountaineers | Ujfalvy | 1.668 | 5- 5.66 |
| Spaniards, immigrants to U.S. America. | Baxter | 1.668 | 5- 5.66 |
| Berbers of Algeria | Topinard | 1.666 | 5- 5.62 |
| Portuguese immigrants to U.S. America | Baxter | 1 663 | 5- 5.43 |
| Ainos | Rosky | 1.660 | 5- 5.33 |
| Portuguese immigrants to U.S. America Ainos Austrian Germans French working classes Esquimaux of North America Hungarians (military statistics) | Novara | 1.658 | 5- 5·27 5- 5·24 |
| French working classes | De Quatrerages . | 1.657 1.654 | 5- 5-10 |
| Esquimaux of North America | Various | 1.652 | 5- 5.04 |
| Hungarians (military statistics) | Scheiber and Beddoe. | 1.650 | 5- 4.93 |
| Caucasians | DHOLDO | 1.646 | |
| New Guinea, various tribes | Various | 1.645 | 5- 4.76 |
| Hindoos | Shortt | | 5- 4.68 |
| | Novara | 1.640 | |
| Ruthenians | Showtt | 1.639 | 5- 4.50 |
| Dravidians | Shortt | 1.638 | |
| Cingalese | Vangang | 1.631 | 5- 4.37 |
| Austrian Roumanians | Novara | 1.630 | 5- 4:17 |
| Chinese | An. di Statist., 1879 . | | 5- 4.00 |
| Italians (conscripts, 1.620) | Novara | 1.625 | 5- 3.98 |
| Fuegans | Majer and Kopernicki | 1.623 | 5- 3.88 |
| Fuegans Polish Jews Poles Finns (Beddoe, 5 ft. 5.81) Papuans | Majer and Kopernicki | 1.622 | 5- 3.87 |
| Poles | | 1-617 | 5- 3.60 |
| Papuans | Various . | 1.606 | 5- 3.20 |
| Japanese | Mrs. Ayrton | 1.604 | 5- 3.11 |
| Aymaras Indians, Peru | Forbes | 1.601 | 5- 3.00 |
| Pernyians | D'Orbigny | 1.600 | 5- 3 00 |
| Peruvians | Various | 1.593 | 5- 270 |
| Journal Committee 1 | Raffles, Crawfurd, &c. | 1.583 | 5- 2.34 |
| Malays | Bailey | A 44 A 45 | 5- 0.42 |

TABLE VI. (continued).

| Race or Nationality | Authority | Mètres | Ft. in. |
|------------------------------------|----------------|--|--|
| Lapps | Horch | 1·500 1·492 1·482 1·448 1·436 1·341 | 4- 11·2 4- 10·7 4- 10·3 4- 9·00 4- 8·53 4- 4·78 |
| Difference between the tallest and | shortest races | •421 | 1- 4.55 |
| Average stature of man according | to the above | 1.658 | 5- 5:25 |

Special Subjects of Inquiry.

- 36. In the sheet of instructions issued by the Committee observations were asked for to illustrate the physical differences of:
 - a. Persons engaged in different occupations.

b. Persons bred and living in towns, or country.

c. Natives of parts of the British Isles differing ethnologically, geologically, or in climate.

d. Boys and men whose intellect and industry are above or below the

average.

e. The general characteristics of men noted for athletic power.

f. The rate of growth in persons of both sexes bred in town and

country, and engaged in different occupations.

The following table shows some of the extreme variations in stature which occur, and which are associated with different occupations and conditions of life, illustrative of the above subjects of inquiry.

Table VII.—Showing the Stature and Weight of Adult Males (age 23-50 years) under different conditions of life.

| and Western March March 1981 | Number | Ft | in. | lbs. |
|--|--------|----|---|-------|
| Scotch Agricultural Population, Galloway | 75 | 5 | 10.5 | 173.6 |
| Metropolitan Police | 192 | 5 | 10.1 | 185.7 |
| renows of the Royal Society. | 98 | 5 | 9.76 | |
| Yorkshire Fishermen, Flambro' | 68 | 5 | 8.71 | 166.8 |
| Athletes (running, jumping, and walking) | 89 | 5 | | 143.7 |
| ocotch Lead-miners, Wenlockhead | 92 | 5 | 8.43 | 163.9 |
| London Fire Brigade | 69 | 5 | 7.40 | 160.8 |
| Juliam Coal-miners . | 51 | 5 | | 152.4 |
| dinburgh and Glasgow Town Population | 32 | 5 | 6.35 | 137.2 |
| Velsh Lead-miners, Cardigan. | 328 | 5 | CONTRACTOR OF THE PARTY OF THE | 155.2 |
| Sheffield Town Population | 100 | 5 | | 142.5 |
| oristor fown Population . | 300 | 5 | | 142.4 |
| unatics, General Population | 1,409 | 5 | | 147.9 |
| Priminals, General Population | 2,315 | 5 | | 140.4 |
| Hertfordshire Labourers | 174 | 5 | | 145.0 |
| diots and Imbeciles | 19 | 5 | 4.87 | 123.0 |

37. The influence of town life and town occupations on the physique of the population in districts in which the race differs little, and the climatic

conditions are the same, is seen by comparing the agricultural population of Ayrshire with that of Glasgow and Edinburgh, where the average difference in stature amounts to 4·15 inches, and in weight to 36·4 lbs., in favour of the country folk. A similar, though not so great a difference, exists in Yorkshire, where the fishermen of Flamborough exceed the artisans of Sheffield in stature by 2·91 inches, and in weight by 24·3 lbs. On the other hand, the population of London exceeds that of the adjoining county of Hertfordshire in stature by 1·57 inches, and in weight by 7·9 lbs. Quetelet observed the same condition in Belgium, where the towns showed a higher stature than the country districts; and he concluded that the greater ease and better food attainable in towns were more favourable to physical development than the hard manual labour and poor fare of the agricultural districts. It is probable that Quetelet compared different classes together, or that the towns in Belgium hold an exceptional position, like London to the adjoining districts in England.

38. As an example of the predominance of race over occupation, the stature and weight of the Scotch lead-miners of Wenlockhead, and the Welsh lead-miners of Cardiganshire, are given in the table. The occupation of lead-mining in both districts is in a great measure hereditary, and has probably been followed under similar conditions in Scotland and Wales for many generations, yet the Scotch exceed the Welsh lead-miners in stature by 2·13 inches, and in weight by 8·7 lbs. The stature and weight of the Durham coal-miners, and of the town populations of Glasgow, Sheffield, and Bristol, are given in this table, as they have been referred to above as influencing the averages of their respective counties, and placing them in an exceptional position as to the racial origin of their inhabitants.

39. One of the objects the Committee has had in view has been 'to ascertain the physical differences of boys and men whose intellect and industry are above or below the average'; but no returns of this kind have been received, except some referring to criminals and lunatics, and those have been introduced here as the most convenient place for their consideration:—

TABLE VIII.—Showing the STATURE and WEIGHT of Adult Male Criminals and Lunatics, compared with that of the General Population.

| | Height Ages | | | | Weight | | | |
|------------------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| Classes | | | | | Ages | | | |
| Crasses | 20 to 25 | 25 to 35 | 35 to 45 | 45 to 55 | 20 to 25 | 25 to 85 | 35 to 45 | 45 to 55 |
| General— | inches | inches | inches | inches | lbs. | lbs, | lbs. | lbs. |
| Average population | 67.5 | 67.9 | 67.9 | 67.9 | 146.2 | 156 | 162 | 163.8 |
| Class 3: country labourers . | 67.2 | 67.5 | 67.5 | 67.8 | 149.5 | 157.4 | 161.2 | 166.4 |
| Class 4: town arti- | 66.5 | 66.6 | 66.9 | 66-6 | 139- | 147.3 | 154.1 | 148.6 |
| Criminals | 65.2 | 65.6 | 65 7 | 65.8 | 136-9 | 140- | 141-4 | 143.4 |
| Lunatics | 65.7 | | | | 147-9 | | | |

40. When compared with the general population, lunatics show a deficiency of stature of 1.96 inches, and of weight 10.3 lbs.; and criminals of 2.06 inches and 17.8 lbs., indicating a deficiency of physical as well as mental stamina in both these unfortunate classes of society. In respect to complexion lunatics show an excess of 5 per cent. of light eyes with dark hair, and criminals of 10 per cent. of dark eyes with dark hair over the general population.

TABLE IX.—Showing the COMPLEXION of Adult Male Criminals and Lunatics, compared with that of the General Population.

| | | No. of observations | Eyes light | | | 1 | Eyes da | Eyes light brown, green, or exceptional, with hair light or dark | Total | |
|-------------------------|---|---------------------|---------------|--------------|--------------|--------------|--------------|--|--------------|-----|
| | | | Hair light | Hair dark | Hair red | Hair dark | Hair fair | Hair red | Eyes l | |
| | | | per cent. | per cent. | per cent. | per cent. | per cent. | per cent. | per cent. | |
| England— General. | | 5,669 | 39.6 | 20.4 | 4.0 | 29.9 | 1.7 | -7 | 3.7 | 100 |
| Criminal | | 2,315 | 40.1 | 13.6 | 1.1 | | 1.000 | | | 100 |
| Lunatic. | | 1,409 | 42.3 | 20.3 | 1.5 | 38.1 | 1.0 | .6 | 5.9 | - |
| Littletic. | | 1,100 | 120 | 20 5 | 1.9 | 31.8 | 1.8 | •4 | 1.9 | _ |
| Total . | | 9,393 | 40.1 | 18.9 | 2.7 | 32.2 | 1.5 | .6 | 4. | _ |
| Wales- | | AP THE | | D DV | 11/11/19 | | | 00000 | | |
| General. | | 704 | 34.4 | 19.9 | 9.8 | 26.4 | 4.7 | 1.9 | 0.8 | 700 |
| Criminal | | 46 | 37. | 17.4 | 0.0 | 45.6 | 4.1 | 1.3 | 3.5 | 100 |
| Lunatic. | | 150 | 34.7 | 27.3 | 3.3 | | - | - | - | - |
| Little City | | 100 | 011 | 213 | 9.9 | 28.7 | 2. | - | 4. | - |
| Total . | | 900 | 34.6 | 21. | 8.2 | 27.8 | 4. | 1. | 3.4 | _ |
| Scotland- | | | | | | | PAIN S | | | |
| General. | | 1,261 | 46.3 | 24.5 | 5.2 | 01.0 | | 100 | | |
| Criminal | : | 194 | 44.3 | 20.1 | 2.6 | 21.2 | .9 | 1. | .9 | 100 |
| Lunatic. | | 342 | 47.4 | 30.7 | | 30. | .5 | 1.5 | 1. | - |
| | | 012 | 11 1 | 30.1 | 1.4 | 17:3 | 1.4 | 1.2 | .6 | - |
| Total. | | 1,797 | 46.3 | 25.2 | 4.2 | 21.4 | 1. | 1.1 | -8 | _ |
| Ireland- | | | | | | | | | | |
| General | | 285 | 49.8 | 10.0 | 0 - | 00 - | | | 1 | |
| Criminal | | 215 | 49.8 | 18.2 | 3.5 | 23.5 | 1.1 | 1.8 | 2.1 | 100 |
| Lunatic | | 29 | 51.7 | 18·6 24·1 | .5 | 28.7 | .2 | .2 | 7. | - |
| | | | 01.1 | 21.1 | 7. | 17.2 | - | - | - | - |
| Total. | | 529 | 47.4 | 19. | 2.5 | 25.3 | .7 | 1.1 | 4. | _ |
| Total United Kingdom | } | 12,619 | 41. | 19.8 | 3.4 | 30.1 | 1.5 | .7 | 3.5 | _ |
| 41 Agan | | 4 | | | | | | - 1 | | |

41. As an example of the relation of high mental to physical qualities the stature of ninety-eight Fellows of the Royal Society is given. Their average stature is slightly above (0.38 inch) that of the professional classes of this country, to which the majority of them belong.

42. As an example of high physical qualities as developed by training, the measurements of eighty-nine professional and amateur athletes are given. Their average stature exceeds that of the general population from which they are drawn by 0.68 inch, while their average weight falls short of that standard by 14.5 lbs. The ratio of weight to stature is, in the athletes, 2.100 lbs., and in the general population 2.323 lbs., for each inch of stature. Thus, a trained athlete whose stature is 5 feet 7 inches should weigh 10 stones, while an untrained man of the same height should weigh 11 stones.

43. The statures of the Metropolitan Police and the London Fire Brigade are given as selected men of the working classes. The former exceed the criminal class, with whom they have to deal, in stature by 4.5 inches, and in weight by 45.3 lbs. The men of the Fire Brigade are selected for their activity, and general fitness to meet sudden and trying demands on their physical and mental energies. The data referring to them may be accepted, therefore, as typical of the best physique which can be obtained for an English army, and of which our army should con-

sist at its best.

Complexion as determined by the Colour of the Eyes and Hair.

44. The difficulty of determining the prevailing complexion of a race, or of the mixed population of a country or a district, by the colour of the hair, as is generally done, and of basing a classification on it, is greater than at first sight appears. Not only do the various shades run imperceptibly into each other, but observers differ in their appreciation of the different shades when viewed under similar conditions, and the prevailing colour of a district determines the relative value of others. Thus a person living among a dark-haired race would consider brown hair as fair, while another person living among a light-haired people would consider it dark, or at any rate not fair in the same sense as the former would. Objections of this kind do not apply to the eyes, as the colour of the iris is due to the anatomical disposition of pigment in front of or behind that structure. In brown and the so-called black eyes a layer of brown pigment covers the front of the iris and hides the deeper structures, and itself determines the colour; while in blue and grey eyes this layer of pigment is wanting, and the colour is due to the dark pigment (the choroid) situated behind the iris, the blue colour in various degrees resulting from the greater translucency of a thin, and the grey from a thick membrane. The marriage, moreover, of fair and dark persons often produces an intermediate shade in the colour of the hair in the children, but only occasionally produces an intermediate change in the colour of the eyes, the rule being that they are blue or brown like one of the parents. The cross between the blue and brown eye should properly be called green (the deeper blue showing through an imperfect layer of yellow brown pigment), but from popular prejudice to this term, eyes of this mixed colour are generally recorded as brown grey, light brown or light hazel.1

45. For these reasons the classification adopted in this Report is based on the colour of the eyes, and with the object of more clearly defining the two prevailing shades of complexion in this country, namely the 'fair' as characterised by light eyes and light hair, and the 'dark' by dark eyes

¹ See the Report for 1880, p. 134, for a further discussion of this subject.

and dark hair, the mixed or neutral eyes are eliminated, and the dark hair is separated from the former, and the light hair from the latter class. The combinations of blue eyes and light red hair, and of brown eyes and dark red hair, are given in separate columns, but the result is not satisfactory, as many cases of light red have doubtless been returned

as fair hair, and of dark red as dark brown hair.

46. In the instructions issued by the Committee observers were requested to return the colours of eyes as grey, light blue, blue, dark blue, light brown, brown, dark brown, green, and black; and the colour of the hair as very fair, fair, golden, red, red brown, light brown, brown, dark brown, black brown, and black, and some chromo-lithographic sheets as tests 1 for the colour of the hair were at first issued; but the system was found to be too complicated for ordinary observers to follow, and they were left to record the colours of both hair and eyes according to the popular meaning of the above terms. An examination of the returns shows that in many cases wide limits have been given to such words as fair, golden, and brown at one end of the scale, and of dark brown and black at the other, which has necessitated the concentration of the data to eliminate errors of observation, and what may be called the 'personal equation' of the colour-sense in different observers. In the Report of the Committee for 1880 a table is given of the colour of eyes and hair according to the above scale, of boys and men of the professional classes from ten to fifty years of age, but, apart from its including too wide a range of ages, it is not so well adapted for showing the relative prevalence of complexions as the one now given.

47. The following grouping of the counties according to the prevalence of fair complexion, or, what is the same thing, according to the degree of nigrescence, shows that certain large districts—much larger than the county boundaries—are occupied by inhabitants of similar racial origin, or who have been subject to conditions of life which have reduced them to similar shades of complexion. The division of the percentages into five degrees is, of course, quite arbitrary, and sometimes two counties, only divided from each other by a decimal, and belonging therefore to the same group, may be represented by a different number. The exact per-

centages are given in Table III.

48. In this classification the men with dark eyes and light hair are combined with those having neutral eyes (green) and light or dark hair, because they are few in number, and because this peculiar complexion is probably due to crossing of the light and dark stocks, and the persistence of one feature of the parent in the eyes and of the other in the hair. The fact that men with dark eyes and light hair are more frequently found in the south-western counties of England, where the light and dark races meet and overlap each other, supports this view of their mixed origin. This complexion, moreover, is common in childhood, but disappears as age advances. According to Table XI. it diminishes in males from 13 per cent., during the first five years of life to 1 per cent., at forty-five years of age, and in females from 16.4 per cent. to 2 per cent. during the same period.

¹ These test-sheets proved not to be well suited for the purpose for which they were intended. The colours were not well graduated, and did not possess the sheen or gloss of the natural hair, on which so much of the variation of the colour depends. On the subject of colour-scales, see the *Bulletins* of the Society of Anthropology of Paris, 3rd S. vi. pp. 91, 92.

TABLE A.—Classification of the Counties of Great Britain and the Provinces of Ireland according to the prevalence of Fair Complexion or the degree of Nighescence of Adult Males.

| The same of the sa | |
|--|--|
| Neutral eyes, with light and dark hair No. per cent, 1 = 0 to 2 2 = 2 - 4 3 = 4 - 6 4 = 6 - 8 5 = 8 upwards | |
| Degree of nigrescence | 4 999899 L 888888 C 8 C 8 C 8 8 8 8 8 C 8 C 8 C |
| Dark eyes, with dark hair No. per cent. 1 under 20 2 = 20-25 3 = 25-30 4 = 30-35 5 = above 35 | on പയപപരായ പക്രാ രായയയെ പയ രാക്യ കുക്കുക്കുക |
| Fair eyes, with dark hair No. per cent. 1 = 10 to 15 2 = 15 - 20 3 = 20 - 25 4 = 25 - 30 5 = 30 upwards | н фафыца пт в ффверы фвер вазафа |
| Fair eyes, with fair hair No. per cent. 1 = above 50 2 = 45-50 3 = 40-45 4 = 35-40 5 = 30-35 | |
| | English and Scotch East Border Group. Central Irish Group. Scotch High. Scotch High. Scotch East Scotch High. Scotch High. Scotch High. Scotch East Scotch East Scotch High. Scotch Group. North-East Scotch Group. Scotch Group. North English Scotch Group. North English Scotch Group. North English Scotch Group. Scotch Group. North English Group. North English Scotch Group. Sc |

| N. W. CO | The state of the state of | | | | |
|--|--|---|----------------------------|--|---|
| 4140 | 00000000000000000000000000000000000000 | 10 to 01 H 10 01 t | 0 10 | 01 00 00 11 | 10 4 10 10 4 |
| | | - | | | , |
| 9 9 10 | 01 01 01 00 01 00 00 00 00 00 00 00 00 0 | 8 6 6 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 | 8 | 112 | 112111 |
| | | | | | |
| | 10 10 4 4 10 80 41 4 41 80 41 | 8 70 83 44 63 70 4 | H 67 | 10 10 10 10 | 47001470 |
| 1 | | A Comment | | | |
| | | | | | |
| 8181 | H 21 24 25 24 4 H 25 24 25 24 | Ø1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 | 000 | 00 01 01 01 | 01 01 10 00 41 |
| | | | | | |
| | | | | | |
| 00 00 00 4 4 | 000000000000000000000000000000000000000 | 00 00 00 00 00 00 | 60 | 41044 | 44101010 |
| 1 | | | | | |
| | | omery | | | |
| | | Montgomer: | | | |
| | · · · · · · · · · · · · · · · · · · · | nd lbr | | | |
| nshire | excel | neth, | | | |
| ingdo . | ingha | Merion | | shire | ilesex |
| Norfolk Cambridge and Huntingdonshire Northamptonshire Bedfordshire | Kent Sussex Surrey Oxfordshire and Buckingham Hampshire Wiltshire Gloucestershire (apparently exceptiona Dorsetshire Somersetshire Cornwall | Denbigh and Flintshire Carnarvon, Anglesea, Me Cardiganshire Brecon and Radnor Glamorgan, Caermarthen Shropshire Hereford and Monmouth | | tland | Mide |
| Norfolk Cambridge and H Northamptonshire Bedfordshire | e and in the control of the control | Angline Radr Caer Caer | | e nd Ru re iire | re and |
| Norfolk Cambridge a Northampton Bedfordshire | Kent . Sussex . Surrey . Oxfordshire Hampshire Gloucesters! Dorsetshire Somersetshi Devonshire Cornwall | igh ar rvon, gansh n and organ, shire ord ar | er | Inshir ster al ickshi stersh | n . nire rdshii |
| Norfolk Cambrid Northan Bedfords | Kent | Denbigh and Flintshire Carnarvon, Anglesea, Merioneth, and Cardiganshire Brecon and Radnor Glamorgan, Caermarthen, and Pembri Shropshire Hereford and Monmouth | Munster | Lincolnshire Leicester and Rutlandshire Warwickshire | London . Berkshire Hertfordshire and Middlesex Suffolk . Essex |
| 7 | | | | | - |
| English Fen Country Group. | sh . | Welsh Group. | est roup. | glish | ondon and Home Counties Group. |
| Inglish F Country Group. | South-west English Group. | elsh (| South-west Irish Group. | Mid-English Group. | London and Home Coun Group. |
| H | Ĭ. | D 11 | So | ME | Log Gr. |

TABLE XI.—Showing the COLOUR of EYES and HAIR of both Males.

| ١. | | | Carlotte State of the State of | 3 1 2 4 | | | .co.o. | | | | | | | | 100 |
|-----|---------------|--------------------------|---|--|-----------------------|---|--|---------------------------------------|---------------------------------|----------|----------------|----------|---------------|----------|----------------|
| | | | | Three light | | Eyes neutral | | Prox dark | | | Eyes | | | Hair | 1 |
| 100 | Age | Num- ber of obser- | Light bl | Eyes light ne, blue, da ey, grey, di | ark blue, ark grey | Green, brown- grey, l'ght brown | | Eyes dark hazel, dark black | brown, | to grey | to light brown | to black | Fair to brown | to dark | Brown to black |
| | oirth- day | va- tions | | Hair . | | Hair | | Hair | | Blue | 1 to] | Brown | ir to | Light to | uwo |
| | | | | Dark | | | Brown, | | | t. | rees | | 9 | 7 | Br |
| | | | Very fair, fair, light brown, brown | brown, black brown, black | Golden, red | Very fair, fair, red, brown, black | dark brown, black brown, black | Very fair, fair, light brown | Red, auburn, red brown | Light | Mixed-Green | Dark- | Light- | Red- | Dark |
| - | | | per cent. | per cent. | per cent. | per cent. | per cent. | per cent. | per cent. | pe | er cen | it. | pe | er cei | at. |
| î. | Birth | 40 | 100 | - | _ | _ | - | 20 | _ | 100 | - | - | - | - | - |
| | 1/2 | 29 | 62\ | -\ | -\ | 24\ | -\ | 14\ | -1 | 62 | 24 | 14 | 76 | - | -1 |
| | 1 | 5 | 60 | 20 | - | 20 | - | - | - | 80 | 20 | - | 60 | - | 20 |
| 1 | 2 | 3 | | | | | | | | 100 | - | - | 100 | - | 31 |
| 1 | 3 | 64 | 50 | 6 | 8 | 12 | 16 | 6 | 2 | 64 | 12 | 24 | 56 | 10 | 22 |
| | 4 | 101 | 52) | 2) | 7/ | 4) | 15) . | 19) | 1) | 61 | 4 | 35 | 71 | 8 | 17 21 |
| | 5 | 197 | 52 | 5) | 7) | 7 | 16) | 12 | 1) | 64 | 7 | 29 | 64 58 | 5 | 24 |
| - | 6 | 222 | 51 | 5 | 4 | 13 | 19 | 7 7 7.4 | 1 1-1-0 | 60 | 14 | 25 | 58 | 6 | 29 |
| | 7 | 265 | 51 -51.4 | 5 5.6 | 5 4.6 | 14 -10.8 | 17 -19-2 | 7 7.4 | 1 1 | 58 | 13 | 29 | 53 | 6 | 28 |
| 4 | . 8 | 270 | 47 | 6 7 | 5 | 13 7 | 22 | 5) | 1) | 65 | 7 | 28 | 61 | 3 | 29 |
| 8 | 9 | 340 251 | 56/ | 12 | 2) | 4) | 22 / 24 \ | 3\ | 3 | 66 | 4 | 30 | 55 | 5 | 36 |
| ı | 10 | 265 | 52 | 11 | 5 | 4 | 20 | 5 | 1 | 70 | 4 | 26 | 59 | 6 | 31 |
| 1 | 12 | 352 | 50 -51.2 | 14 -12-8 | 2 3.2 | 11 -6.2 | 20 -21.4 | 2 -3.6 | 1 -1.6 | 66 | 11 | 23 | 52 | 3 | 34 |
| ı | 13 | 464 | 48 | 12 | 4 | 6 | 23 | 5 | 2 | 64 | 6 | 30 | 53 | 6 | 35 |
| | 14 | 378 | 52 | 15) | 3) | 6) | 20) | 3) | 1) | 70 | 6 | 24 | 55 | 4 | 35 |
| ı | 15 | 253 | 53 | 14\ | 3\ | 10 | 17\ | 2\ | 1) | 70 | 10 | 20 | 55 | 4 | 31 |
| 1 | 16 | 278 | 43 | 17 | 5 | 11 | 20 | 3 | 1 | 65 | 11 | 24 | 46 | 6 | 37 |
| i | 17 | 345 | 40 -43.8 | 14 -14.2 | 4 -4.2 | 12 -11-2 | 25 -22-6 | 4 32 | 1 -0.8 | 58 | 12 | 30 | 44 | 5 | 39 |
| 1 | 18 | 448 | 44 | 13 | 3 | 11 | 26 | 3 | - | 60 | 11 | 29 | 47 | 3 | 39 |
| | 19 | 454 | 39) | 13) | 6) | 12) | 25) | 4) | 1) | 58 | 12 | 30 | 43 | 7 | 38 |
| | 20 | 331 | 42 | 14) | 6) | 8 | 26 | 3 | 1) | 62 | 8 | 30 | 45 | 7 | 40 |
| - | 21 | 281 | 48 | 18 | 3 | 6 | 22 | 2 | 1 | 69 | 6 | 25 | 50 | 4 | 40 |
| 1 | 22 | 257 | 39 -42-2 | 15 -16.4 | 3 -3.2 | 9 -8.6 | 32 -27.0 | 1 1.1.8 | 1 -0.8 | 57 | 9 | 34 29 | 40 | 3 | 43 |
| | 23 | 261 | 43 | 17 | 2 | 9 | 26 | 2 | 1 | 62 59 | 9 | 30 | 40 | 9 | 47 |
| | 24 | 236 | 39) | 18) | 2) | 11) | 29) | 1/ | | 63 | 7 | 30 | 44 | 5 | 44 |
| | 25 | 199 | 41) | 17) | 5) | 7 | 27 | 3 | _) | 60 | 6 | 34 | 37 | 4 | 53 |
| | 26 | 183 | 36 | 20 | 4 4.0 | 8 -7-0 | 33 32.0 | 3 20 | 3 -1-2 | 56 | 8 | 36 | 37 | 5 | 50 |
| | 27 | 189 | 34 - 32-8 | 20 - 20-8 | 2 4.2 | 5 | 34 | 2 | 1 | 58 | 5 | 37 | 30 | 4 | 61 |
| | 28 29 | 179 150 | 28 25 | 20 | 7) | 9) | 36 | 1) | 2) | 52 | 9 | 39 | 26 | 9 | 56 |
| | 30-40 | 900 | 34 | 26 | 5 | 6 | 26 | 2 | 1 | 65 | 6 | 29 | 36 | 6 | 52 |
| 1 | 40-50 | 392 | 33 | 34 | 6 | 6 | 20 | 1 | | 73 | 6 | 21 | 34 | 6 | 54 |
| - | 50-60 | 85 | 36 | 22 | 13 | 7 | 20 | 1 | 1 | 71 | 7 | 22 | 37 | 14 | 49 |
| | 60-70 | 32 | 53 | 19 | 6 | 3 | 19 | - | - | 78 | 3 | 19 | 53 | 6 | 38 |
| 1 | 70- | | | | | 1 | | | | | | | | | |
| I | | | | | | | | | | - | | | | | |

Sexes at all Ages of English and Welsh Origin.

Females.

| | | | | | Eyes neutral | | | | | Eye | g | | Hair | |
|---------------|--------------------------|--|--|----------------|---|--|---------------------------------------|---------------------------------|--------------------|----------------|------------|----------|-------------------|---------------------|
| Age | Num- ber o' obser- | Light b | Eyes light | ark blue, | Green, brown- grey, light brown | Brown | Eyes dark , hazel, darl black | | to grey | to light-brown | to black | to brown | Red-Light to dark | Dark-Brown to black |
| birth- day | va- tions | | Hair | | Hair | | Hair | | Blue | n to] | nwo | Bair | ight t | rown |
| | | Very fair, fair, light brown, brown | Dark brown, black brown, black | Golden, red | Very fair, fair, red, brown, black | Brown, dark brown, black brown, black | Very fair, fair, light brown | Red, auburn, red brown | Light-Blue to grey | Mixed-Green | Dark-Brown | Light-1 | Red- L | Dark-B |
| | | per cent. | per cent. | per cent. | per cent. | per cent. | per cent. | per cent. | - pe | er ce | nt. | p | er ce | nt. |
| Birth | 36 | 100 | - | - | - | - | - | - | 100 | 1- | 1- | - | - | 1 - |
| 1/2 | 34 | 59 | 6) | -) | 20\ | 3 | 12 | -) | 65 | 20 | 15 | 71 | - | 9 |
| 1 | 11 | 46 | - | - | - | 30 | 18 | - | 46 | - | 54 | 64 | - | 36 |
| 2 | 15 | 47 -47.8 | 7:3 | | | 33 -21.4 | 13 -16.4 | 7 - 4.0 | 47 | - | 53 | 60 | 7 | 33 |
| 3 | 45 | 47 | 7 | - | 4 | 18 | 22 | 2 | -54 | 4 | 42 | 69 | 2 | 25 |
| 4 | 119 | 40) | 9) | 9) | 5) | 17) | 17) | 3) | 58 | 5 | 37 | 57 | 12 | 26 |
| 5 | 192 | 53 | 7) | 3 | 9) | 19 | 8) | 1) | 63 | 9 | 28 | - 61 | 4 | 26 |
| 6 | 229 | 51 | 10 | 5 | 9 | 17 | 5 | 3 | 66 | 9 | 25 | 56 | 8 | 27 |
| 7 | 223 | 53 -52-6 | 9 - 8.8 | 3 - 3.4 | 9 - 7.6 | 16 -20.2 | 8 - 5.6 | 2 - 1.3 | 65 | 9 | 26 | 61 | 5 | 25 |
| 8 | 178 | 53 | 12 | 3 | 7 | 21 | 3 | 1 | 63 | 7 | 25 | 56 | 4 | 33 |
| 9 | 237 | 53) | 6) | 3/ | 4) | 28) | 4) | 2) | 62 | 4 | 34 | 57 | 5 | 34 |
| 10 | 281 | 60 | 8 | 4\ | 2 | 22 | 3 | 1) | 72 | 2 | 26 | 63 | 5 | 30 |
| 11 | 322 | 52 | 11 | 4 | 4 | 22 | 5 | 2 | 67 | 4 | 29 | 57 | 6 | 33 |
| 12 | 298 | 47 -51.0 | 15 -10.8 | 3 - 3.8 | 6 - 5.6 | 25 -23.8 | 1 - 3.2 | 3 - 1.8 | 65 | 6 | 29 | 48 | 6 | 40 |
| 13 | 251 | 47 | 10 | 5 | 6 | 27 | 4 | 1 | 62 | 6 | 32 | 51 | 6 | 37 |
| 14 | 265 | 49) | 10) | 3) | 10) | 23) | 3) | 2) | 62 | 10 | 28 | 52 | 5 | 33 |
| 15 | 167 | 45 | 12 | 1) | 9/ | 27 | 4 | 2 | 58 | 9 | 33 | 49 | 8 | 39 |
| 16 | 110 | 52 | 17 | 1 | 5 | 20 | 4 | 1 | 70 | 5 | 25 | .56 | 2 | 37 |
| 17 | 47 | 55 -47.0 | 17 -15.0 | | 7 - 7.8 | 17 -23.8 | 4 - 4.4 | | 72 | 7 | 21 | 59 | - | 34 |
| 18 | 64 | 37 | 15 | - | 9 | 31 | 8 | - | 52 | 9 | 39 | 45 | - | 46 |
| 19 | 94 | 46) | 14/ | 4/ | 9) - | 24) | 2) | 1) | 64 | 9 | 27 | 48 | 5 | 38 |
| 21 | 125 | 46 | 10 | 3) | 10) | 25 | 5) | 1) | 59 | 10 | 31 | 51 | 4 | 35 |
| 22 | 53 | 38 | 17 | 3 | 10 | 30 | 2000 | 2 | 58 | 10 | 32 | 38 | 5 | 47 |
| 23 | 31 | 51 -45.0 | 15 -13.0 | 2 - 3.2 | 9 -11.8 | 19 -24.0 | 4 4.0 | | 68 | 9 | 23 | 55 | 2 | 34 |
| 24 | 20 | 45 | 13 | 3 | 10 | 26 | 3 | - | 61 | 10 | 29 | 48 | 8 | 39 |
| 25 | | 45) | 10) | 5) | 20) | 20) | -/ | -/ | 60 | 20 | 20 | 45 | 5 | 30 |
| 26 | | | | | | | 14. | 1 | | | | - | | |
| 27 | - 46 | 38 | 15 | 2 | | 0.1 | 2 | 100 | 1204 | | | | | |
| 28 | | 00 | 10 | 2 | 8 | 31 | 2 | 4 | 55 | 8 | 37 | 40 | 6 | 46 |
| 29 |) | | | Pho 19 | | | | | 1 | | | 17 | | |
| 30-40 | 27 | 44 | 12 | 3 | 11 | 20 | | | | | | | - | |
| 40-50 | 1 | | 14 | 0 | 11 | 30 | - | - | 59 | 11 | 30 | 44 | -8 | 42 |
| 50-60 | | | 1100 | | | | | | | | | | | |
| 60-70 | 20 | 30 | 20 | 15 | 10 | 25 | _ | - | 65 | 10 | 25 | 30 | 15 | 45 |
| | | | | | | | | | WW. | 1.13 | 40.68 | 43/1.5 | 1 63 | 45.00 |

49. In connection with this subject Table XI., showing the colour of eyes and hair in both sexes and at all ages, should be studied, as it shows the comparative worthlessness of the method often resorted to on the Continent of determining the racial elements of a country by examining the complexion of school children of different ages. The first column, referring to males (light eyes and fair hair). shows the gradual darkening of the hair of fair-complexioned children from 56 per cent. at the first five years of life to 33 per cent. at forty-five years; and the second column (light eyes with dark hair) increases during the same period at nearly a corresponding rate, the percentage of dark hair being 9.3 in the first five years and 34 at forty-five years of age. Thus, 56 + 9.3 = 65.3, and 33 + 34 = 67, or only 1.7 per cent. excess of dark hair received from other sources, or due to probable error of observation. In like manner the green and light-brown eyes of the middle column of the table decrease in number, or in other words become darker, and are transferred to the next column (dark eyes and dark hair) as age advances, from 15 per cent. at the first five years to 6 per cent. at forty-five years of age. The fifth column (dark eyes and hair) increases at the expense of the two adjoining columns from 15.5 per cent. at three and four years to 36 per cent. at twenty-nine years, after which age the percentage falls off very rapidly on account of the earlier accession of grey hair in the dark than the fair complexion of the first column, to which the higher percentages become transferred. The low percentage of dark complexion at ages from forty to seventy years does not arise from the elimination of this complexion by advancing age, or by death, but from the fault of the observers not having recorded the original colour of the hair before it became grey, which necessitated the rejection of all such returns in drawing up the table.

50. The table referring to females shows that darkening of the hair and eyes takes place to a much less extent amongst them than among males, and that there is little disposition for the dark hair to turn grey with advancing age. For corresponding periods to those applied to males, the fair-complexioned females in the first column lose 3.8 per cent. of their number, while the second column receives an accession of dark hair of 4.7 per cent. The dark-complexioned (dark eyes and hair) females in the fifth column increase by 8.6 per cent., at the sole expense of the sixth column, by the darkening of the hair. Unlike the males, the column showing the neutral eyes somewhat increases instead of decreases; and this increase appears to have come from the column containing the fair eyes and red hair, or it may be attributed to the difference in the 'colour equation' of some of the observers—women being much more critical, and therefore less consistent, than men in the definition of colours.

Note.—Dr. Beddoe proposes the use of indices of nigrescence for the classification of the colour of hair and eyes. 'That for the hair is got by subtracting the fair and the red from the dark hair plus twice the black, leaving out the neutral browns, thus:—

2 Black (N) + Dk. Br. - Fair - Red = Index.

The black hair is doubled, because its occurrence shows a much greater tendency to melanosity. The index for the eyes is got by subtracting the light from the dark and neglecting the neutral shades, thus:-

Dark - Light = Index.'

CHILDREN AND ADULTS OF BOTH SEXES.

51. A large portion of the statistics collected by the Committee refer to children, and these, together with those referring to the adults already considered in the early part of this Report, have been arranged in Tables XV. to XXV. to show the influence of age, sex, nurture, occupation, and sanitary surroundings on the physical development of the British population. The children are chiefly those of English parents, as few returns have been received from other parts of the kingdom. All classes of the community are represented, from the upper and professional classes whose children attend the Public Schools, like Eton, Marlborough, and Radley, to the poorest town population, whose children are found in the public elementary (or Board) schools, charitable institutions, and industrial schools. The adults also include all classes, from the Universities of Oxford and Cambridge, to town labourers and factory operatives.

52. In deciding upon the arrangement for practical purposes of returns so varied in their origin, and yet consisting in so large a proportion of information derived from special sources, the first consideration has been to establish a classification of the returns according to the media, or influences which have been instrumental in differentiating one class from another. The Committee has adopted the subjoined scheme, prepared by Mr. Roberts, and first brought before the Association in a paper read in the Anthropological Section in 1878. It is based on the principle of collecting into a standard class as large a number of cases as possible which imply the most favourable conditions of existence in respect to fresh air, exercise, and wholesome and sufficient food—in one word, nurture -and specialising into classes which may be compared with this standard those which depart more or less from the most favourable condition. By this means, in respect to social condition, the influence of mental and manual work; in respect to nurture, the influence of food, clothing, &c., on development; in respect to occupation, the influence of physical conditions; and in respect to climate and sanitary conditions, the influence of town and country life may be determined.

53. The classification has been constructed on the physiological and hygienic laws which are familiar to the students of sanitary science, and on a careful comparison of the measurements of different classes of the people, and especially of school children of the age of from eleven to twelve years. This age has been selected as particularly suited to the study of the media, or conditions of life, which influence the development of the human body, as it is subject to all the wide and more powerful agencies which surround and divide class from class, but is yet free from the disturbing elements of puberty and the numerous minor modifying influences, such as occupation, personal habits, &c., which in a measure shape the physique of older boys and adults. The data on which the classification has been based are given below. The most obvious facts which the figures disclose are the check which growth receives as we descend lower and lower in the social scale, and that a difference of five inches exists between the average statures of the best and the worst nurtured classes of children of corresponding ages, and of 31 inches in

adults.

TABLE XII.—Classification of the British Population according to Media — Occupation and other conditions of life.

| | | Selected Classes | | CLASS VI. Policemen. Fire Brigade. Soldiers. Recruits. Lunatics. Criminals. Industrial. schools. |
|---|---------------------|--|-----------------------|--|
| | Bad | Industrial Classes (Sedentary Trades) 10.90 per cent. | In-door Towns | CLASS V. Fractory Operatives. Tailors. Shoemakers. &c. |
| Labouring Classes | rfect | Artisans 26.82 per cent. | In-door Towns | CLASS IV. Workers in " Wood. " Metal. " Stone. " Paper. " Paper. " Ro. Engravers. Photographers. Printers. |
| | Imperfect | Labourers 47.46 per cent. | Out-door Country | CLASS III. Labourers and Workers on Agriculture. ", Gardens. ", Roads. ", Railways. ", Quarries. Porters. Guards. Woodmen. Brickmakers. Labourers, &c., on Water. ", Sailors. ", Fishermen. ", Fishermen. ", Watermen. ", Watermen. ", Watermen. ", Watermen. ", Goal. ", Minerals. |
| g Classes. | Good | Commercial Class (Lower Mid. Classes) 10.36 per cent. | In-door Towns | CLASS II. Teachers in Elementary Schools. Clerks. Shopkeepers. Shopmen. Dealers in Drugs. Wool. Wool. Silk. Cotton. Foods. Herniture. Herriture. |
| Social Condition.*-Non-labouring Classes. | Very Good | Classes † Middle Classes) cent. | In-door Towns | Forces. Forces. Forces. Forces. gineers. gineers. Givil Servants. Authors. Artists. Art |
| Social Cone | Nurture.†—Very Good | Professional Classes † (Upper and Upper Middle Classes) 4-46 per cent. | Out-door § Country | Country- gentlemen. Gentlemen. farmers. Officers of Army and Navy. Clergymen. Lawyers. Doctors. Civil Engineers. Architects. Dentists Author Artists Teache Musici Actors Banker |

Social Condition; (influences of leisure, mental and manual labour).
 Nurture; (influences of food, clothing, nursing, domestic surroundings, &c.)
 Occupation; (influences of external physical conditions, exercise, &c.)
 Percentage of male population, including male children (Census of 1871).
 Climatic and sanitary surroundings.

TABLE XIII.—Table showing the RELATIVE STATURES of Bors of the age of 11 to 12 years, under different social and physical conditions of life. The zigzag line running through the means shows the degradation of stature as the boys are further and further removed from the most favourable conditions of growth.

| Industrial | | - - - - - - - - | 99 | 20-03 | 20.0 |
|------------------------------------|--------------|--|-------|--------------|-------------|
| Military Asylums | | - 12 2 15 2 46 84 84 118 114 10 0 0 0 0 1 10 | 840 | 51.20 | 51.0 |
| and Work- | Towns | 177 20 20 38 88 50 57 77 113 113 | 341 | 51-56 | 51.5 |
| y Schools Factories and Work-shops | Country | 10 10 10 13 13 11 11 11 11 11 11 11 11 11 11 11 | 293 | 52.17 | 52.0 |
| Elementary Schools Artisans | Towns | 22 24 254 286 177 177 177 | 181 | 52-60 | 52.5 |
| Agricultural | Country | 1 | 904 | 53.01 | 53.0 |
| Middle-class Schools Upper Lower | Towns | 20 27 27 27 20 40 11 11 11 11 11 11 11 11 11 11 11 11 11 | 392 | 53-70 | 53.5 |
| Middle-cla Upper | Towns | 25.55.55.55.55.55.55.55.55.55.55.55.55.5 | 294 | 53:85 | 54.0 |
| Public | Country | 22 23 33 33 115 144 165 175 175 175 175 175 175 175 175 175 17 | 150 | 54.98 | 55.0 |
| Total No. | Observations | 66 85 66 118 230 329 361 441 370 367 252 132 102 12 12 11 | 2862 | 52.60 | 52.5 |
| Height | mones | Mean Mean 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 | Total | Average hgt. | Mean height |

Table XIV.—Showing the Relative Stature of Adults of the ages from 25 to 30 years under different social and physical conditions of life.

The horizontal black lines show the mean Height of each class and the degradation from the standard class

| ial Classe | Lunatics, all Classes | 1 1 2 2 2 2 2 2 2 2 | 341. | 65.5 | 65-65 | |
|---------------------------|---|---|-------|-------------|-----------|--|
| Class VI.—Special Classes | Prisoners, I | 1 2 2 2 2 4 2 2 2 2 2 2 1 1 1 1 2 2 2 2 | 491 | 0-99 | 91-99 | |
| Class V. Sedentary | Occupa- tions: Factories, Tailors | 1 4 0 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 | 198 | 65.5 | 65.93 | The second second |
| Class IV. | Artisan Classes living in Towns | | 342 | 999 | 19-99 | |
| Class III. | Labouring Classes, Agriculturl. Miners, Sailors | 11.03 11.03 11.03 11.04 11.11 11.11 11.12 11.13 | 945 | 67.5 | 67-51 | |
| Class II. | 7 7 2 | 1112288 4 8 8 4 8 1 9 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 | 180 | 0.89 | 67-95 | The same of the sa |
| | Class I. Professional Classes | 1 1 2 2 4 2 2 1 1 1 1 1 1 1 1 | 107 | 0-69 | 69-14 | |
| | General Population, all Classes | 10 10 10 10 10 10 10 10 10 10 10 10 10 1 | 1935 | 67.5 | 67-43 | - |
| | - | AND DESCRIPTION OF THE PARTY OF | | | | 1 |
| | Height in inches | 5445446 6 8 6 9 6 4 6 8 6 6 8 6 6 8 6 8 6 8 6 8 6 8 6 8 | Total | Mean Height | Average . | The state of the s |

Infants at Birth. Table XV.

54. The statistics relating to infants at birth have been tabulated separately, because the conditions of measurement differ from those of other children, the stature having been taken in the recumbent position, and the weight without clothing. The parents of the infants were English and Scotch; and although the charitable institutions from which the observations were obtained are situated in London and Edinburgh, persons bred in the country are frequently admitted as inmates, and it is probable, therefore, that the statistics fairly represent the labouring classes. Observations on infants of other classes of society could not be obtained. The statistics refer only to infants presumably born at the full period of gestation, and contain the due proportion of twin births. The table is constructed to show the relative stature and weight of each infant,

and the differences between the sexes.

55. The table is one of great interest to the student examining the physical development and the physical improvement of a race, as it presents the materials with which he has to deal in its earliest and simplest form. According to this table the average length of male infants is 19.52 inches, and of females 19.32 inches, showing a difference of only one-fifth of an inch. The average naked weight of male infants is 7.12 lbs., and of females 6.94 lbs., a difference of about 3 ounces in favour of males. The range of height between the tallest and shortest male infants is 10 inches, while that of boys of 15 years, when the disturbing influences of puberty are present, is 27 inches. This wide range in adolescence becomes contracted in adults to 20 inches. The range of height of female infants is two inches less than that of male infants, which may be due to accidental causes, but which suggests a less disposition to variation in the size in females than in males, and which may be the cause of the greater freedom of female infants from accidents at the time of birth. It has been ascertained that still births occur in this country in the proportion of 140 males to 100 females, and this higher death-rate of male infants has been attributed to their greater size. We have no statistics of the size or weight of still-born infants, although they could be more easily obtained than those of living infants, but the table before us would seem to confirm this view, as the largest surviving infants are those of males. It would appear, therefore, that the physical (and most probably the mental) proportions of a race, and their uniformity within certain limits, are largely dependent on the size of the female pelvis, which acts as a gauge, as it were, of the race, and eliminates the largest infants, especially those with large heads (and presumably more brains), by preventing their survival at birth.2

² To ascertain if there is any difference between the circumference of the skull as compared with that of the pelvis in adults of very different races of man, Mr. Roberts has measured the skulls and pelves of some European and Andamanese

The greater disposition to vary in range of stature of males than females has been already referred to in the Report of the Committee for 1880, p. 141, in connection with Sir Rawson Rawson's analysis of the successive annual measurements of 12 boys and 13 girls made by Professor Bowditch, of Harvard, United States. 'A marked feature in the charts when compared together is the greater regularity and parallelism of the growth of the girls, especially at the earlier periods of life.'

TABLE XV. -Showing the HEIGHT (LENGTH) and WEIGHT of Infants of both Sexes at Birth.

| | | | | | | | Wei | ghti | Weight in lbs. (naked) | (nak | (pa | | | | | | | Total | Total at each | Ave | Average | Ra | Ratio, Weight |
|-----------------------------|-------|----|-----------|-----|-------------------------------|---------|-----------|---------|------------------------|---------------|--------|-------------------------------|----------|------|------|------|-----|-------|---------------|--------|---------|------|------------------|
| Height, nearest inch | 8 4 4 | | 45 | | 5 4 | | 63 | | 727 | Marie Control | 188 | | 16 16 | 1 | 104 | 111 | 434 | hei | height | weight | ght | Hei | + Height |
| | M | FE | M. | 1 2 | M. | F. | M. F | E. N | M. F. | M. | E. | M | Ei. | M. | E. | M. | E | M. | F. | M. | F. | M. | E. |
| - 76 | 1 | | | 1 | | | _ | | 1 | 1 | 1 | - | 1 | 1 | 1 | 1 | 1 | 1 | ł | 9.5 | 1 | ∙354 | 1 |
| | 1 | 1 | 1 | 1 | 1 | 1 | 1 | -1 | 1 | 2 | 00 | 2 | 1. | 1 | 1 | 1 | 1 | 9 | 4 | 0-6 | 8.5 | -391 | .356 |
| | 1 | 1 | 1 | -1 | 1 | 1 | 1 | 00 | 4 8 | 80 | 3 11 | 9 | 22 | 1 | 1 | 1 | 1 | 20 | 19 | 8.5 | 8.1 | 988. | -868 |
| . 16 | 1 | 1 | 1 | 1 | 1 | 1 | 11 1 | | 40 88 | 8 21 | 9 | 1 | 5 | 67 | 1 | 1 | 1 | 7.5 | 09 | 7.7 | 7.7 | 998. | -866 |
| 20 | 1 | 1 | 1 | 1 | 5 | 8 | 32 3 | - | 69 50 | 0 25 | 5 16 | 22 | 5 | 1 | 1 | 1 | 1 | 133 | 1117 | 7.4 | 7.3 | .370 | -366 |
| 19 | 1 | 1 | 1 | 1 | 17 | 17 4 | 45 6 | 64 4 | 44 86 | 7 | 7 12 | 67 | 63 | 1 | 1 | 1 | 1 | 116 | 181 | 6.9 | 6.9 | .363 | .363 |
| | 1 | 1 | 1 | 7 | 18 | 30 4 | 41 4 | 47 1 | 10 23 | 3 | 3 2 | 1 | 1 | 1 | 1 | 1 | 1 | 73 | 109 | 6.4 | 6.9 | .355 | .850 |
| 17 | 1 | 1 | 60 | 9 | | 10 | 00 | 9 | 1 | 00 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 19 | 25 | 2.9 | 2.0 | .828 | -885 |
| 16 | 1 | 1 | 4 | 1 | 63 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 9 | - | 0.9 | 2.2 | -312 | .344 |
| 61 | 57 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 63 | 1 | 3.5 | 1 | -533 | 1 |
| Total at each weight . | 03 | 11 | 6 | 118 | 55 | 66 18 | 133 168 | 891 88 | 8 154 | 99 1 | 9 20 | 14 | 14 | 60 | 1 | 1 | 1 | 451 | 466 | 7-1 | 6-9 | 1 | 1 |
| Average height | 15.0 | 11 | 17.0 17.5 | 7.5 | 18-2 | 18-3 19 | 19-0 19-0 | -0 19-8 | 19.7 | 7 20.5 | 5 20-4 | 4 21.5 | 5 20-5 | 21.7 | 21.0 | 22.0 | 1 | 19-52 | 19-32 | 1 | 1 | 1 | 1 |
| Ratio, Height + Weight 4-28 | | 1 | 177 3 | 88. | 3-77 3-88 3-31 3-82 2-92 2-92 | 82 2 | 95 29 | 92 2.64 | 34 2.6 | 2 2.4 | 4 2.4 | 2.62 2.44 2.45 2.26 2.16 2.06 | 3.16 | 2.06 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |

Nore.—According to this table, the average difference in the length and weight of the two sexes at birth is 3th of an inch and 3th of a pound in favour of males; but as the table contains I female and 8 males below 17 inches and 5 lbs., it is probable that the average for females is a little too high. On the other hand, one male is returned as 24 inches long and another as weighing 11 lbs., but there are no females of corresponding size. The statistics were obtained from the Queen Charlotte's Lying-in Hospital, London, and from the Royal Maternity, Edinburgh.

Growth of Children of both Sexes.

56. Tables XVI. to XXII. show the growth of children of four of the five classes into which the returns have been divided. Class I. comprises the upper and professional classes and their children, and it may be accepted as representing the best physique of this country, and used as a standard with which to compare all other classes. According to the census of 1871 this class constitutes 4.46 per cent. of the population. Class II. consists of the commercial classes, such as clerks and shopkeepers and their children, whose occupations are carried on in towns, and for the most part indoors, and therefore under less favourable conditions to healthy development than the constituents of Class I. Class II. comprises 10.36 per cent. of the population. Class III. represents the labouring classes, such as agricultural labourers, fishermen, miners, and others who follow outdoor healthy occupations, but whose nurture is inferior to the two former classes. This class comprises 47.46 per cent. or nearly half the population of the country. Class IV. represents the mass of our town population engaged as artisans. Their trades, being carried on indoors, and requiring less physical exercise than Class III., place them under less favourable conditions as to sanitary surroundings. This class forms 26.82 per cent., or about a fourth of the population. Class V., comprising persons living in towns and following sedentary occupations under the most unfavourable conditions as to nurture and sanitary surroundings, has been omitted from the tables, as sufficient data have not been received to fairly represent it. This class constitutes 10.90 per cent. of the population.

57. The average stature and weight of each of the four classes have been worked out from the number of observations for each class, but as the several classes constitute different proportions of the general population the average representing the 'general population' has not been worked out from the total number of observations, but is the average of

skeletons in the Museum of the Royal College of Surgeons, with the following results:-

| | | | Average circum- ference of | Average circum- ference | Datio of |
|-----------------------|----|----------|-------------------------------------|-------------------------------|--------------------|
| | 18 | Stature. | Pelvis. | of Head. | Ratio of Pelvis |
| | | Metres. | m.m. | m.m. | to Head. |
| 1 European female | | 1.592 | 430 | 500 | 1 to 1.16 |
| 6 European males | | 1.712 | 410 | 530 | 1-1-29 |
| Female pelvis . | | | 430 Male head | 530 | 1-1.23 |
| 10 Andamanese females | | 1.408 | 348 | 462 | 1-1-33 |
| 7 Andamanese males | | 1.492 | 337 | 477 | 1-1.42 |
| Female pelvis | | | 348 Male head | 477 | 1-1.37 |

Only one European female skeleton was available for these measurements, but it

appeared to be in every respect a normal one.

From these measurements it is obvious that the difference between the circumference of the head and the pelvis in the adult is much less in the large European than in the small Andaman race, and it is not improbable that the relatively small pelvis of the female Andamanese has been instrumental, in some measure, in differentiating that diminutive race. It is probably in this direction we must look for an explanation of the degenerating influences of town life and sedentary occupations, as they, together with the new movement for the higher education of women, favour the productions of large heads and imperfectly developed bodies of women in this and other civilised countries, and a corresponding disproportion between the size of the head and the circumference of the pelvis.

the other four averages, and it is therefore the average of the four classes rather than of all the individuals measured and weighed. The observations referring to adults are fairly representative of the general population as they were received from all parts of the country; but those referring to children were received from schools devoted to the education of special classes of society, and in numbers which did not correspond with their respective percentage proportion of the general population. By adopting the average of the averages of the four classes into which the school children have been distributed according to the occupations of their parents, the inequality of the percentage proportion has been eliminated. Tables and a diagram showing the mean stature, weight, chest-girth, and strength of males, as deduced from all the observations collected by the Committee, are given in the Report of 1881.

58. Tables (XIII., XIV.) have already been given (s. 53) which show the falling off in the average stature of children of the age 11-12 years, and of adults of the age 25-30 years, as the conditions under which they live are less and less favourable to healthy physical development. The children vary to the extent of five inches, and the adults to $3\frac{1}{2}$ inches, and corresponding variations occur in the weights and other physical qualities.

59. Plate X. shows the growth in stature, weight, and strength of individuals of both sexes, and the girth of chest, head, arm, and leg of males as far as they have been recorded in the returns received by the committee. The tracings are made from the averages in the column representing the general population. Similar tracings of the standard class (males) having been given in the Report for 1880.

60. An examination of the curves and tables shows the following facts:-

(1) Growth is most rapid during the first five years of life; the observations, however, at those ages are not sufficient in number or variety to give a trustworthy average.

(2) From birth to the age of five years the rate of growth is the same in both sexes, girls being a little shorter in stature and lighter in weight than boys.

(3) From 5 to 10 years boys grow a little more rapidly than girls, the difference being apparently due to a check in the growth of girls at these ages.

(4) From 10 to 15 years girls grow more rapidly than boys, and at the ages 11½ to 14½ are actually taller, and from 12½ to 15½ years actually heavier than boys. This difference appears to be due to a check in the growth of boys as well as an acceleration in the growth of girls incident on the accession of puberty.

(5) From 15 to 20 years boys again take the lead, and grow at first rapidly, and gradually slower, and complete their growth at about 23 years. After 15, girls grow very slowly, and attain their full stature about the 20th year.

(6) The tracings and tables show a slow but steady increase in stature up to the 50th year, and a more rapid increase in weight up to the 60th year in males, but the statistics of females are too few after the age of 23 to determine the stature and weight of that sex at the more advanced periods of life.

(7) The curve of the chest-girth in males shows an increase at a rate similar to that of the weight up to the age of 50 years, but it appears to have no definite relation to the curve of stature.

(8) The strength of males increases rapidly from 12 to 19 years, and

at a rate similar to that of the weight; more slowly and regularly up to 30 years, after which it declines at an increasing rate to the age of 60 years. The strength of females increases at a more uniform rate from 9 to 19 years, more slowly to 30, after which it falls off in a manner similar to that of males. The curves of strength for the two sexes are not parallel: at 11 years females are weaker than males by 22 lbs., at 20 years of age by 36 lbs.

The Period of Maturity in Man.

61. The Tables do not show distinctly at what period man attains his full stature, and much difference of opinion exists on this subject. Some French writers (Barnard, Allaire, &c.) maintain that growth in height goes on until the 32nd or 35th year, and Dr. Baxter arrives at the same conclusion from the statistics of the United States Army; while most English writers (Danson, Aitken, Roberts, &c.) regard the 25th as the year of mature growth, and Dr. Beddoe places it as early as the 23rd year, admitting, however, that a slight increase may take place after this age. The difference of opinion on this subject arises, no doubt, from the faulty method of relying on the measurements of many different individuals, instead of measuring the same individuals from year to year until growth ceases. The elimination of the weak and ill-developed by death, the difficulty of following the same class, and all the members of the class, through successive years, and the selection of special classes (i.e. recruits whose ages are never certain), invalidate all conclusions as to the period of maturity drawn from statistics of measurements of many different persons; but, allowing for these sources of error, and judging by the run of the curves formed by the means and averages, it is probable that little actual growth takes place after the age of 21, and that it entirely ceases by the 25th year. It is evident, moreover, from Table XVI., that the full stature is attained earlier in the well-fed and most favoured class (Class I.) than in the ill-fed and least favoured classes of the community (Class IV.).

62. It is difficult to understand, moreover, how any increase of stature can take place after the bones of the skeleton have become consolidated, and the epiphyses firmly united to the body of their respective bones; and the last of these unions in the long bones, on which the stature depends, occurs about the 23rd year. In adopting the 23rd year for men and the 20th for women as the ages of the attainment of maturity the committee was influenced by these considerations, and a desire to understate rather than overstate its case, and to embrace as large a number of observations as possible in its tables. In inquiries of this kind there is generally a slight amount of unconscious selection, very small persons being passed over, or having objections to being measured; and any deficiency of this kind will be balanced by the loss of growth which may occur after the age of 23 years. Females attain to maturity earlier than males, and the age of full growth has been fixed three years earlier

for them.

Influence of Advancing Age.

63. The maintenance of the stature throughout life as shown by Table XVI. is a new and unexpected fact, but it is probably due to the survival of the taller and better developed members of the population, and the elimination by disease or death of the smaller and feebler ones. Quetelet

TABLE XVI.—Showing the Average STATURE (without shoes), at all Ages, of different Classes of the Population of Great Britain.

Males.

| Age last | Po All | eneral pulatio Classe and Co | s. | Pro | Class I. dession classes. and Co | | Co | lass II, mmerc llasses. Towns | | La | lass III abourin Classes. Country | | A | ass IV. rtisans owns | |
|---------------|--------------|---------------------------------------|-----------|------------|---|----------------------|------------|--|-----------|------------|--|---------------------|----------|-------------------------------|---------------------|
| Birth- day | No. Obs. | Average Height. Inches | Increase. | No. Obs. | Average Height. Inches. | Increase. Inches. | No. Obs. | Average Height, Inches. | Increase. | No. Obs. | Average Height, Inches. | Increase Inches. | No. Obs. | Average Height. Inches. | Increase, Inches |
| Birth | 451 | 19.52 | _ | _ | _ | - | _ | | | _ | _ | _ | 451 | 19-52 | |
| 0-1 | 2 | 27.00 | - | - | - | - | - | - | - | 2 | - | - | - | - | - |
| 1- | 1 | 33.50 | - | - | - | - | - | - | - | 1 | - | - | - | | - |
| 2- | 5 | 33.70 | - | | - | - | - | - | - | 5 | - | - | - | - | - |
| 3- | 33 | 36.82 | - | - | - | - | - | - | - | 22 | 37:41 | - | 11 | 36-23 | - |
| 4- | 107 | 38.46 | 1.64 | - | - | - | - | - | - | 19 | 39.30 | 1.89 | 88 | 37.63 | 1.40 |
| 5- | 201 | 41.03 | 2.57 | - | - | = | - | - | - | 34 | 42.35 | 3.05 | 167 | 39-72 | 2.09 |
| 6- | 266 | 44.00 | 2.97 | - | - | | 1 | 45.50 | - | 34 | 44.59 | 2.24 | 231 | 41.90 | 2.18 |
| 7- | 307 | 45.97 | 1.97 | - | 77 | | 4 | 47.50 | - | 39 | 45.81 | 1.22 | 264 | 44 60 | 2.70 |
| 8- | 1524 | 47-05 | 1.08 | - | - | | 61 | 47:60 | - | 324 | 47.09 | 1.28 | 1139 | 46:46 | 1-86 |
| 9- | 2278 | 49.70 | 2.65 | 22 | 50.80 | 2.89 | 211 | 50.03 | 2.43 | 485 | 49 11 | 2.02 | 1560 | 48.88 | 2:42 |
| 10- | 1551 | 51.84 | 2.14 | 101 | 53.69 | 1.54 | 331 | 53.76 | 2.01 | 783 597 | 50-93 | 1.82 | 336 | 52-68 | 1.84 |
| 11- | 1766 | 53.50 | 1.66 | 242 | 55.23 | 2.06 | 687 902 | 55.29 | 1.72 | 395 | 53-67 | 1.35 | 194 | 53.72 | 1-96 |
| 12- | 1981 | 54.99 | 1.49 | 490 | 57-29 | 1 79 | 857 | 57.43 | 2.14 | 403 | 55.31 | 1.64 | 614 | 55.81 | 2.09 |
| 13- | 2743 | 56·91 59·33 | 1.92 | 869 966 | 59·08 61·29 | 2.21 | 800 | 59.47 | 2.04 | 9 | 57.94 | 2:63 | 1653 | 58-61 | 2.80 |
| 14- | 3428 | 62.24 | 2.42 | 974 | 63.61 | 2.32 | 544 | 62-19 | 2.72 | 515 | 61.82 | 3.88 | 1465 | 61.36 | 2.75 |
| 15- | 3498 | 64.31 | 2.07 | 1102 | 66.23 | 2.62 | 110 | 64.55 | 2.36 | 177 | 63-62 | 1.80 | 1391 | 62-85 | 1.49 |
| 16- | 2780 2745 | 66.24 | 1.93 | 1852 | 67-81 | 1.58 | 107 | 66.59 | 2.04 | 75 | 65.87 | 2.25 | 711 | 64.70 | 1.85 |
| 17- | 2305 | 66.96 | .72 | 1724 | 68-26 | .45 | 62 | 67-44 | -85 | 148 | 66.53 | -66 | 371 | 65-60 | -90 |
| 18- | 1434 | 67-29 | -33 | 951 | 68 8 | .32 | 63 | 67.55 | •11 | 143 | 66.87 | •34 | 277 | 66-17 | -57 |
| 20- | 880 | 67-52 | .23 | 461 | 69-08 | - | 61 | 67.58 | •03 | 183 | 66.93 | •06 | 175 | 66.50 | •33 |
| 21- | 757 | 67-63 | .11 | 364 | 68 70 | •12 | 51 | 67-79 | .21 | 177 | 67.15 | -22 | 165 | 66.55 | -05 |
| 22- | 558 | 67-68 | -05 | 227 | 68-94 | - | 53 | 67.82 | •03 | 169 | 67:35 | •20 | 109 | 66-60 | -05 |
| 23- | 592 | 67.48 | _ | 114 | 68.73 | •03 | 59 | 67.42 | _ | 274 | 67:38 | -03 | 145 | 66-40 | - |
| 24- | 517 | 67.78 | *05 | 57 | 68.82 | .09 | 62 | 68.09 | .27 | 258 | 67.47 | -09 | 140 | 66.55 | - |
| 25- | 1 | | | 1 | | 1 | S 47 | 67-93 | - | 218 | 67.52 | -05 | 92 | 66.40 | - |
| 26- | 1 | | | | | | 47 | 68.07 | - | 194 | 67.46 | - | 74 | 66-46 | - |
| 27- | -1576 | 67-80 | •07 | 107 | 69-14 | -32 | 27 | 68-13 | *04 | 162 | 67:76 | +24 | 66 | 66-67 | -07 |
| 28- | | | 100 | | | 19.30 | 33 | 67.65 | - | 208 | 67:31 | - | 59 | 66-65 | - |
| 29- | 1 | | | | 100 | 2 | 1 26 | 67.96 | - | 163 | 67.54 | - | 53 | 66.82 | -15 |
| 30-35 | 1 | 00.00 | 1 | *** | 00.01 | .07 | 1 85 | 67.70 | - | 745 | 67.59 | - | 180 | 66-65 | - |
| 35-40 | 1886 | 68-00 | *20 | 52 | 69.61 | .37 | 82 | 68.07 | - | 631 | 67.62 | - | 111 | 67-08 | -26 |
| 40-50 | 1148 | 67.96 | - | 46 | 69.38 | - | 79 | 68.09 | - | 943 | 67.56 | - | 80 | 66.80 | |
| 50-60 | 198 | 67.92 | - | 13 | 69-50 | - | 16 | 67-69 | - | 147 | 68.06 | .30 | 22 | 66.45 | |
| 60-70 | 44 | | - | 5 | 69.10 | - | 3 | 66.16 | - | 34 | 67.88 | - | 2 | 66-50 | _ |
| 70- | 12 | 69.22 | 1.22 | - | - | - | 1 | 68.50 | _ | 11 | 69-95 | 1.89 | _ | | |
| Total Obs. | 37574 | - | - | 10739 | - | - | 5472 | - | - | 8727 | - | - | 12636 | - | - |

TABLE XVII.—Showing the Average STATURE (without shoes), at all Ages, of different Classes of the Population of Great Britain.

Females.

| Birth 466 19 0-1 6 24 1- 9 27 2- 6 32 3- 43 36 | meral alation. Classes. ad Country sequent 19:31 — 4:83 5:52 7:50 2:67 2:33 4:83 6:23 3:90 8:26 2:03 0:55 2:29 2:88 2:33 4:45 1:57 6:60 2:15 | Pro C Town a | Height. | Increase, Increase, Inches | Co | Class II mmerc Classes Towns Average Height — Height — 28.50 — 37.68 | ial | L | Country Average Height Inches 136.78 | ıg | 300 ON 466 6 7 6 | Class IV Artisan Towns Wardish Height 19:31 24:83 27:38 32:00 | s. |
|--|--|-------------------------------------|----------|----------------------------|------------------------------|---|------|-----|--|------|------------------|---|------------------|
| Birth 466 19 0-1 6 24 1- 9 27 2- 6 32 3- 43 36 | 9:31 — 4:83 5:52 7:50 2:67 2:33 4:83 6:23 3:90 8:26 2:03 0:55 2:29 2:88 2:33 4:45 1:57 6:60 2:15 | | 11111111 | 1111111 | - - 1 - 11 12 | 28·50 — 37·68 | 1111 | | 1111 | | 466 6 7 | 19·31 24·83 27·38 | 5·52 2·55 |
| 0-1 6 24 1- 9 27 2- 6 32 3- 43 36 | 4·83 5·52 7·50 2·67 2·33 4·83 6·23 3·90 8·26 2·03 0·55 2·29 2·88 2·33 4·45 1·57 6·60 2·15 | | 11111 | 11111 | 1 — 11 12 | - 37·68 | | _ | - | - | 6 7 | 24·83 27·38 | 2.55 |
| 0-1 6 24 1- 9 27 2- 6 32 3- 43 36 | 7·50 2·67 2·33 4·83 6·23 3·90 8·26 2·03 0·55 2·29 2·88 2·33 4·45 1·57 6·60 2·15 | 111111 | 11111 | 11111 | 1 — 11 12 | - 37·68 | - | _ | - | - | 6 7 | 24·83 27·38 | 2.55 |
| 2- 6 32 3- 43 36 | 2:33 4:83 6:23 3:90 8:26 2:03 0:55 2:29 2:88 2:33 4:45 1:57 6:60 2:15 | | 11111 | 1111 | - 11 12 | - 37·68 | - | - | - | - | 7 | 27:38 | 2.55 |
| 3- 43 36 | 6·23 3·90 8·26 2·03 0·55 2·29 2·88 2·33 4·45 1·57 6·60 2·15 | | 1111 | | - 11 12 | - 37·68 | | | 36:78 | _ | | 1000 | 12 12 12 12 |
| | 8·26 2·03 0·55 2·29 2·88 2·33 4·45 1·57 6·60 2·15 | | 1.1.1 | | 12 | | _ | 0 | 36:78 | | | 10000 | |
| 4- 99 38 | 0:55 2:29 2:88 2:33 4:45 1:57 6:60 2:15 | | - | - | 1000 | | | | OUT FO | - | 24 | 35.33 | 3.33 |
| | 2·88 2·33 4·45 1·57 6·60 2·15 | _ | - | | 12.5 | 38.50 | *82 | 19 | 38-97 | 2.19 | 68 | 37.30 | 1.97 |
| 5- 157 40 | 4·45 1·57 6·60 2·15 | - | | | 10 | 40.00 | 1.50 | 43 | 41.87 | 2.90 | 104 | 39-77 | 2.47 |
| 6- 189 42 | 6.60 2.15 | | | - | 14 | 42.50 | 2.50 | 44 | 43.43 | 1.56 | 131 | 41.84 | 2.07 |
| 7- 173 44 | | | | - | 30 | 44.43 | 1.93 | 47 | 45.35 | 1.92 | 96 | 43.56 | 1.72 |
| 8- 432 46 | Section 1 | - | - | - | 18 | 47.16 | 2.73 | 119 | 47.10 | 1.75 | 295 | 45.55 | 1.99 |
| 9- 499 48 | 8.73 2.13 | | - | - | 42 | 49.90 | 2.74 | 175 | 48.93 | 1.83 | 282 | 47.36 | 1.81 |
| 10- 480 51 | 1.05 2.32 | 11 | 53.41 | - | 52 | 51.44 | 1.54 | 149 | 50.40 | 1.47 | 268 | 48.96 | 1.60 |
| 11- 441 53 | 3.10 2.05 | 22 | 55.04 | 1.63 | 87 | 53.33 | 1.89 | 115 | 52.48 | 2.08 | 217 | 51.54 | 2.58 |
| 12- 225 55 | 5.66 2.56 | 23 | 57.41 | 2:37 | 87 | 55.68 | 2:35 | 22 | 55.59 | 3.11 | 93 | 53.98 | 2.44 |
| 13- 206 57 | 7.77 2.11 | 68 | 59.03 | 1.62 | 66 | 58:47 | 2.79 | 14 | 57-36 | 1.77 | 58 | 56.22 | 2.24 |
| 14- 240 59 | 9.80 2.03 | 79 6 | 60.78 | 1.75 | 86 | 60.62 | 2.15 | 12 | 59-16 | 1.80 | 63 | 58.56 | 2.34 |
| The state of the s | 0.93 1.13 | 70 € | 62.11 | 1.33 | 98 | 61.28 | -66 | _ | _ | _ | 33 | 59-41 | 0.85 |
| | 1.75 .82 | 49 6 | 62.54 | .43 | 82 | 61.56 | 0.28 | = | _ | _ | 5 | 61.16 | 1.75 |
| | 2.52 .77 | 20 6 | 62.83 | .29 | 68 | 62.22 | -66 | _ | _ | _ | _ | _ | _ |
| | 2.44 - | 25 6 | 62.84 | *01 | 37 | 62.05 | - | _ | | _ | | _ | _ |
| A SECTION AND ADDRESS OF THE PARTY OF THE PA | 2.75 -23 | 48 6 | 63.40 | .56 | 50 | 62.10 | _ | - | _ | _ | _ | _ | _ |
| | 2.98 -23 | 59 6 | 88.39 | - | 71 | 62.58 | .36 | - | _ 1 | _ | | _ | _ |
| | 3.03 .05 | 24 6 | 83-63 | .23 | 36 | 62.44 | - | _ | - | _ | _ | _ | |
| CONTRACTOR OF THE PARTY OF THE | 2.87 - | 1000 | 33.23 | - | 40 | 62.22 | - | - | - | _ | _ | _ | _ |
| 23- 24 63 | | | 33.42 | - | 11 | 62.66 | .08 | - | - | - | | _ | _ |
| 24- 21 62- | | THE R. P. LEWIS CO., LANSING, MICH. | 33.60 | - | - | - | | - | - | - | 16 | 61.81 | -65 |
| 25-30 43 621 | 02 - | The second second | 32-97 | - | - | - | - | - | - | - | 24 | 61.08 | _ |
| 30-35 | | 8 6 | 3.25 | - | - | - | - | - | - | - | - | _ | _ |
| 35-4. | | - | - | - | - | - | - | - | - | - | 11 | 60.90 | _ |
| 40-5: -30 61: | 15 _ | 1- | - | - | - | - | - | - | - | - | 5 | 60-60 | _ |
| 50-60 | | - | - | - | - | - | | - | - | - | 1 | 61.50 | - |
| 60-70 | | (- | - | - | - | - | - | - | - | - | 2 | 60.50 | - |
| 70- 1 | | | - | - | - | - | - | - | - | - | 3 | 60.16 | - |
| Total Obs. 4616 - | - - | 556 | - | - 1 | 1009 | - | - | 767 | - | - | 2284 | _ | _ |

TABLE XVIII.—Showing the Average Weight (including clothes), at all Ages, of different Classes of the Population of Great Britain.

Males.

| Age | Pol | eneral pulatio Classe and Co | n. | Pro | Class I. ofession Classes, and Co | | Co | Class II. mmerci Classes. Towns. | | La (| lass III bourin Classes. Country | g | A | lass IV. rtisans. Fowns | |
|---------------|-------------|---------------------------------------|---------------------|------------|--|----------------------|-----------|---|----------------------|------------|---|----------------------|------------|-------------------------------|----------------------|
| Birth- day | No. Obs. | Average Weight, Pounds | Increase. Pounds | No. Obs. | Average Weight. Pounds. | Increase. Pounds. | No. Obs. | Average Weight. Pounds. | Increase. Pounds. | No. Obs. | Average Weight. Pounds. | Increase, Pounds, | No. Obs. | Average Weight, Pounds. | Increase. Pounds. |
| Birth | 451 | 7.1 | _ | _ | _ | _ | _ | _ | _ | _ | _ | _ | 451 | 7-1 | _ |
| 0-1 | | _ | - | _ | - | _ | - | - | - | - | - | - | - | - | - |
| 1- | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2- | 2 | 32.5 | - | - | - | - | - | - | - | 2 | 32.5 | - | - | - | - |
| 3- | 41 | 34.0 | 1.5 | - | - | - | - | - | - | 11 | 33.1 | - | 30 | 35-0 | - |
| 4- | 102 | 37.3 | 3.3 | - | - | - | 1 | 37.5 | - | 15 | 35.8 | 2.7 | 86 | 38.6 | 3.6 |
| 5- | 193 | 39-9 | 2.6 | - | - | - | - | - | - | 29 | 38-9 | 3.1 | 164 | 40-9 | 2.3 |
| 6- | 224 | 44.4 | 4.5 | - | - | - | - | - | - | 35 | 44.2 | 5.3 | 189 | 44.6 | 3.7 |
| 7- | 246 | 49-7 | 5.3 | - | - | - | 4 | 51.3 | 3.8 | 37 | 47.2 | 3.0 | 205 | 50*7 | 6.1 |
| 8- | 820 | 54.9 | 5.2 | - | - | - | 63 | 55.5 | 4.2 | - 286 | 54.8 | 7-6 | 471 | 54.3 | 3.6 |
| 9- | 1425 | 60.4 | 5.2 | - | - | - | 211 | 62.3 | 6.8 | 415 | 60.5 | 5.7 | 799 | 58.3 | 4.0 |
| 10- | 1464 | 67.5 | 7.1 | 92 | 74.0 | - | 370 | 65.2 | 2.9 | 721 | 67.0 | 6.5 | 281 175 | 64.0 | 5.7 |
| 11- | 1599 | 72.0 | 4.5 | 185 | 78.7 | 4.7 | 686 | 68.0 | 2.8 | 553 | 72:2 | 3.7 | 146 | 69·0 73·0 | 4.0 |
| 12- | 1786 | 76.7 | 4.7 | 369 | 84.9 | 6.2 | 905 | 73.2 | 5.2 | 366 328 | 75:9 | 3.8 | 640 | 79-0 | 60 |
| 13- | 2443 | 82.6 | 5.9 | 621 | 91.6 | 6.7 | 854 | 89.5 | 9.4 | 9 | 89-2 | 9-5 | 1396 | 87.3 | 8.3 |
| 14- | 2952 | 92.0 | 9.4 | 748 | 102.2 | 10.6 | 799 | 99.4 | 9.9 | 676 | 100.6 | 11.4 | 1446 | 96.4 | 9.1 |
| 15- | 3118 | 102-7 | 16.3 | 652 834 | 114.3 | 12.1 | 344 55 | 117:2 | 17.8 | 169 | 117.2 | 16-6 | 1177 | 112.2 | 15.8 |
| 16- | 2235 | 119.0 | 11 9 | 1705 | 141.7 | 15.2 | 38 | 128.8 | 11.6 | 80 | 131.5 | 14.3 | 673 | 121.5 | 9-3 |
| 17- | 2496 | 130.9 | 6.5 | 1638 | 146.4 | 4.7 | 39 | 135.1 | 6.3 | 135 | 138-7 | 7.2 | 338 | 129.3 | 7.8 |
| 18- | 2150 | 137·4 139·6 | 2.2 | 940 | 148.5 | 2.1 | 69 | 138-6 | 3.5 | 140 | 140.2 | 1.5 | 289 | 131.1 | 1.8 |
| 19- | 1438 851 | 143.3 | 3.7 | 451 | 152.4 | 3.9 | 52 | 140-1 | 1.5 | 175 | 144.3 | 4.1 | 173 | 136.4 | 5.3 |
| 20- | 738 | 145.2 | 1.9 | 365 | 152.7 | -3 | 52 | 143.9 | 3.8 | 164 | 147.8 | 3.5 | 157 | 136.2 | - |
| 22- | 542 | 146.9 | 1.7 | 215 | 152.8 | -1 | 51 | 145.5 | 1.6 | 167 | 150.6 | 2.8 | 109 | 138-6 | 2.2 |
| 23- | 551 | 147.8 | .9 | 112 | 151.5 | _ | 57 | 146.8 | 1.3 | 279 | 152.8 | 2.2 | 103 | 140-2 | 1.6 |
| 24- | 30 | 148.0 | .2 | 56 | 149-6 | _ | 57 | 147-1 | •3 | 250 | 151.9 | - | 120 | 148.4 | 3.2 |
| 25- | 100 | 2200 | | - | | 1 | /45 | 148.5 | 1.4 | 224 | 154.1 | 1.3 | 61 | 139-9 | - |
| 26- | | 10 | | | | | 46 | 154.1 | 5.6 | 192 | 154.1 | - | 58 | 142-2 | - |
| 27- | -1523 | 152.3 | 4.3 | 115 | 156.3 | 3.5 | - 26 | 149.2 | - | 171 | 156.7 | 2.6 | 56 | 146.9 | 3.5 |
| 28- | | - | | | | | 33 | 156.1 | 2.0 | 213 | 155.1 | - | 50 | 148.0 | 1.1 |
| 29- |) | 11 1 | | | | | 26 | 154.3 | - | 161 | 158.0 | 1.3 | 46 | 148-1 | -1 |
| 30-35 | 964 | 159.8 | 7.5 | 24 | 171 5 | 15.2 | 87 | 158.5 | 24 | 700 | 159.2 | 1.2 | 153 | 150-1 | 20 |
| 35-40 | 220 | 1000000 | 4.5 | 24 | 173.5 | - | 80 | 166-6 | 8.1 | 631 | 160.5 | 1.3 | 105 | 156.5 | -6:4 |
| 40-50 | 000000 | | - | 44 | 172.5 | 1.0 | 72 | 168-6 | 2.0 | 911 | 162:0 | 1.5 | 113 | 151-7 | - |
| 50-60 | 179 | 166-1 | 1.8 | 13 | 174.5 | 2.0 | 16 | 173.4 | 4.8 | 129 | 170-9 | 8.9 | 21 | 145-6 | - |
| 60-70 | 35 | 158.1 | 2.0 | 5 | 164.5 | - | 3 | 165.7 | - | 24 | 170.9 | - | 3 | 180-8 | |
| 70- | 12 | 182.1 | - | - | - | - | 1 | 189.0 | - | 11 | 175.3 | 44 | | - | |
| Total Obs. | 33043 | - | _ | 9208 | - | - | 5142 | - | _ | 8409 | - | - | 10284 | - | - |

TABLE XIX.—Showing the Average Weight (including clothes), at all Ages, of different Classes of the Population of Great Britain.


Females.

| I | 1 | | | | | | 1 | | | 1 . | | | 1 | | |
|---------------|----------|----------------------------------|------------|-------|----------------------------------|---------------------|-------|--------------------------------|------------|-----------|----------------------------------|---------------------|-------|------------------------------|---------------------|
| | Po | General pulation Il Classo | n. | Pr | Class I. ofession Classes. | nal | | Class II mmerci Classes. | ial | L | llass II. abourir Classes. | ıg | Arti | lass IV san Cla | sses. |
| Age | | and Co | | | and Co | | To | owns on | | | intry o | | To | wns on | ly |
| Birth- | | ge ht. | use. | Obe. | ht. | use. | Obs. | ht. | ise. | bs. | br. | ise. | Obs. | ge ds | ds. |
| | No. Obs. | Average Weight. Pounds | Increase. | No. 0 | Average Weight. Pounds | Increase, Pounds | No. 0 | Average Weight, Pounds | Increase. | No. Obs. | Average Weight, Pounds | Increase, Pounds | No. 0 | Average Weight. Pounds | Increase, Pounds |
| | A | 4 | - | A | 97.0 | - | - | - H | - | - | | A- | - | 420 | |
| Birth | 466 | 6.9 | - | - | - | - | - | - | - | - | - | - | 466 | 6.9 | _ |
| 0- | - | - | - | - | - | | - | - | - | - | - | - | - | - | - |
| 1- | 8 | 20.1 | - | - | - | - | 1 | 22.5 | - | - | - | - | 7 | 19.6 | 12.7 |
| 2- | 9 | 25.3 | 5.2 | - | - | - | - | - | - | - | _ | - | 9 | 25.3 | 5.7 |
| 3- | 30 | 31.6 | 6.3 | - | _ | | 11 | 30.9 | - | 8 | 33-0 | - | 22 | 30.8 | 5.2 |
| 4- | 97 | 36.1 | 4·5 3·1 | - | - | _ | 12 | 37.9 | 7.9 | 17 | 34.6 | 1.6 | 68 | 35.8 | 5.0 |
| 6- | 160 | 39·2 41·7 | 2.5 | _ | | _ | 18 | 38.8 | 0.9 | 44 | 38·4 40·5 | 3.8 | 108 | 40·3 43·1 | 4.5 |
| 7- | 148 | 47.5 | 5.8 | 7 | 51.8 | | 31 | 45.4 | 2.6 | 43 | 46.8 | 6.3 | 122 | 46.2 | 2.8 |
| 8- | 330 | 52.1 | 4.6 | 6 | 52.5 | -7 | 12 | 52.5 | 4·0 7·1 | 42 140 | 51.9 | 5.1 | 99 | 51.8 | 5.6 |
| 9- | 535 | 55.5 | 3.4 | 17 | 55.4 | 2.9 | 23 | 55.0 | 2.5 | 209 | 56.5 | 4.6 | 286 | 55.2 | 3.4 |
| 10- | 495 | 62.0 | 6.5 | 37 | 62-9 | 7.5 | 23 | 62.9 | 7.9 | 171 | 61.8 | 5.3 | 264 | 60.5 | 5.3 |
| 11- | 456 | 68:1 | 6.1 | 61 | 69-9 | 7.0 | 41 | 68.5 | 5.6 | 130 | 67.1 | 5.3 | 224 | 66.8 | 6.3 |
| 12- | 419 | 76.4 | 8.3 | 55 | 79.7 | 9.8 | 55 | 77:3 | 8.8 | 126 | 75.7 | 86 | 183 | 74.9 | 8.1 |
| 13- | 209 | 87-2 | 10.8 | 63 | 89-8 | 10.1 | 60 | 88 2 | 10.9 | 21 | 84.0 | 8.3 | 65 | 84.9 | 10.0 |
| 14- | 229 | 96.7 | 9.5 | 75 | 98.8 | 9.0 | 81 | 96.3 | 8.1 | 12 | 94.0 | 10.0 | 61 | 97-7 | 12.8 |
| 15- | 187 | 106:3 | 9.6 | 60 | 107.3 | 8.5 | 91 | 104.1 | 7.8 | | _ | _ | 36 | 107-6 | 9-9 |
| 16- | 128 | 113.1 | 6.8 | 49 | 113.9 | 6.6 | 75 | 112.2 | 8.1 | _ | _ | _ | | | _ |
| 17- | 74 | 115 5 | 2.4 | 14 | 116.8 | 2.9 | 59 | 114.3 | 2.1 | _ | _ | _ | _ | _ | _ |
| 18- | 64 | 121.1 | 5.6 | 26 | 123.1 | 6.3 | 38 | 119.1 | 4.8 | _ | _ | - | _ | _ | _ |
| 19- | 97 | 123.8 | 2.7 | 47 | 125.5 | 2.4 | 50 | 122.1 | 3.0 | - | _ | - | _ | _ | - |
| 20- | 128 | 123.4 | +6 | 58 | 126.6 | 1.1 | 70 | 120.3 | - | - | - | - | - | - | - |
| 21- | 59 | 121.8 | - | 23 | 125.3 | - | 36 | 118.3 | - | - | - | - | - | - | - |
| 22- | 53 | 123.4 | - | 14 | 122.8 | - | 37 | 124.1 | 2.0 | - | - | - | - | - | - |
| 23- | 29 | 124.1 | -7 | 12 | 128.7 | 2.1 | 16 | 119.4 | - | - | - | - | - | - | - |
| 24- | 19 | 120-8 | - | 5 | 120.5 | - | - | - | - | - | - | - | - | - | - |
| 30-35 | | 120-0 | - | 19 | 119.1 | - | - | - | - | - | - | - | - | | - |
| 35-40 | } 23 | 120-8 | - | 8 | 120-6 | - | - | - | - | - | - | - | - | - | - |
| 40-45 | 9 | 118 | - | _ | _ | _ | - | _ | _ | _ | _ | _ | _ | _ | _ |
| 45-50 | - | - | - | _ | - | _ | - | _ | _ | _ | _ | | | - | |
| 50-60 | 3 | 104 | - | - | - | - | _ | _ | _ | _ | _ | - | _ | _ | _ |
| 60-70 | 777 | - | - | _ | - | - | - | - | _ | - | - | - | | _ | - |
| 70- | 3 | 106:0 | - | - | - | _ | - | - | - | - | - | - | - | - | - |
| Total Obs. | 4685 | _ | - | 656 | - | - | 853 | - | - | 963 | - | _ | 2192 | - | _ |

TABLE XX.—Summary Table showing the average STATURE, WEIGHT, and their relation

| Age last birth- | wit | ight hout oes, nches | clot | ight ith thes, lbs. | Chest-g empt in inc | y, | drav | ngth: ving- wer, lbs. | Spar arms a the b in in | ack, | we | | Ratio weight divided chest-g | ht by |
|-----------------------|-------|-------------------------------|----------------|------------------------------|---------------------------|-------|------|--------------------------------|---|---|-------|-------------------|---------------------------------------|----------|
| day | М. | F. | М. | F. | М. | F. | М. | F | М. | F. | М. | F. | M. | F. |
| Birth | 19.52 | 19:31 | 7-1 | 6.9 | 18.25 | 12.65 | _ | | _ | _ | _ | - | - | - |
| 0-1 | 27.00 | 24.88 | _ | - | - | - | - | _ | - | - | - | - | - | - |
| 1-2 | 33.50 | 27.50 | - | - | - | - | - | - | - | - | - | - | - | - |
| 2- | 33 70 | 32.33 | 82.5 | | - | - | - | - | - | - | -96 | - | 1 | - |
| 3- | 36.82 | 36.05 | 34.0 | 81.9 | - | - | - | - | - | - | .92 | 0.87 | 7 | - |
| 4- | 38.46 | 38-13 | 37.3 | 85.5 | _ | - | - | - | - | - | -99 | -93 | - | - |
| 5- | 41.03 | 40.82 | 39.9 | 39.6 | - | - | - | - | - | - | -97 | .97 | - | - |
| 6- | 44.00 | 42.63 | 44.4 | 42.4 | - | - | - | - | - | - | 1.01 | 1.00 | - | - |
| 7- | 45.97 | 44.45 | 49.7 | 46.7 | - | - | - | - | 100000000000000000000000000000000000000 | 100000000000000000000000000000000000000 | 1.08 | 1.05 | - | - |
| 8- | 47.05 | 46.60 | 54.9 | 52.2 | - | - | - | 17.5 | 17.56 | 100000000000000000000000000000000000000 | 1.16 | 1.12 | - | - |
| 9- | 49.70 | 48.73 | 60-4 | 55.5 | - | - | - | | 49.07 | 150 | 1.22 | 1.14 | - | - |
| 10- | 51.84 | 51.05 | 67.5 | 62.0 | 26.10 | - | - | 15.1 | 1000 | 49.92 | 1.30 | 1.21 | 2.59 | - |
| 11- | 53.50 | 53.10 | 72.0 | 68-1 | 26.53 | - | 37.5 | | 51.98 | 1000000 | 1.35 | 1.28 | 2.72 | - |
| 12- | 54.99 | 55.66 | 76.7 | 76.4 | 27.20 | - | 38.7 | | 54.03 | | 1.39 | 1.37 | 2.82 | - |
| 13- | 56.91 | 57.77 | 82.6 | 87.0 | 28.03 | - | 44.2 | | 100000000000000000000000000000000000000 | 58.06 | 1000 | 1.51 | 2.95 | - |
| 14- | 59.33 | 59.80 | 92.0 | 96.7 | 28.46 | - | 17.0 | | | 1000 0000 | 1.55 | 1.62 | 3.23 | - |
| 15- | 62.24 | 60.93 | 102.7 | 104.8 | 29.74 | - | 52.2 | 29.6 | - | 60.79 | 1.65 | 1.72 | 3.46 | - |
| 16- | 64.31 | 61.75 | 119.0 | 112.7 | 31.53 | - | 58.2 | 31.8 | - | 61.66 | 1.85 | 1.82 | 3·78 3·89 | _ |
| 17- | 66.24 | 62.52 | 130.9 | 114.9 | 83.64 | - | 67.8 | 88.9 | - | 62.52 | 2.05 | 1 1 1 1 1 1 1 1 1 | 4.02 | |
| 18- | 66.96 | 62.44 | 137.4 | 117:7 | 34.19 | - | 74.2 | 38.9 | - | | 2.05 | 1.89 | 4.05 | - |
| 19- | 67-29 | | 139.6 | 123.7 | 34.49 | - | 76.4 | 40.8 | - | 62.69 | 100 | 1.96 | 4.09 | - |
| 20- | 67.52 | | 143.3 | 123.2 | 34.98 | | 77.9 | 42.0 | | 62.49 | 10000 | 1 | 4.13 | |
| 21- | 67-63 | | 145.2 | 121.2 | 35.25 | | 80.2 | 41 9 | | 62.35 | 1 | 1 | 4.16 | _ |
| 22- | 67.68 | | 3 - 1 | 124.2 | 35.88 | | 81.7 | 42.9 | | 62.36 | | 1000 | 32000 | |
| 23- | 67.48 | | 1 | 126.4 | 35.62 | | 79.7 | 38.5 | | 62-22 | | | 1000 | |
| 24- | 67.72 | | 148.0 | 120-6 | 35.82 | - | 80.9 | 89.2 | - | OF ZZ | 2.20 | | 1 10 | |
| 25- | 67:75 | | 149.2 | 1 | | - | 1 | | - | - | 2-28 | | 1 | |
| 26- | 67.78 | The same of | 151.7 | 100.1 | 36-18 | | 83.5 | 40.8 | | 62-61 | | | 4.21 | - |
| 27- | 67.70 | | 0.395 | -120.1 | 90.18 | | 99.9 | 40.8 | 1 | 0201 | 2.27 | 1 3 75 | | |
| 28- | 67.70 | | 153.9 | 11 | | | 1 | 1 3 | 1 | 1 | 2.27 | | 1 | 1 |
| 30- | 67.89 | | 154·2 159·8 | 1 | | | | 1 | | | 2.85 | | 1 | |
| 35- | | | | 121.0 | 87.08 | - | 77.5 | 46.5 | 2 - | 62-10 | 2.41 | | 4:37 | - |
| 40- | 68.09 | 11 | 164.8 | 118-6 | | | | 1 | | | 2.39 | | 1 | |
| 50- | 67-99 | -01.15 | | 1 | 37.58 | - | 76.5 | 1 | | 1 | 2.44 | 1.96 | 4.38 | - |
| 60- | 67.41 | | 158-1 | _ | 1 | | 74.6 | 88- | - | 60.29 | | | | |
| 70- | 69-22 | | 182-1 | _ | _ | | - | 1 | | | 2.65 | | 1 | |

Diagram showing the Stature Weight, Chest girth and Strength of both Sexis, is all Ages of the General Population of the United Kingdom.



Illustrating the Report of the Anthropometric Committee.



CHEST-GIRTH, STRENGTH, and SPAN OF ARMS of both Sexes and of all Ages, to each other.

| 1 | Ratio: weight | S | ation of pan of rms to | Dif | ference | between | the two with m | | females | compa | red | Age |
|------|------------------|----------------|------------------------------|-----------------|----------------|-------------------|------------------|--------|-------------------|-------|------------------------|----------------|
| | rength | | neight | Hei | ght | We | ight | Stre | ength | Span | of arms | last Birth- |
| M | F. | M. | F. | Actual | Per cent. | Actual | Per cent. | Actual | Per cent. | Actua | Per cent. | day |
| _ | - | - | _ | inches -0.21 | | lbs. - 0.2 | - 2.81 | _ | - | _ | _ | Birth |
| - | - | - | - | - | - | - | - | - | - | - | - | 0-1 |
| - | - | - | - | - | - | - | 1 250 | - | - | - | - | 1-2 |
| - | - | - | - | - | - | - | - | - | - | - | - | 2- |
| | - | - | - | -0.77 | -2.09 | | - 6.17 | | - | - | - | 3- |
| - | - | - | - | -0.33 | -0.86 | | | | - | - | - | 4- |
| - | - | - | _ | -0.21 | -0.51 | | 1000 | | - | - | - | 5- |
| - | - | | - | -1:37 | -3.11 | The second second | 1000 | | - | - | - | 6- |
| | 2.98 | -2·87 + ·51 | +1.38 | -1.52 | -3.30 | | - 6.04 | | - | + 2.7 | +6.2 | 7- |
| | 3.70 | 63 | - ·10 - ·34 | -0.45 -0.97 | -0.99 -1.95 | | - 4·92 - 8·11 | 1 | - | -1.0 | -2.1 | 8- |
| | 4-11 | -1.20 | -1.13 | -0.79 | -1.52 | | - 8·11 | - | - | -0.6 | -1.2 | 9- |
| 1.92 | 3.87 | -1.52 | 69 | -0.40 | | | - 5.41 | -19.9 | _53·0 | -0.7 | -1.3 | 10- |
| .98 | 4.06 | 96 | 62 | + 0.67 | + 1.22 | - 0.3 | - 0.39 | -19.9 | -51.4 | +0.4 | -0.7 | 11- |
| *87 | 3.90 | -1.40 | + .29 | + 0.86 | +1.51 | + 4.4 | + 5.32 | -21.9 | -31 ⁻⁴ | + 2.5 | +1.8 | 12- |
| .96 | 3.79 | -2.18 | 76 | + 0.47 | + 0.79 | + 4.7 | + 5.11 | -21.5 | -45.7 | +1.9 | +4.5 | 13- |
| -97 | 3.54 | _ | 14 | -1:31 | -2.10 | + 2.1 | + 2.04 | -22.6 | -43·3 | | NAME OF TAXABLE PARTY. | 14- |
| 2:04 | 3.54 | _ | 09 | 100000 | -4.00 | - 6.3 | - 5.30 | -26.4 | -45·8 | | | 15- |
| .93 | 3.46 | - | _ | | -5.61 | -16- | -12.21 | | -50.0 | | | 17- |
| .85 | 3.03 | _ | + .06 | -4.52 | -6.75 | -19.7 | -14.34 | -35.3 | 10000000 | _ | | 18- |
| . 83 | 3.03 | - | 06 | -4.54 | -6.74 | -15.9 | -11.39 | | | | | 19- |
| .84 | 2.93 | - | 49 | -4.54 | -6.72 | -20.1 | | | -46.0 | _ | _ | 20- |
| .81 | 2.89 | - | — ·84 | -4.60 | -6.30 | -24.0 | -16.53 | | -47.7 | _ | _ | 21- |
| .80 | 2.89 | - | - ·52 | -481 | -7.10 | -22.7 | -15.45 | -38.8 | -47.2 | _ | _ | 22- |
| .85 | 3.28 | - | - ·65 | -4.41 | -6.63 | -21.4 | -14:49 | -41.2 | -51.7 | _ | - | 23- |
| 1.88 | 3.08 | - | 48 | -5.02 | -7.41 | -27.4 | -18·51 | -41.7 | -50.1 | _ | _ | 24- |
| | | | | 45 | | | | | | | - | /25- |
| 1,00 | 0.01 | | | | Name of | | | | 2 | | | 26- |
| 1.82 | 2 94 | - | - 41 | -5.82 | -8.50 | -38. | -24.97 | -42.7 | -51.0 | - | - | 27- |
| | | | | | | | | | | | | 28- |
| | | | | | | | | | | | | 29- |
| .09 | 2.61 | - | 1 - 1 | - | - | -41.0 | - 25.81 | -31.3 | -40.4 | - | | 80- |
| 1.13 | 1 | | + .05 | 6.00 | 10-10 |) | 1 111 | | | | | 85- |
| 228 | 3.11 | | - 05 | -6.98 | 10.18 | -44.5 | -27:34 | -37.4 | -50.0 | - | | 40- |
| _ | 011 | _ | | | | , | | | 000 | | | 50- |
| - |) | _ | - | | | | - | - | - | - | - | 60- |
| | | | | 1 | | _ | - | - | - | - | - | 70- |
| | | | | | | | | | | | - | |

has stated that man attains his maximum height at the age of 30 years, and maintains it up to 50 years, after which it begins to recede, and at 90 it has lost three inches. This may be, and probably is, true of individuals if measured from year to year, but it does not appear to be true of the population in the aggregate. The loss of stature resulting from the degeneration and loss of tissue, and the stooping position assumed by old people, is more than counterbalanced by the survival of a greater number of individuals who are above the average in height. The uniform increase in the weight and chest-girth throughout adult life also confirms this view.

Industrial Schools.

64. The statistics referring to Industrial School children of both sexes are given in a separate form, as illustrating the physique of children bred under the most unfavourable conditions of life. Boys of this class of the age of 14 years are nearly seven inches (6.83) shorter of stature and 24\frac{3}{4} lbs. lighter in weight than the 1st or Standard Class of the foregoing tables. The returns sent in by Mr. R. Sutton from the Swinton School, near Manchester, are the most complete in all their details which the Committee has received from any source, and they may be accepted as models of what such returns ought to be.

TABLE XXI.—Comparative Table of Boys and Girls in Industrial Schools.

| and and the | | | | | | | | | | |
|--|--|---------|---|---|--|----------|--|---------|--|---|
| | Numb | | Hei | ght | Wei | ght | Chest | girth | Span of | f Arms |
| Age | Males | Females | Males | Females | Males | Females | Males | Females | Males | Females |
| 16- 15- 14- 13- 12- 11- 10- 9- 8- 7- 6- 5- 4- 3- 2- 1- 6 & under 12 months 0 & under 6 months | 7 58 102 221 205 158 191 100 69 64 46 37 9 5 — | | inches 57·64 55·43 54·46 53·23 51·79 49·11 48·09 47·02 44·61 43·54 41·14 38·63 36·27 34·50 — | inches 55.50 55.00 52.98 51.16 51.48 47:70 46.44 44.68 42.38 41.15 39.22 37.07 35.50 31.95 27.00 26.25 | 1bs. 93·92 85·50 77·35 72·31 67·40 63·19 56·76 52·40 47·13 45·70 40·43 36 68 33 61 30 50 — 11·00 | 1bs, | inches 29·25 28·30 27·29 26·31 25·85 24·17 23·97 22·58 22·16 21·95 21·42 20·50 — — — — — | inches | inches 57·50 57·17 54·72 52·45 50·10 49·15 47·46 45·30 41·23 40·30 38·10 35·00 33·25 — — — | inches 58·50 54·21 53·60 51·28 49·11 47·21 45·41 43·46 41·95 39·50 38·25 35·90 32·50 |
| Total | 1,273 | 601 | | | | | | | | |

TABLE XXII. - Statement of the Percentage Proportion in each Class, as regards Colour of Eves and Hare, of Boys and

and green, with light or dark hair brown 14.2 Eyes 5.5 4.5 2.2 7.0 9.9 per cent. 0.9 11:1 5.6 per cent. hair Eyes dark, with 2.7 2.9 4.0 1.4 2.9 Fair per cent. 25.8 16.2 42.9 50.0 14.5 29.0 33.4 35.6 25.8 Dark 28.1 per cent. Girls, of English Parentage, in Industrial and Workhouse Schools, at each age. Girls 2.2 5.4 5.1 per cent. 13.4 11.1 Red Eyes light, with 5.5 5.2 29 3.3 12.3 1.7 Dark per cent. 11 hair 4.4 4.3 Light 6.09 50.0 56.5 9.99 64.9 0.08 46.0 per cent. 9.99 54.3 48.4 63.7 42.7 Eyes light brown, Number of with light observations 560 67 64 94 99 69 57 37 or dark hair per cent. 9.1 5.5 3.8 7.8 2.4 5.3 6.4 2.1 3.7 2.5 1.5 3.5 3.5 per cent. 4.7 Red 2.4 Eyes dark, with 1.8 4.5 1.2 1.6 1.8 3.5 Fair per cent. 9.87 Dark hair 24.6 24.6 24.5 33.3 25.0 32.3 40.0 20.4 27.4 34.1 44.4 26. Boys 3.0 per cent. 2.9 2.4 12.3 3.8 Red 2.1 Eyes light, with Dark hair 29.6 16.0 10.5 99.0 18.0 10.0 14.3 3.5 per cent. 4.7 8.3 12.1 6.4 7.9 Light hair 43.0 9.67 per cent. 37.7 49.0 47.3 47.0 50.0 54.2 49.5 54.1 49.1 0.09 58.1 44.4 58.1 Number of observations 182 165 134 85 89 168 57 48 61 6 10 1,072 31 31 Ages last birthday Total 15 14 13 12 10 H 6 00 1 9 10 00 03 4

TABLE XXIII.—Comparison of Boys and Girls, at different Ages, in Industrial School at Swinton, near Manchester.

| Age | Obs | o. of erva- | Hei | ght | Wei | ight | | est- rth | | thing | of a Dra | ngth arm. wing wer | Test dis guish distar | cht. dots tin- led at lace of et |
|--|---|--|--|---|--|---|--|---|--|--|---|--|---|--|
| | M. | F. | M. | F. | M. | F. | M. | F. | M. | F. | M. | F. | M. | F. |
| 14 13 12 11 10 9 8 7 6 5 4 8 Total | 6 28 41 22 32 32 24 32 28 12 3 1 | 21 27 29 31 27 25 28 20 19 15 3 — | inches 55·0 52·5 54·0 50·0 48·2 46·7 43·8 43·6 40·7 38·9 35·0 34·6 | inches 54·4 51.1 49·9 49·4 47·0 45·8 44·4 41·2 39·0 38·6 35·0 — | 1bs. 78·7 70·0 65·4 63·1 57·1 52·7 47·0 46·2 39·9 35·8 32·3 28·0 | 1bs. 80·9 71·3 64·6 60·3 55·4 52·0 47·3 42·4 37·2 34·8 29·7 — | inches 28·3 26·6 25·9 25·3 23·6 23·0 22·6 22·2 21·4 20·8 20·0 20·0 | inches 29·0 27·3 27·6 27·5 26·9 26·2 25·7 25·5 21·0 20·6 19·3 — | cubic 189 166 166 153 140 132 117 77 48 38 22 20 — | inches 177 143 138 145 124 126 112 83 54 41 30 — | lbs. 40·0 37·3 36·0 34·2 26·7 21·7 18·4 18·5 11·5 6·4 4·0 — | 1bs. 33·0 27·6 27·6 25·4 19·5 18·0 17·0 12·5 8·5 6·8 4·3 — | ft, in. 27·9 30·9 32·6 32·3 28·4 24·2 22·8 23·8 19·8 16·4 9·6 — | ft. in. 38·1 37·2 36·7 39·0 34·8 31·7 36·5 27·6 34·3 19·9 13·0 — |
| 7 15 | | | C | olour o | of Eye | s and | Hair. | Perc | entage | e propo | ortion | in ea | ch Clas | is. |
| | | | Hair | · H | light, | Hair red | Hai | ir H | s dark | Hair red | - e | ght b green xcepti eyes, v ght or hair | Total | |
| Boys 2 | or (Ir. | nglish ish . nglish ish . | 54·6 65·0 39·8 50·0 | 26 | 2·7 3·7 3·1 3·3 | 1·7 3·1 3·4 6·1 | 20· 15· 20· 23· | 5 5 | ·2 ·5 - | 3·4 7 1·1 1·2 | | 6·3 6·7 9·1 1·2 | | 100 100 100 100 |

Physical Improvement or Degeneracy of the Population.

65. Few statistics are in existence which help to throw light on this subject. It is generally believed that the population in the manufacturing towns of the North of England is rapidly degenerating, but a comparison of the measurements of stature and weight given in the Report of the Factory Commissioners of 1833, and in the Report to the Local Government Board on 'Changes in Hours and Ages of Employment of Children and Young Persons in Textile Factories,' 1873, shows that this is not the case. On the contrary, an examination of Table XXIV., showing these measurements, indicates a slight but uniform increase in stature, and a very large increase in weight, at corresponding ages. The increase in weight amounts to a whole year's gain, and a child of 9 years of age in 1873 weighed as much as one of 10 years in 1833, one of 10 as much as one of 11, and one of 11 as much as one of 12 years in the two periods respectively.

66. As an example of the condition of a class living under most favourable conditions, a table (XXV.) showing the measurements of the boys in the Friends' (Quakers') School at York, extending over a period of

twenty-seven years, is given. Allowing for one or two obvious errors of observation, the general run of the figures is very uniform, the statures remaining stationary, while there is a slight improvement in the weight at the higher ages in the last nine years.

Table XXIV.—Showing the average Stature and Weight of Factory Children at an interval of 40 years, 1833-1873. (Stanway and Roberts.)

STATURE.

| Age | | Bo | ys | | Girls | | | | | | |
|-------|-----|-----------|------|---------|-------|--------|-----|--------------|--|--|--|
| Age | 1 | 888 | 1 | 873 | 1 | 833 | 1 | 873 | | | |
| lo de | No. | Inches | No. | Inches | No. | Inches | No. | Inches | | | |
| 9 | 17 | 48.14 | 126 | 48.30 | 30 | 47.97 | 144 | 48.31 | | | |
| 10 | 48 | 49.79 | 256 | 49.85 | 41 | 49.62 | 201 | 50.33 | | | |
| 11 | 53 | 51.26 | 196 | 51.59 | 51 | 51.15 | 174 | 51.21 | | | |
| 12 | 42 | 53.38 | 175 | 53.30 | 80 | 53.70 | - | - | | | |
| | | and Steam | 1010 | WEIGHT. | | | | The party of | | | |
| | No. | lbs. | No. | lbs. | No. | lbs. | No. | lbs. | | | |
| 9 | 17 | 51.76 | 136 | 58 15 | 30 | 51.31 | 137 | 55.87 | | | |
| 10 | 48 | 57.00 | 247 | 60.19 | 41 | 54.80 | 179 | 60.59 | | | |
| 11 | 53 | 61.84 | 189 | 67.72 | 63 | 59.69 | 180 | 65.37 | | | |
| 12 | 42 | 65.97 | 167 | 69.76 | 80 | 66.08 | | _ | | | |

Table XXV.—Showing the average Stature and Weight of Boys in the York Friends' School, for 27 years, 1853-1879.

| | | | STA | TURE | | WEIGHT | | | | | | |
|--|--|--|--|---|--|--|---|--|---|--|--|--|
| Age last Birth- day | No. of Obs. | 27 yrs. 1853 to 1879 | 9 yrs. 1853 to 1861 | 9 yrs. 1862 to 1870 | 9 yrs. 1871 to 1879 | 27 yrs. 1853 to 1879 | 9 yrs. 1853 to 1861 | 9 yrs. 1862 to 1870 | 9 yrs. 1871 to 1879 | | | |
| 9- 10- 11- 12- 13- 14- 15- 16- 1 - | 13 86 261 585 874 1117 1174 515 36 | inches 51·5 53·3 56·4 57·7 59·9 62·1 64·2 66·1 67·2 | inches 51·4 53·9 56·5 58·0 60·6 62·1 63·9 65·4 | inches 49·7 *51·6 56·1 57·9 59·9 62·3 64·3 66·1 67·0 | inches 53·4 54·7 56·5 57·4 59·6 61·9 64·2 66·3 67·4 | 1bs. 62·9 68·5 79·7 85·8 95·4 106·0 116·6 127·8 136·3 | 1bs. 63·2 71·6 80·3 86·2 96·9 105·8 113·5 122·2 | lbs. *54·2 *61·1 76·1 86·1 95·0 107·0 117·2 126·6 130 0 | 1bs. 70·3 74·2 81·2 85·4 95·0 105·4 117·2 130·2 138·6 | | | |

^{*} These values are too low, due probably to some error of observation. Mr. R. Clark, who furnishes the returns, is unable to account for the discrepancies in these year

CONCLUSION.

67. Attention has been called to some of the principal points of interest in the data collected by the Committee, but in many respects the tables have been left to speak for themselves; and it is not improbable that a study of them will lead some persons to conclusions differing more or less from those given in this Report.¹

68. The original returns, which the Committee recommend may be placed in the charge of the Anthropological Institute for preservation and future examination, comprise many statistics which could not be introduced into this Report on account of the time and labour required

for their analysis and tabulation.

69. The Committee believes that it has laid a substantial foundation for a further and more exhaustive study of the physical condition of a people by anthropometric methods, and that its action will prove it has been useful as an example to other scientific societies and to individuals in stimulating them, as well as directing them, in the methods of making statistical inquiries relative to social questions. The medical officers, managers, or superintendents of many colleges, schools, and charitable institutions have been induced to keep registers of the physical proportions of those under their charge, which will in a few years become valuable records, not only of the physical condition of the inmates of their institutions, but of the sanitary conditions under which they have lived; they will also be available for the further study of the subjects specially treated of in this Report. The Collective Investigation Committee of the British Medical Association propose to carry on the work of this Committee in a direction which it is most needed, namely, by issuing an album in which persons may methodically record at frequent intervals their height, weight, and other physical qualities, together with points in their personal and medical history. The Committee hopes that this habit will be largely adopted and encouraged by the members of the British Association.

70. The Committee has to express its thanks to the numerous contributors to their store of facts, whose names and contributions have been published from time to time in their interim reports, and to numerous friends who, although not contributors themselves, have induced others

to give their assistance.

The statistics relating to eyesight were dealt with in the Report for 1881, and the returns since received are not sufficient to require a further discussion of the subject.

The subject of colour-blindness was taken up by a Special Committee of the Ophthalmological Society after it had been inaugurated by this Committee, and it was given up on that account. The very interesting report of the Special Committee is published in the first volume of the Trans. of the Ophthal. Soc. 1881.

¹ The inquiries relative to *breathing capacity* were abandoned in 1879 on account of the unsatisfactory nature of the returns received previous to that year. The apparatus were faulty.

APPENDIX A.

Specimen of the cards used by the committee for collecting observations, and the instructions for filling them up. The cards are of different colours for the two sexes, and one corner is cut off to make them face one way when arranged by hand. They can be dealt out like playing cards, and much time and trouble is saved in the analysis of their records.

ANTHROPOMETRIC COMMITTEE OF THE BRITISH ASSOCIATION, 22 Albemarle Street, London

(to which address this Card is to be returned after being filled).

| Height is to be taken as without shoes, and weight in ordinary indoor costume. |
|---|
| Span of Arms is the distance between the tips of the middle fingers extended hori- |
| zontally, measured across the back (i.e. back to the wall). |
| Colour of Eyes should be stated as grey, light blue, blue, dark blue, light brown, brown, |
| dark brown, green, or black. |
| Colour of Hair as very fair, fair, golden, red, red brown, light brown, brown, dark |
| brown, black brown, or black. |
| For chest-girth, breathing capacity, strength, colour-blindness, and eyesight, see the |
| paper of instructions. |
| Under Place of Birth state Parish and County; or, if abroad, the name of the Country. |
| Under Occupation state rank or profession. |
| Race should be stated as English, pure English, very pure English, Irish, pure Irish, very pure Irish, Scotch, pure Scotch, very pure Scotch, or mixed Scotch |
| and English, &c. |
| Origin, as countryfolk, pure countryfolk, very pure countryfolk, townfolk, pure townfolk, country birth, T. since boy, &c. |
| |


FOR A SINGLE SET OF OBSERVATIONS.

| Place | Date188 |
|--|--------------------------------|
| Name (or Initials) | Sex |
| Age—years months | |
| Height, without shoes, inches & eighths | Span of arms, inches & eighths |
| Weight, in ordinary indoor costume, lbs. | Strength, drawing power, lbs. |
| Chest-girth, inches and eighths | |
| Colour of Eyes | |
| | Colour-blindness_ |
| Sight Test-types No. 1, read at inches | |
| " No. 10, " feet | Astigmatism |
| Place of Birth { | Occupation |
| Race | Occupation Origin |
| Name and Address of Observer | |

Girth of Chest.—This is the method adopted in the British Army. Make the person stand quite upright, with his shoulders back, and his arms hanging loosely by his side. The measurement must be taken next to the skin, without compressing it. The lower edge of the tape should touch the nipples, and the measurement should be read off in front. Care should be taken that the tape passes horizontally round the chest, because if the measurement is made obliquely, below the blade-bone, it will be erroneous. The person should be required to count ten slowly during the operation, to prevent him from keeping his lungs over-inflated. (If this measurement is made on females, it should be taken below the breasts.)

Strength of Arm.—It is proposed to measure the force that can be exerted by the arm when pulling (as an archer with a bow). A spring balance should be used for this purpose. The right or left arm, whichever is the strongest, should be used to draw, and the other to resist. The resisting arm must be free, and extended straight from the side, as nearly as possible in the line of the shoulders, and the hand of the other arm brought back towards the ear. (A spring balance, or 'arm-testing machine' for testing the drawing power, can be obtained of Herbert &

Sons, 6 West Smithfield, London, E.C., price 18s. 6d.)



The above figure represents the position in which the strength of arm should be tested.

APPENDIX B.

TABLE XXVI., showing the STATURE, CHEST-GIRTH, and WEIGHT of Recruits, is introduced here for future reference and comparison. The figures show that recruits of the age of 18 years may be expected to increase 1 inch in stature, 1½ inch in chest-girth, and 10 lbs. in weight, before they reach the age of 23 years.

Table XXVI.—Stature (barefoot) of Recruits for the Army, 1860-4.

| Height | Age last Birthday | | | | | | | | | | | | |
|-----------------------|-------------------|--------|--------|--------|--------|-------|-------|-------|-------|--|--|--|--|
| without shoes. Inches | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | | | | |
| 72 and upwards | 2 | 19 | 55 | 52 | 52 | 46 | 49 | 59 | 120 | | | | |
| 71- | 2 | 71 | 123 | 113 | 129 | 101 | 102 | 124 | 240 | | | | |
| 70- | 3 | 205 | 259 | 280 | 276 | 261 | 199 | 253 | 527 | | | | |
| 69- | 21 | 519 | 555 | 559 | 508 | 488 | 400 | 455 | 747 | | | | |
| 68- | 67 | 1172 | - 1139 | 988 | 835 | 756 | 609 | 746 | 1135 | | | | |
| 67- | 219 | 2995 | 2159 | 1706 | 1268 | 1108 | 877 | 964 | 1425 | | | | |
| 66- | 871 | 5593 | -3277- | -2292- | -1428- | | 964 | 1019 | 1349 | | | | |
| 65 | _1224 | 5009 | 2504 | 1814 | 1144 | 881 | 608 | 567 | 996 | | | | |
| 64- | 758 | 3968 | 1344 | 1172 | 718 | 603 | 878 | 421 | 850 | | | | |
| 63- | 386 | 534 | 232 | 358 | 123 | 105 | 63 | 65 | 134 | | | | |
| and under 62 | 135 | 78 | 25 | 26 | 17 | 9 | 7 | 7 | 12 | | | | |
| Total | 3683 | 20,163 | 11,672 | 9360 | 6493 | 5667 | 4251 | 4680 | 7537 | | | | |
| Mean | 65.50 | 66-00 | 66.25 | 66.50 | 66.75 | 67:00 | 67.00 | 67:00 | 67.00 | | | | |

CHEST-GIRTH (empty) of Recruits for the Army Anthropometric Committee.

| Chest-girth, empty. | | | | Age | last Birtl | nday | | | |
|---|--|--|---|--|---|--|---|--|--|
| Inches | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 |
| 43- 42- 41- 40- 39- 38- 37- 36- 35- 34- 33- 32- 31- 30- 29- | - - - 2 - 3 10 -26 - 9 7 2 1 | 3 8 8 37 74 155 166 55 11 5 | - - 2 4 12 70 123 -173- 131 37 9 - | 5 9 13 51 80 -128 63 14 2 1 | - 1 9 19 46 51 79 28 1 - | 2 8 14 32 63 89 20 4 — | 3 9 18 24 38 53 11 2 | 1 1 1 4 13 22 45 45 47 13 1 — | 2 2 2 2 5 16 31 41 44 16 3 - 1 |
| Total | 60 | 516 | 561 | 361 | 230 | 182 | 138 | 191 | 164 |
| Mean | 33.5 | 34.0 | 34.5 | 84.75 | 85.0 | 35.5 | 35.5 | 85.5 | 85.5 |

Weight (naked) of Recruits for the Army, 1860-4.

| Weight without clothes. | | Age last Birthday | | | | | | | | | | | | |
|---|---|--|---|---|--|---|--|--|---|--|--|--|--|--|
| lbs. | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | | | | | |
| 170- 160- 150- 140- 130- 120- 110- 100- under 100 | 4 25 75 338 1345 2724 8494 1404 146 | 39 202 871 3674 9965 -18,196 13,912 2734 282 | 69 331 1228 4055 8881 —11,765 5961 985 50 | 101 441 1896 3950 7128 7497 2937 374 19 | 116 472 1409 3411 5078 4391 1695 151 5 | 145 489 1369 3024 -3981 3351 1191 116 2 | 160 484 1199 2587 -3153- 2206 761 50 1 | 177 - 528 1317 2497 -2914- 2266 757 70 1 | 180 489 1218 2290 -2590- 2132 751 107 8 | | | | | |
| Total | 9555 | 49,875 | 83,825 | 23,843 | 16,723 | 13,672 | 10,559 | 10,527 | 9760 | | | | | |
| Mean | 120-0 | 125.0 | 125.0 | 130.0 | 185.0 | 185.0 | 185.0 | 135.0 | 135.0 | | | | | |

APPENDIX C.

Index to the Tables in the several Reports of the Committee, showing the nature of the measurements given in each Table.

In 1879.

Several selected classes; males at each |

Christ's Hospital School; males at each

British Race in England and America, and Belgians; males and females, at each age.

Recruits, British and American armies, at each age.

Stature, weight, and ratio of weight to height.

Stature, weight, chest-girth, and relation to one another, by Sir Rawson Rawson. Stature and weight, with diagrams, by C.

Roberts.

Stature and weight, by C. Roberts.

In 1880.

Schoolboys of several classes, of age 11 |

Standard class; males of ages 10 to 50.

Standard class; males of ages 10 to 50.

Standard class; males of ages 10 to 50. Professional classes; males of ages 10 to 50 Persons of town and country origin; males at each age.

American boys and girls.

Factory children; boys and girls, 1833, 1871 - 3.

Marlborough College; males at each age.

Telegraph messengers; youthsat each age.

Stature, by C. Roberts.

Stature, weight, chest-girth, and strength of arm, with diagram.

Relation of the several measurements to one another.

Mean annual growth.

Colour of eyes and hair, with diagram.

Stature and weight.

Stature and annual growth, with diagrams, by Prof. Bowditch and Sir Rawson Rawson.

Stature and weight, by C. Roberts.

Stature, weight, chest-girth, girth of head, arm, and leg, by the Rev. T. A. Preston, Sir Rawson Rawson and C. Roberts.

Weight, chest-girth, and lifting power, by G. C. Steet.

In 1881.

General population of United Kingdom; | Increase in stature, weight, chest-girth, males at each age.

General population of United Kingdom; males at each age.

Population of different classes; males at each age.

Population of different classes; males from 25 to 50.

Population of different classes; males at each age.

Population of different classes; males at different ages.

Marlborough College; boys at each age.

and strength of arm, with diagram.

Stature, weight, chest-girth, and strength of arm.

Stature and weight.

Relative stature.

On calculation of deciles, quartiles and medians applied to range of stature, weight, and strength of arm, by F. Galton.

On army test of eyesight in each class, with diagram, by Inspector-Gen. Lawson.

On Snellen's tests for eyesight, near and distant vision, and colour-blindness, by the Rev. T. A. Preston and C. Roberts.

In 1883.

 General population of each part of United Kingdom; adult males.

General population; adult males and females.

3. Population of counties; adult males.

4. Population of counties; adult males.

5. Population of several countries, Europe and America; adult males.

 Population of several races and nationalities; adult males.

 Selected classes (British); adult males.

 Criminals and lunatics (British) compared with other classes; adult males.

 Criminals and lunatics (British) compared with other classes; adult males.

 Population of counties of United Kingdom; adult males.

11. Population of English and Welsh origin; males and females at each age.

12. Classification of population according to media.

13. Schoolboys of several classes, of age 11 to 12.

 Population of several classes; males from 25 to 30.

15. Infants (at birth); males and females

 Population of several classes; males at each age.

17. Population of several classes; females at each age.

18. Population of several classes; males at each age.

 Population of several classes; females at each age.

General population; males and females at each age.

21. Industrial Schools; males and females at each age.

 Industrial Schools; males and females at each age.

23. Swinton Industrial School; males and females at each age.

24. Factory children, 1833-73; males and females at each age.

 York Friends' School, 1853-79; males at each age.

26. Recruits (British army), 1860-64; ages 17 to 25,

Stature, weight, chest-girth, and strength.

Relative stature, weight and strength.

Stature, weight, and complexion, with diagram and five maps.

Stature: ratio per 1,000.

Stature: average, medium, and extreme.

Stature.

Stature and weight.

Stature and weight.

Complexion: colour of eyes and hair.

Complexion: degree of nigrescence.

Complexion.

Nurture, occupations, and sanitary surroundings.

Stature (same Table as in 1880).

Relative stature (same Table as in 1881).

Height, length, and weight. Stature.

Stature.

Weight.

Weight.

Stature, weight, chest-girth, strength, and span of arm; relation to each other, and between the sexes.

Stature, weight, chest-girth, and span of arms.

Complexion.

Stature, weight, chest-girth, breathing capacity, strength of arm, sight, and complexion.

Stature and weight.

Stature and weight.

Stature, weight, and chest-girth.

- List of recent Monographs on the subject of Anthropometry published in England and the United States.
- Gould, B.A. . Investigations in the Military and Anthropological Statistics of American Soldiers. United States Sanitary Commission Memoirs, New York, 1869.
- Beddoe, J. (M.D.) . On the Stature and Bulk of Men in the British Isles. Mem. Anthrop. Soc. vol. iii., London, 1869.
 - Notes and Queries on Anthropology for the use of travellers and residents in uncivilised lands. Drawn up by a Committee appointed by the Brit. Assoc., 1874.
- Fergus, Dr. W., Rodwell, G. F., and Preston, Rev. T. A.

 A Series of Measurements made at Marlborough College.

 Jour. Anthrop. Inst., 1874.—A continuation of these measurements, together with observations on eyesight and colour-blindness, made annually to the present time by The Rev. T. A. Preston, in the Report of the Marlborough College Natural History Society.
- Galton, F. . On the Height and Weight of Boys, aged 14 years, in town and country Public Schools. Jour. Anthrop. Inst., 1875.
- Human Faculty, London, 1883. Contains a List of Papers on Anthropometric subjects contributed to various scientific journals and literary magazines by the author.
- Baxter, J. H. (M.D.) Statistics, Medical and Anthropological. Report of the Provost-Marshal-General's Bureau, U.S. Government, Washington, 1875.
- Roberts, C. . The Physical Development and Proportions of the Human Body. St. George's Hospital Reports, 1874-6.
 - ,, . . The Physical Requirements of Factory Children. Jour. Statistical Soc., 1876.
 - " . . A Manual of Anthropometry. London, 1878.
 - " . . The Detection of Colour-blindness and Imperfect Eyesight, drawn up for the use of the Anthropometric Committee.

 London (Bogue, St. Martin's Place), 1880.
- Bowditch, H. P. (M.D.) The Growth of Children. Eighth Annual Report State Board of Health, Mass., U.S., Boston, 1877.
- Peckham, G. W. (M.D.) Milwaukee, The Growth of Children. Sixth Annual Report of the State Board of Health, Wis., U.S., 1882.