

Ovariectomy : removal of an encysted tumour of the left uterine appendages / by George Southam.

Contributors

Southam, George, 1815-1876.
Royal College of Surgeons of England

Publication/Creation

Salford : Printed by William F. Jackson, 1845.

Persistent URL

<https://wellcomecollection.org/works/b5et8r9x>

Provider

Royal College of Surgeons

License and attribution

This material has been provided by This material has been provided by The Royal College of Surgeons of England. The original may be consulted at The Royal College of Surgeons of England. where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>





OVARIOTOMY.

22

18

REMOVAL

OF AN

ENCYSTED TUMOUR


OF THE

LEFT UTERINE APPENDAGES.

BY GEORGE SOUTHAM,


PHYSICIAN TO THE SALFORD ROYAL HOSPITAL AND DISPENSARY, MANCHESTER.

PRESENTED AT THE ANNIVERSARY MEETING OF THE PROVINCIAL MEDICAL AND SURGICAL ASSOCIATION, AT SHEFFIELD, THURSDAY, JULY 31, 1845.



SALFORD.

PRINTED BY WILLIAM F. JACKSON, NEW BAILEY STREET.
1845.



OVARIOTOMY.*

ly in June, Mrs. S., aged 38, called upon me and stated that she was suffering from an ovarian tumour, which having resisted a variety of remedial means, she was extremely desirous of having it removed. The following is a brief sketch of her history:—

She has been married twenty years, never had any family, and within the last few years has generally enjoyed good health. She first noticed an increase in the size of her abdomen eight years ago, but it was not attended with any derangement in her health, or irregularity in the menstrual function, it was attributed to corpulence. Some time having elapsed, and there being evident deterioration in her condition, medical aid was requested. Since then she has been under the care of several experienced practitioners, and various remedies have been resorted to, all of which, with the exception perhaps of retarding the progress of the disease, have proved ineffectual. During the last twelve months the tumour has increased to such a size as to interfere with respiration, especially when she is in the recumbent posture, and she has occasionally suffered from pain in the left inguinal region. Tapping has been frequently recommended to relieve these symptoms, but she has hitherto refused to con-
cede to it. No apparent reason can be given for the disease, as it originated from a blow on the abdomen when about 16 years of age; nor is she aware that the tumour began in either side, having noticed it until a uniform swelling above the pubes presented itself.

On examining the abdomen I found it nearly globular, and very prominent, distended to at least twice the size of a female at the full term of pregnancy, and elevating the cartilages of the ribs to a considerable extent; the parietes were perfectly smooth and natural colour; fluctuation, though of a resisting kind, was very distinct every part, and percussion elicited a dull sound. Change of position produced scarcely any alteration in its form. The uterus, examined *per vaginam et rectum*, seemed to be of the natural size, and was situated higher in the pelvis than usual; the os was quite healthy, and on the abdomen being raised and depressed by an assistant, whilst the finger was placed in immediate contact with it, the uterus might be distinctly felt to rise and fall with an inclination to the left side. In the erect position it could not be made to

* Reprinted from the Provincial Medical and Surgical Journal.

bound away from the finger. There was no protrusion of vaginal parietes, but an elastic swelling could be easily perceived by pressing on the left side of the uterus and upper part of the vagina. In other respects the pelvic cavity was not encroached upon. Though somewhat emaciated, her general health did not appear impaired; appetite good; tongue clean; pulse natural; bowels occasionally constipated, but easily acted on by medicine; catamenia regular though in less quantity than formerly; felt pain at times in the inguinal region, or whenever she lay on the left side, and in consequence of the pressure of the tumour upwards, she has been unable to lie on her back, or in any other position, excepting on the right side, for the last twelve months.

The symptoms clearly indicating encysted dropsy of the uterine appendages, advanced to such a stage as to render surgical interference necessary, and her constitution not being much affected, I considered the case was peculiarly favourable for the operation. At the same time, in giving this opinion, I candidly stated that tapping, which was now absolutely required, though merely palliative, was attended with less risk, and would probably prolong her life in tolerable comfort for some years. Extirpation was, however, the only effectual means of radically removing the disease, but much more dangerous, two out of five of those who had submitted to it having died. Being possessed of good moral courage, and having witnessed the complete restoration to health of others who had submitted to the operation, she stated her willingness to undergo it notwithstanding its concomitant dangers. Considering it would be more satisfactory to her friends to have another opinion, I requested Dr. Radford to accompany me in my next visit, who, after a careful and conducted examination, coincided with my views.

Extirpation being determined upon, it was fixed to take place on Tuesday, the 24th of June, and her residence not being in my immediate vicinity, I advised her to take lodgings near me. The preliminary arrangements having been entered into, such as abstinence from animal food and stimulating liquors, for a few days, the free evacuation of the bowels with castor oil, the temperature of the room raised to 75° Fahrenheit, and the bladder previously emptied, I commenced the operation at four o'clock in the afternoon, in the presence of Drs. Radford and Clay, Dr. Watson, of Liverpool, Messrs. Robertson, Nursaw, Horn, and Winterbottom, surgeons, and Mr. Williamson, my pupil, all of whom promptly rendered whatever assistance was necessary. An exploratory incision, midway between the umbilicus and pubes, was first made, and the peritoneal cavity opened sufficiently to admit the finger. A characteristic membrane of a bluish white and shining surface appearing at the opening at once satisfied me of the existence of a cyst, and the finger introduced between it and the peritoneum, discovering no adhesions in the immediate neighbourhood, I punctured it with a full-sized trocar.

from sixteen to eighteen pints of clear, lemon-coloured, mucilaginous fluid had been evacuated, pressure on the being well sustained during its escape, the canula was withdrawn and using the index finger as a director, the opening was made above and below with a probe-pointed bistoury to the depth of between six and seven inches. Having ascertained by the introduction into the abdominal cavity, that there were no impediments to the extraction of the tumour, it was carefully drawn out, pressure being continued on the abdomen. Finding it was attached to the uterine extremity of the left broad ligament by a small and slightly vascular pedicle, I tied it firmly with a single thread of the strongest dentist's silk. The pedicle was now cut and the tumour being removed, the margins of the wound were immediately approximated to prevent the ingress of air. After a short interval, the wound was again opened to remove what blood had coagulated internally, and to ascertain that the vessels of the broad ligament were firmly secured. The uterus and opposite ovary were examined and found healthy. One end of the ligature being secured to the other was left dependent at the lowest point of the wound, the edges of which were brought together by four interrupted sutures and straps of transparent tissue plaster. Upon these a broad bandage was applied, and the whole being adjusted by a bandage, the patient was lifted into bed, within twenty-five minutes from the commencement of the operation. During the whole time she sustained herself remarkably well, apparently not suffering much pain, whilst the exploratory incision was made, during which some hæmorrhage occurred, from the parietes being freely supplied with vascular tissue. No difficulty was experienced from the intestines being relaxed, as they were remarkably flaccid. There was very little hæmorrhage and no vomiting, but a feeling of faintness was superinduced by the evacuation of the fluid.

Being placed in bed she complained of pain in the left iliac and back; pulse firm, 88. I ordered a grain of the acetate of opium in the form of pill; toast-water in small quantities when thirsty, and the temperature to be kept at 70° Fahrenheit.

At past seven P.M. Pulse soft, 108; slept three-quarters of an hour after the pill; pain in the side continues; with the exception of the retching after taking the pill, there has not been the least disposition to vomit. Says she is both hungry and thirsty. Diet restricted to toast-water.

3 A.M. Has had some pain along the course of the wound on the left side during the last hour; pressure on the abdomen increases it; complains of a feeling of distention in the bowels, without desire to pass flatus, which she is unable to do. Skin hot; no vomiting; tongue assuming a morbid redness; has not seen the last visit. As the pulse was fuller and more frequent, bled fourteen ounces of blood from the arm, which had the effect of bringing it twelve pulsations, and causing faintness. The night

being warm the temperature of the room to be reduced to 67°, and two-thirds of a grain of morphia administered.

Half-past 8 A.M. Parted with flatus frequently during the last three hours, which has relieved the pain in the abdomen. Pulse 116 soft; skin moist, and of the ordinary temperature; passed about twelve ounces of urine since last visit; still thirsty and feels hungry appears to be proceeding very favourably.

11 A.M. Visited with Dr. Radford, and found her in the same comfortable state as when last seen; pulse 108; temperature 68°. To have barley water, with half an ounce of gum dissolved in each pint.

5 P.M. Vomited once since last visit, which she attributed to some tea she had taken; tongue moist, yet redder than natural countenance cheerful, but features slightly contracted; pulse 116 passed eight ounces of urine; thirst continues. To be allowed wine-glassful of soda water occasionally, and a biscuit.

11 P.M. No return of sickness; pulse 106, soft; troubled with flatulence; expresses herself as feeling very comfortable; slept almost the whole of the time since last visit. The bandage being rather tight it was slackened.

26th, 7 A.M. Had a good night, having slept several hours; features still rather contracted, but cheerful; pulse soft, 100; tongue red, rather dry, and furred down the sides; skin moist; thirsty; passed a pint of urine during the night; has taken half a biscuit this morning. Bowels not having been moved since the operation to take half an ounce of castor oil.

11 A.M. Visited with Dr. Radford; the oil having produced no alvine evacuation to have a common enema. The bandage and pad were removed, and the wound, as seen through the transparent plaster, appeared to be healing.

10 P.M. The enema produced a copious evacuation, but previously had severe pains in the abdomen, which has returned at intervals during the afternoon. It appears to be caused by flatulence. Features still contracted, but countenance cheerful; pulse 104, soft; tongue moister, still red and slightly furred; passed urine twice since last visit, which was rather high coloured and turbid; less thirst. To take ten grains of carbonate of soda in aniseed water every two hours, whilst the flatulence continues.

27th, 7 A.M. Passed a comfortable night, the draught having entirely relieved the flatulence; complains of a short cough, attended with expectoration; tongue moist and less red; pulse 100, soft; bowels moved about an hour ago; passed about eight ounces of high coloured urine through the night.

11 A.M. Visited with Dr. Radford; symptoms the same; the cough and expectoration continuing, ordered a mustard poultice to the chest; the day being cold and wet, to increase the temperature of the room to 70°.

Pulv. Ipecac. Co. gr. iij. tert. hor. capiend.

Diet to consist of barley water, biscuit, and tea.

M. No change in the symptoms; examined the wound, looks remarkably healthy; cough continuing, only one ligature removed. To repeat the mustard poultice; and as she is to have barley water, to have sago.

M. Cough quite relieved since the last poultice, and she is very comfortable; skin moist and warm; countenance cheerful; pulse 106; enjoyed her sago. To omit the pills.

28th, 8 A.M. Passed an excellent night, and feels comparatively well; tongue moist and natural in colour: pulse soft, 96; cough; skin of natural temperature and moist; no thirst; passed twelve ounces of urine during the night, which is clear, but high coloured.

M. Visited with Dr. Radford. The same. To be allowed real broth.

M. Had a very comfortable day; feels well enough to sit up; features no longer contracted; pulse 94, soft; passed urine since last visit, but had no evacuation.

Ol. Ricini. ozss. mane. cap.

29th, 10 A.M. Visited with Dr. Radford. Improving. The oil acted once upon the bowels; pulse 92; tongue quite healthy in appearance. The wound being quite healed, except at the lowest part, the three remaining ligatures were removed.

M. The cough returned during the afternoon, but was immediately relieved by a mustard poultice and two pills (pulv. ipecac. co.); pulse 90, soft; bowels moved this afternoon; no flatulence.

30th, 11 A.M. Visited with Dr. Radford, and found her progressing very favourably; passed an excellent night; feels hungry; thinks she ought to be allowed to sit up to-day. Bowels moved last night.

31st. Passed a very comfortable night; pulse 88; was out of bed this morning for the first time; felt very mazy, and had difficulty in balancing herself. To be allowed some chicken broth for dinner, and bread *ad libitum*.

M. The cough returned last night, but was soon relieved by a poultice and the pills; did not sleep very well, owing to a pain in the epigastrium, which she attributed to having eaten too freely of bread; pulse 90; other symptoms those of health. Diet to consist of real broth, fowl, sago, and tea.

M. Continues to improve; pulse 84; bowels moved daily without medicine; appetite good; appears to be suffering only from weakness. To have a mutton chop for dinner, and to sit up for a short time.

M. Gains strength daily; feels the want of some support to her chest when sitting up, although she is very tightly bandaged; is usually wanting food.

M. Doing well; can walk across the room; catamenia appeared last day, but in small quantity.

12th. returned home, a distance of three miles, and bore the journey very well.

I have visited her at intervals up to the present period, and found her, on each occasion, much improved. The ligature* came away on the forty-ninth day, after which the fistulous opening closed. The cicatrix of the entire wound does not measure more than four inches. She has gained flesh, and is now in the enjoyment of perfect health.

DESCRIPTION OF THE DISEASE.

The tumour consisted of one entire cyst, which, when distended with fluid, resembled the shape of an egg, and weighed thirty-one pounds; the larger end occupied the lower portion of the abdomen, and was highly vascular; the smaller one filled the epigastrium, presented no trace of vessels, and was so exceedingly diaphanous, that the colour of the fluid might be clearly distinguished through its parietes. The disease had evidently originated in the fold of the broad ligament, the ovary being merely adherent to it. The structure of the cyst varied in different parts, being half an inch in thickness near its connexion with the broad ligament, and gradually becoming thinner towards the upper portion, where it appeared scarcely capable of sustaining the weight of the contained fluid, and in all probability would have been ruptured with a very slight blow. The interior presented all the characters of a secreting surface, and the whole of the larger extremity was studded with numerous little nodules, about the size of millet seeds, some in clusters, others solitary. Several appeared to have the consistence of cartilage, others were composed of calcareous matter. A small cyst was also found in the ovary, which, in other respects, presented no abnormal characters.

It would be infringing on the objects of this meeting were I to make any lengthened remarks on the numerous disputed points which ovariectomy involves, I shall not therefore trespass on your time with any other observations than what the case calls for.

* On referring to the report of my former patient, (*Medical Gazette*, November, 1843,) it will be found that the tympanitis and obstinate vomiting which supervened on the operation, caused the ends of the ligatures to be drawn within the abdominal cavity; the wound had perfectly healed, and the patient was restored to health, whilst they were still in the abdomen. After several weeks had elapsed, a small abscess appeared at the lower part of the cicatrix, an opening into which gave exit to a quantity of healthy pus and the ligatures. A free discharge was promoted by poultices for a few days, and at the end of a week the wound closed. No constitutional disturbance occurred, and there has not been the slightest interruption to the most perfect state of health since the termination of the report. The catamenia appear with the greatest regularity, and in the same quantity as previously to the commencement of the disease. There is a tendency to corpulency, which is in a great measure checked by her active habits.

culty in the diagnosis of ovarian tumours is perhaps the most
ant objection that can be urged against the operation. This
doubt continue so long as the mere history of the disease
abdominal examination are regarded as sufficient to detect its
ce. In the case before us, although it had been of many
uration, and several experienced practitioners had been con-
no vaginal examination had been made or proposed. In its
stage a circumscribed tumour might probably have been
distinguished through the abdominal parietes, but when the
applied to me the enlargement was so uniform and symme-
accompanied with such distinct fluctuation in every part, that
impossible to say with any degree of certainty whether the
as encysted.

examination *per vaginam et rectum*, cleared up all doubts, not
of the existence of an ovarian tumour, but of its exact nature.
absence of any protrusion of the vaginal parietes, the elevated
n of the uterus in the pelvic cavity, and the inability to cause
ound away from the finger, proved that the fluid could not be
; whilst the projection of the swelling on the left side, the
d influence which raising and depressing the abdomen pro-
on the position of the uterus, with its inclination to one side
than the other, evidently indicated that the disease was con-
with the left uterine appendages. The uniformity in the
inal distension, the distinct fluctuation in every part of it, and
ability to discover any solid matter encroaching on the pelvic
led me to infer that it consisted of one or two cysts only;
the comparative immunity from any great degree of suffering
within the last twelve or eighteen months, the absence in the
y of any previous attacks of peritonitis, together with the
impediment to the action of the bowels, rendered the exist-
f adhesions doubtful.

the operation, the evils of both the major and minor incisions
guarded against, by making the opening no larger than was
ary to ascertain the nature and connexions of the disease, and
nit of its removal after reducing it by paracentesis, without
oning the least violence or displacement to any of the neigh-
g parts. Had the slightest obstacle occurred previous to the
sful termination of the operation, this plan would also have
d me to have receded without much danger to the patient.
stage of it, however, fully confirmed the accuracy of the
osis, and the trifling interference required in the subsequent
ent contributes to make this one of the most successful cases
orded. Medicine having failed in checking the disease, its
erence with respiration, and embarassment to the patient in
ing her ordinary occupations, pointed out the necessity for
al aid. The inadequacy of tapping, save for temporary relief,
avourable nature of the disease, and the healthy condition of
atient, induced me to undertake its extirpation, which I should

hardly have sanctioned at an earlier period ; for in the present state of our experience, the mere existence of an ovarian tumour should not lead us to decide on its removal, until the inconveniences have become so great as to render surgical interference indispensable. At the same time it would be highly improper to delay its performance until the patient's constitution has been invaded, or some important organ implicated. Where the diagnosis is not sufficiently evident, as a preliminary measure, recourse may be had to tapping, which will frequently clear up any uncertainties that may exist, or if the system has become enfeebled from the pressure and weight of the accumulated fluid, it will afford time for its renovation.

From the tumour presenting that form of ovarian disease, for which tapping is the most beneficial, it will perhaps be urged, that the patient ought not to have undergone the risks of the more formidable operation. Life is no doubt often prolonged by paracentesis. But how frequently is the brief interval of relief thus afforded counterbalanced by the increased distress attending its repetition, which, from producing a continual drain on the system, sooner or later shatters the constitution, and after protracting the individual sufferings, terminates in death.

The first object of all surgical operations is of course the preservation of life. If this can be gained more effectually by tapping notwithstanding its disadvantages, it ought to be preferred to extirpation. Looking merely at the average mortality of the cases in which the latter has been undertaken, the risk certainly appears very great. Since, however, in addition to every variety of disease from the simple encysted dropsy, to the most malignant form of carcinoma, it comprises all those in which the operation has been rashly and unnecessarily performed, it cannot be considered applicable to particular cases. Indeed, a perusal of them plainly shows that the encysted form is not only the most favourable for extirpation, but where the cases have been selected in accordance with the rules which direct us as to the propriety of other operations, and no unnecessary violence has been used in removing the tumours, the mortality has been less than that of major operations generally.

The indifference evinced by many to investigate the researches of others, no doubt forms a strong incentive to test the value of a remedy by statistics ; but when this mechanical system of reflecting is allowed to usurp the power of reason or of knowledge, acquired by experience and observation, or "confines itself to counting facts without interpreting them,"* the principles founded on it will be liable to error.

The unqualified opposition to ovariectomy, formed chiefly from statistical data, by some eminent individuals, appears singularly inconsistent since they do not hesitate to recommend operations still more dangerous for the removal of diseases which are also no

* M. Combe. De la Médecine en France, et en Italie. Paris, 1842.

ately destructive to life; I allude to the application of a
to the innominata, and other large arteries, for aneurism,
mer of which, though performed several times by the ablest
as, has, I believe, invariably failed.

In medical education was less general, the weight of a great
might have influenced the estimation of any new doctrine
profession, but the mind of every practitioner is now so well
with information, that the opinions of no man, however
ished, are likely to be received with any favour, unless
l on a proper investigation of the facts which the subject
es.

iotomy, combining as it does so many requisites for its suc-
prosecution, is not calculated to find general favour in the
so divided a profession as the medical.

ie skilful treatment of the physician, must be superadded
t and promptitude in emergencies of the operative surgeon,
as the experience of the accoucheur in uterine and abdominal

. In proportion as the operator is deficient in any of these
attainments, he is liable to incur the blame of rashness or
nity, where, under more favourable auspices, he might have
high encomiums for his efforts in arresting protracted suffer-
disease. The apparent facility of the operation may induce
to undertake it who are not qualified to form a correct
is of the cases in which it is most likely to be beneficial.
Manual dexterity, unaccompanied by a careful consideration
circumstances of the case, and the condition of the patient,
an lead to the establishment of ovariectomy on a firm founda-
On the contrary, the rash and indiscriminate use of the
ay have a tendency to throw discredit on what might other-
ve a valuable resource for eradicating a disease, which, to
t interesting portion of the human race, has hitherto formed
s obstacle to health and the enjoyment of life. Where the
of the operation is followed by such satisfactory results,
egree of risk may justifiably be incurred on the principle
ice of the father of medicine,

“Ad extremos morbos extrema remedia.”

SCENT, SALFORD, AUGUST, 1845.

