

**An account of a particular change of structure in the human ovarium, by Matthew Baillie, M.D. : vide Philosophical Transactions of the Royal Society of London, Vol. LXXIX, for the year 1789, 4to, London.**

**Contributors**

Royal College of Surgeons of England

**Publication/Creation**

[Edinburgh] : [publisher not identified], 1790.

**Persistent URL**

<https://wellcomecollection.org/works/bv9kka6>

**Provider**

Royal College of Surgeons

**License and attribution**

This material has been provided by This material has been provided by The Royal College of Surgeons of England. The original may be consulted at The Royal College of Surgeons of England. where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>


# THE HISTORY OF THE

... of the ...  
... of the ...  
... of the ...  
... of the ...

... of the ...  
... of the ...  
... of the ...  
... of the ...

... of the ...  
... of the ...  
... of the ...  
... of the ...

... of the ...  
... of the ...  
... of the ...  
... of the ...





XX.

*An Account of a particular Change of Structure in the Human Ovarium.* By Matthew Baillie, M. D. Vide *Philosophical Transactions of the Royal Society of London, Vol. LXXIX. for the year 1789.* 4to, London.

THE ovaria in women, Dr Baillie observes, are subject to a great variety of changes from their natural structure. Many of these are exactly similar to what take place in other parts of the body; but there is one change, which he considers as peculiar to them, and the nature of which, he thinks, has not hitherto been well ascertained. This change of structure is a conversion of the natural substance of an ovarium into a fatty mass, intermixed with hair and teeth.

We have already, in a former part of this volume, presented our readers with a singular history



history of this kind, and with the observations and sentiments of an ingenious anatomist respecting it; and we have here before us another case of the same kind, which, though varying in several particulars, is, in its circumstances, not very dissimilar. This case, which occurred to another distinguished anatomical teacher, is here communicated to the public, with those reflections which it suggested to him.

Though many cases of this kind are to be met with in books of dissections, yet few have ventured to offer any conjectures on the mode of their formation. But, in general, they have, Dr Baillie observes, been considered as imperfect attempts at the growth of a foetus in the ovarium, in consequence of the connexion between a male and a female. This conjecture rests, he admits, on circumstances of strong probability; but he thinks it is liable to many powerful objections. And, among other particulars, from the following case, he is led to believe, that the ovaria in women have some power within themselves, of taking on a process which is imitative of generation, without any previous connexion with a male.

In



In a female child about twelve or thirteen years old, which was lately brought to Windmill Street for dissection, Dr Baillie found the right ovarium converted into a substance, doughy to the touch, and about the size of a large hen's egg. Upon cutting into this substance, he found an apparently fatty mass, intermixed with hair and an excrescence of bones. This startled him very much, as he had always been led to believe, that such appearances were a sort of imperfect conception; and, the circumstances altogether being very singular, he was led to pay particular attention to them.

The fatty mass was of a yellowish white colour, in some places more yellow than in others; it was very unctuous to the feeling, and consisted of shortened and separated particles, not having the same coalescence which the fat has generally in the body. It became very soft when exposed to the heat of a fire, and sunk into a portion of paper on which it was spread, so as to make it more transparent. When the paper to which it was thus applied was exposed to the flame of a candle, it burnt with considerable crackling.



The hair with which the fatty substance was mixed, grew out of the inner surface of the capsule containing it, in some places in solitary hairs, but chiefly in small fasciculi, at scattered irregular distances. Besides these, there were loose hairs involved in the fatty mass. The hairs were some of them of considerable length, even to three inches; were fine, and of a light brown colour. They resembled much more the hair of the head than what are commonly found on the pubis, and corresponded very much in colour to the hair of the girl's head.

There arose, also, from the inner surface of the capsule, some vestiges of human teeth. One appeared to be a canine tooth; another to be a small grinder; two others to be incisors; and there was also a very imperfect attempt at the formation of another tooth. These were not fully formed, the fangs being wanting; but, in two of them, the bodies were as complete as they are ever found in common circumstances. They were each of them inclosed in a proper capsule, which arose from the inner surface of the ovarium, and consisted of a thick white opaque membrane.

Attached



Attached to the capsules of three of the teeth, there was a white spongy substance. The membrane of the ovarium itself was of some considerable thickness, but unequal in the different parts. It was very smooth in its inner surface, and more irregular externally. The uterus was smaller than it is commonly at birth; was perfectly healthy in its structure; and, upon opening into its cavity, it exhibited the ordinary appearances of a child's uterus at that period. The left ovarium was very small, corresponding to the state of the uterus. From this it appears, that the uterus had not yet received an increase in bulk, which is usual at the age of puberty. The hymen was entire, such as is commonly found in a child of the same age; and there was just beginning a lanugo upon the labia, not more than is often found on the upper lip of a boy of fifteen years old.

From this singular appearance, the circumstances of which have been now related, Dr Baillie next proceeds to draw some conclusions. It naturally, he observes, strikes the mind as being very different from any irregular substance formed by disease. It took place in a



part of the body which is subservient to generation, and where a complete foetus is sometimes formed. This might seem to lead to the conclusion, that the production of hair and teeth in the ovarium was a sort of imperfect impregnation. But he thinks there are reasons, at least equally strong, for believing that such productions may arise from an action in the ovarium itself, without any stimulus from the application of the male semen.

In the case which has just been related, the uterus was as small as at birth; and the left ovarium corresponded to the state of the uterus. It had not been stimulated, nor did it appear capable of being stimulated, by the application of the male semen. This he considers as a strong circumstance; for a preparation is still preserved in the collection of Windmill Street, made by the late Dr Hunter, where an ovum was formed in one of the Fallopian tubes, and the uterus was enlarged to more than twice its unimpregnated size. From this he concludes, that when impregnation takes place out of the cavity of the uterus, that viscus still takes a share in the action, and undergoes some of the changes of impregnation.



nation. And, to the same conclusion respecting the action of the uterus in the formation of an extra-uterine foetus, he is led also by other preparations, particularly from one, preserved also in Dr Hunter's collection, where a foetus was formed in the ovarium, in which the uterus was increased to more than twice its ordinary size, was very thick and spongy, and had its blood-vessels enlarged in the same manner as an impregnated uterus. But, in the present case, the uterus had undergone no change.

Dr Baillie farther remarks, that we are not to consider the formation of teeth in the ovarium to be a quicker process than it usually is in the head of a foetus. In this instance, however, the teeth having advanced fully as far as they in general are at some months after birth, their growth must, he thinks, have begun at least more than twelve months before the death of the child; which, in his opinion, brings their commencement to an earlier period than impregnation can be believed to have taken place.

From all these circumstances, Dr Baillie is inclined to suppose, that the formation of hair



and teeth was not here the consequence of any connexion with a male, but arose from some action of the ovarium itself, in which the uterus did not participate. And, as some additional confirmation of the probability of this opinion, he observes, that hair is occasionally found in parts of the human body which are altogether unconnected with the function of generation. Hairs have often been found in encysted tumours; and Dr Baillie particularly mentions several preparations of this sort in the collection of Mr John Hunter. All these circumstances, he thinks, clearly prove, that hair may be formed without any species of generation.

But hair is as much a peculiar substance as teeth. If the one, therefore, can be preternaturally formed in the ovarium, no good reason can, he thinks, be assigned, why the other may not also be formed there. The action producing the one, is not, he observes, better understood than that producing the other. Teeth, therefore, he thinks, may probably be formed by a peculiar action taking place in the ovarium, as well as hair.



To add still farther weight to this opinion, he observes, that many of the adult teeth are formed in a child after birth; and that, therefore, their formation depends on an action taking place in the jaw at a particular period, and not on original growth. Teeth are sometimes, also, occasionally formed at an advanced period of life. Both these processes, he observes, take place after the animal has been formed, in consequence of a certain action being excited in a particular part of the body. There is, therefore, he thinks, less difficulty in believing, that the same sort of process may go on in another part of the body not commonly employed in it. And, in his opinion, it is but reasonable to suppose, that the ovaria should have a greater aptitude of taking on a process somewhat similar to generation, than other indifferent parts of the body, as they constitute a part of the organs which are so materially concerned in the real process itself. From all these circumstances, taken collectively, he thinks it is rendered very probable, that the formation of hair and teeth in the ovarium does not necessarily depend, as has been the common opinion, on a connection



between a male and female, but arises from some action within the ovarium itself, which is imitative of generation.

How far the intelligent reader will be disposed to adopt this theory, we will not pretend to conjecture. We must, however, observe, that we readily agree with the ingenious author of this paper, in thinking, that there is no reason for supposing the generation of hair and teeth, in the fatty substance found in the ovarium, in the case related above, was the effect of any connection with a male; but perhaps a conjecture, more simple than any supposed action of the ovarium itself, imitative of generation, may be suggested with respect to their production. Although teeth, in general, appear only after certain periods of life, and in an established succession; yet it is well known, that the rudiments of all of them are discoverable in the jaw of a foetus at a very early period of its existence. The teeth, as the ingenious Mr John Hunter has justly observed, though deriving their growth and nourishment from the system, may yet be viewed as aliens to the body. This observation is no less applicable to the hairs. Each hair, found either  
upon



upon the head, or any other part of the body, derives its origin from a particular bulb, as much as a tooth does from its original rudiment. May we not, therefore, suppose teeth and hair to grow in the ovaria, from some monstrous conformation of parts forming there the rudiments or bulbs of each? and may not this monstrous conformation be peculiar to the ovaria, from that function for which they are naturally intended? For, although generation is still to be viewed as one of the greatest mysteries in the animal œconomy, yet there can be no doubt, that the ovaria have a concern in that function; and that thus they are subservient even to the production of a complete fœtus, furnished both with hair and with teeth. But it is alone from future observations of facts, that even probable evidence, either of this or any other conjecture, can be obtained.



