# Comparative view of mortality from child-birth in London and Dublin / by Medicus U.

#### Contributors

Merriman, Samuel, 1771-1852. Royal College of Surgeons of England

#### **Publication/Creation**

[London] : [publisher not identified], [1816]

#### **Persistent URL**

https://wellcomecollection.org/works/vxr2vmp6

#### Provider

Royal College of Surgeons

#### License and attribution

This material has been provided by This material has been provided by The Royal College of Surgeons of England. The original may be consulted at The Royal College of Surgeons of England. where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org

## For the London Medical and Physical Journal.

### Comparative View of Mortality from Child-birth in London and Dublin; by MEDICUS U. J.M.

IN the twenty-fifth volume of this Journal, a paper was I inserted (page 214), showing the number of deaths in child-bed, taken from the London Bills of Mortality, according to averages calculated upon periods of ten years each. There has lately been published in the Edinburgh Medical and Surgical Journal, No. XLVII. " An Abstract of the Registry kept at the Lying-in Hospital, in Dublin, from the 8th of Dec. 1757 (the day it was first opened) to Dec. 31, 1814;" from which it appears that 78,001 women have been delivered-that the number of children amounted 4 year to 79, 503, of which 41, 523 were boys, and 37, 982 girls, which e Than is in the proportion of about 10 males to 9 females-that the number of plural births were 1,372, or 1 in 57, of which 1,350 were twins, 21 triplets, and 1 quadruplet; the proportion of more than two at a birth being 1 in 3,545. The still-born children amounted to 4,329, which is about 1 in 18: the children dying in the Hospital after birth (I presume, in the first month) were 4,810, about 1 in 16. The deaths of women in child-bed were 835 (not 838 as misprinted), which is in the proportion of 1 in 93.

In order to compare the deaths in child-bed at this Hospital with what occurred during the same periods at London, I have been induced to take averages of ten years each, as has been before done in your Journal; and, if you think proper to publish the same, my calculations are much at your service.

At the Dublin Lying in Hospital.	
Three years and a few days to	Ten years to
Dec. 31, 17601 in 82	Dec. 31, 17601 in 81
10 ys. to Dec. 31, 17701 in 67	17701 in 72
17801 in 94	17801 in 92
17901 in 103	17901 in 107
18001 in 117	18001 in 113
18101 in 102	18101 in 106
4 yrs. to Dec. 31, 1814 1 in 66	Four years 18141 in 110

To what is the enormous increase of mortality in childbed at the Dublin Hospital, during the last four years, to be attributed?

August, 1816.

Day