

The land operculate mollusca collected during the 'Skeat expedition' to the Malay peninsula in 1899-1900 / by E.R. Sykes.

Contributors

Sykes, Ernest Ruthven, 1867-1954.
Royal College of Surgeons of England

Publication/Creation

[London] : [publisher not identified], 1903.

Persistent URL

<https://wellcomecollection.org/works/cjfhsmz7>

Provider

Royal College of Surgeons

License and attribution

This material has been provided by This material has been provided by The Royal College of Surgeons of England. The original may be consulted at The Royal College of Surgeons of England. where the originals may be consulted. The copyright of this item has not been evaluated. Please refer to the original publisher/creator of this item for more information. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. See rightsstatements.org for more information.

(Pl.)

THE LAND OPERCULATE MOLLUSCA COLLECTED
DURING THE "SKEAT EXPEDITION" TO THE
MALAY PENINSULA IN 1899—1900. BY E. R.
SYKES, F.Z.S.

[*From the* PROCEEDINGS OF THE ZOOLOGICAL SOCIETY OF LONDON,
1903, vol. i.]

[Published August 6, 1903.]

THE HISTORY OF THE
CITY OF BOSTON
FROM THE FIRST SETTLEMENT
TO THE PRESENT TIME

BY
JOHN H. COLEMAN
OF THE
CITY OF BOSTON



Digitized by the Internet Archive
in 2015

<https://archive.org/details/b2240644x>

[From the PROCEEDINGS OF THE ZOOLOGICAL SOCIETY OF LONDON,
1903, vol. i.]
(Published August 6, 1903.)

On the Land Operculate Mollusca collected during the
"Skeat Expedition" to the Malay Peninsula in 1899-
1900. By E. R. SYKES, F.Z.S.

(Plate XX.¹)

The species of Land-Mollusks collected by the "Skeat Expedition," though not very numerous, are of considerable interest on account of the welcome addition made to a fauna which is as yet but little known. They include:—

LEPTOPOMA ASPIRANS Benson.

Leptopoma aspirans Benson, Ann. Nat. Hist. ser. 2, vol. xvii. p. 229.

Hab. Biserat, Jalor. It has been recorded from Bukit Pondong and the Kinta Valley, having been originally described from Tenasserim.

LAGOCHILUS KOBELTI, sp. nov. (Plate XX. figs. 13-15.)

Testa modice umbilicata, turbinata, solidula, lirulis numerosis cincta, periostraco brunneo leviter induta; spira conica; anfr. 5½, convexi; apertura modice obliqua, subcircularis, peristomate duplici, incrassatulo et subreflexo, juxta insertionem breviter sed distincte inciso. Alt. 6·8; diam. max. 7·0 millim.

Hab. Biserat, Jalor.

¹ For explanation of the Plate, see p. 199.

The species of *Lagochilus* recorded from the Malay region are very puzzling and I think Dr. Moellendorff was quite right in describing as *L. rollei* the form that I noted (under the name of *L. townsendi*) from Kelantan. The nearest ally of *L. kobelti* appears to be *L. townsendi* Crosse; I have not seen an authentic specimen of that species, but have compared the form now described with the description and figures given by Crosse and with a specimen collected by Herr Grubauer, from whose collections Dr. Moellendorff has recorded *L. townsendi* as the only species found. The shell I now name is a trifle smaller and more elevated in proportion to the breadth, and the umbilicus is narrower. It may be noted that the reference to Crosse's original paper in the 'Journal de Conchyliologie' should be to p. 200 and not p. 208 as given by De Morgan and Dr. Moellendorff in their papers on the Perak fauna.

I have named the form after Dr. Kobelt as a trifling recognition of his recent study of the Cyclophoridae.

DITROPIS CAVERNÆ, sp. nov. (Plate XX. figs. 17-19.)

Testa depresso-conoidea, late umbilicata, olivacea, tenuis, glabra; spira mediocriter elevata, apice eroso, sutura impressa; anfr. 4 (?) convexi, ultimus antice vix descendens, carinatus, carinis duabus supra peripheriam, unica ad peripheriam, et plurimis in regione umbilicali; apertura subovalis, peristomate incrassatulo. Alt. 2.2; diam. max. 1.7 millim.

Hab. In a cave, Biserat, Jalor.

A single specimen only.

CYCLOPHORUS MALAYANUS (Benson).

Cyclostoma malayanum Benson, Ann. Nat. Hist. ser. 2, vol. x. p. 269.

Hab. Gunong Inas, Perak.

Recently when cataloguing (J. Malac. ix. p. 61) a collection of shells from Kelantan, I gave the names of *Cyclophorus saturnus* Pfr. and *borneensis* Metc.: both the forms then recorded have occurred in the present collection, and I have therefore again considered the identifications. Both, according to my present view, are erroneous, and the group is a very difficult one. The present species, which I regard as a form of *C. malayanus*, was then named *C. saturnus*, and the next species was called *C. borneensis*.

CYCLOPHORUS TUBA (Sby.).

Cyclostoma tuba Sowerby, Proc. Zool. Soc. 1842, p. 83.

Hab. Gunong Inas, Perak.

See remarks under the last species; probably the *C. borneensis*, recorded by De Morgan from Perak, also belongs to this species.

PTEROCYCLOS SUBALATUS, sp. nov. (Plate XX. figs. 1, 2.)

Testa late umbilicata, orbiculato-depressa, lineis incrementi notata,

brunnea, strigis corneis picta, fascia unica nigro-brunnea ad peripheriam ornata; anfr. 4½, mediocriter crescentes, convexi, sutura bene notata separati; apertura subcircularis, peristomate indistincte duplicato, ala parva antice angulato. Alt. 8; diam. max. 16 millim.

Hab. Gunong Inas, at 5000 feet.

I thought at first that this might be a form of *P. blandi* Bens., but it appears to be smaller, more elevate and not so widely umbilicated, and to differ in colour. In the two specimens that I have seen the lip is duplicated only on its outer margin, and the wing is small and thin.

RHIOSTOMA JALORENSIS, sp. nov. (Plate XX. figs. 6-8.)

Nearly related to *R. housei* but larger, slightly more depressed, and with the whorls not so tightly coiled. The tube is long and large, reaching the body of the shell, and is bent slightly backwards at the junction. The disjoined portion of the last whorl is much longer than is the case in *R. housei*, as will be seen from the figure, and is more descending. The colour resembles that of *R. housei*, and a dark band is usually present at the periphery. Operculum as usual. Diam. max. 29 millim.

Hab. Limestone Hills and Caves, Biserat, Jalor.

I have been in some doubt as to whether this is not a local race of *R. housei*, but the differences are constant in the specimens examined and, I think, are of specific value. The porcellaneous appearance of the operculum, so often seen in this group, appears to be due to the wearing down of the outer layers.

RHIOSTOMA, sp.

Hab. Kwala Aring, Kelantan.

A single specimen which agrees well with *R. jousseaumei* De Morgan, save that the tube is placed a little further back from the mouth, and is thinner and more cylindrical. It appears safer, however, to await further material ere describing this form.

OPISTHOPORUS PENANGENSIS Stol.

Opisthoporus penangensis Stoliczka, J. As. Soc. Bengal, vol. xlii. pt. 2, p. 265, pl. x. fig. 7.

Hab. Kwala Aring, Kelantan.

A single specimen, which I refer to this species with some doubt.

RHAPHAULUS ASCENDENS, sp. nov. (Plate XX. figs. 11, 12.)

Testa pupoidea, anguste umbilicata, fusco-purpurea, dense costulato-striata, spira bene elevata, apice obtusulo; anfr. 6, plano-convexi, penultimus gibbosus; apertura subcircularis, intus pallide fusca; peristoma pallide corneum, expansum et reflexum, marginibus callo junctis; tubulus brevis, incrassatus, a sutura oblique ascendens. Alt. 18.5; diam. max. 9.5 millim.

Hab. Patalung.

A single specimen "from rotten wood." Recalling in shape *R. perakensis* Smith, the present species is larger and stouter, and the tube slants obliquely upwards instead of descending; as compared with *R. lorraini* Pfr., which also has an ascending tube, the form is not so cylindrical, the whorls are flatter, and the size is much greater.

RHAPHAULUS PERAKENSIS Smith, var. *JALORENSIS*, n. var.
(Plate XX. figs. 9, 10.)

Shell very similar to *R. perakensis* Smith, but a little thinner and paler in colour, and the tube is bent along the outer lip, being attached to the lip, and not diverted behind it as in typical *R. perakensis*. The tube is broad and short. This form may prove to be a distinct species, but bearing in mind the great variation shown by Col. Godwin-Austen (Moll. India, vol. i. p. 202, pl. xlvii.) to exist in *R. blanfordi* Benson, and also considering how little we know of the group, most forms having been described from very few specimens, I have thought it wiser to give only a varietal name.

Hab. Bukit Bisar, on the borders of Jalor, altitude 2000 feet.

A single specimen.

SCHISTOLOMA ANOSTOMA (Benson).

Cyclostoma anostoma Benson, Ann. Nat. Hist. ser. 6, vol. x. p. 269.

Hab. Belimbing, Ligei.

SCHISTOLOMA SECTILABRUM (Gould).

Cyclostoma sectilabrum Gould, Boston Journ. Nat. Hist. vol. iv. p. 459, pl. xxiv. fig. 10.

Hab. Ulu Selama, Perak.

PUPINA LOWI De Morgan.

Pupina lowi De Morgan, Bull. Soc. Zool. France, vol. x. 1885, p. 414, pl. vii. fig. 3 (*lowi* on plate).

Hab. Gunong Inas, Perak.

PUPINA AUREOLA Stoliczka.

Pupina aureola Stoliczka, J. As. Soc. Bengal, vol. xli. pt. 2, p. 267, pl. x. figs. 11, 12.

Hab. Jalor, a single specimen.

Agrees well with Stoliczka's figure 12, but his figure 11 looks as if it might belong to a different species.

ALYCÆUS THIEROTI De Morgan.

Alycæus thieroti De Morgan, Bull. Soc. Zool. France, vol. x. 1885, p. 403, pl. viii. fig. 6; Moellendorff, Proc. Zool. Soc. 1891, p. 342.

Hab. Belimbing, Ligei, a single specimen.

ALYCÆUS DIPLOCHILUS Moellendorff.

Alycæus diplochilus Moellendorff, J. As. Soc. Bengal, vol. lv. pt. 2, 1886, p. 313; Proc. Zool. Soc. 1891, p. 342, pl. xxx. fig. 8.

Hab. Cave near Biserat, Jalor.

ALYCÆUS CONFORMIS Fulton.

Alycæus conformis Fulton, Ann. Nat. Hist. ser. 7, vol. ix. 1902, p. 68.

Hab. Kwala Aring, Kelantan.

ALYCÆUS PERAKENSIS Crosse.

Alycæus perakensis Crosse, J. Conchyl. vol. xxvii. 1879, p. 206, pl. xii. fig. 7.

Hab. Biserat, Jalor, amongst rocks.

OPISTHOSTOMA ANNANDALEI, sp. nov. (Plate XX. figs. 4, 5.)

Testa conico-pyramidalis, perforata, pallide rufescens, spira bene elevata, apice obtuso; anfr. 7, convexi, primi læves, reliqui distante lamellati, ultimus constrictus, retroversus, ascendens, conspicue solutus; apertura rotundata, peristomate leviter incrassato. Alt. 2.2; diam. max. 2.8; diam. min. 1.6 millim.

Hab. Jalor.

A single specimen, found by Mr. Annandale in débris on the floor of a cave. Related to *O. laidlawi* Sykes, from Kelantan, but the present species is larger, and the spire is much more produced, thus becoming more cylindrical in appearance.

DIPLOMMATINA SKEATI, sp. nov. (Plate XX. fig. 3.)

Testa sinistrorsa, vix rimata, ovato-fusiformis, aurantio-rufa, solidiuscula; anfr. 6½, convexi, primi læves, reliqui costulis parvis subremotis regulariter sculpti; sutura bene impressa; apertura fere circularis, peristomate incrassatulo, bene reflexo, subalato; lamella columellaris parva. Alt. 3.1; diam. max. 1.9 millim.

Hab. Gunong Inas, Perak, at about 5000 feet.

Only a single specimen of the shell, the most salient features of which are the inflated whorls, deeply cut suture, and fine, regular costulations.

DIPLOMMATINA LAIDLAWI, sp. nov. (Plate XX. fig. 16.)

Testa sinistrorsa, rimata, ovato-fusiformis, corneo-albida, apice rubello; anfr. 5½, convexi, primi læves, reliqui costulis parvis remotis regulariter sculpti, interstitiis sub lente dense spiraliter striatis; apertura quadrato-circularis, peristomate duplici, expanso, margine columellari sinuato, subalato; lamella columellaris minima, indistincta. Alt. 2; diam. max. 1 millim.

Hab. Gunong Inas, 3000–4500 feet, Perak.

Found "on the under surface of dead leaves, young with adults."

Much smaller than the last species, the costulæ more distant, and densely marked with minute spirals. I can find no trace of these latter in *D. skeati*, but the single specimen was not taken alive.


GEORISSA MONTEROSATIANA Godwin-Austen & Nevill.

Georissa monterosatiana Godwin-Austen & Nevill, P. Zool. Soc. 1879, p. 729, pl. lix. fig. 6.

Hab. Perak.

EXPLANATION OF PLATE XX.

- Figs. 1, 2. *Pterocyclos subalatus*, p. 195.
3. *Diplommatina skeati*, p. 198.
4, 5. *Opisthostoma annandalei*, p. 198.
6, 7, 8. *Rhiostoma jalorensis*, p. 196.
9, 10. *Rhaphaulus perakensis*, var. *jalorensis*, p. 197.
11, 12. *Rhaphaulus ascendens*, p. 196.
13, 14, 15. *Lagochilus kobelti*, p. 194.
16. *Diplommatina laidlawi*, p. 198.
17, 18, 19. *Ditropis cavernæ*, p. 195.



Green del. et lith.

Mintern Brow imp.

MOLLUSCA FROM THE MALAY PENINSULA.

