

**De maxillae inferioris luxatione : theses anatomico-chirurgicae : has
theses, Deo juvante & praeside M. Petro Sue, secundo ... / tueri conabitur
Carolus Daniel Gaultier de Claubry.**

Contributors

Gaultier de Claubry, Charles Daniel, 1757-1821.
Sue, P. 1739-1816.
Royal College of Surgeons of England

Publication/Creation

Parisiis : Typis Michaelis Lambert, 1782.

Persistent URL

<https://wellcomecollection.org/works/fkpkyfmw>

Provider

Royal College of Surgeons

License and attribution

This material has been provided by The Royal College of Surgeons of England. The original may be consulted at The Royal College of Surgeons of England. where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

DE

MAXILLÆ INFERIORIS LUXATIONE

THESES
ANATOMICO-CHIRURGICÆ.

*HAS THESES, Deo juvante & Praefide M. PETRO SUE,
secundo, antiquo Collegii Preposito & nunc Quæstore, antiquo
Scholarum Præticarum Anatomiae & Chirurgiae Professore,
Regiæ Academiæ Chirurgiae Consiliario, in generali Parisiensi
præfecturâ Chirurgo ordinario, necnon Academiarum Mons-
pellensis, Rothomagensis, Divionensis, Lugdunensis & Bur-
digalensis Socio, tueri conabitur CAROLUS DANIEL
GAULTIER DE CLAUBRY, Blesensis, Artium Liberalium
præclaræ Universitatis Parisiensis Magister, & Principis Atre-
batum Comitis, fratr̄is Regis, Chirurgus trimestris.*

*Die Luna, 17^a Junii, anno Salutis 1782, à sesqui-secundâ
post meridiem ad septimam.*

IN REGIIS CHIRURGORUM SCHOLIS,
PRO ACTU PUBLICO
ET
SOLEMNI COOPTATIONE:




PARISIIS,

Typis MICHAELIS LAMBERT, Regii Chirurgiae
Collegii necnon Academiæ Typographi, viâ Cithareâ.

M. D C C. L X X X I I.

Digitized by the Internet Archive
in 2016

<https://archive.org/details/b22414812>



D. O. M.

DE

MAXILLÆ INFERIORIS LUXATIONE.

THESES ANATOMICO-CHIRURGICÆ.

POSITIONES ANATOMIAE.

I.

MAXILLA inferior, sive mandibula, soleæ ad instar ferreæ figurata, in fœtu infanteque bifida, in unicum os, vigente ætate, coalescit. In tres partes dividitur, scilicet medium, quæ dicitur *corpus*, & laterales quæ *rami* nuncupantur.

A 2

I I.

IN corpore duas facies & totidem oras discernere fas est: facies anterior, convexa, in medio *mentum* vocata, prominentem & perpendicularē habet lineam, *Sympysim Maxillæ inferioris* dictam. Facies posterior, concava, leviusculis hinc illinc eminentiis exasperatur. Ex duabus oris, superior, in completo dentium numero, sexdecim protuberat alveolis, quæ græcis φάλναι & φάλναι, præsepiola, à præsepium similitudine, vocantur. Inferior ora Maxillæ, *basis*, duo labra, unum internum, alterum externum profert. In medio prominet linea superficialis, Maxillæ *spina anterior* dicta.

I I I.

PARALLELOGRAMMUM æmulatur uterque Maxillæ ramus: tres in unoquoque margines, tres anguli occurunt: horum inferior asperitatibus pluribus conspergitur, & licet obtusus, propriè tamen *anguli maxillaris* nomine donatur: superiorum alter triangularis & acuminatus, anterior *coronoides*, alter verò, in modum capituli conformatus, ad aures vergens, *condyloides* processus audit. Hic cavitate quâdam glenoïdeâ, in osse temporali excavatâ, recipitur, pro Maxillæ motibus. Cavitas non secus ac condylus cartilagine obducitur: hic verò, breviori collo sustentatus, pluribus ligamentis temporum ossi alligatur, aliis lateralibus externis, altero interno capsulari. Peculiaris insuper in cavitate cartilago, inter-articularis dicta, adnotatur (1). Pro ratione adductionis aut diductionis Maxillæ, vel sub jugo latef-

(1) Vid. accuratam hujus articuli descriptionem à Cl. *Monro* editam in *The Medical Essays*, vol. I, Obs. XI. p. 131, & vol. III. Obs. XIII, p. 261.

cunt processus coronoïdes, vel extrâ illud prosthinent, & à posterioribus condyloïdeis, incisuris semi-circularibus acutis, distinguntur.

I V.

Q U I N Q U E præcipue musculorum paria Maxillæ inferiori annectuntur, scilicet acuto ejusdem processui *temporalis*, inferiori & laterali faciei externæ parti *masseter*, foveæ maxillarique angulo anteriùs *pterigoïdeus internus*, condyli collo *pterigoïdeus externus*, menti tandem *symphysi biventer*. Superiores præterea & inferiores hyoïdis ossis, communesque seu externi laryngis musculi in eodem osse insertionem habent.

V.

IN interiore mandibulæ substantiâ, canalis angustus, osseus, maximâ ex parte continuus, *canalis maxillaris inferior* dictus, secundum ossis figuram flexuose incedit. Duo habet orificia; alterum in parte internâ, asperum, inæquale, viam præbet arteriæ, maxillaris internæ ramo, & nervo tertii quinti patis rami surculo: alterum in parte externâ infrâ primos molares terminatur. Porrò ea est inter hæc duo orificia consensus, ut si setam interno indideris, eam per externum levi operâ protrudere possis.

V I.

DEORSUM, sursùm, antrorsùm, retrorsùm & in latus moveri potest, ad obeunda manductionis & loquelæ munia, Maxilla inferior. Sursùm elevatur & sic clauditur muscularis temporalibus, massetere utroque, & pterigoïdeis internis, eam quoque interdùm ad latera trahentibus. Antrorsùm ducitur pterigoïdeis externis: biventes eam deprimunt, siveque os apertunt. Ex compositâ autem omnium horum muscularum actione variis in universum excentur motus (1).

(1) De iis nec-non de aliis Consul. Act. Reg. Scient. Academ. ann. 1744.

POSITIONES CHIRURGIAE.

I.

NON in aliam quām anteriorem partem propellitur Maxilla; id quod evenit, dūm ejus condyli super transversum ossis jugalis processum promoventur: Nam in posteriorem, dextram, sinistramque partem excidere nequit, quia obstant processus mastoīdei & styloīdei; nec propter ipsos condylos in latus moveri potest Maxilla. Ligamenta adhuc robustiora lateralē vel posteriori dimotioni inexpugnabile objiciunt impedimentum.

I I.

UTRIUSQUE luxationis speciei eadem causæ tūm internæ tūm externæ. Priores sunt ligamentorum debilitas, convulsiones, effrenati risus, oscitationes nimiæ. Postiores habendæ ictus & lapsus. Porrò ex omnibus his causis frequentior oscitatio.

I I I.

QUANDO utrāque parte è sede suâ excidit mandibula, totum mentum inclinatur & in partem anteriorem promovetur. Aper- tum manet os: deprimuntur genæ: saliva copiosè effluit: sic- cantur fauces: loquela deniqvè & deglutitio difficiles admo- dūm evadunt. Si unâ tantūm parte procidit maxilla, versùs latus oppositum convertitur mentum: dentes Maxillæ superioris den- tibus paribus non respondent; sed sub incisivis canini sunt. In uno latere musculorum tensio & tumefactio solummodo obser- vantur,

I V.

PARUM per se noxia Maxillæ luxatio: pessima tamen, si non reponeretur, procrearet symptomata, ut febrim, soporem, inflammationem, convulsionem, vomitum, atque etiam, juxta Hippocratis sententiam, ægri interitum. Sic quando sub osci-

tatione nonnunquam ita deorsum trahitur Maxilla, ut condyli extra sedem propriam propendeant, processus acuti musculos distorquent, pungendo convellunt, partes aliter ac in sanitatis statu trahunt, nervos & ligamenta violenter tendunt; & indè, nisi os opportunè in sedem compellatur, periculosa mox memorata symptomata. Hæc verò incompletæ luxationi rarius succedunt.

V.

MULTIPLICI ratione, si auctoribus fides habeatur, luxatæ maxillæ obtineri potest repositio. Sunt enim qui colaphis Maxillam in sedem suam compellere jubent: tali modo fuisse repositas Maxillæ luxationes testis est indubitatae fidei cel. *Petit*: luxationem verò tunc fuisse incompletam notat. Alii bacillo, veluti vecte, hinc posterioribus Maxillæ inferioris, indè incisivis alterius Maxillæ dentibus applicato, luxatum in sedem suam propulsare nituntur. Maxima autem ex eâ methodo pullulant incommoda, qualia sunt, dentium incisivarum ruptura, nimius dolor, faucium etiam pertimescenda lœsio, si è manibus excidat vectis. Quidam tandem bacillum brevius & cylindricum usque ad posteriorem arcus maxillaris partem protrudunt, mentoque elevato, condylos muscularum retractioni permittunt. Hujuscencheiresis difficilima praxis: bacillum enim sub solis ultimis dentibus continere molaribus ferè impossibile est & impossibilis fit reductio, si aliis dentibus innitatur bacillum.

V I.

TALIBUS methodis non minùs incongruis quam periculis, sequens subjicienda, optato semper coronata successu. Æger in sedili collocatur; sic ut minister à tergo

stans caput ejus firmiter retineat; vel ut is juxta parietem sedeat, subiecto inter parietem & caput ægri scorleo pulvino duro. Caput eò urgendum est, quò sit immobilius. Deinde Chirurgus, utroque pollice, linteolis priùs involuto, posterioribus ægri molaribus imposito, anteriorem Maxillæ partem volâ manuum complectitur: tūm ramos maxillares pollicibus deprimit, eosque postea retroagit, dūm pollices simul & subito in latera conjicit, ne præcipiti muscularum contractione lœdantur. Cautelæ autem, sub repositione, exposcunt, ut nec sursùm nec lateraliter, sed initio, in quantum licet, deorsùm, postea antrorsùm & denique sursùm ducatur Maxilla. Si unâ parte pruperit maxilla, in uno tantum latere lœfo adhibetur hæc praxis: difficultor verò tunc evadit: 1º. propter majorem in hoc casu muscularum tum contractionem tum resistentiam, 2º. propter minorem oris aperturam, quæ impedit ne pollex propè articulationem ducatur. De variis in hujusmodi luxatione inventis & applicatis, à celeberrimo *Lecat*, mediis, vid. diarium Medicinæ, tom. 11, pag. 28.

V I I.

MAXILLA, suum in locum reposita, spleniis & fasciâ quādam, fundâ dictâ, defensivo medicamento imbutâ, retinetur. Longior injicitur habena, utrinque in longitudinem ad medium ferè incisa, cuius media pars mentum, extremitatum verò duæ hominis caput complectuntur; reliquæ super cervicem, & hinc suprà frontem adductæ, ligantur. Futura præcaventur accidentia mediis generalibus & aptâ vietûs ratione; debellantur præsentia pro re natâ.

F I N I S.