

The pebble-band of the Elgin trias and its wind-worn pebbles / by Wm. Mackie.

Contributors

Mackie, William.
Keith, Arthur, Sir, 1866-1955
Royal College of Surgeons of England

Publication/Creation

London : Printed by Spottiswoode, 1901.

Persistent URL

<https://wellcomecollection.org/works/pnb7vk4g>

Provider

Royal College of Surgeons

License and attribution

This material has been provided by This material has been provided by The Royal College of Surgeons of England. The original may be consulted at The Royal College of Surgeons of England. where the originals may be consulted. Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>



The Outcrop Hill
of the succession of
a pure limestone. The
it is present at five w
There is evidence that
it is a debris fan on
lying E.O.B. rocks
truncated edges of the
than 100 feet in thick
out to the north-w
angles in the opposite
two series of rocks
discrepancy of dip
the other local area
it, and quite a mile in
the north-west, is w
also appear.

Another interest
but exclusively of q
evidence of sand-blast
'Pebbly' pebbles
pointing. Some of t
are beautifully polished
and the surfaces so
polishing. The effect
of differential etching.

Inquiries as to be
have hitherto failed
orientation obtained,
found sand-blast as
transferred by the act
was again subjected.

The result of this
based on the micro
Cottis Hill and
arriving at the same
stones, differences in
the limits from what
is the case of the
though the debris had
jected to wind action.



The Pebble-band of the Elgin Trias and its Wind-worn Pebbles.
By W. M. MACKIE, M.A., M.D.

The Cutties Hillock pebble-band, which has figured so largely in the discussion of the succession of the Elgin sandstones, is not, as has generally been contended, a pure localism. Two new openings into the Triassic rocks of the area show that it is present at five widely separated points. Its characters are constant in all. There is evidence that it is basal in position in the Triassic formation, and taking it as a datum line one is enabled to fix the relation of the Triassic to the underlying U.O.R. rocks with some certainty. It shows that the former overlies the truncated edges of the latter beds in a thin cake, which is probably nowhere more than 100 feet in thickness on a surface slightly inclined upwards from the south-east to the north-west, while the U.O.R. rocks steadily dip at almost constant angles in the opposite direction. Other facts definitely ascertained are, that the two series of rocks wherever they occur in proximity invariably show marked discordance of dip and strike, and that the Cutties Hillock area is detached from the other local areas of Triassic rocks, U.O.R. rocks having been traced all round it, and quite a mile intervening between it and the Spynie and Findrassie area to the north-west, in which interval U.O.R. rocks with discordant dip and strike also appear.

Another interest attaches to the pebble-band in that its pebbles, which are all but exclusively of quartz, quartzite, vein quartz, and chert, show unmistakable evidence of sand-blast action.

'Pyramidal pebbles' are common, with surfaces showing different degrees of polishing. Some of them even present strongly concave surfaces and finer depressions beautifully polished. A considerable number show 'flaking' of their edges, and the surfaces so formed have subsequently been subjected to different degrees of polishing. The cherts are beautifully fretted, and exhibit in perfection the results of differential etching.

Inquiries as to definite orientation of the more polished surfaces of the pebbles have hitherto failed to yield results. The author believes that no such definite orientation obtains, and is of opinion that the pebbles had been subjected to continued sand-blast action in some other locality, and were suddenly and forcibly transferred by the action of water to their present position, where many of them were again subjected to further sand-blast action.

The result of the examination of the pebbles supports the author's contention, based on the microscopical characters of their constituent sand-grains, that the Cutties Hillock sandstones are really Triassic sand-dunes. Other reasons for arriving at the same conclusion are: the peculiar undulating bedding of the sandstones, differences in the mode of occurrence as well as ontological differences of the fossils from what obtain in the adjoining areas.

In the case of the other local Triassic areas deposition in water is assumed, though the *débris* had evidently in some cases for a long time previously been subjected to wind action on a land surface.

PRINTED BY

SPOTTISWOODE AND CO. LTD., NEW-STREET SQUARE
LONDON





No copyright or title
page as the pamphlet
consists of 1 page.



