

**The constitution of ergothioneine : a betaine related to histidine / by  
George Barger and Arthur James Ewins.**

**Contributors**

Barger, George, 1878-1939.  
Ewins, Arthur James.  
Royal College of Surgeons of England

**Publication/Creation**

London : Wellcome Physiological Research Laboratories, 1911.

**Persistent URL**

<https://wellcomecollection.org/works/fzjdtjxs>

**Provider**

Royal College of Surgeons

**License and attribution**

This material has been provided by This material has been provided by The Royal College of Surgeons of England. The original may be consulted at The Royal College of Surgeons of England. where the originals may be consulted. Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>



272

CONSTITUTION

DETAINE REL

GEORGE

ARTHUR

1840-1841



*Dracts 1689. (1)*

THE  
CONSTITUTION OF ERGOTHIONEINE:  
A  
BETAINE RELATED TO HISTIDINE

BY  
GEORGE BARGER, M.A., D.Sc.  
AND  
ARTHUR JAMES EWINS, B.Sc.  
*(From the Transactions of the Chemical Society, 1911, Vol. 99)*



*From*

THE WELLCOME PHYSIOLOGICAL RESEARCH LABORATORIES  
BROCKWELL HALL  
HERNE HILL  
LONDON, S.E.









CCLVII.—*The Constitution of Ergothioneine: a Betaine Related to Histidine.*

By GEORGE BARGER and ARTHUR JAMES EWINS.


Two years ago Tanret (*J. Pharm. Chim.*, 1909, [vi], **30**, 145) isolated from ergot a new crystalline base containing sulphur, and having the formula  $C_9H_{15}O_2N_3S$ . The description of the base given by Tanret suggested to us a possible relationship to the amino-acid histidine; thus ergothioneine, as the new base was called by Tanret, is precipitated by mercuric chloride, and forms a compound with silver. These properties, together with the large nitrogen content, were strongly suggestive of the presence of a glyoxaline ring. Since one of us in conjunction with H. H. Dale (*Trans.*, 1910, **97**, 2592) had already isolated another histidine derivative from ergot, namely, 4(or 5)- $\beta$ -aminoethylglyoxaline, a substance of great physiological activity and interest, we prepared a small quantity of ergothioneine according to Tanret's method, and found that, like histidine and many other glyoxaline derivatives, it gives an intense red coloration with sodium *p*-diazobenzenesulphonate, thus still further supporting our surmise with regard to the constitution of this base.

Ergothioneine contains (in addition to the sulphur atom) three carbon atoms and six hydrogen atoms more than histidine. These might well result from the substitution of three hydrogen atoms of histidine by methyl groups in such a manner that ergothioneine would be a new member of the class of methylated amino-acids or betaines, of which a number of examples are now known to occur in plants. This hypothesis proved to be correct; the substance is indeed a betaine, and almost certainly  $\beta$ -2-thiolglyoxaline-4(or 5)-propiobetaine (I). Like other betaines it has no marked physiological action.

On boiling with a concentrated (50 per cent.) aqueous solution of potassium hydroxide, the base is decomposed quantitatively into



trimethylamine and a yellow acid of the composition  $C_6H_6O_2N_2S$ , which represents the whole of the remaining atoms in the molecule, and the constitution of which is represented by (II). On boiling with dilute nitric acid, the sulphur atom of this acid is completely removed, and a new acid,  $\beta$ -glyoxaline-4(or 5)-acrylic acid (III), is formed. This acid on reduction yields  $\beta$ -glyoxaline-4(or 5)-propionic acid (IV), and we were able to identify both the saturated and the unsaturated acid by comparison with synthetic specimens. The various reactions will be best seen from the following scheme:



The only doubtful point remaining was with regard to the position of the sulphur atom. There are two probable positions, namely, attachment to the  $\beta$ -carbon atom of the side-chain or to the carbon atom in the 2-position in the glyoxaline ring, and choice between these two, although decisive, is based on analogy rather than on direct evidence; indeed, synthesis seems to be the only means of supplying complete proof.

The sulphur in ergothioneine reacts in every way like that of the thiolglyoxalines, and quite different from that in cystein, with which it might be expected to show analogies if attached to the  $\beta$ -carbon atom of the side-chain; thus the sulphur in cystein, as is well known, is readily eliminated by boiling with sodium hydroxide, whereas, as was stated above, the sulphur of ergothioneine is not removed by boiling with the strongest solutions of potassium hydroxide. On the other hand, the sulphur atom, like that of the thiolglyoxalines, is readily and quantitatively oxidised by ferric chloride (as recently employed by Pyman) or by bromine water to sulphuric acid, whereas in the case of cystein a sulphonic acid results, in which the sulphur still remains attached to the carbon atom. The only oxidising agent which acts similarly on cystein and ergo-



thioneine is iodine, which oxidises both to a compound, in which two sulphur atoms are directly linked. That obtained from ergothioneine has the constitution (V).<sup>\*</sup> A further argument for considering ergothioneine to be a thiolglyoxaline derivative is the fact that, like other betaines, it is a feeble mono-acid base, whereas we should expect it to be di-acid if it, like histidine, contained the simple glyoxaline ring. The basic properties of the ring are destroyed by the presence of the sulphur atom as in other thiolglyoxalines (compare, for instance, 2-thiol-4(or 5)-aminomethylglyoxaline, which forms a mono-hydrochloride, and was recently described by Pyman, *Trans.*, 1911, **99**, 672).<sup>†</sup>

Treatment of ergothioneine with ferric chloride gave trimethylhistidine (VI), a betaine which may possibly occur in nature.

#### EXPERIMENTAL.

##### *The Action of Boiling 50 per cent. Aqueous Potassium Hydroxide on Ergothioneine.*

0.88 Gram of ergothioneine was boiled with 20 c.c. of a 50 per cent. aqueous solution of potassium hydroxide, the distillate being collected in a known volume of *N*-hydrochloric acid. Distillation was continued until the evolution of alkaline vapours was practically complete. It was found that 3.1 c.c. of *N*-hydrochloric acid had been neutralised, corresponding with 80 per cent. of the theoretical for the evolution of one nitrogen atom. The main bulk of the distillate was evaporated to dryness, and the residue dissolved in a little alcohol and treated with an alcoholic solution of platinic chloride, when trimethylamine platinichloride (m. p. 241°) separated. (Found, Pt=36.9. Calc., Pt=36.9 per cent.)

One nitrogen atom was thus shown to be evolved as trimethylamine. The strongly alkaline residue in the flask was rendered acid to Congo-red, when there was at once precipitated as a yellow, amorphous solid, *β*-2-thiolglyoxaline-4-acrylic acid.

This acid was found to be characterised by its general insolubility.

<sup>\*</sup> This iodide will be described later. It was already obtained by Tanret, who did not, however, appreciate its true significance. It forms black, steel grey or blue mixed crystals with iodine, a peculiar and extremely rare property which has so far only been observed in the case of cholalic acid, narceine, and saponarin, and is closely analogous to the adsorption of iodine by starch. For this reason Tanret, who analysed the compound, did not obtain results in agreement with any simple formula.

<sup>†</sup> Dr. Pyman has since informed us of his recent observation that thiolglyoxalines are sharply differentiated from glyoxalines by the fact that they at once decolorise a dilute cold solution of potassium permanganate; we find that ergothioneine also does this, but histidine does not, nor does cystine.



It is only moderately soluble in pyridine, and almost insoluble in all other ordinary organic solvents. The acid was crystallised for analysis by making a very dilute solution (0.1 per cent.) of the sodium salt, and acidifying. After three or four hours the acid began to separate, and precipitation was complete in about thirty-six hours. The acid obtained in this way separated in clusters of small prisms, which did not melt below  $275^{\circ}$ :

0.1179 gave 0.1805  $\text{CO}_2$  and 0.0420  $\text{H}_2\text{O}$ .  $\text{C}=41.8$ ;  $\text{H}=3.9$ .

$\text{C}_6\text{H}_5\text{O}_2\text{N}_2\text{S}$  requires  $\text{C}=42.3$ ;  $\text{H}=3.5$  per cent.

By oxidation by means of dilute nitric acid as described below, 90 per cent. of the theoretical amount of sulphur was obtained from the filtrate as barium sulphate.

*The Action of Boiling Dilute Nitric Acid on  $\beta$ -2-Thiolglyoxaline-4-acrylic Acid.*

0.4 Gram of amorphous  $\beta$ -2-thiolglyoxaline-4-acrylic acid was added in small portions to 20 c.c. of 10 per cent. aqueous nitric acid. The mixture was kept gently boiling on a sand-bath for some minutes after solution was complete, and then allowed to cool. The crystalline *nitrate* of  $\beta$ -glyoxaline-4(or 5)-acrylic acid separated out, which after washing and drying melted at  $198^{\circ}$  with explosive decomposition. The yield was 0.28 gram:

0.1114 gave 0.1470  $\text{CO}_2$  and 0.0404  $\text{H}_2\text{O}$ .  $\text{C}=36.0$ ;  $\text{H}=4.0$ .

$\text{C}_6\text{H}_6\text{O}_2\text{N}_2\text{HNO}_3$  requires  $\text{C}=35.8$ ;  $\text{H}=3.5$  per cent.

$\beta$ -Glyoxaline-4(or 5)-acrylic acid was most readily obtained from the nitrate by the addition of one equivalent of sodium carbonate (solid) to a concentrated aqueous solution of the salt, when the acid separates at once as a crystalline solid, which after recrystallisation from dilute acetone was quite pure, and melted at  $235\text{--}236^{\circ}$ :

0.1776 gave 0.3474  $\text{CO}_2$  and 0.0712  $\text{H}_2\text{O}$ .  $\text{C}=53.3$ ;  $\text{H}=4.4$ .

$\text{C}_6\text{H}_6\text{N}_2\text{O}_2$  requires  $\text{C}=52.2$ ;  $\text{H}=4.3$  per cent.

$\beta$ -Glyoxaline-4(or 5)-acrylic acid is moderately soluble in cold, but very readily so in hot, water. Its solutions give an intense red colour with sodium *p*-diazobenzenesulphonate.

The *phosphotungstate* is readily soluble in hot water or cold acetone, and crystallises in small, rectangular plates from dilute acetone.

The *picrate* forms golden-yellow prisms, melting at  $213\text{--}214^{\circ}$ , and sparingly soluble in cold water. The melting point of this picrate



remained unchanged when mixed with a synthetic specimen. The latter we obtained in small quantity by the action of trimethylamine on  $\alpha$ -chloro- $\beta$ -glyoxaline-4(or 5)-propionic acid.

*The Reduction of  $\beta$ -Glyoxaline-4(or 5)-acrylic Acid to  $\beta$ -Glyoxaline-4(or 5)-propionic Acid.*

$\beta$ -Glyoxaline-4(or 5)-acrylic acid was dissolved in a little absolute alcohol, and the hot solution treated with five to six atomic proportions of sodium. The solution was then acidified with hydrochloric acid, and the precipitated salt collected. The filtrate and washings were evaporated to dryness, the residue dissolved in 5 per cent. aqueous sulphuric acid, and precipitated with phosphotungstic acid. The precipitate was treated with acetone, filtered from undissolved material, and the soluble phosphotungstate decomposed in the usual manner. On concentrating the final filtrate a crystalline acid separated, which after recrystallisation melted at  $202^{\circ}$ . When mixed with a specimen of synthetic  $\beta$ -glyoxaline-4(or 5)-propionic acid (m. p.  $202^{\circ}$ ), the melting point remained unaltered. The acid obtained on reduction was therefore proved to be  $\beta$ -glyoxaline-4(or 5)-propionic acid.

*The Action of Ferric Chloride on Ergothioneine. Formation of  $\beta$ -Glyoxaline-4(or 5)-propiobetaine (Histidine-betaine).*

One gram of ergothioneine was boiled for one hour with an aqueous solution containing nine molecular proportions of ferric chloride. The iron was removed from solution by sodium carbonate, the filtrate acidified with sulphuric acid (so as to give a 5 per cent. solution of the latter), and precipitated by phosphotungstic acid. The precipitate, which was completely soluble in acetone, was decomposed in the usual way, and the barium and sulphuric acid removed. The filtrate was concentrated, and treated with a hot aqueous solution of picric acid. On cooling, a crystalline picrate separated, which after recrystallisation from water was obtained in deep yellow prisms melting at  $123^{\circ}$ :

0.1340 gave 0.1908  $\text{CO}_2$  and 0.0396  $\text{H}_2\text{O}$ .  $\text{C}=38.4$ ;  $\text{H}=3.3$ .

$\text{C}_9\text{H}_{15}\text{O}_2\text{N}_3(\text{C}_6\text{H}_3\text{O}_7\text{N}_3)_2$  requires  $\text{C}=38.5$ ;  $\text{H}=3.3$  per cent.

The *dipicrate* is very sparingly soluble in cold, fairly readily so in hot, water.

The *picrolonate* prepared in the usual way forms long, thin, orange-yellow needles, melting at  $229$ — $230^{\circ}$ .

The *aurichloride* separated from dilute aqueous hydrochloric acid as large, broad, deep orange-yellow prisms, melting at  $171^{\circ}$ .




In conclusion, we wish to express our indebtedness to Dr. F. L. Pyman for synthetic specimens of  $\alpha$ -chloro- $\beta$ -glyoxaline-4(or 5)-propionic and  $\beta$ -glyoxaline-4(or 5)-propionic acids for purposes of comparison, and to the Wellcome Chemical Works, Dartford, for a supply of ergothioneine.

THE WELLCOME PHYSIOLOGICAL RESEARCH LABORATORIES,  
BROCKWELL HALL, HERNE HILL, LONDON, S.E.







TITLES  
OF  
PUBLISHED PAPERS  
FROM THE  
WELLCOME CHEMICAL RESEARCH  
LABORATORIES

---

FREDERICK B. POWER, PH.D., LL.D., *Director*  
6, King Street, Snow Hill  
LONDON, E.C.

---

- |   |   |
|---|---|
| 1. SOME NEW GOLD SALTS OF HYOSCINE,<br>HYOSCYAMINE AND ATROPINE             | 12. OBSERVATIONS RELATING TO THE<br>CHEMISTRY OF THE BRITISH<br>PHARMACOPOEIA   |
| 2. THE CHARACTERS AND METHODS<br>OF ASSAY OF THE OFFICIAL<br>HYPOPHOSPHITES | 13. MERCUROUS IODIDE  |
| 3. NOTE ON THE MYDRIATIC ALKALOIDS  | 14. THE COMPOSITION OF BERBERINE<br>PHOSPHATE                                   |
| 4. PREPARATION OF ACID PHENYLIC<br>SALTS OF DIBASIC ACIDS                   | 15. A CONTRIBUTION TO THE PHARMA-<br>COGNOSE OF OFFICIAL STROPHAN-<br>THUS SEED |
| 5. A NEW METHOD FOR THE ANALYSIS<br>OF COMMERCIAL PHENOLS                   | 16. THE CHEMISTRY OF THE JABORANDI<br>ALKALOIDS                                 |
| 6. THE ASSAY OF PREPARATIONS CON-<br>TAINING PILOCARPINE                    | 17. A NEW ADMIXTURE OF COMMERCIAL<br>STROPHANTHUS SEED                          |
| 7. PILOCARPINE AND THE ALKALOIDS<br>OF JABORANDI LEAVES                     | 18. RESEARCHES ON MORPHINE—Part II  |
| 8. A NEW GLUCOSIDE FROM WILLOW<br>BARK                                      | 19. THE CONSTITUTION OF PILOCARPINE<br>—Part II                                 |
| 9. THE CONSTITUTION OF PILOCARPINE<br>—Part I                               | 20. THE CHEMISTRY OF THE BARK OF<br>ROBINIA PSEUD-ACACIA, <i>Linn.</i>          |
| 10. THE COMPOSITION AND DETERMINA-<br>TION OF CERIUM OXALATE                | 21. THE ANATOMY OF THE BARK OF<br>ROBINIA PSEUD-ACACIA, <i>Linn.</i>            |
| 11. RESEARCHES ON MORPHINE—Part I   | 22. A SOLUBLE MANGANESE CITRATE<br>AND COMPOUNDS OF MANGANESE<br>WITH IRON      |



# TITLES OF PUBLISHED PAPERS—continued

23. THE CHEMICAL CHARACTERS OF SO-CALLED IODO-TANNIN COMPOUNDS
24. THE CONSTITUTION OF PILOCARPINE—Part III
25. A NEW SYNTHESIS OF  $\alpha$ -ETHYLTRICARBALLYLIC ACID
26. THE CONSTITUENTS OF THE ESSENTIAL OIL OF ASARUM CANADENSE, *Linn.*
27. DERIVATIVES OF GALLIC ACID
28. THE OCCURRENCE OF SALICIN IN DIFFERENT WILLOW AND POPLAR BARKS
29. THE CONSTITUENTS OF COMMERCIAL CHRYSAROBIN
30. THE CONSTITUENTS OF AN ESSENTIAL OIL OF RUE
31. METHYL  $\beta$ -METHYLHEXYL KETONE
32. INTERACTION OF KETONES AND ALDEHYDES WITH ACID CHLORIDES
33. THE ANATOMY OF THE STEM OF DERRIS ULIGINOSA, *Benth.*
34. THE CHEMISTRY OF THE STEM OF DERRIS ULIGINOSA, *Benth.*
35. THE CONSTITUTION OF PILOCARPINE—Part IV
36. PREPARATION AND PROPERTIES OF DIMETHYLGLYOXALINE AND DIMETHYLPYRAZOLE
37. THE ELECTROLYTIC REDUCTION OF PHENO- AND NAPHTHOMORPHOLONES
38. CHEMICAL EXAMINATION OF KÔ-SAM SEEDS (*BRUCEA SUMATRANA*, *Roxb.*)
39. COMPARATIVE ANATOMY OF THE BARKS OF THE SALICACEÆ—Part I
40. THE CONSTITUTION OF CHRYSOPHANIC ACID AND OF EMODIN
41. THE CONSTITUTION OF EPINEPHRINE
42. A LÆVO-ROTATORY MODIFICATION OF QUERCITOL
43. THE CONSTITUENTS OF THE ESSENTIAL OIL OF CALIFORNIAN LAUREL
44. SOME DERIVATIVES OF UMBELLULONE
45. THE CONSTITUENTS OF CHAULMOOGRA SEEDS
46. THE CONSTITUTION OF CHAULMOOGRIC ACID—Part I
47. CHEMICAL EXAMINATION OF CASCARA BARK
48. CHEMICAL EXAMINATION OF GYMNEMA LEAVES
49. THE RELATION BETWEEN NATURAL AND SYNTHETICAL GLYCERYLPHOSPHORIC ACIDS
50. GYNOCARDIN, A NEW CYANOGENETIC GLUCOSIDE
51. PREPARATION AND PROPERTIES OF 1:4:5-TRIMETHYLGLYOXALINE
52. THE CONSTITUTION OF PILOCARPINE—Part V
53. THE CONSTITUTION OF BARBALOIN—Part I
54. THE CONSTITUENTS OF THE SEEDS OF HYDNOCARPUS WIGHTIANA, *Blume*, AND OF HYDNOCARPUS ANTHELMINTICA, *Pierre*
55. THE CONSTITUENTS OF THE SEEDS OF GYNOCARDIA ODORATA, *R.Br.*
56. THE SYNTHESIS OF SUBSTANCES ALLIED TO EPINEPHRINE
57. CHEMICAL EXAMINATION OF GRINDELIA
58. CHEMICAL EXAMINATION OF AETHUSA CYNAPIUM, *Linn.*
59. PREPARATION AND PROPERTIES OF SOME NEW TROPEINES
60. THE CONSTITUENTS OF THE ESSENTIAL OIL FROM THE FRUIT OF PITTOSPORUM UNDULATUM, *Vent.*
61. THE CONSTITUTION OF UMBELLULONE
62. LONDON BOTANIC GARDENS

# TITLES OF PUBLISHED PAPERS—continued

63. CHEMICAL AND PHYSIOLOGICAL EXAMINATION OF THE FRUIT OF CHAILLETIA TOXICARIA
64. CHEMICAL EXAMINATION OF ERIODICTYON
65. THE BOTANICAL CHARACTERS OF SOME CALIFORNIAN SPECIES OF GRINDELIA
66. THE RELATION BETWEEN NATURAL AND SYNTHETICAL GLYCERYL-PHOSPHORIC ACIDS—Part II
67. THE CONSTITUTION OF UMBELLULONE—Part II
68. THE REDUCTION OF HYDROXYLAMINODIHYDROUMBELLULONE-

80. THE CONSTITUTION OF UMBELLULONE—Part III
81. THE CONSTITUENTS OF OLIVE LEAVES
82. THE CONSTITUENTS OF OLIVE BARK
83. CHEMICAL EXAMINATION OF IPOMŒA PURPUREA
84. THE CHARACTERS OF OFFICIAL IRON ARSENATE
85. PREPARATION OF A SOLUBLE FERRIC ARSENATE
86. THE CONSTITUENTS OF THE EXPRESSED OIL OF NUTMEG
87. CHEMICAL EXAMINATION AND

DING  
OLIVE

OF

BARK

THE  
NUM

THE  
ETYL

## WITH THE DIRECTOR'S COMPLIMENTS

BARKS OF BRUCEA ANTIDYSENTERICA, *Lam.* AND BRUCEA SUMATRANA, *Roxb.*

75. CHEMICAL EXAMINATION OF GRINDELIA—Part II
76. CHEMICAL EXAMINATION OF LIPPIA SCABERRIMA, *Sonder* ("Beukess Boss")
77. CHEMICAL EXAMINATION OF THE ROOT AND LEAVES OF MORINDA LONGIFLORA
78. THE CONSTITUENTS OF THE ESSENTIAL OIL OF NUTMEG
79. CHEMICAL EXAMINATION OF MICROMERIA CHAMISSONIS (*Yerba Buena*)

93. THE ACTION OF NITRIC ACID ON THE ETHERS OF AROMATIC HYDROXYALDEHYDES
94. THE SYNTHESIS OF SUBSTANCES ALLIED TO COTARNINE
95. CHEMICAL EXAMINATION OF ELATERIUM AND THE CHARACTERS OF ELATERIN
96. THE TESTS FOR PURITY OF QUININE SALTS
97. THE CONFIGURATION OF TROPINE AND  $\Psi$ -TROPINE AND THE RESOLUTION OF ATROPINE
98. THE CONSTITUENTS OF THE FRUIT OF ECBALLIUM ELATERIUM



## TITLES OF PUBLISHED PAPERS—continued

# TITLES OF PUBLISHED PAPERS—continued

63. CHEMICAL AND PHYSIOLOGICAL EXAMINATION OF THE FRUIT OF CHAILLETIA TOXICARIA
64. CHEMICAL EXAMINATION OF ERIODICTYON
65. THE BOTANICAL CHARACTERS OF SOME CALIFORNIAN SPECIES OF GRINDELIA
66. THE RELATION BETWEEN NATURAL AND SYNTHETICAL GLYCERYL-PHOSPHORIC ACIDS—Part II
67. THE CONSTITUTION OF UMBELLULONE—Part II
68. THE REDUCTION OF HYDROXYLAMINODIHYDROUMBELLULONE-OXIME
69. THE CONSTITUTION OF CHAULMOOGRIC AND HYDNOCARPIC ACIDS
70. THE CONSTITUENTS OF THE ESSENTIAL OIL OF AMERICAN PENNYROYAL
71. THE CONSTITUTION OF HOMOERIODICTYOL
72. THE INTERACTION OF METHYLENE CHLORIDE AND THE SODIUM DERIVATIVE OF ETHYL MALONATE
73. CHEMICAL EXAMINATION OF THE FRUIT OF BRUCEA ANTIDYSENTERICA, *Lam.*
74. CHEMICAL EXAMINATION OF THE BARKS OF BRUCEA ANTIDYSENTERICA, *Lam.* AND BRUCEA SUMATRANA, *Roxb.*
75. CHEMICAL EXAMINATION OF GRINDELIA—Part II
76. CHEMICAL EXAMINATION OF LIPPIA SCABERRIMA, *Sonder* ("Beukess Boss")
77. CHEMICAL EXAMINATION OF THE ROOT AND LEAVES OF MORINDA LONGIFLORA
78. THE CONSTITUENTS OF THE ESSENTIAL OIL OF NUTMEG
79. CHEMICAL EXAMINATION OF MICROMERIA CHAMISSONIS (*Yerba Buena*)
80. THE CONSTITUTION OF UMBELLULONE—Part III
81. THE CONSTITUENTS OF OLIVE LEAVES
82. THE CONSTITUENTS OF OLIVE BARK
83. CHEMICAL EXAMINATION OF IPOMŒA PURPUREA
84. THE CHARACTERS OF OFFICIAL IRON ARSENATE
85. PREPARATION OF A SOLUBLE FERRIC ARSENATE
86. THE CONSTITUENTS OF THE EXPRESSED OIL OF NUTMEG
87. CHEMICAL EXAMINATION AND PHYSIOLOGICAL ACTION OF NUTMEG
88. SOME OBSERVATIONS REGARDING "OLEUROPEIN" FROM OLIVE LEAVES
89. CHEMICAL EXAMINATION OF ERIODICTYON—Part II
90. THE CONSTITUENTS OF THE BARK OF PRUNUS SEROTINA
91. THE CONSTITUENTS OF THE RHIZOME OF APOCYNUM ANDROSÆMIFOLIUM
92. *iso* AMYGDALIN, AND THE RESOLUTION OF ITS HEPTA-ACETYL DERIVATIVE
93. THE ACTION OF NITRIC ACID ON THE ETHERS OF AROMATIC HYDROXYALDEHYDES
94. THE SYNTHESIS OF SUBSTANCES ALLIED TO COTARNINE
95. CHEMICAL EXAMINATION OF ELATERIUM AND THE CHARACTERS OF ELATERIN
96. THE TESTS FOR PURITY OF QUININE SALTS
97. THE CONFIGURATION OF TROPINE AND  $\Psi$ -TROPINE AND THE RESOLUTION OF ATROPINE
98. THE CONSTITUENTS OF THE FRUIT OF EKBALLIUM ELATERIUM



# TITLES OF PUBLISHED PAPERS—*continued*

99. SYNTHESSES IN THE EPINEPHRINE SERIES
100. CHEMICAL EXAMINATION OF JALAP
101. THE CONSTITUENTS OF RUMEX ECKLONIANUS
102. THE CONSTITUENTS OF COLOCYNTH
103. THE CONSTITUENTS OF RED CLOVER FLOWERS
104. CHEMICAL EXAMINATION OF PUMPKIN SEED
105. CHEMICAL EXAMINATION OF WATERMELON SEED
106. CHEMICAL EXAMINATION OF ORNITHOGALUM THYRSOIDES
107. THE CONSTITUENTS OF THE FLOWERS OF TRIFOLIUM INCARNATUM
108. THE CONSTITUENTS OF THE LEAVES OF PRUNUS SEROTINA
109. SYNTHESIS OF COTARNINE
110. NOTE ON GYNOCARDIN AND GYNOCARDASE
111. CHEMICAL EXAMINATION OF THE TUBEROUS ROOT OF IPOMŒA HORSFALLIÆ
112. THE RESOLUTION OF BENZOYLSCINE
113. NOTE ON THE CONSTITUTION OF  $\alpha$ -ELATERIN
114. THE CONSTITUENTS OF LEPTANDRA
115. THE CONSTITUTION OF ERIODICTYOL OF HOMOERIODICTYOL, AND OF HESPERITIN
116. THE SYNTHESIS OF 2 : 4 : 6 - TRIMETHOXYPHENYL 3 : 4 - DIMETHOXYSTYRYL KETONE
117. THE CONSTITUENTS OF GELSEMIUM
118. SYNTHESSES IN THE EPINEPHRINE SERIES—Part II
119. THE ABSORPTION SPECTRA OF SOME SUBSTITUTED PYRAZINES AND THEIR SALTS
120. ACTION OF SODIUM AMALGAM ON METHYLENE ETHERS
121. THE CONSTITUENTS OF THE RHIZOME OF IRIS VERSICOLOR
122. CHEMICAL EXAMINATION OF THE ROOT OF LASIOSIPHON MEISSNERIANUS
123. THE ORIENTATION OF THE NITRO-GROUP IN NITROMYRISTICINIC ACID
124.  $\alpha$ - $p$ -HYDROXY- $m$ -METHOXYPHENYL-AMINE, AND THE RESOLUTION OF  $\alpha$ - $p$ -HYDROXYPHENYLETHYLAMINE
125. THE CONSTITUENTS OF WITHANIA SOMNIFERA
126. THE CONSTITUENTS OF RHUBARB
127. THE CONSTITUENTS OF BRYONY ROOT
128. THE CONSTITUTION OF SCPOLETIN
129. THE CONSTITUENTS OF THE BULB OF BUPHANE DISTICHA
130. SOME DERIVATIVES OF GELSEMINE
131. SYNTHESIS OF 4 : 6 - DIMETHOXY-2 -  $\beta$  - METHYLAMINOETHYL-BENZALDEHYDE
132. CHEMICAL EXAMINATION OF OENANTHE CROCATA
133. THE CONSTITUENTS OF THE SEEDS OF CASIMIROA EDULIS
134. CHEMICAL EXAMINATION OF CALABAR BEANS
135. CHEMICAL EXAMINATION OF THE LEAVES OF ANONA MURICATA
136. CHEMICAL EXAMINATION OF THE ROOT OF IPOMŒA ORIZABENSIS
137. THE CONSTITUENTS OF COMMERCIAL CHRYSAROBIN
138. THE PROPOSED METHOD OF MICRO-SUBLIMATION FOR THE DETECTION OF AESCULIN AND THE IDENTIFICATION OF GELSEMIUM
139. CHEMICAL EXAMINATION OF SCAMMONY ROOT AND OF SCAMMONY
140. CHEMICAL EXAMINATION OF JAMBUL SEEDS
141. THE CONSTITUENTS OF SOME CUCURBITACEOUS PLANTS
142. RESEARCHES ON THE CONSTITUTION OF PHYSOSTIGMINE—PART I





