

An account of a new species of the bark-tree, found in the island of St. Lucia / by George Davidson ; communicated by Donald Monro.

Contributors

Davidson, George.
Monro, Donald, 1727-1802.
Wilson, George, active 1784.
Royal College of Surgeons of England

Publication/Creation

[London] : [Royal Society of London], 1784.

Persistent URL

<https://wellcomecollection.org/works/waxz2766>

Provider

Royal College of Surgeons

License and attribution

This material has been provided by This material has been provided by The Royal College of Surgeons of England. The original may be consulted at The Royal College of Surgeons of England. where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

*An Account of a new Species of the Bark-Tree, found
in the Island of St. Lucia. By Mr. George Davidson; com-
municated by Donald Monro, M. D. Physician to the Army,
F. R. S.*

Read at the Royal Society, June 24, 1784.

DR. DONALD MONRO.

S I R,

HAVING received from my correspondent Mr. DAVID-
SON, surgeon, in the island of St. Lucia, some Bark,
the product of that island, which is said to possess the virtues
of the Jesuit's Bark, and in a much smaller dose, I shall
esteem it a favour if you will lay before the Royal Society the
specimen which I have sent to you with this letter, together
with Mr. DAVIDSON's account of it, if you think they me-
rit that honour.

I have examined the dried specimens very carefully. They
are not so well preserved as I could wish; but I have since seen
much finer in the possession of Sir JOSEPH BANKS, who has
done me the honour to favour me with the following character,
as most distinctive of it from the other species of Cinchona
already described, which he gave me an opportunity of
examining.

A

It

It is undoubtedly a Cinchona, but not the Cinchona officinalis of LINNÆUS; for it differs from it essentially in its bark in several particulars. It has an emetic quality not common to the true bark, breaks more woody and splintery, and is far more nauseous to the taste. Its decoction is of a dull Burgundy colour; and its extract resembles more the bitter of Gentian than that of the Quinquina. I have procured four ounces of it from half a pound of the Bark boiled in water, and herewith send to you a small specimen.

The drawings, which accompany this letter, are exact copies of the specimens which I received; I therefore hope they will not be thought unworthy the acceptance of the Royal Society.

I have the honour to be, &c.

Henrietta-street, Nov. 6, 1783.

G. WILSON.

Botanic character of the Bark-Tree of St. Lucia.

“Cinchona floribus paniculatis, glabris; laciniis linearibus,
“tubo longioribus; staminibus exsertis; foliis ellipticis, glabris.”

Extract of a Letter from Mr. GEORGE DAVIDSON, dated St. Lucia, July 15, 1783.

IT is now about four years since Mr. ALEXANDER ANDERSON discovered in the woods, near the Grand Cul de Sac, some trees resembling, in the botanical characters, the
true

true *Quinquina* of LINNÆUS. He brought the bark, flowers, and seeds, to Dr. YOUNG of the General Hospital, and trial was made of it there; but not being sufficiently dried, its strong emetic and purgative qualities prevented its exhibition.

The publication of Dr. SAUNDERS, which I received about two months ago, mentioning the introduction of a species of bark of a redder colour, and possessing greater powers than the bark formerly in use, induced us here to try the bark of this country. Dr. YOUNG had by him some that was collected in General GRANT's time: on account of the length of time it had been kept, and its being sufficiently dried, he has met with all the success he could wish.

It is manifestly more astringent than the bark, and the bitter is likewise more durable on the palate.

Hitherto I have generally used the cold infusion, either in lime or simple water, in the proportion of one ounce to three pints of the water. I have likewise given it in substance from twenty to thirty grains; but never exceeded the last quantity, for I never found the stomach able to retain more than twenty grains.

Joined with the *Canella alba*, it forms in spirits an agreeable and elegant tincture. I have made a tincture from the seeds, which are infinitely stronger in taste than the bark itself.

(Signed)

GEO. DAVIDSON.

Mr.

Mr. George Davidson's account of the Bark-Tree of the island of St. Lucia.

THE Bark-Tree of this island is nearly about the size of the cherry-tree, seldom thicker than the thigh, and tolerably straight; the wood is light and porous, without any of the bitterness and astringency of the bark itself.

It delights in a shady situation, the north-west aspect of hills, under larger trees; and is generally to be found about the middle of an hill, near some running water.

The leaves are large, oblong, opposite, and plain, preserving (as well as the flowers and seeds) the bitter taste of the bark.

In the beginning of the rainy season (June), the tree puts forth its flowers in small tufts; at first they are white, but afterwards turn purplish. The stamina are five in number, with a single style. The germen is oblong, bilocular, and furrowed on each side. The seeds are many, and of the winged kind. The corolla is monopetalous, with its mouth divided into five long segments.

The soil in general where it grows is a stiff red clay. The bark itself is of a lighter red than that sent out here to the hospital under the name of *red bark*. It inclines more to the colour of cinnamon. The bitterness and astringency appear to be greater than in either of the other barks.

I apprehend, the proper season for obtaining it is about the month of March, before the flowers come out: after-experience will best determine this.

Infused in cold water, in which form, or in lime-water, I generally use it, it forms a very red tincture, possessing the
bitterness

D
E



Digitized by the Internet Archive
in 2016

<https://archive.org/details/b22463288>



bitterness and astringency of the bark very strongly. A few drops of the *Tinctura florum martialium* give it a very black colour, and occasion a copious deposition of a black sediment. It does the same with the spirituous tincture.

With spirits it forms a beautiful red tincture.

Explanation of the references tab. XIX.

- A. A branch of the Cinchona of St Lucia, with the flowers not yet opened.
- B. The entire seed-vessels.
- C. A seed-vessel split.
- D. One of the seeds, of its natural size.
- E. The same magnified.



a new species of the Bark-Trees.
bitterness and stringency of the bark very strongly. A few
drops of the Tincture of Cinchona bark given in a very black
colour, and taken in a copious quantity of a black sediment
It does the best with the patients in fever.
Which I think it is a beautiful red tincture.

Explanation of the figures in the XIX.

- A. A branch of the Cinchona of St. Lucia, with the flowers
in bud.
- B. The same tree, with
C. A few red flowers.
- D. One of the leaves of its natural size.
- E. The same magnified.

