

A universal formulary containing the methods of preparing and administering officinal and other medicines / by R. Eglesfeld Griffith.

Contributors

Griffith, R. Eglesfeld 1798-1850.
Watson, Thomas, Sir, 1792-1882
Thomas, Robert P. 1821-1864.
Royal College of Physicians of London

Publication/Creation

Philadelphia, Pennsylvania : Blanchard and Lea, 1858.

Persistent URL

<https://wellcomecollection.org/works/eny37c4v>

Provider

Royal College of Physicians


License and attribution

This material has been provided by This material has been provided by Royal College of Physicians, London. The original may be consulted at Royal College of Physicians, London. where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.




Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>



615-13 SL/23-5-G-27 SL






SL


615.


70


16

✓









Presented by Thomas H. ...
President of the College
...

UNIVERSITY

LIBRARY

Letter to John Jay
June 1790

Dear Sir

I have the honor to receive

your letter of the 25th

*Presented by Thomas Watson, M.D.
President of the College.
1864.*


UNIVERSAL
FORMULARY.

Presented by James Watson M.
1877 E. H. Johnson of the College
New

THE UNIVERSITY OF CHICAGO
LIBRARY

UNIVERSAL
LIBRARY

FORMULARY



THE UNIVERSITY OF CHICAGO
LIBRARY

A

UNIVERSAL FORMULARY:

CONTAINING THE METHODS OF

PREPARING AND ADMINISTERING

OFFICINAL AND OTHER MEDICINES.

THE WHOLE ADAPTED TO

PHYSICIANS AND PHARMACEUTISTS.

BY

R. EGLESFELD GRIFFITH, M.D.

A NEW EDITION.



CAREFULLY REVISED AND MUCH EXTENDED

BY

ROBERT P. THOMAS, M.D.

WITH ILLUSTRATIONS.

Selecta sunt quæ medicum nobilitant.—LINNÆUS.



PHILADELPHIA:
BLANCHARD AND LEA.
1858.

UNIVERSAL FORMULARY:

CONTAINING THE PRESCRIPTIONS OF

PREPARING AND ADMINISTERING

OFFICIAL AND OTHER MEDICINES

THE NEW YORK

PHYSICIANS AND PHARMACEUTISTS

R. COLLINS, D.D.

Entered, according to Act of Congress, in the year 1854, by

BLANCHARD AND LEA,

in the Clerk's Office of the District Court of the United States for the Eastern
District of Pennsylvania.

COLLINS, PRINTER.

ROYAL COLLEGE OF PHYSICIANS	
CLASS	615.13
NO.	23874
SUBJECT	
DATE	



PHILADELPHIA

BLANCHARD AND LEA

1855

TO

GEORGE B. WOOD, M.D.,

AND

FRANKLIN BACHE, M.D.,

AUTHORS OF

"THE DISPENSATORY OF THE UNITED STATES OF AMERICA,"

THIS WORK

IS RESPECTFULLY DEDICATED,

BY THEIR FRIEND,

R. E. G.

GEORGE R. WOOD, M.D.

GEORGE R. WOOD, M.D.

WILLIAM A. BACH, M.D.

WILLIAM A. BACH, M.D.

THE UNIVERSITY OF THE STATE OF NEW YORK

THE UNIVERSITY OF THE STATE OF NEW YORK

THIS WORK

THIS WORK

IS A REPRINT OF THE ORIGINAL

AS PUBLISHED BY THE AUTHOR

IN 1880

NEW YORK

NEW YORK

P R E F A C E

TO THE NEW EDITION.



IN preparing a new edition of the "Universal Formulary," the Editor has endeavored to carry out, as far as possible, the views of its lamented Author. Great care has been exercised to avoid those errors which almost inevitably occur in a work of this kind, deriving its materials, as it does, from a variety of sources, and from different languages. The additions are numerous, amounting to about seventy pages. Those made to the Formulary proper have the names of the authorities attached to them, and those to the practical remarks, in the latter part of the work, are enclosed in brackets []. The last editions of the United States, the London, and the Dublin Pharmacopœias, as well as of *L'Officine*, an invaluable work on practical Pharmacy, by Dorvault, of Paris, have been freely used. The Editor has also availed himself of the valuable processes and formulæ that have appeared within the past three years both in the Pharmaceutic and Medical Journals, and most of them will be found under the appropriate heads.

Besides the strictly medical formulæ, many have been added from authentic sources for the preparation of essences, perfumes, inks, soaps, varnishes, &c. &c., which the country Pharmaceutist often finds desirable to manufacture for himself.

PHILADELPHIA, *May*, 1854.

P R E F A C E.

THE design of this work is to present a compendious collection of formulæ and pharmaceutic processes, with such additional information as may render it useful to the physician and apothecary; and the principal aim has been to select materials most generally applicable, and of practical utility. The sources from which they have been derived are very numerous, as will be seen by a reference to the various authorities cited.

The introduction contains tables and observations on the weights and measures employed for pharmaceutical purposes in the United States and in foreign countries, and an explanation or vocabulary of the principal abbreviations and Latin terms used by physicians in writing prescriptions, followed by observations on the management of the sick-room, with rules for the administration of the different classes of medicines.

The formulary is arranged alphabetically, according to the pharmaceutic names adopted in the United States Pharmacopœia; but in each formula, the English appellations for the articles composing it are used, and the quantities of these ingredients are expressed in words, and not in the usual pharmaceutic signs.

These innovations may, and probably will, be objected to by many; but we feel convinced that a change has become requisite, and that fewer mistakes would be committed by physicians in writing prescriptions, both in the names of the ingredients and in the quantities, were they given at length, and in common language, instead of in the abbreviated cabalistic terms now used. In France this change has been made, and it is full time that other countries should follow her example.

In this portion of the work, the compiler has derived much important assistance from several of his friends, and is under great obligation to Mr. Wm. Procter, Jr., for numerous formulæ and many useful suggestions. He, also, is much indebted to Dr. Robert Bridges for his attentive revision of the sheets, as well as for some important corrections.

The various tables which follow, it is hoped, will add to the value of the work. The observations and directions on officinal preparations are given in as concise a form as possible; and are, for the most part, condensed from the excellent edition of *Mohr and Redwood's Pharmacy*, as edited by Mr. Wm. Procter, Jr. To this is added a short view of the action of poisons, with the best means of obviating their effects.

To facilitate a reference to the contents of the work, copious indices have been added, not only of the formulæ, but of the diseases for which they have been advised.

In the botanical portion, the author has been obliged to cite, in almost every instance, his "Medical Botany." This has arisen from the fact, that no other work of a general character on medicinal plants has been issued from the American press.

PHILADELPHIA, March, 1850.

TABLE OF CONTENTS.

	PAGE
PREFACES	7, 9
INTRODUCTION	17
WEIGHTS AND MEASURES.....	17
Weights of the United States and Great Britain.....	17
Dublin Weights	20
Foreign Weights.....	20
Measures	24
SPECIFIC GRAVITY.....	28
TEMPERATURES FOR CERTAIN PHARMACEUTICAL OPERATIONS.....	36
HYDROMETRICAL EQUIVALENTS.....	36
SPECIFIC GRAVITIES OF SOME OF THE PREPARATIONS OF THE PHARMACOPŒIAS.....	41
RELATION BETWEEN THE DIFFERENT THERMOMETRICAL SCALES.....	42
EXPLANATION OF PRINCIPAL ABBREVIATIONS USED IN FORMULÆ.....	43
VOCABULARY OF WORDS EMPLOYED IN PRESCRIPTIONS	44
OBSERVATIONS ON THE MANAGEMENT OF THE SICK ROOM.....	55
Ventilation of the Sick Room.....	55
Temperature of the Sick Room.....	56
Cleanliness in the Sick Room.....	57
Quiet in the Sick Room.....	57
Examination and Preservation of the Excretions.....	58
Administration of Medicine.....	58
Furniture of a Sick Room.....	58
Proper use of Utensils for Evacuations.....	59
DOSES OF MEDICINES.....	60
Age.....	61
Sex	61
Temperament.....	61
Idiosyncrasy.....	61
Habit.....	61
State of the System.....	62
Time of Day.....	62
Intervals between Doses.....	63
RULES FOR ADMINISTRATION OF MEDICINES.....	63
Acids.....	64
Antacids.....	64

	PAGE
Antilithics and Lithontriptics.....	64
Antispasmodics.....	64
Anthelmintics.....	65
Cathartics.....	65
Enemata.....	66
Suppositories.....	66
Demulcents or Emollients.....	67
Diaphoretics.....	67
Diluents.....	68
Diuretics.....	68
Emetics.....	68
Emmenagogues.....	69
Epispastics.....	69
Errhines.....	71
Escharotics.....	71
Expectorants.....	71
Narcotics.....	72
Refrigerants.....	72
Sedatives.....	72
Sialagogues.....	72
Stimulants.....	73
Tonics.....	73
MANAGEMENT OF CONVALESCENCE AND RELAPSES.....	74
UNIVERSAL FORMULARY.....	75
DIETETIC PREPARATIONS NOT INCLUDED AMONG THE PREVIOUS PRESCRIPTIONS.....	487
LIST OF INCOMPATIBLES.....	492
POSOLOGICAL TABLE OF THE MOST IMPORTANT MEDICINES.....	496
TABLE OF PHARMACEUTICAL NAMES WHICH DIFFER IN THE UNITED STATES AND BRITISH PHARMACOPŒIAS.....	503
OFFICINAL PREPARATIONS AND DIRECTIONS.....	514
INTERNAL REMEDIES.....	514
Powders.....	514
Pills and Boluses.....	515
Extracts.....	517
Confections, Conserves, Electuaries.....	518
Pulps.....	519
Syrups.....	519
Mellites or Honeys.....	520
Infusions.....	520
Decoctions.....	520
Tinctures.....	521
Wines.....	523
Vinegars.....	524
Mixtures.....	524
Medicated Waters.....	525
Distilled, Essential, or Volatile Oils.....	528
Fixed Oils and Fats.....	533

TABLE OF CONTENTS.

xiii

	PAGE
Alkaloids.....	533
Spirits.....	534
Troches or Lozenges.....	534
Inhalations.....	535
EXTERNAL REMEDIES.....	535
Baths.....	535
Cold Bath.....	535
Cool Bath.....	536
Temperate Bath.....	536
Tepid Bath.....	536
Warm Bath.....	536
Hot Bath.....	537
Shower Bath.....	537
Local Baths.....	537
Vapor Bath.....	538
Warm Air Bath.....	538
Douches.....	538
Medicated Baths.....	539
Affusion.....	539
Sponging.....	540
Fomentations.....	541
Cataplasms, or Poultices.....	541
Lotions, Liniments, Embrocations.....	542
Vesicatories, or Blisters.....	543
Issues.....	543
Setons.....	543
Ointments.....	544
Cerates.....	544
Plasters.....	545
Fumigations.....	546
BLOOD-LETTING.....	547
General Blood-letting.....	548
Venesection.....	548
Arteriotomy.....	550
Topical Blood-letting.....	550
Cupping.....	550
Leeching.....	551
Scarifications.....	552
POISONS.....	553
INDEX OF DISEASES AND THEIR REMEDIES.....	569
INDEX OF PHARMACEUTICAL AND BOTANICAL NAMES.....	587
GENERAL INDEX.....	595

INTRODUCTION.

BEFORE entering upon the main object of the work, some introductory observations are required, on the forms in which remedial substances are administered, the processes to be pursued in their preparation, and on the weights and measures employed in the proper apportionment of the ingredients, and in the regulation of the doses to be administered.

WEIGHTS AND MEASURES.

These vary much in different countries, and even in the same country. Thus, in Great Britain and the United States, there are three standards of weight recognized and employed: the Avoirdupois, the Troy, and the Apothecaries'.

WEIGHTS OF THE UNITED STATES AND GREAT BRITAIN.

AVOIRDUPOIS WEIGHT.

This is the common standard employed in the selling or buying of such articles as are measured by weight, and is also in use among apothecaries in this country, except in the compounding and dispensing of medicines.

1 drachm	Equal to troy grains.
16= 1 ounce	27.34375
256= 16= 1 pound	437.5
3584= 224= 14= 1 stone	7000.
28672= 1792= 112= 8= 1 hundred weight	98000.
473440=35840=2240=160=20=1 ton.....	784000.
	15680000.

In weighing many articles, what is called the short ton, or 2000 pounds, is employed.

TROY WEIGHT

Is used in the sale of gold, silver, platina, and precious stones.

1 grain
24= 1 pennyweight
480= 20= 1 ounce
5760=240=12=1 pound.

The following are the relative proportions between the troy and the avoirdupois:—
One pound troy is equivalent to 0.822857 pound avoirdupois, or 13 ounces, 2 drachms, 1 scruple, 8.7 grains. [13 ounces, 72.5 grains.]

One ounce troy is equivalent to 1 ounce, 1 drachm, 1 scruple, 6.225 grains avoirdupois.

One pound avoirdupois is equivalent to 1.215277 pound troy, or 1 pound, 2 ounces, 4 drachms, and 2 scruples.

One ounce avoirdupois is equivalent to 7 drachms, 17.5 grains troy.

144 pounds avoirdupois are equivalent to 175 pounds troy.

175 ounces troy are equivalent to 192 ounces avoirdupois.

The following table of equivalents in troy and avoirdupois weight, derived from Redwood's edition of Gray's Supplement to the Pharmacopœia, will be found useful in making these calculations.

EQUIVALENTS IN TROY AND AVOIRDUPOIS WEIGHT.

Troy grains.	TROY.				AVOIRDUPOIS.		
	lbs.	oz.	drs.	grs.	lbs.	oz.	grs.
60			1				60
120			2				120
240			4				240
437.5			7	17.5		1	
480		1				1	42.5
875		1	6	35		2	
960		2				2	85
1312.5		2	5	52.5		3	
1440.		3				3	127.5
1750.		3	5	10		4	
1920.		4				4	170
2187.5		4	4	27.5		5	
2400		5				5	212.5
2625.0		5	3	45.		6	
2880.		6				6	255
3062.5		6	3	2.5		7	
3360		7				7	297.5
3500		7	2	20.		8	
3840		8				8	340
3937.5		8	1	37.5		9	
4320.		9				9	382.5
4375.		9	0	55.		10	
4800		10				10	425
4812.5		10		12.5		11	
5250		10	7	30.		12	
5280		11				12	30
5687.5		11	6	47.5		13	
5760.	1					13	72.5
6125.	1	0	6	5.		14	
6562.5	1	1	5	22.5		15	
7000	1	2	4	40.	1		
7680	1	4			1	1	242.5
9600	1	8			1	5	422.5
10500	1	9	7		1	8	
11520	2				1	10	145
14000	2	5	1	20	2		
17280	3				2	7	217.5
21000	3	7	6	0	3		
23040	4				3	4	290
28000	4	10	2	40	4		
28800	5				4	1	362.5
34560	6				4	14	435
35000	6	0	7	20	5		
40320	7				5	12	70
42000	7	3	4	0	6		
46080	8				6	9	142.5
49000	8	6	0	40	7		
51840	9				7	6	215
56000	9	8	5	20	8		
57600	10				8	3	287.5
63000	10	11	2	0	9		

Troy grains.	TROY.				AVOIRDUPOIS.		
	lbs.	oz.	drs.	grs.	lbs.	oz.	grs.
63360	11				9	0	360
69120	12				9	13	432.5
70000	12	1	6	40	10		
74880	13				10	11	67.5
77000	13	4	3	20	11		
80640	14				11	8	140
84000	14	7	0	0	12		
86400	15				12	5	212.5
91000	15	9	4	40	13		
92160	16				13	2	285
97920	17				13	15	357.5
98000	17	0	1	20	14		
103680	18				14	12	420
105000	18	2	6	0	15		
109440	19				15	10	65
112000	19	5	2	40	16		
115200	20				16	7	137.5
119000	20	7	7	20	17		
120960	21				17	4	210
126000	21	10	4	0	18		
126720	22				18	1	282.5
132480	23				18	14	355
133000	23	1	0	40	19		
138240	24				19	11	427.5
140000	24	3	5	20	20		
144000	25				20	9	62.5
147000	25	6	2	0	21		
149760	26				21	6	135
154000	26	8	6	40	22		
155520	27				22	3	207.5
161000	27	11	3	20	23		
161280	28				23	0	280
167040	29				23	13	352.5
168000	29	2	0	0	24		
172800	30				24	10	425
175000	30	4	4	40	25		
178560	31				25	8	59
182000	31	7	1	20	26		
184320	32				26	5	131.5
189000	32	9	6	0	27		
190080	33				27	2	204
195840	34				27	15	276.5
196000	34	0	2	40	28		
201600	35				28	12	149
203000	35	2	7	20	29		
207360	36				29	9	421.5
210000	36	5	4	0	30		
230400	40				32	14	275
280000	48	7	2	40	40		
288000	50				41	2	125
345600	60				49	5	412.5
350000	60	9	1	20	50		
403200	70				57	9	262.5
420000	72	11	0	0	60		

Troy grains.	TROY.				AVOIRDUPOIS.		
	lbs.	oz.	drs.	grs.	lbs.	oz.	grs.
460800	80				65	13	113
490000	85	0	6	40	70		
518400	90				74	0	400.5
560000	97	2	5	20	80		
576000	100				82	4	250.5
630000	109	4	4	0	90		
645120	112				92	2	245
700000	121	6	2	40	100		
784000	136	1	2	40	112		

When applied to the compounding or dispensing of medicines, this standard of weight is known as Apothecaries' weight, and differs from the last in the subdivision of the ounce, viz :—

APOTHECARIES' WEIGHT.

1 grain
 20 = 1 scruple
 60 = 3 = 1 drachm
 480 = 24 = 8 = 1 ounce
 5760 = 288 = 96 = 12 = 1 pound.

To designate these divisions, the following marks are generally used in prescriptions: a grain, gr.; a scruple, ℥; a drachm, ℥; an ounce, ℥; a pound, ℥.

DUBLIN WEIGHTS.

Adopted by the Dublin College in the edition of their Pharmacopœia for 1850 :—

1 grain
 18.22 = 1 scruple
 54.68 = 3 = 1 drachm
 437.5 = 24 = 8 = 1 ounce
 7000. = 384 = 128 = 16 = 1 pound.

FOREIGN WEIGHTS.

FRANCE.—Anterior to the French Revolution of 1789, the scale of weight used was the *poids de marc*, the unit of which was the pound of Charlemagne, which was equivalent to 7561 Troy grains, and was divided as follows :—

POIDS DE MARC.

	Troy grains.	Grammes.
1 grain	0.8203	0.0531
24 = 1 denier	19.687	1.274
72 = 3 = 1 gros	59.070	3.824
576 = 24 = 8 = 1 once	472.542	30.594
4608 = 192 = 64 = 8 = 1 marc	3780.500	244.753
6912 = 288 = 96 = 12 = 1 livre medicinal	5670.750	367.129
9216 = 384 = 128 = 16 = 1 livre marchand or poid de marc	7561.000	489.505

When the decimal system was introduced by the National Assembly, a new series of measures was adopted, termed the metrical, in which the *metre*, or the ten-millionth part of a quarter of the meridian of the earth, is taken as the unit; this is divided into ten parts, each of which is called a *decimetre*, which in turn is divided into ten *centimetres*. A cubic decimetre is assumed as the unit of measures of capacity, and termed a *litre*. The unit of weight is called a *gramme*, and is a cubic centimetre of distilled water at 39.5 Fahr.

METRICAL WEIGHT.

						Troy grains.
1 milligramme.....	=					.015434
10 = 1 centigramme.....	=					.15434
100 = 10 = 1 decigramme.....	=					1.5434
1000 = 100 = 10 = 1 gramme.....	=					15.434
10000 = 1000 = 100 = 10 = 1 decagramme.....	=					154.34
100000 = 10000 = 1000 = 100 = 10 = 1 hectogramme.....	=					1543.4
1000000 = 100000 = 10000 = 1000 = 100 = 10 = 1 kilogram.....	=					15434.

This system of weights, although adopted by the scientific men of France, was not favorably received by the people generally, who obstinately adhered to the old system of the *poids de marc*; and, although many stringent laws were passed, from time to time, to render the metrical system obligatory, the government in 1812 was forced to make an attempt to amalgamate the two systems, by altering the *livre*, and making it equal to half a kilogramme, assuming this as the unit, and calculating the other divisions from it, according to the old nomenclature. The following table shows the equivalent of the two scales and of avoirdupois:—

FRENCH WEIGHTS OF 1812.

French weights of 1812.	Metrical weight. grammes.	lb.	oz.	dr.	grains.
1 livre =	500	= 1	1	10	6.06
$\frac{1}{2}$ " =	250	=	8	13	3.03
$\frac{1}{4}$ " =	125	=	4	6	15.19
2 once =	62.5	=	2	3	7.60
1 " =	31.25	=	1	1	17.47
$\frac{1}{2}$ " =	15.625	=		8	22.40
2 gros =	7.812	=		4	11.20
1 " =	3.906	=		2	5.60
$\frac{1}{2}$ " =	1.9531	=		1	2.80
1 grain =	0.0542	=			0.837

The adoption of this system was not made obligatory upon the pharmaciens by law until the year 1827; and indeed it appears never to have been generally adopted, the greatest confusion having prevailed with regard to the weights and measures used in the preparation of medicine, as well as in commerce generally, up to the year 1840. In July, 1837, a law was passed which definitively abolished the use of all other weights and measures, excepting those of the metrical or decimal system, from and after August, 1840. The METRICAL WEIGHT, therefore, is now the only one permitted to be used throughout France.

The exact proportion of troy weight to the metrical standard has never been fully and exactly ascertained. In the table previously given, the kilogramme has been considered as equivalent to 15434 grains troy, which is the usual estimate of it. Francœur, however, estimates the pound troy as equal to 392.9986 grammes; Matthieu, Legendre, and Duborg, as equal to 373.0956 grammes; Chelius and Houschild to 373.243 grammes. But the most correct estimation appears to be that of Lochman, who considers 9216 grains poid de marc as equivalent to 7555 grains troy; the pound troy as equal to 7026.320 grains marc; and the kilogramme as equal to 15434 grains troy.* The subjoined tables are calculated on this estimation:

VALUE OF TROY WEIGHT IN METRICAL WEIGHT.

One pound.....	=	373.202	grammes
One ounce.....	=	31.10017	"
One drachm.....	=	3.887521	"
One scruple.....	=	1.295840	"
One grain.....	=	0.06479201	"

* By examination at the English mint the gramme is determined to be 15.434 Troy grains.

VALUE OF TRÖY WEIGHT IN MARC WEIGHT.

One pound	=	12 ounces,	1 gros,	42.32	grains
One ounce	=	1 once		9.53	"
One drachm	=		1 "	1.19	"
One scruple	=			24.40	"
One grain	=			1.22	"

The other European States differ much in their medicinal weights from the troy standard; but it is superfluous to examine each in detail, the following synopsis and table being sufficient to show the variations.

1. The medicinal pound, except in Dublin and Turkey, and by the old poids de marc, is in all of them divided into 12 ounces.

2. In all of them the ounce is divided into 8 drachms, except in Naples, where it contains 10.

3. In all, the ounce is divided into 3 scruples.

4. But the value of the scruple differs. It is composed of 24 grains at Bologna, Coni, Lucca, Modena, Parma, Rome, and in Spain, France, Portugal, Tuscany, and Sardinia.

Of 20 grains in Great Britain, (excepting Dublin, where it is 18.22 grains,) the United States, Austria, Bavaria, Holland, Poland, Prussia, and Sweden; at Lubeck, Naples, Nuremberg, and Venice.

5. Consequently, the pound consists of

5760 grains in the United States, Great Britain, Austria, Bavaria, Holland, Poland, Prussia, and Sweden; at Lubeck, Nuremberg, Naples, and Venice.

6400 grains in Turkey.

6912 grains in Spain, Portugal, Tuscany, Sardinia, Bologna, Lucca, Modena, Parma, Rome, and Coni.

7000 grains in Dublin.

7200 grains at Naples.

9216 grains in France by the poids de marc.

The subjoined table shows the differential values between the troy weight and the medicinal weights of the countries of Europe, calculated in grains:—

COUNTRIES, &c.	Value of the pound, 5760 grains.	Value of the ounce, 480 grains.	Value of the drachm, 60 grains.	Value of the scruple, 20 grains.	Value of the grain.
Austria ⁽¹⁾	5118.09	426.51	53.31	17.77	0.89
Bavaria ⁽²⁾	5971.23	497.60	62.20	20.73	1.04
Bologna.....	7920.93	660.08	82.51	27.50	1.38
Coni.....	8392.40	699.37	87.42	29.14	1.46
Corte.....	7807.42	650.62	81.33	27.11	1.36
Holland ⁽³⁾	5732.38	477.70	59.71	19.90	1.00
Lubeck.....	5823.60	485.30	60.66	20.22	1.01
Lucca.....	7711.73	642.64	80.33	26.78	1.34
Modena.....	7576.80	631.40	78.92	26.31	1.32
Monticelli ⁽⁴⁾	8334.64	694.55	86.82	28.94	1.45
Naples ⁽⁵⁾	8377.11	698.09	87.26	29.09	1.46
Nuremberg ⁽⁶⁾	6007.22	500.60	62.58	20.86	1.04

⁽¹⁾ This weight is used in Austria Proper, Bohemia, Moravia, Hungary, Transylvania, Tyrol, and the Lombardo-Venetian kingdom, with the exception of the city of Venice.

⁽²⁾ This weight has been adopted in Greece.

⁽³⁾ This weight is also used in Belgium.

⁽⁴⁾ Monticelli, Castelvetro, and Polesine.

⁽⁵⁾ Kingdom of the Two Sicilies.

⁽⁶⁾ This weight is no longer used in Nuremberg; this city, being now incorporated in the

COUNTRIES, &c.	Value of the pound, 5760 grains.	Value of the ounce, 480 grains.	Value of the drachm, 60 grains.	Value of the scruple, 20 grains.	Value of the grain.
Parma ⁽⁷⁾	7864.55	655.38	81.92	27.31	1.37
Piacenza ⁽⁸⁾	8124.20	677.02	84.63	28.21	1.41
Poland.....	5996.04	499.67	62.46	20.82	1.04
Portugal.....	7494.62	624.55	78.07	26.02	1.30
Prussia ⁽⁹⁾	6128.12	510.68	63.83	21.28	1.06
Rome	7607.72	633.98	79.25	26.42	1.32
Spain.....	7475.46	622.96	77.87	25.96	1.30
Sweden.....	6034.48	502.87	62.86	20.95	1.05
Turin.....	7770.71	647.56	80.94	26.98	1.35
Turkey.....	7436.78	619.73	77.47	25.82	1.29
Tuscany	7597.21	633.10	79.14	26.38	1.32
Venice.....	7136.23	594.69	74.34	24.78	1.24

kingdom of Bavaria, has adopted the Bavarian standard. But the Nuremberg weight is still extensively used in Germany and other countries. In Germany, we find it in the duchies of Saxe-Altenburg, and of Anhalt Bernburg; grand duchy of Baden; city of Bremen; duchies of Brunswick, Saxe Meiningen, and Saxe Coburg Gotha; cities of Frankfort-on-the-Maine and Hamburg; kingdom of Hanover; electorate of Hesse, grand duchy of Hesse, landgraviate of Hesse Homburg; principality of Hohenzollern Sigmaringen; duchy of Nassau; grand duchy of Oldenburg; principalities of Reuss-Plauen and Reuss-Schleitz, of Schwartzburg-Rudolstadt, and Schwartzburg-Sonderhausen; duchy of Sleswick-Holstein: principality of Waldeck; grand duchy of Saxe Weimar; and kingdom of Wirtemberg.

The Nuremberg standard is also adopted in Denmark, Norway, Finland, Russia, and Switzerland.

⁽⁷⁾ Parma, Borgo San-Donnino, Pellegrino, San-Secondo, Fontanellato, and Bussetto.

⁽⁸⁾ Piacenza, Fiorenzola, and Carpaneta.

⁽⁹⁾ This standard has been adopted in the kingdom of Saxony, the principalities of Lippe-Detmold and Lippe-Schaumberg, and the duchy of Anhalt-Dessau. It differs so little from those of Anhalt-Coethen and the grand duchies of Mecklenburg-Schwerin and Mecklenburg-Strelitz, that the standard of these three territories may be considered identical with that of Prussia.

This table needs hardly any explanation: it will enable the pharmacist—in all cases of foreign formulæ—to calculate the foreign weight in troy weight; thus, *e. g.*—one grain troy being equal to 0.89 grain of Austria—the value of one grain Austrian weight in troy weight may be readily ascertained by a simple rule of three example:—

$$0.89 : 1 :: 1$$

$$1 \times 1 = \times 1 = 0.100$$

$$0.89)0.100(1.1235$$

89

110

89

210

178

320

267

530

445

65

The value of these various pounds in grammes is thus calculated by Jourdain :—

The pound of 5760 grains.				
Venice.....	= 301.230	grammes	Piacenza = 317.577 grammes	
Prussia.....	= 350.761	"	Bologna = 325.665 "	
Sweden.....	= 356.227	"	Soragna = 325.800 "	
Nuremberg.....	= 357.843	"	Parma = 328.000 "	
Poland.....	= 358.510	"	Corte..... = 330.400 "	
Bavaria.....	= 360.000	"	Turin..... = 331.961 "	
Lubeck.....	= 369.126	"	Lucca = 334.500 "	
United States and } Great Britain }	= 373.202	"	Rome = 339.073 "	
Holland.....	= 375.000	"	Tuscany = 339.542 "	
Austria.....	= 420.009	"	Modena = 340.457 "	
The pound of 6400 grains.				
Turkey.....	= 321.317	"	Portugal = 344.190 "	
The pound of 6912 grains.				
Coni.....	= 307.370	"	Spain = 345.072 "	
Monticelli.....	= 307.370	"	The pound of 7200 grains.	
			Naples.....	= 320.230 "
			The pound of 9216 grains.	
			France.....	= 489.503 "

LIQUID MEASURES OF THE UNITED STATES AND GREAT BRITAIN.

The liquid measures employed by the apothecaries in the United States, are the wine gallon and its sub-divisions.

WINE OR APOTHECARIES' MEASURE.

(Adopted in the United States Pharmacopœia.)

	Cubic Inches.	Grains Troy.
1 minim, m		0.95
60 = 1 fluidrachm, f. 3.....	= 0.2256	= 56.96
480 = 8 = 1 fluidounce, f. 3.....	= 1.8047	= 455.69
7680 = 128 = 16 = 1 pint, O.....	= 28.875	= 7291.11
61440 = 1024 = 128 = 8 = 1 gallon, Cong. = 231.		= 58328.88

For a long time, the Royal Colleges of Physicians in England, Scotland, and Ireland declined to recognize the use of measures in preparing and dispensing medicines, on the ground that the varying densities of different fluids rendered it difficult to use one common measure for all without risk of serious errors. But as druggists and medical practitioners constantly employed measures, in defiance of the prohibition, and as the practice, besides being attended with great and obvious convenience, was found to be less fraught with danger than had been conceived, the London College in the first instance, and subsequently the other colleges also, recognized a system of measures of their own, founded on the standard measures of the country. The Colleges of London and Edinburgh, adopting the imperial pint of 1826 as the basis, divide it into twenty parts, called fluidounces; each of which corresponds exactly with an avoirdupois ounce of distilled water at 62° F. and 30° bar., and therefore contains 437.5 grains troy. The fluidounce is subdivided into eight parts of 54.6875 grains, termed fluidrachms; and each of these consists of sixty parts, called minims, which therefore amount each in weight to 0.91146 troy grains of distilled water.

[The Dublin College, at the last revision of its Pharmacopœia in 1851, adopted the same standard for measures; so that all the British Colleges now use the Imperial pint and its subdivision of twenty fluidounces.—ED.]

IMPERIAL MEASURE.

(Adopted by all the British Colleges.)

			Grains Troy.	Avoir.
1 minim	=	0.91	
60 =	1 fluidrachm	54.7	
480 =	8 " 1 fluidounce	437.5	= 1 oz.
9600 =	160 " = 20 " = 1 pint	8750.	= 1.25 lb.
76800 =	1280 " = 160 " = 8 " = 1 gallon	70000.	= 10 lbs.

VALUE OF WINE OR APOTHECARIES' MEASURE IN IMPERIAL MEASURE.

Wine Measure.		Imperial Measure.		
	Pints.	Fluidounces.	Fluidrachms.	Minims.
1 gallon = 6	13	2	23
1 pint =	16	5	18
1 fluidounce =	1	0	20
1 fluidrachm =		1	2.5
1 minim =			1.04

VALUE OF IMPERIAL MEASURES IN WINE OR APOTHECARIES' MEASURE.

	Gallon.	Pints.	Fluidounces.	Fluidrachms.	Minims.
1 gallon 1	1	9	5	8
1 pint	1	3	1	38
1 fluidounce			7	41
1 fluidrachm				58
1 minim				0.9

Besides these regular and authorized measures, there are others constantly used in the preparation and administration of medicines, which require notice. These, which have been adopted for convenience, are far from being uniform, but may be used without danger for ordinary purposes. Drs. Wood and Bache estimate the

Teacup	as containing about	four fluidounces.
Wineglass	"	two fluidounces.
Tablespoon	"	a half a fluidounce.
Teaspoon	"	a fluidrachm.

Dr. Christison gives a somewhat different calculation; he considers a

Tumbler	to contain	eight fluidounces.
Breakfast cup	"	do do
Teacup	"	five fluidounces.
Wineglass	"	two fluidounces.
Tablespoon	"	half a fluidounce.
Dessertspoon	"	two fluidrachms.
Teaspoon	"	one fluidrachm.

In both these estimates, the teaspoon is rated too low; they apply very well to the teaspoon formerly used, but not to the much larger kind now in general use, which approaches the dessertspoon in capacity. Quart and pint bottles are also sometimes employed as measures of capacity for medical purposes; but they are very fallacious guides of quantity, as the so-called quart bottle never holds a quart, nor the misnamed pint bottle a pint. The first seldom holds more than twenty-six to twenty-seven fluid ounces, and the latter from thirteen to fourteen.

It is still more common to estimate small quantities or doses of medicine by drops, as representing, and equivalent to, minims. This is so convenient, that it is not likely to be abandoned, though nothing can be more erroneous, as the size of the

drop of the same fluid varies much with the form of the mouth of the bottle, its size, the mode in which the operation is performed, and still more from the nature of the fluid thus attempted to be measured. Numerous experiments have been made on the subject, but can only be considered as approximate estimates, though sufficiently accurate for practical purposes.

Mr. Alsop, of London, some years since, published the result of some experiments made by him, which show the great variation to which this mode of measuring is liable. The following proves that the number of drops required to measure a fluidrachm are widely different when dropped from a large or a small bottle.

One fluidrachm.	Large bottle.	Small bottle.
Diluted sulphuric acid	24 drops	84 drops
Scheele's hydrocyanic acid	35 "	70 "
Distilled water	31 "	54 "
Solution of ammonia	40 "	48 "
Tincture of opium	84 "	135 "
Rectified spirit	100 "	130 "
Tincture of muriate of iron	100 "	150 "

Mr. Durand, of Philadelphia, in an able paper on the subject (*Journ. Phila. Col. of Pharm.*, vol. i. p. 165), says, "The bulk of drops depends not only on the density of the liquids which furnish them, and the cohesion of the constituent particles of that liquid, but also on the shape of the mouth of the vessel from which they are poured. An open vessel with a beak, such as the common graduated measure, affords a larger drop than a bottle with the stopper half drawn out; a mode commonly practiced. That furnished by the dropping tube is still smaller, and is ever liable to vary with the greater or lesser diameter of its extremity. Besides, in every instance, the first drops poured from any vessel are always smaller than those subsequently obtained."

He goes on to say that the following may be considered as established:—

1. That liquids which contain a small proportion of water afford a small drop; while, on the contrary, liquids containing a large quantity of water furnish a large drop. For instance, concentrated acids, ethers, rectified alcohol, fixed and essential oils, &c., which contain a very small proportion of water, yield a smaller drop than diluted acids, weak alcohol, wine, &c.

2. That, among liquids containing a large proportion of water, those which are not charged with remedial substances give a larger and heavier drop than those same liquids containing extraneous bodies in solution. Thus, weak alcohol, wine, vinegar, and water furnish a larger and heavier drop than the tinctures prepared from them.

The following table illustrates his remarks:—

TABLE

Showing the Difference between Minims, Drops, and Grains of various Medicinal Liquid Preparations of the Pharmacopæia of the United States, &c.

PREPARATIONS.	No of drops in 20 minims.	No. of min. in 20 drops.	No. of drops in 20 grains.	No. of grains in 20 drops.
Sulphuric acid	30	13.3	25	16
Sulphuric ether.....	50	8	60	6
Rectified alcohol	46	8.6	57	7.1
Nitric acid	28	14.2	22.2	18
Acetic acid (crystallizable)	40	10	40	10
Muriatic acid	18	22.2	18.1	22

PREPARATIONS.	No. of drops in 20 minims.	No. of min. in 20 drops.	No. of drops in 20 grains.	No. of grains in 20 drops.
Oil of wormseed (chenopod. anthelminticum).....	40	10	50	8
of peppermint, aniseed, sweet almond, olive, palma christi	40	10	43.5	9
of cloves.....	40	10	36	11
of cinnamon	40	10	32	12.5
Copaiba.....	40	10	40	10
Diluted alcohol.....	40	10	42	9.5
Tincture of hydriodate of potassa, cantharides, kino, digitalis, assafetida, sulphuric acid, colchicum, opium, valerian, guaiacum	40	10	43	9.3
of valerian guaiacum (volatile).....	40	10	50	8
of muriate of iron.....	44	9.1	50	8
Wine, Teneriffe.....	26	15.3	25	16
antimonial	24	16.6	26	15.3
of opium (Sydenham's laudanum)	26	15.3	29	13.7
of colchicum root and seeds	25	16	29	13.7
Vinegar, distilled.....	19	21	20	20
of opium (black drop) } of colchicum } of squill..... }	26	15.3	25	16
Water, distilled.....	15	26.6	17.5	24.5
solution of hydrocyanic acid	15	26.6	17.5	24.5
sulphuric acid (1 to 7)	17	23.5	17	23.5
nitric " "	17	23.5	17	23.5
ammonia (strong)	18	22.2	18.5	22
" (weak).....	15	26.6	20	20
hydriodate of potassa	18	22.2	20	20
arsenite of potassa	19	21	20	20

The only mode to obviate these discrepancies is for the physician to order the administration of small quantities of fluid medicines in minims, and not in drops, which would induce the use of a minim measure in every sick room.

FRENCH MEASURES OF CAPACITY—APOTHECARIES' MEASURE

1 millilitre	=	16.2318 minims.
10= 1 centilitre	=	2.7053 fl. dr.
100= 10= 1 decilitre	=	3.3816 fl. ounces
1000= 100= 10= 1 litre	=	2.1135 pints.
10000= 1000= 100= 10= 1 decalitre.....	=	2.6419 gallons.
100000= 10000= 1000= 100= 10= 1 hectolitre	=	26.4190 "
1000000= 100000= 10000= 1000= 100= 10= 1 kilolitre..	=	264.1900 "

Litres.	Eng. cubic inches.	Imperial pints.	Wine pints.	Troy ounces of water.
1 =	61.028 =	1.7608 =	2.1135 =	32.104
2 =	122.056 =	3.5216 =	4.2270 =	64.208
3 =	183.084 =	5.2822 =	6.3405 =	96.312
4 =	244.112 =	7.0430 =	8.4541 =	128.416
5 =	305.140 =	8.8038 =	10.5676 =	160.520
6 =	366.168 =	10.5646 =	12.6811 =	192.624
7 =	427.196 =	12.3253 =	14.7947 =	224.728
8 =	488.224 =	14.0861 =	16.9082 =	256.832
9 =	549.252 =	15.8469 =	19.0217 =	288.936

The former measure in use before the introduction of the metrical system was the *pint*, and its subdivisions; and it still continues to be employed in some parts. It agrees with the metrical measure as follows:—

1 demi-poisson	=	62.50 grammes	=	0 litre	0 decilitre	6.25 centilitre.
1 poisson	=	125	=	0 “	1 “	2.5 “
1 demi-setier	=	250	=	0 “	2 “	5 “
1 chopine	=	500	=	0 “	5 “	0 “
1 pinte	=	1000	=	1 “	0 “	0 “

The subjoined is a list of the principal liquid measures of Europe, with their corresponding value in the metrical measure of France:—

AUSTRIA.—The fundamental unity is the *mass* or *kanne* = 71.3343 Par. cubic inches, or 1.415015 litre. The mass is divided into 3 *seidel*, and 40 mass make one *eimer*.

BADEN.—The fundamental unity is the *mass* = 1.5 decimetre cubic, and accordingly equivalent to 1.5 litre.

BAVARIA.—The fundamental unity is the *mass* = 43 decimal cubic inches of Bavaria, or 1.06921728 litre. The mass is divided into 4 *schoppen*.

DENMARK.—The fundamental unity is the *pott* ($\frac{1}{32}$ part of a cubic foot Danish), equivalent to 0.96529 litre. The pott is divided into four *pegel*, and 2 potts make a *kanne*.

GRAND DUCHY OF HESSE.—The fundamental unity is the *mass*, which corresponds exactly to 2 litres. The mass is divided into 4 *schoppen*.

NAPLES.—The unity is the *barile*, equivalent to 43.6216 litres.

PORTUGAL.—The unity is the *almude*, equivalent to 16.451 litres.

PRUSSIA.—The unity is the *quart*. This measure has a capacity of 64 cubic inches of distilled water, at 61.25 Fahr. and 27° 10' barom., = 57.724 cubic inches of Paris, or 1.145 litre.

ROME.—The unity is the *barile*, equivalent to 58.3416 litres.

RUSSIA.—The unity is the *wedro*, which contains 750 cubic inches of distilled water at 50° Fahr. and 30° barom., = 30 pounds Russian. The wedro is equivalent to 12.28963047 litres, and one litre is accordingly equivalent to 0.081369410776019 wedro. The wedro is divided into 10 *stof*, and the stof into 10 *tschark*.

SPAIN.—The unity is the *arroba*, equivalent to 16.073 litres.

SWEDEN.—The fundamental unity is the *kanna*, which contains 100 cubic decimal inches of pure water at 62° Fahr., *in vacuo*, equivalent to 6.151951 pounds Swedish. One pound Swedish of pure water at 62° Fahr., *in vacuo*, is equal to 0.4250104 litre; the *kanna* is consequently equivalent to 2.6146431552904 litres, or, calculated upon the maximum density of water, 2.617341607126 litres, or, in shorter terms, 2.62 litres. The *kanna* is divided into 2 *stop*, the stop into 4 *quarter*, and the quarter into 4 *jungfrur*.

TURIN.—The unity is the *brenta*, divided into 36 pints, and equivalent to 49.28468 litres.

TUSCANY.—The unity is the *barile*, equivalent to 45.584 litres.

WIRTEMBERG.—The unity is the *mass*, which contains 78.125 cubic inches (Wirtemberg measure), and is equivalent to 1.83705 litre. The mass is divided into four *schoppen*.

SPECIFIC GRAVITY.

The following remarks on Specific Gravity have been taken from Redwood's edition to Gray's Supplement, and comprise in a condensed form much useful information.

The determination of the specific gravity of a body consists in estimating the weight of a given volume of it, as compared with an equal volume of some other body. The bodies usually taken as the standard of comparison are *pure water* for solids and liquids, and *atmospheric air* for gases.

The *specific gravity of a solid* is determined first by weighing it in the ordinary manner with an accurate balance suspended in the air; then attaching a horse-hair, or fine silken thread to the solid body, immersing it in pure distilled water, and weighing it while thus immersed. The weight of the body in air, divided by the difference between its weight in air and its weight in water, will be its specific gravity. Thus a piece of lead is found to weigh 398 grains in air. When immersed in water, its weight is 362.4 grains; and the difference between these two weights, namely, 35.6, is the weight of the volume of water displaced by the lead, or of a volume of water equal to that of the lead. The volume of water being taken as unity, the specific gravity of the lead is found by the following rule of three sum:—

35.6 : 1 :: 398 : 11.176, the specific gravity of the lead.

In taking the specific gravity of a *solid substance lighter than water*, some modification of the process is required; but we have, nevertheless, the same preliminary points to determine; first, the weight of the substance in air; and secondly, the weight of an equal volume of water. This may be illustrated by taking the specific gravity of a piece of wax. The weight of the wax in air is 105.4 grains. On immersing the wax in water, two pressures are exerted—a pressure downwards, equal to the gravity or weight of the wax, and a pressure upwards, equal to the weight of the volume of water displaced by the wax; but the specific gravity of water being greater than that of wax, the upward pressure preponderates, and the wax rises to the surface. Thus, we find that a volume of water equal to that of the wax weighs as much as the wax, and something more. We must ascertain how much more; and this is done in the following manner: Some body heavier than water, and the weight of which in water is known, is attached to the wax, and the two bodies are weighed in water together. A piece of lead may be used for this purpose. The lead, alone, weighs 378 grains in water; with the wax attached to it, the weight in water is 372.4 grains, making a difference of 5.6 grains; and this 5.6 grains is equal to the excess of the upward over the downward pressure on the wax, when immersed in water. Thus, a volume of water equal to that of the wax weighs 5.6 grains more than the wax, or $105.4 + 5.6 = 111$ grains.

Then, $111 : 1 :: 105.4 : 0.949$, the specific gravity of the wax.

It sometimes happens that the solid substance, the specific gravity of which is to be determined, is *in powder*, or *in several small particles*. In such cases, it is found convenient to proceed as in the following method of taking the specific gravity of calomel:—

100 grains of calomel are introduced into a specific gravity bottle, which holds 1000 grains of distilled water; the bottle is filled up with water, and the weight of the contents is found to be 1083.7 grains; deducting the weight of the calomel (100 grains) from this, the remainder (983.7 grains) will be the weight of the water in the bottle, and the difference (16.3 grains) between this and 1000 grains, the weight of the whole contents of the bottle when filled with distilled water, is the weight of a volume of water equal to the volume of the calomel.

Then, $16.3 : 1 :: 100 : 6.03$, the specific gravity of the calomel.

In taking the *specific gravity of substances soluble in water*, other modifications of the process are required. Sometimes the substance may be covered with a thin coating of varnish, so as to protect it from the action of the water. This method answers very well for blue pill, which may be brushed over with a strong tincture of mastic, and then proceeded with as in the case of the lead. In other instances, however, it is necessary to pursue a different course. Thus, any powder that is soluble in water must have its specific gravity taken, in the first instance, with reference to some liquid in which it is not soluble. Spirit of wine, oil of turpentine, or olive oil, may be used in such cases. The process may be illustrated by describing the method of taking the specific gravity of guano in oil of turpentine.

In the first place, the specific gravity of the oil of turpentine is ascertained to be 0.874. Then 100 grains of guano are introduced into a specific gravity bottle, as in the case of the calomel; and the bottle being filled up with oil of turpentine, the weight of the contents is found to be 922.7 grains, from which deducting 100 grains,

the remainder (822.7 grains) will represent the oil not displaced by the guano; and this, deducted from 874 grains, the quantity of oil the bottle is capable of holding, leaves 51.3 grains as the weight of a volume of oil of turpentine equal to that of the guano. Now, $874 : 51.3 :: 1000 : 58.7$, the weight of a volume of water equal to that of the guano.

Then, $58.7 : 1 :: 100 : 1.7$, the specific gravity of the guano.

The methods by which the *specific gravities of liquids* are usually determined may be divided into two classes:—

1st. Those which consist in filling any suitable vessel with the liquid to be estimated, ascertaining the weight of the contents, and dividing this by the weight of the same volume of water.

2d. Those which consist in displacing a portion of the liquid by some solid body floating in it, and estimating the specific gravity according to the weight and volume of the substance immersed, as compared with its immersion in water.

In the first case, the instruments employed are a specific gravity bottle and an ordinary balance.

In the second case, the instruments used may be comprehended under the general terms of *hydrometers* or *aërometers*. These, however, are distinguished from each other, for there are many varieties of them, by different names, according to the particular purpose for which they are respectively intended, or from some peculiarity in their construction.

The *specific gravity bottle* affords the most accurate means of determining the comparative densities of liquids. It consists, usually, of a globular bottle with a flat bottom and a slender neck, which holds exactly 1000 grains of distilled water at a certain fixed temperature. It is very easy at any time to test the accuracy of one of these bottles by a single experiment; and, having ascertained that the bottle is correctly adjusted with regard to distilled water, the indications afforded with any other liquid will be equally trustworthy. The weight in grains of the quantity of any liquid filling such bottle will indicate its specific gravity.

Hydrometers, or *aërometers*, are floating instruments, and their application for the purpose of determining the specific gravities of liquids depends upon the fact, that a body immersed in any liquid, sustains a pressure from below upwards equal to the weight of the volume of the liquid displaced by such body.

The use of hydrometers for determining the specific gravities of liquids has been traced back to a period about 300 years before Christ; an instrument of this kind being described as the invention of Archimedes, the Sicilian mathematician. It subsequently fell into disuse, but was again brought into notice by Basil Valentine.

There are two kinds of hydrometers, which may be taken as the types of all the different varieties in regard to construction:—

1st. Those which are always immersed in the liquids to be tried, to the same depth, and to which weights are added to adjust the instrument to the density of any particular liquid. Of this description are Fahrenheit's, Nicholson's, and Guyton de Morveau's hydrometers.

2d. Those which are always used with the same weight, but which sink into the liquids to be tried to different depths, according to the densities of the liquids. These usually have graduated scales attached to their stems. Of this description are the common glass hydrometers, generally including those of Baumé, Cartier, Gay Lussac, Twaddle, Zannetti, &c., and the specific gravity beads.

Sikes's and Dicus's hydrometers combine the principles of both types, having movable weights and graduated scales.

Hydrometers may also be divided into two classes, as follows:—

First. Those having a general application, for determining the comparative densities of any liquids.

Second. Those intended for special application; as for estimating the comparative strengths of spirits, or the comparative densities of syrups, oils, &c.

Fahrenheit's, Nicholson's, Guyton de Morveau's, and the common glass hydrome-

ters, including Baumé's, Cartier's, Zanetti's, and the specific gravity beads, belong to the first class.

Gay Lussac's, Sikes's, and Dicus's hydrometers, the saccharometer, urinometer, and elæömeter, belong to the second class.

Fahrenheit's hydrometer consists of two glass bulbs blown in a glass tube, like a common hydrometer, excepting that the upper bulb is larger, and the stem, which is small, is terminated at the top in a cup or funnel. It has a mark on the middle of the stem, indicating the point at which the instrument is to be made to float, by means of weights put into the cup.

Nicholson's hydrometer is a modification of Fahrenheit's. It is made of brass, and consists of a hollow globe, to which is fixed a slender stem surmounted by a cup; on the opposite side of the globe is another cup fixed in a kind of stirrup, and loaded, so that this may always form the lowest point of the instrument when immersed in any liquid. There is a mark on the middle of the upper stem, indicating the point at which the instrument is to be made to float. A certain weight is introduced into the cup, to cause the instrument to sink to the proper mark in distilled water. On immersing the hydrometer into any other liquid, more or less weight will have to be put into the cup, according as such liquid is more or less dense than water. Thus the relative densities of liquids are determined.

This instrument is also applicable for taking the specific gravities of solids. If the solid substance be put into the cup, as part of the weight required to sink the hydrometer in distilled water, the weight of the substance in air is ascertained; and if it be then put into the lower cup, immersed in the water, and the instrument again adjusted, its weight in water is ascertained; and from these its specific gravity is calculated.

Guyton de Morveau's hydrometer is similar to Fahrenheit's.

Baumé's hydrometers are used extensively in this country, as well as in France, and are applicable for all kinds of liquids. There are two distinct instruments: one for liquids lighter than water, and the other for liquids heavier than water. The latter, is, for distinction, called the *acidometer* or *saccharometer* (*pèse-acide* or *pèse-sirop*); the former, the *spirit hydrometer* (*pèse-esprit*).

Baumé's acidometer is made in the form of the common hydrometers. It consists of a glass tube terminated at the lower end by two bulbs, the lowest bulb being much smaller than the other, and intended to contain the ballast with which the instrument is loaded. The scale is marked on a slip of paper, or of ivory fixed in the tube, and is adjusted in the following manner: The top of the tube being open, the slip of paper on which the scale is to be marked is put into the stem, and the instrument is then immersed in pure distilled water; quicksilver is now dropped into the lower ball until the instrument sinks so low in the water that only the top of the stem remains above the surface, and a mark is made on the glass denoting exactly the point to which it sinks. The instrument is now taken out of the pure water and put into a solution of fifteen parts of common salt in eighty-five parts of distilled water, this solution being at the same temperature as the water in which the instrument was previously immersed; the point to which it sinks in this solution is to be marked on the stem as before, and the distance between the two marks being taken with a pair of compasses, and transferred to the slip of paper, the first is made the zero or 0, and the other the 15th degree of the scale. This distance being divided into fifteen equal parts or divisions, each division is called a degree, and the scale is completed by adding as many more degrees as the length of the stem will admit of. This being done, the slip of paper is again introduced into its place, and so fixed that the zero (0) of the scale shall be exactly opposite the first mark made on the glass. The end of the stem is now sealed with the flame of a blowpipe.

Baumé's spirit hydrometer is similar in form to the acidometer; but the weight of the instrument, and the scale, are different. In this case, the hydrometer is first immersed, as before, in pure distilled water; but it is made to float, so that the greater

Fig. 1.



Common Hydrometer.

part of the stem shall be above the surface of the water. This point is marked, and the instrument is then transferred to a solution of ten parts of common salt in ninety parts of water, when another mark is made. The distance between these marks is made ten degrees of the scale, which are divided with the compasses, and marked on the slip of paper, as in the other case; the floating point in the solution of salt being made the zero, and the degrees carried upwards from this point.

The temperature at which these instruments were originally adjusted by Baumé, was 10° Reaumur, or 12.5° Centigrade; but those made in England are usually adjusted at 60° Fahrenheit. It is sometimes important to be aware of this difference.

Cartier's hydrometer is much used in France. It is only applicable for liquids lighter than water. This instrument is a modification of Baumé's spirit hydrometer, the form of the instrument being the same, and the same point being taken as the zero of the scale; but the space which, in Baumé's scale, is divided into 32° , is in Cartier's divided into 30° .

It is becoming the common practice in this country to have the scales of hydrometers marked with the specific gravities intended to be indicated, and this is by far the most convenient kind of hydrometer for general use.

Twaddle's hydrometers are much used in Scotland, and occasionally in England. They are made of glass like the common hydrometers, and are sold in sets of six. Each degree on the scale is equal to 0.005 of specific gravity, so that the specific gravity of a liquid is found, with these hydrometers, by multiplying the number of degrees indicated, by 5, and adding 1000. Thus, 10° by Twaddle's hydrometer, $\times 5 + 1000 = 1.050$ specific gravity.

Zanetti's hydrometers, which are made at Manchester, are also sold in sets of six. With these the specific gravity is got by adding a cipher to the number of degrees indicated.

Specific gravity beads, (*fig. 2*) sometimes called *Lovi's beads*, are hollow sealed globes of glass, about the size of small pistol-bullets. Each bead is a small hydrometer, intended to indicate one fixed density, by its remaining half way between the top and the bottom of the liquid into which it is introduced. These beads are sold in sets, each one being marked with the specific gravity, it is to indicate at a certain fixed temperature. They are very useful in making mixtures of any required densities, as, for instance, in making test acids.

Gay Lussac's alcoometre (*fig. 3*) is frequently employed in France; it is adapted only for estimating the strength of spirits. The instrument is made like a common glass hydrometer, the scale of which is divided into 100 parts or degrees. The lowest division marked 0, at the bottom of the scale, denotes the specific gravity of pure water at a temperature of 15° Cent.; and the highest division, at the top of the scale, the specific gravity of absolute alcohol at the same temperature. The intermediate degrees indicate the number of volumes of absolute alcohol in 100 volumes of the spirit tried. The instrument is accom-




FIG. 3.



panied by a table for correcting the numbers marked on the scale, when it is used at any other temperature than that of 15° Cent.


Sikes's hydrometer is used exclusively in the collection of the spirit revenue. It consists of a spherical ball or float, and an upper and a lower stem made of brass;

(*a b c*, *fig. 4*) the upper stem (*b*) has ten principal divisions, numbered 1, 2, 3, &c., which are each subdivided into five parts; the lower stem (*c*) is made conical, and has a pear-shaped loaded bulb at its lower extremity. There are nine movable weights, (*b*, *fig. 5*) having the form of circular discs, and numbered 10, 20, 30, and so on to 90. Each of the circular weights is cut into its centre, so that it can be placed on the inferior conical stem, and slid down to the bulb; but, in consequence of the enlargement of the cone, they cannot slip off at the bottom, but must be drawn up to the thin part for this purpose. The instrument is adjusted to strong spirit, specific gravity .825, at 60° Fahr., this being reckoned as standard alcohol. In this spirit, the instrument floats at the first division, 0, or zero, without a weight. In weaker spirit, having a greater density, the hydrometer will not sink so low; and, if the density be much greater, it will be necessary to add one of the weights to cause the entire immersion of the bulb of the instrument. Each weight represents as many principal divisions of the stem as its number indicates: thus, the heaviest weight, marked 90, is equivalent to 90 divisions of the stem, and the instrument with this weight added floats at 0 in distilled water. As each principal division on the stem is divided into five, the instrument has a range of 500 degrees between *standard alcohol*, specific gravity .825, and water. In using this instrument, it is immersed in the spirit, and pressed down by the hand to 0, till the whole divided part of the stem be wet. The force of the hand required to sink it will be a guide in selecting the proper weight. Having taken one of the circular weights, which is necessary for this purpose, it is slipped on the conical stem. The instrument is again immersed, and pressed down as before to 0, and is then allowed to rise and settle at any point of the scale. The eye is then brought to the level of the surface of the spirit, and the part of the stem cut by the surface, as seen from below, is marked. The number thus indicated by the stem is added to the number of the weight employed, and with this sum at the side, and the temperature of the spirits at the top, the strength *per cent.* is found in a table which accompanies the hydrometer. The strength is expressed in numbers denoting the excess or deficiency *per cent.* of proof spirit in any sample; and the number itself, having its decimal point removed two places to the left, becomes a factor, whereby the gauged contents of a cask or vessel of such spirit being multiplied, and the product being added to the gauged contents if over proof, or deducted from it if under proof, the result will be the actual quantity of proof spirit contained in such cask or vessel.

Dicas's hydrometer is similar in construction to Sikes's; and it is used in a similar manner, with the same result, indicating the relation of the spirit tried to standard proof spirit.


It is the practice in commerce to designate the strength of spirit as so many

FIG. 4.



Sikes's Hydrometer.


FIG. 5.



Sikes's Hydrometer.

degrees above or below proof, the government having fixed upon what is called *proof spirit* as the standard, in comparison with which, the strength of all spirit shall be estimated. The term *proof* is said to have been derived from the ancient practice of trying the strength of spirit by pouring it over gunpowder in a cup, and then setting fire to the spirit; if, when the spirit had burned away, the gunpowder exploded, the spirit was said to be *over proof*; if, on the other hand, the gunpowder failed to ignite, in consequence of the water left from the spirit, it was said to be *under proof*. The weakest spirit capable of firing gunpowder in this way was called *proof spirit*: but it requires a spirit nearly of the strength of what is now called rectified spirit to stand this test. The *standard proof spirit* of the Excise is defined by law (56 Geo. III. cap. 140) to be "*that which, at a temperature of 51° by Fahrenheit's thermometer, weighs exactly twelve-thirteenth parts of an equal measure of distilled water.*" This will have a specific gravity of .923 at 51° Fahr., or about .920 at 60° Fahr. The *standard alcohol* of the Excise is spirit, the specific gravity of which is .825 at 60° Fahr. By "spirit 60 degrees over proof," is understood a spirit, 100 measures of which, added to 60 measures of water, will form *standard proof spirit*, sp. gr., 920. By "spirit 10 degrees under proof," is understood a spirit, 100 measures of which, mixed with 10 measures of standard alcohol, sp. gr. .825, will form *standard proof spirit*.

Saccharometers, (fig. 6) which are hydrometers intended for determining the density of syrups, are usually made and graduated in the same manner as Baumé's



acidometers, and differ only from these in being made smaller; but the scale is sometimes graduated to indicate the proportion of sugar in the solution.

The *urinometer* is a small hydrometer, originally suggested by Dr. Prout, for estimating the density of urine. The scale (*fig. 7*) is divided into 60 degrees, the zero being the point at which it floats in distilled water. The numbers on the scale, added to 1000, the assumed sp. gr. of water, give the specific gravities at the respective points; thus, supposing the number cut by the surface of the liquid to be 30, this indicates a sp. gr. 1030. The letters H. S., on the back of the scale, (*fig. 8*.) signify *healthy standard*, which ranges from 10° to 20° of the scale.

The *elaëometer* (*figs. 9 & 10*) is a very delicate glass hydrometer, intended for testing the purity of olive oil or oil of almonds, by determining their densities. The 0 or zero of the scale is the point at which the instrument floats in *oil of poppy seeds*. The point at which it floats in *pure olive oil* is made the 50th degree, and the space between these two points is divided into 50 equal parts, and numbered accordingly. It floats at 38 or 38½° in *pure oil of almonds*.

The following tables have been drawn up for the purpose of showing the relations between the indications afforded by some of the foregoing instruments.

Relation between Specific Gravities, and Degrees of Baumé's Hydrometer for Liquids heavier than Water.

Sp. gr. Baumé.	Sp. gr. Baumé.	Sp. gr. Baumé.	Sp. gr. Baumé.	Sp. gr. Baumé.
1.000 = 0	1.116 = 15	1.264 = 30	1.454 = 45	1.714 = 60
1.007 1	1.125 16	1.275 31	1.470 46	1.736 61
1.014 2	1.134 17	1.286 32	1.485 47	1.758 62
1.022 3	1.143 18	1.298 33	1.501 48	1.779 63
1.029 4	1.152 19	1.309 34	1.526 49	1.801 64
1.036 5	1.161 20	1.321 35	1.532 50	1.823 65
1.044 6	1.171 21	1.334 36	1.549 51	1.847 66
1.052 7	1.180 22	1.346 37	1.566 52	1.872 67
1.060 8	1.190 23	1.359 38	1.583 53	1.897 68
1.067 9	1.199 24	1.372 39	1.601 54	1.921 69
1.075 10	1.210 25	1.384 40	1.618 55	1.946 70
1.083 11	1.221 26	1.398 41	1.637 56	1.974 71
1.091 12	1.231 27	1.412 42	1.656 57	2.002 72
1.100 13	1.242 28	1.426 43	1.676 58	2.031 73
1.108 14	1.252 29	1.440 44	1.695 59	2.059 74
				2.087 75

Relation between Specific Gravities, and Degrees of Baumé's Hydrometer for Liquids lighter than Water.

Sp. gr. Baumé.	Sp. gr. Baumé.	Sp. gr. Baumé.	Sp. gr. Baumé.	Sp. gr. Baumé.
1000 = 10	0.934 = 20	0.875 = 30	0.823 = 40	0.777 = 50
0.993 11	0.927 21	0.869 31	0.819 41	0.773 51
0.986 12	0.921 22	0.864 32	0.814 42	0.769 52
0.979 13	0.915 23	0.859 33	0.809 43	0.765 53
0.972 14	0.909 24	0.853 34	0.805 44	0.760 54
0.966 15	0.903 25	0.848 35	0.800 45	0.756 55
0.959 16	0.897 26	0.843 36	0.796 46	0.752 56
0.953 17	0.892 27	0.838 37	0.791 47	0.748 57
0.946 18	0.886 28	0.833 38	0.787 48	0.744 58
0.940 19	0.880 29	0.828 39	0.782 49	0.739 59
				0.735 60

TEMPERATURES TO BE OBSERVED IN CERTAIN PHARMACEUTICAL OPERATIONS.

In the fermentation of saccharine solutions, the highest temperature should not exceed 86° (Thomson).

The lowest temperature at which they will ferment is 38° (Thomson).

The process of acetous fermentation is best conducted at a temperature of about 86°.

The temperature requisite to coagulate albumen varies with the state of dilution. If the quantity of albumen be so great that the liquid has a slimy aspect, a heat of 145° or 150° suffices, but in a very dilute condition boiling is required (Fownes).

In the London and U. S. Pharmacopœias :

When a boiling heat is directed, a temperature is meant of 212° Fahr.

When a gentle heat is directed, a temperature is meant of from 90° to 100°.

The specific gravities of substances ordered in the London Pharmacopœia are to be taken at a temperature of 62°.

A water bath is that by which any substance contained in a proper vessel is exposed either to hot water, or the vapor of boiling water. A sand bath is made of sand, to be gradually heated, in which anything is placed contained in a proper vessel.

Syrups are to be kept in a place where the temperature never exceeds 55°.

Vegetables, shortly after they have been gathered, those excepted which ought to be fresh, are to be lightly strewed, and dried as quickly as possible with a gentle heat (90° to 100°). They are to be kept afterwards in proper vessels, excluded from the access of light and moisture.

In the Dublin Pharmacopœia :

By the term superior heat is meant some degree between 200° and 212°.

When a medium heat is directed, a temperature is meant between 100° and 200°.

When an inferior heat is directed, a temperature is meant between 90° and 100°.

In the process of digestion, an inferior heat is to be applied, unless it should be otherwise directed. In the process of maceration, a heat should be applied between 60° and 90°.

In the Dublin, Edinburgh, and U. S. Pharmacopœias :

Whenever mention occurs of the specific gravity of any body, its temperature is supposed to be at 60°.

HYDROMETRICAL EQUIVALENTS.

Sp. gr. at 60° Fahr.	100 parts contain of Alcohol sp. gr. 796. By weight.		1000 parts contain of standard Alc. sp. gr. 825	Sikes.	Baumé.	Cartier.	Per cent. of Alcohol, sp. gr. 796 by volume. Gay Lussac.
796	100	0			46.5	43.48	100
797	99.5	.5					99.75
798	99	1			46	43.06	99.50
799	98.67	1.33					99.25
800	98.33	1.67					99
801	98	2					98.75
802	97.67	2.33			45	42.14	98.50
803	97.33	2.67				42	98.28
804	97	3					98.15
805	96.67	3.33					98

Sp. gr. at 60° Fah.	100 parts contain of		1000 parts contain of standard Alc. sp. gr. 825	Sikes.	Baumé.	Cartier.	Per cent. of Alcohol, sp. gr. 796 by volume. Gay Lussac.
	Alcohol sp. gr. 796.	Water. By weight.					
806	96.33	3.67					97.80
806.5	96.17	3.83			44	41.22	97.70
807	96	4					97.60
808	95.5	4.5				41	97.40
809	95	5					97.29
809.5	94.89	5.10					97.10
810	94.67	5.33					97
811	94.33	5.67			43	40.34	96.75
812	94	6					96.50
813	93.67	6.33				40	96.25
814	93.33	6.67					96
815	93	7					95.75
816	92.5	7.5			42	39.40	95.50
817	92	8					95.25
818	91.67	8.33					95
818.6	91.5	8.5				39	94.90
819	91.33	8.67					94.75
820	91	9					94.50
821	91.5	9.5			41	38.46	94.25
822	90	10					94
823	89.67	10.33				38	93.75
824	89.33	10.67					93.50
825	89	11	1000	63 O. P.	40	37.55	93.25
826	88.5	11.5	993	62			93
827	88	12	988.5	61.5		37	92.6
828	87.67	12.33	984	61	39.5		92.3
829	87.33	12.67	979.5	60.5	39	36.63	91
830	87	13	975	60			91.7
831	86.5	13.5	970.5	59.5	38.5	36.17	91.35
832	86	14	966	59		36	91
833	85.67	14.33	961.5	58.3			90.65
834	85.33	14.67	957	58	38	35.72	90.3
835	85	15	953	57.5			90
836	84.67	15.33	949	57	37.5	35.26	89.7
837	84.33	15.67	944.5	56.5			89.35
837.6	84.25	15.75	942.5	56.3		35	89.20
838	84	16	940	56			89
839	83.5	16.5	936	55.5	37	34.80	88.75
840	83	17	932	55			88.5
841	82.67	17.33	928	54.5	36.5		88.25
842	82.33	17.67	924	54		34.94	88
843	82	18	920	53.5		34	87.65
844	81.67	18.33	916	53	36	33.88	87.3
845	81.33	18.67	912	52.5			87
846	81	19	908	52			86.7
847	80.5	19.5	903	51	36.5	33.42	86.35
848	80	20	898	50			86
849	79.67	20.33	893	49.5			85.65
850	79.33	20.67	888	49	35	33	85.3
851	79	21	883	48.5			85
852	78.5	21.5	878	48	34.5	32.43	84.7
853	78	22	873	47.5			84.35

Sp. gr. at 60° Fah.	100 parts contain of Alcohol sp. gr. 796. By weight.		1000 parts contain of standard Alc. sp. gr. 825.	Sikes.	Baumé.	Cartier.	Per cent. of Alcohol, sp. gr. 796 by volume. Gay Lussac.
		Water.					
854	77.5	22.5	868	47 O. P.			84
855	77	23	862.5	46.5	34	32.04	83.65
856	76.5	23.5	857	46			83.3
857	76	24	853	45.5	33.5	31.58	83
858	75.67	24.33	849	45			82.7
859	75.33	24.67	844.5	45			82.35
860	75	25	840	45	33	31.13	82
861	74.67	25.33	836.5	44.5		31	81.7
862	74.33	25.67	833	44			81.3
862.5	74.16	25.84	830.5	43.75	32.5	30.76	80
863	74	26	828	43.5			80.8
864	73.5	26.5	823	43			80.3
865	73	27	818	42.5	32	30.21	79.95
866	72.5	27.5	813	42			79.6
867	72	28	810	41			79.3
867.5	71.83	28.17	808.5	40.5	31.5	29.78	79.15
868	71.67	28.33	807	40			79
869	71.33	28.67	802.5	39.5			78.65
870	71	29	798	39	31	29.29	78.3
871	70.5	29.5	792.5	38.5			78
872	70	30	787	38		29	77.7
873	69.5	30.5	781.5	37	30.5	28.83	77.35
874	69	31	776	36			77
875	68.67	31.33	772	35			76.5
876	68.33	31.67	768	34	30	28.38	76
877	68	32	762.5	33			75.65
877.5	67.75	32.25	759.25	32.5		28	75.5
878	67.5	32.5	757	32			75.3
878.5	67.25	32.75	753.75	31.5	29.5	27.91	75
879	67	33	751.5	31			74.8
880	66.5	33.5	746	30			74.3
881	66	34	742	29.5	29	27.44	74
882	65.5	34.5	738	29			73.7
883	65	35	733.5	28.5		27	73.35
883.5	64.83	35.17	731.25	28.25	28.5	26.99	73.17
884	64.67	35.33	729	28			73
885	64.33	35.67	724	27.5			72.5
886	64	36	719	27	28	26.53	72
887	62.67	36.33	714	26			71.5
888	63.33	36.67	709	25			71
889	63	37	704	24.5	27.5	26.07	70.65
890	62.5	37.5	699	24			70.3
891	62	38	694	23			69.8
892	61.5	38.5	689	22	27	25.61	69.3
893	61	39	644.5	21			69
894	60.67	39.33	680	20			68.7
895	60.33	39.67	675.5	19.5			68.35
895.5	60.16	39.84	673.25	19.25	26.5	25.15	68.17
896	60	40	671	19		25	68
897	59.5	40.5	666.5	18			67.65
898	59	41	662	17	26	24.69	67.3
899	58.5	41.5	655.5	16			67
900	58	42	649	15			66.5

Sp. gr. at 60° Fah.	100 parts contain of		1000 parts contain of standard Alc. sp. gr. 825	Sikes.	Baumé.	Cartier.	Per cent. of Alcohol, sp. gr. 796 by volume. Gay Lussac.
	Alcohol sp. gr. 796.	Water. By weight.					
900.5	57.75	42.25	647	14.75 O.P	25.5	24.23	66.52
901	57.5	42.5	645	14.5			66.35
901.5	57.25	42.75	643	14.25		24	66.17
902	57	43	641	14			66
903	56.5	43.5	636	13	25	23.77	65.5
904	56	44	631	12			65
905	55.5	44.5	626	11.5			64.5
906	55	45	621	11	24.5	23.31	64
907	54.5	45.5	616.5	10.5			63.65
908	54	46	612	10		23	63.3
909	53.5	46.5	607	9	24	22.85	62.65
910	53	47	602	8			62.3
911	52.5	47.5	595.5	7.5			61.9
912	52	48	591	7	23.5	22.39	61.5
913	51.67	48.33	586	6			61
914	51.33	48.67	581	5			60.5
915	51	49	576	4	23	21.94	60
916	50.5	49.5	571	3			59.6
917	50	50	560.5	2			59.3
918	49.67	50.33	562	1	22.5	21.48	59
919	49.33	50.67	554	.5			58.5
920	49	51	550	Proof			58
921	48.5	51.5	545	1 U. P.	22	21.02	57.5
922	48	52	540	2			57
923	47.5	52.5	535.5	3	21.5	20.56	56.5
924	47	53	531	4			56
925	46.5	53.5	526	5			55.5
926	46	54	521	6	21	20.10	55
927	45.5	54.5	515.5	6.5			54.5
928	45	55	510	7			54
929	44.5	55.5	505	8			53.5
929.5	44.25	55.75	502.5	8.5	20.5	19.64	53.25
930	44	56	500	9			53
931	43.67	56.33	495.5	10			52.5
932	43.33	56.67	489	11			52
933	43	57	484	12	20	19.18	51.5
934	42.5	57.5	479	13		19	51
935	42	58	472.5	14			50.5
936	41.5	58.5	468	15	19.5	18.72	50
937	41	59	462	16			49.5
938	40.5	59.5	456	17			49
939	40	60	450	18	19	18.26	48.5
940	39.5	60.5	444	19			48
940.5	39.25	60.75	441	19.5		18	47.63
941	39	61	438	20			47.25
942	38.5	61.5	432	21	18.5	17.80	46.5
943	38	62	426.5	22			45
944	37.5	62.5	421	23			45.5
945	37	63	416	23.5	18	17.35	44.75
946	36.5	63.5	411	24			44
947	36	64	399	25		17	43.5
948	35.5	64.5	397	26	17.5	16.89	43
949	35	65	389.5	27			42.25

Sp. gr. at 60° Fah.	100 parts contain of		1000 parts contain of standard Ale. sp. gr. 825	Sikes.	Baumé.	Cartier.	Per cent. of Alcohol, sp. gr. 796 by volume. Gay Lussac.
	Alcohol sp. gr. 796.	Water. By weight.					
950	34.5	65.5	382	28 U. P.			41.5
951	34	66	376	29.5	17	16.43	40
952	33.5	66.5	370	31			40.5
953	33	67	364	32.5			39.75
954	32.5	67.5	358	34			39
955	32	68	352	35	16.5	16.3	38.5
956	31.5	68.5	346	36		16	38
957	31	69	339.5	37.5			37.25
958	30	70	333	39			36.5
959	29.5	70.5	324	40.5	16	15.51	35.75
960	29	71	315	42			35
961	28.5	71.5	307.5	43.5			34.5
962	28	72	300	45	15.5	15	34
963	27	73	292.5	46.5			33
964	26	73.5	285	48			32
965	26	74	277.5	49.5	15	14.59	31
966	25.5	74.5	270	51			30
967	25	75	261.5	52.5			29
968	24	76	253	54			28
968.5	23.75	76.25			14.5	14.13	27.5
969	23.5	76.5	244.5	55.5			27
970	23	77	236	57			26
971	22.5	77.5	227	58.5			25
972	22	78	218	60	14	13.67	24
973	21	79	209	62			23
974	20	80	200	64			22
975	19	81	195	66	13.5	13.21	21
976	18.5	81.5	190.5	68			20
977	18	82	183.5	70			19
978	17	83	175	72	13	12.76	18
979	16	84	163	73.5			17
980	15.5	84.5	150	75			16
981	15	85	143	76			15
982	14	86	135	77	12.5	12.30	14
983	13.5	86.5	128	78.5			13
984	13	87	120	80			12
985	12.5	87.5	112	81			11.25
986	12	88	105	82	12	11.84	10.5
987	11	89	98	83.5			9.75
988	10	90	90	85			9
989	9	91	82	87	11.5	11.38	8
990	8	92	75	89			7
991	7	93	67.5	90.5			6.5
992	6	94	60	92			6
993	5.5	94.5	52.5	93.5	11	10.92	5
994	5	95	45	95			4
995	4	96	37.5	95.5			3.5
996	3.5	96.5	30	96	10.5	10.46	3
997	3	97	22.5	97			2
998	2	98	15	98			1
999	1	99	7.5	99			.5
1000	0	100	0	100	10	10	0

SPECIFIC GRAVITIES OF SOME OF THE PREPARATIONS ORDERED IN THE PHARMACOPŒIAS.

The London Pharmacopœia directs the Specific Gravity to be taken at a temperature of 62° Fahr. The Edinburgh, Dublin, and U. S. Pharmacopœias at 60° Fahr.

		Sp. Gr.
Acetum Destillatum	Edin.	1.005
.....	London	1.0065
Acidum Aceticum	London	1.048
.....	Edin.	1.063
.....	to
.....	1.065
.....	U. S.	1.041
..... Glaciale	Dublin	1.065
..... Forte	Dublin	1.066
..... Dilutum	Dublin	1.006
..... Hydrochloricum	London, U. S.	1.160
.....	Edin.	1.170
.....	Dublin	1.176
..... Dilutum	Edin.	1.050
.....	London	1.043
.....	U. S.	1.046
.....	Dublin	1.045
..... Hydrocyanicum	Dublin	0.997
..... Nitricum	London, U. S.	1.420
..... Purum	Edin., Dublin,	1.500
..... Dilutum	London	1.082
.....	U. S.	1.070
.....	Edin.	1.077
.....	Dublin	1.092
..... Phosphoricum Dilutum	London	1.064
..... Sulphuricum	London, U. S.	1.845
.....	Edin.	1.840
.....	Dublin	1.846
..... Sulphuricum Dilutum	London	1.103
.....	Edin., U. S.	1.090
.....	Dublin	1.084
Æther	London, U. S.	0.750
Æther Sulphuricus	Edin.	0.735
.....	Dublin	0.750
Alcohol	London	0.838
.....	U. S.	0.835
.....	Edin.	0.796
.....	Dublin	0.795
..... Dilutum	U. S.	0.935
Aqua Destillata	U. S. L. E. D.	1.000
..... Ammoniae	Edin.	0.960
.....	Dublin	0.950
..... Acetatis	Edin.	1.011
..... Calcis Muriatis (Calcii Chloridi)	Dublin	1.225
Aqua Potassæ	Edin.	1.072
.....	Dublin	1.068
..... Carbonatis	Dublin	1.301
..... Sodæ Carbonatis	Dublin	1.026

Liquor Ammoniae	London, U. S.	Sp. Gr. 0.960
Fortior	London, U. S.	0.882
Barii Chloridi	Dublin	1.088
Plumbi Diacetatis	London	1.260
.....	U. S.	1.267
Potassae	London	1.063
.....	U. S.	1.056
Carbonatis	London	1.473
Oleum Æthereum	London	1.050
.....	U. S.	1.096
Spiritus Ætheris Nitrici	London, U. S.	0.834
.....	Edin.	0.847
Sulphurici	Edin.	0.809
Ammoniae	Edin.	0.845
.....	U. S.	0.831
Aromaticus	U. S. London	0.918
Fœtidus	London	0.861
Rectificatus	London	0.838
.....	Edin.	0.838
.....	Dublin	0.840
Tenuior	London	0.920
.....	Edin.	0.912
.....	Dublin	0.920
Tinctura Ferri Sesquichloridi	London	0.992
.....	Dublin	1.237

RELATION BETWEEN DIFFERENT THERMOMETRICAL SCALES.

The thermometer always used in this country and England is that of Fahrenheit; it is also used in parts of Germany.

In this instrument, the range between the freezing and boiling points of water is divided into 180°, and as the greatest possible degree of cold was supposed to be that produced by mixing snow and salt together, it was made the zero. Hence, the freezing point became 32°, and the boiling point 212°.

The Centigrade thermometer places the zero at the freezing point of water, and divides the range between the freezing and boiling points into 100°. This scale has long been used in Sweden, under the title of Celsius's thermometer, and is generally adopted on the Continent.

Reaumur's thermometer, which was formerly used in France, divides the space between the freezing and boiling points of water into 80°, and places the zero at the freezing point. It is now little employed, except in the north of Germany.

De Lisle's thermometer is used in Russia. The graduation begins at the boiling point, and increases towards the freezing point. The boiling point is marked 0°, and the freezing point 150°.

To reduce Centigrade degrees to those of Fahrenheit.

RULE.—Multiply by 9, divide the product by 5, and add 32.

$$\text{Thus, } \overset{\text{Cent.}}{40} \times 9 \div 5 + 32 = \overset{\text{Fahr.}}{104}.$$

To reduce Fahrenheit's degrees to those of Centigrade.

RULE.—Subtract 32, multiply by 5, and divide by 9.

$$\text{Thus, } \overset{\text{Fahr.}}{104} - 32 \times 5 \div 9 = \overset{\text{Cent.}}{40}.$$

To reduce Reaumur's degrees to those of Fahrenheit.

RULE.—Multiply by 9, divide by 4, and add 32.

$$\text{Thus, } \overset{\text{Reaum.}}{32} \times 9 \div 4 + 32 = \overset{\text{Fahr.}}{104}.$$

To reduce Fahrenheit's degrees to those of Reaumur.

RULE.—Subtract 32, multiply by 4, and divide by 9.

$$\text{Thus, } \overset{\text{Fahr.}}{104} - 32 \times 4 \div 9 = \overset{\text{Reaum.}}{32}.$$

To reduce Reaumur's degrees to those of Centigrade.

RULE.—Multiply by 5, and divide by 4.

$$\text{Thus, } \overset{\text{Reaum.}}{32} \times 5 \div 4 = \overset{\text{Cent.}}{40}.$$

To reduce Centigrade degrees to those of Reaumur.

RULE.—Multiply by 4, and divide by 5.

$$\text{Thus, } \overset{\text{Cent.}}{40} \times 4 \div 5 = \overset{\text{Reaum.}}{32}.$$

EXPLANATION OF THE PRINCIPAL ABBREVIATIONS OCCURRING IN PHARMACEUTICAL FORMULÆ.

R. Recipe—Take.

F. S. A. Fiat secundem artem—Let it be made or prepared according to the rules of the art.

M. Misce—Mix.

M. S. D. Misce, signa, da—Mix the medicine, and deliver it afterwards, with the requisite instruction, to the patient (or nurse) in writing.

M. F. P. Misce fiat pulvis—Mix to form a powder.

M. F. Mist. Misce fiat mistura—Mix to form a liquid mixture.

M. F. Pil. Misce fiant pilulæ—Mix to form pills.

Div. Divide—Divide.

Sol. Solve—Dissolve.

Fasc. j. Fasciculus*—An armful.

Man. j. Manipulus—A handful, a gripe.

Pugil. j. Pugillus or Pugillum—A pinch.

Cyat. j. Cyathus—A glassful.

Cochl. j. Cochlear or cochleare—A spoonful.

Gutt. Gutta—A drop.

No. 1, 2, 3, &c. The number of pieces or parts.

Ana, or āā. Of each.

P. Ae. Partes æquales—Equal parts.

Q. S. Quantum sufficit—As much as will suffice.

Q. L. Quantum libet—As much as you like.

Q. V. Quantum volueris—As much as you like.

℔. Libra—A pound.

℥. Uncia—An ounce.

ʒ. Drachma—A drachm or dram.

ʒ. Scrupulus—A scruple.

* The terms *fasciculus*, *manipulus*, and *pugillus* are understood differently by authors, as seen in the following table.

	Linnaeus.	Gelger.	Paris.
Pugillus.....	ʒj.....	ʒss ad ʒj.....	ʒj ad ʒij.
Manipulus.....	ʒiv.....	ʒiv.....	ʒj ad ʒijss
Fasciculus.....	ʒvj.....	ʒj.....	

Gr. Granum—A grain
 Pil. Pilula—A pill.
 Pot. Potio—A potion.
 Pulv. Pulvis—A powder.
 Tinct. Tinctura—A tincture.
 Ext. Extractum—An extract.
 Chart. Chartula—A small paper.
 Collyr. Collyrium—An eye water.
 Collutor. Collutorium—A mouth-wash.
 Cong. Congius—A gallon.
 O. Octarius—A pint.
 f℥. Fluiduncia—A fluidounce.
 fʒ. Fluidrachma—A fluidrachm.
 ℥. Minimum—A minim.
 Decoct. Decoction—A decoction.
 Garg. Gargarisma—A gargle.
 Haust. Haustus—A draught.
 Infus. Infusum—An infusion.
 Mass. Massa—A mass.
 Mist. Mistura—A mixture.
 Ss. Semis—A half.

The use of the Latin language for prescriptions and for the accompanying directions is universal in Europe, except in France, where, although the prescription itself may be in Latin, the directions are usually given in French; this is also the case in the United States, where few, if any, physicians give the directions for the use and mode of administration of a formula in any other language than English. As, however, it often becomes of importance to refer to the prescriptions of foreign practitioners, the following table, from Mowbray's *Conspectus*, is introduced:—

A VOCABULARY

Of the Words most frequently occurring in the Prescriptions of Physicians.

A.

A, āā, or ana, of each ingredient; more promptly expressed by the genitive case plural (singulorum) of each; for instance:

R.—Aquæ cinnamomi, tinct. rhei, āā fʒij, means

Take of cinnamon-water, and tincture of rhubarb, of each two fluidrachms.

Abdomen, the belly. Gen.—inis, of the belly. Dat.—ini, to the belly.

Absente febre, while the fever is off.

Accuratè (adverb), accurately. Accuratè pensi, weighed with the utmost exactness.

Accuratè misceantur, mix very completely.

Aciditas, sharpness. Ad gratam aciditatem, make it just acid enough to be palatable, and not to sour.

Acmè, the height of a fever or any other disease.

Ad, to. Ad duas vices, at twice taking. Ad libitum, freely, as you like.

Adde, add. Addantur, let there be added. Addendus, to be added. Addendo, adding.

Admoveatur, let there be applied—antur, plural, when more than one is to be applied.

Adstante febre, while the fever is on.

Æger, a sick person, a patient. Ægra, a female patient.

Aggrediente febre, when the fever is coming on.

Aggressus, an attack. Aggressus febris, the attack of a fever.

Agitato vasè, shaking the phial.

Albus, white.

Aliquot, some. Aliquoties, sometimes.

Alter, altera, alterum, the other.

Alternus, alternate. Alternis horis, every second hour. Alternis diebus, every alternate day.

Aluta, leather. Extende super alutam mollem, spread on soft leather.

Alvus, the belly, the bowels. Adstrictâ alvo, when costive.

Amplus, large. Cochleare amplum, a tablespoonful.

Antè, before. Ut antè, as before.

Antemeridianus, in the forenoon; any time before twelve o'clock at noon.

Apparatus, any sort of preparation, instruments, or, in short, everything that is requisite to be had in readiness for performing any sort of operation. Also, the more delicate term for a bladder and pipe for clysters.

Aperiens, opening, gentle purging.

Applicetur, let there be applied.

Aqua, water. Aquæ, of water. Aqua bulliens, boiling water. Aqua fervens, hot water.

Armatus, armed. Fistula armata, an apparatus for clysters; a pipe and bladder.

Auris, the ear. Auri, to the ear. Aures, ears. Auribus, to the ears.

Aut, or.

B.

Benè, well.

Bibat, let him drink.

Biduum, two days. Omni biduò vel triduò, every two or three days.

Bihorium, the space of two hours. Omni bihorio, every two hours.

Bis, twice; bis, terve, twice or thrice.

Bullio, to boil; bulliens, boiling. Aquæ bullientis, of boiling water.

Bulliant, let them boil.

C.

Calefactus, made warm.

Capiat, let (the patient) take. Capt. cochl. iij. magn., three tablespoonfuls to be taken.

Catharticus, purging.

Cautè, cautiously.

Cerevisia, beer. Cerevisia Londinensis, porter. Cerevisiæ Fermentum, yeast.

Charta, paper. Charta cærulea, blue paper. Chartula, a little piece of paper. Cola trans chartam, filter through paper.

Cibus, food.

Circa, about. Circiter, about.

Citò, soon, quickly. Citissimè (the superlative degree), as quick as possible.

Clausus, clausa, clausum, covered. Vase clauso, in a covered vessel.

Cochleare, a spoonful. Cochlearia, spoonfuls. Coch. magnum, amplum, a table-spoonful. Co. infantulorum, coch. modicum, child's spoonful, a deserts-
spoonful. Co. parvum, a teaspoonful.

Coctio, a boiling. Sub finem coctionis, towards the end of the boiling, when almost boiled.

Cæruleus, blue. Charta cærul., blue paper. Unguentum cæruleum, mercurial ointment.

Cola (imperat. of Colo), strain, to filter.

Colatus, strained, filtered. Colato liquori, to the strained liquor.

Colatura, a straining. Colaturæ, to the strained liquor.

Colatus, let be strained. Colentur, let them be strained.

Collum, a neck.

Comp., Compositus, compound.

Continuo, to continue. Continuantur remedia, go on with the same medicines as last prescribed.

Contundo, to bruise. Contusus, bruised.

Contusio, a bruise, a contusion.

Coquo, to boil. Coque, boil. Coque parùm, boil a little while.

Coquantur, let them be boiled.

Cor, the heart. Scrobiculus cordis, the pit of the stomach.
 Coxa, coxendix, the hip.
 Cras, to-morrow; c. m., cras mane, early to-morrow morning; c. m. s., cras mane sumendus, to be taken early to-morrow morning.
 Crastinus (adj.), of to-morrow. In usum crastinum, for to-morrow's use.
 Cribrum, a sieve. Trans cribrum, through a sieve.
 Cujus, of which. Cujus capiat, or sumat, of which (the patient is to) take.
 Cum, with.
 Cyathus, a cup. Cyathus vinarius, a wineglass. Cyath. theat, a teacupful.

D.

Da, give. Detur, dentur (plural), let be given.
 De, of. De quo, or quâ, of which. De die, in a day.
 Deauratus, gilt. Deaurentur pilulæ, let the pills be gilt.
 Debilitas, weakness.
 Decem, ten. Decimus, the tenth.
 Debitus, due. Ad debitam spissitudinem, to a proper degree of thickness, as to consistence.
 Decubitus, lying down. Horâ decubitûs, at bedtime.
 Deglutio, to swallow. Deglutiatur, is swallowed—etur, let be swallowed.
 Dein, deinde, then; afterwards.
 Dejectio, a depositing, or putting down; also a going to stool; as, post duas dejectiones alvi, after two motions.
 Dejitio, to deposit. Donec alvus bis dejiciat, until the patient shall have had two stools.
 Dexter, the right. Manus dextra, the right hand. Auri dextro, to the right ear.
 Dictus, spoken of, said.
 Dies, a day. Die, in a day; as, bis die, twice a day. Diebus, in days; as, tertiis diebus, every third day. Alternis diebus, every alternate day.
 Dilutum, diluted.
 Dimidium, the half. Dimidius, a, um (adjective), half.
 D. P., Directione propriâ, with its proper direction.
 Diu, a long time. Tere diu, rub for a long time.
 Diuturnus, long continued. Diuturnâ trituratione, by long-continued rubbing, or grinding in the mortar. Diuturna coctione, by long boiling.
 Dolor, pain. Dolores, pains. Doloris, of the pain.
 Dolens, pained. Parti dolenti, to the pained part.
 Donec, until. Dum, whilst.
 Durante dolore, while the pain continues.

E.

Eadem, eandem, the same. Eodem, in the same.
 Effunde, pour out.
 Effervescentia, the effervescence.
 Effervescet, effervesces.
 Ejusdem, of the same, the genitive case of idem.
 Enema, a clyster. Enemata, clysters.
 Erit, shall be.
 Evanescio, evanui, to disappear. Evanuerit, shall have disappeared.
 Exhibe, exhibeatur, give. Exhibendus, to be given.
 Extendo, to extend, also, to spread. Extende super alutam, spread it on leather.
 Extensus, a, um, spread.

F.

Fac, make. Fac in pilulas xij, make into twelve pills.
 Farina, flour. Farina seminis lini, linseed meal.
 Febris, a fever. Febre durante, while the fever is on.
 Femur, a thigh.

Femoribus, to the thighs. Femoribus internis, to the inner sides of the thighs.
 Fervens, boiling. Fervidus, fervida, hot.
 Ferventis, of boiling. R aq. ferventis, take of boiling water.
 Finis, the conclusion. Sub finem coctionis, when almost boiled enough.
 Fiat, make (the singular number). Fiant, make (plural).
 Fontana, fountain. Aqua fontana, spring water.
 Formula, prescription.
 Fetus, a fomentation.
 Fuerit, shall have been; as, donec alvus soluta fuerit, until a motion is (or shall have been) procured.

G.

Gargarisma, a gargle.
 Gelatina, jelly. Gelatina ribesiorum, currant jelly. Gelatina quovis, any sort of jelly.
 Globulus, a little ball. Globuli Gascoigni, Gascoign's ball. Donec globuli evanuerint, until the globules (of quicksilver) totally disappear (so that they cannot be seen even with a microscope.)
 Gradatim, by slow degrees.
 Gratus grata, gratum, agreeable, pleasant. Ad gratam aciditatem, so as to make it pleasantly acid without being too sour. In quovis grato vehiculo, in any agreeable vehicle.
 Gutta, a drop. Guttæ, drops. Guttas, drops.
 Guttatim, drop by drop.

H.

Hac, this. Hac nocte, this night. Hanc, this. Sumat hanc, let him take this.
 Hactenus, hitherto, heretofore, up to the present day.
 Harum, of these. Harum pilularum sumat tres, of these pills let him or her take three.
 Haustus, a draught.
 Hebdomada, a week.
 Heri, yesterday. Ut heri, as yesterday.
 Hesternus, of yesterday. Hesternæ nocte, last night.
 Hirudo, a leech. Hirudines, leeches.
 His, in these, to these. His adde, add to these.
 Hora, an hour. Horæ, of an hour. Horæ (plural) hours.
 H.S. (horâ somni), at the hour of rest.
 H.S.S. (horâ somni sumendus), to be taken at bedtime. Hora decubitus, at the hour of going to rest, bedtime.
 Horâ vespertinâ, in the evening.
 Horæ unius spatio, in the space of one hour.
 Horæ $\frac{1}{4}$, horæ quadrante, quarter of an hour.
 Horis intermediis, at intermediate hours, when two medicines are to be taken. Horis intermediis means that one is to be given exactly at midtime from the other: suppose a draught is ordered (to be taken every six hours), and a powder horis intermediis, that is every six hours intermediately, then a draught will be taken at six o'clock and at twelve, and a powder at three and at nine.
 Hujusmodi, of this sort, like these.
 H. p. n., Haustus purgans noster, a formula of purging draught made according to a practitioner's own private Pharmacopœia, and is prepared so as to keep a long time without spoiling, and thus avoid the trouble of preparing it every time a draught is wanted. Mitt. H. p. n. \mathfrak{z} ij ad ij Vices c. m. s. Mitte Haustus purgantis nostri uncias duas, ad duas Vices cras mane sumendus; send two ounces of our purging draught, to be taken to-morrow morning, at twice, that is, half at first, and the remaining half in an hour if the first do not operate.

I.

Idoneus, proper, appropriate.
 Idem, eadem, the same. Ejusdem, of the same, the genitive case of idem.
 Imponatur (sing.), let there be put on—nantur (plural).
 Imprimis, first.
 In, in. In die, in a day.
 Indiès (an adverb), every day, daily.
 Indicaverit, shows, indicates.
 Infusio, infusum, an infusion.
 Infunde, infundatur, infuse.
 Inter, between.
 Injectio, injection.
 Inquietudo, restlessness. Urgente inquietudine, if restless.
 Injiciatur, throw in, throw up. Injiciatur enema, let a clyster be administered.
 Injiciendus, injiciendum, to be administered.
 Instar (an adverb), as big as. Sumat molem instar mucis moschatae, the bigness of a nutmeg to be taken.
 Intermedius, intermediate.
 Internus, the inner side.

J.

Jam, already. Jampridem, jamdudum, some time ago, heretofore.
 Jugulum, the throat.
 Jusculum, broth. Jusculum ovillum, mutton broth. Jus bovinum, beef tea.
 Juxta, near to.

L.

Lac, milk. Lactis, of milk. Lacte, in milk.
 Lævigatus, levigated.
 Lana, flannel. Lana nova, new flannel.
 Languor, faintness, lowness. In languoribus, in the fainting fits, when low and faint.
 Latus, the side. Lateris, of the side. Lateri, to the side.
 Latus dolens, the pained side. Lateri dolenti, to the pained side.
 Latus, lata, latum (adjective), broad.
 Lectus, a bed. In lecto, in bed.
 Liber, a book.
 Libra, a pound. Libris, libras, pounds.
 Linteum, lint; also linen.
 Liquesco, to liquefy, to melt. Donec liquescat, till it melts.
 Liquidus, liquid. Sedes liquidæ, loose stools. In quovis liquido, in any liquid.
 Londinensis, of London. Pharmacopœia Londinensis, the London Pharmacopœia.
 Lumborum, of the loins.
 Libet (a verb impersonal), it pleases. Ad libitum, just as you please.

M.

Magnus, magna, magnum, large. Magnum cochleare, a tablespoonful.
 Major, greater, larger. Cochlearia duo majori, two tablespoonfuls.
 Maximus, the greatest. Maximâ curâ, with the greatest care.
 Maximè, chiefly.
 Malleolus, the ankle. Malleolus internus, the inner ankle.
 Manè, in the morning. Manè primo; valde mane, very early in the morning.
 Manus, a hand. Manu calefacta, with a warm hand.
 Massa, a mass. Massa pilularis, a mass fit for forming into pills.
 Matutinus, in the morning or forenoon.
 Mediocris, middle-sized. Pilulæ mediocres, middle-sized pills. Cochleare mediocre, a dessertspoonful, a pap-spoonful, or a child's spoonful. Mediocris also means indifferent, as to quality.
 Medius, middle. Media nocte, in the middle of the night.

Melior, melius, better.
 Minatur, minaretur, threatens. Minante, threatening.
 Minimus, very small. Cochleare minimum, a teaspoonful.
 Minutum,* a minute.
 Misce, mix. Bene misceatur, let it be well mixed.
 Mite, send.
 Mittatur (singular), let it be sent. Mittantur (plural), let them be sent.
 Mittatur sanguis, take blood away; *i. e.*, bleed the patient.
 Mistura, mixtura, a mixture.
 Mitigatio, mitigation, alleviation. Donec dolor is mitigatio sit, until the pain is easier.
 Mitigatus, a, um, mitigated, lessened.
 Modicus, middle-sized.
 Modus, a manner. Modo præscripto, in the manner directed.
 Moles, a mass, a lump, a piece. Sumat molem instar nucis moschatæ, let him (or her) take the bigness or size of a nutmeg.
 Mollis, molle, soft.
 Molestus, troublesome. Molestor, to trouble, to be troublesome. Molestante dolore, when the pain is troublesome. Molestante tussi, when the cough is troublesome.
 Mora, delay. Sine morâ, without delay.
 Mos, moris, manner. More solito, in the usual manner; also, in the same manner as I am in the habit of prescribing it to other people.
 Mortarium, a mortar. Mortario aheneo, in the brass mortar. Mortario marmoreo, in the marble mortar. Mort. vitreo, in the glass mortar

N.

Narthecium, a gallipot.
 Nates, the buttocks.
 Ne, lest, also, do not; as, ne tradas sine nummo, do not deliver the medicine without the money.
 Necnon, also.
 Niger, nigra, nigrum, black.
 Ni, nisi, unless.
 Nihil, nothing.
 Nisus, an endeavor, an attempt, a straining, a motion, a straining to vomit, or go to stool.
 Nimis, nimium, too much.
 Nodulus, a little knot. Nodulo ligati, tied up in a piece of clean rag.
 Nomen, nominis, a name. Signetur nomine proprio, write its common name upon the label.
 Novem, nine.
 Nonus, the ninth.
 Novus, nova, novum, new.
 Novissimè, very lately, the last of all.
 Nocte, at night. Noctes, nights. Nocte maneque, night and morning. Alternis noctibus, every second night.
 Nuper, lately. Nuperrime, very lately.
 Nucha, the nape of the neck. Nuchæ, to the nape.
 N.M., Nux moschata, a nutmeg. Sumat magnitudinem nucis moschatæ, take the bigness of a nutmeg.

* The word minutum, for a minute, is very barbarous Latin; we believe there is no such word; but the right Latin for a minute, Sexagesima pars horæ, is as long and as tiresome to write as "Semivitreous oxide of lead" for the simple word "Litharge."

Numeri.	Numbers.
1. or j. unus, una, unum, one. unius, of one.	7. or vij. septem, seven. septimus, seventh. septimana, or, 7 mana, a week.
2. ij. duo, duæ, two. duorum, of two. duobus, in two, to two.	8. viij. octo, eight. octavus, eighth.
3. iij. tres, tria, three. tribus, in three, to three. trium, of three. ter, three times.	9. ix. novem, nine, nonus, ninth.
4. iv. quatuor, four. quartus, a, um, fourth. quater, four times.	10. x. decem, ten. decimus, tenth.
5. v. quinque, five. quintus, fifth.	11. xj. undecim, eleven. 12. xij. duodecim, twelve.
6. vj. sex, six. sextus, sixth.	20. xx. viginti, twenty. 24. xxiv. viginti quatuor, twenty-four.

Obstante, hindering, preventing.

Occasio, occasion, opportunity.

Octo, eight.

Octavus, eighth. Octava quaque horâ, every eighth hour.

Olim, some time ago.

Olla, a pot, a gallipot.

Omnis, all. Omni mane, every morning. Omni horâ, every hour. Omni bihorio, every two hours. Omni nocte, every night. Omni $\frac{1}{4}$ horâ, Omni quadrante horæ, every quarter of an hour. Tere omnia, rub all together.

Omnino, quite, wholly, entirely.

Optimè, very well, as well as possible.

Optimus, ma, mum, best.

Opus, need, occasion. Si opus sit, or fuerit, if it be necessary.

Ovillum juscum, mutton broth.

Ovum, an egg. Vitell. ovi, the yolk of an egg. Ovorum, of eggs.

P.

Pannus, a rag. Pannus linteus, a linen rag. Pannus laneus, Pannus è lanâ, a piece of flannel.

Paroxysmus, a paroxysm, a fit, a convulsion fit.

Pars, a part. Partes, parts. Parti dolenti, to the pained part.

Partitus, parted. Partitis vicibus means that you are not to give a medicine all at once, but divide the dose according to the directions most commonly previously given; for example, if a purging or emetic draught, half or a third (as the case may be) to be taken at first, and the other half or third at a certain distance of time, if the former quantity be not found sufficient to produce the desired effect.

Parvus, little. Cochleare parvum, a teaspoonful.

Parum, a little. Parumper, a little.

Pauculum, paucillum, paululum, a little.

Pectus, the breast. Pectoris, of the breast.

Pediluvium, a bath for the feet.

Per, by, or through.

Pergo, to go on with. Pergat in usu medicinarum, continue the medicines as before.

Peractus, { completed, perfected, quite done, gone through with; as, Peractâ opo-

Perfectus, { ratione emetici, after the emetic has quite done operating.

Perfricetur, let it be rubbed. Perfricandus, to be rubbed.

Pensus, weighed. Accuratè pensi, exactly weighed.

Perpetuus, perpetual. *Fiat perpetuum*, keep it open (when it refers to a blister).
 Pluvialis, also *pluviatilis aqua*, rain-water.
 Pharmacopœia, the dispensatory.
 Pollex, the thumb. *Pollex pedis*, the great toe.
 Pomeridianus, postmeridianus, in the evening or afternoon, time of the day.
 P. R. N., *Pro re natâ*, occasionally, according as circumstances may occur, according as the symptoms may require.
 Poculum, a cup.
 Pomum, an apple. *Pomi*, of an apple.
 Pone, behind. *Pondere*, by weight.
 Porro, moreover. *Post*, after. *Postea*, then, afterwards.
 P. M., *Post meridiem*, afternoon, in the afternoon, after twelve o'clock at noon.
 Potus, drink, any kind of beverage.
 Postulet, postulent, may require, demand.
 Præcipue, especially.
 Præparatus, prepared.
 Præparo, to prepare. *Præparentur*, let them be prepared.
 Primus, ma, mum, the first. *Primò*, first of all. *Primo mane*, very early in the morning.
 Prior, prius, the former, the first.
 Priusquam, before that.
 Pro, for. *Pro re natâ*, p. r. n., occasionally, &c. *Pro ratione*, according to, or, in proportion to; as, *pro ratione ætatis*, according to the age of the patient.
 Proprius, a, um, proper. *S. N. P.*, *Signetur nomine proprio*, mark it with its proper direction.
 Pruritus, an itching. *Pruriens*, itching. *Dolichos pruriens*, cowhage.
 Psora, the itch.
 Pulvis, a powder. *Pulveres*, powders. *Pulveribus*, in powders.
 Purificatus, a, um, purified.
 Pyxis, pyxidis, a pill-box, or lozenge-box.

Q.

Quadrans, *quadrantis*, *quadrante*, quarter. *Omni quadrante horæ*, every quarter of an hour.
Quacum, with which.
Quamprimum, as soon as possible, without the least delay.
Quartus, a, um, the fourth.
Quâqua, every. *Quâvis* (fœm), with any.
Quater, four times. *Quatuor*, four. *Quarta pars*, a fourth part.
Quaque (at the end of any Latin word), and.
Quem, *quam*, *quod*, which (the accusative case).
Qui, *quæ*, *quod*, which (the nominative case).
Quibus, to which, with which.
Quibusdam, to or with some. *Cum guttis quibusdam*, with a few drops.
Quilibet, *quælibet*, *quodlibet*, gen. *Cujuslibet*, abl. *Quolibet*, any.
Quiescat, goes to rest, is easier.
Quinque, five. *Quintus*, a, um, the fifth. *Quinques*, five times.
Quinquina (cinchona), Peruvian bark.
Q. S.—*q. s.*, *Quantum sufficiat*, as much as may be sufficient.
Quorum, *quarum*, of which. *Quos*, *quas*, which. *Quocum*, *quacum*, with which.
Quovis, with any.

R.

Ratio, a reason, also a proportion. *Pro ratione ætatis*, according to the age of the patient. *Pro ratione doloris*, according to the urgency of the pain.
Raucedo, hoarseness.
Recipe, take. *Recipe* (taken substantively), a prescription.

Regio, a region (an anatomical term for certain parts of the body); as, *regio epigastrica*, the epigastric region; *regio lumborum*, the region of the loins; appl. *emplastr. regioni umbilicali*, to the umbilical region or parts in the neighborhood of the navel.

Redigo, to reduce. *Redige*, reduce. *Redigetur*, may be reduced. *Redigatur*, let it be reduced.

Redactus, a, um, reduced. In *pulv. redact.*, reduced to powder.

Refrikerit—pl. int—the subjunctive future of *refrigesco*, to cool.

Reliquus, a, um, the rest, the remaining part.

Remedium, a remedy. *Continuandur remedia*, go on with the same medicines as before.

Repetatur, let be repeated, repeat.

Repetendus, to be repeated.

Respondeo, *responderit*, shall have answered. *Donec alvus ad sedes ij vel iij responderit*, until two or three stools shall have been procured.

Ribes, currants. *Gelatina ribesiorum*, currant jelly.

Ruber, *rubra*, *rubrum*, red.

Rubus idæus, raspberry.

Retinendus, to be retained or kept.

S.

S.V.R., *Spiritus vinosus rectificatus*, rectified spirit of wine.

S.V.Ten., or *tenuis*, proof spirit.

Sæpe, often; *sæpius*, oftener; *sæpissime*, very often.

Saltem, at least.

Sanguis, blood—in *inis*, of blood. *Sanguinis missura*, blood-letting.

Saphena vena, the ankle vein.

Scapula, the shoulder blade. *Inter scapulas*, between the shoulders.

Scilicet, to wit, namely.

Scrobiculus cordis, the pit of the stomach.

Secundus, a, um (adjective), second.

Secundum (preposition), according to.

S.A., *Secundum artem*, according to art; that is, you are to use your own ingenuity to do it in the most proper and scientific way.

Sed, but.

Sedes, a stool—plural, stools.

Semi, or *semis*, half. *Semihora*, half an hour. *Semi drachma*, half a drachm.

Sesqui, one and a half, as *sesquihora*, an hour and a half. *Sesquiuncia*, or *sescuncia*, an ounce and a half. *Sesquidrachma*, a drachm and a half. Remember well to attend to the difference of these two words, for many young men, by not knowing that *sesqui* means one and a half, but confounding it with *semi*, have made bad mistakes.

Semper, always.

Semel, once.

Septem, seven.

Septimana, a week, seven days. *Septimus*, seventh.

Sequens, following.

Serum lactis, whey. In *sero lactis vinoso*, in wine whey.

Serum is also the watery part of the blood which separates from the red part, or *crassamentum*, on standing until cold.

Sex, six; *sextus*, sixth.

Si, if. *Sive*, or; whether.

Signatura, a label or direction.

Signetur, let it be marked, directed, written upon.

S.N.P., *Signetur nomine proprio*, mark it with the name it is usually known by.

Simul, together; as, *terantur simul*, let them be rubbed together. *Simul ac*, at the same time that.

Sine, without. *Sine morâ*, without delay.

- Singultus, hiccup.
 Singulus, a, um, each; in singulis, in each; singulorum, of each.
 Sinister, tra, tram, the left. Auri sinistro, to the left ear.
 Satis, thirst; si sitiatur, if thirsty.
 Solitus, accustomed.
 Solus, alone; only.
 Solutus, a, um, dissolved, also loosened; as, donec alvus soluta fuerit, until a stool is procured.
 Solve, dissolve.
 Somnus, sleep. Hora somni, at bed-time.
 Spina, the spine, the backbone; also, a thorn.
 Spissus, thick. Spissitudo, thickness of consistence.
 Statim, directly, immediately.
 Stent, let them stand. Stet, let it stand.
 Sternutatorius pulvis, sneezing powder, snuff.
 Stupa, tow.
 Sub, subter, under. Sub finem coctionis, when the boiling is almost finished. Sub, prefixed to a word, implies diminution, or a process not completed; also, in many words has the same signification as the termination *ish* in English words, as sub-niger, blackish, not quite black. Subtepidus, lukewarm. For its meaning when applied to terms of chemistry, as subcarbonas, submurias, the pupil must study his chemical nomenclature.
 Subactus, a, um, subdued, dissolved.
 Subitus, a, um, Subitaneus, sudden. Subito (adv.), suddenly.
 Subtepidus, a little warm, lukewarm.
 Subige, dissolve it, make it unite. Subdue quicksilver with lard or balsam of sulphur.
 Subinde, frequently, now and then.
 Subtilis, subtile, reduced to fine powder. Pulv. subtilissimus, the very finest powder.
 Sumat, let him take. Sumatur, sumantur, let it be taken, take.
 Sudor, sweat.
 Superbibito, to drink after taking anything; as, chamomile tea or warm water after an emetic; or a cup of water or any liquid medicine to wash down a dose of any sort of pills.
 Superinfundo, to pour upon.
 Supra, above. Supradictus, above mentioned.
 Syncope, a fainting fit.

T.

- Tabellæ, tabulæ, lozenges.
 Talis, such. Sumat talem, let him take such a one as this.
 Talus, the ankle.
 Tam, so. Tamen, yet.
 Tactus, the touch.
 Tænia, the tapeworm.
 Tempus, temporis, time.
 Tempora, the temples. Temporibus, to the temples. Tempori dextro, to the right temple. Tempori sinistro, to the left temple.
 Temperies, temperamentum, temperament, degree of heat.
 Tenacitas, tenacity. Ad debitam tenacitatem, of a proper degree of tenacity or consistence.
 Teneo, to hold. Tenendus, to be held.
 Tenuis, weak, thin, small, slender.
 Tepefactus, warmed, made warm.
 Ter, three times, thrice. Ter quaterve, three or four times.
 Teres, teretis, round, taper; also, teres is a name for the long and round worm infesting the human body, qu. d. vermis teres.
 Tergum, the back. A tergo, behind.
 Tero, to rub. Tere (imperative), rub. Terendus, to be rubbed. Terantur, let them be rubbed.

Tertius, tertia, tertium, the third.
 Testacea, the prepared powders made of oyster-shells, egg-shells, crabs' claws, &c.
 Thorax, the chest. Thoracis, of the chest.
 Thus, frankincense.
 Tinea capitis, scald head.
 Torrefactus, toasted.
 Tres, tria, three; tribus, in three, to three.
 Triduum, three days.
 Trituratio, a grinding. Trituratus, triturated, ground. Tritus, ground.
 T.O., Tinctura Opii, what is commonly called Laudanum.
 T.O.C., Tinctura Opii Camphorata, Camphorated Tincture of Opium, called, formerly, Paregoric Elixir. This tincture is now called by the London College, in the late reform of the Pharmacopœia, edit. 1809, Tinctura Camphoræ Composita.
 Trans, through. Cola trans chartam, filter through filtering-paper.
 Tussis, a cough. Tussi molestante, when the cough is troublesome.

U.

Ultimus, ultima, mum, the last. Ultimó præscript, which was last prescribed.
 Umbilicus, the navel.
 Unà (an adverb), together.
 Uncia, an ounce. Unciam cum semisse, an ounce and a half.
 Undecim, eleven.
 Unus, una, unum, one. Unius, of one. Uni, to one.
 Urgente tussi, when the cough is troublesome.
 Urgeo, to urge, to oppress, to be troublesome or painful.
 Usus, use. Pergat in usu remediorum, continue the use of the medicine as before.
 Usque ad, up to, as far as.
 Ut, as, that, so that; in the same manner as.
 Utendus, to be used.
 Utatur, let him make use of.
 Uterque, utraque, utrumque, both. Utriusque, of both. Utrique, to both, to either.
 Admoveantur hirundines ij tempori utrique, apply two leeches to each temple.
 Utriuslibet, of whichever of the two, or more, the patient likes best.

V.

Vaccinatio, the act of inoculating for the cowpock.
 Variola vaccinæ, the cowpock.
 Vaccinum lac, cow's milk.
 Valde, very, very much.
 Valeo, to avail; si non valeat, if it does not answer.
 Variola, the smallpox.
 Varicella, the chickenpox.
 Vas, a vessel. Vasis, of a vessel. Vase clauso, in a covered vessel. Agitato vase, shaking the vessel.
 Ve, vel, or; either: -ve is never at the beginning, but the end, of a word.
 Vertebrae, the joints of the neck, back, or loins; the vertebrae, altogether, form that column of bones which is called the spine.
 Verus, true, real, genuine.
 Vena, vein. Vena saphena, the ankle vein.
 Venæsectio, bleeding. Fiat venæsectio, bleed him.
 V. S. B., Venæsectio brachii, bleeding in the arm. Fiat venæsectio in venâ saphena, bleed the patient in the ankle vein — or, it may be understood, to bleed wherever you can find the best vein at the top of the foot to get blood from.
 Vespere, in the evening.
 Vespertinus, in the evening, as horâ vi^a, vespertina, at six o'clock in the evening.
 Vehiculum, a vehicle; that is, whatever liquid or any other eatable or drinkable we take a medicine in, as barley-water, whey, jelly, or panada, &c.
 Viginti, twenty. Vicesimus, vigesimus, the twentieth.

Vinosus, vinarius, of wine. Cyathus vinarius, a wine-glass.
 Vices, turns; ad duas vices sumendus, to be taken at twice; that is, half to be taken at first, and the other half in some time after.
 Vicibus partitis. See Partitus in P.
 Vice, in the room of. Vix, scarcely, hardly. Ut vix sentiatur, so that it can scarcely be perceived.
 Vitellus, the yolk of an egg.
 V. O. S., Vitello ovi Solut; = dissolved in the yolk of an egg.
 Vitrum, a glass. Vitreous, made of glass.
 Vires, strength; si vires permittant, if the strength will bear it.
 Vomitio, a vomiting. Vomitione urgente, when the vomiting is troublesome.
 Vultus, the countenance.

OBSERVATIONS ON THE MANAGEMENT OF THE SICK ROOM.

In the treatment of disease, as much depends on the proper management of the patient, or, as it is usually called, *nursing*, as on the remedial powers of medicines appropriate to the exigencies of the case. How many, in fact, have owed their lives to the sedulous and skilful attention of their nurse or friends, wholly independent of professional assistance! and, on the other hand, what numbers have been sacrificed to a want of knowledge of the proper government of a sick room, in spite of the most judicious treatment on the part of the physician! This most important subject has been strangely overlooked by both lecturers and writers on the practice of medicine. We are acquainted with but two treatises on the subject, in our language, that can be referred to with confidence—the general observations prefixed to “A practice of Physic,” by the late Dr. Dewees, and “The domestic Management of the Sick Room,” by Dr. A. T. Thomson, of both of which we shall freely avail ourself in the following pages.

VENTILATION IN THE SICK ROOM.

Where it is possible, the room in which the patient is confined should be large and lofty, and in all cases provided with a chimney; the upper sashes of the windows should be capable of being let down, which is not always the case; in a word, the room should be of such a construction, as to permit the freest ventilation and renewal of the air without injury to the patient. Fortunately, for the comfort as well as for the benefit of the sick, a great change has taken place as regards their treatment in this respect; at one time, and that not many years since, it was considered essential to their recovery that air should be carefully excluded from their apartments, and every means was taken to accomplish this result; but a more rational mode of treatment is now generally adopted, though physicians occasionally meet with individuals who obstinately adhere to the idea that all access of the outer air is injurious to the sick.

The air of a sick chamber soon becomes impure, and must be changed by the admission of fresh air from without, and not, as is too frequently attempted, by the use of disinfecting agents in the chamber itself; which, although they may mask or destroy offensive odors, only tend to vitiate the air still more. It may be confidently asserted, that no disinfecting or fumigating agent is capable of rendering the air of a sick chamber so pure and salutary as it can be made by proper attention to cleanliness and free ventilation.

This latter can in most cases be accomplished without risk to the patient, by proper management, and should never be neglected. It should be regulated by the season of the year, the state of the atmosphere, and the character of the disease. In winter, in consequence of the draught occasioned by the fire, a renewal of the air of the room takes place to a greater or less degree; but a sick chamber should, if possible, be heated by an open fire, and not by a stove, as the air is thus more rapidly changed, and the unpleasant emanations caused by a large surface of heated iron are obviated.

The state of the weather should also influence the mode of ventilation, for it must be obvious that, if it is damp, it would be injurious, if admitted into a sick room directly or in large quantities. The character of the disease should always be considered; as a general rule, all acute affections require more ventilation than those of a chronic nature, except such as are attended with offensive discharges. No precise rules, however, can be laid down on the subject of ventilation as respects the mode in which it is to be attained; this must, in a great measure, be left to the discretion and good sense of the attendants.

Ventilation is especially required in infectious and contagious disorders, as, in close, ill-ventilated apartments, the power of infection is greatly augmented, and is, as it were, concentrated. The infectious miasm is greatly weakened or diluted by a free access of air, and its powers much augmented by a damp, close atmosphere. Next to ventilation, and of equal importance, is the

TEMPERATURE OF THE SICK ROOM.

There is a very general, but erroneous, opinion among nurses, that a sick person should be kept very warm, to prevent his taking cold, and, consequently, the temperature of the chamber is maintained at a fever heat, to the great inconvenience and positive detriment of the patient. The best general temperature of a sick room is from 60° to 70° F., to be regulated by the thermometer rather than by the sensations of the sick person, as these are oftentimes morbid and deceptive, but at the same time are not to be wholly disregarded; thus, where the temperature is agreeable to the sensations of the attendants, and the patient, notwithstanding, complains of chilliness, it should be increased a few degrees.

In febrile complaints, nothing conduces more to the comfort, and nothing is more proper than a reduced temperature, as it materially aids the physician in relieving the morbid heat of the patient's body, and the increased action of the arterial system. Thus, in a patient in the hot stage of fever, it is surprising to see the almost immediate relief experienced by the admission of cool air into the chamber; a tranquil state ensues, replacing his former jactitation and restlessness; his skin becomes cooler; his respiration, from being hurried and laborious, becomes calm and gentle; his pulse less frequent and active; and, in many cases, a placid and refreshing sleep comes on; or he breaks out into a general perspiration.

Where, on the other hand, the temperature of the room is maintained at a temperature equal to, or nearly equal to that of his body, every morbid symptom will be much aggravated, and his condition rendered still more deplorable. But, whilst a moderately depressed temperature is beneficial in many diseases, it exercises a decidedly injurious effect in others; in affections of the lungs, even a slight change in the thermometer will often cause an attack of cough and an augmentation of the symptoms.

During convalescence, the air of a sick chamber should be frequently renewed, and the temperature kept at about 60° to 65°; but no sudden transitions permitted, as nothing is more essential than a guarded care against extremes and rapid changes of heat and cold.

It is of equal importance that a proper attention should be paid to the temperature of the patient as regards the amount of his covering, for constant errors are committed by nurses in this respect. When a sick person complains of feeling cold, it is always proper to provide him with additional covering: but, if reaction takes place, and heat of skin ensues, it is seldom thought advisable by the attendants to remove any of the now superfluous clothes, "as he might take cold;" the consequence is that a febrile condition is produced, with a dry, hot skin, without a sign of perspiration. If some of the bedclothes be removed, however, this will soon make its appearance to the great relief of the patient. Every nurse should be aware that there is a sweating temperature, and that, when this is transcended, perspiration will not take place, and will cease if it be present. Attention to a simple rule on this point will obviate any difficulty,—to add covering when the patient complains of chilliness, and to remove it when he experiences too much heat.

CLEANLINESS IN THE SICK ROOM.

Nor is attention to cleanliness of less importance, for, although much is accomplished, in preventing a foul and offensive atmosphere, by proper ventilation, this is not sufficient, and is not always practicable. The observance of this is one of a nurse's most important duties; and she should be careful to remove anything that is capable of emitting an unpleasant smell, or of giving out an injurious gas, as speedily as possible. All evacuations are to be taken from the room at once, and never suffered to taint the air, under any pretence whatever. The bedclothes, as well as the personal clothing of the patient, should be changed as frequently as circumstances will allow, and no dirt suffered to accumulate in any part of the room. All articles used by the patient in taking his food, medicine, &c., should be cleansed as soon after they have been employed as possible. No slop-basin or slop-pail should ever be allowed to remain in a sick chamber.

A patient's face and hands should be frequently wiped with a towel moistened with cold water, or vinegar and water, in all cases in which there is no danger of causing chilliness, and he should be allowed to rinse his mouth and clean his teeth, several times a-day; when he is unable to perform the latter office for himself, it should be done by the nurse. This is very grateful to patients in fever, when the tongue is dry or coated with fur or tenacious mucus.

Whilst every means of ensuring cleanliness in the sick room should be practiced, it is to be accomplished with as little noise and bustle as possible; otherwise it becomes annoying and injurious to the patient. It is not requisite that the room should be swept, nor that the furniture should be dusted, every day. What is done should always be performed in the morning, as the patient is usually better able to bear the little noise and bustle necessarily attendant on these operations, after a night's rest and quiet. When the carpet is swept, it should be sprinkled with moist tea leaves a short time previous, to prevent dust from rising.

QUIET IN THE SICK ROOM.

In most diseases, and especially in those attended with fever, nothing is more annoying and distressing to the patient than noise; hence, the utmost quiet should be strictly maintained. All unnecessary conversation should be avoided, as a sick room is an unfit place for gossiping. If conversation be carried on, however, it should be in such a tone of voice that the patient may comprehend it, for nothing is more injudicious or hurtful to a sick person, especially if of a nervous temperament, than the mode usually adopted by the attendants of conversing in whispers, as it leads him to imagine that it refers to something they are afraid or unwilling to communicate to him, and hence, he fatigues himself by endeavoring to ascertain the purport of conversation, or gives way to despondency; on this account, it is better that all talking that is required should be carried on in a moderate tone of voice, and not in whispers, as much less likely to disturb the patient or to excite his attention.

In some cases, especially in those of a nervous character, on the contrary, cheerful conversation is of much benefit to the patient, provided it is not continued so long as to fatigue him.

A great fault in many nurses is being always in a bustle, "putting things to rights," or "fixing the room," thus constantly disturbing the patient, and preventing him from sleeping; nothing is more annoying to a sick person than this bustle or this constant recurrence of noises, which, however insignificant in themselves, render him nervous and impatient, from their repetition. Most of these may be prevented by an attention to order and method on the part of the nurse.

One common source of annoyance to a sick person arises from a frequent opening and shutting of his door, more especially if the lock or hinges are not in good order. In many cases, the door may be kept permanently open, and the patient protected from the light and draught of air by means of a screen. Where this cannot be done, the door should never be opened, except where it is absolutely required, and

it is by all means to be avoided whilst the patient is asleep. One means of avoiding a too frequent opening and shutting of the door is to permit no person to enter the room except such as are required to assist in attendance on the patient.

EXAMINATION AND PRESERVATION OF THE EXCRETIONS.

This is of much importance in many cases, as it enables the physician to judge with some degree of certainty of the condition of the patient and of the progress of the disease. It should never be trusted to a nurse, as it is very rarely that their account can be depended upon. In all cases where the physician deems it necessary, the nurse should be directed to preserve the excretions for his inspection; but they should never be retained in the sick room.

ADMINISTRATION OF MEDICINE, &c.

However skilful the physician may be, and however judicious his treatment, it is rendered perfectly nugatory, if the remedies he orders are not administered according to his directions, as the cure of the patient depends in many cases on the regular application of the prescribed remedies. It is, therefore, the duty of the attendant on the sick to follow implicitly the directions of the physician, as well in exactly complying with his orders, as in doing nothing that she has not been ordered to do. At the same time, there are exceptions to this rule, in which a suspension of the remedy, or a deviation from the order of the physician, is not only allowable, but is absolutely required. Thus, from idiosyncrasy or some other cause, the remedy in the doses ordered may have no effect, or may produce one widely different from that intended or expected. In such cases, it is evident that a strict adherence to the directions of the physician would be productive of evil; but he should be immediately apprised of the circumstance.

Nurses, however, more frequently commit errors in the administration of drinks and nourishment than in that of the prescribed medicines. In the former, they are apt to imagine that they are as good, if not better judges than the physician, and, in consequence, thwart the best directed treatment. The nurse should as scrupulously adhere to the directions of the physician as regards drink and nourishment, as with the doses and times of administration of medicines; nor should the exhibition of food or drink be left to her discretion more than that of remedies.

One error is very common, that of giving them too largely or too frequently; this should be guarded against, as far as possible, by the physician prescribing the quantity and quality of food or drink, as well as the times of giving it, with as much precision as he prescribes doses of medicine and the periods of their exhibition. In fact, in the treatment of the sick, as little as possible should be left to the discretion and judgment of the nurse or attendant of the patient.

FURNITURE OF A SICK ROOM.

Where a disease is of a serious character, or appears likely to be protracted, it becomes of importance that the chamber the patient is to occupy, if circumstances will admit of it, should be selected and arranged in such a manner as will most minister to his comfort, and aid his recovery. It should, if possible, be large and airy, with a northern aspect, so as to avoid the glare of the sun; but, in towns, a room exposed to the sun is preferable to one fronting on the street, in which the patient is constantly disturbed by the noise of passing vehicles, &c. As mentioned under the head of ventilation, it should always be provided with an open chimney, that a due circulation of air may take place.

No article of furniture that is not required should be suffered to remain in the room, as the dusting and arranging it will cause unnecessary noise and bustle, annoying to the patient. Two tables are wanted in most cases: one of a small size, to stand by the side of the bed, to hold such articles as are in immediate use, as the medicines he is taking, the spoon or glass in which they are administered, &c.; but nothing should be permitted to remain upon it, except articles frequently wanted.

The other table, which should be large, and adapted for the reception of medicines not in immediate use, such as spare spoons and glasses, should contain a drawer, supplied with a roll of old and soft linen, a sponge, scissors, a spatula, a roll of muslin and one of flannel bandage, some lint, some adhesive plaster, a pincushion well supplied with pins; in fact, all such articles as may be needed. It may appear unnecessary to allude to these things; but it has happened to every medical man, to observe that, without such previous preparation, much of his time is wasted in the search for what is wanted.

Where a second table is inconvenient, the top of a chest of drawers can be employed instead, for bottles, glasses, &c.; and one drawer should be appropriated to the reception of the various articles above enumerated; another should contain towels, a free supply of which should always be at hand; but none are ever to be used for the reception of dirtied articles of clothing, which should always be at once removed from the room. The washstand should be provided with additional basins, one of which should be of small size; and a large pitcher of water should be always kept in the room.

If there is a looking-glass in the room, it should be placed in such a position as not to be in view of the patient. A large sofa, or a small additional bed, is constantly required, as in certain cases where the patient is unable to sit up, he finds much comfort in being removed to it; it is also required when his bed is to be made. An invalid, or easy, chair is also productive of much benefit in certain cases; and where it can be commanded, it should be of such a construction as will admit of changes, so as to vary the position of the patient at pleasure.

What is called a French bedstead, without curtains, is the best adapted to the sick room. The four post bedstead, when used, should not be furnished with curtains, as these tend to exclude the air, and to retain smells and a vitiated atmosphere. In all cases, a mattress is preferable to a feather-bed; it may be placed over the latter, when greater softness is required. The pillows should be elastic, but so firm as not to permit the head of the patient to sink in them, as this prevents coolness and a due circulation of air. The bed-coverings should be as light as is consistent with warmth and comfort; hence, Marseilles quilts should not be used, as from their weight they oppress the patient, without affording as much warmth as a light blanket.

The change of bedclothes, and shaking up the bed or mattress, must be regulated by the character of the disease and the condition of the patient. In febrile complaints, it should be done, if the strength of the patient will admit, twice in the twenty-four hours. When there is much restlessness, it will be found advantageous to move him night and morning to a large sofa, or to another bed; this tends to promote sleep, and to insure a proper airing of the beds. When he is thus moved, the clothes of the bed he has left should be turned down, and fully exposed to the air, which will render so frequent a change of the sheets less necessary than would otherwise be required. When there is only one bed, and in febrile diseases, the sheets which have been used at night are to be replaced by others in the morning, and freely exposed to the air during the day, and again used at night. When the patient is too much debilitated to be moved from bed to bed, he should be merely removed to the other side of his bed, and the one he has left properly arranged by changing the linen, beating up the bed, &c.; but these changes are never to be made whilst he is in a perspiration.

If a fire in the room is required, it should not be made use of to prepare the food or drinks of the patient; this should be done elsewhere, as well to avoid the smell emitted by the heated articles, as the noise and bustle caused in their preparation.

PROPER USE OF UTENSILS FOR EVACUATIONS, &c.

As it is of the utmost importance to the patient that he should be spared all unnecessary fatigue, some precautions are to be used to prevent this during his evacuations; and one of the duties of a nurse is to enable him to perform those offices with the least expenditure of strength. When the patient is feeble, he should never be allowed to rise to fulfil the calls of nature, and therefore a bed pan and urinal should always

be provided; by means of these, the evacuations can be performed with ease, and without exertion. Many persons, it is true, have an aversion to using them; but this difficulty can generally be overcome by a proper representation of their advantages. One of the objections commonly made is that the use of a bed-pan gives them pain in the back; and this is often the case, where the patient is not properly supported; but can always be prevented by placing a pillow in the hollow of the back, before the pan is used.

In the case of giving drinks or nourishment, much unnecessary fatigue may be spared to the patient. Owing to the form of the vessels usually employed to administer drinks, without the sick person is raised in bed to receive them, much of the fluid is apt to be spilt on the bedclothes, to his great discomfort. Hence, a tumbler, teacup, or bowl should never be employed for this purpose, where the patient is too feeble to raise himself without inconvenience. The sick cup, as it is called, should therefore be employed for this purpose, as it spares all fatigue, and prevents any spilling of fluid on the patient or on the clothes.

When the medicine or food is taken in spoonful quantities, and in cases of nauseous medicines to be administered to children, the medicinal spoon will be found very useful. This is a spoon with a hollow handle, with an opening at its extremity; the bowl is provided with a hinged lid, but is open at the apex. When a fluid is poured into it, and the lid closed, the pressure of the air at the opening at the end of the bowl will prevent any of the fluid escaping, as long as the orifice at the end of the handle is closed, but is projected with some force when this opening is free. By means of this contrivance, medicine or food may be conveniently administered without the patient being obliged to be raised in bed, or in spite of resistance on the part of a child.

DOSE OF MEDICINES.

The doses of medicinal substances, being specific as regards each other, can only be learned by experience; but their remedial or active powers do not always increase with an increase of the dose: thus, if a purgative dose of calomel be ten grains, an increased quantity of the medicine will not induce an equivalent increase in the purgation, though it may cause other consequences. The dose in which a medicine is given often determines its specific action; two grains of ipecacuanha will usually quiet the stomach, or act as a diaphoretic, whilst thirty will produce active emesis; and, in like manner, almost every medicine displays different powers, according to the dose in which it is administered.

Dr. Paris states "that powerful doses are disposed to produce local rather than general effects." This is proved by experience, in many cases, but does not hold good in all: thus, many of the active poisons cause general effects in both large and small doses, and their virulent action is in general proportionate to the quantity taken. In most of what are termed the alterative medicines, however, the alterative action is only displayed when they are administered in small, but long-continued doses, and does not take place when they are taken in such quantities as to excite much local action. It is erroneous to suppose that a medicine is devoid of power, or does not exert an influence on the system, because no marked effect immediately follows its exhibition: for, as is observed by Dr. Barlow, "it is very possible that practitioners often err, especially in the treatment of chronic maladies, from requiring an obvious effect from each dose administered; where it is ascertained that a medicine actually possesses inherent powers, the slow and almost imperceptible exercise of these powers should not be despised. There is often more wisdom in seconding the efforts of nature than in superseding them." In fully admitting the truth of this remark, we are far from advocating the doctrine of Hahnemann, or his invisible doses; nor does it, in fact, accord with his views, for he gives a long catalogue of symptoms, caused by the administration of the minutest quantity of the articles of his *materia medica*.

So many circumstances influence the effects of medicines, that it is almost impossible to say what dose will produce identical effects in different individuals; but it

has been sufficiently ascertained that, in the generality of patients, we may reasonably expect certain results from certain doses. The circumstances that influence the action of medicines, and the doses in which they are to be given, are :—

AGE.—This exercises much influence; the young require smaller doses of a medicine to produce the desired effect than an adult; and the aged, although less susceptible to impressions than formerly, are unable to bear any forcible shock on their system. Various formulas or tables have been published for the graduation of doses to the respective ages; but none of them are as generally adopted as that attributed to Gaubius, which is as follows :—

Dose for a person in adult age being	1 drachm or 1.
That for a person from 14 to 21 years would be	2 scruples or $\frac{2}{3}$.
“ 7 to 14 “	$\frac{1}{2}$ drachm or $\frac{1}{2}$.
“ 4 to 7 “	1 scruple or $\frac{1}{3}$.
“ 4 “	15 grains or $\frac{1}{4}$.
“ 3 “	10 grains or $\frac{1}{6}$.
“ 2 “	8 grains or $\frac{1}{8}$.
“ 1 “	5 grains or $\frac{1}{12}$.

This table, although mainly correct, is often found to be erroneous, when applied to particular cases. Dr. Paris, in his *Pharmacologia*, gives a formula devised by Dr. Young, which is simple, and has been found to be generally correct; this is :—

For children under twelve years, the doses of most medicines must be diminished in the proportion of the age, to the age increased by twelve; thus:

At two years to $\frac{1}{7}$, viz :

$$\frac{2}{2+12} = \frac{1}{7};$$

At 21 the full dose to be given.

But no scheme can be devised, founded on age alone, to which there are not many exceptions. Thus, children require proportionably larger doses of castor oil or calomel to affect them than adults, whilst, and especially when very young, they are exceedingly susceptible to the action of opium and its preparations.

SEX.—It is generally admitted that females require smaller doses of medicines than males, and, as a general rule, this may be correct; but so much depends upon habits of life, temperament, &c., that the exceptions are almost as numerous as the examples.

TEMPERAMENT also exercises much influence on the doses of certain articles of the materia medica; an individual of a nervous temperament is unable to bear with impunity doses of opium that scarcely act on the phlegmatic; the sanguineous are readily affected by the action of stimuli, &c.

IDIOSYNCRASY, or constitutional peculiarity, has more influence than either sex or temperament on the action of medicines. Some of these anomalies are very striking, and almost incredible, and evince the necessity of particular inquiries being made by the physician of every patient with whose constitution he is unacquainted. In some cases, opium, or any of its preparations, cannot be administered without producing the most distressing symptoms; in others, the smallest doses of calomel will cause salivation; whilst in others again it may be profusely given without inducing any effect on the salivary glands. In some persons, the mere vicinity to the poison oak (*Rhus radicans*) will bring on an unpleasant inflammation of the skin; whilst on other individuals this plant has no action whatever. It is often found that copaiba will cause a peculiar eruption, and the smell of ipecacuanha induce a distressing dyspnoea. So tartar emetic will be productive of ptyalism in some individuals even when applied to the surface of the body; blisters of cantharides will cause strangury in some patients, &c. In fact, these idiosyncrasies are so numerous that it would require a volume to notice in detail those already recorded.

HABIT exercises an equal influence with idiosyncrasy in modifying the action of medicines. The usual consequence of a long-continued use of any remedial or stimu-

lating substance, is to diminish the susceptibility of the organs on which it acts to its impression. Hence, to obtain its effects, it must be given in gradually increased quantities. In many cases, however, instead of a diminished, an increased action is caused by repetition of certain substances.

The influence of habit, in diminishing the susceptibility of the system to stimuli, is liable to fewer exceptions as regards vegetable than mineral substances, as may be daily remarked in the use of opium and tobacco. Dr. Christison accounts for this effect of habit in diminishing the influence of vegetable substances, by an increased power of the stomach in decomposing them; but it must also be attributed to a diminished susceptibility in the nervous system to their influence. But mineral medicines appear, in almost every case, to act with increased power by repetition, though the converse has been observed in some cases.

STATE OF THE SYSTEM.—It is well known that, in certain deviations from a healthy or normal condition, a remarkable change is often observed in the action of remedies on the system. Thus, in severe pain, opium can be administered in doses that cannot be borne with impunity in a state of health; but this is still more the case in tetanus, in which scruple doses, frequently repeated, have been administered without producing any marked effects. This modifying influence of disease has been noticed with regard to many other remedies, as in mercury, which can be given in large and repeated doses, in a febrile condition, without causing salivation. Where there is a diminished sensibility of a part, or a comatose condition, the pain, or irritation caused by the application of stimulating or vesicating articles, not being felt by the patient, there is a risk of inducing a sloughing and destruction of the part to which they are applied, if care be not taken to remove them in time; so, also, the application of ammonia to the nostrils, to relieve fainting, has caused violent irritation, and even inflammation of the trachea, the patient being unable, from his condition, of appreciating or expressing his sensation. Dr. Christison notices several fatal cases of bronchitis from this cause.

TIME OF DAY at which medicines are administered also influences their action on the system. As a general rule, no medicine should be given soon after a meal, though, in the case of some of the irritants, as arsenic, corrosive sublimate, iodine, &c., it is often found most expedient to administer them whilst the stomach is filled with food, to avoid their local action on that organ. Dr. Paris observes, under this head: "Evacuating medicines ought to be exhibited late at night, or early in the morning; thus, substances of tardy operation, as calomel, and the resinous purgatives, should be given at night, while saline purgatives, senna, &c. may be given in the morning. It would seem that during sleep the bowels are not so irritable, and, consequently, not so easily acted upon, which allows time for the full solution of the substance." In all cases, the exhibition of purgatives should be so timed that their operation may take place during the day, so as to avoid an undue disturbance of the patient at night. Other directions will be found in a subsequent part of the work.

Numerous other causes exercise much influence on the action of medicines, as climate, the imagination of the patient, the tissue or organ to which they are applied, &c.

Nor should the *variable activity* of a medicine be overlooked; this has occasioned more accidents than is generally supposed. Some remedies, and generally those of a vegetable nature, lose much of their activity by age, and, therefore, require to be given in large doses to produce the desired effect; hence, where a physician has been induced to increase the usual dose of some article, which in a recent or unaltered state is active, and even poisonous, but, from age or other deteriorating cause, has become weakened, serious accidents may occur, if he persist in administering the same doses of a parcel which is fresh, or which is obtained from another apothecary. In such cases, the safest plan is always to commence with the usual dose of the medicine, and to gradually increase it, if this be found necessary.

It should also be borne in mind that certain medicines have the power of accumulating in the system, or, in other words, of not displaying their usual effects when given in small and repeated doses, until a certain quantity has been taken, when, on a sudden, symptoms resembling those caused by an overdose of the article, make their

appearance. This has been remarked in relation to mercury, arsenic, and several other of the metallic medicines, as well as iodine, digitalis, &c. In giving remedies of this character, the physician should always be cautious, and not repeat the dose of them too frequently, because no sensible effect is produced at the expected time.

INTERVALS BETWEEN DOSES.—As every medicine produces a specific and definite action on the system, which, in most cases, continues for a certain time, and then ceases, it becomes necessary to renew or maintain the medicinal impression as long as is required to counteract the morbid condition of the system. In some cases, however, as in the case of an emetic or purgative given for the purpose of merely evacuating the stomach and bowels, the single effect produced may be sufficient, and no repetition of the dose is required. In most attacks of disease, on the other hand, the physician is obliged to prescribe medicines in such quantities as to produce a certain impression, and to repeat the doses at such intervals as will keep up the influence of this impression. Where, from a want of knowledge of the time during which the remedy he has administered will continue to exercise its full action on the system, he postpones a repetition of it for too long a period, although a fresh impression is made by each dose, but little real good is the result, and, from a want of continuity in the remedial impression, no progress is made towards a cure. Hence, a physician should be acquainted with the usual duration of the action of the medicine he is administering, and order a repetition of the doses so as to keep the system under its influence. A nurse should, therefore, in all cases, follow his directions implicitly, as regards the periods at which medicines are to be given, for, by inattention to this rule, injustice is done to the physician, and, what is of more importance, the life of the patient is endangered. Where the diffusible stimulants are administered, especially those of an evanescent character, it is of much consequence that their influence should be kept up; and hence a neglect on the part of a nurse in giving them at the prescribed periods might cause a fatal collapse of the patient. It should be observed, however, that, except in cases where it is necessary thus to maintain the vital powers of a patient, this strict observance of the prescribed intervals between the doses of medicines is not intended to extend to the night, so as to disturb the invalid, as, in most diseases, sleep is one of the best restoratives; but advantage is to be taken of those moments when he is awake, timing the doses as nearly as possible to the designated periods.

RULES FOR THE ADMINISTRATION OF MEDICINES.—As has been stated, fluid medicines are usually administered in drops, teaspoonfuls, tablespoonfuls, &c.—an approximative measurement, which, although not precise or accurate, is sufficiently so in most cases, though not suited to medicines of a very active nature. As the generality of medicines are nauseous, it is necessary to modify, as far as possible, their disagreeable taste, which renders them so repugnant to patients, and often causes much difficulty in their administration, especially to children; but this modification must not be of such a character as to alter their properties. As regards young children, much difficulty will be obviated by the use of the medicine-spoon already described; this will also be found useful in cases of insanity, where patients refuse to take medicine or nourishment.

It is often found that patients experience much difficulty in swallowing pills, especially when these are of small size. The usual mode is to place the pill on the tongue, and to take a mouthful of water, when, on making an effort to swallow, both pass down the throat together; this plan is generally successful; but some persons appear to be unable to take a pill in this manner, and gorge themselves with fluid without effecting the deglutition of the pilular body; in such cases, by enveloping the pill in some bread, or some conserve or sweetmeat, so as to form a bolus, the difficulty is obviated.

Mixtures containing insoluble ingredients should always be shaken up before they are administered; otherwise, the first doses will contain but little of the active ingredients, whilst the last portions will, on the other hand, be more powerful than was intended. Where the article administered is of a volatile nature, it should be given

the moment it is poured out, and the bottle containing the remainder immediately closed.

As each of the classes of medicinal articles requires the observance of certain rules to insure its full benefit, these will be briefly noticed in an alphabetical order.

ACIDS.—These are given to fulfil various indications; some acting as tonics, some as refrigerants, &c. As those most generally used are corrosive, they require to be diluted with some bland fluid, so as not to exercise too powerful a local action on the tissues with which they come in contact; nor should they be kept in a metallic vessel, nor given by means of a metal spoon. As most of them act on the enamel of the teeth in a very rapid and energetic manner, the mouth should be well rinsed with water after the administration of each dose; or they may be taken by being sucked up by means of a small glass tube, thus avoiding any contact with the teeth: this observation applies to many of the acid or super-salts; the usual solution of the sulphate of quinine, which contains free sulphuric acid, is known to act on the teeth very rapidly, and, from a constant repetition of the doses, to injure them in a serious manner; and hence requires to be given with precaution.

ANTACIDS are medicines intended to neutralize an undue or morbid acidity of the stomach or intestines. Their action is chemical, and they produce an innoxious compound which takes the place of the morbid fluid that causes the uneasiness or pain. In many cases, they act most efficaciously, when an emetic or cathartic has been previously administered. To obtain a permanent benefit from them, they should be combined with tonics, aromatics, &c.; as, when given alone, the relief afforded by them is but transient.

They are most advantageously administered early in the morning, or some time previous to a meal during the day. When the salt resulting from the union of the antacid and the morbid acid is purgative, the former is better given at bed-time; thus, the combination of the solution of potassa, of the carbonates of potassa and soda, or of magnesia, forms salts with the gastric acid that act with some energy on the bowels; but this is not the case with chalk mixture or lime water.

The patient should be restricted to light animal food and well boiled vegetables, avoiding sweets, acids, pastry, and, in fact, everything liable to increase the aced condition of the stomach.

ANTILITHICS AND LITHONTRIPTICS are medicines supposed to possess the power of obviating or dissolving urinary calculi. That there are remedial agents capable of removing or relieving that morbid condition of the system on which a lithic acid diathesis depends, there can be no doubt; but it is very questionable if we possess remedies capable of dissolving calculi existing in the urinary organs. Among the antilithic remedies, those most generally employed are magnesia and its salts, the alkalies, soap, several of the acids, some astringent diuretics, many bitter tonics, &c. In cases of a lithic acid diathesis, however, much depends on the regulation of the diet of the patient; thus, when there is a superabundant secretion of uric acid, the food should contain as little azote as possible, and, therefore, be mainly of a vegetable nature; but, at the same time, care must be taken that the healthy action of the stomach is not impaired. There should be an avoidance of all indigestible substances, as fresh bread, pastry, salt meats, and even soups of all kinds; nor should malt liquors or acid wines be allowed. Where the white sand deposits or the phosphates occur, an acid system of diet is beneficial, with abstinence from all alkalies, soda water, &c.; whilst lemonade, the acid wines, and fruits in moderation can be allowed with advantage.

ANTISPASMODICS are remedies which have been considered to be endowed with the power of allaying spasm, and of calming or entirely removing irregular actions of the nervous system. As spasmodic action of the muscular apparatus and a morbid condition of the nervous centres may arise from a variety of causes, it has been found that the most opposite modes of treatment have been required; thus, in some cases, the most efficient antispasmodic is the lancet, whilst in others the morbid symptoms are best alleviated by the most active diffusible stimuli. It is not our intention to enter into the discussion of the mooted question of the *modus operandi* of antispasmodics,

or whether there exists a class of remedies which are endowed with the power of exerting a direct control over spasmodic action. It is, however, shown by experience that the influence of this class of remedial agents is very limited in their operation and very various in their effects, and that to be efficient much discrimination is requisite on the part of the physician in the selection of the article best calculated to attain the desired effect. In general, what are termed antispasmodics have some of the properties of the narcotics, but differ from them in not causing stupor or insensibility. They are mostly substances possessed of a strong odor, and of a more or less volatile nature; and hence also partake of the properties of stimulants.

ANTHELMINTICS are medicines which are capable of expelling or destroying worms situated in the intestinal canal. Several kinds of worms are apt to infest the human intestines, and often give rise to distressing and even fatal symptoms. The medicines calculated to dislodge them are of various kinds: some act mechanically, as active purgatives, powder of tin, cowhage, &c.; some by being poisonous to the worms, as pink root, pomegranate bark, the vegetable bitters, &c.; and to this class belong, in all probability, the acrid, volatile, and resinous vermifuges, as turpentine, camphor, garlic, &c.; and the mineral remedies, as calomel, the preparations of iron, sulphur, &c.; some seem to act by asphyxiating the worms, as the fatty oils, the mucilages, &c.

Vermifuge medicines should be given in as simple a form as possible, and in as large doses as is compatible with their nature, and the age and condition of the patient; and hence the plan of administering them in the form of lozenges, biscuits, &c., is erroneous. One of the most certain of these remedies is the pink root (*spigelia*), either in the form of simple infusion, or united with senna or savin; in the first case, it should be followed by the administration of a brisk cathartic. In consequence of its sometimes causing temporary loss of sight, tremors, &c., which, although seldom dangerous, are apt to frighten the relations and friends of the patient, it is better, especially in cases of children, to administer the infusion in the evening, so that its effects may not be observed. The worm-seed oil is also another native anthelmintic of great power, but has the objection of having so disagreeable an odor and taste as to prevent its general use; the least unpleasant mode of giving it is in emulsion with mint or cinnamon water.

Whatever vermifuge is employed, and whatever success attends its administration, the treatment should not be confined to the mere expulsion of the worms; but the condition of the stomach and bowels should be attended to, and their tone rendered healthy by the use of tonics, and a well-directed and generous diet.

CATHARTICS are those medicines which increase in a greater or less degree the peristaltic action of the intestines. Cathartics have been divided into *laxatives*, which are characterized by acting mildly, without stimulating in any great degree the vessels of the intestines, or exciting a general disturbance of the system; and *purgatives*, which, in addition to evacuating the contents of the bowels, also promote secretions from their mucous coat, giving rise to copious watery stools. Some purgatives have a more violent action, occasioning nausea, faintness, pain, tenesmus, &c.; and some, acting in an overdose as acrid poisons, causing great irritation, and even inflammation of the intestines. But there is another and important character in the action of purgatives, which is that different portions of the intestinal canal are most affected by different cathartic substances; thus, calomel acts, in a marked manner, on the upper portions of the tube, as is shown by the increase of the biliary matters in the evacuations; gamboge also manifests its power on the stomach by often causing vomiting, but also, like the other drastics, causes much irritation in the large intestines; aloes, again, exerts very little action on the upper part of the canal, but spends its force on the lower portion; in fact, each purgative substance is characterized by a peculiarity of action as regards the part of the intestinal canal affected by it, and the nature of the evacuation it produces; and hence, it is of importance, in prescribing an article of this class, that it be suited to the impression wished to be made, and not, as is too frequently the case, selected at random. No class of remedies has been productive of more injury than the present, from their being resorted to without judgment, and from being persevered in for too long a time; for, inde-

pendent of the irritation they create and keep up in the bowels, they impair digestion, and too often lay the foundation of fatal organic diseases. Purgative medicines may be exhibited at any period during the day; but in all cases the patient should not be exposed to the influence of cold air; if, however, the circumstances are not urgent, it is better to administer them late at night or early in the morning, so as to prevent the disturbance and loss of rest of the patient, and also to secure their retention by the stomach; when given on an empty stomach, they also act more speedily and effectually than when this viscus is filled with food. To promote their action, and to obviate their griping effect, warm diluents, as chicken water, gruel, &c., should be freely taken after the first discharge. Where they operate too much, this inordinate action is to be checked by laudanum given by the mouth, or in the form of injections.

Many of these medicines are very nauseous, and so repugnant to the patient as to induce nausea and vomiting, and, therefore, require the addition of correctives. The taste of rhubarb is best disguised by being given in milk, and that of castor oil by the froth of porter. The disagreeable taste of infusion of senna is much lessened if made with cold water, or, if made in the usual manner, by the addition of strong coffee, black tea, or by a few grains of cream of tartar. Aloes is rendered more palatable by the addition of extract of liquorice. Magnesia is mixed more perfectly with water or milk, if it is poured on the surface of the fluid, and permitted to sink as it becomes saturated, than by the opposite course. The neutral salts are rendered more palatable by adding a small portion of lemon juice to their solution, &c.

When a cathartic is given in the form of pill, its operation is always less speedy than when administered in powder or mixture; and it should be borne in mind that the drastic purgatives operate more mildly, and even effectually, in combination than when given alone, as is instanced in the compound cathartic pill.

Enemata or Clysters.—This mode of purgation is now generally employed in our large towns; but a great prejudice exists among many persons against its use, from a fastidious and mistaken delicacy. In consequence of the improved apparatus now to be procured, the administration of these remedies is attended with much less difficulty and exposure than formerly, when the pipe and bag or common syringe were employed; and, therefore, much of the objection to their use is obviated.

Clysters are of almost indispensable utility when it is necessary to evacuate the bowels as speedily as possible, or when the stomach will not bear the administration of a purgative by the mouth, and also in cases where it is of importance to make a direct application to the lower bowels, as in dysentery, colic, &c. Where it is merely wished to open the bowels, an injection of tepid water will often be found sufficient; where this is not found sufficient, the addition of table salt, sweet oil, or molasses, will, in most cases, induce a full discharge. In all cases, the patient should be directed to retain the injection for as long a time as possible, and not to attempt to empty his bowels immediately after the reception of the medicine. Though clysters seldom can be forced beyond the great curve of the colon, and hence their local action is confined to the lower part of the bowels, still, in many cases, their purgative influence extends to the small intestines, and their administration is followed by full and copious evacuations. They are also directed for other purposes than evacuating the bowels, for, from the intimate connection of the rectum with the other pelvic viscera, impressions made upon it are speedily transmitted to the adjoining parts, and hence, remedies can be directed by this channel with much advantage. In irritation of the bladder or uterus, an anodyne injection into the rectum will often afford much relief. In diseases of the lower bowels, also, clysters are of almost indispensable utility, as also in the dislodgement of ascarides seated in the rectum; nor are they less beneficial in those cases of sudden sinking of the powers of life where deglutition is impossible, and yet a prompt stimulating impression is requisite to save the patient; under such circumstances, clysters of some of the diffusible stimuli have proved of the greatest benefit.

Suppositories.—These are substances introduced into the rectum to induce a discharge of its contents. Their use is principally confined to the mere object of re-

lieving costiveness, for which purpose a piece of soap of a conical shape, and of about an inch in length, is the most generally employed; but in some cases, articles of a more stimulating character are necessary. In cases of costiveness in children, the introduction of the end of a small roll of paper, moistened with oil, for a few moments into the rectum, will, in many cases, be found sufficient. The objection to their use is the irritation they often create in the rectum. Sometimes suppositories of opium are found useful in those cases in which anodyne enemata are usually employed, and may be used where patients have a repugnance to the use of clysters, or where the soreness of the rectum prevents the introduction of the pipe of a syringe.

In all cases where enemata or suppositories are employed, they must be introduced with care; otherwise serious injury may be done to the mucous coat of the rectum; instances have occurred where, from a careless insertion of the pipe of a syringe, the gut has been seriously affected, giving rise to fistula, &c. In most cases, the injection for a child under one year is about an ounce, which is to be gradually increased to five or six ounces, as it advances in age; to a youth, from ten to fifteen years old, a pint may be given; whilst an adult requires a pint and a half to a quart. Where, however, the injection is of an anodyne nature, to restrain discharges or to alleviate pain, the laudanum should be mixed with not more than half an ounce to an ounce of bland mucilage, so as not to excite contraction of the bowels by the stimulus of quantity.

DEMULCENTS OR EMOLLIENTS are medicinal agents which diminish tone or irritation in the tissues to which they are applied, and thereby cause a relaxation of the parts, or sheathe and protect them from the action of substances capable of acting on them injuriously.

The term *Emollients* is generally employed to designate remedies which are applied externally to soften and relax external surfaces, and, by contiguous sympathy, the deeper seated parts; these effects have been referred by some to a physical, and by others to a vital, agency. They consist principally of bland watery mixtures, or of simple water, at a somewhat elevated temperature, of oleaginous substances, or those combined in various forms, as fomentations, poultices, &c., which will be more particularly noticed in a subsequent part of the work.

Demulcents are bland substances of various kinds, administered internally, to prevent the action of irritating matters on mucous membranes, or to soothe or lubricate these parts when irritated or inflamed. They are useful in irritation and inflammation of the stomach and bowels, in like conditions of the air-passages and lungs, as well as in affections of the urinary apparatus. When prescribed in cases of catarrh, and other morbid conditions of the respiratory organs, they are usually directed to be taken *ad libitum*; but they should be swallowed slowly, for their main efficacy arises from their lubricating effect on the parts with which they come in contact; and this soothing influence is extended by sympathy to the bronchial membranes and pulmonic structure. They should be taken after a fit of coughing, for, as, by that spasmodic effort, the acrid secretion of the inflamed mucous surface is thrown off, they are enabled to come in direct contact with the irritated part, and thus exercise their soothing influence more efficiently.

DIAPHORETICS are those medicinal agents which promote and increase cutaneous transpiration. These articles are numerous, and of a very heterogeneous character. In general, warm diluents, aided by external heat, are very effectual means of promoting perspiration; whilst, in some cases, this excretion is better promoted by the use of cool liquids, or even by the sudden and temporary application of cold to the surface. They are in almost every case relative agents; and, to produce the desired effect, should be given in certain states of the system, and with certain precautions. Thus, during the exhibition of these remedies, the patient should be confined to bed, and covered with light blankets; his shirt should be of flannel or muslin, as these are good non-conductors of heat. Their action is much promoted by the free use of diluents, given either warm or cold, according to the temperature of the patient's skin. During the continuance of the perspiration, great care should be taken that

it is not suddenly checked from any cause. When it is wished to moderate the sweating, the patient may be removed to another bed, covered with a warm blanket, his clothes changed, and his body dried with dry towels, care being taken that he is not exposed to cold air. During the exhibition of diaphoretics, no medicines of a purgative or diuretic character should be administered, as the action of the latter is incompatible with that of the former, besides causing a frequent exposure to cold during their operation.

DILUENTS are watery fluids which increase the fluidity of the blood, and render the secretions and excretions less acrid or viscid. The best diluent is water, which is to be given either alone or variously flavored, according to the taste of the patient or the circumstances of the case. The temperature at which a diluent is given has much influence on its action. Thus, water at 60° or 70° F. merely dilutes; whilst at 45° to 60° it acts somewhat as a tonic, and at 70°, and to as high a temperature as it can be drunk, it is stimulant. Merely tepid water will often act as an emetic, whilst at 90° or 100° it will quiet the stomach and relieve nausea. In a cool or cold state, diluents diminish the temperature of the body and moderate an undue excitement of the pulse, and are hence called *refrigerants* or *temperants*, and principally consist of aqueous solutions of many of the acids, or of some of the neutral salts.

DIURETICS are medicines which increase or promote the secretion of urine: this they do by increasing the quantity of fluid taken into the stomach, or by stimulating the kidneys to increased action. They are always inconstant in their effects, and cannot be relied upon to produce the desired effect in all cases. To obtain their effect, the surface of the patient's body should be kept cool; otherwise, they are apt to operate as diaphoretics. Hence, the patient should remain out of bed, and the medicine be administered during the day-time; or, if confined to bed, the clothing should be as light as is compatible with comfort. The action of diuretics is much aided by a free use of the cooling diluents.

EMETICS are those medicines which invert the natural action of the stomach, so as to cause an ejection of its contents through the mouth, or vomiting. When the patient is plethoric, blood-letting should always precede the administration of an emetic, as it not only promotes its operation, but prevents any injurious rush of blood to the head. When it is required, a full dose of the emetic substance is to be administered; but, in the generality of cases, it is better to give it in divided doses, frequently repeated, until the desired effect is produced. It often happens, when a physician orders an emetic, that he will be asked not to give a strong one, as the patient is too weak, &c.; but it should be known that, unless the medicine causes full vomiting, it will create more uneasiness and debility than when it acts in the desired manner. When vomiting takes place, it should be encouraged by draughts of tepid water or weak chamomile tea, to be repeated after each discharge; but the use of these fluids is not to be carried too far, because, when the stomach is gorged with liquids, the action of vomiting is often not fully performed, or is wholly checked. When the operation of an emetic is too violent, the best means of checking it is by the administration of laudanum, or the application of fomentations or warm poultices to the epigastrium; if these fail, a spice plaster or a sinapism to the same part will generally be found effectual. If the stomach continues irritable, drink, food, or medicine is to be given in very small quantities. There is always much thirst, which patients are very apt to endeavor to alleviate by freely taking some fluid; but this in most cases being rejected, the evil is only increased; the best plan to relieve it is to frequently wash the mouth with some cool drink, swallowing as little as possible. When the stomach is inactive, and a full dose of an emetic substance does not cause vomiting, it is always unsafe to repeat the dose, as by such a course inflammation may be produced. When the stomach thus resists the action of one article, carried to a due extent, it is better to have recourse to another, or to endeavor to produce emesis by titillating the fauces and throat by means of a feather. Emetics should always be given at the time ordered by the physician, and, as a general rule, on an empty stomach, in the morning; but it is sometimes of benefit to administer them

in the evening, as, from the exhaustion produced by their operation, a disposition to sleep is caused.

EMMENAGOGUES are medicines which excite or promote the catamenial discharge; but there are, strictly speaking, few remedies which have this specific power on the uterus, the majority of them acting by their influence on the system generally, or on parts contiguous to the uterus—ergot being almost the only article which exercises a direct power on this organ, and that rather in increasing its expulsive energy than in promoting the menstrual function, though it has been advantageously employed for the latter purpose. To insure the successful administration of this class of remedial agents, the system must be previously prepared for their use by invigorating it, if there are relaxation and debility; and by an opposite course, if there is an undue degree of arterial action. In a majority of cases, the restoration of the discharge is rather attributable to a proper regulation of the system than to any specific power in the emmenagogue administered.

EPISPASTICS are external applications which are capable of inflaming the skin, and causing an effusion of serum beneath the cuticle. These effects can be produced by a variety of means; but that most generally employed is the Spanish fly, or cantharides. The mode of applying a blister, and of dressing the sore it occasions, is very simple; but, without special directions from the physician, many nurses and attendants on the sick are strangely ignorant on the subject. The best preparation of cantharides for blistering purposes is the cerate of Spanish flies, or common blistering plaster; this will be always found effectual, if properly prepared, and of good materials. It is best spread on soft leather, though linen, muslin, or even stout paper will answer. To obtain a prompt action from a blister, it should be spread thick; powdered flies should not be sprinkled on its surface, for, if the plaster is well made, it requires no such addition; added to which, the powder is apt to adhere to the skin, and to give the patient much uneasiness. Before applying a blister, the part on which it is to be placed should be well washed with soap and water, and thoroughly dried, the hair shaved off, and the skin irritated by being rubbed with a coarse towel, or, if this is not sufficient, with warm vinegar, spirits of turpentine, &c. The plaster is to be secured to the part by means of a bandage; or, what is better, by means of strips of adhesive plaster, or by coating the margin of the leather or cloth on which the blister is spread with the adhesive plaster. Whenever either of the last modes of securing a blister is resorted to, it is necessary, to secure its adhesion, that the skin be perfectly dry. The plaster should be in perfect contact with the skin; but should not, as is too often the case, be subjected to much pressure, for this retards vesication, and is a source of uneasiness to the patient.

The period generally required for a blister to draw is twelve hours; but this is liable to many exceptions: thus, for children, delicate females, and for persons with tender skins, a much shorter time is sufficient. The best plan, when the patient complains of pain and burning, is to lift a corner of the plaster and examine the condition of the skin beneath, and, if it be found vesicated or fully reddened, to remove the blister; or, if the skin is not sufficiently inflamed, to permit the blister to remain till this result is attained. When the skin is sufficiently reddened, but not vesicated, the application of a warm poultice, or of a dressing of basilicon ointment, will cause a copious effusion of serum. In cases of children and irritable persons, or those liable to strangury, this should always be done. To guard against this latter, it has been advised to interpose a piece of gauze, fine muslin, or tissue paper between the skin and the plaster: these should be moistened with oil. It is very doubtful, however, whether this plan is of any real advantage.

In many persons, the action of a blister disposes to sleep; but in very sensitive individuals it causes much irritation; this may be often obviated by the addition of a few grains of extract of hyoscyamus, or of belladonna, to the plaster. When it is requisite to cause a blister as rapidly as possible, the plaster of Spanish flies must be replaced by other means. In such cases, paper, moistened with oil of cantharides, or with the ethereal extract, will be found useful. The application of a pledget, wet with water of ammonia, will also act very rapidly. When a still more rapid effect is

desired, the direct application of heat to the part may be resorted to, by means of a disk of polished metal, heated in boiling water, or by applying a piece of wet blotting paper to the skin, and passing a hot flatiron over it.

Some persons have a great dread of blisters, and object to the application of a large one; but, when the full effect of blistering is required, a large one should always be applied, for the pain and irritation caused by a large blister are very little greater than a small one occasions, whilst the benefit is more decided. Still, as the quantity of the discharge is in a great measure equivalent to the size of the vesication, large blisters depress more than small ones; therefore, when they are intended merely as counter-irritants, small blisters are to be preferred. The size and shape of blisters vary greatly, according to the part to which they are to be applied, and the age of the patient. Dr. Dewees gives the following as the general size of blisters:—

For the legs or thighs, from 7 to 8 inches long by 3 to 3½ broad.

For the back, from 7 to 8 inches long by 4 to 4½ broad.

For the thorax, from 8 to 9 inches long by 7 to 8 broad.

For the stomach, from 8 to 9 inches long by 6 to 7 broad.

For the abdomen, from 10 to 11 inches long by 8 to 10 broad, if designed to cover the whole surface.

For the ears, from the peculiarity of the shape, the size cannot well be defined.

For the temples, from an inch to an inch and a half in diameter; they are usually circular.

These are the sizes for adults; for younger persons they must, of course, be proportionably smaller.

The dressing of a blister requires some skill on the part of a nurse; and much pain and inconvenience will be saved to the patient by the exercise of a little dexterity in the operation. Some trouble will be avoided by having everything necessary for the operation in readiness, as the plasters spread, bandages ready, &c. Each of the vesicles is to be carefully snipped, with the points of a pair of sharp scissors, at their most depending part, and the serum evacuated; but when the vesications are very small and numerous, this should not be attempted, as it would occasion unnecessary fatigue to the patient. In such case, the larger vesicles only are to be opened, and the smaller left till a subsequent dressing, when it is probable they will be found much increased in size. The cuticle should not be removed from the surface of the blistered part, except when it is wished to keep up a continued discharge, in which case the cuticle is best removed by the application of a hot poultice. When it is wished to heal the blistered surface, the dressing should be of simple cerate, thinly spread on a linen rag, and renewed twice a day. When the object is to maintain the discharge for a short time, the dressing should consist of basilicon ointment. When a more permanent impression is desired, some more stimulating application, as savin ointment, must be used. A blistered surface should seldom be washed, as this often causes chilliness and fatigue, and is generally useless. Some persons become faint when a blister is dressed for the first time: this should not excite alarm, as it is a mere sympathetic effect. The plaster for the dressings should not be larger than the blistered surface, as it causes irritation of the sound skin without aiding the discharge. Blisters sometimes, instead of healing properly, become painful and inflamed, and assume the appearance of an eroding or a phagedenic sore; this is best remedied by the application of a soft bread and milk poultice, having its surface anointed with fresh lard or cold cream. If this does not succeed, a lotion of lime-water and linseed oil will often be found effectual; oxide of zinc ointment will likewise prove useful in some cases. When a blistered surface causes much itching, and renders the patient restless, washing the irritated surface with an infusion of flaxseed or slippery elm will, in most cases, relieve the uneasiness. One of the most distressing consequences, arising from the application of a blister, is the production of strangury, which invariably takes place in some persons whenever a blister is used; this is remedied by a free exhibition of emollient and mucilaginous diluents, opiates by the mouth and as an enema, external fomentations to the pubic region, &c.; in some cases, a clyster of solution of sulphate of soda will act very efficaciously. To prevent its occurrence, it

has been found of benefit to incorporate camphor and opium with the epispastic ointment, say twelve grains of the former and four of the latter, for a blister six inches square. It is said that boiling the flies in water, previous to their formation into an ointment, will deprive them of the property of causing strangury.

ERRHINES are medicines which, snuffed up the nostrils, cause an increased discharge of the nasal mucus, and generally the convulsive action of sneezing. They are now seldom prescribed, though undoubtedly useful in some cases; but they should never be employed in persons disposed to apoplexy, or in those affected with hernia, prolapsus uteri, &c.

ESCHAROTICS are topical agents, which, by a chemical or mechanical action, are capable of eroding or disorganizing the solid tissues of the part to which they are applied: those which act mechanically are actual cauteries, as a heated iron, moxa, &c.; those which act chemically are caustics, as fused potassa, nitrate of silver, and chloride of zinc. In cauterizing with a heated iron, this should be at a white heat, as, at this temperature, it occasions less pain to the patient, from its causing an immediate death of the part to which it is applied. In applying it to any part, the surface should be protected by some non-conductor of heat, but not by wet paper or cloth, as the sudden extrication of steam will produce a blistered surface around the burn, and will much increase the pain. The hot iron is rarely employed in this country, except to arrest hemorrhage; in leech bites, where the usual means of stopping the bleeding are ineffectual, the application of the end of a heated wire to the wound will, in most cases, be found effectual. Another mode of applying the actual cautery is by means of moxa. This term is used for a small cone or cylinder of some slowly burning substance, applied to the skin, set fire to, and suffered to burn slowly for a length of time proportionate to the effect intended to be produced; this is a very painful operation, and seldom employed in the United States. The chemical cauterants are numerous, and of various degrees of activity. In using the most powerful of them, caustic potassa, some precautions are necessary, owing to its great deliquescence; when it is employed to form an issue, layers of adhesive plaster should be placed on the skin, with a hole of the proper size cut in their centre, through which the caustic can be applied without danger of its spreading. The chloride of zinc, which is an excellent cauterant, is generally employed by forming it into a paste with flour. The chloride of antimony, though very powerful, from being a fluid, is difficult to manage, but has been advantageously used in cases of poisoned wounds.

EXPECTORANTS are medicines which promote the expulsion of mucus or other matters from the air-passages. It is an ill-defined class of remedies, and is extremely uncertain in its action; but we cannot, in this place, examine the propriety of retaining a division of the materia medica, the articles composing which act in an obscure and by no means regular manner, and in almost all cases by a remote or secondary influence of a nauseating or sedative character. At the same time, daily experience proves that much benefit is derived, in diseases of the respiratory organs, from the administration of substances which are usually considered expectorants, in whatever way they may primarily act on the system. No class of remedies has been more productive of injury than that of so-called cough medicines; nor in any one has quackery exercised a more unbounded sway. It is too common for persons to resort to their use in cough and catarrh without consulting a physician, thus often aggravating a trifling irritation until it becomes a serious, and oftentimes a fatal, inflammation; for, as many of the articles composing them, especially those most frequently employed in domestic and empirical practice, are stimulants, much mischief is apt to follow their indiscriminate use. The following rules should be observed in the exhibition of expectorants—and should always be enforced on the patient. The surface of the body should be kept moderately warm, so as to promote a gentle diaphoresis, and nothing done that may interfere with the primary operation of the remedy on the nervous, lymphatic, or muscular systems, by the secondary effect of which the lungs are benefitted; hence, the use of any that excite diuresis, and more especially

of purging, should be avoided; nor should the patient be allowed to indulge his appetite in food or drinks of a stimulating character.

NARCOTICS are medicines which lessen the manifestation of vital phenomena dependent upon the nervous system, especially deadening sensibility and diminishing the motor power: their full operations produce sleep, or even coma. The objects most generally in view in the administration of these articles are the production of sleep, or the alleviation of pain. No medicines are so much modified in their action by idiosyncrasy and habit. This does not extend to them as a class, but to each individually; hence, it is of importance that a physician should ascertain from a patient, before prescribing a narcotic, whether he has been long or frequently in the habit of using it, or whether it causes any unpleasant effects upon him, so that some other article of the class may be substituted. As regards the time at which a narcotic should be given, it depends upon the cause for which it is to be administered. Thus, if prescribed to allay pain, it may be given at any time, and repeated as long as the pain remains; if to procure sleep, it is best given at night, and so late as to prevent the patient from being disturbed before it produces its full effect; otherwise, it is apt to exercise a stimulating action. In many persons, the secondary effect of opium is to cause much nausea and headache after its narcotic action has ceased, which are best relieved by a cup of strong coffee, or by vegetable acids. When, however, this or any other narcotic produces these consequences, its use should be relinquished, and some other substituted.

REFRIGERANTS are remedies which are supposed to diminish the heat of the body without causing a marked diminution of sensibility or nervous energy, and they are thought to do this, whether administered internally or applied externally; but it is very doubtful whether any medicines act as internal refrigerants independently of the coldness of the medium in which they are administered, except in a secondary manner. Diaphoretics act as refrigerants by inducing an increase of superficial evaporation; sedatives, by lessening the force and rapidity of the circulation; purgatives, and especially the saline, by the copious watery evacuations they produce. The great refrigerant is cold water, whether applied topically or given internally. As a topical application, it is often mixed with fluids of greater volatility, that, by a more rapid evaporation, a proportionate abstraction of heat may take place.

SEDATIVES are those remedial agents which diminish muscular tonicity, and the manifestation of nervous influence, even to the production of syncope, or a suspension, for a certain time, of the functions of the systems affected. Some act generally; others more locally. Blood-letting and tartar emetic are examples of the first; hydrocyanic acid, of the second. Most of the articles of this class are capable, if given in too large doses, or too long continued, of producing a depression of the vital powers which may induce death; and hence require to be administered with much care and judgment.

SIALAGOGUES are medicines which induce an increased secretion from the salivary glands. The modes in which the articles composing this class act are various. Some are merely local irritants, as pellitory, horseradish, &c.; and, when chewed, produce such a local stimulation of the salivary glands as to produce a copious effusion of saliva and mucus. Another and more important section includes such articles as evince their sialagogue powers in a secondary manner, which does not take place until a powerful impression has been made on the system generally; to this belong mercury, gold, &c., and their preparations. In the first of these divisions, their medicinal powers depend wholly on sialagogue effects; hence, they are merely useful as counter-irritants; the local stimulation, and consequent increased secretion produced by their action, often acting advantageously in relieving inflammations and congestions of adjoining parts. The curative value of internal sialagogues, however, is not dependent on the salivary flow they occasion, for though, in many cases, this may be very great, yet it is not essential to their remedial action, and may, in fact, rather be considered as a proof that their full constitutional influence has been attained.

In administering mercury, with a view to the production of its constitutional effect, it should always be recollected that, if it be permitted to exert its sialagogue powers to any great extent, it always causes great distress to the patient, without producing a beneficial effect on the disease; and its operation should, therefore, be closely watched. The condition of the gums should be examined daily, and, as soon as they display evidence of the action of the remedy, it should be discontinued, or given at longer intervals. The constitution of the patient should be sedulously attended to, as mercurials are apt to produce unpleasant consequences in delicate, nervous individuals, and are badly borne by the aged, who have suffered from previous attacks of illness. It is always dangerous to push them to salivation in children, as it frequently produces sloughing of the gums and cheeks. When too great a degree of salivation has been induced by the mercurials, it is of importance to check it as speedily as possible. This is to be attempted by cool air, and by washing the mouth with an infusion of some vegetable astringent, a solution of opium, or one of acetate of lead; this latter is very efficacious, but has the property of rendering the teeth black. Marked benefit has also been derived from a weak solution of chloride of soda; much reliance is placed by some practitioners on the preparations of sulphur, especially the soluble sulphurets, both as a wash in solution, and administered as a powder. An infusion of the smooth sumach (*rhus glabrum*), as well as a strong solution of borax, has likewise been found very useful as a local application. When the inflammation is severe, the administration of cathartics, especially the saline, is requisite, and in some cases blisters and leeches. Where a tendency to slough exists, the application of diluted pyroligneous acid will be found advantageous.

STIMULANTS are remedies which increase the vital activity of the system, or of a single organ, the first being termed *general*, and the latter *local* stimulants. Much discrimination and caution are required in the administration of articles of this class, because, if given when inflammation is present, they are liable to create more mischief than benefit; but they are called for when, on the decline of that condition of an organ, or organs, a state of relaxation or torpidity exists. In this state of things, a gentle stimulation materially assists the functions, and is productive of much benefit.

TONICS are closely allied to stimulants, and some of them act in this manner; but most of them do not cause any sensible excitation of the system, whilst, at the same time, they gradually impart tone and activity to the vital powers. With the exception of iron, most tonics act primarily on the nervous system, and secondarily on the muscular, on which latter their power is most distinctly marked. Iron appears to act in a different manner, by altering or improving the condition of the blood, and consequently increasing the tone of the system. One of the most valuable properties of many articles of this class is that of curing diseases of a periodic type; in what manner, it is difficult to explain, as the laws that govern periodicity are but little understood.

In prescribing the antiperiodic tonics in fevers, as cinchona and its salts, their use should be confined to the intermissions, and, in most cases, in as large doses as the stomach will bear with impunity, so timing the doses that one may be taken a short time before the expected paroxysm. Quinia and its preparations, although eminently antiperiodic, are not possessed of the tonic properties of the bark itself, and, in the large doses recommended of late, are apt to induce unpleasant affections of the head, as vertigo, ringing in the ears, headache, and mental delusions. In administering arsenic as an antiperiodic, its effects should be carefully watched, and be discontinued when it causes constitutional symptoms, as vertigo, and oedema; nor should it be given in debilitated habits, and where the pulse is weak and feeble. It is better, in all cases, to commence its use in small doses, and to increase them until the stomach is somewhat affected, when the quantity is to be diminished, or even discontinued.

MANAGEMENT OF CONVALESCENCE AND RELAPSES.

The management of a patient, after the cessation of an attack of disease, and when he is declared convalescent, is of equal importance with the treatment during the continuance of the malady; and yet it is one strangely neglected by the physician, and is too often trusted to nurses and friends. In all recoveries from sickness, the organs, or parts which have been affected, remain for some time in an excitable condition, and liable to recurrence of the disease on the slightest provocation; and it is of much consequence to inquire what causes operate most frequently in thus producing a relapse. It will be found that they generally consist in the indulgence in food, either in too great quantity, or of an improper character, or in the too early or imprudent exposure to the weather, or in over-exertion of any kind.

Diet. — The stomach, in common with every other part of the body, suffers from a diminution of its normal vigor and power of digestion, and, as a necessary consequence, when food of too nutritious a character is taken into it, it is unable to digest it properly; hence, part of it remains in an undigested state, oppressing the weakened organ, and is either rejected by vomiting, or causes diarrhoea; or, if the stomach is capable of converting it into nourishment, an undue stimulation of that organ ensues, which will result in fever. This latter state of things is also induced by the administration of stimulating drinks. No animal substance, in any form, should be allowed during the continuance of acute febrile disease, nor immediately after its cessation. When it is permitted, it should be at first as little stimulating as possible, and in small quantities at a time; but is to be gradually increased in quantity and quality as the patient gains strength. An invalid should be very cautious in the use of certain articles, considered by nurses and others as peculiarly nourishing and proper for the sick, namely, animal jellies of all kinds. These are exceedingly indigestible, even in a state of perfect health—added to which, they are, in most cases, rendered highly stimulating by the addition of wine or spices.

Exercise. — Another frequent cause of a relapse is over-exertion, from the common, but erroneous, notion that a convalescent will not recover his strength, except he takes as much exercise as he can bear. Properly regulated exertion is highly serviceable; but it should never be carried so far as to produce exhaustion, and should be pursued for some time in doors, before it be attempted in the open air: the latter, at first, should always take place in a carriage, that can be opened or closed at will; the patient may then attempt short walks in the open air; but, in all cases, it is of importance that he is not unduly fatigued, as otherwise injury instead of benefit will be the result. One of the most serious errors, committed with regard to exertion, is that of permitting a convalescent to sit up too frequently, or for too long a time, under the mistaken notion of giving him strength. A patient should never be allowed to sit up longer than is agreeable to his feelings, and never so long as to produce a sense of fatigue.

Air. — Great care must be taken that an invalid is not exposed to cold or damp air, especially if his skin be disposed to moisture; to guard him from sudden changes of temperature, he should wear flannel next to his skin, and have his feet well protected by warm stockings. In every convalescence, from whatever disease, country air is far more invigorating than the vitiated atmosphere of towns; but care must be exercised in the selection of the location, as to its freedom from malaria, &c., for it is far preferable that he should remain within the precincts of the city, than expose himself to the influence of causes which act with peculiar activity on a system debilitated by previous disease.

UNIVERSAL FORMULARY.

A

ABSINTHIUM.

WORMWOOD.

This name is applied to several species of *Artemisia*, all, however, possessing the same properties. That recognized in the U.S. Pharmacopœia, and most generally employed, is *A. absinthium*, a native of Europe, and commonly cultivated in our gardens.

Ser. Syst. Syngen. super. *Nat. Syst.* Astera-
ceæ.

Linn. Sp. Pl. 1188. Griffith, Med. Bot. 405.

The parts used, are the tops or extremities of the branches. These have a grayish-white color; a soft, silky feel, a fragrant odor, and a very bitter taste. They impart their properties to water or alcohol.

Wormwood has been employed as a stimulating tonic, in various forms of dyspepsia, in amenorrhœa, chronic leucorrhœa, gout, vermiform affections, &c. It is given in a variety of forms.

INFUSION OF WORMWOOD.

R. Wormwood, one ounce.
Boiling water, one pint.

Infuse. Dose, one to two fl. ounces. *Brande.*

EXTRACT OF WORMWOOD.

R. Wormwood, one part.
Boiling water, six parts.

Boil, express, and evaporate to a proper consistence. Dose, ten to twenty grains. Tonic and stomachic. *Giordano.*

CLYSTER OF WORMWOOD.

R. Wormwood, }
Rue, } each, three drachms.
Savine, }
Boiling water, one pint.

Boil, strain, and add to each half pint
Castor oil, half an ounce.

Foy.

TINCTURE OF WORMWOOD.

R. Wormwood, one part
Alcohol, eight parts.

Digest with a gentle heat for five days, express, and filter. Tonic, stomachic, and vermifuge. Dose, from twenty drops to two fl. drachms.

Guibourt.

ALKALINE TINCTURE OF WORMWOOD.

R. Wormwood, }
Tansy, } each, eight parts.
Centaury, }
Bog Bean, }
Alcohol, one hundred and twenty parts.
Carbonate of Potassa, one part.

Digest for six days, and filter. Dose, one or two fl. drachms. In dyspepsia and flatulent colic.

Hoffmann.

SYRUP OF WORMWOOD.

R. Wormwood, two ounces.
Boiling water, one pint.

Infuse for 12 hours, strain, add to the filtered liquor twice its weight of sugar, and make a syrup. Dose, a tablespoonful. *Par. Codex.*

WINE OF WORMWOOD.

R. Wormwood, two parts.
White wine, fifty parts.

Digest for four days, express, and filter. A stomachic bitter and anthelmintic. Dose, one fl. ounce. *Taddei.*

OIL OF WORMWOOD.

R. Tops of wormwood, one part.
Olive oil, eight parts.

Heat for two hours on a water-bath, stirring often, express, and filter. As an embrocation to the abdomen, in cardialgia, diarrhœa, &c.

Wurtem. Ph

(75)

ESSENTIAL OIL OF WORMWOOD.

R. Tops of wormwood, at will.
Water, sufficient.
Distil, and collect the oil that floats on the product. Dose, two to four drops several times a-day. *Par. Cod.*

COMPOUND SPIRIT OF WORMWOOD.

R. Wormwood, four pounds.
Juniper, eight ounces.
Cinnamon, two ounces.
Angelica root, half an ounce.
Alcohol, eighteen pints.

Macerate for fourteen days, distil twelve pints, pour back and re-distil ten pints. *Guibourt.*

RATAFIA OF WORMWOOD.

R. White sugar,
Water, each, forty ounces.
Dissolve, and add
Orange-flower water, six fl. ounces,
previously beaten with
white of egg, one.
Then add
Compound spirit of
wormwood, fifty-six fl. ounces.
Heat on a water-bath, cool, and filter. *Guibourt.*

ACACIA.

GUM ARABIC.

This gum is derived from several species of *Acacia*, as *A. vera*, *A. senegal*, *A. arabica*, &c.
Sex. Syst. Polygam. monœcia. *Nat. Syst.* Fabaceæ.

Willdenow. *Sp. Pl.* iv. 1056. Griffith, *Med. Bot.* 270.

In tears or fragments of various sizes, often rounded; colorless or of a yellowish tint, semi-transparent, inodorous, of a slightly sweetish mucilaginous taste. Wholly soluble in water, forming a mucilage. Insoluble in alcohol. Demulcent and nutritive.

MUCILAGE OF GUM ARABIC.

R. Powdered gum Arabic, four ounces.
Boiling water, half a pint.
Add the water gradually to the gum, and rub together, till the mucilage is formed. Strain. Principally used to suspend insoluble powders or to diffuse oily and resinous substances through aqueous fluids. *U.S. Ph.* 1850.

MIXTURE OF GUM ARABIC.

R. Mucilage of gum Arabic, three fl. ounces.
Sweet almonds, ten drachms.

Sugar, five drachms.
Water, (Imp.) two pints.

Blanch the almonds, beat them to pulp with the sugar and mucilage, add the water gradually, constantly stirring. Strain. Demulcent and emollient. *Ed. Ph.*
Used to appease cough and to allay irritation.

GUM ARABIC MIXTURE.

R. Mucilage of gum Arabic, six fl. ounces.
Oil of sweet almonds, three fl. drachms.
Syrup of saffron, one fl. ounce.
Solution carbonate ammonia (*Lond.*), half fl. drachm.
Mix, and make emulsion. A tablespoonful occasionally, in catarrh. *Ainslie.*

GUM LOZENGES.

R. Gum Arabic, four ounces.
Starch, one ounce.
Sugar, one pound.
Pulverize, mix, and form a consistent paste with sufficient rose-water, to form lozenges of convenient size. A useful article to allay irritation of the throat in catarrh. *Ed. Ph.*

PECTORAL PASTE OF GUM ARABIC.

R. Gum Arabic, two pounds.
Dissolve in water, and add
Sugar, one pound and a half.
Evaporate to a proper consistence, adding
Orange-flower water, three fl. drachms.
Employed to allay irritation, like the last. *Foy.*

PATE DE GUIMAUVE.

R. Gum Arabic,
powdered, two pounds, four ounces.
White sugar, two pounds.
Whites of eggs, twenty.
Double orange-flower
water, six fl. ounces.

Dissolve the gum and sugar in a sufficiency of water, strain, next day decant, evaporate by a gentle heat to consistence of honey, add the whites of eggs and stir briskly, continuing the evaporation; when almost complete add the orange-flower water, and continue to stir; pour on a marble slab, and keep in a tin box. *Cules.*

GUM ARABIC PASTE.

R. Gum Arabic, one pound.
Dissolve on a water-bath, in water, eight fl. ounces.
Evaporate to the consistence of honey.
Add
Whites of eggs, six.
Orange-flower water, two ounces.

Beat well together. When sufficiently consistent, run into moulds.

Used for the same purposes as last. *Beral.*

SYRUP OF GUM ARABIC.

R. Gum Arabic, two ounces.
Sugar, fifteen ounces.
Water, eight fl. ounces.

Dissolve the gum in the water without heat, then the sugar with a gentle heat, and strain. A good addition to pectoral and other mixtures. *U. S. Ph. 1851.*

COMPOUND SYRUP OF GUM ARABIC.

R. Syrup of gum Arabic, four fl. ounces.
Sulphate of morphia, one grain.
Oil of sassafras, one drop.
Hoffman's anodyne, half fl. drachm.
Mix. Known as Jackson's cough syrup, and much used in catarrhs, &c. Dose, a tablespoonful every three hours. *Jackson.*

LINCTUS OF GUM ARABIC.

R. Gum Arabic, one part.
Clarified honey, three parts.
Incorporate the gum with the honey. The dose is a teaspoonful occasionally, to relieve irritation of the throat in catarrh. *Beral.*

WENDT'S MIXTURE OF GUM ARABIC.

R. Mucilage gum Arabic,
Linden-flower water, equal parts.
Found useful in doses of a teaspoonful every hour in the diarrhoea of infants. *Radius.*

TRONCHIN'S LOZENGES OF GUM ARABIC.

R. Powdered gum Arabic, eight ounces.
Precipitated sulphuret of antimony,
Anise, each four scruples.
Extract of liquorice, two ounces.
Extract of opium, twelve grains.
White sugar, two pounds.
Mix, and form lozenges of six grains, one to be used occasionally in catarrh and bronchial affections. *Tronchin.*

ACETOSELLA.

WOOD SORREL.

Sex. Syst. Decand. Pentag. *Nat. Syst.* Oxalidaceæ. *Lindley.*

Oxalis acetosella. Linn. Torrey and Gray, Fl. i. 211. Griffith, Med. Bot. 208.

A native of Europe and the United States. The whole plant is used. It is agreeably acid, from its containing a binoxalate of potassa. It

is refreshing and cooling, and is said to be useful in scorbutic affections.

EXTRACT OF WOOD SORREL.

R. Wood sorrel, one part.
Bruise in a mortar, express the juice, and pour on the marc,
Water, three parts.
Boil for an hour, express, evaporate, adding the juice, to the consistence of an extract. Acid and bitter. Dose, a scruple to half a drachm. *Pideret.*

ACETUM.

VINEGAR.

Impure and very dilute acetic acid, containing about five per cent. of real acid. It is the result of acetous fermentation of any fluid which is susceptible of vinous fermentation. It is principally obtained in Europe from malt or weak wine, and in this country from cider. It should be distilled for pharmaceutical use.

It is employed internally as a refrigerant and diuretic, and is a grateful addition to diluent drinks in febrile affections. It is also sometimes used as a clyster. Externally it is beneficial as a fomentation, and likewise forms a useful addition to gargles and collyria.

U. S. Dis

VINEGAR MIXTURE.

R. Vinegar, three fl. ounces.
Honey, two fl. ounces.
Water, three pints.
Mix. As a cooling drink in fevers. *Ammon.*
R. Vinegar, one fl. ounce.
Acetic ether, one fl. drachm.
Syrup of raspberries, three fl. ounces.
Water, two pints.
Mix. As a refreshing drink in fevers. *Augustin.*

MIXTURE OF VINEGAR AND CARDAMOMS.

R. Vinegar, one fl. drachm.
Comp. tincture of cardamoms,
Simple syrup, each, half a n. ounce.
Water, ten fl. ounces.
Mix. To be taken in such portions as the stomach can bear. In sick headache.

VINEGAR PASTILLES.

R. Powdered white sugar, four parts
Heat moderately, and add
Vinegar, one part.
Make pastilles. *Neuman.*

GARGLE OF VINEGAR.

R. Vinegar, two parts.
Honey,
Barley water, each, eight parts.
Mix. As a mouth-wash or gargle in inflamed fauces. *Soubeiran.*

R. Vinegar, two fl. drachms.
Muriate of ammonia, one drachm.
Honey, one and a half fl. ounces.
Water, twelve fl. ounces.
Mix. As a gargle in inflamed fauces. *Ratier.*

SYRUP OF VINEGAR.

R. Vinegar, eleven fl. ounces.
Sugar, fourteen ounces.
Boil together. *Ed. Ph., 1841.*

Mixed with water, this forms a pleasant and refreshing drink in febrile complaints. Dose, half to one fluidounce.

VINEGAR CATAPLASM.

R. Vinegar, one part.
Flour, three parts.
Make cataplasm. An astringent and antiseptic application. *Beral.*

DISTILLED VINEGAR.

Distil a gallon of vinegar by means of a sand-bath, from a glass retort into a glass receiver. Discontinue the process when seven pints shall have been distilled, and keep these for use. *U. S. Ph.*

ACIDUM ACETICUM.

ACETIC ACID. (*Sp. Gr. 1.041.*)

R. Acetate of soda, in powder, one pound.
Sulphuric acid, half a pound.
Red oxide of lead, one drachm.

Pour the acid into a glass retort, gradually add the acetate of soda, and by means of a sand-bath, at a gentle heat, distil the acetic acid, till the residuum becomes dry. Mix the distilled liquid with the red oxide of lead and again distil, with a moderate heat, to dryness. *U. S. Ph., 1840.*

DILUTED ACETIC ACID.
(*Sp. Gr. 1.004.*)

R. Acetic acid, (*sp. gr. 1.041*), a pint.
Distilled water, seven pints.
Mix. *U. S. Ph.*

AROMATIC VINEGAR.

R. Acetic acid, one pint.
Camphor, two ounces.
Oil of lavender, one drachm.
Oil of cinnamon, twenty drops.
Oil of cloves,
Oil of rosemary, each, thirty drops.

Mix. *Gray.*

R. Acetic acid, one pint.
Oil of cloves, one drachm.
Oil of rosemary, two scruples.
Oil of bergamot,
Oil of cinnamon, each, half a drachm.
Oil of pimento, twenty-four grains.
Oil of lavender, one scruple.
Oil of neroli, ten drops.
Camphor, one ounce.
Alcohol, half an ounce.

Mix. *Cooley.*

Aromatic vinegar is used as a pungent and reviving perfume in fainting, &c. As it is corrosive, it should not be allowed to come in contact with the skin or clothes. It is also prepared in the dry way, as follows:

R. Crystallized acetate of soda, one drachm.
Sulphuric acid, twenty drops.
Oil of lemons,
Oil of cloves, each, three drops.

Mix, and keep in a glass-stoppered bottle. *Gray.*

R. Dried rosemary, one ounce.
Origanum, each, half an ounce.
Lavender, half a drachm.
Bruised cloves,
Acetic acid, (1.068), one pint and a half.

Macerate for seven days, strain, express, and filter. *Edin. Ph., 1839.*

AROMATIC SPIRIT OF VINEGAR.

R. Strong acetic acid, one pint.
Camphor, two ounces.
Oil of lavender, nine minims.
Oil of cinnamon, twenty minims.
Oil of cloves, half a drachm.

Mix. To be kept in a glass-stoppered bottle. *Paris Cod.*

ACETIC ACID CATAPLASM.

R. Rye meal, three parts.
Diluted acetic acid, one part.
Mix. Used as an application to ill-conditioned ulcers. *Taddei.*

ACETIC ACID CLYSTER.

R. Barley water, six fl. ounces.
Diluted acetic acid, one to three fl. ounces.

Mix. The proportion of diluted acid may be varied according to the indication. Has been advised in obstinate constipation, ileus, &c.
Swediasur.

PROPHYLACTIC VINEGAR.

R. Dried rosemary, and sage, each, one ounce.
Dried lavender flowers, half an ounce.
Bruised cloves, half a drachm.
Distilled vinegar, two pounds.

Macerate for seven days, press, and filter.

Ed. Ph., 1817.

This is intended as a substitute for the complicated formula formerly used, for *Thieves' vinegar*, at one time so celebrated as a preservative against the plague. It has no advantages over it.

VINEGAR OF THE FOUR THIEVES.

R. Wormwood, }
Roman wormwood, } each, one
Rosemary, } ounce and
Sage, } a half.
Mint, }
Rue, }
Lavender flowers, } two ounces.
Garlic, }
Calamus, }
Cinnamon, } each, two drachms.
Cloves, }
Nutmegs, }
Distilled vinegar, eight pints.

Digest by a moderate heat, in a closely stopped matrass, for twelve days, strain, express, and filter, then add one ounce of camphor dissolved in alcohol.
Lewis.

ACIDUM ACETICUM EMPYREUMATICUM.

PYROLIGNEOUS ACID. (*Ed. Ph.*)

Pyroligneous acid (*Sp. Gr.* 1.034) is an impure acetic acid, obtained by a destructive distillation of wood. It contains creosote, empyreumatic oil, &c., and has been employed in gangrene to correct the fetor and to promote the separation of sloughs, and also as an application to unhealthy ulcers.

U. S. Dispens. Dunglison, N. R. 6th. ed. p. 13.

MOUTH WASH.

R. Pyroligneous acid, half a fl. ounce.
Cinnamon water, four fl. ounces.
Syrup of mulberries, two fl. ounces.

Mix. Used as a mouth-wash in cancrum oris. It should be kept in an opaque glass vessel to prevent decomposition.
Phæbus.

INJECTION OF PYROLIGNEOUS ACID.

R. Pyroligneous acid, two fl. drachms.
Distilled water, six fl. ounces.

Mix. In purulent discharges from the ear.
Buchanan.

CATAPLASM OF PYROLIGNEOUS ACID.

R. Bran, half a pound.
Linseed meal, one ounce.

Mix well, and add Pyroligneous acid, q. s., to make a cataplasm. As an application to foul ulcers.
Reece.

ACIDUM ACETICUM CAMPHORATUM.

CAMPHORATED ACETIC ACID.

This is a mixture of camphor and acetic acid, which might very properly have been placed under the head of acetic acid. But as the *Edin.* and *Dub.* Pharmacopœias recognize it by the above title, we place it separately for the convenience of reference.

CAMPHORATED ACETIC ACID.

R. Camphor, half an ounce.
Acetic acid, six and a half fl. ounces.

Reduce the camphor to powder by means of alcohol, and dissolve it in the acid. The proportions in the *Dublin Pharmacopœia* are one ounce to ten.
Ed. Ph. 1841.

CAMPHORATED VINEGAR.

R. Camphor, thirty parts.
Vinegar, twelve hundred and fifty parts.

Powder the camphor with a small quantity of strong acetic acid, add the vinegar and set aside for a few days. Then filter.
Codex.

R. Camphor, thirty parts.
Distilled vinegar, one thousand parts.

Dissolve.
Raspail.

The above substitutes for *Henry's aromatic spirit of vinegar*, are very pungent perfumes.

ACIDUM ARSENIOSUM.

WHITE ARSENIC.

Arsenious acid is found in the shops, in the form of semi-transparent or opaque lumps, or in that of a heavy white powder. It has no smell, and its taste is slightly acrid. It is volatile at a red heat, giving off white fumes of a garlic-like odor. Extremely poisonous and corrosive, even in minute doses. Dose, $\frac{1}{30}$ th to $\frac{1}{12}$ th of a grain.
U. S. Dis.

ARSENICAL POWDER.

R. White arsenic, one part.
Calomel, one hundred and ninety-nine parts.

Mix. Recommended in obstinate cases of lepra and cancerous ulcers. *Dupuytren.*

COMPOUND POWDER OF WHITE ARSENIC.

R. White arsenic, six drachms.
Powdered dragon's blood, two drachms.
Animal charcoal, four scruples.
Cinnabar, three ounces.

Mix, and triturate well. Has been used in the same description of cases. *Van Mons.*

R. White arsenic, one part.
Kino, eight parts.
Cinnabar, sixteen parts.

Mix, and triturate well. *Ratier.*
Employed in the French hospitals as a caustic to cancerous sores.

ARSENICAL OINTMENT.

R. White arsenic, one drachm.
Lard,
Spermaceti cerate, each, six drachms.

Melt the cerate and lard by a gentle heat, and add the arsenic; triturate well in a glass mortar till perfectly united. *Soubeiran.*

ARSENICAL CERATE.

R. Powdered white arsenic, one scruple.
Simple cerate, one ounce.

Mix the arsenic with the cerate softened by heat, and triturate well.

SOLUTION OF ARSENIC.

R. White arsenic, one scruple.
Distilled water, two fl. ounces.

Dissolve. This and the foregoing ointments have been employed as applications to cancerous sores, but are always dangerous from the poisonous symptoms induced by the absorption of the arsenic.

ARSENICAL PILLS.

R. White arsenic, two grains.
Powdered opium, three grains.
White soap, eight grains.
Powdered liquorice root, q. s.

Mix well and divide into twenty pills. Each of these pills contains $\frac{1}{10}$ th of a grain of arsenic. They have been given in doses of one, or two, three times a-day, according to circumstances, in diseases of an intermittent character. *Ellis.*

ARSENIC AND PEPPER PILLS.

R. White arsenic, one grain.
Black pepper, twelve grains.

Gum Arabic, two grains.
Distilled water, q. s.

Triturate the arsenic and pepper for a long time, add the gum and water, and rub well together. Make twelve pills; each of which will contain $\frac{1}{12}$ th of a grain of arsenic.

Guibourt.

R. White arsenic, fifty-five grains.
Powdered black pepper, nine drachms.
Conserve of roses, sufficient.

Mix, and make eight hundred pills. One daily in chronic psoriasis. *Cazenave and Schedel.*

PILLS OF ARSENIC AND OPIUM.

R. Arsenious acid, two grains.
Powdered opium, eight grains.
Soap, one scruple.

Beat together and divide into twenty-four pills. One to be taken three times a-day. Have been found useful in intermittent fevers, periodical headaches, neuralgia, and lepra.

A. T. Thomson.

ACIDUM BENZOICUM.

BENZOIC ACID.

R. Benzoin, a pound.

Put it into a suitable vessel on a sand-bath, gradually increase the heat, and sublime as long as anything ascends. Wrap the sublimate in bibulous paper, and press, to separate the oil, and again sublime. *U. S. Ph.*

Benzoic acid is in the form of white, feathery hexagonal crystals, when pure having no odor, but usually with an agreeable smell from the presence of oil. The taste is acid and acrid. It is converted into hippuric acid, and voided by the urine when taken by man, and has been recommended in the uric acid diathesis and gout, but does not appear to be of any benefit. It has, however, been serviceable in catarrh of the bladder, and where there is a secretion of granular mucus with phosphates. It is also beneficial as a stimulating expectorant. Dose, ten to fifteen grains.

MIXTURE OF BENZOIC ACID
AND COPAIBA.

R. Benzoic acid, one drachm.
Copaiba, half fl. ounce.
White of egg, q. s.
Camphor water, seven fl. ounces.

Mix. Dose, two tablespoonfuls three times a-day. Recommended in chronic dysuria and vesical irritation. *Walker.*

POWDER OF BENZOIC ACID AND IPECACUANHA.

R. Benzoic acid, two scruples.
Ipecacuanha, six grains.

Golden sulphur of
antimony, three grains.
White sugar, one drachm.

Mix, and triturate well. Divide into six powders. Dose, one, four times a-day in barley water. Said to be useful in asthenic pneumonia. *Phæbus.*

MIXTURE OF BENZOIC ACID.

R. Benzoic acid, one scruple.
Sulphur, each, six grains.
Ipecacuanha, six ounces.
Honey, six ounces.
Syrup of seneka, one ounce.
Syrup of squills, each, one ounce.

Mix. Dose, a teaspoonful, three or four times a-day, in the chronic catarrh of elderly persons. *St. Marie.*

ACIDUM BORACICUM.

BORACIC ACID.

This is obtained from the native springs in Tuscany; or it may be procured from borax. Dose, ten grains to a drachm.

R. Powdered borax, forty parts.
Boiling water, one hundred parts.

Dissolve and add

Muriatic acid, twenty-five parts.

Collect the acid, which crystallizes on cooling, on a filter, drain, wash with cold water, and dry at 234° F. If not pure, dissolve and recrystallize. *Wackenroder.*

If sulphuric acid be used, as is generally advised, it is almost impossible to free the boracic acid from a trace of it.

MIXTURE OF BORACIC ACID.

R. Boracic acid, ten grains.
Oil of sweet almonds, one fl. drachm.
Syrup of lemons, each, one fl. drachm.

Mix. Advised in cardialgia; to be repeated until a cessation of the pain. *Pierquin.*

COLLUTORY WITH BORACIC ACID.

R. Cloves, } each,
Galangal, } two drachms.
Ginger, }
Peruvian bark, }
Gum lac, one and a half drachms.
Benzoin, }
Storax, each, half a drachm.
Alcohol, six fl. drachms.
Acetic acid, one and a half fl. ounces.
Vinegar, five fl. ounces.

Macerate, press, and filter, then add

Boracic acid, half a drachm.

This has been recommended as a mouth wash in a scorbutic condition of the gums. When used it is to be diluted with water.

Phæbus.

ACIDUM CARBONICUM.

CARBONIC ACID.

This stimulating gas is readily absorbed by water, and advantage has been taken of the fact by retaining it in a liquid form under pressure. The mineral or soda water of the shops is water saturated with carbonic acid gas.

CARBONIC ACID WATER.

By means of a forcing-pump, throw into a suitable receiver, nearly filled with water, a quantity of carbonic acid equal to five times the bulk of the water. *U. S. Ph.*

Carbonic acid is readily obtained from marble by means of dilute sulphuric acid.

This water has a sharp, pungent, and agreeable taste. It is much used as a common drink in doses of half a pint.

ACIDUM CITRICUM.

CITRIC ACID.

This exists naturally in the juices of many fruits; to obtain it pure, boiling lemon or lime juice is to be saturated with chalk, when carbonic acid is given off, and an insoluble citrate is formed; this is to be washed and decomposed by boiling with dilute sulphuric acid; insoluble sulphate of lime precipitates, and the citric acid remains in solution and crystallizes on evaporation. To purify, dissolve in water and recrystallize. *U. S. Dispens.*

Dose, five to twenty grains.

ARTIFICIAL LEMON JUICE.

R. Citric acid, one ounce.
Distilled water, fourteen fl. ounces.
Oil of lemons, five drops.

Mix. *Beasley.*

LOZENGES OF CITRIC ACID.

R. Citric acid, three drachms.
White sugar, one pound.
Oil of lemons, sixteen drops.

Triturate well, and add

Mucilage of tragacanth, q s.

Make lozenges of twelve grains each.

Cottureau.

SYRUP OF CITRIC ACID.

R. Citric acid, two drachms.
Oil of lemons, four minims.
Syrup, two pints

Mix the citric acid and oil of lemons with an ounce of the syrup, then add rest of syrup, and dissolve by a gentle heat. *U. S. Ph.*

SYRUP OF CITRIC ACID.

R. Citric acid, five drachms.
Water, ten fl. drachms.
Simple syrup, two pounds.
Tincture of fresh lemon-peel, one fl. drachm.

Dissolve the acid in the water, mix with the syrup at a boiling heat, and, when cold, add the tincture. *Soubéiran.*

DRY LEMONADE.

R. Citric acid, one or two drachms.
White sugar, four ounces.
Oil of lemons, eight drops.

Mix well. A spoonful to a tumbler of water. *Gray.*

CITRATED EFFERVESCING POWDERS.

R. Citric acid, nine drachms.
Divide into eighteen powders.

R. Bicarbonate of soda, eleven drachms.
or,

Bicarbonate of potash, thirteen dr'sms.
Divide into eighteen powders. *Dub. Ph.*

An acid and an alkaline powder are dissolved in separate portions of water, the two solutions are mixed and drank in a state of effervescence. An excellent refrigerant.

ACIDUM GALLICUM.

GALLIC ACID.

It is not certain that this acid exists as such in nature, or whether it arises from the decomposition of tannic acid. It is usually prepared from galls. Many processes have been devised.

R. Decoction of galls, at will.
Expose to action of air in a loosely covered vessel for some months; it will grow mouldy, and become covered with a glutinous pellicle, and gallic acid will be deposited on the sides of the vessel and on the under surface of the pellicle; collect, dissolve, and recrystallize. *Scheele.*

This is analogous to the process official in the U. S. Ph.

R. Decoction of galls, at will.
Sulphuric acid, sufficient

to precipitate. Wash the precipitate with and dissolve by aid of heat in diluted sulphuric acid, boil for a few minutes, let cool, and collect the crystals. *Liebig.*

Gallic acid is a powerful astringent, and has been found useful in hemorrhages and fluxes, as well as in checking night sweats in phthisis.

Dose five to ten grains.

PILLS OF GALLIC ACID.

R. Gallic acid, q. s.
Extract gentian, sufficient
to form pills of two to five grains each. One to be given every three or four hours.
Useful in menorrhagia, hematuria, &c. *Dunghison.*

INJECTION OF GALLIC ACID.

R. Gallic acid, one scruple to one drachm.
Water, two pints.
Mix. Found beneficial in leucorrhœa. *Dunghison.*

ACIDUM HYDROCYANICUM DILUTUM.

HYDROCYANIC ACID.

R. Ferrocyanuret of potassium, two ounces.
Sulphuric acid, one ounce and a half.
Distilled water, sufficient.

Mix the acid with four fl. ounces of the water and pour the mixture, when cool, into a glass retort. To this add the ferrocyanuret, previously dissolved in ten fl. ounces of the water. Pour eight fl. ounces of distilled water into a cooled receiver, and having attached the retort, distil on a sand-bath, with a moderate heat, six fl. ounces. Lastly, add to the product, five fl. ounces of distilled water, or as much as will render the acid of such strength, that 100 grains will be accurately saturated by 12.7 grains of nitrate of silver. *U. S. Ph.*

EXTEMPORANEOUS HYDROCYANIC ACID.

R. Cyanuret of silver, fifty grains and a half.
Muriatic acid, forty-one grains.
Distilled water, one fl. ounce.

Mix the acid with the water, and add the cyanuret, and shake in a well-stopped bottle. Let settle, decant, and keep for use. *U. S. Ph.*

This acid should be kept in closely-stopped bottles, protected from the light. It is a transparent, volatile liquid, of a cooling and then somewhat irritating taste, and a peculiar smell. It is the most active poison known, and must be used with extreme caution. It is used as an anodyne and antispasmodic, in many diseases. The dose is from one to two drops mixed with gum water or syrup, always beginning with the smallest quantity and gradually increasing.

MIXTURE OF HYDROCYANIC ACID.

R. Medicinal hydrocyanic acid, one fl. drachm.
Distilled water, one pint.
Sugar, one ounce and a-half.
Mix. A dessertspoonful twice a-day, gradually increasing the dose. *Magendie.*

R. Powdered gum Arabic, half an ounce.
Water, seven fl. ounces and a-half.

Dissolve, and add

Syrup of tolu, half fl. ounce.
Diluted hydrocyanic acid, twelve drops.

Mix. A tablespoonful every three hours in the cough of phthisis. *S. G. Morton.*

R. Medicinal hydrocyanic acid, half a drachm.

Powdered sugar-candy, one ounce and a half.

Syrup of red cabbage,

" Mallow, each, two ounces.

" Balsam of tolu,

" Maidenhair, each, one ounce.

" Poppies,

" Cinnamon, each, two drachms.

Mix. A dessertspoonful occasionally. *Pierquin.*

SYRUP OF HYDROCYANIC ACID.

R. Medicinal hydrocyanic acid, four grains and a half.
Clarified syrup, one ounce.

Mix. This contains $\frac{9}{17}$ of a grain of anhydrous acid. *Magendie.*

JULEP OF HYDROCYANIC ACID.

R. Medicinal hydrocyanic acid, fifteen drops.

Hoffman's anodyne, two fl. ounces.

Syrup of marsh mallow, three fl. ounces.

Mix. A dessertspoonful every two hours. *Pierquin.*

R. Medicinal hydrocyanic acid, two to four drops.

Syrup of peppermint, one fl. ounce.

Infusion of linden, four fl. ounces.

Mix. A dessertspoonful every hour. *Foy.*

INJECTION OF HYDROCYANIC ACID.

R. Medicinal hydrocyanic acid, one part.
Distilled water, four parts.

In gonorrhœa. *Foy.*

LOTION OF HYDROCYANIC ACID.

R. Diluted hydrocyanic acid, half fl. ounce.

Alcohol, one fl. ounce.

Distilled water, ten fl. ounces and a half.

Mix. As lotion, in impetigo. *A. T. Thomson.*

R. Medicinal hydrocyanic acid, two fl. drachms.

Lettuce water, two pints.

Mix. In hepatic affections. *Magendie.*

R. Diluted hydrocyanic acid, half fl. drachm.

Bicarbonate of soda, two drachms.

Milk, eight fl. ounces.

Mix. In milky scall. *A. T. Thomson.*

ACIDUM HYDRIODICUM.

HYDRIODIC ACID.

R. Tartaric acid, two hundred and sixty-four grains.

Iodide of potassium, three hundred and thirty grains.

Dissolve each in one fl. ounce and a half of distilled water. Mix the solutions, shake and let settle, filter, and add distilled water to make up measure of six and a quarter fl. ounces.

Buchanan.

Has the same properties as iodine, of which each fl. drachm contains five grains; dose, at first a few drops, gradually increased to half a fl. ounce three times a-day.

ACIDUM HYDROSULPHURICUM.

SULPHURETTED HYDROGEN.

R. Sulphuret of iron,
Sulphuric acid, equal parts.

Pulverize the sulphuret, and gradually pour on it the acid diluted with three times its weight of water, and collect the gas; or it may be passed through water to saturation. *Van Mons.*

This gas is an active poison, but has been administered in colica pictonum and mercurial affections. It is, however, more employed externally in cutaneous diseases.

HYDROSULPHURETTED BATH.

R. Sulphuret of potassium, four ounces.
Water, one pint.

Dissolve, and add
Muriatic acid, two drachms.

Pour the whole into the bath. Advised in chronic diseases of the skin, rheumatism, and certain cases of paralysis.

Cadet de Gassicourt.

ARTIFICIAL SULPHURETTED WATER

R. Carbonate of soda, twelve grains.

Liquid hydrosulphuric acid, two pints.

Water, six pints

To be kept in well-closed bottles. Has been recommended in colica pictonum. *Foy.*

HYDROSULPHURETTED LOTION.

R. Sulphuret of
potassium, twenty-four parts.
Water, two hundred and fifty parts.

Dissolve, and add

Sulphuric acid, one part.

Used as a wash in chronic diseases of the skin.
Dupuytren.

ACIDUM LACTICUM.

LACTIC ACID.

R. Sugar, six pounds.
Tartaric acid, half an ounce.
Boiling water, twenty-six pounds.

Mix, set aside for a few days, then add

Old stinking cheese, eight ounces,
well diffused in

Curdled acid skimmed
milk, eight pounds.
Powdered chalk, three pounds.

Place in a warm situation, so as to keep the mixture at a temperature of 86° to 95°. Stir often; in eight or ten days it will solidify into a stiff paste of lactate of lime; now add twenty pounds of boiling water, and half an ounce of caustic lime, boil for half an hour, and filter through linen. Evaporate the liquid to the consistence of syrup, and set aside for four days; remove the lactate of lime, express, agitate with one-tenth of cold water, and express, repeating this operation two or three times. Now dissolve in twice its weight of boiling water, and for every pound of the lactate, add three and a-half ounces of sulphuric acid diluted with an equal weight of water. Filter the hot liquid through a bag, and boil it with one pound and three-eighths of carbonate of zinc, for every pound of sulphuric acid, for a quarter of an hour. Filter whilst boiling hot, and let stand to crystallize; remove these crystalline crusts of lactate of zinc, and wash them with cold water. Then dissolve them in seven and a half parts of boiling water, and pass through the solution a current of sulphuretted hydrogen, till sulphuret of zinc no longer separates. Filter, boil the liquid, to expel the excess of sulphuretted hydrogen, and evaporate on a water-bath to the consistence of syrup.

Bensch.

The juice of beets is to be permitted to ferment for two months in a warm place, then evaporate to the consistence of syrup, treat with alcohol, evaporate, dissolve the residuum in water, saturate, filter, concentrate, and crystallize. Dissolve the crystals in water, purify by boiling with animal charcoal, filter whilst hot, treat with baryta and then with sulphuric acid; the lactic acid will be set free, and may be concentrated in vacuo.

Cotteneau.

Lactic acid is in the form of a colorless syrup, inodorous, but of a very sour taste. It attracts

moisture from the air. It has been recommended by Magendie in dyspepsia, and may, perhaps, be useful in a phosphatic diathesis. Dose, five grains.

LOZENGES OF LACTIC ACID.

R. Lactic acid, two fl. drachms.
Powdered white sugar, one ounce.
Gum tragacanth, q. s.
Oil of vanilla, four drops.

Mix, and form pastilles of half a drachm each, of which three to six may be taken during the day. They should be kept dry. *Magendie.*

LEMONADE OF LACTIC ACID.

R. Lactic acid, one to four fl. drachms.
Simple syrup, two fl. ounces.
Water, two pints.

Mix. Dose, a cupful several times a-day.

Magendie.

ACIDUM MURIATICUM.

MURIATIC ACID.

Prepared by distilling common salt in a glass retort with sulphuric acid and water. It has a density of 1.16, a suffocating odor, an acrid and sour taste. It is very volatile. It gives a curdy-white precipitate with nitrate of silver. It is never given internally except in a diluted form.

DILUTED MURIATIC ACID.

(*Sp. Gr.* 1.046.)

R. Muriatic acid, four fl. ounces.
Distilled water, twelve fl. ounces.

Mix. Dose, twenty drops in sweetened water.
U. S. Ph.

GASEOUS MURIATIC ACID.

R. Common salt, two parts.
Sulphuric acid, three parts.

Mix. The muriatic acid fumes that arise are considered to be disinfectant, but are much inferior to chlorine. *Swedjaur.*

MURIATIC ACID BATH.

R. Muriatic acid, ten fl. ounces.
Water, fifty gallons.

Mix. Found useful in some chronic diseases of the skin. *Soubeyran.*

MURIATIC ACID GARGLE.

R. Muriatic acid, two fl. drachms.
Honey, two fl. ounces.
Barley water, one pint.

Mix. Useful in the angina of scarlet fever, and in ulceration of the mouth and throat.

Ratier.

MURIATIC ACID LOTION.

- R. Muriatic acid, one part.
Water, sixteen parts.

Mix. Advised as a wash for chilblains, and also found beneficial in lepra and other skin diseases. *Foy.*

MURIATIC ACID PEDILUVIUM.

- R. Muriatic acid, two fl. ounces.
Water, eight pints.

Mix. Advised by Scott and others in chronic enlargements of the liver and spleen. *Beral.*

MURIATIC ACID DRAUGHT.

- R. Muriatic acid, ten to twenty drops.
Barley water, eight ounces.

Mix. Dose, a tablespoonful three or four times a day. Has been advised in stone in the bladder. *Ellis.*

MURIATIC ACID LINIMENT.

- R. Balsam Peru, one drachm.
Spermaceti,
White wax,
Muriatic acid, each, two drachms.
Olive oil, two fl. ounces.
Water, six fl. ounces.

Rub well together. To be applied twice a day. *Ratier.*

ACIDUM NITRICUM.

NITRIC ACID.

Nitric acid of the specific gravity 1.42.

DILUTED NITRIC ACID.

- R. Nitric acid,
(*sp. gr.* 1.42), one fl. ounce.
Distilled water, six fl. ounces.

Mix. Dose, twenty to forty drops, in sweetened water, three times a day. *U. S. Ph.*

The *sp. gr.* of diluted nitric acid is 1.07.

COLLUTORY OF NITRIC ACID.

- R. Nitric acid, one fl. scruple.
Honey of roses,
Syrup of mulberries, each, half fl. ounce.

Mix. To touch obstinate venereal ulcers in the mouth. *Phæbus.*

FOMENTATION OF NITRIC ACID.

- R. Nitric acid, one part.
Distilled water, ninety-six parts.

Mix. Used to destroy the fetid smell of foul ulcers; also as a wash in itch. *Soubeiran.*

LOTION OF NITRIC ACID.

- R. Nitric acid, half to one fl. drachm.
Laudanum, one fl. drachm.
Rose water, six fl. ounces.

Mix. To wash venereal ulcers. *Phæbus.*

NITRIC ACID MIXTURE.

- R. Nitric acid, two fl. drachms.
Raspberry syrup, three fl. ounces.
Sugar, three ounces.
Water, two pints.

A wineglassful, with as much Seltzer water, in fetid breath. *Cadet.*

- R. Nitric acid, half fl. drachm.
Water, one pint and a half.
Syrup of cinnamon, one fl. ounce and a half.

Mix. In syphilis, chronic hepatitis, &c. A spoonful every two or three hours. *Augustin.*

- R. Nitric acid, two fl. scruples.
Opium, two grains.
Water, two fl. ounces.
Syrup of cinnamon, half fl. ounce.

Mix. A spoonful every hour, in barley water. In dysentery, cholera, &c. *Ammon.*

- R. Nitric acid, half fl. drachm.
Gum Arabic,
Sugar, each, three drachms.
Water, eight fl. ounces.

Mix. A small tablespoonful in water, as occasion may require. As a tonic, to arrest colliquative sweats, &c. *Ellis.*

OINTMENT OF NITRIC ACID.

- R. Olive oil, one pound.
Lard, four ounces.
Nitric acid, five fl. drachms and a half.

Melt the oil and lard together in a glass vessel; when, on cooling, they begin to stiffen, add the acid, and stir. Used as an application to porrigo, psora, &c. *Dub. Ph., 1826.*

LINIMENT OF NITRIC ACID.

- R. Nitric acid, two fl. ounces.
Oil of turpentine, three fl. ounces.
Theriac, three drachms.
Honey, one ounce.
Alcohol, six fl. ounces.

Mix. As a rubefacient, has been used in cholera. *Foy.*

ACIDUM NITRO-MURIAT-
TICUM.

NITRO-MURIATIC ACID.

R. Nitric acid, by measure, one part.
Muriatic acid, " two parts.
Mix in a refrigerated bottle, and keep in a cool,
dark place. *Dub. Ph.*
Dose, three or four drops.

NITRO-MURIATIC ACID BATH.

R. Nitro-muriatic acid, six fl. ounces.
Water, three gallons.
Mix. Recommended by Scott as a foot-bath in
chronic hepatitis.

ACIDUM OXALICUM.

OXALIC ACID.

This acid exists in many plants in combina-
tion with lime or potash, but is usually obtained
by the action of nitric acid on sugar or starch.
It is in the form of colorless, transparent, pris-
matic crystals, having no odor, but a very acid
taste. This acid is an active poison in large
doses. It is not used in medicine in this coun-
try, but is employed in France.

LOZENGES OF OXALIC ACID.

R. Powdered oxalic acid, one drachm.
White sugar, eight ounces.
Gum tragacanth, two scruples.
Water of orange-peel, five fl. drachms.
Oil of lemons, eight drops.
Mix, and form lozenges of ten grains each.
Cottureau.

ACIDUM PHOSPHORICUM.

PHOSPHORIC ACID.

R. Bones, calcined to
whiteness, five parts.
Sulphuric acid, three parts.
Water, thirty parts.

Mix, and boil for an hour, constantly stirring,
strain, evaporate the fluid to consistence of
honey, pour on an oiled slab, and when cold
break in pieces. *Van Mons.*

This affords not pure phosphoric acid, but an
acid phosphate of lime, yet is the formula re-
cognized by the Batavian, Russian, Prussian,
Saxon, &c., Pharmacopœias.

R. Phosphorus, one part.
Water, two parts.
Nitric acid, eight parts.

Cut the phosphorus in small pieces, and add it
very gradually to the acid, placed on a sand-

bath; when all the phosphorus has been changed
into an acid, evaporate to the consistence of
syrup. *Guibourt.*

DILUTED PHOSPHORIC ACID.

R. Phosphorus, six drachms.
Nitric acid,
(*sp. gr.* 1.42), four fl. ounces.
Distilled water, eight fl. ounces.

Add the phosphorus gradually to the nitric acid
and water in a retort, on a sand-bath, apply
heat, and distil six fl. ounces. Pour these back
into the retort and again distil six fl. ounces,
which reject. Evaporate the residue in a pla-
tinum crucible to two ounces. When cold, add
as much distilled water as will make a pint
(*Imp.*) and mix. Its *sp. gr.* is 1.064.

London Ph.

This acid is stated to be useful in nervous dis-
orders, in lithiasis with phosphatic deposits, in
morbid ossifications, in diabetes, to allay thirst,
&c., and as a local application in caries. The
dose of the dilute acid is from ten drops to a fl.
drachm in gum water.

PILLS OF PHOSPHORIC ACID.

R. Diluted phosphoric acid,
Assafoetida, each, three drachms.
Powdered calamus, q. s.
Make one hundred and eighty pills. Dose, five
to ten, three times a-day.

Recommended in caries.

Phæbus.

FOMENTATION OF PHOSPHORIC ACID.

R. Diluted phosphoric acid, one ounce.
Decoction of chamomile, eight ounces.
Mix. Employed in caries, and as an injection in
fistulas. *Augustin.*

PHOSPHORIC TINCTURE OF MYRRH.

R. Tincture of myrrh, half a fl. ounce.
Diluted phosphoric acid, thirty drops.
Mix. Used in caries of the teeth or bones.
Augustin.

PHOSPHORIC ACID LEMONADE.

R. Diluted phosphoric acid, twenty-four
drops.
Simple syrup, two ounces.
Water, two pints.
Mix. Dose, about an ounce, to relieve thirst in
diabetes. *Ratier.*

ACIDUM SULPHURICUM.

SULPHURIC ACID.

DILUTE SULPHURIC ACID (*Sp. Gr.* 1.09).

R. Sulphuric acid, one fl. ounce.
Water, thirteen fl. ounces.

Add the acid gradually to the water in a glass vessel, and mix them.

U. S. Ph.

Dose, ten to twenty drops.

ELIXIR OF VITRIOL.

R. Sulphuric acid, three and a half fl. ounces.
Ginger, bruised, one ounce.
Cinnamon, bruised, one ounce and a half.
Alcohol, sufficient.

Gradually add the acid to a pint of alcohol. Place the ginger and cinnamon in a percolator, and pour alcohol upon them until a pint of tincture is obtained. Then mix the diluted acid and the tincture. Dose, ten to twenty drops in sweetened water.

U. S. Ph.

SULPHURIC ACID AND ALCOHOL.

R. Sulphuric acid, four ounces.
Acetic acid,
Alcohol, each, two pounds.

Mix. Vulnerary, antiseptic, and astringent. Dose, internally, twenty or thirty drops, in an appropriate vehicle. Externally, as a lotion to foul ulcers and contusions, and to arrest bleeding.

Campana.

SULPHURIC ACID AND NITRIC ETHER.

R. Sulphuric acid,
Nitric ether, equal parts.

Mix very gradually. Dose, five to ten drops in an ounce of water. Found beneficial in spasmodic attacks, sinking, and hemorrhages.

Vogler.

SULPHURIC ACID LEMONADE.

R. Sulphuric acid, one part.
Simple syrup, thirty-two parts.
Water, one hundred and sixty parts.

Mix. As a cooling drink in hemorrhages, and also found useful as a preventive of the effects of lead.

Beral.

SULPHURIC ACID LINIMENT.

R. Sulphuric acid, two drachms.
Olive oil, two and a half ounces.
Oil of turpentine, one ounce.

Mix. In gentle frictions to chilblains, where the skin is not broken.

Foy.

SULPHURIC ACID OINTMENT.

R. Olive oil, eight parts.
Add gradually, constantly stirring,
Sulphuric acid, five parts.

After standing for twenty-four hours, wash well in tepid water, till it will not redden litmus paper. Has been used in itch and paralysis, in frictions.

Brugnatelli.

ACIDUM TANNICUM.

TANNIC ACID.

R. Powdered galls,
Sulphuric ether, of each a sufficient quantity.

Put into a glass adapter, loosely closed at its lower end with carded cotton, sufficient galls to fill about one-half of it, pressing the powder slightly. Then fit the adapter accurately to a receiver, fill it with sulphuric ether, previously washed with water, and close the upper end slightly with a cork. The liquid which passes into the receiver separates into two portions, the lower being the smaller and denser. Pour on additional ether until the lower stratum of liquid in the receiver no longer increases. Separate this, and evaporate to dryness in a capsule by a moderate heat, and reduce to powder.

U. S. Ph.

Tannic acid is of a yellowish-white color, of a powerfully astringent taste, soluble in water, less so in alcohol or ether. It is a strong astringent, and has been successfully used in hemorrhages and other fluxes. The dose is from two to ten grains.

PILLS OF TANNIC ACID.

R. Tannic acid, six grains.
Gum Arabic, twelve grains.
White sugar, seventy-two grains.

Triturate well, and add

Simple syrup, sufficient to make pills of four grains each. Dose, one to four, morning and evening.

Beral.

R. Tannic acid, eight or twelve grains.
Mucilage, sufficient

to make eight pills. One to be given every two or three hours, in gonorrhœa.

Ellis.

PILLS OF TANNIC ACID AND OPIUM.

R. Tannic acid, half a drachm.
Extract of

opium, three-quarters of a grain.
Conserve of roses, sufficient

to make twenty pills. One every hour, in uterine hemorrhage.

Dumars.

GARGLE OF TANNIC ACID.

R. Tannic acid, one drachm.
Honey of roses, two ounces.
Rose water, two fl. ounces.
Distilled water, eight fl. ounces.

Mix. As a gargle, to arrest mercurial salivation.

Beral.

INJECTION OF TANNIC ACID.

R. Tannic acid, half a drachm.
Distilled water, eight fl. ounces

Dissolve. In gleet and leucorrhœa.

Beral.

MIXTURE OF TANNIC ACID.

- R. Tannic acid, twelve grains.
 Syrup of rhatany,
 Syrup of gum, each, one fl. ounce.
 Camphor water, four fl. ounces.

Mix. Six to twelve spoonfuls a-day, in diarrhoea. *Beral.*

OINTMENT OF TANNIC ACID.

- R. Tannic acid,
 Distilled water, each, two drachms.
 Dissolve and rub well with
 Lard, twelve drachms.

ACIDUM TARTARICUM.

TARTARIC ACID.

Is prepared from bi-tartrate of potash, by saturating the excess of acid with chalk, and decomposing the insoluble tartrate of lime by means of dilute sulphuric acid, evaporating and crystallizing.

It is cooling and diuretic in doses of a drachm or more.

SYRUP OF TARTARIC ACID.

- R. Tartaric acid, ten parts.
 Distilled water, twenty parts.

Dissolve and mix with

Boiling syrup, five hundred parts.

Codex.

LOZENGES OF TARTARIC ACID.

- R. Powdered tartaric acid, three drachms.
 White sugar, one pound.
 Oil of lemons, sixteen drops.
 Mucilage of tragacanth, sufficient.

Make into lozenges of twelve grains.

Used to allay thirst. *Cottureau.*

LEMONADE OF TARTARIC ACID.

- R. Tartaric acid, one drachm.
 White sugar, one ounce.
 Water, twelve fl. ounces.

Dissolve. For an agreeable acidulous drink.

Foy.

EFFERVESCING POWDERS.

- R. Tartaric acid, one ounce.

Divide into sixteen powders.

- R. Bicarbonate of soda, one ounce and fifty-four grains.

or,

Bicarbonate of potash, one ounce and one hundred and sixty grains.

Divide into sixteen powders.

An acid and an alkaline powder are separately dissolved in water, and the mixed solutions are taken in a state of effervescence.

Ed. Ph.

ACIDUM VALERIANICUM.

VALERIANIC ACID.

- R. Valerian, forty pounds.
 Distilled water, three hundred and twenty pints.

Distil, collect the oil, continuing the distillation as long as the water is acid. Agitate the oil with milk of lime; nearly saturate the acid-water by milk of lime, first adding that already used, and then fresh; and, lastly, add lime-water to excess. Concentrate till a pellicle appears, then decompose by nitric acid in a long, narrow flask with a ground stopper, decant the valerianic acid, which floats on the liquid, and distil it with a gentle heat till the distilled fluid ceases to be oily.

P. L. Bonaparte.

- R. Valerian, at will.

Boil the root for three or four hours with rather more than its bulk of water, in which an ounce of carbonate of soda is dissolved for every pound of the root, replacing the water as it evaporates. Express strongly; boil the residue twice with the same quantity of water, expressing each time. Mix the liquids, add two fl. drachms of strong sulphuric acid for every pound of the root, and distil till three-fourths of the liquid have passed over. Neutralize this by carbonate of soda, concentrate, decompose the valerianate of soda by sulphuric acid, and separate the free valerianic acid by distillation.

T. and H. Smith.

Valerianic acid may also be obtained from the valerianate of soda, which is now official in the Dublin Pharmacopœia. (1854.)

ACONITUM.

ACONITE. MONKSHOOD.

Many species of Aconitum are official in the various pharmacopœias, but the only one recognized by the U. S. Ph. is the *A. napellus*, a native of Europe.

Sext. Syst. Polyand. Trigyn. *Nat. Syst.* Ranunculacæ.

Linn. Sp. Pl. 751. *Griffith, Med. Bot.* 90.

The whole plant is possessed of highly deleterious properties, but the root is the most powerful. The leaves are also made use of. They have a somewhat nauseous odor, and a bitterish, acrid taste, followed by a peculiar tingling and burning of the lips. They owe their properties to the presence of a peculiar principle called *Aconitia*.

COMPOUND POWDER OF ACONITE.

- R. Powdered aconite leaves,
Precip. sulph. of
antimon., each, one grain.
Carbonate of magnesia, one scruple.
Mix. As an anodyne and diaphoretic in gout
and rheumatism. *Vogler.*

EXTRACT OF ACONITE.

- R. Recent leaves of aconite, one pound.
Moisten with water, bruise in a stone mortar,
express the juice, and evaporate to the proper
consistence. *U. S. Ph.*

- R. Recent leaves of aconite, a sufficient
quantity.

Beat to a pulp, express the juice; subject the
residuum to displacement with rectified spirit,
as long as the liquid is colored; unite the juice
and the tincture; filter; distil off the spirit, and
evaporate to a proper consistence on a vapor-
bath. *E. Ph.*

Both are uncertain preparations; when good,
they should cause a numbness and tingling of
the lips, in a few minutes after taking. Dose,
one to two grains, to be gradually increased.

ALCOHOLIC EXTRACT OF ACONITE.

- R. Aconite leaves, one pound.
Diluted alcohol, four pints.
Make a tincture by displacement; distil off
the alcohol, and evaporate. *U. S. Ph.*

Dose, one-sixth to one-half of a grain.

PILLS OF EXTRACT OF ACONITE.

- R. Alcohol. extract of aconite, one grain.
Powdered liquorice, twelve grains.
Syrup, sufficient
to form a consistent mass. Divide into six
pills. Dose, one every three or four hours.
Turnbull.

ACONITE PLASTER.

- R. A small portion of the alcoholic ex-
tract spread over the surface of
common adhesive plaster.

This has been highly recommended in neu-
ralgia, over the painful part. *Curtis.*

- R. Coarsely powdered aconite
root, four ounces.
Alcohol, sp. gr. .835, sufficient.
Adhesive plaster, three ounces and a
half.

Moisten the aconite root with six ounces of
alcohol, and permit it to macerate twenty-four
hours; then put it in a small displacer, and

when properly packed, pour on gradually suffi-
cient alcohol to make a pint of tincture. Distil
off three-fourths of the alcohol, evaporate the
residue on a water-bath to a thick, syrupy con-
sistence, then add the plaster previously liqui-
fied, and stir constantly, until it is properly
incorporated with the soft resinous extract and
cools.

This plaster should be spread in a thin stratum
on skin or oiled silk. *W. Procter, Jr.*

ACONITE OINTMENT.

- R. Alcohol. extract of aconite, one part.
Lard, two parts.
Recommended as a friction in neuralgia.
Turnbull.

COMPOUND WINE OF ACONITE.

- R. Alcohol. extract of
aconite, one drachm.
Antimonial wine, one ounce.

Make a solution. Dose, fifteen or twenty drops
every three hours, gradually increasing till
some effect is produced.

Found beneficial in chronic rheumatism,
toothache, neuralgia, &c. *Richter.*

TINCTURE OF ACONITE LEAVES.

- R. Aconite leaves, four ounces.
Diluted alcohol, two pints.

Macerate for fourteen days, and filter; or pre-
pare by displacement. Dose, ten to twenty
drops three times a-day. The effects should
be carefully watched. In rheumatism, neu-
ralgia, &c. It is also employed as an embro-
cation to the painful spots, by means of a small
piece of sponge attached to a handle.

U. S. Ph.

TINCTURE OF ACONITE ROOT.

- R. Bruised aconite root, one pound.
Alcohol, two pints.

Macerate for fourteen days, express, and filter.
Or it may be prepared by displacement. Dose,
five to eight drops. *U. S. Ph.*

LINIMENT OF ACONITE ROOT.

- R. Powdered aconite root, four ounces.
Glycerin, two fl. drachms.
Alcohol, sufficient.

Macerate the aconite with half-a-pint of alcohol
for twenty-four hours, then pack it in a small
displacer, and add alcohol gradually, until a
pint of tincture has passed. Distil off twelve
fluid ounces, and evaporate the residue to twelve
fluid drachms. To this add two fluid drachms
of alcohol and the glycerin, and mix them.

W. Procter, Jr.

Brush over a piece of lint or muslin, of the size of the part to be treated, with the liniment, apply this to the surface, and cover it with a piece of oiled silk and a bandage.

AMMONIATED EXTRACT OF ACONITE.

R. Alcohol. extract of
aconite, one drachm.
Water of ammonia, ten drops.
Evaporate gently, to expel excess of ammonia.

AMMONIATED ACONITE OINTMENT.

R. Ammon. extract of
aconite, one drachm.
Lard, three drachms.
Rub well together.

Used for the same purposes as the aconite ointment, but more active and pungent.

Turnbull.

MIXTURE OF ACONITE.

R. Tincture of aconite
leaves, one fl. drachm.
Carbonate of soda, one drachm and
a half.
Sulphate of mag-
nesia, one ounce and a half.
Water, six fl. ounces.

Mix. In gastralgia. A tablespoonful, when the pain is urgent.

Fleming.

ACONITIA.

ACONITINE.

R. Aconite root, dried and
bruised, two pounds.
Alcohol, three gallons.
Diluted sulphuric
acid, }
Solution of am- } each, sufficient.
monia, }
Purified animal
charcoal, }

Boil the aconite with a gallon of the alcohol, for an hour, in a retort attached to a receiver. Pour off the tincture, and repeat the operation a second and third time. Then express, mix the tinctures, filter, and distil off the alcohol. Evaporate the residue by a water-bath to consistence of an extract. Dissolve this in water, and filter. Evaporate the solution with a gentle heat to consistence of syrup. Add to it the diluted acid mixed with distilled water. Drop in the solution of ammonia, and dissolve the precipitated aconitia in diluted sulphuric acid mixed with water. Then add the animal charcoal, occasionally shaking, for a quarter of an hour. Lastly, filter, and, having again dropped in the solution of ammonia, wash the precipitate and dry it.

U. S. Ph.

Not used internally, but successfully employed externally in neuralgia, gout, and rheumatism.

ACONITINE OINTMENT.

R. Aconitine, two grains.
Alcohol, six drops.
Rub well together, and add,
Lard, one drachm.
Turnbull.

R. Aconitine, sixteen grains.
Olive oil, half a drachm.
Lard, one ounce.
Incorporate thoroughly. Turnbull.

These ointments are employed as frictions in neuralgia, over the painful parts. A small portion, not exceeding the size of a pea, is to be used at a time. The operation to be repeated three or four times a-day.

ACONITINE LOTION.

R. Aconitine, eight grains.
Alcohol, two fl. ounces.
Dissolve. Used by means of a friction-sponge over the affected part. Never to be employed where the skin is broken or abraded.
Turnbull.

ADIANTUM.

MAIDENHAIR.

Two species of this genus, *A. pedatum* and *A. capillus veneris* have been much used in Europe as remedies in pectoral affections. They are both bitter and aromatic, but the former is the most active, and is a native of the United States.

Sex. Syst. Cryp. Fil. Nat. Syst. Polypodiaceæ.

INFUSION OF MAIDENHAIR.

R. Maidenhair, three drachms.
Licorice root, two drachms.
Water, sufficient
to obtain two pints of infusion. As a drink in catarrh.

SYRUP OF MAIDENHAIR.

R. Maidenhair, four ounces.
Boiling water, three pints.
Infuse for two hours, strain, and add
Sugar, four pounds.
Clarify with white of egg, and add
Maidenhair, two ounces.
Digest for two hours, and strain. Cottereau.

This syrup is known under the name of *syrup of capillaire*, and is much employed in Europe as a pectoral. Dose, a tablespoonful.

COMPOUND SYRUP OF MAIDENHAIR.

R. Maidenhair,	five ounces.
Marsh mallow,	two drachms.
Asparagus root,	one ounce.
Liquorice root,	two ounces.
Water,	two pints.

Boil to a pint and a half, express, strain, clarify and add

Sugar, thirty-two ounces.

Make syrup. *Wurtemberg Ph.*

Dose, a tablespoonful.

ÆTHER ACETICUS.

ACETIC ETHER.

R. Alcohol (sp. gr. .835),	three thousand parts.
Acetic acid (10°),	two thousand parts.
Sulphuric acid,	six hundred and twenty-five parts.

Mix the alcohol and acetic acid in a glass retort, add gradually the other acid, distil in a sand-bath four thousand parts, rectify this product with a small quantity of carbonate of potassa to obtain three thousand parts. *Codex.*

This ether is milder, more agreeable and diaphoretic, than the other ethers. It is used in low fevers, spasmodic vomiting, and cardialgia, in doses of ten drops to a drachm.

SPIRIT OF ACETIC ETHER.

R. Acetic ether,	one part.
Alcohol,	three parts.

Mix. This is employed as a substitute for Hoffmann's anodyne, in about the same doses. *Giordano.*

MIXTURE OF ACETIC ETHER.

R. Acetic ether,	
Ethereal tincture of valerian,	
Tincture of opium,	equal parts.

Mix. Dose, from ten to thirty drops, in hysteria and hypochondriasis. *Radius.*

ÆTHER HYPONITROSUS.

NITROUS ETHER.

R. Rectified spirit,	fifteen fl. ounces.
Pure nitric acid (1.500),	seven fl. ounces.

Put the spirit with a little clean sand into a two-pint matrass, fitted with a cork, through which

are passed a safety-tube, terminating an inch above the spirit, and another tube leading to a refrigerator. The safety-tube being filled with pure nitric acid, add through it gradually three and a half fl. ounces of the acid. When the ebullition which ensues is nearly over, add the rest of the acid gradually, half a fl. ounce at a time, waiting till the ebullition which slowly arises has subsided, and cooling the refrigerator with a stream of water. The ether which distils over, being received in a bottle, is to be agitated first with a little milk of lime, till it ceases to redden litmus paper, and then with half its volume of a concentrated solution of muriate of lime. *Edin. Ph.*

This ether is stimulant, antispasmodic, carminative, and diuretic. The dose is from ten to forty drops. Its sp. gr. is 0.899.

SWEET SPIRIT OF NITRE.

R. Hyponitrous ether (0.899),	one part.
Rectified spirit,	four parts.
Mix. (Sp. gr. 0.847.)	<i>Edin. Ph.</i>
R. Nitrate of potassa, in powder,	two pounds.
Sulphuric acid,	a pound and a half.
Alcohol,	nine pints and a half.
Diluted alcohol,	one pint.
Carbonate of potassa,	one ounce.

Mix the nitrate of potassa and the alcohol in a glass retort, and, having gradually added the acid, digest with a gentle heat for two hours; then raise the heat and distil a gallon. To the distilled liquor add the diluted alcohol and carbonate of potassa, and again distil a gallon. (Sp. gr. 0.834.) *U. S. Ph.*

Diuretic, diaphoretic, and carminative. Dose, half a fl. drachm to two fl. drachms.

MIXTURE OF SWEET SPIRIT OF NITRE.

R. Sweet spirit of nitre,	one fl. drachm.
Hoffmann's anodyne,	
Aromatic ammoniated alcohol, each,	two fl. drachms.
Mint-water,	six fl. ounces.

Mix. Dose, a tablespoonful occasionally.

Brera.

MIXTURE OF HYPONITROUS ETHER.

R. Powdered ipecacuanha,	half a drachm.
Boiling water,	sufficient

to obtain six ounces of infusion. Strain and add

Hyponitrous ether, one fl. drachm.
Extract of juniper berries, one ounce.

Mix. Dose, a tablespoonful every two hours, in the dropsical swellings consecutive to scur latina. *Radius.*

ETHER HYDROCYANICUS.

HYDROCYANIC ETHER.

R. Cyanuret of potassium,
Sulpho-vinate of barytes,
equal parts.

Mix, and put in a glass retort, distil by a moderate heat. The product separates into two strata, the lighter of which is impure hydrocyanic ether; this is to be separated and agitated with four or five times its bulk of water at 120° to 140° F., again agitated with a little water, decanted, and placed in contact with chloride of lime for twenty-four hours, and then distilled.

Sp. gr. 0.78.

Magendie.

This preparation is said to resemble hydrocyanic acid in its therapeutical effects, but is less active. Its smell is, however, penetrating and offensive. The dose is one to three drops, in mucilage or emulsion, in obstinate or convulsive coughs.

ETHER MURIATICUS.

MURIATIC ETHER.

R. Muriatic acid,
Alcohol,
equal parts.

Distil in a Wolff's apparatus, and preserve the product contained in the second receiver, surrounded by snow and salt.

Cottureau.

It has much the same properties as the other ethers. Dose, thirty to forty drops.

SPIRIT OF MURIATIC ETHER.

R. Muriatic ether,
Alcohol,
equal parts.
Mix. Dose, half a teaspoonful.

Cottureau.

MIXTURE OF MURIATIC ETHER.

R. Spirit of muriatic ether, half a fl. drachm.

Parsley water,

Syrup of rhubarb, each, one fl. ounce.

Mix. A teaspoonful every hour for young children as a diuretic.

Wendt.

ÆTHER (U. S.)—ÆTHER SULPHURICUS.—(Ed. D.)

SULPHURIC ETHER.

This is commonly known under the name of ether. It is an energetic diffusible stimulant, and is much employed as an excitant and antispasmodic in what are termed nervous disorders. It is also considered to be antihelmintic. In vapor it has been used in small

quantities as an inhalation in chronic irritations of the lungs, and of late under the name of *letheon* to induce unconsciousness to pain in surgical operations. The dose of the fluid is from half a fl. drachm to two drachms. When employed as an inhalation it should be purified. (Sp. gr. .750.)

RECTIFIED SULPHURIC ETHER.

R. Sulphuric ether, fourteen fl. ounces.
Potassa, half an ounce.
Distilled water, eleven fl. ounces.

Dissolve the potassa in two fl. ounces of the water, and add the ether to the solution, shaking well; then distil at 120° F. twelve ounces of rectified ether. Shake the distilled fluid with nine fl. ounces of water, and set them by for the water to subside, then pour off the ether and keep in a well-closed bottle.

ETHEREAL OIL.

(OIL OF WINE.)

R. Alcohol, two parts.
Sulphuric acid, four parts.

Mix. Distil till a black froth begins to arise remove the retort from the fire, when cool, add water to the fluid in the receiver, and remove the oil that may float on the surface. Agitate this with a solution of potassa, and separate the ethereal oil.

Van Mons.

SPIRIT OF SULPHURIC ETHER.

R. Sulphuric ether, one pint.
Alcohol, two pints.
Mix. *Ed. Ph.*

HOFFMANN'S ANODYNE.

R. Sulphuric ether, half a pint.
Alcohol, one pint.
Ethereal oil, three fl. drachms.

Mix. *U. S. Ph.*
Dose, a teaspoonful in sweetened water.

LOTION OF SULPHURIC ETHER.

R. Sulphuric ether, two fl. ounces.
Vinegar, six fl. ounces.
Rose water, four fl. ounces.
Distilled water, two pints.

Mix. As a lotion to painful inflammatory tumors.

Pierquin.

SYRUP OF SULPHURIC ETHER.

R. Sulphuric ether, one fl. ounce.
Syrup, one pint.

Mix in a glass vessel, having a stop-cock at the lower part; shake occasionally for a week, and draw off when clear into small bottles.

Par. Cod.

Dose, half an ounce to an ounce.

R. White sugar, twenty-one ounces.
Distilled water, six fl. ounces.
Sulphuric ether, two fl. ounces.

Mix, and pour into a closed flask, agitate for some minutes, and add

Distilled water, six ounces.

Agitate occasionally for five days, filter in a covered funnel. *Paton.*

MIXTURE OF SULPHURIC ETHER AND CAMPHOR.

R. Camphor mixture, seven fl. ounces.
Sulphuric ether,
Syrup of saffron, each, half fl. ounce.

Mix. Dose, a tablespoonful.

MIXTURE OF SULPHURIC ETHER AND TURPENTINE.

R. Sulphuric ether, one fl. drachm.
Oil of turpentine, two fl. drachms.
Sugar, four drachms.
Water, two fl. ounces.

Mix. Two fl. drachms every quarter of an hour in poisoning by *nux vomica*. *Orfila.*

ETHER TEREBINTHINATUS.

TEREBINTHINATED ETHER.

R. Alcohol, two pounds.
Spirit of turpentine, half a pound.

Mix, and add gradually

Nitric acid, two pounds.

Distil off one-half at a gentle heat. Dose, from twenty to forty drops, in syrup or mucilage. Used internally and externally in cases of biliary calculi, jaundice, engorgements of the liver, and rheumatism. *Cadet.*

ALETRIS.

STAR GRASS.

Aletris farinosa. Linn. Bigelow. Am. Med. Bot. iii. 50. Griffith, Med. Bot. 623.

Sex. Syst. Hexandria monogynia. *Nat. Syst.* Hæmodoraceæ. Brown.

A native plant, with an intensely bitter root, which is the officinal portion. The powder is tonic in doses of ten grains.

TINCTURE OF ALETRIS.

R. Aletris root, bruised, three ounces.
Diluted alcohol, two pints.

Make tincture by displacement. Dose, half an ounce to an ounce in colic, and in smaller doses in chronic rheumatism.

ALLIUM.

GARLIC.

Several species of *Allium* have been employed in medicine, but the most important, and that recognized by the Pharmacopœia, is *A. sativum*. Linn. Griffith, Med. Bot. 653. The part used is the bulb, which is composed of several bulbets or *cloves* (*spicæ*), surrounded by a dry, white, thin capsular membrane. These cloves have a peculiar odor, and an acrid and pungent taste. The properties depend on the presence of a volatile oil, and are lost on drying.

Sex. Syst. Hex. monog. *Nat. Syst.* Liliacæ.

Garlic is a stimulating expectorant; diuretic and diaphoretic when given internally, and acts as an irritant, or even vesicant, when applied externally. The dose in substance is from half a drachm to a drachm, or more.

SYRUP OF GARLIC.

R. Fresh garlic, six ounces.
Diluted acetic acid, one pint.
Sugar, two pounds.

Macerate the garlic in ten ounces of acid, in a glass vessel for four days, and express. Then add remainder of acid to dregs, and again express. Add sugar and form syrup. *U. S. Ph.*

As an expectorant in chronic catarrhs, well suited for children. Dose, for a child, about a teaspoonful.

CATAPLASM OF GARLIC.

Bruised cloves of garlic, mixed with common bread and milk poultice, in different proportions. Used as a revulsive, but less active than the mustard cataplasm.

LINIMENT OF GARLIC.

R. Garlic, equal parts.
Lard, equal parts.

To be well rubbed together. Has been recommended as a revulsive in infantile nervous and convulsive disorders. *Gassicourt.*

CAPILLARY LOTION.

R. Garlic, three or four cloves.
Alcohol, two pints.

Macerate for thirty-six hours, filter, and add of Burdock, eight ounces.

The head to be sponged with this every evening, for some weeks. It is said to be efficient to promote the growth of hair. *Phabus.*

ALOE.

ALOES.

Under this name are included the inspissated juices of several species of Aloe, and especially of *A. vulgaris*, *A. socotrina*, and *A. spicata*.

Sex. Syst. Hexandria monogynia. *Nat. Syst.* Liliacæ.

Pereira, Mat. Med. ii. 113. Griffith, Med. Bot. 649.

Three varieties of aloes are found in the shops, the Cape, Socotrine, and Hepatic. The second, when genuine, is the most esteemed, but the first is by far the most abundant, and, when good, answers every purpose for which the drug is used.

U. S. Dispens. art. Aloe.

POWDER OF ALOES.

R. Aloes, six grains.
White sugar, one drachm.
Pulverize very finely. Recommended to be blown in the eye, to remove films and specks on the cornea.

Radius.

POWDER OF ALOES AND CANELLA.

R. Aloes, one pound.
Canella bark, three ounces.
Powder separately, and mix. Dose, five to fifteen grains. A popular emmenagogue, known as *Hiera Picra*.

U. S. Ph.

COMPOUND POWDER OF ALOES.

R. Aloes, an ounce and a half.
Guaiacum resin, one ounce.
Compound powder of cinnamon, half an ounce.

Powder the aloes and resin separately, and then mix them with the compound powder of cinnamon. Dose, ten to twenty grains. Used as a purgative and diaphoretic.

London Ph.

EMMENAGOGUE POWDER.

R. Iron rust, six grains.
Powdered aloes, two grains.
Magnesia, thirty grains.

Mix. For a single dose, to be repeated three times a day.

Brera.

ALOE PILLS.

R. Powdered aloes,
Soap, each, an ounce.

Form a mass with water, to be divided into two hundred and forty pills. Dose, one to three as a laxative, five or more as a purgative.

U. S. Ph.

COMPOUND PILLS OF ALOES.

R. Powdered aloes, half a drachm.
" rhubarb, one drachm.
Oil of cloves, four drops.
Soap, eight grains.
Syrup of rhubarb, sufficient.

Rub well together, and form forty pills. In tardy menstruation, one at night or oftener if required, so as to open bowels but not to purge.

Devees.

R. Powdered aloes, one ounce.
Extract of gentian, half an ounce.
Oil of caraway, forty minims.
Molasses, sufficient.

Beat together till incorporated. *London Ph.*

Dose, five to fifteen grains. A valuable purgative in habitual costiveness.

ALOE AND ASSAFŒTIDA PILLS.

R. Powdered aloes, }
Assafœtida, } each, half an ounce.
Soap, }

Beat with water to form a mass; divide into one hundred and eighty pills. Dose, two to five. Useful in costiveness with flatulency.

U. S. Ph.

ALOE AND MYRRH PILLS.

R. Powdered aloes, two ounces.
Powdered myrrh, one ounce.
Saffron, half an ounce.
Syrup, sufficient.

Beat together, to form a mass; divide into four hundred and eighty pills.

U. S. Ph.

Dose, from three to six. A well-known cathartic and emmenagogue, under the name of Rufus's pills.

ALOE AND IRON PILLS.

R. Barbadoes aloes, two parts.
Sulphate of Iron, three parts.
Aromatic powder, six parts.
Conserve of red roses, eight parts.

Pulverize the aloes and sulphate of iron, mix the whole ingredients and beat into a mass, and divide into five grain pills. Dose, one to three.

Edin. Ph.

A useful emmenagogue in chlorosis and atonic amenorrhœa.

CHAPMAN'S APERIENT PILLS.

R. Aloes, sixteen grains.
Powdered rhubarb, twenty-four grains.
Mastic, twelve grains.

Mix, and make twelve pills. *Chapman.*

CHAPMAN'S ANTI-DYSPEPTIC PILLS.

R. Aloes, half a drachm.
Powdered ipecacuanha, ten grains.
Mastic, thirty grains.
Oil of fennel, eight drops.

Mix, and make twenty pills. *Chapman.*

MITCHELL'S APERIENT PILLS.

R. Aloes, sixteen grains.
Powdered rhubarb, thirty-two grains.
Calomel, two grains.
Tartar emetic, one grain.
Mix, and make sixteen pills.

J. K. Mitchell.

Two or three pills act as an aperient.

HOOPER'S PILLS.

R. Aloes, four hundred parts.
Crystallized Sulphate of iron, two hundred parts.
Extract of black hellebore, } each, one hundred parts.
Myrrh, }
Soap, }
Powdered canella, }
Powdered ginger, each, fifty parts.

Beat into a mass with water, and divide into pills of two and a half grains.

Much used as a cathartic, and an emmenagogue. *Journ. Phil. Col. Pharm. v. 25.*

ANDERSON'S PILLS.

R. Aloes, seven hundred and eighty-seven parts.
Soap, one hundred and thirty-one parts.

Colocynth,
Gamboge, each, thirty-three parts.
Oil of aniseed, sixteen parts.

Reduce the aloes, colocynth, and gamboge to a very fine powder, add oil of aniseed and soap, and beat into a mass with water; divide into three grain pills. A mild purgative.

Journ. Phil. Col. Pharm. v. 25.

LADY WEBSTER'S PILLS.

R. Aloes, six drachms.
Mastich,
Red rose-leaves, each, two drachms.
Syrup of wormwood, sufficient

to form a mass. Divide into three grain pills; used as a laxative in costiveness, dependent on impaired digestion. Dose, one to three. A better form is to substitute powdered rhubarb for the rose-leaves. *Cooley.*

MORRISON'S PILLS.

No. 1. R. Aloes,
Cream of tartar, equal parts.
Mucilage, sufficient

to form mass.

No. 2. R. Aloes, three parts.
Gamboge, two parts.

Colocynth, one part.
Cream of tartar, four parts.

Powder finely, and add
Simple syrup, sufficient
to form mass. Dose, five to ten grains. These
purge actively, especially No. 2. *Cooley.*

FULLER'S PILLS.

R. Aloes, half a drachm.
Senna,
Myrrh, each, one scruple.
Assafetida,
Galbanum, each, ten grains.
Saffron,
Mace, each, five grains.
Sulphate of iron, two scruples.
Simple syrup, sufficient

to form mass. Dose, five to fifteen grains, used as an antispasmodic and aperient. *Cooley*

JAMES'S ANALEPTIC PILLS.

R. Antimonial powder,
Resin of guaiacum,
Aloes and myrrh pill, equal parts.
Syrup, sufficient.

Form a mass, and divide into four grain pills. A diaphoretic purgative. *Cooley.*

PETERS'S PILLS.

R. Aloes, }
Jalap, } each, two drachms.
Scammony, }
Gamboge, }
Calomel, one drachm.

Beat into a mass with alcohol. A powerful purgative. Three pills, containing three grains each, constitute a full dose. *Cooley.*

SPEEDIMAN'S PILLS.

R. Aloes,
Myrrh,
Rhubarb, each, one ounce.
Extract of chamomile, half an ounce.

Beat into a mass with syrup, and divide into four grain pills. A good tonic and stomachic purge. *Burnett.*

SPLENETIC PILLS.

R. Strained aloes,
Gum ammoniac, each, an ounce and a half.
Myrrh,
Bryony, each, two drachms.

Beat into a mass, and divide into four grain pills. Dose, three to five. Extolled in amenorrhœa and hypochondriasis. *Saunders*

DUCHESNE'S PILLS.

R. Aloes,
 Gum ammoniac, each, twenty-four grains.
 Myrrh, six grains.
 Mastich,
 Carbonate of potassa, each, two grains.
 Saffron, one grain.
 Simple syrup, sufficient
 to make a mass. Dose, from ten to twenty grains. Prescribed in engorgements of the abdominal viscera, supervening on intermittent fevers.
Van Mons.

ANTICHLOROTIC PILLS.

R. Aloes,
 Iron rust, each, half a drachm.
 Gum ammoniac, one drachm.
 Extract of dandelion, sufficient
 to form mass. Divide into pills of three grains each. Dose, three to six, morning and evening, in chlorosis and amenorrhœa.
Radius.

BICKER'S PILLS.

R. Iron rust, two drachms.
 Sulphur, }
 Myrrh, } each, one drachm.
 Aloes, }
 Beef gall, sufficient
 to form mass. Divide into four grain pills. Six morning and evening.
Radius.

WHYTT'S PILLS.

R. Chloride of iron,
 Aloes,
 Extract of horehound, each, half a drachm.
 Assafetida, one drachm and a half.
 Form a mass. Divide into two grain pills. Dose, four to five, three times a day, in leucorrhœa and hysteria with constipation.
Radius.

BARTHEZ'S PILLS.

R. Aloes, half a drachm.
 Myrrh, one drachm.
 Musk, twenty grains.
 Camphor, ten grains.
 Balsam Peru, sufficient
 to form mass. Dose, eight grains, three times a day. Advised in amenorrhœa and chlorosis.
Pierquin.

PITSCHAFT'S ECCOPROTIC PILLS.

R. Strained aloes,
 Sulphate of quinine, each, one scruple.

Mix, and divide into twenty pills. Dose, one at bedtime. Found useful in torpor of the large intestines.
Radius.

FRANK'S PILLS.

R. Aloes,
 Jalap, each, four parts.
 Rhubarb, one part.
 Syrup of wormwood, sufficient
 to form mass. Divide into three grain pills
 Dose, one to four during the day.
 These are the *Grains de Santé*, so much employed as dinner pills in Europe.
Foy.

ALOEES AND RHUBARB PILLS.

R. Powdered aloes,
 Rhubarb, each, half a drachm.
 Soap, sufficient
 to form mass. Divide into twenty-five pills. Three or four occasionally in dyspepsia with costiveness.
Ellis.

GRIFFITHS'S PILLS.

R. Powdered rhubarb, one drachm and a half.
 Sulphate of iron, half a drachm.
 Soap, two scruples.
 Water, sufficient
 to form mass. Divide into forty pills.
 A favorite remedy with the late Dr. S. P. Griffiths, to remove costiveness and impart tone to the bowels. Three or four, to be taken at bedtime.

APERIENT PILLS.

R. Aloes,
 Rhubarb, each, one drachm.
 Ipecacuanha, six grains.
 Soap, one scruple.
 Form mass with water, and divide into sixty pills. Dose, one at bedtime, as an aperient; two or three as a purgative.
Ellis.

PILLS OF ALOES AND BLUE MASS.

R. Blue mass, one scruple.
 Powdered aloes, twenty-five grains.
 Incorporate, and divide into fifteen pills. Dose, one every two hours till they operate. Useful in constipation where there is a deficiency of bile.
Ellis.

LAXATIVE ELECTUARY.

R. Aloes, eight grains.
 Cream of tartar, two drachms.
 Honey, sufficient
 to form electuary. For a single dose. Advised in amenorrhœa attributed to abdominal engorgement.
Brera.

ANTHELMINTIC SUPPOSITORY.

R. Aloes, half an ounce.
Common salt, three drachms.
Flour, two ounces.
Honey, sufficient
to form a firm paste, to be divided into twelve
suppositories. Used in cases of ascarides.
Foy.

COMPOUND DECOCTION OF ALOES.

R. Extract of liquorice, seven drachms.
Carbonate of potassa, one drachm.
Powdered aloes
" myrrh,
Saffron, each, one drachm and a half.
Compound tincture of cardamom,
seven fl. ounces.
Distilled water, one pint and a half.
Boil down the liquorice, carbonate of potassa,
aloes, myrrh, and saffron with the water to a
pint, strain, and add the compound tincture of
cardamom. Dose, half a fl. ounce to two fl.
ounces. A mild cathartic, tonic, antacid, and
emmenagogue. *Lond. Ph.*

CLAUDER'S ELIXIR.

R. Carbonate of potassa,
Muriate of ammonia, each, one ounce.
Elder-flower water, a pint and a half.
Make a solution and add
Strained aloes,
Myrrh, each, one ounce.
Saffron, two drachms.
Digest for twenty-four hours, and filter. Dose,
half a drachm to a drachm. Has been recom-
mended in obstructions of the abdominal viscera,
in amenorrhœa, constipation, scurvý, &c.
Pideret.

DETERSIVE INJECTION.

R. Strained aloes, ten grains.
Muriate of ammonia, four grains.
Honey of roses, one ounce.
Fennel water, six ounces.
Make solution, and filter. Stated to be useful
in chronic discharges from the urethra, to be in-
jected three or four times a day. *Soubeiran.*

ANTHELMINTIC CLYSTER.

R. Powdered aloes, one drachm.
Barley water, one pint.
Mix. Very effectual against ascarides in the
rectum. *Radius.*

ALOETIC MIXTURE.

R. Strained aloes,
Myrrh, each, one drachm.
Cream of tartar, half an ounce.
Myrrh water, six ounces.

Make a solution, and filter. Dose, a table
spoonful night and morning, to provoke hemor-
rhoids. *Foy.*

ALKALINE MIXTURE OF ALOES.

R. Aloes, two ounces and a half.
Bicarbonate of soda, six ounces.
Compound spirit of
lavender, two fl. ounces.
Water, four pints.
Mix. Macerate for two weeks, and filter.
Dose, from one fl. drachm to one fl. ounce, half
an hour after meals, for persons of a costive
habit. *Mettauer.*

WINE OF ALOES.

R. Aloes, one ounce.
Cardamom,
Ginger, each, one drachm.
Wine, one pint.
Macerate for fourteen days, occasionally agi-
tating, then filter. *U. S. Ph.*
Purgative, in doses of half an ounce to two
ounces; stomachic and tonic, in doses of one to
two drachms.

ALKALINE WINE OF ALOES.

R. Aloes,
Myrrh,
Saffron, each, one ounce.
Carbonate of potassa, two ounces.
Wine, two pints.
Digest for ten days, and filter. Bitter, tonic
and stimulant. Recommended in dyspepsia
with pyrosis, in doses of an ounce. *Swediaur.*

BALSAMIC WINE OF ALOES

R. Aloes, }
Myrrh, } each, half an ounce.
Olibanum, }
Angelica, }
Balsam Peru, one ounce.
Storax, two ounces.
Benzoin, three ounces.
Flowers of hypericum, four handfuls.
Wine, four pints.
Macerate for fourteen days, and strain. Dose
half an ounce.

TINCTURE OF ALOES.

R. Powdered aloes, one ounce.
Liquorice, three ounces.
Alcohol, half a pint.
Distilled water, one pint and a half.
Macerate fourteen days, and filter. Dose, two
drachms to an ounce. Purgative and sto-
machic. *U. S. Ph.*

TINCTURE OF ALOES AND MYRRH.

R. Powdered aloes, three ounces.
Saffron, one ounce.
Tincture of myrrh, two pints.
Macerate for fourteen days, and filter.

U. S. Ph.

R. Tincture of aloes, three fl. ounces.
Tincture of myrrh, four fl. ounces.
Tincture of saffron, three fl. ounces.

Mix.

Soubeiran.

Long celebrated under the name of *elixir proprietatis*. It is purgative, stomachic, and emmenagogue. The dose is half a drachm to a drachm. Well suited to cold, torpid habits.

ETHEREAL TINCTURE OF ALOES.

R. Myrrh, one ounce and a half.
Spirit of sulphuric ether, one pound.

Digest for four days, and add

Aloes, one ounce and a half.
Saffron, one ounce.

Again digest for four days, and filter.

Edin. Ph. 1817.

This tincture has been highly esteemed as stomachic, vermifuge, and emmenagogue, in doses of half a drachm to a drachm.

BOERHAAVE'S ELIXIR.

R. Aloes,
Myrrh,
Saffron, each, one ounce.
Tartrate of potassa, two ounces.
Alcohol, fourteen ounces.
Distilled water, eight ounces.

Macerate for three days, and filter. Han. Ph.

This has been highly praised in visceral obstructions. The dose is from half a drachm to a drachm.

STOUGHTON'S ELIXIR.

R. Aloes,
Cascarilla, each, one drachm.
Rhubarb, four drachms.
Wormwood, }
Germander, } each, six drachms.
Gentian, }
Orange peel, }
Alcohol, two pints.

Macerate for four days, and filter. Foy.

Stimulant, tonic, and stomachic, in doses of twenty or thirty drops.

ELIXIR OF GARUS.

R. Saffron, one ounce.
Myrrh, two ounces.
Aloes, ten ounces.

Cinnamon,

Nutmeg, each, half an ounce.

Orange-flower water, one pound.

Alcohol, sixteen pounds.

Macerate for two days and distil off eight pounds, to which add

Syrup of maidenhair, two pounds.

And color with caramel, dissolved in

Orange-flower water, eight ounces.

This preparation can scarcely be considered as a medicine, but is rather an agreeable liquor.

Soubeiran.

OINTMENT OF ALOES.

R. Powdered aloes, two drachms.
Lard, one ounce.

Triturate well.

Has been recommended as a friction to the abdomen as a vermifuge in children.

Soubeiran.

ALTHÆA.

MARSH MALLOW.

Several species of Mallow are employed in medicine, but that which is officinal is *A. officinalis*, a herbaceous perennial, with pale purplish flowers, a native of Europe, in moist situations.

Sex. Syst. Monadelph. Polyand. *Nat. Syst.* Malvaceæ.

Linn. Sp. Pl. 966. Griffith, Med. Bot. 161.

The parts used are the roots and leaves. These are inodorous, and have a rapid mucilaginous taste. They are much employed in Europe as demulcent and emollient, but their place is supplied in this country by other articles.

DECOCTION OF MARSH MALLOW.

R. Marsh mallow-root, dried, four ounces.

Raisins, stoned, two ounces.

Boiling water, five pints.

Boil down to three pints. Strain, and set aside until the dregs have subsided, and decant.

Ed. Ph.

A good demulcent drink.

SYRUP OF MARSH MALLOW.

R. Marsh mallow root, thirty parts.
Cold water, one hundred and eighty parts.

Macerate for twelve hours, strain without expression, and add

Syrup, one thousand parts.

Evaporate to the proper consistence, and strain.

Codex.

A good demulcent. Much used in France.

MARSH MALLOW PASTE.

R. Marsh mallow root, powdered,
 Wheat flour, each, three parts.
 White sugar, six parts.
 Mucilage of gum Arabic, two parts.
 Mix, and form paste. *Beral.*
 A good demulcent in irritation of the throat.

MARSH MALLOW LOZENGES.

R. Powdered mallow root, one part.
 White sugar, seven parts.
 Mucilage of tragacanth, sufficient.
 Form into lozenges of sixteen grains each.
Cottureau.

MARSH MALLOW OINTMENT.

R. Flaxseed,
 Mallow root, each, one pound.
 Water, sufficient
 to make a thick mucilage; strain, and add
 Lard, melted, six pounds.
 Evaporate, and add
 Resin, eight ounces.
 Turpentine,
 Yellow wax, each, one pound.
 Melt the whole together. *Beral.*

POWDER OF MARSH MALLOW.

R. Powdered mallow root,
 " liquorice root, each,
 three ounces.
 " nitre, half an ounce.
 " camphor, one drachm.
 Mix, and divide into thirty powders; one to be
 taken three times a day. *Foy.*

ALUMINA.

ALUMINA.

The hydrate of alumina is found naturally in an impure state in the form of different clays, boles, &c., and can be prepared in a pure state by the decomposition of alum.

HYDRATE OF ALUMINA.

R. Alum, any quantity.
 Distilled water, sufficient
 to dissolve the salt. Add gradually a solution of carbonate of potassa, digest at a moderate temperature for a short time, until a precipitate no longer takes place, wash this carefully, and dry. *Sax. Ph.*

RUST'S ASTRINGENT.

R. Armenian bole,
 Carbonate of magnesia,
 White sugar, each, two drachms.
 Oil of mace, two drops
 Rhubarb, one drachm.
 Mix, and triturate well. *Radius.*

Dose, a teaspoonful three times a-day, in chronic gonorrhœa, vaginal and uterine catarrh and slight menorrhagia.

ELECTUARY FOR THE TEETH.

R. White bole, an ounce and a half.
 Gum lac, half an ounce.
 Cinnamon, a drachm and a half.
 Syrup of pinks, sufficient
 to form soft paste. Triturate well together and keep in closed boxes. Used as a dentifrice, but is also beneficial in scorbutic gums. *Swediaur.*

ALUMINÆ ACETAS.

ACETATE OF ALUMINA.

R. Alum, a sufficient quantity.
 Dissolve in water, precipitate by means of aqua ammoniæ, filter, and dissolve the moist precipitate in acetic acid; again filter, and evaporate.

This has been recommended diluted with water in chronic diarrhœa, and mixed with syrup of poppies in slight cases of hemoptysis. Dose, three to five grains. *Van Mons.*

ALUMINÆ SULPHAS.

SULPHATE OF ALUMINA.

R. Alum, a sufficient quantity.
 Dissolve in water, precipitate by aqua ammoniæ, filter, and dissolve the moist precipitate in dilute sulphuric acid, and again filter; evaporate and crystallize.

DETERGENT WASH.

R. Sulphate of alumina, two drachms.
 Water, half a pint.
 Mix. As a wash to foul ulcers. *Pennypacker.*

M. Gannal has found that a solution of this salt, made with about a pound to the quart of water, will preserve a body fresh for a long time, if injected into the blood-vessels; where it is only wished to preserve the body for a month or six weeks, an enema of one quart, and an injection of about a quart into the œsophagus is sufficient. *Dunglison, N. R.*

ALUMEN.

ALUM.

Several varieties of alum are known in commerce, all containing more or less sulphate of iron. Most of that used in this country is of domestic manufacture.

Alum is employed both internally and externally as an astringent, in doses of ten grains to a scruple or more.

BURN'T ALUM.

R. Alum, any quantity.
Melt the alum in an iron or earthen vessel, and continue a moderate heat till it ceases to bubble and becomes dry, then rub it into powder.

U. S. Ph.

Astringent, and a mild escharotic. It is much used to repress the growth of exuberant granulations or proud flesh. Dose, five to ten grains.

POWDER OF ALUM.

R. Alum, one drachm.
Gum Arabic, half a drachm.

Triturate, and divide into four powders, one to be taken every three hours in atonic hemorrhage. *Radius.*

R. Alum, a drachm and a half.
Catechu, one drachm.
Armenian bole, one ounce.

Triturate well. To arrest hemorrhages. Dose, a teaspoonful. *Van Mons.*

R. Alum, four ounces.
Kino, one ounce.

Triturate well. This is the *Saccharine Alum* of the continental writers. Dose, ten to twenty grains, in hemorrhages or diarrhœa. *Ed. Ph.*

R. Alum, thirty grains.
Powdered opium, three grains.

Mix, and make six powders, one to be taken every four hours. *A. T. Thomson.*

R. Alum, one drachm.
Powdered opium, four grains.
" cinnamon, one scruple.

Mix and triturate, divide into four powders, one to be taken every four hours, in menorrhagia or in diarrhœa. *Radius.*

R. Alum, two drachms.
Powdered opium, three grains.
" kino, one scruple.
Sugar of milk, one drachm.

Mix, triturate, and divide into six powders. One to be taken every three hours, in fluxes. *Phæbus.*

ALUM ERRHINE.

R. Alum,
Armenian bole, each, one drachm.
Kino, half a drachm.

Red oxide of iron, two drachms.
Mix, and triturate. As an errhine in epistaxis. *Radius.*

GARGLES OF ALUM.

R. Alum,
Nitrate of potassa, each, three ounces.
Cream of tartar, four ounces.
Acetic acid, four pounds.

Dissolve, evaporate to dryness, and powder the residuum. Half an ounce, dissolved in eight ounces of water, forms a gargle which has been highly praised in inflammation of the fauces and tonsils. *Wirtem. Ph.*

R. Alum, two drachms.
Water, four fl. ounces.

Dissolve. Said to be useful where the breath is offensive. *Cavarra.*

ODONTALGIC SOLUTION OF ALUM.

R. Finely powdered alum, two drachms.
Spirit of nitric ether, seven fl. drachms.

Dissolve. Stated to be an almost infallible cure for toothache. *Blake.*

BOLUSES OF ALUM.

R. Alum,
Extract of Peruvian bark,
Nutmeg, each, half a scruple.
Simple syrup, sufficient

to make a bolus. Prescribed in uterine and other hemorrhages, and repeated as may be required. *Ellis.*

R. Alum, five grains.
Extract of rhatany, eighteen grains.
Conserve of roses, half a drachm.
Syrup of rhatany, sufficient

to make ten boluses. Dose, one every three or four hours. *Foy.*

ALUM PILLS.

R. Alum, six grains.
Extract of opium, one grain.
Catechu, six grains.

Mix, and form into six pills. One to be given every two to four hours. In passive hemorrhages and atonic mucous discharges. *Ellis.*

R. Alum, two drachms.
Kino,
Honey of roses, each, one drachm.

Make pills, of six grains each. Dose, one to six a day, in passive hemorrhages. *Foy.*

R. Alum,
Extract of Peruvian bark,
Muriate of iron and ammonia,
Aromatic powder, each, a drachm
and a half.

Oil of cinnamon, twelve drops.

Mix, and make pills of two grains. Dose, four to six, morning and evening, in passive hemorrhages and mucous discharges. *Augustin.*

ASTRINGENT PILLS OF ALUM.

R. Alum, equal parts.
Catechu, sufficient
Extract of gentian, sufficient
to make pills of two grains each. Dose, four every three hours. Useful in diarrhœa, and especially in leucorrhœa. *Radius.*

PILLS OF ALUM AND BENZOIC ACID.

R. Alum, one scruple.
Benzoic acid, five grains.
Gum Arabic,
White sugar, each, ten grains.

Mix, with sufficient water to form thirty-five pills, to be taken in two days. Have been thought useful in phthisis pulmonalis. *Augustin.*

ELECTUARIES OF ALUM.

R. Alum, one drachm.
Catechu,
Extract of Peruvian bark, each, two drachms.
Conserve of roses, six drachms.
Simple syrup, sufficient

to make an electuary. Dose, one drachm every four hours. In chronic diarrhœa, leucorrhœa, and passive hemorrhages. *St. Marie.*

R. Alum, one drachm.
Extract of logwood, half an ounce.
Balsam Peru, six drops.
Water of sage, sufficient.

Make an electuary. Astringent and antiseptic. Recommended in sponginess of the gums. *Phœbus.*

ALUM COLLYRIUM.

R. Alum, twelve to twenty grains.
Rose water, four fl. ounces.
Dissolve. As an application in chronic ophthalmia. *Ellis.*

ALUM CURD.

R. Powdered alum, half a drachm.
White of egg, one.

Agitate well till a coagulum is formed. To be applied on a rag to inflamed eyes. To be removed when it becomes warm. *Ellis.*

SOLUTIONS OF ALUM.

R. Burnt alum,
Sulphate of zinc, each, half an ounce.
Distilled water, two pints.

Dissolve.

R. Alum, one drachm
Sulphate of zinc, half a drachm.
Borax, four grains.
Rose water, six ounces.

Dissolve. Used as an astringent in bruises, slight hemorrhages, and mucous discharges.

Cadet de Gassicourt.

R. Alum, half a drachm.
Armenian bole, six drachms.
Vinegar,
Red wine, each, half an ounce.

Mix the powdered bole in the solution of the alum in the vinegar, and add the wine. Used as a local application in epistaxis. *Swediaur.*

GARGLES OF ALUM.

R. Alum, one drachm.
Wine, one pound.
Tincture of bark, four fl. drachms.
" myrrh, two fl. drachms.
Honey of roses, two ounces.
Laudanum, one fl. scruple.

Mix. As a gargle in scurvy of the gums.

Foy.

R. Alum, one drachm.
Infusion of red roses,
Barley water, each, three fl. ounces.

Add to the solution

Honey of roses, two fl. ounces.
Ratier.

R. Powdered oak bark, one ounce.
Boiling water, a pint and a half.

Evaporate to one pint, filter, and add

Alum, half a drachm.
Brandy, two fl. ounces.

Used as a gargle in inflammation of the mouth and throat. *Augustin.*

INJECTIONS OF ALUM.

R. Alum, one drachm.
Rose water, two fl. ounces.

Mix. In chronic gonorrhœa.

Foy.

R. Alum,
Sulphate of iron, each, twelve grains.
Honey of roses, one ounce
Barley water, five fl. ounces.

Mix. In chronic gonorrhœa.

Radius.

R. Stramonium leaves, half an ounce.
Boiling water, two pints.

Infuse for half an hour, strain, and add to infusion

Alum, two drachms, to an ounce.
In cancerous affections of uterus. *Foy.*

ALUM INJECTION.

R. Infusion of flaxseed, fifteen fl. ounces.
Alum, two drachms.
Tincture of kino, one ounce.
Mix. As an injection in cauliflower excrescence of the uterus. *Clarke.*

JULEP OF ALUM.

R. Alum, two drachms.
Sulphuric acid, ten drops.
Essence of citron, six drops.
Syrup of lemon, two fl. ounces.
Water, three fl. ounces.
Mix. A tablespoonful every hour, in lead colic. *Radius.*

ALUM LINIMENT.

R. Alum, half an ounce.
Whites of eggs, four.
Spirit of camphor, two ounces.
Mix. As an application to bed sores. *Augustin.*

LOTIONS OF ALUM.

R. Alum, two drachms.
Sulphate of zinc, each, two drachms.
Plantain water, two pints.
Mix. As a wash to wounds and bleeding ulcers. *Cadet de Gassicourt.*
R. Alum, three drachms.
Muriate of ammonia, one drachm.
Liquid hydrosulphuric acid, one scruple.
Rose water, half a pint.
Mix. As a wash, in obstinate cutaneous affections. *Alibert.*

ALUM WHEY.

R. Boiling milk, one pint.
Powdered alum, a drachm and a half.
Mix, separate the curd, and add to the whey
White sugar, one ounce.
Dissolve. In passive hemorrhages. A wine-glassful occasionally, to be taken cold. *Foy.*

AROMATIC ALUM WHEY.

R. Powdered alum, one drachm.
Sugar of milk, half a drachm.
Powdered cinnamon, fifteen grains.
Cow's milk, a pint and a half.

Boil, and strain after coagulation. To be taken in cupful doses, cold, in hemorrhages. *Niemann.*

ALUM DRAUGHTS.

R. Alum, one to two drachms.
Syrup of gum, two ounces.
Distilled water, four fl. ounces.
Mix. Dose, a wineglassful every six hours, in lead colic and uterine hemorrhages. *Guibourt.*
R. Alum, two drachms.
Wine, four fl. ounces.
Gum tragacanth, a drachm.
Catechu, each, eight fl. ounces.
Water, eight fl. ounces.
Mix. One or two spoonfuls every hour in lead colic. *Radius.*

ANGLO-SAXON OINTMENT.

R. Red lead, one pound.
Olive oil, one pound.
White wax, each, one pound.
Powdered amber, one pound.
Burnt alum, two drachms.
Camphor, each, two drachms.
Heat the oil until it becomes of a reddish-brown color, add the red lead and continue the heat; when the mass has the consistence of a plaster, add the amber, and finally, when it cools, the alum and camphor. Highly spoken of as a dressing to foul ulcers. *Niemann.*

OINTMENT FOR CHILBLAINS.

R. Sweet almonds, blanched, half a pound.
Honey, six ounces.
Camphor, }
Flour of mustard, } each, two ounces.
Burnt alum, }
Olibanum, }
Yolks of eggs, three.

Triturate well. This paste is to be thinned in a little water, and rubbed on the inflamed parts night and morning, and then washed off in tepid water, and afterward the parts well dried with a linen cloth. *Suediaur.*

RUST'S OINTMENT.

R. Alum, a drachm and a half.
Camphor, a scruple to half a drachm.
Opium, each, a scruple to half a drachm.
Balsam Peru, one drachm.
Lead ointment, half an ounce.
Triturate well together. Found useful as an application to frosted limbs. *Phabus.*

PILE OINTMENT.

R. Powdered alum, four scruples.
Simple cerate, five ounces and a half.
Mix, and triturate well. As an application to
painful hemorrhoids. *Taddei.*

AMMONIA.

AMMONIA.

This, under ordinary circumstances, exists as a gas, which is readily taken up by water or alcohol. Water absorbs nearly 800 times its volume of the gas at 32° F., and this solution, possessing the properties of the gas, is usually employed.

SOLUTION OF AMMONIA.

R. Pulverized muriate of ammonia,
Quicklime, each, one pound.
Distilled water, one pint.
Water, nine fl. ounces.

Break the lime in pieces, pour the water upon it in a proper vessel, cover this and permit it to rest until the lime slacks and is cold. Mix this with the muriate of ammonia in a mortar, and introduce the mixture into a glass retort. Place the retort on a sand-bath, and adapt to it a receiver connected by means of a glass tube with a quart bottle containing the distilled water. Apply heat, and continue as long as ammonia comes over. Remove the fluid in the bottle, and add so much water as will raise it to the sp. gr. 0.960, and keep in small well-closed bottles. The *liq. ammon. fort.* of the Pharmacopœia has a sp. gr. of 0.882, and the above *liq. ammoniæ* that of 0.960; one fl. ounce of the former with two of water will form the latter.

U. S. Ph.

LINIMENT OF AMMONIA.

R. Solution of ammonia, one fl. ounce.
Olive oil, two fl. ounces.

Mix. Used as a rubefacient, and counter-irritant. *U. S. Ph.*

COMPOUND LINIMENT OF AMMONIA.

R. Stronger water of ammonia, five fl. ounces.
Tincture of camphor, two fl. ounces.
Spirit of rosemary, one fl. ounce.

Mix well. More powerful than the simple liniments. *Edin. Ph.*

GRANVILLE'S LOTION.

(*Milder.*)

R. Water of ammonia (.882), four fl. drachms.

Spirit of rosemary, three fl. drachms.
Spirit of camphor, (3j to Oj),
one fl. drachm.

Mix.

(*Stronger.*)

R. Water of ammonia (.882),
five fl. drachms
Spirit of rosemary, two fl. drachms.
Spirit of camphor, one fl. drachm.

Mix. As a counter-irritant. *Gray.*

GARGLE OF AMMONIA.

R. Water of ammonia (.960), half fl. ounce.

To be added to a strained decoction of

Mallows,
Dry figs, each, six drachms.
Cow's milk, sixteen ounces.

Employed as a resolvent in certain cases of angina. *Pringle*

INJECTION OF AMMONIA.

R. Water of ammonia, one scruple.
Cow's milk, one ounce.

Mix. As an emmenagogue this injection is to be used three times a day. It induces a mucous flow, often followed by the appearance of the catamenia. It must be used with great caution, and made less stimulating at first.

Lavagna

MIXTURE OF AMMONIA.

R. Water of ammonia,
Carbonate of potassa,
each, two drachms.
Cinnamon water, four fl. ounces.

Mix. A spoonful every two or three hours, in cardialgia and acidity of the stomach.

Jourdan.

MIXTURE OF AMMONIA AND ETHER.

R. Water of ammonia,
Sulphuric ether, each, two fl. drachms.
Peppermint water, four fl. ounces.
Syrup of cinnamon, one fl. ounce.

Mix. A spoonful every hour. Said to be very useful in bites of venomous snakes. *Radius*

AMMONIACAL SINAPISM.

R. Flour of mustard, one ounce and a half
Rye flour, half an ounce.
Water of ammonia, sufficient

to form a cataplasm.

This is more active than the simple mustard poultice.

Phæbus.

LOTION OF AMMONIA.

R. Water of ammonia,
Spirit of thyme,
Spirit of camphor, equal parts.

Mix. As a lotion to the forehead, or applied to the temples, in compresses, in sick headache.

Swedjaur.

FETID SPIRIT OF AMMONIA.

R. Assafetida, in small pieces, one ounce
and a half.

Rectified spirit, one pint and a half.

Macerate for twenty-four hours, then distil off the spirit, and mix the product with

Stronger solution of ammonia, three
fl. ounces.

Dub. Ph.

Employed as a stimulant and antispasmodic in doses of half a drachm.

STEER'S OPODELDOC.

R. Alcohol, eight pints.
Soap, twenty ounces.
Water of ammonia, four fl. ounces.
Camphor, eight ounces.
Oil of rosemary,
" monarda, each, one fl. ounce.

Dissolve the soap in the alcohol with a gentle heat, add the other articles, suffer the impurities to subside, and pour into phials whilst warm.

Phil. Coll. Pharm.

LIQUID OPODELDOC.

R. Dry Spanish soap, sixty parts.
Camphor, fifteen parts.
Alcohol, five hundred parts.
Oil of thyme, four parts.
" rosemary, three parts.
Solution of ammonia, thirty parts.

Mix, dissolve, and filter.

Giseke.

PLASTER OF AMMONIA.

R. Stronger water of ammonia, one ounce.
Camphor, two ounces.
Opium, fifteen grains.
Ammoniac, }
Saffron, } each, half an
Gum plaster, } ounce.
Galbanum plaster, }

Mix well. For corns. A thick layer of this plaster is to be spread on a piece of linen or kid-skin, of the exact size of the corn, as it will blister the sound skin.

La Foret.

SULPHURETTED AMMONIACAL LINIMENT.

R. Water of ammonia, one ounce.
Camphor, half a drachm.
Sulphuret of potassium, four drachms.
Olive oil, six ounces.

Mix. Recommended as an application in tinea capitis.

Bories

TEREBINTHINATE AMMONIACAL LINIMENT.

R. Oil of chamomile, }
" henbane, } each, two
Essence of turpentine, } fl. ounces.
Water of ammonia,
Tincture of capsicum, each,
Camphor, one ounce.
 half an ounce.

Mix well. A very stimulating liniment, found beneficial in asthenic gout and chronic rheumatism.

Ammon.

SPIRIT OF AMMONIA.

R. Alcohol, twenty fl. ounces.
Quicklime, one pound.
Muriate of ammonia, one pound.
Water, nine fl. ounces

Prepare as directed for water of ammonia.

U. S. Ph.

AROMATIC SPIRIT OF AMMONIA.

R. Muriate of ammonia, five ounces.
Carbonate of potassa, eight ounces.
Bruised cinnamon,
" cloves, each, two drachms.
Lemon peel, four ounces.
Alcohol,
Water, each, five pints.

Mix and distil seven pints and a half.

U. S. Ph.

Both of these spirits are stimulant and antispasmodic, and are given in doses of ten to thirty drops in sweetened water.

AMMONIÆ ACETAS.

ACETATE OF AMMONIA.

SOLUTION OF ACETATE OF AMMONIA.

R. Diluted acetic acid, two pints.
Carbonate of ammonia, a sufficient quantity.

Add the carbonate gradually to the acid, stirring constantly until all effervescence ceases.

U. S. Ph.

A valuable diaphoretic in febrile and inflammatory diseases, when aided by warmth; otherwise it acts as a diuretic. Dose, half a fl. ounce to a fl. ounce every three or four hours. It is also a good refrigerant lotion. It is known as the *spirit of Mindererus*.

CATAPLASM WITH ACETATE OF AMMONIA.

R. Extract of hemlock, one ounce.
 " of henbane, half an ounce.
 Powder of belladonna, one drachm.
 Solution of acetate of ammonia, sufficient
 to form a cataplasm. Has been found useful
 in ulcerated cancerous sores. *Radius.*

COLLYRIUM OF ACETATE OF AMMONIA.

R. Solution of acetate of ammonia,
 Rose water, equal parts.
 In chronic ophthalmia. *Saunders.*

GARGLE OF ACETATE OF AMMONIA.

R. Solution of acetate of ammonia,
 Honey of roses, each, one fl. ounce.
 Elder water, eight fl. ounces.
 Mix. Recommended by Wendt in scarlatina
 anginosa with ulcerations. *Phæbus.*

MIXTURE OF ACETATE OF AMMONIA.

R. Solution of acetate of
 ammonia, two fl. ounces.
 Cinnamon water, one fl. ounce.
 Antimonial wine, one fl. drachm.
 Distilled water, two fl. ounces.
 Mix. A tablespoonful every two or three hours.
Ellis.

R. Camphor mixture,
 Solution of acetate of
 ammonia, each, half a fl. ounce.
 Antimonial wine,
 Laudanum, each, twenty drops.

Mix. To be taken at bedtime, or oftener, in
 which latter case the laudanum to be diminished.
Ellis.

R. Liquid acetate of
 ammonia, half fl. ounce.
 Camphor mixture, seven fl. drachms.
 Antimonial wine, fifteen drops.
 Syrup of saffron, half a fl. drachm.

Mix. To be taken three times a-day. *Burke.*

AMMONIÆ ARSENIAS.

ARSENATE OF AMMONIA.

R. Arsenic acid, one ounce.
 Stronger water of ammonia, sufficient

to saturate the acid. To be left to evaporate and
 crystallize spontaneously. Dose, $\frac{1}{20}$ th to $\frac{1}{12}$ th
 of a grain. *Cottrean.*

SOLUTION OF ARSENATE OF AMMONIA.

R. Arsenate of ammonia, eight grains.
 Water, eight ounces.
 Angelica water, four drachms.

Mix. Has been recommended, in doses of
 twenty to thirty drops, in obstinate cutaneous
 affections. *Soubeiran.*

AMMONIÆ CARBONAS.

CARBONATE OF AMMONIA.

R. Muriate of ammonia, one pound.
 Dried chalk, one pound and a half.
 Pulverize them separately, then mix them
 thoroughly, and sublime into a cooled receiver.
U. S. Ph.

AROMATIC CARBONATE OF AMMONIA.

R. Carbonate of ammonia, one pound.
 Oil of lavender, three ounces.
 " verbena, half an ounce.
 Grind together, and sublime with a gentle
 heat.

R. Carbonate of ammonia, half an ounce.
 Oil of peppermint,
 " cajeput, each, twelve drops.

Mix, and keep in a closed bottle. *Phæbus.*

R. Powd. carb. ammon., one ounce.
 Stronger solution of
 ammonia, half a fl. ounce.

Oil of rosemary,
 Oil of bergamot, each, ten drops.

Mix, and while moist, put in a wide-mouthed
 bottle, which is to be well closed.

All these are used as smelling salts in faint-
 ness, &c.

POTION OF CARBONATE OF AMMONIA.

R. Carbonate of ammonia, ten grains.
 Cinnamon water, one fl. ounce.
 Tincture of allspice, half a fl. drachm.
 Syrup of saffron, half a fl. ounce.

Mix. To be taken before going to bed.
Phæbus.

R. Comp. tinct. of
 ammonia (*Lond.*), two fl. drachms
 and a half.

Camphor water, seven fl. ounces.
 Tincture of castor, one fl. drachm.
 Simple syrup, three fl. drachms.

Mix. Two spoonfuls in an hysterical attack.
Ainslie.

PILLS OF CARBONATE OF AMMONIA.

R. Carbonate of ammonia, }
 Powdered capsicum, } each, one
 " cloves, } scruple.
 " mace, }
 Oil of caraway, five drops.
 Extract of gentian, twelve grains.
 Simple syrup, sufficient
 to form twenty pills. One every two hours in
 gout in the stomach. *Parrish.*

MIXTURE OF CARBONATE OF AMMONIA.

R. Carbonate of ammonia, }
 Powdered white } each, one drachm
 sugar, } and a half.
 Pow. gum Arabic, }
 Compound spirit of
 lavender, two fl. drachms.
 Mint water, four fl. ounces.

Mix. A tablespoonful every two or three hours.
 If required, forty to fifty drops of laudanum
 may be added to the mixture. *Ellis.*

Useful in low states of the system.

DRAUGHT WITH CARBONATE OF AMMONIA.

R. Oil of valerian, three drops.
 Carbonate of ammonia, ten grains.
 Cinnamon water, two fl. ounces.

Mix. To be taken every four hours, in depression
 of the spirits and nervous headache. *Ellis.*

MIXTURE OF CARBONATE OF AMMONIA
AND GINGER.

R. Powdered ginger, fifteen grains.
 Carbonate of ammonia, eight grains.
 Spirit of cinnamon, two fl. drachms.
 Water, one and a half fl. ounces.

Mix. To be taken in gout or cramp in the
 stomach. *Ellis.*

DROPS OF CARBONATE OF AMMONIA.

R. Carbonate of ammonia, twenty parts.
 Oil of lavender, one part.
 Alcohol, four parts.

Distil. Dose, thirty or forty drops in nervous
 headache. *Cottureau.*

OINTMENT OF CARBONATE OF AMMONIA.

R. Carbonate of ammonia, one part.
 Simple cerate, eight parts.

Mix. Used in frictions to the throat in croup.
Foy.

PLASTER OF CARBONATE OF AMMONIA.

R. Carbonate of ammonia,
 Opium, each, fifteen grains.

Camphor, half a drachm.
 Oil of cajeput, twenty drops.
 Galbanum plaster, half an ounce.

Mix. Said to be useful as an application to the
 pit of the stomach to arrest vomiting in sea-
 sickness. *Phœbus.*

CAMPHORATED AMMONIACAL LINI-
MENT.

R. Liquid carbonate of
 ammonia, four parts.
 Oil of olives, sixteen parts.
 Spirit of camphor, three parts.

Mix. As a revulsive embrocation. *Swediaur.*

AMMONIACAL OINTMENT.

R. Carbonate of ammonia, one drachm.
 Rose ointment, one ounce.
 Oil of jasmine, four drops.

Mix. Recommended as a friction, three or
 four times a day, in engorgements of the
 mammae. *Augustin.*

GONDRET'S AMMONIACAL OINTMENT.

R. Suet,
 Hog's lard, each, one part.
 Stronger water of ammonia, two parts.

Melt the fats and incorporate the ammonia.
Soubeiran.

R. Hog's lard, seven drachms.
 Oil of almonds, a drachm and a half.
 Stronger water of
 ammonia, five drachms.

Melt the lard, mix it with the oil, pour into a
 wide-mouthed bottle with a glass stopper, add
 the ammonia, and agitate well. Keep in a cool
 place.

Rubbed on the skin it causes rubefaction,
 and, if covered with a compress, speedily vesi-
 cates. A useful rubefacient and counter-irritant.
Gondret.

COMPOUND TINCTURE OF AMMONIA.

R. Mastic, two drachms.
 Alcohol, nine fl. drachms.
 Oil of lavender, fourteen minims.
 Stronger water of ammonia, one pint.

Dissolve the mastic in the alcohol, and filter;
 add the other ingredients, and agitate them
 well.

Well known under the name of *eau de luce*
 as a remedy in bites of venomous reptiles. It
 is a good antispasmodic, in doses of from ten to
 forty drops; and also forms a stimulating em-
 brocation. *Lond. Ph.*

AROMATIC AMMONIATED ALCOHOL.

R. Ammoniated alcohol,
 (spirit of ammonia), one pint.

Oil of rosemary,
 " lemons, each, two fl. drachms.
 " cloves,
 " cinnamon, each, half a
 fl. drachm.

Mix the oils, add the ammoniated alcohol, and pour in as much water as will prevent empyreuma in distilling; distil a pint with a gentle heat. *U. S. Ph., 1830.*

MIXTURE OF AMMONIATED ALCOHOL.

R. Ammoniated alcohol,
 (spirit of ammonia), half a fl.
 ounce.
 Oil of amber, one drop.
 " mace, two drops.
 " aniseed, }
 " cloves, } each, three drops.
 " cinnamon, }

Mix. Dose, ten to thirty drops in sweetened gum water, in anorexia and flatus.

Ammon.

ANISATED AMMONIATED ALCOHOL.

R. Alcohol, twenty-four parts.
 Oil of aniseed, one part.
 Dissolve, and gradually add, constantly stirring,
 Spirit of ammonia, six parts.
 Keep in well-closed bottles. *Jourdan.*

LOTION IN HICCUGH.

R. Anisated ammoniated alcohol,
 one fl. drachm.
 Spirit of lavender, two fl. ounces.
 Recommended by Siebold to be rubbed on the thorax and abdomen of infants to check hiccough. *Radius.*

MIXTURE OF ANISATED AMMONIATED ALCOHOL.

R. Extract of liquorice, one drachm.
 Barley water, three fl. ounces and
 a-half.
 Anisated ammoniated alcohol, half a
 fl. drachm.
 Simple syrup, half a fl. ounce.
 Mix. A teaspoonful occasionally in chronic catarrh. *Hufeland.*
 R. Anisated ammoniated alcohol,
 one fl. drachm.
 Syrup of mallows, one fl. ounce.
 Extract of henbane, one scruple.

Mix well. A dessertspoonful three times a day, in chronic bronchitis or in the second stages of whooping-cough. *Radius.*

VICAT'S ANODYNE MIXTURE.

R. Ammoniated alcohol, half a fl. ounce.
 Diluted alcohol, one fl. ounce.
 Opium, two scruples.
 Camphor, one scruple.

Digest for three days, often stirring, and filter. Used on cotton to fill the cavity of carious teeth, and as a lotion to the temples in headache. *Spielman.*

AMMONIÆ BICARBONAS.

BICARBONATE OF AMMONIA.

R. Sesquicarbonate of ammonia, at will.
 Pulverize, spread out on paper, and expose to the air for twenty-four hours. Then enclose in well-stopped bottles. *Dub. Ph.*
 This is a white powder of less disagreeable taste than the common carbonate. It is used for the same purposes. Dose, five to fifteen grains.

AMMONIÆ CITRAS.

CITRATE OF AMMONIA.

R. Fresh lemon juice, one fl. ounce.
 Carbonate of ammonia, sufficient
 to saturate.
 White sugar, one drachm.
 Distilled water, four fl. ounces.

Mix.

A pleasant saline diaphoretic in febrile disorders, resembling the solution of the acetate of ammonia in its effects. Dose, a tablespoonful, as may be required. *Ellis.*

SOLUTION OF CITRATE OF AMMONIA.

R. Citric acid, three ounces.
 Water, [Imp.] one pint.
 Dissolve, and add
 Sesquicarbonate of ammonia,
 two ounces and a half.
 Or sufficient to saturate. *Lond. Ph.*

AMMONIÆ HYDRIODAS.

HYDRIODATE OF AMMONIA.

R. Iodine, one hundred parts,
 Iron filings, fifty parts.
 Water, five hundred parts.

Mix, and heat till the liquid becomes almost colorless; filter, and add carbonate of ammonia, as long as any precipitate takes place, filter, evaporate to dryness, redissolve in four or five times its weight of water; filter, evaporate and crystallize. *Beasley.*

OINTMENT OF HYDRIODATE OF
AMMONIA.

R. Hydriodate of ammonia, one scruple.
Lard, one ounce.
Mix. *Beasley.*

R. Hydriodate of ammonia, one drachm.
Lard, one ounce.
Mix. *Gibert.*

As an application to scrofulous tumors and cutaneous affections.

AMMONIÆ HYDROSULPHAS.

HYDROSULPHATE OF AMMONIA.

R. Water of ammonia, four fl. ounces.

Pass hydrosulphuric acid (obtained from the sulphuret of iron, and sulphuric acid diluted with four times its weight of water), through the water of ammonia in a suitable apparatus, to perfect saturation. To be kept in a well-stopped bottle. Sp. gr. 0.999. *Dub. Ph.*

A powerful sedative, lessening the action of the circulatory system to a great degree, and causing nausea, vomiting, vertigo, &c.

Used in diabetes to check the morbid appetite, by Cruikshank, Rollo, and others. Dose, five or six drops, in a tumbler full of water, three or four times a day, gradually increasing the dose until some vertigo is induced.

SULPHURETTED HYDROSULPHATE OF
AMMONIA.

R. Quicklime, three parts.
Sulphur, each, three parts.

Triturate together, adding water sufficient to form a paste, and incorporate

Sulphate of ammonia, seven parts dissolved in water. Let stand, decant, wash the residuum, rubbing it with a small portion of water, unite the solutions, and filter.

Van. Mons.

Known as *Boyle's fuming liquor*, and recommended in venereal complaints, in phthisis, rheumatism, and gout, in doses of three to four drops in water, augmenting the dose until vertigo ensues.

WILKINSON'S LINIMENT.

R. Boyle's fuming liquor, ten grains.
Chalk, one scruple.
Flowers of sulphur,
Lard,
Tar, each, half an ounce.

Mix well. Advised in chronic diseases of the skin. To be rubbed on a moderate extent of the surface at a time. *Phæbus.*

SULPHURETTED SYRUP.

R. Boyle's fuming liquor, three drachms.
Sulphuret of antimony

and soda, one drachm.

Simple syrup, an ounce and a half.

Mix. Has been prescribed in croup, in the dose of a teaspoonful every hour. *Swediaur.*

AMMONIÆ MURIAS.

SAL AMMONIAC.

POWDER OF SAL AMMONIAC.

R. Sal ammoniac, two drachms.
Camphor, six grains.
Powdered arnica, half a drachm.
White sugar, six drachms.

Mix. Dose, a teaspoonful, three or four times a day. Said to be efficacious as an emmenagogue, and also in smaller doses in chronic catarrh. *Radius.*

BOLUS OF SAL AMMONIAC.

R. Sal ammoniac,
Extract of liquorice, each, one scruple.
Extract of taraxacum, sufficient

to form a bolus. One to be taken every three or four hours, in schirrhous of the prostate, bladder, or rectum. *Radius.*

LOTIONS OF SAL AMMONIAC.

R. Sal ammoniac, two scruples.
Vinegar, one fl. ounce.
Water, one pint.

Mix. *Augustin.*

R. Sal ammoniac, one ounce.
Vinegar,
Alcohol, each, four fl. ounces.

Mix. In strains, bruises, and external inflammation. *Foy.*

FOMENTATION OF SAL AMMONIAC.

R. Powdered bistort,
" pomegranate root,
each, two ounces.
Red wine, one pound.

Heat, digest for an hour, strain, and add

Sal ammoniac, two drachms.

Useful as a fomentation, in strains and bruises. *Foy.*

R. Sal ammoniac, one ounce.
Spirit of rosemary, one pound.

Mix. Said to be useful in engorgement of the mammae. The breast to be covered with cloths dipped in the fluid. *Foy.*

LINIMENT WITH SAL AMMONIAC.

R. Sal ammoniac, one ounce.
 Camphor, half an ounce.
 White soap, six drachms.
 Alcohol, two pounds.
 Digest, and filter. Useful as an application to
 contusions. *Niemann.*

COLLUTORIUM OF SAL AMMONIAC.

R. Pellitory, two drachms.
 Lavender water,
 Vinegar, each, two fl. ounces.
 Opium, two grains.
 Sal ammoniac, one drachm.
 Macerate for a few days, and filter.
Cadet de Gassicourt.

DRAUGHT OF SAL AMMONIAC.

R. Sal ammoniac, five grains.
 Camphor, three grains.
 Gum Arabic, one drachm.
 Parsley water, four fl. ounces.
 Mix. To be taken every two hours in paralytic
 retention of urine. *Radius.*

CATAPLASM WITH SAL AMMONIAC.

R. Sal ammoniac,
 Henbane, each, half an ounce.
 Linseed meal, an ounce and a half.
 Boiling water, sufficient
 to make a cataplasm. Advised in an engorge-
 ment of milk in the breasts. *Radius.*

WASH OF SAL AMMONIAC.

R. Bran, an ounce and a half.
 Mallow leaves, six drachms.
 Water, sufficient
 to make sixteen ounces of solution. Boil, strain,
 and add to decoction
 Sal ammoniac, two drachms.
 Tincture of benzoin, half a fl. ounce.
Radius.

EMULSION OF SAL AMMONIAC.

R. Sweet almonds, one ounce.
 Orange-flower water, two fl. ounces.
 Rose water, eight fl. ounces.
 Sal ammoniac, one drachm.
 Tincture of benzoin, two fl. drachms.
 Rub down the almonds, previously blanched,
 with the rose water, and add the other articles.
 Dose, a tablespoonful. *Phæbus.*

AMMONIÆ NITRAS.

NITRATE OF AMMONIA.

R. Diluted nitric acid, at will.
 Carbonate of ammonia, sufficient
 to saturate. Evaporate by a gentle heat, till a
 pellicle forms; set aside to crystallize. *Pereira.*
 Soluble in two parts of cold water. In doses
 of one to two scruples, it reduces the frequency
 of the pulse and heat of the skin, and also acts
 as a diuretic.

MIXTURE OF NITRATE OF AMMONIA.

R. Carbonate of ammonia, one scruple.
 Nitric acid, sufficient
 to saturate; add
 Syrup of mallows, one fl. ounce.
 Distilled water, two fl. ounces.
 Dose, a dessertspoonful, every two hours.
Radius.

AMMONIÆ PHOSPHAS.

PHOSPHATE OF AMMONIA.

R. Phosphoric acid, at will.
 Saturate with
 Carbonate of ammonia,
 and set by to crystallize. *Niemann.*
 Diuretic and discutient. Recommended by
 Dr. Buckler, of Baltimore, in gout and rheuma-
 tism, as a solvent of uric acid calculus, and in
 diseases dependent on a lithic acid diathesis.
 Dose, twenty to thirty drops of the saturated
 solution, three times a-day.
 R. Powdered calcined bone, ten pounds.
 Sulphuric acid, six pounds.
 Mix in a stone-ware vessel, adding one gallon
 of water, digest for three or four days, frequently
 stirring, add a gallon of boiling water, strain,
 adding more boiling water, till it passes almost
 tasteless. Concentrate to one gallon, saturate
 with carbonate of ammonia, filter, concentrate
 by gentle heat, not above 100° F., and let crys-
 tallize. After removing each crop of crystals,
 add more ammonia, and concentrate. *C. Ellis.*

SOLUTION OF PHOSPHATE OF AMMONIA.

R. Phosphate of ammonia, half an ounce.
 Water, six fl. ounces.
 Dissolve. Dose, a tablespoonful three times a
 day. *C. Ellis.*

AMMONIÆ SUCCINAS.

SUCCINATE OF AMMONIA.

R. Succinic acid, one part
 Water, four parts

Put in a capsule, and add sufficient water of ammonia, till perfectly saturated, with a slight excess of alkali; filter, evaporate, and crystallize.
Guibourt.

IMPURE SUCCINATE OF AMMONIA.

R. Spirits of hartshorn, at will.
Saturate with a solution of succinic acid, filter, evaporate, and crystallize.
Bavan Ph.

SPIRIT OF SUCCINATE OF AMMONIA.

R. Spirit of hartshorn, two fl. ounces.
Sal ammoniac, four ounces.
Carbonate of potassa, twelve ounces.
Powdered amber, six ounces.
Oil of lemons, two fl. drachms.

Mix the sal ammoniac, oil, and amber, add the carbonate of potassa to three pints of alcohol, mixed with a pint of water, and distil on a sand-bath. Dose, thirty to fifty drops, as an antispasmodic.
Niemann.

R. Mastich, two drachms.
Alcohol, nine fl. drachms.

Dissolve, decant, and add
Oil of lavender, fourteen minims.
Oil of amber, four minims.

Stronger solution of ammonia, one pint.
Mix. *Lond. Ph.*

This is the old "*eau de luce*," a powerful stimulant in cases of fainting.

MIXTURE OF SUCCINATE OF AMMONIA.

R. Liquid succinate of ammonia,
Elixir of liquorice, each, two parts.
Wine of opium,
Antimonial wine, each, one part.

Mix. Useful in hooping cough, in doses of ten to forty drops, every two hours.
Augustin.

SPIRIT OF AMMONIA AND SUCCINIC ACID.

R. Succinic acid, one part.
Hot water, eight parts.

Saturate the solution with carbonate of ammonia, and filter. Antispasmodic, sudorific.
Van Mons.

AMMON. SUCCINIC ACID, AND ETHER.

R. Ammon. succinic acid,
Sulphuric ether, equal parts.

Mix. Much praised in gout and rheumatism. Dose, twenty to forty drops in sugar and water, two or three times a-day.
St. Marie.

MIXTURE OF SUCCINATE OF AMMONIA.

R. Liquid succinate of ammonia, five fl. drachms.
Tincture of castor,
Anisated ammon. alcohol, each, three fl. drachms.
Tincture of mace, two fl. drachms.
" nutmeg, one fl. drachm.

Mix. Fifteen to twenty-five drops, three or four times a-day, in hysteria.
Augustin.

LINIMENT OF SUCCINATE OF AMMONIA.

R. Liquid succinate of ammonia, one fl. ounce.
Camphorated oil, two fl. ounces.

Mix. As a liniment to the neck in angina.
St. Marie.

R. Camphor, two scruples.
Oil of turpentine, two drachms.

Dissolve, and add
Soft-soap, one ounce.
Ointment of bayberries, two drachms.
Succinate of ammonia, fifteen grains.

Mix. As a friction in chronic rheumatism, especially sciatica.
Cadet.

AMMONIÆ SULPHAS.

SULPHATE OF AMMONIA.

R. Diluted sulphuric acid, at will.
Carbonate of ammonia, sufficient to saturate; filter, and crystallize. *Soubéiran.*
Laxative and stimulant. Dose, from a scruple to half a drachm.

AMMONIACUM.

AMMONIAC.

This gum-resin is found in the shops, either in whitish or yellowish tears, or in yellowish or brownish masses. It has an unpleasant odor, and a nauseous and somewhat bitter taste. It is the product of *Ferula tingitana*, Linn.; and of *Dorema ammoniacum*, Don.

Pereira, Mat. Med. ii. 489. Griffith, Med. Bot. 325, 331.

Its effects on the system are somewhat like those of assafetida, and appear to be most marked on the bronchial mucous membrane. It also is considered to be emmenagogue. The dose is from six to twelve grains. It is employed externally as a resolvent.

AMMONIAC PILLS.

R. Ammoniac, one drachm.
Syrup of gum, sufficient to make thirty-six pills.
Soubéiran.

- R. Ammoniac,
Myrrh, each, two drachms.
Extract of horehound, one drachm.
" liquorice, three drachms.

Mix. Make two grain pills. Dose, four to ten every three hours. *Augustin.*

- R. Ammoniac, one ounce.
Sagapenum, two ounces.
Elixir proprietatis, sufficient

to make four grain pills. As a remedy in obstructed menstruation. *Bories.*

- R. Ammoniac, two drachms.
Soap, one drachm.
Oil of juniper, twenty drops.

Mix, and form pills of two grains. *Augustin.*

KLEIN'S PILLS.

- R. Ammoniac,
Extract of centaury,
each, half an ounce.
Soap, one ounce.
Oil of amber, ten drops.

Triturate, and make pills of two grains. *Spielmann.*

PILLS OF AMMONIAC AND RHUBARB.

- R. Ammoniac,
Soap,
Rhubarb, each, two drachms.

Triturate, and make with water three grain pills. Dose, two pills, three to four times a-day. *Tissot.*

COMPOUND AMMONIAC PILL.

- R. Ammoniac, one drachm.
Blue pill, fifteen grains.
Powdered squill, six grains.
Simple syrup, sufficient.

Mix, and make sixteen pills. One, three times a-day, in asthmatic cough with hepatic derangement. *Ainslie.*

DE HAEN'S PILLS.

- R. Ammoniac,
Pil. aloes and myrrh,
each, one drachm.
Soap,
Extract of conium, each, a drachm and a half.

Triturate, and form into three grain pills. Dose, four a day, in chlorosis and amenorrhœa. *St. Marie.*

AMMONIAC MIXTURE.

- R. Ammoniac, two drachms.
Water, half a pint.

Rub the ammoniac with the water gradually added, until they are thoroughly mixed. *U. S. Ph.*

The dose is from one to two tablespoonfuls.

COMPOUND AMMONIAC MIXTURE.

- R. Ammoniac, one drachm.
Distilled water, four fl. ounces.

Make a mixture, and add

- Oxymel of squills, two fl. ounces.
Paregoric elixir, half a fl. ounce.

Mix. A tablespoonful every two hours, as a stimulating expectorant. *Ellis.*

- R. Ammoniac mixture, five fl. ounces.
Oxymel of squills, half fl. ounce.
Antimonial wine, twenty-six minims.
Distilled vinegar, three fl. drachms.

Mix. A tablespoonful occasionally in cough, or humoral asthma. *Ainslie.*

- R. Ammoniac,
Assafetida,
Soap,
Powdered valerian,
Flowers of arnica,
Tartar emetic, } each,
two drachms.
eighteen grains.

Mix, and make two-grain pills. In doses of four or five, three times a-day. In amaurosis, arising from abdominal affections. *Richter.*

AMMONIAC AND NITRIC ACID MIXTURE.

- R. Ammoniac, two and a half scruples.
Diluted nitric acid, two fl. drachms.
Water, eight fl. ounces.

Add the acid to the water, then gradually rub the gum with the acid compound, in a composition mortar.

A tablespoonful, two or three times a day. It is highly beneficial in chronic catarrh, especially in that of elderly persons. *Ellis.*

MIXTURES OF AMMONIAC.

- R. Ammoniac mixture, three fl. ounces.
Tincture of castor, half a fl. drachm.
Syrup of tolu, half a fl. ounce.
Laudanum, twenty to thirty drops.
Cinnamon water, one fl. ounce.

Mix. A dessertspoonful every three or four hours in troublesome cough. *Meigs.*

- R. Ammoniac,
Gum Arabic, each, half an ounce
Infusion of hyssop, six fl. ounces.
Syrup of capillaire (maidenhair),
two fl. ounces

Rub the ammoniac and gum with the infusion until a solution is formed.

In tablespoonful doses, occasionally, in catarrhs.
Radiis.

R. Ammoniac mixture, four fl. ounces.
Wine of ammoniac,
Paregoric elixir, each, half fl. ounce.
Syrup of tolu, one fl. ounce.

Mix. In teaspoonful doses, occasionally, as an expectorant.

AMMONIAC PLASTER.

R. Ammoniac, five ounces.
Diluted acetic acid, half a pint.

Dissolve the ammoniac in the acid and strain; evaporate in a water-bath to a proper consistence.
U. S. Ph.

Useful as an application to indolent and scrofulous tumors to promote suppuration.

PLASTER OF AMMONIAC WITH MERCURY.

R. Ammoniac, one pound.
Mercury, three ounces.
Olive oil, one fl. drachm.
Sulphur, eight grains.

To the heated oil, add the sulphur gradually, stirring constantly, then rub the mercury with the compound, until the globules disappear; lastly, add the ammoniac previously melted and strained, and with a gentle heat rub them well together.
U. S. Ph.

More active than the simple ammoniac plaster. Used as a discutient to tumid glands and other indolent swellings, especially syphilitic. Sometimes salivates.

GUM PLASTER.

R. Litharge plaster, four ounces.
Ammoniac,
Galbanum,
Beeswax, each, half an ounce.

Melt the gum resins together and strain; melt also the plaster and wax; add the two mixtures together and thoroughly incorporate. *Edin. Ph.*

A good digestive and suppurative for indolent tumors

AMMONIAC PLASTER WITH HEMLOCK.

R. Strained ammoniac, three ounces.
Extract of hemlock, two drachms.

Melt, and add
Solution subacetate of lead, one drachm.

Stir till cold. *Ph. Chem.*

AMYGDALA.

ALMONDS.

AMYGDALA AMARA.

BITTER ALMONDS.

AMYGDALA DULCIS.

SWEET ALMONDS.

The almond-tree, *Amygdalus communis*, is about the size and much resembles the peach-tree in appearance. It is a native of Persia and Syria, and is extensively cultivated in the southern parts of Europe. There are several varieties, the most important of which are the sweet and the bitter: the former having a sweet, bland kernel; and the latter, one that has the bitter taste of the peach kernel.

Sex. Syst. Icosand. monog. *Nat. Syst.* Amygdalaceæ.

Linn. Sp. Pl. 677. Griffith, Med. Bot. 284.

Sweet almonds act as a demulcent; but the bitter have some of the properties of hydrocyanic acid.

BUTTER OF ALMONDS.

R. Blanched almonds, ten drachms.
White sugar, twelve drachms.
Orange-flower water, two drachms.

Triturate till reduced to a homogeneous paste.
Guibourt.

A spoonful, stirred in a tumbler of water, forms an extemporaneous syrup of orgeat, which latter preparation is difficult to preserve.

ALMOND PASTE.

R. Blanched almonds, eight ounces.
Oil of almonds,
Lemon juice, each, a sufficient quantity.

Diluted alcohol, five ounces.

Form a paste. *Taddei.*

R. Blanched almonds,
Rice flour, each, eight ounces.
Bean flour, sixteen ounces.
Myrrh, two ounces.
Camphor, one ounce and a half.
White soap, twenty ounces.

Form a paste. *Spielmann.*

Used as a cosmetic.

ALMOND POWDER.

R. Blanched almonds, two pounds.
Rice flour,
Orris root, each, four ounces.
Benzoin,
Carbonate of potass,
Spermaceti, each, one ounce.

Oil of rhodium,
 " lavender,
 " cloves, each, thirty drops.

Form a powder. Used as a cosmetic.

ALMOND EMULSIONS.

R. Blanched almonds, one ounce.
 Pound in a marble mortar, and gradually add
 Water, sixteen ounces.
 Strain. *Beral.*

R. Blanched almonds, two ounces.
 White sugar, half an ounce.

Pound in a marble mortar, and add
 Boiling water, twelve fl. ounces.
 Strain, and add
 Orange-flower water, two drachms.

ALMOND CONFECTION.

R. Blanched almonds, one ounce.
 Powdered gum Arabic, one drachm.
 " sugar, half an ounce.

Pound well in a marble mortar till thoroughly mixed. *U. S. Ph., 1830.*

One drachm triturated with an ounce of distilled water forms an excellent emulsion, which is nutritive and emollient.

COMPOUND ALMOND LOTION.

R. Sweet almonds, one ounce.
 Bitter " four drachms.

Form an emulsion with
 Cherry brandy, ten fl. ounces.

Add
 Corrosive sublimate, six grains.
 Tincture of benzoin, six drachms.
 Lemon juice, four drachms.

Mix. As a lotion for pustular eruptions on the face. The mixture should be shaken before it is used. *Siemerling.*

SYRUP OF ORGEAT.

R. Sweet almonds, one pound.
 Bitter " five ounces.
 White sugar, six pounds.
 Orange-flower water, eight fl. ounces.
 Water, three pints.

Blanch the almonds, beat them into a paste with some of the water and the sugar, mix this paste with the rest of the water, press, and strain; dissolve the remainder of the sugar in the emulsion by means of a water-bath, add the orange-flower water, and strain. *Paris Cod.*

SYRUP OF ORGEAT.

R. Sweet almonds, twelve ounces.
 Bitter almonds, four ounces.
 Sugar, six pounds.
 Water, sufficient.
 Orange-flower water, four fl. ounces.

Blanch the almonds, pound thoroughly in a marble mortar (after having soaked them in cold water), mix gradually with one pint and a half of water, and strain with expression. Repeat the process with the same quantity of water, and make up the strained liquor to three pints, in which dissolve the sugar with a gentle heat. When cold, add the orange-flower water and mix. *Ambrose Smith.*

The U. S. Ph. for 1850 directs an almond syrup, which only differs from the above in not containing any orange-flower water. All of these syrups are demulcent and nutritive. But they are chiefly used as flavoring ingredients in cough mixtures.

ARTIFICIAL MILK.

R. Blanched almonds, No. 2.
 Make an emulsion with
 Boiling water, four fl. ounces.
 Cow's milk, six fl. ounces.
 White sugar, one drachm.

Strain. As a substitute for breast-milk for infants. *Rosenstein.*

ANODYNE MILK.

R. Blanched almonds, four ounces.
 Melon seed,
 Squash seed, each, one ounce.
 Lettuce seed,
 Poppy seed, each, one drachm.

Triturate well, adding gradually
 Barley water, sufficient.
 Strain and express; to be taken in glassful doses, to which are to be added two drachms of sugar. *Pierquin.*

MILK OF ROSES.

R. Almonds, one ounce.
 Rose water, five fl. ounces.
 Alcohol, one fl. ounce.
 Soap, half a drachm.
 Attar of roses, two drops.

Blanch the almonds, dry them well with a cloth, beat them to a paste, add the soap, triturate well, then add the rose-water and alcohol, rub well together, and strain through fine linen. This keeps well. *Bateman*

R. Blanched almonds, eight ounces.
 Rose water, three pints.
 Windsor soap,
 White wax,
 Oil of almonds, each, half an ounce

Rectified spirit,	twelve ounces.
Oil of bergamot,	half an ounce.
“ lavender,	one drachm.
Attar of roses,	half a drachm.

Beat the almonds with the rose-water, so as to form an emulsion; melt the soap, wax, and almond oil together by a gentle heat, and rub the mixture with the emulsion; strain; dissolve the essential oils in the spirit, and mix with the strained emulsion. *Redwood.*

GOULARD'S LOTION.

R. Emulsion of bitter almonds,	two hundred parts.
Corrosive sublimate,	
Sal ammoniac, each,	one part.

Foy.

ALMOND CREAM.

R. Marrow pomatum,	
Almond oil, each,	two ounces.

Melt by a gentle heat, and add
Oil of jasmine, two drachms,
stirring till cold. As an application to the hair. *Bateman.*

OIL OF ALMONDS.

The kernels of almonds are to be rubbed and shaken in a bag to remove the reddish dust that adheres to them, then ground in a mill or pounded in a stone mortar. The marc is to be subjected to pressure in a linen bag, and the oil filtered.

COLD CREAM.

R. Rose water,	one fl. ounce.
Oil of almonds,	two fl. ounces.
Spermaceti,	half an ounce.
White wax,	one drachm.

Melt together the oil, spermaceti, and wax, by means of a water-bath, add the rose-water, and stir constantly until the mixture is cold.

U. S. Ph.

ALMOND EMULSION.

R. Sweet almonds,	half an ounce.
Gum Arabic,	half a drachm.
Sugar,	two drachms.
Water,	half a pint.

Blanch the almonds, beat them with the gum and sugar till thoroughly mixed; then rub the mixture with the water gradually added, and strain. *U. S. Ph.*

A good demulcent, in irritation or inflammation of mucous membranes, and may be freely used.

AROMATIC ALMOND EMULSION.

R. Blanched almonds,	
Simple syrup,	

Orange-flower water, each,	one ounce.
Water,	fourteen ounces.

Rub down the almonds with a little of the water, incorporate the paste with the remainder of the fluid, and strain. *Beral.*

GREEN LINCTUS.

R. Pistachio nuts (<i>Pistacia Vera</i>),	No. 14.
Syrup of violets,	one ounce.
Oil of almonds,	half an ounce.
Gum tragacanth,	sixteen grains.
Tincture of saffron,	twenty drops.
Orange-flower water,	two drachms.
Water,	four ounces.

Make an emulsion.

Soubiran.

Employed as a demulcent cough mixture. Dose, a tablespoonful.

EMOLLIENT CLYSTER.

R. Oil of almonds,	
Brown sugar, each,	one ounce.
Barley water,	half a pint.

Mix. *Saunders.*

BITTER ALMOND WATER.

R. Oil of bitter almonds,	sixteen minims.
Carbonate of magnesia,	one drachm.
Water,	two pints.

Rub the oil with the carbonate, then with the water gradually added, and filter through paper.

The full dose is a tablespoonful.

DIURETIC WATER.

R. Bitter almonds,	four ounces.
Dried elder flowers,	six ounces.
Peach kernels,	four ounces.
Cherry kernels,	two ounces.
Wine,	eight pints.

Digest for two days and distil off one-half.

Palat. Ph.

Said to act as a diuretic in doses of half an ounce, but requires to be used with much caution.

OIL OF BITTER ALMONDS.

R. Bitter almonds,	twenty pounds.
Water,	a sufficient quantity.

Pound the almonds, express the oil, let the cake macerate in water for twenty-four hours, distil as long as the product is odorous, separate the oil, redistil the water, and again separate the oil. *Cottureau.*

Very poisonous, especially when fresh; principally employed as a flavoring ingredient, but should always be used with great caution. Dose, half a drop.

LOCKSTADT'S PILLS.

- R. Sulphate of quinine, three grains.
Aromatic powder, ten grains.
Oil of bitter almonds, one drop.
Extract of gentian, sufficient.

Make ten pills. The whole to be taken at a dose, before an expected paroxysm of intermittent fever. *Phabus.*

WHITE LINCTUS.

- R. Blanched sweet almonds, No. 12.
White sugar, six drachms.
Gum tragacanth, twelve grains.
Orange-flower water, one fl. drachm.
Water, four fl. ounces.

Make an emulsion. *Guibourt.*

Used as a demulcent, in catarrhs and bowel-complaints. Dose, a tablespoonful.

- R. Blanched almonds, four drachms.
White sugar, six drachms.
Oil of almonds, three fl. drachms.
Orange-flower water, two fl. drachms.
Gum tragacanth, sixteen grains.
Water, four fl. ounces.

Make an emulsion. *Giordano.*

MIXTURES OF OIL OF ALMONDS.

- R. Oil of almonds,
Simple syrup,
Water, equal parts.

Make an emulsion. *Ratier.*

- R. Oil of almonds, one ounce.
Gum Arabic,
White sugar, each, half an ounce.
Water, eight fl. ounces.

Make an emulsion. *Foy.*

- R. Oil of almonds,
Gum Arabic,
Orange-flower water, each, half an ounce.
Syrup of mallows, one ounce.
Water, three ounces.

Make an emulsion. *Paris Cod.*

- R. Oil of almonds, one ounce.
Yolk of egg, one.
Syrup of mallows, one ounce.
Orange-flower water, two drachms.
Water, two ounces.

Make an emulsion. *Guibourt.*

- R. Oil of almonds,
Syrup of orgeat, each, seven drachms.
Gum Arabic, two drachms.
Bitter almond water, half an ounce.

Make an emulsion. Dose, a teaspoonful. *Radius.*

- R. Oil of sweet almonds,
Syrup of tolu, each, one fl. ounce.
Distilled water, six fl. ounces.
Solution subcarbonate potassa, sufficient.

Make emulsion. A tablespoonful two or three times a day in catarrh. *Ainslie.*

COSMETIC LINIMENT.

- R. Oil of almonds, one ounce.
Balsam of Mecca, one drachm.
Sub. carb. of potassa, half a drachm.
Rose water, four fl. ounces.

Rub the balsam with the oil, add the potassa, triturate for ten minutes, and gradually introduce the rose water.

MACASSAR OIL.

- R. Oil of almonds, at will.
Alkanet root, sufficient
to color.
Oil of cinnamon, sufficient
to scent. *Steph. and Church.*

BANDOLINE.

- R. Oil of almonds, one ounce.
White wax, one drachm.
Tincture of mastich, three drachms.
Oil of bergamot, one drachm.

Melt the oil and wax together by a moderate heat, then add the tincture of mastic and oil of bergamot, and strain.

To fix and retain the hair in form. *Redwood.*

AMYLUM.

STARCH.

MUCILAGE OF STARCH.

- R. Starch, four drachms.
Water, one pint.

Triturate the starch with the water gradually added, then boil for a few minutes. *Lond. Ph.*

Mainly employed as a demulcent injection in irritation of the rectum, &c., and forms a good vehicle for laudanum and other active remedies when given in enemata.

COMPOUND POWDER OF STARCH.

- R. Liquorice root, one ounce.
Starch,
Gum Arabic, }
" tragacanth, } each, two ounces.
Quince seed,
Poppy seed, }

Blanched almonds, four ounces.
 Sugar candy, twelve ounces.
 Triturate well. *Van Mons.*

Has been advised in spitting of blood, catarrh, and diarrhoea, in doses of half a drachm to a drachm.

STARCH LOZENGES.

R. Starch, half an ounce.
 Orris root, each, one ounce.
 Extract of liquorice, half an ounce.
 Saffron, one pound.
 White sugar, one pound.
 Mix, and form into lozenges. *Pideret.*

R. Starch, one ounce.
 Gum Arabic, two ounces.
 White sugar, one pound.
 Benzoic acid, half a drachm.
 Rose water, sufficient
 to form a paste. To be divided into lozenges. *Swediaur.*

These lozenges are somewhat stimulating, and are best suited to chronic catarrhs.

STARCH JELLY.

R. Starch, six ounces.
 Water, six pounds.
 Boil till reduced to three pounds, and add
 Wine, four ounces.
 Lemon juice, one ounce.
 Simple syrup, two ounces.
 Mix well, and permit to cool. *Swediaur.*
 A nourishing diet for convalescents.

MIXTURE OF STARCH AND SUET.

R. Cow's milk, six pounds.
 Mutton suet, two ounces.
 Boil by a gentle heat; constantly stirring, and add
 Starch, a spoonful.
 Boil a short time, and add
 White sugar, a sufficiency.
 Recommended in wineglassfuls in dysentery. *Pringle.*

AMYLI IODIDUM.

IODIDE OF STARCH.

R. Iodine, twenty-four grains.
 Triturate with a little water, and add gradually
 Powdered starch, one ounce.
 Continue the trituration till the mixture assumes a deep and uniform color. *Buchanan.*
 Dose, half a drachm, to be gradually increased to four drachms. Given in cases requiring the use of iodine.

ANEMONE.

ANEMONE.

A genus of perennial herbs with radical leaves. They are all acrid and stimulating. The species most generally employed, are the *A. pratensis* and *A. pulsatilla*, both natives of Europe; but it is probable that the *A. patens* of this country has the same properties.

The *A. pratensis* was at one time in much repute in amaurosis and chronic diseases of the skin.

Sex. Syst. Polyand. polygyn. *Nat. Syst.* Ranunculaceæ.

Lindley, Med. Flor. 2. Griffith, Med. Bot. 80.

EXTRACT OF ANEMONE.

R. Fresh anemone, a sufficient quantity.
 Pound in a mortar with a little water, express the juice, and evaporate in a water-bath. *Beral.*
 The dose is from one grain to four, gradually increasing. It has been advised in doses of half a grain to a grain, in combination with sugar of milk, in hooping cough.

ANEMONE PILLS.

R. Powder of anemone, one drachm.
 Extract of anemone, sufficient.
 Make sixty pills. Dose, one to five, three times a-day, gradually increasing.
 Said to have proved useful in amaurosis.

COLLYRIUM OF ANEMONE.

R. Anemone, three drachms.
 Water, sufficient
 to make six ounces of infusion; add
 Corrosive sublimate, one grain.
 To be dropped in the eye, in opacity of the cornea and incipient cataract. *Radius.*

WATER OF ANEMONE.

R. Fresh anemone, four parts.
 Water, sufficient
 to prevent empyreuma. Distil six parts. *Van Mons.*
 This is the best preparation, as it contains the active principle. It is somewhat caustic, and has been recommended to remove freckles on the skin.

ANGELICA.

ANGELICA.

The only species of this genus that is officinal in our Pharmacopœia, is *A. atro-purpurea*. This possesses much the same properties as the Garden Angelica, and is popularly used in flatulent colic and cardialgia. In Europe, the *A. officinalis* is generally used; and, as both species appear to possess the same properties, they may be indifferently employed.

Sex. Syst. Pentand. monog. *Nat. Syst.* Apiaceæ.

Torrey and Gray, Flor. 1, 64. Griffith, Med. Bot. 322.

COMPOUND TINCTURE OF ANGELICA.

R. Aniseed,
Angelica seed, each, half a pound.
Proof spirit, one gallon.

Water sufficient to prevent empyreuma. Macerate for twenty-four hours, and distil a gallon.

A stimulant, used as a stomachic and carminative. Dose, half a drachm to four drachms. *Dublin Ph.* 1826.

TINCTURE OF ANGELICA.

R. Angelica root, one part.
Alcohol, eight parts.
Water, sixteen parts.
Distil six parts. *Van Mons.*

Employed as a carminative and alexipharmic, in doses of one or two fl. drachms.

WATER OF ANGELICA.

R. Angelica root, one part.
Water, six parts.
Distil three parts. *Cottureau.*

ERRHINE MIXTURE.

R. Water of angelica,
" orange flowers,
" roses, equal parts.
Mix. *Pierquin.*

This has been recommended to destroy the fetid smell in ozæna.

CONSERVE OF ANGELICA.

R. Fresh root of angelica, one part.
Water, eight parts.

Macerate for a few days, boil, clarify the decoction, and add

Sugar, one part and a half.
Cook the root in the syrup, and preserve in this state, or dry. *Giordano.*

ESSENCE OF ANGELICA.

R. Angelica root, one part.
Diluted alcohol, eight parts.
Water, sixteen parts.

Distil off six parts. Stomachic and carminative. *Van Mons.*

ANGUSTURA.

ANGUSTURA BARK.

This is recognized by the U. S. Pharmacopœia as the product of *Galipea officinalis*, but is

also obtained from one or more other species, all natives of South America. It is compact, in flat or rolled, but seldom quilled pieces, of a light gray color externally, and of a yellowish-fawn within. It is very fragile, with a resinous fracture. The odor is faint but disagreeable; the taste is bitter and somewhat aromatic, leaving a sensation of pungency. Its powder is pale yellow.

Sex. Syst. Heptand. monog. *Nat. Syst.* Rutaceæ.

Hancock, Trans. Med. Soc. 1849. Griffith, Med. Bot. 192.

It is a stimulating tonic, in large doses acting on the stomach and bowels. It is more used in chronic diarrhœa, and a debilitated condition of the stomach and bowels, than as a febrifuge. Dose, ten to thirty grains.

INFUSION OF ANGUSTURA.

R. Bruised angustura bark, half an ounce.
Boiling water, one pint.

Macerate for two hours in a covered vessel, and strain. *U. S. Pharm.*

The dose is two fl. ounces, every two to four hours.

TINCTURE OF ANGUSTURA.

R. Powdered angustura bark, four and
a half ounces.

Proof spirit, two pints.
Macerate for fourteen days, then filter. *Ed. Pharm.*

This contains the active principles of the bark, and may be given in the dose of one or two fl. drachms.

ELECTUARY OF ANGUSTURA.

R. Powdered angustura, half an ounce.
" canella, half a drachm.
Honey, a sufficient quantity.

Make an electuary. *Foy.*

This has been employed with benefit in chronic diarrhœa and dysentery, in doses of about a drachm.

MIXTURE OF ANGUSTURA.

R. Infusion of angustura, six fl. ounces.
Cinnamon water, half fl. ounce.
Laudanum, twenty drops.

Mix. Three spoonfuls a day. *Radius.*

ANISUM.

ANISE.

This is the fruit of *Pimpinella anisum*, a perennial plant, cultivated largely in some parts of Europe, and often met with in gardens in this country. The seeds are of a grayish-green color, of an ovate, compressed form, with five

pale, narrow ridges. They are aromatic, and have a pleasant, sweetish taste.

Sex. Syst. Pentand. digyn. *Nat. Syst.* Apiaceæ.

Linn. Sp. Pl. 378. Griffith, Med. Bot. 319.

Anise is an aromatic stimulant, and is used in various forms, in colic and dyspepsia, and to prevent the griping of some purgatives. The oil is officinal, but is almost entirely superseded by that of *Illicium anisatum*, which is nearly identical.

SPIRIT OF ANISEED.

R. Aniseed, bruised, ten ounces.
Proof spirit, one gallon.
Water, two pints.

Mix, and distil a gallon by a gentle fire.

Lond. Pharm. 1836.

R. Oil of anise, three fl. drachms.
Proof spirit, one gallon.

Mix. Dose, a teaspoonful. *Lond. Ph.* 1851.

ESSENCE OF ANISEED.

R. Oil of aniseed, one fl. ounce.
Rectified spirit, nine fl. ounces.

Mix. Dose, twenty to thirty drops, on sugar.

Dub. Ph.

ANISEED WATER.

R. Essence of aniseed, one fl. ounce.
Distilled water, four pints.

Mix with agitation, and filter through paper.

**Dub. Ph.*

Dose, a tablespoonful.

ANISE LOZENGES.

R. White sugar,
Anise water, each, two ounces.

Evaporate to the consistence of honey, and add a mixture of

White sugar, four ounces.
Oil of anise, half a drachm.

Pour on a cold marble slab. *Cottureau.*

Used as a carminative and anti-dyspeptic.

MIXTURE OF ANISEED.

R. Powdered aniseed,
" fennel-seed,
Extract of liquorice, each,
one ounce.

Syrup of mallows, twelve ounces.

Mix. *Hanover Pharm.*

Said to be an excellent carminative, especially for children, in doses of about a teaspoonful.

TINCTURE OF ANISEED.

R. Spirit of aniseed, sixteen ounces.
Elixir proprietatis, one ounce.
Aromatic confection, half an ounce.
Pink flowers, one drachm.

Filter, after sufficient maceration.

Wirttemberg Pharm.

Stomachic, carminative, and pectoral. Dose, one to two drachms.

OIL OF ANISE MIXTURE.

R. Oil of anise, twelve minims.
Sugar, one drachm.
Tincture of ginger, two fl. drachms.
Peppermint water, six fl. ounces.

Mix. Dose, two or three spoonfuls, as a carminative. *Ainslie.*

ANTHEMIS.

CHAMOMILE.

Several species of *Anthemis* are employed in medicine, but the most important and the only one recognized in the U. S. Pharm., is *A. nobilis*. This is an herbaceous perennial, a native of Europe, where it is also extensively cultivated, and is also to be found in gardens in this country. The part used is the flowers; these, as found in the shops, are large, roundish, of a yellowish white color, a peculiar, somewhat balsamic, odor, and a warm, bitter, aromatic taste.

Sex. Syst. Syngen. super. *Nat. Syst.* Astéraceæ.

Linn. Sp. Pl. 1260. Griffith, Med. Bot. 401.

Chamomile is an aromatic bitter tonic, in small doses, but emetic in large ones; the usual mode of administration is in infusion.

INFUSION OF CHAMOMILE.

R. Chamomile, half an ounce.
Boiling water, one pint.
Macerate for ten minutes in a covered vessel, and strain. *U. S. Pharm.*

When cold, it is tonic in doses of one or two fl. ounces, several times a day. As an aid to the operation of emetics, it should be given in a tepid state, and largely.

COMPOUND DECOCTION OF CHAMOMILE.

R. Chamomile, half an ounce.
Fennel seed, two drachms.
Water, one pint.

Boil for a short time, and strain.

Dublin Pharm. 1826.

EXTRACT OF CHAMOMILE.

R. Chamomile, one pound.
Water, one gallon.

Boil down to four pints, filter whilst hot, and evaporate in the vapor-bath to proper consistence. *Edin. Pharm.*

A mild, bitter tonic, in doses of ten to twenty grains.

MIXTURE OF CHAMOMILE.

R. Infusion of chamomile, four fl. ounces.
Syrup of orange-flowers, one fl. ounce.
Ethereal tincture of valerian, one fl. drachm.
Mix. *Foy.*

This has proved useful in hysteria, and other nervous diseases, especially where the stomach is disordered. Dose, a dessertspoonful.

R. Extract of chamomile,
Gum Arabic, each, two drachms.
Chamomile water, five fl. ounces.
Tamarind pulp,
Syrup of chamomile, each, one ounce.

Mix. *Phæbus.*
Said to be useful in diarrhœa, in tablespoonful doses.

SYRUP OF CHAMOMILE.

R. Water of chamomile, two pounds.
Syrup of mallows, half a pound.
Extract of liquorice, one ounce.

Cadet de Gassicourt.

Has been recommended in chronic cough and affections of the lungs, in doses of one or two ounces, to be taken during the day.

R. Coarsely powdered chamomile, one ounce.
Cold water, twelve fl. ounces.
Coarsely powdered sugar, twenty ounces.

Make an infusion by displacement, of the chamomile and water. Remove the residue, and substitute the sugar in its place; on this pour the infusion, till entirely dissolved. Dose, a tablespoonful. *E. Parrish.*

COMPOUND CHAMOMILE PILLS.

R. Assafetida, one scruple and a half.
Extract of chamomile, one drachm.
Powdered rhubarb, one scruple.

Make mass, and divide into thirty pills. Three, twice a day in flatulent dyspepsia.

Ainslie.

ANTIMONIUM.

ANTIMONY.

A metal of a silvery blue color, very brilliant, of a lamellar texture, brittle and easily pulver-

ized. Metallic antimony or Regulus of antimony, is not officinal in our Pharmacopœia, but is the basis of a great number of very important preparations.

ANTIMONII CHLORIDUM

CHLORIDE OF ANTIMONY.

R. Sulphuret of antimony, one part.
Muriatic acid, five parts.

Dissolve by aid of a gentle heat, let stand, decant, evaporate, and then distil almost to dryness. *Soubeyran.*

SOLUTION OF TERCHLORIDE OF ANTIMONY.

R. Precipitated sulphuret of antimony, one pound.
Muriatic acid, four pints.

Add, dissolve by a gentle and increasing heat, and finally boil for fifteen minutes. Strain and boil down to two pints. Sp. gr. 1.470.

Dub. Ph.

The butter of antimony is used as a caustic, to destroy fungous flesh and to cauterize poisoned wounds.

NITRO-MURIATIC OXIDE OF ANTIMONY.

R. Prepared sulphuret of antimony, twenty parts.
Muriatic acid, one hundred parts.
Nitric acid, one part.

Digest the sulphuret with the acids, mixed in a glass, avoiding the fumes, with a gradually increasing heat. Boil for an hour, and pour the liquor, when cooled and filtered, into a gallon of water. Wash the precipitate with plenty of water, till the latter no longer reddens litmus paper, then dry the oxide on bibulous paper.

Dub. Ph., 1826.

R. Liquid chloride of antimony, at will.
Distilled water, sufficient.

Agitate together, wash the precipitate that forms with cold water, and dry by a gentle heat.

Van Mons.

This preparation, which is known as the *Powder of Algaroth*, is of very uncertain action, and is not used, except in making tartar emetic. Dose, two or three grains.

OINTMENT OF CHLORIDE OF ANTIMONY

R. Liquid chloride of antimony,
Corrosive sublimate, each, one drachm.
Powdered savine, two drachms.
Lard, six drachms.

Mix. To destroy venereal excrescences.

Radius.

ANTIMONII SULPHURETUM.

SULPHURET OF ANTIMONY.

The sulphuret or crude antimony is now seldom used as a medical agent, but is the basis of almost all the preparations of that metal. Some forms of exhibition are, however, still in favor, on the continent of Europe.

PRECIPITATED SULPHURET OF ANTIMONY.

- R. Sulphuret of antimony, six ounces.
 Solution of potassa, four pints.
 Distilled water,
 Diluted sulphuric acid, each, a sufficient quantity.

Mix the sulphuret with the solution of potassa and twelve pints of the water, boil by a gentle heat for two hours, frequently stirring, and adding more water, to keep up the original quantity. Strain, while hot; and gradually add the sulphuric acid as long as a precipitate is produced; wash well with hot water, dry, and powder. *U. S. Ph.*

Dose, one to five grains.

PILLS OF SULPHURET OF ANTIMONY.

- R. Sulphuret of antimony, one ounce.
 Guaiacum, two drachms.
 Extract of fumitory, a sufficient quantity.

Mix, and make pills of two grains each.

Radius.

Much employed in Germany, as an alterative in diseases of the skin.

- R. Sulphuret of antimony, half a drachm.
 Extract of opium, five to eight grains.
 " guaiacum, two drachms.

Make fifty pills.

Foy.

Advised in gouty affections; nine pills being given three times a-day.

LOZENGES OF SULPHURET OF ANTIMONY.

- R. White sugar, one hundred and ninety-two parts.
 Sulphuret of antimony, eight parts.
 Mucilage of gum Arabic, one part.

Mix, and make lozenges of twelve grains, each containing half a grain of the sulphuret. *Beral.*

Much employed in Europe, as an alterative in diseases of the skin, in rheumatism, gout, &c.

OINTMENT OF SULPHURET OF ANTIMONY.

- R. White wax, four ounces.
 Olive oil, twelve ounces.
 Melt together, and add

- Powdered charcoal, two ounces.
 Precipitated sulphur,
 Sulphuret of antimony, each, one ounce.

Stir well until cold.

Radius.

Used as an application in tinea capitis.

KERMES MINERAL.

- R. Crystallized carbonate of soda, 125 parts.
 Water, 1280 parts.
 Sulphuret of antimony, 6 parts.

Boil the antimony for an hour in the solution of soda, filter the hot liquor into warm earthen pans, cool very slowly, wash the precipitate with cold water, dry at a moderate temperature, and keep in well-closed bottles. *Paris Cod.*

Dose, one to three grains.

KERMESINE POWDER.

- R. Kermes mineral, two grains.
 Sugar of milk,
 Gum Arabic, each, one drachm.
 Divide into six powders. *Radius.*

Given as a diaphoretic and expectorant, in the dose of one every four hours.

COMPOUND KERMESINE POWDER.

- R. Kermes mineral, half a drachm.
 Ipecacuanha, one grain.
 Prepared chalk,
 Gum Arabic, each, twelve grains.

Mix, and divide into three powders; one to be taken three times a-day, in a spoonful of tea.

St. Marie.

This formula has been much praised in whooping-cough; the proportions are intended for a child of about twelve years of age.

LOZENGES OF KERMES MINERAL.

- R. Kermes mineral,
 Powdered squill, each, sixty grains.
 Extract of opium, one drachm.
 Ipecacuanha, two drachms.
 White sugar, three ounces.
 Mucilage of tragacanth, a sufficient quantity.

Mix, and make four hundred lozenges. *Foy.*

These are expectorant and calming in catarrh. One is to be taken every two hours.

- R. Kermes mineral, one drachm.
 Powdered gum Arabic, eight ounces.
 Extract of opium, twelve grains.
 " liquorice, two ounces.
 Sugar, thirty-two ounces.

Oil of aniseed, six drops.
Mucilage of tragacanth, sufficient.
Mix, and form ten grain lozenges. Known as
Tronchin's lozenges. Used in catarrh.

Van Mons.

R. Benzoic acid, two drachms.
Sugar, thirty-two ounces.
Powdered orris root, four drachms.
" gum Arabic, two ounces.
" starch, four ounces.
Water, four fl. ounces.

Mix, and make fifteen grain lozenges.

Vandamme.

EMULSION WITH KERMES MINERAL.

R. Kermes mineral, ten grains.
Oxymel of squills, one ounce.
Emulsion of gum Arabic, four ounces.

Mix. A tablespoonful occasionally. *Radius.*

MIXTURE WITH KERMES MINERAL.

R. Kermes mineral, two grains.
Gum Arabic, one drachm.
Syrup, four fl. ounces.
" of grapes, one fl. ounce.

Mix. A tablespoonful occasionally in catarrh, when expectoration is difficult. *Pierquin.*

GOLDEN SULPHURET OF ANTIMONY.

R. Mother water of kermes mineral, at will.

Add gradually,
Acetic acid,

till a precipitate is no longer formed. Wash this well in cold water, and dry by a gentle heat.
Dose, one to ten grains. *Paris Cod.*

POWDER OF GOLDEN SULPHURET OF ANTIMONY.

R. Golden sulph. of antimony,
Camphor, each, a grain and a half,
to two grains.

Nitrate of
potassa, half a drachm to one
drachm.

Gum Arabic,
White sugar, each, one drachm.

Triturate well, and divide into six powders.

Radius.

The dose is one, ever two hours, in peripneumonia, after a reduction of the primary symptoms.

R. Golden sulph. of antimony, eight grains.
Opium, two grains.

Powdered Peruvian bark,
" chamomile,
each, four scruples.

Mix, and divide into eight powders; to be taken during the apyrexia, of intermittent fever.

Radius.

MIXTURE OF SULPHURET OF ANTIMONY.

R. Golden sulph. of antimony, six grains.
Extract of lettuce, one scruple.
Syrup of seneka, two fl. ounces.

Mix. A teaspoonful every two hours.

Phæbus.

SULPHURET OF ANTIMONY AND LIME.

R. Golden sulph. of antimony, one part.
Quicklime, three parts.

Triturate together, and add

Boiling water, twenty-four parts.

Dry with a gentle heat, constantly stirring, and keep in a well-closed bottle. *Batav. Ph.*

Emetic and resolvent. Much praised in gout, rheumatism, scrofula, &c. Dose, one to six grains.

PILLS OF SULPHURET OF ANTIMONY AND LIME.

R. Sulphuret of antimony and lime, half a drachm.
Mucilage of tragacanth, one drachm.

Mix, and make sixty pills; five, thrice a-day.

Augustin.

PLASTER OF SULPHURET OF ANTIMONY AND LIME.

R. Yellow wax, one drachm and a half.
Melt, and mix

Pitch, one ounce;

and incorporate in the cooling mixture

Sulphuret of antimony and lime, five scruples.

As a plaster in sciatica, chronic headache, &c. It sometimes occasions pustules.

Radius.

ANTIMONII SULPHAS.

SULPHATE OF ANTIMONY.

R. Antimony, two parts.
Sulphuric acid, three parts.

Heat in an earthen vessel, stirring from time to time, leave the mixture on the fire till it has assumed a grayish-white color, wash carefully, to remove the excess of acid, and dry the powder. *Van Mons.*

ANTIMONII ET POTASSÆ
TARTRAS.

TARTAR EMETIC.

POWDER OF TARTAR EMETIC.

R. Tartar emetic, one grain.
Sugar, thirty grains.
Mix, and divide into ten powders. One every three or four hours, as a diaphoretic in fevers, after bleeding. *A. T. Thomson.*

COMPOUND POWDER OF TARTAR EMETIC.

R. Tartar emetic, half a drachm.
Sulphate of potassa, one ounce.
Powdered liquorice, one ounce and a half.

Mix well. Two scruples contain one grain of tartar emetic. *Beasley.*

R. Tartar emetic, one grain.
Kermes mineral, two grains.
Powdered orris root, one scruple.
" gum Arabic,
" sugar, each, one drachm.

Mix, and divide into twenty-four powders. One every hour, as an expectorant. *Cadet.*

POWDER OF TARTAR EMETIC AND
QUININE.

R. Tartar emetic, three grains.
Sulphate of quinine, ten grains.

Mix, and divide into six powders. One, every two hours, in apyrexia of obstinate intermittents. Both vomits and purges. *Gola.*

POWDER OF TARTAR EMETIC AND
PHOSPHATE OF LIME.

R. Tartar emetic, one grain.
Phosphate of lime, thirty-two grains.

Mix. To be divided into four powders; one to be taken in the evening, in chronic dysury, produced by checked perspiration. *Swediaur.*

POWDER OF TARTAR EMETIC AND IPECA-
CUANHA.

R. Tartar emetic, two grains.
Ipecacuanha, two scruples.
Starch, half a drachm.

Mix, and divide into three powders; one every quarter of an hour, till vomiting is induced. *Radius.*

BOLUS OF TARTAR EMETIC.

R. Tartar emetic, six grains.
Powdered Peruvian bark, six drachms.
Extract of juniper, sufficient.

Mix, and make sixty boluses. To be taken in twenty-four hours in quartan intermittents. *Laennec.*

PILLS OF TARTAR EMETIC AND CAM-
PHOR.

R. Tartar emetic, two grains.
Powdered opium, six grains.
" camphor, thirty-six grains.
Alcohol, three drops.
Conserve of roses, sufficient.

Powder the camphor with the alcohol, add the other powders, mix and incorporate with the conserve, and divide into twelve pills. One every fourth hour as a diaphoretic. *Ellis.*

PILLS OF TARTAR EMETIC AND OPIUM.

R. Tartar emetic,
Opium, each, one grain and a half.
Powdered tragacanth,
Conserve of roses, each, sufficient.

Mix, and make twenty-five pills. Two, night and morning, in chronic pulmonary catarrh. *Parrish.*

R. Tartar emetic, twelve grains.
Opium, ten grains.
Crumb of bread,
Gum Arabic, each, sufficient.

Mix, and make pills of half a grain. Dose, one increased to three in chronic rheumatism. *Broussais.*

PILLS OF TARTAR EMETIC AND
GUAIAIACUM.

R. Tartar emetic, one grain.
Powdered guaiacum, half a drachm.
Pill of aloes and myrrh, half a drachm.
Molasses, sufficient.

Mix, and make sixteen pills. *Beasley.*

SOLUTION OF TARTAR EMETIC.

R. Tartar emetic, four grains.
Sugar, one drachm.
Distilled water, four fl. ounces.

Dissolve. A tablespoonful every ten or fifteen minutes till it operates. *Ellis.*

SOLUTION OF TARTAR EMETIC AND
SQUILL.

R. Tartar emetic, two grains.
Oxymel of squill, half an ounce.
Water, two fl. ounces and a half.

Mix. One-half to be taken, and if it does not vomit in fifteen minutes, the remainder. *Augustin.*

EMULSION OF TARTAR EMETIC.

R. Tartar emetic, five grains.
Camphor, half a drachm.

Blanched almonds,
Syrup, each, one ounce.
Water, ten fl. ounces.

Make an emulsion with the almonds and water, mix the camphor rubbed with a few drops of alcohol, and add the tartar emetic; when dissolved, add the syrup; emetic and expectorant, according to the dose. *Van Mons.*

CLYSTER OF TARTAR EMETIC.

R. Tartar emetic, one to two scruples.
Tepid solution of gum Arabic, one pint.

To overcome obstinate constipation. To be used with much caution. *Chapman.*

R. Tartar emetic, three to eight grains.
Infusion of arnica, twelve fl. ounces.
Dissolve. In apoplexy and cerebral affections. *Foy.*

LOTION OF TARTAR EMETIC.

R. Tartar emetic, one scruple.
Water, one pint.
Dissolve. In cutaneous affections. *Pierquin.*

LOTION OF TARTAR EMETIC AND CORROSIVE SUBLIMATE.

R. Tartar emetic, one drachm.
Corrosive sublimate, five grains.
Compound spirit of lavender, one fl. drachm.
Water, one fl. ounce.

Mix. Wet the finger with the solution, and rub on the part for five or ten minutes; it will cause a pustular eruption in a few hours. *Hannay.*

LOTION OF TARTAR EMETIC AND CAMPHOR.

R. Tartar emetic, one drachm.
Boiling water, one pint.
Tincture of camphor, half a fl. ounce.
Mix. As a counter-irritant on the chest, in pulmonary complaints. *Augustin.*

OINTMENT OF TARTAR EMETIC.

R. Tartar emetic, one drachm.
Lard, one ounce.

Mix. When applied to the skin, it occasions a pustular eruption. *Hannay.*

R. Tartar emetic, two drachms.
Sugar, one drachm.
Cinnabar, five grains.
Spermaceti ointment, nine drachms.

Mix. *Jenner.*

R. Tartar emetic, two drachms.
Lard, one ounce.
Mix. *U. S. Ph.*

COMPOUND TARTAR EMETIC OINTMENT.

R. Tartar emetic, half a drachm.
Sal ammoniac, one drachm.
Camphor, twenty-five grains.
Musk, ten grains.
Lard, one ounce.

Mix. As a counter-irritant, in chronic diseases of the liver. *Fabre.*

PLASTER WITH TARTAR EMETIC.

R. Tartar emetic, one part.
Burgundy pitch, seven parts.
Mix, and spread. *Beral.*

Or, it may be made by sprinkling tartar emetic on a pitch, or adhesive plaster. *Ellis.*

It takes some time to produce its effect, and the eruption that ensues is very irritating and painful.

R. Resin plaster, one ounce.
Resin, four drachms.
Venice turpentine, three drachms.

Melt together by a gentle heat, and when nearly cold, add

Powdered tartar emetic, one drachm.

To be applied to the nape of the neck in scarlatina in children, also in rheumatism. *Niemann.*

AMMONIACAL LINIMENT WITH TARTAR EMETIC.

R. Ammoniacal liniment, one fl. ounce.
Tartar emetic, one drachm.
Mix. To indolent tumors, &c. *Ellis.*

ANTIMONIAL WINE.

R. Tartar emetic, one scruple.
Wine, ten fl. ounces.
Dissolve. *U. S. Ph.*

Dose, as emetic for children, from thirty drops to a fl. drachm, every fifteen minutes, till it operates.

COLLYRIUM WITH ANTIMONIAL WINE.

R. Antimonial wine, half fl. ounce.
Rose water, four fl. ounces.
Balsam of Fioraventi, two drops.

Mix. Recommended in chronic ophthalmia. *Spielmann.*

MIXTURE OF ANTIMONIAL WINE AND AMMONIAC.

R. Ammoniac, one scruple.
 Oxymel of squill, half fl. ounce.
 Antimonial wine, one fl. drachm.
 Syrup of liquorice, one fl. ounce and a half.

Mix. A spoonful every two hours, as an expectorant. *Phæbus.*

MIXTURE OF ANTIMONIAL WINE AND LAUDANUM.

R. Antimonial wine, three fl. drachms.
 Laudanum, one fl. drachm.

Mix. Fifteen drops, every half hour or hour in rheumatic diarrhœa. *Monro.*

MIXTURE OF ANTIMONIAL WINE AND BITTER SWEET.

R. Antimonial wine, half fl. drachm.
 Syrup of marsh mallow, three fl. drachms.
 Fennel water, one fl. ounce and a half.

Extract of bitter sweet, fifteen grains.

Cream of tartar, forty-five grains.

Mix. A teaspoonful every hour, in whooping cough in young children. *Phæbus.*

SYRUP OF ANTIMONIAL WINE.

R. Antimonial wine, } each,
 Sal ammoniac, } two drachms.
 Ammoniac, }
 Oxymel of squill, one ounce.
 Syrup of mallow, half a pound.

Mix. In pectoral affections, to facilitate expectoration and to keep the bowels open. A spoonful every hour. *Cadet.*

PULVIS ANTIMONIALIS.

ANTIMONIAL POWDER.

R. Sulphuret of antimony, powdered,
 Horn shavings, equal parts.

Mix, and put them in a red-hot iron pot, and stir constantly till they acquire an ash-gray color, and no vapor arises. Pulverize the product, put it in a crucible with a perforated cover, and expose this to a heat gradually increased to whiteness, which is to be kept up for two hours. When cold, reduce the product to a fine powder. *Ed. Ph.*

Emetic and diaphoretic; in doses of three to eight grains, diaphoretic; in larger, emetic and purgative, but very uncertain in its operation.

JAMES'S POWDER.

R. Tartar emetic, one scruple.
 Prepared burnt hartshorn,
 Oxide of antimony, each, five scruples.

Mix, and divide into twenty-one grain powders. Said to be the formula by which the original powder was prepared by Dr. James. (*Vid. Phil. Journ. Pharm., vi. 282.*)

PILLS OF ANTIMONIAL POWDER AND CALOMEL.

R. Antimonial powder, ten grains.
 Powdered opium,
 Calomel, each, two grains.
 Conserve of roses, sufficient.

Mix, and make four pills. Two to be taken at bed-time in acute rheumatism. *Ellis.*

APOCYNUM.

DOGSBANE.

Two species of this genus are officinal in the U. S. Pharm. *A. androsæmifolium* and *A. cannabinum*, possessed of much the same properties; these are emetic, cathartic, and sometimes diuretic. The latter is rather the more powerful.

Sex. Syst. Pentand. digyn. *Nat. Syst.* Apocynaceæ.

A. ANDROSÆMIFOLIUM.

DOGSBANE.

Bigelow, Med. Bot. ii. 148. Griffith, Med. Bot. 449.

A native of most parts of the U. S. Emetic and diaphoretic. Dose, 40 grains as an emetic; where it is wished to act on the skin, five to ten grains combined with a grain of opium.

A. CANNABINUM.

INDIAN HEMP.

Griscom, Am. Journ. Med. Sci. xii. 55. Griffith, Med. Bot. 450.

Found in many parts of the United States. Emetic, in doses of fifteen to thirty grains.

DECOCTION OF INDIAN HEMP.

R. Root of Indian hemp, half an ounce.
 Water, a pint and a half.

Boil to a pint. *Griscom.*

This has been found useful in dropsy, in doses of one or two fl. ounces, two or three times a-day, acting as a hydragogue purgative.

EXTRACT OF INDIAN HEMP.

R. Powdered root of Indian hemp, one pound.
Water, one gallon.

Macerate for twenty-four hours, boil down to one quart, strain, and evaporate to a proper consistence. Dose, three to five grains. Acts like the decoction, but is not as efficient.

ARALIA.

Most of the species of this genus are medicinal, but two only are officinal.

Sex. Syst. Pentand. pentag. *Nat. Syst.* Araliaceæ.

A. NUDICAULIS.

FALSE SARSAPARILLA.

Torrey and Gray, Fl. i. 646. Griffith, Med. Bot. 344.

A mild, stimulating diaphoretic, the root has been employed as a substitute for sarsaparilla; best given in infusion.

A. SPINOSA.

ANGELICA TREE.

Torrey and Gray, Fl. i. 647. Griffith, Med. Bot. 345.

The bark is a stimulating diaphoretic; a decoction has been found useful in rheumatic, syphilitic, and cutaneous affections; as has also a tincture.

ARCTIUM.

BURDOCK.

One species of this genus is officinal in our Pharmacopœia, the *A. lappa*. (*Lappa minor*).

Sex. Syst. Syngen. æqual. *Nat. Syst.* Asteraceæ.

Linn. Sp. Pl. 1143. Griffith, Med. Bot. 411. (*Lappa*.)

This is a native of Europe, but generally naturalized in the United States. The parts used are the root and seeds. They are diaphoretic and diuretic, especially the latter. Dose, half a drachm to a drachm.

INFUSION OF BURDOCK.

R. Burdock root, one ounce.
Boiling water, two pints.

Infuse for six hours, and strain. *Cottureau.*

Used as a diaphoretic, in rheumatic pains, &c.

DECOCTION OF BURDOCK.

R. Burdock root, two ounces.
Water, three pints.

Boil down to two pints, and strain. *Wood.*

Dose, a wineglassful.

ARGENTUM.

SILVER.

A solid, white, brilliant, very ductile metal. It is wholly soluble in nitric acid. The only officinal preparations in our pharmacopœia are the nitrate, oxide, and cyanuret; but several others are employed in Europe.

ARGENTI CHLORIDUM.

CHLORIDE OF SILVER.

R. Solution of nitrate of silver, at will.

Add gradually a solution of common salt, wash well the resulting precipitate, and dry by a gentle heat in the shade.

POWDER OF CHLORIDE OF SILVER.

R. Chloride of silver, one grain.
Powdered orris root, two grains.

Triturate well, and divide into ten parts. Used in syphilis, in friction on the tongue. *Serre.*

CHLORIDE OF SILVER AND AMMONIA.

R. Liquid ammonia, at will.

Saturate, by the aid of heat, with recently precipitated and well-washed chloride of silver. Filter whilst hot, and crystallize.

PILLS OF CHLORIDE OF SILVER AND AMMONIA.

R. Chloride of silver and ammonia, one grain.

Powdered orris root, two grains.

Conserve of linden flowers, sufficient.

Mix, and make fourteen pills. *Serre.*

ARGENTI CYANURETUM.

CYANIDE OR CYANURET OF SILVER.

R. Nitrate of silver,
Ferrocyanuret of potassium, each, two ounces.
Sulphuric acid, one ounce and a half.
Distilled water, sufficient.

Dissolve the nitrate in a pint of water, and put the solution in a glass receiver. Dissolve the ferrocyanuret in ten fl. ounces of water, and pour the solution into a retort adapted to the receiver. To the solution in the retort add the sulphuric acid diluted with four fl. ounces of water, and distil until the liquid that passes produces no longer a precipitate in the receiver. Then wash and dry the precipitate. *U. S. Ph*

Used in syphilis, in doses of one-twelfth to one-tenth of a grain.

OINTMENT OF CYANIDE OF SILVER.

R. Cyanide of silver, ten grains.
Lard, one ounce.
Mix. *Serre.*

Employed as an application to ulcerations of the cornea, irritable ulcers, &c.

ARGENTI IODIDUM.

IODIDE OF SILVER.

R. Solution of nitrate of silver, at will.
Add gradually a
Solution of iodide of potassium.
Wash the precipitate, and dry by a gentle heat. *Patterson.*
Dose, one or two grains.

OINTMENT OF IODIDE OF SILVER.

R. Iodide of silver, ten grains.
Lard, one ounce.
Mix. *Serre.*

Used in the same cases as the ointment of the cyanide.

ARGENTI OXIDUM.

OXIDE OF SILVER.

R. Nitrate of silver, four ounces.
Water, half a pint.
Solution of potassa, one pint and
a half.

Dissolve the nitrate in the water, and add the potassa as long as precipitation occurs. Wash and dry the precipitate, and keep it in opaque, well-stopped bottles. *U. S. Ph.*

Used as a substitute for the nitrate, in doses of about half a grain twice a-day, in diseases of the stomach, of undue secretion, &c.

PILLS OF OXIDE OF SILVER.

R. Oxide of silver, six grains.
Powdered liquorice, twelve grains.
Syrup, sufficient.
Make twelve pills. Dose, one pill three times daily, in pyrosis.

OINTMENT OF OXIDE OF SILVER.

R. Oxide of silver, twenty grains.
Lard, one ounce.
Rub well together. *Serre.*

Used as an application to irritable ulcers, &c.

ARGENTI NITRAS.

FUSED NITRATE OF SILVER.

R. Silver, in small pieces, one ounce.
Nitric acid (sp. gr. 1.42), seven
fl. drachms.
Distilled water, two fl. ounces.

Mix the acid with the water, and dissolve the silver in the mixture, on a sand-bath; then gradually increase the heat, so as to dry the salt. Melt this in a crucible, and continue the heat till ebullition ceases, then immediately cast into moulds. *U. S. Ph.*

NITRATE OF SILVER.

R. Silver, in small pieces, one ounce.
Nitric acid, seven fl. drachms.
Distilled water, two fl. ounces.

Mix the acid with the water, and dissolve the silver in the mixture on a sand-bath. Pour off the solution, evaporate to one-half, and crystallize. Pour off the supernatant liquid, again evaporate and crystallize. Repeat the process a third time. Dry the crystals in a glass funnel and preserve in an opaque well-stopped bottle. *U. S. Ph.*

Used as a tonic and antispasmodic, in doses of a fourth of a grain, gradually increased.

SOLUTION OF NITRATE OF SILVER.

R. Nitrate of silver, one drachm.
Distilled water, one fl. ounce.
Dissolve and strain. Keep in the dark. *London Ph.*

Chiefly intended as a test liquor.

PILLS OF NITRATE OF SILVER.

R. Crystallized nitrate of silver, one grain.
Crumb of bread, one drachm.
Make sixteen pills. *Guibourt.*

Each pill contains a sixteenth of a grain of the nitrate.

R. Nitrate of silver, ten grains.
Opium, four grains.
Extract of gentian, " liquorice, each, a drachm
and a half.

Make pills of a grain each. These each contain a twentieth of the nitrate. *Brera.*

R. Nitrate of silver, two grains.
Crumb of bread, sufficient.

Mix well, and divide into four pills. One to be taken every six hours.

Advised in chronic epilepsy and other spasmodic disorders. To prevent the blue tinge of the skin, so often produced by a continued use of this salt of silver, eight drops of diluted nitric acid in a fl. ounce of water, should be taken after each pill. *A. T. Thomson.*

MACKENSIE'S SOLUTION.

- R. Nitrate of silver, one scruple.
 Distilled water, one fl. ounce.
 Dissolve. *Mackensie.*

Used to wash the throat and fauces, in affections of those parts.

NITRATE OF SILVER COLLYRIUM.

- R. Nitrate of silver, ten grains.
 Laudanum, half a fl. drachm.
 Distilled water, one fl. ounce.
 Mix. *Foy.*
 Employed as a wash in purulent ophthalmia.

ANTI-EPILEPTIC MIXTURE.

- R. Nitrate of silver, twelve grains.
 Laudanum, twenty drops.
 Mucilage of gum Arabic, one ounce.
 Mix. *Radius.*
 Given in epilepsy, in doses of twenty drops, twice a-day.

HAIR DYE.

- R. Silver, two drachms.
 Steel filings, half an ounce.
 Nitric acid, one ounce.
 Rain water, eight fl. ounces.
 Dissolve, and strain. To be applied by means of a fine brush. *Bateman.*

INDELIBLE INK.

- R. Carbonate of soda, half an ounce.
 Distilled water, four ounces.
 Mix, and make mordant.
 R. Nitrate of silver, five scruples.
 Gum Arabic, two drachms.
 Sap green, one scruple.
 Distilled water, one fl. ounce.
 Mix, and make ink. *Gray.*

INDELIBLE INK WITHOUT MORDANT.

- R. Nitrate of silver, one ounce.
 Crystallized carbonate of soda, one ounce and a half.
 Tartaric acid, eight scruples.
 Strong solution of ammonia, two fl. ounces.
 Archil, half fl. ounce.
 White sugar, six drachms.
 Powdered gum Arabic, ten drachms.
 Distilled water, sufficient.

Dissolve the nitrate and carbonate in separate portions of the water, mix the solutions, collect and wash the precipitate on a filter, rub it

whilst still moist in a porcelain mortar with the tartaric acid, till effervescence has ceased, add the ammonia to dissolve the tartrate of silver, then mix the archil, sugar, and gum, adding as much water as will make six fl. ounces. *Redwood.*

OINTMENT OF NITRATE OF SILVER.

- R. Nitrate of silver, two grains.
 Lard, two drachms.
 Mix well. As an application in acute ophthalmia. *Velpeau.*
 R. Nitrate of silver, four parts.
 Lard, thirty parts.
 Or, eight parts of the salt to the same proportion of lard, or twelve parts to thirty. Used as an application to white-swelling. *Jobert.*
 R. Nitrate of silver, one part.
 Lard, seven and a half parts.
 Mix, to smear bougies, in cure of gonorrhœa. *Macdonald.*

COMPOUND OINTMENT OF NITRATE OF SILVER.

- R. Nitrate of silver, ten to twenty grains.
 Oxide of zinc ointment, half an ounce.
 Balsam of Peru, one drachm.
 Triturate thoroughly together. To heal venereal ulcers, and to promote cicatrization. *Fricke.*
 R. Nitrate of silver, three grains.
 Solution of subacetate of lead, five drops.
 Lard, one drachm.
 Mix well. In chronic ophthalmia. *Guthrie.*

ARMORACIA.

HORSERADISH.

This is a species of *Cochlearia*, the *C. armoracia*. A native of many parts of Europe, and much cultivated both there and in this country, for the sake of its roots, so well known as a condiment. The root is the part used in medicine. It has a pungent odor, and a warm, acrid taste.

Sex. Syst. Tetrad. silic. *Nat. Syst.* Brassicaceæ.

Linn. Sp. Pl. 904. *Griffith, Med. Bot.* 131.

As a remedial agent, it is an active stimulant, and when applied to the skin acts as a rubefacient.

INFUSION OF HORSERADISH.

- R. Fresh horseradish, sliced,
 Mustard seed, bruised, each, one ounce
 Boiling water, one pint

Macerate for two hours in a covered vessel, and strain.

U. S. Ph.

It is sometimes used in paralytic, scorbutic, and dropsical affections, attended with debility. The dose is two fl. ounces, three or four times a-day.

COMPOUND SPIRIT OF HORSERADISH.

R. Sliced horseradish,
Dried orange peel, each,
twenty ounces.
Bruised nutmeg, five drachms.
Proof spirit, one gallon.
Water, two pints.

Mix, and distil a gallon by a gentle heat.

London Ph.

Employed as a diuretic, in dropsy with debility. Dose, one to four fl. drachms.

MIXTURE OF HORSERADISH.

R. Infusion of horseradish, four fl. ounces.
Syrup of cinchona, one fl. ounce.
Extract of fumitory, one drachm.

Mix. *Foy.*

In spoonful doses, as an antiscorbutic.

COMPOUND HORSERADISH CATAPLASM.

R. Bruised horseradish, six drachms.
" mustard seed, one ounce.
Flaxseed meal, two ounces and a half.
Vinegar, sufficient.

Mix, and make cataplasm, to be applied to the feet as revulsive in diseases of the head.

Ainslie.

COMPOUND INFUSION OF HORSERADISH.

R. Horseradish, one ounce.
Boiling water, one pint.

Infuse for an hour in a close vessel; on cooling, add

Simple syrup, one ounce and a half.

A wineglassful, several times a-day, in dropsy, especially when arising from a granular affection of the kidneys.

ARNICA.

WOLFSBANE.

Several species of this genus are medicinal, but one only is officinal, *A. montana*, a native of Europe; but it is likely that the *A. nudicaule* of this country is possessed of the same powers.

Sex. Syst. Syngen. super. *Nat. Syst.* Astera-
raceæ.

Linn. Sp. Pl. 1245. *Griffith, Med. Bot.* 407.

Arnica is a stimulant, acting with much energy on the cerebro-spinal system, and is also an active irritant of the stomach and bowels. In Germany, where it is much employed, it is given to fulfil a variety of indications.

COMPOUND POWDER OF ARNICA.

R. Powdered arnica root,
" serpentaria,
" sugar, each, two drachms.
Oil of peppermint, ten drops.

Mix, and divide into sixteen powders.

Augustin.

Dose, a powder, every two hours, in the diarrhoea complicating typhoid fevers.

EXTRACT OF ARNICA.

R. Flowers of arnica, one part.
Water, eight parts.
Alcohol, one part.

Mix, and macerate for three days at a gentle heat, express, distil off the alcohol, and evaporate the residue to a proper consistence.

Wurtzburg Ph.

Dose, ten grains to a scruple, in chronic rheumatism and paralysis.

INFUSION OF ARNICA.

R. Flowers of arnica, one ounce.
Boiling water, one pint.

Infuse for half an hour, and strain.

Used in the same cases as the last. Dose, half an ounce to an ounce.

Cottreau.

COMPOUND INFUSION OF ARNICA.

R. Flowers of arnica, one drachm.
" chamomile, half an ounce.

Peppermint, two drachms.
Boiling water, nine fl. ounces.

Mix, macerate, and strain.

Copenh. Ph.

Dose, one ounce.

FOMENTATION OF ARNICA.

R. Flowers of arnica, half an ounce.
Boiling vinegar, sufficient

to obtain six ounces; add

Carbonate of ammonia, two drachms.

Radius.

Used as a warm fomentation in œdema of the scrotum.

DECOCTION OF ARNICA.

R. Flowers of arnica, one ounce.
Water, three pints.

Boil till reduced to two pints, strain, and add
Syrup of ginger, two ounces.

Swedjaur.

Dose, one to two ounces, every two hours, in
aphonia, paralysis, rheumatism, &c.

FOMENTATION OF ARNICA AND RUE.

R. Flowers of arnica, two ounces.
Rue, one ounce.
Boiling water, sufficient

to obtain twelve ounces when strained, after
one hour of maceration. *Radius.*

Used as a fomentation in contusions.

TINCTURE OF ARNICA.

R. Flowers of arnica, one part.
Alcohol, sixteen parts.

Digest, express, and filter. *Beral.*

Dose, thirty drops, several times a-day.

ARSENICUM.

ARSENIC.

Metallic arsenic is not employed in medicine;
but several of its combinations are of much im-
portance.

ARSENICI CHLORIDUM.

SOLUTION OF CHLORIDE OF ARSENIC.

R. Arsenious acid, in small
pieces, half a drachm.
Muriatic acid, one and a half fl.
drachms.
Water, twenty fl. ounces.

Boil the arsenic in the muriatic acid, diluted
with an ounce of water, until dissolved; then
add sufficient water to make one pint (Imp.).

Lond. Ph.

This is much used in Great Britain in lepra
and chorea, in doses of three to ten minims,
thrice daily.

Each fl. ounce represents one grain and a
half of arsenious acid.

ARSENICI IODIDUM.

IODIDE OF ARSENIC.

R. Finely-powdered metallic
arsenic, one drachm.
Iodine, five drachms.

Triturate together, and introduce into a small
flask or retort; place this on a sand-bath, and
apply gentle heat till liquefaction is produced.

The vessel should be nearly full, to prevent the
formation of much iodine vapor, and to enable
the operator to bring the fixed mass in contact
with every part of it, so as to include any
iodine that may have been sublimed. If no
iodine odor is perceptible, and the contents
assume a reddish-yellow color, and crystallize
on the sides of the vessel, the operation is com-
plete. *W. Procter.*

This corresponds with the officinal formula.

Dose, one-tenth of a grain, gradually in-
creased to a quarter, three times a-day. Given
in cancer, and obstinate cutaneous affections.

IODIDE OF ARSENIC PILLS.

R. Iodide of arsenic, one grain.
Extract of conium, twenty grains.
Mix, and make ten pills. *Thomson.*

Used in lepra, and obstinate eruptions.

IODIDE OF ARSENIC OINTMENT.

R. Iodide of arsenic, three grains.
Lard, one ounce.
Triturate well. *Bielt.*

Employed in cases of lepra, and other cutane-
ous affections; but requires much caution.

LIQUOR ARSENICI ET HYDRAR- GYRI IODIDI.

SOLUTION OF IODIDE OF ARSENIC AND MERCURY.

DONOVAN'S SOLUTION.

R. Iodide of arsenic,
Red iodide of mercury,
each, thirty-five grains.
Distilled water, half a pint.

Rub the iodides with half a fl. ounce of water,
then add rest of water, heat to boiling point
and filter. *U. S. Ph.*

Each fl. drachm contains one-eighth of a
grain of arsenic, and one-fourth of a grain of
deutoxide of mercury. This preparation is said
to be useful in obstinate cutaneous diseases, as
lepra, lupus, &c. The dose is from five minims
to half a fl. drachm two or three times a-day.

DRAUGHT OF SOLUTION OF IODIDE OF MERCURY AND ARSENIC.

R. Solution iodide of mercury and arsenic,
two fl. drachms.
Distilled water, three fl. ounces and a
half.
Syrup of ginger, half fl. ounce.

Mix, and divide into four draughts. One to be
taken night and morning. Not to be given in a
metallic spoon. *Donovan.*

ARSENICI TERSULPHURETUM.

ORPIMENT.

DEPILATORY OF ORPIMENT.

R. Orpiment,	one ounce.
Quicklime,	one pound.
Starch,	ten ounces.
Water,	sufficient

to form a soft paste.

Foy.

This is spread on the part where it is desired to remove the hair, and washed off when it begins to dry. It is the *rusma* of the Turks.

DELCROIX'S DEPILATORY.

(*Poudre Subtile.*)

R. Orpiment,	four parts.
Quicklime,	thirty parts.
Powdered gum,	sixty parts.

Used like the preceding to remove superfluous hair.

POWDER OF ORPIMENT.

R. Orpiment,	one ounce.
Quicklime,	twelve ounces.
Jasmine powder,	ten ounces.
Powder of palm soap,	four ounces.

Mix well.

Morfit.

This, when applied as a depilatory, is mixed with a little water.

Both these, although efficient for the purposes designed, are dangerous, and should never be used, except with extreme caution.

LOTION OF ORPIMENT.

R. Orpiment,	two drachms.
Verdigris,	each, half a drachm.
Aloes,	
Myrrh,	
Rose water,	three fl. ounces.
White wine,	six fl. ounces.

Mix.

Van Mons.

Used as an application to fungous ulcers.

ARTEMISIA.

A genus of bitter herbs, several of which are used as medicinal agents to fulfil various indications; some as tonics, (*A. absinthium*), some as moxas, (*A. moxa*), and some as anthelmintics.

Sex. Syst. Syn. pol. super. *Nat. Syst.* Asteraceæ.

1. A. ABROTANUM.

SOUTHERNWOOD.

A native of the south of Europe and the Levant, and generally cultivated in gardens.

Its odor is aromatic and peculiar, and its taste bitter. It is used as a tonic, antispasmodic, and vermifuge.

Linn. Sp. Pl. 1185. Griffith, Med. Bot. 406.

INFUSION OF SOUTHERNWOOD.

R. Southernwood,	six drachms.
Boiling water,	one pint.

Digest for two hours, and strain. Given in hysteria and in dysmenorrhœa. Dose, one fl. ounce.

Taddei.

CLYSTER OF SOUTHERNWOOD.

R. Southernwood,	half an ounce.
Boiling water,	one pint.

Digest for half an hour, and, to six or eight fl. ounces of strained infusion, add

Olive oil,	one ounce.
------------	------------

Said to be effectual in cases of worms, and especially ascarides.

Radius.

2. A. SANTONICA.

SEMEN CONTRA.

Under this name are included several species, as the *A. santonica*, *contra*, *judaica*, &c. They are principally from Barbary and the Levant. The parts used are the flowers, buds, peduncles, &c. Semen contra is much employed in Europe as an anthelmintic, but is seldom used in this country. The dose in powder is from ten to thirty grains. Its active principle is called santonine.

POWDER OF SEMEN CONTRA.

R. Semen contra,	
Extract of tansy, each,	six grains.
Oxide of iron,	four grains.
Oil of valerian,	one drop.

Mix.

Augustin.

ELECTUARY OF SEMEN CONTRA.

R. Semen contra,	ten grains.
Sulphate of iron,	four grains.
Jalap,	
Honey, each,	one scruple.

Mix.

Rosenstein.

ANTHELMINTIC MIXTURE OF SEMEN CONTRA.

R. Infusion of semen contra,	four fl. ounces.
Syrup of senna,	one fl. ounce.

Mix.

Foy.

POWDER OF SANTONINE.

R. Santonine,	six grains.
Sugar of milk,	fifteen grains.

Divide into six powders and give one night and morning, to a child five years old, for lumbrici.

Thomas.

ARUM.

INDIAN TURNIP.

Several species of Arum are used in medicine, but the only one officinal in the U. S. Pharm., is *A. triphyllum*; a native plant, with large, perennial cormus; this is white, fleshy, and extremely acrid in a fresh state, but becomes edible and bland when dried, and kept for any time; and the secula obtained from it resembles arrowroot, in appearance.

Sext. Syst. Mon. polyand. *Nat. Syst.* Araceæ.

Bigelow, Am. Med. Bot. 1, 52. Griffith, Med. Bot. 616.

The powder of the recently-dried root may be given in doses of ten grains, in an emulsion of gum Arabic, sugar, and water, several times a-day. It has been found useful in chronic catarrh, chronic bronchitis, rheumatism, &c.

ASARUM.

There are two plants recognized by this name, one by the London College, and the other by the U. S. Pharm.

Sext. Syst. Dodecand. monog. *Nat. Syst.* Aristolochiaceæ.

1. ASARUM EUROPÆUM.

ASARABACCA.

A native of Europe, of which the leaves and root are used. The first have a feeble aromatic odor, when rubbed, and a somewhat spicy taste; the latter has a strong, penetrating odor, and an acrid, bitter, and nauseous taste; this is sometimes used abroad, as a stimulant emetic, and anthelmintic; the leaves, as the basis of various errhine powders.

Linn. Sp. Pl. 633. Griffith, Med. Bot. 527.

COMPOUND POWDER OF ASARABACCA.

R. Dried leaves of asarabacca, one ounce.
Lavender flowers, one drachm.

Rub together to a fine powder.

Dublin Ph., 1826.

Has been found useful as an errhine, in some cases of headache, toothache, &c. Five or six grains to be snuffed up the nostrils at night.

2 ASARUM CANADENSE.

WILD GINGER.

Bigelow, Med. Bot. 1, 150. Barton, Veg. Mat. Med. p. 85.

This species is common to most parts of the U. S. The leaves are said by Drs. Barton and Bigelow, not to be emetic, but other authorities state that a teaspoonful of this powder never fails to act on the stomach. Like those of the asarabacca, they are eminently errhine. The root is an aromatic stimulant, with active diaphoretic properties, and may be used as a substitute for serpentaria.

INFUSION OF WILD GINGER.

R. Root of wild ginger, half an ounce.
Boiling water, one pint.

Digest for an hour in a covered vessel, and strain. In all cases where Virginia snake-root is indicated. Dose, one to two fl. ounces.

ASCLEPIAS.

Sext. Syst. Pentand. digyn. *Nat. Syst.* Asclepiadaceæ.

Many species of this large genus are employed in medicine, and it is probable that the whole of them are endowed with active properties.

1. A. TUBEROSA.

PLEURISY ROOT.

A native of most parts of the United States. The root is large, and irregularly tuberous; the taste of it is bitter, nauseous, and somewhat acrid. It is an active diaphoretic, and, in large doses, purgative. It has attained much popular reputation in the treatment of diseases of the respiratory organs, and more especially of pleurisy. The dose of the powder is from a scruple to a drachm, several times a-day. An infusion or decoction is preferable to the powder.

INFUSION OF PLEURISY ROOT.

R. Bruised pleurisy root, one ounce.
Boiling water, two pints.

Digest for two hours. Dose, three or four fl. ounces, warm, to be given every two or three hours, until it operates. Chapman.

2. A. INCARNATA.

SWAMP SILK WEED.

A native of many parts of the U. S., usually growing in wet places. The root is the officinal portion. It is emetic and cathartic in full doses of half a drachm to a drachm, and expectorant and alterative in small ones.

3. A. SYRIACA.

SILK WEED.

Very common in the U. S. The root appears to have much the same properties as the last, and also some anodyne powers.

ASPARAGUS.

ASPARAGUS.

Several species of *Asparagus* have been used in medicine, but the only one that is now employed is the *A. officinalis*. This is a native of Europe, and is generally cultivated there and in this country.

Sex. Syst. Hexand. monog. *Nat. Syst.* Liliaceæ.

Linn. Sp. Pl. 448. Stokes, Bot. Mat. Med. ii. 252. Griffith, Med. Bot. 654.

The parts used are the roots and young shoots, (turiones). These are diuretic, aperient, and deobstruent, and have been thought to exercise a sedative influence on the heart. Dr. Wood states (*U. S. Dispens.*) that the berries are more efficient than the shoots.

EXTRACT OF ASPARAGUS SHOOTS.

R. Clarified juice of asparagus, sufficient.
Evaporate by a mild heat to a proper consistence.
Soubeiran.

EXTRACT OF ASPARAGUS ROOTS.

R. Roots of asparagus, sufficient.
Bruise them, add sufficient water to cover them, express, strain, and evaporate by a mild heat.
Soubeiran.

These extracts are given in doses of a scruple to a drachm, as diuretics.

SYRUP OF ASPARAGUS SHOOTS.

R. Juice of asparagus shoots, one pint.
White sugar, two pounds.
Dissolve by means of a water-bath, and strain.
Cottureau.

Advised in pulmonary catarrh, and too great action of the heart, in doses of one to four ounces during the day.

DECOCTION OF ASPARAGUS ROOTS.

R. Asparagus roots, one ounce.
Water, two pints.
Boil, and strain. *Radius.*
To be taken by cupfuls in dropsies.

DIURETIC MIXTURE.

R. Asparagus roots, one ounce.
Water, one pint.
Boil for an hour, strain, and add

Acetate of potassa, twenty grains.
Honey of squills, half an ounce.
Foy.

Used as the last, and in the same diseases.

ASSAFÆTIDA.

ASSAFÆTIDA.

A gum resin or inspissated juice from an umbelliferous plant, a native of Persia, usually thought to be *Ferula assafetida*, but now shown by Dr. Falconer to be a *Narthez*, which, though similar to *Ferula*, is distinct from it. (Royle, Mat. Med., 407.)

Sex. Syst. Pentand. digyn. *Nat. Syst.* Apiaceæ.

Royle, Mat. Med., 407. Griffith, Med. Bot. 326.

Assafetida, as found in the shops, is in masses of a whitish, reddish, or violet hue, composed of adhering tears. Odor, fetid and alliaceous; taste, bitter and somewhat acrid; it forms an emulsion with water.

Is stimulant and antispasmodic, and also emmenagogue and anthelmintic. It is used in spasmodic and convulsive diseases, as hysteria, chorea, hooping-cough, flatulent colic, &c.

Dose, ten grains.

ASSAFÆTIDA PILLS.

R. Assafetida, one ounce and a half.
Soap, half an ounce.
Beat with water into a mass, and divide into two hundred and forty pills; each of which contains three grains of the gum resin. *U. S. Ph.*
R. Assafetida, one drachm and a half.
Powdered orris root,
Mucilage of gum Arabic, each, sufficient.

Beat together, and divide into thirty pills. One to be given occasionally in the hooping-cough of children. *Kapp.*

R. Assafetida, a drachm and a half.
Powdered ginger, half a drachm.
Syrup, sufficient.

Mix, and make thirty pills. Three to be taken every three hours. Have been advised in palsy. *A. T. Thomson*

PILLS OF ASSAFÆTIDA AND LACTUCARIUM.

R. Assafetida, three drachms.
Lactucarium, two scruples and a half.
Mix, and divide into eighty pills. Two or three every hour in hooping-cough of adults. *Radius.*

PILLS OF ASSAFÆTIDA AND IRON

R. Assafetida, } each, half
Sulphate of iron, } an ounce.
Extract of chamomile, }

Mix well, and divide into one hundred and eighty pills. Two or more, three or four times a-day, in hypochondria and hysteria.

Syffret.

R. Black oxide of iron, half a drachm.
Assafetida, a drachm and a half.
Oil of tansy, ten drops.
Extract of wormwood, sufficient.

Mix, and divide into ninety pills. Six to be taken three times a day. Said to be useful against lumbrici and ascarides.

Phabus.

PILLS OF ASSAFETIDA AND MUSK.

R. Assafetida, two drachms.
Musk,
Camphor, each, one drachm.
Ambergris, half a drachm.

Beat well together, and divide into one hundred pills. Two to be taken three times a-day. Said to have proved useful in angina pectoris, hysteria, &c.

Sainte Marie.

PILLS OF ASSAFETIDA AND OPIUM.

R. Assafetida, half an ounce.
Powdered opium,
“ ipecacuanha, each,
four grains.
Oil of peppermint, eight drops.
Alcohol, sufficient.

Beat well together, and divide into one hundred and twenty pills. Ten to be taken three times a day in chronic ischuria, consecutive on gonorrhœa.

Radius.

ASSAFETIDA MIXTURE.

R. Assafetida, two drachms.
Water, half a pint.

Rub together, adding the water by degrees, till well mixed.

U. S. Ph.

R. Assafetida, one drachm.
Sugar, six drachms.
Rose water, five fl. ounces.

Make an emulsion, and add

Hoffmann's anodyne, thirty drops.

Taken in spoonful doses, in spasmodic asthma, and hysteria.

Bories.

ASSAFETIDA MIXTURE.

R. Assafetida, half a drachm.
Solution of acetate of ammonia,
half a fl. ounce.
Pennyroyal water, three fl. ounces.

Mix. One or two spoonfuls in hooping-cough.

Miller.

R. Assafetida mixture,
five and a half fl. ounces.

Compound spt. of lavender,
half fl. ounce.

Aromatic spt. of ammonia,
two fl. drachms.

Mix. Three spoonfuls a-day in hysteria.

Ainslie.

R. Assafetida mixture, two fl. ounces.
Opium, two grains.
Infusion of chamomile, four fl. ounces.

Mix. As an injection in convulsive attacks.

R. Assafetida, half a drachm.
Yolk of egg, one.

Rub together, and gradually add

Water, eight fl. ounces.

Strain. To form two injections. Said to be useful in the hooping-cough of children.

Soubciran.

MIXTURE OF ASSAFETIDA AND OXYMEL OF SQUILL.

R. Assafetida, one scruple.
Powdered digitalis, six grains.
Oxymel of squill, two fl. ounces.
Linden-flower water, four fl. ounces.

Rub well together. Said to be found useful in the dry cough consequent on deranged menstruation.

Pierquin.

Dose, a tablespoonful.

MIXTURE OF ASSAFETIDA AND TOLU.

R. Assafetida, one drachm.
Water, four fl. ounces.

Make an emulsion, and add

Tincture of tolu, half a fl. ounce.

“ opium, forty to fifty drops.

Mix well. A teaspoonful every two hours, in hooping-cough, and a dessertspoonful or more to an adult.

Ellis.

SYRUP OF ASSAFETIDA.

R. Assafetida, one ounce.
Boiling water, one pint.
Sugar, two pounds.

Rub the assafetida with a part of the boiling water till an uniform paste is made. Then gradually add the rest of the water, strain, and add the sugar, applying a gentle heat to dissolve it.

Dose, a tablespoonful, which contains seven grains and a half.

Richd. Peltz.

TINCTURE OF ASSAFETIDA.

R. Assafetida, four ounces.
Alcohol, two pints.

Macerate for fourteen days, and filter.

U. S. Ph

Has all the properties of the gum resin, but is more stimulating. Dose, a fl. drachm.

ALKALINE TINCTURE OF ASSAFETIDA.

R. Assafetida,
Carbonate of potassa, each,
two ounces.
Diluted alcohol, one pint.

Macerate for three days, by a mild heat, and filter. Anti-hysteria, &c. Useful in nervous disorders. Dose, about twenty drops.

Wirtzburg Dis.

AMMONIATED TINCTURE OF ASSAFETIDA.

R. Ammoniated alcohol,
sixteen fl. ounces.
Assafetida, one ounce.

Macerate for twenty-four hours, in a closed vessel, and then distil sixteen ounces.

Van Mons.

Stimulant and anti-hysteria. Dose, five to sixty drops.

TINCTURE OF ASSAFETIDA AND SOOT.

R. Assafetida, one drachm.
Wood soot, two drachms.
Alcohol, three fl. ounces.

Digest for six days, and filter. *Guibourt.*

Highly praised in hysteria, flatulence, and convulsions caused by dentition. Dose, ten to thirty drops in an appropriate vehicle.

ASSAFETIDA PLASTER.

R. Assafetida,
Lead plaster, each, one pound.
Galbanum,
Yellow wax, each, half a pound.
Alcohol, three pints.

Dissolve the assafetida and galbanum in the alcohol, by means of a water-bath; strain while hot, and evaporate to consistence of honey, add lead plaster and wax melted together; stir well, and evaporate to due consistence. *U. S. Ph.*

A good application to the stomach, in hysteria with flatulence, and to the chest in whooping-cough.

DEWEES'S CARMINATIVE.

R. Carbonate of magnesia, half a drachm.
Tincture of assafetida, sixty drops.
" opium, twenty drops.
Sugar, one drachm.
Distilled water, one fl. ounce.

Mix well. *Deweese.*

Very useful in the flatulent colic, diarrhoea, &c., of infants. Dose, twenty-five drops to a child of two to four weeks old, increasing the dose for those of a more advanced age.

PILLS OF ASSAFETIDA AND ALOES.

R. Assafetida,
Powdered aloes, } each, half an ounce.
Soap,

Beat with sufficient water to form a mass; divide into one hundred and eighty pills.

U. S. Ph.

A good purgative in cases of costiveness with flatulence and debility of digestive organs. Dose, two to five.

TINCTURE OF ASSAFETIDA AND CASTOR.

R. Tincture of assafetida,
" castor, each, one fl. ounce.
Aromatic spirit of ammonia, one fl. drachm.

Mix. In hysteria, &c. Dose, about a teaspoonful. *Ellis.*

COMPOUND ASSAFETIDA MIXTURE.

R. Assafetida, one drachm.
Peppermint water, three fl. ounces.
Make a solution, and add
Ammoniated tincture of valerian, two fl. drachms.
Tincture of castor, three fl. drachms.
Sulphuric ether, one fl. drachm.

Mix. Useful in hysteria, in doses of a tablespoonful, largely diluted, every second hour.

Ellis.

ASSAFETIDA ENEMA.

R. Assafetida, one drachm.
Hot water, two fl. ounces.
Triturate together. In flatulent colic, and against ascarides. *Foy.*

ATROPIA.

ATROPINE.

Take recently-dried root of belladonna, exhaust it, by alcohol, 40° Cartier; add to the tincture a quantity of slaked lime equal to one-twentieth of the weight of the root. After twenty-four hours' contact, filter; acidify slightly with sulphuric acid, and again filter. Two-thirds of the alcohol is then removed by distillation. Evaporate the residue to one-twelfth of the weight of the root employed. Carbonate of potassa is then added until the liquid begins to be rendered opaque by a grayish-brown precipitate, carefully avoiding an excess. The liquid is again filtered, carbonate of potassa added, till it ceases to precipitate, and after twenty-four hours the precipitate is collected on a filter and dried. It is then dissolved in concentrated alcohol, treated with animal charcoal and filtered, evaporated and crystallized. The dose is the fortieth or thirtieth of a grain.

Soubiran.

SOLUTION OF ATROPIA.

- R. Atropia, (one, two, or three grains, as the strength may wish to be increased), one grain.
 Nitric or acetic acid, sufficient.
 Water, one fl. ounce.

Dissolve.

As an application to the eyes for dilating the pupil. One or two drops will generally dilate the pupil, if dropped between the lids.

SYRUP OF ATROPIA.

- R. Atropia, one part.
 Syrup, ten thousand parts.
- Dissolve the atropia in one hundred parts of water acidulated with one part of muriatic acid, then mix with the syrup. *Bouchardat.*

TINCTURE OF ATROPIA.

- R. Atropia, fifteen grains.
 Alcohol, (85 per cent.), ten fl. drachms.
- Mix. Dose, one to three drops. *Bouchardat.*

SULPHATE OF ATROPIA.

- R. Diluted sulphuric acid, two fl. drachms.
 Water, half a fl. ounce.
- Mix, and gradually add Atropia, seven scruples and a half; or sufficient to saturate. Filter and evaporate, that crystals may form. *Lond. Ph.*
- Chiefly used externally, in the form of an ointment.

AURANTIUM.

ORANGE.

This fruit is the product of the *Citrus Aurantium*, a native of Asia, but now cultivated in all warm climates; two varieties occur, the one with bitter, the other with sweet fruit; these are considered distinct by Risso.

Sex. Syst. Polydelph. icosand. *Nat. Syst.* Aurantiaceæ.

Lind. Fl. Med. 163. *Griffith, Med. Bot.* 165.

Almost every part is employed in medicine, but only the rind of the fruit is officinal in the U. S. Pharm.

AURANTII CORTEX.

ORANGE PEEL.

POWDER OF ORANGE PEEL AND RHUBARB.

- R. Powdered orange peel,
 " rhubarb,
 Bitartrate of potassa, each, one ounce.

Mix well. Dose, one or two teaspoonfuls a-day. In dyspepsia, with a costive condition of the bowels. *Radius.*

CONFECTION OF ORANGE PEEL.

- R. Fresh orange peel, grated, one pound
 White sugar, three pounds.
- Beat together, till thoroughly mixed. *U. S. Ph.*
- As a vehicle or adjunct to powders.

ELECTUARY OF ORANGE PEEL.

- R. Orange peel, half an ounce.
 Willow bark, two drachms.
 Syrup of orange peel, sufficient.
- Make an electuary. To be taken in divided doses, during the apyrexia of intermittent fever. *Phæbus.*

TINCTURE OF ORANGE PEEL.

- R. Dried orange peel, three ounces and a half.
 Proof spirit, (Imp.), two pints.
- Digest for fourteen days, and filter. *Lond. Ph.*
- Used principally as an addition to infusions, decoctions, &c.

SPIRIT OF ORANGE PEEL.

- R. Orange peel, three ounces.
 Alcohol, sixteen fl. ounces.
 Water, four fl. ounces.
- Digest for four days, and distil sixteen parts. *Cottureau.*
- Used for the same purposes as the last.

ELIXIR OF ORANGE PEEL.

- R. Orange peel, one ounce.
 Cascarilla, half an ounce.
 Water of citron peel,
 " wormwood,
 Alcohol, each, half a pint.
- Digest for four days, and filter. *Moscatti.*
- Dose, from four to six fl. drachms, as a carminative and stomachic.

COMPOUND INFUSION OF ORANGE PEEL.

- R. Dried orange peel, half an ounce.
 Fresh lemon peel, two drachms.
 Bruised cloves, one drachm.
 Boiling distilled water, one pint.
- Macerate for a quarter of an hour, and strain. *Lond. Ph.*
- This preparation is used as a stomachic, in the dose of one or two fl. ounces.

SYRUP OF ORANGE PEEL.

R. Bruised orange peel, two ounces.
Boiling water, one pint.
Macerate for twelve hours, strain, and add
Sugar, two pounds and a half.
Make syrup. *U. S. Ph.*

Used principally as an agreeable drink mixed with water.

R. Recently-dried sweet orange peel, two ounces.

Powder, and put in a displacement apparatus, and pour on it a mixture of two parts alcohol, and one of water, until six fl. ounces are obtained. Pour this on thirty-two ounces (av.) of coarsely-powdered sugar, and spread on paper, until the alcohol has evaporated. Then form it into a syrup, with sixteen ounces of water, merely carrying the heat to the boiling point, strain, and bottle whilst hot.

W. Procter.

AURANTII FOLIA.

ORANGE LEAVES.

ELECTUARY OF ORANGE LEAVES.

R. Orange leaves,
Valerian, each, half a drachm.
Syrup of orange peel, sufficient.
Make an electuary. *Radius.*

A teaspoonful occasionally as an antispasmodic.

JELLY OF ORANGE LEAVES.

R. Orange leaves, an ounce and a half.
Boiling water, two pints.
Digest for twenty-four hours, strain, and add
Powdered salep, two drachms.
Boil to a jelly, and add
Wine,
Syrup of orange flowers, each, one fl. ounce.
In teaspoonful doses, in marasmus. *Radius.*

AURANTII FLORES.

ORANGE FLOWERS.

CONFECTION OF ORANGE FLOWERS.

R. Orange flowers, one part.
Syrup, three parts.
Evaporate to proper consistence. *Taddei.*

PASTILLES OF ORANGE FLOWERS.

R. Sugar,
Orange-flower water, each, two ounces.

Dissolve and evaporate to consistence of honey; add a mixture of

Sugar, four ounces.
Oil of orange flowers, half a drachm.

Make pastilles. *Cottreau.*

ORANGE-FLOWER WATER.

R. Orange flowers, ten parts.
Water, thirty parts.
Distil off twenty parts. *Guibourt.*

R. Orange flowers, ten pounds.
Proof spirit, seven fl. ounces.
Water, two gallons.

Mix, and distil a gallon. *London Ph. 1836.*

SYRUP OF ORANGE FLOWERS.

R. Orange-flower water, one part.
Sugar, two parts.

Dissolve in a close vessel, and evaporate to proper consistence. *Soubeiran.*

OIL OF ORANGE FLOWERS.

R. Orange flowers, one part.
Water, three parts.

Distil, and separate the oil that floats on the product. *Taddei.*

AURUM.

GOLD.

Gold is used as a remedial agent in its pure state, but only in an infinitely divided form, in which it is stated that it acts on the system, and to be equally efficacious as its salts, but its action is much milder. It has been prescribed in various forms of syphilitic diseases, in doses of a quarter of a grain to a grain, three or four times a-day.

POWDER OF GOLD.

R. Gold leaf, one part.
Sulphate of potassa, eight to ten parts.

Triturate thoroughly, wash out the sulphate of potassa, and sift through fine gauze. *Par. Cod.*

To be applied in friction on the gums and tongue.

R. Dilute solution nitro-muriate of gold, at will.

Solution sulphate of iron, sufficient to precipitate; collect this, and dry. *Tromsdorff.*

R. Pulverized gold, six grains.
Starch, sixty-six grains.

Mix, and divide into twelve powders. One to be given four times a-day. *Riecke.*

R. Pulverized gold,
Lycopodium, each, two grains.

Mix. To be rubbed, during the day, on the tongue. *Riecke.*

SYRUP OF GOLD.

R. Pulverized gold, twenty-four grains.
Syrup, one ounce.

Mix. As an application to chancres on the fauces. *Bories.*

OINTMENT OF GOLD.

R. Pulverized gold, one drachm.
Lard, half an ounce.

Triturate well. As a dressing to venereal ulcers, and also to a surface having the cuticle removed; when the surface becomes dry, substitute the ointment of chloride of gold.

Legrand.

AURI CHLORIDUM.

CHLORIDE OF GOLD.

R. Leaf gold, one part.
Nitro-muriatic acid, three parts.

Dissolve by aid of a sand-bath, and evaporate till vapors of chlorine are given off, and set aside to crystallize. Dose, one-twentieth to one-sixteenth of a grain. *Par. Cod.*

R. Pure gold, one part.
Nitro-muriatic acid, (made with three parts muriatic acid, one part nitric acid, and one of water,) two parts.

Put the gold in a porcelain capsule, and add the acid; cover with a plate of glass, and place on a salt-water bath. Continue heat till nitrous fumes are no longer evolved. Raise the glass by means of a fold of paper, and continue heat, till, on introducing a glass rod, the adhering chloride of gold becomes solid on cooling. Remove from fire, and let crystallize. *Leval.*

POWDER OF CHLORIDE OF GOLD.

R. Chloride of gold, one part.
Powdered orris root, two parts.

Mix. In frictions on the gums, in the dose of a fifteenth to a tenth of a grain. *Foy.*

BOLUS OF CHLORIDE OF GOLD.

R. Chloride of gold, half a grain.
Extract of aconite, twelve grains.

Mix, and make six boluses. One to be taken three times a-day. *Brera.*

PILLS OF CHLORIDE OF GOLD.

R. Chloride of gold, ten grains.
Powdered liquorice, three drachms.
Syrup, sufficient.

Mix, and make one hundred and fifty pills. One, thrice a-day. *Ellis.*

TINCTURE OF CHLORIDE OF GOLD.

R. Leaf gold, one drachm.
Nitro-muriatic acid, one ounce.

Dissolve, and add

Oil of rosemary, two ounces.
Alcohol, eight ounces.

Mix. Dose, ten drops. *Spielmann.*

CAUSTIC OF CHLORIDE OF GOLD.

R. Chloride of gold, one ounce.
Nitro-muriatic acid, one fl. ounce.

Mix. Used as a caustic to the surface of cancerous ulcers, applied by means of a dossil of lint. *Recamier.*

COLLYRIUM OF CHLORIDE OF GOLD.

R. Chloride of gold, two grains.
Distilled water, six fl. ounces.

Dissolve. As a collyrium in ophthalmia.

Jahn.

OINTMENT OF CHLORIDE OF GOLD.

R. Chloride of gold, eight grains.
Lard, half an ounce.

Mix. To relieve rheumatic pains. *Legrand.*

AURI ET SODII CHLORIDUM.

CHLORIDE OF GOLD AND SODIUM.

R. Chloride of gold, eighty-five parts.
Chloride of sodium, sixteen parts.

Dissolve in a small quantity of water, evaporate to a pellicle, and let crystallize. *Par. Cod.*
Dose, one-sixteenth of a grain.

R. Leaf gold, four parts.
Nitro-muriatic acid, sufficient

to dissolve. Evaporate to dryness, and add

Chloride of sodium, one part.
Water, thirty-two parts.

Evaporate to one-half, and set aside to crystallize. *Figuier.*

POWDER OF CHLORIDE OF GOLD AND SODIUM.

R. Chloride of gold and sodium, one part.

Lycopodium, two parts.
Mix. *Cadex.*

R. Chloride of gold and sodium, one part.

Sugar of milk, sixteen parts.
Mix. *Radius.*

R. Chloride of gold and sodium,
three parts.
Powdered orris root, nine parts.
Mix. Three grains contain about three-fourths
of a grain of the chloride, and are used for
thirty frictions, for the weakest doses, and three
for the strongest. *Legrand.*

PILLS OF CHLORIDE OF GOLD AND SODIUM.

R. Chloride of gold and sodium,
ten grains.
Potato starch, four grains.
Gum Arabic,
Distilled water, each, one drachm.
Mix, and make one hundred and twenty pills.
Each contains one-twelfth of a grain of chlor-
ide. *Soubeiran.*

R. Chloride of gold and sodium,
four grains.
Extract of bitter sweet, one drachm.
" aconite, ten grains.
Powdered marsh mallow,
Distilled water, each, sufficient.
Mix, and make eighty pills. Dose, three thrice
a day, in dropsy, incontinence of urine, and
amaurosis, &c. *Groetzner.*

R. Chloride of gold and sodium,
one grain.
Extract of mezereon, one drachm.
Mix, and make sixty pills. *Magendie.*

LOZENGES OF CHLORIDE OF GOLD AND SODIUM.

R. Chloride of gold and sodium,
five grains.
Powdered sugar, one ounce.
Mucilage of tragacanth, sufficient.
Rub the chloride in a glass mortar with the
sugar, add the mucilage, and make sixty
lozenges. *Soubeiran.*

SOLUTION OF CHLORIDE OF GOLD AND SODIUM.

R. Chloride of gold and sodium,
two grains.
Distilled water, one ounce.
Dissolve. Ten drops, every two hours, in dropsy
and incontinence of urine. *Radius.*

SYRUP OF CHLORIDE OF GOLD AND SODIUM.

R. Chloride of gold and sodium,
one grain.
Syrup, six ounces.
Dissolve. Dose, from half an ounce to an
ounce. *Bories.*

COMPOUND SYRUP OF CHLORIDE OF GOLD AND SODIUM.

R. Peruvian bark,
Gentian, each, three ounces.
Mezereon, one ounce and a half.
Foxglove, two drachms and a half.
Water, one pint and a half.

Boil down to one pint, and pour the boiling
decoction on

Bitter orange peel, one ounce.
Cloves, half an ounce.

After a sufficient infusion, strain, and add
Chloride of gold and
sodium, four grains.
A spoonful, night and morning. *Bories.*

OINTMENT OF CHLORIDE OF GOLD AND SODIUM.

R. Chloride of gold and
sodium, one scruple.
Lard, one ounce.
Mix well. *Foy.*
R. Chloride of gold and sodium, three
to four grains.
Lard, half an ounce.
Mix thoroughly. *Groetzner.*

AURI CYANIDUM.

CYANIDE OF GOLD.

R. Leaf gold, one part.
Nitro-muriatic acid, six parts.
Dissolve, and evaporate to dryness. Dissolve in
Distilled water, eight parts.
Reduce on a water-bath to one-fourth, and add
very gradually

Cyanide of potassium, half a part.
Distilled water, twenty-four parts.
Agitate, let rest, and separate the cyanide of gold.
Soubeiran.

R. Fresh precipitated oxide of gold,
(washed, but not dried), at will.
Diluted hydrocyanic acid, sufficient.

Boil till solution assumes a beautiful yellow
tint, evaporate to dryness on a water-bath.
Desfosses.

Dose, from eighteenth to tenth of a grain.

POWDER OF CYANIDE OF GOLD.

R. Cyanide of gold, one grain.
Lycopodium, fifteen grains.
Mix, and divide into sixteen powders. One
every day, as friction on the gums and tongue.
Radius.

PILLS OF CYANIDE OF GOLD.

R. Cyanide of gold, one grain.
Powdered liquorice, thirty-one grains.
Mucilage, sufficient.

Mix, and make sixteen pills. *Radius.*

R. Cyanide of gold, one grain.
Extract of mezereon, three grains.
Powdered marsh mallow, sufficient
to make fifteen pills. Dose, one pill twice
a-day.

LOZENGES OF CYANIDE OF GOLD.

R. Cyanide of gold, one grain.
Chocolate, sufficient
to make sixteen lozenges. *Soubeiran.*

SOLUTION OF CYANIDE OF GOLD.

R. Cyanide of gold, three grains.
Diluted alcohol, eight fl. ounces.
A teaspoonful, twice a-day, gradually increasing
the dose. *Christien.*

AURI IODIDUM.

IODIDE OF GOLD.

R. Solution of chloride of gold, at will.
Solution of iodide of potas-
sium, sufficient

to precipitate; wash the precipitate with alco-
hol, and dry it. *Par. Cod.*

R. Solution chloride of gold, at will.
“ hydriodate of am-
monia, sufficient

to precipitate. Wash with alcohol, and dry. *Millet.*

Used like the other preparations, and in the
same doses.

AURI OXIDUM.

OXIDE OF GOLD.

R. Pure gold, one part.
Mercury, six parts.

Make an amalgam; triturate with double the
weight of sulphur and calcine. *Wirt. Ph.*

R. Leaf gold, half an ounce.
Nitre, ten ounces.
Common salt, five ounces.
Alum, eight ounces.

Triturate to extinction, dissolve in lime-water,
precipitate by solution of potassa; wash the
precipitate, and calcine. *Spielmann.*

R. Chloride of gold, one part.
Distilled water, forty parts.

Dissolve, and add
Fresh calcined magnesia, four parts.

Boil; wash the precipitate with distilled water,
then with nitric acid diluted with twenty parts
of water, and again with water. Dry in the
shade. *Par. Cod.*

Dose, from a tenth to three-fourths of a
grain.

POWDER OF OXIDE OF GOLD.

R. Oxide of gold, two drachms
Sulphuret of antimony, half an ounce.
Oil of cinnamon, eight drops.

Triturate together. *Spielmann.*

R. Oxide of gold, four scruples.
Prepared hartshorn, one ounce.

Triturate together. Used in malignant fevers
and small-pox. *Spielmann.*

PILLS OF OXIDE OF GOLD.

R. Oxide of gold, five grains.
Extract of mezereon, two drachms.

Mix, and make sixty pills. *Magendie.*

AURUM AMMONIATUM.

AMMONIATED GOLD.

R. Gold leaf, two drachms.
Nitro-muriatic acid, one ounce.

Dissolve, and add

Water of ammonia, sufficient
to precipitate. Wash and dry, with great cau-
tion. *Niemann.*

This is what is termed fulminating gold, and
explodes at 400° F. It has been used in fevers,
&c., as a diaphoretic, but has produced unplea-
sant consequences.

PILLS OF AMMONIATED GOLD.

R. Ammoniated gold,
Calomel,
Extract of squill, each, half a drachm.
Powdered rhubarb, two drachms.
Conserve of juniper, sufficient.

Mix, and make pills of two grains. *Plenck.*

AURUM STANNO-PARATUM.

PURPLE OF CASSIUS.

R. Chloride of gold, one part.
Distilled water, two hundred parts.

Add to this solution, very gradually,

Pure tin, one part.
Nitro-muriatic acid, three parts.

Dissolve without heat, and add

Distilled water, one hundred parts,

as long as any precipitate falls. Wash this and dry by a gentle heat. *Par. Cod.*, 1839.

This is used like the other preparations of gold.

AVENA SATIVA.

OAT.

The common oat, although generally cultivated from a very early age, is not known in a wild state.

Sex. Syst. Triand. digyn. *Nat. Syst.* Graminaceæ.

Linn. Sp. Pl. 61. *Griffith, Med. Bot.* 662.

Though, for the most part used as food for horses, it contains much nutritive matter, and forms an important article of diet in some parts of Europe. It is also employed in medicine as a bland, nutritious, and somewhat laxative nutriment in inflammatory diseases. When the grains are merely freed from their husk and coarsely broken, they are called groats.

GROAT GRUEL.

R. Groats, three ounces.

Wash well in cold water, and then put into

Fresh water, four pints.

Boil slowly to one-half, and strain through a fine sieve. *A. T. Thomson.*

R. Groats, five drachms.
Liquorice root, three drachms.

Boil the groats in sufficient water to have one quart of decoction, add the liquorice, and after a sufficient infusion, strain. *Cottureau.*

OATMEAL GRUEL.

R. Oatmeal, two ounces.
Water, one pint and a half.

Rub the meal in a basin, with the back of a spoon, in some of the water, pouring off the fluid after the grosser particles have subsided, but whilst the milkiness remains; repeat this with fresh water. Stir well, and unite the washings, and boil until a soft, thick mucilage is formed. *A. T. Thomson.*

Both these preparations may be flavored according to circumstances.

COMPOUND DECOCTION OF OATMEAL.

R. Groats, six ounces.
Red saunders, chipped, one ounce.
Chicory root, one ounce and a half.
Water, twelve pints.

Boil down one-third, and add to the strained decoction

Nitre, half an ounce.
Sugar, two ounces.

Dissolve. Advised in gout, nephritis, &c., to be

taken hot or tepid, morning and evening, for some weeks, in doses of six or eight fl. ounces. *Wirt. Ph.*

FLUMMERY OF OATMEAL.

R. Oatmeal or groats, a quart.

Rub for a considerable time with two quarts of hot water, and let the mixture stand till it becomes sour, then add another quart of hot water, and strain through a hair sieve. Let stand till a white sediment is deposited, decant the fluid, and wash the sediment with cold water. This is now to be boiled with fresh water, till it forms a mucilage, stirring the whole time. It is a very light and somewhat nutritious food, during early convalescence. *A. T. Thomson.*

POWDER FOR A CATAPLASM.

R. Linseed meal, one part.
Oatmeal, two parts.
Dub. Ph., 1826.

R. Powder for a cataplasm, at will.
Boiling water, sufficient.
Mix for a poultice. *Dub. Ph.*, 1826.

OATMEAL POULTICE.

R. Oatmeal, sufficient.
Stir gradually into water kept boiling, until it is sufficiently consistent. *A. T. Thomson.*

AZEDARACH.

AZEDARACH—(PRIDE OF CHINA.)

The *Melia Azedarach* is a tree, a native of many parts of Asia, and naturalized in the warmer parts of Europe and the United States. *Sex. Syst.* Hexand. monog. *Nat. Syst.* Meliaceæ.

Linn. Sp. Pl., 550. *Griffith, Med. Bot.*, 179.

The parts used are the bark of the root, the berries, and the leaves. The former only is officinal in the U. S. Pharm. It is cathartic and emetic; in large doses somewhat narcotic; it is also an efficient anthelmintic.

DECOCTION OF AZEDARACH.

R. Azedarach, four ounces.
Water, two pints.
Boil to a pint, and strain. *Wood.*
Dose, one fl. ounce, every two or three hours, for a child.

OINTMENT OF AZEDARACH BERRIES.

R. Pulp of berries,
Lard, each, one ounce.
Rub well together. Said to be useful as an application in tinea capitis.

B.

BALLOTA LANATA.

HAIRY HOREHOUND.

Several species of *Ballota* have, at different times, been used in medicine, but this is the only one that appears to possess active properties. *Sex. Syst.* Didynam. gym. *Nat. Syst.* Lamiaceæ.

The whole plant is used, with the exception of the root, and is stated to be an excellent diuretic, and useful in dropsical cases.

DECOCTION OF HAIRY HOREHOUND.

R. Hairy horehound, one ounce and a half.
Boiling water, two pints.

Boil down to one pint. Dose, a cupful twice a day, gradually increasing the quantity.

Rehmann.

BALSAMUM PERUVIANUM.

BALSAM OF PERU.

This balsam is the product of *Myrospermum Peruiferum*, a large tree, indigenous to South America, where it is called *quinquino*.

Sex. Syst. Decand. monog. *Nat. Syst.* Fabacæ.

Linn. Sp. Pl. 233. Griffith, Med. Bot. 248.

The balsam is a thick, viscid fluid, of a dark, reddish-brown color, an agreeable, fragrant odor, and a warm, bitterish taste, occasioning a pungent sensation in the throat. It is stimulating, tonic, and expectorant, and has been much used in pectoral complaints, &c., and also externally as an application to indolent ulcers. The dose is about half a fl. drachm.

PILLS OF BALSAM OF PERU.

R. Balsam of Peru, one drachm.
Extract of bitter polygala, two drachms.
Marsh mallow, sufficient

to make one hundred and twenty pills. Ten, four times a day, in chronic mucous discharges.

Schubert.

ACOUSTIC BALSAM.

R. Balsam of Peru, half a drachm.
Narcotic oil,
Onion juice, each, one ounce.

Mix. A dossil of cotton saturated with this oil, is to be introduced into the deaf ear, provided there is no inflammation or violent pain.

R. Beef gall, three fl. drachms.
Balsam of Peru, one fl. drachm.

Mix. To be occasionally dropped into the ear to correct a fetid discharge, syringing it also, daily, with a weak solution of soap and water.

Hugh Smith.

LINIMENT FOR CHILBLAINS.

R. Balsam of Peru, half a drachm.
Muriatic ether,
Laudanum, each, two drachms.

Mix. As a friction. *Henschel.*

MAMILLARY LOTION.

R. Balsam of Peru, one drachm.
Yolk of egg, one.
Spirit of wild thyme, three ounces.

Mix. The sore nipple is to be bathed with this, and then sprinkled with a powder, composed of one drachm of Peruvian bark, and two drachms of gum Arabic.

Iverg.

BALSAM OF PERU MIXTURE.

R. Balsam of Peru, two drachms.
Yolk of egg, one.
Extract of cinchona, two drachms.
Honey of roses, three ounces.

Mix. Two dessertspoonfuls, four times a day in chronic mucous discharges.

St. Marie.

R. Balsam of Peru, half a drachm.
Mucilage of gum Arabic, sufficient.
Cinnamon water,
Water, each, half a fl. ounce.

Mix. To be taken three or four times a day, as an expectorant in chronic catarrh.

Ellis.

TINCTURE OF BALSAM OF PERU.

R. Balsam of Peru, one part.
Alcohol, eight parts.

Macerate for some days, and filter.

Guibourt.

Principally used as an external application to ulcers, &c.

SYRUP OF BALSAM OF PERU.

R. Tincture of balsam of Peru, one fl. ounce.

Tepid simple syrup, two pounds.

Mix, and agitate briskly. Dose, from one to two drachms.

Lisbon Ph.

BALSAM OF PERU COLLUTORY.

R. Tincture of balsam of Peru, one part.
" guaiacum, four parts.

Mix. A teaspoonful to a glass of water, to rinse the mouth.

Taddei.

LOCATELLI'S BALSAM.

- R. Olive oil, six ounces.
 Yellow wax, four ounces.
 Wine, five fl. ounces.

Melt together by a gentle heat, till all moisture is evaporated, and add

- Venice turpentine, six ounces.
 Balsam of Peru, two drachms.
 Red saunders, half an ounce.

Mix. Formerly used in phthisis, but now employed only as an external application. This is the original formula. *Spielmann.*

OINTMENT OF BALSAM OF PERU.

- R. Balsam of Peru,
 Spermaceti ointment, equal parts.

Mix. As a dressing to painful ulcers. *Radius.*

COMPOUND OINTMENT OF BALSAM OF PERU.

- R. Lard, two ounces.
 White wax, four drachms.

Melt in a water-bath, and add

- Balsam of Peru, two drachms.
 Oil of lavender, twelve minims.

As an application to promote the growth of the hair. *Copland.*

PLASTER OF BALSAM OF PERU.

- R. Powdered carbonate of lead, sixteen parts.
 " litharge, eight parts.
 Rose oil, forty-eight parts.
 White wax, thirty-two parts.

Melt together, and, at close of operation, add

- Balsam of Peru, two parts.

As an application to indolent ulcers. *Foy.*

BAPTISIA TINCTORIA.

WILD INDIGO.

A native plant, found in most parts of the country; becoming black when dried.

Sex. Syst. Decand. monog. Nat. Syst. Fabaceæ.

Torrey and Gray, *Flor. i.* 386. Griffith, *Med. Bot.* 231.

The root, which is the part used, is emetic and purgative when fresh. Has been considered a valuable antiseptic and febrifuge, and has been given with advantage in typhus fever, scarlatina, &c., and also used as a wash to foul ulcers, aphthæ, &c. Most employed in decoction.

DECOCTION OF WILD INDIGO.

- R. Root of wild indigo, one ounce.
 Boiling water, a pint and a half.
 Boil down to a pint. Dose, half a fl. ounce every four to eight hours. *Comstock.*

OINTMENT OF WILD INDIGO.

- R. Contused root of wild indigo, two ounces.
 Lard, six ounces.

Simmer together for an hour and strain. Has been found beneficial as an application to burns and ulcers.

BARIUM.

BARIUM CHLORIDUM.

CHLORIDE OF BARIUM.

- R. Carbonate of baryta, one pound.
 Muriatic acid, twelve fl. ounces.
 Water, three pints.

Mix the acid with the water, and gradually add the baryta; towards close of effervescence, apply a gentle heat, and, when action has ceased, filter, and boil down, for crystals to form.

U. S. Ph.

POWDERS OF CHLORIDE OF BARIUM.

- R. Chloride of barium, two drachms.
 Calomel, ten grains.
 Sulphuret of antimony, six grains.

Mix, and divide into forty-eight powders. Dose, two a-day in syrup, in cutaneous affections.

Swediaur.

PILLS OF CHLORIDE OF BARIUM.

- R. Chloride of barium, two drachms.
 Extract of liquorice, each, half a drachm.

Powdered liquorice root, sufficient.
 Water, each, sufficient.

Mix, and divide into one hundred and twenty pills. *Phæbus.*

Dose, four to eight, three or four times a-day.

- R. Chloride of barium, one drachm.
 Resin of guaiacum, half an ounce.
 Conserve of fumitory, sufficient.

Mix, and make one hundred and eighty-eight pills. One to be taken morning and evening, increased to two, against tapeworm. *Pierquin.*

SOLUTION OF CHLORIDE OF BARIUM.

- R. Chloride of barium, one ounce.
 Distilled water, three fl. ounces.

Dissolve, and filter. *U. S. Ph.*

Has been used in small doses in cancer and scrofula. Dose, five drops, two or three times a-day, cautiously increasing.

COLLYRIUM OF CHLORIDE OF BARIUM.

R. Chloride of barium, ten grains.
Distilled water, one ounce.

Dissolve, filter, and add

Mucilage of quince seeds, two drachms.
Wine of opium, half a drachm.

Radiu

The eyelids are to be washed several times a-day with this, in scrofulous ophthalmia.

MIXTURE OF CHLORIDE OF BARIUM.

R. Chloride of barium,
" iron, each, half a drachm.
Distilled water, one ounce.

Dissolve. *Augustin.*

Dose, twenty to sixty drops.

R. Chloride of barium,
Extract of cicuta, each, half a drachm.
Distilled water, half an ounce.
Antimonial wine, one drachm and
a half.

Phæbus.

Dose, fifteen to twenty drops every three hours, in chronic orchitis.

R. Chloride of barium, one drachm.
Balm water, seven fl. drachms.
Antimonial wine, one fl. drachm.

Dose, twenty to thirty drops four times a-day.
Hufeland.

R. Chloride of barium, four grains.
Distilled water, two fl. ounces.
Extract of hemlock, four grains.
Common emulsion, one pound.
Syrup, one ounce.

Mix. To be taken during the day, in scrofula, and scrofulous phthisis. *Brera.*

BARIUM IODIDUM.

IODIDE OF BARIUM.

R. Iodine, one hundred parts.
Iron filings, thirty parts.
Water, sufficient.

Prepare an iodide of iron, add baryta dissolved in twenty parts of water, as long as a precipitate is formed, heat a moment, filter, evaporate, and crystallize. *Magendie.*

Been used with success in scrofula. Dose, $\frac{1}{2}$ of a grain, three times a-day, cautiously increasing.

POWDER OF IODIDE OF BARIUM.

R. Iodide of barium, one grain.
Powdered cinnamon,
Sugar, each, four scruples.

Mix well, and divide into eight powders. One to be given two or three times a-day in scrofula. *Radiu.*

OINTMENT OF IODIDE OF BARIUM.

R. Iodide of barium, four grains.
Lard, one ounce.

Mix. As a friction in scrofulous swellings. *Bielt.*

BARIUM SULPHURETUM.

SULPHURET OF BARIUM.

R. Sulphate of baryta, eleven parts.
Charcoal, one part.
Oil of turpentine, sufficient.

Triturate the baryta with the coal, moisten the mixture with the turpentine, and heat the whole in a crucible to redness; let cool, and preserve. *Van Mons.*

BARYTA.

BARYTES.

R. Nitrate of baryta, sufficient.
Mix in a platina crucible to a red heat. When the mass has become solid and porous, raise to a white heat, remove from fire, and cool. *Van Mons.*

SOLUTION OF BARYTES.

R. Baryta, one part.
Distilled water, twenty parts.
Dissolve. *Van Mons.*

Has been recommended in scrofula, in doses of four to five drops, in some appropriate vehicle.

BARYTIC LINIMENT.

R. Solution of baryta, one part.
Olive oil, six parts.
Rub together. *Soubéiran.*

Advised as an external application in lepra, and other obstinate cutaneous diseases.

BARYTÆ ACETAS.

ACETATE OF BARYTA.

R. Carbonate of baryta, at will.
Acetic acid, sufficient
to saturate; filter, evaporate, and crystallize. *Van Mons.*

SOLUTION OF ACETATE OF BARYTA.

R. Acetate of baryta, one part.
Distilled water, nine parts.
Dissolve, and filter. In same cases and doses as the solution of baryta. *Hamb. Ph.*

BEBEERINA.

BEBEERINE.

This is an alkaloid obtained from the bark of a tree, growing in British Guiana. It is said to belong to the genus *Nectandra*, and has been named *N. rodai*. The bark is in flat pieces, smooth, grayish, hard, heavy, and brittle, with but little odor, though of a very bitter taste. Bebeerine is extracted from this bark in the form of a sulphate, by a process similar to that used to obtain sulphate of quinia. In this form it contains both bebeerine and siperina, and is in thin, somewhat glittering scales of a brownish-yellow color, forming a yellow powder, soluble in cold water, but often forming a turbid solution, which is rendered clear by a few drops of diluted sulphuric acid. Pure bebeerine can be obtained from this solution as follows:—

Decompose by ammonia, wash the precipitate, and whilst moist, triturate with moist hydrated oxide of lead; dry on a water-bath, exhaust with alcohol, and distil off the spirit, treat the residue with ether; on the evaporation of the ether, bebeerine will be left of a bright canary-yellow color, but in powder appears nearly white.

Dose of the sulphate is one to three grains as a tonic, and five grains to a scruple as a febrifuge.

It is not equal to quinine as an antiperiodic, but is a good substitute for that article.

PILLS OF SULPHATE OF BEBEERINE.

R. Sulphate of bebeerine, two drachms.
Conserve of roses, sufficient.

Mix, and form twenty-four pills—one to three, thrice a-day in intermittent fevers. *Christison*.

SOLUTION OF SULPHATE OF BEBEERINE.

R. Sulphate of bebeerine, half a drachm.
Diluted sulphuric acid,
twenty-five minims.

Syrup,
Tincture of orange peel, each,
one fl. ounce.

Water, four fl. ounces.

Mix. A tablespoonful three times a-day.
Christison.

BECCABUNGA.

BROOKLIME.

Two species of *Veronica* of similar properties are included under this name, the *V. beccabunga* and *V. anagallis*; they are both semi-aquatic plants, indigenous to Europe and to this country.

Sex. Syst. Diand. monog. *Nat. Syst.* Scrophulariaceæ.

Griffith, Med. Bot. 517.

These plants are employed usually in a fresh state, and considered to be antiscorbutic and

alterative. They are usually given in infusion made with one or two handfuls of the herb to a pint of boiling water. They have also been employed in conserve, syrup, &c.

WATER OF BROOKLIME.

R. Brooklime,
Water, each, two parts.
Distil off one part. *Cottureau*.
Dose, one to four ounces.

DECOCTION OF BROOKLIME.

R. Fresh brooklime, three ounces.
Water, one pint.
Boil for fifteen minutes and strain. *Copland*.

SYRUP OF BROOKLIME.

R. Clarified juice of brooklime, one part.
Sugar, two parts.
Dissolve, by means of a water-bath. *Taddei*.

BELLADONNA.

BELLADONNA.

This is the *Atropa Belladonna*, an herbaceous plant, native of Europe, and cultivated in some places in the United States.

Sex. Syst. Pentand. monog. *Nat. Syst.* Solanaceæ.

Linn. Sp. Pl. 260. Griffith, Med. Bot. 486.

The whole plant is officinal in some Pharmacopœias, but the leaves only are ordered in that of the U. S. This plant is a powerful narcotic, owing to the presence of a peculiar principle called *Atropia*. (See page 134.) It has been used in a variety of diseases as an anodyne, antispasmodic, and discutient. The dose of the powdered leaves is one to two grains, daily or twice a-day, gradually increasing. But it is more usually given in the form of an extract.

POWDER OF BELLADONNA.

R. Powdered belladonna root, three grains.
“ ipecacuanha, two grains.
Oxide of zinc, six grains.
Sugar, one drachm.

Rub together, and divide into six powders. One every two hours in epilepsy. *Radiis*.

R. Powdered belladonna root, two grains.
“ ipecacuanha, one grain.
Washed sulphur, thirty-two grains.
Sugar of milk, two drachms.

Rub well together, divide into twenty powders. One every three hours in whooping-cough. *Phæbus*

- R. Powdered belladonna, one grain.
 " nitrate of potassa, twenty-one grains.
 " sugar, nine grains.

Make a powder, to be taken at bedtime.

In chronic rheumatism, extensive ulcerations, mania, and epilepsy. *A. T. Thomson.*

- R. Powdered root of belladonna,
 " ipecacuanha, each, six grains.
 " liquorice root,
 " sugar, each, half a drachm.
 Precipitated sulphur, two scruples.
 Oil of anise,
 " amber, each, three minims.

Mix, and make five to twenty powders.

A. T. Thomson.

- R. Powdered leaves of belladonna,
 one to three grains.

- Musk,
 Camphor, each, five grains.
 Sugar, thirty grains.

Triturate well together, and divide into eight powders. *A. T. Thomson.*

POWDER OF BELLADONNA AND RHUBARB.

- R. Powdered belladonna leaves,
 ten grains.
 " rhubarb, two scruples.

Mix, and divide into ten powders. One powder, two or three times a-day, in obstructions of the liver and spleen. *Radius.*

OPIATED BELLADONNA LEAVES.

- R. Belladonna leaves,
 Water, each, two ounces.
 Extract of opium, one drachm.

Dissolve the opium in the water, wash the leaves with the solution, and dry them by a gentle heat. *Guibourt.*

Said to be very useful in phthisis, the patient to smoke a pinch every morning in a common pipe.

FUMIGATION OF BELLADONNA.

- R. Powdered belladonna, one drachm.
 Boiling infusion of sage, one quart.

The steam to be inhaled. Said to be useful in hooping-cough and phthisis. *Soubeiran.*

EXTRACT OF BELLADONNA.

- R. Belladonna leaves, one pound.
 Bruise in a stone mortar, with the addition of a little water; express the juice, heat to boiling point, strain, and evaporate to proper consistence. *U. S. Ph.*

Dose, half to one grain night and morning, to be gradually increased.

ALCOHOLIC EXTRACT OF BELLADONNA.

- R. Belladonna leaves, one pound.
 Diluted alcohol, four pints.

Moisten the leaves with half a pint of alcohol, let stand for twenty-four hours, put in a displacement apparatus, and add gradually the remainder of alcohol. When this has penetrated the belladonna, add water occasionally so as to keep the powder covered. Stop the process when the liquid that passes forms a precipitate in what has already passed. Distil off alcohol, and evaporate to proper consistence.

U. S. Ph.

Dose, quarter to half a grain, to be gradually increased.

COMPOUND BELLADONNA PILLS.

- R. Extract of belladonna,
 Blue pill,
 Powdered ipecacuanha, each, twelve grains.

Mix, and make twelve pills. One to be taken morning and evening, in cancerous affections. *Ainslie.*

PILLS OF BELLADONNA AND CAMPHOR.

- R. Camphor, three drachms.
 Assafetida, three drachms.
 Extract of belladonna, one drachm.
 Extract of opium, fifteen grains.
 Syrup of gum Arabic, sufficient.

Mix, and make one hundred and twenty pills. Give one pill the first day, two the second, and so on till six are given daily, in hysteria. *Debreyne.*

SOLUTION OF EXTRACT OF BELLADONNA.

- R. Extract of belladonna, three grains.
 Cinnamon water, one fl. ounce.

Make a solution. Dose for a child under one year, two or three drops twice or thrice a day, and an additional drop for each additional year. *Ellis.*

Said to have proved efficacious as a preventive of scarlatina.

INFUSION OF BELLADONNA.

- R. Belladonna leaves, four grains.
 Boiling water, two fl. ounces.

Infuse. Take one-half as a dose. *Paris.*

- R. Belladonna leaves, half a drachm.
 Water, seven fl. ounces.

Infuse, strain, and add

Compound tincture of cardamoms,
 one fl. ounce.

Dose, a tablespoonful. *Saunders*

SYRUP OF BELLADONNA.

R. Extract of belladonna, forty-eight grains.
 Distilled water, two fl. ounces.
 Dissolve and filter, then add
 Syrup, three pounds.
 Boil, and strain. *Cottureau.*
 Said to be useful in whooping-cough, in doses
 of one to three drachms.

GILLET'S SYRUP OF BELLADONNA.

R. Dried belladonna leaves, one ounce.
 Digitalis,
 Common nightshade, each,
 one drachm.
 Water, ten fl. ounces.
 Digest for two days, and filter, then add
 Syrup, two pounds. *Cadet.*
 Advised in whooping-cough, in the dose of a
 teaspoonful every hour or two.

TINCTURE OF BELLADONNA.

R. Belladonna leaves, four ounces.
 Diluted alcohol, two pints.
 Macerate for fourteen days, express, and filter. *U. S. Ph.*
 The dose is from fifteen to thirty drops.

ETHEREAL TINCTURE OF BELLADONNA.

R. Belladonna leaves, dried, one part.
 Sulphuric ether, eight parts.
 Macerate for eight days, and filter. *Guibourt.*

BELLADONNA MIXTURE.

R. Extract of belladonna, eight to twelve grains.
 Wine of ipecacuanha, one fl. drachm.
 Syrup of seneka, half fl. ounce.
 " tolu,
 three and a half fl. ounces.
 Mix. Advised in catarrh, in the dose of a tea-
 spoonful, three or four times a-day, using also
 gum water acidulated with lemon-juice. *Ellis.*
 R. Extract of belladonna, fifteen grains.
 Fennel water, five fl. ounces.
 Alcohol, one fl. drachm.
 Dissolve. Dose, a drop for every year of the
 age of a child, not to exceed fifteen; as a pre-
 ventive of scarlatina. *Maisier.*

LINIMENT OF BELLADONNA.

R. Extract of belladonna, two scruples.
 Sulphuric ether, one fl. drachm.
 Cherry laurel water, two fl. ounces.

As a friction to the abdomen in colica picto-
 num. *Phæbus.*

CLYSTER OF BELLADONNA.

R. Belladonna leaves, twelve to twenty grains.
 Powdered jalap, one scruple.
 Boiling water, half a pint.
 Infuse, and strain. *Pitschaft.*
 This has been recommended in strangulated
 hernia, to overcome spasm.

PLASTER OF BELLADONNA.

R. Resin plaster, three ounces.
 Extract of belladonna, one ounce and a half.
 Melt plaster by means of a water-bath, and add
 the extract; mix well. *U. S. Ph.*
 An efficacious application in neuralgic and
 rheumatic pains.

OINTMENT OF BELLADONNA.

R. Extract of belladonna, two drachms.
 Distilled water, two fl. ounces.
 Lard, two ounces.
 Rub well together. Used by Chaussier as an
 application to the neck of the uterus in cases
 of rigidity. *Ratier.*
 R. Fresh belladonna leaves, one part.
 Lard, two parts.
 Simmer together, express, and strain. Used in
 frictions of one or two drachms, to indolent
 tumors. *Soubéiran.*
 R. Extract of belladonna, one drachm.
 Lard, one ounce.
 Mix. *U. S. Ph.*

BENZOINUM.

BENZOIN

Is the concrete juice of the *Styrax Benzoin*, a
 tree of some size, native of several of the East
 Indian Islands.

Sex. Syst. Decand monog. *Nat. Syst.* *Styracaceæ.*

Dryander, Ph. Trans. lxxvii. 308. *Griffith, Med. Bot.* 439.

There are several varieties of benzoin, but
 the best is in whitish tears, united by a reddish-
 brown connecting medium. It has an aromatic,
 agreeable odor, and a somewhat acrid taste. It
 is stimulant and expectorant, and is much used
 in chronic catarrhs.

BENZOATED LARD.

R. Benzoin in coarse powder, one ounce.
 Lard, twenty-five ounces.
 Heat them together in a water-bath for two or
 three hours, strain without pressure, and stir
 while cooling.

Employed for ointments to be long kept, to prevent them from becoming rancid.

Deschamps.

FUMIGATING POWDER.

R. Powdered olibanum, two pounds.
 " benzoin,
 " storax, each, half a pound.
 " dried roses,
 " lavender flowers, each,
 six ounces.

Mix. A small quantity to be thrown on hot coals, to raise a smoke; to be inhaled in whooping-cough, &c.

Dohrn.

FUMIGATING PASTILLES.

R. Benzoin, ten parts.
 Charcoal, twenty-four parts.
 Nitrate of potassa, one part.
 Sassafras, two parts.
 Mucilage of gum Arabic, sufficient
 to make pastilles, which are to be conical.

Beral.

R. Benzoin, sixteen parts.
 Sandal wood, four parts.
 Laudanum, one part.
 Balsam of tolu, four parts.
 Charcoal, forty-eight parts.
 Nitrate of potassa, two parts.
 Mucilage of tragacanth, sufficient.

Make conical pastilles.

Foy.

POWDER OF BENZOIN.

R. Powdered benzoin,
 " sugar candy, equal parts.

Mix. Said to be useful with camphor water in asthma and chronic catarrh.

Pierquin.

OIL OF BENZOIN.

R. Benzoin, at will.
 Water, a small quantity.

Distil on a sand-bath, and separate the oil that passes over.

Advised as a friction in neuralgic and rheumatic pains.

Swedliaur.

COSMETIC WASH OF BENZOIN.

R. Tincture of benzoin, three fl. drachms.
 Liquid subcarbonate of
 potassa, two fl. drachms.
 Rose water, eight fl. ounces.
 Mix. As a lotion in acne.

Augustin.

COSMETIC EMULSION OF BENZOIN.

R. Almond paste, two drachms.
 Rose water, eight fl. ounces.

Make an emulsion, and add

Tincture of benzoin, three fl. drachms.

Schubarth

MILK OF ROSES.

R. Tincture of benzoin, one fl. drachm.
 Rose water, one pint.
 Mix.

Taddei.

R. Tincture of benzoin, one fl. drachm.
 " of tolu, twenty drops.
 Rose water, one pint.

Mix.

Giannini.

Both these are used as cosmetic washes.

LOTION FOR BURNS.

R. Benzoin, six drachms.
 Storax, four drachms.
 Balsam tolu, two drachms.
 Aloes, one drachm.
 Alcohol, eight fl. ounces.

Make a tincture, to be used as a lotion for burns and scalds, before vesication has taken place.

TINCTURE OF BENZOIN.

R. Benzoin, one part.
 Alcohol, eight parts.

Digest for some days, and filter.

Guibourt.

Used as a stomachic, carminative, &c., in doses of ten to twenty drops.

COMPOUND TINCTURE OF BENZOIN.

R. Benzoin, three ounces
 Purified storax, two ounces.
 Balsam tolu, one ounce.
 Powdered aloes, half an ounce.
 Alcohol, two pints.

Macerate for fourteen days, and filter.

U. S. Ph.

A stimulating expectorant, and also used as an application to indolent ulcers. Dose, twenty to sixty drops.

TURLINGTON'S BALSAM.

R. Benzoin, twelve ounces.
 Liquid storax, four ounces.
 Balsam of Peru, two ounces.
 Myrrh,
 Aloes, each, one ounce.
 Balsam of tolu,
 Extract of liquorice, each, four ounces.
 Angelica root, half an ounce.
 Alcohol, eight pints.

Digest for ten days, and strain. Used as a vulnerary.

Phil Coll. Pharm

BERBERIS.

BARBERRY.

One species of this genus, *B. vulgaris*, has been employed in medicine in Europe. It is a native of Europe, but is naturalized in some parts of the U. S.

Sex. Syst. Hexand. monog. *Nat. Syst.* Berberidaceæ.

Torrey and Gray, Fl. i. 49. Griffith, Med. Bot. 112.

The parts used are principally the berries, which are acidulous, and form a substitute for tamarinds, in the preparation of cooling drinks. The bark of the root is bitter and astringent, and is useful in the treatment of apthous sore mouth, and was at one time much esteemed in the treatment of jaundice.

A bitter, crystallizable principle, called berberin, has been obtained from the root. This is tonic in doses of two or three grains, and laxative in doses of ten to fifteen grains.

LEMONADE OF BARBERRIES.

R. Juice of berries, one part.
Sweetened water, fifteen parts.
Mix. As a refreshing drink in fevers. *Beral.*

INFUSION OF BARBERRY.

R. Barberry bark, one ounce.
Boiling water, one pint.
Macerate for two hours.
Used in jaundice. Dose, one fl. ounce.
Copland.

BISMUTHUM.

BISMUTH.

BISMUTHI SUBNITRAS.

SUBNITRATE OF BISMUTH.

R. Purified bismuth, one ounce.
Nitric acid, (*Sp. Gr.* 1.42), two fl. ounces.
Diluted water, sufficient.
Mix one fl. ounce of water with the acid, and dissolve the bismuth in the mixture; when action has terminated, pour clear solution into three pints of the water, and permit precipitate to subside; pour off supernatant fluid, wash the precipitate well, and dry it by a gentle heat.
U. S. Ph.

Used as a tonic and antispasmodic, especially in certain painful affections of the stomach. Dose, five grains, twice or thrice a day, gradually increasing the quantity.

COMPOUND BISMUTH POWDER.

R. Subnitrate of bismuth, twelve grains.
Powdered ipecacuanha, two grains.
Carbonate of magnesia, two drachms.

Mix, and form twelve powders. One, two or three times a-day, in gastrodynia. *Clarus.*

R. Subnitrate of bismuth, eight grains.
Opium, one grain.
Sugar, two drachms.

Mix, and form four powders. *Ammon.*
Much praised by the author in cholera; one powder to be taken every two hours.

POWDERS OF SUBNITRATE OF BISMUTH.

R. Subnitrate of bismuth, three to six grains.
Sugar, ten grains.

Make six powders. Place one powder on the tongue of a child, (one year old), three or four times daily. Used in the diarrhœa of children. *Trousseau.*

PILLS OF SUBNITRATE OF BISMUTH.

R. Subnitrate of bismuth, one drachm.
Mucilage of gum Arabic, sufficient.
Mix, and make thirty pills. One to be given every two hours in dyspepsia. *Ellis.*

SUBNITRATE OF BISMUTH OINTMENT.

R. Subnitrate of bismuth, one part.
Lard, three parts.
Rub well together. Said to be useful in psora, and other cutaneous eruptions. *Kerksig.*

BISTORTA.

BISTORT.

This is the root of the *Polygonum Bistorta*, a native of Europe, and also found in the northern parts of this country, growing in wet situations. *Sex. Syst.* Octand. trigyn. *Nat. Syst.* Polygonaceæ.

Linn. Sp. Pl. 516. Lindley, Fl. Med. 361.

The part employed, is the root; this is bitter and austere, especially in a fresh state. It is a powerful astringent, but is seldom used in this country. The dose of the powder is twenty or thirty grains, three or four times a-day.

INFUSION OF BISTORT.

R. Bistort, one ounce.
Boiling water, two pints.
Infuse for two hours, and strain. Recommended in passive hemorrhages. *Radius*

ASTRINGENT CLYSTER.

R. Bistort, one ounce.
Poppy heads, two drachms.
Water, one pint.
Infuse, and strain. As an injection, in diarrhœa and chronic dysentery. *Guibourt.*

EXTRACT OF BISTORT.

R. Powdered bistort, one pound.
Water, sufficient.

Exhaust by the process of displacement, and evaporate the filtered fluid. *Pharm. Hosp. Mil.*

MIXTURE OF BISTORT.

R. Powdered bistort, two drachms.
Syrup of quince, one ounce.
Rub together, and add
Tincture of catechu, two fl. drachms.
Water, four fl. ounces.

A spoonful every hour, shaking the bottle each time, in passive hemorrhages, and atonic mucous discharges. *Foy.*

BOLETUS LARICIS.

WHITE AGARIC.

A parasitic fungus growing on the trunk of the larch, in many parts of Europe. It is destitute of smell, but the taste, at first vapid and farinaceous, becomes bitter, acrid, and nauseous. Formerly much used as a purgative, in doses of half a drachm to a drachm. It has lately been thought to be very efficacious in arresting the colliquative sweats in phthisis.

POWDER OF AGARIC.

R. Powdered agaric, twelve grains.
White sugar, two drachms.
Mix well, and divide into six powders. One to be given at night, against colliquative sweats. *Radius.*

POWDER OF AGARIC AND OPIUM.

R. Powdered agaric, eighteen grains.
" opium, three grains.
" gum Arabic, two drachms.
Mix, and divide into nine powders. One to be given at night, against colliquative sweats. *Radius.*

PILLS OF AGARIC AND OPIUM.

R. Powdered agaric, fifteen grains.
Extract of opium, two grains and
 a half.
Mix, and form six pills. One or two at bedtime, in same cases as above noticed. *Rayer.*

BRAYERA ANTHELMINTICA.

KOUSO.

Kousso is the flower of the *Brayera Anthelmintica*, an Abyssinian tree about twenty feet high.

Sex. Syst. Icosand. digyn. *Nat. Syst.* Rosaceæ.

Griffith, Med. Bot. 272.

The flowers are the only parts employed. When dried, they are greenish-yellow; taste feebly acrid, and unpleasant; the odor evolved by boiling is very fragrant. They are exclusively used for expelling the tapeworm.

INFUSION OF KOUSO.

R. Flowers of kousso, half an ounce.
Boiling water, ten fl. ounces.

Macerate for half an hour.

A little lemon-juice to be swallowed, and the infusion being stirred up, the whole is taken, liquid and powder, at two or three draughts, at short intervals, being washed down with cold water and lemon-juice. To promote the operation, tea may be taken. In three or four hours, if the remedy has not operated, a dose of castor oil, or a saline purgative should be administered.

Jon. Pereira.

BROMINIUM.

BROMINE.

This elementary substance has much analogy to iodine in its chemical characters, and in its action on the system. It is a dark red, volatile liquid, having a very caustic taste and a disagreeable odor. It has been employed in bronchocæle, scrofula, chronic cutaneous diseases, and hypertrophy of the heart.

SOLUTION OF BROMINE.

R. Bromine, one part.
Distilled water, forty parts.
Mix. Dose, six drops several times a-day, gradually increasing the dose. *Pourché*

ALCOHOLIC SOLUTION OF BROMINE.

R. Bromine, ten drops.
Alcohol, one ounce.
Mix. As an external application. The strength to be increased five drops to the ounce, daily. *Fournet.*

LOTION OF BROMINE.

R. Bromine, twenty to thirty drops.
Water, one pint.
Mix. For scrofulous ulcers. *Glover*
R. Bromine, four fl. drachms.
Water, five fl. ounces.
Mix. *Pourché.*

BRUCIA.

BRUCINE.

R. Powdered false Augustura bark, at will.

Treat it three times with water, acidulated with muriatic acid, mix the liquids, evaporate, add milk of lime, wash the precipitate, dry, and

treat it with alcohol; evaporate this, and combine the residue with sulphuric acid, dissolve the salt in water, treat with animal charcoal, crystallize, redissolve in water, and precipitate by means of ammonia. *Cottureau.*

A highly poisonous alkaloid, obtained from the bark of the *strychnos nux vomica*, or false Angustura bark. It is white, very bitter, and readily soluble in alcohol, but with difficulty in water. Acts on the system like strychnia, but with less energy, and has been given in same class of diseases. Dose, one-quarter to half a grain.

PILLS OF BRUCIA.

R. Brucia, twelve grains.
Conserve of roses, half a drachm.

Mix, and divide into twenty-four pills. One to be given morning and evening, gradually increasing the dose. *Foy.*

SOLUTION OF BRUCIA.

R. Brucia, six grains.
Distilled water, four fl. ounces.
Sugar, two drachms.

Mix. A tablespoonful, morning and evening. *Magendie.*

TINCTURE OF BRUCIA.

R. Brucia, eighteen grains.
Alcohol (0.847), one fl. ounce.

Dissolve. Six to twenty-four drops in some demulcent drink. *Soubéiran.*

BRUCIÆ ACETAS. MURIAS. ET SULPHAS.

ACETATE, MURIATE, AND SULPHATE OF BRUCIA.

These are all made by the same process, using for each the appropriate acids.

R. Brucia, at will.
Acetic acid, sufficient.

Put the brucia in a porcelain capsule on a water-bath; pour a small quantity of water on it, and then add the acid very gradually, constantly stirring till perfect saturation takes place; filter, evaporate to one-half, and then crystallize. *Cottureau.*

Used for the same purposes as the pure alkaloid, and in like doses.

BRYONIA.

BRYONY.

The roots of two species of bryony are used in medicine, the *B. alba*, and *B. dioica*, the first in England, the latter in other parts of Europe;

but are identical in their sensible properties and action on the system.

Sex. Syst. Monœc. syngen. *Nat. Syst.* Cucurbitaceæ.

Linn. Sp. Pl. 621. Griffith, Med. Bot. 311.

The part used is the root; this is acrid and purgative, causing copious, watery stools, in doses of ten grains to two scruples. It has also some reputation as a cataplasm.

CATAPLASM OF BRYONY.

R. Juice of bryony root,
Crumb of bread, each, sufficient
to form a cataplasm. Used as an application to engorged glands of the neck. *Barthez.*

COMPOUND CATAPLASM OF BRYONY.

R. Rased bryony root, three ounces.
Elder flowers, one ounce.
Ammoniac, half an ounce.
Muriate of ammonia, two drachms.
Conium leaves, two ounces.
Vinegar, sufficient.

Mix, and heat. An application to scrofulous tumors. *Plench.*

WINE OF BRYONY.

R. Bryony root, one ounce.
White wine, one pint.
Boil gently, and filter. To be taken in wine-glassfuls, in dropsies. *Hufeland.*

BUCHU.

BUCHU.

Buchu consists of the leaves of several species of *Diosma*, or more properly *Barosma*, especially of *B. crenata* and *B. seratifolia*. They are small shrubs, natives of the Cape of Good Hope.

Sex. Syst. Pentand. monog. *Nat. Syst.* Rutaceæ.

Linn. Sp. Pl. (*Diosma*) 287. Griffith, Med. Bot. 191.

These leaves have a strong, somewhat aromatic odor, and a bitterish, aromatic taste. They are gently stimulant and diuretic, and are much used in diseases of the urinary organs, and also in some other complaints. The dose, in substance, is from twenty to thirty grains.

INFUSION OF BUCHU.

R. Buchu leaves, one ounce.
Boiling water, one pint.
Macerate for two hours, in a covered vessel, and strain. *U. S. Ph.*

The dose is from one to two fluid ounces.

COMPOUND INFUSION OF BUCHU.

- R. Buchu leaves,
 Uva ursi, each, half an ounce.
 Boiling water, eight fl. ounces.
 Digest for half an hour, strain, and add
 Syrup of seneka, half a fl. ounce.
 One or two spoonfuls every two hours, in atony
 of the bladder, and mucous discharges.

Radiu.

TINCTURE OF BUCHU.

- R. Buchu leaves, five ounces.
 Proof spirit, two pints.
 Macerate for fourteen days, and filter.

Dub. Ph.

Dose, from one to four fl. drachms.

EXTRACT OF BUCHU.

- R. Buchu, in coarse powder, one pound.
 Ether, four fl. ounces.
 Alcohol, twelve fl. ounces.

Displace without maceration, add diluted alcohol until a pint of ethero-alcoholic tincture is obtained; suffer this to evaporate spontaneously; treat the residue in the displacer, with diluted alcohol, till two pints are obtained; evaporate to a syrup; add the product of the first tincture, and with a gentle heat concentrate to the proper consistence.

W. Procter.

FLUID EXTRACT OF BUCHU.

- R. Buchu leaves, eight ounces.
 Alcohol, sixteen fl. ounces.
 Water, sufficient.

Reduce the leaves to a coarse powder, moisten them in a covered vessel, with twelve fl. ounces of the alcohol, macerate for six hours, and introduce into a displacer. When the clear fluid has ceased to pass, add the remaining alcohol, mixed with four fl. ounces of water, until the displaced liquid amounts to twelve fl. ounces which is to be set aside until reduced to six fl. ounces by spontaneous evaporation. The residue in the displacer is then to be treated with a pint of cold water by maceration for twelve hours, and subjected to pressure till a pint is obtained. Evaporate this to ten fl. ounces, mix with six fl. ounces of the tincture, agitate for several days, filter, or strain. Dose, one to two teaspoonfuls.

W. Procter.

BUXUS.

Box.

The box is a shrub or small tree, native of the south of Europe, but now generally cultivated in this country.

Sex. Syst. Monœc. triand. *Nat. Syst.* Euphorbiacæ.

The parts used are the wood and leaves. The first is sudorific and possesses somewhat the properties of guaiacum; the latter are purgative, but neither are now used. The following preparation, however, has obtained some celebrity in the cure of gonorrhœa, and in epilepsy.

OIL OF BOX.

- R. Rasped box-wood, sufficient.
 Distil, separate the oil, and rectify it.

Dose, four to five drops, three times a-day.

Wirtemberg Ph.

C.

CACAO.

COCOA. CHOCOLATE NUTS.

These are the seeds of *Theobroma Cacao*, a small tree, indigenous to South America. They are also produced from some other sources.

Sex. Syst. Polyadelph. pentand. *Nat. Syst.* Byttneriacæ.

Linn. Sp. Pl. 1100. Lindley, Flor. Med. 138.

The nuts are of an almond shape, and consist of a white, sweetish, somewhat oleaginous substance, covered by a leathery-like shell. They are principally used as an article of food, but also are employed in medicine.

COMPOUND COCOA POWDERS.

PALAMOU.

- R. Roasted cocoa, eight ounces.
 Rice flour,
 Potato starch, each, two pounds
 Red saunders, powdered, one ounce

Mix. A drachm to an ounce, boiled in water as a restorative diet.

Soubéiran.

RACAHOUT DES ARABES.

- R. Roasted cocoa, two ounces.
 Potato starch, five ounces.

Powdered salep, one ounce.
 Sugar, eight ounces.
 Vanilla, sufficient.
 Mix. Two or three spoonfuls boiled with eight fl. ounces of water, form a nutritious diet for the convalescent. *Foy.*

WAKAKA DES INDES.

R. Roasted cocoa, two ounces.
 Sugar, five ounces and a half.
 Powdered cinnamon, two drachms.
 " vanilla, thirty-six grains.
 Ambergris, three grains.
 Musk, one grain and a half.
 Mix. Half a drachm, boiled with milk or water. More stimulating than the last. *Guibourt.*

COMPOUND COCOA PASTE CHOCOLATE.

VANILLA CHOCOLATE.

R. Cocoa paste, six pounds.
 Sugar, ten pounds.
 Vanilla, eleven drachms.
 Triturate thoroughly together, and form cakes. *Cottureau.*

AROMATIC CHOCOLATE.

R. Cocoa,
 Sugar, each, sixteen ounces.
 Powdered cinnamon, half an ounce.
 Cloves, two drachms.
 Cardamom,
 Vanilla, each, one drachm.
 Triturate together, and form cakes. *Weigleb.*

SALEP CHOCOLATE.

R. Cocoa paste,
 Sugar, each, ten pounds.
 Powdered salep, ten ounces.
 Triturate well, and form cakes of half an ounce each. *Cadet.*

WHITE CHOCOLATE.

R. Sugar, six pounds.
 Rice flour, one pound, twelve ounces.
 Potato starch, eight ounces.
 Gum Arabic, four ounces.
 Tincture of vanilla, half fl. ounce.
 Butter of cocoa, eight ounces.
 Boiling water, sufficient.

Triturate well, into a stiff paste. *Cottureau.*
 These afford a good article of diet, for convalescents, and debilitated persons, when boiled with water or milk. The last is the least stimulating

CHOCOLATE CREAM.

R. Boiling milk, sixteen parts.
 Sugar,
 Yolk of egg, each, two parts.
 Ground cocoa, one part.
 Mix, and boil gently. *Beral.*

BUTTER OR OIL OF COCOA.

R. Cocoa, roasted, at will.
 Reduce to a paste in a warm iron mortar, triturate on a stone slab, add a fifth part boiling water, place in a bag, and express between two heated metal plates; melt the product, and, on cooling, separate the water. This article affords a good medium for making suppositories. *Guibourt.*

BUTTER OF COCOA MIXTURE.

R. Butter of cocoa, three ounces.
 Oil of almonds, one ounce and six drachms.
 Syrup of red poppies, one ounce.
 Orange-flower water, four drachms.
 Mix. As a soothing demulcent in catarrh. In spoonful doses. *Cadet.*

BUTTER OF COCOA OINTMENT.

R. Butter of cocoa,
 Oil of almonds, equal parts.
 Melt over a water-bath. *Guibourt.*

CADMIUM.

CADMIUM.

A soft, crystallizable, ductile, volatilizable metal, not used in medicine in its metallic state, but affording the following remedial salt.

In its general effects, cadmium resembles the zinc salts.

CADMII SULPHAS.

SULPHATE OF CADMIUM.

R. Oxide of cadmium, one ounce.
 Sulphuric acid, sufficient.
 Dissolve, evaporate, and crystallize. *Cottureau.*
 It may also be obtained from the carbonate of cadmium by the addition of sulphuric acid.
 Used in solution and ointment, as an application in chronic ophthalmia.

SOLUTION OF SULPHATE OF CADMIUM.

R. Sulphate of cadmium, two grains.
 Laudanum, twenty drops.
 Rose water, four fl. ounces.
 Mix. As a wash in chronic ophthalmia. *Guibourt.*

R. Sulphate of cadmium, eight grains.
Water, one fl. ounce.
Dissolve. Used as a wash in otorrhœa.
Lincke.

OINTMENT OF SULPHATE OF CADMIUM.

R. Sulphate of cadmium,
one or two grains.
Lard, one drachm.
Mix. As an application in spots on the cornea.
Radiis.

CAHINCA.

CAHINCA.

This is the root of *Chiococca Anguifuga*, a shrub growing in Brazil. Its effects are those of a diuretic, and it has chiefly been employed in dropsical diseases. Dose, twenty to forty grains.

EXTRACT OF CAHINCA.

R. Powdered cahinca, one part.
Diluted alcohol, seven parts.
Macerate twelve hours, then introduce into a displacer, exhaust, and evaporate to consistence of an extract.
Dose, ten to twenty grains.

DECOCTION OF CAHINCA.

R. Bruised cahinca, two drachms.
Water, one pint and a half.
Boil to one-half, and strain.
Dose, a tablespoonful. *Von Langsdorff.*

CALAMINA.

CALAMINE.

This is a native impure carbonate of zinc, found in large quantities in England and Germany. The pure carbonate will be described under the head of *Zinc*.

PREPARED CALAMINE.

R. Calamine, at will.
Heat to redness and pulverize; then reduce to a very fine powder, as directed for prepared chalk. *U. S. Ph.*
Used externally, as a mild astringent and absorbent, to excoriations and ulcerations, and also as an ingredient of cerates, &c.

CALAMINE CERATE.

(*Turner's Cerate.*)

R. Prepared calamine,
Yellow wax, each, three ounces.
Lard, one pound.

Melt the wax and lard, and as they thicken, stir in the calamine.

This cerate is much used for excoriations, ulcerations, &c.

CALAMUS.

CALAMUS.

This is the root of *Acorus Calamus*, an indigenous plant found in most parts of the U. S. in wet situations. It also grows in Europe and Asia.

Sex. Syst. Hexand. monog. *Nat. Syst.* Oron-tiaceæ.

Linn. Sp. Pl. 462. Griffith, Med. Bot. 620.

The part employed is the root; this has a fragrant odor, and a warm, bitterish, aromatic taste. It is a stimulant tonic, and aromatic. The dose, in substance, is from a scruple to a drachm.

ELECTUARY OF CALAMUS.

R. Powdered calamus,
two drachms and a half.
" valerian,
one drachm and a half.
Syrup of orange peel, two ounces.
Mix. A teaspoonful every two hours.
Shubarth.

INFUSION OF CALAMUS.

R. Calamus, one ounce.
Boiling water, one pint.
Infuse for a quarter of an hour, and strain.
Dose, a wineglassful or more.

COMPOUND INFUSION OF CALAMUS.

R. Calamus, ten drachms.
Boiling water, one pint.
Infuse and strain, then add
Peppermint water, two fl. ounces.
Muriatic ether,
one drachm and a half.
Syrup, one ounce.
Mix. Dose, a tablespoonful, in disordered diges-tion. *Augustin.*

TINCTURE OF CALAMUS.

R. Calamus, bruised, one part.
Alcohol, (0.847) eight parts.
Macerate for eight days, and filter. *Guibourt.*

COMPOUND TINCTURE OF CALAMUS

R. Contused calamus,
" ginger, } each,
" coriander, } one ounce
Black pepper,
Alcohol, two pints and a half

Macerate for four days, and filter. Dose, forty to fifty drops as a stomachic and carminative.

Niemann.

- R. Contused calamus,
 " gentian, each, three ounces.
 " geum, two ounces and a half.
 " angelica,
 one ounce and a half.
 " ginger, half an ounce.
 " fennel, two ounces.
 Alcohol, twelve pints.

Macerate for six days, and filter. A teaspoonful in wine. *Tromsdorff.*

CALCIUM.

CALCIUM.

CALCII CHLORIDUM.

CHLORIDE OF CALCIUM.

- R. Chalk, five ounces.
 Muriatic acid,
 Water, each, ten fl. ounces.

Mix the acid and water, and gradually add the chalk. When all action ceases, filter, and evaporate to dryness. Fuse, and pour out on a stone slab; when cool, break in fragments, and preserve in well-stopped bottles.

Lond. Ph., 1836.

CATAPLASM OF CHLORIDE OF CALCIUM.

- R. Chloride of calcium,
 " soda, each, half an ounce.
 Water, half a pint.
 Linseed meal, sufficient.

Make a cataplasm. In scrofulous and white swellings. *Swediaur.*

SOLUTION OF CHLORIDE OF CALCIUM.

- R. Chloride of calcium, eight ounces.
 Water (imp. meas.), twelve fl. ounces.

Dissolve, and filter. *Edin. Ph., 1841.*

Dose, thirty drops two or three times a-day, gradually increased; to be given in milk, or some demulcent. Used in scrofula, goitre, tabes mesenterica, &c.

MIXTURE OF CHLORIDE OF CALCIUM.

- R. Chloride of calcium, one drachm.
 Extract of henbane, ten grains.
 Syrup of liquorice, one ounce.
 Water, six ounces.

Mix. A spoonful four times a-day in scrofula. *Phæbus.*

- R. Chloride of calcium, one drachm.
 Almond mixture, seven fl. ounces.
 Syrup of gum Arabic, one fl. ounce.

Mix. A teaspoonful every three hours. *Gräfe.*

PILLS OF CHLORIDE OF CALCIUM.

- R. Chloride of calcium, one drachm.
 Extract of opium, nine grains.
 Mucilage of gum Arabic, sufficient.

Mix, and make fifty-four pills. One every two or three hours, in gonorrhœa, gradually increasing the dose, until eight, ten, or twelve are taken every hour. *Gräfe.*

CALX CHLORINATA.

CHLORINATED LIME.

CHLORIDE OF LIME.

- R. Hydrate of lime, one pound.
 Chlorine gas, sufficient.

Pass the chlorine gas over the lime, spread in a proper vessel, until it is saturated. *Lond. Ph.*

It is desiccant and disinfectant, and is also used to ill-conditioned ulcers, burns, chilblains, to some cutaneous affections, &c.

PRESERVATIVE LIQUID.

- R. Water, sixteen parts.
 Chlorinated lime, four parts.
 Alum, two parts.
 Nitre, one part.

Mix. Said to be very efficient for the preservation of anatomical preparations. *Reboulet.*

LOZENGES OF CHLORINATED LIME.

- R. Chlorinated lime, two drachms.
 Sugar, eight ounces.
 Starch, one ounce.
 Tragacanth, one drachm.
 Cochineal, three grains.

Rub well together, and make lozenges of three grains each. One to be taken three or four times a-day, in cases of bad breath.

Deschamps.

COLLUTORY OF CHLORINATED LIME.

- R. Chlorinated lime, fifteen to thirty grains.

Mucilage of gum Arabic, one fl. ounce.

Syrup of orange peel, half fl. ounce.

Mix. To be applied by means of a piece of sponge or camel's hair-brush, to ulcers in the mouth. *Angelot.*

- R. Chlorinated lime, three drachms.
 Distilled water,
 Alcohol, each, two fl. ounces.
 Oil of roses, four drops.

Dissolve, and filter. A teaspoonful in a glass of water, to correct fetid breath. *Chevallier.*

CHLORINATED LIME DENTIFRICE.

- R. Chloride of lime, four grains.
 Powdered red coral, two drachms.

A toothbrush, slightly wetted, to be dipped in this powder, and rubbed on the teeth.

Magendie.

COMPOUND INJECTION OF CHLORINATED LIME.

R. Chloride of lime, two drachms.
Decoction of rhatany, thirteen fl. ounces.

Dissolve, and filter. As an injection in ozena, three or four times a-day. *Detmold.*

OINTMENT OF CHLORINATED LIME.

R. Chlorinated lime, one drachm.
Lard, one ounce.

Rub together. In scrofulous swellings. *Cima.*

R. Chlorinated lime, half a drachm.
Lard, one ounce.

Rub together. In goitre. *Werneck.*

R. Chlorinated lime,
Borate of soda, each, one drachm.
Lard, one ounce.

Rub together thoroughly. In chilblains. *Trusen.*

R. Chlorinated lime, one drachm.
Powdered digitalis, two drachms.
Vinegar, two scruples.
Lard, one ounce.

Rub together. In indolent glandular tumors. *Phæbus.*

LINIMENT OF CHLORINATED LIME.

R. Chlorinated lime, half a drachm.

Rub in a glass mortar, adding gradually,

Rose water, one fl. ounce.

and when quite clear,

Oil of almonds, one fl. ounce.

As an application in tinea capitis. *Trusen.*

CALCII IODIDUM.

IODIDE OF CALCIUM.

R. Iodide of iron, at will.

Precipitate with an excess of slaked lime, evaporate to dryness, dissolve in distilled water, filter, evaporate, and crystallize. *Magendie.*

PILLS OF IODIDE OF CALCIUM.

R. Iodide of calcium, ten grains.
Extract of savine, twelve grains.

Mix, and divide into four pills. One every four hours. In amenorrhœa, with scrofula. *Brera.*

R. Iodide of calcium, ten grains.
Extract of aconite, six grains.

Mix, and divide into six pills. One every four hours. In chronic bronchitis and tubercular phthisis. *Brera.*

CALCII SULPHURETUM.

SULPHURET OF CALCIUM.

R. Quicklime, seven parts.
Sulphur, four parts.

Pulverize, and beat for two hours in a covered crucible. *Guibourt.*

R. Quicklime, three parts.
Sulphur, one part.
Water, five parts.

Mix, and boil gently; evaporate to dryness, stirring. Keep in tight bottles. *Codex.*

Stimulant and diaphoretic. Dose, five to ten grains.

COMPOUND PILLS OF SULPHURET OF CALCIUM.

R. Sulphuret of calcium, one drachm.
Powdered cinnamon,
Extract of aconite, each, fifteen grains.

“ mallows, sufficient.

Mix, and make sixty pills. Dose, four, three or four times a-day, in gout and chronic rheumatism. *Phæbus.*

LINIMENT OF SULPHURET OF CALCIUM.

R. Sulphuret of calcium, one ounce.
Oil of juniper, two drachms.
Dippel's animal oil, ten drops.

Mix well. As an embrocation in gout. *Augustin.*

CALCII OXIDUM.

CALX. LIME.

R. Chalk, one pound.

Break into small pieces, and calcine in a strong fire for an hour. *Lond. Ph. 1836.*

LIME WATER.

R. Lime, four ounces.
Distilled water, one gallon.

Slake the lime with a little of the water, add the remainder, stir well, and let stand for three hours. Keep in well-stopped bottles. *U. S. Ph.*

COMPOUND LIME WATER.

R. Rasped guaiacum wood, half a pound.

Contused liquorice root, one ounce

“ sassafras bark, half an ounce.

“ coriander seeds, three drachms.

Lime water, (wine meas.) six pints

Macerate for two days in a closed vessel, and filter. *Dub. Ph. 1826.*

LINIMENT OF LIME WATER AND OIL.

R. Linseed oil, three fl. ounces.
Lime water, six fl. ounces.

Make a liniment. As an application to burns and scalds. *Ellis.*

R. Lime water,
Linseed oil, each, two fl. ounces.

Mix. Used like the preceding, under the name of *Carron oil*. *U. S. Ph.*

LINIMENT OF LIME WATER AND ALCOHOL.

R. Alcohol, two fl. ounces.
Lime water, half a pint.

Mix. In same cases as the last. *Ellis.*

LINIMENT OF LIME WATER AND OPIUM.

R. Lime water,
Linseed oil,
Laudanum, equal parts.

Mix. In the same cases, and as an embrocation to allay pain. *Augustin.*

LINIMENT FOR SORE BREASTS.

R. Lime water,
Almond oil, each, three drachms.
Extract of opium, one grain.

Mix. The breasts are to be covered with lint dipped in this mixture. *Niemann.*

LINIMENT OF LIME AND SULPHUR.

R. Lime,
Sulphur, each, two ounces.
Olive oil, sufficient.

Mix. As a friction in scabies. *Giannini.*

LIME WATER AND CARBONATE OF POTASSA.

R. Carbonate of potassa, two drachms.
Lime water, two pints.

Dissolve. As an antilithic; a wineglassful every two hours, in same quantity of new milk. *Ellis.*

LIME WATER AND MILK.

R. Lime water,
New milk, equal parts.

Mix. As an antacid, and to remove sickness of the stomach. Dose, one to two table-spoonfuls. *Ellis.*

ANTACID MIXTURE.

R. Lime water, four fl. ounces.
Solution of potassa,
Syrup of orange peel, each,
one fl. ounce.

Mix. Dose, a spoonful in a cup of water. Found useful in dyspepsia and heartburn. *Cooley.*

LIME OINTMENT.

R. Lime, one ounce.
Linseed oil, three ounces.
Mucilage of quince seeds, two ounces.

Mix well. In burns and scalds. *Radius.*

R. Slaked lime, one drachm.
Carbonate of soda, two drachms.
Extract of opium, ten grains.
Lard, two ounces.

Rub well together. In obstinate cutaneous affections. *Bielt.*

R. Slaked lime,
Laudanum, each, half a drachm.
Cucumber ointment, four drachms.

Rub well together. As an application to hemorrhoidal tumors. *Guibourt.*

R. Lime,
Sulphur, each, two ounces.
Muriate of soda, one ounce.
Lard, one pound.
Olive oil, two pounds.

Mix well. As a friction in itch. *Ferrara. Ph.*

CALCIS CARBONAS.

CARBONATE OF LIME.

PRECIPITATED CARBONATE OF LIME.

R. Water of chloride of calcium, five parts.
Carbonate of soda, three parts,
dissolved in four times its weight of distilled water. Mix. Wash the precipitate three times, and dry. *Dub. Ph.*

PREPARED CHALK.

R. Chalk, at will.
Add a little water, and rub to fine powder. Throw into a vessel of water, and stir; pour off liquid while yet turbid, into another vessel, and permit chalk to subside; pour off water, and dry the precipitate. *U. S. Ph.*

PREPARED OYSTER SHELL.

R. Oyster shell, at will.
Free it from extraneous matter, wash with boiling water, reduce to powder, and proceed as with chalk. *U. S. Ph.*

As an antacid in bowel affections. In doses of from ten to forty, or more grains, often repeated.

AROMATIC POWDER OF CHALK.

- R. Prepared oyster shell, two drachms.
 Magnesia, one drachm.
 Powdered cinnamon, one scruple.
 Mix.

COMPOUND POWDER OF CHALK.

- R. Prepared chalk, half a pound.
 Cinnamon, four ounces.
 Tormentil,
 Gum Arabic, each, three ounces.
 Long pepper, half an ounce.

Beat separately into fine powder, and mix.

Lond. Ph.

Stimulant, astringent, and antacid; useful in diarrhoea, with acidity, but without inflammation. Dose, ten to twenty grains.

COMPOUND POWDER OF CHALK WITH OPIUM.

- R. Compound powder of chalk,
 six ounces and a half.
 Powdered opium, four scruples.

Mix well.

Lond. Ph.

In diarrhoea of adults, in doses of ten or twenty grains, after each evacuation.

DENTIFRICE POWDER WITH CARBONATE OF LIME.

- R. Powdered cuttlefish bone,
 " Peruvian bark,
 " myrrh, equal parts.

Mix.

Pierquin.

- R. Powdered cuttlefish bone, six ounces.
 " red coral, two ounces.
 " dragon's blood,
 " red saunders,
 " orris root, each, one ounce
 and a half.
 " cinnamon,
 " alum, each, half an ounce.

Mix.

Par. Cod.

LOZENGES OF CHALK.

- R. Prepared chalk, four ounces.
 Gum Arabic powdered, one ounce.
 Nutmeg, " one drachm.
 Sugar, " six ounces.

Rub together till mixed; mix with sufficient water to form mass; divide into lozenges of ten grains each.

U. S. Ph.

As a mild antacid astringent, in diarrhoea.

DENTIFRICE ELECTUARY.

- R. Powdered red coral, four ounces.
 " cuttlefish bone, one ounce.

- Powdered cinnamon, one ounce.
 " cochineal, half an ounce.
 " alum, half a drachm.
 Oil of cloves, six drops.
 Honey, ten ounces.

Mix.

Taddei.

CHALK MIXTURE.

- R. Prepared chalk, half an ounce.
 Sugar,
 Powdered gum Arabic, each, two
 drachms.

Cinnamon water,

Water, each, four fl. ounces.

Rub them together until thoroughly mixed.

U. S. Ph.

A tablespoonful, as may be required.

- R. Prepared chalk, one drachm and
 a half.

White sugar,

Gum Arabic, each, one drachm.

Oil of cinnamon, two drops.

Tincture of opium, forty to fifty drops.

Distilled water, four fl. ounces.

Mix. A tablespoonful, every two hours, in diarrhoea and dysentery.

Ellis

COMPOUND CHALK MIXTURE.

- R. Chalk mixture, five fl. ounces.
 Aromatic confection, one drachm.
 Solution carbonate ammonia,

one fl. drachm.

Laudanum, twenty minims.

Mix. A tablespoonful, occasionally, in diarrhoea.

Ainslie

CALCIS PHOSPHAS.

PHOSPHATE OF LIME.

- R. Burnt bones, in powder, one part.
 Diluted muriatic acid,
 Water, each, two parts.

Digest together for twelve hours, and filter. Add water of caustic ammonia, sufficient to precipitate phosphate of lime. Wash this, and dry.

Dub. Ph. 1826.

BURNT HARTSHORN.

R. Burn pieces of hartshorn in an open vessel till quite white, and prepare them as directed for chalk. *Lond. Ph. 1836.*

Has been thought efficacious in mollities osium and rachitis, but probably inert. Dose, twenty grains, or more.

SYRUP OF PHOSPHATE OF LIME.

- R. Phosphate of lime, one hundred and twenty-eight grains.
 Glacial phosphoric acid, half an ounce.
 Sugar, seven and a half ounces.
 Water, four fl. ounces.
 Essence of lemon, twelve drops.

Mix the phosphate with the water, and heat in a sand bath, gradually adding the phosphoric acid. Add water enough to compensate for the evaporation, then dissolve the sugar, and when cold add the essence of lemon. Each teaspoonful contains two grains of phosphate and four grains of phosphoric acid. *A. B. Durand.*

DENTIFRICE POWDER OF PHOSPHATE OF LIME.

- R. Phosphate of lime, eighteen parts.
 White bole,
 Gum lac, each, twelve parts.
 Cinnamon, one part.
 Myrrh, two parts.

Powder well, and mix. *Van Mons.*

DENTIFRICE ELECTUARY OF PHOSPHATE OF LIME.

- R. Powdered phosphate of lime, two ounces and a half.
 " gum lac, one ounce.
 " cinnamon, one drachm and a half.
 " cloves, one scruple.
 Syrup of lemon, sufficient.

Make a soft electuary. *Van Mons.*

CALENDULA.

MARIGOLD.

This is the officinal name of *Calendula officinalis*, a plant indigenous to Europe, and often cultivated in gardens in this country as a pot herb.

Sex. Syst. Syngen. necess. *Nat. Syst.* Asterales.

Linn. Sp. Pl. 1304. *Lindley, Flor. Med.* 466.

The parts used are the herb, flowers, and seed. Their smell is peculiar, and somewhat aromatic; the taste is feebly bitter, and a little acrid. It was formerly much used, but had fallen into disuse, when, of late years, it has again been brought forward as a remedy in cancerous affections.

EXTRACT OF MARIGOLD.

- R. Marigold, one part.
 Tepid water, four parts.

Macerate for twenty-four hours, boil for a quarter of an hour, express, boil residue with four

parts of water, express, mix two decoctions, and evaporate to proper consistence. *Guibourt.*

PILLS OF THE EXTRACT OF MARIGOLD.

- R. Extract of marigold,
 " cicuta,
 Subcarbonate of iron, each, two drachms.
 Powdered marigold, one drachm and a half.

Mix well, and divide into pills of two grains each. Five or six are to be taken three times a-day in schirrhous and cancerous affections. *Rust.*

LOTION OF EXTRACT OF MARIGOLD.

- R. Extract of marigold,
 " cicuta, each, three drachms.
 Cherry-laurel water, two ounces.
 Tincture of opium, half an ounce.

Mix. As a lotion to cancerous ulcerations. *Rust.*

INFUSION OF MARIGOLD.

- R. Marigold flowers, three drachms.
 Boiling water, sufficient
 to obtain five ounces; infuse, and add to the strained liquid
 Syrup of orange peel, half an ounce.

Dose, two spoonfuls, three times a-day. *Radius.*

LINIMENT OF MARIGOLD.

- R. Marigold flowers, three drachms.
 Water, five fl. ounces.
 Boil to three ounces, strain, and add
 Extract of marigold, one drachm.
 Tar, two ounces.
 Pyroligneous acid, half an ounce.
 Gum Arabic, sufficient.

Mix. As an application to cancerous ulcers. *Radius.*

CALOTROPIS.

MUDAR.

The *Calotropis gigantea* is a native of the East Indies, and is now naturalized in several of the West India Islands.

Sex. Syst. Pentand. monog. *Nat. Syst.* Asclepiadaceae.

Brown, Tr. Wern. Soc. 1. 29. *Griffith, Med. Bot.* 456.

The part used is the bark of the root; this is of a whitish color, inodorous, but with a bitter, nauseous taste. It is purgative, alterative, and diaphoretic, and has been highly spoken of in the treatment of elephantiasis and lepra. Dose, three grains to thirty; in the latter dose it is emetic and purgative.

INFUSION OF MUDAR.

R. Mudar, three drachms.
Boiling water, eight ounces.
Infuse, and strain. As an alterative, one to two fl. drachms; as an emetic, two fl. ounces.
Cassanova.

PILLS OF MUDAR.

R. Powdered mudar, one drachm.
Honey, sufficient.
Beat into mass, and divide into pills of four grains. Two or three to be taken daily as an alterative in lepra, &c.
Cassanova.

OIL OF MUDAR.

R. Powdered mudar, one drachm.
Olive oil, seven ounces.
Infuse for half an hour over a water-bath, and strain. As an application to leprosy sores.
Cassanova.

CAMPHORA.

CAMPHOR.

A peculiar white, crystallizable, semi-transparent, volatile product of several plants; but that used in medicine is obtained from the *Camphora officinarum*, an evergreen tree, of considerable size, indigenous to China, and other Asiatic countries.

Ser. Syst. Eneand. monog. *Nat. Syst.* Lauraceæ.

Nees. Laurin. 88. Griffith, Med. Bot. 553.

It has narcotic, diaphoretic, sedative properties, and is employed in a variety of diseases, and used externally, as an anodyne. The dose is from three to ten grains. Camphor can be powdered by trituration, with a few drops of alcohol.

CAMPHOR POWDER.

R. Camphor, nine grains.
Sugar, one drachm.
Mix, and divide into nine powders. *Augustin.*

COMPOUND CAMPHOR POWDER.

R. Powdered camphor,
Myrrh, each, two drachms.
Peruvian bark,
Chamomile, each, half an ounce.
Charcoal, one ounce.
Mix. Dose, two scruples, or more, in gangrene.
Rust.

R. Powdered camphor,
Benzoic acid, each, six grains.
Sugar, one drachm.
Mix, and divide into four powders. As an antispasmodic and sedative.
Saunders.

R. Powdered gum Arabic, one ounce.
" orange peel,
" sugar, each, two drachms.
" camphor, ten grains.
" opium, five grains.
Mix. A teaspoonful every hour. *Ammon.*
Said to have been useful in cholera.

CAMPHOR PILLS.

R. Powdered camphor, }
Sugar, } each,
Starch, } one scruple.
Crumb of bread, }
Mix, and make twenty pills. *Augustin.*

COMPOUND CAMPHOR PILLS.

R. Powdered camphor, twenty-four grains.
" musk, eight grains.
" opium, two grains.
Syrup, sufficient.
Beat into a mass, and divide into twelve pills. In hospital gangrene. *Dupuytren.*

PILLS OF CAMPHOR AND LACTUCARIUM.

R. Camphor,
Lactucarium, each, fifty grains.
Mix, and make twenty pills. Four to six daily, as an aphrodisiac. *Ricord.*

PILLS OF CAMPHOR AND MUSK.

R. Camphor, one scruple.
Musk, ten grains.
Ammoniac, two scruples.
Opium, four grains.
Mix, and divide into four-grain pills. Dose, four or five in a day, in nervous disorders. *Richard.*

CAMPHOR WATER.

R. Camphor, two drachms.
Alcohol, forty minims.
Carbonate of magnesia, four drachms.
Distilled water, two pints.
Rub the camphor with the alcohol, then with the magnesia, and afterwards, with the water gradually added, and filter. *U. S. Ph.*
Dose, one or two tablespoonfuls, to be repeated as occasion may require.

CAMPHOR WATER AND LAUDANUM.

R. Camphor water, four fl. ounces.
Compound spirit of lavender, two fl. drachms.
Tincture of opium, forty to fifty drops
Mix. A tablespoonful, every two hours, in diarrhoea and dysentery. *Ellis.*

CAMPHOR WATER AND NITRIC ACID.

(Hope's Mixture.)

R. Camphor water, four fl. ounces.
 Nitric acid, four drops.
 Tincture of opium, forty to fifty drops.

Mix. In the same dose, and in the same complaints, as the last. *Ellis.*

R. Nitrous acid, one fl. drachm.
 Laudanum, forty drops.
 Camphor water, eight fl. ounces.

Mix. Dose, one or two fl. ounces every three hours, in dysentery. *Hope.*

R. Camphor water, ten fl. drachms.
 Laudanum, forty drops.
 Spirit of sulphuric ether,

one fl. drachm.

Syrup of poppies, one fl. drachm.

Mix. To prevent a paroxysm of intermittent fever. To be given just before its accession.

Gregory.

CAMPHOR WATER AND HOFFMANN'S ANODYNE.

R. Camphor water, four fl. ounces.
 Hoffmann's anodyne, two fl. drachms.

Mix. A dessertspoonful, every hour or two, in some demulcent drink. In nervous affections and sleeplessness in fevers. *Ellis.*

INFUSION OF CAMPHOR.

R. Camphor, one ounce.
 Boiling water, eight fl. ounces.

Pour the water on the camphor, let stand till cold, and keep in well-stopped bottle. Dose, a tablespoonful, every hour or two, in nervous pervigilium and nervous irritability. *Ellis.*

CAMPHOR MIXTURE.

R. Powdered camphor, one drachm.
 " gum Arabic,
 " white sugar, each,

one drachm and a half.

Tincture of opium, forty drops.

Mint water, four fl. ounces.

Mix. A tablespoonful every two hours, in low conditions of the system. *Ellis.*

R. Camphor water, three fl. ounces.
 Compound spirit of lavender, one fl. ounce.
 Sugar, one drachm.

Mix. Give a tablespoonful every two hours, in diarrhoea and cholera-morbus, adding ten drops of laudanum, when there is much pain. This is the celebrated mixture of *Dr. Jos. Parrish.*

R. Powdered camphor, one scruple.
 " gum Arabic, one drachm.
 " sugar, sufficient.
 Cinnamon water, one fl. ounce.

Mix. To relieve the pain in dysmenorrhœa. One-half to be given as soon as pain is felt; if not relieved in an hour or two, the remainder to be taken. *Devees.*

R. Powdered camphor, half a drachm.
 " gum Arabic, two drachms.
 " sugar, three drachms.

Vinegar, half an ounce.

Water, six ounces.

Mix. As a stimulant. *Phabus.*

EMULSION OF CAMPHOR.

R. Camphor,
 Olive oil, each, one drachm.
 Powdered gum Arabic, half an ounce.
 Water, four fl. ounces.

Triturate the camphor and oil together, until uniformly mixed; add the gum, and then the water gradually, triturating till the emulsion is formed. *W. Procter.*

CAMPHOR AND MILK.

R. Camphor, one drachm.
 Boiling milk, four fl. ounces.

Dissolve. To be used as a stimulant. *Ellis.*

CAMPHOR WITH MYRRH.

R. Powdered camphor, one drachm.
 " myrrh, half a drachm.
 " sugar, two drachms.
 Water, six fl. ounces.

Mix. A tablespoonful every two hours. *Ellis.*

TINCTURE OF CAMPHOR.

R. Camphor, four ounces.
 Alcohol, two pints.

Dissolve. *U. S. Ph.*

Chiefly used as an anodyne embrocation. Dose, ten to twenty drops.

TINCTURE OF CAMPHOR AND SAFFRON.

R. Camphor, one ounce.
 Saffron, one scruple.
 Alcohol, one pint.

Macerate for a day, and filter. *Guibourt.*

As a lotion, or fomentation.

ETHEREAL TINCTURE OF CAMPHOR.

R. Camphor, one ounce.
 Sulphuric ether, eight fl. ounces.

Dissolve. *Swediaur.*

Dose, twenty to thirty drops, in wine, as a stimulant and antispasmodic.

MIXTURE OF TINCTURE OF CAMPHOR.

R. Tincture of camphor,
Goulard's extract, equal parts.

Mix. As a lotion, several times a-day, to parts disposed to ulcerate by constant decubitus.

Tott.

MIXTURE OF CAMPHOR AND CHLOROFORM.

R. Camphor, one drachm.
Chloroform, half a fl. drachm.

Mix, and add
Yolk of egg, one.

Triturate, and add gradually
Water, six fl. ounces.

As a stimulant. Dose, two fl. drachms to fl. half ounce.

Keating.

MIXTURE OF CAMPHOR AND ETHER.

R. Camphor, one scruple.
Sulphuric ether, two drachms.

Dissolve, and add
Laudanum, twenty drops.
Cinnamon water, six ounces.

Mix. A spoonful every three or four hours as a stimulant.

Augustin.

CAMPHOR LINIMENT.

R. Camphor, half an ounce.
Olive oil, two fl. ounces.

Dissolve the camphor in the oil. *U. S. Ph.*
As an anodyne embrocation.

COMPOUND CAMPHOR LINIMENT.

R. Camphor, two ounces and a half.
Solution of ammonia, seven fl. ounces and a half.
Spirit of lavender, one pint.

Mix solution of ammonia with the spirit; then, from a glass retort, by a slow fire, distil a pint; dissolve camphor in the distilled fluid.

Lond. Ph., 1836.

A rubefacient and anodyne embrocation.

R. Camphor, one drachm.
Oil of chamomile,
Wine of opium, each, two drachms.
Oil of hyoseyamus, one ounce.

Mix. *Augustin.*

LINIMENT OF CAMPHOR AND VINEGAR.

R. Tincture of camphor, three fl. ounces.
Acetic acid, one fl. drachm.

Mix. As a lotion or embrocation. *Ellis.*

ETHEREAL CAMPHOR LINIMENT.

R. Soft soap,
Alcohol, each, one ounce.

11

Dissolve and add

Camphor, half an ounce,
Dissolved in
Sulphuric ether, one fl. ounce.
As an embrocation. *Saunders.*

WARD'S ESSENCE FOR HEADACHE.

R. Camphor, two ounces.
Rectified spirit, fourteen ounces.
Water of ammonia, two ounces.
Oil of lavender, half an ounce.

Mix. As an application to the forehead, &c., in headache. *Redwood.*

CAMPHOR WATER MIXTURE.

R. Camphor water,
Solution of acetate of ammonia, each, half a fl. ounce.
Antimonial wine,
Tincture of opium, each, twenty drops.

Mix. To be taken at bedtime, as a stimulating diaphoretic. *Ellis.*

CAMPHOR CLYSTER.

R. Camphor, one drachm.
Olive oil, two ounces.

Mix. An excellent enema in cases of ascari-
rides, to be used for three or four successive
nights. *Ellis.*

R. Common clyster, two pints.
Camphor, two drachms.
Yolk of egg, one.

Mix. A fourth part to be injected at a time to be employed as occasion may require in low fevers. *Chaussier.*

CAMPHOR OINTMENT

R. Lard,
Suet,
Oil of bayberries, } each,
Wax, } half an ounce.

Melt together, and add

Camphor, one drachm.
Said to be efficacious as an application to chil-
blains. *Radius.*

R. Fresh butter, three drachms.
Spermaceti, two drachms.
White wax, three ounces.

Melt together, and add

Powdered camphor,
Oxide of zinc, each, three grains.

To be applied in chronic inflammation of the eyelids, to the affected parts, at night. *Maternite*

R. Camphor, one drachm.
 Basilicon ointment, one ounce.
 Soap, half an ounce.
 Powdered mustard, one scruple.
 Mix. As an external application in lumbago.
Ferriar.

CAMPHOR COLLYRIUM.

R. Camphor water, six fl. ounces.
 Solution acetate of ammonia, two fl. ounces.
 Mix. A mild astringent and stimulant in inflammation of the eyes.
Ellis.

LINIMENT OF CAMPHOR AND TURPENTINE.

R. Oil of turpentine, fifteen parts.
 Camphor, one part.
 Mix. As a stimulating embrocation. *Beral.*

NITRATE OF CAMPHOR.

R. Nitric acid, at will.
 Camphor, sufficient
 to saturate. Decant. Ten drops every two hours on sugar, or in syrup, in chronic pleurisy. Also used as a friction in rheumatism and palsy.
Van Mons.

CAMPHORATED COUGH MIXTURE.

R. Camphor, half a scruple.
 Mucilage of gum Arabic, three ounces.
 Syrup of mallows, one ounce and a half.
 Mix. A tablespoonful every two hours.
Saunders.

CANELLA.

CANELLA.

This is the bark of the *Canella alba*, a large tree, native of Jamaica and other West India islands.

Sex. Syst. Dodecand. monog. *Nat. Syst.* Meliaceæ.

Swartz. Trans. Linn. Soc. 1. 96. Griffith, Med. Bot. 181

The part used is the bark, which is of a pale orange-yellow color, with an aromatic odor, and a warm, bitterish, pungent taste. It is principally used as an adjuvant to other remedies. The dose, as a stimulant, is from ten grains to a scruple.

POWDER OF CANELLA AND ALOES.

R. Aloes, one pound.
 Canella, three ounces.
 Rub into a fine powder, and mix. This is much used in amenorrhœa. Dose, ten to twenty grains.
U. S. Ph.

R. Powdered aloes, one drachm and a half.
 " canella, eighteen grains.
 " serpentaria, twelve grains.
 Mix, and divide into six powders. Give one powder every three hours, in syrup. This and the preceding are generally termed *hiera picra*.
Ellis.

TINCTURE OF HIERA PICRA.

R. Powder of aloes and canella, one ounce and a half.
 Brandy, one pint.
 Macerate ten days.
 A popular and most efficient remedy in amenorrhœa. Dose, a teaspoonful.
Ellis.

CANNABIS.

HEMP.

The *Cannabis sativa* is an annual plant, a native of Asia, but generally cultivated in Europe and the United States. The East Indian plant is more powerful in its action on the system than that growing in more temperate climates. This alone is used in medicine; and it is designated in the U. S. Pharm. as the *Cannabis sativa—variety Indica*.

Sex. Syst. Diœc. pentand. *Nat. Syst.* Cannabinaceæ.

Linn. Sp. Pl. 1457. Griffith, Med. Bot. 572.
 The parts used are the seeds, the leaves, and the stems; the two latter are covered with a resinous exudation, which is narcotic and intoxicating. They have been recommended in a variety of diseases of a formidable character.

EXTRACT OF HEMP.

R. Dried hemp tops, at will.
 Boil in alcohol until all the resinous matter is dissolved; strain, and evaporate to proper consistence.
O'Shaughnessy.
 Has been given with success in tetanus, hydrophobia, cholera, &c. Dose, one to ten grains. To be increased as occasion may require.

R. Dried hemp tops, one pound.
 Rectified spirit, one gallon.
 Boil, distil off the spirit, and evaporate the extract by a gentle heat. *Bengal Dispensatory.*
 Dose, half a grain to a grain, in painful and spasmodic affections.

PURIFIED EXTRACT OF HEMP.

- R. Commercial extract of
hemp, one ounce.
Rectified spirit, four fl. ounces.

Dissolve, let the dregs subside, decant and evaporate the clear solution on a water-bath, to the consistence of an extract. Dose, half a grain.
Dub. Phar.

TINCTURE OF HEMP.

- R. Purified extract of
hemp, half an ounce.
Rectified spirit, (Imp.) half a pint.
Dose, five to twenty drops, in neuralgia, &c.
Dub. Ph.

- R. Extract of hemp, twenty-four grains.
Diluted alcohol, one fl. ounce.
Dissolve. In tetanus, one drachm every half hour, till some effect is produced. In cholera, ten drops every half hour; in other diseases, every two hours.
O'Shaughnessy.

EMULSION OF HEMP-SEED.

- R. Hemp-seed, bruised, six drachms.
Gum Arabic, one drachm.
Parsley water, five fl. ounces.
Rub together so as to form emulsion, strain, and add

Syrup of balsam of Peru, one ounce.
A spoonful every hour in gonorrhœa. *Radius.*

RESIN OF HEMP.

- R. Dried hemp-tops, bruised, at will.
Macerate several times in warm water, and then in a solution of carbonate of soda; afterwards, wash well with water, pressing after each operation. Dry, and digest in rectified spirit, to which milk of lime, containing an ounce of lime to each pound of hemp-tops, has been added. Filter, add a slight excess of sulphuric acid; again filter, distil off most of the spirit, add to the residue three or four times its bulk of water, evaporate the rest of the spirit, decant the water, wash and dry. Dose, two-thirds of a grain.
Smith.

CANTHARIS.

SPANISH FLY.

The *Cantharis vesicatoria* is an insect inhabiting those parts of Europe in which the vine grows naturally. It has a fetid odor and an acrid taste. The body is oblong, of a golden-green color. Spanish flies are stimulant, diuretic, rubefacient, and vesicant. They are used internally in dropsies, and in various disorders of the urinary organs; externally, to irritate or vesicate. Several species are natives of the United States, which are not inferior in their

powers to the foreign insect. Cantharidin, their active principle, is soluble in ether, chloroform, the oils, and boiling alcohol. Dose of the flies, one to two grains.

POWDER OF CANTHARIDES AND SAVINE.

- R. Powdered cantharides, two grains.
" savine, one drachm.

Mix, and divide into four powders. One to be taken at night, as an emmenagogue, watching its effects closely. *Ellis.*

POWDER OF CANTHARIDES AND CAMPHOR.

- R. Powdered cantharides, four grains.
" camphor, eight grains.
Sugar of milk, three drachms.

Mix, and divide into eight powders, one to be given twice a day as a diuretic. *Augustin.*

- R. Powdered cantharides, eighteen grains.
" opium,
" camphor, each,
thirty-six grains.

Mix, and form into thirty-six pills. One to be taken at bedtime, in cases of impotency from general debility. *Ellis.*

EXTRACT OF CANTHARIDES.

- R. Cantharides, at will.
Digest, in four or five times their weight of alcohol; filter, repeat operation, unite the two tinctures, and evaporate. *Soubiran.*

PILLS OF CANTHARIDES AND IRON.

- R. Powdered cantharides, one scruple.
Subcarbonate of iron, two scruples.
Extract of liquorice, one drachm.
Form a mass, and divide into pills of one grain each. Dose, three or four, three times a-day, gradually augmenting the dose. As a diuretic. *Meissner.*

PILLS OF CANTHARIDES AND CAPSICUM.

- R. Powdered cantharides, five grains.
" capsicum, eight grains.
" camphor, one scruple.
" guaiacum, one drachm.
Tincture of colocynth, sufficient.

Form mass, and divide into eighty pills. Dose, one, gradually increasing to four or five, twice a-day, in paraplegia. *Sundelin.*

INFUSION OF CANTHARIDES.

- R. Cantharides, one scruple.
Boiling water, sufficient
to obtain three ounces of infusion, after digesting for half an hour, and straining. *Soubiran.*

ACETIC INFUSION OF CANTHARIDES.

R. Powdered cantharides, two ounces and a half.

Acetic acid, two pints.

Digest for fourteen days and filter. Recommended for blistering the scalp without removing the hair. *Mettauer.*

EMULSION OF CANTHARIDES.

R. Cantharides, one scruple.
Sweet almonds, six drachms.
Sugar, one ounce.

Beat together, and gradually rub up with

Lime water, ten ounces.

Strain. A spoonful every two or three hours in neuralgia, and in catarrh of the bladder. *Berends.*

R. Cantharides, half a drachm.
Sweet almonds, one ounce.
Sugar, half an ounce.

Beat together, and make an emulsion with

Hot water, ten fl. ounces.

Strain. A spoonful, every two or three hours, in whooping-cough. *Hufeland.*

VINEGAR OF CANTHARIDES.

R. Powdered cantharides, two ounces.
Acetic acid, twenty fl. ounces.

Macerate for eight days, express, and filter. *Lond. Ph., 1836.*

As an epispastic, or as a rubefacient lotion.

TINCTURE OF CANTHARIDES.

R. Cantharides, bruised, one ounce.
Diluted alcohol, two pints.

Macerate for fourteen days, express, and filter. *U. S. Ph.*

A very convenient mode of administering cantharides. Dose, from ten to twenty drops. Also useful as a rubefacient.

CAMPHORATED TINCTURE OF CANTHARIDES.

R. Tincture of cantharides, half an ounce.
Camphor, three drachms.
Alcohol, two ounces.

Mix. As a rubefacient. *Pierquin.*

COMPOUND TINCTURE OF CANTHARIDES.

R. Cantharides, one drachm.
Mustard, }
Black pepper, } each, half an ounce.
Camphor, }
Garlic, one bulb.
Vinegar, six ounces.
Alcohol, twelve ounces.

Macerate for some days, express, and filter. Said to be very effectual as a rubefacient lotion in rheumatism, &c. *Foy.*

R. Tincture of cantharides,
Extract of rosemary, each,
one drachm.

Solution of carbonate of
potassa, one drachm and a half.

Water, four fl. ounces.

Mix. As a lotion to promote the growth of hair, where it has been lost from an eruption, or from fever. *Sachse.*

TINCTURE OF CANTHARIDES AND GUA-
IACUM.

R. Powdered cantharides, one drachm.

" guaiacum, forty-five grains.

Contused rhubarb, one drachm and a half.

" shell-lac, half a drachm.

Alcohol, twelve fl. ounces.

Digest, and strain. Dose, from thirty to fifty drops, night and morning, in some demulcent. In gleet. *Ellis.*

ETHEREAL TINCTURE OF CANTHARIDES.

R. Powdered cantharides, one part.
Acetic ether, sixteen parts.

Digest for eight days, express, and filter. *Foy.*
As a rubefacient and vesicant lotion; it is very energetic.

R. Powdered cantharides, three ounces.
Spirit of nitric ether, two pints and a half.

Digest for eight days, and filter.

Used for the same purposes as the last.

Mettauer.

ETHEREAL EXTRACT OF CANTHARIDES.

R. Powdered cantharides, at will.
Sulphuric ether, sufficient.

Treat the powder with the ether in a displacement funnel, properly arranged, until exhausted, and distil off the ether. Flies yield 8 per cent. of this extract. It is employed for increasing the activity of blistering plasters, and for admixture with wax to make blistering tissue-paper. *Soubéiran*

CAMPHORATED ETHEREAL TINCTURE
OF CANTHARIDES.

R. Powdered cantharides, one part.
Nitrate of camphor, three parts.
Alcohol, fifty-six parts.

Digest and express, and filter at the end of a few days. Advised in dropsy, in the dose of ten to twenty drops, in white wine. *Van Mons.*

OIL OF CANTHARIDES.

R. Cantharides, one part.
Olive oil, eight parts.
Digest for six hours on a water-bath, express,
and filter. *Guibourt.*

An active vesicant.

R. Cantharides, one part.
Oil of rosemary, eight parts.
Digest for fifteen days, express, and filter.
More active than the preceding. *Beral.*

R. Powdered cantharides, one part.
Chloroform,
Castor oil, each, one and a half parts.

Mix, let stand in a close vessel for some hours,
and then pass through a displacement ap-
paratus.

This oil vesicates promptly, when applied to
a part, and covered with oiled silk. *E. Dupuy.*

LINIMENT OF CANTHARIDES.

R. Powdered cantharides, one ounce.
Oil of turpentine, half a pint.
Digest for three hours over a water-bath, and
strain. *U. S. Ph.*

An active stimulant when applied to the
skin; used with success in the low stages of
typhus fevers. Requires caution in its use.

LINIMENT OF CANTHARIDES AND CAM-
PHOR.

R. Camphor, three drachms.
Dissolve in
Liniment of cantharides, half fl.
ounce.

Add
Strong mercurial ointment,
Simple ointment, each, one ounce.
Mix, and form liniment. In low stages of
fever, where the object is to arouse the system,
and to affect it with mercury also as an ap-
plication to tumors *Ellis.*

DECOCTION OF CANTHARIDES IN TUR-
PENTINE.

R. Cantharides in powder, four ounces.
Oil of turpentine, twelve ounces.
Mix in a flask, place this in a salt-bath, and
heat for four hours; then dissolve with oil of
turpentine until twelve fl. ounces are obtained.
Dr. Hartshorne.

LINIMENT OF CANTHARIDES AND SOAP.

R. Tincture of cantharides, half an
ounce.
Liquid ammonia, an ounce and a half.
Turpentine soap, two ounces.
Spirit of turpentine, one ounce.
" of camphor, eight ounces.
Mix. A powerful rubefacient. *Niemann.*

CERATE OF CANTHARIDES.

R. Powdered cantharides, one pound.
Yellow wax,
Resin, each, seven ounces.
Lard, ten ounces.

Melt the wax, resin and lard, add the flies, and
keep the mixture fluid for half an hour over a
water-bath; remove from the fire and stir till
cold. *U. S. Ph.*

This is the common blistering plaster.

CANTHARIDES OINTMENT.

R. Cantharides, in powder, two ounces.
Distilled water, half a pint.
Resin cerate, eight ounces.

Boil down the cantharides in the water to one-
half, strain, add cerate, and evaporate to proper
consistence. *U. S. Ph.*

As a stimulating dressing to blisters.

R. Spanish flies, in fine powder,
Olive oil, each, six ounces.
Oil of turpentine, three ounces.
Yellow wax,
Resin, each, four ounces.

Mix the first three ingredients, place on a
water-bath, and continue heat till the turpentine
has nearly evaporated, stirring occasionally.
Then add the wax and resin, previously melted
together, and heat till well incorporated; re-
move from fire, and stir till cold. *W. Procter.*

R. Powdered Spanish flies, four ounces.
Boiling water, twenty fl. ounces.
Macerate for twelve hours, and evaporate by a
gentle heat to the consistence of syrup; add
Yellow wax, four ounces.

Resin,
Olive oil,
Spirit of turpentine, each, one ounce.
Alcohol, two ounces.

Incorporate well by stirring. Said to be active,
to spread well, and not to deteriorate by keep-
ing. *Martin.*

R. Powdered cantharides, one ounce.
Resinous ointment, seven ounces.
Add the flies to the melted ointment, and stir
briskly, as it concretes on cooling. *Ed. Ph.*

OINTMENT OF EXTRACT OF CANTH-
RIDES.

R. Alcoholic extract of cantharides,
eight grains.
Oil of roses, one drachm.
Beef marrow, two ounces.
Oil of lemon, forty minims.

Melt the marrow, add the extract and stir in the
essential oils.

To promote the growth of the hair. *M. Cap.*

PLASTER WITH CANTHARIDES.

- R. Purified Burgundy pitch, } each,
 " elemi, } sixty
 Oil of Spanish flies, } parts.
 Yellow wax,
 Spanish flies,
 Sulphuric ether, each, one hundred
 and twenty-five parts.
 Powdered camphor, twenty parts.

Put the cantharides in a bottle, add the ether, cork, and let rest for eight days. Melt by a gentle heat the pitch, wax, elemi, and oil, add the Spanish flies, and keep in fusion for two hours, stirring occasionally; finally, mix the camphor. *Houdbine.*

COMPOUND CANTHARIDES PLASTER.

- R. Venice turpentine, four ounces and
 a half.
 Burgundy pitch,
 Cantharides, each, three ounces.
 White mustard seed,
 Black pepper, each, two drachms.
 Bees-wax, one ounce.
 Verdigris, half an ounce.

Melt wax and Burgundy pitch, add turpentine, and while hot, add other ingredients in fine powder. Stir till cold. *Ed. Ph., 1841.*

A powerful and speedy vesicant.

CAMPHORATED CANTHARIDES PLASTER.

- R. Plaster of melilot, two ounces.
 Melt and add
 Powdered cantharides, six drachms.
 Camphor, half a drachm.
 As a vesicant. *Spielmann.*

ISSUE OINTMENT.

- R. Powdered Spanish flies, half an ounce.
 Rose water, two fl. ounces.
 Boil, and add
 Tartar emetic, fifteen grains.
 Evaporate to one-half, strain, and add
 Olive oil, three ounces.
 White wax, one ounce and a half.
 Spermaceti, one ounce.

Boil till all moisture is evaporated. Used as a stimulating application to issues. *Physick.*

CANTHARIDES POMATUM.

- R. Powdered Spanish flies, one drachm.
 Alcohol, one ounce.

Macerate, and filter. Ten parts of this tincture are to be well incorporated with ninety parts of lard. Said to be very effectual in preventing the hair from falling off. *Dupuytren.*

ODONTALGIC PLASTER WITH CANTHARIDES.

- R. Pitch,
 Resin, each, one ounce.
 Melt together, and add
 Storax,
 Cantharides, each, two ounces.
 As an irritant plaster to the cheek in cases of toothache. *Van Mons.*

STIMULATING LINIMENT.

- R. Tincture of cantharides, one fl. drachm.
 Soap liniment, six fl. drachms.
 Mix. In indolent chilblains. *Wardrop.*

AROMATIC CANTHARIDES LINIMENT.

- R. Tincture of cantharides, }
 Oil of thyme, } equal parts.
 Solution of ammonia, }
 Mix. A powerful rubefacient. *Augustin.*

BLISTERING COURT PLASTER.

- R. Cantharides, one drachm and a half.
 Acetic ether, two ounces.
 Infuse for eight days, decant, and add
 Resin, two drachms.
 Spread on court plaster. As a mild vesicant. *Cadet.*

ETHER CANTHARIDALIS.

- R. Powdered cantharides, one part.
 Sulphuric ether, two parts.
 Digest for three days, and express. The product is a green oil containing cantharidin, resin, and coloring matters. If the undiluted oil be brushed a few times over a part, it causes vesication. *Oettinger.*

TAFFETA CANTHARIDALIS.

- R. Ether cantharidalis,
 Sulphuric ether, each, one ounce.
 Purified turpentine,
 Colophony, each, two drachms.
 Mix. A piece of taffeta is stretched on a frame, and twice painted over with a brush dipped in the solution; the process is repeated the next day, and again the third day, always passing the brush in the same direction. After a few days the taffeta is covered with a solution of isinglass. *Oettinger.*

COLLODIUM CANTHARIDALE.

- R. Ether cantharidalis,
 Collodion, equal parts.
 Mix. This mixture is to be applied twice on the part to be blistered. *Oettinger.*

BLISTERING CLOTH.

- R. Aqueous extract of cantharides, four parts.
Gelatin in solution, one part.
Mix and brush over a piece of waxed cloth spread in a frame. When the first layer dries, add a second and a third, always passing the brush in the same direction.
Paris Pharm. Society.

SHAMPOO LIQUID.

- R. Rum, three quarts.
Alcohol, one pint.
Water, one pint.
Tincture of cantharides, half an ounce.
Carbonate of ammonia, half an ounce.
Carbonate of potassa, one ounce.
Dissolve the carbonates in the water, and add the solution to the other ingredients mixed together, and shake up well. To remove dandruff from the hair, by moistening it with the mixture, rubbing, so as to form a lather, and then washing with cold water.
Morfit.

CAPSICUM.

CAYENNE PEPPER.

Several species of capsicum are used under the above official name, as *C. baccatum*, *C. frutescens*, &c., but that usually recognized is *C. annuum*, a native of Asia, but now generally cultivated in Europe and America.

Sex. Syst. Pentand. monog. *Nat. Syst.* Solanaceæ.

Linn. Sp. Pl. 270. Griffith, Med. Bot. 497.

Capsicum has a pungent smell, and a hot, bitterish, acrid taste. It is an active stimulant, and rubefacient. Dose, five to ten grains in powder or pill.

CAYENNE PEPPER PILLS.

- R. Powdered cayenne pepper, one drachm.
Crumb of bread, sufficient.
Distilled water, each, sufficient.
Rub into mass, and form twelve pills. One, three or four times a-day. In debility of the stomach, especially from intemperance. *Ellis.*
R. Powdered cayenne pepper, one scruple.
Extract of gentian, one drachm.
Powdered gentian, sufficient.
Form mass, and divide into sixty pills. In chronic dyspepsia. *Radius.*

CAYENNE PEPPER LOZENGES.

- R. Cayenne pepper, three drachms.
Ginger, two drachms.
Horseradish, each, two drachms.

- Sugar, one ounce.
Conserve of roses, one drachm.
Mix, and make thirty lozenges. One to be chewed occasionally in catarrhal deafness.
Fosbroke.

CAYENNE PEPPER SYRUP.

- R. Simple syrup, two pints.
Tincture of cayenne pepper, one fl. ounce.

Pour the tincture on the heated syrup, and evaporate it from the surface of the latter, until the vapor ceases to ignite on the approach of flame. Then mix immediately. *A. Smith.*

INFUSION OF CAYENNE PEPPER.

- R. Cayenne pepper, half an ounce.
Boiling water, one pint.
Infuse for two hours, and strain. Dose, half a fl. ounce. *U. S. Ph.*

CAYENNE PEPPER GARGLE.

- R. Tincture of cayenne pepper, half fl. ounce.
Rose water, eight fl. ounces.
Mix.

- R. Powdered cayenne pepper, two tablespoonfuls.
Common salt, a teaspoonful.
Boiling vinegar, half a pint.
" water, each, half a pint.
Infuse for an hour, and strain. In bad cases of scarlatina. *Stephens.*

- R. Powdered cayenne pepper, half an ounce.
Magnesia, one drachm.
Boiling water, eight ounces.
" vinegar, each, eight ounces.
Infuse, and strain. Used in same cases, and in same manner as above. *Headley.*

TINCTURE OF CAYENNE PEPPER.

- R. Cayenne pepper, one ounce.
Diluted alcohol, two pints.
Macerate for fourteen days, and filter. *U. S. Ph.*
As a stimulant in low states of fever with gastric insensibility. Dose, one to two fl. drachms. Also used in a diluted state as gargle.

CONCENTRATED TINCTURE OF CAYENNE PEPPER.

- R. Cayenne pepper, four ounces.
Rectified spirit, twelve fl. ounces.
Macerate for seven days and strain. Used as an embrocation for toothache and chilblains. *A. Turnbull.*

TINCTURE OF CAYENNE PEPPER AND CANTHARIDES.

- R. Cayenne pepper, one drachm.
 Cantharides, ten drachms.
 Alcohol, one pint.
 Digest for two days, and filter. Dose, ten drops.
Coze.

VINEGAR OF CAYENNE PEPPER.

- R. Cayenne pepper, one part.
 Vinegar, six parts.
 Digest for some days, and filter. In doses of half a drachm to a drachm, in passive dropsies.
Beral.

EXTRACT OF CAYENNE PEPPER.

- R. Powdered cayenne pepper, eight ounces.
 Diluted alcohol, one pint and a half.
 Macerate for six days, then place in a percolator, pour on diluted alcohol until four pints of tincture are obtained, and evaporate in a water-bath to the consistence of an extract. About two ounces of extract are obtained. Dose, one or two grains made into pills. *W. C. Bakes.*

CERATE OF CAYENNE PEPPER.

- R. Extract of cayenne pepper, one drachm.
 Simple cerate, one ounce.
 Rub together. This acts as a rubefacient in twenty or thirty minutes. *W. C. Bakes.*

CAYENNE PEPPER CATAPLASM.

- R. Powdered cayenne pepper, one ounce.
 Ground mustard,
 Common soap, each, three ounces.
 Alcohol, sufficient.
 Mix. An active rubefacient. *Ammon.*

LOTION OF CAYENNE PEPPER.

- R. Tincture of cayenne pepper,
 “ camphor, each, four fl. ounces.
 Solution of ammonia, two fl. ounces.
 Mix. A very powerful rubefacient. *Ammon.*

CARBO ANIMALIS.

ANIMAL CHARCOAL.

This article is prepared by charring animal substances, usually bones. It requires purification for internal use.

PURIFIED ANIMAL CHARCOAL.

- R. Animal charcoal, one pound.
 Muriatic acid,
 Water, each, twelve fl. ounces.

Mix the acid and water, and pour on charcoal; digest for two days, occasionally stirring. Decant, and wash the charcoal several times with water and dry it. *U. S. Ph.*

- R. Lean of beef or mutton, two parts.
 Broken bones, one part.
 Char by a moderate fire in a coffee-roaster, let cool and pulverize. *Weise.*

Has been given with advantage in scrofulous diseases, chronic glandular swellings, &c., in doses of ten to twenty grains.

POWDER OF ANIMAL CHARCOAL.

- R. Animal charcoal, four grains.
 Sugar, one drachm.
 Mix. To be taken morning and evening, in scrofula. *Weise.*
 R. Animal charcoal, six grains.
 Burnt sponge, twelve grains.
 Powdered liquorice root, half a drachm.

Mix, and form six powders, one to be taken morning and night, in schirrhous goitre. *Riecke.*

- R. Animal charcoal, four grains.
 Powdered liquorice root, four scruples.

Mix, and divide into eight powders, one to be taken morning and evening, dry; a little water drank afterwards. After the eight powders are taken, increase the dose, by half a grain at a time, to four grains. In schirrhous tumors of the breasts. A spare diet to be observed. *Weise.*

- R. Animal charcoal, one scruple.
 Subcarbonate of iron,
 Sugar, each, two scruples.

Mix, and divide into ten powders. One three or four times a-day in incipient hypertrophy of the womb or ovaries. *Clarus.*

OINTMENT OF ANIMAL CHARCOAL.

- R. Animal charcoal, one part.
 Lard, two parts.
 Rub together. *Giordano.*

- R. Animal charcoal, half an ounce
 Ointment of mallows, six drachms

Mix. As a friction to glandular engorgements
Radiis

CARBO LIGNI.

CHARCOAL

Is employed in medicine as an antiseptic and absorbent, and has been found useful in dyspepsia, &c. It also enters into the composition of poultices, &c.

CHARCOAL DENTIFRICE POWDER.

R. Powdered charcoal, four parts.
 " Peruvian bark, two parts.
 " myrrh, one part.
 Mix. *Ph. Slesv.*

R. Powdered charcoal, three drachms.
 " Peruvian bark, two drachms.
 " calamus, one drachm.
 " myrrh, half a drachm.
 Oil of cloves, three drops.
 Mix. *Sartori.*

R. Powdered charcoal, }
 " red saunders, } each, one
 " orange peel, } drachm.
 " Peruvian bark, }
 " cremor tartar, }
 Oil of bergamot, twenty drops.
 Mix. *Schubarth.*

POWDER OF CHARCOAL AND QUASSIA.

R. Powdered charcoal, six drachms.
 " quassia,
 Magnesia, each, one drachm.
 Mix. Dose, a teaspoonful, two or three times
 a-day, in pyrosis. *Heim.*

CHARCOAL LOZENGES.

R. Powdered charcoal,
 " sugar, each, one ounce.
 " vanilla, one drachm.
 " chocolate, three ounces.
 Mucilage of tragacanth, sufficient.
 Beat together, and form lozenges of eighteen
 grains. For fetid breath. *Chevallier.*

CHARCOAL ELECTUARY.

R. Powdered charcoal,
 Carbonate of soda, each, one drachm.
 Electuary of senna, two ounces.
 Mix. Two or three teaspoonfuls a-day in obsti-
 nate constipation. *Radius.*

R. Powdered charcoal, }
 " orris root, } each, one
 " Peruvian bark, } scruple.
 Tincture of myrrh, half a drachm.
 Clarified honey, sufficient.
 Mix. Used as a dentifrice. *L. W. Sachs.*

OINTMENT OF CHARCOAL.

R. Powdered charcoal,
 Storax ointment, each, one ounce.
 Camphor,
 Myrrh, each, two drachms.
 Spirit of turpentine sufficient.

Rub well together. As a dressing to fetid
 ulcers. *Rust.*

CHARCOAL CATAPLASM.

R. Freshly prepared charcoal, sufficient.
 Mix with

Simple cataplasm in a tepid state.

Dub. Ph.

As an application to foul and gangrenous
 ulcers.

R. Powdered charcoal, half an ounce.
 Cataplasm of chamomile, five pounds.

Mix. As above. *Brera.*

CHARCOAL SUPPOSITORY.

R. Cork charcoal,
 Wax, each, one part.
 Fresh butter, two parts.
 Mix. As a suppository in hemorrhoids. *Cadet.*

CARBO MINERALIS.

MINERAL COAL.

ANTHRACOKALI.

R. Carbonate of potassa, six ounces.
 Lime, three and a half ounces.
 Water, four pints.

Proceed as in making liquor potassa, and con-
 centrate the clear solution, by boiling in an
 iron pot, till an oily-looking liquid remains,
 then stir in five ounces of finely-powdered
 mineral coal; remove from the fire, and con-
 tinue stirring till the whole is reduced to an
 uniform powder, which is to be immediately
 put in small, well-stopped bottles. *Polya.*

Much recommended in cutaneous diseases,
 scrofula, chronic rheumatism, &c. Dose, two
 grains, twice or thrice a-day.

POWDER OF ANTHRACOKALI.

R. Anthracokali, two grains.
 Powdered liquorice, five grains.
 Mix. Two to four times a-day. *Polya.*

COMPOUND POWDER OF ANTHRACOKALI.

R. Anthracokali, two grains.
 Washed sulphur, three to five grains.
 Powdered liquorice, two to three
 grains.

Mix. In psora, &c. *Polya.*

R. Anthracokali, two grains.
 Calomel, one-sixth of a grain.
 Powdered liquorice, three grains.

Mix. In syphilitic eruptions. *Polya.*

R. Anthracokali,
Golden sulphuret of
antimony, each, half a grain.
Powdered liquorice, three grains.
Mix. *Polya.*

SULPHURETTED ANTHRACOKALI.

R. Prepared like the simple anthracokali, adding four drachms of sulphur to the coal, and dissolving, &c., as above.
The dose and uses are the same. *Polya.*

CARBONIS IODIDUM.

IODIDE OF CARBON.

R. Tincture of iodine, two fl. ounces.
Solution of potassa, sufficient.

Add the solution of potassa to the tincture, till the latter loses its color, then add

Distilled water, sufficient
to precipitate the iodide, separate and dry by a gentle heat.

OINTMENT OF IODIDE OF CARBON.

R. Iodide of carbon, half a drachm.
Simple ointment, six drachms.

Rub together. As an application to enlarged glands, and obstinate cutaneous affections.

Litchfield.

CARDAMINE.

CUCKOO FLOWER.

This is the *Cardamine Pratensis*, a small herbaceous plant, a native of Europe, and also found in the more northern parts of America.

Sex. Syst. Tetrad. siliquos. *Nat. Syst.* Brassicaceæ.

Linn. Sp. Pl. 915. Baker, Med. Trans. 1. 442.

The part employed is the flowers, which are said to be diuretic and antispasmodic. It is not used in this country, and appears to be little deserving of notice.

CARDAMOMUM.

CARDAMOM.

Under the name of Cardamom a variety of aromatic capsules have been used in medicine, all possessing the same properties and somewhat the same appearance. That recognized by the U. S. Pharm. is the product of the *Elettaria Cardamomum*, a native of Malabar.

Sex. Syst. Monand. monog. *Nat. Syst.* Zingiberaceæ.

Maton, Trans. Linn. Soc. x. 254. Griffith, Med. Bot. 633.

The part used is the seeds; these are angular, somewhat rough, of a reddish or brownish colour, with an agreeable fragrant odour, and a warm, aromatic taste. They are seldom used alone, being chiefly employed in compound preparations.

AROMATIC POWDER.

R. Ginger,
Cinnamon, each, two ounces.
Cardamom seeds,
Nutmeg, grated, each, one ounce.

Mix, and rub together into fine powder.

U. S. Ph.

Stimulant and carminative. Dose, ten to thirty grains, in dyspepsia with flatulence.

AROMATIC CONFECTION.

R. Aromatic powder, five ounces and a half.

Powdered saffron, half an ounce.
Syrup of orange peel, six ounces.
Clarified honey, two ounces.

Rub together, and beat into a mass.

U. S. Ph.

Used as a stimulant to the stomach, in doses of ten to sixty grains.

TINCTURE OF CARDAMOM.

R. Cardamom, bruised, four ounces.
Diluted alcohol, two pints.

Macerate for fourteen days, express, and filter.

U. S. Ph.

A pleasant aromatic stimulant. Dose, one to two fl. drachms.

COMPOUND TINCTURE OF CARDAMOM.

R. Powdered cardamom, six drachms.
" caraway, two drachms.
Bruised cochineal, one drachm.
" cinnamon, five drachms.

Raisins, five ounces.

Diluted alcohol, two pints and a half.

Macerate for fourteen days, express, and filter.

Lond. Ph.

An agreeable aromatic and carminative, in the dose of one or two fl. drachms.

ETHEREAL EXTRACT OF CARDAMOM.

R. Cardamom in powder, at will.
Sulphuric ether, sufficient.

Treat, till twice the weight of the seed is obtained. Evaporate spontaneously until deprived of ether. This extract is fluid, and consists of volatile and fixed oil. It may be used for aromatizing powders, or as an addition to pills.

W. Procter

CAROTA.

CARROT.

The seeds and root of the common carrot, *Daucus carota*, a native of Europe, but now extensively naturalized in the United States, and generally cultivated as an esculent in both countries.

Sex. Syst. Pentand. digyn. *Nat. Syst.* Apiaceæ.

Linn. Sp. Pl. 348. Griffith, Med. Bot. 337.

The parts used are the seeds and root; the former are aromatic, and have a pungent, bitterish taste; the root of the wild carrot has a strong smell, and an acrid, disagreeable taste; both these are similar in their remedial effects, being aromatic and diuretic. The root of the cultivated variety is milder, and is not used internally, but has much reputation as an external application to foul ulcers, &c.

INFUSION OF CARROT SEEDS.

R. Carrot seed, half an ounce—one ounce.
Boiling water, one pint.

Infuse. To be taken during the day as a diuretic, in dropsy and nephritic complaints.

CATAPLASM OF CARROT ROOT.

R. Root of garden carrot, at will.
Scrape down to a pulp. As an application to foul and cancerous ulcers. *Wood.*

R. Root of garden carrot, at will.
Boil till soft, and mash. *Dub. Ph.*

This acts merely as an emollient, and is not possessed of any virtues beyond those of the common poultice.

EXTRACT OF CARROT ROOT.

R. Clarified juice of carrot root, at will.
Evaporate on a water-bath to the consistence of honey. *Swediaur.*

As an application to ulcerated cancers.

CARROT OINTMENT.

R. Carrot root, grated, one pound.
Lard, twenty ounces.
Yellow wax, two ounces.

Heat them together till the water of vegetation is driven off, and the fat has acquired a yellow color; then strain for use. *W. Procter.*

CARTHAMUS.

SAFFLOWER.

This is the florets of *Carthamus tinctorius*, a native of the Levant and Egypt, but cultivated in other parts of Europe and in the United States.

Sex. Syst. Syngen. æqual. *Nat. Syst.* Astera-cææ.

Linn. Sp. Pl. 1162. Griffith, Med. Bot. 410.

The seeds were formerly much used in medicine, but are at present seldom resorted to. The florets are somewhat stimulant, and, in warm infusion, diaphoretic, and are used as a substitute for saffron.

INFUSION OF SAFFLOWER.

R. Safflower, two drachms.
Boiling water, one pint

Infuse, and strain.

Used in domestic practice in the exanthemata, to bring out the eruption.

CARUM.

CARAWAY.

Caraway seeds are the product of *Carum carui*, an umbelliferous plant, a native of Europe, and cultivated both there and in this country.

Sex. Syst. Pentand. digyn. *Nat. Syst.* Apiaceæ.

Linn. Sp. Pl. 378. Griffith, Med. Bot. 318.

The seeds, the only part used, are of a brownish color, with a fragrant but peculiar odor, and an aromatic, warm taste. They are a pleasant aromatic and carminative. Dose, in powder, from a scruple to a drachm.

CARAWAY WATER.

R. Caraway seeds, bruised, one pound and a half.
Water, two gallons.
Mix. Distil a gallon. *Lond. Ph.*

INFUSION OF CARAWAY.

R. Caraway seeds, two drachms
Boiling water, one pint.
Infuse, and strain. *Wood*
In flatulent colic of infants.

SPIRIT OF CARAWAY.

R. Bruised caraway, twenty-two ounces.
Proof spirit, one gallon.
Water, two pints.
Mix, and distil one gallon. *Lond. Ph., 1836.*
R. Oil of Caraway, two fl. drachms.
Proof spirit, one gallon.
Dissolve. *Lond. Ph., 1851*

As a carminative. In doses of one to two fl. drachms.

OIL OF CARAWAY.

R. Caraway seeds, at will.
 Water, sufficient.
 Macerate for some time, distil, and separate the oil. *Van Mons.*

ESSENCE OF CARAWAY.

R. Oil of caraway, one fl. ounce.
 Rectified spirit, nine fl. ounces.
 Mix with agitation. *Dub. Ph.*
 Dose, about twenty drops.

EMBROCATION OF CARAWAY.

R. Oil of caraway,
 " peppermint, each, one scruple.
 " olives, one ounce.
 Wine of opium, half a fl. drachm.
 Mix. As an embrocation on the abdomen of infants in colic. *Radius.*

CARYOPHYLLUS.

CLOVES.

Cloves are the unexpanded flowers of *Caryophyllus aromaticus*, a small tree, a native of the Molucca islands, and also cultivated in many other parts of the East Indies.

Sex. Syst. Icosand. monog. *Nat. Syst.* Myrtaceæ.

Linn. Sp. Pl. 735. *Griffith, Med. Bot.* 298.

Cloves are somewhat nail-shaped, of a brown color, with a strong fragrant smell, and a permanent, pungent, aromatic taste. They are highly stimulating, and are given as a carminative, and excitant of the gastric functions, in the dose of five to ten grains, but are more used as an adjunct to various officinal compounds.

INFUSION OF CLOVES.

R. Bruised cloves, two drachms.
 Boiling water, one pint.
 Macerate for two hours in a covered vessel, and strain. *U. S. Ph.*
 Dose, from one to two fl. ounces.

CLOVE WATER.

R. Bruised cloves, one part.
 Water, eight parts.
 Digest for twelve hours, and distil twelve parts.
 As a stimulant and carminative. *Cottureau.*

SPIRIT OF CLOVES.

R. Bruised cloves, one part.
 Alcohol, eight parts.
 Macerate for some days, and distil eight parts. *Guibourt.*

TINCTURE OF CLOVES.

R. Bruised cloves, one part.
 Alcohol, eight parts.
 Treat with one-half the alcohol at a time, and filter. *Beral.*

WINE OF CLOVES.

R. Bruised cloves,
 " mace, each, one drachm.
 Red wine, one pint.
 Boil, and strain. As a fomentation. *Saunders.*

OIL OF CLOVES.

R. Bruised cloves, at will.
 Water, sufficient.
 Macerate for some time, distil, return the product, and redistil several times, separate the oil, which sinks to the bottom. *Van Mons.*
 Dose, two to six drops.

AROMATIC BALSAM OF CLOVES.

R. Oil of cloves,
 " nutmeg, each, one scruple.
 Spirit of juniper berries, two ounces.
 Mix. As a stimulating friction. *Bories.*

ODONTALGIC MIXTURE.

R. Oil of cloves, twelve drops.
 Tincture of pimpinella,
 Sulphuric ether, each, two drachms.
 Aromatic tincture, one drachm.
 Mix. To be applied to the carious tooth. *Augustin.*

CLOVE PLASTER.

R. Yellow wax, twenty-two parts.
 Suet, twenty-four parts.
 Melt, and add
 Powdered olibanum, sixteen parts.
 " cloves, eight parts.
 Oil of nutmeg, six parts.
 " peppermint, one part.
 Mix. *Van Mons.*
 As an application to the stomach, or over a pained part.

CLOVE CATAPLASM.

R. Rye flour, six or eight spoonfuls.
 Red wine, sufficient
 to make a cataplasm; add
 Powdered cloves,
 Grated nutmeg, each, a teaspoonful
 Mix well. As an application to the stomach, to check nausea or pain. *Radius.*

CLOVE BAG.

- R. Powdered cloves, two drachms.
 " marjoram, half an ounce.
 " rosemary, one ounce.

Mix, and quilt between folds of linen, dip in Cologne water, and apply to the pit of the stomach, to check nausea. *Phæbus.*

CASCARILLA.

CASCARILLA.

This is the bark of *Croton eleuthera*, a shrub found in several of the West India islands, and also, perhaps, derived from other species.

Sex. Syst. Monœc. monadelph. *Nat. Syst.* Euphorbiacæ.

Swartz. H. Ind. Oc. 11. 1183. Griffith, Med. Bot. 596.

In rolled fragments, whitish externally, and of a chocolate color within, of an aromatic odor, and a spicy, bitter taste. It is an aromatic tonic, and is useful in debilitated conditions of the stomach and bowels. Dose, in powder, from a scruple to half a drachm.

COMPOUND CASCARILLA POWDER.

- R. Powdered cascarilla, ten grains.
 " opium,
 " ipecacuanha,
 each, one grain.

Mix. To be taken every five hours in obstinate diarrhœa. *Radius.*

- R. Powdered cascarilla, one drachm.
 " hartshorn,
 " cuttlefish bone, each, four drachms.
 " amber, two drachms.

Mix. One drachm to be taken in a glass of sugar and water, at bedtime, against nocturnal pollutions. *Pierquin.*

INFUSION OF CASCARILLA.

- R. Bruised cascarilla, one ounce.
 Boiling water, one pint.

Macerate for two hours in a covered vessel, and strain. *U. S. Ph.*

Dose, two fl. ounces.

ALKALINE INFUSION OF CASCARILLA.

- R. Bruised cascarilla, three ounces.
 Carbonate of potassa, two drachms.
 Boiling water, sixteen fl. ounces.

Macerate for two hours, and filter. Dose, a spoonful, two or three times a-day, as an antacid and tonic. *Palat. Ph.*

EXTRACT OF CASCARILLA.

- R. Bruised cascarilla, one part.
 Alcohol, five parts.

Treat the cascarilla several times with the alcohol, distil, and evaporate the residue. Dose, from ten grains to a drachm. *Guibourt.*

MIXTURE OF EXTRACT OF CASCARILLA.

- R. Extract of cascarilla, one drachm.
 White sugar, six drachms.
 Oil of chamomile, twenty drops.
 Tragacanth, half a drachm.
 Wine of opium, twenty-five drops.
 Cinnamon water, two fl. ounces.
 Peppermint water, four fl. ounces.

Mix. A spoonful every hour in dysentery.

Augustin.

TINCTURE OF CASCARILLA.

- R. Powdered cascarilla, five ounces.
 Proof spirit, two pints.

Macerate fourteen days, and filter. *Lond. Ph.*

Dose, a fluidrachm.

COMPOUND WINE OF CASCARILLA.

- R. Powdered cascarilla, one ounce and a half.
 Contused orange peel, one ounce.
 " cinnamon, two drachms.
 White wine, twenty-seven fl. ounces.

Digest for twenty-four hours, and filter. As a stomachic and tonic, in doses of two ounces, four times a-day. *Phæbus.*

CASSIA FISTULA.

PURGING CASSIA.

This is the fruit or pod of the *Cassia fistula*, a large tree, a native of Egypt, and many parts of Asia, and cultivated in most tropical regions.

Sex. Syst. Decand. monog. *Nat. Syst.* Fabacæ.

Linn. Sp. Pl. 540. Griffith, Med. Bot. 253.

The pods are a foot or more in length, cylindrical, of a dark brown color, with two longitudinal furrows on one side, and one on the other. They are internally divided into numerous cells, each containing one seed enveloped in a soft pulp. This latter, which is the part used, has a faint nauseous smell, and a sweet, mucilaginous taste. It is laxative, in doses of one or two drachms, and purgative in those of one or two ounces.

PULP OF PURGING CASSIA.

- R. Purgive cassia, bruised, sufficient.
 Pour boiling water on it to soften the pulp, strain, and evaporate to proper consistence.

U. S. Pharm.

Dose, one drachm.

CONFECTION OF CASSIA.

R. Cassia pulp, half a pound.
 Manna, two ounces.
 Tamarind pulp, one ounce.
 Syrup of roses, eight fl. ounces.

Mix well, and evaporate to proper consistence.
Lond. Ph. 1836.

Slightly laxative.

MIXTURE OF CASSIA PULP.

R. Pulp of cassia, one ounce.
 Infusion of rhubarb, a drachm and
 a half.

Syrup of roses,
 Manna, each, one ounce.
 Mix well. A mild laxative. *Pierquin.*

CASSIA MARILANDICA.

AMERICAN SENNA.

This species is an indigenous perennial plant, with showy, bright yellow flowers, growing in moist places and on the banks of streams.

Sex. Syst. Decand. monog. *Nat. Syst.* Fabaceæ.

Linn. Sp. Pl. 541. *Griffith, Med. Bot.* 260.

The leaves, which are the officinal portion, are usually exposed for sale in small packages, like the other herbs put up by the Shakers. They have a faint, unpleasant odor, and a nauseous taste like senna, of which they possess all the qualities, but are not quite as active.

INFUSION OF AMERICAN SENNA.

R. American senna, one ounce and a half.
 Coriander seed, bruised, one drachm.

Boiling water, one pint.
 Macerate in a covered vessel one hour, and strain.

Dose, a wineglassful.

CASTOREUM.

CASTOR.

Castor is a peculiar substance obtained from membranous follicles in the common beaver or *Castor fiber*, situated between the organs of generation and the anus. It is in the form of solid, unctuous masses, contained in small, wrinkled, brownish sacs. It has a strong, unpleasant smell, and a bitter, acrid taste. It is stimulant and antispasmodic, and has been much praised in the neuroses. In this country it is not much used. Dose, from ten to twenty grains.

COMPOUND POWDER OF CASTOR.

R. Castor, twelve grains.
 Powdered cascarilla, one scruple.
 Magnesia, one scruple and a half.

Mix, and divide into three powders. *Saunders.*

R. Castor, }
 Powdered valerian, } each,
 Sugar, } ten grains.

Mix, and divide into three doses *Phabus.*
 As antispasmodics.

BOLUS OF CASTOR.

R. Assafetida, }
 Valerian, } each, half an ounce.
 Castor, }
 Amber, }
 Camphor, } one scruple.
 Syrup, } sufficient.

Mix, and form boluses of ten grains each. Said to be efficacious in hysteria, neuralgia, and vertigo. *Foy.*

CASTOR PILLS.

R. Assafetida, one drachm.
 Galbanum,
 Myrrh, each, half a drachm.
 Castor, fifteen grains.
 Tincture of valerian, sufficient.

Beat together, and divide into seventy-two pills. Dose, three to eight, three times a-day, in hysteria. *Phabus.*

PILLS OF CASTOR AND SUCCINIC ACID.

R. Castor, one drachm.
 Succinic acid, half a drachm.
 Extract of gentian, sufficient.

Beat together, and form twenty-four pills. Three, morning and night, in hysteria. *Ellis.*

TINCTURE OF CASTOR.

R. Bruised castor, two ounces.
 Alcohol, two pints.

Digest for seven days, express, and filter.

U. S. Ph.

Dose, from thirty drops to two fl. drachms.

ETHEREAL TINCTURE OF CASTOR.

R. Bruised castor, two ounces.
 " saffron, one ounce.
 Spirit of sulphuric ether, twelve ounces.

Digest for twenty-four hours, and filter. As an antispasmodic. In doses of thirty drops.

Spielmann.

AMMONIATED TINCTURE OF CASTOR.

R. Bruised castor, two ounces and a half.
 Contused assafetida, ten drachms.
 Spirit of ammonia, two pints.

Digest for seven days, in a closed vessel, express, strain, and filter. *Ed. Ph.*

A powerful stimulant, and antispasmodic, in cases of spasm of the stomach, hysteria, &c. Dose, thirty drops to two fl. drachms.

COMPOUND TINCTURE OF CASTOR.

R. Assafetida, two drachms.
 Opium, half a drachm.
 Oil of amber, one drachm.
 Castor, half an ounce.
 Alcohol, four ounces.

Digest for four days, express, and filter. Dose, thirty to forty drops in wine, in hysteria.

Ferrara. Ph.

COMPOUND PLASTER OF CASTOR.

R. Wax plaster, sixteen ounces.
 Soap, four ounces.

Melt together, and add, on cooling,

Camphor, one ounce.
 Castor, half an ounce.

Incorporate well. Highly praised in headache, rheumatism, &c. *Giordano.*

COMPOUND SPIRIT OF CASTOR.

R. Castor, half an ounce.
 Assafetida, two drachms.
 Oil of amber, one drachm.
 " savine,
 " rue, each, half a drachm.
 Alcohol, ten ounces.

Digest, distil, and add

Empyreumat. carb. of ammonia, two ounces.
 Camphor, one drachm.

Distil again. Dose, twenty to forty drops.

Cottureau.

CATALPA.

CATALPA.

The *Catalpa cordifolia* is a beautiful native tree, principally found in the southern and south-western States.

Sex. Syst. Diand. monog. *Nat. Syst.* Bignoniaceæ.

Elliot, Bot. i. 24. Lindley, Flor. Med. 499.

The bark is said to be vermifuge, but the part used in medicine is principally the pods. These have been found efficacious in asthma.

DECOCTION OF CATALPA PODS.

R. Catalpa pods, half an ounce.
 Water, sufficient

to obtain eight ounces of decoction; add

Oxymel of squills, half an ounce.

R. Catalpa pods, half an ounce.
 Seneka, two drachms.
 Water, sufficient

to obtain eight ounces of decoction; add

Oxymel of squills, one ounce.

In spoonful doses, in chronic, nervous asthma.

Brera.

R. Catalpa pods, three or four.
 Water, twelve fl. ounces.

Boil down to six ounces, to be given in two doses, morning and night. *Antomachi.*

CATARIA.

CATNEP.

This is the leaves, or whole herb of *Nepeta Cataria*, a perennial herbaceous plant, a native of Europe, and naturalized in the United States.

Sex. Syst. Didynam. gymnos. *Nat. Syst.* Lamiales.

Linn. Sp. Pl. 797. Griffith, Med. Bot. 512.

It has a peculiar, somewhat unpleasant odor, and a bitterish, aromatic taste. It is stimulant, carminative, &c., and is used in the flatulent colic of infants, and as an emmenagogue, &c.

INFUSION OF CATNEP.

R. Catnep, two drachms.
 Boiling water, eight fl. ounces.

Infuse in a covered vessel. Dose, a tea-spoonful occasionally, whilst hot, for an infant in flatulent colic; to be taken freely, as a diaphoretic, or emmenagogue.

CATECHU.

CATECHU.

Catechu is an extract procured from the wood of *Acacia catechu*, and also from other trees. The *Acacia catechu* is a native of the East Indies, and is now cultivated in some of the West India islands.

Sex. Syst. Polygam. monœc. *Nat. Syst.* Fabaceæ.

Linn. Sp. Pl. 409. Griffith, Med. Bot. 268.

Catechu comes in masses of various forms, of a rusty-brown color externally, but paler within. It is inodorous, but has a bitter and astringent taste, followed by a sensation of sweetness. It principally consists of tannin and extractive. It is tonic and astringent, and is used in cases where astringents are required. The dose is from ten grains to half a drachm, repeated as required.

COMPOUND POWDER OF CATECHU.

- R. Powdered catechu, fifteen grains.
 " salts, two grains.

Mix, and make powder; to be taken after each liquid stool, in diarrhœa from a weakened condition of the bowels.

A. T. Thomson.

- R. Catechu,
 Kino, each, two ounces.
 Cinnamon,
 Nutmeg, each, half an ounce.

Pulverize, mix, and pass through a fine sieve.

An aromatic astringent. Dose, fifteen to twenty grains.

Dub. Ph.

- R. Powdered catechu,
 " cascarilla,
 " gum Arabic, } equal parts.
 Aromatic powder,

Mix. Dose, a scruple every two hours, in chronic diarrhœa.

Augustin.

BOLUS OF CATECHU.

- R. Powdered catechu, twenty-four grains.
 Extract of opium, two grains.
 Conserve of roses, sufficient.

Mix, and form two boluses, one to be taken morning and evening, in chronic diarrhœa.

Foy.

PILLS OF ALUM AND CATECHU.

- R. Alum, six grains.
 Extract of opium,
 Catechu, each, one grain.

Mix, and divide into six pills, one to be given every two to four hours. In passive hemorrhage and atonic mucous discharges.

Ellis.

COMPOUND CATECHU BOLUS.

- R. Powdered catechu, one scruple.
 Confection of opium, twelve grains.
 Aromatic confection, sufficient.

Make a bolus, to be taken twice a-day, in inordinate flow of the menses.

Babington.

- R. Powdered catechu, twelve parts.
 " alum, six parts.
 " opium, two parts.
 Syrup of red roses, sufficient.

Form a mass, and divide into pills of six grains each. One or two a-day, in the decline of gonorrhœa.

Foy.

PILLS OF CATECHU AND LIQUORICE.

- R. Catechu, half a drachm.
 Gum Arabic,
 Extract of liquorice, each, two drachms.

- Mastich, one drachm.
 Syrup of mallows, sufficient.

Form mass, and divide into pills of three grains each. Four, three times a-day, in chronic catarrh.

Van Mons.

LOZENGES OF CATECHU.

- R. Catechu, one ounce.
 Sugar, half an ounce.
 Cloves, one scruple.
 Essence of citron, half a scruple.
 Mucilage of tragacanth, sufficient.

Beat into mass, and divide into lozenges of ten grains each. One, occasionally, to arrest fetor of breath.

Brunswick Ph.

ELECTUARY OF CATECHU.

- R. Catechu,
 Kino, each, four ounces.
 Cinnamon,
 Nutmeg, each, one ounce.
 Opium, diffused in a little sherry wine, one drachm and a half.
 Syrup of red roses, reduced to the consistence of honey, one pint and a half.

Pulverize the solids, mix the opium and syrup, add the powders, and beat into a mass.

Ed. Ph.

In diarrhœa and chronic dysentery, in doses of half a drachm to a drachm.

- R. Catechu,
 Balsam of tolu, each, one drachm.
 Peruvian bark, one ounce.
 Syrup of red roses, sufficient.

Beat together. One drachm morning and evening, in leucorrhœa and gonorrhœa.

Brera.

INFUSION OF CATECHU.

- R. Catechu, one drachm.
 Boiling water, one pint.
 Digest for six hours, and strain.

Guibourt.

COMPOUND INFUSION OF CATECHU.

- R. Powdered catechu, half an ounce.
 Bruised cinnamon, one drachm.
 Boiling water, one pint.

Macerate for an hour in a covered vessel, and strain.

U. S. Ph.

Dose, from one to three fl. ounces, three or four times a-day.

TINCTURE OF CATECHU.

- R. Catechu, three ounces.
 Cinnamon, bruised, two ounces.
 Diluted alcohol, two pints.

Macerate for fourteen days, express, and filter.

U. S. Ph.

A pleasant astringent. Dose, thirty drops to three fl. drachms.

CATECHU COLLUTORY.

R. Catechu,
Myrrh, each, eight parts.
Balsam of Peru, one part.
Spirit of cochlearia,
Alcohol, each, twenty-four parts.

Mix, and digest for four days, and filter. As a mouth-wash, in a spongy condition of the gums. *Saunders.*

MIXTURE OF CATECHU AND LOGWOOD.

R. Extract of logwood, three drachms.
Tincture of catechu, two fl. drachms.
Water, seven fl. ounces.

Mix. Two tablespoonfuls, every three or four hours, in chronic stages of diarrhœa and dysentery. *Ellis.*

ANTI-EMETIC MIXTURE.

R. Catechu, one drachm.
Columbo, thirty grains.
Canella, twenty grains.
Boiling water, four fl. ounces.

Mix, and digest for eight hours, strain, and add

Syrup of red roses, one fl. ounce.

In spoonful doses. *Pierquin.*

INJECTION OF CATECHU.

R. Catechu,
Myrrh, each, one drachm.

Dissolve in

Lime water, four fl. ounces,

and strain. As an injection, in chronic leucorrhœa and gonorrhœa.

AROMATIC PASTILLES OF CATECHU.

(*Cachou Aromatisé.*)

R. Extract of liquorice,
Water, each, three ounces and a half.

Dissolve on a water-bath, and add

Powdered catechu, four hundred and sixty-two grains.

" gum Arabic, two hundred and thirty-one grains.

Evaporate to the consistence of an extract, and incorporate

Powdered mastich, }
" cascarilla, } each,
" charcoal, } thirty grains.
" orris root, }

Reduce to proper consistence, remove from fire, and add

Oil of peppermint, thirty drops.

Tincture of ambergris,

" musk, each, ten drops.

Mix, and form into one grain pills. Used to aromatize the breath. *Gray*

CEANOTHUS.

NEW JERSEY TEA.

Several species of *Ceanothus* have medical properties, but the only one in use in this country, is the *C. Americanus*, a suffruticose plant, found in most parts of the United States.

Sex. Syst. Pentand. trigyn. *Nat. Syst.* Rhamnaceæ.

Linn. Sp. Pl. 284. *Griffith, Med. Bot.* 218.

The leaves and root are bitter and astringent, but the root is the most active. They have been used in a variety of cases, to which astringents are applicable.

DECOCTION OF NEW JERSEY TEA.

R. New Jersey tea root, two drachms.

Water, one pint.

Boil for a quarter of an hour, and strain. Said by Ferrein to be efficacious in gonorrhœa and syphilis; and by Dr. Hubbard, in dysentery and also as a gargle in aphthous sore mouth, and in ulcerations of the fauces.

CENTAUREA BENEDICTA.

BLESSED THISTLE.

An annual plant, native of the south of Europe, and become naturalized in some parts of the United States.

Sex. Syst. Syngen. frust. *Nat. Syst.* Astera-cææ.

Linn. Sp. Pl. 1296. *Griffith, Med. Bot. (Cnicus.)* 409.

The whole herb is used; it has a faint nauseous odor, and a very bitter taste. It is used as a tonic, diaphoretic, or emetic. Dose of powder, as tonic, a scruple to a drachm.

INFUSION OF BLESSED THISTLE.

R. Blessed thistle, three drachms.

Boiling water, one quart.

Infuse for half an hour, and strain. As a diaphoretic. *Par. Cod.*

R. Blessed thistle, half an ounce.

Cold water, one pint.

Infuse, and strain. As a tonic, in the dose of two fl. ounces. *Wood.*

WINE OF BLESSED THISTLE.

R. Blessed thistle, one ounce and a half.

Columbo, three drachms.

Cinnamon, two drachms.

Wine, two pints

Macerate for twenty-four hours, and strain.
Dose, one to four spoonfuls in the morning, as a
stomachic. *Wirtzburg Ph.*

CENTAURIUM.

CENTAURY.

The herbs or flowering heads of *Erythraea centaurium*, a small annual plant, indigenous to many parts of Europe.

Sex. Syst. Pentand. monog. *Nat. Syst.* Gentianaceæ.

Linn. (*Chironia*,) *Sp. Pl.* 332. *Griffith, Med. Bot.* 459.

This plant is very bitter, and is analogous in its properties to gentian. In this country, its use has been superseded by the American centaury or sabbatia. Dose of the powder, from thirty grains to a drachm.

PORTLAND POWDER.

R. Centaury,
Germander,
Gentian,
Round aristolochia,
Ground pine, equal parts.

Powder, and mix. Once highly celebrated in the cure of gout. Dose one drachm.

EXTRACT OF CENTAURY.

R. Centaury, one part.
Hot water, eight parts.

Macerate in one-half the water, express, and strain; repeat the process with the remainder of the water, unite the liquids, and evaporate.

Guibourt.

PILLS OF EXTRACT OF CENTAURY.

R. Extract of centaury, one drachm.
Myrrh, two drachms.
Balsam of Peru, one scruple.

Mix, and form into pills of three grains. Four, three times a-day, as a tonic and stomachic.

St. Marie.

COMPOUND WINE OF CENTAURY.

R. Centaury, }
Apocynum, } each, one ounce.
Madder, }
White wine,
Water, each, a pint and a half.

Boil down to two-thirds, and add to the strained liquid

Syrup of orange peel, two ounces.

A cupful, morning and evening, in jaundice.

Beral.

MIXTURE OF EXTRACT OF CENTAURY.

R. Extract of centaury, one drachm.
Bitter almonds, two drachms.
Chamomile water, two ounces.

Rub well together, and strain. As a febrifuge.
Foy.

CERA.

WAX.

Cera Flava, or yellow wax, is a concrete substance, deposited by the honey bee, *Apis mellifica*.

Cera Alba, or white wax, is prepared from the yellow, by melting it, pouring into thin layers, wetting and turning frequently, and exposing to the air and light on frames.

Many plants yield a concrete principle called vegetable wax, which corresponds in many particulars with that of the bee.

Wax is much employed in pharmacy in the formation of cerates and plasters. It has also been used as a medicine, in the dose of a teaspoonful.

WAXED CLOTH.

R. White wax, eight parts.
Olive oil, four parts.
Turpentine, one part.

Melt together, and spread upon linen or muslin stretched in a frame. Used for making blistering cloth, and vesicating taffetas.

WAX PLASTER.

R. Yellow wax, three pounds.
Suet, each, one pound.
Resin, one pound.

Melt with a moderate heat, and stir till cold. Used to promote discharges from blistered surfaces.

Lond. Ph., 1836.

ROSE LIP SALVE.

R. White wax, one ounce.
Oil of almonds, two ounces.
Alkanet root, one drachm.

Melt and digest till well colored, strain, and add Otto of roses, six drops.

Par. Codex.

SIMPLE CERATE.

R. Lard, eight ounces.
White wax, four ounces.

Melt together, and stir constantly till cool.

U. S. Ph.

R. White wax, three parts.
Spermaceti, one part.
Olive oil, six parts.

Heat together gently and stir till cold. *Ed. Ph.*

CETACEUM.

SPERMACETI.

A peculiar concrete substance, obtained from the head of *Physeter macrocephalus*, or spermaceti whale. It is white, semi-transparent, friable, soft, somewhat oily to the touch, insipid, but with a faint odor. It has been used as a demulcent, and also enters into the composition of several ointments and cerates.

COMPOUND POWDER OF SPERMACETI.

R. Spermaceti, two drachms.
Nitrate of potassa, one drachm.
Orris root,
Sugar, each, one ounce.
Powder well, and mix. A teaspoonful, three or four times a-day, in catarrh. *Radiis.*

SPERMACETI MIXTURE.

R. Spermaceti, two drachms.
Sugar, three drachms.
Paregoric elixir, half fl. ounce.
Water, eight fl. ounces.

Rub spermaceti and sugar together with the yolk of an egg; add the water and paregoric gradually. A tablespoonful, several times a-day. In catarrh. *Ellis.*

R. Spermaceti, half an ounce.
Gum Arabic, two drachms.
Syrup, half an ounce.
Water of bitter almonds, one drachm.
Water, two ounces and a half.

Rub together, and strain. A dessertspoonful every two hours. In catarrh. *Phæbus.*

R. Spermaceti, two drachms.
Olive oil, one drachm.
Powdered gum Arabic, half an ounce.
Water, four fl. ounces.

Triturate the spermaceti with the oil, until reduced to a paste, then add the gum, and lastly the water, gradually. *W. Procter.*

R. Spermaceti, two drachms.
Yolk of egg, one.
Powdered gum Arabic, two drachms.
Paregoric elixir, six fl. drachms.
Antimonial wine, four fl. drachms.
White sugar, three drachms.
Water, six fl. ounces.

Mix. Rub the spermaceti with the yolk of egg, then add the other ingredients separately; when well incorporated, introduce the water gradually. A tablespoonful every two or three hours to an adult. Useful in catarrh, especially of measles. *Dewees.*

SPERMACETI OINTMENT.

R. Spermaceti, six drachms.
White wax, two drachms.
Olive oil, three fl. ounces.

Mix over a gentle fire, and stir till cold.

Lond. Ph.

Mild dressing for open surfaces.

OINTMENT OF SPERMACETI AND ROSE WATER.

R. Rose water, one fl. ounce.
Oil of almonds, two fl. ounces.
Spermaceti, half an ounce.
White wax, one drachm.

Melt together, by a water-bath, the oil, spermaceti and wax; add rose water, and stir till cold. *U. S. Ph.*

A pleasant and cooling application to irritable surfaces, well known as *cold cream*.

COLD CREAM WITHOUT SPERMACETI.

R. White wax, one ounce.
Oil of almonds, four fl. ounces.
Rose water, two fl. ounces.
Borax, half a drachm.
Oil of roses, five minims.

Dissolve the wax in the oil of almonds, by a gentle heat, also dissolve the borax in the rose water, and add the solution to the heated mixture, stirring till cold, then add the oil of roses. *Turnbull.*

SPERMACETI CERATE.

R. Spermaceti, one ounce.
White wax, three ounces.
Olive oil, six fl. ounces.

Melt wax and spermaceti together, then add the oil previously heated, and stir till cold. *U. S. Ph.*

An excellent dressing for blisters, wounds, &c.

SPERMACETI LIP SALVE.

R. Spermaceti,
White wax, each, half an ounce.

Melt, and add

Oil of almonds, one ounce.
" lavender, twenty drops.
" bergamot, ten drops.
" cloves, two drops.

Rub well together. *Niemann.*

R. Spermaceti, one ounce and a half.
White wax, nine drachms.
Oil of almonds, twelve ounces.
Alkanet root, two ounces.
Otto of roses, one drachm.

Digest the first four ingredients in the heat of a water-bath for four hours, strain, and add the otto of roses. *Gray.*

SPERMACE TI LINIMENT.

- R. Spermaceti,
 White wax, each, half a drachm.
 Oil of almonds, one ounce.
 White lead,
 Litharge, each, twenty grains.
 Gum lac, sufficient.

Mix. As a cosmetic, but the presence of the lead requires it to be used with caution.

Pierquin.

SULTANA OINTMENT.

- R. Spermaceti, eight parts.
 Butter of cacao, sixteen parts.
 Oil of almonds, thirty-two parts.
 Balsam of Peru, one part.

Melt together, and add
 Orange-flower water, one part.
 Stir constantly till cold. *Niemann.*

- R. Spermaceti,
 White wax, each, one drachm and
 a half.
 Oil of almonds, three ounces.
 Melt together, and add
 Rose water, two ounces.
 Tincture of benzoin, thirty drops.
 Mix well. *Taddei.*

CETRARIA.

ICELAND MOSS.

This is a lichen found in great abundance in the northern regions of both continents.

Sex. Syst. Cryptog. lichen. *Nat. Syst.* Lichenaceæ.

When dried it is inodorous, but has a bitter, mucilaginous taste. It is demulcent, tonic, and nutritious, and is useful in pulmonary and other complaints, in which the local disease is attended with a debility of the digestive organs, or of the general system. Dose of the powder, thirty grains to a drachm.

DECOCTION OF ICELAND MOSS.

- R. Iceland moss, half an ounce.
 Water, one pint and a half.
 Boil down to a pint; express, and strain. *U. S. Ph.*

To be taken during the twenty-four hours.

- R. Iceland moss, one ounce and a half.
 Water, one quart.

Macerate the moss in a weak solution of carbonate of potassa, before subjecting it to decoction. Boil, and strain. *Van Mons.*

In this preparation the bitter principle is extracted by the potassa, and the result is a mere mucilaginous demulcent.

ICELAND MOSS JELLY.

- R. Iceland moss, two parts.
 Water, twenty parts.
 Boil down to six parts and strain, then add
 Sugar, four parts.
 Set by to cool. *Augustin.*

ICELAND MOSS MIXTURE.

- R. Iceland moss, two ounces.
 Hartshorn shavings, one ounce.
 Water, one quart.

Boil down to a pint, strain, and add
 Wine of opium, fifteen drops.
 In phthisis, to be taken during the day. *Brera.*

- R. Iceland moss, two drachms.
 Powdered salep, ten grains.
 " cinnamon, eight grains.
 Water, twelve fl. ounces.

Boil to six ounces, and add
 Syrup, sufficient.
 Said to be useful in phthisis, leucorrhœa, and
 hooping-cough, in doses of four ounces, three or
 four times a-day. *Augustin.*

- R. Decoction of Iceland
 moss, seven fl. ounces and a half.
 Diluted sulphuric acid, one fl. drachm.
 Syrup, four fl. drachms.
 Laudanum, fifty drops.

Mix. A wineglassful, three times a-day, in
 phthisis. *A. T. Thomson.*

ICELAND MOSS CHOCOLATE.

- R. Fine chocolate, four pounds.
 Sugar, two pounds.
 Powdered Iceland
 moss, one pound and a half.
 Tragacanth,
 Cinnamon, each, four ounces.
 Water, sufficient.

Rub into an uniform paste. *Cadet.*

CETRARINE.

- R. Iceland moss,
 in coarse powder, one pound.
 Alcohol, (.883,) four pounds.

Boil for half an hour, permit to cool, till no va-
 pors are given off, express, and add to the fluid
 Muriatic acid, three drachms.

and four times its bulk of distilled water. Let
 rest for a night in a closed matrass; then de-
 cant, throw deposit on filter, and press; while
 still moist, wash with alcohol or ether; then
 treat with boiling alcohol, filter, and permit ce-
 trarine to precipitate. *Herberger.*

POWDER OF CETRARINE.

R. Cetrarine,
 Gum Arabic, each, two grains.
 White sugar, half a scruple.
 Useful in intermittent fevers, in doses of eight grains every two hours during the apyrexia.
 Muller.

CHELIDONIUM.

CELANDINE.

The herb of *Chelidonium majus*, a perennial herbaceous plant, native of Europe, but generally naturalized in this country.

Sex. Syst. Polyand. polyg. *Nat. Syst.* Papaveraceæ.

Linn. Sp. Pl. 723. Griffith, Med. Bot. 130.

Exudes, when broken, an orange-colored, fetid juice. Its taste is extremely bitter and acrid, leaving a burning sensation. The root is the most powerful. It has been used as a drastic hydragogue, and also in scrofula, &c. The dose of the dried root is half a drachm to a drachm; of the fresh juice, thirty to forty drops.

EXTRACT OF CELANDINE.

R. Juice of celandine, at will.
 Boil so as to coagulate; strain, evaporate, and at the close of evaporation add the coagulum.
 Van Mons.

Dose, five to fifteen grains, as a hydragogue purgative.

PILLS OF CELANDINE.

R. Extract of celandine,
 Ammoniac, each, one drachm
 and a half.
 Soap, one drachm.
 Mix, and form pills of four grains. Two or three a-day, in engorgements of the viscera, especially of the spleen.
 Schubarth.

EXTRACT OF CELANDINE MIXTURE.

R. Extract of celandine, two drachms
 and a half.
 " henbane, one scruple.
 Sulphate of potassa, one ounce.
 Tartar emetic, one grain.
 Elder water, six fl. ounces.
 Oxy-mel of squills, one fl. ounce.
 Mix. A dessertspoonful every two hours, as a hydragogue.
 Augustin.

CHENOPODIUM.

WORMSEED.

This is the fruit of *Chenopodium anthelminticum*, a native plant, found in most parts of the country.

Sex. Syst. Pentand. digyn. *Nat. Syst.* Chenopodiaceæ.

Linn. Sp. Pl. 320. Griffith, Med. Bot. 537.

The fruit is in small grains, of a greenish-yellow color, with a bitterish, aromatic, pungent taste, and a nauseous peculiar smell. Besides the seeds, the expressed juice of the whole plant is used, and the essential oil. They are all efficient anthelmintics. The first are given in the form of an electuary; the second in tablespoonful doses; the oil is given in the dose of five to ten drops to a child three years old.

DECOCTION OF WORMSEED.

R. Fresh leaves of wormseed, one ounce.
 New milk, one pint.
 Orange peel, two drachms.
 Boil, and strain. Dose, a wineglassful twice a-day.

OIL OF WORMSEED.

R. Bruised wormseed, at will.
 Water, sufficient
 to cover the seed; infuse for some hours, distil, and separate the oil. More commonly, however, the whole herbaceous portion is subjected to distillation.

Dose, from four to eight drops for a child, night and morning, for three or four days, to be followed by a cathartic.
 Coxe.

WORMSEED OIL MIXTURE.

R. Oil of wormseed, one drachm.
 Sugar, "
 Gum Arabic, each, one drachm
 and a half.

Mix, and add

Mint water, two and a half fl. ounces.

A teaspoonful, four times a-day for two days, to a child, to be followed by a purge.
 Ellis.

R. Oil of wormseed, one fl. drachm
 and a half.
 Castor oil, three fl. ounces.
 Oil of aniseed, ten drops.

Mix and add

Syrup of rhubarb, or

" of senna, one fl. ounce.

Mix. Dose, a teaspoonful night and morning, to a child two years old. *Am. Jour. Pharmacy.*

CHIMAPHILA.

PIPSISSEWA.

The leaves of *Chimaphila umbellata*, an evergreen plant, found in the northern portions of both continents.

Sex. Syst. Decand. monog. *Nat. Syst.* Pyrolaceæ.

Linn. (*Pyrola*) Sp. Pl. 568. Griffith, Med. Bot. 421.

Pipsissewa has a peculiar and rather pleasant

odor, especially when fresh, and an astringent, bitter, and somewhat sweetish taste. It is tonic, diuretic, and astringent, and has been much used in nephritic complaints, dropsy, &c.

DECOCTION OF PIPSISSEWA.

R. Pipsissewa, one ounce.
Water, one pint and a half.

Boil down to a pint, and strain. *U. S. Ph.*

To be taken during the day, in dropsy and chronic affections of urinary organs.

COMPOUND DECOCTION OF PIPSISSEWA.

R. Decoction of pipsissewa, one pint.
Solution of carbonate of potassa, two drachms.

Mix. Four tablespoonfuls, three times a-day, in same cases as above. *Foy.*

R. Pipsissewa, half an ounce to one ounce.
Water, twelve fl. ounces.

Boil to six ounces, and add

Gin, two ounces.

Let stand for two hours, and strain. Two tablespoonfuls four times a-day, in dropsy.

Radius.

R. Pipsissewa, six drachms.
Water, twelve ounces.

Boil to six ounces, and add at close

Senna, two drachms.

Strain. A tablespoonful every two hours as a diuretic. *Radius.*

EXTRACT OF PIPSISSEWA.

R. Pipsissewa, one pound.
Water, one gallon.

Macerate for twenty-four hours, boil down to two pints, strain, and evaporate to proper consistence. Dose, ten to thirty grains.

PIPSISSEWA PILLS.

R. Extract of pipsissewa,
Resin of guaiacum,
each, one drachm and a half.
Precipitated sulph. of
antimony, twelve grains.

Beat together, and make pills of two grains. Dose, five, twice a-day, in arthritic affections.

PIPSISSEWA SYRUP.

R. Powdered pipsissewa, four ounces.
Water, eight fl. ounces.

Macerate for thirty-six hours, percolate so as to obtain a pint of fluid, evaporate one-half, and add

Sugar, twelve ounces.

One to two tablespoonfuls at a time. *W. Procter.*

PIPSISSEWA BEER.

R. Pipsissewa, half a pound.
Water, one gallon.

Boil, strain, and add

Brown sugar, one pound.

Powdered ginger, half an ounce.

Yeast, sufficient

to produce fermentation. When this has taken place, keep in well-stopped bottles. Dose, a half-tumblerful, three or four times a-day, in strumous affections, especially of the joints.

Dr. I. Parrish.

CHIRETTA.

CHIRETTA.

This is the herb and root of *Agathotes chirayta*, a native of the northern mountainous parts of India.

Sex. Syst. Pentand. digyn. *Nat. Syst.* Gentianaceæ.

Don, Phil. Mag., 1836. Griffith, Med. Bot. 462.

The whole plant is bitter, and the root most so; it is inodorous. It appears to have the same properties as gentian, and is applicable to the same cases. Dose, of powder, is twenty grains.

INFUSION OF CHIRETTA.

R. Chiretta, four drachms.
Boiling water, one pint.

Infuse for two hours, and strain. *Ed. Ph.*

Dose, one to two fl. ounces as a simple tonic bitter.

TINCTURE OF CHIRETTA.

R. Chiretta, five ounces.
Proof spirit, two pints.

Macerate for fourteen days, and filter. Dose, a teaspoonful. *Dub. Ph.*

CHLORINUM.

CHLORINE.

This is an elementary, gaseous fluid, of a greenish-yellow color, and a peculiar, penetrating smell. It has been used as an inhalation in a diluted state, in diseases of the lungs, and also as a fumigation in diseases of the liver, &c., and also as a disinfectant.

CHLORINE.

R. Peroxide of manganese, one part.
Common salt, three parts.

Sulphuric acid,

Water, each, two parts.

Mix the acid and water, pour on the other substances in powder, and collect the gas under a receiver. *Paris Cod.*

CHLORINE WATER.

- R. Muriate of soda,
(common salt), one drachm.
Sulphuric acid, two fl. drachms.
Red oxide of lead, three hundred and
fifty grains.
Water, eight fl. ounces.

Rub the salt and oxide of lead together, put them into the water, add the acid, and agitate occasionally till the red oxide becomes almost white. Allow the insoluble matter to subside before using the liquid. *Ed. Ph.*

To be kept in well-stopped bottles, and in the dark. Chlorine water is stimulant, and antiseptic, and has been used in a variety of diseases. (See *Dunlison's New Rem.*, 6th edit., p. 190, *et seq.*) The dose is from one to four fl. drachms, properly diluted.

CHLORINE COLLUTORY.

- R. Chlorine water, half an ounce.
Syrup of mallows, one ounce.
Mix. In aphthous sore mouth. *Radius.*
R. Chlorine water, half an ounce.
Infusion of sage, six ounces.
Honey of roses, one ounce.
Mix. In same cases. *Radius.*

CHLORINE GARGLE.

- R. Tragacanth, twelve grains.
Distilled water, four fl. ounces.
Chlorine water,
Syrup, each, half an ounce.
Mix. As a gargle in ulceration, and chronic inflammation of the mouth, and fauces. *Foy.*

CHLORINE INJECTION.

- R. Chlorine water, half a drachm.
Distilled water, one ounce and a half.
Extract of opium, fifteen grains.
Mix. As an injection in ulcers of the uterus. *Phæbus.*

CHLORINE CLYSTER.

- R. Chlorine water, two drachms.
Starch, four drachms.
Extract of opium, half a grain.
Water, six ounces.
Mix. As an enema in the diarrhœa of consumptive patients. *Cadet.*

CHLORINE MIXTURE.

- R. Chlorine water, two to four drachms.
Decoction of mallows, two ounces and a half.
Mix. A teaspoonful every hour, in softening of the stomach in children. *Radius.*

- R. Chlorine water, two ounces.
Syrup, six ounces.
Mix. A spoonful every two hours, in typhoid fevers. *Svediaur.*

- R. Chlorine water, two drachms.
Strawberry water, one ounce and a half.
Quince mucilage,
Syrup of mallows, each, six drachms.
Mix. A teaspoonful every two hours, to children in scarlatina. *Radius.*

CHLORINATED OIL.

- R. Olive oil, at will.
Pass a current of chlorine through it; at the end of two or three days, wash it with cold water. As an application to tinea, and lepra. *Deimann.*

CHLORINE OINTMENT.

- R. Chlorine water, one part.
Lard, eight parts.
Triturate well together. Employed in itch. *Augustin.*

CHLORINE LINIMENT.

- R. Chlorine water, one fl. drachm.
Olive oil, one fl. ounce.
Mix well. In itch, tinea, and herpes. *Deimann.*
R. White wax, two drachms.
Melt by a gentle heat, and add
Almond oil, sufficient
to make a liniment; on cooling, add
Chlorine water, one drachm and a half.
As an application to foul ulcers. *Ludwig.*

CHLOROFORMUM.

CHLOROFORM.

This has received various names, as *Chloride of Carbon*, *Chloride of Formyl*, *Chloric Ether*, *Terchloride of Formyl*, &c. It is a transparent, heavy fluid, of a peculiar, fragrant, ethereal, apple-like odor, and an intensely sweet taste. In large doses it is a narcotic poison; in medicine, a stimulant, sedative, antispasmodic anodyne, and anæsthetic; for which latter purpose it has attained much celebrity. The usual dose, when inhaled, is a fl. drachm, to be increased in a few minutes, if no effect is produced. The best plan of inhaling is to twist a handkerchief into the form of a bird's nest, wet this with the chloroform, and apply it to the mouth and nose. The dose, internally, is from five to twenty minims.

- R. Chlorinated lime, ten pounds.
 Water, three gallons and a half.
 Alcohol, two pints.

Mix the lime with the water and alcohol, distil into a cold receiver, and when the temperature reaches 176° F., remove the fire, and allow the distillation to proceed spontaneously; taking care to apply heat if the process slackens, so long as the liquid comes over of a sweet taste. Remove the heavier layer of liquid in the receiver, wash it with water, then with a weak solution of carbonate of soda, agitate it with powdered chloride of calcium, and by a water-bath distil off eleven-twelfths of the liquid.

U. S. Ph.

- R. Powdered chloride
 of lime, four pounds.
 Water, twelve pounds.
 Alcohol, twelve fl. ounces.

Mix, in a capacious still or retort, and distil as long as a dense liquid, which sinks in the water that comes over with it, is produced. Separate this, agitate it with sulphuric acid, and distil it from carbonate of baryta.

Dumas.

EMULSION OF CHLOROFORM.

- R. Chloroform, six drachms.
 Syrup, half an ounce.
 Yolk of egg, one.
 Distilled water, four fl. ounces and a half.

Mix, by rubbing the yolk of the egg with the syrup, then adding the chloric ether and water so as to form an emulsion.

Moxon.

LINIMENT OF CHLOROFORM.

- R. Soap liniment, two fl. ounces.
 Chloroform, one fl. drachm.
 Mix. As an application in neuralgic pains, over the affected part.

Tuson.

CHONDRUS.

IRISH MOSS.—CARRAGEEN.

The fronds of *Chondrus crispus*, a sea-weed growing on rocks and stones in the seas of Northern Europe, especially on the southern and western coasts of Ireland.

Sex. Syst. Cryptog. fuci. Nat. Syst. Ceramiales.

Greville, Alg. Brit. 129. Duglison, New Rem. 6th edit., p. 210.

It is of a yellowish, or purplish color, translucent, nearly tasteless, and scentless. It is nutritive and demulcent, and affords a useful article of diet to convalescents. It has also been recommended in scrofula, and bowel and pectoral affections. Before using, it should be soaked in cold water to remove any salt or other impurities.

DECOCTION OF CARRAGEEN.

- R. Carrageen, sliced, half an ounce.
 Water, three pints.
 Boil for a quarter of an hour, express, and strain.

Beral.

- R. Carrageen, sliced, half a drachm.
 New milk, nine fl. ounces.

Boil to five fl. ounces, strain, and add

- Sugar, half an ounce to an ounce.
 Bitter almond water, one scruple.

To be taken during the day.

Gräfe.

- R. Carrageen, four scruples.
 New milk, twenty-four fl. ounces.

Boil for ten minutes, express, and add

- Orange-flower water, one ounce and a half.

CARRAGEEN JELLY

- R. Carrageen, two drachms.
 New milk, one pint.

Boil to consistence of jelly, strain, and add

- Sugar, one ounce.
 Bitter almonds, two.

Radius.

- R. Mucilage of carrageen, five ounces.
 Sugar, four ounces.

Evaporate to eight ounces, by a gentle heat; strain, and add

- Essence of citron, fifty drops.

Beral.

CARRAGEEN MIXTURE.

- R. Carrageen, half a drachm.
 Water, sufficient

to obtain six ounces of mucilage; strain, and add

- Phosphate of soda, one drachm and a half.

- Syrup of opium, three drachms.

A spoonful, every two hours, in hemoptysis.

Clarus.

CICHORIUM.

SUCCORY.

The *Cichorium intybus* is a perennial herbaceous plant, indigenous to Europe, but naturalized in many places in this country.

Sex. Syst. Syngen. equal. Nat. Syst. Astéraceæ.

The whole plant is bitter, the root the most so. It is tonic, aperient, and deobstruent. The dried and torrefied root is used in Europe to mix with, or as a substitute for coffee.

EXTRACT OF SUCCORY.

R. Juice of succory, at will.
Evaporate to proper consistence. *Guibourt.*
Dose, from ten grains to half a drachm.

INFUSION OF SUCCORY.

R. Dried succory, half an ounce.
Boiling water, two pints.
Infuse and strain, and add
Syrup of maidenhair, one drachm.
Cottureau.

COMPOUND SYRUP OF SUCCORY.

R. Juice of succory,
" dandelion,
" fumitory, each, six ounces.
Sugar, one pound.
Boil to proper consistence, and strain. Dose,
one to two ounces. As a diuretic. *St. Marie.*

CIMICIFUGA.

BLACK SNAKEROOT.

This is the root of *Cimicifuga racemosa*, a native, found in many parts of the United States, in shady places.

Sex. Syst. Polyand. di-pent. *Nat. Syst.* Ranunculaceæ.

Torrey and Gray, Fl. 1. 36. Griffith, Med. Bot. 92.

The root is thick, contorted, dark brown, with numerous radicles. The odor is peculiar and unpleasant, and the taste bitter and somewhat astringent. It is acro-narcotic, and is used in rheumatism, dropsy, pectoral affections, chorea, &c. The dose of the powder, a drachm, several times a-day.

DECOCTION OF BLACK SNAKEROOT.

R. Black snakeroot (contused),
one ounce.
Water, one pint.
Boil for a short time, and strain. Dose, one to two fl. ounces. *Wood.*

TINCTURE OF BLACK SNAKEROOT.

R. Bruised black snakeroot, four ounces.
Alcohol, one pint.
Digest for fourteen days, and filter. Dose, twenty drops, three or four times a-day. *Carson.*
Used in rheumatic cases.

CINCHONA.

PERUVIAN BARK.

This is the bark of several species of *Cinchona*, all natives of South America. There

are numerous varieties of this article used in medicine, but the state of our knowledge on the subject does not permit a reference of them to a particular species of the tree, except in a few cases. At the revision of the U. S. Pharmacopœia, in 1850, the yellow bark was referred to *C. Calisaya*, and the pale bark to *C. Condaminea*, and *C. Micrantha*. The sources of the red bark remain undetermined. (See Griffith, *Med. Bot.* 368.) Peruvian bark owes its properties to the presence of one or more alkaloids, called quinia, cinchonina, &c.

It is a bitter tonic and antiperiodic, and is used in a variety of diseases, and is given in a multitude of forms. The dose in powder as a febrifuge is a drachm, frequently repeated; as a tonic from ten to thirty grains. Its use is now much superseded by that of quinia.

COMPOUND POWDER OF PERUVIAN BARK.

R. Powdered Peruvian bark,
Cremor tartar, each, one ounce.
Powdered cloves, one drachm.
Mix. A drachm and a half every two hours, in the apyrexia of fevers. *Ellis.*

R. Powdered Peruvian bark,
half an ounce.
Virginia snakeroot, one drachm.
Bicarbonate of soda, two scruples.

Mix, and divide into four powders, one to be given every two hours, in apyrexia of obstinate intermittents. *Ellis.*

R. Powdered Peruvian bark,
" uva ursi, each, one or two drachms.
" opium, three grains.

Mix, and divide into six powders; one to be taken three or four times a-day, followed by a draught of two ounces of lime water. Antilithic. *Ferriar.*

R. Powdered Peruvian bark, one ounce.
Tartar emetic, two grains.
Powdered opium, one grain.

Mix, and divide into eight powders; one to be taken every two hours. *Brera.*

R. Powdered Peruvian bark,
Sulphate of magnesia, each, six drachms.

Mix, and divide into four powders. One every two hours, where purging is required, and at the same time a tonic impression to be made. *Ellis.*

R. Powdered Peruvian bark,
thirty grains.
Aromatic powder, ten grains.

Make a powder, to be taken in a cup of milk every three hours. In convalescence from fevers. *A. T. Thomson*

POWDER OF PERUVIAN BARK AND
ARNICA.

- R. Powdered Peruvian bark, } equal
 " arnica root, } parts.
 " camphor, }
 Mix. As an application to gangrenous ulcers. *Cadet.*

POWDER OF PERUVIAN BARK AND
CASCARILLA.

- R. Powdered Peruvian bark, two drachms.
 " cascarilla, twelve grains.
 Mix. In apyrexia of intermittents. *Foy.*

POWDER OF PERUVIAN BARK AND
CAMPHOR.

- R. Powdered Peruvian bark, half an ounce.
 " camphor, one scruple.
 " gum Arabic, two drachms.
 Mix. As an application to atonic ulcers. *Gen. Ph.*

PERUVIAN BARK DENTIFRICE POW-
DERS.

- R. Powdered Peruvian bark, one ounce and a half.
 " red saunders, half an ounce.
 Oil of bergamot,
 Oil of cloves, each, twelve drops.
 Mix. *Hufeland.*
 R. Powdered Peruvian bark, three ounces.
 " cremor tartar,
 " sage leaves,
 " myrrh, each, half an ounce.
 " catechu, six drachms.
 Oil of cloves, sixteen drops.
 Mix. *Hesse Ph.*
 R. Powdered Peruvian bark,
 " charcoal, each, one drachm and a half.
 " cinnamon, half a drachm.
 Mix. *Soubiran.*
 R. Powdered Peruvian bark, one ounce.
 " orris root,
 " sage leaves,
 " myrrh, each, half an ounce.
 Mix. *Fulda Disp.*
 R. Powdered Peruvian bark, two ounces.
 " muriate of ammonia, half an ounce.
 " orris root, one ounce.

- Powdered catechu,
 " myrrh, each, six drachms.
 Oil of cloves, twelve drops.
 Mix. *Pideret.*

POWDER OF PERUVIAN BARK AND
VALERIAN.

- R. Powdered Peruvian bark, two drachms.
 " valerian, one drachm.
 Mix. Divide into four powders. In nervous disorders. *Ratier.*

POWDER OF PERUVIAN BARK AND
ISINGLASS.

- R. Powdered Peruvian bark, one ounce.
 Isinglass, one drachm and a half.
 Mix, and divide into sixteen powders. Dose, at first, from three to six a-day, in passive menorrhagia, leucorrhœa, chronic diarrhœa, &c. *Brera.*

PERUVIAN BARK MIXTURE.

- R. Powdered Peruvian bark, half an ounce.
 Confection of opium, one drachm.
 Lemon juice, two fl. drachms.
 Port wine, four fl. ounces.
 Mix. A wineglassful every two hours, in the apyrexia of intermittent fever. *Meigs.*
 R. Powdered Peruvian bark, two drachms.
 Wine of opium, sixty drops.
 Nitrate of potassa, five grains.
 Infusion of gentian, six fl. ounces.
 Mix. An ounce every two hours. *Pierquin.*
 R. Powdered Peruvian bark, two drachms.
 Compound tincture of bark, one fl. ounce.
 Decoction of red bark, three fl. ounces.
 Syrup, half fl. ounce.
 Mix. A tablespoonful every one or two hours in the apyrexia. *Ellis.*
 R. Powdered Peruvian bark, one ounce and a half.
 Magnesia, six drachms.
 Saffron, one scruple.
 Confection of opium, two drachms.
 White wine,
 Water, each, twelve fl. ounces.
 Mix. To be taken in divided doses, during the day, in apyrexia. *Giordano.*
 R. Powdered Peruvian bark, one ounce.
 Alcohol, eight ounces.
 Sulphuric ether, two drachms.
 Mix. A spoonful every hour, or two hours. *Augustin.*

R. Powdered Peruvian bark, one ounce.
 " ginger, twenty grains.
 Red wine, eight fl. ounces.
 Sugar, one ounce.

Mix. To be taken during the apyrexia.

Radius.

EXTRACT OF YELLOW PERUVIAN BARK.

R. Yellow bark, in coarse
 powder, one pound.
 Alcohol, four pints.
 Water, sufficient.

Macerate the bark in the alcohol for four days, and obtain tincture by displacement. When the liquid measures four pints, set aside and obtain six pints of infusion. Distil off alcohol from tincture, and evaporate infusion to the consistence of thin honey; mix, and evaporate to proper consistence. The extract of red bark is prepared in the same way. *U. S. Ph.*

Dose, ten to thirty grains.

VINOUS EXTRACT OF PERUVIAN BARK.

R. Powdered Peruvian bark, one part.
 White wine, eight parts.
 Digest for three days, express, strain, and evaporate to proper consistence. *Hesse Ph.*

SALT OF BARK.

R. Bruised yellow bark, two pounds.
 Distilled water, one gallon.
 Muriatic acid, half a fl. ounce.

Boil and strain. Then boil the residue twice in an equal quantity of acidulated water. Mix the decoctions, filter, and add lime, while precipitation occurs. Wash the precipitate, exhaust it with hot alcohol, and evaporate by a water-bath, to a pilular consistence.

Dose, two to five grains.

C. Ellis.

COMPOUND BOLUS OF EXTRACT OF PERUVIAN BARK.

R. Extract of Peruvian
 bark, twenty grains.
 Nitrate of potassa, thirty grains.
 Confection of orange peel, sufficient.
 Mix, and form five boluses. *Pierquin.*

COMPOUND PILLS OF EXTRACT OF PERUVIAN BARK.

R. Extract of Peruvian bark, one drachm.
 " opium, one grain.
 Camphor, twelve grains.
 Powdered Peruvian bark, sufficient.
 Beat into a mass, and form twelve pills. One to be taken three or four times a-day. *Ellis.*

R. Extract of Peruvian bark,
 " rhubarb,
 " gentian,
 " blessed thistle, equal parts.

Beat into mass, and form pills of four grains. Dose, one or two, three times a-day. *St. Marie.*

R. Extract of Peruvian bark,
 " gentian, each, one drachm.
 Sulphate of iron, half a drachm.
 Powdered myrrh, one drachm.
 Oil of caraway, ten drops.
 Syrup of ginger, sufficient.

Beat well together, and divide into forty pills. Three to be taken three times a-day. In the apyrexia of intermittents. *Thomson.*

R. Extract of Peruvian bark,
 two drachms.
 Powdered alum, one drachm.
 Syrup, sufficient.

Mix, and divide into thirty-six pills. Four to be taken every four or six hours. In passive hemorrhages. *A. T. Thomson.*

FLUID EXTRACT OF PERUVIAN BARK.

R. Coarsely-powdered
 yellow bark, eight ounces.
 Diluted alcohol, sufficient.

Macerate the bark in sufficient alcohol to cover it, for twelve hours; then by displacement obtain four pints of tincture. Evaporate the tincture on a water-bath to nine fl. ounces, then add fourteen ounces of sugar. Continue the heat till dissolved, and strain. Dose, a teaspoonful. *A. B. Taylor.*

DECOCTION OF YELLOW PERUVIAN BARK.

R. Yellow bark, bruised, one ounce.
 Water, one pint.
 Boil for ten minutes, and strain while hot. Dose, two fl. ounces. *U. S. Ph.*

The decoction of red bark is prepared in the same manner.

R. Red Peruvian bark, bruised, one ounce.
 Water, one pint.
 Boil for ten minutes, and, while hot, add
 Virginia snakeroot, half an ounce.
 Orange peel, two drachms.

Infuse for half an hour, near the fire, in a covered vessel. Dose, a wineglassful every hour. *Ellis.*

R. Decoction of Red Peruvian
 bark, six fl. ounces.
 Compound tincture of
 bark, one fl. ounce.
 Aromatic sulphuric acid, ten drops
 Mix. A tablespoonful every hour or two. *Ellis.*

DECOCTION OF PERUVIAN BARK MIXTURE.

R. Decoction of Peruvian bark, six ounces.
Tincture of Peruvian bark, one ounce.
Diluted sulphuric acid, one drachm.
Syrup of orange peel, half an ounce.

Mix. A tablespoonful every hour or two. *Foy.*

R. Decoction of Peruvian bark, eight ounces.
Infusion of arnica flowers, six ounces.
Camphor, two scruples.
Syrup of tolu, one ounce.
Emulsion of gum, four ounces.

Mix. To be taken in divided doses, during the day. *Brugnatelli.*

DECOCTION OF PERUVIAN BARK AND CASCARILLA.

R. Peruvian bark, one ounce.
Cascarilla, two drachms.
Water, twelve fl. ounces.

Boil to eight ounces, express, and strain, and add, when cold,

Sulphuric ether, two drachms.
Mix. *Copenhagen Ph.*

DECOCTION OF PERUVIAN BARK AND RHATANY.

R. Peruvian bark,
Rhatany, each, half an ounce.
Water, sufficient

to obtain nine ounces of decoction; strain, and add

Vinous extract of Peruvian bark, four scruples.
Compound tincture of Peruvian bark, one fl. drachm.
Citron water, one fl. ounce.

Mix. In chronic diarrhœa. *Brera.*

COMPOUND DECOCTION OF PERUVIAN BARK.

R. Peruvian bark, one ounce.
Virginia snakeroot, half an ounce.
Orange peel, two drachms.
Powdered cloves, one drachm.
Carbonate of potassa, one drachm.

Mix. Put the mixture into a proper vessel, with three half pints of water, boil down gently, to one pint; let settle, decant. Dose, a wineglassful every hour, for eight hours, in the apyrexia. *Dewees.*

INFUSION OF YELLOW PERUVIAN BARK.

R. Yellow Peruvian bark, bruised, one ounce.
Boiling water, one pint.

Infuse for two hours, in a covered vessel, and strain. *U. S. Ph.*

Dose, two fl. ounces, as occasion may require. The infusion of red bark is prepared in the same way.

INFUSION OF PERUVIAN BARK WITH LIME WATER.

R. Bruised Peruvian bark, two ounces.
Lime water, two pints.

Infuse, and strain. A wineglassful to be given three or four times a-day. *Ellis.*

INFUSION OF PERUVIAN BARK AND VALERIAN

R. Bruised Peruvian bark, one ounce and a half.
Bruised valerian, half an ounce.
Boiling water, two pints.

Infuse for twelve hours, and strain. *Fulda Disp.*

INFUSION OF PERUVIAN BARK AND SERPENTARIA.

R. Bruised Peruvian bark, ten drachms.
Boiling water, one pint.

Infuse for four hours, then boil for half an hour, then add

Virginia snakeroot, two drachms.
Infuse for two days, and strain. *Saunders.*

INFUSION OF PERUVIAN BARK AND MAGNESIA.

R. Powdered Peruvian bark, one ounce.
Magnesia, one drachm.
Boiling water, twelve fl. ounces.

Infuse for twelve hours, express, and strain. Said to be more astringent than other infusions. *Van Mons.*

COMPOUND INFUSION OF PERUVIAN BARK.

R. Peruvian bark,
Juniper berries, each, one drachm.
Orange peel,
Cinnamon, each, two drachms.
Canella, half a drachm.
Ipecacuanha, fifteen grains.
Boiling water, seven fl. ounces.

Infuse, and strain, then add

Extract of juniper berries, one drachm and a half.

In spoonful doses, in atonic dropsies. *Beries.*

R. Root of viper's grass, one ounce.
Contused lemon seeds, two drachms.
Water, six pints.

Boil to four pints; pour boiling liquid on
Bruised Peruvian bark, two ounces.
Contrayerva, one ounce.
Infuse for a quarter of an hour, strain, and add
Honey of elder, three ounces.
As a stimulating diuretic. *Span. Ph.*

COMPOUND INFUSION OF PERUVIAN BARK.

R. Powdered red bark, one ounce.
Aromatic sulphuric acid, one fl. drachm.
Water, one pint.
Macerate for twelve hours, and strain. *U. S. Ph.*
Dose, two fl. ounces

ELECTUARY OF PERUVIAN BARK.

R. Powdered Peruvian bark, one ounce.
Syrup of orange peel, sufficient.
A teaspoonful every hour. *Radiis.*

ASTRINGENT ELECTUARY OF PERUVIAN BARK.

R. Powdered Peruvian bark,
" orange peel, each, six drachms.
" crabs' eyes, two drachms.
Confection of red roses,
" rose hips, each, six drachms.
Syrup of catechu, sufficient.
Mix. Dose, one drachm, two or three times a-day, in chronic diarrhoea. *Saunders.*

ELECTUARY OF PERUVIAN BARK AND CATECHU.

R. Powdered Peruvian bark, one ounce.
" catechu, one drachm.
Balsam of tolu, each, one drachm.
Syrup of comfrey, sufficient.
Divide into ten doses, one to be taken twice or three times a-day, in some appropriate vehicle.
Much praised in hemoptysis. *Pierquin.*

ELECTUARY OF PERUVIAN BARK AND TIN.

R. Powdered Peruvian bark, one ounce.
" tin, half an ounce.
" valerian, each, sufficient.
Syrup, sufficient.
Mix. Dose, a drachm, morning and evening, in epilepsy. *Cadet.*

ELECTUARY OF PERUVIAN BARK AND CLOVES.

R. Powdered Peruvian bark, two ounces.
Cloves, one drachm.
Simple syrup, sufficient
to form electuary. A piece, the size of a nutmeg, every two hours in, the apyrexia of intermittent fever. *Dewees.*

ELECTUARY OF PERUVIAN BARK AND IRON.

R. Powdered Peruvian bark, six drachms.
Protoxide of iron,
Confection of opium, each, two drachms.
Syrup of cinnamon, sufficient.
Mix. Dose, a drachm, morning and evening, in ascites, after the evacuation of the fluid. *Cadet.*

ELECTUARY OF PERUVIAN BARK AND SULPHUR.

R. Powdered Peruvian bark, one drachm.
" sulphur, two drachms.
" crabs' eyes, four grains.
" spermaceti, each, one drachm.
Extract of opium, and a half.
Powdered mushrooms, four drachms.
Confection of roses, sufficient.
Syrup of milfoil, sufficient.
Mix. Dose, two scruples, three times a-day. Highly praised in phthisis. *Cadet.*

TINCTURE OF PERUVIAN BARK.

R. Powdered yellow bark, six ounces.
Diluted alcohol, two pints.
Macerate for fourteen days, express, and filter. *U. S. Ph.*

Generally employed as an adjuvant to other preparations of bark, in the dose of one to four fluidrachms.

COMPOUND TINCTURE OF PERUVIAN BARK.

(HUXHAM'S TINCTURE.)

R. Powdered red bark, two ounces.
Bruised orange peel, one ounce and a half.
" Virginia snakeroot, three drachms.
Saffron, cut, one drachm.
Red saunders, rasped, each, twenty fl. ounces.
Diluted alcohol, sufficient.
Macerate for fourteen days, express, and filter. *U. S. Ph.*

Mix. As a friction, three times a-day, in weakness of the back. *Augustin.*

PERUVIAN BARK BEER.

R. Bruised Peruvian bark, one and a half ounces.
Cinnamon, two drachms.
Grated nutmeg, seven drachms.
Yeast, two ounces.
Sugar, twenty-five ounces.
Water, one hundred fl. ounces.

Mix, and expose to a warm temperature. After fermentation, decant and strain. Three or four glasses to be taken during the apyrexia of intermittents. *Ferrara Ph.*

COMPOUND SYRUP OF PERUVIAN BARK.

R. Powdered Peruvian bark, two drachms.
Powdered seneka, half a drachm.
" liquorice, one drachm.
Contused poppy heads, three.
" ipecacuanha, one scruple.
Ground ivy, } each, a handful.
Hyssop, }
Pennyroyal, }
Sugar, one pound.
Water, sufficient.

Boil, and make a syrup. Three spoonfuls a-day, in hooping-cough. *Bories.*

WINE OF PERUVIAN BARK.

R. Powdered Peruvian bark, one part.
White wine, twelve parts.
Digest for twenty-four hours, often stirring, and filter. *Cottreaux.*
R. Red wine, twenty parts.
Extract of Peruvian bark, one part.
Mix. *Niemann.*

WINE OF PERUVIAN BARK AND IRON.

R. Bruised Peruvian bark, one ounce and a half.
Oxide of iron, half an ounce.
Powdered cinnamon, two drachms.
Sugar, one ounce.
White wine, two pints.
Mix, and digest for two days, and filter. Dose, an ounce, two or three times a-day, as a tonic.

WINE OF PERUVIAN BARK AND CALAMUS.

R. Bruised Peruvian bark, Calamus, each, one ounce and a half.
Quassia, } each, six drachms.
Cinnamon, }
Elder flowers, }

Alcohol, three pints.
Water, eighteen pints.

Macerate, express, strain, and add

Tincture of chloride of iron, twelve fl. ounces.
Orange-flower water, one pint and a half.
Syrup, six ounces

Mix. Dose, two drachms, morning and evening, in leucorrhœa and gonorrhœa. *Cadet*

PERUVIAN BARK MIXTURE.

R. Peruvian bark, six drachms.
Confection of opium, three drachms.
Cremor tartar, two drachms.
Water, six fl. ounces.

Mix. A tablespoonful every hour, in the apyrexia. *Devees.*

SYRUP OF PERUVIAN BARK.

R. Extract of bark, fourteen drachms.
White wine, two pounds.
White sugar, three pounds.

Dissolve the extract in the wine, filter, and form a syrup with the sugar, with the aid of a very moderate heat. *Monchon.*

CINCHONIA.

CINCHONIA.

An alkaloid existing in Peruvian bark; most abundant in the pale kinds. It is white, translucent, and crystallizable, but little soluble in water; very bitter, and has much the same properties as quinia.

CINCHONIA.

R. Pale bark, one thousand parts.
Muriatic acid, sixty-four parts.
Water, twelve thousand parts.
Quicklime, one hundred parts.

Exhaust the bark by three successive boilings with a third part of the acid and water, each time; unite the decoctions, add the lime diffused in water, wash, and dry the precipitate; treat it with alcohol, filter whilst hot, distil in a water-bath, evaporate to dryness, digest in cold alcohol, dissolve the residue in boiling alcohol, with the addition of some animal charcoal, filter whilst hot, and crystallize. *Par. Cod*

PILLS OF CINCHONIA.

R. Cinchonia, one scruple.
Confection of roses, sufficient.
Mix, and form twenty-four pills. Dose, two to eight. *Cadet.*

MURIATE OF CINCHONIA.

R. Cinchonia, at will.
 Muriatic acid, sufficient
 to saturate. Filter, evaporate, and crystallize.
Giordano.

KINATE OF CINCHONIA.

R. Alcoholic solution of sulphate
 of cinchonia, at will.
 Aqueous solution of kinate
 of lime, sufficient
 to produce full precipitation; filter, evaporate, re-
 dissolve, and crystallize. *Magendie.*

SULPHATE OF CINCHONIA.

R. Cinchonia, at will.
 Diluted sulphuric acid, sufficient
 to dissolve the cinchonia. Evaporate, and crys-
 tallize. *Guibourt.*

PILLS OF SULPHATE OF CINCHONIA.

R. Sulphate of cinchonia, half a drachm.
 Confection of roses, sufficient.
 Mix, and make thirty pills. Dose, one to eight.
Cadet.

BOLUS OF CINCHONIA.

R. Sulphate of cinchonia, three grains.
 Crumb of bread,
 Honey,
 Liquorice, each, sufficient.
 Mix, and form a bolus. One every two hours.
Brera.

SYRUP OF CINCHONIA.

R. Sulphate of cin-
 chonia, forty-eight grains.
 Syrup, one pint.
 Mix. A tablespoonful is a dose. *Cadet.*
 R. Sulphate of cin-
 chonia, twenty-four grains.
 Orange-flower water, two fl. drachms.
 Syrup, twelve fl. ounces.
 Mix. As above. *Giordano.*

WINE OF CINCHONIA.

R. Sulphate of cin-
 chonia, twenty-four grains.
 Madeira wine, two pints.
 Make solution, and filter. Dose, one to four
 ounces. *Cottureau.*

TINCTURE OF CINCHONIA.

R. Sulphate of cinchonia, twelve grains.
 Alcohol, one ounce.
 Dissolve. Dose, a teaspoonful. *Foy.*

MIXTURE OF SULPHATE OF CINCHONIA.

R. Sulphate of cinchonia, three grains.
 Peppermint water, three ounces.
 Syrup of capillaire, one ounce.
 Mix. To be taken in four doses, during the
 day. *Cadet.*

CINNAMOMUM.

CINNAMON.

This is the bark of *Cinnamomum Zeylanicum*,
 and *C. aromaticum*, and also, perhaps, of other
 species; all natives of various parts of the tro-
 pical regions of Asia. That most commonly
 used in this country, is what is termed cassia
 in commerce, and is imported from China.

Sex. Syst. Enneand. monog. *Nat. Syst.* Lau-
 raceæ.

Nees, Laurineæ. 52. Griffith, Med. Bot. 555.
 It is in tubular pieces, of a light brownish-red
 color, of an aromatic, warm, agreeable taste,
 and of a fragrant, peculiar smell. It is a warm
 cordial stimulant. Dose, in powder, ten grains
 to a scruple.

COMPOUND POWDER OF CINNAMON.

R. Powdered cinnamon,
 " ginger, each, two ounces.
 Cardamom seeds,
 Grated nutmeg, each, one ounce.
 Rub together. *U. S. Ph.*

Stimulant, and carminative. Dose, ten to
 thirty grains, in debility of the stomach, with
 flatulence.

R. Powdered cinnamon, one ounce.
 " cloves, } each, three
 " ginger, } drachms.
 " nutmeg, }
 " mace, two drachms.
 " red saunders, half an ounce.
 " sugar, thirty-two ounces.

Mix well. Dose, from ten grains to a drachm.
Wurtemberg Ph.

R. Powdered cinnamon, twenty grains.
 " cloves, twelve grains.
 " vanilla, six grains.
 Rice flour, six drachms.
 Sugar, half an ounce.

Mix. Dose, six grains to a scruple. *Spielmann.*

CINNAMON LOZENGES.

R. Powdered sugar,
 Cinnamon water, each, six ounces.
 Boil to syrup, and add
 Powdered sugar, one pound.
 Oil of cinnamon, one drachm and a
 half.

Mix, make lozenges, and dry. *Taddei.*

COMPOUND ELECTUARY OF CINNAMON.

R. Confection of orange peel, three ounces.
 " nutmeg, one ounce and a half.
 " ginger, six drachms.
 Powdered cinnamon, half an ounce.
 Syrup of orange peel, sufficient.
 Mix. *Swediaur.*

INFUSION OF CINNAMON.

R. Bruised cinnamon, half an ounce.
 Boiling water, two pints.
 Infuse. In dyspepsia and nervous colics.
Radius.

COMPOUND VINEGAR OF CINNAMON.

R. Cinnamon, } each, one drachm.
 Cloves, }
 Nutmeg, }
 Peppermint, half an ounce.
 Lavender,
 Rosemary, each, one ounce.
 Vinegar, eight fl. ounces.
 Diluted alcohol, twenty fl. ounces.
 Mix, and distil twenty-eight ounces. Used as an aromatic. *Van Mons.*

COMPOUND WINE OF CINNAMON.

R. Powdered cinnamon, } each, half a drachm.
 Cloves, }
 Mace, }
 Cardamom, }
 White wine, two pints.
 Boil, filter, and add
 Sugar, ten ounces.
 As a cordial stomachic and stimulant. *Augustin.*

TINCTURE OF CINNAMON.

R. Bruised cinnamon, three ounces.
 Diluted alcohol, two pints.
 Macerate for fourteen days, express, and filter.
U. S. Ph.
 Dose, one to four fl. drachms.

COMPOUND TINCTURE OF CINNAMON.

R. Bruised cinnamon, one ounce.
 " cardamom seeds, half an ounce.
 " ginger, three drachms.
 Diluted alcohol, two pints.
 Macerate for fourteen days, express, and filter.
U. S. Ph.
 A warm, aromatic tincture, useful in flatulence, gastrodynia, &c. Dose, one to two fl. drachms.

R. Bruised cinnamon, one ounce.
 " cloves, } each,
 " nutmeg, } half an ounce.
 Cut saffron,
 Bruised calamus, one drachm.
 " mace, two drachms.
 Fresh lemon peel,
 " orange peel, each, one drachm.
 Diluted alcohol, one pint and a half.
 Digest, express, and filter. Dose, fifty to eighty drops, as a stomachic. *Wirttemberg Ph.*

TINCTURE OF CINNAMON GARGLE.

R. Tincture of cinnamon, one part.
 Syrup of currants, two parts.
 Peppermint water, sixteen parts.
 Mix. In malignant sore throat. *Foy.*

ETHEREAL TINCTURE OF CINNAMON.

R. Bruised cinnamon, three drachms.
 " cardamom, half an ounce.
 " angelica, one drachm and a half.
 " long pepper, one drachm.
 Sulphuric ether, five fl. ounces.
 Diluted alcohol, ten fl. ounces.
 Macerate for eight days, express, and filter.
 Stomachic and carminative. Dose, thirty to fifty drops. *Niemann.*

SPIRIT OF CINNAMON.

R. Oil of cinnamon, two fl. drachms.
 Proof spirit, one gallon.
 Dissolve.
 Dose, one fl. drachm. *Lond. Ph.*

ESSENCE OF CINNAMON.

R. Oil of cinnamon, one fl. ounce.
 Rectified spirit, nine fl. ounces.
 Mix with agitation. *Dub. Ph.*
 Dose, ten drops.

WATER OF CINNAMON.

R. Oil of cinnamon, half fl. drachm.
 Carbonate of magnesia, one drachm.
 Distilled water, two pints.
 Rub the oil with the magnesia, and both with the water, and filter. *U. S. Ph.*
 Mostly employed as a vehicle or adjuvant to other medicines.

COMPOUND CINNAMON WATER.

R. Sage, } each, two parts.
 Rosemary, }
 Peppermint, }
 Lavender, }

Fennel,
Cinnamon, each, one part.
Water, one hundred and twenty parts.
Mix, and distil seventy-two parts. In spoonful doses. *Saxon Ph.*

OIL OF CINNAMON.

R. Bruised cinnamon, at will.
Water, sufficient.
Distil, and separate the oil. *Par. Cod.*

CINNAMON SPICE PLASTER.

R. Olibanum, three ounces.
Yellow wax, half an ounce.
Powdered cinnamon, six drachms.
Oil of pimento,
" lemons, each, two drachms.

Melt the wax and olibanum, and add the cinnamon, rubbed with the oils. *Coze.*

COCCULUS.

COCCULUS INDICUS.

This is the fruit of *Anamirta cocculus*, a woody vine, native of the East Indies.

Sex. Syst. Diœc. dodecand. *Nat. Syst.* Menispermaceæ.

Wight and Arnott, 1. 446. Griffith, Med. Bot. 105.

The berries are about the size of a pea, roundish, with a thin, dry, blackish, external coat, enclosing another of a lighter color, containing an oily, bitter, white kernel. They are inodorous, but of a permanent, bitter taste. They are acro-narcotic, and owe their properties to the presence of a peculiar principle, called *picrotoxin*. They are seldom used in medicine.

OINTMENT OF COCCULUS INDICUS.

R. Cocculus indicus, at will.
Separate the kernels, beat them in a mortar; first alone, then with a little lard; finally, add lard equal to five times the weight of kernels. *Ed. Ph.*

Used for the destruction of vermin, and in the treatment of ringworms and scabies of the scalp.

OINTMENT OF PICTROTOXIN.

R. Picrotoxin, six grains.
Lard, one ounce.
Mix well. In tinea. *Radius.*

COCCUS.

COCHINEAL.

Cochineal is a small, hemipterous insect, inhabiting Mexico and some parts of South America, living principally on different species

of cactus, and especially the *Opuntia cochinillefera*. It is in grains of a reddish-black color, often covered with a white powder. It has a feeble odor, and a bitterish, somewhat acidulous taste. The powder is of a reddish-purple color, tinging fluids of a deep red. It is said to be somewhat anodyne and antispasmodic, but is principally used as a coloring agent.

TINCTURE OF COCHINEAL.

R. Powdered cochineal, one part.
Diluted alcohol, eight parts.
Macerate eight days, express, and filter. Advised in convulsive coughs. Dose, twenty drops to a fl. drachm. *Beral.*

MIXTURE OF COCHINEAL AND CARB. POTASSA.

R. Carbonate of potassa, one scruple.
Powdered cochineal, half a scruple.
Sugar, one drachm.
Distilled water, four fl. ounces.
Make mixture. Dose, for children, a teaspoonful, every two or three hours. A highly popular remedy in whooping-cough. *Ellis.*

MIXTURE OF COCHINEAL AND COMMON SALT.

R. Common salt, three drachms.
Powdered cochineal, fifteen grains.
Divide into six doses. One to be taken every morning, the last followed by a purge. As a vermifuge. *Rush.*

COCHLEARIA.

SCURVY GRASS.

Scurvy grass is the *Cochlearia officinalis*, an herbaceous plant, native of the northern parts of Europe, and cultivated there and in the United States as a garden herb.

Sex. Syst. Tetradyn. silic. *Nat. Syst.* Brassicaceæ.

Griffith, Med. Bot. 131.

The whole herb is used; it has a penetrating and acrid smell when bruised, and a pungent, bitter taste. It is antiscorbutic, aperient, and diuretic.

ELECTUARY OF SCURVY GRASS.

R. Conserve of scurvy
grass, one ounce and a half.
Calamus, one drachm.
Syrup of orange peel,
Aromatic sulphuric acid,
each, sufficient.
Mix. A teaspoonful every two hours, as an antiscorbutic. *Augustin.*

GARGLE OF SCURVY GRASS.

- R. Spirit of scurvy grass, one drachm.
 Syrup of Peruvian bark, half an ounce.
 Infusion of water cress, eight ounces.
 Mix. As a gargle in scurvy. *Foy.*

CODEIA.

CODEIA.

An alkaloid obtained from opium, and bearing the same relation to morphia that cinchonia does to quinia. It is precipitated by tannin, and is soluble in ether, but is not thrown down from a dilute solution of its salts by ammonia. It acts somewhat like morphia, but produces great itching of the skin.

CODEIA.

- R. Opium, at will.
 Macerate in a sufficient quantity of water, decant, treat with chloride of calcium, filter, evaporate, and crystallize; decompose by ammonia, filter, concentrate fluid, crystallize; dissolve the crystals in water, add a slight excess of caustic potassa, dissolve the precipitate in alcohol and ether, and crystallize. Dose, one to two grains, in neuralgia, &c. *Cottureau.*

- R. Mother water of morphia, (by Gregory's process), at will.
 Evaporate and crystallize; dissolve the crystals, and re-crystallize. Treat with solution of potassa, which dissolves the morphia and precipitates the codeia; wash the latter with a little water, dissolve in hot ether, and suffer the solution to evaporate spontaneously.

SYRUP OF CODEIA.

- R. Codeia, twenty-four grains.
 Distilled water, four fl. ounces.
 Sugar, eight ounces.
 Reduce codeia to fine powder, triturate with one-third of water allow to settle and decant; treat residue with the remainder of water, at two operations. Treat in a matrass on water-bath till quite dissolved, add sugar, and dissolve. *Cap.*

Dose, one drachm, night and morning; gradually increased in neuralgic irritation of the stomach, hooping-cough, &c

MURIATE OF MORPHIA AND CODEIA.

- R. Opium, at will.
 Treat with water, add a strong solution of chloride of calcium, filter with animal charcoal, evaporate, and crystallize. *Magendie.*
 Dose, half a grain to a grain.

COFFEA.

COFFEE.

Coffee is the product, principally, of *Coffea Arabica*, a small tree, native of the eastern coast of Africa, but now cultivated in most tropical regions. The varieties are very numerous, and in some cases probably are derived from other species.

Sex. Syst. Pentand. monog. *Nat. Syst.* Cinchonaceæ.

Linn. Sp. Pl. 245. Griffith, Med. Bot. 361.

Raw coffee has febrifuge properties, and has been used with success in the treatment of intermittents. In a roasted state it acts on the nervous system, producing wakefulness, and also somewhat excites the circulatory system. It is said to be useful in periodic asthma, and is an efficient anti-emetic. Its active principle, *caffein*, is crystallizable.

DECOCTION OF RAW COFFEE.

- R. Raw coffee, one ounce.
 Water, eighteen fl. ounces.
 Boil to two-thirds, strain, and add
 Lemon juice, two fl. ounces.
 To be taken warm, during the apyrexia. *Foy.*

VINEGAR OF COFFEE.

- R. Roasted coffee, ground, three ounces.
 Vinegar, twelve ounces.
 Boil, strain, and add
 Sugar; one ounce and a half.
 Two spoonfuls occasionally, in poisoning by opium, after the evacuation of the poison. *Pierquin.*

SYRUP OF COFFEE.

- R. Coffee, roasted, four ounces.
 Water, two pints.
 Refined sugar, three pounds.
 Infuse the coffee in a pint and a half of the water for six hours, boil in a water-bath, let settle, decant, and add remainder of the water. Let stand for some hours; decant, add sugar, form syrup, and strain. Dose, from one ounce to two. *Ferrari.*

CITRATE OF CAFFEIN.

- R. Caffein, at will
 Solution of citric acid, sufficient
 to saturate. Dissolve at a temperature of 112° F., evaporate, and crystallize.
 This salt is very soluble. Dose, one to three grains.

PILLS OF CITRATE OF CAFFEIN.

- R. Citrate of caffein, eight grains.
 Dog grass, (*Triticum repens*), fifteen grains.

Mix, and divide into ten pills. Dose, one pill every two hours in sick headache. *Hannon.*

SYRUP OF CITRATE OF CAFFEIN.

R. Citrate of caffein,
two and a half drachms.
Simple syrup, four fl. ounces.

Dissolve.

Dose, a teaspoonful every two hours, in sick headache. *Hannon.*

COLCHICUM.

COLCHICUM.

The *Colchicum autumnale*, or meadow saffron, is a bulbous plant, a native of most parts of Europe, flowering in the autumn, and putting forth its leaves, and perfecting its seed the succeeding spring.

Sex. Syst. Hexand. monog. *Nat. Syst.* Melanthaceæ.

Linn. Sp. Pl. 485. Griffith, Med. Bot. 644.

The parts used are the bulb and the seeds. The first is of a brown color externally, white within, rounded on one side, flattened on the other, of scarcely any smell, but an acrid, bitter taste; the seeds are small, rounded, of a brownish-yellow color, and of the same taste as the bulb. They owe their properties to the presence of colchicia.

Colchicum is principally used in the various forms of gout and rheumatism. It is an active irritant, and in large doses, acro-narcotic. The dose of the powdered bulb is two to eight grains, every four or six hours; that of the seeds much the same.

EXTRACT OF COLCHICUM BULB.

R. Fresh colchicum bulb, one pound.

Bruise in a stone mortar, express juice, and evaporate to proper consistence. *Lond. Ph.*

Dose, one or two grains.

ACETIC EXTRACT OF COLCHICUM BULB.

R. Fresh colchicum bulb, one pound.

Acetic acid, three fl. ounces.

Bruise, gradually adding acid, express, and evaporate, without straining, in porcelain vessel to proper consistence. *Lond. Ph.*

R. Coarsely powdered colchicum

root, (dried,) one pound.

Acetic acid, four fl. ounces.

Water, a sufficient quantity.

Mix the acid with a pint of water, and pour on the root. Transfer to a percolator, and add water until the liquid passes tasteless. Then evaporate to proper consistence. *U. S. Ph.*

Dose, one to two grains, two or three times a-day.

COLCHICUM PILLS.

R. Powdered colchicum, three grains.
Soap, sufficient.

Make three pills, to be taken during the day, gradually increasing to five or six. *Ritton.*

COMPOUND COLCHICUM PILLS.

R. Acetic extract of colchicum, three grains.

Dover's powder,

Compound extract of colocynth, each, one grain.

Mix, and make a pill. In gout, after having taken, for some days, thirty to forty drops of colchicum wine, morning and evening. *Halford.*

LARTIGUE'S PILLS.

R. Extract of colchicum root, two grains.

" foxglove, one grain.

" col. comp., twenty grains.

Mix, and form five pills. One at night, in gout. *Lartigues.*

WINE OF COLCHICUM ROOT.

R. Colchicum root, bruised, one pound.

Sherry wine, two pints.

Macerate for fourteen days, express, and filter. *U. S. Ph.*

WINE OF COLCHICUM SEED.

R. Colchicum seed, bruised, four ounces.

Sherry wine, two pints.

Macerate for fourteen days, express, and filter. *U. S. Ph.*

The dose of the first, which is a saturated wine of the root, is from ten drops to half a fl. drachm; for the second, one to two fl. drachms.

R. Colchicum seed, bruised, three ounces.

Sherry wine, deprived of its

spirit by evaporation, two pints.

Macerate for six days, and filter. *Taddei.*

This is said to be the formula of the *Eau medicinale of Husson.*

TINCTURE OF COLCHICUM SEED.

R. Colchicum seed, bruised, four ounces.

Diluted alcohol, two pints.

Macerate for fourteen days, express, and filter. *U. S. Ph.*

Dose, from half a fl. drachm to two fl. drachms. Sometimes used as an embrocation in gout, rheumatism, and neuralgia.

ETHEREAL TINCTURE OF COLCHICUM SEED.

R. Colchicum seed, bruised, four ounces.

Spirit of nitric ether, two pints.

Digest for ten days, and filter. The advantage

claimed for this preparation is a greater tendency to act on the kidneys.

Dose, twenty to thirty drops. *Mettauer.*

COMPOUND TINCTURE OF COLCHICUM SEED.

R. Colchicum seed, bruised, five ounces.
Aromatic spirit of ammonia, two pints.
Macerate for seven days, express, and filter.
Lond. Ph.

More stimulating than the last. Dose, thirty drops to a fl. drachm.

TINCTURE OF COLCHICUM FLOWERS.

R. Dried colchicum flowers, one ounce.
Diluted alcohol, one pint.
Macerate for eight days, express, and filter.
Advised in acute rheumatism. *Bushell.*

VINEGAR OF COLCHICUM ROOT.

R. Dried colchicum root, two ounces.
Diluted acetic acid, two pints.

Macerate the root in the acid for seven days, express, and let dregs subside; pour off and add alcohol. *U. S. Ph.*

As a diuretic in dropsy; also given in gout, rheumatism, &c. Dose, thirty drops to two fl. drachms.

OXYMEL OF COLCHICUM.

R. Fresh colchicum root, sliced, one ounce.
Distilled vinegar, one pint.
Clarified honey, two pounds.

Macerate the colchicum in the vinegar for two days, express, and strain; add honey, and boil to consistence of syrup. *Dub. Ph. 1826.*

Dose, one fl. drachm twice a-day.

VINEGAR OF COLCHICUM SEED.

R. Colchicum seed, one part.
Distilled vinegar, four parts.
Macerate a month, and filter. *Beral.*

SYRUP OF COLCHICUM.

R. Vinegar of colchicum root, sixteen parts.
Sugar, twenty-six parts.
Boil, by a gentle fire. *Swediaur.*

MIXTURE OF COLCHICUM AND MAGNESIA.

R. Magnesia, one drachm.
Sugar, sufficient.
Gum Arabic, each, sufficient.
Distilled water, four fl. ounces.
Wine of colchicum root, forty drops.

Mix. A tablespoonful every two hours, till it operates. In gout and rheumatism. *Ellis.*

MIXTURE OF COLCHICUM AND SULPHATE OF MAGNESIA.

(SCUDAMORE'S MIXTURE.)

R. Sulphate of magnesia, one to two ounces.
Mint water, ten fl. ounces.
Vinegar of colchicum,
Syrup of saffron, each, one fl. ounce.
Magnesia, eight scruples.

Mix. Dose, one to three tablespoonfuls, every two hours, till four to six evacuations are procured in the twenty-four hours. In gout. *Scudamore.*

COMPOUND WINE OF COLCHICUM.

R. Tincture of colchicum seed, one scruple to half a fl. drachm.
Wine of colchicum seed, half a fl. ounce.

Mix. Fifteen to forty drops, three times a-day. *Weber.*

TINCTURE OF COLCHICUM MIXTURE.

R. Tincture of colchicum seed,
" guaiacum, each, three fl. drachms.

Mix. Thirty to forty drops, three times a-day, in chronic rheumatism. *Radius*

TINCTURE OF COLCHICUM AND DIGITALIS.

R. Tincture of colchicum seed,
" digitalis, each, two fl. drachms.

Nitric ether, one scruple.

Mix. Twenty drops, morning and evening, on sugar, in hydrothorax. *Hildebrand.*

COLCHICUM MIXTURE.

R. Wine of colchicum seed, thirty drops.
Denarcotized laudanum, twenty-five drops.
Sugar, thirty grains.
Water, one fl. ounce.

Mix. Found useful in acute rheumatism. To be taken at night. *Dewees.*

COLCHICUM MIXTURE.

R. Wine of colchicum seed, forty drops.
Acetated tincture of opium, twenty drops.
Sugar, thirty grains.
Water, one fl. ounce.

Mix. In gout, after inflammation is allayed. To be taken at bedtime. *Dewees*

R. Expressed juice of colchicum flowers, two parts.
Brandy, one part.

Mix, and let rest for a few days: decant, and keep for use in well-stopped bottles. *Wilson.*

Said to be identical in every respect to the *Eau medicinale de Husson.*

MIXTURE OF COLCHICUM AND ELATERIUM.

R. Elaterium, one grain.
Spirit of nitric ether, two fl. ounces.
Tincture of squills,
Oxymel of colchicum, each, half a fl. ounce.
Syrup, one fl. ounce.
Mix. A teaspoonful, three or four times a-day, in hydrothorax. *Ferriar.*

VINEGAR OF COLCHICUM MIXTURE.

R. Vinegar of colchicum,
Syrup, each, half a fl. ounce.
Carbonate of magnesia, one drachm and a half.
Peppermint water, four ounces.
Mix. In tablespoonful doses, as a sudorific in gout, &c. *Foy.*

MIXTURE OF COLCHICUM AND SQUILLS.

R. Oxymel of colchicum,
" squills,
Wine of tobacco, each, half fl. ounce.
Mix. A teaspoonful four times a-day. As a diuretic in dropsy. Requires caution. *Ferriar.*

MIXTURE OF WINE OF COLCHICUM.

R. Wine of colchicum root, half fl. drachm.
Carbonate of magnesia, one drachm.
Cinnamon water,
Distilled water, each, three fl. ounces.
Mix. A tablespoonful, three times a-day. As a diuretic. *Ellis.*

MIXTURE OF COLCHICUM AND AMMONIA.

R. Oxymel of colchicum,
Solution acetate of ammonia, each, two fl. ounces.
Parsley water, six fl. ounces.
Mix. A spoonful, every two hours, as a diuretic. *Augustin.*

LINIMENT OF COLCHICUM AND CAMPHOR.

R. Tincture of colchicum root,
" camphor, each, three fl. ounces.
Mix. As an embrocation, in gout, rheumatism, and neuralgia. *Leycock.*

COLLINSONIA.

HORSE-BALM.

The *Collinsonia Canadensis* is a native plant, found in most parts of this country, in woods; it is from two to three feet in height.

Sex. Syst. Diand. monog. *Nat. Syst.* Lamiales.

Linn. Sp. Pl. 39. *Griffith, Med. Bot.* 513.

The whole plant has a strong, peculiar, unpleasant odor, and a warm, pungent taste: these are most developed in the root; this is knotty and hard. It is diuretic, diaphoretic, and tonic; and is popularly used in diseases of the bladder, leucorrhœa, and dropsy, and externally as a fomentation to wounds, ulcers, &c.

DECOCTION OF HORSE-BALM.

R. Horse-balm root, bruised, two ounces.
Water, two pints.
Boil for half an hour, and strain. Dose, a wineglassful, every two hours, in dropsy.

TINCTURE OF HORSE-BALM.

R. Bruised horse-balm root, two ounces.
Diluted alcohol, one pint.
Macerate for fourteen days, express, and filter. Dose, a teaspoonful.

OIL OF HORSE-BALM.

R. Horse-balm, at will.
Water, sufficient.
Distil, return product, and redistil; collect product. Dose, five to ten drops.

COLLODIUM.

See GOSSYPIMUM.

COLOCYNTHIS.

COLOCYNTH.

Colocynth is the dried pulp of the fruit of *Citrullus colocynthis*, a cucurbitaceous vine found in many parts of Asia and Africa. The fruit is a round pepo, of the size and color of an orange, with a thin, but hard rind, and containing numerous seeds, enveloped in a white, spongy pulp.

Sex. Syst. Monœc. monadelph. *Nat. Syst.* Cucurbitaceæ.

Royle, Mat. Med. 396. *Griffith, Med. Bot.* 306.

It occurs in the shops in white, round, light balls, composed of a spongy, dried pulp, enclosing numerous seeds. This spongy substance has a faint odor, but an intensely bitter and nauseous taste. It is a powerful drastic and hydragogue purgative, and is seldom given alone. Dose, five to ten grains.

POWDER OF COLOCYNTH.

- R. Colocynth, one to three grains.
 Gum Arabic, }
 Liquorice, } each, five grains.
 Sugar, }
- Mix. As a hydragogue purgative. *Augustin.*

EXTRACT OF COLOCYNTH.

- R. Sliced colocynth, one pound.
 Distilled water, two gallons.
- Mix, and boil over a slow fire for six hours, occasionally adding more water. Strain while hot, and evaporate to proper consistence. *Ed. Ph.*

- R. Sliced colocynth, three pounds.
 Distilled water, half a gallon.
- Macerate for thirty-six hours, occasionally pressing with the hand. Express strongly, strain, and evaporate to proper consistence. *Lond. Ph.*

Dose, from five grains to half a drachm.

- R. Colocynth pulp, at will.
 Diluted alcohol, nine grains.
- Macerate the pulp in a sufficiency of menstruum to cover it for twenty-four hours, displace till exhausted, and evaporate to the proper consistence. This extract is more active than the aqueous preparation. *W. Procter.*

COMPOUND EXTRACT OF COLOCYNTH.

- R. Colocynth sliced, six ounces.
 Powdered aloes, twelve ounces.
 " scammony, four ounces.
 " cardamom, one ounce.
 Castile soap, three ounces.
 Diluted alcohol, one gallon.

Macerate the colocynth in the alcohol, with a mild heat for four days, express, and filter; then add the aloes, scammony, and soap; evaporate to proper consistence, and add the cardamom. *U. S. Ph.*

A valuable and safe cathartic. In small doses, laxative. Dose, five to twenty grains.

COMPOUND CATHARTIC PILLS.

(ANTIBILIOUS PILLS.)

- R. Compound extract of colocynth, powdered, half an ounce.
 Extract of jalap, in powder,
 Calomel, each, three drachms.
 Gamboge, in powder, two scruples.
- Mix, and with water form mass, to be divided into one hundred and eighty pills. *U. S. Ph.*

A most excellent cathartic, when it is wished to act on the biliary organs. Dose, one to three or four pills. Each pill contains one grain of calomel.

COMPOUND PILLS OF COLOCYNTH.

- R. Aloes, eight parts.
 Scammony, each, four parts.
 Powdered colocynth,
 Sulphate of potassa,
 Oil of cloves, each, one part.
 Rectified spirit, sufficient.

Pulverize the aloes, scammony, and sulphate of potassa, together; mix the colocynth with them, add oil of cloves, and with the rectified spirit beat into a mass, to be divided into five grain pills. *Ed. Ph.*

Dose, five to fifteen grains.

- R. Compound extract of colocynth, one scruple.
 Resin of jalap, six grains.
 Compound powder of scammony,
 Calomel, each, ten grains.
 Tartar emetic, one grain.
 Castile soap, five grains.
 Oil of cinnamon, four drops.

Beat well together, and form fifteen pills. Dose, one to three, at bedtime. *Meigs.*

- R. Extract of colocynth, two drachms.
 Resin of jalap, one drachm.
 Soap, one drachm and a half.
 Guaiacum, three drachms.
 Tartar emetic, eight grains.
 Oil of juniper,
 " rosemary, each, four drops.
 Syrup of buckthorn, sufficient.

Mix well, and divide into four-grain pills. Dose, one to three. *Barclay.*

PILLS OF COLOCYNTH AND HENBANE.

- R. Extract of colocynth, two parts.
 " henbane, one part.

Beat together, with a few drops of rectified spirit, and divide into five-grain pills. *Ed. Ph.*

Dose, five to twenty grains.

COLOCYNTH CLYSTER.

- R. Extract of colocynth, one scruple.
 Common salt, three drachms.
 Syrup of dogtooth-grass, one ounce and a half.
 Infusion of chamomile, five ounces.

In cerebral affections. *Phæbus.*

COLOCYNTH MIXTURE.

- R. Colocynth, one drachm.
 Boiling water, six ounces.
- Boil for ten minutes, strain, and add, when cold,
 Hoffmann's anodyne, one drachm.
 Syrup of orange peel, one ounce

Mix. A spoonful, three times a-day, in ascites, or hydrothorax. *Augustin.*

TINCTURE OF COLOCYNTH.

R. Colocynth, eight parts.
Star anise, one part.
Alcohol, ninety-six parts.

Macerate for three days, and filter. Dose, fifteen to twenty drops. *Van Mons.*

TINCTURE OF COLOCYNTH MIXTURE.

R. Tincture of colocynth, two drachms.
Solution of antimoniated soap, (*Cod. Hamb.*), six drachms.

Mix. Dose, twenty to thirty drops a-day, in lepra and obstinate cutaneous affections. *Heim.*

COLOMBA.

COLUMBO.

Columbo is the root of *Cocculus palmatus*, a climbing plant, a native of Mozambique.

Sex. Syst. Diœc. hexand. *Nat. Syst.* Menispermaceæ.

Hooker, Bot. Mag. 2970-71. Griffith, Med. Bot. 103.

As found in the shops, the root is in round slices, externally of a brown, wrinkled appearance, internally yellow. It is somewhat aromatic, and has a very bitter taste. It is an excellent bitter tonic, with no astringency. It is useful in diseases where the pure bitters are required, and generally agrees with the stomach. Dose, in powder, is from ten to thirty grains.

POWDER OF COLUMBO AND IRON.

R. Powdered columbo, }
Subcarbonate of iron, } each, one
Powdered rhubarb, } drachm.
" ginger, }

Mix, and make twelve powders; one to be taken every four hours. *A. T. Thomson.*

POWDER OF COLUMBO AND MAGNESIA.

R. Powdered columbo, ten grains.
Magnesia, two scruples.

Mix for a dose. In cardialgia. *Brugnatelli.*

POWDER OF COLUMBO AND TARTRATE OF IRON.

R. Tartrate of iron and potassa, two scruples.
Powdered columbo, half a drachm.

Mix, and divide into four powders. One every three or four hours, in syrup. *Ellis.*

COMPOUND PILLS OF COLUMBO.

R. Powdered columbo, four scruples.
Opium, four grains.
Oil of peppermint, ten drops.
Syrup of pinks, sufficient.

Beat into mass, and form thirty pills. Two, three times a-day, in spasmodic vomiting.

St. Marie.

R. Powdered columbo, one drachm.
" rhubarb, two scruples.
Extract of chamomile, two drachms.
Oil of caraway, five drops.
Syrup of saffron, sufficient.

Form mass, and divide into pills of four grains. Four to be taken a-day, in mania with amenorrhœa. *Augustin.*

INFUSION OF COLUMBO.

R. Bruised columbo, half an ounce.
Boiling water, one pint.

Macerate two hours in a covered vessel, and strain. *U. S. Ph.*

Dose, two fl. ounces, three or four times a-day.

INFUSION OF COLUMBO AND GINGER.

R. Bruised columbo, one ounce.
" ginger, two drachms.
Boiling water, one pint.

Infuse, and strain. A wineglassful cold, every two hours, in chronic diarrhœa. *Ellis.*

INFUSION OF COLUMBO, RHUBARB, &c.

R. Bruised caraway, "
" columbo, "
" rhubarb, each, one scruple.
Boiling water, eight fl. ounces.

Digest for two hours, and strain.

R. To strained liquid, three and a half fl. ounces;

Add

Tincture of rhubarb, one fl. drachm.
Syrup of ginger, two fl. drachms.

Mix. Dose, a teaspoonful to a tablespoonful, in diarrhœa. *Ellis.*

MIXTURE OF COLUMBO.

R. Columbo, half a drachm.
Boil in

Water, three to five fl. ounces.

Strain, and add

Carbonate of potassa, ten grains.
Lemon juice, three fl. drachms.
Tincture of opium, twelve drops.

Mix. A tea to a tablespoonful, every hour, as an anti-emetic. *Ellis.*

COMPOUND DECOCTION OF COLUMBO.

R. Columbo,
 Quassia, each, two drachms.
 Orange peel, one drachm.
 Rhubarb, one scruple.
 Subcarbonate of potassa, half a drachm.
 Water, twenty fl. ounces.
 Boil down to a pint, strain, and add
 Tincture of lavender, half fl. ounce.

Coxe.

EXTRACT OF COLUMBO.

R. Bruised columbo, two parts.
 Alcohol, three parts.
 Water, nine parts.
 Macerate for twenty-four hours, distil off the
 alcohol, and evaporate to proper consistence.

Cottureau.

TINCTURE OF COLUMBO.

R. Columbo, bruised, four ounces.
 Diluted alcohol, two pints.
 Macerate for fourteen days, and filter. *U. S. Ph.*
 Dose, one to four fl. drachms.

MIXTURE OF COLUMBO AND CASCARILLA.

R. Powdered columbo, half an ounce.
 Water, ten fl. ounces.
 Boil to six ounces, add, towards close,
 Cascarella, two drachms.
 Strain, and add
 Tincture of orange peel, two drachms.
 Syrup of cinnamon, one ounce.
 A spoonful every hour, in chronic diarrhoea.

Berends.

MIXTURE OF COLUMBO AND SALEP.

R. Extract of columbo, one drachm.
 Decoction of salep, three ounces.
 Fennel sugar, two scruples.
 Mix. Two spoonfuls a-day, in the diarrhoea of
 children.

Wendt.

COMPTONIA.

SWEET FERN.

A shrubby, indigenous plant, found in the
 northern and middle states.

Sex. Syst. Monœc. triand. *Nat. Syst.* Myrica-
 cœæ.

Aiton, Hort. Kew. iii. 334. Griffith, Med. Bot.
 584.

It is aromatic, especially when bruised. It
 is a mild astringent, and is used in bowel com-
 plaints, in the form of decoction.

CONIUM.

HEMLOCK.

This is an umbelliferous, poisonous plant, the
Conium maculatum, a native of Europe, but natu-
 ralized in many parts of this country, growing
 in waste places.

Sex. Syst. Pentand. digyn. *Nat. Syst.* Api-
 acœæ.

Linn. Sp. Pl. 349. Griffith, Med. Bot. 339.

Both the seeds and the leaves are used. The
 latter have a strong, unpleasant, narcotic odor,
 and a somewhat bitterish taste; the seeds pos-
 sess these qualities in a less degree. They are
 both narcotic, but neither stimulant nor seda-
 tive, and are given in a variety of complaints,
 to alleviate, especially in malignant tumors, in
 which they are thought to exert a curative in-
 fluence. The dose, in powder, is from three to
 four grains of the leaves; and of the seeds
 rather less; twice a-day, gradually increasing.
 The active principle, *conia*, is seldom used.

POWDER OF HEMLOCK.

R. Powdered hemlock, five grains.
 " liquorice, six grains.

Make a powder, to be taken three times a-day.
 In scirrhus affections, scrofula, old ulcers, &c.

FECULA OF HEMLOCK.

R. Hemlock juice, at will.
 Place on the fire, remove the coagulum, wash
 it well, dry, and pulverize. Dose, five to twenty
 grains. *Van Mons.*

EXTRACT OF HEMLOCK.

R. Fresh hemlock leaves, one pound.
 Bruise in a stone mortar, sprinkling on them a
 little water; express the juice, heat to boiling
 point, strain, and evaporate to proper con-
 sistence. *U. S. Ph.*

Dose, two grains, morning and evening,
 gradually increased till it affects the system.

ALCOHOLIC EXTRACT OF HEMLOCK.

R. Hemlock leaves, in coarse
 powder, one pound.
 Diluted alcohol, four pints.

Moisten powder with half a pint of the alcohol,
 and allow to macerate for twenty-four hours;
 place in percolating apparatus and add re-
 mainder of alcohol; when this has disappeared,
 add water so as to keep powder covered. Stop
 process when the liquid passing, causes precipi-
 tate in that previously obtained. Distil off al-
 cohol, and evaporate to proper consistence.

U. S. Ph

Dose, one to two grains, twice a-day, gradu-
 ally increasing.

PILLS OF HEMLOCK AND IPECACUANHA.

R. Extract of hemlock,
Dover's powder, each, ten grains.
Mix, and form five pills. Two at bedtime, in
pulmonary irritation, with rheumatic, and other
local pains. *Ellis.*

R. Extract of hemlock, five drachms.
Powdered ipecacuanha, one drachm.
Molasses, sufficient.
Mix. Dose, four or five grains. *Lond. Ph.*

PILLS OF HEMLOCK AND CALOMEL.

R. Extract of hemlock, two scruples.
Calomel, fifteen grains.
Mix, and form fifteen pills. One, three times
a-day, in syphilitic affections. *Ellis.*

PILLS OF HEMLOCK AND DANDELION.

R. Extract of hemlock, }
" dandelion, } each, one
Gum Arabic, } drachm.
Soap, }
Honey, } sufficient.
Mix, and form three-grain pills. In glandular
and other tumors. *Brera.*

INFUSION OF HEMLOCK.

R. Hemlock leaves, half an ounce.
Boiling water, one pint.
Infuse and strain. As a wash to cancerous
and malignant ulcers. *Ellis.*

HEMLOCK COLLUTORY.

R. Hemlock leaves, two drachms.
Poppy seed, one drachm.
Henbane seed, half a drachm.
Milk, half a pint.
Boil slightly, and strain. In rheumatic tooth-
ache. *Phæbus.*

TINCTURE OF HEMLOCK.

R. Hemlock leaves, four ounces.
Diluted alcohol, two pints.
Macerate for fourteen days, express, and filter.
U. S. Ph.
Dose, thirty drops to a fl. drachm.

ETHEREAL TINCTURE OF HEMLOCK.

R. Hemlock leaves, one part.
Sulphuric ether, four parts.
Macerate for two days, and filter. *Soubéiran.*

OINTMENT OF HEMLOCK.

R. Fresh hemlock leaves,
Prepared lard, each, one pound.

Boil till the leaves become crisp, then express
through linen. *Lond. Ph.*

R. Juice of hemlock leaves, one part.
Lard, four parts.

Boil till moisture is expelled, and express through
a cloth. As a dressing to painful ulcers.
Svediaur.

OIL OF HEMLOCK.

R. Fecula of hemlock, one part.
Olive oil, four parts.

Mix, and strain. As a friction to the abdomen
in ileus, and to the anus in hemorrhoids.
Van Mons.

HEMLOCK PLASTER.

R. Elemi, two parts.
White wax, one part.

Melt together, and add
Alcoholic extract of hem-
lock, nine parts.

As an application to painful ulcers and malign-
ant tumors. *Guibourt.*

R. Juice of hemlock, twelve parts.
Powdered hemlock, six parts.

Evaporate to a pasty consistence, and add to a
melted mixture of

Yellow wax, ten parts.
Turpentine, four parts.
Olive oil, six parts.

Used as above. *Pideret.*

COMPOUND HEMLOCK PLASTER.

R. Yellow wax, eight parts.
Resin, sixteen parts.
Fecula of hemlock, forty-eight parts.

Melt together, and add a melted mixture of
Ammoniac, four parts.
Turpentine, one part.

Stir well till cold. Used as above. *Van Mons.*

R. Hemlock plaster, one ounce.
Burgundy pitch, half an ounce.
Diachylon, each, half an ounce.

Mix, and spread, then sprinkle with from six to
twelve grains of tartar emetic, according to the
age of the patient. Said to be very useful in
hooping-cough.

R. Soap plaster, one ounce.
Extract of hemlock, half an ounce.
Powdered belladonna, two drachms.
Mix. *Hamburg Cod.*

CATAPLASM OF HEMLOCK.

R. Powdered hemlock,
Linseed meal, each, one ounce.
Fecula of hemlock,
Boiling water, each, sufficient.

Mix. As an application to cancerous and other painful ulcers. *Guibourt.*

R. Hemlock leaves, a handful.
Henbane leaves, half a handful.
Lard, half an ounce.
Rose oil, one ounce.
Decoction of mallows, sufficient.

Mix, and boil to the consistence of a cataplasm. *Pierquin.*

R. Powdered hemlock, two ounces.
Carrot juice, one pound.
Tincture of opium and saffron, three drachms.
Linseed meal, sufficient.

Make a cataplasm. In cancerous ulcers. *Pideret.*

MIXTURE OF HEMLOCK AND PAREGORIC.

R. Extract of hemlock, one drachm.
Paregoric elixir,
Syrup of tolu, each, half fl. ounce.
Rose water, four fl. ounces.

Mix. From half to a teaspoonful once in four hours, to a child a year old, watching the effect, in pertussis. *Pearson.*

SUPPOSITORY OF HEMLOCK.

R. Extract of hemlock,
White wax, each, one part.
Butter of cocoa, four parts.

Mix. In painful hemorrhoids and spasms of the rectum. *Beral.*

SOLUTION OF CONIA.

R. Conia, two parts.
Distilled water, two hundred parts.
Alcohol, thirteen parts.

Mix. Used in scrofulous ophthalmia, and intolerance of light, by frictions around the orbits several times a-day. *Fronmuller.*

CONTRAYERVA.

CONTRAYERVA.

Contrayerva is the root of *Dorstenia contrayerva*, and other species, all natives of South America, Mexico, and the West Indies. They are perennial, dwarf, herbaceous plants, usually growing in high, rocky places.

Sex. Syst. Tetrand. monog. *Nat. Syst.* Moraceae.

Linn. Sp. Pl. 124. Griffith, Med. Bot. 577.

The root is oblong, hard, rough, and solid, of a reddish-brown color, with numerous, long, yellowish fibres. It has an aromatic odor, and a warm, pungent, somewhat bitterish taste. It is a stimulant, tonic, and diaphoretic, and has been found useful in low states of the system. The dose of the powder is about half a drachm.

COMPOUND POWDER OF CONTRAYERVA.

R. Powdered contrayerva, six drachms.
Virginia snakeroot, two drachms.
Prepared chalk, one ounce.

Mix. Dose, a teaspoonful, two or three times a-day, as a tonic. *Spielmann.*

EXTRACT OF CONTRAYERVA.

R. Contrayerva, bruised, one part.
Boiling water, six parts.

Infuse for three days, boil, strain, express, and evaporate to proper consistence. Dose, one to two scruples. *Palat. Cod.*

DECOCTION OF CONTRAYERVA.

R. Contrayerva, bruised, one drachm.
Water, one pint.

Boil for ten minutes, and strain. *Taddei.*

GARGLE OF CONTRAYERVA.

R. Contrayerva, half an ounce.
Figs, one ounce.
Water, eighteen fl. ounces.

Mix, and boil down to two-thirds. *Saunders.*

TINCTURE OF CONTRAYERVA.

R. Contrayerva, one part.
Diluted alcohol, eight parts.

Digest in one-half the alcohol for four days, strain and digest with the other half of the alcohol; strain, and filter the united liquids. Dose, thirty to forty drops. *Guibourt.*

COMPOUND TINCTURE OF CONTRAYERVA.

R. Contrayerva,
Virginia snakeroot, each,
one ounce and a half.

Juniper berries, one ounce.
Spirit of ammonia, six fl. ounces.
Alcohol, one pint.

Macerate for four days, express, and filter. Dose, forty to eighty drops, as a stimulating diaphoretic. *Lip. Dispen.*

CONTRAYERVA MIXTURE.

R. Powdered contrayerva, eight grains.
Cinnamon water, one fl. ounce and a half.

Nutmeg water, two fl. drachms.
Syrup of cinnamon, three fl. drachms.

Mix. A spoonful as a stomachic. *Swediaur.*

R. Gum tragacanth,
Comp. contrayerva powder, each,
one scruple.
Distilled water, one fl. ounce

Cinnamon water, three fl. drachms.
Syrup of saffron, one fl. drachm.
Mix. As a quieting draught for a woman after delivery.
Merriman.

CONVOLVULUS PANDURATUS.

WILD POTATO.

This is a native plant, with a large perennial root, found in sandy soils in most parts of the United States.

Sex. Syst. Pentand. monog. *Nat. Syst.* Convolvulaceæ.

Linn. Sp. Pl. 219. *Griffith, Med. Bot.* (Ipomœa) 476.

The root, as found in the shops, is in circular pieces, of a yellowish-brown color, scarcely any smell, and a bitterish, somewhat acrid taste. It is feebly cathartic, but is said to be a good diuretic in calculous complaints. Dose, as purgative, forty grains. As a diuretic, it is given in decoction.

COPAIBA.

COPAIBA—COPAIVA.

This is the juice of *Copaifera officinalis* and other species, all large trees, natives of South America, and principally of Brazil.

Sex. Syst. Decand. monog. *Nat. Syst.* Fabaceæ.

Linn. Sp. Pl. 557. *Griffith, Med. Bot.* 264.

This juice is clear and transparent, of a pale yellowish color; a peculiar and somewhat unpleasant odor, and a bitterish, pungent, nauseous taste. It is rather less consistent than olive oil when fresh, but, by age, becomes thicker, and may even assume the solidity of a resin. It is stimulant, diuretic, and cathartic, and is principally used in morbid discharges from mucous surfaces, especially those of the genito-urinary organs. The dose is from twenty drops to a fl. drachm, three times a-day.

BOLUSES OF COPAIBA.

R. Copaiba, two drachms.
Gum Arabic, one ounce.
Powdered liquorice, sufficient.

Mix, and form mass. Dose, a scruple to two scruples. *Radius.*

R. Copaiba, one ounce.
Calcined magnesia, one drachm.
Opium, four grains.
Powdered cubebs, three drachms.
Syrup of poppies, sufficient.

Mix, and divide into boluses of half a drachm. Two to be taken at night, in gonorrhœa.

Henschel.

PILLS OF COPAIBA.

R. Copaiba, two ounces.
Magnesia, (freshly prepared), one drachm.

Mix, and set aside, till it concretes into a pilular mass; this to be divided into two hundred pills. *U. S. Ph.*

Dose, two to six, two or three times a-day.

R. Copaiba,
Powdered cubebs, each, one drachm.
Mucilage of gum Arabic, sufficient.

Mix, and divide into three-grain pills. Five to ten a-day. *Hildenbrand.*

R. Copaiba, one ounce.
Dragon's blood, half an ounce.
Calomel, one drachm.
Conserve of roses, four ounces.

Mix, and make pills of six grains. Dose, from three to five a-day. *Foy.*

PILLS OF COPAIBA AND CUBEBS.

R. White wax, two drachms.
Melt, by a gentle heat, and add
Copaiba, half an ounce.
Powdered cubebs, one ounce.

Incorporate well, and divide into two hundred and forty pills. *Berens.*

PILLS OF COPAIBA, CUBEBS, AND TURPENTINE.

R. Oil of copaiba,
" cubebs,
" turpentine, each, one fl. drachm.
Magnesia, two drachms.

Mix, and form sixty pills.

This is said to be an excellent preparation. It was communicated by Mr. Procter, who is unacquainted with the author of it.

COPAIBA CONFECTION.

R. Turpentine, one ounce.
Copaiba, half an ounce.
Mix well, in a warmed mortar, and add
Mucilage of gum Arabic, one ounce.
Conserve of roses, four ounces.

Dose, two to three drachms, three times a-day, in obstinate gonorrhœa. *Swediaur.*

R. Copaiba,
Powdered cubebs, each, four drachms
and a half.
Yolk of egg, one.
Conserve of roses, half an ounce.

Mix. A teaspoonful three or four times a-day. *Voght.*

R. Copaiba,
Powdered cubebs, each, two ounces.
" alum, one ounce.
Opium, five grains.

Incorporate well. One drachm to be taken in the pulp of a prune, night and morning, and rapidly increased to two drachms. Is very effectual in gonorrhœa. *Berton.*

EMULSION OF COPAIBA.

- R. Copaiba,
Mucilage gum Arabic, each,
two ounces.
Water, twelve fl. ounces.
Rub the copaiba gradually with the mucilage
in a mortar, add the water by degrees, con-
stantly rubbing. *Beral.*

INJECTION OF COPAIBA.

- R. Copaiba, two drachms.
Mucilage of gum
Arabic, half an ounce.
Lime water, six fl. ounces.
Mix well. As an injection in ulcers of the
rectum, vagina, or urethra. *Abernethy.*
R. Copaiba, half an ounce.
Yolk of egg, sufficient.
Lime water, six ounces.
Honey of roses, three ounces.
Mix. As an injection in fistulous ulcers.
Plenck.

COPAIBA MIXTURE.

- R. Copaiba,
Sweet spirit of nitre, each,
half a fl. ounce.
Powdered gum Arabic,
" sugar, each, one drachm.
Compound spirit of
lavender, two fl. drachms.
Tincture of opium, one fl. drachm.
Distilled water, four fl. ounces.
Mix. A tablespoonful three times a-day in go-
norrhœa. *Chapman.*
R. Copaiba,
Powdered cubebs, each, half an ounce.
Gum Arabic, two drachms.
Paregoric elixir, half a fl. ounce.
Sugar, two drachms.
Water, eight fl. ounces.
Mix. A tablespoonful every three hours.
Ellis.

- R. Copaiba, half an ounce.
Mint water, }
Alcohol, } each, half a fl. ounce.
Syrup, }
Sweet spirit of nitre, half a fl. drachm.
Mix. To be taken in four doses in a day, fol-
lowed by demulcent drinks. In chronic gonor-
rhœa. *Ellis.*

- R. Copaiba,
Sweet spirit of nitre, each,
half a fl. ounce.
Compound spirit of lavender,
Laudanum, each, one fl. drachm.

- Powdered gum Arabic, one drachm.
Water, three fl. ounces.
Mix. A tablespoonful three times a-day in go-
norrhœa. *Chapman.*

- R. Copaiba half an ounce.
Mucilage of gum Arabic, two ounces.
Clarified honey, six drachms.
Solution of potassa, two drachms.
Essence of lemon, half a drachm.
Rose water, five fl. ounces.
Mix. Two to three spoonfuls, three times
a-day. *Stephenson.*

- R. Copaiba, one drachm and a half.
Powdered kino, one scruple.
Mucilage of gum Arabic, two drachms.
Syrup of rhatany, one ounce.
Decoction of chamomile, four ounces.
Mix. A spoonful every hour or two. *Foy.*

- R. Copaiba, half a drachm.
Tincture of allspice,
Syrup of mallows, each, two drachms.
Mucilage of gum Arabic, sufficient.
Water, one ounce.
Mix. To be taken morning and evening.
Saunders.

- R. Copaiba, }
Syrup of lemon, } each, one
Mint water, } ounce.
Orange-flower water, }
Sulphuric acid, one drachm.
Tragacanth, sufficient.
Mix. A spoonful, morning and evening.
Delpsch.

- R. Copaiba, }
Balsam of Tolu, } each, half an
Powdered gum Arabic, } ounce.
Elixir of vitriol, twenty drops.
Distilled water, six fl. ounces.
Mix. A tablespoonful, in chronic whooping-
cough. *Ellis.*

- R. Copaiba, one drachm.
Carbonate of potassa, thirty grains.
Extract of Peruvian
bark, twenty grains.
Gum Arabic, twelve grains.
Aloes, two grains.
Syrup of capillaire, two ounces.
Cinnamon water,
Mint water, each, three fl. ounces.
Mix. Dose, a wineglassful, several times a-day,
as an anthelmintic. *Pierquin.*

COPAHINE-MEGE.

- R. Oxidized copaiba, eighty parts.
Powdered cubebs,
Carbonate of soda, each, eight parts.
Calcined magnesia, five parts.

Mix, and let the mixture stand till solidified, and then make into small masses, which are to be covered with sugar. Dose, four or five, three times a-day. Oxidized copaiba is prepared by treating copaiba with nitric acid, and then washing with water.

Joscau.

TINCTURE OF COPAIBA.

R. Copaiba, one part.
Alcohol, eight parts.
Digest for several days, and filter. Dose, thirty to sixty drops. *Guibourt.*

ALKALINE TINCTURE OF COPAIBA.

R. Copaiba, one ounce.
Carbonate of potassa, one drachm.
Alcohol, four fl. ounces.
Digest, and filter. Dose, twenty-five to fifty drops. *Brunswick Ph.*

COMPOUND TINCTURE OF COPAIBA.

R. Copaiba, one ounce.
Guaiacum, two drachms.
Oil of sassafras, half a drachm.
Alcohol, four and a half fl. ounces.
Digest with a gentle heat; filter. *Foy.*
Dose, a drachm, in some infusion.

R. Copaiba, one ounce and a half.
Balsam Peru, half an ounce.
Carbonate of potassa, one drachm and a half.
Cut saffron, two drachms.
Alcohol, sixteen fl. ounces.
Mix, and digest for three days, and filter. Two spoonfuls, two or three times a-day, in water or in wine. *Wirtemberg Ph.*

OIL OF COPAIBA.

R. Copaiba, one ounce.
Water, one pint and a half.
Distil, return product several times, and redistil; separate oil. *Ed. Ph.*
Dose, ten to thirty drops, rubbed up with mucilage and water.

COPAIBA CLYSTER.

R. Copaiba, one to four drachms.
Extract of opium, one grain.
Yolk of egg, one.
Decoction of mallows, four to six fl. ounces.
Make an emulsion, as a clyster in gonorrhœa. *Guibourt.*

COPAIBA PASTE.

R. Sweet almonds, six drachms.
Mallow paste, one drachm.
Catechu, half a drachm.
Copaiba, three drachms.
Rub together into a paste. Dose, a spoonful. *Phœbus.*

COPTIS.

GOLDTHREAD.

The root of *Coptis trifolia*, a small evergreen plant, found in the more northern parts of both continents, in wet and boggy situations.

Sex. Syst. Polyand. polyg. *Nat. Syst.* Ranunculaceæ.

Salisbury, Linn. Trans. viii. 305. Griffith, Med. Bot. 87.

The roots, which are the parts used, are long, slender, orange-yellow, with no smell, but a strong, and purely bitter taste. It is a simple tonic bitter, with no astringency. It is used as a stomachic, and as a local application to aphthous sore mouth. Dose, in powder, ten to thirty grains.

INFUSION OF GOLDTHREAD.

R. Goldthread, one ounce.
Boiling water, one pint.
Infuse for an hour, and strain. Dose, half fl. ounce to two fl. ounces. *Dunglison.*

TINCTURE OF GOLDTHREAD.

R. Goldthread, one ounce.
Diluted alcohol, one pint.
Macerate for a week, and filter. Dose, one fl. drachm to three fl. drachms. *Wood.*

CORIANDRUM.

CORIANDER.

This is the fruit of *Coriandrum sativum*, an annual plant, a native of the south of Europe, but naturalized in many other parts of that quarter of the world, and also extensively cultivated.

Sex. Syst. Pentand. digyn. *Nat. Syst.* Apiaceæ.

Linn. Sp. Pl. 367. Griffith, Med. Bot. 341.

The fruit is somewhat globular, of a grayish, or brownish-yellow color, and often separated into two portions, or mericarps. It has a pleasant aromatic smell and taste. It is principally used as an adjuvant to other articles, to disguise their taste, or to modify their griping qualities. Dose, from a scruple to a drachm.

COMPOUND POWDER OF CORIANDER.

R. Powdered coriander, one drachm and a half.
Ivory filings, }
Burnt hartshorn, } each, one scruple.
Prepared chalk, }
Powdered cinnamon, half a scruple.
Sugar of roses, sufficient.
Mix well. A teaspoonful, after eating, as a stomachic. *Pierquin.*

R. Powdered coriander,
 " rhubarb,
 " columbo, each, ten grains.
 Mix. To be taken at bedtime, in dyspepsia
 with flatulence and costiveness. Stomachic.
Ainslie.

TINCTURE OF CORIANDER.

R. Coriander, one part.
 Diluted alcohol, eight parts.
 Macerate for a week, and filter. *Beral.*

COMPOUND WATER OF CORIANDER.

R. Coriander, eight ounces.
 Citron water, one fl. ounce.
 Nutmeg, }
 Storax, } each, half an ounce.
 Benzoin, }
 Vanilla, three drachms.
 Alcohol, forty-eight fl. ounces.

Macerate for two days, and distil. Dose, half
 fl. ounce, as a stomachic and carminative.
Spielmann.

CORNU.

HARTSHORN.

This is officinally the horn of the *Cervus elaphas*, or European stag; but that of our native species has the same properties. It is found in the shops, in the form of shavings, which are of a yellowish-white color, friable, and without smell or taste. They consist principally of gelatine and phosphate of lime, and are employed to make a jelly, which is not superior to any other animal jelly.

HARTSHORN JELLY.

R. Hartshorn shavings, four ounces.
 Water, eight ounces.
 Muriatic acid, one drachm.

Beat together for ten minutes, wash in two or three waters, boil with fresh water for half an hour, express, strain, and add

Sugar, four ounces.

Boil down sufficiently for the mixture to jelly, on cooling. *Ferrez.*

COMPOUND JELLY OF HARTSHORN.

R. Hartshorn jelly, eight ounces.
 Paste of sweet almonds, one ounce.
 Sugar, half an ounce.
 Orange-flower water, one drachm.
 Essence of citron, twelve drops.

Mix, and heat; then permit to cool. *Guibourt.*

BURNT HARTSHORN.

R. Burn pieces of hartshorn in an open vessel, till quite white; powder, and prepare like prepared chalk.

Lond. Ph.

Dose, half a drachm to two drachms.

DENTIFRICE OF BURNT HARTSHORN.

R. Powdered burnt hartshorn, one ounce.
 Orris root, two drachms.
 Dragon's blood, one drachm.
 Oil of roses, two drops.

Mix. *Steph. & Church.*

CORNUS.

DOGWOOD.

CORNUS CIRCINATA.

ROUND-LEAVED DOGWOOD.

A large shrub, native of the United States, with a bitter, astringent, aromatic bark.

CORNUS FLORIDA.

DOGWOOD.

A small tree found in most parts of the United States, flowering early in the spring. The bark, especially of the root, has a bitter, astringent, somewhat aromatic taste, and a faint odor.

CORNUS SERICEA.

SWAMP DOGWOOD.

A large shrub, like the others, a native of this country, with a bark of the same qualities. *Sex. Syst.* Tetrand. monog. *Nat. Syst.* Cornaceæ.

The barks of these shrubs and trees are tonic and astringent, and are said to have much the same properties as Peruvian bark, and may be used as a substitute for it. They are, however, much inferior. The dose of the powder is from a scruple to a drachm.

DECOCTION OF DOGWOOD.

R. Dogwood bark, bruised, one ounce.
 Water, one pint.

Boil for half an hour, and strain. A wine-glassful every hour, in the apyrexia of intermittents. *U. S. Ph.*

R. Dogwood bark,
Sassafras bark, each, six ounces.
Water, eight pints.
Boil, and strain. Given as above. *Niemann.*

PILLS OF ROUND-LEAVED DOGWOOD.

R. Extract of round-leaved dog-
wood, one drachm and a half.
Powdered ginger, ten grains.
Dried carbonate of soda, ten grains.
Mix, and form twenty-four pills. *Reece.*

WINE OF ROUND-LEAVED DOGWOOD.

R. Extract of round-leaved dog-
wood, three drachms.
White wine, one pint and a half.
Digest for fourteen days, and filter. *Reece.*

COTULA.

MAYWEED.

A small annual syngenesious plant, a native of Europe, and extensively naturalized in the United States. It is the *Maruta cotula* of botanists. (*Anthemis cotula*. Linn.)

Sex. Syst. Syng. frust. Nat. Syst. Asteraceæ.
De Candolle, Prod. vi. 13. Griffith, Med. Bot. 399.

The whole herb has a strong, unpleasant smell, and a bitter, warm taste. It has much the same properties as chamomile as an internal remedy, and may be used as a substitute for it; but its disagreeable odor renders it nauseous to most persons. It is also employed in amenorrhœa, and as an antispasmodic. It is usually given in infusion.

INFUSION OF MAYWEED.

R. Mayweed, half an ounce.
Boiling water, one pint.

Macerate for a quarter of an hour, in a covered vessel, and strain. Used in same cases, and in same manner as infusion of chamomile.

CATAPLASM OF MAYWEED.

R. Mayweed, fresh, at will.
Bruise, or cut very fine. When applied as a cataplasm, it produces redness and vesication as rapidly as Spanish flies. *Ashley.*

CREASOTUM.

CREASOTE.

A peculiar substance, analogous to the volatile oils, existing in the products of the distillation of wood. It is a colorless, oleaginous fluid, of a burning, caustic taste, and of an odor resembling that of smoked meat. It is corrosive,

but, in a diluted state, is an effectual preserver of animal substances. It is irritant, narcotic, styptic, &c., and has been employed in a variety of diseases, both internally and as a local application.

CREASOTE.

R. Tar, at will.
Distil, changing the recipients several times, till the residue has the consistence of pitch; agitate the heavy oil that passes with a little concentrated sulphuric acid; then mix it with its own volume of water, and rectify in small retorts; dissolve the product which sinks to the bottom in a hot solution of potassa, aided by a gentle heat; when it cools, add a slight excess of sulphuric acid, separate the oil, and distil it in small quantities at a time; again rectify it two or three times. Dose, one to two drops, mixed with water.

CREASOTE MIXTURE.

R. Creasote, six drops.
Powdered mallow,
Sugar, each, one drachm.
Mix well, and divide into three powders, to be taken during the day. In hemoptysis and phthisis. *Radius.*

CREASOTE PILLS.

R. Creasote, one drachm.
Powdered mallow root, sufficient
to make one hundred and ten pills. Said to be beneficial in acute rheumatism. Four or five to be taken, morning and evening. *Riech.*

R. Creasote, one drachm.
Extract of liquorice,
one drachm and a half.
Water, sufficient.

Beat into mass, and make pills of two grains. Two, morning and evening, in phthisis and laryngitis. *Radius.*

R. Creasote, ten drops.
Powdered liquorice,
Mucilage of gum Arabic,
each, sufficient.

Mix, and form twenty pills. Dose, one, three times a-day. In bronchitis, neuralgia, &c. *Ellis.*

R. Creasote, one drachm.
Extract of liquorice,
Galbanum, each, half a drachm.
Powdered mallows, two drachms.

Form mass, and divide into one hundred and twenty pills. Six, four times a-day, in phthisis. *Riech.*

ALCOHOLIC SOLUTION OF CREASOTE.

R. Creasote, one part.
Alcohol, sixteen parts.

Mix. A drop or two placed in the cavity of an aching tooth, will afford relief. *Marinus.*

R. Creasote, one fl. drachm.
Alcohol, two fl. ounces.

Mix. Used with water as a mouth-wash, to correct fetor of the mouth, and to check caries of the teeth. *Buchner.*

R. Creasote, one fl. drachm.
Alcohol, one fl. ounce and a half.
Tincture of cochineal, six fl. drachms.
Oil of mint, thirty-six drops.

Mix. Used as above. *Ringhini.*

CREASOTE LOTION.

R. Creasote, half a drachm.
Water, five ounces.

Mix. As a lotion in psora, &c. *Radius.*

R. Creasote, ten drops.
Vinegar, two fl. drachms.
Water, two fl. ounces.

Mix. As an application to phagedenic ulcerations and chancres. *Short.*

CREASOTE COLLUTORY.

R. Creasote, three drops.
Compound spirit of lavender, twenty drops.
Distilled water, half fl. ounce.

Mix. *Black.*

R. Creasote, four drops.
Distilled water, two fl. ounces.

Mix. In irritation of the gums. *Fremanger.*

R. Creasote, half a fl. drachm.
Powdered gum Arabic, one ounce and a half.
Camphor water, two fl. ounces and a half.

Mix. To be used every two hours in aphthous ulceration of the mouth. *Magendie.*

CREASOTE MIXTURE.

R. Creasote, sixteen drops.
Acetic acid, each, sixteen drops.
Compound spirit of juniper, each, one fl. ounce.
Syrup, each, one fl. ounce.
Water, fourteen fl. ounces.

Mix the creasote with the acid, gradually add the water, and then the spirit and syrup. *Ed. Ph.*

Dose, one fl. ounce.

R. Creasote, two to four drops.
Mucilage of gum Arabic, one ounce.
Infusion of salep, five ounces.
Sugar, one drachm.

Mix. A spoonful every two hours in the colicque diarrhœa of phthisis. *Radius.*

R. Creasote, five drops.

Mucilage of gum Arabic, three fl. ounces.
Syrup of mallows, one fl. drachm.

Mix. A spoonful every three hours in hemoptysis. *Santoni.*

R. Creasote, one drop.
Camphor water,
Compound infusion of gentian, each, six fl. drachms.

Mix. To check vomiting. *Santoni.*

CREASOTE OINTMENT.

R. Creasote, half a fl. drachm.
Lard, one ounce.

Melt the lard, add the creasote, and stir till cold. *U. S. Ph.*

As an application in scaly, cutaneous eruptions, ulcers, &c.

R. Simple cerate,
Oil of almonds, each, one ounce.
Creasote, thirty drops.

Mix. As an application in scrofulous caries. *Fremanger.*

COMPOUND OINTMENT OF CREASOTE.

R. Creasote,
Solution of subacetate of lead, each, ten drops.
Extract of opium, one grain and a half.
Lard, one ounce.

Mix. As an application to chilblains. *Devergie.*

R. Creasote, half a drachm.
Purified animal charcoal, one drachm.
Alcohol, one fl. drachm and a half.
Spermaceti ointment, one ounce and a half.

Mix. Used as an application to burns, to be diluted, if necessary. *Sutro.*

CROCUS.

SAFFRON.

Saffron is the stigmas of *Crocus sativus*, a bulbous-rooted plant, native of Greece, Asia Minor, &c., and extensively cultivated in many parts of Europe.

Sex. Syst. Triand. monog. *Nat. Syst.* Iridaceæ.

Linn. Sp. Pl. 50. Griffith, Med. Bot. 626.

Saffron has a peculiar, sweetish, fragrant odor, a warm, bitter taste, and is of an orange red color. It was at one time much used as an excitant, narcotic aromatic and emmenagogue, but is seldom employed in this country, except as a coloring ingredient in compound preparations. Dose, from ten to thirty grains.

PILLS OF SAFFRON.

R. Saffron, } each, one drachm.
 Myrrh, }
 Sulphur, }
 Inspissated bile, sufficient.
 Mix, and form one hundred and fifty pills.
 Dose, twelve a-day, as an emmenagogue.

Phæbus.

TINCTURE OF SAFFRON.

R. Saffron, cut fine, two ounces.
 Proof spirit, two pints.
 Digest for fourteen days, express, and filter.
 Dose, one to two fl. drachms.

Ed. Ph.

COMPOUND TINCTURE OF SAFFRON.

R. Saffron, one ounce.
 Myrrh, two ounces.
 Socotrine aloes, ten ounces.
 Cinnamon, } each, half an ounce.
 Mace, }
 Nutmeg, }
 Orange-flower water, one pint.
 Alcohol, sixteen pints.
 Digest for two days, and distil off eight pints.

Cottureau.

ELIXIR DE GARUS.

R. Compound tincture of saffron, eight pints.
 Syrup of maidenhair, ten pints.
 Mix, color with caramel, dissolved in Orange-flower water, eight fl. ounces.
 Used as a stomachic and carminative.

Foy.

INFUSION OF SAFFRON.

R. Saffron, two drachms.
 Boiling water, two pints.
 Infuse, and strain. Advised in debility of the digestive organs.

Beral.

SYRUP OF SAFFRON.

R. Saffron, five drachms.
 Boiling water, one pint.
 White sugar, three pounds.
 Alcohol, two and a half fl. ounces.
 Macerate the saffron in the water for twelve hours, strain, and add the sugar. When the syrup is cool add the spirit.
 Slightly stimulant.

Lond. Ph.

SAFFRON COLLYRIUM.

R. Saffron, one part.
 Boiling water, two hundred parts.
 Infuse, strain, and add Tincture of opium, one part.

Phæbus.

SAFFRON OINTMENT.

R. Cut saffron, one scruple.
 Camphor, two scruples.
 Oil of henbane, two drachms.
 Goulard's cerate, six drachms.
 Mix. As an application to painful hemorrhoids.

Spielmann.

CUBEBA.

CUBEBS.

Cubebs are the berries or fruit of *Piper cubeba*, a climbing perennial vine, native of many parts of the East Indies.

Sex. Syst. Diand. trigyn. *Nat. Syst.* Piperaceæ.

Linn. Sp. Pl. 90. *Griffith, Med. Bot.* 567.

Cubebs somewhat resemble black pepper, but are of a lighter color, and are furnished with a short stalk. The shell is thin and hard, and the seed round, white, and oily. The taste is acrid, peppery, and camphoraceous; the odor aromatic and pleasant. Cubebs are stimulant, with a special action on the genito-urinary organs, and have been much used in gonorrhœa, leucorrhœa, &c., and have also been found useful in piles, bronchitis, &c. The dose of the powder is from ten grains to three drachms, according to circumstances.

POWDER OF CUBEBS.

R. Powdered cubebs, half a drachm to one drachm.
 " ipecacuanha, eighth of a grain.
 Oil of peppermint, half a drop.
 Mix. To be taken four times a-day, in gonorrhœa.

Schmidt.

POWDER OF CUBEBS AND ERGOT.

R. Powdered ergot, two scruples.
 " cubebs, one ounce.
 " cinnamon, half a drachm.
 " sugar, one drachm.

Mix, and divide into eight powders. One to be given three or four times a-day. In gonorrhœa, gleet, leucorrhœa, &c.

Ryan.

POWDER OF CUBEBS AND HEMLOCK.

R. Powdered cubebs, half a drachm to a drachm.
 Extract of hemlock, half a grain.
 Prepared oyster shell, five grains.
 Mix. To be taken four times a-day in gonorrhœa.

Phæbus.

POWDER OF CUBEBS AND ALUM.

R. Powdered cubebs, two ounces.
 " alum, half an ounce.
 Mix, and divide into nine powders. One to be taken three times a-day.

Ricord.

CUBEBA LOZENGES.

(SPITTA'S LOZENGES.)

- R. Powdered cubebs, two drachms.
 Balsam of tolu, six grains.
 Mix, and add
 Extract of liquorice, one ounce.
 Syrup of balsam of Peru, one drachm.
 Gum Arabic, sufficient.

Rub well together, and form lozenges of ten grains. One of these, permitted gradually to melt in the mouth, alleviates the obstruction in the nose, in coryza. *Spitta.*

ELECTUARY OF CUBEBS.

- R. Powdered cubebs, half an ounce.
 Clarified honey, sufficient.

Mix. Three or four teaspoonfuls a-day, in gonorrhœa, mucous discharges from the bladder, &c. *Radius.*

- R. Powdered cubebs,
 Copaiba, each, two ounces.
 Powdered alum, one ounce.
 Extract of opium, five grains.

Rub together. Dose, one drachm, night and morning, in the pulp of a prune, rapidly increasing the dose to two drachms. In gonorrhœa. *Beral.*

INJECTION OF CUBEBS.

- R. Powdered cubebs, one ounce.
 Water, one pint.
 Boil, and strain. Add
 Extract of belladonna, one scruple.
 In gonorrhœa and leucorrhœa. *Soubiran.*

CLYSTER OF CUBEBS.

- R. Powdered cubebs, one to four drachms.
 Decoction of mallows, six ounces.
 Mix. To be administered night and morning, in gonorrhœa. *Foy.*

FLUID EXTRACT OF CUBEBS.

- R. Powdered cubebs, one pound.
 Sulphuric ether, sufficient.

Introduce the powder into a displacer, insert the lower end into a closely-fitting bottle, add the ether carefully, cover the top of the filter with wet bladder pierced with several pin-holes. The flow should be very gradual. Introduce the tincture into a large retort, and distil, by means of a water-bath, into a well-cooled receiver, till five-sixths have passed over. Continue the evaporation at a heat below 120° F., till the extract is of proper consistence. This is now officinal in the U. S. Ph. The dose is from ten to twenty minims. *W. Procter.*

EMULSION OF FLUID EXTRACT OF CUBEBS.

- R. Fluid extract of cubebs, two drachms.
 Powdered gum Arabic, half an ounce.
 " sugar, one drachm.
 Water, three ounces and a half.
 Mix. A tablespoonful is equal to two drachms of cubebs. *W Procter.*

ALCOHOLIC ETHEREAL EXTRACT OF CUBEBS.

- R. Powdered cubebs, at will.
 Exhaust by ether, by means of a displacement apparatus, act on residue with diluted alcohol, and distil the two tinctures separately. Evaporate the residue of the alcoholic tincture by means of a water-bath; add both together, and let the remaining ether evaporate spontaneously. *Labelonye.*

LOZENGES OF ALCOHOLIC-ETHEREAL EXTRACT OF CUBEBS.

- R. Extract of cubebs, eight ounces.
 Alcohol, two pints.

Dissolve, and add
 Powdered sugar, one pound.
 Oil of peppermint, eighteen drops.

Mix, and allow alcohol to evaporate by a gentle heat; reduce to powder, and add

- Mucilage tragacanth, sufficient.
 Mix, and divide into lozenges of six to eighteen grains. *Labelonye.*

CUBEBA MIXTURE.

- R. Powdered cubebs, two drachms.
 Carbonate of soda, half a drachm.
 Mucilage of gum Arabic, six fl. drachms.
 Mint water, six fl. ounces.

Mix. A tablespoonful every hour. *Fosbroke.*

- R. Powdered cubebs, two drachms.
 Subnitrate of bismuth, half a drachm.
 Mucilage of gum

Arabic, half fl. ounce.
 Syrup, six fl. drachms.
 Water, six fl. ounces.

Mix. A tablespoonful four times a-day. *Fosbroke.*

Both these are recommended in the various affections of mucous membranes.

- R. Powdered cubebs, two drachms.
 Wine, two ounces.
 Essence of bergamot, one drop.

Mix. To be taken every hour or two. *Pierquin.*

TINCTURE OF CUBEBS.

- R. Bruised cubebs, four ounces
 Diluted alcohol, two pints

Macerate for fourteen days, express, and filter.

U. S. Ph.

Dose, one to two drachms, as a stomachic and carminative, or in advanced stage of gonorrhœa.

R. Powdered cubebs, four ounces.
Spirit of nitric ether, two pints.

Digest for eight days and filter. Serviceable in sub-acute inflammations of bladder, uterus, and of the mucous lining of the stomach and intestines.

Mettauer.

SYRUP OF ALCOHOLIC-ETHEREAL EXTRACT OF CUBEBS.

R. Extract of cubebs, three ounces.
Mucilage of gum Arabic, sufficient.
Peppermint water, one pint.
White sugar, two pounds.

Mix. A teaspoonful, several times a-day, in a glass of water.

Labelonye.

OIL OF CUBEBS.

R. Powdered cubebs, one part.
Water, two parts.

Distil, and collect the oil.

Giordano.

Dose, ten to twelve drops, gradually increased. Given in emulsion, or in gelatin capsules.

MIXTURE OF OIL OF CUBEBS AND COPAIBA.

R. Oil of cubebs, one scruple.
Copaiba, two ounces.

Mix. A teaspoonful, four times a-day.

Beral.

CUCUMIS.

CUCUMBER.

This is the fruit of the *Cucumis sativus*, a vine-like plant, cultivated in most parts of the civilized world.

Sex. Syst. Mon. monadelph. *Nat. Syst.* Cucurbitaceæ.

De Cand. Prod. iii. p. 300.

CUCUMBER OINTMENT.

R. Green cucumbers, (fit for the table), seven pounds.
Lard, twenty-four ounces.
Veal suet, fifteen ounces.

Wash and grate the unpared cucumbers, and express the juice. Melt the suet, add the lard, and strain, stirring constantly; as it thickens, add one-third of the juice, and beat with a wooden spatula. The part that separates by standing is decanted, and the other two-thirds are consecutively incorporated, and decanted in the same manner. It is usual to keep the ointment in glass jars, covered with rose water, to prevent access of air.

W. Procter, Jr.

CUNILA.

DITTANY.

The whole herb of *Cunila mariana*. A native plant, found in most places in the United States. *Sex. Syst.* Diand. monog. *Nat. Syst.* Lamiales.

Linn. Sp. Pl. 30. *Griffith, Med. Bot.* 509.

It has a powerful, aromatic odor, and a warm, pungent taste. It is stimulant, carminative, sudorific, and emmenagogue.

INFUSION OF DITTANY.

R. Dittany, half an ounce.
Boiling water, one pint.

Infuse and strain. Used warm, as a diaphoretic and emmenagogue.

OIL OF DITTANY.

R. Dittany, at will.
Water, sufficient.

Distil, and separate the oil. As a carminative, in doses of two or three drops, with sugar.

CUPRUM.

COPPER.

A brilliant, ductile, malleable metal, of a reddish color, having an unpleasant taste and smell. It is not used in its pure state in medicine, but its salts afford numerous remedial preparations.

CUPRI ACETAS.

ACETATE OF COPPER.

R. Pulverized verdigris, at will.
Acetic acid, sufficient.

Dissolve, filter, evaporate, and crystallize.

Van Mons.

CUPRI SUBACETAS PRÆPARATUM.

VERDIGRIS.

PREPARED SUBACETATE OF COPPER.

R. Verdigris, in powder, at will.
Reduce to powder by trituration in a mortar, and separate the finer parts for use by a sieve.

Dub. Ph.

POWDER OF VERDIGRIS AND SAVINE.

R. Verdigris, equal parts.
Savine, equal parts.

Mix. To be dusted on sores with fungous excrescences.

Ellis.

PILLS OF ACETATE OF COPPER.

R. Acetate of copper, fifteen grains.
Boiling water, sufficient
to dissolve; add
Opium, five grains.
Extract of liquorice, one drachm.
Powdered liquorice, sufficient.
Mix, and make one hundred and eighty pills.
Dose, three to ten, three times a-day. *Phabus.*

VERDIGRIS AND ALUM.

R. Verdigris, }
Nitro, } each, eight parts.
Alum, }
Melt together, and add
Camphor, one to two parts.
Much used in Europe as a basis for ophthalmic
solutions, under the name of *Beer's divine stone*.
Radiis.

OPHTHALMIC WASHES.

R. Beer's divine
stone, eight to twelve grains.
Water, four fl. ounces and a half.
Wine of opium,
a fl. scruple to one fl. drachm.
Mix. In chronic ophthalmia. *Benedict.*
R. Beer's divine stone,
three to four grains.
Elder water, one fl. ounce.
Tincture of opium, twelve drops.
Solution of subacetate of lead,
five drops.
Mix. Used like the last. *Rust.*

COMPOUND VERDIGRIS LOTION.

R. Verdigris,
Burnt alum, each, two drachms.
Honey, half an ounce.
White wine, one pint.
Mix. As a wash for indolent venereal or
scorbutic ulcers. *Scherf.*

METZ'S BALSAM.

R. Linseed oil,
Olive oil, each, six ounces.
Oil of laurel berries, one ounce.
Turpentine, two ounces.
Melt by a gentle heat, and add
Powdered aloes, two drachms.
" verdigris, three drachms.
" white vitriol,
one drachm and a half.
Pour into a bottle, and add
Oil of juniper, half an ounce.
" cloves, one drachm.
Mix. As a dressing to wounds and ulcers.
Guibourt.

VERDIGRIS OINTMENT.

R. Verdigris, in fine powder, one drachm.
Simple ointment, fifteen drachms.
Melt the ointment, and add the verdigris, and
stir till cold. *U. S. Ph.*
Used as a mild escharotic and stimulant to
fungous ulcers, to warts, corns, &c., and obsti-
nate cutaneous affections.
R. Verdigris, two drachms.
Oxide of zinc,
Camphor, dissolved in
alcohol, each, six drachms.
Triturate well, and incorporate with a melted
mixture of
Lard,
Suet, each, two ounces.
and stir till cold.
Recommended in scrofulous ophthalmia.
Swediaur.

VERDIGRIS PLASTER.

R. Galbanum, one ounce.
Pitch, half an ounce.
Litharge plaster, two drachms.
Melt, and add
Verdigris,
Muriate of ammonia, each, one ounce.
This is a good application to corns and warts.
Saxon Ph.

EGYPTIAN OINTMENT.

R. Verdigris, five parts.
Purified honey, sixteen parts.
Strong vinegar, seven parts.
Burnt alum, half a part.
Mix, and melt by a gentle heat, stirring con-
stantly. This is thought to form an excellent
detergent application to ulcers. *Giordano.*

LINIMENT OF VERDIGRIS.

R. Powdered verdigris, one ounce.
Vinegar, seven fl. ounces.
Honey, fourteen ounces.
Dissolve the verdigris in the vinegar, strain,
gradually add the honey, and boil down to
proper consistence. *Lond. Ph.*
Used like the last, and also much diluted as
a gargle in venereal sore throat.

CUPRI MURIAS.

MURIATE OF COPPER.

R. Chloride of potassium, seven parts
Sulphate of
copper, eleven and a half parts

Pulverize, mix, and gradually add twelve parts boiling water; on cooling, filter, separate the sulphate of potassa that forms, then permit the crystals of the muriate to form, separate, and dry.

Van Mons.

CUPREOUS ETHER.

R. Chloride of barium, ten parts.
Sulphate of copper, twelve parts.
Sulphuric ether, six parts.

Triturate the salts together, add the ether, and decant when all the muriate of copper is dissolved. Has been recommended in small doses in epilepsy.

Van Mons.

MURIATE OF COPPER AND AMMONIA.

R. Muriate of copper,
" ammonia, equal parts.

Dissolve in water, add liquid ammonia, drop by drop, as long as any precipitate takes place, and is again dissolved by the addition of more ammonia; filter, and evaporate.

Augustin.

Used in epilepsy, in doses of two to ten grains.

SOLUTION OF AMMONIACAL MURIATE OF COPPER AND MERCURY.

R. Copper filings, two drachms.
Calomel, two ounces.

Dissolve the copper in two fl. ounces liquid ammonia, and the calomel in two fl. ounces muriatic acid, with fifteen drops of nitric acid; mix the solutions in such proportions that the precipitate that first forms is re-dissolved.

Radius.

KOECHLIN'S DROPS.

R. Solution of ammoniacal
muriate of copper and
mercury, two fl. drachms.
Distilled water, twenty fl. ounces.

Mix. Much praised in obstinate venereal affections, scrofula, &c. Dose, a teaspoonful after each meal.

Augustin.

CUPRUM AMMONIATUM.

AMMONIATED COPPER.

R. Sulphate of copper, half an ounce.
Carbonate of ammonia, six drachms.

Rub together in a glass mortar till effervescence ceases, wrap in bibulous paper, and dry with a gentle heat.

U. S. Ph.

Has been used in epilepsy, chorea, hysteria, &c. Dose, a quarter to half a grain twice a-day, and gradually increased.

SOLUTION OF AMMONIATED COPPER.

R. Ammoniated copper, one drachm.
Distilled water, one pint.

Dissolve and filter.

Lond. Ph.

As a lotion to foul and indolent ulcers, or internally, in same cases as the powder.

INJECTION OF AMMONIATED COPPER.

R. Ammoniated copper, five grains.
Rose water, eight fl. ounces.
Mix. In gonorrhœa.

Ellis.

POWDER OF AMMONIATED COPPER AND BELLADONNA.

R. Ammoniated copper,
Powdered belladonna
root,
Sugar, two to four grains.
one to four grains.
one drachm.

Mix, and form six powders, one to be taken every two hours, in epilepsy.

Hildebrand.

PILLS OF AMMONIATED COPPER.

R. Ammoniated copper, two grains.
Crumb of bread, sufficient.

Mix, and form four pills. One to be taken twice a-day, in epilepsy, gradually increasing the dose.

A. T. Thomson.

R. Ammoniated copper, fifteen grains.
Crumb of bread, two scruples.
Sugar, one scruple.
Liquid ammonia, sufficient.

Mix, and form thirty pills. Dose, one, three times a-day, gradually increased.

Van Mons.

COMPOUND PILLS OF AMMONIATED COPPER.

R. Ammoniated copper,
Opium, each, ten grains.
Extract of dandelion,
Powdered mallow, each, two scruples.

Mix, and form fifty pills. Dose, five, twice a-day, in diabetes.

Radius.

GARGLE OF AMMONIATED COPPER.

R. Ammoniated copper, eight grains.
Savine water, six fl. ounces.

Mix. In chronic sore throat.

Kopp.

ointment of AMMONIATED COPPER.

R. Solution of ammoniated
copper, one fl. drachm.
Simple cerate, melted, one ounce.

Mix well. As a stimulant to indolent ulcers.

Swedjaur.

CUPRI SULPHAS.

SULPHATE OF COPPER.

POWDER OF SULPHATE OF COPPER.

R. Sulphate of copper, twelve grains.
Sugar of milk, four scruples.

Mix, and divide into four powders. Much praised in croup—given as follows: one is to be administered at once; another dissolved in three spoonfuls of water, one of which is to be taken every hour. *Radius.*

R. Sulphate of copper, four grains.
Sugar of milk, eight scruples.

Mix, and divide into eight powders. Two a-day in obstinate diarrhœa. *Elliotson.*

COMPOUND POWDER OF SULPHATE OF COPPER.

R. Sulphate of copper, }
" zinc, } equal parts.
Alum, }
Carbonate of lead, }
Armenian bole, }

Melt the first three substances together, pulverize, add the other two articles, and sift.

This is a powerful astringent, and has proved useful as a styptic. *Cottureau.*

PILLS OF SULPHATE OF COPPER.

R. Sulphate of copper, four grains.
Extract of cinchona, thirty-two grains.

Mix, make mass, and divide into sixteen or twenty pills. One to be taken three times a-day, in obstinate intermittents. *Ellis.*

R. Sulphate of copper, one drachm.
Ipecacuanha, each, sufficient.
Syrup, sufficient.

Form mass, and divide into pills of five grains each. Dose, two to four, every two or three days, in the morning, before eating, in phthisis. *Foy.*

R. Sulphate of copper, six grains.
Powdered calamus, two drachms.
Extract of liquorice, sufficient.
Water, each, sufficient.

Form mass, and divide into ninety-six pills. Four to seven, two or three times, in the apyrexia of obstinate intermittents. *Adair.*

PILLS OF SULPHATE OF COPPER AND OPIUM.

R. Sulphate of copper, two grains.
Opium, four grains.
Conserve of roses, sufficient.

Mix, and make sixteen pills. One, three times a-day, in obstinate intermittents. *Chapman.*

COMPOUND PILLS OF SULPHATE OF COPPER.

R. Sulphate of copper, two grains.
Alcohol. ext. cascarilla,

seventeen grains.

Opium, two grains

Syrup of ginger, sufficient.

Mix, and make eight pills. One, three times a-day, in epilepsy. *Ainslie.*

ELECTUARY WITH SULPHATE OF COPPER.

R. Sulphate of copper, one scruple.
Opium, one grain.

Armenian bole,

Catechu, each, one drachm and a half.

Syrup, sufficient.

Mix. In obstinate diarrhœa. *Saunders.*

GARGLE WITH SULPHATE OF COPPER.

R. Sulphate of copper, twenty grains.

Infusion of sage, six fl. ounces.

Tincture of myrrh,

" catechu,

" kino, each, one drachm.

" pimperella, five drachms.

Honey, six drachms.

Mix well. In obstinate salivation. *Kopp.*

COLLYRIUM OF SULPHATE OF COPPER.

R. Sulphate of copper,

Armenian bole, each, eight grains.

Camphor, two grains.

Boiling water, eight fl. ounces

Rub well together, permit to settle, and strain. As an application in purulent ophthalmia of infants. *Ware.*

R. Sulphate of copper, six grains.

Camphor, one drachm.

Boiling water, eight fl. ounces.

Rub the camphor with the water, strain, and add sulphate of copper. As a substitute for the above. *Ellis.*

SULPHATE OF COPPER LOTION.

R. Sulphate of copper, one ounce and a half.

Alum, five drachms and a half.

Sage leaves, five ounces.

Vinegar, one pint

Solution of muriate of

ammonia, two pints

Boil together for half an hour. Used as an application, in a tepid state, to swellings and contusions. *Purmann*

INJECTION OF SULPHATE OF COPPER.

R. Sulphate of copper, six grains.
 Distilled water, six fl. ounces.
 Tincture of opium, one fl. drachm.

Dissolve. As an injection in chronic gonorrhoea. *Ellis.*

R. Sulphate of copper, six grains.
 Distilled water, four fl. ounces.

Dissolve, and add
 Solution subacetate of lead, twenty drops.

As a wash and injection in phymosis. *Swediaur.*

OINTMENT OF SULPHATE OF COPPER.

R. Sulphate of copper,
 Calamine, each, five grains.
 Camphor, two grains.
 Fresh butter, two drachms.

Triturate well together. A small portion to be applied to the edges of the eyelids, in the evening, in psorophthalmia. *Rust.*

R. Sulphate of copper,
 Verdigris, each, two drachms.
 Alum, half an ounce.
 Corrosive sublimate, two scruples.
 Lard, one ounce.
 Burgundy pitch, one pound.

Melt the pitch and lard by a gentle heat, and add the other ingredients, stirring till cold. As a dressing to venereal and fungous ulcers. *Augustin.*

R. Powdered sulphate of copper,
 " catechu, each, four drachms.
 " alum, nine drachms.
 " resin, four ounces.

Olive oil, sufficient.

Make ointment. As an application to indolent and ill-conditioned ulcers. *Kerr.*

SULPHATE OF COPPER STYPTIC.

R. Sulphate of copper, three grains.
 Sulphuric acid, twenty drops.
 Water, two ounces.

Dissolve. In epistaxis. Twenty to forty drops to be taken in water, every hour. *Thatcher.*

CYDONIUM.

QUINCE SEEDS.

These seeds are the product of *Cydonia vulgaris*, a small tree, native of some parts of Europe, but extensively cultivated in this country. The fruit is much used for the purpose of making preserves.

Sex. Syst. Icosand. pentag. *Nat. Syst.* Poma-ceæ

Persoon, Enchirid. ii. 40. Griffith, Med. Bot. 291.

The seeds are inodorous, insipid, and abounding with mucilage; one drachm rendering six ounces of water viscid. Used like the other bland mucilages.

DECOCTION OF QUINCE SEEDS.

R. Quince seeds, two drachms.
 Distilled water, one pint.

Boil over a slow fire, for ten minutes, and strain. *Lond. Ph.*

R. Quince seeds, one ounce.
 Boiling water, six fl. ounces.

Digest on hot coals, for two hours; strain.

Both these are used in ophthalmia, &c.

QUINCE SEED MIXTURE.

R. Mucilage of quince seeds, one ounce.
 Yolks of eggs, two.

Honey of roses, three ounces.

Mix. A teaspoonful occasionally, in cough and hoarseness. *Sainte Marie.*

BANDOLINE FOR THE HAIR.

R. Mucilage of quince seeds, eight fl. ounces.

Cologne water, or Brandy, eight fl. ounces.

Mix.

Used as an application to the hair, to give gloss and smoothness.

CYMINUM.

CUMIN.

Cumin seeds are produced by *Cuminum cyminum*, an umbelliferous plant, a native of Egypt, but cultivated in southern Europe.

Sex. Syst. Pentand. digyn. *Nat. Syst.* Apiaceæ.

Linn. Sp. Pl. 365. *Lindley, Fl. Med.* 51.

These seeds have a peculiar, strong, heavy odor, and a warm, bitterish, aromatic taste. They are carminative and stimulant, are seldom used internally, but enter into the composition of some warm plasters.

CUMIN PLASTER.

R. Cumin,
 Caraway,
 Laurel berries, each, three ounces.
 Burgundy pitch, three pounds.
 Yellow wax, three ounces.
 Olive oil,

Water, each, one fluidounce and a half.

Melt the pitch and wax together, add the other ingredients, stirring well, and evaporate to a proper consistence. *Lond. Ph.*

D.

DELPHINIUM.

LARKSPUR.

The *Delphinium consolida* is a native of Europe, but has become partially naturalized in this country, and is also much cultivated in gardens as an ornamental flower.

Sex. Syst. Polyand. trigyn. *Nat. Syst.* Ranunculaceæ.

Linn. Sp. Pl. 748. Griffith, Med. Bot. 88.

Several parts of this plant have been employed medicinally, as the flowers, seeds, and root. They owe their properties to the presence of an alkaloid, called delphinia. The flowers are said to be diuretic, vermifuge, and emmenagogue; and the seeds and root to be diuretic, &c.

TINCTURE OF LARKSPUR SEEDS.

R. Larkspur seeds, one ounce.
Diluted alcohol, one pint.
Macerate for some days, and filter. Dose, ten to twenty drops, three times a-day, in spasmodic asthma. *Augustin.*

DELPHINIA.

DELPHINIA.

This alkaloid is prepared from the seeds of different species of *Delphinium*, but generally from those of the *D. staphisagria*.

R. Larkspur seeds, at will.
Water, sufficient.

Boil, repeat the decoction with another portion of water, till all soluble portions of the seeds are dissolved. Concentrate the united decoctions, add calcined magnesia, and filter after a short ebullition. Wash the precipitate with cold water, and dry it; digest it in alcohol on a water-bath, and permit to crystallize.

Magendie.

This is used in the same cases as veratria, in doses of a quarter to half a grain, to the extent of two to three grains a-day. It is also employed in ointment, or in solution in alcohol.

SOLUTION OF DELPHINIA.

R. Delphinia, one scruple.
Rectified spirit, two fl. ounces.
Dissolve. For outward use. *Turnbull.*

OINTMENT OF DELPHINIA.

R. Delphinia, ten to thirty grains.
Olive oil, one drachm.
Rub together, and add
Lard, one ounce.
Mix well. *Turnbull.*

DIANTHUS.

PINK.

The only species that is officinal is *D. caryophyllus* or clove pink, a native of the south of Europe, but generally cultivated in gardens, for the beauty and fragrance of its flowers.

Sex. Syst. Decand. digyn. *Nat. Syst.* Caryophyllaceæ.

Linn. Sp. Pl. 587. Woodville, Med. Bot. i. 205.

The parts used are the flowers: these should be of a dark red color, and very aromatic. They are principally employed to form a syrup, which is used as a vehicle for other medicines.

SYRUP OF CLOVE PINK.

R. Petals of clove pink, one pound.
Boiling water, two pints.
Infuse for twelve hours, strain, and add twice the weight of sugar. *Guibourt.*

DIGITALIS.

FOXGLOVE.

The foxglove is a native of Europe, but is cultivated in this country, both as an ornamental plant and for medicinal purposes.

Sex. Syst. Didynam. angios. *Nat. Syst.* Scrophulariaceæ.

Linn. Sp. Pl. 868. Griffith, Med. Bot. 520.

The parts used are the leaves: these should be dried in the dark, and always kept from the light. When good, they are of a dull green color, a feeble, narcotic odor, and a bitter, unpleasant taste. Digitalis is narcotic, sedative, and diuretic, and, in large doses, poisonous. It is given to fulfil many indications; as a diuretic, as a sedative, and antispasmodic, to reduce the force of the circulation, &c. It is given in substance in the dose of one grain, two or three times a-day, gradually increasing the quantity until some effect is produced on the system.

POWDER OF FOXGLOVE.

R. Powdered foxglove, one grain.
" cinnamon, four grains.
" sugar, ten grains.

Mix. For a single dose, to be repeated two or three times a-day. *Radius.*

R. Foxglove, a quarter to one grain.
Sulphate of
quinia, a half to two grains.
Fennel seed, six grains.
Sugar of milk, ten grains.

Mix. To be taken three or four times a-day, in phthisis. *Gunther.*

- R. Powdered foxglove, ten grains.
 " calomel, six grains.
 " valerian, two scruples.
 " assafetida,
 " castor, each, half a drachm.
 " sugar, one drachm.

Mix, and divide into twenty-four powders. One, morning and evening, in the convulsions attendant on hydrocephalus. *Pierquin.*

- R. Powdered foxglove, fifteen grains.
 " nitre, three drachms.
 " cremor tartar, half an ounce.

Mix, and divide into six powders. One to be taken every two hours. As a diuretic in dropsies. *Brera.*

- R. Powdered foxglove, six grains.
 Golden sulphuret of antimony,
 Calomel, each, three grains.
 Sugar, one drachm.

Mix, and divide into six powders. One, every three hours in peripneumonia. *Radius.*

- R. Powdered foxglove, eight grains.
 Extract of opium, four grains.
 Sugar, one ounce.
 Oil of peppermint, four drops.

Mix, and divide into eight powders. One in the evening, in palpitation of the heart. *Sainte Marie.*

EXTRACT OF FOXGLOVE.

- R. Fresh juice of foxglove, at will.
 Evaporate to proper consistence, without removing fecula. *Lond. Ph. 1836.*
 Dose, from half a grain to two grains.

BOLUS OF FOXGLOVE.

- R. Powdered foxglove, fifteen grains.
 Calomel, eight grains.
 Root of juniper, one drachm.
 Mix, and form four boluses. One, every four hours in encephalitis. *Brera.*

- R. Powdered foxglove, twelve grains.
 Calomel, six grains.
 Tartar emetic, two grains.
 Purified honey,
 Powdered liquorice root, each,
 sufficient.

Mix, and form four boluses. One, every four hours, in hemoptysis and engorgement of the abdominal viscera. *Brera.*

PILLS OF FOXGLOVE.

- R. Powdered foxglove, twelve grains.
 Calomel, six grains.
 Opium, two grains.
 Conserve of roses, sufficient.

Mix, and form twelve pills. One to be taken every eight hours, in hydrothorax and ascites. *Ellis.*

- R. Powdered foxglove, four grains.
 " camphor, twelve grains.
 Extract of henbane, eighteen grains.

Mix, and make six pills. Two to be taken at bedtime. Used in maniacal and spasmodic affections. *A. T. Thomson.*

- R. Powdered squill, four grains.
 " myrrh, one scruple.
 " foxglove, ten grains.
 Calomel, six grains.

Triturate together, and add

- Assafetida, half a drachm.
 Extract of gentian, sufficient.

Beat into mass, and divide into fifteen pills. One night and morning, as a diuretic in dropsies. *Paris.*

- R. Powdered foxglove,
 " opium, each, six grains.
 Conserve of roses, sufficient.

Mix, and make twelve pills. One to be taken every four hours. In asthma, &c. *Ellis.*

- R. Powdered foxglove, one drachm.
 Extract of hemlock,
 " rhubarb,
 Aloes, each, half a drachm.
 Ammoniac, four scruples.
 Oxymel of squill,
 Powdered liquorice root,
 each, sufficient.

Mix, and make pills of six grains. Three to be taken morning and evening. In dropsy, &c. *Brera.*

PILLS OF FOXGLOVE AND SQUILL.

- R. Powdered foxglove,
 " squill, each, one part.
 Aromatic electuary, two parts.
 Confection of roses, sufficient.

Beat them into a mass, and divide into pills of four grains each. *Ed. Ph.*

Dose, one or two pills.

INFUSION OF FOXGLOVE.

- R. Foxglove leaves, one drachm.
 Boiling water, half a pint.
 Tincture of cinnamon, one fl. ounce.

Macerate the foxglove with the water, in a covered vessel, for two hours; strain, and add the tincture. *U. S. Ph.*

The dose is half a fl. ounce, twice a-day, till the system is affected.

MIXTURE OF FOXGLOVE AND ACETATE OF POTASSA.

R. Infusion of foxglove, four fl. ounces.
Tincture of foxglove, one fl. drachm.
Acetate of potassa, one drachm.
Tincture of opium, ten drops.
Mix. A dessertspoonful three or four times a-day; as a diuretic. *Ellis.*

MIXTURE OF FOXGLOVE AND ACETATE OF LEAD.

R. Powdered foxglove, three to five grains.
Solution of subacetate of lead, five drops.
Syrup of orange flowers, one ounce.
Infusion of poppies, six ounces.
Mix. In spoonful doses, in hypertrophy of the heart. *Foy.*

MIXTURE OF FOXGLOVE AND TARTARIC ACID.

R. Infusion of foxglove, half an ounce.
Tartaric acid, one scruple.
Carbonate of soda, twenty-four grains.
Sweet spirit of nitre, one drachm.
Tincture of squill, four drops.
Mint water, two ounces.
Mix. As a diuretic, in ascites. To be taken twice or thrice a-day. *Dewees.*

EXPECTORANT MIXTURE OF FOXGLOVE.

R. Foxglove leaves, thirty grains.
Boiling water, sufficient
to obtain four ounces of strained infusion; add
Syrup of gum Arabic, three ounces.
Kermes mineral, six grains.
Syrup of mallows, one ounce.
Mix. To be taken, in teaspoonful doses, in pneumonia and pleurisy. *Brera.*

TINCTURE OF FOXGLOVE.

R. Foxglove leaves, four ounces.
Diluted alcohol, two pints.
Macerate for fourteen days, and filter.
U. S. Ph.
Dose, from ten to twenty drops, two or three times a-day.

MIXTURE OF TINCTURE OF FOXGLOVE.

R. Tincture of foxglove, one fl. drachm.
" opium, sixty drops.
Distilled water, two fl. ounces.
Mix. A teaspoonful, two or three times a-day, in hemoptysis and incipient phthisis. *Ellis.*

ETHEREAL TINCTURE OF FOXGLOVE.

R. Foxglove leaves, one part.
Sulphuric ether, four parts.
Macerate for two days, and decant. Dose, from ten to twenty drops. *Soubairan.*
R. Foxglove leaves, one ounce and a half.
Spirit of nitric ether, two pints.
Digest for ten days, and filter. Dose, twenty drops to a fluidrachm, in some diuretic infusion, in dropsies. *Mettauer.*

VINEGAR OF FOXGLOVE.

R. Foxglove leaves, half an ounce.
Vinegar, four fl. ounces.
Infuse for twenty-four hours, strain, and add
Sugar, five ounces.
A teaspoonful, several times a-day. Highly spoken of in incipient phthisis. *Martius.*

OINTMENT OF FOXGLOVE.

R. Bruised foxglove leaves, one part.
Lard, two parts.
Melt over a slow fire, until all moisture is driven off, and strain. As an application to chronic ulcers. *Soubairan.*

PLASTER OF FOXGLOVE.

R. Foxglove leaves, two drachms.
Vinegar, one fl. ounce.
Macerate, and evaporate to consistence of an extract, and add
Calomel, ten grains.
Elemi ointment, eight scruples.
Mix. In chronic glandular swellings. *Sundelin.*

SYRUP OF FOXGLOVE.

R. Foxglove leaves, four ounces.
Water, sufficient.
Sugar, one pound.
Bruise the foxglove, add sufficient water to render quite moist, exhaust, by process of displacement, evaporate in a water-bath to ten ounces, add sugar, and form syrup. One teaspoonful is equivalent to fifteen grains of digitalis, or about two ounces of infusion. *Duhamel.*

DIGITALINA.

DIGITALIN.

R. Powdered foxglove, at will.
Macerate for twenty-four hours in a sufficiency of sulphuric ether, decant, repeat the operation several times, the last on a water-bath. Unite, and filter the tinctures, and distil off most of the ether, over a water-bath. Treat the residue with distilled water. Add, very gradually finely-powdered litharge, till there is no acid

reaction. Evaporate to dryness, treat with sulphuric ether, and evaporate to crystallizing point. *Magendie.*

Dose, from one-fortieth to one-twentieth of a grain.

PILLS OF DIGITALIN.

R. Digitalin, three quarters of a grain.
Powdered gum Arabic,
Mucilage of gum Arabic, each,
sufficient.

Mix, and make twenty pills. Dose, one to four daily, in hypertrophy of the heart. *Bouchardat.*

R. Digitalin, three quarters of a grain.
Powdered squill,
“ scammony, each, seventy-five grains.

Syrup of gum Arabic, sufficient.

Rub well together, and divide into one hundred pills. Give two pills, then four, and afterwards six daily, in dropsy with disordered circulation. *Falken.*

GRANULES OF DIGITALIN.

R. Digitalin, fifteen and a half grains.
Sugar, one ounce and a half.
Water, sufficient.

Mix, and form one thousand granules. Dose, from four to six a-day. *Homolle.*

DIOSPYROS.

PERSIMMON.

The Persimmon, or *Diospyros Virginiana*, is a native tree, found in the middle and southern States; most frequently in the latter.

Sex. Syst. Diœc. octand. *Nat. Syst.* Ebenaceæ.

Linn. Sp. Pl. 1510. Griffith, Med. Bot. 435.

Several parts of the Persimmon have been used in medicine, as the bark and the unripe fruit. They are both powerful astringents, and have been employed with success in bowel complaints and hemorrhages. The bark may be given in infusion; the fruit in syrup or vinous tincture.

INFUSION OF PERSIMMON BARK.

R. Persimmon bark,
bruised, half an ounce.
Boiling water, one pint.

Macerate for two hours, and strain.

Used in doses of a fl. ounce, in intermittents, and as a gargle in ulcerated sore throat.

WINE OF PERSIMMONS.

R. Crushed green persimmons, one pound.
Port wine, one pint and a half.

Macerate for fourteen days, strain, and filter. Dose, two to four fl. drachms a-day, as an astringent. *Beasley.*

DIRCA.

LEATHERWOOD.

The *Dirca palustris* is an indigenous shrub, found in boggy situations, with long, tough roots.

Sex. Syst. Octand. monog. *Nat. Syst.* Thymelacææ.

Linn. Amœn. iii. 12. Griffith, Med. Bot. 561.

The part used is the bark, which is very fibrous; it causes redness and vesication, when applied to the skin; when chewed, it induces salivation. It has been given in infusion, or decoction, as a sudorific, and expectorant, and also in the same class of diseases as mezereon.

DRACONTIUM.

SKUNK CABBAGE.

A perennial plant, with very large leaves, indigenous to the United States, growing in wet situations. It is the *Symplocarpus foetidus*.

Sex. Syst. Tetrand. monog. *Nat. Syst.* Oroniaceæ.

Linn. Sp. Pl. 1372 (Dracontium). Griffith, Med. Bot. 619.

The root is officinal, but the seeds are equally powerful. They both have a very acrid taste when fresh. The leaves, when bruised, have a nauseous smell, and are a good stimulating application to blisters. The root and seeds are stimulant, antispasmodic, and narcotic; and have been employed with success in asthma, hysteria, &c. The dose in powder is from two to twenty grains, several times a-day.

INFUSION OF SKUNK CABBAGE-ROOT.

R. Skunk cabbage root, one ounce.
Boiling water, one pint.

Infuse in a covered vessel for an hour, and strain. Dose, an ounce to two ounces.

TINCTURE OF SKUNK CABBAGE ROOT.

R. Skunk cabbage root, sliced, one ounce.
Alcohol, six fl. ounces.

Macerate for fourteen days, and strain. *Turner.*

TINCTURE OF SKUNK CABBAGE SEED.

R. Skunk cabbage seed, two drachms.
Alcohol, four fl. ounces.

Macerate for fourteen days, and strain. *Turner.*

DULCAMARA.

BITTERSWEET.

The Bittersweet, or *Solanum dulcamara*, is a climbing shrub, indigenous to Europe, and naturalized extensively in this country.

Sex. Syst. Pentand. monog. *Nat. Syst.* Solanaceæ.

Linn. Sp. Pl. 264. Griffith, Med. Bot. 480.

The officinal portions are the small stems. These have a heavy, unpleasant odor, when fresh, but are scentless when dry. They have a somewhat bitter taste, followed by a peculiar sweetness. The properties of Bittersweet are those of a narcotic, diuretic, and diaphoretic. It has been used in chronic rheumatism, asthma, &c., and especially in chronic cutaneous disorders. The dose, in substance, is from thirty grains to a drachm; but it is seldom given in this form.

DECOCTION OF BITTERSWEET.

R. Bittersweet, bruised, one ounce.
Water, one pint and a half.

Boil down to a pint, and strain. *U. S. Ph.*

Dose, from one to two fl. ounces, three or four times a-day, gradually increased, till a pint is taken in the twenty-four hours. In chronic eruptions, &c.

COMPOUND DECOCTION OF BITTERSWEET.

R. Bittersweet, half an ounce.
Liquorice root, }
Burdock, } each, two drachms.
Sassafras bark, }
Guaiacum, }

Water, two pints.
Boil down to sixteen ounces, and strain. Dose, one to two fl. ounces, several times a-day, in chronic rheumatism, and venereal affections.

Augustin.

EXTRACT OF BITTERSWEET.

R. Bittersweet, in coarse powder, one pound.
Water, sufficient.

Mix the powder with a pint of the water, let it stand for twenty-four hours, then put in a displacement apparatus, and add water gradually, till the liquor that passes is but slightly impregnated with the properties of the bittersweet. Heat the filtered fluid, strain, and evaporate to proper consistence.

U. S. Ph.

Dose, from five to ten grains.

R. Freshly-dried bittersweet, eight ounces.
contused, four pints.
Diluted alcohol,

Digest by a gentle heat, for fourteen days, express, filter, and evaporate on water-bath, to a proper consistence. Dose, five to eight grains.

J. W. Simes.

SYRUP OF BITTERSWEET.

R. Coarsely powdered bitter-sweet, four ounces.
Water, twelve fl. ounces.
Alcohol, four fl. ounces.

Mix the fluids, pour on the powder in a displacer, until one pound of tincture is obtained, adding water to displace the diluted alcohol. Evaporate to half a pint, add fifteen ounces of sugar, and make a syrup.

W. Procter.

Dose, a tablespoonful.

PILLS OF EXTRACT OF BITTERSWEET.

R. Extract of bittersweet, one ounce.
Crude antimony,
Powder of bittersweet, each, half an ounce.

Mix, and form pills of two grains each. Dose, from fifteen to twenty, twice a-day, in obstinate cutaneous affections.

Radius.

MIXTURE OF BITTERSWEET.

R. Extract of bittersweet, three drachms.
" seneka, two drachms.
" hemlock, one drachm.

Antimonial wine,

Cinnamon water, each, one fl. ounce.

Mix. Forty to eighty drops, four times a-day, in scrofulous swelling, spasmodic cough, &c.

Augustin.

R. Extract of bittersweet, half a scruple.
Infusion of liquorice,
Decoction of mallows,
each, one ounce and a half.
Wine of opium, two drops.
Oxymel, three drachms.

Mix. A teaspoonful, in chronic catarrh in children.

Radius.

E.

ELATERIUM.

ELATERIUM

Is a peculiar substance deposited by the juice of the fruit of *Ecballium elaterium*, a native of the south of Europe, growing in waste places.

Sex. Syst. Monœc. monad. *Nat. Syst.* Cucurbitaceæ.

Griffith, Med. Bot. 305.

Elaterium is in light, friable, thin, somewhat curled flakes or fragments, of a pale, grayish-green color. The taste is acrid, and bitterish; but the odor is very slight. The dose is from one to two grains, of the common commercial kind; of the pure and genuine, not more than one-eighth to a quarter of a grain.

ELATERIUM PILLS.

R. Elaterium, ten grains.
Extract of gentian, three drachms.

Mix, and form pills of four grains each. One or two, twice a-day, as a hydragogue purgative.
Saunders.

R. Elaterium, six grains.
Extract of gentian, nine grains.
Soap, each, nine grains.

Mix, and form twelve pills. One to four in obstinate constipation.
Radius.

R. Elaterium, half a drachm.
Aloes, two drachms.
Gamboge, each, two drachms.
Ammoniac, two ounces.
Tincture of wormwood, sufficient.

Mix, and form pills of four grains each.
Swed. Ph.

R. Elaterium, fifteen grains.
Sulphate of potassa, one scruple.
Soap, one drachm.
Powdered ginger, one drachm and a half.

Rub the elaterium and sulphate of potassa together, and then with the ginger and soap, adding sufficient water to form mass, and make sixty pills. One or two every hour, until full evacuations are procured.
Sprague.

ELATERIUM MIXTURE.

R. Elaterium, one grain.
Spirit of nitric ether, two fl. ounces.
Tincture of squill, half a fl. ounce.
Oxymel of colchicum, each, half a fl. ounce.
Syrup, one fl. ounce.

Mix. A teaspoonful three or four times a-day, in a little water. In ascites, or hydrothorax.
Ferriar.

R. Elaterium, one grain.
Parsley water, six fl. ounces.
Oxymel of squill, half a fl. ounce.
Syrup of buckthorn, each, half a fl. ounce.
Root of juniper, one ounce.
Aromatic tincture, one drachm.
Spirit of nitric ether, two fl. drachms.

Mix. Used as above, and in the same cases.
Hufeland.

OIL OF ELATERIUM.

R. Fruit of elaterium, one part.
Oil of olives, two parts.

Digest in a water-bath, for some days, then evaporate all moisture, and strain. As an application for sore nipples, frosted limbs, and painful hemorrhoids.
Lond. Ph.

ELATERINA.

ELATERIN.

R. Juice of ecballium elaterium, at will.
Treat with water, dissolve the residue in alcohol, evaporate to consistence of syrup, purify the crystals that form by washing with ether.
Dose, one-sixteenth of a grain.

TINCTURE OF ELATERIN.

R. Elaterin, one grain.
Alcohol, one ounce.
Nitric acid, four drops.

Mix. Dose, twenty to forty drops, in cinnamon water.
Morris.

ELEMI.

ELEMI.

Elemi is a resinous exudation of various species of plants, principally belonging to the natural order of Amyridaceæ. It occurs in masses of various consistence and color, but usually diaphanous; it has a terebinthinate and somewhat aromatic odor, and a warm, unpleasant taste. It has the usual properties of the turpentine, but is only employed as an external application. It is seldom used in this country, but is extensively employed in Europe.

OINTMENT OF ELEMI.

R. Elemi, one pound.
Common turpentine, ten ounces.
Suet, two pounds.
Olive oil, two fl. ounces.

Melt the elemi with the suet; remove from fire, and add turpentine and oil, and strain.

Lond. Ph. 1836.

As a gentle stimulant to flabby ulcers, &c.

R. Elemi,
Venice turpentine,
each, one ounce and a half.
Suet, two ounces.
Balsam of tolu, one ounce.

Melt together, and strain. As a dressing to indolent and ill-conditioned ulcers. *Niemann.*

ELEMI CAUTERY PLASTER.

R. Elemi,
Spermaceti, each, five parts.
Turpentine, six parts.
White wax, ten parts.

Melt together, and spread on paper. To keep up discharge from issues. *Soubéiran.*

EMETIA.

EMETINE.

This is a peculiar alkaloid, found in the various kinds of ipecacuanha, and to which they owe their active properties. There are two varieties, the colored or impure, and the white or pure. The latter is seldom met with in this country.

IMPURE EMETINE.

R. Ipecacuanha, one part.
Alcohol (.835), four parts.

Macerate for some days, express, and filter; treat the residue with three parts of alcohol as above, unite the tinctures, distil, dissolve the residue in four parts of cold water, filter, evaporate to the consistence of honey, and finish the drying by a gentle heat. *Cottéreau.*

It is reddish-brown, inodorous, bitter, and deliquescent. Dose, half a grain, to one grain.

PURE EMETINE.

R. Impure emetine, at will.

Dissolve in water, treat with magnesia, wash with cold water, treat several times with alcohol, on a water-bath, evaporate the solution to dryness, dissolve the residues in very dilute sulphuric or acetic acid, boil with a little animal charcoal, add an alkaline solution to saturate the acid, dry the precipitate, dissolve it in alcohol, and evaporate to dryness. *Cottéreau.*

It is yellowish, or white, pulverulent, not deliquescent. Dose, 1-16th to 1-4th of a grain.

EMETINE MIXTURE.

R. Impure emetine, four grains.
Infusion of orange
leaves, two fl. ounces.
Syrup of orange flowers, half a fl.
ounce.

Mix. A dessertspoonful every half hour, to produce vomiting. *Magendie.*

R. Pure emetine, dissolved in
nitric acid, one grain.
Infusion of linden, three fl. ounces.
Syrup of mallow, one fl. ounce

Mix. Given as the last. *Foy.*

EMETINE LOZENGES.

R. Impure emetine, thirty-two grains.
Sugar, two ounces.
Mucilage of tragacanth, sufficient.

Rub together, and form lozenges of ten grains. One will usually vomit a child, and three or four an adult. *Magendie.*

R. Impure emetine, thirty-two grains.
Sugar, four ounces.
Mucilage of tragacanth, sufficient.

Mix, and form lozenges of nine grains each. One occasionally as an expectorant. *Guibourt.*

SYRUP OF EMETINE.

R. Impure emetine, sixteen grains.
Dissolve in a little water, filter, and add
Syrup, one pound.

R. Pure emetine, four grains.
Syrup, one pound.

Mix. The dose of these syrups is a teaspoonful *Soubéiran.*

ERGOTA.

ERGOT.

Ergot is the diseased seeds of the common rye, but the cause of this morbid growth has given rise to much controversy. The most general opinion, now is, that it is owing to a fungus, but authors are not agreed as to the character of this. The most recent investigator, Mr. Quekett, calls it *Ergotetia arborifaciens*. It is also a question whether the peculiar properties of ergot depend on the fungus, or on the morbid grain. Ergot is one of the few articles that act specifically on the uterus, increasing its contractile energy; and its principal use is as a stimulant to uterine action in labor, and also to check hemorrhage from that organ; but it has been employed in a variety of other diseases, and especially, externally, as a styptic. The dose, in substance, to aid labor, is from fifteen to twenty grains, to be repeated every twenty minutes, till the desired effect is produced, or till a drachm is taken.

INFUSION OF ERGOT.

R. Ergot, one drachm.
Boiling water, four fl. ounces.
Infuse. One-third as a dose, to be repeated every twenty minutes. *Wood.*

R. Ergot, half a drachm.
Boiling water, one fl. ounce and a half.

Infuse for half an hour, and strain. To be repeated every four hours. *Ramsbotham.*

DECOCTION OF ERGOT.

R. Ergot, one ounce.
Water, two pints and a half.
Boil to two pints, and strain. A tablespoonful every quarter of an hour.

WINE OF ERGOT.

R. Bruised ergot, two ounces.
Sherry wine, one pint.
Macerate for fourteen days, express and filter. *U. S. Ph.*

Dose, in labor, two to three fl. drachms; for other purposes, one to two fl. drachms.

SYRUP OF ERGOT.

R. Ergot, one ounce and a half.
White wine, two ounces.
Macerate for eight days, express, and filter. Treat the residue three times with water, mix all the decoctions together, and add

Sugar, one pound.
Evaporate to consistence of syrup, and add the vinous tincture. Dose, one to two fl. ounces. *Foy.*

R. Ergot, twenty grains.
Extract of opium, three-quarters of a grain.
Syrup, eight fl. ounces.
Mix. Dose, one fl. ounce, occasionally, in engorgements of the uterus. *Lisfranc.*

TINCTURE OF ERGOT.

R. Ergot, two ounces and a half.
Diluted alcohol, one pint.
Macerate for fourteen days, and strain. Dose, twenty drops to a fl. drachm. As a sedative. *Guy's Hospital.*

R. Powdered ergot, eight ounces.
Proof spirit, two pints.
Macerate for two weeks, strain, express, and filter. Dose, a teaspoonful. *Dub. Ph.*

R. Powdered ergot, two ounces.
Spirit of nitric ether, one pint.
Digest for ten days and filter. Used in uterine hemorrhage. Dose, one fl. drachm. *Mettauer.*

COMPOUND POWDER OF ERGOT.

R. Powdered ergot, two scruples.
" cubebs, one ounce.
" cinnamon, half a drachm.
" sugar, one drachm.

Mix, and divide into eight powders; one to be taken, three or four times a-day, in leucorrhœa or gleet. *Ryan.*

ERGOT MIXTURE.

R. Powdered ergot, half a drachm.
Syrup, half a fl. ounce.
Mint water, one fl. ounce.

Mix. One-third to be taken at a dose, and repeated every twenty minutes; to aid labor. *Soubeiran.*

R. Powdered ergot, one drachm.
Syrup, three fl. ounces.
Tincture of opium, twenty drops.
Essence of bergamot, six drops.

Mix. A spoonful, to be repeated as may be required. *Pierquin.*

ERGOT CLYSTER.

R. Powdered ergot, one to three drachms.
Water, twelve fl. ounces.
Boil for ten minutes, and strain. As an enema, to aid labor. *Foy.*

COMPOUND ERGOT PILLS.

R. Powdered ergot, half a drachm.
Extract of gentian, one drachm.
Rub together, and make fifteen pills. In dysmenorrhœa. One pill to be taken three times a-day. *Devees.*

ERGOT INJECTION.

R. Powdered ergot, half an ounce.
Boiling water, half a pint.
Infuse, and strain. As an injection in erythema of the vagina and urethritis. *Desruelles.*

EXTRACT OF ERGOT, OR ERGOTINE.

R. Powdered ergot, at will.
Water, sufficient.

Exhaust the ergot by the process of displacement; heat the solution on a water-bath, filter, and evaporate to consistence of syrup; add alcohol to precipitate gummy principles; permit to rest, decant, and evaporate to consistence of soft extract. *Bonjean.*

Said to be very efficacious in hemorrhages, in doses of two grains every two hours; also applied as a styptic to bleeding vessels.

ERGOTINE MIXTURE.

R. Extract of ergot, fifteen grains.
Syrup of orange flowers, one fl. ounce.
Water, three fl. ounces.

Mix. A tablespoonful, every quarter of an hour, in uterine hemorrhage. Dose to be increased, if requisite. *Bonjean.*

PILLS OF ERGOTINE.

R. Extract of ergot, one drachm.
Powdered liquorice root, sufficient.
Mix, and make fifty pills. Dose, six to ten,
a-day. *Bonjean.*

OIL OF ERGOT.

R. Powdered ergot, at will.
Sulphuric ether, sufficient.
Exhaust ergot by process of displacement;
evaporate result by a gentle heat. *Wright.*

Dose, from twenty to fifty drops, in some appropriate vehicle, in parturition, &c. In doses of ten drops, every three hours, in diarrhoea, gastric irritability, &c.

HYDRO-ALCOHOLIC EXTRACT OF ERGOT.

R. Powdered ergot,
Sulphuric ether, equal weights.

Extract the oil from the ergot by the ether in a percolator. Digest the residue with six times its weight of water, for two days, at 167° F. Decant, evaporate to one-fourth, and add alcohol as long as a precipitate takes place. Filter the liquid and evaporate to consistence of honey. Treat the residue of the ergot with its original weight of alcohol, digest for three days, express, and filter. Finally mix the aqueous and alcoholic products, and evaporate to the consistence of a thick extract. *Stickel.*

Dose, five to ten grains.

ERIGERON.

1. ERIGERON CANADENSE.

CANADA FLEABANE.

A common native plant, with numerous white flowers, found in waste places, in the northern and middle States.

Sex. Syst. Syngen. super. *Nat. Syst.* Asteraceæ.

Willd. Sp. Pl. iii. 1954. Torrey & Gray, Fl. ii. 175.

The whole plant is used; it has a rather pleasant smell, and a bitterish, acrid, somewhat astringent taste. It is said to be diuretic, tonic, and astringent, and to be useful in dropsies and diarrhoea. Dose, in substance, thirty grains to a drachm.

INFUSION OF CANADA FLEABANE.

R. Canada fleabane, one ounce.
Boiling water, one pint.
Infuse. Dose, two to four fl. ounces. *Dupuy.*

EXTRACT OF CANADA FLEABANE.

R. Canada fleabane, one pound.
Water, one gallon.

Boil down to four pints, filter, and evaporate to proper consistence. Dose, five to ten grains.

2. ERIGERON ANNUUM.

FLEABANE.

This species, which has also been called *E. heterophyllum*, is very closely allied to, and identical in properties with, *E. strigosum* (*E. Philadelphicum*). It is also a common plant in the United States.

Persoon, Synop. ii. 431. Griffith, Med. Bot. 394.

The whole plant is used. It has a peculiar, but not unpleasant odor, when bruised, and an astringent, bitterish taste. It is much celebrated as a diuretic, especially in dysuria of children.

INFUSION OF FLEABANE.

R. Fleabane, one ounce.
Boiling water, one pint.

Infuse. Dose, two to four fl. ounces, every three or four hours. *Dewees.*

ERYNGIUM.

1. ERYNGIUM MARITIMUM.

SEA HOLLY.

This plant is a native of Europe, growing on sandy beaches.

Sex. Syst. Pentand. digyn. *Nat. Syst.* Apiaceæ.

Linn. Sp. Pl. 337. Griffith, Med. Bot. 315.

It is thought to be diuretic, and was also esteemed aphrodisiac. The part used is the root, which has a sweet, agreeable taste, and an aromatic odor.

2. ERYNGIUM AQUATICUM.

BUTTON SNAKEROOT.

A native species, growing in wet places, from Virginia to Florida.

Linn. Sp. Pl. ed. 2. p. 336. Torrey & Gray, Fl. 1. 604.

The root is bitter, aromatic, and pungent. It is diaphoretic and expectorant, and is useful in the same cases to which senega is appropriate. Used to some extent in South Carolina.

EUPATORIUM.

BONESET.

This, the *E. perfoliatum*, is a native plant, found in abundance in most parts of the United States, in moist situations. Several other native species are almost identical in their properties, as *E. teucrifolium*, *E. purpureum*, &c.

Sex. Syst. Syngen. æqual. *Nat. Syst.* Asteraceæ.

The parts used are the leaves and flowers. These have a faint odor, and a peculiar, bitter taste. It is tonic and diaphoretic, and, in large doses, emetic and laxative. Dose, in substance, as a tonic, twenty to thirty grains.

INFUSION OF BONESET.

R. Boneset, one ounce.
Boiling water, one pint.
Infuse for two hours, in a covered vessel, and strain. *U. S. Ph.*

As a diaphoretic, to be taken freely, warm; as a tonic, a fl. ounce, cold.

DECOCTION OF BONESET.

R. Boneset, one ounce.
Water, one pint and a half.
Boil down to a pint.

Dose, a wineglassful, or more, as an emetic and cathartic.

COMPOUND INFUSION OF BONESET.

R. Boneset, half an ounce.
Sage, each, one drachm.
Cascarilla, one drachm.
Boiling water, one pint and a half.

Infuse till cold, and strain. A wineglassful, every three or four hours. In hectic fever.

Ellis.

EUPHORBIA.

SPURGE.

1. EUPHORBIA COROLLATA.

BOWMAN'S ROOT.

A native plant, with numerous white flowers, found in many parts of the United States, in dry, sandy soils.

Sex. Syst. Dodecand. trigyn. *Nat. Syst.* Euphorbiaceæ.

Linn. Sp. Pl. 258. *Griffith, Med. Bot.* 592.

The root, which is the part used, is large, white, inodorous, and almost tasteless. It is a safe and certain emetic, in doses of fifteen or twenty grains, and diaphoretic in that of three or four grains.

2. EUPHORBIA IPECACUANHA.

WILD IPECACUANHA.

This is also a native plant, with numerous procumbent stems, and variously shaped leaves.

Linn. Sp. Pl. 653. *Griffith, Med. Bot.* 592.

The root, the part used, is large, white, almost inodorous, and has a somewhat sweet taste. It is an energetic emetic, but, like the last species, apt to act on the bowels. Dose, ten to fifteen grains.

COMPOUND POWDER OF WILD IPECACUANHA.

R. Powdered wild ipecacuanha,
" opium, each, six grains.
Sulphate of potassa, forty-eight grains.

Triturate well, and divide into six powders. As a substitute for Dover's powder, to which it is equal, if not superior. *W. P. C. Barton.*

3. EUPHORBIA HYPERICIFOLIA.

BLACK PURSLANE.

A small, procumbent, native plant, with leaves often marked with a purple spot.

Linn. Sp. Pl. 652. *Griffith, Med. Bot.* 593.

The whole plant is used, and is said to be astringent and narcotic. It is inodorous, but has a sweetish, somewhat austere taste. It has been recommended in diarrhœa, fluor albus, &c.

INFUSION OF BLACK PURSLANE.

R. Dried black purslane, half an ounce.
Boiling water, one pint.

Infuse for half an hour, and strain. Dose, in dysentery and diarrhœa, a tablespoonful, every hour, till the morbid symptoms yield, then less frequently; in menorrhagia and leucorrhœa, a wineglassful, three times a-day. *Zollickoffer.*

4. EUPHORBIA LATHYRIS.

MOLE PLANT.

A tall species, native of Europe, but often cultivated in this country.

Linn. Sp. Pl. 655. *Lindley, Flor. Med.* 194.

The seeds, which are the officinal portion, afford a purgative oil, which is tasteless, and almost inodorous. It acts on the bowels, in doses of from four to twelve drops, but is apt to cause nausea and vomiting.

MIXTURE OF OIL OF EUPHORBIA.

R. Oil of euphorbia, eight drops.
Powdered gum Arabic, one drachm.
Sugar, two ounces.
Distilled water, three fl. ounces.

Mix. As a purgative draught. *Pichonnier.*

EUPHORBIIUM.

EUPHORBIIUM

Is the concrete resinous juice obtained from several succulent species of euphorbia, principally natives of Africa. It is in the form of tears, or small, irregular masses, of a yellowish, or reddish color. The taste is at first slight,

but soon becomes acrid and burning; the odor is very faint. It is a violent emetic and purgative, and is not used internally; as an external application it is employed in Europe, and also as a sternutatory.

STERNUTATORY OF EUPHORBIIUM.

R. Powdered euphorbium, one drachm.
 " white hellebore, four scruples.

Mix. Very violent in its action. *Pierquin.*

OIL OF EUPHORBIIUM.

R. Euphorbium, one part.
 Olive oil, ten parts.
 Digest for ten days, and filter. As a friction in paralysis. *Van Mons.*

PLASTER OF EUPHORBIIUM.

R. Pitch, three ounces.
 Yellow wax, each, one ounce.
 Turpentine, one ounce.
 Melt together, and add

Ammoniac,	}	each, one ounce.
Olibanum,		
Mastich,		
Calamine,		
Euphorbium,	}	each, two ounces.
Pyrethrum,		
Common salt,		

Stir, till cold. A stimulating and rubefacient application, known as *capuchin ointment*.
Wirttemberg Ph.

EUPHRASIA.

EYEBRIGHT.

The *Euphrasia officinalis* is a small, annual plant, a native of Europe, and the northern parts of Asia and America.

Sex. Syst. Didynam. gymnos. *Nat. Syst.* Scrophulariaceæ.

Linn. Sp. Pl. 841. *Lindley, Flor. Med.* 506.

The whole plant is used. It is slightly bitter and aromatic, and, at one time, was much used in diseases of the eyes. Dr. Kranichfeld says it is peculiarly beneficial in catarrhal ophthalmia, and also in cough and other catarrhal affections.

F.

FERRUM.

IRON.

A solid, hard, malleable, ductile metal, of a silvery-white color, and fibrous texture, attracted by the magnet, of a slight styptic taste, and a faint, but peculiar odor, when rubbed. Oxidizable in the air, and when heated to whiteness, burning with great brilliancy. It is employed medicinally in a great number of forms, both in the metallic state, in those of oxides and salts. All these are powerfully tonic and astringent.

FERRI FILUM.

IRON WIRE.

FERRI RAMENTA.

IRON FILINGS.

These should always be perfectly pure, when used internally, and in a state of minute division.

FERRI PULVIS.

PREPARED METALLIC IRON.

R. Iron filings, at will.

Pound in an iron mortar, with great force, sift through a fine sieve, and porphyryze, until all metallic brilliancy has disappeared, and again, sift. Keep in a perfectly dry place. *Paris Cod.*

Dose, from five to twenty grains.

R. Spread oxide of iron in a tube, heat the tube, and cause a stream of hydrogen gas to pass through it, till the iron is reduced.

Soubeiran.

R. Place precipitated carbonate of iron on layers of iron gauze, in a tube of wrought iron; pass a stream of hydrogen through it, heat to a dull red, and maintain this for some hours; keeping up a small current of gas, till cool.

W. Procter

Dose, two to five grains every three hours.

POWDER OF METALLIC IRON.

R. Prepared metallic iron, ten grains.
 Sugar, one drachm.

Mix. To be given, four times a-day, in chlorosis and amenorrhœa. *Brera*

COMPOUND POWDER OF METALLIC IRON.

- R. Prepared metallic iron,
Powdered guaiacum, each, half a drachm.
Prepared oyster shell, half an ounce.
Sugar, three drachms.

Mix. Divide into ten powders; one, twice a-day, in scrofula. *Galis.*

- R. Prepared metallic iron,
Powdered cinnamon, each, half a drachm.
Sugar, three drachms.

Mix, and divide into twelve powders. One every three hours, as a tonic. *Guibourt.*

- R. Prepared metallic iron, one grain.
Carbonate of magnesia, five grains.
Powdered cinnamon, two grains.
" sugar, one scruple.

Mix. One half in the morning, the other in the evening, in rachitis. *Hufeland.*

- R. Prepared metallic iron,
Powdered anise, Castor, each, two drachms.
Powdered cinnamon, " nutmeg, each, one drachm.

Mix, and divide into twenty-four powders. One, morning and evening, in chlorosis. *St. Marie.*

BOLUS OF PREPARED IRON.

- R. Prepared metallic iron,
Powdered myrrh, each, ten grains.
Aromatic powder, two grains.
Powder of rosemary, half a drachm.
Syrup of orange peel, sufficient.

Make a bolus. To be taken morning and evening, in amenorrhœa. *Augustin.*

PILLS OF PREPARED IRON.

- R. Prepared metallic iron, two ounces.
Extract of wormwood, sufficient.

Mix, and make six-grain pills. Three or four, morning and evening, in chlorosis, dyspepsia, &c. *Soubiran.*

- R. Prepared metallic iron, one drachm.
Powdered black hellebore, one scruple.
Extract of gentian, two drachms.
Syrup of saffron, sufficient.

Mix, and form sixty pills. Two, three times a-day, as an emmenagogue. *St. Marie.*

- R. Prepared metallic iron, half an ounce.
Powdered colombo, four scruples.
" rhubarb, " cinnamon, each, two scruples.

Extract of quassia, sufficient.

Mix, and form pills of three grains each. Three to four a-day, as a tonic in debility. *Jahn.*

AROMATIC MIXTURE OF IRON.

- R. Powdered Peruvian bark, one ounce.
" colombo, three drachms.
Bruised cloves, two drachms.
Iron filings, half an ounce.

Digest for three days in a covered vessel, with as much peppermint water as will yield a mixture of twelve ounces, after filtration; then add compound tincture of cardamom, three ounces, and tincture of orange peel, three drachms. *Dub. Ph.*

As a tonic, in doses of one or two fl. ounces.

FERRI ACETAS.

ACETATE OF IRON.

- R. Subcarbonate of iron, one part.
Acetic acid, six parts.

Digest for three days, and filter. *Dub. Ph. 1826.*
Dose, from ten to twenty-five drops, in water.

TINCTURE OF ACETATE OF IRON

- R. Acetate of potassa, two parts.
Sulphate of iron, one part.
Rectified spirit, twenty-six parts.

Rub the acetate and sulphate together, dry by a moderate heat, and triturate with the spirit. Digest in a closed bottle for seven days, and decant. *Dub. Ph. 1826.*

Dose, from thirty drops to a teaspoonful, in some convenient vehicle.

ALCOHOLIC TINCTURE OF ACETATE OF IRON.

- R. Liquid acetate of iron, one part.
Diluted alcohol, seven parts.

Digest for two days, and filter. Dose, ten to fifty drops. *Guibourt.*

ETHEREAL TINCTURE OF ACETATE OF IRON.

- R. Liquid acetate of iron, nine ounces.
Acetic ether, one ounce.
Alcohol, two ounces.

Mix. Dose, twenty to thirty drops. *Spielmann.*

ACETATE OF IRON AND ALUM.

- R. Sulphate of iron, two parts.
Alum, one part.
Boiling water, ten parts.

Filter the solution, and gradually add solution of subacetate of lead till precipitation ceases; let rest, and filter; evaporate to consistence of a jelly. As a marking ink; it resists alkalies, but not acids. *Guibourt.*

FERRUM AMMONIATUM.

AMMONIATED IRON.

- R. Subcarbonate of iron, three ounces.
 Muriatic acid, ten fl. ounces.
 Muriate of ammonia, two pounds and a half.
 Distilled water, four pints.

Mix the subcarbonate with the acid, in a glass vessel, and digest for two hours; then add the muriate of ammonia, previously dissolved in the water, filter, and evaporate to dryness.

U. S. Ph.

Said to be useful in amenorrhœa, scrofula, epilepsy, &c. Dose, four to twelve grains, several times a-day.

COMPOUND POWDER OF AMMONIATED IRON.

- R. Ammoniated iron,
 Powdered quassia, each, two drachms.
 Powdered chamomile, half an ounce.

Mix, and divide into twenty-four powders. One, four times a-day, as a febrifuge.

Wurtzburg Ph.

- R. Ammoniated iron, five grains.
 Ammoniac, three grains.
 Powdered seneka, five grains.
 " liquorice, one drachm.

Mix, and divide into twelve powders. One, every three hours, in gonorrhœa.

Foy.

BOLUS OF AMMONIATED IRON.

- R. Ammoniated iron, five grains.
 Aromatic powder, twelve grains.
 Mucilage, sufficient.

Mix, and make bolus. One to be taken twice a-day, in chlorosis and scrofula.

Ainslie.

PILLS OF AMMONIATED IRON.

- R. Ammoniated iron,
 Galbanum, each, one drachm.
 Assafetida, two drachms.
 Castor, one scruple.
 Tincture of valerian, sufficient.

Beat into mass, and form pills of three grains. Two, morning and evening, in atonic nervous disorders.

Radius.

- R. Ammoniated iron,
 Aromatic powder,
 Extract of Peruvian bark,
 Alum, each, half a drachm.
 Oil of cinnamon, twelve drops.

Beat into mass, and form pills of two grains. Five to ten, morning and evening, in asthenic menorrhagia.

Formey.

MIXTURE OF AMMONIATED IRON.

- R. Powdered galanga,
 " rhubarb, each,
 half a drachm.
 Boiling water, sufficient

to obtain three ounces of strained solution; add

- Ammoniated iron, six grains.
 Peppermint sugar, four grains.

Mix. A dessertspoonful, every three hours, in indigestion and chronic diarrhœa of children.

Wendt.

TINCTURE OF AMMONIATED IRON.

- R. Ammoniated iron, four ounces.
 Proof spirit,
 Water, each, one pint.

Dissolve, and filter. Dose, forty to sixty drops.

Lond. Ph.

FERRI AMMONIO-CITRAS.

AMMONIO-CITRATE OF IRON.

- R. Crystallized citric acid, four ounces.
 Clean iron filings, two ounces.
 Distilled water,
 Solution of ammonia, each, sufficient.

Dissolve the citric acid in twenty times its weight of water, add the iron and apply a gentle heat till effervescence ceases, renewing the water as it evaporates; filter, add the ammonia in slight excess, and evaporate on a water-bath to a syrupy consistence; then spread out in thin layers, and dry by a gentle heat.

Redwood.

The advantages of this salt, are its solubility and freedom from unpleasant taste. Dose, five to ten grains.

- R. Liquid citrate of iron, one ounce.
 Water of ammonia, sufficient
 to saturate the excess of acid.

Beral.

FERRI ARSENIAS.

ARSENIATE OF IRON.

- R. Solution of sulphate of iron, at will.
 Add gradually a solution of arseniate of potassa, collect, and wash the precipitate. Dose, one-twelfth of a grain.

PILLS OF ARSENIATE OF IRON.

- R. Arseniate of iron, three grains.
 Extract of hops, two drachms.
 Powdered mallow, half a drachm.
 Syrup, sufficient.

Beat together, and form forty-eight pills. One or two to be given, daily, in scrofulous, herpetic, and cancerous affections.

Bielt.

OINTMENT OF ARSENIATE OF IRON.

- R. Arseniate of iron, half a drachm.
 Phosphate of iron, two drachms.
 Spermaceti cerate, six drachms.

Triturate together. As an application to cancerous ulcers. To be used with great caution.

Carmichael.

FERRI BROMIDUM.

BROMIDE OF IRON.

- R. Bromine, one part.
 Distilled water, twenty parts.
 Iron filings, sufficient.

Boil on a sand-bath; when the solution becomes greenish, filter, and evaporate to dryness.

Magendie.

PILLS OF BROMIDE OF IRON.

- R. Bromide of iron, twelve grains.
 Conserve of roses, eighteen grains.
 Gum Arabic, twelve grains.

Mix, and form twenty pills. Two, in the morning and evening, as a tonic and alterative.

Magendie.

- R. Bromide of iron, one drachm.
 Extract of liquorice, sufficient.

Mix, and make sixty pills. One or two, in the morning and evening, in scrofula, &c. *Radiis.*

OINTMENT OF BROMIDE OF IRON.

- R. Bromide of iron, one drachm.
 Bromine, twelve grains.
 Lard, one ounce.

Mix. In frictions on the inside of the thighs, in amenorrhœa. *Cadet.*

FERRI CARBONAS.

CARBONATE OF IRON

From the avidity with which carbonate of iron absorbs oxygen from the air and loses its carbonic acid, it is a very unstable preparation; and therefore what is usually prescribed under this name is merely a subcarbonate, or sesquioxide. By combination with saccharine matter, it, however, is protected from the oxidizing process, to a very great extent.

SACCHARINE CARBONATE OF IRON.

- R. Sulphate of iron, four ounces.
 Carbonate of soda, five ounces.
 Pure sugar, two ounces.
 Water, four pints.

Dissolve the sulphate and carbonate, each in half the water, add the solutions together, collect the precipitate, divest it of as much water as possible, and immediately triturate it with the sugar. Dry it at a low temperature.

Ed. Ph.

Dose, from five to thirty grains.

PILLS OF CARBONATE OF IRON.

(VALLET'S MASS.)

- R. Sulphate of iron, eight ounces.
 Carbonate of soda, ten ounces.
 Clarified honey, three ounces.
 Sugar, two ounces.
 Syrup, sufficient.
 Boiling water, two pints.

Dissolve the sulphate and carbonate, each in a pint of water, adding to each solution one fl. ounce of syrup; mix the two solutions in a bottle just large enough to hold the mixture, close accurately with a stopper, and set by to let the carbonate of iron subside; decant, wash precipitate with water sweetened with syrup, in the proportion of a fl. ounce to the pint, until the washings cease to be saline; express, in flannel, as much of the water as possible, and mix with the honey. Heat over a water-bath, to proper consistence. This is known as Vallet's carbonate of iron. Dose, from ten to thirty grains, in the course of the day, where iron is indicated.

U. S. Ph.

SUBCARBONATE OF IRON.

- R. Sulphate of iron, eight ounces.
 Carbonate of soda, nine ounces.
 Boiling water, one gallon.

Dissolve the sulphate and carbonate, each in four pints of the water, mix the solutions, stir, set by for powder to subside, decant, wash the precipitate with hot water, wrap in bibulous paper, and dry with gentle heat. *U. S. Ph.*

Dose, five to thirty grains, or even more, several times a-day.

POWDER OF SUBCARBONATE OF IRON.

- R. Subcarbonate of iron, one drachm.

Divide into six powders. One every six hours, in syrup. In chorea, neuralgia, &c. *Ellis.*

MIXTURE OF SUBCARBONATE OF IRON.

- R. Powdered myrrh, one drachm.
 Sulphate of iron, one scruple.
 Carbonate of potassa, one drachm.
 Sugar, two drachms.
 Water, six fl. ounces.

Mix. Dose, a tablespoonful, according to circumstances. *Ellis.*

COMPOUND IRON MIXTURE.

- R. Myrrh, one drachm.
 Carbonate of potassa, twenty-five grains.

Rose-water, seven fl. ounces and a half.
Powdered sulphate of iron, one scruple.
Spirit of lavender, half a fl. ounce.
White sugar, one drachm.

Rub the myrrh with the rose-water, gradually added, mix with the spirit of lavender, sugar, and carbonate of potassa, and, lastly, with the iron; pour into a well-stopped bottle, and keep closed. *U. S. Ph.*

These are well known under the name of Griffith's anti-hectic mixture. Dose, one or two fl. ounces, two or three times a-day, in chlorosis, debility of the gastric organs, &c.

R. Subcarbonate of iron,
Carbonate of potassa,
each, four ounces.
Black hellebore, two ounces.
Water, sufficient.

Boil, after macerating for some days, stirring from time to time; then let settle, decant, and evaporate to consistence of an extract, and add to this

Spirit of scurvy-
grass, three times its weight,
and filter. Forty to sixty drops, in hypochondria, melancholia, &c. *Palat. Ph.*

ARTIFICIAL CHALYBEATE WATER.

R. Water, fifty pints.
Carbonate of lime, five drachms.
" magnesia, ten drachms.
Black oxide of iron, two drachms.
Sulphate of magnesia, six drachms.
Common salt, one drachm.

Mix, and condense a thousand cubic inches of carbonic acid in the solution. *Swed. Jour.*

COMPOUND PILLS OF SUBCARBONATE OF IRON.

R. Powdered foxglove, half a drachm.
" yew leaves,
Carbonate of iron, each, one drachm.
Ergot, two drachms.
Syrup, sufficient.

Mix, and make two hundred and fifty pills. Two to eight a-day, with an infusion of savine, in chlorosis and amenorrhœa. *Neumann.*

R. Powdered myrrh, two drachms.
Carbonate of soda,
Sulphate of iron, each, one drachm.
Syrup, sufficient.

Rub the myrrh with the carbonate of soda, then add the sulphate of iron, and rub well; beat with syrup into a mass, and divide into eighty pills. *U. S. Ph.*

Dose, two to six pills, three times a-day.

R. Sulphate of iron,
Bicarbonate of potassa, each, two drachms.
Powdered mallow root, thirty-six grains.

Gum Arabic, eighteen grains.

Beat with water into a mass, and form forty-eight pills.

R. Sulphate of iron,
Myrrh, each, two scruples.
Carbonate of potassa,
Soap, each, half a drachm.

Beat together, and form forty pills. Dose, two, three times a-day, as an emmenagogue. *Ellis.*

R. Subcarbonate of iron,
Extract of conium, each, one drachm.

Mix, and divide into twenty-four pills. Two to be taken twice a-day. Given in fluor albus and scrofula. *A. T. Thomson.*

IRON WINE.

R. Carbonate of iron,
one ounce and a half.
Contused orange peel,
" gentian, each, half an ounce.
Port wine, two pints.

Mix, and digest at a moderate heat for three days, repeatedly agitating. *Chapman.*

IRON PLASTER.

R. Subcarbonate of iron, three ounces.
Lead plaster, two pounds.
Burgundy pitch, half a pound.

Add the subcarbonate to the other ingredients previously melted together, and stir till cold.

U. S. Ph.

As a strengthening plaster, in pains, and want of power in the muscles

FERRI CARBURETUM.

CARBURET OF IRON.

This article, well known as Plumbago, or Black Lead, has been highly recommended, of late years, in cutaneous affections, and is recognized as officinal by many of the pharmacopœias of continental Europe.

PREPARED CARBURET OF IRON.

R. Pulverized carburet of iron,
one pound.

Boil in water for an hour, decant, and digest in eight ounces of water, mixed with two ounces of nitric and muriatic acids; digest for twenty-four hours, decant, and wash well with water, and dry. Dose, from five to fifteen grains, two or three times a-day. *Pruss. Ph.*

POWDER OF CARBURET OF IRON.

R. Powdered carburet of iron,
Sugar, each, half a drachm.
Mix, and divide into six powders. One to be
taken every hour, in lepra and lichen. *Radiis.*

PILLS OF CARBURET OF IRON.

R. Powdered carburet of iron,
Extract of bittersweet, each,
one drachm.
Calomel, twelve grains.
Golden sulphuret of
antimony, six grains.
Extract of liquorice, sufficient.
Mix, and form sixty pills. Dose, five, three
times a-day. *Niemann.*

ELECTUARY OF CARBURET OF IRON.

R. Powdered carburet of iron,
Flowers of sulphur,
Crude antimony, each, half an ounce.
Black oxide of mercury, one scruple.
Wild violet,
Extract of pulsatilla,
" bittersweet, each,
two drachms.
Camphor, one drachm.
Syrup of orange peel, sufficient.
Mix. To be used in eight days, in teaspoonful
doses. *Phæbus.*

MIXTURE OF CARBURET OF IRON.

R. Powdered carburet of iron,
Sulphur, each, one drachm.
Triturate together, and add
Barley water, one pint.
To be taken in four doses. In psora, &c. *Brera.*

OINTMENT OF CARBURET OF IRON.

R. Powdered carburet of iron,
half an ounce.
Flowers of zinc, one drachm.
Lard, one ounce.
Triturate together. As an application in her-
petic affections. *Brera.*
R. Powdered carburet of iron, one ounce.
Flowers of sulphur, four scruples.
Rose ointment, six ounces.
Mix. *Taddei.*

PLASTER OF CARBURET OF IRON.

R. Powdered carburet of iron, one part.
Soap plaster, four parts.
Triturate together. *Giordano.*

FERRI CHLORIDUM.

CHLORIDE OF IRON.

R. Subcarbonate of iron, at will.
Muriatic acid, sufficient
to dissolve the carbonate; evaporate to dryness
by a water-bath. *Cottureau.*

R. Peroxide of iron, five ounces.
Muriatic acid, nineteen fl. ounces.
Boil for ten minutes, then evaporate to fifteen
ounces; filter. Place under a bell-glass, with a
vessel containing caustic potassa; in a few days
mammillary granules will form; separate these,
and preserve in a glass-stopped bottle. *Beral.*

CHLORIDE OF IRON AND ACETATE OF
LEAD.

R. Liquid chloride of iron, four ounces.
Evaporate to one-half, and add
Acetate of lead, four ounces.
Dry, and pulverize. Four to twelve grains in
hemorrhages. *Cadet de Gassicourt.*

TINCTURE OF CHLORIDE OF IRON.

R. Subcarbonate of iron, half a pound.
Muriatic acid, one pint.
Alcohol, three pints.
Pour the acid on the subcarbonate; when the
effervescence has ceased, apply a gentle heat,
and continue it, stirring occasionally, until the
carbonate is dissolved; then filter and add the
alcohol to the clear solution. *U. S. Ph.*

An active and energetic preparation; tonic,
diuretic, styptic, &c. Dose, ten to thirty drops,
gradually increased, two or three times a-day.

R. Muriatic acid, two ounces (troy).
Protocarbonate of
iron, sufficient to saturate.
Honey, two ounces and a half.
Alcohol, sufficient.

Saturate the acid with the carbonate, add the
honey and sufficient alcohol to make nineteen
fl. ounces; after standing six hours, filter.
W. Procter.

SYRUP OF CHLORIDE OF IRON.

R. Chloride of iron, one part.
Syrup, twenty-three parts.
Dissolve. Dose, one to four drachms. *Beral.*

CHLORIDE OF IRON MIXTURE.

R. Saffron, half a drachm.
Water, four fl. ounces.
Infuse for an hour, strain, and add
Chloride of iron, half a drachm.
Syrup of valerian, two ounces.
Mix. Dose, a tablespoonful several times a-day,
in hemorrhage or chlorosis. *Cadet de Gassicourt.*

R. Mallow root, two drachms.
Water, sufficient
to obtain two ounces of decoction; strain, and add

Gum Arabic, two drachms.
Chloride of iron, half a scruple.
Syrup of mallow, six drachms.

Mix. Two teaspoonfuls, every hour or two, in softening of the stomach. *Radiis.*

R. Tincture of chloride of iron,
Compound tincture of
aloes, each, half a fl. ounce.
Tincture of castor, two fl. drachms.

Mix. Ten to thirty drops, three times a-day, in infusion of chamomile, as an emmenagogue. *Ellis.*

ETHEREAL, OR BESTUCHEFFE'S TINCTURE OF CHLORIDE OF IRON.

R. Sesquichloride of iron, one drachm.
Sulphuric ether, one fl. ounce.
Alcohol, three fl. ounces.

Mix. *Prus. Ph., 1846.*

The sesquichloride of iron may be obtained pure and neutral, by passing chlorine through a solution of protochloride of iron, until a solution of the red ferrocyanide of potassium no longer produces a blue precipitate; then evaporate by a water-bath, and crystallize. *Fr. Mayer.*

R. Sesquichloride of iron, one part.
Sulphuric ether, four parts.

Mix, and add
Alcohol, eight parts.
Van Mons.

R. Tincture of chloride of iron,
Sulphuric ether, equal parts.

Mix. *Guibourt.*

R. Chloride of iron, one part.
Hoffmann's anodyne, seven parts.

Mix. *Beral.*

These are active preparations, and much used in Germany. Dose, twenty to thirty drops.

FERRI CITRAS. CITRATE OF IRON.

R. Crystals of citric acid, three parts.
Hydrated oxide of iron (dry), two parts.

Distilled water at 180°, twelve parts.

Dissolve, filter, and evaporate to consistence of syrup, spread in thin layers, and dry. *Duhamel.*

The dose is five to eight grains.

LIQUID CITRATE OF IRON.

R. Citrate of iron, four ounces.
Water, twelve fl. ounces.

Dissolve, with the aid of heat, and evaporate until the solution measures half a pint.

W. Procter.

SYRUP OF CITRATE OF IRON.

R. Liquid citrate of iron, one ounce
Syrup, fifteen ounces.
Spirit of citron, two drachms.

Mix. Dose, two drachms to an ounce. *Beral.*

WINE OF CITRATE OF IRON.

R. Liquid citrate of iron, eight scruples.
Rhenish wine, sixteen ounces.

Digest for two or three days, and filter. *Beral.*

AROMATIC WINE OF CITRATE OF IRON.

R. Iron filings, one ounce.
Lemon juice, three fl. ounces.
Contused gentian, half an ounce.
" cinnamon, two drachms.
Rhenish wine, sixteen ounces.

Digest for twenty-four hours, and decant. The dose is a drachm to half a fl. ounce, two or three times a-day. *Hamburgh Cod.*

R. Iron filings, four ounces.
Bitter oranges, four.

Beat them together, and at the end of two days, add to the mixture

Madeira wine, ten fl. ounces.

Spirit of orange peel, two fl. ounces.

Digest, and then express and filter. Dose, half a drachm to two drachms. *Batav. Ph.*

Said to be one of the best compounds of iron.

TINCTURE OF CITRATE OF IRON.

R. Liquid citrate of iron, two ounces.
Diluted alcohol, thirteen fl. ounces.
Spirit of citron, one fl. ounce.

Mix. Dose, ten to fifty drops. *Beral.*

SYRUP OF CITRATE OF IRON.

R. Moist protocarbonate of iron,
(prepared as directed for
Vallet's mass, and washed
with sweetened water), at will.

Dissolve in a slight excess of citric acid, dissolved in water, and evaporate to dryness to make the syrup.

R. Citrate of iron, one ounce.
Syrup, five fl. ounces.

Dissolve. Dose, thirty drops to a teaspoonful. *E. Parrish*

CITRATE OF IRON AND QUINIA.

R. Liquid citrate of iron, four parts
Solution of citrate of quinia, one part

Mix, and evaporate. As a tonic, in doses of five grains, three times a-day.

Beral.

CITRATE OF IRON AND MAGNESIA.

R. Hydrated oxide of iron, two parts.
Citric acid (in solution), three parts.

Mix, dissolve, saturate the liquor with carbonate of magnesia, and evaporate to dryness.

This salt is soluble. The dose is from two to five grains.

Van der Corput.

SYRUP OF CITRATE OF IRON AND MAGNESIA.

R. Citrate of iron and magnesia, eight parts.

Orange-flower water, fifteen parts.

Simple syrup, one hundred and eighty parts.

Mix.

Van der Corput.

FERRI FERROCYANURETUM.

PRUSSIAN BLUE.

R. Sulphate of iron, four ounces.
Sulphuric acid,

three fl. drachms and a half.

Nitric acid, six fl. drachms, or sufficient.

Ferrocyanuret of potassium, four ounces and a half.

Water, two pints.

Dissolve the sulphate in a pint of the water, add the sulphuric acid, and boil. Add the nitric acid in small portions, boiling for a few moments after each addition, till a dark color is no longer produced, and allow to cool. Dissolve the ferrocyanuret in the rest of the water, and add the solution by degrees, to the first liquid, stirring each time. Filter, wash the precipitate with boiling water, until the washings are tasteless. Dry, and powder.

U. S. Ph.

Tonic, febrifuge, and alterative. Dose, three to five grains, several times a-day, gradually increased till some effect is produced.

POWDER OF PRUSSIAN BLUE.

R. Prussian blue, twelve grains.
Sugar, one drachm.

Mix, and divide into twelve powders. One, every two hours. In epilepsy.

Radius.

COMPOUND POWDER OF PRUSSIAN BLUE.

R. Prussian blue, twelve grains.
White pepper,
Mustard, each, half an ounce.

Mix, and divide into twelve powders. One, every quarter of an hour, in the apyrexia of intermittents.

Foy.

R. Prussian blue,
Powdered guaiacum, each, half to one drachm.
Mix, and divide into twelve powders. One, three times a-day, in intermittents.

Ellis.

PILLS OF PRUSSIAN BLUE.

R. Prussian blue, eighteen to thirty-six grains.

Ammoniac,
Extract of dandelion,
Rhubarb, each, one drachm.

Mix, and make fifty-four pills. Four to six, twice a-day. In diseases of the ganglionic system.

Radius.

MIXTURE OF PRUSSIAN BLUE.

R. Prussian blue, one ounce.
Parsley water, six ounces.
Cherry-laurel water, two drachms.

Mix. A teaspoonful, three times a-day. In stone and gravel.

Radius.

ointment OF PRUSSIAN BLUE.

R. Cod-liver oil, one ounce.
Prussian blue, twenty-four grains.
Cyanide of mercury, eight grains.
Oil of cherry-laurel, four drops.

Mix. As an application in strumous ophthalmia.

Canon de Villards.

BLUE INK.

R. Pure Prussian blue, six parts.
Oxalic acid, one part.

Triturate with a little water, to a perfectly smooth paste; then dilute with water to a proper consistence. A little gum may be added to prevent the fluid from spreading.

Mohr.

FERRI GALLAS.

GALLATE OF IRON.

This is not used in medicine, but, with the tannate of iron, it forms the basis of writing ink.

WRITING INK.

R. Aleppo galls, bruised, twelve pounds.
Water, six gallons.

Boil in a copper vessel for an hour, adding water, to supply loss by evaporation, and strain; boil residue with four gallons of water, for half an hour, and again boil with two gallons of water; mix decoctions, and add

Sulphate of iron, four pounds.
Contused gum

Arabic, three pounds and a half.

Agitate, permit to settle, strain through fine hair sieve, and keep in close vessels.

Lewis.

- R. Powdered galls, eight pounds.
 Rasped logwood, four pounds.
 Gum Arabic, three pounds.
 Sulphate of iron, four pounds.
 " copper,
 Sugar candy, each, one pound.
 Ammoniac, two ounces.
 Water, sixteen pints.
 Alcohol, half a pint.

Macerate for forty-eight hours, with a gentle heat, let rest for ten days, and decant into bottles. *Hunt.*

- R. Powdered galls, fifty parts.
 Water, eight hundred parts.

Digest for twenty-four hours, strain, and add
 Sulphate of iron,
 Gum Arabic, each, twenty-five parts.

When clear, add a solution of

- Muriate of ammonia, eight parts.
 Gum Arabic, two parts.
 Boiling water, sixteen parts.
 Oil of lavender, one part.

Said to be indelible. *Guibourt.*

- R. Bruised galls, forty ounces.
 Gum Arabic, ten ounces.
 Sulphate of iron, nine ounces.
 Soft water, forty-five fl. ounces.

Macerate for three weeks, often stirring. A very permanent ink. *English Exchequer.*

FERRI IODIDUM.

IODIDE OF IRON.

- R. Iodine, two ounces.
 Iron filings, one ounce.
 Distilled water, one pint and a half.

Mix the iodine with a pint of the water, in a glass vessel, gradually add the iron filings, constantly stirring. Heat till liquid becomes of a light greenish color, filter, and wash residuum with remainder of water, boiling hot. Evaporate liquid to dryness, in an iron vessel, at a heat not above 212° . Keep in a well-closed bottle. *U. S. Ph.*

Tonic, alterative, diuretic, and emmenagogue. Dose, three grains, gradually increased to eight, or more.

LOZENGES OF IODIDE OF IRON.

- R. Iodide of iron, thirty grains.
 Saffron, half a drachm.
 Mucilage of tragacanth,
 Infusion of cinnamon, each, sufficient.

Mix and make two hundred and forty lozenges. Fifteen to twenty a-day, gradually augmenting, in amenorrhœa and leucorrhœa. *Pierquin.*

PILLS OF IODIDE OF IRON.

- R. Iodide of iron, one drachm.
 Bromide of sodium, half a drachm.
 Extract of liquorice, sufficient.

Mix, and form pills of two grains. One to three, morning and evening, in scrofula, &c. *Wernick.*

PILLS OF PROTO-IODIDE OF IRON.

- R. Crystallized sulphate of iron, 24.7 grains.
 Iodide of potassium, 32.4 grains.
 Gum tragacanth, 4.6 grains.
 Sugar, 15.4 grains.
 Syrup of marsh mallow,
 Powder of marsh mallow, each, sufficient.

Reduce the sulphate to a fine powder, then the iodide of potassium; triturate the mixture, then add the gum, sugar, and syrup, and, if necessary, the powder of marsh mallow. Mix, and make thirty-six pills. Each contains .8 of a grain of dry iodide, or 1.09 of the hydrated iodide of iron. Keep in closed bottles, in a dry place. *Callond.*

- R. Sulphate of iron, one drachm.
 Iodide of potassium, four scruples.
 Powdered tragacanth, ten grains.
 Powdered sugar, half a drachm.

Beat into a mass with syrup, and divide into forty pills. *U. S. Ph.*

- R. Iodine,
 Iron, reduced by hydrogen,
 each, half a drachm.

Rub the iron and iodine to a fine powder, then add

- Honey, sufficient

to give the consistency of molasses, and rub till the mixture assumes a greenish tinge, then add

- Extract of cinchona, two scruples.
 Powdered liquorice root, sufficient

to form a mass. Divide into thirty-two pills. Dose, one pill three or four times a-day.

D. F. Wright.

SOLUTION OF IODIDE OF IRON.

- R. Iodide of iron, one drachm
 Distilled water, one fl. ounce

Mix. Six, to ten drops, three times a-day, in some cold water. *Ellis*

- R. Iodide of iron, twenty-four grains.
 Distilled water, one fl. ounce.

Dissolve. A teaspoonful is a dose; this contains three grains. *Ellis.*

OFFICIAL SOLUTION OF IODIDE OF IRON.

- R. Iodine, two ounces.
 Iron filings, one ounce.
 Powdered sugar, twelve ounces.
 Distilled water, sufficient.

Mix the iodine with five fl. ounces of the water, in a porcelain or glass vessel, gradually add the iron filings, constantly stirring. Heat gently till the mixture becomes of a light-greenish color, filter, and add sugar. Wash on a filter, and allow to pass till the whole of the filtered liquid amounts to twenty fl. ounces. *U. S. Ph.*

Dose, from thirty to fifty drops.

SYRUP OF IODIDE OF IRON.

- R. Dry iodine, two hundred grains.
 Fine iron wire, well
 cleaned, one hundred grains.
 White sugar, in
 powder, four ounces and a half.
 Distilled water, six fl. ounces.

Boil the iodine, iron, and water, together, in a glass matrass; at first gently, to avoid the expulsion of iodine vapor, afterwards briskly, till about two fl. ounces remain. Filter this quickly, while hot, into a matrass containing the sugar, dissolve with a gentle heat, and add distilled water to make up six fl. ounces. Twelve minims contain one grain of iodide of iron.

Ed. Ph.

- R. Iodine, two ounces.
 Iron by hydrogen, half an ounce.
 Water, sufficient.
 Sugar, twelve ounces.

Powder the iodine, add two fl. ounces of water, then the iron very gradually, filter the solution into a bottle containing the sugar, wash the residual impurities, pour the washings into the bottle, and add water sufficient to make the whole measure twenty fl. ounces; shake the bottle and dissolve the sugar. *B. J. Crew.*

SYRUP OF PROTO-IODIDE OF IRON.

- R. Sulphate of iron, one drachm and a half.
 Iodide of potassium, two drachms.
 Water, half an ounce.
 Syrup, one ounce and a half.

Rub the salts together, dissolve them in the water, add the syrup, and bottle at once.

W. Procter.

WINE OF IODIDE OF IRON.

- R. Sulphate of iron, eight parts.
 Iodide of potassium, ten and a half parts.
 White wine, three hundred and twenty parts.

Pulverize the two salts, adding a few drops of wine, triturate for a few moments, add the rest of the wine, and filter. To be kept in glass-stopped bottles. Thirty-two parts contain one of the iodide. *Callond.*

- R. Iodide of iron, half an ounce.
 Claret wine, one pint.

Dissolve. A dessertspoonful, morning and evening. *Soubeiran.*

TINCTURE OF IODIDE OF IRON.

- R. Iodide of iron, one drachm.
 Alcohol,
 Distilled water, each, one fl. ounce.

Mix. Thirty drops to a fl. drachm, twice a-day. *Poy.*

TINCTURE OF PROTO-IODIDE OF IRON.

- R. Sulphate of iron, eight parts.
 Iodide of potassium, ten and a half parts.
 Alcohol (.842), one hundred and sixty parts.

Triturate the salts, separately, and then together; add alcohol, and filter. To be kept in glass-stopped bottles. Sixteen parts contain one of iodide of iron. *Callond.*

MIXTURE OF IODIDE OF IRON.

- R. Iodide of iron, sixteen grains.
 Compound tincture
 of gentian, one fl. ounce
 Distilled water, seven fl. ounces.

Mix. Two tablespoonfuls, two or three times a-day. *Ashwell.*

INJECTION OF IODIDE OF IRON.

- R. Iodide of iron, half an ounce.
 Distilled water, one pint.

Dissolve. As an injection, or lotion, in amenorrhœa, leucorrhœa, &c. *Soubeiran.*

- R. Iodide of iron, three grains.
 Distilled water, six fl. ounces.

Dissolve. In gonorrhœa; strength to be increased if required. *Ricord.*

SYRUP OF IODIDE AND CHLORIDE OF IRON.

- R. Iodine, three hundred and eighty-four grains.

Muriatic acid
 (1.16), four ounces and a half.
 Powdered sugar, twelve ounces.
 Iron filings,
 Distilled water, each, sufficient.

Mix the acid with an ounce and a half of clean iron-filings; allow to stand; agitate occasionally; when neutralized, filter. Mix the iodine with three ounces of water in a flask, add half the weight of iron filings, agitate till all the iodine is combined, and filter. Mix the solutions, add the sugar, and as much water as will make a pint. *W. Procter.*

SYRUP OF IODIDE AND CHLORIDE OF IRON MIXTURE.

R. Syrup of iodide and chloride of iron, two drachms.
Syrup of orange peel, four drachms.
Infusion of cascarrilla, four fl. ounces.
Mix. One-fourth to be taken twice a-day. *Battley.*

BATH OF IODIDE OF IRON.

R. Iodide of iron, two ounces.
Water, two hundred pints.
In leucorrhœa, amenorrhœa, chlorosis, &c. The quantity of iodide may be increased gradually to double, for adults. *Pierquin.*

OINTMENT OF IODIDE OF IRON.

R. Iodide of iron, one drachm and a half.
Lard, one ounce.
Triturate together. A piece as large as a hazel-nut to be rubbed on the inside of the thigh, morning and evening, in leucorrhœa and amenorrhœa; also beneficial in glandular swellings. *Pierquin.*

FERRI LACTAS.

LACTATE OF IRON.

R. Lactate of lime, one hundred parts.
Boiling water, five hundred parts.
Dissolve.

Pure sulphate of iron, sixty-eight parts.
Distilled water, five hundred parts.

Dissolve.

Mix the two solutions; filter, add an excess of lactic acid, heat in a water-bath, stirring constantly; filter, to separate the sulphate of lime; evaporate rapidly either in an iron vessel or porcelain one, with the addition of some iron filings; filter, and crystallize; wash the crystals with alcohol, and dry. *Lepage.*

Used in same cases as other preparations of iron. Dose, one or two grains, frequently repeated.

LOZENGES OF LACTATE OF IRON.

R. Lactate of iron, one drachm.
Sugar, one ounce and a half.
Mucilage of gum Arabic, sufficient.
Mix, and make lozenges of twelve grains. *Cap.*

PILLS OF LACTATE OF IRON.

R. Lactate of iron, equal parts.
Powdered mallow, sufficient.
Clarified honey, sufficient.
Make pills of two grains each. *Cap.*

SYRUP OF LACTATE OF IRON.

R. Lactate of iron, one drachm.
White sugar, twelve ounces and a half.
Boiling distilled water, six fl. ounces and a half.
Triturate the lactate with half an ounce of the sugar, dissolve quickly in the boiling water, pour the solution into a matrass, place on a sand-bath, add the rest of the sugar gradually; when dissolved, filter; and when cold, pour into well-stopped bottles.
Dose, from two to four fl. drachms. *Cap.*

FERRI MALAS.

MALATE OF IRON.

R. Black oxide of iron, at will.
Juice of sour apples, sufficient.
Heat the juice to the boiling point; add the oxide, as long as any is dissolved; cool, filter, and evaporate to dryness. Dose, eight grains to a scruple. *Van Mons.*

SOLUTION OF MALATE OF IRON.

R. Malate of iron, one ounce.
Water, twelve fl. ounces.
Dissolve and filter. As an astringent, in doses of a fl. drachm. *Spielmann.*

TINCTURE OF MALATE OF IRON.

R. Malate of iron, one part.
Cinnamon water, three parts.
Alcohol, each, three parts.
Macerate for three days, and filter. Dose, forty to sixty drops. *Austrian Ph.*

MIXTURE OF MALATE OF IRON.

R. Extract of Peruvian bark, gentian, each, one drachm.
Peppermint water, Chamomile water, each, one fl. ounce.
Malate of iron, half a drachm.
To be taken in divided doses, in dyspepsia, &c. *Augustin.*

BITTER INFUSION OF IRON.

R. Iron filings, three ounces.
Bruised ginger, gentian, each, one ounce.
" orange peel, half an ounce.

Infuse in a pint of strong old cider for a month, frequently agitating, and filter.

Dose, half a drachm to one drachm, thrice daily, as a tonic.

FERRI NITRAS.

NITRATE OF IRON.

R. Red oxide of iron, four drachms.
Nitric acid, six fl. drachms.

Dissolve, and add

Water, one fl. ounce.

Filter. Dose, six to twelve drops, in a convenient vehicle. In dyspepsia, and calculous complaints. *Swediaur.*

SYRUP OF PROTO-NITRATE OF IRON.

R. Iron wire (card teeth), two ounces.
Nitric acid (*sp. gr.* 1.42), three fl. ounces.
Water, thirteen fl. ounces.
Powdered sugar, two pounds.

Put the iron in a wide-mouthed bottle, kept cool by standing in cold water, and pour upon it three fl. ounces of water. Mix the acid with ten fl. ounces of water, and add it gradually to the iron, agitating frequently, until the acid is saturated. Filter the solution into a bottle containing the sugar, and marked to contain thirty fl. ounces, adding water enough to measure that quantity. Dissolve the sugar, strain, pour into suitable phials, and seal them. *W. Procter, Jr.*

SOLUTION OF PER-SESQUINITRATE OF IRON.

R. Iron wire (cut small), one ounce and a half.
Nitric acid, three fl. ounces.
Water, fifteen fl. ounces.

Dissolve, decant, and filter; add

Muriatic acid, one fl. drachm,

diluted with sufficient water to make the whole thirty fl. ounces. Dose, ten drops several times a-day, gradually increasing. Has been of great benefit in chronic diarrhoea, &c. *Kerr.*

R. Iron wire, in pieces, one ounce.
Nitric acid (*sp. gr.* 1.42), three fl. ounces.
Distilled water, sufficient.

Mix the acid with a pint of water, add the iron, and agitate occasionally, until gas ceases to come off; then filter the solution, and add sufficient water to make it measure thirty fl. ounces. *U. S. Ph.*

FERRI OXIDUM HYDRATUM.

HYDRATED OXIDE OF IRON.

R. Sulphate of iron, four ounces.
Sulphuric acid, three fl. drachms and a half.
Nitric acid, six fl. drachms, or sufficient.
Solution of ammonia, sufficient.
Water, two pints.

Dissolve the sulphate in the water, gradually add the sulphuric acid, boil, add nitric acid very gradually, boiling after each addition, till no dark color is produced. Filter, allow to cool, add solution of ammonia, in excess, stirring well. Wash precipitate with water, as long as washings give a precipitate with chloride of barium. Keep in close bottles, under water.

U. S. Ph.

Used in large doses, as an antidote to arsenic.

SOLUTION OF TERSULPHATE OF IRON.

R. Crystallized sulphate of iron, sixty-four ounces.
Sulphuric acid, seven fl. ounces.
Nitric acid (*sp. gr.* 1.38), twelve fl. ounces.
Water, sufficient.

Powder the sulphate of iron, mix the acids and five fl. ounces of water, put on a sand-bath, and add the powdered sulphate gradually, stirring until effervescence ceases. The dense solution is then to be diluted with water until it measures four pints and a half, and filtered through muslin.

When the hydrated oxide is wanted for use, an equal measure of commercial solution of ammonia causes the requisite precipitation. Thus:

R. Of above solution of tersulphate of iron, half a pint.
Water, two pints.

Mix, and add

Of solution of ammonia, half a pint.

Stir the mixture, and then throw the whole on a piece of strong muslin; express the liquor until the oxide of iron remains in the cloth of a pasty consistence. Finally, wash the oxide with water. *W. Procter, Jr.*

The above formula is offered to the pharmacist as an *instantaneous means* of preparing the hydrated sesquioxide of iron, in cases of poisoning by arsenic. The solution of tersulphate of iron should always be kept on hand.

FERRI OXIDUM NIGRUM

BLACK OXIDE OF IRON.

R. Sulphate of iron, six ounces
Sulphuric acid, two fl. drachms, two fl. scruples

Pure nitric
acid, four fl. drachms and a half.
Stronger water of am-
monia, four fl. ounces and a half.
Boiling water, three pints.

Dissolve half the sulphate in half the water, add the sulphuric acid, and boil; add the nitric acid, gradually, boiling briskly after each addition. Dissolve the rest of the sulphate in the remainder of the water, mix the two solutions, and immediately add the ammonia, stirring briskly. Collect the precipitate on a muslin filter, wash in water till the washings scarcely give a precipitate with nitrate of baryta; dry at a low temperature. *Ed. Ph.*

R. Wash blacksmiths' scales with water; dry; separate metallic iron by means of a magnet; reduce to powder, and proceed as for prepared chalk. *Dub. Ph., 1826.*

R. Iron filings, any quantity; place in an earthenware pan, wet thoroughly with water; in an hour heap up on an iron plate; twelve hours afterwards moisten with hot water, and repeat this in twelve hours; and, a day afterwards, triturate with cold water, decant the turbid water, let it settle, collect the deposit on a filter, and, when perfectly drained, dry by a mild heat. *Par. Cod.*

These do not give identical products, but they are sufficiently alike to be used for the same purposes.

POWDER OF BLACK OXIDE OF IRON.

R. Black oxide of iron, three parts.
Aromatic powder, six parts.
Sugar, eight parts.

Mix. Dose, thirty to forty grains, twice a-day, in leucorrhœa and rachitis. *Suediaur.*

R. Black oxide of iron,
Colombo, each, one drachm.
Rind of bitter orange, half an ounce.
Cinnamon, one scruple.

Mix, and divide into nine powders. Three to be given a-day, in chlorosis and amenorrhœa. *Brera.*

PILLS OF BLACK OXIDE OF IRON.

R. Black oxide of iron, eight grains.
Powdered saffron,
" valerian, each, sixteen grains.
Syrup of wormwood, sufficient.

Mix, and form eight pills; four to eight a-day, in amenorrhœa. *Foy.*

R. Black oxide of iron,
Powdered canella, each, one drachm.
Extract of ox bile,
" centaury, each, three drachms.

Mix, and form pills of four grains. Four to six a-day, in engorgements of the abdominal viscera. *Pierquin.*

ELECTUARY OF BLACK OXIDE OF IRON.

R. Black oxide of iron, half an ounce.
Carbonate of potassa, two scruples.
Prepared chalk,
Ginger, each, two drachms.
Syrup of orange-peel, sufficient.

Mix well. Dose, half a drachm, two or three times a-day, in chlorosis and leucorrhœa. *Suediaur.*

R. Black oxide of iron,
Soap, each, one drachm.
Extract of hemlock, twelve grains.
" squill, one scruple.
Oxymel, sufficient.

Mix well. Dose, half a drachm to a drachm, in dropsy, with engorgement of the abdominal viscera. *Brera.*

FERRI OXIDUM RUBRUM.

RED OXIDE OF IRON.

R. Sulphate of iron, any quantity; heat till water of crystallization is expelled; then roast by a powerful heat, till all acid is driven off; wash, till all traces of acid disappear; dry. *Dub. Ph., 1826.*

Astringent and tonic, and has been advised in neuralgia, &c., in doses of five to fifteen grains, but is inferior to the subcarbonate.

PILLS OF RED OXIDE OF IRON.

R. Red oxide of iron,
Powder of pot marigold,
Extract of pot marigold,
each, one drachm.
Mucilage of gum Arabic, sufficient.

Mix, and form ninety pills. Highly spoken of in cancerous affections; five to eight to be taken, three times a-day. *Rust.*

RED OXIDE OF IRON PLASTER.

R. Litharge plaster, two pounds.
Frankincense, half a pound.
Red oxide of iron, three ounces.

Melt the plaster and frankincense together, add the oxide, and stir well. *Dub. Ph., 1826.*

- R. Peroxide of iron, one ounce.
 Burgundy pitch, two ounces.
 Litharge plaster, eight ounces.

Melt the pitch and plaster, and add the iron, stirring constantly. *Dub. Ph.*, 1850.

As a strengthening plaster, in muscular relaxation and weakness.

- R. Lead plaster, twenty-four parts.
 Resin, six parts.
 Olive oil,
 Yellow wax, each, three parts.
 Red oxide of iron, eight parts.

Triturate the oxide with the oil, and add to a mixture of the other substances, melted together. *Guibourt.*

FERRI PHOSPHAS.

PHOSPHATE OF IRON.

- R. Sulphate of iron, five ounces.
 Phosphate of soda, six ounces.
 Water, one gallon.

Dissolve the sulphate and phosphate, each, in four pints of the water; mix the solutions, and let settle; decant, wash the precipitate with hot water, and dry with a gentle heat. *U. S. Ph.*

Used with advantage in amenorrhœa, and certain cases of dyspepsia. Dose, five to ten grains.

FERRI SULPHAS.

SULPHATE OF IRON.

- R. Iron wire, in small pieces, twelve ounces.
 Sulphuric acid, eighteen ounces.
 Water, one gallon.

Mix the acid and water, and add the iron; heat till effervescence ceases. Pour off the solution, add half a drachm of sulphuric acid, filter, the end of the funnel touching the bottom of the receiving vessel. Evaporate in a matrass, set aside to crystallize in a covered vessel, drain crystals in a funnel, dry, and keep in well-closed bottles. *U. S. Ph.*

Astringent and tonic. Dose, from one to five grains.

DRIED SULPHATE OF IRON.

- R. Sulphate of iron, at will.
 Heat over a moderate fire, in a porcelain vessel, till converted into a grayish-white mass, cool, and powder. *Ed. Ph.*

Three grains are equal to five of the undried.

POWDER OF SULPHATE OF IRON.

- R. Powdered sulphate of iron, four grains.
 " ginger, ten to sixteen grains.

Mix. To be taken twice a-day, in amenorrhœa, chlorosis, &c. *Saunders.*

- R. Powdered sulphate of iron, half an ounce.
 Powdered tansy, three ounces.

Mix. Dose, a drachm. As a vermifuge. *Dan. Ph.*

PILLS OF SULPHATE OF IRON.

- R. Dried sulphate of iron, two parts
 Extract of dandelion, five parts.
 Conserve of red roses, two parts.
 Powdered liquorice root, three parts.

Beat together, and form into five-grain pills. *Ed. Ph.*

- R. Sulphate of iron, two drachms.
 Extract of wormwood, half an ounce.
 Syrup of saffron, sufficient.

Beat into mass, and divide into one hundred and fifty pills. Four to five, three times a-day. In chlorosis, leucorrhœa, &c. *Werthof.*

- R. Sulphate of iron, } each,
 Kino, } one drachm
 Gentian, } and a half.
 Extract of gentian, }
 Turpentine, three drachms.
 Powdered mallow root, sufficient.

Beat into mass, and make two hundred and seventy pills. Ten, four times a-day, in secondary gonorrhœa. *Walch.*

BLAND'S ANTI-CHLOROTIC PILLS.

- R. Sulphate of iron, in fine powder, one ounce.
 Carbonate of potassa, in dry powder, one ounce.
 Mucilage of tragacanth,
 Powdered liquorice root, each, sufficient.

To make ninety-nine pills. Triturate well, and divide into pills. To be taken, one in the morning and evening, for the first three days, a third pill at noon for the next three days, and so increased to nine a-day. *Bland.*

PILLS OF IRON AND RHUBARB.

- R. Dried sulphate of iron, four parts
 Extract of rhubarb, ten parts.
 Conserve of red roses, five parts.

Beat into mass, and form five-grain pills.
 Tonic and laxative, in dose of two or three.

COMPOUND SULPHATE OF IRON PILLS.

- R. Sulphate of iron, one scruple
 Powdered senna, half a scruple

Powdered jalap,
 “ cream of tar-
 tar, each, half a scruple.
 “ ginger, twelve grains.
 Syrup, sufficient.
 Form mass, and divide into twenty-five pills. *Ellis.*

These are known as *Hooper's pills*, as are also the following:—

R. Barbadoes aloes, eight ounces.
 Sulphate of iron, four ounces.
 Extract of black hellebore,
 Myrrh,
 Soap, each, two ounces.
 Powdered canella,
 “ ginger, each, one ounce.
 Water, sufficient.

Beat into mass, and divide into two and a half grain pills. *Phil. Coll. Pharm.*

As laxative and emmenagogue.

R. Dried sulphate of iron, one scruple.
 Powdered aloes, two scruples.
 “ cloves, five grains.
 Venice turpentine, sufficient.

Form mass, and divide into twenty pills. One, three times a-day, as an emmenagogue. *Ellis.*

SOLUTION OF SULPHATE OF IRON.

R. Sulphate of iron,
 Alum, each, six ounces.
 Water, forty-eight ounces.

Dissolve, filter, and add

Sulphuric acid, four ounces.

In hemorrhages, in doses of ten or twelve drops in water. *Spielmann.*

SULPHATE OF IRON INJECTION.

R. Sulphate of iron, half to one drachm.
 Sage water, four ounces.
 Mucilage gum Arabic, half a drachm.

Mix. As an injection, in nasal and uterine hemorrhages. *Berends.*

COMPOUND IRON MIXTURE.

R. Myrrh, one drachm and a half.
 Oil of partridge berry, six drops.
 “ nutmeg, two drops.
 Carbonate of potassa, two scruples.
 White sugar, one ounce.
 Sulphate of iron, half a drachm.
 Distilled water, seven fl. ounces.

Rub down the myrrh and the oils with some of the water added very gradually, then add the carbonate of potassa and sugar, and lastly, the sulphate of iron, dissolved in the rest of the water, and immediately bottle. *Bond.*

MIXTURE OF SULPHATE OF IRON.

R. Sulphate of iron, one scruple.
 Sulphuric acid, four to six drops.
 White sugar, one drachm.
 Water, four fl. ounces.

Mix. A teaspoonful, with water, every two to four hours, as a tonic. *Ellis.*

R. Sulphate of iron, four grains.
 Elixir of vitriol, twenty drops.
 Distilled water, one fl. ounce.

Mix. A teaspoonful, in a wineglass of water, three times a-day. *Ellis.*

R. Powdered myrrh, one drachm.
 “ gum Arabic, two drachms.
 Syrup, one fl. ounce.
 Infusion of chamomile, six fl. ounces.

Mix well, and add

Powdered sulphate of iron, fifteen grains.
 Cinnamon water, one fl. ounce.

Mix. Two spoonfuls, every three hours, as an emmenagogue. *Frank.*

SYRUP OF SULPHATE OF IRON.

R. Sulphate of iron, two drachms.
 Water, one fl. ounce.

Dissolve, filter, and add

Syrup of gum Arabic, seventeen fl. ounces.

Mix. A tablespoonful, twice a-day, in chlorosis, leucorrhœa, &c. *Soubiran.*

FERRI SULPHURETUM.

SULPHURET OF IRON.

Heat an iron rod to a white heat, apply to it a roll of sulphur, receiving the sulphuret of iron in water; separate it from the sulphur, dry, and keep in a well-stopped bottle. *Dub. Ph.*

Used to make hydrosulphuric acid, by the addition of diluted sulphuric acid.

FERRI ET POTASSÆ TARTRAS.

TARTRATE OF IRON AND POTASSA.

R. Subcarbonate of iron, three ounces.
 Muriatic acid, ten fl. ounces.
 Solution of potassa, five pints and a half.
 Bitartrate of potassa, seven ounces and a half.
 Distilled water, one gallon and a half.

Mix the subcarbonate with the acid, digest for two hours, pour into a gallon of the water, let rest for an hour, decant, add solution of potassa,

wash precipitate, and whilst moist, mix with the bitartrate and half a gallon of the water; keep at a temperature of 140° for thirty hours, often stirring; filter, and evaporate on a water-bath, at same temperature, to dryness.

U. S. Ph., 1840.

R. Sulphate of iron, eight ounces.
Bitartrate of potassa, seven ounces.
Distilled water, half a gallon.

Prepare the hydrated oxide of iron from the sulphate. Mix the bitartrate of potassa with the water, heat to 140°, add the hydrated oxide, frequently stirring, until it is no longer dissolved. Then filter, and evaporate on a water-bath, to the consistence of syrup, and spread it out on plates, that it may dry in the form of scales.

U. S. Ph., 1850.

Dose, from ten grains to half a drachm, in solution, or with an aromatic or bitter, in bolus.

POWDER OF TARTRATE OF IRON AND COLOMBO.

R. Tartrate of iron and
potassa, two scruples.
Powdered colombo, half a drachm.

Mix, and divide into four powders. One, every three or four hours, in syrup. As a tonic.

Ellis.

EFFERVESCENT POWDER OF TARTRATE OF IRON.

R. Subcarbonate of mag-
nesia, two drachms.
Carbonate of potassa, half an ounce.
Tartrate of iron, two scruples.

Teaspoonful in water in divided doses, during the day; to be taken whilst effervescing. *Radius.*

SOLUTION OF TARTRATE OF IRON AND POTASSA.

R. Tartrate of iron and
potassa, one ounce.
Cinnamon water, one pint.

Dissolve; as a lotion or fomentation for bruises. Internally, in leucorrhœa, chlorosis, &c.

Swedjaur.

COMPOUND WINE OF TARTRATE OF IRON.

R. Subcarbonate of
iron, one ounce and a half.
Powdered gentian,
" orange peel, each,
half an ounce.
Red wine, two pints.

Macerate for three days, and give a wine-glassful two or three times a-day. In dyspepsia, &c. *Ellis.*

R. Subcarbonate of iron, two ounces.
Powdered cinnamon, one ounce.
Rhenish wine, two pints.

Macerate for some days, and filter. *Van Mons.*

WINE OF TARTRATE OF IRON.

R. Subcarbonate of iron, one part.
Rhenish wine, twenty-four parts.

Macerate for some days, and filter. Dose, one or two ounces. *Soubiran.*

COMPOUND TINCTURE OF TARTRATE OF IRON.

R. Tartrate of iron wine,
Tincture of calamus,
Hoffmann's balsamic elixir, each,
four fl. drachms.

Mix. Dose, seventy drops, morning and evening, in wine. Said to be an excellent remedy in ascarides. *Dorfmüller.*

FERRI AMMONIO-TARTRAS.

AMMONIO-TARTRATE OF IRON.

R. Tartaric acid, one part.
Boiling water, sufficient

to dissolve; add

Iron filings, three parts.

Digest in a warm place, for two or three days. Add caustic ammonia in slight excess, stir well, pour off the solution, set it aside, decant the clear liquid, and evaporate to dryness; redissolve in distilled water, add a little more ammonia, filter, and evaporate, in shallow, porcelain dishes, by a gentle heat, till it becomes brittle, then chip it off with a blunt-pointed knife. *Aikin.*

R. Tartaric acid, fifty drachms.
Water, one gallon.

Saturate with

Carbonate of ammonia,

and add fifty drachms more of acid. Heat the solution in a water-bath, and add moist hydrated oxide of iron (derived from fifty-three and a half drachms of subcarbonate of iron dissolved in muriatic acid, and precipitated by ammonia.) Digest till the oxide is dissolved; filter, and evaporate to dryness by means of a water-bath.

W. Procter.

Has the same properties as the other preparations of iron, but has the advantage of being readily soluble, of being more palatable, and very permanent. The dose is five grains or more, in pills or solution.

SOLUTION OF AMMONIO-TARTRATE OF IRON.

R. Ammonio-tartrate of iron, half a drachm.
 Distilled water, seven fl. drachms.
 Rectified spirit, one fl. drachm.
 Mix, and dissolve. *Aikin.*

FERRI TANNAS.

TANNATE OF IRON.

R. Tannic acid, ninety parts.
 Boiling water, sufficient
 to dissolve; add gradually
 Subcarbonate of iron,
 four hundred and forty parts.

Agitate till effervescence ceases. Evaporate at 176°, in a porcelain vessel, until it becomes thick, then dry on a plate, in a stove, at 95°.

Benedetti.

Found useful in chlorosis, in doses of two or three grains, thrice a-day, increasing the dose as required.

FERRI VALERIANAS.

VALERIANATE OF IRON.

R. Clean iron filings, at will.
 Put in a wedgewood mortar, add gradually an equal weight of valerianic acid, stirring continually. In an hour, add distilled water, pour the whole into a flask, warm gently, and filter. The surface, exposed to the air, becomes covered with a crystalline layer of valerianate of iron. Collect this on a filter, repeat this as long as crystals are formed.

Dose, one to two grains. *Ruspini.*

R. Convert four ounces of sulphate of iron into the persulphate, (as directed in the formula for the hydrated peroxide of iron, and add water to make the solution measure eight fl. ounces. Then dissolve five ounces, three drachms, of valerianate of soda in half a pint of water, and mix the two solutions. Wash the precipitate on a filter, and dry it by placing it for some days, rolled in bibulous paper, on porous brick.

It should be kept in a well-stopped bottle.

Dub. Ph.

FICUS.

FIGS.

Figs are the fruit of *Ficus carica*, a small tree, a native of Asia, and now cultivated in all warm climates. There are numerous varieties.

Sex. Syst. Polygam. diœc. *Nat. Syst.* Moraceæ.

Linn. Sp. Pl. 1513. Griffith, Med. Bot. 576.

They come to this country in a dry state, from the ports of the Levant. They are nutritious, laxative, and demulcent, and are used in medicine in demulcent decoctions, and sometimes externally, as a suppurative poultice.

DECOCTION OF FIGS.

R. Figs, one ounce.
 Water, one pint.
 Boil, and strain. Demulcent and pectoral.
Cadet.

GARGLE OF FIGS.

R. Figs,
 Mallow root, each, one ounce.
 Milk, two pints.
 Boil away one-fourth. As an emollient gargle in sore throat. *Radius.*

COMPOUND DECOCTION OF FIGS.

R. Figs,
 Raisins, each, two ounces.
 Barley water, two pints.

Boil for a quarter of an hour, and add

Liquorice root, half an ounce.

Infuse, and strain.

Foy.

This is identical with the compound decoction of barley, of the Lond. Ph., and like it, is nutritive, demulcent, and laxative.

COFFEE OF FIGS.

R. Figs, roasted like coffee, two drachms.
 Water, one pint.

Boil gently, and strain.

Radius.

This is said to be an excellent substitute for coffee, and to form an excellent diet in diseases of the chest.

PASTE OF FIGS.

R. Figs, at will.
 Reduce to a pulp, express through a hair-sieve, mix with four times the weight of sugar, concentrate by a gentle heat, and cut into lozenges.
Soubeyran.

FILIX MAS.

MALE FERN.

The Male Fern is the rhizome of *Nephrodium (Aspidium) filix mas*, a native of the temperate regions of the whole world, in shady, rather damp situations.

Sex. Syst. Crypt. fil. *Nat. Syst.* Polypodiaceæ.
 Linn. Sp. Pl. 1551. (Polypodium). Griffith, Med. Bot. 670.

The rhizome is of a brown color, externally, with a feeble, but peculiar odor, and a sweet, bitter, astringent taste. It should always be used as fresh as possible. It is tonic and anthelmintic, but is not much used in this country. The dose, in substance, is from one to three drachms, twice a-day.

POWDER OF MALE FERN.

R. Male fern, one ounce.
Gamboge,
Calomel, each, fourteen grains.
Scammony, eighteen grains.
Mix, and divide into three powders. In tape-worm. *Cadet.*

BOLUS OF MALE FERN.

R. Male fern, half an ounce.
Rhubarb, two drachms.
Semen contra, half an ounce.
Calomel, one drachm.
Syrup of wormwood, sufficient.
Mix, and form twelve boluses. *Spielmann.*

ETHEREAL EXTRACT OF MALE FERN.

R. Powdered male fern, at will.
Exhaust it in a displacement apparatus, with ether, and distil off the ether. Dose, from fifteen to twenty grains, twice a-day.

PILLS OF EXTRACT OF MALE FERN.

R. Ethereal extract of
male fern, one scruple.
Powder of male fern, twelve grains.
Conserve of roses, sufficient.
Mix, and make twelve to sixteen pills. To be taken in the evening, at short intervals. *Peschier.*

OIL OF MALE FERN.

R. Powdered male fern, four parts.
Alcohol, eight parts.
Digest for twenty-four hours, decant, repeat the operation with five parts of alcohol, unite the liquids, distil off the alcohol, mix the residue with three parts of water, distil, and permit to rest, and collect the oil which sinks to the bottom, and wash it with pure water. *Magendie.*

PILLS OF OIL OF MALE FERN.

R. Oil of male fern, twenty-four drops.
Powdered mallow root, three scruples.
Fresh lard, two scruples.
Mix, and divide into twenty-four pills. One-half to be taken in the evening, and the remainder next morning, and, an hour afterwards, a dose of castor oil. *Jourdan.*

ELECTUARY OF MALE FERN.

R. Ethereal extract of
male fern, half a drachm.
Honey of roses, one ounce.
Mix. Half in the evening, the remainder the next morning. *Radius.*

JELLY WITH MALE FERN.

R. Male fern,
Corsica moss, each, six drachms.
Isinglass, one drachm.
Water, eight ounces.
Boil to three ounces, and strain; add
Ethereal extract of
male fern, one scruple.
Syrup of mulberries, one ounce.
To be taken in divided doses, during the day. *Phabus.*

MIXTURE OF MALE FERN.

R. Male fern, half an ounce.
Elecampane, two drachms.
Wormwood, a handful.
Semen contra, one drachm.
Water, sufficient
to obtain a quart of decoction; add
Vermifuge syrup, one ounce.
Mix. *Bories.*
R. Powdered male fern, four drachms.
Balm water, three ounces.
Syrup of gum, one ounce.
Mix. To be taken in the evening, and, next morning an ounce of castor oil. *Radius.*

FÆNICULUM.

FENNEL.

There are several varieties, or species of Fennel, closely resembling each other, and having the same properties; that recognized as officinal by the U. S. Pharm., is the *F. vulgare*, a native of Europe, and often cultivated in the gardens in this country.

Sex. Syst. Pentand. digyn. *Nat. Syst.* Apiaceæ.

Linn. Sp. Pl. 377. *Griffith, Med. Bot.* 321.

The part used is the seeds. These are fragrant, with a sweetish, warm, pleasant, aromatic taste. They are employed as a carminative, and as corrective of other, and more unpleasant remedies. The dose of the powder, is from a scruple to half a drachm.

COMPOUND POWDER OF FENNEL.

R. Powdered fennel seed, }
" anise, } each, one
" lettuce, } ounce and
" white poppy, } a half.
" benne,
" sugar, six ounces.

Mix. Dose, a drachm two or three times a-day. Said to increase the secretion of milk in nursing women. *Niemann.*

INFUSION OF FENNEL.

R. Fennel seed, one drachm.
Boiling water, one pint.

Infuse for two hours, and strain. Given in teaspoonful doses to infants, in windy colic, or used as an enema, for the same purpose. *Taddei.*

FENNEL WATER.

R. Oil of fennel, half a fl. drachm.
Carbonate of magnesia, one drachm.
Distilled water, two pints.

Rub the oil with the magnesia, gradually add the water, constantly rubbing, and filter.

U. S. Ph.

OIL OF FENNEL.

R. Bruised fennel seed, at will.
Water, sufficient.

Mix, and distil. *Guibourt.*

Dose, five to fifteen drops.

ESSENCE OF FENNEL.

R. Oil of fennel, one fl. ounce.
Alcohol, nine fl. ounces.

Mix. Dose, twenty or thirty drops. *Dub. Ph.*

FENNEL OINTMENT.

R. Oil of fennel, thirty drops.
Lard, four ounces.

Triturate together. Said to destroy lice in the hair. *Niemann.*

FRASERA.

AMERICAN COLOMBO.

The *Frasera Walteri* is one of the tallest and most elegant of our native herbaceous plants, sometimes attaining a height of ten feet, with a pyramid of yellowish-white flowers, of three or four feet in length.

Sex. Syst. Tetrand. monog. Nat. Syst. Gentianaceae.

Walter, Fl. Carol. 87. Griffith, Med. Bot. 462.

The root, which is the officinal portion, is large and succulent, and is usually dried in circular sections, like colombo. It has no odor in a dried state, but possesses a sweetish-bitter taste. It is a mild bitter tonic, suited to all cases to which the simple bitters are applicable. The dose, in powder, is from thirty grains to a drachm.

INFUSION OF AMERICAN COLOMBO.

R. American colombo, bruised, one ounce.
Boiling water, one pint.

Infuse for an hour, and strain. *Wood.*

Dose, one to two fl. ounces several times a-day.

TINCTURE OF AMERICAN COLOMBO.

R. Contused American colombo, one ounce.
Diluted alcohol, one pint.

Macerate for fourteen days, and filter. Dose, one to three drachms. *Dunlison.*

FULIGO.

SOOT.

The soot of wood is of a black color, has a peculiar smell, and a bitter, disagreeable, empyreumatic taste. It owes its properties to the presence of pyrogenous oil or resin, and creasote; but also contains various salts, united to acetic acid. It was formerly officinal, but gradually was relinquished in regular practice; but, of late years, has again attracted attention, and has been employed with success in a variety of diseases.

PILLS OF SOOT.

R. Extract of soot, half a drachm.
Comp. galbanum pill, one scruple.
Oil of valerian, ten minims.

Mix well, and form twelve pills; two, thrice a-day, in hysteria. *Neligan.*

R. Soot, one drachm and a half.
Carbonate of ammonia, half a drachm.
Tar, sufficient.

Mix, and make five-grain pills. *Ferriar*

DECOCTION OF SOOT.

R. Soot, two handfuls.
Water, two pints.

Boil for half an hour, and strain. As a lotion, in herpetic, psoric, and venereal ulcers.

Soubeiran.

TINCTURE OF SOOT.

R. Soot, one part.
Diluted alcohol, eight parts.

Macerate for eight days, and filter. Dose, from thirty to sixty drops, as an antispasmodic, in hysteria, &c. *Guibourt.*

MIXTURE OF SOOT.

R. Soot, half an ounce.
 Carbonate of potassa, one ounce and a half.
 Carbonate of ammonia, two drachms.
 Elder-flower water, nine fl. ounces.

Digest for some time by a gentle heat, and filter. Dose, thirty to sixty drops, several times a-day. In dyspepsia, suppressed cutaneous affections, &c. *Dunglison.*

R. Hickory ashes, one quart.
 Soot, six ounces.
 Boiling water, one gallon.

Mix, and digest for twenty-four hours, frequently stirring, and decant. A teacupful, three times a-day, used in dyspepsia. *Ellis.*

SOOT MOUTH-WASH.

R. Extract of soot, one part.
 Vinegar, twelve parts.

Mix. A few drops, in a glass of tepid water, in aphthous sore mouth. *Soubeiran.*

OIL OF SOOT.

R. Soot, four ounces.
 Olive oil, twelve ounces.

Boil, and strain. As an application to ulcers in the mouth, or on the genitals. *Radius.*

SOOT OINTMENT.

R. Soot, one part.
 Lard, four parts.

Triturate together. As an application in tinea, scrofulous sores, &c. *Soubeiran.*

COMPOUND SOOT OINTMENT.

R. Soot, half an ounce.
 Lard, each, half an ounce.
 Extract of belladonna, one drachm.

Triturate together. As an application to cancerous sores. *Radius.*

R. Soot, an ounce and a half.
 Lard, four ounces.
 Sulphate of zinc, six drachms.

Triturate together. In cases of tinea. *Borics.*

R. Soot, Powdered Peruvian bark, each, half an ounce.
 Powdered charcoal, " sulphur, each, one ounce.
 Lard, sufficient

to make an ointment. In tinea. *Radius.*

CATAPLASM OF SOOT.

R. Soot, two ounces.
 Whites of eggs, six.

Mix. As an application to tinea, herpes, and scrofulous ulcers. *St. Marie.*

INJECTION OF SOOT.

R. Decoction of soot, one pint.
 Alum, half an ounce.
 Water, six ounces.

Mix. Advised as an injection, in leucorrhœa. *Soubeiran.*

FULIGOKALI.

FULIGOKALI.

R. Potassa, twenty parts.
 Shining soot, one hundred parts.
 Water, sufficient.

Boil for an hour, cool, dilute with water, filter, evaporate to dryness, and keep in well-closed bottles. *Deschamps.*

Used, both internally and externally, in chronic cutaneous disorders. Dose, two to three grains, several times a-day.

SULPHURETTED FULIGOKALI.

R. Fuligokali, sixty parts.
 Potassa, fourteen parts.
 Sulphur, five parts.

Dissolve the sulphur and potassa, add the fuligokali, and evaporate to dryness.

PILLS OF SULPHURETTED FULIGOKALI.

R. Sulphuretted fuligokali, five drachms.
 Starch, two drachms and a half.
 Tragacanth, eight grains.
 Syrup, sufficient.

Mix, and make one hundred pills, and cover them with two or three coats of tragacanth. *Deschamps.*

OINTMENT OF FULIGOKALI.

R. Fuligokali, sixteen to thirty grains.
 Lard, one ounce.

Triturate together. Used as a detergent, resolvent, and somewhat stimulant application, in obstinate cutaneous affections. *Gibert.*

G.

GALBANUM.

GALBANUM.

This is the concrete juice of an unknown plant, but generally admitted to belong to the Apiaceæ. According to Don, and others, the Asiatic galbanum is the product of *Galbanum officinale*, a native of Persia.

Don, Trans. Linn. Soc. xvi. 603. Griffith, Med. Bot. 333.

Galbanum is in whitish or yellowish tears, or in brownish-yellow, or brownish masses, formed of these tears agglutinated, and containing various impurities. The odor is peculiar, and somewhat balsamic; the taste is hot, bitter, and acrid. It is stimulant, expectorant, and antispasmodic, in doses of ten to twenty grains; and also used externally to indolent swellings, &c.

PURIFIED GALBANUM.

R. Galbanum, one part.
Alcohol, four parts.

Dissolve by means of a water-bath, strain, treat the residue with a fresh quantity of alcohol, unite the two solutions, distil off the alcohol, and evaporate the residue. *Guibourt.*

COMPOUND GALBANUM PILLS.

R. Galbanum, six drachms.
Myrrh, each, two drachms.
Assafetida, sufficient.
Syrup,

Beat into mass, and divide into two hundred and forty pills. *U. S. Ph.*

Dose, ten to twenty grains, in chlorosis and hysteria.

R. Galbanum, two drachms.
Oil of savine, four to twelve drops.
Malate of iron, one drachm and a half.
Powdered valerian, sufficient.

Beat into mass, and divide into one hundred and twenty pills. Two, three times a-day, in amenorrhœa, chlorosis, &c. *Phœbus.*

GALBANUM MIXTURE.

R. Galbanum, two drachms,
Vinegar of squills,
Fennel water, each, two fl. ounces.
Spirit of Mindererus, two fl. drachms.
Nitric ether, one fl. drachm.
Syrup of mallows, half fl. ounce.

Mix. Three or four spoonfuls a-day, in humid asthma. *Jahn.*

TINCTURE OF GALBANUM.

R. Galbanum, in small pieces, two ounces.
Proof spirit, two pints.
Digest for seven days, and filter.

Dub. Ph., 1826.

Antispasmodic, &c. In doses of one to three fl. drachms.

COMPOUND TINCTURE OF GALBANUM.

R. Tincture of galbanum,
" wormwood,
Spirit of rosemary, equal parts.
Mix. Dose, forty to fifty drops. *Augustin.*

GALBANUM PLASTER.

R. Lead plaster, two pounds.
Galbanum, half a pound.
Yellow wax, four ounces.

Melt the galbanum, add the plaster and wax, and melt the whole, and strain. *Dub. Ph., 1826.*

COMPOUND GALBANUM PLASTER.

R. Galbanum, eight ounces.
Turpentine, ten drachms.
Burgundy pitch, three ounces.
Lead plaster, three pounds.

Melt the galbanum and turpentine together, strain, add the pitch, and then the lead plaster, melted; mix well. *U. S. Ph.*

A good application to chronic, scrofulous enlargements of the glands and joints.

R. Galbanum and saffron
plaster, eight parts.
Camphor,
Petroleum,
Subcarbonate of ammonia, each,
one part.

Melt together. In chronic rheumatism. *Augustin.*

R. Galbanum,
Turpentine, each, one part.
Melt, strain, and add a melted mixture of
White wax, four parts.
Spermaceti, two parts.
Oil of almonds, one part.
Mix well. A milder application than any of the above. *Spielmann.*

GALIIUM VERUM.

YELLOW LADIES-BEDSTRAW.

A perennial, herbaceous plant, native of Europe, which is stated to have the power of coagulating milk, and also used to dye yellow.

Sex. Syst. Tetrand. monog. *Nat. Syst.* Rubiaceæ.

Linn. Sp. Pl. 155. *Stokes, Bot. Mat. Med.* i. 202.

The whole plant is used; it has an agreeable odor, and an astringent, acidulous, somewhat acrid taste. It was, at one time, much used in epilepsy and hysteria, and also as an application in cutaneous affections.

INFUSION OF YELLOW LADIES-BED-STRAW.

R. Flowers of yellow ladies-bedstraw, two drachms.
Boiling water, two pints.

Digest for an hour, and strain. To be taken in a tepid state, in wineglassful doses, as a sedative and antispasmodic, in cerebral affections. *Chaussier.*

WINE OF YELLOW LADIES-BEDSTRAW.

R. Expressed juice of yellow ladies-bedstraw, six fl. ounces.
White wine, one to two fl. ounces.

Mix. To be taken in epilepsy, as soon as indications of an attack manifest themselves.

Bories.

GALLA.

GALLS.

Galls are morbid excrescences on *Quercus infectoria*, caused by the puncture of a hymenopterous insect, the *Diplolepis gallæ tinctoriæ*. They come from the Levant ports, and present several varieties, the best of which are dark colored. They are round, hard, solid, brittle, inodorous, and have a bitter, very astringent taste. They contain tannic and gallic acids. They are powerfully astringent. They are more used as external applications, than as internal remedies. Dose, ten to thirty grains, several times a-day.

COMPOUND POWDER OF GALLS.

R. Powdered galls,
" kino, each, half an ounce.
" alum, two drachms.

Mix. As a styptic. *Radius.*

DECOCTION OF GALLS.

R. Galls, half an ounce.
Water, one pint and a half.
Boil to a pint, and strain.

R. Bruised galls, two ounces and a half.
Water, two pints.
Boil to a pint, and strain. *Lond. Ph.*

INFUSION OF GALLS.

R. Powdered galls, one to two drachms.
Boiling water, one pint.
Digest for half an hour, and strain. Used for injections, fomentations, gargles, &c. *Taddei.*

COMPOUND INFUSION OF GALLS.

R. Infusion of galls, four fl. ounces.
Prepared chalk, half an ounce.
Tincture of opium, half fl. drachm.
Powdered gum Arabic, one drachm.
Mix. Dose, a tablespoonful every two hours.
In diarrhœa. *Ellis.*

GARGLE OF GALLS.

R. Infusion of galls, seven fl. ounces.
Alcohol, one fl. ounce.
Mix. As a gargle in relaxation of uvula and soft palate. *Ellis.*

LOTION OF GALLS.

R. Claret wine, one pint.
Common salt, one drachm.
Sulphate of iron, two drachms.
Boil for a few minutes, and add
Oxide of copper, one drachm.
Boil for two minutes. Add

Powdered galls, two drachms.
This is used to color the hair; it is to be washed with this fluid, dried with a warm cloth, and then washed with common water. *Laforest.*

SYRUP OF GALLS.

R. Bruised galls, two drachms.
Brandy, eight fl. ounces.
Mix. Burn out the spirit, and add
Powdered cinnamon,
" mace, each, two drachms.
While hot, add sugar by burning it over the inflamed brandy, till a syrup is formed. Decant. Dose, a tablespoonful every two hours.
In obstinate chronic diarrhœa. *Parrish.*

TINCTURE OF GALLS.

R. Bruised galls, four ounces.
Diluted alcohol, two pints.
Macerate for fourteen days, express, and filter. *U. S. Ph.*
A powerful astringent, in doses of one to three fl. drachms.

R. Bruised galls, one pound.
 Water, two pints.
 Macerate twenty-four hours, and add
 Alcohol, two pints.
 Eau de Cologne, six fl. ounces.
 Filter. As an injection, diluted with water, in
 leucorrhœa and gonorrhœa. *Gibert.*

EXTRACT OF GALLS.

R. Galls, in coarse powder, at will.
 Macerate in sufficient water to cover them for
 twenty-four hours; then introduce the moist
 mass into a displacer, and act on it with water,
 until the galls are nearly exhausted; evaporate
 the infusion to dryness, on a water-bath.

OINTMENT OF EXTRACT OF GALLS.

R. Extract of galls, one drachm.
 Lard, one ounce.
 Triturate the extract with a little water, until it
 is reduced to a uniform syrupy consistence, then
 add the lard, and mix thoroughly. *D. S. Jones.*

OINTMENT OF GALLS.

R. Galls, in very fine powder, one ounce.
 Lard, seven ounces.
 Mix. As an application to piles, prolapsus ani,
 and to indolent ulcers. *U. S. Ph.*

COMPOUND OINTMENT OF GALLS.

R. Galls, in very fine powder, two drachms.
 Powdered opium, one drachm.
 Lard, one ounce.
 Mix well. *Ed. Ph.*
 As an application to irritable piles, &c.

R. Powdered galls,
 Tincture of opium, each, two parts.
 Camphor, one part.
 Lard, eight parts.

Triturate well together. As an application to
 painful piles. *Foy.*

R. Finely powdered galls, one scruple.
 Simple cerate, one ounce.
 Essence of lemon, twenty drops.
 Lead water,
 Laudanum, each, forty drops.

Rub well together. As an application to
 hemorrhoids, after subsidence of inflammation.
Devees.

GAMBOGIA.

GAMBOGE.

This is the concrete juice of several species
 of East Indian trees, probably belonging to the
 genus *Hebradendron*.

Sex. Syst. Polyand. monog. *Nat. Syst.* Clu-
 siaceæ.

Graham, *Comp. Bot. Mag.* ii. 199. Griffith,
Med. Bot. 152.

Gamboge is a gum resin, of a yellow-orange
 color, opaque, brittle, almost inodorous, of an
 insipid taste at first, but soon becoming acrid.
 It is a powerful hydragogue and drastic purga-
 tive, but apt to produce nausea and vomiting.
 Dose, from one to six grains, in pill, emulsion,
 or alkaline solution.

COMPOUND POWDER OF GAMBOGE.

R. Bitartrate of potassa, two drachms.
 Powdered jalap, one drachm.
 " gamboge, six grains.

Mix, and divide into six powders. One to be
 given every two or three hours, in syrup, as a
 hydragogue purgative. *Ellis.*

R. Powdered gamboge, two grains.
 " sulphate of iron, six grains.
 " sugar, one scruple.
 Oil of peppermint, three drops.

Mix. To be taken twice a-day, against tape-
 worm. *Vogt.*

COMPOUND PILLS OF GAMBOGE.

R. Powdered gamboge, one drachm.
 " aloes, one drachm and
 a half.
 " ginger, half a drachm.
 Soft soap, two drachms.

Beat into mass. *Lond. Ph.*
 Dose, ten to twelve grains, as a purgative.

R. Powdered gamboge, ten grains.
 " guaiacum,
 Blanched almonds, each, one drachm.
 Syrup, sufficient.

Beat together, and form twenty-eight pills.
 Two to four, as a gentle purgative. *Saunders.*

ALKALINE SOLUTION OF GAMBOGE.

R. Powdered gamboge, one drachm.
 Carbonate of potassa, three drachms.
 Water, six fl. drachms.

Dissolve. Dose, fifteen drops, every three hours,
 as a hydragogue in dropsies. *Van Mons.*

GAMBOGE MIXTURE.

R. Powdered gamboge, four grains.
 Spirit of nitric ether, one fl. drachm.
 Tincture of senna, two fl. drachms.
 Mint water,
 Syrup of buckthorn, each,
 half a fl. ounce

Mix. As a hydragogue purgative, in dropsy,
 especially in hydrothorax. *Ferriar*

- R. Gamboge, two scruples.
 Tartrate of potassa, one ounce.
 White sugar, two drachms.
 Water, six ounces.

Make a solution. As a purgative, in ascites. A tablespoonful, every two or three hours, until it operates. *Dewees.*

GAMBOGE AND ELATERIN MIXTURE.

- R. Gamboge, four grains.
 Elaterin, half a grain.
 Sweet spirit of nitre, one ounce.
 Water, four ounces.

Mix. As a hydragogue purgative, in ascites. A tablespoonful, to be taken every two or three hours, until free purgation is induced. *Dewees.*

ALKALINE TINCTURE OF GAMBOGE.

- R. Gamboge, one part.
 Carbonate of potassa, two parts.
 Diluted alcohol, twenty-four parts.
 Digest for some days, and filter. *Scherf.*

AMMONIACAL TINCTURE OF GAMBOGE.

- R. Gamboge, thirty-six grains.
 Spirit of ammonia, four fl. ounces.
 Digest for eight days, and filter. Dose, one to two fl. drachms. *Swediaur.*

GAULTHERIA.

PARTRIDGE BERRY.

The leaves of *Gaultheria procumbens*, a small, indigenous, evergreen plant, found in most parts of the United States, in dry, sandy situations.

Sex. Syst. Decand. monog. *Nat. Syst.* Ericaceæ.

Kalm, Amœn. Acad. iii. 14. Griffith, Med. Bot. 424.

These leaves have a peculiar, aromatic taste and odor, with a little astringency. They are stimulant, and somewhat tonic and astringent; and are much used in the form of tea, in bowel diseases, and as an emmenagogue.

OIL OF PARTRIDGE BERRY.

- R. Leaves of partridge berry, at will.
 Water, sufficient.
 Distil, and collect the oil that passes. *Coxe.*

Principally used to flavor mixtures, &c. In large doses, is poisonous. Dissolved in diluted alcohol, is in common use as a carminative and stimulant.

PARTRIDGE-BERRY WATER.

- R. Oil of partridge berry, sixteen drops.
 Carbonate of magnesia, one drachm.
 Water, one pint.

Triturate the oil, magnesia, and a little of the water together, then add the rest of the water, and filter.

GENTIANA.

GENTIAN.

The roots of several species of gentian are used in medicine; but that which is most generally recognized as officinal, is *Gentiana lutea*, a tall, perennial plant, native of the mountains of Europe.

Sex. Syst. Pentand. digyn. *Nat. Syst.* Gentianaceæ.

Linn. Sp. Pl. 329. Griffith, Med. Bot. 460.

The root is of a grayish-brown color externally, and yellowish within; it has a slightly sweetish, but extremely bitter taste, and a faint, but peculiar odor. It is a pure and simple bitter tonic, with no astringency. The dose of the powder is from ten to forty grains, but is seldom given in this form.

COMPOUND POWDER OF GENTIAN.

- R. Powdered gentian,
 " cascarilla,
 " orange peel, each,
 one drachm
 Peppermint sugar, three drachms.

Mix. Dose, thirty grains, several times a-day, as a stomachic. *Augustin.*

EXTRACT OF GENTIAN.

- R. Gentian, in coarse powder, one pound.
 Water, sufficient.

Exhaust the powder by means of displacement. Heat the filtered liquid to the boiling point, strain, and evaporate to proper consistence.

U. S. Ph.

Dose, ten to thirty grains, as a tonic; also as a basis for various metallic pills.

FLUID EXTRACT OF GENTIAN.

- R. Powdered gentian, sixteen ounces.
 Water, sufficient.
 French brandy, six fl. ounces.

Macerate the gentian for twelve hours in two and a half pints of water, introduce into a percolator, add water, and pass five pints of infusion. Evaporate this to ten fl. ounces on a water-bath, add the brandy, and strain through cotton flannel. Dose, half a fl. drachm.

W. Procter, Jr.

COMPOUND GENTIAN PILLS.

- R. Powdered aloes, one drachm.
 Extract of gentian, half a drachm.
 Oil of caraway, ten drops.
 Syrup, sufficient.

Beat into mass, and form pills of four grains. Two or three for a dose, as a purgative. *Ellis.*

Extract of gentian,
Powdered rhubarb,
Soap, equal parts.
Water, sufficient.

Beat into mass, and form four-grain pills. Tonic and laxative. *Fulda Dispen.*

INFUSION OF GENTIAN.

R. Contused gentian, two drachms.
Boiling water, one pint.
Infuse for twelve hours, and strain. *Beral.*

COMPOUND INFUSION OF GENTIAN.

R. Bruised gentian, half an ounce.
" orange peel,
" coriander, each, one drachm.
Diluted alcohol, four fl. ounces.
Water, twelve fl. ounces.

Pour on the alcohol, and, in three hours, add the water, macerate for twelve hours, and strain. *U. S. Ph.*

Dose, a fl. ounce, as a tonic.

R. Bruised gentian,
" calamus, each, one ounce.
Centaurry, six drachms.
Rosemary, two drachms.
Boiling water, four pints.

Infuse for twelve hours, and strain. Used as an emmenagogue, in doses of three ounces, twice a-day. *Spielmann.*

INFUSION OF GENTIAN WITH RHUBARB.

R. Bruised gentian, one drachm.
" rhubarb, two drachms.
Boiling water, twelve fl. ounces.

Macerate for an hour, strain, and add

Carbonate of ammonia, two scruples.

Dose, one to two fl. ounces, in dyspepsia, chronic rheumatism, &c. *Steph. and Church.*

MIXTURE OF GENTIAN.

R. Carbonate of magnesia, one drachm.
Infusion of gentian, six fl. ounces.

Mix. A wineglassful, three times a-day, as an antilithic, where uric acid abounds. *Brande.*

R. Extract of gentian, two drachms.
Carbonate of potassa, two scruples.
Mint water, five fl. ounces.
Tincture of orange peel, five fl. drachms.
Sugar, one ounce.

Mix. A spoonful, three times a-day, as a stomachic and carminative. *Berends.*

WINE OF GENTIAN.

R. Gentian, coarsely-powdered, half an ounce.
Yellow bark, coarsely-powdered, one ounce.
Orange peel, two drachms.
Canella, in coarse powder, one drachm.
Proof spirit, four fl. ounces and a half.
Sherry wine, thirty-six fl. ounces.

Digest the root and barks in the spirit, for twenty-four hours; add wine, digest for seven days, express, strain, and filter. *Ed. Ph.*

A stomachic bitter, in doses of four to eight fl. drachms.

SYRUP OF GENTIAN.

R. Gentian, in coarse powder, forty-eight parts.
Cold water, sufficient.
Syrup, fifteen hundred parts.

Put the gentian in a displacement apparatus, and make a strong infusion; filter this, and add to boiling syrup, of such a consistence, as, after the addition, it will make 30° Baumé. *Tauvel.*

SYRUP OF EXTRACT OF GENTIAN.

R. Aqueous extract of gentian, two drachms.
Water, eight fl. ounces.
Sugar, sixteen ounces (av.).

Dissolve the extract in the water, add the sugar, and form a syrup with a gentle heat, and strain. *W. Procter.*

TINCTURE OF GENTIAN.

R. Gentian, one part.
Diluted alcohol, four parts.

Macerate for six days, and filter. Dose, one to two fl. drachms. *Cottureau.*

ACIDULATED TINCTURE OF GENTIAN.

R. Compound tincture of gentian, four fl. ounces.
Elixir of vitriol, half fl. drachm.

Mix. A teaspoonful, in sugar and water, three or four times a-day. *Ellis*

COMPOUND TINCTURE OF GENTIAN.

R. Bruised gentian, two ounces.
Orange peel, one ounce.
Bruised cardamom, half an ounce.
Diluted alcohol, two pints.

Macerate for fourteen days, express, and filter. *U. S. Ph.*

An excellent bitter tonic, in doses of one fl drachm.

AMMONIACAL TINCTURE OF GENTIAN.

R. Bruised gentian, one ounce.
Carbonate of ammonia, two drachms.
Diluted alcohol, two pints.

Macerate the root in the alcohol for eight days, filter, and add the ammonia. Dose, one to four drachms, in scrofula. *Taddei.*

ALKALINE TINCTURE OF GENTIAN.

R. Bruised gentian, one ounce.
Carbonate of soda, three drachms.
Diluted alcohol, two pints.

Macerate for eight days, and filter. Used in same cases, and same doses, as the former. *Guibourt.*

MIXTURE OF TINCTURE OF GENTIAN.

R. Tincture of gentian, one fl. ounce and a half.
Tincture of rhubarb,
" cinnamon, each, half fl. ounce.
" opium, ten drops.

Mix. A dessertspoonful, every hour, in diarrhoea. *Foy.*

R. Extract of gentian,
" cascarilla, each, two drachms.
Peppermint water, four fl. ounces.

Mix. In spoonful doses, as a tonic and stomachic. *St. Marie.*

GENTIAN AND SULPHURIC ACID MIXTURE.

R. Compound infusion of gentian, five fl. ounces.
Compound tincture of gentian, one fl. ounce.
Diluted sulphuric acid, one fl. drachm.

Mix. A tablespoonful, three times a-day, in dyspepsia. *Brande.*

GENTIANIN.

GENTIANIN.

R. Powdered gentian, at will.

Macerate in cold ether, and repeat till exhausted; unite the tinctures, and distil; treat the residue several times with cold alcohol, distil the tinctures, evaporate the residue to dryness; mix it with an excess of magnesia, treat with sulphuric ether, then with oxalic acid, then again with the ether, and distil. *Magendie.*

Dose, one to two grains.

PILLS OF GENTIANIN.

R. Gentianin, five grains.
Conserve of roses,
Powdered liquorice, each, sufficient.
Mix, and make six pills.

SYRUP OF GENTIANIN.

R. Gentianin, sixteen grains.
Syrup, one pound.
Dissolve, and boil for a short time. Dose, half an ounce to an ounce. *Borics.*

TINCTURE OF GENTIANIN.

R. Gentianin, five grains.
Alcohol, one ounce.
Mix. Dose, from one to four drachms. *Cadet.*

GERANIUM.

CRANESBILL.

This is the root of *Geranium maculatum*, a native, perennial plant, found, in most parts of the country, in woods and shady places.

Sex. Syst. Monadelph. decand. *Nat. Syst.* Geraniaceæ.

Linn. Sp. Pl. 955. *Griffith, Med. Bot.* 209.

The root is in small rough pieces, of a dark-brown color externally, and flesh-colored within. It is inodorous; the taste is astringent, but not bitter. The dose is from ten to thirty grains, as an astringent, in all cases in which kino or rhatany is found useful.

DECOCTION OF GERANIUM.

R. Geranium, one ounce.
Water, one pint and a half.
Boil down to a pint. Dose, from one to two fl. ounces. *Wood*

EXTRACT OF GERANIUM.

R. Bruised geranium root, one pound.
Water, one gallon.
Boil down to one-half, strain, and evaporate to due consistence. This extract is very similar to that of Rhatany, and may be given in the same cases, and in the same doses.

GEOFFROYA INERMIS.

CABBAGE-TREE BARK.

The Cabbage-Tree is a native of the West India Islands; it is a large tree, having a smooth, gray bark.

Sex. Syst. Diadelph. decand. *Nat. Syst.* Fabaceæ.

Linn. Sp. Pl. 1043. *Griffith, Med. Bot.* 247. (Andira.)

The bark is in long, fibrous pieces, of a brownish ash color externally, and yellowish within; it has a resinous fracture, an unpleasant odor, and a sweetish, mucilaginous, bitterish taste. It is a powerful vermifuge, in doses of a scruple to half a drachm. The best form of administration is in syrup.

DECOCTION OF CABBAGE-TREE BARK.

R. Cabbage-tree bark, one ounce.
Water, two pints.

Boil down to one pint, and strain. Dose, one to two fl. ounces. *Ellis.*

EXTRACT OF CABBAGE-TREE BARK.

R. Cabbage-tree bark, one part.
Water, eight parts.

Boil for a quarter of an hour, strain; add four parts of water, boil, and strain; evaporate the united decoctions to a proper consistence. Dose, three grains. *Belg. Ph.*

MIXTURE OF CABBAGE-TREE BARK.

R. Cabbage-tree bark, half an ounce.
Water, six ounces.

Boil for half an hour, strain, and add
Tincture of cabbage-tree
bark, one ounce.
Syrup of orange peel, half an ounce.
A tablespoonful, morning and evening. *Niemann.*

GEUM.

AVENS.

1. GEUM RIVALE.

WATER AVENS.

This plant is a native of Europe and the United States, in wet, boggy situations.

Sex. Syst. Icosand. polyg. *Nat. Syst.* Rosaceæ.

Linn. Sp. Pl. 717. English Bot. 1. 106.

Root reddish or purplish, hard and brittle, inodorous, of an astringent, bitterish taste. It is tonic and astringent, and has been used advantageously in passive hemorrhages, leucorrhœa, &c. The dose, in powder, is from a scruple to a drachm, three times a-day.

DECOCTION OF WATER AVENS.

R. Water avens root, one ounce.
Water, one pint and a half.

Boil down to a pint, and strain. To be given in doses of one or two fl. ounces.

2. GEUM VIRGINIANUM.

WHITE AVENS.

Also a native of the United States, growing in woods and along streams. Flowers small and white.

Linn. Sp. Pl. 719. Griffith, Med. Bot. 279.

The root is of a brown color, somewhat aromatic, and of a bitterish, astringent taste. It has the same properties as the *G. rivale*, and is given in the same cases and doses.

3. GEUM URBANUM.

AVENS.

A native of Europe, growing in woods and waste places. The flowers are small, and of a yellow color.

Linn. Sp. Pl. 716. Eng. Bot. i. 1400.

The root consists of a short caudex with many long fibres. When fresh, it has somewhat the odor of cloves, but when dry, is almost inodorous. The taste is bitter and astringent. Like the other species, it is tonic and astringent.

COMPOUND POWDER OF AVENS.

R. Powdered avens root,
“ gum Arabic, each,
 three drachms.
“ kino,
“ cinnamon, each,
 one drachm.
“ sugar, half an ounce.

Mix. A teaspoonful every two or three hours, in diarrhœa. *Vogel.*

MIXTURE OF AVENS.

R. Bruised avens, one ounce.
Water, twelve fl. ounces.

Boil to eight ounces, and to the strained cold decoction, add

Extract of avens, two drachms.
Sulphuric ether, one drachm.

Mix. Two spoonfuls, every two hours. As a febrifuge. *Radius.*

GILLENIA.

INDIAN PHYSIC.

This is the root of *Gillenia trifoliata*, and *G. stipulacea*, native herbaceous plants, found in most parts of the country, in shady, and rather marsh situations.

Sex. Syst. Icosand. pentag. *Nat. Syst.* Rosaceæ.

Mœnch. Sp. Pl. 286. Griffith, Med. Bot. 282.

The roots are about as thick as a quill, wrinkled longitudinally, and irregularly undulated; a light brown color, and consisting of a thick cortical portion, and a slender liguous

centre. The bark is of a disagreeable, bitter taste. It is a mild and certain emetic, and forms a good substitute for ipecacuanha. The dose is from twenty to thirty grains, to be repeated in half an hour, if it does not produce the desired effect.

GLYCERINA.

GLYCERIN.

R. Lead plaster, recently prepared, and fluid,

Boiling water, each, one gallon.

Mix, stir briskly for fifteen minutes, cool, and pour off the supernatant liquid. Evaporate this to the specific gravity 1.15, and pass a current of sulphuretted hydrogen slowly through it while a black precipitate is thrown down. Filter, boil, and evaporate until the liquid ceases to lose weight. *U. S. Ph.*

In a pure state it is a nearly colorless, viscid fluid, and is uncrystallizable. *Sp. gr.* 1.25.

It has been used with success in lotions, poultices, baths, &c., to render them emollient and soothing; in diseases of the skin, as pityriasis, impetigo, chapped hands, nipples, face, &c.

LOTION OF GLYCERIN.

R. Glycerin, one fl. drachm.
Borax, ten grains.
Rose water, seven fl. drachms.

Mix. Used as a lotion in lichen, three times a-day; also in chapped hands. *Clymer.*

GLYCERIN OINTMENT.

R. Spermaceti, half an ounce.
White wax, one drachm.
Oil of almonds, two fl. ounces.
Glycerin, one fl. ounce.

Melt the wax and spermaceti with the oil of almonds at a moderate heat, add the glycerin, and rub until well mixed and cold.

Used to protect and soothe inflamed surfaces. *J. H. Ecky.*

R. White wax,
Spermaceti, each, one ounce and a half.
Lard, five ounces.
Glycerin, one fl. ounce.

Melt the wax and spermaceti, add the lard and stir till the mixture thickens, then add the glycerin and mix thoroughly.

Used as a substitute for spermaceti cerate. *J. Laidley.*

GLYCERIN PASTE.

R. Gum Arabic, one ounce.
Boiling water, two fl. ounces.
Glycerin, two fl. drachms.

Dissolve the gum in the water, add the glycerin, and strain if necessary. *P. B. Goddard.*

GLYCYRRHIZA.

LIQUORICE ROOT.

Several species of this genus have sweet and demulcent roots, but the only officinal species is the *G. glabra*, a native of the south of Europe, and of some parts of Asia and Africa, and also successfully cultivated in more northern regions.

Sex. Syst. Diadelph. decand. *Nat. Syst.* Fabaceæ.

Linn. Sp. Pl. 1046. *Griffith, Med. Bot.* 236.

The root is in long pieces of various thicknesses, of a grayish-brown color externally, and yellowish within; it is inodorous, and of a sweet, mucilaginous taste, with a slight degree of acidity. It is an excellent demulcent, and is much used in affections of the mucous membranes, and as an addition to the more irritating or nauseous remedies, to abate their acrimony or conceal their taste.

EXTRACT OF LIQUORICE.

R. Fresh liquorice root, bruised,
two pounds and a half.
Boiling distilled water, two gallons.

Macerate for twenty-four hours, boil down to a gallon, and strain, evaporate to proper consistence. *Lond. Ph.*

REFINED LIQUORICE.

R. Crude liquorice, at will.
Dissolve in water, strain, and evaporate. These two articles are useful demulcents, and additions to cough mixtures. *U. S. Disp.*

LIQUORICE PASTE.

R. Extract of liquorice, one pound.
Cold water, four pints.

Dissolve, strain, and add

Gum Arabic, two pounds.
Sugar, one pound.

Evaporate to proper consistence. *Soubeiran.*

R. Gum Arabic, two pounds.
Extract of liquorice, nine ounces.
Sugar, one pound.
Water, sufficient.

Evaporate to consistence of syrup, and add

Powdered orris root,
" elecampane, each half an ounce.

Continue to evaporate, then add

Oil of aniseed, two scruples.
Spielmann.

LOZENGES OF LIQUORICE.

R. Extract of liquorice,
Gum Arabic, each, six ounces.
Sugar, one pound.

Dissolve in sufficiency of boiling water, evaporate on water-bath, to proper consistence for lozenges.
Ed. Ph.

LOZENGES OF LIQUORICE AND OPIUM.

R. Powdered opium, half an ounce.
" liquorice,
" gum Arabic,
" sugar, each, ten ounces.
Oil of anise, one fl. drachm.

Mix the powder well, add the oil of anise, form mass with water, and divide into lozenges of six grains.
U. S. Ph.

Each lozenge contains one-tenth of a grain of opium.

R. Powdered liquorice,
" gum Arabic,
" sugar, each, two drachms.
" opium, six grains.
Oil of anise, four drops.

Mix, and with sufficient water form mass, and divide into sixty lozenges. These are known as *Dr. Wistar's cough lozenges*.
Ellis.

Both the above are demulcent and anodyne, and are well suited to allay cough, where opium is permissible.

MIXTURE OF LIQUORICE.

R. Powdered liquorice, one drachm.
Chamomile water, one fl. ounce.
Syrup of mallow, four fl. ounces.

Mix. In spoonful doses, to allay cough. *Foy.*

COMPOUND MIXTURE OF LIQUORICE.

R. Powdered liquorice, } each, half
" gum Arabic, } an ounce.
Sugar,
Paregoric, two fl. ounces.
Antimonial wine, one fl. ounce.
Spirit of nitric ether, half a fl. ounce.
Water, twelve fl. ounces.

Rub the liquorice, gum and sugar with the water gradually poured on them; then add the other ingredients and mix.
U. S. Ph.

R. Powdered liquorice,
" gum Arabic, each,
two drachms.
Boiling water, four fl. ounces.

Mix, and dissolve, then add
Spirit of nitric ether,
Antimonial wine, each, one fl. drachm.
Tincture of opium, forty to fifty drops.
A tablespoonful occasionally, in catarrh. *Ellis.*

R. Powdered liquorice,
" gum Arabic, each,
two drachms.
Boiling water, four fl. ounces.

Dissolve, and add

Antimonial wine, two fl. drachms.
Tincture of opium, twenty minims.

Mix. A tablespoonful, occasionally. *U. S. Disp.*
These three mixtures are well known under the name of *Brown mixture*.

SYRUP OF LIQUORICE.

R. Liquorice root, two ounces.
Maidenhair, one ounce.
Hyssop, half an ounce.
Water, two pints.

Infuse for twenty-four hours, boil to one-half, and add to strained decoction,

Honey, eight ounces.
Sugar, sixteen ounces.
Rose water, four fl. ounces.

Make a syrup. As a cough mixture, in spoonful doses.
Wirtem. Ph.

GOSSYPIMUM.

COTTON

Is the down or hair attached to the seeds of *Gossypium herbaceum*, and other species of the genus. It consists of fine, white filaments, having neither odor nor taste, and is insoluble in water, alcohol, ether, the oils, &c., but is soluble in strong alkaline solutions.

Sex. Syst. Monadelph. polyand. *Nat. Syst.* Malvaceæ.

Linn. Sp. Pl. 845. Griffith, Med. Bot. 162.

It is principally used for the fabrication of cloth, but is also employed in medicine, as an application to recent burns and scalds, to erysipelas, and to parts affected with rheumatism. It should be applied in the form of thin sheets. The seeds furnish much oil, and the root is said to be an active emmenagogue.

DECOCTION OF COTTON ROOT.

R. Inner part of cotton root, four ounces.
Water, one quart.

Boil down to a pint. Said to be as effectual as ergot in promoting uterine contraction. Dose, a wineglassful, every twenty or thirty minutes.
Bouchelle.

COLLODION.

This preparation, which is a solution of gun-cotton in ether, is used as a dressing to wounds; by its evaporation, it unites them closely, preserves them from the action of the air, is impervious to moisture; being transparent, it permits the progress of the wound to be examined, &c.

R. Nitric acid (1.45),
Sulphuric acid (commercial), each,
one fl. ounce.
Cleaned and bleached cotton,
two drachms

Saturate the cotton thoroughly with the acids, and macerate for twelve hours. Wash the cotton; dry rapidly, by artificial heat, in the shade, and dissolve it in

Sulphuric ether (official),
one pint and a half.

E. Parrish. W. Livermore.

R. Finely-carded cotton, half an ounce.
Powdered nitrate of potassa,
ten ounces.

Sulphuric acid, eight fl. ounces and
a half.

Ether, two pints and a half.
Alcohol, one fl. ounce.

Add the sulphuric acid to the nitre in a wedge-wood mortar, and rub them until uniformly mixed; add the cotton, and by means of the pestle and a glass rod, imbue it thoroughly with the mixture for four minutes. Transfer the cotton to a vessel containing water, and wash it carefully until the washings cease to have an acid taste, or to be precipitated by chloride of barium. Separate the fibres of cotton, dry by a gentle heat and dissolve it by agitation in the ether previously mixed with the alcohol, and strain. It should be kept in closely-stopped bottles. *U. S. Ph.*

GRANATUM.

POMEGRANATE.

The *Punica granatum*, a small tree, a native of, and cultivated in warm climates, and much esteemed for its subacid fruit, affords two articles employed medicinally, the rind of the fruit, and the bark of the root.

Sex. Syst. Icosand. monog. *Nat. Syst.* Myrtaceæ.

Linn. Sp. Pl. 676. *Griffith, Med. Bot.* 294.

The rind, as found in the shops, is in fragments of a dry, hard, brittle consistence; of a yellowish, or reddish-brown color, inodorous, and of an astringent, somewhat bitter taste. The bark of the root is in small pieces, of an ash-gray or yellowish color, externally, yellow within; brittle, with scarcely any odor, and an astringent, but not bitter taste. The rind is a good astringent, in doses of ten to thirty grains; and the bark of the root, an efficient vermifuge, especially in a fresh, or recently-dried state.

DECOCTION OF POMEGRANATE RIND.

R. Pomegranate rind, two ounces.
Distilled water, a pint and a half.
Boil to a pint, and strain. *London Ph.*
Dose, one fl. ounce. As an astringent.

GARGLE OF POMEGRANATE RIND.

R. Pomegranate rind, half an ounce.
Red roses, two drachms.
Boiling water, six fl. ounces.

Infuse, strain, and add

Clarified honey, one fl. ounce.
Alum, two drachms.

Used as a gargle, in inflammation of the fauces and throat. *Ellis.*

INJECTION OF POMEGRANATE RIND.

R. Pomegranate rind, one part.
Water, thirty-two parts.
Boil away one-half. *Beral.*

DECOCTION OF POMEGRANATE ROOT.

R. Bark of pomegranate root, two ounces.
Water, one pint and a half.
Macerate for two hours, and then boil to a pint. To be taken in three doses. As a vermifuge, in cases of tænia. *Foy.*

EXTRACT OF POMEGRANATE ROOT.

R. Bark of pomegranate root, at will.
Diluted alcohol, sufficient.
Macerate for a week, distil off the alcohol, and evaporate the residue to proper consistence. *Soubéiran.*

ELECTUARY OF EXTRACT OF POMEGRANATE ROOT.

R. Extract of pomegranate root, six drachms.
Gum tragacanth, sufficient.
Lemon juice, two fl. ounces.
Linden water, three fl. ounces.
Mix. One-half to be taken, and half an hour afterwards the remainder. *Radius.*

MIXTURE OF EXTRACT OF POMEGRANATE ROOT.

R. Extract of pomegranate root, six drachms.
Lemon juice,
Mint water,
Linden water, each, two fl. ounces.
Mix. One-fourth to be taken every quarter of an hour. *Radius.*

BOLUS OF POMEGRANATE ROOT.

R. Powdered bark of pomegranate root, one drachm.
Powdered red Peruvian bark, two drachms.
Copaiba, sufficient.
Mix, and form twelve boluses. One, morning and evening, in chronic mucous discharges. *Radius.*

R. Powdered bark of pomegranate root, one drachm.
Assafetida, half a drachm.

Croton oil, four drops.
Syrup, sufficient.
Mix, and make fifteen boluses. Five a-day, in the treatment of tænia. Foy.

GRATIOLA.

HEDGE HYSSOP.

Most of the species of this genus have active properties, but only one of them has been employed in medicine; the *G. officinalis* of Europe. A native species, the *G. aurea*, is possessed of identical properties, which, in both cases, depend on the presence of *veratria*.

Sex. Syst. Diand. monog. *Nat. Syst.* Scrophulariaceæ.

Linn. Sp. Pl. 24. Griffith, *Med. Bot.* 518.

The whole plant is used; it is almost inodorous, but has a bitter, nauseous taste. It is a drastic cathartic and emetic, with some diuretic properties. It has been stated that it forms the basis of the *Eau medicinale de Husson*. The dose of the powder is from ten to twenty grains.

COMPOUND POWDER OF HEDGE HYSSOP.

R. Powdered hedge hyssop, two scruples.
Calomel, five grains.
Assafetida, half a drachm.
Oil of peppermint, three drops.

Mix, and divide into two powders. One to be taken every three hours, in tænia. *Augustin*.

WINE OF HEDGE HYSSOP.

R. Dried herb of hedge hyssop, one ounce.

Contused root of hedge hyssop, half an ounce.
White wine, sixteen fl. ounces.

Macerate for eight days, with a gentle heat, and strain. A teaspoonful, at bedtime, drinking after it half a pint of beef tea. If necessary, repeat next evening. *Perkins*.

Said to be the original recipe for the *Eau medicinale*.

GUAIAIACUM.

GUAIAIACUM.

The *Guaiacum officinale*, a large tree, a native of the West Indies and some parts of South America, affords two medicinal products; the wood and resin, or concrete juice.

Sex. Syst. Decand. monog. *Nat. Syst.* Zygo-phylleæ.

Linn. Sp. Pl. 546. Griffith, *Med. Bot.* 203.

The wood (*lignum vitæ*) is very hard, of a greenish-brown color, of a bitterish, somewhat acrid taste, and of a faint smell, except when

heated. The resin is of a greenish, or dark olive color, of a feeble odor, and of an acrid taste. They both are stimulant and alterative, and have been much used in chronic rheumatism, gout, obstinate cutaneous affections, amenorrhœa, &c. The dose of the resin is from ten to thirty grains.

COMPOUND DECOCTION OF GUAIAIACUM WOOD.

R. Guaiacum shavings, three ounces.
Raisins, two ounces.
Sassafras root, rasped,
Liquorice root, bruised,
each, one ounce.
Water, eight pints.

Boil the guaiacum and raisins in the water, down to five pints, adding the other ingredients, towards close of operation. Strain. *Ed. Ph.*

This is the old decoction of the woods.

From one to two pints may be taken, during the day, in divided doses, as an alterative in chronic rheumatism and cutaneous diseases.

R. Rasped guaiacum wood, one ounce.
" sassafras, half an ounce.
Contused liquorice root, one drachm.
Water, five pints.

Boil for two hours, strain, and add

Cinnamon water, three ounces.

A wineglassful, several times a-day, as an alterative, &c. *Augustin*.

R. Rasped guaiacum wood, two drachms.
Water, sufficient

to obtain one pint of strained decoction. After an hour's boiling, add

Wine of colchicum seed, two drachms.

A wineglassful every two hours, in chronic pains of the joints. *Radius*.

EXTRACT OF GUAIAIACUM WOOD.

R. Powdered guaiacum wood, at will.
Exhaust with boiling water, by means of a displacement apparatus, and evaporate.

Van Mons.

OIL OF GUAIAIACUM WOOD.

R. Guaiacum wood, at will.
Distil on a sand-bath, separate the oil, and rectify it. Dose, four or five drops, in gonorrhœa; also as a friction in rheumatism.

Taddei.

COMPOUND OIL OF GUAIAIACUM.

R. Oil of guaiacum, three ounces.
Balsam of Peru, one ounce.
Ammoniated alcohol, half an ounce

Mix. In caries of the teeth, and as a dressing to malignant ulcers. *Swedjaur*

COMPOUND POWDER OF GUAIAIACUM.

R. Powdered resin of
 guaiacum, two drachms.
 Cream of tartar, half an ounce.
 Sugar, six drachms.

Mix. Three teaspoonfuls a-day, in gout.
Phæbus.

R. Powdered resin of guaiacum,
 " nitrate of potassa, each,
 one drachm.
 " ipecacuanha, three grains.
 " opium, two grains.

Mix, and divide into six powders. One, every three hours, as a stimulating diaphoretic in acute rheumatism, after reduction of inflammation.
Ellis.

PILLS OF GUAIAIACUM AND TURPENTINE.

R. Powdered resin of
 guaiacum, one drachm.
 Venice turpentine, sufficient.

Mix, and make fifteen pills. One, three times a-day, in gleet and leucorrhœa.
Ellis.

PILLS OF GUAIAIACUM, ALOES, &C.

R. Powdered guaiacum resin,
 one drachm.
 " aloes, thirty-six grains.
 " rhubarb, two drachms.
 Canada balsam, sufficient.

Form mass, and divide into forty-eight pills. One, every three hours, as a stimulating cathartic.
Ellis.

PILLS OF GUAIAIACUM, ANTIMONY, &C.

R. Powdered resin of guaiacum,
 Golden sulphuret of
 antimony, each, half a drachm.
 Calomel, ten grains.
 Extract of dandelion, sufficient.

Form mass, and divide into three-grain pills. Three to four, three times a-day, in painful affections of the joints.
Hildebrand.

PILLS OF GUAIAIACUM, SULPHUR, &C.

R. Powdered resin of
 guaiacum, half an ounce.
 Washed sulphur, one ounce.
 Crude antimony, one drachm
 and a half.

Extract of blessed thistle, sufficient.

Form mass, and make pills of two grains. Ten, three times a-day, in gout, rheumatism, and obstinate cutaneous affections.
Quarin.

MIXTURE WITH GUAIAIACUM, &C.

R. Powdered resin of
 guaiacum, two drachms.
 Powdered nitrate of
 potassa, one drachm and a half.
 Powdered gum Arabic, one drachm.
 " tartar emetic, one grain.
 " liquorice, one scruple.
 Distilled or mint water,
 eight fl. ounces.

Mix. A tablespoonful, three or four times a-day, in rheumatism, &c.
Ellis.

GUAIAIACUM MIXTURE.

R. Resin of guaiacum, three drachms.
 Sugar, half an ounce.
 Powdered gum Arabic, two drachms.
 Cinnamon water, one pint.

Rub the resin with the sugar, and gum Arabic, and gradually add the cinnamon water while rubbing.
Lond. Ph.

One to three tablespoonfuls, two or three times a-day.

MIXTURE OF GUAIAIACUM AND BITTER-SWEET.

R. Powdered resin of
 guaiacum, two drachms.
 Powdered gum Arabic, three drachms.
 Extract of bitter sweet, three drachms.
 Syrup of mallow, one fl. ounce.
 Distilled water, seven fl. ounces.

Mix. A tablespoonful every three hours, in chronic rheumatism.
Berends.

TINCTURE OF GUAIAIACUM.

R. Powdered resin of
 guaiacum, half a pound.
 Alcohol, two pints.

Macerate for fourteen days, and filter.
U. S. Ph.

Dose, one to three fl. drachms, three or four times a-day, in gout and chronic rheumatism; to be given in milk or mucilage.

DEWEES'S TINCTURE.

R. Powdered resin of
 guaiacum, four ounces.
 Carbonate of soda, or
 potassa, one drachm and a half.
 Powdered pimento, one ounce.
 Diluted alcohol, one pint.

Digest for a few days. Add volatile spirit of ammonia, if required, in proportion of one or two drachms to each four ounces of tincture. Highly praised in idiopathic amenorrhœa, in doses of a teaspoonful, three times a-day, in milk or wine.
Dewees.

AMMONIATED TINCTURE OF GUAIAIACUM.

R. Powdered resin of
guaiacum, four ounces.
Aromatic spirit of
ammonia, one pint and a half.
Macerate for fourteen days, and filter.

U. S. Ph.

Dose, one to two fl. drachms, two or three times a-day, in chronic rheumatism.

AMMONIATED TINCTURE OF GUAIAIACUM AND COPAIBA.

R. Ammoniated tincture of
guaiacum, one fl. ounce.
Copaiba, half fl. ounce.

Mix. A teaspoonful, two or three times a-day, in amenorrhœa with general debility. *Ellis.*

AMMONIATED TINCTURE OF GUAIAIACUM AND PAREGORIC.

R. Ammoniated tincture of
guaiacum, one fl. ounce.

Camphorated tincture of
opium, one fl. ounce.
Mix. Two teaspoonfuls every two hours, in
misplaced and retrocedent gout. *Ellis.*

ODONTALGIC MIXTURE OF GUAIAIACUM.

R. Tincture of guaiacum, six fl. drachms.
Oil of valerian, one fl. ounce.
Syrup of scurvy grass,
Compound tincture of
benzoin, each, two fl. drachms.
Tincture of opium, one fl. drachm.

Mix. A teaspoonful mixed with hot water, to be held in the mouth, in toothache. *Carus.*

MIXTURE OF TINCTURE OF GUAIAIACUM AND HENBANE.

R. Tincture of guaiacum, two fl. drachms.
" henbane, one fl. drachm.

Mix. Twenty to thirty drops, morning and evening, in spasm of the stomach and neuralgia. *Radius.*

H.

HÆMATOXYLON.

LOGWOOD.

This is the heart wood of *Hæmatoxylon campechianum*, a middle-sized tree, a native of Campeachy, and naturalized in several of the West India islands. It is principally used for dyeing purposes, but is also employed in medicine.

Sex. Syst. Decand. monog. *Nat. Syst.* Fabaceæ.

Linn. Sp. Pl. 549. Griffith, Med. Bot. 252.

It comes in logs of a dark-yellowish color externally, and deep red internally; for medical use, it is chipped or rasped. It has a feeble but peculiar odor, and a sweetish, somewhat astringent taste. It is used as a mild astringent in chronic diseases, and relaxed conditions of the bowels.

INFUSION OF LOGWOOD.

R. Rasped logwood, half an ounce.
Boiling water, one pint.
Infuse for two hours, and strain. A table-spoonful every two or three hours, in cholera and diarrhœa of children. *Ellis.*

DECOCTION OF LOGWOOD.

R. Rasped logwood, one ounce.

Water, two pints.

Boil down to a pint, and strain. *U. S. Ph.*

A good astringent in diarrhœa, especially in children. Dose for an adult, two fl. ounces; for a child about two years of age, two or three fl. drachms, several times a-day.

EXTRACT OF LOGWOOD.

R. Rasped logwood, one pound.
Water, one gallon.

Boil to four pints, strain while hot, then evaporate to proper consistence. *U. S. Ph.*

Dose, from ten to thirty grains.

ELECTUARY OF EXTRACT OF LOGWOOD.

R. Extract of logwood,
" liquorice, each, two drachms.
" Peruvian bark, three drachms
" cascarilla, one drachm.
Mucilage of quince-seed, sufficient.

Mix. A teaspoonful, three times a-day, in chronic diarrhœa, &c. *Radius.*

MIXTURE OF EXTRACT OF LOGWOOD.

R. Extract of logwood, three drachms.
Boiling water, seven fl. ounces.

Dissolve, strain, and add

Tincture of cinnamon, six fl. drachms.

" catechu, two fl. drachms.

Mix. Dose, one fl. ounce, every six hours.

Beasley.

R. Extract of logwood, three drachms.

Tincture of catechu, two fl. drachms.

Water, seven fl. ounces.

Mix. Two spoonfuls, every three or four hours.

In diarrhœa and dysentery.

Ellis.

R. Extract of logwood, three drachms.

Spirit of cinnamon, one fl. ounce and
a half.

Water, seven fl. ounces.

Tincture of kino, two fl. drachms.

Mix. In the same doses, and in the same diseases, as the last.

Swediaur.

HAMAMELIS.

WITCH HAZEL.

The *Hamamelis Virginica* is a large native shrub, found along streams in most parts of the country, presenting several varieties.

Sex. Syst. Tetrand. digyn. *Nat. Syst.* Hamamelidaceæ.

Pursh, Fl. Am. i. 116. Griffith, Med. Bot. 350.

The parts used are the bark and leaves. These are bitter and astringent, leaving a sensation of sweetness. They are considered sedative, astringent, and tonic, and useful in bowel affections and hemorrhages, and externally as an application to tumors, painful hemorrhoids, &c. They are used in infusion, decoction, or poultice.

HEDEOMA.

PENNYROYAL.

Hedeoma pulegioides is a small, annual, aromatic plant, abundant in most parts of the United States, growing in dry, sterile situations.

Sex. Syst. Diand. monog. *Nat. Syst.* Lamiaceæ.

Persoon, Synop. ii. 131. Griffith, Med. Bot. 308.

The whole plant is used. It has a warm, pungent taste, and a powerful aromatic odor. It is a stimulating aromatic, used to obviate nausea, and relieve flatulence, and also, in domestic practice, as an emmenagogue.

INFUSION OF PENNYROYAL.

R. Pennyroyal, two drachms.
Boiling water, sufficient

to afford six ounces of strained infusion. To be taken freely, in a warm state, at bedtime, the feet having been previously bathed in hot water, in amenorrhœa.

OIL OF PENNYROYAL.

R. Pennyroyal, at will.
Water, sufficient.

Distil, and collect the oil. Dose, two to ten drops, in flatulent colic and nausea.

HELENIUM.

SNEEZEWORD.

Sneezewort, or *Helenium autumnale*, is an indigenous plant, found in wet situations, in most parts of the United States, flowering in the autumn.

Sex. Syst. Syngen. super. *Nat. Syst.* Asteraceæ.

Torrey & Gray, Fl. ii. 384. Griffith, Med. Bot. 398.

This plant is almost inodorous, but has a bitter, somewhat pungent, acrid taste. It is tonic, diaphoretic, and powerfully errhine; this latter property is most developed in the flowers, and especially the central florets; these, when powdered, may be used in those cases to which errhines are applicable.

HELIANTHEMUM.

FROSTWEED.

The *Helianthemum Canadense* is a native perennial plant, growing in many parts of the United States, in dry, sandy soils.

Sex. Syst. Polyand. monog. *Nat. Syst.* Cistaceæ.

Mich. Fl. i. 308. Darlington, Fl. Cest. 313.

It has little or no odor, but an astringent, somewhat aromatic, bitterish taste. It is tonic and astringent, and has proved beneficial in scrofula.

HELLEBORUS.

BLACK HELLEBORE.

Several species of *Helleborus* are possessed of almost identical medical properties, but the only one used in this country, is *H. niger*, so called on account of the color of its roots. It is a native of mountain woods, in many parts of Europe, and is cultivated as an ornamental flowering plant.

Sex. Syst. Polyand. polyg. *Nat. Syst.* Ranunculaceæ.

Linn. Sp. Pl. 783. Griffith, Med. Bot. 85.

The parts used are the small fibres, or roots, which are about as thick as straw, and black; they proceed from a rhizome or caudex, and have a somewhat nauseous odor, and a bitter, unpleasant, acrid taste. Black hellebore is a drastic hydragogue purgative; having, also,

considerable emmenagogue powers. The dose in powder is from ten to twenty grains, as a purge; two or three grains, as an alterative.

COMPOUND BLACK HELLEBORE PILLS.

R. Powdered black hellebore, fifteen grains.
Calomel, five grains.
Powdered ipecacuanha, three grains.
Syrup of ginger, sufficient.
Mix, and make four pills. Two to be taken every four hours, in dropsy, till full purgation is caused. *Ainslie.*

EXTRACT OF BLACK HELLEBORE.

R. Powdered black hellebore, at will.
Exhaust by means of the displacement process, with cold diluted alcohol. Distil off the alcohol, and evaporate to proper consistence. *U. S. Ph.*

Dose, ten to fifteen grains, as a drastic purge.

VINO-ALCOHOLIC EXTRACT OF BLACK HELLEBORE.

R. Powdered black hellebore, two pounds.
Carbonate of potassa, half a pound.
Diluted alcohol, eight pints.
Macerate for twelve hours, express, and pour on the residuum,
White wine, eight pints.
Digest for twenty-four hours, express, mix the two tinctures, and evaporate. *Cottureau.*
Dose, ten to fifteen grains.

COMPOUND PILLS OF EXTRACT OF BLACK HELLEBORE.

R. VINO-alcoholic extract of black hellebore,
Extract of myrrh, each, two ounces.
Powder of blessed thistle, one ounce.
Beat together, and form pills of one grain. Ten to twenty a-day, in amenorrhœa, dropsy, &c. *Radius.*

These pills are much used in Europe, under the name of *Bacher's pills*.

R. Extract of black hellebore,
Assafetida,
Ammoniac,
Soap, each, two drachms.
Rhubarb, sufficient.
Beat into mass, and form pills of two grains.
Dose, ten to twelve, morning and evening, as a purgative and emmenagogue. *Augustin.*

TINCTURE OF BLACK HELLEBORE.

R. Bruised black hellebore, four ounces.
Diluted alcohol, two pints.

Macerate for fourteen days, express, and filter. *U. S. Ph.*

Used as an emmenagogue, in doses of thirty drops to a fl. drachm, night and morning, watching its action.

MIXTURE WITH TINCTURE OF BLACK HELLEBORE.

R. Tincture of black hellebore, half a fl. ounce.
Tincture of myrrh, one fl. ounce.
" Spanish flies, two fl. drachms.

Mix.

Thirty drops, three times a-day, in sugar and water, as an emmenagogue.

COMPOUND WINE OF BLACK HELLEBORE.

R. Bruised black hellebore, one ounce.
" wormwood, a handful.
White wine, four pints.

Macerate for three days, express, and filter. Two to three spoonfuls, in the morning, fasting. In dropsy. *Brunner.*

OINTMENT OF BLACK HELLEBORE.

R. Powdered black hellebore, one to two drachms.
Lard, one ounce.
Mix. As an application to obstinate herpetic eruptions. *Soubeiran.*

HEMIDESMUS.

INDIAN SARSAPARILLA.

The *H. indicus* is a native of many parts of India. It is a climbing plant, and has been long used in the East as an efficient medicinal agent; but was almost unknown in Europe, or this country, until about 1819.

Sex. Syst. Pentand. digyn. *Nat. Syst.* Asclepiadaceæ.

Brown, Hort. Kew. ii. 75. Griffith, Med. Bot. 453.

The root, which is the part used, is long, tortuous, rugose, with longitudinal furrows; it is brownish, externally, and has a peculiar and somewhat aromatic odor, and a bitterish taste. It has the properties of sarsaparilla; and like it, is given in infusion, decoction, &c. These are made in the same manner as those of sarsaparilla, and given in the same doses, and in similar diseases.

SYRUP OF INDIAN SARSAPARILLA.

R. Indian sarsaparilla, half a pound.
Boiling water, one pint

Digest in a covered vessel, with a gentle heat, for three or four hours, strain, add twice the weight of sugar, and make syrup. *Beasley.*

MIXTURE OF INDIAN SARSAPARILLA.

R. Indian sarsaparilla, ten drachms.
Extract of liquorice, ten grains.
Distilled water, ten fl. ounces.

Digest for twelve hours, strain, heat to 180°, and again strain. One-third three times a-day.

Beasley.

R. Syrup of Indian sarsaparilla, five ounces.

Solution of potassa, half to one fl. drachm.

Orange-flower water, one fl. ounce.

Mix.

One fl. ounce, thrice a-day, in barley-water. For gonorrhœa. *Bellinarye.*

INFUSION OF INDIAN SARSAPARILLA.

R. Indian sarsaparilla, two ounces.
Lime water, one pint.

Infuse in a close vessel for twelve hours. Dose, a wineglassful. *Ashburner.*

DECOCTION OF INDIAN SARSAPARILLA.

R. Indian sarsaparilla, two ounces.
Water, one pint and a half.

Boil down to one pint. To be taken during the day, in wineglassful doses. *Pereira.*

HEPATICA.

LIVERWORT.

The *H. triloba* is a small native plant, found in most parts of the United States, and also in the northern regions of Europe and Asia, having three-lobed leaves, and presenting two marked varieties: one with the lobes of the leaves rounded, the other with them acute.

Sex. Syst. Polyand. polyg. *Nat. Syst.* Ranunculaceæ.

Torrey & Gray, Fl. i. 14. Griffith, Med. Bot. 81.

The whole herb is used. It is inodorous, and has a mucilaginous, slightly astringent and bitterish taste. It is a very mild, demulcent tonic and astringent, and was much employed at one time as a deobstruent in herpetic affections, and also as a remedy in diseases of the lungs. Its powers, however, are very slight. It is given in infusion; to be taken freely.

HERACLEUM.

MASTERWORT.

This is the root of *Heracleum lanatum*, a very large, perennial plant, found in many parts of the United States, in waste places.

Sex. Syst. Pentand. digyn. *Nat. Syst.* Apiaceæ.

Torrey & Gray, Fl. i. 632. Griffith, Med. Bot. 335.

The part used is the root; this resembles the parsnip in appearance, but has a rank, unpleasant odor, and a pungent, acrid taste. It is said to be diuretic, expectorant, and antispasmodic, and has proved useful in epilepsy, attended with a disordered condition of the digestive organs, in doses of two or three drachms daily. It is also used in strong decoction, in dyspepsia with flatulence and cardialgia.

HEUCHERA.

ALUM ROOT.

Most of the species of *Heuchera* are possessed of identical properties, but the only one recognized by the U. S. Pharmacopœia is *H. Americana*. This is an indigenous plant, growing in shady, rocky situations, in most parts of the country.

Sex. Syst. Pentand. digyn. *Nat. Syst.* Saxifragaceæ.

Torrey & Gray, Fl. i. 578. Griffith, Med. Bot. 313.

The root is rugose, irregular, yellowish, almost inodorous, and of a strong, styptic taste. It is a powerful astringent, and may be employed in such cases as require medicines of this class.

HIPPOCASTANUM.

HORSE CHESTNUT.

The horse chestnut, or *Æsculus hippocastanum*, is a beautiful and lofty tree, a native of the central parts of Asia, but extensively cultivated in Europe, and the United States.

Sex. Syst. Heptand. monog. *Nat. Syst.* Sapindaceæ.

Linn. Sp. Pl. 488. Griffith, Med. Bot. 213.

The part used is the bark; this is light, brittle, of a brownish-red externally, of a yellow-brown within. It has a somewhat aromatic odor, and an astringent, bitter taste. It is a somewhat active astringent, and has been much praised in intermittent fevers. The dose of the powder is two to four scruples, every three hours, till an ounce and a half is taken.

COMPOUND POWDER OF HORSE CHESTNUT.

R. Powdered horse chestnut bark,
" willow bark, each, half an ounce.

" gentian,

" calamus,

" cloves, each, two drachms.

Mix.

Hufeland.

DECOCTION OF HORSE CHESTNUT
BARK.

R. Horse chestnut bark, one ounce and a half.
Water, thirty fl. ounces.
Boil down to ten ounces, adding towards close of operation,
Liquorice root, one drachm.
Strain. A cupful every two hours. *Niemann.*

COMPOUND DECOCTION OF HORSE
CHESTNUT.

R. Horse chestnut bark,
Willow bark, each, half an ounce.
Calamus,
Root of water avens,
each, two drachms.
Water, sixteen fl. ounces.
Boil down to one-half. *Spielmann.*
R. Horse chestnut bark, one ounce and a half.
Water, eighteen fl. ounces.
Boil down to one-half, strain, and add
Sulphuric ether, one to two drachms.
Syrup of orange peel, one ounce.
Mix. To be used during the apyrexia. *Phæbus.*

EXTRACT OF HORSE CHESTNUT.

R. Horse chestnut bark, at will.
Water, sufficient.
Exhaust the bark in a displacement apparatus, and evaporate to proper consistence. Dose, five to fifteen grains. *Van Mons.*

HORDEUM.

BARLEY.

There are several species of *Hordeum*, but those usually cultivated in this country are *H. vulgare* and *H. distichon*, the latter of which is recognized as officinal. The native country of these is unknown, but they have been cultivated from the earliest ages.

Sex. Syst. Triand. digyn. *Nat. Syst.* Graminacæ.

Linn. Sp. Pl. 125. Griffith, Med. Bot. 664.

The seeds are used in various forms; in that of meal, malt, pearl barley, &c. They are one of the mildest and least irritating of the cerealia, and are much used in decoction, &c., as a nutritive and demulcent drink.

PREPARED BARLEY MEAL.

R. Barley meal, at will.
Tie it in a linen or cotton cloth, and boil it for twelve hours, then let it cool, remove the outer crust, and pulverize the centre. A useful diet, boiled with milk, in bowel diseases.
Hanover Ph.

R. Barley meal, twelve ounces.
Sugar, four ounces.
Powdered cinnamon, half a drachm.

Mix, and place in a proper vessel, covering with wheat dough, put in an oven and bake, remove, cool, and pulverize. Half an ounce to two ounces, cooked with water or milk, forms an excellent diet, in debilitated conditions of the system. *Taddei.*

BARLEY SUGAR.

R. Decoction of barley, a pint and a half.
Sugar, two pounds.
Boil to proper consistence, and form lozenges or rolls. As a demulcent in catarrh. *Giordano.*

DECOCTION OF BARLEY.

R. Pearl barley, two ounces.
Water, four pints and a half.
Wash the barley in cold water, drain, pour on it half a pint of the water, boil for a short time, drain off this water, add the remainder in a boiling state, and boil down to one-half, and strain. *U. S. Ph.*
As a nutritive and demulcent drink in febrile and inflammatory diseases.

COMPOUND DECOCTION OF BARLEY.

R. Decoction of barley, two pints.
Sliced figs, two ounces and a half.
Bruised liquorice root, five drachms.
Stoned raisins, two ounces and a half.
Water, one pint.
Mix, and boil down to two pints, and strain. *Lond. Ph.*

A demulcent, nutritive, and somewhat laxative drink.

BARLEY WATER WITH NITRATE OF
POTASSA.

R. Decoction of barley, one pint.
Nitrate of potassa, two drachms.
Lemon juice, one fl. ounce.
Mix. To be used warm as a diaphoretic drink. *Ellis.*

HUMULUS.

HOPS.

By this is meant the strobiles of the *Humulus lupulus*, a climbing vine, a native of Europe, and probably of this country. It is extensively cultivated for its aments or strobiles, which are largely employed in the preparation of malt liquors, and also in medicine.

Sex. Syst. Diœc. pentand. *Nat. Syst.* Cannabinacæ.

Linn. Sp. Pl. 1457. Griffith, Med. Bot. 574

Hops consist of thin, somewhat translucent, leaf-like scales, of a greenish-yellow color, having, near their base, two small, round, dark seeds. Their odor is strong and peculiar, somewhat narcotic, and fragrant; their taste is bitter, aromatic, and somewhat astringent. These properties depend on a peculiar secretion, called Lupuline. Hops are tonic, and slightly narcotic, and are used in various conditions of the system. The dose, in substance, is from half a drachm to a drachm, but it is seldom administered in this form.

INFUSION OF HOPS.

R. Hops, half an ounce.
Boiling water, one pint.

Macerate for two hours in a covered vessel, and strain. *U. S. Ph.*

Dose, one to two fl. ounces, in dyspepsia, nervous tremors, &c.

EXTRACT OF HOPS.

R. Hops, at will.
Diluted alcohol, sufficient.

Treat in a displacement apparatus, distil off the alcohol, and evaporate the residue. Dose, from ten to thirty grains. *Cotteneau.*

HOP MIXTURE.

R. Extract of hops, two drachms.
Water of hops, seven fl. ounces.
Tincture of hops, half fl. ounce.
Syrup of orange peel, one fl. ounce.

Mix. A tablespoonful every hour, as a tonic and stomachic. *Niemann.*

TINCTURE OF HOPS.

R. Hops, five ounces.
Diluted alcohol, two pints.

Macerate for fourteen days, express, and filter. *U. S. Ph.*

Dose, from one to three fl. drachms, as a tonic, and narcotic, especially in the wakefulness and tremors of drunkards.

ALKALINE TINCTURE OF HOPS.

R. Hops, one ounce.
Centaury, each, one ounce.
Peel of bitter orange, two drachms.
Carbonate of potassa, one scruple.
Diluted alcohol, eighteen fl. ounces.

Macerate for eight days, express, and filter. Dose, half an ounce to an ounce.

CATAPLASM OF HOPS.

R. Hops, one pound.
Flaxseed meal, two ounces.
Beer, sufficient.

Mix. As an application to bruises and indolent tumors. *Radius.*

OINTMENT OF HOPS.

R. Hops, two ounces.
Lard, ten ounces.

Digest for some hours, by a gentle heat, express, and strain; recommended to relieve the pain of cancerous sores. *Swedjaur.*

HYDRARGYRUM.

MERCURY.

Mercury is a brilliant, silver-white, fluid metal, having neither taste nor smell. It becomes solid at -39° F., and boils at 665° . Mercury, in masses, does not appear to act on the system, but when in a state of great division, it produces marked and peculiar effects. It acts as an alterative, deobstruent, sialagogue, &c. It has been employed in almost all diseases, in some of its preparations, each of which has some peculiarity of action.

PURIFIED MERCURY.

R. Mercury, three pounds.
Pure muriatic acid, half fl. ounce.
Distilled water, two fl. ounces.

Place the mercury in a glass retort, and distil off two-thirds. Boil this for a few minutes with the acid and water, remove the acid by repeated washing with water and decantation, pour the metal into a capsule and dry it by the application of heat. *Dub. Ph.*

MERCURY WITH ANTIMONY.

R. Mercury, four parts.
Crude antimony, three parts.
Sulphur, two parts.

Triturate together till the mercury disappears. As a diaphoretic and alterative. Dose, one to four grains. *Giordano.*

R. Mercury, one ounce.
Washed flowers of sulphur, one ounce.
Powdered sulphuret of antimony, three ounces.

Rub them in a warm stone mortar, with a little water or sulphuret of ammonium, until the globules of mercury disappear. *Cod. Hamb. 1845.*

This preparation is much used in France and Germany, under the name of *antimonial ethiops*.

COMPOUND MERCURIAL POWDER.

R. Powdered resin of guaiacum, one drachm.
Mercury with antimony, one scruple.
Magnesia, each, one scruple.

Mix. To be taken in two days, in divided doses, in chronic exanthematous affections. *Hufeland.*

R. Mercury with antimony,
twenty-four grains.
 Subcarbonate of soda, eighteen grains.
 Powdered sassafras,
 Sugar, each, one drachm.
 Mix, and divide into six powders. One to be
 taken daily, in chronic diseases of the skin.
Sundelin.

PILLS OF MERCURY AND ANTIMONY.

R. Mercury, one drachm.
Crude antimony, four scruples.
Flowers of sulphur, two scruples.
Extract of opium, one scruple.

Triturate till mercury disappears, with
Syrup of mallow, sufficient.

Divide into one hundred and fifty pills. Dose,
five or six a-day. Highly spoken of by Huxham,
in venereal pains, scrofula, &c. *Baldinger.*

℞. Mercury with antimony, one drachm.
 Extract of hemlock, two drachms.
 Soap, half an ounce.
 Galbanum,
 Extract of ox gall, each, half a drachm.
 Beat into mass, and form pills of two grains.
 Dose, eight to ten, three times a-day, in scirrhus
 of the pylorus. *Radiis.*

MERCURY WITH CHALK.

R. Mercury, three ounces.
Prepared chalk, five ounces.
Rub together till all the globules disappear.
U. S. Ph.

A mild preparation, well suited as an alterative in complaints of children. Dose, five grains to half a drachm, twice a-day, for adults; two or three grains for a child.

MERCURY AND CHALK.

R. Mercury,	three ounces.
Resin,	six drachms.
Prepared chalk,	five ounces.
Alcohol,	sufficient.

Make a paste with the resin, and a small quantity of the alcohol; then add the mercury, which may be extinguished in a short time; add the chalk and alcohol gradually, so as to keep up the pasty consistence; then add sufficient alcohol to dissolve out the resin, and wash the powder on a filter, and dry.

MERCURY AND CHALK WITH IPECAC-
UANHA.

R. Mercury with chalk, one scruple.
Powdered ipecacuanha, ten grains.
Mix, and divide into six powders. One, night
and morning, in syrup. In dyspepsia with
biliary derangement. *Paris.*

MERCURY AND CHALK LINIMENT.

R. Mercury,
Prepared chalk, each, half an ounce.
Triturate till globules disappear, and add
Honey of roses, two ounces.
Mix well. As a dressing to venereal ulcers.
Bories.

MERCURY AND GUM.

R. Mercury, one part.
Gum Arabic, two parts.
Make a mucilage with a quarter of the gum, rub the mercury with it till extinguished, add the remainder of the gum, with as much water as is required, mix well, dry by a gentle heat, and pulverize. *Paris Cod.*

PILLS OF MERCURY AND GUM WITH
HEMLOCK.

R. Mercury,	one part.
Gum Arabic,	two parts.
Syrup of violets,	four parts.
Triturate till the mercury is extinguished, and add	
Extract of hemlock,	one part.
Powdered liquorice,	sufficient.
Mix well, and form pills of two grains. Two to four, twice a-day, as an alternative. <i>Plenck.</i>	

MERCURIAL MUCILAGE OF GUM ARABIC.

R. Mercury,	one part.
Gum Arabic,	two parts.
Water,	sufficient.

Triturate till the mercury is extinguished. This is a good preparation of mercury; it is given in the dose of two teaspoonfuls, in syrup or mucilage, morning and night, as an alterative or anthelmintic.

Beral.

MERCURIAL GARGLE.

R. Mercury,	half a drachm.
Gum Arabic,	three drachms.
Syrup of poppies,	half an ounce.
Calomel,	six grains.
Triturate till mercury is extinguished, and add	
Decoction of clematis, (Vir-	
gin's bower), twenty-six fl. ounces.	
Honey of roses,	one ounce.
Essence of myrrh,	one drachm.
Mix. As a gargle in syphilitic angina, and	
ozæna. <i>Plenk.</i>	

MERCURIAL LOTION.

R. Mercury,	one drachm.
Gum Arabic,	four drachms.
Syrup of poppies,	sufficient

Rub together till globules disappear, and add
 gradually, constantly rubbing,

Boiling milk,	eight fl. ounces.
---------------	-------------------

As a lotion in gonorrhœal ophthalmia, ulcers on the penis, and also as a gargle in venereal sore throat.

Plenck.

MERCURY WITH MAGNESIA.

- R. Purified mercury,
Manna, each, two parts.
Carbonate of magnesia, one part.

Rub the mercury with the manna, adding a little water, till globules disappear; add one-eighth part of the magnesia, still rubbing, and when mixed, sixteen parts of hot water, and agitate; let sediment subside; then decant, and repeat washing, till all the manna is removed; mix the residue with remainder of magnesia, and dry.

Dub. Ph., 1826.

A mild mercurial, well suited as an alterative in children, especially when there is constipation. Dose, three to four grains.

BLUE PILLS.

- R. Mercury, one ounce.
Confection of roses, one ounce and a half.
Powdered liquorice root, half an ounce.

Rub the mercury with the confection, till all the globules disappear, add the liquorice root, and beat into mass. Divide into four hundred and eighty pills.

U. S. Ph.

One of the mildest and best of the mercurial preparations. From five to fifteen grains as a purgative; one pill every night, or every other night, as an alterative; one pill, two or three times a-day, as a laxative. Each pill contains one grain of mercury.

BLUE PILL WITH JALAP.

- R. Blue pill,
Powdered jalap,
" aloes, each, fifteen grains.
Mix, and form twelve pills; three at night, as a purgative.

Ellis.

BLUE PILL WITH RHUBARB.

- R. Blue pill, nine grains.
Powdered rhubarb,
Bicarbonate of soda, each, twelve grains.
Aromatic syrup of rhubarb, sufficient.

Beat into mass, and form twelve pills. One, twice or thrice a-day, as an alterative, in hepatic derangement.

Hartshorne.

BLUE PILL WITH COLOCYNTH.

- R. Blue pill,
Compound extract of colocynth,
each, five grains.
Oil of caraway, two drops.

Mix, and make two pills. A very active purgative.

Ellis.

BLUE PILL AND QUINIA.

- R. Blue pill,
Sulphate of quinia,
Powdered aloes, each, twelve grains.
Aromatic syrup of rhubarb, sufficient.

Beat into mass, and form twelve pills. One, twice to four times a-day, as a tonic alterative in deranged conditions of the liver, consequent to fevers.

Ellis.

COMPOUND MERCURIAL PILLS.

- R. Blue pill, five grains.
Powdered ipecacuanha, two grains.
Camphor, one grain and a half.
Syrup of ginger, sufficient.

Mix, and make two pills. One, morning and evening, in hepatitis, till mouth is affected.

Ainslie.

- R. Blue pill, four grains.
Compound powder of squill, six grains.
Powdered ipecacuanha, two grains.
Syrup of ginger, sufficient.

Mix, and make three pills. To be taken in a day; in hepatic obstruction, threatening dropsy.

Ainslie.

- R. Blue pill,
Antimonial powder, each, two and a half grains.
Opium, half a grain.
Syrup of ginger, sufficient.

Make a pill. To be taken at bedtime; in venereal blotches, attended with diarrhœa.

Ainslie.

ABERNETHY'S PILLS.

- R. Blue pill, ten grains.
Powdered jalap, twenty grains.
Syrup of buckthorn, sufficient.

Mix well, and divide into six pills. Two at night, with a wineglassful of infusion of senna, in the morning.

Cooley.

MERCURIAL MIXTURE

- R. Mercury, six drachms.
Syrup of poppies, one ounce.
Triturate till globules disappear, and add

Orange-flower water, two fl. ounces.
Rose water, one fl. ounce.

Mix well. Dose, a spoonful, morning and night, in gastrodynia and spasmodic vomiting.

Cadet de Gassicourt.

MERCURIAL OINTMENT.

- R. Mercury, two pounds.
Lard, twenty-three ounces.
Suet, one ounce.

Rub the mercury with the suet and a little of the lard, till the globules disappear; add the remainder of the lard, mixing well. *U. S. Ph.*

This is the strong mercurial ointment. It can be weakened by the addition of lard, as may be wished. Used as an inunction to mercurialize the system, by rubbing about a drachm on the inside of the thighs, twice a-day.

R. Mercury, two pounds.
Lard, rendered rancid by exposure in a damp, divided state, four ounces.
Lard, nineteen ounces.
Suet, one ounce.

Triturate the mercury and rancid lard, until the globules disappear; then add the lard and suet, and triturate until the texture of the mass is uniform. *W. Procter.*

CAMPHORATED MERCURIAL OINTMENT.

R. Mercurial ointment, one ounce.
Camphor, one drachm.

Mix. Used like the former; also to disperse indolent swellings.

MERCURIAL CATAPLASM.

R. Mercurial ointment, half an ounce.
Camphor, two drachms.
Boiling milk, three ounces.
Crumb of bread, sufficient.

Mix, and form cataplasm. *Radius.*

MERCURIAL CERATE.

R. Mercurial ointment,
Simple cerate, equal parts.

Mix well. As a dressing to venereal ulcers. *Guibourt.*

COMPOUND MERCURIAL CERATE.

R. Mercurial ointment,
Soap cerate, each, four ounces.
Camphor, one ounce.

Rub well together. *Lond. Ph.*

As an application to disperse indolent swellings.

COMPOUND MERCURIAL LINIMENT.

R. Mercurial ointment,
Lard, each, four ounces.
Camphor, one ounce.
Rectified spirit, one fl. drachm.
Solution of ammonia, four fl. ounces.

Rub the camphor with the spirit, then with the lard and mercurial ointment; lastly, add gradually the solution of ammonia, and mix well. *Lond. Ph.*

As a stimulating liniment, in chronic swellings, glandular tumors, &c.

OPIATED MERCURIAL LINIMENT.

R. Mercurial ointment, four parts.
Oil of sweet almonds, forty-eight parts.
Tincture of opium, three parts.

Mix well. Advised in inflamed ulcers of the glans penis, to be applied two or three times a-day. *St. Marie.*

COMPOUND MERCURIAL OINTMENT.

R. Mercurial ointment, two ounces.
Lard, twelve ounces.
Powdered stavesacre, three ounces.

Melt the lard, and add the other ingredients, mixing well. As an application to destroy lice. *Giordano.*

R. Mild mercurial ointment, eight parts.
Soft soap, two parts.
Camphor, one part.

Rub well together. Recommended in perioritis and engorgement of the testicles. *Swediaur.*

MERCURIAL AND BELLADONNA PLASTER.

R. Mercurial ointment, one ounce.
Ammoniac, six drachms.
Extract of belladonna, four drachms.
Hydrocyanic acid, thirty drops.

Make mass with ammoniac and extract, with a little water; mix mercurial ointment and acid, and rub the whole together. Useful as an application to scirrhus and scrofulous tumors.

Med. Chirurg. Pharm.

OINTMENT OF MERCURY AND BELLADONNA.

R. Strong mercurial ointment, thirty parts.
Extract of belladonna, four parts.
" opium, one part.
Balsam of Peru, sufficient.

Mix well. As an application to painful hemorrhoidal tumors. *Mignot.*

MERCURIAL PLASTER.

R. Mercury, six ounces.
Olive oil,
Resin, each, two ounces.
Lead plaster, one pound.

Melt the oil and resin together, and, when cool, rub the mercury with them till the globules di-

appear; gradually add the lead plaster, previously melted, and mix well. *U. S. Ph.*

As an application to buboes, venereal nodes, &c.

R. Mercurial ointment, one ounce.
Oil of turpentine,
Camphor, each, two drachms.
Simple cerate, one ounce.

Mix well. Used as a rubefacient application over the region of the liver, when blisters cannot be used. *Ellis.*

HYDRARGYRI ACETAS.

ACETATE OF MERCURY.

ACETATE OF PROTOXIDE OF MERCURY.

R. Protonitrate of mercury, one part.
Distilled water, six parts.

Dissolve the salt in water, acidified with a little nitric acid, add a solution of acetate of soda, or potassa, filter, wash, and dry the precipitate. *Beral.*

PILLS OF ACETATE OF MERCURY.

R. Red oxide of mercury, one pound.
Distilled vinegar, eight pints.

Dissolve, and triturate a pint of this solution with Flake manna, two pounds, rubbing the mixture for a long time; dry before the fire, often stirring; when of a proper consistence, form pills of a grain and a half. *Guibourt.*

These pills were highly esteemed under the name of *Keyser's anti-venereal pills*. They contain, at first, the acetate of the deutoxide, but this gradually changes to the acetate of the protoxide, then to an oxide; therefore, the following have been substituted.

R. Acetate of mercury,
Flake manna,
Powdered gum Arabic, each, one scruple.
Rose water, sufficient.

Beat into mass, and form twenty pills. As a sialagogue, three to be taken at night, or one three times a-day. *Ellis.*

PILLS OF ACETATE OF MERCURY AND OPIUM.

R. Acetate of mercury,
Opium,
Camphor, each, thirty grains.
Syrup of poppies, sufficient.
Mix, and make thirty pills. *Carmichael.*

SOLUTION OF ACETATE OF MERCURY.

R. Acetate of mercury, ten or twelve grains.
Rose water, five fl. ounces.
Dissolve. As a lotion in obstinate cutaneous affections. *Niemann.*

LINIMENT OF ACETATE OF MERCURY.

R. Acetate of mercury, one part.
Olive oil, two parts.
Lard, six parts.

Rub the salt with a little of the lard, and add gradually the remainder, and then the oil, constantly rubbing. Used as an application in herpes. *Van Mons.*

HYDRARGYRUM AMMONIATUM.

WHITE PRECIPITATE.

R. Corrosive sublimate, six ounces.
Distilled water, one gallon.
Solution of ammonia, eight fl. ounces.

Dissolve the corrosive sublimate in the water with the aid of heat, and when cold add the solution of ammonia, frequently stirring. Wash the precipitate thoroughly, and dry it. *U. S. Ph.*

This is seldom, if ever, used, except as an external remedy.

OINTMENT OF WHITE PRECIPITATE.

R. White precipitate, one drachm.
Simple ointment, one ounce and a half.
Softened the ointment over a gentle fire, and mix the white precipitate. *U. S. Ph.*

As an application to cutaneous eruptions.

HYDRARGYRI BORAS.

BORATE OF MERCURY.

R. Calomel, twenty-two parts.
Borate of soda, twenty-six parts.
Triturate together; in a quarter of an hour add a little water, then gradually, more, constantly rubbing, permit to settle, decant, wash the precipitate till the washings are insipid and dry. *Van Mons.*

The borate of mercury is said to resemble calomel in its action. Dose, two grains a-day, gradually augmenting.

HYDRARGYRI BROMIDUM.

BROMIDE OF MERCURY.

There are two bromides of mercury; one the protobromide, answering to calomel; the other the bibromide, like corrosive sublimate, an irritant poison.

PROTOBROMIDE OF MERCURY.

R. Solution of bromide of potassium,
at will.

Add a weak solution of protonitrate of mercury, as long as it causes a precipitate; wash this, and dry by a gentle heat. Dose, one grain a-day, gradually increased, as an alterative; four or five grains as a purgative. *Magendie.*

BI-BROMIDE OF MERCURY.

R. Bromine,
Mercury, each, equal parts.

Mix, and sublime. Dose, one-twentieth of a grain, gradually increased to a fourth. In syphilis. *Magendie.*

ETHEREAL SOLUTION OF BI-BROMIDE OF MERCURY.

R. Bi-bromide of mercury, one grain.
Sulphuric ether, one fl. drachm.

Dissolve. Dose, ten to twenty drops, in barley-water. In syphilis and lepra. *Werneck.*

HYDRARGYRI CHLORIDUM CORROSIVUM.

CORROSIVE SUBLIMATE.

Corrosive sublimate is a very energetic and poisonous preparation, requiring much care in its exhibition. It will produce the usual effects of the other mercurials, but is less apt to salivate. It is much used in secondary syphilis, and as an alterative, and also as an external application, as a stimulant, and escharotic. The best antidote, when poisonous doses have been taken, is albumen, followed by an emetic.

SOLUTION OF CORROSIVE SUBLIMATE.

R. Corrosive sublimate,
Muriate of ammonia, each, ten grains.
Distilled water (*imp. meas.*), one pint.

Dissolve. *Lond. Ph.*

A fl. ounce contains half a grain of the mercurial salt. The dose is from one to two fl. drachms, in some mucilaginous drink.

ALCOHOLIC SOLUTION OF CORROSIVE SUBLIMATE.

R. Corrosive sublimate, one to two grains.
Diluted alcohol, four fl. ounces.

Dissolve. A tablespoonful, night and morning, in decoction of sarsaparilla. *Ellis.*

POWDER OF CORROSIVE SUBLIMATE AND COPPER.

R. Corrosive sublimate,
Sulphate of copper, each, ten grains.
Nitrate of silver, six grains.

Mix. Applied to venereal excrescences, previously moistened with water. *Kruzer.*

POWDER OF CORROSIVE SUBLIMATE AND ZINC.

R. Corrosive sublimate,
Sulphate of zinc, equal parts.

Mix. In onychia maligna, sprinkled on the sore, and covered with lint soaked in tincture of myrrh. *Perkins.*

PILLS OF CORROSIVE SUBLIMATE.

R. Corrosive sublimate, five grains.
Distilled water, thirty to forty drops.
Confection of roses, one scruple.
Powdered liquorice, sufficient.

Dissolve the corrosive sublimate in the water, and add the other articles, and rub well together. Make forty pills. One, three to four times a-day, in syphilis. *Ellis.*

COMPOUND PILLS OF CORROSIVE SUBLIMATE.

R. Corrosive sublimate, half a grain.
Extract of Peruvian bark, ten grains.
" opium, half a grain.
Powdered Peruvian bark, sufficient.

Beat together, and make two pills. One, morning and evening. Each contains a quarter of a grain of the mercurial salt. They require much caution in their use. In syphilis. *Dupuytren.*

R. Corrosive sublimate, six grains.

Dissolve in

Distilled water, sufficient,
and add

Extract of hemlock, one drachm.
Powdered hemlock, sufficient.

Beat into mass, and make forty-eight pills, to be given like the above. Each pill contains an eighth of a grain of corrosive sublimate. *Ellis.*

CORROSIVE SUBLIMATE MIXTURE.

R. Corrosive sublimate, four grains.
Muriate of ammonia, eight grains.
Diluted alcohol, two fl. ounces.

Mix. A teaspoonful twice a-day, in barley-water, in syphilis. *Ainslie.*

R. Corrosive sublimate, eight grains.
Muriatic acid, twenty-four drops.
Comp. tincture of cardamom, eight fl. ounces.

Mix. Dose, two drachms morning and evening, in a wineglassful of sugar and water. In syphilis. *Guibourt.*

R. Corrosive sublimate, two grains.
Distilled water, six fl. ounces.
Spirit of cinnamon,
Syrup, each, one fl. ounce.

Mix. One or two spoonfuls twice or thrice a-day, in venereal cases. *Ellis.*

LOTION OF CORROSIVE SUBLIMATE.

R. Corrosive sublimate, three to six grains.
Distilled water, sufficient.

Dissolve, and add

Extract of hemlock, two drachms.
" chamomile, each, one fl. drachm.

Tincture of opium, one fl. drachm.
Honey of roses, one ounce.

Mix. As an application to venereal ulcers in the throat, and on the labia pudendi. *Rust.*

R. Corrosive sublimate, one drachm.
Distilled water, six fl. ounces.

Dissolve, and add

Muriate of ammonia, two drachms.
Nitrate of potassa, half an ounce.

Dissolve. As a wash in itch. *Good.*

COSMETIC LOTION WITH CORROSIVE SUBLIMATE.

R. Blanched sweet almonds, one ounce.
" bitter almonds, four drachms.

Cherry-laurel water, ten fl. ounces.

Make an emulsion, and add

Corrosive sublimate, six grains.
Tincture of benzoin, six fl. drachms.
Lemon juice, four fl. drachms.

Mix. As a wash for eruptions on the face, to be used morning and evening, previously to be shaken. It must be used with caution.

Cadet de Gassicourt.

LOTION OF CORROSIVE SUBLIMATE AND COPPER.

R. Corrosive sublimate, two grains.
Acetate of copper, six grains.
Distilled water, two pints.

Dissolve. As a wash in obstinate porrigo. *Augustin.*

LOTION OF CORROSIVE SUBLIMATE AND CAMPHOR.

R. Corrosive sublimate, half a drachm.
Camphor, one drachm.
Diluted alcohol, one ounce.

Dissolve. As an application to destroy condylomata. *Phæbus.*

BATEMAN'S MERCURIAL LOTION.

R. Corrosive sublimate, two grains.
Compound spirit of lavender, one fl. ounce.
Distilled water, four fl. ounces.

Dissolve. As a lotion in obstinate cutaneous eruptions. *Bateman.*

ANTACRID TINCTURE.

R. Powdered resin of guaiacum, one ounce.
Canada balsam, each, one scruple.
Corrosive sublimate, one scruple.
Oil of sassafras, two fl. drachms.
Alcohol, eight fl. ounces.

Dissolve the mercurial salt in one-half of the alcohol, and then add the remainder, and the other ingredients; after due a digestion, filter. Ten to twenty drops, morning and evening, in wine or water, in syphilis. *Ellis.*

COLLYRIUM OF CORROSIVE SUBLIMATE.

R. Corrosive sublimate, two grains.
Opium, ten grains.
Rose water, four fl. ounces.

Dissolve, and add

Mucilage of quince-seed, half an ounce.

Mix. *Van. Mons.*

INJECTION OF CORROSIVE SUBLIMATE.

R. Corrosive sublimate, three grains.
Water of rosemary, three fl. ounces.
Distilled water, each, three fl. ounces.

Dissolve. As an injection in fistula lachrymalis. *Beer.*

R. Corrosive sublimate, one drachm.
Alcohol, one fl. ounce.

Dissolve. Add from five to twenty drops to a solution of

Sulphate of zinc, five to ten grains;

In

Water, four fl. ounces.

A drachm to be injected into the urethra, three times a-day, in gonorrhœa. *Whately.*

YELLOW WASH.

R. Corrosive sublimate, one to three grains.
Lime water, one fl. ounce.
Mix. *Ellis.*

R. Corrosive sublimate, one scruple.
Carbonate of potassa, one drachm.

Rub together, and add gradually,

Distilled water, one pint.

Mix. *Span. Ph.*

These preparations are employed as lotions to venereal and phagedenic ulcers, and should be shaken up when used.

COSMETIC WASH.

R. Blanched bitter almonds, six ounces.
Beef tea, sixty-four ounces.

Make an emulsion, and add

Corrosive sublimate, half an ounce.

Lemon juice, ten ounces.

Solution of carbonate of potassa, half an ounce.

Incorporate gradually

Whites of eggs, six ounces.

Strain, and add

Camphor, rubbed with
mucilage gum Arabic,
half a drachm.

Mix well. It is said to be an effectual wash for eruptions on the face, but must be used with extreme caution, and very seldom. *Van Mons.*

HYDRARGYRI CHLORIDUM MITE.

CALOMEL.

Calomel is prepared by several modes, on the large scale, and is seldom or never made by the apothecary. The several processes will be found, with judicious observations on each, in the U. S. Dispensatory. It is used in almost every disease, either as a purgative, alterative, anthelmintic, &c. As a purgative it is peculiar, in not producing effects in proportion to the dose.

POWDER OF CALOMEL AND ANTIMONY.

R. Calomel,
Golden sulphuret of
antimony, equal parts.

Triturate together. This powder has been much celebrated under the name of *Plummer's Alterative*, as a deobstruent and alterative. The dose is from five to ten grains a-day, in divided doses. It should be used recently prepared, as in a short time it changes to sulphuret of mercury, and oxide of antimony. *Guibourt.*

POWDER OF CALOMEL AND JALAP.

R. Calomel, five grains.
Powdered jalap, ten grains.

Mix. As a purgative, to be mixed with syrup or molasses. *Ellis.*

R. Calomel, three grains.
Powdered jalap,
Sugar, each, ten grains.

Mix. Make a powder to be taken at night, or early in the morning, in bilious fevers, and obstructed bowels. *A. T. Thomson.*

POWDER OF CALOMEL, ANTIMONY, AND HENBANE.

R. Calomel, one grain.
Golden sulphuret of
antimony, three grains.
Powdered extract of
henbane, one grain.
Powdered sugar, half a drachm.

Mix. To be taken night and morning, in spasmodic diseases; is said to have proved very useful in nyctalopia. *Phæbus.*

POWDER OF CALOMEL, JALAP, AND RHUBARB.

R. Calomel,
Powdered jalap,
" rhubarb, each, five grains.
Oil of cinnamon, one drop.

Mix. As a purgative, to be given in syrup or molasses. *Ellis.*

POWDER OF CALOMEL AND FOXGLOVE.

R. Calomel, three grains.
Powdered foxglove, four grains.
Sugar, one drachm.

Mix, and divide into twelve powders. Two a-day, in chronic hydrocephalus. *Berends.*

POWDER OF CALOMEL, NITRATE OF POTASSA, &c.

R. Calomel, six grains.
Nitrate of potassa, one drachm.
Tartar emetic, half a grain.

Mix, and divide into six powders. One, every two hours, as a diaphoretic in febrile affections. *Ellis.*

POWDER OF CALOMEL AND OPIUM.

R. Calomel, sixteen grains.
Powdered opium, four grains.
" ipecacuanha, eight grains.

Mix, and divide into eight powders. One to be taken every hour or two. In dysentery. *Chapman.*

POWDER OF CALOMEL AND GAMBAGE

R. Calomel, five grains.
Powdered gamboge,
three to six grains.

Mix. As an anthelmintic. *Ellis*

R. Calomel,
Powdered gamboge,
" jalap,
" rhubarb,
" cinnamon, each,
two drachms.

Mix. Dose, five to twenty grains.
A. T. Thomson.

ANTHELMINTIC PURGATIVE.

R. Calomel, three grains.
Compound powder of
scammony, twelve grains.

Mix. To be taken at once, in cases of lumbrici.
A. T. Thomson.

POWDER OF CALOMEL AND PINK ROOT.

R. Calomel, four grains.
Powdered pink root, ten grains.

Mix. To be taken two mornings in succession;
also on afternoon of second day, followed by a
mild purgative; as an anthelmintic for children
over four years of age. Ellis.

DRY COLLYRIUM OF CALOMEL.

R. Calomel,
Powdered sugar, each, half a drachm.
" opium, ten grains.

Mix. To be blown into the eye, in ulcers of
the cornea. Radius.

CALOMEL PILLS.

R. Calomel, half an ounce.
Powdered gum Arabic, one drachm.
Syrup, sufficient.

Mix the calomel and gum, then beat with syrup
into mass, and divide into two hundred and forty
pills. U. S. Ph.

A very convenient form to give calomel,
whether as purgative, alterative, &c.; each pill
contains one grain of calomel.

PILLS OF CALOMEL, QUINIA, &c.

R. Calomel, six grains.
Powdered opium, three grains.
Sulphate of quinia, twelve grains.
Syrup, sufficient.

Beat into mass, and form twelve pills. One,
night and morning, as an alterative in condi-
tions following bilious fever. Ellis.

PILLS OF CALOMEL AND ACETATE OF LEAD.

R. Acetate of lead, half a drachm.
Calomel, five grains.
Confection of roses, sufficient.

Form mass, and divide into ten pills. One to be
given every two to four hours, in hematemesis,
&c. Ellis.

PILLS OF CALOMEL AND DANDELION.

R. Calomel, four grains.
Extract of dandelion, eighteen grains.

Mix. To be taken in a day, in divided doses, in
abdominal obstructions. Brera.

PILLS OF CALOMEL, SQUILL, &c.

R. Plummer's powder, twelve grains.
Ammoniac, two grains.
Extract of dandelion, three grains.
Powdered squill, half a drachm.

Beat together, and form pills of three grains.
Dose, five, three or four times a-day, in engorge-
ments of the abdominal viscera, with anasarca
supervening on intermittent fever. St. Marie

PILLS OF CALOMEL AND IRON.

R. Calomel,
Golden sulphuret of antimony,
Sulphate of iron,
Myrrh, each, one drachm.
Syrup, sufficient.

Beat into mass, and form pills of three grains.
Dose, four, morning and evening, in the same
cases as above. Swediaur.

PILLS OF CALOMEL AND CATECHU.

R. Powdered catechu,
Copaiba, each, three drachms.
Calomel, one scruple.
Syrup of comfrey, sufficient.

Beat into mass, and make one hundred and fifty
pills. Four, thrice a-day, in leucorrhœa or gon-
orrhœa. St. Marie.

COMPOUND CALOMEL PILLS.

R. Calomel,
Oxysulphuret of anti-
mony, each, two drachms.
Powdered guaiacum resin,
half an ounce.
Molasses, two drachms.

Rub the calomel with the antimony, and then
with the other ingredients, till well incorporated.
Lond. Ph. 1836.

These pills are known as *Plummer's pills*, and
have been much employed in chronic rheuma-
tism, and obstinate cutaneous affections, espe-
cially when there is a syphilitic taint. The dose
is from three to six grains, twice a-day.

R. Calomel, two scruples.
Precipitated sulphuret
of antimony, one drachm.
Guaiacum, two drachms.
Copaiba, sufficient.

Mix, and make sixty pills. Three at night, in venereal herpes, till mouth is affected; also using frequent tepid baths. *Ainslie.*

COMPOUND CATHARTIC PILLS.

R. Powdered compound extract of colocynth, half an ounce.
Extract of jalap,
Calomel, each, three drachms.
Gamboge, two scruples.

Mix, and beat into a mass with water. To be divided into one hundred and eighty pills. *U. S. Ph.*

As a cathartic or laxative, in cases where there is derangement of the liver. Dose, as a purgative, three pills; as a laxative, one pill.

PILLS OF CALOMEL AND COLOCYNTH.

R. Compound extract of colocynth, forty-eight grains.
Calomel, one scruple.

Mix, and divide into twenty pills. Two or three will act as a cathartic. *Ellis.*

PILLS OF CALOMEL AND ANTIMONY.

R. Calomel, ten grains.
Golden sulphuret of antimony, one scruple.
Extract of pot marigold,
" hemlock, each, two drachms.

Mix, and beat into mass, and form pills of two grains. Dose, five, thrice a-day, in chronic indurations. *Rust.*

R. Calomel, six grains.
Kermes mineral, twelve grains.
Syrup of elder, sufficient

to make six pills. One every two hours, in the decline of peripneumonia, and in the cough of children arising from worms. *Brera.*

PILLS OF CALOMEL AND GUAIAIACUM.

R. Calomel,
Resin of guaiacum, each, two drachms.
Powdered mallow, four ounces.
Syrup, sufficient.

Form mass, and make pills of four grains. Dose, four or five a-day, in syphilis. *Alibert.*

PILLS OF CALOMEL, JALAP, &C.

R. Calomel,
Resin of jalap,
Compound extract of rhubarb,
Soap, equal parts.

Form mass, and make pills of one grain. Dose, ten to twelve grains, as a purgative and anthelmintic. *Par. Cod.*

PILLS OF CALOMEL AND OPIUM.

R. Calomel, two grains.
Opium, half a grain.

Mix. Thrice a-day, in neuralgia of the face. *Leslie.*

ELECTUARY WITH CALOMEL, &C.

R. Calomel, ten grains.
Powdered rhubarb,
" semen contra,
" valerian, each, two drachms.

Conserve of worm-

wood, one ounce and a half.

Oxymel of squill, sufficient.

Mix. A drachm to half an ounce, in epilepsy complicated with worms. *Swediasur.*

CALOMEL OINTMENT.

R. Calomel, one drachm.
Rose ointment, three drachms.

Mix. As an application in herpes. *Dupuytren.*

R. Calomel, one drachm.
Sublimed sulphur, two drachms.
Lard, one ounce.

Mix well. In obstinate cutaneous affections. *Fouquier.*

OINTMENT OF CALOMEL AND SQUILL.

R. Calomel,
Powdered squill, each, half a drachm.
Lard, two drachms.
Oil of roses, four drops.

Mix. In chronic swellings of the joints. *Dupuytren.*

OINTMENT OF CALOMEL AND ACETATE OF COPPER.

R. Calomel,
Acetate of copper, each, one scruple.
Lard, eleven drachms

Mix well. As an application in porrigo. *Cadet de Gassicourt.*

OINTMENT OF CALOMEL, ALUM, &C.

R. Calomel, two drachms.
Burnt alum,
Litharge, each, half an ounce
Oil of turpentine, two fl. drachms.
Simple ointment, one ounce and a half.

Mix well. As an application to tinea capitis, to be applied at night, and washed off in the morning. *Ellis.*

CALOMEL AND CAMPHOR OINTMENT.

R. Calomel, half a drachm.
Camphor, twelve grains.
Lard, one ounce.

Mix. As an application in lichen. *Bielt.*

HYDRARGYRI ET QUINIAE
CHLORIDUM.

CHLORIDE OF MERCURY AND QUINIA.

R. Bichloride of mercury, one part.
Muriate of quinia, three parts.

Dissolve separately in the smallest quantity of water, and mix the solutions. Collect the precipitate, and dry by a gentle heat. *McDermott.*

PILLS OF CHLORIDE OF MERCURY AND
QUINIA.

R. Chloride of mercury
and quinia, fifteen grains.
Opium, six grains.
Crumb of bread, sufficient.

Mix well, and make thirty pills. One, thrice a-day, to produce salivation. *Hamilton.*

HYDRARGYRI CYANURETUM.

CYANURET OF MERCURY.

R. Ferrocyanuret of iron, four ounces.
Red oxide of mercury, three ounces.
Distilled water, three pints.

Put the ferrocyanuret of iron and the oxide of mercury, well powdered and mixed, into a glass vessel, and pour on them two pints of the water. Boil, and stir, and if in half an hour a blue color remains, add a small portion of oxide of mercury, and continue boiling until the mixture becomes of a yellowish color; then filter, wash the residue in a pint of distilled water, and again filter; mix the solutions, and evaporate, and crystallize. Purify these by dissolving in distilled water, filtering and evaporating. *U. S. Ph.*

It is given in doses of a sixteenth to an eighth of a grain, in syphilis, chronic inflammation of the thoracic and abdominal organs, &c., and also is employed externally in porrigo, and other cutaneous affections.

SOLUTION OF CYANURET OF MERCURY.

R. Cyanuret of mercury, four grains.
Distilled water, eight fl. ounces.

Dose, half a drachm to a drachm. *Chaussier.*

COMPOUND PILLS OF CYANURET OF
MERCURY.

R. Cyanuret of mercury, six grains.
Opium, twelve grains.
Crumb of bread, one drachm.
Honey, sufficient.

Mix, and make ninety-six pills. One, morning and evening. *Guibourt.*

R. Cyanuret of mercury, eighteen grains.
Muriate of ammonia,
Extract of aconite, each,
three drachms.

“ box, one ounce and a half.

Oil of anise, one scruple.

Beat together, and make four hundred pills. Two, morning and evening. *Cadet.*

COMPOUND TINCTURE OF CYANURET OF
MERCURY.

R. Cyanuret of mercury, eighteen grains.
Water, fourteen fl. ounces.
Alcohol, ten fl. ounces.
Muriate of ammonia,
Extract of aconite, each,
three drachms.

“ box, one ounce and a half.

Oil of sassafras, twenty-four drops.

Dissolve the cyanuret in the water, add the ammonia, extract, and alcohol, let stand for some hours, filter, and add essential oil.

GARGLE OF CYANURET OF MERCURY.

R. Cyanuret of mercury, ten grains.
Barley water, one pint.
Honey of roses, one ounce.

Mix. As a gargle. *Brera.*

R. Sarsaparilla, half an ounce.
Water, sixteen fl. ounces.

Boil, and add towards close of operation

Fresh hemlock, two drachms.

Strain, and add to eight ounces of filtered liquid,

Cyanuret of mercury, two grains.

As a gargle. *Muller.*

OINTMENT OF CYANURET OF MERCURY.

R. Cyanuret of mercury, twelve grains.
Lard, one ounce

Rub well together. As an application to venereal ulcers. *Fo3*

R. Cyanuret of mercury, sixteen grains
Lard, one ounce.

Oil of lemon, fifteen drops.

Rub together. As an application to moist tetter. *Bielt.*

HYDRARGYRI IODIDUM.

GREEN IODIDE OF MERCURY.

R. Mercury,	one ounce.
Iodine,	five drachms.
Alcohol,	sufficient.

Rub the mercury and iodine together, adding sufficient alcohol to form a soft paste, and continue rubbing till globules disappear. Dry the iodide in the dark, with a gentle heat, and keep from light, in a well-stopped bottle. *U. S. Ph.*

The dose is about a grain a-day, gradually increased to three or four. Used in scrofula, and scrofulous syphilis.

POWDER OF IODIDE OF MERCURY.

R. Iodide of mercury,	one to eight grains.
Magnesia,	one drachm.

Mix. Divide into twelve powders. One, three times a-day. *Radius.*

PILLS OF IODIDE OF MERCURY.

R. Iodide of mercury,	five grains.
Confection of roses,	sufficient.

Mix, and make thirty pills. One, three times a-day. *Ellis.*

R. Iodide of mercury,	one drachm.
Confection of dog-rose,	three drachms.
Powdered ginger,	one drachm.

Beat together. Dose, five to ten grains. *Lond. Ph.*

R. Iodide of mercury,	one grain.
Extract of juniper,	twelve grains.
Powdered liquorice,	sufficient.

Mix, and make eight pills. Two, morning and evening, gradually increasing to double the number. *Magendie.*

COMPOUND PILLS OF IODIDE OF MERCURY.

R. Iodide of mercury,	six grains.
Extract of opium,	four grains.
Lactucarium,	twenty-four grains.
Extract of guaiacum,	forty-eight grains.

Beat together, and form forty-eight pills. In syphilis of children; one pill to a child six months old, and two to four at more advanced ages. *Ricord.*

R. Iodide of mercury,	half a drachm.
Extract of guaiacum,	one drachm.
“ lettuce,	two scruples.
“ sarsaparilla,	sufficient.

Mix, and make seventy-two pills. One, and then two daily. *Bielt.*

R. Iodide of mercury,	
Extract of lettuce, each,	half a drachm.
“ hemlock,	one drachm.

Mix, and make thirty pills. *Ricord.*

R. Iodide of mercury,	two scruples.
Aloes,	} each,
Sulphate of iron,	
Myrrh,	
Oil of savine,	twenty drops.

Mix, and make twenty-four pills. One, thrice a-day, in amenorrhœa. *Barbour.*

OINTMENT OF IODIDE OF MERCURY.

R. Iodide of mercury,	one scruple.
Lard,	one ounce and a half.

Rub well together. In scrofulous swellings, indolent glandular tumors, &c. *Soubeyran.*

R. Iodide of mercury,	one ounce.
White wax,	two ounces.
Lard,	six ounces.

Melt the wax and lard, and stir in the iodide. *Lond. Ph.*

As a dressing to scrofulous ulcers, &c.

R. Iodide of mercury,	six grains.
Acetate of morphia,	eight grains.
Lard,	one ounce.

Rub well together. As an application to obstinate glandular swellings. *Pelletan.*

R. Iodide of mercury,	one drachm.
Soap,	half a drachm.
Rose water,	two drachms.
Rose ointment,	six drachms.

Mix. *Riecke.*

HYDRARGYRI IODIDUM RUBRUM.

RED IODIDE OF MERCURY.

R. Corrosive sublimate,	one ounce.
Iodide of potassium,	ten drachms.
Distilled water,	two pints.

Dissolve the corrosive sublimate in a pint and a half, and the iodide in half a pint, of the water, and mix the solutions. Filter, and wash the collected precipitate with distilled water. Dry by a gentle heat, and keep in a well-stopped bottle. *U. S. Ph.*

Much more active than the iodide. Used in the same cases, in doses of a sixteenth, gradually increased to the fourth of a grain.

PILLS OF RED IODIDE OF MERCURY.

R. Red iodide of mercury,	one grain.
Extract of juniper,	twelve grains.
Powdered liquorice,	sufficient.

Mix, and make eight pills. One, morning and evening. *Guibourt.*

R. Red iodide of mercury,	five grains.
Syrup,	sufficient.

Rub well together, and then with

Crumb of bread,
Sugar, each, sufficient
to make sixty pills. Two, morning and evening,
gradually increasing. *Radius.*

WASH OF RED IODIDE OF MERCURY.

R. Red iodide of mercury, twelve grains.
Distilled water, six fl. ounces.
Mix. As a lotion to scrofulous ulcers. *Radius.*

TINCTURE OF RED IODIDE OF MERCURY.

R. Red iodide of mercury, twenty grains.
Alcohol (.837), one fl. ounce and
a half.
Dissolve. Dose, five to ten drops, in distilled
water. *Foy.*

ETHEREAL TINCTURE OF RED IODIDE OF MERCURY.

R. Red iodide of mercury, twenty grains.
Sulphuric ether, one ounce and a half.
Dissolve. Rather more powerful than the last,
and therefore to be given in smaller doses. *Magendie.*

OINTMENT OF RED IODIDE OF MER- CURY.

R. Prepared like the ointment of the
iodide. *Lond. Ph.*

It is more active than the ointment of the
iodide.

R. Red iodide of mercury, one scruple.
Lard, one ounce and a half.
Rub well together. As a dressing to obstinate
venereal ulcers. *Soubeiran.*

R. Red iodide of mercury, fifteen grains.
Lard, one ounce.
Essence of bergamot, twenty drops.
Mix well. As an application to chronic cuta-
neous affections. *Foy.*

HYDRARGYRI NITRAS.

NITRATE OF MERCURY.

NITRATE OF THE PROTOXIDE OF MER- CURY.

Mercury,
Nitric acid, equal parts.
Dissolve, with aid of a gentle heat, then boil
until a yellow sediment is formed; decant, and
permit to crystallize. *Guibourt.*

PILLS OF PROTONITRATE OF MERCURY.

R. Protonitrate of mercury, ten grains.
Extract of liquorice, forty grains.
Beat into mass with a little water, and form
sixty pills. *Foy.*

OINTMENT OF NITRATE OF MERCURY.

R. Mercury, one ounce.
Nitric acid, fourteen fl. drachms.
Fresh neat's foot oil, nine fl. ounces.
Lard, three ounces.

Dissolve the mercury in the acid, melt the oil
and lard together, and when they become thick,
on cooling, add the solution and mix well. *U. S. Ph.*

A stimulant and alterative application, used
in most cases diluted with lard. Employed in
various skin diseases, &c. It is known under
the name of *Citrine ointment*.

OINTMENT OF NITRATE OF MERCURY AND LEAD.

R. Mercury, four parts.
Lead, half a part.

Dissolve separately in
Nitric acid, sufficient.

Then mix

Oil of almonds, twenty-four parts;
Lard, forty-eight parts;

first with the mercurial, and then with the sa-
turnine solution; rub well together. *Van Mons.*

Advised by Armstrong in porrigo favosa.

LINIMENT OF NITRATE OF MERCURY.

R. Ointment of nitrate of
mercury, two ounces and a half.
Simple cerate, seven ounces and a half.
Olive oil, five fl. ounces and a half.
Mix well. *Beasley.*

R. Ointment of nitrate of mercury,
Almond oil, equal parts.
Triturate together till perfectly mixed. *Sir H. Halford.*

ACID NITRATE OF MERCURY.

R. Mercury, two ounces.
Pure nitric acid, one and a half fl.
ounces.

Water, one ounce and a half.

Mix the acid with the water, dissolve in them
the mercury by the aid of heat, and evaporate
to the bulk of two ounces and a half, (Imp.
meas.). *Dub. Ph.*

R. Mercury, by weight, four parts.
Nitric acid (1.321), by weight,
eight parts.

Dissolve, and evaporate the solution to nine parts. *Par. Codex.*

This solution is much used as a caustic. The part to which it is applied becomes white, and in a few days a yellow scab falls off.

HYDRARGYRI OXIDUM NIGRUM.

BLACK OXIDE OF MERCURY.

R. Calomel, four ounces.
Potassa, each, one pint.
Water, one pint.

Dissolve the potassa in the water, let settle and decant; add the calomel, stirring well till the black oxide is formed, decant, wash the oxide with distilled water, and dry with a gentle heat. *U. S. Ph.*

Alterative, purgative, and sialagogue. Dose, one quarter of a grain to two grains.

BLACK WASH.

R. Calomel, one drachm.
Lime water, four fl. ounces.
Mix well. As a lotion to venereal and phagedenic ulcers. *Ellis.*

R. Calomel, half a drachm.
Powdered opium, two drachms.
Lime water, two fl. ounces.

Mix. As a dressing for chancres. *Rust.*

PILLS OF BLACK OXIDE OF MERCURY.

R. Black oxide of mercury, one drachm.
Confection of roses, three drachms.
Powdered chamomile, half a drachm.

Mix. As a substitute for blue pill. *Tyson.*

OINTMENT OF BLACK OXIDE OF MERCURY.

R. Black oxide of mercury, one part.
Lard, sixteen parts.

Subject to a temperature of 300° to 320° for an hour, stirring continually, remove, and stir till cold. As a substitute for mercurial ointment. *Donovan.*

HYDRARGYRI OXIDUM RUBRUM.

RED OXIDE OF MERCURY.

RED PRECIPITATE.

R. Mercury, thirty-six ounces.
Nitric acid, eighteen fl. ounces.
Water, two pints.

Dissolve the mercury with a gentle heat, in the acid and water, and evaporate to dryness. Rub to powder, and heat in a shallow vessel, as long as red vapors arise. *U. S. Ph.*

R. Purified mercury, at will.
Put in an open glass vessel, with narrow mouth and broad bottom; expose to heat of 600°, till converted into red scales. *Dub. Ph., 1826.*

R. Corrosive sublimate, four ounces.
Solution of potassa, twenty-eight fl. ounces.
Distilled water, six pints.

Dissolve the corrosive sublimate in the water, and add the solution of potassa. Let settle, decant, wash the precipitate with distilled water, and dry with a gentle heat. *Lond. Ph., 1836.*

These preparations are seldom used internally in this country, as they are harsh and uncertain in their operation, but are much employed externally as stimulants and escharotics.

PILLS OF RED OXIDE OF MERCURY.

R. Red oxide of mercury, one grain.
Powdered opium, each, one grain.
Oil of cloves, two drops.

Mix, and make three pills. One, every night for a week, in venereal complaints. *Ellis.*

R. Red oxide of mercury, three grains.
Acetate of morphia, one grain.
Powdered mallow root, sufficient.
Water, each, sufficient.

Mix, and form twenty pills. One, morning and evening. *Radius.*

POWDER OF RED OXIDE OF MERCURY.

R. Red oxide of mercury, half a drachm.
Burnt alum, each, half a drachm.
Powdered savine, two drachms.

As an application to condyloma, whitlow, &c. *Phæbus.*

OINTMENT OF RED OXIDE OF MERCURY.

R. Red oxide of mercury, in fine powder, one drachm.
Simple ointment, one ounce.

Soften the ointment by a gentle heat, and add oxide of mercury and mix well. *U. S. Ph.*

An efficacious, stimulating application to foul and indolent ulcers, psorophthalmia, &c. Where it is too stimulating, dilute with lard.

R. Red oxide of mercury, one drachm.
Venice turpentine, one ounce.

Mix. As an application to indolent ulcers, and also used in itch. *Ellis.*

OINTMENT OF RED OXIDE OF MERCURY AND SULPHUR.

R. Red oxide of mercury, one part.
Mercury, each, one part.
Precipitated sulphur, sixteen parts

Triturate till globules disappear, and add

Lard, thirty-two parts.
Rub well together. As a friction in the cure of itch. *Swedliaur.*

BROWN OINTMENT OF RED OXIDE OF MERCURY.

R. Red oxide of mercury, one part.
Basilicon ointment, six parts.
Rub well together. Used in the treatment of indolent venereal ulcers. *Saunders.*

OINTMENT OF RED OXIDE OF MERCURY AND TIN.

R. Red oxide of mercury, two drachms.
Amalgam of equal parts of mercury and tin, four drachms.
Rose ointment, one ounce.
Rub well together, and add
Oil of peppermint, twenty drops.
Highly spoken of in hemorrhoidal tumors. *Cadet de Gassicourt.*

OINTMENT OF RED OXIDE OF MERCURY AND CINNABAR.

R. Red oxide of mercury, one ounce.
Cinnabar, one drachm.
Lard, one pound.
Rub well together. Employed in chronic inflammations of the eyelids. *Giordano.*

OINTMENT OF RED OXIDE OF MERCURY AND ZINC.

R. Red oxide of mercury, one part.
Sulphate of zinc, two parts.
Lard, ninety-six parts.
Rub well together. In scrofulous ophthalmia, as an application to the edges of the eyelids. *Dupuytren.*

OINTMENT OF RED OXIDE OF MERCURY AND LEAD.

R. Red oxide of mercury,
Prepared calamine,
Burnt alum,
Acetate of lead, each, one drachm.
Corrosive sublimate, twelve grains.
Rose ointment, one ounce.
Rub well together. Celebrated in the treatment of herpetic affections. *Guibourt.*

HYDRARGYRI PHOSPHAS. PHOSPHATE OF MERCURY.

R. Solution of nitrate of mercury, at will.
Add to it a solution of phosphate of soda, as long as any precipitate is formed; decant, wash the precipitate well, and dry it. *Van Mons.*

This has been highly praised in secondary syphilis. The dose is half a grain to a grain, combined with sugar or some aromatic powder.

PILLS OF PHOSPHATE OF MERCURY.

R. Phosphate of mercury, nine grains.
Tartar emetic, one grain.
Opium, six grains.
Conserve of roses, sufficient.
Mix, and make nine pills. One in the evening, at bedtime. *Radius.*

HYDRARGYRI SULPHAS.

PERSULPHATE OF MERCURY.

R. Mercury, ten ounces.
Sulphuric acid, six fl. ounces.
Expose them to heat in a glass vessel, and increase the heat, until the mass becomes white and perfectly dry. *Dub. Ph., 1850.*

Not used as a remedy, but serves to form corrosive sublimate, calomel, and turpeth mineral.

YELLOW SULPHATE OF MERCURY.

TURPETH MINERAL.

R. Persulphate of mercury, one part.
Warm water, twenty parts.

Rub together in an earthenware mortar, and pour off the supernatant liquid; wash the yellow powder with warm distilled water, as long as the washings are precipitated by caustic potassa. Dry. *Dub. Ph., 1826.*

R. Mercury, four ounces.
Sulphuric acid, six ounces.

Mix them in a glass vessel, and boil on a sand-bath till a dry white mass remains. Rub this into powder, and throw it into boiling water. Pour off the supernatant liquor, wash the yellow precipitate repeatedly in hot water, and dry it. *U. S. Ph.*

Turpeth mineral is alterative, errhine, and emetic. The dose, as an alterative, is from a quarter to half a grain; as an errhine, one grain, mixed with five grains of some bland powder; as an emetic, two to five grains.

COMPOUND POWDER OF SULPHATE OF MERCURY.

R. Sulphate of mercury, one grain.
Powdered asarabacca, one drachm and a half.
Mix, and divide into eight powders. One to be used as an errhine, twice a-day, in gutta serena, ozæna, &c. *Ellis.*

OINTMENT OF SULPHATE OF MERCURY.

R. Sulphate of mercury, one part.
Lard, eight parts.

Rub well together. As an application to herpes, porrigo, &c. *Soubeiran.*

R. Sulphate of mercury, two drachms.
Chloride of lime, three drachms.
Oil of almonds, six drachms.
Lard, two ounces.

Melt together the oil and lard, and mix the two powders. More stimulating than the last. Used in obstinate cutaneous affections. *Chevallier.*

HYDRARGYRI SULPHURETUM NIGRUM.

BLACK SULPHURET OF MERCURY.
ETHIOPS MINERAL.

R. Mercury, one pound.
Sulphur, each, one pound.

Rub together till all globules disappear. *U. S. Ph.*

Employed as an alterative in cutaneous affections and glandular swellings. Dose, from five to thirty grains, or even more, several times a-day.

POWDER OF BLACK SULPHURET OF MERCURY.

R. Black sulphuret of mercury, four drachms.
Prepared chalk, two ounces.
Powdered amber, one ounce.

Mix well, and divide into sixty-four powders. One, and afterwards two a-day, in a glass of sugar and water. In scrofulous affections. *St. Marie.*

R. Black sulphuret of mercury, one drachm.
Prepared oyster shell, each, one drachm.
Powdered orange peel, half a drachm.

Mix, and make ten powders. Three a-day, as an anthelmintic. *Phabus.*

PILLS OF BLACK SULPHURET OF MERCURY.

R. Black sulphuret of mercury, two drachms.
Extract of bitter-sweet, each, two drachms.
Powdered mallow, sufficient.

Mix, and form one hundred and twenty pills. Four to five, three times a-day. In obstinate cutaneous affections. *Radius.*

R. Black sulphuret of mercury, one scruple.
Extract of dandelion, three drachms.
Ammoniac, two drachms.

Powdered squill, half a drachm.
Syrup, sufficient.

Beat together, and make pills of three grains. Dose, five, thrice a-day. In ascites with engorgement of the abdominal viscera. *St. Marie.*

R. Black sulphuret of mercury, half an ounce

Guaiacum,
Soap, each, two drachms.
Golden sulphuret of antimony, two scruples.

Extract of horehound, sufficient.

Beat together, and form pills of two grains. Ten, morning and evening. In chronic rheumatism. *Baldinger.*

HYDRARGYRI SULPHURETUM RUBRUM.

RED SULPHURET OF MERCURY.
CINNABAR.

R. Mercury, forty ounces.
Sulphur, eight ounces.

Melt the sulphur, and mix the mercury with it over the fire. When mass begins to swell remove from fire, and cover the vessel, to prevent combustion; when cool, powder, and sub-
U. S. Ph.

At one time, cinnabar was much used as an alterative and anthelmintic, but it is now seldom employed except for fumigation.

Dose, five to ten grains.

POWDER OF RED SULPHURET OF MERCURY.

R. Red sulphuret of mercury, half an ounce.
Powdered sugar candy, three ounces.
Oil of cinnamon, half a drachm.

Rub well together. At one time celebrated as cordial, stomachic, and analeptic, in doses of a scruple. *Swediaur.*

PILLS OF RED SULPHURET OF MERCURY.

R. Red sulphuret of mercury, half a drachm.
Powdered Virginia snakeroot, each, half a drachm.
Syrup, sufficient.

Mix, and form twelve pills. Two to be taken thrice a-day; alterative and diaphoretic; useful in obstinate cutaneous affections. *Ellis.*

R. Red sulphuret of mercury, one drachm.
Extract of wormwood, one drachm.
" hemlock, each, one drachm.

Beat together, and form pills of two grains. Dose, seven, morning and evening, in venereal glandular swellings. *Augustin.*

FUMIGATING POWDER.

R. Red sulphuret of mercury, four drachms.
 Olibanum, two drachms.
 Mix. To be thrown on a red hot iron, and the diseased parts exposed to the fumes. In herpes, and venereal pustules. *Foy.*

CERATE OF RED SULPHURET OF MERCURY.

R. Powdered red sulphuret of mercury, fifteen grains.
 Yellow resin, half an ounce.
 Yellow wax,
 Lard, each, half a pound.

Melt the resin, wax, and lard together, and add the sulphuret. As a dressing to ill-conditioned ulcers. *Swediaur.*

OINTMENT OF RED SULPHURET OF MERCURY.

R. Red sulphuret of mercury, one drachm and a half.
 Muriate of ammonia, half a drachm.
 Lard, two ounces.
 Rose water, one fl. drachm.

Rub well together. Advised in pityriasis, to diminish the itching. *Radiis.*

R. Red sulphuret of mercury, one drachm.
 Camphor, one scruple.
 Simple cerate, one ounce.

Rub well together. Used in the treatment of herpes. To be applied at least twice a-day. *Alibert.*

HYDRARGYRI TARTRAS.

TARTRATE OF MERCURY.

R. Black oxide of mercury, at will.
 Tartaric acid, sufficient.

Boil together, in a due proportion of water, until the oxide is dissolved; filter, and evaporate to the point of crystallization; wash the crystals. *Swediaur.*

Dose, one to two grains, twice a-day.

TARTRATE OF MERCURY AND POTASSA.

R. Boiling saturated solution of nitrate of mercury, at will.

Add, drop by drop, a boiling saturated solution of bitartrate of potassa, as long as a precipitate is formed; decant, wash the precipitate well with cold water, dissolve in boiling water, and set aside to crystallize. *Van Mons.*

Dose, one to three grains. Once very celebrated as *Pressavin's vegeto-mercurial liquor.*

HYDRASTIS.

YELLOW ROOT.

This is the root of *Hydrastis Canadensis*, a native plant, more common in the western than in the eastern States. The flowers are yellow and fugacious, and are succeeded by red fruit, resembling raspberries, but not eatable.

Sex. Syst. Polyand. polygyn. *Nat. Syst.* Ranunculaceæ.

Linn. Sp. Pl. 784. *Griffith, Med. Bot.* 82.

The root is contorted, rugose, and of a bright yellow color. It has a strong, somewhat narcotic odor, and a very bitter taste. It is a powerful tonic bitter, and is also used in decoction as a wash in chronic ophthalmia. It is given in powder or decoction.

HYOSCYAMUS.

HENBANE.

Several species of *Hyoscyamus* are medicinal, but the only one that is officinal, is the *H. niger*, a native of Europe, and now naturalized in some parts of the United States. It has a lurid appearance, and a rank, unpleasant odor.

Sex. Syst. Pentand. monog. *Nat. Syst.* Solanaceæ.

Linn. Sp. Pl. 257. *Griffith, Med. Bot.* 484.

All parts of the plant are active, but the leaves are generally employed, though the seeds also are officinal. The dried leaves have but little taste or smell. The seeds are small, roundish, somewhat reniform, of a grayish color; of an unpleasant smell, and an oily, somewhat bitter taste. They both owe their properties to an alkaloid, called *hyoscyamia*. They are anodyne and hypnotic, and are used to relieve pain, to procure sleep, and to quiet an excited nervous system. The dose of the powdered leaves is from five to ten grains; of the seeds rather less.

AQUEOUS EXTRACT OF HENBANE.

R. Henbane leaves, one pound.
 Bruise them in a stone mortar, adding a little water; express, heat the juice to boiling point, strain, and evaporate. *U. S. Ph.*

Dose, two grains, twice a-day, to be gradually increased till the system is affected.

ALCOHOLIC EXTRACT OF HENBANE.

R. Henbane leaves, in coarse powder, one pound.
 Diluted alcohol, four pints.

Moisten the henbane with half a pint of the alcohol, and let stand for twenty-four hours; then put in a displacement apparatus, and exhaust, with the remainder of the alcohol, adding sufficient water at the close, to drive off all the spirit. Distil off the alcohol, and evaporate the residue to proper consistence. *U. S. Ph.*

Dose, as for the last, and in the same manner.

FLUID EXTRACT OF HENBANE.

R. Henbane leaves, in coarse
powder, eight ounces.
Diluted alcohol, sufficient.
Sugar, eight ounces.

Pour one pint of the diluted alcohol on the powder, macerate twenty-four hours, put in a percolator and pass three pints of tincture. Evaporate the tincture to ten fl. ounces, dissolve the sugar in it while hot, and when cold, two fl. ounces of alcohol (.835), or sufficient to measure a pint. *C. A. Smith.*

R. Extract of henbane, half an ounce.
Water, ten fl. ounces.
Sugar, eight ounces.

Rub the extract with the water, add the sugar, dissolve, and then add sufficient alcohol to measure a pint. *W. Procter, Jr.*

Dose, a teaspoonful.

COMPOUND POWDER OF HENBANE.

R. Powdered extract of henbane,
Oxide of zinc, each, ten grains.
Magnesia, half a drachm.
Sugar of milk, two drachms
and a half.

Mix, and divide into ten powders. One every three hours, as a sedative, in nervous attacks. *Vogler.*

COMPOUND PILLS OF HENBANE.

R. Extract of henbane,
" opium,
" belladonna,
" hemlock, each, one ounce.

Beat together, and form pills of one grain. Said to be useful in neuralgia of the face, in doses of one to six a-day. *Pierquin.*

R. Extract of henbane, half a drachm.
" opium, two grains.
Powdered foxglove,
" ipecacuanha, each,
five grains.

Beat together, and make thirty pills. One every three hours, in hooping-cough. *Phæbus.*

R. Extract of henbane,
" valerian,
Oxide of zinc, equal parts.

Beat together, and form pills of three grains. In neuralgia, in doses of one to ten. *Foy.*

R. Extract of henbane, two scruples.
Powdered myrrh, one drachm
and a half.
" squill, half a drachm.
Distilled water, sufficient.

Beat together, and make thirty pills. Two, night and morning, in catarrh and phthisis, with debility of the pulmonary organs. *Paris.*

PILLS OF HENBANE AND OPIUM.

R. Extract of henbane,
" hemlock, each,
fifteen grains.

Powdered opium, four grains.

Beat into mass, and make ten pills. One at night, as an anodyne. *Ellis.*

PILLS OF HENBANE AND IPECACUANHA.

R. Extract of henbane, ten grains.
Powdered ipecacuanha, five grains.

Mix, and make ten pills. One, every hour or two, in irritability of the bowels, with flatulence. *Ellis.*

INFUSION OF HENBANE.

R. Leaves of henbane, half an ounce.
Boiling water, one pint.

Infuse. As a lotion to painful ulcers. *Ellis.*

COMPOUND INFUSION OF HENBANE.

R. Henbane leaves,
Poppy heads,
Mallow, each, one ounce.
Boiling water, four pints.

Infuse for an hour, and strain. As a fomentation to painful tumors, &c. *Radius.*

MIXTURE OF HENBANE.

R. Oil of almonds, half an ounce.
Powdered gum Arabic, two drachms.
Oil of fennel, one drop.
Extract of henbane, six grains.
Sugar, half an ounce.
Water, two fl. ounces.

Rub well together. A spoonful every two hours, in catarrh. *Foy.*

MIXTURE OF HENBANE AND ANTIMONY.

R. Extract of henbane, ten grains.
Antimonial wine, two fl. drachms.
Dissolve. In hooping-cough, in doses of five to ten drops. *Augustin.*

TINCTURE OF HENBANE.

R. Henbane leaves, four ounces.
Diluted alcohol, two pints.

Macerate for fourteen days, express, and filter. *U. S. Ph.*

Dose, a fl. drachm.

ETHEREAL TINCTURE OF HENBANE.

R. Henbane leaves, one part.
Sulphuric ether, six parts.

Macerate for eight days, express, and filter. Dose, ten to twelve drops, three or four times a-day. *Hesse Ph.*

MIXTURE OF HENBANE AND SQUILL.

R. Tincture of squill, ten drops.
Nitric acid, five drops.
Extract of henbane, three grains.
Water, one fl. ounce and a half.

Mix. In spasmodic asthma; to be repeated every three or four hours. *Bree.*

LINIMENT OF HENBANE.

R. Extract of henbane, half a drachm.
White soap, two drachms.
Linseed oil, six ounces.

Mix. As an embrocation to glandular swellings. *Dupuytren.*

OINTMENT OF HENBANE.

R. Extract of henbane, one drachm.
Lard, one ounce.

Rub together. As an application to painful hemorrhoids. *Taddei.*

COMPOUND OINTMENT OF HENBANE.

R. Leaves of henbane,
" stramonium,
" bittersweet,
" elder, each, a handful.

Lard, two pounds.

Heat together till all moisture is evaporated, and strain. As an application to frost-bites, chilblains, &c. *Spielmann.*

R. Fresh poplar buds, four ounces.
Lard, twelve ounces.

Leaves of henbane,
" poppy,
" belladonna,
" black nightshade, each, one ounce.

Heat together till all moisture is evaporated. As a soothing and anodyne ointment to painful local affections. *Cottreau.*

This is much used in Europe, under the name of *Unguentum populeum*.

I.

ICHTHYOCOLLA.

ISINGLASS.

This is the swimming bladder of several kinds of fish. The best, however, is procured from various species of sturgeon, inhabiting the more southern parts of Russia. It is a pure and nutritious gelatine, and is rather employed as food, and in the arts, than in medicine, in which it is only used as the basis of court-plaster.

COURT-PLASTER.

R. Isinglass, one ounce.
Water, eight fl. ounces.

Macerate for twenty-four hours, and add
Alcohol, eight fl. ounces.

When perfectly dissolved, brush over strained silk, or sarsnet, several times, alternating with a layer of tincture of Peru or benzoin. *Cottreau.*

R. Isinglass, nine drachms.
Dissolve in water, and add hot filtered mixture of

Alcohol (0.921), twelve fl. ounces.
Tincture of benzoin, two fl. ounces.

Apply this to strained silk, till six coats are given, permitting each to dry before applying another, then brush over twice with a mixture of

Tincture of benzoin, six fl. ounces.
Liquid turpentine, four ounces. *Virey.*

R. Isinglass, one ounce.
Water, two fl. ounces.

Mix, and allow to stand till the isinglass is quite soft, then add

Rectified spirit,
three fl. ounces and a half;

mixed with

Water, one fl. ounce and a half.

Expose to heat of water-bath, till dissolved, and apply to oiled silk, nailed to a board, by means of a soft brush; apply four coats. *Liston.*

R. Fasten a piece of fine muslin, linen, or silk, to a flat board, and give it a thin coat of flour paste; over this, when dry, two coats of colorless gelatine are to be applied. Said to be superior to the usual court-plaster. *Deschamps.*

R. To one quart of jelly (obtained by boiling two beef's feet in sufficient water to cover them, and removing the fat from the surface), add one ox-gall; boil, and strain through a coarse cloth. Then add, while warm, two fl. ounces

of laudanum, and the same quantity of tincture of camphor. Stir well together, and apply the composition to silk stretched on frames, two or three times with a paint-brush, and carefully dry. This is far less apt to irritate the skin than common court-plaster.

Bullock.

DIAMOND CEMENT.

R. Isinglass, softened in water, and dissolved in the smallest quantity of diluted alcohol;
Of the solution, two ounces.
Ammoniac, ten grains;
and whilst still liquid, add
Solution of mastic, made with half a drachm, in three drachms of alcohol.
Stir well together. *Redwood.*

INDIGO.

INDIGO.

A tinctorial material, obtained from several species of *Indigofera*. It is found in commerce, in the form of small, solid, brittle masses, having scarcely any taste or odor, of a deep blue color, but assuming a coppery lustre when rubbed. It is principally used in the arts, but has also been employed as a remedial agent, in various spasmodic diseases. The dose is about a scruple, two or three times a-day, rapidly increased to a drachm or more.

POWDER OF INDIGO.

R. Powdered indigo, half an ounce.
Aromatic powder, half a drachm.
Mix. A teaspoonful, three times a-day, in epilepsy. *Radius.*

ELECTUARY OF INDIGO.

R. Powdered indigo, half an ounce.
Aromatic powder, half a drachm.
Syrup, one fl. ounce.
Mix. To be taken in divided doses, during the day. *Phæbus.*

PILLS OF INDIGO.

R. Powdered indigo, seventy-five grains.
Assafetida, fifteen grains.
Castor, seven grains.
Mix, and form twenty pills. One, every hour. *Podreca.*

SULPHATE OF INDIGO.

R. Indigo, Bengal, in powder, four ounces
Sulphuric acid, twenty-four ounces.
Dissolve by maceration and agitation, and add water until the solution measures four pints.

INULA.

ELECAMPANE.

The root of *Inula helenium*, a large herbaceous plant, a native of Europe, and cultivated in gardens in this country, where it has also become naturalized in some places.

Sex. Syst. Syngen. super. *Nat. Syst.* Astera-ceæ.

Linn. Sp. Pl. 881. Griffith, Med. Bot. 397.

The root is thick, fleshy, and branched. As dried and found in the shops, it is in slices of a yellowish-gray color, with a peculiar and aromatic odor, and an unpleasant, bitterish taste, with an after sensation of pungency. It is tonic and expectorant, and also acts in some cases as a diuretic and diaphoretic. The dose of the powder is from a scruple to a drachm.

DECOCTION OF ELECAMPANE.

R. Elecampane root, half an ounce.
Water, one pint.
Boil. Dose, a fl. ounce every hour or two. *Wood.*

COMPOUND PILLS OF ELECAMPANE.

R. Extract of elecampane, one drachm.
Powdered foxglove,
“ ipecacuanha, each, ten grains.
Opium, five grains.
Beat into mass, and form pills of two grains. One every hour, in chronic catarrh. *Heim.*

EXTRACT OF ELECAMPANE.

R. Elecampane root, at will.
Water, sufficient.
Make a concentrated infusion, strain, and evaporate to the proper consistence. Dose, from a scruple to a drachm. *Cottreau*

MIXTURE OF ELECAMPANE.

R. Extract of elecampane, two drachms.
Antimonial wine, one fl. ounce.
Syrup of orange flowers, half a fl. ounce
Mix. Dose, twenty to thirty drops, every three hours, in asthma and chronic catarrh. *Jahn.*

COMPOUND OXYMEL OF ELECAMPANE.

R. Elecampane root, one ounce.
Orris root, half an ounce.
Water, thirty-six fl. ounces.

Boil down to two-thirds, strain, and add

Honey, sixteen ounces.
Ammoniac, one ounce.
Vinegar, eight fl. ounces.

Evaporate to the consistence of honey. A spoonful occasionally, in catarrh and humid asthma. *Spielmann.*

IODINIUM.

IODINE.

Iodine is an elementary non-metallic body, having many of the properties of chlorine. It exists in many marine vegetables and animals, and also in some mineral bodies and salt springs; but is principally obtained from kelp, or the ashes of seaweeds. It is soft, friable, and opaque; of a bluish-black color and metallic lustre, and has a strong and penetrating odor and an acrid taste. It is employed as a stimulant or alterant in morbid actions of the absorbent and glandular systems, both externally and internally, and, in over doses, acts as an irritant poison.

The dose, in substance, is from the eighth of a grain to a grain.

PILLS OF IODINE.

R. Iodine, six grains.
Extract of gentian, one drachm.
Mucilage of gum Arabic, sufficient.

Mix, and form twenty-four pills. These are said to be useful in mercurial or scorbutic salivation. *Radius.*

TINCTURE OF IODINE.

R. Iodine, one ounce.
Alcohol, one pint.

Dissolve. *U. S. Ph.*

Sixteen minims, or thirty-five drops, contain one grain of iodine. Dose, from ten to twenty drops, three times a-day; to be increased.

SATURATED TINCTURE OF IODINE.

R. Iodine, two scruples.
Alcohol, one fl. ounce.
Compound spirit of lavender, two fl. drachms.

Mix. Five to ten drops, twice a-day, gradually increasing. In amenorrhœa, leucorrhœa, gonorrhœa, &c. *Durand.*

COMPOUND TINCTURE OF IODINE.

R. Iodine, half an ounce.
Iodide of potassium, one ounce.
Alcohol, one pint.

Mix, and form a tincture. Dose, five to fifteen drops, as may be required. *U. S. Ph.*

ETHEREAL TINCTURE OF IODINE.

R. Iodine, two scruples.
Sulphuric ether, one fl. ounce and a half.

Dissolve, and filter. *Magendie.*

R. Iodine, six grains.
Sulphuric ether, two fl. drachms.

Dissolve. Dose, ten drops, two or three times a-day, much diluted. In diseases of the skin or glands. *Ellis.*

R. Iodine, at will.
Sulphuric ether, sufficient

to make a saturated solution. This is very useful as a counter-irritant, and is applied by a camel-hair pencil. The part is then to be covered with a pledget of lint to check evaporation. *R. Thomson.*

MIXTURE OF IODINE AND IODIDE OF POTASSIUM.

R. Iodine, three grains.
Iodide of potassium, six grains.
Distilled water, one fl. ounce.

Dissolve. Dose, six to twenty drops, thrice a-day, in a draught of cold water. In glandular diseases and affections of the mucous membranes. *Morton.*

BATH WITH IODINE.

For adults. 1. 2. 3.

R. Iodide of potassium, 231 grs. 308 grs. 370 grs.
Iodine, 123 " 154 " 184 "
Water, twenty fl. ounces.

Dissolve, and add to bath of sixty gallons.

For children.

R. Iodide of potassium, 77 grs. 92 grs. 107 grs.
Iodine, 38 " 46 " 61 "
Water, ten fl. ounces.

Dissolve, and add to bath of sixty gallons. Used in scrofulous affections. *Lugol.*

SOLUTION OF IODINE.

For internal use. 1. 2. 3.

R. Iodine, $\frac{1}{2}$ gr. $\frac{2}{3}$ gr. 1 gr.
Commonsalt, 12 grs. 12 grs. 12 grs.
Distilled water, Oj. Oj. Oj.

For external use.

R. Iodine, 2 grs. 3 grs. 4 grs.
Distilled water, Oj. Oj. Oj.

As a lotion. *Lugol.*

LOTION OF IODINE.

- R. Tincture of iodine, half a fl. ounce.
Iodide of iron, twelve grains.
Chloride of antimony, half an ounce.

Mix. As an application to corns, by means of a camel's-hair pencil, after the corn has been pared. *Henderson.*

- R. Tincture of iodine, one fl. drachm.
Black drop, two fl. drachms.
Oil of almonds,
Lime water, each, two fl. ounces.
Mix carefully. As a lotion for chilblains. *Cadet.*

- R. Iodine, one or two grains.
Alcohol, one fl. drachm.
Water, one pint.

Mix. To wash scrofulous ulcers. *Foy.*

MIXTURE OF IODINE.

- R. Tincture of iodine, ninety drops.
Decoction of Peruvian bark, ten fl. ounces.

Mix. Two spoonfuls, three times a-day. *Radius.*

- R. Tincture of iodine, ten drops.
Water of soapwort, four fl. ounces.
Syrup of orange-flowers, one fl. ounce.

Mix. In spoonful doses. In the treatment of scrofulous ulcers. *Foy.*

CLYSTER OF IODINE.

- R. Gum Arabic, four drachms.
Water, six fl. ounces.

Dissolve, and add

- Tincture of iodine, five drops.

In amenorrhœa, dysmenorrhœa, &c. *Cadet.*

GARGLE OF IODINE.

- R. Tincture of iodine, one to two fl. drachms.
Tincture of opium, one fl. drachm.
Water, six fl. ounces.

Mix. As a gargle, in ulceration of the tonsils. *Ross.*

OINTMENT OF IODINE.

- R. Iodine, one scruple.
Iodide of potassium, four grains.
Water, six minims.
Lard, one ounce.

Rub the iodine and iodide with the water, and then with the lard, till thoroughly mixed.

U. S. Ph.

In goitre, scrofulous swellings, &c.

- R. Iodine, one scruple.
Rose ointment, one ounce.

Mix. To be applied to the tonsils, night and morning, by means of a brush. Not to be used till all inflammation is subdued. *Cerchiari.*

- R. Iodine, one part.
Lard, sixteen parts.

Rub together carefully. *Guibourt.*

COMPOUND OINTMENT OF IODINE.

- R. Iodine, half a drachm.
Iodide of potassium, one drachm.
Alcohol, one fl. drachm.
Lard, two ounces.

Rub the iodine and iodide of potassium with the alcohol, and then with the lard, till thoroughly mixed. *U. S. Ph.*

Used in same cases as the iodine ointment, but is more active.

IODINE OINTMENT WITH OIL OF TOBACCO.

- R. Iodine, twelve grains.
Iodide of potassium, four scruples.
Oil of tobacco, fifty drops.
Lard, two drachms.

Mix. To relax rigid muscles. *Dover.*

IODINE OINTMENT AND MERCURY.

- R. Mercurial ointment, sixteen parts.
Iodine, three parts.

Mix. In frictions, in ovarian dropsy. *Radius.*

LINIMENT OF IODINE.

- R. Liniment of opium, one fl. ounce.
Tincture of iodine, one fl. drachm.

Mix. *Manson.*

CATAPLASM OF IODINE.

- R. Tincture of iodine, half fl. ounce.
Flaxseed meal, one ounce.
Oatmeal, three ounces.
Distilled water, sufficient.

Make a cataplasm. Used as an application to scrofulous tumors and goitre.

Dunglison, N. Rem.

IODINE PLASTER.

- R. Iodine, one drachm.
Iodide of potassium, one scruple.
Water, half a fl. drachm.

Triturate in a mortar, and add to

- Burgundy pitch, three ounces.

Wax,

- Suet, each, half an ounce

Melt together, and triturate till well mixed.

IPECACUANHA.

IPECACUANHA.

The root of *Cephalis ipecacuanha*, a small herbaceous plant, a native of Brazil.

Sex. Syst. Pentand. monog. *Nat. Syst.* Rubiaceæ.

Richard, Bull. Med. iv. 92. Griffith, Med. Bot. 357.

This root is about the size of a large quill, of an irregular, twisted shape, apparently formed of small, unequal rings, about a line in height, separated by narrow interstices; when broken, it presents two distinct parts, a thin ligneous centre, and a thick cortical layer of a brownish color, and somewhat resinous appearance, having an herbaceous, sub-acrid, somewhat bitter taste, and a nauseous odor. Externally it presents some varieties of color, but is usually brown. It is emetic, and sometimes purgative in full doses, and diaphoretic and expectorant in small, but repeated quantities. As an emetic, the dose is fifteen to twenty grains; as a nauseant, two to four grains; as an expectorant, one to two grains.

POWDER OF IPECACUANHA AND OPIUM.
(DOVER'S POWDER.)

R. Ipecacuanha,
Opium, each, in powder, one drachm.
Sulphate of potassa, one ounce.
Triturate thoroughly. *U. S. Ph.*

R. Ipecacuanha,
Liquorice root,
Extract of opium, each,
in powder, one ounce.
Sulphate of potassa,
Nitrate of potassa, each, four ounces.

Triturate well. *Par. Cod.*

This latter formula approaches nearest to the original formula of Dover, but is not superior to the first, now generally used. Dose, five to ten grains, as a diaphoretic.

DOVER'S POWDER, (Original).

R. Opium, one ounce.
Saltpetre,
Vitriolated tartar, each, four ounces.
Ipecacuanha, one ounce.
Liquorice, one ounce.

"Put the saltpetre and tartar into a red-hot mortar, stirring them with a spoon until they have done flaming. Then powder them very fine. After that slice in your opium; grind these to a powder, and then mix the other powders with them." *Dover.*

POWDER OF IPECACUANHA AND TARTAR
EMETIC.

R. Powdered ipecacuanha, one scruple.
Tartar emetic, one grain.

Mix. As an emetic, or in divided doses, as a nauseant and sudorific. *Ellis.*

POWDER OF IPECACUANHA AND RHUBARB.

R. Powdered ipecacuanha,
" rhubarb, each, one scruple.

Mix. As an emeto-cathartic in full doses; or tonic alterative, in doses of three or four grains. *Beasley.*

POWDER OF IPECACUANHA AND MURIATE
OF AMMONIA.

R. Powdered ipecacuanha, a quarter grain.
Muriate of ammonia, twelve grains.
Extract of liquorice,
Sugar of milk, each, six grains.

Mix. To be taken every two hours, as an expectorant, on the decline of pneumonia. *Kopp.*

POWDER OF IPECACUANHA AND ANTI-
MONY.

R. Ipecacuanha,
Golden sulphuret of
antimony, each, one grain.
Camphor, three grains.
Gum Arabic, one scruple.

Triturate well. To be taken every two hours, in deficient expectoration in pneumonia. *Phæbus.*

POWDER OF IPECACUANHA AND CAR-
BONATE OF SODA.

R. Powdered ipecacuanha,
" opium, each, one grain.
Carbonate of soda, twelve grains.

Mix, and rub well together. To be taken every eight hours, in spasmodic asthma, whooping-cough, &c. *A. T. Thomson.*

POWDER OF IPECACUANHA AND NITRE.

R. Dover's powder, one drachm.
Nitrate, one ounce.

Mix. To be taken in divided doses. *Beasley.*

R. Powdered ipecacuanha, six grains.
" nitrate of potassa, one scruple and a half.
" myrrh, twelve grains.

Mix, and divide into four powders; one to be taken every four hours. Found useful in asthma, and the earlier stages of phthisis. *A. T. Thomson.*

POWDER OF IPECACUANHA AND CALO-
MEL.

R. Powdered ipecacuanha, six grains.
" opium, three grains.

Calomel, one grain and a half.
 Powdered nitre, half a drachm.
 Mix, and make six powders. One, every three
 or four hours, as a diaphoretic. *Ellis.*

POWDER OF IPECACUANHA AND MYRRH.

R. Powdered ipecacuanha, six grains.
 " myrrh, twelve grains.
 " nitre, half a drachm.

Mix, and make four powders. One, every four
 hours, as a stimulating expectorant. *Paris.*

POWDER OF IPECACUANHA AND TRAGACANTH.

R. Powdered ipecacuanha, five grains.
 Compound tragacanth
 powder, ten grains.

Mix, make a powder, to be taken every four or
 six hours. Useful at the commencement of
 febrile diseases, after clearing the stomach and
 bowels. *A. T. Thomson.*

EXTRACT OF IPECACUANHA.

R. Coarsely-powdered
 ipecacuanha, two parts.
 Proof spirit, seven parts.

Treat by displacement; distil off the spirit and
 evaporate, on water-bath, to proper consistence.
Paris Cod.

BOLUS OF IPECACUANHA.

R. Powdered ipecacuanha, two grains.
 Conserve of opium, one scruple.
 Prepared chalk, sufficient.

Mix, and make eight boluses. Two, night and
 morning, in dysentery. *St. Marie.*

R. Powdered ipecacuanha, two grains.
 Aromatic powder, ten grains.
 Syrup of cinnamon, sufficient.

Mix, and make two boluses. One, in the even-
 ing, in dyspepsia. *Cadet.*

PILLS OF IPECACUANHA AND OPIUM.

R. Dover's powder, three parts.
 Confection of roses, one part.

Mix, and make pills of four grains each.
Ed. Ph.

PILLS OF IPECACUANHA AND SQUILL.

R. Dover's powder, three drachms.
 Powdered squill,
 Ammoniac, each, one drachm.
 Molasses, sufficient.

Mix, and form mass. Dose, five to ten grains,
 as a diaphoretic. *Lond. Ph.*

PILLS OF IPECACUANHA AND FOXGLOVE.

R. Powdered ipecacuanha,
 " foxglove, each, half a
 drachm.

" opium, five grains.
 Extract of elecampane, sufficient.

Mix, and make sixty pills. One, every three
 hours, in spasmodic asthma. *Heim.*

PILLS OF IPECACUANHA AND CENTAURY.

R. Powdered ipecacuanha,
 Extract of centaury, each,
 fifteen grains.

Powdered rhubarb, half a drachm.
 Mix, and make thirty pills. One, night and
 morning, in dyspepsia. *Radius.*

LOZENGES OF IPECACUANHA.

R. Powdered ipecacuanha, half an ounce.
 " sugar, fourteen ounces.

Arrow root, four ounces.
 Mucilage of tragacanth, sufficient.

Mix the powder thoroughly, and form into a mass
 with the mucilage, and divide into lozenges of
 ten grains each. *U. S. Ph.*

A good expectorant in catarrh. Each con-
 tains a quarter of a grain of ipecacuanha.

LOZENGES OF IPECACUANHA AND CAMPHOR.

R. Powdered ipecacuanha, fifteen grains.
 " camphor,

" sugar, each, one drachm.
 Mucilage of tragacanth, sufficient.

Mix, and make sixty lozenges. *Beasley.*

R. Powdered sugar, five hundred parts.
 Flake manna, one hundred and
 twenty-five parts.

Powdered ipecacuanha, eighteen parts.
 Lactucarium, eight parts.
 Powdered squill, four parts.
 Mucilage of tragacanth, sufficient.

Make into a uniform paste, and divide into loz-
 enges of ten grains. Five or six a-day, in ca-
 tarrh, &c. *Grunn.*

CONFECTION OF IPECACUANHA.

R. Powdered ipecacuanha, twelve grains.
 Flowers of sulphur, one scruple.
 Powdered orris, one drachm.
 Syrup of mallow,
 Manna, each, two ounces

Mix. A teaspoonful, two or three times a-day,
 in hooping-cough. *Bories.*

DECOCTION OF IPECACUANHA.

- R. Bruised ipecacuanha, two drachms.
Water, twelve ounces.

One-third of the water is to be used at a time, so as to make three decoctions; these are to be united, and should amount to six ounces. Celebrated in dysentery; one-third to be taken at a dose.

Soubéiran.

MIXTURE OF IPECACUANHA.

- R. Powdered ipecacuanha, half a drachm.
Tartar emetic, one grain.
Tincture of squill, one fl. ounce.
Distilled water, seven fl. ounces.

Mix. Four tablespoonfuls; then two every ten minutes, till vomiting is produced.

Ellis.

- R. Powdered ipecacuanha, five grains.
Syrup of poppies, one drachm.
Spirit of cinnamon, two drachms.
Chalk mixture, thirteen drachms.

Mix. As a draught in dysentery.

Hooper.

- R. Powdered ipecacuanha, two grains.
Carbonate of soda, ten grains.
Syrup of poppies, one fl. ounce.
Mint water, six fl. ounces.

Mix. In spoonfuls, to check spasmodic vomiting.

St. Marie.

- R. Ipecacuanha, one drachm.
Senna, two drachms.
Boiling water, six fl. ounces.

Infuse for twelve hours, strain, and add

Oxymel of squill,
Syrup of hyssop, each, one ounce.

Mix. In spoonfuls, in hooping-cough.

Guibourt.

SYRUP OF IPECACUANHA.

- R. Ipecacuanha, in coarse powder, one ounce.
Diluted alcohol, one pint.
Sugar, two pounds and a half.
Water, sufficient.

Macerate the ipecacuanha in the alcohol for fourteen days, and filter. Evaporate to six fl. ounces, filter, and add water to make a pint. Lastly, add the sugar and proceed as for syrup.

U. S. Ph.

As emetic for children, one to two fl. drachms.
As expectorant, five to twenty minims.

- R. Bruised ipecacuanha, (troy), six ounces.
Alcohol, one pint and a half.
Water, one pint.

Mix to form a tincture. Digest for ten days, filter, and add one pint of water by way of displacement; evaporate in a water-bath to two pints; add immediately

Refined sugar, three pounds and a quarter (troy),
and boil. *Coggeshall.*

COMPOUND SYRUP OF IPECACUANHA.

- R. Ipecacuanha, one drachm.
Orris root, two drachms.
Red Peruvian bark, half an ounce.
Seneka,
Iceland moss, each, two ounces.
Sugar, two pounds.
Boiling water, two pints.

Digest for two hours, strain, and evaporate to one pint; then add the sugar, and make a syrup. In spoonful doses, as an expectorant.

Courtois.

- R. Syrup of ipecacuanha,
" poppies, each, two ounces.
" orange flowers, one ounce.

Oxymel of squill, one ounce and a half.

Mix. Two spoonfuls every hour, in hooping-cough.

Cadet.

TINCTURE OF IPECACUANHA.

- R. Bruised ipecacuanha, two ounces.
Spirit of nitric ether, two pints.

Digest for eight days, and filter. Used as a diaphoretic in febrile affections.

Mettauer

WINE OF IPECACUANHA.

- R. Bruised ipecacuanha, two ounces.
Sherry wine, two pints.

Macerate for fourteen days, express, and filter.

U. S. Ph

Emetic and diaphoretic. Dose, for an adult, as emetic, one fl. ounce; for a child, one fl. drachm.

COMPOUND WINE OF IPECACUANHA.

- R. Ipecacuanha, four parts.
Sugar, two parts.
Diluted alcohol, twenty-four parts.

Digest for eight days, and add

Sherry wine, ninety-six parts.
Star anise, one part.
Sugar, four parts.

Filter at the end of six days.

Niemann.

WINE OF IPECACUANHA AND TARTAR EMETIC

- R. Wine of ipecacuanha, one fl. ounce.
Tartar emetic, one grain.

Mix. A teaspoonful, every ten or fifteen minutes, till vomiting is induced.

Ellis.

ALKALINE WINE OF IPECACUANHA.

- R. Wine of ipecacuanha, forty minims.
Tincture of opium, eight drops.

Carbonate of soda, sixteen grains.
Syrup, two fl. drachms.
Water, fourteen fl. drachms.
Mix. A teaspoonful, to children, in hooping-cough
Pierson.

ELIXIR OF IPECACUANHA.

R. Powdered ipecacuanha,
Balsam of tolu, each, half an ounce.
Flowers of benzoin,
Opium,
Saffron, each, two drachms.
Oil of anise, one drachm.
Camphor, two scruples.
Alcohol, two pints.

Digest, express, and filter. Dose, one to two drachms, as a stimulating sudorific. *Cadet.*

IRIS FLORENTINA.

FLORENTINE IRIS. ORRIS.

The Florentine iris is a native of the southern parts of Europe, and affords a root, or rhizome, known as orris root.

Sex. Syst. Triand. monog. *Nat. Syst.* Iridaceæ.

Linn. Sp. Pl. 55. Griffith, Med. Bot. 625.

The rhizome is horizontal, knotty, and fragrant; in a recent state it is acrid, and somewhat bitter; this acrimony disappears in a great measure on desiccation. It is seldom used in medicine, but forms an ingredient in numerous tooth-powders, &c.

DENTIFRICE WITH ORRIS ROOT.

R. Cream of tartar, two ounces.
Powdered orris root, one ounce.
" myrrh,
" kino, each, half an ounce.

Mix well. *Redwood.*

R. Powdered Castile soap,
" orris root, each, two ounces.
" cuttlefish bone,
" prepared chalk, each, three ounces.

Oil of cloves,
Essence of lemon, each, twenty drops.

Mix. *Redwood.*

J.

JALAPA.

JALAP.

Jalap is the root of the *Ipomœa jalapa*, a native of Mexico.

Sex. Syst. Pentand. monog. *Nat. Syst.* Convolvulaceæ.

Coxe, Am. Journ. Med. Sci., 1830. Griffith, Med. Bot. 474.

The root, when dried, is in pieces of various forms and sizes; usually entire, but sometimes in slices. The epidermis is thin and brown; within it is grayish, with brown, concentric rings. The powder is of a brownish-yellow, with a peculiar, unpleasant odor, and a sweetish, sub-acrid, nauseous taste. It is an active purgative, in doses of ten to thirty grains.

POWDER OF JALAP AND CALOMEL.

R. Powdered jalap, ten grains.
Calomel, five grains.
Mix. To be given in syrup or molasses. As a purgative. *Ellis.*

R. Powdered jalap, twelve grains.
Calomel, three grains.
Sulphate of potassa, seven grains.

Mix; make a powder, to be taken at night; a useful purge in diminishing the action of the liver. *A. T. Thomson.*

POWDER OF JALAP AND CREAM OF TARTAR.

R. Powdered jalap, one drachm.
Cream of tartar, six drachms.
Mix, and divide into six powders. One every three hours, as a hydragogue purgative. *Ellis.*

R. Jalap, two ounces.
Cream of tartar, one ounce.
Sugar, thirteen ounces.
Oil of orange peel, two drachms.

Mix. Dose, one to two drachms. *Cadet.*

R. Powdered jalap, one ounce.
Cream of tartar, two ounces.

Mix. *U. S. Ph.*
Dose, thirty grains to a drachm.

POWDER OF JALAP AND IPECACUANHA.

R. Jalap, fifteen grains.
Ipecacuanha, five grains.
Oil of cinnamon, two drops.

Mix. *Paris.*

POWDER OF JALAP AND SCAMMONY.

R. Jalap, one drachm

Scammony, one scruple.
 Ipecacuanha, ten drachms.
 Mix, and divide into powders of eight grains.
 In mucous discharges. *Dumas.*

COMPOUND POWDER OF JALAP.

R. Powdered jalap, three ounces.
 Cream of tartar, six ounces.
 Powdered ginger, two drachms.

Mix. Half a drachm to one drachm, as a hydragogue purgative. *Lond. Ph.*

EXTRACT OF JALAP.

R. Jalap, in coarse powder, one pound.
 Alcohol, four pints.
 Water, sufficient.

Macerate the jalap in the alcohol for four days, then exhaust by the process of displacement; first by alcohol, then by water. Distil off the alcohol from the tincture, and evaporate the infusion, till both are of the consistence of thin honey, mix them, and evaporate to proper consistence. *U. S. Ph.*

Dose, from ten to twenty grains.

RESIN OF JALAP.

R. Coarsely-powdered jalap, at will.

Moisten thoroughly with rectified spirit, put in a displacement apparatus and exhaust; distil off the spirit, and evaporate to due consistence. *Ed. Ph.*

Dose, four to twelve grains.

ALKALINE EXTRACT OF JALAP.

R. Powdered jalap, two pounds.
 Alcohol (.921), one gallon.

Digest for some days, strain, and filter; evaporate slowly on the water-bath to a pilular consistence, adding gradually,

Carbonate of potassa, twelve drachms.

Dose, three to nine grains. *Durand.*

COMPOUND PILLS OF JALAP.

R. Jalap,
 Rhubarb,
 Aloes, each, one ounce.
 Soap, two ounces.
 Syrup of ginger, sufficient.

Mix, and form mass. Dose, twenty to thirty grains. *Foy.*

PILLS OF JALAP AND CALOMEL.

R. Resin of jalap,
 Soap, each, one drachm.
 Oil of orange peel, six drops.
 Mix, and make sixty pills. *Alibert.*

ELECTUARY OF JALAP.

R. Powdered jalap,
 Black sulphuret of mercury, each, two drachms.
 Scammony, one drachm.
 Resin of jalap,
 Squill, each, one drachm and a half.
 Syrup of buckthorn, sufficient.

Mix. Dose, half a drachm to a drachm, in dropsy or lead colic. *Cadet.*

R. Powdered jalap, two drachms.
 Cream of tartar, one ounce and a half.
 Ginger, one scruple.
 Molasses, two ounces and a half.

Mix. Dose, two drachms. *Beasley.*

R. Powdered jalap,
 Cream of tartar,
 Nitre, each, half an ounce.
 Confection of senna, one ounce.
 Simple syrup, sufficient.

Mix. A bolus the size of a hazlenut, three or four times a-day, in hemorrhoids. *Ellis.*

EMULSION OF JALAP.

R. Resin of jalap, nine grains.
 Soap, three grains.
 Olive oil, one ounce and a half.

Triturate well. Said to be useful in dysentery. A tablespoonful, occasionally. *Vogel.*

MIXTURE OF JALAP.

R. Infusion of senna, six fl. ounces.
 Extract of jalap, half a drachm.
 Tartrate of potassa, four drachms.
 Oil of ginger, twenty minims.

Mix. *Beasley.*

SOAP OF JALAP.

R. Resin of jalap,
 Soap, equal parts.

Dissolve in Alcohol, sufficient.

Evaporate to pilular consistence. Dose, ten to fifteen grains. *Pruss. Ph.*

TINCTURE OF JALAP.

R. Powdered jalap, six ounces.
 Diluted alcohol, two pints.

Macerate for fourteen days, express, and filter. *U. S. Ph.*

Seldom given alone, but added to cathartic mixtures, in portions of one to two fl. drachms.

COMPOUND TINCTURE OF JALAP.

R. Jalap, six drachms.
 Black hellebore, three drachms.

Juniper berries, half an ounce.
 Rasped guaiacum, half an ounce.
 Brandy, twenty-four fl. ounces.
 Digest for three days, express, and filter.

Ed. Ph., 1744.

R. Jalap, eight ounces.
 Scammony, two ounces.
 Turbeth, one ounce.
 Brandy, twelve pints.

Digest for twelve hours, on a water-bath, filter, and add

Syrup of senna, four pounds.
 Mix. *Giordano*.

MIXTURE OF TINCTURE OF JALAP.

R. Tincture of jalap, two fl. drachms.
 Vinegar of squill, one fl. drachm.
 Mint water, one fl. ounce.

Mix. To be taken three times a-day. *Ellis*.

JUGLANS CINEREA.

BUTTERNUT.

The Butternut, or White Walnut, is an indigenous tree, with numerous, nearly horizontal, branches, found in many places in the United States.

Sex. Syst. Monœc. polyand. *Nat. Syst.* Juglandaceæ.

Linn. Sp. Pl. 1415. *Griffith, Med. Bot.* 589.

The part used is the inner bark, especially of the root; this is fibrous, has little odor, but a bitter, somewhat acrid taste. It is, at first, white, but gradually changes to a dark brown. It is a mild cathartic, and is well suited to cases of costiveness and bowel affections. It is usually given in extract.

EXTRACT OF BUTTERNUT.

R. Butternut, in coarse powder, one pound.
 Water, sufficient.

Mix the butternut with a pint of the water, macerate for twenty-four hours, introduce into a displacement apparatus, and exhaust. Heat the fluid to a boiling point, strain, and evaporate to proper consistence. *U. S. Ph.*

Dose, from five to ten grains, as a laxative; from twenty to thirty, as a purgative.

JUNIPERUS.

JUNIPER.

The common Juniper, *J. communis*, is an erect, evergreen shrub, sometimes attaining a large size, found in Europe, and Northern Asia and America. The tops and the berries are officinal.

Sex. Syst. Dicc. monadelph. *Nat. Syst.* Pinaceæ.

Linn. Sp. Pl. 1470. *Griffith, Med. Bot.* 608.

The berries are oblong globular, marked with three radiating furrows at top, and contain three seeds. They are of a purplish-black color, have a sweetish, somewhat terebinthinate taste, and a peculiar, aromatic odor. They are used as stimulating diuretics, especially as adjuvants to more active remedies. The tops have a balsamic odor, and a resinous, bitterish taste, with the same medicinal properties as the berries.

EXTRACT OF JUNIPER.

R. Bruised juniper berries, one pound.
 Distilled water, three pints.

Macerate at 77° to 86° F. for twenty-four hours; strain with gentle pressure, add more water to the berries, and let stand for twelve hours. Filter, and evaporate to proper consistence. Dose, one to three drachms. *Par. Cod.*

INFUSION OF JUNIPER.

R. Juniper berries, bruised, one ounce.
 Boiling water, one pint.

Infuse, and strain. To be taken during the twenty-four hours. *Radius*.

COMPOUND INFUSION OF JUNIPER.

(CIDER MIXTURE.)

R. Juniper berries, bruised,
 Mustard seed,
 Ginger, each, half an ounce.
 Horseradish, bruised,
 Parsley root, bruised, each,
 one ounce.
 Cider, two pints.

Infuse, and strain. A wineglassful, two or three times a-day, in general dropsy. *Parrish*.

R. Juniper berries, bruised, one ounce and a half.
 Boiling water, one pint.

Infuse in a close vessel for two hours, strain, and add

Cream of tartar, one ounce.

Mix. A teacupful, several times a-day, with forty drops of sweet spirit of nitre. In dropsy. *Ellis*.

R. Juniper berries, bruised, two ounces and a half.
 Boiling water, one pint.

Infuse for two hours, and strain; add

Compound spirit of juniper, ten fl. drachms.

Dose, two fl. ounces, three times a-day. *Beasley*.

R. Juniper berries, bruised, four ounces.
 Boiling water, twelve fl. ounces.
 Holland gin, four fl. ounces

Mix, and macerate for twelve hours; then throw on a cotton cloth, and express. Then let sufficient water be added to the dregs, and pressed out to make a pint. The advantages of this preparation are greater strength, and permanency.

W. Procter, Jr.

ELECTUARY OF JUNIPER.

R. Extract of juniper,
Clarified honey, each, one pound.
Juniper berries,
Cinnamon,
Orange peel, each, half an ounce.
Mix. Stomachic and diuretic. Dose, one to four drachms.

Saxon Ph.

MIXTURE WITH OIL OF JUNIPER.

R. Oil of juniper, half a drachm.
Nitric ether,
Tincture of fox-glove, each, three drachms.
Mix. Twenty to thirty drops, every three hours.

Hufeland.

R. Oil of juniper, twenty-five drops.
Sugar,
Gum Arabic, each, two drachms.
Distilled water, four fl. ounces.

Mix. A tablespoonful, every hour or two, as a diuretic.

Ellis.

COMPOUND SPIRIT OF JUNIPER.

R. Juniper berries, bruised, one pound.
Caraway, bruised,
Fennel, bruised, each, one ounce and a half.
Diluted alcohol, one gallon.
Water, two pints.

Macerate in the alcohol, for twenty-four hours, add the water, and with a slow fire, distil one gallon.

U. S. Ph., 1840.

R. Oil of juniper, one fl. drachm. and a half.
" caraway,
" fennel, each, ten minims.
Diluted alcohol, one gallon.

Dissolve the oils in the alcohol.

U. S. Ph., 1850.

Dose, two to four fl. drachms, as an addition to diuretic mixtures, &c., in debilitated cases of dropsy.

LINIMENT OF JUNIPER.

R. Oil of juniper, one ounce and a half.
Oil of anise, six drops.
Lard, two ounces.

Mix well; as an application in tinea capitis.

Sully.

R. Spirit of juniper, two ounces.
Oil of cloves,
Concrete oil of nutmeg, each, half a drachm.

Mix. As a friction in ecchymosis, and indolent tumors.

Augustin.

JUNIPERUS VIRGINIANA.

RED CEDAR.

The tops of the red cedar are officinal. This is an evergreen tree, sometimes attaining a great size, with small, fleshy, closely imbricated leaves. It is found in all parts of the United States, but is most abundant and vigorous in the south.

Sex. Syst. Diœc. monadelph. Nat. Syst. Pinaceæ.

Linn. Sp. Pl. 1471. Griffith, Med. Bot. 609.

The leaves or tops have a peculiar and somewhat aromatic odor, a bitterish and somewhat acrid taste. They have much the same properties as those of savine, but are not as active, though often used as a substitute for them.

CERATE OF RED CEDAR.

R. Powdered red cedar leaves, one ounce.
Resin cerate, six ounces.

Mix the powder with the cerate previously softened. Used as a substitute for savine cerate, but is much less effectual.

Wood.

K.

KINO.

KINO.

This is an extract obtained from several different plants, but that most used and esteemed is the product of *Pterocarpus marsupium*, a large tree, indigenous to some parts of India.

It is usually in small, angular fragments, of a

dark, brownish-black color, opaque and shining; of a brittle and friable consistence, an astringent and slightly bitter taste; adhering to the teeth when chewed, and tinging the saliva of a red color. It is a powerful astringent, and much used in the treatment of mucous discharges, &c., and as a local application in gonorrhœa, to indolent ulcers, relaxed uvula, &c. The dose in powder is from ten to thirty grains.

COMPOUND POWDER OF KINO.

R. Powdered kino, fifteen drachms.
 " opium, one drachm.
 " cinnamon, four drachms.
 Rub well together. Dose, five to twenty grains.
Lond. Ph.

As an anodyne astringent, in diarrhœa and hemorrhage of the bowels.

R. Powdered kino,
 Sugar of milk, each, one scruple.
 Mix. To be taken every three or four hours, in cardialgia.
Radius.

COMPOUND PILLS OF KINO.

R. Powdered kino, one scruple.
 " opium, two grains.
 Mucilage of gum Arabic, sufficient.
 Mix, and make four pills. One, every four hours, in diarrhœa, pyrosis, &c.
Ellis.

R. Powdered kino, two scruples.
 " camphor, thirty grains.
 Aromatic powder, one scruple.
 Oxide of zinc, ten grains.
 Mucilage of gum Arabic, sufficient.
 Mix, and make twenty pills. Two, morning and evening, in diabetes.
Augustin.

ELECTUARY OF KINO.

R. Powdered kino, six drachms.
 " alum,
 " cinnamon, each, two drachms.
 Syrup, sufficient.

Mix into an electuary. One drachm, two or three times a-day, in diarrhœa, chronic dysentery, &c.
Swediaur.

INFUSION OF KINO.

R. Kino, two drachms.
 Boiling water, eight fl. ounces.
 Infuse, and strain when cool. Dose, a fl. ounce.
Wood.

INJECTION OF KINO.

R. Kino, one to two drachms.
 Alum, twenty to thirty grains.
 Boiling water, two pints.
 Infuse for an hour, and strain. As an injection in chronic urethritis.
Swediaur.

SYRUP OF KINO.

R. Kino, two ounces.
 Tepid water, three pints.
 Dissolve, and add
 Sugar, four pounds.
 Clarify, and evaporate to consistence of syrup.
Taddei.

TINCTURE OF KINO.

R. Powdered kino, three ounces and a half.
 Rectified spirit, two pints.
 Macerate for fourteen days, and filter.
Lond. Ph.

R. Powdered kino, six drachms.
 Diluted alcohol, sufficient.
 Mix the kino with an equal bulk of sand, and pour diluted alcohol on it, in a percolator, until eight fl. ounces of tincture are obtained.
U. S. Ph.

Dose, one to two fl. drachms, generally as an adjuvant to astringent mixtures.

KRAMERIA.

RHATANY.

The roots of several species of *Krameria* are possessed of analogous properties, but the only one that is officinal, is that of *K. triandra*, a small shrub found in Peru, in dry, sandy places. The stem is procumbent, and much branched, with sessile, hairy leaves, and dark red flowers. The root is long and branching.

Sex. Syst. Tetrand. monog. *Nat. Syst.* Polygalaceæ.

Ruiz & Pavon, Fl. Per. i. 93. Griffith, Med. Bot. 227.

The dried root is of a dark, reddish-brown color, with no odor, but a bitter and astringent taste. The small roots are the most powerful. It is somewhat tonic, and powerfully astringent. It is used for most purposes to which the vegetable astringents are applied. The dose in powder is from twenty to thirty grains.

COMPOUND POWDER OF RHATANY.

R. Extract of rhatany,
 Catechu, each, ten grains.
 Alum, four grains.
 Powdered liquorice, ten grains.
 Mix. To be taken three or four times a-day, in diarrhœa, &c.
Vogt.

DENTIFRICE OF RHATANY.

R. Extract of rhatany, one to two drachms.
 Myrrh, one drachm.
 Powdered orris, six drachms.
 Oil of cinnamon, two drops.
 " myrrh, ten drops.
 Balsam of Peru, ten grains.
 Mix well.
Phæbus.

EXTRACT OF RHATANY.

R. Prepared like extract of gentian, by percolation with cold water.
U. S. Ph.
 Dose, ten to twenty grains.

PILLS OF RHATANY.

- R. Extract of rhatany, one scruple.
 Powdered kino, sufficient.
 Mix, and make eight pills. One, every hour or two, to check mucous discharges. *Ellis.*
 R. Extract of rhatany, one drachm.
 " rhubarb, two scruples.
 Ginger, one scruple.
 Mix, and make twenty-four pills. *Reece.*

ELECTUARY OF RHATANY.

- R. Powdered rhatany, one ounce.
 " ginger, one drachm.
 Syrup of orange peel, sufficient.
 Mix. A teaspoonful every two or three hours. *Ferrara Ph.*

INFUSION OF RHATANY.

- R. Bruised rhatany, one ounce.
 Boiling water, one pint.
 Infuse for four hours, and strain. *U. S. Ph.*
 Dose, one to two fl. ounces.

COLLUTORY OF RHATANY.

- R. Extract of rhatany, half a drachm.
 Mucilage of quince seed, two drachms and a half.
 Sage water, one ounce and a half.
 Mix. As a mouth-wash in aphthæ, &c. *Radius.*

CLYSTER OF RHATANY.

- R. Extract of rhatany, one scruple.
 Mucilage of quince seed, three drachms.
 Infusion of sage, one fl. ounce and a half.
 Mix. To check diarrhœa, &c. *Ammon.*

INJECTION OF RHATANY.

- R. Extract of rhatany, half an ounce.
 Tincture of catechu, " kino, each, one drachm and a half.

To be added to a decoction of

- Rhatany, ten drachms.
 Water, three pints,
 reduced to two-thirds, and strained. As an injection in leucorrhœa, and prolapsus of the uterus. *Cadet.*

MIXTURE OF RHATANY.

- R. Extract of rhatany, one drachm.
 Syrup of poppies, " Rose water, each, two fl. ounces.
 Mix. A teaspoonful occasionally, in passive hemorrhage and chronic dysentery. *Fouquier.*
 R. Extract of rhatany, one drachm.
 Chamomile water, six fl. ounces.
 Elixir of vitriol, half a drachm.
 Mix. As the last. *Hildebrand.*

SYRUP OF RHATANY.

- R. Extract of rhatany, two ounces.
 Water, one pint.
 Sugar, two pounds and a half.
 Dissolve the extract in the water and filter; add the sugar, and form syrup. *U. S. Ph.*
 Dose, half a fl. ounce.

TINCTURE OF RHATANY.

- R. Powdered rhatany, six ounces
 Diluted alcohol, two pints.
 Macerate for fourteen days, express, and filter. *U. S. Ph.*
 Dose, one to two fl. drachms.

COMPOUND OINTMENT OF RHATANY.

- R. Burgundy pitch, eight parts.
 Venice turpentine, two parts.
 White wax, one part.
 Extract of rhatany, two parts.
 Alum, one part.

Melt the first three ingredients over a gentle fire, and add the rhatany and alum, in fine powder, and incorporate well.

Useful where an astringent application is indicated.

I.

L A C.

MILK.

The milk of several animals has been used medicinally, but that generally employed is from the cow. It is mostly employed in the form of whey, and of sugar of milk.

SIMPLE WHEY.

- R. Milk, three pints.
 Prepared rennet, sufficient.
 Cover up the mixture, and place before the fire, till a uniform curd is formed; divide this with a spoon, and strain. A good diluent in febrile

affections; when boiled down to one-half, it is nutritive.

A. T. Thomson.

VINEGAR WHEY.

R. Milk, one pint.
Vinegar, half fl. ounce.

Boil a few minutes, and separate the curd.

Ellis.

TAMARIND WHEY.

R. Boiling milk, one pint.
Tamarinds, two ounces.

Boil till coagulation is effected, and strain.

Plenck.

CREAM OF TARTAR WHEY.

R. Boiling milk, one quart.
Cream of Tartar, half an ounce.

Boil until the curd separates, and strain.

Plenck.

ALUM WHEY.

R. Clear whey, one pint.
Alum, two drachms.

Dissolve. A glassful occasionally, in passive hemorrhages.

Foy.

AROMATIC ALUM WHEY.

R. Powdered alum, one drachm.
Sugar of milk, half a drachm.
Powdered cinnamon, fifteen grains.
Milk, two pints.

Boil, and strain when coagulated. In passive uterine hemorrhages. To be taken during the day.

Niemann.

MUSTARD WHEY.

R. Bruised mustard seed, half an ounce.
Milk, one pint.

Boil till the milk is coagulated, and strain. A cupful, as a diuretic, in dropsical affections.

A. T. Thomson.

WINE WHEY.

R. Boiling milk, half a pint.
White wine, one to two fl. ounces.
Sugar, sufficient.

Boil for ten minutes, constantly stirring, then strain. A good mode of giving wine as a stimulant. Dose must depend on circumstances.

Ellis.

POWDER OF MILK.

R. Milk, two pints.
Powdered carbonate of soda, half a drachm.
Water, one fl. ounce.
Sugar, one pound.

Dissolve the soda in the water, add the solution to the milk, gently heat, and stir. When evaporated to one-third, add the sugar gradually, and stir. Remove from fire, pour into plates, and dry in an oven. When dry, powder, and keep in well-stopped bottles. One to two ounces, mixed with a quart of water, forms an agreeable drink, and is a good substitute for milk.

Legrip.

ARTIFICIAL GOATS' MILK

R. Fresh suet, one ounce.
Cut in small pieces, and tie in a muslin bag, and boil in

Milk, two pints.
Sugar candy, two drachms.

A good diet in scrofulous emaciation, and in latter stages of phthisis.

A. T. Thomson.

MIXTURE OF MILK AND SUET.

R. Sheeps' suet, two ounces.
Milk, one pint.
Starch, half an ounce.

Boil slowly for half an hour. As a drink, in dysentery and cholera infantum.

Ellis.

MILK AND SODA-WATER.

R. Milk, a teacupful.

Heat nearly to boiling, and add

Sugar, a teaspoonful.

Put in a large tumbler, and add

Soda-water, sufficient.

A good mode of taking milk when there is much acidity of stomach.

A. T. Thomson.

SUGAR OF MILK.

R. Whey, at will.

Clarify with white of egg, and evaporate on water-bath, to crystallization, and set aside in a cool place.

Redwood.

Cooling and laxative, in doses of two drachms to one ounce.

SYRUP OF MILK.

R. Skimmed milk, twelve pounds.

Evaporate to one-half, and add

Sugar, nine pounds.

Cherry-laurel water, three ounces.

Guibourt.

MIXTURE OF SUGAR OF MILK AND ICELAND MOSS.

R. Iceland moss, four drachms.

Water, one pint and a half.

Reduce to one-half by boiling, strain, and add

Sugar of milk, two drachms.

Milk, one pint.
Syrup of asparagus, three ounces.
A cupful, occasionally, in chronic catarrh.
Cadet.

MIXTURE OF SUGAR OF MILK AND GUM ARABIC.

R. Sugar of milk,
Sugar, each, two pounds.
Gum Arabic, three pounds.
Extract of dog-grass, seven ounces.
Mix. Half an ounce dissolved in a quart of water, forms a refreshing and demulcent drink.
Cadet.

SUGAR OF MILK DRAUGHT.

R. Sugar of milk, half an ounce.
Boiling water, two pounds.
Lemon juice, sufficient.
Mix. Recommended in cholera, to appease the thirst.
Ammon.

CHINA CEMENT.

R. Curd of milk, dried and powdered, ten ounces.
Quicklime, powdered, one ounce.
Camphor, two drachms.

Mix, and keep in closely-stopped bottles. When used, a portion is to be mixed with a little water into a paste, to be applied quickly. Used to cement broken glass, china, &c.
Cooley.

LACCA.

LAC

Is a resinous substance obtained from several kinds of East Indian plants, but particularly from the *Croton lacciferum*. Many varieties are known in commerce, but the principal are seed, shell, and stick lac, of which the purest is the shell. This is in thin fragments of various shades of yellow, or reddish color; brittle, inodorous, and insipid. It is somewhat astringent, and was formerly used in medicine, but is now mainly employed in the arts, for the manufacture of varnishes, sealing wax, &c.

WHITE OR BLEACHED LAC.

R. Shell lac, at will.
Boil in solution of potassa, pass chlorine gas through the solution, and malaxate in hot water.
Redwood.

AQUEOUS SOLUTION OF LAC.

R. Shell lac, five ounces.
Borax, one ounce.
Water, one pint.

Boil together, and strain. Used as a varnish or a vehicle for colors. Mixed with lampblack, it forms an ink that will resist the action of acids.
Cooley.

CEMENT OF SHELL LAC.

R. Shell lac, at will.
Alcohol, sufficient
to make a solution of the consistence of molasses. Exceedingly tenacious.
Cooley.

RED SEALING WAX.

R. Shell lac, two pounds.
Venice turpentine, one pound.
Vermilion, or best dichromate of lead, one pound and a half.
Melt the lac and turpentine together, and stir in the vermillion.
Gray.

The other colored waxes are made in the same manner, using the appropriate pigment.

LACTUCA ELONGATA.

WILD LETTUCE.

The Wild Lettuce is an indigenous, herbaceous perennial, from three to six feet high, with pale yellow flowers, growing in woods and fertile soils.

Sex. Syst. Syngen. æqual. *Nat. Syst.* Astera-ceæ.

Willd. Sp. Pl. iii. 1525. Big. Seq. 241.

Considered to resemble the *Lactuca virosa* in medical properties. An extract made from it may be given in doses of five to fifteen grains.

LACTUCA.

LETTUCE.

This well known plant is commonly cultivated for the table as a salad. It is erect and leafy, the leaves being ovate, semi-amplexicaul, and entire or toothed. The whole abounding in a bitter, milky juice.

It is anodyne and sedative, but is seldom used in this country except in the form of the extract, or lactucarium; but is employed to fulfil many indications in Europe.

MIXTURE OF LETTUCE AND DANDELION.

R. Fresh leaves of lettuce,
" root of dandelion,
" " chicory, each,
three ounces.

Beat the whole into a pulp, adding, gradually, Peppermint water, three fl. ounces.

Express, and filter. To be taken, during the day, in hypochondriasis, with insomnia.
Vogt.

EXTRACT OF LETTUCE.

(THRIDACE.)

R. Juice of the stalks of lettuce, at will.
Evaporate in the stove by a gentle heat, on plates. *Par. Cod.*

SYRUP OF THRIDACE.

R. Thridace, two drachms.
Water, two fl. ounces.
Add to the solution
Boiling syrup, one pound.
Evaporate, and strain. *Par. Cod.*

LACTUCARIUM.

R. Collect the milky juice that flows from incisions made in the stalks of lettuce, and allow it to harden.
Dose, five to eight grains.

PILLS OF LACTUCARIUM.

R. Lactucarium, twelve grains.
Make six pills; one every two hours, till sleep is procured. *Ellis.*
R. Lactucarium, twelve grains.
Conserve of elder berries,
Extract of liquorice, each, sufficient.
Mix, and make four pills. One, every three hours, in obstinate coughs, without expectoration. *Brera.*

MIXTURE OF LACTUCARIUM.

R. Lactucarium, thirty grains.
Decoction of Iceland moss, two ounces.
Mucilage, half an ounce.
Syrup, one ounce.
Mix. Two spoonfuls every two hours, in spasmodic cough, insomnia, hysteria, &c. *Brera.*

SYRUP OF LACTUCARIUM.

R. Lactucarium, seventeen grains.
Distilled water, three hundred grains.
Simple syrup, half a pint.
Extract the lactucarium, with the water at the boiling temperature, add the solution to the syrup, and evaporate it to half a pint. *Monchon.*
R. Powdered lactucarium, sixty-four grains.
Carbonate of potassa, thirty-two grains.
Sugar, four ounces.
Water, sufficient.
Grind the lactucarium with the carbonate of potash, till well mixed; add enough water to

moisten completely; stand aside for twelve hours, and then slowly obtain two fl. ounces by percolation; add the sugar, and dissolve with a gentle heat. Each fl. drachm contains two grains of lactucarium. *T. S. Wiegand.*

TINCTURE OF LACTUCARIUM.

R. Powdered lactucarium, four ounces.
Proof spirit, two pints.
Act by displacement or digestion. *Ed. Ph.*
Dose, thirty minims to two fl. drachms.

LOZENGES OF LACTUCARIUM.

R. Powdered lactucarium, two drachms.
" sugar, six ounces.
" gum Arabic,
" liquorice, each, five ounces.
Tincture of tolu, half an ounce.
Mix, and make lozenges of ten grains. *Ed. Ph.*
As expectorant, in catarrhal affections.

WATERY EXTRACT OF LETTUCE.

R. Fresh lettuce leaves, one pound.
Bruise them in a stone mortar, adding a little water, express the juice, and evaporate, without straining, to a proper consistence. *Lond. Ph.*
Dose, five to fifteen grains.
R. Stalks and leaves of lettuce, when old and yellow, at will; macerate in water for twenty-four hours, boil for two hours, drain through a sieve, and evaporate in shallow vessels by exposure.
Dose, three to four grains. *Probat.*

POWDER OF EXTRACT OF LETTUCE.

R. Extract of lettuce, two grains.
Powdered foxglove, half a grain.
" sugar, twelve grains.
Mix. To be taken every two hours, in hydrothorax. *Hufeland.*

MIXTURE OF EXTRACT OF LETTUCE.

R. Extract of lettuce, two scruples.
Tincture of foxglove, half an ounce.
Cinnamon water, one drachm and a half.
Mix. Thirty drops to one fl. drachm, every two hours. *Radius.*

AROMATIC TINCTURE OF LETTUCE.

R. Leaves of lettuce, one ounce.
Cinnamon, one drachm.
Alcohol,
Water, each, four fl. ounces.
Mix. Macerate for a week, express, and filter. Fifteen to thirty drops, in dropsy. *Niemann.*

WATER OF LETTUCE.

R. Fresh lettuce, ten parts.
 Water, twenty parts.
 Distil off ten parts. *Par. Cod.*
 Dose, half a fl. ounce to two fl. ounces.

SYRUP OF LETTUCE.

R. Water of lettuce, one pint.
 Sugar, two pounds.
 Make syrup. *Par. Cod.*

MIXTURE OF WATER OF LETTUCE.

R. Water of lettuce,
 " linden, each, two fl. ounces.
 " orange flowers, two fl. drachms.
 Syrup of poppies, one fl. ounce.
 Mix. A spoonful every two hours. *Foy.*

LAURUS.

LAUREL. BAY TREE.

The berries (*Lauri bacca*) and leaves (*Lauri folia*) of the *Laurus nobilis*, or bay tree, are used in medicine. This laurel is a small, evergreen tree, indigenous to the countries bordering on the Mediterranean.

Sex. Syst. Enneand. monog. *Nat. Syst.* Lauracæ.

Linn. Sp. Pl. 529. Griffith, Med. Bot. 550.

The leaves are fragrant, and have an aromatic, bitterish taste. The berries are small, oval, of a deep blue, when fresh, of a brownish-black, when dried, aromatic, and with an acrid, bitterish taste; they contain a concrete, greenish oil. Neither the leaves nor berries, nor their products, are now much used in medicine; they are stimulant and narcotic.

OIL OF LAUREL.

R. Fresh laurel berries, at will.
 Crush, warm the pulp for a few moments by a gentle heat, and subject to pressure. *Guibourt.*
 Used as a friction, in swelling and pain in the joints, colic, &c.

LAUREL OINTMENT.

R. Suet, eight ounces.
 Melt, and add
 Laurel oil, one pound.
 Essence of turpentine, one ounce and a half.
Port. Ph.

There are a variety of formulas for this ointment, which is much used under the name of *Nervine ointment*, or *balsam*.

R. Fresh laurel leaves,
 " berries, each, one pound.
 Lard, two pounds.
 Digest the bruised leaves and berries with the lard, till all moisture is driven off, and express. *Par. Cod.*

Both these are used as stimulating frictions.

LAURO-CERASUS.

CHERRY-LAUREL.

This is the officinal appellation of the leaves of the *Prunus lauro-cerasus*, a small evergreen tree, a native of Asia Minor, but cultivated in the temperate parts of Europe, both for ornament, and for medicinal purposes.

Sex. Syst. Icosand. monog. *Nat. Syst.* Drupacæ.

Linn. Sp. Pl. 678. Griffith, Med. Bot. 289.

The leaves are ovate-oblong, smooth, and shining, with scarcely any smell when entire, but, when bruised, giving out a strong bitter-almond odor; they have an astringent, bitter taste, with a flavor like the peach kernel. They become scentless by desiccation. They possess properties analogous to those of hydrocyanic acid, and some of the preparations from them are used for similar purposes. From the plant not being cultivated in this country, these preparations are not in use.

CHERRY-LAUREL WATER.

R. Fresh leaves of cherry-laurel, one pound.
 Water, two pints and a half.
 Compound spirit of lavender, one ounce.

Chop the leaves, mix them with the water, and distil off one pint (Imp.); agitate this well; if any milkiness remain, filter, then add the spirit of lavender. *Ed. Ph.*

A sedative narcotic, similar to diluted hydrocyanic acid, but of uncertain strength. Dose, thirty minims to a fl. drachm.

R. Essential oil of cherry-laurel, half a drachm.
 Diluted hydrocyanic acid, six drachms.
 Distilled water, twelve fl. ounces.
 Rub together, and filter. *Hæmle.*

INFUSION OF CHERRY-LAUREL.

R. Fresh leaves of cherry-laurel, four ounces.
 Boiling water, two pints.
 Infuse, strain, and add
 Clarified honey, four ounces.
 Used as a lotion in cancer of the lips, and in malignant ulcers. *Cadet.*

OINTMENT OF CHERRY-LAUREL.

- R. Essential oil of cherry-laurel, one drachm.
Lard, one ounce.
Mix. As an application to alleviate pain of cancers. *Soubiran.*

CHERRY-LAUREL WATER LOTION.

- R. Distilled cherry-laurel water, four ounces.
Sulphuric ether, one ounce.
Extract of belladonna, two drachms.
Mix. Said to be very beneficial as a lotion in cases of neuralgia. *Roux.*

MIXTURE OF CHERRY-LAUREL WATER.

- R. Cherry-laurel water, sixty drops.
Peppermint water, two fl. ounces.
Infusion of quassia, four fl. ounces.
Mix. In dysentery, &c., in doses of a table-spoonful, three times a-day. *Augustin.*

CERATE OF CHERRY-LAUREL.

- R. Cherry-laurel water, twelve parts.
Oil of almonds, sixteen parts.
White wax, four parts.
Melt the oil and wax together, and mix the cherry-laurel water, stirring well. As an application to burns. *Roux.*

LAVANDULA.

LAVENDER.

Several species of *Lavandula* are used in medicine, but the only one recognized by the U. S. Ph. is the *Lavandula vera*. This is a small shrub, a native of the south of Europe, and commonly cultivated in our gardens; it has opposite, sessile, narrow leaves, and spikes of small, blue flowers.

Sex. Syst. Didyn. gymnos. *Nat. Syst.* Lamiales.

Linn. Sp. Pl. 800. Griffith, Med. Bot. 501.

The part used is the flowers; these have a strong, fragrant odor, and an aromatic, warm, bitterish taste. Lavender is an aromatic stimulant, and much used in nervous debility, but is mostly employed as a perfume.

COMPOUND POWDER OF LAVENDER.

- R. Lavender, Benzoin, Cloves, Sal ammoniac, each, one ounce.
Petals of red roses, four ounces.
Pomegranate flowers, Chamomile flowers, each, two ounces.

- Pot marigold flowers, two ounces.
Peppermint flowers, six drachms.
Myrrh, four drachms.
Cologne water, two drachms.
Solution of ammonia, eighteen drops.
Oil of cinnamon, six drops.

Mix well. To perfume rooms and drawers. *Guibourt.*

SPIRIT OF LAVENDER.

- R. Fresh lavender, two pounds.
Alcohol, one gallon.
Water, two pints.

Mix, and distil a gallon by a slow fire. *U. S. Ph.*

LAVENDER-WATER.

- R. Oil of lavender, one fl. ounce.
Diluted alcohol, fifteen fl. ounces.

Mix. *Gray.*

- R. Alcohol, five gallons.
Oil of lavender, twenty ounces.
" bergamot, five ounces.
Essence of ambergris, half an ounce.

Mix. *Brande.*

- R. Oil of lavender, four fl. ounces.
Essence of musk, two fl. ounces.
Oil of bergamot,

ten fl. drachms and a half.

Oil of cloves, five drachms.

" roses, one drachm.

" origanum, half a drachm.

" rosemary, half a drachm.

Alcohol, ten pints.

Water, two pints.

Mix. *Gray.*

These are used for perfumes.

- R. Flowering tops of lavender, two pounds.

Water, sufficient.

Distil four pounds by steam. *Par. Codex.*

COMPOUND SPIRIT OF LAVENDER.

- R. Spirit of lavender, three pints.
Spirit of rosemary, one pint.
Bruised cinnamon, one ounce.
" cloves, two drachms.
" nutmeg, half an ounce.
Rasped red saunders, three drachms.

Macerate for fourteen days, express, and filter. *U. S. Ph.*

As a stimulant and carminative, in doses of thirty drops to a fl. drachm.

OINTMENT OF LAVENDER.

R. Oil of lavender,
 " nutmeg,
 Butter of cocoa, equal parts.
 Mix. To increase the growth of hair.

ETHEREAL TINCTURE OF LAVENDER.

R. Oil of lavender, one part.
 Sulphuric ether, seven parts.
 Mix. Dose, five to ten drops. *Beral.*

COMPOUND ESSENCE OF LAVENDER.

R. Oil of lavender, forty-eight drops.
 " cloves, thirty-two drops.
 " orange peel, sixteen drops.
 " bergamot, eight drops.
 Sweet spirit of nitre, eight drops.
 Oil of yellow sandal wood,
 " neroli,
 " roses, each, two drops.
 " cinnamon, one drop.
 Diluted alcohol, one ounce.
 Dissolve, and add
 Honey water, eight ounces.
 Essence of ambergris and
 musk, one ounce.
 Mix. As a perfume. *Redwood.*

LIMON.

LEMON.

The Lemon is the fruit of the *Citrus limonum*, a native of Asia, but now generally cultivated in all warm climates.

Sex. Syst. Polydelph. icosand. *Nat. Syst.* Aurantiaceæ.

Risso, Ann. Mus. xx. 201. Griffith, Med. Bot. 168.

The parts used in medicine are, the juice and the outer rind; the latter has a fragrant odor, and an aromatic, bitter taste; the juice, as is well known, is very acid, with a peculiar, grateful and refreshing taste.

ESSENCE OF LEMON.

R. Pure oil of lemon, one fl. ounce.
 Deodorized alcohol, eight fl. ounces.
 Exterior rind of
 lemon (fresh), half an ounce.

Mix. Macerate for forty-eight hours, and filter; used for flavoring mixtures, and for pastry. *W. Procter.*

R. Thin outer rinds of lemons, at will.
 Alcohol, sufficient.

Put the rinds in a bottle, add sufficient alcohol to cover, and macerate two days. Filter.

For pastry the flavor of this preparation is more delicate, than if made from the oil usually found in the shops. *Francis.*

LOZENGES OF OIL OF LEMON.

R. Oil of lemon, one drachm.
 Sugar, twelve ounces.
 Lemon water, sufficient.

Boil one-third of the sugar, with the lemon water, to the consistence of honey, by a gentle heat, add the rest of the sugar and the oil, and form lozenges. *Cottreau.*

LOZENGES OF LEMON JUICE.

R. Lemon-juice, two ounces and a half.
 Sugar, sixteen ounces.
 Essence of lemon, one scruple.
 Mix, and form lozenges. *Radius.*

COLLUTORY OF LEMON JUICE.

R. Lemon juice,
 Sugar, each, one part.
 Claret wine, sixteen parts.
 As a mouth-wash in scurvy. *Brera.*

LEMONADE.

R. Lemon juice, four ounces.
 Fresh lemon peel, half an ounce.
 Sugar, four ounces.
 Boiling water, three pints.
 Mix, and let cool. *Ellis.*
 R. Lemons, three.
 Boiling water, two pints.
 Sugar, two ounces.

Cut the lemons in slices, pour the water on them, and add the sugar at the end of an hour, and strain. *Ratier.*

COMPOUND LEMONADE.

R. Lemons, two.
 Peel them, remove the seeds, and beat them with
 Toasted bread, two ounces.
 Adding, gradually,
 Infusion of malt, one pint and a half.
 Strain, and add
 Syrup of mulberries, one ounce.
 Wine, six ounces. *Cadet.*

LEMON-PEEL WATER.

R. Pare the rind of one lemon, previously rubbed with half an ounce of loaf sugar; put both into a jar, and add a quart of boiling water; when cold, decant, and add one tablespoonful of lemon juice

A. T. Thomson.

LEMON SYRUP.

R. Strained lemon juice, one pint.
 Sugar, two pounds.
 Mix, and form syrup. *U. S. Ph.*

R. Oil of lemon, half a fl. drachm.
 Citric acid, one ounce.
 Simple syrup, one gallon.

Rub the oil with some powdered sugar, then, with a portion of the syrup, dissolve the citric acid in two ounces of water, and mix the whole.
A. Smith.

SYRUP OF LEMON PEEL.

R. Fresh lemon peel, three ounces.
 Boiling water, one pint.

Infuse for three days, and add
 Syrup, three pounds.
Guibourt.

R. Fresh lemon peel, three ounces.
 Boiling water, one pint and a half.

Infuse for twenty-four hours, strain, and add
 Sugar, two pounds,
 and make syrup. *Francis.*

VINOUS SYRUP OF LEMON PEEL.

R. Fresh lemon peel, half a pound.
 Wine, a pint and a half.

Macerate for twenty-four hours, express, and add

Syrup, one pound and a half.

Boil, strain, and add
 Oil of lemon, rubbed
 with sugar, fifteen drops.
Sard. Ph.

TINCTURE OF LEMON.

R. Fresh lemon peel, three ounces and
 a half.

Proof spirit, two pints.

Macerate for seven days, express, and filter.

Dub. Ph.
 Dose, one fl. drachm.

SPIRIT OF LEMON.

R. Fresh lemon peel, one part.
 Alcohol, four parts.
 Water, two parts.

Macerate for two days, and then distil four parts.
Taddei.

ODORIFEROUS SPIRIT OF LEMON.

R. Oil of lemon,
 " bergamot,
 " lavender, each, one part.
 Acetic ether, four parts.

Mix. Used as a perfume. *Sax. Ph.*

COLOGNE WATER.

R. Oil of bergamot,
 " lemon,
 " cedrat, each, three ounces.
 " rosemary,
 " neroli,
 " lavender, each,

one ounce and a half.

" cinnamon, six drachms.

Alcohol, twenty-four pounds.

Eau de melisse, three pounds.

Spirit of rosemary, two pounds.

Dissolve the oils in the alcohol, add the other articles, and, after eight days, distil four-fifths.

Codex.

R. Balm, two pounds.

Rosemary, eight ounces.

Lemon peel, six ounces.

Nutmeg,

Cloves,

Cinnamon,

Coriander, each, two ounces.

Oil of bergamot, one ounce.

Alcohol, six pounds.

Water, eight pounds.

Mix, and distil by a gentle fire. *Niemann.*

R. Oil of neroli,

" lemon,

" bergamot,

" orange,

" rosemary, each, twelve drops.

Cardamom, one drachm.

Alcohol, one pint.

Mix, and distil two-thirds. Said by Tromsdorff to be the formula used at Cologne. *Niemann.*

R. Oil of bergamot,

" lemon,

" cedrat,

" neroli, each, four drachms.

" cinnamon,

" rosemary, each, one drachm.

Alcohol, three pounds.

Spirit of rosemary, three ounces.

Digest for a few days, and distil almost to dryness; add to the product

Balm water, nine ounces.

Spirit of jasmine,

" orris, each, four drachms.

Giordano.

R. Oil of bergamot, three ounces.

" lemon, two ounces.

" lavender, three drachms and
 a half.

" neroli, two drachms and a half.

" origanum, two drachms.

" rosemary, one drachm.

Essence of vanilla, two drachms.
 Musk, ten grains.
 Alcohol, thirteen pints.
 Rose water, two pints.
 Orange-flower water, one pint.
 Mix, and after fourteen days, filter. *Gray.*

R. Essence of bergamot, two ounces.
 " rosemary,
 " mint, each, one drachm.
 " lemon, two drachms.
 " neroli, ten grains.
 Balm water, two ounces and a half.

Mix. *Ferrara Ph.*

R. Oil of bergamot,
 " lemon, each, one drachm.
 " neroli, half a drachm.
 " cedrat, twenty drops.
 Honey water, one ounce.
 Alcohol, one pint.

Mix. *Gray.*

R. Oil of neroli,
 " orange,
 " lemon,
 " marjoram,
 " rosemary,
 " each, one drachm and a half.
 " cinnamon, four drops.
 Civet, three grains.
 Ambergris, two grains.
 Alcohol, four pounds and a half.

Macerate for a month, and filter. *Swedisch.*

R. Oil of bergamot, three drachms.
 " lemon, two drachms.
 " lavender, twenty-five drops.
 " neroli, fifteen drops.
 " origanum, ten drops.
 Alcohol, two pints.

Mix. *Gray.*

UNPARALLELED WATER.

R. Oil of lemon, half an ounce.
 " bergamot, two drachms and a half.
 " cedrat, two drachms.
 Alcohol, seventy-two ounces.
 Hungary water, eight ounces.

Mix. *Guibourt.*

LINUM.

FLAXSEED.

Flaxseed is the product of *Linum usitatissimum*, or common flax, now cultivated in most parts of the world, but whose native country is unknown.

Sex. Syst. Pentand. pentag. *Nat. Syst. Lina-*
ceæ.

Linn. Sp. Pl. 397. *Griffith, Med. Bot.* 206.

Both the seeds, and the oil extracted from them are officinal. The first are small, oval, shining, and of a brown color. They are devoid of smell, and have a mucilaginous taste when unbroken, but an oily one when chewed. When steeped in water, they afford a viscid, inodorous and almost tasteless mucilage. The oil, which is obtained by expression, is laxative and emollient, but is principally used in the arts.

COMPOUND MEAL OF FLAXSEED.

R. Flaxseed meal, }
 Barley meal, } equal parts.
 Rye meal, }

Mix. To make cataplasm. *Cottrecau.*

EMOLLIENT CATAPLASM.

R. Compound meal of flaxseed, four ounces.
 Water, sufficient.
 Boil to proper consistence. *Par. Cod.*

CATAPLASM OF FLAXSEED.

R. Boiling water, ten fl. ounces.
 Flaxseed meal, sufficient.
 Mix, and make cataplasm. *Lond. Ph.*

CATAPLASM OF FLAXSEED MEAL AND BRAN.

R. Flaxseed meal,
 Bran, equal parts.
 Boiling water, sufficient.
 Pour the water on the bran and meal, and stir constantly until the poultice has a pulpy consistence. *Ellis.*

INFUSION OF FLAXSEED.

R. Flaxseed, half an ounce.
 Liquorice root, bruised, two drachms.
 Boiling water, one pint.
 Macerate for two hours, in a covered vessel, and strain. *U. S. Ph.*

A pleasant demulcent drink, in inflammatory affections of the lungs, bladder, &c.

FLAXSEED MIXTURE.

R. Flaxseed, one drachm.
 Boiling water, six fl. ounces.
 Infuse, strain, and add
 Syrup of poppies, half an ounce.
 Orange-flower water, two fl. drachms.
 Mix. *St. Marie.*

CLYSTER OF FLAXSEED OIL.

- R. Flaxseed oil, two ounces.
 " infusion, eight ounces.
 Mix. *Swediaur.*

FLAXSEED OIL LINIMENT.

- R. Flaxseed oil, twelve fl. ounces.
 Lime water, seven fl. ounces.
 Mix. As an application to burns and scalds. *Ainslie.*

- R. Flaxseed oil,
 Lime water, equal measures.
 Mix.

This is generally called *Carron oil*, and is much used for burns and scalds. *Ed. Ph.*

LIRIODENDRON.

TULIP TREE BARK.

The tulip tree, or *Liriodendron tulipifera*, is one of the most magnificent of our native trees. It is found in most parts of the United States, and is popularly known as the poplar.

Sex. Syst. Polyand. polyg. *Nat. Syst.* Magnoliaceæ.

Linn. Sp. Pl. 755. Griffith, Med. Bot. 98.

The officinal portion is the bark; this, when deprived of its epidermis, is of a yellowish-white color, having a faint odor, but a bitter, pungent, aromatic taste. It is a stimulating tonic, and has been used as a febrifuge, as well as in dyspepsia, chronic rheumatism, &c. The dose in powder is from half a drachm to two drachms.

INFUSION OF TULIP TREE BARK.

- R. Tulip tree bark, one ounce.
 Boiling water, one pint.

Infuse, and strain. Dose, one to two fl. ounces, as a stimulating tonic and diaphoretic. *Wood.*

TINCTURE OF TULIP TREE BARK.

- R. Tulip tree bark, bruised, four ounces.
 Diluted alcohol, one pint.
 Macerate for a week, express, and filter. Dose, about a fl. drachm. *Beasley.*

LOBELIA.

LOBELIA—INDIAN TOBACCO.

Several species of *Lobelia* are medicinal, but the only one recognized as officinal, is the *L. inflata*, a small annual plant, found in most parts of the United States, having numerous small blue flowers, on leafy racemes. The leaves are ovate, sessile, serrate, and hairy.

Sex. Syst. Pentand. monog. *Nat. Syst.* Lobeliaceæ.

Linn. Sp. Pl. 1006. Griffith, Med. Bot. 418.

The whole herb is officinal, but the root and seed vessels are the most active. When dried, it has a somewhat unpleasant odor, and an acrid, burning, nauseous taste, resembling that of tobacco. It is emetic, and, in small doses, expectorant and sudorific. It also possesses narcotic properties. It is principally used in asthma; it has also been used in other diseases of the respiratory organs, &c. Dose in substance, as an emetic, from five to twenty grains, as an expectorant, one to three grains.

INFUSION OF LOBELIA.

- R. Lobelia, one ounce.
 Boiling water, one pint.

Infuse. A fl. ounce every half hour, till vomiting ensues. *Ellis.*

TINCTURE OF LOBELIA.

- R. Lobelia, four ounces.
 Diluted alcohol, two pints.

Macerate for fourteen days, express, and filter.

U. S. Ph.

Dose, as emetic, half fl. ounce; as antispasmodic in asthma, one to two fl. drachms, every two or three hours; and as an expectorant, ten to forty drops.

ETHEREAL TINCTURE OF LOBELIA.

- R. Lobelia, in coarse powder, five ounces.
 Spirit of sulphuric ether, two pints.

Form tincture by displacement. *Ed. Ph.*

Dose, the same as alcoholic tincture.

- R. Lobelia, one pound.
 Alcohol, four pints.
 Spirit of nitric ether, four pints.
 " sulphuric ether, four ounces.

Macerate for fourteen days in a dark place, and filter. Dose, five to twenty minims.

Whitelaw.

MIXTURE OF LOBELIA.

- R. Tincture of lobelia, one drachm.
 Decoction of mallow, six ounces.
 A spoonful, every two or three hours, as an expectorant. *Radius*

ACETIC EXTRACT OF LOBELIA.

- R. Lobelia seed, bruised, eight ounces.
 Diluted alcohol, four pints.
 Acetic acid, one fl. ounce.

Macerate the seed in the diluted alcohol, to which the acid has been added, for forty-eight hours; then displace until four pints of tincture are obtained, using diluted alcohol to expel the last portion, and then evaporate to the consistence of an extract. *W. Procter.*

FLUID EXTRACT OF LOBELIA.

- R. Bruised lobelia tops, eight ounces.
 Acetic acid, one fl. ounce.
 Diluted alcohol, three pints.
 Alcohol, six fl. ounces.

Macerate the lobelia in a pint and a half of the diluted alcohol, mixed with the acid, for twenty-four hours; introduce into a percolator, and obtain three pints of tincture; evaporate this in a water-bath to ten fl. ounces, strain, add the alcohol, and filter. A fl. drachm is equal to half a fl. ounce of the tincture. *W. Procter.*

VINEGAR OF LOBELIA.

- R. Lobelia leaves, in powder, four ounces.
 Diluted acetic acid, one and a half pints.

Macerate for twelve hours, and displace until twenty-four fluid ounces are obtained—using diluted acetic acid to expel the last portion.

W. Procter.

SYRUP OF LOBELIA.

- R. Vinegar of lobelia, six fl. ounces.
 Sugar, twelve ounces (av.).
 Dissolve the sugar in the vinegar, by aid of heat, remove the scum which rises, and strain.

W. Procter.

ANOTHER FORM.

- R. Fluid extract of lobelia, two fl. ounces.
 Simple syrup, ten fl. ounces.
 Mix. *W. Procter.*

LUPULINA.

LUPULIN

Is the powder attached to the strobiles of *Humulus lupulus*, and has all the properties of hops. It is a yellowish powder, mixed with portions of the scales of the hop strobiles. It is tonic, and somewhat narcotic. It is obtained by rubbing, or threshing and sifting the strobiles, or by washing these several times in different portions of water, mixing the washings, and letting the lupulin deposit, when it is to be dried. The dose is from six to ten grains.

POWDER OF LUPULIN.

- R. Lupulin, one part.
 Sugar, two parts.
 Mix. Dose, ten to twenty grains. *Foy.*

PILLS OF LUPULIN.

- R. Lupulin, at will.
 Rub into mass in a warm mortar, and divide into two-grain pills. *Magendie.*
 R. Lupulin, two drachms.
 Gum tragacanth,
 Water, each, sufficient.
 Make mass, and divide into pills of two grains.

EXTRACT OF LUPULIN.

- R. Lupulin, four ounces.
 Alcohol, eight fl. ounces.
 Mix in a percolator, and allow to stand for an hour. Displace until two pints of filtered liquor are obtained; then set aside in a shallow dish for spontaneous evaporation. One drachm of lupulin yields two scruples of extract. *Livermore.*

TINCTURE OF LUPULIN.

- R. Lupulin, four ounces.
 Alcohol, two pints.
 Macerate for fourteen days, and filter. *U. S. Ph.*
 Dose, one to two fl. drachms, in sweetened water.

SYRUP OF LUPULIN.

- R. Tincture of lupulin, one part.
 Syrup, seven parts.
 Mix. Dose, half an ounce to an ounce. *Foy.*

EMULSION WITH LUPULIN.

- R. Tincture of lupulin, four scruples.
 Syrup of red poppies,
 " of orange flowers,
 each, one ounce.
 Emulsion of almonds, four ounces.
 Water, ten ounces.
 Mix. *Beral.*

OINTMENT OF LUPULIN.

- R. Lupulin, one part.
 Lard, three parts.
 Digest, on a water-bath, for five or six hours, and strain. To appease the pain of cancerous ulcers. *Soubéiran.*

M.

MACIS.

MACE.

A lacinated and reticulated, smooth, thin, flexible membrane, of a saffron-yellow color, which is the arillus investing the shell of the nutmeg, which, in taste and odor, it much resembles. It is an active, aromatic stimulant, but is more used for culinary purposes than in medicine. The dose is from ten grains to a scruple.

TINCTURE OF MACE.

R. Mace, one part.
Alcohol (.851), eight parts.
Macerate for eight days, express, and filter.

Beral.

Dose, from thirty to forty drops, as a carminative and stomachic.

OIL OF MACE.

This is prepared by beating nutmegs to a paste, which is to be enclosed in a bag and then exposed to the vapor of water, and afterwards expressing the oil with heated plates.

It is a fat oil mixed with a volatile oil, of a firm consistence and fragrant odor. *Redwood.*

CARMINATIVE DROPS.

R. Oil of mace, half a drachm.
Nitric ether, a drachm and a half.
Mix. In flatulent colic, ten or twelve drops on sugar. *Radius.*

NERVINE BALSAM.

R. Oil of mace, four ounces.
Beef marrow, four ounces.
Melt together, and add
Oil of rosemary, two drachms.
" cloves, one drachm.
Camphor, one drachm.
Balsam of tolu, two drachms.
Dissolved in
Rectified spirit, four drachms.
As a liniment in rheumatism, &c. *Redwood.*

MAGNESIUM.

This metal is not used, but many of its compounds are largely employed.

MAGNESIA.

MAGNESIA.

R. Carbonate of magnesia, any quantity.
Put in an earthen vessel, and expose it to a red heat for two hours, or till the carbonic acid is wholly expelled. *U. S. Ph.*

HEAVY MAGNESIA.

R. Mix solutions of one hundred and twenty-three parts of crystallized sulphate of magnesia, and one hundred and forty-four parts of crystallized carbonate of soda; evaporate to dryness, and calcine till the carbonic acid is expelled. Dissolve out the sulphate of soda, wash, and dry.

R. Phillips.

POWDER OF MAGNESIA AND RHUBARB.

R. Powdered rhubarb, one scruple.
Magnesia, ten grains.
Oil of cinnamon, one drop.

Mix. As a purgative, to be given in sugar and water. *Ellis.*

POWDER OF MAGNESIA AND SULPHUR.

R. Precipitated sulphur,
Magnesia, each, half an ounce.

Mix. A teaspoonful, three or four times a-day, as an aperient. *Ellis.*

POWDER OF MAGNESIA AND ORANGE PEEL.

R. Magnesia, four drachms.
Powdered orange peel,
" fennel-seed,
each, one drachm.
Sugar, two drachms.

Mix. Ten or twelve grains, three times a-day, as a stimulant to the digestive organs. *Foy.*

TROCHES OF MAGNESIA.

R. Magnesia, four ounces.
Sugar, one pound.
Grated nutmeg, one drachm.
Mucilage of tragacanth, sufficient.

Rub the first three ingredients together until thoroughly mixed, then form a mass with the mucilage, and divide into troches weighing ten grains. *U. S. Ph.*

MAGNESIA MIXTURE.

R. Magnesia, one drachm.
Water of ammonia, one fl. drachm.
Spirit of cinnamon, three fl. drachms.
Water, five and a half fl. ounces.

Mix. Recommended by Dr. James, in the cardialgia of pregnant women. Two or three teaspoonfuls, to be taken as occasion may require. *Sims.*

- R. Magnesia, thirty grains.
 Syrup of ginger, two drachms.
 Peppermint
 water, two fl. ounces and a half.
 Compound spirit of
 lavender, half fl. drachm.
 Spirit of caraway, half fl. ounce.
 Mix. A spoonful, every hour, as an antacid. *Foy.*

MAGNESIA AND GENTIAN.

- R. Magnesia, one drachm.
 Infusion of gentian, six fl. ounces.
 Mix. A wineglassful, three times a-day, in cases of uric acid diathesis. *Brande.*

MAGNESIA AND RHUBARB MIXTURE.

- R. Magnesia, half a drachm.
 Powdered rhubarb, two grains.
 " sugar, one drachm.
 Essence of peppermint, six drops.
 Distilled water, one fl. ounce and a half.
 Mix. In bowel complaints of children. A teaspoonful, every two hours, till it operates. *Ellis.*

MAGNESIÆ ACETAS.

ACETATE OF MAGNESIA.

- R. Carbonate of magnesia,
 one hundred and twenty parts.
 Acetic acid, sufficient
 to saturate. Evaporate till the mixture weighs three hundred parts. It forms a syrupy fluid. One ounce of this solution mixed with three ounces of syrup of oranges, constitutes the weaker solution, and one ounce and a half to three ounces of syrup, forms the stronger solution. *Renault.*

MAGNESIÆ CARBONAS.

CARBONATE OF MAGNESIA.

SOLUTION OF MAGNESIA.

- R. Sulphate of magnesia, seven drachms.
 Bi-carbonate of soda, nine drachms.
 Water, twenty fl. ounces.
 Carbonic acid gas, six volumes.

Dissolve the salts in the water, and pass the gas through the mixture. The result will be bi-carbonate of magnesia in solution. *Phabus.*

DINNEFORD'S FLUID MAGNESIA.

- R. Howard's magnesia,
 seventeen and one-half grains.
 Distilled water, one fl. ounce.

Introduce into a cylindrical tinned vessel a mixture in these proportions, and force into it carbonic acid for five hours and a half, during the whole of which time the cylinder is kept revolving. *Pereira.*

DEWEES'S CARMINATIVE.

- R. Carbonate of magnesia, half a drachm.
 Tincture of assafetida, forty drops.
 " opium, twenty drops.
 Sugar, one drachm.
 Distilled water, one fl. ounce.
 Mix. In flatulent colic, diarrhoea, &c., of children. Dose, twenty drops and upwards, according to age. *Deweese.*

DALBY'S CARMINATIVE.

- R. Carbonate of magnesia, two scruples.
 Oil of peppermint, one drop.
 " nutmeg, two drops.
 " aniseed, three drops.
 Tincture of castor, thirty drops.
 " assafetida, fifteen drops.
 Spirit of pennyroyal, fifteen drops.
 Compound tincture of
 cardamom, thirty drops.
 Peppermint water, two fl. ounces.
 Mix. *Paris.*

- R. Carbonate of potassa, half an ounce.
 " magnesia, twelve ounces.
 Laudanum, six fl. ounces.
 Oil of peppermint,
 " caraway,
 " fennel, each, two fl. scruples.
 Sugar, thirty-two ounces.
 Water, ten pints.

Triturate the oils with the sugar and magnesia, then add the remainder. *Phil. Coll. Pharm.*

CARBONATE OF MAGNESIA MIXTURE.

- R. Carbonate of magnesia, half a drachm.
 Sulphate of magnesia, three drachms.
 Aromatic spirit of
 ammonia, one fl. drachm.
 Tincture of rhubarb, half fl. ounce.
 " henbane, half fl. drachm.
 Mint water, four fl. ounces.

Mix. As a carminative cathartic. A tablespoonful, two or three times a-day. *Meigs.*

MAGNESIA AND COLCHICUM.

- R. Carbonate of magnesia, one drachm.
 Sugar,
 Gum Arabic, each, sufficient.
 Wine of colchicum root, forty drops.
 Distilled water, four fl. ounces.

Mix. In gout and rheumatism. A tablespoonful, every two hours, till it operates. *Ellis.*

MIXTURE OF MAGNESIA AND CAMPHOR.

R. Magnesia, one drachm.
 Camphor, half a drachm.
 Sugar,
 Gum Arabic, each, two drachms.
 Sulphuric ether, half fl. drachm.
 Distilled water, four fl. ounces.

Mix. In flatulency and irritable stomach. A tablespoonful, four or five times a-day. *Ellis.*

MAGNESIÆ CITRAS.

CITRATE OF MAGNESIA.

R. Citric acid, at will.
 Dissolve in water, and add
 Carbonate of magnesia, sufficient
 to saturate; wash the powder, and dry by a gentle heat. Laxative, but not as active as the sulphate. *Beasley.*

SOLUBLE CITRATE OF MAGNESIA.

R. Crystallized citric acid,
 one hundred grains.
 Calcined magnesia, thirty-five grains.
 Water, fifteen drops.

Dissolve the acid with the water, then gradually add the magnesia; or omit the water and melt the acid in a sand-bath in its own water of crystallization, and incorporate the magnesia with it. The mixture soon hardens, and may be pulverized for use.

This formula is modified from one given by Dervault, of Paris. *Parrish & Smith.*

SOLUTION OF CITRATE OF MAGNESIA.

R. Soluble citrate of magnesia,
 one ounce.
 Water, eight fl. ounces.

Dissolve, transfer to a suitable bottle, and add

Syrup, one and a half fl. ounces.
 Bi-carbonate of potassa, forty grains.

Cork immediately and secure.

Parrish & Smith.

R. Carbonate of magnesia,
 two hundred and ninety-two grains.
 Crystallized citric acid,
 four hundred and forty-six grains.
 Water, ten fl. ounces.
 Lemon syrup, two fl. ounces.

Dissolve one hundred and thirty-eight grains of the carbonate in two fl. ounces of water, holding in solution one hundred and seventy grains of citric acid, and pour it into a twelve ounce mineral water bottle. The remainder of the magnesia is then triturated with the rest of the water, and poured into the bottle, and one hundred and eighty-five grains of the acid added,

and the bottle immediately corked and tied over. As soon as, with occasional agitation, the opaque fluid becomes slightly milky, remove the cork, filter the solution, and pour it back into the bottle, with two ounces of lemon syrup, and ninety-one grains of citric acid; cork and wire. Each ounce contains one drachm of the citrate. Dose, from a half to a whole bottle.

Rabourdin.

R. Carbonate of magnesia, five drachms.
 Citric acid, seven and a half drachms.
 Syrup of citric acid, two fl. ounces.
 Water, sufficient.

Dissolve the citric acid in four fl. ounces of water, and add to the solution four drachms of the carbonate of magnesia, previously rubbed with three fl. ounces of water. When the reaction has ceased, filter the solution into a strong glass bottle of the capacity of twelve fl. ounces, into which the syrup of citric acid has been introduced. Rub the remaining carbonate of magnesia with two fl. ounces of water, pour into the bottle, cork, and secure with twine.

U. S. Ph.

CITRATE OF MAGNESIA WATER.

R. Subcarbonate of magnesia,
 five drachms and a quarter.
 Citric acid, five drachms and a half.
 Lemon syrup, two drachms and a half.
 Water, sufficient

to fill a Seidlitz-water bottle. Dissolve the acid in one-sixth of the water, triturate the carbonate with the remainder, and add to it half the acid solution; and, on the ceasing of the effervescence, pour it into the bottle with the syrup; add the acid solution, and cork instantly. As a purgative draught. *Bardet.*

MAGNESIÆ SULPHAS.

SULPHATE OF MAGNESIA.

[EPSOM SALTS.]

CHELTENHAM SALTS.

R. Sulphate of magnesia,
 Chloride of sodium, each, four parts.
 Sulphate of soda, three parts.

To be well dried before being powdered and mixed. Dose, half an ounce, or more, in solution. *Cooley.*

The compound saline powder of the *Ed. Phar.* differs from this, in the substitution of sulphate of potash for the sulphate of soda.

SEIDLITZ WATER.

R. Sulphate of magnesia, two drachms
 Chloride of magnesium,
 eighteen grains.
 Soda water, twenty fl. ounces.

Dissolve. As a purgative.

Foy.

R. Sulphate of magnesia, two drachms to one ounce. Dose, two to three drachms, in six ounces of water. *Christison.*

Water, one pint.

Dissolve, and charge the solution with three volumes of carbonic acid gas. *Beasley.*

PURGATIVE EMULSION.

R. Sulphate of magnesia, two drachms.
Flake manna, one ounce.
Simple emulsion, four fl. ounces.

Dissolve. A tablespoonful, every two hours. *Wendt.*

SULPHATE OF MAGNESIA AND TARTAR EMETIC.

R. Sulphate of magnesia one ounce.
Tartar emetic, one grain.
Flake manna, one ounce.
Lemon juice, half an ounce.
Water, eight fl. ounces.

Make a solution, and strain. A tablespoonful, every hour, till it operates. *Dewees.*

SULPHATE OF MAGNESIA AND COFFEE.

R. Sulphate of magnesia, one ounce.
Powdered roasted coffee, two drachms and a half.
Water, sixteen ounces.

Boil in an earthen vessel, for two minutes; remove from fire, and let infuse for some minutes; then filter, or strain. This will destroy the bitter taste of the salt. *Combes.*

CLYSTER WITH SULPHATE OF MAGNESIA.

R. Sulphate of magnesia, two ounces.
Olive oil, one fl. ounce.
Tepid water, one pint.

Mix. One-half to be injected, and if it does not produce the desired effect, the remainder to be administered. *Ellis.*

SULPHATE OF MAGNESIA AND SULPHURIC ACID.

R. Sulphate of magnesia, sufficient to saturate

Water, seven fl. ounces.

Add to solution,

Diluted sulphuric acid, one fl. ounce.

Dose, a tablespoonful, in a wineglassful of water, every hour, till it operates. *Henry.*

SULPHATE OF MAGNESIA AND ROCHELLE SALT.

R. Sulphate of magnesia,
Tartrate of potassa and soda,
equal parts.

SULPHATE OF MAGNESIA AND NITRIC ACID.

R. Sulphate of magnesia, half an ounce.
Tincture of jalap, one fl. drachm.
Nitric acid, two drops.
Mint water, two fl. ounces.

Mix. To be taken for a draught. *Ellis.*

MAGNESII SULPHURETUM.

SULPHURET OF MAGNESIUM.

R. Pure magnesia, five parts.
Sulphur, four parts.

Fuse together. Dose, five to ten grains. *Jourdain.*

SYRUP OF SULPHURET OF MAGNESIUM.

R. Sulphuret of magnesium, half an ounce.
Fennel-water, six ounces.

Dissolve, strain, and add

Sugar, fifteen ounces.

Said to be useful in chronic exanthemata, hooping-cough, &c. Dose, a spoonful, occasionally. *Radius.*

MAGNESIÆ PHOSPHAS.

PHOSPHATE OF MAGNESIA.

R. Magnesia, at will.
Diluted phosphoric acid, sufficient

to saturate; filter, and evaporate to dryness. *Niemann.*

Stated to be useful in rachitis, in doses of ten to twenty grains.

MAGNESIÆ TARTRAS.

TARTRATE OF MAGNESIA.

R. Solution of tartaric acid, at will.
Carbonate of magnesia, sufficient to saturate.

Evaporate the solution to dryness, in a water-bath. Used by Rademacher, in painful chronic maladies of the spleen. Dose, one scruple to one drachm. *Percira.*

APERIENT EFFERVESCING MAGNESIA.

- R. Carbonate of magnesia, one part.
Sulphate of magnesia, }
Bi-carbonate of soda, } each,
Tartrate of soda and } two parts.
potassa,
Tartaric acid,

Drive off the water of crystallization by heat, reduce to powder, mix thoroughly, and enclose in dry, strong bottles, which are to be well corked, and sealed with wax. Dose, a teaspoonful, in half a tumbler of water, drunk in a state of effervescence. *Durand.*

MAGNESIÆ BI-TARTRAS.

BI-TARTRATE OF MAGNESIA.

- R. Tartaric acid, one hundred and twenty-five parts.
Distilled water, two thousand parts.
Dissolve, and add gradually,
Carbonate of magnesia, one hundred and fifty-seven parts.
Evaporate and crystallize. *Ph. Hanov.*

MAGNOLIA.

The barks of several species of Magnolia are employed as stimulating tonics in the United States, but more especially that of *M. Glauca*. It is a stimulating, bitter tonic, with some diaphoretic powers, and the decoction has been used with some success in intermittent fevers and rheumatism, as has also a tincture of the cones. Dose, in powder, half a drachm to a drachm.

TINCTURE OF MAGNOLIA.

- R. Recently-dried bark, or cones of magnolia, four ounces.
Diluted alcohol, one pint.
Macerate for a week, express, and filter. In chronic rheumatism. *Beasley.*

MALTUM.

MALT.

Malt is barley germinated by warmth and moisture, and then subjected to such a degree of heat as to destroy the vital principle. It contains sugar, gum, and hordein, and is principally used in the manufacture of malt liquors.

INFUSION OF MALT.

- R. Ground malt, one pint.
Scalding water, three pints.

Infuse for two hours, strain, and add sugar or lemon juice, if required. *Ellis.*

Much prescribed by the late Dr. Parrish, as a drink in inflammatory fevers.

EMOLLIENT CLYSTER.

- R. Ground malt,
Powdered mallow root,
Pearl barley, each, two drachms.
Water, one quart.
Boil down to a pint, and strain. *Phabus.*

MALVA.

COMMON MALLOW.

Several species of Mallow are recognized as officinal in the European Pharmacopœias, and, although they are very similar in their properties, that most generally employed is the *M. sylvestris*, the leaves and flowers being the officinal parts. They are emollient and demulcent, but are seldom or never employed in this country.

COMPOUND DECOCTION OF MALLOW.

- R. Dried mallow, one ounce.
Dried chamomile, half an ounce.
Water, one pint.
Boil for a quarter of an hour, and strain. Employed for fomentations and enemata. *Lond. Ph., 1836.*

MANGANESIUM.

MANGANESE.

This metal is never employed in medicine, but several of its oxides and salts have obtained some celebrity.

MANGANESII CARBONAS.

CARBONATE OF MANGANESE.

- R. Peroxide of manganese, at will.
Wash in very dilute muriatic acid, dissolve in strong muriatic acid, and evaporate to dryness. Dissolve part in water, and precipitate with carbonate of soda; wash precipitate, and digest it with a solution of the rest of the salt. Filter, and precipitate by carbonate of soda; wash, and dry. *Beasley.*

PILLS OF CARBONATE OF MANGANESE AND IRON.

- R. Crystallized sulphate of iron, seventy-five parts.
Sulphate of manganese, twenty-five parts.

Crystallized carbonate
of soda, one hundred and twenty
parts.
Honey, sixty parts.
Water, sufficient.

Mix well, and divide into pills of three grains.
Burin Dubuisson.

MANGANESII MURIAS.

MURIATE OF MANGANESE.

R. Muriatic acid, at will.
Carbonate of manganese, sufficient

to saturate; evaporate, and crystallize. Preserve in closely-stopped bottles. A solution in water has been praised as a gargle in aphthous sore throat.
Jourdain.

PILLS OF MURIATE OF MANGANESE.

R. Muriate of manganese, two scruples.
Gum Arabic,
Extract of liquorice, each, one scruple.

Mix, and form pills of two grains. Advised in obstinate cutaneous affections, in doses of from five to ten.
Augustin.

MANGANESII IODIDUM.

IODIDE OF MANGANESE.

PILLS OF IODIDE OF MANGANESE.

R. Iodide of potassium,
Sulphate of manganese, equal parts.
Honey, sufficient.

Form mass, and divide into four-grain pills, which are to be kept in a well-stopped bottle. Dose, one pill, gradually increased to six pills, daily.
Hannon.

SYRUP OF IODIDE OF MANGANESE.

R. Sulphate of manganese,
sixteen drachms.

Iodide of potassium,
nineteen drachms.

Sugar,
Water, each, sufficient.

Dissolve each of the salts in three fl. ounces of water containing two drachms of syrup; mix, and after precipitation, filter the solution into a bottle containing twelve ounces of sugar; add water to make a pint, and shake the bottle till the sugar is dissolved. Each fl. ounce contains one drachm of iodide of manganese. Dose, ten drops to half a fl. drachm. *W. Procter, Jr.*

R. Iodide of potassium, 330 grains.
Tartaric acid, 264 grains.

Dissolve each in one and a half fl. ounces of water; mix, filter, and saturate the remaining hydriodic acid with carbonate of manganese. Then filter, and add sufficient syrup to make six fl. ounces. Each fl. ounce contains fifty grains of the iodide of manganese. *Livermore.*

SYRUP OF IODIDE OF IRON AND MANGANESE.

R. Iodide of potassium, 1000 grains.
Proto-sulphate of iron, 630 "
Proto-sulphate of manganese, 210 "
Clean iron filings, 100 "
Powdered sugar, 4800 "
Distilled water, sufficient.

Rub the sulphates and iodide separately to powder, mix with the iron-filings, add half a fl. ounce of water, and rub to an uniform paste; add the same quantity of water a second, and a third time, at intervals of fifteen minutes, and rub. Place the sugar in a bottle, and drain the dense solution into it through a filter, adding water slowly to the magma, until the solution of the iodides is displaced, and the water measures twelve fl. ounces. Lastly, agitate the bottle till the sugar is dissolved. Each fl. ounce contains fifty grains of the iodides, in the proportion of three parts of iodide of iron to one of iodide of manganese. Dose, ten to twenty drops. *W. Procter, Jr.*

MANGANESII OXIDUM.

BLACK OXIDE OF MANGANESE.

POWDER OF OXIDE OF MANGANESE.

R. Oxide of manganese, two grains.
Magnesia, one scruple.

Mix. To be taken every three hours, in chlorosis. *Brera.*

BOLUS OF OXIDE OF MANGANESE.

R. Oxide of manganese, fourteen grains.
Extract of savine,
Aloes, each, ten grains.

Mix, and divide into six boluses; to be taken during the day, as an emmenagogue. *Niemann.*

OINTMENT OF OXIDE OF MANGANESE.

R. Oxide of manganese, one part.
Lard, two parts.
Giordano.

OINTMENT OF OXIDE OF MANGANESE AND SULPHUR.

R. Oxide of manganese,
Sulphur, each, one ounce.

Soap, one ounce.
Lard, three drachms.
Mix. *Beasley.*

Both these ointments have been recommended in porrigo, and other skin diseases.

MANGANESII PHOSPHAS.

PHOSPHATE OF MANGANESE.

R. Sulphate of manganese, four ounces.
Phosphate of soda, five ounces.
Water, sufficient.

Dissolve the salts severally in two pints of water, mix the solutions, wash the precipitate until the sulphate of soda is removed, press in bibulous paper, and dry. *W. Procter, Jr.*

MANGANESII SULPHAS.

SULPHATE OF MANGANESE.

R. Black oxide of manganese, ten parts.
Pounded coal, one part.

Ignite in a gas retort. Dissolve the protoxide thus formed in sulphuric acid, with the addition at the end of a little hydrochloric acid; then evaporate the sulphate to dryness, and heat again to redness in the gas retort. By this process the iron present is rendered insoluble, while the sulphate of manganese may be dissolved out and crystallized. *Graham.*

R. Carbonate of manganese, at will.
Diluted sulphuric acid, sufficient
to saturate. Evaporate and crystallize. Dose, one to two drachms, in half a pint of water; to be taken before breakfast, as a cholagogue cathartic. *Beasley.*

MANNA.

MANNA.

This is the concrete juice of several species of *Fraxinus*, but principally of the *F. ornus*, or *Ornus Europæa*; a small tree found in the south of Europe.

Sex. Syst. Diand. monog. *Nat. Syst.* Oleaceæ.

Linn. Sp. Pl. 1510. *Griffith, Med. Bot.* 445.

There are several varieties of manna—*flake*, common or in *sorts*, and *fat*. The first is the best and purest. It consists of sugar, a peculiar principle called mannite, and a viscous, nauseous matter. Manna is a gentle laxative, but sometimes causes flatulence and pain. Dose, for an adult, one to two ounces; for children, one to four drachms.

LOZENGES OF MANNA.

R. Mallow root,
Water, each, four ounces.

Boil a short time, and add
Manna, twelve ounces.

When dissolved, strain and add
Sugar, six pounds.
Opium, dissolved in
water, twelve grains

Evaporate to the consistence of an electuary and add

Orange-flower water, three ounces.
Essence of bergamot,
" citron, each, four drops.

Evaporate to proper consistence, pour on a marble slab, and divide into lozenges. *Guibourt.*

SYRUP OF MANNA.

R. Manna, six ounces.
Water, twelve fl. ounces.

Dissolve, and strain; then add

Sugar, one pound.
Evaporate to syrup. *Pidereau.*

EMULSION OF MANNA.

R. Sweet almonds, half an ounce.
Manna, two ounces.
Syrup of peach flowers, one fl. ounce.
Infusion of liquorice, four fl. ounces.
Orange-flower water, half fl. ounce.

Make an emulsion. *Radius.*

R. Manna,
Oil of almonds, each, one ounce.
Carbonate of potassa, twelve grains.
Cinnamon water,
Water, each, three fl. ounces.

Rub the first three ingredients together, and gradually add the waters. Two fl. ounces to be taken daily, in divided doses, in inflammation of the kidneys. *Babington.*

MANNA MIXTURE.

R. Sulphate of magnesia, one ounce.
Manna, six drachms.
Tincture of
senna, one fl. drachm and a half.
Boiling water, five fl. ounces.

Mix. As a purge. *Ainslie.*

R. Manna, one ounce.
Sulphate of
soda, one ounce and a half.
Water, six fl. ounces.

Mix. *Augustin*

R. Manna, four ounces
Senna, one ounce
Cream of tartar, half an ounce.
Anise, half an ounce.
Boiling water, four pints

Digest for some time, and strain. *Span. Pharm.*

MANNITE.

MANNITE.

- R. Common manna, six pounds.
 Distilled water, three pints.
 White of egg, one.

Mix, and boil for a few minutes; strain when cold; press impure mannite in a cloth; mix it with its weight of water, and again press. Dissolve in boiling water, with animal charcoal, filter, evaporate to a pellicle, and set aside to crystallize.

Ruspini.

- R. Manna, one part.
 Boiling water, three parts.

Dissolve, and add subacetate of lead to separate gummy and resinoid matters; filter through muslin, and precipitate excess of lead by dilute sulphuric acid. Filter through paper, and concentrate by gentle heat to consistence of syrup. Pour the hot syrup into twice its bulk of cold alcohol, which will precipitate the mannite as the solution cools.

C. T. Bonsall.

Dose, from two drachms to two ounces.

MARANTA.

ARROWROOT.

Arrowroot is the fecula of the roots, or rhizomes of many plants, but that in common use is derived from the *Maranta arundinacea*, a native of the West Indies.

Sex. Syst. Monand. monog. *Nat. Syst.* Marantaceæ.

Linn. Sp. Pl. 2. Griffith, Med. Bot. 637.

West Indian arrowroot is a light, white powder, tasteless and inodorous. It has a firm feel, and crackles when pressed. It is nutritious and demulcent, and is an article of diet well suited to the sick and convalescent, especially in bowel complaints. It is prepared for use by adding to it a certain proportion of boiling water, usually about a pint to a tablespoonful of the fecula.

WATER ARROWROOT.

- R. Rub the arrowroot with a little cold water, till well mixed; then pour boiling water over it, stirring constantly; afterwards boil for five minutes. Sweeten with white sugar, and flavor with lemon juice, or some aromatic; if wine be added, none of the astringent kinds should be used.

Thomson.

MILK ARROWROOT.

- R. Arrowroot, a tablespoonful.
 Sweet milk,
 Boiling water, each, half a pint.
 Proceed as with water arrowroot.

Ellis.

- R. Milk, fifteen fl. ounces.
 Sugar, two ounces.
 Arrowroot, one ounce.

Proceed as above.

Beral.

ARROWROOT PUDDING.

- R. Rub a tablespoonful of arrowroot with a little cold water, and add to it, stirring constantly, a pint of boiling milk; with this mix one egg and three teaspoonfuls of sugar, previously beaten together. This may be boiled or baked, and forms a good diet in convalescence.

Thomson.

BEEF-TEA ARROWROOT.

This is made in the same manner as the water arrowroot, except that beef tea is used in the boiling state instead of water, and that the mixture is to be boiled twenty minutes instead of five.

Thomson.

MARRUBIUM.

HOREHOUND.

Horehound, *M. vulgare*, is a small, herbaceous plant, a native of Europe, and naturalized in some parts of this country.

Sex. Syst. Didynam. gymnos. *Nat. Syst.* Lamiales.

Linn. Sp. Pl. 816. Griffith, Med. Bot. 512.

The whole herb is officinal. It has a strong and peculiar odor in the fresh state, which is almost lost on drying. Its taste is bitter, and somewhat acrid. It is tonic, emmenagogue, antispasmodic, and, in large doses, laxative. The dose of the powder is from thirty grains to a drachm.

EXTRACT OF HOREHOUND.

- R. Powdered horehound, sufficient.
 Exhaust with water by the process of displacement, and evaporate to proper consistence. The dose is from a scruple to half a drachm.

COMPOUND DECOCTION OF HOREHOUND.

- R. Dried horehound, one ounce.
 Bruised liquorice root,
 Flaxseed, each, half an ounce.
 Boiling water, one pint and a half.

Macerate for four hours, and strain. Dose, from one to two fl. ounces, in catarrh.

Steph. & Church.

PECTORAL MIXTURE.

- R. Extract of horehound,
 " couch grass, each,
 two drachms.

Decoction of dandelion, ten fl. ounces.
Oxymel of squill,
Syrup of fennel, each, two fl. ounces.
Mix. A tablespoonful occasionally. *Augustin.*

SYRUP OF HOREHOUND.

R. Dried horehound, one ounce.
Horehound water, two pints.
Digest for two hours, express, and add
Sugar, four pounds.
Form a syrup. *Cottureau.*

HOREHOUND CANDY.

R. Expressed juice of horehound, one pint.
Sugar, ten pounds.
Boil to proper consistence, pour into moulds, or on a marble slab, and divide. *Cooley.*
Much used to allay irritation of throat, in catarrh.

MASTICHE.

MASTICH.

Mastich is a concrete, resinous exudation from the *Pistacia lentiscus*; a small tree, a native of most of the countries bordering on the Mediterranean.

Sex. Syst. Diœc. pentand. *Nat. Syst.* Anacardiaceæ.

Linn. Sp. Pl. 1455. Griffith, Med. Bot. 186.

Mastich occurs in yellowish, semi-transparent, brittle grains or tears, of a mild, agreeable smell, and a resinous but not unpleasant taste. It is much used by the Turks as a masticatory to sweeten the breath and to strengthen the gums. It is more employed in the arts as the basis of varnishes, than in medicine.

MASTICH DENTIFRICE.

R. Powdered mastich,
Prepared chalk, each, two drachms.
Powdered orris root, half an ounce.
Oil of roses, three drops.
Mix. *Phæbus.*

AROMATIC MASTICH COLLUTORY.

R. Mastich, two drachms.
Balsam of Peru, half a drachm.
Gum Arabic, two drachms.
Orange-flower water, six fl. ounces.
Make an emulsion, and add
Tincture of myrrh, two fl. drachms.
Phæbus.

ANTI-ODONTALGIC PASTE.

R. Mastich,
Sandarach, each, two drachms.

Kino, ten grains.
Opium, two grains.
Oil of rosemary, eight drops.
Spirit of horseradish, sufficient.
Mix, and form a paste. Used to fill decayed teeth. *Gassicourt.*

DINNER PILLS.

R. Aloes, six drachms.
Mastich,
Red roses, each, two drachms.
Syrup of wormwood, sufficient.
Beat into mass, and divide into three-grain pills. Dose, two pills as a laxative.

ETHEREAL TINCTURE OF MASTICH.

R. Mastich, four parts.
Sulphuric ether, one part.
Dissolve. To fill the cavity of carious teeth, for which purpose a small dossil of cotton is saturated with it, and introduced into the cavity. *Soubeyran.*

PICTURE VARNISH.

R. Mastich, three hundred and sixty parts.
Venice turpentine, forty-five parts.
Camphor, fifteen parts.
Spirit of turpentine, one thousand parts.
Dissolve with heat. *Redwood.*

CRYSTAL VARNISH.

R. Mastich, three ounces.
Alcohol, one pint.
Dissolve. Used to fix pencil drawings. *Cooley.*

MATICO.

MATICO.

This is the leaves of a Peruvian plant, or plants; in most cases of the *Piper angustifolium* (*Aritanthe elongata*).

Sex. Syst. Diand. trigyn. *Nat. Syst.* Piperaceæ.

The leaves, as found in commerce, are strongly veined, or reticulated; of a greenish-ash color; mixed with fragments of the stem and flower spikes; of a somewhat aromatic odor, and a warm, spicy taste. They have attained much celebrity as a hemostatic, and have also been given in mucous and other discharges. As a styptic, they are applied in substance to the bleeding part.

INFUSION OF MATICO.

R. Matico, half an ounce.
Boiling water, half a pint.
Infuse for one hour. Dose, two tablespoonfuls, two or three times a-day, or oftener. In hemorrhagic and other discharges. *Dub. P.*

INFUSION OF MATICO AND SENNA.

R. Matico,
 Senna, each, two drachms.
 Boiling water, one pint.
 Infuse, and strain. Dose, one fl. ounce and a-half, repeatedly. *Watmough.*

DECOCTION OF MATICO.

R. Matico, one ounce.
 Water, one pint.
 Boil for ten or fifteen minutes, and strain.
 Given as above. *Jeffreys.*

TINCTURE OF MATICO.

R. Matico, two and a half ounces.
 Diluted alcohol, one pint.
 Macerate for fourteen days, and strain. *Lane.*
 Dr. Jeffreys directs three ounces of matico.
 Dose, from thirty to sixty drops, in water. In hemorrhagic discharges, &c.

R. Powdered matico leaves, eight ounces.
 Proof spirit, two pints.
 Macerate for fourteen days, express, and filter.
 Dose, one fl. drachm. *Dub. Ph.*

EXTRACT OF MATICO.

R. Matico, at will.
 Treat by maceration and displacement with a mixture of three parts of alcohol (.835) and two parts of water. Evaporate the tincture on a water-bath to consistence of an extract.
 Dose, ten to twenty grains.

SYRUP OF MATICO.

R. Matico, four ounces.
 Diluted alcohol, one pint.
 Make tincture by displacement, evaporate to one-half; add
 Sugar, sufficient,
 and form syrup. *Ruschenberger.*
 Used in same cases as the tincture. Dose, one to two fl. drachms.

OINTMENT OF MATICO.

R. Powdered matico, three drachms.
 " opium, three grains.
 Lard, one ounce.
 Mix. As an application to hemorrhoids. *Young.*

MATRICARIA.
GERMAN CHAMOMILE.

This is the flowers of *M. chamomilla*, a small, annual, herbaceous plant, native of many parts of Europe.

Sex. Syst. Syngen. super *Nat. Syst.* Astera-
 ceæ

Linn. Sp. Pl. 1256. *Stokes, Bot. Mat. Med.* iv. 238.

The flowers, which are the officinal part, are smaller than those of the true chamomile. They have a strong, penetrating, unpleasant odor, which is much diminished by drying. Their taste is bitter and somewhat nauseous. The medical properties are much the same as those of chamomile, and it is given in the same manner.

INFUSION OF GERMAN CHAMOMILE.

R. Flowers of German cham-
 omile, six drachms.
 Boiling water, eight fl. ounces.

Infuse for two hours, and strain. Dose as a tonic, a tablespoonful every two hours. An extract and the oil are much used in Germany, for the same purposes as we employ common chamomile. *Walther.*

MEL.

HONEY.

This is a peculiar fluid, prepared from flowers by the *Apis mellifica*, or honey-bee.

In a recent state it is fluid, but by age concretes into a soft, granular mass. It has a peculiar, somewhat aromatic odor, and a sweet taste, followed by faint acidity. It resembles sugar in its properties, but is more laxative. It is principally used in medicine as a vehicle.

CLARIFIED HONEY.

R. Honey, at will.
 Melt by means of a water-bath, and remove the scum. *U. S. Pharm.*

PREPARED HONEY.

R. Clarified honey, half a pint.
 Diluted alcohol, one pint.
 Prepared chalk, half an ounce.

Mix the honey and alcohol, add the chalk, and let the mixture stand for two hours, occasionally stirring. Heat to boiling, filter, and evaporate on water-bath, so that when cold the sp. gr. will be 1.32. *U. S. Pharm., 1840.*

OXYMEL.

R. Honey, five pounds.
 Strong acetic acid, seven fl. ounces.
 Distilled water, eight fl. ounces.
 Mix the acid and water with the honey previously heated. *Lond. Ph.*

R. Honey, two pounds.
 Distilled vinegar, one pint.
 Boil to consistence of syrup, removing scum. *Dub. Pharm., 1826.*

HYDROMEL.

R. Honey, two parts.
Water, thirty-two parts.
Dissolve, and strain. A refreshing and slightly laxative drink. *Foy.*

PECTORAL HYDROMEL.

R. Washed Iceland moss, two ounces.
Hyssop leaves, half an ounce.
Hydromel, two pounds.
Macerate for five days, strain, and add
Sugar, three ounces.
In catarrh, &c., one or two spoonfuls occasionally. *Foy.*

EXPECTORANT MIXTURE.

R. Honey,
Fresh butter, each, two ounces.
Mix, and melt over a gentle fire. A spoonful occasionally. *St. Marie.*

CEROMEL.

R. Yellow wax, one ounce.
Honey, four ounces.
Melt the wax, and add the honey, stirring well.
As an application to indolent ulcers. *Van Mons.*

HONEY WATER.

R. Honey, two parts.
Washed sand, three parts.
Introduce into a retort, and distil on a sand-bath, and remove the oil that floats on the product. Aperient and diuretic; in doses of a scruple to half a drachm. *Spielmann.*

R. Rectified spirit, eight pints.
Rose water, two pints.
Orange-flower water, two pints.
Oil of cloves, half an ounce.
" lavender, half an ounce.
" bergamot, two ounces.
" sandal wood, one drachm.
Honey, one ounce.
Tincture of saffron, one ounce.
Musk, ten grains.
Macerate for a week, and filter. Used as a perfume. *Gray.*

R. Honey,
Coriander, each, eight ounces.
Fresh lemon peel, one ounce.
Cloves, six drachms.
Nutmeg,
Benzoin,
Storax, each, half an ounce.
Vanilla, three drachms.
Rose water,
Orange-flower water, each, five ounces.
Alcohol, forty-eight ounces.

Distil by means of a water-bath. Sometimes amber or musk is added. This is chiefly used for the toilette. *Guibourt*

MELISSA.

BALM.

Several species have been used in medicine, but the only one that is officinal in this country is *M. officinalis*, a small herbaceous plant, which is a native of the south of Europe, and is very generally cultivated in our gardens.

Sex. Syst. Didynam. gymnos. *Nat. Syst.* Lamiales.

Linn. Sp. Pl. 827. *Stokes, Bot. Mat. Med.* iii. 365.

The whole herb is used; it has an agreeable odor, somewhat like that of lemons, and an aromatic, slightly bitter taste. Its medicinal properties are very slight, but it communicates an agreeable flavor to infusions.

INFUSION OF BALM.

(BALM TEA.)

R. Balm, two to four drachms.
Boiling water, six fl. ounces.

Infuse for two hours and strain. Used to favor or restore the eruption in the exanthemata, in doses of a wineglassful. *Walther.*

BALM WATER.

R. Fresh balm, twelve pounds.
Water, sufficient.
Mix, and distil one gallon. *Codex.*

COMPOUND SPIRIT OF BALM.

R. Fresh balm, one pound and a half.
Lemon peel, four ounces.
Cinnamon,
Cloves,
Nutmegs, each, two ounces.
Angelica, dried,
Coriander, each, one ounce.
Alcohol, eight pounds.

Macerate for four days, and distil.

Used as a perfume

Codex.

R. Fresh mint, in
flower, thirteen ounces.
Sage, three ounces and a half.
Angelica, eighteen drachms.
Hyssop,
Marjoram, each, fourteen drachms.
Thyme, thirteen drachms.
Cinnamon,
Coriander, each, twelve drachms.

Rosemary,
Cloves,
Nutmeg, each, ten drachms.
Anise, half an ounce.
Lemon peel, one drachm.
Alcohol (.927),
 one hundred and forty ounces.

Macerate for two days, distil, and rectify.

Guibourt.

This form of the compound spirit of balm is known as the *Eau des Carmes*. It is used as a perfume, stomachic, and stimulant.

R. Compound spirit of balm,
 sixteen parts.

Spirit of mint,
" rosemary, each, twelve parts.
" sage, nine parts.
" thyme, eight parts.

Mix. This is known as *Eau de Dardel*, and is used as above. *Guibourt.*

ANTI-HYSTERIC WATER.

R. Fresh balm, one pound.
Laurel berries,
Cinnamon, each, one ounce.
Myrrh, half an ounce.
Castor, two drachms.
White wine, twelve pounds.

Digest for some time, and distil off one-half. As a stimulant, and antispasmodic in hysteria. *Wirtemberg Ph.*

MENTHA PIPERITA.

PEPPERMINT.

Many species of *Mentha* are used in medicine, but two only are recognized in our pharmacopœia; the *M. piperita* and *M. viridis*; both natives of Europe, and naturalized in this country.

Sex. Syst. Didynam. gymnos. *Nat. Syst.* Lamiaceæ.

Smith, Eng. Bot. x. 687. Griffith, Med. Bot. 502

The whole plant is used; it has a peculiar, aromatic odor, and a balsamic, pungent, camphorated taste, followed by a sensation of coolness. It is aromatic, carminative, and stimulant, and is much employed to expel flatus, obviate nausea, &c.

OIL OF PEPPERMINT.

R. Peppermint, at will.
Water, sufficient.

Distil, and collect the oil that floats on the product. *Par. Cod.*

WATER OF PEPPERMINT.

R. Oil of peppermint, half a fl. drachm.
Carbonate of magnesia, one drachm.
Distilled water, two pints.

Rub the oil with the carbonate of magnesia, and then with the water gradually added, and filter. *U. S. Ph.*

Dose, a tablespoonful.

SPIRIT OF PEPPERMINT.

R. Oil of peppermint, three fl. drachms.
Proof spirit, one gallon.

Dissolve. *Lond. Ph.*

ESSENCE OF PEPPERMINT.

R. Oil of peppermint, two fl. ounces.
Alcohol, one pint.

Mix. *U. S. Ph.*

Dose, from ten to twenty drops.

PEPPERMINT LOZENGES.

R. Oil of peppermint, one drachm.
Water of peppermint, sufficient.
Sugar, twelve ounces.

Mix, and form pastilles weighing nine grains. *Cottureau.*

CARMINATIVE MIXTURE.

R. Essence of peppermint, one fl. ounce.
Peppermint water, four fl. ounces.
Syrup of mallow, three fl. ounces.
Oil of cinnamon, two drops.
" peppermint, three drops.

Mix. A spoonful every two hours, in convulsive hiccough. *Alibert.*

R. Magnesia, half a drachm.
Peppermint

water, two and a half fl. drachms.

Compound spirit of
lavender, half fl. drachm.

Spirit of caraway, four fl. drachms.

Syrup of ginger, two fl. drachms.

Mix. A teaspoonful occasionally, as an antacid and carminative. *Paris.*

MENTHA VIRIDIS.

SPEARMINT.

This species, like the last, although a native of Europe, has become extensively naturalized in many parts of the United States; principally in low, wet situations. It is less powerfully aromatic and pungent than peppermint, but is more agreeable in odor and taste to most persons. It has the same properties, and its preparations are the same as of that article.

INFUSION OF MINT.

R. Dried spearmint, three drachms.
Boiling water, half a pint.
Infuse for fifteen minutes, and strain. *Dub. Ph.*

COMPOUND INFUSION OF MINT.

R. Dried spearmint, two drachms.
Boiling water, sufficient
to afford six fl. ounces of strained infusion.

Add

Sugar, two drachms;
Oil of spearmint, three drops;

Dissolved in

Compound tincture of
cardamom, half fl. ounce.

To allay nausea, &c. Dose, one to two fl. ounces. *Dub. Ph., 1826*

R. Dried mint, two ounces.
Red roses, four scruples.
Boiling water, one pint.
Diluted sulphuric acid, two fl. drachms.

Macerate for half an hour, strain, and add

Sugar, one ounce and a half;

and dissolve. *Guy's Hospital.*

WATER OF SPEARMINT.

R. Oil of spearmint, half a fl. drachm.
Carbonate of magnesia, one drachm.
Water, two pints.

Rub the oil with the carbonate, then gradually with the water, and filter. Dose, a table-spoonful. *U. S. Ph.*

ARQUEBUSADE WATER.

R. Dried mint,
" angelica tops, each, one pound.
" wormwood, half a pound.
Angelica seed, five ounces.
Oil of juniper, half a drachm.
Spirit of rose-
mary, three pints and a half.
Rectified spirit, five gallons.
Water, four gallons.

Mix, and distil six gallons. Much celebrated as a vulnerary, for contusions, and for cleansing and healing ulcers and wounds, especially those caused by firearms. *Redwood.*

ESSENCE OF SPEARMINT.

R. Oil of spearmint, two fl. ounces.
Alcohol, one pint.

Dissolve. Dose, twenty to thirty drops. *U. S. Ph.*

SPIRIT OF SPEARMINT.

R. Oil of spearmint, three fl. drachms
Proof spirit, one gallon
Dissolve. Dose, half a fl. drachm. *Lond. Ph.*

MENYANTHES.

BUCKBEAN.

The *M. trifoliata*, or buckbean, is an aquatic herbaceous plant, with ternate leaves, a native of both Europe and North America.

Sex. Syst. Pentand. monog. *Nat. Syst.* Gentianaceæ.

Linn. Sp. Pl. 207. *Griffith, Med. Bot.* 464.

All parts of the plant are medicinal. They are very bitter, but have very little odor. In small doses, the buckbean is tonic and astringent, in large ones cathartic, and even emetic. As a tonic, the dose of the powdered leaves, or root, is from twenty to thirty grains.

INFUSION OF BUCKBEAN.

R. Buckbean, one ounce.
Boiling water, one pint.

Macerate for two hours, and strain. Dose, one to two fl. ounces. *Saunders.*

EXTRACT OF BUCKBEAN.

R. Clarified juice of buckbean, at will.

Evaporate to the proper consistence. *Guibourt.*
Dose, ten to fifteen grains.

COMPOUND PILLS OF BUCKBEAN.

R. Extract of buckbean,
" valerian, each, two drachms.
Soap, half an ounce.
Rhubarb, one drachm and a half.
Syrup, sufficient.

Mix, and make pills of two grains. Ten, three times a-day, in a costive condition of the bowels. *Vogel*

MIXTURE OF BUCKBEAN, FUMITORY, &c.

R. Extract of buckbean,
" fumitory,
" couch grass, each,
half an ounce.
Chamomile water, ten fl. ounces
Compound infusion of
horseradish, four fl. ounces.
Sulphuric acid, half a drachm

Mix. Two tablespoonfuls a-day as an antiscorbutic. *Sella*

DIURETIC MIXTURE.

R. Extract of buckbean, half an ounce.
 Vinegar of squill, two fl. ounces.
 Tincture of assafetida, half fl. ounce.
 Mix. Dose, thirty-drops, three times a-day, as
 a diuretic. *Augustin.*

MEZEREUM.

MEZEREON.

This is the bark of several species of *Daphne*; two species of which are officinal in the U. S. Pharm., the *D. mezereum*, and *D. gnidium*, both shrubs indigenous to Europe. The first being the most generally used.

Sex. Syst. Octand. monog. *Nat. Syst.* Thymelacæe.

Linn. Sp. Pl. 509. Griffith, Med. Bot. 560.

The officinal portion is the bark. This, as found in the shops, is in long, narrow strips, of a greyish color and fibrous texture. It is almost inodorous, with a sweetish taste at first, soon becoming extremely acrid. It is used as an external irritant and stimulant, and internally as an alterative diaphoretic.

DECOCTION OF MEZEREON.

R. Mezereon, two drachms.
 Water, three pints.
 Boil till reduced to a quart. Add
 Liquorice root, half an ounce.
 Strain. Four to eight fl. ounces a-day. *Thomson.*

In syphilitic affections, especially when there are night-pains and eruptions.

COMPOUND DECOCTION OF MEZEREON.

R. Mezereon, two drachms.
 Bittersweet, half an ounce.
 Burdock, two ounces.
 Water, four pints.

Boil down to three pints, and add

Liquorice root, two drachms.

Strain. In the same cases as above, and in obstinate diseases of the skin. *Van Mons.*

EXTRACT OF MEZEREON.

R. Mezereon, three parts.
 Alcohol, five parts.
 Digest, and repeat the operation twice, with the same proportion of alcohol, each time; unite the tinctures; filter, and distil off three-fourths of the spirit; filter the residue, retaining what is left on the filter. *Ham. Pharm.*

GREEN OIL OF MEZEREON.

R. Extract of mezereon, at will.
 Sulphuric ether, sufficient.

Macerate, express, and permit the ether to evaporate. *Guibourt.*

MEZEREON OINTMENT.

R. Dried bark of garou (*daphne gnidium*), one hundred and twenty-five parts.
 Lard, four hundred and fifty parts.
 White wax, forty-five parts.

Slice the bark, moisten with alcohol, and bruise it well. Digest on a sand-bath with the lard for twelve hours, express, cool, and remove the deposit. Melt the ointment with the lard, and stir till cold.

This is the celebrated *pommade épispastique au garou*. *Codex.*

R. Sliced mezereon, four ounces.
 Lard, fourteen ounces.
 White wax, two ounces.

Moisten the bark with alcohol, beat in an iron mortar to a fibrous mass, digest on a salt-water bath, with the lard and wax previously melted together, for twelve hours; strain with strong expression, and cool slowly to permit the undissolved matter to subside. *U. S. Ph.*

R. Extract of mezereon, one part.
 Alcohol,
 White wax, each, eight parts.
 Lard, seventy-two parts.

Dissolve the extract in the alcohol; add the wax and lard; heat to expel the alcohol, and strain. *Guibourt.*

R. Green oil of mezereon, one part.
 White wax, sixteen parts.
 Lard, one hundred and sixty parts.
 Mix, and melt by means of a water-bath.

Used as a stimulating application to blistered surfaces, and to indolent ulcers.

MONARDA.

HORSEMINT.

Almost all the species of *Monarda* are possessed of medicinal properties, but one only is recognized as officinal, the *M. punctata*. This is a native, perennial, herbaceous plant, with yellow flowers, spotted with brown; usually growing in sandy soils.

Sex. Syst. Diand. monog. *Nat. Syst.* Lamiacæe.

Linn. Sp. Pl. 126. Griffith, Med. Bot. 510.

The whole plant is aromatic, and abounds in a pungent, volatile oil. It is used in infusion, for flatulent colic, and as an emmenagogue.

OIL OF HORSEMINT.

R. Fresh horsemint, at will.
 Water, sufficient.

Distil, and collect the oil that floats on the product.

Internally, as a carminative, in doses of two or three drops, with sugar and water. Externally, as a rubefacient, in low states of fever, rheumatism, &c. In most cases it must be diluted.

OIL OF HORSEMENT LINIMENT.

- R. Oil of horsemint, half an ounce.
Tincture of camphor, two ounces.
Laudanum, two drachms.
Mix. As a rubefacient application. *Atlee.*

MONESIA.

MONESIA.

This is the extract of an unknown tree, perhaps the *Chrysophyllum glycyphlaeum*, coming from South America. It is in the form of hard, thick cakes, having at first a sweet taste, which soon becomes astringent and acrid. It is of a dark brown color, friable, and soluble in water. It has been used with success in various discharges, especially of a chronic character; in chronic bronchitis, &c., in doses of two to ten grains, frequently repeated; also as an application to atonic ulcers.

PURIFIED MONESIA.

- R. Monesia, bruised, one pound.
Boiling water, six pints.
Infuse for twenty-four hours, stirring occasionally; decant, and evaporate by a water-bath. *Beasley.*

MONESIA MIXTURE.

- R. Monesia, two scruples.
Water, seven and a half fl. ounces.
Compound tincture of cardamom, half a fl. ounce.
Mix, and dissolve. Dose, a tablespoonful. *Neligan.*

SYRUP OF MONESIA.

- R. Monesia, one drachm.
Water, one fl. drachm.
Boiling syrup, twelve fl. ounces.
Mix. Dose, half a fl. ounce. *Derosne.*

COMPOUND SYRUP OF MONESIA.

- R. Hot syrup of monesia, sixteen fl. ounces.
Orange-flower water, half a fl. ounce.
Extract of poppies, sixteen grains.
Mix. As above. *Derosne.*

TINCTURE OF MONESIA.

- R. Monesia, one ounce.
Diluted alcohol, nine and a half fl. ounces.
Water, two fl. ounces.
Macerate, and decant. *Donovan.*

- R. Monesia, half an ounce.
Alcohol, two fl. ounces.
Water, seven and a half fl. ounces. *St. Ange.*

As above. Used in injections; half a drachm to a drachm, to six fl. ounces of water.

MONESIA OINTMENT.

- R. Monesia, one drachm.
Lard, one ounce.
Mix. *St. Ange.*
R. Monesia,
Water, each, one part.
White wax, two parts.
Oil of almonds, four parts.
Mix. *Derosne.*
As an application to indolent ulcers.

MORA.

MULBERRIES.

Two species of *Morus* produce fruit which appears to have identical properties; *M. nigra*, a native of Europe, which is officinal in the foreign Pharmacopœias; and *M. rubra*, a native of the United States.

Sex. Syst. Monœc. tetrand. Nat. Syst. Moraceæ.

Mulberries are refreshing and laxative, and form the basis of a grateful drink in febrile cases.

SYRUP OF MULBERRIES

- R. Strained juice of mulberries, one pint.
Sugar, two and a half pounds.
Rectified spirit, two and a half fl. ounces.
Dissolve with a gentle heat, set aside for twenty-four hours, remove the scum, decant, and add the spirit. *Lond. Ph.*
As an addition to cooling drinks in fever.

ROB OF MULBERRIES.

- R. Strained juice of mulberries, at will.
Evaporate to the consistence of honey. *Austr. Ph.*
Used as a detersive application to ulcers, and as an addition to gargles.

MORPHIA.

MORPHIA.

- R. Opium, sliced, one pound.
Distilled water, }
Alcohol, } each, sufficient.
Animal charcoal, }
Solution of ammonia, six fl. ounces
Macerate the opium with four pints of the water, for twenty-four hours; and having worked it with the hand, digest for twenty-four hours and

strain. Operate on the residue twice, in the same manner. Mix the infusions, evaporate to six pints, and filter; then add five pints of alcohol, and afterwards three fl. ounces of the solution of ammonia mixed with half a pint of alcohol. Let rest for twenty-four hours, add the rest of the ammonia, mixed as before, and set aside for twenty-four hours, to crystallize. Purify the crystals by boiling them with two pints of alcohol, till dissolved, and filtering while hot through animal charcoal, and recrystallizing.

U. S. Ph.

R. Mix a concentrated infusion of opium with milk of lime (in which the lime is one-fourth the weight of the opium used); heat the mixture to boiling; filter while boiling hot, through linen, and add an excess of muriate of ammonia. As it cools, the morphia is precipitated.

Mohr.

R. Muriate of morphia, one ounce.
Distilled water, one pint.

Dissolve, and add

Solution of ammonia, five fl. drachms, mixed with one fl. ounce of distilled water; shake together; wash the precipitate with distilled water, and dry by a gentle heat.

Lond. Ph., 1836.

One-sixth of a grain is about equal in power to a grain of opium.

PILLS OF MORPHIA.

R. Morphia, one grain.
Conserve of roses, sufficient.
Mix, and make six pills.

Magendie.

MORPHIA DRAUGHT.

R. Morphia, a quarter of a grain.
Syrup of poppies, one drachm.
Distilled water, one fl. ounce.

Mix. At bedtime.

Brera.

INJECTION OF MORPHIA.

R. Morphia, one grain.
Almond oil,
Oil of lilies, each, half an ounce.

Mix.

R. Morphia, two grains.
Yolk of egg, one.
Oil of chamomile,
" poppies, each, one ounce.

Mix. To ease pain in earache, and used in acute gonorrhœa and hemorrhoids.

Brera.

MORPHIÆ ACETAS.

ACETATE OF MORPHIA.

R. Powdered morphia, one ounce.
Distilled water, half a pint.
Acetic acid, sufficient.

Mix the morphia with the water, and drop in the acid, constantly stirring, till the morphia is saturated and dissolved. Evaporate to consistence of syrup on a water-bath. Dry by a gentle heat, and powder.

U. S. Ph.

Dose, one-eighth to a quarter of a grain.

SOLUTION OF MORPHIA AND IPECACUANHA.

R. Acetate of morphia, one scruple.
Diluted acetic acid, one fl. drachm.
Water, two fl. ounces.
Wine of ipecacuanha, four fl. ounces.
Diluted alcohol, ten fl. ounces.

Mix. Dose, a teaspoonful, containing about one-sixth of a grain of the acetate, and one grain of ipecacuanha.

COMPOUND POWDER OF ACETATE OF MORPHIA.

R. Acetate of morphia, one grain.
Powdered foxglove, six grains.
Extract of pulsatilla, twelve grains.
Sugar, one drachm.

Mix, and form thirty-six powders. Four a-day, in the cough and sleeplessness of consumptive patients.

Cadet.

BOLUS OF ACETATE OF MORPHIA.

R. Acetate of morphia, one grain.
Olive oil, ten drops.
Crumb of bread,
Honey, each, sufficient.

Mix, and make six boluses.

Brera.

PILLS OF ACETATE OF MORPHIA.

R. Acetate of morphia, one grain.
Conserve of roses, sixteen grains.

Mix, and make eight pills.

Mialhe.

R. Acetate of morphia, one grain.
Golden sulphuret of antimony,
Extract of aconite, each, two grains.
Powdered liquorice,
Honey, each, sufficient.

Mix, and make eight pills.

Brera.

R. Acetate of morphia, one grain.
Powdered digitalis, six grains.
" camphor, ten grains.
" gum Arabic, eight grains.
Syrup of tolu, sufficient.

Beat into mass, and divide into six pills. One may be taken every three hours.

A. T. Thomson.

R. Acetate of morphia, fifteen grains.
 Ammonio-sulphate of copper, half a drachm.
 Inspissated bile,
 Powdered quassia, each, four scruples.
 Mix, and make one hundred pills. Five,
 morning and evening, in diabetes mellitus.
Brendt.

SOLUTION OF ACETATE OF MORPHIA.

R. Acetate of morphia, sixteen grains.
 Acetic acid, two fl. drachms.
 Distilled water, six fl. drachms.
 Mix. Dose, six to twenty-four drops.
Dunglison.

R. Acetate of morphia, eighty-two grains.
 Rectified spirit, five fl. ounces.
 Water, fifteen fl. ounces.

Dissolve, and filter if not clear. This is intended to be of the same strength as laudanum.
Dub. Ph.

R. Morphia, four grains.
 Distilled vinegar, two fl. drachms.
 " water, five fl. drachms.
 Rectified spirit, one fl. drachm.
 Mix. Dose, twenty minims.
Manch. Hosp.

ALCOHOLIC SOLUTION OF ACETATE OF MORPHIA.

R. Acetate of morphia, sixteen grains.
 Alcohol, one fl. ounce.
 Dissolve.
Cottureau.

SYRUP OF ACETATE OF MORPHIA.

R. Acetate of morphia, four grains.
 Dissolve in a small portion of water, and a few drops of acetic acid, and add to
 Syrup, sixteen ounces.
Par. Cod.

MIXTURE WITH ACETATE OF MORPHIA.

R. Solution of acetate of morphia, twenty drops.
 Lactucarium, ten grains.
 Infusion of chamomile, five ounces.
 Syrup of marsh mallow, half an ounce.
 Mix. Dose, a spoonful.
Beral.

CLYSTER WITH ACETATE OF MORPHIA.

R. Starch, one drachm.
 Hot water, one pint.
 Acetate of morphia, one grain.
 Mix. In chronic diarrhoea.
Cadet.

OINTMENT OF ACETATE OF MORPHIA.

R. Acetate of morphia, six to eight grains.
 Lard, one to two drachms.

Mix. As a friction in violent rheumatic pains.
Foy.

R. Acetate of morphia, four grains.
 Mercurial ointment,
 Simple ointment, each, two drachms.

Mix. In frictions on the labia, twice a-day, in cancer of the uterus.
Hildenbrand.

MORPHIÆ BIMECONAS.

BIMECONATE OF MORPHIA.

R. Meconic acid, two hundred grains.
 Boiling water, sufficient.

Dissolve, and add

Morphia, sufficient
 to saturate. (About 310 grains). Evaporate
 to dryness.
Squire.

Dose, one-fourth of a grain.

SOLUTION OF BIMECONATE OF MORPHIA.

R. Bimeconate of morphia, ten grains.
 Rectified spirit, one fl. drachm.
 Distilled water, thirteen fl. drachms.

Mix. About the strength of laudanum.
Beasley.

MORPHIÆ CITRAS.

CITRATE OF MORPHIA.

R. Morphia, sixteen grains.
 Crystallized citric acid, eight grains.
 Distilled water, one ounce,

colored with

Tincture of cochineal, two drachms.

Mix. Dose, six to thirty drops, in the twenty-four hours.
Magendie.

MORPHIÆ HYDRIODAS.

HYDRIODATE OF MORPHIA.

R. Muriate of morphia, two parts.
 Iodide of potassium, one part.

Make a strong solution of each, and mix; wash the precipitate in a little cold water, press in bibulous paper, redissolve in hot water, and let crystallize.
A. T. Thomson.

MORPHIÆ MURIAS.

MURIATE OF MORPHIA.

R. Morphia, in powder, one ounce
 Distilled water, half a pint
 Muriatic acid, sufficient

Mix the morphia with the water, drop in the acid, constantly stirring, till the morphia is saturated and dissolved. Evaporate by means of a water-bath to crystallization. Dry upon bibulous paper. *U. S. Ph.*

Dose, one-eighth to a quarter of a grain.

SOLUTION OF MURIATE OF MORPHIA.

R. Muriate of morphia, one drachm and a half.
Rectified spirit, five fl. ounces.
Distilled water, fifteen fl. ounces.

Mix the spirit and water, and dissolve the muriate in the mixture with the aid of heat.

Ed. Ph.

Eighteen minims contain one-sixth of a grain of the muriate, equal to a grain of opium.

SYRUP OF MURIATE OF MORPHIA.

R. Muriate of morphia, one grain.
Syrup, four ounces.

Mix. Dose, a teaspoonful. As an expectorant. *Cadet.*

MIXTURE OF MURIATE OF MORPHIA.

R. Muriate of morphia, sixteen grains.
Acetic acid, four drops.
Distilled water, one fl. ounce.
Alcohol, one fl. drachm.

Mix. Dose, six to twenty drops. *Cadet.*

COMPOUND SYRUP OF MURIATE OF MORPHIA.

R. Muriate of morphia, two grains.
Syrup of pinks, ten ounces.
" balm, four ounces.
" orange flowers, two ounces.

Mix. Dose, half an ounce to an ounce. *Cadet.*

LOZENGES OF MURIATE OF MORPHIA.

R. Muriate of morphia, one scruple.
Tincture of tolu, half a fl. ounce.
Sugar, twenty-five ounces.

Dissolve the muriate in a little hot water, mix it and the tincture with the sugar, beat into a mass with mucilage, and divide into lozenges of fifteen grains. Each lozenge contains a fortieth of a grain of the muriate. *Ed. Ph.*

LOZENGES OF MURIATE OF MORPHIA AND IPECACUANHA.

R. Made as the last, with the addition of one drachm of ipecacuanha.

Ed. Ph.

MORPHIÆ NITRAS.

NITRATE OF MORPHIA.

R. Morphia, at will.
Dilute nitric acid, sufficient.

Saturate, dissolve, evaporate, and let crystallize. *Giordano.*

MORPHIÆ PHOSPHAS.

PHOSPHATE OF MORPHIA.

Make like the last, using dilute phosphoric acid instead of nitric.

MORPHIÆ SULPHAS.

SULPHATE OF MORPHIA.

R. Morphia, in powder, one ounce.
Distilled water, half a pint.
Diluted sulphuric acid, sufficient.

Mix the morphia with the water, drop in the acid, constantly stirring till the morphia is saturated and dissolved. Evaporate on water-bath and let crystallize. Dry crystals on bibulous paper. *U. S. Ph.*

Dose, one-eighth to a quarter of a grain.

BOLUS OF SULPHATE OF MORPHIA.

R. Sulphate of morphia, two grains.
Oil of almonds, sufficient.
Sulphate of iron, four grains.
Crumb of bread,
Honey, each, sufficient.

Mix, and make eight boluses.

R. Sulphate of morphia, one grain.
Ipecacuanha, three grains.
Oil of almonds, twelve drops.
Extract of aconite, two grains.
Liquorice powder,
Honey, each, sufficient.

Mix, and make six boluses. One, every three or four hours. *Brera.*

PILLS OF SULPHATE OF MORPHIA.

R. Sulphate of morphia, two grains.
Cyanide of potassium, four grains.
Mucilage, sufficient.

Mix, and make twenty-four pills. One every six hours, in neuralgia. *Rougier.*

R. Sulphate of morphia, one grain.
Olive oil, sufficient.
Ipecacuanha, three grains.
Nux vomica, two grains.
Crumb of bread,
Honey, each, sufficient.

Mix, and make six pills. One, every two hours. *Brera.*

SOLUTION OF SULPHATE OF MORPHIA.

R. Sulphate of morphia, eight grains.
Distilled water, half a pint.

Dissolve. *U. S. Ph.*

One fl. drachm contains an eighth of a grain.

R. Sulphate of morphia, four grains.
Distilled water, one fl. ounce.

Dissolve. One fl. drachm contains half a grain.
Copland.

SYRUP OF SULPHATE OF MORPHIA.

R. Sulphate of morphia, one grain.
Dissolve in a little water, and add to

Syrup, four ounces.

Each ounce contains one quarter of a grain of sulphate. *Cadet.*

MIXTURE OF SULPHATE OF MORPHIA.

[MAGENDIE'S SOLUTION.]

R. Sulphate of morphia, sixteen grains.
Acetic acid, four drops.
Distilled water, one ounce.
Alcohol, one drachm.

Mix. Dose, six to twenty drops. *Magendie.*

LOTION OF SULPHATE OF MORPHIA AND BORAX.

R. Sulphate of morphia, six grains.
Borax, half an ounce.
Rose water, eight fl. ounces.

Mix. As an application in pruritus vaginæ, washing first with tepid soap and water. *Meigs.*

MORPHIÆ TARTRAS.

TARTRATE OF MORPHIA.

R. Morphia, at will.

Saturate with

Solution of tartaric acid, sufficient.

Evaporate and crystallize. *A. T. Thomson.*

MOSCHUS.

MUSK.

This is a peculiar concrete substance obtained from the *Moschus moschiferus*, a small animal of the deer kind, inhabiting the mountainous regions of central Asia. The musk is secreted in the male, in an oval sac, situated near the generative organs. It is found in commerce in these sacs; it is concreted or granular; of a brownish color; soft and greasy to the touch; of a powerful, penetrating odor, and of a bitter, unpleasant, somewhat acrid taste. From its high price it is very liable to adulteration. It is antispasmodic and stimulant, and was formerly much used in spasmodic diseases of all kinds, as well as a stimulant in low states of the system. The dose is from five to ten grains.

POWDER OF MUSK.

R. Musk, three grains.
Opium, half a grain.
Gum Arabic, one scruple.
Sugar, two drachms.

Triturate together, and divide into six powders. One, every three hours, in whooping-cough.

Augustin.

R. Musk, sixteen grains.
Valerian, twenty-four grains.
Camphor, eight grains.

Mix. As an antispasmodic, in hysteria, &c.; in doses of three to twelve grains. *Guibourt.*

This is called by Jourdain, *Tonquin powder*, but this name belongs rather to the following:

R. Musk, sixteen grains.
Cinnabar, twelve grains.

Mix. For a single dose. *Spielmann.*

Once celebrated as a remedy in hydrophobia.

MUSK BOLUS.

R. Musk, five to ten grains.
Camphor, five grains.
Syrup, sufficient.

Make a bolus. Antispasmodic. *Ellis*

R. Musk, ten grains.
Carbonate of ammonia, each, ten grains.
Conserve of roses, sufficient.

Make a bolus. One, every three hours. In mortification, attended with spasmodic action.

Ellis.

MUSK PILLS.

R. Musk, eight grains.
Opium, two grains.
Camphor, twenty-four grains.
Syrup, sufficient.

Mix, and form six pills. To be taken during the day, in hospital gangrene. *Dupuytren.*

R. Musk, ten grains.
Camphor, one scruple.
Ammoniac, two scruples.
Opium, four grains.

Mix, and make pills of four grains. Four or five in the twenty-four hours, in nervous disorders. *Richard.*

R. Musk, twelve grains.
Castor, twenty-four grains.
Assafetida, thirty grains.
Tincture of valerian, sufficient.

Make twenty-four pills. Four, three times a-day, in nervous complaints. *Radius*

MUSK MIXTURE.

- R. Musk, two drachms.
 Sugar,
 Gum Arabic, each, one drachm.
 Distilled water, six fl. ounces.

Mix. A tablespoonful, every hour or two, in low fevers. *Ellis.*

- R. Musk mixture, six fl. ounces.
 Paregoric elixir, half fl. ounce.
 Ammon. tinct. of valerian, one fl. drachm.

Mix. A teaspoonful, three or four times a-day, in pertussis in children. *Ellis.*

- R. Musk, two grains.
 Fennel water, six fl. ounces.
 Laudanum, ten drops.
 Syrup of poppies, two drachms.

Mix. A spoonful, every hour, in trismus. *Cheselden.*

- R. Musk mixture,
 Camphor water, each, three fl. ounces and a half.

Syrup of ginger,
 Spirit of sulphuric ether, each, two fl. drachms.

Mix. A tablespoonful, every three or four hours, in low fevers. *Ainslie.*

MUSK CLYSTER.

- R. Musk, ten grains.
 Valerian, half an ounce.
 Starch, half a drachm.
 Boiling water, eight ounces.

Mix. As an antispasmodic and excitant. *Radius.*

- R. Musk, twelve grains.
 Sugar, two scruples.
 Spirit of ammonia, thirty drops.
 Infusion of flaxseed, four fl. ounces.

Mix. For children with convulsions. *Ellis.*

TINCTURE OF MUSK.

- R. Musk, one part.
 Alcohol, twelve parts.

Digest for twelve days, and filter. Dose, thirty to sixty drops. *Guibourt.*

MUCUNA.

COWHAGE.

This is the bristles of the pods of *Mucuna pruriens*, a perennial, twining plant, native of the warmer parts of America, bearing curved brown pods, covered with short bristles, which, when dry, readily separate.

Sex. Syst. Diadelph. decand. *Nat. Syst.* Fabaceæ.

De Candolle, Prod. ii. 405. Griffith, Med. Bot. 242.

The pod is shaped somewhat like the italic S, and is covered with brown, bristly hairs, which adhere to any substance coming in contact with them. These hairs or spicula are possessed of anthelmintic powers, but whether they act mechanically or not, has not been fully ascertained, though the probability is that they do.

ELECTUARY OF COWHAGE.

- R. Cowhage, two scruples.
 Syrup, half an ounce.

Mix. A teaspoonful every morning, fasting, for three days, to be followed by a dose of castor oil; in cases of lumbrici. *Correa.*

- R. Cowhage, one drachm.
 Honey, sufficient

to make electuary. To be given as above. *Ellis.*

OINTMENT OF COWHAGE.

- R. Cowhage, seven and a half grains.
 Lard, one ounce.

Mix thoroughly.

This is used as a counter-irritant by rubbing it on the skin for ten or twenty minutes. *Blatin.*

MYRISTICA.

NUTMEG.

The nutmeg is the kernel of the fruit of *Myristica moschata*, a small tree, a native of the Molucca islands, and now cultivated in various tropical regions.

Sex. Syst. Diœc. monadelph. *Nat. Syst.* Myristicaceæ.

Thunberg, Act. Holm. 1782. Griffith, Med. Bot. 109.

The nutmeg is an oval, oblong nut, of a lightish-brown color externally, and of a reddish-grey with red veins, internally; of an agreeable, fragrant odor, warm, aromatic taste, and unctuous feel. It is principally used for culinary purposes, but is also employed in medicine as a stimulant, and to disguise the taste of nauseous remedies.

VOLATILE OIL OF NUTMEG.

- R. Grated nutmeg, at will.
 Water, sufficient.

Distil, and separate the oil. Dose, one or two drops.

AROMATIC POWDER.

- R. Cinnamon, two ounces.
 Ginger, each,
 Cardamom seed,
 Nutmeg, grated, each, one ounce.

Rub together into very fine powder. *U. S. Ph.*
Stimulant and carminative; dose, ten to thirty grains.

GOELIS'S ANTIHECTIC POWDER.

R. Burnt hartshorn,
Powdered nutmeg,
Roasted laurel berries, each, one part.
Liquorice, three parts.

Mix, and make a powder. Advised in the hectic attendant on scrofulous affections, in doses of ten grains. *Augustin.*

STIMULATING CLYSTER.

R. Powdered nutmeg, one drachm.
Powdered columbo, one drachm.
" salep, one scruple.
Infusion of flaxseed, four fl. ounces.

Mix. As a stimulating enema. *Ammon.*

ESSENCE OF NUTMEG.

R. Volatile oil of nutmeg, one fl. ounce.
Alcohol, nine fl. ounces.
Mix, with agitation. Dose, twenty drops. *Dub. Ph.*

SPIRIT OF NUTMEG.

R. Nutmeg, bruised, two ounces.
Diluted alcohol, one gallon.
Water, one pint.
Mix, and distil one gallon, with a slow fire. *U. S. Ph.*

Dose, one to two fl. drachms; principally used to flavor other medicines.

COMPOUND SPIRIT OF NUTMEG.

R. Nutmeg, two ounces.
Lemon-peel, }
Orange-peel, } each, one ounce.
Spear-mint, }
Balm, }
Diluted alcohol, four pints.
Water, twelve pints.
Distil three pints. *Van Mons.*

As a stomachic, and an external application to contusions.

MYRRHA.

MYRRH.

Myrrh is the concrete juice of *Balsamodendron myrrha*, a small shrubby tree, with spiny branches, a native of Arabia, &c.

Sex. Syst. Octand. monog. *Nat. Syst.* Amyridaceæ.

Nees, Offic. Planz. liv. 17. *Griffith, Med. Bot.* 171.

Myrrh occurs in tears, or in agglutinated masses, of various shades of color; the best is of a reddish-yellow color, somewhat translucent, having a peculiar, aromatic odor, and a warm, bitter taste. It is a stimulating tonic. It is given in a variety of diseases, and is used externally as an application to foul ulcers, aphthæ &c. The dose is from ten to thirty grains usually in combination.

POWDER OF MYRRH AND IPECACUANHA.

R. Powdered myrrh, twelve grains.
" ipecacuanha, six grains.
" nitre, half a drachm.

Mix, and divide into four powders. One, every fourth hour. Stimulating expectorant. *Paris.*

EMMENAGOGUE POWDER.

R. Myrrh, twelve grains.
Saffron, three grains.
Oil of cloves, one drop.
Rub into powder. *Augustin.*

PILLS OF MYRRH AND ZINC.

R. Sulphate of zinc, ten grains.
Powdered myrrh, one drachm and a half.
Conserve of roses, sufficient.

Mix, and form twenty pills. Two, twice a-day. In pertussis. *Paris.*

PILLS OF MYRRH AND SULPHATE OF IRON.

R. Myrrh,
Sulphate of iron, each, two scruples.
Carbonate of potassa,
Soap, each, half a drachm.
Rub into mass, and form forty pills. Two, thrice a-day. In amenorrhœa. *Ellis.*

PILLS OF MYRRH AND SQUILL.

R. Myrrh, one drachm and a half.
Dried squill, half a drachm.
Extract of henbane, two scruples.
Distilled water, sufficient.
Rub into mass, and form thirty pills. Two, night and morning. In catarrh and phthisis. *Paris.*

PILLS OF MYRRH AND CANADA BALSAM.

R. Myrrh, three drachms.
Canada balsam, one drachm and a half.
Opium, half a drachm.

Mix, and make pills of two grains. Two to four, every hour, in ulcerated phthisis. *Augustin.*

ALKALINE SOLUTION OF MYRRH.

R. Myrrh, two ounces.
 Carbonate of soda, one drachm.
 Boiling water, eight fl. ounces.
 Macerate for two days and filter. *Swedidaur.*

COMPOUND EXTRACT OF MYRRH.

R. Myrrh, two ounces.
 Gum Arabic, two drachms.
 Triturate together; add sufficient water to make a thick emulsion, and mix well with

Extract of couchgrass, four ounces.
 Advised in phthisis, and ulcers of the uterus.
 Dose, one to three drachms, mixed with water, two or three times a-day. *Swedidaur.*

MYRRH COLLUTORY.

R. Lime water, one fl. ounce and a half.
 Tincture of myrrh, two fl. drachms.
 Honey of roses, two drachms.

Mix. *Pharm. Chirurg.*

R. Tincture of myrrh, two fl. drachms.
 Infusion of sage, six fl. ounces.
 Honey of roses, two ounces.

Mix *Augustin.*

MYRRH GARGLE.

R. Tincture of myrrh,
 Vinegar, each, two fl. ounces.
 Honey, one ounce.
 Infusion of contrayerva,
 one pint and a half.

Mix. In putrid sore throat, scarlatina, &c. *Fothergill.*

R. Honey of roses, one ounce and a half.
 Barley water, twelve fl. ounces.
 Tincture of myrrh, six fl. drachms.
 Vinegar, one fl. ounce.

Mix. As a gargle in putrid sore throat. *Ainslie.*

STIMULATING INJECTION.

R. Myrrh, one ounce.
 Quicklime, two ounces.
 Water, two pints.

Infuse for a few days, and decant. As an injection in fistulous ulcers. *St. Marie.*

TINCTURE OF MYRRH.

R. Myrrh, bruised, four ounces.
 Alcohol, three pints.

Macerate for fourteen days, and filter. *U. S. Ph.*

TINCTURE OF MYRRH AND HELLEBORE.

R. Tincture of myrrh, one fl. ounce.
 " black hellebore,
 half a fl. ounce.

Tincture of Spanish flies,

two fl. drachms.

Mix. As an emmenagogue, in doses of thirty drops, three times a-day, in a little sugar and water. *Ellis.*

GRIFFITH'S MYRRH MIXTURE.

R. Myrrh, one drachm.
 Sulphate of iron, one scruple.
 Carbonate of potassa, one drachm.
 Sugar, two drachms.
 Water, six fl. ounces.

Make mixture. As a tonic in phthisis, in tablespoonful doses, according to circumstances. *Ellis.*

R. Myrrh,
 Sugar, each, one drachm.
 Carbonate of potassa,
 twenty-five grains.

Rub together, and add, gradually,
 Rose water,
 seven and a half fl. ounces.
 Spirit of lavender, half a fl. ounce.

Then add
 Crystallized sulphate of iron,
 rubbed to powder, one scruple.

Mix. *Ellis.*

COMPOUND MIXTURE OF MYRRH.

R. Powdered myrrh, one drachm.
 Carbonate of potassa, half a drachm.
 Sulphate of iron, twelve grains.
 Mucilage gum Arabic, two fl. drachms.
 Decoction of liquorice,
 six and a half fl. ounces.

Spirit of allspice, one fl. ounce.
 Rub the myrrh, potassa, sulphate of iron, and mucilage, well together; add gradually the other ingredients. Dose, a tablespoonful twice or thrice a-day. *Babington.*

Nearly the same as Griffith's mixture, and used in the same cases.

OIL OF MYRRH.

R. Myrrh, two parts.
 Washed sand, three parts.
 Distil, separate the oil that passes, and rectify it. *Wirt. Ph.*

MYRRH PLASTER.

R. Balsam of Peru,
 Camphor,
 Powdered myrrh, each,
 one ounce and a half.
 Lead plaster, thirty-two ounces.

Triturate the first three ingredients together and when intimately mixed, add the melted plaster, and stir until chilled; then form into rolls of half a pound each.

N.

NAPHTHALINA.

NAPHTHALINE

Is a product of the distillation of coal tar, and is deposited from the rectified oil of coal tar, in white, shining, concrete crystals, and may be purified by dissolving in alcohol and recrystallizing. It is an active expectorant in doses of eight to thirty grains.

SYRUP OF NAPHTHALINE.

R. Naphthaline, sixteen grains.
Dissolve in smallest quantity of hot alcohol, and triturate with

Syrup, four ounces.

Dose, a tablespoonful every two hours.
Dupasquier.

OINTMENT OF NAPHTHALINE.

R. Naphthaline, one scruple.
Lard, five drachms.

Mix. As an application in dry tetter, lepra, psoriasis, &c.
Emery.

LOZENGES OF NAPHTHALINE.

R. Naphthaline, five scruples.
Sugar, twenty ounces.
Oil of aniseed,
Mucilage of tragacanth, each, sufficient.

Mix, and form lozenges of fifteen grains. Expectorant. One occasionally, to the amount of twenty a-day.
Dupasquier.

NARCOTINA.

NARCOTINE

Is a crystallizable, white, tasteless, inodorous principle, existing in opium. It forms bitter salts with the acids. It has been used with success as an antiperiodic, in doses of three grains, three times a-day.

It may be obtained by treating opium, or the extract of opium, with ether, or by the following process:—

R. Residue of opium, exhausted by water, at will.

Dry it, powder it coarsely, add cold acetic acid, express, and filter; add an excess of ammonia, wash the precipitate with cold water, dissolve in boiling alcohol, decolorize by means of animal charcoal, let cool, and crystallize.
Pereira.

BOLUS OF NARCOTINE.

R. Narcotine, one scruple.
Oil of almonds six drops.

Crumb of bread,

Honey, each, sufficient

Mix, and make eight boluses. One, three times a-day, in the apyrexia of intermittents.

Brera.

MIXTURE OF NARCOTINE.

R. Narcotine, one drachm.
Lemon syrup, one fl. ounce.
Lemon water, two fl. ounces.

Mix. Dose, a dessertspoonful.
Brera.

NARCOTINÆ MURIAS.

MURIATE OF NARCOTINE.

R. Opium, two pounds.
Alcohol, twenty pounds.

Rub together, gradually adding the spirit, till the opium is exhausted, decant, and press residue; to solution add ammonia, till turbid. Distil off three-fourths of alcohol, and let the product cool; wash the deposit with water, and then with a drachm of muriatic acid mixed with a quart of water. Filter the solution, and evaporate to dryness. Dose, about the same as of narcotine.
O'Shaughnessy.

[See Dunglison's New Remedies, 6th edit. p. 536.]

NUX VOMICA.

NUX NOMICA

Is the seed of the *Strychnos nux vomica*, a moderate-sized tree, a native of many parts of the East Indies, having a very bitter wood.

Sex. Syst. Pentand. monog. *Nat. Syst.* Loganiaceæ.

Linn. Sp. Pl. 271. *Griffith, Med. Bot.* 469.

The seeds are flat, peltate, with narrow annular striæ, somewhat downy on the surface; they are of a horny consistence, inodorous, and of a bitter, acrid, somewhat nauseous taste. *Nux vomica* is a violent excitant of the cerebro-spinal system, and in large doses an active poison. In small doses, frequently repeated, it is tonic, diuretic, and even laxative. It owes its energetic properties to the presence of strychnia and brucia. It is principally used in paralysis, in doses of about five grains, three or four times a-day.

POWDER OF NUX VOMICA.

R. Powdered *nux vomica*, three grains
Gum Arabic,
Sugar, each, twelve grains.

Mix. Advised in chronic dysentery. One, every day.
Soubeiran.

R. Powdered nux vomica, eighteen grains.
 " ipecacuanha,

twenty-four grains.

Powdered rhubarb, one drachm.

Prepared chalk, two scruples.

Oil of peppermint, two drops.

Mix. To be divided into twelve powders. One every three or four hours, in cardialgia, &c.

Vogt.

EXTRACT OF NUX VOMICA.

R. Nux vomica, one pound.
 Alcohol, sufficient.

Expose the nux vomica to steam, till softened, slice, dry, and reduce it to powder. Place this in a displacement apparatus, add alcohol till it comes away without bitterness. Distil off the alcohol from the tincture, and evaporate to a proper consistence.

U. S. Ph.

Dose, half a grain to a grain.

TINCTURE OF NUX VOMICA.

R. Powdered nux vomica, one part.
 Alcohol, four parts.

Macerate for fourteen days, and filter.

Cottureau.

Dose, five to twenty drops, in paralysis.

R. Nux vomica, rasped, eight ounces.
 Alcohol, two pints.

Macerate for fourteen days, express, and filter.

Dose, five to fifteen drops. *U. S. Ph.*

R. Powdered nux vomica, two ounces.
 Spirit of nitric ether, two pints.

Digest ten days, and filter. Dose, half a fl. drachm, in seminal debility.

Mettauer.

COMPOUND TINCTURE OF NUX VOMICA.

R. Extract of nux vomica, twenty-four grains.
 Camphor, one drachm.
 Tincture of pellitory, one fl. ounce.

Mix. Dose, twenty drops, four times a-day, in paralysis. *Radius.*

R. Tincture of nux vomica,
 " Spanish flies, each,

one fl. drachm.

Phosphoric ether, one fl. scruple.

Mix. Thirty drops, three or four times a-day, in paralysis. *Radius.*

PILLS OF NUX VOMICA.

R. Powdered nux vomica, thirty grains.
 Conserve of roses, sufficient.

Mix, and make ten pills; one, twice or thrice a-day, in paralysis, closely watching the effects.

Ellis.

PILLS OF EXTRACT OF NUX VOMICA.

R. Extract of nux vomica, one scruple.
 " liquorice, seven scruples.

Mix, and make eighty pills. Two to four, two or three times a-day, in paralysis. *Radius.*

COMPOUND PILLS OF NUX VOMICA.

R. Extract of nux vomica, six grains.
 Black oxide of iron, one drachm.

Mix, and make twenty-four pills. Three a-day, in atonic incontinence of urine. *Mondiere.*

PILLS OF NUX VOMICA AND ALOES.

R. Extract of nux vomica, ten grains.
 Pills of aloes and myrrh, four scruples.

Mix well, and form thirty-six pills. One or two, night and morning. *Copland.*

MIXTURE OF NUX VOMICA.

R. Extract of nux vomica, ten grains.
 Elder water, six fl. ounces.
 Mucilage of gum Arabic,
 Syrup of mallow, each, one fl. ounce.

Mix. A spoonful every two hours, in chronic dysentery. *Ammon.*

LOTION OF NUX VOMICA.

R. Extract of nux vomica, eight grains.
 Alcohol, two fl. ounces.
 Stronger water of ammonia, half fl. ounce.

Mix. As a stimulating lotion to paralyzed limbs. *Radius.*

O.

OLEUM ANIMALE EMPYREUMATICUM.

DIPPEL'S ANIMAL OIL.

R. Oil of hartshorn, at will.
Distil off one-fourth, by a slow heat, on a sand-bath, keeping it protected from the light.

Van Mons.

Antispasmodic, diaphoretic, and anodyne, in doses of five or ten drops; poisonous in large doses.

LINIMENT OF DIPPEL'S OIL.

R. Dippel's oil, one drachm.
Oil of almonds, six drachms.
Turpentine soap, half an ounce.

Mix. As an exciting liniment, also as an application to the abdomen, in cases of worms.

Sundelin.

TINCTURE OF DIPPEL'S OIL.

R. Dippel's oil, one part.
Sulphuric ether, fifteen parts.

Dissolve. Fifteen to thirty drops, as a stimulant and antispasmodic.

Beral.

MIXTURE OF DIPPEL'S OIL.

R. Dippel's oil, one drachm.
Hoffmann's anodyne, three drachms.

Dissolve. Twenty to thirty drops, four times a-day, in chorea and tetanus.

Radius.

OLEUM CAJUPUTI.

OIL OF CAJEPUT.

This is a fluid, transparent, green volatile oil, of an aromatic and pleasant odor, and a warm, pungent taste; obtained from the *Melaleuca cajuputi*, a small tree, a native of the Molucca Islands.

Sex. Syst. Polyadelph. icosand. Nat. Syst. Myrtaceæ.

Maton, Lond. Pharm. 1800. Griffith, Med. Bot. 296.

It is also furnished by other species. It is an active stimulant, when given internally, in doses of a few drops, and acts as a rubefacient when applied to the skin.

MIXTURE OF OIL OF CAJEPUT.

R. Oil of cajeput, half a drachm.
Dippel's animal oil, one drachm.

Mix. As a stimulant. Dose, five to fifteen drops.

Augustin.

CAJEPUT LINIMENT.

R. Cajeput oil,
Camphor, each, three drachms.
Soft soap, two ounces.
Alcohol, one pint.
Water of ammonia, two fl. ounces.

Mix. As an embrocation. Chapman.

OLEUM MORRHUÆ.

COD LIVER OIL.

This, which is also called *Oleum jecoris aselli*, is procured from the livers of several kinds of fish, but principally from the Cod (*Gadus Morrhua*), though that from the Ray is said to be preferable. There are several varieties; one, clear and limpid, the other dark-colored, and somewhat acrid, which is said by many writers to be the most powerful. The pale oil has a faint fishy smell, and a bland taste. It has obtained much celebrity as an alterative, in strumous affections; in chronic gout, rheumatism, and cutaneous diseases, &c. The dose for an adult, is from half a tablespoonful to three tablespoonfuls, two or three times a-day.

MIXTURE OF COD LIVER OIL.

R. Cod liver oil, one fl. ounce.
Gum Arabic,
Sugar, each, two drachms.
Cinnamon, or mint water, four fl. ounces.

Mix. One or two tablespoonfuls, morning and evening. Ellis.

R. Cod liver oil, four fl. ounces.
Solution of carbonate of potassa, half a fl. ounce.
Syrup of orange-peel, half a fl. ounce.
Peppermint water, seven fl. ounces.

Mix. Dose, one and a half fl. ounces to three fl. ounces. Beasley.

R. Cod liver oil, one fl. ounce.
Solution of carbonate of potassa, two fl. drachms.
Syrup of orange-peel, one fl. ounce.
Oil of calamus, three drops.

Mix. Dose, one to two fl. drachms, night and morning, for rickets in children. Fehr.

R. Cod liver oil,
Syrup of orange-peel,
Aniseed water, each, one fl. ounce
Oil of calamus, three drops

Mix. Three spoonfuls a-day, in rachitis, and gouty swellings. Phœbus

R. Cod liver oil, half a fl. ounce.
 Solution of potassa, forty drops.
 Peppermint water, half a fl. ounce.

Mix for a draught. This should be followed by a teaspoonful of lemon-juice to liberate the oil on the stomach. *Percival.*

SYRUP OF COD LIVER OIL.

R. Cod liver oil, eight parts.
 Powdered gum Arabic, five parts.
 Simple syrup, four parts.

Make an emulsion, and add

Sugar, twenty-four parts.

Dissolve by gentle heat, and add

Orange-flower water, two parts. *Duclo.*

LINIMENT OF COD LIVER OIL.

R. Cod liver oil, one fl. ounce.
 Water of ammonia, half a fl. ounce.

Mix. *Brach.*

Cod liver oil, four fl. drachms.

Lead water, two fl. drachms.

Yolk of egg, three drachms.

Mix. As an application to ulcers, &c. *Brefeld.*

OINTMENT OF COD LIVER OIL.

R. Cod liver oil, three fl. ounces.
 Spermaceti, six drachms.
 White wax, two drachms.

Melt together, and stir till cold. *Beasley.*

R. Caustic soda, one ounce.
 Water, two fl. ounces and a half.

Dissolve, and add

Cod liver oil, eight fl. ounces.

Agitate briskly. *Deschamps.*

R. Cod liver oil, ten parts.
 Lead water, five parts.
 Lard, ten parts.

Mix. *Brefeld.*

COMPOUND OINTMENT OF COD LIVER OIL.

R. Cod liver oil, one fl. drachm.
 Red oxide of mercury, four grains.
 Simple cerate, two scruples.

Mix. *Cunier.*

R. Cod liver oil, three drachms.
 Extract of wood soot, two drachms.
 Citrine ointment, one drachm.
 Beef marrow, six ounces.

Mix. *Carron.*

In opacities of the cornea, and scrofulous ophthalmia.

COMPOUND OIL OF COD LIVER.

R. Cod liver oil, one fl. drachm.
 Walnut oil, two fl. drachms.

Mix. A drop or two to be introduced between the eyelids, in opacities of the cornea. *Radius.*

OLEUM OLIVÆ.

OLIVE OIL.

The oil of the fruit of *Olea Europæa*, a small tree, originally from Syria, but now extensively cultivated in the countries bordering on the Mediterranean.

Sex. Syst. Diand. monog. *Nat. Syst.* Oleaceæ.

Linn. Sp. Pl. 11. *Griffith, Med. Bot.* 442.

Good olive oil is an unctuous fluid, of a pale yellow or greenish color; almost inodorous, and of a bland taste. It is principally used as an article of food, but is also employed in medicine as a demulcent, emollient, and laxative, and in the composition of liniments, ointments, &c. The dose, as a laxative, is about a fl. ounce.

OLIVE OIL MIXTURE.

R. Syrup of gum, four fl. ounces.
 Olive oil, half fl. ounce.

Mix. As a laxative. *Radius.*

R. Olive oil, eight fl. ounces.
 Aromatic spirit of ammonia, two fl. drachms.

Mix. Three tablespoonfuls, night and morning, as an anthelmintic. *Ellis.*

R. Olive oil, one fl. ounce.
 Solution carbonate of potassa, half fl. drachm.
 Mint water, seven fl. ounces.

Mix. *Guy's Hospt.*

R. Olive oil, one fl. ounce.
 Water of carbonate of ammonia, one fl. drachm.
 Mint water, seven fl. ounces.

Mix. *Guy's Hospt.*

R. Olive oil mixture, eight fl. ounces.
 Manna, one ounce and a half.

Mix. Dose, three tablespoonfuls as a laxative. *St. Bart. Hospt.*

OLIVE OIL CLYSTER.

R. Common salt, one tablespoonful.
 Olive oil,
 Molasses, each, two tablespoonfuls.
 Warm water, one pint.

Mix. *U. S. Dispens.*

R. Manna, one ounce.
Compound decoction of
chamomile, ten fl. ounces.
Dissolve, and add
Olive oil, one fl. ounce.
Sulphate of magnesia half an ounce.
Mix. *Dub. Ph., 1826.*

OLEUM RICINI.

CASTOR OIL.

The oil of the seeds of *Ricinus communis*, a perennial tree in tropical countries, but an annual herbaceous plant, in temperate latitudes. The seeds are ovate, compressed, bean-like, of a greyish-ash color, marbled with reddish-brown, not unlike the dog-tick in appearance. *Sex. Syst. Monœc. monad. Nat. Syst. Euphorbiacæ.*

Linn. Sp. Pl. 1430. Griffith, Med. Bot. 599.

The oil, which is generally obtained by expression, is a thick, viscous, colorless fluid, with a faint, but unpleasant odor, and a mild, but nauseous taste, followed by a slight sensation of acidity. It is a mild, but prompt cathartic, acting rather as an evacuant than as an excitant of the alvine secretions. The dose is about a fl. ounce; for infants, from one to four fl. drachms. Its disgusting taste is best disguised by mixing it with froth of porter.

OLEAGINOUS MIXTURE.

R. Powdered gum Arabic, two drachms.
Sugar, one drachm.
Rub together with a little mint water, and add gradually,
Castor oil, one ounce.
Mint water, four fl. ounces.

Triturate well. A tablespoonful, every hour or two hours, till it acts. *Ellis.*

In some cases, as in dysentery, it is of benefit, to add thirty or forty drops of laudanum to the mixture.

R. Castor oil,
Mucilage of gum Arabic,
Syrup of orgeat, each, two ounces.
Water, ten fl. ounces.
Make an emulsion. Dose, as last. *Beral.*

R. Castor oil, one ounce.
Yolk of egg, one.
Water of orange-peel, two fl. drachms.
Water, two fl. ounces.
Make an emulsion. *Radius.*

R. Castor oil, one ounce.
Yolk of egg, one.
Orange-flower water, half an ounce.

Simple syrup, half an ounce.
Water, two ounces.
Make an emulsion. *Cottreau.*

R. Castor oil, one ounce.
Yolk of egg, one.
Peppermint water, half a fl. ounce.
Water, two fl. ounces.
Syrup, one fl. ounce.

Mix the yolk with a little water, add the oil gradually, rubbing briskly in a mortar, then add slowly the remainder of the waters and syrup. *Codex.*

R. Castor oil, eleven drachms.
Powdered tragacanth, half a drachm.
White sugar, seventy-five grains.
Water, two and a half fl. ounces.
Syrup of orange-flowers, six fl. drachms.

Triturate the tragacanth with the sugar; add the syrup, and rub well in a mortar, until the mucilage begins to thicken, then add the oil, and continue rubbing till it is homogeneous, adding the water gradually, during the process. *Manne.*

EMULSION OF CASTOR OIL SEED.

R. Castor oil seed, half an ounce.
Anise water, four fl. ounces.
Sugar, two drachms.

Deprive the seed of their exterior coat, triturate them to a uniform pulp with a little water, and the sugar, and finally add the remainder of the water gradually, and strain through a coarse cloth. The seed are more acrimonious than the oil, wherefore it is better to commence this preparation in small doses. Dose, a teaspoonful, to be gradually increased to a tablespoonful.

W. Procter, Jr.

ANTHELMINTIC EMULSION.

R. Castor oil,
Mucilage of gum
Arabic, each, one ounce.
Syrup of Corsica moss,
Water of semen contra,
" chamomile, each, two fl. ounces.

Emulsion of sweet
almonds, eight fl. ounces.

Mix, and make an emulsion. As an anthelmintic. Dose, a fl. ounce. *Beral.*

CASTOR OIL CLYSTER.

R. Castor oil, one ounce and a half.
Yolks of eggs, two.
Infusion of chamomile, six fl. ounces.
Mix. *Radius*

MIXTURE OF CASTOR OIL AND ETHER.

R. Castor oil, one ounce.
Sulphuric ether, two drachms.
Mix. A spoonful every two hours, advised as an anthelmintic, in cases of tape-worm.

Radius.

BANDOLINE.

R. Castor oil, two ounces.
Spermaceti, one drachm.
Arnotta, half a drachm.
Oil of bergamot, one drachm.
Otto of roses, five drops.

Mix, melt by a moderate heat, and strain. To stiffen, and keep hair in form. Redwood.

OLEUM TEREBINTHINÆ.

OIL OF TURPENTINE.

This is usually known as *Spirits of Turpentine*, and is the volatile oil obtained from the turpentine afforded by several species of *Pinus*. It is limpid, colorless, of a strong, penetrating, peculiar odor, and of a warm, pungent, somewhat bitterish taste. It is stimulant, diuretic, anthelmintic, and cathartic, and externally, rubefacient.

PURIFIED OIL OF TURPENTINE.

R. Oil of turpentine, eight parts.
Alcohol, one part.
Agitate together, and pour off the spirit, and repeat the process several times. Nimmo.

LOTION FOR CHILBLAINS.

R. Oil of turpentine, four parts.
Sulphuric acid, one part.
Olive oil, ten parts.
Mix. To be applied to the affected part, night and morning. Gassicourt.

OIL OF TURPENTINE AND HONEY.

R. Oil of turpentine, two fl. drachms.
Honey, one fl. ounce.
Mix. A teaspoonful night and morning, in warm tea. In sciatica. Ellis.

LOTION FOR TOOTHACHE.

R. Oil of turpentine, one fl. drachm and a half.
Oil of cloves,
Oil of cajeput, each, half a fl. drachm.
Balsam of Peru, two drachms.
Opium, two scruples.

To be well rubbed together. As an application to the face, in cases of toothache. Beasley.

TURPENTINE MIXTURE.

R. Oil of turpentine, one hundred and twenty drops.
Powdered gum Arabic,
" sugar, each, two drachms.
Laudanum, sixty drops.
Compound spirit of
lavender, two fl. drachms.
Mint water, five fl. ounces.
Mix. A tablespoonful every two hours, in low forms of fever, &c. Ellis.

VERMIFUGE EMULSION.

R. Oil of turpentine, six fl. drachms.
Gum Arabic, two drachms.
Chamomile water, six fl. ounces.
Sulphuric ether, two fl. drachms.
Mix. Two spoonfuls, night and morning, in cases of tapeworm. Radius.

TURPENTINE MIXTURE.

R. Oil of turpentine, one fl. ounce.
Powdered gum Arabic,
" sugar, each, two drachms.
Mint water, four fl. ounces.
Mix. A tablespoonful every two hours till it operates. An active purgative. Frank.
R. Oil of turpentine, three fl. drachms.
Yolks of eggs, two.
Syrup of mint, two fl. ounces.
" orange flowers,
Ether, each, one fl. ounce.
Tincture of cinnamon, half a fl. drachm.
Mix. A spoonful, three times a-day, in neuralgia and rheumatism. Foy.

R. Oil of turpentine, one fl. ounce.
Yolk of egg, one.
Triturate together, and add, gradually,
Emulsion of almonds, four fl. ounces.
Syrup of orange, two fl. ounces.
Compound spirit of
lavender, four fl. drachms.
Oil of cinnamon, four drops.
One fl. ounce, three times a-day, advised as a purge in iritis. Carmichael.

SPIRIT OF TURPENTINE MIXTURE.

R. Honey,
Oil of turpentine,
Ammoniated tincture of
guaiacum, each, two drachms.
Oil of cloves,
" lemon, each, three drops.
Mix. A teaspoonful, twice or three times a-day, in sciatica and lumbago. Copland.

TURPENTINE CLYSTER.

- R. Oil of turpentine, one fl. ounce and a half.
Yolk of egg, one.
Tepid infusion of flaxseed, one pint.
Mix. *Ellis.*
- R. Oil of turpentine, one fl. ounce.
Yolk of egg, one.
Decoction of barley, nineteen fl. ounces.
Lond. Ph.

TURPENTINE AND ETHER.

- R. Oil of turpentine,
Sulphuric ether, equal parts.
- Mix. Dose, twenty to forty drops, in honey or syrup, in biliary calculi, and as an external application in rheumatism. *Cottureau.*

TURPENTINE LINIMENT.

- R. Oil of turpentine,
“ olives, each, two fl. ounces.
Tincture of camphor, one fl. ounce.
Water of ammonia, one fl. drachm.
- Mix. As an external rubefacient. *Ellis.*
- R. Soft soap, two ounces.
Camphor, one ounce.
Oil of turpentine, sixteen fl. ounces.
- Mix. A powerful rubefacient. *Lond. Ph.*
- R. Oil of turpentine, half a pint.
Resin cerate, one pound.
- Melt the cerate, and add the turpentine.
U. S. Ph.

This is known as *Kentish's ointment*, and is much used as an application to burns and scalds.

ACETIC TURPENTINE LINIMENT.

- R. Oil of turpentine, three fl. ounces.
Acetic acid, five fl. drachms.
Rose water, two and a half fl. ounces.
Essence of lemon, four fl. scruples.
Yolk of egg, one.
- Mix. As an external embrocation and liniment in phthisis. *Stokes.*

SULPHURIC TURPENTINE LINIMENT.

- R. Oil of turpentine, three fl. ounces.
Sulphuric acid, one fl. drachm.
Olive oil, three fl. ounces.
- Mix. *Brodie.*

OPIATED TURPENTINE LINIMENT.

- R. Oil of turpentine, one fl. ounce.
“ chamomile, two fl. ounces.
Laudanum, one fl. drachm.
- Mix. As a lotion in neuralgia. *Recamier.*

STARKEY'S SOAP.

- R. Dry carbonate of potassa,
Oil of turpentine,
Turpentine, equal parts.
- Mix the potassa with the oil, then add the turpentine, and triturate till the mixture is of the consistence of honey. Used in dropsy, and in gonorrhœa. Dose, eight to ten grains. *Guibourt.*

DIURETIC WINE.

- R. Oil of turpentine, two fl. drachms.
Lemon juice, one fl. ounce.
Wine, four fl. ounces.
- Mix. For a single dose. *Pierquin.*

OLEUM TIGLII.

CROTON OIL.

This oil is procured, for the most part, from the seed of the *Croton tiglium*, but also from those of two or three other species. They are all natives of India, and the adjoining parts of Asia. The *C. tiglium* is a moderate-sized shrub, bearing a somewhat triangular nut, containing three ovoid seeds, of a reddish-brown color, having an oleaginous kernel, which affords, on pressure, the oil in question.

Sex. Syst. Monœc. monadelph. *Nat. Syst.* Euphorbiacæ.

Linn. Sp. Pl. 1426. *Griffith, Med. Bot.* 597.

The oil, when pure and fresh, is nearly colorless, or yellowish; but when kept for some time, becomes of a reddish-brown or orange color. It has a faint, but peculiar smell, and an acrid and hot taste. It is a powerful hydragogue purgative, and has been much used in dropsy, apoplexy, and visceral obstructions; when applied externally, it causes irritation and inflammation of the skin, followed by a pustular eruption; and has been successfully employed in rheumatism, neuralgia, and bronchial and pulmonary affections. Dose for an adult, one to two drops.

BOLUS OF CROTON OIL.

- R. Croton oil, one drop.
Powdered gum Arabic, half a drachm.
Syrup of orange flowers, sufficient.
- Mix, and make four boluses. Two to four to be taken in the morning. *Foy.*

PILLS OF CROTON OIL.

- R. Croton oil, six drops.
Soap, half a drachm.
Oil of caraway, eight drops.
Powdered liquorice root, sufficient.
- Mix, and make twelve pills. Dose, one or more. *Reece.*

R. Croton oil, six drops.
 Pills of aloes and myrrh, one drachm and a half.
 Soap, one scruple.
 Powdered liquorice root, sufficient.

Mix, and make thirty pills. Dose, two to three, or more. *Copland.*

R. Croton oil, one drop.
 Crumb of bread, sufficient.

Mix, and make four pills. One, every hour, until they operate. *Ellis.*

R. Croton oil, two drops.
 Soap, two grains.
 Gum Arabic, sufficient.

Mix, and make four pills. *Foy.*

COMPOUND CROTON OIL PILLS.

R. Powdered scammony,
 Powdered aloes, each, sixty-four parts.
 Croton oil, three parts.
 Alcohol, four parts.

Dissolve the oil in the alcohol; add the solution, gradually, to a mixture of the powders, and beat into a mass. Divide into pills of five grains. One to three, for children of fourteen years of age; three to five, for adults. *Beral.*

PILLS OF CROTON OIL AND QUINIA.

R. Croton oil soap, three grains.
 Sulphate of quinia, one drachm.
 Extract of dandelion, sufficient.

Mix, and make twenty pills. One for a dose. *Caventou.*

PILLS OF CROTON OIL AND BLUE MASS.

R. Croton oil soap, three grains.
 Extract of henbane,
 Blue mass, each, twenty-four grains.
 Oil of pimento, twelve minims.

Mix, and make twelve pills. Two at bedtime. *Neligan.*

LOZENGES OF CROTON OIL.

R. Vanilla chocolate, two drachms.
 Sugar, one drachm.
 Starch, one scruple.
 Croton oil, five drops.

Mix, and make thirty lozenges. *Soubairan.*

SOAP OF CROTON OIL.

R. Croton oil, two parts.
 Solution of caustic soda, one part.

Mix; put into paper moulds; in a few days, slice, and keep in well-stopped bottles. Dose, one to three grains, in pills. *Foy.*

TINCTURE OF CROTON OIL.

R. Croton oil, sixteen drops.
 Alcohol, one ounce.

Macerate for six or eight days, and filter. Dose, fifteen to twenty-five drops. *Soubairan.*

R. Croton oil, four drops.
 Tincture of myrrh, one fl. ounce.

Mix, digest, and filter. Dose, one to two fl. drachms. *Bateman.*

R. Croton oil, eight drops.
 Alcohol, one fl. ounce.

Mix, digest, and filter. Dose, half to one fl. drachm. *Nimmo.*

EMULSION OF CROTON OIL.

R. Croton oil, three drops.
 Almond oil, half a fl. ounce.
 Powdered gum Arabic, two drachms.

Triturate well, and gradually add
 Syrup of orange flowers, one fl. ounce.
 Chamomile water, five fl. ounces.

A tablespoonful every two hours, till it operates. *Phabus.*

R. Croton oil, one drop.
 Yolk of egg, two drachms.

Orange-flower water,
 Mint water, each, one ounce.

Make an emulsion. *Foy.*

MIXTURE OF CROTON OIL.

R. Croton oil, one or two drops.
 Mucilage of gum Arabic,
 Distilled water, each, one fl. ounce.

Mix. A teaspoonful every two hours, until it operates. *Ellis.*

R. Croton oil, two drops.
 White sugar, two drachms.
 Gum Arabic, half a drachm.

Tincture of cardamom,
 half a fl. drachm.

Distilled water, one fl. ounce.

Mix. Dose, two dessertspoonfuls every three or four hours. As it is agreeable to the taste, it is suited for children, but in smaller doses. *Beral.*

R. Tincture of croton oil,
 twenty-five drops.

Mucilage of gum Arabic, one drachm.
 Water, one ounce.

Mix. In the morning, fasting. *Foy.*

SAPONACEOUS SOLUTION OF CROTON OIL.

R. Croton oil, eight drops.
 Potassa, six grains.

Distilled water, two fl. drachms.

Mix. From three to six drops may be given for a dose. *Ellis.*

LINIMENT OF CROTON OIL.

R. Croton oil, one fl. ounce.
Oil of turpentine, seven fl. ounces.
Mix with agitation. *Dub. Ph.*

R. Croton oil, one part.
Olive oil, five parts.
Mix. *Pereira.*

R. Croton oil, one fl. drachm.
Oil of turpentine, one fl. ounce.
Mix. *Corrigan.*

R. Croton oil, four drops.
Carbonate of soda, ten grains.
Spirit of mint, half an ounce.

In friction, in rheumatism. *Foy.*

R. Croton oil,
Solution of potassa, each,
half fl. ounce.

Mix, and agitate.

R. The above solution, thirty minims.
Rose water, one fl. ounce.

Mix. To be used twice a-day, till pustules appear. *J. Allen.*

EMBROCACTION OF CROTON OIL.

R. Croton oil, twenty minims.
Tartar emetic, one scruple.
Solution of potassa, one fl. drachm.
Water, eight fl. drachms.

Mix. To keep up a mild eruption on the skin. *Morris.*

CERATE OF CROTON OIL.

R. Lard, two and a half parts.
Wax, half a part.

Melt together, and when nearly cold, mix with them

Croton oil, one part.
Caventou.

R. Soap cerate, four parts.

Melt, and when semifluid, add

Croton oil, one part.
Beasley.

PLASTER OF CROTON OIL.

R. Lead plaster, four parts.

Melt, and when nearly cold, add

Croton oil, one part.

Spread on linen for an adhesive and irritating plaster. *Bouchardat.*

R. Lead plaster, eight parts.

Melt, and when nearly cold, add

Croton oil, twenty parts.

To be spread as above. A very active counter irritant. *Bouchardat.*

OINTMENT OF CROTON OIL.

R. Croton oil, ten minims.
Lard, half an ounce.

Mix. *Ainslie.*

OPIUM.

OPIUM.

Opium is the inspissated juice of the unripe capsules of the *Papaver somniferum*, and presents many varieties, as the Turkey or Smyrna, the East Indian or Bengal, the Egyptian, &c.; of which the first is the best, and affords the largest proportion of morphia.

Opium contains various peculiar principles, several of which are officinal, and are treated of under their respective titles. Turkey opium is in flattened, rounded masses, of half a pound to two pounds in weight, covered externally with the seed-vessels of some species of dock. The texture is soft; the color is pale brown; the odor is strong and narcotic, and the taste bitter and acrid.

Opium is stimulant, in small and repeated doses, narcotic in large; and also antispasmodic, diaphoretic, sedative, and anodyne. It is used to fulfil a variety of indications; to procure sleep, to lull pain, to check morbid discharges to alleviate cough, &c. The medium dose is one grain, but in spasm, &c., it is given in much larger doses.

EXTRACT OF OPIUM. (AQUEOUS.)

R. Opium, one pound.
Water, five pints.

Cut the opium into small fragments, macerate it for twenty-four hours in a pint of water, break the pieces down with the hand, and express; add another pint of water to the residuum, macerate for twenty-four hours, and again express; repeat this process till all the water has been used. Filter the several infusions, unite them, and evaporate on a water-bath to due consistence. *U. S. Ph*

Dose, half a grain.

EXTRACT OF OPIUM. (AQUEO-ALCOHOLIC.)

R. Opium, one part.
Alcohol, four parts.

Cut the opium in small pieces, and digest it in the alcohol for two days, at a gentle heat; express, and treat the residue with four parts of warm water; express, unite the solutions, and evaporate to due consistence. *Taddei.*

EXTRACT OF OPIUM. (ACETOUS.)

R. Opium, one ounce
Distilled vinegar, two pints.

Cut the opium into small pieces, digest in the vinegar for two days, on a sand-bath, stirring from time to time, decant, filter, and evaporate to due consistence. *Soubéiran.*

EXTRACT OF OPIUM. (ALCOHOLIC.)

R. Opium, at will.
Alcohol, sufficient.

Reduce the opium to small pieces, digest in the alcohol in a closed vessel, by a gentle heat, often stirring, filter, and distil off the alcohol, till the residue is of due consistence.

Antwerp Ph.

EXTRACT OF OPIUM. (VINOUS.)

R. Opium, one part.
White wine, four parts.

Reduce the opium to small pieces, macerate it in the wine for twenty-four hours, occasionally stirring; express. Macerate the residue in two more parts of wine, and express; unite the solutions, and evaporate to proper consistence.

Par. Cod.

EXTRACT OF OPIUM. (DENARCOTIZED.)

R. Aqueous extract of opium, at will.

Rub it with a little water, put it in a flask, add sulphuric ether, agitate, and decant; repeat the process with other portions of ether, as long as anything is taken up, and evaporate the residuum to a pilular consistence. *Robiquet.*

R. Aqueous extract of opium, four parts.
Resin, one part.

Beat together, and add

Boiling water, sixteen parts.

Boil till reduced one-half, add as much cold water as has been boiled away, filter, and evaporate. *Limousin—Lamothe.*

EXTRACT OF OPIUM. (BY FERMENTATION.)

R. Opium, one part.
Water, eight parts.

Mix, and add

Yeast, sufficient.

Let ferment for a week, at a temperature of 68° to 70°; then dilute with water, filter, boil till all vinous odor is dissipated, and evaporate to proper consistence. *Deyeux.*

R. Opium, two ounces.

Dissolve in a mixture of

Quince juice,
Water, each, sixty ounces.

Digest for some days, with a gentle heat, strain, and add

White sugar, four ounces.

Let ferment for a month, stirring from time to time, then strain, and evaporate. *Langolst.*

EXTRACT OF OPIUM. (ROASTED.)

R. Powdered opium, one part.

Heat it on a flat dish, over a moderate fire, constantly stirring, as long as fumes are given off. Treat it twice with six times its weight of cold water, filter, and evaporate. *Guibourt.*

The last three forms of Extract, whilst containing the full proportion of morphine, are deprived of the irritating and virose principles of opium.

ELIXIR OF OPIUM.

R. Opium, ten drachms.
Water, sufficient.
Alcohol, (95 p. ct.), four fl. ounces.

Reduce the opium to a thin pulp with water, let it stand in a cool place for two days, transfer to a glass funnel containing filtering paper, add a superstratum of water equivalent to the bulk of the whole mass, and filter twelve fl. ounces of the solution; to which add the alcohol.

E. Dupuy.

R. Powdered opium, ten drachms.
Ether,
Alcohol, each, four fl. ounces.
Water, sufficient.

Macerate the opium in half a pint of water for two days, and express; macerate the dregs twice, successively, in six fl. ounces of water; mix, and strain the liquors; evaporate to two fl. ounces, and agitate several times with the ether. Separate the ether by a funnel, evaporate the solution of opium to dryness, dissolve the extract in half a pint of water, place on a filter, add sufficient water to make twelve fl. ounces, filter, and add the alcohol. *W. Procter, Jr.*

These elixirs are about the same strength as laudanum. Dose, twenty-five drops.

COMPOUND POWDER OF OPIUM AND CHALK.

R. Powdered opium, six grains.
" cinnamon, one drachm.
" long pepper, eight grains.

Prepared chalk,
one drachm and a half.

Mix, and divide into twelve powders. One, three or four times a-day. In diarrhoea. *Ellis.*

POWDER OF ROASTED OPIUM.

R. Powdered opium, at will.

Put it in a flat dish; moisten it with white wine, expose to a moderate heat, constantly stirring till it is perfectly dry; repeat the operation till the opium is one-half reduced, moisten with vinegar, and rub into paste; dry, and pulverize. As an astringent in hemorrhages and mucous discharges, in doses of one or two grains, with some bitter extract. *Giordano.*

POWDER OF OPIUM, CAMPHOR, &C.

R. Powdered opium, ten grains.
 " camphor, two scruples.
 Carbonate of ammonia, four scruples.
 Starch, fifteen grains.
 Mix, and make eight powders. One, every hour or two, as an antispasmodic. *Swedjaur.*

POWDER OF OPIUM AND NITRE.

R. Powdered opium, four grains.
 " nitre,
 Sugar of milk, each, one drachm and a half.
 Mix, and make six powders. As an antispasmodic. *Phæbus.*

POWDER OF OPIUM AND SULPHUR.

R. Powdered opium,
 " camphor, each, two grains.
 " sulphur,
 " sugar, each, half a drachm.
 Mix, and make four powders. One, every three hours, in lead colic. *Hildebrand.*

POWDER OF OPIUM AND MUSK.

R. Powdered opium, two grains.
 Musk, five grains.
 Magnesia, four grains.
 Sugar of milk, ten grains.
 Mix. To be taken every two to four hours, in delirium tremens. *Vogt.*

POWDER OF OPIUM AND ANTIMONY.

R. Dover's powder,
 James's powder, each, four grains.
 Mix. To be taken every four hours, in obstinate rheumatic pains. *Brera.*

PILLS OF OPIUM.

R. Powdered opium, one drachm.
 Soap, twelve grains.
 Beat into a mass with water, and divide into sixty pills. *U. S. Ph.*

PILLS OF OPIUM, HENBANE, AND HEMLOCK.

R. Powdered opium, four grains.
 Extract of henbane,
 " hemlock, each, fifteen grains.
 Mix, and divide into ten pills. One at night, when an anodyne is required. *Ellis.*

PILLS OF OPIUM AND SULPHURET OF ANTIMONY.

R. Extract of opium, ten grains.
 Precip. sulphuret of antimony, twelve grains.
 Nitrate of potassa, twenty-four grains.
 Syrup, sufficient.
 Mix, and make six pills. One at bedtime, to cause perspiration, and to ease pain in rheumatism. *Recamier.*

PILLS OF OPIUM AND FOXGLOVE.

R. Powdered opium,
 " foxglove, each, six grains.
 Conserve of roses, sufficient.
 Mix, and make twelve pills. One, every four hours, in asthma, &c. *Ellis.*

AROMATIC PILLS OF OPIUM.

R. Extract of opium, }
 Saffron, } each, one
 Powdered cinnamon, } drachm.
 " nutmeg,
 " cardamom, }
 Syrup of orange flowers, sufficient.
 Mix, and make pills of three grains. Known as *Oriental Pills*, and considered to be aphrodisiac. One to three, at bedtime. *Cadet.*

PILLS OF OPIUM AND CAMPHOR.

R. Extract of opium, three grains.
 Camphor, six grains.
 Syrup, sufficient.
 Mix, and make six pills. One to three a-day, as an anodyne, and antispasmodic. *Foy.*

PILLS OF OPIUM AND BUTTER OF CACAO.

R. Butter of cacao,
 Powdered gum Arabic, each, forty-eight grains.
 Extract of opium, twelve grains.
 Syrup of ipecacuanha, sufficient.
 Mix, and make pills of five grains. One in the evening, as an anodyne and expectorant. *Petit*

PILLS OF OPIUM AND MUSK.

R. Extract of opium, twelve grains.
 " valerian,
 Musk, each, twenty-four grains.
 Mix, and make sixteen pills. One, then two, then three a-day, in hysteria. *Fou.*

PILLS OF OPIUM AND SULPHATE OF ZINC.

R. Extract of opium, one grain.
Sulphate of zinc, four grains.
Syrup of gum, sufficient.

Mix, and make four pills. Two a-day, in painful mucous discharges from the urethra or vagina. *Foy.*

PILLS OF OPIUM, HEMLOCK, AND CALOMEL.

R. Extract of opium, eighteen grains.
" hemlock, one drachm.
Calomel, thirty-six grains.
Syrup of mallow, sufficient.

Mix, and make thirty-six pills. Two to six a-day, as an alterative and sedative, in organic affections. *Foy.*

R. Opium, four grains.
Calomel, six grains.
Tartar emetic, one grain.
Extract of hemlock, one scruple.

Mix, and make eight pills; two to be taken at bedtime, in rheumatic pains. *Brande.*

PILLS OF OPIUM AND ACETATE OF LEAD.

R. Extract of opium, one grain.
Acetate of lead, four grains.
Powdered henbane, eight grains.

Mix, and make eight pills. One, morning and evening, in epilepsy. *Recamier.*

R. Acetate of lead, twenty-four grains.
Powdered opium, three grains.
Syrup, sufficient.

Mix, and make twelve pills. One every three hours, in hemorrhages, dysentery, and cholera. *Chapman.*

PILLS OF OPIUM AND ACETATE OF MERCURY.

R. Extract of opium, .
Acetate of mercury, .
Camphor, each, twelve grains.
Syrup of poppies, sufficient.

Mix, and make thirty pills. One, morning and evening, in syphilis. *Carmichael.*

PILLS OF OPIUM, NITRATE OF SILVER, &C.

R. Extract of opium, seventy-two grains.
Nitrate of silver, six grains.
Musk, forty-eight grains.
Camphor, ninety-six grains.

Mix, and make ninety-six pills. One, morning and evening, gradually increasing the dose, in epilepsy, paralysis, &c. *Foy.*

PILLS OF OPIUM, CASTOR, &C.

R. Opium, half a grain.
Castor, six and a half grains.
Powdered digitalis, one grain.
Syrup, sufficient.

Make two pills. One to be taken twice or thrice a-day. In spasmodic asthma and dyspnoea. *A. T. Thomson.*

PILLS OF OPIUM AND LIQUORICE.

R. Powdered opium, ten grains.
Extract of liquorice, one drachm.

Mix, and make eighty pills. One, occasionally, as an expectorant. *Wirt. Ph.*

CONFECTION OF OPIUM.

R. Powdered opium, four drachms and a half.
Aromatic powder, six ounces.
Clarified honey, fourteen ounces.

Rub the opium with the aromatic powder, add the honey, and beat together till thoroughly mixed. *U. S. Ph.*

As a stimulant narcotic, in atonic gout, flatulent colic, &c. It contains one grain of opium in thirty-six grains of the mass.

R. Powdered catechu, four ounces.
" kino, three ounces.
" nutmeg, .
" cinnamon, each, one ounce.

Opium, dissolved in
wine, one drachm and a half.
Syrup of red roses, twenty-seven ounces.

Mix. Each drachm contains rather less than half a grain of opium. *Soubeiran.*

ANTI-ODONTALGIC MASS.

R. Opium, two grains.
White wax, two drachms.
Mastich, one drachm.
Oil of almonds, three drachms.
" cloves, twelve drops.
Cochineal, eight grains.

Rub into a uniform mass. To fill carious teeth. *Clarus.*

R. Opium, five grains.
Oil of cloves, three drops.
Extract of henbane, five grains.
" belladonna, ten grains.
Powdered pellitory, sufficient.

Mix, and make a consistent mass. Used as above. *Rust.*

BALSAM FOR THE TOOTHACHE.

R. Opium, one scruple.
Oil of turpentine, one drachm and a half.

Oil of cloves,
" cajeput, each, half a drachm.
Balsam of Peru, two drachms.

Mix. *Beasley.*

R. Opium,
Camphor, each, two scruples.

Dissolve in a little alcohol, and add

Oil of cloves, one drachm.
" nutmeg, six drachms.
Guaiacum, two drachms.

Mix. *Van Mons.*

TOOTHACHE DROPS.

R. Opium,
Camphor, each, ten grains.
Alcohol, sufficient.

Oil of cloves,
" cajeput, each, one drachm.

Mix. *Copland.*

ANTIDYSENTERIC OPIATE.

R. Purified opium, four grains.
Ipecacuanha, half a drachm.
Tormentilla, one drachm.

Syrup of whortleberries,
Conserve of red
roses, each, six drachms.

Mix. Dose, one drachm, every hour. *Quarin.*

PLASTER OF OPIUM.

R. Powdered opium, two ounces.
Burgundy pitch, three ounces.
Lead plaster, one pound.
Boiling water, four fl. ounces.

Melt together the pitch and plaster; then add the opium mixed with the water, and boil to the proper consistence. *U. S. Ph.*

As an application in rheumatic and other pains

PLASTER OF OPIUM AND CAMPHOR.

R. Powdered opium,
" camphor,
" soap, each, one drachm.

Laudanum, sufficient
to make a plaster. *Ellis.*

R. Opium,
Camphor, each, half a drachm.
Lead plaster, sufficient.

Melt and mix. For local pains. *Paris.*

R. Opium, one drachm.
Simple plaster, two ounces.
Galbanum, one ounce.
Oil of caraway, two drachms.

Melt the last three ingredients, and add the opium. As an application to the abdomen, in flatulent colic, diarrhoea, and dysentery.

Swediaur.

LINIMENT OF OPIUM.

R. Castile soap, six ounces.
Opium, one ounce and a half.
Camphor, three ounces.
Oil of rosemary, six fl. drachms.
Rectified spirit, (Imp.) two pints.

Macerate the soap and opium in the spirit for three days; filter; add the oil and camphor, and agitate briskly. *Ed. Ph.*

As an embrocation in rheumatic pains, sprains, &c.

LINIMENT OF OPIUM AND OIL OF CHAMOMILE.

R. Opium,
Oil of chamomile, each, half a drachm.
" almonds, two drachms.

Mix. As a friction around the eyes, in spasm of the eyelids. *Weller.*

ANTI-OTITIC MIXTURE.

R. Opium, four grains.
Saffron, ten grains.
Myrrh, half a drachm.
Juice of mallow, half an ounce.
Oil of almonds, two ounces.

Triturate well together, and strain. As an injection into the ear, in pain in that organ.

Pierquin.

ANODYNE OINTMENT.

R. Opium,
Saffron, each, one scruple.
Yolk of egg, one.
Poplar ointment,
Acetate of lead ointment, each, one ounce.

Mix. As an application to painful hemorrhoids. *Fulda Ph.*

R. Powdered opium, one scruple.
Lard, one ounce.

Rub together. *Lond. Ph.*

R. Syrup of opium, one ounce.
Lard, three ounces.
Essence of roses, four drops.

Mix. For chapped lips. *Pierquin*

R. Opium, half a drachm.
 Extract of hemlock, one drachm.
 Basilicon ointment, half an ounce.

Mix. As an application to gangrenous ulcers.
Carus.

R. Opium, ten grains.
 Alum, fifteen grains.
 Lard, half an ounce.

Mix. In the treatment of ulcerated, atonic buboes.
Simon.

OINTMENT OF OPIUM AND TAR.

R. Powdered opium, two drachms.
 Tar ointment, one ounce.

Mix. As an application to hemorrhoids. *Ellis.*

CERATE OF OPIUM.

R. Opium, half a drachm.
 Yolk of egg, one.

Mix well, and add
 Simple cerate, one ounce.
 Triturate well together. *Lagneau.*

SYRUP OF OPIUM.

R. Extract of opium, one part.
 Water, sixteen parts.
 Syrup, five hundred parts.

Dissolve the opium in the water, filter, add it to the syrup, heated to boiling, and strain.

Par. Cod.

SUCCINATED SYRUP OF OPIUM.

R. Syrup of opium, one ounce.
 Spirit of amber, two grains.

Mix. *Soubéiran.*

ANODYNE COLLYRIUM.

R. Extract of opium, ten grains.
 Camphor, six grains.
 Boiling water, twelve fl. ounces.

Rub the opium and camphor together, and add the water, and strain. *Ellis.*

R. Extract of opium, four grains.
 Rose water, four ounces.

Dissolve, and strain. *Par. Cod.*

OPIUM FOMENTATION.

R. Extract of opium, two drachms.
 Boiling water, one pint.

Dissolve, and strain. As a fomentation in pruriginous affections. *Radius.*

INJECTION OF OPIUM.

R. Opium, twelve grains.
 Solution of subacetate of lead, twelve drops.

Water, nine ounces.
 Mix. As an injection in gonorrhœa.

Girtenner.

R. Extract of opium, six grains.
 " belladonna, one drachm and a half.

Decoction of wild lettuce, one pound.

Dissolve. As an injection in neuralgia, and hemorrhage of the urethra and vagina. *Foy.*

R. Extract of opium, one and a half grains.
 Distilled water, five fl. drachms.

Dissolve. As an injection in coryza, and other inflammations of the nasal, mucous membrane.

Deschamps.

He directs one nostril to be closed by the finger, and the liquid to be drawn up into the other, &c.

CLYSTER OF OPIUM.

R. Powdered opium, two grains.
 Mucilage of gum Arabic, half fl. ounce.

Tepid milk, two fl. ounces.

Mix. *Ellis.*

R. Flaxseed, one ounce.
 Boiling water, six ounces.

Infuse for an hour, strain, and add

Extract of opium, two grains.
 Dissolve. *Spielmann.*

SUPPOSITORY OF OPIUM.

R. Powdered opium, two grains.
 Soap, four grains.

Mix. *Ellis.*

LINCTUS WITH OPIUM.

R. Extract of opium, one grain.
 " Peruvian bark, four grains.

Camphor, six grains.
 Sugar, one drachm.

White linctus, four ounces.

Mix. In bronchitis, when the cough is violent. *Foy.*

MIXTURE OF OPIUM AND SYRUP OF POPPIES.

R. Opium, two to three grains.
 Syrup of poppies, one ounce.

Chamomile water, six ounces.
 Mix. A spoonful every half hour, in dysentery. *Swedliaur.*

MIXTURE OF OPIUM AND LIME WATER.

R. Extract of opium, one grain.
 Lime water,

Oil of almonds, each, three fl. drachms.
 Mix. For the treatment of sore nipples, to be applied on dossils of lint. *Sibergundi.*

MIXTURE OF OPIUM AND CINNAMON
WATER.

R. Powdered opium, ten grains.
Sugar, one drachm.
Cinnamon water, six fl. ounces.

Mix. A tablespoonful every two hours, in tetanus and colica pictonum. *Ellis.*

WATER OF OPIUM.

R. Opium, in small pieces, one pound.
Water, six pounds.

Macerate for forty-eight hours, and distil one pound. It is given in the dose of one drachm, in syrup. *Foy.*

MURIATE OF OPIUM.

R. Powdered opium, one ounce.
Muriatic acid, one ounce.
Distilled water, twenty ounces.

Mix, and shake the mixture, frequently, for fourteen days, strain, and filter. Dose, from twenty to forty drops. Said not to cause headache. *Nichol.*

VINEGAR OF OPIUM. BLACK DROP.

R. Powdered opium, eight ounces.
Coarsely-powdered nutmeg, one ounce and a half.
Saffron, half an ounce.
Sugar, twelve ounces.
Diluted acetic acid, sufficient.

Digest the opium, nutmeg, and saffron, with a pint and a half of diluted acetic acid, on a sand-bath, with a gentle heat, for forty-eight hours, and strain. Digest the residue with an equal quantity of diluted acetic acid, in the same manner, for twenty-four hours. Put the whole into a displacement apparatus, and return the filtered liquor as it passes, until it comes away clear. When the filtration has ceased, pour diluted acetic acid gradually on the residue in the filter, until the whole quantity of filtered liquor amounts to three pints. Lastly, add the sugar, and, by means of a water-bath, evaporate to three pints and four fl. ounces. *U. S. Ph.*

Dose, about six minims or ten drops, which are nearly equivalent to one grain of opium.

R. Opium, two ounces.
Nutmegs, grated, three drachms.
Saffron, one drachm.
Distilled vinegar, one pound.

Boil together for a quarter of an hour, then add an ounce of sugar, and half an ounce of yeast; let this mixture ferment for six weeks, strain, and evaporate to four fl. ounces. (*Sp. gr. 1.2.*) One ounce is equivalent to half an ounce of opium. Dose, one or two drops.

Cod. Hamb., 1845.

LANCASTER BLACK DROP.

R. Opium, half a pound.
Verjuice, three pints.
Bruised nutmeg, one ounce and a half.
Saffron, half an ounce

Boil to a proper consistence; add two ounces of yeast, and let stand in a warm place for six or eight weeks, and then in the open air till of the consistence of syrup, then decant, filter, and bottle; adding a little sugar to each bottle. Dose, six to ten drops. *Armstrong.*

HOULTON'S BLACK DROP.

R. Opium, two ounces and a half.
Diluted acetic acid, thirty-two ounces.
Digest for six days with a gentle heat, filter, and evaporate to an extract; macerate in Rectified spirit, five fl. ounces,
Distilled water, thirty-five ounces,
for eight days, and filter. *Beasley.*
About the strength of laudanum.

GUY'S HOSPITAL BLACK DROP.

R. Powdered opium, eight ounces,
Juice of crab apples, two pints,
Boil gently for half an hour, decant, and boil residue with one pint more of the juice, for a quarter of an hour; express, and strain; mix the two liquors, and add
Bruised nutmeg, one ounce.
Saffron, half an ounce.
Yeast, half a fl. ounce.

Ferment for some days; macerate for fourteen days; filter, and evaporate by a water-bath to consistence of thin syrup. Dose, two to ten minims. *Beasley.*

ROUSSEAU'S BLACK DROP.

R. Opium, four ounces.
Honey, twelve ounces.
Hot water, five pounds.
Yeast, two drachms.

Dissolve the opium and honey separately in the hot water, mix, and add the yeast; keep at about 86° F. for a month; express; filter, distil off sixteen ounces, and evaporate residue to ten ounces; add to it four and a half ounces of strong spirit; mix, and filter. Seven drops are equivalent to one grain of opium. *Beasley.*

PORTER'S BLACK DROP.

R. Opium, four ounces.
Citric acid, two ounces

Beat together in a mortar, and add

Boiling distilled water, one pint.
Triturate well together, let stand for twenty-four hours, and filter. Dose, from six to twenty-four drops. *Redwood.*

ACETATED TINCTURE OF OPIUM.

- R. Powdered opium, two ounces.
Vinegar, twelve fl. ounces.
Alcohol, half a pint.

Rub the opium with the vinegar, then add the alcohol, macerate for fourteen days, express, and filter through paper.

Dose, ten minims or twenty drops, which are equivalent to a grain of opium. *U. S. Ph.*

MIXTURE WITH BLACK DROP.

- R. Houlton's black drop, ten drops.
Spirit of nitric ether, half a drachm.
Distilled water, one ounce.

Mix. To be taken at once. *Foy.*

SEDATIVE MIXTURE.

- R. Opium, two grains.
Distilled vinegar, half an ounce.
Plantain water, six ounces.
Syrup of white poppy, one ounce.

Mix, and filter. In spoonful doses, in hæmoptysis with spasms. *Pierquin.*

WINE OF OPIUM.

- R. Powdered opium, two ounces.
Bruised cinnamon, one drachm.
" cloves, each, one drachm.
Sherry wine, one pint.

Macerate for fourteen days, agitating occasionally, express, and filter. *U. S. Ph.*

Dose, about twenty drops, equivalent to a grain of opium.

- R. Extract of opium, two ounces.
Cinnamon water, ten ounces.
Alcohol, two ounces.
White wine, four ounces.

Mix, and macerate for four days, and filter. *Brugnatelli.*

Much stronger than the last. Dose, ten to fifteen drops.

- R. Acetous extract of opium, six drachms.
Sherry wine, ten ounces.
Brandy, two ounces.

Mix; macerate for four days, and filter; each drachm contains four grains of opium. *Lalouette.*

LAUDANUM OF SYDENHAM.

- R. Opium, two ounces.
Saffron, one ounce.
Bruised cinnamon, one drachm.
" cloves, each, one drachm.
Sherry wine, one pint.

Infuse them together in a bath-heat for two or three days, till the tincture becomes of a due consistence, and after straining it off, set it by for use. Dose, sixteen or eighteen drops. *Rush's Sydenham, p. 155.*

COLLYRIUM OF WINE OF OPIUM.

- R. Decoction of flaxseed, four ounces.
Saffron, one drachm.
Wine of opium, one drachm.

Macerate the saffron in the flaxseed decoction, strain, and add the wine of opium. *Foy.*

- R. Acetate of copper, three grains.
Dissolve in

Rose water, eight fl. ounces;
and add,

Wine of opium, one fl. drachm.
In chronic ophthalmia. *Foy.*

FOMENTATION OF WINE OF OPIUM.

- R. Opium, one ounce.
Wine, two pints.

Boil down to one pint. As an anodyne application, in gouty and rheumatic pains, &c. *Pierquin.*

ODONTALGIC DROPS.

- R. Wine of opium,
Hoffmann's anodyne,
Oil of peppermint, equal parts.

Mix. In frictions on the cheek, and applied to carious teeth, on cotton. *Dobberan.*

MIXTURE OF WINE OF OPIUM.

- R. Wine of opium, ten drops.
Cinnamon water, one ounce.
Balm water, two ounces.
Tincture of castor, twenty drops.
Syrup of opium, half an ounce.

Mix. In spoonful doses, in uterine colic. *Augustin.*

- R. Cascarilla,
Columbo, each, two drachms.
Boiling water, sufficient

to obtain seven ounces of strained infusion.

Add to this

Wine of opium,
Sulphuric ether, each, twenty drops.
To be taken by degrees, in chronic diarrhœa. *Brera.*

TINCTURE OF OPIUM. LAUDANUM.

- R. Opium, powdered, two ounces and a half.
Diluted alcohol, two pints.

Macerate for fourteen days, and filter. *U. S. Ph.*

Dose, thirteen minims or twenty-five drops, equivalent to a grain of opium.

BATTLE'S SEDATIVE DROPS.

R. Hard extract of opium, three ounces.
Boiling distilled water, thirty ounces.
Dissolve, filter, when cold, and add

Rectified spirit, six ounces,
and water sufficient to make up two pints.

Cooley.

Dose, twenty drops.

SMITH'S CONCENTRATED LAUDANUM.

R. Denarcotized opium, four ounces.

Dissolve in alcohol, filter, evaporate to consistence of an extract, redissolve in water, and evaporate the filtered solution to twelve ounces; add

Rectified spirit, twenty-two drachms.
Distilled water, sufficient
to make up sixteen ounces. Dose, three to five drops.

Beasley.

COMPOUND TINCTURE OF OPIUM.

R. Extract of liquorice,
Opium, each, half an ounce.
Carbonate of potassa, one drachm.
Water, three pints.

Boil down to one pint, filter, and evaporate to twelve ounces; then add

Spirit of pimento, five fl. ounces.
Powdered cochineal, half a drachm.

Let rest for some time, and filter.

Med. Chirurg. Rev.

AMMONIATED TINCTURE OF OPIUM.

R. Benzoic acid,
Chopped saffron, each, six drachms.
Sliced opium, half an ounce.
Oil of anise, one drachm.
Spirit of ammonia, (Imp.) two pints.

Digest for seven days, and filter. *Ed. Ph.*

Also called *Scotch Paregoric*. Eighty minims should contain one grain of opium.

CAMPHORATED TINCTURE OF OPIUM.
PAREGORIC.

R. Powdered opium,
Benzoic acid, each, one drachm.
Oil of anise, one fl. drachm.
Clarified honey, two ounces.
Camphor, two scruples.
Diluted alcohol, two pints.

Macerate for fourteen days, and filter.

U. S. Ph.

Half a fl. ounce contains rather less than a grain of opium. Dose, for an infant, five to twenty drops; for an adult, one to two fl. drachms.

BATEMAN'S PECTORAL DROPS.

R. Diluted alcohol, four gallons.
Rasped red saunders, two ounces.

Digest for twenty-four hours, filter, and add

Powdered opium,
" catechu,
Camphor, each, two ounces.
Oil of anise, four fl. drachms.

Digest for ten days. About as strong as camphorated tincture of opium, or two grains of opium to the fl. ounce. *Phil. Coll. Pharm.*

GODFREY'S CORDIAL.

R. Tincture of opium, one pint and a half.
Sugar-house molasses, sixteen pints.
Alcohol, two pints.
Water, twenty-six pints.
Carbonate of
potassa, two ounces and a half.
Oil of sassafras, four fl. drachms.

Dissolve the carbonate of potassa in the water; add the molasses; heat over a gentle fire, till they simmer; remove the scum; add the laudanum and oil of sassafras, previously mixed together. *Phil. Coll. Pharm.*

Contains rather more than one grain of opium to the fl. ounce.

COMPOUND PILLS OF SOAP.

R. Powdered opium, half an ounce.
Soap, two ounces.

Beat with water into a pilular mass. Dose three to five grains. *U. S. Ph.*

TINCTURE OF OPIUM AND SOAP.

R. Opium, half an ounce.
Soap, two ounces.
Alcohol, sixteen ounces.

Digest for three days on a water-bath, filter, and dissolve in the liquor

Camphor, six drachms.
Oil of rosemary, one drachm.

Principally used in frictions, in pains in the limbs, &c., but also given internally, in doses of thirty to fifty drops, in wine. *Turin Ph.*

SUCCINATED TINCTURE OF OPIUM.

R. Opium, forty grains.
Camphor,
Amber, each, half a drachm.
Alcohol, six ounces.

Digest for four days on a water-bath, and filter. As a friction in spasmodic attacks. *Dumas*

SWEDIAUR'S TINCTURE OF OPIUM.

R. Extract of opium, one drachm.
 Distilled water, four drachms.
 Alcohol, half a drachm.
 Dissolve, and filter. Five drops are equivalent
 to a grain of opium. *Swediaur.*

WARNER'S TINCTURE OF OPIUM.

R. Opium,
 Soap, each, six drachms.
 Nutmeg, one drachm.
 Camphor, four drachms.
 Saffron, forty grains.
 Spirit of ammonia, nine ounces.
 Digest for ten days, agitating occasionally, then
 filter. *Van Mons.*

LETTSON'S ELIXIR.

R. Opium,
 Benzoic acid,
 Saffron, each, two drachms.
 Camphor, four scruples.
 Oil of anise, one drachm.
 Ipecacuanha,
 Balsam of tolu, each, half an ounce.
 Alcohol, two pounds.
 Macerate for ten days, and filter. Dose, five
 to twenty drops, in hooping-cough. *Augustin.*

CLYSTER OF LAUDANUM.

R. Tincture of opium, half a drachm.
 Infusion of flax-
 seed, two to four fl. ounces.
 Mix. *Ellis.*

R. Decoction of starch, four fl. ounces.
 Tincture of opium, thirty minims.

Mix. In obstinate vomiting, strangury from
 blisters, painful affections of the genito-urinary
 apparatus, tenesmus, &c. *Lond. Ph.*

CLYSTER OF LAUDANUM AND VALERIAN.

R. Tincture of opium, five to six drops.
 Infusion of valerian, three ounces.
 Mix, in spasms, in children. *Swediaur.*

OPIATE LINIMENT.

R. Olive oil, two fl. ounces.
 Tincture of opium, two fl. drachms.
 Solution of acetate of
 lead, half a fl. ounce.

Mix. *Ellis.*

R. Sulphuric
 ether, two fl. drachms and a half.
 Tincture of cam-
 phor, two fl. ounces and a half.
 Laudanum, half a fl. ounce.

Mix; as an embrocation in flatulent colic.

Ainslie.

R. Tincture of opium,
 Spirit of sulphuric ether,
 Tincture of camphor,
 each, two fl. ounces.

Mix. As an embrocation in rheumatism, tooth-
 ache, earache, &c. *Ellis.*

LINIMENT OF LAUDANUM AND LIME
WATER.

R. Tincture of
 opium, two fl. drachms and a half.
 Lime water, one fl. ounce.
 Oil of almonds, two fl. ounces.

Mix. To be applied on lint, four times a-day,
 on painful syphilitic pustules. *Phabus.*

LAUDANUM OINTMENT.

R. Tincture of opium, one drachm.
 Spermaceti ointment, two ounces.

Mix. As an application, morning and evening,
 to painful hemorrhoids. *Brera.*

BROWN MIXTURE.

R. Paregoric, one ounce.
 Antimonial wine, half an ounce.
 Powdered gum Arabic,
 Extract of liquorice,
 each, three drachms
 Water, six ounces.

Mix well. A tablespoonful, every three or four
 hours, in catarrh. *Dewees.*

OPIATE LINCTUS.

R. Tincture of opium, two fl. drachms.
 Diluted sulphuric
 acid, two fl. drachms and a half.
 Molasses, eight fl. ounces.
 Water, three fl. ounces.

Mix. A teaspoonful, occasionally, to quiet
 cough. *Beasley.*

ANODYNE DRAUGHT.

R. Camphor water, nine drachms.
 Nitrate of potassa, six grains.
 Comp. spirit of ether, one drachm
 Tincture of opium,
 ten to twelve minims

Syrup of poppies, two drachms
 Mix. To be taken at bedtime. *Copland*

R. Tincture of opium, twelve minims.
 Water, one fl. ounce.
 Pimento water, three drachms.
 Syrup of poppies, one drachm.

Mix. *Beasley.*

R. Tincture of
 opium, fifteen to twenty-five drops.
 Syrup of poppies, two fl. drachms.
 Spirit of cinnamon, one fl. drachm.
 Distilled water,
 one fl. ounce and a half.
 Mix. Ellis.

MIXTURE OF LAUDANUM AND TARTAR
 EMETIC.

R. Laudanum, one fl. drachm.
 Tartar emetic, four grains.
 Camphor water, eight fl. ounces.

Mix. In delirium tremens, and the advanced
 stages of low fevers. Dose, half a fl. ounce to
 one fl. ounce. Graves.

WISTAR'S COUGH LOZENGES.

R. Powdered gum Arabic,
 " extract of liquorice,
 " sugar, each, two drachms.
 " opium, six grains.
 Oil of anise, four drops.

Mix, and add
 Distilled water, sufficient.

Make into mass, and divide into sixty lozenges.
 One, three or four times a-day. Ellis.

ECLECTIC DOVER'S POWDER.

R. Powdered opium, half a drachm.
 " camphor, two drachms.
 " ipecacuanha, one drachm.
 Cream of tartar, one ounce.

Mix thoroughly. Dose, ten grains.
Am. Jour. Phar, 1854.

OPOPONAX.

OPOPONAX.

A gum resin obtained from the *Opoponax*
chironium, a tall, parsnip-like plant, a native of
 the warm countries of Europe and the Levant.

Ser. Syst. Pentand. digyn. *Nat. Syst.* Apia-
 cæ.

De Candolle, iv. 170. Griffith, Med. Bot. 323.

It occurs in tears and irregular lumps, or
 fragments, of a reddish-yellow color. Its odor
 is strong, peculiar, and unpleasant, and its taste
 bitter and acrid. It is seldom used in this
 country, but was formerly much esteemed in a
 variety of diseases. The dose is from ten to
 thirty grains.

TINCTURE OF OPOPONAX.

R. Opoponax, one part.
 Alcohol, five parts.
 Macerate for some days, and filter. Beral.

COMPOUND TINCTURE OF OPOPONAX.

R. Round birthwort,
 Long birthwort,
 Orris root, each, half an ounce.
 Opoponax,
 Sagapenum, each, two drachms
 Guaiacum, four scruples.
 Cloves, two drachms.
 Camphor, three drachms.
 Alcohol, ten ounces.

Macerate for twenty-four hours, and filter. As
 an application to foul venereal ulcers. Brera.

EMULSION OF OPOPONAX.

R. Opoponax,
 Soap, each, one drachm.
 Yolk of egg, two drachms.
 Syrup of worm-
 wood, one ounce and a half.
 Fennel water, three fl. ounces.

Make an emulsion. A teaspoonful every hour
 as a vermifuge. Bories.

ORIGANUM.

MARJORAM.

The article so designated in the U.S. Pharm.
 is the herb of the *Origanum vulgare*, a native
 of Europe, but extensively naturalized in the
 United States.

Ser. Syst. Didynam. gymnos. *Nat. Syst.* La-
 miacæ.

Linn. Sp. Pl. 834. Griffith, Med. Bot. 511.

The dried herb has an aromatic, agreeable
 odor, and a hot, pungent taste, depending on the
 presence of a volatile oil. It is somewhat tonic
 and stimulating, and has been used in the form
 of infusion as a diaphoretic and emmenagogue.
 The oil is employed in stimulating liniments.

OIL OF MARJORAM.

R. Marjoram, at will.
 Water, sufficient.

Mix, distil, and separate the oil in the receiver.

Much of the imported oil is obtained from
 another plant (*Thymus vulgaris*), which is frau-
 dulently substituted for the genuine. Dose, one
 to three drops. This enters into the composi-
 tion of the official opodeldoc.

ORYZA.

RICE.

The seed or grain of *Oryza sativa*, an annual
 plant, said to be a native of Ethiopia, but now
 extensively cultivated in most warm countries.

Ser. Syst. Hexand. digyn. *Nat. Syst.* Gra-
 minacæ.

Linn. Sp. Pl. 475. Griffith, Med. Bot. 660.
This grain as found in commerce is deprived of its cuticle, is white, dry, hard, inodorous, and of a farinaceous taste. It is very nutritive, and, in the form of a decoction, emollient.

RICE WATER.

R. Rice, well washed, two ounces.
Water, two quarts.
Boil for an hour and a half, then add sugar and flavoring, as may be required. *Ellis.*

R. Rice, five drachms.
Water, sufficient

to obtain a quart of decoction; add

Liquorice root, three drachms.
Let stand for some time, and strain, adding sugar, &c., at will. *Cottreau.*

RICE GRUEL.

R. Ground rice, one ounce.
Cinnamon, one drachm.
Water, one quart.
Boil for forty minutes, adding the aromatic near the close. Strain, and sweeten. *Ellis.*

MUCILAGE OF RICE.

R. Rice, one ounce.
Macerate it for three hours in
Tepid water, one quart.
Then boil slowly for an hour, and strain. *A. T. Thomson.*

RICE JELLY.

R. Rice, sufficient.
Macerate in as much water as will cover it, boil slowly, adding water as it evaporates, until the rice is reduced to a pap; sweeten and flavor, and pass through a fine sieve. On cooling, it becomes a moderately-consistent jelly. A good diet in dyspepsia, &c.

OVUM.

EGG.

OIL OF EGGS.

R. Yolks of eggs, at will.
Heat gently till the moisture is dissipated, introduce into a displacer, and exhaust by ether, distil the product in a water-bath, heat the residue, till the albuminous matter coagulates, then strain. *Par. Cod.*

Was, at one time, much esteemed as an application to hemorrhoids, chaps, excoriations, &c., but it is now seldom employed.

EMULSION OF EGGS.

R. Yolks of eggs, two.
Powdered sugar, one ounce.
Boiling water, fourteen ounces.
Triturate the eggs and sugar in a marble mortar, gradually adding the water. *Beral.*

EMOLLIENT CLYSTER OF EGGS.

R. Yolks of eggs, two.
Decoction of bran, one pound.
Mix. *Pierquin.*

RESTORATIVE CLYSTER OF EGGS.

R. Yolk of egg, one.
White wine, two fl. ounces.
Beef tea, without salt, eight fl. ounces.
Mix. *Radius.*

MIXTURE WITH EGGS.

R. Yolk of egg, one.
Cream, six fl. ounces.
Cinnamon barley-water, six fl. drachms.
Sugar, one ounce.
Mix. To be taken in teaspoonful doses, in the convalescence of children. *Foy.*

R. Yolk of egg, one.
Water, two pints.
Make an emulsion, and add
Common salt, half a drachm.
Mix. In mesenteric atrophy of children. *Hufeland.*

EGG AND BRANDY MIXTURE.

R. Brandy,
Cinnamon water, each, four fl. ounces.
Yolks of eggs, two.
Sugar, half an ounce.
Oil of cinnamon, two minims.
Mix. *Lond. Ph.*
As a stimulant in the sinking stage of fevers.

EGG AND WINE MIXTURE.

R. Yolks of eggs, two.
Oil of cinnamon, twenty drops.
Mix, and add
Madeira wine,
Cinnamon water, each, three fl. ounces.
Distilled water, two fl. ounces.
Sugar, two drachms.
Mix. Three or four tablespoonfuls for a dose. In convalescence from low fevers. *Ellis.*

LINIMENT OF EGGS.

R. Yolk of egg, one.
Flaxseed oil, two ounces.

Mix well. As an application to burns. *Radiis.*

R. White of egg, one.
Flaxseed oil, three ounces.

Mix well. As the last. *Mynsicht.*

CERATE OF EGGS.

R. Yolk of egg, one part.
Simple cerate, two parts.
Mix well. *Foy.*

R. Yolk of a hard-boiled egg, one.
Yellow wax, half an ounce.
Almond oil, one ounce and a half.

Melt the wax and oil together, and add the egg, triturating them thoroughly together. As an application to burns. *Soubeiran.*

P.

PAPAYER.

POPPY HEADS.

The ripe capsules of *Papaver somniferum*. These capsules are of a more or less globular form, crowned by a radiated, persistent stigma; of a light brown color, a papery and brittle texture, inodorous, but of a slightly bitter taste. They are analogous in properties, but in an inferior degree, to opium. They contain numerous small, oleaginous seeds, which are used in many preparations.

DECOCTION OF POPPY HEADS.

R. Poppy heads, sliced, four ounces.
Water, four pints.

Boil for a quarter of an hour, and strain.

Lond. Ph.

As a soothing and anodyne fomentation, in painful tumors and inflammations.

SYRUP OF POPPIES.

R. Poppy capsules, bruised,
and deprived of seeds, three pounds.
Sugar, five pounds.
Boiling water, five gallons.
Rectified spirit, five fl. ounces.

Boil the capsules in the water down to two gallons; press strongly. Then boil the strained liquor to four pints, and strain, while hot. Set aside for twelve hours, for the dregs to subside; boil the clear liquor down to two pints; in this dissolve the sugar, and lastly add the spirit.

Lond. Ph.

Dose, one fl. drachm, as a sedative and hypnotic.

R. Alcoholic extract of
poppy, four drachms.
Water, four fl. ounces.

Dissolve, and add

Boiling syrup, four pounds.
Evaporate to the proper consistence. *Par. Cod.*

R. Poppy heads, sixteen ounces.
Diluted alcohol, four pints.
Sugar, thirty ounces.

Deprive the heads of their seeds; bruise them thoroughly, macerate them in twice their weight of diluted alcohol for two days, express powerfully, add the remainder of the alcohol, and after twenty-four hours, again express. Evaporate the liquids to one pint, strain, and add the sugar, and dissolve by the aid of a gentle heat. *W. Procter.*

SUBSTITUTE FOR SYRUP OF POPPIES.

R. Sulphate of morphia, four grains.
Water, one fl. ounce.

Dissolve, and mix with

Syrup, fifteen fl. ounces.

Each fl. ounce contains one-quarter of a grain of the sulphate of morphia. *Wood.*

EXTRACT OF POPPY HEADS.

R. Poppy heads, bruised, fifteen ounces.
Boiling water, one gallon.

Macerate for twenty-four hours, boil down to four pints, strain, and evaporate over a vapor-bath, to proper consistence. *Ed. Ph.*

The dose is from five to ten grains.

PECTORAL SYRUP.

R. Dates, two pounds.
Jujube, one pound.
Liquorice root, half a pound.

Mallow root,

Maidenhair,

Poppy heads, each, four ounces

Water, sixteen pints

Boil, strain, and add

Sugar, eight pounds

Evaporate to the consistence of syrup. Dose, one to two ounces. *Gassicourt.*

SEDATIVE INJECTION.

- R. Simple emulsion, five fl. ounces.
Decoction of poppy heads, one pint.
White of egg, one drachm.

Mix. As an injection in acute gonorrhœa.
Gassicourt.

SEDATIVE MIXTURE.

- R. Flaxseed, two drachms.
Poppy head, one.
Water, sufficient

to obtain eight ounces of infusion; add

Yolk of egg, one.

Mix well. In painful diarrhœa. *St. Marie.*

PAREIRA.

PAREIRA BRAVA.

This is the root of *Cissampelos pareira*, a climbing shrub, with large, cordate, villous leaves, a native of the West Indies and South America.

Sex. Syst. Diœc. monand. *Nat. Syst.* Menispermaceæ.

Linn. Sp. Pl. 1473. *Griffith, Med. Bot.* 106.

The root, which is the officinal portion, as found in the shops, is usually in large, crooked pieces, of a dark color, externally, and yellowish within; it has at first a sweetish, and somewhat aromatic taste, but leaves a bitterish, unpleasant impression; the odor is very slight. It is tonic and alterative, and acts specifically on the bladder, diminishing irritability, and diminishing mucous secretion. The dose, in substance, is from thirty grains to a drachm.

INFUSION OF PAREIRA BRAVA.

- R. Pareira brava, six drachms.
Boiling water, one pint.

Macerate for two hours, and strain. *Edin. Ph.*

Dose, one to two fl. ounces, in irritation and chronic inflammation of urinary passages.

DECOCTION OF PAREIRA BRAVA.

- R. Pareira brava, four drachms.
Water, one pint and a half.

Boil to a pint, and strain. Dose, from eight to twelve fl. ounces, during the day. *Brodie.*

- R. Pareira brava, sliced, ten drachms.
Water, one pint and a half.

Boil to a pint, and strain. Dose, two fl. ounces.
Lond. Ph.

EXTRACT OF PAREIRA BRAVA.

- R. Powdered pareira brava, one pound.
Water, sufficient.

Mix the powder with half a pint of the water; introduce into a displacement apparatus, and

pour on water, till the powder is exhausted. Heat the fluid to the boiling point; strain, and evaporate to proper consistence. *Edin. Ph.*

Dose, ten grains to half a drachm.

TINCTURE OF PAREIRA BRAVA.

- R. Pareira brava, two ounces.
Diluted alcohol, one pint.

Digest for seven days, and filter. *Brodie.*

Dose, fifty to sixty drops

PAULLINIA.

PAULLINIA, (GUARANA)

Is a preparation from the seeds of *Paullinia sorbilis*, a climbing plant, indigenous to Brazil. The seeds are dried, pounded, mixed with cocoa and cassava, and with water, formed into a paste, which is dried. This preparation is of a brown color, hard, light, inodorous, and of a somewhat astringent taste. It is highly esteemed in Brazil, in diseases of the bowels and bladder, and has been used in Europe, as a tonic in these, and in chlorosis, &c., with much success.

Sex. Syst. Octand. trigyn. *Nat. Syst.* Sapindaceæ.

Martius, Mat. Med. Bras. 59.

LOZENGES OF PAULLINIA.

- R. Paullinia, five drachms and a half.
Vanilla sugar, one pound and a half.
Water, sufficient.

Mix, and form lozenges of ten grains each.

Gavrelle.

SYRUP OF PAULLINIA.

- R. Paullinia, two drachms and a half.
Simple syrup, two pints.

Mix. Dose, half a fl. ounce. *Gavrelle.*

PILLS OF PAULLINIA.

- R. Paullinia, sufficient.

Make into pills of a grain and a half. Five to ten, as occasion may require. *Gavrelle.*

ALCOHOLIC EXTRACT OF PAULLINIA.

- R. Powdered paullinia, at will
Alcohol, sufficient.

Introduce into a displacement apparatus, and pass the alcohol through till the powder is exhausted; distil off the spirit, and evaporate to proper consistence. Eight to ten grains, during the day. *Dechastelus.*

CHOCOLATE WITH PAULLINIA.

- R. Paullinia, one ounce.
Chocolate, sixteen ounces.

Mix, and form a paste. As a restorative, in diseases of debility, chlorosis, &c. *Beasley.*

PETROLEUM.

PETROLEUM.

A liquid bitumen, of a fluid consistence, of a brownish-black, or reddish-brown color, having a bituminous odor, and an acrid, strong taste. It is found in various parts of the world, issuing from the earth in the form of springs. Many of these exist in the United States, as on the shores of Seneca Lake, on the Kenhawa, &c. These products are light-colored and more liquid than the Barbadoes and other foreign varieties, and are known as Seneca oil.

Petroleum is a stimulating antispasmodic, and sudorific, with some anthelmintic properties. The dose is from half a drachm to a drachm.

BRITISH OIL.

R. Oil of turpentine,
 " flaxseed, each, eight fl. ounces.
 " amber, four fl. ounces.
 " juniper, four fl. drachms.
 Barbadoes petroleum, three fl. ounces.
 Seneca oil, one fl. ounce.

Mix. As a rubefacient liniment.
Phil. Coll. Pharm.

R. Oil of turpentine, eight fl. ounces.
 Barbadoes petroleum, four fl. ounces.
 Oil of rosemary, four fl. drachms.
 Mix. *Gray.*

EMBROCATION OF PETROLEUM.

R. Petroleum, half an ounce.
 Oil of turpentine, one drachm.
 Spirit of juniper, four ounces.

Mix. As a friction to the loins, in atony of the urinary passages and dropsy. *Radius.*

R. Petroleum, half an ounce.
 Spirit of lavender, one ounce.
 Laudanum, two drachms.

Mix. As a lotion to frozen limbs. *Phæbus.*

R. Petroleum,
 Alcohol, each, one ounce.
 Mix. *Saunders.*

R. Petroleum, one drachm and a half.
 Spirit of ammonia, two ounces.

Mix. As a remedy for chilblains. *Radius.*

OINTMENT OF PETROLEUM.

R. Petroleum, three drachms.
 Camphor, one scruple.
 Simple ointment, one ounce and a half.

Mix. *Radius.*

PLASTER OF PETROLEUM.

R. Resin, half an ounce.
 Camphor, one drachm.
 Opium, half a drachm.
 Petroleum, sufficient.

Mix. As an application in chronic rheumatism.
Radius.

ANTHELMINTIC MIXTURE OF PETROLEUM.

R. Petroleum, half an ounce.
 Tincture of assafetida, six drachms.

Mix. Forty drops, three times a-day, in cases of tape-worm. *Schwartz*

DIURETIC MIXTURE OF PETROLEUM.

R. Petroleum, two drachms
 Tincture of squill, one drachm.
 Spirit of juniper, six drachms.

Mix. Forty drops, three times a-day, in dropsy
Vogt.

PETROSELINUM.

PARSLEY.

Parsley, or *Petroselinum sativum*, is a biennial herbaceous plant, with a fusiform root; a native of southern Europe, but generally cultivated in our gardens, for the sake of its leaves, which are used for culinary purposes.

Sex. Syst. Pentand. digyn. *Nat. Syst.* Apiaceæ.

Hoffmann, Umb. 1, tom. 1. Griffith, Med. Bot. 318.

The root, which is the officinal portion, is fusiform, white, fleshy, of a pleasant odor and a somewhat sweetish, aromatic taste, becoming inert by drying; it should therefore be used in the recent state. It is diuretic and slightly aperient.

INFUSION OF PARSLEY ROOT.

R. Parsley root, one ounce.
 Boiling water, one pint.

Infuse and strain. A teacupful, with a drachm of sweet spirit of nitre, every three or four hours. *Ellis.*

In strangury from blisters; and highly spoken of by Dr. Chapman, in dropsies.

OIL OF PARSLEY.

R. Parsley seed, one part
 Water, four parts.
 Distil, and separate the oil. *Brunswick Ph.*

Carminative, and also said to be very efficacious in gonorrhœa, in doses of three or four drops, three times a-day.

PHLORIDZINA.

PHLORIDZINE.

A bitter principle obtained from the bark of the apple, pear, cherry, and plum-trees, especially from that of the root. It is white, very bitter, and is said to be a powerful febrifuge. It is obtained by boiling the fresh bark of the root for two hours in sufficient water to cover it, decanting the decoction, and renewing the process. The two decoctions are united, permitted to stand for twenty-four hours, when a deposit of phloridzine will take place, and, by evaporation, an additional portion will be obtained. This impure product is to be treated with distilled water and animal charcoal, to purify it.

POWDER OF PHLORIDZINE.

R. Phloridzine, ten to fourteen grains.
Sugar, one drachm.
Mix. In the apyrexia of intermittent fever.
De Koninck.

PHOSPHORUS.

PHOSPHORUS.

An elementary substance, usually in the form of thin sticks, which are translucent, of a pale-yellowish color, and flexible consistence. It has no perceptible taste, but an alliaceous odor. It is principally made from bone ashes, which consist mainly of phosphate of lime. It is a highly powerful, diffusible stimulant, and ought never to be given in substance, and its effects should be always closely watched. It has been given in cases of extreme prostration of the vital powers.

PHOSPHORATED OIL.

R. Phosphorus, one part.
Almond oil, sixteen parts.
Dissolve, and add

Oil of bergamot, sufficient to flavor.

Mix. Dose, five to ten drops in some mucilaginous liquid. *Magendie.*

R. Phosphorus, dried and sliced, six grains.
Oil of almonds, one ounce.

Digest by the aid of warm water and agitation, and when cold, carefully decant the oil from any undissolved phosphorus. Dose, five to fifteen drops. *Prus. Ph.*

Has been advised in gout, chronic rheumatism, &c., and also externally, as a friction.

PHOSPHORATED ETHER.

R. Phosphorus, one part.
Sulphuric ether, fifty parts.

Macerate for a month in a bottle covered with black paper; decant into small bottles similarly prepared. Dose, five to ten drops, in some emollient liquid, every four hours. *Par. Cod.*

R. Phosphorus, two grains.
Oil of peppermint, half a drachm.

Add to the solution

Sulphuric ether, half fl. ounce.
Mix well. Two to six drops, every four hours, on sugar. In epilepsy, paralysis, &c. *Augustin.*

PHOSPHORATED CERATE.

R. Phosphorated ether, five parts.
Simple cerate, twenty-four parts.
Mix. As a friction in obstinate cutaneous affections. *Foy.*

PHOSPHORATED OIL OF TURPENTINE.

R. Phosphorus, two grains.
Oil of turpentine, three fl. drachms.
Mix, and dissolve. Dose, ten to twelve drops, in obstinate intermittents; to be given in a cupful of oatmeal gruel. *Hufeland.*

PHOSPHORATED LINIMENT.

R. Phosphorus, six grains.
Oil of almonds, one ounce.
Caustic ammonia,
Camphor, each, two grains.
Mix. *Augustin.*

PHOSPHORATED OINTMENT.

R. Phosphorated ether, one drachm.
Lard, one ounce.
Rub together. *Guibourt.*
R. Phosphorus, ten grains.
Camphor, two scruples.
Lard, one ounce.
Mix carefully, and make an ointment. Used in palsies, by rubbing in twice daily. *Hufeland.*

PHOSPHORUS PASTE.

[RAT'S BANE.]

R. Phosphorus, eight parts.
Lukewarm water, } each, one hundred
Rye meal, } and eighty parts.
Melted butter, }
Sugar, one hundred and twenty parts.
Liquefy the phosphorus in the water; pour into a mortar, and add the meal; when cold, add the butter and sugar. An active poison for rats and mice. *Simon.*

PHYTOLACCA.

POKE.

Both the berries and the root of the Poke, or *Phytolacca decandra*, are officinal. It is a large, herbaceous plant, with a perennial root, and

bearing numerous clusters of dark purple berries.

Sex. Syst. Decand. decagyn. *Nat. Syst. Phytolaccaceæ.*

Linn. Sp. Pl. 631. *Griffith, Med. Bot.* 535.

The root, as found in the shops, is in transverse slices, of a light-brown color externally; and exhibiting on the cut surfaces numerous concentric rings. The taste is somewhat saccharine, followed by a sensation of acrimony; the odor of the fresh root is somewhat like that of ginseng, but this disappears on drying. It is emeto-cathartic, with some narcotic properties. It has been used as an alterative, in syphilis, rheumatism, and chronic eruptions. The berries have also been praised in the same complaints. The dose of the powdered root, is from one to five grains, as an alterative; from ten to thirty as an emetic.

OINTMENT OF POKE.

R. Powdered root, or leaves
of poke, one drachm.
Lard, one ounce.
Rub together. As an application in tinea capitis, psora, &c. *Wood.*

TINCTURE OF POKE-BERRIES.

R. Bruised poke-berries, four ounces.
Diluted alcohol, one pint.
Macerate for fourteen days, and filter. Dose, a teaspoonful, in chronic rheumatism.

PIMENTA.

PIMENTO. ALLSPICE.

This, which is also called *Jamaica pepper*, is the unripe berries of *Eugenia pimenta*, a beautiful evergreen tree, which is indigenous to the West Indies and South America.

Sex. Syst. Icosand. monog. *Nat. Syst.* Myrtaceæ.

De Candolle, Prod. iii. 285. *Griffith, Med. Bot.* 300.

The berries, which are the officinal part, are collected before they are ripe, and dried in the sun. In this state they are round, brown, somewhat rough, and a little larger than a peppercorn. They have an aromatic, agreeable odor, and a powerful, clove-like taste. Pimento is a warm, aromatic stimulant, but is more employed as a condiment than in medicine. The dose, in powder, is from ten to forty grains.

BOLUS OF ALLSPICE.

R. Powdered allspice,
" cinnamon, four grains.
" saffron, each, four grains.
" opium, one grain.
Conserve of roses, sufficient.
Mix, and make two boluses. In chronic diarrhoea. *Foy.*

WATER OF ALLSPICE.

R. Allspice, bruised, one pound.
Water, two gallons.
Distil a gallon. *Lond. Ph.*

Used as a carminative. Dose, one to two fl. ounces.

SPIRIT OF ALLSPICE.

R. Allspice, bruised, three ounces.
Diluted alcohol, one gallon.
Water, sufficient

to prevent empyreuma. Macerate for twenty-four hours; distil off a gallon. Used in same cases as other aromatic spirits, in doses of one to two fl. drachms. *Dub. Ph., 1826.*

R. Oil of pimento, two fl. drachms.
Diluted alcohol, one gallon.
Dissolve the oil in the alcohol. *U. S. Ph.*

TINCTURE OF ALLSPICE.

R. Allspice, one part.
Alcohol, five parts.
Macerate for fifteen days, and filter. Dose, from ten to twelve drops. *Swediaur.*

ESSENCE OF ALLSPICE.

R. Oil of pimento, one fl. ounce.
Rectified spirit, nine fl. ounces.
Mix with agitation. Dose, thirty drops, as a carminative. *Dub. Ph.*

PIPER.

BLACK PEPPER.

This article is the berries of *Piper nigrum*, a perennial, shrubby vine, from eight to twelve feet high, a native of India, and extensively cultivated in many parts of tropical Asia.

Sex. Syst. Diand. trigyn. *Nat. Syst.* Piperaceæ.

Linn. Sp. Pl. 40. *Griffith, Med. Bot.* 564.

The fruit, which is the officinal part, is gathered before it is quite ripe, and by the drying of the pulp over the grayish-white seed, becomes wrinkled and black; in this state it is known as *black pepper*; if permitted to ripen, and deprived of the skin by maceration, and dried, the fruit forms what is called *white pepper*, which is less active than the black.

Black pepper is a warm, carminative stimulant, but is principally used as a condiment. It has, however, been used in gonorrhœa, &c., and in the treatment of intermittent fevers, and also as an external application, in the form of ointment, to tinea capitis. The dose is from ten to twenty grains.

CONFECTION OF BLACK PEPPER.

- R. Powdered black pepper,
 " elecampane, each,
 " fennel seed, three pounds.

Honey,
 Sugar, each, two pounds.

Rub the first three ingredients and the sugar together, and keep in a covered vessel. When to be used, add the honey, and incorporate thoroughly, (or seven ounces of the powder with two ounces of the honey). *Lond. Ph.*

This, which is similar to *Ward's paste*, so celebrated in the cure of piles, must be used for a length of time, to be of service. Dose, from one to two drachms, two or three times a-day.

ELECTUARY OF BLACK PEPPER.

- R. Powdered black pepper,
 Conserve of orange-peel, each, one ounce.
 Syrup of orange peel, sufficient.

Make an electuary. As a stomachic and carminative. A drachm, twice a-day. *Saunders.*

RESINOUS OIL OF BLACK PEPPER.

- R. Black pepper, at will.

Grind it, and exhaust by means of alcohol (.835); distil off the alcohol. Treat the resinous residue with solution of potassa, which removes all but the piperine. The alcoholic liquid is then to be saturated with diluted muriatic acid, and the fluid oleo-resin that precipitates, is, after being washed with water, ready for use.

W. Procter.

FLUID EXTRACT OF BLACK PEPPER.

- R. Black pepper, in powder, one pound.
 Ether, sufficient.

Put the powder in a percolator, and pour ether gradually upon it until two pints of tincture pass. Distil off one pint and a half on a water-bath, and expose the residue in a shallow vessel till all the ether is evaporated, and the deposition of piperine has ceased. Lastly, separate the piperine by expression.

Dose, one or two drops. *U. S. Ph.*

VOLATILE OIL OF BLACK PEPPER.

- R. Powdered black pepper, at will.
 Water, sufficient.

Distil, and separate the oil from the product. This has the odor, but not the pungency of the fruit.

OINTMENT OF BLACK PEPPER.

- R. Powdered black pepper, four ounces.
 Lard, one pound.

Mix. As an application in *tinea capitis*.
Dub. Ph., 1826.

PLASTER OF BLACK PEPPER.

- R. Powdered black pepper,
 Galbanum, each, three drachms.
 Pitch, two ounces.
 Oil of laurel berries, sufficient.

Mix. As a stimulating plaster, in rheumatism, &c. *Augustin.*

CATAPLASM OF BLACK PEPPER.

- R. Flour of mustard, half a pound.
 Powdered black pepper,
 " ginger, each, one drachm.
 Boiling water, sufficient

to make a soft paste. To be applied to the pit of the stomach in colic, &c.; or to the feet, as a revulsive. *Ellis.*

PIPER LONGUM.

LONG PEPPER.

This is the dried spikes of the *Piper longum*, or *Chavica officinarum*, a climbing shrub, with diœcious flowers, a native of some of the East India islands, and cultivated in Sumatra and Java.

Linn. Sp. Pl. 41. Griffith, Med. Bot. 566.

The officinal portion is the fruit catkins, dried in the sun. These are aromatic; of a pungent, fiery taste; of a greyish-brown color; cylindrical. This pepper has nearly the same properties as the black, but it is little used in this country.

CATAPLASM OF LONG PEPPER.

- R. Powdered long pepper,
 " ginger, each, half an ounce.
 White of egg, sufficient.

Mix well into a paste. As a rubefacient application in pleurisy. *Foy.*

PIPERINUM.

PIPERINE.

A peculiar principle obtained from *Piper nigrum*. It is white, and crystallizes in four-sided prisms. It was thought to be an alkali, but the experiments of Pelletier show that it is more analogous to the resins. It has been much praised in Italy, as a febrifuge; and is also spoken of with favor by physicians of other countries. It has been supposed that its acrid taste and remedial properties were dependent on an admixture of acrid oil, but Dr. Christison attempts to prove that such is not the case.

R. Alcoholic extract of black pepper, at will.

Treat it with water, containing a hundredth of caustic potassa; wash the residue in cold water; dissolve in alcohol; filter, and permit to evaporate and crystallize. *Guibourt.*

Dose, from two to ten grains.

PILLS OF PIPERINE.

R. Piperine, twenty-four grains.
Crumb of bread, sufficient.

Mix, and make twelve pills. One, every two hours. *Meli.*

R. Piperine, twelve grains.
Extract of gentian, sufficient.

Mix, and make twelve pills. One, every hour, during the apyrexia of intermittents. *Ellis.*

PILLS OF PIPERINE AND MERCURY.

R. Blue pill, one grain.
Piperine,
Sulphate of quinia, each, two grains.
Syrup, sufficient.

Mix, and make a pill. *Hartle.*

One, three times a-day, for a few days, after the paroxysm has been checked by piperine.

TINCTURE OF PIPERINE.

R. Piperine, one part.
Alcohol, seven parts.
Dissolve, and filter. *Beral.*

PISCIDIA ERYTHRINA.

JAMAICA DOGWOOD.

This is a small West Indian tree, with a hard, heavy wood, and a narcotic bark, employed to intoxicate fish.

Sex. Syst. Diadelph. decand. *Nat. Syst.* Fabaceæ.

Linn. Sp. Pl. 993. *Griffith, Med. Bot.* 246.

The part used is the bark of the root, which appears to be a stimulating narcotic; causing excitement of the system, copious perspiration, and profound sleep. Applied topically, to allay pain.

TINCTURE OF JAMAICA DOGWOOD.

R. Bruised Jamaica dogwood, one ounce.
Alcohol, four fl. ounces.

Digest for seven days, and filter. Full dose, as a narcotic, one fl. drachm. *Hamilton.*

PIX BURGUNDICA.

BURGUNDY PITCH.

This is the prepared concrete juice of *Abies excelsa*, a lofty tree, a native of Europe and northern Asia.

23

Sex. Syst. Monœc. monadelph. *Nat. Syst.* Pinaceæ.

De Candolle, Fl. Fr. iii. 375. *Griffith, Med. Bot.* 606.

It is also obtained from *A. picea*.

When pure, it is hard, brittle, opaque, of a yellowish, or brownish-yellow color, and of a weak, turpentine-like taste and odor. The *Thus* of the London College, is the concrete juice of the *Spruce Fir*, and *Pinus palustris*, as it exudes. It is in solid, brittle tears, of a brownish-yellow color, and emits an agreeable odor when burned.

Both are used as ingredients in plaster.

PREPARED THUS OR FRANKINCENSE.

R. Frankincense, one pound.
Water, sufficient to cover.

Liquefy by heat, strain through a hair sieve, and when cold pour off the water. Used for making plasters. *Lond. Ph.*

PITCH PLASTER.

R. Burgundy pitch, two pounds.
Prepared Thus, one pound.
Resin,
Wax, each, four ounces.
Expressed oil of nutmeg, one ounce.
Olive oil,
Water, each, two fl. ounces.

Melt the thus, pitch, resin, and wax together, and add the olive oil, oil of nutmegs, and water; mix, and evaporate to a proper consistence. *Lond. Ph.*

As a rubefacient plaster in pectoral and rheumatic affections, &c. It often causes a serous discharge, and much irritation.

R. Burgundy pitch, six ounces.
Wax, half an ounce.
Turpentine, one drachm.
Melt, and mix. *Guy's Hosp.*

BURGUNDY PITCH PLASTER.

R. Burgundy pitch, six pounds.
Yellow wax, half a pound.

Melt, and stir constantly till they thicken. *U. S. Ph.*

WARMING PLASTER.

R. Burgundy pitch, three pounds and a half.
Cerate of Spanish flies, half a pound.

Melt together on a water-bath, and stir constantly till they thicken. *U. S. Ph.*

AROMATIC PLASTER.

R. Resin of spruce fir, three ounces.
Yellow wax, half an ounce.
Powdered cinnamon, six drachms.
Oil of pimento,
" lemon, each, two drachms.

Melt the resin and wax together, and strain. When they begin to thicken, on cooling, mix in the cinnamon, previously rubbed with the oils, and make a plaster. *Dub. Ph.*, 1826.

A good local stimulant, when applied to the region of the stomach, allaying nausea and vomiting, and relieving gastric uneasiness.

COMPOUND PITCH PLASTER.

R. Burgundy pitch, two parts.
White wax, four parts.
Turpentine,
Balsam of tolu, each, one part.
Mix and melt. *Beral.*

R. Burgundy pitch, twenty-four parts.
Resin,
Yellow wax, each, four parts.
Resin of spruce fir, twelve parts.
Oil of nutmeg, one part.
Melt together. *Lisbon Ph.*

PITCH CERATE.

R. Burgundy pitch, six parts.
Suet, four parts.
White wax, three parts.
Melt together. *Beral.*

PILLS OF BURGUNDY PITCH.

R. Burgundy pitch, five drachms.
Mucilage, sufficient.
Mix, and make one hundred pills. Six to eight, three times a-day, in cutaneous affections. *Ulrich.*

PIX CANADENSIS.

CANADA, OR HEMLOCK PITCH.

This is the prepared concrete juice of the *Abies Canadensis*, a lofty tree, found in the more northern parts of the United States, in Canada, &c., and also in the mountainous portions of the Middle States.

Sex. Syst. Monœc. monadelph. *Nat. Syst.* Pinacœ.

Mich. N. A. Sylv. iii. 185. Griffith, Med. Bot. 606.

The resin exudes spontaneously, and hardens on the bark, from which it is separated by boiling in water. It, however, requires a purification, by melting and straining, before it is fit for use. In this state, it is hard, brittle, of a dark yellowish-brown color, which becomes darker by exposure to the air, of a peculiar, but faint odor, and a scarcely perceptible taste.

It is a mild rubefacient, much resembling Burgundy pitch in its action and powers, and is employed for the same purposes.

The volatile oil obtained from the tops by distillation, and called the *oil of hemlock*, is possessed of emmenagogue properties. It has been used to produce abortion.

PIX LIQUIDA.

TAR.

This is an impure turpentine, obtained by the slow combustion of various species of the pine tribe, and more especially, of the *pinus palustris*. It is made in large quantities in North Carolina, and in various other places in America and Europe.

It is of a brownish-black color, of a tenacious consistence, has a peculiar empyreumatic odor, and a somewhat bitter, resinous, subacid taste.

Its properties are analogous to those of the turpentines. It is used both internally and externally. The dose is from half a drachm to one drachm, several times a-day.

TAR PILLS.

R. Tar, one drachm.
Powdered elecampane, sufficient.
Mix, form a mass, and divide into twenty pills. *Beasley.*

R. Tar, one drachm.
Wheat flour, sufficient.
Mix, form a mass, and divide into twenty pills. *G. B. Wood.*

R. Tar, two scruples.
Liquorice powder, one scruple.
Mix, and make sixteen pills. *Seymour.*

TAR WATER.

R. Tar, two pints.
Water, one gallon.
Mix, stirring with a wooden rod for fifteen minutes; when the tar has subsided, strain, and keep in well-stopped bottles. *Dub. Ph.* 1826

It is stimulant and diuretic, and was once much used in a variety of diseases, especially in those of the lungs. The dose is from one to two pints, in divided doses, in the course of the day. It is also used as a lotion in cutaneous diseases.

WINE OF TAR, OR TAR BEER.

R. Water, three quarts.
Wheat bran, one quart.
Tar, one pint.
Honey, half a pint.

Simmer together for three hours, and when cool, add a pint of brewers' yeast, and let it stand for thirty-six hours. Then bottle. Dose, a tablespoonful. *Duhamel.*

R. Ground malt,
Honey,
Tar, each, one pound.
Yeast, half a pint.
Water, sufficient.

Keep the malt, honey, and three quarts of water, at 150° F. for three hours, and when cool add

the yeast. Set aside for thirty-six hours, then decant, and to the clear solution add the tar, stirring constantly. Shake up occasionally for a week, then filter, and bottle for use.

W. Procter, Jr.

R. Strong beer, one gallon.
Tar, sufficient to saturate.

Mix, and allow them to macerate for three days, with occasional agitation. Dose, a tablespoonful.

Robinson.

TAR OINTMENT.

R. Tar, one pound.
Suet, each,

Melt the suet with a moderate heat, add the tar, and stir till cold.

U. S. Ph.

A stimulant application to various cutaneous eruptions, as psoriasis and tinea capitis.

COMPOUND TAR OINTMENT.

R. Tar ointment,
Cerate of subacetate of
lead, each, half a pound.

Mix. *St. Bart's Hosp.*

R. Tar ointment,
Sulphur ointment, equal parts.

Mix. *Guy's Hosp.*

PIX NIGRA.

BLACK PITCH.

This is the solid black mass that remains after the evaporation of tar. It has a shining fracture. It is gently stimulant and tonic, and has been used internally in some cutaneous diseases, and in piles. The dose is from ten grains to a drachm, in piles. It is also employed externally.

OINTMENT OF BLACK PITCH.

R. Black pitch,
Wax,
Resin, each, eleven ounces.
Olive oil, (Imp.) one pint.

Melt together, and strain. *Lond. Ph.*

Used for the same purposes as tar ointment.

R. Black pitch,
Resin,
Yellow wax, each, one part.
Olive oil, four parts.

Melt the two resins together, add the wax, and then the oil.

Guibourt.

R. Black pitch, sixteen parts.
Yellow wax, twelve parts.
Lard, ten parts.
Olive oil, two parts.

Melt together. *Pipon.*

PLASTER OF BLACK PITCH.

R. Black pitch, }
Resin, } each, two parts.
Suet, }
Turpentine, }
Yellow wax, three parts.

Melt together. *Niemann.*

R. Black pitch,
Resin,
Wax, each, eight parts.
Suet, one part.

Melt together. *Wirtem. Ph.*

PILLS OF BLACK PITCH.

R. Black pitch, one drachm.
Powdered gum Arabic, half a drachm.

Mix, and divide into twenty pills. Dose, two every night, in piles. *Wardleworth.*

PLATINUM.

PLATINA.

PLATINI BICHLORIDUM.

BICHLORIDE OF PLATINA.

Made by dissolving platinum in nitro-muriatic acid, and evaporating the solution to dryness by a gentle heat. *Beasley.*

It, as well as a concentrated solution, is of a deep red color. It is very deliquescent and soluble in water. It is poisonous in the dose of fifteen grains, but has been used as an alterative in syphilis, in doses of one-eighth to one fourth of a grain.

MIXTURE OF BICHLORIDE OF PLATINUM

R. Bichloride of platinum, two to four grains.
Sugar, two drachms
Distilled water, three fl. ounces.

Dissolve. Dose, a tablespoonful, three times a-day. *Hoefer.*

PILLS OF BICHLORIDE OF PLATINUM.

R. Bichloride of platinum, seven and a half grains.
Guaiaacum, one drachm.
Powdered liquorice root, sufficient.

Mix, and form twenty pills. One pill three times a-day. *Hoefer.*

OINTMENT OF BICHLORIDE OF PLATINUM.

R. Bichloride of platinum, one drachm
Extract of belladonna, two drachms.
Lard, four ounces.

Rub well together. As an application to indolent ulcers. *Hoefer.*

SODII CHLORO-PLATINAS.

CHLOROPLATINATE OF SODIUM.

Mix a solution of six parts of chloride of sodium with a solution of seventeen parts of bichloride of platina; evaporate, and crystallize. The crystals are of a deep yellow color, soluble in water and alcohol. They are similar in their medical properties to the analogous salt of gold. Dose, one grain. *Beasley.*

INJECTION OF CHLOROPLATINATE OF SODIUM.

R. Chloroplatinate of sodium, half a drachm.
Decoction of poppies, eight fl. ounces.
Dissolve. Used as an injection in gonorrhœa. *Hoefel.*

PLUMBUM.

LEAD.

A soft, bluish-grey, malleable metal, with a perceptible taste, and a peculiar smell when rubbed. It is not officinal in its metallic state, but its preparations are much employed, as sedatives and astringents.

PLUMBI ACETAS.

ACETATE OF LEAD.

POWDER OF ACETATE OF LEAD.

R. Acetate of lead,
Powdered opium, each, six grains.
Sugar, twenty-four grains.

Mix. Three grains, morning and evening, to check night sweats and diarrhœa, in phthisis. *Foy.*

R. Calomel, one or two grains.
Acetate of lead, half to one grain.

Mix, and divide into four powders. One, every three hours, in cholera infantum of infants. *Ellis.*

R. Acetate of lead, two to three grains.
Opium, a quarter of a grain.
Acetate of soda, three grains.
Sugar of milk, five grains.

Mix To be taken every hour, in hemorrhages. *Phœbus.*

PILLS OF ACETATE OF LEAD.

R. Acetate of lead,
Powdered mallow, each, one drachm.
Simple syrup, sufficient.

Mix, and make thirty-six pills. Four to five a day, to check sweats in phthisis. *Radius.*

R. Acetate of lead, half a drachm.
Calomel, five grains.
Conserve of roses, sufficient.

Mix, and make ten pills; one, every two to four hours, in hematemesis. *Ellis.*

R. Acetate of lead, thirty grains.
Powdered colchicum, twenty grains.
" opium, three grains.
Mucilage of gum Arabic, sufficient.

Mix thoroughly, and form into ten pills. One to be taken every six hours. Used in active hemorrhages, washed down with a draught of one drachm of distilled vinegar to one fl. ounce of water. Also, given in phthisis, after bleeding, one twice a-day. *A. T. Thomson.*

R. Acetate of lead, one scruple.
Opium, one grain.
Conserve of roses, sufficient.

Mix, and make twelve pills. One every hour at first, then every two hours, in cholera. *Graves.*

R. Acetate of lead, twelve grains.
Opium, six grains.
Conserve of roses, sufficient.

Mix, and make six pills. One to be repeated according to circumstances, in hemoptysis, &c. *Ellis.*

R. Acetate of lead, six grains.
Powdered mallow,
Extract of seneka, each, one drachm.

Mix, and make sixty pills. Two to five, several times a-day, in hemoptysis. *Koop.*

RUBEFACIENT PASTE.

R. Acetate of lead, one ounce.
Bi-sulphate of potassa, three ounces.
Water, sufficient.

Rub into a paste. It acts powerfully on the skin. *Clarus.*

COLLYRIUM OF SUBACETATE OF LEAD.

R. Solution of subacetate of lead, twelve drops.
Wine of opium, forty drops.
Rose water, four fl. ounces.

Mix. *Ellis.*

R. Acetate of lead, one scruple.
Lime water, twelve fl. ounces.
Catechu, three drachms.
Honey of roses, two ounces.

Dissolve, and filter. This is much more astringent than the last. *Augustin.*

SOLUTION OF SUBACETATE OF LEAD.

[GOULARD'S EXTRACT.]

R. Acetate of lead, sixteen ounces.
Powdered litharge, nine ounces and a half.
Distilled water, four pints.

Boil together, for half an hour, adding distilled water, so as to preserve the measure, filter, and keep in close-stopped bottles. *U. S. Ph.*

The *diluted solution*, or *lead-water*, is made by mixing two fl. drachms of the above solution to a pint of distilled water.

ACETATE OF LEAD CERATE.

Acetate of lead, four parts.
Soap, one part and a half.
Mix thoroughly in a heated mortar, and add
White wax, three parts.
Previously melted with
Olive oil, three parts.
Mix well. *Van Mons.*

PLASTER OF ACETATE OF LEAD.

R. Acetate of lead, one drachm.
Sulphur, half an ounce.
Resin, two ounces.
Melt together. As an application to venereal buboes. *Bories.*

PILE OINTMENT.

R. Acetate of lead, fifteen grains.
Burnt cork, half an ounce.
Fresh butter, two ounces.
Triturate well together. *Bories.*

ACETATE OF LEAD OINTMENT.

R. Acetate of lead,
Extract of belladonna, each, one part.
Lard, six parts.
Mix. As an application to fissures of the anus. *Foy.*

R. Acetate of lead, in fine powder, one ounce.
Simple ointment, twenty ounces.
Mix thoroughly. *Ed. Ph.*

SUBACETATE OF LEAD CATAPLASM.

R. Common cataplasm, four ounces.
Solution of subacetate of lead, one fl. ounce.
Muriate of ammonia, half a drachm.
Mix. As an application to indolent tumors. *Radius.*

SUBACETATE OF LEAD COLLYRIUM.

R. Solution of subacetate of lead, five drops.
Elder water, half a fl. ounce.
Laudanum, ten drops.
Mix. *Rust.*

SUBACETATE OF LEAD GARGLE.

R. Solution of subacetate of lead, half a fl. drachm.
Barley water, one pint.
Simple syrup, one ounce.
Mix. *Radius.*

SUBACETATE OF LEAD INJECTION.

R. Solution of subacetate of lead, two fl. drachms.
Distilled vinegar, eight fl. ounces.
Rose water, twenty-four fl. ounces.
Mix. In leucorrhœa. *Young.*

LINIMENT FOR MILK ABSCESS.

R. Olive oil, two ounces.
Lead water, one drachm.
Sulphuric ether, two drachms.
Laudanum, one drachm.
Mix. A rag moistened with this liniment to be frequently applied to the inflamed part. *Dewees.*

R. Acetate of lead, one drachm.
Distilled vinegar, two ounces.
Dissolve, and add
Alcohol, one ounce.
Distilled water, five ounces.
Mix. To be applied as above. *Clarke.*

ACETATE OF LEAD LOTION.

R. Diluted solution of subacetate of lead, one fl. ounce.
Spirit of camphor, three fl. ounces.
Water, one pint.
Mix. As an application to old ulcers. *Ellis.*

ACETATE OF LEAD CATAPLASM.

R. Crumb of bread, one pound.
Milk, four pounds.
Boil together, and add
Solution of subacetate of lead, one fl. ounce.
Spielmann.

VEGETO-MINERAL WATER OF GOULARD.

R. Solution of subacetate of lead, one part.
Alcohol (.921.), four parts.
Distilled water, forty-eight parts.
Mix. *Tadder.*
Used in fomentations, lotions, and cataplasms, and may always be replaced by the diluted solution of the subacetate.

ACETATE OF LEAD GARGLE.

R. Acetate of lead, sixty grains.
 Extract of opium, six grains.
 Rose water, eight fl. ounces.
 Syrup of mulberries, one ounce.
 Mix. *Pierquin.*

ACETATE OF LEAD MIXTURE.

R. Acetate of lead, half an ounce.
 Sulphate of iron, three drachms.
 Vinegar,
 Alcohol, each, two fl. ounces.
 Rose water, six fl. drachms.

Dissolve the acetate in the vinegar with a gentle heat; add the sulphate, and then the alcohol, mixed with the rose water. Highly praised in the sweats and colliquative diarrhoea of phthisis; in gonorrhoea and nocturnal emissions.

Gormann.

GOULARD'S BALSAM.

R. Oil of turpentine, at will.
 Heat, and gradually add
 Acetate of lead, sufficient.

Constantly stirring, till no more will dissolve; let rest, and decant while hot. Used as an application to eroding and painful ulcers.

Van Mons.

SUBACETATE OF LEAD CERATE.

[GOULARD'S CERATE.]

R. Solution of subacetate
 of lead, two fl. ounces and a half.
 White wax, four ounces.
 Olive oil, nine fl. ounces.
 Camphor, half a drachm.

Melt the wax, and mix with eight fl. ounces of the oil; remove from fire, and when it begins to thicken, gradually add solution of subacetate of lead; stir till cool, then add camphor, dissolved in rest of oil, and mix. *U. S. Ph.*

INJECTION OF SUBACETATE OF LEAD
AND LIME WATER.

R. Solution of subacetate
 of lead, thirty-four drops.
 Lime water, four fl. ounces.
 Olive oil, two fl. drachms.

Mix, and shake whenever used. In inflammation of the prostate and urethra. *Foy.*

LINIMENT OF SUBACETATE OF LEAD.

R. Solution of subacetate of lead,
 Laudanum,
 Honey of roses, each, two fl. drachms.
 Conserve of roses, one ounce.

Mix

Guy's Hosp.

SUBACETATE OF LEAD OINTMENT.

R. Cerate of subacetate of lead,
 Poplar ointment, each, one ounce.
 Powdered camphor, two drachms.

Mix. In inflamed acne and other cutaneous affections. *St. Marie.*

R. Solution of subacetate of lead,
 White wax, each, one drachm.
 Olive oil, one ounce.

Mix. As an application in ophthalmia. *Armstrong.*

SUBACETATE OF LEAD PLASTER.

R. Olive oil, fourteen fl. ounces.
 Yellow wax, twenty-four ounces.

Melt together, and add

Solution of subacetate
 of lead, one fl. ounce.
 Camphor, rubbed in a
 little oil, half a drachm.

Mix well, and spread on linen or skin. It has been much praised in rheumatic pains, and indolent ulcers. *Van Mons.*

PLUMBI CARBONAS.

CARBONATE OF LEAD.

[WHITE LEAD—CERUSE.]

CAMPHORATED POWDER OF CARBONATE
OF LEAD.

R. Carbonate of lead, ten drachms.
 Starch, two drachms.
 Sarcocolla,
 Gum Arabic,
 " tragacanth, each, one drachm.
 Camphor, half a drachm.

Triturate together. As an application to ulcers. *Giannini.*

OINTMENT OF CARBONATE OF LEAD.

R. Carbonate of lead, two ounces.
 Simple ointment, one pound.

Powder the carbonate very fine, add it to the ointment previously melted, and mix. *U. S. Ph.*

As an application to excoriated or ulcerated surfaces.

R. Carbonate of lead, six ounces.
 Camphor, one ounce.
 Olibanum, two ounces.
 Rose oil, three ounces.
 Whites of eggs, seven.

Triturate together. Same uses as above. *Harrer.*

- R. Carbonate of lead, one ounce.
 Litharge, two drachms.
 Armenian bole, one ounce.
 Honey of roses, half an ounce.
 Lard, sufficient.

Triturate together. As a dressing to phagedenic and cancerous ulcers. *Swedjaur.*

- R. Carbonate of lead, one ounce.
 Opium, one drachm.
 Lard, half an ounce.
 Anodyne balsam, sufficient.

Mix. As an application in facial neuralgia. *Fouquier.*

CAMPHORATED CARBONATE OF LEAD OINTMENT.

- R. Carbonate of lead, one ounce.
 Simple ointment, five ounces.
 Camphor, two scruples.

Mix. *Beasley.*

PLASTER OF CARBONATE OF LEAD.

- R. Carbonate of lead, one pound.
 Olive oil, two pounds.
 White wax, three ounces.
 Water, two pounds.

Melt the wax and oil together, add the carbonate of lead, and stir till mixed; then add the water and boil; malaxate, and melt again.

Par. Cod.

- R. Carbonate of lead, one pound.
 Olive oil, two pints.
 Yellow wax, four ounces.
 Lead plaster, one pound and a half.
 Powdered orris root, nine ounces.

Boil together the oil and carbonate of lead, adding a little water, and constantly stirring, till they are perfectly incorporated, then add the wax and plaster; when these are melted, mix in the orris, and stir well.

U. S. Ph., 1830.

As a substitute for *Mahy's plaster*, and a good application to inflamed and excoriated surfaces, as bed sores, &c.

PLUMBI CHLORIDUM.

CHLORIDE OF LEAD.

- R. Acetate of lead, nineteen ounces.
 Chloride of sodium, six ounces.

Dissolve the two salts separately, the former in three pints of boiling, distilled water, the latter in one pint of distilled water; mix the solutions, and wash the precipitate with distilled water.

Lond. Ph., 1836.

OINTMENT OF CHLORIDE OF LEAD.

- R. Chloride of lead, one drachm.
 Simple cerate, one ounce.
 Mix. As an application in cancerous ulcerations. *Tuson.*

LOTION OF CHLORIDE OF LEAD.

- R. Chloride of lead, one drachm.
 Water, one pint.
 Dissolve: as a wash to cancerous ulcerations, and in painful neuralgic tumors. *Tuson.*

PLUMBI IODIDUM.

IODIDE OF LEAD.

- R. Solution of iodide of potassium,
 " acetate of lead, each, sufficient.

Add one solution gradually to the other, till there is no longer a precipitate, wash this in cold water, and dry it. *Cottureau.*

- R. Iodide of potassium,
 Nitrate of lead, each, one ounce.
 Water, a pint and a half.

Dissolve the salts separately, each in one-half of the water; mix the solutions, collect the precipitate on a filter of linen or muslin, and wash it with water. Boil the powder in three gallons of water, acidulated with three fl. ounces of pyroligneous acid. Let any undissolved matter subside, maintaining the temperature near the boiling point, and pour off the clear liquor, from which the iodide of lead will crystallize on cooling. *Ed. Ph.*

This is a preferable formula to that made with the acetate of lead, as above, or that of the *Lond. Ph.*

PILLS OF IODIDE OF LEAD.

- R. Iodide of lead, four grains.
 Conserve of roses, one scruple.

Mix, and form twelve pills. One, night and morning, gradually increased, in scrofulous affections. *Foy.*

- R. Iodide of lead, half a drachm.
 Conserve of roses, sufficient.

Mix, and form one hundred and twenty pills Administered as above. *Gassicourt*

POWDER OF IODIDE OF LEAD.

- R. Iodide of lead,
 Powdered foxglove, each, six grains.
 Alcoholic extract of
 stramonium, two grains
 Sugar, one drachm.

Mix, and divide into twenty-four powders. One four times a-day. *Gassicourt*

OINTMENT OF IODIDE OF LEAD.

- R. Iodide of lead, one ounce.
Lard, eight ounces.

Mix. As an application to indolent and scrofulous swellings. *Lond. Ph.*

PLUMBI OXIDUM SEMIVITREUM.

LITHARGE.

LEAD PLASTER. DIACHYLON.

- R. Litharge, in fine powder, five pounds.
Olive oil, one gallon.
Water, two pints.

Boil together over a gentle fire, stirring constantly, till a plaster is formed. *U. S. Ph.*

LEAD CERATE.

- R. Lead plaster, six parts.
White wax, half a part.
Melt, mix, and add
Olive oil, two parts and a half.
Van Mons.

LEAD OINTMENT.

- R. Olive oil, eight parts.
Lard, }
Butter, }
Mutton fat, } each, two parts.
Yellow wax, }
Litharge, }

Heat the oil, fats, and wax together, till they emit fumes; gradually add the litharge; stir constantly, till the mixture assumes a dark-brown color, and add

- Pitch, one part.
Strain. *Foy.*

This preparation is much used in France, under the name of *Onguent de la Mere*, as a maturant to buboes, abscesses, &c.

COMPOUND LEAD OINTMENT.

- R. Lead plaster, three pounds.
Olive oil, eighteen fl. ounces.

Melt together, and add
Prepared chalk, six ounces.
Diluted acetic acid, six fl. ounces,
and stir till cold. *Lond. Ph.*

- R. Fresh root of figwort, three ounces.
Leaves of henbane, three handfuls.
Fresh butter, one pound and a half.
Litharge, two ounces.
Yolks of eggs, two.

Boil the root and leaves in the butter, till all moisture is driven off; express, and strain; add the litharge: stir till cold, and add the yolks of eggs, and incorporate well. *Saxon Ph.*

Celebrated as an application to painful hemorrhoids.

DIAPALMA PLASTER.

- R. Lead plaster, eight pounds.
White wax, eight ounces.

Melt by a gentle heat, and add
Sulphate of zinc, four ounces,
dissolved in a little water; continue the heat, constantly stirring till all moisture is evaporated. *Guibourt.*

GAUTHIER'S PLASTER.

- R. Diapalma plaster, twelve parts.
Olive oil,
White wax, each, one part.
Turpentine, two parts.

Melt the first three articles together, with a gentle heat, and add the turpentine. *Guibourt.*

BAYNTON'S ADHESIVE PLASTER.

- R. Lead plaster, one pound.
Resin, six drachms.
Melt together, and spread on muslin. *Beasley*

LEAD CATAPLASM.

- R. Flaxseed meal,
Arrowroot, each, two ounces.
Water, eleven fl. ounces.
Lead ointment, one ounce.

Mix, and heat to a proper consistence. As a maturing application. *Beral.*

BANYER'S OINTMENT.

- R. Litharge, two ounces.
Burnt alum,
Calomel, each, one ounce and a half.
Lard, two pounds.
Venice turpentine, half a pound.

Mix. As an application in porrigo. *Cazenave & Schedel.*

PLUMBI OXIDUM RUBRUM.

RED LEAD.

NUREMBERG PLASTER.

- R. Red lead, eight ounces.
Olive oil, one pound.

Mix, and heat till it assumes a brown color, and add

Resin, half an ounce.
Yellow wax, one ounce and a half.
Camphor, two drachms.
Mix, and stir till well incorporated.

Hessian Ph.

PLUMBI SACCHARAS.

SACCHARATE OF LEAD.

R. Mix one part of sugar with two parts of nitric acid, diluted with ten of water, and apply heat as long as any reaction takes place; neutralize with chalk, filter, and add to the filtered solution acetate of lead as long as any precipitate is found; wash this, and dry it.

Beasley.

NITRO-SACCHARATE OF LEAD.

R. Saccharate of lead, at will.
Dissolve in nitric acid, diluted with nineteen parts of water, filter, evaporate, and set aside to crystallize.

Hoskins.

SOLUTION OF NITRO-SACCHARATE OF LEAD.

R. Nitro-saccharate of lead, five grains.
Saccharic acid, twenty-five drops.
Water, five ounces.
Dissolve. Proposed as a solvent for phosphatic calculi.

Hoskins.

PLUMBI TANNAS.

TANNATE OF LEAD.

R. Strong infusion of galls, at will.
Solution of acetate of lead, sufficient.
Add the solution of the acetate, drop by drop, till it no longer causes a precipitate; wash this, and dry it.

Fantonetti.

LINIMENT OF TANNATE OF LEAD.

R. Decoction of oak bark, eight fl. ounces.
Solution of acetate of lead, sufficient.
Add the solution of lead to the decoction by degrees, till it causes no precipitate; wash the precipitate, and add
Alcohol, two drachms.
Recommended in bed sores.

Sundelin.

OINTMENT OF TANNATE OF LEAD.

R. Recent tannate of lead, twelve parts.
Lard, thirty parts.
Mix. As an application to bed sores.

Tott

R. Decoction of oak bark, six ounces.
Solution of acetate of lead, one ounce and a half.

Mix. Collect the precipitate, and mix whilst moist,

Lard, one ounce.
Camphor, ten grains.

As an application to excoriations.

Sundelin.

PODOPHYLLUM.

MAY APPLE.

This is the root or rhizome of the *Podophyllum peltatum*, a native, herbaceous plant, growing in damp situations, having two large peltate, palmate leaves.

Sex. Syst. Polyand. monog. *Nat. Syst.* Berberidaceæ.

Linn. Sp. Pl. 722. *Griffith, Med. Bot.* 115.

The dried root is about as thick as a quill, of a blackish-brown color externally, and somewhat corrugated; within, it is of a dirty white; it has a faint, but unpleasant odor, and a bitterish, somewhat sweetish taste. It is a purgative of the same character as jalap, and may be given in the same combinations. Dose, ten grains to a scruple.

EXTRACT OF MAY APPLE.

R. May-apple root, in coarse powder, one pound.
Alcohol, four pints.
Water, sufficient.

Macerate the May-apple root for four days in the alcohol, and then obtain four pints of tincture by displacement with the water; then, by same means, obtain six pints of infusion; distil off the alcohol from tincture, and evaporate the infusion, till each are of the consistence of thin honey. Mix, and evaporate to a due consistence. Dose, five to fifteen grains.

U. S. Ph.

PODOPHYLLIN.

R. May-apple root, one pound.
Alcohol,
Animal charcoal, in coarse powder, each, a sufficient quantity.

Reduce the May-apple root to powder, moisten it with half its weight of alcohol, introduce into a displacer, and pour on alcohol until four pints of tincture are obtained. Distil off the alcohol, treat the residual extract with ether to remove fixed oil, boil the residue with water, and dissolve what remains in alcohol, 42° B.; treat it with purified animal charcoal, and allow the decolorized solution to evaporate spontaneously. The dry, amorphous mass, is podophyllin, and, in cathartic power, six grains of it are equal to eight of jalap resin.

J. R. Lewis

An impure podophyllin has been prepared by W. S. Merrill, by precipitating the resin from a concentrated alcoholic tincture, by the addition of water.

PILLS OF PODOPHYLLIN AND MAY APPLE.

R. Podophyllin, ten grains.
Extract of May apple, half a drachm.
Mix and divide into ten pills. Dose, one or two pills. *Thomas.*

PILLS OF PODOPHYLLIN AND IRON.

R. Podophyllin,
White turpentine, each, half a drachm.
Carbonate of iron, one drachm.
Mix well, and divide into thirty pills. *Ecl. Med. Jour.*

COMPOUND PILLS OF PODOPHYLLIN.

R. Podophyllin,
Scammony,
Gamboge, each, in powder, one drachm.

Rub together for half an hour, and add
Castile soap, half a drachm.
Beat into a mass, and divide into one hundred and twenty pills. *Ecl. Med. Jour.*

POTASSIUM.

POTASSIUM

Is a soft, silver-white metal, readily oxidizable by the action of the air; when thrown on water, it takes fire and burns with a rose-colored flame, combining with oxygen, and forming potassa which is dissolved in the water. It must be kept in liquids, into whose composition no oxygen enters, as naphtha, &c. It is not used in medicine, but its numerous compounds are highly important.

R. Fused carbonate of potassa, one pound.
Iron filings, three quarters of a pound.
Powdered charcoal, one pound.

Mix, and introduce into a gun-barrel or iron bottle, placed in a furnace so made that the flame of a very strong fire may surround it, and maintain every part at a uniform heat approaching to whiteness; the gun-barrel or bottle is to be connected by an iron tube with a copper receiver partly filled with naphtha and surrounded by ice; the heat is to be gradually raised until the requisite temperature is reached, when the potassium will distil over. *Curaudau.*

POTASSII BROMIDUM.

BROMIDE OF POTASSIUM.

R. Bromine, two ounces.
Iron filings, one ounce.
Carbonate of potassa, two ounces and one drachm.
Distilled water, four pints.

To a pint and a half of the water, first add the iron and then the bromine. Set them aside for half an hour, occasionally stirring. Apply a gentle heat, and when the liquid becomes greenish, add the carbonate of potassa dissolved in a pint and a half of the water. Strain, and wash what remains in a pint of boiling distilled water, and again strain. Evaporate the mixed liquors to crystallization. *U. S. Ph.*

Alterative and resolvent; used both externally and internally in bronchocoele and scrofula, in doses of two to ten grains, three times a-day.

SOLUTION OF BROMIDE OF POTASSIUM.

R. Bromide of potassium, twelve grains.
Lettuce water, three ounces.
Syrup of marsh mallow, one ounce.
Mix. To be given in tablespoonful doses, in the twenty-four hours, in scrofula. *Foy.*

R. Bromide of potassium, one to two scruples.
Syrup of orange peel, one fl. ounce.
Water, three fl. ounces.
Mix. Dose, a tablespoonful. *Ellis.*

OINTMENT OF BROMIDE OF POTASSIUM.

R. Bromide of potassium, thirty-four grains.
Lard, one ounce.
Mix. In frictions on scrofulous swellings, and to tinea and tetter. *Magendie.*

COMPOUND OINTMENT OF BROMIDE OF POTASSIUM.

R. Bromide of potassium, twenty-four grains.
Liquid bromine, six to twelve drops.
Lard, one ounce.
Mix. *Magendie.*

POTASSII CHLORIDUM.

CHLORIDE OF POTASSIUM.

R. Carbonate of potassa, at will.
Dissolve in a sufficient quantity of water, filter, and add muriatic acid to saturation; concentrate the solution by evaporation, and cool slowly, that crystals may form. *Guibourt.*

Formerly known as the *Salt of Sylvius*, and used in fever to excite perspiration, and as a diuretic, in doses of from one to two scruples.

SOLUTION OF CHLORINATED POTASSA.

R. Carbonate of potassa, one part.
Water, ten parts.

Dissolve, and pass a current of chlorine gas through the solution, till it is saturated, and preserve in well-stopped bottles. This is known as *Eau de Javelle*, and is principally used for bleaching, but may be employed as a disinfectant, like Labarraque's liquid. *Guibourt.*

POTASSII CYANURETUM.

CYANURET OF POTASSIUM.

R. Ferrocyanuret of potassium, dried, eight ounces.
Carbonate of potassa, dried, three ounces.

Mix the salts and throw the mixture into a deep iron crucible, heated to redness; keep up the temperature till effervescence ceases, and the fused mass concretes of a white color upon a warm glass rod dipped into it. Then pour the liquid carefully into a shallow dish to solidify, stopping before the salt becomes contaminated with the precipitated iron. Break up the mass while yet warm, and preserve it in well-stopped bottles. *U. S. Ph.*

Dose, one-eighth of a grain.

R. Ferrocyanuret of potassium, dried, eight ounces.
Pure carbonate of potassa, three ounces.

Charcoal, in fine powder, one ounce.

Mix well, fuse in an iron vessel until effervescence ceases, and the fusion is tranquil; pour out the fused mass on an iron plate, and cool. Powder when cool, dissolve in alcohol (.900), and separate and dry the crystals which form on cooling, and preserve them in well-closed bottles. *Wöhler.*

PILLS OF CYANURET OF POTASSIUM.

R. Cyanuret of potassium,
Starch mixed with syrup of gum, each, half a grain.

Mix, and make two pills. Dose, one morning and evening. In spasmodic affections of the chest. *Foy.*

SOLUTION OF CYANURET OF POTASSIUM.

R. Cyanuret of potassium, twenty-two grains.
Diluted alcohol, nine fl. drachms.

Dissolve. Equal to hydrocyanic acid. *Laming.*

MIXTURE OF CYANURET OF POTASSIUM.

R. Cyanuret of potassium, seven grains and a half.
Distilled water, one pint.
Sugar, one ounce and a half.

Mix. A tablespoonful, night and morning, in pectoral affections. *Magendie.*

R. Cyanuret of potassium, four grains.
Distilled water, two ounces.
Syrup, three drachms.

Mix. A teaspoonful, every three or four hours, in asthma and chronic catarrh. *Cadet.*

R. Tincture of castor,
Musk,
Nitrate of potassa, each, four grains.
Cyanuret of potassium, two grains.
Linden-flower water, eight fl. ounces.

Mix. To be taken in divided doses, in twenty-four hours, in chorea. *Fouquier.*

R. Lettuce water, two ounces.
Cyanuret of potassium, half a grain to a grain and a half.

Syrup of marsh mallow, one ounce.
Mix. A tablespoonful, every two hours. *Magendie.*

SYRUP OF CYANURET OF POTASSIUM.

R. Cyanuret of potassium, seven grains and a half.
Simple syrup, one pound.

Mix. *Magendie.*

LOTION OF CYANURET OF POTASSIUM.

R. Cyanuret of potassium, one to four grains.
Distilled water, one ounce.

Mix. *Foy.*

R. Cyanuret of potassium, eight grains.
Distilled water, one ounce.

Mix. To be applied by means of compresses, in cases of neuralgia, articular rheumatism, &c. *Foy.*

R. Cyanuret of potassium, ten grains.
Emulsion of bitter almonds, six ounces.

Mix. As an application to chronic eruptions attended with much itching. *Cazenave.*

OINTMENT OF CYANURET OF POTASSIUM.

R. Cyanuret of potassium, one to four grains.
Lard, one ounce.

Mix. *Lombard.*

R. Cyanuret of potassium, twelve grains.
 Oil of almonds, two drachms.
 Cold cream, two ounces.
 Mix. To be rubbed on the skin to relieve
 neuralgic pains. *Cazenave.*

POTASSII IODIDUM.

IODIDE OF POTASSIUM.

R. Potassa, six ounces.
 Iodine, in powder, sixteen ounces.
 Charcoal, " two ounces.
 Boiling water, three pints.

Dissolve the potassa in the water, add the iodine gradually until in excess, stirring after each addition till the solution becomes colorless. Evaporate to dryness, stirring in the charcoal towards the close; rub to powder and heat to dull redness, maintaining the temperature for fifteen minutes. Cool, dissolve in water, filter, evaporate and crystallize. The mother liquor yields more crystals by evaporation. *U. S. Ph.*

R. Iodine, sixteen ounces.
 Distilled water, eight pints.
 Sulphuret of barium, sufficient.
 Sulphate of potassa, twelve ounces.

Mix the iodine with the water, and gradually add the powdered sulphuret, until the solution becomes colorless, constantly stirring. Heat to the boiling point, add the sulphate of potassa, boil for a quarter of an hour, and filter. Evaporate to crystallization. *Mohr.*

R. Iodine, one hundred grains.
 Carbonate of potassa, seventy-five grains.
 Water, two drachms.
 Iron filings, thirty grains.

Mix, and heat slightly, then to redness; the resulting red powder is to be treated with water; filtered, and evaporated to dryness. *Pypues.*

Dose, two to fifteen grains, but has been given in still larger quantities.

PILLS OF IODIDE OF POTASSIUM.

R. Iodide of potassium, five drachms.
 Distilled water, six drachms.
 Crumb of bread, sufficient.

Mix, and make three hundred pills. In goitre, leucorrhœa, &c. *Pierquin.*

R. Iodide of potassium, fifteen grains.
 Burnt sponge,
 Extract of bittersweet, each,
 five drachms.
 Distilled water, sufficient.

Mix, and make one hundred and eighty pills. Six, twice a-day, in scrofula and goitre. *Vogt.*

LOZENGES OF IODIDE OF POTASSIUM.

R. Iodide of potassium, one drachm.
 Sugar, three ounces.
 Mucilage of tragacanth, sufficient.

Mix, and make lozenges of twelve grains. One to six, or more, a-day. *Giordano.*

R. Iodide of potassium, seventy-five grains.
 Powdered coffee, half a drachm.
 " sugar, four ounces.
 Mucilage of gum, made with
 a strong infusion of coffee, sufficient.

Mix, and make three hundred lozenges. Each contains a quarter of a grain of the iodide. In goitre, scrofula, &c. *Pierquin.*

SOLUTION OF IODIDE OF POTASSIUM.

R. Iodide of potassium, thirty-six grains.
 Distilled water, one ounce.

Dissolve. Dose, ten minims. *Magendie.*

R. Iodide of potassium, one drachm.
 Distilled water, one ounce.

Dissolve. Fifteen drops, gradually increased, three times a-day, in ovarian dropsy. *Elliotson.*

R. Iodide of potassium, four ounces.
 Distilled water, eight fl. ounces.

Dissolve. Ten minims contain five grains of the iodide. *Guy's Hosp.*

INJECTION OF IODIDE OF POTASSIUM.

R. Iodide of potassium, one to three grains.
 Distilled water, one pint.

Dissolve. To stimulate fistulous sinuses in scrofulous patients. *Foy.*

MIXTURE OF IODIDE OF POTASSIUM.

R. Iodide of potassium, one to four drachms.
 Lettuce water, eight fl. ounces.
 Mint water, two fl. drachms.
 Syrup of marsh mallow, one fl. ounce.

Mix. *Foy.*

R. Iodide of potassium, four drachms.
 Lettuce water, eight fl. ounces.
 Orange-flower water, one fl. drachm.
 Tincture of fox-glove, one to two fl. drachms.
 Syrup of marsh mallow, one fl. ounce and a half.

Mix. *Magendie.*

In hypertrophy of the heart. A teaspoonful, morning and evening.

R. Iodide of potassium, two drachms.
Distilled water, sixteen fl. ounces.
Syrup, two fl. ounces.
Mix. Two or three tablespoonfuls a-day.

Cazenave.

R. Iodide of potassium, half a drachm.
Syrup of ginger, one fl. ounce.
Water, five fl. ounces.

Mix. Dose, a tablespoonful, three times a-day.

Ellis.

R. Quassia,
Gentian, each, one drachm.
Boiling water, sixteen fl. ounces.

Macerate for an hour, strain, and add

Iodide of potassium, thirty-six grains.
Carbonate of potassa, two drachms.

Mix. A tablespoonful, three times a-day.

Cumming.

R. Iodide of potassium, three grains.
Medicinal hydro-
cyanic acid, ten to twelve drops.
Lettuce water, four fl. ounces.
Syrup of marsh mallow, one fl. ounce.

Mix. A teaspoonful every hour, in phthisis.

Magendie.

R. Iodide of potassium, two grains.
Sulphate of magnesia, half an ounce.
Tartar emetic, half a grain.
Distilled water, six fl. ounces.

Mix. A teaspoonful, three or four times a-day, in scrofula.

Radius.

SYRUP OF IODIDE OF POTASSIUM.

R. Iodide of potassium, one drachm.
Simple syrup, one pound.

Dissolve.

Cadet.

R. Iodide of potassium, twenty grains.
Peppermint water, two fl. drachms.
Simple syrup, ten fl. ounces.

Mix.

Giordano.

R. Iodide of potassium, two drachms.
Simple syrup, six fl. ounces.

Mix.

Cazenave.

IODIDE OF POTASSIUM AND SARSAPARILLA.

R. Iodide of potassium, one drachm.
Decoction of sarsaparilla, two pints.
Syrup of orange-peel, two fl. ounces.

Mix. To be taken in glassfuls, during the day.

Magendie.

OINTMENT OF IODIDE OF POTASSIUM.

R. Iodide of potassium, half a drachm.
Lard, one ounce and a half.

Mix.

Magendie.

The strength of this ointment may vary much, from containing a twenty-fourth of the salt, to an eighth or more, according as it may be required. In goitre, scrofulous swellings, &c.

R. Iodide of potassium, in
powder, one drachm.
Boiling water, one fl. drachm.
Lard, one ounce.

Dissolve the iodide in the water, and mix the solution with the lard.

U. S. Ph.

R. Iodide of potassium, half a drachm.
Solution of potassa, two drops.
Lard, one ounce and a half.
Wax, three drachms.

Mix.

Coindet.

R. Spermaceti, half an ounce.
Olive oil, six drachms.
White wax, two drachms.
Iodide of potassium, four scruples.
Oil of lemon,

“ roses, each, three drops.

Mix. This ointment does not change color by keeping.

Kallhofert.

R. Iodide of potassium, one drachm.
Magnesia, six grains.

Rub together with a few drops of water, and add

Rose water ointment, one ounce.

Mix well.

Hamb. Ph.

R. Iodide of potassium,
Carbonate of soda, each,
half a drachm.

Rose water ointment, six drachms.

Mix well. In chronic enlargement of the testicle.

Walthen.

OINTMENT OF IODIDE OF POTASSIUM AND OPIUM.

R. Iodide of potassium, one drachm and a half.
Lard, one ounce and a half.
Laudanum, one fl. drachm.

Mix.

A. T. Thomson.

R. Iodide of potassium, half a drachm.
Extract of opium, ten grains.
Simple cerate, one ounce.

Mix. As a dressing to cancerous ulcers.

OINTMENT OF IODIDE OF POTASSIUM AND MERCURY.

R. Iodide of potassium, twelve to twenty grains.
Mercurial ointment, half an ounce.

Mix. In various diseases of the skin.

Blarius

- R. Ointment of iodide of potassium,
 Mercurial ointment,
 Oil of henbane, each, two ounces.
 " juniper, one drachm.

Mix. *Schonlein.*

When iodide of potassium and mercurial ointment are mixed, the globules soon appear; but if the iodide be artificially dried, previously, a uniform ointment is produced.

- R. Iodide of potassium, three parts.
 Iodide of mercury, two parts.
 Camphor, two parts.
 Galen's cerate, thirty-two parts.

Mix. Used as a friction in acute hydrocephalus, in doses of half a drachm to a drachm. *Gelfin.*

OINTMENT OF IODIDE OF POTASSIUM AND MORPHIA.

- R. Iodide of potassium, fifteen grains.
 Muriate of morphia, half a drachm.
 Lard, ten ounces.

Mix. In painful tumors of the breasts, to be applied twice a-day. *Chomet.*

LINIMENT OF IODIDE OF POTASSIUM.

- R. Iodide of potassium, one ounce.
 Alcohol, four ounces.

Dissolve.

- R. Animal soap, one ounce and a half.
 Alcohol, four fl. ounces.

Dissolve. Add the two solutions together, aromatize with a few drops of oil of lavender, and pour into wide-mouthed bottles. *Boudet.*

- R. White soap, seven drachms.
 Diluted alcohol, two fl. ounces.

Dissolve by a gentle heat, and whilst warm, add, Iodide of potassium, four drachms, dissolved in

- Diluted alcohol, four fl. drachms.

Beasley.

- R. Iodide of potassium, four drachms.
 Diluted alcohol, two ounces.

Dissolve.

- Curd soap, six drachms.
 Diluted alcohol, two ounces.

Dissolve by a gentle heat, and add the solution of iodide of potassium, and aromatize. *Foy.*

These are used as liniments in goitre.

- R. White soap, ten drachms.
 Oil of almonds, ten drachms.
 Iodide of potassium, one drachm.
 Water, one drachm.

Dissolve the iodide in the water, and add it to the soap and oil, melted together. *Guibourt.*

PLASTER OF IODIDE OF POTASSIUM.

- R. Iodide of potassium, one ounce.
 Prepared frankincense, six ounces.
 Wax, six drachms.
 Olive oil, two fl. drachms.

Melt the frankincense and wax, add the iodide previously rubbed with the oil, and stir till cool. *Lond. Ph.*

SOAP OF IODIDE OF POTASSIUM.

- R. Camphor, one drachm.
 Tincture of benzoin, three drachms.

Add to the solution, triturating well,

- Iodide of potassium, two drachms.
 Goulard's extract, four drachms.

Add to the mixture,

- Oil of almonds, thirty-four drachms.
 Solution of potassa, two ounces.
 Essence of lavender, half a drachm.

Mix well. As an application to chilblains. *Cadet.*

POTASSII SUPER-IODIDUM.

IODURETTED POTASSIUM.

- R. Iodide of potassium, twenty parts.
 Iodine, six parts.

Mix, and triturate together till the mass is homogeneous, and of a dark brown color. *Giordano.*

To have a perfectly saturated salt, iodine in excess is to be added to a solution of the iodide until no more is dissolved. In the following preparations there is a mixture of the iodide and superiodide.

IODURETTED WATER.

- R. Iodide of potassium, six grains.
 Iodine, one grain.
 Water, two pints.

Dissolve. To be used as a drink at meals.

Magendie.

LUGOL'S IODURETTED WATERS.

No. 1.

- R. Iodide of potassium, one grain and a half.
 Iodine, three-quarters of a grain.
 Distilled water, eight fl. ounces.

Dissolve.

No. 2.

- R. Iodide of potassium, two grains.
 Iodine, one grain.
 Distilled water, eight fl. ounces.

Dissolve.

No. 3.

R. Iodide of potassium, two grains and a half.
Iodine, one grain and a quarter.
Distilled water, eight fl. ounces.

Dissolve. The first is for young children, to be taken in divided doses, in three days; the second for those of more advanced age, in two days; the third for adults, in one day. *Lugol.*

R. Iodide of potassium, six grains.
Iodine, three grains.
Water, sixteen fl. ounces.

Dissolve. In poisoning by the vegetable alkalis, in wineglassfuls, after the stomach has been emptied. *Bouchardat.*

COMPOUND SOLUTION OF IODIDE OF POTASSIUM.

R. Iodide of potassium, ten grains.
Iodine, five grains.
Distilled water, twenty fl. ounces.

Dissolve. Dose, two to six fl. drachms.

Lond. Ph.

R. Iodide of potassium, one ounce and a half.
Iodine, six drachms.
Distilled water, one pint.

Dissolve. Dose, six drops, three times a-day, in sweetened water. *U. S. Ph.*

IODURETTED BATHS.

R. Iodide of potassium, four scruples.
Iodine, two scruples.
Water, ten fl. ounces.

Dissolve. To be added to a sufficient quantity of water. For children.

R. Iodide of potassium, six drachms.
Iodine, three drachms.
Water, twenty fl. ounces.

Dissolve. To be added to a sufficient quantity of water. For adults. *Foy.*

COLLYRIUM OF IODURETTED POTASSIUM.

R. Iodide of potassium, twenty-four grains.
Iodine, one to two grains.
Rose water, six fl. ounces.

Dissolve. To be used four times a-day, in scrofulous ophthalmia. *Ryan.*

IODURETTED INJECTION.

R. Iodide of potassium, four grains.
Iodine, two grains.
Distilled water, sixteen fl. ounces.

Dissolve.

R. Iodide of potassium, eight grains.
Iodine, four grains.
Distilled water, sixteen fl. ounces.

Dissolve. To stimulate fistulous sinuses.

Guibourt.

SYRUP OF IODURETTED POTASSIUM.

R. Ioduretted potassium, twenty grains.
Peppermint water, two fl. drachms.
Simple syrup, two fl. ounces.

Dissolve the biniodide in the peppermint water, and add the solution to the syrup. *Giordano.*

LOTION OF IODURETTED POTASSIUM.

R. Iodide of potassium, two to four grains.
Iodine, one to two grains.
Distilled water, eight fl. ounces.

Dissolve. As an application in scrofulous ophthalmia, fistulas, &c. *Lugol.*

CAUSTIC SOLUTIONS OF IODURETTED POTASSIUM.

R. Iodide of potassium, two ounces.
Iodine, one ounce.
Distilled water, three fl. ounces.

Dissolve. *Guibourt.*

R. Iodide of potassium, one ounce.
Iodine, half an ounce.
Distilled water, six fl. ounces.

Dissolve. *Soubeiran.*

These two preparations are used to touch the eyelids in scrofulous ophthalmia.

R. Iodide of potassium, one ounce.
Iodine, each, one ounce.
Distilled water, two fl. ounces.

Mix. *Guibourt.*

R. Iodide of potassium, one ounce.
Iodine, half an ounce.
Distilled water, each, one ounce.

Mix. *Cadet.*

These are used to touch the surfaces of scrofulous ulcers.

COMPOUND TINCTURE OF IODINE.

R. Iodide of potassium, one ounce.
Iodine, half an ounce.
Alcohol, one pint.

Dissolve. *U. S. Ph.*

R. Iodide of potassium, two ounces.
Iodine, one ounce.
Alcohol, (Imp.) two pints.

Dissolve. *Lond. Ph.*

Dose, from fifteen to thirty drops.

MIXTURE OF IODURETTED POTASSIUM.

R. Iodide of potassium, four drachms.
 Iodine, two grains.
 Orange-flower water,
 Mint water, each, three fl. ounces.

Mix. A dessertspoonful, three times a-day, in epilepsy. *Magendie.*

R. Iodide of potassium, half a drachm.
 Iodine, half a grain.
 Syrup of poppies, half a fl. ounce.
 Distilled water, half a pint.

Mix. Two tablespoonfuls, three times a-day, in syphilis combined with scrofula. *Tyrell.*

COMPOUND OINTMENT OF IODINE.

R. Iodide of potassium, one drachm.
 Iodine, half a drachm.
 Alcohol, one fl. drachm.
 Lard, two ounces.

Rub the iodide and iodine with the alcohol, and then with the lard, until they are thoroughly mixed. *U. S. Ph.*

R. Iodide of potassium, twenty-four grains.
 Iodine, twelve grains.
 Lard, two ounces.

Mix. *Lugol.*

R. Iodide of potassium, two drachms.
 Iodine, eighteen grains.
 Lard, two ounces.

Mix. *Lugol.*

R. Iodide of potassium, two drachms and a half.
 Iodine, twenty-one grains.
 Lard, two ounces.

Mix. *Lugol.*

R. Iodide of potassium, two drachms and a half.
 Iodine, twenty-four grains.
 Lard, two ounces.

Mix. In scrofulous ulcers, &c. *Lugol.*

OINTMENT OF BINIODIDE OF POTASSIUM AND OPIUM.

R. Iodide of potassium, one drachm.
 Iodine, fifteen grains.
 Rousseau's laudanum, two drachms.
 Lard, two ounces.

Mix. As an application to scrofulous ulcers. *Foy.*

PLASTER OF IODURETTED POTASSIUM.

R. Iodide of potassium,
 Iodine, each, ten grains to one scruple.
 Mercurial or soap plaster, two ounces.

Mix. As an application to syphilitic and gouty swellings. *Ebers.*

POTASSII HYDRARGYRO-
IODIDUM.

HYDRARGYRO-IODIDE OF POTASSIUM.

R. Iodide of potassium, eight grains,
 dissolved in ten or fifteen minims of water.

Red iodide of mercury, eleven grains.

Mix. This contains twenty grains of the hydrargyro-iodide, and is not decomposed by water. *Channing.*

R. Bichloride of mercury, one equivalent.

Dissolve in a solution of

Iodide of potassium, four equivalents.

Evaporate to dryness, treat with alcohol, and evaporate to crystallization. *Amb. Smith.*

Dose, one-twelfth of a grain, three times a-day; in many cases, a much smaller quantity is required to be given.

SOLUTION OF HYDRARGYRO-IODIDE OF
POTASSIUM.

R. Iodide of potassium, three grains and a half.
 Red iodide of mercury, four grains and a half.
 Distilled water, one fl. ounce.

Dissolve first, the iodide of potassium, and then the mercurial salt in the water. The compound salt amounts to about eight grains. The dose is from two to five drops, three times a-day, much diluted, in dyspepsia, enlargement of the spleen, dropsy, &c. *Channing.*

R. Iodide of potassium,
 Red iodide of mercury, eight grains.
 each, eight grains.
 Distilled water, eight fl. ounces.

Mix. Dose, two fl. drachms and upwards, in the twenty-four hours. *Puche.*

R. Hydrargyro-iodide of potassium, twelve grains.
 Water, sixteen fl. ounces.

Mix. *Lamothe.*

TINCTURE OF HYDRARGYRO-IODIDE OF
POTASSIUM.

R. Hydrargyro-iodide of potassium, one grain.
 Diluted alcohol, one fl. ounce.

Dissolve. Ten drops, three times a-day. *Channing.*

PILLS OF HYDRARGYRO-IODIDE OF
POTASSIUM.

R. Red iodide of mercury,
 Iodide of potassium, each, eight grains.

Sugar of milk, sixty-four grains.
Mucilage of gum Arabic, sufficient.
Mix, and make thirty-two pills. *Puche.*
R. Iodide of potassium,
Red iodide of mercury,
each, six grains.
Opium, twelve grains.
Mix, and make twenty-four pills. *Mialhe.*

OINTMENT OF HYDRARGYRO-IODIDE OF POTASSIUM.

R. Red iodide of mercury, seven grains.
Iodide of potassium, two scruples.
Lard, one ounce.
Mix. To be applied to tumors, two or three times a-day. *Hildreth.*
R. Hydrargyro-iodide of potassium, one scruple.
Lard, one ounce.
Mix. *Lamothe.*

POTASSII SULPHO-CYANURETUM.

SULPHO-CYANURET OF POTASSIUM.

R. Ferro-cyanuret of potassium, three parts.
Sulphur, one part.
Pack in a crucible, heat to redness, for an hour, treat with alcohol, and evaporate to crystallization. *Van Mons.*
R. Prussian blue, three parts.
Sulphuret of potassium, one part.
Put the mixture into a covered crucible, and heat to a dull redness for half an hour; treat with alcohol, filter the solution, evaporate, and crystallize. *Van Mons.*
R. Digest an aqueous solution of cyanuret of potassium with sulphur, of which it will take up one-third.
Filter, and evaporate. *Beasley.*

POTASSII SULPHURETUM.

SULPHURET OF POTASSIUM.

LIVER OF SULPHUR.

R. Sulphur, one ounce.
Carbonate of potassa, two ounces.
Rub the carbonate of potassa, previously dried, with the sulphur; melt the mixture in a covered crucible over the fire; then pour it out, and, when cold, put it in a bottle, which is to be well stopped. *U. S. Ph.*
A stimulant expectorant, and diaphoretic, in small doses, poisonous in large; used externally in many cutaneous diseases. Dose, from two to ten grains, several times a-day.

HYPOSULPHITED SULPHURET OF POTASSIUM.

R. Sublimed sulphur, one part.
Solution of potassa, three parts.
Mix, and heat on a sand-bath, until it marks 39° B. *Guibourt.*
It consists of three parts of sulphuret, and one of hyposulphite of potassium.

FERRO-SULPHURET OF POTASSIUM.

R. Carbonate of potassa,
Sulphur, each, one ounce.
Black oxide of iron, two drachms.
Mix, and melt in a crucible, pour out on a slab of marble, break in pieces, and keep in a well-closed bottle. Dose, three to four grains. *Ferrara Ph.*

SULPHURET OF POTASSIUM AND CREAM OF TARTAR.

R. Sulphuret of potassium, four scruples.
Cream of tartar, four drachms.
Mix, and divide into twenty-four powders. Dose, one every four hours, in a glass of sweetened water. Said to be efficacious in mercurial salivation. *Bories.*

BOLUS OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium, three grains.
Conserve of elder berries, sufficient.
Mix. Make six boluses; one every three hours. In mercurial salivation. *Brera.*
R. Sulphuret of potassium, six grains.
Black oxide of iron, three grains.
Extract of quassia, ten grains.
Burnt sponge, sufficient.
Mix, and make a bolus. One, morning and evening, in goitre and glandular affections. *Phabus.*

PILLS OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium, one drachm.
Extract of liquorice, sufficient.
Mix, and make thirty pills. Two to five, several times a-day.

COMPOUND PILLS OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium,
Ammoniac,
Extract of dandelion, each,
one drachm.
Soap,
Rhubarb, each, half a drachm.
Opium, three grains.
Mix, and make pills of two grains. *Radius.*

ELECTUARY OF SULPHURET OF POTASSIUM.

R. Butter of cocoa, two drachms.
 Oil of almonds, half an ounce.
 Melt together, and add, triturating well,
 Sulphuret of potassium, ten grains.
 Sugar, three drachms.

As an alterative for children, in teaspoonful doses. *Phæbus.*

SYRUP OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium, eight grains.
 Distilled water, sixteen grains.

Dissolve, and add
 Simple syrup, one ounce.
Par. Cod.

A teaspoonful, for croup in children.

COMPOUND SYRUP OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium, one ounce.
 Fennel water, sixteen ounces.
 Simple syrup, thirty-eight ounces.

Mix. Once much celebrated as *Willis's syrup*, in croup, &c. *Giordano.*

SYRUP OF HYPOSULPHITED SULPHURET OF POTASSIUM.

R. Hyposulphited sulphuret of potassium, sixteen grains.
 Simple syrup, one ounce.

Mix. *Guibourt.*

SOLUTION OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium, one part.
 Water, sufficient
 to make a solution of 30° B. *Soubeiran.*

WATER OF SULPHURET OF POTASSIUM.

R. Washed sulphur, one part.
 Water of caustic potassa, eleven parts.
 Boil for ten minutes, and filter. Keep in well-stopped bottles. *Dub. Ph. 1826.*

This is not analogous to a solution of sulphuret of potassium, as it contains much hypsulphite of potassa. Dose, ten minims to one fl. drachm, three times a-day.

TINCTURE OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium, four ounces.
 Diluted alcohol, sixteen ounces.
 Digest for twenty-four hours, and strain.

Quincy.

R. Carbonate of potassa, one ounce.
 Sulphur, two ounces.

Melt together, and digest with
 Alcohol, one pound.

Filter. *Saxon. Ph.*

Recommended in itch, in doses of sixty drops.

LOTION OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium, one ounce.
 Water, half a pint.

Dissolve. As a wash in herpetic and other cutaneous eruptions. *Ellis.*

R. Sulphuret of potassium, one to two parts.
 Water, sixteen parts.

Dissolve. *Foy.*

COMPOUND LOTION OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium, half an ounce.
 Soap, one ounce.
 Alcohol, four fl. ounces.
 Tincture of myrrh, half a fl. ounce.
 Lime water, one pint.

Mix. As an application in tinea capitis. *Ellis.*

R. Sulphuret of potassium, one to two ounces.
 Water, one pint.

Dissolve.

R. Muriatic acid, one to two fl. ounces.
 Distilled water, two pints.

Mix an ounce of each solution with four ounces of warm water. As an application in psora. *Alibert.*

R. Sulphuret of potassium, three drachms.

Soap, one drachm and a half.
 Lime water,

seven and a half fl. ounces.

Diluted alcohol, two fl. ounces.

Mix. *Burns.*

Known as *Barlow's lotion*, and used in various cutaneous diseases.

R. Sulphuret of potassium, two drachms.
 Soap, two drachms and a half.

Lime water, seven fl. ounces.

Alcohol, one fl. drachm.

Mix. *Bielt.*

As a lotion in porrigo.

R. Sulphuret of potassium, four ounces.
 Water, one pint and a half.

Sulphuric acid, half an ounce.

Mix. As a lotion in itch, to be used morning and evening. *Dupuytren.*

SULPHURET OF POTASSIUM-BATH.

R. Sulphuret of potassium, four ounces.
Water, twelve ounces.
Dissolve, and mix with the water of a bath.

Beral.

R. Sulphuret of potassium, one ounce.
Common salt, two ounces.
Carbonate of soda, four drachms.
Leaves of sage, one to two handfuls.
Water, six quarts.

Boil for twenty-four hours. In fomentations, douches, &c., in spina ventosa and scrofula.

St. Marie.

R. Sulphuret of potassium, four ounces.
Water, two hundred pints.

Mix, and add

Glue, two pounds,
dissolved in ten pints of boiling water.

Foy.

INJECTION OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium, one drachm.
Distilled water, eight to twelve ounces.

Dissolve. In gonorrhœa.

Wedekind.

MIXTURE OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium,
one drachm and a half.

Bicarbonate of potassa, ten grains.

Oil of peppermint, one to two drops.

Syrup of orange peel, two ounces.

Mucilage, one ounce.

Mix. A teaspoonful, every two hours, as an alterative in scrofula.

Lockstaedt.

R. Sulphuret of potassium, one scruple.
Carbonate of potassa, ten grains.

Peppermint water, two ounces.

Syrup of saffron, one ounce.

Mix. A spoonful, every two hours, in cutaneous diseases.

Phæbus.

R. Sulphuret of potassium, one scruple.
Water, one ounce.

Syrup of cinnamon, two drachms.

Mix. To be taken in three doses, in mercurial diseases.

Dzondi.

R. Sulphuret of potassium, one scruple.
Kermes mineral, ten grains.

Syrup of seneka, two ounces.

Anisated ammonia, one scruple.

Mix. A spoonful, every hour in croup.

Hagen.

R. Sulphuret of potassium,
Sugar, each, one drachm.

Water, one pint.

Mix. Has been advised as an antidote in poisoning by arsenic.

Augustin.

OINTMENT OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium,
Carbonate of soda, each,
three drachms.

Lard, three ounces.

Mix. As an application in tinea capitis.

Alibert.

R. Soap, one pound.

Water, one ounce.

Soften by means of a water-bath, and add

Sulphuret of potassium, three ounces.

Oil of pinks, two pounds.

Mix well.

Jadelot.

R. Sulphuret of potassium, six ounces.

Water, two ounces.

Dissolve, and add

Soap, two pounds.

Oil of pinks, four pounds.

" thyme, two drachms.

Mix.

Giordano.

R. Powdered soap, two ounces.

Sulphuret of potassium,

Water, each, one ounce.

Mix well, and add, gradually,

Olive oil, five ounces.

This liniment alters rapidly when exposed to the air.

Beral.

COMPOUND PLASTER OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium,
Powdered hemlock, each,
two drachms.

Camphor,

Turpentine, each, four drachms.

Soap, half a drachm.

Yellow wax, one ounce.

Simple plaster, four ounces.

Melt and mix well. As a dressing to tumefied lymphatic glands.

Bories

SOAP OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium, one part.
Boiling water, two parts

Dissolve, and add

Yellow wax, one part.

Evaporate, stirring continually.

Bavar. Ph.

POTASSA.

CAUSTIC POTASSA.

R. Solution of potassa, one gallon.

Evaporate quickly, in a clean iron vessel, over the fire, till ebullition ceases, and the potassa

melts. Pour into moulds, and keep in well-stopped bottles. *U. S. Ph.*

A powerful escharotic, used to form issues and in opening abscesses.

SOLUTION OF CAUSTIC POTASSA.

R. Caustic potassa,
one drachm and a half.
Distilled water, two fl. ounces.

Dissolve. As a rubefacient in tetanus, to be applied to the spine. *Jos. Hartshorne.*

POTASSA WITH LIME.

R. Caustic potassa,
Lime, each, one ounce.
Rub them together, and keep in a well-stopped bottle. *Lond. Ph.*

Used as the above, but is slower in producing an effect. It is to be made into a paste with a little alcohol.

SOLUTION OF POTASSA. (*Sp. Gr.* 1.056.)

R. Carbonate of potassa, one pound.
Lime, half a pound.
Boiling distilled water, one gallon.

Dissolve the carbonate of potassa in half a gallon of the water; pour a little of the water on the lime, and when slaked, add the remainder. Mix the hot liquors, and boil for ten minutes, stirring continually; set the mixture aside in a covered vessel, till clear. Decant, and keep in well-stopped bottles of green glass. *U. S. Ph.*

R. Caustic potassa, one scruple.
Water, one fl. ounce.

Dissolve. This solution very nearly represents the above. *W. Procter.*

Dose, ten to fifteen minims.

LITHONTRIPTIC SOLUTION OF CAUSTIC POTASSA.

R. Calcined carbonate of
potassa, half a pound.
Burnt oyster shells,
one pound and a half.
Boiling water, eight pints and a half.

Mix, and let rest for twenty-four hours, and then filter. One fl. drachm three times a-day, in flaxseed tea, adding fifteen grains of magnesia to each dose. *Saunders.*

SAVIARD'S STIMULANT LOTION.

R. Caustic potassa, two drachms.
Camphor, two scruples.
Sugar, two ounces.
Water, two pints.

Mix. As an application to indolent ulcers. *Foy.*

COLLYRIUM OF CAUSTIC POTASSA.

R. Caustic potassa, one grain.
Distilled water, one ounce.

Dissolve. A drop or two to be introduced into the eye, to remove specks on the cornea.

Gimbernat.

R. Caustic potassa, one grain.
Extract of opium, four grains.
Distilled water, four fl. ounces.

Dissolve, and filter. In chronic ophthalmia. *Hamb. Ph.*

MIXTURE OF CAUSTIC POTASSA.

R. Tincture of caustic potassa,
half an ounce.

Volatile tincture of
guaiacum, two drachms.
Laudanum, half a drachm.

Mix. Twenty drops, three times a-day, in gout. *Græfe.*

INJECTION OF CAUSTIC POTASSA.

R. Caustic potassa, half a grain.
Chamomile water,
one ounce and a half.
Laudanum, five drops.

Mix. As an injection into the ear, in deafness. *Rust.*

R. Caustic potassa, two grains.
Distilled water, one ounce.

Dissolve. As an injection, at the commencement of gonorrhœa. *Girtanner.*

POTASSÆ ACETAS.

ACETATE OF POTASSA.

R. Acetic acid, one pint.
Carbonate of potassa, sufficient
to saturate. To be added gradually; filter;
evaporate to dryness, by means of a sand-bath;
keep in closely-stopped bottles. *U. S. Ph.*

A diuretic, in doses of a scruple to a drachm;
as a laxative, in doses of two or three drachms.

SCILLITIC ACETATE OF POTASSA.

R. Carbonate of potassa, half an ounce.
Vinegar of squills, twelve fl. ounces.
Evaporate to the consistence of honey, and add

Alcohol (.842), six fl. ounces.

Decant, after digesting for a few days. Dose, forty to sixty drops, in some demulcent fluid, in dropsy and asthma. *Keup.*

BOLUS OF ACETATE OF POTASSA.

R. Acetate of potassa, one scruple.
Conserve of pot marigold, sufficient.

Mix. To be taken in the morning, for some days, to retard the secretion of milk.

St. Marie.

COMPOUND PILLS OF ACETATE OF POTASSA.

R. Acetate of potassa,
Ammoniac,
Pills of aloes and
myrrh, each, one part.
Soap, two parts.
Simple syrup, sufficient.

Mix, and make pills of five grains. *Van Mons.*

CONSERVE WITH ACETATE OF POTASSA.

R. Acetate of potassa, half an ounce.
Sulphate of soda, one drachm.
Juice of scurvy grass,
" fumitory,
" dandelion, each, two ounces.
Sugar, sufficient.

Mix, and form conserve. A teaspoonful, two or three times a-day, in obstructions of the bowels. *Bories.*

R. Acetate of potassa,
Powdered burnt
sponge, each, two drachms.
Calomel, twelve grains.
Sulphuret of antimony,
one drachm and a half.
Jalap, sixteen grains.
Sulphur, one drachm.
Simple syrup, sufficient.

Mix. Dose, twelve to twenty-four grains, twice a-day, in scrofula. *Baumes.*

LIQUID ACETATE OF POTASSA.

R. Acetate of potassa, one part.
Distilled water, two parts.
Dissolve. *Amster. Ph.*

R. Acetate of potassa, at will.
Expose to the action of the air, till it deliquesces, and filter. *Pideret.*

Dose, forty to eighty drops, in an appropriate vehicle, as a diuretic.

MIXTURE WITH LIQUID ACETATE OF POTASSA.

R. Liquid acetate of potassa, two ounces.
Extract of Peruvian bark,
two drachms.
" cascarilla, one drachm.
Tincture of aloes
and myrrh, one ounce.

Mix. In intermittent fevers, complicated with obstruction and dropsy; in doses of thirty drops, every two hours, during the apyrexia. *Bories.*

R. Liquid acetate of potassa, four ounces.
Extract of centaury, two drachms.
Sulphuric ether, twenty drops.
Syrup of kermes, two ounces.

Mix. In dropsy and obstructions. In doses of thirty drops. *Saunders.*

TINCTURE OF ACETATE OF POTASSA.

R. Acetate of potassa, two ounces.
Alcohol, four fl. ounces.
Digest for some days, and filter. Dose, thirty to sixty drops. *Bruns. Ph.*

MIXTURE OF ACETATE OF POTASSA.

R. Acetate of potassa,
Extract of dandelion,
" hemlock, each,
half an ounce.

Fennel water, six ounces.
Syrup of marsh mallow, one ounce.

Mix. Two teaspoonfuls, every four hours, in jaundice. *Quarin.*

R. Carbonate of potassa, fifteen grains.
Distilled vinegar, sufficient

to dissolve; add to the solution

Water, half an ounce.
Mint water, one ounce.
Pimento water,
Vinegar of squill, each, one drachm.
Simple syrup, two drachms.

Mix. Dose, half a drachm to a drachm, every six hours, in dropsy. *Milman.*

R. Arnica root, one drachm.
Boiling water, sufficient

to obtain four ounces of infusion; add

Carbonate of potassa, saturated
with vinegar of squill,
half a drachm.
Syrup of orange flowers,
half an ounce.

Mix. A spoonful every two hours, in dropsy. *Wendt.*

R. Acetate of potassa,
Oxymel of squill, each, one drachm.
Linden water, four drachms.
Wine of opium, fifteen drops.
Syrup of marsh mallow, one ounce

Mix. In spoonful doses, as a diuretic and sedative. *Pierquin.*

POTASSÆ ARSENITIS LIQUOR.

ARSENICAL OR FOWLER'S SOLUTION.

R. Arsenious acid, in small fragments,
Pure carbonate of
potassa, each, sixty-four grains

Distilled water, sufficient.
Compound spirit of lavender, half a fl. ounce.

Boil the arsenious acid and carbonate of potassa with twelve fl. ounces of the water, in a glass vessel, till the acid is entirely dissolved. To the solution, when cold, add the spirit of lavender, and afterwards, sufficient distilled water to make it fill exactly the measure of a pint. *U. S. Ph.*

Used in the same cases as arsenious acid, especially in intermittent fever. Each fl. drachm contains half a grain of arsenious acid. Dose, about ten drops, two or three times a-day.

MIXTURE OF FOWLER'S SOLUTION.

R. Fowler's solution, sixty drops.
Laudanum, thirty drops.
Compound spirit of lavender, one fl. drachm.
Cinnamon water, four fl. ounces.

Mix. Dose, a tablespoonful, for an adult, a teaspoonful for a child, every two or three hours. When given in this form, Fowler's solution is less apt to disagree with the stomach than if administered in the undiluted state. *Ellis.*

MIXTURE OF IODINE AND ARSENIC.

R. Lugol's solution, two fl. drachms.
Fowler's solution, one fl. ounce.
Mix. Dose, five drops, three times daily. *Ellis.*

POTASSÆ ARSENIAS.

ARSENIATE OF POTASSA.

R. White arsenic,
Nitre, each, one ounce.

Pulverize separately, mix, and introduce into an earthenware retort; heat to redness, as long as nitrous fumes are evolved; let cool, dissolve the residuum in four pints of boiling distilled water, evaporate, and crystallize. *Dub. Ph., 1806.*

Dose, one-sixteenth to one-eighth of a grain.

MIXTURE OF ARSENIATE OF POTASSA.

R. Arseniate of potassa, one-fifth of a grain.
Mint water, three ounces.
Simple syrup, half an ounce.

Mix. In spoonful doses, in the apyrexia of intermittents. *Foy.*

POTASSÆ BORAS.

BORATE OF POTASSA.

R. Boracic acid, six parts.
Bicarbonate of potassa, five parts.

Mix, and heat to redness in a crucible, dissolve the residuum in water, filter, and evaporate to dryness. Dose, a few grains in calculous disorders. *Beasley.*

POTASSÆ BORO-TARTRAS.

BORO-TARTRATE OF POTASSA.

R. Powdered bitartrate of potassa, four ounces.
Boracic acid, one ounce.
Water, three pints.

Mix, and put in a silver basin, and boil till most of the water is evaporated, and continue the evaporation by a gentle heat, constantly stirring. When the matter has become thick, take it up in portions, flatten them, and dry in a stove; reduce them to powder, and keep in well-closed bottles. *Par. Cod.*

R. Bitartrate of potassa, twelve ounces.
Borate of soda, six ounces.
Tartaric acid, three drachms.

Dissolve in water, clarify with white of egg, and proceed as directed by the Paris Codex. *Cambornac.*

R. Borax, one part.
Boiling water, ten parts.

Dissolve, and add Purified cream of tartar, two and one-half parts.

Mix, and evaporate on a vapor-bath until it becomes a tenacious mass; place on paper, dry by a gentle heat, and rub to powder. *Wackroder.*

This is known as *soluble cream of tartar*. It is purgative in about the same dose as cream of tartar.

POWDER OF BORO-TARTRATE OF POTASSA.

R. Cream of tartar, one ounce.
Borax, three drachms.
Sugar, two ounces.

Mix, and divide into three powders; one to be taken, in a glass of water, every half hour. *Pierquin.*

COMPOUND PILLS OF BORO-TARTRATE OF POTASSA.

R. Boro-tartrate of potassa, one drachm and a half.
Extract of buckbean, two ounces.
Seneka snake-root, two drachms.
Colchicum, one drachm and a half.

Mix, and make pills of two grains. Dose, six to twelve every two hours, as a hydragogue purgative. *Augustin.*

SOLUTION OF BORO-TARTRATE OF POTASSA.

R. Cream of tartar,
Borax, each, one drachm.
Boiling water, one pint.
Dissolve. *Bories.*

MIXTURE OF BORO-TARTRATE OF POTASSA.

R. Boro-tartrate of potassa, four to six drachms.
Juniper water,
Parsley water, each, three ounces.
Dissolve, and add
Nitric ether, two drachms.
Syrup of poppies, one ounce.
Mix. Two spoonfuls every two hours, as a diuretic. *Niemann.*
R. Boro-tartrate of potassa, one ounce.
Mint water, eight ounces.
Nitric ether, two drachms.
Oxymel of squill, one ounce.
Mix. Two spoonfuls every two hours, as a diuretic. *Selig.*
R. Boro-tartrate of potassa, half an ounce.
Tartar emetic, one grain.
Water, four ounces.
Mix. In spoonful doses, in congestion of the head. *Radius.*

BORO-TARTRATE OF POTASSA AND MAGNESIA.

R. Boro-tartrate of potassa, one part.
Carbonate of magnesia, one-fourth part.
Water, six parts.
Dissolve the boro-tartrate in the water, saturate with the magnesia, evaporate carefully, till reduced to a tenacious paste, which divide into small masses, and dry in a stove. *Mailliere Renault.*

MIXTURE OF BORO-TARTRATE OF POTASSA AND MAGNESIA.

R. Boro-tartrate of potassa and magnesia, thirty parts.
Citric acid, two parts.
Syrup of lemon, sixty parts.
Warm water, three hundred parts.
Dissolve. *Mailliere Renault.*

POTASSÆ CARBONAS.
CARBONATE OF POTASSA.

R. Impure carbonate of potassa, three pounds.
Water, two pints and a half.

Dissolve the impure carbonate in the water, filter, pour into a clean iron vessel, and evaporate over a gentle fire, till the solution thickens; then remove from the fire, and stir constantly, with an iron spatula, till the salt granulates. *U. S. Ph.*

Used as an antacid, and diuretic, &c. Dose, from five to twenty grains.

PURE CARBONATE OF POTASSA.

R. Bitartrate of potassa, two pounds.
Nitrate of potassa, one pound.
Rub them separately into powder, mix, and throw the mixture into a brass vessel, heated nearly to redness, that they may undergo combustion; from the residue prepare the pure carbonate as directed for the carbonate. *U. S. Ph., 1840.*
R. Bicarbonate of potassa, one pound.
Put it in an iron crucible, heat gradually to redness for half an hour; when cold, remove the carbonate, dissolve in water, filter, and evaporate to dryness in a porcelain crucible. *U. S. Ph., 1850.*

SOLUTION OF CARBONATE OF POTASSA.

R. Carbonate of potassa, one pound.
Distilled water, twelve fl. ounces.
Dissolve, and filter. *U. S. Ph.*
Dose, ten minims to a fl. drachm, properly diluted.

POWDER OF CARBONATE OF POTASSA.

R. Carbonate of potassa, ten grains.
Powdered chamomile,
" myrrh, each, twenty-four grains.
Mix. To be taken every hour, in the apyrexia of intermittents. *Foy.*

R. Carbonate of potassa, one part.
Powdered gum Arabic, eight parts.
Triturate together for a long time. Dose, half a drachm to a drachm, dissolved in water, in engorgement of the bowels. *Guibourt.*

COLLYRIUM OF CARBONATE OF POTASSA.

R. Carbonate of potassa, two to six grains.
Water, one ounce.
Mix. To be dropped in the eye every two to four hours, in specks on the cornea. *Himley.*

R. Carbonate of potassa, twenty-five grains.

Veal broth, filtered,
Celandine water, each, two ounces
Digest for twenty-four hours on a sand-bath, filter, and add
Tincture of aloes, twenty-four drops.
In specks on the cornea. *Bories*

TINCTURE OF CARBONATE OF POTASSA.

R. Carbonate of potassa, one part.
Calcine it with a strong heat in a crucible for two hours, pour it in a heated mortar, pulverize it rapidly, and pour on the hot powder,

Alcohol, four parts.

Digest for fifteen days in a stove, often agitating, and filter. *Guibourt.*

This is a weak alcoholic solution of caustic potassa.

COLLUTORY OF CARBONATE OF POTASSA.

R. Alcohol, four ounces.
Essence of mint, twenty drops.
" roses, eight drops.
Cochineal,
Carbonate of potassa, each, ten grains.

Macerate for twenty-four hours, and filter. Much esteemed as a mouth-wash and gargle, under the name of "*Oriental Water*." A teaspoonful is to be used, mixed with a glass of water. *Delabarre.*

EMULSION WITH CARBONATE OF POTASSA.

R. Carbonate of potassa, one scruple.
Yolk of egg, three drachms.
Mucilage, one drachm.
Oil of almonds,
Cherry water, each, one ounce.
Linden water, two ounces.

Mix. A teaspoonful, every hour, in the colic of children. *Rosenstein.*

FOMENTATION WITH CARBONATE OF POTASSA.

R. Carbonate of potassa,
Soap, each, one ounce.
Elder water, two pounds.

Dissolve, filter, and add

Sal ammoniac, two drachms.

As an application to bruises, &c. *Cadet.*

LINIMENT WITH CARBONATE OF POTASSA.

R. Solution of carbonate of potassa, two ounces.
Olive oil, four ounces.
Yolks of eggs, two.

Mix. As an application to chaps. *Plenck.*

LOTION OF CARBONATE OF POTASSA.

R. Carbonate of potassa, three drachms.
Water, four ounces.

In pruritus vaginæ. *Trousseau.*

R. Carbonate of potassa, one ounce.
Rose water, one pound.

Dissolve, and filter. As an application to chilblains. *Brugnatelli.*

R. Carbonate of potassa, three drachms.
Common salt, two drachms.
Rose water, eight ounces.
Orange-flower water, two ounces.

Mix. As a lotion in sunburn and tan.

Sundelin.

MIXTURE OF CARBONATE OF POTASSA.

R. Carbonate of potassa, two drachms.
Antimonial wine, one fl. drachm.
Laudanum, forty drops.
Compound spirit of lavender, two fl. drachms.
Distilled water, four fl. ounces.

Mix. A tablespoonful, every hour or two, as an expectorant. *Ellis.*

R. Carbonate of potassa, one scruple.
Lemon juice, four fl. drachms.
Cinnamon water, seven fl. drachms.
Wine of ipecacuanha, fifteen to twenty drops.

Mix. To be taken every three or four hours, as a diaphoretic. *Burke.*

R. Carbonate of potassa, one scruple.
Powdered gum Arabic, half a drachm.
Oil of mint, two drops.
Laudanum, ten drops.
Water, three ounces.

Mix. A tablespoonful for a dose, as may be required, in sickness of the stomach. *Ellis.*

R. Carbonate of potassa, one drachm.
Peppermint water, four ounces.
Hoffmann's anodyne, two scruples.
Simple syrup, half an ounce.

Mix. Dose, a tablespoonful in asthma, with acidity of the stomach. *Augustin.*

R. Carbonate of potassa,
White sugar, each, one drachm.
Compound spirit of lavender, two fl. drachms.
Laudanum, forty drops.
Mint water, four fl. ounces.

Mix. A tablespoonful, every hour or two, in sickness of the stomach, with acidity. *Ellis.*

R. Aromatic spirit of ammonia, one fl. drachm.
Carbonate of potassa, two drachms.
Cinnamon water, four fl. ounces.

Mix. A teaspoonful occasionally, in pyrosis. *Ellis.*

R. Carbonate of potassa,
Ipecacuanha,
Calomel, each, one drachm and a half.
Water, two pints.

Boil down to one-half in an earthen vessel. Two spoonfuls a-day, in a quart of infusion of sarsaparilla. *St. Marie.*

R. Carbonate of potassa, one drachm.
Elder-flower water, six ounces.
Syrup of marsh mallow, one ounce.

Mix. A spoonful every hour, in angina, accompanied with aphtha, in catarrhal fevers. *Radius.*

R. Carbonate of potassa, one scruple.
Cochineal, half a drachm.
Hyssop water, four ounces.
Syrup of poppies,
" tolu, each, half an ounce.
Orange-flower water,
Syrup of ipecacuanha, each, one ounce.

Mix. A spoonful, every two hours, in whooping-cough. *Bories.*

COMPOUND TINCTURE OF CARBONATE OF POTASSA.

R. Carbonate of potassa,
Cinnamon water, each, one pound.
Opium, two ounces.
Vinous cinnamon water, four ounces.

Digest in a water-bath for three weeks, often agitating; and add to the filtered solution,

Gum Arabic, two ounces.
Carbonate of ammonia, one ounce.
Cinnamon water, six ounces.

In syphilis. *Niemann.*

OINTMENT OF CARBONATE OF POTASSA.

R. Carbonate of potassa, one ounce.
Sulphur, two ounces.
Lard, four ounces.

Mix. As a friction in itch. *Brera.*

POTASSÆ BICARBONAS.

BICARBONATE OF POTASSA.

R. Carbonate of potassa, four pounds.
Distilled water, ten pints.

Dissolve the carbonate in the water, and pass carbonic acid through the solution, till it is fully saturated. Then filter and evaporate, that crystals may form, taking care that the heat does not exceed 160° F. Pour off the supernatant liquor, and dry the crystals on bibulous paper. Carbonic acid is obtained from marble, by the addition of dilute sulphuric acid. *U. S. Ph.*

Properties are the same as those of the carbonate, but it generally agrees better with the stomach. Dose, twenty grains to half a drachm.

EFFERVESCING POWDERS OF BICARBONATE OF POTASSA.

R. Bicarbonate of potassa,
six hundred and forty grains

Divide into sixteen powders.

R. Tartaric acid, one ounce.

Divide into sixteen powders.

Keep the acid and alkaline powders in papers of different colors. *Ed. Ph.*

Usually made with the bicarbonate of soda.

EFFERVESCING SOLUTION OF BICARBONATE OF POTASSA.

R. Bicarbonate of potassa, one drachm.
Distilled water, one pint (Imp.).

Dissolve, and pass into it carbonic acid, compressed by force, more than is sufficient for saturation; keep in a well-stopped vessel. *Ed. Ph.*

R. Bicarbonate of potassa,
twenty grains.

Mineral water, (of the shops,) half a pint.

Mix and drink in a state of effervescence.

Pereira.

MIXTURE OF BICARBONATE OF POTASSA.

R. Bicarbonate of
potassa, ten to fifteen grains.
Seltzer water, six fl. ounces.

Dissolve. To be taken three or four times a-day, as an antilithic. *Ellis.*

R. Bicarbonate of potassa, one drachm.
Mint water, six fl. ounces.

Dissolve. A spoonful every hour—has been recommended in cholera. *Ammon.*

POTASSÆ ET AMMONIÆ CARBONAS.

CARBONATE OF POTASSA AND AMMONIA.

R. Carbonate of potassa, four parts.
" ammonia, one part.
Water, sufficient

to dissolve the two salts. Pass a current of carbonic acid through the solution to saturation, heat gently, and let crystallize.

In doses of half a drachm, in half an ounce of mint water, four times a-day, in diabetes, dyspepsia, gravel, &c. *Swediaur*

POTASSÆ CHLORAS.

CHLORATE OF POTASSA.

- R. Carbonate of potassa, two parts.
Quicklime, one part.

Mix and expose to a current of chlorine gas. When saturated, heat the mixture gently, digest it in water, and separate the chlorate from the filtered liquid, by crystallization. *Graham.*

- R. Caustic potassa, one part.
Water, sufficient

to form a solution of sp. gr. 1.110.

- Lime, five and one-half parts.

Mix and heat to temp. of 122° F.; then pass chlorine through the mixture to saturation. Evaporate nearly to dryness, dissolve in boiling water, filter, and crystallize. *F. C. Calvert.*

Has been used in scurvy, chronic hepatitis, syphilis, &c., in doses of fifteen to twenty grains.

POWDER OF CHLORATE OF POTASSA.

- R. Chlorate of potassa, six to eight grains.
Sugar, one scruple.

Mix. To be given two to four times a-day, in phthisis, to diminish the febrile state. *Radiis.*

STEVENS'S SALINE POWDER.

- R. Chlorate of potassa, seven grains.
Chloride of sodium, one scruple.
Bicarbonate of soda, half a drachm.

Mix. For a dose. *Stevens.*

Recommended in cholera.

SOLUTION OF CHLORATE OF POTASSA.

- R. Chlorate of potassa, one drachm.
Distilled water, twelve fl. ounces.

Dissolve. *Copland.*

As a lotion to indolent ulcers.

- R. Chlorate of potassa, half a drachm.
Syrup, two and a half drachms.
Water, twelve and a half drachms.

Mix. To be given in the course of the day, in tablespoonful doses, in cancrum oris. *H. Hunt.*

- ℞. Chlorate of potassa, one drachm and a half.
Distilled water, five fl. ounces.

Dissolve. A tablespoonful, every two hours, in obstinate rheumatism, tic-douloureux, &c. *Knod.*

POTASSÆ CITRAS.

CITRATE OF POTASSA.

- R. Citric acid, ten ounces.
Bicarbonate of potassa, fourteen ounces.
Water, two pints.

Dissolve the acid in the water, and gradually add the bicarbonate; when effervescence ceases, filter the solution, if necessary, through paper, and evaporate to dryness, observing to stir constantly, as soon as the salt begins to granulate. Then rub it in a mortar, pass it through a coarse sieve, and put it in bottles, which must be closely stopped. *U. S. Ph.*

- R. Carbonate of potassa, three drachms.
Fresh lemon juice, sufficient

to saturate. Let stand for twenty-four hours, filter, and evaporate to dryness. *Van Mons.*

EFFERVESCING DRAUGHT.

- R. Carbonate of potassa, two drachms.
Distilled water, four fl. ounces.

Dissolve.

- R. Fresh lemon juice,
Distilled water, each, two fl. ounces.

Mix. Add two tablespoonfuls of the diluted lemon juice, to one of the alkaline solution, and let the mixture be taken in a state of effervescence.

An excellent diaphoretic. *Ellis.*

SOLUTION OF CITRATE OF POTASSA.

(NEUTRAL MIXTURE.)

- R. Citric acid, half an ounce.
Oil of lemon, two minims.
Water, half a pint.
Bicarbonate of potassa, sufficient.

Rub the citric acid with the oil of lemon, and then with the water, till dissolved; lastly, add the bicarbonate of potassa gradually, till the acid is perfectly saturated, then filter. *U. S. Ph.*

This neutral mixture is used where fresh lemon juice cannot be procured; it is as efficacious, but is not as agreeable.

- R. Citrate of potassa, three drachms.
Distilled water, four fl. ounces.
Oil of lemon, two drops.
White sugar, two drachms.

Mix. A tablespoonful to be taken every two hours. In same cases as the neutral mixture. *Ellis.*

COMPOUND NEUTRAL MIXTURE.

- R. Lemon juice, one fl. ounce.
Carbonate of potassa, sufficient

to saturate; add

- Mint water, seven fl. ounces.
Tartar emetic, one grain.
Syrup, half an ounce.

Mix. *Lisbon Ph.*

MIXTURE OF CITRATE OF POTASSA AND PERUVIAN BARK.

R. Lemon juice, one fl. ounce and a half.
Carbonate of potassa, one drachm.
Tincture of Peruvian bark,

one fl. ounce.

Cinnamon water, three fl. ounces.

Mix. A tablespoonful every two hours. To promote insensible perspiration while taking bark. *Ellis.*

POTASSÆ IODAS.

IODATE OF POTASSA.

R. Iodine, at will.
Caustic potassa, sufficient.

Dissolve the iodine in the alkali, till the solution begins to become colored. Evaporate to dryness; treat the residue with alcohol, and preserve the residue. The alcohol takes up the iodide of potassium. *Ferrara Ph.*

R. Iodide of potassium, one part.
Fuse in a capacious crucible, and gradually add to the fused salt, after removing from the fire,

Chlorate of potassa,
one and a half parts.

Wash the mass with warm water, which leaves the iodate undissolved. *Beasley.*

SYRUP OF IODATE OF POTASSA.

R. Iodate of potassa, twenty grains.
Peppermint water, two drachms.
Simple syrup, ten ounces.

Mix. *Giordano.*

POTASSÆ NITRAS.

NITRATE OF POTASSA.

(NITRE — SALTPETRE.)

POTASSÆ NITRAS PURUM.

PURIFIED NITRATE OF POTASSA.

R. Nitrate of potassa, four pounds.
Distilled water, five pints.

Dissolve the nitre in two pints of boiling water, and stir the solution till it cools. Decant, drain the crystals, and wash with the remainder of the water. Finally dry in an oven. *Dub. Ph.*

Dose, five to ten grains.

FUSED NITRATE OF POTASSA.

SAL PRUNELLE.

R. Nitrate of potassa, at will.
Melt in a crucible, and cast it into moulds. *Guibourt.*

R. Nitrate of potassa,
two pounds and a half.

Melt in a crucible, adding gradually
Sulphur, half an ounce.

After the deflagration, and the mixture is somewhat cooled, pour it into a basin, and turn this in all directions, so that the mass may cool in a third and uniform layer. When cool, break this in pieces. *Span. Ph.*

This preparation will contain some sulphate of potassa.

POWDER OF NITRATE OF POTASSA AND ORRIS ROOT.

R. Nitrate of potassa, one drachm.
Spermaceti, two drachms.

Sugar,
Orris root, each, one ounce.

Mix. A teaspoonful, in catarrhal affections. *Augustin.*

NITROUS POWDERS.

R. Powdered nitre, one drachm.
Tartar emetic, one grain.

Calomel, four grains.

Mix, and divide into eight powders. One every two hours. As a diaphoretic, &c., in fevers. *Dewees.*

R. Nitrate of potassa, two ounces.
Bitartrate of potassa, four drachms.

Tartar emetic, four grains.

Mix. Dose, ten to thirty grains. *Univer. Coll. Hosp.*

POWDER OF NITRATE OF POTASSA AND SQUILL.

R. Nitrate of potassa, fifteen grains.
Powdered squill,

" pimento, each, ten grains.

Mix. Dose, ten grains, two to three times a-day, as a diuretic. *Swedjaur.*

POWDER OF NITRATE OF POTASSA AND CAMPHOR.

R. Nitrate of potassa, ten grains.
Camphor, four to eight grains.

Gum Arabic, twenty-four grains.

Mix, and triturate well; one-third to one-half at a dose. *Foy.*

PILLS OF NITRATE OF POTASSA.

R. Nitrate of potassa, six drachms.
Powdered gum Arabic, three drachms.

" liquorice root,

" marsh mallow, each,
three ounces.

Simple syrup, sufficient

Mix, and make pills of five grains. In inflammation of the urethra and dysuria; five or six, three times a-day. *Foy.*

PILLS OF NITRATE OF POTASSA AND CAMPHOR.

- R. Nitrate of potassa, four parts.
Camphor,
Conserve of roses, each, two parts.
Mix, and make pills of four grains. Two to ten
a-day, in gonorrhœa. *Guibourt.*

NITRATED EMULSION.

- R. Nitrate of potassa, two drachms.
Sugar of milk, one ounce.
Extract of henbane, half a scruple.
Emulsion of almonds, one pound.
Mix, and dissolve. A spoonful every hour, in
gonorrhœa. *Phœbus.*

COOLING LOTION OF NITRATE OF POTASSA.

- R. Nitrate of potassa,
Sal ammoniac, each, one part.
Water, forty-eight parts.
Dissolve, and add
Vinegar, four parts.
As a lotion and application, by means of com-
presses, to contusions and ecchymoses. *Hep. Ph.*

NITRATE OF POTASSA GARGLE.

- R. Nitrate of potassa, seven drachms.
Barley water, fourteen fl. ounces.
Oxymel, one fl. ounce and a half.
Mix. As a gargle, in inflammatory sore throat.
Ainslie.

MIXTURE OF NITRATE OF POTASSA.

- R. Nitrate of potassa, eight grains.
Tincture of digi-
talis, fifteen to twenty drops.
Water, ten fl. drachms.
Sweet spirit of nitre,
Syrup of roses, each, half a fl. drachm.
Mix. To be taken twice a-day, as a diuretic.
Burke.

- R. Nitrate of
potassa, one drachm and a half.
Mucilage of
gum Arabic, two fl. ounces.
Antimonial wine, forty minims.
Syrup of orange peel, half a fl. ounce.
Water, four fl. ounces.
Mix. One fl. ounce, three times a-day, in re-
mittent fever, with hot skin. *Ainslie.*

- R. Nitrate of potassa, eight to ten grains.
Water, eleven fl. drachms.
Solution of tartar emetic, fifteen drops.

Simple syrup,
Sweet spirit of nitre, each,
half a fl. drachm.

Mix. To be taken twice a-day, as a diapho-
retic. *Burke.*

- R. Extract of chamomile, one drachm.
Nitrate of potassa, half an ounce.
Chamomile water, four fl. ounces.

Mix. A spoonful every two or three hours, as
an antispasmodic. *Radius.*

- R. Nitrate of potassa, half an ounce.
Barley water, one pound.
Syrup of marsh mallow, six ounces.
Oil of almonds, four ounces.

Mix. A glassful every four hours, in dysury
and strangury. *Cadet.*

- R. Nitrate of potassa, two drachms.
Decoction of asparagus, two pounds.
Oxymel of squill, half an ounce.

Mix. To be taken in divided doses, as a diu-
retic. *Brera.*

- R. Juniper berries, bruised, two ounces.
Boiling water, one pint.

When cold, strain, and add

- Nitrate of potassa, two drachms.
Syrup of ginger, one fl. ounce.

Dose, one ounce to an ounce and a half, every
three or four hours, in dropsy. *Hartman.*

POTASSÆ OXALAS.

OXALATE OF POTASSA.

The article usually sold as oxalate of potassa,
salt of sorrel, &c., is the quadroxalate of potassa.
It is not much used in medicine, but is em-
ployed in the arts, to remove ink and iron stains
from linen and cotton; to bleach the straw for
bonnets, &c.

- R. Oxalic acid, one part.
Carbonate of potassa, sufficient
to saturate; add to the solution three parts more
of the acid; evaporate, and crystallize. *Cooley.*

POWDER OF OXALATE OF POTASSA.

- R. Oxalate of potassa, twenty grains.
Tartrate of potassa,
Sulphate of potassa, each, one drachm.
Scammony, fifteen grains.
Red saunders, ten grains.

Mix. To be taken in the morning, in intermit-
tent fevers. Two doses are stated to be suffi-
cient. *Giordano.*

LOZENGES OF OXALATE OF POTASSA.

- R. Oxalate of potassa,
one drachm and a half.

White sugar, eight ounces.
Gum tragacanth, two scruples.
Lemon water, five drachms.
Oil of lemon, eight drops.
Mix, and make lozenges of twelve grains.
Guibourt.

POTASSÆ SILICAS.

SILICATE OF POTASSA.

R. Powdered quartz or flint, one part.
Subcarbonate of potassa, two parts.
Mix, and fuse in a crucible; when cool, dissolve in water; filter, and evaporate to dryness. *Ure.*

SOLUTION OF SILICATE OF POTASSA.

R. Silicate of potassa, ten to fifteen grains.
Distilled water, six to eight fl. ounces.
Dissolve. To be taken twice a-day, to remove gouty concretions. *Ure.*

SOLUBLE GLASS.

R. Subcarbonate of potassa, seventy parts.
Subcarbonate of soda, fifty-four parts.
Silix, one hundred and ninety-two parts.
Melt together. The resulting glass is soluble in boiling water. The solution forms a fine, transparent, elastic varnish. *Dobereiner.*

POTASSÆ SULPHAS.

SULPHATE OF POTASSA.

R. Residuum of the preparation of nitric acid, two pounds.
Boiling water, two gallons.
Expel the excess of acid by heating the salt in a crucible; boil the remainder in the water, till a pellicle forms; filter the solution; set aside to crystallize; pour off the water, and dry the crystals. *Lond. Ph., 1836.*

A mild cathartic, in doses of one drachm; but in doses of four or five drachms it acts as an irritant.

COMPOUND SALINE POWDER.

R. Sulphate of potassa, three ounces.
Muriate of soda, four ounces.
Sulphate of magnesia, each, four ounces.
Dry the salts separately, with a gentle heat; then triturate them well together, and preserve in glass vessels. *Ed. Ph.*

Aperient, in doses of two or three drachms, dissolved in half a pint of carbonic acid water. To be taken before breakfast.

POWDER OF SULPHATE OF POTASSA AND RHUBARB.

R. Sulphate of potassa, one drachm.
Powdered rhubarb, half a drachm.
" chamomile, one drachm.

Mix, and divide into six powders; one, twice a-day, in sugar and water, in dyspepsia and torpor of the bowels. *Ellis.*

R. Sulphate of potassa, two ounces.
Powdered rhubarb, one ounce.
Sal ammoniac, half an ounce.

Mix. Half a drachm, in same cases as the last. *Pfaff.*

PILLS OF SULPHATE OF POTASSA.

R. Sulphate of potassa, two drachms.
Powdered rhubarb, two scruples.
Oil of fennel, six drops.
Extract of blessed thistle, sufficient.
Mix, and make sixty pills. Dose, five or six a-day, as a purgative. *Phabus.*

MIXTURE WITH SULPHATE OF POTASSA.

R. Centaury, one ounce.
Chamomile, each, one ounce.
Water, sufficient
to obtain three pints of infusion; add
Sulphate of potassa,
Honey, each, two ounces.
Three wine-glassfuls a-day, in fevers. *Swediaur.*

EFFERVESCING DRAUGHT OF SULPHATE OF POTASSA.

R. Sulphate of potassa, one drachm.
Carbonate of soda, each, one drachm.
Dissolve separately, in two fl. ounces of water, each; mix, and take whilst effervescing. *Barker.*

POTASSÆ BISULPHAS.

BISULPHATE OF POTASSA.

R. Salt remaining after distillation of nitric acid, two pounds.
Sulphuric acid, one pound.
Boiling water, (Imp.) six pints.

Dissolve the salt in the water, add the acid, and mix. Boil down the solution, and set aside, that crystals may form. *Lond. Ph., 1836.*

Aperient and tonic, in doses of one to two drachms, properly diluted.

DISINFECTING POWDER.

R. Acid sulphate of
potassa, four hundred and ten parts.
Subacetate of lead, seventy parts.
Manganese, thirty parts.
Reduce these, separately, to fine powder, and,
when wanted, mix in a proper vessel. *Keist.*

POTASSÆ ET AMMONIÆ
SULPHAS.

SULPHATE OF POTASSA AND AMMONIA.

R. Bisulphate of potassa, one part.
Boiling water, two parts.
Dissolve, and add
Solution of ammonia, sufficient
to saturate; and crystallize. *Van Mons.*

POTASSÆ ET MAGNESIÆ
SULPHAS.

SULPHATE OF POTASSA AND MAGNESIA.

R. Sulphate of
potassa, three and a half parts.
Sulphate of magnesia, ten parts.
Boiling water, sufficient.
Filter and crystallize.

POTASSÆ SULPHAS CUM
SULPHURE.

SULPHATE OF POTASSA WITH SULPHUR.

R. Nitrate of potassa,
Sulphur, equal parts.
Mix, and throw in small successive portions
into a red-hot crucible; when cool, reduce to
powder, and keep in well-closed bottles. *Ed. Ph.*

Has much the same properties as the sulphate
of potassa. Dose, half a drachm to a drachm.

POTASSÆ TARTRAS.

TARTRATE OF POTASSA.

SOLUBLE TARTAR.

R. Carbonate of potassa, sixteen ounces.
Bitartrate of potassa,
three pounds, or sufficient.
Boiling water, one gallon.

Dissolve the carbonate of potassa in the water,
then gradually add the bitartrate in fine pow-
der to the solution, till it is perfectly saturated,

and boil. Filter the liquor, evaporate till a pel-
licle forms, and set aside to crystallize. Pour
off the liquid, and having dried the crystals on
bibulous paper, keep them in closely-stopped
bottles. *U. S. Ph.*

A mild, cooling purgative, in doses from a
drachm to an ounce.

POWDER OF TARTRATE OF POTASSA AND
RHUBARB.

R. Tartrate of potassa, two drachms.
Powdered rhubarb,
" orange peel, each,

one drachm.

Oil of cajeput, one scruple.

Mix. A teaspoonful, three times a-day, in ob-
structions of the portal system. *St. Marie.*

MIXTURE OF TARTRATE OF POTASSA.

R. Tartrate of potassa,
six to eight drachms.

Infusion of chicory, twenty ounces.

Manna, two ounces.

Mix. To be taken in the morning, in four por-
tions, every day or every second day, for a fort-
night, in chronic affections of the liver. *Grant.*

R. Tartrate of potassa, one ounce.

Extract of soapwort, half an ounce.

Balm water, six ounces.

Mix. Two spoonfuls, night and morning, in
all forms of hemorrhoidal disease. *Radius.*

R. Tartrate of potassa, half an ounce.

Extract of centaury, two drachms.

Water, eight fl. ounces.

Mix. Two spoonfuls every hour or two, in ob-
structions of the liver. *Svediaur.*

R. Tartrate of potassa, one drachm.

Nitrate of potassa, half a drachm.

Manna, one ounce.

Decoction of dandelion, six fl. ounces.

Mix. Two spoonfuls, every two hours, in drowsy
consecutive to scarlet fever. *Phabus.*

POTASSÆ BITARTRAS.

BITARTRATE OF POTASSA.

CREAM OF TARTAR.

This is cathartic, diuretic, and cooling. The
dose is from one to two drachms, as an ape-
rient; and from half an ounce to one ounce, as
a hydragogue purgative.

DENTIFRICE OF BITARTRATE OF PO-
TASSA.

R. Bitartrate of potassa, six parts.
Powdered shell-lac, eight parts.

Powdered cuttlefish bone,
 " orris root, each, eight parts.
 " cloves,
 " myrrh,
 " mastich, each, two parts.
 Triturate well together. *Guibourt.*
 R. Bitartrate of potassa,
 Rhatany, equal parts.
 Orris root, sufficient
 to give an agreeable odor. Triturate well
 together. *Ferrara Ph.*

POWDER OF BITARTRATE OF POTASSA.

R. Bitartrate of potassa, three ounces.
 Nitrate of potassa, three drachms.
 Sugar, four ounces.
 Mix. Dose, one to two drachms, in whey, as a
 laxative. *St. Marie.*
 R. Bitartrate of
 potassa, a drachm and a half.
 Powdered squill, two grains.
 " digitalis, one grain.
 " ginger, five grains.

Mix. Make a powder, to be taken every eight
 hours. In ascites and anasarca.

A. T. Thomson.

R. Bitartrate of potassa, thirty grains.
 Powdered squill, two grains.
 " ginger, four grains.

Mix. Make a powder, to be taken every eight
 hours. Useful in ascites. *A. T. Thomson.*

R. Bitartrate of potassa, six drachms.
 Powdered jalap, one drachm.

Mix, and divide into six powders. Give one
 powder every three hours, in dropsical cases,
 requiring purging. *Chapman.*

ELECTUARY OF BITARTRATE OF POTASSA.

R. Bitartrate of potassa, one ounce.
 Powdered ginger,
 Conserve of roses, each, one drachm.
 Syrup of orange peel, sufficient.

Mix. To be taken in spoonful doses, as a hy-
 dragogue purgative. *Monro.*

OXYMEL OF BITARTRATE OF POTASSA.

R. Bitartrate of potassa, one part.
 Clarified honey, two parts.

Mix. As a purgative in bilious fever and drop-
 sies. *Swediaur.*

SOLUTION OF BITARTRATE OF POTASSA.

R. Bitartrate of potassa, one ounce.
 Water, two pints.

Dissolve. To be taken freely during the day,
 as a diuretic. *Ellis.*

COMPOUND SOLUTION OF BITARTRATE
 OF POTASSA.

R. Bitartrate of potassa, two drachms.
 Manna, two ounces.
 Water, eight ounces.
 Lemon juice, half an ounce.

Mix, and clarify with the white of an egg; in-
 fuse a small quantity of orange peel in the fluid,
 and strain on cooling. A pleasant laxative.

Taddei.

MIXTURE OF BITARTRATE OF POTASSA.

R. Bitartrate of potassa, two scruples.
 Antimonial wine,
 Vinegar of squill, each, half a drachm.
 Parsley water, one ounce and a half.
 Syrup of seneka, six drachms.

Mix. A teaspoonful every two hours, in dropsy
 supervening on scarlatina. *Vogt.*

R. Bitartrate of potassa, half an ounce.
 Barley water, two pounds.

Dissolve, and add

Brandy, one to two ounces.

Much praised as a diuretic in dropsies.

Milman.

R. Bitartrate of potassa, one ounce.
 Borax, two drachms.
 Boiling water, sufficient

to dissolve. To ten ounces of the cooled solu-
 tion add

Nitrate of potassa. two drachms.
 Oxymel, two ounces.

Mix. *Beasley.*

R. Bitartrate of potassa,
 Extract of hore-
 hound, each, two drachms.
 Balm water, three ounces.

Mix. One half to be taken morning and eve-
 ning, in hemorrhoids, with constipation.

Augustin.

IMPERIAL DRINK.

R. Bitartrate of potassa, half an ounce.
 Lemon, cut into slices, one.
 White sugar, half a pound.
 Water, three pints.

Mix. Let stand for half an hour, and strain.
 A pleasant, cooling drink. *Gray*

POTASSÆ ET AMMONIÆ
 TARTRAS.

TARTRATE OF POTASSA AND AMMONIA.

R. Bitartrate of potassa, one pound.
 Boiling water, sufficient

to dissolve; add gradually,

Solution of ammonia, sufficient
to saturate the excess of the acid; filter, and
evaporate, so that crystals may form.

Hamb. Ph.

SOLUTION OF TARTRATE OF POTASSA AND AMMONIA.

R. Solution of carbonate of
ammonia, six ounces.
Add, gradually,
Bitartrate of potassa, sufficient
to saturate; let stand for some hours, and filter.
Diuretic, in doses of a drachm. Leipsic Ph.

MIXTURE OF TARTRATE OF POTASSA AND AMMONIA.

R. Tartrate of potassa and
ammonia, half an ounce.
Fennel water, six fl. ounces.
Extract of dandelion,
Clarified honey, each, one ounce.
Mix. A spoonful every hour, as a diuretic.
Phæbus.

PRINOS.

BLACK ALDER.

This, which is also known as *Winter berry*, is an indigenous shrub, found in most parts of the country, principally in low, moist situations; flowering in June, and bearing numerous scarlet berries, which remain on the bush after the fall of the leaves.

Sex. Syst. Hexand. monog. Nat. Syst. Aquifoliaceæ.

Linn. Sp. Pl. 471. Griffith, Med. Bot. 434.

The officinal portion is the bark, which, when dried for use, is in slender pieces, of a greenish-white color internally, and of an ash-grey, mixed with brown, externally; brittle, inodorous, and of a bitter, astringent taste. It is tonic and astringent, and has been used with success in diarrhœa, intermittent fevers, &c.; The dose is from thirty grains to a drachm, three or four times a-day.

DECOCTION OF BLACK ALDER.

R. Black alder, two ounces.
Boiling water, three pints.
Boil down to a quart, and strain. One gill,
every two hours. W. P. C. Barton.

PRUNUM.

PRUNES.

Prunes are the dried fruit of various varieties of *Prunus domestica*, or cultivated Plum tree. They are principally derived from the south of

France, and are much used as an article of dessert; but are also employed in medicine, as a laxative, either alone or in combination, especially with senna, as in the confection of senna, &c., in the form of pulp.

PULP OF PRUNES.

R. Prunes, at will.
Soften the prunes in the vapor of boiling water, and having removed the stones, beat the remainder in a mortar, and pass it through a hair sieve.
U. S. Ph.

PRUNUS VIRGINIANA.

WILD-CHERRY BARK.

This is the bark of *Cerasus serotina*, though the U. S. Pharm. still retains the old name of the tree as the designation for the medicine. The *Cerasus serotina* is found in most parts of the United States, and in some situations attains a very large size.

Sex. Syst. Icosand. monog. Nat. Syst. Drupacæ.

De Candolle, Prod. ii. 540. Griffith, Med. Bot. 288.

The bark of both the root and branches is used, but the former is to be preferred. As dried for use, it is in pieces of various sizes, deprived of epidermis; of a reddish-brown color, and readily pulverized; when fresh it has an odor of bitter almonds, which is much diminished on drying, but reappears on maceration in water; the taste is bitter and aromatic. It is tonic, but it also exercises a sedative action on the circulatory and nervous systems, and is much used in a variety of diseases. The dose in powder is half a drachm to two drachms.

INFUSION OF WILD-CHERRY BARK.

R. Wild-cherry bark,
bruised, half an ounce.
Cold water, one pint.
Macerate for twenty-four hours, and strain. It may also be prepared by displacement.
U. S. Ph.

Dose, two fl. ounces, three or four times a-day.

R. Powdered wild-cherry bark, one ounce.
Orange peel, two drachms.
Water, one pint.

Macerate the bark for six hours, and then add the orange peel. Dose, a wineglassful. Ellis.

SYRUP OF WILD-CHERRY BARK.

R. Powdered wild-cherry
bark, four ounces.
Water, twelve fl. ounces.

Macerate for two days; put in a displacement apparatus; add water till twelve fl. ounces are obtained, returning the first portions till it comes away clear, then add

Sugar, twenty-four ounces,
and form syrup. *W. Procter.*

Dose, from a fl. drachm to a fl. ounce.

R. Same as the above, but with half the quantity of sugar.

D. S. Jones.

R. Wild-cherry bark, in coarse powder, five ounces.
Sugar, two pounds.
Water, sufficient.

Macerate the bark for twenty-four hours in a close vessel, transfer to a percolator, and gradually pour on water until a pint of filtered liquor is obtained. To this add the sugar, in a bottle, and shake occasionally until dissolved.

U. S. Ph.

Dose, a tablespoonful.

PYRETHRUM.

PELLITORY.

This is the root of *Anacyclus pyrethrum*, a small plant with perennial roots and annual stems; a native of the countries bordering on the Mediterranean.

Sex. Syst. Syngen. super. *Nat. Syst.* Astera-ceæ.

De Candolle, Prod. vi. 15. Griffith, Med. Bot. 402.

The dried root is about the size of the little finger, with a thick, brown bark, marked with black, shining points; it is inodorous; its taste is at first slight, but afterwards extremely acrid, leaving a burning, pricking sensation in the mouth. It is a powerful local irritant, and is principally used as a masticatory and sialagogue; but has also been employed in intermittents, palsies, &c.

LOZENGES OF PELLITORY.

R. Powdered pellitory,
" mastich, each, one drachm.

Mucilage of tragacanth, sufficient.

Mix, and make lozenges of twelve grains each.
As a masticatory in toothache. *Foy.*

TINCTURE OF PELLITORY.

R. Pellitory, one part.
Alcohol, four parts.

Macerate for eight days, and filter. *Par. Cod.*

COMPOUND TINCTURE OF PELLITORY.

R. Pellitory, four drachms.
Camphor, three drachms.
Opium, one drachm.
Oil of cloves, two drachms.
Alcohol, six fl. ounces.

Macerate for eight days, and filter. As a sialagogue in toothache. *Brande.*

R. Pellitory, one ounce.
Para cress flowers, four ounces.
Italian elecampane leaves, one ounce.
Alcohol, eight fl. ounces.

Macerate for fifteen days, express, and filter. Much celebrated as an odontalgic, under the name of *Paraguay Roux*. *Gray.*

LINIMENT OF PELLITORY.

R. Tincture of pellitory, six fl. drachms.
Camphorated oil, half a fl. ounce.
Water of ammonia, half a fl. drachm.

Mix. As a lotion for chilblains, and in rheumatic pains. *Radius.*

PLASTER OF PELLITORY.

R. Yellow wax, three ounces.
Turpentine, two ounces and a half.

Melt together, and add

Ammoniac, }
Sagapenum, } each,
Galbanum, }
Powdered pellitory, } one ounce.
" mustard, }

Mix well. As a rubefacient application in rheumatism, &c. *Cadet.*

EXTRACT OF PELLITORY.

R. Pellitory, ground, one pound.
Alcohol, two pints.
Ether, half a pint.

Mix the ether with a pint of the alcohol, pour it gradually on the powder, and put it in a displacer; when it ceases to drop, add the remainder of the alcohol, and finally sufficient diluted alcohol, to displace two pints and a half in all. Allow the ethero-alcoholic tincture to evaporate spontaneously, or with a moderate heat, until a soft extract is obtained.

Employed to destroy the sensibility of the nerves of teeth, previous to plugging, or for toothache. *W. Procter*

Q.

QUASSIA.

QUASSIA

Under the name of Quassia, the wood of two different plants has been used,—that of *Q. amara*, and of *Q. excelsa*, the latter being alone recognized in the U. S. Pharm., under the name of *Simaruba excelsa*. This is a large tree found in many of the West India Islands, and known as the *bitter ash*.

Sex. Syst. Pentand. monog. *Nat. Syst.* Simarubaceæ.

Lindley, Flor. Med. 208. Griffith, Med. Bot. (Picræna), 200.

It is found in commerce, in billets of various sizes, having a smooth, brittle bark. The wood is white, but becomes yellowish on exposure to the air; it has scarcely any smell, but possesses an intense, permanent bitter taste. It is a pure and powerful tonic, and is much used for that purpose. It is seldom given in substance.

INFUSION OF QUASSIA.

R. Rased quassia, two drachms.
Cold water, one pint.

Macerate for twelve hours, and strain.

U. S. Ph.

Dose, two fl. ounces, three or four times a-day.

COMPOUND INFUSION OF QUASSIA.

R. Quassia,
Virginia snakeroot,
Orange-peel, each, half an ounce.
Boiling water, two pints.

Infuse and strain. A teacupful, cold, three times a-day. *Ellis.*

EXTRACT OF QUASSIA.

R. Rased quassia, one pound.
Water, sufficient.

Mix the quassia with a pint of water, let stand for twenty-four hours, transfer to a percolator, and pour water upon it until exhausted. Heat the filtered liquor to the boiling point, strain, and evaporate to the proper consistence.

U. S. Ph.

Dose, three to five grains. A very efficient, bitter tonic.

TINCTURE OF QUASSIA.

R. Rased quassia, two ounces.
Diluted alcohol, two pints.

Macerate for fourteen days, express, and filter; or it may be made by displacement. *U. S. Ph.*

Dose, two fl. drachms.

COMPOUND TINCTURE OF QUASSIA.

R. Bruised cardamom,
" cochineal, each, half an ounce.
Powdered cinnamon,
Chipped quassia, each, six drachms.
Raisins, seven ounces.
Diluted alcohol, (Imp.) two pints.

Digest for seven days, strain, express residue, and filter. Dose, one or two fl. drachms.

Ed. Ph.

WINE OF QUASSIA.

R. Rased quassia, half an ounce.
Orange-peel, two drachms.
Wine, one pint and a half.

Macerate for twenty-four hours, express, and filter. Dose, three fl. drachms to a fl. half ounce, twice a-day. *Radius.*

QUERCUS.

OAK BARK.

Several species of oak are recognized as official in the Pharmacopœias, viz., the *Q. alba*, or *White oak*, and *Q. tinctoria*, or *Black oak*, in the U. S.; *Q. pedunculata*, *European White oak*, in the Lond. and Edin.; and *Q. robur*, *European oak*, in the Dublin Pharmacopœia.

Sex. Syst. Monœc. polyand. *Nat. Syst.* Corylaceæ.

Griffith, Med. Bot. 585.

The part used is the bark; this, in all the species, is astringent and tonic, and has been used in a variety of diseases, but more generally as an external application than as an internal remedy. The dose is from thirty grains to a drachm.

COMPOUND POWDER OF OAK BARK.

R. Powdered oak bark, one scruple.
Calamus,
Gentian, each, five grains.

Mix. To be taken every three hours, in apyrexia of intermittents. *Augustin.*

EXTRACT OF OAK BARK.

R. Powdered oak bark, one pound.
Water, one gallon.

Boil down to one-half, express, and filter. Evaporate at a heat of 200° F., until it begins to thicken, then reduce by a heat of 100° F., to the proper consistence. *Dub. Ph., 1826.*

The dose is from ten to forty grains.

DECOCTION OF OAK BARK.

R. Oak bark, one ounce and a half.
Water, two pints.
Boil down to one pint, and strain. *Dub. Ph.*

The *U. S. Ph.*, for decoction of white oak bark, orders one ounce of the bark, and water a pint and a half, to be boiled down to a pint. The dose is a wineglassful, as an astringent in chronic diarrhœa, &c.

GARGLE OF OAK BARK.

R. Decoction of oak bark, one pint.
Add

Alum, half a drachm.
Brandy, two fl. ounces.

As a gargle in chronic sore throat, with relaxed uvula; and as an injection in leucorrhœa. *Ellis.*

CATAPLASM OF OAK BARK.

R. Powdered oak bark, }
" catechu, } each, one ounce.
Barley meal,
Water, sufficient.

Boil to the proper consistence. *Foy.*
As an application in gangrene and mortification.

CONFECTION OF ACORNS.

R. Powdered acorns, three ounces.
" red coral,
" catechu, each,
one ounce and a half.

Confection of dog rose, ten ounces.
Syrup of red roses, sufficient.

Mix. One drachm every four hours, in chronic diarrhœa. *Bories.*

ACORN COFFEE.

R. Acorns, roasted, six drachms.
Coffee, roasted, two drachms.
Water, one pint.

Boil for a short time, and strain. Two or three cupfuls a-day, in scrofula and rachitis. *Augustin.*

QUINIA.

QUININE. QUINIA.

R. Sulphate of quinia, one part.
Boiling water, thirty parts.

Dissolve. Add water of ammonia sufficient to precipitate the quinia, wash the precipitate in warm distilled water, and dry it.

TINCTURE OF QUINIA.

R. Quinia, one part.
Alcohol, seven parts.
Dissolve. Dose, twenty to forty drops. *Beral.*

IMPURE QUINIA.

R. Yellow bark, one hundred parts.
Muriatic acid, five parts.
Water, five hundred parts.

Boil, decant, and repeat process three times with the same proportions of acid and water; unite decoctions, precipitate with milk of lime, wash and dry the precipitate, treat it several times with boiling alcohol, mix, and filter the solutions, and distil off the spirit. *Trousseau.*

This is said to be as active as the sulphate, and to be less unpleasant to the taste. It is given in the same doses.

TINCTURE OF IMPURE QUINIA.

R. Impure quinia, one ounce.
Alcohol,
Distilled water, each, twelve fl. ounces.
Mix. *Piorry.*
Dose, a tablespoonful.

AMORPHOUS QUINIA, OR CHINOIDINE.

R. Mother waters of sulphate of quinia, at will.
Solution of carbonate of potassa, sufficient to precipitate. Wash and dry this precipitate; dissolve in sulphuric ether, decant, and evaporate by a gentle heat. *Neligan.*

QUINIÆ ACETAS.

ACETATE OF QUINIA.

R. Quinia, two parts.
Distilled water, three parts.
Heat, and add as much acetic acid as will dissolve the quinia, and render the solution somewhat acid. Filter whilst boiling, and set aside to crystallize. *Cottreau.*

Acts like the other salts of quinia, and is in no way superior to them. Dose, one to ten grains, according to circumstances.

QUINIÆ ARSENIAS.

ARSENIATE OF QUINIA.

R. Arsenic acid, one drachm and a half.
Distilled water, six fl. ounces.
Quinia, five drachms.

Boil till solution takes place. Filter, let crystallize, and purify by recrystallization. In intermittents; dose, one-tenth to one-fourth of a grain. *Bourieres.*

QUININÆ ARSENIS.

ARSENITE OF QUINIA.

R. Sulphate of quinia, five hundred grains.
Water, acidulated with sulphuric acid, sufficient to dissolve.

Precipitate by solution of ammonia; collect, wash, and press the precipitate. Dissolve it in eight fl. ounces of alcohol, and add seventy-two grains of arsenious acid, heat together, and filter. The arsenite crystallizes on cooling. *Soubéiran.*

DI-ARSENITE OF QUINIA.

R. Arsenious acid, ten grains.
Carbonate of potassa, five grains.
Distilled water, five fl. drachms.

Boil for half an hour, adding water to make up the loss by evaporation, so that each fl. drachm may contain two grains of arsenic. Add

Sulphate of quinia, two scruples; previously dissolved in boiling water. Collect the precipitate on a filter, wash and dry it.

Dose, one-third of a grain, in chronic cutaneous affections. *Kingdon.*

QUININÆ CITRAS.

CITRATE OF QUINIA.

R. Quinia, two parts.
Distilled water, three parts.

Heat, and add sufficient citric acid to acidulate the mixture; when a perfect solution is effected, filter, and set aside to crystallize. *Magendie.*

The dose is the same as that of the sulphate: it is used in similar cases.

SYRUP OF CITRATE OF QUINIA.

R. Acid citrate of quinia, thirty six grains.
Simple syrup, one pint.

Mix. Half fl. ounce to one fl. ounce, in the twenty-four hours. *Magendie.*

QUININÆ ET FERRI CITRAS.

CITRATE OF QUINIA AND IRON.

(See *Ferri et Quinix Citras*, Page 233.)

QUININÆ FERROCYANAS.

FERROCYANATE OF QUINIA.

R. Sulphate of quinia, one hundred parts.
Ferrocyanide of potassium, thirty-one parts.
Distilled water, twenty-five hundred parts.

Boil for a few minutes, then let the solution cool; separate the oily compound, and wash it with a little water. Dissolve it in boiling alcohol, and let crystallize. *Par. Cod.*

Said to be more efficacious than any other salt of quinia. Dose, about the same as the sulphate.

MIXTURE OF FERROCYANATE OF QUINIA.

R. Ferrocyanate of quinia, four grains.
Alcohol, one fl. drachm.

Dissolve, and add

Camphor water, seven fl. drachms.

To be taken as required, shaking the phial.

Donovan.

PILLS OF FERROCYANATE OF QUINIA.

R. Ferrocyanate of quinia, twenty-four grains.
Mucilage of gum Arabic, sufficient.

Mix, and make twelve pills. Two for a dose.

Donovan.

QUININÆ HYDRIODAS.

HYDRIODATE OF QUINIA.

IODIDE OF QUINIA.

R. Add, by drops, a solution of twenty-four parts of iodide of potassium, in eight parts of water, to a strong solution of twenty parts of sulphate of quinia; wash the precipitate quickly, and dry in the shade.

Righini.

In obstinate intermittents and scrofulous affections.

BIN-IODIDE OF QUINIA.

R. Sulphate of quinia, one part.
Dissolve in boiling water, and add

Iodide of potassium, two parts, dissolved in water. Evaporate on a sand-bath to one-third, and allow the residue to cool; separate and preserve the resinous deposit.

Used in scrofulous enlargements of the glands. Dose, half a grain to one grain. *Kingdon.*

QUININÆ ET FERRI IODIDUM.

IODIDE OF QUINIA AND IRON.

R. Pour a strong solution of acid sulphate of quinia into a fresh solution of iodide of iron; collect the precipitate; dry by pressing between blotting paper, and keep it from the air.

Bouchardat.

QUININÆ HYDRIODAS IODURETA.

IODURETTED HYDRIODATE OF QUINIA.

R. Add a solution of iodide of iron, containing a slight excess of iodine, to an acid solution of quinia. Treat the precipitate with boiling alcohol, and filter whilst hot, and set aside to crystallize.
Bouchardat.

These preparations are useful where an alterative and tonic are required.

QUININÆ ET HYDRARGYRI CHLORIDUM.

CHLORIDE OF QUINIA AND MERCURY.

R. Corrosive sublimate, one part.
Muriate of quinia, three parts.

Dissolve separately, in the smallest quantity of water; mix the solutions; collect the precipitate, and dry by a gentle heat. *McDermott.*

PILLS OF CHLORIDE OF QUINIA AND MERCURY.

R. Double chloride of quinia and mercury, fifteen grains.
Opium, six grains.
Crumb of bread, sufficient.

Mix, and form thirty pills. One, three times a-day, to produce salivation. *Hamilton.*

QUININÆ KINAS.

KINATE OF QUINIA.

R. Alcoholic solution of sulphate of quinia, at will.
Aqueous solution of kinate of lime, sufficient
to precipitate; filter, evaporate, redissolve, and crystallize. *Magendie.*

PILLS OF KINATE OF QUINIA.

R. Kinate of quinia,
Powdered black pepper,
Extract of worm-wood, each, one drachm.

Mix, and make sixty pills. Three, every two or three hours, in obstinate intermittents. *Ronander.*

QUININÆ LACTAS.

LACTATE OF QUINIA.

R. Lactic acid, at will.
Quinia, sufficient

to saturate; leave the solution to evaporate spontaneously, in a shallow vessel, in a warm room, till crystals are formed.

Dose, three to ten grains. *Bouchardat.*

PILLS OF LACTATE OF QUINIA.

R. Lactate of quinia, half a drachm.
Extract of juniper, sufficient.
Mix, and make twenty pills. Two to six a-day, in intermittents. *Bouchardat.*

MIXTURE OF LACTATE OF QUINIA.

R. Lactate of quinia, seven grains.
Mint water, five fl. drachms.
Syrup of cloves, one fl. ounce.
Water, three and a half fl. ounces.
Mix. To be taken in divided doses during the apyrexia of intermittents. *Bouchardat.*

SYRUP OF LACTATE OF QUINIA.

R. Lactate of quinia, fifteen grains.
Dissolve in
Water, one fl. ounce
Add
Sugar, two ounces.
Form syrup. A teaspoonful, in the intermittents of children. *Bouchardat.*

QUININÆ MURIAS.

MURIATE OF QUINIA.

R. Sulphate of quinia, one ounce and a half.
Chloride of barium, half an ounce.

Dissolve separately, in boiling distilled water; mix the solutions, filter, and evaporate to crystallization. *Guibourt.*

R. Chloride of barium, five drachms.
Boiling water, one pint.

Dissolve, and gradually add
Sulphate of quinia, two ounces.
Boil for a few minutes, filter whilst hot, and dry the crystals. *Prus. Ph.*

R. Diluted muriatic acid, at will.
Quinia, sufficient

to saturate; evaporate, and crystallize. Dose, from half a grain to a grain. *Cottureau*

MIXTURE OF MURIATE OF QUINIA.

R. Muriate of quinia, twelve grains.
Diluted muriatic acid, five minims.
Distilled water, seven fl. ounces.
Syrup of orange flowers, one fl. ounce.
Mix. Dose, one fl. ounce. *Neligan.*

- R. Muriate of quinia, eight grains.
 Fennel water, five fl. ounces.
 Muriatic ether, one fl. drachm.
 Sugar, half an ounce.

Mix. A tablespoonful every two hours.

Radius.

- R. Muriate of quinia, one scruple.
 Peppermint water, half a fl. ounce.

Mix. Twenty to sixty drops, every two hours, in intermittents of children.

Radius.

QUININÆ NITRAS.

NITRATE OF QUINIA.

- R. Diluted nitric acid, at will.
 Quinia, sufficient
 to saturate; boil with animal charcoal, filter,
 evaporate, and let crystallize. *Taddei.*

QUININÆ PHOSPHAS.

PHOSPHATE OF QUINIA.

- R. Quinia, two parts.
 Water, three parts.
 Boil, and add
 Phosphoric acid, sufficient
 to saturate; filter while hot, and let crystallize. *Turin. Ph.*

QUININÆ SULPHAS.

SULPHATE OF QUINIA.

- R. Yellow bark, in coarse
 powder, four pounds.
 Muriatic acid, three fl. ounces.
 Lime, in powder, five ounces.
 Water, five gallons.
 Sulphuric acid, }
 Alcohol, } each, sufficient.
 Animal charcoal, }

Boil the bark in one-third of the water, with one-third of the muriatic acid, and strain through linen. Repeat this process twice. Mix the decoctions, and, whilst hot, gradually add the lime, mixed with two pints of water, stirring constantly, till the quinia is precipitated. Wash precipitate with distilled water, press, dry, and digest in boiling alcohol; decant, and repeat till alcohol is no longer rendered bitter. Mix the liquors, and distil off the alcohol, till a brown, viscid mass remains; add to this half a gallon of distilled water, heat to boiling, and add as much sulphuric acid as will dissolve the impure quinia: then add one ounce and a half of ani-

mal charcoal, boil for two minutes, filter while hot, and set aside to crystallize. If the solution be entirely neutral, acidulate slightly with sulphuric acid; if too acid, add more animal charcoal. Separate the crystals, dissolve them in boiling water a little acidulated with sulphuric acid, add a little animal charcoal, and recrystallize; place the crystals in bibulous paper, and dry by a gentle heat. Treat the mother waters with solution of ammonia, and proceed with the precipitate as before. *U. S. Ph.*

POWDER OF SULPHATE OF QUINIA.

- R. Sulphate of quinia, three to twelve grains.
 White sugar, two drachms.
 Mix, and divide into six powders. *Radius.*

POWDER OF SULPHATE OF QUINIA AND TARTAR EMETIC.

- R. Sulphate of quinia, ten grains.
 Tartar emetic, three grains.
 Mix, and divide into six powders. One, every two hours, in the apyrexia of obstinate intermittents. *Gola.*

POWDER OF SULPHATE OF QUINIA AND SODA.

- R. Sulphate of quinia, one to two grains.
 Carbonate of soda, four to five grains.
 Sugar, one scruple.
 Mix, and divide into six powders. One, morning and evening, in scrofulous ophthalmia. *Ammon.*

POWDER OF SULPHATE OF QUINIA AND TARTARIC ACID.

- R. Tartaric acid, fifteen grains.
 Sulphate of quinia, one grain and a half.
 Mix, and add
 Bicarbonate of soda, eighteen grains.
 Sugar, half a drachm.
 Mix in water, for one dose. In the apyrexia of intermittents. *Meireu.*

POWDER OF SULPHATE OF QUINIA AND MORPHIA.

- R. Sulphate of quinia, two to six grains.
 " morphia, half to one grain.
 Mix, and divide into four powders. *Magendie.*

COMPOUND POWDER OF SULPHATE OF QUINIA.

- R. Sulphate of quinia, half a grain.
 Powdered fox-glove, a quarter to one grain.

Powdered fennel, six grains.
 Sugar of milk, ten grains.
 Mix. To be taken three or four times a-day, in the hectic fever of phthisis. *Gunther.*

R. Sulphate of quinia, three grains.
 Opium, one grain.
 Gum Arabic
 Sugar, each, six grains.

Make a powder. To be taken just before the paroxysm of malignant intermittents. *Neumann.*

R. Sulphate of quinia, two grains.
 " iron, one grain.
 Powdered fennel, one scruple.
 Oil of chamomile, one drop.

Mix. To be taken every three hours, in obstinate intermittents. *Phæbus.*

R. Sulphate of quinia, quarter of a grain.
 Chocolate, seven grains.
 Sugar of milk, two grains.

Mix. To be taken every three hours, in debility of the stomach. *Kopp.*

PILLS OF SULPHATE OF QUINIA.

R. Sulphate of quinia, one ounce.
 Powdered gum Arabic, two drachms.
 Honey, sufficient.

Mix the sulphate and gum, and beat with the honey, into a mass, and divide into four hundred and eighty pills. *U. S. Ph.*

Each pill contains one grain of the sulphate.

R. Sulphate of quinine, fifteen grains.
 Extract of chamomile, fifteen grains.

Mix, and make six pills. To be taken during the apyrexia of intermittent fevers. *Elliotson.*

R. Sulphate of quinia, ten grains.
 Extract of wormwood, sufficient.

Beat into a mass, and make six pills. *Doreault.*

R. Sulphate of quinia, twelve grains.
 Powdered tragacanth, one grain.

Rub together, add sufficient water to form a plastic mass, and divide into six pills.

E. Parrish.

R. Sulphate of quinia, twenty grains.
 Elixir of vitriol, fifteen drops.

Drop the acid into the sulphate, and rub with a spatula until the mass assumes a pilular consistence. Make six pills. *E. Parrish.*

PILLS OF SULPHATE OF QUINIA AND GENTIAN.

R. Sulphate of quinia, one scruple.
 Extract of gentian, two scruples.

Mix, and make twenty pills.

COMPOUND PILLS OF SULPHATE OF QUINIA.

R. Calomel, six grains.
 Powdered opium, three grains.
 Sulphate of quinia, twelve grains.
 Syrup, sufficient.

Beat into mass, and divide into twelve pills. One, night and morning, as an alterative. *Ellis.*

R. Blue pill mass, } each,
 Sulphate of quinia, } twelve grains.
 Powdered aloes, }
 Aromatic syrup of rhubarb, sufficient.

Beat into mass, and divide into twelve pills. One, three or four times a-day. This, or the foregoing combination, is suited to the condition following bilious remittent or intermittent fevers. *Ellis.*

R. Sulphate of quinia, twelve grains.
 Extract of gentian, one scruple.
 Compound rhubarb pill, two scruples.
 Blue pill mass, six grains.

Mix, and make twelve pills. One, three times a-day. *Ryan.*

R. Sulphate of quinia, ten to fifteen grains.

Dry phosphoric acid, two scruples.
 Powdered mallow root, four scruples.
 Extract of centaury, sufficient.

Mix, and make sixty pills. Three or four pills, two or three times a-day, in nervous debility with tendency to abortion. *Radius.*

R. Powdered camphor, one scruple.
 Sulphate of quinia, two scruples.
 Pill of aloes and

myrrh, one drachm and a half.
 Syrup of ginger, sufficient.

Mix, and make forty pills. One, twice a-day. *Copland.*

SYRUP OF QUINIA.

R. Sulphate of quinia, thirty-two grains.
 Distilled water, two fl. drachms.
 Aromatic sulphuric acid, six drops.

Dissolve, and add to

Syrup, sixteen fl. ounces. *Par. Cod.*

R. Sulphate of quinia, sixteen grains.
 Syrup of ginger, two fl. ounces

Mix. Dose, a teaspoonful. *Ellis.*

SYRUP OF QUINIA AND COFFEE.

R. Ground roasted coffee, four ounces.
 Boiling water, two pints, four fl. ounces

Make an infusion, let cool, and add

Sugar, four pounds.
Dissolve by means of a water-bath, and add
Sulphate of quinia, one drachm,
dissolved in a little water, acidulated with sulphuric acid. *Bories.*

MIXTURE OF SULPHATE OF QUINIA.

R. Sulphate of quinia, twenty grains.
Sulphuric acid, one drop.
White sugar, one drachm.
Cinnamon water, two fl. ounces and a half.

Mix. A teaspoonful every hour, in the apyrexia of intermittents. *Ellis.*

MIXTURE OF SULPHATE OF QUINIA AND COFFEE.

R. Strong infusion of coffee, five fl. ounces.

Sulphate of quinia, twenty-four grains.

Sugar, four drachms.

Mix. Dose, a tablespoonful. The coffee conceals the bitterness of the quinia. *Beasley.*

TINCTURE OF QUINIA.

R. Sulphate of quinia, one scruple.
Alcohol, half a fl. ounce.

Dissolve. Dose, ten to twenty drops. *Ellis.*

COMPOUND TINCTURE OF QUINIA.

R. Sulphate of quinia, forty-eight grains.
Compound tincture of orange-peel,

five and a half fl. ounces.

Elixir of vitriol, forty-five drops.

Mix. Dose, half a fl. drachm to two fl. drachms. *Copland.*

R. Sulphate of quinia, three hundred and twenty grains.

Tincture of orange-peel, forty fl. ounces.

Digest for seven days, and filter. Each fl. drachm contains one grain of the sulphate. *Lond. Ph.*

WINE OF QUINIA.

R. Sulphate of quinia, twelve grains.
Madeira wine, two pints.

Dissolve. Dose, one to two fl. ounces. *Magendie.*

AROMATIC WINE OF QUINIA.

R. Sulphate of quinia, eighteen grains.
Citric acid, fifteen grains.
Orange wine, twenty-four fl. ounces.

Mix. *Collier.*

LINIMENT OF SULPHATE OF QUINIA.

R. Sulphate of quinia, half a drachm.
Tartar emetic, six grains.
Extract of opium, twelve grains.
Spirit of camphor, eighteen fl. drachms.

Mix. Three fl. drachms to be rubbed on the epigastrium, three times a-day, in intermittents. *Schuster.*

OINTMENT OF SULPHATE OF QUINIA.

R. Sulphate of quinia, one ounce.
Alcohol, sufficient
Sulphuric acid, each, to dissolve.

Lard, four ounces.

Rub together. Half an ounce to be rubbed into the groins, in malignant intermittents. *Antonini.*

PLASTER OF SULPHATE OF QUINIA.

R. Lead plaster, six drachms.
Resin, two drachms

Melt together, and add

Sulphate of quinia, one drachm

Oil of cajeput,

Camphor, each, one scruple.

Mix. To be applied to the epigastrium as a prophylactic in cholera. *Ammon.*

ENEMA OF SULPHATE OF QUINIA.

R. Sulphate of quinia, twelve to twenty grains.
Flaxseed tea, four fl. ounces.

Mix. To be used every four to six hours. *Ellis.*

DENTIFRICE OF SULPHATE OF QUINIA.

R. Sulphate of quinia, four grains.
Prepared coral, one ounce.
Carmine lake, eight grains.
Essence of myrrh, two drops.

Mix. *Pelletier.*

GARGLE OF SULPHATE OF QUINIA.

R. Sulphate of quinia, twelve grains.
" copper, sixteen grains.
Elixir of vitriol, one fl. drachm.
Water, eight fl. ounces.

Mix. To be used three or four times a-day, in obstinate sore throat. *Hartshorne.*

QUININÆ SULPHO-TARTRAS.

SULPHO-TARTRATE OF QUINIA.

R. Sulphate of quinia, four drachms.
Tartaric acid, four drachms and a-half.
Distilled water, two fl. ounces.

Dissolve. Half a fl. drachm to a drachm, in the course of the day. *Righini.*

MIXTURE OF SULPHO-TARTRATE OF QUINIA.

R. Sulphate of quinia, six grains.
Tartaric acid, three grains.
Syrup, one fl. ounce.
Mix. Dose, a teaspoonful. *Casorati.*

QUININÆ TANNAS.

TANNATE OF QUINIA.

R. Sulphate of quinia, one part.
Water, twenty parts.
Diluted sulphuric acid, sufficient.
Dissolve, filter, and add
Infusion of galls, sufficient
to precipitate; collect precipitate, wash and dry it. *Greek Ph.*

Found useful in intermittent neuralgia.

IMPURE TANNATE OF QUINIA.

R. Powdered Peruvian bark, one part.
Vinegar, six parts.
Macerate for twenty-four hours; then boil, and decant. Repeat the process with fresh vinegar. Mix the decoctions, filter when cold, and add infusion of galls as long as precipitation takes place. Collect, wash, and dry the precipitate.

Dose, similar to that of the sulphate. *Buchner.*

QUININÆ TARTRAS.

TARTRATE OF QUINIA.

R. Quinia, two parts.
Water, three parts.
Mix, boil, and add
Tartaric acid, sufficient
to dissolve the quinia. Filter whilst hot, and let crystallize. *Par. Cod.*

QUININÆ VALERIANAS.

VALERIANATE OF QUINIA.

R. Fresh precipitated quinia, three parts.
Valerianic acid, one part.
Water, sixty parts.
Mix; boil; filter while hot, and let crystallize; dry the crystals under 122° F. *Wettstein.*

This may also be prepared by double decomposition, between the muriate of quinia and the valerianate of soda.

Said to be more powerful than the sulphate, and not to cause nervous symptoms. It is given in solution, pill, enema, &c.

PILLS OF VALERIANATE OF QUINIA.

R. Valerianate of quinia, twelve grains.
Powdered tragacanth, six grains.
Water, sufficient.

Rub together, and divide into eight pills.

Dose, one pill every hour, in hemicrania.

Thomas

R.

RESINA.

RESIN.

Resin, or, as it is generally called, *Rosin*, is the residuum after the distillation of the volatile oil from various species of the Pine tribe. In this state it is *Yellow resin*, or *Colophony*, which, when melted and agitated with water, becomes of a whitish color, and forms *White resin*. Resin, when pure, is yellowish-brown, inclining to olive or brown; it is solid, brittle, of a smooth and shining fracture; of faint odor, and a somewhat acrid taste. It is principally used in medicine as an ingredient in plasters and ointments.

HÆMOSTATIC POWDER.

R. Powdered resin, four parts.
" gum Arabic,
" charcoal, each, one part.

Mix. As an application to check external bleeding. *Guibourt.*

R. Resin, two parts.
Gum Arabic, one part.
Charcoal, half a part.

Pulverize well, and mix. Has proved useful in checking bleeding from wounds, &c. *Bonnefoux.*

RESIN CERATE, OR BASILICON OINTMENT.

R. Resin, five ounces
Lard, eight ounces.
Yellow wax, two ounces.

Melt together; strain through linen; and stir till cool. *U. S. Ph.*

As a stimulant application to blistered surfaces, burns, ulcers, &c.

COMPOUND RESIN CERATE.

R. Resin,	} each,	one pound.
Suet,		
Yellow wax,		
Turpentine,		half a pound.
Flaxseed oil,		half a pint.

Melt together; strain through linen, and stir till cold. *U. S. Ph.*

Known as *Deshler's Salve*; is rather more stimulating than the last.

RESIN, OR ADHESIVE PLASTER.

R. Powdered resin,	half a pound.
Lead plaster,	three pounds.

Melt the plaster by a gentle heat, and add the resin, mixing well. *U. S. Ph.*

PLASTER OF ST. ANDREW.

R. White resin,	eight ounces.
Elemi,	two ounces.
Venice turpentine,	
Oil of bay laurel, each,	one ounce.

Melt together, and strain. An adhesive plaster. *Guibourt.*

PLASTER OF VIGO WITH MERCURY.

R. Lead plaster,	two pounds, eight ounces.	
Yellow wax,		
Resin, each,	two ounces.	
Ammoniac,	} each,	five drachms.
Bdellium,		
Olibanum,		
Myrrh,		
Saffron,		three drachms.
Turpentine,		two ounces.
Liquid storax,		six ounces.
Mercury,		twelve ounces.
Oil of lavender,		two drachms.

Powder the gum-resins and saffron, and rub the mercury with the storax and turpentine, in an iron mortar, until globules disappear. Melt the plaster with the wax and resin, and add the powders and the oil of lavender. When cool, add the mercurial mixture and incorporate thoroughly. *Par. Cod.*

Used to prevent pitting in small-pox, by spreading it on linen or leather, and covering the exposed part.

RHAMNUS.

BUCKTHORN.

The berries and juice of the *R. catharticus*, are possessed of active purgative properties. This plant is a native of Europe, and is occasionally met with in this country. European practitioners hold it in high esteem.

Sex. Syst. Pentand. monog. *Nat. Syst.* Rhamnaceæ.

Linn. Sp. Pl. 279. *Griffith, Med. Bot.* 217.

Sometimes the berries are used, at others, their expressed juice. Dose of the dried berries one drachm.

EXTRACT OF BUCKTHORN.

R. Expressed juice of buckthorn berries,	at will.
--	----------

Permit the juice to undergo a slight fermentation, and then evaporate to the proper consistence. Dose, one scruple. *Beasley.*

SYRUP OF BUCKTHORN.

R. Juice of buckthorn berries, four pints.	
Sliced ginger,	
Powdered pimento, each,	six drachms.
Sugar,	four pounds.
Alcohol,	six fl. ounces.

Set the juice aside for three days, for the dregs to subside, and strain. Add the ginger and pimento to a pint of the juice, macerate four hours with a gentle heat, and strain. Boil down the remainder of the juice to a pint and a half. Mix the liquors, dissolve the sugar in them, and add the spirit. Dose, a tablespoonful. *Lond. Ph.*

RHEUM.

RHUBARB.

Rhubarb is the root of various species of *Rheum*, which grow in the deserts of Tartary, and in the central parts of Asia. Several species are cultivated in England, France, and Germany, yielding the *European variety* of rhubarb.

Sex. Syst. Enneand. trigyn. *Nat. Syst.* Polygonaceæ.

Lind. Fl. Med. 358. *Griffith, Med. Bot.* 539.

Three kinds of Rhubarb are usually found in commerce, the Russian or Turkey, the European, and the Chinese; the first of which is the most valuable, but the last is the most generally used, and, when good, it answers every purpose. All are purgatives, with some tonic and astringent powers, and are much used in various forms of disease. The dose, in substance, to produce a full effect, is from twenty to thirty grains. They are given in a vast variety of forms.

POWDER OF RHUBARB AND MAGNESIA.

R. Powdered rhubarb,	one scruple.
Magnesia,	ten grains.

Mix. To be given in syrup, or sugar and water. *Ellis.*

POWDER OF RHUBARB AND CHALK.

R. Powdered rhubarb, fifteen grains.
Compound chalk powder, twenty-four grains.
Mix. In the evening, in cardialgia. *Foy.*

POWDER OF RHUBARB AND SULPHATE OF POTASSA.

R. Powdered rhubarb, one drachm.
" sulphate of potassa, two drachms.
Mix. Ten grains to a drachm, every morning. *Fordyce.*

COMPOUND POWDERS OF RHUBARB.

R. Magnesia, }
Cream of tartar, } each, half an
Powdered rhubarb, } ounce.
" chamomile, }
Oleo-sacch. of fennel, }

Mix. A teaspoonful, twice or thrice a-day, in obstructions of the abdominal viscera. *Selle.*

R. Powdered rhubarb, thirty grains.
" sulphate of potassa, one drachm.
" chamomile, each,

Mix, and divide into six powders. One, twice a-day, in dyspepsia with torpor of the bowels. *Ellis.*

R. Powdered rhubarb, four ounces.
Magnesia, one pound.
Powdered ginger, two ounces.

Mix well, and keep in well-closed bottles. *Ed. Ph.*

A good antacid laxative. Dose, five grains to a drachm, according to age.

ROASTED RHUBARB.

R. Coarsely-powdered rhubarb, at will.
Place it in a shallow iron dish, heat regularly, so as to brown the powder, which should be so constantly stirred, that the influence of the heat should be uniform throughout; when the color has changed to brown, the process is to be stopped, and the rhubarb reduced to fine powder, for use. *W. Procter.*

R. Powdered rhubarb, at will.
Heat in an iron vessel, constantly stirring, till it becomes almost black, then smother it in a covered jar. Dose, five to ten grains, as an astringent in diarrhœa. *Hoblyn.*

PILLS OF RHUBARB.

R. Powdered rhubarb, six drachms.
Soap, two drachms.

Beat them with water, so as to form a mass, and divide into one hundred and twenty pills. *U. S. Ph.*

Each pill contains three grains of rhubarb.

COMPOUND RHUBARB PILLS.

R. Powdered rhubarb, one ounce.
" aloes, six drachms.
" myrrh, half an ounce.
Oil of peppermint, half a fl. drachm.
Water, sufficient.

Beat together into mass, and divide into two hundred and forty pills. *U. S. Ph.*

A warm, tonic laxative, useful in costiveness with debility of the stomach. Dose, two to four a-day.

R. Powdered rhubarb, one drachm and a half.
Sulphate of iron, half a drachm.
Soap, two scruples.
Distilled water, sufficient.

Beat into mass, and divide into forty pills. In similar cases as last, three or four to be taken at bedtime. *Griffitts.*

PILLS OF RHUBARB AND IPECACUANHA.

R. Powdered rhubarb, one scruple.
" ipecacuanha, ten grains.
Opium, three grains.
Oil of cinnamon, five drops.
Gum Arabic, sufficient.

Triturate together, and divide into ten pills. One to be given every two or three hours, in dysentery, to relieve tormina and tenesmus. *Chapman.*

PILLS OF RHUBARB AND CARAWAY.

R. Powdered rhubarb, two drachms.
Syrup, one fl. drachm.
Oil of caraway, ten minims.
Mix, and divide into forty pills. *Kitchener*

PILLS OF RHUBARB AND CHAMOMILE.

R. Powdered rhubarb, }
" aloes, } each,
" myrrh, } one drachm.
Extract of chamomile, }
Oil of chamomile, twelve drops.

Mix, and divide into sixty pills. These are known as *Speediman's pills*. *Beasley.*

R. Powdered rhubarb, }
" ginger, each, half a drachm.
Extract of chamomile, one drachm.

Beat together, and divide into thirty pills; three to be taken before each meal. Have been recommended in dyspepsia and chlorosis. *A. T. Thomson*

PILLS OF RHUBARB AND IRON.

R. Dried sulphate of iron, four parts.
 Extract of rhubarb, ten parts.
 Conserve of red roses, five parts.
 Beat into a mass, and divide into five-grain pills. Dose, two pills, as a tonic and laxative.

Ed. Ph.

PILLS OF RHUBARB AND SODA.

R. Powdered rhubarb, } each, one drachm
 Carbonate of soda, } and a half.
 Extract of gentian, }
 Mix, and make sixty pills. *Guy's Hosp.*
 R. Powdered rhubarb, } each, one
 Dried carbonate of soda, } scruple.
 Extract of gentian, }
 Calomel, three grains.
 Mix, and make twenty pills. Two, occasionally, in dyspepsia. *Ellis.*

PILLS OF RHUBARB AND OX GALL.

R. Inspissated ox gall,
 Ammoniac,
 Powdered rhubarb, equal parts.
 Mix, and form pills of two grains each. *Leipsic Ph.*
 In constipation, deficiency of bile, &c.

LOZENGES OF RHUBARB.

R. Powdered rhubarb,
 Cream of tartar, each, two drachms.
 Fresh orange peel, half a drachm.
 Sugar, dissolved in
 orange-flower water, four ounces.
 Mix, and make lozenges of eighteen grains. *Beral.*

SUPPOSITORY OF RHUBARB.

R. Extract of rhubarb, half a drachm.
 Soap, three drachms.
 Powdered rhubarb, sufficient.
 Mix, and make three suppositories. *Radius.*

ELECTUARY OF RHUBARB.

R. Powdered rhubarb,
 one drachm and a half.
 Sulphate of potassa, one drachm.
 Cream of tartar, half an ounce.
 Pulp of tamarinds, two ounces.
 Mix. A teaspoonful. *Saunders.*

INFUSION OF RHUBARB.

R. Bruised rhubarb, one drachm.
 Boiling water, half a pint.
 Infuse for two hours, and strain. *U. S. Ph.*

One or two fl. ounces, every three or four hours, till it operates.

R. Bruised rhubarb, one ounce.
 Boiling water, eighteen fl. ounces.
 Infuse for twelve hours, in a covered vessel; add

Spirit of cinnamon, two fl. ounces,
 and strain. Dose, as above. *Ed. Ph.*

ALKALINE INFUSION OF RHUBARB.

R. Bruised rhubarb, two drachms.
 Carbonate of potassa, one drachm.
 Boiling water, half a pint.
 Infuse for four hours, strain, and add
 Tincture of cinnamon, half a fl. ounce. *Copland.*

EXTRACT OF RHUBARB.

R. Powdered rhubarb, fifteen ounces.
 Proof spirit, one pint.
 Distilled water, seven pints.
 Macerate for four days by a gentle heat; strain; set aside to settle; decant, and evaporate to proper consistence. *Lond. Ph.*
 Dose, ten to thirty grains.

FLUID EXTRACT OF RHUBARB

R. Coarsely-powdered rhubarb,
 eight ounces.
 Sugar, five ounces.
 Tincture of ginger, half a fl. ounce.
 Oil of fennel,
 Oil of anise, each, four minims.
 Diluted alcohol, sufficient.
 Mix the rhubarb with an equal bulk of coarse sand, add twelve fl. ounces of diluted alcohol, and allow the mixture to stand for twenty-four hours. Place in a percolator, and add diluted alcohol until the rhubarb is exhausted. Evaporate the resulting tincture on a water-bath to five fl. ounces, add the sugar, dissolve it, and add the tincture of ginger, holding the oils in solution. *U. S. Ph.*

Dose, half a fl. drachm.

FLUID EXTRACT OF RHUBARB AND SENNA.

R. Coarsely-powdered senna,
 twelve ounces.
 " " rhubarb,
 four ounces.
 Bicarbonate of potassa, half an ounce.
 Sugar, eight ounces.
 Tincture of ginger, one fl. ounce.
 Oil of cloves, eight minims.
 " aniseed, sixteen minims.
 Water,
 Alcohol, each, sufficient.

Mix the senna and rhubarb, pour upon them two pints of diluted alcohol, and macerate for twenty-four hours. Place in a percolator, and add a mixture of one part of alcohol with three parts of water, until one gallon of tincture is obtained. Evaporate this, on a water-bath, to eleven fl. ounces; dissolve in it the sugar and bicarbonate, strain, and add the tincture of ginger, holding the oils in solution. The whole should measure one pint.

Dose, one teaspoonful.

W. Procter, Jr.

SYRUP OF RHUBARB.

R. Rhubarb, bruised, two ounces.
Boiling water, one pint.
Macerate for twenty-four hours, and strain; then add

Sugar, two pounds,
and make syrup. U. S. Ph., 1840.

R. Coarsely-powdered rhubarb, two ounces.
Alcohol, half a pint.
Water, one pint and a half.
Sugar, two pounds.

Mix the alcohol and water, and pour four fl. ounces on the rhubarb, previously mixed with coarse sand, and macerate for four hours. Transfer to a percolator, and add the remainder of the alcohol and water. Evaporate the resulting tincture to thirteen fl. ounces, add the sugar, and make a syrup. U. S. Ph., 1850.

Mild cathartic. Dose, for young children, one to two fl. drachms.

AROMATIC SYRUP OF RHUBARB.

R. Rhubarb, bruised, two ounces and a half.
Cloves, bruised,
Cinnamon, bruised, each, half an ounce.
Nutmeg, bruised, two drachms.
Diluted alcohol, two pints.

Macerate for fourteen days, and strain; evaporate on a water-bath to one pint, and add, (while it is still hot),

Syrup, six pints,
previously heated. U. S. Ph.

A warm laxative in cases of children. Dose, one fl. drachm, in bowel complaints, to be repeated every two hours.

SYRUP OF RHUBARB AND SENNA.

R. Bruised rhubarb, one ounce.
Senna, two ounces.
Fennel seed,
Bruised cinnamon, each, two drachms.
Boiling water, two pints and a half.
Macerate for twelve hours, strain, and add
Sugar, three pounds.
Make syrup. Ed. Ph., 1744.

TINCTURE OF RHUBARB.

R. Rhubarb, bruised, three ounces.
Cardamom, bruised, half an ounce.
Diluted alcohol, two pints.

Macerate for fourteen days, express, and filter. U. S. Ph.

Dose, as purgative, half to one fl. ounce; as stomachic and tonic, one to two fl. drachms.

TINCTURE OF RHUBARB AND GENTIAN.

R. Rhubarb, bruised, two ounces.
Gentian, bruised, half an ounce.
Diluted alcohol, two pints.

U. S. Ph.

Macerate for fourteen days, express, and filter. In flatulent colic, &c. Dose, as above.

R. Rhubarb, bruised, two ounces.
Gentian, bruised, half an ounce.
Virginia snakeroot,
one drachm and a half.

Diluted alcohol, two pints.

Macerate for three days, express, and filter.

Van Mons.

TINCTURE OF RHUBARB AND ALOES.

R. Rhubarb, bruised, ten drachms.
Aloes, powdered, six drachms.
Cardamom, bruised, half an ounce.
Diluted alcohol, two pints.

Macerate for fourteen days, express, and filter.

U. S. Ph.

This was formerly known as *Elixir sacrum*, and was much used.

TINCTURE OF RHUBARB AND SENNA.

R. Rhubarb, bruised, one ounce.
Senna, two drachms.
Coriander, bruised,
Fennel, bruised, each, one drachm.
Red saunders, rasped, two drachms.
Saffron,
Liquorice, each, half a drachm.
Raisins, stoned, half a pound.
Diluted alcohol, three pints.

Macerate for fourteen days, express, and filter.

U. S. Ph.

Well known and much used under the name of *Warner's gout cordial*, as a stomachic and slight purgative. Dose, from a half to one fl. ounce.

WARNER'S CORDIAL.

R. Contused rhubarb, one ounce.
Senna, half an ounce.
Saffron, one drachm.
Fennel seed,
Coriander, each, two drachms

Liquorice, four drachms.
 Raisins, stoned, one pound.
 Brandy, three pints.
 Mix, and macerate for a week, and strain.
 Dose, half a wineglassful, in atonic gout, with flatulence. *Chapman.*

TINCTURE OF RHUBARB AND ANISEED.

R. Rhubarb, bruised,
 Liquorice root, bruised, each, two ounces.
 Aniseed, bruised,
 Sugar, each, one ounce.
 Diluted alcohol, two pints.
 Macerate for fourteen days, express, and filter. *Copland.*

Used as the above tincture.

SWEET TINCTURE OF RHUBARB.

R. Coarsely-powdered
 rhubarb, eight ounces.
 Powdered liquorice root, three ounces.
 Bruised aniseed, three ounces.
 " cardamom, half an ounce.
 " orange peel, one ounce.
 Diluted alcohol, sufficient
 to make one gallon of tincture, by displacement.

Dose, a tablespoonful. *A. B. Taylor.*

R. Bruised rhubarb, one ounce and a half.
 Bruised liquorice root,
 " aniseed, each, six drachms.
 Sugar, one ounce and a half.
 Diluted alcohol, two pints.
 Macerate for two weeks, and filter. Or prepare the tincture by displacement. *D. S. Jones.*

ALKALINE TINCTURE OF RHUBARB.

R. Rhubarb, bruised, one ounce and a half.
 Carbonate of potassa, three drachms.
 Boiling water, twelve fl. ounces.
 Macerate for twelve hours, strain, and add
 Tincture of cinnamon, two fl. ounces. *Prus. Ph.*

A mild purgative, where there is acidity of stomach, in doses of one-half to two fl. ounces.

WINE OF RHUBARB.

R. Rhubarb, bruised, two ounces.
 Canella, bruised, one drachm.
 Diluted alcohol, two fl. ounces.
 Wine, one pint.

Macerate for fourteen days, occasionally agitating, express, and filter. *U. S. Ph.*

A warm, cordial laxative, in doses of one to four fl. drachms.

R. Rhubarb, bruised, two ounces and a half.
 Cardamom, bruised, half an ounce.
 Saffron, two drachms.
 Wine, two pints.
 Alcohol, eight fl. ounces.

Macerate for ten days, express, and filter.

Par. Cod.

Used as the last.

WINE OF RHUBARB AND GENTIAN.

R. Rhubarb, bruised, one ounce.
 Gentian, " two drachms.
 Canella, " one drachm.
 Wine, one pint.

Macerate for three days, express, and filter.

Swed. Ph.

COMPOUND WINE OF RHUBARB.

R. Rhubarb, bruised,
 Orange peel, bruised, each, half an ounce.
 Wine, two pints.

Macerate for twenty hours, strain, and add

Oleo-sacch. of mace, one ounce.

Hoffmann's anodyne, one drachm.

As a stomachic, a tablespoonful three times a-day. *Phæbus.*

RHUBARB MIXTURE.

R. Bruised rhubarb, one drachm.
 " fennel, one scruple.
 Sulphate of magnesia, half an ounce.
 Manna, one ounce.
 Boiling water, six fl. ounces.

Infuse and strain, when cold. As a mild purgative; one-half to be taken at a dose. *Radius.*

R. Infusion of rhubarb, one fl. drachm and a half.
 Spirit of ammonia, two fl. scruples.
 Cinnamon water, one fl. ounce and a half.
 Syrup of opium, six fl. drachms.

Mix. A teaspoonful every two hours, in the chronic diarrhœa of young children. *Vegt.*

R. Powdered rhubarb, one drachm.
 Carbonate of soda, two drachms.
 Tincture of orange peel, one fl. ounce and a half.
 Decoction of liquorice, ten fl. ounces and a half.

Mix. Dose, half a fl. ounce to one fl. ounce, three times a-day. *Guy's Hosp.*

R. Powdered rhubarb, one drachm.
 Magnesia, one drachm and a half.

- Powdered ginger, one scruple.
 Peppermint water, one pint.
 Mix. Dose, half a fl. ounce. *Gregory.*
 R. Powdered rhubarb, two scruples.
 Tartrate of potassa, one ounce.
 Peppermint water, six fl. ounces.
 Tincture of senna,
 Syrup of ginger, each, half a fl. ounce.
 Mix. Dose, one fl. ounce. *Brande.*

RHŒAS.

RED POPPY.

The *Papaver rhœas*, or Red Poppy, is a native of Europe, where it is found in great abundance in cultivated grounds, and has become naturalized in many places in this country.

Sex. Syst. Polyand. monog. *Nat. Syst.* *Papaveraceæ.*

Linn. Sp. Pl. 726. *Woodville, Med. Bot.* 387.

The parts used are the petals; these have a mucilaginous, somewhat bitter taste, and a narcotic smell, which latter is lost on drying. Their action on the system is slight; but they are used in Europe in the preparation of a syrup, prescribed as an anodyne in the catarrhal affections of children.

SYRUP OF RED POPPY.

- R. Red poppy petals, one pound.
 Water, one pint.
 Heat the water on a water-bath, gradually add the petals, remove from the fire, macerate for twelve hours, express, and strain; then add
 Sugar, three pounds,
 and make syrup. When cold, add
 Rectified spirit,
 two and a half fl. ounces.
Lond. Ph.

INFUSION OF RED POPPY.

- R. Red poppy petals, one drachm.
 Boiling water, one pint.
 Infuse and strain. To be taken freely, in catarrhal affections. *Coltereau.*

COMPOUND INFUSION OF RED POPPY.

- R. Red poppy petals, two ounces.
 Diluted sulphuric acid, fifteen drops.
 Sugar, two ounces.
 Decoction of barley, one pint.
 Infuse, and strain. Used as above. *St. Marie.*

ROSA CANINA.

DOG ROSE.

This shrub is a native of Europe, growing in hedges and thickets.

Sex. Syst. Icosand. polyg. *Nat. Syst.* *Rosaceæ.*

Linn. Sp. Pl. 704. *Woodville, Med. Bot.* 493.

The fruit, which is the officinal portion, is smooth, oval, red, and of a pleasant, acidulous taste. It is principally used in confection.

CONFECTION OF DOG ROSE.

- R. Pulp of fruit of dog rose, one pound.
 Sugar, twenty ounces.
 Gradually add the sugar to the rose-pulp, and rub together till well mixed. *Lond. Ph.*

Has been advised in diarrhœa and dysentery, as an astringent; but is principally used as a pill basis.

ROSA CENTIFOLIA.

HUNDRED-LEAVED ROSE.

This species, of which there are innumerable varieties, is cultivated in every garden, but its native country is unknown, though probably it is from Persia.

The petals are the only part used; these are fragrant, and have an acidulous, somewhat bitterish taste. They are slightly laxative, but are principally used for making rose water

ROSE WATER

- R. Fresh hundred-leaved
 rose petals, eight pounds
 Water, two gallons.
 Mix, and distil one gallon. *U. S. Ph.*
 R. Oil of roses, twenty minims.
 Distilled water, (Imp.) half a gallon.
 Agitate together, and filter. *Dub. Ph.*

OINTMENT OF ROSE WATER.

- R. Rose water, one fl. ounce
 Oil of almonds, two fl. ounces.
 Spermaceti, half an ounce.
 White wax, one drachm.
 Melt the last three ingredients together, on a water-bath, and stir in the rose water till cold. *U. S. Ph.*
 R. Oil of sweet almonds, two fl. ounces.
 Spermaceti, six drachms.
 Glycerin, four fl. drachms.
 Oil of roses,
 " bergamot, each, two drops.

Melt the spermaceti with a gentle heat, stir in the oil of almonds gradually, remove the mixture from the fire, stir constantly, adding the glycerin, and finally incorporate the volatile oils. *Jos. Laidley*

Much used, under the name of *Cold cream*, as a cooling application to irritated surfaces.

ROSE LOZENGES.

R. Powdered sugar,
 Rose water, each, six ounces.
 Evaporate gently, to a thick syrup, and add
 Coarsely-powdered sugar, one pound.
 Oil of roses, one drachm and a half.
 When dissolved, pour in drops, on a cold, oiled,
 marble slab, and dry. *Cottureau.*

COLLUTORY OF ROSE WATER.

R. Rose water, three fl. ounces.
 Cream,
 Whites of eggs,
 Syrup of violets, each, one fl. ounce.
 Mix. *St. Marie.*

OIL OF ROSES.

R. Fresh rose leaves, at will.
 Water, sufficient
 to moisten the leaves; distil, and collect the oil
 that floats on the product. *Guibourt.*

ESSENCE OF ROSES.

R. Pounded rose leaves,
 Alcohol (.874), each, ten parts.
 Infuse for twenty-four hours, by a gentle heat,
 and distil. *Taddei.*

ROSE OIL.

R. Pounded rose leaves, one part.
 Olive oil, four parts.
 Digest with gentle heat for four days, express,
 and strain; repeat process with fresh leaves, a
 second and third time; separate the water, and
 filter. *Soubeyran.*
 As an application to chilblains and irritated
 surfaces.

ROSA GALICA.

RED ROSE.

This species is a native of the south of Europe, and is generally cultivated in our gardens. The varieties are very numerous.

Linn. Sp. Pl. 704. Griffith, Med. Bot. 273.

The parts used in medicine are the petals of the unexpanded flowers, deprived of their white claws, and dried. In this state they are of a purplish red color, of an agreeable odor, and bitterish, astringent taste. They have mild astringent properties, but are more used as a vehicle, than for their own powers.

CONFECTION OF ROSES.

R. Red roses, in powder, four ounces.
 Powdered sugar, thirty ounces.

Clarified honey, six ounces.
 Rose water, eight fl. ounces.

Rub the roses with the rose water, heated to 150°; add gradually the sugar and honey, and beat together. *U. S. Ph.*

Principally used as a pill basis.

ACID INFUSION OF ROSES.

R. Red roses, half an ounce.
 Boiling water, forty-eight fl. ounces.
 Diluted sulphuric acid, three fl. drachms.

Digest in a glass vessel for half an hour; express, strain, and add

Sugar, one ounce and a half.

As a cooling drink, mixed with water.

Dub. Ph., 1826.

R. Red roses, half an ounce.
 Boiling water, two pints and a half.
 Diluted sulphuric acid, three fl. drachms.

Sugar, one ounce and a half.

Pour the water on the roses, in a glass vessel, add the acid, macerate for half an hour, strain, and add the sugar. *U. S. Ph.*

HONEY OF ROSES.

R. Red roses, two ounces.
 Clarified honey, two pints.
 Boiling water, one pint and a half.

Macerate the roses in the water for two hours, and strain; add the honey, and evaporate to proper consistence. *U. S. Ph., 1840.*

R. Coarsely-powdered red roses, two ounces.
 Clarified honey, twenty fl. ounces.
 Boiling water, twelve fl. ounces.

Macerate the roses for four hours in eight fl. ounces of the water, and express; macerate the residue in four fl. ounces of boiling water for half an hour, and again express. Mix the last infusion with four fl. ounces of the first and with the honey, and evaporate to one pint. Add the reserved infusion and strain. *U. S. Ph., 1850.*

As an addition to gargles.

ELECTUARY OF ROSES.

R. Confection of roses, sixteen parts.
 Syrup of tolu, four parts.
 " poppies, one part.

Mix. Used as an astringent in bowel affections, in doses of a teaspoonful. *Foy.*

TINCTURE OF ROSES.

R. Red roses, five ounces.
 Alcohol, two fl. ounces.
 Rose water, eight fl. ounces.

Digest for four days, express, and filter; digest residuum with half a pint of alcohol for three days; express, filter, and mix the liquors.

Squire.

Used principally as a perfume.

WINE OF ROSES.

R. Red roses, one part.
Red wine, sixteen parts.
Infuse for half an hour, and strain. As an application to indolent ulcers. *Beral.*

COMPOUND VINEGAR OF ROSES.

R. Red roses, one ounce and a half.
Sumach leaves, one ounce.
Wine of quince, four fl. ounces.
Vinegar of red wine, ten fl. ounces.

Macerate for ten days, express, and filter. As an application to contusions, and as a styptic. *Van Mons.*

SYRUP OF RED ROSES.

R. Red roses, two ounces.
Boiling water, one pint.
Sugar, twenty ounces.

Macerate the rose leaves in the water for twelve hours, strain, add sugar, and form a syrup. *Ed. Ph.*

A very mild astringent, of a fine red color.

ROSMARINUS.

ROSEMARY.

This is the *Rosmarinus officinalis*, a small evergreen shrub, a native of the south of Europe, and generally cultivated in gardens elsewhere.

Sex. Syst. Diand. monog. *Nat. Syst.* Lamiaceæ.

Linn. Sp. Pl. 23. *Griffith, Med. Bot.* 507.

The officinal portions are the flowering extremities. These have a fragrant odor, and a bitter, pungent taste. Rosemary is stimulant and carminative, but is principally employed as an aromatic addition to lotions and liniments, and as an ingredient in articles of perfumery.

AROMATIC BATH.

R. Rosemary, }
Thyme, } each, half a pound.
Sage, }
Origanum, }
Mint, }
Boiling water, six pints.

Macerate for twelve hours, strain, and add
Essence of soap, four ounces.
Muriate of ammonia, two ounces.

Pour the whole into water sufficient for a bath. *Foy.*

In chronic rheumatism, cutaneous affections, dyspepsia, &c.

AROMATIC FOMENTATION.

R. Rosemary, half an ounce.
Red wine,
Water, each, three fl. ounces.
Infuse, and express. As a fomentation in contusions. *Augustin.*

AROMATIC VINEGAR.

R. Rosemary, one ounce.
Sage, each, one ounce and a half.
Lavender, one ounce and a half.
Cloves, half a drachm.
Vinegar, two pints.
Infuse for eight days, and strain. As a lotion in contusions, sprains, &c. *Spielmann.*

ESSENCE OF ROSEMARY.

R. Oil of rosemary, one fl. ounce.
Rectified spirit, nine fl. ounces.
Mix with agitation. Dose, twenty to forty drops. *Dub. Ph.*

TINCTURE OF ROSEMARY.

R. Rosemary, one part.
Spirit of rosemary, four parts.
Macerate, express, and filter. *Bruns. Ph.*

OIL OF ROSEMARY.

R. Rosemary, at will.
Water, sufficient
to cover. Distil, and collect the oil that floats on the product. Principally used in stimulating liniments, but sometimes given as carminative in doses of two or three drops. *U. S. Ph.*

ROSEMARY OIL.

R. Rosemary, one part.
Olive oil, three parts.
Heat till the water has evaporated, express, and strain. As a stimulating application. *Span. Ph.*

SPIRIT OF ROSEMARY.

R. Oil of rosemary, four drachms.
Alcohol, one gallon.
Dissolve the oil in the alcohol. *U. S. Ph.*
As an ingredient in lotions, and liniments, &c.

HUNGARY WATER.

R. Rosemary, three parts
Lavender, one part
Diluted alcohol, six parts.
Water, twelve parts.
Mix, and distil three-fourths. Used as a perfume. *Van Mons*

R. Spirit of lavender, seven fl. ounces.
 Spirit of rosemary, four fl. ounces.
 Mix. Used as a substitute for the above.

Pereira.

R. Fresh rosemary, in blossom, four pounds.
 Fresh sage, in blossom, six ounces.
 Ginger, two ounces.
 Cut into pieces, and add
 Alcohol, twelve pounds.
 Water, two pints.
 Distil eleven pints by a gentle heat.
Wirt. Pharm.

RUBIA. MADDER.

The *Rubia tinctorum*, or Madder, is a perennial plant, a native of the south of Europe, and extensively cultivated in many countries for the sake of its roots, which are used in dyeing reds.

Sex. Syst. Tetrand. monog. *Nat. Syst.* Rubiaceæ.

Linn. Sp. Pl. 153. *Griffith, Med. Bot.* 382.

The root is the officinal portion, and as found in commerce, is in long, cylindrical pieces, about as thick as a quill, of a deep red or brown color. It has a peculiar and unpleasant odor, and a bitterish, astringent taste. It was much used formerly as a diuretic, but is at present seldom prescribed, except as an emmenagogue, for which purpose it is highly spoken of by many practitioners.

COMPOUND POWDER OF MADDER.

R. Powdered madder,
 " sugar, each, two drachms.
 " sulphate of potassa,
 half an ounce.

Mix well. Dose, ten grains to a scruple, in rachitis. *Radius.*

DECOCTION OF MADDER.

R. Powdered madder, one ounce.
 Boiling water, one pint.

Simmer for fifteen minutes, and add

Bruised cloves, one drachm.
 Strain when cool. A wineglassful every three hours, a short time previous to the expected time of the menstrual discharge. *Dewees.*

R. Powdered madder, one ounce.
 " mace, two drachms.
 Water, one pint and a half.

Boil down to one pint, strain, and add

Aromatic tincture, two fl. drachms.
 Syrup of citric acid, two fl. ounces.

Dose, three fl. ounces three or four times a-day, in amenorrhœa, chlorosis, &c. *Swed. Pharm.*

R. Powdered madder, half an ounce.
 " hops, one drachm.
 English walnut leaves, three.
 Water, two pints.

Boil down to a pint and a half, strain, and add
 Tincture of tartrate of iron,
 one fl. drachm.

Advised in scrofula, in doses of two ounces, night and morning. *St. Marie.*

RUMEX. DOCK.

Various species of dock have been used at different times, but the *R. Britannica*, or water dock, and the *R. obtusifolius*, or blunt-leaved dock, are alone officinal in this country.

Sex. Syst. Hexand. trigyn. *Nat. Syst.* Polygonaceæ.

Willd. Sp. Pl. ii. 250. *Griffith, Med. Bot.* 545.

All of the species of dock are cooling, aperient, and slightly diuretic, but they are seldom employed.

DECOCTION OF DOCK ROOT.

R. Bruised water dock-root, one ounce.
 Boiling water, one pint.

Boil for fifteen minutes and strain. This has proved efficacious in ichthyosis. Dose, one fl. ounce. *A. T. Thomson.*

RUBUS TRIVIALIS. DEWBERRY.

RUBUS VILLOSUS. BLACKBERRY.

These two species of shrubby plants are very common in the United States, and are well known for their pleasant fruit.

Sex. Syst. Icosand. polyg. *Nat. Syst.* Rosaceæ.

Torrey and Gray, i. 454, 455. *Griffith, Med. Bot.* 275, 276.

The juice of the fruit is employed in the form of a syrup. The officinal portions are the roots. These are identical in their effects; they have a very faint odor, and a bitter, astringent taste. The smaller are to be preferred. They are efficient, but mild astringents, and have been found very useful in a relaxed condition of the bowels.

DECOCTION OF BLACKBERRY ROOT.

R. Small roots of blackberry, one ounce.
 Water, one pint and a half.

Boil down to one pint, and strain. Dose, one or two fl. ounces, three or four times a-day, in diarrhœa, &c.

SYRUP OF BLACKBERRIES.

R. Juice of blackberries, twenty ounces.
Sugar, three pounds.

Boil, and strain. Said to be very useful in dysentery.

R. Blackberries, thirty quarts.

Mash, strain through unbleached muslin, and express the remainder. Then heat in a proper vessel

Sugar, (av.) sixty-four pounds.

Water, two and one-quarter gallons.

Make a syrup, and stir in the blackberry juice, continuing the heat until the syrup has boiled two or three minutes. The syrup should mark 31° Baum. while boiling. Remove from the fire, skim and strain. As this syrup is rather insipid, its flavor may be improved by adding aromatics. *Ambrose Smith.*

RUTA.

RUE.

The only species recognized in medicine is the *Ruta graveolens*, a small shrub with glaucous leaves and yellow flowers, a native of the south of Europe, but generally cultivated in gardens.

Sex. Syst. Decand. monog. *Nat. Syst.* Rutaceæ.

Linn. Sp. Pl. 523. *Griffith, Med. Bot.* 189.

The leaves are the part usually employed; they have a nauseous odor, and a bitter, acrid, disagreeable taste. They are stimulant and antispasmodic, and have been much used as an emmenagogue and vermifuge. The dose of the powder is from ten to twenty grains, two or three times a-day.

CONFECTION OF RUE.

R. Powdered rue,
Caraway,
Laurel berries, each,
one ounce and a half.

Sagapenum, half an ounce.

Black pepper, two drachms.

Clarified honey, sixteen ounces.

Mix. *Lond. Ph.*

It is used as an antispasmodic, in the form of an enema made with a scruple to a drachm, mixed with half a pint of warm, mucilaginous fluid.

EXTRACT OF RUE.

R. Dried rue, two pounds.
Alcohol (.927), seven pints.

Exhaust the rue by the process of displacement, distil off the alcohol, and evaporate the residue. *Wirt. Ph.*

MIXTURE OF RUE AND SQUILL.

R. Juice of rue leaves, one ounce.
Oxymel of squill, half an ounce.

Mix. A teaspoonful occasionally, in hysteric affections. *Pierquin.*

OIL OF RUE.

R. Fresh leaves, at will.
Water, sufficient.

Mix, distil, and separate the oil. Dose, one or two drops. *Lewis's Disp.*

R. Dried rue, sixty parts.
Olive oil, five hundred parts.

Digest for two hours on a sand-bath, agitating from time to time, express, and filter.

Par. Cod.

SYRUP OF RUE.

R. Coarsely-powdered sugar, fifteen ounces.

Tincture of rue, two fl. ounces.

Distilled water, seven fl. ounces.

Dissolve, and strain. *Beral.*

TINCTURE OF RUE.

R. Expressed juice of rue,
Alcohol (.847), each, four fl. ounces.

Mix, and filter at the end of twenty-four hours. *Beral.*

MIXTURE OF RUE.

R. Rue,
Savine,

Wormwood, each, three drachms.

Boiling water, one pint.

Infuse, strain, and add

Castor oil, half an ounce.

As an injection, in cases of ascarides. *Radins*

S.

SABADILLA.

CEVADILLA.

This article is the seed of one or more plants belonging to the natural order of Melanthaceæ; the U. S. Ph. attributes it to *Veratrum sabadilla*, and the Lond. Ph. to *Helonias (Asagraea) officinale*.

The Cevadilla of the shops consists of follicles, seeds, &c. The first are ovate, oblong, acuminate, of a pale, yellowish-brown color, and a thin, papery consistence. The seeds are small, curved, acuminate, shining, wrinkled, and of a blackish-brown color. They have very little odor, but their taste is bitter and acrid. These seeds are a drastic emeto-cathartic, and have been used as an anthelmintic, and in nervous diseases, &c.; but are principally used in the preparation of veratria. The dose is from five to ten grains, in powder; but it should always be given with much caution.

COMPOUND POWDER OF CEVADILLA.

R. Powdered cevadilla, two grains.
Sulphate of iron, one grain.
Powdered semen contra,
Magnesia, each, ten grains.
Mix. To be given four times a-day, in cases of tenia. *Radiis.*

TINCTURE OF CEVADILLA.

R. Seeds of cevadilla, bruised, at will.
Alcohol, sufficient
to cover them; digest for ten days, and filter.
As a lotion in rheumatism. *Turnbull.*

EXTRACT OF CEVADILLA.

R. Tincture of cevadilla, at will.
Evaporate to proper consistence. Dose, one-sixth of a grain, as a substitute for veratria in tic douloureux, &c. *Turnbull.*

OINTMENT OF CEVADILLA.

R. Powdered cevadilla, four ounces.
" mustard,
" pellitory, each, two ounces.
Lard, thirty-two ounces.
Mix well. To destroy lice. *Par. Cod.*

OINTMENT OF EXTRACT OF CEVADILLA.

R. Extract of cevadilla, fifteen to twenty grains.
Lard, one ounce.
Mix well. As a substitute for veratria ointment, but not as efficient. *Turnbull.*

CAPUCHIN POWDER.

R. Powdered cevadilla,
" stavesacre,
" parsley seed,
" tobacco, equal parts.
Mix. To destroy vermin in the hair.

Niemann.

SABBATIA.

AMERICAN CENTAURY.

The *Sabbatia angularis* is a small annual, or biennial plant, growing in abundance in various parts of the United States, having numerous rose-colored flowers, which appear in July and August.

Sex. Syst. Pentand. monog. *Nat. Syst.* Gentianaceæ.

Pursh, Flor. Am. i. 137. Griffith, Med. Bot. 458.

The herbaceous portion of the plant is official. It is a pure bitter, with no astringency or aroma.

INFUSION OF AMERICAN CENTAURY.

R. American centaury, one ounce.
Boiling water, one pint.
Infuse; and strain, when cool. Dose, a wine-glassful, every two hours, as a tonic. *Chapman*

TINCTURE OF AMERICAN CENTAURY.

R. American centaury, one part.
Diluted alcohol, six parts.
Digest for four days, express, and filter.
Dose, a fl. drachm to two fl. drachms.

EXTRACT OF AMERICAN CENTAURY.

R. American centaury, one part.
Diluted alcohol, eight parts.
Subject the herb to two or three macerations in the alcohol; heat to boiling; unite the tinctures, distil off the alcohol, and evaporate the residue.
Dose, five to ten grains.

SABINA.

SAVINE.

Savine is the tops of *Juniperus sabina*, a small tree or shrub, a native of the south of Europe, somewhat resembling the red cedar in its foliage.

Sex Syst. Diœcia monadelph. *Nat. Syst.* Pinaceæ.

Linn. Sp. Pl. 1472. *Griffith, Med. Bot.* 608.

The tips of the branches are officinal. These, when dried, have a bitter, acrid taste, and an unpleasant odor. Savine is an active stimulant, acting powerfully on the skin and uterus. It also acts as an external irritant. The dose of the powder is from five to fifteen grains, three or four times a-day.

POWDER OF SAVINE AND GINGER.

R. Powdered savine,
" ginger, each, one drachm.
Sulphate of potassa, two drachms.
Mix, and divide into six powders. One, to be taken twice a-day, in amenorrhœa. *Ellis.*

POWDER OF SAVINE AND SPANISH FLIES.

R. Powdered savine, one drachm.
" Spanish flies, two grains.
Mix, and divide into four powders. One, at bedtime, in amenorrhœa. *Ellis.*

POWDER OF SAVINE AND VERDIGRIS.

R. Powdered savine,
" verdigris, equal parts.
Mix. As an escharotic to venereal warts. *J. Hunter.*

POWDER OF SAVINE AND PINKROOT.

R. Powdered pinkroot,
" senna, each, two scruples.
" savine, twelve grains.
Mix, and divide into six powders. Anthelmintic. One, every morning, till three are taken, then a dose of castor oil. If required, repeat. *Ellis.*

PILLS OF SAVINE.

R. Extract of savine, half an ounce.
Powdered savine, two drachms.
Oil of savine, sufficient.
Mix, and form pills of two grains each. Four, three times a-day, as an emmenagogue. *Niemann.*

EXTRACT OF SAVINE.

R. Powdered savine, two pounds.
Alcohol, seven pints.
Exhaust the savine by the process of displacement; distil off the alcohol, and evaporate the residue, on a water-bath. *Par. Cod.*

OIL OF SAVINE.

R. Savine, at will
Water, sufficient to cover.
Distil, and collect the oil that floats on the product. *Guibourt.*

A powerful stimulant emmenagogue, an rubefacient. Dose, two to five drops.

INFUSION OF SAVINE.

R. Savine, one drachm.
Boiling water, eight fl. ounces.
Infuse for half an hour, and strain. Dose, half a fl. ounce. *Pereira.*
R. Savine, one drachm.
Camphor, six grains.
Boiling water, five fl. ounces.
Infuse, and strain. *Horn.*

TINCTURE OF SAVINE.

R. Savine, one part.
Alcohol, four parts.
Macerate for ten days, and filter. Dose, ten to fifteen drops. *Soubeiran.*

COMPOUND TINCTURE OF SAVINE.

R. Extract of savine, one ounce.
Tincture of castor, sixteen fl. ounces.
Tincture of myrrh, eight fl. ounces.
Digest till dissolved. As an emmenagogue. *Lond. Ph., 1788.*

R. Compound tincture of
savine, one fl. ounce.
Tincture of black hellebore,
half a fl. ounce.
" castor, two fl. drachms.
Mix. Forty drops, three times a-day. *Ellis.*

SAVINE CERATE.

R. Powdered savine, two ounces
Resin cerate, one pound.
Melt the cerate, and mix in the savine. *U. S. Ph.*
As a dressing to keep up the discharge of blisters, &c.

SAVINE OINTMENT.

R. Powdered savine,
Lard, equal parts.
Mix. As an application to venereal warts. *Radius.*
R. Fresh savine, bruised, half a pound
White wax, three ounces.
Lard, one pound.
Melt the lard and wax together, mix in the savine, and express through linen. *Lond. Ph.*

SACCHARUM.

SUGAR.

Sugar is procured from the *Saccharum officinarum* and other species, and is used in several forms, as molasses, brown, and refined sugar, the latter of which only, is recognized in the U. S. Pharm.

SYRUP.

R. Sugar, two pounds and a half.
Water, one pint.

Dissolve the sugar in the water, by the aid of heat, remove any scum that may form, and strain while hot. U. S. Ph.

CAMEL OR BURNT SUGAR.

R. Sugar, at will.

Place in a shallow vessel, and heat over a gentle fire, till it assumes a dark-brown color. Used as a coloring for spirits, &c. Gray.

VERMIFUGE SUGAR.

R. Ethiops mineral, two parts.
Quicksilver, three parts.

Rub together until the globules are extinguished; add

Sugar, seven parts,
and mix thoroughly. Soubeiran.

SACCHARATED POWDERS.

Under this name, M. Beral has proposed the mixture of sugar with various active medicines, all the ingredients being reduced to powder. The peculiarity of the preparation is that sugar enters it in large proportion, while the true medical agent is in small quantity.

SACCHARATED POWDER OF DIGITALIS.

R. Fresh leaves of digitalis,
deprived of midribs and
nerves, one part.

Place in bibulous paper, and expose to the air in the shade for twelve hours. Then mix carefully with

White sugar, three parts.

Dry the mixture with a gentle heat, pulverize, and preserve in opaque bottles.

In the same manner may be prepared saccharated powders of fresh *aconite* leaves, *belladonna*, *hemlock*, *henbane*, *savine*, *stramonium*, and *rue*.

Dorvault.

SACCHARATED POWDER OF JALAP.

R. Tincture of jalap, sixty parts.
Sugar, in small lumps,
five hundred parts.

Pour the tincture on the sugar, and permit the alcohol to evaporate spontaneously, or on a sand-

bath. The process of evaporation may be accelerated by reducing the sugar to a coarse powder, twenty-four hours after the addition of the tincture. The active principle is retained in the sugar.

In the same manner may be prepared saccharated powders of—

Belladonna,	Mace,
Castor,	Myrrh,
Cinnamon,	Nutmeg,
Cinchona,	Rhubarb,
Cloves,	Saffron,
Henbane,	Squill,
Ipecacuanha,	Tolu,
&c. &c.	Vanilla,
	&c. &c.

This mode of obtaining the active principles of medicines in a soluble form, may be resorted to, advantageously, in making lozenges. Dorvault.

OLEO-SACCHARATED POWDERS.

R. Any volatile oil, one part.
Sugar, seventy-two parts.

Rub together. For convenience of preparation, one drop of the oil to one drachm of sugar, is sufficiently accurate for all practical purposes. By this admixture, the volatile oil becomes soluble in water. Dorvault.

SAGAPENUM.

SAGAPENUM.

This is a gum resin, derived most probably from some plant belonging to the *Apiaceæ*. It is in irregular masses, composed of cohering fragments, of a yellowish-brown, olive, or reddish-yellow color; of a waxy consistence, of a somewhat alliaceous odor, and a pungent, bitterish, unpleasant taste. It is a mild stimulant, somewhat resembling *assafetida* in its properties, but inferior to it. It has been used as an emmenagogue and antispasmodic, but it is now seldom prescribed internally, but is still employed as an external application as a discutient.

PREPARED SAGAPENUM.

R. Sagapenum, one pound.
Water, sufficient to cover.

Heat the sagapenum with the water until they are mixed. Strain through a hair sieve, and evaporate on a water-bath, stirring constantly.

Lond. Ph.

COMPOUND SAGAPENUM PILLS.

R. Sagapenum, one ounce.
Aloes, half a drachm.
Syrup of ginger, sufficient.

Beat together. Dose, ten grains.

Lond. Ph., 1836.

As a stimulant, antispasmodic laxative, in flatulent colic, &c.

SAGO.

SAGO.

This is the prepared fecula of the pith of various species of Palmaceæ and Cycadaceæ. It is prepared in two forms, meal and pearl sago, the latter of which is most generally employed. This is in small, hard, whitish or brownish grains, inodorous, and of a mucilaginous taste. It is insoluble in cold water, but by long boiling forms a glutinous solution. It is nutritive, easily digestible, and forms an excellent article of diet for the sick and convalescent.

SAGO MUCILAGE.

R. Sago, one ounce.
Water, one pint.

Macerate by a gentle heat, for two hours, then boil for fifteen minutes, stirring continually, till the grains are perfectly dissolved; add sugar or flavoring, according to circumstances.

A. T. Thomson.

SAGO POSSET.

R. Sago, two ounces.
Water, two pints.

Macerate for two hours, by a gentle heat, then boil, till a mucilage is formed; then rub half an ounce of sugar on the rind of a lemon, add this and one fl. drachm of tincture of ginger to half a pint of sherry wine, pour the whole into the sago mucilage, and boil for five minutes. Useful in convalescence to restore strength. A wineglassful to be taken every four or five hours.

A. T. Thomson.

SAGO MILK.

R. Sago, one ounce.
Water, one pint.

Macerate for half an hour, then add

Milk, one pint and a half.

Boil slowly, till the sago is perfectly dissolved.

A. T. Thomson.

SALICINUM.

SALICIN.

R. Decoction of willow bark, at will.
Treat with slaked lime, filter, and evaporate to consistence of syrup; add alcohol, again filter, evaporate, and let crystallize; wash the crystals with cold water.

Cottureau.

Dose, four to six grains every three hours, in apyrexia of intermittents.

PILLS OF SALICIN.

R. Salicin,
Extract of gentian,
each, twelve grains.
Liquorice powder, sufficient.
Mix, and make six pills.

Foy.

R. Salicin, twenty-four grains.
Mucilage of gum Arabic, sufficient
to make eight pills. Dose, one pill, every three hours, in the apyrexia of intermittent fevers.

Ellis.

COMPOUND SALICIN PILLS.

R. Salicin, one scruple.
Compound rhubarb pill, two scruples.
Mix, and make twelve pills.

Beasley.

COMPOUND POWDER OF SALICIN.

R. Salicin, two scruples.
Aromatic powder, one drachm.
Mix, and divide into twelve powders.

Neligan.

R. Salicin, fifteen grains.
Tartar emetic, one grain.
Powdered sugar, fifty grains.

Mix, and divide into ten powders. One, thrice a-day.

Krombolz.

SALIX.

WILLOW.

Many species of willow have been used in medicine, and are recognized in the different Pharmacopœias; that admitted by the U. S. Ph. is the *Salix alba*, a small tree, a native of Europe, but extensively cultivated in the United States.

Sex. Syst. Diœcia diand. Nat. Syst. Salicaceæ.

Linn. Sp. Pl. 1449. Lindley, Flor. Med. 318.

The officinal portion is the bark; this is in thin, flexible, fibrous pieces, of a brown color. It has a peculiar, aromatic odor, and an astringent, bitter taste. It is tonic and astringent, and has been used as a substitute for Peruvian bark.

EXTRACT OF WILLOW BARK.

R. Powdered willow bark, two parts.
Distilled water, one part.

Macerate twelve hours, transfer to a percolator, exhaust, raise the liquid to the boiling point, strain, and evaporate to the consistence of an extract.

Dorvault.

COMPOUND POWDER OF WILLOW BARK.

R. Powdered willow bark,
" horse-chestnut bark,
" gentian,
" calamus,
" avens root, equal parts.

In intermittent fevers, in drachm doses.

Hufeland

WILLOW BARK DENTIFRICE.

R. Powdered willow bark,
 " charcoal, each,
 half an ounce.
 Extract of myrrh, two drachms.
 Balsam of Peru, half a scruple.
 Oil of cinnamon, three drops.
 Triturate well together. *Phœbus.*

DECOCTION OF WILLOW BARK.

R. Contused willow
 bark, one ounce and a half.
 Water, fifteen fl. ounces.
 Boil down to one-half. *Niemann.*

OINTMENT OF WILLOW LEAVES.

R. Expressed juice of willow
 leaves, two fl. ounces.
 Lard, sufficient.
 Mix, and heat till moisture is driven off. Useful
 as a dressing to foul ulcers. *Hufeland.*

SALVIA.

SAGE.

Though many species of *Salvia* possess analogous remedial properties, one only is recognized by the U. S. Ph.,—the *Salvia officinalis*, a perennial plant, native of the south of Europe, but generally cultivated in our gardens, flowering in June, at which time it should be collected.

Sex. Syst. Diand. monog. *Nat. Syst.* Lamiaceæ.

Linn. Sp. Pl. 34. Griffith, Med. Bot. 505.

The leaves are the officinal portion. These have a strong, fragrant odor, and a warm, bitterish, aromatic, somewhat astringent taste. Sage is stimulant, with some astringent and tonic powers. It is principally used in the composition of gargles, and is also said to abate the excessive sweats in hectic fever. The dose, in powder, is from twenty to thirty grains.

INFUSION OF SAGE.

R. Sage, half an ounce.
 Boiling water, two pints.
 Infuse for half an hour, and strain. Sugar and lemon juice may be added, according to circumstances. As a drink in fevers. *Ellis.*

COMPOUND INFUSION OF SAGE.

R. Sage,
 Boneset, each, half an ounce.
 Cascarilla, one drachm.
 Water, one pint and a half.
 Infuse till cold, and strain. Dose, a wineglassful every three or four hours. In hectic fever. *Ellis.*

GARGLE OF SAGE.

R. Infusion of sage, one pint.
 Diluted sulphuric acid, two drachms.
 Honey of roses, one ounce.

Mix. In relaxation of the uvula, &c. *Radius.*

R. Infusion of sage, two pints.
 Tincture of Peruvian bark,
 Syrup of mulberries, each, half a fl. ounce.
 Spirit of horseradish, one drachm.

Mix. More active than the last. *Cadet.*

VINEGAR OF SAGE.

R. Sage, one part.
 Vinegar, six parts.
 Mix, and distil four parts. *Beral.*
 As a gargle, mixed with water.

SAMBUCUS.

ELDER FLOWERS.

The species recognized by the U. S. Ph., is the *Sambucus Canadensis*, which appears to be perfectly identical in properties with the European kind, the *S. nigra*.

Sex. Syst. Pentand. trigyn. *Nat. Syst.* Caprifoliaceæ.

Willd. Sp. Pl. 1494. Griffith, Med. Bot. 353.

The officinal portion is the flowers, but the berries, bark, and leaves are also used. The flowers have a faint but peculiar odor, when dried, and a bitterish taste. They are slightly stimulant and diaphoretic. The berries are diaphoretic and laxative; the bark acts as a hydragogue purgative, as do also the leaves.

ELDER-FLOWER WATER.

R. Fresh elder flowers, ten pounds.
 Water, (Imp.) two gallons.
 Mix, and distil a gallon. *Lond. Ph.*

Principally used as a flavoring ingredient in mixtures and emulsions.

VINEGAR OF ELDER FLOWERS.

R. Elder flowers, one part.
 Vinegar of red wine, twelve parts.
 Macerate for eight days, express, and filter. As a gargle. *Soubeiran.*

GARGLE OF ELDER FLOWERS.

R. Elder flowers, one ounce.
 Boiling water, one pint.
 Infuse, strain, and add
 Nitrate of potassa, half an ounce.
 Tincture of burnet, three fl. drachms.
 Oxymel, two fl. ounces.
 Mix. *Saunders*

EXTRACT OF ELDER BERRIES.

R. Elder berries, one pound.
Bruise them in a mortar; express the juice, and evaporate to due consistence. *Dub. Ph.*, 1788.

Has been praised in rheumatic, gouty, and eruptive affections; in doses of one to four drachms.

MIXTURE OF EXTRACT OF ELDER BERRIES.

R. Extract of elder berries,
Pulp of prunes,
Syrup of red
poppies, each, two ounces.
Nitrate of potassa, one drachm.

Mix. Two or three spoonfuls a-day, in asthma. *St. Marie.*

DECOCTION OF ELDER BARK.

R. Elder bark, three ounces.
Water, two pints.
Boil to one-half. Half a pint, morning and evening, in dropsy. *Sydenham.*

OINTMENT OF ELDER LEAVES.

R. Fresh elder leaves, three pounds.
Lard, four pounds.
Suet, two pounds.
Boil the leaves in the lard till crisp; express; strain; add the suet, and melt. *Dub. Ph.*, 1826.

As a cooling application.

OINTMENT OF ELDER FLOWERS.

R. Elder flowers,
Lard, equal parts.
Melt, and continue heat, till all moisture is driven off, and express. *Beral.*
Used as the last.

SANGUINARIA.
BLOODROOT.

This is the rhizome of the *Sanguinaria Canadensis*, popularly known as bloodroot, or puccoon. It is a small, herbaceous perennial, flowering at the very commencement of the spring; at which time the leaf is small, but attains a large size during the summer.

Sex. Syst. Polyand. monog. *Nat. Syst.* Papaveraceae.

Linn. Sp. Pl. 723. *Griffith, Med. Bot.* 127.

The root is horizontal, abrupt, often contorted, of a reddish-brown color, externally, and of a bright orange-red within. When dried it is somewhat flattened, much wrinkled and twisted, of a reddish-brown color externally, and of a bright orange within, becoming brown by exposure. The powder is brownish-red. It has a faint narcotic odor, and a bitterish, acrid taste, which is very persistent. It is an acrid

emetic, with narcotic and stimulant properties. Dose, as an emetic, from ten to twenty grains; for other purposes, from one to five grains.

PILLS OF BLOODROOT.

R. Powdered bloodroot, one drachm.
Conserve of roses, sufficient.

Mix, and make thirty pills. One to two as an alterative, &c., five to ten as an emetic.

INFUSION OF BLOODROOT.

R. Bloodroot, half an ounce.
Boiling water, one pint.

Infuse for two hours. Dose, half to one fl. ounce. *Beasley.*

TINCTURE OF BLOODROOT.

R. Bloodroot, bruised, four ounces.
Diluted alcohol, two pints.

Macerate for fourteen days, express, and filter. *U. S. Ph.*

As emetic, three or four fl. drachms; as expectorant, alterative, &c., thirty to sixty drops.

R. Bruised bloodroot, two ounces.
Spirit of nitric ether, two pints.

Digest eight days, and filter.

This is a valuable expectorant and diaphoretic, in doses of half a fl. drachm to one fl. drachm. *Mettauer.*

COMPOUND TINCTURE OF BLOODROOT.

R. Bloodroot,
Lobelia,
Skunk-cabbage root,
Asarabacca,
Pleurisy root, each,
in coarse powder, one ounce.

Place them in a vessel, and cover with

Boiling water or vinegar, one pint; and cover tightly. When cold, add

Alcohol, three pints.
Macerate fourteen days, and filter. Used as an emetic. *Ecl. Med. Jour.*

SYRUP OF BLOODROOT.

R. Coarsely-powdered bloodroot, eight ounces.
Acetic acid, four ounces.
Water, five pints.
Sugar, (troy) two pounds.

Mix two fl. ounces of the acetic acid with a pint of water and macerate the root for three days. Transfer to a percolator, and displace with the remainder of the water, previously mixed with the other half of the acetic acid. Evaporate on a water-bath to eighteen fl. ounces, add the sugar, and form a syrup.

T. S. Wiegand

SANTALUM.

RED SAUNDERS.

This is the wood of *Pterocarpus santalinus*, a lofty tree, found in the East Indies. It is imported in logs or billets.

Sex. Syst. Diadelph. decand. *Nat. Syst.* Fabaceæ.

Linn. Suppl. 318. Griffith, Med. Bot. 245.

Red saunders is a mild astringent and tonic; but it is chiefly, if not exclusively, employed to impart color.

SAPO.

SOAP.

Soaps are combinations of animal or vegetable oils, with one or more of the alkalies; they are principally used for washing, but some of them are official.

SAPO DURUS.

HARD SOAP.

This is officinal under the name of *Sapo*, in the U. S. and Lond. Phs., and is a combination of olive oil and soda.

SPANISH, OR CASTILE SOAP.

This is the officinal soap of the U. S. Ph., and presents two varieties, the white and the marbled, the first of which only should be used.

SAPO MOLLIS.

SOFT SOAP.

This soap is a combination of olive or other vegetable oils and potassa, or animal oils with the same alkali.

SAPO VULGARIS.

COMMON SOAP

Is a hard soap, made from tallow and caustic soda.

Besides these officinal soaps, there are many others, which are used medicinally, or for cleansing purposes.

ALMOND SOAP.

R. Caustic solution
of soda, one thousand parts.
Oil of almonds,
two thousand one hundred parts.

Mix, and keep the mixture for some days at a temperature of 68° F., stirring from time to time, until it acquires the consistence of a soft paste; place in moulds until dry. It should not

be used for medicinal purposes, until it has been exposed to the action of the air for a month or two.
Par. Cod.

BEEF-MARROW SOAP.

R. Purified beef
marrow, five hundred parts.
Caustic solution of soda,
two hundred and fifty parts.
Common salt, one hundred parts.
Water, one thousand parts.

Put the marrow, with hot water, into a porcelain vessel, and heat, until it is melted; then add the solution of caustic soda, by degrees, constantly stirring, till saponification is complete; then add the salt, slightly stirring, collect the soap which rises, drain it, melt it by a gentle heat, pour into moulds, and allow it to solidify.

Par. Cod.

SOAP OF TURPENTINE. STARKEY'S SOAP.

R. Dry carbonate of potassa,
Oil of turpentine,
Venice turpentine, equal parts.

Triturate the potassa with the oil, and then with the turpentine, until the mass has attained a proper consistence. Preserve in an earthenware vessel.
Par. Cod.

Was supposed to be a corrector of the injurious effects of opium, hellebore, &c. It is now principally given in gonorrhœa and dropsy. Dose, eight to ten grains.

AROMATIC SOAP.

R. Concentrated solution of
soap, six ounces.
Oil of bergamot,
" lavender,
" rosemary, each, two drachms.

Mix. Employed in baths, as a tonic and antispasmodic.
Niemann.

CAMPHORATED SOAP.

R. White soap sixteen ounces.
Boiling water, eight ounces.
Olive oil, six ounces.
Powdered camphor, one drachm.

Dissolve the soap in the water, evaporate gently to the consistence of a soft paste, add the camphor incorporated in the oil, mix well, and pour into moulds.

Said to be useful in chaps and excoriations.

Wetzler.

COSMETIC SOAP POWDER.

R. White soap, twelve ounces.
Carbonate of potassa, two ounces.
Powdered orris root, three ounces.
" horse-chestnuts,
two pounds.

Oil of lavender,
 " bergamot, each, forty drops.
 " lemon, thirty drops.
 " cloves, ten drops.
 Sugar, half an ounce.

Mix, and form a powder. *Niemann.*

ARSENICAL SOAP.

R. Arsenious acid, two pounds.
 Carbonate of potassa, twelve ounces.
 Camphor, five ounces.
 White soap, two pounds.
 Powdered lime, eight ounces.

Reduce each to powder, and mix. Used as a preservative for specimens of natural history against the attacks of insects. Known as *Beconi's arsenical soap*. *Gannal.*

SOAP CERATE.

R. Solution of subacetate
 of lead, two pints.
 Soap, six ounces.
 White wax, ten ounces.
 Olive oil, one pint.

Boil the saturnine solution with the soap, over a slow fire, to the consistence of honey; transfer to a water-bath, and evaporate until all the moisture is dissipated; add the wax, previously melted with the oil, and mix. *U. S. Ph.*

Soap cerate is cooling and sedative.

CAMPHORATED SOAP LINIMENT.

OPODELDOC.

R. Common soap, sliced, three ounces.
 Camphor, one ounce.
 Oil of rosemary,
 " origanum, each, one fl. drachm.
 Alcohol, one pint.

Digest the soap in the alcohol, on a sand-bath, until dissolved; add the camphor and oils, dissolve, and pour into broad-mouthed bottles. *U. S. Ph.*

CAMPHORATED TINCTURE OF SOAP.

SOAP LINIMENT.

R. Soap, in shavings, four ounces.
 Camphor, two ounces.
 Oil of rosemary, half a fl. ounce.
 Water, four fl. ounces.
 Alcohol, two pints.

Mix the alcohol and water, and digest the soap in the mixture, on a water-bath, till dissolved; filter, and add the camphor and oil. *U. S. Ph.*

SAPONINE. (*For cleaning gloves.*)

R. Powdered
 soap, two hundred and fifty parts.
 Solution of chlorinated
 potassa, one hundred and sixty-five parts.

Solution of ammonia, ten parts.

Water,
 one hundred and fifty-five parts.

Mix, and form a paste: a small portion rubbed on a glove with a piece of flannel will cleanse it. *Duvignan.*

GREASE BALLS.

R. Fuller's earth,
 two pounds four ounces.
 Carbonate of soda,
 Soap, each, eight ounces.

Add,
 Yolks of eggs, eight.

Well beaten with
 Ox gall, eight ounces.
 Levigate thoroughly, form into cakes or balls, and dry. *Lenormand.*

BOLUS OF SOAP.

R. White soap, two scruples.
 Oil of caraway, two or three drops.
 Syrup, sufficient.

Mix, and make two boluses. They are purgative. *Swediaur.*

PILLS OF SOAP.

R. White soap, one hundred and twenty-five parts.

Marsh mallow root,
 powdered, sixteen parts.
 Nitrate of potassa, four parts.

Beat together till well incorporated, and divide into four-grain pills. *Par. Cod.*

PILLS OF SOAP AND OX GALL.

R. Powdered white soap, two drachms.
 Extract of ox gall, one drachm.

Mix, and incorporate
 Powdered guaiacum,
 Calomel, each, half a drachm.
 Powdered guaiacum wood, sufficient.

Mix, and make four-grain pills. In gout, one to two, morning and evening. *Vicq D'Azir.*

COMPOUND SOAP PILLS.

R. White soap, three drachms
 Ammoniac,
 Rhubarb, each, one drachm.

Aloes, ten grains.
 Assafetida,
 Saffron, each, thirty-six grains.
 Mix, and make three-grain pills. Purgative
 and alterative; four to six a-day. *Recamier.*

R. Powdered opium, half an ounce.
 Soap, two ounces.

Beat into a pilular mass, with water. Dose,
 three to five grains. *U. S. Ph.*

SOAP PLASTER.

R. Soap, sliced, four ounces.
 Lead plaster, three pounds.

Rub the soap into a semi-fluid state with water;
 then mix it with the plaster, previously melted,
 and boil to proper consistence. *U. S. Ph.*

ELECTUARY OF SOAP, &c.

R. White soap, one drachm and a half.
 Venice turpentine, one drachm.
 Seneka,
 Marsh mallow, each, two drachms.
 Rob of juniper, sufficient.
 Mix. Four teaspoonfuls a-day. *Vogt.*

ESSENCE OF SOAP.

R. White soap, twenty-four parts.
 Distilled water, thirty-two parts.
 Alcohol (.923), sixty-four parts.
 Carbonate of potassa, one part.
 Essence of lemon, sufficient.

Dissolve the soap in the water and alcohol; add
 the potassa and essence, and filter. Used for
 the toilette. *Soubeiran.*

CAMPHORATED ESSENCE OF SOAP.

R. White soap, three parts.
 Camphor, one part.
 Spirit of rosemary, sixteen parts.

Dissolve the camphor, and then the soap, in the
 spirit. As an embrocation in rheumatic pains,
 &c. *Guibourt.*

ETHEREAL SOLUTION OF SOAP.

R. Beef marrow soap, one part.
 Acetic ether, five parts.

Dissolve by aid of a water-bath. Used as an
 embrocation in rheumatic pains, &c. *Foy.*

CAMPHORATED ACETIC BALSAM OF SOAP

R. Common soap, one drachm.
 Camphor, each, one ounce.
 Acetic ether, one ounce.
 Oil of thyme, ten drops.

Dissolve the soap in the ether with the aid of
 heat; add the camphor, then the oil, and filter.
 Used as above. *Cottureau.*

CATAPLASM OF SOAP.

R. Common soap, four drachms.
 Roasted onion,
 Mustard, each, two ounces.
 Water, sufficient.

Heat together, and mix into a cataplasm. As
 a maturing application to boils, abscesses,
 &c. *Foy.*

SOAP SUPPOSITORY.

R. Soap, two ounces.
 Common salt, one ounce
 Honey, sufficient.

Mix, and form into conical suppositories, and
 oil them on the surface. *Spielmann.*

CLYSTER OF SOAP.

R. Barley water, six ounces.
 White soap, one drachm.
 Honey, six drachms.

Mix, and dissolve. *Brera.*

SOAP OF COD LIVER OIL.

R. Cod liver oil, two ounces.
 Caustic soda, two drachms.
 Water, five drachms.

Dissolve the soda in the water, and mix the
 solution with the oil. *Deschamps.*

IODURETTED SOAP OF COD LIVER OIL.

R. Soap of cod liver oil, one ounce.
 Iodide of potassium, one drachm.
 Water, one drachm.

Dissolve the iodide in the water, and add it to
 the soap. *Deschamps.*

SOAP WITH SULPHUR.

R. Soap, four ounces.
 Sulphur, each, half a drachm.
 Oil of bergamot, half a drachm.
 Water, sufficient.

Beat together. As an application in itch.
Frank.

ITCH OINTMENT.

R. Brown soap, one ounce.
 Common salt,
 Sulphur, each, half an ounce.
 Alcohol, one fl. drachm.
 Vinegar, two fl. drachms.
 Chloride of lime, half a drachm.

Rub well together. One-fourth to be used
 night and morning, as a friction. It is effectual,
 cheap, and inoffensive. *Emory.*

SARSAPARILLA.

SARSAPARILLA.

This is the officinal name for the roots of several species of *Smilax* found in Mexico and various parts of South America. These roots are known in commerce by the names of the places from whence they are shipped, as Para, Honduras, Jamaica, Vera Cruz, &c.

Sarsaparilla, as found in commerce, is in packages, composed of dried roots, several feet in length, about the thickness of a quill, more or less wrinkled, of an ash-grey to a dark-brown color externally, and white to brownish within. The odor is slight but peculiar, the taste is at first mucilaginous, but finally somewhat acrid. It is considered alterative and tonic, but its real action is not well understood. The dose, in powder, is half a drachm to a drachm, three or four times a-day.

POWDER OF SARSAPARILLA AND PERUVIAN BARK.

R. Powdered sarsaparilla, one ounce.
 " Peruvian bark, three drachms.
 Carbonate of soda, two drachms.
 Mix, and divide into sixteen powders. One, thrice a-day, as an alterative. *Cline.*

INFUSION OF SARSAPARILLA.

R. Sarsaparilla, bruised, one ounce.
 Boiling water, one pint.
 Digest for two hours, in a covered vessel, and strain. *U. S. Ph.*
 One to four fl. ounces, three times a-day.

ALKALINE INFUSION OF SARSAPARILLA.

R. Sarsaparilla, bruised, twelve ounces.
 Liquorice root, bruised, one ounce and a half.
 Solution of potassa, one fl. ounce and a half.
 Boiling water, five pints and a half.
 Macerate for twenty-four hours, and strain.
 Dose, from eight fl. ounces to one pint daily.
 As an alterative. *St. Geo. Hosp.*

DECOCTION OF SARSAPARILLA.

R. Sliced sarsaparilla, five ounces.
 Distilled water, four pints.
 Boil down to two pints, and strain. *Lond. Ph.*
 Dose, four to six fl. ounces a-day.

COMPOUND DECOCTION OF SARSAPARILLA.

R. Sarsaparilla, sliced, six ounces.
 Bark of sassafras, sliced,
 Guaiacum wood, rasped,

Liquorice root, bruised, one ounce.
 each,
 Mezereon, sliced, three drachms.
 Water, four pints.

Macerate for twelve hours; then boil for a quarter of an hour, and strain. *U. S. Ph.*

As an alterative and diaphoretic, in secondary syphilis, cutaneous affections, &c. Dose, four to six fl. ounces three times a-day.

FELTZ'S DECOCTION OF SARSAPARILLA.

R. Sarsaparilla, bruised, three ounces.
 Isinglass, half an ounce.
 Crude antimony (tied in a rag), three drachms.
 Water, five pints.
 Boil to two pints and a half, and strain. *Beasley.*

VINACHE'S DECOCTION OF SARSAPARILLA.

R. Sarsaparilla, bruised,
 China root, bruised,
 Guaiacum wood, rasped, each, one ounce and a half.
 Crude antimony (in a rag), two ounces.
 Water, six pints.
 Macerate for twelve hours, boil to three pints, and add
 Sassafras bark, sliced,
 Senna, each, half an ounce.
 Infuse for an hour, strain, let settle, and decant. *Foy.*

LISBON DIET DRINK.

R. Guaiacum wood, rasped, one ounce.
 Sarsaparilla, bruised, three ounces.
 Mezereon, sliced, half an ounce.
 Crude antimony (in a rag), two ounces.
 Water, twelve pints.
 Boil down to eight pints, and add
 Red saunders, rasped,
 White sandal, rasped, each, three ounces.
 Rose wood, rasped,
 Sassafras bark, sliced, each, one ounce.
 Liquorice root, sliced, half an ounce.
 Infuse for four hours, strain, and add syrup according to taste. *Foy.*
 Dose, a pint to two pints a-day.
 R. Sarsaparilla, bruised, four ounces.
 Dried walnut peel, four ounces.
 Guaiacum, rasped, one ounce and a half.

Crude antimony (in a rag), half an ounce.
 Water, four pints.
 Boil down to three pints. *Pearson.*

ZITTMAN'S DECOCTION.

R. Sarsaparilla, bruised, twelve ounces.
 Spring water, forty-eight pints.
 Macerate for twenty-four hours, and introduce,
 tied up in a rag,
 Styptic powder (four parts alum
 and one of kino),

one ounce and a half.
 Calomel, half an ounce.
 Cinnabar, one drachm.
 Boil down to sixteen pints, and add

Aniseed,
 Fennel seed, each, half an ounce.
 Senna, three ounces.
 Liquorice root, one ounce and a half.

Strain, and put aside as *Strong decoction*. Add
 to the dregs of the strong decoction,

Sarsaparilla, bruised, six ounces.
 Spring water, forty-eight pints.

Boil down to sixteen pints, and add

Lemon peel, }
 Cinnamon, } each, three drachms.
 Cardamom, }
 Liquorice root, six drachms.

Strain, and label, *Weak decoction*. To be given
 freely in the treatment of syphilitic and cuta-
 neous affections, in chronic rheumatism, &c.

Walther.

JAUPERAND'S DECOCTION.

R. Bruised sarsaparilla,
 " China root,
 Crude antimony, each, two ounces.
 Carbonate of potassa, one drachm.
 Senna,
 Sassafras, sliced, each, one ounce.
 Rhubarb,
 Peruvian bark, each, half an ounce.
 Water, two gallons.

Boil by a gentle fire for eight hours, till twelve
 pints of decoction are obtained. Let stand for
 some time, and strain. Advised in scrofula, two
 fl. ounces, three times a-day. *Bories.*

EXTRACT OF SARSAPARILLA.

R. Sarsaparilla, in coarse powder,
 one pound.
 Diluted alcohol, four pints.

Make a tincture by displacement, filter, distil
 off the alcohol, and evaporate to the consistence
 of an extract. *U. S. Ph.*

Dose, from ten to twenty grains, three or four
 times a-day.

FLUID EXTRACT OF SARSAPARILLA.

R. Sarsaparilla, bruised, sixteen ounces.
 Liquorice root, bruised,
 Guaiacum wood, rasped,
 Sassafras bark, sliced, each,

two ounces.
 Mezereon, sliced, six drachms.
 Diluted alcohol, eight pints.

Digest for fourteen days, strain, express, and
 filter. Evaporate on a water-bath to twelve fl.
 ounces, add twelve ounces of sugar, and remove
 from fire when this is dissolved. Dose, one fl.
 drachm, three or four times a-day.

W. Hodgson, Jr.

The only difference between this and the offi-
 cinal formula of the U. S. Ph., is the omission
 of the guaiacum wood in the latter.

ESSENCE OF SARSAPARILLA.

R. Sarsaparilla, bruised, ten ounces.
 Distilled water, six pints.

Macerate at a temperature of 120° for six hours
 and strain; repeat with same quantity of water.
 Mix the liquors, and evaporate in china vessels
 at 160°.

If reduced to ten fl. ounces (or to nine fl.
 ounces, with one fl. ounce of spirit added), one
 fl. drachm mixed with seven fl. drachms, is
 equal to the decoction of usual strength. If re-
 duced to five fl. ounces, one fl. drachm is equal
 to two fl. ounces of decoction. *Hening.*

R. Alcoholic extract of sarsa-
 parilla, one pound.
 Sherry wine, three pints.

Dissolve and filter. Half a fl. ounce to one fl.
 ounce, mixed with a quart of water, is equal to
 the decoction. *Beral.*

COMPOUND ESSENCE OF SARSAPARILLA.

R. Sarsaparilla, bruised, eight ounces.
 Boiling water, sufficient

to exhaust the root, by successive macerations;
 unite the liquors, and evaporate to ten fl. ounces;
 strain, and add, when cool,

Alcohol (.842),
 Tincture of guaia-
 cum, each, half a fl. ounce.
 White wine, one fl. ounce.
 Essence of sassafras, twelve drops.
 Extract of liquorice, two drachms.

Filter. Dose, a spoonful, morning and evening,
 in some convenient vehicle. *Cadet.*

SYRUP OF SARSAPARILLA.

R. Sarsaparilla, sliced, fifteen ounces.
 Boiling water, one gallon.

Macerate for twenty-four hours, boil down to
 four pints, strain, and add

Sugar, fifteen ounces,
and evaporate to proper consistence. *Ed. Ph.*
Dose, a tablespoonful.

R. Alcoholic extract of sarsaparilla, six ounces.
Water, four pints.

Dissolve by means of a water-bath, strain, and add

Sugar, eight pounds.
Dissolve whilst hot. *Beral.*

Each fl. ounce is equal to fifteen grains of the extract.

SARSAPARILLA SYRUP FOR MINERAL WATER.

R. Sarsaparilla,
Liquorice root, each,
finely bruised, two pounds (av.).
Sugar, thirty pounds (av.).
Oil of anise, } each, forty
" wintergreen, } drops.
" sassafras, }
" cinnamon, } five drops.
Water, sufficient.

Digest the roots in two gallons of warm water, for twelve hours; then transfer to a percolator, and pass two gallons of infusion. In this dissolve the sugar by the aid of heat, and to the syrup, when cool, add the oils, previously rubbed up with a little sugar. *A. Smith.*

R. Sarsaparilla, bruised,
Liquorice root, " each, one pound.
Cinnamon, "
Sassafras root, " each, six ounces.
Cloves, }
Aniseed, } each, two ounces.
Coriander, }
Red saunders,
Cochineal, each, one ounce and a half.
Alcohol, two pints.
Water, two gallons.

Digest together for four days, strain, and make a syrup with twenty-seven pounds (av.) of sugar. *A. Smith.*

COMPOUND SYRUP OF SARSAPARILLA.

R. Sarsaparilla, bruised, two pounds.
Guaiacum wood, rasped, three ounces.
Hundred-leaved roses, }
Senna, } each, two
Liquorice root, } ounces.
bruised, }
Oil of sassafras,
Oil of anise, each, five minims.
Oil of partridge-berry, three minims.
Diluted alcohol, ten pints.
Sugar, eight pounds.

Macerate the first five ingredients in the alcohol for fourteen days, express, and filter; evaporate the tincture on a water-bath, to four pints; filter, add the sugar, and form a syrup; then, having rubbed the oils with a little of the syrup, mix well with the remainder.

U. S. Ph.

R. Sarsaparilla, two pounds.
Bittersweet,
Pipsissewa, each, half a pound.
Guaiacum,
Liquorice root, each, four ounces.
Sassafras,
Partridge-berry-leaves,
each, two ounces.
Sugar, twelve pounds.

Reduce the ingredients to coarse powder, macerate them in diluted alcohol, for two days, put in a displacement apparatus, and displace slowly until two gallons of liquid have passed. Evaporate on a water-bath till reduced to six pints, then add the sugar, and form a syrup. *Staples.*

Dose, half a fl. ounce, three or four times a-day.

SYRUP OF LAFFECTEUR.

R. Sarsaparilla, bruised,
Marsh-reed grass, each, thirty ounces.
Borage flowers, eight ounces.
Senna,
White roses, each, two ounces.
Sugar,
Honey, each, six pounds.
Water, eighteen pints.

Boil the first two ingredients in half the water for an hour, strain off the decoction, and repeat the process with the remainder of the water, and boil two hours; towards close, add the senna, rose leaves, and borage; strain, mix the decoctions, add the sugar and honey, and boil to the consistence of syrup. Dose, six tablespoonfuls, early in the morning. *Ellis.*

R. Sarsaparilla, bruised, nine pounds.
Guaiacum wood, rasped,
China root, bruised, each, six pounds.
Yellow Peruvian bark,
bruised, three pounds.

Macerate for twenty-four hours in

Water, one hundred and forty pints.

Boil down to one-third, express, and strain. Repeat twice, with same proportions of water; strain, unite the three decoctions, and add

Purified molasses, thirty pounds.

Evaporate to the consistence of syrup, and pour boiling, on

Sassafras, rasped, five pounds.

Aniseed, four ounces.

Borage flowers, one pound and a half.

Finally strain.

Cadet.

One to three ounces a-day, in syphilitic and cutaneous disorders.

SYRUP OF GESNOUIN.

R. Syrup of sarsaparilla, four pounds.
 " senna, three ounces.
 Extract of borage, five ounces.
 Conserve of elder berries, one ounce.

Mix, with heat. Four to six spoonfuls, early in the morning, using the infusion of sarsaparilla for a drink. *Foy.*

SARSAPARILLA BEER.

R. Sarsaparilla, bruised, two pounds.
 Bark of guaiacum, powdered, eight ounces.
 Guaiacum wood, rasped, Aniseed,
 Liquorice root, each, four ounces.
 Mezereon, bark of root, two ounces.
 Molasses, two pounds.
 Cloves, bruised, half an ounce.
 Boiling water, four gallons.

Mix and shake, thrice a-day. Let ferment; when fully fermented, to be taken in doses of a small tumblerful, two or three times a-day. *Hancock.*

SASSAFRAS.

SASSAFRAS.

Two parts of the Sassafras, *Laurus sassafras* (*Sassafras officinale*), are officinal; the pith of the young branches and the bark of the root.

Sassafras is a medium-sized tree, found in most parts of the United States, growing in woods and open places, flowering before the appearance of the leaves.

Sex. Syst. Enneand. monog. *Nat. Syst.* Lauraceæ.

Nees, Pl. Med. i. 131. Griffith, Med. Bot. 551.

The pith is in slender, cylindrical pieces, very spongy and light, with a mucilaginous and somewhat aromatic taste, affording a clear mucilage on the addition of water. The bark is in small fragments, of a reddish-brown color, brittle, of a fragrant odor, and a pleasant aromatic taste. It is stimulant and diaphoretic.

INFUSION OF SASSAFRAS BARK.

R. Sassafras bark, half an ounce.
 Boiling water, one pint.

Macerate for six hours, and strain. As a stimulating diaphoretic. *Niemann.*

COMPOUND INFUSION OF SASSAFRAS BARK.

R. Sassafras bark, half an ounce.
 Liquorice root, two drachms.
 Boiling water, one pint.

Infuse. One-half, morning and evening. In scrofulous complaints. *Hufeland.*

INFUSION OF SASSAFRAS PITH.

R. Sassafras pith, one drachm.
 Water, one pint.

Macerate for three hours, and strain. As a soothing application in ophthalmia, and as a demulcent drink in diseases of the bowels, lungs, and bladder. *U. S. Ph.*

TINCTURE OF SASSAFRAS.

R. Sassafras bark, sliced, one part.
 Alcohol, six parts.

Macerate for three days, express, and filter. Dose, one fl. drachm. *Beral.*

COMPOUND TINCTURE OF SASSAFRAS.

R. Sassafras bark, bruised, four ounces.
 Sage, two ounces.
 Nutmeg, bruised, one ounce.
 Cinnamon, bruised, half an ounce.
 Star anise, three drachms.
 Alcohol, two pints.

Macerate for three days, express, and filter. *Wirt. Ph.*

As a stomachic and carminative.

OIL OF SASSAFRAS.

R. Bruised sassafras bark, at will.
 Water, sufficient.

Distil, and collect the oil which sinks to the bottom of the product. *Guibourt.*

Dose, two to ten drops, as a carminative and stimulant.

SCAMMONIUM.

SCAMMONY.

This is the concrete juice of the root of *Convolvulus scammonia*, a twining perennial plant, a native of Syria. It has a large, succulent root, affording a milky juice, which concretes by exposure to the air.

Sex. Syst. Pentand. monog. *Nat. Syst.* Convolvulaceæ.

Linn. Sp. Pl. 218. Griffith, Med. Bot. 477.

The best, or virgin scammony is light, resinous when broken, friable, not effervescing with an acid, of various shades of color, from dark ash to dark olive; forming an emulsion with water. The odor is peculiar and unpleasant, resembling old cheese; the taste is bitterish and acrid. It is an active purgative, causing pain, and usually operating with harshness when given alone, but much milder in combination. The dose is from five to fifteen grains.

POWDER OF SCAMMONY AND CREAM
OF TARTAR.

R. Powdered scammony,
Cream of tartar, equal parts.
Mix. *Ed. Ph.*
Dose, fifteen to thirty grains.

POWDER OF SCAMMONY AND SOOT.

R. Powdered scammony, one drachm.
Soot, one drachm and a half.
Powdered resin, two drachms.
Mix. Dose, twenty to thirty grains. *Beasley.*

COMPOUND POWDER OF SCAMMONY.

R. Scammony,
Hard extract of jalap, each,
two ounces.
Ginger, half an ounce.
Rub separately into fine powder, and mix.
Dose, ten to twenty grains. *Lond. Ph.*

PILLS OF SCAMMONY.

R. Powdered scammony, fifteen grains.
Sugar, ten grains.
Rub together, and add
Oil of caraway, four minims.
Make ten pills. Dose, three pills. *Copland.*

PILLS OF SCAMMONY AND OX GALL.

R. Powdered scammony, two drachms.
Inspissated ox gall, three drachms.
Extract of gentian, half an ounce.
Mix, and make one hundred and sixty-two pills.
Dose, four to six a-day. *Cadet.*

COMPOUND PILLS OF SCAMMONY.

R. Powdered scammony,
Extract of henbane,
Powdered gamboge,
Compound extract of colocynth,
Soap, each, twelve grains.
Water, sufficient.
Beat into mass, and form twelve pills. Dose,
two to three. *Guy's Hosp.*

R. Scammony, twenty-four grains.
Aloes,
Gamboge, each, twelve grains.
Ginger, one scruple.
Molasses, sufficient.
Rub together, and divide into twelve pills.
St. Bart's Hosp.

CONFECTION OF SCAMMONY.

R. Powdered scam-
mony, one ounce and a half.
Cloves, bruised,
Powdered ginger, each, six drachms.
Oil of caraway, half fl. drachm.
Syrup of roses, sufficient.

Rub the dry ingredients into a fine powder.
When the confection is to be used, add the
syrup, and rub well; then the oil of caraway,
and mix. Dose, twenty to forty grains.

Lond. Ph.

Dose, about half a drachm.

R. Powdered scammony,
" jalap, each, two drachms.
" squill,
" one drachm and a half.
" resin of jalap,
half a drachm.
Syrup of buckthorn, sufficient.

Mix, and form confection. A hydragogue pur-
gative, in dose of ten grains to a scruple. *Foy.*

EXTRACT OR RESIN OF SCAMMONY.

R. Scammony, in fine powder, at will,
Boil in successive portions of proof spirit, till
the spirit ceases to dissolve anything; filter,
distil off the spirit, pour off the watery solution
from the resin, wash this well with boiling
water, and dry in a heat not above 240°.

Ed. Ph.

Dose, five to ten grains.

EMULSION OF SCAMMONY.

R. Resin of scammony, seven grains.
Unskimmed milk, three fl. ounces.
Triturate the resin with a little of the milk,
and gradually add the rest. *Ed. Ph.*
The whole at a dose.

COMPOUND EMULSION OF SCAMMONY.

R. Scammony, seven grains.
Sugar, two drachms.
Triturate, and add gradually
New milk, three fl. ounces.
Cherry-laurel water,
three or four drops.
Take as a dose. *Planche*

SCAMMONY MIXTURE.

R. Resin of scammony, twelve grains.
Spirit of rosemary, one fl. drachm
Orange-flower water,
Syrup of peach-
flowers, each, one fl. ounce
Mix. *Cadet.*

SCAMMONY OIL.

R. Powdered scammony, one scruple.
Oil of almonds, one ounce.
Rub well together. Dose, half an ounce.

Van Mons.

ELIXIR OF SCAMMONY.

R. Scammony, two drachms.
Proof spirit, eight fl. ounces.
Heat, and then set fire to the spirit, and add
Sugar, four ounces.
When it is dissolved, extinguish the flame, and add

Syrup of violets, two fl. ounces.
Filter. It will give ten ounces, containing twelve grains of scammony to the ounce.

Guibourt.

SOAP OF SCAMMONY.

R. Scammony, one drachm.
Soap, two drachms.
Alcohol (.874), two fl. ounces.
Dissolve by means of a water-bath.

Guibourt.

PURGATIVE PASTILLES.

R. Scammony, six grains.
Tincture of senna, forty drops.
Carbonate of magnesia, five scruples.
White sugar, eight scruples.
Powdered liquorice, eight grains.
Gum tragacanth, five grains.
Oil of aniseed, one drop.
Syrup of violets, sufficient

to make eight pastilles. These are known as *Pastilles de Santé*, and are useful in cases of acid eructation and flatulence. One or two, taken early in the morning, act as a laxative; by repeating the dose every quarter of an hour, full purgation ensues.

Delvincourt.

PURGATIVE CHOCOLATE.

R. Chocolate, five drachms.
Scammony resin, six grains.
Calomel, one and a half grains.
Sugar, half a drachm.

Triturate the scammony and sugar together in a mortar; then add the calomel, and, when thoroughly mixed, add the chocolate, previously softened by the heat of a water-bath. Make into forms. To be taken dry, and followed by some bland liquid.

Mialhe.

SCILLA.

SQUILL

Is the bulb of *Scilla maritima*, a perennial plant growing near the sea-coast of the countries bordering on the Mediterranean. The

bulb is large, pyriform, and tunicated; there are two varieties, the red and the white, but identical in their medical qualities.

Sex. Syst. Hexand. monog. *Nat. Syst.* Liliaceæ.

Linn. Sp. Pl. 442. *Griffith, Med. Bot.* 652.

The bulb is sliced and dried for use; in this state it is in white, or yellowish-white, semi-transparent pieces, which are flexible when moist, but brittle when dry; of a mucilaginous, bitter taste, but scarcely any odor.

Squill is expectorant, diuretic, and in large doses emetic, and even purgative. Dose, as a diuretic or expectorant, one grain two or three times a-day; as an emetic, six to twelve grains.

POWDER OF SQUILL AND NITRE.

R. Powdered squill, six grains.
" nitre, one drachm.

Mix, and divide into six powders. One, three times a-day, in sugar and water. As a diuretic.

Ellis.

POWDER OF SQUILL AND IPECACUANHA.

R. Powdered squill, twelve grains.
" ipecacuanha, twenty-four grains.

Mix, and divide into six powders. One, every two hours, in chronic catarrh.

Foy.

POWDER OF SQUILL AND CREAM OF TARTAR.

R. Powdered squill, one drachm.
Cream of tartar, nine drachms.

Mix. Twenty to thirty grains, two or three times a-day, as a diuretic.

Guy's Hosp.

POWDER OF SQUILL AND SULPHUR.

R. Powdered squill, one part.
Sulphur, two parts.
Powdered sugar, three parts.

Mix. Dose, from five to twenty grains, according to age, in catarrhs.

Guibourt.

POWDER OF SQUILL AND CALOMEL.

R. Powdered squill, three grains.
Calomel,
Powdered gamboge, each, half a grain.

Mix. To be taken in the evening. As a diuretic.

A. Cooper.

COMPOUND POWDER OF SQUILL.

R. Powdered squill,
" foxglove, each, one grain.
Oil of juniper, two drops.
Borate of soda,
Powdered liquorice, each, one scruple.
Powdered cinnamon, two grains.

Mix. To be repeated, two or three times a-day, as a diuretic. *Radiis.*

R. Powdered squill,
" foxglove, each, two grains.
Resin of jalap, six grains.
Acetate of potassa, thirty grains.
Sulphate of soda, two drachms.

Mix. To be taken early in the morning, as a purgative and diuretic. *Bories.*

PILLS OF SQUILL AND CALOMEL.

R. Calomel,
Powdered squill, each, twelve grains.
Conserve of roses, sufficient.

Mix, and form twelve pills. One, night and morning. In dropsy, arising from visceral degeneration. *Ellis.*

PILLS OF SQUILL AND AMMONIAC.

R. Powdered squill, thirty grains.
Ammoniac, one drachm and a half.
Extract of hemlock, thirty grains.

Mix, and form thirty pills; two, twice a-day. In asthma and chronic catarrh. *Ellis.*

PILLS OF SQUILL AND CROTON OIL.

R. Compound pills of squill, two scruples.
" extract of colocynth, two scruples.

Croton oil, six minims.

Mix, and form eighteen pills. Three, twice a-week, in dropsy. *Selwyn.*

PILLS OF SQUILL AND IPECACUANHA.

R. Powdered squill, half an ounce.
" ipecacuanha, two drachms.
Extract of opium, fifteen grains.
Butter of cocoa, one ounce.
Syrup of gum, sufficient.

Beat together, and make pills of four grains. One, three times a-day, as an expectorant. *Cadet.*

R. Powdered squill, twelve grains.
" ipecacuanha, twelve grains.
Extract of dandelion, three scruples.
Blue pill, ten grains.
Simple syrup, sufficient.

Mix, and make twenty-four pills. Two, to be taken morning and evening, in dropsy. *Ainslie.*

PILLS OF SQUILL AND HENBANE.

R. Powdered squill, half a drachm.
Extract of henbane, two scruples.
Myrrh, one drachm and a half.
Water, sufficient.

Beat together, and make thirty pills. Two, morning and evening, as an expectorant. *Paris.*

COMPOUND PILLS OF SQUILL.

R. Calomel, three grains.
Fresh squill,
Ammoniac, each, one scruple.
Dover's powder, half a drachm.
Conserve of roses, sufficient.

Make mass, and divide into thirty pills. One, three or four times a-day. In troublesome catarrh. *Latham.*

R. Powdered squill, half a drachm.
Calomel, twenty-four grains.
Tartar emetic, six grains.
Powdered opium, eighteen grains.
Syrup, sufficient.

Beat together, and make twenty-four pills. One at night, in catarrh. *Ellis.*

R. Powdered squill, one drachm.
" ginger, two drachms.
" ammoniac, each, three drachms.
Soap, three drachms.
Syrup, sufficient.

Beat into mass, and form one hundred and twenty pills. *U. S. Ph.*

Five to ten grains, three or four times a-day, as a stimulant expectorant.

R. Powdered squill, fifteen grains.
" ammoniac, one drachm and a half.
" seneka, two drachms.
Golden sulph. of antimony, half a drachm.

Extract of dandelion, sufficient.

Mix, and make pills of two grains. Diuretic. *Berends.*

R. Powdered squill, twenty-four grains.
Calomel, sixteen grains.
Soap, one drachm and a half.
Galbanum, two drachms.
Extract of dandelion, three drachms and a half.
Essence of orange, sufficient.

Mix, and make pills of two grains. Three or four a-day, as purgative and diuretic. *Pideret.*

COMPOUND SYRUP OF SQUILL.

R. Bruised squill,
" seneka, each, four ounces.
Tartar emetic, forty-eight grains.
Water, four pints.
Sugar, three pounds and a half.

Pour the water on the squill and seneka, boil to one-half, strain, and add the sugar; evaporate to three pints, and, whilst hot, add the tartar emetic. *U. S. Ph.*

R. Squill,
Seneka, in coarse powder, each,
four ounces.
Tartar emetic, forty-eight grains.
Alcohol, half a pint.
Water, sufficient.
Sugar, three pounds and a half.

Mix the alcohol with two pints and a half of the water, and macerate the squill and seneka in the mixture for twenty-four hours. Put the whole into a displacement apparatus, and add as much water as may be necessary to obtain three pints of filtered liquor. Boil this for a few minutes, evaporate to one-half, and strain; then add the sugar, and evaporate to three pints; lastly, dissolve the tartar emetic in the syrup, whilst hot. *U. S. Ph.*

Known as *Coze's Hive Syrup*. Dose, as an expectorant, twenty to thirty drops, for adults; for children, five to ten drops; in croup, ten drops to a fl. drachm, repeated till vomiting is produced.

R. Extract of seneka and squill,
one pound.

Clarified honey, at 160°, six pounds.

Mix, and add
Tartar emetic, sixteen grains
to each sixteen fl. ounces of the syrup.
J. H. Ecky.

R. Oxymel of squill,
one fl. ounce and a half.
Syrup of ipecacuanha,
" poppies, each, two fl. ounces.
" orange-flowers,
half an ounce.

Mix. Dose, half to one fl. ounce in whooping-cough. *Cadet.*

WINE OF SQUILL.

R. Squill, one ounce.
White wine, sixteen fl. ounces.
Macerate for twelve days, express, and filter.
Par. Cod.

COMPOUND WINE OF SQUILL.

R. Bruised squill, one ounce.
" orange-peel,
" calamus, each, three drachms.
" juniper berries, two drachms.
White wine, four pints.

Macerate for three days, express, filter, and add
Oxymel of squill, two fl. ounces.

Mix.
R. Dried squill, bruised, } each,
Rhubarb, } one drachm.
Juniper berries, }
Cinnamon, three drachms.
Zedoary, two drachms.

Carbonate of
potassa, one drachm and a half.
White wine, two pints.
Macerate, express, and filter. Dose, two to four small wineglassfuls a-day, as a diuretic.
Taddei.

BITTER WINE OF SQUILL.

R. Squill,
Root of swallow-
wort, each, half an ounce.
Peruvian bark,
Canella,
Lemon-peel, each, two ounces.
Angelica, half an ounce.
Wormwood,
Balm, each, one ounce.
Juniper berries,
Mace, each, half an ounce.
White wine, eight pints.

Digest by a gentle heat, for four days, express, and filter. As a diuretic, in doses of one to three fl. ounces. *Cottureau.*

TINCTURE OF SQUILL.

R. Squill, four ounces.
Diluted alcohol, two pints.
Macerate for fourteen days, express, and filter.
It may also be prepared by percolation. *U. S. Ph.*

Dose, as diuretic or expectorant, twenty to forty drops.

ALKALINE TINCTURE OF SQUILL.

R. Squill, two ounces.
Solution of potassa, two fl. drachms.
Diluted alcohol, twelve fl. ounces.
Macerate for three days, express, and filter.
Van Mons.

TINCTURE OF SQUILL AND ELATERIUM.

R. Tincture of squill,
Vinegar of colchi-
cium, each, two fl. drachms.
Spirit of nitric ether, one fl. ounce.
Elaterium, one grain.
Mix. Dose, fifteen minims to one fl. drachm.
St. Bart.'s Hosp.

COMPOUND TINCTURE OF SQUILL AND BENZOIN.

R. Squill,
Orris root,
Elecampane, each, three ounces.
Benzoin, two drachms.
Liquorice root,
Aniseed,
Myrrh, each, four scruples.

Ammoniac, two scruples.
Saffron, eighteen grains.
Diluted alcohol, one pint, six fl. ounces.
Macerate for fifteen days, express, and filter.
Celebrated as *Wedel's elixir*. Dose, forty to sixty drops, in catarrh, asthma, &c. *Cadet.*

ETHEREAL TINCTURE OF SQUILL.

R. Squill, one ounce and a half.
Muriatic ether,
Spirit of juniper, each, half a pound.
Water, sufficient.
Distil one pound. Dose, ten to twenty drops, in water, as a diuretic. *Augustin.*

R. Bruised squill, four ounces.
Spirit of nitric ether, two pints.
Digest for eight days, and filter. Dose, half a fl. drachm, as an expectorant and diuretic. *Mettauer.*

MIXTURE OF SYRUP OF SQUILL, &c.

R. Syrup of squill, half fl. ounce.
Paregoric elixir, two fl. drachms.
Antimonial wine, one fl. drachm.
Powdered gum Arabic, half a drachm.
Distilled water, three fl. ounces.
Mix. A teaspoonful, every two or three hours, as an expectorant; suited to children. *Ellis.*

R. Syrup of squill,
Peppermint water, each, two fl. ounces.
Paregoric elixir,
Compound spirit of lavender, each, half fl. ounce.
Syrup, one fl. ounce.
Mix. A spoonful, three or four times a-day. A valuable expectorant. *Christison.*

MIXTURE OF OXYMEL OF SQUILL AND VALERIAN.

R. Powdered valerian, two drachms.
Oxymel of squill, one fl. ounce.
Laudanum, twenty drops.
Water, one fl. ounce.
Mix. A teaspoonful, every hour, after an emetic, in croup. *Kimbel.*

MIXTURE OF EXTRACT OF SQUILL AND RHUBARB.

R. Extract of squill, sixteen grains.
Liquid acetate of potassa, half an ounce.
Parsley water, four fl. ounces.
Syrup of rhubarb, one fl. ounce.
Mix. As a diuretic. *Radius.*

MIXTURE OF OXYMEL OF SQUILL AND AMMONIAC.

R. Oxymel of squill, six fl. drachms.
Ammoniac, one drachm.
Extract of elecampane, twenty-four grains.
Syrup of hyssop, six fl. drachms.
Hyssop water, four fl. ounces.
Mix. A tablespoonful every hour, as an expectorant. *Foy.*

MIXTURE OF VINEGAR OF SQUILL.

R. Vinegar of squill, one fl. drachm.
Barley water, five fl. ounces.
Syrup of saffron, one fl. ounce.
Mix. A tablespoonful, three or four times a-day, in troublesome cough. *Ainslie.*

MIXTURE OF VINEGAR OF SQUILL AND AMMONIA.

R. Vinegar of squill, two fl. drachms.
Solution of acetate of ammonia,
Compound spirit of horseradish, each, one fl. ounce and a half.
Diluted pimento water, four fl. ounces and a half.
Mix. Dose, two fl. ounces, three times a-day. As a diuretic. *St. Bart's Hosp.*

MIXTURE OF OXYMEL OF SQUILL AND IPECACUANHA.

R. Ipecacuanha, one drachm and a half.
Lemon peel, two drachms.
Cream of tartar, half an ounce.
Water, four fl. ounces.
Boil, strain, and add
Oxymel of squill,* half a fl. ounce.
Mix well; as an expectorant. *Cadet.*

MIXTURE OF OXYMEL OF SQUILL AND MARSH MALLOW.

R. Oxymel of squill,
Syrup of marsh mallow,
Mucilage of gum Arabic, equal parts.
Mix. As an expectorant; in spoonful doses. *Foy.*

EMULSION WITH SQUILL.

R. Oil of almonds,
Mucilage of gum Arabic, each, one fl. ounce.
Wine of squill, eight fl. scruples.
" ipecacuanha, four fl. scruples.
Syrup of tolu, two fl. ounces.
Water of hyssop, four fl. ounces.
Water, eight fl. ounces.
Make an emulsion. *Beral.*

OINTMENT OF SQUILL.

R. Squill, three ounces.
 Solution of potassa, sufficient.
 Reduce to mucilage by boiling, and add
 Lard, sufficient
 to form an ointment. As a friction to indolent
 tumors. *Hufeland.*

EXTRACT OF SQUILL.

R. Squill, one pound.
 Proof spirit, four pints.
 Macerate for some days, express, strain and
 filter; digest the residue with two pints of
 spirit, express, strain, and filter. Distil off the
 spirit, and evaporate to proper consistence.
 Dose, one to three grains. *Par. Cod.*

ACETIC EXTRACT OF SQUILL.

R. Bruised squill, one pound.
 Acetic acid, three fl. ounces.
 Distilled water, one pint.
 Mix; digest with a gentle heat for forty-eight
 hours, express by strong pressure, and evaporate,
 without straining, to the proper consistence.
 Dose, half a grain. *Niblett.*

ELECTUARY OF SQUILL.

R. Oxymel of squill, two fl. ounces.
 Cream of tartar, three ounces.
 Mix. Dose, two drachms. *Beasley.*
 R. Powder of squill and nitre,
 Ammoniac,
 Tartrate of potassa, each, two drachms.
 Muriate of ammonia, one scruple.
 Syrup of cinnamon, sufficient.
 Mix. Dose, two drachms, three times a-day, in
 dropsy. *Radius.*

VINEGAR OF SQUILL.

R. Bruised squill, four ounces.
 Diluted acetic acid, two pints.
 Macerate the squill with the diluted acetic acid
 in a close glass vessel for seven days; express,
 let settle, and pour off the clear liquid. *U. S. Ph.*

Diuretic and expectorant. Dose, thirty minims
 to a fl. drachm.

OXYMEL OF SQUILL. (*Sp. Gr.* 1.32).

R. Vinegar of squill, two pints.
 Clarified honey, one pint and a half.
 Mix, and evaporate by water-bath to the proper
 consistence. *U. S. Ph.*
 Dose, one to two fl. drachms.

SYRUP OF SQUILL.

R. Vinegar of squill, one pint
 Sugar, two pounds.
 Mix, and form syrup. *U. S. Ph.*
 As expectorant. Dose, about a fl. drachm.

AROMATIC SYRUP OF SQUILL.

R. Vinegar of squill, half a pound.
 Syrup of cinnamon,
 " ginger, each, one pound.
 Mix. As a stimulating expectorant. *Swediaur.*

SCOPARIUS.

BROOM.

This is the fresh tops of the *Cytisus scopari-*
us, a shrub, native of many parts of Europe,
 and cultivated in this country, with numerous,
 bright yellow flowers.

Sex. Syst. Diadelph. decand. *Nat. Syst.* Fa-
 baceæ.

Link. enum. ii. 241. *Griffith, Med. Bot.* 233.
 The officinal portion is the tops of the branch-
 es, but the seeds are also used. The branches
 have a bitter, nauseous taste, and a strong, pe-
 culiar odor. It is a diuretic and cathartic, and
 even emetic, and has been advantageously used
 in dropsy.

INFUSION OF BROOM.

R. Broom tops, one ounce.
 Boiling water, one pint.
 Digest for four hours in a covered vessel, and
 strain. *Lond. Ph., 1836.*
 Dose, two to four fl. ounces three times a-day.

DECOCTION OF BROOM.

R. Broom tops,
 Juniper tops, each, half an ounce.
 Cream of tartar, two drachms and
 a half.
 Water, one pint and a half.
 Boil down to a pint, and strain. *Ed. Ph.*

COMPOUND DECOCTION OF BROOM.

R. Broom tops,
 Juniper berries,
 Dandelion root, each, half an ounce.
 Water, one pint and a half.
 Boil down to a pint, and strain. *Lond. Ph.*
 Dose, same as infusion.

EXTRACT OF BROOM.

R. Tops of broom, at will

Boil, in eight times their weight of water, down to one-half; express, let settle, and strain. Evaporate by vapor-bath to proper consistence.

Dub. Ph., 1826.

Dose, a scruple to a drachm.

CONSERVE OF BROOM.

R. Broom flowers, one part.
Sugar, two parts.
Rub together. Dose, a drachm to two drachms, in gout, rheumatism, &c. *Van Mons.*

SCROPHULARIA NODOSA.

FIGWORT.

A semi-aquatic plant, native of many parts of Europe, flowering in July.

Sex. Syst. Didyn. angios. *Nat. Syst.* Scrophulariaceæ.

Linn. Sp. Pl. 863. *Griffith, Med. Bot.* 518.

The leaves are the part used; they have a rank, unpleasant odor, and a bitter, disagreeable taste. They have been used as a tonic, diuretic, and anthelmintic, but are principally used as an external discutient application.

OINTMENT OF FIGWORT.

R. Fresh leaves of figwort, two pounds.
Prepared lard, each, two pounds.
" suet, one pound.
Boil till the leaves are crisp, express, and strain. *Dub. Ph.*, 1826.

As an application to piles, painful swellings, and cutaneous eruptions.

OIL OF FIGWORT.

R. Leaves of figwort, three parts.
Olive oil, one part.
Infuse, express, and strain. Used as above. *Van Mons.*

SENEGA.

SENEKA.

The officinal name for the root of *Polygala senega*, a native, perennial plant, found in most parts of the United States, on hill sides and in dry woods, flowering from June to August.

Sex. Syst. Diadelph. octand. *Nat. Syst.* Polygalaceæ.

Linn. Sp. Pl. 990. *Griffith, Med. Bot.* 225.

The root is contorted, knotted, tapering, branched with a ridge or carina. It has a thick, resinous, greyish-yellow, cortical layer, which is the active portion. It has a peculiar, but unpleasant odor; and a mucilaginous, sweetish taste, which becomes acrid and irritating. Its action is very various, being emetic, sudorific, diuretic, and expectorant, and even emme-

nagogue and purgative. It is principally used as a stimulating expectorant and diuretic. The dose, in powder, is from ten to twenty grains; but it is seldom given in this form.

COMPOUND POWDER OF SENEKA.

R. Powdered seneka, half an ounce
Cream of tartar, six drachms.
Magnesia, two drachms.
Mix. A teaspoonful, twice a-day, in pannus of the eye. *Radius.*

PILLS OF SENEKA.

R. Powdered seneka, six drachms.
Extract of dandelion, sufficient.
Mix, and make pills of two or three grains. Six to twelve, three times a-day, in hypopyon and pannus. *Radius.*

INFUSION OF SENEKA.

R. Seneka, ten drachms.
Boiling water, (Imp.), one pint.
Infuse for four hours in a covered vessel, and strain. *Ed. Ph.*
Dose, one to three fl. ounces.

COMPOUND INFUSION OF SENEKA.

R. Seneka, two drachms.
Boiling water, six fl. ounces.
Infuse, strain, and add
Syrup of tolu, one fl. ounce.
Ammoniac, half a drachm.
Mix well. A tablespoonful, every two hours. In catarrh. *Ellis.*

DECOCTION OF SENEKA.

R. Seneka, bruised, one ounce.
Water, one pint and a half.
Boil down to one pint, and strain. *U. S. Ph.*
Dose, a tablespoonful, every three hours.

R. Seneka, bruised, one ounce.
Liquorice root, half an ounce.
Distilled water, one pint and a half.
Boil down to a pint, and strain. Dose, a tablespoonful, in pectoral diseases. *Ellis.*

R. Bruised seneka, one ounce.
Orange peel, half an ounce.
Water, one pint.

Simmer by a slow fire, till reduced one-third; add orange peel; cover till cold, then strain. Four fl. ounces during the day, at first, and then increase, as an emmenagogue. *Chapman*

SYRUP OF SENEKA.

R. Bruised seneka, four ounces.
Water, one pint.
Sugar, one pound.

Boil the seneka with the water, to one-half; strain, add the sugar, and form syrup.

U. S. Ph.

R. Coarsely-powdered seneka,
four ounces.
Alcohol, half a pint.
Water, one pint and a half.
Sugar, fifteen ounces.

Mix the alcohol and water, pour half a pint of the liquid on the seneka, macerate for twelve hours, transfer to a percolator, pour on the remaining liquid, evaporate the resulting tincture to half a pint, add the sugar, and form a syrup.

U. S. Ph.

Dose, one to two fl. drachms, in pectoral complaints.

EMULSION WITH SENEKA.

R. Bruised seneka, half an ounce.
Water, nine fl. ounces.

Boil down one-third, strain, and add

Camphor, one scruple.
Gum Arabic, two drachms.

Make an emulsion, and add

Compound tincture of ammonia,
two fl. drachms.

A tablespoonful, every two hours, as a stimulating expectorant.

Phæbus.

PECTORAL JELLY WITH SENEKA.

R. Bruised seneka, two drachms.
Water, twelve fl. ounces.

Boil down to nine fl. ounces, strain, and add

Isinglass, sufficient.
Syrup of orange flowers, one ounce.

Form a jelly. A tablespoonful, occasionally.

Radiis.

MIXTURE WITH SENEKA.

R. Seneka, bruised, two drachms.
Water, sufficient

to obtain eight fl. ounces of strained decoction; add

Camphor, one drachm.

Mix. A tablespoonful, occasionally. *Brera.*

R. Infusion of seneka, four fl. ounces.
Syrup of ipecacuanha, one fl. ounce.
Oxymel of squill, three fl. drachms.
Tartar emetic, one grain and a half.

Mix. A tablespoonful every quarter of an hour, till vomiting is produced. In croup.

Jadelot.

R. Bruised seneka,
one drachm and a half.

Boiling water, sufficient
to obtain four fl. ounces of the strained decoction; add

Antimonial wine, two fl. scruples.
Syrup of marsh mallow, one fl. ounce.

A small spoonful, occasionally, in the second stage of croup, as an expectorant. *Wendt.*

EXTRACT OF SENEKA AND SQUILL.

R. Seneka,
Squill, each, in coarse powder,
two pounds.
Diluted alcohol, sufficient

to cover; transfer to a displacement apparatus pass the alcohol through till nearly tasteless; strain, distil off spirit, and evaporate to four pounds.

R. Extract of seneka and squill,
one pound.
Honey, six pounds.

Mix at temperature of 160° F., and add sixteen grains of tartar emetic to every pint of the syrup.

This is proposed as a substitute for *Coxe's Hive Syrup*. The dose is the same as that of the official preparation. *J. H. Ecky.*

POLYGALIC ACID.

R. Seneka, in powder, sixteen ounces.
Alcohol (35° B.), six pints.
Ether, sufficient.
Purified animal charcoal, two ounces.

Boil the seneka in four pints of the alcohol for fifteen minutes; suffer it to cool, then throw the whole on a displacement filter, and when the liquid ceases to pass, add more alcohol, until six pints of tincture are obtained. Distil off the alcohol till of a syrupy consistence, wash this with ether, to remove fatty matter, and throw the residue into several times its bulk of concentrated alcohol. After standing for twenty-four hours, the polygalic acid will be precipitated as a light brown powder, which is to be collected on a filter, washed with cold, strong alcohol, and dried. It may be rendered more pure by boiling it with alcohol and the animal charcoal, filtering and collecting the powder.

W. Procter, Jr.

SYRUP OF POLYGALIC ACID.

R. Polygalic acid, eight grains.
Boiling water, half a fl. ounce.
Syrup of gum Arabic, one fl. ounce and a half.

Dissolve and mix.

Dose, a teaspoonful.

W. Procter, Jr.

SENNA.

SENNA.

Under this name is included the leaves of several species of *Cassia*, as *C. acutifolia*, *C. elongata*, *C. obovata*, &c., forming what are termed

Alexandria, Tripoli, India, and Mecca sennas. They are all perennial plants, growing in Africa and Asia.

Ser. Syst. Decand. monog. *Nat. Syst.* Fabaceæ.

Christison, Dispen. 858. Griffith, Med. Bot. 254.

The leaves only are officinal in the U. S. and British Pharmacopœias, but the pods are recognized by several of the European authorities. The leaves have a faint, but unpleasant odor, and a somewhat bitter, very nauseous taste. Senna is an active purgative, usually causing nausea and colic when given alone. It is seldom administered in substance; the dose is from half a drachm to two drachms.

COMPOUND POWDER OF SENNA.

R. Powdered senna,
Cream of tartar, each, two ounces.
Powdered scammony, half an ounce.
“ ginger, two drachms.

Mix. Dose, a scruple to a drachm.
Lond. Ph., 1788.

POWDER OF SENNA AND GUAIAIACUM.

R. Powdered senna,
“ guaiacum, equal parts.

Mix. Dose, two teaspoonfuls, twice or thrice a-day. In jaundice. *Augustin.*

ALCOHOLIC EXTRACT OF SENNA.

R. Senna, one part.
Alcohol, five parts.

Heat slowly on a water-bath, to boiling point, and let cool; after twenty-four hours, express and strain; repeat process with same quantity of alcohol, unite the tinctures, distil off the alcohol, and evaporate. *Guibourt.*

PILLS OF SENNA.

R. Powdered senna, one drachm.
Extract of dandelion, sufficient.

Mix, and make thirty pills. Dose, five to eight. *Hufeland.*

CONFECTION OF SENNA.

[LENITIVE ELECTUARY.]

R. Senna, eight ounces.
Coriander, four ounces.
Liquorice root, bruised, three ounces.
Figs, one pound.
Pulp of prunes,
“ tamarinds,
“ purging cassia, each,
half a pound.
Sugar, two pounds and a half.
Water, four pints.

Rub the senna and coriander together, and separate ten ounces with a sieve. Boil the residue with the figs and liquorice root in the water, to one-half, express, and strain. Evaporate by a water-bath to a pint and a half, add the sugar and form a syrup; then rub the pulps with the syrup, add the sifted powder, and mix well. *U. S. Ph.*

A good laxative. Dose, about two drachms, taken at bedtime.

COMPOUND CONFECTION OF SENNA.

R. Confection of senna, two ounces.
Powdered jalap, one drachm.
Cream of tartar, two drachms.
Powdered ginger,

one drachm and a half.

Syrup, sufficient.

Mix. Dose, one drachm. *Beasley.*

ELECTUARY OF SENNA AND RHUBARB.

R. Powdered rhubarb,
“ senna, each, two drachms.
“ fennel, three drachms.
“ liquorice, half an ounce.

Pulp of purging cassia,

“ tamarinds, each, two ounces.

Syrup of male fern, sufficient.

Mix. Dose, two drachms to half an ounce. *Van Mons.*

ELECTUARY OF SENNA AND CREAM OF TARTAR.

R. Powdered senna, three drachms.
Cream of tartar, one drachm.
Manna, three ounces and a half.
Pulp of purging cassia,

“ tamarinds,

each, one ounce and a half.

Syrup of violets, sufficient.

Mix. Dose, about two drachms. *Van Mons.*

ELECTUARY OF SENNA AND FIGS.

R. Powdered senna, half an ounce.
Pulp of figs, sufficient.

Make an electuary. A piece the size of a hazel nut, night and morning, to obviate costiveness. *Ellis.*

ELECTUARY OF SENNA AND SULPHUR.

R. Powdered senna,
Sulphur,
Cream of tartar, each, one ounce.
Manna, six ounces.
Pulp of tamarinds, eight ounces.
Syrup of senna, sufficient.

Mix. Said to be useful in hemorrhoids. Dose, two to three drachms. *Swediaur.*

COMPOUND WINE OF SENNA.

R. Senna,	one ounce.
Rhubarb,	six drachms.
Cloves,	
Saffron, each,	one drachm.
Sherry wine,	two pints.

Macerate for five days, often agitating, and decant. As a tonic purgative. Dose, about two fl. ounces. *Pierquin.*

R. Senna,	four ounces.
Coriander,	
Fennel, each,	two drachms.
Sherry wine,	two pints and a half.

Digest for three days; add

Stoned raisins,	three ounces.
-----------------	---------------

Macerate for twenty-four hours, express, and strain. *Swed. Ph.*

COMPOUND TINCTURE OF SENNA.

R. Senna,	three ounces and a half.
Caraway, bruised,	three drachms and a half.
Cardamom, bruised,	one drachm.
Stoned raisins,	five ounces.
Proof spirit,	two pints.

Macerate for fourteen days, express, and filter. This is the old *Elixir Salutis*. *Lond. Ph.*

A warm, stomachic purgative, in atonic gout, &c., in doses of two fl. drachms to a fl. ounce.

TINCTURE OF SENNA AND JALAP.

R. Senna,	three ounces.
Jalap, in powder,	one ounce.
Coriander, bruised,	
Caraway, bruised, each,	half an ounce.
Cardamom, bruised,	two drachms.
Sugar,	four ounces.
Diluted alcohol,	three pints.

Macerate for fourteen days, express, and filter. *U. S. Ph.*

Similar to the last; used in same cases, and in like doses.

TINCTURE OF SENNA AND GENTIAN.

R. Senna,	four ounces.
Gentian, bruised,	
Orange peel, bruised, each,	one ounce.
Proof spirit,	sixty-four ounces.

Infuse for some days, often stirring, express, and filter. *Swediaur.*

INFUSION OF SENNA.

R. Senna,	one ounce.
Coriander, bruised,	one drachm.
Boiling water,	one pint.

Macerate for an hour, in a covered vessel, and strain. *U. S. Ph.*

Dose, about four fl. ounces.

COMPOUND INFUSION OF SENNA.

R. Senna,	half an ounce.
Sulphate of magnesia,	
Manna, each,	one ounce.
Fennel seed,	one drachm.
Boiling water,	half a pint.

Macerate in a covered vessel till cool, and strain. Dose, one-third, every four or five hours, till it operates. *Wood.*

BLACK DRAUGHT.

R. Senna,	
Mint, each,	ten drachms.
Boiling water,	two pints.

Macerate for an hour, strain, and add

Sulphate of magnesia,	eight ounces.
-----------------------	---------------

Dose, two to four fl. ounces. *Guy's Hosp.*

INFUSION OF SENNA AND TAMARINDS.

R. Senna,	one drachm.
Tamarinds,	one ounce.
Coriander, bruised,	one drachm.
Brown sugar,	half an ounce.
Boiling water,	eight fl. ounces.

Infuse for four hours in a covered vessel, and strain. Dose, two to four fl. ounces. *Ed. Ph.*

INFUSION OF SENNA AND COFFEE.

R. Senna,	two drachms.
Roasted coffee,	one drachm.
Boiling water,	
Hot milk, each,	three fl. ounces.

Infuse for twelve hours, and strain. To be taken at once, in the morning. *Foy.*

R. Senna,	ten grains.
Hot infusion of coffee,	
" milk, each,	at will.

Mix, and, when cool, strain, and add

Sugar,	sufficient.
--------	-------------

An agreeable purge for children. The amount of senna to be increased according to the age. *Guersant and Blake.*

R. Senna,	
Sulphate of magnesia,	
Coffee, roasted and	
ground, each,	half an ounce.
Boiling water,	four fl. ounces.

Digest for half an hour, strain, and add

Sugar,	one and a half ounces.
--------	------------------------

As efficient as the "black draught," but much more agreeable. *Combes.*

R. Coffee, roasted and ground,
Senna, each, one ounce.
Boiling water, half a pint.

Make an infusion, and evaporate to one ounce, and add

Simple syrup, three fl. ounces.
Mix. *Combes.*

INFUSION OF SENNA AND BUCKTHORN.

R. Senna, two drachms.
Sulphate of soda, half an ounce.
Boiling water, four fl. ounces.
Syrup of buckthorn, one fl. ounce.

Infuse, cool, and strain. *Ellis.*

INFUSION WITH TINCTURE OF SENNA.

R. Infusion of senna, six fl. drachms.
Tincture of senna, two fl. drachms.
Manna, two drachms.
Spear-mint water,
Distilled water, each, one fl. ounce and a half.

Mix. One-half at a dose; to be repeated, if it does not operate. *Abernethy.*

ANTHELMINTIC EMULSION.

R. Infusion of senna, ten fl. drachms.
Syrup of buckthorn, one fl. drachm.
Confection of scammony, two scruples.
Copaiba, thirty minims.
Spirit of turpentine, six fl. drachms.

Mix, and make an emulsion. Very efficient as an anthelmintic; to be taken early in the morning. *Med. Chirur. Rev.*

SENN A MIXTURE.

R. Infusion of senna, one fl. ounce.
Tincture of jalap, one fl. drachm.
Sulphate of magnesia, four drachms.
Syrup of ginger, one fl. drachm.

Mix. As a purgative draught. *Ainslie.*

INFUSION OF SENNA AND RHUBARB.

R. Senna, six drachms.
Manna, one ounce.
Bruised rhubarb,
" cardamom, each,

Boiling water, two drachms.
one pint.

Infuse for one hour, and strain. Half a tea-cupful every hour, till it operates. *Ellis.*

INFUSION OF SENNA AND LEMON JUICE.

R. Senna, one ounce and a half.
Fresh lemon-peel, one ounce.

Lemon juice, one fl. ounce.
Boiling water, sixteen fl. ounces.
Infuse for two hours, and strain. *Lond. Ph., 1746.*

SYRUP OF SENNA.

R. Senna, two ounces.
Fennel-seed, bruised, one ounce.
Boiling water, one pint.
Sugar, fifteen ounces.

Digest the senna and fennel seed in the water, with a gentle heat, for an hour; strain, add the sugar, and form a syrup. *U. S. Ph*

Dose, one to two fl. drachms, to children.

FLUID EXTRACT OF SENNA.

R. Senna, one pound.
Alcohol, four pints.
Boiling water, eight pints.
Sugar, ten ounces.
Oil of fennel, four drops.

Pour the boiling water on the senna in a covered vessel, and set aside until cold; add the alcohol, and macerate for twelve days; strain with pressure, decant, filter, and distil off the alcohol; evaporate on a water-bath, to twelve fl. ounces, to which add the sugar, and dissolve by a gentle heat; when cold, add the oil of fennel, and mix. *C. Ellis, 1835.*

Dose, two fl. drachms.

R. Senna, well
bruised, sixteen ounces (Troy).
Diluted alcohol, sufficient.
Oil of caraway,
Oil of anise, each, sixteen drops.
Sugar, twelve ounces.

Macerate the senna in two pints of the diluted alcohol for twenty-four hours; displace slowly with the same menstruum until four pints pass. Evaporate this on a water-bath to ten fl. ounces, and add the sugar; when this is dissolved, remove from the fire, and add the oils, dissolved in diluted alcohol.

Duhamel, modified by Procter.

R. Senna, in coarse
powder, two pounds and a half.
Sugar, twenty ounces.
Oil of fennel, one fl. drachm.
Compound spirit of
ether, two fl. drachms.
Diluted alcohol, four pints.

Mix the senna with the diluted alcohol, and macerate for twenty-four hours; introduce into a a percolator, and pour in a mixture of two parts of water and one part of alcohol, until one gallon and a half shall have passed. Evaporate this on a water-bath, to twenty fl. ounces, filter, add the sugar, dissolve, and then add the compound spirit of ether, holding the oil of fennel in solution. *U. S. Ph.*

Dose, half a fl. ounce.

R. Senna, fifteen pounds.
Boiling water, sufficient.
Exhaust the senna by process of displacement, evaporate the fluid in vacuo, to ten pounds, and add ten pounds of molasses, previously boiled to a candy; add further, twenty-four fl. ounces of rectified spirit, and water sufficient to make fifteen pints. *Duncan.*

Dose, two drachms.

SYRUP OF SENNA AND RHUBARB.

R. Syrup of senna, four fl. ounces.
" rhubarb, two fl. ounces.
" cinnamon, one fl. ounce.
Mix. *Van Mons.*

SYRUP OF SENNA AND CIDER.

R. Cider, four pounds.
Senna, four ounces and a half.
Digest for a night, express, and strain, then add Sugar, three pounds.
Clarify, and form a syrup. *Sard. Ph.*

SERPENTARIA.

VIRGINIA SNAKEROOT.

Virginia snakeroot is furnished by several species of *Aristolochia*, as the *A. serpentaria*, *A. hirsuta*, and *A. reticulata*, all natives of the United States, especially the western portions.

Sex. Syst. Gynand. hexand. *Nat. Syst.* Aristolochiaceæ.

Willd. Sp. Pl. iv. 159. Griffith, Med. Bot. 529.

The root, which is the officinal portion, is very similar in all the above-mentioned species. It is in tufts of slender, long, matted fibres, attached to a knotty, rugged head. These fibres are brittle, of a yellowish or brownish color, with an aromatic odor, and a pungent, bitter, camphorated taste. It is a stimulating tonic, and may also act as a diaphoretic, and diuretic. The dose of the powder is from ten to thirty grains.

COMPOUND PILLS OF VIRGINIA SNAKEROOT.

R. Powdered Virginia snakeroot, twenty-four grains.
Camphor, forty-eight grains.
Conserve of roses, sufficient.
Mix, and make eighteen pills; to be taken during the apyrexia, in malignant intermittents. *Foy.*

ELECTUARY OF VIRGINIA SNAKEROOT.

R. Powdered Virginia snakeroot,
" contrayerva, each, one drachm.

Aromatic confection, one drachm.
Preserved ginger, six drachms.
Syrup of parsley, sufficient.
Make electuary. Half a drachm, every four hours, as a febrifuge. *Cadet.*

FLUID EXTRACT OF VIRGINIA SNAKEROOT.

R. Virginia snakeroot,
Sugar, each, in powder, eight ounces
Water,
Alcohol, each, sufficient.

Macerate the snakeroot for a day or two in a pint of alcohol, introduce into a percolator, and add diluted alcohol until four pints of tincture pass. Evaporate the tincture to twelve fl. ounces, dissolve the sugar in it, and strain through flannel.

Dose, half a fl. drachm, equivalent to fifteen grains of the root. *J. C. Savery.*

R. Virginia snakeroot, bruised, twelve ounces.
Water,
Alcohol, each, sufficient.

Mix the snakeroot with twelve fl. ounces of alcohol, macerate for twenty-four hours, transfer to a percolator, and add alcohol until a pint and a half of tincture pass. Evaporate spontaneously to six fl. ounces.

Displace the root with water until three pints of infusion have passed; evaporate on a water-bath to six fl. ounces. Finally, mix the tincture and infusion, and filter.

Each fl. ounce represents one ounce of the root. *A. B. Taylor.*

INFUSION OF VIRGINIA SNAKEROOT.

R. Virginia snakeroot, half an ounce.
Boiling water, one pint.

Macerate for two hours in a covered vessel, and strain. *U. S. Ph.*

Dose, one to two fl. ounces, every two hours, in low forms of fever.

COMPOUND INFUSION OF VIRGINIA SNAKEROOT.

R. Virginia snakeroot,
Contrayerva, each, five drachms.
Boiling water, one pint.

Macerate for two hours, strain, and add

Tincture of Virginia snakeroot, two fl. ounces.

Mix. Dose, a tablespoonful. *Beasley.*

TINCTURE OF VIRGINIA SNAKEROOT.

R. Virginia snakeroot, bruised, three ounces.
Diluted alcohol, two pints.

Macerate for fourteen days, express, and filter.
Dose, one to two fl. drachms. *U. S. Ph.*

MIXTURE OF VIRGINIA SNAKEROOT
AND ALLSPICE.

R. Comp. infus. of Virginia
snakeroot, twelve fl. ounces.
Tincture of allspice, four fl. ounces.

Mix. Four spoonfuls, every six hours, in retro-
cedent and atonic exanthemata.

Saunders.

ACETATED MIXTURE OF VIRGINIA
SNAKEROOT AND ALLSPICE.

R. Add vinegar, two fl. ounces, to sixteen
fl. ounces of last-mentioned mixture.

Dose, as above, in petechial typhus. *Swediaur.*

INFUSION OF VIRGINIA SNAKEROOT
AND ETHER.

R. Virginia snakeroot, six drachms.
Boiling water, eight fl. ounces.

Infuse, and strain; when cold, add
Sulphuric ether, two fl. drachms.

A tablespoonful, every hour. *Richter.*

WINE OF VIRGINIA SNAKEROOT AND
VANILLA.

R. Virginia snakeroot, six drachms.
Vanilla, two drachms.
Sherry wine, one pint.

Macerate for four hours, and add to the strained
liquid,

Camphor, half a drachm.
Acetic ether, one fl. drachm.
Syrup of cinnamon, half a fl. ounce.

Dose, two spoonfuls an hour, in low states of the
system. *Vogt.*

TINCTURE OF VIRGINIA SNAKEROOT
AND BALSAM OF PERU.

R. Virginia snakeroot, six drachms.
Proof spirit, five fl. ounces.
Water, sufficient

to obtain six fl. ounces of strained fluid; after
one hour of infusion, add

Camphor, half a drachm.
Balsam of Peru, one drachm.
Gum Arabic, two drachms.

Make an emulsion, and add
Acetic ether, one fl. drachm.

A spoonful, every hour, in cases requiring sti-
mulation. *Phæbus.*

SESAMUM.

BENNE.

Two species of Sesamum, the *S. orientale*,
and the *S. Indicum*, afford leaves and seeds that
are recognized as officinal. They are both na-
tives of Asia, and are cultivated in the West
Indies and in the southern parts of the United
States.

Sex. Syst. Didyn. angios. *Nat. Syst.* Pedu-
liaceæ.

Linn. Sp. Pl. 883, 884. *Griffith, Med. Bot.*
515.

The seeds are small, inodorous, and afford a
bland oil, which may be used for all the pur-
poses of olive oil. The leaves abound in a
gummy substance which forms, with water, a
rich, bland mucilage, useful in all cases requir-
ing the use of demulcents.

INFUSION OF BENNE.

R. Fresh leaves of benne, a handful.
Cool water, one pint.

Infuse till a mucilage is formed. When dried
leaves are used, the water should be hot.

Wood.

OIL OF BENNE.

This is obtained from the seeds by expression.
It is inodorous, of a sweet, bland taste, and
keeps well without becoming rancid. It is
used as an application to promote softness of
the skin. *Redwood*

SIMARUBA.

SIMARUBA.

This is the bark of the root of *Simaruba offi-
cinalis*, a large tree, native of Jamaica, and
several parts of South America.

Sex. Syst. Decand. monog. *Nat. Syst.* Sima-
rubaceæ.

De Cand. Prod. i. 733. *Griffith, Med. Bot.*
198.

The bark is in long pieces, much rolled or
quilled. It is fibrous, tough; of a greyish color
externally, and yellowish within. It is inodor-
ous, but extremely bitter. It has the properties
of the pure bitters, but is apt to act on the sto-
mach and bowels.

INFUSION OF SIMARUBA.

R. Simaruba bark, bruised,
three drachms

Boiling water, one pint.

Macerate for two hours, and strain. *Ed. Ph.*

Dose, two fl. ounces.

COMPOUND INFUSION OF SIMARUBA.

R. Simaruba,
Wormwood, each, two drachms.
Boiling water, one pint.

Infuse for fifteen minutes ; strain, and add
 Syrup of gentian, one fl. ounce.
 To be taken in wineglassful doses. *Foy.*

SINAPIS.

MUSTARD.

The seeds of two species of *Sinapis* are used in medicine, those of *S. nigra* and *S. alba*, both annual plants indigenous to many parts of Europe, and cultivated in our gardens.

Sex. Syst. Tetrady. siliq. *Nat. Syst.* Brassicaceæ.

Linn. Sp. Pl. 933. *Griffith, Med. Bot.* 133.

Black mustard seeds are small, globular, of a deep brown color, having a peculiar smell when bruised, which becomes pungent on the flour being mixed with water ; their taste is bitterish and pungent. White mustard seeds are larger, of a yellowish color, and less pungent. These seeds act as a laxative ; when in the state of flour, they operate as an emetic, and when applied to the skin as a rubefacient, or even vesicant.

CATAPLASM OF MUSTARD.

R. Boiling water, ten fl. ounces.
 Flaxseed meal,
 Flour of mustard, each,
 two ounces and a half.

Mix the powders, and gradually add the water, stirring till a cataplasm is formed. *Lond. Ph.*

R. Flour of mustard, two ounces.
 Crumb of bread, four ounces.
 Common salt, half an ounce.
 Vinegar, sufficient.

Mix. *Ammon.*
 Used as rubefacients and revulsives.

COMPOUND CATAPLASM OF MUSTARD.

R. Flour of mustard, eight ounces.
 Powdered white pepper,
 " ginger, each, one drachm.
 Oxymel, sufficient.

Mix. *Foy.*

R. Flour of mustard, four ounces.
 Yeast, one ounce and a half.
 Muriate of ammonia, one drachm.
 Rue, half an ounce.
 Vinegar, sufficient.

Mix. *Span. Ph.*

Used as above.

R. Flour of mustard, four ounces.
 Hot water, sufficient.

Mix. As a rubefacient. *Ellis.*

MUSTARD GARGLE.

R. Flour of mustard, one drachm.
 Water of angelica,
 " hyssop, each, six fl. ounces.
 Spirit of horseradish, half a fl. ounce.
 " muriate of ammonia,
 four fl. scruples.
 Honey of roses, four fl. ounces.
 Mix. *Spielmann.*

MUSTARD FOOT BATH.

R. Flour of mustard, four ounces.
 Hot water, one gallon.
 Mix. *Beral.*

MUSTARD WHEY.

R. Milk, one pint.
 Water, two pounds.
 Bruised mustard-seed,
 one ounce and a half.

Boil till curdled, and strain. *Beasley.*

R. Milk, one pint.
 Bruised mustard seed, one ounce.

Boil till curdled, and strain, then add

Sherry wine, six fl. ounces.

A draught every hour or two, in low states of the system. *Ellis.*

MUSTARD OINTMENT.

R. Flour of mustard, three ounces.
 Oil of almonds, half a fl. ounce.
 Lemon juice, sufficient.

Mix. In sun-burn, &c *Frank.*

COMPOUND MUSTARD OINTMENT.

R. Resin ointment, one ounce.
 Soap, half an ounce.
 Camphor, two scruples.
 Flour of mustard, one scruple.

Mix. *Ferriar.*

LOTION OF MUSTARD.

R. Flour of mustard, one ounce.
 Express the oil, and add to the mass
 Oil of turpentine, eight ounces.

Macerate for three days, express, and strain.
 As a friction in paralysis. *Niemann.*

FIXED OIL OF MUSTARD.

A fixed oil is obtained in Philadelphia from mustard, by subjecting the bruised seed to powerful expression in an hydraulic press.

This oil is used as a substitute for sweet oil, for dressing salad, &c.

VOLATILE OIL OF BLACK MUSTARD.

R. Bruised black mustard seed, twenty pounds.
Cold water, sufficient to cover.
Distil, and separate the oil. *Magendie.*
A very active rubefacient.

TINCTURE OF OIL OF MUSTARD.

R. Volatile oil of mustard, twelve parts.
Alcohol, two hundred and fifty parts.
Dissolve. A good rubefacient. *Soubéiran.*

LINIMENT OF MUSTARD.

R. Lard,
Oil of almonds, each, twelve ounces.
Yellow wax, four ounces.
Melt by a gentle heat, pour into a heated mortar, and when cooling, add
Oil of lavender, three ounces.
Carbonate of ammonia, half an ounce.
Tincture of mustard, two fl. ounces.
Camphor, one ounce.
Triturate the ammonia with the oil of lavender; dissolve the camphor in the tincture, and add both to the melted mass, incorporating well.
As an application to chilblains. *Beral.*

SODIUM.

SODIUM.

SODII BROMIDUM.

BROMIDE OF SODIUM.

R. Solution of bromide of iron, at will.
" carbonate of soda, sufficient
to precipitate; filter, and evaporate. *Magendie.*

OINTMENT OF BROMIDE OF SODIUM.

R. Bromide of sodium, thirty-four grains.
Lard, one ounce.
Mix. As an application to obstinate cutaneous eruptions. *Magendie.*

SODII CHLORIDUM.

SODÆ MURIAS.

COMMON SALT.

POWDER OF COMMON SALT AND COCHINEAL.

R. Powdered common salt, three drachms.
" cochineal, fifteen grains.

Mix, and divide into six powders. Anthelmintic.
One every morning, following the last dose by some purgative. *Rush.*

COMPOUND SALINE POWDER.

R. Common salt, four ounces.
Sulphate of magnesia, each, three ounces.
Sulphate of potassa, three ounces.
Dry the salts separately, and pulverize; rub them well together, and keep in well-closed vessels. *Ed. Ph.*

As a laxative, in doses of two or three drachms, in a half pint of carbonic acid water, early in the morning.

BATH OF COMMON SALT AND GELATINE.

R. Common salt, one pound.
Water, four pints.
Pour the solution in a bath, and add
Flanders glue, two pounds;
dissolved in
Water, six pints.
Recommended in scrofulous affections. *Foy.*

FOMENTATION OF COMMON SALT.

R. Common salt, two ounces.
Water, six fl. ounces.
Vinegar,
Brandy, each, three fl. ounces.
Mix. As a fomentation to bruises. *Vogt.*

CLYSTER WITH COMMON SALT.

R. Common salt, one ounce.
Barley water, half a pint.
Add to the solution
Olive oil, one fl. ounce.
Mix. *Mid. Hosp.*

R. Common salt, one to two ounces.
Tepid flaxseed infusion, one pint.

Dissolve, and add
Castor, or olive oil,
Molasses, each, two fl. ounces.

One-half to be used at a time; the remainder in half an hour afterwards, if needed. *Ellis.*

CLYSTER OF COMMON SALT AND ARNICA.

R. Common salt, one ounce.
Infusion of arnica, twelve fl. ounces.
Mix. Said to be useful in apoplexy and paralysis. *Foy*

MIXTURE OF COMMON SALT AND LEMON JUICE.

R. Lemon juice, at will.
Common salt, sufficient
to saturate; filter. A tablespoonful every hour,
in apyrexia of intermittents. *Bories.*

OINTMENT OF COMMON SALT.

R. Common salt, one drachm.
Water, sufficient to dissolve.
Lard, one ounce.

Rub together till smooth. The strength to be gradually increased from one drachm of the salt to four. To inflamed eyelids. *Tavignot.*

COMPOUND OINTMENT OF COMMON SALT.

R. Common salt, one ounce and a half.
Water, sufficient to dissolve.
Simple cerate, three ounces.
Rose-water ointment, one ounce.

Rub together. In tinea capitis, as a friction, morning and evening. *St. Marie.*

R. Common salt, two drachms.
Nut oil, one drachm.
Ox gall, one ounce and a half.

Digest together for thirty-six hours, and triturate well. As a friction to scrofulous tumors. *Roncalli.*

SODII SULPHURETUM.

SULPHURET OF SODIUM.

R. Sulphur, two hundred parts.
Carbonate of potassa, dried,
two hundred and sixty-seven parts.

Mix; heat in a matrass till melted; on cooling, detach, and keep in a well-closed vessel. *Cottureau.*

BATH WITH SULPHURET OF SODIUM.

R. Sulphuret of sodium, ten ounces.
Solution of common
salt and gelatine,
(see page 431,) four fl. ounces.

Mix with the water of the bath, at the moment the patient enters it. In chronic cutaneous affections. *Cadet.*

MIXTURE OF SULPHURET OF SODIUM AND SAL AMMONIAC.

R. Sulphuret of sodium, one drachm.
Muriate of ammonia, fifty grains.

Dissolve each in six ounces of water, mix the solutions, after having filtered them. As a lotion in itch, &c. *Van Mons.*

LOTION WITH SULPHURET OF SODIUM.

R. Sulphuret of sodium, three ounces.
Soap, one ounce and a half.

Dissolve in

Alcohol, one pint.

Add to the solution

Lime water, eight pints.

In tinea capitis; the head to be washed with it every day, without removing the hair.

LINIMENT OF SULPHURET OF SODIUM.

R. Sulphuret of
sodium, four parts and a half.
Sulphur,
Oil of almonds, each,

one part and a half.
Lard, six parts and a half.

Rub the powders with the lard, and gradually add the oil. As a lotion in chronic cutaneous diseases. *Van Mons.*

OINTMENT OF SULPHURET OF SODIUM.

R. Sulphuret of sodium, three drachms.
Lard, one ounce and a half.

Rub well together. As an application in itch. *Swedliaur.*

SODA.

CAUSTIC SODA.

R. Solution of soda, at will.

Evaporate in a silver capsule, till it will solidify, on cooling; pour into moulds. *Guibourt.*

SOLUTION OF SODA.

R. Crystallized carbonate
of soda, twenty-five parts.
Quicklime, six parts.

Boil with a sufficient quantity of water; strain, and evaporate, till it marks 36° of the areometer. *Guibourt.*

R. Carbonate of soda, thirty-one ounces.
Lime, nine ounces.
Boiling water, (Imp.) one gallon.

Dissolve the carbonate in half a gallon of the water; sprinkle a little water on the lime in an earthen vessel, and when it is slaked, add the remainder of the water. Mix the two solutions, and shake them frequently until cold. Allow the carbonate of lime to subside; pour off the supernatant liquid, (which should have the *sp. gr.* 1.061,) and keep in well-stopped green-glass bottles. *Lond. Ph.*

SODÆ ACETAS.

ACETATE OF SODA.

R. Carbonate of soda, at will.
Distilled vinegar, sufficient
to saturate; evaporate to *sp. gr.* 1.276, and set
aside to crystallize. *Dub. Ph.*, 1826.

At the present time, the acetate of soda is
chiefly prepared by the manufacturer of crude
pyroligneous acid, in the process of obtaining
the officinal acetic acid.

Diuretic or cathartic, according to dose,
which varies from one scruple to four drachms.

COMPOUND PILLS OF ACETATE OF
SODA.

R. Acetate of soda, five drachms.
Camphor,
Nitrate of potassa, each, two drachms.
Conserve of elder berries, sufficient.
Mix, and make one hundred and twenty pills.
Dose, two, morning and evening, to check the
secretion of milk. *Bories.*

R. Acetate of soda,
Powdered rhubarb,
Inspissated ox gall, each, one drachm.
Powdered gum Arabic, half a drachm.
Mix, and make pills of two grains. Dose, ten
to fifteen, twice a-day, as a laxative. *Augustin.*

SODÆ ARSENIAS.

ARSENIATE OF SODA.

R. White arsenic, one hundred parts.
Nitrate of
soda, one hundred and sixteen parts.

Pulverize, and introduce into a luted retort;
heat in a reverberatory furnace, break the re-
tort, dissolve the salt in water, add carbonate
of soda, till the solution is neutral, or slightly
alkaline; filter, evaporate, and crystallize.

In intermittents, and some diseases of the
skin. Dose, one-sixteenth to one-eighth of a
grain. *Par. Cod.*

SOLUTION OF ARSENIATE OF SODA.

(HEINECKE'S SOLUTION.)

R. Arseniate of soda, six grains.
Peppermint
water, two fl. ounces and a half.

Add to the solution

Wine of cinnamon, half a fl. ounce.
Laudanum, one fl. drachm.

Dose, forty to fifty drops, four times a-day.

Heinecke.

PEARSON'S SOLUTION.

R. Crystallized arseniate of soda, one grain.
Distilled water, one fl. ounce.
Dissolve. Dose, twenty drops. *Pearson.*

SODÆ BORAS.

BORAX.

Borax is found native in several parts of
Europe, and in Peru. But the chief supplies
of it are obtained from the shores of certain
lakes in Persia and Thibet.

It is a mild diuretic. The dose is about
thirty grains.

COMPOUND POWDER OF BORAX.

R. Powdered borax, one drachm.
" saffron, ten grains.
Oil of cinnamon, two drops.
Mix, for two doses. As an emmenagogue. *Wedel.*

R. Powdered borax, half an ounce.
" castor,
" saffron, each, one drachm and a half.

Oil of cinnamon, eight drops.
" amber, six drops.

Mix, and form a powder. Dose, from a scruple
to half a drachm, to facilitate parturition, expel
placenta, &c. *Wirt. Ph.*

LOZENGES OF BORAX.

R. Powdered borax, two drachms.
" sugar, half an ounce.
Mucilage, sufficient.
Mix, and make thirty lozenges. One, occa-
sionally, in aphthous sore mouth. *Phabus.*

COLLUTORY OF BORAX.

R. Borax, two drachms.
Water,
Tincture of myrrh, each, one fl. ounce.
Honey of roses, two ounces.
Mix. *Swediaur.*

R. Borax, three drachms.
Mucilage of quince
seeds, eight fl. ounces.
Honey of roses, two ounces.
Mix. *Bahi.*

R. Borax, three drachms.
Infusion of roses,
one fl. ounce and a half.
Honey of roses, two ounces.
Mix. *Pringle*

COLLYRIUM OF BORAX.

R. Borax, one drachm.
 Laudanum, half fl. drachm.
 Rose water, four fl. ounces.

Mix. *Fricke.*

R. Borax, half a drachm.
 Sugar, one drachm.
 Rose water, two fl. ounces.

Mix. *Richard.*

GARGLE OF BORAX.

R. Borax, two drachms.
 Oxymer, half fl. ounce.
 Water, eleven fl. ounces.

Mix. *Beasley.*

R. Borax, one drachm.
 Tincture of myrrh, half fl. ounce.
 Clarified honey, one fl. ounce.
 Rose water, four fl. ounces.

Mix. *Ellis.*

In aphthous sore mouth, ptyalism, &c.

LOTION OF BORAX.

R. Borax, half a drachm.
 Orange-flower water,
 Rose water, each, half fl. ounce.

Dissolve. In pruritus. *Cadet.*

R. Borax, half an ounce.
 Sulphate of morphia, six grains.
 Rose water, eight fl. ounces.

Mix. Used as a lotion in pruritus vulvæ.

C. D. Meigs.

HONEY OF BORAX.

R. Powdered borax, one drachm.
 Clarified honey, one ounce.

Mix. *Dub. Ph.*

In aphtha.

VINEGAR OF BORAX.

R. Distilled vinegar, two fl. ounces.
 Borax, one drachm.

Dissolve. As lotion to ringworm of the scalp.
Christison.

MIXTURE OF BORAX.

R. Borax, two drachms.
 Chamomile water, four fl. ounces.
 Syrup of orange peel, half fl. ounce.
 Liquid succinate of ammonia,

two fl. drachms.

Wine of opium, half a fl. drachm.

Mix. As an emmenagogue, and to facilitate parturition. Dose, two spoonfuls, every half hour. *Swediaur.*

OINTMENT OF BORAX.

R. Powdered borax, fifteen grains.
 Fresh butter, two drachms.

Rub together. Applied to scaly eruptions.
Radius.

R. Powdered borax, one drachm.
 Lard, one ounce.

Rub together. An excellent application to painful hemorrhoidal tumors, and to cracked nipples.
Pereira.

R. Powdered borax, two drachms.
 Rose ointment, one ounce.

Mix. For chilblains. *Cadet.*

SODÆ CARBONAS.

CARBONATE OF SODA.

DRIED CARBONATE OF SODA.

R. Carbonate of soda, at will.

Heat in a clean iron vessel, till perfectly dried, stirring constantly; rub into powder. *U. S. Ph.*

Dose, three to ten grains.

SOLUTION OF CARBONATE OF SODA.

R. Carbonate of soda, one ounce and a half.

Distilled water, (Imp.) one pint.

Dissolve. The *sp. gr.* is 1.026. *Dub. Ph.*

Dose, half a fl. ounce to one ounce diluted, two or three times a-day, as antacid, &c.

DIURETIC SOLUTION WITH CARBONATE OF SODA.

R. Carbonate of soda, three drachms.
 Nitre, one drachm and a half.

Honey, one ounce and a half.

Decoction of couchgrass, three pints.

Mix. A glassful every hour, in dropsy.

Cadet.

MIXTURE OF CARBONATE OF SODA AND IPECACUANHA.

R. Carbonate of soda, twelve grains.
 Wine of ipecacuanha, twenty drops.

Laudanum, four drops.

Distilled water, one fl. ounce.

Mix. A teaspoonful, every two or three hours, in hooping-cough of children. *Ellis.*

MIXTURE OF CARBONATE OF SODA AND CHAMOMILE.

R. Carbonate of soda, one drachm.
 Chamomile water, three fl. ounces.

Syrup of gentian, one fl. ounce.
Tincture of Peruvian bark,
one fl. drachm.
Mix. A spoonful, occasionally, in scrofula.
Foy.

PILLS OF SODA AND RHUBARB.

R. Powdered rhubarb,
Dried carbonate of soda,
Extract of gentian, each, one scruple.
Calomel, three grains.
Mix, and make twenty pills. Two, occasionally,
in dyspepsia. *Ellis.*

PILLS OF SODA AND IPECACUANHA.

R. Carbonate of soda, one drachm.
Bitter almonds, half a drachm.
Ipecacuanha, three grains.
Extract of madder,
twenty-four grains.
Water, sufficient.
Beat into mass, and form sixty pills. Three,
morning and evening, in obstinate cutaneous
diseases. *Foy.*

CARBONATE OF SODA PILLS.

R. Extract of chamomile, one drachm.
Powdered rhubarb, one scruple.
Carbonate of soda, half a drachm.
Oil of caraway, ten minims.
Syrup of ginger, sufficient.
Mix, and make twenty-four pills, two to be
taken thrice a-day, in dyspepsia. *Ainslie.*

PILLS OF CARBONATE OF SODA.

R. Dried carbonate of soda, one drachm.
Soap, two scruples.
Water, sufficient.
Beat together, and make thirty pills. Three,
thrice a-day. In calculous affections. *Ellis.*

MIXTURE OF CARBONATE OF SODA AND QUASSIA.

R. Carbonate of soda,
two drachms and a half.
Rasped quassia,
one drachm and a half.
Boiling water, one pint.
Infuse for one hour, and strain. Two to four
spoonfuls a-day, in dyspepsia. *Augustin.*

MIXTURE OF CARBONATE OF SODA AND GENTIAN.

R. Carbonate of soda, four scruples.
Compound infusion of gentian,
three fl. ounces.

Cinnamon water, three fl. ounces.
Tincture of cardamom, half a fl. ounce.
Mix. A tablespoonful, every two or three hours,
in acidity and flatulence. *Ellis.*

LOTION OF CARBONATE OF SODA.

R. Saturated solution of carbonate
of soda, one ounce.
Warm water, two pints.
Mix. As a lotion in pruritus. *Radius.*

INJECTION OF CARBONATE OF SODA.

R. Carbonate of soda, one drachm.
Soap, two ounces.
Water, twelve fl. ounces.
Advised as an injection, in cases of uric acid
calculi. *Bories.*

POWDER OF CARBONATE OF SODA AND RHUBARB.

R. Dried carbonate of soda, one ounce.
Rhubarb, four drachms.
Ginger, one scruple.
Mix. In cardialgia, and dyspepsia. *Beasley.*
Dose, ten to twenty grains.

POWDER OF CARBONATE OF SODA AND MERCURY.

R. Dried carbonate of soda, five drachms.
Calomel, one drachm.
Compound chalk powder, ten drachms.
Mix. Dose, eight to sixteen grains. *Guy's Hosp.*

OINTMENT OF CARBONATE OF SODA.

R. Carbonate of soda, two drachms.
Wine of opium, one fl. drachm.
Lard, one ounce.
Mix. In cutaneous affections attended with
pain. *Soubéiran.*

R. Carbonate of soda,
Sulphuret of
sodium, each, one drachm.
Lard, one ounce.

Mix. In tinea capitis, having removed the
crusts. *Alibert.*

R. Impure carbonate of soda,
three drachms.
Slaked lime, two drachms.
Lard, two ounces.
Mix. As the last. *Cadet.*

R. Carbonate of soda, two drachms.
Slaked lime, one drachm.
Opium, two grains.
Lard, two ounces.
Mix. In prurigo. *Biett.*

SODÆ BICARBONAS.

BICARBONATE OF SODA.

R. Crystallized carbonate of soda, at will.
Saturate in a proper apparatus, with carbonic acid gas. *U. S. Ph.*

R. Carbonate of soda, one part.
Water, two parts.

Dissolve, and pass a current of carbonic acid through the solution; let crystallize. *Cottreueau.*

EFFERVESCING SOLUTION OF SODA.

R. Bicarbonate of soda, one drachm.
Water, one pint.

Dissolve, and saturate it with carbonic acid.
Keep in a well-closed vessel. *Lond. Ph.*

EMULSION WITH BICARBONATE OF SODA.

R. Bicarbonate of soda, sixteen grains.
Nitro, four grains.
Syrup of marsh mallow,
" violets, each, one fl. ounce.
Lettuce water, two fl. ounces.
Emulsion of almonds,
twelve fl. ounces.

Mix. In nephritic complaints. *Beral.*

EFFERVESCING POWDERS OF SODA.

R. Bicarbonate of soda,
one ounce, fifty-four grains.
Tartaric acid, one ounce.

Powder each, separately, and divide each into sixteen powders, and preserve in papers of different colors. *Ed. Ph.*

Known under the name of *Soda powders.*

R. Bicarbonate of soda, twenty-six grains.
Powdered ginger, five grains.
White sugar, two drachms.
Essence of lemon, one drop.

Mix, and put in a white paper.

R. Tartaric acid, half a drachm.
Put in a blue paper. When used, dissolve the contents of each paper in four fl. ounces of water, mix the solutions, and drink in a state of effervescence.

Known as the *Ginger beer powders.* *Pereira.*

POWDER OF BICARBONATE OF SODA AND MAGNESIA.

R. Magnesia, half an ounce.
Bicarbonate of soda,
Powdered ginger, each, one scruple.

Mix. A small teaspoonful, occasionally. *Ellis.*

ANTI-EMETIC MIXTURE.

R. Bicarbonate of
soda, one drachm and a half.
Powdered gum Arabic, two drachms.
Oil of spearmint, four drops.
White sugar, two drachms.
Carbonic acid water, four fl. ounces.

Mix. A tablespoonful occasionally, to calm the stomach, or prevent vomiting. *Dewees.*

LOZENGES OF BICARBONATE OF SODA.

R. Bicarbonate of soda, one ounce.
Powdered gum Arabic, half an ounce.
" sugar, three ounces.
Mucilage, sufficient.

Beat together, and form lozenges. *Ed. Ph.*
Antacid and antilithic.

R. Bicarbonate of soda, two ounces.
Sugar, thirty-four ounces.
Oil of peppermint, eighteen drops.
Mucilage, thirty-six drachms.

Mix, and make lozenges of eighteen grains. *Magendie.*

R. Bicarbonate of soda, four parts.
Sugar, sixty parts.
Chocolate, eight parts.
Mucilage, nine parts.

Make lozenges of eighteen grains. *Beral.*

R. Bicarbonate of soda, four drachms.
Sugar, ten ounces.
Balsam of tolu, one drachm.
Mucilage of tragacanth, sufficient.

Make lozenges of twenty grains. *Cadet.*

MIXTURE OF BICARBONATE OF SODA AND COPAIBA.

R. Bicarbonate of soda, one drachm.
Almond emulsion, four fl. ounces.
Copaiba, two fl. drachms.
Laudanum, sixty drops.

Mix. A tablespoonful occasionally, in acute pain in the bladder. *Ellis.*

SODÆ CHLORAS.

CHLORATE OF SODA.

R. Tartaric acid, five ounces.
Boiling water, two pints.

Dissolve, and add
Carbonate of soda,
forty-five drachms and a half.

Also, dissolve

Chlorate of potassa,
four ounces and fifteen grains.
Boiling water, sixteen fl. ounces.

Mix the solutions whilst boiling, filter, evaporate, and crystallize. *Hamb. Ph.*

LOTION OF CHLORATE OF SODA.

R. Chlorate of soda, two drachms.
Water, four fl. ounces.
In pruritus. *Darling.*

GARGLE OF CHLORATE OF SODA.

R. Chlorate of soda,
one to three scruples.
Barley water, three ounces.
Honey of roses, one ounce.
Mix. *Radius.*

SODA CHLORINATA.

CHLORINATED SODA.

R. Dried carbonate of soda,
nineteen parts.
Water, one part.

Mix and place in a suitable vessel, and pass in chlorine gas to saturation. *Christison.*

This article is more generally used in the liquid state, prepared in the following manner:

SOLUTION OF CHLORINATED SODA.

R. Chlorinated lime, one pound.
Carbonate of soda, two pounds.
Water, one gallon and a half.

Dissolve the carbonate of soda in three pints of the water, with the aid of heat. Add to the remainder of the water the chlorinated lime, in powder, in small portions at a time. Set aside for some hours, to settle; decant clear solution, and add that of the carbonate of soda; decant, and strain through linen, and keep in opaque bottles. *U. S. Ph.*

R. Carbonate of soda, one part.
Water, ten parts.
Dissolve, and pass a current of chlorine through the solution. *Guibourt.*

This is known as *Labarraque's Disinfecting Solution*; it is chiefly employed as a disinfectant, but is also used internally, in doses of thirty drops to one fl. drachm, in low fevers, &c.

DILUTED SOLUTION OF CHLORINATED SODA.

R. Solution of chlorinated soda,
half a fl. drachm.
Distilled water, four fl. ounces.
To be taken in spoonful doses, in the apyrexia of intermittents. *Gouze.*

R. Solution of chlorinated soda, one part.
Water, twelve to sixteen parts.
As a lotion to foul and cancerous ulcers. *Foy.*

GARGLE OF CHLORINATED SODA.

R. Solution of chlorinated soda,
one fl. drachm.
Hydromel, one fl. ounce.
Distilled water, seven fl. ounces.
Mix. *Beral.*

INJECTION OF CHLORINATED SODA.

R. Solution of chlorinated soda,
one fl. ounce.
Water, sixteen fl. ounces.
Mix. Used as a vaginal injection in fetid discharges from malignant diseases. *Pereira.*

CATAPLASM OF CHLORINATED SODA.

R. Boiling water, six fl. ounces.
Flaxseed meal, four ounces and a half.
Solution of chlorinated soda,
two fl. ounces.

Add the meal gradually to the water, constantly stirring; then mix in the chlorinated soda.

Applied to foul and sloughing ulcers.

Lond. Ph.

SODÆ HYPOSULPHIS.

HYPOSULPHITE OF SODA.

R. Carbonate of soda, eight ounces.
Distilled water, one pint.

Dissolve, and mix with the solution

Sulphur, one ounce.

Then pass an excess of sulphurous acid gas into the liquid; boil for a few minutes in a matrass; filter, evaporate to one-third, and set aside to crystallize. *Par. Cod.*

Used in diseases of the skin, in doses of ten grains to one drachm.

R. Carbonate of soda,
seven hundred and thirty parts.
Sulphur, forty-five parts.
Water, fifteen hundred parts.

Mix the sulphur with a small portion of the water, and add the soda, dissolved in the rest. Introduce into two two-necked bottles, so as to fill them about two-thirds.

R. Clean iron filings,
fifteen hundred parts.
Sulphuric acid (1.84),
three thousand parts

Put into a flask, let cool, place on a sand-bath, and convey the sulphurous acid gas, as it is generated, by rather large tubes, first into a

washing bottle, and then through the above solution in the two-necked bottles. Continue the process for ten or twelve hours, mix the contents of the two bottles, filter, and evaporate to crystallization.

Legrip.

R. Dried carbonate of soda, in powder, one pound.
Sulphur, five ounces.

Mix and heat in a porcelain vessel until the sulphur melts; stirring the mass to bring every portion in contact with the air. Dissolve the sulphate of soda, thus formed, in water; filter, boil the solution with sulphur; filter, evaporate, and crystallize.

Walchner.

Principally used in daguerreotyping.

SYRUP OF HYPOSULPHITE OF SODA.

R. Hyposulphite of soda, one ounce.
Water, twelve fl. ounces.
Sugar, twenty-three ounces.

Dissolve the salt in the water, add the sugar, and form syrup. Dose, from one to two ounces, twice a-day.

Mouchon.

BATH OF HYPOSULPHITE OF SODA.

R. Hyposulphite of soda, one to four ounces.
Water, one gallon.

Dissolve, and mix with the water of a bath; adding vinegar while the patient is in the bath, to liberate the sulphurous acid and sulphur.

Pereira.

SODÆ NITRAS.

NITRATE OF SODA—CUBIC NITRE.

This salt is found in immense beds in Peru, whence it is imported for the manufacture of nitric acid, chrome yellow, sulphuric acid, &c.

It crystallizes in rhomboidal prisms; its taste is sharp, cooling, and bitter.

It is sometimes employed by the pyrotechnist, but it is seldom used in medicine.

SODÆ PHOSPHAS.

PHOSPHATE OF SODA.

R. Powdered burnt bones, ten pounds.
Sulphuric acid, six pounds.
Carbonate of soda, sufficient.

Mix the powdered bone with the acid in an earthen vessel, add a gallon of water, and stir. Digest for three days, occasionally adding a little water, then pour in a gallon of boiling water, and strain; adding boiling water till the liquid passes almost tasteless. Let settle, decant, and boil down to a gallon. After settling, pour

it into an iron vessel, heat, and gradually add the carbonate of soda dissolved in hot water, as long as there is effervescence; then filter, and let crystallize. More crystals will be obtained by adding carbonate of soda in excess, and again evaporating to crystallization. Keep in a well-stopped bottle.

U. S. Ph.

A mild purgative, in doses of four drachms to one ounce.

R. Acid phosphate of lime, at will.
Add carbonate of soda to saturation; let settle, decant, evaporate, and crystallize.

Guibourt.

COMPOUND POWDER OF PHOSPHATE OF SODA.

R. Phosphate of soda, two drachms.
Carbonate of soda, one drachm.
Fennel sugar, half an ounce.

Mix. Two to three teaspoonfuls a-day, to aid digestion.

Berends.

MIXTURE OF PHOSPHATE OF SODA.

R. Phosphate of soda, half an ounce.
Decoction of carrageen, six fl. ounces.
Syrup of orgeat, half an ounce.

Mix. As a purgative.

Radius.

R. Phosphate of soda, half an ounce.
Syrup of marsh mallow, each, half an ounce.
Water, four fl. ounces.

Mix. As a purgative.

Augustin.

SODÆ SULPHAS.

SULPHATE OF SODA.

[GLAUBER'S SALT.]

POWDER OF SULPHATE OF SODA AND TARTAR EMETIC.

R. Sulphate of soda, two ounces.
Tartar emetic, one grain.
Opium, each, one grain.
Gum Arabic, one grain.
Powdered liquorice, one drachm.

Mix. To be taken in water, in divided doses, during the day, as a purgative and diuretic.

Swedjaur.

POWDER OF SULPHATE OF SODA AND NITRE.

R. Dried sulphate of soda, eighteen drachms.
Nitrate, half a drachm.
Tartar emetic, one grain.

Mix. One-third, as a dose, in water or broth.

Beasley

POWDER OF SULPHATE OF SODA AND
OPIUM.

R. Sulphate of soda, four scruples.
Powdered opium, two grains.
Mix. In hemorrhages and inflammations, after
bleeding. *Radiis.*

COMPOUND PILLS OF SULPHATE OF
SODA.

R. Sulphate of soda, half a drachm.
Bitter almonds, one drachm.
Ipecacuanha, two grains.
Extract of madder, sufficient.
Beat into mass, and make sixty pills. Three,
morning and evening, in obstinate cutaneous
affections. *Foy.*

ELECTUARY WITH SULPHATE OF SODA.

R. Dried sulphate of soda, half an ounce.
Pulp of tamarinds, one ounce.
Syrup of lemon juice, sufficient.
Mix. Two spoonfuls every hour, till it operates.
Phæbus.

COMPOUND SOLUTION OF SULPHATE OF
SODA.

R. Sulphate of soda, fourteen drachms.
Acetate of potassa, twenty-four grains.
Nitro, eighteen grains.
Tartar emetic, quarter of a grain.
Water, two pints.
Dissolve, and filter; in wineglassfuls, in habitual
constipation. *Soubeiran.*

EMULSION WITH SULPHATE OF SODA.

R. Sulphate of soda, one ounce.
Oil of almonds, one fl. ounce.
Yolks of eggs, two.
Syrup of manna, two fl. ounces.
Fennel water, six fl. ounces.
Mix. Dose, two spoonfuls. *Phæbus.*

LEMONADE WITH SULPHATE OF SODA.

R. Sulphate of soda, one ounce and a half.
Lemon juice, half an ounce.
Sugar, three ounces.
Water, sixteen fl. ounces.
Mix. *Radiis.*

CLYSTER OF SULPHATE OF SODA.

R. Sulphate of soda, one ounce.
Barley water, six fl. ounces.
Dissolve, and add
Oil of flaxseed, two ounces.
Swedjaur.

LOTION OF SULPHATE OF SODA.

R. Sulphate of soda, six ounces.
Soap, three drachms.
Lime water, eighteen fl. ounces.
Rectified spirit, six fl. drachms.
Mix. In chronic cutaneous affections.
Van Mons.

SUPPOSITORY OF SULPHATE OF SODA.

R. Sulphate of soda,
Soap, each, half an ounce.
Honey, sufficient.
Mix, and make four suppositories. *Phæbus.*

SODÆ BISULPHAS.

BISULPHATE OF SODA.

R. Dried sulphate of soda, ten parts.
Sulphuric acid, seven parts.
Heat gently in a crucible.
Purgative in doses of two to six drachms.
Beasley

BISULPHATE OF SODA AND MAGNESIA.

R. Bisulphate of soda, at will.
Hot water, sufficient.
Dissolve, add magnesia to saturation; filter, and
crystallize.
Laxative and diuretic, in doses from half a
drachm to two drachms, several times a-day.
Van Mons.

SODÆ ET POTASSÆ TARTRAS.

TARTARIZED SODA. ROCHELLE SALT.

R. Carbonate of soda, one pound.
Cream of tartar, sixteen ounces.
Boiling water, five pints.
Dissolve the carbonate of soda in the water, and
add gradually the cream of tartar; filter, evapo-
rate to a pellicle, and set aside to crystallize.
Decant, and dry crystals on bibulous paper.
U. S. Ph.

Mild purgative, in doses of half an ounce to
an ounce.

POWDER OF TARTARIZED SODA AND
RHUBARB.

R. Tartarized soda, ten grains.
Powdered rhubarb,
three to seven grains.
Mix. To be taken every morning for two weeks,
in enlargement of the mesentery, in children.
Fordyce.

SEIDLITZ POWDERS.

- R. Tartarized soda, two drachms.
Bicarbonate of soda, two scruples.

Mix, and dissolve in half a tumblerful of water,
Powdered tartaric acid, twenty-five grains.

Dissolve in the same quantity of water, as above, and mix.

To be taken during effervescence. *Ellis.*

MIXTURE OF TARTARIZED SODA.

- R. Tartarized soda, half an ounce.
Sulphate of potassa, two drachms.
Infusion of succory, one pint.
Oxymel of squill, four fl. ounces.

Mix. Four spoonfuls, in the morning, with an hour's interval. In visceral obstructions.

St. Marie.

- R. Tartarized soda, one ounce.
Extract of henbane, six grains.
Syrup of chamomile, one fl. ounce.
Water, six fl. ounces.

Mix. A spoonful every hour, in constipation.

Phæbus.

WHEY WITH TARTARIZED SODA.

- R. Root and leaves of dandelion,
Fumitory,
Water-cress,
Chervil, each, a handful.
Clarified whey, four pints.

Boil for five minutes, express, strain, and add
Tartarized soda, two drachms.
Honey, six drachms.

To be taken every morning, early, in four doses, with the interval of an hour. Much praised in visceral obstructions. *St. Marie.*

CLYSTER WITH TARTARIZED SODA.

- R. Tartarized soda,
Extract of dandelion, each,
five drachms.

Infusion of elder flowers, five fl. ounces.

Dissolve. *Radius.*

SODÆ VALERIANAS.

VALERIANATE OF SODA.

- R. Bichromate of potassa,
in powder, (avoir.) nine ounces.
Fusel oil, (Imp.) four fl. ounces.
Sulphuric acid,
(Imp.) six fl. ounces and a half.
Water, (Imp.) half a gallon.
Solution of caustic soda,
(Imp.) one pint.

Dilute the sulphuric acid with ten fl. ounces of water; and dissolve the bichromate in the remainder of the water. When both solutions are cool, place them in a matrass, add the fusel oil, shake well together, until the temperature of the mixture falls to 80° or 90° F. Connect the matrass with a condenser, apply heat, and distil half a gallon. Saturate the distillate with the caustic soda, remove the oil from the surface, and evaporate until the aqueous vapor is given off. Withdraw the heat, and as the valerianate of soda concretes, divide it while warm into fragments, and keep in a well-stopped bottle.

Dub. Ph.

The valerianate of soda is not used in medicine, but it constitutes the source from which all the other valerianates are obtained by double decomposition.

SPIGELIA.

PINKROOT.

This is the root of the *Spigelia marilandica*, a beautiful, herbaceous, perennial plant, a native of the southern and southwestern parts of the United States.

Sex. Syst. Pentand. monog. *Nat. Syst.* Loganiaceæ.

Linn. Sp. Pl. 249. *Griffith, Med. Bot.* 466.

The root consists of numerous small, blackish fibres, forming a dense bunch, and arising from a short rhizome. The odor is faint, and the taste sweetish, bitter, and unpleasant. It is a safe, and tolerably certain anthelmintic, with slight purgative and narcotic properties. It is given in powder, in the dose of one to two drachms for an adult, ten to twenty grains for a child; to be repeated morning and evening, for a few days, and followed by an active purgative.

PINKROOT WITH CALOMEL.

- R. Powdered pinkroot, ten grains.
Calomel, four grains.

Mix. To be taken two mornings in succession; and on the afternoon of the second day, followed by a purgative. For children above four years of age. *Ellis.*

PINKROOT WITH SAVINE.

- R. Powdered pinkroot,
" senna, each, two scruples.
" savine, twelve grains.

Mix, and divide into six powders. One, every morning, for three days, followed by a purgative. *Ellis.*

INFUSION OF PINKROOT.

- R. Pinkroot, half an ounce.
Boiling water, one pint.

Macerate for two hours, in a covered vessel, and strain. *U. S. Ph.*

Four fl. drachms to a fl. ounce, to children; four to eight fl. ounces to adults; morning and evening, followed by a purgative.

COMPOUND INFUSION OF PINKROOT. (WORM TEA.)

- R. Pinkroot, half an ounce.
Senna, two drachms.
Manna, one ounce.
Savine, two scruples.
Fennel seed, two drachms.
Boiling water, one pint.

Infuse. Dose, one tablespoonful for a child, two years old, three times a-day. *Ellis.*

- R. Pinkroot, one ounce.
Bruised rhubarb, one drachm.
Senna, two drachms.
Semen contra, one drachm.
Manna, two drachms.
Coriander, half a drachm.
Boiling water, one quart.

Infuse. A small teacupful, three times a-day. *Hays.*

FLUID EXTRACT OF PINKROOT.

- R. Pinkroot, sixteen ounces.
Diluted alcohol, sufficient.
Sugar, twelve ounces.

Reduce the root to powder, macerate it in twice its weight of the diluted alcohol for twenty-four hours, then displace until four pints of tincture are obtained. Evaporate to ten fl. ounces, add the sugar, heat till dissolved. The whole should measure one pint. *W. Procter.*

COMPOUND FLUID EXTRACT OF PINKROOT.

- R. Bruised pinkroot, four ounces.
Senna, three ounces.
Savine, one drachm.
Manna, one ounce.
Sugar, eight ounces.
Alcohol, half a pint.
Boiling water, two pints.

On the pinkroot, senna, and savine, pour the boiling water, and cover till cool. Then add the alcohol, and macerate for twenty-four hours; transfer to a displacement apparatus, and, having displaced, evaporate the product, at a low heat, to twelve fl. ounces, in which dissolve the manna and sugar, and evaporate to one pint. One fl. ounce contains the strength of two drachms of pinkroot and a drachm and a half of senna. Dose, for a child, a teaspoonful. *T. Estlack.*

- R. Pinkroot, sixteen ounces (av.).
Senna, eight ounces.
Powdered sugar, twenty-four ounces.
Carbonate of potassa, one ounce.

- Oil of caraway,
Oil of anise, each, half a drachm.
Diluted alcohol, sufficient.

Reduce the pinkroot and senna to a coarse powder, mix them with two pints of diluted alcohol, and macerate for two days. Put in a displacer, gradually add more diluted alcohol, until five pints have passed. Evaporate this in a water-bath to twenty fl. ounces, and add the carbonate of potassa. Triturate the oils with a portion of the sugar, then with the remainder; mix with the evaporated fluid, and apply a gentle heat. The whole should measure two pints.

Dose, from half a teaspoonful, to a child of one to two years of age, to a tablespoonful for an adult. *W. Procter.*

FLUID EXTRACT OF PINKROOT AND SENNA.

- R. Pinkroot, in coarse powder, one pound.
Senna, in coarse powder, six ounces.
Sugar, one pound and a half.
Carbonate of potassa, six drachms.
Oil of caraway,
“ anise, each, half a fl. drachm.
Diluted alcohol, sufficient.

Mix the pinkroot and senna with two pints of diluted alcohol, allow the mixture to stand for two days, then transfer to a percolator, and add diluted alcohol until half a gallon of tincture is obtained. Evaporate this on a water-bath, to one pint; then add the carbonate of potassa, and, after the sediment has dissolved, the sugar previously triturated with the oils. Lastly, dissolve the sugar, with a gentle heat.

Dose, one fluidrachm for a child three years old. *U. S. Ph.*

SPIRÆA.

HARDHACK.

The U. S. Pharm. indicates the root of *Spiræa tomentosa* as officinal; but this portion is less active than the leaves and flowers. It is a beautiful shrub, with spikes of rose-colored flowers, and leaves of a dark-green color above, and white beneath.

Sex. Syst. Icosand. pentagyn. *Nat. Syst.* Rosaceæ.

Linn. Sp. Pl. 701. *Griffith, Med. Bot.* 280.

The leaves and flowers are usually sold in packages; they have an odor not unlike that of black tea, and a bitter, very astringent taste. The Hardhack is a tonic and astringent, of considerable powers, and has the advantage of seldom being offensive to the stomach.

DECOCTION OF HARDHACK.

- R. Hardhack, one ounce
Water, one pint and a half

Boil down to one pint. Dose, from one to two fl. ounces.

Wood.

EXTRACT OF HARDHACK.

R. Hardhack, at will.
Water, sufficient.

Exhaust by the process of displacement, and evaporate the product by means of a water-bath to proper consistence. Dose, five grains to a scruple.

Griffith.

SPONGIA.

SPONGE.

Sponge is the horny skeletons of small polymorphous marine animals, found attached to submerged rocks in the seas of most warm climates. As found in the shops, it is in the form of a light, porous mass, of a yellowish-brown color, and very elastic, and readily imbibing fluids.

It is used for the purpose of cleansing the surfaces of wounds and ulcers, and, in a compressed state, to form tents to dilate sinuses, &c. When burnt, or reduced to charcoal, it has been given successfully in some diseases; probably owing its efficacy to the presence of iodine.

BURNT SPONGE.

R. Sponge, at will.

Cut into pieces, separate any extraneous matters by beating, then burn in a closed iron vessel till it becomes black and friable, and rub into a fine powder.

U. S. Ph.

Used in goitre, glandular swellings of a scrofulous nature, &c.; in doses of one to three drachms.

POWDER OF BURNT SPONGE.

R. Burnt sponge, ten grains.
Powdered rhubarb, four grains.

Mix. To be taken morning and evening. In scrofula.

Hulse.

COMPOUND POWDER OF BURNT SPONGE.

R. Burnt sponge, six drachms.
Powdered cinnamon,

“ ginger, each, one drachm.

Sulphate of potassa, two drachms.

Sugar, five drachms.

Mix.

Greek Ph.

Dose, a teaspoonful, three times a-day, in scrofula.

R. Burnt sponge, four drachms.
Carbonate of magnesia,

Nitre,

White sugar, each, two drachms.

Rub into powder. Dose, as above.

Clarus.

BOLUS OF BURNT SPONGE.

R. Burnt sponge, one scruple.
Sulphate of potassa, fifteen grains.
Balsam of sulphur (*Lond. Ph.*), ten drops.
Syrup, sufficient.

Mix, and form two boluses. One, morning and evening, in scrofula.

Cadet.

ELECTUARY OF BURNT SPONGE.

R. Burnt sponge, half an ounce.
Syrup of orange peel, one ounce and a half.

Mix. Two to four spoonfuls a-day, in goitre, &c.

Radius.

STANNUM.

TIN.

POWDER OF TIN.

R. Tin, at will.

Melt in an iron vessel, and, while cooling, stir till reduced to powder, which is then to be sifted.

U. S. Ph.

As an anthelmintic, in doses of half an ounce, mixed with syrup or molasses, for several mornings, followed by an active purgative.

ELECTUARY OF TIN.

R. Powder of tin, one ounce.
Extract of wormwood,
Powdered jalap, each, one drachm.
Compound syrup of chicory, sufficient.

Mix. To be taken in twelve doses.

Foy.

OINTMENT OF TIN.

R. Amalgam of tin, half an ounce.
Rose ointment, one ounce.
Red precipitate, two drachms.
Oil of peppermint, twenty drops.

Rub well together. Said to be useful in hemorrhoids.

Brera.

STANNI CHLORIDUM.

CHLORIDE OF TIN.

R. Tin, at will.
Muriatic acid, sufficient.

Dissolve, and crystallize.

Van Mons.

Used as an antispasmodic and a vermifuge. Dose, one-eighth to one-half of a grain.

STANNI SULPHURETUM.

SULPHURET OF TIN.

R. Tin, three parts.
Sulphur, one part.

Heat in a crucible, as long as a flame arises; then cool, and pulverize. As a vermifuge, in doses of ten to twenty grains. *Van Mons.*

COMPOUND POWDER OF SULPHURET OF TIN.

R. Sulphuret of tin, four parts.
Powdered jalap, two parts.
Assafetida, one part.

Mix. Twelve to twenty grains, three times a-day, as a vermifuge. *Port. Ph.*

AURUM MUSIVUM—MOAIC GOLD.

R. Tin, twelve parts.

Melt at a low heat in an earthen crucible; add

Mercury, six parts,

and make an amalgam; add

Sulphur, seven parts.

Sal ammoniac, six parts.

Rub well together, introduce into a matras, and heat moderately until sulphuretted hydrogen ceases to be given off. When cold, separate and preserve the upper yellow layer of the product. *Par. Cod.*

Employed under the name of *bronze powder* for ornamental work. It was formerly used as a vermifuge.

STANNI OXIDUM.

OXIDE OF TIN.

R. Tin, at will.

Keep it melted in an open vessel, constantly stirring till it is reduced into a grey powder, and sift. *Swed.aur.*

Has been recommended in tape-worm, in doses of five or six grains, several times a-day.

STAPHISAGRIA.

STAVESACRE.

The seeds of *Delphinium staphisagria*, a beautiful annual plant, with terminal racemes of blue flowers; a native of the south of Europe.

Sex. Syst. Polyand. trigyn. *Nat. Syst.* Ranunculaceæ.

Linn. Sp. Pl. 750. Lindley, Med. Flor. 9.

The seeds are large, somewhat triangular, of a brown color, a slight but unpleasant odor, and an acrid, bitter, nauseous taste. Their active principle, *delphinia*, is occasionally employed. They are principally used to destroy lice in the hair, but also for other purposes.

DECOCTION OF STAVESACRE.

R. Stavesacre, one ounce.
Water, one pint and a half.

Boil down to a pint, and strain. As a lotion in itch. *Swed.aur.*

VINEGAR OF STAVESACRE.

R. Stavesacre, one part.
Vinegar, sixteen parts.

Macerate and express. As a wash to destroy vermin in the hair. *Beral.*

OINTMENT OF STAVESACRE.

R. Stavesacre, two parts.
Lard, three parts.

Suet, each, three parts.
Melt for some time, and strain. To destroy lice. *Bruns. Ph.*

STATICE.

MARSH ROSEMARY.

The root of the *Statice Caroliniana* is occasionally employed in medicine. This little plant is indigenous in the United States, growing in salt marshes near the sea-coast.

Sex. Syst. Pentand. pentagyn. *Nat. Syst.* Plumbaginaceæ.

Nuttall, Gen. i. 206. Griffith, Med. Bot. 525.

Marsh rosemary is a powerful astringent, and is much used in the New England States, in the treatment of diarrhœas, aphthous, and ulcerative affections of the mouth and fauces, and in other cases requiring the use of astringents. Dose, ten to thirty grains.

DECOCTION OF MARSH ROSEMARY.

R. Root of marsh rosemary, bruised, one ounce.

Boiling water, one pint.

Boil for fifteen minutes, and strain. Dose, a wineglassful.

STILLINGIA.

QUEEN'S ROOT.

The root of *Stillingia sylvatica* is the part used. The Queen's root, or Queen's delight, as it is often termed, attains a height of two or three feet. It is found in the pine barrens of the Middle States.

Sex. Syst. Monœc. monad. *Nat. Syst.* Euphorbiaceæ.

Willden. Sp. Pl. iv. 588.

In large doses (twenty to thirty grains), it is emetic and cathartic; in doses of three to five grains, it is an alterative.

DECOCTION OF QUEEN'S ROOT.

R. Bruised Queen's root, one ounce.
Boiling water, twenty fl. ounces.
Boil down to a pint, and strain. Dose, one or two fl. ounces. *G. B. Wood.*

STRAMONIUM.

STRAMONIUM.

THORN-APPLE—JAMESTOWN WEED.

Several parts of the *Datura stramonium* are used in medicine, as the root, leaves, and seeds. It is an annual herbaceous plant, found in the United States, having a rank, fetid odor, and bearing large, funnel-shaped, white flowers.

Sex. Syst. Pentand. monog. *Nat. Syst.* Solanaceæ.

Linn. Sp. Pl. 255. Griffith, Med. Bot. 490.

The leaves have a fetid, narcotic odor, and a bitter, nauseous taste; the odor is lost on drying, but the taste remains. The seeds are small, reniform, of a brownish-black color, inodorous, of a bitter, somewhat acrid taste. Stramonium is a powerful narcotic, and is much used in a variety of affections, both internally and externally. The dose of the powdered leaves is from two to three grains; of the seeds about a grain.

PILLS OF STRAMONIUM SEED.

R. Powdered stramonium seed, ten grains.
" camphor, one drachm.
" savine, five scruples.
Extract of seneka, four scruples.
Mix, and make pills of two grains. Dose, six, three times a-day. In rheumatism. *Vogt.*

COMPOUND STRAMONIUM PILLS.

R. Extract of stramonium, one drachm.
Soap, two drachms.
Powdered gum Arabic, one scruple.
" liquorice, two scruples.
Mucilage of tragacanth, sufficient.
Make mass, and divide into sixty pills. One, night and morning, in asthma. *Halford.*

EXTRACT OF STRAMONIUM SEED.

R. Powdered stramonium seed, one pound.
Diluted alcohol, sufficient.

Rao the powder with half a pint of the alcohol, then introduce into a displacement apparatus, and exhaust by means of diluted alcohol. Distil the filtered liquor, and evaporate the residue to a proper consistence. *U. S. Ph.*

Dose, a quarter to half a grain, twice a-day, gradually increasing.

EXTRACT OF STRAMONIUM LEAVES.

R. Stramonium leaves, one pound.
Bruise in a stone mortar, with a little water. Express, and heat the juice to the boiling point; strain, and evaporate to proper consistence.

U. S. Ph.

Dose, one grain, twice a-day, gradually increasing till it produces its effects.

MIXTURE OF EXTRACT OF STRAMONIUM SEED.

R. Extract of stramonium seed, twelve grains.
Antimonial wine, half a fl. ounce.
Dissolve. Ten drops, every three hours, in an infusion of balm, as an antispasmodic. *Hufeland.*

ANTI-RHEUMATIC LOTION.

R. Stramonium seed, bruised, one ounce.
Alcohol, one pint.
Infuse, strain, and add
Opium, in powder, one ounce.
Camphorated alcohol, two fl. ounces.
Mix. In frictions to the diseased part. *Pierquin.*

SYRUP OF STRAMONIUM.

R. Sugar, fifteen ounces.
Tincture of stramonium, two fl. ounces.
Distilled water, seven fl. ounces.
Mix, and form syrup without heat, and strain. *Beral.*

R. Stramonium seed, bruised, one ounce.
Vinegar, one pound.
Infuse for two days, strain, and add
Sugar, two pounds.
Dissolve. *Wirt. Ph.*

TINCTURE OF STRAMONIUM SEED.

R. Stramonium seed, bruised, four ounces.
Diluted alcohol, two pints.
Macerate for fourteen days, express, and filter. Or by the process of displacement. *U. S. Ph.*
Dose, from twenty to forty drops, two or three times a-day, increasing till it affects the system.

TINCTURE OF STRAMONIUM LEAVES.

R. Filtered juice of stramonium leaves, Alcohol (.847), equal parts.
Mix, and filter at the end of twenty-four hours. *Beral.*

ETHEREAL TINCTURE OF STRAMONIUM.

R. Stramonium seed, one part.
Sulphuric ether, four parts.
Macerate for a week, and filter. Dose, two to three drops. *Soubiran.*

WINE OF STRAMONIUM.

R. Stramonium seed, bruised, two ounces.
Wine, eight fl. ounces.
Alcohol, one fl. ounce.
Macerate for some days, and filter. Dose, six drops to a fl. scruple. *Van Mons.*

OINTMENT OF STRAMONIUM.

R. Fresh stramonium leaves, one pound.
Lard, three pounds.
Wax, half a pound.
Boil the leaves in the lard till they become crisp, then strain through linen; afterwards add the wax, previously melted, and stir till cold. *U. S. Ph., 1840.*

R. Extract of stramonium leaves, one drachm.
Lard, one ounce.
Moisten the extract with a little water, and then rub it with the lard. *U. S. Ph., 1850.*
A valuable anodyne application to painful hemorrhoids, and tumors.

COMPOUND OINTMENT OF STRAMONIUM.

R. Bark of the root of bitter-sweet, Stramonium leaves, Hemlock leaves, Deadly nightshade leaves, Yellow dock root, each, two ounces.
Bruise the roots and leaves, and simmer them in spirit; then add
Lard, one pound,
and simmer gently till the leaves are crisped. Express through linen, and add
Venice turpentine, two ounces.
Stir well together. An useful application to indolent and glandular swellings. *Ecl. Med. Jour.*

OIL OF STRAMONIUM.

R. Stramonium leaves, one part.
Olive oil, two parts.
Heat till all moisture is driven off, then express, and strain. *Cottureau.*

STYRAX.

STORAX

Is the concrete juice of *Styrax officinale*, a small tree, a native of the warm parts of Europe, Syria, &c. It is procured by making incisions in the bark.

Sex. Syst. Decand. monog. *Nat. Syst.* *Styracaceæ.*

Linn. Sp. Pl. 635. *Griffith, Med. Bot.* 437.
Several kinds of Storax are employed; that in grains, in mass, in calamita, and the liquid. It has a fragrant odor, and a warm, aromatic taste. It is a stimulating expectorant. Dose, ten to twenty grains, twice a-day.

PURIFIED STORAX.

R. Storax, sufficient.
Alcohol, each, sufficient.
Dissolve, and strain; distil off the alcohol by a gentle heat, till the storax is of proper consistence. *U. S. Ph.*

COMPOUND PILLS OF STORAX.

R. Purified storax, three drachms.
Powdered opium, Saffron, each, one drachm.
Beat together. Five grains contain one of opium. *Lond. Ph.*

SYRUP OF STORAX.

R. Purified storax, three ounces.
Alcohol, one fl. ounce.
Water, one pint.
Digest for twelve hours on a water-bath, and add
Sugar, two pounds.
Orange-flower water, one fl. ounce.
Mix. *Giordano.*

STRYCHNIA.

STRYCHNINE—STRYCHNIA.

R. Nux vomica, rasped, four pounds.
Powdered lime, six ounces.
Muriatic acid, three fl. ounces.
Alcohol, Diluted sulphuric acid, Solution of ammonia, Purified animal charcoal, Water, each, sufficient.

Digest the nux vomica in two gallons of water, acidulated with a fl. ounce of muriatic acid, for twenty-four hours; boil for two hours, express, and strain through linen. Boil residue twice successively, in the same quantity of acidulated water, and proceed as before. Mix the decoctions, and evaporate to consistence of thin syrup; add lime, previously mixed with a pint of water, and boil for ten minutes, often stirring. Pour mixture into a double linen bag, and wash well with water; press, dry, and powder the precipitate. Treat the powder repeatedly with boiling alcohol, till it loses its bitterness; mix the solutions, and distil off the alcohol in a water-bath. Mix the residue with

water, apply heat, and drop in sufficient diluted sulphuric acid to dissolve the strychnia; treat with animal charcoal at boiling temperature; filter, evaporate, and crystallize. Dissolve the crystals in water, and add solution of ammonia to precipitate the strychnia. Dry the strychnia on bibulous paper. *U. S. Ph.*

R. Powdered nux vomica, nine pounds.
Water,
Sulphuric acid, each, sufficient.
Powdered quicklime, ten ounces.
Alcohol, fifteen pints.

Mix the nux vomica with sufficient water to form a thin paste, and keep at the temperature of 70 or 80°, until gas ceases to form; express, and boil the residue in several successive portions of water, and express. Set liquid aside to deposit; decant, and evaporate to three gallons. Add nine ounces of quicklime, and after six hours' contact, express strongly; heat the liquid to the boiling point, and add a slight excess of sulphuric acid; separate the liquid by decantation, and evaporate to four pints; add one ounce of quicklime, and act as before, adding this precipitate to the former; dry, and powder; digest in five pints of alcohol diluted with five pints of water, at a gentle heat; separate precipitate, and boil in five pints of alcohol twice; mix, and filter solutions; distil off four-fifths, and set aside to crystallize. *Molyn.*

Very active and dangerous. Dose, one-sixteenth to one-tenth of a grain, at first, carefully watching the effects, and slowly increasing.

COMPOUND POWDER OF STRYCHNIA.

R. Strychnia, one grain.
Black oxide of iron,
Sugar,
Gum Arabic, each, one drachm.
Rub well together, and divide into twelve powders. *Brera.*

PILLS OF STRYCHNIA.

R. Strychnia, two grains.
Conserve of roses, thirty-six grains.
Mix, and make twenty-four pills. One or two, morning and evening, in paralysis. *Magendie.*
R. Strychnia, two grains.
Extract of valerian, sufficient.
Mix well, and make thirty-two pills. One, early in the morning, for five days; then one, morning and evening, in amaurosis. *Furnari.*

STIMULANT PILLS.

R. Strychnia, one grain.
Acetic acid, one minim.
Crumb of bread, one scruple.
Mix thoroughly, and make ten pills. One to be taken every six hours, in the paralysis arising from lead. *A. T. Thomson.*

TINCTURE OF STRYCHNIA.

R. Strychnia, three grains.
Alcohol (.842), one fl. ounce.
Dissolve. Dose, from six to twenty-four drops, twice a-day. *Magendie.*

MIXTURE OF STRYCHNIA.

R. Strychnia, one grain.
Distilled water, two fl. ounces.
White sugar, two drachms.
Acetic acid, three drops.
Mix. One fl. drachm contains one-sixteenth of a grain of strychnia. *Beasley.*

OINTMENT OF STRYCHNIA.

R. Strychnia, sixteen grains.
Lard, one ounce.
Rub well together. As a friction on paralyzed parts. *Sandras.*

STRYCHNIA COLLYRIUM.

R. Strychnia, two grains.
Diluted acetic acid, one drachm.
Distilled water, one ounce.
Mix. A few drops of this applied to the eye, several times a-day, is stated to be beneficial in amaurosis. *Henderson.*

LINIMENT OF STRYCHNIA.

R. Strychnia, thirty grains.
Olive oil, one ounce and a half.
Rub well together. Ten drops to be rubbed on the temples, in amaurosis. *Neligan.*

STRYCHNINÆ ACETAS.

ACETATE OF STRYCHNIA.

R. Strychnia, at will.
Acetic acid, sufficient
to dissolve; evaporate, and crystallize. *Griffith.*

SOLUTION OF ACETATE OF STRYCHNIA.

R. Acetate of strychnia, three grains.
Alcohol, one fl. drachm.
Cinnamon water, seven fl. drachms.
Dissolve. Five drops twice a-day. *Radiis.*
R. Strychnia, one grain.
Distilled vinegar, one fl. drachm.
" water, nine fl. drachms.
Mix. Dose, one fl. drachm; containing one-tenth of a grain. *A. T. Thomson.*

TINCTURE OF ACETATE OF STRYCHNIA.

R. Acetate of strychnia, one grain and a half.
Alcohol, half fl. ounce.

Dissolve. Five to twenty drops, twice a-day, in syphilitic pains in the bones. *Fricke.*

STRYCHNIAE IODAS.

IODATE OF STRYCHNIA.

R. Strychnia, at will.
Iodic acid, sufficient
to saturate; dissolve in boiling alcohol, filter, and crystallize. *Magendie.*

R. Solution of muriate of strychnia, at will.
Solution of iodate of soda, sufficient
to precipitate; treat as the last. *Jourdan.*

Dose, one-eighth of a grain, in pill. One, morning and evening, gradually increasing. In paraplegia.

STRYCHNIAE MURIAS.

MURIATE OF STRYCHNIA.

R. Strychnia, at will.
Muriatic acid, sufficient
to saturate; evaporate, and crystallize. *Cottureau.*

Dose, one-eighth of a grain.

STRYCHNIAE NITRAS.

NITRATE OF STRYCHNIA.

R. Strychnia, at will.
Nitric acid, sufficient
to saturate, with the aid of heat; filter, while hot, evaporate, and crystallize. *Giordano.*

This is as active as strychnia, and is used in similar cases. Dose, one-eighth of a grain.

OINTMENT OF NITRATE OF STRYCHNIA.

R. Nitrate of strychnia, one grain and a half.
Lard, two drachms.
Rub well together. Used as a friction on paralyzed parts. *Wendt.*

STRYCHNIAE SULPHAS.

SULPHATE OF STRYCHNIA.

R. Strychnia, at will.
Sulphuric acid, sufficient
to saturate at a gentle heat; filter, whilst hot, evaporate, and crystallize. *Cottureau.*

SYRUP OF SULPHATE OF STRYCHNIA.

R. Sulphate of strychnia, four grains.
Syrup, one pint.

Mix carefully.

Each fl. ounce contains one-quarter of a grain of the sulphate. *Trousseau.*

SUCCINUM.

AMBER.

This is a solid body, of various shades of yellow; it is translucent, or opaque, and of a vitreous fracture; brittle, tasteless, inodorous, except when heated; then exhaling a peculiar, penetrating, but rather agreeable smell. It is a resin found in a fossil state, in various parts of the world. Not much used in medicine, but extensively employed in the arts.

FUMIGATING POWDER.

R. Amber,
Mastich,
Juniper berries, each, four parts.
Cascarilla, one part.
Powder, and mix. *Scherf.*

OIL OF AMBER.

R. Powdered amber, at will.
Mix, with an equal weight of sand, in a glass retort, which is to be only half filled. Then distil on a sand-bath, gradually increasing the heat. Separate the oil from the product, and keep in well-closed bottles. *U. S. Ph.*

RECTIFIED OIL OF AMBER.

R. Oil of amber, one pint.
Water, six pints.
Mix in a glass retort; distil till four pints of water, with the oil, have passed over; then separate the oil, and keep it in well-stopped bottles. *U. S. Ph.*

Stimulant and antispasmodic; in doses of five to fifteen drops; also used externally.

ARTIFICIAL MUSK.

R. Oil of amber, one part.
Add gradually
Nitric acid, two parts.
Let react for twenty-four hours, and wash in cold water. *Van Mons.*

R. Strong nitric acid,
three drachms and a half.

Add very gradually to

Rectified oil of amber, one drachm, in a large, glass vessel. When action has ceased, permit to rest for twenty-four hours; then remove the upper or resinous portion and

wash it well in cold and then in hot water, till all acid is removed.

Williams.

Useful in those cases in which musk is applicable.

TINCTURE OF ARTIFICIAL MUSK.

R. Artificial musk, one part.
Alcohol, eight parts.

Dissolve with a gentle heat.

Forty drops as an antispasmodic. Useful in doses of five or six drops, in hooping-cough in children.

Van Mons.

R. Artificial musk, two drachms.
Alcohol, eight ounces.

Dissolve, and filter. Dose, from twenty-five drops to a drachm, according to age.

Williams.

EMULSION OF ARTIFICIAL MUSK.

R. Artificial musk, twelve grains.
Blanched almonds, four.

Triturate well together, and gradually add

Water, six fl. ounces.

Dose, for a child of two years, two teaspoonfuls. Valuable in hooping-cough.

Hufeland.

LINIMENT OF OIL OF AMBER.

R. Oil of amber, two fl. drachms.
" olives, half fl. ounce.
Laudanum, two fl. drachms.
Brandy, three fl. ounces.

Mix. To be rubbed between the shoulders, in hooping-cough and infantile convulsions.

Parrish.

R. Oil of amber,
" cloves, each, half an ounce.
" olives, one ounce.

Mix.

Beasley.

R. Oil of amber, one drachm.
" nutmeg, two drachms and a half.

Mix. As a friction in hysteria, and headache.

Wirt. Ph.

TINCTURE OF AMBER.

R. Powdered amber, one ounce.
Diluted alcohol, sixteen ounces.

Digest for six days, and filter.

Par. Cod.

Dose, forty to sixty drops.

ETHEREAL TINCTURE OF AMBER.

R. Oil of amber, one part.
Sulphuric ether, fifteen parts.

Mix. Dose, fifteen to thirty drops.

Beral.

ALKALINE TINCTURE OF AMBER.

R. Powdered amber, two ounces.
Solution of carbonate
of potassa, half an ounce.
Diluted alcohol, eight fl. ounces.

Digest for some days, and filter. Dose, twenty to forty drops.

Spielman.

BALSAM OF AMBER.

R. Essence of amber, two fl. ounces.
" myrrh, one fl. ounce.
Oil of turpentine, half fl. ounce.

Mix. Digest at a gentle heat. As an antispasmodic friction.

Radius.

R. Powdered amber, one ounce
Oil of turpentine, two ounces.

Digest at a gentle heat.

Bate.

EAU DE LUCE.

R. Oil of amber, two drachms.
White soap,
Balsam of Mecca, each, fifteen grains.
Rectified spirit, six ounces.

Macerate for eight days, and filter. To one fl drachm of this, add two fl. ounces of water of ammonia. Used in the treatment of snake bites, as an antispasmodic, &c.

Redwood.

MIXTURE OF OIL OF AMBER.

R. Rectified oil of amber, eighty drops
Gum Arabic,
Sugar, each, half an ounce.
Tincture of tolu, two fl. drachms.
Distilled water, four fl. ounces.

Mix. A tablespoonful, every two or three hours, in spasmodic cough.

Ellis.

SULPHUR.

SULPHUR.

Three officinal forms of sulphur are admitted in the U. S. Ph.; viz., *Sublimed Sulphur*, *Washed Sulphur*, and *Precipitated Sulphur*.

SUBLIMED SULPHUR.

R. Sulphur, at will.

Heat at a temperature of 500° to 600° F., in an earthen vessel, and condense the fumes in a large receiver.

WASHED SULPHUR.

R. Sublimed sulphur, at will.

Wash thoroughly with water, as long as any acidity is present.

U. S. Ph.

PRECIPITATED SULPHUR.

- R. Sublimed sulphur, one pound.
 Lime, one pound and a half.
 Water, two gallons.
 Muriatic acid, sufficient.

Slake the lime with a little water, mix it with the sulphur, add the rest of the water, boil for two or three hours, occasionally adding water, and filter. Dilute the liquid with an equal bulk of water, and add sufficient muriatic acid to precipitate the sulphur. Wash the precipitate till the washings are tasteless, and dry it.

U. S. Ph.

Dose, one drachm.

POWDER OF SULPHUR AND CAMPHOR.

- R. Washed sulphur,
 Sugar, each, half a drachm.
 Camphor, six grains.

Mix, and divide into six powders. One, every hour, in mercurial cachexy. *Radius.*

POWDER OF SULPHUR AND LIQUORICE.

- R. Sulphur, one drachm.
 Powdered fennel seed, two drachms.
 " liquorice, four drachms.

Mix. A teaspoonful occasionally, in catarrh. *Phæbus.*

POWDER OF SULPHUR AND ORRIS ROOT.

- R. Washed sulphur,
 eight to eighteen grains.
 Sugar of milk, one scruple.
 Powdered orris root,
 one to three drachms.

Mix, and divide into eight powders. One, every two hours, in milk, in the catarrh of children. *Kopp.*

COMPOUND POWDER OF SULPHUR.

- R. Sulphur, one ounce.
 Powdered fennel seed, one drachm.
 " gum Arabic,
 " sugar, each, two ounces.

Mix. Dose, half a drachm to a drachm, in dysentery. *Van Mons.*

- R. Sulphur, half an ounce.
 Powdered liquorice, one ounce.
 " orris root, two drachms.
 " benzoin, one scruple.
 " sugar, two ounces.

Oil of fennel,
 " anise, each, ten drops.

Mix. Dose, from a scruple to half a drachm. *Wirt. Ph.*

- R. Sulphur, two drachms.
 Cream of tartar, six drachms.
 Powdered fennel seed, one drachm.
 " sugar, half an ounce.

Mix. A spoonful two or three times a-day, as a laxative, in constipation. *Radius.*

- R. Sublimed sulphur,
 one drachm and a half.

Brown sugar,
 Cream of tartar, each, two drachms.

Mix, and make a powder. To be taken twice a-day, in itch, impetigo, or acne. *Ainslie.*

POWDER OF SULPHUR AND CREAM OF TARTAR.

- R. Sublimed sulphur, half an ounce.
 Cream of tartar, one ounce.

Mix with molasses. A teaspoonful, four or five times a-day, to children, in cutaneous affections. *Ellis.*

POWDER OF SULPHUR AND MAGNESIA.

- R. Precipitated sulphur,
 Magnesia, each, half an ounce.

Mix. A teaspoonful, four or five times a-day, as an aperient. *Ellis.*

POWDER OF SULPHUR AND ANTIMONY.

- R. Washed sulphur, two drachms.
 Golden sulphuret of antimony,
 Camphor, each, eight grains.
 Sugar, two scruples.

Mix, and make a powder. *Hufeland.*
 As a diaphoretic.

SULPHUR ELECTUARY.

- R. Sublimed sulphur, two ounces.
 Cream of tartar, one ounce.
 Clarified honey (by weight),
 one ounce.

Syrup of ginger,
 " saffron, each, half a fl. ounce.

Rub all the ingredients in a mortar until thoroughly mixed. Dose, two drachms, as a laxative. *Dub. Ph.*

- R. Washed sulphur,
 Cream of tartar, each, half an ounce.
 Simple, or lemon syrup, sufficient.

Mix, and make electuary. To keep the bowels open in hemorrhoids of pregnant women. A teaspoonful, at night. *Devees*

- R. Precipitated sulphur,
 one drachm and a half
 Orange marmalade, half an ounce

Pulp of tamarinds, two ounces.
 Sugar, one ounce.
 Mix. A tablespoonful three times a-day, in hemorrhoids. *Radiu.*

R. Washed sulphur, one ounce and a half.

Confection of senna, two ounces.
 Nitre, one drachm.
 Syrup of orange peel, sufficient.

Mix. One or two drachms, once or twice a-day, in hemorrhoidal affections. *Ellis.*

R. Sulphur,
 Burnt sponge, each, two drachms.
 Prepared oyster-shell, one drachm.
 Conserve of roses, two ounces and a half.

Mix. A teaspoonful, three or four times a-day, as an alterative, in scrofulous affections. *Vogt.*

COMPOUND ELECTUARY OF SULPHUR.

R. Sublimed sulphur, half an ounce.
 Cream of tartar, one drachm and a half.
 Confection of senna, one ounce.
 Simple syrup, sufficient.

Mix, and make electuary. A teaspoonful, at bedtime, for hemorrhoids. *Ainslie.*

LINCTUS WITH SULPHUR.

R. Washed sulphur, half a drachm.
 Benzoic acid, fifteen grains.
 Syrup of violets, one ounce and a half.
 Oxymel, one ounce.

Mix. A tablespoonful, occasionally, as an expectorant. *Bories.*

ELECTUARY OF SULPHUR.

R. Washed sulphur, half an ounce.
 Confection of senna, one ounce and a half.
 Syrup, sufficient.

Mix. A teaspoonful, morning and evening, in hemorrhoids. *Lewis's Disp.*

SULPHUR MIXTURE.

R. Precipitated sulphur, two drachms.
 Laudanum, twenty drops.
 Water, four fl. ounces.

Mix. One spoonful, three times a-day, in mercurial disease. *Radiu.*

R. Precipitated sulphur, two drachms.
 Fennel water, three fl. ounces.
 Cinnamon water, one fl. ounce.
 Syrup of opium, half a fl. ounce.

Mix. A spoonful every two hours, in diabetes mellitus. *Augustin.*

R. Washed sulphur, one scruple.
 Mucilage, seven drachms.
 Sugar, half an ounce.
 Rose water, one fl. drachm.

A teaspoonful, every hour, in pectoral affections of young children. *Phabus.*

BALSAM OF SULPHUR.

R. Washed sulphur, two ounces.
 Olive oil, eight ounces.

Heat the oil, and gradually stir in the sulphur, till it assumes the consistence of a thick balsam. *Lond. Ph. 1746.*

Used as an external application to foul ulcers.

ETHEREAL BALSAM OF SULPHUR.

R. Terebinthinated balsam of sulphur,
 Sulphuric ether, each, half a drachm.
 Dippel's animal oil, six drachms.

Mix. In flatulent colic. *Augustin.*

TEREBINTHINATED BALSAM OF SULPHUR.

R. Sulphur, one part.
 Oil of turpentine, three parts.

Digest together in a sand-bath, till the oil be saturated with the sulphur. Then separate the balsam from the undissolved sulphur.

As an external application to ulcers, &c. *Ed. Ph., 1722.*

OINTMENT OF SULPHUR.

R. Sulphur, one pound.
 Lard, two pounds.

Mix. *U. S. Ph.*

As an application in itch.

COMPOUND SULPHUR OINTMENT.

R. Sulphur, one ounce.
 Ammoniated mercury,
 Benzoic acid, each, one drachm.
 Oil of bergamot,
 Sulphuric acid, each, one fl. drachm.
 Nitrate of potassa, two drachms.
 Lard, half a pound.

Melt the lard, add the other ingredients, and stir till cold. *U. S. Ph.*

As an application in itch, tinea capitis, crusta lactea, &c.

R. Sulphur, half a pound.
 Powdered white hellebore, two ounces.
 Nitrate of potassa, one drachm.
 Soft soap, half a pound.
 Lard, one pound and a half.
 Oil of bergamot, thirty minims.

Mix. *Lond. Ph., 1836.*

Used as the last, but more irritating.

R. Sulphur, two parts.
Carbonate of potassa, one part.
Lard, eight parts.
Mix. In itch. *Foy.*

R. Sulphur, two ounces.
Powdered muriate of ammonia, two drachms.
Oil of mint, one drachm.
Lard, four ounces.
Mix. In cases of inveterate itch. *Dewees.*

R. Washed sulphur, ten drachms.
Carbonate of potassa, half an ounce.
Cinnabar, one ounce.
Oil of bergamot, half a fl. ounce.
Lard, ten ounces.
Mix. In itch. *Bateman.*

OINTMENT OF SULPHUR AND CAMPHOR.

R. Washed sulphur, half a drachm.
Camphor, one scruple.
Rose ointment, one ounce.
Oil of roses, three drops.
Mix. As an application in psorophthalmia. *Allen.*

OINTMENT OF SULPHUR AND SOAP.

R. Sulphur, one part.
Soft soap, two parts.
Mix. In itch. *Radius.*

OINTMENT OF SULPHUR AND ZINC.

R. Sulphur, two ounces.
Sulphate of zinc, one ounce.
Oil of bayberries, each, six ounces.
Lard, six ounces.
Mix. In tinea capitis. *Sax. Ph.*
R. Sulphur, four parts.
Oxide of zinc, three parts.
Oil of bayberries, six parts.
Lard, twelve parts.
Mix. As last. *Van Mons.*

LINIMENT OF SULPHUR AND SOAP.

R. Soap, three ounces.
Water, six ounces.
Dissolve with a gentle heat, and add
Sulphur, three ounces.
Used as a lotion in itch. *Lugol.*

CERATE OF SULPHUR.

R. Sulphur, two parts.
Cold cream, seven parts.
Oil of almonds, one part.
Mix. *Par. Cod.*

SULPHURIS CARBURETUM,
VEL,
CARBONIS BI-SULPHURETUM.

BI-SULPHURET OF CARBON.

R. Sulphuret of iron, eleven parts.
Charcoal, three parts.
Mix, and introduce into a stone retort, furnished with a glass tube, dipping into water. Separate the sulphuret which collects at the bottom of the water in the recipient, and re-distil it from chloride of lime. *Van Mons.*

R. Sulphur, at will.

Pass the vapor slowly through charcoal, heated to redness, in a porcelain tube, collect the product in a receiver, and purify by another distillation. *Cottreau.*

This fluid is extremely volatile, and hence has been suggested as an anæsthetic agent. But thus far, experiments with it have not shown any superiority over other and safer liquids. In fact, they rather prove the reverse. It is used internally as a sudorific in rheumatism; dose, two to three drops. Also used externally.

DROPS OF SULPHURET OF CARBON.

R. Sulphuret of carbon, one fl. drachm.
Alcohol, half a fl. ounce.
Mix. Four to six drops, every two hours, in rheumatism. *Wutzer.*

MIXTURE OF SULPHURET OF CARBON.

R. Sulphuret of carbon, one scruple.
Cow's milk, six fl. ounces.
Sugar, two drachms.
Mix. A tablespoonful four times a-day, in hypertrophy of the stomach and contraction of the œsophagus. *Clarus.*

LINIMENT OF SULPHURET OF CARBON.

R. Sulphuret of carbon, half an ounce.
Oil of almonds, one ounce.
Mix. As an embrocation in gouty nodes. *Mansfield.*

R. Sulphuret of carbon, one drachm.
Camphorated oil, one ounce.
Beasley.

R. Camphor, two drachms.
Dissolve in
Sulphuret of carbon, half a fl. ounce
And add

Alcohol, one fl. ounce.
As an embrocation in rheumatism. *Lampadius*

- R. Sulphuret of carbon, two fl. drachms.
 Camphorated ammon.
 liniment, two fl. ounces.
 Mix. As an embrocation in rheumatism.
Wutzer.

SULPHURIS IODIDUM.

IODIDE OF SULPHUR.

- R. Iodine, four ounces.
 Sulphur, one ounce.

Rub the iodine and sulphur together, in a porcelain or glass mortar. Put the mixture into a matrass, close the orifice slightly, and apply a gentle heat, so as to darken the mass, but not melt it. When uniformly dark, increase the fire so as to melt the iodine; then incline the vessel in different directions, to return to the mass any portions that have been condensed on the surface of the vessel; lastly, allow the matrass to cool, break it, and put the iodide into well-stopped bottles. *U. S. Ph.*

Used as an external application, in cutaneous affections.

POWDER OF IODIDE OF SULPHUR

- R. Powdered iodide of sulphur, ten grains.
 Powdered gum Arabic, sufficient.
 Mix, and divide into six powders. One, morning and evening, for an adult. Useful in porrigo, prurigo, impetigo, and tinea; also in nocturnal incontinence of urine. *Escolar.*

OINTMENT OF IODIDE OF SULPHUR.

- R. Iodide of sulphur, half a drachm.
 Lard, one ounce.

Rub the iodide with a little of the lard, then add the remainder, and mix. *U. S. Ph.*

- R. Iodide of sulphur, five parts.
 Lard, ninety-six parts.

Mix. *Foy.*

The strength is to be varied according to circumstances. Much used in chronic cutaneous diseases.

T.

TABACUM.

TOBACCO.

Tobacco is the leaves of *Nicotiana tabacum*, an annual, herbaceous plant, a native of the warmer parts of America, and extensively cultivated in the United States, and elsewhere. There are numerous varieties, but, in all of them, the leaves are the part that is used.

Sex. Syst. Pentand. monog. *Nat. Syst.* Solanaceæ.

Linn. Sp. Pl. 258. *Griffith, Med. Bot.* 493.

Tobacco, as found in commerce, consists of the dried leaves, which are packed in bundles; it has a narcotic, penetrating odor, and a bitterish, nauseous, acrid taste. It has the powers of a sedative narcotic, with those of an emetic and diuretic; it also acts as an errhine and sialagogue. In large doses it acts as a poison. It is employed to produce relaxation in spasmodic affections, as a diuretic, &c.

COMPOUND POWDER OF TOBACCO.

- R. Powdered tobacco,
 " valerian, each, two drachms.
 Oil of lavender,
 " marjoram, each, three drops.
 Mix. Used as a cephalic snuff. *Boeli.*

- R. Powdered tobacco, two grains.
 Tartar emetic, one grain.
 Powdered sugar, two drachms.
 " gum Arabic, half a drachm.

Mix, and form twenty powders. One, every two hours, in whooping-cough. *Petschaft.*

EXTRACT OF TOBACCO.

- R. Cut tobacco, four ounces.
 Water, two pints.

Boil, and let simmer for two or three hours, strain, and evaporate to consistence of an extract. For external use in neuralgia. *Chippendale.*

PILLS OF TOBACCO.

- R. Extract of tobacco, one drachm.
 Liquorice, half an ounce.
 Vinegar of squill, sufficient.
 Mix, and make one hundred and eighty pills.
 Dose, one to three in dropsy. *Van Mons.*

WINE OF TOBACCO.

- R. Tobacco, cut, one ounce.
 Wine, one pint.

Macerate for fourteen days, express, and filter.
U. S. Ph.
 From ten to twenty minims, as a diuretic.

TINCTURE OF TOBACCO.

R. Cut tobacco, one ounce.
 Diluted alcohol, one pint.
 Digest for three days, express, and filter. Dose,
 ten minims. *Augustin.*

INFUSION OF TOBACCO.

R. Tobacco, one drachm.
 Boiling water, one pint.
 Macerate for an hour, in a covered vessel, and
 strain. *U. S. Ph.*

Used as an enema; one-half only should be
 used at a time; employed in strangulated her-
 nia, obstinate colic, &c. Great caution must
 be used, as dangerous effects have followed its
 administration.

MIXTURE OF TOBACCO.

R. Tobacco, one drachm.
 Boiling water, two fl. ounces.
 Infuse for twenty minutes, filter, and add
 Alcohol, two fl. drachms.
 Thirty to fifty drops, twice or thrice a-day, in
 hydrothorax and dysuria. *Fowler.*

LOTION OF TOBACCO.

R. Tobacco, two drachms to half an ounce.
 Boiling water, one pint.
 Infuse. As a lotion in psora, but must be used
 with great caution. *Ellis.*

CATAPLASM OF TOBACCO.

R. Tobacco, one ounce.
 Beat up with water, and form a cataplasm. To
 be applied to the throat in croup, and in spasm
 of glottis, &c. *Ellis.*

OINTMENT OF TOBACCO.

R. Fresh tobacco, cut, one ounce.
 Lard, one pound.
 Boil over a gentle fire, till the leaves become
 friable, and strain through linen. *U. S. Ph.*
 In irritable ulcers, tinea capitis, &c. To be
 used with caution.

R. Powdered tobacco,
 Sulphur, each, four ounces.
 Powdered white hellebore, two ounces.
 Oleander leaves, one ounce.
 Common salt, ten drachms.
 Rose ointment, two pounds.
 Mix. As an application in psora, &c. *Taddei.*

R. Extract of tobacco, one drachm.
 Simple cerate, one ounce.
 Mix. As a friction in neuralgia. *Chippendale.*

OIL OF TOBACCO.

R. Kentucky tobacco, at will
 Introduce it into an earthen or iron retort, con-
 nected with a receiver, which should be fur-
 nished with a tube, to conduct the incondensable
 gases to a chimney; heat the retort to the tem-
 perature of 600° F., or thereabout, until the
 empyreumatic oil ceases to distil; separate the
 black tarry product from the acid liquor in the
 receiver, and preserve it for use.

NICOTIA. NICOTINE.

R. Kentucky tobacco, at will.
 Cut to pieces, and extract it with water; evapo-
 rate to an extract, exhaust this with alcohol;
 distil off the alcohol, treat the soft extract with
 an excess of potassa in water; agitate this with
 ether, in separate portions, till the impure nico-
 tine is removed; unite the ethereal liquids, add
 an excess of pulverized oxalic acid, wash the
 oxalate of nicotine which precipitates with ether;
 then treat it with an aqueous solution of potassa,
 and again agitate the liquid with ether to re-
 move the alkaloid, which is obtained, by distil-
 ling off the ether, in the form of a light brown,
 syrupy liquid, and consists of nicotine, ether,
 water, and some ammonia. The last three
 bodies may be removed by heating the nicotine
 in a retort to the temperature of 284° F., in a
 current of hydrogen, for twelve hours; then, by
 raising the heat to 356° F., the nicotine distils
 over pure. *Schloessing.*

Nicotine is too strong for internal adminis-
 tration.

TAMARINDUS.

TAMARINDS.

Tamarinds are the preserved fruit of the
Tamarindus Indica, a large tree, a native of the
 East Indies, and extensively cultivated in the
 tropical parts of America. Those brought here,
 are principally derived from the West Indies.

Sex. Syst. Monadelph. triand. *Nat. Syst.* Fa-
 baceæ.

Linn. Sp. Pl. 48. *Griffith, Med. Bot.* 262.

The preserved pods, as they come to us, are
 in a dark-colored, adhesive mass, formed of
 pulp, fragments of the pods, seeds, and syrup,
 of a sweet acidulous taste. They are cooling
 and laxative; and, when mixed with water,
 form a grateful drink in febrile diseases.

PULP OF TAMARINDS.

R. Tamarinds, at will
 Digest in a small quantity of water, then pass
 through a sieve. *U. S. Ph.*

ELECTUARY OF TAMARINDS.

R. Pulp of tamarinds, one ounce and a half.
 Cream of tartar, half an ounce.
 Syrup of raspberries, sufficient.
 Mix. Two teaspoonfuls, morning and evening,
 as a laxative. *Radius.*

INFUSION OF TAMARINDS.

R. Pulp of tamarinds, one ounce.
 Boiling water, two pints.
 Infuse for one hour, and strain. As a cooling
 drink in fevers. *Cotteneau.*

TAMARIND WHEY.

R. Milk, two pints.
 Tamarinds, two ounces.
 Boil, and strain. As a cooling drink.
Pereira.

TANACETUM.

TANSY.

Tansy, or *Tanacetum vulgare*, is a perennial herbaceous plant, with numerous yellow flowers; a native of Europe, and generally cultivated in our gardens, and has also become naturalized in some places.

Sex. Syst. Syngen. super. *Nat. Syst.* Astera-
 raceæ.

Linn. Sp. Pl. 1184. Griffith, Med. Bot. 406.

The whole plant is officinal; it has a peculiar and strong odor, diminished by drying; and a warm, bitter, aromatic taste. It is an aromatic bitter, and has been praised in intermittents, hysteria, amenorrhœa, as an anthelmintic, &c. As a vermifuge, the seeds are to be preferred.

INFUSION OF TANSY.

R. Fresh tansy, one ounce.
 Boiling water, one pint.
 Infuse, and strain. Used internally, and as an
 injection against ascarides. *Niemann.*

EXTRACT OF TANSY

R. Tansy, one pound.
 Alcohol, one pint.
 Water, eight pints.
 Digest for three days, express, distil off the
 alcohol, and evaporate to proper consistence.
 Dose, six to twenty grains. *Giordano.*

OIL OF TANSY.

R. Tansy, at will.
 Water, sufficient
 to cover. Distil, and separate the oil.
Guibourt.
 Dose, one to two drops, as a vermifuge.

PILLS OF TANSY.

R. Oil of tansy, one fl. scruple.
 Extract of English walnut,
 two drachms.
 Powdered marsh mallow, sufficient.
 Mix, and make sixty pills. Three to five every
 two hours, as a vermifuge. *Radius.*

COMPOUND TINCTURE OF TANSY.

R. Tansy, one ounce.
 Wormwood,
 Rhubarb, each, three ounces.
 Sherry wine, two fl. ounces.
 Diluted alcohol, twenty fl. ounces.
 Digest for eight days, and filter. Dose, one to
 two fl. drachms, two or three times a-day, as a
 vermifuge. *Van Mons.*

TAPIOCA.

TAPIOCA

Is a fecula obtained from the root of *Janipha manihot*, a shrub-like, herbaceous plant, a native of the tropical parts of America, where it is largely cultivated under the name of Cassava, and forms an important article of food.

Sex. Syst. Monœc. monadelph. *Nat. Syst.*
 Euphorbiaceæ.

Kunth. ii. 85. Griffith, Med. Bot. 601.

Tapioca is found in the shops, in the form of irregular, rough, white grains, having little odor or taste, swelling up in hot water, and affording a bland and nutritious diet for the sick and convalescent.

TAPIOCA JELLY.

R. Tapioca, two tablespoonfuls.
 Water, one pint.
 Boil gently for an hour, or till it becomes
 gelatinous; flavor with sugar, wine, &c., ac-
 cording to circumstances. *Ellis.*

TAPIOCA PUDDING.

R. Yolks of eggs, two.
 Sugar, half an ounce.
 Beat together, and stir the mixture with
 Tapioca mucilage, one pint.
 Bake in a slow oven. The mucilage should be
 made with milk, instead of water.
A. T. Thomson.

TARAXACUM.

DANDELION.

This officinal article is the root of *Taraxacum dens-leonis*, a small, herbaceous plant, with a perennial, fusiform root. It is a native of Europe, but has become naturalized in this country.

Sex. Syst. Syngen. æqual. *Nat. Syst.* Asteraceæ.

Haller, i. 23. Griffith, Med. Bot. 414.

The root is fusiform, of a light-brown color externally, succulent, inodorous, and of a mucilaginous, bitterish taste. It is aperient, diuretic, and somewhat tonic, and thought to be resolvent in engorgements of the liver.

DECOCTION OF DANDELION.

R. Bruised dandelion root, four ounces.
Water, a pint and a half.
Boil down to a pint, and strain. *Lond. Ph.*
A wineglassful, two or three times a-day, as a diuretic, &c.

INFUSION OF DANDELION.

R. Bruised dandelion, two ounces.
Boiling water, one pint.
Macerate for two hours in a covered vessel, and strain. *U. S. Ph.*
Dose, a wineglassful, three times a-day.

COMPOUND INFUSION OF DANDELION.

R. Infusion of dandelion, four fl. ounces.
Extract of dandelion, two drachms.
Carbonate of soda, half a drachm.
Tartrate of potassa, three drachms.
Tincture of rhubarb, three fl. drachms.
“ henbane, twenty drops.
Mix. One-third part, three times a-day, in dropsical and visceral affections. *Meigs.*

FLUID EXTRACT OF DANDELION.

R. Dandelion root, fresh, thirty-two ounces.
Slice it, and reduce to a pulp. Mix this with one-sixth its bulk of alcohol, macerate for twenty-four hours, and express strongly. Add a pint of water containing a little alcohol, and again express. Evaporate the mixed products to twelve fl. ounces, add four fl. ounces of alcohol, and filter.
A teaspoonful is equal to half a drachm of the extract obtained from the expressed juice. *W. Procter.*

EXTRACT OF DANDELION.

R. Dandelion, gathered in September, five pounds.
Slice the dandelion; bruise it in a stone mortar, sprinkling on it a little water, until reduced to a pulp. Then express the juice, strain, and evaporate in a shallow dish over a water-bath, constantly stirring, to the proper consistence. *U. S. Ph.*
Dose, a scruple to a drachm, three times a-day, alone, or dissolved in cinnamon or mint water.

R. Bruise the recent root, collected in September; add one-fourth its bulk of alcohol; allow the whole to macerate twelve hours, express powerfully, add a little diluted alcohol, and again express. Evaporate the mixed liquids in a water-bath to the proper consistence.

W. Procter.

PREPARED JUICE OF DANDELION

R. Fresh dandelion root, gathered in September or October, twenty pounds.
Alcohol (.835), four pints.

Slice the roots, and reduce to a pulpy mass; then add the alcohol, and mix thoroughly. Set aside in suitable vessels. When wanted for use, express the pulpy mass in a screw press, and add sufficient sugar to the juice to form a syrup. *W. Procter.*

FLUID EXTRACT OF DANDELION AND SENNA.

R. Senna, two pounds.
Torrefied dandelion root, one pound.
German chamomile, quarter of a pound.
Sugar, twenty ounces.
Carbonate of potassa, or carb. of soda, one ounce.
Oil of winter-green, half a drachm.
Alcohol, two ounces.
Water, half a gallon.

Powder the dry plants, and mix them with the water, holding the alkaline carbonate in solution. Let the mixture stand for twelve hours, then introduce into a percolator, and add water until a gallon of liquid shall have passed. Evaporate on a water-bath to twenty ounces, add the sugar, filter, and when cold, add the alcohol, holding the oil of winter-green in solution. Dose, a teaspoonful to a tablespoonful. *E. Dupuy.*

PILLS OF EXTRACT OF DANDELION.

R. Extract of dandelion, half a drachm.
Powdered liquorice, sufficient.
Mix, and make eight pills. One, three times a-day. In dropsy, and diseases of the urinary apparatus. *Ellis.*
R. Extract of dandelion, Soap, each, one ounce.
Liquid acetate of potassa, sufficient.
Mix, and make pills of four grains. Four to six in a day, as diuretic, &c. *St. Marie.*

PILLS OF DANDELION AND BLUE MASS.

R. Extract of dandelion, half a drachm.
 Blue pill, five to ten grains.
 Powdered uva ursi, sufficient.

Mix, and make ten pills. One, thrice a-day.
 In dropsy connected with liver disease. *Ellis.*

MIXTURE OF DANDELION.

R. Extract of dandelion, two ounces.
 Peppermint water, six fl. ounces.
 Clarified honey, one ounce.

Mix. Two spoonfuls, every three hours.

Richter.

R. Extract of dandelion, half a drachm.
 Decoction of dandelion, nine fl. drachms.
 Spirit of nitric ether, one fl. drachm.
 Syrup of ginger, two fl. drachms.

Mix. To be taken three times a-day, as a diuretic in anasarca and ascites. *Sprague.*

R. Cream of tartar, two drachms.
 Bruised cloves, two scruples.
 Sugar, one ounce.
 Decoction of dandelion, one pound.

Macerate for two hours, and strain. One-fourth to be given every six hours, in dropsy.

Sprague.

CLYSTER OF DANDELION.

R. Dandelion root, bruised, three ounces.
 Bran, one ounce.
 Water, sixteen fl. ounces.

Boil down to eight fl. ounces; add, at close,

Chamomile,
 Valerian, each, two drachms.

Strain; for two injections. Much praised in obstructions of the bowels. *Berends.*

TEREBINTHINA.

TURPENTINE.

Turpentine is the juice of various species of the Pine tribe, and is also procured from trees of other orders. That used in the United States is obtained from several species of *Pinus* and *Abies*, but chiefly from the *P. palustris*.

Sex. Syst. Monœc. monadelph. *Nat. Syst.* Pinacœ.

Willden. Sp. Pl. iv. 499. Griffith, Med. Bot. 604.

The officinal turpentine consists, essentially, of a volatile oil and a resin. All the turpentines are possessed of much the same properties; they have an aromatic odor, and a somewhat pungent and bitterish taste. They are stimulant, diuretic, anthelmintic, and, in large doses, purgative. They are also used externally as rubefacients.

TURPENTINE PILLS.

R. Turpentine, one drachm.
 Divide into fifteen pills. One, four or five times a-day, in affections of the urinary organs.

Ellis.

PILLS OF TURPENTINE AND GUAIAIACUM.

R. Powdered guaiacum, one drachm.
 Venice turpentine, sufficient.

Mix, and make fifteen pills. One, thrice a-day; in gleet and leucorrhœa. They sometimes cause strangury.

Ellis.

PILLS OF TURPENTINE AND MAGNESIA.

R. Turpentine, fourteen drachms.
 Calcined magnesia, thirty-six grains.

Mix, and at the end of twelve hours make pills of six grains. In diseases of the urinary organs.

Monchon.

PILLS OF TURPENTINE AND RHUBARB.

R. Turpentine, two ounces.
 Extract of rhubarb, three drachms.
 Camphor, two drachms.

Mix, and make pills of four grains. Dose, three, thrice a-day, in leucorrhœa and gonorrhœa.

St. Marie.

PILLS OF TURPENTINE AND MYRRH.

R. Turpentine,
 Spermaceti, each, half an ounce.
 Powdered myrrh, two drachms.
 " olibanum, sufficient.

Mix, and make pills of three grains. One every three hours, in chronic catarrh.

Cadet.

PILLS OF TURPENTINE AND JALAP.

R. Turpentine, one drachm.
 Soap of jalap, half a drachm.
 Extract of henbane, four grains.
 Calomel, eight grains.

Make pills of three grains. Dose, four every three hours, for two days. As an anthelmintic.

Augustin.

ELECTUARY OF TURPENTINE.

R. Turpentine,
 Soap,
 Powdered rhubarb, each, one drachm.
 Syrup, sufficient.

Mix. Three teaspoonfuls a-day, in dropsy.

Radius.

TURPENTINE CLYSTER.

R. Venice turpentine, half a fl. ounce.
 Yolk of egg, one.

Infusion of flaxseed, ten fl. ounces.
 Rapeseed oil, one fl. ounce.
 Mix. In colic, and obstinate constipation.
Ph. Noscom. Ed.

LINIMENT OF TURPENTINE.

R. Yellow wax, one part.
 Melt, and add
 Turpentine,
 Oil of turpentine, each, one part.
 Mix. Much praised in chronic ulcers on the legs.
Van Mons.

R. Yellow wax, half a pound.
 Olive oil, one pound.
 Red saunders, two ounces.
 Turpentine, one pound.
 Melt together, strain, and add
 Camphor, two drachms.
 As an application to contusions and ulcerations.
Van Mons.

R. Oil of turpentine,
 Olive oil, each, one ounce and a half.
 Diluted sulphuric acid, three drachms.
 Mix. As an embrocation in rheumatism.
Chapman.

OINTMENT OF TURPENTINE.

R. Turpentine, two pounds.
 Simple ointment, nine ounces.
 Melt together, with a gentle heat, constantly stirring.
Austrian Ph.

COMPOUND OINTMENT OF TURPENTINE.

R. Mucilage, one pound.
 Fresh butter, two pounds.
 Olive oil, three pounds and a half.
 Melt together, and add
 Turpentine, three ounces.
 Mix well. As an application in engorgement of the breasts, in nephritic pains, &c.
Swediaur.

TURPENTINE PLASTER.

R. Turpentine, one ounce.
 White of egg,
 Wheat flour,
 Sugar,
 Honey,
 Brandy,
 Olive oil, each, one ounce.
 Mix the turpentine and honey, gradually adding the other ingredients, and rub well. As an application to painful ulcers.
St. Marie.

COMPOUND TURPENTINE PLASTER.

R. Turpentine, eight parts.
 Suet, twenty-four parts.
 Yellow wax, thirty-two parts.
 Essence of mint,
 " cloves, each, one part.
 Oil of mace, six parts.
 Powdered benzoin, eight parts.
 " olibanum, sixteen parts.
 Mix, with the assistance of heat. Useful in diarrhœa, dyspepsia, flatulence, and colic.
Ph. Sued.

TURPENTINE MIXTURE.

R. Turpentine, two drachms.
 Yolk of egg, one.
 Rub together, and add
 Extract of rhubarb,
 Sal prunelle, each, one drachm.
 Then add gradually
 Infusion of liquorice, ten fl. ounces.
 Syrup of marsh mallow, one fl. ounce.
 Mix. A tablespoonful every hour, in gonorrhœa.
Bories.

TURPENTINE MIXTURE.

R. Turpentine, one ounce.
 Mint water, four fl. ounces.
 Gum Arabic, sufficient.
 Simple syrup, one ounce.
 Extract of belladonna, one grain.
 Mix, and make emulsion. Used in gonorrhœa, after the reduction of the inflammatory stage; when it acts too powerfully on the bowels, it must be intermitted for a few days.
Ebriart.

TESTA PRÆPARATA.

PREPARED OYSTER-SHELL.

(See Page 156.)

TILIA EUROPÆA.

LINDEN.

The linden is a handsome tree, a native of the north of Europe, and much cultivated in the United States for ornament and shade.

Sex. Syst. Polyand. monog. *Nat. Syst.* Tiliaceæ.

Linn. Sp. Pl. 733. *Lindley, Flor. Med.* 147.

The part used is the flowers, which, when dried, have a faint but agreeable odor, and a sweetish, mucilaginous taste. They are antispasmodic, and diaphoretic, and are much used in France. It is probable that the flowers of our native species have the same properties.

INFUSION OF LINDEN FLOWERS.

R. Linden flowers, two drachms.
Boiling water, two pints.
Infuse for half an hour, and strain. A mild
antispasmodic. *Par. Cod.*

COMPOUND INFUSION OF LINDEN FLOWERS.

R. Linden flowers,
Chamomile flowers,
Orange leaves, each, two drachms.
Boiling water, two pints.
Infuse for half an hour, strain, and add
Syrup, two fl. ounces.
A small cupful, occasionally, as an antispas-
modic. *Foy.*

BALSAMUM TOLUTANUM.

BALSAM OF TOLU.

This is the concrete juice of *Myrospermum toluiferum*, a large tree, (of which little is known), growing in several parts of South America.

Sex. Syst. Decand. monog. *Nat. Syst.* Fa-
baceæ.

De Candolle, *Prod.* ii. 95. Griffith, *Med.*
Bot. 250.

As first obtained, the balsam is soft and tena-
cious, but it gradually becomes hard and brit-
tle; then it is transparent, shining, of a yellow-
ish, or reddish-brown color, of a fragrant odor,
and a sweetish, warm taste. It is a stimulating
expectorant. The dose is from ten to thirty
grains.

LOZENGES OF TOLU.

R. Balsam of Tolu, one ounce.
Alcohol, one fl. ounce.
Dissolve and add
Water, two fl. ounces.
Heat in a water-bath, and filter; add,
Tragacanth, four scruples.
Sugar, sixteen ounces.
Beat into a paste, and make lozenges of twelve
grains. *Par. Cod.*

TINCTURE OF TOLU.

R. Balsam of Tolu, three ounces.
Alcohol, two pints.
Macerate till dissolved, then filter. *U. S. Ph.*
A highly stimulating expectorant. Dose, one
to two fl. drachms.

COMPOUND TINCTURE OF TOLU.

R. Balsam of Tolu, two ounces.
Balsam of Peru, one ounce.

Benzoic acid,
Saffron, each, half an ounce.
Alcohol, twenty-four fl. ounces.
Digest for three days, and filter. *Spielman.*

TINCTURE OF TOLU WITH FOXGLOVE.

R. Tincture of
Tolu, one fl. ounce and a half.
Elixir of vitriol, half a fl. drachm.
Tincture of foxglove, one fl. drachm.
Antimonial wine, two fl. drachms.
Clarified honey, one fl. ounce and a
half.
Powdered liquorice, half a drachm.
Distilled water, six fl. ounces.
Mix. A tablespoonful, according to circum-
stances. *Ellis.*

EMULSION OF TOLU.

R. Balsam of Tolu, three drachms.
" Mecca, eight drops.
Sweet almonds, half an ounce.
Barley water, eighteen fl. ounces.
Sugar, six drachms.
Make an emulsion. A spoonful occasionally,
in chronic catarrh, &c. *Spielman.*

SYRUP OF TOLU.

R. Tincture of Tolu, one fl. ounce and a
half.
Water, one pint.
Sugar, two pounds and a half.
Mix the tincture with the sugar, and expose the
mixture in a shallow dish to a gentle heat until
the alcohol has evaporated; then add the water,
heat gradually till the sugar is dissolved, and
strain. *U. S. Ph.*
R. Tincture of Tolu, two fl. ounces.
Carbonate of magnesia, two drachms.
Sugar, (avoir.) one pound and a half.
Water, twelve fl. ounces.
Rub the tincture with the carbonate and two
ounces of the sugar, gradually add the water,
and filter; then dissolve the remainder of the
sugar, by a gentle heat, in the filtered liquid.
This affords a clear, transparent syrup. *J. D. Finley.*

R. Tincture of
Tolu, one and a half fl. ounces.
Sugar, (troy) two and a half pounds.
Water, one pint.
Mix the tincture with one pound of sugar, in a
shallow dish, and allow the alcohol to evaporate
spontaneously. Then add the remainder of the
sugar, and dissolve it in twelve fl. ounces of the
water. Beat up the white of an egg with the
remaining four ounces of water, add it to the
syrup, boil for a minute or two, and strain
through flannel. *J. Laidley.*

R. Balsam of Tolu, ten drachms.
Boiling water, (Imp.) one pint.
Sugar, two pounds and a half.
Boil the balsam in the water for half an hour, occasionally stirring; strain when cold, add the sugar, and form syrup. *Lond. Ph.*

MIXTURE OF TOLU AND MORPHIA.

R. Oxymel of
squill, one ounce and a half.
Syrup of Tolu, half an ounce.
Acetate of morphia, half to one grain.
Mix. A teaspoonful, as occasion may require, in catarrh. *Ellis.*

MIXTURE OF TOLU AND BELLADONNA.

R. Syrup of Tolu, three fl. ounces and a half.
" seneka, half a fl. ounce.
Extract of bella-donna, eight to twelve grains.
Wine of ipecacuanha, one fl. drachm.
Mix. A teaspoonful, three or four times a-day, in cough. *Ellis.*

MIXTURE OF TOLU AND ALMOND EMULSION.

R. Almond emulsion, fifteen parts.
Syrup of Tolu, one part.
Mix. In cough. *Beral.*

MIXTURE OF TOLU AND COPAIBA.

R. Balsam of Tolu,
Copaiba,
Powdered gum
Arabic, each, half an ounce.
Elixir of vitriol, twenty drops.
Distilled water, six fl. ounces.
Mix. A tablespoonful occasionally, in chronic hooping-cough. *Ellis.*

TOLU MIXTURE.

R. Tincture of Tolu, one ounce.
Laudanum, two drachms.
Tincture of foxglove, one drachm.
Mix. In the chronic cough of hæmoptysis. Forty to fifty drops, every three or four hours. *Dewees.*

MIXTURE OF TOLU AND OPIUM.

R. Balsam of Tolu, one drachm.
Powdered gum
Arabic, one drachm and a half.
Water, four fl. ounces.
Syrup of opium, half a fl. ounce.
Make an emulsion. *Niemann.*

INHALATION OF TOLU.

R. Balsam of Tolu, one ounce.
Boiling water, one pint.
Mix. The vapor to be inhaled. *Ellis.*

TORMENTILLA.

TORMENTIL.

The root of *Potentilla tormentilla*, a small, perennial plant, with a large, woody, dark-brown root, a native of many parts of Europe, growing in barren spots.

Sex. Syst. Icosand. polygyn. *Nat. Syst.* Rosaceæ.

Sibthorp, Fl. Ox. 162. Griffith, Med. Bot. 278.

The root, when dried, is in irregular pieces, which are knotty, tuberculated, and of a dark, reddish-brown color. Its odor is slightly aromatic and its taste very astringent. It is a simple and somewhat active astringent. Dose, thirty grains to a drachm.

COMPOUND POWDER OF TORMENTIL.

R. Powdered tormentil,
" gum Arabic,
each, three ounces.
Armenian bole, six ounces.
Powdered cinnamon, four ounces.
" long pepper, half an ounce.
Mix. *Geneva Ph.*

EXTRACT OF TORMENTIL.

R. Tormentil, one part.
Water, eight parts.
Boil, and strain; repeat with same quantity of water, mix the two decoctions, strain, and evaporate to the proper consistence.
Dose, from twenty to thirty grains. *Amst. Ph.*

DECOCTION OF TORMENTIL.

R. Tormentil, bruised, two ounces.
Distilled water, one pint and a half.
Boil down to a pint, and strain. *Lond. Ph.*
Dose, one to two fl. ounces, three or four times a-day.

GARGLE OF TORMENTIL.

R. Tormentil, one ounce.
Water, twelve fl. ounces.
Boil down one-third, strain, and add
Alum, one drachm.
Honey, one ounce.
Mix. *Radius*

TOXICODENDRON.

POISON OAK.

The leaves of *Rhus toxicodendron*, a shrubby plant, a native of the United States, are the parts used. This plant sometimes becomes a vine, whence it has received the name of *Rhus radicans*, or Poison vine. Both varieties grow in woods, along fence rows, &c., and are extremely poisonous to some persons, whilst other individuals are not affected by them.

Sex. Syst. Pentand. trigyn. *Nat. Syst.* Anacardiaceæ.

Linn. Sp. Pl. 381. *Griffith, Med. Bot.* 184.

The leaves are inodorous, and of a disagreeable, acrid taste. They are narcotic and stimulant, acting like the acro-narcotic poisons in large doses. They have been used with success in obstinate cutaneous affections, chronic rheumatism, &c. The dose is half a grain, gradually increased.

EXTRACT OF POISON OAK.

R. Leaves of poison oak, one part.
Alcohol, four parts.

Infuse, strain, distil off the spirit, and evaporate to proper consistence. *Par. Cod.*

Dose, one grain, gradually increased till some effect is induced.

PILLS OF POISON OAK.

R. Extract of poison oak,
Camphor, each, fifteen grains.
Extract of arnica,
“ valerian, each, thirty grains.

Powdered calamus, sufficient.

Beat into mass, and make thirty pills. One, three times a-day, in hemiplegia. *Brera.*

TINCTURE OF POISON OAK.

R. Clarified juice of poison oak,
Alcohol, equal parts.

Mix, and filter at end of twenty-four hours. Dose, five drops, gradually increased.

TRAGACANTHA.

TRAGACANTH.

A gummy exudation from *Astragalus verus*, and other species of *Astragalus*. All of them are small shrubs, with thorny petioles, found in the countries bordering on the Levant, in Persia, &c.

Sex. Syst. Diadelph. decand. *Nat. Syst.* Fabaceæ

Lindley, Flor. Med. 247. *Griffith, Med. Bot.* 239.

Tragacanth is in contorted, vermicular pieces, rounded or flattened, of a whitish, or whitish-

yellow color; semi-translucent and corneous; tough, except when quite dry. It is inodorous, and of a bland, mucilaginous taste. It swells up and forms a paste or mixture with water, but does not dissolve. It is demulcent and nutritive

COMPOUND TRAGACANTH POWDER.

R. Powdered tragacanth,
“ gum Arabic,
“ starch, each,

one ounce and a half.

Sugar, three ounces.

Mix well. *Lond. Ph.*

Demulcent. Dose, from thirty grains to a drachm.

MUCILAGE OF TRAGACANTH.

R. Tragacanth, one ounce.
Boiling water, one pint.

Macerate for twenty-four hours, occasionally stirring, triturate till uniform, and strain forcibly through linen. *U. S. Ph.*

Principally used as a basis for more active medicines.

PASTE OF TRAGACANTH.

R. Tragacanth, thirty-two parts.

Macerate in

Water, one thousand parts.

Also, dissolve

Isinglass, forty-eight parts,

in

Water, eight hundred parts.

Strain through a linen cloth, and mix with the macerated gum; then add

Syrup, one thousand parts.

Evaporate by a gentle fire, to the consistence of a soft paste, and gradually add

Orange-flower water, sixty-four parts.

Evaporate to proper consistence on a water-bath. As an expectorant. *Foy.*

R. White tragacanth, sixty-four parts.

Isinglass, ninety-six parts.

Water, three thousand parts.

Simple syrup, two thousand parts.

Orange-flower water,

one hundred and twenty-eight parts.

Bruise the tragacanth, and place it in a vessel for two days with five-sixths of the water, and dissolve the isinglass in the rest of the water; strain both solutions; heat the syrup to boiling, and mix the whole, and continue heat, till the mixture is of the consistence of soft paste, stirring well. Remove from fire; add orange-flower water, and evaporate on water-bath, till sufficiently firm; pour on marble slab, and divide into lozenges. *Mouchon.*

TRIOSTEUM.

FEVER-ROOT.

The root of *T. perfoliatum*, is placed on the secondary list of the U. S. Pharmacopœia.

Sex. Syst. Pentand. monog. *Nat. Syst.* Caprifoliaceæ.

Linn. Sp. Pl. 395. Griffith, Med. Bot. 352.

Fever-root or fever-wort is found in most parts of the United States. The root is of a yellowish or brownish color externally, and it has a nauseous odor, and a bitter taste. It is a mild cathartic, in doses of ten to fifteen grains.

EXTRACT OF FEVER-ROOT.

R. Powdered fever-root, one pound.
Diluted alcohol, four pints.

Moisten the powder with half a pint of the diluted alcohol, macerate for twenty-four hours, transfer to a percolator, and add gradually the remainder of the alcohol. Add water until the tincture passes tasteless. Then distil off the alcohol, and evaporate to the consistence of an extract. *Thomas.*

Dose, five to eight grains.

TUSSILAGO.

COLTSFOOT.

Coltsfoot is a small, perennial plant, with large, radical, cordate leaves; and flowers on simple, leafless scapes. It is the *Tussilago farfara*, and is a native both of Europe and North America.

Sex. Syst. Syngen. super. *Nat. Syst.* Astera-cææ.

Linn. Sp. Pl. 1214. Griffith, Med. Bot. 393.

The whole herb is used, but the leaves and flowers should be preferred. The latter have an agreeable smell; the former are inodorous, but have a rough, bitterish, mucilaginous taste. Both are demulcent, and have been much used in pectoral affections.

DECOCTION OF COLTSFOOT.

R. Flowers of coltsfoot, one ounce.
Water, two pints.

Boil down to a pint, and strain. *Beasley.*

Dose, a wineglassful.

COMPOUND DECOCTION OF COLTSFOOT.

R. Flowers of coltsfoot, six ounces.
Figs,
Raisins,
Jujubes, each, two ounces.
Water, twelve pints.

Boil down one-third, adding

Liquorice root, two ounces.

Strain. As a demulcent drink. *Taddei.*

SYRUP OF COLTSFOOT.

R. Flowers of coltsfoot, one pound.
Boiling water, two pints.

Macerate for twelve hours, express, strain, and add

Sugar, four pounds.

Make syrup. *Par. Cod.*

U.

ULMUS CAMPESTRIS.

ELM BARK.

This is a large tree, a native of Europe, having strong, spreading branches, and a rough, cracked bark.

Sex. Syst. Pentand. digyn. *Nat. Syst.* Ulma-cææ.

Linn. Sp. Pl. 327. Lindley, Flor. Med. 307.

The part used is the inner bark; this is thin, tough, of a brownish-yellow color, inodorous, of a mucilaginous, slightly astringent taste. It is demulcent, and somewhat astringent, and also acts as an alterative.

DECOCTION OF ELM BARK.

R. Elm bark, bruised,
two ounces and a half.

Distilled water, two pints.
Boil down to one pint, and strain. *Lond. Ph.*

From four to six fl. ounces, two or three times a-day, in cutaneous affections.

COMPOUND INFUSION OF ELM BARK.

R. Elm bark,
Burdock root,
Dock root,
Bittersweet,
Fumitory, each, two drachms
Boiling water, one pint

Infuse for four hours, strain, and add

Syrup of sarsaparilla, one fl. ounce

To be taken in twenty-four hours, in divided doses, in chronic cutaneous diseases. *Cadet*

ULMUS FULVA.

SLIPPERY ELM.

A very lofty tree, found in the United States, especially west of the mountains, growing in dry, elevated situations.

Sex. Syst. Pentand. digyn. *Nat. Syst.* Ulmaceæ.

Mich. N. Am. Sylv. iii. 89.

The inner bark is the part used. This is in long, flat strips, of a tawny color on the outer surface, and reddish on the inner, of a peculiar, though feeble odor, and a mucilaginous taste. It is a good demulcent, and is much used in diseases of the mucous membranes. It is also very nutritious.

INFUSION OF SLIPPERY ELM BARK.

R. Slippery elm bark, bruised, one ounce.
Boiling water, one pint.
Macerate for two hours, and strain. *U. S. Ph.*
To be used freely as a demulcent.

CATAPLASM OF SLIPPERY ELM BARK.

R. Ground slippery elm bark, at will.
Hot water, sufficient.
Make a cataplasm. A soothing application to irritable ulcers, &c.

UVA URSI.

UVA URSI—BEARBERRY.

Uva ursi is the leaves of *Arctostaphylos uva ursi*, a small, evergreen shrub, inhabiting the northern parts of both continents, and the high mountains in more southern latitudes.

Sex. Syst. Decand. monog. *Nat. Syst.* Ericaceæ.

Sprengel, ii. 287. Griffith, Med. Bot. 425.

The leaves, when dried, have a somewhat hay-like smell; their taste is astringent and bitterish. They are astringent and tonic, with some special action on the urinary organs. They are much used in diseases of the genito-urinary organs. Dose, in powder, from a scruple to a drachm, three or four times a-day.

COMPOUND POWDER OF UVA URSI.

R. Powdered uva ursi,
" Peruvian bark, each,
two drachms.
" opium, three grains.
Mix, and divide into six powders. One, twice a-day, with lime water, in nephritic complaints, and as an antilithic. *Ferriar.*

R. Powdered uva ursi, one drachm and a half.
Bicarbonate of soda, one drachm.
Mix, and divide into twelve powders. One, thrice a-day, in diseases of the kidney and bladder. *Ellis.*

R. Powdered uva ursi, half an ounce.
" gum Arabic, two drachms.
" jalap, one drachm.
" sugar, half an ounce.
Oil of orange-peel, six drops.
Mix. Dose, one drachm, in nephritic colic. *Quarin.*

DECOCTION OF UVA URSI.

R. Uva ursi, one ounce.
Water, twenty fl. ounces.
Boil down to a pint, and strain. *U. S. Ph.*
Dose, from one to two fl. ounces, three or four times a-day.

EXTRACT OF UVA URSI.

R. Uva ursi, two pounds and a half.
Boiling water, two gallons.
Macerate for twenty-four hours, boil down to a gallon, strain while hot, and evaporate to a proper consistence. *Lond. Ph.*
Dose, five to thirty grains.

MIXTURE OF UVA URSI.

R. Uva ursi, one ounce and a half.
Milfoil, one ounce.
Water, three pints.
Boil down to two pints, and add, at close,
Liquorice, one ounce.
Strain, and add
Syrup of cinnamon, two ounces.
To be taken in divided doses, in passive hemorrhages of lungs and bladder. *Radius.*

SYRUP OF UVA URSI.

R. Uva ursi, eight ounces.
Boiling water, two pints.
Infuse the finely-bruised leaves in the water for three hours, put in a displacement filter, and add water sufficient to obtain two pints of infusion. Evaporate to one pint, and add two pounds (av.) of sugar, and make syrup. *W. Procter.*

V.

VALERIANA.

VALERIAN.

Several species of Valerian are used in medicine, but the only one officinal is the *Valeriana officinalis*, a handsome, perennial, herbaceous plant, with white or rose-colored flowers, in terminal corymbs.

Sex. Syst. Triand. monog. *Nat. Syst.* Valerianaceæ.

Linn. Sp. Pl. 45. Griffith, Med. Bot. 384.

The part used is the root; this consists of many long, slender fibres, arising from a tuberculated rhizome; the color is yellowish, or brownish; the odor is strong and peculiar, and the taste is bitter and aromatic. It is an active antispasmodic, and is much used in nervous diseases. The dose is from half a drachm to a drachm, two or three times a-day.

COMPOUND POWDER OF VALERIAN.

R. Powdered valerian, one drachm.
 " orange leaves, one scruple.
 Muriate of ammonia, two grains.
 Oil of cajeput, four drops.
 Mix. A teaspoonful, four times a-day, in hot tea. In epilepsy. *Niemann.*

R. Powdered valerian, one ounce.
 Oxide of zinc, one scruple.
 Musk, ten grains.
 Mix. As an antispasmodic, in teaspoonful doses. *Brera.*

R. Powdered valerian, one scruple.
 Aromatic powder, ten grains.
 Mix. To be taken three or four times a-day. In hysteria, hemicrania, chlorosis, &c. *A. T. Thomson.*

BOLUS OF VALERIAN AND IRON.

R. Powdered valerian, one drachm.
 Carbonate of iron, ten grains.
 Mucilage of gum Arabic, sufficient.
 Mix, and make bolus. One, three times a-day, as antispasmodic. *Ellis.*

BOLUS OF VALERIAN AND SULPHATE OF POTASSA.

R. Powdered valerian, two drachms and a half.
 Sulphate of potassa, eighteen grains.
 Syrup of orange-peel, sufficient.
 Mix, and make eighteen boluses. *Cadet.*

COMPOUND PILLS OF VALERIAN.

R. Powdered valerian, half a drachm.
 Castor,
 Oxide of zinc, each, one scruple.
 Mix, and make eighteen pills. Three, thrice a-day. *Dupuytren.*

ELECTUARY OF VALERIAN.

R. Powdered valerian,
 " orange-leaves, each, six drachms.
 Syrup of wormwood, sufficient.
 Mix. Two or three teaspoonfuls a-day. *Swediaur.*

R. Powdered valerian, two ounces.
 " Peruvian bark, half an ounce.
 Carbonate of ammonia, two drachms.
 Ginger syrup, sufficient.
 Mix. Dose, a drachm every hour or two, in periodic hemicrania. *Donald Monro.*

INFUSION OF VALERIAN.

R. Valerian, half an ounce.
 Boiling water, one pint.
 Macerate for an hour in a covered vessel, and strain. *U. S. Ph.*
 Dose, two fl. ounces, three or four times a-day.

COMPOUND INFUSION OF VALERIAN.

R. Valerian, one ounce and a half.
 Decoction of Peruvian bark, two pints.
 Make an infusion, strain, and add

Syrup of chamomile, one fl. ounce and a half.
 Camphor, twenty grains.
 Mix. One to two fl. ounces, three times a-day. *Saunders*

TINCTURE OF VALERIAN.

R. Valerian, bruised, four ounces.
 Diluted alcohol, two pints.
 Macerate for fourteen days, express, and filter. Or prepare by displacement. *U. S. Ph.*
 Dose, from one to four fl. drachms.

ETHEREAL TINCTURE OF VALERIAN.

R. Coarsely-powdered valerian, one ounce.
 Spirit of sulphuric ether, eight ounces.

Macerate for eight days, and filter. Dose, thirty to sixty drops. *Prus. Ph.*

AMMONIATED TINCTURE OF VALERIAN.

R. Valerian, bruised, four ounces.
Aromatic spirit of ammonia, two pints.

Macerate for fourteen days, express, and filter. This may also be made by displacement.

U. S. Ph.

Dose, one to two fl. drachms, in milk, or some mucilaginous fluid.

COMPOUND AMMONIATED TINCTURE OF VALERIAN.

R. Valerian, one ounce.
Cloves, one drachm.
Mace, each, one drachm.
Lemon-peel, one drachm and a half.
Aromatic spirit of ammonia, twenty-four fl. ounces.

Macerate for a week, express, and filter. Dose, a half to one fl. drachm. *Van Mons.*

COMPOUND TINCTURE OF VALERIAN.

R. Valerian, two ounces.
Castor, each, one ounce.
Saffron, one ounce.
Peppermint water, twenty fl. ounces.
Alcohol, each, twenty fl. ounces.

Macerate for a week, express, and filter. Dose, a half to one fl. drachm. *Wirt. Ph.*

WINE OF VALERIAN.

R. Valerian, one ounce.
Sherry wine, one pint.

Macerate for twenty-four hours, express, and filter. A spoonful, as may be required.

Niemann.

SYRUP OF VALERIAN.

R. Bruised valerian, one pound.
Water, eight pints.

Macerate for twelve hours; then distil off one pint and a half; strain, and filter the remainder. Mix with eight pounds of syrup; evaporate to six and a half pounds, and add the distilled fluid. *Par. Cod.*

MIXTURE OF VALERIAN AND AMMONIA.

R. Powdered valerian, one scruple.
Carbonate of ammonia, ten grains.
Cinnamon water, two fl. ounces.

Mix. As a draught, every fourth hour, in a nervous headache, &c. *Ellis.*

MIXTURE OF VALERIAN AND HOFFMANN'S ANODYNE.

R. Valerian, six drachms.
Boiling water, eight fl. ounces.

Make an infusion, strain, and add

Cinnamon water, two fl. ounces.
Hoffmann's anodyne, two fl. drachms.
Syrup, one fl. ounce.

Mix. A spoonful, as required.

Foy.

TINCTURE OF VALERIAN AND HOFFMANN'S ANODYNE.

R. Tincture of valerian, one fl. ounce.
Hoffmann's anodyne, each, one fl. ounce.

Mix. A teaspoonful, as may be required.

Ellis.

OIL OF VALERIAN.

R. Bruised valerian, at will.
Water, sufficient.

Distil, and separate the oil from the product. Dose, three to five drops.

MIXTURE OF OIL OF VALERIAN AND AMMONIA.

R. Oil of valerian, eight drops.
Aromatic spirit of ammonia, one fl. drachm.
Water, four fl. ounces.
Sugar, two drachms.

Mix. A tablespoonful every two or three hours. *Ellis.*

EXTRACT OF VALERIAN.

R. Coarsely-powdered valerian, at will. Moisten with twice its weight of alcohol (.834), let macerate for three days, put into a displacement apparatus, and exhaust with alcohol (.915). The first product, on being distilled over a water-bath, affords a dark-green resin, containing most of the volatile oil, and valerianic acid. The product of the distillation is to be diluted with distilled water, to reduce it to (.935), and the valerian exhausted by additional spirit. Mix the tinctures, filter and distil. The residue is to be evaporated and mixed with the first product. One grain is equal to five of valerian. *Burin.*

FLUID EXTRACT OF VALERIAN.

R. Valerian, in coarse powder, eight ounces.
Ether, four fl. ounces.
Alcohol, twelve fl. ounces.
Diluted alcohol, sufficient.

Mix the ether and alcohol, and, having incorporated the valerian with one-half of the mixture, introduce the mass into a percolator, and

gradually pour in the remainder; then add diluted alcohol until one pint of tincture shall have passed. Place this product in a shallow vessel, and allow it to evaporate spontaneously to five fl. ounces.

Again pour diluted alcohol on the mass in the percolator until ten fl. ounces of tincture are obtained. Mix this with the five fl. ounces of ethereal tincture (adding a little alcohol to dissolve any oleo-resinous matter deposited); allow the mixture to stand, with occasional agitation, for four hours; then filter, and add sufficient alcohol to make the whole measure one pint.

Dose, one fl. drachm.

U. S. Ph.

R. Valerian,	eight ounces.
Alcohol,	eight fl. ounces.
Sulphuric ether,	four fl. ounces.
Water,	sufficient.

Bruise the valerian finely, put it in a displacement apparatus, and add the alcohol and ether, mixed; remove the last portions of the tincture by water, and set the solution aside in a warm room to evaporate, till all the ether and half the alcohol have escaped. The residue will amount to four fl. ounces, of a greenish color. Proceed to displace with the water, setting aside the first eight fl. ounces of the product, and proceed with the process till the root is exhausted. Evaporate this to four fl. ounces; mix all the solutions. The dose is about a teaspoonful.

This is the original formula, on which that of the *U. S. Pharmacopœia* is based.

E. T. Ellis.

VALERIANIC ACID.

R. Coarsely-powdered valerian root,	one pound.
Carbonate of soda,	one ounce,
dissolved in	
Water,	one pint and a half.

Boil for three hours, replacing the water as it evaporates, and express strongly; repeat the process twice with the same quantity of water. Mix the decoctions, add two fl. drachms of sulphuric acid to neutralize the soda, and distil three-fourths of the liquid. Neutralize the distilled liquid with carbonate of soda, and concentrate; then add sulphuric acid to decompose the valerianate of soda, and collect the valerianic acid by distillation. *T. and H. Smith.*

Valerianic acid is now prepared on the large scale, in the process for manufacturing valerianate of soda (*see page 440*), from fusel oil, by the reaction between it, sulphuric acid, and bichromate of potassa.

R. Bichromate of potassa,	two pounds and three quarters.
Hot water,	four pounds and a half.
Mix in a retort, and add, very slowly, a cooled mixture of	

Fusel oil,	one pound.
Sulphuric acid,	four pounds.
Water,	two pounds.

30

Distil. About nine ounces of oily valerianic acid are obtained. *Gruneberg.*

VANILLA.

VANILLA.

Under this name are included the seed pods of several species of *Vanilla*, but that most generally known is the *V. aromatica*, a climbing plant, a native of Mexico and South America.

Sex. Syst. Gynand. monand. *Nat. Syst.* Orchidaceæ.

Swartz, *Occident*, iii. 1518. *Flore Medicale*, vi. 345.

The part used is the pod; this is long, slender, cylindrical, compressed, of a blackish-brown color, formed of two valves, and filled with a brown pulp, containing a great quantity of small, blackish, shining seeds. The odor, is peculiar but pleasant, the taste aromatic, hot, and pungent. It is principally used in this country for flavoring purposes, but is esteemed by European writers as an aromatic stimulant, increasing the power of the muscular system, and exciting the sexual feelings. It has been given in hysteria, rheumatism, impotence, &c. The dose is from eight to twelve grains.

VANILLA SUGAR.

R. Tincture of vanilla,	one ounce.
Sugar,	eight ounces.
Mix well, dry in a stove, and pulverize.	<i>Beral.</i>

VANILLA LOZENGES.

R. Vanilla,	one ounce.
Sugar,	seven ounces.
Gum tragacanth,	half a drachm.
Water,	half an ounce.
Mix, and make lozenges of twelve grains.	<i>Beral.</i>

TINCTURE OF VANILLA.

R. Vanilla,	one part.
Alcohol,	four parts.
Macerate for fourteen days, express, and filter.	
Dose, forty drops.	<i>Cottureau.</i>

SPIRIT OF VANILLA.

R. Vanilla,	one part.
Alcohol,	
Water, each,	twelve parts.
Mix, and distil twelve parts.	<i>Soubiran.</i>

VANILLA ARROWROOT.

R. Milk,	fifteen ounces.
Sugar,	
Vanilla sugar,	
Arrowroot, each,	one ounce.
Mix, and boil, constantly stirring.	<i>Beral</i>

VANILLA MILK.

- R. Milk, sixteen ounces.
 Vanilla sugar, half an ounce.
 Dissolve, and strain. *Beral.*

ESSENCE OF VANILLA.

- R. Vanilla, nine ounces.
 Spirit of abelmoschus
 (ambrette), two pints.
 Cloves, thirty grains.
 Musk, seven grains.

Used as a perfume, and for flavoring. *Cooley.*

FLUID EXTRACT OF VANILLA.

- R. Vanilla, one ounce.
 Sugar, three ounces.
 Syrup,
 Water, each, half a pint.
 Deodorized alcohol, one fl. ounce.

Cut the vanilla in thin, transverse slices, triturate it with the sugar, till reduced to a moderately-fine powder, add the syrup with two ounces of the water, put the mixture into a strong pint-bottle, cork, and tie it over; place it in a vessel of water, which is then to be heated to the boiling point, and kept there for half an hour; remove the cork, and strain. The residue is then to be mixed with the remainder of the water and the alcohol, the mixture returned to the bottle, and heated as before, strained, and mixed with the first product.

W. Procter.

VERATRIA.

VERATRIA.

- R. Cevadilla, bruised, two pounds.
 Alcohol, three gallons.
 Sulphuric acid,
 Solution of ammonia,
 Purified animal charcoal,
 Magnesia, } each,
 sufficient.

Boil the cevadilla in a gallon of the alcohol, in a retort with a receiver attached, for an hour, and pour off the liquor. To the residue, add another gallon of alcohol, together with the distilled portion, boil for an hour, and pour off the liquor; repeat a third time. Express, mix, and strain the liquors, and distil off the alcohol, on a water-bath. Boil the residue three or four times in water, acidulated with sulphuric acid, mix, and strain the liquors, and evaporate to the consistence of syrup. Add magnesia in slight excess, shake frequently, express, and wash residue. Repeat expression and washing two or three times. Dry the residue, digest it with gentle heat, several times in alcohol, straining after each digestion. Distil off alcohol from

the mixed liquors, boil the residue for fifteen minutes in water, with a little sulphuric acid and purified animal charcoal, and strain. Thoroughly wash the residue, mix the washings with the strained liquor, evaporate gently to consistence of syrup, add as much solution of ammonia as will precipitate the veratria, separate this and dry it.

U. S. Ph.

Veratria is a violent and active poison. It is given internally in minute doses, but is more generally used externally, in gout, rheumatism, nervous affections, &c. The dose is from one-twelfth to one-sixth of a grain.

PILLS OF VERATRIA.

- R. Veratria, half a grain.
 Syrup of gum,
 Powdered gum Arabic, each,
 sufficient.

Mix, and make six pills. Dose, one, to be increased to three. *Magendie.*

PILLS OF VERATRIA AND HENBANE.

- R. Veratria, one grain.
 Extract of henbane,
 Powdered liquorice, each,
 twelve grains.

Mix, and make twelve pills. One, thrice a-day. *Turnbull.*

TINCTURE OF VERATRIA.

- R. Veratria, four grains.
 Alcohol, one fl. ounce.

Dissolve. Dose, from ten to twenty drops in a wineglassful of water, in dropsy, gout, rheumatism, &c. *Magendie.*

LOTION OF VERATRIA.

- R. Veratria, one scruple to one drachm.
 Alcohol, two fl. ounces.

Mix. As an embrocation; to be used with caution. *Turnbull.*

LINIMENT OF VERATRIA.

- R. Veratria, eight grains.
 Alcohol,
 Soap liniment, each,
 four fl. drachms.

Mix well. *Brande.*

OINTMENT OF VERATRIA.

- R. Veratria, five to twenty grains.
 Lard, one ounce.

Mix. A piece about the size of a hazelnut to be rubbed for five or fifteen minutes, over the seat of the disease, in rheumatism, &c.

Turnbull.

R. Veratria, one scruple.
 Rub with
 Olive oil, one drachm.
 Add
 Spermaceti ointment, seven drachms.
 Mix. As the last. *Turnbull.*

OINTMENT OF VERATRIA AND OPIUM.

R. Veratria, half a drachm.
 Powdered opium, one drachm.
 Lard, one ounce and a half.
 Mix. As a friction, in rheumatism. *Dunghison.*

OINTMENT OF VERATRIA AND IODINE.

R. Veratria, one scruple.
 Iodide of potassium, half a drachm.
 Lard, one ounce.
 Mix. *Jourdan.*

VERATRIÆ MURIAS.

MURIATE OF VERATRIA.

R. Veratria, at will.
 Muriatic acid, sufficient
 to saturate; filter, evaporate, and crystallize. *Giordano.*

VERATRIÆ NITRAS.

NITRATE OF VERATRIA.

VERATRIÆ SULPHAS.

SULPHATE OF VERATRIA.

Are made in the same manner, using the appropriate acids. Dose of each, about one-eighth of a grain.

SOLUTION OF SULPHATE OF VERATRIA.

R. Sulphate of veratria, one grain.
 Distilled water, two fl. ounces.
 Dissolve. Dose, one to two fl. drachms, in a mixture, as a substitute for the wine of colchicum or the *Eau medicinale*. *Cadet.*

VERATRUM ALBUM.

WHITE HELLEBORE.

A perennial, herbaceous plant, with a fleshy, fusiform rhizome, having numerous fibres. It is a native of the mountainous parts of Europe. *Sex. Syst.* Polygam. monœc. *Nat. Syst.* Melanthaceæ.

Linn. Sp. Pl. 1479. Griffith, Med. Bot. 643. The root, which is the officinal part, is in cylindrical, somewhat conical pieces, with numerous radicles; these are corrugated, and of a

blackish-brown color. Their odor is slight, and their taste at first sweetish, and then bitter and acrid. White hellebore is an active irritant, causing violent vomiting and purging. Externally, it acts like the other powerful acids. Internally, in small doses, it has been used in many diseases; but it is now principally employed externally in the treatment of cutaneous affections, &c. Dose, from a grain to two grains, gradually increased.

POWDER OF WHITE HELLEBORE.

R. Powdered white hellebore, three grains.
 " starch, one scruple.
 Mix. As an errhine in amaurosis, and lethargic affections. *Radius.*

DECOCTION OF WHITE HELLEBORE.

R. White hellebore, ten drachms.
 Distilled water, two pints.
 Boil down to a pint, and add
 Rectified spirit, three fl. ounces,
 and strain. *Lond. Ph., 1836.*
 As a wash, in cutaneous diseases.

WINE OF WHITE HELLEBORE.

R. Bruised white hellebore, four ounces.
 Wine, one pint.
 Macerate for fourteen days, express, and filter. *U. S. Ph.*
 Dose, ten minims, gradually increased, two or three times a-day, in gout and rheumatism.

TINCTURE OF WHITE HELLEBORE.

R. Bruised white hellebore, four ounces.
 Diluted alcohol, one pint.
 Macerate for eight days, express, and filter. *Ed. Ph.*
 Dose, the same as of the wine, and in same cases.

OINTMENT OF WHITE HELLEBORE.

R. Powdered white hellebore, two ounces.
 Lard, eight ounces.
 Oil of lemon, twenty minims.
 Mix. *U. S. Ph.*
 As a friction in itch, &c. To be used with caution on children.

COMPOUND OINTMENT OF WHITE HELLEBORE.

R. Powdered white hellebore, two ounces.
 " nitre, one drachm.
 Soft soap, six ounces
 Sulphur, each, one pound and a half.
 Lard, one pound and a half.
 Mix. As a friction in itch. *Phæbus.*

VERATRUM VIRIDE.

AMERICAN OR GREEN HELLEBORE.

A tall, perennial plant, with a thick, fleshy rhizome, having a tunicated top, and numerous fibrous radicles. It is a native of the United States, growing in damp meadows, and wet situations.

Sex. Syst. Polygam. monœc. *Nat. Syst.* Melanthaceæ.

Aiton, Hort. Kew. iii. 422. Griffith, Med. Bot. 643.

The root, in a fresh state, has an unpleasant odor, which disappears on drying. Its taste is bitter and acrid. Its properties are much the same as those of the white hellebore, but is stated not to purge, and to rapidly reduce the frequency and force of the pulse. Dose, one grain, gradually increasing.

TINCTURE OF AMERICAN HELLEBORE.

R. American hellebore, bruised,
six ounces.

Diluted alcohol, one pint.

Macerate for fourteen days, express, and filter.
Dose, half to one fl. drachm. *Osgood.*

R. Dried root of American hellebore,
in powder, eight ounces.
Alcohol (.835), sixteen ounces.

Macerate for two weeks, and express. Dose, for an adult, eight drops, every three hours. Each dose to be increased one or two drops, until nausea, vomiting, or a reduction of the frequency of the pulse takes place; then to be reduced to one-half in all cases. *Norwood.*

WINE OF AMERICAN HELLEBORE.

R. American hellebore, bruised,
six ounces.

White wine, fourteen fl. ounces.

Diluted alcohol, two ounces.

Macerate for fourteen days, express, and filter.
Dose, as of tincture. *Osgood.*

The recent root should be used in the two preparations of Dr. Osgood.

EXTRACT OF AMERICAN HELLEBORE.

R. Fresh root of American hellebore,
at will.

Bruise well, express the juice, and evaporate to proper consistence, by a gentle heat, or by exposure to the sun. Dose, from a fourth to half a grain. *Osgood.*

OINTMENT OF AMERICAN HELLEBORE.

R. Powdered extract of American
hellebore, one drachm.
Simple cerate, one ounce.
Oil of lemon, three minims.

Mix. *Osgood.*

PILLS OF AMERICAN HELLEBORE.

R. Extract of American hellebore,
ten grains.
Opium, five grains.
Soap, fifteen grains.
Mucilage of gum Arabic, sufficient.
Mix, and make thirty pills. One, every three
or four hours, in rheumatism, &c. *Osgood.*

MIXTURE OF AMERICAN HELLEBORE.

R. Tincture of American hellebore,
“ bloodroot,
Camphorated tincture of opium,
equal parts.
Mix. Dose, fifty to sixty minims, in catarrh.
Osgood.

VIOLA.

VIOLET.

Two species of violet, viz: *V. pedata* and *V. odorata*, are sometimes used in medicine. The former is indigenous to this country, and is official in the U. S. Pharm. The latter is a small, perennial plant, with fragrant, blue flowers, a native of Europe, but generally cultivated in our gardens.

Sex. Syst. Pentand. monog. *Nat. Syst.* Violaceæ.

Linn. Sp. Pl. 1324. Griffith, Med. Bot. 140.

The flowers of both species are of a blue color, almost inodorous when dried, and of a slightly bitter taste. They are principally used as a demulcent, and as a laxative for infants.

CONSERVE OF VIOLETS.

R. Violet flowers, fresh, one part.
Sugar, three parts.
Rub together. *Soubiran.*

SYRUP OF VIOLETS.

R. Fresh violet flowers, one pound.
Boiling water, two pints and a half.
Infuse for twenty-four hours, strain, and dissolve
Sugar, seven pounds and a half,
and form syrup. *Ed. Ph.*
A gentle laxative for infants, in doses of one
to two fl. drachms.

HONEY OF VIOLETS.

R. Flowers of violets,
Boiling water, each, six pounds.
Infuse, strain, and add
Honey, six pounds.
Evaporate to a proper consistence. *Wirt. Ph.*

OIL OF VIOLETS.

R. Fresh violet flowers, fifteen ounces.
Almond oil, five pounds.
Infuse, with three equal quantities of the flowers,
and strain. Used as an article of perfumery.
Taddei.

W.

WINTERA.

WINTER'S BARK.

This is derived from an evergreen tree, found in the southern parts of South America, the *Drimys Winteri*, and also from other species, growing in Peru, &c.

Sex. Syst. Polyand. tetragyn. *Nat. Syst.* Magnoliaceæ.

Forster, Gen. 84. Griffith, Med. Bot. 100.

The bark is in quilled pieces of some length; of a pale, yellowish, or dull, reddish gray, with darker spots externally, and of a dark cinnamon

internally. It has an aromatic odor, and a warm, spicy taste. It is a stimulating, aromatic tonic. The dose, in powder, is about half a drachm.

TINCTURE OF WINTER'S BARK.

R. Bruised Winter's bark, one part.
Diluted alcohol, eight parts.

Digest with one-half of the alcohol, and repeat with the other. Unite the two tinctures, and filter.

X.

XANTHORRHIZA.

YELLOW-ROOT.

The root of *Xanthorrhiza apiifolia*, a plant indigenous to the United States, is occasionally employed in medicine.

Sex. Syst. Pentand. polyg. *Nat. Syst.* Ranunculaceæ.

Willd. Sp. Pl. i. 1568. Griffith, Med. Bot. 95.

The root and bark of this little shrub are possessed of tonic properties, and may be used in cases where the pure bitters are indicated.

INFUSION OF YELLOW-ROOT.

R. Bruised yellow-root, one ounce.
Boiling water, one pint.

Macerate in a covered vessel for two hours, and strain. Dose, as a tonic, one fl. ounce. *Thomas.*

EXTRACT OF YELLOW-ROOT.

R. Coarsely-powdered yellow-root, eight ounces.
Diluted alcohol, two pints.

Moisten the powder with six fl. ounces of diluted alcohol, and macerate twenty-four hours; then transfer to a percolator, and add gradually the rest of the diluted alcohol. When this has

passed, continue the percolation with water until the powder is exhausted. Distil off the alcohol, and evaporate to the consistence of an extract. *Thomas.*

Dose, five grains.

XANTHOXYLUM.

PRICKLY-ASH.

The bark of *Xanthoxylum fraxineum*, a shrub about ten feet high, growing in the Middle and Western States, is recognised in the U. S. Pharm.

Sex. Syst. Diœc. pentand. *Nat. Syst.* Xanthoxylaceæ.

The bark is possessed of stimulant and irritant properties. It is more used in domestic, than in regular practice.

DECOCTION OF PRICKLY-ASH.

R. Contused bark of prickly-ash, one ounce.
Water, three pints.

Boil down to a quart.

One pint to be taken in divided doses, during the twenty-four hours, in chronic rheumatism.

G. B. Wood.

Z.

ZINCUM.

ZINC.

ZINCI OXIDUM.

OXIDE OF ZINC.

- R. Sulphate of zinc, one pound.
 Carbonate of ammonia, six ounces and a half.
 Distilled water, three gallons.

Dissolve the sulphate of zinc and carbonate of ammonia, separately, in twelve pints of the water, strain, and mix the solutions. Wash the precipitate well with water, and expose to a strong heat, to drive off the carbonic acid.

U. S. Ph., 1840.

- R. Precipitated carbonate of zinc, one pound.

Expose to a strong heat in a shallow vessel, so as to drive off the carbonic acid.

U. S. Ph., 1850.

Tonic and antispasmodic, and given in nervous and spasmodic affections, in doses of two to ten grains and upwards, several times a-day. Also used externally.

POWDER OF OXIDE OF ZINC.

- R. Oxide of zinc, three grains.
 White sugar, five grains.

Mix. One powder every four hours. Useful in gastric or spasmodic cough. *A. T. Thomson.*

COMPOUND POWDER OF OXIDE OF ZINC.

- R. Oxide of zinc, six grains.
 Extract of henbane, one grain.
 Oil of valerian, one drop.
 Powdered valerian, half a drachm.

Mix, and divide into nine powders. One, thrice a-day, in nervous disorders. *Radiis.*

- R. Oxide of zinc, one drachm.
 Powdered opium, ten grains.
 " Peruvian bark, " fennel, each, one ounce.

Mix, and divide into fifty powders. One, every two hours, in whooping-cough. *Augustin.*

POWDER OF OXIDE OF ZINC AND COLOMBO.

- R. Oxide of zinc, four grains.
 Powdered colombo, thirty grains.

Mix, and divide into four powders. One, every three hours, in dyspepsia. *Brera*

PILLS OF OXIDE OF ZINC.

- R. Oxide of zinc, two scruples.
 Conserve of roses, sufficient.

Mix, and make ten pills. One, three or four times a-day, in epilepsy, chorea, &c. *Ellis.*

COMPOUND PILLS OF OXIDE OF ZINC.

- R. Oxide of zinc, five grains.
 Extract of cascarilla, six grains.
 Syrup of ginger, sufficient.

Mix, and make three pills; one to be taken three times a-day, in epilepsy. *Ainslie.*

- R. Oxide of zinc, one scruple.
 Powdered valerian, thirty grains.
 Castor, four grains.
 Syrup, sufficient.

Mix, and make twelve pills. To be taken during the day, in epilepsy, &c. *Foy.*

LOTION WITH OXIDE OF ZINC.

- R. Oxide of zinc, one ounce.
 Elder-flower water, one fl. ounce and a half.

Agitate together. In pustular erysipelas. *Augustin.*

OINTMENT OF OXIDE OF ZINC.

- R. Oxide of zinc, one ounce.
 Lard, six ounces.
 Mix. *U. S. Ph.*

A mild, astringent application, in chronic ophthalmia, and in excoriation and ulceration.

OINTMENT OF IMPURE OXIDE OF ZINC.
(TUTTY OINTMENT.)

- R. Prepared tutty, one drachm.
 Lard, five drachms.
 Rub together. *Pereira.*

OINTMENT OF OXIDE OF ZINC AND OPIUM.

- R. Oxide of zinc, half a drachm.
 Fresh butter, one drachm.
 Powdered opium, one grain.
 Mix. In chronic ophthalmia. *Hencke.*

OINTMENT OF OXIDE OF ZINC AND
CALOMEL.

- R. Oxide of zinc, half a drachm.
Calomel,
Powdered aloes, each, one grain.
Fresh butter, two drachms.
Mix. In scrofulous ophthalmia. *Scarpa.*

CERATE OF OXIDE OF ZINC AND
LYCOPodium.

- R. Oxide of zinc,
Lycopodium, each, fifteen grains.
Simple cerate, half an ounce.
Mix. In ulceration of the breasts, tetter, &c. *Hufeland.*

ZINCI CHLORIDUM.
CHLORIDE OF ZINC.

- R. Zinc, in small
pieces, two ounces and a half.
Nitric acid,
Prepared chalk, each, one drachm.
Muriatic acid, sufficient.

Add the zinc and muriatic acid together, in a glass vessel, and dissolve; strain, add the nitric acid, and evaporate to dryness. Dissolve in water, add the chalk, let rest for twenty-four hours, filter, and again evaporate. *U. S. Ph.*

- R. Solution of chloride of barium, at will.
Solution of sulphate of zinc, sufficient
to precipitate sulphate of baryta. Filter and
evaporate the fluid to obtain crystals of chloride
of zinc. *Righini.*

Principally used externally, as a caustic and
stimulant; also given internally, in very small
doses, in the neuroses, &c.

CHLORIDE OF ZINC. (*Disinfectant.*)

- R. Granulated zinc, four pounds (av.)
Muriatic acid, four pounds, or sufficient.
Water, two gallons, two pints.

Add the acid gradually to the zinc, in a glass
or porcelain vessel, till it is dissolved, taking
care there is no excess of acid; strain through
a coarse cloth, and add the water. This con-
tains one part of the chloride in twelve, and is
of the strength for a disinfectant. *E. Parrish.*

ETHEREAL TINCTURE OF CHLORIDE OF
ZINC.

- R. Chloride of zinc, four drachms.
Alcohol, one fl. ounce.
Sulphuric ether, two fl. ounces.
Mix. Five drops every four hours, in a little
sugar and water, gradually increasing the dose,
in chorea, epilepsy, &c. *Hufeland.*

LOTION OF CHLORIDE OF ZINC.

- R. Chloride of zinc, eight grains.
Aloes, two scruples.
Distilled water, four fl. ounces.
Mix. As an application to atonic and foul
ulcers. *Vogt*

INJECTION OF CHLORIDE OF ZINC.

- R. Liquid chloride
of zinc, twenty-four drops.
Water, four fl. ounces.
Mix, and filter. A small quantity to be injected
a short distance into the urethra, two or three
times a-day, in gleet. *Gaudriot.*

VAGINAL SUPPOSITORY OF CHLORIDE OF
ZINC.

- R. Liquid chloride of zinc, five drops.
Sulphate of morphia, half a grain.
Mix with three drachms of the following paste:
Mucilage of tragacanth, six parts.
White sugar, three parts.
Starch, nine parts.
Make a suppository. To be used every day, or
every other day. *Guadriot.*

CANQUOIN'S PASTE OF CHLORIDE OF
ZINC.
No. 1.

- R. Chloride of zinc, one part.
Wheat flour, or sulphate
of lime, four parts.
Water, sufficient
to make a paste. Mix.

No. 2.

- R. Chloride of zinc, one part.
Wheat flour, three parts.
Water, sufficient.
Mix.

No. 3.

- R. Chloride of zinc, one part.
Wheat flour, two parts.
Water, sufficient.
Mix.

The chloride and flour are to be carefully and
quickly mixed; add the water to half the quan-
tity, so as to form a soft paste, and mix with this
the rest of the powder. Form into cakes of
half a line to four lines in thickness. To remain
on twenty-four hours, or more (six to ten are
often sufficient). To produce an eschar in can-
cer, lupus, &c. *Canquoin.*

- R. Chloride of zinc, one part.
Chloride of antimony, half a part.
Flour, two and a half parts.
Water, sufficient.

To be mixed as above, so as to be moulded into any desired form. As a caustic, in nodulated cancerous tumors. *Canquoin.*

Powdered opium may be added to any of the above, to mitigate the pain.

ZINCI CYANURETUM.

CYANURET OF ZINC.

R. Sulphate of zinc, one part.
Distilled water, ten parts.

Dissolve; add

Solution of cyanuret of potassium, sufficient to form precipitate, being added by drops. Filter, wash the precipitate, and dry. *Kunze.*

Used in spasmodic affections, &c., in doses of a twelfth to a quarter of a grain.

POWDER OF CYANURET OF ZINC.

R. Cyanuret of zinc, three grains.
Magnesia, twenty-four grains.
Powdered cinnamon, twelve grains.

Mix, and divide into twelve powders. One, every four hours, in nervous affections, &c. *Guibourt.*

PILLS OF CYANURET OF ZINC.

R. Cyanuret of zinc, fifteen grains.
Extract of liquorice, two drachms.

Mix, and make sixty pills. One, three times a-day. *Kopp.*

ZINCI FERROCYANURETUM.

FERROCYANURET OF ZINC.

R. Ferrocyanuret of potassium, sixty parts.

Sulphate of zinc, eighty-three parts.

Dissolve separately in boiling water, add the solutions together; collect the precipitate of ferrocyanuret of zinc, wash it well, and dry. *Schindler.*

Given in doses of one grain, twice or thrice a-day, gradually increasing the quantity, in nervous affections, nocturnal pains, &c.

POWDER OF FERROCYANURET OF ZINC.

R. Ferrocyanuret of zinc, six grains.
Magnesia, four grains.
Powdered cinnamon, three grains.

Mix. Every four hours, in nervous affections, cramp of the stomach, &c. *Radius.*

The quantity of the salt of zinc is too large, but it is thus ordered in various authors.

COMPOUND PILLS OF FERROCYANURET OF ZINC.

R. Ferrocyanuret of zinc, fifteen grains.
Powdered valerian, half a drachm.
Extract of valerian, sufficient.

Mix, and make sixty pills. Two, morning and evening, in chorea. *Rosensteil.*

R. Ferrocyanuret of zinc, ten grains.
Magnesia, two scruples.
Oil of valerian, twenty drops.
Syrup of chamomile, five scruples.
Powdered valerian, ten scruples.

Mix, and make into three-grain pills. Six, three times a-day, gradually increasing. *Vogt.*

SOLUTION OF FERROCYANURET OF ZINC.

R. Ferrocyanuret of zinc, four grains.
Distilled water, two fl. ounces.

Mix. Dose, a tablespoonful, four times a-day, in chorea. *Radius.*

MIXTURE OF FERROCYANURET OF ZINC.

R. Ferrocyanuret of zinc, eight to twenty grains.

Powdered gum Arabic, two drachms.

Cherry water, three fl. ounces.

Wine of opium, one fl. scruple.

Cherry-laurel water, half a fl. ounce.

Mix. As a collyrium, in scrofulous and rheumatic ophthalmia. *Ammon.*

ZINCI IODIDUM.

IODIDE OF ZINC.

R. Iodine, one hundred and seventy parts.
Powdered zinc, twenty parts.

Heat in a matrass till the iodide sublimes. *Magendie.*

R. Iodine, two parts.
Powdered zinc, one part.

Digest in a little water, and agitate till the solution is colorless. Evaporate with a gentle heat. *Beasley.*

Used externally.

OINTMENT OF IODIDE OF ZINC.

R. Iodide of zinc, one drachm.
Lard, one ounce.

Mix. As a friction to tumors, &c. *Ure.*

COLLYRIUM OF IODIDE OF ZINC.

R. Iodide of zinc, four grains.
Distilled water, six fl. ounces.

Dissolve.

SOLUTION OF IODIDE OF ZINC.

R. Iodide of zinc, ten to thirty grains.
Distilled water, one fl. ounce.

Dissolve. To be applied, by means of a sponge tied to a quill, to enlarged tonsils; to be followed, after some days' perseverance, by a saturated solution applied by a camel's hair brush. *T. Ross.*

SYRUP OF IODIDE OF ZINC.

R. Iodine, four drachms.
Powdered zinc, two drachms.
Water, four fl. ounces.

Agitate till colorless; then filter into

Syrup, twelve fl. ounces.

Mix. *A. T. Thomson.*

R. Iodine, twelve drachms, two scruples.
Finely granulated zinc, one ounce.
Sugar, (avoir.) one pound.
Water, nine fl. ounces.

Add the zinc and iodine to the water in an evaporating dish, and heat gently until the solution becomes colorless. Place the sugar in a wide-mouthed bottle, filter the hot solution into it, and add sufficient water to make a pint of syrup. Dissolve the sugar by a gentle heat.

Each fl. ounce contains one drachm of iodide of zinc. *A. B. Taylor.*

ZINCI ACETAS.

ACETATE OF ZINC.

R. Acetate of lead, one pound.
Granulated zinc, nine ounces.
Distilled water, three pints.

Dissolve the acetate in the water, and filter. Add the zinc, and agitate occasionally, in a stopped bottle, till the liquid gives no precipitate with a solution of iodide of potassium. Filter, evaporate by a gentle heat to one-fifth, acidulate the solution by acetic acid, and set aside to crystallize. Decant, and dry the crystals on bibulous paper. If they are colored, dissolve in distilled water, heat, and drop into the hot solution precipitated carbonate of zinc, in successive portions, until a small quantity of the liquid passes colorless on filtration; filter, acidulate with a few drops of acetic acid, evaporate, and crystallize. *U. S. Ph.*

Used as an external application.

COLLYRIUM OF ACETATE OF ZINC.

R. Sulphate of zinc,
Acetate of lead, each, six grains.
Rose water, four fl. ounces.

Mix, and filter. *Ellis.*

R. Acetate of zinc, fifteen to thirty grains.
Distilled water, twelve fl. ounces.
Dissolve. *Ware.*

R. Sulphate of zinc, one drachm.
Acetate of lead, half a drachm.
Camphor, one scruple.
Rose water, twelve fl. ounces.
Mix, and filter the solution. *Spielmann.*

INJECTION OF ACETATE OF ZINC.

R. Acetate of zinc, eight grains.
Rose water, four fl. ounces.
Dissolve. In gonorrhœa. *Ellis.*

LOTION OF ACETATE OF ZINC.

R. Acetate of zinc, four scruples.
Distilled vinegar, sixteen fl. ounces.
Dissolve. *Beral.*

TINCTURE OF ACETATE OF ZINC.

R. Sulphate of zinc,
Acetate of potassa, each, one part.
Rub together, and add
Diluted alcohol, sixteen parts.
Macerate for a week, stirring occasionally, and filter. *Dub. Ph., 1826.*
As a collyrium and injection, properly diluted.

ZINCI CARBONAS.

CARBONATE OF ZINC.

PRECIPITATED CARBONATE OF ZINC.

R. Sulphate of zinc,
Carbonate of soda, each, one pound.
Boiling water, one gallon.

Dissolve the salts separately in four pints of water, and mix the solutions; wash the precipitated carbonate, first by decantation, and afterwards on a filter; then press, and dry it. *U. S. Ph.*

CERATE OF CARBONATE OF ZINC.

R. Precipitated carbonate of zinc, two drachms.
Simple ointment, ten drachms.
Mix them. *U. S. Ph., 1850*
R. Prepared carbonate of zinc,
Yellow wax, each, half a pound.
Lard, two pounds.

Melt the lard and wax together, and when, on cooling, they begin to thicken, add the carbonate of zinc, and stir till cold. This latter cerate is now official under the name of ceratum calaminæ. (See page 153.)

U. S. Ph., 1840

Much used as a dressing to excoriations, shallow ulcerations, &c.

PLASTER OF CARBONATE OF ZINC.

- R. Simple plaster,
Powdered carbonate of zinc, each,
two pounds.
Yellow wax, five ounces.
Suet,
Turpentine, each,
one ounce and a half.
Melt, and mix thoroughly. *Van Mons.*

- R. Prepared carbonate of zinc,
White lead, each,
one ounce and a half.
Yellow wax, four ounces.
Olive oil, six ounces.
Melt the wax with the oil, and add the two powders, stirring well. To diminish a superabundant suppuration in ulcers. *Cadet.*

ZINCI SULPHAS.

SULPHATE OF ZINC.

- R. Zinc, in small pieces, four ounces.
Sulphuric acid, six ounces.
Distilled water, four pints.

Introduce the zinc and water into a glass vessel, and add the sulphuric acid by degrees; when all effervescence has ceased, filter, boil to a pellicle, and set aside to crystallize. *U. S. Ph.*

It is astringent, tonic, and, in large doses, a prompt and effective emetic. Dose, as a tonic, one to two grains; as emetic, ten to thirty.

PILLS OF SULPHATE OF ZINC.

- R. Sulphate of zinc, ten grains.
Conserve of roses, sufficient.
Mix, and make twenty pills. One, three times a-day. In hooping-cough, &c. *Ellis.*

PILLS OF SULPHATE OF ZINC AND TURPENTINE.

- R. Sulphate of zinc, two drachms.
Venice turpentine, sufficient.
Mix, and make sixty pills. As a tonic. *Beasley.*

PILLS OF SULPHATE OF ZINC AND MYRRH.

- R. Sulphate of zinc, ten grains.
Powdered myrrh,
one drachm and a half.
Conserve of roses, sufficient.
Mix, and make twenty pills. From two to four pills a-day, in hooping cough. *Paris.*

COMPOUND PILLS OF SULPHATE OF ZINC.

- R. Sulphate of zinc, one grain.
Extract of gentian, eight grains.
Syrup of ginger, sufficient.
Mix, and make two pills, to be taken during the day, with two fl. drachms of infusion of chamomile, in epilepsy. *Ainslie.*

SOLUTION OF SULPHATE OF ZINC.

- R. Sulphate of zinc, two grains.
Chamomile water, three fl. ounces.
Dissolve. A spoonful, two or three times a-day, in chronic dysentery. *Radius.*

COLLYRIUM OF SULPHATE OF ZINC.

- R. Sulphate of zinc, six grains.
Acetate of lead, ten grains.
Laudanum, one fl. drachm.
Water, five fl. ounces.
Mix. As an application to the eyes in ophthalmia, after reduction of the inflammation. *Gregory.*

- R. Sulphate of zinc, six grains.
Mucilage of quince seed, one fl. ounce.
Distilled water, six fl. ounces.
Spirit of camphor, five drops.
Mix, and strain. *Scarpa*
To be used as the last.

COLLYRIUM OF SULPHATE OF ZINC AND CAMPHOR.

- R. Sulphate of zinc, one scruple.
Camphor water,
Rose water, each, eight fl. ounces.
Mix. *Ainslie.*

FOMENTATION WITH SULPHATE OF ZINC.

- R. Sulphate of zinc,
one drachm and a half.
Decoction of pomegranate bark,
four pints.
" Peruvian bark, two pints.
Laudanum, one fl. ounce.
Mix. As an application to external piles, and in vaginal gonorrhœa. *Brera.*

GARGLE OF SULPHATE OF ZINC.

- R. Sulphate of zinc, one drachm.
Honey, half a fl. ounce.
Tincture of myrrh,
Brandy, each, one fl. ounce.
Rose water, six fl. ounces.
Mix. In ulcerations of gums, &c., occasioned by excessive salivation. *Ellis.*

INJECTION OF SULPHATE OF ZINC.

- R. Sulphate of zinc, one drachm.
Camphor water, two fl. ounces.
Water, two pints.
Mix. *Swediaur.*

- R. Sulphate of zinc, ten grains.
Powdered gum Arabic, two drachms.
Laudanum, one fl. drachm.
Distilled water, eight fl. ounces.
Mix. In gonorrhœa. *Ellis.*

LOTION OF SULPHATE OF ZINC.

- R. Sulphate of zinc, sixteen grains.
Rose water, sixteen fl. ounces.
Mix. *Beral.*

- R. Sulphate of zinc, half a drachm.
Water, half a pint.
Lead water, each, half a pint.
Mix. *Beasley.*

- R. Sulphate of zinc, four scruples.
Distilled vinegar, one pint.
Mix. *Beral.*

SULPHATE OF ZINC MIXTURE.

- R. Sulphate of zinc, one grain.
Decoction of cascarilla, twelve fl. drachms.
Simple syrup, four fl. drachms.
Mix, to be divided into four doses, one to be taken three times a-day, in hooping-cough. *Ainslie.*

ZINCI VALERIANAS.

VALERIANATE OF ZINC.

- R. Fresh valerian root, one hundred parts.
Water, five hundred parts.
Sulphuric acid, ten parts.
Bichromate of potassa, six parts.

Macerate the coarsely-powdered valerian in the water, to which have been added the sulphuric acid and the bichromate, for twenty-four hours; then introduce the mixture into a still, and apply heat, returning the first fourth of the product to the still; the process is then to be continued, until the distilled water ceases to redden litmus paper. The product is then to be placed in a large matrass, and an excess of pure hydrated carbonate of zinc is to be added, and allowed to digest at 190° F., for three or four hours, on a sand-bath, until saturated. The solution is then to be filtered and evaporated over a naked fire, till reduced to fifty parts, then placed on plates, and suffered to evaporate in a drying-room, or by a moderate temperature. *Lefort.*

- R. Contused valerian, thirty-two ounces.
Water, eight pounds.
Sulphuric acid, three ounces.

Macerate for two days, and distil as long as the product reddens litmus paper. Expose to the air for a month, then put it in a matrass with two hundred and twenty-five grains of recently-prepared oxide of zinc, and digest on a sand-bath for eight or ten hours, at 175° F., agitating occasionally; filter whilst hot, evaporate to three-fourths, and dry the residue on earthen plates. *Buisson.*

- R. Valerianate of soda, two ounces and a half.
Sulphate of zinc, two ounces and seven drachms.
Water, two pints.

Dissolve each salt in one-half of the water, and raise both solutions to the temperature of 200° F.; then mix them, and skim off the crystals as they are formed. A farther quantity of crystals can be obtained by concentrating the mixture to four fl. ounces. Wash the crystals with cold water to separate the sulphate of soda, then drain on a filter, and dry. *Dub. Ph.*

Has been recommended as a powerful tonic and antispasmodic, in neuralgia and hysteria, but has not been sufficiently tried to ascertain its true powers. Dose, one to two grains.

PILLS OF VALERIANATE OF ZINC.

- R. Valerianate of zinc, nine grains.
Tragacanth, thirty grains.
Mix, and make twelve pills. One, morning and evening, in nervous headache, &c. *Devey.*

MIXTURE OF VALERIANATE OF ZINC.

- R. Valerianate of zinc, one grain and a half.
Distilled water, five fl. drachms.
Syrup, one fl. ounce.
Mix. A tablespoonful every half hour. *Devey.*

ZINGIBER.

GINGER.

The rhizome of *Zingiber officinale*, a small herbaceous, perennial plant, a native of the East Indies.

Sex. Syst. Monand. monog. *Nat. Syst.* Zingiberaceæ.

Roscoe, Trans. Linn. Soc. viii. 348. Griffith, Med. Bot. 629.

The rhizome, or root, is creeping, knotty, lobated or palmated, and fleshy; with an agreeable, pungent, camphoraceous odor, and an aromatic, acrid, burning taste. Two kinds are met

with in commerce, the black and the white; the first consists of the roots, scalded in hot water and dried; the second is also composed of the roots, but deprived of their epidermis previous to desiccation. Ginger contains a volatile oil, a soft, acrid resin, some gum, various salts, and the usual vegetable constituents. It is stimulating, somewhat acrid, aromatic, and stomachic. It is given in doses of ten grains to a scruple, or more.

INFUSION OF GINGER.

R. Bruised ginger, half an ounce.
Boiling water, one pint.
Macerate for two hours in a covered vessel, and strain.

Dose, a wineglassful.

U. S. Ph.

GINGER LOZENGES.

R. Powdered ginger, two ounces.
" sugar, fourteen ounces.
Gum tragacanth,
Water, each, twelve drachms.
Mix well, and form lozenges of sixteen grains each.

Soubéiran.

GINGER BEER.

R. Sugar, three pounds.
Bruised ginger, two ounces.
Cream of tartar, one ounce.
Lemons, sliced, four.
Yeast, eight fl. ounces.
Boiling water, four gallons.

Pour the water on the four first-named ingredients, infuse for two hours, and strain; then add the yeast, and, when fermentation has commenced, pour into strong bottles, and wire down the corks.

Redwood.

R. White sugar, twenty pounds.
Lemon juice, eighteen fl. ounces.
Honey, one pound.
Bruised ginger, twenty-two ounces.
Water, eighteen gallons.

Boil the ginger in three gallons of the water for half an hour; add the sugar, lemon juice, and honey, with the remainder of the water, and strain; when cold, add the white of an egg, and half a fl. ounce of essence of lemon; let rest for four days, and bottle.

Pereira.

SYRUP OF GINGER.

R. Tincture of ginger, four fl. ounces.
Water, four pints.
Sugar, ten pounds.

Mix the tincture with four pounds of sugar, and expose the mixture, in a shallow dish, to a gentle heat, until the alcohol has evaporated. Add the residue of the sugar, and then the water, in a covered vessel; heat gradually, till the sugar is dissolved, and strain.

U. S. Ph.

Mr. Joseph Laidley proposes, as an improvement, the addition of the whites of two or three eggs, beat up with a small quantity of water, to the syrup. The whole is then to be boiled for two minutes, and strained through flannel. By this method the syrup is rendered beautifully clear.

J. Laidley.

R. Tincture of ginger, two fl. ounces.
Carbonate of magnesia, two drachms.
Sugar, (avoir), one pound and a half.
Water, twelve fl. ounces.

Rub the tincture with the carbonate of magnesia and two ounces of the sugar, in powder, gradually add the water, and filter. The remainder of the sugar is then dissolved in the filtered liquid by means of a gentle heat.

J. D. Finley.

TINCTURE OF GINGER.

R. Ginger, bruised, eight ounces.
Alcohol, two pints.

Macerate for fourteen days, express, and filter through paper. This tincture may also be prepared by displacement.

U. S. Ph.

Dose, half to one fl. drachm.

OIL OF GINGER.

R. Ginger, one part.
Water, six parts.

Digest for five or six days; distil, and collect the oil.

ETHEREAL EXTRACT OF GINGER.

R. Ginger, four ounces.
Sulphuric ether, six ounces.

Treat the powdered ginger with the ether, in a displacement apparatus, and evaporate the product by means of a water-bath. One part corresponds to sixteen of ginger.

Beral.

TINCTURE OF ETHEREAL EXTRACT OF GINGER.

R. Ethereal extract of ginger, one part.
Alcohol (.825), fifteen parts.
Dissolve.

Beral.

SYRUP OF ETHEREAL EXTRACT OF GINGER.

R. Ethereal extract of ginger, one scrup.
Syrup, one ounce.

Dissolve, and strain, after twenty-four hours' digestion.

Beral.

ESSENCE OF GINGER.

R. Jamaica ginger, four ounces.
French brandy, one pint.

Powder the ginger coarsely, moisten it with the brandy, put it in a displacer, and pour the remainder of the brandy on it; when it ceases to pass, add diluted alcohol, till one pint has passed in all.

R. Bruised Jamaica

ginger, twelve pounds.
Alcohol, two gallons and a half.

Digest fourteen days, express, strain, and reduce by distillation, to one gallon. Cool, and filter. A most excellent preparation. *Cooley.*

R. Powdered ginger,
" animal char-

coal, equal parts.
Alcohol, sufficient.

Moisten the powders for twenty-four hours, with sufficient alcohol to cover them; then

transfer to a percolator, and return the first runnings two or three times. Change the receiver, and gradually pour on alcohol, as required, until as much essence is obtained as there was ginger employed. *Cooley.*

SPICE PLASTER.

R. Powdered ginger, two ounces.

" cloves,

" cinnamon, each, one ounce.

" red pepper, two drachms.

Tincture of ginger, half an ounce.

Honey, sufficient.

Mix the powders; add the tincture, and sufficient honey to make of proper consistence for a stiff cataplasm. *W. Procter.*

ADDENDA.

Under this title will be found a few articles belonging to the Formulary proper, together with a large number of practical receipts, which could not have been introduced, with propriety, under any of the officinal heads.

FLEMING'S TINCTURE OF ACONITE.

R. Powdered aconite root,
(troy) sixteen ounces.

Rectified spirit, sixteen fl. ounces.

Macerate for four days, then pack in a percolator, and add rectified spirit until twenty-four fl. ounces of tincture are obtained.

Dose, two to three minims, as an anodyne. *Pereira.*

ACOUSTIC BALSAM.

R. Compound tincture of benzoin,
Tincture of castor,

" opium, each, one fl. ounce.

Essential oil of assafetida, five drops.

Mix. *Cooley.*

R. Tincture of ambergris,
" assafetida,

" castor,

" opium, each, one fl. ounce.

Terebinthinated balsam of sulphur,

Oil of rue, each, fifteen drops.

Mix. One or two drops are poured into the ear, or a piece of wool moistened therewith is introduced, in cases of atonic deafness. *Baumé.*

ANATOMICAL INJECTIONS.

I. FOR PRESERVING THE BODY.

A. Molasses Injection.

R. Boiling water, two gallons.
Common salt, sufficient

to saturate.

Nitre, two to four ounces.

Molasses, four ounces.

Mix together, and boil for a few minutes. The proportions of nitre and molasses are variable. This injection penetrates well, and it will preserve the body long enough for dissecting purposes.

B. Chloride of Zinc Injection.

R. Commercial muriatic acid, ten pounds.

Add gradually

Scrap zinc, more than sufficient

to neutralize the acid.

This solution is extremely concentrated and acid. Where the body is to be kept during the summer months, the Editor deems it advisable to throw the undiluted solution into the aorta, until the fluid oozes from an incision made into the cellular tissue of one of the toes. About half a gallon is usually required to effect this object. In winter, one to three pints, diluted with water, will answer.

As a mere preservative, nothing can surpass this injection; but it is open to the serious objection of bleaching the tissues.

C. Nitrate of Lead Injection.

R. Nitric acid, ten pounds.
Powdered litharge, more than sufficient

to saturate.

Mix. One to two pints of this fluid may be advantageously added to a solution of salt containing a little nitre. The mixture forms a good preservative injection; giving a rusty, rather than a bleached appearance to the tissues.

D. Method of Preserving a Corpse.

R. Corrosive sublimate, sufficient to saturate

Alcohol, three pints.

This mixture is to be thrown into the aorta. To be followed daily for three successive days by the same quantity of a saturated solution of the acetate of alumina. The surface of the body is to be frequently washed with the latter solution. By this method, the natural appearance of the body may be retained for a week.

W. R. Grant.

E. Gannal's Injection.

R. Sulphate of alumina, two pounds.
Acetate of lead, half a pound.

Dissolve the salts, separately, in water, mix the solutions, and filter.

The body is to be injected with six pints of this fluid, by one of the large arteries. *Dorvault.*

II. FOR ARTERIAL AND VENOUS PREPARATIONS.

A. For Coarse objects.

R. Tallow,
Resin,
Yellow wax, equal parts.

Melt over a slow fire, and add sufficient red lead, or vermilion to color.

R. Wax, sixteen ounces.
Resin, eight ounces.
Turpentine varnish, eight ounces.
Vermilion, one ounce.

Melt the wax and resin, stirring in the varnish and vermilion.

This injection is liable to melt in warm weather, thus giving a flattened appearance to the blood vessels.

R. Tallow, two pounds.
Magnesia, half an ounce.
Vermilion, one ounce.

Melt the tallow, and stir in the magnesia and vermilion.

This, like the two former injections, is thrown in hot. The following is used cold, and hence is called the *cold paint injection*.

R. White lead, well ground, two pounds
Turpentine varnish, twelve drachms.
Linseed oil, six ounces.

Grind the white lead and varnish together, add the oil, and rub to the consistence of cream; in which state it is to be thrown into the arteries.

Dublin Dissector.

B. For Fine Preparations.

R. Lead, five parts.
Melt, and add
Bismuth, eight parts.
Tin, three parts.
Mercury, two parts.

Melt. This liquid amalgam is used for injecting the kidneys and other organs. *Francis.*

R. Glue, one ounce.

Dissolve, and add

Strong size, one pint.
Vermilion, or other coloring matter, one ounce and a half.

Mix. *Francis.*

R. Canada balsam, one pound.
Vermilion, sufficient to color.

Mix intimately. *Dorvault.*

C. For Preserving Specimens.

In the anatomical museums of this country, it is customary to use alcohol in the preservation of specimens.

For the brain, nerve-matter, and ligaments, strong alcohol is employed. For other parts, a mixture of two parts of alcohol, and one of water is found to answer.

The following are possessed of much merit, as preservative fluids:

R. Saturate water with sulphurous acid, and add a little creasote.

Cooley.

R. Powdered corrosive sublimate, two drachms.
Alcohol, one pint.

Mix, dissolve, and decant after twenty-four hours. *Francis*

Goadby's Solutions.

	No. 1.	No. 2.	No. 3.
Salt,	4 oz.	4 oz.	8 oz.
Alum,	2 oz.	2 oz.	
Corrosive sublimate,	2 grs.	3 grs.	2 grs.
Water,	2 pts.	4 pts.	2 pts.

Mr. Goadby usually employs No. 1; but for delicate preparations that might be injured by a concentrated solution, he substitutes No. 2. As alum decomposes osseous matter, No. 3 is used when the tissues contain carbonate of lime.

	No. 4.	No. 5.
Salt,	8 oz.	8 oz.
Corrosive sublimate,		15 grs.
Arsenious acid,	15 grs.	15 grs.
Water,	2 pints.	2 pints.

These solutions are used for old and soft preparations. *Dorvault.*

FLUID EXTRACT OF BLACK SNAKE-ROOT.

R. Black snakeroot, powdered,
(troy) eight ounces.

Alcohol, one pint.
Diluted alcohol, sufficient.

Macerate the root in the alcohol for twenty-four hours, then transfer to a percolator, add diluted alcohol until a pint of tincture is obtained, and allow this to evaporate spontaneously to one fl. ounce. Displace the dregs with diluted alcohol, until half a pint of tincture has passed. Evaporate the latter tincture, on a water-bath, to four fl. ounces, add six ounces of sugar, dissolve, and stir in the concentrated alcoholic tincture. The whole measures half a pint. *W. J. Watson.*

Dose, half a teaspoonful.

In practice this is found to be too consistent. It is therefore better to concentrate the tinctures to a pint, instead of half a pint.

R. Black snakeroot, in coarse powder,
(troy) sixteen ounces.

Alcohol, one pint.
Ether, half a pint.
Diluted alcohol, sufficient.

Introduce the powder into a percolator for volatile liquids, add the alcohol and ether; allow the percolation to take place by drops, and, as soon as the menstruum disappears above, add diluted alcohol until the filtered tincture measures one pint and a half. Let this evaporate in a capsule to half a pint. Add diluted alcohol to the dregs in the percolator until two pints more of tincture are obtained. Evaporate this on a water-bath to eight fl. ounces, mix gradually with the first product, let stand for a few hours, filter, and add alcohol, if necessary, to make a pint. *W. Procter, Jr.*

EXTRACT OF BLACK SNAKEROOT.

A solid extract of the root can readily be prepared by the above process, by continuing the evaporation on a water-bath. Eight grains of this extract represent a drachm of the root.

W. Procter, Jr.

BLACKET'S TINCTURE OF BELLADONNA.

R. Extract of belladonna, ten drachms.
Proof spirit, one pound.

Macerate, dissolve, and strain.

Dose, two or three minims, as an anodyne.

Lond. Ph. Jour.

BLACKING.

No. 1.

R. Ivory black, twelve ounces.
Treacle, four ounces.
Sperm oil, one ounce.
Vinegar, two pints.
Oil of vitriol, (by weight) two ounces.

Mix the first three ingredients, and add the vinegar gradually, stirring after each addition; add the oil of vitriol very carefully, stirring constantly until the effervescence ceases.

Lond. Ph. Jour.

No. 2.

R. Ivory black, twelve ounces.
Treacle, each, twelve ounces.
Sperm oil, three ounces.
Oil of vitriol, each, four pints.
Vinegar, four pints.

Mix the ivory black, treacle, and vinegar together; then mix the sperm oil and oil of vitriol separately, and add them to the other mixture.

Redwood.

No. 3.—(Liquid.)

R. Ivory black, in fine powder, one pound.
Treacle, three-fourths of a pound.
Sweet oil, two ounces.
Beer,
Vinegar, each, one pint.

Rub the first three together until uniformly mixed, then add the beer and vinegar. *Cooley.*

No. 4.—(Paste.)

R. Ivory black, in powder, one pound and a quarter.
Treacle, one pound.
Sweet oil, two ounces.

Rub together till well mixed, then add
Vinegar, sufficient
to form a paste. *Cooley.*

BLACKING FOR GRATES.

R. Asphaltum, four pounds.
Melt and add
Linseed oil, two pounds.
Oil of turpentine, one gallon.
Mix. *Gray.*

BOOT COMPOSITION.

R. Boiled linseed oil, one pint.
Oil of turpentine,
Black resin,
Wax, each, three ounces

Melt the wax and resin together, and stir in the boiled oil; then remove the pot from the fire, and when it has cooled a little, add the turpentine, and mix. *Cooley*

- R. Caoutchouc, two drachms.
 Mineral naphtha, two ounces.
 Asphaltum,
 Ivory black, each, half an ounce.
 Oil of turpentine, one ounce.

Dissolve the caoutchouc in the naphtha, and the asphaltum in the turpentine; mix the two solutions, and add the ivory black.

Lond. Ph. Jour.

Both of these compositions are reputed to be water-proof.

BUG POISON.

No. 1.

- R. Alcohol, one hundred and twenty parts.
 Camphor, two parts.
 Oil of turpentine, four parts.
 Corrosive sublimate, one part.
 Mix and dissolve. *Dorvault.*

No. 2.

- R. Coal-tar naphtha,
 Oil of turpentine, equal parts.
 Mix. *Redwood.*

No. 3.

For Roaches and Ants.

- R. Arsenic, in powder, one part.
 Mashed potatoes, three parts.
 Mix.

No. 4.

For Roaches, Rats, Mice, &c.

- R. Phosphorus, eight parts.
 Tepid water,
 Rye meal,
 Butter, each, one hundred and eighty parts.
 Sugar, one hundred and twenty-five parts.

Liquefy the phosphorus in the tepid water in a bottle, and mix it in a mortar with the rye meal; when cold, add the butter and sugar, and mix them all thoroughly together. *Gray.*

No. 5.

For Flies, &c.

- R. Rasped quassia, two drachms.
 Boiling water, one pint.
 Boil together for a few minutes, strain, and add
 Molasses, four fl. ounces.
 Mix. *Dorvault.*

CEMENTS.

Diamond Cement.

- R. Isinglass, two drachms.
 Sprinkle it with water, and, when soft, dissolve it in the smallest possible quantity of proof spirit by a moderate heat.

- R. Mastich, one drachm.
 Rectified spirit, three fl. drachms.

Dissolve. Mix the two solutions, and stir in one drachm of powdered ammoniac rubbed down with a little water. Keep the cement in a bottle. When wanted for use, plunge the bottle in warm water to soften the cement, which is applied in thin layers on the fractured surfaces, and the pieces are to be held in firm contact till cool.

Used for setting jewelry, mending china, &c.
Lond. Pharm. Jour.

Cement for China and Stoneware.

Gelatine is allowed to swell in cold water, the jelly is warmed, and so much freshly-slaked lime is added as is necessary to thicken the mass. A thin coating of this cement is applied warm to the gently-heated surface, and the pieces are held together under strong pressure. *Heller.*

- R. Whites of eggs, at will.
 Finely-powdered quicklime, sufficient
 to thicken. Used for mending china, &c.; but it does not resist moisture.

- R. Shellac, at will.
 Alcohol, sufficient
 to form a liquid of the consistence of molasses.
 This is an excellent cement.

Cement for the Laboratory.

- R. Resin, five pounds.
 Dry red ochre, in fine powder,
 Wax, each, one pound.
 Plaster of Paris, two to four ounces.
 Melt the resin and wax, add the ochre, and, lastly, the plaster. Mix well together. *Singer.*

Luting for Bottles, &c.

Melt common resin and yellow wax together, and add red ochre in small quantities, and when of the proper consistence, boil for six or eight minutes; add a small quantity of oil of turpentine, and stir well with a spatula. The proportions of the different constituents vary, the luting being more or less brittle, or elastic, as the ochre prevails. *Francis.*

- R. Flour,
 Whiting, each, ten parts.
 Common salt, one part.
 Mix and add sufficient water to make a paste. This is used by the distillers of spirituous liquors. *Francis.*

- R. Sal ammoniac,
 Whiting, equal parts.
 Water, sufficient
 to form a paste. This forms a good lute. *Francis.*

R. Common clay,
Whiting, equal parts.
Water, sufficient
to form a paste. This lute will stand a high heat. Francis.

COLD CREAM.

R. Rose water, half a fl. ounce.
Oil of almonds, two fl. ounces.
Spermaceti, half an ounce.
White wax, one drachm.

Melt the oil, spermaceti, and wax, by means of a water-bath; then add the rose water, and stir till cold. This keeps better than the officinal ointment B. Canavan.

COLORS FOR SHOW-BOTTLES.

Blue.

R. Sulphate of copper,
Alum, each, one ounce.
Water, two pints.
Sulphuric acid, half an ounce.
Mix, and dissolve.

R. Prussian blue, ten grains.
Oxalic acid, twenty grains.
Water, one pint.
Dissolve.

R. A. Dissolve ammonio-sulphate of copper, and ammonio-nitrate of nickel (prepared by dissolving nickel in diluted nitric acid, and adding ammonia in excess), in water.

B. Dissolve the sulphate of indigo in water.

C. Dissolve Prussian blue in diluted muriatic acid, and dilute with water.

Purple.

R. Verdigris, two drachms.
Spirit of hartshorn, four ounces.
Water, one pint and a half.

Mix.

R. Sulphate of copper, one ounce.
Carbonate of ammonia, one ounce and a half.
Water, two pints and a half.

Mix.

R. Infusion of logwood, two pints.
Spirit of hartshorn, sufficient
to produce the desired hue.

Mix.

Lilac.

R. Add carbonate of ammonia to a solution of nitrate of cobalt, until the precipitate first formed is re-dissolved; adding ammonio-sulphate of copper to bring out the desired tint.

R. Dissolve zaffre (impure oxide of cobalt) in muriatic acid, filter, and add carbonate of ammonia in excess; to this add ammonio-sulphate of copper, until the required color is produced.

Orange.

R. Dissolve bichromate of potassa in water till the desired tint is obtained.

A little sulphuric acid is sometimes added.

Yellow.

R. Bichromate of potassa, six drachms.
Carbonate of potassa, four drachms.
Water, sixteen ounces.

Mix.

Red.

R. A. Color the spirit of hartshorn, or an aqueous solution of sal ammoniac with cochineal.

B. Add vinegar to red-beet liquor, or to red-cabbage liquor.

C. Dissolve carmine in the solution of ammonia, and dilute with water.

Green.

R. A. Dissolve three ounces of verdigris in sulphuric acid, and add four pints of water.

B. Dissolve one ounce of nickel in six ounces of nitric acid, and add five pints of water.

C. Dissolve ammonio-sulphate of copper in water, and add bichromate of potassa, until the required color is produced.

FLUID EXTRACT OF CUBEBS.

R. Powdered cubebs, one pound.
Sulphuric ether, }
Alcohol, } each, sufficient.
Water, }

Pack the cubebs in a displacement funnel, cover with a mixture of equal parts of alcohol and ether, close the top and bottom of the funnel, and let stand for twenty-four hours. Then remove the cork, and, after the dropping has ceased, add alcohol until a pint of tincture is obtained. Set this aside in a shallow vessel, and permit it to evaporate spontaneously to one-half. Obtain a second pint of tincture by displacing with diluted alcohol, and permit three-fourths of it to evaporate spontaneously. Obtain a third pint by displacing with water, and allow it to evaporate one-fourth. To the three products, amounting to one pint and a half, add another pint of fluid obtained by again displacing the cubebs with water. Transfer the two pints and a half of fluid to a proper bottle.

and dissolve in it sixteen ounces of sugar, making the whole measure three pints.

One fl. drachm is equal to a scruple of the powder. *B. Canavan.*

DENTIFRICE.

No. 1.

R. Powdered tormentilla, eighty parts.
 " orris root, forty parts.
 Cream of tartar, twenty parts.
 Fused chloride of calcium, four parts.

Reduce the chloride rapidly to powder, and mix it with the other ingredients. *Hainault.*

No. 2.

R. Powdered Peruvian bark, one ounce.
 " myrrh, } each, half an
 " rhatany, } ounce.
 " orris root, }

Mix carefully.

No. 3.

R. Powdered Peruvian bark,
 " myrrh, each, half an ounce.
 " charcoal, two ounces.

Mix.

DEPILATORIES.

No. 1.

R. Sulphuret of sodium, three parts.
 Quick lime, in powder, ten parts.
 Starch, ten parts.

Mix. Rub a little of this powder with water, apply it to the part, and remove the hair in a minute or two with a wooden knife. *F. Boudet.*

No. 2.

R. To a strong solution of sulphuret of barium, add sufficient powdered starch to make a paste. It is used like the preceding. *Redwood.*

EAU DE PAGLIARI.

R. Tincture of benzoin, eight fl. ounces.
 Alum, one pound.
 Water, ten pounds.

Mix. This preparation is said to possess an extraordinary power of coagulating the blood. *Sedillot.*

MIXTURES FOR COLORED FIRES.

Red Fire.

R. Dried nitrate of strontia, seventy-two parts.
 Sulphur, twenty parts.
 Gunpowder, eight parts.
 Coal dust, two parts.

Mix. This mixture is liable to spontaneous explosion. *Redwood.*

R. Chlorate of potassa, sixty-one parts.
 Sulphur, sixteen parts.
 Carbonate of strontia, twenty-three parts.

Mix the powders with the hand. *Marchand.*

Yellow Fire.

R. Chlorate of potassa, sixty-one parts.
 Sulphur, sixteen parts.
 Dry soda, twenty-three parts.

Mix. *Marchand.*

R. Dried nitrate of soda, seventy-four and a half parts.
 Sulphur, nineteen and a half parts.
 Charcoal, six parts.

Mix. *Gray.*

Blue Fire.

R. Nitre, five parts.
 Sulphur, two parts.
 Metallic antimony, one part.

Mix. *Gray.*

R. Chlorate of potassa, sixty-one parts.
 Sulphur, sixteen parts.
 Strongly-calcined alum, twenty-three parts.

Mix. *Marchand.*

Green Fire.

R. Chlorate of potassa, seventy-three parts.
 Sulphur, seventeen parts.
 Boracic acid, ten parts.

Mix. *Marchand.*

Violet Fire.

R. Chlorate of potassa, sixty parts.
 Sulphur, sixteen parts.
 Carbonate of potassa, twelve parts.
 Alum, twelve parts.

Mix. *Marchand.*

White Fire.

R. Nitre, forty-six and a half parts.
 Sulphur, twenty-three parts.
 Gunpowder, twelve and a half parts.
 Zinc powder, eighteen parts.

Mix. *Gray.*

LIQUID GLUE.

R. Glue, two pounds and a half.
 Water, two pints and a quarter.

Dissolve in a water-bath, and add gradually Nitric acid (*sp. gr.* 1.32), seven ounces.

Remove from the fire, and stand aside to cool. This glue keeps well, and is used for a variety of purposes. *Dumoulin.*

MARINE GLUE.

R. Caoutchouc, two to four parts.
Coal-tar naphtha, thirty-four parts.
Mix, dissolve by heat, and add

Powdered shellac, sixty-four parts.
Heat gently, and stir till uniformly mixed; then pour out on plates to harden.

When used, it is necessary to heat the cement to 248° F. and apply it with a brush.

The union made by this cement is very durable. *Dorevault.*

INKS.

BLACK INK.

R. Chipped logwood, twenty-two pounds.
Boiling water, sufficient
to yield fourteen gallons of filtered decoction.
When cold, add, very gradually,

Yellow chromate of potassa, sufficient
to obtain the required color.

This ink does not corrode steel pens.

Runge.

R. Powdered extract of
logwood, (avoir.) twelve ounces.
Bichromate of potassa, half an ounce.
Water, five gallons.

Dissolve the ingredients separately in water, and mix them.

In a short time, the ink will be fit for use.

W. H. Pile.

BLUE INK.

R. Dissolve Prussian blue in an aqueous solution of oxalic acid.

Lond. Pharm. Jour.

R. Dissolve indigo in sulphuric acid, and add water to produce the proper shade. As much potassa is to be added as the liquid will bear without dropping its color.

Cooley.

RED INK.

R. Pure carmine, twelve grains.
Water of ammonia, three ounces.

Dissolve, and add

Powdered gum, eighteen grains.

Mix.

Where expense constitutes an objection, half a drachm of powdered drop-lake may be substituted for the powdered carmine.

Buchner's Repert.

R. Brazil wood, four ounces.
Alum, two ounces.
Distilled water, two pints.

Boil for fifteen minutes, and strain; then add
Powdered gum Arabic, one ounce.
Mix, and bottle for use. *Gray.*

INDELIBLE INK.

R. Nitrate of silver, two drachms.
Gum Arabic, one drachm.
Distilled water, one fl. ounce.
Indian ink, sufficient to color.

Mix, and dissolve.

R. Carbonate of soda, in
crystals, two ounces.
Gum Arabic, two drachms.
Water, four fl. ounces.

Mix.

The place to be marked is moistened with the last solution, and dried. It is then to be marked with a pen dipped in the first solution, and afterwards exposed to the rays of the sun; and washed after twenty-four hours. *U. S. Disp.*

FLUID EXTRACT OF IPECACUANHA.

R. Powdered ipecacuanha, (troy) eight ounces.
Alcohol (.835), twelve fl. ounces.

Mix, and set aside for twelve hours. Then add sufficient alcohol to make it of the consistence of syrup, and introduce into a displacer, in which it gradually settles down as the alcohol percolates. A piece of muslin is then to be laid on the surface, and alcohol is to be added until the filtered liquid measures half a gallon. Reserve the first half pint that comes through; then distil and evaporate the remainder to eight fl. ounces, to which the reserved half pint is to be added.

A fl. drachm represents half a drachm of the root. *Jos. Laidley.*

SYRUP OF IPECACUANHA.

R. Fluid extract of ipecacuanha, eight fl. ounces.
Simple syrup, four pints.
Mix, and evaporate to three pints. Then add
Simple syrup, four pints.
Water, one pint.

Mix.

If the solution should not be clear, it may be clarified by beating the white of an egg with the water before adding to the syrup, then boiling for a few minutes, and straining.

Jos. Laidley.

PREPARATIONS OF IRON.

SACCHARINE CARBONATE OF IRON AND MANGANESE.

R. Finely-powdered sulphate of iron, three ounces, one drachm.
Carbonate of soda, five ounces

Sulphate of man-
ganese, one ounce, one scruple.
Powdered
sugar, two ounces and a half.

Dissolve each of the first three ingredients in a pint and a half of water, and mix the solutions. Collect the precipitate on a cloth filter, and wash it immediately with cold water; squeeze out as much water as possible, and, without delay, triturate the pulp with the sugar. Dry it at a temperature of 120° F.

Dose, five grains, gradually increased to a scruple, three times a-day. *T. S. Speer.*

SYRUP OF PROTO-NITRATE OF IRON.

R. Nitric acid (*sp. gr.* 1.42),
two fl. ounces.

Carbonate of lime, sufficient
to saturate. Filter, and wash the filter with sufficient water to make the solution measure three fl. ounces.

R. Proto-sulphate of iron,
in crystals, fourteen hundred and
eighty-five grains.

Water, ten fl. ounces.

Mix, dissolve, and filter. Mix the two solutions, and pour immediately on a muslin filter, stretched across one end of a glass percolator. Allow the filtrate to fall upon two pounds of sugar. Dissolve without heat, and add water to make the syrup measure thirty fl. ounces. Each fl. ounce contains thirty-two grains of the proto-nitrate of iron. *T. Lancaster.*

COMPOUND SYRUP OF PHOSPHATE OF IRON.

R. Proto-sulphate
of iron, four drachms and two
scruples.

Crystallized phosphate
of soda, seven drachms and a half.

Phosphate of potassa, two drachms.

Freshly-precipitated phos-
phate of lime, four drachms.

Glacial phosphoric acid, one ounce.

Sugar, in coarse powder, eight ounces.

Water, sufficient.

Dissolve the sulphate of iron, and five and a half drachms of the phosphate of soda, severally, in three fl. ounces of the water, and mix the solutions. Wash the precipitate with (cold) boiled water, mix it with the phosphate of lime and half a pint of water, in a porcelain capsule, apply heat, gradually add the phosphoric acid, continuing the heat until a clear solution is obtained, and dissolve in it seven (troy) ounces of the sugar. Then dissolve the phosphate of potassa, two drachms of the phosphate of soda, and an ounce of sugar, in a fl. ounce of water; acidulate the solution with phosphoric acid, and add it to the syrupy solution first obtained.

Then add forty drops of muriatic acid to remove the cloudiness of the mixture.

A teaspoonful contains one and two-fifths grains of proto-phosphate of iron, two and a half grains of phosphate of lime, one and one-fifth grains of each of the alkaline phosphates, and four and a half grains of free phosphoric acid; which may be considered the dose.

T. S. Wiegand.

JACKSON'S PECTORAL LOZENGES.

R. Powdered ipecacuanha, ten grains.

Precipitated sulphuret of
antimony, five grains.

Muriate of morphia, six grains.

Powdered gum Arabic, } each, eleven

" sugar, } drachms.

" liquorice, }

Oil of sassafras, four drops.

Tincture of Tolu, four fl. drachms.

Syrup, sufficient.

Mix, and divide into two hundred lozenges.

S. Jackson.

JACKSON'S PECTORAL SYRUP.

R. Sassafras pith, one drachm.

Gum Arabic, one ounce.

Water, one pint.

Macerate twelve hours, then add

Sugar, twenty-one ounces.

Dissolve without heat, strain, and add

Muriate of morphia, eight grains.

Mix carefully, and add water sufficient to make the whole measure two pints. Dose, a teaspoonful to a tablespoonful, every three hours.

S. Jackson.

KIRKLAND'S NEUTRAL CERATE.

R. Lead plaster, eight ounces.

Olive oil, four ounces.

Melt, and, while fluid, add

Prepared chalk, four ounces.

Stir, until sufficiently cooled, then add

Acetic acid, four fl. ounces.

Acetate of lead, three drachms.

Mix. *Paris.*

LEDOYEN'S DISINFECTING LIQUID.

R. Nitrate of lead, (prepared by
dissolving litharge in di-
luted nitric acid), one part.

Water, eight parts.

Mix, and dissolve. Used to correct noxious
effluvia, &c. *Pereira.*

MEDICATED WATERS.

R. Any essential oil, two drachms.

Precipitated chalk, one ounce.

Alcohol, two fl. ounces.
Distilled water, one gallon.

Rub the oil with the chalk, then add the spirit gradually, and when completely dissolved, pour in the water by degrees; let it stand three minutes, and filter through close paper.

Mr. John Fordred affirms that chalk is decidedly preferable to the carbonate of magnesia, in preparing medicated waters.

Lond. Pharm. Jour.

EXTEMPORANEOUS PREPARATION OF MEDICATED WATERS.

R. Any essential oil, two drops.
Carbonate of magnesia, six grains.
Rub together, and add gradually
Water, one fl. ounce.

Filter the solution through bibulous paper.

PROCESS FOR COATING PILLS.

R. Flaxseed, one part.
White sugar, three parts.
Distilled water, sufficient.

Pour the boiling water on the seed to extract the mucilage, separate the thick mucilage, add the sugar, and carefully drive off the whole of the moisture by evaporation. Reduce the dried mucilage to powder, in which the pills (having been previously moistened,) are to be rolled until covered with a layer of the compound.

Calloud.

R. Butter of cacao, at will.
Melt. Throw the prepared pills into the melted butter, then remove them with a perforated skimmer, and roll them in powdered sugar, or sugar of milk.

Calloud.

PASTILLES DE PARIS.

R. Powdered cubebs, one ounce and a half.
" gum Arabic, one ounce.
" liquorice, " "
" sugar, each, four ounces.
Oil of lemon, fifty drops.
Mix, and make lozenges of eight grains.

POMATUM.

R. Beef marrow, seventeen drachms.
Nervine balsam, (see page 305), seventeen drachms.
Rose oil, two drachms.
Melt together, and add
Alcoholic extract of cantharides, six grains,
dissolved in a little alcohol. *Dupuytren.*
R. Beef marrow, three to four ounces.
Castor oil, one ounce.

Melt, and strain the marrow, add the oil, and, as the mixture cools, beat continually until it assumes a white, frothy appearance; flavoring with extract of orange.

PUTTY.

Whiting made into a paste with boiled linseed oil.

SEALING WAX.

Red.

R. Shellac, two pounds.
Venice turpentine, one pound.
Vermilion, or best dichromate of lead, one pound and a half.

Melt the shellac and turpentine together, and add the pigment, as the mixture cools.

Redwood.

Black

Is made in the same way as the red, only substituting the best lamp-black for vermilion, or dichromate of lead.

In like manner, other colors are imparted by varying the pigment, and using chromate of lead, verdigris, green verditer, &c. *Redwood.*

Bottle Wax.

R. Black resin, six pounds.
Bees' wax, half a pound.
Ivory, or lamp-black, one pound and a half.

Melt the resin and wax, and stir in the ivory-black.

Venetian red, red lead, or bole, may be substituted for lamp-black. *Redwood.*

SOLDERS.

Solder for Tin Plate.

R. Tin, two parts.
Lead, one part.
Mix.

Solder for Pewter.

R. Tin, ten parts.
Lead, five parts.
Bismuth, one part.
Mix.

Solder for Iron, Copper, and Brass.

R. Copper, equal parts.
Zinc, equal parts.
Mix.

Solder for Zinc and Lead.

R. Lead, two parts.
Tin, one part.
Mix.

Solder for Silver.

R. Silver,	five parts.
Brass,	six parts.
Zinc,	two parts.
Mix.	<i>Gray's Supp.</i>

Solder for Gold.

R. Gold,	twelve parts.
Silver,	two parts.
Copper,	four parts.
Mix.	<i>Francis.</i>

FELT SPLINTS.

Dissolve three pounds of shellac in two quarts of alcohol, in a tinned vessel, with a tight cover to prevent evaporation. Spread a piece of woollen cloth on a board, and with a clean brush saturate both sides of the cloth with the solution. Hang it up until thoroughly dried; then lay it again upon the board, and apply a second coat of the solution to one side only of the cloth. Dry again, and apply a third coat to the same side. While the last coat is yet fresh, fold the cloth so that the side having three coats shall be applied to itself. Then, with a hot flat-iron, smooth, and press the surfaces together. When cold, a slight rubbing with sand-paper makes it fit for use.

It becomes a firm, almost unyielding board; but exposure to a moderate heat will make it pliant, so that it can easily and accurately be adapted to any surface. *Dr. F. H. Hamilton.*

TRACING PAPER.

Paper brushed over with a thin varnish of colorless Damara resin, the varnish being allowed to soak through the paper without any apparent coating of it remaining on the surface. *Redwood.*

VARNISHES.

Varnish for coating Metals.

R. Copal,	
Oil of rosemary, each,	one part.
Alcohol,	two to three parts.
Mix.	It should be applied hot, and, when dry, it will be found very hard and durable. <i>Schindler.</i>

Varnish for Leather.

R. Oil of turpentine, saturated with caoutchouc,	six parts.
Copal,	
Oil of rosemary, each,	two parts.
Mix.	This varnish should be applied somewhat fluid, and always dried at a high temperature. <i>Schindler.</i>

R. Alcohol,	twenty gallons.
Shellac,	thirty-five pounds.
Resin,	twenty pounds.
White turpentine,	ten pounds.
Powdered lamp-black,	four pounds.

Bruise the shellac and resin, put them in the alcohol, and stir ten or twelve times during the day, which will prevent the formation of lumps, and will nearly dissolve it in one day. Next day, add the turpentine, and stir five or six times; the third day, stir in the powdered and sifted lamp-black.

No heat is employed.

E. S. Frey.

Varnish for Furniture.

R. Dissolve fused copal in oil of turpentine.

If the copal has not been kept a sufficient time in a state of fusion, the varnish made with it remains soft for some time after it is dry, and afterwards peels off. *Schindler.*

Amber Varnish.

R. Amber,	one pound.
Place it in an iron pot, and render it semifluid by heat; then add	
Pale boiled oil,	ten ounces.
Mix, remove from the fire, and add	
Oil of turpentine,	one pint.
Stir well together.	<i>Cooley.</i>

Japan Varnish.

R. Oil of turpentine,	eight ounces.
" lavender,	six ounces.
Camphor,	one drachm.
Bruised copal,	two ounces.
Mix and dissolve. Used for jappanning tin.	<i>Cooley.</i>

Lac Varnish (Aqueous).

R. Pale shellac,	five ounces.
Borax,	one ounce.
Water,	one pint.
Digest at nearly the boiling point until dissolved; then strain. A good vehicle for water colors, inks, &c.	<i>Cooley.</i>

Lac Varnish.

R. Shellac,	ninety parts.
Venice turpentine,	four parts.
Alcohol,	five hundred parts.
Digest until dissolved, and strain.	<i>Gray.</i>

Copal Varnish.

A variety of methods is resorted to, for making copal varnish. One of the cheapest is to dissolve copal in drying linseed oil, near the boiling point, and, when sufficiently cool, to dilute the mixture with oil of turpentine.

Cooley.

DIETETIC PREPARATIONS.

NOT INCLUDED AMONG THE PREVIOUS PRESCRIPTIONS.

TOUS LES MOIS.

This is a variety of arrowroot, prepared from the rhizomes of a species of *Canna*, and is to be employed in the same manner, and in same cases, as arrowroot.

MUCILAGE OF SALEP.

Salep is the prepared bulbs of several species of orchidaceous plants. It is more nutritive than arrowroot or sago, and hence is adapted for the convalescent, rather than for the sick. These roots are powdered with much difficulty. The mucilage is prepared by dissolving the powder in boiling water, constantly stirring, and adding to the solution, sugar and milk. *Soubeiran* orders powdered salep, half an ounce, to be boiled with a sufficient quantity of water till dissolved, four ounces of sugar added, and to be aromatized according to taste.

BISCUIT JELLY.

White biscuit, four ounces, to be boiled down in four pints of water, to one-half, strained, and evaporated to one pint; one pound of white sugar, four ounces of port wine, and one drachm of cinnamon water added, and the whole well mixed.

It has been found useful in debility of the digestive organs.

A. T. Thomson.

BREAD JELLY.

Cut a French roll into slices, toast these on both sides, and boil in a quart of water, until the whole forms a jelly, adding more water if required; strain, and flavor.

This is nutritious, and may be made more so by using broth, wholly deprived of fat, instead of water.

Cooley.

PANADA.

Stale wheat bread, one ounce; cinnamon, one drachm; water, one pint.

Cover up and let stand for an hour, beat up and boil for ten minutes, adding a little grated nutmeg and sugar. Wine may be added, if required.

Ellis.

COMPOUND SALEP POWDERS.

Powdered salep, tragacanth, and sago, each, four ounces; cochineal, half a drachm; prepared oyster-shells, one ounce.

These are to be carefully mixed and divided into powders, of one drachm each. Stir one of these into a pint of milk, and boil for ten or fifteen minutes. To be taken freely in diarrhoea and dysentery.

Ellis.

These are known as *Castanella's powders*, for which the following is used as a substitute.

Powdered gum Arabic, tragacanth, arrowroot, sago, and tapioca, each, two drachms. Mix them well together, boil in a pint of milk, flavored with nutmeg or cinnamon.

Used as a diet in dysentery, diarrhoea, &c.

VEGETABLE BROTH.

Take two potatoes, one carrot, one turnip, and one onion.

Slice them, and boil in a quart of water for an hour, adding more water from time to time, so as to keep up the original quantity; flavor with salt, and a small portion of potherbs; strain. When advisable, a small quantity of mushroom catsup added to this broth greatly improves its taste.

A good substitute for animal food, when the last is inadmissible.

Ellis.

GLOUCESTER JELLY.

Rice, sago, pearl barley, hartshorn shavings, eryngium root, each, one ounce; boil in three pints of water to one pint, and strain.

This is very nutritive, dissolved in broth, milk, or wine.

A. T. Thomson.

ALMOND JELLY.

Blanched sweet almonds and white sugar, each, one ounce; water, four ounces; make an emulsion; strain, and add melted hartshorn jelly, half a pound; orange-flower water, one drachm; and essence of lemon, two or three drops.

Cooley.

BOILED FLOUR.

Take of fine flour a pound, tie it up in a linen cloth as tight as possible, and after frequently dipping it into cold water, dredge the outside with flour, till a crust is formed round it, which will prevent the water soaking into it, whilst boiling.

Boil for a long time, and permit to cool, when it will become a hard, dry mass.

This is to be grated and prepared like arrow-root. A good diet for children, in diarrhoea, &c. *Ellis.*

BEEF TEA.

Take of lean beef, cut into shreds, one pound; water, one quart.

Boil for twenty minutes, removing any scum that arises. When it has become cold, strain. *Ellis.*

Take half a pound of good rump steak, cut it into thin slices, and spread these in a hollow dish; sprinkle a little salt over them, and pour upon the whole a pint of boiling water.

Cover the dish and place it near the fire for half an hour; then remove to a pan and boil for fifteen minutes; strain through a fine sieve.

The quantity of water is too small for the strength of the tea for invalids, but is sufficient to extract all the soluble parts of the beef, and the tea can be reduced to the proper strength by the addition of boiling water. *A. T. Thomson.*

ESSENCE OF BEEF.

Take of lean beef, sliced, a sufficient quantity to fill the body of a porter bottle, cork up loosely, and place it in a pot of cold water, attaching the neck, by means of a string, to the handle of the pot.

Boil for an hour and a half to two hours, then decant the liquid and skim it.

To this preparation may be added spices, salt, wine, brandy, &c., according to the taste of the patient and nature of the disease. *Ellis.*

MUTTON TEA.

Take one pound of good mutton, freed from the fat, and cut into thin slices; pour over it a pint and a half of boiling water, in the same manner as directed for beef tea; but it requires to be boiled for half an hour, previous to straining.

If the invalid desires the addition of barley, an ounce of clean pearl barley, washed, and macerated in boiling water for an hour, may be boiled with the mutton tea, and the undissolved barley separated by straining. *A. T. Thomson.*

VEAL TEA.

This is to be made in the same manner as beef tea, using a pound of the fillet of veal, freed from fat and sliced, and a pint and a half of boiling water; boiling for half an hour. It may also be made with the same quantity of the fleshy part of a knuckle of veal. By boiling this latter down to one-half, and straining, the decoction will gelatinize, and, if poured into small cups, will keep good for several days.

By the addition of an equal quantity of boiling water to this jelly, it is fit for use in a few minutes. *A. T. Thomson.*

CHICKEN WATER.

Take half a chicken, divested of all fat, and break the bones; add to this half a gallon of water.

Boil for half an hour, strain, and season with salt. *Ellis.*

Take a small chicken, free it from the skin, and from all the fat between the muscles; divide it longitudinally into halves; remove the lungs, liver, &c. then cut it, bones and muscles, into thin slices, and put these into a pan with a sufficient quantity of salt; add a quart of boiling water, cover the pan, and simmer slowly for two hours; strain through a fine sieve.

A. T. Thomson.

CALVES' FEET JELLY.

Take two calves' feet, and add to them one gallon of water, and boil down to one quart; strain, and when cold, remove all fat; then add the whites of six or eight eggs, well beaten; a pint of wine, half a pound of loaf sugar, and the juice of four lemons, and mix well.

Boil for a few minutes, constantly stirring; then strain through a flannel bag. The wine may be omitted if necessary. *Ellis.*

TOAST WATER.

Toast thoroughly a slice of stale bread, put it in a jug and pour over it a quart of water, which has been boiled and cooled, and in two hours decant; a small piece of orange or lemon peel put into the jug with the bread, improves the flavor greatly.

This forms a good drink in febrile affections. *A. T. Thomson.*

APPLE WATER.

Slice two large apples, and pour over the slices, a pint of boiling water.

Let stand for an hour, and decant; if necessary, sweeten with a little white sugar.

A. T. Thomson.

It is also prepared by boiling for an hour, ten ounces of sliced apple in two pints of water, and straining.

Soubeiran.

LEMON-PEEL WATER.

Pare the rind of one lemon, which has been previously rubbed with half an ounce of refined loaf sugar, put the peeling and sugar into a jar, and pour over them a quart of boiling water. When cold, pour off the fluid, and add a tablespoonful of lemon juice.

If wine be not improper, a glass of sherry may be added, instead of the lemon juice.

A. T. Thomson.

BRAN TEA.

Fresh wheat bran, one pound; water, three quarts.

Boil down to one quart, strain, and add sugar, honey, or molasses, according to the taste of the patient.

CHICKEN PANADA.

Take the white meat of a boiled or roasted chicken, free it from the skin, and cut it into small morsels; pound these in a mortar with an equal quantity of stale bread, and a sufficiency of salt, adding gradually some of the water in which the chicken has been boiled, or some beef tea, until the whole forms a thin, fluid paste.

Put this into a pan, and boil for ten minutes, continually stirring.

A. T. Thomson.

MUTTON BROTH.

Take a pound of mutton, freed from fat, put it into a pan with three pints of water, and simmer for two hours. Take three carrots, the same number of turnips, peel, and cut them in slices, boil them for half an hour in a quart of water, throw them on a colander to drain off the water, and having boiled two onions sliced in a pint of water, and poured off the water, add the whole of these vegetables to the mutton liquor; after removing the meat, season with salt and a little celery seed.

Simmer slowly for four hours, put in the meat again, and continue to simmer for another

hour. The meat may be served with the broth. A palatable and very nutritive dish for convalescents.

A. T. Thomson.

FOWL WITH RICE.

Free a young fowl from the skin and the fat on the exterior of the body, and simmer it in good beef tea, till it is very tender.

Season with salt, and, having boiled some rice, add it to the liquor before the fowl is dished.

A. T. Thomson.

WATER SOUCHY.

Take two small, fresh flounders, boil them in a quart of water to one-third, or long enough to reduce the fish to a pulp; strain the liquor through a sieve, and, having cut off the fins of four other small flounders, put the latter into the liquor, with a sufficiency of salt, a few grains of Cayenne pepper, and a small quantity of chopped parsley, and boil till the fish is perfectly done.

The fish and liquor are to be eaten together. Few dishes are more relished by convalescents from fever.

A. T. Thomson.

Scale, gut, and wash two perch; put salt in the water; when it boils, put in the fish, with an onion cut into slices and separated into rings, and a handful of parsley, picked and washed clean.

When the fish are done, put them in a soup-dish, and pour the liquor over them.

Hofland.

MULLED WINE.

Take a quarter of an ounce of bruised cinnamon, half a nutmeg, grated, and ten bruised cloves; infuse them in half a pint of boiling water for an hour, strain, and add half an ounce of white sugar.

Pour the whole into a pint of hot port or sherry wine.

A good cordial and restorative in the low stages of fever, or in the debility of convalescence from fevers.

A. T. Thomson.

COLD CUSTARD.

Take the yolk and white of an egg, and a tablespoonful of sugar; beat together, till the tenacity of the white of the egg is entirely destroyed; add gradually, constantly stirring, half a pint of cold water, two teaspoonfuls of rose-water, and a little grated nutmeg.

A wineglassful to be taken every two or three hours.

Devoes

MOLASSES POSSET.

Put in a saucepan a pint of best molasses, a teaspoonful of powdered white ginger, and a quarter of a pound of fresh butter.

Simmer for half an hour, stirring frequently. Then stir in the juice of two lemons, or two tablespoonfuls of vinegar, cover the pan, and let it stand by the fire five minutes.

A. T. Thomson.

CHICKEN JELLY.

Cut a chicken into small pieces, bruise the bones, and put the whole into a stone jar, with a cover that fits water-tight. Set the jar in a large kettle of boiling water, and keep it boiling for three hours.

Then strain, and season with salt, pepper, and mace, or with loaf sugar and lemon juice, according to circumstances and taste.

A. T. Thomson.

CHOCOLATE.

Put milk and water on to boil; then scrape the chocolate fine, one or two squares to a pint, as will best suit the stomach; when the mixture of milk and water boils, take it off the fire, throw the chocolate into it, mill it well, and serve it up with the froth.

The sugar may be mixed with the scraped chocolate, or added afterwards. It should never be made before it is wanted, as heating it again injures the flavor, and causes a separation of the oil.

Cooley.

CHOCOLATE MILK.

Dissolve an ounce of scraped chocolate in a pint of boiling new milk.

Cooley.

COFFEE MILK.

Boil a dessertspoonful of ground coffee in a pint of milk, for a quarter of an hour; then clear it with white of egg or isinglass, let it boil for a few minutes, and set it by the side of the fire to fine. Sweeten according to taste.

This is a suitable breakfast for those of spare habit, and disposed to affections of the lungs.

Cooley.

RICE CUSTARD.

Boil half a cupful of the best ground rice in a pint of milk, until dissolved; then mix it with a quart of cream; flavor with nutmeg, mace, and a little brandy.

Cooley.

FRUMENTY.

Bruised wheat, boiled in water until quite soft; drain, thin with milk, sweeten, and flavor according to taste.

The bruised wheat boiled to a paste, and, when cold, eaten with milk, in the evening, for some time, will often relieve costiveness.

GINGER BEER.

Take of white sugar, three pounds; bruised ginger, two ounces; cream of tartar, one ounce; lemons, sliced, four; boiling water, four gallons; yeast, eight ounces.

Pour the water on the four first-named ingredients, and infuse for two hours; then strain, add the yeast, and, when fermentation has continued for some hours, put into stone bottles and tie down the corks.

Gray.

SPRUCE BEER.

Take of sugar, six pounds; essence of spruce, four ounces; boiling water, ten gallons; yeast, eight ounces.

Add the water to the sugar and essence, ferment with the yeast, and bottle in the same way as ginger beer.

Gray.

MOLASSES BEER.

Take of molasses, fourteen pounds; hops, a pound and a half; water, thirty-six gallons; yeast, a pound.

Boil the hops in the water, add the molasses, and ferment.

Gray.

CAUDLE.

Into a pint of thin gruel, put, while it is boiling hot, the yolk of an egg, beaten with sugar, and mixed with a tablespoonful of cold water, a glass of wine, and some nutmeg.

Mix well together.

A nourishing, restorative mixture, given during convalescence.

Gray.

BARLEY MIXTURE.

Take of pearl barley, sliced figs, stoned raisins, each, two ounces and a half; liquorice root, sliced and bruised, five drachms; water, five pints and a half.

Clean the barley by washing, boil it in four and a half pints of the water, down to two pints; add the figs, raisins, and liquorice root, with the remaining pint of water; boil down to two pints, and strain.

This is nutritive and demulcent.

Gray.

ARTIFICIAL ASSES' MILK.

Boil together a pint of water and an ounce of hartshorn shavings, until reduced to a jelly; add two ounces of white sugar; when cold, mix with a pint of new milk, and a teaspoonful of syrup of Tolu.

To be taken freely, as a nutritive beverage.

Cooley.

ISINGLASS BLANCMANGE.

Isinglass, a quarter of a pound; rose water, half a pint; milk, two quarts; milk of almonds, half a pint.

Boil to a proper consistence, and permit to cool.

Hoffmann.

RICE BLANCMANGE.

Ground rice, two ounces; milk, one

pint; lump sugar, three ounces; a little lemon peel and cinnamon.

Dissolve the rice in the milk, by boiling, reduce it to a proper consistence, then add the spice and sugar; boil for a few minutes, strain, and let cool. The rice should be rubbed up with a little water, before adding it to the milk, to prevent it from being in lumps.

Cooley.

SODA CAKES.

Flour, one pound; bicarbonate of soda, a quarter of an ounce; sugar and butter, each, half a pound; make a paste with milk; add candied orange, lemon, or citron, or the fresh peel, grated, according to taste.

They may be made with the same quantity of carbonate of magnesia, instead of the soda, and the candied peel omitted. In dyspepsia, with acidity.

Cooley.

LIST OF INCOMPATIBLES.

A complete list of all the incompatibles to each medicinal agent would swell the catalogue to an inordinate degree. The following is only intended to present the chief incompatibles.

It should also be borne in mind that clinical experiments have not yet been sufficiently numerous, to determine whether the substances capable of producing precipitation in vegetable solutions, are really incompatible with the active medicinal principles of the plants in question.

- | | |
|--|---|
| <p>ABSINTHIUM, with sulphates of iron and zinc, acetates of lead, nitrate of silver, tartar emetic.</p> <p>ACACIA, with Goulard's extract, alcohol, nitric acid, muriated tincture of iron.</p> <p>ACIDUM ACETICUM, with alkalies, alkaline and earthy carbonates.</p> <p>ACIDUM CITRICUM, with mineral acids, acetates of lead, nitrate and acetate of mercury, alkalies, alkaline sulphurets.</p> <p>ACIDUM ARSENIOSUM, with magnesia, lime water, hydrosulphate of potassa, hydrated peroxide of iron, astringent vegetable infusions and decoctions.</p> <p>ACIDUM GALLICUM, with lime water, alkaline carbonates, acetate of lead, sulphate of copper, nitrate of silver, iodide of iron, sulphate of iron, tartar emetic, solution of opium, &c.</p> <p>ACIDUM HYDROCYANICUM, with mineral acids, metallic oxides, chlorine, &c.</p> <p>ACIDUM MURIATICUM, with alkalies and their carbonates, alkaline earths, metallic oxides, sulphuret of potassium, tartrate of potassa, and most metallic salts, especially those of silver.</p> <p>ACIDUM NITRICUM, with the metallic oxides, the salifiable bases, the essential oils, &c.</p> <p>ACIDUM NITRO-MURIATICUM, with oxides, earths and alkalies, the sulphurets, &c.</p> <p>ACIDUM PHOSPHORICUM, with the soluble salts of lime, baryta, and lead.</p> <p>ACIDUM SULPHURICUM, with the earths,</p> | <p>alkalies, and their carbonates, the sulphurets, &c.</p> <p>ACIDUM TANNICUM, with per salts of iron, albumen, gelatine, alkalies, alkaline earths, and carbonates, tartar emetic, acetate of lead, vegetable alkaloids, &c.</p> <p>ACIDUM TARTARICUM, with alkalies and their carbonates, and the alkaline earths and carbonates.</p> <p>ÆTHER HYPONITROSUS, with alcoholic solution of caustic potassa.</p> <p>ÆTHER MURIATICUS, with solution of caustic potassa.</p> <p>ALUMEN, with the alkalies and alkaline carbonates, lime, magnesia, acetate of lead, infusion of galls, &c.</p> <p>AMMONIA, with acids, mineral salts, alum, &c.</p> <p>AMMONIÆ ACETAS, with alkalies, strong acids, corrosive sublimate, nitrate of silver, alkaline earths, &c.</p> <p>AMMONIÆ CARBONAS, with acids, caustic potassa and soda, magnesia, alum, chloride of calcium, bitartrate and bisulphate of potassa, the salts of iron, bichloride of mercury, salts of lead, sulphate of zinc, &c.</p> <p>AMMONIÆ MURIAS, with sulphuric and nitric acids, salts of lead and silver, potassa, soda, the carbonates of potassa and soda, lime, &c.</p> <p>ANGUSTURA, with sulphates of iron and copper, nitrate of silver, tartar emetic, acetates of lead, bichloride of mercury, potassa, infusion of galls, &c.</p> |
|--|---|

- ANTHEMIS, with solution of gelatine, infusions containing gallic acid, salts of iron, nitrate of silver, salts of lead, bichloride of mercury, &c.
- ANTIMONII SULPHURETUM, with nitric and nitro-muriatic acids.
- ANTIMONII ET POTASSÆ TARTRAS, with alkalies and earths, and their carbonates, strong acids, hydrosulphurets, lime water, chloride of calcium, salts of lead, soaps, infusions containing gallic acid, rhubarb, &c.
- ARMORACIA, with carbonates of the alkalies, bichloride of mercury, nitrate of silver, vegetable bitters and astringents, &c.
- ARGENTI NITRAS, with the fixed alkalies, sulphuric, muriatic, and arsenious acids, and their salts, lime, the chlorides and sulphurets, astringent vegetable infusions, solutions of the salts of mercury and copper, &c.
- ARNICA, with sulphates of iron and zinc, acetate of lead, mineral acids, &c.
- AURANTII CORTEX, with infusion of bark, sulphate of iron, lime-water, &c.
- BARYTÆ MURIAS, with the alkaline and earthy carbonates, alum, nitrate of silver, &c.
- BENZOINUM, with the acids and alkalies.
- BISTORTA, with salts of iron, gelatine, &c.
- BELLADONNA, with caustic alkaline solutions, tannin, vegetable astringents, &c.
- BISMUTHI NITRAS, with the alkalies, vegetable astringents, &c.
- CALAMUS, with acetate of lead.
- CALCII CHLORIDUM LIQUOR, with the soluble sulphates, carbonates of soda and potassa, and carbonate of magnesia, &c.
- CALCIS LIQUOR, with the mineral and acetic, phosphoric, tartaric, and citric acids, muriate of ammonia, the alkaline carbonates, soap, the vegetable astringents, alum, sulphates of iron and zinc, sulphate of magnesia, chlorides of mercury, nitrate of silver.
- CALCIS CARBONAS, with the acids and acidulous salts, alum, muriate of ammonia.
- CAPSICUM, with corrosive sublimate, acetate of lead, nitrate of silver, sulphates of iron, zinc, and copper, carbonates of the alkalies.
- CARDAMOMUM, with the acids, sulphate of iron, bichloride of mercury, &c.
- CARYOPHYLLUS, with tartar emetic, sulphates of iron and zinc, &c.
- CASCARILLA, with lime water, sulphates of iron and zinc, infusions containing tannic or gallic acid.
- CASSIA FISTULA, with alcohol.
- CATECHU, with alkalies, the salts of iron, gelatine, &c.
- CINCHONA, with strong acids, alkalies, sulphates of iron and zinc, nitrate of silver, tartar emetic, lime, magnesia, &c.
- COCCUS, with acetate of lead, sulphates of zinc and iron.
- COLCHICUM, with acids, which render the vinous tincture drastic; alkalies, on the contrary, render it milder in its operation.
- COLOCYNTHIS, with fixed alkalies, sulphate of iron, nitrate of silver, acetate of lead, &c.
- COLOMBA, with ammonia, lime water, mineral acids, muriate of iron, nitrate of silver, acetate of lead, isinglass.
- CONIUM, with the strong acids, alkalies, tannin, &c.
- CONTRAYERVA (tincture), with water.
- COPAIBA, with the mineral acids.
- COPTIS TRIFOLIA, with nitrate of silver, and acetate of lead.
- CRETA PRÆPARATA, with acids and acidulous salts, alum, muriate of ammonia.
- CUPRUM AMMONIATUM, with acids, potassa and soda, lime water.
- CUPRI SULPHAS, with fixed alkalies, ammonia, and its carbonate, bichloride of mercury, arsenite of potassa, nitrate of silver, acetate of lead, vegetable infusions, &c.
- CYDONIA, with acids, most metallic salts, alcohol.
- DIGITALIS, with sulphate of iron, infusion of Peruvian bark, acetate of lead, tannin, and vegetable astringents.

- FERRUM AMMONIATUM, with acids, the fixed alkalies, lime-water, astringent infusions.
- FERRI IODIDUM, with fixed alkalies, lime-water, vegetable astringents, &c.
- FERRI ET POTASSÆ TARTRAS, with the mineral acids, alkalies, vegetable astringents.
- FERRI SUBCARBONAS, with the mineral acids, acidulous salts, &c.
- FERRI CHLORIDI, TINCTURA, with alkalies and their carbonates, lime-water, carbonate of lime, magnesia, and its carbonate, solution of gum, vegetable astringents.
- FERRI SULPHAS, with nitric acid, fixed alkalies, and their carbonates, lime-water, nitrates of potassa and of silver, borate of soda, acetate of lead, iodide of potassium, vegetable astringents, &c.
- GALLA, with alkalies, the carbonates of the alkalies, lime-water, sulphates of iron and zinc, acetate of lead, tartar emetic, bichloride of mercury, gelatin, vegetable alkaloids, &c.
- GUAIACI TINCTURA, with water, the mineral acids, spirit of nitric ether, earthy and metallic salts, &c.
- GRANATUM (*Cortex*), with sulphate of iron, &c.
- HÆMATOXYLON, with mineral acids, alum, sulphates of iron and copper, tartar emetic, acetate of lead.
- HYDRARGYRUM AMMONIATUM, with muriatic and other acids, the fixed alkalies and protochloride of tin.
- HYDRARGYRI CHLORIDUM MITE, with the alkalies, lime, muriate of ammonia, chlorides of potassium and sodium, carbonates of the alkalies, nitric acid, salts of iron, lead, and copper, iodide of potassium, soaps, &c.
- HYDRARGYRI CHLORIDUM CORROSIVUM, with alkalies and their carbonates, carbonates of lime, tartar emetic, sulphuret of potassium, soap, iron, copper, lead, and their salts, nitrate of silver, albumen, gelatin, gluten, milk, vegetable astringents, fixed oils, &c.
- HYDRARGYRI OXIDUM RUBRUM, with the mineral acids.
- HYDRARGYRI OXIDUM NIGRUM, with the mineral and acetic acids.
- HYDRARGYRI IODIDUM, with the mineral acids, chloride of sodium.
- HYDRARGYRI IODIDUM RUBRUM, with the mineral acids, iodide of potassium, chloride of sodium.
- HYDRARGYRUM CUM CRETA, with the mineral and acetic acids, acidulous salts, alum, &c.
- HYOSCYAMUS, with acetate of lead, nitrate of silver, sulphate of iron, tannin, and the vegetable astringents.
- IODINE, with starch, and magnesia
- IPECACUANHA, with the vegetable astringents, acetate of lead, &c.
- KINO, with the salts of iron, acetate of lead, the mineral acids, gelatin, tartar emetic, &c.
- KRAMERIA, with salts of iron, acetate of lead, gelatin, mineral acids, &c.
- MAGNESIA, with acids and acidulous salts, muriate of ammonia, metallic salts.
- MAGNESIÆ CARBONAS, with acids and acidulous salts, metallic salts, lime-water, muriate of ammonia, bitartrate of potassa.
- MAGNESIÆ SULPHAS, with ammonia, acetate of lead, chloride of calcium, lime-water, potassa and soda, and their carbonates, &c.
- MENTHA, with sulphate of iron, nitrate of silver, acetate of lead, &c.
- MORPHIÆ ACETAS, with alkaline carbonates, ammonia, vegetable astringents, all articles incompatible with infusion of opium, except acetate of lead.
- MORPHIÆ MURIAS, with acetate of lead, tannin, ammonia, alkaline carbonates, &c.
- MOSCHUS, with mineral acids, bichloride of mercury, sulphate of iron, nitrate of silver, infusion of bark, &c.
- OPIUM, with ammonia, carbonates of potassa and soda, nitrate of silver, acetate of lead, salts of copper, iron, and zinc, astringent infusions, &c.

- POTASSA, with acids and acidulous salts, earthy and metallic salts, ammonia and its salts, &c.
- POTASSÆ ACETAS, with mineral acids, sulphates of soda and magnesia, tartaric acid, most metallic and earthy salts, &c.
- POTASSÆ ARSENITIS, LIQUOR, with lime-water, acids, chlorides of iron and calcium, sulphate of magnesia, alum, sulphates of iron and copper, iodide of iron, nitrate of silver, vegetable astringents.
- POTASSÆ BITARTRAS, with strong acids, lime-water, ammonia, carbonates of potassa and soda, magnesia, sulphate of magnesia, &c.
- POTASSÆ CARBONAS, with acids, metallic salts, lime-water, sulphate of magnesia, muriate of ammonia, alum, calomel, &c.
- POTASSÆ CITRAS, with salts of lime, lead, and silver, sulphuric and other acids.
- POTASSÆ NITRAS, with alum, sulphate of magnesia, metallic sulphates, sulphuric acid, muriatic acid (when heated).
- POTASSÆ SULPHAS, with tartaric acid, acetate of lead, nitric and muriatic acids, chloride of calcium, bichloride of mercury, nitrate of silver, &c.
- POTASSII SULPHURETUM, with acids, and most metallic salts.
- POTASSÆ TARTRAS, with most acids, lime-water, acetate of lead, nitrate of silver, chloride of calcium.
- POTASSII IODIDUM, with acetate of lead, bichloride of mercury, tartaric acid, metallic salts, all acids and acidulous salts, except bitartrate of potassa.
- PIMENTA, with alum, ammonia, alkaline carbonates, salts of iron, copper, zinc, and silver, vegetable astringents.
- PIPER NIGRUM, with vegetable astringents.
- PLUMBI ACETAS, with the alkalies and their carbonates, tartaric, citric, and sulphuric acids, tartrate and bitartrate of potassa, alkaline and metallic sulphates, alum, borax, lime-water, vegetable astringents, infusion of opium, the soaps, milk.
- PLUMBI IODIDUM, with solution of potassa, sulphuric acid.
- QUASSIA, with nitrate of silver, acetate of lead, &c.
- QUININÆ SULPHAS, with tartaric acid, tartrate of potassa, alkalies and their carbonates, lime-water, infusion of galls.
- RHEUM, with strong acids, lime-water, sulphates of iron and zinc, tartar emetic, bichloride of mercury, vegetable astringents.
- ROSA GALLICA, with sulphates of iron and zinc, gelatin, lime-water.
- SALIX, with lime-water, sulphate of iron, alkaline carbonates, solution of isinglass.
- SALVIA, with salts of iron.
- SAPO (*and liniments, &c., containing it*), with sulphates of lime and magnesia, chloride of calcium, alum, metallic salts, lime-water.
- SARSAPARILLA, with infusion of galls, lime water, acetate of lead.
- SCILLA, with lime-water, alkaline carbonates, nitrate of silver, acetate of lead, &c.
- SENNA, with strong acids, carbonates of the alkalies, lime-water, tartar emetic.
- SERPENTARIA, with acetate of lead.
- SODÆ CARBONAS, with acids, bitartrate of potassa, acidulous, metallic and earthy salts, lime-water.
- SODÆ PHOSPHAS, with mineral acids, lime, magnesia, &c.
- SODÆ SULPHAS, with salts precipitated by sulphuric acid, carbonates of potassa, acetate of potassa.
- SODÆ ET POTASSÆ TARTRAS, with most acids and acidulous salts, acetate of lead, nitrate of silver.
- SPIRITUS ÆTHERIS NITRICI, with sulphate of iron, tincture of guaiacum, alkaline and earthy carbonates, especially those with an excess of acid.
- TAMARINDUS, with the salts of potassa, the alkaline carbonates, lime-water, tartar emetic.
- TARAXACUM, with corrosive sublimate, sulphate of iron, nitrate of silver, acetate of lead, infusion of galls, &c.

THEA, with salts of iron, gelatin, lime-water, &c.	UVA URSI, with salts of iron, gelatin, tartar emetic.
TORMENTILLA, with solution of isinglass, salts of iron, alkalies, &c.	ZINCI OXIDUM, with acids, acidulous salts, potassa, soda, and ammonia.
TRAGACANTH, with sulphates of iron and copper, acetate of lead, alcohol, &c.	ZINCI SULPHAS, with potassa, soda, and ammonia, and their carbonates, the hydrosulphurets, milk, mucilages, astringent vegetable infusions, &c.
ULMUS, with alcoholic tinctures, if added in quantity.	

POSOLOGICAL TABLE

OF THE

MOST IMPORTANT MEDICINES.

Absinthium, Infusion, f℥j to ij. Extract, grs. x to xx. Tincture, gtt. xx to f℥ij. Oil, gtt. ij to iv.	Æther, Muriatic, gtt. xxx to lx. Sulphuric, f℥ss to j. spirit, f℥j to iij. compound, f℥ss to ij.
Acetosella, Extract, ℥j to ℥ss.	Allium, Syrup, f℥j.
Acidum Arseniosum, gr. $\frac{1}{16}$ to $\frac{1}{8}$.	Aloe, Powder, gr. v to x. and Canella, grs. v to xx.
Benzoicum, grs. x to xx.	Wine, f℥j. to f℥ss.
Boracicum, grs. v to xx.	Tincture, f℥j. to f℥ss. and Myrrh, f℥ss to ij.
Gallicum, grs. ij to x.	Alumen, grs. v to xx.
Hydrocyanicum (medicinal), gtt. j to ii.	Ammonia, Water, gtt. v to xx.
Hydriodicum, gtt. v to x.	Acetate, Solution of, f℥ss to j.
Muriaticum, gtt. v to x. dilat. gtt. xx to xxx.	Carbonate, grs. v to x.
Nitricum, gtt. ij to vj. dilat. gtt. xx to xxx.	Tincture comp., gtt. x to xl.
Phosphoricum, dilat. gtt. x to f℥j.	Spirit of, gtt. v to xx. aromatic, gtt. x to xx.
Sulphuricum, gtt. ij to v. dilat. gtt. x to xxx. aromatic, gtt. v to xx.	Citrate, Solution of, f℥ss to j.
Tannicum, gr. j to iij.	Hydrosulphate, gtt. v to vj.
Tartaricum, ℥j to ij.	Muriate, gr. j to v.
Aconitum, Powder, gr. j to ij. Extract, gr. j. alcoholic, gr. $\frac{1}{8}$ to $\frac{1}{4}$.	Nitrate, ℥ss to j.
Tincture of root, gtt. v to vj.	Phosphate, grs. x to xl.
Tincture of leaves, gtt. x to xv.	Succinate, Spirit, gtt. xx to l.
Æther, Acetic, gtt. x to f℥j.	Sulphate, ℥j. to ℥ss.
Hyponitrosus, gtt. x to lx.	Ammoniacum, grs. v to xv. mixture, f℥ss to j.
Nitric, Spirit of, f℥ss to ij.	Amylum, Iodide, ℥ss to j.
Hydrocyanic, gtt. ij to iij.	Angelica, Tincture, f℥j to ij.
	Angustura, Powder, grs. x to ℥j. Infusion, f℥ij. Tincture, f℥j to ij.

<p>Anthemis, Infusion, f 3j to ij. Extract, grs. x to xx. Syrup, f 3ss.</p> <p>Anthracokali, grs. ij to iv.</p> <p>Antimonium, grs. x to xx. Sulphuret, grs. v to x. precipitated, grs. j to iij. Kermes mineral, gr. $\frac{1}{2}$ to x. Potassa et, Tartrate of, gr. $\frac{1}{8}$ to j. Wine, gtt. xxx to f 3j. Powder, grs. iij to x.</p> <p>Apocynum, Decoction, f 3j to ij. Extract, grs. iij to v.</p> <p>Argentum, Chloride, gr. $\frac{1}{10}$ to gr. ij. and Ammonia, gr. $\frac{1}{4}$ to $\frac{1}{10}$. Cyanide, gr. $\frac{1}{12}$ to $\frac{1}{8}$. Iodide, gr. $\frac{1}{2}$ to j. Nitrate, gr. $\frac{1}{4}$ to ij. Oxide, gr. $\frac{1}{4}$ to ij.</p> <p>Armoracia, Infusion, f 3j to ij. Spirit. comp. f 3j to iv.</p> <p>Arnica, Powder, grs. v to x. Extract, grs. v to x. Infusion, f 3ss to j. Tincture, gtt. xxx to l.</p> <p>Arsenicum, Iodide, gr. $\frac{1}{10}$ to $\frac{1}{8}$. Donovan's Solution, gtt. v to xx.</p> <p>Arum, Powder, grs. x.</p> <p>Asclepias Tuberosa, Infusion, f 3ij to iv.</p> <p>Asparagus, Extract, 3j to 3j. Syrup, 3j to ij.</p> <p>Assafoetida, grs. ij to x. Mixture, f 3ss to j. Tincture, f 3j. ammoniated, gtt. v to f 3j.</p> <p>Atropia, gr. $\frac{1}{30}$.</p> <p>Aurum, gr. $\frac{1}{4}$ to j. Chloride, gr. $\frac{1}{20}$ to $\frac{1}{16}$. Sodium et, Chloride, gr. $\frac{1}{20}$ to $\frac{1}{10}$. Cyanide, gr. $\frac{1}{18}$ to $\frac{1}{10}$. Iodide, gr. $\frac{1}{20}$ to $\frac{1}{10}$. Oxide, gr. $\frac{1}{10}$ to $\frac{1}{4}$.</p> <p>Azederach, Decoction, f 3ss to j.</p> <p>Ballotta Lanata, Decoction, f 3j to ij.</p> <p>Balsamum Peruvianum, f 3ss.</p> <p>Balsamum Tolutanum, gr. x to xxx. Syrup, f 3ss. Tincture, f 3j.</p> <p>Baptisia Tinctoria, Decoction, f 3ss.</p> <p>Barium, Chloride, Solution, gtt. v. Iodide, gr. $\frac{1}{8}$.</p> <p>Bebeerina, Sulphate, grs. v to 3j.</p> <p>Belladonna, gr. j to ij.</p>	<p>Belladonna, Extract, gr. $\frac{1}{4}$. alcoholic, gr. $\frac{1}{2}$ to j. Syrup, 3j to ij. Tincture, gtt. xv to xxx.</p> <p>Benzoin, Compound Tincture, gtt. x to xx.</p> <p>Bismuthum, Subnitrate, grs. ij to x.</p> <p>Bistort, Infusion, f 3j to ij.</p> <p>Boletus Laricis, gr. j to iij.</p> <p>Brominium, Solution, gtt. v to x.</p> <p>Brucia, gr. $\frac{1}{8}$ to $\frac{1}{2}$. Tincture, gtt. v to xx.</p> <p>Buchu, gr. xx to xxx. Infusion, f 3ij.</p> <p>Cahinea, 3j to 3j.</p> <p>Calamus, Infusion, f 3j to ij.</p> <p>Calcium, Chloride, Solution, gtt. xx to xxx. Iodide, grs. ij to iij.</p> <p>Calx. Lime-water, f 3ss to ij.</p> <p>Calcis Carbonas, Precipitated, grs. x to xx. Creta præparata, gr. x to xxx. Testa præparata, gr. x to xxx. Phosphate, 3j to ij.</p> <p>Calendula, Extract, grs. ij to v.</p> <p>Calomel, as an alterative, gr. $\frac{1}{10}$ to $\frac{1}{2}$. as a purgative, gr. v. to xij.</p> <p>Calotropis, Infusion, f 3j to f 3j. Powder, grs. iij to xx.</p> <p>Camphora, grs. iij to x. Water, f 3ss to j. Tincture, gtt. v to xx.</p> <p>Canella, grs. x to 3j.</p> <p>Cannabis, Extract, gr. $\frac{1}{2}$ to iij. Tincture, gtt. x to xl.</p> <p>Cantharis, gr. $\frac{1}{2}$ to j. Tincture, gtt. xx to f 3j.</p> <p>Capsicum, grs. v to x. Infusion, f 3ss. Tincture, f 3ss to ij.</p> <p>Carbo Animalis, grs. x to xx. Ligni, 3ss to j. Mineralis, (Anthracokali,) grs. ij to iv.</p> <p>Cardamomum, Tincture, f 3j to ij.</p> <p>Carota, Infusion, seeds, f 3ij to iv.</p> <p>Carum, Spirit, f 3j to ij.</p> <p>Caryophyllus, Powder, grs. v to x. Infusion, f 3j to ij. Tincture, f 3j to ij. Oil, gtt. ij to v.</p> <p>Cascarilla Powder, 3j to 3ss. Infusion, f 3j to ij. Extract, grs. x. to xx. Tincture f 3j.</p>
---	--

- Cassia Fistula, ʒij to ʒss .
 Castoreum, grs. v to xx.
 Tincture, gtt. xxx to fʒij .
 Catalpa, Decoction, fʒij to ij .
 Catechu, grs. x to ʒj .
 Infusion, comp. fʒj to ij .
 Tincture, gtt. xxx to fʒij .
 Centaurea Benedicta, ʒj to ʒj .
 Infusion, fʒj to ij .
 Cetraria, ʒss to j.
 Decoction, fʒij .
 Cetrarine, grs. ij to v.
 Chelidonium, Extract, grs. v to xv.
 Chenopodium, Juice, fʒss .
 Oil, gtt. iv to x.
 Chimaphila, Decoction, fʒij to iv.
 Extract, grs. x to xxx.
 Chiretta, ʒj .
 Chlorinium, Water, fʒj to iv.
 Chloroformum, gtt. v to xx.
 Cimicifuga, Decoction, fʒj to ij .
 Tincture, gtt. xx.
 Cinchona, Powder, ʒss to j.
 Extract, grs. x to xxx.
 Decoction, fʒj .
 Infusion, fʒij .
 Tincture, fʒj to iv.
 compound, fʒj to iv.
 Cinchonia, grs. ij to x.
 Sulphate, gr. ij to x.
 Cinnamomum, Powder, grs. x to ʒj .
 comp. grs. x to xxx.
 Tincture, fʒj to iv.
 Water, fʒss .
 Codeia, Syrup, fʒj .
 Morphia and, Muriate, gr. $\frac{1}{4}$ to $\frac{1}{2}$.
 Colchicum, Powder, grs. ij to viij.
 Extract, bulb, gr. j. to ij .
 Wine, root, gtt. x. to xx.
 seeds, fʒj .
 Tincture, seeds, fʒss to ij .
 Colocynthis, Powder, grs. v to x.
 Extract, grs. v to x.
 " comp. grs. v to xv.
 Colomba, Powder, grs. x to xxx.
 Infusion, fʒij .
 Tincture, fʒj to iv.
 Extract, grs. j to v.
 Conium, Powder, grs. ij to v.
 Extract, gr. j to ij .
 " alcoholic, gr. j to ij .
 Tincture, gtt. xx to fʒj .
 Contrayerva, Powder, ʒss .
 Extract, ʒj .
 Tincture, gtt. xxx to xl.
 Copaiba, grs. xx to ʒj .
 Tincture, gtt. xxx to fʒj .
 Copaiba, Oil, gtt. x to xxx.
 Coptis, Powder, grs. x to xxx.
 Infusion, fʒss to ij .
 Tincture, fʒj to ij .
 Cornus, Powder, ʒj to ʒj .
 Creasotum, gtt. j to ij .
 Crocus, grs. x to xxx.
 Cubeba, Powder, grs. xx to ʒij .
 Tincture, fʒj to ij .
 Oil, gtt. v to x.
 Fluid extract, gtt. x to xxx.
 Cupri sulphas, (as emetic), gr. j to ij .
 Cuprum Ammoniatum, gr. $\frac{1}{4}$ to $\frac{1}{2}$.
 Delphinium, Tincture, gtt. x to xx.
 Delphinia, gr. $\frac{1}{4}$ to $\frac{1}{2}$.
 Digitalis, Powder, gr. $\frac{1}{2}$ to j.
 Infusion, fʒss .
 Tincture, gtt. x to xx.
 Digitalina, gr. $\frac{1}{16}$ to $\frac{1}{8}$.
 Diosma, Powder, grs. xx to xxx.
 Infusion, fʒj to ij .
 Tincture, fʒj to ij .
 Extract, fluid, fʒj to ij .
 Diospyros, Infusion, fʒj .
 Dulcamara, Powder, grs. xxx to ʒj .
 Decoction, fʒj to ij .
 Extract, grs. v to x.
 Elaterina, gr. $\frac{1}{16}$.
 Tincture, gtt. xx to xl.
 Elaterium (common), gr. j to ij .
 Clutterbuck's, gr. $\frac{1}{8}$ to $\frac{1}{4}$.
 Emetia, gr. $\frac{1}{16}$ to j.
 Syrup, fʒj .
 Ergota, Powder, grs. x to xxx.
 Infusion, fʒj .
 Wine, fʒij to ij .
 Tincture, gtt. xx to fʒj .
 Extract (Ergotine), gr. ij .
 Oil, gtt. xx to l.
 Erigeron Annuum, Infusion, fʒij to iv.
 Erigeron Canadense, Infusion, fʒij to iv.
 Extract, grs. v to x.
 Eupatorium, Powder, grs. xx to xxx.
 Infusion, fʒj .
 Euphorbia Corollata, Powder, grs. x to xx.
 Ipecacuanha, Powder, grs. x to xv.
 Hypericifolia, Infusion, fʒss to ij .
 Lathyris, Oil, gtt. iv to xii.
 Ferrum, Ammoniated, grs. iv to xii.
 Tincture, gtt. xl to fʒj .

Ferri Pulvis, gr. ij to x.	Granatum, Decoction (rind), f 3j.
Ferrum, Acetate, gtt. x to xxv.	“ (bark), f 3ij to iv.
Tincture, gtt. xxx to f 3j.	Gratiola, Wine, f 3j.
Arseniate, gr. $\frac{1}{16}$ to $\frac{1}{10}$.	Guaiacum, Resin, grs. x to ʒj.
Bromidum, grs. j to iij.	Wood, decoction, f 3j to ij.
Carbonate, grs. x to 3ij.	Tincture, f 3j to ij.
“ Saccharine, grs. x to xx.	Ammoniated, f 3j to ij.
Carburet, grs. v to xv.	Hæmatoxylon, Infusion, f 3ss to ij.
Chloride, gr. j to ij.	Decoction, f 3ss to ij.
Tincture, gtt. x to xxx.	Extract, grs. x to xxx.
Citrate, grs. iv to viij.	Helleborus, Powder, grs. x to ʒj.
“ and Quinia, grs. v to x.	Extract, grs. v to xv.
Ferrocyanuret, grs. iij to v.	Tincture, gtt. xxx to f 3j.
Iodide, grs. ij to iij.	Humulus, Infusion, f 3j to ij.
Solution, gtt. x to f 3j.	Tincture, f 3j to iij.
Syrup (Lond.), f 3j.	Hydrargyrum,
Lactate, grs. ij to iv.	with Antimony, gr. j to iv.
Malate, grs. v to ʒj.	“ Chalk, grs. v to ʒj.
Nitrate, gtt. vj to xii.	“ Magnesia, grs. v to ʒj.
Persesquintrate, gtt. x to xii.	Blue pill, grs. v to x.
Oxide, Black, grs. v to xx.	Acetate, gr. j.
“ Red, grs. v to xv.	Borate, gr. ij.
Phosphate, grs. v to x.	Bromide, gr. j to v.
Sulphate, grs. ij to v.	Bibromide, gr. $\frac{1}{10}$ to $\frac{1}{10}$.
“ dried, gr. j to iij.	Chloride, Corrosive, gr. $\frac{1}{16}$ to $\frac{1}{4}$.
Tartrate, grs. v to x.	Chloride, Mild, as an alterative, gr. $\frac{1}{10}$ to $\frac{1}{2}$.
Wine, f 3j.	as a purgative, gr. v. to xij.
Ammonio, grs. v.	Cyanuret, gr. $\frac{1}{16}$ to $\frac{1}{8}$.
Tannate, gr. ij to iij.	Iodide, gr. $\frac{1}{4}$ to j.
Valerianate, grs. j to ij.	“ Red, gr. $\frac{1}{16}$ to $\frac{1}{10}$.
Wine, f 3j.	Oxide, Black, gr. j to ij.
Filix Mas, 3j to ij.	“ Red, gr. $\frac{1}{2}$ to j.
Extract, grs. xv to xx.	Phosphate, gr. $\frac{1}{2}$ to j.
Frasera, Powder, grs. xxx to 3j.	Sulphate, gr. $\frac{1}{4}$ to v.
Infusion, f 3j to ij.	Sulphuret, Black, grs. v to x.
Fuligo, Tincture, gtt. xxx to f 3j.	Sulphuret, Red, grs. x to xv.
Fuligokali, gr. ij to iij.	Tartrate, gr. j to ij.
Galbanum, grs. x to xx.	Hippocastanum, Powder, ʒj to iv.
Galla, Powder, grs. v to xxx.	Hyoseyamus, Powder, grs. ij to v.
Infusion, f 3j to ij.	Extract, gr. j.
Tincture, f 3j to iij.	Tincture, gtt. xx to f 3ss.
Syrup, f 3ss.	
Gambogia, gr. j to vj.	Indigum, Powder, ʒj to ij.
Solution, alkaline, gtt. xv.	Iodinium, gr. $\frac{1}{8}$ to j.
Gaultheria, Oil, gtt. ij to x.	Tincture, gtt. x to xx.
Gentiana, Powder, grs. x to xl.	Compound tincture, gtt. xv to xxx.
Extract, grs. x to ʒj.	Lugol's Solution, gtt. v to x
Infusion, f 3j.	
Tincture, comp., f 3j to ij.	
Geranium, grs. x to xxx.	
Extract, grs. x to xx.	
Geoffroya, ʒj to 3ss.	
Decoction, f 3j to ij.	
Gillenia, grs. xx to xxx.	

Ipecacuanha, Powder, grs. xv to xxx.	Mentha Viridis, Infusion, f 3j to ij.
“ comp. grs. v to x.	Oil, gtt. ij.
Syrup, f 3j to ij.	Menyanthes, Powder, grs. xx to xxx.
Wine, f 3j to f 3j.	Infusion, f 3j to ij.
Jalapa, Powder, grs. x to xxx.	Extract, grs. x to xv.
“ comp., 3ss to j.	Mezereum, Decoction, f 3j to iv.
Extract, grs. v to x.	Monarda, Oil, gtt. j to ij.
“ alkaline, grs. iij to ix.	Monesia, grs. ij to x.
Soap, grs. x to xv.	Syrup, f 3ss.
Tincture, f 3j to ij.	Morphia, gr. $\frac{1}{2}$ to $\frac{1}{4}$.
Juglans Cinerea, Extract, grs. v to xxx.	Acetate, gr. $\frac{1}{2}$ to $\frac{1}{4}$.
Juniperus, Extract, 3j to iij.	Bimeconate, gr. $\frac{1}{4}$.
Infusion, f 3ij to iv.	Muriate, gr. $\frac{1}{2}$ to $\frac{1}{4}$.
Oil, gtt. iij to vj.	Sulphate, gr. $\frac{1}{2}$ to $\frac{1}{4}$.
Kino, Powder, grs. x to xx.	Moschus, grs. v to x.
Tincture, f 3ss to ij.	Tincture, gtt. xxx to f 3j.
Krameria, Powder, grs. xx to xxx.	Mucuna, Electuary, 3i to ij.
Extract, grs. x to xx.	Myroxylon, f 3ss.
Infusion, f 3i to ij.	Syrup, f 3ss to j.
Syrup, f 3ss.	Myrrha, grs. x to xxx.
Tincture, f 3j to ij.	Tincture, f 3ss to j.
Lactucarium, grs. iij.	Narcotina, grs. ij.
Tincture, f 3ss to ij.	Muriate, grs. ij to iv.
Lauro-Cerasus, Water, f 3ss to j.	Nux vomica, Powder, grs. iij to v.
Liriodendron, Powder, 3ss to ij.	Extract, gr. $\frac{1}{2}$ to ij.
Infusion, f 3j to ij.	Tincture, gtt. v to xx.
Tincture, f 3j to ij.	Oleum Animale Empyreumatic. gtt. v to x.
Lobelia, Powder, grs. v to x.	Cajuputi, gtt. ij to v.
Tincture, f 3ss to j.	Morrhuae, f 3ss to ij.
Lupulina, grs. v to x.	Olivae, f 3j.
Tincture, f 3i to ij.	Ricini, f 3ss to j.
Macis, grs. x to 3j.	Terebinthinae, gtt. v to f 3i.
Tincture, gtt. xxx to xl.	Tiglii, gtt. i to ij.
Magnesia, 3j.	Opium, gr. j.
Carbonate, 3j to 3j.	Confection, gr. x.
Citrate, Solution, f 3iv to xij.	Extract, gr. ss.
Sulphate, 3j.	Vinegar, Black drop, gtt. vii to x.
Sulphuret, Syrup, f 3ss.	Lancaster “ “ gtt. vj to x.
Phosphate, grs. x to xx.	Houlton's “ “ gtt. xx.
Magnolia, Powder, 3ss to j.	Rousseau's “ “ gtt. iv.
Manganesii, Carbonate, gr. x to 3j.	Porter's “ “ gtt. v to xx.
Muriate, grs. iv to x.	Tincture, gtt. xxv.
Oxide, grs. ij to iv.	“ acetated, gtt. xx.
Sulphate, 3ss to ij.	“ Smith's, gtt. iij to v.
Manna, 3j to ij.	“ Ammoniated, f 3ss to j.
Mannite, 3ij to 3ij.	“ Camphorated, f 3j to iij.
Marrubium, Extract, 3i to 3ss.	“ Swediaur's, gtt. v.
Matico, Infusion, f 3j.	Wine, gtt. xx.
Tincture, gtt. xxx to f 3j.	Elixir, gtt. v to xx.
Syrup, f 3j to ij.	Opoponax, grs. x to xx.
Mentha Piperita, Oil, gtt. i to iij.	Papaver, Syrup, f 3ij to f 3j.
Essence, gtt. x to xx.	Pareira Brava, Powder, grs. xxx to 3j.
Water, f 3ss.	Infusion, f 3j to ij.

Pareira Brava, Decoction, f 3j to ij.	Quinia, as an anti-intermittent, gr. viij to xx.
Extract, grs. x to 3ss.	Amorphous, gr. j to iv.
Tincture, gtt. l to lx.	Acetate, gr. j to ij.
Paullinia, grs. x to xx.	Arseniate, gr. 1/8.
Petroleum, 3ss to j.	Citrate, gr. j to ij.
Petroselinum, Infusion, f 3ij to iv.	Ferrocyanate, gr. j to ij.
Phloridzina, grs. v to xv.	Mercury and, Chloride, gr. 1/2.
Phosphorus, gr. 1/16.	Kinate, gr. j to ij.
Phytolacca, Powder, grs. x to xxx.	Lactate, gr. j to ij.
Piper Nigrum, Confection, 3j to ij.	Muriate, gr. j. to ij.
Fluid extract, gtt. j to iij.	Sulphate, gr. j to ij.
Piperinum, gr. j to ij.	Valerianate, gr. j to ij.
Piscidia Erythrina, Tincture, f 3j.	
Pix liquida, 3ss to j.	Rheum, Powder, grs. x to xxx.
Platini Bichloridum, gr. 1/8 to 1/4.	Roasted, grs. v to x.
Plumbum, Acetate, gr. j to iv.	Infusion, f 3j to ij.
Iodide, gr. 1/2 to j.	Extract, grs. x to xx.
Podophyllum, Powder, grs. x to xx.	" Fluid, gtt. xv to f 3j.
Extract, grs. v to xv.	Syrup, f 3j to f 3ss.
Podophyllin, grs. j to ij.	" Aromatic, f 3j to f 3ss.
Potassium, Bromide, grs. ij to x.	Tincture, f 3j to f 3ss.
Chloride, 3j to ij.	" and Senna, f 3j to f 3ss.
Cyanuret, gr. 1/8 to 1/4.	Wine, f 3j to iv.
Iodide, gr. j to x.	Rubia, Powder, 3ss.
Iodo-hydrargyrate, gr. 1/12.	Decoction, f 3ij.
Potassa, Acetate, 3ss to ij.	Rubus, Decoction, f 3j to ij.
Arsenite, solution, gtt. x.	Ruta, Powder, grs. x to xx.
Arseniate, gr. 1/16.	Oil, gtt. ij to iij.
Borate, gr. v to x.	
Boro-tartrate, 3ij to 3ss.	Sabadilla, Powder, grs. ij to v.
Carbonate, gr. v to xx.	Extract, gr. 1/8.
Bi-carbonate, gr. xv to xxx.	Sabbatia, Infusion, f 3ij to iv.
Chlorate, gr. x to xx.	Sabina, Powder, grs. v to x.
Citrate, solution, f 3ss.	Oil, gtt. ij to v.
Nitrate, gr. v to x.	Salicin, grs. iv to vj.
Silicate, gr. x to xv.	Salvia, Infusion, f 3ij to iv.
Sulphate, 3iv to v.	Sanguinaria, Powder, grs. x to xx.
Bisulphate, 3j to ij.	Tincture, f 3ss to ij.
" with Sulphur, 3ss to j.	Sarsaparilla, Powder, 3ss to j.
Tartrate, 3j to 3j.	Decoction, f 3iv to vj.
Bitartrate, 3j to 3ss.	Extract, grs. x to xx.
and Ammonia, Tartrate, 3j.	" Fluid, f 3j.
Prinos, Powder, grs. xxx to 3j.	Syrup, f 3ss to j.
Decoction, f 3ij.	" Compound, f 3ss to j.
Prunus Virginiana, Powder, grs. xxx to 3ij.	Sassafras, Infusion, f 3j to ij.
Infusion, f 3ij.	Oil, gtt. ij to v.
Syrup, f 3j to f 3j.	Scammonium, Powder, grs. v to x.
	Confection, grs. xx to xxx.
	Resin, grs. v to x.
Quassia, Infusion, f 3j.	Scilla, Powder, grs. j to ij.
Extract, grs. ij to v.	Syrup, f 3j.
Tincture, f 3j to ij.	" Compound, gtt. x to f 3j.
Quercus, Powder, grs. xxx to 3j.	Tincture, gtt. xx to xl.
Extract, grs. x to xx.	Extract, gr. ss to ij.
Decoction, f 3ij to iij.	Vinegar, f 3ss to j.
Quinia, as a tonic, gr. j to ij.	Oxymel, f 3j to ij.

- Scoparius, Infusion, f 3j to iv.
Extract, 3j to 3j.
- Senega, Powder, grs. x to xx.
Infusion, f 3j to ij.
Decoction, f 3ss.
Syrup, f 3j to ij.
- Senna, Powder, 3ss to ij.
Confection, 3ij.
Infusion, f 3iv.
Syrup, f 3j to ij.
Extract, fluid, f 3ss.
- Serpentaria, Powder, grs. x to xx.
Infusion, f 3j to ij.
Tincture, f 3j to ij.
- Simaruba, Infusion, f 3ij.
- Sodii Chloridum, Powder, grs. x to 3ss.
- Soda, Acetate, 3j to 3iv.
Arseniate, gr. $\frac{1}{16}$ to $\frac{1}{8}$.
Borate, grs. xx to xxx.
Carbonate, grs. x to xx.
Bi-carbonate, grs. xv to xxx.
Hydrosulphate, grs. x to 3j.
Phosphate, 3iv to 3j.
Sulphate, 3ss to j.
Tartrate, 3ss to j.
and Potassa, Tartrate, 3ij to 3ss.
- Spigelia, Powder, 3j to ij.
Infusion, f 3iv to f 3j.
Comp. f 3j to ij.
Extract, Fluid, 3j to 3ss.
- Spiraea, Decoction, f 3j to ij.
Extract, grs. v to 3j.
- Spongia, Burnt, 3j to ij.
- Stannum, Powder, 3ss.
Chloride, gr. ij.
Sulphuret, grs. x to xx.
Oxide, grs. v. to vj.
- Stramonium, Powder, gr. ij to iij.
(seeds), gr. j.
Extract (seeds), gr. $\frac{1}{4}$ to $\frac{1}{2}$.
(leaves), gr. j.
Tincture (seeds), gtt. xx to xxx.
- Styrax, grs. x to xx.
- Strychnia, gr. $\frac{1}{16}$ to $\frac{1}{10}$.
Acetate, solution, gtt. v.
Iodate, gr. $\frac{1}{8}$.
- Succinum, Oil, gtt. v to xv.
Tincture, gtt. xl to lx.
- Sulphur, Precipitated, 3ss to ij.
Electuary, 3j to ij.
- Sulphuris Carburetum, gtt. ij to iij.
- Tabacum, Wine, m x to xx.
Tincture, m x to xx.
- Tanacetum, Extract, grs. v to 3j.
- Taraxacum, Decoction, f 3j to iij.
- Taraxacum, Extract, 3j to 3j.
" Fluid, f 3j.
- Terebinthina, grs. ij to v.
Oil, gtt. v to f 3j.
- Testa præparata, gr. x to xxx.
- Tolutanum, grs. x to xxx.
Tincture, f 3j to ij.
Syrup, f 3j to f 3ss.
- Tormentilla, Powder, grs. xxx to 3j.
Decoction, f 3j to ij.
- Toxicodendron, Powder, gr. $\frac{1}{2}$ to ij.
Extract, gr. j.
- Tussilago, Decoction, f 3ij to iv.
Syrup, f 3j to f 3ss.
- Ulmus Campestris, Decoction, f 3ij to iv.
- Ulmus Fulva, Decoction, f 3iv to vj.
- Uva Ursi, Powder, 3j to 3j.
Decoction, f 3j to ij.
Extract, grs. v. to xxx.
- Valeriana, Powder, 3ss to j.
Electuary, 3j to ij.
Infusion, f 3ij.
Tincture, f 3j to iv.
Ammon. f 3j to ij.
Wine, f 3j to iv.
Oil, gtt. ij to v.
Extract, grs. ij to x.
Fluid, f 3j to ij.
- Vanilla, Powder, grs. v to x.
Tincture, gtt. xxx to xl.
- Veratria, gr. $\frac{1}{12}$ to $\frac{1}{8}$.
Tincture, gtt. v to x.
- Veratrum Album, Powder, gr. j to ij.
Wine, m x.
Tincture, m x.
- Veratrum Viride, Powder, gr. j.
Tincture, f 3ss to j.
Norwood's, gtt. viij.
Wine, f 3ss to j.
Extract, gr. $\frac{1}{4}$ to $\frac{1}{2}$.
- Viola Odorata, Syrup, f 3j to ij.
- Wintera, Powder, 3ss to j.
- Zinci Oxidum, grs. ij to x.
Chloridum, Solution, gtt. v.
Tincture, gtt. v.
Cyanuretum, gr. $\frac{1}{12}$ to $\frac{1}{4}$.
Ferrocyanuretum, gr. j.
Sulphas, gr. j to xxx.
Solution, f 3ss.
Valerianas, gr. j to ij.
- Zingiber, Powder, gr. x to 3j.
Tincture, f 3j to ij.
Syrup, f 3ss.

TABLE OF PHARMACEUTICAL NAMES,

WHICH

DIFFER IN THE UNITED STATES, AND THE LONDON, EDINBURGH,
AND DUBLIN PHARMACOPŒIAS.

UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Absinthium,	idem,	idem,	
Acacia,	idem,	Gummi acaciæ,	Acacia vera, et A. verek.
Acetum,	idem,	Acetum gallicum,	Acetum gallicum.
Acetum destillatum,	idem,	idem,	
Acidum aceticum,	idem,	idem,	A. aceticum glaciale, et A. aceticum forte.
Acidum arseniosum,	idem,	Arsenicum album,	Arsenici oxydum album venale.
Acidum hydrocyanicum dilutum,	idem,	A. hydrocyanicum,	A. hydrocyanicum dilutum.
Acidum muriaticum,	A. hydrochloricum,	A. muriaticum, et m. purum,	A. muriaticum purum, et A. muriaticum venale.
Acidum muriaticum dilutum,	A. hydrochloricum dilutum,	A. muriaticum dilutum,	idem
Acidum nitricum,	idem,	A. nitricum, et A. nitricum purum,	A. nitricum venale.
Acidum sulphuricum,	idem,	A. sulphuricum, et A. sulphuricum purum,	A. sulphuricum venale, et A. sulphuricum purum.
Aconiti folia, et radix,	Aconiti radix; A. folium,	Aconitum,	Aconitum.
Adeps,	idem,	Axungia,	Adeps suillus,—Axungia.
Æther,	idem,	Æther sulphuricus,	Æther sulphuricus.
Alcohol,	Spiritus rectificatus,	idem,	idem.
Alcohol dilutum,	Spiritus tenuior,	idem,	idem.
Aloe,	A. Barbadosensis, A. Hepatica, A. Socotrina,	A. Barbadosensis, A. Indica, A. socotrina,	A. hepatica.
Althæa flores, et radix,	Althæa,	A. flores et radix,	
Alumen exsiccatum,	idem,	idem,	Alumen siccatum
Ammonia murias,	Ammonia hydrochloras,	Ammonia murias,	idem.

UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Ammoniaë carbonas,	Ammoniaë sesquicarbonas,	Ammoniaë carbonas,	Ammoniaë sesquicarbonas.
Amygdala dulcis,	Amygdala, var. dulcis,	Amygdala dulcis,	Amygdala, var. dulcis.
Angustura,	Cusparia,	idem,	
Antimonii et potassæ tartras,	Antimonii potassio-tartras,	Antimonium tartarizatum,	idem.
Antimonii sulphuretum præcipitatum,	Antimonii oxysulphuretum,	Antimonii sulphuretum aureum,	Antimonii sulphuretum præcipitatum.
Antimonii sulphuretum,	Antimonii tersulphuretum,	Antimonii sulphuretum,	idem.
Aqua camphoræ,	Mistura camphoræ,	idem,	idem.
	Aqua pulegii,	idem,	Aqua menthæ pulegii.
Argentum,	idem,	idem,	Argentum purificatum.
Argenti nitras, et argenti nitras fusus,	Argenti nitras,	idem,	Argenti nitras fusus.
Aurantii cortex,	idem,	idem,	Aurantium
Barii chloridum,		Barytæ murias,	Barii chloridum.
Bismuthi subnitras,	Bismuthi nitras,	Bismuthum album,	Bismuthi subnitras.
Buchu,	idem,	Bucku,	Buchu.
Calamus,		Calamus aromaticus,	
Calcii chloridum,		Calcis murias,	Calcii chloridum.
Calx,	idem,	idem,	Calx recens usta.
Carbo ligni,	Carbo,	Carbo ligni,	idem.
Carota,	idem,	Dauci radix,	Daucus carota.
Carum,	Carui,	idem,	idem.
Caryophyllus,	Caryophyllum,	Caryophyllus,	idem.
Cassia fistula,	Cassia,	Cassiae pulpa,	
Cassiae fistulæ pulpa,	Cassia præparata,	Cassiae pulpa,	
Cera flava,	Cera,	Cera flava,	idem.
Ceratum cantharidis,	Emplastrum cantharidis,	idem,	idem.
Ceratum cetacei,	idem,	Ceratum simplex,	Unguentum cetacei.
Ceratum plumbi subacetatis,	Ceratum plumbi compositum,		
Ceratum resinæ,	idem,	Unguentum resinosum,	Unguentum resinæ.
Ceratum sabinæ,	Unguentum sabinæ,	Ceratum sabinæ,	Unguentum sabinæ.
Ceratum saponis,	Ceratum saponis compositum,		
Ceratum simplex,	Ceratum,		
Cetraria,	idem,	idem,	Lichen Islandicus.
Chimaphila,	idem,	Pyrola,	idem.
Cinchona pallida,	idem,	Cinchona cinerea,	{ Cinchona Condaminea.
		Cinchona coronæ,	
			{ Cinchona micrantha.
Cinnamomum,	idem,	idem et Cassiæ cortex,	Cinnamomum.

UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Coccus,	idem,	Cocci,	Coccus cacti.
Colchici radix,	Colchici cormus,	idem,	Cochicum autum- nale. Cormus.
Colchici semen,	idem,	Colchici semina,	Colchicum autum- nale. Semina.
Colomba,	Calumba,	idem,	idem.
Confectio aromati- ca,	Confectio amygda- læ,	Conserva amygda- larum,	Confectio aromati- ca.
Confectio aurantii corticis,	idem,	Electuarium aroma- ticum,	
Confectio opii,	Confectio aurantii,	Conserva aurantii,	
	idem,	Electuarium opii,	
Confectio rosæ,	Confectio piperis,	Electuarium pipe- ris,	Confectio piperis ni- gri.
	Confectio rosæ,	Conserva rosæ,	Confectio rosæ.
	Confectio rosæ ca- ninæ,	Conserva rosæ fruc- tus,	
Confectio sennæ,	idem,	Electuarium sennæ,	Confectio sennæ.
Conii folia,	Conium,	idem,	idem.
Copaiba,	idem,	idem,	Copaiva balsam.
Coriandrum,	idem,	idem,	Coriander.
Cubeba,	idem,	Cubebæ,	Cubeba.
Cuprum ammonia- tum,	Cupri ammonio-sul- phas,	Cuprum ammonia- tum,	Cupri ammonio-sul- phas.
Cupri subacetas,	Ærugo,	idem,	Cupri subacetas.
	Cuminum,	Cuminum,	
	Decoctum aloës compositum,	Decoctum aloës,	Decoctum aloës compositum.
Decoctum cetrariæ,	idem,		Decoctum lichenis islandici.
Decoctum chima- philæ,	idem,		Decoctum pyrolæ.
Decoctum cinchonæ flavæ,	Decoctum cincho- næ,	idem,	
Decoctum cinchonæ rubræ,	idem,	Decoctum cincho- næ,	
	Decoctum hordei compositum,	Mistura hordei,	
Decoctum quercûs albæ,	Decoctum quercûs,	idem,	idem.
	Decoctum sarsæ,	Decoctum sarzæ,	Decoctum sarsapa- rillæ.
Decoctum sarsapa- rillæ compositum,	Decoctum sarsæ compositum,	Decoctum sarzæ compositum,	Decoctum sarsapa- rillæ compositum.
	Decoctum scoparii compositum,	Decoctum scoparii,	
Elaterium,	Extractum elaterii,	Elaterium,	idem.
Emplastrum ammo- niaci cum hydrar- gyro,	idem,	Emplastrum ammo- niaci et hydrar- gyri,	Emplastrum ammo- niaci cum hydrar- gyro.
Emplastrum galba- ni compositum,	Emplastrum galba- ni,		

UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Emplastrum picis cum cantha- ride,			Emplastrum calefa- ciens.
Emplastrum plum- bi,	idem,	Emplastrum lithar- gyri,	idem.
Emplastrum resi- næ,	idem,	Emplastrum resino- sum,	Emplastrum resi- næ.
	Extractum aloës,		Extractum aloës aquosum.
Extractum bella- donnæ,	idem,	idem,	Succus spissatus belladonnæ.
Extractum canna- bis,			Extractum cannabis Indicæ.
Extractum cincho- næ flavæ,	Extractum cincho- næ,	idem,	
Extractum cincho- næ rubræ,	idem,	Extractum cincho- næ,	
Extractum conii,	idem,	idem,	Succus spissatus co- nii.
Extractum hyoscy- ami,	idem,	idem,	Succus spissatus hy- oscyami.
	Extractum lupuli	idem,	Extractum hu- muli.
Extractum opii,	idem,	idem,	Extractum opii aquosum.
Extractum sarsapa- rillæ,	Extractum sarsæ,		Extractum sarsa- parillæ.
Extractum sarsapa- rillæ fluidum,	Extractum sarsæ liquidum,	Extractum sarsæ fluidum,	Extractum sarsapa- rillæ fluidum.
Extractum stramo- nii seminis,	Extractum stramo- nii.	idem,	
Ferri filum, et Ferri ramenta,	Ferrum in fila trac- tum,	Ferri filum, et Ferri limatura,	Ferri fila, et Ferri scobs.
Ferri et potassæ tartras,	Ferri potassio-tar- tras,	Ferrum tartariza- tum,	idem.
Ferri oxidum hy- dratum,		Ferrugo,	Ferri peroxydum hydratum.
	Ferri carbonas cum saccharo,	Ferri carbonas sac- charatum,	idem.
Ferri subcarbonas,	Ferri sesquiox- idum,	Ferri oxidum ru- brum,	Ferri carbonas.
Ferrum ammoni- um,	Ferri ammonio- chloridum,		
Ficus,	idem,	Fici,	Ficus carica.
Filix mas,		Filix,	
Galla,	idem,	Gallæ,	idem.
Gambogia,	Cambogia,	idem,	idem.
Glycyrrhiza,	idem,	Glycyrrhizæ radix,	Glycyrrhiza.
Granati fructûs cor- tex,	Granatum,		
Granati radicis cor- tex,	Granati radix,	idem,	Punica granatum.
Guaiaci lignum,	idem,	idem,	Guaiacum offi- nale.

UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Guaiaci resina,	Guaiacum,	idem,	idem.
Hæmatoxylon,	Hæmatoxylum,	Hæmatoxylon,	Hæmatoxylum.
Humulus,	Lupulus,	idem,	Humulus lupulus.
Hydrargyri chloridum corrosivum,	Hydrargyri bichloridum,	Sublimatus corrosivus,	Sublimatum corrosivum
Hydrargyri chloridum mite,	Hydrargyri chloridum,	Calomelas,	idem.
Hydrargyri iodidum,	idem,		Hydrargyri iodidum viride.
Hydrargyri iodidum rubrum,	idem,	Hydrargyri biniodidum,	Hydrargyri iodidum rubrum.
Hydrargyri oxidum rubrum,	Hydrargyri nitricoxidum,	Hydrargyri oxidum rubrum,	Hydrargyri oxidum rubrum.
Hydrargyri sulphas flavus,			Hydrargyri oxidum sulphuricum.
Hydrargyri sulphuretum nigrum,	Hydrargyri sulphuretum cum sulphure,		Hydrargyri sulphuretum nigrum.
Hydrargyri sulphuretum rubrum,	Hydrargyri bisulphuretum,	Cinnabaris,	
Hydrargyrum ammoniatum,	Hydrargyri ammonio-chloridum,	Hydrargyri precipitatum album,	Hydrargyri ammonio-chloridum.
Hyoseyami folia,	Hyoseyamus,	idem,	idem.
Infusum angusturæ,	Infusum cuspariæ,	idem,	Infusum angusturæ.
Infusum armoraciæ,	Infusum armoraciæ compositum,		
	Infusum aurantii compositum,	Infusum aurantii,	Infusum aurantii compositum.
Infusum buchu,	idem,	Infusum bucku,	Infusum buchu.
Infusum catechu compositum,	idem,	Infusum catechu,	Infusum catechu compositum.
Infusum cinchonæ flavæ,	Infusum cinchonæ,	idem,	
Infusum colombæ,	Infusum calumbæ,	idem,	idem.
Infusum gentianæ compositum,	idem,	Infusum gentianæ,	Infusum gentianæ compositum.
Infusum humuli,	Infusum lupuli,		
Infusum lini compositum,	idem,	Infusum lini,	
Infusum rosæ compositum,	idem,	Infusum rosæ,	Infusum rosæ acidum.
Infusum sennæ,	Infusum sennæ compositum,	Infusum sennæ,	Infusum sennæ compositum.
Infusum tabaci,	Enema tabaci,	idem,	idem.
Inula,	idem,		Inula helenium.
Iodinium,	idem,	Iodineum,	Iodinium.
Juniperus,	idem,	Juniperi cacumina,	Juniperus communis. Cacumina.

UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
		Juniperi fructus,	Juniperus communis. Baccæ.
	Lactuca,		Lactuca sativa.
Limon,	Limones,	idem,	idem.
Limonis cortex,	Limorum cortex,	idem,	idem.
Linimentum terebinthinæ,	idem,	Linimentum terebinthinatum,	Linimentum terebinthinæ.
Linum,	Lini semen,	Lini farina et semina,	Linum.
Liquor ammoniæ,	Ammoniæ liquor,	Ammoniæ aqua,	Ammoniæ liquor.
Liquor ammoniæ acetatis,	idem,	Ammoniæ acetatis aqua,	Ammoniæ acetatis liquor.
Liquor ammoniæ fortior,	Ammoniæ liquor fortior,	Ammoniæ aqua fortior,	Ammoniæ liquor fortior.
	Liquor ammoniæ sesquicarbonatis,	Ammoniæ carbonatis aqua,	
Liquor arsenici et hydrargyri iodidi,			Arsenici et hydrargyri hydriodatis liquor.
Liquor barii chloridi,		Solutio barytæ muriatis,	Barii chloridi liquor.
Liquor calcii chloridi,		Calcis muriatis solutio,	Calcii chloridi liquor.
Liquor calcis,	idem,	Aqua calcis,	Liquor calcis.
	Liquor cupri ammonio-sulphatis,	Cupri ammoniati solutio,	Cupri ammoniati aqua.
Liquor ferri iodidi,	Syrupus ferri iodidi,	Ferri iodidi syrupus,	Syrupus ferri iodidi.
Liquor ferri nitratis,			Ferri pernitratis liquor.
Liquor iodinii compositus,		Iodinei liquor compositus,	
	Liquor morphiæ hydrochloratis,	Morphiæ muriatis solutio,	Morphiæ muriatis liquor.
Liquor plumbi subacetatis,	Liquor plumbi diacetatis,	Plumbi diacetatis solutio,	Plumbi subacetatis liquor.
Liquor plumbi subacetatis dilutus,	Liquor plumbi diacetatis dilutus,		Plumbi subacetatis liquor compositus.
Liquor potassæ,	idem,	Potassæ aqua,	Potassæ causticæ liquor.
Liquor potassæ arsenitis,	idem,	Liquor arsenicalis,	idem.
Liquor potassæ carbonatis,	idem,		Potassæ carbonatis liquor.
	Liquor sodæ,		Sodæ causticæ liquor.
Liquor sodæ chlorinatae,	idem,		Sodæ chlorinatae liquor.
	Manganesii binoxidum,	Manganesii oxidum,	Manganesii per-oxidum.
Maranta,	idem,	idem,	Maranta arundinaceæ.
Marmor,		idem,	Marmor album.

UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Mezereum,	idem,	Mezereon,	idem.
Mistura amygdalæ,	idem,	Mistura acaciæ, Mistura amygdala- rum,	Emulsio Arabica. Mistura amygdalæ
Morphiæ murias,	Morphiæ hydro- chloras,	Morphiæ murias,	idem.
Mucilago acaciæ,	Mistura acaciæ, Decoctum amyli,	Mucilago, Mucilago amyli,	Mucilago acaciæ. idem.
Mucuna,	idem,	idem,	Dolichos.
Oleum amygdalæ, Oleum bergamii, Oleum cajuputi, Oleum cari, Oleum cinnamomi,	Amygdalæ oleum, Cajuputi, Oleum carui, Cinnamomi oleum,	Bergamotæ oleum, Cajuputi oleum, idem, idem et Cassiæ oleum,	Cajeputum. idem. Oleum cinnamomi.
Oleum limonis, Oleum lini,	Limonum oleum, Lini oleum,	idem, idem,	Citrus limonum. Linum usitatissi- mum.
Oleum morrhuæ, Oleum myristicæ, Oleum olivæ, Oleum ricini, Oleum rosæ, Oleum terebinthi- næ, Oleum tigllii,	Morrhua oleum, Olivæ oleum, Ricini oleum, Terebinthinæ oleum, Tigllii oleum, Oleum pulegii,	Myristicæ oleum, idem, idem, Rosæ oleum, idem, Crotonis oleum, idem,	Oleum morrhua. Oleum myristicæ. Oleum olivæ. Oleum ricini. Oleum rosæ. Oleum terebinthi- næ. Croton oil. Oleum menthæ pu- legii.
Ovum,	Ovi albumen et vi- tellus,	Ovum,	idem.
Pilulæ aloës,	Pilulæ aloës cum sapone,	Pilulæ aloës,	
Pilulæ aloës et myrrhæ,	Pilula aloës cum myrrhâ, Pilula hydrargyri chloridi composi- ta, Pilula colocynthi- dis composita,	Pilulæ aloës et myrrhæ, Pilulæ calomelanos compositæ, Pilulæ colocynthi- dis,	Pilulæ aloës cum myrrhâ. idem.
Pilulæ galbani com- positæ,	Pilula galbani com- posita, Pilula cambogiæ composita,	Pilulæ assafœtidæ, Pilulæ cambogiæ,	Pilulæ colocyn- thidis compositæ. Pilulæ assafœtidæ compositæ. Pilulæ gambogiæ compositæ.
Pilulæ opii,		Pilulæ opii, sive thebaicæ,	
Pilulæ saponis com- positæ, Pilulæ scillæ com- positæ,	Pilula saponis com- posita, Pilula scillæ com- posita, Pilula styracis com- posita,	Pilulæ scillæ, Pilulæ styracis,	Pilulæ saponis com- positæ. Pilulæ scillæ com- positæ.
Piper,	Piper nigrum, Pix,	idem, Pix arida,	idem.
Plumbi carbonas,	idem,	idem,	idem et Ceresa

UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Plumbi oxidum semivitreum,	Plumbi oxidum,	Lithargyrum,	idem.
Potassæ carbonas impurus,			Lixivus cinis.
Potassæ carbonas,	idem,	idem,	Potassæ carbonas à lixivo cinere.
Potassæ carbonas purus,		Potassæ carbonas purum,	idem.
Potassii ferrocyanuretum,	Potassii ferrocyanidum,	idem,	idem.
Potassa,	Potassæ hydras,	Potassa,	Potassa caustica.
Potassa cum alce,	idem,	idem,	Potassa caustica cum calce.
Potassii sulphuretum,	idem,	idem,	Hepar sulphuris.
Prunum,	idem,	Pruna,	idem.
Pruni pulpa,	Prunum præparatum,		
Pulvis aromaticus,	Pulvis cinnamomi compositus,	Pulvis aromaticus,	idem.
	ulvis cretæ compositus cum opio,	Pulvis cretæ opiatum,	idem.
Pulvis ipecacuanhæ et opii,	Pulvis ipecacuanhæ compositus,	idem,	idem.
		Pulveres effervescentes,	Pulveres effervescentes tartarizati.
Quercus alba et Q. tinctoria,	Quercus,	Quercus cortex,	Quercus.
Quiniæ sulphas,	Quinæ disulphas,	Quinæ sulphas,	idem.
	Rhamni succus,	Rhamni baccæ,	
	Rhœas,	idem,	Papaver rhœas.
	Rosæ canina,	Rosæ fructus,	
Saccharum,	idem,	Saccharum purum,	Saccharum purificatum.
		Saccharum commune,	Brown sugar.
	Sacchari fæx,	idem,	Theriaca.
Salix,		Salicis cortex,	
Santalum,	Pterocarpus,	idem,	
Sapo,	idem,	Sapo durus,	idem.
	Sapo mollis,	idem,	idem.
Sarsaparilla,	Sarsa,	Sarza,	Sarsaparilla.
Sassafras radicis cortex,	Sassafras,	idem,	Sassafras root.
Scoparius,	idem,	Scoparium,	idem.
Senna,	Senna Alexandrina et Senna Indica,	idem,	Senna.
Sinapis,	idem,	Sinapi,	Sinapis alba, S. nigra.
			idem.
Sodæ boras,	Borax,	idem,	Sodæ carbonas crystallizatum.
Sodæ carbonas,	idem,	idem,	idem.
Sodæ carbonas exsiccatus,	Sodæ carbonas exsiccata,	Sodæ carbonas siccatum,	

UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Sodii chloridum,	idem,	Sodæ murias,	Sodii chloridum.
Sodæ et potassæ tartras,	Sodæ potassio-tar- tras,	Potassæ et sodæ tartras,	Sodæ et potassæ tartras.
Spiritus ætheris compositus,	idem,		Spiritus æthereus oleosus.
Spiritus ætheris ni- trici,	idem,	idem,	Spiritus æthereus nitrosus.
Spiritus lavandulæ compositus,	Tinctura lavandulæ composita, Spiritus menthæ piperitæ,	Spiritus lavandulæ compositus, Spiritus menthæ,	idem.
Stramonii folia, Stramonii semen,	Stramonii folium, idem,	Stramonium,	Stramonium. The seeds.
Styrax purificata, Sulphur,	Styrax præparata, idem,	Extractum styracis, idem,	Sulphur sublima- tum.
Sulphur lotum,		Sulphur sublima- tum,	
Sulphuris iodidum,	idem,		Sulphur iodatum.
Syrupus,	idem,	Syrupus simplex,	idem.
Syrupus aurantii corticis,	Syrupus aurantii,	idem,	idem.
Syrupus limonis,	Syrupus limonum, Syrupus rosæ,	idem, Syrupus rosæ centi- foliæ,	
	Syrupus sarsæ, idem,	Syrupus sarzæ, idem,	idem.
Syrupus tolutanus,			
Tamarindi pulpa,	Tamarindus præ- paratus,	Tamarindus,	
Tamarindus,	idem,	idem,	Pulp of tamarinds.
Taraxacum,	idem,	idem,	Taraxacum dens- leonis.
Terebinthina Cana- densis,		Balsamum Cana- dense,	
Tinctura aconiti radicis,	Tinctura aconiti,		Tinctura radicis aconiti.
Tinctura aloës et myrrhæ,	Tinctura aloës com- posita,	Tinctura aloës et myrrhæ,	
Tinctura belladon- næ,	idem,		Tinctura foliorum belladonnæ.
Tinctura campho- ræ,	Spiritus camphoræ,	Tinctura bucku, Tinctura campho- ræ,	Tinctura buch. idem.
Tinctura catechu,	Tinctura catechu composita,	Tinctura catechu,	idem.
Tinctura colchici seminis,	Tinctura colchici,	idem,	Tinctura seminum colchici.
Tinctura colombæ,	Tinctura calumbæ,	idem,	idem.
Tinctura ferri chlo- ridi,	Tinctura ferri ses- quichloridi,	Ferri muriatis tinc- tura,	Tinctura ferri ses quichloridi.
Tinctura gallæ,	idem,	Tinctura gallarum,	Tinctura gallæ.
Tinctura guaiaci ammoniata,	Tinctura guaiaci composita,	Tinctura guaiaci ammoniata,	
Tinctura humuli,	Tinctura lupuli,		

UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Tinctura iodinii,		Tinctura iodinei,	
Tinctura iodinii composita,	idem,		idem.
Tinctura lupuli- næ,		Tinctura lupuli,	
Tinctura olei men- thæ piperitæ,			Essentia menthæ piperitæ.
Tinctura olei men- thæ viridis,			Essentia menthæ viridis.
Tinctura opii cam- phorata,	Tinctura camphoræ composita,	Tinctura opii cam- phorata,	idem.
Tinctura saponis camphorata,	Linimentum sapo- nis,	idem,	idem.
Tinctura sennæ et jalapæ,		Tinctura sennæ composita,	
Tinctura valerianæ ammoniata,	Tinctura valerianæ composita,	Tinctura valerianæ ammoniata,	
Trochisci glycyrrhi- zæ et opii,		Trochisci opii,	
	Tussilago,		Tussilago farfara.
Unguentum anti- monii,	Unguentum anti- monii potassio- tartratis,	Unguentum anti- moniale,	Unguentum anti- monii tartarizati.
Unguentum can- tharidis,	idem,	Unguentum infusi cantharidis,	Unguentum can- tharidis.
	Ceratum canthari- dis,	Unguentum can- tharidis,	
Unguentum cupri subacetatis,		Unguentum ærugi- nis,	Unguentum cupri subacetatis.
Unguentum gallæ,			Unguentum galla- rum.
	Unguentum gallæ compositum.	Unguentum gallæ et opii,	
Unguentum hy- drargyri ammo- niati,	Unguentum hy- drargyri ammo- nio-chloridi,	Unguentum præci- pitati albi,	
Unguentum hy- drargyri nitratis,	idem.	Unguentum citri- num,	Unguentum hy- drargyri nitratis vel Unguentum citrinum.
Unguentum hy- drargyri oxidi rubri,	Unguentum hy- drargyri nitrico- oxidi,	Unguentum oxidi hydrargyri,	Unguentum hy- drargyri oxydi rubri.
Unguentum iodinii compositum,	idem,	Unguentum iodinei,	Unguentum iodinei compositum.
	Ceratum plumbi acetatis,	Unguentum plumbi acetatis,	idem,
Unguentum sim- plex,		idem,	Unguentum ceræ albæ.
Unguentum zinci oxidi,	Unguentum zinci,	idem,	Unguentum zinci oxydi.
Uva passa,	Uva,	Uvæ passæ,	idem.
Veratrum album,	Veratrum,	idem.	

UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Vinum album,	Vinum xericum,	Vinum album,	Vinum album Hispanicum.
Vinum antimonii,	Vinum antimonii potassio-tartratis,	Vinum antimoniale,	Antimonii tartarizati liquor
Vinum colchici radicis,	Vinum colchici,	idem,	
Vinum veratri albi,	Vinum veratri,		
Zinci carbonas præcipitatus,			Zinci carbonas.
Zinci oxidum,	idem,	idem,	Zinci oxydum.

OFFICINAL PREPARATIONS AND DIRECTIONS.

INTERNAL REMEDIES.

POWDERS — These are of two kinds: *simple* and *compound*. The first are prepared by pulverization; and the second by the mixture of two or more simple powders, except where one of the ingredients is employed to facilitate the more minute division of the others, as in the case of the powder of ipecacuanha and opium. Many of the most important articles used in this form are powdered by grinding and stamping, by persons who make it a special business. When this operation is performed by the apothecary, or medical practitioner, it is most frequently accomplished by means of the pestle and mortar, and the sieve: but in some cases, a stone slab and muller are required; whilst in others, the article is merely rubbed through a sieve. Whenever a substance cannot be dried completely, without an alteration of its properties, recourse must be had to an intermedium, by which the moisture may be absorbed, or its state of aggregation modified. Thus, sugar is the best intermedium in pulverizing vanilla or nutmeg. When camphor is to be pulverized, the addition of a small quantity of alcohol will much facilitate the operation. In other cases, the intermedium should be of so hard a consistence as to assist in breaking down the substance to be pulverized; thus, gold-leaf is best reduced to powder by rubbing it with sulphate of potassa, and afterwards removing this latter by means of water.

The rules to be observed in the preparation of powders are:—

1. Operate, if possible, on perfectly dry articles, and in dry weather.
2. Adapt the nature of the mortar, and the mode of operating, to the nature of the substance: thus, woods and barks should be pulverized in an iron mortar; sugar, alum, and nitre, in one of marble; corrosive sublimate in one of glass.
3. The mortar should be provided with a cover to prevent particles from being forced out by the action of the pestle, and also to arrest the escape of the finer particles, which would otherwise be diffused through the atmosphere, causing a loss of the product, and an annoyance to the operator.
4. Separate, from time to time, by aid of a sieve, the pulverized portions, returning the coarser particles to the mortar; and repeat this alternate pulverization and sifting until the process is completed.

Compound Powders.—1. Each substance is to be pulverized separately, and the quantity ordered in the prescription then weighed off; otherwise, the due proportions will not be maintained.

2. Pulverize soft substances with those which are the reverse, which will thus serve as an intermedium.

3. Pass all the substances through the same sieve, as this will render them more homogeneous, and will also prevent unnecessary loss.

Some substances, however well dried, are reduced to powder with great difficulty by the usual modes of pulverization; as, for instance, nux vomica and jalap, which require to be exposed to the steam of boiling water for some time, and then rapidly dried.

Few articles, except those containing volatile constituents, are deteriorated in their properties by being pulverized. Some, on the contrary, become more active by this process, from their less active portions being separated; for instance, the powder

of ipecacuanha, if properly prepared, and the ligneous part rejected, is far stronger than the root—all the efficient properties existing in the cortical portion.

Some other processes are employed to obtain certain powders: viz. Precipitation, elutriation, and granulation.

Precipitation.—Tartar emetic is obtained in a perfectly fine powder, by precipitation from a concentrated aqueous solution by means of alcohol; so, also, the precipitated carbonate of lime is prepared by the action of a solution of carbonate of soda on the solution of chloride of calcium.

Elutriation.—This is a method by which the finer particles of a powder are separated from the coarser. It is performed by diffusing the powder through water, permitting the heavier portions to subside, then withdrawing the fluid, holding the finer particles in suspension, into another vessel, and allowing them to subside. This process is employed in making prepared chalk, &c.

Granulation.—This is used to obtain certain metals in a finely divided state; as zinc, tin, iron, &c.

Zinc is granulated either coarsely or finely: the first form is procured by melting the metal in an iron ladle, and pouring it slowly, in a fine stream, into cold water; the fine powder is obtained by rubbing the melted metal in an iron mortar. This metal loses its ductile properties at about 400° , and becomes so brittle as to be pulverized without difficulty. It does not melt, except at a temperature of 773° , so that the melting is many degrees above its brittle point; hence, by beginning the trituration at the melting point, it gradually reaches that at which it is most readily pulverized. The mortar should be thick and well heated before the fluid metal is poured into it. After being granulated, it should be properly sifted.

The granulation of tin may be effected in the same manner; but a more common mode is to pour the melted metal into a strong wooden box, provided with a closely-fitting lid, and agitating till the tin is cold; by this means, it is rapidly converted into powder and small grains, which can be separated from each other by elutriation or sifting.

Iron is granulated by filing, or by means of a pestle and mortar (see page 227); but the best means to procure metallic iron in a finely divided state is by reducing it from the state of the carbonate by means of hydrogen gas (see page 227).

PILLS AND BOLUSES.—Pills are small globular masses, of a semi-solid consistence, of the weight of half a grain to six grains, having as a base a powdery extract, &c., and rendered of the proper consistence by a syrup, pulp, &c. A bolus differs from a pill merely in being much larger, and generally less solid, weighing from six to twelve grains, and even more.


No form of medicinal agents is more frequently employed than that of the pill; not only because of the facility with which it is administered, and its comparatively little taste, but because this form answers so excellent a purpose in the preservation of certain compounds.

The article, or articles, to be administered in this mode, should first be formed into a consistent, moderately-firm mass—sufficiently plastic to admit of its being moulded into shape, without adhesion to the moulding instrument, and yet of so firm a consistence as to retain the form given to it. To accomplish this, the following rules must be attended to:—

1. All the constituents, which can be pulverized, must be reduced to the state of fine powder, and then thoroughly mixed together.
2. If soft ingredients enter into the composition, they must be triturated with the harder articles, which thus serve as an intermedium.
3. No deliquescent salt should enter into the composition of officinal pills; otherwise they will become altered and decomposed.
4. The nature of the excipient should be suited to that of the constituents: thus, syrups are to be used for most vegetable powders; soap for fatty matters; calcined magnesia for copaiba and turpentine. In some cases, no excipient is required, as for most of the gum resins. Extracts, also, when of proper consistence, can be made

into pills without any addition. Mucilage, which is often ordered in the formation of various pills, is only suited to those which are to be used in a short time after

FIG. 11.



COUNTER DRAWER CONTAINING A PILL-MACHINE.

A. Pill-machine. B. Pill-cutter. C. Roller. D. D. Sides of the drawer.

they are made; but is not suited to such as are to be kept, because pills, into which it enters as an excipient, become so hard as to pass through the bowels almost unaltered.

FIG. 12.




Fig. 12. Water-bath, in which is placed the vessel, *a*, holding a solution of one part of gelatine in two parts of water. *b*. Waste pipe to carry off the steam. *c*. Gas burner.

5. The excipient should be gradually added, and the mass beaten and triturated till it is perfectly homogeneous.

6. When the pilular mass is properly made, the next operation consists in dividing it into pills of equal size, which is effected by dividing it with a spatula into morsels of the requisite weight, and rolling them between the fingers; or, in a more effectual, rapid, and cleanly manner, by means of a pill-machine. (See fig. 11.)

7. To prevent any adhesion of the pills after they are made, and to prevent any disagreeable taste, they are, in most cases, covered with an inert powder: as, powdered liquorice-root, starch, orris-root, lycopodium, and frequently with magnesia; but this latter is wholly unsuited to some preparations—for instance, to pills of calomel—as decomposition will be caused. In Europe, they are often coated with gold or silver leaf; but this plan is seldom adopted in this country. Where, from their nauseous taste, or other circumstances, it becomes expedient to cover them,

this is best done by means of gelatine. Each pill, being stuck on the point of a thin wire, four or five inches in length, is dipped into a solution of gelatine, (a, fig. 12,) so as to coat it completely; and the wire is then inserted into a pin-cushion, or a vessel containing fine sand, (fig. 13,) and left till the gelatine is firm, which occurs in about a quarter of an hour, when the pill can be transferred to a pan, (fig. 14,) to dry.

FIG. 13.




FIG. 14.



COATING PILLS WITH GELATINE.

Fig. 13. Vessel partly filled with sand, into which the pins holding the coated pills are stuck.

Fig. 14. Pan for holding the partially-dried pills.

from the evaporation of a solution, maceration, digestion, or infusion, or of an expressed juice. They are generally of a soft consistence, of a dark color, and of an odor and taste analogous to those of the substance from which they are derived. They have been classed, according to the fluid employed in obtaining them; as watery, alcoholic, hydro-alcoholic, ethereal, vinous, acetic, &c. Some, however, are made from the expressed juices of plants, without any intermedium. The great object, in selecting one or more of these menstrua, is to employ that which takes up the largest proportion of the active ingredients of the root, bark, &c.; for the most advantageous solvent for one constituent may be wholly inefficient for others. Thus, alcohol is the proper menstruum for resinous substances, water for bitter extractive, vinegar or diluted acids, for articles containing the alkaloids, &c.; but, in many cases, an extract, containing all the active properties of the drug, can be obtained only by a combination or a succession of different solvents.


Extracts made by inspissating the expressed juices of plants have been considered to contain the active principles of the vegetable in a less altered form than by any other mode. But it has been shown that such is far from being the case, as, in some instances, the expressed juice does not contain the active constituents. Thus, in aconitum, much of the efficient principle is left in the residuum; added to which, in some extracts made in this manner, the medicinal ingredients are destroyed, or much deteriorated, in the evaporation. This is obviated, in a great measure, by allowing the expressed juice to evaporate spontaneously at ordinary temperatures; or, by carrying on this process in vacuo. It has been found that extracts made by the latter method have more of the aroma and taste of the original article, keep better, and are more efficient. When inspissation of the natural juices of the plant is not practicable, a solution of the active ingredients is to be obtained by the use of certain menstrua—as water and alcohol—according to the substance to be acted upon. When the active principles are soluble in water, that fluid is to be used as the vehicle of extraction; when resinous, alcohol is to be employed; when oleo-resinous, ether will be found the best menstruum. When it is desired that all the principles soluble in the various menstrua should be reunited in the extract, the vegetable substance is first to be subjected to the action of one, and then to another, of the solvents; and the solutions thus obtained are to be mixed together, and properly inspissated. These solutions are obtained by maceration, digestion, infusion, and displacement. Decoction is seldom to be employed, as it affords extracts containing much inert matter, and, of course, of inferior quality. Of late years, the process of displacement has been much used in the formation of extracts, whatever the menstruum selected, as it affords a solution of the soluble principles in a much smaller quantity of fluid; and hence, prevents the injurious effects of too long an exposure to heat in the subsequent evaporation. (See Tinctures.)

With regard to the mode of evaporating extracts, much caution is requisite, as the various constituents of these bodies are very liable to undergo decomposition

EXTRACTS.—This name is given to all products resulting

when exposed to the action of heat and atmospheric air. Sometimes the evaporation, especially of aqueous solutions, is performed over the naked fire; but this plan

FIG. 15.



WATER BATH.

This is made of tinned-iron, or copper, in such a manner, as to leave a space between the two rings forming its wall, into which water can be introduced through the orifice *a*. The temperature of the water can be raised by a stove heat, or by the spirit lamp (represented beneath) above 212° F., by causing the mouth of the tube *b* to dip into a cup of mercury.

The great objection is the additional expense it entails on the operation. Spontaneous evaporation has also been successfully employed; though necessarily slow, it affords excellent products. It is always advantageous to aid this process by employing artificial heat to a certain degree, by conducting it in a drying room or warm closet, and by causing a current of dry air, heated to about 100°, to pass over the surface of the evaporating fluid. Extracts made in this manner keep well, not undergoing the spontaneous decomposition so detrimental to extracts in general. When alcoholic solutions are to be concentrated, distillation should always be used, as not only can the alcohol be recovered, but, from the process being carried on out of the atmospheric air, one great source of injury to the product is avoided.

The consistence of extracts varies according to their ingredients, and the purposes to which they are to be applied. In addition to the fluid extracts, properly so-called, two different degrees of consistence are recognized in the Pharmacopœias: one so soft as to be readily made into pills; the other so hard as to be pulverizable. These different forms of extracts are useful because of the readiness with which they can be formed into pills or powders, when prescribed. But, in many cases, extracts undergo a change, some becoming soft and others hard; and one of the difficult parts of an apothecary's business is to keep these preparations of an uniform consistence. They should be preserved in glazed earthenware pots, protected as much as possible from the action of the air. Notwithstanding every precaution that can be taken, a diminution of activity will, sooner or later, take place; hence, these preparations should be examined from time to time, and, if necessary, be renewed. The safest plan is to renew those made from recent plants once a year.

CONFECTIONS, CONSERVES, ELECTUARIES. — As there is no essential difference in these preparations, they may be advantageously treated under one head. They are simple and compound: the first being merely a mixture of a recent vegetable substance and sugar, beaten into an uniform mass; the second, of the same substances, but in greater numbers. In consequence of the greater simplicity of prescriptions, of late years, the number of preparations of this kind have greatly diminished, and they are employed more as vehicles for other medicines than for their own remedial powers. The following rules are to be observed in their preparation and preservation:—

is liable to many objections, more particularly towards the close of the operation, when a small excess of heat may cause such a change in the character of the ingredients as to deprive them of all useful properties; and it is from this cause that so many of the extracts found in the shops are wholly worthless. Other and far better methods consist in the use of water and steam-baths (see Fig. 15), which prevent all danger of burning the extract. In the inspissation, when conducted in open vessels, it is of importance that the evaporation should be carried on rapidly, that the solution may be exposed to the action of the air for as short a time as possible. Evaporation in vacuo is always to be preferred when practicable, as the process can not only be carried on at a lower temperature, but the deteriorating influence of the air is avoided.

1. The powders to be incorporated are to be in a state of fine division, and the pulps used, perfectly homogeneous, and not too fluid.
2. The mixture should be intimate, and the consistence of the resulting mass should be solid enough to prevent a separation of the ingredients; and yet sufficiently soft to allow of its being swallowed without being masticated.
3. If the mixture swells up, and gives off carbonic acid, it is to be well rubbed in a mortar, to break down any sugar that may have crystallized, and any hard lumps that may have formed, so as again to form an uniform mass.
4. They should be preserved in earthenware or porcelain pots, well covered, and kept in a moderately moist situation.

PULPS are simple medicinal preparations, of a soft consistence, formed of the tissues of vegetable substances, or some of their thin parts; they differ from extracts in containing insoluble as well as soluble principles. They are to be prepared by rubbing the fruit, or other articles from which they are made, through a wire sieve, so as to separate the vegetable fibre from the soft parts by which it is surrounded. When the fruit is hard or dry, it is to be softened by means of boiling water, previous to expression through the sieve. The expressed pulp, if not sufficiently consistent, is to be properly evaporated. Pulps are to be preserved in the same manner as confectious. Very few of these preparations are now used in medicine.

SYRUPS.—These are liquid, viscous medicines, consisting of a concentrated solution of sugar in aqueous fluids.

All fluids susceptible of dissolving more than their weight of sugar can be formed into syrups. These syrups are of two kinds: simple and compound. Simple syrup is prepared by dissolving sugar in pure water; and compound syrups are obtained by dissolving the sugar in solutions of various substances, formed by infusion, decoction, expression, &c. The former is usually made in this country from refined sugar, and not, as in Europe, from the impure and unrefined qualities of this article. All syrups require to be perfectly filtered, so as to be limpid, and they should have a certain viscosity of consistence, and be capable of being preserved without entering into fermentation, or crystallization. These latter properties depend on their not containing the proper proportion of sugar—an excess being deposited in a crystalline form, and a deficiency causing the solution to run into fermentation. The best mode of ascertaining the proper point of concentration is by means of the specific gravity at different temperatures. The specific gravity of well-prepared simple syrup is, when boiling, about 1.261, and when cold 1.319; but the proper degree of concentration is more readily obtained by means of Baumé's hydrometer, (see pages 31, 34). This should stand at about 30° in boiling syrup, and at 35° when it is cold. Other modes are also employed, which, although sufficiently accurate in the hands of an experienced operator, are not to be generally depended upon. They are derived from the degree of viscosity acquired by the syrup, as shown by the time required for the parts of a drop to re-unite, and by the length of the thread which a drop will produce before detaching itself, when poured from a spoon or ladle. When the syrup, on cooling, presents a crystalline pellicle, it is a proof that the evaporation has been carried too far; but, when the sugar has been mixed with an acid, or when the process has been too much prolonged, the sugar loses its power of crystallization, however much the syrup is concentrated, and, therefore, does not form a pellicle.

The compound syrups, if kept any time, are liable to various alterations, depending on their nature, and the degree of care used in their preparation. Thus, the acid syrups, as the syrup of lemons, when too concentrated, deposit a copious white precipitate; and, in some cases, solidify entirely. By heating them, they again become liquid; but again let fall a precipitate on cooling. This deposit is analogous to grape sugar, and is caused by the action of the acid on the sugar. When the sugar bears too small a proportion to the liquid, syrups are apt to run into fermentation. Even when the sugar is in proper proportion, this change often takes place, if the solution contains much amylaceous or extractive vegetable matter. Even when too much

concentrated, they may also undergo this change, from part of the sugar being deposited in a crystalline state; and the crystal, attracting the sugar necessary to the preservation of the syrup, reduces its strength, and renders it liable to the same change as though it was originally too weak.

Syrups, especially those containing the juices of fruits, should be bottled whilst hot, and, when cold, well stopped and sealed; and these, as well as all other kinds, should be kept in a temperature not exceeding 60° F. Various plans have been devised to preserve syrups; but the best is to prepare them only in such quantities as will be used within a short time. The addition of chlorate of potassa, as advised by Macculloch, and of sugar of milk, as advised by Chereau, has proved useful; but the best mode is that of Mr. Durand, viz., adding about one drachm of Hoffmann's anodyne to each pint of syrup; this appears to have the property of arresting or preventing any tendency to fermentation.

MELLITES, OR HONEYS, are liquid, viscous medicines, somewhat analogous to syrups, but in which the sugar is replaced by honey. Like syrups, they may be divided into simple and compound, or medicated. Their preparation, medical properties, modes of administration and preservation, are very similar to those of the syrups, and do not, therefore, require further notice. Though formerly much used, they are now seldom prescribed, as they are often found to disagree with the stomach, especially if made with the honey of commerce in an unpurified state.

INFUSIONS are aqueous solutions, made by treating vegetable products with cold or hot water, but never carried to ebullition. They are seldom made by the apothecary, who merely furnishes the medicinal ingredients; whilst the preparation is usually confided to the nurse or other attendant on the patient. This plan, although more economical than when the preparation is compounded by the apothecary, often renders the prescription of the physician of little avail, from the infusion being erroneously made. They are generally prepared by pouring boiling water on the ingredients, and macerating in a tightly-covered vessel until the liquid cools; or the vessel is kept for some time at a low heat before a fire. In most cases, an infusion should be strained or filtered before being used; but, in general, nurses are content merely to decant them for use. The vessels, in which infusions are made, are usually of stone or earthenware. When it is an object to retain the heat as long as possible, polished metallic vessels, provided they are not acted upon by the infusion, are to be preferred, as, from their inferior radiating powers, they retain heat better than earthenware. Infusions are mostly made for extemporaneous use, as they are exceedingly liable to decomposition, and consequently cannot be kept long without spoiling. They may, however, be preserved for some time by pouring them, whilst hot, into bottles, so as to fill them, and forcing in stout corks. If the operation be properly performed, so as to perfectly exclude the air, the infusion will keep for some months. An infusion made with cold water, by the process of displacement, has less tendency to spoil than one made in the usual manner with hot water.

DECOCTIONS differ from infusions in the circumstance that the substance to be acted upon is subjected to the solvent action of the menstruum at its boiling point, and continued for a longer or shorter period, according to the solubility of the substance, or its constituents. By decoction, all the principles soluble in water can be obtained; and even many substances not properly soluble in that fluid are diffused through it, and held in suspension. On the other hand, many substances are injured or destroyed by decoction, especially when their active principles are volatile, and when, during ebullition, chemical changes take place, by which the active constituents are rendered insoluble, or are decomposed. In fact, protracted ebullition is seldom required since the process of displacement has become known, as it has been found that water at a common temperature, by this method, will exhaust most vegetable substances more effectually than the same fluid at 212° by decoction. There are some exceptions; as all emulsive preparations, and certain gummy solutions, require

ebullition: As in the case of infusions, decoctions are only made extemporaneously, for they readily decompose by keeping. They should be prepared in closed vessels, and generally in those of earthenware. Copper vessels should be used with great care, as many vegetable decoctions corrode them, especially if permitted to cool in them. Iron vessels are not to be employed when the decoction contains tannin or gallic acid. Zinc is very rapidly acted upon by many vegetable substances, and should, therefore, be avoided. In all cases, the vessels employed should be deep and narrow, rather than broad and shallow, to obviate as much as possible the influence of the air on the product.

TINCTURES are solutions of vegetable, animal, and, in some cases, of mineral substances in spirituous fluids. The spirit employed is alcohol, either diluted or undiluted, either pure or medicated; and, in some cases, ether. The form of tincture is one much used in pharmacy: it presents the active principles of drugs in a small volume; it can be preserved in an unaltered state for a long time; and is, in most cases, well adapted to unite with other substances in extemporaneous prescriptions. Tinctures are made by maceration, or by displacement.


Maceration is an operation in which, by the action of a fluid at common temperatures, continued for a certain length of time, a solution of the principles of an organic substance in that fluid is obtained. As before stated, in making tinctures, the strength of the spirituous menstruum employed is varied according to circumstances. If the substance to be acted upon is resinous, alcohol is to be employed; if it contains also matters soluble in water, and insoluble in pure alcohol, this article, in a diluted state, is to be used. The period of maceration varies from a few hours to several weeks; and, during the process, the vessel in which it is performed should be kept closed, to prevent evaporation. When the spirit has remained upon the substance for the directed period, it should be decanted, instead of being left, as is too often the case, standing on the dregs until it is used. This practice is erroneous, and may be attended with evil consequences; for the characters and properties of a tincture which has been suffered to remain too long in contact with the solid ingredients, will often be found to differ considerably from what they would have been if the process had been terminated at the indicated time. When a tincture is made by maceration, it requires to be filtered, after being separated from the dregs; otherwise, it will be turbid, and deposit much insoluble matter on standing.

Displacement.—This process is of comparatively late introduction in the making of tinctures, &c. For a great proportion of these preparations, it is decidedly to be preferred to maceration; but for others it has not proved so satisfactory. Mohr and Redwood observe, on this point:—

“When tinctures are made in large quantities, displacement is never likely to supersede maceration, on account of any practical advantages it may possess. If the prescribed directions be duly attended to, the process of maceration is unexceptionable. The process is more simple than the other; the mode of operating is more uniform, it is, in fact, always the same; it requires less of skill and dexterity in conducting it; it requires less constant attention during its progress, which, in operating on large quantities, is a consideration; and, finally, the apparatus required is less complicated.

“When only small quantities of tincture are made at a time, and kept in stock, the adoption of the process of displacement will often be found convenient and advantageous. It offers the means of making a tincture in two or three hours, which, by the other process, would require as many weeks. The process being thus completed in so short a time (for the quantity contemplated might be made at one operation), it would not be so likely as the other to experience neglect during the performance of it, or a deviation from the prescribed instructions; the product would, therefore, be more uniform. Finally, in many cases, the tincture and spirit may be removed from the dregs more completely, in operating on small quantities, by this process, than by the other.”

VARIOUS FORMS OF DISPLACEMENT-APPARATUS.



DISPLACEMENT APPARATUS.

DISPLACER FOR
VOLATILE
LIQUIDS.

GLASS DISPLACER.

GLASS DISPLACER.

CONTINUOUS
DISPLACEMENT.

Fig. 16. Apparatus for tinctures. A. The glass adapter, closed by a diaphragm at the lower end, on which is placed some clean sand. The adapter is fitted by a perforated cork into the mouth of the bottle B. C is a tube drawn to a capillary point to permit the escape of air.

Fig. 19. A is an ordinary tin percolator having the rim c soldered around it, to form a water-joint when the lid d is placed in it. a is a perforated diaphragm, and e a tin tube, through which the atmosphere of the bottle B communicates with that above.

Figs. 20 and 21. Common glass chimneys with muslin diaphragms across the lower end.

Fig. 22 shows an arrangement for continuous displacement. Whenever the fluid in the percolator sinks below the mouth of the inverted bottle, the contents of the latter will run out and maintain the supply.

Several forms of displacement-apparatus are used in this country, but the one most frequently employed is that of Boullay. It is a vessel nearly cylindrical, made of tin, with its lower end funnel-shaped. It is made of various sizes; with the cylinder 12 inches in length by $2\frac{1}{2}$ in diameter, or 14 by 4, or 17 by 6. A movable metallic plate, pierced with numerous holes, and provided with a knob or handle in the centre, and fitted accurately to the lower part of the cylinder, is to be placed in the projecting supports. On this, a thin stratum of carded cotton, covered with the powder to be operated upon, is to be laid, and the process carried on as directed above. This apparatus should be furnished with a stopcock at the lower end. Soubeiran has made an improvement on this filter by adapting to it a tin receiver, provided with a cock at the lower part, so that the filtered liquor can be drawn off at will. The substance to be exhausted must be reduced to various degrees of division in different instances; but, for the most part, the form of coarse powder is the most suitable, because, when the powder is very fine, the fluid is apt to be arrested in its passage, after becoming saturated with the soluble principles. In general, it is proper to soak the powder for a few hours with a sufficiency of the spirit, to make it into a stiff pulp, before putting it into the cylinder; otherwise, the spirit may make narrow passages for itself, and leave a part of the material unacted on; but, in some instances, no such preliminary step is necessary, and here, as in the case of tincture of myrrh, the best procedure is to introduce the powder in the dry state, and to pour the spirit over it. Before the spirit is poured over the pulp or powder, it must be packed in the cylinder. The degree of pressure to be used, which varies in

different instances, is to be learned only from experience in each case; and it may be observed, that this is the operation on the correct performance of which the success of the process chiefly depends, and the only one which requires some skill and nicety of management. If the material be loosely packed, the spirit will pass through too quickly, and will not be sufficiently saturated; if too firm pressure be employed, percolation will take place too slowly, or not at all. On the whole, the firmness of the packing should be inversely as the solvent power of the spirit upon the solid materials; but to this rule there are many exceptions. Since the fluid, for an obvious reason, passes after a time with increasing speed, it is occasionally necessary to press down the matter in the cylinder as the process advances; but neither does this rule always apply. In some operations, especially those on the large scale, it is found useful to promote the displacement by pressure. This may be produced simply by supplying the solvent through a tube several feet high, adapted to the upper end of the cylinder, and terminating at the top in a funnel. A still better method, as involving the use of less spirit, consists in the use of compressed air. By means of a condensing syringe, or a column of water or mercury in a tube, air is condensed in the bottle below; and, when the material and spirit have been introduced into the percolator, the top is tightly secured by a screw, and the compressed air admitted by a stopcock. This method answers excellently on the small scale with a column of mercury.

The solution which first passes through in this process is always in a state of high concentration. In general, it is a simple solution of the soluble ingredients of the crude drug in the fluid employed. But sometimes the solvent, if compound, is resolved into its component parts; and the fluid which passes through is only one of these, holding the soluble parts of the drug in solution. Thus, if diluted alcohol be poured over powder of myrrh, in the cylinder, the fluid which first drops into the receiver is a solution of oily consistence, composed of resin and volatile oil dissolved in alcohol. And, if powder of galls be acted on in like manner, by aqueous sulphuric ether, two layers of fluid are obtained—a highly concentrated solution of tannin in the water of the ether, and a weak solution of the same principle in pure ether.

Displacement is accomplished, in the manner here described, with little or no intermixture of the liquid above, with that which is below, in the pulpy mass. If, after the pulp has been allowed to drain till drops cease to fall, the quantity of spirit retained by it be ascertained—that is, by subtracting what has dropped from what has been used to make the pulp,—and the same quantity be poured gently over the mass, repeatedly, as soon as the dropping caused by each successive addition ceases, a series of liquids will be obtained by the simple displacement of one another in the pulp; and the last portion of spirit used may be obtained by pouring over the pulp the same quantity of water. Hence, one of the many advantages possessed by this method of making tinctures, over the ordinary mode by maceration and expression, is that no part of the tincture is lost by being left behind in the residuum.

Even when a strict attention is paid to the prescribed formula, the resulting tinctures may be found to vary in medicinal strength; for any variation in the activity of the drug will influence the tincture. These preparations, although retaining their properties better than other vegetable solutions, will gradually undergo changes, from age and exposure, either losing their color, depositing insoluble matter, or, in some cases, as in that of the tincture of kino, becoming less astringent, and gradually gelatinizing.

WINES are tinctures in which the menstruum is wine, Sherry wine is ordered to be employed by the U. S. and British Pharmacopœias. The solvent power of wines on vegetable products depends on the water and alcohol they contain; the other principles found in them are injurious to it as an excipient; the mucilage disposing to fermentation, and the coloring matters and tannin often decomposing the active principles of the vegetable substances dissolved in it. On this account, these preparations are now going out of use, and mixtures of alcohol and water, of the proper

strength, substituted as solvents. These can always be made of definite strength, which can seldom be attained when wine is used.

Medicated wines are made by maceration or displacement; but the latter process is seldom used. In whatever way they are prepared, they are very liable to undergo a change, and hence should be made in small quantities only, and be kept in well-closed bottles, and in a cool and dark place.

VINEGARS are solutions of certain parts of vegetables in distilled vinegar or diluted acetic acid. Few of these preparations are officinal; but, in some cases, the state of solution in distilled vinegar is the best that can be desired, at least as far as energy is concerned. This is particularly the case when the activity of the medicinal substance depends on the presence of an alkaloid, which, by uniting with the acetic acid, forms a salt that is readily soluble; and, therefore, may be more perfectly extracted than by other menstrua.

As even distilled vinegar contains vegetable matter, rendering it liable to decomposition, it has been found advantageous to substitute acetic acid, properly diluted. In consequence of this liability to become decomposed, medicated vinegars should be made in small quantities, and be renewed at short intervals. In many cases, the acetates of the vegetable alkaloids, dissolved in water or diluted alcohol, may be substituted for medicated vinegars, producing all their good effects, and attended with but few of their disadvantages.

MIXTURES are liquid medicines containing two or more ingredients, generally of extemporaneous preparation, though a few are recognized as officinal. They are in very constant use in the administration of remedies. The ingredients which usually enter into their composition are salts, and other solid bodies which are soluble, or readily miscible with aqueous fluids; also tinctures, spirits, syrups, decoctions, &c.

In making a mixture, the objects to be attained are the perfect solution of the soluble ingredients, and an equal diffusion of those which are not so, through the excipient, as well as the thorough admixture of all the constituents, so that each dose shall be similarly composed. The combination of articles in a mixture is effected either by agitation, or by rubbing the ingredients together in a mortar. When volatile substances enter into a mixture, they should be added last. Insoluble powders, which are easily diffused through a fluid, may be mixed with the liquid ingredients, by merely shaking them together in the bottle. When, however, powders do not mix readily with liquids, they must be triturated in a mortar with a small quantity of the fluid, gradually adding more, until they are of the consistence of thin paste. In the same manner, extracts and electuaries are to be rubbed down with a portion of the excipient previous to putting them in the bottle.

Draughts differ from mixtures only in being designed to be taken at a single dose. This is not a common mode of prescribing medicines, but in some cases it is advantageous, as, when it is wished to apportion the doses of a medicine accurately, or when the medicine is liable to undergo a change from the action of air.

Emulsions are mixtures containing substances of an oleaginous or resinous nature suspended in aqueous fluids by the aid of gum, syrup, yolk of egg, or any viscid matter. It is a mechanical compound, and the union of its ingredients is promoted and rendered more or less permanent by the addition of an alkali. Emulsions are made by triturating certain oleaginous seeds with water, or by mixing the oil procured from them with the necessary ingredients; and also from gum resins in the same manner. Sometimes they are made with oils and an alkali. In making an emulsion of the oils or oleo-resins with mucilage, the former should be gradually added to the latter; by adding the mucilage to the oil, a good emulsion cannot be formed. Mucilage is preferable to an alkali in making many emulsions, as those with castor oil or copaiba; an alkali is preferable in forming an emulsion of almond or olive oil; but an emulsion formed with either of these agents alone will often separate, if the other be added. The presence of soluble salts in an emulsion is also apt to cause a separation of the oil. Much spirit will produce the same effect in emulsions

made with mucilage, and an acid in those made with an alkali. Some substances cannot be formed into good emulsions either with an alkali or with mucilage. This is the case with spermaceti and oil of turpentine. In such cases, the yolk of an egg is the best agent for effecting the admixture, by rubbing down the ingredients in a mortar.

Volatile oils are most readily made into an emulsion by being previously mixed with one of the terebinthinate fixed oils. Scammony is formed into an emulsion by means of milk; but resin of jalap will not unite with this fluid, and is best emulsified by triturating it with almonds and water. Emulsions should be made with cold water, and strained.

Mr. W. Procter, Jr., gives the following valuable directions for making emulsions (*Phil. Journ. Pharm.* xv. 11): "In making an emulsion, a good deal depends even on so slight a circumstance as the form and material of the mortar and pestle. An emulsion may, indeed, be made in a smooth porcelain mortar, but the process is unquestionably more successful and easy in a Wedgewood one, and still more so in the old-fashioned marble mortar, which is generally used to this day by the druggists and apothecaries in England. But whatever be the composition of the mortar, it is essential that it should be perfectly *round* at the bottom; not flattish, as is sometimes the case; and the pestle should be so formed as in its motion to leave no hollows between its base and the concave surface of the mortar. Otherwise, the emulsion will most probably be imperfect. The mucilage or other viscid substance should always be put into the mortar before anything else: the oil (or balsam) may then be very gradually rubbed in, taking care not to add it more quickly than it can be subdued by the pestle; and if, during this part of the manipulation, the mixture should begin to assume a breaking or curdling appearance at the edges, a few drops of water must be immediately incorporated with it, before adding the remainder of the oil. For want of this precaution, I have often known an emulsion suddenly to lose its tenacious consistence in the mortar, and it is then in vain to endeavor to restore it. After the oil is thoroughly incorporated, some care is requisite to avoid separating it again by too hasty an effusion of the water or other fluid of the mixture; and, if any alcoholic or acid liquid is to be added, it must be at the very end of the process. Indeed, an acid liquid, even a slightly acescent syrup, will often entirely destroy the emulsion. Mixtures of copaiba are frequently spoiled by the addition of sweet spirit of nitre; which might be avoided by first diluting it with one or two portions of water."

MEDICATED WATERS.—This term is used by the U. S. Pharmacopœia to designate all preparations of water impregnated with medicinal substances, not appertaining to other classes of preparations. It includes the **DISTILLED WATERS** of the British Colleges, and is a more appropriate designation, because many of the latter preparations are not distilled waters, but are made with an essential oil united to the water by trituration or agitation, no distillation being employed.

By Distillation.—There are two modes of obtaining medicated waters by this process; 1st, by distilling the water directly off the substances; 2d, by employing the essential oil already separated from the vegetable, and distilling it over with the water. In most cases, where the fresh plant can be procured, it is to be preferred. Many of these substances lose their volatile oil and fragrant properties by drying; but, in some instances, the oil is retained, notwithstanding desiccation. Many which lose the oil by the process of drying retain it fully for a length of time, when preserved by being beat into a pulp with common salt; and, when kept in this state, afford very good distilled waters.

The material which supplies the volatile oil is, in general, simply mixed with the water in a state of fine division; and this is probably the best mode, where heat is used in such a way as to exclude the risk of empyreuma. If it does not require to be finely divided, as in the case of fresh leaves and flowers, it may be put conveniently into a net-bag, which, suspended in the middle of the still, may be withdrawn with facility, when its contents are exhausted. Some manufacturers use steam,


instead of water, for obtaining distilled waters: that is, the material to be distilled is spread over a fine gauze partition, or a plate perforated with numerous small holes, and steam is driven through the mass.

When the vegetable substance to be exhausted is a bark, wood, or other solid matter, it must be reduced to a state of moderately fine division. But this is not generally necessary in the case of leaves or flowers, because boiling water breaks down the cells in which the volatile oil is contained. When leaves, however, are thick and leathery, as in the instance of the cherry-laurel, the process is facilitated by chopping them down; and, in most cases, where leaves are large, it is difficult to get a sufficient quantity into the still without cutting them into pieces. In preparing the finer kinds of distilled waters, it is necessary to clean the materials carefully, to remove all decayed leaves or flowers, or those infested by insects, and sometimes also to separate the leaf-stalks, or the green claw of the petals.

Heat may often be applied directly to the vessel; but in this way empyreuma is apt to be occasioned, especially in large operations, in consequence of the solid matters remaining fixed at the bottom. To avoid this, it is usual to apply the heat, in limited operations, through the medium of a solution of hydrochlorate of lime, which raises a temperature between 212° and 270° , according to its strength—or by means of an oil-bath, with a thermometer to regulate the temperature; and, on the great scale, it is best applied by means of steam admitted under pressure into a space surrounding the still.—Another cause of the empyreumatic taint of some distilled waters is the formation of a species of mucilaginous substance, at the expense of the volatile oil. This substance, which forms chiefly when the distillation is pushed too fast, or too far, and is seen encrusting globules of volatile oil, undissolved in the water, is apt to deposit itself on the side of the still, above the boiling materials, where it is afterwards decomposed by the heat. This fact explains the well-known observation, that the finest distilled waters are obtained by gentle distillation, and by abstaining from complete exhaustion of the materials. A still greater improvement is to prepare them with the vacuum-still, in the same way as is now often practiced in making extracts.

[Although a minute description of the process of distillation cannot be given in a work of this kind, yet the mode of conducting it on a small scale will be better understood by reference to the accompanying figures from *Mohr, Redwood, and Procter's Pharmacy*.

FIG. 23.




PHARMACEUTICAL STILL, SEEN IN SECTION.

enables the operator to inspect the progress of the distillation, and to stir the contents of the still when necessary. This opening is stopped with a cork, or a tin cap. *e* is a funnel-tube into which a current of cold water runs during distillation, the warm water running off by the tube on the opposite side.

In using this apparatus, the water-joint should be two-thirds filled with water, the materials introduced, and the head adjusted and filled with water.

Fig. 23 represents a pharmaceutical still, holding about two gallons, made of tinned iron, and intended to fit in the top of a cylinder stove. A is the boiler; B the head, on the inner surface of which the condensation occurs; C the neck communicating with the recipient. *a a* is a rim, soldered around the mouth of the boiler, so as to form a water-joint; *c c* is a circular rim, soldered on the base of the head, in such a manner that the upper part forms a gutter for conducting the condensed fluid from the base of the condensing cone, *d d*, to the neck, C, whilst the lower part projects below into the double rim of the boiler, *a a*, to form the water-joint. *b* is an opening corresponding to the tubulure of a retort, which

FIG. 24.



DISTILLATION OF SPIRITS.

retort stand consists of the foot *x*, of the rod *z*, of the Gay-Lussac holder *a*, and of the rings *e e e*.

FIG. 26.



GAS FURNACE AND FLASK FOR DISTILLATION.

FIG. 25.




Fig. 24 represents Mohr's still, which is very convenient for limited operations. *a*, the retort; *b*, the connecting tube; *c*, the refrigerator, through which passes the condensing tube, represented at *a*, Fig. 25; *d*, the stand for supporting *c* the tub of cold water; *f*, the refuse water.


Figs. 26 and 27. A very simple and convenient arrangement is represented in these figures.

Fig. 26. *a*, funnel for introducing fresh liquid into the flask during distillation. *b*, the connecting tube.

Fig. 27. Liebig's condenser, consisting of the tube *b*, through which a glass tube is seen to pass. This connects by one end with the connecting tube of the flask, and from the other the distillate drops out.

By placing a vessel holding water, and provided with a stop-cock, on one of the rings *e e e*, and by turning the cock, a continuous stream of water is carried through the tube *c* to the lower end of the condenser *b*, then upwards, surrounding the glass tube until it escapes by the pipe *d*. The

FIG. 27.



LIEBIG'S CONDENSER.

[One of these three forms of apparatus will be found very convenient in conducting the distillatory processes of the shop.—*Editor*.]

The other mode of obtaining medicated waters by distillation is by employing the volatile oil, previously separated from the plant, for the vegetable itself. This mode of operating affords a product less liable to change than the above; but it is not so aromatic as when obtained from the fresh plant.

Distilled waters, however carefully they may be kept, are apt, sooner or later, to lose their aroma; and some of them even become mouldy, and acquire thereby an unpleasant odor. They have been thought to keep better with the addition of about a fortieth part of rectified spirit; which may be either put into the still with the water, or added afterwards to the distilled fluid. But the advantages of this addition, although sanctioned by the authority of the Dublin Pharmacopœia, are doubted by practical men. It is believed that the most effectual precaution for preserving them is to prepare them with extremely pure, natural waters, such as snow, rain, or very fine spring water (Müller), free, especially, of any unusual proportion of carbonic acid; and to keep them in black, orange, or red bottles, instead of bottles of clear glass (Hanle). A better mode of procedure is to re-distil the water as soon as any change is perceived in it; this restores its original odor, and renders it less subject to alteration.

By Admixture.—Another method of making medicated waters is by impregnating the water with the volatile oil, by triturating them together with the addition of carbonate of magnesia, and filtering to remove the latter. This affords an excellent product, as it is pure and permanent. The addition of the carbonate of magnesia is merely to enable the operator to produce such a minute division of the oil as will enable water to act on it more efficiently. Other powders act in the same way, and are preferred by some manufacturers.

DISTILLED, ESSENTIAL, OR VOLATILE OILS.—Volatile oils are contained in vegetable cells, generally peculiar, and often so large as to be distinct to the naked eye. Sometimes they exist in such abundance that the oil may be obtained by mere expression. The oils of lemon, orange, bergamot, and citron are prepared in this way by the manufacturer; and from many other substances, such as the unripe germen of rue, and the undeveloped corolla of the clove-tree, oils may be squeezed out by pressure with the nail. In some rare instances, as in that of the Liquid Borneo-Camphor from the *Dryobalanops Camphora*, and that of the Laurel-oil of Guiana from a species of *Ocotea*, volatile oil is obtained largely, and of considerable purity, by exudation from incisions. Much more frequently, however, it is exuded spontaneously, or from incisions, as a turpentine in combination with resin, or as a gum-resin in union with both gum and resin. Most frequently of all, it cannot be obtained by any of these modes, but adheres with more or less force to the flowers, leaves, fruit, bark, or wood, which contain it. In that case, it is sometimes destroyed or dispersed when the plant is dried, more especially if the organ which contains it is the flower; but often, when contained in the leaf, and very generally, if contained in the seed, bark, or wood, it is retained in part, or altogether, under desiccation, and even under long keeping. In some instances, it appears that the volatile oil obtained from plants does not exist ready formed, but is produced, on bruising or distilling them with water, through the reaction of other principles on one another. Of this mode of production, two remarkable examples exist—the volatile oils of the bitter almond and black mustard seed; and it is probable that the oils of cherry-laurel leaves, peach leaves, and of the leaves of other amygdalaceous plants, are similarly circumstanced.

A few volatile oils used in medicine are obtained by expression, such as the oils of orange, lemon, and bergamot, already mentioned. When oils exude along with resin in the form of turpentine, they may be separated from the resin by heat alone, as they are volatilizable at about the temperature of 400°. But thus obtained, they cannot be of fine quality, because the heat required is too near that at which resins, as well as the oils themselves, undergo decomposition. In general, therefore, this plan is not followed, for pharmaceutic purposes; and volatile oils are obtained from turpentine by distilling them with water. For, although they do not enter into

ebullition at the temperature of boiling water, their vapors pass over in large quantity with steam; and, condensing along with it, form distilled waters—with volatile oils either floating on the water, or, more rarely, sinking to the bottom. The volatile oils of many vegetable substances are to be separated in this way alone, because the other principles contained along with them in the crude substances, are empyreumatized by the higher temperature which is required to disengage the oil without the co-operation of watery vapor. In some instances, it has been thought advantageous to substitute for water in the still a strong solution of common salt, because a somewhat higher temperature is required to boil it. There seems an advantage, in this change, for the rectification of oils previously obtained from the raw materials in which they reside; but the advantage in the case of distillation from crude vegetable substances is doubtful.

The method of distilling volatile oils differs little from that described above for preparing distilled waters. The same precautions must be observed in applying heat, and for the same reasons. The formation of mucilaginous matter at the expense of the oil, towards the close of the distillation, is shown by the globules of oil coming over enveloped in a fine pellicle of it. The quantity of water must be proportionally less, or rather, the same portion of water is to be used with successive portions of the material which yields the oil; otherwise, a material loss is sustained by solution of the oil in the water. In some instances, where the oil exists in low proportion, and is of great value, the distilled fluid should be left at rest for some time, and exposed to as low a temperature as can be commanded.

The mixed vapors which pass over condense into a milky-looking fluid, which, after standing some time in the receiver, separates into two portions, one a solution of a part of oil in water, and the other of the oil itself, which occupies the upper or lower part, according as it is lighter or heavier than the water.

Chevallier gives the following rules for the distillation of volatile oils:—

1. To operate upon large quantities, in order to obtain a greater product, and of better quality.
2. To conduct the distillation rapidly.
3. To divide the substances minutely, in order to facilitate the extrication of the oil.
4. To employ only sufficient water to prevent the plant from burning.
5. For substances whose oil is heavier than water, to saturate the water in the still with common salt, to raise the boiling point, and thus to enable the vapor to carry over more of the oil.

FIG. 28.





FIG. 29.




Figs 28 and 29. Various forms of pipettes for separating liquids, filling vials, &c. (See p. 530.)

6. To employ, when possible, water which had already been distilled from off the same substances, and had thus become saturated with oil.

7. For oils naturally fluid, to cool the liquid in the refrigeratory frequently; but to retain it at 80° to 90° for those oils which easily become solid.

After the distillation, the oil should be separated from the water; for this purpose, several ingenious instruments have been devised, a few of which are represented in figures 28 to 33 inclusive.

FIG. 30.




SEPARATING-FUNNEL.

FIG. 31.




SEPARATOR.

FIG. 32.



SEPARATOR.

FIG. 33.



SEPARATOR FOR HEAVY OR LIGHT OILS.

Fig. 30. The ordinary separating funnel, having a perforated ground-glass stopper.

Fig. 31 is a vessel for separating ether, or other volatile liquids from heavier ones.

Fig. 32 answers very well when the quantity of the fluids is very small. By inclining it, the heavier liquid escapes by the orifice *a*.

Fig. 33 is a separator for either heavy or light oils. The former can be drawn off by the orifice of the tube *b*; the latter by the tubule *d*.

The following table, from *Christison's Dispensatory*, showing the proportion of volatile oil obtained from the chief medicinal vegetable substances, according to the most recent experiments, will be found useful. The data are chiefly extracted or calculated from experiments by M. Raybaud in the *Journal de Pharmacie*, xx.—by Dr. Martius in *Repertorium für die Pharmacie*, xxxix.—by Dr. Bley in the same work, xlviii.—by M. Dann and by M. Voelter in the same work, lv.;—and a few have been added from experiments of Dr. Christison himself. The numbers represent the number of ounces obtained from 100 pounds avoirdupois. The letters before the figures refer to the authority for each.

	Author.	Ounces in 100 lbs. av.
Amygdalus communis. Bitter almond	Ra.	0.38
Amygdalus communis. Bitter almond	Vo.	7.70
Amygdalus communis. Bitter almond	Duflos.	0.8 to 4.80
Angelica Archangelica—dried root	Ra.	4.50
Anthemis nobilis—fresh flowers raised at Grasse	Ra.	0.75
Anthemis nobilis—dried flowers do.	Ra.	1.38
Anthemis nobilis—dried flowers, long kept, Germany.....	Bl.	4.50
Anthemis nobilis—flowers freshly dried	Steer.	5.33
Anthemis nobilis—flowers 12 months dried	Steer.	3.00
Apium graveolens—dried fruit.....	Ra.	9.00
Apium petroselinum—fresh herb, after flowering	Ra.	3.38
Apium petroselinum—dry fruit, France	Ra.	12.0
Apium petroselinum—dry fruit, Germany	Da.	30.0
Artemisia absinthium—fresh herb, Paris	Ra.	2.0
Artemisia absinthium—dried herb, recent, Germany	Ma.	16.0
Artemisia absinthium—dried herb, a year old, Germany	Bl.	8.75
Artemisia absinthium—dried herb, 3 years old, Germany	Ma.	5.0
Artemisia — ? Wormseed of commerce	Ra.	3.0
Artemisia — ? Levant wormseed	Vo.	10.8
Calamus aromaticus—fresh root, Germany.....	Ma.	16.0
Calamus aromaticus—recently dried, Germany	Bl.	17.5
Calamus aromaticus—long dried, Germany.....	Da.	14.3
Carum carui—dried fruit of French commerce	Ra.	50.12

	Author.	Ounces in 100 lbs. av.
Carum carui—dried fruit of German commerce	Ma.	66.5
Carum carui, do. do.	Da.	46.6
Carum carui, do. do.	Vo.	70.0
Caryophyllus aromaticus. Cloves, <i>Bourbon</i>	Ra.	144.0
Caryophyllus aromaticus. Cloves, <i>Cayenne</i>	Ra.	152.0
Caryophyllus aromaticus. Cloves, do.	Bl.	125.0
Caryophyllus aromaticus. Cloves, <i>Molucca</i> : French commerce	Ra.	148.0
Caryophyllus aromaticus. Cloves, do. English commerce	Ra.	112.5
Caryophyllus aromaticus. Cloves, average, German commerce	Vo.	226.0
Caryophyllus aromaticus. Cloves, finest, do.	Da.	250.0
Caryophyllus aromaticus. Cloves, do.	Steer.	272.0
Cinnamomum zeylanicum—cinnamon of commerce	Ra.	1.56
Cinnamomum cassia—cassia bark of commerce	Ra.	12.0
Citrus aurantium—Sweet orange flowers, 1 May, <i>Nice</i>	Ra.	5.0
Citrus vulgaris—Bitter orange flowers, 7 May, <i>Nice</i>	Ra.	5.9
Citrus vulgaris, do. do. 12 May, <i>Carmet</i>	Ra.	4.12
Citrus vulgaris, do. do. 16 July, <i>Paris</i>	Ra.	0.9
Citrus vulgaris, do. do. 14 Dec., <i>Paris</i>	Ra.	6.5
Citrus aurantium—rind of 100 oranges, by expression	Ra.	2.5
Citrus aurantium, do. by distillation	Ra.	2.75
Citrus vulgaris—rind of 100 oranges, by expression	Ra.	4.0
Citrus vulgaris, do. by distillation	Ra.	4.25
Citrus limetta—rind of 100 limes, by distillation	Ra.	2.12
Citrus bergamium—rind of 100 bergamots, by distillation	Ra.	2.9
Citrus limonum—rind of 100 lemons, by expression	Ra.	1.9
Citrus limonum, do. by distillation	Ra.	1.4
Cochlearia armoracia—fresh seeds	Ra.	0.9
Coriandrum sativum—dry fruit of French commerce	Ra.	2.3
Coriandrum sativum—dry fruit of German commerce	Da.	9.0
Croton Eleutheria—casarilla bark	Bl.	5.62
Cuminum cyminum—dry fruit of French commerce	Ra.	44.0
Cuminum cyminum—dry fruit of German commerce	Bl.	32.5
Daucus carota—dry fruit	Ra.	0.66
Daucus carota—fresh root	Ra.	0.14
Dracocephalum moldavicum—flowering herb	Ra.	2.10
Drimys Winteri—Winter's bark (probably, however, only <i>Canella alba</i>)	Ra.	0.50
Eugenia pimenta—pimenta berries, <i>Jamaica</i>	Ra.	12.38
Fœniculum officinale—dry fruit of French commerce	Ra.	33.0
Fœniculum officinale—dry fruit of German commerce	Ma.	56.6
Fœniculum officinale, do. do.	Bl.	83.0
Fœniculum officinale, do. do.	Da.	60.4
Fœniculum officinale—flowering herb, <i>Grasse</i>	Ra.	4.9
Fœniculum officinale—herb after flowering, <i>Grasse</i>	Ra.	6.0
Galipea officinalis—Cusparia-bark of commerce	Ra.	1.5
Genista canariensis—Rhodium wood	Ra.	0.47
Geum urbanum—dry roots	Ra.	0.53
Hyssopus officinalis—flowering herb, <i>Grasse</i>	Ra.	5.30
Illicium anisatum—star anise-fruit	Ra.	34.21
Illicium anisatum—star anise-fruit	Da.	25.5
Juniperus communis—green berries, 12 Sept.	Ra.	3.9
Juniperus communis—ripe berries, 1 Dec., <i>France</i>	Ra.	7.75
Juniperus communis, do. fresh, <i>Germany</i>	Do.	15.5
Juniperus communis, do. a year old, <i>Germany</i>	Ma.	10.8
Juniperus communis, do. a year old, <i>Germany</i>	Bl.	16.25
Juniperus sabina—fresh twigs, 5 March, <i>Grasse</i>	Ra.	19.05
Juniperus sabina—fresh twigs, 2 Oct., <i>Paris</i>	Ra.	14.25
Juniperus sabina—dried twigs, recent, <i>Germany</i>	Ma.	40.0
Juniperus sabina—dried twigs, a year old, <i>Germany</i>	Ma.	25.0
Larix cedrus—fresh cedar wood, <i>Paris</i>	Ra.	0.3
Larix cedrus—cedar wood of commerce	Ra.	4.25
Laurus nobilis—fresh leaves, 26 Jan., <i>Paris</i>	Ra.	5.25
Laurus nobilis—leaves some years dried, <i>Germany</i>	Bl.	4.10
Laurus nobilis, { fresh leaves, } poor soil, low site	Chr.	7.33
Laurus nobilis, { early in Oct. } poor soil, high site	Chr.	6.9
Laurus nobilis, { near <i>Edinb.</i> } very fine soil, low site	Chr.	17.12
Lavandula vera—flowering herbs, 2 Aug., <i>Grasse</i>	Ra.	11.5
Lavandula vera, do. 2 Aug., <i>Grasse</i> , north exposure	Ra.	9.12

	Author.	Ounces in 100 lbs. av.
Lavandula vera—flowering herb, 26 July, <i>Souzeillas</i>	Ra.	9.0
Lavandula vera—herb after flowering, 26 Sept., <i>Souzeillas</i>	Ra.	15.0
Lavandula spica—fresh herb, 24 July, <i>Paris</i>	Ra.	7.62
Lavandula spica—fresh herb, 4 Aug., <i>Grasse</i>	Ra.	12.5
Lavandula stæchas—dry spikes	Ra.	6.43
Ligusticum levisticum—fresh herb, <i>Paris</i>	Ra.	1.12
Melissa officinalis—fresh flowering herb	Ra.	0.25
Mentha piperita—fresh tops in flower, <i>Grasse</i>	Ra.	6.25
Mentha piperita—fresh tops in flower, <i>Paris</i>	Ra.	3.40
Mentha piperita—dried tops in flower, <i>Germany</i>	Bl.	15.62
Mentha piperita—dried tops in flower, <i>Germany</i>	Ma.	21.0
Mentha pulegium—fresh flowering herb	Ra.	1.0
Myristica moschata—mace of commerce, finest.....	Vo.	154.0
Myristica moschata, do. fine.....	Bl.	125.0
Myristica moschata, do. worm-eaten.....	Bl.	65.6
Myristica moschata—nutmegs of commerce, fine	Bl.	108.25
Myristica moschata, do. worm-eaten.....	Bl.	64.1
Myrtus communis—fresh leaves, September 20, <i>Grasse</i>	Ra.	4.5
Myrtus communis—fresh leaves, September 6, <i>Paris</i>	Ra.	2.5
Origanum majorana—fresh flowering herb, August 3, <i>Grasse</i>	Ra.	8.5
Origanum majorana do. August 3, <i>Paris</i>	Ra.	4.4
Origanum vulgare do. September 15, <i>Paris</i>	Ra.	0.4
Pimpinella anisum—dry fruit of French commerce	Ra.	35.12
Pimpinella anisum—dry fruit, new, German commerce	Ma.	37.5
Pimpinella anisum—dry fruit, old do.	Ma.	27.0
Pimpinella anisum—dry fruit of do.	Vo.	25.0
Pimpinella anisum—dry fruit of German commerce.....	Da.	43.75
Piper cubeba—Cubebs of French commerce	Ra.	19.5
Piper nigrum—White pepper of French commerce.....	Ra.	16.0
Piper nigrum—Black pepper of do.	Ra.	18.12
Prunus lauro-cerasus—fresh leaves, November 23, <i>Paris</i>	Ra.	2.12
Prunus lauro-cerasus, { fresh leaves } undeveloped, June 7	Chr.	10.13
Prunus lauro-cerasus, { from the same } half-grown, June 7	Chr.	7.20
Prunus lauro-cerasus, { plants: near } full-grown, 8 weeks on tree,		
Prunus lauro-cerasus, { Edinburgh. } July 15.....	Chr.	4.96
Prunus lauro-cerasus, { } 12 months on tree, June 2...	Chr.	1.04
Prunus lauro-cerasus, { fresh leaves of the } 3 months on the tree.	Chr.	7.04
Prunus lauro-cerasus, { same plant, 1 Sept. } 15 months on the tree	Chr.	2.24
Prunus lauro-cerasus, { 1836, Edin. }		
Renealmia cardamomum—lesser cardamoms	Ra.	11.42
Rosa centifolia—fresh flowers, <i>Grasse</i>	Ra.	0.25
Rosmarinus officinalis—fresh flowering herb, <i>Grasse</i>	Ra.	5.0
Rosmarinus officinalis, do. <i>Paris</i>	Ra.	3.5
Ruta graveolens—fresh flowering herb, 20 July, <i>Grasse</i>	Ra.	4.12
Ruta graveolens, do. 28 July, <i>Paris</i>	Ra.	0.63
Ruta graveolens—flowering herbs, newly dried, <i>Germany</i>	Bl.	4.4
Ruta graveolens—dried seeds, <i>South of France</i>	Ra.	19.0
Salvia officinalis, v. <i>minor</i> —fresh herb, 12 Mar., <i>Grasse</i>	Ra.	6.0
Salvia officinalis, v. <i>minor</i> —fresh herb, 14 June, <i>Paris</i>	Ra.	2.5
Salvia officinalis, v. <i>major</i> —fresh herb, 12 Mar., <i>Grasse</i>	Ra.	4.0
Salvia officinalis, v. <i>major</i> —fresh herb, 14 June, <i>Paris</i>	Ra.	3.05
Santalum album—sandal-wood of commerce	Ra.	5.0
Sinapis nigra—black-mustard-seed, <i>Germany</i> , 12 months old.....	Da.	3.9
Sinapis nigra—black-mustard-seed, <i>Germany</i> , fresh.....	Da.	5.0
Sinapis nigra—black-mustard-seed, <i>France</i> , fresh.....	Da.	7.75
Sinapis nigra—black-mustard-seed, <i>France</i>	Vo.	9.1
Tanacetum vulgare—fresh flowering herb, 9 July, <i>Grasse</i>	Ra.	1.2
Tanacetum vulgare—fresh flowering herb, 25 July, <i>Paris</i>	Ra.	5.8
Tanacetum vulgare—fresh tops, <i>Germany</i>	Da.	5.0
Tanacetum vulgare—dried flowering herb, <i>Germany</i>	Bl.	15.6
Thuya occidentalis, { fresh } aged, stunted tree; exposed. Oct. 21	Chr.	10.8
Thuya occidentalis, { twigs } aged, vigorous; sheltered. Oct. 21....	Chr.	10.25
Thuya occidentalis, { near } young, vigorous; exposed. Oct. 9.....	Chr.	18.25
Thuya occidentalis, { Edin. } young, vigorous; exposed; fine soil,		
Thuya occidentalis, { } Sept. 26	Chr.	26.40
Thymus serpyllum—fresh flowering herb, 6 Aug. <i>Grasse</i>	Ra.	5.0
Thymus serpyllum—fresh flowering herb, 5 July, <i>Paris</i>	Ra.	0.9

	Author.	Ounces in 100 lbs.av.
Thymus vulgaris—fresh flowering herb, 16 Aug., <i>Grasse</i>	Ra.	6.5
Thymus vulgaris—fresh flowering herb, 13 July, <i>Paris</i>	Ra.	3.75
Valeriana officinalis—dry root, a year old, <i>Germany</i>	Bl.	30.16
Valeriana officinalis—the root, <i>Germany</i>	Da.	15.0
Valeriana officinalis—the root, <i>Germany</i>	Vo.	10.5
Verbena odorata—fresh flowering herb, <i>Paris</i>	Ra.	3.1
Zingiber officinale—dry root of commerce	Ra.	10.8

Volatile oils should be preserved in dark bottles, carefully closed and nearly full. When kept for any time, they are apt to undergo certain changes, becoming dark-colored, losing their agreeable odor, and growing thick and clammy. It is then necessary to re-distil them with a certain quantity of water, by which the undecomposed portion is again obtained in a pure state. Another mode is to agitate with recently heated animal charcoal; this restores their clearness, and in a great measure their fragrance.

FIXED OILS AND FATS.—Fixed oils are obtained from the seeds of vegetables, by expression; by boiling the bruised seeds in water, and removing the oil that rises to the surface; and by dissolving out the oil by means of a menstruum.

By Expression.—This is accomplished in several ways: By means of the wedge press, or by a screw or hydraulic press, the former being the more generally employed. In almost all cases, it becomes requisite to heat the seeds gently, to render the oil more liquid; but care must be taken that the heat is not too great; otherwise, the oil will become of a dark color, and acquire an unpleasant taste. It is by this method that linseed and castor oils are obtained. The oil, as it comes from the press, is seldom pure or fit for use; it is freed from these impurities by boiling with water, and separating the pure oil. These oils are often colored, owing, in most cases, to heating the seeds too much; this is peculiarly the case with linseed oil, which is seldom seen of a light color.

By Decoction.—This is effected by boiling the bruised seeds in water, and skimming off the oil as it rises to the surface. This plan is also employed in making cod-liver oil, as well as in the manufacture of castor oil in the West Indies.

By Solution.—This affords a very pure oil, but is seldom employed on account of expense. It is, however, useful in facilitating the extraction of some of the thick and viscid oils. Thus, croton oil is more readily obtained by mixing the ground seeds with half their weight of alcohol, and, after letting the mixture stand for some time, submitting it to pressure, and distilling off the spirit from the product.

Solution in ether is also useful in some cases, when expense is not an object; oil of ergot is best obtained in this manner.

The solid oils, as oil of cocoa, &c., require the aid of heat in their expression, and the ground material containing them is therefore placed between heated plates in the press.

Animal Fats.—Formerly, the fats of many animals were employed in pharmacy; but, at present, those principally used are lard, suet, and beef's marrow. To render these fit for pharmaceutic purposes, the crude material is to be cut into small pieces, freed as much as possible from all extraneous substances, and placed in a boiler with water, and heated until it is fused; the heat is continued till the water has evaporated, and the fluid fat is then strained. These fats are best preserved by being run into glazed jars, and kept from the action of the air.

ALKALOIDS.—The vegetable alkaloids may be obtained by a variety of processes; but these invariably comprehend decomposition of the alkaloidal salt in the crude drug, either by the superior affinity of an alkali, earth, or alkaline carbonate, or by double decomposition with some compound salt whose base forms an insoluble salt with the acid in the drug. Active neutral principles—such as narcotin from opium,

piperin from white pepper, picrotoxin from *cocculus indicus*, and elaterin from *elaterium*—may be obtained through the agency of such simple solvents as water, rectified spirit, and sulphuric ether, used singly or successively. But no alkaloid can be thus isolated.

The solutions from which vegetable alkaloids are prepared, are, in general, obtained best by the method of percolation, or displacement. Some experience is required to apply this process in all cases with success. But, when well performed, it is greatly superior, in general, to any other mode of extracting the active matters of vegetable drugs, in cases in which the liquid used is spirituous or ethereal; and it is often not less advantageous in the instance of water, as well as acidulous fluids. The precautions for applying it successfully have been considered under the head of tinctures.

SPIRITS are alcoholic solutions of volatile principles obtained by distillation.

When spirit is distilled with aromatic vegetables which contain volatile oil, the oil, for the most part, rises with the spirituous vapor, and condenses along with it in a state of solution. In some cases, the volatile oil rises with the vapor of strong spirit, so that alcohol may be employed for the purpose; and this is occasionally necessary for keeping the oil of the distilled spirit in solution. In other instances, the oil does not begin to pass over until watery vapor also passes in considerable proportion with the spirituous vapor; so that diluted alcohol is required for the process.

For most medicinal purposes, but especially the present, the menstruum should be made by diluting alcohol with water.

The best apparatus for preparing distilled spirits is that represented at page 527, fig. 24. It has been proposed to prepare them by distillation in a vacuum-still; but many volatile oils will not rise with spirit-vapor at the low temperature at which spirit boils in a vacuum.

These preparations, like distilled waters, may be obtained either from crude vegetable substances, or from their volatile oils. They are seldom, however, of such fine aroma when prepared in the latter as in the former way, unless care be taken to employ those volatile oils only which have been recently as well as carefully distilled.

Essences differ from spirits in being volatile substances dissolved in alcohol; but this name is generally applied to alcoholic solutions of the volatile oils.

TROCHES, OR LOZENGES, are small, dry, solid masses, consisting of powders incorporated with sugar and mucilage. Some writers include under this name *drops* and *pastes*, making the following distinction between them:—

Lozenges, when the principal basis is sugar, and when the ingredients are combined without the aid of heat.

Drops, when the principal basis is sugar, but when the ingredients are combined with the aid of heat.

Pastes, when the principal basis is a vegetable juice or pulp, and when the mixture is of a soft consistence.

Lozenges are much more employed in Europe than in this country as a mode of administering medicines. In making them, the sugar is employed in a powdered state; the more active ingredients added in powder, or in a liquid state; and the whole mixed into a paste by the addition of mucilage. The mucilage generally used is that of tragacanth, as being more tenacious than that of gum Arabic. After the ingredients are properly incorporated, the paste is rolled out into a uniform sheet on a marble slab, previously sprinkled with some powdered starch to prevent adhesion, and cut into small cakes by means of a punch. These cakes are then to be placed on sieves, and kept in a drying-room until they have become perfectly dry and hard, when they are to be sifted, and kept in well-closed bottles.

Drops are formed from coarsely-powdered sugar, to which the flavoring or medicinal ingredients are added in a liquid state, so as to moisten the sugar. The mix-

ture is then melted in a proper vessel, over a clear fire, after which the melted mass should be allowed to drop in small portions on a marble slab or greased metallic plate, and when cold, be removed, and kept like lozenges.

Pastes are usually formed of inspissated vegetable juices, or decoctions, with the addition of sugar, gum, &c. These are to be brought to a proper consistence, and the mixture poured into flat, shallow moulds, or rolled out, as in the case of lozenges, and divided into pieces of the desired size.

INHALATIONS.—These have, at different times, been much in vogue as a means of combating disease, especially complaints of the respiratory organs. Within the last few years, much interest has been excited relative to this mode of administering medicinal agents, from the remarkable anæsthetic effects caused by the inhalation of the vapor of ether and chloroform. The simplest form of using either of these is by pouring the requisite quantity on a hollow sponge, and applying this over the mouth of the patient. Various forms of inhaling apparatus have been invented, which will be found described in *Mohr and Redwood's Pharmacy*, 535–9, with directions for their use.

EXTERNAL REMEDIES.

BATHS.—By the term bath is meant the complete or partial immersion of the body in a fluid or gaseous medium, differing, in some circumstances, from that to which it has been accustomed. In order to obtain the full efficacy of a bath, without injurious consequences, its temperature should be regulated, and particularly prescribed by the physician ordering it. The following temperatures are those given by Dr. Forbes in the *Cyclopædia of Practical Medicine*:—

Cold Bath	between 33° and 60° F.
Cool Bath	" 64° and 75° F.
Temperate Bath	" 75° and 85° F.
Tepid Bath	" 85° and 92° F.
Warm Bath.	" 92° and 98° F.
Hot Bath	" 98° and 112° F.

The effects of these different baths on the system are very dissimilar, according to their temperature, and the time during which the patient is subjected to their influence.

COLD BATH.—When a person plunges into a cold bath, he is first sensible of a sudden sensation of cold upon the surface, accompanied by an oppression of breathing, causing this function to be performed in convulsive gasps. This is called the *shock*, and is caused by a rapid contraction of the cutaneous capillaries, and a retrocession of the blood to the lungs and other internal organs. In a short time, the difficulty of breathing disappears, the temperature becomes agreeable, and if the person now leaves the water, a warmth of the surface comes on, termed the *glow*, succeeded by a sense of invigoration of the whole system. Should the person remain in the water for too long a time, another train of symptoms manifest themselves; the sensation of cold soon attains to an unpleasant degree of chilliness, followed by rigors; a bluish tint is perceptible on the surface of the body; the blood accumulates in the internal organs; and, on leaving the water, there is no reaction, or a very feeble one, the surface remaining cold, the extremities benumbed; and headache, difficult respiration, often pain in the chest, ensue, with a sense of depression and lassitude. The use of proper means will often remove these symptoms; but they may lead to a variety of diseases of the internal organs.

The objects, in prescribing a cold bath, are the production of a sudden and powerful impression on the nervous system, and the tonic influence it exercises

when followed by due reaction. In the first of these, it has been found useful in certain affections where there is a derangement of the functions of sensation, of motion or sensation unattended with a congestive or inflammatory condition of the internal organs; but it is more frequently ordered to fulfil the second indication.

It is always contra-indicated when, from debility, the system does not react so as to produce a glow; when there is a tendency to congestion of the cerebral vessels, or any serious organic affection of the heart, lungs, or kidneys. In all cases, it is advantageous, before taking the cold bath, to take such exercise as will raise the circulation, without occasioning fatigue or perspiration, for reaction is almost certain to follow the immersion, except when the person has remained in the water too long a time. The period of immersion should not exceed five minutes.

COOL BATH.—The action and uses of this are similar to the last, but are less powerful. It is, therefore, better calculated for those who are much debilitated.

TEMPERATE BATH.—As the temperate bath is of a temperature closely approaching that of the body, the shock and subsequent reaction are almost wanting. It is, therefore, much more employed for purposes of comfort and cleanliness than as a remedial agent. In delicate persons, it should always be used instead of the cold or cool bath, and is always better suited to very young children than lower temperatures.

TEPID BATH.—This is intermediate in operation between the temperate and the warm bath, and varies in effects and uses according to the temperature. In perfect health, it should not be used as an habitual indulgence; but, for the purposes of cleanliness, an occasional recurrence to it allows of a more perfect ablution than can be effected by cooler baths. It is better to use it about noon, when the first process of digestion of the morning meal is over, and immediately afterwards to take brisk exercise in the open air. In cases of fatigue and febrile irritation, from over exertion or a long journey, the tepid bath is generally found very beneficial. It is also serviceable to persons of sedentary habits, &c. In all such cases, however, it is not to be employed immediately after a meal, or when the individual is unduly excited, either mentally or corporeally. In one class of complaints, those dependent on gastric irritation, the tepid, or even the warm bath proves of much service.

WARM BATH.—The first effect of a warm bath is to produce a sensation of heat upon the surface, and to increase the pulse in quickness and fulness, though in most cases to diminish its tenseness. The cutaneous circulation more especially becomes affected, and the body is increased in bulk, as shown by the increased pressure of ligatures, or of rings upon the fingers. The secondary effects, when the immersion is continued for some time, are muscular relaxation, sometimes to a considerable degree; even after leaving the bath, a disposition to lassitude continues for some time, with a tendency to perspiration.

The remedial effects of a warm bath depend on its temperature, the time a patient remains in it, and the subsequent treatment. The medium time for remaining in the bath is from twenty to twenty-five minutes; but this must be regulated by the effect produced. It is beneficial in incipient catarrh, in some congestions of the internal organs, chronic rheumatism, and in spasmodic affections, especially those of children; but is contra-indicated in active fever, or when there is congestion, or a determination of blood to the head.

In the convulsions of children, its effects are remarkably beneficial, as it not only relaxes spasm, and relieves for the moment, but soothes nervous irritation. In cases where the convulsions are severe, it will be found advantageous to apply cold water to the head. When a warm bath is administered to a child, care must be taken not to expose it to the cold air for the purpose of drying its body; the best plan is to envelop it in a warm blanket, and to place it in bed at once. By this plan, it is not

liable to take cold, which is a common objection to the use of the warm bath for children.

HOT BATH.—From its temperature being above that of the body, the hot bath is far more stimulating than the preceding, as evinced by the excitement of the pulse, the sensations of fulness in the head and throbbing of the cerebral vessels. Its use is principally confined to cases where it is wished to arouse nervous energy and vital action, as in Asiatic cholera, &c.; or where there is a sudden retrocession of cutaneous diseases. It has also proved useful in certain forms of rheumatism, and paralysis. As the intention is mainly to induce excitement, the patient is not to be exposed to its action long enough to cause exhaustion.

Whatever description of bath is ordered to be used, the original temperature is to be maintained during the whole time the patient remains in the water. At the end of some minutes, therefore, the heat should be tested by a thermometer, and, if requisite, hot water added. The sensations of the bather are always a fallacious criterion.

SHOWER BATH.—This is a modification of the cold affusion, being attended with the same effects, but in a less degree. The short duration of it renders it less refrigerant than the cold bath, and causes the primary shock to be the most important part of the influence it exercises, which may be modified by increasing or diminishing the temperature of the water, or that of the height from which the fluid falls. It is used either cold or tepid. The former is most advantageous when the powers of the system are sufficient to cause a reaction, that results in a glow on the surface, soon after using the bath; but, where this is not the case, as in debilitated individuals, the temperature is to be increased. It is adapted to those cases where a powerful impression is wished to be made upon the nervous system, as in chorea, hysteria, &c., and is also of much benefit to persons subject to determination to the head. In such cases, it is found advantageous to make the patient stand in hot water at the time of taking the bath. The reaction following its use is much promoted by using friction to the surface. The improved and portable shower baths, now to be procured, obviate many of the objections to its use which formerly existed. For children, the best and most convenient apparatus that can be employed is one described by Dr. A. T. Thomson. (*Domestic Management of the Sick Room.*)

“It consists of a hollow vessel made of tin, with a perforated bottom. The body of the vessel is of a bell-shape, with a hollow tube rising from the top, and terminating in a broad perforated rim. When the bath is to be used, it must be sunk in a bucket of water, until it is completely submerged; the air is thus driven out of the bath, through the tube, and the bath filled with water. The thumb of an attendant is then to be placed on the opening in the centre of the rim, and the bath raised from the bucket of water. The pressure of air upon the holes in the bottom retains the water in the bath; and, on raising the thumb from the upper orifice, the whole is rapidly discharged. In using it, the child must be placed in an empty tub, and the bath, being held over his head, is then to be discharged; and the child immediately afterwards dried, with friction. When salt water is used for this bath, the hair should be kept dry, by means of an oil skin cap.”

LOCAL BATHS.—These are most frequently used warm. The most common are the *hip-bath* and *foot-bath*. They are principally employed as revulsives. The first has been found a valuable remedy in diseases of the womb, and in irritations of the pelvic organs. Where it is employed merely to soothe pain, the temperature should not exceed 80° to 90°, and the patient is to remain in it for some time; but when it is intended to excite the uterus to greater action, it should be as hot as can be borne by the patient, though the continuance in it should not exceed ten to fifteen minutes. The hot foot-bath should be of as high a temperature as can be borne, so as to redden the skin of the immersed parts effectually. The vessel used should be sufficiently

deep to allow the legs to be immersed nearly to the knees. It is a valuable remedy in the early stages of catarrh, and local congestions of the head, chest, or abdomen; and in the dyspnoea accompanying hypertrophy of the heart, it has often proved very beneficial. It may be rendered more stimulating by the addition of common salt, carbonate of potassa, or flour of mustard. In torpid states of the liver, a mixture of nitric and muriatic acids, in the proportion of about an ounce of each to every gallon of water, often proves highly useful.

VAPOR BATH.—This consists in either wholly enveloping the patient in the steam of hot water, or merely his body, or some of its parts. A much more elevated temperature can thus be borne than where the body is immersed in water, and its action is more confined to the skin; hence, although less stimulating, it is more diaphoretic than the hot bath. Where the vapor is inhaled, the heated surface being extended, it is more powerful in its effects. It has proved of great benefit where an active revulsion to the surface is indicated, and also in the treatment of cutaneous diseases. It can be applied with little trouble. All that is required is that the patient be seated on a chair with a vessel of hot water placed beside him, and the whole enveloped with a blanket, to be thrown over his head if the vapor is to be breathed, or pinned around his neck where this is not the case. The steam soon surrounds his body, and causes a copious perspiration; and, should it cease too soon to be evolved, its generation may be restored by dropping a heated brick or stone into the water. Where the vapor is not respired, it may be used of a higher temperature.

THE WARM AIR BATH is more stimulating than the vapor bath, producing—especially where the warm air is also breathed—general uneasiness, heat of skin, excitement of the pulse, until a general perspiration ensues. It is readily administered even to persons in bed. The bedclothes should be elevated by a light frame, and the heated air be admitted by means of a tube. It has been found beneficial in chronic rheumatism, and was at one time much praised in the collapsed stage of cholera.

DOUCHES consist in the forcible application of a stream of hot or cold water to a part of the body, in such a manner that the force of the stream of the fluid shall aid the stimulant effect. The hot douche may be formed by pouring from a height hot water from a tea-kettle, or through a tube about an inch and a half in diameter, so as to cause the fluid to strike forcibly upon the affected part of the body, whilst at the same time percussion is employed; or, in other words, the part is beaten by means of a caoutchouc bottle, stuffed with wool, and affixed to a cane handle.

A temperature of 160° is readily borne in douching. When the disease is chronic rheumatism, or when deep-seated pains are to be removed, the douching should be continued for either half or three quarters of an hour at a time; and the patient should be afterwards placed in bed between blankets, in order to encourage sweating. The most decisive advantage, in enlarged, stiff, and painful joints, results from this mode of douching.

In cases of complicated fevers, and in some other diseases, in which inflammation of the brain occurs, the *cold douche* is frequently ordered. The mode of applying it consists in pouring a stream of cold water on the shaved head. The patient should be raised in bed, and the head held over a basin, or other vessel, whilst a stream of cold water is directed on the crown, and the fall of the water gradually increased. After this, the head should be merely dried, but not rubbed. It not only soothes the patient, but often induces sleep, although it usually causes a most uncomfortable feeling at first; but relief is so quickly experienced, that its repetition is generally requested by the invalid.

A *vapor douche* is very stimulating, and when of a high temperature, and continued for any time, will cause vesication. It may be used as a counter-irritant instead of moxa.

MEDICATED BATHS are such as contain medicinal agents in solution in the water, and are used both topically and generally. Saline, acid, and alkaline substances, with iodine and sulphur, are the materials most commonly employed. Medicated baths are sometimes natural, as those of sea water, and certain mineral springs; others consist of solutions of various articles, artificially prepared. Of the medicated baths, sea water is the most generally used, more, however, for pleasure, than for remedial purposes. It is not only stimulating, on account of its saline constituents, and its use followed by a perfect glow, but these constituents also exert a medicinal action on the system, in certain diseases of debility. The nitro-muriatic acid bath has been found useful in diseases of the liver; and alkaline, ioduretted, and alkaline sulphuretted baths, are advantageously employed in many forms of cutaneous disease.

Medicated *vapor* baths are prepared by impregnating aqueous vapor with the volatile principles of medicinal plants, though it is very doubtful whether they produce any effects that cannot be obtained from the simple vapor bath. But the fumes of sulphur, chlorine, camphor, mercury, &c., in combination with aqueous vapor, have been found to exert powerful effects on the system, and to be very effectual in the cure of certain diseases.

AFFUSION.—This consists in pouring or dashing a quantity of water on the body, or a portion of the body, from an inconsiderable height, most generally for the purpose of reducing its temperature; but it is also employed as a revulsive agent, and to give an impulse to the nervous system. Affusions are made with cold or tepid water. Cold affusion is one of the most powerful general means of abstracting caloric from the body, we possess; it not only lessens the temperature of the surface, but it diminishes the action of the heart. It has been successfully employed in the treatment of fevers, but its use requires much judgment, as respects the time of its application, which, according to Currie and others, is, "when there is no sense of chilliness present, when the heat of the skin is steadily above what is natural, and when there is no general or profuse perspiration." It should not be employed either in the cold or sweating stage of fever, or in the hot stage where the heat does not much exceed the natural standard. From inattention to these circumstances, and from patients being injured by its effects, affusion is not at present often resorted to in the treatment of fevers. In scarlatina and some other of the exanthemata, it has been employed with benefit, to reduce the morbid heat of skin. It is, however, more relied upon in inflammatory diseases of the brain; in these cases, the water is to be poured upon the head, inclined over a basin or tub, by means of a pitcher. In children, it is sufficient to squeeze a large sponge, previously saturated with cold water, at some height above the head. Where a general affusion is deemed necessary, the patient should be taken out of bed, his head having been previously shaved; and, being stripped naked, and placed on a stool in an empty tub, from *three to five* gallons of water, at a temperature not under 40° Fahr., should be thrown over him. This affusion should be repeated until he feels cold, or rather until a rigor or shivering, or even chattering of the teeth, is experienced. He should then be dried, placed in bed, and a little warm wine and water administered to him, to aid the reaction, and the consequent perspiration.

But in cases where the invalid faints on raising him into the erect position; where he feels chilly, although the thermometer indicates a high febrile temperature; where the skin is bedewed with perspiration; where the patient is a woman, and the monthly change is present; or where the operation is dreaded, it ought not to be employed. The evening is the best time for using the cold affusion.

Cool affusion is sometimes used instead of the cold, and is preferable for weak, irritable individuals. Tepid affusion is also beneficial in certain cases; especially where there is a fear that perfect reaction will not take place after the application of cold water; or where disease of the pulmonary organs exists. It has been found very efficacious in scarlatina, as also in hectic fever. Warm affusion has likewise been used; but its effects are far more temporary than those of the warm bath.

SPONGING.—The object of sponging the body, either with cold or with tepid water, is to reduce the heat of the surface by means of evaporation, with the view of softening the pulse and diminishing its frequency; of rendering the skin perspirable; of relieving headache, checking delirium, and promoting sleep. In order to perform it well, the patient should be taken out of bed; and, being undressed and placed in a chair, two or three persons should, at the same time, pass large sponges, wet with simple cold or tepid water, or vinegar and water, rapidly over different portions of the naked body, until the whole of it shall have been successively sponged, and a chill has come over the patient, who is then to be dried and placed in bed.

This appears to be a simple process; but it requires both caution and judgment to determine on its use, and to render it not only salutary, but free from danger. For example, if, notwithstanding the continuance of great heat and dryness of the skin, a sensation of chilliness comes over the patient when the sponging is about to be done, the operation should be delayed; neither should it be done if the smallest tendency to perspiration, indicated by moisture in the axillæ, or on the palms of the hands, displays itself. But, if it has been done, and if partial relief has been obtained, if the heat of the skin returns, and the pulse rises, it may again be resorted to and repeated.

But, besides the advantages to be obtained from *sponging*, during the presence of disease, it is also productive of great benefit in warding off diseases, or in a prophylactic point of view. The sponging, however, under such circumstances, is to be performed in a different manner from that which has been described.

In those predisposed to asthma and to bronchitis, known in ordinary language by the term *cold*, or *cough*, and in those, also, who are predisposed to *croup*, nothing is so serviceable in warding off the paroxysm of the former disease, or in preventing the attack of the latter malady, as cold and tepid sponging of the chest and the trunk of the body. It is also equally serviceable during dentition in infants, and as a general tonic in delicate conditions of the system. In these cases, it should be followed by gentle friction over the surface of the body.

Sponging the body, as a prophylactic, or as a tonic, is always most efficacious when it is performed whilst the patient is still in bed. Let him sit up; or, if unable to do so, let him be raised into the sitting position, keeping the lower limbs under the bed-clothes. The night-shirt or night-dress is then to be stripped down, and a large towel or sheet put round the waist, to prevent the bed-clothes from being wetted. The naked trunk of the body, and the upper part of the arms, are then to be moistened with a sponge, or a piece of flannel, dipped either in cold or tepid water, or vinegar and water, or salt water, as the case may demand; after which, in drying the skin, if water only has been used, brisk friction should be employed, so as to cause a general glow upon the surface. In some cases, salt and water, or vinegar and water, are preferable to the simple water. The best proportions are two ounces of *salt*, or the same quantity of common *vinegar*, to a pint of water. When these adjuncts are used, friction is not required; indeed, when salt water is employed, the skin should be dried with a soft, old towel, fitted to soak up the water only, and to leave the saline particles upon the surface.

The best time for using such spongings is just before getting out of bed in the morning. They may be used in winter as well as in summer. In efficacy, they are certainly nearly equal to the shower-bath, which may be resorted to in vigorous habits; but, when the habit is delicate, and not equal to the reaction, on which the beneficial influence of the shower-bath rests, sponging is preferable. It should not be used when perspiration is present.

Sponging is as beneficial as the shower-bath, in rendering the habit less susceptible of cold; and, when conjoined with exercise in the open air, and proper regimen, not only asthma, coughs, and catarrhs, may be warded off by its employment, but the predisposition to tubercular consumption may be lessened, in a very remarkable degree, by its daily employment.

FOMENTATIONS.—These may be regarded as a species of local bathing. Although the decoctions usually employed are useful in aiding warmth, by their soothing or sedative influence, yet they are secondary objects in the application of fomentations, the intention being to convey heat, combined with moisture, to the part fomented. Flannel cloths, wrung out of boiling water, by means of two sticks turned in opposite directions, form the best fomentations. If they be shaken up, and laid lightly over the part, they involve a considerable quantity of air, which, being a bad conductor, retains the heat in them for a considerable time. In every process of fomenting, there should be two flannels, each three yards long, with the ends sewed together, to admit of the boiling water being wrung out of them; and the one flannel should be got ready whilst the other is applied. The fineness or the coarseness of the flannel is not a matter of indifference: the coarser it is, the less readily does it conduct heat, and the longer it retains its warmth; therefore, it is more efficient for fomenting. White flannel also retains the heat longer than colored flannel.

Stuping is a variety of fomentation useful in many cases, but especially in affections of the eyes. The patient should be his own operator. He should sit up in bed, and should place, in the hollow of his hand, a small piece of flannel, wrung out of boiling water, and hold the hand at such a distance under the eyes that the vapors may rise to them—changing the flannel as often as it ceases to give out warm vapor. If narcotic or sedative additions are directed, they should be poured hot upon the flannel, each time it is changed.

CATAPLASMS, or POULTICES, are modifications of fomentations; they generally consist of pulpy substances capable of absorbing much moisture, and of such consistence as to be applied accurately to any surface, however irregular. Their action, in most cases, depends upon the liquids with which they are moistened and the heat retained by the mass. Cataplasms may be emollient, medicated, or revulsive.

Emollient Cataplasms are usually made of bread and water, linseed meal, Indian meal, slippery elm, &c., and water; or of mashed vegetable substances, such as carrots, and marshmallows.

As warmth and moisture are the principal effects of emollient poultices, an excellent substitute for them is lint, dipped in warm water, and laid over the diseased part; the lint should be covered with a towel, in order to prevent evaporation. This mode is much neater and more cleanly than any poultice, and where merely warmth and moisture are required, it will answer all the purposes intended. A peculiar fabric, called *spongio-piline*, consisting principally of sponge, has lately been invented in England, which is said to answer very well, when properly imbued with hot water, as a substitute for the simple emollient poultice.

The addition of a little lard is useful, if the poultice does not require to be frequently changed. When the object, however, is to promote suppuration, the poultice should be frequently renewed, in which case the addition of the lard is unnecessary.

Poultices should never be heavy, nor very bulky, but they should be frequently repeated. They are useful, in all cases of inflammation which cannot be backed, to assist the suppurative process and the advancement of the matter to the surface. After an abscess is opened, the poultices should still be continued for a few days. Poultices may be used, also, as fomentations in colic, and in inflammation of the bowels. In such cases, however, as the chief object is to allay pain and to soothe, they should be made with a decoction of poppy-heads, instead of water.

Medicated Cataplasms are those in which, to the simple emollient poultice, are added other ingredients, intended to exercise an influence on the part independent of mere warmth and moisture; as, for instance, the fermented poultice, in which yeast forms a constituent; the charcoal poultice, and various others enumerated in the Formulary, all intended to fulfil certain indications.

Revulsive Cataplasms are those which, by inducing a local irritation or inflammation of the parts to which they are applied, act as derivatives. To this class belong mustard poultices, or *Sinapisms*, Cayenne pepper, garlic poultices, &c.

Mustard poultices are made by rubbing the flour of mustard, of the shops, with

water, into the consistence of a paste. When it is wished to have a weaker preparation, wheaten, rye, or linseed flour is to be added to the mustard in such proportions as are deemed sufficient. The water used should be tepid, or merely warm; a boiling temperature is injurious, as it tends to coagulate the albumen of the mustard, and thus to prevent the extrication of the volatile oil, which is the active principle. Vinegar is often used, but never increases the energy of the preparation, and, in the case of the black mustard, diminishes its power. This poultice should be spread thickly on a rag, as soon as it is prepared, and covered with gauze or tissue paper, in order to prevent adhesion to the skin. In a few minutes after it is applied, the sinapism causes a sensation of heat, which increases to almost intolerable burning. After some time, the pain subsides, and is followed by a sense of weight and throbbing of the arteries. If the sinapism be left on, however, the burning pain again returns, and becomes so great, that few persons can bear the action of a mustard cataplasm longer than three quarters of an hour.

In delicate, sensitive people, the sinapism need not remain applied above six or eight minutes, as the effect continues some time after its removal. If it remain too long on, gangrene may supervene; this cannot happen unless the invalid be insensible, as too much pain is felt to permit the lengthened application of a sinapism; but, in cases of stupor or insensibility, or in low fevers, a sinapism, if left on longer than an hour, may cause both vesication and gangrene. If the redness only is required, the pain may be moderated by mixing with the paste a drachm of tincture of opium, or two drachms of tincture of henbane, neither of which, although they allay the pain, diminishes the rubefacient activity of the sinapism.

As a substitute for the mustard poultice, where mere external irritation is required, the capsicum or red pepper will answer admirably, and it has the advantage of not vesicating: it may be used as a poultice, by mixing the powder with bread and milk, or Indian meal, or as a lotion mixed with warm spirits.

LOTIONS, LINIMENTS, EMBROCATIONS, are medicines of a more or less fluid character, which are employed as external applications. They may be used as mere fomentations to soothe pain and remove inflammation, as a mode of introducing medicinal substances into the system, or as revulsive agents, according to their nature and composition. Liniments, which generally contain oily constituents, are usually aided by friction of the part to which they are applied, either with the hand alone, or with some article capable of exciting some irritation of the skin, as a piece of flannel or a soft flesh-brush. In many cases, in which liniments are found beneficial, the advantage obtained from them is attributable rather to the friction than to any medicinal power in the lotion itself.

Frictions, whether simple or conjoined with liniments, are frequently important aid to medical treatment. Simple friction is performed by the hand alone, or with a piece of flannel, a hair glove, or a flesh-brush. If it be properly performed—namely, by short, brisk strokes with the tips of the fingers, and with great celerity, when the naked hand is the agent; and if it be continued for an hour or upwards, and repeated several times a-day—its influence in reducing swelled glands and swellings of the joints, as well as in alleviating rheumatic pains, is very great; but, besides being well performed, the friction should be continued for half an hour, in order to render it useful.

Friction, when performed in a gentle, slow, and equable manner, by producing a continued repetition of an agreeable impression on the nervous system, acts as a soother of pain, independent of any aid from opiate liniments, and also induces sleep. The beneficial influence depends, in a great measure, on its transferring the attention of the sufferer from the seat of pain, and confining it to the mild and agreeable sensation impressed by the friction. In performing it, the operator should sit by the side of the bed, and, introducing the hand under the bed-clothes, rub the legs or the arms gently, with equally lengthened, but slow movements. When the invalid is a child, its influence is more powerful when aided by a monotonous, but a

soft tune, which, although it operates upon a distinct sense, yet, by combination, renders the friction more soporific.

When friction is intended to stimulate the surface, it is necessary to aid its influence by liniments containing ammonia, or camphor, or some other excitant or acrid substance; but these should be ordered by the medical attendant. In applying them, the same method should be adopted as when simple friction is employed; but the hand of the operator must be guarded by a glove from the influence of the acrimony of the liniment.

Friction is also used to introduce various substances—namely, mercurials, iodine, opium, &c.,—into the body through the medium of the skin. The rubbing, in such cases, ought to be brisk, and applied with sufficient force to abrade slightly the cuticle; and continued until the substance, which is usually in the form of an ointment, disappears, owing to the absorption by the skin induced by the friction. In using such frictions, the hand of the operator should be guarded by a glove; otherwise, he is likely to suffer salivation, or some other unpleasant result, from the introduction of the medicinal agent into his own system.

VESICATORIES, or BLISTERS, have already been noticed in the introduction (p. 69), and do not require to be further commented upon.

ISSUES are small ulcers artificially established for the purpose of procuring and keeping up a discharge of pus. They form an important derivative remedy in many diseases, and are to be formed by the following methods: the simplest is by the application of a small blister, which is to be dressed with some epispastic or irritating ointment. It is, however, difficult to prevent the sore from healing, and the discharge is seldom sufficiently copious. They are also formed by making incisions through the skin, and inserting issue peas into the wound. The best and most effectual plan is by the use of caustic; this is done by means of a piece of leather, spread with adhesive plaster, through which a hole is made of the size of the intended issue; this is to be warmed and applied to the skin, so as to apply closely. The aperture is then to be filled with the caustic, in the form of a paste, and covered with another piece of leather, spread as above. These are to be left on until the skin is destroyed, and then to be removed and a poultice applied until the eschar separates. The late Dr. Wistar was in the habit of recommending the application of a small blister, and of rubbing the vesicated surface with caustic potassa for a few minutes. This is an expeditious and excellent method, as it is effectual and gives but little pain. The issue is kept open either by the introduction of peas, or a piece of lint smeared with the ointment of savine, into the wound. The peas operate by mechanical irritation; and, as they swell from the action of the heat and the moisture of the ulcer, they tend also to enlarge the wound. The peas should be changed at least once in twenty-four hours. Instead of the peas, it has been advised to use a small block of hard wood, with the lower surface cut into lozenges; these, pressing on the surface of the sore, occasion a copious discharge, and it will be found more convenient of application than peas. The very irritation, however, by which the discharge is maintained, is apt to cause fungous growths from the bottom of the wound, which should be kept down; this is to be done by dressing the issue occasionally with some escharotic, as burnt alum or red precipitate. Issues should be washed twice a-day with warm water; otherwise, they are apt to become very offensive.

SETONS are wounds kept in a suppurating state by means of some foreign body, which prevents their healing. They are made by passing a seton needle, armed with a skein of silk, thread, or a slip of gum elastic, through a folded portion of the skin, withdrawing the needle and leaving the silk in the wound. If a seton needle cannot be procured, the puncture may be made with a thumb-lancet or bistoury, and the silk passed through the wound by means of an eyed probe. The edges of the wound are apt, however, after it has been discharging for some length of time, to become

zallous, and the discharge to cease. When this occurs, the irritating substance should be smeared with some stimulating ointment. It should be recollected that much mischief may result from suddenly suppressing or removing either an issue or a seton. If this takes place, purgatives should be administered, and the discharge of the issue or the seton checked as gradually as possible.

OINTMENTS are simple or compound fatty bodies of a soft consistence, which are applied to the skin by inunction. They are more consistent than liniments, but softer than cerates. Simple ointment consists of one part of wax and four of lard, melted together and thoroughly mixed. This preparation is the basis of many compound ointments. Compound ointments are of three kinds: one consisting of simple ointment, combined mechanically with various substances; the second, formed by dissolving the active ingredients in melted ointment or lard; the third, made by uniting substances to the fatty matters that induce a chemical action between them.

Compound ointments of the first class are mixtures of the fatty basis with various solid and liquid substances; these are incorporated by trituration in a mortar, or on a stone slab. When hard substances enter into the combination, they require to be previously reduced to a fine powder, or, if they are soluble, to be rubbed into a paste with an appropriate menstruum, and then admixed. When extracts form constituents of an ointment, if they are not of a soft consistence, they should be rendered so by trituration with water before they are mixed with the fatty body. The mixture of two or more ointments may be effected by rubbing them together on a stone slab by means of a spatula.

Compound ointments of the second kind are made by boiling recent vegetable substances in the fatty basis, until all the water of vegetation is driven off; the heat should then be reduced to about that of boiling water, when the ointment is to be strained. Those of the third kind, as citrine ointment, are prepared by adding the constituents gradually together, aided by heat and constantly stirring.

Ointments are preserved by keeping them in glazed jars, with the surface covered with tin foil. According to M. Deschamps, the admixture of a twenty-fifth part of benzoin with the fatty matter used to make the ointment, will prevent or greatly retard the process of decomposition (see p. 146). In the case of simple ointment, and those for highly irritable surfaces, this plan answers well; where it is inexpedient, the ointments should be renewed at short intervals.


CERATES consist of a basis of wax and a fatty matter, with which other substances are incorporated. They are of such a consistence as to be softened, but not melted, by the warmth of the body. The general directions for making them are similar to those for ointments.

Many preparations of the present class are made by simply mixing or triturating their component parts together. But when resins, wax, spermaceti, or concrete oils are to be mixed either together or with fluid oils, it is better to unite them with the aid of heat, which not only liquefies the solids, but likewise renders them much more soluble in the liquids. Brisk agitation is required while the mixture concretes on cooling; otherwise, the solid ingredients will separate either mechanically or by crystallization. Even in purifying lard or suet by fusion and filtration, it is right to stir the mass as it cools; otherwise, the stearin or solid oil is apt to separate in part from the elain or liquid oil, so that an irregular mixture of the two principles is obtained. A gentle heat is sufficient for making ointments, cerates, and the like, because most of the ingredients are easily fusible; and a strong heat must be avoided, for in that case acrid acids are engendered, which may alter the properties of the preparation. Hence the vapor-bath is generally used for applying heat; and it ought always to be employed in operations on a large scale.

When the cerate is intended to irritate and not to soothe, a high temperature may be employed, which, in some cases, is advantageous, as in the cerate of cantharides, by more effectually dissolving out the active principle.

PLASTERS are more consistent than cerates, adhesive at the temperature of the body, and requiring the aid of heat to soften them sufficiently to be spread. They are of two kinds; one consisting of a compound of olive oil and litharge, or an oleo-margarate of lead, having the common lead plaster as a basis, united to resinous or other substances; the other, composed wholly of resinous matters, or these mixed with fatty substances or wax.

FIG. 34.



APPARATUS FOR SPREADING PLASTERS.

[Fig. 34 represents a convenient apparatus for spreading plasters. *a c e* is a solid block of wood, rather larger than the intended plaster, and having the upper surface *c* made slightly convex; *b d* is a tinned sheet-iron lid to cover the block, to which it is fastened by the hinge *r*, and farther secured by the clasps seen at *d*; *n* is an oblong opening in the cover of the exact size of the plaster; *m* is a frame for marking the leather into squares. The leather, thus prepared, is placed on the convex surface *c*, the lid *b d* is turned down and fastened by the clasps, as seen in the figure *i h*. The plaster melted in the pan *k* is then poured on the leather *i*, and, by means of the iron instrument *g*, is spread uniformly over its surface. When cool, it is separated from the frame by passing the sharp-pointed instrument *l* around its inner margins.]

In making plasters, it is necessary, in most cases, to operate at a temperature that will not volatilize or carbonize the ingredients; hence, a water-bath or steam heat is preferred. In the case of the burnt plaster (*onguent de la mere*) of the Paris Codex, sufficient heat is employed to cause the fats to fume. Plasters should be made in metallic vessels, much larger than will contain the substances to be operated upon, as the mass increases in bulk by the action of heat causing an extrication of vapors. The water ordered in making lead plaster should be added at the commencement of the process; otherwise, when poured on the melted mass, it may cause a projection of portions of the heated materials on the operator. The ingredients should be thoroughly stirred together during the process of melting. When this operation is concluded, the plaster is to be removed, in small portions at a time, into a vessel constantly supplied with cold water; and, when sufficiently consistent, rolled into cylinders on a wet marble, and again placed in cold water to harden. To render plasters of a lighter color, they are kneaded and pulled under water; but this process should not be practiced on compound plasters containing substances soluble in water; nor should these plasters be cooled in water, but suffered to chill on a marble slab until fit to be made into rolls. Some of the officinal plasters belong to the

second class, and are made by melting resinous matters with the addition of certain oleaginous substances.

Plasters are preserved by enveloping the rolls with paper, to exclude the air as much as possible. When kept for any length of time, they are apt to become hard and brittle, and to lose their color. When this is the case, they should be re-melted by a gentle heat, and sufficient oil added to restore their proper consistence.

Spreading plasters for use, requires skill and address on the part of the operator. They are spread on various textures, but most generally on linen or cotton cloth, or leather. The shape and size must be regulated by the part to which they are to be applied. The plaster is to be spread on the leather or cloth by means of a heated spatula of a peculiar form; this is brought in contact with the end of a roll, which, becoming liquefied, can then be evenly diffused over the surface. Where the plaster is sufficiently adhesive to maintain its place when applied to the body, it is to be spread so as to leave about half an inch of margin uncovered; but where it does not possess this quality in sufficient degree, it should be surrounded with an adhesive margin, which is to be prepared before spreading the body of the plaster. Full and particular directions for spreading plasters will be found in Procter's edition of *Mohr and Redwood*, p. 516, *et seq.*, and to which we would refer those who wish for clear and accurate directions on the subject.

FUMIGATIONS are extrications of vapors or gases, designed to modify the air of a sick chamber, and to mask any unpleasant odors that may be present, or to produce a medicinal effect on those parts of the body with which they are brought in contact.

Fumigations, for the purpose of obviating or masking unpleasant odors in a sick room, should never be employed to supersede ventilation and cleanliness; for most of them, instead of purifying the air, only render it more unfit for respiration. They are generally made by burning pastilles, sugar, juniper berries, benzoin, &c., so as to create an odoriferous smoke. As disinfecting agents, they are utterly useless; and are relics of an ancient custom of burning frankincense, and other odorous substances in vitiated air, to overcome the feter which is more or less present. They disguise unpleasant odors; but they accomplish nothing more. The infection remains not only unaltered by the diffusion of the most powerful aromatic vapors, but its deleterious properties are sometimes augmented by them.

The fumes of burning camphor, and the vapors arising from its tincture, have more pretensions than either of the above to the name of a useful article of fumigation. It is much employed and confided in on the Continent; but the experience both of American and British physicians does not warrant the opinion that it possesses any power of destroying infection or contagion. Nothing is more ridiculous than the custom, which was at one time very general, and which is still continued to a certain extent—namely, that of carrying a camphor-bag about the person, as a protection against infectious diseases.

Vinegar is, not without reason, regarded as possessing some chemical influence in decomposing infectious and contagious matters; and, consequently, it is almost invariably sprinkled over the floor of the rooms of those suffering under infectious diseases; or the vapor of hot vinegar is diffused through their apartments. It is thought to be still more salubrious, and a more powerful disinfectant, when it holds camphor or aromatic oils in solution; hence the great popularity of the preparations called *Aromatic Vinegar* (see p. 78) and *Thieves' Vinegar* (see p. 79). The repute of the latter is founded upon a story, that four thieves, who plundered the dead bodies during the plague at Marseilles, with perfect security, on being questioned respecting the cause of this impunity, confessed, on the condition of their lives being spared, that they attributed it solely to the use of *Aromatic Vinegar*.

Vinegar, in this state of combination, is extremely agreeable and refreshing, both to the invalid and the attendants of the sick room. The benefit which it produces depends upon a certain degree of stimulus imparted to the sensitive nerves, which are generally in a low condition in an infectious atmosphere; but, as a chemical

agent, its powers are too feeble to be followed by much benefit. It is, however, always refreshing, and is much better adapted for overpowering the unpleasant odors of the sick room than any of the former substances.

The most efficacious fumigations that can be employed, are those of chlorine. This disinfecting gas is extricated from a mixture of common salt, black oxide of manganese, and sulphuric acid (see p. 182). Instead of chlorine disengaged in this mode, it is more common to use the chloride of lime, the chloride of soda, or the chloride of zinc; solutions of either of these, placed in shallow vessels in a room, will give off chlorine in a gradual manner, by a decomposition of their constituents. Some difference of opinion exists as to the mode in which this gas acts. Dr. A. T. Thomson is of opinion that it operates by decomposing the infected air:—

“With regard to the mode in which chlorine operates, the most probable opinion is, that it decomposes the infectious matter, which is a compound. One principle of this compound seems to be hydrogen; another, ammonia; and a third, a fetid volatile oil; besides the unknown matter of infection. The chlorine unites with the hydrogen, and forms hydrochloric acid, which combines with the second component of the infectious medium—namely, ammonia—and forms sal ammoniac; the withdrawing of both of which from the air causes the precipitation of the volatile oil; whilst the chlorine also neutralizes the real matter of infection, whatever it may be; and thus the vitiated atmosphere is purified.”

Although chlorine is capable of destroying fetid and unpleasant odors in the sick room, it is liable to many objections. However extricated, it is apt to create irritation of the respiratory organs, and coughing, when respired; it corrodes all metallic substances, and must, therefore, be used with caution; from the chlorides evolving the gas more slowly, they are, in most cases, preferable to the immediate extrication of chlorine from salt and manganese.

Other acid fumigations have also been found useful as disinfectants; as those of nitrous acid, so highly recommended by Smith and Carmichael, and considered by Christison even as superior to chlorine. Sulphurous acid fumes, produced by burning sulphur, have likewise been recommended. Both are capable of destroying unpleasant odors and emanations; but the same objections apply against their use in a sick chamber—that of causing much irritation of the organs of respiration; hence they are unfitted to cases where there is any disease or weakness of these parts.

Fumigations, used as medicinal agents, are much more prescribed on the Continent of Europe than in Great Britain or the United States. They are sometimes employed as inhalations—as emollient vapors in the treatment of bronchial affections; and balsamic vapors in affections of the chest. They are also used externally, as the fumes of benzoin in rheumatic and gouty complaints; those of aloes in deficient menstruation; those of sulphur and mercury in various diseases of the skin.

The effects of fumigations vary according to the temperature at which the vapors are used; when it is high, and the whole body of the patient is exposed to it, it produces much the same train of symptoms as those caused by a vapor-bath. When the fumigation is made with substances which give off no watery vapors, the phenomena that result are analogous to those of a heated air-bath, but more marked, especially those which occur in the skin and subcutaneous tissue.

BLOOD-LETTING.

Various methods are practiced for this purpose. General bleeding is performed by opening a vein, called venesection; or by puncturing an artery, which has received the name of arteriotomy. Topical blood-letting is practiced by opening the small vessels of a part by means of slight incisions, or what is called scarification, by the application of leeches, and by cupping.

GENERAL BLOOD-LETTING. — *Venesection* is one of the most frequently employed of the different modes of detracting blood. It is usually practiced on one of the veins of the arm. A band or ligature is to be applied around the arm above the point at which the vein is to be opened, so as to obstruct the passage of the blood to the heart; this causes an enlargement and turgidity of the vessel below the ligature. This bandage should never be applied so tightly as to prevent the passage of blood in the arteries of the part. The vein is to be opened by means of a common or thumb lancet, or a spring lancet. In using the first, the vein is to be kept from rolling by means of the thumb of the left hand placed a small distance below where the puncture is to be made; but care should be taken not to alter the relative position of the skin and vein, because, in resuming their position after the lancet is inserted, the aperture through the skin will no longer correspond with that in the vein, in consequence of which a thrombus or ecchymosis will form—a very common accident in venesection. The lancet is to be pushed into the vein, and when its point is within the cavity of the vessel, it is to be carried forward a little, so as to enlarge sufficiently the opening. The arm is to be kept extended after the operation until the necessary quantity of blood is taken. If the blood should not flow as freely as is desired, the patient should exert the muscles of the arm by grasping a stick, or moving the fingers.

The spring-lancet is much used in some parts of the United States, and is almost universally employed in Philadelphia. This has arisen, in great measure, from its being preferred to the common lancet by Dr. Physick, as well as by Dr. Dorsey, whose reasons for this preference are thus given in his "Elements of Surgery."

"In a country, situated like the United States, where every surgeon, except those residing in the large cities, is compelled to be his own cutler, at least so far as to keep his instruments in order, the spring-lancet has a decided preference over the lancet; the blade of this can with great ease be sharpened by any man of common dexterity, and if not very keen, it does no mischief, whereas a dull lancet is a most dangerous instrument; and no one can calculate with certainty the depth to which it will enter: to sharpen a lancet, is regarded by the cutler as one of his nicest and most difficult jobs; it is one to which few surgeons are competent.

"The *safety* of using the fleam is demonstrated by daily experience; there is no country in which venesection is more frequently performed than in the United States, and, *perhaps, none where fewer accidents from the operation* have occurred; of these few, I beg leave to state that all the aneurisms produced by bleeding, which I have seen, have been in cases where the lancet was used. I have since, however, met with an exception to this statement. I have seen the brachial artery opened by a spring-lancet, but it was by an old barber, half blind, and very clumsy."

"The manner of using the spring-lancet differs in nothing from the operation with the common lancet, excepting that the surgeon must place the instrument in such a situation over the vein that, when the spring is touched, the orifice into the vein will have a proper size and direction. Dexterity in this is very readily and speedily acquired. In point of *facility* in its use, it has a great advantage over the lancet.

"Among the advantages of the spring-lancet, *economy* is not the least. A country practitioner who is constantly employing the English lancets, and who is particular in using none but the best, must necessarily consume half the emolument derived from the operation in the purchase of his instruments. One spring-lancet, with an occasional new blade, will serve him all his life."

After a sufficient quantity of blood has been drawn, the flow is to be stopped by removing the ligature and placing a finger on the orifice, when any blood that has soiled the arm is to be wiped off, and the edges of the orifice carefully brought together, that they may unite by the first intention. To accomplish this, let the bleeder, with the thumb of that hand which holds the arm, push the skin towards the orifice, while he draws it on the other side to the same point with the compress; thus the skin will be thrown into folds at the wound, over which he is immediately

to apply the compress, which should be broad, to keep the skin better together, and thick, to make the compression more certain.

When blood-letting is ordered, everything required for the operation should be at hand. The ligature for tying the arm, so as to obstruct the flow of blood in the veins below it, should be a soft old ribbon; and the same may be employed for securing the pledget over the orifice, when the flow of the blood is to be stopped. The pledget, or compress, should be a piece of linen, folded into a square form, about an inch in diameter, and comprehending three or four thicknesses of the material. It is always proper to use a graduated basin for receiving the blood, that the exact quantity drawn may be ascertained. If the invalid be bled in bed, he should sit erect; if he be up, he should be placed erect in a chair. When the operator is dexterous, there is little or no necessity for guarding either the bed or the clothes of the patient from the blood; as the pressure of the thumb of the operator, placed below the point where the vein is to be punctured, should not be relaxed until the utensil for receiving the blood be conveniently placed for that purpose. A basin with a little tepid water, and a clean, soft towel, should be ready for washing and drying the arm, before the compresses and the bandage for preventing the further escape of the blood be applied.

When a nurse or an attendant is not fully instructed in her duty, there is always considerable anxiety displayed to provide smelling salts and other means to obviate fainting. But, in many cases, much of the benefit of blood-letting depends on the fainting which it causes; and the propriety of checking that effect should be left entirely to the physician. No person should be present at this operation, nor should any one attempt to hold the basin for receiving the blood who is liable to become sick or faint at the sight of blood.

The arm should not be used for some hours after a vein has been opened in it; but, if everything has gone on well, the bandage may be removed at the end of two days. When this has been applied too tightly, the forearm and hand are apt to swell and become painful, in which case the bandage is to be loosened. If, on removing the bandage, it is found that the orifice, instead of being closed, presents an inflamed appearance, with its lips swelled and red, the arm is to be kept perfectly quiet, the edges of the wound brought in contact, and some cool application made to the part, to relieve the inflammation. In those cases where the inflammation extends beyond the orifice, and assumes an erysipelatous appearance, with a hard and painful tumor around the wound, a compress is to be applied upon the vein at the inflamed part, to make the two sides adhere together; if they do not adhere, simple contact will still be sufficient to prevent suppuration in this part; or, if inflammation has gone so far as to make the surgeon suspect that suppuration has taken place, then the compress must be put upon that part of the vein just above the suppuration.

The remedy for inflammation of the veins consequent on venesection, which answers best, is the application of a blister over the inflamed part. As soon as the inflammation commences, a small plaster of simple cerate, spread on linen, should be applied on the orifice, and over this, a blister large enough to cover the whole inflamed part, extending three or four inches from the orifice in every direction. This remedy was first proposed, and made use of, by Dr. Physick, and has been employed by many physicians with great success, even after extensive inflammation of the vein has taken place.

Sometimes it becomes expedient to open the external jugular vein. To do this, the head should be laid on one side, and the vein compressed by the operator's thumb near the clavicle, and the opening made in that part of the vein which lies over the sterno-cleido-mastoid muscle.* The blood soon ceases to escape after the pressure is removed, but a small strip of adhesive plaster is necessary.

[* In bleeding from the jugular vein, it is generally advisable to compress both of the external jugulars. This is easily done by placing the thumb of the left hand upon one jugular vein, and the forefinger of the same hand upon the other. If this point is not attended to, the operation may fail, in consequence of the free anastomosis across the neck, and the passage of the blood downwards upon the opposite side.]

In children, especially where they are very fat, a vein in the leg or foot can often be found, especially about or above the malleolus, when none are perceptible in the arm. To promote their distension, and to increase the bleeding when they are opened, it is found useful to immerse the limb in warm water.

In bleeding young children from the arm, when they are restless and fretful, it is a good plan to attach the child's arm to the left forearm of the operator, by means of ligatures at the wrist and centre of the arm; by this means, complete control is obtained over the motion of the child's arm, and the venesection can be accomplished with comparative ease.

ARTERIOTOMY is the opening of an artery to detract blood. It is much more limited than venesection in its application, being practiced only upon the temporal and posterior auricular arteries; and, in most cases, is restricted to the former only. In this, the anterior branch is preferred, where it passes above the exterior angle of the eyebrow. In performing the operation, the vessel is to be partially divided transversely, and not longitudinally. When as much blood has been drawn as is deemed necessary, the incision is to be deepened so as to divide the artery entirely. On the consequent retraction of the cut extremities, the bleeding will generally cease; if it should not, a graduated compress, maintained in place by a bandage around the head, is to be applied.

TOPICAL BLOOD-LETTING.—This consists of the abstraction of blood by cupping, leeching, or scarification.

Cupping is performed in the following manner: The skin being softened by means of a sponge and warm water, a small bell-like glass, known as a cupping-glass, having the air contained in it rarefied by being passed over the flame of a lamp, or by other means, is immediately applied to the part, all hair and other extraneous substances being previously removed: from the formation of a vacuum beneath the cup, the pressure of the air on the surrounding surface causes that portion included in the cup to swell, and the vessels to become turgid. When this has taken place, the cup is removed and several incisions made by means of a scarificator, an instrument containing numerous lancets, which, by means of a spring, can make a corresponding number of incisions at the same moment; the depth of these incisions can be regulated by means of a screw which protrudes or withdraws the lancet, according to the vascularity of the part, the quantity of blood to be drawn, &c. When a sufficient quantity of blood has collected in the cup, it is to be removed by introducing the nail of one of the fingers under the edge, by which means, air being allowed to enter, the cup becomes detached. The part being washed with warm water to remove any clots of blood, the cup is again to be applied as above, and the operation continued till a sufficient quantity of blood is obtained. Sometimes, especially when applied to the scalp, the cups fill so rapidly with blood, as to become detached almost immediately on being applied.

For the operation of cupping, a basin of hot water, sponges, and clean soft towels are to be provided. It is generally considered as a severe and painful operation; but this is not the case, if the operator understands his business. This is readily ascertained by observing the manner in which the cups are applied, and the rapidity with which they are filled. A good cupper does not exhaust much of the air in the cup before applying it, but simply passes its mouth rapidly over the flame of a lamp; for, when it is held over the flame even for a few seconds, the compression of the edge of the cup upon the skin is so great, that it operates as a ligature, and checks the flow of the blood to the scarified part; hence very little blood is procured. A good cupper, also, removes the cups without spilling the blood which they contain; and the whole operation is completed in a short time.

In our large towns, it is but seldom that physicians, nurses, or those attending in a sick room, are called upon to perform the operation of cupping, as professional cuppers are readily to be obtained; but in small towns or villages, this is not the case, and the duty devolves either upon the physician or on those acting as nurses; and

hence some directions are necessary to those unaccustomed to this task. Few are found sufficiently expert to exhaust the air in the cup by means of the lamp; and even among the professional cuppers in this city this mode is very rare, though, when properly performed, it is by far the best. The usual plan is to rarify the air in the cup by means of a small cone of paper, dipped in spirits of wine, or strong brandy; this is ignited and thrown into the cup, which is instantly to be applied to the intended spot. Where the proper cupping-glasses and scarificator are not to be had, wine-glasses or very small tumblers may be substituted for the first, and small incisions by means of a thumb-lancet will answer the purpose of the latter. A very convenient apparatus may be obtained at most of the surgical-instrument makers, consisting of cups and an exhausting syringe fitted to them, by which even the most inexperienced nurse can perform the operation of cupping.

The cicatrices of the scarifications leave a permanent mark; on which account, females should not be cupped upon the nape of the neck; but when blood is to be drawn from the head, the glasses should be applied behind the ears, and a portion of hair removed in such a manner that it may be covered by what remains.

When cupping-glasses are applied without any scarification being practiced, the operation is called *dry cupping*, and is much used to cause a speedy irritation of the skin for the relief of oppression of breathing, local pains, &c. To obtain the full benefit from this operation, the cups should be suffered to remain upon the part until they cause an exudation of a small quantity of serum, or great irritation of the part. The application of dry cups has been found extremely beneficial in poisoned wounds; they act not only by abstracting the poison, but also, by the pressure they exercise on and around the part, in preventing the absorption of it. The experiments of Dr. Barry in England, and of Drs. Pennock and Rodrigue in this country, show unequivocally the value of this plan of treatment.

Leeching consists in the application of leeches to any vascular part of the body. These are annulated animals, having an extensile, soft body, terminated at each extremity with a muscular disk, the anterior of which is furnished below with a mouth having three crescentic jaws, armed with numerous sharp teeth. To draw blood, the leech first renders the part tense by means of the anterior disk, which acts like a cupping-glass, and then makes a triangular wound by a saw-like motion of its tri-radiate jaws; it then draws the blood by suction, until it is completely gorged, when it detaches itself and falls off.

Leeches are seldom properly applied or managed. The part to which they are intended to be applied should be washed with a little soap and warm water, then with simple cold water, and, lastly, it should be well dried. If the part be hot and inflamed, the leeches should be put, for a few minutes, into tepid water; and this should be done, also, when they are to be applied in the mouth, or to any part of the body warmer than the general surface; but, at all times, before they are applied, they should be dried between the folds of a clean soft towel. The easiest and best mode of applying them is, first, to place the number to be used in a hollow made with the points of the fingers in a towel folded like a napkin; then, so to turn the towel and the leeches upon the part where it is intended they should fix, that the towel will cover them. The hand must be kept over the towel, to prevent their escape, until they all bite, which usually happens in a few minutes; after which, the towel may be removed. By this method, twenty or thirty leeches can be applied more rapidly and with less trouble than two, when each leech is separately applied. Another plan is to put the leeches into a pill-box or small glass, and apply this over the part to which it is designed that they should attach themselves. If this plan, however, cannot be pursued, owing to the nature of the part to which they are to be applied—as, for example, the inner or outer angle of the eye—then the simplest method is to scratch the skin with the point of a needle, and to apply the leech to the spot moistened with blood. When they are to be applied within the mouth, or any open cavity, each leech should be put into a large quill, with its head towards the open end of the quill, which should be applied to the part, and retained upon it until the leech is fixed, when the quill may be gently withdrawn; but a thread

should be tied round the tail of the leech when it is to be applied within the mouth, to prevent it from being swallowed—an accident which has occasionally happened, and has been productive of serious consequences.

Leeches should never be forcibly detached, as their teeth are apt to separate, and, being left in the wound, to cause an erysipelatous inflammation of the part. They should be permitted to drop spontaneously, which being the result of a temporary suffocation (*asphyxia*), all muscular energy ceases in the animal, and, the teeth shrinking, it drops off entire. A bread-and-water poultice, not too hot, should then be laid over the bites, to encourage the bleeding. The invalid should be kept warm in bed, when it is necessary to abstract a large quantity of blood. In general, the bites soon cease to bleed; but, in some instances, a copious flow takes place; and therefore, to prevent exhaustion, the poultice should be frequently examined. This exhaustion is more likely to occur in children than in adults; and, for the same reason, leeches should not be applied upon children late in the evening, unless they are very urgently required. It is also proper, in young patients, to select for their application a part which admits of pressure; for example, the thorax and temples, where the bones are covered with a thin layer of soft parts. Sometimes the hemorrhage continues so as to become alarming; in such cases, where pressure is unavailing or cannot be practiced, creasote or some of the styptics applied to the bites will be found useful. Powdering the spot with rye flour has also proved effectual. When these means fail, cauterizing the bites by means of a fine point of nitrate of silver, or with a red-hot probe or knitting-needle generally proves successful.

Scarifications are slight incisions made in a vascular part, in order to relieve the engorged capillaries; they are practiced in inflammations of the conjunctiva, and in those of the tonsils, with much benefit in many cases. The operation is very simple, consisting merely in dividing the surface of the inflamed parts to a slight depth with the point of a lancet or scalpel, and encouraging the bleeding by means of warm water.

POISONS.

POISONS are usually defined to be substances of an animal, vegetable, or mineral nature, which, when administered in a small quantity, are capable of producing deleterious effects on the animal economy, and, in some instances, of causing the same consequences when applied to the surface of the body. But this definition is not strictly accurate, and is not applicable to all cases of poisoning; nor is it, perhaps, possible to give in a few words such an one as would include all poisonous agents. Poisons are usually arranged according to their action on the system; but, as the alphabetical plan has been pursued in the main body of this work, it will be continued in the present instance. For full information on the subject, the reader is referred to the work of A. S. Taylor, "On Poisons in relation to Medical Jurisprudence and Medicine."

[GENERAL ANTIDOTE for poisoning, in which the nature of the poison is unknown:—

R. Calcined magnesia, Pulverized charcoal, Sesqui-oxide of iron,	} equal parts, in a sufficient quantity of water.
--	---

This preparation is perfectly innocent, and is very likely to be efficacious, for its ingredients, though simple, are antidotes to the most common and active poisons. *Bull. de Thèrap.*]

ACID, ACETIC.

Symptoms.—Great heat and burning pain in the stomach, convulsions, death.

Morbid Appearances.—Mouth and fauces brownish; lingual papillæ enlarged; œsophagus lined with a brownish adventitious membrane; stomach livid and even blackened; vessels much injected.

Antidotes.—Magnesia or its carbonates mixed into a cream with water; soap, and the alkalies.

ACID ARSENIOS, or ARSENIC. A metallic, corrosive poison.

Symptoms.—A metallic, austere taste; a great flow of saliva; nausea and vomiting; fainting, great thirst; a sensation of heat in the stomach, which, in many cases, rejects the mildest fluids; much griping, tenesmus, and purging; the stools being dark-colored and very offensive; the urine scanty and high-colored; the pulse small, frequent, and often intermittent; distressing palpitation of the heart, with labored respiration and cold sweats; prostration of strength; sometimes paralysis of the extremities, delirium, convulsions, &c.

Morbid Appearances generally confined to the stomach and intestines; in the stomach, in the form of intense inflammation, but not of erosion or abrasion; the inflammation is also evident in the upper intestines, but slight in the colon, though often violent in the rectum. The morbid appearances of other organs various. Sometimes no morbid phenomena to be found.

Tests.—In the solid state: it is entirely sublimed by heat. If mixed with charcoal, and heated in a suitable test-tube, deoxidated arsenic will be obtained in form of a metallic coating inside the tube; and this may be re-converted into arsenious acid by urging it in various directions along the tube with the aid of a minute spirit.

lamp flame; the facets of the crystals thus formed (on the cooler situations) will be seen in some places with the unassisted eye, but more distinctly by means of a four-power lens.

In solution: ammoniacal nitrate of silver produces a lemon-colored (*arsenite of silver*) precipitate. Ammoniacal sulphate of copper throws down a fresh, grass-green (*arsenite of copper*) precipitate. Transmission of sulphuretted hydrogen produces a bright yellow (*sesquisulphuret of arsenic*) precipitate. Lime-water precipitates a white (*arsenite of lime*) powder; but this test is not one to be relied upon.

Placed with zinc, and diluted sulphuric acid, in Marsh's apparatus, the arseniuretted hydrogen thence arising, when lighted, will deposit metallic arsenic on a piece of glass held within the flame.

[*Reinsch's Test.*—Acidulate the suspected liquid with muriatic acid, and boil copper wire or foil in it for ten minutes. The arsenic is deposited on the copper as a white alloy, from which it can be separated as arsenious acid, by subjecting the copper, cut into shreds, to a low red heat in the bottom of a small glass tube.]

The precipitates referred to, if washed and dried, may be treated as directed above for arsenic in the dry or solid condition.

In testing suspected matters obtained from the stomach, these (and, in cases of death, the stomach itself) must be cut or broken down, and boiled during, at least, three-quarters of an hour: if not sufficiently fluid, add distilled water. Strain, and with addition of a small quantity of potassa, again boil during a quarter of an hour, and filter. If this liquor manifests either alkaline or acid reaction, neutralize with potassa, or with acetic acid, as may be required; then acidulate it faintly with hydrochloric acid. Solution of nitrate of silver will remove the acid; and solution of muriate of soda will precipitate any excess of silver. The liquor may now be experimented on with the above tests.

Treatment.—Abundant draughts of sweet milk, gruel, decoctions of starch, or oily mixtures; tickling the fauces, &c., to induce vomiting; the stomach-pump; emetics of sulphate of zinc. Hydrated sesquioxide of iron (newly prepared, *v. process*, p. 238,) in large doses is the only antidote of reliance.

[Light magnesia may be used with advantage, until the hydrated peroxide of iron can be procured.]

Afterwards combat any inflammatory symptoms by the usual means, and let the patient subsist, for a long time, wholly on the blandest diet.

All arsenical poisons have much the same action, and similar means are to be used for their detection and for counteracting their effects.

ACID, BORACIC.—This is said to be very virulent; but no instances have been recorded of its fatal effects on man.

Tests.—Not very soluble in water; soluble in alcohol, which, when inflamed, burns with a rich green color; soluble in the caustic alkalies.

ACID, CARBONIC.—This gas is freely liberated in respiration, combustion, and fermentation, as also in the calcination of lime; and is found in coal mines, wells, and cellars.

Symptoms.—Great drowsiness, giddiness, difficulty of respiration, loss of muscular power and sensibility, and coma. The whole body, but especially the face, appears swollen and livid; eyes are unusually prominent, and retain their brilliancy for some time.

Tests.—It extinguishes a taper if the proportion be above twelve or fifteen per cent.; lime-water, or a solution of subacetate of lead, is precipitated white by it.

Treatment.—Remove the person into the open air, and elevate the head; dash cold water over him, apply stimulating frictions to the thorax and extremities, and use artificial inflation of the lungs; as soon as the patient can swallow, stimulants may be cautiously administered; but, if there are signs of oppression of the brain, trepansection is to be performed.

ACID, CITRIC.—In large doses, acts as an irritant poison; but no fatal case in the human subject has been recorded.

ACID, HYDROCYANIC or PRUSSIC.—An extremely active, sedative poison.

Symptoms.—When the dose is large, almost immediate death ensues; in smaller quantities, it causes pain in the head, stupor, nausea, faintness, vertigo, and loss of sight; followed by difficulty of respiration, dilated pupils, a small vibrating pulse, and syncope, which will end in death, if curative means are not employed.

Morbid Appearances.—None; but a strong odor of the acid is exhaled from the stomach.

Tests.—The (bitter almond) peculiarity of its odor. When a little potassa is mixed with liquids containing this poison, and solution of the sulphate, or sesquichloride of iron added, a greyish-green precipitate is thrown down—which deepens to a Prussian blue tint, on addition of a few drops of sulphuric acid. The nitrate of silver produces a white (*cyanide of silver*) precipitate; which, after being washed and dried, and then held on a watch-glass over a flame, burns with a fresh rose-color, cyanogen being at the same time evolved.

[*Sulphur Test.*—Place two drops of a solution of hydrosulphate of ammonia, containing an excess of sulphur, in the centre of a watch-glass, and invert it accurately over the vessel containing the poisoned liquid. Remove the glass in three or four minutes, and dry the moistened spot gently over a spirit-lamp. Let a drop of water fall on the white film, and then a drop of the perchloride of iron. If prussic acid be present, a blood-red solution (sulpho-cyanate of iron) is produced; and this red color is discharged by the addition of one or two drops of a solution of corrosive sublimate.]

When a mixture is to be examined, containing matters from the stomach, &c., if alkaline, it must first be neutralized by addition of sulphuric acid, then one-eighth part cautiously distilled therefrom into a receiver immersed in some frigorific mixture; and the product may then be tested by nitrate of silver, &c., as above.

Treatment.—The internal remedy best calculated to act chemically on this poison, is carbonate of potassa in solution, quickly succeeded by watery solutions of sulphate of iron (with the intention of decomposing the acid, and forming the less injurious substance named *Prussian blue*); but the instances of success by these, or, indeed, by any other kinds of treatment, are extremely rare. Inhalation of chlorine gas, ammonia, &c.; artificial respiration; exhibition of energetic stimulants, such as brandy, liquid ammonia; chlorine-water, chloride of soda and of lime, have all been extolled, and may be tried. Mustard poultices to the stomach, and cowhage applied over the chest; venesection at the neck, temples, &c.; and affusion of cold water to the head and spinal region, are more likely to prove timely excitant and effective remedies. Employment of the stomach-pump, emetics, &c., seems a most improbable means of relief—or, rather, a waste of time.

ACID, MURIATIC or HYDROCHLORIC. — A corrosive mineral poison.

Symptoms.—Sensation of burning in the throat and stomach, styptic taste in the mouth, much thirst; the eyes red and sparkling; the pulse frequent and tense; the skin hot and dry; the tongue red and glazed; the lips black; vomiting of blood and yellow matter, having the peculiar smell of the acid; cold sweats, delirium, &c. Orfila also says, a thick white fume, having the penetrating odor of the acid, issues from the mouth.

Morbid Appearances.—Mouth, fauces, throat, and stomach highly inflamed, of a deep red color; mucous membrane sometimes detached or destroyed, sometimes perforated; contents of the stomach yellow or dark green.

Tests.—Characteristic acid taste. The presence of muriatic acid in small proportion, is at all times demonstrable in the juices of the stomach; therefore proof of its presence in *excess* is required in cases of poisoning. When a piece of glass, wetted with this acid, is held close to *liquor ammoniæ*, so as to let the vapors of both

meet, white (*muriate of ammonia*) fumes are formed. Solution of nitrate of silver produces a white (*chloride of silver*), curdy-looking precipitate.

Treatment.—The immediate administration of soap, magnesia, soda, or potassa, mixed in bland demulcent drinks, to be followed by the free use of emollients and mucilages. If patient survives the first effects of the poison, employ antiphlogistic means to overcome the supervening inflammation.

ACID, NITRIC, or AQUAFORTIS.—A corrosive mineral poison.

Symptoms.—Much the same as those produced by the last-mentioned poison.

Morbid Appearances.—In cases where death has occurred soon after the ingestion of the poison, the most striking appearance is a layer of yellow matter on all parts reached by the acid. The lips, chin, and hands of the person are also often stained with orange-colored spots. Perforations of the stomach are found in most cases.

Tests.—The orange-colored fumes that are given off, when it is boiled with copper filings, are characteristic. It reddens morphia, and blackens a solution of protosulphate of iron. A fluid containing it, on the addition of carbonate of potassa forms nitre, which may be known by its deflagration, or by powdering a small portion, and placing it in a glass tube with some fine copper filings, moistening with water, and adding a few drops of sulphuric acid—when, if it be nitre, there will be an evolution of the orange-colored fumes of nitrous acid.

Treatment.—The administration of magnesia, or chalk, in some bland fluid, then the free use of demulcents, and subsequently the employment of antiphlogistic means to subdue inflammation.

ACID, OXALIC.—A corrosive vegetable poison.

Symptoms.—Burning heat of the stomach, nausea; sometimes vomiting, but at others, ineffectual efforts to discharge the contents of the stomach; great prostration of strength, violent pain, spasmodic respiration, convulsions, &c. When the patient survives the first effects of the poison, all the symptoms of violent inflammation of the alimentary canal are developed. In a diluted form, it appears to cause paralysis of the heart or symptoms of cerebro-spinal disease.

Morbid Appearances.—Tongue and fauces usually covered with a viscid white coat; the stomach containing a dark-brown mucous fluid, and its substance of almost a gelatinous consistency. In some cases, no traces of the action of the poison are perceptible; but, when death does not occur for some time, the alimentary canal is found inflamed.

Tests.—In solid state: has the characteristic sour taste of most acids; and produces the same changes as they do on vegetable blues. When placed in water, a crackling sound accompanies the solution of its crystals.

In solution: Solution of chloride of lime produces a white (*oxalate of lime*) precipitate; which is insoluble in hydrochloric acid, but soluble in nitric acid. Solution of nitrate of silver also throws down a white precipitate, which, on being carefully dried by aid of heat, on bibulous paper, acquires a brownish outline, and towards the end shows some slight explosions.

Treatment.—Mixtures of magnesia, chalk, whiting, or plaster scraped from off the inside wall of an apartment, mixed with water (the amount of the latter being as small as convenient for swallowing); abundant drinks *subsequently*. No alkalies should be employed.

ACID, SULPHURIC.—A corrosive mineral poison.

Symptoms.—An austere styptic taste in the mouth; a sensation of burning heat in the throat and stomach, followed by nausea, vomiting, and much fetor of the breath. The matters vomited contain both venous and arterial blood. Signs of great inflammation of the abdominal viscera soon manifest themselves, with difficult respiration, a croupy cough, and a small, frequent, irregular pulse; great anxiety

and restlessness, convulsive motions of the muscles of the face and lips; sometimes a papulous eruption precedes death.

Morbid Appearances.—These are not always to be found, except in the fauces and larynx, which, in most cases, present evidences of the highly corrosive action of the poison. The stomach is sometimes found to contain a quantity of dark grumous matter, and is much distended with fetid gas—its membranes ulcerated, dark-colored, and having numerous corroded spots, and even perforations.

Tests.—The acid may be in a concentrated or a diluted state. If in the former, any organic matter placed in contact with it is blackened and charred; when mixed with an equal bulk of water, much heat is evolved; when boiled with copper filings or mercury, sulphurous acid gas is evolved. When the acid is in a diluted state, the best test is nitrate of baryta, which causes a dense white precipitate of sulphate of baryta; this can be verified by calcining it for some minutes with an equal weight of charcoal, wrapped in platina foil, then introducing the residue into a glass tube and adding a few drops of muriatic acid. This will cause an extrication of sulphuretted hydrogen, which can be recognized by its odor, and by its blackening carbonate or acetate of lead.

Treatment.—The same as for the other mineral acids.

ACID, TARTARIC.—A corrosive vegetable poison.

Symptoms.—Very analogous to those caused by oxalic acid.

Morbid Appearances are likewise similar to those produced by oxalic acid.

Tests.—When heated on platina foil, it burns with a pale reddish flame, and exhales a peculiar acrid vapor, leaving much carbonized matter. When a solution is treated with lime-water, it affords a white precipitate, soluble in an excess of the acid; when treated with caustic potassa, it affords a granular precipitate of the bitartrate.

Treatment.—The same as for oxalic acid.

ACETATE OF COPPER. *See* Copper.

ACETATE OF LEAD. *See* Lead.

ACETATE OF MORPHIA. *See* Opium.

ACONITE. *See* Vegetable Poisons.

ALCOHOL.—A narcotico-acrid poison.

Symptoms.—These vary according to the dose. In small quantities, there is mere excitement. In large doses, much excitement, with delirium, confusion of intellect, followed by somnolency; nausea and vomiting, and even coma and apoplexy. When an undue portion has been swallowed, it often proves instantly fatal; it may also prove fatal by occasioning or aggravating other diseases. The effects of an habitual use of it are diseases of the viscera, and various mental and nervous disorders. The symptoms of poisoning with alcohol may be mistaken for those of epilepsy or apoplexy.

Morbid Appearances are inflammation, softening, &c., of the mucous membrane of the stomach, congestion of the cerebral vessels, and sanguineous or serous extravasation in the brain or lungs.

Tests.—Odor of the contents of the stomach, or of matters ejected from it. The chemical proof consists in removing the contents of the stomach, mixing them with distilled water, filtering and distilling in union with carbonate of potassa or soda; the product is mixed with fused chloride of sodium, and again distilled. Alcohol will be found in the receiver.

Treatment.—Withdraw the contents of the stomach, as speedily as possible, by a stomach-pump; afterwards an emetic of salt and water should be given in large doses, and repeated, at short intervals, till the stomach is well cleared. The cold

affusion is useful in some cases. Ammonia may be employed as a stimulant, and general symptoms obviated by blood-letting; but this must be employed with great caution.

ALKALIES.

AMMONIA.—A corrosive poison.

Symptoms.—Excoriation of the mouth and fauces; burning sensation in the throat and stomach, usually followed by vomiting and purging, the ejected matters being often bloody. When the quantity taken is large, an immediate feeling of strangulation ensues, attended with convulsions. If the result is fatal, it quickly follows the ingestion of the poison. The inhalation of ammonia by the nostrils, when too freely used in cases of fainting, has caused the same symptoms as when taken into the stomach.

Morbid Appearances.—Marks of high inflammation of the parts with which the poison has come in contact.

Tests.—The pungent odor; its alkaline reaction—but which is dissipated by heat. By causing a yellow precipitate with a mixture of arsenious acid and nitrate of silver; by producing a rich violet-blue solution with the salts of copper; [by yielding a yellow precipitate with the bichloride of platinum; by giving a white precipitate with the bichloride of mercury;] and forming white fumes with chlorine or hydrochloric acid.

Treatment.—The immediate administration of vinegar or one of the vegetable acids, and afterwards the copious use of demulcents. When ammoniacal vapor has been inhaled, the patient should inspire the vapor of vinegar.

POTASSA.—A corrosive mineral poison.

Symptoms.—An acrid, caustic, urinous taste in the mouth; a sensation of burning heat in the throat; nausea, and sometimes vomiting of bloody matters. The surface cold and clammy; the pulse quick and feeble; hypercatharsis, and violent colicky pains.

Morbid Appearances.—Strong marks of inflammation in the alimentary canal, softening, erosion of the mucous coat, and, in some cases, perforation of the stomach.

Tests.—Alkaline reaction; precipitation of nitrate of silver in the form of a dark-colored oxide. Carbonic acid water causes no precipitate. A concentrated solution affords a canary-yellow precipitate with bichloride of platinum. [A solution of tartaric acid causes a white precipitate of cream of tartar.]

Treatment.—Give vinegar and the diluted vegetable acids; to be followed by a free use of demulcents, or oleaginous mixtures.

SODA.—A corrosive mineral poison.

Symptoms and Morbid Appearances.—Analogous to those following the use of potassa.

Tests.—Caustic soda in solution is not precipitated by bichloride of platinum or tartaric acid; its alkaline nature can be ascertained by the usual tests. [Antimoniate of potassa affords a white precipitate, when added to the salts of soda. Soda tinges the outer flame of the blow-pipe yellow.]

Treatment.—The same as for potassa.

ANTIMONY.—Metallic antimony is not regarded as a poison; but its vapor is said to cause unpleasant symptoms when respired.

ANTIMONY, TARTARIZED, generally known as TARTAR EMETIC, is an irritant metallic poison.

Symptoms.—Nausea and severe vomiting, hiccough, cardialgia, burning heat at the epigastrium, severe colic and purging, small and rapid pulse, cold skin, syncope,

difficult respiration, vertigo, insensibility to external stimulants, painful cramps in the lower extremities, and great prostration of strength.

Morbid Appearances.—The stomach and intestines often much inflated with gas; their mucous membrane red, and covered with a slightly adhering viscid layer; the peritoneum of a dark brick-red hue; the membranes of the brain displaying marks of inflammation.

Tests.—In a solid state, add charcoal, introduce into a test-tube and expose to heat; metallic antimony will be found of a greyish-black lustre. In the state of solution, diluted nitric acid causes a white precipitate. Sulphuretted hydrogen throws down a reddish-orange precipitate. Solutions containing tannin cause a copious, curdled, whitish-yellow precipitate. Should matters from the stomach be present, the solution must be well agitated with a small portion of muriatic and tartaric acid; then filtered previous to being experimented upon.

Treatment.—Encourage vomiting by free administration of warm water and other diluents, or employ the stomach-pump, if necessary. Infusions, tinctures, or extracts, containing tannin, are very useful by decomposing the poison.

BARYTA, AND PREPARATIONS OF.—The only preparations of baryta that have caused death are the CARBONATE and CHLORIDE.

Symptoms.—Those of irritation, combined with an affection of the brain and nervous system, as vertigo, convulsions, and paralysis.

Morbid Appearances.—Evidences of inflammation of the mucous coat of the stomach.

Tests.—In solution: sulphuric acid throws down a profuse dense white (*sulphate of baryta*) precipitate. Solution of nitrate of silver also produces a white (*chloride of silver*) precipitate with the chloride of barium. Sulphuretted hydrogen produces no change of appearance; which circumstance, being peculiar to salts of barium, is valuable as a *negative test*.

Treatment.—Sulphates of magnesia or soda in solution in water, if the patient is seen early; they are, however, of little use where the carbonate has been taken. In this case, a mixture of sulphate of magnesia and diluted vinegar, together with emetics and the stomach-pump, should be used.

BELLADONNA. See Vegetable Poisons.

BISMUTH.—The only preparation that has caused death is the SUBNITRATE.

Symptoms.—Burning pain in the throat, nausea, vomiting and diarrhoea, accompanied by coldness of the extremities, and spasms.

Morbid Appearances.—Inflammation of the fauces and œsophagus, redness of the stomach and intestines.

Tests.—In substance: Dissolves in nitric acid, without change of color, and without effervescing, which distinguishes it from a *carbonate*. Sulphuric acid produces no precipitate, thus disproving the possible idea of its being a preparation of *lead*. Dissolve it in nitric acid, and add solution of potassa so long as any precipitate is thrown down; gather this in a filter, and, after washing and drying it, place it on charcoal, and treat it as directed for preparations of lead, when its reduction to the metallic condition will be effected.

[The nitrate is decomposed by being poured into a large quantity of cold water, yielding an insoluble subnitrate. The latter salt is blackened by exposure to sulphuretted hydrogen gas.]

Treatment.—Induce vomiting by tickling the fauces, &c., and give continuous full draughts of sweet milk.

BROMINE.—An irritant, corrosive mineral poison.

Symptoms.—Irritation and inflammation of the parts with which it comes in contact, whether in substance or in vapor.

Morbid Appearances.—Fauces, œsophagus, and stomach inflamed and corroded, mucous membrane gelatinized, duodenum of a yellow color, and thickened; the parts retaining a strong smell of the poison.

Tests.—Its color, odor, and volatility. All solutions containing it are rendered colorless by the addition of potassa. On being evaporated, the residue is to be incinerated at a low temperature, the ash dissolved in distilled water, filtered, and chlorine passed through the solution. On the addition of ether, the bromine is taken up by that fluid. [With nitrate of silver it affords a yellowish-white (*bromide of silver*) precipitate.]

Treatment.—The immediate and free use of albumen or starch, and magnesia.

BRUCIA. *See* Vegetable Poisons.

CALOMEL. *See* Mercury.

CAMPHOR. *See* Vegetable Poisons.

CANTHARIDES.—An acrid and corroding animal poison.

Symptoms.—A burning sensation in the throat; violent pain in the stomach and bowels; nausea, vomiting, and purging—the ejections being frequently bloody and purulent; great heat and irritation of the urinary organs, sometimes the most painful priapism; pulse quick and hard; and convulsions, tetanus, delirium, and syncope.

Morbid Appearances.—Inflammation and erosion of the stomach; if in substance, fragments of the flies, adhering to the mucous coat, or mixed with the contents of the stomach; marks of inflammation in the intestines and urinary organs: these are most evident when death does not soon follow the ingestion of the poison.

Tests.—The characteristic green, shining appearance of the fragments of the flies, and the character of the symptoms.

Treatment.—The promotion of vomiting by means of warm demulcents; copious dilution, bleeding, the warm bath, opiate frictions, enemata of mutton broth, laudanum, &c. Camphor, though not an antidote, alleviates some of the most distressing symptoms.

CARBONIC ACID GAS.

Symptoms.—Great drowsiness; difficulty of respiration; loss of sensibility; the countenance of a livid or deep leaden color.

Morbid Appearances.—Whole body, but especially the face, swollen, which is also usually livid, and the features distorted; eyes generally prominent, and retaining their brilliancy. Vessels of the brain and lungs in a state of congestion.

Tests.—Expose a vessel, filled with lime-water, to the deleterious atmosphere, when, if carbonic acid be present, it will assume a milky appearance, and let fall a white precipitate. A lighted candle will detect the presence of carbonic acid gas (if in excess) by being immediately extinguished in the tainted atmosphere; it must, for this purpose, be held near the floor. But the utmost caution should be observed in introducing lighted tapers into close situations *supposed* to contain this gas, lest an explosive hydrocarbon may, instead, be met with.

Treatment.—Remove the patient from the situation or apartment in which the poison has been inhaled; or open the doors, windows, &c., for admission of pure air. Continued application of cold water to the head and neck. Bleeding may be necessary; or cupping at back of the neck. Cautious inhalation of steam from warm water placed in a convenient vessel.

CHLORINE.—An acrid gaseous poison.

Symptoms.—Great and painful constriction of the glottis, cough, sensation of suffocation, alternating with asphyxia; if the case is protracted, inflammation of the

larynx and pneumonic inflammation; odor of the gas perceptible in the breath and in the vomited matters.

Morbid Appearances.—Marks of irritation and inflammation in all the parts with which the poison has come in contact.

Tests.—A solution of this gas has a yellowish color, and a strong, peculiar smell, which latter is evolved on boiling. It gives a white precipitate with nitrate of silver, which is insoluble in muriatic acid.

Treatment.—The best remedy is a free use of albumen. Magnesia with mucilaginous fluids may also be freely exhibited. [Inhalations of ammonia, ether, and the vapor of warm water, and even of sulphuretted hydrogen, have been recommended. The last article must be employed with great care.]

CHROME.—The only form of this metal that requires notice is the

BICHROMATE OF POTASSA.—An acrid mineral poison.

Symptoms.—Burning heat in the mouth, throat, and stomach; excessive and painful vomiting of bloody mucus; convulsions, palsy, &c.

Morbid Appearances.—Softening and abrasion of the mucous coat of the stomach and intestines.

Tests.—In substance, by its orange-red color. In solution, by affording a rich red precipitate with nitrate of silver, a bright yellow with acetate of lead, a dingy green with sulphuretted hydrogen.

Treatment.—The use of the carbonates of potassa or soda to neutralize the excess of chromic acid, followed by the administration of emetics. [It would also be advisable to employ chalk or magnesia, in connexion with milk or the albumen of eggs.]

COCCULUS INDICUS. See Vegetable Poisons.

COLCHICUM AUTUMNALE. See Vegetable Poisons.

CONIUM MACULATUM. See Vegetable Poisons.

COPPER.—This metal is not poisonous in a metallic state, but becomes highly so when oxidized, or in combination.

The most common preparations of copper are the ACETATE (*verdigris*), SULPHATE (*blue vitriol*), the CARBONATE, and AMMONIURET.

Symptoms.—Violent headache, vomiting, pain in the bowels, cramps in the lower extremities, a peculiar and permanent coppery taste, diarrhœa, convulsions, palsy, and insensibility.

Morbid Appearances.—Marks of inflammation in the stomach and intestines. When the case has been protracted, there is often a green tinge of the lining membrane, and a jaundiced appearance of the skin.

Tests.—The Sulphate in a solid state presents a bright blue color, and leaves an astringent metallic impression on the tongue.

In solution: the transmission of sulphuretted hydrogen affords a brownish-black (*sulphuret of copper*) precipitate. Solution of ammonia precipitates a blue substance, which, on further addition of ammonia, becomes dissolved; but when only a small amount of the poison is present, no precipitation occurs, a clear violet-colored solution being then at once presented. Ferrocyanide of potassium throws down a reddish-brown (*ferrocyanide of copper*) precipitate. A piece of polished zinc or iron (as the blade of a table-knife) acquires a plating of copper, if placed in a liquid containing the above salt. The tests for sulphuric acid will indicate its presence here as the combining acid.

The Acetate: Same tests as preceding, excepting the last.

The Ammonio-sulphate: has a brilliant violet-blue color, and ammoniacal odor. Solution of arsenious acid affords a fresh apple-green precipitate; this experiment is, conversely, that for arsenious acid.

Treatment.—Vomiting to be promoted by copious draughts of warm water, milk, or mucilaginous fluids; various antidotes have been highly praised. Sugar was formerly recommended, but has not been as successful as was supposed. Albumen and milk form an insoluble compound with copper, provided they are in large excess. The protosulphuret of iron and iron filings have also been employed with advantage; but their action is too slow. The hydrated oxide of iron has been successfully administered where the arsenite of copper had been taken.

CORROSIVE SUBLIMATE. *See* Mercury.

DIGITALIS. *See* Vegetable Poisons.

IODINE.

Symptoms.—In large doses, this substance causes heat and constriction of the fauces, nausea, offensive eructations, pain in the stomach, retching, colic, diarrhoea, quickness of the pulse, tremblings, great thirst, salivation, convulsions, &c.

Morbid Appearances.—Distension and inflammation of the stomach and bowels; sphacelation in some places; and, in protracted cases, increased size and paleness of the liver.

Tests.—In the solid form, it is in bluish-grey scales, having somewhat the odor of chlorine. When heated in a tube, it gives off violet fumes; when added to the mucilage of starch, it strikes a deep blue color. When it is mixed with other matters, pass a stream of sulphuretted hydrogen through the mixture, boil it, saturate with potassa, add mucilage of starch, filter, add nitric acid, and the color will indicate iodine.

Treatment.—Promote vomiting, resort to the free administration of amylaceous and mucilaginous fluids, and treat the inflammatory symptoms as they occur.

IODIDE OF POTASSIUM.

Symptoms.—Uneasiness of the stomach, followed by nausea, and a burning sensation in the stomach, vomiting, cephalalgia, vertigo, and tremors.

Morbid Appearances.—Stomach contracted; mucous membrane, with ecchymosed spots, and some abrasions; marks of inflammation in the intestines.

Tests.—Solid state: heat with the protoxide of manganese; moisten the powder with a mixture of equal parts of sulphuric acid and water; heat in a tube, and the purple vapor of iodine makes its appearance.

In solution: add to the filtered fluid an equal quantity of mucilage of starch, and a few drops of nitric acid, and the characteristic blue color will be produced. Test the urine in the same manner. It may also be detected in the solids of the body, by drying them, incinerating and lixiviating, filtering, and using starch and nitric acid.

Treatment.—The same as for iodine.

IRON.—Not poisonous in a metallic state; but some of its preparations act as irritants, especially the sulphate and the muriate.

Symptoms.—Nausea, vomiting, and purging; sensation of heat in the stomach and bowels; dejections of a black color, &c.

Morbid Appearances.—Softening of the mucous coat of the stomach and intestines; the whole extent of the alimentary canal of a black color; marks of inflammation.

Tests.—For the sulphate: ferrocyanide of potassium will give a greenish-blue precipitate, becoming a deep blue by exposure to the air; hydrosulphuret of ammonia will give a black precipitate; and gallic or tannic acid will give a bluish-black precipitate. The presence of sulphuric acid can be ascertained by means of nitrate of baryta.

Treatment.—In poisoning by either salt, magnesia or the alkaline carbonates should be freely given.

LEAD.—In a metallic state, lead does not appear to be poisonous; but most of its compounds are, especially the acetate, chloride, carbonate, and oxide.

Symptoms.—Obstinate constipation, violent colic with retraction of the abdomen; vomiting, pulse small and hard, laborious breathing, and tremors, gums with a blue tinge, paralysis of the extremities, &c.

Morbid Appearances.—The appearances on dissection are very various, but are principally those of inflammation.

Tests.—In a solid state: Mixed with potassa and charcoal, and exposed to a red heat (or placed in a suitably sized hollow in a piece of charcoal, and exposed to a blow-pipe flame), metallic lead is formed.

In solution: Sulphuric acid, and the sulphates, throw down a white (*sulphate of lead*) precipitate. Solution of carbonate of soda gives a white (*carbonate of lead*) deposit. Solution of chromate of potassa a rich yellow (*chromate of lead*) precipitate. [Solution of iodide of potassium affords a yellow (*iodide of lead*) precipitate.] Transmission of hydrosulphuric acid produces a blackish (*sulphuret of lead*) precipitate. Zinc precipitates lead from a solution of any of its salts.

To render the *carbonate* amenable to the foregoing tests, it must first be dissolved in a sufficiency of nitric acid diluted with distilled water.

Treatment.—Administer a solution of sulphate of zinc in full doses; induce and continue vomiting. If an insoluble salt of lead is the poison, almond or olive oil must at first be given in considerable quantity. The free exhibition of the sulphate of soda, or magnesia, or of milk, or albumen, to be followed by an emetic. In lead-colic, the treatment consists of active purgatives, followed by full doses of opium or Dover's powder. Inflammatory symptoms are counteracted by antiphlogistic means.

LOBELIA.—See Vegetable Poisons.

MERCURY.—In a metallic state, mercury exercises but a slight influence on the body, but in a state of vapor it is capable of causing violent symptoms. All the salts of mercury are poisonous; but the most important is corrosive sublimate.

Symptoms.—The symptoms caused by corrosive sublimate resemble those produced by arsenic, but, from the salt being more soluble, they are more immediate and violent; there is a more marked taste, the evacuations are more frequently bloody, and there is a whitened condition of the epithelium of the mouth. There are three varieties of poisoning with mercury. In the first, the leading symptoms are, violent irritation of the alimentary tube; namely, vomiting, purging, pain at the pit of the stomach, and irritation in the throat; metallic, styptic taste, corrosion of the mouth, tongue, and palate; constriction of the throat, and difficulty of swallowing. Blood evacuated both by vomiting and by stools, suppression of urine, countenance flushed, tumid, and bloated. In the second variety, salivation and sloughing of the mouth succeed to the irritation and inflammation. In the third, mercurial erethism comes on, and is not preceded by the symptoms of local irritation. The first variety arises from the more soluble salts of mercury, in large doses; the second, from the same preparations, but in smaller doses, and more diluted; the third, by the more insoluble and refractory compounds.

Morbid Appearances.—These are similar to those attendant on irritative or corrosive poisoning. There are, frequently, shrivelling of the tongue, and enlargement of its papillæ and root. In some cases, red and black spots in the cavities of the heart. Corrosion, ulceration, and disorganization of the mucous coat of the stomach and intestines. An inflamed condition of the urinary organs is also frequently observed.

Tests.—Corrosive sublimate in a solid state: Is sublimed when heated in a test-tube; and the acrimonious fumes speedily condense into a crystalline, semi-transparent mass. Placed in a test-tube, and lime-water, potassa, or soda, added in solution, a yellow (*peroxide of mercury*) precipitate is thrown down. Triturated with

an equal weight of well-dried carbonate of potassa, metallic globules of mercury will become visible.

In solution: Ammonia throws down a white (*ammonio-chloride of mercury*) precipitate. Solution of proto-chloride of tin affords a (*calomel*) precipitate, which, at first, is white, but acquires a leaden color on adding more of the test; when this precipitate (after being well agitated) is dried, minute globules of quicksilver are formed. Transmission of sulphuretted hydrogen produces a (*bisulphuret of mercury*) precipitate, which at first is leaden-colored, then black. Solution of iodide of potassium affords an intensely brilliant scarlet (*biniodide of mercury*) precipitate. Corrosive sublimate may be reduced to the metallic state through galvanic influence, applied as follows: A drop of the suspected fluid being laid on a piece of polished gold, and both it and the gold touched at the same instant by a point of iron (as a thick needle, or the end of a penknife), a small silvery coating of mercury soon becomes apparent on the gold. Being freely soluble in sulphuric ether, addition of this fluid is of much service when the poison is found mingled with organic or other matters.

The Bicyanide: When heated in a tube, cyanogen gas is evolved, which, when inflamed at the mouth, burns with a rose-red flame, with a blue halo.

Calomel: Is sublimed by heat. When heated along with potassa, or lime-water, it assumes, at first, a black appearance, and then quicksilver globules are observed.

The Biniodide: On cautious application of heat, it sublimes in red-colored crystals, which soon change to yellow, and subsequently to a dusky hue. On exposure to a sufficient degree of heat, iodine fumes are disengaged. When mixed with potassa (in equal weights) and heated in a test-tube, decomposition occurs; metallic mercury being sublimed, and iodide of potassium deposited in the tube.

Red Precipitate: Exposed to heat in a test-tube, metallic mercury is sublimed; evolution of oxygen being at the same time demonstrable.

The Persulphate: Treated in the same manner as the foregoing, sublimation of the mercury takes place; and sulphurous acid gas is perceptibly evolved.

Treatment.—If vomiting does not already exist, it must be produced by the exhibition of emetics. Various antidotes have been suggested for this poison. The best appears to be albumen of eggs in continuous large doses, and infusion of catechu afterwards; sweet milk; mixtures of flour with water (for the sake of the gluten) in successive cupfuls; iron filings; emetics of tartrate of antimony and potassa have been recommended, to combat ptyalism, when present; but to check excessive salivation, small doses of the chlorate of potassa appear to be the most successful.

[The hydrated proto-sulphuret of iron has been proposed by Mialhe, and confirmed by Orfila, who says, if it be taken immediately after the ingestion of the corrosive sublimate, it destroys completely the poisonous properties of the latter.]

MORPHIA. *See* Vegetable Poisons. Opium.

NUX VOMICA. *See* Vegetable Poisons.

OPIUM. *See* Vegetable Poisons.

PHOSPHORUS.—A corrosive poison.

Symptoms are protracted in their appearance. A hot alliaceous taste in the mouth; acrid burning sensation in the throat and stomach; nausea and vomiting. The vomited matters are of a dark color, and emit white fumes; pulse small and frequent; sometimes violent convulsions.

Morbid Appearances.—A general inflammatory appearance of the stomach and intestines, with sphacelated spots in various parts.

Tests.—The peculiarity of its odor; its highly inflammable property, when only moderately heated, and, when rubbed on the skin, its property of rendering the latter luminous in the dark.

Treatment.—Mixture of magnesia and cold water, in repeated draughts. Pro-

mote continuous vomiting by tickling application of a feather to the fauces. The exhibition of oil is injurious, as it tends to dissolve the phosphorus.

STRYCHNIA. See Narcotico-irritants, under the head of Vegetable Poisons.

TIN.—This is not poisonous in a metallic state; but some of its salts, as the chlorides, are violent irritants.

Symptoms.—An austere metallic taste; constriction of the throat, impeded respiration, violent vomiting, cramp of the stomach and violent colic, purging, pulse small and quick, convulsions, sometimes paralysis, asphyxia.

Morbid Appearances.—Inflammation and erosion of the stomach and intestines.

Tests.—Chloride of gold gives a deep purple-brown, almost black precipitate. Nitrate of silver affords a white precipitate, insoluble in nitric acid. Sulphuretted hydrogen throws down a deep chocolate-brown precipitate, even in diluted solutions.

Treatment.—Copious draughts of milk; then excite vomiting. Inflammatory symptoms to be combated by antiphlogistics, and nervous irritation by opiates and antispasmodics.

VEGETABLE POISONS.—These are very numerous, and possessed of the most different qualities; some being mere irritants, others narcotic irritants—whilst another class are pure narcotics.

IRRITANTS.—This includes a great number of vegetable substances, the principal of which are ALOES, ARUM, CAPSICUM, COLOCYNTH, CREASOTE, CROTON SEEDS, ELATERIUM, EUPHORBIIUM, GAMBOGE, JALAP, MEZEREON, CASTOR SEEDS, SAVIN, SCAMMONY, &c.

Symptoms.—Many of these articles act especially on the bowels, and in moderate doses are efficaciously used as purgatives. In large doses, they cause hypercatharsis, and much irritation of the stomach and bowels.

Morbid Appearances.—These are various degrees of inflammation of the stomach and bowels.

Tests.—None by means of reagents. The rich yellow color of *gamboge* may sometimes prove a means of its detection; and familiar acquaintance with others of the above substances may lead to their recognition by smell, taste, &c.

Treatment.—The expulsion of the poison by means of emetics, and, when this is effected, copious demulcent drinks in a warm state; succeeded by liquid cordial, stimulant, or opiate medicines suited to particular circumstances.

NARCOTICS.—This is a small class of the vegetable poisons, the most important of which are OPIUM, HYDROCYANIC ACID, (see page 555), and HYOSCYAMUS.

OPIUM.

Symptoms.—A dark, suffused countenance, drowsiness, stupor, perfect insensibility, followed by delirium or profound coma; then a pallid countenance, deep and stertorous breathing, cold sweats, slow and full pulse, cold and livid skin, suspension of all the secretions except perspiration; at length, the pulse becomes frequent, feeble, and thread-like; and, sometimes, convulsions, particularly in children.

Morbid Appearances.—Occasionally redness of the stomach and intestines, fluidity of the blood, engorgement of the lungs; but the only post-mortem appearance that appears to occur in all cases is turgescence of the vessels of the brain, sometimes with effusion of water on its surface or into the ventricles. But this condition of the brain does not furnish of itself any evidence of poisoning by opium, as it is frequently found to arise from a variety of other causes.

Tests.—Opium in substance: its peculiar taste, color, and odor (especially the latter quality when arising from opium in a softened or moist state, and more so if it be moderately heated; as when just obtained from the stomach).

In solution: Nitric acid throws down a red-colored (*nitrate of morphia*) precipitate. Tincture of the chloride of iron produces a (*meconate of iron*) deposit, which is also red. Tannic acid yields a white (*tannate of morphia*) precipitate.

When the suspected fluid has been procured from the stomach, it usually contains a mixture of organic substances; these must be broken down, and the whole rendered sufficiently fluid by the addition of distilled water *when necessary*, and then be slightly acidulated with acetic acid. The liquor thus obtained, when well agitated, and filtered, must be evaporated to consistence of syrup. And, after being digested with alcohol, it must be boiled, and again concentrated to nearly the consistence of syrup. When this alcoholic extract (as it has been called) is dissolved in distilled water, the solution of ammonia, cautiously added, produces a precipitate of morphia, which acquires a deep-red color changing to yellow, when nitric acid is dropped into it, and a blue color on the addition of tincture of chloride of iron. When the last named precipitate is removed by filtration, and acetate of lead added to the supernatant (*meconate of ammonia*) liquor, a (*meconate of lead*) deposit is afforded; which, when suspended in water, is decomposed by the transmission of sulphuretted hydrogen; a dark precipitate being thereby thrown down. The fluid remaining from this last process acquires a deep cherry-red hue on the addition of the tincture of chloride of iron.

Morphia in the solid state: Is only slightly soluble in water. Nitric acid turns it first to an orange, and then to a deep orange-red color. Chloride of iron gives a blue color, which is destroyed by acids. Iodic acid is decomposed by morphia, and iodine is set free. To show this, the acid is to be mixed with starch; on the addition of morphia, the iodine combines with the starch, which becomes of a blue color, if the quantity be large, or of a reddish or purplish tint, if it be small.

Treatment.—The stomach to be evacuated by means of the stomach-pump, as speedily as possible. This is to be pursued until the water used no longer has the smell or taste of opium. When a stomach-pump cannot be procured, or if the patient shall have swallowed solid opium, emetics of sulphate of zinc should be administered, until the stomach is freed; strong decoctions of tea or coffee, or other vegetable astringents, are found useful in the intervals. Cold affusions on the head, chest, and spine, have been used with great success. Flagellation and other means of arousing the patient from a state of lethargy must also be employed. Bleeding should not be resorted to until all the poison has been eliminated, as the abstraction of blood tends to promote absorption, and, even then, with much caution. The administration of vinegar is also objectionable. The best liquid that can be given is a strong decoction of coffee. The various antidotes that have been recommended are useless.

[The evidence, in favor of the successful employment of artificial respiration in cases that were apparently desperate, has become so strong and unquestionable, that no practitioner is justifiable in abandoning his case until this measure has been faithfully tried. Active stimulation is often required.]

HYOSCYAMUS.

Symptoms.—Sickness, stupor, dimness of sight, and delirium, followed by coma and much dilatation of the pupils; pulse at first hard, but becoming weak and tremulous; petechiæ often making their appearance before death.

Morbid Appearances.—Inflammation of the stomach, bowels, and brain.

Tests.—There are no tests that can be relied upon, except a recognition of the plant, and the nature of the symptoms.

Treatment.—The speedy evacuation of the poison, by means of emetics and purgatives, and the subsequent use of acidulous drinks.

[The Editor doubts the propriety of the latter recommendation, notwithstanding the high European authority in its favor, because the use of acidulous drinks will, most certainly, result in the formation of soluble salts with the remnants of the hyoscyamia that may be in the stomach; thereby adding to the danger. He would suggest, instead, the free use of vegetable astringents in solution.]

NARCOTICO-IRRITANTS.—These are very closely allied in their effects to the last class, but have a more direct action on the spinal marrow and nerves, as shown by the more frequent occurrence of convulsions and paralysis. They differ much from

each other in their action on the system; most of them, however, owing their properties to the presence of an alkaloidal principle. The most important are: DIGITALIS, VERATRUM, CONIUM, COLCHICUM, LOBELIA, ACONITUM, BELLADONNA, STRAMONIUM, TABACUM, NUX VOMICA, &c.

Symptoms.—These are various, but, in general, vertigo, coma, delirium, paralysis, or convulsions, with disturbance and pain in the stomach and intestines, are observed. Those belonging to the nux vomica tribe have marked effects on the spinal marrow, causing tetanus and convulsions, but seldom coma or delirium; whilst, on the other hand, squill and foxglove produce symptoms of narcotism, preceded by vomiting, and other signs of irritant action on the stomach.

Morbid Appearances.—These, like the symptoms, are very various. In some cases, there is inflammation of the stomach and intestines; in others, this is wholly wanting. Where there have been symptoms of cerebral disturbance, traces of congestion of the brain are usually discernible, but are not found in every case.

Tests.—Most of these poisons, as before said, owe their deleterious powers to the presence of an alkaloidal principle, which is exceedingly difficult to detect by chemical tests; hence the strongest evidence is to be derived from an inspection of the fragments of the plant itself, if it has been administered in substance; but this mode of discrimination fails where the poison has been taken in the form of extract, infusion, or decoction. In such cases, the only reliance is on the symptoms and concomitant circumstances.

[Conia, the active principle of conium, when liberated from its combinations by potassa, is volatile, and exhales a strong mouse-like or urinous odor. With the vapor of muriatic acid, it forms dense white fumes. The odor of conia is so characteristic, that it can scarcely be confounded with any other poisonous agent.

Veratria (obtained from veratrum), affords a reddish-yellow solution with nitric acid, an intensely-red one with concentrated sulphuric acid, and a white precipitate with tannic acid. When applied to the nose, it causes severe sneezing.

Brucia (found in the seeds of the nux vomica) is colored red by nitric acid, and this color changes to violet by the addition of proto-chloride of tin. With chlorine it gives a red color. Sulphuric acid first reddens brucia, and then turns it yellow and green.

Pure strychnia (also found in the nux vomica) may be recognized by rubbing a small portion with a few drops of sulphuric acid, containing one-hundredth of its weight of nitric acid. No change ensues; but the addition of a very small quantity of the per-oxide of lead, or of bichromate of potassa, changes the solution to a blue color, then to a red, and in the course of a few hours to a yellow color. Commercial strychnia generally affords a red color, changing to a yellow, with nitric acid, owing to the presence of brucia.]

Treatment.—This consists in the prompt use of emetics, or the stomach-pump, followed by the administration of purgatives. No antidote can be relied upon; but, as tannic acid decomposes these alkaloids, the free use of decoctions or infusions containing it may be resorted to with advantage.

ZINC.—This has no poisonous properties in a metallic state, but several of its salts possess active powers. Of these, the sulphate or white vitriol is the most common, and therefore the most likely to occasion unpleasant consequences.

Symptoms.—Pain in the abdomen, violent vomiting and diarrhœa, quick pulse, paleness and contraction of the features, coldness of the extremities. There is always a very austere taste in the mouth.

Morbid Appearances.—Marks of inflammation in the stomach and intestines.

Tests.—Ammonia gives a white precipitate, soluble in an excess of the alkali. Chromate of potassa affords an orange-red deposit, the chromate of zinc.

Treatment.—The free exhibition of warm water with milk, albumen, magnesia, &c. Infusions containing tannic acid. Where the poison has entered the bowels, emollient clysters are to be given.

TABLE OF POISONS AND ANTIDOTES.

The following table, from Taylor, "On Poisons," gives a succinct view of the most approved antidotes for the several poisons noticed in the foregoing pages:—

NON-METALLIC POISONS.

<i>Poisons.</i>	<i>Antidotes.</i>
Mineral Acids. { Sulphuric, Nitric, Muriatic, Nitro-Muriatic,	{ Magnesia mixed with water or milk; carbonate of lime; compound chalk powder; soda, potassa; the fixed oils.
Vegetable Acids. { Oxalic, Tartaric, Binoxalate of potassa,	{ Carbonate of lime. (Chalk or whiting.) Carbonate of lime. Sulphate of lime and water. Carbonate of soda in solution.
Salts. { Bitartrate of potassa, Potassa, soda, ammo- nia, and their car- bonates,	{ Vinegar, lemon juice, citric acid, oil.
Alkalies. { Baryta and its soluble salts, Carbonate of baryta,	{ Sulphate of soda, potassa, magnesia or lime. Mixture of sulphate of magnesia and vinegar.
Salts. { Alum,	{ Carbonate of soda or ammonia.

METALLIC POISONS.

Arsenic and soluble arsenites.	{ Hydrated per-oxide of iron; hydrated magnesia.
Corrosive sublimate and salts of mer- cury.	{ Mixture of oil and lime water.
Soluble salts of lead.	{ Albumen, gluten, or flour diffused in water; milk.
Carbonate of lead.	{ The alkaline, or earthy sulphates. Sulphate of magnesia and vinegar.
Soluble salts of copper.	{ Albumen, gluten, flour diffused in water; milk.
Tartar emetic.	{ Decoctions and tinctures containing tan- nic acid.
Chloride of antimony.	{ Magnesia.
Salts of tin.	{ Carbonate of soda; magnesia.
Sulphate or acetate of zinc.	{ Milk; carbonate of soda; magnesia.
Sulphate of iron.	{ Milk; carbonate of soda; magnesia.
Nitrate of silver.	{ Carbonate of soda or ammonia. Chloride of sodium.

NARCOTIC POISONS.

Opium; hyoscyamus.	{ Emetics; stomach pump; cold affusion. Strong decoction of coffee; electro- magnetism; tannic acid.
Prussic acid.	{ Ammonia; chlorine; cold affusion.

MODES OF DISTINGUISHING SOME OF THE VEGETABLE ALKALOIDS, WHEN IN POWDER.

Treat the powder with nitric acid; this is colored red by *Brucia*, *Delphia*, *Morphia*, and by the *Strychnia* of commerce, but not the pure. If the reddened acid becomes violet on the addition of protochloride of tin, it is *Brucia*; if it becomes black and carbonaceous, it is *Delphia*. If the powder is fusible without decomposition, and decomposes iodic acid, it is *Morphia*; if it is not fusible without decomposition, and does not decompose iodic acid, it is *Strychnia*. If the powder strikes a green with nitric acid, it is *Solanina*; if insoluble in ether, and does not redden nitric acid, it is *Emetia*; if soluble in ether, does not redden nitric acid, but melts and volatilizes with heat, it is *Atropia*; if thus affected by ether or nitric acid, but does not volatilize, it is *Veratria*.

INDEX

OF

DISEASES AND THEIR REMEDIES.

- ABSCCESS.** Lead ointment, 360. Lead cataplasm, 360. Caustic potassa, 371. Soap cataplasm, 412.
- ABSCCESS, MAMMARY.** Liniment acetate lead, 357. Oint. iod. potass. and morphia, 366.
- ACIDITY OF STOMACH.** Ammonia, 103. Lime water and milk, 156. Prepared chalk, 156. Alkaline infusion cascarilla, 173. Milk and soda water, 295. Magnesia mixt., 305. Carbonate potass., 375. Alkaline tincture rhubarb, 398. Purgative pastilles, 418. Carbonate soda, 434. Lozenges, soda, 436.
- ACNE.** Wash of benzoin, 147. Ointment of subacetate lead, 358. Compound powder sulphur, 449.
- AMAUROSIS.** Ammoniac mixture, 111. Anemone, 116. Chloride of gold and sodium, 138. Comp. powder sulphate mercury, 278. Pills strychnia, 446. Collyrium and liniment strychnia, 446. Powder white hellebore, 467.
- AMENORRHŒA.** Hiera picra and other preparations of aloes, 94, 98, 162. Rufus's pills, 94. Injection ammonia, 103. Muriate ammonia, 108. De Haen's pills, 111. Pills of ammoniac, 111. Iodide of calcium, 155. Cantharides and savine, 163. Compound pills of columbo, 200. Mayweed, 208. Saffron, 210. Infusion dittany, 212. Metallic iron, 228. Bolus metallic iron, 228. Ammoniated iron, 229. Ointment bromide iron, 230. Pills carbonate iron, 230. Mixture chloride iron, 233. Lozenges iodide iron, 235. Ointment iodide iron, 237. Black oxide iron, 239. Phosphate iron, 240. Sulphate iron, 240. Hooper's pills, 95, 241. Compound galbanum pills, 247. Compound infusion gentian, 251. Compound tincture guaiacum, 258. Ammoniated tincture guaiacum, 259. Infusion pennyroyal, 260. Bacher's pills, 261. Tincture black hellebore, 261. Compound pills iodide mercury, 275. Saturated tincture iodine, 284. Clyster iodine, 285. Oxide of manganese, 310. Horehound, 312. Myrrh and sulphate iron, 325. Tincture myrrh and hellebore, 326. Madder, 402. Decoction madder, 402. Savine and ginger, 405. Savine pills, 405. Oil savine, 405. Seneka, 423. Compound infusion senna, 426. Mixture of borax, 434.
- ANASARCA.** Calomel and squill, 272. Cream of tartar, 383. Mixture dandelion, 456.
- ANÆSTHETIC REMEDIES.** Sulphuric ether, 92. Chloroform, 183.
- ANGINA PECTORIS.** Assafetida and musk, 133. Succinate of ammonia, 110.
- ANGINA SYPHILITIC.** Mercurial gargle, 265.
- APHONIA.** Decoction arnica, 128.
- APHTHÆ.** Decoction wild indigo, 142. Decoction barberry, 148. New Jersey tea, 177. Chlorine, 183. Infusion goldthread, 206. Collutory creasote, 209. Soot mouth-wash, 246. Collutory rhatany, 294. Mixture carbonate potassa, 377. Lozenges borax, 433. Collutory borax, 433. Honey borax, 434.

ARSENIC, POISONING BY. Hydrated oxide of iron, 238. Mixture of sulphuret of potassium, 371.

ASCARIDES. Suppository aloes, 97. Injection aloes, 97. Clyster Southern wood, 130. Assafetida and iron, 133. Clyster assafetida, 134. Clyster camphor, 161. Wormseed, 181. Compound tincture tartrate of iron, 242. Mixture of rue, 403. Mixture senna, 427. Infusion tansy, 454.

ASCITES. Electuary Per. bark and iron, 189. Colocynth mixture, 199. Pill foxglove, 218. Foxglove and tartaric acid, 219. Elaterium mixture, 222. Gamboge mixture, 250. Black sulphuret of mercury, 279. Cream of tartar, 383. Mixture dandelion, 456.

ASTHMA. Ammoniac pill, 111. Ammoniac mixture, 111. Assafetida mixture, 133. Benzoin, 147. Catalpa, 175. Coffee, 195. Tincture larkspur, 217. Pill foxglove, 218. Skunk cabbage, 220. Galbanum mixture, 247. Henbane and squill, 282. Mixture elecampane, 283. Oxymel elecampane, 284. Ipecacuanha and carbonate soda, 286. Ipecacuanha and nitre, 286. Ipecacuanha and foxglove, 287. Tincture lobelia, 303. Opium and foxglove, 337. Opium and castor, 338. Solution cyanide potass., 363. Scillitic acetate potassa, 372. Mixture of carbonate potassa, 376. Mixture extract elder berries, 409. Squill and ammoniac, 419. Compound tincture squill and benzoin, 420. Compound stramonium pills, 444.

ATROPHY. Mixture eggs, 346.

BED SORES. Alum liniment, 102. Mixture of tincture camphor, 161. Carbonate lead plaster, 359. Liniment and ointment tannate of lead, 361.

BLADDER, DISEASES OF. Benzoic acid, 80; and copaiba, 80. Sal ammoniac draught, 109. Emulsion cantharides, 164. Buchu, 151. Electuary of cubebs, 211. Infusion of flaxseed, 301. Pareira brava, 348. Paullinia, 348. Mixture carbonate soda and copaiba, 436. Compound powder uva ursi, 462.

BONES, PAINS OF. Decoction mezereon, 318. Tincture acetate strychnia, 447

BRONCHITIS. Anisated ammoniated alcohol, 107. Nitrate of silver, 126. Indian turnip, 131. Iodide calcium, 155. Creasote pills, 208. Monesia, 319. Linctus opium, 340.

BUBO. Mercurial plaster, 267. Anodyne ointment, 340. Plaster acetate lead, 357. Lead ointment, 360. Lead cataplasm, 360.

BURNS, SCALDS. Ointment wild indigo, 142. Lotion benzoin, 147. Solution chlorinated lime, 154. Lime water and oil, 156. Compound creasote ointment, 209. Carded cotton, 255. Cerate cherry-laurel, 299. Liniment flaxseed oil, 303. Kentish's ointment, 333. Liniment eggs, 347. Basilicon ointment, 393.

CALCULOUS AFFECTIONS. Draught muriatic acid, 85. Phosphate of ammonia, 109. Lime water and carbonate potassa, 156. Peruvian bark, 185. Wild potato, 204. Mixture Prussian blue, 234. Nitrate of iron, 238. Magnesia and gentian, 251. Solution nitro-saccharate lead, 361. Solution caustic potassa, 372. Borate of potassa, 374. Mixture bicarbonate potassa, 377. Pills carbonate soda, 435. Lozenges bicarbonate soda, 436. Injection carbonate soda, 435. Uva ursi, 462.

CALCULI, BILIARY. Turpentine and ether, 333.

CANCER. Arsenical powder, 80. Solution of arsenic, 80. Injection alum, 101. Acetate ammonia, 105. Iodide arsenic, 129. Chloride gold, 137. Chloride barium, 142. Compound pills belladonna, 145. Extract marigold, 158. Liniment marigold, 158. Carrot ointment, 171. Hemlock, 201. Cataplasm hemlock, 202. Arseniate of iron, 230. Pills red oxide iron, 239. Compound soot ointment, 246. Hop ointment, 264. Infusion cherry-laurel, 298. Ointment cherry-laurel, 299. Ointment turpentine, 457. Ointment carbonate lead, 359.

- Ointment chloride lead, 359. Lotion chloride lead, 359. Ointment iodide potass. and opium, 365. Solution chlorinated soda, 437. Chloride zinc, 471.
- CANCERUM ORIS. Pyroligneous acid, 79. Solution chlorate potassa, 378.
- CARDIALGIA. Oil of wormwood, 75. Mixture boracic acid, 81. Acetic ether, 91. Ammonia, 103. Lime-water and potass., 156. Columbo and magnesia, 200. Compound powder of kino, 293. Magnesia mixture, 305. Nux vomica, 328. Rhubarb and chalk, 395. Carbonate soda and rhubarb, 435.
- CARIES. Phosphoric acid, 86. Creasote ointment, 209.
- CATARACT. Anemone, 116. Solution of atropia, 135.
- CATARRH. Gum mixture, lozenges, &c., 76. Mixture benzoic acid, 81. Infusion and syrup of maidenhair, 90, 91. Hydrocyanic ether, 92. Garlic, 93. Marsh mallow, 98. Ammoniac mixture, 111. Ammoniac and nitric acid, 111. Ammoniated anisated alcohol, 107. Almond emulsion, 115. Starch lozenges, 116. Lozenges kermes mineral, 120. Mixture kermes mineral, 121. Tronchin's lozenges, 121. Tartar emetic and opium, 122. Arum, 131. Syrup asparagus, 132. Butter cocoa, 152. Catechu and liquorice, 176. Powder spermaceti, 179. Spermaceti mixture, 179. Mixture bittersweet, 221. Emetia lozenges, 223. Barley sugar, 263. Prepared liquorice, 254. Mixture henbane, 281. Compound pills elecampane, 283. Mixture elecampane, 283. Compound oxymel elecampane, 284. Ipecacuanha lozenges, 287. Lozenges ipecacuanha and camphor, 287. Sugar of milk and Iceland moss, 295. Lozenges lactucarium, 297. Horehound syrup and candy, 312-13. Pectoral hydromel, 315. Mixture balsam Peru, 141. Myrrh and squill, 325. Solution cyanuret potassium, 363. Nitrate potassa and orris root, 379. Syrup red poppies, 399. Squill and ipecacuanha, 418. Seneka and prep., 423-4. Sulphur and liquorice, 449. Sulphur and orris root, 449. Turpentine and myrrh, 456. Emulsion balsam tolu, 458. Mixture tolu and morphia, 459. Green hellebore, 468. Pastilles de Paris, 485.
- CEPHALALGIA. Mixture vinegar and cardamom, 77. Lotion ammonia, 104. Carbonate ammonia, 106. Asarabacca, 131. Ward's essence, 161. Castor plaster, 175. Sneezewort, 260. Liniment oil amber, 448. Mixture valerian and ammonia, 464. Valerianate zinc, 475.
- CEREBRAL AFFECTIONS. Clyster tartar emetic, 123. Horseradish, 128. Clyster colocynth, 199. Bolus of foxglove, 218. Infusion galium, 248. Mixture borotartar potass., 374. Clyster common salt and arnica, 431.
- CHANCER. Syrup gold, 137. Ointment gold, 137. Creasote lotion, 209. Mercurial lotion, 265. Mercurial liniment, 267. Black wash, 277.
- CHAPS. Glycerin, 254. Anodyne ointment, 339. Liniment carbonate potassa, 376. Cold cream, 114, 179, 481. Ointment elder leaves, 409. Camphorated soap, 410.
- CHILBLAINS. Muriatic acid lotion, 85. Sulphuric acid liniment, 87. Alum ointment, 102. Solution chloride lime, 155. Camphor ointment, 161. Liniment of cantharides, 166. Tincture of Cayenne pepper, 167. Compound ointment creasote, 209. Ointment henbane, 282. Lotion iodine, 285. Liniment balsam Peru, 141. Turpentine lotion, 332. Embrocation petroleum, 349. Soap iodide potass., 366. Lotion carbonate potass., 376. Liniment pellitory, 385. Rose oil, 400. Mustard liniment, 431. Borax ointment, 434.
- CHLOROSIS. Pills aloes and iron, 94. Barthez's pills, 96. De Haen's pills, 111. Metallic iron, 227. Compound powder metallic iron, 228. Pills metallic iron, 228. Bolus ammoniated iron, 229. Pills carbonate iron, 231. Tincture chloride iron, 232. Bath iodide iron, 237. Black oxide iron, 239. Electuary black oxide iron, 239. Sulphate iron, 240. Bland's pills, 240. Syrup sulphate iron, 241. Tartrate of iron and potass., 242. Tannate iron, 243. Compound galbanum pills, 247. Oxide manganese, 310. Chocolate with paullinia, 348. Compound rhubarb pills, 395. Decoction madder, 402. Compound powder valerian, 463.

CHOLERA. Mixture nitric acid, 85. Liniment nitric acid, 85. Compound bismuth powder, 148. Camphor powder and mixture, 159-60. Extract hemp, 162. Infusion logwood, 259. Sugar milk draught, 296. Opium and acetate lead, 338. Compound pill acetate lead, 356. Mixture bicarbonate potassa, 377. Stevens's saline powder, 378. Plaster sulphate quinia, 392.

CHOLERA INFANTUM. Logwood, 259. Milk and suet, 295. Acetate lead, 356. Infusion benne, 429.

CHOREA. Arsenic, 129. Ammoniated copper, 214. Carbonate iron, 230. Mixture Dippel's animal oil, 329. Mixture cyanide potass., 363. Oxide zinc, 470. Ethereal tincture chloride zinc, 471. Ferrocyanuret zinc, 472. Black snakeroot, 185, 479.

COLIC. Alkaline tincture wormwood, 75. Aniseed, 117. Dewees' carminative, 134, 306. Clyster assafetida, 134. Infusion caraway, 171. Embrocation caraway, 172. Clove plaster, 172. Infusion catnep, 175. Infusion cinnamon, 193. Infusion fennel, 245. Oil pennyroyal, 261. Oil laurel, 298. Mace, 305. Dalby's carminative, 306. Plaster opium and camphor, 339. Opium liniment, 344. Cataplasm black pepper, 352. Emulsion carbonate potassa, 376. Tincture rhubarb and gentian, 397. Compound sagapenum pills, 406. Terebinthinate balsam sulphur, 450. Infusion tobacco, 453. Clyster turpentine, 456. Turpentine mixture, 457. Ginger, 476. Spice plaster, 477.

COLICA PICTONUM. Sulphuretted water, 83. Sulphuric acid lemonade, 87. Alum mixture, 101. Alum julep, 102. Liniment belladonna, 146. Electuary of jalap, 290. Opium and sulphur, 337. Opium and cinnamon water, 341.

CONDYLOMATA. Lotion corrosive sublimate and camphor, 270. Red oxide mercury, 277.

CONSTIPATION. Acetic acid clyster, 79. Preparations aloes, 94. Assafetida and aloes, 94. Mitchell's pills, 95. Anderson's pills, 95. Webster's pills, 95. Morrison's pills, 95. Griffitt's pills, 96. Tartar emetic clyster, 123. Pills of aloes and assafetida, 134. Charcoal electuary, 169. Pills elaterium, 222. Extract butternut, 291. Compound pills buckbean, 317. Compound rhubarb pills, 395. Rhubarb and ox gall, 396. Senna, 424-8. Compound solution sulphate of soda, 439. Rochelle salts, 439. Compound powder sulphur, 449. Clyster turpentine, 456.

CONTUSIONS. Sulph. acid and alcohol, 87. Lotion sal ammoniac, 108. Liniment sal ammoniac, 109. Arnica and rue, 129. Lotion sulphate copper, 215. Cataplasm hops, 264. Arquebusade water, 317. Compound spirit nutmeg, 325. Fomentation carbonate potass., 376. Lotion nitrate potass., 380. Wine and vinegar of roses, 401. Aromatic fomentation, 401. Aromatic vinegar, 401. Opodeldoc, 411. Soap liniment, 411. Fomentation common salt, 431. Turpentine liniment, 457.

CONVULSIONS. Garlic, 93. Assafetida mixture, 133. Powder of foxglove, 218. Musk clyster, 324. Liniment oil amber, 448.

CORNEA, AFFECTIONS OF. Aloes, 94. Anemone, 116. Ointment cyanide silver, 126. Ointment iodide silver, 126. Ointment sulphate cadmium, 153. Powder calomel and sugar, 272. Compound ointment cod liver oil, 330. Collyrium caustic potass., 372. Collyrium carbonate potass., 375.

CORNS. Plaster ammonia, 104. Verdigris ointment, 213. Verdigris plaster, 213. Lotion iodine, 285.

CORYZA. Lozenges cubebs, 211. Injection opium, 340.

COUGH. Mixture gum Arabic, 76. Jackson's cough syrup, 77, 484. Hydrocyanic ether, 92. Ammoniac, 111. Almond emulsion, 114. Green linctus, 114. White linctus, 115. Tartar emetic, 122. Antimonial wine and ammoniac, 124. Assafetida and squill, 133. Camphorated cough mixture, 162. Tincture cochineal, 194. Mixture quince seeds, 216. Liquorice, 254. Wistar's cough lozenges, 255. Barley sugar, 263. Powder of ipecacuanha and myrrh, 287. Lactu-

carium, 297. Syrup of muriate of morphia, 322. Lozenges of naphthaline, 327. Brown cough mixture, 255, 344. Opiate linctus, 344. Wistar's lozenges, 345. Pectoral syrup, 347. Squill, 418-22. Seneka and prep, 423-4. Storax, 445. Mixture oil amber, 448. Linctus with sulphur, 450. Mixture tolu and belladonna, 459. Oxide zinc, 470. Jackson's lozenges, 484. Pastilles de Paris, 485.

CRAMP IN THE STOMACH. Ammoniated tincture of castor, 175. Ferrocyanide zinc, 472.

CROUP. Carbonate ammonia ointment, 106. Sulphuretted syrup, 108. Sulphate copper, 215. Syrup sulphuret potass., 370. Compound syrup sulphuret potass., 370. Mixture sulphuret potass., 371. Compound syrup squill, 419. Oxy-mel squill and valerian, 421. Mixture seneka, 423. Cataplasm tobacco, 453.

CUTANEOUS DISEASES. Hydrosulphuretted bath, 83. Muriatic acid bath, 84. Lotion alum, 102. Arseniate ammonia, 105. Hydriodate ammonia, 107. Sulphuretted hydrosulphate ammonia, 108. Sulphuret antimony, 120. Lotion tartar emetic, 123. Iodide arsenic, 129. Donovan's solution, 129. Chloride barium, 142. Solution bromine, 149. Lime ointment, 156. Anthracokali, 169. Ointment iodide carbon, 170. Creasote ointment, 209. Decoction bittersweet, 221. Extract bittersweet, 221. Mixture soot, 245. Fuligokali, 246. Decoction guaiacum, 257. Guaiacum and sulphur, 258. Compound mercurial powder, 264. Solution acetate mercury, 268. Ointment white precipitate, 268. Bateman's lotion, 270. Plummer's pills, 272. Calomel ointment, 273. Ointment red iodide mercury, 276. Ointment nitrate mercury, 276. Ointment sulphate mercury, 279. Black sulphuret mercury, 279. Red sulphuret mercury, 279. Compound decoction mezereon, 318. Cod liver oil, 329. Phosphorated cerate, 350. Burgundy pitch, 354. Tar water, 354. Lotion cyanide potass., 363. Ointment iodide potass. and mercury, 365. Ioduretted water, 366. Sulphuret potass., 369. Lotion sulphuret potass., 370. Barlow's lotion, 370. Mixture sulphuret potass., 371. Aromatic bath, 401. Extract elder berries, 409. Sarsaparilla, 413-16. Ointment bromide sodium, 431. Sulphuret sodium, 432. Arseniate soda, 433. Soda and ipecacuanha, 435. Ointment carbonate soda, 435. Hyposulphite soda, 437. Compound pills sulphate soda, 439. Lotion sulphate soda, 439. Sulphur and cream of tartar, 449. Poison oak, 460. Decoction elm bark, 461. Decoction white hellebore, 467. Iodide zinc, 472. Arsenical pills, 80. Lotion of hydrocyanic acid, 83. Hydrosulphuretted bath, 83. Ointment of nitric acid, 85. Goulard's lotion, 114. Safflower, 171. Glycerin, 254. Ethereal tincture of iodine, 284. Muriate of manganese, 310. Fowler's solution, 374. Mixture of iodine and arsenic, 374. Di-arsenite of quinia, 388. Zittman's decoction, 414. Syrup of Laffecteur, 415. Iodide of sulphur, 452.

DEAFNESS. Cayenne pepper lozenges, 167. Acoustic balsam, 477. Injection caustic potassa, 372.

DEBILITY. Preparations cacao, 151. Camphor mixture, 160. Hartshorn jelly, 207. Preparations iron, 228-243. Barley meal, 263. Clyster eggs, 346. Mixture eggs, 346. Chocolate and paullinia, 348.

DELIRIUM TREMENS. Opium and musk, 337. Laudanum and tartar emetic, 345.

DIABETES. Phosphoric acid, 86. Hydrosulphate ammonia, 108. Compound pills ammoniated copper, 214. Compound pills kino, 293. Pills acetate morphia, 321. Carbonate potassa and ammonia, 377. Sulphur mixture, 450.

DIARRHŒA. Tannic acid, 88. Acetate alumina, 99. Alum, 100. Angustura, 117. Mixture chamomile, 119. Antimonial wine and laudanum, 124. Arnica, 128. Dewees' carminative, 134. Clyster bistort, 148. Compound powder chalk, 157. Lozenges chalk, 157. Chalk mixture, 157. Camphor water and laudanum, 159. Camphor water and nitric acid, 160. Compound cascarilla

powder, 173. Catechu, 176. Chlorine clyster, 183. Peruvian bark and rhatany, 188. Electuary Peruvian bark, 189. Infusion columbo and ginger, 200. Columbo and cascarilla, 201. Creasote mixture, 209. Sulphate copper, 215. Electuary sulphate copper, 215. Wine persimmons, 220. Infusion black purslane, 226. Mixture ammoniated iron, 229. Persesquinitrate iron, 238. Compound infusion galls, 248. Syrup galls, 248. Geranium, 252. Decoction pomegranate rind, 256. Infusion logwood, 259. Electuary logwood, 259. Compound powder kino, 293. Electuary kino, 293. Compound powder rhatany, 293. Monesia, 319. Injection acetate morphia, 321. Opium and chalk, 336. Roasted opium, 336. Plaster opium and camphor, 339. Mixture wine of opium, 342. Sedative mixture, 342-348. Bolus allspice, 351. Acetate lead, 356. Acetate lead mixture, 358. Decoction oak bark, 387. Confection acorns, 387. Roasted rhubarb, 395. Rhubarb mixture, 398. Confection dog rose, 399. Electuary roses, 400. Decoction blackberry root, 402. Infusion benne, 429. Hardhack, 441. Compound turpentine plaster, 457. Tormentil, 459. Infusion slippery elm, 462. Mixture of gum Arabic, 77. Subnitrate of bismuth, 148. Columbo and salep, 201. Sweet fern, 201. Oil of ergot, 225. Tincture of gentian, 252. Avens, 253. Barley, 263. Magnesia and rhubarb, 306. Marsh rosemary, 443.

DISINFECTION. Prophylactic vinegar, 79. Gaseous muriatic acid, 84. Chloride of lime, 154. Chlorine, 182. Solution chloride potass., 363. Solution chloride soda, 437.

DROPSY. Indian hemp, 124. Infusion horseradish, 127. Compound spirit horseradish, 128. Decoction asparagus, 132. Extract asparagus, 132. Chloride gold and sodium, 138. Hairy horehound, 141. Wine bryony, 150. Camphorated ethereal tincture cantharides, 164. Vinegar cayenne pepper, 168. Infusion carrot seeds, 171. Celandine, 181. Pipsissewa, 182. Black snakeroot, 185. Vinegar colchicum, 197. Colchicum and squill, 198. Horsebalm, 198. Colocynth, 198. Powdered foxglove, 218. Pills foxglove, 218. Infusion foxglove, 218. Elaterium, 222. Canada fleabane, 225. Gamboge, 249. Black hellebore, 261. Bacher's pills, 261. Compound wine black hellebore, 261. Jalap and cream of tartar, 289. Compound powder jalap, 290. Compound infusion juniper, 291. Preparations juniper, 291. Mustard whey, 295. Aromatic tincture lettuce, 297. Starkey's soap, 333. Embrocation petroleum, 349. Mixture petroleum, 349. Infusion parsley root, 349. Solution iodide potassium, 364. Scillitic acetate potassa, 372. Liquid acetate potassa, 373. Mixture acetate potassa, 373. Borotartrate potassa, 375. Nitrate potassa and squill, 380. Mixture nitrate potassa, 380. Cream of tartar, 382. Tartrate potassa and ammonia, 383. Decoction elder bark, 409. Confection scammony, 417. Squill, 418-22. Squill and nitrate potassa, 418. Squill and cream of tartar, 418. Infusion and decoction broom, 422. Acetate soda, 433. Solution carbonate soda, 434. Pills tobacco, 452. Wine tobacco, 452. Decoction dandelion, 455. Extract dandelion, 455. Electuary turpentine, 456. Veratria, 466. Sweet spirit of nitre, 91. Infusion of buchu, 150. Extract and decoction of cahinca, 153. Infusion of Peruvian bark, 188. Ethereal tincture of foxglove, 219. Digitalin, 220. Phosphate of iron, 240. Electuary of jalap, 290.

DROPSY, OVARIAN. Ointment iodine and mercury, 285. Solution iodide potass., 364.

DYSENTERY. Mixture nitric acid, 85. Starch jelly, 116. Starch and suet, 116. Angustura, 117. Clyster bistort, 148. Camphor water and laudanum, 159. Camphor water and nitric acid, 160. Cascarilla mixture, 173. Electuary catechu, 176. Infusion black purslane, 226. Mixture extract logwood, 260. Calomel and opium, 271. Decoction ipecacuanha, 288. Mixture ipecacuanha, 288. Emulsion jalap, 290. Electuary kino, 293. Mixture rhatany, 294. Milk and suet, 295. Mixture cherry-laurel, 299. Injection morphia, 320. Nux vomica, 327. Mixture nux vomica and aloes, 328. Oleaginous mixture, 331.

- Opium and acetate lead, 338. Opiate, 339. Plaster opium and camphor, 339. Opium and syrup poppies, 340. Pills of rhubarb and ipecacuanha, 395. Confection dogrose, 399. Syrup blackberries, 403. Compound powder sulphur, 449. Infusion slippery-elm, 462. Compound pills sulphate zinc, 474.
- DYSMENORRHEA.** Infusion southern wood, 130. Camphor mixture, 160. Compound pills ergot, 224. Clyster iodine, 285.
- DYSPEPSIA.** Wormwood, 75. Aloes pills, 94–96. Aniseed, 118. Orange peel and rhubarb, 135. Cayenne pepper pills, 167. Infusion cinnamon, 193. Powder coriander, 207. Pill prepared metallic iron, 228. Mixture malate iron, 237. Nitrate iron, 238. Mixture soot, 246. Infusion gentian and rhubarb, 251. Mixture gentian and sulphuric acid, 252. Masterwort, 262. Infusion hops, 264. Mercury, chalk, and ipecacuanha, 265. Bolus ipecacuanha, 287. Ipecacuanha and centaury, 287. Rice jelly, 346. Solution hydrargyro-iodide potass., 368. Sulphate potassa and rhubarb, 381. Rhubarb, 395, 398. Soda and rhubarb, 435. Mixture carbonate soda and quassia, 435. Compound turpentine plaster, 457. Oxide zinc and columbo, 470. Alkaline tincture of wormwood, 75. Lactic acid, 84. Alkaline wine of aloes, 97. Chamomile pills, 119. Subnitrate of bismuth, 148. Calamus, 153. Antacid mixture, 156. Aromatic powder, 170. Wine of tartrate of iron, 242. Magnesia and orange peel, 305.
- DYSPNŒA.** Opium and castor, 338.
- DYSURIA.** Benzoic acid and copaiba, 80. Tartar emetic and phosphate lime, 122. Infusion fleabane, 225. Pills nitrate potassa, 379. Mixture tobacco, 453.
- EAR, AFFECTIONS OF.** Pyroligneous acid, 79. Acoustic balsam, 141. Injection morphia, 320. Anti-otitic mixture, 339. Opium liniment, 344.
- ECCHYMOSIS.** Juniper liniment, 292. Lotion nitrate potassa, 380.
- ELEPHANTIASIS.** Mudar, 158.
- ENGORGEMENTS, VISCERAL.** Pills black oxide iron, 239. Electuary black oxide iron, 239. Calomel and dandelion, 272. Calomel and squill, 272. Sulphate mercury, 278. Compound powder rhubarb, 395. Tartrate soda and rhubarb, 439. Mixture tartrate soda, 440. Tartrate soda whey, 440. Compound infusion dandelion, 455.
- EPILEPSY.** Nitrate silver, 126–127. Belladonna, 144. Oil box, 151. Electuary Peruvian bark, 189. Muriate copper, 214. Ammoniated copper, 214. Ammoniated copper and belladonna, 214. Compound pills sulphate copper, 215. Ammoniated iron, 229. Prussian blue, 234. Wine galium, 248. Masterwort, 262. Calomel and opium, 273. Indigo, 283. Opium and nitrate silver, 338. Phosphorated oil, 350. Mixture biniodide potassium, 324. Compound powder valerian, 463. Oxide zinc, 470. Compound pills oxide zinc, 470. Ethereal tincture chloride zinc, 471. Compound pills sulphate zinc, 474.
- EPISTAXIS.** Alum, 100. Sulphate copper styptic, 216. Injection sulphate iron, 241.
- EXCORIATION.** Cold cream, 114, 179, 481. Carbonate lead ointment, 359. Plaster carbonate lead, 359. Ointment tannate lead, 361. Rose oil, 401. Camphor soap, 410. Ointment oxide zinc, 470. Cerate carbonate zinc, 473.
- EXANTHEMATA.** Compound mercurial powder, 264. Sulphuret magnesium, 308. Balm, 315. Mixture Virginia snakeroot and allspice, 429. Compound infusion elm bark, 461.
- EYE, AFFECTIONS OF.** Atropia, 135. Camphor ointment, 161. Camphor collyrium, 162.
- FACE, ERUPTIONS ON, BLOTCHES, &c.** Wash sal ammoniac, 109. Almond paste, 112. Almond powder, 112. Compound almond lotion, 113. Milk roses, 113, 147. Goulard's lotion, 357. Cosmetic liniment, 115. Water anemone, 116. Wash of benzoin, 147. Emulsion benzoin, 147. Spermaceti liniment, 180.

Sultana ointment, 180. Emulsion corrosive sublimate, 270. Cosmetic wash, 270-1. Lotion carbonate potassa, 376. Beef marrow soap, 410. Aromatic soap, 410. Cosmetic soap powder, 410. Essence of soap, 412. Lotion oxide of zinc, 470.

FAINTING. Aromatic vinegar, 78-79. Aromatic carbonate of ammonia, 105. Succinate ammonia, 110.

FEVER. Vinegar mixture, 77. Syrup vinegar, 78. Effervescing powders, 82, 88. Tartaric acid, 88. Sweet spirit of nitre, 91. Acetate ammonia, 105. Citrate ammonia, 107. Nitrate ammonia, 109. Tartar emetic, 122. Mixture extract centaury, 178. Boneset, 226. Compound powder ammoniated iron, 229. Mixture avens, 253. Calomel and jalap, 271. Calomel and nitrate potassa, 271. Dover's powder, 286. Ipecacuanha and tragacanth, 287. Infusion malt, 309. Citrate potassa, 378. Nitrate potassa, 379. Nitrous powders, 379. Mixture nitrate potassa, 380. Mixture sulphate potassa, 381. Infusion of sage, 408. Tamarinds, 453.

FEVER, HECTIC. Compound infusion boneset, 226. Gœlis's powder, 325. Compound powder sulphate quinia, 390. Compound infusion sage, 408.

FEVER, INTERMITTENT. Pills arsenic and opium, 80. Lockstadt's pills, 115. Golden sulphuret antimony, 121. Tartar emetic and quinine, 122. Electuary orange-peel, 135. Bebeerine, 144. Camphor water and ether, 160. Cetrarine, 181. Preparations Peruvian bark, 185, 191. Cinchonia, 191. Decoction coffee, 195. Dogwood, 207. Pills sulphate copper, 215. Sulphate copper and opium, 215. Persimmon, 220. Compound powder Prussian blue, 234. Horsechestnut, 262. Tulip-tree bark, 303. Magnolia, 309. Narcotina, 327. Phloridzine, 350. Phosphorated oil turpentine, 350. Piperine, 353. Mixture acetate potassa, 373. Arsenical solution, 373. Arseniate potassa, 374. Mixture citrate potassa and bark, 379. Oxalate potassa, 380. Quassia, 386. Oak bark, 386. Quinia and salts, 387-393. Salicin, 407. Willow bark, 407. Virginia snakeroot, 428. Common salt and lemon juice, 432. Arseniate soda, 433. Solution chlorinated soda, 437.

FEVER, TYPHUS. Carbonate ammonia, 106. Powder of oxide of gold, 139. Wild indigo, 142. Clyster camphor, 161. Liniment cantharides, 165. Tincture cayenne pepper, 167. Musk mixture, 324. Turpentine mixture, 332. Laudanum and tartar emetic, 345. Egg and brandy mixture, 346. Egg and wine mixture, 346. Mixture Virginia snakeroot and allspice, 429. Wine Virginia snakeroot and vanilla, 429. Tincture Virginia snakeroot and balsam Peru, 429. Mustard whey, 430. Chlorinated soda, 437.

FISSURES, ANUS. Ointment acetate lead, 357.

FISTULÆ. Injections copaiba, 205. Injection corrosive sublimate, 270. Injection myrrh, 326. Injection iodide potass., 364. Injection iodine, 367. Compound lotion iodide potass., 367.

FLATULENCE. Ammoniated alcohol, 107. Tincture angelica, 117. Aniseed, 118. Chamomile pills, 118. Tincture assafetida and soot, 134. Assafetida plaster, 134. Elixir orange-peel, 135. Tincture benzoin, 147. Calamus, 153. Aromatic powder and confection, 170. Compound tincture cardamom, 170. Spirit caraway, 171. Preparations cloves, 172. Cinnamon, 192. Coriander, 206. Elixir de Garus, 210. Oil of dittany, 212. Infusion of fennel, 245. Oil of partridge-berry, 250. Pennyroyal, 260. Henbane and ipecacuanha, 281. Lavender, 299. Magnesia and camphor, 307. Peppermint, 316. Nutmeg, 324. Confection opium, 338. Water allspice, 351. Electuary black pepper, 352. Compound tincture sassafras, 416. Purgative pastilles, 418. Mixture carbonate soda and gentian, 435. Compound turpentine plaster, 457. Ginger, 476.

FÆTOR ORIS. Mixture nitric acid, 85. Alum, 100. Lozenges chloride of lime, 154. Charcoal lozenges, 169. Lozenges catechu, 176. Pastilles catechu, 177. Creasote, 209.

- FROST, EFFECTS OF.** Muriatic acid lotion, 85. Sulphuric acid liniment, 87. Rust's ointment, 102. Oil elaterium, 222. Compound ointment henbane, 282. Embrocation petroleum, 349.
- FUNGOUS FLESH.** Burnt alum, 100. Chloride antimony, 119. Lotion orpiment, 130. Verdigris and savine, 212. Verdigris ointment, 213.
- GANGLIONIC SYSTEM, DISEASES OF.** Cataplasm of bryony, 150. Prussian blue, 234.
- GANGRENE.** Compound camphor pills, 159. Peruvian bark and arnica, 186. Cataplasm Peruvian bark, 190. Musk pills, 323. Anodyne ointment, 340. Cataplasm oak bark, 387.
- GASTRALGIA.** Mixture aconite, 90. Subnitrate bismuth, 148. Syrup codeia, 195.
- GASTRODYNIA.** Oxide silver, 126. Compound bismuth powder, 148. Compound tincture cinnamon, 193. Oil of érgot, 225. Mercurial mixture, 266.
- GLANDS ENLARGED.** Plaster ammoniac with mercury, 112. Cataplasm bryony, 150. Ointment chloride lime, 155. Animal charcoal, 168. Ointment of iodide of carbon, 170. Hemlock and dandelion, 202. Plaster foxglove, 219. Ointment iodide iron, 237. Compound galbanum plaster, 247. Compound mercurial liniment, 267. Ointment iodide mercury, 275. Red sulphuret mercury, 279. Liniment henbane, 282. Ethereal tincture iodine, 284. Mixture iodine and iodide potassium, 284. Sulphuret potassium, 369. Compound plaster sulphuret potassium, 371. Bin-iodide of quinia, 388. Burnt sponge, 442.
- GLEET.** Tannic acid, 87. Tincture cantharides and guaiacum, 164. Cubebs and ergot, 210. Compound powder ergot, 224. Pills guaiacum and turpentine, 258. Injection chloride zinc, 471.
- GLOTTIS, SPASM OF.** Cataplasm tobacco, 453.
- GOITRE.** Solution chloride calcium, 154. Animal charcoal, 168. Iodine ointment, 285. Cataplasm iodine, 285. Iodide potass., 364. Ointment iodide potass., 365. Liniment iodide potass., 366. Sulphuret potass., 369. Burnt sponge, 442.
- GONORRHOEA.** Injection of hydrocyanic acid, 83. Pills of tannic acid, 87. Alumina, 98. Alum, 101. Nitrate silver, 127. Assafetida and opium, 133. Oil box, 131. Pills chloride calcium, 154. Emulsion hemp, 163. Compound bolus catechu, 176. Electuary catechu, 176. Wine Peruvian bark and calamus, 191. Copaiba and its preparations, 204-6. Cubebs, 210-12. Injection ammoniated copper, 214. Injection sulphate copper, 215. Compound powder ammoniated iron, 229. Injection iodide iron, 236. Pills sulphate iron, 240. Tincture galls, 249. Oil guaiacum, 257. Mixture of Indian sarsaparilla, 262. Injection corrosive sublimate, 270. Calomel and catechu, 272. Saturated tincture iodine, 284. Injection opium, 340. Sedative injection, 348. Oil parsley, 349. Injection of chloro-platinate of sodium, 356. Acetate lead mixture, 358. Injection sulphuret potass., 371. Injection caustic potassa, 372. Pills nitrate potassa and camphor, 380. Nitrated emulsion, 380. Starkey's soap, 410. Turpentine and rhubarb, 450. Turpentine mixture, 457. Injection acetate zinc, 473. Fomentation sulphate zinc, 474. Injection sulphate zinc, 475.
- GOUT.** Aconite, 89. Ammoniacal liniment, 104. Carbonate ammonia, 106. Phosphate ammonia, 109. Ammoniacal succinic acid, 110. Golden sulphuret antimony, 120, 121. Sulphuret calcium, 155. Camphorated ether, 161. Portland powder, 178. Pills pipsissewa, 182. Colchicum, 196. Wine hedge hyssop, 257. Pills guaiacum and sulphur, 258. Ammoniacal tincture guaiacum, 259. Magnesia and colchicum, 306. Confection opium, 338. Phosphorated oil, 350. Mixture caustic potassa, 372. Solution silicate potassa, 381. Warner's cordial, 397. Extract elder berries, 409. Pills soap and ox gall, 411. Cor-

- serve broom, 423. Compound tincture senna, 426. Liniment sulphuret carbon, 451. Turpentine mixture, 457. Veratria, 466. Wine white hellebore, 467.
- GRAVEL. Mixture Prussian blue, 234. Carbonate potassa and ammonia, 377.
- GUMS, AFFECTIONS OF. Boracic acid, 81. Alumina, 99. Electuary alum, 101. Gargle alum, 101. Collutory catechu, 177. Collutory creasote, 209. Gargle sulphate zinc, 474.
- HEMATEMESIS. Calomel and acetate of lead, 272. Pills acetate lead, 356.
- HEMOPTYSIS. Powder starch, 116. Carrageen mixture, 184. Electuary Peruvian bark and catechu, 189. Creasote mixture, 208, 209. Bolus foxglove, 218. Mixture tincture foxglove, 219. Sedative mixture, 342. Compound pills acetate lead, 356. Tolu mixture, 459. Mixture uva ursi, 462.
- HEMORRHAGES. Gallic acid, 82. Alum, 100. Alum whey, 102. Bistort, 148. Alum and catechu, 176. Extract of Peruvian bark, 187. Sulphate copper, 215. Compound powder ergot, 224. Extract ergot, 224. Chloride iron and acetate lead, 232. Tincture chloride iron, 232. Solution sulphate iron, 241. Compound powder galls, 248. Avens, 253. Compound powder kino, 293. Mixture rhatany, 294. Alum whey, 295. Matico, 313. Roasted opium, 336. Opium and acetate lead, 338. Compound pills acetate lead, 356. Acetate lead, 356. Mixture bistort, 149. Hæmostatic powder, 393. Powder sulphate soda and opium, 439. Mixture uva ursi, 462. Eau de Pagliari, 482.
- HEMORRHOIDS. Aloetic mixture, 97. Alum ointment, 103. Lime ointment, 156. Charcoal suppository, 169. Extract Peruvian bark, 187. Hemlock oil, 202. Suppository hemlock, 203. Saffron ointment, 210. Oil elaterium, 222. Ointment galls, 249. Compound gall ointment, 249. Witch hazel, 260. Ointment mercury and belladonna, 267. Ointment red oxide mercury and tin, 278. Ointment henbane, 282. Electuary jalap, 290. Ointment matico, 314. Injection morphia, 320. Anodyne ointment, 339. Ointment opium and tar, 340. Laudanum ointment, 344. Confection black pepper, 352. Ward's paste, 352. Pile ointment, 357. Lead ointment, 360. Mixture tartrate potassa, 382. Mixture bitartrate potassa, 383. Ointment figwort, 423. Electuary senna and sulphur, 425. Ointment tin, 442. Ointment stramonium, 445. Sulphur electuary, 449. Fomentation sulphate zinc, 474.
- HAIR, LOSS OF, &c. Garlic, 93. Almond cream, 114. Macassar oil, 115. Fixature, 115, 216, 332. Hair dye, 127. Compound tincture cantharides, 164. Pomatum cantharides, 165, 166. Shampoo liquid, 167. Pomatum Peruvian bark, 190. Lotion galls, 248. Compound ointment balsam Peru, 142. Ointment of lavender, 300.
- HAIR, TO REMOVE. Orpiment powder, 130. Various depilatories, 482.
- HEART, AFFECTIONS OF. Syrup asparagus, 132. Bromine, 149. Foxglove and acetate lead, 219. Digitalin, 220. Mixture iodide potass., 364.
- HEMICRANIA. Pills of valerianate of quinia, 393. Compound powder valerian, 463. Electuary valerian, 463.
- HEMIPLEGIA. Pills poison oak, 460.
- HEPATITIS. Mixture nitric acid, 85. Bath nitromuriatic acid, 86. Compound mercurial pill, 266. Chlorate potassa, 378. Dandelion, 455.
- HERNIA. Clyster belladonna, 146. Infusion tobacco, 453.
- HERPES. Chlorine liniment, 183. Arseniate iron, 229. Ointment carburet iron, 232. Decoction soot, 245. Ointment black hellebore, 261. Liniment acetate mercury, 268. Compound calomel pill, 272. Calomel ointment, 273. Ointment red oxide mercury and lead, 278. Ointment sulphate mercury, 279. Red sulphuret mercury, 280. Ointment red sulphuret mercury, 280. Lotion sulphuret potassium, 370.

HICCOUGH. Ammoniated alcohol, 107. Assafetida, 133. Carminative mixture, 316.

HOOPING-COUGH. Succinated ammonia, 110. Extract anemone, 116. Kermes' mineral, 120. Antimonial wine and bittersweet, 124. Assafetida pills and mixture, 132, 133. Assafetida and tolu, 133. Assafetida plaster, 134. Belladonna, 144, 146. Fumigating powder, 147. Emulsion cantharides, 164. Tincture bark and cantharides, 190. Syrup Peruvian bark, 191. Cochineal and carbonate potass., 194. Confection ipecacuanha, 287. Mixture ipecacuanha, 288. Compound syrup ipecacuanha, 288. Alkaline wine ipecacuanha, 289. Syrup sulphuret magnesium, 308. Myrrh and zinc, 325. Lettsom's elixir, 344. Mixture carbonate potassa, 377. Compound syrup squill, 420. Carbonate soda and ipecacuanha, 434. Tincture artificial musk, 448. Liniment oil of amber, 448. Compound powder tobacco, 452. Mixture balsam tolu and copaiba, 459. Sulphate zinc, 474. Mixture sulphate zinc, 475. Oxide of zinc, 470. Ammoniated alcohol, 107. Syrup of codeia, 195. Hemlock plaster, 202. Copaiba mixture, 205. Pills of henbane, 281. Mixture of henbane and antimony, 281. Powder ipecacuanha and carbonate of soda, 286. Powder of musk, 323. Musk mixture, 324.

HYDROPHOBIA. Extract hemp, 162. Tonquin powder, 323.

HYDROCEPHALUS. Powder foxglove, 218. Calomel and foxglove, 271. Ointment iodide potass. and mercury, 365.

HYDROTHORAX. Tincture colchicum and foxglove, 197. Colchicum and elaterium, 198. Mixture colocynth, 200. Pills foxglove, 218. Mixture elaterium, 222. Gamboge mixture, 249. Extract lettuce, 297. Mixture tobacco, 453.

HYPOCHONDRIA. Assafetida, 132. Mixture lettuce and dandelion, 296.

HYPOPION. Pills seneka, 423.

HYSTERIA. Mixture acetic ether, 91. Whytt's pills, 96. Carbonate ammonia, 105. Succinate ammonia, 110. Mixture chamomile, 119. Infusion southern wood, 130. Assafetida and iron, 132. Assafetida mixture, 133. Alkaline tincture assafetida, 134. Tincture assafetida and soot, 134. Assafetida and castor, 134. Belladonna and camphor, 145. Bolus castor, 174. Castor pills, 174. Skunk cabbage, 220. Soot pills, 245. Compound galbanum pills, 247. Mixture lactucarium, 297. Anti-hysteric water, 316. Musk, 323. Pills opium and musk, 337. Tincture opium and asarabacca, 343. Mixture rue and squill, 403. Liniment oil of amber, 448. Compound powder valerian, 463. Vanilla, 465. Valerianate zinc, 475.

ILEUS. Hemlock oil, 202.

IMPETIGO. Lotion hydrocyanic acid, 83. Glycerine, 254. Compound powder sulphur, 449. Powder iodide sulphur, 452. Ointment oxide zinc and opium, 470.

IRITIS. Turpentine mixture, 332.

ISCHURIA. Assafetida and opium, 133.

ISSUES. Ointment of cantharides, 166. Elemi plaster, 223.

ITCH. Sulphuric acid ointment, 87. Lime-water and sulphur, 156. Lime ointment, 156. Chlorine ointment, 183. Lotion corrosive sublimate, 270. Ointment red oxide mercury and sulphur, 277. Tincture sulphuret potass., 370. Compound lotion sulphuret potass., 370. Ointment carbonate potassa, 377. Soap and sulphur, 412. Decoction stavesacre, 443. Compound powder sulphur, 449. Sulphur ointment, 450. Compound sulphur ointment, 450. Ointment white hellebore, 467. Sulphuret sodium, 432.

JAUNDICE. Terebinthinated ether, 93. Barberry, 148. Compound wine centaury, 178. Acetate of potassa, 373. Senna and guaiacum, 425.

JOINTS, AFFECTIONS OF. Pipsissewa beer, 182. Compound decoction guaiacum, 257. Pills guaiacum and antimony, 258. Ointment calomel and squill, 273. Oil laurel, 298.

KIDNEYS, AFFECTIONS OF. Emulsion manna, 311. Carbonate soda, 435. Ointment turpentine, 457. Uva ursi, 462.

LABOR. Contrayerva mixture, 203. Ergot and its preparations, 224-5. Decoction cotton root, 255. Compound powder borax, 433.

LARYNGITIS. Creasote pills, 208. Liniment croton oil 335. Plaster croton oil, 335.

LEUCORRHEA. Injection gallic acid, 82. Tannic acid, 87. Whytt's pills, 96. Alum, 101. Electuary catechu, 176. Peruvian bark and isinglass, 186. Wine Peruvian bark and calamus, 191. Cubebs and ergot, 210. Compound powder ergot, 224. Infusion black purslane, 226. Compound pills carbonate iron, 231. Lozenges iodide iron, 235. Bath iodide iron, 237. Ointment iodide iron, 237. Black oxide iron, 239. Electuary black oxide iron, 239. Sulphate iron, 240. Syrup sulphate iron, 241. Solution tartrate iron and potass., 242. Injection soot, 246. Tincture galls, 248. Avens, 253. Compound powder guaiacum, 258. Calomel and catechu, 272. Saturated tincture iodine, 284. Injection rhatany, 294. Injection subacetate lead, 357. Iodide potass., 364. Decoction oak bark, 387. Pills turpentine and guaiacum, 456.

LEPRA. Pills arsenic and opium, 80. Iodide arsenic, 129. Donovan's solution, 129. Barytic liniment, 143. Mudar, 159. Chlorine oil, 183. Mixture tincture colocynth, 200. Carburet iron, 232. Ethereal solution bibromide mercury, 269. Ointment naphthaline, 327.

LICHEN. Carburet iron, 232. Ointment glycerine, 254. Ointment calomel and camphor, 274. Ointment cyanuret mercury, 274.

LIPS, EXCORIATION OF. Spermaceti lip salve, 179.

LIVER, AFFECTIONS OF. Lotion hydrocyanic acid, 83. Muriatic acid pediluvium, 85. Terebinthinated ether, 93. Ointment tartar emetic, 123. Belladonna, 145. Blue pill and jalap, 266. Blue pill and quinia, 266. Compound mercurial pill, 266. Compound cathartic pills, 273. Calomel and jalap, 273, 289. Tartrate of potassa and rhubarb, 382. Mixture tartrate potassa, 382.

LUMBAGO. Camphor ointment, 162. Spirit turpentine mixture, 332.

LUNGS, AFFECTIONS OF. Syrup chamomile, 119. Tartar emetic, 122. Tartar emetic and opium, 122. Lotion tartar emetic, 123. Ointment tartar emetic, 123. Syrup antimonial wine, 124. Fig coffee, 243. Syrup asparagus, 132. Compound tincture benzoin, 147. Hemlock and ipecacuanha, 202. Compound pills henbane, 281. Ipecacuanha and antimony, 286. Myrrh and ipecacuanha, 325. Naphthaline, 327. Tar water, 354. Solution cyanuret potass., 363. Decoction and syrup seneka, 423. Storax, 445. Balsam tolu, 458. Colts-foot, 461.

MAMMÆ, AFFECTIONS OF. Ointment carbonate ammonia, 106. Fomentation sal-ammoniac, 108. Cataplasm sal-ammoniac, 109. Lime water and oil, 156. Charcoal, 168. Ointment turpentine, 457. Ointment oxide zinc and lycopodium, 471.

MANIA. Compound pills columbo, 200. Pills foxglove, 218.

MARASMUS. Jelly orange leaves, 136.

MEASLES. Infusion safflower, 171. Spermaceti mixture, 179.

MENORRHAGIA. Pills gallic acid, 82. Pills tannic acid and opium, 87. Alum, 100. Compound bolus catechu, 176. Peruvian bark and isinglass, 186.

- Ergot, 224-5. Infusion black purslane, 226. Pills ammoniated iron, 229. Injection sulphate iron, 241. Aromatic alum whey, 295.
- MERCURIAL DISEASE. Mixture sulphuret potass., 371. Sulphur mixture, 450.
- MILK, TO INCREASE SECRETION OF. Compound powder fennel, 245.
- MILK, TO RETARD SECRETION OF. Bolus acetate potassa, 372. Pills acetate soda, 433.
- MOUTH, AFFECTIONS OF. Vinegar gargle, 78. Muriatic acid gargle, 84. Collutory nitric acid, 85. Alum, 100. Chlorine gargle, 154, 183. Collutory balsam Peru, 141. Collutory carbonate potassa, 376. Boracic acid wash, 81. Sal-ammoniac, 109. Phosphate of lime, 158. Decoction of New Jersey tea, 177. Goldthread, 206. Creasote mixture, 209. Myrrh, 326.
- MOLLITIES OSSIUM. Phosphate lime, 157.
- MUSCLES, RIGIDITY OF. Ointment iodine and oil tobacco, 285.
- NAUSEA. Clove cataplasm, 172. Clove bag, 173. Pennyroyal, 260. Compound infusion mint, 317. Spice plaster, 477. Mixture carbonate potassa, 376.
- NEPHRITIS. Compound decoction oatmeal, 140. Infusion carrot seed, 171. Emulsion bicarbonate soda, 436. Compound ointment of turpentine, 457. Compound powder of uva ursi, 462.
- NEURALGIA. Pills of arsenic and opium, 80. Aconitia, 90. Tannate of quinia, 393. Aconite plaster and lotion, 89. Plaster belladonna, 146. Oil benzoin, 147. Emulsion cantharides, 164. Bolus castor, 174. Liniment chloroform, 184. Tincture colchicum, 196. Colchicum and camphor, 198. Creasote pills, 208. Carbonate iron, 230. Mixture tincture guaiacum, 259. Calomel and opium, 273. Compound pills henbane, 281. Lotion cherry-laurel, 299. Sulphate morphia, 322. Turpentine mixture, 332. Opiated turpentine liniment, 333. Carbonate lead ointment, 358. Lotion chloride lead, 359. Lotion cyanuret potassium, 363. Ointment cyanuret potass., 363. Solution chlorate potassa, 378. Extract cevadilla, 404. Extract tobacco, 452. Ointment tobacco, 453. Veratria, 466. Valerianate zinc, 475.
- NERVOUS DISORDERS. Camphor and musk, 159. Camphor water and Hoffmann's anodyne, 160. Peruvian bark and valerian, 186, 190. Pills ammoniated iron, 229. Infusion hops, 264. Compound powder henbane, 281. Musk, 323. Wild cherry bark, 384. Compound pills sulphate quinia, 391. Oxide zinc, 470. Cyanuret zinc, 472. Ferrocyanuret zinc, 472. Chloride of zinc, 471.
- NIPPLES, EXCORIATED. Oil elaterium, 222. Glycerine, 254. Mammillary lotion, 141. Opium and lime water, 340. Borax, 434.
- NODES, VENEREAL. Mercurial plaster, 267.
- NUX VOMICA, POISONING BY. Sulphuric ether and turpentine, 93.
- NYCTALOPIA. Calomel, antimony, and henbane, 271.
- ŒDEMA. Fomentation arnica, 128.
- ONYCHIA. Corrosive sublimate and zinc, 269. Red oxide mercury, 277.
- OPHTHALMIA. Alum collyrium, 101. Alum curd, 101. Acetate ammonia, 105. Antimonial wine, 123. Nitrate silver, 127. Ointment nitrate silver, 127. Chloride gold, 137. Chloride barium, 143. Sulphate cadmium, 152. Camphor collyrium, 162. Solution of conia, 203. Verdigris and alum, 213. Verdigris ointment, 213. Collyrium sulphate copper, 215. Decoction quince-seeds, 216. Eyebright, 227. Ointment Prussian blue, 234. Ointment red oxide of mercury and zinc, 278. Yellow root, 280. Wine opium, 342. Collyrium acetate lead, 357. Collyrium subacetate lead, 357. Ointment subacetate lead, 358. Collyrium biniodide potass., 367. Compound lotion iodide potass., 367. Collyrium caustic potass., 372. Sulphate of quinia and soda, 390. Ointment oxide

- zinc, 470. Ointment oxide of zinc and calomel, 471. Mixture ferrocyanuret zinc, 472. Collyrium acetate zinc, 473. Collyrium sulphate zinc, 474.
- OPHTHALMIA, GONORRHOEAL. Mercurial lotion, 265.
- OPIUM, POISONING BY. Vinegar coffee, 195.
- ORCHITIS. Mixture chloride barium, 143.
- OTORRHOEA. Sulphate of cadmium, 153.
- OZÆNA. Angelica, 117. Chloride lime, 155. Mercurial gargle, 265. Compound powder sulphate mercury, 278.
- PALPITATION. Syrup asparagus, 132. Powder foxglove, 218. Wild cherry bark, 384.
- PANNUS. Compound powder seneka, 423. Pills seneka, 423.
- PARALYSIS. Infusion horseradish, 127. Extract arnica, 128. Assafetida, 132. Brucia, 150. Nitrate camphor, 162. Cantharides and Cayenne pepper, 163. Oil of euphorbium, 227. Nux vomica, 328. Tincture nux vomica, 328. Pills opium and nitrate silver, 338. Phosphorated ether, 350. Lotion mustard, 430. Bath common salt and arnica, 431. Strychnia, 446. Iodate strychnia, 447.
- PERIPNEUMONIA. Golden sulphuret antimony, 120. Powder foxglove, 218. Pills calomel and antimony, 273.
- PERIOSTITIS. Compound mercurial ointment, 267.
- PHYMOSIS. Solution sulphate copper, 216.
- PHTHISIS. Mixture hydrocyanic acid, 82. Alum and benzoic acid, 101. Mixture chloride barium, 143. Belladonna, 145. Iodide calcium, 155. Iceland moss, 180. Carrageen, 184. Electuary Peruvian bark and sulphur, 189. Creasote, 208. Pills sulphate copper, 215. Foxglove, 217-19. Compound pills of henbane, 281. Ipecacuanha and nitrate potassa, 286. Artificial goat's milk, 295. Locatelli's balsam, 141. Myrrh and Canada balsam, 325. Compound extract myrrh, 326. Griffith's mixture, 326. Acetic turpentine liniment, 333. Acetate lead, 356-358. Mixture iodide potassium, 364. Chlorate potassa, 378. Acetate of morphia, 320. Wine of tar, 354. Wild cherry bark, 384.
- PLEURITIS. Infusion of pleurisy root, 131. Nitrate of camphor, 162. Mixture foxglove, 219. Cataplasm long pepper, 352.
- PNEUMONIA. Benzoic acid and ipecacuanha, 80. Mixture foxglove, 219. Powder ipecacuanha and muriate ammonia, 286. Powder ipecacuanha and antimony, 286.
- PORRIGO. Ointment nitric acid, 85. Lotion corrosive sublimate and copper, 270. Ointment calomel and acetate copper, 273. Ointment nitrate mercury and lead, 276. Ointment sulphate mercury, 279. Ointment oxide manganese, 310. Banyer's ointment, 360. Compound lotion sulphuret potass., 370. Powder iodide sulphur, 452.
- PRURIGO. Fomentation opium, 340. Ointment carbonate soda, 435. Powder iodide sulphur, 452.
- PRURITUS. Lotion carbonate potassa, 376. Lotion borax, 434. Lotion carbonate soda, 435. Lotion chlorate soda, 437.
- PSORIASIS. Arsenic and pepper pills, 80. Ointment nitric acid, 85. Ointment subnitrate bismuth, 148. Compound powder anthracokali, 169. Decoction soot, 245. Naphthaline ointment, 327. Ointment poke, 351. Tar ointment, 355. Compound lotion sulphuret potassium, 370. Lotion tobacco, 453. Ointment tobacco, 453.
- PSOROPHTHALMIA. Camphor ointment, 161. Ointment sulphate copper, 216. Ointment red oxide mercury, 277. Ointment red oxide mercury and cinnabar, 278. Ointment common salt, 432. Compound sulphur ointment, 451.

PYROSIS. Alkaline wine aloes, 97. Oxide of silver, 126. Charcoal and quassia, 169. Compound pills kino, 293. Mixture carbonate potassa, 376.

RACHITIS. Phosphate lime, 157. Black oxide iron, 239. Phosphate magnesia, 308. Mixture cod liver oil, 329. Acorn coffee, 387. Compound powder madder, 402.

RECTUM, DISEASES OF. Starch mucilage, 115. Suppository hemlock, 203. In jection copaiba, 205.

RHEUMATISM. Aconite, 89. Aletris, 93. Ammoniacal liniment, 104. Phosphate ammonia, 109. Golden sulphuret antimony, 120. Tartar emetic and opium, 122. Plaster tartar emetic, 123. Antimonial powder, 124. Burdock, 125. Extract arnica, 128. Ointment chloride gold, 137. Sulphuret lime, 155. Nitrate camphor, 162. Compound tincture cantharides, 164. Plaster castor, 175. Black snakeroot, 185. Colchicum, 196. Tincture colchicum flowers, 197. Creasote pills, 208. Compound decoction bittersweet, 221. Compound galbanum plaster, 247. Hedge hyssop, 257. Decoction guaiacum, 257. Oil guaiacum, 257. Compound powder guaiacum, 258. Guaiacum mixture, 258. Guaiacum and bittersweet, 258. Tincture guaiacum, 258. Plummer's pills, 272. Black sulphuret mercury, 279. Nervine balsam, 305. Magnesia and colchicum, 306. Tincture magnolia, 309. Oil of horsemint, 319. Turpentine mixture, 332. Opium and antimony, 337. Opium plaster, 339. Fomentation wine opium, 342. Opium liniment, 344. Plaster petroleum, 349. Phosphorated oil, 350. Plaster black pepper, 352. Pitch plaster, 353. Plaster subacetate lead, 358. Solution chlorate potassa, 378. Plaster pellitory, 385. Tincture cevadilla, 404. Camphorated essence soap, 412. Zittman's decoction, 414. Conserve broom, 423. Pills stramonium seed, 444. Lotion stramonium, 444. Bisulphuret carbon, 451. Turpentine liniment, 457. Poison oak, 460. Veratria, 466. Wine white hellebore, 467. Green hellebore, 468. Hydrosulphuretted bath, 83. Liniment of croton oil, 335. Tincture of poke, 350. Prickly ash, 469.

RINGWORM. Ointment cocculus indicus, 194. Vinegar borax, 434.

SALIVATION. Sulphuretted hydrogen, 83. Muriatic acid gargle, 84. Tannic acid, 87. Gargle sulphate copper, 215. Pills iodine, 284. Sulphuret potass. and cream of tartar, 369. Gargle borax, 434.

SCARLATINA. Mixture hyponitrous ether, 91. Acetate ammonia, 105. Plaster tartar emetic, 123. Belladonna, 145. Cayenne pepper gargle, 167. Chlorine mixture, 183. Myrrh gargle, 326. Mixture tartrate potassa, 382. Mixture cream of tartar, 383.

SCIATICA. Liniment succinated ammonia, 110. Plaster sulphuret antimony, 121. Oil turpentine and honey, 332. Oil turpentine mixture, 332.

SCHIRRHUS. Bolus sal ammoniac, 108. Extract marigold, 158. Charcoal, 168. Hemlock, 201. Mercury and antimony, 265. Ointment mercury and belladonna, 267.

SCROFULA. Hydriodate ammonia, 108. Chloride barium, 143. Iodide barium, 143. Baryta, 143. Compound cataplasm bryony, 150. Solution chloride calcium, 154. Chloride lime, 155. Animal charcoal, 168. Anthracokali, 169. Hemlock, 201. Hydriodic acid, 83. Mixture of sulphuret of antimony, 121. Pipsissewa beer, 182. Carrageen, 184. Soot, 245, 246. Burnt sponge, 442. Koechlin's drops, 214. Mixture bittersweet, 221. Compound powder metallic iron, 228. Ammoniated iron, 229. Bromide iron, 230. Compound pills carbonate iron, 231. Lozenges iodide iron, 235. Ammoniated tincture gentian, 252. Frost-weed, 260. Ointment mercury and belladonna, 267. Mercury and antimony, 264. Iodide mercury, 275. Ointment iodide mercury, 275. Binioidide mercury, 275. Black sulphuret mercury, 279. Iodine bath, 284. Mixture iodine, 285. Cod liver oil, 330. Iodide lead, 359. Ointment iodide lead, 360. Solution bromide potassium, 362. Ointment bromide potassium, 362. Ointment iodide potassium, 365. Mixture iodide potassium, 365. Bath sulphu-

- ret potassium, 371. Mixture sulphuret potassium, 371. Conserve acetate potassa, 373. Acorn coffee, 387. Hydriodate quinia, 388. Decoction madder, 402. Jauperand's decoction, 414. Compound infusion sassafras, 416. Compound ointment common salt, 431. Bath common salt and gelatine, 431. Carbonate soda and chamomile, 434. Sulphur electuary, 449.
- SCURVY. Wood sorrel, 77. Clauder's elixir, 97. Alumina, 99. Mixture horseradish, 128. Brooklime, 144. Scurvy grass, 194. Collutory lemon juice, 300. Mixture buckbean, 317. Chlorate potassa, 378.
- SEA SICKNESS. Plaster carbonate ammonia, 106.
- SLEEPLESSNESS. Camphor water and Hoffmann's anodyne, 160. Tincture hops, 264. Mixture lactucarium, 297. Acetate morphia, 320. Preparations opium, 336. Syrup poppies, 347. Jamaica dogwood, 353.
- SNAKE BITES. Ammonia and ether, 103. Eau de luce, 106.
- SORE THROAT, VENEREAL. Mercurial lotion, 265.
- SPASMODIC DISEASES. Sulphuric acid and ether, 87. Fuller's pills, 95. Succinated ammonia, 110. Nitrate silver, 126. Electuary orange leaves, 136. Camphor, 159. Ethereal tincture camphor, 160. Extract of hemp, 162. Castor, 174. Chloroform, 183. Mayweed, 208. Pills foxglove, 218. Tincture galbanum, 247. Calomel, antimony, and henbane, 271. Anti-hysterical water, 316. Musk, 323. Dippel's animal oil, 329. Opium and its preparations, 336. Succinated tincture opium, 343. Clyster opium and valerian, 344. Cyanide potass., 363. Mixture nitrate potassa, 380. Confection rue, 403. Extract stramonium, 444. Purified oil amber, 447. Artificial musk, 447. Eau de Luce, 448. Compound powder of tobacco, 452. Balsam amber, 448. Infusion linden, 458. Valerian, 463. Oxide zinc, 470. Cyanuret zinc, 472.
- SPERMATORRHEA. Compound powder cascarilla, 173. Nux vomica, 328. Mixture acetate lead, 358.
- SPINA VENTOSA. Bath sulphuret potass., 371.
- SPLEEN, DISEASES OF. Celandine, 181. Solution hydrargyro-iodide potass., 368. Tartrate of magnesia, 308.
- SPRAINS. Lotion sal-ammoniac, 108. Compound spirit of nutmeg, 325. Liniment opium, 339. Aromatic vinegar, 78.
- STOMACH, AFFECTIONS OF. Oxide of silver, 126. Subnitrate of bismuth, 148. Cayenne pepper pills, 167. Chlorine mixture, 183. Tincture chloride iron, 233. Mixture tincture guaiacum, 259. Aromatic plaster, 477. Compound powder sulphate quinia, 390. Mixture sulphuret carbon, 451.
- STRANGURY. Clyster laudanum, 344. Infusion parsley, 349. Mixture nitrate potassa, 380.
- SWEATING, PROFUSE. Mixture nitric acid, 85. Agaric, 149. Acetate lead, 356. Mixture acetate lead, 358.
- SYPHILIS. Mixture nitric acid, 85. Sulphuretted hydrosulphate ammonia, 108. Chloride silver, 125. Cyanuret silver, 125. Orpiment, 130. Preparations gold, 136-139. Compound powder anthracokali, 169. Hemlock and calomel, 202. Anthracokali, 169. Bittersweet, 221. Liniment of laudanum and lime water, 344. Sarsaparilla, 413-16. Syrup of Laffecteur, 415. Koechlin's drops, 214. Mercury and its preparations, 264-280. Opiated acetate mercury, 268, 338. Bichloride platinum, 355. Chloroplatinate soda, 356. Mixture biniodide potass., 368. Compound tincture carbonate potassa, 377. Chloride quinia and mercury, 389. Zittman's decoction, 414. Acetate of strychnia, 447.
- TAPEWORM. Chloride barium, 142. Kousoo, 149. Male fern, 244. Powder of hedge hyssop, 257. Compound powder gamboge, 249. Decoction pomegranate root, 256. Anthelmintic emulsion, 332. Mixture petroleum, 349. Compound powder cevadilla, 404. Oxide tin, 443.
- TEETH, DISEASES OF. Electuary of alumina, 99. Alum and ether, 100. Vicat's mixture, 107. Chloride lime, 154. Cantharides plaster, 166. Tincture of Cayenne pepper, 167. Oil cloves, 172. Plaster castor, 175. Chlorine oil, 183.

- Hemlock collutory, 202. Solution creasote, 208. Compound oil guaiacum, 257. Mixture guaiacum, 258-259. Mastich paste, 313. Turpentine lotion, 332. Anti-odontalgic mass, 313, 338. Balsam and drops for toothache, 339. Odontalgic drops, 342. Pellitory and preparations, 385. Willow bark, 408.
- TESTICLE, ENGORGED.** Compound mercurial ointment, 267. Ointment iodide potassium, 365.
- TETANUS.** Extract and mixture of hemp, 162-163. Mixture Dippel's animal oil, 329. Opium and cinnamon water, 341. Solution caustic potassa, 372.
- THROAT, AFFECTIONS OF.** Nitrate silver, 127. Cayenne pepper gargle, 167. Gargle Peruvian bark, 190. Liniment verdigris, 213. Gargle ammoniated copper, 214. Gargle figs, 243. Gargle galls, 248. Gargle pomegranate rind, 256. Muriate manganese, 310. Gargle myrrh, 326. Gargle nitrate potassa, 380. Gargle sulphate quinia, 392. Gargle sage, 408. Vinegar sage, 408. Vinegar elder flowers, 408. Opodeldoc, 411. Sulphate of zinc, 474. Vinegar gargle, 78. Alum gargle, 101. Cinnamon gargle, 193. Gargle of persimmon, 220. Gargle of cyanuret of mercury, 274.
- TINEA CAPITIS.** Lotion hydrocyanic acid, 83. Ammoniacal liniment, 104. Sulphuret antimony, 120. Ointment azedarach, 140. Liniment chloride lime, 155. Picrotoxin, 194. Compound soot ointment, 246. Ointment calomel and alum, 273. Ointment poke, 351. Ointment black pepper, 352. Tar ointment, 355. Ointment black pitch, 355. Ointment bromide potass., 362. Compound lotion sulphuret potass., 370. Ointment sulphuret potass., 371. Compound ointment common salt, 432. Lotion sulphuret sodium, 432. Ointment carbonate soda, 435. Compound sulphur ointment, 450. Ointment sulphur and zinc, 451. Sulphuret carbon, 451.
- TONSILS INFLAMED, &c.** Gargle alum, 100. Gargle iodine, 285. Ointment iodine, 285. Iodide of zinc, 473.
- TENESMUS.** Clyster, laudanum, 344. Pills rhubarb and ipecacuanha, 395.
- TRISMUS.** Musk mixture, 324.
- TUMORS AND SWELLINGS.** Lotion of sulphuric ether, 92. Hydriodate of ammonia, 108. Ammoniac plaster, 112. Ointment of chlorinated lime, 155. Plaster of ammoniac and mercury, 112. Gum plaster, 112. Ammoniacal liniment and tartar emetic, 123. Liniment of cantharides and camphor, 165. Hemlock plaster, 202. Hop cataplasm, 264. Camphorated mercurial ointment, 267. Compound mercurial liniment, 267. Pills of calomel and antimony, 273. Decoction of poppy heads, 347. Cataplasm of subacetate of lead, 357. Ointment of iodide potassium and morphia, 366. Opodeldoc, 411. Soap liniment, 411. Ointment of squill, 422. Stramonium ointment, 445. Ointment of iodide of zinc, 472.
- ULCERS.** Nitric acid, 85. Sulphate alumina, 99. Burnt alum, 100. Anglo-Saxon ointment, 102. Ointment oxide silver, 126. Ointment nitrate silver, 127. Chloride lime, 154. Ointment charcoal, 169. Charcoal cataplasm, 169. Chlorine liniment, 183. Peruvian bark and camphor, 186. Cerate and cataplasm Peruvian bark, 190. Hemlock infusion and ointment, 202. Verdigris lotion, 213. Metz's balsam, 213. Egyptian ointment, 213. Ammoniated copper, 214. Ointment ammoniated copper, 214. Ointment sulphate copper, 216. Ointment foxglove, 219. Ointment soot, 246. Ointment galls, 249. Compound oil guaiacum, 257. Ointment red oxide mercury, 277. Lotion henbane, 282. Infusion cherry-laurel, 298. Ceromel, 315. Mezereon ointment, 318. Ointment monesia, 319. Rob of mulberries, 319. Tincture balsam Peru, 141. Ointment balsam Peru, 142. Plaster balsam Peru, 142. Cod liver oil, 330. Ointment bichloride platinum, 355. Lotion acetate lead, 357. Goulard's lotion, 357. Ointment subacetate lead, 358. Compound powder carbonate lead, 358. Ointment carbonate lead, 358. Solution iodide potass., 364. Ointment biniodide potass., 368. Saviard's lotion, 372. Solution chlorate potassa, 378. Resin cerate, 393. Wine roses, 401. Ointment willow leaves, 408. Balsam sulphur, 450. Turpentine liniment, 457. Turpentine plaster, 457. Cataplasm slippery

- elm, 462. Lotion chloride zinc, 471. Plaster carbonate zinc, 474. Acetic acid cataplasm, 78. Pyroligneous acid cataplasm, 79. Sulphuric acid, 87. Ointment of cyanuret of silver, 126. Calamine cerate, 153. Carrot cataplasm, 171. Copaiba, 205. Creasote lotion, 209. Ointment of elemi, 223. Oil of soot, 246. Lotion of corrosive sublimate, 270. Yellow wash, 270. Lotion of biniodide of mercury, 276. Cerate of red sulphuret of mercury, 280. Mixture of iodine, 285. Arquebusade water, 317. Tincture of opoponax, 345. Chlorinated soda, 437. Ointment tobacco, 453.
- ULCERS VENEREAL.** Mercurial cerate and liniment, 267. Lotion corrosive sublimate, 270. Yellow wash, 270. Ointment cyanuret mercury, 274. Ointment red iodide mercury, 276. Black wash, 277. Compound tincture opoponax, 345. Ointment biniodide potass. and opium, 368.
- URETHRITIS.** Injection kino, 293. Injection subacetate lead and lime water, 358. Pills nitrate potassa, 379. Chloride of zinc, 471. Detersive injection, 97. Ergot injection, 224. Opium and sulphate of zinc, 338.
- URINARY ORGANS, AFFECTIONS OF.** Pipsissewa, 182. Buchu, 151. Clyster laudanum, 340. Pareira brava, 348. Embrocation petroleum, 349. Extract dandelion, 455. Turpentine pills, 456. Pills turpentine and magnesia, 456. Compound powder uva ursi, 462.
- URINE, INCONTINENCE OF.** Chloride gold and sodium, 138. Compound pills nux vomica, 328. Powder iodide sulphur, 452.
- URINE, RETENTION OF.** Sal-ammoniac, 108.
- UTERUS, DISEASES OF.** Alum, 101, 102. Animal charcoal, 168. Chlorine injection, 183. Syrup ergot, 224. Injection rhatany, 294. Ointment acetate morphia, 321. Mixture wine opium, 342. Belladonna ointment, 146.
- UVULA, RELAXATION OF.** Gargle galls, 248. Gargle oak bark, 387. Gargle sage, 408.
- VAGINA, DISEASES OF.** Injection copaiba, 205. Lotion of morphia and borax, 323. Injection opium, 340. Suppository chloride zinc, 471.
- VERMIN, TO DESTROY.** Ointment cocculus indicus, 194. Ointment fennel, 245. Ointment cevadilla, 404. Capuchin powder, 404. Vinegar stavesacre, 443. Ointment stavesacre, 443. Compound mercurial ointment, 267.
- VERTIGO.** Bolus castor, 174.
- VOMITING, TO CHECK.** Mixture catechu, 177. Compound pills columbo, 200. Mixture columbo, 200. Creasote mixture, 209. Clyster laudanum, 344. Aromatic plaster, 353. Anti-emetic mixture, 436. Carbonate potassa, 376.
- VOMITING, SPASMODIC.** Mercurial mixture, 266. Mixture ipecacuanha, 288. Compound pills of columbo, 200.
- WARTS.** Escharotic powder alum, 100. Ointment chloride antimony, 119. Verdigris ointment, 213. Verdigris plaster, 213. Corrosive sublimate and copper, 269. Savine and verdigris, 405. Savine ointment, 405.
- WEAKNESS OF BACK.** Lotion Peruvian bark, 190. Iron plaster, 231. Plaster red oxide iron, 239.
- WHITE SWELLING.** Ointment nitrate silver, 127. Chloride calcium, 154.
- WORMS.** Tincture and wine wormwood, 75. Ointment aloes, 98. Semen contra, 130. Santonine, 130. Assafetida and iron, 133. Azedarach, 140. Wormseed, 181. Cochineal and common salt, 194. Mixture copaiba, 205. Male fern, 244. Cabbage-tree bark, 253. Mercurial mucilage, 265. Calomel and gamboge, 271. Calomel and jalap, 271. Black sulphuret mercury, 279. Cowhage, 324. Dippel's animal oil, 329. Mixture olive oil, 330. Anthelmintic emulsion, 331. Castor oil and ether, 332. Emulsion opoponax, 345. Savine and pinkroot, 407. Anthelmintic emulsion, 427. Pinkroot, 440-41. Powder tin, 442. Sulphuret tin, 443. Tansy, 454. Turpentine and jalap, 456. Tincture and wine of wormwood, 75. Powder of sulphate of iron, 240. Decoction of pomegranate, 256. Calomel and pinkroot, 272. Vermifuge sugar, 406.

INDEX

OF

PHARMACEUTICAL AND BOTANICAL NAMES

- Abies canadensis*, 354
excelsa, 353
picea, 353
Abietis resina, 353
Absinthium, 75
Acacia, 76
arabica, 76
catechu, 175
senegal, 76
vera, 76
Acetosella, 77
Acetum, 77
Acidum aceticum, 78
empyreumaticum, 79
camphoratum, 79
arseniosum, 79
benzoicum, 80
boracicum, 81
carbonicum, 81
citricum, 81
gallicum, 82
hydrocyanicum, 82
hydriodicum, 83
hydrosulphuricum, 83
lacticum, 84
muriaticum, 84
nitricum, 85
nitro-muriaticum, 86
oxalicum, 86
phosphoricum, 86
sulphuricum, 86
tannicum, 87
tartaricum, 88
valerianicum, 88, 465
Aconitia, 90
Aconitum, 88
napellus, 88
Acorus calamus, 153
Adiantum, 90
capillus veneris, 90
pedatum, 90
Æsculus hippocastanum, 262
Æther aceticus, 91
hydrocyanicus, 92
hyponitrosus, 91
muriaticus, 92
sulphuricus, 92
terebinthinatus, 93
Agathotes chirayta, 182
Aletris, 93
farinosa, 93
Allium, 93
sativum, 93
Aloe, 93
socotrina, 93
Aloe spicata, 93
vulgaris, 93
Althæa, 98
officinalis, 98
Alumina, 99
Aluminæ acetas, 99
sulphas, 99
Alumen, 100
Ammonia, 103
Ammoniæ acetas, 104
arsenias, 105
carbonas, 105
bicarbonas, 107
citras, 107
hydriodas, 107
hydrosulphas, 108
urias, 108
nitras, 109
phosphas, 109
succinas, 109
sulphas, 110
Ammoniacum, 110
Amygdala, 112
amara, 112
dulcis, 112
Amygdalus communis, 112
Amylum, 115
Amyli iodidum, 116
Anacyclus pyrethrum, 385
Anamirta cocculus, 194
Anemone, 116
pratensis, 116
pulsatilla, 116
Angelica, 116
atropurpurea, 116
officinalis, 116
Angustura, 117
Anisum, 117
Anthemis, 118
nobilis, 118
cotula, 208
Anthracoali, 169
Antimonialis pulvis, 124
Antimonium, 119
Antimonii chloridum, 119
et potassæ tartras, 122
sulphas, 121
sulphuretum, 120
Apis mellifica, 314
Apocynum, 124
androsæmifolium, 124
cannabinum, 124
Aralia nudicaulis, 125
spinosa, 125
Arctium, 125

- Aretium lappa, 125
 Arctostaphylos uva ursi, 462
 Argentum, 125
 Argenti chloridum, 125
 cyanuretum, 125
 iodidum, 126
 oxidum, 126
 nitras, 126
 Aristolochia hirsuta, 428
 reticulata, 428
 serpentaria, 428
 Armoracia, 127
 Arnica, 128
 montana, 128
 nudicaule, 128
 Arsenicum, 129
 Arsenici iodidum, 129
 chloridum, 129
 et hydrargyri iodidum, 129
 tersulphuretum, 130
 Artanthe elongata, 313
 Artemisia, 130
 absinthium, 75
 abrotanum, 130
 contra, 130
 judaica, 130
 moxa, 130
 santonica, 130
 Arum, 131
 triphyllum, 131
 Asagraea officinalis, 404
 Asarum, 131
 canadense, 131
 europæum, 131
 Asclepias, 131
 incarnata, 131
 syriaca, 132
 tuberosa, 131
 Asparagus, 132
 officinalis, 132
 Aspidium filix mas, 243
 Assafoetida, 132
 Astragalus verus, 460
 Atropa belladonna, 144
 Atropia, 134
 Aurantium, 135
 Aurantii cortex, 135
 flores, 136
 folia, 136
 Aurum, 136
 ammoniatum, 139
 stanno-paratum, 139
 Auri chloridum, 137
 cyanidum, 138
 et sodii chloridum, 137
 iodidum, 139
 oxidum, 139
 Avena sativa, 140
 Azedarach, 140

 Ballota lanata, 141
 Balsamodendron myrrha, 325
 Balsamum Peruvianum, 141
 Balsamum toluatanum, 458
 Baptisia tinctoria, 142
 Barium, 142
 Barii chloridum, 142
 iodidum, 143
 sulphuretum, 143
 Barosma crenata, 150
 serratifolia, 150
 Baryta, 143
 Barytæ acetas, 143
 Bebeerina, 144
 Beccabunga, 144

 Belladonna, 144
 Benzoinum, 146
 Berberis, 148
 vulgaris, 148
 Bismuthum, 148
 Bismuthi subnitras, 148
 Bistorta, 148
 Boletus laricis, 149
 Brayera anthelmintica, 149
 Brominium, 149
 Brucia, 149
 Bruciæ acetas, 150
 murias, 150
 sulphas, 150
 Bryonia, 150
 alba, 150
 dioica, 150
 Buchu, 150
 Buxus, 151
 sempervirens, 151

 Cacao, 151
 Cadmium, 152
 Cadmii sulphas, 152
 Cahinea, 153
 Calamus, 153
 Calamina, 153
 Calcis carbonas, 156
 phosphas, 157
 Calcii chloridum, 154
 iodidum, 155
 oxidum, 155
 sulphuretum, 155
 Calcium, 154
 Calendula, 158
 officinalis, 158
 Calotropis, 158
 gigantea, 158
 Calx, 155
 chlorinata, 154
 Camphora, 159
 officinarum, 159
 Canella, 162
 alba, 162
 Cannabis, 162
 sativa, 162
 Cantharis, 163
 vesicatoria, 163
 Capsicum, 167
 annuum, 167
 baccatum, 167
 frutescens, 167
 Carbo animalis, 168
 ligni, 168
 mineralis, 169
 Carbonis iodidum, 170
 sulphuretum, 451
 Cardamine, 170
 pratensis, 170
 Cardamomum, 170
 Carota, 171
 Carthamus, 171
 tinctorius, 171
 Carum, 171
 carui, 171
 Caryophyllus, 172
 aromaticus, 172
 Cascarilla, 173
 Cassia acutifolia, 424
 elongata, 424
 fistula, 173
 marilandica, 174
 obovata, 424
 Castor fiber, 174
 Castoreum, 174

- Catalpa, 175
 cordifolia, 175
 Cataria, 175
 Catechu, 175
 Ceanothus, 177
 americanus, 177
 Centaurea benedicta, 177
 Centaurium, 178
 Cephaelis ipecacuanha, 286
 Cera, 178
 flava, 178
 alba, 178
 Cerasus serotina, 384
 Cetaceum, 179
 Cetraria, 180
 Chavica officinarum, 352
 Chelidonium, 181
 majus, 181
 Chenopodium, 181
 anthelminticum, 181
 Chimaphila, 181
 umbellata, 181
 Chiococca anguifuga, 153
 Chiretta, 182
 Chlorinum, 182
 Chloriformum, 183
 Chondrus, 184
 crispus, 184
 Chrysophyllum, 319
 Chicorium, 184
 intybus, 184
 Cimicifuga, 185
 racemosa, 185
 Cinchona, 185
 calisaya, 185
 condaminea, 185
 micrantha, 185
 Cinchonia, 191
 Cinnamomum, 192
 aromaticum, 192
 zeylanicum, 192
 Cissampelos pareira, 348
 Citrullus colocynthis, 198
 Citrus aurantium, 135
 limonum, 300
 Cocculus indicus, 194
 palmatus, 200
 Coccus, 194
 Cochlearia, 194
 armoracia, 127
 officinalis, 194
 Codeia, 195
 Coffea, 195
 arabica, 195
 Colchicum, 196
 autumnale, 196
 Collinsonia, 198
 canadensis, 198
 Collodium, 255
 Colocynthis, 198
 Colomba, 200
 Colophony, 393
 Comptonia, 201
 Conium, 201
 maculatum, 201
 Contrayerva, 203
 Convolvulus panduratus, 204
 scammonia, 416
 Copaiba, 204
 Copaisera officinalis, 204
 Coptis, 206
 trifolia, 206
 Coriandrum, 206
 sativum, 206
 Cornu, 207
 circinata, 207
 florida, 207
 sericea, 207
 Cotula, 208
 Creasotum, 208
 Creta præparata, 156
 Crocus, 209
 sativus, 209
 Croton eleutheria, 173
 lacciferum, 296
 tigilium, 333
 Cubeba, 210
 Cucumis, 212
 sativus, 212
 Cuminum cyminum, 216
 Cunila, 212
 mariana, 212
 Cuprum, 212
 ammoniatum, 214
 Cupri acetas, 212
 subacetas præparatum, 212
 murias, 213
 sulphas, 215
 Cydonium, 216
 Cydonia vulgaris, 216
 Cyminum, 216
 Cytisus scoparius, 422

 Daphne gnidium, 318
 mezereum, 318
 Datura stramonium, 444
 Daucus carota, 171
 Delphinium, 217
 consolida, 217
 staphisagria, 217, 443
 Delphinia, 217
 Dianthus, 217
 caryophyllus, 217
 Digitalis, 217
 purpurea, 217
 Digitalina, 219
 Diosma, 150
 Diospyros, 220
 virginiana, 220
 Diplolepis gallæ tinctoriæ, 248
 Dirca, 220
 palustris, 220
 Dorema ammoniacum, 110
 Dorstenia contrayerva, 203
 Dracontium, 220
 Drimys winteri, 469
 Dulcamara, 221

 Ecbalium elaterium, 222
 Elaterina, 222
 Elaterium, 222
 Elemi, 222
 Elettaria cardamomum, 170
 Emetia, 223
 Ergota, 223
 Ergotetia abortifaciens, 223
 Erigeron, 225
 annuum, 225
 canadense, 225
 heterophyllum, 225
 philadelphicum, 225
 strigosum, 225
 Eryngium, 225
 aquaticum, 225
 maritimum, 225
 Erythraea centaurium, 178
 Eugenia pimenta, 351
 Eupatorium, 225
 perfoliatum, 225

- Eupatorium purpureum, 225
 teucriifolium, 225
 Euphorbia, 226
 corollata, 226
 hypericifolia, 226
 ipeacacuanha, 226
 lathyris, 226
 Euphorbium, 226
 Euphrasia, 227
 officinalis, 227
 Ferrum, 227
 ammoniatum, 229
 Ferri acetas, 228
 ammonio-tartras, 241
 ammonio-citras, 229
 arsenias, 229
 bromidum, 230
 carbonas, 230
 carburetum, 231
 chloridum, 232
 citras, 233
 et potassæ tartras, 241
 ferrocyanuretum, 234
 filum, 227
 gallas, 234
 iodidum, 235
 lactas, 237
 malas, 237
 nitras, 238
 oxidum hydratum, 238
 nigrum, 238
 rubrum, 239
 phosphas, 240
 pulvis, 227
 ramenta, 227
 sulphas, 240
 sulphuretum, 241
 tannas, 243
 valerianas, 243
 Ferula assafoetida, 132
 tingitana, 110
 Ficus, 243
 carica, 243
 Filix mas, 243
 Foeniculum, 244
 vulgare, 244
 Fraxera, 245
 walteri, 245
 Fraxinus ornus, 311
 Fuligo, 245
 Fuligokali, 246
 Gadus morrhua, 329
 Galbanum, 247
 officinale, 247
 Galipea officinalis, 117
 Galium verum, 248
 Galla, 248
 Gambogia, 249
 Gaultheria, 250
 procumbens, 250
 Gentiana, 250
 lutea, 250
 Gentianin, 252
 Geranium, 252
 maculatum, 252
 Geoffroya inermis, 252
 Geum, 253
 rivale, 253
 urbanum, 253
 virginianum, 253
 Gillenia, 253
 trifoliata, 253
 stipulacea, 253
 Glycerina, 254
 Glycyrrhiza, 254
 glabra, 254
 Gossypium, 255
 herbaceum, 255
 Granatum, 256
 Gratiola, 257
 aurea, 257
 officinalis, 257
 Guaiacum, 257
 officinale, 257
 Hæmatoxylon, 259
 campechianum, 259
 Hamamelis, 260
 virginica, 260
 Hebradendron, 249
 Hedeoma, 260
 pulegioides, 260
 Helenium, 260
 autumnale, 260
 Helianthemum, 260
 canadense, 260
 Helleborus, 260
 niger, 260
 Helonias officinale, 404
 Hemidesmus, 261
 indicus, 261
 Hepatica, 262
 triloba, 262
 Heracleum, 262
 lanatum, 262
 Heuchera, 262
 americana, 262
 Hippocastanum, 262
 Hordeum, 263
 distichon, 263
 vulgare, 263
 Humulus, 263
 lupulus, 263, 304
 Hydrargyrum, 264
 ammoniatum, 268
 Hydrargyri acetas, 268
 boras, 268
 bromidum, 268
 chloridum corrosivum, 269
 mite, 271
 cyanuretum, 274
 et quiniæ chloridum, 274
 iodidum, 275
 rubrum, 275
 nitras, 276
 oxidum nigrum, 277
 rubrum, 277
 phosphas, 278
 sulphas, 278
 sulphuretum nigrum, 279
 rubrum, 279
 tartras, 280
 Hydrastis canadensis, 280
 Hyoscyamus, 280
 niger, 280
 Ichthyocolla, 282
 Illicium anisatum, 118
 Indigofera, 283
 Indigum, 283
 Inula, 283
 helenium, 283
 Iodinium, 284
 Ipecacuanha, 286
 Ipomœa jalapa, 289
 Iris florentina, 289
 Jalapa, 289

- Janipha manihot*, 454
Juglans cinerea, 291
Juniperus, 291
 communis, 291
 sabina, 404
 virginiana, 292

Kino, 292
Kousso, 149
Krameria, 293
 triandra, 293

Lac, 294
Lacca, 296
Lactuca, 296
 elongata, 296
Lactucarium, 297
Lappa minor, 125
Lauri baccæ, 298
 folia, 298
Lauro-cerasus, 298
Laurus, 298
 nobilis, 298
 sassafras, 416
Lavandula, 299
 vera, 299
Limon, 300
Linum, 302
 usitatissimum, 302
Liquor arsenici et hydrargyri iodidi, 129
Liriodendron, 303
 tulipifera, 303
Lobelia, 303
 inflata, 303
Lupulina, 304

Macis, 305
Magnesia, 305
Magnesiæ acetat, 306
 carbonas, 306
 citras, 307
 phosphas, 306
 sulphas, 307
 tartras, 308
 bi-tartras, 309
Magnesiæ sulphuretum, 308
Magnesium, 305
Magnolia, 309
 glaucæ, 309
Maltum, 309
Malva, 309
 sylvestris, 309
Manganesii carbonas, 309
 iodidum, 310
 urias, 310
 oxidum, 310
 phosphas, 311
 sulphas, 311
Manganesium, 309
Manna, 311
Mannite, 312
Maranta, 312
 arundinacea, 312
Marrubium, 312
 vulgare, 312
Maruta cotula, 208
Mastiche, 313
Matico, 313
Matriearia, 314
 chamomilla, 314
Mel, 314
Melaleuca cajuputi, 329
Melia azedarach, 140
Melissa, 315
 officinalis, 315

Mentha piperita, 316
 viridis, 316
Menyanthes, 317
 trifoliata, 317
Mezereum, 318
Monarda, 318
 punctata, 318
Monesia, 319
Mora, 319
Morphia, 319
Morphiæ acetat, 320
 bimeconas, 321
 citras, 321
 hydriodas, 321
 urias, 321
 nitras, 322
 phosphas, 322
 sulphas, 322
 tartras, 323
Morus nigra, 319
 rubra, 319
Moschus, 323
 moschiferus, 323
Mucuna, 324
 pruriens, 324
Myristica, 324
 moschata, 324
Myrospermum toluiferum, 458
 peruiferum, 141
Myrrha, 325

Naphthalina, 327
Narcotina, 327
Narcotinæ urias, 327
Narthex, 132
Nectandra rodæi, 144
Nepeta cataria, 175
Nephrodium filix mas, 243
Nicotiana tabacum, 452
Nux vomica, 327

Olea europæa, 330
Oleum animale empyreumaticum, 329
 cajuputi, 329
 jecoris aselli, 329
 morruæ, 329
 olivæ, 330
 ricini, 331
 terebinthinæ, 332
 tiglii, 333
Opium, 335
Opoponax, 345
 chironium, 345
Opuntia cochinillifera, 194
Origanum, 345
 vulgare, 345
Ornus europæa, 311
Oryza, 345
 sativa, 345
Ovum, 346
Oxalis acetosella, 77

Papaver, 347
 rhœas, 399
 somniferum, 335, 347
Pareira, 348
 brava, 348
Paullinia, 348
 sorbilis, 348
Petroleum, 349
Petroselinum, 349
 sativum, 349
Phloridzina, 350
Phosphorus, 350
Phytolacca, 350

- Phytolacca decandra, 350
 Physeter macrocephalus, 179
 Pieræna excelsa, 386
 Pimenta, 351
 Pimpinella anisum, 117
 Pinus, 332
 palustris, 456
 Piper, 351
 angustifolium, 313
 cubeba, 210
 longum, 352
 nigrum, 351
 Piperinum, 352
 Piscidia erythrina, 353
 Pistacia lentiscus, 313
 Pix burgundica, 353
 canadensis, 354
 liquida, 354
 nigra, 355
 Platini bichloridum, 355
 Platinum, 355
 Plumbi acetas, 356
 carbonas, 358
 chloridum, 359
 iodidum, 359
 oxidum semivitreum, 360
 rubrum, 360
 saccharas, 361
 tannas, 361
 Plumbum, 356
 Podophyllin, 361
 Podophyllum, 361
 peltatum, 361
 Polygala senega, 423
 Polygonum bistorta, 148
 Potassa, 371
 Potassæ acetas, 372
 arsenitis liquor, 373
 arsenias, 374
 bicarbonas, 377
 bisulphas, 381
 bitartras, 382
 loras, 374
 borotartras, 374
 carbonas, 375
 chloras, 378
 citras, 378
 et ammoniæ carbonas, 377
 sulphas, 382
 tartras, 383
 et magnesiæ sulphas, 382
 iodas, 379
 nitras, 379
 purum, 379
 oxalas, 380
 silicas, 381
 sulphas, 381
 sulphas cum sulphure, 382
 Potassæ tartras, 382
 Potassii biniodidum, 366
 bromidum, 362
 chloridum, 362
 cyanuretum, 363
 hydrargyro-iodidum, 368
 iodidum, 364
 super-iodidum, 366
 sulpho-cyanuretum, 369
 sulphuretum, 369
 Potassium, 362
 Potentilla tormentilla, 459
 Prinos, 384
 verticillatus, 384
 Prunum, 384
 Prunus domestica, 384
 lauro-cerasus, 298
 Prunus virginiana, 384
 Pterocarpus marsupium, 292
 santalinus, 410
 Punica granatum, 256
 Pyrethrum, 385
 Quassia, 386
 amara, 386
 excelsa, 386
 Quercus, 386
 alba, 386
 infectoria, 248
 pedunculata, 336
 robur, 386
 tinctoria, 386
 Quinia, 387
 Quiniæ acetas, 387
 arsenias, 387
 arsenis, 388
 citras, 388
 et ferri citras, 388
 iodidum, 388
 et hydrargyri chloridum, 388
 ferrocyanas, 388
 hydriodas, 388
 iodureta, 389
 kinas, 389
 lactas, 389
 urias, 389
 nitras, 390
 phosphas, 390
 sulphas, 390
 sulpho-tartras, 392
 tannas, 393
 tartras, 393
 valerianas, 393
 Resina, 393
 Rhamnus, 394
 catharticus, 394
 Rheum, 394
 Rhoas, 399
 Rhus toxicodendron, 469
 Ricinus communis, 331
 Rosa canina, 399
 centifolia, 399
 gallica, 400
 Rosmarinus, 401
 officinalis, 401
 Rubia, 402
 tinctorum, 402
 Rubus trivialis, 402
 villosus, 402
 Rumex, 402
 britannica, 402
 obtusifolius, 402
 Ruta, 403
 graveolens, 403
 Sabadilla, 404
 Sabbatia, 404
 angularis, 404
 Sabina, 404
 Saccharum, 406
 officinarum, 405
 Sagapenum, 406
 Sago, 407
 Salicinum, 407
 Salix, 407
 alba, 407
 Salvia, 408
 officinalis, 408
 Sambucus, 408
 canadensis, 408
 nigra, 408
 Sanguinaria, 409
 canadensis, 409

Santalum, 410
 Sapo, 410
 durus, 410
 mollis, 410
 vulgaris, 410
 Sarsaparilla, 413
 Sassafras, 416
 officinale, 416
 Scammonium, 416
 Scilla, 418
 maritima, 418
 Scoparius, 422
 Scrophularia nodosa, 423
 Senega, 423
 Senna, 424
 Serpentaria, 428
 Sesamum, 429
 indicum, 429
 orientale, 429
 Simaruba, 429
 officinalis, 429
 Sinapis, 430
 alba, 430
 nigra, 430
 Smilax, 413
 Soda, 432
 Sodæ acetas, 433
 arsenias, 433
 boras, 433
 carbonas, 434
 bicarbonas, 436
 chloras, 436
 chlorinata, 437
 nitras, 438
 et potassæ tartras, 439
 hyposulphis, 437
 murias, 431
 phosphas, 438
 sulphas, 438
 bisulphas, 439
 valerianas, 440
 Sodium, 431
 Sodii bromidum, 431
 chloridum, 431
 chloro-platinas, 356
 sulphuretum, 432
 Solanum dulcamara, 221
 Spigelia, 440
 marilandica, 440
 Spiræa, 441
 tomentosa, 441
 Spongia, 442
 Stanni chloridum, 442
 oxidum, 443
 sulphuretum, 443
 Stannum, 442
 Staphisagria, 443
 Statice, 443
 caroliniana, 443
 Stramonium, 444
 Strychnia, 445
 Strychniæ acetas, 446
 iodas, 447
 murias, 447
 nitras, 447
 sulphas, 447
 Strychnos nux vomica, 327
 Styrax, 445

Styrax benzoin, 146
 officinale, 445
 Succinum, 447
 Sulphur, 448
 Sulphuris carburetum, 451
 iodidum, 452
 Symplocarpus foetidus, 220
 Tabacum, 452
 Tamarindus, 453
 indica, 45
 Tanacetum, 454
 vulgare, 454
 Tapioca, 454
 Taraxacum, 454
 dens leonis, 454
 Terebinthina, 456
 Testa præparata, 156
 Theobroma cacao, 151
 Tilia Europæa, 457
 Tolutanum balsamum, 458
 Tormentilla, 459
 Toxicodendron, 460
 Tragacantha, 460
 Triosteum, 461
 perfoliatum, 461
 Tussilago, 461
 farfara, 461
 Ulmus campestris, 461
 fulva, 462
 Uva ursi, 462
 Valeriana, 463
 officinalis, 463
 Vanilla, 465
 aromatica, 465
 Veratria, 466
 Veratriæ murias, 467
 nitras, 467
 sulphas, 467
 Veratrum album, 467
 sabadilla, 404
 viride, 468
 Veronica anagallis, 144
 beccabunga, 144
 Viola, 468
 odorata, 468
 pedata, 468
 Wintera, 469
 Xanthorrhiza, 469
 apiifolia, 469
 Xanthoxylum, 469
 fraxineum, 469
 Zincum, 470
 Zinci acetas, 473
 carbonas, 473
 chloridum, 471
 cyanuretum, 472
 ferro-cyanuretum, 472
 iodidum, 472
 oxidum, 470
 sulphas, 474
 valerianas, 475
 Zingiber, 475
 officinale, 475

GENERAL INDEX.

A

Abbreviations in pharmaceutical formulæ, 43

Acetate of alumina, 99

— ammonia, 104

— baryta, 143

— brucia, 150

— copper, 212

— iron, 228

— lead, 356

— magnesia, 306

— mercury, 268

— morphia, 320

— potassa, 372

— quinia, 387

— soda, 433

— strychnia, 446

— zinc, 473

Acetic acid, 78-553

— cataplasm, 78

— clyster, 79

— diluted, 78

Acetic ether, 91

Acetic infusion of cantharides, 164

Acid, acetic, 78-553

— camphorated, 79

— arsenious, 79-553

— benzoic, 80

— boracic, 81-554

— carbonic, 81-554-560

— citric, 81-555

— gallic, 82

— hydriodic, 83

— hydrocyanic, 82-555

— hydrosulphuric, 83

— lactic, 84

— muriatic, 84-555

— nitric, 85-556

— nitro-muriatic, 86

— oxalic, 86-556

— phosphoric, 86

— polygalic, 424

— pyroligneous, 79

— sulphuric, 86-556

— tannic, 87

— tartaric, 88-557

— valerianic, 88-465

Acids, 64

Aconite, 88

— extract, 89

— alcoholic, 89

— ammoniated, 90

— pills of, 89

— liniment, 89

— mixture, 90

— ointment, 89

— ammoniated, 90

— plaster, 89

— powder, compound, 89

— saccharated powder, 406

— tincture of leaves, 89

Aconite, Fleming's tincture, 477

— tincture of root, 89

— wine, compound, 89

Aconitine, 90

— lotion, 90

— ointment, 90

Acorn coffee, 387

Acoustic balsam, 141-477

Adhesive plaster, 394

— Baynton's, 360

Administration of medicines, 58

Ærometers, 30

Affusion, 539

Agaric, white, 149

— pills, with opium, 149

— powder, 149

— with opium, 149

Air in convalescence, 74

Alcohol, 557

— ammoniated, 106

— aromatic, 106

— mixture, 107

— Vicat's anodyne, 107

— anisated, ammoniated, 107

— lotion, 107

— mixture, 107

Algaroth, powder of, 119

Alkaloids, 533-568

Allspice, 351

— bolus, 351

— essence, 351

— spirit, 351

— tincture, 351

— water, 351

Almonds, 112

— bandoline, 115

— bitter, 112

— butter, 112

— clyster, emollient, 114

— cold cream, 114

— confection, 113

— cream, 114

— emulsion, 113-114

— aromatic, 114

— jelly, 487

— linctus, green, 114

— white, 115

— liniment, cosmetic, 115

— Lockstadt's pills, 115

— lotion, compound, 113

— Goulard's, 114

— Macassar oil, 115

— milk anodyne, 113

— artificial, 113

— of roses, 113

— mixtures, 115

— oil, 114

— of bitter, 114

— paste, 112

- Almond powder, 112
 — soap, 410
 — sweet, 112
 — syrup of orgeat, 113
 — water of bitter, 114
 — diuretic, 114
 Aloes, 93
 — clyster, 97
 — decoction, compound, 97
 — electuary, 96
 — elixir, Boerhaave's, 98
 — Clauder's, 97
 — Garus's, 98
 — Stoughton's, 98
 — injection, 97
 — mixture, 97
 — alkaline, 97
 — ointment, 98
 — pills, 94
 — Anderson's, 95
 — antichlorotic, 96
 — aperient, 96
 — and rhubarb, 96
 — Barthez's, 96
 — Bicker's, 96
 — Chapman's, 94
 — compound, 94
 — Duchesne's, 96
 — Frank's, 96
 — Fuller's, 95
 — Griffith's, 96
 — Hooper's, 95
 — James's, 95
 — Lady Webster's, 95
 — Mitchell's, 95
 — Morrison's, 95
 — Peter's, 95
 — Pittschart's, 96
 — Speediman's, 95
 — splenetic, 95
 — Whytt's, 96
 — with assafetida, 94
 — blue mass, 96
 — iron, 94
 — myrrh, 94
 — powder, 94
 — and canella, 94, 162
 — compound, 94
 — emmenagogue, 94
 — suppository, 97
 — tincture, 97
 — and myrrh, 98
 — ethereal, 98
 — wine, 97
 — alkaline, 97
 — balsamic, 97
 Alum, 100
 — saccharine, 100
 — bolus, 100
 — burnt, 100
 — collyrium, 101
 — curd, 101
 — draught, 102
 — electuary, 101
 — errhine, 100
 — gargle, 100, 101
 — injection, 101, 102
 — julep, 102
 — liniment, 102
 — lotion, 102
 — ointment, 102
 — Anglo-Saxon, 102
 — chilblain, 102
 — pile, 103
 — Rust's, 102
 Alum, pills, 100
 — with benzoic acid, 101
 — powder, 100
 — root, 262
 — solution, 101
 — odontalgic, 100
 — whey, 102, 295
 — aromatic, 102, 295
 Alumina, 99
 — hydrate, 99
 — acetate, 99
 — astringent, Rust's, 99
 — electuary, 99
 — sulphate, 99
 — wash, detergent, 99
 Amber, 447
 — artificial musk, 447
 — emulsion, 448
 — tincture, 448
 — balsam, 448
 — eau de luce, 448
 — liniment, 448
 — mixture of oil, 448
 — oil, 447
 — rectified, 447
 — powder, fumigating, 447
 — tincture, 448
 — alkaline, 448
 — ethereal, 448
 — varnish, 486
 American centaury, 404
 — extract, 404
 — infusion, 404
 — tincture, 404
 — columbo, 245
 — infusion, 245
 — tincture, 245
 — hellebore, 468
 — extract, 468
 — mixture, 468
 — ointment, 468
 — pills, 468
 — tincture, 468
 — wine, 468
 — senna, 174
 — infusion, 174
 Ammonia, 103, 558
 — gargle, 103
 — Granville's lotion, 103
 — injection, 103
 — liniment, 103
 — terebinthinate, 104
 — compound, 103
 — sulphuretted, 104
 — lotion, 104
 — mixture, 103
 — and ether, 103
 — and muriate copper, 214
 — opodeldoc, 104
 — Steer's, 104
 — plaster, 104
 — solution, 103
 — spirit, 104
 — aromatic, 104
 — fetid, 104
 — sinapism, 103
 — water, 103
 — acetate, 104
 — cataplasm, 105
 — collyrium, 105
 — gargle, 105
 — mixture, 105
 — solution, 104
 — arseniate, 105
 — solution, 105

- Ammonia carbonate, 105
 aromatic, 105
 draught, 106
 with ginger, 106
 drops, 106
 liniment, 106
 mixture, 106
 ointment, 106
 Gondret's, 106
 pills, 106
 plaster, 106
 potion, 105
 tincture, compound, 106
 bicarbonate, 107
 chloride, and silver, 125
 pills of, 125
 citrate, 107
 solution, 107
 hydriodate, 107
 ointment, 108
 hydrosulphate, 108
 liniment, 108
 sulphuretted, 108
 syrup, 108
 muriate, 108
 bolus, 108
 cataplasm, 109
 collutory, 109
 draught, 109
 emulsion, 109
 fomentation, 108
 liniment, 109
 lotion, 108
 powder, 108
 wash, 109
 nitrate, 109
 mixture, 109
 phosphate, 109
 solution, 109
 succinate, 109
 impure, 110
 liniment, 110
 mixture, 110
 spirit, 110
 with ether, 110
 succinic acid, 110
 sulphate, 110
 Ammoniac, 110
 mixture, 110
 compound, 111
 with nitric acid, 111
 pills, 110
 compound, 111
 De Haen's, 111
 Klein's, 111
 with rhubarb, 111
 plaster, 112
 gum, 112
 with hemlock, 112
 mercury, 112
 Ammoniated alcohol, 106
 anisated, 107
 lotion, 107
 mixture, 107
 aromatic, 106
 mixture, 107
 Vicat's, 107
 gold, 139
 pills, 139
 iron, 229
 bolus, 229
 mixture, 229
 pills, 229
 powder, 229
 tincture, 229
 Ammonio-citrate of iron, 229
 tartrate of iron, 242
 solution, 243
 Amorphous quinine, 387
 Anatomical injections, 477
 preservative, 477 ZZZ
 arterial and venous, 478
 Anderson's pills, 95
 Anemone, 116
 collyrium, 116
 extract, 116
 pills, 116
 water, 116
 Angelica, 116
 conserva, 117
 essence, 117
 errhine mixture, 117
 tincture, 117
 compound, 117
 tree, 125
 water, 117
 Angustura, 117
 electuary, 117
 infusion, 117
 mixture, 117
 tincture, 117
 Animal charcoal, 168
 fats, 533
 oil, (Dippel's) 329
 Aniseed, 117
 essence, 118
 lozenges, 118
 mixture, 118
 oil, 118
 spirit, 118
 tincture, 118
 water, 118
 Antacids, 64
 Anthelmintic purgative, 272
 Anthelmintics, 65
 Anthracokali, 169
 powder, 169
 compound, 169
 sulphuretted, 170
 Antibilious pills, 199
 Antidote general, 553
 Anti-dysenteric opiate, 339
 Antilithics, 64
 Antimonial ethiops, 264
 powder, 124
 James's, 124
 pills, with calomel, 124
 wine, 123
 Antimony, 119, 558
 chloride, 119
 solution, 119
 nitro-muriatic oxide, 119
 chloride ointment, 119
 sulphuret, 120
 golden, 121
 powder, 121
 Kermes mineral, 126
 emulsion, 121
 lozenges, 120
 mixture, 121
 Kermesine powder, 120
 compound, 120
 lozenges, 120
 mixture, 121
 with lime, 121
 pills, 121
 plaster, 121
 ointment, 120
 pills, 120
 precipitated, 120

- Antimony sulphate, 121
 — tartrate and potassa, 122-558
 Anti-odontalgic mass, 338
 — paste, 313
 Antispasmodics, 64
 Apothecaries' measure, 24
 — weight, 20
 Apple water, 489
 Approximative measures, 25
 Aqua fortis, 85-556
 Arnica, 128
 — decoction, 128
 — extract, 128
 — fomentation, 128, 129
 — infusion, 128
 — — compound, 128
 — powder, 128
 — — compound, 128
 — tincture, 129
 Aromatic bath, 401
 — soap, 410
 — vinegar, 78-401
 Arrowroot, 312
 — beef tea, 312
 — milk, 312
 — pudding, 312
 — vanilla, 465
 — water, 312
 Arseniate of ammonia, 105
 — iron, 229
 — potassa, 374
 — quinia, 387
 — soda, 433
 — solution, 433
 Arsenic, 79-129-553
 — chloride, 129
 — solution, 129
 — iodide, 129
 — ointment, 129
 — pills, 129
 — solution with mercury, 129
 — draught, 129
 — tersulphuret, 130
 Arsenical solution, 373
 — soap, 411
 Arsenious acid, 79-553
 — cerate, 80
 — ointment, 80
 — pills, 80
 — with opium, 80
 — pepper, 80
 — powder, 80
 — compound, 80
 Arsenious acid solution, 80
 Arsenite of potassa, solution, 373
 — of quinia, 388
 Arteriotomy, 550
 Artificial asses' milk, 491
 — milk, 113
 — musk, 447
 — sulphuretted water, 83
 Asarabacca, 131
 — powder, compound, 131
 Asparagus, 132
 — decoction of roots, 132
 — extract of roots, 132
 — — of shoots, 132
 — mixture, 132
 — syrup of shoots, 132
 Assafetida, 132
 — Dewees's carminative, 134
 — enema, 134
 — mixture, 133
 — — compound, 134
 — — with oxymel squill, 133
 — — tolu, 133
 — pills, 132
 — with aloes, 134
 — iron, 132
 — lactucarium, 132
 — musk, 133
 — opium, 133
 — plaster, 134
 — syrup, 133
 — tincture, 133
 — alkaline, 134
 — ammoniated, 134
 — and castor, 134
 — and soot, 134
 Asses' milk, artificial, 491
 Atropine, 134
 — solution, 135
 — syrup, 135
 — tincture, 135
 — sulphate, 135
 Aurum musivum, 443
 Avens, 253
 — mixture, 253
 — powder, compound, 253
 — water-, 253
 — decoction, 253
 — white, 253
 Avoirdupois weight, 17
 Azedarach, 140
 — decoction, 140
 — ointment, 140

B

- Balm, 315
 — infusion, 315
 — spirit, compound, 315
 — tea, 315
 — water, 315
 — antihysteria, 316
 Balsam acoustic, 141, 477
 — amber, 448
 — cloves, aromatic, 172
 — Goulard's, 358
 — Locatelli's, 142
 — Metz's, 213
 — nervine, 305
 — of Peru, 141
 — collutory, 141
 — liniment, 141
 — lotion mamillary, 141
 Balsam of Peru, mixture, 141
 — ointment, 142
 — — compound, 142
 — pills, 141
 — plaster, 142
 — syrup, 141
 — tincture, 141
 — of tolu, 458
 — soap, camphorated acetic, 412
 — sulphur, 450
 — ethereal, 450
 — terebinthinated, 450
 — toothache, 339
 — Turlington's, 147
 Bandoline, 115, 216, 332
 Banyer's ointment, 360
 Barberry, 148

- Barberry infusion, 148
 lemonade, 148
 Barium, 142
 chloride, 142
 collyrium, 143
 mixture, 143
 pills, 142
 powders, 142
 solution, 142
 iodide, 143
 ointment, 143
 powder, 143
 sulphuret, 143
 Barley, 263
 decoction, 263
 compound, 263
 with nitrate of potassa, 263
 meal, 263
 sugar, 263
 mixture, 490
 Barlow's lotion, 370
 Barytes, 143, 559
 liniment, 143
 solution, 143
 acetate solution, 143
 Basilicon ointment, 393
 Bateman's drops, 343
 Baths, 535
 Bath, aromatic, 401
 cold, 535
 common salt and gelatine, 431
 cool, 536
 douche, 538
 foot, 537
 hip, 537
 hot, 537
 hydrosulphuretted, 83
 hyposulphite soda, 438
 iodide of iron, 237
 iodine, 284
 ioduretted, 367
 local, 537
 medicated, 539
 muriatic acid, 84
 nitro-muriatic acid, 86
 shower, 537
 sulphuret of potassium, 371
 sodium, 431
 temperate, 536
 tepid, 536
 vapor, 538
 warm, 536
 air, 538
 Baumé's hydrometer, 31
 table of agreement with sp.
 gr., 35
 Bay-tree, 298
 Baynton's plaster, 360
 Bearberry, 462
 Bebeerine, 144
 pills of sulphate, 144
 solution of sulphate, 144
 Beconi's soap, 411
 Beef essence, 488
 tea, 488
 arrowroot, 312
 Beef marrow soap, 410
 Beer, ginger, 476, 490
 molasses, 490
 Peruvian bark, 191
 pipsissewa, 182
 sarsaparilla, 416
 spruce, 490
 tar, 354
 Beer's divine stone, 213
 Belladonna, 144
 clyster, 146
 extract, 145
 alcoholic, 145
 solution, 145
 fumigation, 145
 infusion, 145
 liniment, 146
 mixture, 146
 ointment, 146
 opiated, 145
 pills, compound, 145
 and camphor, 145
 plaster, 146
 powder, 144
 and rhubarb, 145
 saccharated, 406
 syrup, 146
 Gillet's, 146
 tincture, 146
 ethereal, 146
 Blackett's, 479
 Benne, 429
 infusion, 429
 oil, 429
 Benzoic acid, 80
 and copaiba mixture, 80
 and ipecacuanha powder, 80
 mixture, 81
 Benzoin, 146
 balsam, Turlington's, 147
 emulsion, 147
 lard, benzoated, 146
 lotion, 147
 milk of roses, 147
 oil, 147
 pastilles, fumigating, 147
 powder, 147
 fumigating, 147
 tincture, 147
 compound, 147
 wash, 147
 Bestucheffe's tincture, 233
 Bibromide of mercury, 269
 Bicarbonate of ammonia, 107
 of potassa, 377
 of soda, 436
 Bichloride of platinum, 355
 Bichromate of potassa, 561
 Bimeconate of morphia, 321
 Biniodide of potassium, 366
 of quinia, 388
 Biscuit jelly, 487
 Bismuth, 148-559
 subnitrate, 148
 ointment, 148
 pills, 148
 powder, compound, 148
 Bistort, 148
 clyster, astringent, 148
 extract, 149
 infusion, 148
 mixture, 149
 Bisulphate of potassa, 381
 soda, 439
 and magnesia, 439
 Bitartrate of magnesia, 309
 of potassa, 382
 Bitter almonds, 112
 Bittersweet, 221
 decoction, 221
 compound, 221
 extract, 221
 pills, 221
 Bittersweet mixture, 221

- Bittersweet mixture and antimonial wine, 124
 — syrup, 221
 Black alder, 384
 — decoction, 384
 — drop, 341
 — Guy's Hospital, 341
 — Houlton's, 341
 — Lancaster, 341
 — Porter's, 341
 — Rousseau's, 341
 — draught, 426
 — hellebore, 260
 — extract, 261
 — vino-alcoholic, 261
 — pills, 261
 — ointment, 261
 — pills, compound, 261
 — tincture, 261
 — mixture, 261
 — wine, compound, 261
 — ink, 483
 — oak, 386
 — oxide of iron, 238
 — manganese, 310
 — mercury, 277
 — pepper, 351
 — cataplasm, 352
 — confection, 352
 — electuary, 352
 — fluid extract, 352
 — oil, resinous, 352
 — volatile, 352
 — ointment, 352
 — plaster, 352
 — Ward's paste, 352
 — pitch, 355
 — ointment, 355
 — pills, 355
 — plaster, 355
 — purslane, 226
 — infusion, 226
 — snakeroot, 185
 — decoction, 185
 — extract, 479
 — fluid, 479
 — tincture, 185
 — sulphuret of mercury, 279
 — wash, 277
 Blackberry, 402
 — decoction, 402
 — syrup, 403
 Blackett's tincture belladonna, 479
 Blacking, 479
 — for grates, 479
 Blancmange isinglass, 491
 — rice, 491
 Blessed thistle, 177
 — infusion, 177
 — wine, 177
 Blistering cloth, cantharides, 167
 — court plaster, 166
 Blisters, 70
 Blood-letting, 547
 — general, 548
 — topical, 550
 Bloodroot, 409
 — infusion, 409
 — pills, 409
 — syrup, 409
 — tincture, 409
 — compound, 409
 Blue ink, 483
 — pills, 266
 — and colocynth, 266
 — jalap, 266
 — and quinine, 266
 Blue pills and rhubarb, 266
 Boiled flour, 488
 Bolus, 515
 — acetate of morphia, 320
 — potassa, 372
 — allspice, 351
 — alum, 100
 — ammoniated iron, 229
 — burnt sponge, 442
 — castor, 174
 — catechu, 176
 — chloride of gold, 137
 — cinchonia, 192
 — copaiba, 204
 — croton oil, 333
 — foxglove, 218
 — ipecacuanha, 287
 — iron, metallic, 228
 — male fern, 244
 — muriate ammonia, 108
 — musk, 323
 — narcotine, 327
 — oxide of manganese, 273
 — Peruvian bark, extract of, 187
 — pomegranate root, 256
 — sal-ammoniac, 108
 — soap, 411
 — sulphate of morphia, 322
 — sulphuret of potassium, 369
 — tartar emetic, 122
 — valerian and iron, 463
 — sulphate of potassa, 463
 Boneset, 225
 — decoction, 226
 — infusion, 226
 — compound, 226
 Boot composition, 479
 Boracic acid, 81-554
 — collutory, 81
 — mixture, 81
 Borate mercury, 268
 — potassa, 374
 — soda, 433
 Borax, 433
 — collutory, 433
 — collyrium, 434
 — gargle, 434
 — honey, 434
 — lotion, 434
 — lozenges, 433
 — mixture, 434
 — ointment, 434
 — powder, compound, 433
 — vinegar, 434
 Boullay's filter, 522
 Bowman's root, 226
 Box, 151
 — oil, 151
 Bran tea, 489
 Brass, solder for, 485
 Brayera, 149
 Bread jelly, 487
 British oil, 349
 Bromide of iron, 230
 — mercury, 268
 — potassium, 362
 — sodium, 431
 Bromine, 149-559
 — lotion, 149
 — solution, 149
 — alcoholic, 149
 Brooklime, 144
 — decoction, 144
 — syrup, 144
 — water, 144
 Broom, 422

Broom, conserve, 423
 — decoction, 422
 — compound, 422
 — extract, 422
 — infusion, 422
 Broth, mutton, 489
 — vegetable, 487
 Brown mixture, 255-344
 Brucia, 149
 — acetate, 150
 — muriate, 150
 — pills, 150
 — solution, 150
 — sulphate, 150
 — tincture, 150
 Bryony, 150
 — cataplasm, 150
 — compound, 150
 — wine, 150
 Buchu, 150
 — extract, 151
 — fluid, 151
 — infusion, 150
 — compound, 151
 — tincture, 151
 Buckbean, 317

Buckbean extract, 317
 — infusion, 317
 — mixture, 317
 — diuretic, 318
 — pills, compound, 317
 Buckthorn, 394
 — extract, 394
 — syrup, 394
 Bug poison, 480
 Burdock, 125
 — decoction, 125
 — infusion, 125
 Burgundy pitch, 353
 — pills, 354
 — plaster, 353
 Burnt alum, 100
 — hartshorn, 157-207
 — sponge, 442
 — sugar, 406
 Butter of almonds, 112
 — cocoa, 152
 — mixture, 152
 — ointment, 152
 Butternut, 291
 — extract, 291
 Button snakeroot, 225

C

Cabbage-tree bark, 252
 — decoction, 253
 — extract, 253
 — mixture, 253
 Cachou aromatisé, 177
 Cadmium, 152
 — sulphate, 152
 — ointment, 153
 — solution, 152
 Caffein, citrate, 195
 — pills, 195
 — syrup, 196
 Cahinca, 153
 — decoction, 153
 — extract, 153
 Cajeput oil, 329
 — liniment, 329
 — mixture, 329
 Cakes, soda, 491
 Calamine, 153
 — prepared, 153
 — cerate, 153
 Calamus, 153
 — electuary, 153
 — infusion, 153
 — compound, 153
 — tincture, 153
 — compound, 153
 Calcium, 154
 — chloride, 154
 — cataplasm, 154
 — mixture, 154
 — pills, 154
 — solution, 154
 — iodide, 155
 — pills, 155
 — sulphuret, 155
 — liniment, 155
 — pills, compound, 155
 Calomel, 271
 — collyrium, dry, 272
 — electuary with, 273
 — pills, 272
 — and acetate of lead, 272
 — antimony, 273
 — catechu, 272

Calomel, pills, and colocynth, 273
 — dandelion, 272
 — guaiacum, 273
 — jalap, 273
 — iron, 272
 — opium, 273
 — quinine, 272
 — squill, 272
 — cathartic, compound, 273
 — compound, 272
 — ointment, 273
 — with acetate of copper, 273
 — alum, 273
 — camphor, 274
 — squill, 273
 — powder with antimony, 271
 — and henbane, 271
 — anthelmintic, 272
 — with foxglove, 271
 — gamboge, 271
 — jalap, 271
 — and rhubarb, 271
 — nitrate of potassa, 271
 — opium, 271
 — pink-root, 272
 Calves'-feet jelly, 488
 Camphor, 159
 — and chloroform, 161
 — ether, 161
 — Hoffmann's anodyne, 160
 — laudanum, 159
 — milk, 160
 — nitric acid, 160
 — clyster, 161
 — collyrium, 162
 — cough mixture, 162
 — essence, Ward's, 161
 — emulsion, 160
 — infusion, 160
 — liniment, 161
 — and turpentine, 162
 — vinegar, 161
 — compound, 161
 — ethereal, 161
 — mixture, 160

- Camphor mixture and myrrh, 160
 ——— Parrish's, 160
 ——— nitrate, 162
 ——— ointment, 161
 ——— pills, 159
 ——— and lactucarium, 159
 ——— musk, 159
 ——— tartar emetic, 122
 ——— compound, 159
 ——— powder, 159
 ——— compound, 159
 ——— tincture, 160
 ——— and saffron, 160
 ——— ethereal, 160
 ——— mixture, 161
 ——— water, 159
 ——— mixture, 161
 Camphorated acetic acid, 79
 ——— soap, 410
 ——— vinegar, 79
 Canada fleabane, 225
 ——— extract, 225
 ——— infusion, 225
 ——— pitch, 354
 Canella, 162
 ——— and aloes, powder, 162
 Cantharides, 163, 560
 ——— cerate, 165
 ——— cloth, blistering, 167
 ——— court-plaster, blistering, 166
 ——— decoction, 165
 ——— emulsion, 164
 ——— extract, 163
 ——— ethereal, 164
 ——— infusion, 163
 ——— acetic, 164
 ——— liniment, 165
 ——— aromatic, 166
 ——— and camphor, 165
 ——— soap, 165
 ——— stimulating, 166
 ——— oil, 165
 ——— ointment, 165
 ——— of extract, 165
 ——— issue, 166
 ——— pills with capsicum, 163
 ——— iron, 163
 ——— plaster, 166
 ——— camphorated, 166
 ——— compound, 166
 ——— odontalgic, 166
 ——— pomatum, 166
 ——— powder with camphor, 163
 ——— savine, 163
 ——— shampoo liquid, 167
 ——— tincture, 164
 ——— camphorated, 164
 ——— compound, 164
 ——— ethereal, 164
 ——— camphorated, 164
 ——— with guaiacum, 164
 ——— vinegar, 164
 Cantharidale, collodium, 166
 ——— ether, 166
 ——— taffeta, 166
 Capuchin powder, 404
 Caramel, 406
 Caraway, 171
 ——— embrocation, 172
 ——— essence, 172
 ——— infusion, 171
 ——— oil, 172
 ——— spirit, 171
 ——— water, 171
 Carbon, iodide, 170
 ——— ointment, 170
 Carbon, bisulphuret of, 451
 ——— drops, 451
 ——— liniment, 451
 ——— mixture, 451
 Carbonate of ammonia, 105
 ——— iron, 230
 ——— and manganese, 483
 ——— lead, 358
 ——— lime, 156
 ——— magnesia, 306
 ——— manganese, 309
 ——— potassa, 375
 ——— and ammonia, 377
 ——— soda, 434
 ——— zinc, 473
 Carbonic acid, 81-554-560
 ——— water, 81
 Carburet of iron, 231
 ——— sulphur, 451
 Cardamom, 170
 ——— confection, 170
 ——— extract, ethereal, 170
 ——— powder, aromatic, 170
 ——— tincture, 170
 ——— compound, 170
 Carminative, Dalby's, 306
 ——— Dewees's, 134-306
 ——— drops, 305
 Carrageen, 184
 ——— decoction, 184
 ——— jelly, 184
 ——— mixture, 184
 Carron oil, 156-303
 Carrot, 171
 ——— cataplasm, 171
 ——— extract, 171
 ——— infusion of seeds, 171
 ——— ointment, 171
 Cartier's hydrometer, 32
 Cascarella, 173
 ——— extract, 173
 ——— mixture, 173
 ——— infusion, 173
 ——— alkaline, 173
 ——— powder, compound, 173
 ——— tincture, 173
 ——— wine, compound, 173
 Cassia, purging, 173
 ——— confection, 174
 ——— mixture, 174
 ——— pulp, 173
 Castanella's powders, 487
 Castile soap, 410
 Castor oil, 331
 ——— bandoline, 332
 ——— clyster, 331
 ——— emulsion, 331
 ——— seeds, 331
 ——— mixture, 331
 ——— with ether, 332
 Castor, 174
 ——— bolus, 174
 ——— pills, 174
 ——— plaster, compound, 175
 ——— powder, compound, 174
 ——— saccharated, 406
 ——— spirit, compound, 175
 ——— tincture, 174
 ——— ammoniated, 175
 ——— compound, 175
 ——— ethereal, 174
 ——— with succinic acid, 174
 Catalpa, 175
 ——— decoction, 175
 Cataplasms, 541
 ——— acetate of lead, 357

- Cataplasm, acetate ammonia, 105
 — acetic acid, 78
 — black pepper, 352
 — bryony, 150
 — compound, 150
 — carrot-root, 171
 — cayenne pepper, 168
 — charcoal, 169
 — chloride of calcium, 154
 — chlorinated soda, 437
 — clove, 172
 — emollient, 302
 — flaxseed, 302
 — and bran, 302
 — garlic, 93
 — hemlock, 202
 — hops, 264
 — horseradish, 128
 — iodine, 285
 — lead, 360
 — long pepper, 352
 — mayweed, 208
 — mercurial, 267
 — mustard, 430
 — compound, 430
 — oak bark, 387
 — oatmeal, 140
 — Peruvian bark, 190
 — powder for, 140
 — pyroligneous acid, 79
 — sal ammoniac, 109
 — slippery elm, 462
 — soap, 412
 — soot, 246
 — subacetate of lead, 357
 — tobacco, 453
 — vinegar, 78
 Catechu, 175
 — bolus, 176
 — compound, 176
 — collutory, 177
 — electuary, 176
 — infusion, 176
 — compound, 176
 — injection, 177
 — lozenges, 176
 — mixture, anti-emetic, 177
 — with logwood, 177
 — pastilles, aromatic, 177
 — pills with alum, 176
 — liquorice, 176
 — powder, compound, 176
 — tincture, 176
 Cathartics, 65
 Catnep, 175
 — infusion, 175
 Caudle, 490
 Caustic chloride of gold, 137
 — potassa, 371
 — soda, 432
 Caustics, 71
 Cayenne pepper, 167
 — cataplasm, 168
 — cerate, 168
 — extract, 168
 — gargle, 167
 — infusion, 167
 — lotion, 168
 — lozenges, 167
 — pills, 167
 — syrup, 167
 — tincture, 167
 — with cantharides, 168
 — vinegar, 168
 Celandine, 181
 — extract, 181
 Celandine, mixture, 181
 — pills, 181
 Celsius's thermometer, 42
 Cements, 480
 — diamond, 283-480
 — for china, 296-480
 — for the laboratory, 480
 — shell lac, 296
 Centaury, 178
 — extract, 178
 — mixture, 178
 — pills, 178
 — Portland powder, 178
 — wine, compound, 178
 — American, 404
 Centigrade thermometer, 42
 Cerate, 544
 — acetate of lead, 357
 — arsenical, 80
 — carbonate of zinc, 473
 — cayenne pepper, 168
 — cherry-laurel, 299
 — croton oil, 335
 — eggs, 347
 — Goulard's, 358
 — Kirkland's, 484
 — lead, 360
 — mercurial, 267
 — compound, 267
 — opium, 340
 — oxide of zinc and lycopodium, 471
 — Peruvian bark, 190
 — phosphorated, 350
 — pitch, 354
 — red cedar, 292
 — red sulphuret of mercury, 280
 — resin, 393
 — compound, 394
 — savine, 405
 — simple, 178
 — soap, 411
 — spermaceti, 179
 — subacetate of lead, 358
 — sulphur, 451
 Ceromel, 315
 Ceruse, 358
 Cetrarine, 180
 — powder, 181
 Cevadilla, 404
 — capuchin, 404
 — extract, 404
 — ointment, 404
 — ointment, 404
 — powder, compound, 404
 — tincture, 404
 Chalk, 156
 — mixture, 157
 — prepared, 156
 Chalybeate water, artificial, 231
 Chamber utensils, 59
 Chamomile, 118
 — decoction, compound, 118
 — extract, 118
 — pills, compound, 119
 — infusion, 118
 — mixture, 119
 — syrup, 119
 — American, 208
 — German, 314
 Charcoal, animal, 168
 — ointment, 168
 — powder, 168
 — purified, 168
 — cataplasm, 169
 — electuary, 169
 — lozenges, 169

- Charcoal ointment, 169
 — powder, dentifrice, 169
 — and quassia, 169
 — suppository, 169
 — mineral, 169
 — wood, 168
 Cheltenham salts, 307
 Cherry-laurel, 298
 — cerate, 299
 — infusion, 298
 — ointment, 299
 — water, 298
 — lotion, 299
 — mixture, 299
 Chicken jelly, 490
 — panada, 489
 — water, 488
 Chinoidine, 387
 Chiretta, 182
 — infusion, 182
 — tincture, 182
 Chlorate of potassa, 378
 — soda, 436
 Chloride of antimony, 119
 — barium, 142
 — calcium, 154
 — gold, 137
 — and sodium, 137
 — iron, 232
 — lead, 359
 — lime, 154
 — potassium, 362
 — quinia and mercury, 274, 389
 — pills, 274, 389
 — silver, 125
 — sodium, 431
 — tin, 442
 — zinc, 471
 Chlorinated lime, 154
 — soda, 437
 Chlorine, 182, 560
 — clyster, 183
 — collutory, 183
 — gargle, 183
 — injection, 183
 — liniment, 183
 — mixture, 183
 — oil, 183
 — ointment, 183
 — water, 183
 Chloroform, 183
 — emulsion, 184
 — liniment, 184
 Chloroplatinate of sodium, 356
 — injection of, 356
 Chocolate, 490
 — aromatic, 152
 — cream, 152
 — Iceland moss, 180
 — milk, 490
 — paullinia, 348
 — purgative, 418
 — salep, 152
 — vanilla, 152
 — white, 152
 Chrome, 561
 Cider mixture, 291
 Cinchonia, 191
 — bolus, 192
 — kinate, 192
 — muriate, 192
 — pills, 191
 — sulphate, 192
 — pills, 192
 — mixture, 192
 — syrup, 192
 Cinchonia tincture, 192
 — wine, 192
 Cinnabar, 279
 Cinnamon, 192
 — electuary, compound, 193
 — essence, 193
 — gargle, 193
 — infusion, 193
 — lozenges, 192
 — oil, 194
 — plaster, spiced, 194
 — powder, compound, 192
 — saccharated, 406
 — spirit, 193
 — tincture, 193
 — compound, 193
 — ethereal, 193
 — vinegar, compound, 193
 — water, 193
 — compound, 193
 — wine, compound, 193
 Citrate of ammonia, 107
 — solution, 107
 — caffein, 195
 — iron, 233
 — and quinia, 233
 — and magnesia, 234
 — magnesia, 307
 — morphia, 321
 — potassa, 378
 — quinia, 388
 Citrated effervescing powders, 82
 Citric acid, 81, 555
 — lozenges, 81
 — syrup, 81, 82
 Citrine ointment, 276
 Cleanliness in sick room, 57
 Cloth, waxed, 178
 Cloves, 172
 — bag, 173
 — balsam, 172
 — cataplasm, 172
 — infusion, 172
 — mixture, odontalgic, 172
 — oil, 172
 — plaster, 172
 — saccharated powder, 406
 — spirit, 172
 — tincture, 172
 — water, 172
 — wine, 172
 Clysters, 66
 — acetate of morphia, 321
 — acetic acid, 79
 — anthelmintic, of aloes, 97
 — assafetida, 134
 — belladonna, 146
 — bistort, 148
 — camphor, 161
 — castor oil, 331
 — chlorine, 183
 — colocynth, 199
 — common salt, 431
 — and arnica, 431
 — copaiba, 206
 — cubebs, 211
 — dandelion, 456
 — egg, 346
 — emollient, oil of almonds, 114
 — malt, 309
 — ergot, 224
 — flaxseed oil, 363
 — iodine, 285
 — laudanum, 344
 — and valerian, 344
 — musk, 324

- Clyster, olive oil, 330
 — opium, 340
 — Peruvian bark, 190
 — rhatany, 294
 — soap, 412
 — southernwood, 130
 — stimulating, with nutmeg, 325
 — sulphate of magnesia, 308
 — quinia, 392
 — soda, 439
 — tartar emetic, 123
 — tartarized soda, 440
 — turpentine, 456
 — oil, 333
 — wormwood, 75
 Coating pills, 485, 516
 Cocculus indicus, 194
 — ointment, 194
 — picrotoxin, 194
 Cochineal, 194
 — tincture, 194
 — with carbonate of potassa, 194
 — common salt, 194
 Cocoa, 151
 — butter, 152
 — ointment, 152
 — mixture, 152
 — chocolates, 152
 — aromatic, 152
 — cream, 152
 — salep, 152
 — vanilla, 152
 — white, 152
 — powders, compound, 15
 — palamoud, 151
 — racahout, 151
 — wakaka, 152
 Codeia, 195
 — and muriate of morphia, 195
 — syrup, 195
 Cod liver oil, 329
 — compound, 330
 — liniment, 330
 — mixture, 329
 — ointment, 330
 — compound, 330
 — soap, 412
 — ioduretted, 412
 — syrup, 330
 Coffee, 195
 — decoction, 195
 — figs, 243
 — milk, 490
 — syrup, 195
 — vinegar, 195
 — acorn, 387
 Colchicum, 196
 — extract bulb, 196
 — acetic, 196
 — liniment, with camphor, 198
 — mixture, 197
 — with ammonia, 198
 — elaterium, 198
 — magnesia, 197
 — squill, 198
 — sulphate magnesia, 197
 — oxymel, 197
 — pills, 196
 — compound, 196
 — syrup, 197
 — tincture, flowers, 197
 — seed, 196
 — ethereal, 196
 — and foxglove, 197
 — compound, 197
 — mixture, 197
 Colchicum, vinegar, root, 197
 — seed, 197
 — mixture, 198
 — wine, compound, 197
 — mixture, 198
 — root, 196
 — seed, 196
 Cold bath, 535
 — cream, 114, 179, 481
 — without spermaceti, 179
 — custard, 489
 Collodion, 255
 Collutory, balsam of Peru, 141
 — boracic acid, 81
 — borax, 433
 — carbonate of potassa, 376
 — catechu, 177
 — chlorinated lime, 154
 — chlorine, 183
 — creasote, 209
 — hemlock, 202
 — lemon juice, 300
 — mastich, aromatic, 313
 — myrrh, 326
 — nitric acid, 85
 — pyroligneous acid, 79
 — rhatany, 294
 — rose-water, 400
 — sal-ammoniac, 109
 — soot, 246
 Collyrium, acetate ammonia, 105
 — of lead, 356
 — zinc, 473
 — alum, 101
 — anemone, 116
 — anodyne, 340
 — antimonial wine, 123
 — ioduretted potassium, 367
 — borax, 434
 — calomel, dry, 272
 — camphor, 162
 — carbonate of potassa, 375
 — caustic potassa, 372
 — chloride of barium, 143
 — gold, 137
 — corrosive sublimate, 270
 — iodide of zinc, 472
 — nitrate of silver, 127
 — saffron, 210
 — strychnia, 446
 — subacetate of lead, 357
 — sulphate of copper, 215
 — zinc, 474
 — and camphor, 474
 — wine of opium, 342
 Colocynth, 198
 — clyster, 199
 — extract, 199
 — compound, 199
 — mixture, 199
 — pills, and henbane, 199
 — compound, 199
 — powder, 199
 — tincture, 200
 — mixture, 200
 Cologne water, 301
 Colors for show-bottles, 481
 Colored fires, 482
 Columbo, 200
 — decoction, compound, 201
 — extract, 201
 — infusion, 200
 — and ginger, 200
 — rhubarb, 200
 — mixture, 200
 — and cascarrilla, 201

- Columbo, mixture, and salep, 201
 — pills, compound, 200
 — powder and iron, 200
 — magnesia, 200
 — tartrate of iron, 200
 — tincture, 201
 — American, 245
 Coltsfoot, 461
 — decoction, 461
 — compound, 461
 — syrup, 461
 Common salt, 431
 — bath, with gelatine, 431
 — clyster, 431
 — with arnica, 431
 — fomentation, 431
 — mixture with lemon juice, 432
 — ointment, 432
 — compound, 432
 — powder, compound, 431
 — with cochineal, 431
 Confections, 518
 Confection acorns, 387
 — almonds, 113
 — aromatic, 170
 — black pepper, 352
 — cassia, 174
 — copaiba, 204
 — dog rose, 399
 — ipecacuanha, 287
 — opium, 338
 — orange flowers, 136
 — peel, 135
 — roses, 400
 — rue, 403
 — scammony, 417
 — senna, 425
 — compound, 425
 Conia, solution, 203
 Conserves, 518
 Conserve of acetate of potassa, 373
 — angelica, 117
 — broom, 423
 — violets, 468
 Contrayerva, 203
 — decoction, 203
 — extract, 203
 — gargle, 203
 — mixture, 203
 — powder, compound, 203
 — tincture, 203
 — compound, 203
 Copahine-Mège, 205
 Cool bath, 536
 Copaiba, 204
 — boluses, 204
 — clyster, 206
 — confection, 204
 — emulsion, 205
 — injection, 205
 — mixture, 80, 205
 — oil, 206
 — paste, 206
 — pills, 204
 — and cubebs, 204
 — cubebs and turpentine, 204
 — tincture, 206
 — alkaline, 206
 — compound, 206
 Copal varnish, 486
 Copper, 212, 561
 — solder for, 481
 — acetate, 212
 — ammoniated, 214
 — gargle, 214
 — injection, 214
 — Copper, ammoniated, ointment, 214
 — pills, 214
 — compound, 214
 — powder, with belladonna, 214
 — solution, 214
 — muriate, 213
 — and ammonia, 214
 — cupreous ether, 214
 — Koechlin's drops, 214
 — solution, with mercury, 214
 — subacetate, 212
 — balsam, Metz's, 213
 — liniment, 213
 — lotion, compound, 213
 — ointment, 213
 — Egyptian, 213
 — pills, 213
 — with alum, 213
 — plaster, 213
 — powder, with savine, 212
 — wash, ophthalmic, 213
 — sulphate, 215
 — collyrium, 215
 — electuary, 215
 — gargle, 215
 — lotion, 215
 — injection, 216
 — ointment, 216
 — pills, 215
 — and opium, 215
 — compound, 215
 — powder, compound, 215
 — styptic, 216
 Cordial, Godfrey's, 343
 — Warner's, 397
 Coriander, 206
 — powder, compound, 206
 — tincture, 207
 — water, compound, 207
 Corpse, preservation of, 478
 Corrosive sublimate, 269
 — collyrium, 270
 — injection, 270
 — lotion, 270
 — Bateman's, 270
 — cosmetic, 270
 — with camphor, 270
 — copper, 270
 — mixture, 269
 — pills, 269
 — compound, 269
 — powder with copper, 269
 — zinc, 269
 — solution, 269
 — alcoholic, 269
 — tincture, antacid, 270
 — wash, cosmetic, 271
 — yellow, 270
 Cotton, 255
 — decoction of root, 255
 Court plaster, 282
 Cowhage, 324
 — electuary, 324
 — ointment, 324
 Coxe's hive syrup, 420-424
 Cranesbill, 252
 — decoction, 252
 — extract, 252
 Cream, almond, 114
 — chocolate, 152
 — cold, 114-179-481
 — without spermaceti, 179
 — of tartar, 382
 — soluble, 374
 Creasote, 208

Creasote, collutory, 209
 — lotion, 209
 — mixture, 208, 209
 — ointment, 209
 — compound, 209
 — pills, 208
 — solution, alcoholic, 208
 Croton oil, 333
 — bolus, 333
 — cerate, 335
 — embrocation, 335
 — emulsion, 334
 — liniment, 335
 — lozenges, 334
 — mixture, 334
 — ointment, 335
 — pills, 333
 — compound, 334
 — with blue mass, 334
 — quinine, 334
 — plaster, 335
 — soap, 334
 — solution, saponaceous, 334
 — tincture, 334
 Cubebs, 210
 — clyster, 211
 — electuary, 211
 — fluid extract, 211-481

Cubebs, extract, emulsion, 211
 — alcoholic ethereal, 211
 — lozenges, 211
 — alcoholic ethereal extract, 211
 — mixture, 211
 — oil and copaiba, 212
 — oil, 212
 — powder, 210
 — and alum, 210
 — ergot, 210
 — hemlock, 210
 — syrup, alco-ether. extract, 212
 — tincture, 211
 Cubic nitre, 438
 Cuckoo flower, 170
 Cucumber, 212
 — ointment, 212
 Cumin, 216
 — plaster, 216
 Cupping, 550
 Curd, alum, 101
 Custard, cold, 489
 — rice, 490
 Cyanuret of gold, 138
 — mercury, 274
 — potassium, 363
 — silver, 125
 — zinc, 472

D

Dalby's carminative, 306
 Dandelion, 454
 — clyster, 456
 — decoction, 455
 — extract, 455
 — fluid, 455
 — and senna, 455
 — pills, 455
 — with blue mass, 456
 — infusion, 455
 — compound, 455
 — juice, 455
 — mixture, 456
 De Lisle's thermometer, 42
 Decoctions, 520
 — aloes, compound, 97
 — arnica, 128
 — asparagus roots, 132
 — azedarach, 140
 — barley, 263
 — compound, 263
 — bittersweet, 221
 — compound, 221
 — black alder, 384
 — snakeroot, 185
 — blackberry root, 402
 — boneset, 226
 — brooklime, 144
 — broom, 422
 — compound, 422
 — burdock, 125
 — cabbage-tree bark, 253
 — cahina, 153
 — cantharides in turpentine, 165
 — carrageen, 184
 — catalpa, 175
 — chamomile, compound, 118
 — coffee, raw, 195
 — coltsfoot, 461
 — compound, 461
 — columbo, compound, 201
 — contrayerva, 203
 — cotton-root, 255

Decoction, dandelion, 455
 — dock root, 402
 — dogwood, 207
 — elder bark, 409
 — elecampane, 283
 — elm bark, 461
 — ergot, 224
 — figs, 243
 — compound, 243
 — galls, 248
 — geranium, 252
 — guaiacum wood, compound, 257
 — hairy horehound, 141
 — hardhack, 441
 — horehound, compound, 312
 — horse balm, 198
 — horsechestnut bark, 263
 — compound, 263
 — Iceland moss, 180
 — Indian hemp, 124
 — sarsaparilla, 262
 — ipecacuanha, 288
 — logwood, 259
 — madder, 402
 — common mallow, 309
 — marsh rosemary, 443
 — marshmallow, 98
 — matico, 314
 — mezereon, 318
 — compound, 318
 — New Jersey tea, 177
 — oak bark, 387
 — oatmeal, compound, 140
 — Pareira brava, 348
 — yellow Peruvian bark, 187
 — and cascarilla, 188
 — rhatany, 188
 — compound, 188
 — pipsissewa, 182
 — compound, 182
 — pomegranate rind, 256

Decoction, pomegranate root, 256
 — poppy heads, 347
 — prickly ash, 469
 — queen's root, 444
 — quince seed, 216
 — sarsaparilla, 413
 — compound, 413
 — Feltz's 413
 — Jauperand's 414
 — Vinache's, 413
 — Zittmann's, 414
 — seneka, 423
 — soot, 245
 — stavesacre, 443
 — tormentil, 459
 — uva ursi, 462
 — water avens, 253
 — white hellebore, 467
 — wild indigo, 142
 — willow bark, 408
 — woods, 257
 — wormseed, 181
 Deleroix's depilatory, 130
 Delphinia, 217
 — ointment, 217
 — solution, 217
 Demulcents, 67
 Dentifrice, 482
 — bitartrate potassa, 382
 — burnt hartshorn, 207
 — charcoal, 169
 — chlorinated lime, 154
 — electuary, 157, 158
 — mastich, 313
 — orris-root, 289
 — Peruvian bark, 186
 — phosphate of lime, 158
 — rhatany, 293
 — sulphate of quinia, 392
 — willow bark, 408
 — with carbonate of lime, 157
 Depilatories, 482
 — orpiment, 130
 Deshler's salve, 394
 Dewberry, 402
 Dewees's carminative, 134, 306
 — tincture of guaiacum, 258
 Diachylon, 360
 Diapalma plaster, 360
 Diaphoretics, 67
 Di-arsenite of quinia, 388
 Dietetic preparations, 487
 Diet-drink, Lisbon, 413
 Diet in convalescence, 74
 Digitalin, 219
 — pills, 220
 — granules, 220
 Digitalis, 217
 Diluents, 68

Dinner pills, 313
 Dinneford's fluid magnesia, 306
 Dippel's animal oil, 329
 — liniment, 329
 — mixture, 329
 — tincture, 329
 Disinfecting liquid of Ledoyen, 484
 Displacement, 521
 Distillation, 525, 529
 Distilled oils, 528
 — waters, 525
 Dittany, 212
 — infusion, 212
 — oil, 212
 Diuretics, 68
 Dock, 402
 — decoction, 402
 Dog rose, 399
 — confection, 399
 Dogwood, 207
 — decoction, 207
 — pills, 208
 — Jamaica, 353
 — round-leaved, 207
 — wine, 208
 — swamp, 207
 Dogsbane, 124
 Dolichos, 324
 Donovan's solution, 129
 Dose of medicines, 60
 Douche, 538
 Dover's powder, 286
 — eclectic, 345
 Drops, 26-534
 — Bateman's, 343
 — Battley's, 343
 — black, 341
 — carbonate ammonia, 106
 — carminative, 305
 — Koechlin's, 214
 — odontalgic, 342
 — sulphuret carbon, 451
 — table of, 26
 — toothache, 339
 Draughts, 524
 — alum, 102
 — anodyne, 344
 — black, 426
 — effervescing, 377, 378, 381
 — carbonate ammonia, 106
 — and ginger, 106
 — morphia, 320
 — muriatic acid, 85
 — sal-ammoniac, 109
 — solution, iodide of arsenic and mercury,
 129
 — sugar of milk, 296
 — sulphate potassa, effervescing, 381

E

Eau des Carmes, 316
 — de Dardel, 316
 — de Javelle, 363
 — de luce, 106-448
 — medicinale de Husson, 196-198-257-467
 — de Pagliari, 482
 Effervescing draught, 377-378-381
 — powders, 82-88-377-436
 Egg, 346
 — cerate, 347
 — clyster, emollient, 346
 — restorative, 346

Egg, emulsion, 346
 — liniment, 347
 — mixture, 346
 — with brandy, 346
 — wine, 346
 — oil, 346
 Eläometer, 35
 Elaterin, 222
 — tincture, 222
 Elaterium, 222
 — mixture, 222
 — oil, 222

- Elaterium, pills, 222
 Elder, 408
 — decoction of bark, 409
 — extract of berries, 409
 — mixture, 409
 — gargle, 408
 — ointment, 409
 — leaves, 409
 — vinegar, 408
 — water of flowers, 408
 Elecampane, 283
 — decoction, 283
 — extract, 283
 — mixture, 283
 — oxymel, compound, 284
 — pills, compound, 283
 Electuaries, 518
 Electuary, alumina, 99
 — dentifrice, 157, 158
 — laxative, 96
 — for teeth, 99
 — of alum, 101
 — angustura, 117
 — bitartrate of potassa, 383
 — black oxide of iron, 239
 — pepper, 352
 — burnt sponge, 442
 — calamus, 153
 — calomel, 273
 — carburet of iron, 232
 — catechu, 176
 — charcoal, 169
 — cinnamon, compound, 193
 — cowhage, 324
 — cubebs, 211
 — indigo, 283
 — jalap, 290
 — juniper, 292
 — kino, 293
 — lenitive, 425
 — logwood, extract of, 259
 — male fern, 244
 — orange leaves, 136
 — peel, 135
 — Peruvian bark, 189
 — astringent, 189
 — with catechu, 189
 — cloves, 189
 — iron, 189
 — sulphur, 189
 — tin, 189
 — pomegranate root, 256
 — rhatany, 294
 — rhubarb, 396
 — roses, 400
 — scurvy grass, 194
 — semen contra, 130
 — senna and cream of tartar, 425
 — figs, 425
 — rhubarb, 425
 — sulphur, 425
 — soap, 412
 — squill, 422
 — sulphate of copper, 215
 — soda, 439
 — sulphur, 449, 450
 — compound, 450
 — sulphuret of potassium, 370
 — tamarinds, 454
 — tin, 442
 — turpentine, 456
 — valerian, 463
 — Virginia snakeroot, 428
 Elemi, 222
 — caustery plaster, 223
 — ointment, 222
- Elixir, Boerhaave's, 98
 — Clauder's, 97
 — Garus's, 98, 210
 — ipecacuanha, 289
 — Lettsom's, 344
 — opium, 336
 — orange peel, 135
 — salutaris, 426
 — scammony, 418
 — Stoughton's, 98
 — vitriol, 87
 — Wedell's, 421
 Elm bark, 461
 — decoction, 461
 — compound infusion, 461
 — slippery, 462
 Elutriation, 515
 Embrocation, caraway, 172
 — croton oil, 335
 — petroleum, 349
 Embrocations, 542
 Emetic, tartar, 122, 558
 Emetics, 68
 Emetin, 223
 — impure, 223
 — lozenges, 223
 — mixture, 223
 — pure, 223
 — syrup, 223
 Emmenagogues, 69
 Emollients, 67, 541
 Emulsions, 524
 — almond, 113, 114
 — aromatic, 114
 — anthelmintic, 331, 427
 — artificial musk, 448
 — bicarbonate soda, 436
 — benzoin, cosmetic, 147
 — camphor, 160
 — cantharides, 164
 — carbonate of potassa, 376
 — castor-oil seed, 331
 — chloroform, 184
 — copaiba, 205
 — croton oil, 334
 — cubebs, ethereal extract, 211
 — egg, 346
 — hemp seed, 163
 — jalap, 290
 — kermes mineral, 121
 — lupulin, 304
 — manna, 311
 — nitrated, 380
 — opoponax, 345
 — purgative, 308
 — sal ammoniac, 109
 — scammony, 417
 — compound, 417
 — seneka, 424
 — squill, 421
 — sulphate soda, 439
 — tartar emetic, 122
 — tolu, 458
 — vermifuge, 332
 Enemata, 66
 Epispastics, 69
 Epsom salts, 307
 Ergotine, 224
 Ergot, 223
 — clyster, 224
 — decoction, 22
 — extract, 224–225
 — infusion, 223
 — injection, 224
 — mixture, 224
 — extract, 224

- Ergot, oil, 225
 — pills, compound, 224
 — extract, 225
 — powder, compound, 224
 — syrup, 224
 — tincture, 224
 — wine, 224
 Errhine, alum, 100
 — euphorbium, 227
 Errhines, 71
 Escharotics, 71
 Essences, 534
 Essence of allspice, 351
 — angelica, 117
 — aniseed, 118
 — caraway, 172
 — cinnamon, 193
 — fennel, 245
 — ginger, 476
 — lavender, compound, 300
 — lemon, 300
 — nutmeg, 325
 — peppermint, 316
 — roses, 400
 — rosemary, 401
 — sarsaparilla, 414
 — ——— compound, 414
 — soap, 412
 — ——— camphorated, 412
 — spearmint, 317
 — vanilla, 466
 — Ward's, for headache, 161
 Essential oils, 528
 Ether, acetic, 91
 — mixture, 91
 — spirit, 91
 — cantharidalis, 166
 — cupreous, 214
 — hydrocyanic, 92
 — muriatic, 92
 — ——— mixture, 92
 — ——— spirit, 92
 — nitrous, 91
 — ——— mixture, 91
 — ——— sweet spirit, 91
 — ——— ——— mixture, 91
 — phosphorated, 350
 — sulphuric, 92
 — ——— ethereal oil, 92
 — ——— Hoffmann's anodyne, 92
 — ——— lotion, 92
 — ——— mixture with camphor, 93
 — ——— ——— turpentine, 93
 — ——— rectified, 92
 — ——— spirit, 92
 — ——— syrup, 92
 — terebinthinated, 93
 Ethereal oil, 92
 Ethiops mineral, 279
 Euphorbium, 226
 — oil, 227
 — plaster, 227
 — sternutatory, 227
 Examination of excretions, 58
 Exercise in convalescence, 74
 Expectorants, 71
 Extracts, 517
 Extract of aconite, 89
 — ——— alcoholic, 89
 — ——— ammoniated, 90
 — American centaury, 404
 — ——— hellebore, 468
 — anemone, 116
 — arnica, 128
 — asparagus roots, 132
 — ——— shoots, 132
 — Extract of belladonna, 145
 — ——— alcoholic, 145
 — bistort, 149
 — bittersweet, 221
 — black hellebore, 261
 — ——— alcoholic, 261
 — black pepper, fluid, 352
 — black snakeroot, 479
 — ——— fluid, 479
 — broom, 422
 — buchu, 151
 — ——— fluid, 151
 — buckbean, 317
 — buckthorn, 394
 — butternut, 291
 — cabbage-tree bark, 253
 — cahinea, 153
 — Canada fleabane, 225
 — cantharides, 163
 — ——— ethereal, 164
 — cardamom, 170
 — carrot root, 171
 — cascarella, 173
 — cayenne pepper, 168
 — celandine, 181
 — centaury, 178
 — cevadilla, 404
 — chamomile, 118
 — colchicum bulb, 196
 — ——— acetic, 196
 — colocynth, 199
 — ——— compound, 199
 — columbo, 201
 — contrayerva, 203
 — cubebs, fluid, 211-481
 — ——— alcoholic-ethereal, 211
 — dandelion, 455
 — ——— fluid, 455
 — ——— and senna, 455
 — elder berries, 409
 — elecampane, 283
 — ergot, 224, 225
 — fever-root, 461
 — foxglove, 218
 — galls, 249
 — gentian, 250
 — ——— fluid, 250
 — geranium, 252
 — ginger, ethereal, 476
 — Goulard's, 356
 — guaiacum wood, 257
 — hardhack, 442
 — hemlock, 201
 — ——— alcoholic, 201
 — hemp, 162
 — ——— purified, 163
 — henbane, alcoholic, 280
 — ——— aqueous, 280
 — ——— fluid, 281
 — hops, 264
 — horehound, 312
 — horse-chestnut, 263
 — Indian hemp, 125
 — ipecacuanha, 287
 — ——— fluid, 483
 — jalap, 290
 — ——— alkaline, 290
 — juniper, 291
 — lettuce, 297
 — ——— watery, 297
 — liquorice, 254
 — lobelia, acetic, 303
 — ——— fluid, 304
 — logwood, 259
 — lupulin, 304
 — male fern, 244

Extract of marigold, 158
 — matico, 314
 — may apple, 361
 — mezereon, 318
 — myrrh, 326
 — nux vomica, 328
 — oak bark, 386
 — opium, acetous, 335
 — — alcoholic, 336
 — — aqueo-alcoholic, 335
 — — aqueous, 335
 — — by fermentation, 336
 — — denarcotized, 336
 — — roasted, 336
 — — vinous, 336
 — pareira brava, 348
 — paullinia, alcoholic, 348
 — pellitory, 385
 — Peruvian bark, 187
 — — fluid, 187
 — — vinous, 187
 — pink-root, compound, 441
 — — fluid, 441
 — — and senna, 441
 — pipsissewa, 182
 — poison oak, 460
 — pomegranate-root, 256
 — poppy heads, 347
 — quassia, 386
 — rhatany, 293

Extract of rhubarb, 396
 — fluid, 396
 — and senna, 396
 — rue, 403
 — sarsaparilla, 414
 — — fluid, 414
 — savine, 405
 — scammony, 417
 — seneka, with squill, 424
 — senna, alcoholic, 425
 — — fluid, 427
 — squill, 422
 — — acetic, 422
 — stramonium leaves, 444
 — — seed, 444
 — succory, 185
 — tansy, 454
 — tobacco, 452
 — tormentil, 459
 — uva ursi, 462
 — valerian, 464
 — — fluid, 464
 — vanilla, fluid, 466
 — Virginia snakeroot, fluid, 428
 — willow bark, 407
 — wood sorrel, 77
 — wormwood, 75
 — yellow root, 469
 Eyebright, 227

F

Fahrenheit's thermometer, 42
 False sarsaparilla, 125
 Fats, 533
 Fecula of hemlock, 201
 Felt splints, 486
 Fennel, 244
 — essence, 245
 — infusion, 245
 — oil, 245
 — ointment, 245
 — powder, compound, 244
 — water, 245
 Fern, male, 243
 — bolus, 244
 — electuary, 244
 — extract, ethereal, 244
 — — pills, 244
 — jelly, 244
 — mixture, 244
 — oil, 244
 — — pills, 244
 — powder, 244
 Ferrocyanate of quinia, 388
 Ferrocyanuret of iron, 205
 — — zinc, 472
 Ferro-sulphuret of potassium, 369
 Fever-root, 461
 — extract, 461
 Figs, 243
 — coffee, 243
 — decoction, 243
 — — compound, 243
 — gargle, 243
 — paste, 243
 Figwort, 423
 — oil, 423
 — ointment, 423
 Filter, Boullay's, 522
 Fires, colored, 482
 Fixed oils, 533

Flaxseed, 302
 — cataplasm, 302
 — — with bran, 302
 — clyster of oil, 303
 — infusion, 302
 — liniment of oil, 303
 — meal, cataplasm, 302
 — — compound, 302
 — mixture, 302
 Fleabane, 225
 — Canada, 225
 — — infusion, 225
 Fleming's tincture of aconite, 477
 Flies, Spanish, 163
 Florentine orris, 289
 Flour, boiled, 488
 Fluid extract of black pepper, 352
 — black snakeroot, 479
 — buchu, 151
 — cubebs, 211, 481
 — dandelion, 455
 — — and senna, 455
 — gentian, 250
 — henbane, 281
 — ipecacuanha, 483
 — lobelia, 304
 — Peruvian bark, 187
 — pinkroot, 441
 — — and senna, 441
 — — compound, 441
 — rhubarb, 396
 — — and senna, 396
 — sarsaparilla, 414
 — senna, 427
 — valerian, 464
 — vanilla, 466
 — Virginia snakeroot, 428
 Flummery of oatmeal, 140
 Fomentations, 541
 Fomentation, arnica, 128

Fomentation, arnica, with rue, 129
 — aromatic, 401
 — carbonate of potassa, 376
 — common salt, 431
 — nitric acid, 85
 — opium, 340
 — phosphoric acid, 86
 — sal ammoniac, 108
 — sulphate of zinc, 47
 — wine of opium, 342
 Foreign weights, 20
 Fowl, with rice, 489
 Fowler's solution, 373
 — mixture, 374
 — iodine and arsenic, 374
 Foxglove, 217
 — bolus, 218
 — extract, 218
 — infusion, 218
 — mixture, expectorant, 219
 — with acetate of lead, 219
 — ——— potassa, 219
 — ——— tartaric acid, 219

Foxglove, ointment, 219
 — pills, 218
 — — and squill, 218
 — plaster, 219
 — powder, 217
 — — saccharated, 406
 — syrup, 219
 — tincture, 219
 — — ethereal, 219
 — — mixture, 219
 — — vinegar, 219
 Frankincense, 353
 Frictions, 542
 Frost-weed, 260
 Frumenty, 490
 Fuligokali, 246
 — ointment, 246
 — — sulphuretted, 246
 — — pills, 246
 Fumigations, 546
 Fumigation with belladonna, 145
 Furniture of sick room, 58
 — varnish, 486

G

Galbanum, 247
 — mixture, 247
 — pills, compound, 247
 — plaster, 247
 — — compound, 247
 — purified, 247
 — tincture, 247
 — — compound, 247
 Gallic acid, 82
 — injection, 82
 — pills, 82
 Galls, 248
 — decoction, 248
 — extract, 249
 — — ointment, 249
 — gargle, 248
 — infusion, 248
 — — compound, 248
 — lotion, 248
 — ointment, 249
 — — compound, 249
 — powder, compound, 248
 — syrup, 248
 — tincture, 248
 Gallate of iron, 234
 Gamboge, 249
 — pills, compound, 249
 — powder, compound, 249
 — mixture, 249
 — — with elaterin, 250
 — solution, alkaline, 249
 — tincture, alkaline, 250
 — — ammoniacal, 250
 Gannal's injection, 478
 Gargle, ammonia, 103
 — acetate of ammonia, 105
 — — lead, 358
 — alum, 100, 101
 — ammoniated copper, 214
 — borax, 434
 — chlorate of soda, 437
 — cayenne pepper, 167
 — chlorinated soda, 437
 — chlorine, 183
 — cinnamon, tincture, 193
 — contrayerva, 203
 — cyanuret of mercury, 274
 — elder flowers, 408

Gargle, figs, 243
 — galls, 248
 — iodine, 285
 — mercurial, 265
 — muriatic acid, 84
 — mustard, 430
 — myrrh, 326
 — nitrate of potassa, 380
 — oak bark, 387
 — Peruvian bark, 190
 — pomegranate rind, 256
 — sage, 408
 — scurvy grass, 195
 — subacetate of lead, 357
 — sulphate of copper, 215
 — — quinia, 392
 — — zinc, 474
 — tannic acid, 87
 — tormentil, 459
 — vinegar, 78
 Garlic, 93
 — cataplasm, 93
 — liniment, 93
 — lotion, capillary, 93
 — syrup, 93
 Gauthier's plaster, 360
 Gay Lussac's alcoomètre, 32
 Gentian, 250
 — extract, 250
 — fluid, 250
 — syrup, 251
 — infusion, 251
 — — compound, 251
 — — with rhubarb, 251
 — mixture, 251
 — pills, compound, 250
 — powder, compound, 250
 — syrup, 251
 — tincture, 251
 — — acidulated, 251
 — — alkaline, 252
 — — ammoniacal, 252
 — — compound, 251
 — — mixture, 252
 — — with sulphuric acid, 252
 — — wine, 251
 Gentianin, 252
 — pills, 252

- Gentianin, syrup, 252
 tincture, 252
 German chamomile, 314
 infusion, 314
 Ginger, 475
 beer, 476-490
 powders, 436
 essence, 476
 extract, ethereal, 476
 syrup, 476
 tincture, 476
 infusion, 476
 lozenges, 476
 oil, 476
 spice plaster, 477
 syrup, 476
 tincture, 476
 Glass, soluble, 381
 Glauber's salts, 438
 Gloucester jelly, 487
 Glue, liquid, 482
 marine, 483
 Glycerin, 254
 lotion, 254
 ointment, 254
 paste, 254
 Goadby's solutions, 478
 Godfrey's cordial, 343
 Gold, 136
 ointment, 137
 powder, 136
 syrup, 137
 solder for, 486
 ammoniated, 139
 pills, 139
 chloride, 137
 bolus, 137
 caustic, 137
 collyrium, 137
 ointment, 137
 pills, 137
 powder, 137
 tincture, 137
 with sodium, 137
 lozenges, 138
 ointment, 138
 pills, 138
 powder, 137
 solution, 138
 syrup, 138
 compound, 138
 cyanide, 138
 lozenges, 139
 pills, 139
 powder, 138
 solution, 139
 Gold, iodide, 139
 oxide, 139
 pills, 139
 powder, 139
 purple of Cassius, 139
 Golden sulphuret of antimony, 121
 Goldthread, 206
 infusion, 206
 tincture, 206
 Gondret's ointment, 106
 Goulard's balsam, 358
 cerate, 358
 extract, 356
 Granulation, 515
 Granules of digitalin, 220
 Granville's lotion, 103
 Gravity, specific, 28
 Grease balls, 411
 Griffith's mixture, 231, 326
 Griffith's pills, 96
 Groat gruel, 140
 Gruel, oatmeal, 140
 rice, 346
 Guaiacum, 257
 decoction, compound, 257
 extract, 257
 mixture, 258
 with bittersweet, 258
 oil, 257
 compound, 257
 pills with aloes, 258
 antimony, 258
 sulphur, 258
 turpentine, 258
 tartar. emetic, 122
 powder, compound, 258
 tincture, 258
 ammoniated, 259
 Dewees's, 258
 and paregoric, 259
 copaiba, 259
 mixture, odontalgic, 259
 with henbane, 259
 Guarana, 348
 Gum Arabic, 76
 linctus, 77
 lozenges, 76, 77
 mixture, 76
 Wendt's, 77
 mucilage, 76
 paste, 76
 paté de guimauve, 76
 syrup, 77
 compound, 77
 plaster, 112
 tragacanth, 460

H

- Hair dye, 127
 Hairy horehound, 141
 decoction, 141
 Hardhack, 441
 decoction, 441
 extract, 442
 Hartshorn, 207
 burnt, 207
 dentifrice, 207
 jelly, 207
 compound, 207
 Hedge hyssop, 257
 powder, compound, 257
 wine, 257
 Heinecke's solution, 433
 Hellebore, American, 468
 black, 260
 white, 467
 Hemlock, 201
 cataplasm, 202
 collutory, 202
 extract, 201
 alcoholic, 201
 fecula, 201
 infusion, 202
 mixture, with paregoric, 203
 oil, 202, 354
 ointment, 202

Hemlock, pills with calomel, 202
 — pills with dandelion, 202
 — ipecacuanha, 202
 — pitch, 354
 — plaster, 202
 — compound, 202
 — powder, 201
 — saccharated, 406
 — suppository, 203
 — tincture, 202
 — ethereal, 202

Hemp, 162
 — emulsion, 163
 — extract, 162
 — purified, 163
 — resin, 163
 — tincture, 163
 — Indian, 124

Henbane, 280-566
 — extract, alcoholic, 280
 — aqueous, 280
 — fluid, 281
 — infusion, 281
 — compound, 281
 — liniment, 282
 — mixture, 281
 — with antimony, 281
 — squill, 282
 — ointment, 282
 — compound, 282
 — pills, compound, 281
 — and ipecacuanha, 281
 — opium, 281
 — powder, compound, 281
 — saccharated, 406
 — tincture, 281
 — ethereal, 281

Hiera picra, 162
 — tincture, 162

Hive syrup, Coxe's, 420-424

Hoffmann's anodyne, 92

Holly, sea, 225

Honeys, 314-520
 — borax, 434
 — clarified, 314
 — ceromel, 315
 — hydromel, 315
 — mixture, expectorant, 315
 — oxymel, 314
 — pectoral, 315
 — prepared, 314
 — water, 315
 — roses, 400
 — violets, 468

Hooper's pills, 95-241

Hope's mixture, 160

Hops, 263
 — cataplasm, 264
 — extract, 264
 — infusion, 264
 — mixture, 264
 — ointment, 264

Hops, tincture, 264
 — alkaline, 264

Horehound, 312
 — candy, 313
 — decoction, compound, 312
 — extract, 312
 — hairy, 141
 — mixture, pectoral, 312
 — syrup, 313

Horse balm, 198
 — decoction, 198
 — oil, 198
 — tincture, 198
 — chestnut, 262
 — decoction, 263
 — compound, 263
 — extract, 263
 — powder, compound, 262

Horsemint, 318
 — liniment, 319
 — oil, 318

Horseradish, 127
 — cataplasm, compound, 128
 — infusion, 128
 — compound, 128
 — mixture, 128
 — spirit, compound, 128

Hot bath, 537

Hundred-leaved rose, 399

Hungary water, 401

Huxham's tincture, 189

Hydrargyro-iodide of potassium, 368

Hydrated oxide of iron, 238

Hydriodate of ammonia, 107
 — ointment, 108
 — morphia, 321
 — quinia, 388
 — ioduretted, 389

Hydriodic acid, 83

Hydrochloric acid, 84, 555

Hydrocyanic acid, 82, 555
 — extemporaneous, 82
 — injection, 83
 — julep, 83
 — lotion, 83
 — mixture, 82
 — syrup, 83
 — ether, 92

Hydromel, 315
 — pectoral, 315

Hydrometers, 30

Hydrometrical equivalents, 36

Hydrosulphate of ammonia, 108
 — sulphuretted, 108

Hydrosulphuric acid, 83
 — artificial sulphuretted water, 83
 — bath, hydrosulphuretted, 83
 — lotion, hydrosulphuretted, 84

Hyposulphited sulphuret of potassium, 369

Hyposulphite of soda, 437

I

Iceland moss, 180
 — cetrarine, 180
 — powder, 181
 — chocolate, 180
 — decoction, 180
 — jelly, 180
 — mixture, 180

Idiosyncrasy, 61

Imperial drink, 383
 — measure, 25

Incompatibles, table of, 492

Indelible inks, 127, 483

Indian hemp, 124
 — decoction, 124
 — extract, 125
 — physic, 254
 — sarsaparilla, 261
 — decoction, 262
 — infusion, 262
 — mixture, 262

- Indian sarsaparilla syrup, 261
 — tobacco, 303
 — turnip, 131
 Indigo, 283
 — electuary, 283
 — pills, 283
 — powder, 283
 — sulphate, 283
 — wild, 142
 — decoction, 142
 — ointment, 142
 Infusions, 520
 Infusion, American centaury, 404
 — columbo, 245
 — senna, 174
 — angustura, 117
 — arnica, 128
 — compound, 128
 — balm, 315
 — barberry, 148
 — belladonna, 145
 — benne, 429
 — bistort, 148
 — black purslane, 226
 — blessed thistle, 177
 — bloodroot, 409
 — boneset, 226
 — compound, 226
 — broom, 422
 — buchu, 150
 — compound, 15
 — buckbean, 317
 — burdock, 125
 — calamus, 153
 — camphor, 160
 — Canada fleabane, 225
 — cantharides, 163
 — acetic, 164
 — caraway, 171
 — carrot seeds, 171
 — cascarrilla, 173
 — alkaline, 173
 — catechu, 176
 — compound, 176
 — catnep, 175
 — Cayenne pepper, 167
 — chamomile, 118
 — cherry-laurel, 298
 — chiretta, 182
 — cinnamon, 193
 — cloves, 172
 — columbo, 200
 — with ginger, 200
 — rhubarb, 200
 — dandelion, 455
 — dittany, 212
 — elm bark, 461
 — ergot, 223
 — fennel, 245
 — flaxseed, 302
 — fleabane, 225
 — foxglove, 218
 — galls, 248
 — compound, 248
 — gentian, 251
 — compound, 251
 — with rhubarb, 251
 — German chamomile, 314
 — ginger, 476
 — goldthread, 206
 — hemlock, 202
 — henbane, 281
 — compound, 281
 — hops, 264
 — horseradish, 127
 — compound, 128
 Infusion, Indian sarsaparilla, 262
 — iron, bitter, 237
 — juniper, 291
 — compound, 291
 — kino, 293
 — linden flowers, 458
 — compound, 458
 — lobelia, 303
 — logwood, 259
 — maidenhair, 90
 — malt, 309
 — marigold, 158
 — matico, 313
 — compound, 314
 — mayweed, 208
 — mint, 317
 — compound, 317
 — mudar, 159
 — orange-peel, compound, 135
 — pareira brava, 348
 — parsley-root, 349
 — pennyroyal, 260
 — persimmon bark, 220
 — Peruvian bark, 188
 — compound, 188, 189
 — with lime water, 188
 — magnesia, 188
 — serpentaria, 188
 — valerian, 188
 — pinkroot, 440
 — compound, 441
 — pleurisy-root, 131
 — quassia, 386
 — compound, 386
 — red poppy, 399
 — compound, 399
 — rhatany, 294
 — rhubarb, 396
 — alkaline, 396
 — roses, acid, 400
 — safflower, 171
 — saffron, 210
 — sage, 408
 — compound, 408
 — sarsaparilla, 413
 — alkaline, 413
 — sassafras bark, 416
 — compound, 416
 — pith, 416
 — savine, 405
 — seneka, 423
 — compound, 423
 — senna, 426
 — compound, 426
 — with buckthorn, 427
 — coffee, 426
 — lemon juice, 427
 — rhubarb, 427
 — tamarinds, 426
 — tincture of senna, 427
 — simaruba, 429
 — compound, 429
 — skunk cabbage, 220
 — slippery elm, 462
 — southern wood, 130
 — succory, 185
 — tamarinds, 454
 — tansy, 454
 — tobacco, 453
 — tulip tree bark, 303
 — valerian, 463
 — compound, 463
 — Virginia snakeroot, 428
 — compound, 428
 — with ether, 429
 — yellow ladies' bedstraw, 248

- Infusion, yellow root, 469
 — wild cherry bark, 384
 — ginger, 131
 — wormwood, 75
 Inhalations, 535
 Inhalation, balsam of tolu, 459
 Injections, anatomical, 477
 — chloride of zinc, 477
 — Gannal's, 478
 — molasses, 477
 — nitrate of lead, 478
 Injection, acetate of zinc, 473
 — aloes, 97
 — alum, 101, 102
 — ammonia, 103
 — ammoniated copper, 214
 — carbonate of soda, 435
 — catechu, 177
 — caustic potassa, 372
 — chloride of zinc, 471
 — chlorinated lime, compound, 155
 — soda, 437
 — chlorine, 183
 — chloroplatinate of sodium, 356
 — copaiba, 205
 — corrosive sublimate, 270
 — cubebs, 211
 — deterative, 97
 — ergot, 224
 — gallic acid, 82
 — hydrocyanic acid, 83
 — iodide of iron, 236
 — potassium, 364
 — ioduretted, 367
 — kino, 293
 — kousso, 149
 — morphia, 320
 — opium, 340
 — pomegranate rind, 256
 — pyroligneous acid, 79
 — rhatany, 294
 — sedative, 348
 — soot, 246
 — stimulating, 326
 — subacetate of lead, 357
 — and lime water, 358
 — sulphate of copper, 216
 — iron, 241
 — zinc, 475
 — sulphuret of potassium, 371
 — tannic acid, 87
 Ink, blue, 234-483
 — indelible, 127-483
 — without mordant, 127
 — red, 483
 — writing, 234-483
 Inspissated juices, 517
 Iodate of potassa, 379
 — strychnia, 447
 Iodide of arsenic, 129
 — and mercury, 129
 — barium, 143
 — gold, 139
 — iron, 235
 — lead, 359
 — manganese, 310
 — mercury, 275
 — red, 275
 — potassium, 364, 562
 — quinia, 388
 — silver, 126
 — starch, 116
 — sulphur, 452
 — zinc, 472
 Iodine, 284, 562
 — bath, 284
 — cataplasm, 285
 — clyster, 285
 — gargle, 285
 — liniment, 285
 — lotion, 285
 — mixture, 285
 — with iodide of potassium, 284
 — ointment, 285
 — compound, 285
 — with mercury, 285
 — oil of tobacco, 285
 — pills, 284
 — plaster, 285
 — solution, 284
 — tincture, 284
 — compound, 284
 — ethereal, 284
 — saturated, 284
 Ioduretted potassium, 366
 — water, 367
 Ipecacuanha, 286
 — bolus, 287
 — confection, 287
 — decoction, 288
 — elixir, 289
 — extract, 287
 — fluid, 483
 — lozenges, 287
 — and camphor, 287
 — mixture, 288
 — pills, with centaury, 287
 — foxglove, 287
 — opium, 287
 — squill, 287
 — powder with antimony, 286
 — benzoic acid, 80
 — calomel, 286
 — carb. soda, 286
 — muriate ammonia, 286
 — myrrh, 287
 — nitre, 286
 — opium, 286
 — rhubarb, 286
 — tartar emetic, 122, 286
 — tragacanth, 287
 — saccharated, 406
 — syrup, 288, 483
 — compound, 288
 — tincture, 288
 — wild, 226
 — powder, 226
 — wine, 288
 — alkaline, 288
 — compound, 288
 — with tartar emetic, 288
 Irish moss, 184
 Iris, Florentine, 289
 Iron, 227, 562
 — preparations of, 483
 — solder for, 485
 — filings, 227
 — mixture, compound, 230
 — powder, 227
 — wire, 227
 — acetate, 228
 — and alum, 228
 — tincture, alcoholic, 228
 — ethereal, 228
 — ammoniated, 229
 — bolus, 229
 — mixture, 229
 — pills, 229
 — powder, compound, 229
 — tincture, 229
 — arseniate, 229
 — ointment, 230

Iron, arseniate, pills, 229
 bromide, 230
 ointment, 230
 pills, 230
 carbonate, 230
 artificial chalybeate water, 231
 pills, 230
 plaster, 231
 saccharine, 230
 and manganese, 483
 subcarbonate, 230
 mixture, 230
 pills, comp., 231
 powder, 230
 wine, 231
 carburet, 231
 electuary, 232
 mixture, 232
 ointment, 232
 pills, 232
 plaster, 232
 powder, 232
 prepared, 231
 chloride, 232
 and acetate of lead, 232
 mixture, 232
 syrup, 232
 tincture, 232
 ethereal, 233
 citrate, 233
 ammonio-, 229
 and magnesia, 234
 syrup, 234
 quinia, 233
 liquid, 233
 syrup, 233
 tincture, 233
 wine, 233
 aromatic, 233
 ferrocyanuret, blue ink, 234
 mixture, 234
 ointment, 234
 pills, 234
 powder, 234
 compound, 234
 Prussian blue, 234
 gallate, 234
 writing ink, 234
 iodide, 235
 bath, 237
 injection, 236
 lozenges, 235
 mixture, 236
 ointment, 237
 pills, 235
 solution, 235
 officinal, 236
 syrup, 236
 and chloride of iron, 236, 237
 tincture, 236
 wine, 236
 lactate, 237
 lozenges, 237

Iron, lactate, pills, 237
 syrup, 237
 malate, 237
 bitter infusion, 237
 mixture, 237
 solution, 237
 tincture, 237
 metallic, prepared, 227
 bolus, 228
 mixture, 228
 pills, 228
 powder, 228
 compound, 228
 nitrate, 238
 per-sesquinitrate, solution, 238
 proto-nitrate, syrup, 238, 484
 oxide, black, 238
 electuary, 239
 pills, 239
 powder, 239
 hydrated, 238
 red, 239
 pills, 239
 plaster, 239
 phosphate, 240
 compound syrup, 484
 proto-iodide, 235
 pills, 235
 syrup, 236
 tincture, 236
 sulphate, 240
 dried, 240
 injection, 241
 mixture, 241
 compound, 241
 pills, 240
 Bland's, 240
 compound, 240
 with rhubarb, 240
 powder, 240
 solution, 241
 syrup, 241
 tersulphate, solution, 238
 sulphuret, 241
 tannate, 243
 tartrate, and potassa, 241
 powder, and columbo, 242
 effervescent, 242
 solution, 242
 tincture, compound, 242
 wine, 242
 compound, 242
 ammonio-, 242
 solution, 243
 valerianate, 243
 Irritants, 565
 Isinglass, 282
 blanc mange, 491
 court plaster, 282
 diamond cement, 283
 Issues, 543
 ointment, 166

J

Jackson's pectoral lozenges, 484
 syrup, 77-484
 Jalap, 289
 electuary, 290
 emulsion, 290
 extract, 290
 alkaline, 290
 mixture, 290

Jalap, pills, compound, 290
 with calomel, 290
 powder, compound, 290
 saccharated, 406
 with calomel, 289
 cream of tartar, 289
 ipécacuanha, 289
 scammony, 289

Jalap, resin, 290
 — soap, 290
 — tincture, 290
 — compound, 290
 — mixture, 291
 Jamaica dogwood, 353
 — tincture, 353
 — pepper, 351
 James's powder, 124
 Jamestown weed, 444
 Japan varnish, 486
 Jelly, almond, 487
 — biscuit, 487
 — bread, 487
 — calves' feet, 488
 — carrageen, 184
 — chicken, 490
 — Gloucester, 487
 — hartshorn, 207
 — compound, 207

Jelly, Iceland moss, 180
 — male fern, 244
 — orange leaves, 136
 — pectoral, 424
 — rice, 346
 — starch, 116
 — tapioca, 454
 Juices, inspissated, 517
 Juice of dandelion, 455
 Julep, alum, 102
 — hydrocyanic acid, 83
 Juniper, 291
 — electuary, 292
 — extract, 291
 — infusion, 291
 — compound, 291
 — liniment, 292
 — mixture of oil, 292
 — spirit, compound, 292

K

Kentish ointment, 333
 Kermes mineral, 120
 — lozenges, 120
 Kermesine powder, 120
 Keyser's pills, 268
 Kinate of cinchonia, 192
 — quinia, 389
 Kino, 292
 — electuary, 293
 — infusion, 293

Kino, injection, 293
 — pills, compound, 293
 — powder, compound, 293
 — syrup, 293
 — tincture, 293
 Kirkland's neutral cerate, 484
 Kouso, 149
 — infusion, 149
 Krameria, 293

L

Labarraque's solution, 437
 Lac, 296
 — bleached, 296
 — cement, 296
 — sealing wax, 296
 — solution, aqueous, 296
 — varnish, 486
 Lactate of iron, 237
 — quinia, 389
 Lactic acid, 84
 — lemonade, 84
 — lozenges, 84
 Lactucarium, 297
 — lozenges, 297
 — mixture, 297
 — pills, 297
 — syrup, 297
 — tincture, 297
 Ladies' bed straw, yellow, 248
 — infusion, 248
 — wine, 248
 Lady Webster's pills, 95
 Lancaster black drop, 341
 Lard, benzoated, 146
 Larkspur, 217
 — tincture, 217
 Lartigue's pills, 196
 Laudanum, 342
 — Smith's, 343
 — Swediaur's, 344
 — Sydenham's, 342
 Laurel, 298
 — oil, 298
 — ointment, 298

Laurel, cherry, 298
 Lavender, 299
 — essence, compound, 300
 — ointment, 300
 — powder, compound, 299
 — spirit, 299
 — compound, 299
 — tincture, ethereal, 300
 — water, 299
 Lead, 356, 563
 — solder for, 485
 — acetate, 356
 — cataplasm, 357
 — cerate, 357
 — gargle, 358
 — Goulard's balsam, 358
 — liniment, 357
 — lotion, 357
 — mixture, 358
 — ointment, 357
 — pile, 357
 — paste, rubefacient, 356
 — pills, 356
 — plaster, 357
 — powder, 356
 — carbonate, 358
 — ointment, 358
 — camphorated, 359
 — plaster, 359
 — powder, camphorated, 358
 — chloride, 359
 — lotion, 359
 — ointment, 359
 — iodide, 359

- Lead, iodide, ointment, 360
 — pills, 359
 — powder, 359
 — litharge, 360
 — cataplasm, 360
 — cerate, 360
 — ointment, 360
 — ————— Banyer's, 360
 — ————— compound, 360
 — ————— plaster, 360
 — ————— Baynton's, 360
 — ————— diapalma, 360
 — ————— Gaultier's, 360
 — red, 360
 — ————— plaster, Nuremberg, 360
 — saccharate, 361
 — ————— nitro, 361
 — ————— solution, 361
 — subacetate, 356
 — ————— cataplasm, 357
 — ————— cerate, 358
 — ————— collyrium, 356, 357
 — ————— gargle, 357
 — ————— Goulard's vegeto-mineral
 — ————— water, 357
 — ————— injection, 357
 — ————— with lime, 358
 — ————— liniment, 358
 — ————— ointment, 358
 — ————— plaster, 358
 — ————— solution, 356
 — tannate, 361
 — ————— liniment, 361
 — ————— ointment, 361
 Leather wood, 220
 — ————— varnish for, 486
 Leeching, 551
 Ledoyen's disinfecting fluid, 484
 Lemon, 300
 — collutory of juice, 300
 — Cologne water, 301
 — essence, 300
 — juice, artificial, 81
 — lozenges of juice, 300
 — ————— oil, 300
 — spirit, 301
 — ————— odoriferous, 301
 — syrup, 301
 — ————— vinous of peel, 301
 — tincture, 301
 — water of peel, 300-489
 — ————— unparalleted, 302
 Lemonade, 300
 — ————— barberries, 148
 — ————— compound, 300
 — ————— dry, 82
 — ————— lactic acid, 84
 — ————— phosphoric acid, 86
 — ————— sulphuric acid, 87
 — ————— tartaric acid, 88
 — ————— with sulphate of soda, 439
 Lenitive electuary, 425
 Lettsom's elixir, 344
 Lettuce, wild, 296
 — ————— extract, 297
 — ————— lactucarium, 297
 — ————— lozenges, 297
 — ————— mixture, 297
 — ————— pills, 297
 — ————— syrup, 297
 — ————— tincture, 297
 — ————— thridace, 297
 — ————— syrup, 297
 — ————— watery, 297
 — ————— mixture, 297
 — ————— powder, 297
 Lettuce, mixture, with dandelion, 296
 — ————— syrup, 298
 — ————— tincture, 297
 — ————— water, 298
 — ————— mixture, 298
 Lignum vitæ, 257
 Lime, 155
 — ————— ointment, 156
 — ————— water, 155
 — ————— compound, 155
 — ————— and carbonate of potassa, 156
 — ————— milk, 156
 — ————— liniment, 156
 — ————— with alcohol, 156
 — ————— oil, 156
 — ————— opium, 156
 — ————— sulphur, 156
 — ————— mixture, antacid, 156
 — ————— carbonate, 156
 — ————— dentifrice, 157
 — ————— electuary, 157
 — ————— lozenges, 157
 — ————— mixture, 157
 — ————— compound, 157
 — ————— powder, aromatic, 157
 — ————— compound, 157
 — ————— with opium, 157
 — ————— precipitated, 156
 — ————— prepared chalk, 156
 — ————— oyster shells, 156
 — ————— chlorinated, 154
 — ————— collutory, 154
 — ————— dentifrice, 154
 — ————— injection, 155
 — ————— liniment, 155
 — ————— lozenges, 154
 — ————— ointment, 155
 — ————— preservative liquid, 154
 — ————— phosphate, 157
 — ————— electuary, dentifrice, 158
 — ————— hartshorn, burnt, 157
 — ————— powder, dentifrice, 158
 — ————— syrup, 158
 Linctus, gum Arabic, 77
 — ————— green, 114
 — ————— opiate, 344
 — ————— opium, 340
 — ————— sulphur, 450
 — ————— white, 115
 Linden, 457
 — ————— infusion, 458
 — ————— compound, 458
 Liniments, 542
 Liniment, acetate of mercury, 268
 — ————— aconite root, 89
 — ————— alum, 102
 — ————— amber, oil of, 448
 — ————— ammonia, 103
 — ————— camphorated, 106
 — ————— compound, 103
 — ————— sulphuretted, 104
 — ————— terebinthinate, 104
 — ————— barytic, 143
 — ————— belladonna, 146
 — ————— cajeput oil, 329
 — ————— camphor, 161
 — ————— compound, 161
 — ————— ethereal, 161
 — ————— turpentine, 162
 — ————— with vinegar, 161
 — ————— camphorated ammoniacal, 106
 — ————— soap, 411
 — ————— cantharides, 165
 — ————— aromatic, 166
 — ————— soap, 165
 — ————— stimulating, 166

- Liniment, cantharides, with camphor, 165
 carbonate of potassa, 376
 ammonia, 106
 for chilblains, 141
 chlorinated lime, 155
 chlorine, 183
 chloroform, 184
 cod-liver oil, 330
 colchicum and camphor, 198
 cosmetic, 115
 croton oil, 335
 Dippel's oil, 329
 eggs, 347
 flaxseed oil, 303
 for milk abscess, 357
 sore breasts, 156
 garlic, 93
 henbane, 282
 horsemint, 319
 hydrosulphate ammonia, 108
 iodide of potassium, 366
 iodine, 285
 juniper, 292
 laudanum and lime water, 344
 lime and sulphur, 156
 water and alcohol, 156
 oil, 156
 opium, 156
 marigold, 158
 mercurial, compound, 267
 opiated, 267
 mercury and chalk, 265
 muriatic acid, 85
 mustard, 431
 nitrate of mercury, 276
 nitric acid, 85
 opiate, 344
 opium, 339
 with oil of chamomile, 339
 pellitory, 385
 phosphorated, 350
 sal ammoniac, 109
 soap, 411
 spermaceti, 180
 strychnia, 446
 subacetate of lead, 358
 succinate of ammonia, 110
 sulphate of quinia, 392
 sulphur and soap, 451
 sulphuret of calcium, 155
 carbon, 451
 sodium, 432
 sulphuric acid, 87
 tannate of lead, 361
 tartar emetic, ammoniacal, 123
 turpentine, 457
 acetic, 333
 oil, 333
 opiated, 333
 sulphuric, 333
 veratria, 466
 verdigris, 213
 Wilkinson's, 108
 Lip salve, 178, 179
 Liquid glue, 482
 Liquorice root, 254
 extract, 254
 refined, 254
 lozenges, 254
 with opium, 255
 mixture, 255
 compound, 255
 paste, 254
 syrup, 255
 Lisbon diet drink, 413
 List of incompatibles, 492
 Litharge, 360
 Lithontriptics, 64
 Liver of sulphur, 369
 Liverwort, 262
 Lobelia, 303
 extract, acetic, 303
 fluid, 304
 infusion, 303
 mixture, 303
 syrup, 304
 tincture, 303
 ethereal, 303
 vinegar, 304
 Local baths, 537
 Locatelli's balsam, 142
 Lockstadt's pills, 115
 Logwood, 259
 decoction, 259
 extract, 259
 electuary, 259
 mixture, 260
 infusion, 259
 Long pepper, 352
 cataplasm, 352
 Lotions, 542
 Lotion, anti-rheumatic, 444
 Barlow's, 370
 Bateman's, 270
 capillary, 93
 for burns, 147
 chilblains, 332
 hiccup, 107
 toothache, 332
 Granville's, 103
 Goulard's, 114
 hydrosulphuretted, 84
 mammillary, 141
 mercurial, 265
 Saviard's, 372
 of acetate of lead, 357
 zinc, 473
 aconitine, 90
 almond, compound, 113
 alum, 102
 ammonia, 104
 ammoniated alcohol, 107
 borax, 434
 bromine, 149
 carbonate of potassa, 376
 soda, 435
 Cayenne pepper, 168
 cherry-laurel water, 299
 chlorate of soda, 437
 chloride of lead, 359
 zinc, 471
 corrosive sublimate, 270
 and camphor, 270
 copper, 270
 cosmetic, 270
 creasote, 209
 cyanuret of potassium, 363
 galls, 248
 glycerin, 254
 hydrocyanic acid, 83
 ioduretted potassium, 367
 iodine, 285
 marigold, extract of, 158
 muriatic acid, 85
 mustard, 430
 nitrate of potassa, 380
 nitric acid, 85
 nux vomica, 329
 orpiment, 130
 oxide of zinc, 470
 Peruvian bark, 190
 sal ammoniac, 108

Lotion, sulphate of copper, 215
 ——— morphia and borax, 323
 ——— soda, 439
 ——— sulphuret of potassium, 370
 ——— compound, 370
 ——— sodium, 432
 ——— zinc, 475
 ——— sulphuric ether, 92
 ——— tartar emetic, 123
 ——— and camphor, 123
 ——— corrosive sublimate, 123
 ——— tobacco, 453
 ——— veratria, 466
 ——— verdigris, compound, 213
 Lozenges, 534
 ——— of anise, 118
 ——— balsam of tolu, 458
 ——— bicarbonate of soda, 436
 ——— borax, 433
 ——— catechu, 176
 ——— Cayenne pepper, 167
 ——— chalk, 157
 ——— charcoal, 169
 ——— chloride of gold and sodium, 138
 ——— chlorinated lime, 154
 ——— cinnamon, 192
 ——— citric acid, 81
 ——— croton oil, 334
 ——— cubebs, 211
 ——— extract, 211
 ——— cyanide of gold, 139
 ——— emetine, 223
 ——— ginger, 476
 ——— gum, 76
 ——— iodide of iron, 235
 ——— potassium, 364
 ——— ipecacuanha, 287
 ——— and camphor, 287
 ——— Jackson's, 484

Lozenges of kermes mineral, 120
 ——— lactate of iron, 237
 ——— lactic acid, 84
 ——— lactucarium, 297
 ——— lemon juice, 300
 ——— oil, 300
 ——— liquorice, 254
 ——— and opium, 255
 ——— magnesia, 305
 ——— manna, 311
 ——— marsh mallow, 99
 ——— muriate of morphia, 322
 ——— and ipecacuanha, 322
 ——— naphthaline, 327
 ——— oxalate of potassa, 380
 ——— oxalic acid, 86
 ——— paullinia, 348
 ——— pellitory, 385
 ——— peppermint, 316
 ——— rhubarb, 396
 ——— rose, 400
 ——— Spitta's, 211
 ——— of starch, 116
 ——— tartaric acid, 88
 ——— Tronchin's gum, 77-121
 ——— of vanilla, 465
 ——— Wistar's, 255-345
 Lugol's ioduretted water, 366
 ——— solution, 366
 Lunar caustic, 126
 Lupulin, 304
 ——— emulsion, 304
 ——— ointment, 304
 ——— pills, 304
 ——— powder, 304
 ——— syrup, 304
 ——— tincture, 304
 Luting for bottles, 480

M

Macassar oil, 115
 Mace, 305
 ——— balsam, nervine, 305
 ——— drops, carminative, 305
 ——— oil, 305
 ——— saccharated powder, 406
 ——— tincture, 305
 Maceration, 521
 Mackenzie's solution, 127
 Madder, 402
 ——— decoction, 402
 ——— powder, compound, 402
 Magendie's solution of morphia, 323
 Magnesia, 305
 ——— aperient effervescing, 309
 ——— Dinneford's fluid, 306
 ——— heavy, 305
 ——— mixture, 305
 ——— and gentian, 306
 ——— rhubarb, 306
 ——— powder and orange peel, 305
 ——— rhubarb, 305
 ——— sulphur, 305
 ——— troches, 305
 ——— acetate, 306
 ——— bitartrate, 309
 ——— carbonate, 306
 ——— carminative, Dalby's, 306
 ——— Dewees's, 306
 ——— mixture, 306
 ——— with camphor, 307

Magnesia, carbonate, mixture with colchicum, 306
 ——— solution, 306
 ——— citrate, 307
 ——— soluble, 307
 ——— solution, 307
 ——— water, 307
 ——— phosphate, 308
 ——— sulphate, 307
 ——— Cheltenham salts, 307
 ——— clyster, 308
 ——— emulsion purgative, 308
 ——— mixture, with coffee, 308
 ——— nitric acid, 308
 ——— Rochelle salt, 308
 ——— Seidlitz water, 307
 ——— sulphuric acid, 308
 ——— tartar emetic, 308
 ——— tartrate, 308
 Magnesium, sulphuret, 308
 ——— syrup, 308
 Magnolia, 309
 ——— tincture, 309
 Mahy's plaster, 359
 Maidenhair, 90
 ——— infusion, 90
 ——— syrup, 90
 ——— compound, 91

- Malate of iron, 237
 Male fern, 243
 — bolus, 244
 — electuary, 244
 — extract, ethereal, 244
 — pills, 244
 — jelly, 244
 — mixture, 244
 — oil, 244
 — pills, 244
 — powder, 244
 Mallow, common, 309
 — decoction, 309
 Malt, 309
 — clyster, 309
 — infusion, 309
 Management of convalescence, 74
 — sick room, 55
 Manganese, 309
 — carbonate, 309
 — pills, with iron, 309
 — iodide, 310
 — pills, 310
 — syrup, 310
 — and iron, 310
 — muriate, 310
 — pills, 310
 — oxide, black, 310
 — bolus, 310
 — ointment, 310
 — compound, 310
 — powder, 310
 — phosphate, 311
 — sulphate, 311
 Manna, 311
 — emulsion, 311
 — lozenges, 311
 — mixture, 311
 — syrup, 311
 Mannite, 312
 Marine glue, 482
 Marjoram, 345
 — oil, 345
 Marsh's test, 554
 Marsh mallow, 98
 — decoction, 98
 — lozenges, 99
 — ointment, 99
 — paste, 99
 — powder, 99
 — syrup, 98
 — rosemary, 443
 — decoction, 443
 Marigold, 158
 — extract, 158
 — lotion, 158
 — pills, 158
 — infusion, 158
 — liniment, 158
 Masterwort, 262
 Mastich, 313
 — collutory, 313
 — dentifrice, 313
 — dinner pills, 313
 — paste, anti-odontalgic, 313
 — tincture, ethereal, 313
 — varnish, crystal, 313
 — picture, 313
 Matico, 313
 — decoction, 314
 — extract, 314
 — infusion, 313
 — with senna, 314
 — ointment, 314
 — syrup, 314
 — tincture, 314
 May apple, 361
 — extract, 361
 — pills and podophyllin, 362
 — podophyllin, 361
 Mayweed, 208
 — cataplasm, 208
 — infusion, 208
 Meal, barley, 263
 — flaxseed, compound, 302
 Measures, 24
 — approximative, 25
 — drops, 26
 — foreign, 28
 — French, 27
 — imperial, 25
 — wine, 24
 Medicated waters, 484-525
 — baths, 539
 Mellites, 520
 Mercury, 264-563
 — purified, 264
 — cataplasm, 267
 — cerate, 267
 — compound, 267
 — gargle, 265
 — liniment, 267
 — compound, 267
 — opiated, 267
 — lotion, 265
 — mixture, 266
 — mucilage, 265
 — ointment, 266
 — camphorated, 267
 — compound, 267
 — and belladonna, 267
 — pills, 266
 — and antimony, 265
 — colocynth, 266
 — jalap, 266
 — quinia, 266
 — rhubarb, 266
 — Abernethy's, 266
 — compound, 266
 — plaster, 267
 — and belladonna, 267
 — powder, compound, 264
 — acetate, 268
 — liniment, 268
 — pills, 268
 — with opium, 268
 — solution, 268
 — ammoniated, 268
 — ointment, 268
 — and antimony, 264
 — chalk, 265
 — liniment, 265
 — with ipecacuanha, 265
 — gum, 265
 — pills, with hemlock, 265
 — magnesia, 266
 — borate, 268
 — bromide, 268
 — bibromide, 269
 — protobromide, 269
 — solution, ethereal, 269
 — calomel, 271
 — chloride, and quinia, 274
 — pills, 274
 — chloride, mild, 271
 — corrosive, 269
 — corrosive sublimate, 269
 — cyanuret, 274
 — gargle, 274
 — ointment, 274
 — pills, compound, 274
 — solution, 274

- Mercury, cyanuret, tincture, compound, 274
 — iodide, green, 275
 — ointment, 275
 — pills, 275
 — compound, 275
 — powder, 275
 — red, 275
 — ointment, 276
 — pills, 275
 — tincture, 276
 — ethereal, 276
 — wash, 276
 — nitrate, 276
 — acid, 276
 — liniment, 276
 — ointment, 276
 — with lead, 276
 — pills, 276
 — oxide, black, 277
 — wash, 277
 — ointment, 277
 — pills, 277
 — red, 277
 — ointment, 277
 — brown, 278
 — with cinnabar 278
 — lead, 278
 — sulphur, 277
 — tin, 278
 — zinc, 278
 — pills, 277
 — powder, 277
 — phosphate, 278
 — pills, 278
 — sulphate, yellow, 278
 — ointment, 279
 — powder, compound, 278
 — persulphate, 278
 — sulphuret, black, 279
 — pills, 279
 — powder, 279
 — red, 279
 — cerate, 280
 — ointment, 280
 — pills, 279
 — powder, 279
 — fumigating, 280
 — tartrate, 280
 — and potassa, 280
 Metals, varnish for coating, 486
 Method of displacement, 521
 Metrical weights, 21
 — measures, 27
 Metz's balsam, 213
 Mezereon, 318
 — decoction, 318
 — compound, 318
 — extract, 318
 — oil, 318
 — ointment, 318
 Milk, 294
 — anodyne, 113
 — arrowroot, 312
 — artificial, 113
 — asses', 491
 — goats', 295
 — cement, china, 296
 — chocolate, 490
 — coffee, 490
 — mixture, with suet, 295
 — soda water, 295
 — powder, 295
 — roses, 113, 147
 — sago, 407
 — Milk, sugar of, 295
 — draught, 296
 — with gum Arabic, 296
 — Iceland moss, 295
 — syrup, 295
 — vanilla, 466
 — whey alum, 295
 — aromatic, 295
 — cream of tartar, 295
 — mustard, 295
 — simple, 294
 — tamarind, 295
 — vinegar, 295
 — wine, 295
 Mindererus, spirit, 105
 Mineral, Kermes, 120
 Mint, 316, 318
 Mixtures, 524
 Mixture, acetate of ammonia, 105
 — lead, 358
 — morphia, 321
 — potassa, 373
 — acetic ether, 91
 — aconite, 90
 — aloes, alkaline, 97
 — aloetic, 97
 — American hellebore, 468
 — ammonia, 103
 — and ether, 103
 — ammoniac, 111
 — and nitric acid, 111
 — compound, 111
 — ammoniated alcohol, 107
 — iron, 229
 — angustura, 117
 — anisated ammoniated alcohol, 107
 — aniseed, 118
 — antacid, 156
 — anthelmintic, 130, 349
 — anti-emetic, 177, 436
 — anti-epileptic, 127
 — anti-otitic, 339
 — antimonial wine and ammoniac, 124
 — bittersweet, 124
 — laudanum, 124
 — anodyne, Vicat's, 107
 — arseniate of potassa, 374
 — asparagus, diuretic, 132
 — assafetida, 133
 — and oxymel of squill, 133
 — tolu, 133
 — compound, 134
 — avens, 253
 — balsam of Peru, 141
 — tolu, 459
 — and almond emulsion, 459
 — belladonna, 459
 — copaiba, 459
 — morphia, 459
 — opium, 459
 — barley, 490
 — belladonna, 146
 — benzoic acid, 80
 — and copaiba, 80
 — bicarbonate of potassa, 377
 — soda and copaiba, 436
 — bichloride of platinum, 355
 — biniodide of potassium, 368
 — bistort, 149
 — bitartrate of potassa, 383
 — bittersweet, 221
 — black drop, 342
 — hellebore, 261
 — boracic acid, 81
 — borax, 434

- Mixture, borotartrate of potassa, 375
 ————— and magnesia, 375
 ————— brown, 255, 344
 ————— buckbean, 317
 ————— diuretic, 318
 ————— butter of cocoa, 152
 ————— cabbage-tree bark, 253
 ————— camphor, 160, 161
 ————— and chloroform, 161
 ————— ether, 161
 ————— myrrh, 160
 ————— Parrish's, 160
 ————— water, 161
 ————— camphorated cough, 162
 ————— carbonate of ammonia, 106
 ————— and ginger, 106
 ————— magnesia, 306
 ————— and camphor, 307
 ————— colchicum, 306
 ————— potassa, 376
 ————— soda and chamomile, 434
 ————— gentian, 435
 ————— ipecacuanha, 434
 ————— quassia, 435
 ————— carburet of iron, 232
 ————— carminative, 316
 ————— carrageen, 184
 ————— cascarilla, 173
 ————— cassia pulp, 174
 ————— castor oil and ether, 332
 ————— catechu and logwood, 177
 ————— caustic potassa, 372
 ————— celandine, 181
 ————— centaury, 178
 ————— chalk, 157
 ————— compound, 157
 ————— chamomile, 119
 ————— cherry-laurel water, 299
 ————— chloride of barium, 143
 ————— calcium, 154
 ————— iron, 232
 ————— chlorine, 183
 ————— cider, 291
 ————— citrate of potassa and Peruvian bark, 379
 ————— cochineal and carbonate of potassa, 194
 ————— common salt, 194
 ————— cod-liver oil, 329
 ————— colchicum, 197
 ————— and ammonia, 198
 ————— elaterium, 198
 ————— magnesia, 197
 ————— sulphate of magnesia, 197
 ————— tincture, 197
 ————— with foxglove, 197
 ————— vinegar, 198
 ————— with squill, 198
 ————— wine, 198
 ————— colocynth, 199
 ————— tincture, 200
 ————— colored fire, 482
 ————— columbo, 200
 ————— and cascarilla, 201
 ————— salep, 201
 ————— common salt and lemon juice, 432
 ————— contrayerva, 203
 ————— copaiba, 205
 ————— corrosive sublimate, 269
 ————— creasote, 208, 209
 ————— croton oil, 334
 ————— cubeb, 211
 ————— oil and copaiba, 212
 ————— cyanuret of potassium, 363
 Mixture, dandelion, 456
 ————— Dippel's oil, 329
 ————— diuretic, 349
 ————— eggs, 346
 ————— and brandy, 346
 ————— wine, 346
 ————— elaterium, 222
 ————— elder, 409
 ————— elecampane, 283
 ————— emetine, 223
 ————— ergot, 224
 ————— ergotine, 224
 ————— errhine, 117
 ————— expectorant, 315
 ————— extract of lettuce, 297
 ————— ferrocyanuret of quinine, 388
 ————— zinc, 472
 ————— flaxseed, 302
 ————— Fowler's solution, 374
 ————— foxglove and acetate lead, 219
 ————— potassa, 219
 ————— tartaric acid, 219
 ————— expectorant, 219
 ————— tincture, 219
 ————— galbanum, 247
 ————— gamboge, 249
 ————— and elaterin, 250
 ————— gentian, 251
 ————— and sulphuric acid, 252
 ————— tincture, 252
 ————— guaiacum, 258
 ————— and bittersweet, 258
 ————— paregoric, 259
 ————— copaiba, 259
 ————— henbane, 259
 ————— odontalgic, 259
 ————— gum Arabic, 76
 ————— Wendt's, 77
 ————— hemlock and paregoric, 203
 ————— henbane, 281
 ————— and antimony, 281
 ————— squill, 282
 ————— Hope's, 160
 ————— hops, 264
 ————— horseradish, 128
 ————— hydrocyanic acid, 82
 ————— hyponitrous ether, 91
 ————— Iceland moss, 180
 ————— Indian sarsaparilla, 262
 ————— iodide of iron, 236
 ————— and chloride, syrup of, 237
 ————— potassium, 364
 ————— and sarsaparilla, 365
 ————— iodine, 285
 ————— and arsenic, 374
 ————— iodide of potassium, 284
 ————— ipecacuanha, 288
 ————— iron, aromatic, 228
 ————— compound, 230-241
 ————— jalap, 290
 ————— tincture, 291
 ————— Kermes mineral, 121
 ————— lactate of quinia, 389
 ————— lactucarium, 297
 ————— laudanum and tartar emetic, 345
 ————— lettuce and dandelion, 296
 ————— extract, 297
 ————— water, 298
 ————— liquorice, 255
 ————— compound, 255
 ————— lobelia, 303
 ————— logwood, extract, 260
 ————— magnesia, 305
 ————— and gentian, 306

- Mixture, magnesia and rhubarb, 306
 — malate of iron, 237
 — male fern, 244
 — manna, 311
 — mercurial, 266
 — milk and suet, 295
 — monesia, 319
 — muriate of morphia, 322
 — — quinia, 389
 — muriatic ether, 92
 — musk, 324
 — myrrh, compound, 326
 — — Griffith's, 326
 — narcotine, 327
 — neutral, 378
 — — compound, 378
 — nitrate of ammonia, 109
 — — potassa, 380
 — nitric acid, 85
 — nux vomica, 328
 — odontalgic, 172
 — oil of almonds, 114
 — — amber, 448
 — — anise, 118
 — — cajeput, 329
 — — euphorbia, 226
 — — juniper, 292
 — — mole plant, 226
 — — valerian, 464
 — oleaginous, 331
 — olive oil, 330
 — opium and cinnamon water, 341
 — — lime water, 340
 — — syrup of poppies, 340
 — pectoral, 312
 — Peruvian bark, 186-188-191
 — phosphate of soda, 438
 — pomegranate root, extract, 256
 — Prussian blue, 234
 — quince seed, 216
 — rhatany, 294
 — rhubarb, 398
 — rue, 403
 — — and squill, 403
 — scammony, 417
 — Scudamore's, 197
 — sedative, 342-348
 — semen contra, 130
 — seneka, 424
 — senna, 427
 — soot, 246
 — spermaceti, 179
 — squill, syrup, 421
 — — extract, and rhubarb, 421
 — — oxymel and ammoniac, 421
 — — — ipecacuanha, 421
 — — — marsh mallow, 421
 — — — valerian, 421
 — — vinegar, 421
 — — — and ammonia, 421
 — starch and suet, 116
 — stramonium, 444
 — strychnia, 446
 — subcarbonate of iron, 230
 — succinate of ammonia, 110
 — sugar of milk and gum Arabic, 296
 — — Iceland moss, 295
 — sulphate of cinchonia, 192
 — — iron, 241
 — — magnesia and coffee, 308
 — — — nitric acid, 308
 — — Rochelle salt, 308
 — — sulphuric acid, 308
 Mixture, sulphate of magnesia and tartar emetic, 308
 — — morphia, 323
 — — potassa, 381
 — — quinia, 392
 — — — and coffee, 392
 — — — zinc, 475
 — — sulpho-tartrate of quinia, 393
 — — sulphur, 450
 — — sulphuret of antimony, 121
 — — — and lime, 121
 — — carbon, 451
 — — potassium, 371
 — — sodium, and sal-ammoniac, 432
 — — sulphuric ether and camphor, 93
 — — — turpentine, 93
 — — sweet spirit of nitre, 91
 — — tannic acid, 88
 — — tartarized soda, 440
 — — tartrate of potassa, 382
 — — — and ammonia, 384
 — — tobacco, 453
 — — turpentine, 457
 — — — oil, 332
 — — — and ether, 333
 — — uva ursi, 462
 — — valerianate of zinc, 475
 — — valerian and ammonia, 464
 — — — Hoffmann's anodyne, 464
 — — vinegar, 77
 — — — and cardamoms, 77
 — — Virginia snakeroot, acetated, 429
 — — — and allspice, 429
 — — wine of opium, 342
 — — wormseed oil, 181
 Mode of administration of medicine, 58
 Molasses beer, 490
 — — posset, 490
 Mole plant, 226
 — — mixture of oil, 226
 Monesia, 319
 — — mixture, 319
 — — ointment, 319
 — — purified, 319
 — — syrup, 319
 — — — compound, 319
 — — tincture, 319
 Monkshood, 88
 Morphia, 319
 — — draught, 320
 — — injection, 320
 — — pills, 320
 — — acetate, 320
 — — — bolus, 320
 — — — clyster, 321
 — — — mixture, 321
 — — — ointment, 321
 — — — pills, 320
 — — — powder, compound, 320
 — — — solution, 320, 321
 — — — — alcoholic, 321
 — — — — with ipecacuanha, 320
 — — — syrup, 321
 — — — bimeconate, 321
 — — — — solution, 321
 — — — citrate, 321
 — — — hydriodate, 321
 — — — muriate, 321
 — — — lozenges, 322
 — — — — and ipecacuanha, 322
 — — — mixture, 322
 — — — solution, 322
 — — — syrup, 322
 — — — — compound, 322
 — — — nitrate, 322

Morphia, phosphate, 322
 — sulphate, 322
 — bolus, 322
 — lotion, 323
 — mixture, 323
 — pills, 322
 — solution, 323
 — syrup, 323
 — tartrate, 323
 Mosaic gold, 443
 Mucilage, gum Arabic, 76
 — mercurial, 265
 — rice, 346
 — sago, 407
 — salep, 487
 — starch, 115
 — tragacanth, 460
 Mudar, 158
 — infusion, 159
 — oil, 159
 — pills, 159
 Mulberries, 319
 — syrup, 319
 — rob, 319
 Mulled wine, 489
 Muriate of ammonia, 108
 — brucia, 150
 — cinchonia, 192
 — copper and ammonia, 214
 — manganese, 310
 — morphia, 321
 — and codeia, 195
 — narcotina, 327
 — opium, 341
 — quinia, 389
 — strychnia, 447
 — veratria, 467
 Muriatic acid, 84, 555
 — bath, 84
 — diluted, 84
 — draught, 85
 — gargle, 84
 — gaseous, 84
 — liniment, 85
 — lotion, 85
 — pediluvium, 85

Muriatic ether, 92
 Musk, 323
 — artificial, 447
 — emulsion, 448
 — tincture, 448
 — bolus, 323
 — clyster, 324
 — mixture, 324
 — pills, 323
 — powder, 323
 — tincture, 324
 Mustard, 430
 — cataplasm, 430
 — compound, 430
 — foot-bath, 430
 — gargle, 430
 — liniment, 431
 — lotion, 430
 — oil, fixed, 430
 — volatile, 431
 — tincture, 431
 — ointment, 430
 — compound, 430
 — whey, 295, 430
 Mutton tea, 488
 — broth, 489
 Myrrh, 325
 — collutory, 326
 — extract, compound, 326
 — gargle, 326
 — injection, 326
 — mixture, compound, 326
 — Griffith's, 231, 326
 — oil, 326
 — pills and Canada balsam, 325
 — squill, 325
 — sulphate of iron, 325
 — zinc, 325
 — plaster, 326
 — powder, emmenagogue, 325
 — and ipecacuanha, 325
 — saccharated, 406
 — solution, alkaline, 326
 — tincture, 326
 — and hellebore, 326

N

Naphthaline, 327
 — lozenges, 327
 — ointment, 327
 — syrup, 327
 Narcotics, 72, 565
 Narcotico-irritants, 566
 Narcotine, 327
 — bolus, 327
 — mixture, 327
 — muriate, 327
 Nervine balsam, 305
 Neutral mixture, 378
 New Jersey tea, 177
 — decoction, 177
 Nicotina, 453
 Nitrate of ammonia, 109
 — mixture, 109
 — camphor, 162
 — iron, 238
 — mercury, 276
 — acid, 276
 — morphia, 322
 — potassa, 379
 — quinia, 390

Nitrate of silver, 126
 — strychnia, 447
 — veratria, 467
 Nitre, 379
 — cubic, 438
 — fused, 379
 — purified, 379
 — sweet spirit, 91
 Nitric acid, 85, 556
 — collutory, 85
 — diluted, 85
 — fomentation, 85
 — liniment, 85
 — lotion, 85
 — mixture, 85
 — ointment, 85
 Nitrous ether, 91
 — powders, 379
 Nitro-muriatic acid, 86
 — bath, 86
 — saccharate of lead, 361
 Norwood's tincture, 468
 Nuremberg plaster, 360
 Nutmeg, 324

Nutmeg, clyster, 325
 ——— essence, 325
 ——— oil, 324
 ——— powder, anti-hectic, 325
 ——— ——— aromatic, 324
 ——— ——— saccharated, 406
 ——— spirit, 325
 ——— compound, 325
 Nux vomica, 327
 ——— extract, 328

Nux vomica, extract, pills, 328
 ——— lotion, 328
 ——— mixture, 328
 ——— pills, 328
 ——— and aloes, 328
 ——— compound, 328
 ——— powder, 327
 ——— tincture, 328
 ——— compound, 328

O

Oak bark, 386
 ——— acorn coffee, 387
 ——— cataplasm, 387
 ——— confection, 387
 ——— decoction, 387
 ——— extract, 386
 ——— gargle, 387
 ——— powder, compound, 386
 Oat, 140
 ——— cataplasm, 140
 ——— decoction, 140
 ——— flummery, 140
 ——— gruel, groat, 140
 ——— powder for cataplasm, 140
 Oatmeal gruel, 140
 Observations on management of the sick room, 55
 Official preparations and directions, 514
 Oil of almond, 114
 ——— mixture, 115
 ——— bitter, 114
 ——— amber, 447
 ——— rectified, 447
 ——— anise, 118
 ——— mixture, 118
 ——— benne, 429
 ——— benzoin, 147
 ——— black pepper, resinous, 352
 ——— ——— volatile, 352
 ——— box, 151
 ——— cajeput, 329
 ——— cantharides, 165
 ——— caraway, 172
 ——— carron, 303
 ——— castor, 331
 ——— chlorine, 183
 ——— cinnamon, 194
 ——— cloves, 172
 ——— cod liver, 329
 ——— copaiba, 206
 ——— croton, 333
 ——— cubeb, 212
 ——— dittany, 212
 ——— eggs, 346
 ——— elaterium, 222
 ——— ergot, 225
 ——— euphorbium, 227
 ——— fennel, 245
 ——— figwort, 423
 ——— ginger, 476
 ——— guaiacum, 257
 ——— compound, 257
 ——— hemlock, 202, 354
 ——— horsebalm, 198
 ——— horsemint, 318
 ——— laurel, 298
 ——— macassar, 115
 ——— mace, 305
 ——— male fern, 244
 ——— marjoram, 345
 ——— mezereon, green, 318

Oil of mudar, 159
 ——— mustard, fixed, 430
 ——— ——— volatile, 431
 ——— myrrh, 326
 ——— nutmeg, 324
 ——— olive, 330
 ——— orange flowers, 136
 ——— parsley, 349
 ——— partridge berry, 250
 ——— pennyroyal, 260
 ——— peppermint, 316
 ——— rose, 400
 ——— rosemary, 401
 ——— roses, 400
 ——— rue, 403
 ——— sassafras, 416
 ——— savine, 405
 ——— scammony, 418
 ——— soot, 246
 ——— stramonium, 445
 ——— tansy, 454
 ——— tobacco, 453
 ——— turpentine, 332
 ——— and ether, 333
 ——— ——— honey, 332
 ——— ——— clyster, 333
 ——— ——— emulsion, 332
 ——— ——— liniment, 333
 ——— ——— acetic, 333
 ——— ——— opiated, 333
 ——— ——— sulphuric, 333
 ——— lotion, 332
 ——— for toothache, 332
 ——— mixture, 332
 ——— purified, 332
 ——— soap, Starkey's, 333
 ——— wine, 333
 ——— valerian, 464
 ——— violets, 468
 ——— wine, 92
 ——— wormseed, 181
 ——— wormwood, 75
 ——— essential, 76
 ——— British, 349
 ——— Dippel's, 329
 ——— distilled, 528
 ——— essential, 528
 ——— table of, 530
 ——— fixed, 533
 ——— phosphorated, 350
 ——— volatile, 528
 Ointments, 544
 Ointment, ammoniacal, 106
 ——— Gondret's, 106
 ——— Anglo-Saxon, 102
 ——— anodyne, 339
 ——— arsefical, 80
 ——— Banyer's, 360
 ——— basilicon, 393
 ——— citrine, 276
 ——— Egyptian, 213

- Ointment, for chilblains, 102
 ———— issue, 166
 ———— itch, 412
 ———— Kentish,
 ———— pile, 103,
 ———— Rust's, 102
 ———— sultana, 180
 ———— of acetate of lead, 357
 ———— ———— morphia, 321
 ———— aconite, 89
 ———— ———— ammoniated, 90
 ———— aconitine, 90
 ———— aloes, 98
 ———— alum, 102
 ———— American hellebore, 468
 ———— ammoniated copper, 214
 ———— animal charcoal, 168
 ———— arseniate of iron, 230
 ———— azedarach, 140
 ———— balsam of Peru, 142
 ———— ———— compound, 142
 ———— belladonna, 146
 ———— bichloride of platinum, 355
 ———— biniodide of potassium, 368
 ———— ———— and opium, 368
 ———— black hellebore, 261
 ———— ———— oxide of mercury, 277
 ———— ———— pepper, 352
 ———— ———— pitch, 355
 ———— borax, 434
 ———— bromide of iron, 230
 ———— ———— potassium, 362
 ———— ———— ———— compound, 362
 ———— ———— sodium, 431
 ———— butter cocoa, 152
 ———— calomel, 273
 ———— ———— and acetate of copper, 273
 ———— ———— alum, 273
 ———— ———— camphor, 274
 ———— ———— squill, 273
 ———— camphor, 161
 ———— cantharides, 165
 ———— carbonate of ammonia, 106
 ———— ———— lead, 358
 ———— ———— ———— camphorated, 359
 ———— ———— ———— potassa, 377
 ———— ———— ———— soda, 435
 ———— carburet of iron, 232
 ———— carrot, 171
 ———— cevadilla, 404
 ———— ———— extract, 404
 ———— charcoal, 169
 ———— cherry-laurel, 299
 ———— chloride of antimony, 119
 ———— ———— gold, 137
 ———— ———— ———— and sodium, 138
 ———— ———— ———— lead, 359
 ———— chlorinated lime, 155
 ———— chlorine, 183
 ———— cocculus indicus, 194
 ———— cod liver oil, 330
 ———— ———— compound, 330
 ———— common salt, 432
 ———— ———— compound, 432
 ———— cowhage, 324
 ———— creasote, 209
 ———— ———— compound, 209
 ———— croton oil, 335
 ———— cucumber, 212
 ———— cyanide of silver, 126
 ———— cyanuret of mercury, 274
 ———— ———— potassium, 363
 ———— delphinia, 217
 ———— elder flowers, 409
 ———— ———— leaves, 409
 ———— elemi, 222
 Ointment of fennel, 245
 ———— figwort, 423
 ———— foxglove, 219
 ———— fuligokali, 246
 ———— galls, 249
 ———— ———— compound, 249
 ———— ———— extract, 249
 ———— glycerin, 254
 ———— gold, 137
 ———— hemlock, 202
 ———— henbane, 282
 ———— ———— compound, 282
 ———— hops, 264
 ———— hydrargyro-iodide of potassium, 369
 ———— hydriodate of ammonia, 198
 ———— iodide of arsenic, 129
 ———— ———— barium, 143
 ———— ———— carbon, 170
 ———— ———— iron, 237
 ———— ———— lead, 360
 ———— ———— mercury, 275
 ———— ———— potassium, 365
 ———— ———— ———— and opium, 365
 ———— ———— ———— mercury, 365
 ———— ———— ———— morphia, 366
 ———— ———— ———— silver, 126
 ———— ———— ———— sulphur, 452
 ———— ———— ———— zinc, 472
 ———— iodine, 285
 ———— ———— compound, 285, 368
 ———— ———— with mercury, 285
 ———— ———— ———— oil of tobacco, 285
 ———— laudanum, 344
 ———— laurel, 298
 ———— lavender, 300
 ———— lead, 360
 ———— ———— compound, 360
 ———— lime, 156
 ———— lupulin, 304
 ———— marsh mallow, 99
 ———— matico, 314
 ———— mercurial, 266
 ———— ———— camphorated, 267
 ———— ———— compound, 267
 ———— ———— and belladonna, 267
 ———— of mezereon, 318
 ———— monesia, 319
 ———— mustard, 430
 ———— ———— compound, 430
 ———— naphthaline, 327
 ———— nitrate of bismuth, 148
 ———— ———— mercury, 276
 ———— ———— ———— and lead, 276
 ———— ———— ———— silver, 127
 ———— ———— ———— compound, 127
 ———— ———— ———— strychnia, 447
 ———— nitric acid, 85
 ———— opium and tar, 340
 ———— oxide of antimony, 119
 ———— ———— manganese, 310
 ———— ———— ———— and sulphur, 310
 ———— ———— ———— silver, 126
 ———— ———— ———— zinc, impure, 470
 ———— petroleum, 349
 ———— phosphorated, 350
 ———— of pierotoxin, 194
 ———— poke, 351
 ———— Prussian blue, 234
 ———— red iodide of mercury, 276
 ———— ———— oxide of mercury, 277
 ———— ———— ———— and basilicon, 278
 ———— ———— ———— cinnabar, 278
 ———— ———— ———— lead, 278

- Ointment of red oxide of mercury and sulphur, 277
 ————— tin, 278
 ————— zinc, 278
 ————— sulphuret of mercury, 280
 ————— rhatany, compound, 294
 ————— rose water, 399
 ————— saffron, 210
 ————— savine, 405
 ————— soot, 246
 ————— compound, 246
 ————— spermaceti, 179
 ————— and rose water, 179
 ————— squill, 422
 ————— stavesacre, 443
 ————— stramonium, 445
 ————— strychnia, 446
 ————— subacetate of lead, 358
 ————— sulphate of cadmium, 153
 ————— copper, 216
 ————— mercury, 279
 ————— quinia, 392
 ————— sulphur, 450
 ————— and camphor, 451
 ————— soap, 451
 ————— zinc, 451
 ————— compound, 450
 ————— sulphuret of antimony, 120
 ————— potassium, 371
 ————— sodium, 432
 ————— sulphuric acid, 87
 ————— tannate of lead, 361
 ————— tannic acid, 88
 ————— tar, 355
 ————— compound, 355
 ————— tartar emetic, 123
 ————— compound, 123
 ————— tin, 442
 ————— tobacco, 453
 ————— turpentine, 457
 ————— compound, 457
 ————— tutty, 470
 ————— veratria, 466
 ————— and iodine, 467
 ————— opium, 467
 ————— verdigris, 213
 ————— white hellebore, 467
 ————— compound, 467
 ————— white precipitate, 268
 ————— wild indigo, 142
 ————— willow leaves, 408
 ————— zinc, oxide, 470
 ————— and calomel, 471
 ————— opium, 470
 Oleaginous mixture, 331
 Oleo-saccharated powders, 406
 Olive oil, 330
 ————— clyster, 330
 ————— mixture, 330
 Onguent de la mère, 360
 Opiate, anti-dysenteric, 339
 Opium, 335, 565
 ————— balsam, 339
 ————— cerate, 340
 ————— clyster, 340
 ————— collyrium, 340
 ————— confection, 338
 ————— draught, anodyne, 344
 ————— drops, 339
 ————— elixir, 336
 ————— Lettsom's, 344
 ————— extract, acetous, 335
 ————— alcoholic, 336
 ————— aqueous, 335
 ————— aqueo-alcoholic, 335
 ————— denarcotized, 336
 Opium, extract, fermented, 336
 ————— roasted, 336
 ————— vinous, 336
 ————— Godfrey's cordial, 343
 ————— fomentation, 340
 ————— injection, 340
 ————— linctus, 340, 344
 ————— liniment, 339
 ————— with oil of chamomile, 339
 ————— lozenges, Wistar's, 345
 ————— mass, anti-odontalgic, 338
 ————— mixture, anti-otitic, 339
 ————— brown cough, 344
 ————— with cinnamon water, 341
 ————— lime water, 340
 ————— syrup of poppies, 340
 ————— sedative, 342
 ————— muriate, 341
 ————— ointment, 339
 ————— and tar, 340
 ————— pills, 337, 343
 ————— aromatic, 337
 ————— with acetate of lead, 338
 ————— mercury, 338
 ————— butter of cocoa, 337
 ————— camphor, 337
 ————— castor, 338
 ————— foxglove, 337
 ————— hemlock and calomel, 338
 ————— henbane, 337
 ————— liquorice, 338
 ————— musk, 337
 ————— nitrate of silver, 338
 ————— soap, 343
 ————— tartar emetic, 122
 ————— sulphate of zinc, 338
 ————— sulphuret of antimony, 337
 ————— plaster, 339
 ————— with camphor, 339
 ————— powder, compound, with chalk, 336
 ————— Dover's, 286
 ————— eclectic, 345
 ————— roasted, 336
 ————— with antimony, 337
 ————— camphor, 337
 ————— musk, 337
 ————— nitre, 337
 ————— sulphur, 337
 ————— suppository, 340
 ————— syrup, 340
 ————— succinated, 340
 ————— tincture, 342
 ————— acetated, 342
 ————— ammoniated, 343
 ————— and soap, 343
 ————— tartar emetic, 345
 ————— Bateman's, 343
 ————— Battley's, 343
 ————— camphorated, 343
 ————— clyster, 344
 ————— and valerian, 344
 ————— compound, 343
 ————— liniment, 344
 ————— and lime water, 344
 ————— ointment, 344
 ————— Smith's, 343
 ————— succinated, 343
 ————— Swediaur's, 344
 ————— Warner's, 344
 ————— vinegar, or black drop, 341
 ————— Guy's hospital, 341
 ————— Houlton's, 341
 ————— Lancaster, 341
 ————— mixture, 342
 ————— Porter's, 341
 ————— Rousseau's, 341

Opium, water, 341
 — wine, 342
 — collyrium, 342
 — drops, odontalgic, 342
 — fomentation, 342
 — mixture, 342
 — Sydenham's, 342
 Opodeldoc, liquid, 104, 411
 — Steer's, 104
 Opoponax, 345
 — emulsion, 345
 — tincture, 345
 — compound, 345
 Orange, 135
 — flowers, 136
 — confection, 136
 — oil, 136
 — pastilles, 136
 — syrup, 136
 — water, 136
 — leaves, 136
 — electuary, 136
 — jelly, 136
 — peel, 135
 — confection, 135
 — electuary, 135
 — elixir, 135
 — infusion, compound, 135
 — powder, with rhubarb, 135
 — spirit, 135

Orange peel, syrup, 136
 — tincture, 135
 Orgeat, syrup of, 113
 Oriental pills, 337
 — water, 376
 Orpiment, 130
 — depilatory, 130
 — lotion, 130
 — powder, 130
 Orris root, 289
 — dentifrice, 289
 Oxalate of potassa, 380
 Oxalic acid, 86, 556
 — lozenges, 86
 Oxide, antimony, nitro-muriatic, 119
 — gold, 139
 — iron, hydrated, 238
 — lead, 360
 — manganese, 310
 — mercury, 277
 — silver, 126
 — tin, 443
 — zinc, 470
 Oxymel, 314
 — bitartrate of potassa, 383
 — colchicum, 197
 — elecampane, compound, 284
 — squill, 422
 Oyster-shell, prepared, 156

P

Palamoud, 151
 Panada, 487
 — chicken, 489
 Paregoric, elixir, 343
 — Scotch, 343
 Paraguay-roux, 385
 Pareira brava, 348
 — decoction, 348
 — extract, 348
 — infusion, 348
 — tincture, 348
 Parrish's camphor mixture, 160
 Parsley, 349
 — infusion, 349
 — oil, 349
 Partridge-berry, 250
 — oil, 250
 — water, 250
 Pastes, 534
 Paste, almond, 112
 — chloride of zinc, 471
 — cocoa, compound, 152
 — copaiba, 206
 — de guimauve, 76
 — fig, 243
 — glycerin, 254
 — liquorice, 254
 — marshmallow, 99
 — anti-odontalgic, 313
 — pectoral, of gum Arabic, 76
 — phosphorus, 350
 — tragacanth, 460
 — rubefacient, 356
 — Ward's, 352
 Pastilles, catechu, 177
 — de Paris, 485
 — de santé, 418
 — fumigating, 147
 — orange flowers, 136
 — purgative, 418
 — vinegar, 77

Paullinia, 348
 — chocolate, 348
 — extract, alcoholic, 348
 — lozenges, 348
 — pills, 348
 — syrup, 348
 Pearson's solution, 433
 Pediluvium, mustard, 430
 — muriatic acid, 85
 Pellitory, 385
 — extract, 385
 — liniment, 385
 — lozenges, 385
 — plaster, 385
 — tincture, 385
 — compound, 385
 Pennyroyal, 260
 — infusion, 260
 — oil, 260
 Pepper, black, 351
 — cubebs, 210
 — long, 352
 — Jamaica, 351
 — white, 351
 Peppermint, 316
 — essence, 316
 — lozenges, 316
 — mixture, 316
 — oil, 316
 — spirit, 316
 — water, 316
 Persimmon, 220
 — infusion of bark, 220
 — wine of fruit, 220
 Persulphate of mercury, 278
 Peruvian bark, 185
 — beer, 191
 — cataplasm, 190
 — cerate, 190
 — clyster, 190
 — decoction of yellow, 187

- Peruvian bark, decoction of, compound, 188
 mixture, 188
 with cascarilla, 188
 rhatany, 188
 electuary, 189
 astringent, 189
 with catechu, 189
 cloves, 189
 iron, 189
 sulphur, 189
 tin, 189
 extract, 187
 fluid, 187
 bolus, 187
 pills, 187
 vinous, 187
 gargle, 190
 infusion of yellow, 188
 compound, 188, 189
 with limewater, 188
 magnesia, 188
 snakeroot, 188
 valerian, 188
 lotion, 190
 mixture, 186, 191
 pomatum, 190
 powder, compound, 185
 dentifrice, 186
 with arnica, 186
 camphor, 186
 cascarilla, 186
 isinglass, 186
 valerian, 186
 saccharated, 406
 salt, 187
 syrup, 191
 compound, 191
 tincture, 189
 compound, 189
 ammoniated, 190
 with cantharides, 190
 gentian, 190
 snakeroot, 190
 valerian, 190
 wine, 191
 and calamus, 191
 iron, 191
 Petroleum, 349
 anthelmintic mixture, 349
 British oil, 349
 diuretic mixture, 349
 embrocation, 349
 ointment, 349
 plaster, 349
 Pewter, solder for, 485
 Pharmaceutical names, table of, 503
 Phloridzine, 350
 powder, 350
 Phosphate of ammonia, 109
 solution, 109
 iron, 240
 lime, 157
 manganese, 311
 magnesia, 308
 mercury, 278
 morphia, 322
 quinia, 390
 soda, 438
 Phosphoric acid, 86
 diluted, 86
 fomentation, 86
 lemonade, 86
 pills, 86
 tincture of myrrh, 86
 Phosphorated ether, 350
 liniment, 350
 Phosphorated oil, 350
 turpentine, 350
 ointment, 350
 Phosphorus, 350, 564
 cerate, 350
 paste, 350
 Picrotoxin, 194
 Pills, 515
 Abernethy's, 266
 acetate of copper, 213
 lead, 356
 mercury, 268
 and opium, 268
 morphia, 320
 potassa, compound, 373
 soda, compound, 433
 aconite, extract, 89
 agaric and opium, 149
 aloes, 94
 and assafetida, 94
 blue mass, 96
 iron, 94
 myrrh, 94
 rhubarb, 96
 Anderson's, 95
 antichlorotic, 96
 aperient, 94, 96
 Barthez's, 96
 Bicker's, 96
 Chapman's anti-dyspeptic, 94
 compound, 94
 dinner, 313
 Duchesne's, 96
 Frank's, 96
 Fuller's, 95
 Griffitt's, 96
 James's analeptic, 95
 Hooper's, 95, 241
 Lady Webster's, 95
 Mitchell's, 95
 Morrison's, 95
 Peters's, 95
 Pitschaft's eccoprotic, 96
 Rufus's, 94
 Speediman's, 95, 305
 splenitic, 95
 Whytt's, 96
 alum, 100
 and benzoic acid, 101
 astringent, 101
 American hellebore, 468
 ammoniac, 110
 and rhubarb, 111
 compound, 111
 De Haen's, 111
 Klein's, 111
 ammoniated copper, 214
 compound, 214
 gold, 139
 iron, 229
 anemone, 116
 antibilious, 199, 273
 antimonial powder and calomel, 124
 arsenate of iron, 229
 arsenic and opium, 80
 pepper 80
 arsenical, 80
 assafetida, 132
 and aloes, 134
 lactucarium, 132
 iron, 132
 musk, 133
 opium, 133
 Bacher's, 261
 balsam of Peru, 141
 belladonna, compound, 145

- Pills, belladonna and camphor, 145
 — bichloride of platinum, 355
 — bittersweet, extract, 221
 — black hellebore, 261
 — ———— extract of, 261
 — black oxide of mercury, 277
 — ———— iron, 239
 — ———— pitch, 355
 — Bland's antichlorotic, 240
 — bloodroot, 409
 — blue, or mercurial, 266
 — ———— and colocynth, 266
 — ———— jalap, 266
 — ———— quinine, 266
 — ———— rhubarb, 266
 — ———— compound, 266
 — borotartarate of potassa, compound, 374
 — bromide of iron, 230
 — brucia, 150
 — buckbean, compound, 317
 — burgundy pitch, 354
 — calomel, 272
 — ———— and acetate of lead, 272
 — ———— antimony, 273
 — ———— catechu, 272
 — ———— colocynth, 273
 — ———— dandelion, 272
 — ———— guaiacum, 273
 — ———— iron, 272
 — ———— jalap, 273
 — ———— opium, 273
 — ———— quinine, 272
 — ———— squill, 272
 — ———— compound, 272
 — ———— cathartic, 273
 — camphor, 159
 — ———— compound, 159
 — ———— and lactucarium, 159
 — ———— musk, 159
 — cantharides and capsicum, 163
 — ———— iron, 163
 — carbonate of ammonia, 106
 — ———— iron, 230
 — ———— compound, 231
 — ———— and manganese, 309
 — ———— soda, 435
 — carburet of iron, 232
 — castor, 174
 — ———— and succinic acid, 174
 — catechu and alum, 176
 — ———— liquorice, 176
 — cathartic, compound, 199, 273
 — Cayenne pepper, 167
 — celandine, 181
 — centaury, extract, 178
 — chamomile, compound, 119
 — chloride of barium, 142
 — ———— calcium, 154
 — ———— gold, 137
 — ———— and sodium, 138
 — ———— mercury and quinia, 274, 389
 — ———— silver and ammonia, 125
 — citrate of caffeine, 195
 — colchicum, 196
 — ———— compound, 196
 — colocynth, compound, 199
 — ———— and henbane, 199
 — columbo, compound, 200
 — copaiba, 204
 — ———— and cubebs, 204
 — ———— and turpentine, 204
 — corrosive sublimate, 269
 — ———— compound, 269
 — creasote, 208
 — croton oil, 333
 — ———— compound, 334
 Pills, croton oil, and blue mass, 334
 — ———— quinia, 334
 — cyanuret of gold, 139
 — ———— zinc, 472
 — ———— mercury, compound, 274
 — ———— potassium, 363
 — dandelion, 455
 — ———— and blue mass, 456
 — digitalin, 220
 — dinner, 313
 — dogwood, round-leaved, 208
 — elaterium, 222
 — elecampane, compound, 283
 — ergot, extract of, 225
 — ———— compound, 224
 — ferrocyanate of quinia, 388
 — ferrocyanuret of zinc, compound, 472
 — foxglove, 218
 — ———— and squill, 218
 — fuligokali, sulphuretted, 246
 — galbanum, compound, 247
 — gallic acid, 82
 — gamboge, compound, 249
 — gentian, compound, 250
 — gentianin, 252
 — guaiacum and aloes, 258
 — ———— antimony, 258
 — ———— sulphur, 258
 — ———— turpentine, 258
 — hemlock and calomel, 202
 — ———— dandelion, 202
 — ———— ipecacuanha, 202
 — henbane, compound, 281
 — ———— and ipecacuanha, 281
 — ———— opium, 281
 — hydrargyro-iodide of potassium, 368
 — iodide of arsenic, 129
 — ———— calcium, 155
 — ———— iron, 235
 — ———— lead, 359
 — ———— manganese, 310
 — ———— mercury, 275
 — ———— compound, 275
 — ———— potassium, 364
 — iodine, 284
 — indigo, 283
 — ipecacuanha and centaury, 287
 — ———— foxglove, 287
 — ———— opium, 287
 — ———— squill, 287
 — iron, prepared, 228
 — jalap and calomel, 290
 — ———— compound, 290
 — Keyser's, 268
 — kinate of quinia, 389
 — kino, compound, 293
 — lactate of iron, 237
 — ———— quinia, 389
 — lactucarium, 297
 — Lartigue's, 196
 — Lockstadt's, 115
 — lupulin, 304
 — male fern, extract, 244
 — ———— oil, 244
 — marigold, extract, 158
 — mercury and antimony, 265
 — ———— gum, with hemlock, 265
 — morphia, 320
 — mudar, 159
 — muriate of manganese, 310
 — musk, 323
 — myrrh and Canada balsam, 325
 — ———— squill, 325
 — ———— sulphate of iron, 325
 — ———— zinc, 325
 — nitrate of bismuth, 148

- Pills, nitrate of potassa, 379
 — and camphor, 380
 — silver, 126
 — nux vomica, 328
 — and aloes, 328
 — compound, 328
 — extract, 328
 — opium, 337
 — and acetate of lead, 338
 — mercury, 338
 — butter of cacao, 337
 — camphor, 337
 — castor, 338
 — foxglove, 337
 — hemlock and calomel, 338
 — henbane, 337
 — liquorice, 338
 — musk, 337
 — nitrate of silver, 338
 — sulphate of zinc, 338
 — sulphuret antimony, 337
 — aromatic, 337
 — oxide, black, of iron, 239
 — gold, 139
 — red, of iron, 239
 — silver, 126
 — zinc, 470
 — compound, 470
 — paullinia, 348
 — Peruvian bark, extract, 187
 — phosphate of mercury, 278
 — phosphoric acid, 86
 — piperine, 353
 — and mercury, 353
 — pipsissewa, 182
 — Plummer's, 272
 — podophyllin, compound, 362
 — and iron, 362
 — and may apple, 362
 — poison oak, 460
 — proto-nitrate of mercury, 276
 — Prussian blue, 234
 — red iodide of mercury, 275
 — oxide of mercury, 277
 — rhatany, 294
 — rhubarb, 395
 — compound, 395
 — and caraway, 395
 — chamomile, 395
 — ipecacuanha, 395
 — iron, 396
 — ox-gall, 396
 — soda, 396
 — saffron, 210
 — sagapenum, compound, 406
 — salicine, 407
 — compound, 407
 — savine, 405
 — scammony, 417
 — and ox-gall, 417
 — compound, 417
 — seneka, 423
 — senna, 425
 — soap, 411
 — and ox-gall, 411
 — compound, 343, 411
 — soda and ipecacuanha, 435
 — rhubarb, 435
 — soot, 245
 — squill and ammoniac, 419
 — calomel, 419
 — croton oil, 419
 — ipecacuanha, 419
 — henbane, 419
 — compound, 419
 — storax, 445
- Pills, stramonium, 444
 — compound, 444
 — strychnia, 446
 — stimulant, 446
 — sulphate of bebeerine, 144
 — cinchonia, 192
 — copper, 215
 — and opium, 215
 — compound, 215
 — iron, 240
 — and rhubarb, 240
 — compound, 240
 — morphia, 322
 — potassa, 381
 — quinia, 391
 — compound, 391
 — with gentian, 391
 — soda, compound, 439
 — zinc, 474
 — compound, 474
 — with myrrh, 474
 — turpentine, 474
 — sulphuret of antimony, 120
 — and lime, 121
 — calcium, compound, 155
 — mercury, black, 279
 — red, 279
 — potassium, 369
 — compound, 369
 — tannic acid, 87
 — and opium, 87
 — tansy, 454
 — tar, 354
 — tartar emetic and camphor, 122
 — guaiacum, 122
 — opium, 122
 — tobacco, 452
 — turpentine, 456
 — and guaiacum, 456
 — jalap, 456
 — magnesia, 456
 — myrrh, 456
 — rhubarb, 456
 — valerian, compound, 463
 — valerianate of quinia, 393
 — zinc, 475
 — veratria, 466
 — and henbane, 466
 — Virginia snakeroot, compound, 428
- Pimento, 351
 Pink, 217
 — syrup, 217
 Pinkroot, 440
 — and calomel, 440
 — savine, 440
 — extract, fluid, 441
 — compound, 441
 — infusion, 440
 — compound, 441
- Piperine, 352
 — pills, 353
 — and mercury, 353
 — tincture, 353
- Pipsissewa, 181
 — beer, 182
 — decoction, 182
 — compound, 182
 — extract, 182
 — pills, 182
 — syrup, 182
- Pitch, black, 355
 — ointment, 355
 — plaster, 355
 — Burgundy, 353
 — cerate, 354
 — pills, 354

- Pitch, Burgundy, plaster, 353
 aromatic, 353
 compound, 354
 Canada, 354
 Plasters, 545
 Plaster, acetate of lead, 357
 aconite, 89
 adhesive, 394
 Baynton's, 360
 ammonia, 104
 ammoniac, 112
 with hemlock, 112
 mercury, 112
 assafetida, 134
 balsam of Peru, 142
 belladonna, 146
 biniodide of potassium, 368
 black pepper, 352
 pitch, 355
 Burgundy pitch, 353
 cantharides, 166
 camphorated, 166
 compound, 166
 odontalgic, 166
 carbonate of ammonia, 106
 lead, 359
 zinc, 474
 carburet of iron, 232
 castor, compound, 175
 cinnamon, spice, 194
 clove, 172
 court, 282
 blistering, 166
 croton oil, 335
 cumin, 216
 diapalma, 360
 elemi, cautery, 223
 euphorbium, 227
 foxglove, 219
 galbanum, 247
 compound, 247
 Gaulthier's, 360
 gum, 112
 hemlock, 202
 compound, 202
 iodine, 285
 ioduretted potassium, 368
 iron, 231
 lead, 360
 Mahy's, 359
 mercurial, 267
 and belladonna, 267
 myrrh, 326
 Nuremberg, 360
 opium, 339
 and camphor, 339
 pellitory, 385
 petroleum, 349
 pitch, 353
 aromatic, 353
 compound, 354
 red oxide of iron, 239
 resin, 394
 soap, 412
 spice, 194-477
 St. Andrew's, 394
 subacetate of lead, 358
 sulphate of quinia, 392
 sulphuret of antimony and lime, 121
 potassium, compound, 371
 tartar emetic, 123
 turpentine, 457
 compound, 457
 verdigris, 213
 vigo with mercury, 394
 warming, 353
 Plaster, wax, 178
 Platina, 355
 bichloride, 355
 mixture, 355
 ointment, 355
 pills, 355
 Pleurisy root, 131
 infusion, 131
 Plummer's alterative, 271
 pills, 272
 Podophyllin, 361
 pills, compound, 362
 and iron, 362
 may apple, 362
 Poids de Marc, 20
 Poison-oak, 460
 extract, 460
 pills, 460
 tincture, 460
 Poison for rats, &c., 480
 Poisons, 553
 Poke, 350
 ointment of, 351
 tincture, 351
 Polygalic acid, 424
 Poinatum, 485
 cantharides, 166
 Peruvian bark, 190
 Pomegranate, 256
 bolus of root, 256
 decoction of rind, 256
 root, 256
 extract of root, 256
 electuary, 256
 mixture, 256
 gargle of rind, 256
 injection of rind, 256
 Pommade epispastique au garou, 318
 Poppy heads, 347
 decoction, 347
 extract, 347
 injection, sedative, 348
 mixture, sedative, 348
 syrup, 347
 substitute for, 347
 pectoral, 347
 red, 399
 Portland powder, 178
 Posological table, 496
 Posset, molasses, 490
 sago, 407
 Potassa, 371, 558
 acetate, 372
 scillitic, 372
 bolus, 372
 conservé, 373
 liquid, 373
 mixture, 373
 mixture, 373
 pills, compound, 373
 tincture, 373
 arsenate, 374
 mixture, 374
 arsenite, solution, 373
 bicarbonate, 377
 effervescing powders, 377
 solution, 377
 mixture, 377
 bichromate, 561
 bisulphate, 381
 disinfecting powder, 382
 bitartrate, 382
 dentifrice, 382
 electuary, 383
 imperial drink, 383
 mixture, 383

- Potassa, bitartrate, oxymel, 383
 powder, 383
 solution, 383
 compound, 383
 borate, 374
 borotartrate, 374
 and magnesia, 375
 mixture, 375
 mixture, 375
 pills, compound, 374
 powder, 374
 solution, 375
 carbonate, 375
 and ammonia, 377
 collutory, 376
 collyrium, 375
 emulsion, 376
 fomentation, 376
 liniment, 376
 lotion, 376
 mixture, 376
 ointment, 377
 powder, 375
 pure, 375
 solution, 375
 tincture, 376
 compound, 377
 caustic, 371
 collyrium, 372
 injection, 372
 lotion, stimulant, 372
 mixture, 372
 solution, 372
 lithontriptic, 372
 chlorate, 378
 powder, 378
 compound, 378
 Stephens's saline, 378
 solution, 378
 citrate, 378
 effervescing draught, 378
 mixture, 379
 solution, 378
 compound, 378
 iodate, 379
 syrup, 379
 nitrate, 379
 emulsion, 380
 fused, 379
 gargle, 380
 lotion, 380
 mixture, 380
 pills, 379
 with camphor, 380
 powder, 379
 with camphor, 379
 orris root, 379
 squill, 379
 nitrate, purified, 379
 oxalate, 380
 lozenges, 380
 powder, 380
 silicate, 381
 soluble glass, 381
 solution, 381
 sulphate, 381
 and ammonia, 382
 magnesia, 382
 sulphur, 382
 draught, 381
 mixture, 381
 pills, 381
 powder, compound, 381
 and rhubarb, 381
 tartrate, 382
 and ammonia, 383
- Potassa, tartrate, and ammonia, mixture, 384
 solution, 384
 mixture, 382
 powder, and rhubarb, 382
 with lime, 372
 solution, 372
- Potassium, 362
 ioduretted, 366
 bath, 367
 collyrium, 367
 injection, 367
 lotion, compound, 367
 mixture, 368
 ointment, 368
 and opium, 368
 plaster, 368
 solution, compound, 367
 caustic, 367
 syrup, 367
 tincture, compound, 367
 water, 366
 Lugol's, 366
 bromide, 362
 ointment, 362
 compound, 362
 solution, 362
 chloride, 362
 solution, 363
 cyanuret, 363
 lotion, 363
 mixture, 363
 ointment, 363
 pills, 363
 solution, 363
 syrup, 363
 hydrargyro-iodide, 368
 ointment, 369
 pills, 368
 solution, 368
 tincture, 368
 iodide, 364, 562
 and sarsaparilla, 365
 injection, 364
 liniment, 366
 lozenges, 364
 mixture, 364
 ointment, 365
 and mercury, 365
 morphia, 366
 opium, 365
 pills, 364
 plaster, 366
 soap, 366
 solution, 364
 syrup, 365
 sulpho-cyanuret, 369
 sulphuret, 369
 and cream of tartar, 369
 bath, 371
 bolus, 369
 electuary, 370
 ferro-sulphuret 369
 hyposulphited, 369
 syrup, 370
 injection, 371
 lotion, 370
 compound, 370
 mixture, 371
 ointment, 371
 pills, 369
 compound, 369
 plaster, compound, 371
 soap, 371
 solution, 370

- Potassium, ferro-sulphuret, syrup, 370
 ————— compound, 370
 ————— tincture, 370
 ————— water, 370
- Potato, wild, 204
- Potion of carbonate of ammonia, 105
- Poudre subtile, 130
- Poultices, 541
- Powders, 514
 ————— compound, 514
 ————— effervescing, 88
 ————— nitrous, 379
 ————— oleo-saccharated, 406
 ————— saccharated, 406
 ————— simple, 514
- Powder, acetate of lead, 356
 ————— morphia, compound, 320
 ————— aconite, compound, 89
 ————— agaric, 149
 ————— and opium, 149
 ————— algaroth, 119
 ————— almond, 112
 ————— aloes, 94
 ————— and canella, 94, 162
 ————— compound, 94
 ————— emmenagogue, 94
 ————— alum, 100
 ————— escharotic, 100
 ————— ammoniated copper and belladonna, 214
 ————— iron, 229
 ————— animal charcoal, 168
 ————— anthracokali, 169
 ————— compound, 169
 ————— antihectic, 325
 ————— antimonial, 124
 ————— arnica, compound, 128
 ————— aromatic, 170, 324
 ————— arsenical, 80
 ————— arsenic, compound, 80
 ————— asarabacca, 131
 ————— avens, compound, 253
 ————— belladonna, 144
 ————— and rhubarb, 145
 ————— benzoic acid and ipecacuanha, 80
 ————— benzoin, 147
 ————— bicarbonate of potassa, 377
 ————— soda and magnesia, 436
 ————— bismuth, compound, 148
 ————— bitartrate of potassa, 383
 ————— black oxide of iron, 239
 ————— sulphuret of mercury, 279
 ————— borax, compound, 433
 ————— borotartrate of potassa, 374
 ————— bronze, 443
 ————— burnt sponge, 442
 ————— compound, 442
 ————— calomel and antimony, 271
 ————— jalap, 271
 ————— foxglove, 271
 ————— nitrate of potassa, 271
 ————— opium, 271
 ————— gamboge, 271
 ————— pinkroot, 272
 ————— antimony and henbane, 271
 ————— jalap and rhubarb, 271
 ————— camphor, 159
 ————— compound, 159
 ————— cantharides and camphor, 163
 ————— savine, 163
 ————— capuchin, 404
 ————— carbonate of lead, camphorated, 358
 ————— potassa, 375
 ————— soda, effervescing, 436
 ————— and mercury, 435
 ————— rhubarb, 435
- Powder, carburet of iron, 232
 ————— cascarilla, compound, 173
 ————— castor, compound, 174
 ————— catechu, compound, 176
 ————— cevadilla, compound, 404
 ————— chalk, aromatic, 157
 ————— compound, 157
 ————— with opium, 167
 ————— charcoal and quassia, 165
 ————— chlorate of potassa, 373
 ————— chloride of barium, 142
 ————— gold, 137
 ————— and sulfur, 137
 ————— silver, 125
 ————— cinnamon, compound, 192
 ————— cocoa, compound, 151
 ————— colocynth, compound, 199
 ————— columbo and iron, 200
 ————— magnesia, 200
 ————— tartrate of iron, 200
 ————— common salt and cochineal, 431
 ————— compound, 431
 ————— contrayerva, compound, 203
 ————— coriander, compound, 206
 ————— corrosive sublimate and copper, 263
 ————— zinc, 269
 ————— cubebs, 210
 ————— and alum, 210
 ————— ergot, 210
 ————— hemlock, 210
 ————— cyanide of gold, 138
 ————— disinfecting, 382
 ————— Dover's, 286
 ————— eclectic, 345
 ————— emmenagogue, 325
 ————— ergot, compound, 224
 ————— extract of lettuce, 297
 ————— fennel, compound, 244
 ————— foxglove, 217
 ————— fumigating, 147, 280, 447
 ————— galls, compound, 248
 ————— gamboge, compound, 249
 ————— gentian, compound, 250
 ————— ginger beer, 436
 ————— gold, 136
 ————— golden sulphuret of antimony, 121
 ————— guaiacum, compound, 258
 ————— hæmostatic, 393
 ————— hedge hyssop, 257
 ————— hemlock, 201
 ————— henbane, compound, 281
 ————— horse-chestnut, compound, 262
 ————— indigo, 283
 ————— iodide of barium, 143
 ————— lead, 359
 ————— mercury, 275
 ————— sulphur, 452
 ————— ipecacuanha and antimony, 286
 ————— calomel, 286
 ————— carbonate of soda, 286
 ————— muriate of ammonia, 286
 ————— myrrh, 287
 ————— nitre, 286
 ————— opium, 286
 ————— rhubarb, 286
 ————— tartar emetic, 286
 ————— tragacanth, 287
 ————— jalap and calomel, 289
 ————— cream of tartar, 289
 ————— ipecacuanha, 289
 ————— scammony, 289
 ————— compound, 290
 ————— James's, 124
 ————— kermesine, 120
 ————— compound, 120

- Powder, kino, compound, 293
 — lavender, compound, 299
 — lupulin, 304
 — madder, compound, 402
 — magnesia and orange peel, 305
 — rhubarb, 305
 — sulphur, 305
 — male fern, 244
 — marsh mallow, 99
 — mercurial, compound, 264
 — metallic iron, 227
 — compound, 228
 — milk, 295
 — musk, 323
 — myrrh and ipecacuanha, 325
 — nitrate of potassa and camphor, 379
 — orris root, 379
 — squill, 379
 — compound, 379
 — nux vomica, 327
 — oak bark, compound, 386
 — opium and antimony, 337
 — camphor, 337
 — chalk, compound, 336
 — musk, 337
 — nitre, 337
 — sulphur, 337
 — roasted, 336
 — orange peel and rhubarb, 135
 — orpiment, 130
 — oxalate of potassa, 380
 — oxide of gold, 139
 — manganese, 310
 — Peruvian bark, and arnica, 186
 — camphor, 186
 — cascarilla, 186
 — isinglass, 186
 — valerian, 186
 — compound, 185
 — phloridzine, 350
 — phosphate of soda, compound, 438
 — pinkroot and calomel, 440
 — savine, 405, 440
 — Portland, 178
 — Prussian blue, 234
 — compound, 234
 — red oxide of mercury, 277
 — sulphuret of mercury, 279
 — rhatany, compound, 293
 — rhubarb and chalk, 395
 — magnesia, 394
 — sulphate of potassa, 395
 — compound, 395
 — sal ammoniac, 108
 — salicin, compound, 407
 — saline, compound, 307
 — Stevens's, 378
 — santonine, 130
 — sarsaparilla and Peruvian bark, 413
 — savine and ginger, 405
 — pinkroot, 405, 440
 — Spanish flies, 405
 — verdigris, 405
 — scammony and cream of tartar, 417
 — soot, 417
 — compound, 417
 — Seidlitz, 440
 — semen contra, 130
 — seneka, compound, 423
 — senna, and guaiacum, 425
 — compound, 425
 — soap, cosmetic, 410
 — spermaceti, compound, 179
 — squill and calomel, 418
 — cream of tartar, 418
 — ipecacuanha, 418
 Powder, squill and nitre, 418
 — sulphur, 418
 — compound, 418
 — starch, compound, 115
 — strychnia, compound, 446
 — subcarbonate of iron, 230
 — subnitrate of bismuth, 148
 — sulphate of copper, 215
 — compound, 215
 — iron, 240
 — mercury, compound, 278
 — potassa, and rhubarb, 381
 — compound, 381
 — quinia, 390
 — and morphia, 390
 — soda, 390
 — tartar emetic, 390
 — tartaric acid, 390
 — compound, 390
 — soda and opium, 439
 — nitre, 438
 — tartar emetic, 438
 — sulphur and antimony, 449
 — camphor, 449
 — cream of tartar, 449
 — liquorice, 449
 — magnesia, 449
 — orris root, 449
 — compound, 449
 — sulphuret of tin, 443
 — tartar emetic, 122
 — and ipecacuanha, 122
 — phosphate lime, 122
 — quinine, 122
 — compound, 122
 — tartarized soda and rhubarb, 439
 — tartrate of iron and columbo, 242
 — effervescent, 242
 — potassa and rhubarb, 382
 — tin, 442
 — tobacco, 452
 — tormentil, 459
 — tragacanth, compound, 460
 — uva ursi, compound, 462
 — valerian, compound, 463
 — verdigris and savine, 212
 — wild ipecacuanha, 226
 — willow bark, compound, 407
 — white hellebore, 467
 — zinc, oxide, 470
 — and columbo, 470
 — compound, 470
 — cyanuret, 472
 — ferro-cyanuret, 472
 Precipitation, 515
 Preservative liquid, 154
 Pressavin's vegeto-mercurial liquor, 280
 Prickly ash, 469
 — decoction, 469
 Pride of China, 140
 Process for coating pills, 485
 Proof spirit, 34
 Prophylactic vinegar, 79
 Proper use of utensils, 59
 Prunes, 384
 — pulp, 384
 Prussian blue, 234
 — ink, 234
 — mixture, 234
 — ointment, 234
 — pills, 234
 — powder, 234
 — compound, 234
 Prussic acid, 82, 555

Pudding, arrowroot, 312
 ———— tapioca, 454
 Pulp of purging cassia, 173
 ———— tamarinds, 453
 Pulp, 519
 Purging cassia, 173
 Purified animal charcoal, 168

Purple of Cassius, 139
 Putty, 485
 Pyroligneous acid, 79
 ———— cataplasm, 79
 ———— injection, 79
 ———— mouth wash, 79

Q

Quassia, 386
 ———— extract, 386
 ———— infusion, 386
 ———— compound, 386
 ———— tincture, 386
 ———— compound, 386
 ———— wine, 386
 Queen's root, 443
 ———— decoction, 444
 Quiet in sick room, 57
 Quince seed, 216
 ———— bandoline, 216
 ———— decoction, 216
 ———— mixture, 216
 Quinia, 387
 ———— amorphous, 387
 ———— impure, 387
 ———— tincture, 387
 ———— tincture, 387
 ———— acetate, 387
 ———— and chloride mercury, 389
 ———— pills, 389
 ———— arseniate, 387
 ———— arsenite, 388
 ———— di-arsenite, 388
 ———— citrate, 388
 ———— and iron, 233
 ———— syrup, 388
 ———— ferro-cyanate, 388
 ———— mixture, 388
 ———— pills, 388
 ———— hydriodate, 388
 ———— iodide, 388
 ———— with iron, 388
 ———— bin-iodide, 388
 ———— ioduretted hydriodate, 389
 ———— kinate, 389
 ———— pills, 389
 ———— lactate, 389

Quinia, lactate, mixture, 389
 ———— pills, 389
 ———— syrup, 389
 ———— muriate, 389
 ———— mixture, 389
 ———— nitrate, 390
 ———— phosphate, 390
 ———— sulphate, 390
 ———— dentifrice, 392
 ———— enema, 392
 ———— gargle, 392
 ———— liniment, 392
 ———— mixture, 392
 ———— with coffee, 392
 ———— ointment, 392
 ———— pills, 391
 ———— and gentian, 391
 ———— compound, 391
 ———— plaster, 392
 ———— powder, 390
 ———— and morphia, 390
 ———— soda, 390
 ———— tartar emetic, 390
 ———— tartaric acid, 390
 ———— compound, 390
 ———— syrup, 391
 ———— and coffee, 391
 ———— tincture, 392
 ———— compound, 392
 ———— wine, 392
 ———— aromatic, 392
 ———— sulpho-tartrate, 392
 ———— mixture, 393
 ———— tannate, 393
 ———— impure, 393
 ———— tartrate, 393
 ———— valerianate, 393
 ———— pills, 393

R

Racahout, 151
 Rat poison, 350, 480
 Ratafia of wormwood, 76
 Reaumur's thermometer, 42
 Red cedar, 292
 ———— cerate, 292
 ———— ink, 483
 ———— iodide of mercury, 275
 ———— lead, 360
 ———— oxide of iron, 239
 ———— mercury, 277
 ———— poppy, 399
 ———— infusion, 399
 ———— compound, 399
 ———— syrup, 399
 ———— precipitate, 277
 ———— roses, 400
 ———— saunders, 410
 ———— sulphuret of mercury, 279

Refrigerants, 72
 Reinsch's test, 554
 Resin, 393
 ———— cerate, 393
 ———— compound, 394
 ———— hemp, 163
 ———— jalap, 290
 ———— plaster, 394
 ———— St. Andrew's, 394
 ———— vigo with mercury, 394
 ———— powder, hæmostatic, 393
 ———— scammony, 417
 Rhatany, 293
 ———— clyster, 294
 ———— collutory, 294
 ———— dentifrice, 293
 ———— electuary, 294
 ———— extract, 293
 ———— infusion, 294

Rhatany, injection, 294
 — mixture, 294
 — ointment, compound, 294
 — pills, 294
 — powder, compound, 293
 — syrup, 294
 — tincture, 294
 Rhubarb, 394
 — electuary, 396
 — extract, 396
 — fluid, 396
 — with senna, 396
 — infusion, 396
 — alkaline, 396
 — lozenges, 396
 — mixture, 398
 — pills, 395
 — compound, 395
 — and aloes, 96
 — caraway, 395
 — chamomile, 395
 — ipecacuanha, 395
 — iron, 396
 — oxgall, 396
 — soda, 396
 — powder and chalk, 395
 — magnesia, 394
 — sulphate of potassa, 395
 — compound, 395
 — saccharated, 406
 — roasted, 395
 — suppository, 396
 — syrup, 397
 — aromatic, 397
 — and senna, 397
 — tincture, 397
 — alkaline, 398
 — sweet, 398
 — and aloes, 397
 — aniseed, 398
 — gentian, 397
 — senna, 397
 — Warner's cordial, 397
 — wine, 398
 — and gentian, 398
 — compound, 398
 Rice, 345
 — blanchmange, 491

Rice, custard, 490
 — gruel, 346
 — jelly, 346
 — mucilage, 346
 — water, 346
 Rob of mulberries, 319
 Rochelle salt, 439
 Rose, dog, 399
 — confection, 399
 — hundred-leaved, 399
 — collutory, 400
 — essence, 400
 — lozenges, 400
 — oil, 400
 — water, 399
 — ointment, 399
 — red, 400
 — confection, 400
 — electuary, 400
 — honey, 400
 — infusion, acid, 400
 — syrup, 401
 — tincture, 400
 — vinegar, compound, 401
 — wine, 401
 Rosemary, 401
 — aromatic bath, 401
 — fomentation, 401
 — vinegar, 401
 — essence, 401
 — Hungary water, 401
 — oil, 401
 — spirit, 401
 — tincture, 401
 — marsh, 443
 Rosin, 393
 Round-leaved dogwood, 207
 Rue, 403
 — confection, 403
 — extract, 403
 — mixture, 403
 — and squill, 403
 — oil, 403
 — powder, saccharated, 406
 — syrup, 403
 — tincture, 403
 Rules for the administration of medicine, 58
 Rust's astringent, 99

S

Saccharate of lead, 361
 Saccharated powders, 406
 Saccharine alum, 100
 — carbonate of iron, 230
 — and manganese, 483
 Saccharometer, 31, 34
 Safflower, 171
 — infusion, 171
 Saffron, 209
 — collyrium, 210
 — elixir, 210
 — infusion, 210
 — ointment, 210
 — pills, 210
 — powder saccharated, 406
 — syrup, 210
 — tincture, 210
 — compound, 210
 Sagapenum, 406
 — prepared, 406
 — pills, compound, 406
 Sage, 408

Sage, gargle, 408
 — infusion, 408
 — compound, 408
 — vinegar, 408
 Sago, 407
 — milk, 407
 — mucilage, 407
 — posset, 407
 Sal-ammoniac, 108
 — bolus, 108
 — cataplasm, 109
 — collutory, 109
 — draught, 109
 — emulsion, 109
 — fomentation, 108
 — liniment, 109
 — lotion, 108
 — powder, 108
 — wash, 109
 Sal prunelle, 379
 Salep, mucilage, 487
 — powder, compound, 487
 Salicin, 407

- Salicin, pills, 407
 — compound, 407
 — powder, compound, 407
 Salt, common, 431
 — of sorrel, 380
 Saltpetre, 379
 Saponine, 411
 Sarsaparilla, 413
 — beer, 416
 — decoction, 413
 — compound, 413
 — Feltz's, 413
 — Jauperand's, 414
 — Vinache's, 413
 — Zittman's, 414
 — essence, 414
 — compound, 414
 — extract, 414
 — fluid, 414
 — infusion, 413
 — alkaline, 413
 — Lisbon diet drink, 413
 — powder, with Peruvian bark, 413
 — syrup, 414
 — compound, 415
 — for mineral water, 415
 — Gesnouin's, 416
 — Laffeteur's, 415
 — Indian, 261
 — decoction, 262
 — infusion, 262
 — mixture, 262
 — syrup, 261
 Sassafras, 416
 — bark, 416
 — infusion, 416
 — compound, 416
 — oil, 416
 — tincture, 416
 — compound, 416
 — pith, 416
 — infusion, 416
 Saviard's lotion, 372
 Savine, 404
 — cerate, 405
 — extract, 405
 — infusion, 405
 — oil, 405
 — ointment, 405
 — pills, 405
 — powder with cantharides, 405
 — ginger, 405
 — pinkroot, 405
 — verdigris, 405
 — saccharated, 406
 — tincture, 405
 — compound, 405
 Scammony, 416
 — chocolate, 418
 — confection, 417
 — elixir, 418
 — emulsion, 417
 — compound, 417
 — extract, 417
 — mixture, 417
 — oil, 418
 — pastilles, 418
 — de santé, 418
 — pills, 417
 — compound, 417
 — with ox-gall, 417
 — powder with cream of tartar, 417
 — soot, 417
 — compound, 417
 — resin, 417
 — soap, 418
 Scarifications, 552
 Scudamore's mixture, 197
 Scurvy grass, 194
 — electuary, 194
 — gargle, 195
 Sea holly, 225
 Sealing wax, 485
 — red, 296
 Sedatives, 72
 Seidlitz powders, 440
 — water, 307
 Semen contra, 130
 — electuary, 130
 — mixture, 130
 — powder, 130
 — of santonine, 130
 Semi-vitrified oxide of lead, 360
 Seneka, 423
 — decoction, 423
 — emulsion, 424
 — extract with squill, 424
 — infusion, 423
 — compound, 423
 — jelly, 424
 — mixture, 424
 — pills, 423
 — polygalic acid, 424
 — syrup, 424
 — powder, compound, 423
 — syrup, 423
 Senna, 424
 — American, 174
 — infusion, 174
 — confection, 425
 — compound, 425
 — electuary with cream of tartar, 425
 — figs, 425
 — rhubarb, 425
 — sulphur, 425
 — emulsion, 427
 — extract, alcoholic, 425
 — fluid, 427
 — infusion, 426
 — compound, 426
 — with buckthorn, 427
 — coffee, 426
 — lemon juice, 427
 — rhubarb, 427
 — tamarinds, 426
 — tincture, 427
 — mixture, 427
 — pills, 425
 — powder, compound, 425
 — and guaiacum, 425
 — syrup, 427
 — and cider, 428
 — rhubarb, 428
 — tincture, compound, 426
 — with gentian, 426
 — jalap, 426
 — wine, compound, 426
 Setons, 543
 Shampoo liquid, 167
 Show-bottles, colors for, 481
 Shower bath, 537
 Sialagogues, 72
 Sikes's hydrometer, 32
 Silicate of potassa, 381
 Silk weed, 132
 Silver, 125
 — solder for, 486
 — chloride, 125
 — and ammonia, 125
 — pills, 125
 — powder, 125
 — cyanide, 125

- Silver, cyanide, ointment, 126
 — iodide, 126
 — ointment, 126
 — nitrate, 126
 collyrium, 127
 hair dye, 127
 indelible ink, 127
 mixture, 127
 ointment, 127
 compound, 127
 pills, 126
 solution, 126
 Mackensie's, 127
 oxide, 126
 ointment, 126
 pills, 126
 Simaruba, 429
 — infusion, 429
 compound, 429
 Sinapisms, 541
 Sinapism of ammonia, 103
 Skunk cabbage, 220
 — infusion, 220
 — tincture, 220
 Slippery elm, 462
 — cataplasm, 462
 — infusion, 462
 Snakeroot, black, 185
 — decoction, 185
 — extract, fluid, 479
 — tincture, 185
 — Virginia, 428
 electuary, 428
 extract, fluid, 428
 infusion, 428
 compound, 428
 mixture, with allspice, 429
 acetated, 429
 ether, 429
 pills, compound, 428
 tincture, 428
 with balsam Peru, 429
 wine, with vanilla, 429
 Sneezewort, 260
 Soap, 410
 — almond, 410
 — aromatic, 410
 — arsenical, 411
 — balsam, camphorated acetic, 412
 — beef marrow, 410
 — bolus, 411
 — camphorated, 410
 liniment, 411
 — castile, 410
 — cataplasm, 412
 — cerate, 411
 — clyster, 412
 — cod-liver oil, 412
 ioduretted, 412
 — common, 410
 — croton oil, 334
 — electuary, 412
 — essence, 412
 camphorated, 412
 — grease balls, 411
 — hard, 410
 — iodide of potassium, 366
 — itch ointment, 412
 — jalap, 290
 — liniment, 411
 — pills, 411
 and ox-gall, 411
 compound, 411
 — plaster, 412
 — powder, cosmetic, 410
 41
 Soap, saponine, 411
 — scammony, 418
 — soft, 410
 — solution, ethereal, 412
 — Spanish, 410
 — Starkey's, 333, 410
 — sulphuret of potassium, 371
 — suppository, 412
 — tincture, camphorated, 411
 — turpentine, 410
 — with sulphur, 412
 Soda, 432, 558
 — powders, 436
 — solution, 432
 effervescing, 436
 — acetate, 433
 pills, compound, 433
 — arseniate, 433
 solution, 433
 — bicarbonate, 436
 emulsion, 436
 lozenges, 436
 mixture, 436
 and copaiba, 436
 powder, 436
 and magnesia, 439
 — bisulphate, 439
 and magnesia, 439
 — borate, 433
 — cakes, 491
 — carbonate, 434
 dried, 434
 injection, 435
 lotion, 435
 mixture with chamomile, 434
 gentian, 435
 ipecacuanha, 434
 quassia, 435
 ointment, 435
 pills, 435
 and ipecacuanha, 435 wP
 rhubarb, 435
 powder, with mercury, 435
 rhubarb, 435
 solution, diuretic, 434
 — caustic, 432
 — chlorate, 436
 gargle, 437
 lotion, 437
 — chlorinated, 437
 cataplasm, 437
 gargle, 437
 injection, 437
 solution, 437
 diluted, 437
 — hyposulphite, 437
 bath, 438
 syrup, 438
 — nitrate, 438
 — phosphate, 438
 mixture, 438
 powder, compound, 438
 — sulphate, 438
 clyster, 439
 electuary, 439
 emulsion, 439
 lemonade, 439
 lotion, 439
 pills, compound, 439
 powder, with nitre, 438
 opium, 439
 tartar emetic, 438
 solution, compound, 439
 suppository, 439
 — tartarized, 439
 clyster, 440

- Soda, tartarized, mixture, 440
 ————— powder, with rhubarb, 439
 ————— Seidlitz, 440
 ————— whey, 440
 ————— valerianate, 440
 Sodium, 431
 ————— bromide, 431
 ————— ointment, 431
 ————— chloride, 431
 ————— bath, with gelatine, 431
 ————— clyster, 431
 ————— and arnica, 431
 ————— fomentation, 431
 ————— mixture, with lemon juice, 432
 ————— ointment, 432
 ————— compound, 432
 ————— powder, compound, 431
 ————— and cochineal, 431
 ————— chloroplatinate, 356
 ————— injection, 356
 ————— sulphuret, 432
 ————— bath, 432
 ————— liniment, 432
 ————— lotion, 432
 ————— mixture with sal-ammoniac, 432
 ————— ointment, 432
 Solders, 485
 Solder for brass, copper, and iron, 485
 ————— gold, 486
 ————— lead, 485
 ————— pewter, 485
 ————— silver, 486
 ————— tin, 485
 ————— zinc, 485
 Soluble glass, 381
 ————— tartar, 382
 Solution, arsenical, 80, 373
 ————— Donovan's, 129
 ————— Fowler's, 373
 ————— Heinecke's, 433
 ————— Labarraque's, 437
 ————— Mackenzie's, 127
 ————— Magendie's, 323
 ————— Pearson's, 433
 ————— of acetate ammonia, 104
 ————— baryta, 143
 ————— mercury, 268
 ————— morphia, 321
 ————— alcoholic, 321
 ————— strychnia, 446
 ————— alum, 101
 ————— odontalgic, 100
 ————— ammonia, 103
 ————— citrate, 107
 ————— arseniate of ammonia, 105
 ————— soda, 433
 ————— arsenic, 80
 ————— arsenite of potassa, 373
 ————— ammoniated copper, 214
 ————— muriate of copper and mercury, 214
 ————— ammonio-tartrate of iron, 243
 ————— atropine, 135
 ————— barytes, 143
 ————— belladonna, extract, 145
 ————— bicarbonate of potassa, effervescing, 377
 ————— bimeconate of morphia, 321
 ————— bitartrate of potassa, 383
 ————— compound, 383
 ————— borotartrate of potassa, 375
 ————— bromide of mercury, ethereal, 269
 ————— potassium, 362
 ————— bromine, 149
 ————— alcoholic, 149
 Solution of brucia, 150
 ————— carbonate of potassa, 375
 ————— soda, 434
 ————— diuretic, 434
 ————— effervescing, 436
 ————— caustic potassa, 372
 ————— chlorate of potassa, 378
 ————— chloride of arsenic, 129
 ————— barium, 142
 ————— calcium, 154
 ————— gold and sodium, 138
 ————— chlorinated soda, 437
 ————— diluted, 437
 ————— potassa, 363
 ————— citrate of ammonia, 107
 ————— magnesia, 307
 ————— potassa, 378
 ————— conia, 203
 ————— corrosive sublimate, 269
 ————— alcoholic, 269
 ————— creasote, alcoholic, 208
 ————— croton oil, saponaceous, 334
 ————— cyanide of gold, 139
 ————— cyanuret of mercury, 274
 ————— potassium, 363
 ————— delphinia, 217
 ————— ferrocyanuret of zinc, 472
 ————— gamboge, alkaline, 249
 ————— hydrargyro-iodide of potassium, 363
 ————— iodide of iron, 235
 ————— official, 236
 ————— potassium, 364
 ————— caustic, 367
 ————— compound, 367
 ————— zinc, 473
 ————— iodine, 284
 ————— lac, 296
 ————— magnesia, 306
 ————— malate of iron, 237
 ————— morphia and ipecacuanha, 320
 ————— muriate of morphia, 322
 ————— myrrh, alkaline, 326
 ————— nitrate of silver, 126
 ————— nitro-saccharate of lead, 361
 ————— potassa, 372
 ————— lithontriptic, 372
 ————— persesquinitrate of iron, 238
 ————— phosphate of ammonia, 109
 ————— silicate of potassa, 381
 ————— soap, ethereal, 412
 ————— soda, 432, 436
 ————— subacetate of lead, 356
 ————— sulphate of bebeerine, 144
 ————— cadmium, 152
 ————— iron, 241
 ————— morphia, 323
 ————— soda, compound, 439
 ————— veratria, 467
 ————— zinc, 474
 ————— sulphuret of potassium, 370
 ————— tartar emetic, 122
 ————— and squill, 122
 ————— tartrate of iron and potassa, 242
 ————— potassa and ammonia, 384
 ————— terechloride of antimony, 118
 ————— tersulphate of iron, 238
 Soot, 245
 ————— cataplasm, 246
 ————— decoction, 245
 ————— injection, 246
 ————— mixture, 246
 ————— mouth wash, 246
 ————— oil, 246
 ————— ointment, 246
 ————— compound, 246
 ————— pills, 245

- Soot, tincture, 245
 Southernwood, 130
 — clyster, 130
 — infusion, 130
 Spanish flies (see Cantharides)
 Spearmint, 316
 — arquebusade water, 317
 — essence, 317
 — infusion, 317
 — compound, 317
 — spirit, 317
 — water, 317
 Specific gravity, 28
 — beads, 32
 — table, 41
 Spermaceti, 179
 — cerate, 179
 — liniment, 180
 — lip salve, 179
 — mixture, 179
 — ointment, 179
 — and rosewater, 179
 — sultana, 180
 — powder, compound, 179
 Spice plaster, 194, 477
 Spirits, 534
 Spirit of acetic ether, 91
 — allspice, 351
 — ammonia, 104
 — aromatic, 104
 — and succinic acid, 110
 — fetid, 104
 — aniseed, 118
 — balm, compound, 315
 — caraway, 171
 — castor, compound, 175
 — cinnamon, 193
 — cloves, 172
 — horseradish, compound, 128
 — juniper, compound, 292
 — lavender, 299
 — compound, 299
 — lemon, 301
 — odoriferous, 301
 — Mindererus, 105
 — muriatic ether, 92
 — nutmeg, 325
 — compound, 325
 — orange peel, 135
 — peppermint, 316
 — rosemary, 401
 — spearmint, 317
 — succinated ammonia, 110
 — sulphuric ether, 92
 — sweet of nitre, 91
 — turpentine, 332
 — vanilla, 465
 — vinegar, aromatic, 78
 — wormwood, compound, 76
 Spitta's lozenges, 211
 Splints, felt, 486
 Sponge, 442
 — burnt, 442
 — bolus, 442
 — electuary, 442
 — powder, 442
 — compound, 442
 Sponging, 540
 Spongio-piline, 541
 Spruce beer, 490
 Spurge, 226
 Squill, 418
 — electuary, 422
 — emulsion, 421
 — extract, 422
 — acetic, 422
 — Squill, extract, and rhubarb, 421
 — ointment, 422
 — oxymel, 422
 — mixture, with ammoniac, 421
 — ipecacuanha, 421
 — marsh mallow, 421
 — valerian, 421
 — pills and ammoniac, 419
 — calomel, 419
 — croton oil, 419
 — henbane, 419
 — ipecacuanha, 419
 — compound, 419
 — powder with calomel, 418
 — cream of tartar, 418
 — ipecacuanha, 418
 — nitre, 418
 — sulphur, 418
 — compound, 418
 — saccharated, 406
 — syrup, 422
 — aromatic, 422
 — compound, 419
 — mixture, 421
 — tincture, 420
 — alkaline, 420
 — and elaterium, 420
 — benzoin, compound, 420
 — ethereal, 421
 — vinegar, 422
 — mixture, 421
 — and ammonia, 421
 — wine, 420
 — compound, 420
 — bitter, 420
 Star grass, 93
 — tincture, 93
 Starch, 115
 — iodide, 116
 — jelly, 116
 — lozenges, 116
 — mixture, with suet, 116
 — mucilage, 115
 — powder, compound, 115
 Starkey's soap, 333, 410
 Stavesacre, 443
 — decoction, 443
 — ointment, 443
 — vinegar, 443
 Sternutatory, of euphorbium, 227
 Stevens's saline powder, 378
 Stimulants, 73
 Storax, 445
 — pills, compound, 445
 — purified, 445
 — syrup, 445
 Stramonium, 444
 — extract of leaves, 444
 — seeds, 444
 — lotion, 444
 — mixture, 444
 — oil, 445
 — ointment, 445
 — compound, 445
 — pills, seed, 444
 — compound, 444
 — powder, saccharated, 406
 — syrup, 444
 — tincture of leaves, 444
 — seed, 444
 — ethereal, 445
 — wine, 445
 Strychnia, 445, 565
 — acetate, 446
 — solution, 446
 — tincture, 447

- Strychnia, collyrium, 446
 iodate, 447
 liniment, 446
 mixture, 446
 muriate, 447
 nitrate, 447
 ointment, 447
 ointment, 446
 pills, 446
 stimulant, 446
 powder, compound, 446
 sulphate, 447
 syrup, 447
 tincture, 446
 Styptic of sulphate of copper, 216
 Stuping, 541
 Subacetate of copper, 212
 lead, 356
 Subcarbonate of iron, 230
 Subnitrate of bismuth, 148
 Succinate of ammonia, 109
 Succory, 184
 extract, 185
 infusion, 185
 syrup, compound, 185
 Sugar, 406
 barley, 263
 burnt, 406
 milk, 295
 syrup, 406
 vanilla, 465
 vermifuge, 406
 Sulphate of alumina, 99
 ammonia, 110
 antimony, 121
 atropia, 135
 brucia, 150
 cadmium, 152
 cinchonia, 192
 copper, 215
 indigo, 283
 iron, 240
 magnesia, 307
 manganese, 311
 morphia, 322
 potassa, 381
 and ammonia, 382
 magnesia, 382
 sulphur, 382
 quinia, 390
 soda, 438
 strychnia, 447
 veratria, 467
 zinc, 474
 Sulpho-cyanuret of potassium, 369
 tartrate of quinia, 393
 Sulphur, 448
 balsam, 450
 etheral, 450
 terebinthinated, 450
 cerate, 451
 electuary, 449, 450
 compound, 450
 iodide, 452
 ointment, 452
 powder, 452
 linetus, 450
 liniment, with soap, 451
 liver, 369
 mixture, 450
 ointment, 450
 compound, 450
 with camphor, 451
 soap, 451
 zinc, 451
 powder, compound, 449
 Sulphur, powder, with antimony, 449
 camphor, 449
 cream of tartar, 449
 liquorice, 449
 magnesia, 449
 orris root, 449
 precipitated, 449
 sublimed, 448
 washed, 448
 Sulphuret of antimony, 120
 arsenic, 130
 barium, 143
 calcium, 155
 carbon, 451
 iron, 241
 magnesium, 308
 mercury, black, 279
 red, 279
 potassium, 369
 sodium, 432
 tin, 443
 Sulphuretted anthracokali, 170
 hydrogen, 83
 water, artificial, 83
 hydrosulphate of ammonia, 108
 Sulphuric acid, 86, 556
 aromatic, 87
 and alcohol, 87
 nitric ether, 87
 diluted, 86
 lemonade, 87
 liniment, 87
 ointment, 87
 ether, 92
 ethereal oil, 92
 Hoffmann's anodyne, 92
 lotion, 92
 mixture, with camphor, 93
 turpentine, 92
 rectified, 92
 spirit, 92
 syrup, 92
 Suppositories, 66
 Suppository, anthelmintic, 97
 charcoal, 169
 hemlock, 203
 opium, 340
 rhubarb, 396
 soap, 412
 sulphate of soda, 439
 chloride of zinc, 471
 Swamp dogwood, 207
 silkweed, 131
 Swediaur's tincture, 343
 Sweet almonds, 112
 fern, 201
 spirit of nitre, 91
 violet, 468
 Sydenham's laudanum, 342
 Sylvius, salt of, 363
 Syrups, 519
 Syrup, 406
 acetate of morphia, 321
 antimonial wine, 124
 asparagus shoots, 132
 assafetida, 133
 atropine, 135
 balsam of Peru, 141
 tolu, 458
 belladonna, 146
 Gillet's, 146
 biniodide of potassium, 367
 bittersweet, 221
 blackberry, 403
 bloodroot, 409
 brooklime, 144

- Syrup, buckthorn, 394
 — capillaire, 90
 — cayenne pepper, 167
 — chamomile, 119
 — chloride of gold and sodium, 138
 — compound, 138
 — iron, 232
 — cinchonia, 192
 — citrate of caffein, 196
 — iron, 233
 — and magnesia, 234
 — quinia, 388
 — citric acid, 81, 82
 — clove pink, 217
 — cod-liver oil, 330
 — codeia, 195
 — coffee, 195
 — colchicum, 197
 — coltsfoot, 461
 — cubebs, alcoholic ethereal extract, 212
 — cyanuret of potassium, 363
 — emetine, 223
 — ergot, 224
 — foxglove, 219
 — galls, 248
 — garlic, 93
 — gentian, 251
 — extract, 251
 — gentianin, 252
 — ginger, 476
 — ethereal extract, 476
 — gold, 137
 — gum, 77
 — compound, 77
 — hive, 420
 — horehound, 313
 — hydrocyanic acid, 83
 — hydrosulphate of ammonia, 108
 — hyposulphite of soda, 438
 — hyposulphited sulphuret of potassium 370
 — Indian sarsaparilla, 261
 — iodate of potassa, 379
 — iodide and chloride of iron, 236
 — of iron, 236
 — manganese, 310
 — and iron, 310
 — potassium, 365
 — zinc, 473
 — ipecacuanha, 288, 483
 — compound, 288
 — Jackson's pectoral, 484
 — kino, 293
 — lactate of iron, 237
 — quinia, 389
 — lactucarium, 297
 — lemon, 301
 — peel, 301
 — vinous, 301
 — lettuce, 298
 — liquorice, 255
 — lobelia, 304
 — lupulin, 304
 — maidenhair, 90
 — compound, 91
 — manna, 311
 — marsh mallow, 98
 Syrup, matico, 314
 — milk, 295
 — monesia, 319
 — compound, 319
 — mulberries, 319
 — muriate of morphia, 322
 — compound, 322
 — naphthaline, 327
 — opium, 340
 — succinated, 340
 — orange flowers, 136
 — peel, 136
 — orgeat, 113
 — paullinia, 348
 — pectoral, 347
 — Peruvian bark, 191
 — compound, 191
 — phosphate of iron, 484
 — lime, 158
 — pipsissewa, 182
 — polygalic acid, 424
 — poppy heads, 347
 — substitute for, 347
 — proto-iodide of iron, 236
 — proto-nitrate of iron, 238, 484
 — red roses, 401
 — poppy, 399
 — rhatany, 294
 — rhubarb, 397
 — and senna, 397
 — aromatic, 397
 — rue, 403
 — saffron, 210
 — sarsaparilla, 414, 415
 — for mineral water, 415
 — compound, 415
 — Gesnouin's, 416
 — Laffecteur's, 415
 — seneka, 423
 — senna, 427
 — and cider, 428
 — rhubarb, 428
 — squill, 422
 — aromatic, 422
 — compound, 419
 — storax, 445
 — stramonium, 444
 — succory, compound, 185
 — sulphate of iron, 241
 — morphia, 322
 — quinia, 391
 — and coffee, 391
 — strychnia, 447
 — sulphuret of magnesium, 308
 — potassium, 370
 — compound, 370
 — sulphuretted, 108
 — sulphuric ether, 92
 — tartaric acid, 88
 — thridace, 297
 — uva ursi, 462
 — valerian, 464
 — vinegar, 78
 — violets, 468
 — wild-cherry bark, 384
 — Willis's, 370
 — wormwood, 75

T

Table of avoirdupois weights, 17, 18

- distilled oils, 530
 — drops and minims, 26
 — foreign weights, 22

Table of hydrometrical equivalents, 36

- incompatibles, 492
 — pharmaceutical names, 503
 — poisons and antidotes, 568

- Table, posological, 496
 — of sp. grs. and Baumé's hydrometer, 35
 — of preparations, 41
 Taffeta cantharidalis, 166
 Tamarinds, 453
 — electuary, 454
 — infusion, 454
 — pulp, 453
 — whey, 295, 454
 Tannate of iron, 243
 — lead, 361
 — quinia, 393
 Tannic acid, 87
 — gargle, 87
 — injection, 87
 — mixture, 88
 — ointment, 88
 — pills, 87
 — and opium, 87
 Tansy, 454
 — extract, 454
 — infusion, 454
 — oil, 454
 — pills, 454
 — tincture, compound, 454
 Tapioca, 454
 — jelly, 454
 — pudding, 454
 Tar, 354
 — beer, 354
 — ointment, 355
 — compound, 355
 — pills, 354
 — water, 354
 — wine, 354
 Tartaric acid, 88, 557
 — effervescing powders, 88
 — lemonade, 88
 — lozenges, 88
 — syrup, 88
 Tartar emetic, 122, 558
 — bolus, 122
 — clyster, 123
 — emulsion, 122
 — liniment, with ammonia, 123
 — lotion, 123
 — and camphor, 123
 — corrosive sublimate, 123
 — ointment, 123
 — compound, 123
 — pills and camphor, 122
 — guaiacum, 122
 — opium, 122
 — plaster, 123
 — powder, 122
 — and ipecacuanha, 122
 — phosphate of lime, 122
 — quinine, 122
 — compound, 122
 — solution, 122
 — and squill, 122
 — wine, 123
 — collyrium, 123
 — mixture, with ammoniac, 124
 — bitter sweet, 124
 — laudanum, 124
 — syrup, 124
 Tartarized soda, 439
 Tartrate of antimony and potassa, 122
 — iron and potassa, 241
 — magnesia, 308
 — mercury and potassa, 280
 Tartrate of morphia, 323
 — potassa, 382
 — and ammonia, 383
 — quinia, 393
 — soda and potassa, 439
 Tea, balm, 315
 — beef, 488
 — bran, 489
 — mutton, 488
 — veal, 488
 — worm, 441
 Temperature of sick room, 56
 Temperatures in pharmaceutical operations, 36
 Tepid and temperate baths, 536
 Terebinthinated ether, 93
 Thermometrical scales, 42
 Thieves' vinegar, 79
 Thorn apple, 444
 Thridace, 297
 Thus, 353
 Tin, 442, 565
 — solder for, 485
 — chloride, 442
 — electuary, 442
 — ointment, 442
 — oxide, 443
 — powder, 442
 — sulphuret, 443
 — powder, compound, 443
 Tinctures, 521
 Tincture, acetate of iron, 228
 — alcoholic, 228
 — ethereal, 228
 — morphia, 321
 — potassa, 373
 — strychnia, 447
 — zinc, 473
 — aconite leaves, 89
 — root, 89
 — Fleming's, 477
 — aloes, 97
 — ethereal, 98
 — and myrrh, 98
 — allspice, 351
 — amber, 448
 — alkaline, 448
 — ethereal, 448
 — American centaury, 404
 — columbo, 245
 — hellebore, 468
 — ammonia, compound, 106
 — ammoniated iron, 229
 — angelica, 117
 — compound, 117
 — angustura, 117
 — aniseed, 118
 — antacid, 270
 — arnica, 129
 — assafetida, 133
 — alkaline, 134
 — ammoniated, 134
 — and castor, 134
 — soot, 134
 — atropine, 135
 — balsam of Peru, 141
 — tolu, 458
 — compound, 458
 — with foxglove, 458
 — belladonna, 146
 — Blackett's, 479
 — ethereal, 146
 — benzoin, 147
 — compound, 147
 — Bestucheffe's, 233
 — black hellebore, 261
 — snakeroot, 185

- Tincture, bloodroot, 409
 — compound, 409
 — brucia, 150
 — buchu, 151
 — calamus, 153
 — compound, 153
 — camphor, 160
 — and saffron, 160
 — ethereal, 160
 — cantharides, 164
 — and guaiacum, 164
 — camphorated, 164
 — ethereal, 164
 — compound, 164
 — ethereal, 164
 — carbonate of potassa, 376
 — compound, 377
 — cardamom, 170
 — compound, 170
 — cascarilla, 173
 — castor, 174
 — ammoniated, 175
 — compound, 175
 — ethereal, 174
 — catechu, 176
 — Cayenne pepper, 167
 — and cantharides, 168
 — cevadilla, 404
 — chiretta, 182
 — chloride of gold, 137
 — iron, 232
 — ethereal, 233
 — zinc, ethereal, 471
 — cinchonia, 192
 — cinnamon, 193
 — compound, 193
 — ethereal, 193
 — citrate of iron, 233
 — cloves, 172
 — cochineal, 194
 — colchicum flowers, 197
 — seed, 196
 — compound, 197
 — ethereal, 196
 — and digitalis, 197
 — colocynth, 200
 — columbo, 201
 — contrayerva, 203
 — compound, 203
 — copaiba, 206
 — alkaline, 206
 — compound, 206
 — coriander, 207
 — croton oil, 334
 — cubebs, 211
 — cyanuret mercury, compound, 274
 — Dippel's oil, 329
 — elaterin, 222
 — ergot, 224
 — foxglove, 219
 — ethereal, 219
 — galbanum, 247
 — compound, 247
 — galls, 248
 — gamboge, alkaline, 250
 — ammoniacal, 250
 — gentian, 251
 — acidulated, 251
 — alkaline, 252
 — ammoniacal, 252
 — compound, 251
 — gentianin, 252
 — ginger, 476
 — ethereal extract, 476
 — goldthread, 206
 — guaiacum, 258
 — Tincture, guaiacum, ammoniated, 259
 — Dewees's, 258
 — hemlock, 202
 — ethereal, 202
 — hemp, 163
 — henbane, 281
 — ethereal, 281
 — hiera piera, 162
 — hops, 264
 — alkaline, 264
 — horse balm, 198
 — Huxham's, 189
 — hydrargyro-iodide of potassium, 368
 — iodide of iron, 236
 — iodine, 284
 — compound, 284, 367
 — ethereal, 284
 — saturated, 284
 — ipecacuanha, 288
 — jalap, 290
 — compound, 290
 — Jamaica dogwood, 353
 — kino, 293
 — lactucarium, 297
 — larkspur seeds, 217
 — lavender, ethereal, 300
 — lemon, 301
 — lettuce, aromatic, 297
 — lobelia, 303
 — ethereal, 303
 — lupulin, 304
 — mace, 305
 — magnolia, 309
 — malate of iron, 237
 — mastich, ethereal, 313
 — matico, 314
 — monesia, 319
 — musk, 324
 — artificial, 448
 — mustard, 431
 — myrrh, 326
 — and hellebore, 326
 — nux vomica, 328
 — compound, 328
 — opium, 342
 — acetated, 342
 — ammoniated, 343
 — and soap, 343
 — Bateman's, 343
 — camphorated, 343
 — compound, 343
 — Smith's, 343
 — succinated, 343
 — Swediaur's, 343
 — Warner's, 343
 — opoponax, 345
 — compound, 345
 — orange peel, 135
 — pareira brava, 348
 — pellitory, 385
 — compound, 385
 — Peruvian bark, 189
 — ammoniated, 190
 — and cantharides, 190
 — gentian, 190
 — snakeroot, 190
 — valerian, 190
 — compound, 189
 — phosphoric, of myrrh, 86
 — piperine, 353
 — poison oak, 460
 — poke, 351
 — proto-iodide of iron, 236
 — quassia, 386
 — compound, 386
 — quinia, 387

- Tincture, quinia, impure, 387
 — red iodide mercury, 276
 — — — — — ethereal, 276
 — rhatany, 294
 — rhubarb, 397
 — — — — — and aloes, 397
 — — — — — aniseed, 398
 — — — — — gentian, 397
 — — — — — senna, 397
 — — — — — alkaline, 398
 — — — — — sweet, 398
 — rosemary, 401
 — roses, 400
 — rue, 403
 — saffron, 210
 — — — — — compound, 210
 — sassafras, 416
 — — — — — compound, 416
 — savine, 405
 — — — — — compound, 405
 — senna and jalap, 426
 — — — — — gentian, 426
 — — — — — compound, 426
 — skunk cabbage root, 220
 — — — — — seed, 220
 — snakeroot, black, 185
 — — — — — Virginia, 428
 — — — — — and balsam Peru, 429
 — soap, camphorated, 411
 — soot, 245
 — squill, 420
 — — — — — alkaline, 420
 — — — — — and benzoïn, 420
 — — — — — elaterium, 420
 — — — — — ethereal, 421
 — stargrass, 93
 — stramonium, ethereal, 445
 — — — — — leaves, 444
 — — — — — seed, 444
 — strychnia, 446
 — sulphate of quinia, 392
 — — — — — compound, 392
 — sulphuret of potassium, 370
 — tansy, compound, 454
 — tartrate of iron, compound, 242
 — tobacco, 453
 — tulip-tree bark, 303
 — valerian, 463
 — — — — — ammoniated, 464
 — — — — — — — — — compound, 464
 — — — — — compound, 464
 — — — — — ethereal, 463
 — — — — — and Hoffmann's anodyne, 464
 — vanilla, 465
 — veratria, 466
 — white hellebore, 467
 — Winter's bark, 469
 — wormwood, 75
 — — — — — alkaline, 75
 Toast water, 488
 Tobacco, 452
 — — — — — cataplasm, 453
 — — — — — extract, 452
 Tobacco, infusion, 453
 — — — — — lotion, 453
 — — — — — mixture, 453
 — — — — — nicotina, 453
 — — — — — oil, 453
 — — — — — ointment, 453
 — — — — — pills, 452
 — — — — — powder, compound, 452
 — — — — — tincture, 453
 — — — — — wine, 452
 Tolu, balsam of, 458
 — — — — — emulsion, 458
 — — — — — inhalation, 459
 — — — — — lozenges, 458
 — — — — — mixture and belladonna, 459
 — — — — — — — — — almond emulsion, 459
 — — — — — — — — — copaiba, 459
 — — — — — — — — — morphia, 459
 — — — — — — — — — opium, 459
 — — — — — powder, saccharated, 406
 — — — — — syrup, 458
 — — — — — tincture, 458
 — — — — — — — — — compound, 458
 — — — — — — — — — and foxglove, 458
 Tonics, 73
 Tonquin powder, 323
 Tormentil, 459
 — — — — — decoction, 459
 — — — — — extract, 459
 — — — — — gargle, 459
 — — — — — powder, compound, 459
 Tous les mois, 487
 Tracing paper, 486
 Tragacanth, 460
 — — — — — mucilage, 460
 — — — — — paste, 460
 — — — — — powder, compound, 460
 Troches, 534
 — — — — — of magnesia, 305
 Tronchin's lozenges, 77, 122
 Troy weight, 17
 Tulip-tree bark, 303
 — — — — — infusion, 303
 — — — — — tincture, 303
 Turlington's balsam, 147
 Turner's cerate, 153
 Turpentine, 456
 — — — — — clyster, 456
 — — — — — electuary, 456
 — — — — — liniment, 457
 — — — — — mixture, 457
 — — — — — oil, 332
 — — — — — ointment, 457
 — — — — — — — — — compound, 457
 — — — — — pills, 456
 — — — — — — — — — with guaiacum, 456
 — — — — — — — — — jalap, 456
 — — — — — — — — — magnesia, 456
 — — — — — — — — — myrrh, 456
 — — — — — — — — — rhubarb, 456
 — — — — — plaster, 457
 — — — — — — — — — compound, 457
 Tutty ointment, 470
 Twaddle's hydrometer, 32

U

- Unguentum populeum, 282
 Urinometer, 35
 Unparalleled water, 302
 Uva ursi, 462
 — — — — — decoction, 462
 Uva ursi, extract, 462
 — — — — — mixture, 462
 — — — — — powder, compound, 462
 — — — — — syrup, 462

V

Valerian, 463
 — bolus, with iron, 463
 — sulphate of potassa, 463
 — electuary, 463
 — extract, 464
 — fluid, 464
 — infusion, 463
 — compound, 463
 — mixture, with ammonia, 464
 — Hoffmann's anodyne, 464
 — oil, 464
 — with ammonia, 464
 — pills, compound, 463
 — powder, compound, 463
 — syrup, 464
 — tincture, 463
 — ammoniated, 464
 — compound, 464
 — compound, 464
 — ethereal, 463
 — with Hoffmann's anodyne, 464
 — wine, 464
 Valerianate of iron, 243
 — quinia, 393
 — soda, 440
 — zinc, 475
 Valerianic acid, 88, 465
 Vallet's mass, 230
 Vanilla, 465
 — arrowroot, 465
 — essence, 466
 — extract, fluid, 466
 — lozenges, 465
 — milk, 466
 — powder, saccharated, 406
 — spirit, 465
 — sugar, 465
 — tincture, 465
 Vapor bath, 538
 Varnishes, 486
 Varnish, amber, 486
 — copal, 486
 — crystal, 313
 — furniture, 486
 — Japan, 486
 — lac, 486
 — leather, 486
 — metal, 486
 — picture, 313
 Veal tea, 488
 Vegetable broth, 487
 Venesection, 547
 Ventilation of sick room, 55
 Veratria, 466

Veratria, liniment, 466
 — lotion, 466
 — muriate, 467
 — nitrate, 467
 — ointment, 466
 — with iodine, 467
 — opium, 467
 — pills, 466
 — with henbane, 466
 — sulphate, 467
 — solution, 467
 — tincture, 466
 Verdigris, 212
 Vermifuge sugar, 406
 Vesicatories, 69
 Vinegar, 77
 — cataplasm, 78
 — distilled, 78
 — gargle, 78
 — mixture, 77
 — and cardamom, 77
 — pastilles, 87
 — syrup, 78
 — whey, 295
 — aromatic, 78, 401
 — spirit, 78
 — borax, 434
 — camphorated, 79
 — cantharides, 164
 — Cayenne pepper, 168
 — cinnamon, compound, 193
 — coffee, 195
 — colchicum root, 197
 — seed, 197
 — elder flowers, 408
 — four thieves, 79
 — foxglove, 219
 — lobelia, 304
 — opium, 341
 — prophylactic, 79
 — rosemary, aromatic, 401
 — roses, compound, 401
 — sage, 408
 — squill, 422
 — stavesacre, 443
 Vinegars, 524
 Violet, 468
 — conserve, 468
 — honey, 468
 — oil, 468
 — syrup, 468
 Virginia snakeroot, 428
 Vocabulary of words used in prescriptions, 44
 Volatile oils, 528

W

Wakaka, 152
 Walnut, white, 291
 Warm bath, 536, 538
 Ward's essence, 161
 — paste, 352
 Warner's cordial, 397
 — tincture, 343
 Wash, black, 277
 — cosmetic, 147, 271
 — detergent, 99
 — ophthalmic, 213

Wash, red iodide of mercury, 276
 — sal-ammoniac, 109
 — yellow, 270
 Water, allspice, 351
 — ammonia, 103
 — anemone, 116
 — angelica, 117
 — aniseed, 118
 — anti-hysteria, 316
 — apple, 489
 — arquebusade, 317

- Water, arrowroot, 312
 — avens, 253
 — balm, 315
 — barley, with nitrate potassa, 263
 — bitter almonds, 114
 — brooklime, 144
 — camphor, 159
 — and laudanum, 159
 — nitric acid, 160
 — Hoffmann's anodyne, 160
 — caraway, 171
 — chalybeate, artificial, 231
 — cherry laurel, 298
 — chicken, 488
 — chlorine, 183
 — cinnamon, 193
 — compound, 193
 — citrate of magnesia, 307
 — clove, 172
 — Cologne, 301
 — coriander, compound, 207
 — distilled, 525
 — diuretic, 114
 — elder flower, 408
 — fennel, 245
 — Goulard's vegeto-mineral, 357
 — honey, 315
 — Hungary, 401
 — ioduretted, 366
 — lavender, 299
 — lemon peel, 300, 489
 — lettuce, 298
 — lime, 155
 — and carbonate of potassa, 156
 — milk, 156
 — compound, 155
 — medicated, 484, 525
 — opium, 341
 — orange flower, 136
 — oriental, 376
 — partridge berry, 250
 — peppermint, 316
 — rice, 346
 — rose, 399
 — Seidlitz, 307
 — souchy, 489
 — spearmint, 317
 — sulphuret of potassium, 370
 — sulphuretted, artificial, 83
 — tar, 354
 — toast, 488
 — unparalleled, 302
 Wax, 178
 — sealing, 485
 — cloth, 178
 — cerate, 178
 — plaster, 178
 — rose lip-salve, 178
 Wedell's elixir, 421
 Weights and measures, 17
 — apothecaries', 20
 — avoirdupois, 17
 — foreign, 20
 — French, 20
 — metrical, 21
 — troy, 17
 Wendt's mixture, 77
 Whey, 294
 — alum, 102, 295
 — aromatic, 102, 295
 — cream of tartar, 295
 — mustard, 295, 430
 — tamarind, 295, 454
 — tartarized soda, 446
 — vinegar, 295
 — wine, 295
 White avens, 253
 — hellebore, 467
 — decoction, 467
 — ointment, 467
 — compound, 467
 — powder, 467
 — tincture, 467
 — wine, 467
 — lead, 358
 — precipitate, 268
 — ointment, 268
 — walnut, 257
 Wild cherry bark, 384
 — infusion, 384
 — syrup, 384
 — ginger, 131
 — infusion, 131
 — indigo, 142
 — ipecacuanha, 226
 — powder, compound, 226
 — lettuce, 296
 — potato, 204
 Wilkinson's liniment, 108
 Willis's syrup, 370
 Willow bark, 407
 — decoction, 408
 — dentifrice, 408
 — extract, 407
 — ointment, 408
 — powder, compound, 407
 Wine of aconite, compound, 89
 — aloes, 97
 — alkaline, 97
 — balsamic, 97
 — American hellebore, 468
 — black hellebore, 261
 — blessed thistle, 177
 — bryony, 150
 — cascarilla, compound, 173
 — centaury, compound, 178
 — cinchonia, 192
 — cinnamon, compound, 193
 — citrate of iron, 233
 — aromatic, 233
 — cloves, 172
 — colchicum root, 196
 — seed, 196
 — compound, 197
 — dogwood, round-leaved, 208
 — ergot, 224
 — gentian, 251
 — hedge hyssop, 257
 — iodide of iron, 236
 — ipecacuanha, 288
 — alkaline, 288
 — and tartar emetic, 288
 — compound, 288
 — iron, 231
 — opium, 342
 — Sydenham's, 342
 — Peruvian bark, 191
 — and calamus, 191
 — iron, 191
 — persimmons, 220
 — quassia, 386
 — quinia, 392
 — aromatic, 392
 — rhubarb, 398
 — and gentian, 398
 — compound, 398
 — roses, 401
 — senna, compound, 426
 — squill, 420
 — bitter, 420
 — compound, 420
 — stramonium, 445

Wine of sulphate of quinia, 392
 ——— aromatic, 392
 ——— tar, 354
 ——— tartrate of iron, 242
 ——— compound, 242
 ——— tobacco, 452
 ——— valerian, 464
 ——— Virginia snakeroot, 429
 ——— white hellebore, 467
 ——— wormwood, 75
 ——— yellow ladies' bedstraw, 248
 ——— antimonial, 123
 ——— diuretic, 333
 ——— measure, 24
 ——— mulled, 489
 ——— whey, 295
 Wines, 523
 Winter's bark, 46
 ——— tincture, 469
 Wistar's cough lozenges, 255, 345

Witch hazel, 260
 Wood sorrel, 77
 ——— extract, 77
 Worm tea, 441
 Wormseed, 181
 ——— decoction, 181
 ——— oil, 181
 ——— mixture, 181
 Wormwood, 75
 ——— clyster, 75
 ——— essential oil, 76
 ——— extract, 75
 ——— infusion, 75
 ——— oil, 75
 ——— ratafia, 76
 ——— spirit, compound, 76
 ——— syrup, 75
 ——— tincture, 75
 ——— alkaline, 75
 ——— wine, 75

Y

Yellow ladies' bedstraw, 248
 ——— infusion, 248
 ——— wine, 248
 ——— root, 280, 469

Yellow root, extract, 469
 ——— infusion, 469
 ——— sulphate of mercury, 278
 ——— wash, 270

Z

Zinc, 470, 567
 ——— solder for, 485
 ——— acetate, 473
 ——— collyrium, 473
 ——— injection, 473
 ——— lotion, 473
 ——— tincture, 473
 ——— carbonate, 473
 ——— cerate, 473
 ——— plaster, 474
 ——— precipitated, 473
 ——— chloride, 471
 ——— disinfectant, 471
 ——— injection, 471
 ——— lotion, 471
 ——— paste, Canquoin's, 471
 ——— suppository, 471
 ——— tincture, ethereal, 471
 ——— cyanuret, 472
 ——— pills, 472
 ——— powder, 472
 ——— ferrocyanuret, 472
 ——— mixture, 472
 ——— pills, compound, 472
 ——— powder, 472
 ——— solution, 472
 ——— iodide, 472
 ——— collyrium, 472
 ——— ointment, 472
 ——— solution, 473

Zinc, iodide, syrup, 473
 ——— oxide, 470
 ——— cerate, with lycopodium, 471
 ——— lotion, 470
 ——— ointment, 470
 ——— with calomel, 471
 ——— opium, 470
 ——— tutty, 470
 ——— pills, 470
 ——— compound, 470
 ——— powder, 470
 ——— compound, 470
 ——— with columbo, 470
 ——— sulphate, 474
 ——— collyrium, 474
 ——— with camphor, 474
 ——— fomentation, 474
 ——— gargle, 474
 ——— injection, 475
 ——— lotion, 475
 ——— mixture, 475
 ——— pills, 474
 ——— compound, 474
 ——— with turpentine, 474
 ——— myrrh, 474
 ——— solution, 474
 ——— valerianate, 475
 ——— mixture, 475
 ——— pills, 475
 Zittmann's decoction, 414

100	101
102	103
104	105
106	107
108	109
110	111
112	113
114	115
116	117
118	119
120	121
122	123
124	125
126	127
128	129
130	131
132	133
134	135
136	137
138	139
140	141
142	143
144	145
146	147
148	149
150	151
152	153
154	155
156	157
158	159
160	161
162	163
164	165
166	167
168	169
170	171
172	173
174	175
176	177
178	179
180	181
182	183
184	185
186	187
188	189
190	191
192	193
194	195
196	197
198	199
200	201

Y

202	203
204	205
206	207
208	209
210	211
212	213
214	215
216	217
218	219
220	221
222	223
224	225
226	227
228	229
230	231
232	233
234	235
236	237
238	239
240	241
242	243
244	245
246	247
248	249
250	251
252	253
254	255
256	257
258	259
260	261
262	263
264	265
266	267
268	269
270	271
272	273
274	275
276	277
278	279
280	281
282	283
284	285
286	287
288	289
290	291
292	293
294	295
296	297
298	299
300	301

Z

302	303
304	305
306	307
308	309
310	311
312	313
314	315
316	317
318	319
320	321
322	323
324	325
326	327
328	329
330	331
332	333
334	335
336	337
338	339
340	341
342	343
344	345
346	347
348	349
350	351
352	353
354	355
356	357
358	359
360	361
362	363
364	365
366	367
368	369
370	371
372	373
374	375
376	377
378	379
380	381
382	383
384	385
386	387
388	389
390	391
392	393
394	395
396	397
398	399
400	401
402	403
404	405
406	407
408	409
410	411
412	413
414	415
416	417
418	419
420	421
422	423
424	425
426	427
428	429
430	431
432	433
434	435
436	437
438	439
440	441
442	443
444	445
446	447
448	449
450	451
452	453
454	455
456	457
458	459
460	461
462	463
464	465
466	467
468	469
470	471
472	473
474	475
476	477
478	479
480	481
482	483
484	485
486	487
488	489
490	491
492	493
494	495
496	497
498	499
500	501

