Pharmacopoeia of St. Bartholomew's Hospital.

Contributors

Moore, Norman, 1847-1922 Royal College of Physicians of London

Publication/Creation

Edinburgh: Ballantyne Press, 1900.

Persistent URL

https://wellcomecollection.org/works/nxay2vb6

Provider

Royal College of Physicians

License and attribution

This material has been provided by This material has been provided by Royal College of Physicians, London. The original may be consulted at Royal College of Physicians, London. where the originals may be consulted. Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



THE


PHARMACOPŒIA


ST. BARTHOLOMEW'S HOSPITAL


1900.

615.11(47)

615.11(42)








Norman Moore

PHARMACOPŒIA

OF

St. Bartholomew's Mospital



PRINTED AT THE BALLANTYNE PRESS
EDINBURGH AND LONDON
1900

₩OYAL C	LIBRARY
CLASS	615.11 (42)
ACCN.	3751
SOURCE	
DATE	

PREFACE

The only important changes in the present edition of the Hospital Pharmacopæia are such as are necessary to make it accord with the British Pharmacopæia of 1898.

In the latter, where the strength of Preparations has been altered, it has been done, for the most part, with the object of arranging for greater uniformity of dose.

In order to render this uniformity more striking to the eye and more easily remembered, the alphabetical order of the Posological Table is partially broken through in the case of the Extracts, the Liquid Extracts, and the Tinctures, which are separated into divisions.

The Extracts are printed in two divisions:

- 1. Those with the usual dose not exceeding 1 grain.
- 2. Those with the usual dose above 1 grain.

The Liquid Extracts in three divisions :-

- 1. Those with the usual dose less than 5 minims.
- 2. Those with the usual dose between 5 and 15 minims.
- 3. Those with the usual dose from $\frac{1}{2}$ to 1 fluid drachm.

The Tinctures in two divisions:

- 1. Those with dose not usually exceeding 15 minims.
- 2. Those with dose usually from ½ to 1 fluid drachm.

The information thus conveyed, together with the notes as to the Alkaloidal Strengths in certain cases, will probably be found of more value to the Prescriber than a comparison of the present doses with those of a past and obsolete Pharmacopæia.

CONTENTS

				P	AGE
MEDICAL OFFICERS					7
FORMULÆ ALPHABETICALLY AR	RANG	ED			9
TABLE OF PROPORTIONATE DOSE	ES .				55
POSOLOGICAL TABLE					56
HYPODERMIC MEDICATION .					75
LIST OF B.P. PREPARATIONS FO	R EX	TERN	AL	USE	77
LIST OF B.P. SUPPOSITORIES .					80
MEASURES AND WEIGHTS .					81
DIETARY					84



Medical Officers of the Yospital

MR. HOLDEN.
SIR THOMAS SMITH, BART.

Consulting Ophthalmic Surgeon.
MR. POWER.

Physicians.

DR. CHURCH.

DR. GEE.

SIR DYCE DUCKWORTH, M.D.

DR. HENSLEY.

SIRT. LAUDER BRUNTON, M.D., F.R.S.

Assistant Physicians.

DR. NORMAN MOORE.

DR. SAMUEL WEST.

DR. ORMEROD.

DR. HERRINGHAM.

DR. TOOTH.

Physician-Accoucheur.

DR. CHAMPNEYS.

Assistant Physician-Accoucheur.

DR. GRIFFITH.

Ophthalmic Surgeons.

MR. VERNON.

MR. JESSOP.

Dental Surgeons.

MR. PATERSON.

MR. ACKERY.

Surgeons.

MR. WILLETT.

MR. LANGTON.

MR. MARSH.

MR. BUTLIN.

MR. WALSHAM.

Assistant Surgeons.

MR. HARRISON CRIPPS.

MR. BRUCE CLARKE.

MR. BOWLBY.

MR. LOCKWOOD.

MR. D'ARCY POWER.

Aural Surgeon.

MR. CUMBERBATCH.

Medical Officer in Charge of Electrical Department.

DR. LEWIS JONES.

Pathologist.

DR. ANDREWES.

Assistant Dental Surgeons.

MR. READ.

MR. ACKLAND.



PHARMACOPŒIA

[The letters "St.B.P." are appended when the British Pharmacopæia has a preparation of the same name but different composition.]

Aqua Carui (St.B.P.).				
Spirit of Caraway				2 fl. drachms.
Distilled Water				to 1 pint.
Aqua Caryophylli.				
Spirit of Cloves				2 fl. drachms.
Distilled Water				to 1 pint.
Aqua Cassiæ Ligneæ				
Spirit of Cassia Ligne	ea			2 fl. drachms.
Distilled Water				
Aqua Glycyrrhizæ.				
Extract of Liquorice				1 ounce.
Rectified Spirit				1 fl. ounce.
Distilled Water				to 1 pint.
Aqua Menthæ Piperi	itæ (S	t.B.P	.).	
Spirit of Peppermint				2 fl. drachms.
Distilled Water				to 1 pint.
Aqua Rosmarini.				
Spirit of Rosemary				2 fl. drachms.
Distilled Water				to 1 pint.

Balneum Calidum.

Water at 98° F. to 106° F. ... a sufficiency.

Balneum Tepidum.

Water at 85° F. to 92° F. ... a sufficiency.

Balneum Frigidum.

Water at 65° F. or reduced by gradual addition of ice to 40° F. or below ... a sufficiency.

Balneum Alkalinum.

Carbonate of Sodium, 4 ounce for each gallon of water used.

Balneum Furfuris.

Bran, 2 ounces for each gallon of water used.

Mix the Bran with a small quantity of boiling water, and add to sufficient water to give the required quantity.

Balneum Iodi.

Strong Solution of Iodine, & ounce for each gallon of water used.

Balneum Sulphureum.

Sulphurated Potash, 4 ounce for each gallon of water used.

Confectio Ferri Redacti.

 Reduced Iron
 ...
 ...
 30 grains.

 Treacle
 ...
 ...
 1 ounce.

Dose. $-\frac{1}{2}$ to 2 teaspoonfuls.

Confectio Jalapæ.	
Compound Powder of Jalap	$\frac{1}{2}$ ounce.
Oil of Caraway	4 minims.
Glycerine, by weight	$\frac{1}{2}$ ounce.
Dose. $-\frac{1}{2}$ to 2 teaspoonfuls.	
Confectio Potassii Tartratis Acidi.	
Acid Tartrate of Potassium	240 grains.
Ginger, in powder	
Treacle, by weight	7
Dose.— $\frac{1}{2}$ to 2 teaspoonfuls.	
Confectio Sulphuris (St.B.P.).	
	200 amaina
Sublimed Sulphur Acid Tartrate of Potassium	4.5
Treacle, by weight	
Dose.—1 to 2 teaspoonfuls.	
Dose.—1 to 2 teaspoonfuts.	
Confectio Sulphuris Composita.	
Confection of Sulphur	dounce.
Guaiacum Resin, in powder	
Rhubarb Root, in powder	
Nutmeg, in powder	
Treacle, by weight	
Dose.—2 to 4 teaspoonfuls.	
Decoctum Cinchonæ.	
Red Cinchona Bark	1½ ounce.
Distilled Water	1 pint
Dose.—1 fluid ounce.	*

Decoctum Hemidesmi Compositum.

Hemidesmus Root		 	$2\frac{1}{2}$ ounces.
Sassafras Root		 	¹ / ₄ ounce.
Guaiacum Wood		 	dounce.
Mezereon Bark		 	$\frac{1}{8}$ ounce.
Extract of Liquorice	e	 	dounce.
Boiling Distilled Wa	ater	 	$1\frac{1}{2}$ pints.

Digest for one hour, boil for ten minutes, cool, strain, and make up to one pint.

Dose.—1 fluid ounce.

*Emplastrum Belladonnæ (St.B.P.).

Green Extract	of Bellado	nna	 1 part.
Resin Plaster) -61-		O monto
Soan Plaster	or each		 2 parts.

Melt the plasters in a water bath, add the extract, and mix thoroughly.

Emulsio Iodoformi.

Iodoform	 	 1 ounce.
Rectified Spirit	 	 q.s.
Glycerine	 	 7 ounces.
Distilled Water	 	 2 ounces.

Enema Asafetida.

Tincture of Asafetida	 	13	fl.	ounce.
Mucilage of Starch	 	4	fl.	ounces.

Enema Fellis Bovini.

Purified Ox Bile	 	 30 grains.
Soap Enema	 	 1 pint.

^{*} This differs from the Emplastrum Belladonnæ, B.P., 1898, now prepared with a Liquid Extract of the Root.

_				
]	Enema Glycerini.			
	Glycerine			1-4fl.drachms.
	May be used diluted with ab	out an	equa	l quantity of
	water			
]	Enema Nutriens—1.			
	Essence of Beef			3 fl. ounces.
	Milk			1 fl. ounce.
	Mix at a temperature of 95° F.	Win	ne, bro	undy, or opium
	to be added a	s orde	red.	
]	Enema Nutriens—2.			
	Essence of Beef			3 fl. ounces.
	One Egg.			
E	Beat up the egg, and mix with the brandy, or opium		_	
7	Enema Nutriens Peptonia	70 † 11 m	1	
-	To either of the above, at			ture of 150° F
	add one fluid drachm of		upera	
		dano	r Pane	
		-		creaticus: allow
	the mixture to cool to 98	-		creaticus: allow
1		-		creaticus: allow
1	the mixture to cool to 98	5° F. 1	oefore	ereaticus: allow using.
	the mixture to cool to 98 Enema Olei Olivæ. Olive Oil at 90° F	5° F. 1	oefore	ereaticus: allow using.
	the mixture to cool to 98 Enema Olei Olivæ. Olive Oil at 90° F Enema Olei Ricini.	5° F. 1	oefore 	e using.
	the mixture to cool to 98 Enema Olei Olivæ. Olive Oil at 90° F Enema Olei Ricini. Castor Oil	5° F. 1	efore	e using. 1 to 1 pint. 2 fl. ounces.
	the mixture to cool to 98 Enema Olei Olivæ. Olive Oil at 90° F Enema Olei Ricini.	5° F. 1	efore	e using. 1 to 1 pint. 2 fl. ounces.
I	the mixture to cool to 98 Enema Olei Olivæ. Olive Oil at 90° F Enema Olei Ricini. Castor Oil	5° F. 1	efore	e using. 1 to 1 pint. 2 fl. ounces.
I	the mixture to cool to 98 Enema Olei Olivæ. Olive Oil at 90° F Enema Olei Ricini. Castor Oil Mucilage of Starch, tepid	5° F. 1		e using. 1 to 1 pint. 2 fl. ounces. to 1 pint.

Extractum Malti cum Oleo Morrhuæ. Extract of Malt 2 parts Cod Liver Oil 1 part. Dosc.—1 to 4 teaspoonfuls. *Fomentum Acidi Borici. Boric Acid 6 drachms. Boiling Water to 1 pint. *Fomentum Belladonnæ. Green Extract of Belladonna ... 1 drachm. Boiling Water ... 1 pint. *Fomentum Boroglyceridi. Boroglyceride 1 drachm. Boiling Water... ... 1 pint. Boroglyceride is prepared by heating together in a water bath 2 parts of Boric Acid and 3 parts of Glycerine. *Fomentum Papaveris. Bruised Poppy Capsules ... 2 ounces. Water ... 1 pint. Boil the Poppies in 2 quarts of water for 15 minutes, strain, and use the strained fluid hot as a fomentation. *Fomentum Papaveris cum Potassii Chlorate. Chlorate of Potassium ... dounce. Hot Poppy Fomentation ... I pint. ...

^{*} Temperature when used to be 120°-130° F.

F	omentum Terebinthinæ.			
	Oil of Turpentine			2 fl. drachms.
	Sprinkle on Flannel dipped i			
	wrung out; when applied			
	Gutta-Percha			
G	argarisma Acidi Carboli	ci.		
	Glycerine of Carbolic Acid			10 minims.
	Distilled Water			
G	argarisma Acidi Hydroc	hlor	ici.	
	Diluted Hydrochloric Acid			10 minims.
	Decoction of Cinchona			to 1 fl. ounce.
G	argarisma Acidi Tannici	i.		
	Glycerine of Tannic Acid			1 fl. drachm.
	Distilled Water			to 1 fl. ounce.
G	argarisma Aluminis.			
	Alum			10 grains.
	Distilled Water	**		to 1 fl. ounce.
1G	targarisma Aluminis Con	npos	itum.	
	Alum			
	Tincture of Myrrh (St.B.P.			
L	Distilled Water			to 1 fl. ounce.
10	argarisma Boracis.			
	Borax			
	Tincture of Myrrh (St. B.P.			
	Distilled Water			to 1 fl. ounce.

Gargarisma Boracis et P	otassi	ii Ch	loratis.
73			3 grains.
			6 grains.
Tincture of Myrrh (St. B. I			
Distilled Water			
2720022000 110000			oo z ni odnooi
Gargarisma Capsici.			
Tincture of Capsicum			10 minims.
Diluted Sulphuric Acid			6 minims.
Decoction of Cinchona			to 1 fl. ounce.
Gargarisma Chlori.			
Solution of Chlorine of ful	ll stren	igth	
or diluted as may be or	rdered		1 fl. ounce.
See p. 35, under Liqu	or Chl	ori.	
Gargarisma Potassii Chlo	ratis		
Chlorate of Potassium			10 grains.
Tincture of Myrrh (St.B.P			
	.)		o minims.
Distilled Water			
Distilled Water			
Distilled Water Gargarisma Potassii Perr			to 1 fl. ounce.
Gargarisma Potassii Perr Solution of Permanganate	nanga	nati	to 1 fl. ounce.
Gargarisma Potassii Perr Solution of Permanganate	nanga	nati	to 1 fl. ounce.
Gargarisma Potassii Perr Solution of Permanganate Potassium	nanga of	nati	to 1 fl. ounce.
Gargarisma Potassii Perr Solution of Permanganate Potassium	nanga of	nati	to 1 fl. ounce. s. 10 minims.
Gargarisma Potassii Perr Solution of Permanganate Potassium Distilled Water Guttæ Acidi Borici et O	nanga of pii.	nati	to 1 fl. ounce. s. 10 minims. to 1 fl. ounce.
Gargarisma Potassii Perr Solution of Permanganate Potassium Distilled Water Guttæ Acidi Borici et Og Boric Acid	nanga of	nati	to 1 fl. ounce. s. 10 minims.
Gargarisma Potassii Perr Solution of Permanganate Potassium Distilled Water Guttæ Acidi Borici et O	nanga of pii.	nati	to 1 fl. ounce. s. 10 minims. to 1 fl. ounce. 5 grains.

Guttæ Atropinæ Sulphatis.						
Sulph	ate of Atropir	ne			1 grain.	
-	ed Water				to 1 fl. ounce.	
Guttæ A	Atropinæ et	Cocai	næ.			
Sulpha	ate of Atropin	ie			2 grains.	
Hydro	chloride of C	ocaine			5 grains.	
Distill	ed Water				to 1 fl. ounce.	
Guttæ A	Atropinæ et	Zinci	Sulp	hati	s.	
Sulpha	ate of Atropir	ne			1 grain.	
Sulph	ate of Zinc				1 grain.	
Distill	ed Water				to 1 fl. ounce.	
Guttæ C	ocainæ.					
Hydro	chloride of C	ocaine			5 grains.	
	2 777 /				to 1 fl. ounce.	
M 11 O						
Guttæ C	cainæ et (Olei R	icini.			
Cocair					20 grains.	
Castor	· Oil				to 1 fl. ounce.	
Guttæ I	Homatropina	e.				
Hydro	bromide of H	omatro	opine		2 grains.	
	ed Water				to 1 fl. ounce.	
Guttæ Homatropinæ et Cocainæ.						
Homa	tropine				10 grains.	
Cocair					10 grains.	
Almor					to 1 fl. ounce.	
Guttæ Opii.						
	_	pium			2 fl. drachms.	
					to 1 fl. ounce.	
		1000		1000	I in ounce.	

Guttæ Physostigminæ.			0		
Sulphate of Physostigmine					
Distilled Water			to 1 fl. ounce.		
Guttæ Zinci.					
Sulphate of Zinc			2 grains.		
Distilled Water			to 1 fl. ounce.		
*TT					
*Haustulus Ammonii Chlor	riai et	1p			
Chloride of Ammonium			$2\frac{1}{2}$ grains.		
Wine of Ipecacuanha			3 minims.		
Syrup of Tolu			10 minims.		
Camphor Water			to 1 fl. drachm.		
*Haustulus Belladonnæ Co	mposi	tus.			
Tincture of Belladonna			1 minim.		
Tincture of Squill			1 minim.		
Wine of Ipecacuanha			1 minim.		
Syrup of Tolu			10 minims.		
Distilled Water			to 1 fl. drachm.		
*Haustulus Bismuthi Comp	ositus	5.			
Oxycarbonate of Bismuth			2 grains.		
Bicarbonate of Sodium			3 grains.		
Compound Tincture of Care	damom	s	10 minims.		
Caraway Water (St. B. P.)					
*Haustulus Carminativus.					
Bicarbonate of Sodium			2 grains.		
Aromatic Spirit of Ammoni					
Glycerine			10 minims.		
Caraway Water (St. B.P.)					
* Where the word Haustulus is used draught suitable for a child					

For table of variation of doses at different ages, see page 55.

*Haustulus Cinchonæ. Compound Tincture of Cinchona Chlorate of Potassium Diluted Nitro-Hydrochloric Acid Glycerine Caraway Water (St.B.P.)		2 grains.2 minims.5 minims.
*Haustulus Ferri cum Oleo Morri Solution of Citrate of Iron and Ammonium of each Cod Liver Oil		
*Haustulus Ferri cum Oleo Morri Solution of Citrate of Iron a Ammonium Cod Liver Oil Extract of Malt Saccharated Solution of Lime	and 	20 minims. 10 minims. 20 minims.
*Haustulus Magnesii Sulphatis c Sulphate of Magnesium Compound Tincture of Rhubarb Caraway Water (St.B.P.)		5 grains. 10 minims.
*Haustulus Olei Ricini. Castor Oil Mucilage of Acacia Dill Water		10 minims.
*Haustulus Potassii Bromidi Con Bromide of Potassium Tincture of Belladonna Syrup Distilled Water * See Foot-note, p. 18		 1½ grains. 1 minim. 15 minims.

*Haustulus Santonini et Olei Ricini.

Santonine, in powder ... 1 grain.

Castor Oil ... of each ... 15 minims.

Liquorice Water ... to 1 fl. drachm.

Haustus Acidi Gallici.

Glycerine of Gallic Acid ... 25 minims.

Rectified Spirit 15 minims.

Distilled Water ... to 1 fl. ounce.

Contains 5 grains of Gallic Acid.

Haustus Acidi Hydrocyanici.

Diluted Hydrocyanic Acid ... 4 minims.

Bicarbonate of Sodium ... 5 grains.

Distilled Water ... to 1 fl. ounce.

**Haustus Acidi Nitro-hydrochlorici.

Diluted Nitro-Hydrochloric Acid ... 10 minims.

Infusion of Orange Peel ½ fl. ounce.

Chloroform Water ... to 1 fl. ounce.

Haustus Aconiti cum Colchico et Quinina.

Tincture of Aconite ... 7 minims.

Tincture of Colchicum Seeds ... 6 minims.

Sulphate of Quinine ... 1 grain.

Diluted Sulphuric Acid ... 1½ minims.

Chloroform Water ... to 1 fl. ounce.

* See Foot-note, p. 18.

^{**} When a double asterisk is printed against any Haustus, it signifies that such draught may in reduced quantity be considered suitable for children.

**Haustus Ætheris Ammoniatus.	
Spirit of Ether	$\frac{1}{2}$ fl. drachm.
Aromatic Spirit of Ammonia .	
Distilled Water	
**Haustus Albus.	
Carbonate of Magnesium	15 grains.
Sulphate of Magnesium	
Peppermint Water (St.B.P.)	to 1 fl. ounce.
**Haustus Ammonii Acetatis.	
Solution of Acetate of Ammonium	4 fl. drachms.
Distilled Water	to 1 fl. ounce.
Haustus Ammonii Acetatis cun	Camphora.
	- Constitution of the
Solution of Acetate of Ammonium Camphor Water	2 fl. drachm.
Solution of Acetate of Ammonium	2 fl. drachm.
Solution of Acetate of Ammonium	2 fl. drachm. to 1 fl. ounce.
Solution of Acetate of Ammonium Camphor Water **Haustus Ammonii Acetatis cur	2 fl. drachm. to 1 fl. ounce. n Scilla.
Solution of Acetate of Ammonium Camphor Water	2 fl. drachm. to 1 fl. ounce. n Scilla. 20 minims.
Solution of Acetate of Ammonium Camphor Water **Haustus Ammonii Acetatis cur	2 fl. drachm. to 1 fl. ounce. n Scilla. 20 minims.
Solution of Acetate of Ammonium Camphor Water	2 fl. drachm. to 1 fl. ounce. n Scilla. 20 minims.
Solution of Acetate of Ammonium Camphor Water	2 fl. drachm. to 1 fl. ounce. n Scilla. 20 minims. to 1 fl. ounce.
Solution of Acetate of Ammonium Camphor Water	2 fl. drachm. to 1 fl. ounce. n Scilla. 20 minims. to 1 fl. ounce.
Solution of Acetate of Ammonium Camphor Water **Haustus Ammonii Acetatis cur Vinegar of Squill Draught of Acetate of Ammonia **Haustus Ammonii Chloridi. Chloride of Ammonium	2 fl. drachm. to 1 fl. ounce. n Scilla. 20 minims. to 1 fl. ounce. 10 grains. 11 fl. drachm.
Solution of Acetate of Ammonium Camphor Water	2 fl. drachm. to 1 fl. ounce. n Scilla. 20 minims. to 1 fl. ounce. 10 grains. 11 fl. drachm.
Solution of Acetate of Ammonium Camphor Water	2 fl. drachm. to 1 fl. ounce. n Scilla. 20 minims. to 1 fl. ounce. 10 grains. 11 fl. drachm.
Solution of Acetate of Ammonium Camphor Water	2 fl. drachm. to 1 fl. ounce. n Scilla. 20 minims. to 1 fl. ounce. 10 grains. 11 fl. drachm. to 1 fl. ounce.
Solution of Acetate of Ammonium Camphor Water **Haustus Ammonii Acetatis cur Vinegar of Squill Draught of Acetate of Ammonia **Haustus Ammonii Chloridi. Chloride of Ammonium Aromatic Spirit of Ammonia Chloroform Water Haustus Ammoniaci Fetidus. Fetid Spirit of Ammonia	2 fl. drachm. to 1 fl. ounce. n Scilla. 20 minims. to 1 fl. ounce. 10 grains. 11 fl. drachm. to 1 fl. ounce.

^{**} See Foot-note, p. 20.

Haustus Antimonii cum Antimonial Wine Sulphate of Magnesium Distilled Water Contains \(\frac{1}{8} \) grain of Tar			30 minims. 60 grains. to 1 fl. ounce.
**Haustus Bismuthi. Oxycarbonate of Bismuth Bicarbonate of Sodium Chloroform Water			10 grains. 10 grains. to 1 fl. ounce.
**Haustus Calumbæ Alkali Bicarbonate of Sodium Tincture of Calumba Spirit of Chloroform Infusion of Orange Peel			10 grains. 20 minims. 5 minims. to 1 fl. ounce.
**Haustus Cascaræ Sagrada Liquid Extract of Cascara Aromatic Spirit of Ammor Syrup of Ginger Glycerine Peppermint Water	nia 		
Spirit of Chloroform		:	10 minims. 5 minims. to 1 fl. ounce.
Haustus Chlori. Glycerine †Solution of Chlorine of full or diluted as ordered	strengt	h,	o 1 fl. ounce.
** See Foot-note † See p. 35, under Lie		ori.	

**Haustus Chloroformi Comp	positus.			
Chloroform		1 minim.		
Diluted Hydrocyanic Acid		3 minims.		
Solution of Acetate of Morp	ohine	3 minims.		
Tincture of Cascarilla		20 minims.		
Diluted Nitric Acid		10 minims.		
Glycerine		60 minims.		
Distilled Water		to 1 fl. ounce.		
**Haustus Cinchonæ Acidus				
Spirit of Chloroform		10 minims.		
Aromatic Sulphuric Acid		10 minims.		
Decoction of Cinchona		to 1 fl. ounce.		
Haustus Colchici.				
Tincture of Colchicum Seed	s	10 minims.		
Carbonate of Magnesium		10 grains.		
Peppermint Water (St. B. P.)		to 1 fl. ounce.		
Haustus Copaibæ.				
Copaiba		15 minims.		
Spirit of Nitrous Ether		15 minims.		
Solution of Potash		5 minims.		
Mucilage of Gum Acacia		60 minims.		
Camphor Water		to 1 fl. ounce.		
Haustus Copaibæ et Cubeb	æ.			
Cubebs, in powder		30 grains.		
Copaiba Draught		to 1 fl. ounce.		
**Haustus Cretæ Aromaticus.				
Aromatic Powder of Chalk		30 grains.		
Cassia Water		to 1 fl. ounce.		
** See Foot-note,	p. 20.			

**Haustus Cretæ Aromaticus cum Opio.

Aromatic Powder of Chalk and
Opium 20 grains.
Cassia Water to 1 fl. ounce.

Contains ½ grain of Opium.

**Haustus Effervescens Ammoniatus.

Carbonate of Ammonium ... 15 grains.

Distilled Water ... to 1 fl. ounce.

This Solution to be added to the following:—

Citric Acid 17 grains.

Distilled Water ... to 1 fl. ounce.

To be taken during effervescence.

Haustus Ferri Alkalinus.

Citrate of Iron and Ammonium ... 5 grains.

Bicarbonate of Sodium ... 10 grains.

Spirit of Chloroform ... 5 minims.

Infusion of Calumba ... to 1 fl. ounce.

**Haustus Ferri et Ammonii Citratis.

Citrate of Iron and Ammonium ... 8 grains.

Carbonate of Ammonium ... 2 grains.

Chloroform Water to 1 fl. ounce.

Haustus Ferri cum Nuce Vomica.

Solution of Perchloride of Iron ... 10 minims.

Tincture of Nux Vomica ... 3 minims.

Diluted Phosphoric Acid ... 5 minims.

Chloroform Water ... to 1 fl. ounce.

	Haustus Ferri et Potassii Iodi	di.	
	Tartarated Iron		5 grains.
	Iodide of Potassium		
	Peppermint Water (St.B.P.)		
	Haustus Ferri cum Quassia.		
	Solution of Perchloride of Iron		15 minims.
	Infusion of Quassia		to 1 fl. ounce.
*	*Haustus Ferri Salinus.		
	Citrate of Potassium		20 grains.
	Draught of Iron and Ammoniu		
	Citrate		to 1 fl. ounce.
*	*Haustus Filicis Maris.		
	Liquid Extract of Male Fern		1 fl. drachm.
			20 grains.
	~		20 minims.
	Peppermint Water (St.B.P.)		
			Tar outloo.
*	*Haustus Flavus.		
	Caramel		a sufficiency.
	Distilled Water		to 1 fl. ounce.
	Haustus Gentianæ cum Rheo A	lk	alinus.
	Bicarbonate of Sodium		10 grains.
	Oil of Peppermint		
	G : 11 C GI 1 C		5 minims.
	*		2 fl. drachms.
	Compound Infusion of Gentian		to 1 fl. ounce.
	** See Foot-note, p. 20.		

Haustus Gentianæ cum Rheo et Potassii Bromido. Bromide of Potassium 10 grains. Alkaline Draught of Gentian and Rhubarb ... to 1 fl. ounce. Haustus Hemidesmi cum Potassii Iodido. Iodide of Potassium ... 5 grains. Compound Decoction of Hemidesmus to 1 fl. ounce. Haustus Hydrargyri Perchloridi. Solution of Perchloride of Mercury 1 fl. drachm. ... a sufficiency. Caramel Chloroform Water ... to 1 fl. ounce. Contains \(\frac{1}{10}\) grain of Perchloride of Mercury. Haustus Hydrargyri Perchloridi cum Potassii Todido. Iodide of Potassium ... 5 grains. Draught of Perchloride of Mercury 1 fl. ounce. Contains about 10 grain of Red Iodide of Mercury. **Haustus Ipecacuanhæ Ammoniatus. Wine of Ipecacuanha .. 10 minims. Carbonate of Ammonium 5 grains. ... 1 fl. drachm. Syrup of Tolu Distilled Water ... to 1 fl. ounce. Haustus Ipecacuanhæ Emeticus. Wine of Ipecacuanha Oxymel of Squill ... of each ... 1/2 fl. ounce.

** See Foot-note, p. 20.

Haustus Menthæ Sulphuricus.

Diluted Sulphuric Acid ... 10 minims.

Syrup of Red Poppy (St.B.P.) ... 30 minims.

Peppermint Water (St.B.P.) ... to 1 fl. ounce.

Haustus Menthæ Sulphuricus cum Ferri Sulphate.

Sulphate of Iron ... 1 grain.

Spirit of Chloroform ... 10 minims.

Sulphuric Mint Draught ... to 1 fl. ounce.

Haustus Menthæ Sulphuricus cum Magnesii Sulphate.

Sulphate of Magnesium ... 60 grains. Sulphuric Mint Draught ... to 1 fl. ounce.

**Haustus Menthæ Sulphuricus cum Papavere et Scilla (Haustus pro Tussi).

Syrup of Poppy Capsules ... 30 minims.

Tincture of Squill ... 10 minims.

Sulphuric Mint Draught ... to 1 fl. ounce.

Haustus Morphinæ Hydrochloridi.

Solution of Hydrochloride of Morphine 20 minims.

Distilled Water ... to 1 fl. ounce.

Contains nearly & grain of Hydrochloride of Morphine.

^{**} See Foot-note, p. 20.

Haustus Morphinæ Hydrochloridi	cum Ferro.
Solution of Hydrochloride of Mor-	
	10 minims.
Solution of Perchloride of Iron	15 minims.
Diluted Hydrochloric Acid	3 minims.
Chloroform Water	to 1 fl. ounce.
Contains nearly 1 grain of Hydrochloride	of Morphine.
**Haustus Olei Morrhuæ.	
Cod Liver Oil	
Cod Liver Oil } each !	2 fl. drachms.
Refined Sugar, in powder	30 grains.
Caraway Water (St.B.P.)	to 1 fl. ounce.
Haustus Opiatus.	
Tincture of Opium	10 minims.
Chloroform Water t	to 1 fl. ounce.
Contains the soluble matter of nearly 3 grain	n of Opium.
Haustus Plumbi Acetatis.	
Acetate of Lead	grains.
Oxymel	l fl. drachm.
Distilled Water t	o 1 fl. ounce.
**Haustus Potassii Citratis.	

Citrate of Potassium

Distilled Water

20 grains.

to 1 fl. ounce.

1	Haustus Potassii Citratis Effervescens.
	Bicarbonate of Potassium 20 grains.
	Distilled Water to 1 fl. ounce.
	This Solution to be added to the following:—
	Citric Acid, in powder 14 grains.
	Distilled Water to 1 fl. ounce.
	To be taken during effervescence.
	Haustus Potassii Chloratis.
	Chlorate of Potassium 10 grains.
	Distilled Water to 1 fl. ounce.
	Haustus Potassii cum Hyoscyamo.
	Nitrate of Potassium 10 grains.
	Bicarbonate of Potassium 20 grains.
	Tincture of Henbane 30 minims.
	Chloroform Water to 1 fl. ounce.
	Haustus Potassii Iodidi Ammoniatus.
	Iodide of Potassium 5 grains.
	Carbonate of Ammonium 2 grains.
	Distilled Water to 1 fl. ounce.
*	Haustus Quininæ Sulphatis.
	Sulphate of Quinine 1 grain.
	Diluted Sulphuric Acid 1½ minims.
	Distilled Water to 1 fl. ounce.
	Haustus Quininæ Sulphatis cum Ferro.
	Sulphate of Quinine 1 grain.
	Sulphate of Iron ½ grain.
	Spirit of Chloroform 10 minims.
	Sulphuric Mint Draught to 1 fl. ounce.
	** See Foot-note, p. 20.

Haustus Quininæ Iodido.	Sulphatis	cu	ım Potassii		
Sulphate of Quinine			1 orgin		
Iodide of Potassium			0		
Diluted Sulphuric A	oid		5 grains. 1½ minims.		
TO 1 1111 3 TYPE 1			to 1 fl. ounce.		
Distilled Water .			to 1 n. ounce.		
**Haustus Rhei Comp	ositus.				
Compound Rhubarb	Powder		30 grains.		
Caraway Water (St.I	B.P.)		to 1 fl. ounce.		
Haustus Salol.					
Salol			8 grains.		
Mucilage of Acacia.			2 fl. drachms.		
Cassia Water .			to 1 fl. ounce.		
Hanatua Saanarii Car	nnogitug				
Haustus Scoparii Con			00 1		
Tartrate of Potassiur			20 grains.		
Spirit of Juniper .			30 minims.		
Infusion of Broom .			to 1 fl. ounce.		
Haustus Senegæ Am	moniatus.				
Carbonate of Ammor	nium		5 grains.		
Tincture of Squill			15 minims.		
Infusion of Senega			$\frac{1}{2}$ fl. ounce.		
Chloroform Water					
**Haustus Sennæ Compositus (House Physic).					
Sulphate of Magnesi	um		1 ounce.		
Spirit of Peppermint			10 minims.		
Extract of Liquorice			1 drachm.		
Infusion of Senna			to 1 fl. ounce.		

^{**} See Foot-note, p. 20.

Haustus Serpentariæ Ammo	niatus.	
Carbonate of Ammonium		5 grains.
Tincture of Serpentary		
		10 minims.
Peppermint Water (St.B.P.)		to 1 fl. ounce.
Haustus Sodii Salicylatis Co	mposit	us.
Salicylate of Sodium .	_	
Aromatic Spirit of Ammonia		20 minims.
Chloroform Water		
Haustus Sodii Salicylatis et	Quinir	ıæ.
Salicylate of Sodium		10 grains.
Sulphate of Quinine		1 grain.
		to 1 fl. ounce.
Haustus Sodii Tartratis Effe	ervesce	ns.
Bicarbonate of Sodium		20 grains.
Distilled Water		
This Solution to be added to		
Tartaric Acid, in powder		17 grains.
W. I. III W. Market		to 1 fl. ounce.
To be taken during e	ffervescer	ace.
Haustus Strychninæ et Acid	li Phosi	phorici.
Solution of Hydrochlori		
Strychnine		
Diluted Phosphoric Acid .		

Contains about ¹/₃₆ grain of Hydrochloride of Strychnine.

Infusion of Quassia ... to 1 fl. ounce.

... 15 minims.

Spirit of Chloroform

Haustus Terebinthi	næ.			
Oil of Turpentine				20 minims.
Mucilage of Gum A				2 fl. drachms.
Caraway Water (St				to 1 fl. ounce.
•				
Haustus Valerianæ	cum I	Iyosc	yamo).
Ammoniated Tinct	ure of	Valeria	an	30 minims.
Tincture of Henbar	ne			20 minims.
Camphor Water				to 1 fl. ounce.
Haustus Zinci Emet	cicus.			
Sulphate of Zinc				30 grains.
Distilled Water				to 1 fl. ounce.
*Toisetie Asidi Desie				
*Injectio Acidi Boric				
·Boric Acid				
Water				to 1 pint.
*Injectio Aluminis	et Zin	ci.		
Alum				40 grains.
Sulphate of Zinc				
Water				
*Injectio Bismuthi.				
Subnitrate of Bism				
Water				to 1 pint.
*T-: II-rdnongram	: Todi	1; D.,	hni	
*Injectio Hydrargyr				
Red Iodide of Mero				
Iodide of Potassiun				20 grains.
Water				to 1 pint.

^{*} These Injections are intended for vaginal use; for others, see under Lotions, p. 36.

*	Injectio Hydrargyri Perchloridi.
	Perchloride of Mercury 5 grains.
	Glycerine $\frac{1}{2}$ fl. ounce.
	Water to 1 pint.
*	Injectio Potassii Permanganatis.
	Solution of Permanganate of Potas- \1 ft ounce
	Solution of Permanganate of Potas- sium $\int_{-1}^{1} fl$. ounce.
	Water to 1 pint.
×	Injectio Sodii Bicarbonatis.
	Bicarbonate of Sodium 60 grains.
	Water to 1 pint.
*	Injectio Zinci Sulphocarbolatis.
	Sulphocarbolate of Zinc 120 grains.
	Water to 1 pint.
	For Hypodermic Injections see p. 75.
	Linctus.
	Acetic Acid 2 minims.
	Syrup of Poppy Capsules 36 minims.
	Vinegar of Squill 36 minims.
	Tragacanth, in powder 6 grains.
	Confection of Hips 100 grains.
	Water to 1 fl. ounce.
	$Dose1 \ teaspoonful.$
	Linctus cum Ipecacuanha.
	Wine of Ipecacuanha 30 minims.
	Linctus to 1 fl. ounce.
	Dose.—1 teaspoonful, containing nearly 4 minims of
	Ipecacuanha Wine.
	* See Foot-note, p. 32.

Linctus Scillæ Opiatus.

Compound Tincture of Camphor
Oxymel of Squill ... equal parts.
Syrup of Tolu

Dose.—1 teaspoonful, containing the soluble matter of $\frac{1}{12}$ grain of Opium.

Linimentum Belladonnæ (St.B.P.).

Belladonna Root, in coarse powder 20 ounces.

Camphor 1 ounce.

Methylated Alcohol ... to 30 fl. ounces.

Linimentum Calcis (St.B.P.) (Carron Oil).

Solution of Lime Linseed Oil equal parts.

Linimentum Cantharidis.

Vinegar of Cantharides ... 6 fl. drachms.

Glacial Acetic Acid ... 40 minims.

Spirit of Rosemary ... 20 minims.

Glycerine ... 6 fl. drachms.

Distilled Water ... to 4 fl. ounces.

Linimentum Cantharidis cum Ammonia.

Tincture of Cantharides ... 80 minims.

Strong Solution of Ammonia ... 160 minims.

Spirit of Rosemary ... 20 minims.

Glycerine ... 160 minims.

Almond Oil ... 160 minims.

6 fl. drachms.

Distilled Water ... to 4 fl. ounces.

Linimentum Iodi Dilutum. Tincture of Iodine Strong Solution of Iodine Contains about 37 grains of Iodine in a fluid ounce.
Linimentum Opii (St.B.P.). Tincture of Opium 1 fl. ounce. Ammoniated Liniment of Soap to 4 fl. ounces Amount of Opium 3ths that of B.P. Liniment.
Linimentum Plumbi Subacetatis. Solution of Subacetate of Lead ½ fl. ounce. Olive Oil to 4 fl. ounces
Linimentum Potassii Iodidi.2 drachms.Iodide of Potassium2 drachms.Solution of Ammonia $\frac{1}{2}$ fl. ounce.Ammoniated Liniment of Soapto 4 fl. ounces
Linimentum Saponis cum Ammonia. Soft Soap
Liquor Chlori. Chlorate of Potassium

Add the Acid to the Chlorate in a large bottle; when the Chlorine given off has displaced the air, add the water gradually, corking and shaking the bottle after each addition.

*Liquor Ferri et An	nmoni	i Citra	atis.	
Citrate of Iron and	Ammo	nium		12 grains.
				to 1 fl. ounce.
Lotio Acidi Borici				
Boric Acid				10 grains.
Distilled Water				
Lotio Acidi Borici	cum G	lycer	ino.	
Boric Acid				15 grains.
Glycerine				
Distilled Water				
Lotio Acidi Carboli	ci cun	n Iodo).	
Carbolic Acid, lique	fied			12 minims.
Tincture of Iodine				
Distilled Water				
Lotio Acidi Nitrici	cum	Opio.		
Diluted Nitric Acid				3 minims.
Opium Lotion				
Lotio Acidi Tannici	Sulp	hurosa	a.	
Glycerine of Tannic	_			1 fl. drachm.
				1 fl. drachm.
-Distilled Water				to 1 fl. ounce.
Lotio Aluminis (Three	Stren	gths).		
Alum, in powder				4, 8, or 16 grs.
				to 1 fl. ounce.
* This has in former ed				

Lotio Ammonii Chlo	ridi.			
Chloride of Ammonia	um			15 grains.
Diluted Acetic Acid				20 minims.
Methylated Spirit				20 minims.
Distilled Water				to 1 fl. ounce.
Lotio Argenti Nitra	tis (V	aried S	Strengt	ths).
Nitrate of Silver			from	1 to 15 grains.
Distilled Water				to 1 fl. ounce.
Lotio Boracis.				
Borax				10 grains.
Distilled Water				to 1 fl. ounce.
Lotio Calaminæ Con	nposi	ta.		
Prepared Calamine				1 ounce.
Oxide of Zinc				dounce.
Solution of Subaceta	te of	Lead		1 fl. drachm.
Glycerine				1 fl. drachm.
Lime Water				to 1 fl. ounce.
Lotio Cupri Sulphat	is.			
Sulphate of Copper				2 grains.
Distilled Water				to 1 fl. ounce.
Lotio Ferri Sulphati	S.			
Sulphate of Iron				3 grains.
Distilled Water				to 1 fl. ounce.
Lotio Opii.				
Liquid Extract of Or	oium			1 fl. drachm.
Distilled Water		***		to 1 fl. ounce.

Lotio Plumbi.				
Strong Solution of Su	baceta	te of I	ead	6 minims.
Distilled Water				to 1 fl. ounce.
Lotio Plumbi cum O	pio,			
Liquid Extract of Op				
Lead Lotion				to 1 fl. ounce.
Tatio Cala Chianina				
Lotio Sodæ Chlorina				
Solution of Chlorinate				
Distilled Water				to 1 fl. ounce.
Lotio Sodii Chloridi,				
Chloride of Sodium				15 grains.
Distilled Water				to 1 fl. ounce.
Lotio Sulphuris Comp	posita			
Precipitated Sulphur				
Spirit of Camphor				
Lime Water				to 1 fl. ounce.
Lotio Zinci Composit	a.			
Sulphate of Zinc				1 grain.
Acetate of Zinc				
Tincture of Catechu				
Liquid Extract of Op	ium			30 minims.
Distilled Water		* * *		to 1 fl. ounce.
Lotio Zinci Oxidi.				
Oxide of Zinc				30 grains.
Glycerine				$\frac{1}{2}$ fl. drachm.
Lime Water				to 1 fl. ounce.

Lotio Zinci Sulphat	tis (Thre	ee Stre	engths).
Sulphate of Zinc				1, 2, or 4 grains.
Distilled Water				to 1 fl. ounce.
Lotio Zinci Sulpho	carbola	tis.		
Sulphocarbolate of	Zinc			4 grains.
Distilled Water				to 1 fl. ounce.
Oleum Acidi Carbo	lici.			
Carbolic Acid, liqu	efied			12 minims.
				to 1 fl. ounce.
Contains nearly $2\frac{1}{2}$				
Pasta Acidi Salicy	lici.			
Salicylic Acid				1 part.
Glycerine	***			9 parts.
Pasta Calcis et Po	tassæ 1	Alcoh	olica.	
Caustic Potash, in	powder			5 parts.
Slaked Lime				6 parts.
Rectified Spirit				a sufficiency.
Pasta Zinci Chloric	di.			
Chloride of Zinc, in Flour	n powde	r }		equal parts.
Glycerine		***		a sufficiency.
Pigmentum Acidi	Chromi	ci (Va	rying	Strengths).
				10-180 grains.
Distilled Water				to 1 fl. ounce.

Pigmentum Acidi Tannic	ei.		
Tannic Acid			
Glycerine			1 fl. ounce.
This is three and a half times Tannic		ngth	of Glycerine of
1 annic	Acu.		
Pigmentum Belladonnæ.			
Green Extract of Bellado Glycerine	nna (equal parts.
diyeemie			
Pigmentum Chloral Com	positu	m.	
Hydrate of Chloral			
Menthol of ex	ach		1 part.
Camphor			
Pigmentum Ferri Perchl	oridi (Two	Strengths).
Perchloride of Iron, cryst	tallised		60-120 grains.
Glycerine			
Distilled Water			
Pigmentum Hydrargyri			
Cyanide of Mercury			15 grains.
Distilled Water			to 1 fl. ounce.
Pigmentum Iodi.			
Iodine	***		30 grains.
Glycerine			
Heat in a water bath till	the Iodin	ne is	dissolved.
Pilula Aloes et Nucis V	omicæ		
Pill of Barbadoes Aloes			4½ grains.
Extract of Nux Vomica			‡ grain.
Syrup of Glucose			a sufficiency.

70.17	9					
D1 111	0	Λ -	- 71	αm	7 77	\sim
Pilu	La			UL	111	OU.
the war for				_ L		

Atropine $\frac{1}{100}$ grain.Rectified Spirit $\frac{1}{20}$ minim.Liquorice Powder2 grains.Syrup of Glucosea sufficiency.

Pilula Belladonnæ.

Green Extract of Belladonna $\frac{1}{2}$ grain.

Liquorice Powder ... 2 grains.

Syrup of Glucose ... a sufficiency.

Pilula Calcii Sulphidi (Three Strengths).

Sulphide of Calcium ... $\frac{1}{8}$, $\frac{1}{4}$, or $\frac{1}{2}$ grain. Sugar of Milk ... 2 grains. Glycerine of Tragacanth ... a sufficiency.

Pilula Camphoræ et Hyoscyami.

Camphor ... $2\frac{1}{2}$ grains. Extract of Henbane

Pilula Cannabis Indicæ (Three Strengths).

Extract of Indian Hemp ... $\frac{1}{4}$, $\frac{1}{2}$, or 1 grain. Liquorice Powder ... a sufficiency.

Pilula Cascaræ Sagradæ.

Extract of Cascara Sagrada ... $2\frac{1}{2}$ grains. Liquorice Powder ... a sufficiency.

^{*} Smaller sizes of these pills $(\frac{1}{2}, \frac{1}{4}, \frac{1}{8})$ are also kept for children, to be broken up at time of taking, and mixed with syrup or fruit paste.

Pilula Cascaræ Sagradæ et Nu	cis	Vomicæ.
Extract of Cascara Sagrada		$2\frac{1}{2}$ grains.
Extract of Nux Vomica		d grain.
Capsicum Fruit, in powder		
Liquorice Powder		a sufficiency.
Pilula Codeinæ.		
Codeine		½ grain.
Liquorice Powder		1 grain.
Glycerine of Tragacanth		a sufficiency.
Pilula Creosoti.		
Creosote		1 minim.
Hard Soap		
Liquorice Powder		
Pilula Creosoti et Galbani.		
Creosote		1 minim.
Compound Pill of Galbanum		
Compound Rhubarb Pill		
Pilula Elaterii Composita.		
Elaterium		‡ grain.
Subchloride of Mercury		1½ grain.
Compound Extract of Colocynth		2 grains.
Capsicum Fruit, in powder		$\frac{1}{2}$ grain.
Syrup of Glucose		a sufficiency.
Pilula Euonymi Composita.		
Dry Extract of Euonymus		1 grain.
		1½ grain.
Ipecacuanha, in powder		½ grain.
Extract of Henbane		

Pi	lula Ferri Sulphatis.			
	Dried Sulphate of Iron			1 grain.
	Ginger, in powder			
	Extract of Gentian			
Pi	lula Hydrargyri et Colo	cynth	idis	
	Mercurial Pill			
	Compound Extract of Colo			
Pi	lula Hydrargyri Colocy	nthidi	s e	t Hvoscvami.
	Mercurial Pill			
	Compound Extract of Colo			
	Extract of Henbane			
				- 8
Pi	lula Hydrargyri et Dig	italis	Con	nposita.
	Mercurial Pill			1 grain.
	Powdered Digitalis			1 grain.
	Powdered Squill			1 grain.
	Extract of Henbane			
P	ilula Hydrargyri et Rhe	ei.		
	Mercurial Pill			2 grains.
	Compound Rhubarb Pill			
P	ilula Hydrargyri et Scil	læ.		
	Mercurial Pill			3 grains.
	T 7 7 7 111			1½ grain.
· D				20
P	ilula Hydrargyri cum C			
	Mercury with Chalk			3 grains.
	Gentian Root, in powder Syrup of Glucose			a sufficiency.
	by rup or Grucose			•

^{*} See Foot-note, p. 41.

*Pilula Hydrargyri cum Creta Composita. Mercury with Chalk 2½ grains. Compound Powder of Ipecacuanha 2½ grains. Syrup of Glucose a sufficiency. Contains ¼ grain each of Opium and Ipecacuanha.
*Pilula Hydrargyri cum Creta et Quininæ.
Mercury with Chalk 3 grains.
Sulphate of Quinine 1 grain.
Syrup of Glucose a sufficiency.
Syrup of Grucose a sumciency.
*Pilula Hydrargyri cum Creta et Rhei.
Mercury with Chalk 2½ grains.
Rhubarb, in powder 2½ grains.
Syrup of Glucose a sufficiency.
Pilula Hydrargyri Iodidi Viridis et Opii.
Green Iodide of Mercury ½ grain.
Opium ½ grain.
Syrup of Glucose a sufficiency.
Pilula Hydrargyri Subchloridi et Colocynthidis.
Subchloride of Mercury 1 grain.
Compound Extract of Colocynth 4 grains.
Pilula Hydrargyri Subchloridi Colocynthidis et Hyoscyami.
Subchloride of Mercury 1 grain.
Compound Extract of Colocynth 3 grains.
Extract of Henbane 1 grain.

* See Foot-note, p. 41.

Pilula Hydrargyri Subchloridi et Jalapæ.

(House Pill.)

Subchloride of Mercury ... 1 grain.

Jalap ... 3 grains.

Syrup of Glucose ... a sufficiency.

Pilula Hydrargyri Subchloridi et Rhei.

Subchloride of Mercury ... 1 grain.

Rhubarb ... 4 grains.

Syrup of Glucose ... a sufficiency.

Pilula Hydrargyri Subchloridi et Scammonii.

Subchloride of Mercury ... 1 grain.

Scammony ... 3 grains.

Syrup of Glucose ... a sufficiency.

Pilula Rhei et Nucis Vomicæ.

Compound Rhubarb Pill ... 3½ grains.

Extract of Nux Vomica ¼ grain.

Capsicum Fruit, in powder ½ grain.

Pilula Santonini.

Santonine ... 3 grains.

Extract of Gentian ... a sufficiency.

Pilula Zinci Oxidi et Belladonnæ.

^{*} See Foot-note, p. 41.

Pilula Zinci Valerianatis Composita Valerianate of Zinc Compound Galbanum Pill Compound Rhubarb Pill	1 grain. 3 grains.
Potus Potassii Tartratis Acidi (Imp	erial Drink).
Acid Tartrate of Potassium Half a fresh Lemon	60 grains.
Sugar	a sufficiency.
Boiling Water	1 pint.
Infuse, stirring occasionally till cold, t	then strain.
Pulvis Acidi Tannici et Boracis.	
Tannic Acid	3 grains.
Borax	
1-2 grains for Laryngeal Insuffla	
Pulvis Acidi Tannici et Morphinæ.	
Tannic Acid	3 grains.
Hydrochloride of Morphine	
Borax	
1-2 grains for Laryngeal Insuffla	
Pulvis Iodoformi et Morphinæ.	
Hydrochloride of Morphine	3 grains.
Iodoform	
Borax	to 2 drachms.
1-2 grains for Laryngeal Insuffla	tion.
Dulyia Marphine at Paragia	
Pulvis Morphinæ et Boracis.	
Hydrochloride of Morphine	10 grains.

Pulvis Hydrargyri S	ubchl	oridi	et Z	inci.
Subchloride of Mercu	ary			1 part.
Oxide of Zinc				2 parts.
For	extern	al use.		
Pulvis Iodoformi et	Acidi	i Bori	ci.	
Iodoform				1 part.
Boric Acid				
	extern			
Pulvis Stramonii Con	mposi	tus.		
Stramonium Leaf, po	wdere	d		1 ounce.
Anise Fruit				$\frac{1}{2}$ ounce.
Tobacco Leaf				30 grains.
Nitrate of Potassium				$\frac{1}{2}$ ounce.
From 10 to 30 grains to	be bus	rnt and	the s	moke inhaled.
Sapo Liquidus.				
Soft Soap				2 ounces.
Methylated Spirit				
spiritus Anethi.				
Oil of Dill				1 fl. ounce.
Rectified Spirit				to 50 fl. oz.
piritus Anisi (St.B.)	P.).			
Oil of Anise				1 fl. ounce.
Rectified Spirit				to 50 fl. oz.
This is \frac{1}{5} the str	ength o	of the 1	B.P. S	Spirit.
piritus Carui.				
Oil of Caraway				1 fl. ounce.
TO 7 CI				to 50 fl. oz.

Spiritus Caryophylli. Oil of Cloves 1 fl. ounce. Rectified Spirit to 50 fl. oz.
Spiritus Cassiæ Ligneæ. Oil of Cassia 1 fl. ounce. Rectified Spirit to 50 fl. oz.
Suppositorium Acidi Tannici et Belladonnæ.Tannic Acid 3 grains.Green Extract of Belladonna ½ grain.Oil of Theobroma to 15 grains.
Suppositorium Acidi Tannici Belladonnæ et Opi Tannic Acid 5 grains. Green Extract of Belladonna ½ grain. Opium 1 grain.
Oil of Theobroma to 15 grains. Suppositorium Belladonnæ (St.B.P.). (Three Strengths). Green Extract of Belladonna ½,1, or 2 grains. Oil of Theobroma to 15 grains.
Suppositorium Belladonnæ et Morphinæ. Green Extract of Belladonna ½ grain. Hydrochloride of Morphine ¼ grain. Oil of Theobroma to 15 grains.
Syrupus Papaveris (Syrup of Poppy Capsules). Poppy Capsules, bruised and freed from seeds 2½ pounds. Treacle 2 pounds. Distilled Water at 170° F 3 gallons. Macerate for 12 hours, draw off and express the liquor,

S	yrupus Rhœado	os (S	St.B.P	.) (Syr	up of	Red Poppy).
	Dried Red Popp	y Pe	etals			13 ounces.
	Treacle					
	Boiling Distille					
	Infuse the petals					
	express to					
T	inctura Myrrh	æ (S	t.B.P.).		
	Myrrh in coarse	e pov	vder			4 ounces.
	Methylated Spi	-				1 pint.
	1		L			1
U	Inguentum Aci	di T	annic	i.		
	Tannic Acid					1 drachm.
	*Mixed Paraffin					to 1 ounce.
			/61	D D)		
PU	Inguentum Atr	opin	æ (St.	B. P.).		
	Atropine					1 grain.
	Oleic Acid					20 grains.
ı	Soft Paraffin					to 1 ounce.
	This is about	the	strengt	h of the	B.P.	Ointment.
U	Inguentum Atr	opin	æ et	Acidi	Bori	ci.
ı	Atropine					½ grain.
1	Boric Acid					60 grains.
1	Soft Paraffin					to 1 ounce.
t	Jnguentum Atr	opin	æ et	Cocain	ıæ.	
1	Atropine					½ grain.
1	Classia					2½ grains.
1	Oleic Acid					20 grains.
1	Soft Paraffin					
1						
1	* Soft Paraffin, with					ard Paramn to be

f fit consistency, the amount varying with time of year.

Unguentum A	tropina	e et	Hydrai	rgyri	Oxidi Flavi
Atropine					$\frac{1}{2}$ grain.
Yellow Oxide	e of Merc	cury			4 grains.
Oleic Acid			* * *		20 grains.
Soft Paraffin					to 1 ounce.
Unguentum A	Atropina	e et	Iodofo	rmi.	
Atropine					1 grain.
Iodoform					60 grains.
Oleic Acid					20 grains.
Soft Paraffin	1				to 1 ounce.
Unguentum (Cocainæ	(St	B.P.).		
Cocaine					5 grains.
Oleic Acid					20 grains.
Soft Paraffin	1				to 1 ounce.
This is less t	han $\frac{1}{3}$ the	stre	ngth of th	ie B.I	P. Ointment.
Unguentum (Cocainæ	Cor	npositu	m.	
Cocaine					
Hydrate of	Chloral				60 grains.
Menthol					60 grains.
Soft Paraffin	n				to 1 ounce.
Unguentum (Cocainæ	et	Hydrai	gyri	Oxidi Flavi
Cocaine					
Yellow Oxid	le of Mer	cury			4 grains.
Oleic Acid					20 grains.
Soft Paraffin	n				to 1 ounce.
Unguentum					
Prepared Control Prepared La			٠		equal parts.

U	nguentum Diachyli. Lead Plaster Mixed Paraffin equal parts.
U	nguentum Ferri Sulphatis. Sulphate of Iron, in fine powder 10 grains. Prepared Lard to 1 ounce.
U	nguentum Hydrargyri Ammoniati Compositum. Ammoniated Mercury 10 grains. Solution of Coal Tar 1 fl. drachm. Mixed Paraffin to 1 ounce.
U	nguentum Hydrargyri Nitratis Compositum. Carbolic Acid 60 grains. Ointment of Nitrate of Mercury ½ ounce. Ointment of Sulphur to 2 ounces.
U	nguentum Hydrargyri Oleatis (St.B.P.). (5, 10, or 15 p.c.)
	Oleate of Mercury 1, 2, or 3 parts. Benzoated Lard to 20 parts.
U	nguentum Hydrargyri Oxidi Flavi (St.B.P.). (Two Strengths.)
	Yellow Oxide of Mercury 2 or 4 grains. Mixed Paraffin to 1 ounce. About $\frac{1}{4}-\frac{1}{2}$ the strength of the B.P. Ointment.
	nguentum Hydrargyri Oxidi Rubri dilutum.

Unguentum Hydrargyri et Plumb Acetate of Lead Subchloride of Mercury Oxide of Zinc Ointment of Nitrate of Mercury Soft Paraffin		10 grains. 10 grains. 20 grains. 20 grains.
Unguentum Hydrargyri Sulphidi (Two Strengths.)	Ru	ıbri.
Red Sulphide of Mercury of each Creosote Prepared Lard		2 minims.
Unguentum Physostigminæ. Sulphate of Physostigmine Soft Paraffin		
Unguentum Picis cum Plumbo. Ointment of Tar Ointment of Acetate of Lead		equal parts.
Unguentum Simplex. Yellow Wax Prepared Lard		1 part. 3 parts.
Unguentum Styracis. Prepared Storax, liquefied Prepared Lard		½ ounce. 1 ounce.

τ	Inguentum Sulphuris Composit	um.	
	Sublimed Sulphur		30 grains.
	Red Sulphide of Mercury		10 grains.
	Ammoniated Mercury		10 grains.
	Olive Oil		2 fl. drachms.
	Creosote		4 minims.
	Prepared Lard		1 ounce.
τ	Inguentum Zinci Carbolatum.		
	Carbolic Acid, liquefied		1 fl. drachm.
	Ointment of Zinc		1 ounce.
7	Zanor Asidi Carbalisi		
'	apor Acidi Carbolici.		20 : :
	Carbolic Acid, liquefied		20 minims.
	Water at 150° F		1 pint.
7	Zapor Benzoini.		
	Compound Tincture of Benzoin		1 fl. drachm.
	Water at 150° F		
	Water at 150 F		i pint.
7	7apor Camphoræ.		
	Spirit of Camphor		15 minims.
	Alcohol, 60 per cent		to 1 fl. drachm.
	Water at 150° F		1 pint.
7	7apor Creosoti.		
			10 minima
	Creosote Light Carbonate of Magnesium (Diffused in) Water		4 grains
	(Diffused in) Water		to 1 fl. dwo above
	TIT T MOD IS		
	Water at 150° F.		1 pint.

Vapor Pini.

(Oil of Pine			 5 minims.
Oil of Pine Light Carbonate of (Diffused in) Water	Magn	esium	 4 grains.
(Diffused in) Water			 to 1 fl. drachm.
Water at 150° F.			 1 pint.

Vapor Terebenæ.

(Terebene, pure			 5 minims.
Light Carbonate of	Magn	esium	 4 grains.
(Diffused in) Water			 1 fl. drachm.
Water at 150° F.			 1 pint.

SCALE OF DOSES AT DIFFERENT AGES.

Half the dose for an adult may be considered suitable for a child of about 12, half of this for a child of 4, half of this again for a child of 2, half of this again for a child 9 months to 1 year.

Thus-

AGE
 Adult
 12
 4
 2
 9 m. to 1 yr.

 DOSE
 1

$$\frac{1}{2}$$
 $\frac{1}{8}$
 $\frac{1}{10}$

For intermediate ages, when the dose for an adult is a draught of 1 fluid ounce, the following may be a sufficient guide:—

AGE.				D	OSE.
9 m. to 1	year		 	½ fl.	drachm.
2	years	 	 	1	,,
3	"	 	 	$1\frac{1}{2}$	11
4	11	 	 	2	,.
8	"	 	 	3	. 11
12	11	 	 	4	11
17	11	 	 	6	11
Ac	lult		 	1 fl.	ounce.

POSOLOGICAL TABLE

The names of substances and preparations not contained in the British Pharmacopæia, 1898, are printed in italics; doses not therein given are printed in clarendon type.

Acetani	ilidum (Antifebr	in)			1 to 3 gr.
	Ipecacuanhæ				10 ,, 30 min.
Acetum	Scillæ				
Acidum	Aceticum				5,, 15,,
_	— dilut	um			½ ,,. 2 fl. dr.
_	Arseniosum				$\frac{1}{60}$,, $\frac{1}{15}$ gr.
-	Benzoicum				5 ,, 15 ,,
_	Boricum				
_	Carbolicum				
	— liq	uefacti	ım		
	Citricum				
	Gallicum				
_	Hydrobromicun				
	Hydrochloricun				
_	Hydrocyanicum			-	
_	Nitricum dilutu				
_	Nitro-hydrochle		*		5 ,, 20 ,,
_	Phosphoricum of				
	Salicylicum		-		5 ,, 20 gr.
	Sulphuricum Ar				5 ,, 20 min.
	Sulphuricum di			_	5 ,, 20 ,,
_	Sulphurosum (5				30 ,, 60 ,,
	Tannicum				
_	Tartaricum				

Æther			 10 to 30 min.
(Single Dose)			 40 ,, 60 ,,
- Aceticus			 20 ,, 40 ,,
(Single dose)			 60 ,, 90 ,,
Aloe Barbadensis			 2 ,, 5 gr.
— Socotrina			 2 ,, 5 ,,
Aloinum			 $\frac{1}{2}$,, 2 ,,
Alumen			 5 ,, 10 ,,
Ammoniacum			 5 ,, 15 ,,
Ammonii Benzoas			 5 ., 15 ,,
— Bromidum			 5 ,, 30 ,,
- Carbonas			 3 ,, 10 ,,
— Chloridum			 5 ,, 20 ,,
— Iodidum			 3 ,, 20 ,,
- Phosphas			 5 ,, 20 ,,
— Salicylas			 3 ,, 20 ,,
Amyl Nitris			 $\frac{1}{2}$,, 1 min.
— For Inhalatio			 2 ,, 5 ,,
Antimonii Oxidum			 1 ,, 2 gr.
Antimonium Sulphuratur	n		 1 ,, 2 ,,
- Tartaratum			
		essant	1 1
	(Emet		 1,, 2,,
Apomorphinæ Hydrochlo	ridum		 $\frac{1}{10}$,, $\frac{1}{4}$,,
Agna Anothi			 $\frac{1}{2}$,, 2 fl. oz.
— Anisi			 1 7
- Aurantii Flores			 1 0
Camphoræ			 1 0
— Carui			 1 0
- Chloroformi			 1 0
- Cinnamomi			 1 1
- Fæniculi			 1 1
- Laurocerasi			
			 $\frac{1}{2}$,, 1 fl. dr.

Aqua Menthæ Piperitæ		$\frac{1}{2}$ to 2 fl. oz.
— Menthæ Viridis		$\frac{1}{2}$,, 1 ,,
— Pimentæ		$\frac{1}{2}$,, 2 ,,
Rosæ		½ ,, 2 ,,
— Sambuci		$\frac{1}{2}$,, 2 ,,
Argenti Nitras		$\frac{1}{4}$,, $\frac{1}{2}$ gr.
— Oxidum		$\frac{1}{2}$,, 2 ,,
Arsenii Iodidum		$\frac{1}{20}$,, $\frac{1}{5}$,,
Asafetida		5 ., 15 ,,
Atropina		$\frac{1}{200}$,, $\frac{1}{100}$,,
Atropinæ Sulphas		$\frac{1}{200}$,, $\frac{1}{100}$,,
Balsamum Peruvianum		5 ,, 15 min.
— Tolutanum		5 ,, 15 gr.
Benzol		5,, 10 min.
Bismuthi Benzoas		2 ,, 20 gr.
— Carbonas		5 ,, 20 ,,
— Oxidum		5 ,, 20 ,,
— Salicylas		5 ,, 20 ,,
— Subnitras		5 ,, 20 ,,
Borax (Sodæ Biboras)		5 ,, 20 ,,
Bromal Hydras		2 ,, 5 ,,
Bromoform		$\frac{1}{2}$,, 2 min.
· Bromol (Tribromophenal)		$\frac{1}{2}$,, 2 gr.
Butyl-Chloral Hydras		5 ,, 20 ,,
Caffeina		1 ,, 5 ,,
Caffeinæ Citras		2 ,, 10 ,,
— Effervescens		60 ,, 120 ,,
Calcii Carbonas Præcipitatus	·	10 ,, 60 ,,
— Chloridum		5 ,, 15 ,,
— Hypophosphis		3 ,, 10 ,,
- Phosphas		5 ,, 15 ,,
— Salicylas		2 ,, 20 ,,
Calx Sulphurata	***	‡ ,, 1 ,,

Cambogia			 	$\frac{1}{2}$ to	2 gr.
Camphora			 	2 ,,	5 ,,
Carbo Ligni			 	60 ,,	120 .,
Catachy			 	5 ,,	15 ,,
Cerii Oxalas			 	2 ,,	10 ,,
Chloral Hydras			 	5 ,,	20 ,,
Chloralimidum .			 	20 ,,	40 ,,
Chloroformum .			 	1 ,,	5 min.
Cocaina			 4.4	$\frac{1}{20}$,,	1 gr.
Cocainæ Hydrochlo	ride		 	1 ,,	
Codeina				1 ,,	2 ,,
Codeinæ Phosphas				$\frac{1}{4}$,,	2 ,,
011110			 	2 ,,	5 ,,
Confectio Piperis .			 		120 ,,
— Rosæ Gal	licæ		 		
- Sennæ .			 	60 ,,	120 ,,
- Sulphuris	5		 	60 ,,	120 ,,
Conii Folia			 	2 ,,	8 ,,
Copaiba			 	30 ,,	60 nim.
Creosotum			 	1 ,,	5 ,,
Creta Præparata .			 	10 ,,	60 gr.
Cubebæ Fructus .			 	30 ,,	60 ,,
Cupri Sulphas (Astr	ingent	5)	 	1 ,,	2 ,,
— — (Eme					
Cusso			 	1 ,,	$\frac{1}{2}$ oz.
Decoctum Aloes Co					2 fl. oz.
- Granati (Cortici	S	 	$\frac{1}{2}$,,	1 ,,
- Hæmator	kyli		 	1/2 ,,	2 ,,
Digitalis Folia	**		 	$\frac{1}{2}$,,	2 gr.
Elaterinum			 	1 ,,	1 ,,
Elaterium			 	1 10 ,,	$\frac{1}{2}$,,
Ergota				20 ,,	60 ,,
Ethyl-Carbamate (U:	rethane	2)	 	10 ,,	60 ,,
Eucalypti Gummi.			 	2 ,,	5 ,,

Extractum	Belladonnæ A	Alcoho	licum	(Al-			
	kaloids 1	p.c.)			1 to	1	gr.
_	Belladonnæ					1	,,
_	Cannabis Inc	dicæ			1,,	1	,,
	Colchici					1	
_	Nucis Vomi	icæ (S	trych	nine			
	5 p.c.)				1,,	1	,,
	Opii (Morphi	ne 20 p	o.c.)		1,,	1	, ,
	Physostigma					1	,,
_	Stramonii					1	,,
-	Strophanthi					1	,,
Extractum	Aloes Barbad	lensis			1 ,,	4	,,
	Anthemidis				2 ,,		
	Cascaræ Sag	radæ					
	Colocynthidi	s Com	positu	m	2 ,,		
_	Convallariæ				2 ,,	8	,,
_	Ergotæ				2 ,,	8	,,
_	Euonymi Sic	cum			1 ,,	2	,,
	Gentianæ				2 ,,	8	,,
	Glycyrrhizæ				5 ,,	60	
_	Hyoscyami V	Viride .			2 ,,	8	,,
_	Jalapæ				2 ,,	-	
_	Krameriæ				5 ,,	15	,,
	Malti						
	Rhei				2 ,,		
	Taraxaci				5 ,,	15	,,
	Viburni Prui	rifolii			2 ,,	10	"
Extractum	Belladonnæ l	Liquid	um (A	lka-			
	loids 3 p.	.c.)			$\frac{1}{2}$,,	2	min.
_	lpecacuanhæ						
	loids $2-2\frac{1}{4}$ p.	15.0	-			-	,,
-	_	(En	netic)		15 ,,	20	,,

Ex	tractum			micæ nine 1½	-		1 to	3	min.
Ex	tractum	Cimic	fugæ	Liquid	um		5 ,,	30	,,
	_	Cinch	onæ]	Liquidu	m (A	lka-			
		lo	ids 5 p	o.c.)			5 ,,	15	,,
	-	Ergota	e Liqu	nidum			10 ,,	30	, ,
		Hama	melidi	s Liqui	dum		5 ,,	15	,,
	_	Hydra	stis L	iquidun	n		5 ,,	15	,,
	-	Jabora	andi L	iquidur	n		5 ,,	15	,,
		Opii	Liqui	dum	(Morp	hin:			
		34	p.c.)				5 ,,	30	,,
Ex	tractum	Coce	Liquid	lum			$\frac{1}{2}$,,	1 f	d.dr.
	_	Filicis	Liqui	idum			$\frac{3}{4}$,,	$1\frac{1}{2}$,,
	_	Glycy	rrhizæ	Liquid	um		$\frac{1}{2}$,,	1	,,
	-	Pareir	æ Liq	uidum			$\frac{1}{2}$,,	2	, ,
	_	Sarsæ	Liqui	dum			2 ,,	4	, ,
	_	Tarax	aci Lie	quidum			$\frac{1}{2}$,,	2	,,
fe	l Bovinu	m Pur	ificatu	m			5 ,,	15	gr.
Fe	rri Arsei	nas					16 :	1	,,
-	- Carb	onas Sa	acchar	atus			10 ,,	30	,,
-	et Ar	nmonii	Citra	s			5 ,,	10	,,
-	et Qu	ininæ	Citras				5 ,,	10	,,
-	- Iodia	lum					1 ,,	5	,,
-	- Phos	phas					5 ,,	10	,,
-	_			***			1 .,	5	,,
-				ıs			$\frac{1}{2}$,,	3	,,
fe	rrum Re						1 ,,	5	,,
	— Ta	rtaratu	m				5 ,,	10	3.2

Glusidum (Saccharin) \frac{1}{2} , \text{, 2 m} Glycerinum 1 , 2 fl.dr. Glycerinum Acidi Borici (30 p.c.) 15 , 30 min. — Carbolici (1 in 5) 5 , 10 , 10 , 20 , 10 , 10 , 20 , 10 , 10	Galban	um			 5 to	15 gr.	
Glycerinum Acidi Borici (30 p.c.) 15,, 30 min. — — Carbolici (1 in 5) 5,, 10, — — Tannici (1 in 5) 10,, 20, — Aluminis (1 in 6) \frac{1}{2},, 1fl.dr. — Boracis (1 in 7) \frac{1}{2},, 1, — Pepsini 1,, 2, Guaiaci Resina 5,, 15 gr. Guaiacol 1,, 5 min. Guarana 10,, 60 gr. Homatropinæ Hydrobromidum \frac{1}{30}, \frac{1}{10}, — Perchloridum \frac{1}{32}, \frac{1}{16}, — Perchloridum \frac{1}{32}, \frac{1}{10}, — Subchloridum \frac{1}{2}, \frac{5}{10}, Hydrargyrum cum Creta 1,, 5,, Hyoscinæ Hydrobromidum \frac{1}{200}, \frac{1}{100}, Hyoscyaminæ Sulphas \frac{1}{200}, \frac{1}{100}, Infusum Aurantii \frac{1}{2}, 1 fl. oz. — Aurantii Compositum \frac{1}{2}, 1 fl. oz. — Aurantii Compositum \frac{1}{2}, 1 fl. oz. — Calumbæ \frac{1}{2}, 1 fl. oz. — Caryophylli \frac{1}{2}, 1 fl. oz. — Chiratæ \frac{1}{2}, 1 fl. oz. — Cinchonæ Acidum \frac{1}{2}, 1 fl. oz. — Ergotæ 1,, 2							
Glycerinum Acidi Borici (30 p.c.) 15 ,, 30 min.							r.
— — Carbolici (1 in 5) 5 ,, 10 ,, 20 ,	-						
— Aluminis (1 in 6) 1/2 , 1 ft. dr. — Boracis (1 in 7) 1/2 , 1 ft. dr. — Pepsini 1 , 2 , 1 — Pepsini 1 , 2 , 1 Guaiaci Resina 5 , 15 gr. Guaiacol 1 , 5 min. Guarana 10 , 60 gr. Homatropinæ Hydrobromidum 1/3 0 , 1/2 0 , 10 Hydrargyri Iodidum Rubrum 1/3 2 , 1/1 0 , 10 — Perchloridum 1/3 2 , 1/1 0 , 10 — Subchloridum 1/3 2 , 1/1 0 , 10 — Subchloridum 1/3 2 , 1/1 0 , 10 Hydrargyrum cum Creta 1 , 5 , 10 Hydrargyrum cum Creta 1 , 5 , 10 Hyoscinæ Hydrobromidum 1/2 0 , 1/10 0 , 10 Hyoscyaminæ Sulphas 1/2 0 , 1/10 0 , 10 Hyoscyaminæ Sulphas 1/2 0 , 1/10 0 , 10 Infusum Aurantii 1/2 1 , 1 ft. oz. — Aurantii Compositum 1/2 1 , 1 , 1 — Buchu 1 , 2 , 1 , 1 — Calumbæ 1/2 1 , 1 , 1 — Cascarillæ 1/2 1 , 1 , 1 — Chiratæ 1/2 1 , 1 , 1 — Cinchonæ Acidum 1/2 1 , 1 , 1 — Cuspariæ 1 , 2 , 2 , 1 — Ergo	_						-
— Aluminis (1 in 6) 1/2 ,, 1 fl. dr. — Boracis (1 in 7) 1/2 ,, 1 ,, 1 ,, 2 ,, 1 — Pepsini 1 ,, 2 ,, 2 ,, 1 Guaiaci Resina 5 ,, 15 gr. Guaiacol 1 ,, 5 min. Guarana 10 ,, 60 gr. Homatropinæ Hydrobromidum 1/3 0 ,, 1/2 0 ,, 10	_						
— Boracis (1 in 7) 1/2 , 1 , 1 , 2 , 3 — Pepsini 1 , 2 , 2 , 3 Guaiaci Resina 5 , 15 gr. Guaiacol 1 , 5 min. Guarana 10 , 60 gr. Homatropinæ Hydrobromidum 1/3 0 , 1/2 0 , 1	_						r.
— Pepsini 1, 2, 1 Guaiaci Resina 5, 15 gr. Guaiacol 1, 5 min. Guarana 10, 60 gr. Homatropinæ Hydrobromidum 1 1, 5 min. Hydrargyri Iodidum Rubrum 1 1, 5 min. — Perchloridum 1 1, 5 min. — Perchloridum 1 1, 5 min. — Perchloridum 1 1, 5 min. — Subchloridum 1 1, 5 min. — Hydrargyrum cum Creta 1 1, 5 min. — Hydrargyrum cum Creta 1 1, 5 min. — Hydrargyrum cum Creta 1 1, 5 min. — Aurantii Compositum 1 1, 5 min. — Buchu 1 1, 5 min. — Calumba 1 1, 2 min. — Calumba 1 1, 2 min. — Caryophylli 1 1, 1 min. — Cinchona 1	_					_	
Guaiaci Resina 5 , 15 gr. Guaiacol 1 , 5 min. Guarana 10 , 60 gr. Homatropinæ Hydrobromidum \[\frac{1}{80} \] \[\frac{1}{20} \] \[\]							
Guarana 1, 5 min. Guarana 10, 60 gr. Homatropinæ Hydrobromidum $\frac{1}{80}$, $\frac{1}{20}$, Hydrargyri Iodidum Rubrum $\frac{1}{32}$, $\frac{1}{16}$, — Perchloridum $\frac{1}{32}$, $\frac{1}{16}$, — Subchloridum $\frac{1}{2}$, 5 , Hydrargyrum cum Creta 1 ,, 5 , Hyoscinæ Hydrobromidum $\frac{1}{200}$, $\frac{1}{100}$, Hyoscyaminæ Sulphas $\frac{1}{200}$, $\frac{1}{100}$, Infusum Aurantii $\frac{1}{2}$, 1 fl. oz. — Aurantii Compositum $\frac{1}{2}$, 1 fl. oz. — Buchu $\frac{1}{2}$, 1 m — Calumbæ $\frac{1}{2}$, 1 m — Caryophylli $\frac{1}{2}$, 1 m — Cascarillæ $\frac{1}{2}$, 1 m — Cinchonæ Acidum $\frac{1}{2}$, 1 m — Cuspariæ $\frac{1}{2}$, 1 m — Digitalis $\frac{1}{4}$, $\frac{1}{2}$, $\frac{1}{2}$, — Ergotæ $\frac{1}{4}$, $\frac{1}{2}$, $\frac{1}{2}$, — Gentianæ Compositum $\frac{1}{2}$, 1 m — Krameriæ $\frac{1}{2}$, 1 m	Guaiaci						
Guarana 10 , 60 gr. Homatropinæ Hydrobromidum $\frac{1}{80}$, $\frac{1}{20}$, $\frac{1}{10}$, $\frac{1}{100}$, $$							1.
Homatropinæ Hydrobromidum $\frac{1}{80}$,, $\frac{1}{20}$,, Hydrargyri Iodidum Rubrum $\frac{1}{32}$,, $\frac{1}{16}$,, — Perchloridum $\frac{1}{32}$,, $\frac{1}{16}$,, — Subchloridum $\frac{1}{2}$,, $\frac{5}{2}$,, Hydrargyrum cum Creta 1 ,, $\frac{5}{2}$,, Hyoscinæ Hydrobromidum $\frac{1}{200}$,, $\frac{1}{100}$,, Hyoscyaminæ Sulphas $\frac{1}{200}$,, $\frac{1}{100}$,, Infusum Aurantii $\frac{1}{2}$,, 1 fl. oz. — Aurantii Compositum $\frac{1}{2}$,, 1 ,, — Buchu $\frac{1}{2}$,, 1 ,, — Calumbæ $\frac{1}{2}$,, 1 ,, — Caryophylli $\frac{1}{2}$,, 1 ,, — Cascarillæ $\frac{1}{2}$,, 1 ,, — Chiratæ $\frac{1}{2}$,, 1 ,, — Cinchonæ Acidum $\frac{1}{2}$,, 1 ,, — Digitalis $\frac{1}{4}$,, $\frac{1}{2}$,, — Ergotæ $\frac{1}{4}$,, $\frac{1}{2}$,, — Gentianæ Compositum $\frac{1}{2}$,, 1 ,, — Krameriæ $\frac{1}{2}$,, 1 ,,	Guaran	a					
Hydrargyri Iodidum Rubrum $\frac{1}{32}$, $\frac{1}{16}$, $\frac{1}{100}$, $\frac{1}{100}$, $\frac{1}{100}$, $\frac{1}{100}$, $\frac{1}{100}$, $\frac{1}{100}$, 1	Homati	opinæ Hydrobro	midun	a	 1199000		
— Subchloridum 1/2 , 5 , 5 , . Hydrargyrum cum Creta 1 , 5 , . Hyoscinæ Hydrobromidum 1/2 00 , 1/100 , . Hyoscyaminæ Sulphas 1/2 00 , 1/100 , . Hyoscyaminæ Sulphas 1/2 00 , 1/100 , . Infusum Aurantii 1/2 , 1 fl. oz. — Aurantii Compositum 1/2 , 1 , . — Buchu 1 , · 2 , . — Calumbæ 1/2 , 1 , . — Caryophylli 1/2 , 1 , . — Cascarillæ 1/2 , 1 , . — Chiratæ 1/2 , 1 , . — Cinchonæ Acidum 1/2 , 1 , . — Cuspariæ 1 , , 2 , . — Digitalis 1 , , 2 , . — Ergotæ 1 , , 2 , . — Gentianæ Compositum 1/2 , 1 , . — Krameriæ 1/2 , 1 , .	Hydrar	gyri Iodidum Ru	brum		 The state of the s		
Hydrargyrum cum Creta 1 ,, 5 ,, Hyoseinæ Hydrobromidum $\frac{1}{200}$, $\frac{1}{100}$,, Hyoseyaminæ Sulphas $\frac{1}{200}$, $\frac{1}{100}$,, Infusum Aurantii $\frac{1}{2}$, 1 fl. oz. — Aurantii Compositum $\frac{1}{2}$, 1 ,, — Buchu 1 ,, 2 ,, — Calumbæ $\frac{1}{2}$,, 1 ,, — Caryophylli $\frac{1}{2}$,, 1 ,, — Cascarillæ $\frac{1}{2}$,, 1 ,, — Chiratæ $\frac{1}{2}$,, 1 ,, — Cinchonæ Acidum $\frac{1}{2}$,, 1 ,, — Cuspariæ 1 ,, 2 ,, — Digitalis $\frac{1}{4}$,, $\frac{1}{2}$,, — Ergotæ 1 ,, 2 ,, — Gentianæ Compositum $\frac{1}{2}$,, 1 ,, — Krameriæ $\frac{1}{2}$,, 1 ,,		Perchloridu	m		 $\frac{1}{32}$,,	1 16 ,,	
Hyoscinæ Hydrobromidum $\frac{1}{200}$, $\frac{1}{100}$, Hyoscyaminæ Sulphas $\frac{1}{200}$, $\frac{1}{100}$, Infusum Aurantii $\frac{1}{2}$, 1 fl. oz. — Aurantii Compositum $\frac{1}{2}$, 1 , 1 , — Buchu 1 ,, 2 , — Calumbæ $\frac{1}{2}$, 1 , 1 , — Caryophylli $\frac{1}{2}$, 1 , 1 , — Cascarillæ $\frac{1}{2}$, 1 , 1 , — Chiratæ $\frac{1}{2}$, 1 , 1 , — Cinchonæ Acidum $\frac{1}{2}$, 1 , 1 , — Digitalis $\frac{1}{4}$, $\frac{1}{2}$, $\frac{1}{2}$, — Ergotæ 1 ,, 2 , — Gentianæ Compositum $\frac{1}{2}$, 1 , 1 , — Krameriæ $\frac{1}{2}$, 1 , 1 ,	_	Subchloridu	m		 $\frac{1}{2}$,,	5 ,,	
Hyoscinæ Hydrobromidum $\frac{1}{200}$, $\frac{1}{100}$, Hyoscyaminæ Sulphas $\frac{1}{200}$, $\frac{1}{100}$, Infusum Aurantii $\frac{1}{2}$, 1 fl. oz. — Aurantii Compositum $\frac{1}{2}$, 1 , 1 , — Buchu 1 ,, 2 , — Calumbæ $\frac{1}{2}$, 1 , 1 , — Caryophylli $\frac{1}{2}$, 1 , 1 , — Cascarillæ $\frac{1}{2}$, 1 , 1 , — Chiratæ $\frac{1}{2}$, 1 , 1 , — Cinchonæ Acidum $\frac{1}{2}$, 1 , 1 , — Digitalis $\frac{1}{4}$, $\frac{1}{2}$, $\frac{1}{2}$, — Ergotæ 1 ,, 2 , — Gentianæ Compositum $\frac{1}{2}$, 1 , 1 , — Krameriæ $\frac{1}{2}$, 1 , 1 ,	Hydrar	gyrum cum Creta	a		1 ,,	5 ,,	
Infusum Aurantii $\frac{1}{2}$, 1 fl. oz. — Aurantii Compositum $\frac{1}{2}$, 1 , 1 ,, — Buchu 1 , 2 ,, — Calumbæ $\frac{1}{2}$, 1 ,, — Caryophylli $\frac{1}{2}$, 1 ,, — Cascarillæ $\frac{1}{2}$, 1 ,, — Chiratæ $\frac{1}{2}$, 1 ,, — Cinchonæ Acidum $\frac{1}{2}$, 1 ,, — Cuspariæ 1 , 2 ,, — Digitalis $\frac{1}{4}$, $\frac{1}{2}$,, — Ergotæ 1 , 2 ,, — Gentianæ Compositum $\frac{1}{2}$, 1 ,, — Krameriæ $\frac{1}{2}$, 1 ,,	Hyoscin	næ Hydrobromid	um				
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	Hyoscy	aminæ Sulphas			 1 200 ,,	1 100 ,,	
— Buchu 1,, 2, 3 — Calumbæ ½, 1 — Caryophylli ½, 1 — Cascarillæ ½, 1 — Chiratæ ½, 1 — Cinchonæ Acidum ½, 1 — Cuspariæ 1, 2 — Digitalis ½, ½ — Ergotæ 1, 2 — Gentianæ Compositum ½, 1 — Krameriæ ½, 1	Infusun	a Aurantii			 $\frac{1}{2}$,,	1 fl. oz	
— Calumbæ $\frac{1}{2}$,, 1 ,, — Caryophylli $\frac{1}{2}$,, 1 ,, — Cascarillæ $\frac{1}{2}$,, 1 ,, — Chiratæ $\frac{1}{2}$,, 1 ,, — Cinchonæ Acidum $\frac{1}{2}$,, 1 ,, — Cuspariæ $\frac{1}{2}$,, 1 ,, — Digitalis $\frac{1}{4}$,, $\frac{1}{2}$,, — Ergotæ $\frac{1}{4}$,, $\frac{1}{2}$,, — Gentianæ Compositum $\frac{1}{2}$,, 1 ,, — Krameriæ $\frac{1}{2}$,, 1 ,,	_	Aurantii Compo	ositum		 $\frac{1}{2}$,,	1 ,,	
— Caryophylli $\frac{1}{2}$,, 1 — Cascarillæ $\frac{1}{2}$,, 1 — Chiratæ $\frac{1}{2}$,, 1 — Cinchonæ Acidum $\frac{1}{2}$, 1 — Cuspariæ 1,, 2 — Digitalis $\frac{1}{4}$,, $\frac{1}{2}$ — Ergotæ $\frac{1}{2}$,, 1 — Gentianæ Compositum $\frac{1}{2}$,, 1 — Krameriæ $\frac{1}{2}$,, 1	_	Buchu			 1 ,,	2 ,,	
— Cascarillæ $\frac{1}{2}$,, 1 ,, — Chiratæ $\frac{1}{2}$,, 1 ,, — Cinchonæ Acidum $\frac{1}{2}$,, 1 ,, — Cuspariæ 1 ,, 2 ,, — Digitalis $\frac{1}{4}$,, $\frac{1}{2}$,, — Ergotæ 1 ,, 2 ,, — Gentianæ Compositum $\frac{1}{2}$,, 1 ,, — Krameriæ $\frac{1}{2}$,, 1 ,,	_	Calumbæ			 $\frac{1}{2}$,,	1 ,,	
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	_	Caryophylli			 $\frac{1}{2}$,,	1 ,,	
— Cinchonæ Acidum $\frac{1}{2}$, 1 ,, 2 ,, — Cuspariæ 1 ,, 2 ,, — Digitalis $\frac{1}{4}$, $\frac{1}{2}$ — Ergotæ 1 ,, 2 ,, — Gentianæ Compositum $\frac{1}{2}$ 1 — Krameriæ $\frac{1}{2}$., 1	-	Cascarillæ			 $\frac{1}{2}$,,	1 ,,	
— Cuspariæ $\frac{1}{4}$,, $\frac{1}{2}$,, — Ergotæ $\frac{1}{2}$,, $\frac{1}{2}$,, — Gentianæ Compositum $\frac{1}{2}$,, $\frac{1}{2}$,, — Krameriæ $\frac{1}{2}$,, $\frac{1}{2}$,,	_	Chiratæ			 $\frac{1}{2}$,,	1 ,,	
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	_	Cinchonæ Acid	um		 $\frac{1}{2}$,.	1 ,,	
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	_	Cuspariæ			 1 ,,	2 ,,	
— Gentianæ Compositum $\frac{1}{2}$,, $\frac{1}{2}$	_	Digitalis		* *//	 丰 ,,	$\frac{1}{2}$,,	
— Krameriæ $\frac{1}{2}$,, 1 ,,	_	Ergotæ			 1 ,,	2 ,,	
	_	Gentianæ Comp	positur	n	 $\frac{1}{2}$,,	1 ,,	
— Lupuli $\frac{1}{2}$,, 2 ,,		Krameriæ			 $\frac{1}{2}$,,	1 ,,	
	_	Lupuli		***	 $\frac{1}{2}$,,	2 ,,	

Infusum Quassiæ		$\frac{1}{2}$ to 1 fl. oz.
— Rhei		$\frac{1}{2}$,, 1 ,.
— Rosæ Acidum		1 1
— Scoparii		1 9
— Senegæ		1 1
— Sennæ		1 1
(Single dose)		1 0
- Serpentariæ		1 1
— Uvæ Ursi		1 1
odoformum		1 9
pecacuanhæ Radix (Expectora		1 0
— (Emetic)		15 20
Talapa		= 00
Talapæ Resina		0 ~
Kino		
Liquor Ammoniæ		
Dani!	***	0
- Ammonii Acetatis	***	1, - ,,
		The state of the s

	***	2 ,, 8 min.
Arsenici Hydrochloricus	s (1 p.c.)	2 ,, 8 ,,
- Arsenii et Hydrargyi	ri Iodidi	
(each 1 p.c.)		5 ,, 20 ,,
- Atropinæ Sulphatis (1 p	.c.)	$\frac{1}{2}$ 1
- Bismuthi et Ammonii C	itratis	½ 1 fl. dr.
- Calcis		1 4 fl. oz
— Saccharatus		20 60 min.
- Calumbæ Concentratus		1 fl. dr
- Chiratæ Concentratus		$\frac{1}{2}$,, 1 ,,
- Cuspariæ Concentratus		1 1
- Ethyl Nitritis		20, 60 min
- Ferri Acetatis		5 15
Perchloridi		5 ,, 15 ,,
		0 ,, 10 ,,

Liquor	Ferri Pernitratis				5 to	15	min	
_	Hydrargyri Percl	hloridi	(1 gr	. in				
	1 fl. oz.)				$\frac{1}{2}$,,	1	fl. d	r.
_	Hydrogenii Perox							
	Krameriæ Concen							
	Magnesii Carbona							
	Morphinæ Acetat							
	- Hydro	A STATE OF THE STA						
_	— Tartra							
_]	Pancreatis							r.
	Potassæ (6 per ce				1.5			
	Potassii Permang							
	Quassiæ Concentr		-				,,	
	Rhei Concentratu							
	Sarsæ Compositus							
	Senegæ Concentra							
	Sennæ Concentrat						,,	
	Serpentariæ Conc				_			
	Sodæ Chlorinatæ							
	Sodii Arsenatis (1				(7.75)		,,	
	Strychninæ Hydro	-					,,	
	Thyroidei		_					
	Frinitrini (1 p.c.)							
	arbonas							
	itras							
	_ Effervescens							
Lupulin	um				2 ,,	5	,,	
	ia Levis							
C	(Single dose)							
_	Ponderosa				5 ,,	30	,,	
	(Single dose)							
Magnesi	ii Carbonas Levis							
		ngle do						

	Magnesii Carbonas Por	nderosu	Q	5 to 20 cm
	integration caroonas ro		le dose)	 5 to 30 gr.
	- Sulphas			30 ,, 60 ,,
	(Single de			 30 ,,120 ,,
		ervesce		 $\frac{1}{4}$,, $\frac{1}{2}$ OZ.
	En			 60 ,.240 gr.
	Menthol		e dose)	 $\frac{1}{2}$,, 1 oz.
	Methylacetanilide (Exalg			 $\frac{1}{2}$,, 2 gr.
	Mistura Ammoniaci			 $\frac{1}{2}$,, $\frac{2}{2}$,,
				 $\frac{1}{2}$,, 1 fl. oz.
	- Amygdalæ			 2), - ,,
	— Creosoti			 $\frac{1}{2}$,, 1 $-$,,
	- Cretæ			 $\frac{1}{2}$., 1 ,,
	— Ferri Composi	ta		 $\frac{1}{2}$,, 1 ,,
	— Guaiaci			 $\frac{1}{2}$,, 1 ,,
				 1 ,, 2 ,,
	- Senna Compos			 1,, 2,,
	- Spiritus Vini (fallici		 1 ,, 2 ,,
	Morphinæ Acetas			 $\frac{1}{8}$,, $\frac{1}{2}$ gr.
	 Hydrochlori 	dum	* * *	 $\frac{1}{8}$,, $\frac{1}{2}$,,
	— Tartras		***	 $\frac{1}{8}$,, $\frac{1}{2}$,,
	Moschus			 5 ,, 10 ,,
	Naphthol			 3 ,, 10 ,,
	Nux Vomica			 1 ,, 4 ,,
(Oleum Anethi			 $\frac{1}{2}$,, 3 min.
	— Anisi			 $\frac{1}{2}$,, 3 ,,
	- Anthemidis			 $\frac{1}{2}$,, 3 ,,
	— Cajuputi			 $\frac{1}{2}$,, 3 ,,
	— Carui			 $\frac{1}{2}$,, 3 ,,
	— Caryophylli			 1, 3,
	— Cassice			 $\frac{1}{2}$,, 3 ,,
	— Cinnamomi			 $\frac{1}{2}$,, 3 ,,
	- Copaibæ			 5 ,, 20 ,,
	— Coriandri			 1 9
				2 ,, 5 ,,

Oleum	Crotonis					1/2	to	1	min.
	Cubebæ					5	,,	20	,,
_	Eucalypti					$\frac{1}{2}$,,	3	,,
	Juniperi					$\frac{1}{2}$,,	3	,,
_	Lavandulæ					1/2	,,	3	,,
	Limonis					$\frac{1}{2}$,,	3	,,
_	Menthæ Pip	peritæ				12	,,	3	,,
-	Viridis					$\frac{1}{2}$,,	3	,,
	Morrhuæ					1	,,	4	fl. dr.
	Myristicæ					$\frac{1}{2}$,,	3	min.
	Phosphorat	um				1	,,	5	,,
	Pimentæ					1/2	,,	3	,,
_	Ricini					1	,,	8	fl. dr.
_	Rosmarini					$\frac{1}{2}$, ,	3	min.
_	Santali					5	,,	30	,,
_	Terebinthin	æ				2	,,	10	,,
_	_	(Anth	elmint	ic)		3	,,	4	fl. dr.
Opium						12	,,	2	gr.
Oxyme	1					1	,,	2	fl. dr.
_	Scillæ					$\frac{1}{2}$,,	1	,,
Paralde	ehydum ·					$\frac{1}{2}$,,	2	fl. dr.
Pepsin	um					5	,,	10	gr.
Phenac	etinum					5	,,	10	,,
Phenaz	onum (Anti	pyrin,	Analge	sine)		5	,,	20	,,
Phosph	orus				1	00	,,	$\frac{1}{20}$,, .
Physos	tigminæ Sul	phas			7	10	,,	$\frac{1}{20}$,,
Picroto	xinum				1	00	,,	$\frac{1}{25}$,,
Pilocar	pinæ Nitras				3	1 2 0	,,	$\frac{1}{2}$,,
Pilula A	Aloes Barba	densis				4	,,	8	,,
_	— et Asa	fetidæ				4	,,	8	,,
_	— et Fer	ri	• • •			4	,,	8	,,
_	— et Myı	rhæ				4	,,	8	,,
_	- Socotr	inæ				4	,,	8	,,

Pilula	a Cambogiæ	Compo	sita			1 +0	0
		dis Con	mnos	ita			8 gr.
	_	et. I	Hvoso	vami		4 ,,	
	Ferri	001	15030	уашт			8 ,,
_	Galbani Co	mnosit					15 .,
_	Hydrargyri	прови	va			4 ,,	
		Subch	lorid	Comp		4 ,,	0.00
_	Ipecacuanh	æ cum	Scill	On 5	osna	4 ,,	
_	Phosphori (2 nc	BUIL	a (Op. 5	p.c.)	4 ,,	
	Plumbi cun	Opio	(On	191 n a	١	1 ,,	
-	Quininæ Su	Inhatis	(Op.	122 p.C	•)	2 ,,	
_	Rhei Compo	neita	· · · ·		• • •	2 ,,	
_	Saponis Con	mnosit	2 (00	20		4 ,,	
_	Scammonii	Comp	acito				5 ,,
_	Scillæ Com	posite	osita				8 ,,
	izina	posita					8 ,,
Plumb	oi Acetas					-	10 ,,
Podon	hylli Resina						5 ,,
Potass	sii Acetas		• • •			1,,	
_						10 ,,	
	Bromidun					1 0 ,,	
	Carbonas	1					30 ,,
	Chloras	• • • •				5 ,,	20 ,,
	Citras					õ ,,	15 ,,
						10 ,,	40 ,,
	Iodidum Nitras	• •	* * *			5 ,,	20 ,,
		• • • •				5 ,,	20 ,,
	Permanga					1 ,,	3 ,,
	Salicylas					5 ,,	30 ,,
	Sulphas					10 ,,	40 ,,
	Tartras					30 ,, 2	40 ,,
Pulnic	- Acid					20 ,,	60 ,,
Luivis	Antimoniali	S				3 ,,	6 ,,

Pulvis Catechu Composi	tus		10 to 40 gr.
— Cinnamomi Comp			
- Cretæ Aromaticus			10 ,, 60 ,,
— Elaterini Compos			
— Glycyrrhizæ Comp			
— Ipecacuanhæ Com	777		
(Dover's Powder, Opium			
Pulvis Jalapæ Composite	the state of the s		20 to 60 gr.
- Kino Compositus			5 ,, 20 ,,
- Opii Compositus (-	-	2 ,, 10 ,,
- Rhei Compositus	_	-	20 ,, 60 ,,
- Scammonii Comp			10 ,, 20 ,,
* - Sodæ Tartaratæ E			
- Tragacanthæ Com	positu	S	 20 ,, 60 ,,
Quininæ Hydrobromas			 1 ,, 5 ,,
 Hydrochloridur 			 4 40
	Acida	ım	 1 ,, 10 ,,
— Salicylas			 2 ,, 6 ,,
- Sulphas			 1 ,, 10 ,,
Resorcin			 5 ,, 15 ,,
Rhei Radix			 3 ,, 10 ,,
(Single do	ose)		 15 ,, 30 ,,
Salicinum			 5 ,, 20 ,,
Salol			 5 ,, 15 ,,
Santoninum			 2 ,, 5 ,,
Scammoniæ Resina			 3 ,, 8 ,,
Scammonium			
			1 ,, 3 ,,
Senna			10 ,, 30 ,,
Soda Tartarata			 120 ,,240 ,,

^{*} Tartarated Soda 120 gr., and Sodium Bicarbonate 40 gr., with 38 gr. of Tartaric Acid, commonly known as Seidlitz Powder.

Sodii Arsenas	$\frac{1}{40}$ to $\frac{1}{10}$ gr.
— Benzoas	5 ,, 30 ,,
— Bicarbonas	5 ,, 30 ,,
— Bromidum	5 ,, 30 ,,
— Carbonas	5 ,, 30 ,,
- Exsiccatus	3 ,, 10 ,,
— Chloridum	5 ,,20 ,,
Emetic	60 ,,240,,
Citro-Tartras Effervescens	60 ,,120 ,,
— Hypophosphis	3 ,, 10 ,,
— Hyposulphis	5 ,, 20 ,,
— Iodidum	5 ,, 20 ,,
— Nitris	1 ,, 2 ,,
- Phosphas	30 ,,120 ,,
(Single dose)	$\frac{1}{4}$,, $\frac{1}{2}$ oz.
- Effervescens	60 ,,120 gr.
(Single dose)	$\frac{1}{4}$, $\frac{1}{2}$ oz.
- Salicylas	10 ,, 30 gr.
- Sulphas	00 100
(Single dose)	$\frac{30}{4}$,, $\frac{1}{2}$ oz.
- Effervescens	
(Single dose)	"
- Sulphis	$\frac{1}{4}$,, $\frac{1}{2}$ oz.
- Sulphocarbolas	5 ,, 20 gr.
Spiritus Ætheris	
(Single dose)	
- Ætheris Compositus	2
(Single dose)	20 ,, 40 m.
- Nitrosi	2
(Single dose)	20 ,, 40 m.
- Ammoniæ Aromaticus	and the same of th
	20 ,, 40 m.
- (Single dose) - Fetidus	
	20 ,, 40 m.
(Single dose)	$1, 1\frac{1}{2}$ fl. dr.

Spiritus Anisi (10 p.c.)	 5 to 20 min.
- Armoraciæ Compositus	 1 ,, 2 fl. dr.
— Cajuputi (10 p.c.)	 5 ,, 20 min.
— Camphoræ (10 p.c.)	 5 ,, 20 ,,
— Chloroformi (5 p.c.)	 5 ,, 20 ,,
(Single dose)	 30 ,, 40 ,,
— Cinnamomi (10 p.c.)	 5 ,, 20 ,,
— Juniperi (5 p.c.)	 20 ,, 60 ,,
— Lavandulæ (10 p.c.)	 5 ,, 20 ,,
— Menthæ Piperitæ (10 p.c.)	 - 00
— Myristicæ (10 p.c.)	 5 ,, 20 ,,
— Rosmarini (10 p.c.)	5 ,, 20 ,,
Strychnina	
Strychninæ Hydrochloridum	 1 1
Succus Belladonnæ	 - 1
— Conii	 1 ,, 2 fl. dr.
— Hyoscyami	 $\frac{1}{2}$,, 1 ,,
— Scoparii	 1 6)
— Taraxaci	 1 ,, 2 ,,
Sulphonal	 10 ,, 30 gr.
Sulphur Præcipitatum	 20 ,, 60 ,,
— Sublimatum	 20 ,, 60 ,,
Syrupus Aromaticus	 $\frac{1}{2}$,, 1 fl. dr.
— Aurantii	 $\frac{1}{2}$,, 1 ,,
Floris	 $\frac{1}{2}$,, 1 ,,
 Calcii Lactophosphatis 	 $\frac{1}{2}$ to 1 ,,
— Cascaræ Aromaticus	 $\frac{1}{2}$,, 2 ,,
— Chloral (1 in 6)	 $\frac{1}{2}$,, 2 ,,
— Codeinæ (2 grs. in fl. oz.)	 $\frac{1}{2}$,, 2 ,,
— Ferri Iodidi	 $\frac{1}{2}$,, 1 ,,
— Phosphatis	 1 ,,
cum Quin.	
Strych.	 $\frac{1}{2}$,, 1 ,,

	Surunus	Hemidesr	ni				1		7	0 1
,	—									fl. dr.
		Limonis						,,		
		Pruni Virg		næ			2	, ,	1	,,
	_	Rhei					2	to	2	,,
		Rhœados					$\frac{1}{2}$,,	1	,,
	_	Rosæ					$\frac{1}{2}$,,	1	,,
	_	Scillæ					$\frac{1}{2}$,,	1	,,
	-	Sennæ					1/2	,,	2	,,
	_	Tolutanus					$\frac{1}{2}$,,	1	,,
	-	Zingiberis					1		1	
7	labellæ	Trinitrini	(Niti	roglyce	erini)		(1)	1	gr.)
7	Cereben	um					5			min.
1	Theobron	nina					1		5	gr.
1	Theobron	ninum Nati	rio-Se	alicylic	um (I	Diuretin)	5	"	15	,,
T	hymol						1	"	2	"
I	hyroid	eum Siccur	n				3	,,	10	"
								,,	10	"
T	inctura	Aconiti					2		5	min.
		(Single	e dos	e)			5	"	15	
			æ (A	lkaloi	ds 1	p.c.)	5	,,	15	,,
	_	Cannabis	Indi	cæ	20	p.c.)	5	,,	15	
	_	Cantharid	is	Control			9	"	5	,,
		(Sing	le do	se)			7	"	15	"
		Capsici		50)			5	"	15	,,
	_	Chloroforn	mi et	Morn	hinm	Com	0	"	19	"
		posita-		morp	шие	Com-				
						71 0				
		Dil	Hede	Λ		$7\frac{1}{2}$ p.c.				
		Morro	h II	Ac.		5 ,,	~			
		Cocci	п. П	yur.		1 ,,	5	,,	15	,,
		Cocci .	omi				5	,,	15	,,
		Colchici S	emin	um			5	,,	15	"
		Convallario	æ		• • • •		5	,,	30	"
		Croci					5	,,	15	,,

Tinctura	Digitalis			5 to	15	min.
_	Ferri Perchloridi			5 ,,	15	,,
_	Gelsemii			5 ,,	15	,,
_	Iodi			2 ,,	5	,,
_	Lobeliæ Ætherea			5 ,,	15	,,
_	Nucis Vomicæ (Str	ych. 1/4 p.c.)	5 ,,	15	,,
_	Opii (Morphine 3					
	1 gr. Op. in 15 n	nin.)		5 ,,	15	,,
	(Single dose)			20 ,,	30	,,
_	Podophylli			5 ,,	15	,,
_	Scillæ			5 ,,	15	,,
_	Stramonii			5 ,,	15	,,
	Strophanthi			5 ,,	15	,,
Tinctura	Aloes			$\frac{1}{2}$,,	1 :	fl. dr.
	(Single dose)			$1\frac{1}{2}$,,		,,
_	Asafetidæ			1/2 ,,	1	,,
_	Aurantii			$\frac{1}{2}$,,		,,
_	Benzoini Composita	a				,,
_	Buchu			$\frac{1}{2}$,,	-	,,
_	Calumbæ			$\frac{1}{2}$,,	1	,,
_	Camphoræ Compos					
	2 gr. in 1 fl. oz.)	50 57 5		$\frac{1}{2}$,,	1	,,
_	Cardamomi Compo	sita		$\frac{1}{2}$,,	1	,,
	Cascaræ Sagradæ			$\frac{1}{2}$,,	1	,,
_	Cascarillæ			1/2 ,,	1	,,
	Castorei			$\frac{1}{2}$,,	1	,,
	Catechu			$\frac{1}{2}$,,	1	,,
_	Chiratæ			1 ,,	1	,,
	Cimicifugæ			$\frac{1}{2}$,,	1	,,
	Cinchonæ (Alkaloid	ds 1 p.c.)			1	,,
		ta (Alk. 1]			1	,,
	Cinnamomi				1	,,

PHARMACOPŒIA.

Ti	nctura	Conii		 $\frac{1}{2}$ to	1	fl. dr.
	_	Cubebæ		 $\frac{1}{2}$,,	1	,,
	_	Ergotæ Ammoniata	ı	 $\frac{1}{2}$,,	1	, ,
	_	Gentianæ Composit	a	 $\frac{1}{2}$,,	1	, ,
	.—	Guaiaci Ammoniata	a	 $\frac{1}{2}$,,	1	,,
	_	Guaranæ		 $\frac{1}{2}$,,	1	,,
	_	Hamamelidis		 $\frac{1}{2}$,,	1	,,
		Hydrastis		 1,,	1	3,
	_	Hyoscyami		 1,,	1	,,
		Jaborandi		 $\frac{1}{2}$,,	1	,,
	_	Jalapæ		 1/2 ,,	1	,,
	_	Kino		 1 ,,	1	,,
	_	Krameriæ		 $\frac{1}{2}$,,	1	,,
		Lavandulæ Compos	sita	 $\frac{1}{2}$,,	1	,,
	_	Limonis		 1 ,,	1	,,
	_	Lupuli		 $\frac{1}{2}$,,	1	,,
	_	Myrrhæ		 $\frac{1}{2}$,,	1	,,
	_	Opii Ammoniata (a		2 //		
				1/2 ,,	1	,,
	_	Pruni Virginianæ		 1 ,,	1	,,
		0		 1 ,,	1	,,
		Quillaiæ		 $\frac{1}{2}$,,	1	,,
		Quininæ (2 p.c.)		 1/2 ,,	1	,,
		— Ammonia		1,,	1	,,
	_	Rhei Composita		 $\frac{1}{2}$,,	1	,,
		(Single dose)		 2 ,,	4	,,
		Senegæ		 1 ,,	1	,,
	_	Sennæ Composita		 1 ,,	1	,,
		(Single dose		 2 ,,	4	,,
		Serpentariæ		 1 ,,	1	,,
	_	Sumbul		 $\frac{1}{2}$,,	1	,,
	-	Tolutana		1 ,,	1	,,
	-	Valeriana Ammoni		1 ,,	1	,,
	_	Zingiberis		 1/2 ,,	1	,,
				- //		11

Trochiscus Acidi Benzoici	 		(½ gr.)
— — Carbolici			(1,,)	
— — Tannici			$(\frac{1}{2},,)$	
 Bismuthi Composit 			(2,,)	
- Boracis	 * * *		(3,,)	
— Catechu	 		(1,,)	
- $Cubebæ$	 		$(\frac{1}{2},,)$	
- Eucalypti Gummi	 		(1,,)	
— Ferri Redacti				
— Guaiaci Resinæ				
- Ipecacuanhæ	 		$(\frac{1}{4},,)$	
— Krameriæ				
- Krameriæ (1 gr.) et				
- Morphinæ				
— Morphinæ $(\frac{1}{36} gr.)$ e			$(\frac{1}{12},,)$	
 Potassii Chloratis 				
— Santonini				
 Sodii Bicarbonatis 				
— Sulphuris				
Vinum Antimoniale (2 gr. Tart. A				
As an Emetic				
— Colchici			30 min.	
_ Ferri				
— — Citratis				
- Ipecacuanhæ (Expector				
- (Emetic)				•
- Quininæ (1 gr. to fl. oz.)				
Zinci Acetas				
- Oxidum				
- Sulphas				
_ (Emetic)				
- Valerianas		1 ,,		

SUBSTANCES USED FOR HYPODERMIC MEDICATION.

The four following formulæ are in the British
Pharmacopæia:—

	Pharmacopæia :
I	njectio Apomorphinæ Hypodermica (1 p.c.).
	Apomorphine Hydrochloride 1 grain.
	Diluted Hydrochloric Acid 1 minim.
	Distilled Water to 110 minims.
	Dose, subcutaneously, 5 to 10 minims.
I	njectio Cocainæ Hypodermica (10 p.c.).
	Cocaine Hydrochloride 10 grains.
	Salicylic Acid å grain.
	Distilled Water to 110 minims.
	Dose, subcutaneously, 2 to 5 minims.
I	njectio Ergotæ Hypodermica (33 p.c.).
	Extract of Ergot 33 grains.
	Phenol 1 grain.
	Distilled Water to 110 minims.
	Dose, subcutaneously, 3 to 10 minims.
I	njectio Morphinæ Hypodermica (5 p.c.).
	Morphine Tartrate 5 grains.
	Distilled Water to 110 minims.
	Dose, subcutaneously, 2 to 5 minims.
S	ome of the Liquors of the British Pharmacopæia may also
	be used for hypodermic injection in doses as follows:-
	Liquor Atropinæ Sulphatis (1 p.c.)
	Dose, subcutaneously, 1 to 3 minims.
	Liquor Morphinæ Acetatis (1 p.c.)
	Dose, subcutaneously, 10 to 30 minims.
	Liquor Morphinæ Hydrochloridi (1 p.c.)
	Dose, subcutaneously, 10 to 30 minims.
	V 2

Liquor Morphinæ Tartratis	(1 p. c.)
Dose, subcutaneously, 10 to	30 minims.
Liquor Strychninæ Hydrochlorid	di (1 p.c.)
Dose, subcutaneously, 1 to	2 minims.

The following are used in the form of discs, to be dissolved in 6 to 10 minims of distilled water at the time of using.

Those printed in brackets are usually kept.

Apomorphinæ Hydroc	hlorid	lum		1	, [10] gr.
Atropinæ Sulphas			[$[\frac{1}{150}], \frac{1}{100}$	$\left[\frac{1}{60}\right]$,
Caffeinæ Sodio-Salicyla	S				$[\frac{1}{2}]$,,
Cocainæ Hydrochlorid	um			6,	$\frac{1}{4}$, $\left[\frac{1}{2}\right]$,
Hyoscinæ Hydrobromi	idum			[200	$], \left[\frac{1}{75}\right] ,,$
Hyoscyaninæ Sulphas				1	$\frac{1}{0}$, $\begin{bmatrix} \frac{1}{20} \end{bmatrix}$,,
Morphinæ Sulphas			$\frac{1}{12}$,	$[\frac{1}{8}, [\frac{1}{6}], [\frac{1}{4}]]$	$], \frac{1}{3}, \frac{1}{2} :,$
Physostigminæ Sulpha	atis				1 0 0 ,,
Pilocarpinæ Nitras				$\left[\frac{1}{10}\right]$	$[\frac{1}{6}], \frac{1}{3},$
Sparteinæ Sulphas					$\frac{1}{2}$,,
Strychninæ Sulphas				$\frac{1}{150}$, $\frac{1}{100}$	$\left[\frac{1}{60}\right]$,
Trinitrini					250 ,,
The following have	0000	sionall.	r hoon	admini	stored

The following have occasionally been administered hypodermically:—

Codeina $\frac{1}{4}$ gr. $\frac{1}{000}$... $\frac{1}{100}$... Also in Oily Solvents :—

Creosotum 1 to 3 min.

Guaiacol 1, 3 ,,

As Stimulants may be used hypodermically:-

Brandy ... $\frac{1}{2}$ to $1\frac{1}{2}$ fl. dr. Ether ... 20 m. to 1 fl. dr.

LIST OF THE MORE IMPORTANT PREPARATIONS OF THE BRITISH PHARMACOPŒIA FOR EXTERNAL USE,

With the approximate percentages by weight of the active ingredients.

Acetum Cantharidis				10 p.c.
Argenti Nitras.				
Induratus.				
- Mitigatus.				
Charta Sinapis.				
Collodium.				
— Flexile.				
Vesicans.				
Emplastrum Ammoniaci o	eum H	ydrai	0179	20 p.c.
— Belladonnæ				aloids) $\frac{1}{2}$,,
Calefaciens				arides) 4,,
 Cantharidis 				35 ,,
 Hydrargyri 				33 ,,
35				15 ,,
— Opii				10 ,,
- Picis				17
— Plumbi)xide) 25 ,,
- Plumbi Iodio				
				(xide) 20 ,,
- Saponis				
Extractum Belladonnæ Li	iquidu	m	(Alka	loids) 3
Glycerinum Acidi Borici				30,,
- Carboli	ci			20 ,,
— — Tannici				20 ,,
				16,,
— Amyli.				
- Boracis				16,,
— Plumbi Subac				25 ,,
Hydrargyri Oleas.				
Lamellæ Atropinæ			(Each d	isc) 1/6000 gr.

Lamellæ Cocainæ				1	cer
- Homatropinæ		• • • •			gr.
= Physostigminæ				$\frac{1}{100}$	
Linimentum Aconiti				1 0 0 0 GG v	
— Ammoniæ				66 I	
— Ammoniæ — Belladonnæ	• • •	 (Tion		25	1)
		-	id Extract)		,,
— Calcis			on of Lime)		"
— Camphoræ				20	,,
			(Camphor)	_	,,
Strong Solution			nia		,,
- Chloroformi				50	,,
				$12\frac{1}{2}$,,
— Hydrargyri				16	,,
— Opii	('.	Cincture	of Opium)	50	,,
— Potassii Iodid	li cun	n Sapon	e	14	,,
— Saponis			(Camphor)	5	,,
— Sinapis		(Oil o	of Mustard)	4	,,
- Terebinthinæ				65	,,
				44	,,
Liquor Epispasticus			antharides)	50	,,
— Iodi Fortis				10	,,
 Picis Carbonis 				20	,,
- Plumbi Subacetatis				13	,,
		is		25	,,
- Potassæ				6	,,
Lotio Hydrargyri Flava		2.500		1 2	,,
- Nigra				2 2 3	
Unguentum Acidi Borici				10	,,
— — Carbolic		•••		4	,,
				2	,,
— Salicylic	1				"
- Aconitinæ				2	"
- Aquæ Rosæ				0	
Atropinæ			(Allrala; 2)	2	,,
— Belladonnæ			(Alkaloid)		"
 Cantharidis 				9	,,

TT						
U	nguentur	n Capsici				20 p.c.
	-	Cetacei				20 ,,
	_	Chrysarobi	ni			4 ,,
	_	Cocainæ				4 ,,
	_	Conii		(8	Succus Co	
		Creosoti				10 ,,
	_	Eucalypti				10
	_	Gallæ				00
	-	Gallæ cum	Opio			10-171
	_	Glycerini Pl	umbi S	Subaceta	atis(Lead	$(Op.) \frac{7}{2},$
	_	Hamamelid	is		(Lia E	-4 \ 10
	_	Hydrargyri				50
	_	_		oniati		50 ,,
	_	_		ositum		10 ,,
	_	_		Rubri		20 ,,
	_	Hydrargyri				4 ,,
			11101ac			10 ,,
	_	_	Oleati	Dilut		2 ,,
				Flavi		
						2 ,,
				Rubri		10 ,,
		Iodi	Subch	loridi		10 ,,
	200.00					4 ,,
		Iodoformi				10 ,,
	_	Paraffini Picia I : 12				
	_	Picis Liquid	æ			70 ,,
	_	Plumbi Acet				4 ,,
	_		onatis			10 ,,
	_		di			10 ,,
	_	Potassii Iod	idi			10 ,,
		Resinæ				27 ,,
	_	Staphisagria				17 ,,
	-	Sulphuris				10 ,,
	_		didi			4
	_	Veratrinæ				9
	-	Zinci				1.5
	-	- Oleatis				10 ,,

LIST OF SUPPOSITORIES OF THE BRITISH PHARMACOPŒIA.

Suppositorium Acidi Carbolici (1 gr.).

- Tannici (3 gr.).
- Belladonnæ (1½ gr. Alk. Extr.)
- Glycerini (70 p.c.).
- Hydrargyri (5 gr.).
- Iodoformi (3 gr.).
- Morphinæ (Morph. Hydr. gr. ⅓).
- Plumbi Compositum (Lead Ac. 3 gr., Op. 1 gr.).

ENGLISH MEASURES AND WEIGHTS.

APOTHECARIES' FLUID MEASURE.

60 Minims (M))	=	1 Fluid Drac	hm	 f.3
8 Drachms		=	1 Fluid Oun	ce	 f.3
20 Ounces		=	1 Pint		 0
8 Pints		=	1 Gallon		 C

A Tea-spoonful should be equivalent to one Drachm, a Dessertspoonful to two, a Table-spoonful to four.

APOTHECARIES' WEIGHT.

	20 Grains	=	1 Scruple	 Э
(60 gr.)	3 Scruples	=	1 Drachm	 3
	8 Drachms	-	1 Ounce	 3
	12 Ounces	=	1 Pound	 tb

The Ounce and the Pound of Apothecaries' Weight may be considered obsolete; the Weights of the British Pharmacopæia being:—

437.5 Grains = 1 Ounce Avoirdupois.

16 Ounces (7000) gr. = 1 Pound Avoirdupois.

METRIC SYSTEM OF WEIGHTS AND MEASURES.

The Unit of Length is the Metre.

The ten-millionth part of the estimated length of a quarter of a terrestrial meridian.

The Unit of Capacity is the Litre.

The volume of a cube, the length of whose edge is one-tenth of a metre.

The Unit of Weight is the Gramme.

The weight of the volume of distilled water (at 4° C., when its density) is greatest) contained in a cube the length of whose edge is one-hundredth of a metre.

Decimal fractional parts are indicated by prefixes derived from the Latin:—

Deci.

Centi.

Milli.

Decimal multiples are indicated by prefixes derived from the Greek:—

Deka.

Hecto.

Kilo.

Thus-

*Millimetre ... = $\frac{1}{1000}$ or .001 metre.

Centimetre ... = $\frac{1}{100}$,, '01

Decimetre $\dots = \frac{1}{10}$, 1 ,

Metre.

Dekametre $\dots = \dots = 10$ metres.

Hectometre ... = ... 100 ,

Kilometre ... = ... 1000 ,,

Millilitre ... = $\frac{1}{1000}$ litre = 1 cubic centimetre.

Centilitre ... = $\frac{1}{100}$,, = 10 .. ,

Decilitre ... = $\frac{1}{10}$,, = 100 ,,

Litre = 1000 ,, ,,

Dekalitre ... = 10 litres.

Hectolitre $\dots = 100$,,

Kilolitre ... = 1000 ,,

^{*}Histologists use the micromillimetre (μ) , which is $\frac{1}{1000}$ of a millimetre.

Milligramme ... = '001 gramme.

Centigramme ... = '01

Decigramme .. = '1

... = Weight of 1 cubic centimetre of Gramme

Dekagramme ... = 10 grammes. [water at 4° C.

Hectogramme.. = 100

Kilogramme ... = 1000 ,,

Relation between English Measures and Weights and those of the Metric System.

1 Metre ... = 39.37079 inches.

1 Gramme = 15.43235 grains.

The following Approximations, sufficiently close for some purposes, may be noted :-

Metre ... 39\frac{3}{2} inches.

Decimetre ... $3\frac{15}{16}$,,

Centimetre ... $\frac{2}{5}$,,

Millimetre Millimetre ... $\frac{1}{25}$,, Litre ... $1\frac{3}{4}$ pints.

Cubic Centimetre ... 17 minims.

Kilogramme ... 354 ounces,

Gramme ... $15\frac{1}{2}$ grains.

Centigramme ... between 1 and 1 grain.

Milligramme d grain.

Foot 305 millimetres.

Inch 25

Pint ... 568 cubic centimetres

Fluid Ounce ... 28

Ounce 28 grammes.

Grain 65 milligrammes.

	No. 1. FULL DIET (Men). [Dieta Carnis.] D. C.	No. 2. FULL DIET (Women). [Dieta Carnis.] D. C.
BREAKFAST { TEA{	Tea. \$\frac{3}{2}\$ pint. Butter. \$\frac{1}{2}\$ oz. Tea. \$\frac{3}{4}\$ pint. Butter. \$\frac{1}{2}\$ oz.	Tea. \$\frac{3}{2}\$ pint. Butter. \$\frac{1}{2}\$ oz. Tea. \$\frac{3}{4}\$ pint. Butter. \$\frac{1}{2}\$ oz.
	Meat, 6 times a week 6 oz. (dressed, without bone) Fish, once a week10 oz. (dressed, without bone)	Meat, 6 times a week 4 oz. (dressed, without bone) Fish, once a week 8 oz. (dressed, without bone)
DURING REMAINDER OF 24 HOURS	Potato	Potato
	Pudding. 4 oz. Milk. 1 pint. Bread. 18 oz.	Pudding. 4 oz. Milk. 1 pint. Bread. 12 oz.

MEAT AND FISH :-

Sunday Roast	Beef. Th
	Mutton. Fri
Tuesday Irish S	Stew. Sat
Wednesday Roast	Mutton.

Thursday Boiled Mutton Friday Fish.
Saturday Roast Mutton.

Patients on admission are to be placed on Spoon Diet

Beef Tea, Beef Essence, Mutton Broth, and Mutton Dry Arrowroot and Cocoa, in powder, to be

It is particularly requested that, in ordering Diets, the Medical Officers ordered for a Patient unless the Medical Officer considers it essential be re-written weekly.

HOSPITAL-DIET SCALE.

No. 3.	No. 4.	No. 5.
CONVALESCENT DIET.	MILK DIET.	SPOON DIET.
Dieta Dimidia. D. D.	$\begin{bmatrix} \textit{Dieta Lactis.} \\ \textit{D. L.} \end{bmatrix}$	Dieta Tenuis. D. T.
Tea ½ pint.	Tea ½ pint.	Tea ½ pint.
Butter $\frac{1}{2}$ oz.	Butter $\frac{1}{2}$ oz.	
Tea ½ pint.	Tea $\frac{1}{2}$ pint:	Tea ½ pint.
Butter $\frac{1}{2}$ oz.	Butter \dots $\frac{1}{2}$ oz.	
Mutton Ghop 4 oz. (dressed, without bone)	Pudding (not Suet) 8 oz.	
Or one of the following:		
Minced or Pounded Meat 4 oz. (dressed, without bone)		
Fish		
Rabbit 4 oz. (dressed, without bone)		
Mashed Potato 6 oz.		
Pudding (not Suet) 8 oz.		
Milk 2 pints.		. Milk 2 pints.
Bread { Men 14 oz. Women 12 oz.	Bread $\{ \begin{array}{l} \text{Men} \dots 14 \text{ oz.} \\ \text{Women} \dots 12 \text{ oz.} \end{array} $	
Women12 oz.	Women12 oz.	

PUDDINGS :-
Rice Four times a week.
Sago Once a week.
Suet Twice a week.
(with Golden Syrup)
Custard Only when specially
andanad

 $\begin{array}{c} \text{CHILDREN'S DIET:-}\\ \text{For Young Children} \left\{ \begin{array}{c} Spoon\ or\ Milk\\ Diet,\\ without\ Tea. \end{array} \right. \\ \text{For Older Children}\ .\ \ \textit{Women's Full}\\ Diet. \end{array}$


until a Diet is ordered by the Physician or Surgeon.


Essence to be supplied only when specially ordered.


kept in the Wards for use when ordered.


will adhere as closely as possible to the above Diet Scale. No extra is to be to the Patient's proper treatment. All Diets, as well as stimulants, are to




















mos.xc.n.Sc.

PAGE #IN

DUE TO TISSUE