

**Management of the organs of digestion, in health and in disease / by  
Herbert Mayo.**

**Contributors**

Mayo, Herbert, 1796-1852.  
University of Glasgow. Library

**Publication/Creation**

London, 1840.

**Persistent URL**

<https://wellcomecollection.org/works/pqe3rqu5>

**Provider**

University of Glasgow


**License and attribution**

This material has been provided by This material has been provided by The University of Glasgow Library. The original may be consulted at The University of Glasgow Library. where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>



No.


*B459- i. 1*


1875.


GLASGOW  
UNIVERSITY  
LIBRARY.

---

*EUING COLLECTION.*







Digitized by the Internet Archive  
in 2016

<https://archive.org/details/b24921968>

NEW YORK

MAYO ON DIGESTION.



LONDON:  
HARRISON AND CO., PRINTERS,  
ST. MARTIN'S LANE.

MANAGEMENT  
OF THE  
ORGANS OF DIGESTION,  
IN  
HEALTH AND IN DISEASE.



BY HERBERT MAYO, F.R.S.

SENIOR SURGEON OF THE MIDDLESEX HOSPITAL,  
FORMERLY ONE OF THE PROFESSORS OF ANATOMY AND SURGERY  
TO THE ROYAL COLLEGE OF SURGEONS.

*THE SECOND EDITION.*


LONDON :  
JOHN W. PARKER, WEST STRAND.

M.DCCC.XL.

MANAGEMENT

ORGAN OF THE DISTRICT

HEALTH AND INDIAN



THE DISTRICT

HEALTH AND INDIAN

THE DISTRICT

THE DISTRICT

## ADVERTISEMENT.

---

IN a volume entitled *The Philosophy of Living*, I have explained the principles which should guide us as to diet, exercise, sleep, clothing, choice of residence, with a view to the preservation of health. But for the same object, other rules require to be observed. These regard the management of the Stomach in circumstances at variance with perfect health, or when its powers have become impaired, or are naturally feeble; and the regulation of the Bowels. The following pages contain a series of instructions upon these points, and are so far supplemental to the former

work. But the sketch, which I have here given, goes further, and includes the consideration of Indigestion generally, and of most of the functional disorders of the bowels; together with a special account of the symptoms and treatment of diseases of the Rectum. Upon the latter subject I had formerly published a work, now out of print.

H. M.

19, *George Street,*  
*Hanover Square.*

---

# CONTENTS.

---

## CHAPTER I.

### MANAGEMENT OF THE STOMACH.

	Page
The process of Digestion . . . . .	1
Different Qualities required in Food . . . . .	3
Kind of Food suited to particular Ages and Con- stitutions . . . . .	4
Want of Moderation, the common error of Diet . . . . .	6
Use of Food as a Stimulant . . . . .	6
Requisite with some, immediately on Rising . . . . .	7
In Plethoric Persons . . . . .	9
Use of Suppers . . . . .	10
Epilepsy produced by Depletion and Exhaustion . . . . .	13
Opposite Sources of Head Disorder . . . . .	16
Diet before and after Surgical Operations . . . . .	21
Nature of Indigestion . . . . .	23
Accidental Indigestion . . . . .	26
Protracted Disorder from Undigested Food . . . . .	28
Bile on the Stomach . . . . .	30
Habitual Indigestion from Weakness of Stomach . . . . .	35
Means applicable to the Treatment of Indigestion . . . . .	41
Diet . . . . .	42
Exercise . . . . .	45
Aperient Medicine . . . . .	45
Tonics . . . . .	46
Heartburn, its Nature and Treatment . . . . .	49
Waterbrash . . . . .	52

	Page
Pain of the Stomach . . . . .	54
— from Undigested Food . . . . .	54
— from Habitual Indigestion . . . . .	54
— from Heightened Sensibility of the Stomach . . . . .	57
— occurring when the Stomach is empty . . . . .	57
— from Spasm . . . . .	59
— from Ulceration . . . . .	60
Vomiting . . . . .	61
Sickness of Pregnancy . . . . .	63
Sea-sickness . . . . .	64
Vomiting following Surgical Operations . . . . .	66
Vomiting, a part of Habitual Indigestion . . . . .	67
Vomiting from Cancer of the Stomach . . . . .	67
Symptomatic Head Disorders . . . . .	69
Symptomatic Chest Affections . . . . .	72

## CHAPTER II.

## MANAGEMENT OF THE BOWELS.

Sources of the Alvine Excretions . . . . .	75
I. OF LOOSE BOWELS.	
Looseness habitual to certain Constitutions . . . . .	80
Liable to be increased and become Disease . . . . .	81
Cases and Treatment . . . . .	83
Simple Diarrhœa . . . . .	88
Febrile Diarrhœa . . . . .	90
Diarrhœa, with absence of Bile . . . . .	90
Cholera . . . . .	91
Asiatic Cholera . . . . .	92
II. OF CONSTIPATION OF THE BOWELS.	
Costiveness natural to strong Constitutions . . . . .	95
Circumstances which encourage it . . . . .	95
Constipation from thin Bowels . . . . .	95
Constipation salutary . . . . .	97

	Page
Constipation from Inattention . . . . .	98
————— from Neglect of Exercise . . . . .	99
————— from Deficiency of Bile . . . . .	99
————— after Residence in Hot Climates . . . . .	99
————— from Affections of the Brain and Spinal Marrow . . . . .	100
————— from Hysteria . . . . .	101
Constipation either Complete or Incomplete . . . . .	102
Relation of Costiveness to Obstruction . . . . .	103
————— to different periods of Life . . . . .	103
Treatment of Costiveness . . . . .	104
Exercise . . . . .	106
Regular times of Relieving the Bowels . . . . .	106
Laxative Diet . . . . .	106
Injections . . . . .	108
Galvanism . . . . .	109
Cold and Hot Bathing . . . . .	109
Laxative Medicine . . . . .	110
Drastic Purgatives . . . . .	112
Mercury . . . . .	112

## CHAPTER III.

## DISEASES OF THE RECTUM.

Causes . . . . .	115
Means of Prevention . . . . .	116

## I. FULNESS OF THE HEMORRHOIDAL VESSELS AND BLEEDING.

Causes of Local Congestion . . . . .	118
Treatment . . . . .	119
Congestion, with Bleeding . . . . .	119
Varieties of this case, and their Treatment . . . . .	120
Bleeding, without Congestion . . . . .	121



II. OF PILES.	Page
Their Nature generally . . . . .	125
External Piles . . . . .	128
Varieties and Treatment . . . . .	128
Anal Piles . . . . .	131
Varieties and Treatment . . . . .	131
Mistaken for Inward Piles . . . . .	134
——— for Abscess . . . . .	134
Not to be opened unnecessarily . . . . .	135
Inward Piles . . . . .	136
Four Varieties . . . . .	137
Treatment of the First . . . . .	138
——— of the Second . . . . .	139
——— of the Third and Fourth . . . . .	142
Changes that supervene spontaneously . . . . .	143
 III. PROLAPSUS.	
Origin and Progress. . . . .	143
Treatment in Children . . . . .	145
——— in Adults . . . . .	145
Use of Instruments . . . . .	146
Cure by operation . . . . .	147
Prolapsus from Weakness of the Sphincter . . . . .	151
 IV. OF FISSURE OF THE RECTUM.	
The Origin commonly Mechanical . . . . .	152
Treatment ordinarily sufficient . . . . .	152
Operation . . . . .	153
 V. CONTRACTION OF THE ANUS.	
Spasm of the Sphincter . . . . .	154
Permanent Narrowing . . . . .	156
 VI. OF STRICTURE OF THE RECTUM.	
Spasmodic Stricture . . . . .	157
Cases . . . . .	158
Permanent Stricture . . . . .	164
Treatment . . . . .	166
Danger attending the injudicious use of Bougies . . . . .	166
Stricture resembling Cancer . . . . .	168

	Page
<b>VII. OF SUBSTANCES IN THE RECTUM PARTIALLY OR COMPLETELY OBSTRUCTING THE PASSAGE.</b>	
Accumulations of Fæces . . . . .	171
Hard Intestinal Concretions . . . . .	176
Hard and Indigestible Substances swallowed . . . . .	178
Bodies directly introduced . . . . .	179
<b>VIII. OF DEFECTS OF THE RECTUM EXISTING AT BIRTH.</b>	
Closure of the Orifice . . . . .	179
Partial . . . . .	179
Complete . . . . .	182
The Rectum deficient . . . . .	183
Case of Restoration . . . . .	185
Wrong termination of the Rectum . . . . .	186
<b>IX. OF LACERATION OF THE INTESTINE.</b>	
Incomplete Laceration . . . . .	187
Laceration of the Sphincter in Labour . . . . .	188
Operations employed to remedy it . . . . .	191
Laceration above the Sphincter . . . . .	192
<b>X. ABSCESS AND FISTULA.</b>	
Diffused Abscesses . . . . .	195
Circumscribed Abscesses . . . . .	196
Fistula . . . . .	197
<b>XI. CANCER OF THE RECTUM.</b>	
Various kinds . . . . .	201
Treatment . . . . .	204
Operation . . . . .	205
Cancerous Polypi . . . . .	206
Discharges from the Rectum . . . . .	206

---

**“THERE** is a wisdom in this beyond the rules of physic : a man’s own observation, what he finds good of and what he finds hurt of, is the best physic to preserve health. \* \* \* \*  
If you make physic too familiar, it will make no extraordinary effect when sickness cometh. I commend rather some diet for certain seasons, than frequent use of physic, except it be grown into a custom ; for those diets alter the body more, and trouble it less.”—BACON.

MANAGEMENT  
OF  
THE ORGANS OF DIGESTION,  
IN HEALTH AND IN DISEASE.

---

CHAPTER I.

MANAGEMENT OF THE STOMACH.

IN the first chapter of the volume\*, to which the present is intended to form a supplement, I have explained the physiology of digestion, and the nature of the changes which food undergoes in the stomach, and have laid down the principles which should regulate the quantity, quality, and frequency of our meals at different ages, and in various states of health. The following is a summary of the observations on diet therein contained.

The process of digestion is the solution of food in the stomach through the chemical action of a sub-acid liquid, called the gastric juice, which is poured out by the lining membrane of the stomach, much in the same manner as the perspiration exudes upon the

\* *The Philosophy of Living.*

surface of the skin on other occasions, and for other purposes. The quantity of gastric juice, which the stomach can furnish at a single meal, depends not only on the quantity of digested food which the system is in want of to make up for previous waste, and to supply the materials of growth, but on a great variety of other causes, which are capable, directly or indirectly, of raising or lowering the powers of digestion. In health, the appetite is the natural index and measure of the ability of the stomach to digest.

Of what we introduce into the stomach, a greater or less quantity is water, pure, or nearly so. Water, and such simple liquids as water containing a very small proportion of wine or ardent spirit, neither are digested, nor are mixed up in the process of digestion, but are carried away from the stomach soon after they have been introduced into it, leaving any food which may have been taken at the same time, to be there chymified. The quantity of simple liquid (within certain limits, of course) taken *at a single meal* is, therefore, of little consequence, as every one knows by experience, who has sat down to dinner thirsty from exercise. But the habitual practice of taking large quantities of liquid is, on the other hand, injurious; liquids fill the stomach and bowels with wind, and lead to the production of a relaxed habit of body, causing either fat, or excessive per-

spirations, or both, and immoderate flow of urine. The less liquid taken daily, the firmer the flesh, the stronger the stomach, the wholesomer the condition of the body.

Liquid largely combined with food, as in soups and broths, taken as a meal, renders it less easy of digestion, or the gastric juice finds a difficulty in getting at and acting upon the food in this state of dilution. But this holds in health only, and during illness the rule is often reversed.

The uses of liquid, as part of our diet in health, are, first to make up for the waste occasioned by excessive perspirations, or alvine discharges; secondly, to form vehicles, by which different stimulants and restoratives may be introduced into the system, such as tea, coffee, and the like, that contribute to our health or comfort.

That which constitutes our food, if for a single meal, must possess *one quality*; if for a succession of meals, it must combine *three*.

The food for a single meal must be *digestible*; for a succession of meals, *digestible, nutritious, stimulant*.

The essence of *digestibleness* is, that the gastric juice should easily penetrate and act upon the food. The food may resist the action of the gastric juice either mechanically or chemically; tough hard meat, especially if swallowed in unchewed morsels, exemplifies the first source of indigestibleness, so

likewise oily food, and broths and soups. The flesh of young animals, and certain rich meats, as pork, duck, salmon, are refractory in the second way.

Different sorts of food are *nutritious* in very different degrees. The flesh of adult warm-blooded quadrupeds furnishes most nutriment; next that of birds, game more than poultry: then the best farinaceous food, arrowroot, wheaten bread, &c.; then fish; finally, inferior farinaceous food, and esculent and mucilaginous vegetables and fruits.

Food must have a *stimulating* quality. If animals are fed on any one kind of concentrated food exclusively, they fatten, indeed, for the first fortnight, but then they begin to lose flesh, waste, and, after a brief period, die. A stimulating quality is imparted to food in four ways,—first, by using aliments that are not simple and concentrated, milk, for instance, instead of sugar, or butter alone:—secondly, by varying our food at different meals:—thirdly, by the use of condiments:—fourthly, by taking with our meals fermented liquors.

In infancy and childhood, what is wanted is food easy of digestion, but not stimulating, nor of the most nutritive quality, for that in early life heats and excites. Food of this description is signified by the term “light food.” Light food goes with frequent meals,—that is to say, the stomach which requires it can consume a moderate quantity only

at a time, and so requires another meal before a long interval. For children with struma in the habit, a diet more nutritious and stimulant is required than for others.

The digestion of convalescents resembles the digestion of children. In age, which is second childhood, frequent but moderate meals again are necessary; but they must not consist in any large proportion of liquid aliment, like those of young children and convalescents: and they require to be both nutritive and stimulant.

In those, whose physical strength is largely drawn upon, in nursing women, in men in training for athletic feats, a diet at the same time nutritive and stimulating, and plentiful, and repeated meals, are necessary.

To grown-up persons, in health, but with no similar calls upon their strength, meals few and sparing, varied in quality, nutritious, but neither too easy of digestion, nor combining adventitious stimulants, and just hearty enough to prevent craving for food at intermediate hours, are to be studied. The general tendency in the strong and the healthy is to overload the stomach: with such the appetite commonly exceeds the powers of digestion; and the rule to be constantly borne in mind in respect to diet (as indeed to everything else) is to observe moderation.

Men can bear stimulants in excess better than



women ; but they require stimulants less. To the health of many women one or two glasses of wine daily is essential, whereas most men would be the better for disusing wine altogether.

These heads, and the views to which they lead, having been fully gone into in the work already referred to, I hence proceed to wind off the thread of practical observations and advice, which I propose should constitute the present chapter.

We may assume it to be true, that men in good health, and not taking violent exercise, for the most part eat too much food, and would be the better for reducing their diet ; and carrying the same principle into derangement of the system and ill-health, we may safely lay it down, that the greater number of complaints may be benefited, if not cured, by abstinence. But physicians have to study exceptions as well as rules : if the latter are the first learnt, and the most obvious, the discrimination of the former, or the distinguishing those cases to which common rules do not apply, is the more difficult part of medicine and surgery. Or, deviations from common rules depend upon other rules of more partial application, which have to be sought when the first are known.

So, in the present case, it is true that digestion is an effort, the toil of an organ, which exhausts its powers, and sympathetically lowers for a time the tone of the system ; that a full meal, to a certain

extent, oppresses the vital economy, which, so overborne, only recovers its forces, after the up-hill labour of digestion has been accomplished. But it is not less true, that food taken in moderation, and when wanted, independently of the nutriment it contains, and before that is conveyed into the circulation, and while digestion is beginning and in progress, directly and immediately strengthens and invigorates.

To give an example,—a very learned and distinguished person told me, in a conversation on the subject of digestion, that his brother, a middle-aged man, not deficient in muscular strength, and leading a country life, is compelled to take his breakfast immediately on rising. If, instead of this, he begins the day with exercise or study, he becomes speedily exhausted, and remains so for the rest of the day. My informant had been struck with his brother's idiosyncrasy, as, in his own case, things are just reversed. To enable *him* to go through his habitual daily mental and bodily exertion, it is necessary that he should eat nothing for three or four hours after rising. His stomach, it appears, does not wake as soon as his brain: that languor of digestion, which sleep induces, is not with him shaken off till some time after rising.

His brother's case, to which I am acquainted with many parallel, is pertinent to my present argument. To mention another:—a gentleman, aged fifty-four, has been under my care, who, with a

pulse which varies from twenty-one to twenty-nine beats in a minute, is in perfect health, with this drawback,—that he has had two epileptic fits, one severe, the other transient, and several threatenings. The worry of business in which he is engaged, the mental excitement and exhaustion consequent upon it, are attended in him with a sensation that he will lose himself, which stimulants and quiet, when at the worst, have removed. In his ordinary habits, this gentleman does not feel himself secure (and as he has learned by experience with reason), unless he takes a cup of tea and a slice of bread and butter in bed before rising, his hour of rising being seven, and his subsequent breakfast-hour nine. After more excitement of business than usual the day before, he adds some brandy to his dose of tea, of the necessity of which his feelings of greater depression and physical alarm on waking now advertise him. Such a case, in which the tone of the heart and of the brain have been singularly lowered, conveys, indeed, a great deal more than I have advanced it to illustrate, but it strikingly exemplifies the principle, that food may be used, in fitting circumstances, as a direct stimulant. (1837.)

This gentleman is now (1840) in stronger and firmer health; his pulse has risen to thirty to thirty-four beats in a minute. A full narrative of his case, which combines several points of interest, is given in a paper on *Instances of slow pulse*,

read before the College of Physicians, and published in the *Medical Gazette* for June, 1838. The causes of slowness of the pulse are very various, but one, unquestionably, is debility arising from insufficient nutrition. In one of the cases quoted in the paper referred to, that of a young lady, to whom no nourishment was given by the mouth for six weeks, the pulse in that time fell to thirty. In a young gentleman who was nourished by injections alone for five days, after the operation of uniting a fissure of the soft palate by suture, the pulse fell in frequency, and became irregular on the fifth day. Another young gentleman, in whose case I was more recently consulted, with a pulse of forty, was thin, delicate, nervous, and alarmed about his health; he went to the sea-side, where taking tonics, and wine and ale, and nourishing food, he became stronger and better; and his pulse has risen to nearly sixty.

The following instance exemplifies the use of food as a stimulant in another habit of body. A gentleman, aged sixty, tall, large in person, who has had two or three threatenings of an apoplectic seizure (for which I have attended him, and which have been averted by prompt and copious depletion, by cupping and aperient medicine), is in the habit of eating, not in excess, but heartily, at breakfast and dinner. But he finds these two nutritious meals insufficient, and after being engaged in business in the fore part of the day, is liable to sensa-

tions of sinking, alternating with palpitation of the heart, unless he eats a meat luncheon, and drinks with it a glass of porter. I am sure that this gentleman could not preserve his health without this third meal; yet his system is full and plethoric, although, at the same time, inclining to relaxation. A person of greater physical vigour would not feel the want of a third meal; a spare person, not of the first strength, would want it, and have no doubt about, and obviously incur no risk in taking it. (1837.) [In the autumn of 1838, this gentleman was attacked with dyspnœa and other symptoms of heart disease, of which he died after a few months' illness.]

It is a rule of dietetics not to load the stomach immediately before sleep. But this has likewise occasionally to yield to the principle I am now illustrating, the use of food as a stimulant.

A gentleman, about forty years of age, when living, in point of diet, as he thought, very carefully, exceeded his strength by superadding literary labour to professional business; at length, night after night, he found himself, on going to bed, chilly, disposed to shiver, his pulse like a thread, with an unpleasant tendency to lose himself, and a feeling as if he would faint, joined with vertigo. These sensations he at first relieved by drinking hot water and sal volatile, and hot water and brandy, upon which the action of the heart became

stronger, a glow came upon the skin, and he would get to sleep, liable, however, to be broken by a return of the same feelings. I advised this gentleman to eat a light supper regularly an hour before going to bed, and to drink three glasses of sherry; when the unpleasant feelings which he had up to that time experienced, left him. Those who have mental exertion to go through in the evening, are unfitted for it by a hearty dinner, while from a light one they may not derive support enough. To such persons a moderate dinner and a light supper are the proper substitute for a single nutritious meal. These rules will not, I trust, appear trivial. The relief gained by attending to them is immense; and though they seem obvious, yet they do not occur to every one by whom they are needed.

The following case exemplifies more generally the injurious effects which may arise from neglect of adequate sustenance.

I had occasion to attend an artist under these circumstances. A few days before, he had suddenly dropped down in his studio, and lost his senses for a few seconds; since then he had been threatened with another such seizure. When I saw him he was pale, his countenance expressive of alarm, his pulse feeble; he had come in from a walk, and was apprehensive of another fainting fit. Upon drinking, at my desire, some brandy and

water, he recovered ; his colour returned, and the faintness wore off. I learned that this gentleman had literally starved himself ; that he entertained an aversion from animal food, except in the form of soup, that he lived upon puddings and vegetables. He had indulged this humour so far, that the appearance of the meat upon his table when his family dined, produced sensations amounting to loathing and nausea. Nevertheless, it was evident that animal food was now necessary for him, as the means of restoring his strength ; and as I found that he would have less reluctance to attempt to take it at an unusual hour, I desired him to try a beefsteak at luncheon, and again at supper, giving up his six o'clock dinner altogether. I likewise prescribed some tonic medicine. Pursuing this plan, he lost his faintnesses, and gained strength.

Every one knows that exhaustion and want of sustenance will produce faintness, dizziness, headach. But unquestionably, in individual cases, the source of these symptoms is often overlooked. It is liable to be overlooked as soon as apprehension of fulness of the cerebral vessels, and of determination of blood to the head, gets possession of the mind of the practitioner. And while overlooked, the treatment adopted on the opposite view is often producing the very affections it is intended to prevent, and the patient's health and constitution are undergoing serious injury. *That* faintness and tendency to

sudden unconsciousness, which great reduction of strength and exhaustion produces, may thus be driven into epilepsy; and vertigo and headach established as habitual ailments. The reader will allow me to exemplify these important considerations by narrating two cases.

I attended a gentleman in his eightieth year, for a local complaint, who formerly being in trade, fell into illness of the following kind. At the age of five-and-thirty, being of a frame slight but strongly built, he was at once living heartily, and confined constantly by business to the house. Having been for some months sensibly out of health through these causes, he was one day taken with weakness of the right leg, and inability to articulate; in an hour his speech returned, but his right leg continued weak. He was now cupped once, and purged repeatedly; after six weeks, he was sent to Bath, where he was again cupped, purged, and restricted to an innutritious diet and water. He returned to town after two months, much reduced in strength. Almost immediately upon his arrival in London, he had an epileptic seizure, and in a day or two a second, and quickly a third. Altogether he had about three hundred fits in the ensuing two years and a half; by which time they ceased, and have never recurred. Shortly after their commencement, his diet was changed, he was allowed meat and wine; with this his



strength improved; and with the improvement in his strength, the tendency to epilepsy lessened. During his recovery, the occurrence of anything which exhausted and lowered him was sure to bring on a fit\*.

It is to be observed of *epileptic* seizures in general, that they more frequently depend upon exhaustion and want of proper support and stimulus, than upon the contrary class of causes. They are likewise, *it is not intended to be denied*, liable to proceed from other causes, from disease of the cranium, from pressure on or alteration of the structure of the brain, from increased cerebral vascularity; the last supposition I believe to be correct, from instances which I have witnessed of epilepsy in persons of a

\* This gentleman gave me the following account of the sensations with which the fits began. There was first a peculiar motion along his right leg, which quickly seemed to mount to the pit of his stomach; from thence it rushed to his head, and he fell senseless. The right leg was the leg which had been weak at the commencement of the attack; and at this time he always walked upon it with a slight halt, not bending the knee. If he tried to walk with his right leg in the usual way, the instant that he lifted it from the ground bending the hip and knee, the peculiar sensation forerunning the epileptic seizure commenced in the leg, and he fell unconscious. On the other hand, when this sensation was beginning, he could sometimes defeat the attack by pressing with his hands above the knee, and forcing the limb strongly against the ground. The limb would then for a short time be shaken by muscular action, and then be tranquil, and the fit be averted.

full habit, attended with fixed local pain in the head; in which the complaint has been benefited by purging and bleeding. But, certainly, epilepsy most frequently arises from debilitating causes, and in a lowered habit of body; so is it often threatened in persons who have sensibly gone beyond their strength, the slight giddiness and faintness experienced by such persons being of this nature. These threatenings may often be removed by stimulants; otherwise they may go on to strong convulsive seizures and total unconsciousness. Epilepsy of this kind may be viewed as cerebral fainting.

The next case is from Dr. Abercrombie. "A lady, aged twenty-five, had been frequently bled on account of symptoms in the head which had supervened on an injury. Considerable relief had followed each bleeding; but the symptoms had soon returned, so as to lead to a repetition of the bleeding at short intervals, and this had been going on for several months. When Dr. Abercrombie saw her, she was stretched upon a couch, her face of the most deathlike paleness, or rather of the paleness of a stucco figure, her pulse very rapid, and as small as a thread, her general weakness extreme. The mass of blood appeared to be reduced to the lowest point that was compatible with life; but she still complained of frequent headach, violent throbbing of the head, confusion and giddiness. It was evident that evacuations could be carried no further;

and on consultation with a very intelligent man, who had the charge of her, it was agreed, as a last experiment, to make trial of the opposite system, *nourishing diet* and tonics. In a fortnight she was restored to very tolerable health."

There is another ground, on which nutritive diet occasionally becomes requisite in the management of persons disposed to apoplexy and palsy. I shall afterwards have to notice the fact of the production of head-symptoms by indigestion, or their occasional occurrence as sympathetic effects of disorder of the stomach. When the stomach is weak, it is sometimes incapable of chymifying the less nourishing kinds of diet; and vegetable and farinaceous food cause flatulence, heart-burn, and distension, that are attended with threatenings of some head-attack. In this class of cases, reduction of the circulation and the avoidance of everything heating, strengthening, or stimulating, may be to be mainly studied; nevertheless a diet containing animal food may be necessary. A gentleman recently consulted me to decide this question for him: one had bid him live on farinaceous food alone; while another told him to eat meat; he wanted a third opinion to turn the scale. He is between sixty and seventy years of age; two years ago, being as now engaged in extensive concerns in business, after several weeks, during which he had observed that his memory of names was impaired,

he was seized with palsy of one side of the body ; for this he was bled, and he gradually got better : his mind entirely recovered. He does not appear to have been, after this seizure, careful in diet ; temperate he was, but he ate hastily, and indifferently of any kind of food ; then he would suffer from indigestion ; and he learnt to know that he had eaten what disagreed with him, by the extreme bodily weakness and failure of muscular power which it caused. Living in this manner, he was taken with apoplexy, and while lying nearly insensible, he vomited with violence a large quantity of intensely sour liquid ; he was freely bled, and recovered in a few weeks ; and he has since had another similar attack. Now he is perfectly well, with the exception of the weakness of the leg and arm left from the first palsy ; and a sense of uneasiness in the head rising to pain through the temples, that is brought on by any sudden bodily movement, or if his sight is distracted by many objects simultaneously, or if his stomach is the least disordered. Otherwise he is fully capable of attending to the details of complicated business, and does not suffer from intellectual exertion. Indigestion with him was marked by distension of the stomach and flatulence ; eating potatoes has of late invariably produced such an indigestion, and the attendant pain in the head ; cauliflower, peas, and other vegetables, he can eat without ill consequences ; of animal food,

boiled salt meat alone sensibly disagrees with him, or disturbs his digestion. I recommended the adoption of a varied diet, of which tender animal food should form a part.

In both of the apoplectic attacks which this patient has suffered, it has been mentioned that he has been freely bled; and in both, the apoplectic coma became deeper after the abstraction of blood. It is not impossible that both the attacks depended upon indigestion, and that remedies appropriate to the latter, and rest, were the treatment principally necessary. This patient, however, without being corpulent, has been rather disposed to fulness of habit, and the vessels upon the temples are full and prominent; so that to a certain extent depletion may have been necessary with him.

I feel that I have to guard against the preceding observations being misinterpreted. For it is certain that in a large proportion of cases, apoplexy, palsy, epilepsy, are the result of increased vascular action in the head, and that a lowered diet, local and general depletion, and other means of reducing or diverting action, are alone advisable, and that no other plan can be safely substituted for them, or is to be thought of. Only it is most important constantly to bear in mind, in connexion with the preceding principle, that head-symptoms may actually begin in weakness, and from the first admit of relief by strengthening means only; that

(which is a case still more liable to be overlooked,) head-symptoms which began in fulness and local action, and have been for a time benefited by reduction, after a period may be kept up by the very means which at first were salutary, and may require contrary measures for their final removal. And that where there is no want of power, and the habit even disposed to plethora, head-attacks may supervene dependent upon other causes than vascular fulness. The variety of causes, upon which the classes of symptoms under consideration, that is to say, threatenings of loss of consciousness, may depend, is curiously extensive. As the third case, they may proceed, as will be by-and-by further illustrated, from an overloaded and deranged stomach alone; or as a fourth case, they may be products of hysteria; or as a fifth, their source may be purely mechanical.

I attended a woman, in whom I had reason to believe a complete obliteration of the cavity of the womb had taken place from inflammation; at all events, there was suppression of the periodical relief. But she was regularly seized with attacks of pain in the head, coldness, numbness, and weakness of one side of the body, which would last three or four days. At the worst, she had a strong epileptic seizure in addition.

The late Dr. Sweatman told me that he once attended a lady who had been compelled to remain

in the horizontal posture for ten years. If she sat up, in a short time she fell into, or began to fall into deliquium, and fainted. But she looked in perfect health, and had borne several children during this period. The opinions of several eminent London physicians had been taken, who, finding nothing wrong in the circulation, had all recommended tonics. Dr. Sweatman begged to see the patient's legs, which he found covered with immensely dilated veins. Into these, when she stood up, all the blood in her body gravitated; and hence she became faint. The legs being bandaged, she was cured at once.

There are, of course, infinite other disorders besides those which manifest themselves by head-symptoms, in which lowering diet and reduction would be prejudicial. It is to be laid down as a truth of universal application in medicine and surgery, that disturbance of the health of the entire body, or of a part, general or specific, takes its most important character in a practical point of view from the strength or weakness of the system attacked, and is most efficiently controlled by remedies calculated in reference to these elements. And it probably may be assumed that the greater number of complaints, especially when beginning in persons of good stamina, have febrile and inflammatory features, or a sthenic cast; whence it follows, that at the commencement of most dis-

orders, a spare diet, and other means of reducing the strength are to be employed. Nevertheless, a large proportion of ailments of every description are found in connexion with a lowered habit and exhausted bodily forces; in which the contrary practice is required. It is, perhaps, less the disease than the constitution and age of the patient that has most to be considered in the plan of treatment. Therefore it does not follow because a vital organ is the seat of organic disease, that the patient is necessarily to be reduced in strength. When the structure of the heart, or of the lungs, or the brain, is materially deranged, it is often necessary to use means to strengthen, and recruit, and stimulate; the weakened organ and system may sometimes even require to be supported at the risk of increasing those diseased conditions of the former.

There are no cases, to the successful issue of which the management of diet more essentially conduces, than those in which capital surgical operations are performed. Here again the general rule is, light food, and systematic depletory preparation. A patient somewhat reduced in strength is in a more favourable state to recover from a grievous bodily injury, than one in rude health. And after a great operation, the first danger to be thought of is that of fever and inflammation. But danger of the opposite character has often to be provided against. In persons of a scro-



fulous habit who suffer amputation, in all the young and delicate, in middle-aged persons worn by long disease, in persons advanced in life, who are compelled to undergo surgical operations, the chance of ultimate recovery may be much increased by the strength of the patient being recruited and fostered through a more generous diet before the operation, and supported by the same means afterwards. I remember hearing narrated the case of a gentleman who had an ambiguous local affection, which at last was thought to be cancerous. He was between fifty and sixty years of age. Every means were tried to disperse the tumour before an operation was recommended; and the last resorted to were strict confinement to the horizontal posture, and a course of mercury. These having failed, the operation, not a very serious one, was now resorted to in his debilitated habit. In a few days erysipelas and nervous irritation supervened, and the patient died, who probably would not have died had his strength been previously recruited by liberal diet, exercise, and the like.

Sometimes abstinence from food is a *mechanical* condition of the success of a surgical operation. This is the case after the operation for congenital fissure of the palate. It is here desirable that the patient should not swallow for the ensuing four or five days; he may however be nourished by enemata of strong broth and beef tea. Preparatorily,

again, in such a case, it is desirable that he should be in strong health and excellent condition of body.

In the observations which I have thus far made, I have considered the regulation of diet in reference rather to the whole system, or to other organs than the stomach; and have supposed no fault in the digestive powers. Let us now look to the cases in which management of the stomach is wanted to disembarass it of unwholesome or indigestible food, or to restore it, when below its proper tone, to health and strength.

Under different circumstances, the stomach presents every degree of failure, from capricious and uncertain digestion, to complete inability to chymify food. Except in the last case, the evil generally admits either of being corrected, or of being greatly mitigated. The stomach may be weak, but a diet suited to it will be digested without difficulty or suffering; as on the other hand, food in excess, or of an unwholesome quality, will disagree with the strongest stomach. When the proper ratio is not observed between the strength of the stomach and the quantity and quality of the food, indigestion arises; or an indigestion may proceed from an accidental cause, and be the complaint of a day, or it may proceed from error in the general habits, or weakness of the organ, and amount to established indisposition. By the term indigestion, the latter state of things is commonly meant.

In health, all that we ought to know of our digestion through sensation is, that satiety has succeeded hunger, and that for a shorter or longer period we are indisposed to exertion. When digestion is imperfect, many sensations, local and general, occur to distress and disturb us. The stomach may be the seat of pain, and of feelings of weight and distension; nausea and loathing of food may take the place of appetite, and retching and vomiting may ensue; the brain and nerves may be disordered, depression of spirits, headach, confusion of thought, disturbed vision, vertigo, fits of insensibility, may follow; the action of the heart may be irregular; the breathing oppressed and embarrassed, with constant troublesome cough.

The condition of the stomach itself, during these distressing symptoms, may be different in cases where the general features are much alike; and it is remarkable that the extent and ultimate importance of the changes in its structure and organization bear no proportion to the sufferings which the patient experiences. Generally speaking, the most grave diseases of the stomach, except towards their close, produce much less physical distress than the lighter.

All the symptoms now to be considered arise from disorder of the functions of the stomach, and depend directly upon its inability to digest, upon

food lying unchanged in the stomach, upon failure or depravation of its secretion, upon exaltation of its natural sensibility; and this functional failure or disturbance may co-exist with no visible alteration of structure, or at all events may depend upon alterations that are so transient as to disappear in death. Ulceration of the stomach, thickening of the stomach, cancer of the stomach, produce more or fewer of the symptoms which I have enumerated; but they produce them secondarily; they produce them when they at length cause in the stomach at times or constantly that functional failure, upon which indigestion depends.

The reader is to understand my object to be to describe the features and treatment of indigestion in the popular acceptation of the term, that is to say, of disorder of the stomach, not disease. Nevertheless I shall be obliged to notice the points at which the two touch, or where the persisting symptoms of indigestion indicate altered structure. I may be permitted to observe that in my *Outlines of Pathology*, I have endeavoured to arrange and classify the diseases of parts, in which determinable lesion of structure exists. To that volume I would refer the reader who wishes to go deeper into the subject of diseases of the stomach, than I have here entered.

The term indigestion, as it has been remarked, is applied to two things: either to a single attack

of disorder of the stomach lasting a few hours or days; or to confirmed and habitual, or frequently-recurring disorder. To commence with exemplifying the first, I may give an example of a common accidental attack of indigestion. A gentleman sent to me, who, having dined with moderation, and drunk three or four glasses of wine, had at twelve retired to bed without any feeling of indisposition. He awoke at one, with nausea and pain at the stomach; he then vomited, but without relief of the pain: the bowels then acted several times. When I saw him about six in the morning, he complained of a dull heavy pain at the epigastrium, pressure upon which gave him great uneasiness: his tongue was a little furred, but he had no pain, nor was there general tenderness of the belly. I considered the pain to arise from undigested food, and therefore mixed for him two tumblers of warm water and salt, which he swallowed and retained in the stomach a few minutes, and then rejected, along with a quantity of half-digested food, containing some leaves of water-cress very little changed. He felt relieved, slept, awoke at ten free from uneasiness, took an aperient draught, and by the afternoon was perfectly well.

To produce such an indigestion, it hardly seems necessary that anything positively unwholesome should be taken. Fish, shell-fish, or raw vegetables, however, often seem to be the food at which the

stomach takes offence. Yet the party may have been in the habit of eating the same kind of food without its having disagreed before; so the fault in such cases commonly is in the stomach, which is accidentally out of order, and then the least digestible part of an ordinary meal resists digestion, and becomes an irritant.

So, when, as in the case described, uneasiness at the stomach and nausea supervene shortly after a meal, it is desirable at once to bring on, or to encourage vomiting. For this purpose, half a teaspoonful of common salt in a tumbler of tepid water, followed by a second within a few minutes, if it be necessary, will generally answer. This remedy has the following advantage: if great part of the offending matter has been thrown up already, the salt and water often does not produce further vomiting, but at once calms the stomach, and subsequently acts as an aperient. If the matter thrown off the stomach is acid, the patient, as soon as the stomach is quiet, should take ten to thirty grains of carbonate of soda in a wine-glass of water, not at once, but by two or three mouthfuls. After a few hours' sleep, a breakfast-cup of arrowroot and milk is one of the best things that can be taken, being at once light, digestible, and nutritious.

It is, however, important to remark, that vomiting is not a remedy to be prescribed indiscriminately. A vomit agrees with good stomachs, and with

young people (so children are the better for emetics at the commencement of most disorders). But persons of weak stamina, or of a weak digestion, are shaken to pieces by vomiting, and suffer for some days afterwards. After the middle period of life, again, the effort of vomiting, generally violent, is extremely distressing. There are some, again, with remarkably strong stomachs, to whose idiosyncrasy this process is abhorrent; and who can bear the roughest discipline of medicine rather than the strain and effort of retching.

The ordinary consequences of neglecting such an attack of indigestion as has formed the theme of these remarks, are, either a smart attack of purging brought on seemingly by undigested food reaching the bowels, or several days' indisposition, impaired appetite, heartburn, furred tongue, and general uneasiness, till the food, either remaining in the stomach has been at length digested, or has gradually passed by the bowels, or has been helped away by purgative medicine. But sometimes the undigested food remains for days without altering its place, or the symptoms changing; and it becomes necessary, though late, to discharge it by vomiting, for fear of some serious illness.

Lieut. Gen.—, aged sixty-eight, consulted me for the following symptoms. Five days previously, after a repast, at which he had eaten not immoderately, but indiscriminately, he had been taken in

the night with uneasiness of the stomach, attended with nausea, but he had not vomited. The pain at the stomach continued through the following day, when he took some aperient medicine, which moved the bowels, but gave him no further relief: the uneasiness at the stomach remained. He then determined to disregard it, expecting that it would wear off. But this plan not succeeding, on the fifth day he thought the disorder too serious to be neglected; day and night there had been a wearying uneasiness of the stomach: the tongue was furred, the appetite and relish of food gone; yet he took food at his regular meals, which he digested. The pain was distinctly seated in the stomach, which was uncomfortable if pressed: the seat of the pain had not varied from the first. I made no doubt that some part of the dinner of the preceding Sunday remained undigested in his stomach. Accordingly I prescribed an emetic of a grain of tartarized antimony, and a scruple of ipecacuanha; the vomiting which followed, was violent and distressing; but when the stomach had been emptied by it, the pain was gone.

To exemplify this case,—undigested food in the stomach,—in its extreme results.—A gentleman approaching seventy years of age, and living much in convivial society, habitually ate and drank to excess. He had had one or two warnings, consisting in attacks of giddiness and threatenings of a



fit, which, however, had passed off and were forgotten. One night, he awoke with shivering and uneasiness at the stomach; which, upon the application of hot flannels, and on his taking stimulants, subsided; but he was ailing the next day; then for ten days he went into society as usual, but he lived more moderately. I knew him; but he had not consulted me. But I had called to ask after his health: when on his way to dine at the — Club, he left a card at my house, with these words in pencil, "Thank God, my dear Mayo, I am well again." I learned that at his club he ate, drank, and sang, but was observed to look ill, and left the table earlier than usual. He went home, was taken worse, and died towards morning. An examination of the body being made, the stomach was found enormously distended with the meals of many days hardly altered. An emetic administered a week before, and abstinence afterwards, would have saved his life.

A common form of temporary indigestion, or disorder of stomach, results from bile, and is closely allied to the preceding instances. It often goes with them, and the same causes which temporarily weaken digestion, produce bile. Excess in diet, unwholesome food, exposure to cold and wet, with some the mere change of the wind to east or north-east, these physical causes, and any mental disturbance, are liable to cause bile to be poured

into the duodenum in large quantities, from whence it regurgitates into the stomach.

Bile in the stomach causes headach, sometimes attended with impairment of sight, either dimness or partial vision, with loss of appetite, &c., and often ends in nausea, with a bitter taste in the mouth, vomiting of bile, either pure or mixed with ingesta, or diluted with mucus.

Persons of the bilious temperament are liable to slight attacks of this description from slight causes. Their skin, and the white of the eye, are then tinged yellow; they experience languor, loss of appetite, and severe headach. Quiet, abstinence, and brisk cathartic medicine, accelerate the departure of the attack, which is disposed to last from half a day to two or three days.

I have known the affection of vision in bilious attacks take three forms. A gentleman, aged twenty-six, who was under my care for a protracted constitutional disease, told me that he was liable, when bilious, to vertigo, attended with a dark shade over the centre of the field of vision gradually diminishing towards the circumference; so that he could not see what he looked at, but only things around, and that dimly. Such an attack might last half an hour. On going off, pain in the head, and sometimes nausea, would supervene. Another gentleman told me that he had, when suffering bilious attacks, often experienced general failure of

vision, so that he could only distinguish that part of a scene to which his eye was immediately turned, all around being obscured by a brown fog. The same gentleman told me that, at other times, he had semivision of objects, sometimes seeing the right half only of objects with both eyes, while in other attacks, he had seen the left half alone. These visual eclipses would last two or three hours, being at their worst about half an hour.

The most acute and severe attacks of bile on the stomach are characterized by bilious vomiting, with pain at the præcordia, coldness, and great depression of strength. At first, the stomach is so irritable, that any attempt to allay the vomiting by medicine only aggravates the complaint: the least excitement of the organ causes more bile to flow into it. Perfect quiet and warmth are the best remedies, joined with the application to the pit of the stomach of equal parts of camphor-liniment and laudanum upon hot flannel. After a little, the stomach becomes tranquillized; and there are several remedies, which in different cases, admit of being then beneficially used. These are, a small quantity,—an ounce, for example,—of hot brandy and water, with ten or fifteen drops of laudanum, or a wine-glass of soda-water, with a tea-spoonful or dessert-spoonful of brandy, or soda-water with seven or ten grains of carbonate of soda in it, or a wine-glass of hot water, with twenty drops of sal-

volatile, combined or not with laudanum. The preference of either of these remedies in any case must be left to the tact of the practitioner; only they are not inert, and where one will agree and do good, the others will often do harm. With those in whom laudanum generally produces head-ach, it had better not be tried at first; when the patient fancies a preference for hot liquid, or the reverse, that indication should be attended to. As general rules, carbonic acid is the wholesomest restorative in such cases; while any quantity of any liquid is sure to re-excite vomiting. Those who have an habitual aversion from anything cold, women, for instance, in whom spasm of the stomach is so produced, yet when rallying from a bilious stomach attack, are often revived and restored to comfortable feeling by small draughts of cold soda water, which in health would disagree with them.

It is sometimes useful in such cases, at their commencement, and when the retching is very violent, and pure bile is being thrown off, to wash out the stomach by a large draught of bland mucilaginous fluid, as tepid barley-water, or by very weak chamomile tea; but in general, this is bad practice, and only protracts the disposition to vomit.

As soon as the stomach is perfectly tranquil, it is desirable to administer a laxative dose, which

carries off the bile from the bowels, and helps to set the system to rights. The infusion of senna with equal parts of the compound decoction of aloës, with a drachm of some neutral salt, and of tincture of senna, and of the compound tincture of cardamoms, will answer this purpose.

A young gentleman lately consulted me, in whom the severe cold of the spring [1837], and the adoption of more sedentary habits than he had been accustomed to, had produced a most troublesome tendency to bile upon the stomach. Abstemiousness, moderation in exercise, abstinence from wine and fermented liquors of all kinds, with doses of aperient medicine twice a week, kept him in health. The least deviation from this strict rule, and headach, sickness, and vomiting of bile at the worst, supervened. Calomel or blue pill added to his purgative dose, invariably made him worse, and had to be scrupulously avoided.

In general, those who suffer from bile, are the better for some form of mercury, followed by an aperient; a fact which seems at variance with the phenomena of the preceding case; but it is not really so. Bilious people are those who have a stronger disposition than others to form bile. In all, but particularly in bilious people, preparations of mercury increase the quantity of bile they form. Now, if the disposition is not great, the bile formed by the mercury is all carried off by the subsequent

purging, and the system is completely cleared of bile for the time. But if the disposition to form bile is greater, anything that heightens it only does greater mischief; the bile is formed there in such quantities that the blood cannot be cleared of it; and the only safety and comfort of the patients consists in adopting the rules which are calculated to prevent the redundancy of the humour on which their habitual indispositions depend.

Habitual indigestion arises from weakness of stomach, either original, or brought on by habitual incaution in diet, or by a sedentary life and neglect of exercise, or by inattention to the bowels, or by residence in hot climates, or by mental causes.

The following case, in which the weakness of stomach appears to have existed from birth, exemplifies many of the ordinary features of dyspepsia, and some of its less common and remote consequences.

Mr. H. R., aged sixty-four, from a boy has had an appetite always moderate, when his digestion has been at the best; but that has been repeatedly disordered in attacks which have lasted several weeks, coming on and leaving him gradually. During these attacks, he has been well and free from all uneasiness, as long as the stomach has been empty; and the immediate effect of a meal has been to produce comfortable bodily feelings; and these, indeed, after a meal, he could prolong

by continuing to excite the stomach by new ingesta; so by drinking wine, and then hot tea, after dinner, he could put off the distress and suffering which would attend the effort at digestion. That suffering consisted in severe pain of the stomach, with a sense of weight and distention, often with heart-burn. These symptoms, as the consequences of his breakfast, would come on about eleven in the forenoon. The same would supervene at seven or eight in the evening, unless put off by the use of stimulants, which seemed at the same time to retard the commencement of digestion. In that case he would go to bed at eleven, but wake after two or three hours' sleep in great suffering. The extent of the pain depended upon the inability of the stomach to digest; sometimes digestion was already half gone through, and interrupted by heartburn only, when an alkaline effervescent draught would quiet the stomach, and digestion be completed. At other times, this means giving no relief, he was compelled, for present ease, to excite vomiting mechanically, when the meal of the preceding evening would be thrown up nearly unchanged; the pain would then cease, and he would sleep and wake in good health.

This gentleman had, at different times, made trial of almost every remedy. Tonics, from the mildest to the strongest, from chamomile tea to quinine, he had found either nugatory or prejudicial.

Two remedies alone had benefited him ; these were just enough of the compound extract of colocynth taken in the afternoon to keep the bowels open ; the other a few grains of magnesia in an effervescing mixture, to relieve flatulence and heartburn. On one occasion, he had thought himself the better for the application of a mustard poultice to the pit of the stomach. His food, when his stomach was the most disordered, consisted of gruel, or soup and bread, or boiled whiting, and he took no wine. As he recovered from each attack, the stomach would be extremely capricious, and one day he would be better for and digest nourishing stimulant food, and then be compelled again to resort to the simplest for a few days.

Living upon these conditions, in the autumn of 1835 the following change took place. Of a sudden, the action of the kidneys was remarkably increased (a symptom, in elderly persons, always pregnant with alarm); in a week, the ancles became swollen and œdematous; in another week, the increased action of the kidneys subsided, but the legs continued to fill with water more rapidly than at first. At the same time, the digestion, which had been very much disordered before the invasion of these new symptoms, became stronger and better than it had been for a considerable period. The œdema of the legs however increased, and the belly began to enlarge. Towards Christ-



mas, the dropsy, which had declared itself, and the swelling of the legs, rendered it difficult for him to move up and down stairs. He had become thin and extenuated in the upper part of his person ; his countenance pinched and yellow ; his appetite, too, now failed him, and his digestion wholly. At this time, he put himself under my care. On examining the belly, which contained a few pints only of fluid, I found, at the upper part, and rather to the right, that is to say, over the place of the lesser or right portion of the stomach, a diffused hardness or boardiness, with tenderness on pressure. I therefore applied eight leeches over this region ; he became faint with the loss of blood, which was thin and serous. The following day I applied a blister over the same surface, and recommended him to take small doses of blue pill and compound extract of colocynth daily, and to live on gruel ; when the blister had healed, the tenderness had gone ; at the same time his countenance had improved, and his appetite. It was my impression that the stomach, at that time, was the seat of chronic inflammation, which the remedies resorted to seemed to have removed or mitigated. Nevertheless, the water went on accumulating in the legs and belly, and the action of the kidneys was less than natural. Yet his strength improved with his digestion. His pulse, though weak, was regular, the action of the heart was, in every respect,

normal; he could sleep on either side, or on his back, and with a single pillow. There was no albumen in the urine. I thought that what was now wanted was the restoration of the action of the kidneys; to produce this effect, he therefore took, but ineffectually, the decoction of broom-tops with the acetate of potass, then the infusion of taraxacum. I then gave him extract of elaterium, combined with blue pill; this acted and forced the kidneys, and diminished the water, but, at the same time, it purged and weakened him, and disordering his stomach, made him, in other respects, worse, and threw him back. In a short time, the tone of the stomach rallied, but the water increased again. There was more dropsy, the œdematous swelling spread up the body, and in the morning his face and eyelids were puffed with fluid. I now determined to give this patient mechanical relief at all events, and accordingly I punctured his legs with a grooved needle, in twenty or thirty points each; the water flowed plentifully, and continued to exude for a week, then ceased, leaving the legs very much reduced in size, and the dropsy certainly lessened. He felt, however, more weak than before, but his digestion was tolerable. What practice was now to be pursued? It was the opinion of two other medical men, who saw him at that time, that blue pill, with squill and digitalis, should now be tried, (the

more likely now to move the kidneys, from the mechanical relief already obtained.) To myself, another opinion appeared preferable; it was, that steel might now be given with advantage; and this I recommended to his adoption, influenced by the following considerations. There were two plausible views of the complaint; the first, and certainly, at the time, the most probable, was that the real complaint was some alteration of structure about the stomach (thickening of the coats of the stomach, perhaps, with induration and enlargement of the lumbar lymphatic glands, or pancreas, or both,) causing pressure on the great veins at the upper part of the abdomen; this supposition accounted for all the symptoms of the case, (especially the co-existence of dropsy and anasarca, without disease of the kidneys and heart.) The second supposition was, that the dropsy proceeded from a feeble circulation and imperfect nutrition, a digestion disordered for half a century, and a thin blood. Now, on the first supposition, the case was incurable; on the second, it perhaps admitted of relief. But the course recommended, of mercury, and squills, and digitalis, was directed to the first alternative, at once to excite the kidneys, and to resolve any inward thickening, (in which process some additional weakening was involved;) steel, on the other hand, was calculated to fulfil the second indication, it might strengthen. The event

showed, as far as such an instance can show, that the choice of the latter course was the right one. The patient took two drachms of steel wine, at first twice a day, afterwards three times a day; in two or three days, the kidneys began to act, and in a few weeks the legs and belly were emptied of water. The patient's strength, in the meantime, slowly increased; and after several fluctuations during the last two years, between ill health and convalescence, the patient is now in the enjoyment of renovated health, and a better digestion than he has generally possessed. Two things deserve to be added. He often finds great benefit from taking at night an effervescing saline with a few drops of laudanum. And the winter before last [1838-9] having fallen and broken his arm, by which accident he was confined to his bed a few days, and to his room six weeks, he seemed to recover a new power of digestion during this confinement, and he has been better ever since.

I shall now, as the most convenient mode of unfolding what has to be laid before the reader, proceed to consider, first the remedies by which indigestion may be alleviated or mitigated; secondly, some prominent features, which give their character to individual cases.

The remedies of use in the cure of indigestion, are either general or special. The former class, as liable to require employment in every case of in-

digestion, I will mention first and separately: the latter may be appropriately described in connexion with the peculiar and special features which the disease often assumes, and to the relief of which they are individually applicable.

The general remedies for indigestion are, regulation of diet, exercise of the mind and body, especially if combined with amusement, aperient and tonic medicines.

The rules of diet for weak stomachs flow directly from the principles already explained. For instance, when at the feeblest, for the principal meal the stomach will bear the lightest food only,—arrowroot, or gruel, or soup with bread, or the most digestible fish simply boiled; when stronger, it may digest game, while as yet incapable of digesting poultry or butchers' meat; afterwards wine, which, in the weaker states, heats and irritates, strengthens and provokes digestion: in each stage, pastry, highly-seasoned dishes, rich food, are to be avoided; vegetables, pastry, and fruit, to be avoided, or most sparingly partaken of. At first the food is to be light, little more than digestible; then digestible and nutritious; afterwards it may combine a stimulating quality besides. But there are cases which go beyond common rules: in some, vegetable diet is the best; in others, the smallest quantities of a single kind of food often repeated; in others nutritive food, but one meal only partaken

of in the twenty-four hours. The following are instances in point.

A gentleman (Dr. Abercrombie mentions) accustomed to moderate, but very comfortable living, had been for many years a martyr to stomach complaints, seldom a day passing in which he did not suffer greatly from pain in the stomach, with flatulence, acidity, and the usual train of dyspeptic symptoms, and in particular he could not touch a bit of vegetable food, without suffering from it severely. He had gone on in this manner for years, when he was seized with complaints in his head, threatening apoplexy, which, after being relieved by the usual means, showed such a tendency to recur, that it has been necessary ever since to restrict him to a diet *almost wholly of vegetables*, and in very moderate quantity. Under this regimen, so different from his former manner of living, he has continued free from any recurrence of the complaints in his head, and has never been known to complain of his stomach.

A female had suffered the gravest stomach symptoms, giving rise to the belief that she laboured under cancer. She had severe pain in the stomach, when the smallest quantity of food was taken, with great tenderness on pressure, and constant vomiting occurring regularly at a certain period after eating. A variety of treatment had been employed without benefit, when Dr. Barlow determined upon trusting

entirely to regimen, by restricting her to a diet, consisting wholly of fresh-made uncompressed curd, of which she was to take a table-spoonful at a time, and to repeat it as often as she found it desirable. On this article she lived for several months, and recovered perfect health.

The gentleman whose case I have recently narrated at some length, suffered a very severe relapse after the influenza of 1837. There were several weeks, during which he found it necessary to give way to the complaint, and to nurse his power of digesting in the following way: he took a single meal only in the twenty-four hours, and that a light but nutritious dinner. This single meal his stomach could generally digest tolerably well. In the morning he took one small cup of tea, and one mouthful of bread. If he exceeded this quantity, which appeared necessary to sustain him for the day, his stomach became in an hour painful, disturbed, flatulent, and did not recover itself till the following morning; his dinner taken in the afternoon then only disordered him still more, causing him a night of pain.

The reader is to gather from these instances, which it would be tiresome to multiply further, that the worst sufferings of indigestion often admit of relief through the simplest means, which, again, are often not to be hit upon at once, but are found out in each individual case by patient study of all

its peculiarities, assisted by a thorough knowledge of the principles on which digestion depends, as well as of all the observed exceptions.

Upon the general efficacy of bodily exercise in this complaint, and (where sufficient exercise cannot be taken) of systematic friction to promote action of the skin, upon the advantage of change of occupation, mental relaxation, and amusement, change of air and scene, upon the injurious effects of fatigue and exhaustion of all kinds,—it is unnecessary to enlarge: these are points which any one's common sense and self observation may regulate.

The next means are the use of aperient medicines. Indigestion often occurs in persons, who, not taking adequate exercise, and confined to the house by business, fall into an habitually costive state of bowels. They lose their appetite; the tongue is furred, the breath heated; they have frequent headach; and experience after their meals a sense of weight, distention, and general uneasiness. In such persons, the use of opening medicine is highly beneficial. But strong purgatives are to be avoided; for though they relieve the bowels, they directly weaken and disorder the stomach. Mild aperients only are therefore to be used, mercury being combined with them when necessary. For example, a dinner-pill (two should be taken) consisting of four grains of extract of rhubarb, and half a grain of carbonate of soda, or three grains of the watery



extract of aloës, with one of mastich ; or at night equal parts of castile soap and compound extract of colocynth, of each four grains ; or the same, with two grains of scammony, or, instead, a drachm of the lenitive electuary ; or, in the morning, a drachm of Epsom salts in peppermint water, alone, or with two drachms of tincture of rhubarb ; or two drachms of Cheltenham salts, in a third of a pint of water. Not to multiply instances of the efficacy of such plans, I cite the following only as an extreme case, rendering the mention of commoner ones unnecessary.

A protracted case of stomach disorder was treated by Dr. Parry, in which there was vomiting so constant, that every thing was rejected, even a teaspoonful of water. The case had gone on in this manner for several weeks, and the patient was reduced to the last degree of emaciation, when Dr. Parry ordered half a grain of aloës to be given every four hours, moistened only with a few drops of liquid. This was retained, and acted gently on the bowels ; and, in less than two days, the complaint entirely subsided. The bowels had been freely moved during the previous treatment, and other remedies in great variety had been employed without any benefit.

It is often advantageous to combine in indigestion aperient medicine with tonics ; or, this suggestion properly applies to the specific cases where tonics

are available. It is much more common to meet with instances where aperient medicines alone are salutary, than in which tonics are advisable alone. Of tonics alone, perhaps, the nitric acid with some bitter is generally the most useful; but all are occasionally valuable: even chamomile tea (a tea-cup of a very weak infusion) taken on rising, sometimes has a sensible effect; gentian, again, columbo, the infusion of orange-peel, with five grains of carbonate of soda or ammonia, are often beneficial. So, again, there are cases, in which a dessert-spoonful of white mustard-seed, swallowed daily, has been of service. Nor are these recommendations to be disregarded, because they are empirical; and the success attending them so far accidental that it is impossible to say beforehand, which of such different remedies, or whether any, will prove of use. Take, for example, the following instance from Dr. Abercrombie; there is no question but the practice was purely tentative of combining in this case a tonic with an aperient, but it was perfectly successful.

A lady, aged about thirty, laboured under the following symptoms in the summer of 1818. She was affected with violent pain in the stomach which seized her immediately after dinner, continued with great violence during the whole evening, and gradually subsided after midnight: it sometimes occurred after breakfast, but more rarely.

The complaint was of two years' standing, during which time a great variety of practice, and every variety of diet had been tried, but with slight and transient benefit. The paroxysms occurred with perfect regularity; she was considerably reduced in flesh and strength, and had a sallow unhealthy look, and her whole appearance gave strong grounds for suspecting organic disease. In the epigastric region no hardness could be discovered, but there was considerable tenderness on pressure at a particular spot. Various remedies were employed during the summer with little advantage: at last, however, she appeared to derive some benefit from lime-water, and returned home in the autumn rather better. But the affection soon recurred, and she returned to Edinburgh as bad as ever. After another trial of various remedies, this severe and intractable affection subsided, under the use of the following simple remedy:—She took two grains of the sulphate of iron three times a day, combined with four grains of the aromatic powder, and one grain of aloës, which was found enough to regulate the bowels. Under the use of this remedy, she was soon free from complaints, and has continued to enjoy good health.

The special features of indigestion, of which each, when most prominent, gives a character of its own to the disease, and requires some particular modification of treatment, (setting aside, for the present

the attendant symptomatic disorders,) are referrible to three heads. Or the stomach may be disordered in three ways, in its secretion, in its mode of sensibility, in its action. In health, the stomach secretes gastric juice proper in quality, and just enough in quantity; it feels nothing when proper food is put into it, and undergoes digestion; and that food when changed into chyme, or digested, it propels by the action of its muscular fibres into the intestines. But in indigestion, the stomach is liable to form a liquor, indigestive, irritating, or inert and profuse in quantity;—it may be the seat of pain and tenderness;—its action may be inverted. Each of these form principal features in different cases of indigestion, which may be studied under the three heads of heartburn or waterbrash, heightened sensibility of the stomach, nausea and vomiting.

*Heartburn.*—The cause of heartburn is superfluous acid in the secretions of the stomach; or it is probable that there is something more than excess of acid, and that the acid formed is different from the wholesome solvent acid of the gastric juice. However this may be, heartburn depends upon the presence of a superfluous quantity of acid in the stomach, which manifests itself by its effects on the sentient surface of the œsophagus. From time to time the acid escapes from the opening of the stomach, regurgitating along the tube of deglu-

tition. But the sensibility of that tube is different from that of the stomach; it is much higher, and the sour liquid, which was not felt by the stomach, immediately excites in the œsophagus a painful sense of acridness.

The proof that it is the *acid* quality of the secretion which produces the sensation of heartburn, lies in the fact, that an alkaline draught immediately relieves it; ten grains to half a drachm of the carbonate of soda or potass, or magnesia, in a wine-glass of water, directly allay or lessen the unpleasant feeling, which if it recurs, again gives way to the same remedy.

Most persons have experienced heartburn; it is the commonest form of indigestion. In general, it occurs under the following law. The stomach has been set wrong by some imprudence in diet; and in the evening, or at night, heartburn is felt. The heartburn is for the time removed by an alkali. But for several evenings, the digestion having once been set wrong, it returns. The disposition to form acid then gradually wears off; recovery is promoted by abstemiousness in diet, and the use of one or two doses of aperient medicine.

Heartburn is of common occurrence in delicate women; they are slightly out of health, languid, of capricious appetite; the heartburn is felt more frequently after breakfast than after dinner, in the latter being prevented or corrected by the stimulus

of wine. In such cases the following common formula, combining a tonic with an alkali, is often beneficial,—six drachms of infusion of orange-peel, joined to the same quantity of infusion of cloves, with from seven to fifteen drops of liquor potassæ, and ten to fifteen drops of the aromatic spirit of ammonia, to be taken twice a day; that is, two hours after breakfast, and an hour before dinner. This remedy pursued for a fortnight, with the occasional use of aperient medicine, will often completely restore the tone of the stomach from this trivial degree of impairment. In place of the infusion of orange-peel, twenty drops of the compound tincture of gentian, in three table-spoonfuls of spring water, may be used, which is the more convenient from its portableness. In general, I know no better or wholesomer tonic of a light description than the above combination, ammonia, namely, with gentian, soda being added or not, according to the quantity of acid formed.

There are some with whom it is beneficial to take a small quantity of alkali *with* their meals; they should take, for instance, five grains of carbonate of soda, and five of magnesia, in a wine-glass of hot water, or milk and water, at breakfast, and the same or a larger quantity in half a tumbler of Seltzer water, at the close of dinner. The formation of acid is thus prevented. The connexion between acid in the stomach and gout is

well known. I am acquainted with a gentleman strongly disposed to gout, whose constitution affords a perfect test of the efficacy of alkalies taken thus at meals. If he omits to take his dose of alkali at dinner, within an hour after the repast, one knuckle of his hand is sure to become red and heated: this he invariably experiences if he neglects adding to his temperate meal the antidote to the acid which it causes.

*Waterbrash.*—There are patients into whose stomachs, when empty, there is poured out a vast quantity of transparent liquid, sometimes tasteless, sometimes sweet, sometimes acid, or of an acrid taste. The complaint is known by the name of waterbrash; it is more common in women than in men, and exists in every degree of severity or mildness. When the quantity of liquid formed is considerable, great extenuation often follows. It is commonly unproductive of pain, and is thrown up without nausea, the mouth filling spontaneously with the liquid, which rises without an effort, or as in animals chewing the cud. Sometimes, however, it is the reverse, and the patient suffers very violent pain at the stomach, which is only relieved upon her bringing up in the same manner a pint or more of limpid and perhaps tasteless fluid; this may happen three or four times in the day. By an observation of Andral's, the complaint appears to arise from a low degree of inflammation of the

glands of the stomach, which become slightly enlarged, and pour out a more or less altered mucus, which is the waterbrash. It is a sort of relaxed inflammation, and accordingly is cured by astringents. The following formulæ are appropriate: ten grains of kino, with a third of a grain of opium, taken in water three times a day; or three grains of alum and five of kino so taken; if the fluid thrown up is acid, an alkali should be combined with the astringent. In the worst case of this disorder, which has fallen under my own notice, the complaint resisted every means that I employed for its relief; when the patient, a labourer from the country, about forty years of age, solicited my attention to another disorder under which he laboured, namely, stricture of the urethra, attended with great heat and pain in passing water. For this, by the ordinary means, he obtained in the course of a few weeks considerable relief. In the mean time, however, the waterbrash had likewise greatly improved, and in a few weeks finally left him. At the time when he became my patient, and for months previously, he used to discharge from the stomach by violent and protracted retching upwards of a quart of brownish acid mucus about once a week. During the preceding days the stomach would slowly swell and become distended. In the mean time he took food and digested it daily. Possibly small doses of cubebs and of



balsam of copaiva and alkalies, which he took for the heat at the neck of the bladder, may have contributed to his recovery.

*Pain of the Stomach.*—One kind of pain of the stomach has been already described. When undigested food is retained in the stomach, it is liable to cause settled pain and uneasiness,—a symptom so far useful, that it leads to the expulsion of the offending substance. The stomach sometimes marks in this manner that a particular kind of food is temporarily unacceptable to it. A gentleman, in the general habit of drinking porter with his dinner, for some days before a severe and alarming attack of vertigo from indigestion, experienced a dull ache in the stomach after drinking porter, which would last for an hour, and subside very slowly; he had every reason to believe that the attack of illness which followed, arose from pressing this then unwholesome liquid upon the reluctant stomach. One sort of stomach-ache thus arises from the presence, in a healthy stomach, of less digestible food.

Another kind of pain of the stomach results from habitual indigestion, and supervenes from one hour to five or six after a meal. The meal is eaten with more or less appetite: after a period, the solvent secretion not being poured out in sufficient quantity, pain more or less severe supervenes; this is accompanied with, and partly caused by

distension from flatulence, sometimes by heart-burn; sometimes it depends upon the mere presence of the unchanged food, and there is little distension and no acidity. This occurrence is well exemplified in a case above narrated, page 35; I have met with it in several other instances in the same degree of aggravation. At the worst, the patient has no repose or cessation of pain till he has mechanically excited vomiting and thrown the food off his stomach, which often is returned unchanged, although eight or nine hours have elapsed since the meal. In lighter seizures, after two or three hours of suffering, the stomach beats the meal; the pain wears off, enough gastric juice having gradually been poured out to dissolve the food.

The treatment of such a case must vary with its features. If there is tenderness of the epigastrium, leeches and hot fomentations should be applied; and if the patient has not an irritable skin, a blister afterwards, over the stomach. By these means, what inflammatory character the indisposition may bear admits of being removed. The same object is promoted by the use of aperient medicine, and by regulating the diet to that which is plain in kind, and extremely moderate in quantity; above all things, if the patient is a good observer of his own symptoms, by attending to the suggestions of his appetite. If there is acidity pre-

sent, as a temporary expedient, an alkaline draught gives great relief; and the best form is the effervescent one,—a draught, for instance, consisting of thirty grains of carbonate of potass in a wine-glass of spring water, with twenty grains of tartaric acid stirred into it. Liquids containing carbonic acid are very salutary to the stomach; whether they act on any chemical principle is not known. There is generally some flatulent distension along with the state of indigestion which I am now describing; the effervescent saline, instead of leaving the stomach loaded with more wind, promotes its immediate expulsion. One of the principal means, however, for allaying this painful indigestion, consists in lowering the sensibility of the stomach by opium. This, in general, should be combined with a purgative;—the dose may consist of half a grain of opium, three of compound extract of colocynth, one of soap; this pill to be taken two or three hours after the meal, the digestion of which is most painful. The aperient is given to obviate the constipating effect of the opium. Opium, however, requires to be used with caution in such cases, lest it disguise some progressive mischief of an inflammatory or ulcerative character in the organ. Opium may be given during the paroxysm of indigestive pain; Batley's liquor opii sedativus is then probably the best form; ten to fifteen drops may be administered in an effervescing saline.

Opium applied externally is sometimes beneficial, either in the form of the emplastrum opii, or upon hot flannel equal parts of laudanum and camphor liniment.

Sometimes the stomach has a morbid sensibility which causes pain the moment anything is introduced into it, the pain continuing till digestion is completed. This form of disorder generally appears not to depend upon inflammatory action, or, at all events, it does not yield to the means which should be efficient against inflammation; so leeches and blisters, and aperient medicine, are commonly of no avail. The remedies which are to be tried, and which are sometimes efficient, must be viewed as purely empirical; not that they are to be undervalued on that account, but rather the more studied, as calculated by their operation to throw light upon the nature of the disorder which they relieve.

The remedies which are likely to be beneficial, are the oxide of bismuth, given in a pill;—to begin with, two grains of oxide of bismuth combined with two grains of extract of chamomile: or kresote, given one drop as a dose in an ounce and a-half of water three times a-day. Or prussic acid, a drop of which to two drops in a wine-glass of water, has again allayed morbid sensibility of the stomach.

A symptom, the opposite of the last, is pain when

the stomach is empty, often attended with craving for food, and relieved by taking it; so three hours after breakfast, as soon as the stomach has got rid of its contents, pain will supervene, that is not relieved till another meal. A patient with a pulsating tumour in the belly, and suffering habitually from indigestion and pain of the present description, was used to relieve that pain, by having recourse to another meal as soon as it recurred; taking thus between eleven and twelve bread and cheese and ale, as a corrector of the craving pain following the complete digestion of his breakfast. His tongue was furred, his complexion loaded, the action of the bowels irregular. By reducing his diet, and prescribing the regular use of aperient medicine, his health greatly improved: at the same time, the early luncheon was forbidden, and a draught of half a drachm of tincture of henbane, seven grains of soda, and ten drops of sal-volatile, in water, taken on the recurrence of the pain of the stomach, was sufficient to remove it.

To give another instance. A. B., aged forty-five, for the last four or five years, had experienced pain and weakness at the chest, which was most distressing in the evening; it went away on going to bed, and was better on her taking food. The catamenia ceased a year ago. During the last half year she has been worse; there has been frequent distension of the stomach, and flatulence,

and loss of appetite; the bowels being confined. I ordered a five-grain pill of blue pill and extract of rhubarb at night, and a light tonic, with ammonia and soda twice a day; this she found of benefit. Afterwards, the bowels being still insufficiently relieved, two grains of hydrargyrum cum cretâ, with two of compound extract of colocynth, were taken every night. This plan being pursued for three weeks, she had become much better, and her appetite had returned; still there was some pain at the stomach, frequently recurring during the day. For this she took a grain of oxide of bismuth, with three of the confectio opii, three times a day, from which she derived additional benefit; when she left London.

The same remedies which I have enumerated for the preceding case, have equally been tried in this, but with uncertain success. Another, that has been of use, is the nitrate of silver, beginning with a grain at a dose, in a pill made with crumb of bread.

There are attacks of pain in the stomach, which are commonly attributed to spasm. They come on suddenly, are of the most acute description, and seldom last long; they are described as suggesting the idea of a knife stuck through the side. Women are more subject to these seizures than men. I suppose that they entirely result from distension with wind; they are allayed at once, by taking a

wine-glassful either of very hot water alone, or hot water with twenty drops of sal-volatile, or with a dessert-spoonful of brandy in it; at the same time, and in which the relief seems to consist, wind breaks off the stomach. Sometimes a few drops of laudanum in hot water, with some cordial, form the best remedy for these seizures\*.

Generally speaking, pain in the stomach does not denote serious disease, but is the result of slight functional disturbance. Nevertheless, obstinate cases of pain at the stomach, attended with

\* The popular objection to laudanum is the danger of falling into the habit of taking it, which is the greater for the false supposition that the dose must be progressively increased. This, indeed, is the case, if the remedy is daily and hourly resorted to, but not otherwise. I have under my care a patient with disease of the womb, whom I am obliged to allow the daily dose of seventy grains of crude opium; the continual increase has been unavoidable, as the remedy is used for constant pain. She says that her memory is not so perfect as it was; but she experiences no confusion of head, nor any form of mental disturbance. When the use of opium is not continual, the same dose, as with aperient medicine, serves a life-time. I am acquainted with a gentleman who, several years ago, received a charge of shot in the calf of his leg; some of the shot have lodged in the tibial nerve, and he experiences constant pain in the foot, only varying in severity. To alleviate this, at one time he used to take 700 drops of laudanum daily; after a time he gradually discontinued taking it. This gentleman is now, from time to time, subject to spasmodic pain at the stomach, after he has taken anything to disorder his digestion. He then has recourse to laudanum, and finds the small dose of seven or eight drops always enough to pacify the nerves of that organ.

distension, with flatulence, recurring irregularly, lasting several days or weeks, and gradually lessening, then renewed, again going away and returning, we sometimes found to depend upon ulceration of the lining membrane of the stomach. This is a complaint of a serious nature; for although it is often recovered from, yet a certain proportion of those attacked with it, after weeks, or months, or years of capricious indigestion, die suddenly, through perforation of the stomach, and the escape of its contents into the cavity of the belly.

Pain of the stomach, therefore, although commonly dependent on trivial causes, should, if it occur frequently, be looked upon with suspicion, and all the points connected with it carefully considered. The accompanying symptom, which gives it the most alarming complexion, is tenderness on pressure, not general soreness of the whole stomach (for whenever the stomach is distended, it is uneasy on pressure), but tenderness always to be found about one spot. When with this there is disinclination to food, the tongue disposed to dryness, occasional nausea and vomiting, one would entertain serious apprehensions of chronic inflammation, thickening, ulceration. In such a case, the greatest abstinence, leeches to the pit of the stomach, blisters, a superficial issue, the milder preparations of mercury, are the appropriate remedies.

III. *Nausea and Sickness*.—There are few sub-



jects in pathology which have more extensive relations than the present. The stomach is the centre of sympathies: no organ can be suddenly deranged, but the stomach is drawn into consensual disorder, and vomiting is the readiest form of disturbed action into which it falls.

To take an extreme instance: in a case, in which I performed amputation at the hip-joint (the patient, I may mention, recovered), for a few hours the utmost prostration existed, the patient hardly was heard to breathe, the pulse was quick and tremulous, no water was secreted, she vomited repeatedly. In injury of the head, or sudden effusion in the brain, vomiting is an almost constant attendant,—upon sudden and violent mental exertion, vomiting occasionally takes place,—the same is produced by violent bodily exertion on a full stomach. If any inward action is going wrong, if a gall-stone is passing through the gall-duct, a calculus descending from the kidney,—vomiting appears to follow as a natural consequence.

The mechanism of vomiting is very simple, and the reverse of what it is popularly supposed to be. In vomiting, the stomach is perfectly passive. It does not act in expelling its contents. It is a flaccid bag, squeezed by external pressure. The muscles of the belly and the diaphragm suddenly and convulsively compress and empty it. This convulsive action depends on the sensation of nausea. The

phenomena are paralleled by those of sneezing; if you look at the sun, or take a pinch of snuff being unaccustomed to it, a sensation is excited, which is relieved by an uncontrollable convulsed expiration. Nausea is the analogous sensation, which is relieved by retching.

The two forms of nausea and vomiting that are most familiar, are those attending early pregnancy and sea-sickness.

The vomiting of pregnancy is very variable. Generally it is first experienced shortly after conception, and terminates with quickening. Sometimes it is hardly experienced at all, in other cases it lasts during the whole of pregnancy. Sometimes it gives little distress, in other cases it has been so constant as to compel the production of premature labour to save the patient.

The vomiting of pregnancy seems, to a certain extent, dependent upon change of posture; the nausea and retching are commonly experienced on rising; the vomiting, however, frequently recurs after breakfast; the sufferer then lies down, becomes tranquil, and afterwards, as the day goes on, her stomach usually recovers its tone. The vomiting sometimes returns at other periods of the day. Preserving the recumbent posture always mitigates the disposition to vomit. The cause, however, of this sickness is in a small part only mechanical. In an operation which I performed for an obstruction which had been caused by inflammation fol-

lowing labour, and which had a prosperous result, on half-dividing the extremity of the womb, the patient became sick. The vomiting of pregnant women is caused by sympathy of the stomach with the womb; it is a sudden indigestion produced by consent with a remote and labouring organ\*.

Vomiting is liable to be produced by uterine congestion, when it obeys the same laws as the vomiting of pregnancy.

In sea-sickness, the nausea and vomiting result from sympathy of the stomach with the senses and brain. It is the feeling of disturbed equilibrium which, in this instance, deranges the organ of digestion. This any one newly at sea may soon ascer-

\* Such is the force of sympathy which the womb exerts, that almost every functional disturbance may be produced through it. Salivation has been produced by pregnancy. In the following case, a disease, which pregnancy arrested, has been equally arrested by salivation.

I attended a patient with Mr. Angus, of Greek-street, who was the subject of violent spasms of the limbs, leading to fits of insensibility, when at the worst. These spasms, at one time, suddenly ceased; soon after, she found that she was pregnant; and her delivery took place nine months all but two days after the sudden cessation of the spasms. Eight days after her confinement, the spasms returned; and when she had suffered with them for some weeks, I was consulted. I advised that some local means should be applied at the back of the neck, where she felt pain that seemed to shoot thence down the arm, and prescribed mercury to affect the mouth; she recovered. She has since had four serious relapses; each time she has completely recovered by taking mercury in large quantities. Her amendment has invariably commenced as soon as the mercury has affected the mouth

tain to be the fact. If you walk or stand upon the deck, all is seen and felt to be unsteady and shifting; the vessel moves even on a calm sea with a gentle undulation, and the masts and sails and lines of cordage now sink, now rise; all is visually unsteady, and you have, at the same time, incessantly to adapt your body to the varying inclination to the horizon of the plane on which you stand; unaccustomed to these impressions, you become giddy and sick. If you have partaken of a meal shortly before sailing, the effect is but the more certain and rapid; the stomach is contending with the ingesta, and all the more readily disordered.

That it is the sense of disturbed equilibrium which produces the sickness, is shown by the following proof. If, instead of maintaining the erect posture, you lie down, or, in other words, assume a posture of more stability, the nausea abates; and if, in addition, you close your eyes, and exclude the visual disturbance or unsteadiness, the nausea is further lessened.

Upon these facts the treatment of sea-sickness turns.

If the voyage is to be a short one, from ten to thirty hours, go on board with an empty stomach, betake yourself to your cot, and lie down. It is better to go on board fatigued, which will enable you to sleep at once; when you wake, preserve the recumbent posture; some nausea may super-

vene, but it will be slight, and soon pass off; if you feel an appetite for food, eat very sparingly of the lightest and plainest food.

If the voyage is to be a longer one, still it should commence in the same manner; and during the sleep into which you fall from fatigue, your senses will become, in a degree, accustomed to the motion of the vessel; and the training will have begun, by which the art is acquired of preserving one's equilibrium on a rocking surface.

Nausea and vomiting are liable to occur after important surgical operations, or grievous bodily accidents, in three forms.

One kind has been already adverted to; it goes with the general prostration, and follows the bodily injury immediately. It is comparable to that which is experienced by those upon whom mercury acts as a poison; they have headach, feeble intermitting pulse, nausea and vomiting. Small quantities of brandy in soda-water, with a few drops of laudanum, form the best remedy. If the vomiting is obstinate, an opiate plaster or liniment to the pit of the stomach will assist.

A second kind is bilious vomiting, the nature of which has been fully noticed at page 33. After the amputation of a limb, in twenty-four or forty-eight hours, the patient will sometimes experience such a seizure.

A third kind, which sometimes follows within

forty-eight hours after surgical operations, is vomiting of large quantities of a transparent grass-green acid and acrid fluid, without the bitterness of bile. This commonly goes with great debility, and is of most unfavourable augury. Effervescing alkaline draughts, with laudanum and a mustard plaster or blister applied over the stomach, and if the bowels have not been relieved, a purgative injection, are the proper remedies.

Vomiting occurring habitually as a mode of indigestion from primary disorder of the stomach, is not less distressing from the suffering it occasions, than alarming from the possible nature of its source. Cancer of the stomach is uniformly attended with this symptom, although in some cases of this fatal disorder, it is much more prominent and constant than in others. In the last stages of the disease, when no ingesta are received into the stomach, the matter thrown up resembles coffee-grounds, and consists of blood changed by the action of the stomach. Uneasiness of the præcordia, loss of appetite, vomiting when food is taken, tenderness on pressure, and sensible tumour or hardness of the epigastrium, are the pathognomic features of cancer of the stomach, under which the patient gradually becomes emaciated, and finally dies of exhaustion and inanition. But fortunately there are few cases absolutely without hope. And it has frequently happened that when the symptoms

have been such as to lead to the greatest apprehension of the worst results, under fortunate treatment they have been dispelled.

To some which I have already given, I will add the following, as exemplifying the efficacy of another plan of treatment.

A young woman in the family of the English consul-general at the Hague, in the spring of 1818, was subject to intractable vomiting, which had gradually supervened in three months. At first, the vomiting took place occasionally only; after a short time, she observed that it occurred after those meals at which she took meat; in time, after every meal, and occasionally when nothing had been taken into the stomach. However, she threw up no blood or coffee-ground fluid; and although there was pain at the præcordia, and tenderness on pressure, there was no hardness. She had become greatly emaciated; a variety of remedies had been tried, and had proved ineffectual. I recommended that she should take three times a day a quarter of a grain of sugar of lead, and a third of a grain of opium, and that a blister should be applied to the pit of the stomach. On the following day the vomiting ceased, and never returned; the lead and opium were however continued for a week, (opening medicine being likewise given,) and a second blister applied.

I have since met with several parallel cases of different degrees of severity, which have yielded to

similar treatment, where the vomiting has been either the only symptom of the case; or where it has been coupled with hysteria.

These, however, are to be viewed as cases of vomiting of very unusual severity; commonly the complaint yields more readily—to abstinence, rest, aperient medicine, opiate applications, and blisters to the pit of the stomach, opium administered internally, creosote.

The principal symptomatic disorders which flow from indigestion affect the head or the chest. Upon these I may add a few brief remarks.

In the severest cases of head seizure from stomach derangement, the patient falls apoplectic; he lies in a state of heavy stupor; the remedy wanted to relieve which is an emetic: the stomach is oppressed with a refractory meal. It requires, indeed, no small exercise of judgment to distinguish such a case; but thus it has made itself known. A person has fallen insensible; the countenance pale, the pulse weak, not full and laborious, the insensibility complete; some stimulant has been given to reanimate expiring life. Then vomiting has followed, an undigested meal has been thrown up, and the patient has recovered.

Some of the less serious forms of head affection from impaired digestion have been already adverted to. The commonest features are excellently described in a paper by the late Dr. Warren, from



which I extract the two following passages as giving a very graphic account of the disorder. The treatment is the cure of indigestion.

“In one form of disorder,” says Dr. Warren, “the pulse is languid and feeble, but not more frequent than natural; the tongue is whitish and slightly coated, the edges of it are of a pale-red colour. The patient perceives a sensation of mistiness before the eyes, and general indistinctness of vision; he feels a dull pain or weight in the head, attended with some confusion, is slightly giddy, and fearful of falling. When on foot, he thinks himself insecure unless in company, but if leaning upon the arm of a companion, sitting still, or moving in a carriage, he is free from giddiness and alarm. If desired to walk towards a distant object, he is reluctant to make the attempt, from a doubt of his ability to reach it without falling; and with the inconsistency of a person under alarm, will, without hesitation, proceed to a greater distance than that which has been proposed to him, provided that it be in the direction of his own house, or lead him to any other situation where he conceives he may be within reach of assistance. These symptoms are attended with an uneasiness or irritation of the stomach, a slight nausea, and often by a sensation of constriction of the fauces, accompanied with a watery secretion from the posterior part of the mouth. Coldness, slight stiff-

ness or numbness of the fingers, are sometimes present, and the other parts of the system are in general affected with a great degree of nervous sensibility."

"In general, restlessness precedes the attack which is followed by uneasiness of the head, and want of the usual distinctness of ideas, oppression of spirits, disinclination, and sometimes incapacity for mental exertion, chillness of the body, coldness and dampness of the hands and feet. Next succeeds pain or dull aching of the head, sometimes of the forehead only, at others affecting the crown of the head, and posterior part of it, attended with a sensation of coldness and tightness of the scalp, slight giddiness, weight, pain, distension and stiffness of the eyeballs. In some cases, as these symptoms increase, they are accompanied by tingling and numbness of the fingers and hand. This numbness is never so severe as entirely to deprive them of feeling; and, as far as I have observed, seldom attacks the whole hand at the same time; but usually first attacks one or two of the fingers, and as they recover, spreads to a neighbouring finger, and having, in this manner, traversed the whole hand, extends to the wrist, after which it ceases altogether. The symptom last mentioned is of short duration, seldom continuing more than a quarter of an hour; but it leaves a slight sensation of stiffness, which does not

entirely cease until the headach has been removed by art, or has finished its natural period. As long as it is present, the hands are damp and cold, the pulse is languid, but neither quicker nor slower than is natural."

Complications of head symptoms with vomiting occasionally display, in the following remarkable manner, the force of sympathy between the head and stomach. Vomiting, with severe pain in the head, is sometimes relieved, when nothing else will relieve it, by cold applied to the forehead.

Of the symptomatic effects of indigestion that are felt upon the chest, the commonest are cough, dry, ringing, a stomach-cough, which the ear recognises, which often goes with heartburn, and appears provoked by it; or again, cough coming on in the morning upon rising, with considerable expectoration of thick mucus, exciting nausea. The more alarming symptoms are disturbance of the action of the heart, which is often irregular, intermitting, seeming to stop with a suffocative sensation, or beating hurriedly with violent palpitation. All the feelings which go with organic disease of the heart may depend on indigestion. And, on the other hand, organic disease of the heart will often go on for an indefinite time, producing none or the slightest symptoms, being sometimes for years not progressive, but strictly stationary.

The treatment of these symptoms cannot be

separated from the treatment of the indigestion which occasions them, excepting thus far. When the action of an organ is once set wrong, it sometimes continues to go wrong merely from habit. So when an indigestion is getting better, there may remain nervous irregularity of the heart's action, that some trivial remedy will allay. Thus, when the patient wakes at night with uneasy sensations at the heart, and under the alarm which these produce, he may obtain relief from such a draught as the following:—half a drachm of tincture of henbane, or of tincture of hops, with fifteen drops of sal volatile, in hot water or in camphor julep. If there is acidity present, five or ten grains of carbonate of magnesia or of soda should be added.

The most remarkable case of chest-affection dependent on indigestion which has come under my observation is the following:

A gentleman, forty-seven years of age, of great strength of mind and body, but who had lived very freely, and who was accustomed to, and disregarded, the common results of intemperance, became at length the subject of peculiar seizures, which attacked him on waking during the night, and in the morning. They used to begin with a tickling sensation about the chest, and a feeling of lightness and cheerfulness; then a sense of constriction supervened, amounting rapidly to pain; it seemed first, he told me, as if the chest was

strictly bandaged, then as if the ribs were pinched together by red-hot screws; the agony being now intense, he suffered, in addition, pain at the wrists, and his hands were numb and powerless. The whole of the attack would occupy three or four minutes; the pain appears to have been frightful; and the patient was left for some time afterwards extremely exhausted. He recovered his health upon a total change of diet, taking no wine or fermented liquor of any kind, or animal food; but eating for dinner, pudding hot with cinnamon, and for breakfast, and in the evening, taking tea only, with bread and butter and ginger. This patient consulted me when he had pursued this system, which had cured him of the attacks above described, so long a time that it had reduced him to a state of debility, and threatened to originate a new form of dyspepsia. Upon resuming ordinary diet, with a moderate use of wine, by my advice, he completely recovered his health, and is now after many years perfectly well.

---

## CHAPTER II.

## MANAGEMENT OF THE BOWELS.

WHEN the food has been converted into chyme by the action of the gastric juice, it passes from the stomach into the small intestines. There the chyme becomes penetrated by the bile and other fluids; after which mixture it gradually separates into two parts; one of which, the chyle, is absorbed by the lacteal vessels, and carried into the blood; while the remainder is propelled onwards into the great intestines, to be thence, after some additional changes of less importance, and some further absorption of its elements, finally eliminated. Such is, in brief, the ordinary theory of the functions of the bowels. It assumes, the reader will observe, that food consists of two parts, recrementitious matter, and refuse. But a doubt may be raised whether this distinction is invariable and essential; or, rather it is most probable that it is not so. Under some circumstances, there is every reason to believe that all the food taken becomes nutriment. When, for example, one is in habits of strong exercise, eating at the same time sparingly, and of wholesome and digestible food alone, it is presumable that the whole of many meals is assimilated.

When we inquire for rules as to the management of the función of the bowels, it becomes evident that in what concerns recrement and nutrition, there is room for none. When the stomach has digested, and the chyme has reached the bowels, the nutriment which it contains is, without our control or knowledge, absorbed and conveyed into the circulation. Rules of management only apply to the getting rid of the refuse part. The importance of this point, however, merges in one of greater consequence. The same channel which provides an escape for the accidental quantity of refuse in the food, (whether less or greater, from the food having been sparing or in excess, digestible or indigestible,) gives vent simultaneously to new secretions. It is to be understood that the alimentary canal forms a sort of inward skin, or has a membranous lining continuous with the outer skin, which pours out its own secretions, as the outer skin pours out perspiration. And as there are special occasions when the flow of perspiration from the external surface relieves and lightens the body, so are there others when secretion from the mucous lining of the intestines serves a parallel purpose. Or we are no doubt to suppose that one use of the intestinal secretions is to dilute the refuse matter passing down, and to cause to separate from it additional nutriment; but it is additionally certain that another object is obtained by these secretions,

the cleansing, namely, or purifying the blood of the matter so discharged. And it is probable that unless our food is indigestible or immoderate, the elimination of that part of the *fæces* which is new secretion is of much more importance to the œconomy than the expulsion of the refuse of the *ingesta*.

It is moreover probable that the larger part of the *fæces* is new secretion, not refuse food. Whence it is that the quantity of *fæces* does not bear an uniform proportion to the quantity of *ingesta*. Another circumstance determines this proportion;—which is the disposition to pour out intestinal secretion, which varies in different persons, just as the same function of the skin varies; there are some who perspire readily and profusely, and others who never display sensible perspiration; so are there some to whom large alvine discharges are natural, others to whom they are not so. The blood has, through one vent or another, to get rid of the noxious elements which accumulate in it in the changing flow of the circulation; that vent may be the skin, the kidneys, the bowels, almost vicariously.

In proportion as the *fæces* consist of intestinal secretion, is their increased quantity capable on the one hand of relieving the system, on the other of lowering it. Suppressed perspiration leads to various kinds of illness, so likewise the suppression



of this function. Excessive perspiration debilitates; much greater debility is produced by excessive action of the bowels. And both upon the same principle. The blood in the one case is not cleared and strained of impurities, on the other, too much material is abstracted from it.

Consistently with, or on following out these views, it will be found to be generally true, that persons with natural looseness of the bowels are of less stamina than those who are habitually constive. And the management of the bowels becomes of the greatest moment; as in it are the readiest and least exceptionable means of altering the quantity and chemical nature of the blood, and of reducing the bodily strength. Both of these objects have continually to be aimed at in medicine and surgery.

There is a collateral ground upon which the management of the intestinal secretions is of a value hardly less than the preceding. All the organs of the body are wonderfully linked together by sympathetic influences. So that one cannot be disturbed without the others being in some degree affected; and often, when several are deranged, a wholesome action excited in one will draw the others into health; so Dr. Currie found in the general disorder of many functions ushering in continued fever,—the temperature raised above the natural standard, the skin hot and dry, the tongue

furred, the digestion suppressed, the pulse frequent, the brain excited,—that by a strong and forcible change impressed on *one* of these functions, all the rest were, as it were, pulled right simultaneously: the cold affusions, which he employed, lowered the temperature of the body, and the other functions fell into their natural rule and order consentaneously, and at once. But the sympathies of the bowels seem to ramify through, and search the whole system; and there is hardly a local complaint, or catenation of general disorder, but admits, to a certain extent, of being drawn towards health, by the judicious exhibition of medicines that act upon the bowels. In all febrile and inflammatory attacks, (excepting such of the latter as have their seat in the bowels themselves,) in all cases of congestion of blood in important organs, in the brain, the lungs, the liver, for example, the mode of relief that is commonly surest and safest is to force copious evacuations from the bowels, combined with bloodletting where bloodletting is advisable, otherwise its preferable substitute. The following case, which has recently been under my care, is an instance so apposite, that I am tempted to introduce it here. A gentleman, aged 53, of temperate habits, short but stout in person, two years ago, had several ulcers upon the legs and thighs; of these he recovered, taking a course of alterative medicine, and the sores being treated in the usual manner.

I recommended him in future to observe the greatest care as to diet, medicine, exercise; and he did so. Nevertheless in the spring of the present year, other ulcers made their appearance similar to the former; for which, when he had doctored them some time himself, he again placed himself under my care, and I gave him the additional directions necessary, and they were improving and I saw him seldom;—when he became short breathed, his belly full of wind, his face livid, his legs blue, and his urine rapidly decreased in quantity, and was high coloured. The least exertion hurried his breathing distressingly; his pulse was extremely frequent. It was evident that he was threatened with effusion on the chest; but none had yet taken place: he could lie flat in bed, on his back, or on either side; his ankles were not swollen. He laboured under a thin heart, and an overloaded circulation. I had him cupped to ten ounces on the chest with slight relief; then after repeated doses of elaterium, producing copious watery purging, the flow of urine was re-established, and he recovered.

#### SECTION I.—OF LOOSE BOWELS.

HABITUAL looseness of the bowels,—their action being repeated, for example, two or three times in the twenty-four hours,—although not commonly met with in persons of the most enduring frames

and hardiest fibre, is yet consistent with the enjoyment of good health, and the attainment of great age. It has this advantage,—it prevents plethora, and confers some exemption from acute inflammatory illnesses. In general, those who are fat, and persons of a highly nervous temperament even when thin and spare, have loose bowels. Persons of both these classes are less capable than others of strong and sustained mental or bodily exertion. It is difficult to contend against nature, but such persons act judiciously, when they try to improve their temperament and habit of body; to which nothing more contributes than moderation in diet, and abstemiousness in regard to liquids, and liquid food.

Habitual looseness of the bowels is not common in children, but many persons, from the age of twenty upwards, fall into it. Through some accidental influence their health goes wrong, and the disturbance of the system tells upon the mucous lining of the intestine, rendering it præternaturally irritable. The disorder shows itself in increased secretion, frequent loose motions, sometimes consisting of mucus alone, sometimes of liquid fecal matter: there is sometimes a general sense of uneasiness in the belly; in other cases local pain, which comes and goes, but recurs about the same spot, often joined with tenderness on pressure. In this condition, fluctuating between worse and better,

a patient may continue for weeks, and months, and years, if not properly treated. The ailment is peculiarly liable to be made worse through atmospheric agency, and the influence of the mind; any strong mental exertions or emotions are sure to increase it; it, therefore, in a remarkable degree, interferes with, and sets a check upon, a life of energetic thought and action, converting a person otherwise in health, and fitted for strong and busy occupation, into a desponding valetudinarian. The management of such cases turns upon the following principles. Every error of diet must be avoided; the stomach is commonly sound, and capable of digesting wholesome and moderate food; fruit, much vegetable of any kind, all raw vegetables, and acids, porter likewise, and ale and cider, are to be shunned; but wine, either Madeira or sherry, taken to the quantity of two or three glasses, is generally found to strengthen without heating the irritable bowels. Of medicines, tonics, especially preparations of iron, the lighter astringents, and opium given in small doses, and with caution, are often serviceable. All aperient medicines, especially preparations of mercury, are to be scrupulously avoided, supposing, of course, that the complaint is not complicated by other local or general disorder requiring the exhibition of such medicines.

The following cases may serve to exemplify different varieties of irritable bowels.

A gentleman about forty years of age described his case to me thus: he had been for three years liable to some ailment of his bowels. His appetite and digestion are tolerably good; but he suffers for weeks together a sense of general inward soreness of the belly; when this is at the worst, he passes mucus from the bowels several times in the day. The complaint is attended with loss of strength, indisposition to exertion, depression of spirits. It seems connected with the state of the skin, and is more influenced by the weather than by any other cause. In warm summer weather, with southerly and westerly winds, the skin acts, he perspires, and he is well or materially better. When an easterly wind sets in, his skin is parched, and all his symptoms recur.

A gentleman, now forty-three years of age, of a sanguine-nervous temperament, enjoyed good health till twenty. At that period he was attacked twice or thrice with inflammation of the eyes, for which he was much purged and bled. This treatment, he thought, weakened him, and gave origin to pains and uneasiness in the bowels, and looseness, which became habitual. Between the ages of twenty-eight and thirty-two, he travelled on the Continent, and enjoyed better health; the irritable state of the bowels left him, and he re-acquired strength. Near the time of his return, however, the complaint re-appeared, brought on, as he

thought, by eating fruit, and drinking light wines. He now became engaged in London in the profession of the law, and being much in court, and his mind anxious about his profession, his ailment gained ground. He was at the worst in the summer of 1833. Joined to frequent purgings, sometimes with mucus, and even blood, on two or three occasions, he experienced acute and exhausting pain in the belly below the navel, which part was sore on pressure; he became extremely weak, and the alvine discharges used to be followed by faintness. He was then on the circuit: medical assistance being called in, leeches were applied to the belly, but he thought this remedy further weakened him; hot fomentations were then used alone, and he took some doses of laudanum; after some days he became better again. On his subsequent return to London, he placed himself under the care of an eminent practitioner, and consecutively, but at intervals, under two others, who, taking different views of his case, treated him thus. The first prescribed calomel, which materially increased the complaint. The second prescribed a course of blue-pill, which appeared to the patient to do him harm, but to agree with him better than the calomel. The third physician prescribed carbonate of iron, which was of considerable benefit to the patient, checked in some degree the habitual looseness, and improved

his strength; as the efficacy of the remedy diminished with use, its dose was greatly increased; at the same time, sulphate of copper with opium was used in conjunction with the iron, and the disorder was certainly checked; but new sensations of uneasiness began in the belly, which felt full, and loaded, and uncomfortable, and the patient became apprehensive of the effects of the quantity of mineral substance he was daily introducing into his inside. He discontinued these medicines, therefore, and then applied to me for advice. I recommended him to try vegetable astringents; and prescribed half a drachm of tincture of catechu, with six drops of laudanum, in a light bitter, to be taken twice a day. The system seemed to agree with him, and in a fortnight he was considerably better, having lost all the uneasiness of the belly, and the looseness only troubling him on days of more than usual anxiety and business, and always being checked by an additional dose of the astringent. It is a year and a half since this gentleman first consulted me. He has gone on for this period with the same medicines; the bitter only having been occasionally changed; and when he has been exhausted by unusual professional labour, ten, or twenty, or thirty drops more of tincture of catechu having been added, and two or three of laudanum to each draught. On one occasion, for a few weeks, I added as a tonic a daily dose of carbonate of



iron, which seemed of use; on another occasion, I prescribed with opium, the sulphate of quinine, but it disagreed. He now retains very good health on the condition of taking the astringent draught twice a day; and has completely recovered his strength and capability of exertion.

About three months ago I saw, at the request of Mr. Marshall, of Greek Street, a person who had laboured under looseness of the bowels for two years. It had commenced without any assignable cause; he had daily several yeasty motions; and from a large stout man had become extenuated, and his legs were dropsical; during the last six months, the complaint had gone on increasing; the remedies ordinarily employed in diarrhoea had been used without success. A parallel treatment to that employed in the last case entirely corrected in a few weeks the disordered state of his bowels.

The following case of chronic looseness of the bowels presented a different character to the preceding.

Colonel ——, aged about fifty, consulted me for these symptoms. He was liable to attacks of swelling and fulness of the lower intestine, attended with discharge of blood; these were occasional only; but he suffered inconvenience daily from looseness of the bowels, which acted not only in the morning once or twice, and again perhaps at night, but in the middle of the day, so

as to incommode and interfere with his usual avocations. He was otherwise in strong health, of large person, inclining to plethora. The ailment of which he complained had originated in India, where he had served many years; and there his health had been uninterrupted, owing, as he most likely justly argued, to the habitual looseness of his bowels, which had been moved two or three times a day, with occasional discharges of blood. He was desirous, now that there was no further occasion for this relief of his system, to get rid of what had become a very troublesome affection. He tried, at my recommendation, some plans of which the first were fruitless; these consisted in the daily use of active aperient medicine; then in the use of the same with an opiate injection after the bowels had been freely moved in the morning; then reduction of diet and wine; the inconvenience still continued; I then advised him to try the hot air bath, and to carry off, by profuse perspirations, which his habit could well sustain, the secretions which he had to spare. This remedy was completely successful at the time; and he has not applied to me since.

Acute attacks of looseness of the bowels, or of diarrhœa, are very common in this country at the beginning of autumn, and are often incurred at every season of the year. Two causes serve to produce them, one unwholesome food, such as fish

or fruit when they disagree, and exposure to cold.

Acute purging ordinarily takes one of the following forms in this country. Simple diarrhœa, febrile diarrhœa, diarrhœa with absence of bile, diarrhœa with excess of bile or cholera.

Simple diarrhœa begins with one or two loose motions without pain; then frequent watery but perfectly fœcal motions, with griping, and flatulence, and tenesmus; the tongue is furred, the relish for food impaired, and the patient is languid and indisposed to exertion. Such an attack, left to itself, will often terminate in twenty-four or thirty-six hours, and occurring in a very full habit, and not being severe, should not be hastily stopped. In others, it is desirable to arrest simple diarrhœa early, for its continuance weakens. It is a common error that the continuance of diarrhœa shows that there is some irritant in the intestines, which nature is trying to dislodge. In ninety-nine cases out of a hundred, this is certainly not the case. When the disorder originates in this cause, the offending matter is early in the attack expelled, and nothing remains but the irritated mucous membrane, which continues to pour out immense quantities of watery secretion. Nevertheless this rule is not without occasional exceptions; and violent watery purging has been cured even by an emetic, that has acted by dislodging an offending substance

from the stomach. In children again, it is generally right (and often all that is necessary,) in an attack of looseness of the bowels, to clear out the inside by an aperient dose, of which a powder consisting of chalk and mercury, soda and rhubarb, is the most useful form, to be repeated twice or three times, if requisite, or to be followed by some cordial and slightly astringent medicine.

The management of simple diarrhœa involves two things; nothing is to be done or taken which has a tendency further to excite the bowels; and the medicines used should be calculated to allay their heightened sensibility, and repress the inordinate secretion. The patient, to fulfil the first indication, should remain at home and at rest, and should take no food. To fulfil the second, nothing is better than to begin with two drachms of compound tincture of rhubarb in peppermint water; and then every succeeding three hours to take six drachms of chalk mixture and the same quantity of peppermint or cinnamon water, with five, ten, or fifteen drops of laudanum; or an ounce of infusion of cloves, and half an ounce of the chalk mixture, with ten grains of opiate confection. The diarrhœa will commonly abate after two such doses, which may, however, be continued if necessary. When the purging is attended with tenesmus, that is, with painful straining, an injection

should be administered (and repeated if necessary) of twenty, thirty, or forty drops of laudanum in a tea-cup of tepid gruel or barley-water, or mucilage of starch.

When diarrhoea sets in with shivering and fever, it requires to be treated with a lighter hand; first because it cannot be certain which is symptomatic, which primary, the diarrhoea or the fever; secondly, because if the complaint is febrile diarrhoea only, the fever would be liable to be heightened by the remedies before prescribed.

Rest in bed, and entire abstinence from food, are the first remedies, and essential; with this an effervescent saline draught, with five grains of carbonate of soda in excess, and three grains of Dover's powder, to be repeated every four hours; often, two grains of blue-pill twice a day are a judicious addition to this treatment.

That form of diarrhoea which is characterized by the absence of bile is known by the white clayey appearance of the first semifluid motions, and the appearance, like chalk and water, of those which follow; the motions have likewise a peculiar and offensive foetor. The appropriate remedy for this attack, in addition to rest and abstinence, is mercury with opium. Three grains of blue-pill, or of chalk and mercury, with seven grains of opiate confection, made into two pills, to be taken every

three hours, will at once control the excessive secretion, and bring back in a few hours the deficient flow of bile.

When looseness, which has commenced in either of the modes described, persists, what is principally to be apprehended is, ulceration of the mucous lining of the bowels. In the obstinate purging of certain fevers, of the last stages of consumption, in protracted dysentery, this change of structure is invariably present; and it is liable to occur in such cases as I am now considering. The remedies which are now appropriate are the strong mineral astringents, lunar caustic, or blue vitriol; of either of which half a grain, with a like quantity of opium, should be taken twice or three times a day. Of the two, the former is perhaps preferable; of the efficacy of which I happened very lately to see a notable instance in an old Frenchman of broken constitution, who was in the Middlesex Hospital for local sloughing, when diarrhœa supervened, that resisted every other means.

The indigenous cholera of this country is characterized by purging of bile, frequently accompanied with vomiting of bile, cramps of the legs and belly, and considerable depression of strength. The things to be done are, to abstain from food, from mercury, and aperient medicine; to calm the stomach by perfect bodily stillness, by small quantities (half a wine-glass) of soda-water, with eight or ten

drops of laudanum in it, given in the intervals of vomiting, by an opiate liniment or plaister applied over the stomach; the only object is to tranquillize. When there is great depression, æther, ammonia, and brandy are necessary,—often at the commencement of the attack.

I will conclude this section with some remarks on the treatment of Asiatic cholera, of which malady the following are the symptoms. After diarrhœa, or sometimes no premonitory symptoms having occurred, sudden vomiting of the contents of the stomach without nausea, and sudden purging without pain; the vomiting and purging frequently recurring, the liquid discharged being like rice-water or gruel. With this, spasms of the muscles of the extremities and belly. In a short time, from one to three or four hours, collapse supervenes; urine is not secreted; the pulse is extinguished; the eyes are sunk; the eyelids are surrounded by a dark circle; the voice is hardly audible; the tongue is cold; the skin cold and passively shrunk, with a dirty look, or livid and blueish, particularly on the extremities; cramps of the limbs constant; the vomiting and purging cease, and life.

In this tremendous malady, everything that has hitherto been extensively tried, appears to have failed. And the general disposition of medical practitioners is to fall back upon that which may

be considered the rational treatment of the disease, namely, repeated doses of calomel and opium; calomel to excite bile, opium to allay the heightened irritability of the intestines. The saline treatment which Dr. Stevens introduced is scientific, but has proved nugatory; and the injection of hot salt and water in the veins (however wonderful in its temporary effects,) has not been successful enough to authorize much reliance on this dangerous experiment upon the living œconomy. But there are other remedies, which however little they promise, certainly deserve to be more fully investigated. These are, lunar caustic with opium, oil of croton, cold affusions; the two first to stimulate the stomach and intestines to healthy action, the third to excite general reaction. It is obvious that these remedies should be tried separately. Some partial success, which each of these means has had, furnishes sufficient ground for recommending their further trial. It is probable, however, that, in its worst forms, the appalling antithesis which Magendie used in describing Asiatic cholera, conveys no exaggeration: "It begins," said he, "where other diseases end—in DEATH."

## SECTION II.—CONSTIPATION OF THE BOWELS.

By constipation is meant deficient action of the bowels, in consequence either of suppression or



diminution of the alvine secretions, or of sluggishness of the muscular structure of the intestinal tube, or both of these causes united.

With most persons the bowels are usually relieved once a day, either on rising or after breakfast, the impulse to this function being increased pressure on the lower bowels through the change of posture on getting up, or through the distension of the stomach by the first meal. But as there are some whose bowels are naturally disposed to act more than once in the twenty-four hours, so are there others to whom nature suggests one motion only in two or three days, or even at longer intervals. These peculiarities of constitution have to be carefully distinguished from the results of bad management of, or inattention to the right performance of, the excretory function. For it would be not less injurious to health, though in a different manner, to force one of naturally slow bowels to daily evacuations, than to allow a person with the usual tendencies this way to pass several days without the natural relief.

There are circumstances, again, that produce temporary constipation, which cannot be considered unhealthy. After purging, whether from medicine or otherwise, the bowels are usually confined for a day or two; nor is it necessary to take means to obviate this. In travelling, costiveness is commonly experienced; here, however, it is often con-

nected with a slight disposition towards feverishness; it is, therefore, prudent, when on a journey, to be moderate in diet. With many a sea-voyage produces continued constipation of the bowels, which is not found inconsistent with perfect health. Violent exercise again, especially if systematically continued, as in training, has an effect of the same kind; in the latter case, the superfluities of the body are carried off by another channel.

As a general observation, it is certain that a disposition to costiveness goes with physical strength. And this is easily accounted for; there is so much the less drain upon the system. But there are instances in which costiveness is associated with weakness; yet in which it is not only salutary and natural, but irremediable. In these cases, the suppression of alvine discharge depends upon organic feebleness of the bowels; which instead of having their proper strength and fabric, are, as it is found upon dissection, thin and structurally weak.

I attended a young person, a few months before her death, for what her friends were obstinate in supposing obstruction from local disease of the lower bowel. A year before, she had had scarlet fever, which left her debilitated, and liable to an occasional spasmodic difficulty of swallowing. She was very tall, thin, and extenuated, and weak; food went against her, so that she took scarcely any; once in a week or ten days, she passed a minute

quantity of dark green fæces. Dr. Merriman afterwards saw this patient. She died, after having gradually wasted to the utmost degree of emaciation. On the inspection of the body, no appearance of disease was found, but the intestines were remarkably *thin*.

The following case exemplifies similar features, namely, irremediable inability of the bowels to secrete, united with general weakness of the frame, not leading, however, as in the former instance, to a fatal atrophy, but allowing a permanent state of subdued health and strength, varied from time to time with alarming indisposition, of which the source is the suppressed intestinal function.

I was requested to see a young medical man, who, I heard, was in a fit. I found him lying on the floor, sensible, but exhausted with suffering; the flexor muscles of the limbs, and the muscles of the abdomen, were in strong spasmodic action. He had been in this state for several hours. Ammonia, and hot brandy and water were given him, and he gradually rallied. This, I learnt, was not the first seizure of the kind which he had experienced. Attacks of a similar description, but of less severity, would come on several times in the year; they were preceded by obstinate costiveness. This patient, now twenty-eight years of age, up to the age of fifteen, enjoyed excellent health. At that age, his bowels fell into the state of costiveness, which has

continued since. He grew up of a slight and delicate frame, physically incapable of much bodily exertion, and indisposed to it by a languor and drowsiness which probably arose from the imperfect action of the bowels. The bowels now act once in five or six days only; what is then passed is healthy; it is only extraordinarily deficient in quantity. With this he has little appetite; and even that he is afraid of indulging, lest it should lead to one of the attacks which I have described. These attacks, it has been mentioned, occur when the bowels have been confined for an unusually long period. The belly then becomes hard, and a little swollen; there is sickness, but nothing is thrown up but what has recently been taken into the stomach. There is a sense of uneasiness and pain above the navel. When at the close of such an attack, the bowels are relieved, the motions which pass are still extremely scanty.

There are other cases, equally rare with those which I have now cited, but of greater interest on account of the probable recovery of the patient, in which costiveness occurs in connexion with weakness and want of nutrition, from functional, and temporary, but salutary inactivity of the bowels. The patient, either after illness, or without anything to lead to it, falls into a state of declining strength, yet no evidence of inward disease can be detected. The bowels, however, are habitually

confined, and the motions scanty and insufficient. In cases of this description, the prescriptive practice is to combine invigorating diet and tonics with aperient medicine. The patient, however, instead of thriving upon this system, is liable to become weaker; but recovers when the bowels are left to themselves, and nature is allowed to recruit herself in her own way.

Ordinary cases of constipation, those in which the affection is prejudicial to health, and requires to be remedied by medical interference, differ among each other, as they depend or not upon an assignable cause, in the extent to which they derange the health, in their specially affecting this or that function, as they amount to complete or incomplete suppression of secretion, as they bear a chronic form, or threaten the invasion of acute disease, as they occur in infancy and youth, in middle life, in age.

The commonest form of constipation is that which results from inattention to regular times of relieving the bowels. Men of literary pursuits are naturally more prone to this error of self-management than practical men. All the personal habits of the latter are necessarily more methodical. So every variety of general and local disorder of the stomach and bowels is more prevalent among the former. When costiveness proceeds from this cause in persons who live temperately, and take sufficient exercise, its usual consequences

are one or other of the local affections of the lower bowels, which will be presently treated of.

A common source of costiveness is neglect of exercise. The circulation, in such cases, becomes loaded, the bowels torpid, the tongue furred, the appetite impaired, the patient suffers depression of spirits, clouded intellect, headach, habitual drowsiness and heaviness, and indisposition to exertion.

Want of bile is a frequent source of costiveness. This itself may proceed from neglect of exercise. But there are persons to whom disorder of the liver is the certain result of any departure from wholesome habits; of these, some, which I have already exemplified, suffer by profuse biliary secretion; others by deficiency of bile. In the latter case, the motions are clay-coloured, adhesive, and the bowels sluggish and inactive from want of the stimulus the bile affords.

Those who have resided long in hot climates are liable to costiveness from deranged biliary secretion. They experience languor, depression, uncertain spirits, uncertain appetite and digestion, joined with a furred and white tongue, uneasiness about the belly, flatulence, irregular bowels, with evacuations always scanty, sometimes firm and clayey, at other times loose, and like pitch. Nothing can be more uncertain than their state of health. When the weather is dry, and not too cold, they are often perfectly well, their spirits cheerful, their digestion

good ; on the same occasions, their bowels act properly. But in damp weather, especially when combined with cold, they rise in the morning unrefreshed, depressed in mind, without appetite, with a sense of weight, and uneasiness about the belly, sometimes with nausea, the bowels deranged as abovementioned. After dinner, the system is temporarily recruited ; the next day, the same sufferings occur. The remarkable influence of calomel and a regular course of aperient medicine, evinces, in such cases, that the disorder turns upon deficient alvine and biliary secretion.

Costiveness again occurs as a natural consequence of disease of the brain or spinal marrow. In children threatened with water in the head, in deranged persons, in those whose spine is affected, this symptom is a characteristic feature. The following is a striking example of the third case. I was consulted in the case of a young lady, one of whose symptoms was obstinate constipation of the bowels, requiring her to take daily from twenty to thirty grains of the compound extract of colocynth. She had been ill four years, and her sufferings had commenced with some pain across the belly, and costiveness. After a fortnight's illness, the constipation yielded ; but one leg became feeble, and the knee of that side was frequently spasmodically bent. This complication of palsy and spasms soon after affected the other leg : afterwards one

hand became feeble and contracted. These symptoms grew upon her; but she retained a remarkably fine complexion, and had all the appearance, as she lay on the sofa, from which she could not move herself, of a person in health. The vertebral column was perfectly straight and even; but she often experienced pain at the lower part of the dorsal portion, and pressure there gave her uneasiness. This circumstance, coupled with the other symptoms, led me to recommend that issues should be made at the lower part of the back. The remedy was followed by considerable relief. The legs seemed less weak, the knees were not so frequently nor so painfully contracted, and the bowels acted with half the usual dose of drastic purgatives. But the improvement was temporary; and, disappointed of obtaining permanent relief, this patient consulted other surgeons. She died six months afterwards. On examining the spinal cord, it was found softened for the length of two inches towards the lumbar end.

That obscure affection of the nervous system which is called hysteria, is often productive of obstinate confinement of the bowels. Costiveness, again, is liable to be caused by the water we drink, and the air we breathe. Chalybeate waters have this property generally, and a newly-painted house acts on some in the same manner. Frequently, however, there is no assignable cause for this dis-



order. The patient only knows that his habit of body has become confined, and begins sooner or later to experience some of the consequent evils.

Nothing is more capricious than the influence of costive bowels on the body generally. With some, the least deficiency of the customary relief produces sensible alteration of the health; with others, total constipation for days leaves the bodily sensations undisturbed. Yet it may be affirmed, that with the exception of certain alterations of structure, and the special results of morbid and atmospheric poisons, there is no local or general ailment which may not be produced by constipation. The organs which most commonly suffer through this cause are, the stomach, the lower intestines, the head;—so indigestion in all its forms, every disease of the rectum, headach, hypochondriasis, fulness of the cerebral vessels, palsy, apoplexy, are often direct consequences of confined bowels.

Constipation may be complete or incomplete, and if the latter, may exist in every degree. When complete, or nearly so, in some cases it will continue for weeks, producing very slight disturbance of the system, a little headach, and loss of appetite, perhaps, and having come on gradually, will gradually and unaccountably go away, seemingly having been little influenced by remedies. When incomplete, it may be temporary, and occur occasionally only; or it may be troublesome for years

or perpetually recur as the commonest feature of a life of ailment and disorder.

That which gives the most serious complexion to confined bowels is, that the symptom may proceed from another cause than costiveness. There may be no lack of intestinal secretion, no general inability of the bowels to propel their contents, but still there may be no passage. Obstruction in one of its numerous forms may be present; and that which seemed at first costiveness, may soon, upon the rising of other symptoms, upon pain, tenderness of the belly, vomiting, supervening, take a more alarming and different character. Into these cases I do not enter on the present occasion; I have endeavoured to point out all their varieties in my "Outlines of Pathology;" here I can only advert to this branching of the subject, and the alliance of obstruction of the bowels with constipation. The latter, however, as I shall have occasion further on to exemplify, sometimes leads directly to the former; one cause of obstruction consisting in accumulation of fæces in the great intestine.

Viewed in reference to different periods of life, costiveness may be represented as common in infancy, and as then deserving prompt attention and correction from its relation to hydrocephalus; as common again in childhood, when still the same suspicion is to be entertained; common in youth

again; in general, however, as of less consequence in youth, and easily set right; often again in youth encouraged by neglect and inattention; and in young women dependent upon hysteria. Constipation in middle age is commonly not serious in degree, but often enough to impair the health, and contribute to a train of slight ailments that are directly attributable to insufficient alvine secretion: among these local uneasiness, or pain frequently recurring and often fixed in one part of the colon, is a common feature. In old age, the complaint either assumes a different character, often running into complete retention of masses of fæces in the intestine, which, weak and sluggish, is no longer stimulated by their presence to expel them; or re-assumes its original importance in relation to the circulation in the brain, leading to fulness of the vessels of the head, and cerebral effusion.

The treatment of costiveness, as it may be inferred from what has been already said, is a point of no small consequence in medicine and surgery; so many and so various are the conditions of the bodily health with which costiveness combines, and so many and so different the means of correcting it appropriate to different cases; so important often the consequences of neglecting this ailment; yet so important at times to disregard and permit its continuance; so important, on many other occasions, to use the principal means that are calcu-

lated to correct this complaint, when it does not exist. Or costiveness is commonly obviated by aperient medicine; but aperient medicine is of not less value in numerous cases when the bowels are not costive. Or it may be more just to say that there are states of the system, in which the ordinary action of the bowels is, relatively to the relief required for health, constipation, and in which health is to be maintained or restored by systematic purging. It has proved a very profitable vein of quackery to employ this principle under a disguise, and to vend medicines, that are merely purgative, as a panacea for every complaint. The success of the imposition has turned upon the fact, that when the system is out of order, and there is no failure of stamina, in three cases out of four it is only necessary to purge, that is to say, to excite great increase of secretion in these organs, and the disorders of the rest subside.

I cannot better describe the treatment of constipation than by enumerating the different means which are available in it, and explaining to what cases each is appropriate.

It should, however, be observed, to begin with, that, independently of the cases already mentioned, where costiveness is directly salutary, and in persons of average health, constipation of the bowels occasionally supervening without sensibly impairing the health had often better not be interfered with

at all. In trifling ailments, Time is the great physician. Health is never stationary, but has its natural ebbs and flows; and in the wholesomest state of the animal œconomy, there are slight disturbances or intervals of less perfect condition of the bodily functions, from which it is best to wait their spontaneous recovery.

Where costiveness proceeds from known causes, the means of removing it are of course evident. If this ailment has supervened on neglect of bodily exercise, bodily exercise should be resumed to cure it. For the strong, an hour's walking or riding before breakfast is not only most salutary, but is likewise compatible with most occupations and professions. For persons of less strength,—for women generally,—exercise in the afternoon is preferable; the same is requisite in addition to morning exercise for the strong. So a late hour of dining, in summer, seven o'clock, for instance, is a wholesome arrangement, allowing from ten to five for business, and leisure for exercise after it.

Where costiveness is joined with inattention to regular times of relieving the bowels, it will often cease upon observing regularity in this respect.

The disposition to costive bowels may often be obviated by using articles of food which are laxative. Fruit generally has this effect; but all fruit is not equally good for the purpose. The best are grapes, currants, strawberries, raspberries, goose-

berries, eaten at breakfast. Next, peaches, nectarines, pears, apples, which may be conveniently eaten as luncheon. Fruit is certainly less wholesome after dinner; but when baked or otherwise cooked, it may be eaten as part of the repast. Cherries and plums are perhaps the least wholesome; they are liable to purge. Currants again, which agree with most people, with some produce heat of the lower intestine and hemorrhoids by their seeds lodging there. All fruit likewise, as nature intended, is best used for a season only; the wholesomest when pursued too long, will disagree either with the stomach or bowels. A gentleman, eighty years of age, who has frequently consulted me, being of a very hale constitution, has costive bowels, which in the early part of the summer he finds easily manageable by eating fruit, principally currants. One day in the middle of August, two or three years ago, he had eaten currants with his breakfast; the bowels had afterwards acted properly, when in the forenoon, with no previous warning, he dropped down in a state of insensibility; he continued insensible for half an hour; then vomited a quantity of half-digested currants, and soon completely recovered, no indisposition beyond languor and some weakness being left.

Vegetables are generally laxative; many find brown bread answer the same purpose. When they agree and answer, these are unobjectionable means. But both are liable to disorder the stomach.

The next means we may consider are such as are purely mechanical. Those in whom fæces are formed in adequate quantity, but are not voided through sluggishness of the muscular structure of the bowel, or because there is not liquid enough, often obtain perfect relief by the use of injections of water. The water should be used cold, to be more stimulating; to render it still more so, half a teaspoonful of salt may be added. The quantity ordinarily requisite is a pint. It is sometimes of course preferable to use warm water. Many persons who are temperate livers, and of sedentary habits, find the daily use of such an injection half an hour before the customary time of relieving the bowels, sufficient to obviate troublesome costiveness, and to cure the local complaints it has produced. But this remedy does not answer with all. In some, it produces uneasiness and a sense of weight and dragging from the loins, with general lassitude. In others, the bowel becomes relaxed, and a disposition to protrusion is formed through it. Nor is the practice of injections wholly free from more serious accidents. Sometimes the intestine has been torn, and the gravest consequences have ensued, through awkwardness in the use of the instrument. The tube of the apparatus is generally made so long and sharp, that it is easily capable of tearing and lacerating; properly it should not exceed an inch and a half in length, and should end in a smooth

sphere, half an inch in diameter, tapering to a neck a third of an inch in diameter.

Electricity has been successfully made use of to produce relief of the bowels.

A gentleman had been under the care of the most eminent physicians in England and Ireland for an obstinate state of the bowels, which was originally ascribed to having slept in a newly painted room. From being of a full habit, he became greatly emaciated, and the complaint went on in this manner for two years. Dr. Cheyne then recommended galvanism, which in about three weeks restored the natural action of the bowels, and he soon recovered perfect health.

Cold-bathing, either the plunge or shower-bath, is an important indirect means of obviating costiveness, where it forms part of hysteria. The hot-bath is of great use in cases which present an acuter character, threatening obstruction.

These serve occasionally. But the means of obviating costiveness most commonly to be resorted to, consist in the use of purgative medicines. The classes of cases to which they particularly apply are those in which the intestinal secretions are not formed in wholesome quantity. But they are sometimes in addition the only resource in cases of the other class. These remedies may be enumerated under the heads of laxatives, drastic purgatives, mercurials.



As laxatives, any of the following medicines may be safely tried, but generally one form is found to suit each case better than another.

It is to be observed that aperient medicine may be taken either in the afternoon or at night, or in the morning early, and that there are circumstances which render in different cases either plan preferable. In persons who are liable to suffer with indigestive pain at night, it is as well that the bowels should be habitually relieved before retiring to rest. For this purpose, two or three grains of compound extract of colocynth, taken in the afternoon, or at tea, are often sufficient. Or a short time before dinner may be taken, in two pills, two grains of extract of chamomile, five of extract of rhubarb, one of carbonate of soda; or in a single pill four grains of the extract of aloes with one of mastich.

To produce relief of the bowels in the morning, there may be taken at bed-time either a spoonful of the lenitive electuary; or, in two pills, three or four grains of compound extract of colocynth and castile soap, of each equal parts. It is sometimes necessary to warm these laxative medicines, which else produce uneasy sensations in the bowels, flatulence, and griping; for this purpose a grain of ginger, or half a drop of the essential oil of cloves or caraways, may be added. With some two or three grains of extract of hyoseyamus render the action of laxative medicines more pleasant.

With some, to whom aperient medicine is necessary, it answers better to have the inside quiet during the night, and in the morning to administer the dose, which may consist either of a drachm of Epsom salts, and two drachms of tincture of rhubarb in peppermint-water, or in infusion of roses; or of one to two drachms of Cheltenham salts in half a tumbler of spring water; or (to combine a tonic) equal parts of infusion of senna and orange-peel, with half a drachm of Epsom or Rochelle salt, and five grains of carbonate of soda; or using the effervescent form, of two drachms of Rochelle salt, with two scruples of carbonate of soda in half a pint of water, five and thirty grains of tartaric acid being stirred into it immediately before taking.

The occasions on which laxative medicines are resorted to are mostly these. Every one living the artificial life of cities, occasionally finds himself out of order, he has lost his usual appetite, his tongue is furred on rising, and he is disposed to expectorate tenacious mucus, with a sense of nausea leading to retching; the bowels are confined or irregular; one or two doses of aperient medicine remove the indisposition. Many suffer from local diseases of the lower intestine, which are aggravated by costiveness, and relieved or cured by the use of aperients. Those again who have met with bodily injuries, fractures of the limbs, for instance, requiring absolute rest, fall into costiveness, and

then into heat and fever, unless protected by laxative medicine. Finally, in almost all chronic, and in many acute ailments, manifest improvement follows upon slightly increasing the natural action of the bowels.

Drastic purgatives are required by some, where aperients suffice for others. Otherwise the proper use of drastics is not to obviate costiveness, but to lower and reduce (when reduction is necessary,) by the quantity of secretion they carry off. So in many disorders, purging is of the greatest utility. As a drastic purgative in costiveness, gamboge, scammony, jalap, compound extract of colocynth, with Castile soap, of each two grains, to be taken at night, form an useful combination; it is better the following morning to administer a saline aperient draught. Castor oil, four to six drachms for an adult, is one of the easiest medicines of this class; but it does not agree with the bilious, and with many is apt to heat the lower intestine; to children and infants it is very salutary. The oil of croton is invaluable in cases of protracted costiveness that have resisted other means; such are often met with in aged persons, sometimes in middle life; a drop in a bread-pill repeated every three hours, will seldom fail to accomplish the desired object.

Mercury is of great but partial use in removing constipation. There are many who cannot bear

the action of this drug, in whom a single dose will produce the greatest languor and depression, or violent bilious disorder, or headach and sickness, or even salivation. The inquiry should, therefore, always be made before mercury is prescribed, whether it is known to disagree with the patient. There are others, again, whom mercury always suits; whom it serves as the best alterative, the best aperient, and, indirectly, the best tonic. Nor is it possible to tell, beforehand, with whom this remedy will agree or disagree; but to scrofulous persons, and to the highly bilious, with the nervous and to persons with weak stomachs, it should be prescribed with caution.

The preparation of mercury used will often make a difference; those whom a grain or two of calomel would lower and distress, are often set right of slight indigestion or costiveness, by two or three grains of chalk and mercury or blue-pill, alone, or joined with three or four grains of extract of henbane, or of rhubarb. These gentler doses are often of the greatest use to scrofulous children, whom more active mercurialization would greatly injure. On the other hand, some children are lowered by chalk and mercury, with whom calomel agrees. Calomel in general agrees with children (and in such should be usually given when purgation is required) who have a disposition to water in the head; it commonly again answers remark-

ably well in the chronic impairment of the stomach and bowels produced by residence in hot climates; and is the first medicine to be thought of in all cases where there is a deficiency of biliary secretion. The *ratio operandi* of medicine is seldom satisfactorily traced; but mercury unquestionably acts on the nervous system, on the heart, on the mouth and salivary glands, on the skin, on the liver, on the bowels. Upon the liver, however, I believe that mercury acts indirectly; and that as regards this organ, its influence is derived from its causing bile to accumulate in the blood. When there is more bile in the blood than natural, one of two events in time must follow; either more bile must escape by the liver or the body will be jaundiced. But it is practically found that a moderate increase of the quantity of bile in the blood commonly leads to the first result, and forces a passage for itself out of the liver instead of into the tissue of other organs. Consistently with this view, I believe that mild cases of jaundice are benefited by mercury, while severer ones are much aggravated.

---

## CHAPTER III.

## DISEASES OF THE RECTUM.

THE lower termination of the alimentary canal is the seat of a variety of troublesome complaints, the disposition to which is referable to the following causes. Above this organ are placed masses of bowels, the circulation in which forms one great system, having its most dependent part here. The blood, therefore, gravitates to the rectum, whenever the body is upright, so that, under the most favourable circumstances, the hemorrhoidal\* veins are exposed to more strain and pressure than most other veins. And when it is recollected that cold, passions of the mind, irregularities in diet, tend to throw the blood inwards upon the bowels, it will easily be understood that there may often be congestion enough of these vessels to determine local swellings, inflammations, effusions.

The main channel, again, for the return of the blood from the bowels, is through the liver, and it is hardly too much to say, that every disturbance of the function of that organ acts as an impediment

\* The veins of the rectum are called hemorrhoidal, from their disposition to bleed.

to the passage of the blood through it, and tends in the same degree to increase the congestion of the vessels of the lower bowel. In addition, the greater sensibility of this part of the canal, the acrid nature of its contents, the peculiar muscular structure with which it terminates,—all contribute to heighten its liability to disease, while its situation exposes it to several kinds of mechanical injury.

There are, however, rules, deducible from these same considerations, the observance of which tends materially to prevent the occurrence of many of the troublesome complaints which are to form the subject of the present chapter.

I. Abstemiousness in diet, and temperance:—which, recommendable on every other ground, are essential on this,—that as their neglect gorges the vessels of the bowels with blood, the effects of which are, as it has been explained, most felt at the lowest part.

II. The prevention of costiveness by adequate means:—of which the various kinds have been described in the preceding chapter. Costiveness is productive of disease of the rectum;—first, by causing general fulness of the abdominal system of vessels; and secondly, by the straining to which it leads.

III. The use of soap and water, every time the bowels are relieved, so as completely to purify the part. This rule, which is one of common cleanliness, is sufficient of itself to prevent half the more trivial ailments about the rectum.

The disorders which are incurred through neglect of these rules, or through original tendency to disease, or casual lesion, together with the different kinds of congenital malformation, will be considered under the following heads. Fulness of the hemorrhoidal vessels and hemorrhage; piles and excrescences; prolapsus, or protrusion and eversion of the bowel; fissure of the mucous membrane; spasm and contraction of the orifice; stricture of the rectum; obstruction of the passage of the bowel by its contents; want of perforation at birth; laceration of the bowel; abscess and fistula; cancer.



SECTION I.—FULNESS OF THE HEMORRHOIDAL  
VESSELS AND BLEEDING.

THESE disorders require to be considered together, as they frequently go together, the second being often at once the effect and in some degree the natural relief of the first. They are liable, however, to occur separately,—that is to say, congestion of the vessels of the rectum is not always followed by bleeding; and, on the other hand, bleeding from the rectum is not necessarily preceded by symptoms of local vascular turgescence. So, it is desirable to look at the subject in three points of view, and to consider separately, hemorrhoidal congestion, hemorrhoidal congestion with bleeding, and bleeding from other causes\*.

1. Any cause whatsoever that disturbs the body or mind, is liable to produce the first affection. The common causes are, errors of diet, neglect of aperient medicine and exercise, or exposure to cold. But without any assignable cause, this complaint often makes its appearance.

The symptoms of hemorrhoidal congestion are sensations of weight, aching, fulness, heat at the fundament; the aching pain sometimes extends to the loins, and round the lower part of the belly,

\* I here pass over the consideration of hemorrhage from ulceration of the bowels in fever, and in phthisis, from the action of poisons, from local injuries.

sometimes down one or both thighs. These symptoms supervene gradually, are at their height in twenty-four hours, and remain troublesome two, three, or four days: then gradually wear off. The proximate cause of these symptoms is local determination of blood, or something approaching towards inflammation. The vessels are turgid, the part hot, as in an inflammation, and there is a disposition to local swelling from effusion of serum and lymph.

The means which relieve an attack of this kind are the following:—rest, and that in the recumbent posture; abstinence from wine and heating food; laxative medicine, such as one or two drachms of lenitive electuary, an ounce and a-half of infusion of rhubarb with three drachms of manna and a drachm of tartrate of potass, or a Seidlitz powder; the local application of cold or warm water, whichever being tried gives the most ease. Generally cold applications are preferable to warm. If the pain is severe, leeches are to be applied.

2. Congestion, with bleeding, is the addition of that symptom to those which have been described. The bleeding is, in some cases, to a certain degree accidental,—that is to say, it only takes place upon the action of the bowels, when the strain upon the turgid hemorrhoidal vessels is increased, and the mucous surface is compressed. If the bleeding is inconsiderable in quantity, the fact of

its occurrence should not modify the treatment of the complaint. The bleeding often manifests itself at the commencement of the attack, so as to appear to the patient to be its cause.

Congestion with hemorrhage sometimes puts on the character of an acute febrile attack. The patient experiences shiverings, followed by pain in the loins, and a symptomatic fever, which, after a few hours, is relieved by a copious discharge of blood from the rectum. Such attacks are not frequent, and their nature is seldom ascertained before the critical hemorrhage occurs, which relieves them. The proper treatment of this case antecedently to the hemorrhage, is the local application of leeches, and saline aperient medicine.

Congestion with hemorrhage often recurs to a slight extent, lasting two or three days, whenever the system is overcharged and disordered. The complaint has then the use of clearing the system, and probably often averts some other attack, and is so far salutary. It is not, however, by any means safe to rely on this mode of relief, and to calculate upon so compounding for a course of indulgence. The same congestion, which leads to salutary bleeding, is liable, at the same time, to lead to structural disease. Occasional hemorrhage from the bowel, if moderate, is least to be interfered with in persons who have recently spit blood, or who show a tendency towards apoplexy.

3. The cases in which the hemorrhage is out of all proportion to the local congestion, and is the only local symptom which presents itself, are of more consequence than those yet adverted to.

The quantity of blood which is sometimes thus lost is enormous: four pints have been passed in a few hours: a pint daily for a month; these are, of course, extreme and rare cases: but to a less, though highly prejudicial extent, such hemorrhages are common, producing an exsanguinated appearance, weakness, swelling of the legs, palpitation of the heart, headach, faintness, vertigo, extreme nervousness, spasmodic difficulty of swallowing, and the whole train of like symptoms.

The source of these great bleedings is the same as of the lesser: there is no rupture of vessels; the blood transudes through their coats, and pours from the whole surface, but commonly from some points more than from others. It is just the same in bleeding from the lungs, or from the nose. No vessel is torn, but the blood passes freely through the thin membranous tubes, which ought to contain it. Such bleedings, again, are in part only local ailments; the blood, indeed, finds vent at one part in preference to another, but that which makes it flow at all, is some general influence, and the place of the hemorrhage is comparatively accidental. Accordingly, in the most remarkable cases of the class, the discharge of blood from one part of the

general inward mucous surface will sometimes cease, and commence upon another: after hemorrhage from the bowels has ceased, bleedings from the nose will supervene: these may alternate with vomitings of blood: these, again, with spitting of blood from the lungs. All these ceasing, the patient may have apoplexy.

But of these different constitutional hemorrhages, that from the bowel is the least locally injurious and troublesome. Blood poured out in the brain causes cerebral apoplexy; in the lungs, pulmonary apoplexy; in the stomach, vomiting and disturbed digestion; from the bowel it passes away leaving no local consequences. In such cases, therefore, it becomes a question, whether any attempt should be made to suppress the flow of blood, or whether pains should not rather be taken to keep up, or even to restore, this comparatively innocuous vent.

There are instances, in which great judgment is required to determine which course should be pursued. When the habit is full, and the disposition to hemorrhage strongly marked, for the time the hemorrhoidal discharge may require to be encouraged. Generally, however, this is not the case; but the two objects may be pursued together,—of suppressing the flow of blood, and of correcting the plethoric condition of the body.

There are two means by which hemorrhage from

the rectum may be stopped. The first is applicable to all cases, in which it is safe to attempt to arrest the bleeding; it consists in the use of cold, and local styptics. These are applied in the following ways:—by bathing the part with cold, and even iced, water, by daily injection of cold water, or of a few ounces of infusion of catechu, or of the same, with a few grains of sulphate of zinc or of alum. Astringent injections are not to be used in a greater quantity than five or six ounces; the best time for their application is before rising, when they are retained longer. Or a suppository of Ward's paste and soap, in equal parts, may be introduced into the rectum at night, or a drachm of the same medicine may be swallowed in pills, two or three times a day.

A gentleman, who has consulted me, has formerly spit blood, and is now liable, on very slight occasions, to pass blood from the bowel: he labours under habitual protrusion of the lining membrane when the bowels act: this, if roughly treated, always bleeds: he, therefore, never uses, after an evacuation, any means to replace the protrusion but washing with alum-water, and in this way obviates the recurrence of hemorrhage.

The blood which flows from the rectum is of a bright scarlet colour: in some cases of protrusion it may be *seen* to exude from the vascular mucous surface. It often appears to flow from one or two

points more than the rest, and spins out freely as if from a ruptured vessel. This has led surgeons, in some cases, to apply a ligature to the bleeding point; nor wholly without advantage, when the patient is becoming daily weaker through loss of blood. For a few days afterwards the bleeding does not return, owing to the slight inflammatory action which supervenes, and lasts till the ligatures are thrown off.

In general, when there is a strong disposition to hemorrhage from the rectum, any mechanical cause will excite it,—the fæces being firmer than natural, for example; and, on the other hand, the use of drastic aperient medicine.

Nevertheless, cases of hemorrhage from the bowel are occasionally met with, unattended with local heat, not excessive in quantity, and in persons of a strong and rigid fibre, where the only successful treatment is repeated purging.

It is needless to add, after what has been said, that those in whom an habitual or periodical hemorrhage from the bowel has been stopped, require to observe the greatest caution as to diet, use of aperient medicine, and the like,—to prevent the circulation becoming over-loaded, and the occurrence of hemorrhage in more important organs.

## SECTION II.—OF PILES.

PILES are soft tumours, generally round, sometimes flattened, sometimes with a broad base, sometimes attached by a neck, varying from the size of a pea to that of a French walnut, which form within the lower intestine, and at or about its orifice.

Sometimes they are situated entirely without the bowel, and are covered by the common integument; these are called *outward piles*, or *skin piles*.

When they form within the bowel above the sphincter muscle, they are of course entirely covered by mucous membrane; these are called *inward*, or *intestinal piles*. They are liable to be forced down when the bowels are relieved, so as to protrude externally; but their appearance is so different from that of outward piles, and their place of attachment and origin is so easily determinable by examination, that it is hardly possible to mistake the one for the other. They are soft, with more or less firmness, and red, blue, or brown in colour, under different circumstances.

Sometimes piles are situated exactly at the margin of the intestine, and are wholly without the sphincter, or nearly so, forming swellings with this peculiarity, that they are half covered with mucous membrane, half with skin. These are marginal or *anal piles*; they are sometimes confounded with inward piles.



These swellings, whatever situation they occupy, consist of the following elements: first, of a growth or production of skin or mucous membrane, which forms their covering; secondly, of a growth of the minute vessels, which goes with the growth of the flap of mucous membrane or integument; thirdly, of dilated veins, which are partly produced, like varicose veins of the legs, by the pressure of the column of venous blood in the intestines, partly by the local increase of vascularity; fourthly, of inflammatory products, lymph or serum, which infiltrate the tissue of the flap of skin or mucous membrane, giving to it more or less firmness. Piles in different cases present one or other of these elements more prominently than the rest. Some are merely little pendulous flaps, others have the highest vascularity in addition, others with some thickness are soft, and are blue with dilated veins, others again are of a firm consistence, whitish if external, brown if internal, from effused lymph or serum.

Piles frequently occur in persons who are liable to the attacks of vascular fulness already described, which are thence often miscalled attacks of piles. But the piles are generally there before the attack, although they swell and become larger during the attack; they continue for a time swollen after the increased heat and pain have subsided; they then gradually become less full and tense, and are not more sensible than the surrounding parts.

Piles have been divided into bleeding piles, and such as do not bleed; the bleeding, however, has little to do specifically with the piles. If they are external they do not bleed (for blood most rarely transudes through the skin); if they are internal, they bleed only in connexion with the disposition to bleed common to the adjacent mucous surface.

Or the tumours called piles bear the following relation to the congestion described in the preceding section. They sometimes arise from it; when they already exist they participate in it, and are the principal seat of the pain; and they probably render the attacks of congestion more frequent than they otherwise would be.

The progress of this complaint presents several varieties. Or this is common to each sort; the lump which forms the pile may come either suddenly or gradually; its condition is always fluctuating, sometimes more uneasiness being present than at other times; but the alternations with the attacks of pain are often remissions only, and there is constantly present greater or less uneasiness. In other cases, and these by far the most frequent, there are intervals of entire absence of pain, in which the lump, now less than before, has not more sensibility than the neighbouring parts; finally, the lump may continue for a lifetime; or may disappear by suppuration, forming a small abscess; or, what is more common, may waste, losing its

firmness and interior hardness by absorption, and becoming only a thin flap of skin or mucous membrane with no disposition to pain or action of any kind.

I shall describe the three varieties of piles separately.

*External piles*, of the kind at least that is most common, are little lumps, generally oblong, and more elevated than thick (resembling, for instance, in shape, the seed of a lupin adhering by its concave edge), and from that size to the magnitude of an inch in length, half an inch or more in height, and the same in breadth. They are laid generally laterally to the orifice of the bowel. Sometimes there is one only, more commonly there are two or three. Sometimes they are flatter. They are covered by the common integument, of which they have the colour. In their quietest state they are soft flaps of skin and membrane, having common sensibility; when angry, they are firm, hard, and exquisitely painful. There is commonly an intermediate state of heat, itching, and irritation, between the quiescent state and the most aggravated condition of these tumours.

There are several modes of treating external piles. When they are unattended with pain, the best way is to leave them alone, only scrupulously using soap and water after each action of the bowels, as the irregularity of the surface makes it

otherwise probable that some impurity would remain, and it is certain that nothing contributes to the growth of these tumours more than this cause. It is unnecessary to repeat that the greatest attention should be paid to the action of the bowels. When the swellings are uneasy by fits only, and this often happens for two or three days together, unfitting the patient for his customary avocations, the best way by far is to remove them by an operation, which is painful indeed, but is over in a moment. They should on no account be removed by ligature, which is slow, very painful, and even uncertain, but are to be cut off with scissors, or with a scalpel. The part has then to be poulticed for two or three days, during which the patient should remain in the recumbent posture. If the patient has not nerve enough to submit to the momentary pain of the operation, the next best thing is to apply pressure. This may be made by means of a metal cylinder, like a ruler, an inch and a half in diameter, on which the patient is to habituate himself to sit. At first the pressure is painful; but use renders it less so, and the tumours often waste under it. When the pain arising from outward piles is constant and severe,—and it sometimes will continue incessantly for months,—the same plan, with cooling lotions, or cold water frequently applied, is still available. But it is far better in such a case to remove the tumours at once. The

operation is not followed by bleeding; nor have I ever known any bad consequences ensue, but, on the contrary, constant and almost immediate relief, even when the swellings have been cut off in their angry stage.

Under the head of external piles, another complaint, rare, indeed, but extremely troublesome, has to be grouped. Its distinctive feature is intolerable itching, which is not constant, but comes on occasionally during the day, and invariably during the night. When the part is examined in its quiescent state, nothing is to be seen which at all accounts for the severity of the patient's sufferings. The skin about the anus is not raised into permanent lumps or ridges, but looks healthy. But it has acquired properties analogous to those of erectile tissue. When the period of heat and irritation supervenes, the marginal skin of the anus becomes swelled out in two or three ridges, one of which, perhaps, extends to, and involves the raphe of the perineum, and sometimes one or two little growths of skin, or pendulous flaps, that were loose and irritable before, shoot up, curling in a distended and angry state. The mode of treating the disorder is by cooling medicine and diet, and the avoidance of all that heats the body; a course of sarsaparilla, combined with an alkali, I have known useful; locally, a wash containing hydrocyanic acid is the best application. This is a complaint occurring in middle

life, and different entirely from one of the most distressing complaints of age—pruritus senilis.

Sometimes crops of arborescent warts grow from the circumference of the anus; these are sore and painful, and can never be perfectly cleansed. The attempt to destroy these with caustic or the ligature, is seldom more than partially successful; they should be taken off with scissors or with the knife, and the trifling bleeding which follows stopped by the application of lunar caustic.

In siphilis, sore and uneasy tumours are liable to form near the anus; they are commonly circular, half an inch in diameter, flat, with a white moist surface, raised a third of an inch; these are spots of psoriasis, which the warmth and moisture of the surface develope to this magnitude and character.

*Anal piles*, or those which are situated exactly at the termination of the intestine, are more troublesome than the preceding kind, and they present several varieties, between which it is very important to distinguish.

The character common to each variety is this, that it is situated on one side of, and exactly at the margin of the orifice of the bowel, so that it has one part towards the cavity of the bowel, and covered with mucous membrane, the other half being outward, and covered by the common integument.

Anal piles vary in magnitude from that of a pea to that of a chesnut. The larger they are, the more painful they are found to be, for this reason: when they acquire any size, the internal part is within the grasp of the sphincter muscle, the pressure of which causes constant uneasiness.

In their quiescent state anal piles are little folds or puckers at the external border of the sphincter, where the mucous lining of the bowel is continuous with the skin; they have little or no hardness, and no sensibility; and in this state are best left alone, care being taken to keep the action of the bowels regular, and to use frequent ablution with a lotion of cold soap and water, two or three times a day.

When this kind or the preceding are attended with troublesome itching, saturnine applications may be used. I have seen much relief obtained from an ointment consisting of an ounce of lard, and two drachms of oxide of bismuth.

In their angry state marginal piles present the following appearances.

The tumour is sometimes a little hard blue knob, bigger than a pea, which is moveable, very tender on pressure, with a constant feeling of stiffness and aching in the part; or with the same sensations, two or three hard blue knobs appear set round the margin of the intestine.

The pain and tenseness of these blue tumours

lead one to expect that they would be relieved by puncturing with a lancet. This, however, is not the case. They are exquisitely sensible; puncturing them gives great pain, and increases the heat and aching of the part. Rest, cold lotions, or warm poulticing (as either is found to be more soothing), laxative medicine, leeches, if required by the severity of the pain, are the proper means of relieving such an attack; the common period of which is three to five days. The swellings at length becoming free from pain gradually subside to the state of flaps or puckers.

In this chronic state, they admit of, and require indeed, another and different treatment. They require treatment, from their disposition to fill again, and give renewed pain for a few days. Advantage should be taken of their chronic and indolent state to endeavour to disperse them.

The means for this purpose are local astringents and pressure. The best local astringents are nut-gall applied in an ointment, a drachm of nut-gall to an ounce of lard, or spermaceti ointment with ten or twenty drops of liquor plumbi to the ounce, or alum-water. To make pressure, the patient should either sit upon a metal cylinder as before described, or should use a short rectum bougie.

Another kind of anal pile is a lump about the size of a chestnut, occupying one side of the aperture of the intestine. It is not very prominent either



towards the cavity of the bowel, or externally, but feels of this size and shape when examined. It is tense, elastic, tender on pressure, and when at the worst, is the seat of constant aching and heat, with occasional throbbing. Rest, abstinence, laxative medicine, are the first remedies to be prescribed in this kind of attack, which often comes on very suddenly. As the tension of the part is considerable, warm poultices are more soothing than cold applications. Abstraction of blood by leeches gives relief, but not to the same extent as in the cases already described.

Such a swelling is occasionally mistaken for what it is not; it is liable to be considered an inward pile, that has been extruded from within the sphincter, and which would be relieved by being forcibly pressed back. Such an operation, however, would necessarily fail of any good result, while it would produce a serious increase of pain. A patient who consulted me for an anal pile of this description, had been told that by this means, if he could bear the pain, he would obtain relief. Luckily the remedy was not tried. I recommended the means already specified, rest, abstinence, laxative medicine, poulticing, and the tumour became softer, lessened, and finally dispersed.

Such a swelling bears a nearer resemblance to an abscess, from which it is occasionally impossible to distinguish it. There is a firm, elastic, heated,

painful, and throbbing lump; it may be a pile consisting of dilated and turgid veins and effused lymph; or matter may be forming in it. Supposing this doubt to exist, the best practice is still to poultice the part for twelve hours, *unless the pain is very urgent and intolerable*. In the latter case, the tumour had better at once be laid open with a lancet. The operation will give relief in either alternative. If there is matter, it is the best practice; and may prevent the formation of a fistula. If there is not matter, the puncture gives vent to clots of blood, which turn out of an inflamed vein, or out of the cellular tissue. The tension is thus removed, and the pain quickly subsides.

I believe that the pain of these swellings would always be relieved, and that they would always disappear sooner, if they were punctured. But there is an objection to doing this, unless the pain renders it necessary. The punctured wound is liable to bleed troublesomely. I attended, with Mr. Beck of Lambeth, a gentleman under these circumstances. A large and painful anal pile had been punctured, and a small clot of blood had escaped from it; but the wound continued to bleed, or the part to fill with blood, which came away partly clotted, partly fluid. This continuing, the patient was so weakened, that he fainted on the night-chair. Examining the part, I found that the hemorrhage was not venous, but proceeded

from an artery which had been divided by the lancet; the vessel was tied, and the bleeding ceased. However, there is very little risk of arterial hemorrhage, if the lancet is not plunged deep into the tumour, (which is quite unnecessary,) but rather used to slice it or make an incision through its entire length, a third of an inch in depth.

*Inward piles* are pendulous tumours, which appear to grow from the inner surface of the bowel above the sphincter. They commonly are situated near the orifice; or in that situation alone do they produce symptoms, and give trouble. These, when swollen, are forced out at each action of the bowels; when shrunk and quiescent, they are not so protruded, and are not felt by the patient.

As it has been explained, these tumours are growths of the mucous membrane, retaining more or less of its natural vascularity, and expanded, as a more or less tense sac, over some dilated veins and effused lymph. But, at first, these tumours are often folds of mucous membrane only; and when it has happened that they have become distended with turgid veins and lymph, the veins will yet again often shrink, and the lymph be absorbed; when the pendulous flap of mucous membrane alone is left. So the essential character of these tumours is, that they are constituted of little bags of the lining membrane of the bowel formed above the sphincter, liable to be more or less vascular, to be

more or less distended with turgid veins and interstitial lymph, and to be at times forced out from within the cavity of the bowel so as to protrude externally.

These folds of the lining membrane of the intestine I have seen assume the following different characters.

Sometimes they are of a florid red colour, soft, with very slight thickening, being enlarged folds of the lining membrane, which are forced out at each motion, and require some pressure to replace them. They have no increased sensibility, and give no pain; but they are troublesome from the care and time required in replacing them, and from the mucus which flows from them, and which continually escapes from the bowel, and from their disposition to bleed.

In other and more common instances, they project, as one or more pendulous tumours, soft, but of some thickness and substance, of a dark red colour, the mucous surface very little changed from the natural appearance. This is the ordinary appearance of inward piles when they are in a tumid and painful state. In the intervals they shrink.

In other instances the tumour, thin, long, and pendulous, appears as if the contents of the part adherent to the intestine had been absorbed, so that a soft, red, vascular knob is protruded from the bowel, with a thin peduncle of less

vascularity, an inch or more in length, which carries its vessels.

In other cases, the tumour is firm and hard, and mahogany coloured, the mucous membrane being stretched over a mass of lymph and compressed vessels, forming a round tumour from half an inch to an inch and a half in diameter; the tumour protruding each time that the bowels act; the pain and uneasiness it occasions being extremely variable.

Each of the preceding varieties which I have described, deserves to be considered separately, in reference to its chronic and habitual state.

The first is frequently called prolapsus of the lining of the bowel; but it is more than mere protrusion: it is protrusion with growth and elongation, and has much more affinity to other cases of inward piles than to true prolapsus. It is generally attended with little pain, but there is a troublesome discharge of mucus, sometimes of blood, and it is some time after an evacuation before the folds are returned. There are two modes of treating this complaint; one is by astringents, the other by the ligature. The first is necessarily palliative only. To give instances of both.

A gentleman, whose case I have before adverted to, formerly spat blood, and is now liable to occasional attacks of congestion of and bleeding from the bowel. Whenever the bowels act, a circular

fold of vascular membrane protudes, which, if roughly touched, bleeds. His constant practice is then to bathe it with alum-water till it returns; and I do not think that any further treatment would be judicious in this case.

A gentleman under forty years of age, who had passed several years in India, applied to me for advice for these symptoms. Every time the bowels acted, two or three small folds of mucous membrane were extruded; they were intensely vascular, not only red, but covered with visible dilated vessels. It took time to return the protrusion, which, however, neither bled nor was painful, but there was a constant discharge of mucus. He had tried cold and astringents, and various medicines, ineffectually. I therefore recommended him to have the principal folds removed by the ligature. The operation was attended with very little pain, and proved completely successful.

The second kind which I have enumerated, thickened and vascular piles, preserving the usual florid colour and surface of the mucous lining of the intestine, seldom require removal by the ligature. These are the kind most frequently met with. In their quiescent state, they have very little sensibility, and they are disposed to shrink and waste under proper treatment. This consists in the use of laxatives and astringents; a drachm of precipitated sulphur, or of lenitive electuary

taken at night, a drachm of Ward's paste taken twice during the day, with moderation in diet, will, in most instances, cause these tumours to waste and shrink, and become mere pendulous slips of membrane. In this case injections of cold water, half an hour before the usual time of relieving the bowels, are again useful. If the tumour, under continued treatment, does not waste, but continues the seat of occasional aching and uneasiness, the ivory bougie recommended by the late Mr. Mackenzie may be resorted to. This instrument is an inch and a half long, having an oval head exceeding three-quarters of an inch in its transverse diameter, tapering to a neck half an inch in thickness; it rests upon a convex plate of ivory, which is pressed by a bandage against the orifice of the bowel when the bougie has been introduced. At first, the use of this instrument is painful, but it becomes less so in a little time; and the pressure it makes diminishes both the size and painfulness of the tumours. All these means failing, the tumours may be removed.

External piles, it has been mentioned, should never be removed by ligature, as the skin which forms their covering is acutely sensible; they should be removed by the knife or scissors, as the operation so done is over in a moment, and there can be no danger of hemorrhage. With inward piles it is the reverse; when removed by excision, occa-

sionally the most troublesome hemorrhage follows ; when removed by the ligature, on the other hand, the pain is seldom very severe ; at all events, it is not more acute than that which removal by a cutting instrument occasions.

In removing inward piles by ligature, the following points should be attended to. If a large tumour is to be tied, it is better to pass a double ligature through its base with a needle, and to tie each half separately. And again, it is important before tying the ligature, to cut into the tumour. These precautions are necessary to allow the ligature to be drawn so tight as completely to strangulate the tumour, and to cause it at once to mortify. If the tumour is only half strangulated, the pain instead of subsiding continues till the fact is discovered, and the ligature sufficiently tightened.

In this form of the complaint, if the ligature is resorted to, it is generally unnecessary, when there are several tumours, to tie more than one or two ; the rest will waste.

This operation is attended with as little danger as any operation. Still it must be borne in mind that there is some. Once in a hundred times, fatal inflammation of the membrane of the belly follows it. Nor is it possible out of many patients with the appearance of good general health, to conjecture on whom the unfortunate lot would fall ; yet out of many, one or more there would be whose



health is wavering, and so insecure, that this weight thrown in the adverse scale is mortal.

Removal by ligature must be considered as the best treatment for the third and fourth kinds of inward piles which I have described. I conclude that there is less risk from the operation in those cases. The tumours have in both a character further altered from the natural structure. The operation again is so far necessary that it is the only mode of getting rid of the complaint; inasmuch, at least as far as my experience has gone, tumours of this kind do not waste, but remain permanently troublesome. All these circumstances being explained to the patient, it rests with him to decide whether he will bear the habitual inconvenience, or incur the very trifling risk attending the operation for its removal.

Looking at the varieties of inward piles in reference to their occasional changes, whether towards aggravation or otherwise, the following features present themselves.

1. They are, of course, liable to be involved in the general congestive state already described, to relieve which the recumbent posture, fomentation and poulticing, abstinence, and laxative medicine, are appropriate.

2. When in this state, they are liable to slough, which is the best thing that can happen. This change has a formidable appearance, but the

sloughing does not go beyond the tumour, which it removes and cures.

3. The congestive state of piles occasionally leads to suppuration in them, when they form little abscesses, which being opened or bursting spontaneously, the tumour afterwards shrinks and disappears.

4. After a severe congestive attack, piles are more likely to shrink and waste, than when they are in a chronic state of uneasiness.

5. Inward piles, when of large size, and casually swollen, sometimes cannot be returned by the patient when they have protruded. In this state they cause extreme pain, and excite inflammation and fever. By gentle pressure, the surface being smeared with cold cream, a surgeon finds no difficulty in returning such a tumour. This is a case in which, the irritation of the temporary strangulation having subsided, removal by the ligature should be strongly recommended.

### SECTION III.—OF PROLAPSUS.

THE lower intestine is liable to protrude, everted as you may evert the finger of a glove. This may result from two causes,—either from habitual straining, that forces the bowel out of the anus, or from relaxation of the sphincter, which deprives it of the natural support.

The cases of protrusion from unusual straining are again of two kinds,—in one there is adventitious local disease: in the other, none. The principal of the adventitious local diseases referred to, are piles, stone in the bladder, and worms. The first is not an uncommon cause of prolapsus in adults: the two last, in children. In general the prolapsus ceases when the adventitious disease is removed. If it should persist, either from the natural adhesion of the bowel to neighbouring parts being relaxed and elongated, or from dilatation of the sphincter, or from both causes united, it has to be treated by one of the methods presently to be mentioned.

The common cause of prolapsus, where there is no antecedent local disease, is costive bowels: the complaint is then one of slow growth. At first the protrusion is little observable, and returns spontaneously, immediately after the bowels have acted; but it goes on increasing; and after some months or years, a length of four, five, or six inches is extruded at every motion, to return which, some time, and the recumbent posture, and pressure, are necessary: add to which, there is a constant flow of mucus, and the bowel is liable to prolapse in walking or riding, or even on standing any length of time: and the constant exposure of the mucous membrane causes it to become sore and excoriated, and, at times, the prolapsed part will swell to a

volume which it is beyond the patient's power to replace, and he lies in considerable pain till the tumour can be returned by a surgeon. Even when the quantity of intestine protruded is not great, nor attended with much soreness or swelling, sometimes hours will pass before the patient can return the part himself. So I have known a patient, as a protection against the inconvenience of prolapsus, habituate himself to relieve the bowels once in forty-eight hours only, and that before retiring to rest, in order that he might have the night to allow the bowel gradually to become replaced.

For different cases of this complaint the treatment must be different.

When it occurs in children (in whom it is frequent) from costive bowels, it is generally only necessary to obviate the constipation by laxative medicine, and the disorder ceases. For this purpose, the lenitive electuary, or precipitated sulphur, or rhubarb with magnesia, are appropriate medicines. When, the bowels having been thus regulated, the complaint persists, the daily use of some astringent injection is advisable. Three or four ounces of lime-water, or of infusion of catechu, or of the same, with a few grains of alum, or of sulphate of zinc, will generally give tone enough to the part to prevent its reprotrusion. These injections should be administered before the child rises.

In grown-up persons, when the complaint de-

pende on costiveness, both the cause and the effect may often be together remedied by the daily use of an injection of cold water. When this is not sufficient, aperient medicine to regulate the bowels, with Ward's paste to stimulate the lower intestine, are the next remedies. The Ward's paste may be taken either in pills, a drachm three times a-day, or may be introduced into the rectum as a suppository made up with equal parts of soap. Or again, stimulating injections may be resorted to.

When, through delay, the complaint has become inveterate, and several inches of the bowel habitually protrude, and the anus has become dilated from the quantity of intestine that is frequently down, the preceding remedies often fail to produce an effect; then the management of the case becomes mechanical, or relief is to be obtained either by the use of different instruments adapted to support the bowel, or by an operation.

The means consist in making pressure upon the anus. This I have known successfully managed by an elastic belt passed over the shoulders and under the perineum. But the commonest expedient is a ball of ivory pressed against the anus by a spring like that of a truss.

In looking for an example of the advantages of this method, I do not find a better one than a case told to me, some years ago, by a gentleman with whom I have the pleasure of being acquainted,

not a medical man, in a conversation about this disease.

“Some years ago I travelled with a young Russian, a fine, well-built, handsome fellow (perhaps he was a cross-breed), who was so full of spirits, that he appeared to me a man under temporary excitement from some gladdening cause, rather than one swimming in the stream of his ordinary disposition. And so it turned out. He had been afflicted with a prolapsus for several years to such a degree, that his life was a misery to him. He could not take the slightest exercise, and was obliged to recline upon a sofa almost the whole day. Every thing had been tried in vain; but one day he fell in with a gentleman, who advised him to get a ball of ivory large enough to cover the anus, and fasten it so as to keep it always in that situation. Immediately he felt the greatest relief; the sphincter by degrees recovered its elasticity; he suffered less and less by the protrusion of the gut when he went to the water-closet, and when I saw him, he was able to take strong exercise without the slightest inconvenience: whereas, before the application of the ball, he could not walk across the room, without bringing on the complaint.”

In place of this expedient there are two operations suited to varieties in the case, which, in young persons, are a preferable alternative to constantly wearing an instrument.

One of these operations consists in pinching up two or three little folds of the mucous lining of the protruded intestine, on different aspects of the gut, and tying them with ligatures, causing the portions of mucous membrane to slough off, leaving little ulcers. The ligatures usually are loose in four or five days, and the ulcers commonly heal in a week. This operation causes the intestine to be sore, and to be temporarily thickened, and, therefore, to have less mobility, and to be less easily displaced than before, and the habit of protrusion is broken.

The other operation consists in pinching up with forceps three or four of the thin folds of skin which converge towards the orifice of the bowel, and cutting them off with scissors; the wounds heal in a few days. The effect of the operation is to narrow and to give due rigidity to the orifice of the intestine.

Both of these operations, which are directed each to a different element of the complaint, are necessary in the worst cases.

The following case, which was attended by Dr. Chalmers of Croydon and myself, and seen in the after-illness by Dr. Arnott, will exemplify the progress and successful result of the first of the two:—

A young lady, twenty years of age, had suffered for several years with headaches, torpid bowels, painful and irregular menstruation, pains in the

back and legs, irritability of the bladder, hysteria. Dr. Chalmers of Croydon, who was consulted, was led to conjecture that there must be some disease of the rectum, a knowledge of which had been kept back from her family. By closely questioning his patient, he at length learned that she laboured under prolapsus ani; that a protrusion took place every time that the bowels acted; that it was of considerable volume, and that it could not be replaced without difficulty. The young lady, it appeared, had suffered from this complaint as long as she could recollect, and as she grew up, the infirmity increasing upon her became a source of perpetual misery, which a false shame prevented her disclosing. The constant fear upon her mind that the complaint would become known, had probably contributed more than the local disorder to cause the train of symptoms under which she suffered. The mass which protruded when the bowels were moved by an enema was of the size of an orange; and as the coats of the bowel which formed it were not at all thickened, I have no doubt that the voluminous folds which came down, were an eversion of as much as six or seven inches of intestine. The sphincter ani was extremely lax: the eversion was ascertained to begin about an inch within it.

The method of treatment pursued, with its effects, was the following:—



A small fold of intestine was pinched up with forceps, and tied with a silk ligature; care was taken to include the mucous and submucous coats alone in the ligature: the whole surface included was less than that of a sixpence. Before finally tightening the ligature, the surface of the little fold was cut with scissors. Three such folds were tied upon opposite aspects of the bowel, and at different distances from the sphincter. The patient hardly felt the operation, so small is the sensibility of the internal parts of the body, unless when inflamed. The parts were then replaced. During the first four days which followed the operation, the patient was not allowed to sit up; and the bowels, which had been well unloaded before, were kept confined, very light and moderate liquid nourishment alone being allowed, and an enema of laudanum having been administered. Upon the fourth day, when the bowels were moved with an enema of warm water, the patient was greatly disappointed at finding that the protrusion returned; yet she remarked that the bowel admitted of being replaced with greater ease than before. Some soreness in the part, however, was now observed; blood too was passed from the bowel; there was irritation of the bladder; and for the first time, a protrusion took place during the act of micturition.

Upon examining the bowel on the sixth day after the operation, it was found to be in the fol-

lowing condition. The mucous membrane, when the bowel was prolapsed, appeared darker and fuller of blood. The little portions of membrane which had been tied had come away; the ligatures, however, had not yet separated, but remained fixed in the shallow ulcers which they had produced: they were divided and removed. From this time the local complaint improved daily: the protrusion became less and less, then did not recur each time the bowels acted, and in a fortnight had entirely ceased.

A strange accident ensued in this young lady's recovery; after a few days she became deranged; but the alienation did not last many weeks, and for some time that I occasionally heard accounts of her, she enjoyed perfect health of body and mind.

There are cases of prolapsus, which result from weakness and relaxation of the sphincter of the nature of palsy; this complaint occurs either alone, or in combination with weakness of the legs. In either case the only local remedy is the palliative one, already described, of the ivory ball and spring.

#### SECTION IV.—OF FISSURE OF THE RECTUM.

FISSURE of the rectum is a small ulcer of the mucous membrane, which is liable to form immediately above the sphincter, and is met with more commonly on the back part of the intestine, than in

front or laterally. The ulcer is generally oblong, a third to three-quarters of an inch in length. The edges are sometimes raised and hard; in other cases, their want of elevation and softness, and the shallowness of the ulcer, render the complaint difficult of detection.

The complaint usually originates in costive bowels, and its commencement often is a transverse rent of the mucous membrane from the pressure of hardened fæces in their expulsion. The crack so formed does not heal, but remains an ulcer.

The patient is sometimes aware at the time that some injury has taken place, from the unusual pain attending the relief of the bowels, and the soreness afterwards. In other cases the symptoms supervene gradually. They consist of pain, aching, soreness within the bowel, brought on by evacuating its contents, and remaining for a longer or shorter period, the pain being in time very severe, and a sense of weight and uneasiness being at last always present.

The simplest treatment is often sufficient to cure this troublesome complaint. It is frequently only necessary to keep the bowels slightly relaxed, the motions semiliquid; and the ulcer will spontaneously heal, when no longer torn open daily by the passage of hardened fæces. But additional means are sometimes necessary: these are, first, the application of ointments to the sore. The best

for the purpose are those which contain the milder preparations of mercury; but they require to be varied, and sometimes other applications succeed when these fail. Where there is great pain and tenderness, suppositories of opium and lard are occasionally more useful. Sometimes the application of lunar caustic, on the other hand, takes off the irritability of the ulcer.

When the ulcer will not yield to these simple means, either of the following trivial operations may be resorted to: the edges of the ulcer may be pared off by means of probe-pointed scissors bent upon the flat introduced on the finger into the anus; or the surface and edge of the ulcer may be divided by turning the edge of a straight probe-pointed bistoury, similarly introduced, against it.

One variety of the complaint that I have met with is the following. The irritation of the ulcer has caused a growth of the part of the surface on which it has been situated, giving rise to a tumour or kind of prolapsus. At each relief of the bowels the fold protrudes, on which the ulcer may be seen. In a case of this nature, I pared the edges of the ulcer, hoping it would heal. This practice failing, I included the whole protruded fold in a ligature; and the patient was well on the separation of the slough.

## SECTION V.—CONTRACTION OF THE ANUS.

CONTRACTION of the anus may be either spasmodic and temporary, the result of undue action of the sphincter muscle, or it may consist in permanent thickening and induration of the orifice of the intestine.

Spasmodic contraction of the anus is most commonly met with as a symptom of fissure of the rectum, and disappears when that complaint is cured; but sometimes it occurs as an independent disorder.

In the former case it forms an aggravation of the disorder; but it disappears on its cure. In the latter case it sometimes occurs as a kind of cramp, which comes on suddenly. The patient, who has gone to bed quite well, awakes in violent pain. The sphincter muscle is hard, and in strong action, so that a small bougie cannot be introduced without difficulty. In some cases these paroxysms recur daily, in others only two or three times a-year. In some, again, the attack of spasm comes on gradually, and after producing uneasiness for several days, gradually wears off; in others, it is sudden in its invasion, and sudden in leaving the patient.

The complaint generally depends upon a confined state of the bowels; and a brisk cathartic at night, with an aperient draught in the morning, will often

be sufficient to cure it. Or the patient finds it enough to use injections of warm water, upon which the spasmodic contraction wears off.

When the pain is very severe, an ounce of tepid water, with twenty or thirty drops of the liquor opii sedativus, should be injected into the bowel, and at the same time some aperient medicine taken. Sometimes the spasm is relieved by extending the circular sphincter muscle, and keeping its fibres on the stretch. The patient for this purpose may introduce a large mould candle into the anus.

Sometimes the sphincter acquires double or treble its natural size and thickness, without any alteration of the texture of the part. I have known this arise from the habit of eating peppered food. In such a case solid motions will not pass; and it is necessary daily to take some laxative, to prevent obstruction and derangement of the bowels. In one case of this kind, which was under my care, the patient underwent severe aggravation of his symptoms on the accidental supervention of fissure of the rectum. He became, however, as well as before, upon going abroad and commencing a course of Carlsbad waters.

There are other cases in which the anus becomes permanently contracted and indurated, constituting a kind of permanent stricture, and generally combining both permanent and spasmodic contraction. The motions are always passed with difficulty and

with pain, and all the common symptoms of stricture of the rectum are present.

In this more aggravated form, the complaint is slow to yield to the simple remedies which have been recommended; and the complicated sufferings which the patient endures, require speedier relief. This is obtained by dividing the sphincter on one side. The incision should be as before, transverse, neither forward nor backward, but *lateral*.

The wound should not be allowed to heal by adhesion, but by granulation. The effect of which is, that for several days the part is relaxed, and the strain and tension are removed; through which the morbid sensibility disappears, and with it the preternatural thickening.

A sort of tic douloureux occasionally is situated at the extremity of the bowel. In some cases the paroxysm of nervous pain occurs very seldom, and is of short duration, in which case no local treatment is requisite. When the pain has been constant, and the ordinary means of allaying neuralgia have failed, I have known it serviceable to perform the operation of dividing the sphincter.

#### SECTION VI.—OF STRICTURE OF THE RECTUM.

ALL the mucous passages in the body are liable to become constricted; the urethra, the gullet, the windpipe, the lachrymal duct, and the rest. They

are narrowed through inflammatory condensation and thickening of their texture. But in addition to structural and permanent narrowing, those mucous canals which are surrounded with muscular fibres, are liable to spasmodic stricture; that is to say, to temporary narrowing produced by spasm of the fibres which surround them: the spasm over, the canal is of the same dimensions as before. The spasm in all these cases depends upon an irritated state of the mucous lining of the canal, which brings on by consent spasm of the surrounding muscular fibres.

To consider spasmodic stricture first.

We may begin by inquiring what part of the lower intestine is the seat of spasmodic stricture? What I have seen of the complaint disposes me to think that no single point is more liable to this affection than another. The cases, however, to which I refer, have been anything but satisfactory. They have left me with the impression that both the rectum, and the sigmoid flexure of the colon, are liable to partial contractions of their muscular tunic, capable of obstructing the passage of the fæces, and of making resistance to the introduction of instruments. This irregular and spasmodic action generally goes with a vitiated state of the secretions; and is more frequently relieved by a regulated diet and aperient medicine, and the use of injections, than by the employment of instruments.



The symptoms which denote the existence of this complaint are, irregular action of the bowels, with a sense of difficulty in emptying the intestine, uneasiness of the lower part of the belly above the left groin before the bowels are relieved, weight and uneasiness about the sacrum, occasional discharge of mucus, the motions sometimes natural, commonly either relaxed or in fragments,—the existence of no other disease being ascertainable.

One of the best instances which I can give of the disorder is the following: it contains a very useful lesson as to its treatment. The patient is a physician, who is now through his judicious self-management restored to perfect health. The extract which I shall quote from a letter, in which at my request he favoured me with an outline of his case, that had several times been the subject of communication with me before, will convey to the reader an idea of the suffering which may attend this kind of disorder.

“In my life,” says the writer of this communication, “I never knew what it was to have a single action of the bowels without the aid of medicine, or to be free for many hours together from all the wretchedness of disorder and of remedies in conjunction, excepting for two short intervals of time, during one of which I trusted simply to the use of injections of warm water, and during the other, when I took the white mustard-seed, and that

with so singular an effect, that for a while I thought I had quite got rid of my complaint. With the exception of these two intervals, I have never been able till lately to say there is in life that which is worth living for, or in other and more proper words, I did not know what it was to wish to live. To say nothing of the medical discipline which I have undergone again and again, I have been examined and treated for stricture of the rectum and of the sigmoid flexure of the colon for years, and for years never passed anything from my bowels larger than a horse-bean, if solid, or of the little finger, if of a softer consistence. Oftentimes have I been quite incapacitated for exertion, and never able to enter upon my professional duties with anything like alacrity or cheerfulness. It is now nearly two years ago since I came to the resolution of abandoning all remedial measures: to leave off at once physic, injections, and the bougie: to take nothing in the shape of food that could by possibility irritate the stomach or bowels, and to leave them to act of and for themselves, when they could no longer retain their contents. I had, as you may suppose, difficulties in bringing about so entire a change. At first I suffered much inconvenience from a sense of fulness in the bowels and in the head. But this I contrived to obviate by the very occasional use of an injection of warm water, determining with myself to overcome the disposition to

contraction by making the contents of the lower bowels the means of dilating them. By a steady perseverance in this course of discipline, I have perfectly recovered; know nothing now of that distress of feeling which for at least twenty years made life burdensome to me; I have seldom or ever occasion to have recourse to medicine, and then only as a man in perfect health would do. I should tell you, that at one time such was the state of the stricture in the rectum, that the largest-sized urethra-bougie alone would pass, and that at another the contraction was so far in the intestine that a bougie of three feet in length was considered necessary to reach it."

It may be instructive to place in contrast with the preceding case another, which I shall again give in the words of my patient, who describes, in the third person, his own case. In this instance the idea of abandoning medicine appears likewise to have been resorted to, but not with an equally good effect; for it led to a very serious attack of obstruction of the bowels. In this instance, again, the use of the bougie has proved salutary. There exists, however, in this patient a point at which the bowel is so constantly found narrowed, that it is more than probable that he labours under permanent stricture of the rectum, in addition to that tendency to general irregular contraction of the adjacent bowel, which constitutes spasmodic stricture.

“A gentleman, now in his fifty-eighth year, who from early youth had been subject to a very irregular action of the bowels, amounting frequently to an alternation of costiveness and dysentery, was, about fifteen years ago, strongly urged by a physician to abstain from medicine, and to let the bowels alone. This experiment was tried with great resolution. In spite of much suffering and increasing feverishness, the patient took no medicine for more than a week. Inflammation however ensued. Intolerable pain in the abdomen, and simultaneous vomiting and purging, reduced him to an alarming state in the middle of the night. Skilful medical assistance was fortunately obtained without loss of time, and the acute symptoms were subdued. The patient's general health however grew worse. His bowels were never at rest. Acrid mucus was incessantly formed, and frequently passed, leaving the sufferer in a state of great weakness. Blood was sometimes observed in the mucus. Scarcely anything was passed without great effort and pain. Spasmodic contractions of the rectum were constantly attendant on every attempt to ease it. Great emaciation and prostration of muscular power took place, as also restlessness at night, amounting sometimes to the most painful startings from sleep. After a few ineffectual attempts to perform a cure, treating the case as one of liver derangement, he confined himself to

the use of the common purgatives for the paroxysms of the complaint, and of a small quantity of rhubarb and ginger before ~~himself~~, for the daily symptoms. ~~though~~ though very slowly, yet he improved from year to year; but owing to the unsettled state of the bowels, he could hardly venture out of his house. By the advice of a friend, he tried, about two years ago, the daily use of lavements by means of Read's syringe. He has used nothing but tepid water. At first the lavement produced great nervous weakness; but this symptom disappeared in a short time. At present he enjoys a certain degree of comfort and ease, which entirely depends on the use of the lavement early in the morning. From a local examination, it has lately been ascertained, that the rectum is contracted to about half an inch diameter, at a distance of about five inches from its termination. The daily passing of a wax bougie, softened by heat, is attended with little or no pain. The distension of the contracted part by this mechanical means relieves the spasmodic contractions, which the patient frequently feels a little above the sigmoid flexure."

I shall conclude my remarks upon the subject of spasmodic stricture of the rectum with the following case, which was communicated to me by Mr. Crosse of Norwich. I will give it in his own words. It is unnecessary to state, that that eminent provincial surgeon was not the practitioner

through whose mismanagement the fatal termination of the case was produced.

The case is one in which the coats of the intestine were remarkably thin, which gave rise to an imperfectness in the action of the bowels (as explained at page 95), which was erroneously considered to proceed from stricture. In every point of view this serious case is full of interest.

“A young woman of delicate frame was supposed to have stricture of the rectum, which led her medical attendant to employ in no very gentle manner a firm bougie. After much difficulty the instrument was made to pass; but the patient in a few hours became very ill, vomited, and was chilly, and in about forty-eight hours died. It was found that the bougie had perforated the coats of the bowel at the sigmoid flexure, about seven inches from the anus, and had entered the peritoneal cavity. The preparation, which is in my collection, shows the rectum to be capacious for an inch or two next the anus; but all the rest of the bowel preserved, being a length of eight or nine inches, is very contracted, so that it would only admit a small instrument half an inch in diameter; and at the same time its coats are very delicate and attenuated, readily allowing the bougie, in the hands of a boisterous surgeon, to perforate them. There is a great abundance of adipose substance and of fatty appendages about the sigmoid flexure

of the colon. The bowel presents no thickening or partial contraction, but a smallness of calibre generally, with remarkable delicacy and thinness of the coats—accounting for the presence of symptoms, during the life of the patient, which might have led to the supposition that stricture existed.”

Permanent stricture of the rectum consists in a partial thickening of the submucous coat of the bowel and of the adjacent cellular texture; through which means a smooth ring is formed, generally from a third to half an inch in depth, which projects into and narrows the channel. Sometimes the thickening does not include the whole circle of the intestine, but a segment only. It is presumable that this thickening results from chronic inflammation.

The ordinary seat of stricture of the rectum is from two and a half to four inches from the orifice of the gut. But sometimes it occurs at a greater distance, at six to seven inches for example; and a contraction of the same nature is occasionally met with in different parts of the colon.

The symptoms of stricture of the rectum are the common and necessary consequences of the excretory canal being narrowed at one part.

The fæces are passed in small and narrow and flattened portions. The quantity voided at a time is inconsiderable, from the effort required to pass it through the stricture. The bowel being thus

insufficiently relieved, the effort has to be repeated frequently during the day; and it is only after many efforts that all its contents are passed. The narrowed portion of the canal is extremely sensible; which is owing partly to the thickening which forms it being originally produced by an inflammation, partly to the irritation of the mucous lining of the stricture, which is occasioned by the more forcible pressure of the fæces against it. When the bowels act, in addition to a sense of obstruction in the part, pain is experienced and tenesmus, and mucus is voided; and at other times a sense of weight and tightness, with general uneasiness, is felt. The habitual confinement of the bowels alternates with periods of looseness and purging. Uterine irritation, irritation of the bladder and urethra, numbness and pain down the thighs and legs, are occasionally concomitant symptoms.

But if these symptoms are common to every affection by which the channel of the rectum is narrowed, in what manner is a case in which they occur proved to be a stricture?

It has been mentioned, that the seat of stricture, in nineteen cases out of twenty, is near the orifice of the bowel; that is to say, within the distance which admits of examination by the finger. In common cases, therefore, no difficulty exists in identifying the disease: the finger may be passed



into the smooth and firm narrow ring which forms the stricture; and direct proof may be thus obtained that this and no other disease is present.

The treatment of ordinary cases of stricture of the rectum is no less simple in theory than satisfactory in its practical results. In stricture of the rectum, as in stricture of the urethra, if a bougie of a size calculated easily to fill the stricture be passed into the narrowed part daily, or every second, third, or fourth day, according to the irritability of the patient, and retained for from ten minutes to a quarter of an hour at each introduction, the pressure of the bougie causes the absorption of the lymph, on which the inflammatory thickening of the walls of the canal depends; the patient is thus enabled gradually to introduce larger and larger instruments, and the channel is at length restored to its original calibre.

A diet carefully regulated, the use of mild aperient medicines, of injections of tepid water, and of anodyne suppositories, are important accessories in the treatment of stricture of the rectum.

In the use of instruments for contraction of the rectum, the point to be constantly the most thought of and studied is gentleness. If any considerable degree of force is used, the bowel will be torn: if pressure is made against the stricture with too large an instrument, the adjacent and sound part of the bowel is certain to give way sooner than the stric-

ture. The canal of the urethra is often torn in this manner by the pressure of an instrument too large to enter the strictured part.

In stricture again of either canal, if by dexterous management an instrument too large is successfully forced through the contracted part, it may yet produce most serious consequences. If such an instrument is forcibly passed into a stricture of the urethra, the patient in a few hours after the operation is seized with shivering, and symptomatic fever follows, which will last several days. If similar violence is done to the rectum, the surgeon being anxious rapidly to complete the dilatation of the stricture, within twenty-four hours shivering supervenes, as in the former instance: but here it has a more serious character; it is probably the precursor of peritonitis, to which the patient may in a few days fall a sacrifice. There is a singular consent between the pelvic mucous passages and the peritoneum. If a stricture either of the vagina or of the rectum be roughly dealt with, peritonitis is liable to ensue; notwithstanding that the violence is done at a part of either canal, which is not covered with peritoneum.

Such violence is not merely mischievous, but it is utterly unjustifiable on any ground: it can be used only through a mistake of the principle of dilatation; the object of which is not mechanically to stretch the narrowed canal, but to excite

by moderate pressure the absorption of that which thickens and contracts it.

Division of a stricture of the rectum has been recommended. I have done this operation with advantage upon some occasions, but my general experience is not in favour of it. It is liable to produce troublesome bleeding; and what is obtained by the division must be kept open by the use of bougies, by which alone in nearly the same time dilatation might have been effected.

It is not always easy to distinguish stricture of the rectum from incipient carcinoma.

A lady about forty-five years of age had suffered severely from piles, which were removed five years ago by the ligature. It was stated to her that they did not grow from the fore-part of the rectum. Some months after this the lady began to feel a tightness and sense of obstruction in the rectum. These sensations gradually became more distressing: much effort and straining were necessary to pass the fæces, which were narrow, flattened, and in fragments. After two years of suffering, this patient applied to me for advice. There was an induration, which began two inches within the rectum, and occupied two-thirds of the circumference of the gut. The central and broadest part of the induration was towards the vagina; at this part it was two-thirds of an inch in depth. The part was sore or tender on pressure. I recommended that the bowels should be

relieved every morning by means of a lavement of tepid water, and that a soft wax bougie should be introduced into the narrowed part every second day. Under this treatment, combined with the occasional use of aperient medicine, a decided amendment took place: the narrowed part yielded to a certain extent, and there was a proportionate alleviation of all the symptoms. But in a short period the patient became worse again; the introduction of the bougie now gave more pain; it was therefore discontinued. The passage was indeed certainly freer, but the induration towards the vagina was not lessened. Under these circumstances I wished Mr. Copeland to see the case in consultation. The impression which the examination left us with was, that the disorder would prove carcinoma.

The plan which the patient followed was slightly modified. The use of the bougie for a time was not resumed. The increased sensibility of the part went away. But it was not long before the patient again complained of the contraction returning; upon which the bougie was again used, but for a shorter period than on the first occasion. Since then at intervals the patient has occasionally had recourse to this remedy again. She is now materially better: the narrowing has lost its doubtful character: the induration is less in extent, and the projecting band has little more than the character

of a thickened fold of mucous membrane. Some discharge of matter *per vaginam* took place, and continued for several weeks, about a year ago. I am disposed to think that it proceeded from the induration, which may have suppurated, and the abscess have broken into the vagina at that time. This account was written four years ago. The lady has suffered no further inconvenience from the complaint. (1837.) Now (1840) this lady is in perfect health.

Stricture of the rectum is occasionally met with in combination with a dilated sac of intestine below it. The stricture with the part above is then borne down as a sort of inward prolapsus into the lower and dilated portion of the rectum. This disease is liable to be overlooked, but when discovered yields, like other stricture, to dilatation by means of the common wax bougie.

Stricture of the rectum is a disease of extremely rare occurrence. It is, however, a common trick of quacks to represent the complaint as frequent, and to inflict a course of treatment by instruments upon patients who have no stricture to remove. Two or three years ago, I saw, in consultation with Mr. Copeland, a lady of rank, who for several years preceding had been under systematic and almost daily treatment for stricture of the rectum; the question proposed for our consideration was, whether this plan required to be continued; it turned

out, on examination, that there was no ground for supposing that stricture had existed, and that, in all human probability, the only complaint ever present had been functional irregularity of the action of the bowels.

SECTION VII.—OF SUBSTANCES IN THE RECTUM PARTIALLY OR COMPLETELY OBSTRUCTING THE PASSAGE.

As the substances which may be impacted in the rectum are of several kinds, and are derived from various sources, so the symptoms which they produce are different, and each kind requires to be considered separately. They are, first, the natural alvine contents unduly accumulated; secondly, hard intestinal concretions; thirdly, hard and indigestible substances that have been swallowed; fourthly, hard substances introduced into the rectum through the anus by design or accident.

I. Accumulations of the natural contents of the rectum in a mass too large to admit of being expelled by the efforts of the bowel, sometimes, but very rarely, take place in adults; they occur rather less unfrequently in children, but are not very uncommon in aged persons.

This ailment is liable to result in children from neglect of attention to the relief of the bowels. If the contents of the lower intestines are not expelled, some absorption of their liquid parts takes

place, the mass is thus lessened, and is rendered more solid and less stimulating, at the same time that, by an inverted action of the intestine, it is forced retrogressively along the canal.

In this manner collections of fæces, drier and firmer than natural, often disposed in little round hard fragments, or *scibalæ*, are liable to accumulate in any part of the great intestine from the rectum upwards, and to form a source of obstruction.

The symptoms which are caused by such accumulations vary in different cases. Sometimes many days pass,—a fortnight or three weeks,—without the ailment being discovered. For this period the bowels have not acted, but the child's appearance has not altered so as to attract observation. Gradually, however, he loses his appetite and colour; has headach, is sick, and vomits, the belly is full, tumid, uneasy. Having learned that these symptoms have been preceded, and are attended by entire costiveness, it is best to begin with unloading the intestine by means of injections. For this purpose, warm water with soap dissolved in it, or barley water, a quart containing a tablespoonful of salt, will commonly answer.

When the lower part of the bowel has been by these means cleared, aperient medicines should be given by the mouth, to unload the upper part of the great intestine, and clear the whole passage. Further means than injections of emptying the

lower bowel are commonly unnecessary; sometimes, however, the accumulation is in so great a mass, and so solid, as to resist the entrance of liquid, and it is necessary to use a scoop in addition to the syringe, in order to loosen and dislodge the obstructing masses.

Mechanical means are required for two different cases under this head. On the one hand, the accumulation often cannot be moved from above, till the mass is diminished by getting rid of the lower portion. On the other hand, great collections of fragments, sufficient entirely to disorder the digestive apparatus, will accumulate in different parts of the great intestine, and yet allow daily, and even frequently, some passage to liquid secretion.

A publican, aged thirty-five, had been bed-ridden for six months. He lay supported in bed in an inclined position: if he lay down he was oppressed and uneasy: if he stood up, he vomited. He was corpulent, and ate voraciously. The bowels would act with medicine, but not without. This resulted from an accumulation in the great intestine, higher than could be reached by examination. The most powerful drastics were given, but without dislodging it. It was then advised to inject very large quantities of tepid water. After nine pints had been slowly and cautiously thrown into the intestine, an immense discharge of scibalæ took place. The injection, but in less quantity, was repeated



daily for a week, and for that period more scibalæ continued to be discharged. After the accumulation had been thus got rid of, an alterative course of medicine and regulated diet perfectly restored the health of the patient.

In young persons it has happened that an accumulation of fæces in the descending part of the colon, producing fulness in that region, and pressure upon the lumbar nerves, has given rise to symptoms exactly resembling those of caries of the lumbar vertebræ, which have disappeared on the exhibition of drastic purgatives; and the patients have in a few days been restored to health out of a state of progressive and alarming indisposition, which appeared utterly hopeless.

In elderly persons the accumulation is often not in scibalous fragments, but the intestine is filled with a mass of cohesive substance, like putty. This is probably less stimulating than ordinary fæces, and the intestine again, from age, is torpid. Elderly persons often have repeated attacks of the nature I am now describing, and after having got over two or three, sink eventually under a return. However, by constant attention to the action of the bowels, the disorder may be often prevented; and when it has decidedly recurred, it often gives way to the prompt use of drastic purgatives. It is wonderful how little inconvenience is caused to the patient by large accumulations of this description.

I attended, with Mr. Drew, of Gower-street, an

elderly lady, who was labouring under this complaint: the belly was large and full, but not tense or tender upon pressure. In conjunction with strong purgative medicines and purgative injections, the hot bath, and venesection to the extent the pulse and her age justified, had been tried, but ineffectually. The complaint gave way, however, under the use of oil of croton, of which she took three drops, a drop every two hours: the bowels then began to act, an immense quantity of fæcal matter was discharged, and she was restored to health. Two years afterwards this lady died of a return of the same complaint.

The following case exemplifies other features of such an attack, and the necessity of using mechanical means to unload the intestine:—

I was requested by Mr. Reid of Bloomsbury-square to see an elderly lady. She had generally enjoyed excellent health, and had been accustomed to take a good deal of exercise. But ten weeks previously she had been attacked with rheumatism, which had confined her to her bed. The bowels during this ailment gradually became inactive, and at length were never moved without medicine. The medicine which agreed best was castor oil, but even this failed to give sufficient relief. The quantity of fæces passed was small, and its expulsion caused some pain of a spasmodic character, which remained for hours afterwards. These pains

occasionally came on at other times, and were accompanied with throbbing, and a sense of stoppage in the rectum. To relieve her sufferings, it was necessary to give opium, which increased the constipation. Upon examining the rectum, Mr. Reid found that it contained a quantity of clayey fæces, which he removed with a scoop. Water was then injected, and more fæces were returned with it. This operation gave the patient temporary relief; but in the course of a few days the pain gradually returned, and became more intense than ever, coming on in paroxysms, especially when the bowels were moved, a thin fluid, however, being all that passed. It was at this period that I saw the patient in consultation with Mr. Reid. The accumulation of fæces in the rectum had returned in a surprising quantity; but there was great tenderness and soreness about the sphincter, which indisposed the patient to allow the trial of any further mechanical relief. However, with care and gentleness, the orifice was gradually dilated, an immense volume of fæces was brought away, and the rectum, which was unusually capacious, was at length emptied. The pain then subsided: a dose of castor oil which was administered, brought away on the following day more fæcal matter that had been accumulated above, and the relief was complete.

II. By intestinal concretions I mean such as are

immediately compounded of substances which have been swallowed as food or medicine. Of these there are, doubtless, many that have not yet been submitted to chemical analysis. Dr. Marcet and Dr. Wollaston examined some, which proved to be concretions of caseous matter. Others have been composed of oat-cake. And magnesia, where taken in large doses and repeatedly, has been consolidated in the bowels into hard masses of formidable magnitude.

Such concretions being of slow increase, cause a gradual invasion of abdominal uneasiness, pain, nausea, vomiting,—finally, local irritation, when they reach the lower bowel. It is then only that their existence is positively ascertainable; when their removal is to be practised by the mechanical means which their shape and resistance render necessary. The following instance, given by a late eminent American surgeon, conveys a just idea of this form of disorder:—

A female, aged thirty-five, had been for some years subject to constipation and repeated attacks of colic; the former had increased, attended with sickness of the stomach, and the latter had become more frequent, from which she only experienced relief when her bowels were moved, a task not accomplished without great difficulty and painful efforts, joined to the use of much purgative medicine. Dr. Bushe was called to visit her in one of

these paroxysms. "I found her," he says, "sallow, emaciated, dejected. From the severe bearing-down pains, together with the sense of weight and fulness in the sacral region which she complained of, I was led to make an examination of the rectum, when I found the mucous membrane slightly protruding, the sphincter excessively irritable, and a large concretion distending the pouch of the rectum. I now apprised her of the nature of the case, and the absolute necessity of removing the foreign body, to which she willingly consented. For this purpose, the patient being placed in a suitable position, I introduced a strong and long pair of lithotomy forceps, with which cautiously laying hold of the concretion, I slowly and steadily extracted it, with no more injury than slight laceration of the mucous membrane, although, on measurement, it proved to be six inches and three-quarters in circumference, and two inches and a-half in length. The bowels were then freely washed out with injections: leeches and fomentations were applied to the anus, the recumbent position was enjoined, and a speedy recovery ensued."

III. Hard and indigestible substances that have been swallowed, and have made their way down to the rectum, have been most commonly portions of bone. They generally pass through the bowels without creating disturbance, and are first suspected through the pain and irritation which they produce

in the rectum. The ordinary period that they take to traverse the length of the intestinal canal is about eight days. Their extraction is a mechanical operation, in which the surgeon must be guided by their shape, magnitude, consistence, position.

IV. Foreign substances directly introduced into the rectum, either require the same management as the cases last described, or in their introduction, produce wounds, which combine with the common features of local injuries those of laceration of the intestine.

#### SECTION VIII.—OF DEFECTS OF THE RECTUM EXISTING AT BIRTH.

THE lower part of the intestine is liable to malformations of various kinds, which are discovered at, or shortly after birth. Many of these admit of being completely repaired: others are irremediable.

These imperfections are referable to three heads,—obstructions of the orifice, or in the intestine above it;—deficiency of part, or of the whole, of the rectum;—improper terminations of the rectum in other parts, or of other parts in the rectum.

I. Congenital obstructions of the bowel are either partial or complete. Partial obstruction is, when the bowel is perfect everywhere but at its opening, and when that, instead of being natural, is a narrow

canal, not much larger than will admit a probe, and incapable of giving passage to other than liquid fæces. This defect of structure is liable to escape observation for many weeks, owing to the fluidness of the alvine evacuations at first; but sooner or later,—that is to say, when the fæces become more solid,—it causes obstruction, uneasiness and distension of the belly, loss of appetite, vomitings, emaciation. As soon as the defect is discovered, it may be rectified. The method to be employed for this purpose depends upon the size of the opening. When that is not extremely small, it will often allow of dilatation by means of a sponge-tent, or conical bougie of waxed cloth. When the aperture is too closely contracted for this, it requires to be enlarged by the division of its border. The operation is unattended with danger, and is, at the same time, much less fretting and troublesome to the infant, than the more tedious process of dilating. Add to which, that the child's health may be suffering from obstruction, and prompt as well as complete relief may be required.

When a proper aperture has been made, it is generally unnecessary to use artificial means to prevent its closing again: the passage of the contents of the bowels will probably be sufficient. The direction in which the incision is to be made, depends upon the place of the existing aperture, in relation to the bowel and to the middle line of the

body. The division of parts should be very trifling in extent, as it is easily enlarged, if that should prove necessary; and the less its extent, the more perfect afterwards will be the voluntary power of closing the intestine.

The following case may serve to exemplify many of the features of the congenital imperfection of the bowel under consideration,—its first oversight, the derangement of health produced by it, the facility and completeness with which it may be remedied.

I was requested, in the spring of the present year (1837), to see an infant six weeks old, which had thriven for the first fortnight after birth, and then had pined and fallen away. A few days before I saw the infant, some kind of farinaceous food had been tried, after which it was noticed that the relief of the bowels was imperfect, and attended with straining and pain. On examination, it was found that the rectum opened by a very small aperture, barely large enough to admit a probe; the aperture was upon the right side, and close upon the raphe of the perinæum. On the other side of that line, and about a quarter of an inch from the aperture described, there was a shallow *cul de sac*, like another attempt to open the intestine.

Having examined the part with a probe, I introduced into the canal a button-pointed bistoury, with which I divided the integuments backwards, for



the extent of half an inch. A quantity of semi-fluid fæces immediately came away; the aperture allowed the extremity of the little finger to be introduced into the sac of the intestine, which seemed to terminate about a third of an inch from the surface. The child from this time improved rapidly, and became strong and healthy. No dressing or plug was introduced; but on the sixth day, when the opening seemed disposed to contract, I again enlarged it to a very slight extent with the bistoury. Since then (now six months) nothing more has been necessary, the aperture remains perfectly free, yet appears to have a sufficient power of closing. (1838.) About three months after this when the child was weaned, the fæces assuming a more solid character passed with such difficulty that considerable accumulation took place, and the belly became enlarged, some fæces, however, passing daily. The child now fell away, and became fretful and ailing; on examination the aperture was found not to be large enough to give passage to the solider fæces. It was therefore enlarged and a short ivory bougie introduced daily for several weeks. The child became again perfectly well, and continues so: the sphincter is apparently not deficient.

Complete obstruction of the bowel may take place either at the anus, or higher up in the intestine.

The former case is known by the absence of an

external aperture, joined to a fulness and prominence where the bowel should open. The imperfection is easily remedied, by puncturing the full and prominent part, and enlarging the puncture into a crucial incision, a quarter of an inch each way. The aperture should be kept open by a pledget of oiled lint, or bougie.

When the external orifice is perfect, sometimes one, sometimes two adhesions of the bowel are met with within: these are suspected when, with a natural opening, there is no discharge of the contents of the bowel. Their existence may be ascertained by an examination of the intestine. These adhesions are said to be sometimes very slight, and to admit of being broken through by pressure with the finger. When of a firmer consistence, constituting strong membranous septa, they require to be perforated with a trocar, and the opening cautiously dilated, or otherwise enlarged.

II. Deficiency of the rectum, partial or complete, is a more serious form of obstruction; but the first kind, in many cases, is still remediable.

When the lower part of the intestine is deficient, its place is occupied by fat and muscular fibre: often, too, the pelvis is unusually contracted. That there exists some considerable deficiency, the surgeon judges from the want of an external perforation, and of that fulness and prominence to the touch which goes with simple imperforation.

Nevertheless, the length of intestine absent is necessarily uncertain: it may be greater, it may be less. Practically, there are two forms only of this case—one, in which the surgeon, under no circumstance of the infant straining and forcing, distinguishes any fulness of the perinæum: the other, in which such an impulse or greater fulness is more or less sensible. The first case is extremely unpromising: however, in both the same method must be adopted; the surgeon must try and find the extremity of the bowel.

For this purpose he is not to use a trocar, but having made a crucial incision with a lancet or scalpel in the perinæum, where the bowel ought to have opened, he is cautiously to seek in the direction of the sacrum for a prominent sac. This can only be safely done by a process of slow and cautious dissection. When at length the *cul de sac*, in which the intestine has terminated, is reached, it has to be punctured, its contents let out, and a short ivory or metal bougie left in the wound to preserve it open, to be from time to time removed to allow the contents of the intestine to escape.

How far in such an operation a surgeon should pursue the search after the terminus of the bowel, must be determined by his anatomical knowledge. He should not, however, too hastily give up the attempt, which if it fails one day, may be afterwards renewed, for two reasons. First, if the in-

testine cannot be found from below, there is, in my opinion, no other justifiable expedient left; and the little patient, after surviving two, three, or four months, with no evacuations, with an enlarging belly, vomitings and gradual emaciation, must sink. The second, that the relief obtained by thus opening the bowel not only preserves life, but sometimes preserves with it the possession of bodily comfort, and as much capability of physical exertion and active habits as those naturally formed enjoy. To what extent this may be the case is exemplified in the following instance.

A young gentleman now in good health, and twenty years of age, asked my advice during the past spring for his further management of such a case in his own person. He had been born with part of the rectum deficient, and an artificial passage had been made. It had been kept open by instruments, the form of which was the following; I give the size from that which he at present uses. It is a cylinder of metal, two inches and a half in length, and something more than half an inch in diameter, one end is rounded, the other fixed to the convex surface of a short thick half cylinder. This part, when the instrument is introduced, rests and is fixed by a bandage against the hollow between the nates. The rounded end of the instrument passes into the cavity of the bowel. Or on examining the artificial passage, it is found to be upwards

of two inches in length, opening at that distance into the extremity of the rectum, which is felt to have a soft, smooth, circular lip of mucous membrane; the character of this circular orifice is quite distinct from the more rigid and callous wall of the canal which leads to it. Wearing the instrument which I have described, this young gentleman has joined in all the active sports of his companions, is a bold rider, and fond of hunting, from which he does not suffer in the least. The circumstance which led him to consult me was, that having used a new instrument, which had been ill cast, and was notched and rough at one point, it had produced considerable soreness in the part of the canal against which it had pressed. The absence of the ordinary sphincter-muscle had been of no inconvenience. The difficulty he had experienced had been in expelling, not in retaining the contents of the intestine. Even if he took aperient medicine, this difficulty was experienced; which, however, was obviated by the use of injections of warm water. (1837.)

III. The third head under the subject of imperfections is the wrong termination of the rectum in other parts, or of other parts in the rectum.

The latter case is essentially irremediable; the former is not necessarily more than a complicated case of deficiency of the rectum. Suppose, for instance, the intestine to want an inch in length, and

to open into the vagina or the bladder at the same part;—there is here the deficiency explained, with its appropriate remedy, under the last head, and in addition, an opening into another cavity. There are then two malformations instead of one. But the additional one renders the first more tractable. If, for instance, the intestine open into the vagina, it is much more easy to make an artificial canal to the extremity of the intestine than without this means of directing the incision. The first step to be taken is to cut such a new canal, and to gain for it proper dimensions and permanency by metal cylinders kept constantly in it. When this has been accomplished, it remains to close the original false aperture by sutures.

#### SECTION IX.—OF LACERATION OF THE RECTUM.

MECHANICAL laceration of the rectum may involve either its inner lining only, or all its coats. The mucous membrane alone may be lacerated, or the bowel may be torn through.

The first case may arise from hard fæcal matter passing, or from the forcible introduction of a foreign body. If the patient is kept at rest a few days, and upon a restricted diet, and the bowel washed out by emollient clysters, the lesion is quickly repaired, and the part heals. If neglected, such a lesion becomes the troublesome complaint

which has been already described as a fissure or superficial ulcer of the rectum.

A rent or laceration of the entire thickness of the intestine may take place either at the orifice, not involving a complete division of the sphincter, or it may include the whole breadth of the sphincter, or, leaving this muscle untouched, it may perforate the rectum above. These injuries are of more frequent occurrence in women than in men, and are in the former of much more consequence. In the latter, they are simple wounds, which heal more or less readily, being attended in different cases with more or less local inflammation, and the like. In women, these injuries acquire a serious character from threatening to establish permanent communication between the intestine and the vagina.

The two first lesions are liable to take place in labour, when the descent of the child is more rapid than the yielding of the external parts.

If the laceration, though complete, is superficial only, and does not include all the fibres of the sphincter, the lesion is of little consequence; the state of the patient ensures the observance of all the conditions necessary for its reparation. These are rest in the recumbent posture, and frequent ablution with warm water. The superficial rent quickly re-unites by granulation.

When the laceration extends through the sphincter-muscle, so as entirely to divide it, a serious evil

naturally follows. The tone of the muscle draws asunder the edges of the rent, which thus are prevented re-uniting: in this state they cicatrize separately, leaving an opening into the vagina of an inch to an inch and a half in depth; the power of retaining the contents of the bowel is of course lost. Fortunately there are modes both of preventing this ill consequence of the accident, and of repairing it when established.

When the laceration is quite recent, that is to say when two or three days only have elapsed, the following practice is to be adopted. The sphincter-muscle is to be divided in one or two places, that is to say, on one or both sides of the anus. By this means, the lateral strain or traction upon the rent of communication between the intestine and vagina is done away with, and there is no impediment to its re-union by granulation. But it is necessary to keep the edges of the rent in a wholesome state by frequent bathing with tepid water, and to prevent their cicatrizing separately by daily touching the edges with lunar caustic, and to prevent the lateral sections of the sphincter from closing too soon by means of pledgets of lint introduced into, and retained in them constantly. A patient on whom I performed this operation eight days after laceration of the sphincter in labour, had entirely recovered in five weeks; the original rent first healing, the lateral sections being then allowed to close. By



the same time the retaining power of the sphincter was perfect. This patient has subsequently borne two children, without again suffering rupture of the sphincter. This practice was originally used by Mr. Copeland in cases of recent laceration of the sphincter. It had previously been used by Dieffenbach as a part of the more elaborate operation requisite when the communication is of long standing. In recent cases, however, Dieffenbach himself does not employ it, but contents himself with sutures alone, which, in the relaxed state of the parts following labour, are generally successful.

When the edges of a laceration through the sphincter have cicatrized without uniting, the means of remedying the defect consist in paring off the edges, and keeping the new surfaces in apposition by means of sutures. It is to render their adhesion surer that Dieffenbach introduced the lateral section of the sphincter into the present operation. The necessity of using it here is, however, very doubtful, and many cases have done well in which it has not been employed. The secret of the success of the operation is the management of the patient before and after it. For several days consecutively aperient medicine should have been given, so as to leave an artificial costiveness for the period necessary for the adhesion to be completed. Or without this condition is rigorously maintained, the operation, however skilfully performed, will probably fail.

The mechanical elements of the operation are, first, providing as large an extent of surface of adhesion as possible by making the sections oblique, and extending them through part of the skin immediately adjacent; secondly, the use of two sets of sutures, one for the mucous membrane of the rectum, the other for the mucous membrane of the vagina: the former are of course to be first introduced and tied, the latter afterwards. The former may be allowed to separate of themselves into the rectum; the latter may be easily removed on the fourth, fifth, or sixth day, according to the degree of adhesion, or otherwise. Thirdly, keeping the bowels undisturbed for five, six, or eight days after the operation. This may be accomplished by previous purging, and by subsequent spare diet; and if fæces still accumulate, by breaking the accumulation down and washing it out, through means of an elastic catheter introduced through the sphincter.

One source of failure against which the surgeon cannot entirely guard is the supervention of inflammation. But some security on this score may be obtained by delaying the operation till the patient's health is in the best order, either through diet and medicine, or temporary change of air, and the like. Likewise by noting the present influences of the weather as they are observable in the wards of hospitals; for it is certain there are some seasons in which the state of the atmosphere strongly disposes

to erysipelas. These times are commonly the setting-in of damp weather. It is desirable, for the same purpose, that the patient should not reside at the time in the middle of a large town, but live in purer air in the suburb.

These precautions are less essential in other operations in which the result is not influenced by the supervention of slight inflammation, which here is fatal to success. Nevertheless, there is still a hope even where this unlucky consequence has followed; the operation which has failed once may prove successful when repeated.

Laceration of all the coats of the intestine above the sphincter has sometimes been produced by the passage of hardened fæces. Dr. Langmore told me an instance (the patient afterwards was under our care for cancer of the breast,) in which such a communication had been so produced. A lady, aged 40, naturally of a constipated habit, and at the time being on a journey, on striving to relieve the bowels, which had not acted for many hours, felt something give way, and the following morning some fæces passed *per vaginam*. On examination, a transverse rent was found within the sphincter, sufficiently large to admit the end of the finger. The only treatment adopted in this case consisted in frequently cleansing the part by injections of warm water, and regulating the state of the bowels by proper medicines. In five weeks, the fæces

had ceased to pass through the communication, which had closed permanently.

When such an aperture loses its disposition to contract, its closure may be promoted by occasionally touching the edge with nitrate of silver. If these means fail, the suture may be employed, the precautions already explained being observed.

The most severe case of this kind of laceration which I have witnessed occurred in an elderly lady, who had been seriously injured some weeks before by the pipe of a syringe improperly used. The rectum had been torn about two inches from the orifice, and had partly sloughed into the vagina; at the same time, there was a large foul cavity higher up between the uterus and the intestine, from which a discharge of matter, and blood, as well as the contents of the bowel, flowed *per vaginam*. This lady's health appeared to be giving way under the spread of this foul and confined abscess, for the relief of which it was necessary to divide the sphincter, and to lay the lower part of the bowel open into the vagina, upon which she recovered.

#### SECTION X.—ABSCESS AND FISTULA.

ABSCESSSES frequently form in the neighbourhood of the rectum. Some may be considered as accidental, and are out of the scope of my present

remarks. These are, first, abscesses from foreign bodies, as a fish-bone, an apple-core, or the like, that have become impacted in the gut immediately above the sphincter. When the presence of such a substance has been ascertained by examination, it has of course to be extracted;—when its removal is followed by the immediate healing of the abscess, or the latter takes one of the characters presently to be described. A second form of accidental abscess at this part is, suppuration taking place in piles; this occurrence has been already noticed as a favourable event. A third kind is the pointing of lumbar abscess by the side of the anus; this case, which it would be out of place to consider here, requires, of course, to be treated on those common principles which regulate surgical practice in cases of chronic scrofulous abscess.

Independently of these accidental cases there is a great disposition towards the formation of abscesses about the anus, attributable to local causes already explained; namely, the dependent situation of the part, and the pressure of the column of venous blood, to which may be added the want of support to the vessels of the part resulting from the laxness of the surrounding tissue. And, as it is certain that many abscesses are often critical; that a feverish condition of the habit, or a gross and overloaded state of the body, or an impoverished state of the blood, will often lead to the formation of matter in

one part or other; so the circumstances above pointed out render the rectum particularly liable to be their seat.

Abscesses about the rectum are of two sorts; one of which may be termed diffused; the other circumscribed. The former, deep-seated at their commencement, gradually work their way outward, towards the perineum, and end in completely isolating the rectum behind and laterally, which lies in a great bag of matter. The latter are small gatherings from the size of a pea to that of a chestnut, on one or other aspect of the bowel, and close to the anus. The former are often attended with great constitutional disturbance, which disappears as soon as they are opened; the latter unattended by serious symptoms at their origin, end by giving rise to a very troublesome local disease, which will be presently considered.

The formation of deep-seated abscess upon the intestine is usually preceded by shivering and fever, furred tongue, pain in the back and head, sometimes delirium. It often happens that there is very trifling uneasiness about the rectum, the only local symptom being irritation of the bladder; the real nature of the disease is thus liable to escape observation for several days, till being suspected, it is at length ascertained by examining the perineum and rectum.

Part of the treatment of this complaint proceeds

on general principles; rest in the recumbent posture, abstinence, and saline aperients, with leeches locally applied, will mitigate all the symptoms. But that from which the greatest relief is to be obtained is the early opening of the abscess, into which a lancet should be introduced as soon as the touch detects that suppuration is commencing, or the length of time makes it probable that the inflammation is pointing. There is another rule respecting such abscesses; when they have matured, and, as then commonly happens, enclose and isolate the rectum in a bag of matter, it is necessary of course to give the freest escape to the matter; but this is to be obtained not by a large incision, but by two or three punctures of the breadth of a lancet-blade on different sides of the anus. So treated, these abscesses speedily contract and heal.

The smaller abscesses that form in the neighbourhood of the rectum present different features. Sometimes they attain the size of an egg without exciting uneasiness enough to lead the patient to observe the complaint; and the abscess bursting is the first intimation he gets of its existence. In the other extreme there is an acute circumscribed inflammation, with considerable tension, pain, heat, aching, and throbbing for several days before matter forms. Intermediately between these extremes every variety is met with. The course and progress of such a phlegmon varies likewise in different instances. If

taken early, and leeches are applied and fomentations, and the patient is kept in the horizontal posture, the diet and action of the bowels being properly regulated, the phlegmon sometimes disperses. Commonly, however, suppuration takes place, and it is then important to open the abscess, as soon as anything like softening is felt. By this means fistula may often be prevented. The part is then to be poulticed, and the patient kept at rest, and in the horizontal posture, that the abscess may dry up and close speedily.

The situation of these abscesses varies. Sometimes they are exactly anal, or are situated at the side of the aperture of the intestine. I have known such an abscess treated as an anal pile, and the patient remain in pain for days, which puncturing with a lancet has at once removed. Sometimes the abscess is situated upon the side of the nates, removed two or three inches from the anus. Sometimes it seems to have more to do with the os coccygis than with the intestine. At other times it forms in the perineum, and threatens rather to implicate the urethra, which is indeed occasionally opened by it; so that the first consequence of the complaint is urinary fistula.

The evil of these abscesses is, that they are liable to become sinuses, or fistulæ. A sinus is a tubular canal in the cellular tissue, commonly just large enough to admit a probe, which secretes, from its



lining membrane, a thin matter, and has a disposition not only to continue indolent, but to extend itself in new directions. Or a sinus originates in an abscess which, instead of closing, contracts to form such a canal. Abscesses occurring in other parts are liable to leave sinuses; more especially abscesses in the groin and in the axilla. When sinuses occur near the anus, they may extend either along the perineum beneath the integument, or along the sacrum, or by the side of the rectum. What is properly termed fistula, exists, when such a tube, at its inner extremity, has an inward communication with a mucous surface. For example, when the tubular canal runs *towards* the urethra only, it is a sinus; when it opens *into* it, it is a urinary fistula; and the same in reference to the rectum. However, the term blind fistula is commonly used to denote a mere sinus running parallel to the intestine and not communicating with it. Occasionally an abscess upon the intestine opens into it, without pointing externally.

*Fistula ani*, therefore, presents three essential varieties; it may have one opening alone, and that by the side of the anus; or its one opening may be into the bowel; or it may open both externally and internally, and allow fæcal matter to escape through it.

There is one common plan of treatment, under which each of these kinds of fistula occasionally

heals. That is, regulation of diet, tonic medicine, and change of air to improve the state of the system, the use of aperients to prevent costiveness, the application of local stimulants.

But generally these means fail, and it becomes requisite to have recourse to a surgical operation; that is to say, unless the inconvenience arising from the fistula is extremely slight, the sinus being narrow, the quantity of moisture discharged inconsiderable, and the patient out of health, or aged.

If the fistula has two openings, a probe-pointed bistoury is passed from the outer opening into the intestine, and being there received upon the forefinger of the left hand, is drawn out, dividing the intervening thickness. The wound is made to heal by granulation from the bottom, through the introduction of a thin piece of lint, replaced as often as it is necessary.

If the fistula has an external opening only, the same operation is requisite, the end of the bistoury being made to perforate the intestine opposite the end of the sinus.

If the fistula has an internal opening only, (when it is often most difficult of detection, soreness only, and occasional pain and slight discharge of matter, indicating it,) a parallel, but slighter operation is to be performed; the bistoury being passed into the sinus from within, and the lining of the bowel being thence divided to the sphincter.

If the fistula open into the intestine some distance from the anus, from two to three inches, the section of so much of the intestine would risk the division of an artery of sufficient magnitude to cause troublesome hemorrhage. In such a case it is best to thread the sinus with a flexible wire, which being carried into the rectum, and brought out at the anus, allows its ends to be twisted together on the outside. Then by tightening the twisted knot, the sinus is led to ulcerate in a direction towards the perineum, and simultaneously to close above. When it has been thus rendered altogether superficial, the common operation may be performed to complete the cure.

Sometimes it happens that the wound made in the operation is slow to heal: in that case stimulant applications are to be used, and tonic medicines, a liberal diet, and change of air, are to be resorted to, to recruit the patient's strength.

Sometimes it happens that fistulæ, which communicate at one or more points with the intestine, burrow further as blind sinuses in one or more directions. In general it is sufficient to lay these open as far as they run beneath the skin, or to their communications with the intestine. At all events when they extend deep, this should be first tried, though it may afterwards prove requisite to lay them open to their extreme depth.

Fistulæ often occur in persons of undermined

constitution,—with disease of the lungs, or liver, and broken health. In such persons, an operation is not to be recommended; as it would only give gratuitous pain, with no reasonable chance of curing the malady.

#### SECTION XI.—CANCER OF THE RECTUM.

FOR consideration under this head, I have reserved a variety of cases, which, differing among themselves in several pathological features, yet agree in their symptoms, in their fatal course, in the treatment which they require.

Anatomically, they so far agree, that they each consist in a local change of structure, in which the natural texture becomes more or less gristly, the lining membrane abraded or ulcerated, and the passage through the intestine narrowed.

Anatomically, they differ thus: in one variety the thickness of the coats of the intestine remains nearly unaltered; only the part becomes a contracted gristly tube. In the other forms the diseased part enlarges in bulk; the increased growth bearing characters which are familiar to the scientific surgeon, under the names of carcinoma, medullary carcinoma, gelatiniform sarcoma, melanoma.

Diseases of the rectum are more frequent in women than in men, a difference, the general causes of which are apparent in the varying conditions

of the uterus and in the shape of the female pelvis. The womb through its increased size in pregnancy interrupts the free return of blood by the hemorrhoidal veins, and mechanically obstructs the action of the bowels. In the unimpregnated state, the periodical congestion of the same organ cannot fail of extending its influence to the vessels of the rectum. The straightness of the sacrum, again, in women, and the general expansion of the pelvic bones at the inferior outlet, deprive the lower part of the bowel of that pressure and support which appear to conduce to the healthy state of the viscera. Something, too, is to be attributed to the less regular relief of the bowels and of the bladder, and their frequent over-distension, which result from the natural, but in this instance, prejudicial delicacy of women. The comparative frequency of *cancer* of the rectum in women depends upon the co-operation of another and a more serious cause, which is the prevalence in women of the cancerous diathesis.

Of the different forms of the disease, that in which there is no enlargement is again the most frequent kind in women; or I have seen many cases of this variety in women, and not one in men. In all the cases of the complaint that I have seen in men, there has been schirrhous enlargement.

The symptoms of cancer of the rectum, slow in their early progress, and hardly observed at first,

are at length developed into local pain, sometimes of a dull aching character, at other times acute and lancinating, with a sense of weight and confinement in the part, numbness and uneasiness extending round the loins to the pubes, numbness of the hips and thighs. These sensations are aggravated upon walking, standing, or sitting, and are mitigated by the recumbent posture. The act of evacuating the bowels increases the pain and distress. The fæces are either liquid, or are passed in small fragments and by repeated efforts: blood, matter, mucus, are expelled with them. The patient, if a female, suffers, in addition, bearing down pains of the womb, irritation of the bladder, pain in making water, incontinence of urine. The difficulty and occasional obstruction of the passage of fæces produce fits of distension and of general pain and tenderness of the abdomen, accompanied with hiccup and vomiting.

The situation of cancer of the rectum depends, in some degree, upon its kind.

That which is without thickening generally commences half an inch or an inch within the anus, and from this point occupies from two or three inches to thirteen or fourteen. Sometimes, but rarely, the hardness and ulceration commence quite at the orifice.

Cancer with thickening is commonly situated two or three inches from the anus; or here is the

lower edge of the mass of disease, which may thence extend to two, three, or five or six inches of the intestine; generally it occupies about three or four inches of the bowel, and involves its entire circumference. Sometimes even to the end it is much more circumscribed, and occupies a small part only of one side of the bowel.

The severity of suffering in cancer of the rectum bears no proportion to the length of intestine implicated in it. There are two ways in which the disease proves fatal. In general the patient consumes gradually, worn out by long-continued suffering. In other instances the disease produces complete obstruction of the bowels, and the patient perishes more quickly. This form of the malady is attended with the severest distress. Complete obstruction of the bowels rarely occurs except in cases where there is a large mass of gristly substance in the intestine. But I have known the disease terminate in fatal obstruction, and with the most aggravated suffering, when the mass occupied no more than the last three inches of the rectum.

The treatment to be pursued in carcinoma of the rectum is the following.

The pain is to be allayed by opiates. Solid opium, laudanum, the acetate and muriate of morphia, the extract of stramonium, are each in their turn available. Opiates act more promptly and efficaciously in this disease when taken into

the stomach, than when applied in injections or as suppositories. If administered in the latter form, the subacetate of lead may be advantageously combined with them.

The bowels in the early stages of the complaint should be regularly relieved. In general a drachm of the lenitive electuary taken over-night, with an injection of tepid water the following morning, will be sufficient for this purpose. A few drops of laudanum may be taken, if necessary, immediately after the action of the bowels.

The canal of the intestine is to be dilated with a bougie, if it is so narrowed by the disease as to obstruct materially the passage of the fæces.

When the canal is obstructed, it is requisite to introduce a long flexible tube past the cancer, and by its means to break down, softening at the same time by water injected through it, the fæces that have accumulated above.

It sometimes happens that the disease is situated at the orifice of the intestine. This case presents two varieties of very unequal importance. In one the disease occupies not more than a square inch of one side of the anus. In the other it occupies the whole circumference of the end of the bowel. In either case the disease, which is curable by no other means, admits of being removed by the excision of the part. But while the operation in the first case is not long about, and is not attended with



severe pain, and always in my experience has done well, and the patient has recovered in a very short space of time, and has afterwards preserved considerable retentive power over the intestine; in the other, the operation has proved tedious and dangerous, and at best has left a serious mutilation. The first of these operations I have done three times with perfect success: the last twice only. In the latter two instances I removed more than an inch and a half of the circumference of the bowel including the sphincter. One of the patients survived two years, and died accidentally of peritonitis but the disease had begun to threaten a return, and there was most troublesome prolapsus. The other died a few weeks after the operation.

Cancer of the rectum occasionally assumes the character of polypus; and the pendulous growth is liable to be forced down whenever the bowels act. Such a tumour has no sensibility; it had better be removed by ligature. In a case of this kind, which was under my care some years ago, the polypus was thus taken away very much to the comfort of the patient. The pendulous growth was not reproduced. But a mass of indurated substance formed upon the fore-part of the bowel within the anus. Part of this I likewise removed by ligature, but the fungus again increased, and the complaint ended fatally.

I have mentioned discharge of blood, mucus, and

pus, as some of the symptoms of cancer of the rectum. But these symptoms are often present, where cancer does not exist; blood, it has been explained, may exude from the mucous membrane without any alteration of the texture of the latter; mucus is poured out in increased quantity upon any irritation of the part; and even pus or puriform liquid may be secreted without any breach of surface. Any great quantity of either of these products being formed, is against the disease being cancer. Plentiful discharge of mucus from the intestines, with narrowed and fragmentary fæces, commonly indicates only spasmodic stricture of the rectum. Even the separation of what seem portions of the mucous membrane itself, being in fact *false membrane* only, and that in considerable quantity, with constant local uneasiness, and general impairment of health, proceeds only from partial subacute inflammation of the lining of the intestine. In other persons the same degree of inflammation, enduring for months, sometimes for years, will be attended with immense secretion of puriform or opaque yellowish liquid, which is passed, almost without the possibility of retaining it, several times during the day. By means of alterative courses of medicine, and the use of astringent injections, and time, this distressing ailment may be mitigated, and sometimes cured.

---

## WORKS BY MR. MAYO.

---

### THE PHILOSOPHY OF LIVING.

For though we Christians do continually aspire and pant after the Land of Promise; yet it will be a token of God's favour towards us, in our journeyings through the world's wilderness, to have our shoes and garments (I mean those of our frail bodies) little worn or impaired.—*Bacon*.

---

### OUTLINES OF PHYSIOLOGY.

FOURTH EDITION, 1837.

The information contained in the present volume is at the same time so complete, so much condensed, and so clearly arranged, that we conceive it ought to be in the hands of every student of physiology, and of every one who wishes to obtain a just view of the structure of the human frame, the uses of its various organs, and the nature of animal bodies generally.—*Edinburgh Medical Review, July, 1834*.

---

### OUTLINES OF PATHOLOGY.

In barely more than a page we find condensed nearly all that is certainly known of the pathology of dropsy. We cannot forego the pleasure of presenting our readers with this passage, &c.

Cases are related illustrating distinctions in the anatomical appearances of cancer, to which succeeds an excellent account of the progress of that disease in the breast. The remarks upon an important practical question are very judicious.

We cannot afford space for noticing the diseases of joints, the description of which, as of other subjects of surgical pathology, the author conducts with even more than usual ability.—*Forbes's Medical Review, January, 1837*.

---

# NEW BOOKS,

PUBLISHED BY

JOHN W. PARKER, WEST STRAND,

LONDON.

---

**WOMAN'S RIGHTS and DUTIES**, considered in their Principles, and their Influence on Society. By a WOMAN. Two Volumes, Post Octavo.

**WOMAN'S MISSION.** EIGHTH EDITION. 3*s.* 6*d.*

“If women could once be made to understand their real mission in this world, and to feel their own importance and responsibility, a surprising change must immediately take place in society, giving it a higher tone and purer spirit.”

**THE YOUNG LADY'S FRIEND.** A Manual of Practical Advice and Instruction to young Females on their entering upon the Duties of Life. Third Edition, 3*s.* 6*d.*

**THE FAMILY HAND-BOOK, or PRACTICAL INFORMATION** in DOMESTIC ECONOMY; including Cookery, Household Management, and all Subjects connected with the Health, Comfort, and Expenditure of a Family. With Choice Receipts. 5*s.*

**THE DOMESTIC GARDENER'S MANUAL;** being an Introduction to Practical Gardening, on Philosophical Principles; to which is added, a NATURALIST'S CALENDAR, and an Appendix on the Operations of Forcing, including the Culture of Vines in Pots. By JOHN TOWERS, C.M.H.S. Second Edition, Enlarged and Improved, Octavo, 13*s.*

**ORIGINAL FAMILY SERMONS;** by ONE HUNDRED AND FIFTY CONTEMPORARY DIVINES OF THE ESTABLISHED CHURCH. Five Volumes, at 6*s.* 6*d.* each.

**SACRED MINSTRELSY;** a COLLECTION of the FINEST SACRED MUSIC, by the best Masters, arranged as Solos, Duets, Trios, &c., and Choruses; and with Accompaniments for the Piano-Forte or Organ. Two handsome Folio Volumes, 2*l.* 2*s.* half-bound.

**LETTERS of EMINENT PERSONS;** Selected and Illustrated, with Biographical Notices, by R. A. WILLMOTT, Author of *Lives of the English Sacred Poets, &c.* Post 8vo. 7s. 6d.

**THE RUBI,** a Tale of the Sea; a Poem, in Six Cantos. By F. W. MANT, late R.N. Foolscap Octavo, 4s. 6d.

**MUSICAL HISTORY, BIOGRAPHY, and CRITICISM;** being a GENERAL SURVEY of MUSIC from the earliest period to the present time. By GEORGE HOGARTH. *A new and enlarged Edition, in Two Volumes.* 10s. 6d.

**ROADS AND RAILROADS, TUNNELS, BRIDGES, and CANALS,** popularly described; with an Account of the VARIOUS VEHICLES and MODES of TRAVELLING IN ALL PARTS OF THE WORLD, from the earliest to the present Time. One Volume, Illustrated by upwards of a Hundred Wood-Cuts. 5s.

**HUMBOLDT'S TRAVELS and DISCOVERIES in AMERICA.** With Engravings, 2s. 6d.

**MUNGO PARK;** his LIFE and TRAVELS, with an Account of his Death, from the Journal of ISAACO, and the substance of later Discoveries relative to his fate. With Engravings. 2s. 6d.

**CAPTAIN COOK;** his LIFE, VOYAGES, and DISCOVERIES. With Engravings. 2s. 6d.

**CHRISTOPHER COLUMBUS;** his LIFE, VOYAGES, and DISCOVERY of the NEW WORLD. With Engravings. 2s. 6d.

**THREE WEEKS in PALESTINE and LEBANON.** With Engravings. 3s.

**TRAVELLING SKETCHES in EGYPT and SINAI,** including a Visit to Mount Horeb, and other localities of the Exodus; Abridged from the French of DUMAS. By A BIBLICAL STUDENT. 3s. 6d.

GERMANY, BOHEMIA, and HUNGARY, visited in 1837. By the Rev. G. R. GLEIG, M. A., Author of *The Subaltern*. Three Vols., Post 8vo., strongly bound in ornamental cloth. 11. 11s. 6d.

THE CITY of TREVES and its ROMAN ANTIQUITIES. From the German of WYTTEBACH. Edited, with NOTES, by DAWSON TURNER, Esq., F.R.S., F.S.A. Illustrated from Drawings made on the spot. Octavo, 8s. 6d.

RESEARCHES IN BABYLONIA, ASSYRIA, AND CHALDÆA; forming part of the Labours of the Euphrates Expedition, and published with the sanction of the Board of Control. By WILLIAM AINSWORTH, Surgeon and Geologist to the Expedition. With Illustrations, Maps, &c. 12s. 6d.

RECREATIONS in PHYSICAL GEOGRAPHY, or the EARTH AS IT IS. By MISS R. M. ZORNLIN. Foolscap Octavo, with numerous Illustrations, 6s.

RECREATIONS in ASTRONOMY, with a Glossary. By Rev. LEWIS TOMLINSON, M. A., with more than Fifty Illustrations, Foolscap Octavo, 4s. 6d.

RECREATIONS in GEOLOGY, with a Preliminary Discourse on the Nature and Advantages of Geology, and a Glossary. By R. M. ZORNLIN. With Illustrations. 4s. 6d.

RECREATIONS in CHEMISTRY. By THOMAS GRIFFITHS, Chemical Lecturer at St. Bartholomew's Hospital. *In the Press*.

EASY LESSONS IN MECHANICS: with Familiar Illustrations of the Practical Application of Mechanical Principles. A new Edition. 3s.

READINGS in SCIENCE; familiar EXPLANATIONS of Appearances and Principles in NATURAL PHILOSOPHY. A new Edition, with many Engravings. 5s.

THE STUDENT'S MANUAL OF NATURAL PHILOSOPHY; comprising Descriptions, Popular and Practical, of the most important Philosophical Instruments, their History, Nature, and Uses. Dedicated, by permission, to the Lord Bishop of Salisbury. By CHARLES TOMLINSON. 10s. 6d.

**An INTRODUCTION to the STUDY of CHEMICAL PHILOSOPHY**; being a preparatory View of the Forces which concur to the Production of Chemical Phenomena. By **J. FREDERIC DANIELL**, F.R.S., Professor of Chemistry in King's College, London; Author of *Meteorological Essays*. 16s.

**LECTURES on ASTRONOMY**, delivered at King's College, London. By **PROFESSOR MOSELEY**, of King's College, London. Numerous Illustrations. 5s. 6d.

**MECHANICS APPLIED TO THE ARTS**. By **PROFESSOR MOSELEY**, of King's College, London. A New Edition, improved. With numerous Engravings. 6s. 6d.

**PRACTICAL GEOLOGY and MINERALOGY**, including the **CHEMISTRY of METALS**. By **JOSHUA TRIMMER**, F.G.S. With numerous Illustrations. *In the Press*.

**A DICTIONARY of MATERIA MEDICA and PRACTICAL PHARMACY**; including a Translation of the Formulæ of the London Pharmacopœia. By **WILLIAM THOMAS BRANDE**, F.R.S.L. and E.; of Her Majesty's Mint; Professor of Chemistry and Materia Medica to the Apothecaries' Company. In a closely printed Octavo Volume, 15s.

**A MANUAL of CHEMISTRY**. By **WILLIAM THOMAS BRANDE**, F.R.S., &c., of Her Majesty's Mint; Professor of Chemistry at the Royal Institution of Great Britain. **THE FIFTH EDITION**, revised and enlarged, and embracing all the newest Facts and Discoveries. *In the Press*.

**THE PHILOSOPHY of the INDUCTIVE SCIENCES**, founded upon their History; containing the Philosophy of Pure Mathematics; the Philosophy of the Mechanical Sciences; of Physic; of Chemistry; of Mineralogy; of Botany; of Zoology; and the Logic of Induction. By **WILLIAM WHEWELL**, B.D., F.R.S., Fellow of Trinity College, and Professor of Moral Philosophy in the University of Cambridge. Two Vols. Octavo, 30s. Also, by the same Author,

**THE HISTORY of the INDUCTIVE SCIENCES**, from the Earliest Times to the Present. Three Vols. Octavo, 2l. 2s.

**ON the FOUNDATION of MORALS;** Four Sermons preached before the University of Cambridge. By the Rev. W. WHEWELL. Second Edition, 3s. 6d.

**FULLER'S HISTORY of the UNIVERSITY of CAMBRIDGE.** By the Rev. M. PRICKETT, M.A., F.S.A., and T. WRIGHT, M.A., F.S.A., Trinity College, Cambridge. A New Edition, with Notes. Octavo, 12s.

**THE LIFE and SERVICES of GENERAL LORD HARRIS, G.C.B.,** during his CAMPAIGNS in AMERICA, the WEST INDIES, and INDIA. By the Right Hon. S. R. LUSHINGTON, Private Secretary to Lord Harris, and late Governor of Madras. Octavo, with an Original Portrait, 13s.

**LIVES of EMINENT CHRISTIANS.** By R. B. HONE, M.A., Vicar of Hales Owen. New Edition. Three Volumes, 4s. 6d. each.

Vol. I.	Vol. II.	Vol. III.
ARCHBISHOP USHER, DOCTOR HAMMOND, JOHN EVELYN, BISHOP WILSON.	BERNARD GILPIN, PHILIP DE MORNAY, BISHOP BEDELL, DOCTOR HORNECK.	BISHOP RIDLEY, BISHOP HALL, The Hon. ROBERT BOYLE.

**LIVES of ENGLISH SACRED POETS.** By R. A. WILLMOTT, Esq., Trinity College, Cambridge. Now complete, in Two Volumes, at 4s. 6d. each.

The FIRST SERIES contains an Historical Sketch of Sacred Poetry, and the Lives of the English Sacred Poets preceding MILTON.

The SECOND SERIES commences with MILTON, and brings down the Lives to that of BISHOP HEBER inclusive.

**THE LIFE of ARCHBISHOP SANCROFT.** By GEORGE D'OYLY, D.D., F.R.S., Rector of Lambeth. Octavo, Second Edition, Revised, with a Portrait. 9s.

**BISHOP BUTLER.** Memoirs of his Life, Character, and Writings. By THOMAS BARTLETT, M.A., One of the Six Preachers of Canterbury Cathedral, and Rector of Kingstone, Kent. With an Original Portrait, 8vo., 12s.



**THE WORKS OF DOCTOR DONNE**, Dean of Saint Paul's in 1619—1631; with a Memoir of his Life, and Critical Notices of his Writings. By HENRY ALFORD, M.A., Vicar of Wymeswold, Leicestershire, and late Fellow of Trinity College, Cambridge. Six Vols., 8vo., 3*l.* 12*s.*

**THE CHURCHMAN'S GUIDE**; a Copious Index of Sermons and other Works, by eminent Church of England Divines; digested and arranged according to their subjects and continued to the present day. By the Rev. JOHN FORSTER, M.A., Incumbent of the Royal Chapel in the Savoy. Octavo, 7*s.*

**GARRICK'S MODE of READING the LITURGY** of the CHURCH of ENGLAND. A New Edition, with Notes, and a PRELIMINARY DISCOURSE on PUBLIC READING. By RICHARD CULL, Tutor in Elocution. Octavo, 5*s.* 6*d.*

**THE RITUAL OF THE UNITED CHURCH** of ENGLAND and IRELAND ILLUSTRATED. With occasional Reference to the Objections of Dissenters. By the Rev. G. W. TYRRELL, A.M., Secretary of the Down and Connor Diocesan Church Education Society. 6*s.*

**A MANUAL of CHRISTIAN ANTIQUITIES**; or, an Account of the Constitution, Ministers, Worship, Discipline, and Customs of the Early Church; with an Introduction, containing a Complete and Chronological Analysis of the Works of the Antenicene Fathers. By the Rev. J. E. RIDDLE, M.A., Author of a *Latin-English Dictionary, Luther and His Times, &c.* In a large Octavo Volume. 18*s.*

**HISTORY of the CHURCH of IRELAND**, from the REFORMATION to the REVOLUTION. By the Right Rev. RICHARD MANT, D.D., Lord Bishop of Down and Connor. Octavo, 17*s.*

**HISTORY of the CHURCH of ENGLAND**, to the REVOLUTION in 1688; embracing Copious Histories of the Thirty-Nine Articles, the Translation of the Bible, and the Compilation of the Book of Common Prayer. By THOMAS VOWLER SHORT, D.D. THIRD EDITION, Octavo, 16*s.*

**ON THE EARLY FATHERS**; an Introductory Lecture to the Course delivered at Cambridge. By the Rev. J. J. BLUNT, B.D., Lady Margaret's Professor of Divinity in that University. Octavo, 2s.

**THE VICTORY of FAITH**, and other SERMONS. By JULIUS CHARLES HARE, M.A., Rector of Herstmonceux, Sussex; and late Fellow of Trin. Coll., Camb. 10s. 6d.

**THE SCRIPTURAL CHARACTER of the ENGLISH CHURCH CONSIDERED**, in a Series of Sermons, with Notes and Illustrations. By the Rev. DERWENT COLERIDGE, Master of Helleston Grammar School, Cornwall. 12s. 6d.

**THE STANDARD of CATHOLICITY**; or, an Attempt to point out in a plain Manner, certain, safe, and leading Principles, amidst the conflicting Opinions by which the Church is at present agitated. By the Rev. G. E. BIBER, LL.D. 14s.

**THE DISTRICT VISITOR'S MANUAL**; a Compendium of Practical Information and of Facts, for the Use of District Visitors; with a PREFACE, by the Rev. THOMAS DALE, M.A., Vicar of St. Bride's, London; and a SERMON, by the Rev. CHRISTOPHER BENSON, M.A., Master of the Temple. Second Edition, 4s. 6d.

**EXPLANATORY NOTES on the PRAYER-BOOK VERSION of the PSALMS**. By WM. KEATING CLAY, B.D., Jesus Coll., Cambridge, and Minor Canon of Ely. 7s. 6d.

**UNIVERSAL MYTHOLOGY**; an Account of the most important Mythological Systems, their Origin and Connexion. With Considerations on the KORAN and the TALMUD. By the Rev. HENRY CHRISTMAS, St. John's College, Cambridge. 7s.

**THE NEW CRATYLUS**; or, CONTRIBUTIONS towards a more ACCURATE KNOWLEDGE of the GREEK LANGUAGE. By the Rev. JOHN WILLIAM DONALDSON, M.A., Fellow of Trinity College, Cambridge. 17s.

**THE GREEK TEXT** of the **ACTS** of the **APOSTLES**; with Notes, Original and Selected; for the Use of Students in the University. By **HASTINGS ROBINSON, D.D., F.A.S.**, Rector of Great Warley. 8s.

**THE CAMBRIDGE GREEK** and **ENGLISH TESTAMENT**; the Greek (from the **THIRD EDITION** of **STEPHENS**), and the English from the Authorized Version, being given in Parallel Columns on the same Page. 8s. 6d.

**THE AULULARIA** of **PLAUTUS**, with Notes by **JAMES HILDYARD, M.A.**, Fellow of Christ's College, Cambridge. 7s. 6d.

**THE MENÆCHMEI** of **PLAUTUS**, with a Glossary and Notes. By the Same. A New Edition. 7s. 6d.

**A LIFE** of **ARISTOTLE**, including a Critical Discussion of some Questions of Literary History connected with his Works. By **J. W. BLAKESLEY, M.A.**, Fellow of Trinity College, Cambridge. Octavo, 8s. 6d.

**STEMMATA ATHENIENSIA**; Tables of Biography, Chronology, and History, to facilitate the Study of the Greek Classics. Quarto, 5s.

**THE FROGS** OF **ARISTOPHANES**, with English Notes, for Schools and Students. By the Rev. **H. P. COOKESLEY**, Editor of *The Plutus* and *The Birds*. 7s.

**SCHLEIERMACHER'S INTRODUCTIONS** to the **DIALOGUES** of **PLATO**; translated from the German, by the Rev. **W. DOBSON, M.A.**, Fel. Trin. Coll., Camb. 12s. 6d.

**DISSERTATIONS ON THE EUMENIDES** OF **ÆSCHYLUS**, with the Greek Text, and Critical Remarks. From the German of **MULLER**. Octavo, 9s. 6d.

THE HISTORY OF THE ARTS OF THE  
ANCIENTS AND MODERNS, FOR THE USE OF  
SCHOOLS AND COLLEGES. BY JOHN W. PARKER, D.D.

THE HISTORY OF THE ARTS OF THE  
ANCIENTS AND MODERNS, IN ENGLISH  
(Saxons), and in Parallel

THE HISTORY OF THE ARTS OF THE  
ANCIENTS AND MODERNS, IN  
Cam-

THE HISTORY OF THE ARTS OF THE  
ANCIENTS AND MODERNS, IN  
A Class

THE HISTORY OF THE ARTS OF THE  
ANCIENTS AND MODERNS, IN  
the

THE HISTORY OF THE ARTS OF THE  
ANCIENTS AND MODERNS, IN  
of the

THE HISTORY OF THE ARTS OF THE  
ANCIENTS AND MODERNS, IN  
with H. E.

THE HISTORY OF THE ARTS OF THE  
ANCIENTS AND MODERNS, IN  
to

THE HISTORY OF THE ARTS OF THE  
ANCIENTS AND MODERNS, IN  
OF

THE HISTORY OF THE ARTS OF THE  
ANCIENTS AND MODERNS, IN  
From

PUBLISHED BY JOHN W. PARKER,

THE GREEK TEXT of the ACTS of  
APOSTLES, with Notes, English and German. By the  
Reverend Wm. Burgon. By GASTON ROBINSON, D.  
D., Bishop of Exeter, &c.

THE CAMBRIDGE GREEK and ENGLISH  
TESTAMENT, the Greek Text, the Textus Receptus, &c.  
and the English Version, being given in  
conjunction with the Latin Vulgate, &c.

THE AULULARIA of PLAUTUS, with Notes  
by JAMES HILLIARD, M.A., Fellow of Trinity College,  
Oxford. 7s. 6d.

THE MENAECHMI of PLAUTUS, with a  
Critical Edition. By the same. 7s. 6d.

A LIFE of ARISTOTLE, including a Critical  
Edition of his Works, of Literary History, &c.  
By Wm. G. W. BLAKESLEY, M.A., Fellow of Trinity  
College, Oxford. 7s. 6d.

STEMMATA ATHENIENSIA; Tables of  
Genealogy, &c., for the Study of  
Greek Literature. 7s. 6d.

THE FROGS of ARISTOPHANES,  
English Verse for Schools and Colleges. By G. W. G.  
BLAKESLEY, Master of The Queen's and The Duke's  
Schools, &c.

SCHLEIERMACHER'S INTRODUCTIONS  
to the WORKS of PLATO, translated into English  
by Wm. B. EYRE, M.A., F.R.S., &c.

DISSERTATIONS ON THE PUMENIDES  
of ARISTOPHANES, with the Greek Text and Critical Remarks,  
by G. W. G. BLAKESLEY. 7s. 6d.

London: JOHN W. PARKER, Paternoster Row, Strand.

