Salivation exploded: : or, a practical essay on the venereal disease, fully demonstrating the inefficacy of salivation, and recommending an approved succedaneum. Illustrated with some remarkable cases, which had withstood three, four, or five salivations, and were afterwards cured, by that safe, easy and certain method, the alterative one. To which is added, a dissertation on gleets and weaknesses, seminal as well as venereal, in both sexes; with observations on diseases of the urethra, and the use of prophylactics. / By Charles Swift, surgeon.

Contributors

Swift, Charles. University of Glasgow. Library

Publication/Creation

London: Printed for S. Bladon, No. 13, Pater-Noster Row; R. Faulder, New-Bond Street; P. Brett, opposite St. Clement's Church, Strand; Walter and Bell, No. 169, Piccadilly, MDCCLXXXIX [1789]

Persistent URL

https://wellcomecollection.org/works/szz9yh9e

Provider

University of Glasgow

License and attribution


This material has been provided by This material has been provided by The University of Glasgow Library. The original may be consulted at The University of Glasgow Library. where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.




Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org






Digitized by the Internet Archive in 2016











SALIVATION EXPLODED:

OR, A

PRACTICAL ESSAY ON THE

VENEREAL DISEASE,

FULLY DEMONSTRATING THE

Inefficacy of Salivation, and recommending an approved Succedaneum.

ILLUSTRATED

With some remarkable Cases, which had withstood Three, Four, or Five Salivations, and were afterwards cured, by that safe, easy and certain Method, the Alterative One.

TO WHICH IS ADDED,

A Differtation on Gleets and Weaknesses, Seminal as well as Venereal, in both Sexes; with Observations on Diseases of the Urethra, and the Use of Prophylactics.

By CHARLES SWIFT, SURGEON.

Opprime dum nova funt, fubiti mala femina morbi, nam mora dat vires. Ovip.

THE SEVENTH EDITION.

LONDON:

Printed for S. BLADON, No. 13, Pater-Noster Row; R. FAULDER, New-Bond Street; P. BRETT, Opposite St. Clement's Church, Strand; WALTER and BELL, No. 169, Piccadilly.


M DCCLXXXIX.

[Price Two Shillings.]

SWIFT (Charles)

Sp. Coll.

64/4679



ADVERTISEMENT.

Notwithstanding the number and variety of publications upon the present subject, the Author slatters himself that his observations (the result of much experience) will not be unacceptable to the Faculty; especially the junior part of it, to whose perusal this Essay is particularly recommended.

Common cases, or such as daily occur, and are not attended with any unusual symptoms, are purposely omitted; and to avoid prolixity, a few only of the other kind are inserted, they being thought sufficient to elucidate the subject.

Recent cases are likewise pretermitted, in order to obviate the idea of relapses, which even the most impartial reader might otherwise have suggested.

Parliament Street, June 26th, 1780. THE TISH ENTENT tide begins state of the or engineer stronger while

PRACTICAL ESSAY

ONTHE

VENEREAL DISEASE, &c.

HE Venereal Disease, beyond all others, having by its peculiar virulence and frequent attacks upon the human body, made the most alarming depredations therein, no temporal subject, I presume, ought sooner to claim our attention, than the thorough investigation and speedy subjection of so formidable an enemy .- Many and various have been the opinions of the learned, relative to the time and place of its first appearancethis circumstance has afforded an ample field for disputation to both ancients and moderns; of whom fome the most eminent, at different periods, have very ingeniously and candidly treated upon the fubject: notwithstanding which, the matter is yet doubtful and likely to remain fo. A knowledge of the precise origin of the disease must however appear of but little importance, when put in competition with that of the most effectual remedy for the radical cure of it. I shall therefore leave such controversies to those who are fonder of, or may be better qualified to adjust them than myself, and rest contented with recommending the most radical and easy method of removing so loathsome a disease. This mode of cure is the most gentle and efficacious, as will appear in the following pages: it will stand the test of the strictest scrutiny, being sounded on the broad basis of long experience and uninterrupted success. In support of this affertion, I shall produce some very remarkable cases, selected from many hundreds, where the usual or general modes of cure had proved abortive.

It has been frequently observed, that the knowledge of a disease is half its cure; in no instance can I suppose this adage to have been more fully verified than in the Venereal Disease; which, Proteus like, appears in so many different forms, that very often it will require no small share of judgment and experience to distinguish it from some other distemper, whose symptoms it may resemble in many respects, or with which

which it may in reality be combined. This being the case we must not wonder at the many errors daily committed by the ignorant, or inexperienced in the profession.

ALMOST innumerable have been the instances of reputed rheumatisms, nervous head-achs, wandering gouts, scurvies, declines, consumptions, &c. &c. which, upon a minute investigation, have been proved to be venereal. A happy circumstance, when this discovery is made early enough to save the patient from an untimely grave; but alas! how frequently does the reverse happen?

This melancholy observation is particularly corroborated in the great number of atrophies, or consumptions, by which the inhabitants of this island are many of them supposed to be destroyed. This fatality has generally been ascribed to the climate, but I am sirmly persuaded, that, upon a strict and impartial enquiry, many other causes may be assigned; and amongst them, Venus will be found to have had no

B 2

fmall

fmall share. I am induced to be of this opinion from the many instances that have fallen under my own immediate inspection, where patients, after having been pronounced to be in the last stages of a confumption, and given over as lost, have been perfectly restored by the use of antivenereals. A circumstance of such importance, most certainly merits the attention of every medical practitioner. Should this admonition take the desired effect, I shall experience the greatest satisfaction in having rendered such an essential service to the community.

In the following sheets it will evidently appear, that the Venereal Disease, in every state or stage, from the mildest to the most inveterate, may be perfectly, easily, and safely cured, without salivation; and that the patient, although labouring under a confirmed pox, will not be confined within doors a single day during the whole time he may be taking such medicines as will effect a radical cure. I shall likewise demonstrate that gleets of every denomination

are curable, provided the origin of each be duly investigated, and in consequence thereof, a suitable remedy not only applied, but regularly persevered in for a length of time, proportionable to the cause: to the want of that perseverance, in my opinion, may often be ascribed the want of success in the general treatment of gleets; for I do not recollect (amongst the great number and variety of my patients for several years past) three cases which did not give way, or were not cured, by properly and strictly persisting in the use of injections or bougies, and some internal alteratives.

To prove not only the inexpediency, but the inefficacy of falivation in venereal maladies, I shall produce several cases, in which its effects were merely palliative; in other words, afforded only a temporary relief, although repeated even to a fourth or fifth time; after which the same patients (not-withstanding the violent shock, and manifest injury their constitutions had thereby sustained) were perfectly and radically cured by that easy gentle method, which I mean to recommend, namely, the alterative one.

For argument fake only, suppose we admit that falivation is efficient and might be depended upon; what man in his fenfes but would give the preference to the method here proposed? When he considers how extremely difgusting and painful the process of falivation is universally allowed to be, the consequent confinement indispenfably necessary therein, and the many other difagreeables that generally or frequently succeed it; such as destruction of the teeth and gums, and the fœtid or stinking breath thereby occasioned: the great relaxation of the falivary glands, or an erofion of their excretory ducts, producing a copious troublesome flow of saliva, perhaps for the remainder of life; together with the derangement of the whole frame, in fuch a degree as feldom to be got the better of, even by those who may have been originally possessed of the best stamina, or strongest constitutions.

ALL these circumstances duly considered, it is rather surprising that salivation has not long since been universally exploded; the only assignable reason must be the want of a proper

proper fuccedaneum; that, however, very happily for mankind, can no longer be pleaded in excuse, as the alterative method is daily gaining ground from the following very cogent reasons, viz. the operation of proper alteratives being so extremely mild and gentle as to admit of being safely administered at any time or season, to persons of all ages, and of each sex: to pregnant women almost the whole time of gestation, even to the last month, from which happy circumstance, the sectus and its mother both receive a cure at the same time.

Two other very material advantages are to be derived from the use of alteratives, first, the certainty of cure, even in the worst, the most inveterate cases, where many repeated salivations have proved inessectual; secondly, not the least confinement being requisite throughout the whole course; which to the mercantile part of the world is a circumstance of no little importance: the patient being able to transact business as well in the open air as elsewhere, and there being no other restriction or alteration

teration respecting his diet, than the omission of acids, he will not be under any apprehension of a discovery taking place in the family.

ANOTHER particular in favour of these medicines, is, that contrary to the effects of most others, they will increase the appetite, and thereby accelerate the cure, during the performance of which, the patient's food cannot be of too nutritious a nature, insomuch that meat may be allowed twice or thrice per day, in preference to gruels, ptysans, or slops of any kind.

The only disadvantage that attends the alterative method, is this; patients being at large the whole time, and capable of purfuing either business or pleasure, some are apt to prefer the latter, and very frequently not only neglect taking the medicine, but commit various excesses, and sometimes contract fresh infections, while we are endeavouring to remove the old one. Indeed, some have been frank enough to confess, that a species of economy had given rise to these irregularities; from a supposition that

that no additional expence would be incurred, they thought it therefore the properest time to gratify such inclinations: not considering that violent exertions of the parts affected must certainly procrastinate the cure, should they even be fortunate enough to escape an increase of the venereal virus.

SALIVATION has none of these obstacles to combat with: for when once a man has completely immerfed himfelf into that delightful state, it will seldom, if ever happen, that he will, fo circumstanced, be possessed of either ability or inclination to deviate from the most rigid injunctions that may be imposed by his doctors, medicinal or divine. In support of this opinion, nothing more will be necessary than a recital of some few of the concomitant symptoms or constant attendants on a perfect falivation: namely, a tumefaction of the head, face, and throat, to a great degree, with much pain and inflammation therein; thus the tongue frequently becomes too large for the mouth, which by this time is excoriated

excoriated and extremely tender, together with the gums, tonfils, and all the neighbouring parts; in fine, the whole mouth and its appendages become one foul ulcer, from which iffues a stench that may be better conceived than described, accompanied with a ptyalism or spitting, in quantity from two to six pints in the twenty-four hours; which agreeable process is to be continued some weeks, and in the course of a few months, perhaps, repeated again and again.

The same advantages as the foregoing, has the old method of curing claps by mercurials and drastic purges, which were generally repeated every two or three days for a considerable length of time, till the discharge put on a more benign aspect; or the patience, or strength of the patient being exhausted, occasioned a discontinuance thereof.

FROM these premises we must readily infer that a person thus constantly harassed by strong rough medicines, could have little relish or inclination to transgress the laws of sobriety.

Con-

Consideratis considerandis, every impartial man most certainly will prefer the alterative mode of cure, to that by salivation.

HAVING premised thus much, now let us turn our thoughts to the most eligible means of obtaining a radical cure for this prevalent disease, it having often with great truth been observed, that as nothing can be done with greater facility, than to cavil or find fault, therefore no merit will be derived from fuch a procedure, unless at the same time it shall be accompanied with fome plan or scheme more likely to effect the purpose in question. To obviate this reflection, I shall not only propose such a remedy as will be most likely to answer, but shall also produce, in support of it, some cases which (from the peculiar fuccess attending them) will go near to call in question the veracity of the man who records them: be that as it may, fuch an idea shall not prevent my giving a faith. ful relation of them. Before, however, we enter upon that part of the business, I shall make fome observations on the means by which those events took place.

C 2

VERY

VERY numerous and different are the prescriptions which at various periods have been recommended, as fully competent to fubdue the venereal poison; amongst the great variety offered for this falutary purpose; many were diametrically opposite to each other, advocates however, were not wanting to extol their manifold virtues: on these occasions, both the vegetable and mineral kingdoms were thoroughly ranfacked and attentively examined, and many of their properties afcertained by a feries of ingenious experiments: and even at the prefent period, we are fometimes under the neceffity of having recourse to both, in the reduction of fo powerful an opponent as the venereal disease. Be it remembered however, that vegetables alone, when properly prepared, will effect a cure in many cases, although in others it must be acknowledged that minerals will likewise be required. The gums will fome of them be found of fingular use, particularly camphor, in every stage of the disease, from a gonorrhoea to a confirmed lues; in the former it will greatly contribute to alleviate the heat of

of urine, as well as correct the virulence; and in the latter it ought never to be omitted, for feveral reasons; in the first place, it is a most powerful alterative; in the next it strengthens the nerves, exhilerates the spirits, and, when joined with minerals, fortifies the stomach and bowels against many of their disagreeable or pernicious effects. It diffuses a general warmth throughout the body, and greatly assists in conducting other medicines towards the cuticular and renal glands.

As to the choice of alterative medicines best adapted to the cure of the venereal disease, I believe none will be sound more essignation actions than a composition of the following viz. camphor, antimony, and sulphur: in some cases it may be necessary to add both mercury and opium. These must be assisted with decoctions of the woods in quantities sufficient to attenuate and saturate the whole mass of contaminated juices. Any particular recipe or prescription of the above, would

^{*} Guaiacum, farfaparilla, China fassafras, fantalum, glycyrrhiza, &c.

be of very little use, as not only the doses, but the quantity of each ingredient must fo frequently be either increased or diminished agreeable to the variation of fymptoms and circumstances; which sometimes require even the omiffion of one or more of those ingredients. The age, fex, temperament or constitution of the patient, as well as the degree of infection, must render it indifpenfably necessary to make the alterations above mentioned. Another material circumstance frequently occurs, namely, the addition of other acrimonious humours; to which particular attention must be paid, or we shall not succeed in our endeavours to remove the venereal one. From hence it must evidently appear, that no particular recipe can be invariably relied upon, in this or any other difeafe.

THE symptoms of a virulent gonorrhoea (commonly termed a clap) are too well known to require any description in this place: let us therefore consider upon the means most eligible for its removal. The cure may be easily and speedily effected, if under-

undertaken when the first symptoms are perceived; particularly that gentle irritation, or trickling fensation in the urinary passage, which so frequently precedes the discharge: the disease when taken at this stage, may be very fafely removed in the space of two days, by applying medicine immediately to the part affected; this cannot be conveniently done in any other form than that of an injection, which should be composed of some gentle detergent that will stop the further progress of the venereal virus, by preventing its abforption. Chancres may likewise be almost as speedily removed, provided the remedy be applied at their very first appearance.

Should this precaution be omitted, the acrid matter will corrode the parts, and produce ulcerations; confequently a longer time will be required to effect a cure: and it may be necessary to administer internals to prevent a flow of humours to the part which might retard the cure, if not entirely frustrate our intentions.

If the complaint be not removed in the early stages, it will soon become confirmed: and then due regard must be paid to the patient's habit of body, mode of living, &c.

IT is with me a matter of furprise, that injections have not been in more general practice; as nothing certainly can be more rational, than to apply the remedy to the part affected. The advantages or the reverse, attending the use of them, must evidently arise from the ingredients of which they are composed: no man sure can be so prejudiced as to declare that his objection is merely to that particular mode of conveying a medicine: to speak impartially; if crude, corrosive, indigested ulcers in every other part of the body require topical applications, why, in the name of common fense, should they be neglected or omitted in parts naturally fo very delicate and fenfible, as the organs of generation are incontestably allowed to be? Neglects of this kind are frequently productive of the most disagreeable confequences; they may, independent of the injuries from absorption, give rise to gleets and and obstructions in the urethra, which if not attended to, may terminate fatally.

THE stillicidium, or running, is very uncertain as to the time of its appearance; sometimes it will happen in six hours after the embrace, in others twice as many weeks have elapsed before it has been perceived: these, I must acknowledge, are not very common cases, but such as have really fallen under my inspection, as will be seen in the sequel: in general however, it appears in less than ten days. Chancres likewise vary as much in this particular.

In the treatment of a confirmed clap, a great deal will depend upon its being simple or compound; in other words, whether it originated from a person who had only a gonorrhæa, or who at the same time was poxed likewise. In the latter case, powerful internals will be absolutely necessary; in the former, a cure may frequently be accomplished by injection only.

CLAPS and chancres are often contracted at the same time, but do not always make their first appearance together; I have seen instances even where some weeks have intervened.

As the composition of injection, like that of every other remedy, must vary according to circumstances, I shall decline giving any particular recipe; one general ingredient however I will venture to recommend, namely, Armenian bole, which from its absorbent and other good qualities will prove highly beneficial both in correcting the venereal acrimony, and defending the found parts of the urethra therefrom.

In the cure of claps, some practitioners have attended too much to the colour of the discharge, others to the consistence of it; both are fallacious, particularly the former; as it will sometimes be of a deep yellow, even to the last day, especially in bilious habits: frequently it is equally high coloured in old gleets of seven or ten years standing, where the party has been entirely free

free from infection all that time, as in some cases has been sufficiently manifested by the healthy state of wives and children. A discharge of similar colour and consistence is often to be found attending the fluor albus, as it is called. From hence will follow the indispensable necessity of examining surther, than just the state of the running: the best criterion will be the aspect of the urethral orisice, and the sensations throughout that tube.

Many families have been rendered unhappy from the length of time this disease fometimes requires to make its first appearance: for a man, after performing a kind of quarantine for three weeks or a month, and no fymptom of infection having shewn itself during that time, would naturally conclude himself entirely free from it, and proceed accordingly; but how great his aftonishment, as well as chagrin, to discover (perhaps a month or two afterwards) the most indisputable signs of his being infected: several instances of this kind have fallen under my inspection, some of which, D 2 I must

I must confess I should not have credited, had not they been accompanied with proofs not to be contested. From this we may infer, that although the virus very often makes the most rapid progress, at other times its operations are as sluggish: this may arise from the degree of infection, or the patient's habit of body, or from both.

CASE I.

A MERCHANT in the city of London, four years fince, applied to me in confequence of great heat in making urine, painful erections, and a discharge of matter from the urethra, so acrid, as to threaten a phymofis, it having already excoriated the greatest part of the glaus penis. Thus fituated, he defired to know if I did not call his diforder a clap; to which I replied in the affirmative, and that it was not of the mildest kind: " Then fir, said he, I shall mention a circumstance, that I think will furprife you, which is, that not a fingle symptom of the disease appeared until two months and three days after it was contracted:" upon this account he deemed it

it not venereal; I affured him however that it was fo, and could be removed only by treating it as fuch. My patient was not of a bad habit of body, nor had his blood been previously heated by intemperance, the chordee was notwithstanding so troublesome, that I expected to have been obliged to use opiates, which I am very feldom under the necessity of doing; injections generally answering the purpose, by speedily removing the inflammation and its confequences: my only objection to opium in these cases, is, the constipation which mostly attends the use of it; although I am not an advocate for the old method by catharties, yet I would not be understood to recommend absolutely the reverse; which must indubitably tend to increase, rather than alleviate fymptoms arising from inflammation. It will be found fully fufficient, that the intestinal evacuations be the fame as at the time of perfect health .- Now to return to my patient; he was much alarmed at the high degree of pain and inflammation, which however was confiderably leffened in the space of two days by means

means of injection and internal alteratives; three weeks continuance of which perfectly cured him, without any confinement or alteration in his usual mode of living, except abstinence from acids and spirituous liquors.

CASE II.

A GENTLEMAN of Dorsetshire, in the month of February, 1778, came to town on purpose to be under my care; his symptoms at that time were a discharge of yellow matter from the urethra, chancres on the glans penis, a large bubo in the left groin, and a fmaller one in the right. Upon my enquiring into particulars, he folemnly declared that he had not the least reason to think himself infected, till upwards of nine weeks after coition, when he was attacked with a trickling fenfation in the urethra, which was followed by a running and heat of urine; chancres foon afterwards appeared, and were fucceeded in a short time by tumefaction of the inguinal glands, which thereupon became extremely painful. He was at first attended by a furgeon in his own neighbourhood, who bled him very freely, gave strong mercurial purges repeatedly, and ordered

dered him to dilute plentifully with barley water, &c. Fomentations and cataplasms were applied to the buboes till suppuration took place, when he proposed to discharge their contents by means of caustic or incision, but the patient not confenting to either, immediately left the country, and foon after applied to me, requesting that I would remove them without opening or fuffering them to break, as he had the greatest aversion to fores of every kind, and much dreaded all chirurgical operations: I told him, that confidering the very forward state in which the buboes at that time were, (the fluctuation of matter being readily felt upon the flightest pressure) I could make no other promise, than to exert my best abilities in endeavouring to effect what he fo ardently defired; observing likewife, that I frequently had been fortunate enough to render those painful operations unnecessary by the use of internal medicines alone; which, in my opinion, are the only ones that can fafely be relied upon.

As to the repelling of buboes (in the vulgar phrase, putting them back) I never make

make the least hefitation of attempting to effect it, for this obvious reason, that a bubo is not a critical abfcefs, of which I am well convinced from daily experience, notwithstanding the general received opinion to the contrary; I have feen many instances where buboes have been carefully brought forward to the most perfect suppuration, liberally opened, and a copious discharge very industriously encouraged for the space of four, five, or fix months; notwithstanding which the parties have remained completely poxed; as the nocturnal pains, foul ulcers, and eruptions fufficiently manifested a few months afterwards, when they became my patients, and were radically cured by the alterative method in half that time. This being a fact not to be controverted, why should suppuration be encouraged, as it must be attended with great pain, filthiness and inconvenience? And after all, not being critical, a regular course of medicines will be indifpenfably necessary: the cicatrix or fcar is likewise a circumstance not the most agreeable, especially to the fair fex .-Without apologizing for this digression, I will

will hasten to my Dorsetshire friend. Entertaining these sentiments, I immediately fat about the discussion of his buboes; for which purpose alteratives were administered internally, and detergent injections and dreffings applied to the urethra and the chancres on the penis: by these means the fymptoms gradually abated, and at the end of a month he thought himself cured; but I advised a continuance of the medicines three or four weeks longer, on account of the high degree of virulence we had had to encounter with: he readily consented to persevere in the use of them, having already experienced fuch falutary effects; and to Thew that he did not think the decoction at all naufeous, he generally drank it at meals in lieu of any other liquor. Thus, in less than two months, without being confined to the house a single day during the whole course, he was perfectly re-established in health, and has remained well ever fince. He was highly pleased at the removal of the buboes without opening, and very generoufly rewarded my fervices.

CASE III.

In the month of July, 1777, a Gentleman requested my advice upon the following fymptoms; an unufual (though not painful) fensation in the urethra, a discharge therefrom, very fmall in quantity, and of colour fo light as fcarce to tinge his linen. I was of opinion that these symptoms were the harbingers or fore-runners of a clap; but he strongly infisted that his disorder could be nothing more than weakness, or a strain, as he had had no venereal connection for eleven weeks previous to this indisposition. I informed him, that should the cure of his difease be undertaken in this early stage, I believed it might be removed in less than a week, if he would strictly adhere to my instructions, but as he seemed so firmly perfuaded that his diforder was not infectious, I advised him to wait only a short time, and he would then be perfectly fenfible of the real nature and tendency of it. Three days only had elapsed, before he paid me a fecond visit, when I found my prognostick sufficiently verified; the urethra and

and glans penis being highly inflamed, great heat in making water, some pain in erection, and a discharge of yellow matter issuing from the urinary canal: he was by this time fully convinced of his error, and requested that I would relieve him as expeditiously as possible.—I told him that his delay of three days would procrastinate the cure more than a week, and in the present state of the disease I judged that it would at least require a fortnight, should even his temperance and regularity not be called in question. However, by the use of injection and a few alterative pills, he was perfectly cured in thirteen days.

CASE IV.

A GENTLEMAN in Westminster sent for me two years since, to give an opinion upon a stillicidium, or discharge from the urethra, which had first made its appearance two days before I saw him; he was quite at a loss to account for it, not having copulated for upwards of two months prior to that period. I enquired whether he felt any irritation or E 2

trickling in the urinary passage, he answered in the affirmative, upon which I pronounced the case to be venereal, although the running was perfectly clear, and the parts entirely free from any external inflammation. My patient, however, feemed to think that I had formed an erroneous opinion; I therefore defired him to defer using medicine of any kind for a day or two, in which time I made no doubt of an alteration in his fentiments. The next morning I was again fent for, and informed, that the parts did not look quite fo cool as on the preceding day, and that his urine felt rather warmer than usual, these symptoms he attributed to some things he had eat or drank the day before.

I REPLIED, that in twenty-four hours I was fully perfuaded he would affign another cause. Before breakfast the next day, he sent for me again, and said that he no longer entertained the least doubt of the disease being infectious, having lately perceived some symptoms of an approaching chordee; the running was now become yellow,

yellow, the parts were inflamed externally as well as within, and he could not make urine without some degree of pain. I prepared for him a cooling injection, of which he used a syringeful night and morning, and as he was very subject to the gout, I administered no other medicines internally, than decoctions of the woods and camphor, by which means, in the space of seventeen days, the cure was very easily completed.

CASE V.

A GENTLEMAN in the county of Kent, put himself under my care on the ninth day of December, 1779; he had enjoyed a very good state of health for some years, till within three days of his application to me, when he discovered a yellow running from the urethra, attended with a heat of urine, and an inflammation of the glans penis and præputium. There was likewise a small chancre on the frænum. After ocular demonstration, I assured him that his disease was venereal, my patient was not of a different opinion (having been frequently

quently infected before) but he expressed great astonishment at its being so; from the circumstance of not having had any venereal intercourse since the fifth day of the preceding September, and having been entirely free of any symptoms of the disease till the fixth day of December; an interval of thirteen weeks and upwards from the time of contracting the disease, to that of its appearance. Upon my interrogating him very closely as to the commencement of this affair, his answer was, that he could have no interest in deceiving me, and that it was impossible he should have been mistaken, having a memorandum in his pocket-book of some particular occurrences of that day, relative to this business, which with him reduced the matter to an absolute certainty. I observed, that probably sometime subsequent to that period he might have been too free in his libations to Bacchus, and be thereby deprived of recollecting his last engagement with Venus. fupposition he obviated, by declaring upon his honour that he had not been inebriated for several months past, on the contrary, he had

[31]

had rather lived abstemiously than otherwise. This gentleman's health was perfectly re-established in three weeks, by the use of internal alterants and external detergents.

THE foregoing cases having been selected from many similar ones, to shew what great length of time the venereal disease sometimes requires to manifest itself; let us now advert to others diametrically the reverse in point of time.

CASE VI.

years ago had been my patient, consulted me in January last; he complained of an irritation in the urethra, and some heat in the glans penis, which appeared rather inflamed externally. He informed me, that his last connection had been only on the preceding evening, at about eleven o'clock, and antecedent to that, he had not committed an act of venery for more than five months, during which time his health had been remarkably good. At six o'clock that morning he was awakened by a sharp pain

in the yard, which foon abated, but did not entirely go off, a disagreeable heat and irritation still remaining. From the symptoms coming on fo early as within feven hours after the embrace, he could not believe his complaint to be venereal, but fupposed that it arose from friction, and the bad wine he had that night drank; I therefore declined to prescribe any medicines, but advised that he should live after his usual manner, and think no more of the difease till it might become more worthy of his attention: which happened two nights afterwards, by the pain he experienced from a chordee; there was likewife a yellow running from the urethra, accompanied with a heat of urine; his doubts by this time being perfectly removed, he immediately entered upon the use of injection and pills, by which he obtained a cure in less than three weeks.

C A S E VII.

A NAVAL officer in the month of March, 1778, was attacked with an unufual warmth in the urinary passage, in less than six hours after coition, and it gradually increased for three

on, attended with a smarting of urine: in this stage of the disease, he placed himself under my care, informing me that he was lately returned from a long voyage, and for several months past had not had an opportunity of getting the insection, he could therefore be certain as to the date of his disorder, having been entirely free from venereal complaints for two years before. In effecting this Gentleman's cure, near a month was required, by reason of the high degree of scurvy with which he was at that time insessed.

C A S E VIII.

A GENTLEMAN in the county of Essex, about six months since, paid me a visit on the following account; having taken a night's lodging with a lady of his acquaintance, the very next morning at breakfast he selt some uneasiness in the urethra, which in the course of a few hours increased, and at the end of sive days terminated in the symptoms of both clap and

pox,

pox, viz. a discharge of virulent matter from the urethra, with heat of urine, &c. and three chancres externally; two on the præputium, and one on the glans penis. He was not much surprised at the early appearance of the disease, as with him it generally happened so; he neglected however to use medicine of any kind, till it had arrived at the state above-mentioned, when he became my patient; and, without surther delay, began upon a course of alteratives and detergents, which effected his cure in less than a month.

To the peculiarities of constitution, as well as degrees of infection, must be attributed the different periods in which particular diseases make their appearance. This circumstance may depend on the disproportion of one or more of the ingredients or materials which compose the human body. Some men are naturally so irritable and instantly to take fire like tinder or touchwood, from the smallest spark; while others of cold phlegmatic habits, are (as is to be expected) quite the reverse.

C A S E IX.

Four years fince a Gentleman in the county of Northampton, made a journey to town, to confult me upon what he called a weakness of a particular kind, as it was accompanied with great inflammation; I immediately suspected what the case was, and upon examination, found it to be a most virulent clap, with an incipient phymofis; he complained likewise of a chordee. I was much furprifed when he told me that his apothecary had not deemed it venereal, but treated it as arifing merely from relaxation, giving large doses of Peruvian bark and elixir of vitriol. My patient faid that the only reason why his disorder had not been judged infectious was, that it did not discover itself till more than fix weeks after coition; from which circumstance the apothecary had declared, that it could not poffibly be venereal. By the use of deterfive injections and alterative pills, he returned home in perfect health at the expiration of about three weeks.

F 2

CASE

CASE X.

In the month of December, 1778, a tradefman in Westminster commenced my patient, in consequence of a virulent gonorrhæa, which he had contracted more than a month before his application to me, and of which he gave me the following account; that in fix hours after copulation, he perceived an unufual fensation in the yard, a kind of trickling or itching in the urinary canal, which for upwards of three weeks had been a little troublesome, but by no means painful, until the 26th day, when a clear colourless discharge issued from the urethra, which now began to be much inflamed, and felt very fore whenever the urine paffed through it; on the 28th day, the running was quite yellow and rather copious; and upon the twenty-ninth he was attacked with a chordee, which induced him to be under my care the next morning; he faid he should certainly have applied fooner, but the inconvenience he experienced previous to the 26th day was fo trifling,

triffing, that he had not entertained the least idea of his complaint being venereal; it was however entirely removed in fifteen days, by the same means as were used in some of the preceding cases.

C A S E XI.

A GENTLEMAN in the county of Surry, fent for me in the month of August, 1778. I found him labouring under a confirmed pox, of which the following were his fymptoms, a violent head-ach, nocturnal pains in the shin bones, with nodes thereon, a large foul ulcer in his throat, upon the left tonfil, a heat in the urethra, and an ob-Aruction therein: upon my requesting to know every particular relative to the rife and progress of the disease; he informed me, that it was contracted about ten months before, and one thing appeared very ftrange to him, namely, his knowing himfelf to be infected in less than twelve hours after the connection, from a very troublesome irritation in the urinary paffage, which feized him about that time, and for fome days encreased, notwithstanding his immediate

application to a furgeon, who ordered almond emulfions, with nitre, manna and falts; and to dilute freely with barley water, in which gum arabic had been diffolved, partial warm baths and venefection were not omitted; in the course of a week or tendays, mercurial ointment was used very liberally for at least a month, with brisk purges at intervals. The running, he faid, was very plentiful for the space of five or fix weeks: he then took balfamics for three weeks, and was thought to be cured; however, in about feven or eight weeks afterwards, he perceived his health to be gradually on the decline, a diminution of appetite, strength and spirits, and sleep interrupted by wandering pains in his head and limbs :- nodes foon after formed on the shins, and an ulcer in the throat. On the appearance of these symptoms, another furgeon was employed, who administered large quantities of the sublimate solution, for a confiderable time, when finding that the disease did not give way, he put himfelf under my care, and was perfectly cured in two months, by the alteratives recommended

[39]

mended in the 13th page, and a few bougies to remove the urethral obstructions.

BE it observed, that no injections had been used in this gentleman's case; to which circumstance, in my opinion, may be attributed the want of success, for had a proper detergent one been applied in the beginning, the venereal matter would not have been absorbed, and consequently the pox, with its disagreeable attendants, been thereby prevented.

C A S E XII.

A GENTLEMAN in Westminster eighteen months since, consulted me in consequence of a boil (as he termed it) seated amongst the hair on the pubes, very near the penis. This ulcer had resisted every application his surgeon had made to it for three or four weeks. From the hard edges and other appearances, I pronounced it to be a chancre; this my patient did not at that time credit, however, not long after he altered his opinion upon the appearance of a bubo in the left groin, which in a few days became very painful, when he paid me a second visit,

visit, and requested to have it repelled: that was performed, and a cure effected by an alterative course in about thirty days.

C A S E XIII.

In the month of November, 1778, a Gentleman, in the county of Middlefex labouring under a confirmed lues venerea, applied to me for its cure; the fymptoms at that time were, a fmall node on the right Thin bone, an ulcer in the nofe, nocturnal pains, and a copper coloured eruption overfpreading a great part of his body. He informed me, that, till within the last twelve months, he had never experienced a fingle fymptom of the venereal difease, and that his present afflictions, if they were venereal, must arise from a clap contracted a year before, which denoted a high degree of malignity, the inflammation being very confiderable and of long duration, attended with heat of urine, chordee, &c. the running he faid was copious, and continued upwards of five weeks, in which time he had been purged every three or four days, had rubbed in a large quantity of mercurial ointment.

ointment, afterwards taken electaries and pills in abundance: in consequence of this process the symptoms had disappeared, and at the end of two months his cure was pronounced. Some few months after, however, he perceived his health to be gradually on the decline, without being able to affign the real cause, not having the least suspicion of venereal matter; his appetite and strength daily decreased, his fleeps were short, interrupted, and afforded little or no refreshment, and at length colliquative fweats enfued: thus circumstanced, he consulted a physician, who, judging his cafe to be a confumption, put him under a course of ass's milk, &c. &c. for a confiderable time, but growing every day worse and worse, he was ordered a change of air, and removed into another county. Finding however no greater relief from the alteration of air, than he had before experienced from medicine; he began to despair, and most earnestly wished for his diffolution. Fortunately, an old acquaintance from town called to fee him, and upon hearing fome of his fymptoms ftrongly

strongly urged him to become my patient, he accordingly did so, and was perfectly cured in the space of three months, merely by the use of alteratives.

C A S E XIV.

A GIRL on the town applied to me upon the tenth of November, 1778, and faid she supposed it would be necessary for her to undergo a thorough repair, not having been free from infection for a confiderable time: her fymptoms then were, an ulcer in the throat, a foul eruption on the abdomen, a brace of small buboes, and a great difcharge of acrid matter from the pudendum. As the was extremely anxious to have the buboes dispersed, I told her that no time should be lost, and consequently advised the immediate use of medicine; she replied, that it must be deferred till next day, as her mistress expected much company that evening, and had engaged her to one gentleman in particular, with whom fhe was to fleep that night, his name having been entered in the book for that purpose some days fince; and for which he had paid the ufual retaining

retaining fee .- I was not furprifed at this account, fimilar ones having occurred before, and particularly this instance; a friend of mine, an Officer in the army, made application to a lady abbefs to know on what night a particular nymph would be difengaged; after inspecting her ledger, she informed him that the lady was disposed of for twenty-two nights, and if he pleased she would take down his name for the twentythird, upon paying the usual compliment .--But to return to my patient; she called upon me again the next day, and without any further delay entered upon a course of alteratives, by means of which her health was perfectly re-established in less than seven weeks. She was under great apprehensions lest the buboes should suppurate and occasion fcars, but was foon made easy on that head, as they dispersed very kindly in a few days, the other symptoms gradually disappeared, and in little more than three weeks she thought herfelf well, it was therefore with fome difficulty that she could be perfuaded to continue the medicines a short time longer, in order to confirm the cure.

G 2

CASE

C A S E XV.

A MECHANIC in Westminster, being employed in the house of a nobleman, became acquainted with the lady's woman, and after many entreaties, perfuaded her to grant him the last favour: in the space of a fortnight however, he was fully convinced of the impropriety of the connection, being at that time obliged to commence my patient, on account of a large chancre feated on the anterior part of the scrotum; it was the most malign that ever came under my inspection, being in fize equal to a shilling, and had the appearance of an efchar, produced by fome potential cautery; the edges were hard and thick, and the scrotum around it much inflamed. The urethra, and every other part of the penis had entirely escaped the infection; from which circumstance he did not suspect the nature of his complaint, till the inguinal glands enlarged and became painful. Proper externals and a course of anti-venereals effected a cure in thirty-two days.

C A S E XVI.

ABOUT two years ago, a Lady applied to me in consequence of a sore throat, which had for fome weeks been extremely troublesome, notwithstanding the variety of gargarisms, &c. that had been used in expectation of its removal. Upon inspecting her throat, I pronounced the ulceration there to be venereal, and was confirmed in that opinion, when she informed me, that her rest of late had been greatly disturbed by nocturnal head-achs; she however was at the same time entirely at a loss to account for these symptoms, the pudendum having never been diseased. The infection, I said, had in this case most probably been conveyed by the tongue, and therefore I defired to know whether she had kissed columbatim; to which she answered in the affirmative, and jocosely observed, that such confequences might be added to the many other cogent reasons for keeping that member within its proper limits. She then entered upon a course of alterants, and was cured in about five weeks.

The infection may be communicated likewise by contaminated semen, without injuring the pudendum in its passage to the uterus, from whence its poisonous particles may be absorbed, and by means of the lymphatics conveyed to the blood, and thereby to every part of the body.

C A S E XVII.

A GENTLEMAN in the vicinity of London, about five years fince, confulted me upon the ill-state of health which himself and lady at that time laboured under. He informed me of his having led a very irregular life for fome years previous to matrimony, that he had been frequently difeased, and recollected having had chancres a few weeks before his marriage, which being removed by mercurial ointment and cathartics, at length he was supposed to be cured, and foon after entered into the holy state: in about five months his wife was thought to be pregnant from the fuppression of the menstrual evacuation, and the family felicitated themselves thereupon; but the other concomitant fymptoms not attending

attending in proper time, the parties were greatly difappointed, and particularly fo, to find that in lieu thereof the lady's health as well as that of her hufband, was evidently on the decline. A flow fever, want of appetite, strength and spirits soon after succeeded, together with that of fleep in the night, being generally drowfy in the day time. Medicines of various kinds had been administered without producing the defired effect; the gentleman had an eruption upon his breaft and other parts of the body, which was encouraged by the faculty, in expectation of its being critical; they were however, greatly mistaken, as none of the other fymptoms disappeared in consequence of it. The lady still continued in a very languid state, notwithstanding the number and variety of medical prescriptions. In this unhappy fituation they remained for fome weeks, when an old bottle companion paid the gentleman a visit, and observing some of the eruption on his forehead, faid, "I'll be d-m--d if you've not got a corona veneris. your case has been most egregiously mistaken; pray lose no more time, and give

me leave to fend my furgeon to you;" he acquiesced, and I waited upon him the next morning, when I found that his disease was most evidently venereal from many other symptoms, as well as the eruption, which was of the true copper colour.

Upon examining the lady, her skin in fome places appeared rather discoloured, (as if nature was labouring to throw off part of the noxious humour that way, but wanted strength to effect it) and she had likewise every symptom of a pocky hectick. The difease had been communicated to her entirely by the contaminated femen of the man, without its having done the least injury to the vagina, or any other part of the pudendum. After having made the neceffary enquiries, I recommended a course of alteratives to both man and wife, by the ufe of which, in less than two months they were perfectly cured, as appears at this time by the healthiness of their offspring. The lady became pregnant about two months after leaving off the medicines, and has borne several children since.

plan pursued by some of our modern debauchees, prior to entering into the connubial state, all the foregoing disagreeables would have been prevented. This prudent, plan or measure, is nothing more than going through a short course of alteratives: which in many instances must be highly necessary, particularly after a series of irregularities, and a variety of insections, which through neglect or improper treatment, may be productive of great mischief in suture.

THE inefficacy of a first or second salivation occurring so very frequently, I do not think those cases worthy of insertion here, I shall therefore immediately proceed to the relation of such as have withstood three, four or sive repetitions of that foul process.

C A S E XVIII.

In the year 1775, a Gentleman in the city of London put himself under my care, on account of a confirmed pox, with which he had been infected some years, notwith-

H

Standing

standing three salivations, and long courses of the folution of corrofive fublimate and farfaparilla decoctions .- He gave the following particulars of his case; that he contracted the infection in the year 1769, which at that time manifested itself in a clap and chancres; his furgeon administered medicines for eight or nine weeks, and then difmissed him as cured. In the course of a few months a foul eruption overspread his body, and ulcers formed in his throat; he again applied to the furgeon, who advised falivation, and foon after put him in a course of spitting for some weeks, and then pronounced his cure to be radical. In less than fix months the difease re-appeared, but with much greater malignity than before: the ulcer in the throat broke out afresh, with the addition of two in the nose, which foon fwelled to fuch a degree, as to obstruct, in some measure, the fight of the left eye, and he was deprived of rest by nocturnal pains. Thus circumstanced, he employed another furgeon, who also salivated him; the spitting he said was continued above three weeks longer than in the preceding falivation, after which he was affured by his medical

medical friend, that the disease was perfectly removed. Bufiness some months afterwards required his taking a journey into the North, where he was alarmed with the return of fome of his former symptoms, nocturnal pains, &c. During his residence in the country, he took a great deal of mercury in various forms and preparations for feveral weeks; which proving to be merely paliative, he returned to town and confulted another furgeon, who recommended a third falivation; this he did not very readily agree to, having experienced its inefficacy twice before. However, being informed that it was the dernier resource, and at the same time entertaining a very high opinion of the abilities and integrity of his furgeon, he at length confented to be falivated a third time; when a copious spitting was continued for a confiderable time, as long as it could with any degree of either fafety or propriety be purfued.

HE was attended likewise by a physician, who, with the surgeon, declared that the disease was entirely vanquished, and that nothing more was necessary than pure air and

nutritious diet to recruit his strength, and finally complete the cure. Indeed they were, he faid, quite positive as to the success of this last falivation, the fymptoms having given way fo readily, and the process in every respect been so satisfactorily conducted throughout. The patient acquiesced in the prognostic, and without further delay removed into the country for the benefit of air, &c. This pleasing delusion lasted not many months, for he had the mortification to perceive that some of the symptoms of his old diforder were gradually coming forward again; he therefore confulted his phyfician once more, by whose direction he took fublimate folution and farfaparilla for some months; these medicines afforded him only a temporary relief, and were not attended with better fuccess than the three falivations which had preceded them, as evidently appeared not many months after by a return of the difease. His patience as well as constitution was by this time nearly exhausted; infomuch that he frequently declared, no one should ever persuade him to make any further trial of medicine, it having fo repeatedly

peatedly proved fallacious. The difeafe however becoming almost insupportable, he was prevailed upon to make one effort more for its removal: in consequence of which he became my patient at the time above-mentioned, when he had feveral foul ulcers on the back, two in his nofe, one on the cheek, and another behind the right ear; his head was almost covered with scabs, as was likewife the fcrotum; from all these issued such naufeous effluvia, that his nearest relations and friends could not remain in the fame room with him long at a time. Pains in his head and limbs prevented him from fleeping, except towards morning or in the day. His appetite, strength and spirits were so much reduced, that he despaired of relief from any thing but a total diffolution. After having attentively given ear to this melancholy detail of repeated disappointments, and examined the ulcers, &c. and made the proper enquiries relative to his former habit of body, or natural temperament previous to the difeafe, I declared as my opinion, that he might be cured by alteratives, if of the most powerful kind, and regularly persevered in for some weeks. He affented, though

though not without first observing, that by this time he had entirely loft all faith in medicine. I prepared for him strong decoctions of the woods, and likewife alterative pills, and defired him to take them in fuch dofes, and as repeatedly as the strength of his stomach would admit of. Relying folely on the effects of internals, I applied no other dreffing to the ulcers than dry lint throughout the whole time. I recommended diet of the most nutritious kind, and that it should be taken in small quantities, frequently repeated, by reason of the very weak state to which the disease and reiterated salivations had reduced him. In little better than a week, the nocturnal pains entirely left him, the scabs foon after fell off, and the ulcers were in good condition; his appetite and spirits daily increased, insomuch that he thought himself well in less than a month, the ulcers being all healed, and every fymptom of the difease having by that time disappeared .- Notwithstanding which I judged it necessary to continue the medicines a month longer, to prevent a relipfe. My patient had no objection to persevere in the use of them for any length of time, having

having already experienced fuch falutary. effects; which, with great pleasure, he contrafted to those of the methods he had formerly purfued; instead of being confined to his chamber, and undergoing one of the greatest punishments (as he expressed it) in the power of the faculty to inflict, namely, that naufeous process a falivation, he had enjoyed moderate exercise in the open air every day the whole time he was under cure, and experienced an increase of appetite, strength and spirits in proportion to the time of using the medicines; and as a proof of their not being unpleasant, he constantly drank the apozem at meals, and at other times when thirsty. Thus was he radically cured in the space of two months, having never experienced any return of the complaint, as he not long fince informed me.

C A S E XIX.

THREE years fince, a Merchant in Westminster employed me to extricate him from the following disagreeables; nodes on the shins, an ulcer in the throat, nocturnal pains, and a scabby eruption on his body, which

which from its stench was extremely offenfive. He was much emaciated, and had little or no appetite. I defired to be informed of particulars, the date of the difeafe, in what manner it first appeared, and what method had been used for its removal. To which he replied, that in the year 1772 he caught the infection, which made its first appearance in a bubo on the right groin, and was not preceded by either chancre or clap, nor had he ever been infected before. The tumour increased very fast and was extremely painful, his furgeon encouraged suppuration, and when that was complete, opened it by caustic. The discharge continued upwards of three months, during which time he rubbed in large quantities of mercurial ointment, and was purged at intervals to prevent a fore mouth; and at the end of about fourteen weeks his cure was pronounced. In a few months however, the disease returned; ulcers having formed in the throat, accompanied with nocturnal pains in his head and limbs, for which he underwent a falivation, and was again supposed to have received a cure; but in less than fix months he was obliged to apply once more

more to his furgeon, the difease having reappeared with the addition of a node on the left shin-bone; he was falivated a fecond time, and was faid to be perfectly cured. After some months the disease again appeared, when he employed another furgeon who perfuaded him to make trial of falivation for the third time, which affording only a temporary relief, he confulted a physician, by whose directions he took medicines for feveral months, until the difease was thought to be removed; being reduced by this process to a very languid state, country air and afs's milk was recommended, but they had not the defired effect; for foon . after the discontinuance of anti-venereals, he gradually relapfed into his former fituation. His medical friends being again confulted, they strongly urged the necessity of a fourth falivation, as the only probable means of fucceeding; by this time he entertained no very high opinion of falivation, but being almost worn out with pain and disappointment, he agreed to it with this proviso, that it should be carried to such a degree, as to terminate in the perfect reestablishment of his health, or in the grand

cure

cure of all mortal complaints. To humour him they acquiefced in the propofal, and he was falivated for the fourth time; the fpitting they affured him was continued as long, and to as great a degree as his strength poffibly could admit of: and every circumstance considered, they had not the least reason to doubt but a radical cure would be the consequence. This defirable prediction however, was by no means verified, as evidently appeared in the course of a few months, the disease returning with seemingly accumulated virulence, which he observed to be the case after each of the preceding falivations. Having finished the narrative, he defired to be informed whether I could render him any fervice, which he was induced to believe might happen, from fome accounts he had lately heard of my having effected cures after the failure of repeated falivations. I told him that I had been fuccessful in that line, and notwithstanding the obstinacy of his disease, I entertained fome hopes of relieving him. After some further conversation, he defired me to administer what I thought proper, and that he would strictly observe my directions.

directions in every respect. Accordingly, I recommended a course of alteratives; and at the same time the most nutritious diet, with moderate exercise in the open air, to be frequently repeated, proportionably to his encrease of strength. Within the first fortnight I sent him three times to a bagnio, in order both to cleanse the skin and promote perspiration. By these means he obtained a persect cure in the space of ten weeks, and has remained healthy ever since.

C A S E XX.

A GENTLEMAN in the county of Cornwall, four years ago, put himself under my care, his case was a confirmed pox, under which he had laboured near eight years, the symptoms at times having been palliated by sive salivations, and a variety of medicines. Upon my requesting to be informed of every particular relative to his complaint, he told me, that being a man of very sew words, he never dealt in long narrations, of which his case would prove one if given in detail; he should therefore mention a few of the most material circumstances, and wrap them

up in very small compass. He said, "That about eight years before, he contracted a fmall chancre on the penis near the frænum, which his furgeon healed in a few days by the use of mercurials both externally and internally. A short time after, a bubo formed in the groin; it suppurated, was opened by caustic, and, after discharging some weeks, healed; he took medicines fix or eight weeks, and was faid to be cured. In the course of a few months he was visited by an ulcer in the throat, nodes, and nocturnal pains. In fine, the inveteracy of his difeafe was fuch, that, from the time above mentioned, to that in which he was speaking, it had withstood five falivations, and great quantities of different medicines administered between each of them: and (as he expreffed it) had he not had the constitution of a horse, the disease, or the methods used for its removal, must in that length of time have destroyed him: but thanks to heaven, his stamina were naturally of the strongest kind. He was now quite emaciated and worn down with nocturnal pains in his head, limbs, &c. There were three nodes

on his ribs, and a large ulcer on the head, from whence feveral pieces of bone had exfoliated, and many more were fo loofe as to require extraction foon after he became my patient; this opening had been made by a caustic for the purpose of discharging a large collection of matter. These were his fymptoms at the time he made application to me; and notwithstanding the measures hitherto adopted had fucceeded no better than as mere palliatives, I despaired not of removing the difease; having so frequently experienced the powerful effects of alterants in the most confirmed cases, where many repeated falivations had proved abortive. This gentleman acquired a radical cure in three months by the use of alteratives: had the exfoliations been compleated fooner, I should not have given him the medicines fo long.

C A S E XXI.

In the year 1777, a Gentleman from the county of York, came up to town to be under my care. He had been afflicted with a venereal complaint almost ever since his birth,

birth, by means of an infected nurse, whose foul milk fo gradually affected him, that fome weeks elapfed before his parents were able to account for his declining state of health; happily however, the cause was discovered, and a check put to the rapid progress of the disease, by means of antivenereals. The growth however of the child was much impeded, as the medicines though continued a long time had only palliated the fymptoms, and faved the life of the patient. When he confulted me, his age did not exceed twenty years, he carried but little flesh, and was low in stature, although his parents were the reverse. Foul eruptions frequently infested many parts of his body, and fometimes afforded a temporary relief from the achs and pains which he generally at other times experienced. His eyes were extremely weak, and often fo much affected by this acrid humour, as to be threatened with a loss of fight. The infection having been thus deeply rooted in the constitution, and in a manner intervowen with it from his infancy, I entertained very slender hopes of totally

municated to my patient, who, nevertheless, entered very chearfully upon a course of my medicines, which some of his acquaintance had frequently given him a very favourable account of. In the space of three weeks he acquired a good appetite; the cutaneous eruption gradually gave way, as did likewise the pain in his limbs, &c. he daily improved in strength and spirits, and at the end of three months appeared to be in high health, and was grown fat. By my advice he continued the medicines three months longer, and was then perfectly cured.

Let this, amongst many other similar instances, serve as a caution to parents how they entrust their infant offspring to mercenaries. Other acrid humours may often be derived from the same unhappy source.

C A S E XXII.

A TRADESMAN in Westminster, about two years and a half since, became my patient, together with his wife and child, who, at that time, were all infected with the venereal disease; the man had a coppercoloured

coloured eruption on his body, and an ulcer in the throat; the woman a bubo in the right groin, and nocturnal pains; and the child a venereal eruption which overspread the greatest part of its body. This complicated scene of distress took its rife from the husband, who informed me, that nine or ten months before, he had made an imprudent connexion; twelve days after which, a fmall ulceration (a common excoriation he supposed it to be, though in reality a chancre*) made its appearance on the glans penis, near the frænum; not suspecting it to be venereal, he applied a piece of dry lint or cotton, which healed it in a few days. On examining the part, I found, that notwithstanding it had skinned over fo readily, yet there remained a fmall callofity, from which all this mischief had originated. In the course of a few months he had reason to suspect himself to be infected, and a short time after the mortification to find that he had communicated the disease to his wife, from whose milk it was conveyed to the child also. An enlargement

^{**} By the patient's account not much larger than a pin's head

of the inguinal gland was the first venereal fymptom that attacked the woman, none of the neighbouring parts having been at all affected; we may therefore conclude, that The received the infection from the contaminated femen of the man. The family furgeon and apothecary having for fome time administered medicines without the defired effect; I was thereupon called in, and found the parties in the fituation as above described. The child at this time was not more than twelve months old, confequently very little fuccess could be expected from the application of medicine immediately to the infant itself, for, let the preparation be ever fo palatable to the taste, or gentle in its effects, a sufficient quantity would not be taken to eradicate a disease of this kind. Therefore, I defired the mother to continue fuckling her child, that the remedy might be conveyed to it through the fame channel that had communicated the difease; for this purpose, I immediately put her under a course of alteratives, of which she could only take small doses on account of the infant, whereby the cure was protracted to near four months, at which period I had the fatisfaction of feeing both mother and child perfectly freed from this loathfome difease. The man's health was re-established in half the time that had been requisite for the cure of his wife and child, by reason of his taking the medicines in their full doses.

I could mention feveral cases similar to the foregoing ones, was I not of opinion, that they will elucidate the subject as well as ten times the number.

N.B. In the treatment of chancres, their callofity should be removed as expeditiously as possible, to prevent the ill effects of absorption which will otherwise ensue.

Excoriations on the glans penis and præputium of men, and on the labia pudendi, vagina, &c. of women, have very frequently been treated in the most erroneous manner; some that were truly venereal, have been totally neglected, whilst others that arose from very different causes, have been so carefully attended to, as almost to endanger

the life of the patient; by the application of strong mercurials, drastic purges, &c. Nor are instances wanting of persons being salivated on the like occasion.

PROPHYLACTICS.

HAVING treated hitherto on curatives only, let us now bestow a few thoughts upon preventives. The doctrine of prevention, although it may not be coeval with the disease, is nevertheless of considerable antiquity: for the accomplishment of so desirable a purpose, a great variety of methods have been proposed; many of them are to the highest degree absurd, and a few only that bear even the semblance of probability, particularly amongst those recommended by the ancients.

During my attendance at the lectures of medical professors several years since, I have frequently been much entertained at the ingenuity with which this subject has been handled. One of these gentlemen has strongly contended, that a mucilaginous or oleaginous application is the only topical

one that can be depended upon; and, in support of this opinion, expatiated on the power of emollient and unctuous remedies in correcting or counteracting the most acrid and corrofive fubftances, by blunting their fpicula, and sheathing the acute angles, &c. thereof. Another professor, perhaps equally learned, and probably not inferior to the former in experience, infifts, that a strong alkali is the only thing which can effect this grand desideratum, and for this reason, that in order to expel the venereal virus as expeditiously as possible, nothing will be fo likely to answer that purpose, as a medicine of this kind, which immediately stimulates the glands to such a degree, as to produce a very copious fecretion of their fluids, and thereby dislodge the enemy without either impediment or delay. With equal plaufibility, acids have been recommended by fome, and spirits by others; not confidering that the pain, &c. arifing therefrom, would frequently injure as much, and perhaps fometimes more, than the difease which they were intended to guard against.

VERY

VERY numerous have been the advocates for mercurial preparations, whilst feveral have given the preference to those of lead, Every man of experience, however, must acknowledge that theories in general (how fpecious foever they may appear) are, the major part of them, extremely defective, when put to the test. I have myfelf feen a number of various experiments made on the present subject; indeed the line of practice in which for many years I have been materially engaged, has enabled me to profecute an enquiry of this kind to as great an extent as most men: and after all, I am perfuaded that very gentle stimulants* will anfwer the purpose more effectually than applications of any other fort; I am led to be of fuch an opinion from the very great fucfess that has attended this mode of prefervation for feveral years past; those of my patients who have made the proper and timely use of a composition of this kind, have not been infected, I believe, once in fifty times upon an average; which confidering all circumstances, is as much as can

^{*} A weak folution of metallic falts.

reasonably be expected. Were I to assert that any medicine or preparation is infallible, I should pay no great compliment to the understanding of my readers, by advancing so palpable a falsehood; suffice it to observe, that some of my friends entertain the highest opinion of this composition, from the great and repeated risks they have ran for many months successively, without receiving the least injury; notwithstanding their sometimes copulating with such as they knew to be infected at the time of action.

To guard against the infection, some practitioners have thought it sufficient to advise the application of unctuous or other substances to the glans penis, in order to obstruct or close up the absorbent vessels or pores thereof, being of opinion, that the enemy always gets admission that way: in opposition to such doctrine, be it remembered, that chancres frequently six upon other parts of the penis, and sometimes attack the scrotum and its vicinities, from which buboes have ensued, as appears in some of the foregoing cases.

GLEETS.

Or gleets issuing from the urethra, there are two species upon which I propose making some observations; namely, the venereal gleet, and the seminal one. The former is the discharge which succeeds a clap; in other words, the running that continues after the removal of infection; this disease arises from the urethral ulcerations not having been healed in proper time, which may be occasioned by various means; for instance, by too frequent repetition of strong catharties; by unctuous applications long continued; by acrimonious humours; in fine, by any thing that relaxes to too great a degree, or for too long a continuance.

THE cure is often protracted by the irregularity of the patient, who sometimes is kind enough to impute it to the mismanagement of his surgeon; and on the other hand, it may so happen, as to be really the case.

Some of the reasons assigned for the difficulty of curing ulcers on the legs, will hold good good in those of the urethra; the situation of both affording a depending orifice, or drain to the humours, which consequently will very readily flow down thereto; and when Nature has been accustomed for a length of time to any particular evacuation, and the outlet thereof, the slux of humours will there be very considerable, and the task of diverting them to any other channel will be proportionably difficult.

ANOTHER difagreable circumstance attending ulcerations of the urethra, is, the frequent flows of urine over them, which at times being highly acrimonious, must interrupt the digestion of the ulcers; and a part of it lodging in them, may corrode in fome degree, unless prevented by proper topical applications, which cannot here be conveyed in a more fuitable form than that of injection. And even after the ulcers are healed it will generally be found expedient to continue injection a week or ten days longer, to confirm the parts, and defend them from the urinous falts, which (should this precaution be omitted) are fometimes

sometimes so acrid as to abrade the new skin, and make the sores discharge afresh. The same thing may happen when the urme is loaded with coarse gravel or stones, which will soon destroy the tender cicatrix; and if neglected but a few days, a dessuction of humours to the part may ensue, and must consequently retard the cure.

So high an opinion do I entertain of injections, that I will venture to affirm, that almost every gleet (unaccompanied with caruncles, or callous edges of those ulcers from whence it proceeds) may be cured by injection, provided the redundant acrimony (if there happens to be any) shall have been previously corrected. Innumerable almost are the instances which might be produced in support of this opinion: a few only of the most remarkable ones will be here introduced to elucidate that point.

GLEETS, attended with great obstructions in the urethra, occasioned by carnosities, caruncles, callosities, funguses, &c. (call them by what names you please) will generally require medicated bougies to re-

move them before we can expect a radical cure of the gleet. They will likewise be necessary in the removal of strictures, and the callous edges of old obstinate ulcers in this part.

As to the composition of bougies, great care must be had that nothing corrosive or very stimulating be introduced therein; on the contrary, it must consist of the mildest digestives; and when formed into bougies, must have a very even polished surface, and be of fuch temper or confistence as readily to adapt themselves to the curvatures, &c. of the urethra, and yet at the same time be endued with fufficient strength to make the necessary pressure. The foregoing caution relative to the composition of bougies, cannot be too much inculcated, when we confider the delicate texture of the urethra and its extreme irritability: no part of the body, I prefume, bearing greater analogy in point of fenfibility: than the urethra to the eye.

In the management of bougies, more attention is necessary than people in general imagine; for instance, if too large a one be introduced,

an hæmorrhage (that very much alarms the patient, though in other respects not prejudicial) or an inflammation ensues, which must retard the cure, should it produce no other disagreeable effect.

THE distention of these parts ought to be made very gradually for the reasons above assigned, as well as to avoid the pain that must otherwise be occasioned. If the end of the bougie be suffered to press against the neck of the bladder, it generally stimulates to such a degree, as to produce a strangury.

A LIGATURE of some kind should be always affixed to a bougie, previous to, or immediately after its introduction, both to prevent it from slipping into the bladder, and to retain it exactly in the proper situation.

How long each bougie may be worn, will depend upon the irritation it occasions; it must not remain in the urethra after it becomes painful, for reasons very obvious. When the urethra has been habituated to

L 2

them

them, they may be retained the whole night, provided fleep be not thereby prevented, or much interrupted.

ANTECEDENT to a course of bougies, it will frequently be necessary to prepare the parts with a cooling injection, and to repeat it during the use of them, as soon as the least symptom of inflammation makes its appearance. Gentle aperients at this time may likewise be serviceable, such as lenitive electuary, sulphur, magnesia, &c. Moderate exercise, and a total abstinence from acids and strong liquors must also be enjoined.

As to the length of time which may be proper to continue the use of bougies, that must depend upon the degree of obstruction in the passage, and the irritability thereof, together with the patient's habit of body, and many other circumstances.

IT will not I believe be disputed, that there may be the abuse, as well as proper use of bougies, for should they be continued long after the disappearance of discharge, a fresh fresh running will be produced, by the very means employed to remove the old one; they are not, however, to be left off abruptly; it must be done gradatim, lest the cicatrix should exceed its due bounds, and thereby lessen the diameter of the urethra.

MEN fometimes want patience under a course of bougie; but the time may be greatly shortened by the assistance of proper injections. Many instances indeed have occurred, where injection has performed a cure after bougie, and every other remedy proved inessectual. Such improvement has of late been made in this respect, that I do not find it necessary to use half the number of bougies which I formerly did, cures now being effected in so much less time.

C A S E XXIII.

In the month of March, 1777, I was applied to by a Gentleman of fortune in the county of Derby, who had laboured under a venereal gleet for more than fix years; in the course of which time he had employed several

feveral of the faculty in his own neighbour. hood, and made journies to town for the opinions and prescriptions of some eminent ones in the metropolis; in confequence whereof, he had used a variety of medicines both external and internal, which afforded him only a temporary relief; the discharge generally returning in ten days or a fortnight afterwards. The difease commenced, he faid, in the year 1770, when he contracted a clap, for the cure of which, his furgeon had purged him pretty brifkly, and used a considerable quantity of mercurial ointment, until the infection was supposed to be removed; then, balfamics, aftringents, &c. &c. were administered; and he afterwards used the cold bath for several weeks fuccessively. All these expedients however proved merely palliative, as the complaint constantly returned foon after their discontinuance. The discharge was copious and of a deep yellow colour; I defired permission to examine the urethra with a bougie, he readily affented, but observed that bougies would be of no service to him, having used them upwards of eleven months without

without any good effect; I informed him that I did not propose his wearing them, unless there should be a very material obftruction in the passage; which upon examination not being the case, I recommended alteratives and an aftringent injection, the former to be continued three weeks or a month, and the latter, double that time-with moderate exercise, and a ftrict abstinence from acids of every kind; and at the same time to keep his body solutive with fulphur or magnefia alba. By thefe means he was radically cured in the space of two months, and has remained well ever fince, as he perfonally informed me in the month of January, 1780.

C. A S E XXIV.

About two years fince a Gentleman in Westminster applied to me for the removal of a gleet, which had troubled him twelve years; during that time, he had at intervals taken a great deal of medicine, and of various kinds. One remarkable circumstance attending this case, was, that notwithstanding the very long continuance of the running, no carnosities

carnofities or obstructions of any fort had formed in the urethra, which I ascertained by the introduction of a bougie, previous to delivering any opinion upon it; when finding the passage perfectly clear, I told my patient that his complaint in all probability might be removed without much difficulty, provided he would live temperately, and adhere strictly to the rules I should prescribe; namely, to avoid acids, strong liquors, and violent exercise; and regularly to use every night and morning a syringeful of injection. These he did; and was perfectly cured in the course of three months.

C A S E XXV.

A NOBLE LORD, in the month of May, 1778, fent for me, on account of a gleet attended with a small obstruction in the urethra: he informed me that his disorder was of four years standing, and had resisted every means prescribed for its removal; that he had taken a variety of medicines, been once salivated, and had used bougies. The discharge was thin, yellow and copious, unaccompanied with either pain or inflammation.

tion. As his Lordship was of a good habit of body, I did not prescribe any internal remedies, but trusted wholly to injection, which in the space of fix weeks effected a radical cure.

N. B. THE same regimen was observed as in the preceding case, and the injection was used three times each day.

C A S E XXVI.

A STATIONER in Westminster, three years ago, applied to me, in consequence of a gleet, with a small caruncle in the urinary canal, which complaint he had laboured under upwards of five years, and during that time had used various medicines, but they had all proved ineffectual; he was however, perfectly cured in less than two months, merely by the use of injection.

C A S E XXVII.

I was confulted about two years fince by a Coal-merchant in the county of Middlefex,

who had then contracted a virulent gonorrhæa, in addition to an old gleet with which
he had been troubled for fome years. He
expressed much concern at this accident,
imagining that the gleet would be a great
impediment to the cure of the gonorrhæa;
but I soon made him easy on that head, by
observing, that probably both complaints
would be removed at the same time, as I
had very frequently been fortunate enough
to cure clap and gleet both together: and
indeed it so happened in this case, at the
end of sive weeks, by means of injection
and an alterative apozem.

C A S E XXVIII.

ABOUT seventeen months ago, an Irish Peer became my patient; his case was a venereal gleet, with which he had been afflicted many years: bougies as well as internal medicines had been repeatedly used without success. Upon examination I sound a small carnosity in the urethra, from which issued a moderate quantity of yellow matter. I proposed injection, but he at first objected to it, alledging

alledging his having tried it without effect; not confidering (as I observed to him) that the composition of injections was as likely to vary as that of any other remedy. I likewife related to him fome instances wherein that mode of cure had been particularly successful after every other had proved the reverse. He thereupon agreed to put it once more to the test, but was obliged to defer it for a few weeks on account of a long journey he was the next day to fet out upon. In the courfe of a month his Lordship returned, and then informed me that my attendance was doubly necessary to him; having, on the journey, added a clap to the old diforder. The running was plentiful, and accompanied with a heat of urine and chordee.

I IMMEDIATELY recommended the use of injection and a few alterative pills, which, with a proper regimen, perfectly cured him of both clap and gleet in the space of thirty-two days.

From the number of cases that I have seen similar to the preceding ones where a M 2 fresh

fresh infection has been engrafted upon an old gleet, and the little difficulty generally experienced in removing both together, I am led to believe, that in many instances, the stimulus produced by the venereal virus, has frequently contributed to accelerate the cure of the gleet; and particularly so, when it sixes immediately upon the old ulcer; whose callous edges may thus be removed more speedily perhaps than by any other means.

A VERY striking instance of this kind will be found in the subsequent case.

C A S E XXIX.

Four years fince, a Linen-draper in the City, after having ineffectually employed feveral gentlemen of the faculty for the removal of a venereal gleet which had troubled him eight years, made application to me for the fame purpose. I examined the urethra with a bougie, and found very little obstruction therein; I then enquired into the different methods of cure that had been adopted by his other furgeons, to which he replied,

that after going through feveral mercurial courses, taking a great quantity of balsamics, and using a variety of astringents internally as well as topically, bougies were at last recommended; and notwithstanding the uninterrupted application of them for some months, and the strictest regimen at the same time being observed, the disease still maintained its ground. Upon further enquiry, I found that my patient was a very temperate man, and rather of a cold phlegmatic habit; I therefore judged that an alteration in his mode of living might be of use, and accordingly prescribed half a pint of good port wine to be taken every day after dinner, and twice or thrice that quantity after supper. I likewise ordered an injection to be used night and morning, and a few internal alteratives. This plan was purfued for a fortnight or three weeks without producing any very material alteration in the disease; his spirits and appetite however were greatly benefited, infomuch that in the course of a week or ten days longer perseverance therein, he was induced once more to pay his devoirs at the shrine of Venus, and there again became a burntburnt-offering, having then acquired both clap and chancres. His reward for stealing fire (if it may be deemed a thest) differed widely from that which Prometheus experienced; for to his great joy and surprise, the old inveterate gleet was radically cured by the means used to remove the gonorrhæa and chancres, in the course of twenty-eight days.

IT may be asked, whether no fuccedaneum, can be invented to answer the purpose as well as venereal matter? To which I should only give this answer, let future experiments and observations ascertain that point. It may likewise be suggested, that the fupposed advantages accruing from this fresh infection, do not absolutely arise therefrom, but from the medicines used in consequence of it, which may fo correct the latent acrimony of the blood and juices, as to effect a cure merely by that circumftance. My reply to this will be, that I have administered the same kind of medicines, and for a longer time in some of these cases, previous to the new infection. It may then perhaps be faid, that they were not continued long enough;

be it so; or at this rate we might go on ad infinitum, and never put a period to our suppositions; we will therefore leave to practice instead of theory, the determination of this matter.

C A S E XXX.

I was confulted a few years ago by the Captain of a marching regiment, who had been afflicted many years with obstructions in the urethra-they were particularly diftreffing to him during the late war, occasioning repeated suppressions of urine, which at length became fo frequent, as to oblige him to carry a fmall catheter, or a bougie generally in his pocket; unluckily, however, he was unprovided in that respect once upon a forced march, when being attacked with a total suppression of urine, and having no affiftance at hand, he was under the necesfity of taking a small twig from a tree, and after properly shaping, forced it through the obstructions into the bladder, made a pasfage for the urine, and thereby faved his life, which had often been in the most imminent danger from the same cause.

AT the time he applied to me the urethra was fo much obstructed with caruncles, that I found great difficulty in paffing the smallest bougie, and therefore recommended the immediate application of them: his answer was, "That if no other means were used, he was very certain no cure would be obtained, having worn bougies at times for some years past;" Ireplied, "That internal remedies were likewife indispensably necessary in his case, by reason of the very high degree of scurvy with which he was most evidently infested; his body being almost covered with pustules, boils and scabs, the rankest of the kind that I had ever feen. Very powerful antifcorbutics were therefore liberally administered during the whole course of bougie, in order to correct that acrimony, which would otherwise have much impeded, if not entirely prevented the formation of a proper cicatrix. The discharge from the urethra was very plentiful for the first three weeks, and from its sharpness sometimes inflamed the glans penis and præputium as well as the urinary duct, but was kept within bounds by the use of emolient and cooling injections. Thus, was the urethra entirely cleared

cleared of obstructions, the body of its acrimonious humour, and in less than four months the patient's health perfectly re-established.

C A S E XXXI.

A MEMBER of the House of Commons about a year ago fent for me on account of an obstruction in the urethra, which had troubled him fome years, and of late produced a frequent inclination to make urine, though it could flow now only in a very fmall stream, and with much difficulty, from the encreased fize of the carnofity. The obstruction was so great, that it would not admit the smallest bougie to pass it; I therefore contented myself with introducing one up to the caruncle, upon which it made a gentle preffure, being properly fixed with a ligature: I gave directions for its being retained in the same position as long as possible, that the bougie might gradually either infinuate itself through this fleshy substance, or else slip past it. My patient persevered in the method I had proposed for the space of near eleven hours, when he had the fatisfaction of perceiving that the obstruction N

obstruction gave way to the bougie. A small hæmorrhage enfued, which in a few minutes ceased of itself-soon afterwards he attempted to make urine, but at first found the paffage more obstructed than usual; however, upon straining, he forced out something, which he imagined was coagulated blood, but on examination, it proved to be the carnofity itself-in shape and fize it nearly refembled a pea, and was composed of a tough membranous substance—the basis of it was very small, as will be readily conceived from its being fo eafily feparated. The unufual frequency of making water immediately ceased upon the removal of the excrescence, which without doubt had so stimulated the parts as to be the occasion of it; upon the fame principle I prefume as any extraneous body would have done. Astringent injections now became necessary to procure a firm cicatrix; bougies were likewise frequently introduced to preserve the natural diameter of the urethra in that particular part; and, in order to confirm the cure, thefe applications were continued about a week or ten days.

CASE

C A S E XXXII.

A TRADESMAN in Westminster, was for feveral weeks incommoded with a frequent defire to void urine, which at length encreafed to fuch a degree as to prevent his attendance on business: thus circumstanced, he became my patient. Upon enquiry, I found that he had not been subject to either gravel or stone, nor was there any reason to fuspect that infection had given rife to the complaint. I then proposed to examine the urethra with a bougie, but was prevented by a wart just within the orifice; that excrescence I removed with scissors, and destroyed its roots with an escharotic. The patient informed me next day that he no longer felt any irritation at the neck of his bladder, nor had any inclination to urine oftener than he usually did before the diforder took place. The parts foon healed; and he has been free from strangury ever fince.

HERE we may observe, that a disease at one end of the urethra, will produce a disease agreeable

agreeable sensation at the other. It frequently happens that a stone in the bladder, when it presses in a particular direction, will create such an itching at the glans penis and præputium, that the patient cannot refrain from rubbing and pulling those parts.

Excrescences in the urethra fometimes affect very materially the feminal ducts, as will appear in some of the following cases.

IT often occurs, even in claps, that the patient thinks his difease seated in one part of the urethra, when it happens to be in another: indeed we cannot wonder at the many instances of very great sympathy in these parts, when we consider their exquisite sensibility.

C A S E XXXIII.

A GENTLEMAN in the county of Norfolk, had many years laboured under a difficulty of making urine, occasioned by excrescences in the urethra, the consequence of a long continued gleet. He had had frequent suppressions

fions of urine after violent exercise, hard drinking, or excesses of any kind; this he deemed merely a temporary inconvenience, it having generally been foon removed by the introduction of a catheter or a bougie: which circumftance rendered him negligent as to the making of proper application for a radical cure. However, he at last suffered very feverely for his inattention, as will appear in the fequel. In the month of June, 1777, being very much heated with exercise and high living, he was attacked by his old complaint, (the suppression of urine) and recourse was immediately had to the former expedients, but the inflammation and obstruction were fo great, that no instrument could be passed into the bladder-his surgeon endeavoured to relieve him by the usual means, fuch as bleeding, warm bathing, terebinthinate and other clysters, anodynes, &c. but they proved ineffectual. After the suppression had continued upwards of thirty hours, he determined to come up to town, though at the distance of more than one hundred miles. The anxiety and pain that he experienced during the journey, may be better

better imagined than described. On his arrival in town he immediately sent for me: but, half an hour before I saw him, the urine had forced its way through the perinæum, a part of which, and of the adjoining scrotum, I found in a mortisted state. The mortistication extended at least six inches in circumference.

From the pale emaciated aspect of my patient, I could not expect any great affiftance from his constitution, which by his own account had been much impaired by a long fuccession of irregularities. His age did not exceed forty years; in the last ten of which he had been frequently attacked with fevere fits of the gout. All these circumstances confidered, I could not form the most favourable prognosis; however, I was particularly careful, that my opinion should not be couched in fuch terms as might deprefs his fpirits, being fully aware of the bad confequences that frequently enfue from the mind's influence upon the body, especially in putrid complaints. Having in these cases no time to lose, I immediately proceeded to scarify the parts as deeply as the nature of them would

After stuping the wound extremely well with a spirituous somentation, I applied warm digestives, and over them (instead of cataplasm of any kind) soft rags moistened with camphorated spirits, and ordered the nurse to dip a sponge in the same every half hour, and squeeze it over the dressings and the neighbouring parts, in order to prevent as much as possible the mortification from extending itself any further. The dressings were often renewed, on account of the urine's slowing so frequently through the wound.

As internal antiseptics were likewise indispensably necessary, I prescribed an electary of Peruvian bark and Virginian snake-root, to be taken as often and in as large doses as his stomach would bear, and that it should be washed down with camphorated julep or good Port wine, of both which I allowed him to take very freely. His liquors were most of them acidulated with elixir of vitriol, to which was added the aromatic tincture, to prevent its offending the stomach, and at the same time to render it more cardiac.

There were likewise prepared for him strong rich foups, and what is called beef-tea, of which he drank plentifully. Opiates were administered the first twelve days, and at fuitable intervals, clysters to keep his bowels in a proper state. By these means the progress of the mortification was soon impeded, and in a short time entirely stopped. The floughs began to separate in the course of twelve days, when I thought it adviseable to attempt regaining the natural passage for the urine, the whole of which constantly flowed through the wound ever fince it had first made an outlet that way. The caruncles in the urethra made so much resistance, that four days elapfed before the smallest bougie could be passed through them; but as foon as that point was gained, a bougie was constantly kept in the urethra both night and day. The urine by this method foon recovered the natural channel, and in less than a fortnight, not a drop of it passed through the wound, which now digested well, and in a few days incarned, and foon after cicatrifed, as kindly as could have been expected. By continuing the bougies fix weeks longer, the urethra

fructions, and the parts firmly healed. Soon afterwards, my patient returned to the country in high fpirits, and in a much better state of health than he had been for many preceding months.

SEMINAL GLEET.

HAVING hitherto made little or no mention of the feminal gleet, I will now offer a few observations upon the origin of that disease, and produce some particular cases, both to illustrate the subject, and recommend a method of cure that has frequently succeeded after every other had proved inessectual.

THAT seminal gleets or weaknesses (call them by which name you please) are of all others, the most difficult to cure, I presume, will not be controverted by any experienced practitioner; this difficulty will be encreased in proportion to the immediate cause of the disease, and the contingencies that may attend it, the habit of body, mode of living, climate, profession or occupation, &c. of the patient,

patient. The difease may arise from those excrescences in the urethra, which we term caruncles, carnosities, &c. and it is sometimes occasioned by the destruction of that natural caruncle, the caput gallinaginis urethræ. It originates likewise from an acrimonious state of the blood and juices. Sometimes it is produced by a general relaxation of the whole frame, from whatever cause soever that may take its rise, whether from a feries of debaucheries, excessive venery, self-pollution, &c. &c.

In some of the foregoing cases, I have demonstrated that an excrescence in the urethra will stimulate the bladder (though at the distance of some inches) to such a degree as to bring on a frequent inclination to evacuate its contents; and when that stimulus happens to fix upon the seminal ducts, the reservoirs, or the testicles themselves, similar effects must naturally be expected. After the seminals have been thus stimulated for a length of time, their retentive faculties must indubitably become so much impaired and weakened, as to admit of very frequent

frequent and involuntary emissions. The fame consequences may also be expected, when the blood is much impregnated with acrimony of any kind, and a confiderable share of it falls upon any particular organ (a gland for instance, or its excretory duct) the irritation thereby produced must excite an unnatural fecretion, as well as expulsion of that particular fluid, and will at the same time alter the quality of it. A long continued repetition of this stimulus gradually relaxes the mouths of the excretory ducts, together with their valves or fphincters, those natural obstacles to involuntary evacuations; in fine, they entirely lofe their original powers of retention, and the fluid is frequently discharging itself, almost as foon as fecerned; or upon the least exertion in making urine or going to stool, supposing the feminal veffels to have been thus vifited by the acrimonious humour.

C A S E XXXIV.

An Officer in the Navy after having laboured under a seminal gleet for six years and upwards, put himself under my care about two years and a half fince. His spirits, strength, appetite, &c. were greatly reduced, infomuch, that he was threatened with a total loss of virility. After interrogating him as to the cause of his disease, I examined the urethra with a bougie, and found it perfectly free from obstruction: I then inspected the colour and confistence of the discharge, and foon perceived that it had loft its natural texture, and feemed to abound with acrimony, which upon further enquiry proved to be fcorbutic. He informed me that he had been much afflicted with the scurvy for some years previous to the appearance of his prefent complaint. Having now investigated the cause of the disease, I had but little difficulty to apprehend in performing the cure of it; as my patient, from his appearance as well as promifes, perfuaded me, that nothing on his part would be wanting for the completion of it. I had not formed a wrong conjecture; for with the greatest regularity and attention he persevered in a course of antiscorbutics, which in a few months entirely removed his gleet, and at the fame time perfectly reestablished his health in every other respect. CASE

C A S E XXXV.

A GENTLEMAN in the county of Middlefex, applied to me two years ago, on account of a feminal gleet, to which at times he had been for some years subject, though at intervals entirely free from it; his vigour therefore was not very much impaired thereby, but as the returns of the complaint, had of late been more frequent than usual, he began to be very apprehensive of consequences. Upon examination, I found that the cause of this gleet might be truly deemed arthritic; it was indeed fufficiently evident from the appearance of the discharge, and the patient's own account of his cafe. The femen was loaded with a cretaceous matter, fimilar to the chalk stones produced by the gout; in proof of which he informed me that he had been afflicted with the gout many years, chalk stones had appeared on his feet, but that no more of them had formed after the cretaceous matter found its way to the fpermatic veffels, and (as he very rationally fupposed,) discharged itself with the semen ever fince that period. Being perfectly fensible that

that the disease originated from the gout, I recommended patience, and a resignation to his sate. This prescription not satisfying him, I promised to consider further upon his case, and exert my best abilities to serve him. After a little reslection, there occurred to my memory the case of a gentleman, who some years since employed me to cure him of the venereal disease, and as he was greatly afflicted with the gout, requested very earnestly that I would not give him the least particle of mercury; I acquiesced therewith, and in the course of a few weeks removed his complaint, merely by the use of vegetables.

Sometime afterwards he informed me, that these medicines had so purified his blood and juices, that the gout had not visited him so frequently as usual, and the paroxysms were much milder and of shorter duration than they formerly had been. He has therefore taken the medicine since that time as an anti-arthritic. The recollection of this circumstance induced me to administer remedies of a similar kind for the cure of this arthritic gleet; and I had soon the satisfaction

of perceiving that the discharge gradually decreased, and was less and less impregnated with the chalk, until it entirely disappeared, which happened in about two months: I advised nevertheless a continuance of the medicines sometime longer in order to prevent a relapse. The medicines used upon this occasion were, diuretics and diaphoretics, in the composition of which, camphor was no inconsiderable ingredient.

BEFORE we difmiss the subject, it may not be improper to mention fome other diforders, to which the organs of generation are liable from this arthritic acrimony. I have been informed, from very good authority, of priapisms originating from the same fource; one in particular, of a gentleman in the county of Surry, which is faid to have continued an incredible length of time. Instances I have myself seen where the gout had produced tumefactions of the testicles, to as high a degree as those in venereal cases, and which, like them, have been removed by discutient fomentations, and emollient cataplasms, assisted with powerful sudorifics -internally.

[104]

internally. In these cases opiates likewise may repeatedly be administered to very great advantage.

C A SE XXXVI.

ABOUT eighteen months fince, I was confulted by an eminent Artist in the vicinity of London, who had for many years been troubled with a gleet and obstructions in the urinary passage, occasioned by a clap which was contracted eleven years before. There had been, ever fince that period, a discharge of yellowish matter that might be faid to ebb and flow, being very copious at one time, and quite the reverse at others; excess of any kind, never failed to encrease it very confiderably. Within the last three years he perceived the urethra to be materially obstructed, and from that time a new symptom made its appearance, namely, a mixture of femen with the old running; in proportion to the encrease of the caruncles, the feminal discharge became more plentiful, infomuch that during the last four or five months, he very feldom made urine or went to stool, without a confi-

a confiderable quantity of feed being evacuated at the fame time. He informed me of his having taken large quantities of the most bracing and strengthening medicines, but with no other effect, than a temporary palliation of the fymptoms. The long continuation of the difease, added to a close attention to bufinefs, made terrible havock with his constitution, reduced him almost to a skeleton, and relaxed his nerves to fuch a degree, as to render life burthenfome in the extreme. The excrefcences had fo choaked up the urethra, that the fmallest bougie could not be passed without great difficulty. I began his cure, by attempting to remove these obstructions, being fully convinced that no medicine could be of use till that was effected. I advised him to wear a bougie both night and day, and for as many hours together as the parts would admit of, without occasioning pain or loss of rest. His bowels were at the same time kept in proper order by the use of gentle aperients: to which circumstance it will always be found indifpenfably neceffary to attend in every complaint of this nature,

P

both

both in order to prevent inflammation, and to obviate any material exertion in the evacuation of the fæces. His diet was of the most nutritious kind, but as he at first had little or no appetite, I gave him strong decoctions of the woods, which foon procured one, and at the same time corrected any latent acrimonious humours: after which, corroborants and restoratives were administered with the greatest success. By these means, at the expiration of four months, he was fo far recovered as to imagine himfelf cured; the medicines however were continued feven or eight weeks longer, by which time his health was perfectly reestablished.

C A S E XXXVII.

Almost fimilar to the preceding case, was that of a Gentleman in the County of Cambridge, who had been afflicted for several years with a seminal gleet, and had taken a great quantity of medicines of different kinds, without receiving any material benefit—a circumstance not to be wondered at, when we consider that the cause of

of the disease had not hitherto been understood, no attention having been paid to a caruncle in the urethra, which at last encreafed fo much, as almost entirely to prevent the paffing of his urine; upon this account it was that he made application to me, having long before given up all expectation of removing the feminal gleet, every effort for that purpose having so repeatedly proved abortive. Luckily however, the stoppage of urine obliged him to have recourse to the only method that could possibly remedy his old diforder. By using bougies, the urethra was cleared of obstruction in the course of a few weeks, in consequence of which, the seminal discharge gradually abated, and with the affistance of some internal bracers, a radical cure was effected in less than five months.

C A S E XXXVIII.

A GENTLEMAN in the western Part of England, became my patient three years fince, in confequence of a feminal weakness or gleet, which had been his companion for thirteen years, and was at length very near

P 2

putting

putting a period to his existence, notwithstanding a long continued use of the most powerful restoratives both in medicine and diet, which had been prescribed for him by several eminent practitioners.

HE had taken aftringents, balfamics, strengtheners and bracers of almost every kind, large quantities of bark, steel, ginfang, &c. &c. nor was the cold bath omitted, which in the beginning, he faid, afforded a temporary relief, but after a few months use entirely lost its effect. The difease had rendered him so weak, as to be fcarce able to walk or ftand upright, and his spirits were likewise proportionably depressed. He informed me, that his present complaints took their rife from a clap contracted upwards of thirteen years fince, that it was not attended with much pain or inflammation, nor were any of the fymptoms fuch, as indicated any high degree of virulence; his furgeon, however, had fupplied him very liberally with mercurials and draftic purges, and had continued them fome weeks, by which means his appetite, strength,

strength, &c. were much impaired, and a gleet enfued, which from a venereal, degenerated into a feminal one. Frequent nocturnal emissions of femen interrupted his fleep, and a discharge of the same kind generally happened whenever he made urine or went to stool. After having given this account, he requested to have my opinion upon his cafe. My answer was; that from the long continuance of the venereal gleet, I fuspected some excrescence had formed in the urethra, and should that be the case, I doubted not of rendering him very effential fervice. He replied, "That he did not imagine there could be any material obstruction in the passage, as his urine flowed almost as well as ever; but that it did not pass quite so freely, which he attributed wholly to the debility of the parts. However, upon examination with a bougie, I found a caruncle about four inches up the urethra, which, in that place diminished its diameter nearly one third part; the rest of the canal was perfeetly free from either stricture or any other obstruction: this discovery gave my patient much pleasure, and at the same time great furprise,

furprise, that such a substance should be formed without his perceiving it; that was a circumstance, I told him, which frequently occurred, and might be very eafily accounted for, by only reflecting upon the very flow, gradual progress of these carnosities; in the present instance above twelve years had elapsed during the formation of a very fmall one. I then advised him to enter upon the use of bougies; and that they should remain in the urethra as many hours as possible, both night and day, provided neither pain nor inflammation enfued, to prevent which, as well as too great efforts in discharging the fæces; his body was kept rather in a laxative state than otherwise, by taking occasionally flos sulphuris, magnesia alba, or electarium lenitivum. In the course of a week he was able to wear the bougies twelve hours in the twenty-four; by which means a copious discharge of matter was produced, and in proportion to that, did the caruncle decrease, and the seminal evacuation abate. In less than a month the carnofity was removed, and with it the stimulus which had operated so powerfully upon the spermatic vessels, &c. At this time I thought corroborative

roborative medicines would take the defired effect, I accordingly administered them in as large doses, and as repeatedly as his stomach would bear, his diet was likewise of the most nutritious kind, and as simple as possible, all high seasoned sauces, &c. being studiously avoided, together with spirituous liquors, and vegetable acids. By the above method this gentleman was restored to perfect health in the space of eight months; he married soon after, and has got an heir to his estate.

In these cases I always recommend animal food, or a meat diet in preference to any other: and when the stomach is too weak to admit of it in substance, strong soups or broths must be extracted from it, and taken very liberally. After meals, a glass or two of good old Madeira or Mountain wine may be drank, and will generally prove beneficial. Moderate exercise will likewise be useful, but great care must be had that it never amounts to satigue. The patient should be thinly clad both night and day; rise early, and go to bed the same. In sine, the same regimen may be observed in these complaints

as is generally recommended in those termed nervous.

ALL excrescences or carnosities in the urethra have not a venereal origin, that, in common with every other glandulous membrane, may certainly produce them from fome other causes; among which, coarse gravel and small rough or sharp stones passing through this tube may fometimes occasion them. An instance of this kind I remember to have feen in a Kentish farmer, who applied to me on account of obstructions in the urethra, which could not be supposed to arise from any venereal cause, as he solemnly declared, that he had never been foinfected. He had however frequently voided gravelor stones with his urine. which, I make not the least doubt, gave rife to the excrescences, from having wounded the membrane, which afterwards healed, but not without producing thefe fungi.

THE glandulous membrane that lines the nose, we know will sometimes send forth excrescences named polypi, independent of any venereal matter whatever.

BEFORE

BEFORE we close our observations upon gleets and weaknesses, let us turn our thoughts upon those which are peculiar to the fair sex.

THE leucorrhæa, fluor albus, or the difease vulgarly termed the whites, though often with very great impropriety, the difcharge being at times as high coloured as that produced by the venereal virus itself. This gleet is the cause of many disorders, and as frequently the consequence of others. It fometimes arises merely from a cacochymia, at others from a profluvium menfium, bad labours, miscarriages, a general relaxation, immoderate venery, acrimonious humours, &c. When this discharge is very copious and of long duration, it produces many difagreeable (if not fatal) effects: it may give rife to the whole tribe of what are called nervous complaints, hysterics, &c. and may fometimes occasion an atrophy or confumption.

It is to be lamented, that they are most subject to this disorder, who apparently are

Q

the least able to sustain it, namely, those of the most delicate frames and constitutions. A sedentary mode of living, improper diet, too much cloathing, bed-warming and teadrinking, contribute greatly to encrease, if not to produce, this disease: the use of strong liquors (I am asraid) may sometimes be added to the catalogue of causes.

From a defire of becoming more delicate, many good constitutions have been greatly injured, (if not entirely ruined) by the too frequent repetition of phlebotomy and strong cathartics.

The pudenda of each fex being generally very much benefited by cold bathing, why, in the name of common fense, is it not more frequently practised? Women, in particular, would derive infinite advantage from washing these parts regularly night and morning with cold water, except at the time of menstruation; as I believe that sew things would tend more to prevent the disease in question than these ablutions, if properly pursued. An immersion of the whole body would

[115]

would certainly be preferable to a partial bath, but as the former is not always practicable, and is generally attended with great trouble and inconvenience, we must content ourselves with the latter, provided it be diligently persisted in.

CERTAIN fects are strictly enjoined to make frequent ablutions, as a ceremony or duty incumbent on the professors of such religions; a custom, however, of this kind, (if viewed only in a temporal light) must be highly beneficial to society.

CAN any good reason be assigned, why the external parts of generation should not be as often washed as our faces, they must certainly require it as frequently, although seldom exposed to view? Courtesans (for more reasons than one) are the strictest observers of this salutary custom.

THE great difficulty attending the cure of the fluor albus, may in some degree be ascribed to the very lax texture and depending situation of the parts subject to this discharge.

Q 2

In the cure of female gleets, we must administer such alteratives as will remove any latent acrimony, and at the fame time recommend the most nutritious diet, together with a proper regulation of the other nonnaturals. After the blood and juices shall have been fufficiently purified by the means abovementioned; then may balfamics and astringents be used to great advantage. Be it remembered also, that we must not rely folely upon internals, for it will generally be found necessary to use topical applications at the same time, viz. lotions and injections, which must be composed of the most powerful aftringents. The cold bath will likewife be of fervice, where unfound vifcera or very great tenderness of lungs do not prohibit the use of it. The alteratives recommended in the former part of this Essay will, upon trial, be allowed the most efficacious ones in the present complaint. Opiates, in many stages of the disease, may likewise be administered to very great advantage, provided they be so corrected, as to obviate any material constipation of body.

[117]

OF all the non-naturals, a due regulation of the passions must be very strictly attended to; nay, it will be found indispensably necessary, as without it no cure will ever be obtained; indeed, it will generally prove one grand essential in the cure of many female complaints.

THAT the vagina uteri, and its vicinities are very liable to great defluctions is fufficiently evident from the difease just now mentioned, which in general does not make its appearance before the age of puberty: fome females however, even in the infant state, experience so great a flow of humours as to produce a discharge, in colour, not unlike the gonorrhæal one, and frequently acrimonious enough, to occasion much inflammation, pain, and tumefaction in these parts. Several instances of this kind have fallen under my inspection, and upon enquiry, I have generally found that the complaint arose from either a scorbutic or a scrophulous humour. Emollient fomentations and cataplasms, assisted with gentle purgatives, and mild alterants, will foon foon abate the fymptoms, which generally disappear in the course of a few days; especially if the discharge has been rather copious in the beginning.

To prevent a relapse, it will be necessary to keep the intestines in a laxative state, which may be easily effected by the use of lac sulphuris, or magnesia alba; the preference however must be given to the former, as it will at the same time prove no insignificant alterative.

THE most obstinate cases are such as derive their origin from a leprous humour, one of these I remember to have seen; it was attended with a very high degree of inflammation, and required venæsection, opiates and cathartics, together with the most powerful internal alteratives, and external emollients. A composition of camphor, antimony and sulphur, was in this case extremely useful; the cure indeed might with justice be wholly ascribed to it; every other medicine having proved ineffectual.

[119]

THE cure of gleets and weaknesses in either sex, will generally require a considerable time, but if the methods I have recommended be put in practice, and properly pursued, I am firmly persuaded that they will very seldom fail of success.

FINIS.

