

Observations on the muscles, and particularly on the effects of their oblique fibres : with an appendix.

Contributors

Monro, Alexander, 1733-1817.
University of Glasgow. Library

Publication/Creation

Edinburgh, 1794.

Persistent URL

<https://wellcomecollection.org/works/btus9am4>

Provider

University of Glasgow


License and attribution

This material has been provided by This material has been provided by The University of Glasgow Library. The original may be consulted at The University of Glasgow Library. where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>



*I Liffing
from
the author*

OBSERVATIONS
ON
THE MUSCLES,
AND PARTICULARLY ON
THE EFFECTS OF THEIR OBLIQUE FIBRES:
WITH
AN APPENDIX,
IN WHICH
THE PRETENSION OF DR GILBERT BLANE,
THAT HE *FIRST* DEMONSTRATED THE SAME EFFECT TO BE
PRODUCED BY OBLIQUE MUSCLES AS BY STRAIGHT
ONES, WITH A LESS PROPORTIONAL DE-
CURTATION OF FIBRES,
IS PROVED TO BE UNFOUNDED.

BY ALEXANDER MONRO, M. D.
PROFESSOR OF MEDICINE, ANATOMY AND SURGERY IN THE UNIVERSITY
OF EDINBURGH; FELLOW OF THE ROYAL COLLEGE OF PHYSICIANS,
AND OF THE ROYAL SOCIETY OF EDINBURGH, AND OF
THE ROYAL ACADEMY OF SURGERY IN PARIS.

EDINBURGH:
PRINTED FOR J. DICKSON, AND E. BALFOUR;
AND T. CADELL, LONDON.

M.DCC.XCIV.

Digitized by the Internet Archive
in 2016

<https://archive.org/details/b24927788>

OBSERVATIONS

ON THE

MUSCLES, &c.

AS it appeared to me, when I first began, in 1759, to deliver in this University a public course of lectures on Anatomy and Surgery, that the structure of the oblique muscles had not been sufficiently examined, nor even the number of them attended to by authors, and that some of their chief purposes or effects had been entirely overlooked by them, I endeavoured then, and in every course of lectures since that time, to direct very particularly the attention of students to those subjects.

I BEGAN with observing, as a material defect in the otherwise very accurate and elegant tables of ALBINUS, as well as in the former systems of VESALIUS, EUSTACHIUS, BIDLOO and COUPER, that the tendinous membranes or aponeuroses, with which many muscles, particularly of the extremities, are covered,

A

vered, and with which the oblique muscles are closely connected, were not delineated, yet that the knowledge of these is not only of use in the practice of surgery, but for understanding the action of the muscles.

I REMARKED, that although in some parts of the body, tendinous membranes, such as those between the cartilages of the ribs, or the aponeuroses palmares, or fasciæ latæ of the thighs, served merely for the defence of the parts, or as sheaths to them, as they were connected to them by the cellular substance only, yet, in general, they served, besides the mere purpose of defence, to furnish a greater extent of surface for the attachment of oblique fleshy fibres.

I SHEWED them, that wherever tendinous membranes run longitudinally on the surfaces of muscles, fleshy fibres, placed obliquely, were found; that in many muscles, as in the semimembranosus, or flexor pollicis longus, fleshy fibres passed obliquely from the inner part of the tendon on one side, to the inner part of the tendon on the other side of the muscle, or such muscles were semi-penniform, (See T. 1. fig. 1.); that in other muscles, as in the rectus extensor cruris, or flexor pollicis pedis longus, a third tendinous membrane was found in the middle of the muscle, between which and the inner parts of the tendons on the two sides of the muscle, the fleshy fibres passed obliquely, and produced a complete penniform appearance, (see T. 1. fig. 2.); and some muscles, as the soleus, might be called compound penniform, because, on cutting them lengthways, we discovered several longitudinal tendinous membranes, to both sides of which oblique fleshy fibres were connected. See T. 1. fig. 3.

I ALLEGED, that the direction, length and number of fibres in such muscles had not been sufficiently attended to by anatomists or by surgeons; and that, in many instances, the breadth of these muscles had been mistaken for their length; that in consequence of such inattention, they would find the celebrated LOUIS * attempting to discard the double incision of the soft parts in the amputation of the thigh, although, from the obliquity and shortness of the muscular fibres which cover the thigh-bone, this improvement of CHESELDEN is more essential than in the amputation of the humerus, where the fleshy fibres, though oblique, are proportionally longer, and of course their retraction greater.

WHERE the fibres of muscles run obliquely, it is evident, and has been observed by BORELLUS and others, that the fibres will be more numerous than if the same space had been covered with longitudinal fibres; and although an oblique fibre will not raise a weight with the same force as a straight fibre, yet the number of the fibres may be so much increased by their obliquity, as to do more than compensate for the loss of force occasioned by the obliquity. Thus, let us suppose a longitudinal muscle to be five inches long, and one inch in breadth, and let us suppose it to contain in its breadth four fibres or ropes, each one-fourth of an inch in diameter, as in T. 2. fig. 1. the force of this muscle may be represented by the number 4.

LET us next suppose these ropes to be cut into pieces, each of which is one inch and a quarter in length, as represented by dotted transverse lines A, B, C, we shall, by doing so, form 16 ropes or fibres.

A 2

LET

* Acad. Roy. de Chirurg. T. 2. p. 357.

LET us next suppose, that these ropes, representing muscular fibres, are laid obliquely, like the hypotenuses of right-angled triangles, of which the bases are equal to one inch, and the height or perpendicular equal to three quarters of an inch, as in T. 2. fig. 2. each such fibre will, as BORELLUS has demonstrated, lose two-fifths of its force. But as there are 16 fibres instead of 4, their force will be as 16 multiplied by 3, to 5 multiplied by 4, or as 48 to 20, or as 12 to 5.

BUT that the mere increase of the number of fibres, or force of the muscles, which alone has been observed by authors, is not the sole purpose of nature, appears from this, that in some places, and particularly between the ribs, oblique fibres are employed, although it is evident, that a greater number of straight fibres, or of fibres perpendicular to the ribs, might have been disposed in the same space.

THE other purposes of oblique muscles, and which had not been perceived by authors, are,

To perform much more extensive motions with the same degree of shortening of the fleshy fibres, than can be performed by straight muscles, or with a less degree of shortening of the fibres, to perform motions of equal extent.

I SHALL now endeavour, in the first place, to demonstrate, that a pair of oblique muscles, placed between the same parallels with two straight muscles, perform, with the same proportional contraction, more extensive motions than the straight muscles can do.

FIRST,

FIRST, As one oblique muscle, so situated, is longer than a straight muscle, if each be shortened one third, or any other proportional part, it is evident, that the place of the insertion of the muscle will move through a greater space, when the oblique muscle acts. Thus, in the annexed figure, (T. 2. fig. 3.) if P, L represent two parallel lines, and AB represents an oblique and AC a straight muscle, it is plain, that when each is shortened one third, and that the place of the insertion is moved directly towards the place of the origin of the muscle, the motion occasioned by the oblique muscle will be proportionally as much greater than that produced by the straight muscle, as the hypotenuse, or line AB, is longer than the perpendicular line AC.

BUT next, let us suppose, (see T. 2. fig. 4.) that the point A cannot be drawn directly towards the point D or E, on account of the connections of the bones, such as the ribs, which the parallel lines PA and DE represent; or suppose, that such bones, when moving, remain parallel to each other; or let us suppose, that two oblique muscles balance each other, so that their insertions, instead of being moved directly towards their origins, are moved in a diagonal line, between the two muscles.

THUS, let PA and DE (T. 2. fig. 4.) be two parallel lines, and let AB and AC represent two straight muscles, and AD and AE two oblique muscles, it is plain, that when the two straight muscles have shortened themselves one third part of their length, their insertion A will be brought down to number 1. But when the oblique muscles AD and AE, by acting together, have brought the point A down to 1, and are in the situation of the dotted lines 1 D and 1 E, they cannot have lost more of their length than the length of the perpendicular A 1,
which

which is shorter than the hypotenuse Ab , or less than the third of the length of the oblique muscle. In fact, they have lost less of their length than $A1$, because the two sides $A1$ and $1E$ of the triangle $A1E$ must be longer than the third side AE ; and therefore oblique muscles can perform as great a degree of motion as straight muscles, without being shortened in the same proportion; or, which is the same thing, if they continue to act till they are shortened in the same proportion, the place of their insertion, A , will descend farther, or through a larger space.

I SHALL now proceed a step farther, and endeavour to demonstrate, that where two oblique muscles balance each other, the motion of their insertion is more extensive than can be produced by two straight muscles of the same length with the oblique muscles.

THUS, in T. 2. fig. 5. let AB and AC represent two straight muscles, and AD and AE two oblique muscles of the same length, and we shall suppose the length of each muscle to consist of any given number of inches or parts, suppose five parts, 1, 2, 3, &c. or I, II, III, &c. and when in action to be capable of shortening itself one fifth part or two fifth parts of its length. Let the two corresponding numbers 1 and I, or 2 and II, be joined by the straight lines $1I$, and $2II$, so as to form the isosceles triangles $A1I$, or $A2II$. When the two straight muscles have acted fully, or shortened themselves one-fifth of their length, the point A will descend to 1 . But when the two oblique muscles have, by their action, brought the point A down to 1 , they have not lost one-fifth of their length; for the dotted lines representing them must be longer than the lines IE or ID , because the angle $1IE$ being equal to the
two

two angles $\angle I A$, and $\angle A I$ of the isosceles triangle $A I I$, must be larger than a right angle, and therefore the side $I E$ must be longer than the side $I E$; that is, the oblique muscles, after bringing the point A down to I , have not lost one-fifth of their length; or if they continue to act till they have lost one-fifth of their length, they will bring the point A lower down than can be done by straight muscles, shortened in the same proportion.

To make this proposition still plainer, if possible, by calculation, I shall suppose the oblique and straight muscles in T. 2. fig. 6. to be each five parts or five inches in length; that the bases of the triangles $B D$ and $C D$ measure four inches; and that the perpendiculars, or altitudes of the triangles, measure three inches; and let it be supposed, that these muscles, in action, can be shortened one-fifth of their length, the straight muscles, on that supposition, can bring A down to I only: But it is evident, that the oblique muscles will not be shortened one-fifth of their length till the point A has descended to D , or to number 3; or the oblique muscles will, with the same degree of contraction, move the point A three times farther than can be done by straight muscles of the same length.

In the next place, we may easily demonstrate, that the extent of the motion produced by the co-operation of oblique muscles, increases with their greater degree of obliquity.

Thus, let us compare the extent of motion, produced by the pair of oblique muscles $A D$ and $A E$, (T. 2. fig. 5.) with that of the still more oblique pair of muscles represented in the same figure by the lines $I D$ and $I E$. Let the muscles

$A D$

AD and AE be supposed to move the point A to number 1, and let the muscles 1 D and 1 E be supposed to move number 1 to number 2, or through a like space. It is evident, that in the triangles 1 I E and 2 II E, the angles 1 I E and 2 II E are equal; but as the angle 2 E II is larger than the angle 1 E I, the angle II 2 E must be less than the angle I 1 E. Hence, as the sides of triangles are longer in proportion to the width of the opposite angles, the side I E will be longer in proportion to 1 E, than the side II E is in proportion to 2 E. The muscular fibres, therefore, AD and AE, in bringing the point A down to number 1, will lose more, in proportion of their length, than the more oblique fibres 1 D and 1 E will do in moving number 1 to number 2.

To prove this by calculation, let us suppose the muscle to be still represented by the hypotenuse of a right angled triangle, five inches in length, and capable of shortening itself one inch, and that one of the other sides measures four inches, and that the third side measures three inches. But let the side 3 form the basis of the triangle, and the side 4 its perpendicular, as in T. 2. fig. 7.

IN this case, the square of the hypotenuse, when it has shortened itself one inch, will be 16, from which deducting 9, the square of the basis, the number 7 remains for the square of the perpendicular. But the square root of that number being more than $2\frac{1}{2}$, the oblique muscles, shortened one-fifth, cannot bring the point A down $1\frac{1}{2}$ inch, or to B, or cannot move the point A half so far as they were shewn to do, when the obliquity was greater, by making the basis 4 inches and the altitude 3 inches.

Or

OR let us, on the other hand, increase the obliquity, as in T. 2. fig. 8. by supposing two right-angled triangles, so constructed, as that their hypotenuses measure 13 inches, their bases 12, and altitude 5 inches, and that the hypotenuses represent two oblique muscles. It is plain, that when these have shortened themselves one inch, or one thirteenth part of their length, they will move the point A through a space of five inches, or five times farther than straight muscles of the same length, shortened in the same proportion, could do.

HENCE, as the obliquity of an oblique muscle is gradually increasing during its action, its force is diminishing; while its effect, of producing extensive motion, is increasing. Thus, a muscle, representing the hypotenuse of a right-angled triangle, whose sides are to each other as the numbers 3, 4, and 5, and the altitude 3, by shortening itself half an inch, does not move its insertion one full inch; but if it is shortened another half inch, its insertion is moved through a space of upwards of two inches more. When it begins to act, it has three-fifths of the strength of a straight muscle of the same size; but when it acts again, after having moved its insertion the space of an inch, it has two-fifths only of the strength of the straight muscle.

To illustrate what I have been demonstrating, I used, after dissecting and demonstrating the recti muscles of the abdomen, to cut their ends off from the ossa pubis, and to apply them to the tops of the ossa ilia, so as to represent oblique muscles; and from this I was led to make the remark, that if both the two external oblique, or the two internal oblique muscles, or all these, acted at once, the obliquity of the one balancing the obliquity of the other, the trunk of the body would be bended

B

straight

straight forwards, and that flexion made by them might be greater than that made by the recti muscles, which, at first sight, seemed to be more suited to the purpose.

IN like manner, I used to take out several of the ribs, with their intercostal muscles; and after shewing the two layers of these muscles laid obliquely, and decussating each other, I used to dissect some portions of the two layers, in such a manner as to represent oblique muscles, with their origins at a distance from each other, but their insertions meeting in a point, or with their insertions, as well as their origins, at a distance from each other.

IN the back part of the spine, I very particularly demonstrated the obliquity of many of the muscles, some of which are called semispinales, because one end of them only is fixed to the spinal processes, and the fibres passing obliquely, the other end of them is fixed to the transverse processes, or other parts of the neighbouring vertebræ.

IN the extremities, I not only carefully demonstrated the obliquity of the fleshy fibres, in the half and whole penniform muscles, but pointed out their connection with their tendinous aponeuroses, the different direction of the tendinous and fleshy fibres, and the uses of the aponeuroses and tendinous sheaths; and that, by means of the sheaths, there was so little difference between the length of the muscles in the bended and extended state of the member, that short fleshy fibres, especially when placed obliquely, could produce a very extensive motion.

BUT

BUT in treating of particular parts, I dwelt chiefly on the structure and effects of the intercostal muscles, as a variety of opinions concerning their operation has, in the course of the last hundred years, been proposed, and as no author had explained the reason of the obliquity of their fibres, nor of their being disposed in two layers of decussating fibres.

THAT their structure might be fully understood, I first laid bare the surface of the external intercostal muscles, and between the next two ribs, I cut off the external intercostals, to shew the internal, as in T. 3. fig. 4.

IN another space, I shewed a small bundle of the external intercostal, decussating a similar bundle of the internal intercostal, and forming a figure like the letter X, but in which the stroke representing the external muscle is more oblique than the other; for the internal intercostals are less oblique than the external. See T. 3. fig. 5.

THEN I dissected small bundles of the external and internal intercostals, with their origins at a distance from each other, but their insertions meeting in a point, in the rib above or in the rib below, so as to form triangles, of which the rib made the base, (see T. 3. fig. 6. and 7.); or I dissected them with their insertions, as well as their origins, at a distance from each other, as in T. 3. fig. 8.

IN the last place, I demonstrated a part of the structure which has not been sufficiently examined by authors; to wit, that the cartilages between the ribs and the sternum, with the exception of the cartilage of the first rib, are not fixed to the sternum in the same manner as to the ribs; for the rib, which

is hollowed, receives the cartilage, and is so firmly united to it, that in a recent subject, they cannot be separated without lacerating the cartilage; but the inner part of the cartilage is tied by a capsular ligament to the edges of the pit in the sternum, and the concave part of the pit is connected by fine cellular threads only to the end of the cartilage, so that the cartilage and sternum may, after cutting the capsular ligament, be separated from each other without tearing the cartilaginous fibres. Hence, when the ribs are moved, the capsular ligament is twisted, and the end of the cartilage rolls upon the sternum. See T. 3. fig. 1, 2, 3. and 9.

AFTER fully explaining the structure, I endeavoured to prove, as Dr HALLER had done, but with some additional arguments, that both rows of intercostal muscles conspired to elevate the ribs, or that they were muscles of inspiration; and that, when the intercostal muscles alone acted, and the ribs were not forcibly kept down, they could have no other effect; and that all the ribs in inspiration were moved upwards uniformly.

THE chief circumstances which prove beyond a doubt, that the two rows of intercostal muscles conspire in elevating the ribs, are,

1. THAT the first rib is so much fixed at both its ends, as to be almost immovable, and its cartilage, instead of being connected to the sternum by a capsular ligament, or articulated with it in the same manner as the cartilages of the other ribs, grows as firmly to the sternum as to the rib. See T. 3. fig. 9.

2. THAT the second rib is more fixed than the third, and the third more fixed than the fourth, and so on downwards.

3. THAT as the ribs, from the first rib downwards, grow gradually longer, and describe portions of larger circles, we may observe, that in general, or when we examine a middle portion of the intercostal muscles, or a portion half-way between the sternum and vertebræ, the insertion of the lower end of the portion is at a greater distance from either end of the lower rib, or from a straight line drawn between the two ends of that rib, than its origin in the rib above is from the two ends of that rib, or from a straight line drawn between them. Hence, whether we consider the head of the rib, connected with the vertebræ as its centre of motion, or whether we consider the rib as moving upon a straight line or axis drawn between its two ends, it follows, that a muscle placed between two ribs acts with a longer lever upon the under rib than upon the upper one, and therefore must elevate the under rib. That the force of this argument might be more readily understood, I have laid leaden probes along each of the seven uppermost ribs of an adult subject, from the vertebræ to the sternum, and have represented their lengths and curvatures in T. IV. The crooked continued lines represent the lengths and curvatures of the different ribs and their cartilages. The straight dotted lines represent the distances between their heads and the sternum. The continued perpendicular line represents the distance of the middle of each rib from a straight line drawn between its two ends. The numbers 1, 2, 3, 4, 5, 6, 7, express first, second, &c. ribs, of which the first is the shortest and innermost, and the seventh the longest and outermost. The other numbers annexed denote eighths of an inch.

4. To

4. To determine the effect of the contraction of any muscle, I apprehend we need only to observe in the dead body what the situation is in which the muscle in question is relaxed. Applying this rule, we shall find that the whole intercostal muscles, internal as well as external, are shortened when we elevate the ribs, and place them in that situation in which we find they are in inspiration.

5. If the internal intercostal muscles had been intended for the depression of the ribs, we certainly should not have found them continued to the sternum, because their anterior ends are fixed above to the edge of the sternum, or so near to the insertion of the cartilage of the upper rib in the sternum, and their inferior ends are, in consequence of their obliquity, fixed to the under rib so much farther from the sternum, that they must act upon the under rib with more advantage of lever, or are intended for its elevation.

On the other hand, if the internal intercostals had been intended for the depression of the ribs, we certainly should have found them continued backwards to the spine, because, from their obliquity, their under end would have been fixed to the vertebræ or nearer to the head of the rib, and their upper end at such a distance from it, that this portion of the muscle would have been better calculated than any other portion of it for the depression of the rib.

6. In a few experiments which I made on living animals, soon after I began to study anatomy, and which I repeated afterwards, particularly in 1770, I saw plainly, as Dr HALLER had done, that both rows of intercostal muscles were in action during inspiration.

AFTER

AFTER proving, that both rows of intercostal muscles conspire in elevating the ribs, I used to point out the fallacy of the demonstrations, by which BAYLE, HAMBERGERUS, and others, have pretended to prove that the internal intercostal muscles depress the ribs. The machine they describe as representing the ribs, vertebræ and sternum, resembles very exactly two wooden rulers, A, B, kept parallel by two pieces of brass, C and D, such as are used for drawing parallel lines; and the two layers of the intercostal muscles are represented by the threads EF and HG, passing obliquely from the one ruler to the other, and decussating each other. See T. 2. fig. 9.

LET C, one of the pieces of brass, represent the vertebræ, and the other piece D the sternum. Let A represent the uppermost rib on the right side of the body, and B the second rib. Let EF represent the external, and HG the internal intercostal muscle.

THEN let C, representing the immoveable vertebræ, be held fast, and let EF be pulled or shortened, they tell us that the second rib B must be more affected than the first, because the lower end of the muscle being at a greater distance from C than the upper end of it, the muscle will act upon the second rib with a longer lever, and therefore that the external intercostal muscles must elevate the rib.

BUT when the internal intercostal muscles, represented by HG, are shortened, they observe that matters will be reversed; and as their origin in the first rib is farther from the vertebræ, or centre of motion, than their insertion in the second rib, that having a longer lever, they must serve to pull the first rib down.

ACCORDINGLY

ACCORDINGLY the rulers, on pulling alternately the threads EF and HG, will be moved alternately upwards and downwards.

BUT to shew the fallacy of this, I need only to add to what has been before observed, that we can perform a full inspiration, without bringing the upper rib, or top of the sternum, upwards or nearer to our head; whereas the pretended demonstration rests entirely on the supposition, that all the ribs, not excepting the first and sternum, have a large play upwards and downwards alternately. Stop the play of the first rib, or suppose it to be fixed in its place, which is the fact, and the boasted demonstration is annihilated.

THE late opinion of SABATIER *, that both rows of intercostal muscles serve for expiration, and that the ribs are elevated by the scaleni and serrati postici superiores, which are fixed to a few only of the upper ribs, scarcely merits a comment. It is refuted by what is above mentioned, and by the want of the external intercostals near to the sternum, and of the internal near to the spine; for intercostals at those places would have served to depress the ribs more powerfully than in any other part of the thorax.

LET us now consider the purpose,

FIRST, Of the obliquity of the fibres in the intercostal muscles, and,

SECONDLY, Of their being disposed in two layers, the fibres of which decussate each other.

IT

* See Anatom. T. 3. p. 465. 7. m.

IT is evident, that the obliquity of the fibres here is not intended to increase their number, or the strength of the muscle, because the fibres would have been more numerous if they had passed directly from the one rib to the other, or had been inserted into the ribs at right angles.

I APPREHEND, therefore, that we are to explain the reasons of the structure in the following manner :

NATURE, in order to give protection to the heart and lungs, has formed the ribs as broad and flat as possible, or left no more space between them than is required for lodging muscles for their motion in respiration. Consistently with this view, as the ribs are fixed at both ends, so that they cannot be moved backwards and forwards, but are confined to motion upwards and downwards, remaining nearly parallel to each other, oblique muscles are preferred to straight ; for if the former can, as I have before demonstrated, perform more extensive motion than the latter, even where both are of the same length, they must have a still greater effect, where the two kinds of muscles are confined between the same parallels.

THUS, suppose the direct distance, or perpendicular drawn from one rib to another, to be represented by three parts, and that the intercostal muscle, in consequence of its obliquity, measures five such parts, and that each of these is capable, when in action, of shortening itself one-fifth part of its length ; it appears from the demonstration, that the oblique muscle can move the rib through a space five times greater than the straight muscle can do.

ON accurate mensuration, I found the length of the intercostal muscle to be one inch and a half, the perpendicular line one inch, and the base about one and one-eighth inch. Hence, calculating on the supposition, that the muscular fibre, in action, shortens itself one-fifth of its length, it will be found, that the intercostal muscles, in consequence of their obliquity, produce a greater motion of the ribs than perpendicular muscles could have done, nearly in the proportion of 35 to 12.

THE only point remaining to be explained, is, why nature hath formed two layers of intercostal muscles decussating each other.

THE purpose of this, I apprehend, is to render the motion of the rib upwards as direct as possible, and to prevent it from being drawn or pressed forwards upon the sternum, or backwards upon the vertebræ, so much as, by its friction, to interrupt the freedom of its motion.

UPON the whole, by the obliquity of the intercostal muscles, the motion of the ribs is very much greater than could have been performed by straight muscles placed between them: At the same time, by their consisting of two layers, or two muscles decussating and balancing each other, the motion of the ribs, upwards and downwards, is as direct, and with as little friction, as if it had been performed by straight or perpendicular muscles.

A P P E N D I X,

In which the Pretension of Dr GILBERT BLANE, that he *first* demonstrated the same Effect to be produced by Oblique Muscles as by Straight Ones with a less proportional Decurtation of Fibres, is proved to be unfounded.

AS Dr GILBERT BLANE, physician in London, who attended my Course of Lectures on Anatomy and Surgery two years, *to wit*, 1769 and 1770, has, in the Croonian Lecture on Muscular Motion, read by him at the Royal Society, November 13. and 20. 1788, and printed in London in 1790, given a geometrical demonstration, that the same effect is produced by oblique muscles, with a less proportional decurtation of fibres, than if the same motion had been performed by a direct power, without acknowledging that he had learned from me any thing upon the subject, I found myself under the disagreeable necessity, after I had treated of it in my usual manner, in my last Course of Lectures, to mention Dr BLANE's publication, and then to read Notes I had written on

the subject, so far back as winter 1759-60, and other notes which happened to be taken from my Lectures, by Mr THOMAS THORBURN in 1770, during the last year Dr BLANE attended my Course; and of which many copies have, since that time, been written and circulated among the students.

DR BLANE, who was informed of this, as I supposed and wished he should be, because his brother was, last winter, one of my pupils, has, since that time, *to wit*, on July 27. 1792, written a letter to Mr BENJAMIN BELL, Surgeon in Edinburgh, on that subject, which Mr BELL thought himself authorised to shew me; and I received it from him on the 21st of August 1792.

THAT what Dr BLANE has published or written on the subject may be fully understood by the Reader, I shall now subjoin the part in question of Dr BLANE's Lecture and Letter, then add the notes I referred to, taken from my Lectures in 1770 by Mr THORBURN, with the attestations of gentlemen distinguished by their genius and learning, who have done me the honour of attending my Courses of Lectures, before, when and since Dr BLANE did so; and I shall conclude with a few Remarks upon the whole.

An

An EXTRACT from THE CROONIAN LECTURE ON MUSCULAR MOTION, read by Dr GILBERT BLANE at the Royal Society, November 13. and 20. 1788, corrected, enlarged and printed at London, 1790, in 4to, pages 55, 56, 57.

“ BUT the advantage, or rather compensation of obliquity, which I mean particularly here to demonstrate, is, that the same effect is produced with a less proportional decurtation of fibres, than if the same motion had been performed by a direct power. BORELLI has estimated geometrically the loss of power from oblique action, but seems to have overlooked this compensation of it, which is not inconsiderable, when we reflect that there is thereby a saving of contraction, and consequently of fatigue. This can be rendered an object of geometrical proof, and I here subjoin a demonstration of it, which I made out, when engaged in the study of anatomy, eighteen years ago.

“ LET the line AB, in the annexed diagram (T. 2. fig. 10.) represent a moveable bone, and the line CD a fixed bone parallel to it. Let FE, perpendicular to these lines, represent a muscle acting in its own direction, and the lines GE, HE represent two muscles acting obliquely, and producing, by a diagonal action, the same effect as the other. If the
“ bone

“ bone AB be brought to the situation ab, by the action of the
 “ muscle FE, the muscle will then be in the situation FK. If
 “ the bone is brought into the same situation by the action of
 “ the muscles GE, HE, these muscles will then be in the si-
 “ tuation GK, HK.

“ THE proposition to be demonstrated is, that the line GK
 “ bears a greater proportion to the line GE, than the line FK
 “ does to the line FE; for FK is to FE as GL is to GE,
 “ (Euc. Elem. B. vi. prop. 2.) and the angle ELK, being less
 “ than a right angle, the angle GLK, which is adjacent to it,
 “ must be greater than a right angle; and the angle GKL be-
 “ ing in the same triangle GLK, must be less than a right
 “ angle. The line GK, therefore, which subtends the greater
 “ angle, is greater than the line GL, subtending the lesser,
 “ and therefore bears a greater proportion to GE. But the
 “ line GL is to GE, as FK is to FE; and therefore GK
 “ bears a greater proportion to GE, than FK does to FE;
 “ that is, the fibres of the muscles acting obliquely, suffer a
 “ less proportional decurtation than those of the muscle act-
 “ ing directly.

“ IT is farther obvious, that the more oblique the action
 “ becomes, the greater saving there will be of contraction;
 “ for in moving the line ab towards CD, the line FK dimi-
 “ nishes in a swifter ratio than the line GK, and when the
 “ former has vanished, the latter is in the situation GF.”

COPY

COPY of a LETTER from Dr GILBERT BLANE, Physician in London, to Mr BENJAMIN BELL, Surgeon in Edinburgh. Dated London, July 27. 1792.

DEAR SIR,

I WAS a good deal hurt at learning, some time ago, that Dr MONRO had made very free with my name in his last Course of Lectures, respecting some Remarks of mine in a Lecture on Muscular motion, before the Royal Society, of which I printed a few copies for my friends. If the Doctor's animadversions were to reach only those to whom I am known, I should not pay much attention to them, as they know me to be incapable of so unworthy a conduct; but as the greater part of his auditors are most probably strangers to me and my character, I cannot entirely overlook this imputation.

THE question is about a matter of fact, which passed one or two and twenty years ago, relative to some remarks on the obliquity of Muscles, which the Doctor alleges I have borrowed from him, without acknowledging it. As I originally owed all my anatomical knowledge to Dr MONRO, there can be no doubt that I was led to consider this subject from that part of the Course which related to the muscles, particularly the intercostals; and, as a Professor in a General Course communicates all that is known on the subject, whether discovered by himself or others, I do not know whether Dr MONRO delivered his remarks on this subject, as his own or not; nor would this make much to the present argument, as it would
be

be equally wrong in me to arrogate the observations of any other.

WHAT I mean to allege in answer, is, 1. That Dr MONRO, during my attendance, never made use of any geometrical demonstration, but only such gross illustrations as were adapted to his mixt audience. 2dly, Dr MONRO's observations were entirely confined to the intercostal muscles, the mechanism of which I have made use of, rather as an exception, than an illustration of my doctrine, the essence of which consists in establishing the universality of oblique motion in muscles, and the advantage of this position in multiplying their number, and saving fatigue, by lessening their proportional decurtation. So far as I know, these remarks are wholly my own. If it should be alleged that they are only an extension or improvement of what was delivered in the Lectures, I am ready to allow this; for, as I have said before, it was from them I derived all my original anatomical knowledge; but if this mode of arguing is adopted, there is an end to all merit and all future research. For what subject is there to which one can turn their thoughts, on which a good deal has not been already said? With regard to the geometrical demonstration, I can adduce a fact which I think cannot fail fully to convince Dr MONRO that it never could be my purpose to defraud him of any right he had to the remarks in question. In summer 1774, there was a small club of students who held weekly meetings at each others rooms, to converse upon medical subjects in Latin, as a preparatory exercise to graduation. I was a member of this club, and one evening I brought with me the mathematical proposition and remarks, which I shewed as my own, and they were canvassed accordingly. I farther remember, that the meeting at which this passed, was held at the apartment of Mr Turner, a gentleman from Yorkshire,

shire, who lodged in the corner of the Lower College, at the house of Mr Innes, who so long and ably assisted Dr MONRO in his dissections. I appeal to any one whether, in an assembly of intelligent students, who had themselves recently attended the Lectures, I could have had the effrontery to pass for my own, what had been a part of the preceding Course of Lectures, of which they were as much masters as myself.

As it would give me great pain to have any altercation with Dr MONRO, to whom I wish to retain that respect which I owe to my old masters, I should be much obliged to you, when you happen to meet the Doctor in consultation or otherwise, if you will bring on this subject; which I think, from what I have said, you can state in such a manner as to satisfy him.

WHEN I first began, in 1759, to deliver, without the assistance of my father, the whole Course of Lectures on Anatomy and Surgery, I had ~~but~~^{very} little time for preparation, as I had been called home on account of my father's indisposition, much sooner than I expected. My notes were therefore very short, on almost every part of my Course. And on this subject I did not, till of late years, collect in one Lecture all the circumstances I have mentioned in the foregoing paper. It appeared to me improper to deliver such a Lecture, before the students had seen the structure of the muscles, and equally improper to delay taking any notice of the effects of the oblique muscles, till I had finished the myology. I therefore used, generally in my third Lecture on myology, after demonstrating the abdominal muscles, to mention the more extensive motion produced by oblique, than by straight muscles. I observed that the oblique muscles of the abdomen concurring, might bend the body farther forwards, or draw farther downwards

D

the

the cartilage ensiformis, than the recti muscles, which were supposed to be, and, at first sight, appeared to be better fitted for that purpose; and that, if the recti muscles had passed from the ensiform cartilage, obliquely outwards, to the tops of the ossa ilia, they would, with the same degree of shortening, have produced a more extensive motion. And sometimes, in order to be better understood, I cut them off from the ossa pubis, and turned their lower ends sideways to the tops of the ossa ilia. Sometimes I added the geometrical and arithmetical demonstration, by drawing with chalk on a black board, fig. 4. and afterwards fig. 6. or converting fig. 4. into fig. 6. by lengthening the perpendiculars. But, generally, I referred these demonstrations till I had shown the intercostals; where not only the obliquity, but decussation of the muscular fibres, fell to be explained.

IN treating various other parts of the myology, and particularly of the muscles on the posterior part of the trunk, I never failed to point out the various effects of the obliquity of the muscles.

WHEN I came to the muscles of the hands and feet, where the obliquity of the fibres, in many of them, is so remarkable, that they have been called penniform, I thought it sufficient to show the structure accurately, and to put them in mind of, without repeating my demonstration.

I SHALL, now, copy verbatim, first, my own notes, on which I lectured in 1759-60; and then those written at different times by other persons.

NOTES,

NOTES, &c.

[The following is a Literal Copy of Notes written by Dr MONRO
in Winter 1759-60.]

1760. LECT. 17. ON LARGE BODY.


Of RESPIRATION.

AFTER describing, in six quarto pages, the motions of the Thorax, and endeavouring to prove that both rows of Intercoastal muscles elevate the ribs, I wrote as follows:

“ The intention of nature in this is very simple and evident. By oblique fibres, a much greater effect is produced than by straight fibres; or two bones may be brought much nearer to each other. By straight fibres you can only approach two bones one third nearer to each other; whereas, by oblique fibres they may be brought close to each other. This is evident by comparing A with C.”*

1760.

* N. B. These three figures are copied exactly by the engraver, from the original manuscript, written in 1760, on the margin of which they are drawn.



1760. Of the Course, v. 8. p. 86, 87.

LECT. 68. of the Course.

REMARKS on the Muscles that move the Head and true
Vertebræ.

Action of each to be Explained.

ALL muscles before and behind, acting at once, pull directly forwards or backwards.


PART of these before and behind, acting at the same time, pull the head to a side, as sterno-cleido-mastoid and splenius with trachelo mastoid; and, as the chief muscles are oblique, the flexions are greater than by straight fibres. Neck moved by above mentioned muscles, with addition of longus colli, transversalis colli, intertransversalis, interspinales, spinales, part of musculus angularis, scaleni, which operate in a similar way, the oblique fibres of severals making the extent of their effect greater.

DURING the years 1762, 1763, 1764, Dr JOHN HAYGARTH, now physician in Chester, attended my lectures, and as physiological as well as practical matters, attracted his notice; I received

ved from him, if I recollect rightly, two letters on the subject in question. And, fortunately, I have found part of the first letter, which I was afraid I had destroyed many years ago with other papers I thought it of no use to preserve. In this fragment, he writes in the following words :

To Dr MONRO, *jun.*

YOUR account of the advantage derived from the united action of both the interco-



stal muscles crossing one another, in procuring a greater extent of motion, at first pleased me excessively, the solution is so extremely simple and beautiful. On further reflection, it appeared to me that this greater extent of motion might be procured from fibres running in one direction, and was owing to the obliquity, not the crossing of the fibres. Let EK and FH represent two bones, AC a muscular fibre connecting them together in an oblique direction. Draw the perpendicular BC. Let AB be two inches long, AC three inches. If the fibre AC contracts one-third of its length, will it not coincide with AB, and bring the points B and C together? Now, if we imagine other oblique fibres EF, GH, will not they—

FROM this fragment it appears that I had demonstrated the great effect of oblique fibres in the intercostal muscles ; but, as I speak in lecturing without notes before me, I either had neglected to explain the reason of there being two layers of intercostal muscles decussating each other ; or the explanation I had given had not been attended to by Dr HAYGARTH.

I ANSWERED this letter, by endeavouring to explain the reason of two such layers, in the way I have done above, and received a second letter from the DOCTOR, expressing his satisfaction with the solution I had given him.

VERY lately I have written to Dr HAYGARTH, and received his answer in the following words :

LETTER from Dr A. MONRO to Dr JOHN HAYGARTH.

DEAR SIR,

As I intend to publish very soon, some Observations on the Muscles, and wish to ascertain the time at which I first demonstrated, that a greater extent of motion can be produced by oblique muscles than by straight ones, I give you this trouble, as I recollect that you paid particular attention to the subject, whilst studying in this University. I have in my manuscript introduced the mention of your name in the following terms :

During the years 1762, &c.

THE favour I beg is, that you will collect from your notes, or recollect from memory what then passed on this subject,


and write it to me as soon as you conveniently can, which will much oblige,

Your most obedient servant,

ALEX. MONRO.

Edinburgh, October 27. 1792.

To Dr John Haygarth.



LETTER from Dr JOHN HAYGARTH to Dr ALEX. MONRO.

DEAR SIR,

Chester, October 31. 1792.

I HAVE fortunately preserved a copy of the letter to which you refer. I attended, and with very high satisfaction, three Courses of your Anatomical Lectures. The letter was dated the 2d of February 1764, during my attendance on the second Course. The fragment you have preserved very exactly corresponds with my copy. "Your account, &c. brought into contact." The remainder of my letter states some difficulties concerning the alterations by which the eye adapts itself to view objects at great and little distances, and concerning scrophulous tumours, as they affect the lymphatic glands.

I WELL remember, that in a very obliging manner, you took the trouble to explain the difficulties I had suggested. A second letter to express my thanks and conviction might very probably be written; but at this distance of time I do not recollect it.

IF my testimony can be of any service to ascertain the time when you demonstrated that a greater extent of motion can be produced by oblique muscles than by straight ones, you are every way well entitled to it. I shall give it with peculiar pleasure.

LETTER from Dr MONRO to Dr HAYGARTH.

DEAR SIR,

I RECEIVED the favour of your answer to my first letter, from which I observe, that I had not been sufficiently explicit in acquainting you that I had lost that part of your letter written in 1764, which followed the extract I sent you.

I AM therefore obliged to trouble you again, to beg,

1. THAT you will send me what followed, to complete the fragment, or that you will send me the sequel of your letter to the words "brought into contact."

2. To

2. To ask whether you recollect the particular demonstrations by mathematical figures and arithmetical calculations, which I then employed in proof of the more extensive motions produced by oblique than by straight muscles.

3. WHETHER you recollect that I applied the demonstrations not only to the intercostal muscles, but to muscles in general, and particularly to the abdominal muscles, the muscles on the back part of the head, and to the penniform muscles of the extremities.

As I have engaged to read a paper on this subject to the Royal Society of this place, you will add much to the obligation, by sending me your answer as soon as you can conveniently.

I am,

Dear Sir,

Your very humble servant,

A. MONRO.

Edinburgh, January 5. 1793.

E

LETTER

LETTER from Dr HAYGARTH to Dr MONRO.

DEAR SIR,

I AM indeed much concerned, that the inaccurate hurry with which my answer to your letter was written, has given you the trouble of writing another. I did not then observe, that the fragment of my former letter in your possession wanted a few words at the conclusion, to render the sense complete.

AFTER "E, F, G, H," add, "will they not bring the
" point I to touch F, and the point K to touch H; and will
" not the two bones be brought into contact?"

ON the farther subject of your enquiry, I have searched with attention the notes taken at your Anatomical Lectures, which I have still preserved, but without success. They are indeed so very imperfect, especially on the anatomy of the muscles, as to afford no decisive evidence on either side of the questions you ask.

NOR, 2^{dly}, At this distance of time can I recollect any demonstrations, or mathematical figures and arithmetical calculations to prove the more extensive motion produced by oblique, than by straight muscles.

3^{dly}, NOR do I remember that you applied such demonstrations to the intercostal, abdominal, occipital, or dorsal muscles, or to the penniform muscles of the extremities.

BUT

BUT I wish particularly to declare, that neither the imperfection of my memory, nor the imperfection of my manuscript, can afford any proof, or any very strong presumption that you did not teach these doctrines in 1762-1765, when I had the pleasure to attend your Lectures; a pleasure which is always recollected with the greatest satisfaction and gratitude, by

Your obliged friend and servant,

JOHN HAYGARTH.

Chester, January 12. 1793.

IN the years 1769 and 1770 Dr GILBERT BLANE attended my Lectures; and in the last of these two years Mr THOMAS THORBURN did so: He likewise attended my Lectures in 1773. About the same period, Mr THORBURN attended several of the other Medical Professors, and wrote down their as well as my Lectures, which, for a number of years afterwards, were copied and sold, or lent out to be copied by the students. And, on the lowest computation, that twenty copies of my Lectures have since that time been written annually by the students, there must be now extant more than four hundred copies of them; for they have been handed down to this time, with fewer corrections and additions than might have been expected.

ON the 8th of November 1774, I purchased a copy of Mr THORBURN's manuscript, written in ten volumes, by Mr JOHN WILSON, who, from being lame, had the conceit of calling himself *Claudero*, and was best known by this name.

He subscribes the last page of the last volume in the following words:

“ F I N I S .

“ Edinburgh, April 29. 1774.

“ CLAUDERO *scripsit.*”

I FOUND that Mr THORBURN, who had no knowledge of Anatomy when he began to write my Lectures, had not attempted to write my demonstrations or descriptions of the parts; neither had he attempted to copy any of those figures which, in this and in many other parts of my Course, I have been in the custom of drawing with chalk upon a black board, in order to render my Lectures more intelligible to the students. In consequence of this, his notes are particularly imperfect wherever the Lecture is connected with figures. Keeping this in view, let us next read what he had written.

MUSCLES of the ABDOMEN.

VOL. II. p. 348. “ With regard to the oblique, their action may be reduced to the greatest simplicity; for the external oblique of the one side has its fibres in the same direction

“ rection with the internal of the other ; so may be considered
 “ as forming one digastric muscle ; and when I draw down
 “ the thorax, or raise the pelvis with a degree of obliquity,
 “ these muscles are employed ; but in other respects their ac-
 “ tion is similar to that of the recti ; for if I act with them
 “ all, the slanting direction of the one balancing that of the
 “ other, the body is moved straight.”

VOL. II. p. 350. “ With regard to the oblique, we find
 “ that the degrees of motion they perform are greater than
 “ could have been made by straight muscles. This is a
 “ circumstance that is entirely overlooked ; yet the influence
 “ of it is very powerful. The obliquity of the one muscle ba-
 “ lances the obliquity of the other, as I have said, when the
 “ body is brought forwards, and here the action is a good deal
 “ more extensive, than it otherwise would have been. Thus
 “ suppose an action to be required by the rectus, the rec-
 “ tus we shall say can shorten itself one-third of its length,
 “ and the thorax would have been drawn so much down ; but
 “ when two such muscles are placed slanting, and the ob-
 “ liquity of the one balances that of the other, the action is
 “ much more extensive. In this part of the body it is not
 “ often that we have occasion for so great an extent of motion,
 “ but between the ribs for instance, where the space left
 “ for motion is very small, this contrivance is very useful
 “ and necessary.”

VOL. II. p. 437. “ The only difficulty here, and which
 “ has puzzled many anatomists, is to account for the obliqui-
 “ ty of the intercostal muscles : And next, why there are
 “ two rows of them ? I apprehend this difficulty entirely
 “ depends upon anatomists not having observed a circum-
 “ stance

“ stance of the utmost moment, and which we noticed before, that wherever extensive motion is required, oblique fibres are employed, the oblique fibres giving a greater play than the straight, bring the two ribs much nearer, and in consequence of there being two rows, the obliquity of the one balances the obliquity of the other, and raises the rib in an equal way, neither pushing it backwards upon the spine, nor drawing it forwards upon the sternum.”

VOL. II. p. 466. “ When we observe in what way the head can be moved, the combined action of the muscles in moving it is very simple. In general, every muscle draws its two points as much as the articulation will allow in a straight line; and all the muscles having slanting fibres twist the head. But if a pair of these is in action at the same time, the obliquity of the one balancing the obliquity of the other, they serve the purpose of a straight muscle. So if I want to draw the occipital bone backwards in a straight line, instead of employing muscles which run very nearly in that direction, we can employ the musculi trapezii, which run in a slanting direction, and the head moves in a diagonal line between them, and by their means we can draw the head much farther back than we could do by a muscle sent directly from the bottom of the neck to the occiput.

“ If I mean to draw the head straight forwards, I employ all the muscles on both sides, whether straight or oblique, and in like manner, when I mean to draw it backwards. But for pulling it sideways, we are obliged to employ an anterior and posterior muscle at the same time. So in order to bring the mastoid process down to the shoulder, I
“ act

“ act with the sterno-mastoideus and the splenius muscle,
 “ which counteract one another as to obliquity, and, by a suc-
 “ cession of these motions, I can make my head describe a
 “ circle.”

VOL. II. p. 470. “ If we mean to raise the scapula, we em-
 “ ploy two muscles which decussate each other, the obliquity
 “ of the one balancing the obliquity of the other, *to wit*, the
 “ upper part of the trapezius and the levator scapulæ. If I
 “ mean to pull it down, to press any thing under my arm, I
 “ employ the lower part of the trapezius; for different por-
 “ tions of a muscle may be employed at different times; and
 “ with it, at the fore part, the serratus anticus major, parti-
 “ cularly the portion of it near to the lower angle, which
 “ move it farther than a straight muscle would have done.
 “ If I mean to bring it backwards, I employ the whole of the
 “ trapezius, but more especially the lower and upper ends,”
 &c.

Mr THOMAS THORBURN, after finishing his studies in this University, settled as a Practitioner of Physic in March, Cambridgeshire, where he died twelve years ago. His son, Mr John Thorburn, who is studying Physic, and has attended my Lectures this and the last winter, is in possession of his father's original manuscript, written in short hand in 1770, which he has extended as accurately as he could, and favoured me with the following exact transcript from it.

LECT.

LECT. 27.—Muscles of the Abdomen.

IN a similar way we explain the action of the oblique muscles, making allowance for the obliquity of their fibres. The external oblique draws down the ribs in a slanting direction, pulls the body, and at the same time turns it over to a side, or twists the body. The internal oblique has a similar effect; nay the external oblique of one side concurs with the internal oblique of the other in the same action, so we may consider them as one digastric muscle, or, they may be considered as a continuation of each other. So the operation is perfectly similar, only, that the right external acts at the same time with the left internal and *vice versa*. Or we may throw all the oblique muscles into action, or a pair, and, the obliquity of the one being balanced by that of the other, we bring the body straight forwards, as by the recti muscles. I would only add, that by the oblique muscles we can perform a greater flexion and motion, than by the recti. The rectus appears better calculated for bending the body forwards, but a much larger motion can be made by the oblique. Take any proportion of contraction, suppose one-third, the rectus muscle acting, the cartilago ensiformis cannot descend lower than this. But, when the oblique muscles act, the one balancing the other, can bring down the cartilago ensiformis nearer to the bones of the pelvis. So they really make a larger motion than straight fibres can do. But in considering the thorax, where the effects of the oblique muscles are more necessary than here, I shall consider this.

LECT. 37.

IT only remains, that we determine the power by which the ribs are moved, and here authors have perplexed themselves very much from not attending to the particular uses of oblique fibres in the intercostal spaces, and from not rightly understanding the use of two rows of intercostal muscles, and the two rows are given because of the obliquity.

I APPREHEND that every difficulty may be easily removed, and one general point of much consequence in our body is explained by it, *viz.* the great effect which oblique muscles have in performing a more extensive motion, than is possible by means of straight fibres.

Now, I don't recollect that this has been explained by any author. I shall draw the abdomen, here is the cartilago eniformis, and here are the ossa pubis,



NEXT these two represent the ossa ilia, and suppose a large flexion of the body is intended, and that I need to draw down the eniform cartilage very near to the ossa pubis. That cannot be done by the muscoli recti, which I next draw. Let us assign any particular length to the recti, that they are 10 inches long, and determine to what degree the common muscles shorten in their action. Suppose one-fifth, or when the recti

F

muscles

muscles act, the cartilago ensiformis is brought down two inches. Now, I next draw two oblique muscles of the exact length that I have supposed the recti, 10 inches long, and I conceive to myself, that these contract in the same proportion, when that is made, where will the ensiform cartilage be brought? It will be brought a great deal lower. When a workman wants to draw a right angle, he makes a triangle, one side of the triangle is eight, and call the other side six, if this angle be right, the long side is exactly 10. Now, if I shorten each of these fibres one-fifth, the obliquity of the one balancing the obliquity of the other, the cartilago ensiformis moves in the diagonal line, and can be brought down opposite the insertion of the muscles, or instead of moving two inches, it may be made to move six, which is three times more than it did formerly. Try the experiment with cords and weights, and you will be convinced that the oblique perform much greater motion. Now, with regard to the intercostal muscles, their large motions depend upon this entirely, yet, the reason has not been given. The space is so small that a straight fibre could not have given sufficient play, hence, there is an obliquity. If there had been one row of oblique fibres only, the ribs would have been drawn either too much forwards or backwards, and the motion would not have been equal. But by throwing in two rows, the greatest motion is performed, and the motion is perfectly equable, the ribs neither moving upon the vertebræ behind, nor the sternum before, but moving parallel to each other. And as the upper ribs are more fixed than the under, when the intercostal muscles are in action, there is a general motion upwards. For, supposing that by any power, the intercostal muscles are thrown into action, the
under

under rib moves towards the upper, and the uppermost of all is fixed by its structure.

LECT. 40. Muscles of the Head.

WE have seen a great number of muscles which are chiefly intended for the motion of the head, but few of them are upon the fore part, because the muscles of the jaws can at pleasure act upon the whole head, and that with great advantage, in consequence of the distance of the jaws from the condyles of the occiput the centre of motion. When we act only with the muscles proper to the head, the different degrees of obliquity of the sterno mastoidei balancing each other, the head is drawn forwards, and so with respect to the muscles behind, and in general, you will observe, that the obliquity extends the motion.

I ACT with the muscles of one side, whilst I bend the head forwards. I turn the face towards that side, and alternate the motion. I give a kind of rotation to the head. In like manner on the back part, I either with the action of all the posterior muscles, draw the head strongly backwards, or acting with the oblique, I give it a rotation; or to take all the motions which we are capable of performing, I bend the head straight forwards by the anterior straight muscles, and back-

wards by the posterior, but at the side I employ an anterior and posterior muscle at the same time. So if I mean to press the ear down upon the shoulders, I employ the sterno mastoideus, and the splenius suppose of the same side, the obliquity of the one, balancing the obliquity of the other, the head is moved in a diagonal between them, and by a succession of these, I can make my head describe a circle.

NEXT let us attend to the motions of the scapula. Authors before WINSLOW, used to speak of the motions of the scapula, upwards, downwards, backwards and forwards. WINSLOW has added to these a rotation of the scapula, and fond of this observation, he carries things too far, rejecting the direct motions. Nay, farther, he errs with respect to the manner of its rotation, conceiving it to roll upon an imaginary axis put through the middle. But from its connection to the clavicle, it is impossible it should do this. We cause the scapula to roll upon the end of the clavicle, which is its centre of motion. Next we roll the clavicle itself upon the top of the sternum, but that I can perform the motions upwards, downwards, backwards and forwards, is so evident, that we all experience it, and can point out the muscles fitted to perform it. First, the trapezius muscle, and joined to this the levator scapulæ are fitted for making the motions of the scapula upwards, the obliquity of the one balancing that of the other, and preventing the rotation. Next, the whole trapezius muscle being at once thrown into action, draws the basis of the scapula towards the spine of the body. If again, I mean to make a motion of the scapula downwards, here are two muscles in our view perfectly fitted for our purpose, the lower share of the trapezius and the serratus anticus major; and to pull the scapula directly forwards,

forwards, I slacken the trapezius, and act with the ferratus anticus major, and balance it with the ferratus minor or lesser pectoral, to pull forward the coracoid process; and by a succession of these, I can move the scapula in a circle.

TRANSCRIBED from the original manuscript lectures, taken in short hand by Mr THOMAS THORBURN in 1770, by me, his son.

JOHN THORBURN.

Edinburgh, December 27. 1792.

As I understood that Mr BENJAMIN BELL and Mr JAMES RUSSELL, Surgeons in Edinburgh, had purchased copies of notes taken from my Lectures, without knowing by whom they were written, I asked the favour of them to furnish me with extracts from their manuscripts on the subjects in question. Mr RUSSELL purchased his copy in the year 1775 or 1776, and Mr BELL has had his copy about six years in his possession.

possession. I found they were both written in the same, following words; but, not being copied by the same hands, the pages of the manuscript are differently numbered.

PAGE $\frac{2}{2} \frac{8}{2} \frac{2}{4}$. I would only add, that by the oblique muscles we can perform a greater flexion and motion than by the recti. Take any proportion of the contraction you will, suppose a third, when the rectus muscle acts, the cartilago enfiformis cannot descend towards the os pubis more than one third; but when the oblique muscles act, the one balancing the other, we can bring the cartilago enfiformis nearer to the bones of the pelvis in a greater proportion.

PAGE $\frac{3}{2} \frac{4}{6} \frac{5}{8}$. It only remains, that we determine the power by which the ribs are moved. And here Authors have perplexed themselves very much, from not attending to the particular use of oblique fibres in the intercostal spaces, and from not rightly understanding the use of two rows of intercostal muscles; and the two rows are given because of the obliquity. I apprehend that every difficulty can be removed with the greatest ease, and one general point of much consequence in our body is explained by it, *viz.* the great effect which oblique muscles have in performing a more extensive motion than is possible by means of straight fibres. Thus, suppose a large flexion of the body is intended, and that I want to draw down the enfiform cartilage very near to the ossa pubis, that can very well be done by the muscle recti. Suppose that these are ten inches long, and that they shorten one fifth, so that when the recti act, the cartilago enfiformis is brought down two inches; but suppose we use two oblique muscles of the same length, and that they contract in the same manner, the enfiform cartilage will now be brought a great deal lower,
the

the obliquity of the one balancing the obliquity of the other, the cartilage moves in the diagonal line between them; and instead of moving two inches only, it will be made to move six, that is three times more than it did formerly.

Now, with respect to the intercostal muscles, their large motion depends upon this entirely, the spaces are so small that a straight fibre could not have given a sufficient play, hence there is an obliquity. Further, if there had been only one row of oblique fibres, the ribs would have been moved either too much forwards or backwards, the motion would not have been equable; but by throwing in two rows, the extent of the motion is preserved, and the motion is perfectly equable, the ribs moving parallel to each other; and as the upper ribs are more fixed than the under, when the intercostal muscles are in action, there is a general motion upwards, the under ribs move towards the upper, and from top to bottom they move in this uniform manner; so that Dr BOERHAAVE was mistaken in supposing that the under ribs were pulled downwards, while the motion of the upper ribs was directed upwards. They all move in the same direction.

At the anterior parts the external intercostals are wanting, and the internal are wanting behind, and the reason of this is perfectly simple.

If the external had been continued to the sternum, the last fibres would have been fixed to a less moveable bone below, and to a rib above; therefore, instead of assisting in raising the ribs, they would have pulled them down, and hence as soon as the intercostals approach so near, that from this obliquity the fibre would have been in danger of influencing the
upper

upper rib more than the under, they are laid aside, they are carried no farther.

IN the internal intercostal muscles, that proportion is reversed. In consequence of their direction and the angle which the ribs make with the vertebræ, they would have come at length to influence the upper rib more than the under, when they would not only be useless, but even hurtful; and from this view, we are enabled to refute an error that has crept in, that the external and internal intercostals are antagonists to each other, that the external elevate the ribs, while the internal depress them. Whereas it appears from this, that they are intended to co-operate; and, without using further arguments, I have seen the co-operation in living animals, and it has been long ago very justly described by Dr HALLER.

Attested by

BENJ. BELL.

EXTRACT

EXTRACT from the MINUTES of the PHYSICAL CLASS of
the ROYAL SOCIETY of Edinburgh.

Edinburgh, 7th January 1793.

Dr MONRO, after having read a paper, entitled, *Observations on the Muscles, and particularly on their Oblique Fibres*, requested of the Society that a committee might be appointed to examine into the dates of certain discoveries or improvements contained in the above paper. The Society accordingly appointed the following members, as a committee for that purpose, viz. Dr GREGORY, Dr RUTHERFORD, Dr DUNCAN, Dr THOMAS SPENS, Mr BENJAMIN BELL, Mr JAMES RUSSELL, Mr Professor STEWART, Mr Professor GREENFIELD, Mr Professor PLAYFAIR. And the Society farther instructed this committee to examine into the dates above mentioned, and to take such measures for ascertaining them as they should judge proper, and to report to the Society.

Extracted from the Minutes of the Royal Society,

By JOHN PLAYFAIR, *Sec. Phys. Clk.*

G

R E P O R T,

REPORT, &c. of the COMMITTEE.

Edinburgh,

The Committee referred to in the above extract met, and had laid before them, by Dr MONRO, a manuscript, being an appendix to his paper, read in the Royal Society, January 7. 1793, in which were contained the following pieces :

A LITERAL copy from original notes of a Lecture in 1760, on Respiration, p. 26, 27. App.

FRAGMENT of a letter from Dr HAYGARTH, Physician in Chester, to Dr MONRO *junior*.

ANOTHER letter from Dr HAYGARTH to Dr MONRO, dated January 12. 1793.

Also passages copied from a manuscript of Dr MONRO's Lectures, by THOMAS THORBURN, the copy of which is dated April 29. 1774. The passages being from the manuscript, Vol. II. p. 348. 437. 466. &c.

Also from the manuscript of the same Mr THORBURN, copied by his son Mr JOHN THORBURN, Lect. 27.—37.—40.

THE

THE Committee having compared all the preceding extracts with the original papers, found them faithfully and correctly copied from them. In evidence of which they have signed this attestation.

D. RUTHERFORD.

J. GREGORY.

AND. DUNCAN.

WM. GREENFIELD.

JAMES RUSSELL.

BENJ. BELL.

TH. SPENS.

DUGALD STEWART.

I SHALL next subjoin a letter I wrote to the following gentlemen, with their answers *.

Mr *Dugald Stewart*, formerly Professor of Mathematics, and at present Professor of Moral Philosophy in the University of Edinburgh.

Dr *Daniel Rutherford*, Professor of Botany in Edinburgh.

Dr *Andrew Duncan*, Professor of the Institutions of Medicine in Edinburgh.

Dr

* Dr *Stewart* attended my Course of Lectures in 1770.

Dr *Rutherford* in 1764, 1765, 1766, 1767.

Dr *Duncan* in 1764, 1766.

Dr *Gregory* in 1768, 1771, 1772.

Mr *Bell* in 1766, 1767, 1768, 1770.

Mr *Russell* in 1771, 1772, 1774, 1775.

Dr *James Home* in 1777, 1778, 1779.

Dr *Hope* in 1783, 1784, 1785, 1786.

Mr *Greenfield* in 1790.

Dr *James Gregory*, Professor of the Practice of Physic in Edinburgh.

Mr *Benjamin Bell* and Mr *James Russell*, Surgeons in Edinburgh.

Dr *James Home*, Physician in Edinburgh.

Dr *Thomas Hope*, Professor of Medicine in the University of Glasgow.

The Reverend Mr *William Greenfield*, Professor of Rhetoric and Belles Lettres in the University of Edinburgh, and at present conjoined with Professor *Robison* in teaching the Class of Natural Philosophy.

S I R,

As I intend to publish very soon some observations on the muscles, and wish to ascertain the time at which I first demonstrated, that a greater extent of motion can be produced by oblique than by straight muscles, I hope you will excuse me for asking the favour of you to answer the following questions :

1. Do you recollect, that when you did me the honour of attending my Lectures, I endeavoured to demonstrate by mathematical figures drawn with chalk upon a board, and by arithmetical calculations, that oblique muscles perform more extensive motions than straight muscles contracted in the same proportion?

2. Do you recollect the particular diagrams or calculations I then employed?

3. Do you recollect that I dissected the two layers of the intercostal muscles in such a manner as to resemble the mathematical figures I had drawn with chalk on the board, and that I applied my demonstrations not only to the intercostal muscles, but to muscles in general, and particularly to the abdominal muscles, the muscles of the back part of the head and trunk of the body, and to the penniform muscles of the extremities, when these several classes of muscles were demonstrated?

4. Do you recollect, that when I explained the action of the abdominal muscles, and compared the action of the oblique muscles with that of the recti, I observed, that if the lower ends of the recti had been fixed to the ossa ilia, instead of the ossa pubis, they would have made a more extensive motion of the cartilago ensiformis, or would have drawn it

lower down or nearer to the ossa pubis, than is possible for them to do in their real situation?

I am, Sir,

Your most humble servant,

ALEX. MONRO.

Edinburgh, February 11. 1793.

To this Letter I received the following Answers.

DEAR SIR,

I was so very young when I attended your Lectures, and my attention has been so much occupied since that time with other studies, that, notwithstanding the great pleasure with which I heard them, I retain a very indistinct remembrance of them at present. I am not able to say any thing satisfactory in answer to the queries you propose. But I recollect, in general, that you remarked, the greater extent of motion produced by oblique than by straight muscles contracted in the same proportion; an observation which might probably have escaped my memory, with many other parts of your Course, if it had not led you to take notice of some very beautiful circumstances in the mechanism of the body. I recollect likewise,

likewise, that you illustrated your ideas on the subject by means of a diagram; but at this distance of time, I have perfectly forgotten your mode of demonstration.

I am, Dear Sir,

Your most obedient humble servant,

DUGALD STEWART.

Argyle Square, 17th February 1793.

DEAR SIR,

I received the letter addressed to me, in which you propose some questions respecting the doctrine you delivered of the action of oblique muscles, and the advantages obtained by the oblique direction of their fibres, when I had the honour of attending your Anatomical Lectures. So many years have elapsed since that time, that many particulars relating to what then passed have slipped from my memory, and I derive but little assistance, upon the present occasion, from the Notes of your Lectures I have preserved, they happen to be so concise on the point in question. What I do clearly recollect to remembrance, I shall freely communicate to you. I recollect perfectly, that you took great pains to demonstrate the direction

tion of the fibres, and also to point out the extent of both the external and internal intercostal muscles; and that afterwards, when treating of the action of these muscles, you observed, that not only both of them served to elevate the ribs, but also that, by reason of the obliquity of their fibres, they must cause a much greater extent of motion of the ribs, by the same proportional contraction, than they could have done had their fibres run directly and perpendicularly from one rib to another.

The diagrams and calculations by which you established the latter observation, I do not precisely recollect; to me, however, I know they were quite satisfactory. One very simple illustration I well remember, to wit, your placing two blow-pipes betwixt two parallel lines, so that their points meeting together, formed an angle at one of the parallels, while the other ends reached to and were kept constantly at the same distance from each other on the other parallel; then gradually approaching the points of this second parallel till they coincided with it. By the successive points of intersection of the blow-pipes in this progress, you remarked might be denoted the gradual contraction of oblique muscular fibres, represented by the blow-pipes, and the motion of any part attached to them corresponding to this contraction; and by the distance betwixt the points of the blow-pipes, as these overlapped each other when coinciding with the parallel, was denoted twice the quantity by which such fibres must have contracted, in order to bring the two parallels together, or to bring bones situated like them into contact, while a fibre placed perpendicularly betwixt them, to effect the same extent of motion, must be supposed to have contracted by a quantity equal to its whole length, or to have been reduced to a

H

mere

mere point. This illustration appeared to me so clear and decisive, that nothing could add to the conviction it conveyed; I was then much more accustomed to geometrical than to anatomical demonstrations, and it at once recalled to me some of the known properties of triangles, and particularly the relative diminution of the secant and tangent, as an angle is diminished. I was so delighted with the beautiful mechanism of the parts I had seen, and with the ingenious explanation you had given of their action, that I well recollect I soon thereafter made a small model to imitate their motion. It consisted of an upright pillar, from which were extended two horizontal arms, one over the other, the lowermost of which was moveable. Threads were attached to this arm, and *were passed through small rings on the uppermost arm*, having such obliquity as the fibres of the different layers of the intercostal muscles; by pulling or shortening the threads, the lowermost arm was drawn upwards, precisely as the ribs are by the contraction of the intercostal muscles.

I do not recollect to what other particular muscles you extended the doctrine above mentioned, but I understood, that you considered it as general, and consequently applicable more or less strictly to every oblique muscle whatever.

I am, with much regard,

Dear Sir,

Your most humble servant,

D. RUTHERFORD.

Edinburgh, February 14. 1793.

DEAR

DEAR SIR,

I received your's yesterday, and am very sorry that my recollection does not enable me to give any positive or precise answer to your questions. I attended your Lectures during the winter sessions 1764-5, and 1766-7. Ever since that period I have been satisfied, that oblique muscles perform more extensive motions than straight ones contracted in the same proportion. But I do not particularly recollect the demonstration you gave of it at that time. Nor do I find any account of it in the short notes which I took from your Lectures. In these, however, I find the following paragraph: "HALLER
 " imagines, that the reason why the intercostals are placed
 " obliquely is, that there may be room for the insertion of a
 " greater number of muscular fibres; but when we reflect,
 " that had they been placed perpendicularly, the ribs could
 " not have been brought by far so near one another, we see
 " a much more satisfactory reason for it."

I remain your's sincerely,

ANDREW DUNCAN.

Edinburgh, 13th February 1793.

St John's Street, Wednesday Evening, 13th February 1793.

DEAR SIR,

I AM very sorry that I have neither any notes of your Lectures when I was your pupil, nor yet such particular remembrance of your doctrine about the purpose and the advantage of oblique muscles, as to enable me to answer, in a satisfactory manner, the queries that you have put to me.

I REMEMBER perfectly the general principle, and the application and illustration of it in the case of the two layers of intercostal muscles; but the more full illustration of it by diagrams, arithmetical calculations, and geometrical reasoning, such as I had the pleasure of hearing from you at the Royal Society last month, had escaped my memory. Perhaps I had attended to them less than I should have done, and forgot them the more readily, because the fundamental proposition is so nearly self-evident, that the very slightest illustration, or even a clear and precise enunciation of it, such as I am sure you would not fail to give, must have convinced me at once of its truth. Observe, I attended your Lectures three different years (if I remember right) between 1768 and 1773, both inclusive.

Your's most truly,

J. GREGORY.

Mr

Mr BENJAMIN BELL, in answer to my first question, answers, That he does ; but in answer to my three questions, says he does not recollect.

DEAR SIR, *Edinburgh, 22d February 1793.*

I WAS lately favoured with a letter from you, requesting me to recollect the account you gave in your Lectures, when I had the honour of attending them, of the advantages procured by the oblique direction of muscular fibres. The subject I well remember was a very favourite one with you, and that you insisted at great length upon the explanation of certain doctrines which you considered as peculiar to yourself. I remember too, that you endeavoured to illustrate your opinions by the delineation of some figures upon a board. But I cannot, at this distance of time, charge my memory with the distinct recollection of what these figures were, neither can I be positive with regard to the nature of these peculiar doctrines. I recollect, however, that certain doubts occurred to me at the time, which I suggested to you in conversation, and which, I think, you then resolved to my satisfaction.

OF the second and third queries I remember nothing distinctly.

RESPECTING

RESPECTING the fourth query likewise, my recollection is not very accurate, though I have some faint remembrance of your attempting to prove that the recti muscles, if their lower end, in place of being inserted in the os pubis, had been inserted into the spines of the ossa ilia, would have had more effect in drawing the sternum down to the pelvis.

THESE are all the circumstances I now recollect about the subjects stated in your queries to me; and although the answer be deficient in many respects, I am yet confident, that, so far as they go, they give a fair statement of the substance of your Lectures.

I am,

Dear Sir,

Your most obedient and humble servant,

JAMES RUSSELL.

DEAR

2

DEAR SIR, *Edinburgh, April 23. 1793.*

I RECEIVED your letter some time ago, and would have answered your queries sooner, but I wished to consult the notes which I took of your Lectures. These I have not as yet been able to find. What I am, therefore, about to state is from memory.

WHEN I had the pleasure of attending your Lectures in the years 1777, 1778, 1779, while treating of the motions of muscles, you stated the common idea of the advantage of oblique muscles over straight muscles, *viz.* that by the obliquity of their direction, room was given for the insertion of a greater number of muscular fibres, and that this increased number of fibres more than counterbalanced the power lost by the obliquity of their direction. But you observed, that there was still a further advantage derived from the obliquity of muscular fibres: that, when two muscles were placed betwixt two parallel lines, the fibres of one of which were oblique, and those of the other perpendicular to these two parallel lines, and both of these muscles contracting in the same proportion to their lengths, that then the quantity of motion performed by the former was greater than that performed by the latter: Or, that the quantity of motion performed by these two kinds of muscles being equal, the degree of contraction of the oblique muscle was less than that of the straight muscle.

You illustrated the above proposition by geometrical figures and a simple arithmetical calculation. But, at this distance
of

of time, it is impossible for me to recollect the particular diagrams or arithmetical figures.

You applied this general theorem to explain the action of the oblique muscles over the whole body; but more especially, you called the attention of your hearers to it, when you treated of the muscles of the abdomen and thorax.

WITH regard to the muscles of the abdomen, you compared the action of the obliquus externus and internus with that of the rectus, and shewed that the two first acting together, performed in the diagonal a motion greater than was actually performed by the rectus.

WHEN treating of respiration, you agreed with HALLER and other physiologists, that the action of the external and internal intercostal muscles was not antagonistical, as had been supposed by many, the one being employed in elevating, and the other in depressing the ribs; but that both of them, acting at the same time, elevated the ribs, and were therefore muscles of inspiration. Besides this, you delivered as an idea peculiar to yourself, that by the obliquity of the fibres of the intercostal muscles, a much greater quantity of motion was performed, than if perpendicular fibres had been used between the same distance of the ribs. Consequently, if their action had been employed, a greater length of muscle must have been necessary to perform the same quantity of motion; and thus a less quantity of bony surface would have been presented as a shield to defend from injuries the important organs which the thorax contains. This you explained, by dissecting the intercostal muscles, and presenting different views of them. You measured them, and the distance of the ribs, and
applying

applying the above proposition to the actual measurement, you shewed the advantage of the present structure. And lastly, you thought, that by employing two oblique muscles instead of one, the ribs were more directly elevated than they would have been by one set of oblique fibres.

WHETHER you applied the above proposition to explain the action and advantages of the penniform muscles, or of the muscles on the back of the head and of the trunk of the body, I cannot so positively assert; but I rather think that you did, at least the idea was not new to me when I lately heard it applied to these particular muscles. I am confident, however, that I cannot be mistaken in what I have stated above, as ideas delivered in your Course of Lectures, when I had the honour of attending them, since at that time they made a strong impression on me, as a most beautiful piece of mechanism in the animal body, by which the greatest quantity of motion was performed by the least power.

I have the honour to be,

S I R,

Your most obedient humble servant,

JAMES HOME.

I

S I R,

S I R,

A FEW days ago I had the honour of receiving your letter, containing several queries. In reply, I have to acquaint you, that I recollect, that, in your Lectures, you demonstrated by mathematical figures, drawn with chalk upon a board, that oblique muscles perform more extensive motions than straight ones contracted in the same proportion; and that you employed arithmetical calculations to estimate the difference.

To the best of my recollection, the diagram you figured was an isosceles triangle, having a line drawn from the apex perpendicular to the base. In this figure two sides of the triangle represented oblique muscles, and the perpendicular drawn from the angle formed by them represented a straight one. At same time, you showed, by dividing the two sides and perpendicular into so many equal portions, and drawing lines from the angles at the base to the perpendicular, that, a decurtation of oblique muscles, suppose of one-fifth part of their length, would cause the apex, which is to be considered as a moveable point, to approach nearer to the base, in the direction of their diagonal, than would be occasioned by a proportional contraction, that is in the present supposition, by a contraction of one-fifth of the length of a straight muscle. From my memory, as well as from the notes I took from some parts of your Lectures, I am certain of your application of these doctrines, to the oblique muscles constituting the two layers of the intercostals. From the latter, I find, you computed that the intercostal muscles, in consequence of their obliquity, produce by their action, nearly three times as much motion of the ribs, as would take place,
were

were they straight and contracted in the same ratio. My memory does not enable me to speak so precisely and decidedly respecting the application of them, to the other oblique muscles of the body. I have no notes from that part of your Course of Lectures in which you treated of their action.

I am,

S I R,

Your obedient servant,

THOMAS CHA. HOPE.

College, Glasgow, February 1793.


P. S. I must apologise to you for being so long in acknowledging and answering your letter. I assure you I have seized the first moment of leisure for writing.

T. C. H.

S I R,

I AM favoured with your letter, in which you propose to me some questions respecting the account which you delivered in your Lectures of the beautiful fact in Physiology, that oblique muscles perform more extensive motions than straight ones which are contracted in the same proportion.


I REMEMBER in general, that you shewed the truth of the proposition with great distinctness, both by means of geometrical figures, and of arithmetical calculations. To the best of my recollection, your account of it was to the following purpose:



AB (see Tab. 2. fig. 11.) represents a muscle, the extremity A is considered as fixed; but the other extremity B, when the muscle is contracted, moves in the straight line BC, to which AB is oblique. From A the fixed extremity of the oblique muscle, let AC be drawn perpendicular to BC; and let the other extremity by means of the contraction, be removed from B to D. In performing this motion, the oblique muscle AB is not only less shortened in proportion to its length, than a straight muscle, such as CB would have been, but it is in fact shortened by a smaller quantity than CB would have been; or the difference between AB and AD is less than BD, and even when another oblique muscle, as *aB*, conspires to produce the motion, the contractions of both the oblique muscles taken together will not only be less in proportion to their lengths than the contraction of CB in proportion to its length; but farther, if AB and *aB* are each of them longer than twice CB, the contractions of both of them together will be less than BD. The figure you employed was, I believe, such as

I have drawn; and if I am not mistaken, you shewed the truth of the proposition by different arithmetical calculations in the following manner: You first took three numbers representing the hypotenuse, and the sides of the right angled triangle ABC; and then supposing the side BC to be diminished by a certain quantity, you computed what would be the length of the hypotenuse.

I CANNOT pretend to recollect the general demonstration which you gave of the proposition. The following one readily occurs, and may possibly be the same that I heard from you, though it is more probable that your's was a simpler and a better one.



ROUND A as a centre, (see Tab. 4. fig. 13.) with AD as a radius, describe the arch DE meeting AB in E; and draw DF meeting AB at right angles in F. The contraction of the oblique muscle is BE; this is always less than BF; BF is always less than BD; and therefore, *a fortiori*, BE is less than BD. But more particularly, the triangle BFD, ACB, are similar; for the angles at C and F are right angles, and the angle ABC is common to both the triangles; therefore $BD : BF :: AB : BC$.

So that in performing the same motion the contraction of the straight muscle would exceed the contraction of the oblique, in a greater proportion than that in which the length of the oblique muscle exceeds the length of the straight one. So that if AB were double of CB, the contraction of the straight muscle would be more than double the contraction of the oblique. In very small contractions, however, the contraction of the straight muscle would be to that of the oblique in the proportion of AB to BC *sensibly*; for if BE is very small in comparison of AB, BF and BE will be *sensibly* equal.

As my knowledge of physiology is extremely slight, I cannot pretend to answer the remaining questions. I remember, however, that I was much delighted with your application of the general proposition to the different oblique muscles, and particularly to the intercostals. Indeed I received very great satisfaction from all that part of your Course which I was so fortunate as to attend; and I often regretted that I had not more leisure to impress your valuable Lectures on my mind by private study.

I have the honour to be, Sir,

Your most obedient humble servant,

WM. GREENFIELD.

Castlehill,
Saturday, 19th February 1793.

Edinburgh,

Edinburgh, October 31. 1793.

THIS evening, after the press was so far advanced, I received from Mr JOHN BELL, surgeon in Edinburgh, a work he, this day, published on the Anatomy of the Bones, Muscles and Joints, in which, page 289. he has, with much candour, done me justice on the subject in question, by printing the following note :

“ I REMEMBER many years ago, to have heard Dr MONRO explain the office of the intercostal muscles by a diagram, deducing from that argument, the more powerful effect of all muscles having oblique fibres.”

Mr JOHN BELL attended my Lectures in the years 1775 and 1776.

ALEX. MONRO.

I SHALL now conclude with the few following remarks :

1. Dr BLANE now says, That there can be no doubt he was led to consider this subject from that part of my Course which related to the muscles, and particularly the intercostals. Why then did he not acknowledge this in his Croonian Lecture?

2. Dr

2. Dr BLANE next ventures to assert, that during his attendance, he (Dr MONRO) never made use of any geometrical demonstration, but only such gross illustrations, as were suited to his mixed audience. On this I would make the following observations: The *first* is, That from the facts already proved, it appears, that I demonstrated this matter, first geometrically, founding on the propositions of Euclid; and, then, arithmetically, in order to make my demonstration more convincing: 2^d, Dr BLANE has thought proper to use the terms Gross Illustrations, of which it is not in his power to give the specific meaning: because, nothing but geometrical demonstration could have led me to perceive the truth of this proposition, or enabled me to prove it to others. And, it appears, that from these illustrations, Dr BLANE was so much convinced of the truth of the doctrine, as to try whether he could hit upon some other mode of proving its truth.

3. Dr BLANE affirms, that my (Dr MONRO's) observations were confined to the intercostal muscles, which the Reader has found to be contrary to fact; for the application of this doctrine was made when I treated of the abdominal muscles; the intercostal muscles; the muscles of the whole trunk; the penniform muscles of the extremities; particular muscles; and so often that there was more danger of my tiring the students with it, than of not inculcating it sufficiently. How all this should have escaped Dr BLANE, during his two years attendance, is truly surprising; especially as he tells us, that he was, at the time, led to consider this subject from that part of my Course which related to the muscles.

4. To support his pretensions to the first discovery of a Geometrical Demonstration, he tells us how well his demonstration

stration was received in a club of students. Some Readers will, perhaps, think it sufficient for me to observe upon this, that these young gentlemen must have been as inattentive as himself: but the real fact may have been, that although Dr BLANE copied very nearly the figure I always employed in my demonstration; yet, by adding to it a number of A's and B's and C's, affecting the style of Euclid, he gave it the appearance of originality. But I appeal to the Reader, whether my demonstration is not more simple and intelligible to a beginner than his is, and equally convincing to a mathematician.

5. NOT contented with asserting what has been disproved, that my observations were entirely confined to the intercostal muscles, he adds, the mechanism of which he has made use of, rather as an exception, than illustration of his doctrine.

IF Dr BLANE could have made good the propriety of this exception, and at the same time persuaded his reader that my gross illustrations, or even demonstrations, were applied to the intercostal muscles only, that is, were misapplied; his remarks would have been, as he phrases it, wholly his own.

BUT, on this point, the Doctor has been singularly unfortunate: for, as the number of the fibres of the intercostal muscles or their strength is lessened, instead of being increased, by their obliquity; it is evident that their obliquity could serve the purpose only of rendering the motion of the ribs more extensive than could have been performed by straight or perpendicular fibres.

H


6. Dr

6. Dr BLANE says, if it should be alleged that his remarks are only an extension or improvement of what was delivered in my Lectures, he is ready to allow this.

BUT I conclude by observing, that Dr BLANE's modification of my Demonstrations is so far from being an extension or improvement of them, that it only shows that oblique muscles can perform a more extensive motion than straight ones placed between the same parallels; whereas I proved, not only that oblique muscles can do so, but that, with the same degree of contraction, they can perform a more extensive motion than straight muscles of equal length.


S I N I S.

Fig. 1.




Vastus medialis

Fig. 2.



Rectus femoris

Fig. 3.



Sartorius

Dr. Beaune says, if it should be alleged that his remarks are only an extension or improvement of what was delivered in my Lectures, he is ready to allow this.

But, to conclude by observing, that Dr. Beaune's modification of my Demonstration is in fact from being an extension or improvement of them, that it only shows that oblique muscles can perform a more extensive motion than straight muscles placed between the same parallels; whereas I proved, that oblique muscles can do so, but that, with the same degree of contraction, they can perform a more extensive motion than straight muscles of equal length.

TABLE II









Fig. 2.



Fig. 4.




Fig. 3.



Fig. 9.




Fig. 5.




Fig. 6.




Fig. 7.



Fig. 8.



TAB. IV



D E S C R I P T I O N

OF A

HUMAN MALE MONSTER,

ILLUSTRATED BY TABLES,

WITH REMARKS;

FROM THE

T R A N S A C T I O N S

OF THE

ROYAL SOCIETY OF EDINBURGH.

DESCRIPTION

OF

HUMAN MADE MONUMENTS

ILLUSTRATED BY TABLES

WITH REMARKS

FROM THE

TRANSACTIONS

OF THE

ROYAL SOCIETY OF EDINBURGH

DESCRIPTION of a HUMAN MALE MONSTER, *illustrated*
by *Tables, with Remarks.* By ALEXANDER MONRO, M. D.
F. R. S. EDIN. *Fellow of the Royal College of Physicians,*
Professor of Medicine, Anatomy and Surgery in the University
of Edinburgh, Fellow of the Royal Academy of Surgery in
Paris, &c. &c.

[Read Nov. 6. 1792.]

THIS monster, of which the mother was delivered by Mr THOMAS ANDERSON, surgeon in Leith, after the birth of a complete child at the full time, had its proper membranes and a placenta, with a short umbilical cord.

THE following parts were wanting in it; to wit, the bones of the head; the brain, with the organs of sight, hearing, smell and taste; the neck; about one half of the ribs; the larynx, trachea and lungs; the heart; the pharynx, œsophagus and stomach, with all the small intestines, except the end of the ilium; the anus; the liver, spleen, pancreas and omenta; the renal glands; terminations of the ureters; the middle part of the urethra; the right testicle; both arms; both patellæ; with several of the bones of the feet and toes.

A ROUND opening (see fig. 1. and 2.) which led to a thimble-like cavity, shut at its bottom, had some distant resemblance to the mouth.

THE

THE soft parts of the trunk were supported by sixteen vertebræ, six ribs, an os sacrum, and two ossa innominata. The legs had each an os femoris, tibia and fibula, with an imperfect number of the bones of the feet. See fig. 2. X. and fig. 4. I. &c. to 16. 17.

THE umbilical cord was connected at nearly the usual height above the ossa pubis. See fig. 1. E.

THE penis, covered with a large preputium, had the usual situation and structure. See fig. 1. F.

THE lower part of the trunk contained an intestinal tube, shut at its beginning, and composed of an upper part, four inches long, resembling the end of the ilium; for it terminated in the side of an intestine, resembling the caput coli, with its appendix vermiformis. From this place, to its lower end, the great intestine measured thirteen inches; and the end of the rectum, which was much contracted, terminated in the back part of the bladder of urine, above its sphincter. The rectum contained viscid semipellucid mucus, but no black stuff, like the meconium. See fig. 2. O. P. Q. R. S. T. U. V. and fig. 3. O. P.

IN the mesentery and mesocolon, there were about a dozen conglobated lymphatic glands, of the usual shape, colour and consistence. See fig. 2. From which it appeared, that the intestines were provided with lacteal vessels; and we therefore cannot doubt, that the other parts of the body were furnished with lymphatic vessels, or that there was an absorbent, as well as circulating system in this monster.

AT the upper part of the trunk, covered by the ribs, there were two kidneys of a large size, with a pelvis and ureter to each. The right ureter was dilated to the size of a goose's quill. The left one was small. Both were shut at their under ends, and had no communication with a small sac, which, in situation and structure, resembled the bladder of urine, and had

had an urachus coming from it. See fig. 3. W. W. Y. and fig. 2. W. X. Y.

THERE was only one testis, situated in the usual manner, on the left side. See fig. 2. Z.

THE prostate gland surrounded, as usual, the neck of the bladder. See fig. 4. X.

THE urethra, which was the common passage for the fœces, as well as for the feminal liquors, and that of the sac resembling the vesica urinaria, was wanting from within an inch of the vesica to within an inch of the extremity of the penis. See fig. 4. V. Y. and fig. 3. F. G.

THE spinal marrow was of a conical shape, with the top or small part of the cone at its upper end, and at its lower end it formed a cauda equina. From its two ends and sides, it sent off eighteen pairs of nerves; which, at their origin and in their progress, were nearly as large as they are in a perfect fœtus, or where the brain and cerebellum are connected with the spinal marrow. See fig. 4. 1. *Sc.* to 16. *n. n.*

THE umbilical cord was nearly proportioned to the bulk of the monster; and, at the umbilicus, consisted of one vein and two arteries, within which I found red blood. The vein was more capacious than both arteries conjoined; and, as soon as it entered the abdomen, was divided into various branches, which were dispersed upon all parts of the body. See fig. 3. *a, b, c, d, e, f, g*; fig. 2. *b, i*; fig. 4. *b, i*.

VESSELS, every where, accompanied the branches of the umbilical vein, corresponding with them in size, as well as situation; and, joining together, formed trunks, from which, at the sides of the pelvis, two vessels were continued, one of them on each side of the vesica urinaria and urachus, to the umbilicus, which they perforated, and then went, along the umbilical cord, towards the placenta, resembling the umbilical arteries. See fig. 3. *b, i, k, l, m*; fig. 4. *k, l*; fig. 2. *b, i*.

B

UNLUCKILY

UNLUCKILY, before I received the monster from Mr ANDERSON, he had entrusted the injection of its placenta to some person, who had managed it so negligently, that nothing, he told me, could be determined as to the distribution or communication of the vessels of the placenta with each other, or with those of the placenta of the complete child, or with those of the mother.

EXPLANATION of the FIGURES, representing the parts of a human Male Monster, of its real size.

FIG. I. represents the fore view of it entire.

- A. B. C. A circular mass, more than two inches thick, which supplies the place of head, trunk and arms.
- D. A thimble-like cavity, somewhat resembling the mouth.
- E. The umbilical cord.
- F. G. The penis and preputium.
- H. I. K. L. M. N. The thighs, legs and feet.

FIG. II. In this figure, at the letters A. B. C. D. F. G. H. I. K. L. M. N. the same parts are represented as in fig. I. The cavity of the abdomen being laid open by a longitudinal incision, we perceive,

- O. P. The small intestine.
- Q. The caput coli, and appendix vermiformis.
- R. S. T. U. V. The great intestine.
- W. X. The right and left ureters.
- Y. The vesica urinaria and urachus.

Z. The

Z. The left testicle, with its spermatic cord, cremaster muscle and vas deferens.

b, i. Two large vessels, at the sides of the pelvis, furnished by the umbilical vein.

E. E. The two umbilical arteries.

FIG. II.* In this figure, the conglobated, lymphatic or lacteal glands of the mesentery are represented.

FIG. III. In this figure, the distribution of the blood-vessels, chiefly, is represented. At the letters A. B. C. F. G. H. I. K. L. M. N. the same parts are represented as in fig. 1. and fig. 2.

O. P. shew the intestines pushed behind the blood-vessels to the left side.

W. W. The kidneys and ureters.

X. The ribs which covered the kidney, drawn towards the right side.

Y. The bladder of urine.

a, b, c, d, e, f, g, The umbilical vein, divided into branches for the several parts of the body.

b, i, k, l, Vessels accompanying the several branches of the umbilical vein.

m, Two vessels resembling the umbilical arteries.

n, n, The sciatic nerves.

FIG. IV. In this figure, the spinal marrow, and nerves connected with it, are chiefly represented.

A. B. C. H. I. K. L. M. N. represent the same parts as the former figures.

V. represents a probe passed from the rectum through the neck of the bladder into the urethra.

Y. A bristle passed from the bladder into the urethra.

B 2

S. The

S. The spinal marrow.

E. The cauda equina.

I. 2. &c. to 16. Nerves sent off from the spinal marrow in pairs.

17. The os sacrum.

n, n, The sciatic nerves.

REMARKS on such MONSTERS.

MONSTERS wanting the head, heart and lungs, and, in almost every other respect, agreeing with that above described, have been mentioned by authors, particularly by MERY and WINSLOW *, and the learned Dr ROEDERER † has given a full description of a monster, in which one small muscular sac only was found, instead of a complete heart, communicating with the continuation of one of two veins which were found in the umbilical cord; but the real course of the blood, or the causes of its motion, appear to have been misapprehended by all these authors.

MERY thinks the blood of the fœtus must have been moved by the motion of the heart of the mother, and considers the want of the heart in such monsters, as a strong confirmation of the opinion he entertained, that there is a circulation of the blood carried on between the mother and the fœtus ‡.

As

* Mem. de l'Acad. 1720 and 1740.

† Act. Got. t. iv. 1754.

‡ MERY, Mem. de l'Acad. des Scien. 1720. 1^{re} Reflexion. "Sa vie n'a pu avoir pour principes que la respiration et le mouvement circulaire du sang de sa mere." And in the Histoire, "Le défaut du cœur prouve que le sang qui a circulé dans ce fœtus ne recevoit pas son impulsion que du cœur de mere." M. MERY a toujours soutenu la circulation reciproque entre la mere et le fœtus, et telle que le fœtus est toujours comme un membre de la mere.

As WINSLOW had not found any red blood in the vessels of the fœtus, nor traced within it the branches of the umbilical vein, but those only, as he supposed, of the vessel he called aorta, and which he thought performed the office of an artery, he is led to the supposition, that, instead of a circulation, there was only a sort of progression of the colourless blood, or lymphatic humour, to the capillary extremities of the arterial ramifications, and that it transfused, by little and little, and very slowly, into the cellular texture of all the parts, and perhaps, at last, passed through the pores of the skin, in the form of moisture *.

Dr ROEDERER † not only applies the term of vena cava to the large vein with which the umbilical vein is joined to the heart, but describes the cava as ascending from the abdomen to the thorax ‡. In like manner, he not only applies the name
aorta

* WINSLOW, Mem. de l'Acad. des Scien. 1740.

P. 588. "La veine ombilicale, s'étant écartée du cordon de son entrée dans le ventre, y formoit un tronc fort court, qui montoit tout droit et s'implantoit à la base du bouton cutané, s'adossant là avec le tronc d'un autre vaisseau de pareille grosseur, qui sortoit de la même base, et qui étant d'abord courbé vers en bas, descendoit derrière les paquets des intestins, à peu près comme le tronc de la portion inférieure de l'aorte, et se distribuoit ensuite en plusieurs branches, de la manière que je dirai ci après."

P. 590. "On ne voyoit pas une goutte, ni aucune apparence de sang rouge dans toute l'étendue du corps de cet enfant ; ni aucun vestige de vaisseaux veineux."

P. 600. "Hors la petite portion de la veine umbilicale après son entrée par le nombril, je n'ai trouvé, dans tout le corps de cette enfant, aucun vaisseaux veineux, ni le moindre vestige soit de tronc, soit des ramifications de veines."

P. 604. "Mais à l'égard de la circulation intrinsèque dans les parties mêmes de ce demi-corps, l'absence ou la privation totale des vaisseaux veineux m'a fait conjecturer, qu'au lieu de circulation proprement dite, il n'y a eu qu'une espèce de progression ou trusion jusqu'aux extrémités capillaires de toutes les ramifications artérielles, et que là ce sang lymphatique transfusoit, peu à peu, et très lentement dans le tissu cellulaire de toutes les parties. — Et, peut-être, passoit par les pores externes de la peau, en manière de moiteur. Je n'avance tout ceci que comme des pures conjectures," &c. &c.

† Com. Soc. R. Sc. Gotting. tom. iv. com. 4.

‡ P. 109. "Duplicem autem umbilicalis funis venam largitur ; altera minor, cum vena cava, ex abdomine ascendente confluit."

aorta to the vessel which accompanies the continuation of the umbilical veins ; but speaks of his aorta as ascending from the thorax to the head *, and sending off the subclavian and the carotid arteries ; and remarks, that canals proper to the latter were wanting †. And he observes, that the aorta, after descending, as usual, between the crura of the diaphragm, gave off the mesenteric, renal, lumbar and iliac arteries ; and that the left iliac artery sent off an umbilical artery ; and concludes his description in the following words : “ Ita, quidem, si arteriæ umbilicalis dextræ, arteriæque celiacæ defectus——excipiatur, vix ab usitata fabrica aberrans arteria aorta in abdomine distribuitur.”

AFTER an elaborate description of the several parts of the monster, Dr ROEDERER proposes the cause of the motion of its humours, in the following words :

P. 189. “ MOTUS qui—humores agitat, causa indagatur. “ Ast aliquis, lentus licet, foetus parasitici humores motus agitavit. A corde, fueto motore, repeti iste motus nequit, neque multum auxilii propulsus in uterum maternum sanguis ferre potest. Præter vero istum, levem, debilemque.—— “ Ipsa vasorum actio, sive contrahendo agat, sive attrahendo, “ vi

* P. 121. “ Arteria magna, quam aortam vocant, ex abdomine in thoracem ascendit. In thorace eandem pene directionem servans, nulloque cum corde canali cõfluens, sola et a corde distincta, iter suum absolvit. Nullus proinde ex aorta arcus formari potest, sed laterales rami ex recto aortæ trunco emittuntur. Sunt isti rami qui descripti sequuntur.

In regione costæ primæ levissime descendentes arteriæ subclaviæ nascuntur ; ex quibus vicissim triplex alia ramorum species oritur, quarum primus ad cervicem, &c. Porro truncus aortæ per semipollicem postquam progressus est in duos ramos dividitur, duas nempe arterias carotides, quæ ad altitudinem laryngis sine insigniori ramo ascendunt.——Ascendit, autem, carotis dextra, &c.——Ad latus tandem laryngis canalis communis in sex omnino ramos dividitur.”

† P. 143. “ Canalis pro arteria carotide deest. Carotis per amplum foramen lacrarum ad cerebrum tendit.”

“vi illa capillaribus tubis familiari, præcipuum humoribus
 “motum impertiri debet.—Accedant forsan et aliæ in fœtu
 “nostro causæ incognitæ, ipsa fortasse a colore excitata fluido-
 “rum agitatio, aliaque.”

BUT as to the direction in which he supposed the humour to be moved, he says nothing, and therefore leaves us to judge of his opinion, from the foregoing description of the blood-vessels.

TO the opinions of all these authors, when fully considered, we shall find insuperable objections.

THUS, without saying in objection to that of MERY, that it is so far from being certain, that there is a circulation of red blood between the mother and fœtus, that the contrary opinion is the most probable, we cannot conceive, although the anastomoses of the uterine with the placental vessels were proved, that the mere impulse of the blood in the minute arteries should have carried the blood, not only into the trunks, but through all the capillary branches of the vessels of the fœtus, and again back from these to the placenta, and from its umbilical arteries into the umbilical veins and veins of the uterus.

THE opinion of WINSLOW is far more unsatisfactory than that of MERY. In the first place, it cannot be applied to the monster described by MERY, or to that before us, where there were two sets of vessels. In the next place, WINSLOW was so far from tracing distinctly the joining of the umbilical vein with the vessel he calls aorta, that he describes it as merely *s'adossant* with the trunk of the aorta*.

3. ALTHOUGH he repeatedly affirms, that there were no venous vessels in any part of the body of the monster, yet his description of the vessels of the kidney will not, when considered, be found to correspond with his general assertion; for he describes a vessel which indeed he calls arterious, but which began

* See p. 588. of Mem. de l'Acad. or Note, p. 221.

began on the fore-part of the belly above the navel, at the place where the small portion of the umbilical vein terminated in the cavity of the cutaneous button, from which various branches were sent into the kidney at its convex part, and from its concave part, different arteries, he says, came out in an extraordinary manner *.

UPON the whole, as the umbilical cord is not said to have been uncommon in size or structure; as there were two sorts of vessels connected with the kidney; as it is so improbable, as to be incredible, that the foetus received arteries without corresponding veins, or that there was merely a protrusion of the humours, and exudation of them, without circulation, I have no doubt that WINSLOW, especially as he did not inject the vessels of the umbilical cord, had mistaken the continuation of the umbilical veins, and the branches of the vessels he calls aorta, for branches of the same vessel; and as the monster he examined agreed very nearly, in all other respects, with that I have described, I apprehend it must have agreed likewise in having two kinds of blood-vessels or arterious and venous canals.

THE learned Dr ROEDERER rejects the opinion of MERY, that the blood of the foetus is circulated by the heart of the mother, and supposes, that capillary attraction, heat, and some activity of the vessels, may contribute to its motion. But as he applies the term aorta, not to the continuation of the umbilical vein, but to the other principal vessel of the monster, and describes

* P. 602. "Ce tronc arteriel qui étoit comme la portion inferieure de l'aorte descendante, au lieu de tenir la route naturelle en arriere le long des vertebres, il en étoit ici très éloigné. Il commençoit sur le devant du ventre au dessus du nombril, à l'endroit où se terminoit la petite portion de la veine ombilicale.—Il jettoit des branches dans la masse du rein par sa convexité. Il sortoit de la concavité plusieurs artères.

describes it as sending branches downwards from the abdomen to the inferior extremities, and upwards from the thorax to the head, and applies the name of carotid arteries to two of these branches, with the additional remark, that the canales carotici were wanting, it will, I apprehend, appear evident from these circumstances, and from what I am about to observe in the next section, that he misunderstood the direction in which the blood was moved and circulated.

Of the Direction of the Blood in this Monster.

As there are two kinds of vessels in the umbilical cord, and likewise within the body of this monster, which we shall call, in the common style, arterious and venous, we cannot doubt, that these communicated with each other, and that the blood was conveyed by them in a circle.

To describe the circle more exactly, we cannot doubt, that the blood was conveyed from the placenta by the umbilical vein into the body of the monster. We next found, that the umbilical vein within the monster was divided into various branches, which could be traced to all its parts, or that these branches performed the office of arteries, or resembled the vena porta hepatica. Contiguous to these branches, we found, every where, other vessels which formed a trunk or large vessel, which, by its situation, resembled our aorta. But we must suppose, that these branches served the purpose of receiving the blood from the extremities of the branches of the umbilical vein, or were in reality venous vessels. From the vessel resembling the aorta in situation, but very different in office, two vessels were sent off, which ran at the sides of the bladder to the umbilicus, and formed the arteries of the umbilical cord and of the placenta, and, in the placenta, must have terminated

in the minute beginnings of the umbilical vein, to complete the circle in which the foetal blood was moved.

THUS, we observe the umbilical vein in the placenta and umbilical cord performing the office of a vein, but its continuation within the body of the monster, performing the office of an artery. On the other hand, we find the vessel we have called aorta, performing the office of a vein within the monster, and that of an artery in the umbilical cord and placenta.

Of the Causes of the Motion of the Blood in this Monster.

IN the monster examined by WINSLOW, which I have endeavoured to shew agreed very nearly with that I have described, no red blood was found in any of the vessels; and therefore we must conclude, that none of the red arteries of the mother anastomosed with the umbilical veins; and even where there is the ordinary structure, it is so far from being certain, that the vessels of the uterus, which convey red blood, anastomose with those of the umbilical cord, that the contrary is the most probable opinion.

IT is therefore very improbable, that the blood in the umbilical vein was pushed on by the heart of the mother.

FURTHER, though we were to admit, that the arteries of the mother anastomosed with the umbilical veins, yet as their communications must be supposed very minute, and the momentum of the blood in them very much broken, we cannot conceive, that it could have been sufficient to push the blood through the terminations of all the branches of the umbilical veins, in the several organs of its body, into the vessel we call aorta, and again from the aorta back to the placenta by the umbilical arteries, and through the minute branches of these to the veins of the mother, and beginnings of the umbilical veins.

WE

WE therefore must conclude, that the circulation of the blood in the placenta and body of the monster, was carried on by a well regulated muscular action of the blood-vessels. In one of the worms, the *echinus esculentus*, I found in the mesentery, which is a principal part of it, two such large vessels without a heart, and which, we can scarcely doubt, resembled our aorta and cava, and circulated its fluid; and in fishes*, the blood which passes through the liver describes three circles, and in all other parts of the fish the blood describes two circles before it returns to the heart; which motion of it we must suppose to be chiefly owing to the muscular action of the vessels, as the force of the heart appears to be as much spent in the gills of the fish as in the lungs of a man.

FROM considering the manner and cause of the motion of the blood in this monster, and comparing with it the motion of the blood in fishes and in the sea-egg, we are, by analogy, led to the following general conclusions:

1. THE arteries contribute much to the circulation of the blood in our bodies.

2. IT is probable that, in man, the veins likewise assist in circulation; and, in particular, there can be no doubt, that the vena portarum, by its action, contributes much to the motion of the blood through our liver.

3. FOR the like reasons, we may conclude, that arterious vessels, independent of the impulse of the heart, may act in such a manner, as to perform the secretion of liquors, to nourish the solids, and to add to their bulk; and particularly, that the branches of the vena portarum change certain parts of the blood into bile.

B 2

Remarks

* See MONRO on Fishes, p. 67. Tab. xliii.

Remarks on the Nervous System of this Monster.

1. As the spinal marrow, and pairs of nerves sent off from it, had nearly the usual size and structure, although the brain, cerebellum, and medulla oblongata, were entirely wanting, we find reason for calling in question the common doctrine of authors, which teaches, that the spinal marrow and nerves derive their origin from the brain and cerebellum, and are dependent upon it as much as the ducts of glands are upon the glands which send liquors into them.

2. FURTHER, as the several parts of this monster were furnished with nerves, and as we have found, that its arteries and veins, by a well-regulated varied and complicated action, circulated the blood, we must suppose, that their muscular fibres were actuated by those nerves. We therefore find in this monster, not only the existence and common appearance of the spinal marrow and nerves connected with it, although the brain and cerebellum were wanting, but we have proof that these, independent of the brain and cerebellum, may actuate the muscular fibres in the vessels of an animal, or that nervous energy, or fluid, as it is commonly called, is not derived from the brain and cerebellum solely; that is, we conclude, that the nerves, as well as the brain and cerebellum, are capable of furnishing nervous energy; and that there is no more reason for believing, that the nerves are derived from the brain, than that the brain is derived from the nerves; or all the parts and branches of the nervous system appear to possess the general power or office of furnishing nervous energy.

Of the Duration of the Life of this Monster.

As in man and similar animals, the direct or indirect influence of respiration seems necessary for the continuance of life, and as the lungs were wanting in this monster, we must suppose, that it could have outlived the separation from the mother for a very short time only. But when we add to this, that, by the ligature of the umbilical cord, a stop would be mechanically put to the circulation of its blood, it is evident, that its life must have terminated with its delivery.

Of the Time at which this Monster must have acquired the Structure which has been described.

As this monster was provided with a distinct placenta and membranes, and its body surrounded with and protected by the liquor amnii; as no vestige appeared of the brain, cerebellum, organs of the senses, or other parts of the head; as nervous threads, proper to this monster, ascended from the upper end of the spinal marrow towards the upper parts of its body; as its system of circulating vessels was complete without a heart, and the manner of their branching different in many respects from the common structure: it must surely appear, to an unprejudiced person, absurd to suppose, with many eminent authors, that such monsters, when first produced, had the ordinary structure, and that this was afterwards altered by pressure and other accidents.

THE like observation may be extended to many other monsters in my possession, I believe I might say to almost all other
monsters

monsters which have been described ; particularly to two, of which I published a description, illustrated with figures, in my work on the Nervous System. In one of them, a human monster, one heart supplied two heads and two trunks. In the other, a kitten, one heart, consisting of two auricles and two ventricles, sent off from its left ventricle one aorta, which supplied one head and two bodies *.

* See Observations on the Nervous System, Tab. viii. ** and Tab. xii.

F I N I S.

Figl



A. Hall del.

J. Smith sc.

