

Modern medical therapeutics : a compendium of recent formulae and specific therapeutical directions, from the practice of eminent contemporary physicians, American and foreign / by George H. Napheys.

Contributors

Napheys, George H. 1842-1876.
Royal College of Physicians of London

Publication/Creation

London : Bailliere, Tindall, and Cox, 1885.

Persistent URL

<https://wellcomecollection.org/works/dpf5z4xw>

Provider

Royal College of Physicians


License and attribution

This material has been provided by This material has been provided by Royal College of Physicians, London. The original may be consulted at Royal College of Physicians, London. where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>




S4126-Q-f-41

S.L. 1
The Royal College of Physicians
of London.

*From the Library of
Sir Andrew Clark, Bart.
Presented by Lady Clark.*

80 D









MODERN
MEDICAL THERAPEUTICS:

A COMPENDIUM OF
RECENT FORMULÆ

AND

SPECIFIC THERAPEUTICAL DIRECTIONS,

FROM THE PRACTICE OF EMINENT CONTEMPORARY PHYSICIANS,
AMERICAN AND FOREIGN.

BY GEORGE H. NAPHEYS, A. M., M. D., ETC.

EDITED BY JOSEPH F. EDWARDS, M. D., AND
D. G. BRINTON, M. D.

EIGHTH EDITION, ENLARGED AND REVISED.

LONDON:
BAILLIERE, TINDALL & COX,
20 KING WILLIAM STREET, STRAND.
1885.



Entered according to Act of Congress, in the year 1885, by

D. G. BRINTON,

In the Office of the Librarian of Congress, at Washington, D. C.

All rights reserved.

ROYAL COLLEGE OF PHYSICIANS LIBRARY	
CLASS	616-08
ACCN.	24482
SOURCE	
DATE	

PREFACE TO THE EIGHTH EDITION.

IN presenting this work to the profession for the eighth time, every part of it has been subjected to a careful revision. All valuable new material which has appeared since the last edition has been incorporated. The recent additions to the *Materia Medica*, large in number, and some of great importance, have been introduced, and their merits stated.

These additions have unavoidably increased the size of the volume, and this in spite of the omission of the section on "Diseases of Children," which it is intended to make the nucleus of a special treatise on the same plan. This brings up the actual increment of new matter to considerably more than a hundred pages, while all that is retained from former editions has been closely scanned, and much of it recast.

A portion of this new matter has never before been published. It has been received in answer to an inquiry sent to a number of eminent physicians as to their most successful methods of treating various diseases, and also from notes of hospital and clinical lectures.

PREFACE TO THE SEVENTH EDITION.

THE present edition of this work is not a mere reprint of the last one. The whole material has been subjected to a close scrutiny, and the most recent writings on materia medica and therapeutics have been searched in order that no addition of moment should be overlooked. Some paragraphs have been dropped, others have been condensed; but the additions, in spite of such condensation, have been so extended that in order to keep the volume within its previous limits, it has been found necessary to select a smaller type for the *Résumé of Remedies*. By this change, the editors have been enabled to add several topics not previously treated, and many new names and references, to those heretofore given. Especial attention has been given to the periodical literature of the last few years, as so many valuable suggestions are sure to lie neglected in journals unless transferred to the more accessible form of a volume.

It will be noted that this volume is confined to diseases strictly *medical*. Surgical diseases are treated in a similar manner in the companion volume on *Surgical Therapeutics*; while diseases of women and obstetrical complications are considered in the third number of the series, the *Therapeutics of Gynecology and Obstetrics*.

PREFACE TO THE SIXTH EDITION.

IN presenting the sixth edition of this work to the medical public, the editor has endeavored to render it worthy the very favorable reception it has had, by subjecting it to a thorough and careful revision. The progress of therapeutics has been very marked within the last few years. Standard treatises on Practice, written by men still living, have, in many instances, become, in great part, obsolete. WATSON and WOOD, AITKEN and TANNER, no longer represent the best therapeutics of the day. Their methods and their remedies have been largely superseded; and the practitioner who would keep abreast of the advances made in his profession must look to a more modern class of writers for his information.

The recognition of this fact has led to many substitutions in the present edition. The monographs and treatises, as well as the numerous medical periodicals of the last year or two, have been carefully examined and collated, so as to render this work in reality what it professes to be—a treatise on *modern* therapeutics. At the same time, the effort has been made to exercise a judicious conversation in regard to long-known and well-tried remedies; and to avoid admitting what is new simply because it is new.

EXTRACT FROM THE PREFACE TO THE FOURTH EDITION.

THIS work stands alone, of its kind, in medical literature. It presents the Art of Therapeutics in all its aspects, and divested of that barren theorizing which has been its bane. Acquainting the reader with the exact treatment of each disease by living clinical teachers and careful practitioners of several countries, it vastly widens his therapeutic resources, prevents him from becoming a routinist, and inspires him with confidence in the remedies he employs.

So far from leading him to become a mere *formularum præscriptor*, the diversity it exhibits teaches him independence of thought, while the care with which the indications calling for particular remedies or combinations are given, awakens and instructs his powers of clinical observation.

But this remark is not intended to belittle the value of definite formulæ. The author has most wisely preserved these whenever possible. By skillful combinations, nauseous articles are rendered palatable, which is often no small matter; ingredients of facile decomposition may be given permanence; the labors of the pharmacist are lightened; and, more than all, the efficiency of pharmaceutical products may be very largely increased.

The intelligent recognition of the last-mentioned fact may be claimed as one of the later advances of medical science. The poly-pharmacy of our forefathers in the profession has disappeared, not giving way to any nonsensical 'law of the single remedy,' as Homœopathy advocates, but to a clear insight into the *synergic* action of remedies, by virtue of

which, a judicious combination of several drugs acts in a given direction more forcible than any one of them singly. It is needless, here, to quote instances of this truth. They may be found in abundance in the pages which follow.




TABLE OF CONTENTS.

	PAGE
PREFACE TO THE EIGHTH EDITION.	iii
PREFACE TO THE SEVENTH EDITION	iv
PREFACE TO THE SIXTH EDITION.	v
PREFACE TO THE FOURTH EDITION	vi
TABLE OF CONTENTS.	ix

I. DISEASES OF THE NERVOUS SYSTEM.

GENERAL THERAPEUTICS OF THE NERVOUS SYSTEM	17
APOPLEXY AND CEREBRAL HYPERÆMIA	21
Notes on Remedies.	30
General Remedies	31
External Remedies.	32
CEREBRAL ANÆMIA	33
Notes on Remedies.	34
CHOREA.	34
Notes on Remedies.	42
External Remedies.	46
EPILEPSY	47
Inhalations,	57
Mechanical and Physical Means.	57
Notes on Remedies.	58
External Remedies.	63
HEADACHES.	64
Bilious or Dyspeptic Headache	65
Congestive or Plethoric Headache.	66
Gouty Headache	68
Hysterical Headache	69
Migraine, Hemicrania, Periodical or Sick Headache.	69
Nervous Headache	72
Neuralgic Headache	75
Rheumatic Headache	76
NOTES ON REMEDIES	77
Bilious and Dyspeptic Headache	77
Congestive or Plethoric Headache	78
Gouty Headache	78
Hysterical and Nervous Headache	78

	PAGE
EXTERNAL REMEDIES	80
Neuralgic Headache	80
Periodic Headache	80
Rheumatic Headache	81
HYPOCHONDRIASIS OR MELANCHOLIA	81
Notes on Remedies	84
INSANITY	84
Notes on Remedies	90
INSOMNIA OR SLEEPLESSNESS	90
Notes on Remedies	96
External Remedies	98
NEURALGIA	98
Notes on Remedies	106
External Remedies	111
NEURASTHENIA	114
Notes on Remedies	115
PARALYSIS	115
Strychnia and its administration	116
Electricity in Paralysis	117
Notes on Remedies	119
External Remedies	120
PROGRESSIVE LOCOMOTOR ATAXIA (POSTERIOR SPINAL SCLEROSIS)	121
Notes on Remedies	123
External Remedies	124
SCIATICA	124
Notes on Remedies	127
External Remedies	129
SPINAL IRRITATION	130
Notes on Remedies	132
SPASMODIC DISEASES	133
TIC DOULOUREUX (FACIAL NEURALGIA)	137
Notes on Remedies	139
External Remedies	140
VERTIGO	141
Notes on Remedies	144

II. DISEASES OF THE RESPIRATORY SYSTEM.

APHONIA	146
Notes on Remedies	148
ASTHMA	149
Notes on Remedies	158
External Measures	163
ACUTE BRONCHITIS	164
Notes on Remedies	169
CHRONIC BRONCHITIS	171
Notes on Remedies	178

	PAGE
External Remedies	183
CATARRH, CORYZA, AND POST-NASAL CATARRH	184
The Abortive Treatment of a Cold	184
General Treatment of Catarrh	187
Post-nasal Catarrh.	189
Radical Treatment of Nasal Catarrh	191
Notes on Remedies.	193
External Remedies.	195
EMPHYSEMA.	196
Pneumatic Treatment	200
Notes on Remedies.	200
External Remedies.	201
EMPYEMA	202
Notes on Remedies.	206
External Measures.	206
HÆMOPTYSIS	206
Notes on Remedies.	210
External Measures.	212
HAY FEVER, HAY ASTHMA, ROSE COLD, AUTUMNAL CATARRH	213
Notes on Remedies	217
LARYNGITIS.	219
Acute Laryngitis.	220
Chronic Laryngitis.	221
Notes on Remedies.	223
External Measures.	225
PHTHISIS	225
Acute Phthisis.	225
Chronic Phthisis.	226
Cod-liver Oil Treatment	234
Treatment by Hypophosphites	236
Calcification of the Lung	238
Specific or Anti-parasitic Treatment.	239
Climate and Travel in	241
TREATMENT OF COMPLICATIONS	243
The Anæmia and Debility	243
The Cough	244
The Diarrhœa.	246
The Dyspnœa	247
The Hæmoptysis.	247
The Laryngitis	247
The Night Sweats	247
Notes on Remedies	250
Inhalations.	254
External Remedies.	255
PLEURISY.	256
Treatment by Compression of the Thorax	257
Notes on Remedies.	258

	PAGE
PNEUMONIA.	259
General Treatment.	261
Notes on Remedies.	266
External Remedies.	269

III. DISEASES OF THE CIRCULATORY SYSTEM.

GENERAL THERAPEUTICS OF HEART DISEASE	271
Treatment of Secondary Affections of the Heart.	275
ANGINA PECTORIS.	281
Notes on Remedies.	284
CARDIAC DILATATION.	285
Notes on Remedies.	287
CARDIAC HYPERTROPHY	288
Notes on Remedies	291
CARDITIS, ENDOCARDITIS AND PERICARDITIS	292
Notes on Remedies	296
Rheumatic Carditis	297
FATTY DEGENERATION OF THE HEART	298
Notes on Remedies	298
FUNCTIONAL PALPITATION AND OTHER DERANGEMENTS	299
Notes on Remedies	301
External Measures	302
IRRITABLE HEART	302
VALVULAR DISEASE OF THE HEART	303
Notes on Remedies	308

IV. DISEASES OF THE DIGESTIVE SYSTEM.

APETITE, LOSS OF	311
BILIARY CALCULUS	313
Notes on Remedies	316
CHOLERA ASIATICA	317
Notes on Remedies	326
General Measures	329
COLIC.	330
Notes on Remedies.	332
CONSTIPATION—HABITUAL	334
Notes on Remedies	342
DIARRHŒA—ACUTE AND CHRONIC	346
Notes on Remedies	352
Acute Diarrhœa	352
Sub-Acute and Chronic Diarrhœa	354
DYSENTERY—ACUTE AND CHRONIC	358
Acute Dysentery	358
Chronic Dysentery	363
Medication by the Rectum	365

	PAGE
Notes on Remedies	367
Acute Dysentery	367
External Measures	371
Chronic Dysentery	371
DYSPEPSIA	373
Cerebral Dyspepsia	378
Climacteric Dyspepsia	379
Notes on Remedies	380
GASTRITIS, GASTRO-ENTERITIS AND ENTERITIS	387
Notes on Remedies	389
External Measures	389
GASTRIC ULCER	390
Notes on Remedies	395
HÆMATEMESIS	397
Notes on Remedies	398
HEPATITIS, HEPATIC ABSCESS	399
Notes on Remedies	403
INTESTINAL WORMS	404
Tænia, Tape Worms	404
Notes on Remedies	407
Ascaris Lumbricoides, Lumbrici, Round Worms	410
Ascaris Vermicularis, Oxyuris, Thread Worms, Seat Worms	413
Notes on Remedies	416
Trichinæ, Trichinosis	416
JAUNDICE	417
Catarrhal Jaundice	417
Notes on Remedies	422
TORPIDITY AND FUNCTIONAL DERANGEMENT OF THE LIVER	423
Notes on Remedies.	424
PYROSIS.	425
Notes on Remedies.	426
VOMITING.	427
Notes on Remedies.	427

V. DISEASES OF THE URINARY SYSTEM.

ALBUMINURIA.	431
Notes on Remedies.	434
BRIGHT'S DISEASE.	436
Notes on Remedies.	442
External Measures.	443
DIABETES INSIPIDUS, POLYURIA, POLYDIPSIA	444
Notes on Remedies.	446
DIABETES MELLITUS.	447
The Dietetic Treatment	447
The Milk Diet Treatment.	451
Notes on Remedies.	452

	PAGE
DROPSY.	455
Diuretics and Cathartics	459
Notes on Remedies.	461
External Measures	463
FUNCTIONAL URINARY DISORDERS	464
Incontinence of Urine	464
Dysuria.	466
Notes on Remedies.	466
HÆMATURIA.	467
Notes on Remedies.	469
NEPHRALGIA (RENAL COLIC).	470
PAROXYSMAL HÆMATINURIA	471
NEPHRITIS	472
Notes on Remedies	474
URÆMIA	476
Notes on Remedies	478


VI. DISEASES OF THE BLOOD.

ANÆMIA	479
Tonic Bath	481
Tonic Inhalation.	481
Notes on Remedies	484
External Measures.	485
CEREBRO-SPINAL MENINGITIS	486
Notes on Remedies	489
DENGUE (BREAK-BONE FEVER)	490
GOUT	493
Mineral Waters	497
Notes on Remedies	498
External Applications	500
HEMORRHAGIC MALARIAL FEVER.	501
INTERMITTENT FEVER (AGUE, CHILLS)	503
The Personal Prophylaxis of Malaria	503
The Anti-Quinine Treatment.	508
The Hypodermic Use of Quinine	514
The Treatment of Enlarged Spleen or Ague Cake	517
Notes on Remedies	519
MILK SICKNESS	528
PURPURA HÆMORRHAGICA.	530
Notes on Remedies.	531
RELAPSING FEVER.	531
ACUTE RHEUMATISM.	533
Salicin and Salicylic Acid Treatment	533
Lemon or Lime Juice Treatment	535
Alkaline Treatment	536
Notes on Remedies	544

	PAGE
Local Applications.	546
CHRONIC RHEUMATISM.	548
Notes on Remedies.	550
TYPHOID FEVER.	551
Turpentine Treatment	558
Antiseptic Treatment	559
Specific or Anti-Parasitic Treatment	559
Notes on Remedies.	561
TYPHO-MALARIAL FEVER	564
TYPHUS FEVER	565
Treatment by Cold Baths.	568
Notes on Remedies	569
VARIOLA—SMALL POX	570
Notes on Remedies.	572
To Prevent Pitting	573
External Measures.	574
YELLOW FEVER.	575
Notes on Remedies.	581

VII. TOXIC DISEASES.

ALCOHOLISM (INTOXICATION ; DELIRIUM TREMENS ; CHRONIC ALCOHOLISM).	583
1. Drunkenness.	583
2. Acute Alcoholism or Delirium Tremens.	586
3. Chronic Alcoholism	589
Notes on Remedies	594
ARSENICISM (CHRONIC ARSENICAL POISONING)	597
OPIISM (OPIUM EATING ; THE OPIUM HABIT)	598
HYDRARGYSM (CHRONIC MERCURIAL POISONING)	600
PLUMBISM (LEAD POISONING).	604



MODERN MEDICAL THERAPEUTICS.

I. DISEASES OF THE NERVOUS SYSTEM.*

The General Therapeutics of the Nervous System—Apoplexy and Cerebral Hyperæmia—Cerebral Anæmia—Chorea—Epilepsy—Headaches—Hypochondriasis or Melancholia—Insanity (Dementia)—Insomnia or Wakefulness—Neuralgia—Neurasthenia—Paralysis—Progressive Locomotor Ataxia (Posterior Spinal Sclerosis)—Sciatica—Spinal Irritation—Spasmodic Diseases—Tic Douleureux—Vertigo.

THE GENERAL THERAPEUTICS OF THE NERVOUS SYSTEM.

The general resources at the command of the physician in treating diseases, are set forth as follows by

DR. E. C. SEGUIN

in a lecture in his *Opera Minora*, 1884:

Means which increase the blood supply. The first is *posture*. By placing the patient on his back, the amount of blood in the spinal cord is increased. In this position, if the limbs and trunk are elevated, anæmia of the brain is counteracted. *Compression of arteries* is another important measure. Tourniquets may be applied to the limbs in extreme cases of cerebral anæmia. Compression of the carotids

* NOTE.—The subjects of Tetanus, Traumatic Neuralgia, Traumatic Paralysis, Rabies and Neuroses of the Skin, are treated of in NAPHEYS' *Surgical Therapeutics*; Hysteria and the Special Nervous Affections of Women in the *Therapeutics of Gynecology and Obstetrics*, edited by Dr. WM. B. ATKINSON.

was used to cut short epileptiform seizures supposed to be due to cerebral hyperæmia. Of *stimulants*, alcohol is the best; food may also be regarded as a stimulant. *Paralyzing the vaso motor nerves*, by producing enlargement of the arteries, is productive of hyperæmia in the arterial district, to which the nerves are distributed. Of means to produce this result, the most effectual is inhalation of *amyl nitrite*, gtt. ij-v. In the epileptic aura it is frequently efficient in checking the attack.

Means which decrease the blood-supply. Of these the most obvious is *venesection*. This is now out of fashion; but with symptoms of violent cerebral hyperæmia in a full-blooded subject, it would be proper to use it. Leeches and wet-cups relieve hyperæmia, not by abstracting blood, but by irritating nerves, and thus acting on the vaso-motor filaments. *Position*, as by raising the part, decreases its blood-supply. In congestion or other affections of the spine, place the patient upon his face. In some cases of insomnia, the patient can sleep in the sitting posture. *Cold*, applied continuously, diminishes the amount of blood in a part. The best method of employing it is to apply ice in block immediately to the skin; or pounded ice in a bladder; or the ether spray. It is indicated in meningitis and in intra-spinal congestion and in inflammation. *Galvanism* stimulates the vaso-motor nerves and produces local anæmia; so also will *ergot*, and in a less degree *belladonna*. These two remedies are therefore largely used in congestions and inflammations of the nerve-centers. *Mental* and *physical rest* are important aids to diminish the blood-supply to the nerve-centers.

Cerebral excitants. *Alcohol* stands first, then *cannabis indica*, and in some constitutions belladonna and opium in small doses. *Emotional excitement* is especially indicated where the mind is occupied in contemplating delusions.

Spinal excitants. Of these the most prominent are *nux vomica*, and its alkaloids, *strychnia* and *brucia*. The indication for their use is simple exhaustion of the nerve-centers, not due to inflammation. They are also valuable in some cases of chorea, epilepsy and spinal irritation. The dose should be small at first, but pushed rapidly to the point of obtaining the physiological effects of the drug. *Quinia* is also entitled to a place in this class of remedies. *Cantharides* seems to act more particularly upon the lower portion of the spinal cord. It is useful in bladder paresis, impotency, and simple genital inertia.

Cerebral Depressants. *Cold* may be employed for this purpose, as above stated. *Bromide of potassium* is a true hypnotic, and valuable in insomnia, delirium, mental anxiety, etc. *Hydrate of chloral* is the best medicine for producing sleep. It does not intoxicate nor much disturb the digestive system. It is invaluable with insane patients. A perfectly safe dose is gr. xx–xxv. The conjoint use of the bromide and chloral is very satisfactory. A drachm of the bromide may be given in the afternoon, followed at bed-time by ℥i–ij of chloral. *Opium* has frequently, especially on females, very unpleasant after-effects. A very small amount of atropia, added to a solution of morphia, will palliate the nauseating and constipating effects of the latter.

Spinal Depressants. The typical medicine of this class is *conium*. It is the antagonist of strychnia, and can be used with benefit in spasmodic affections. This remedy should be given only once a day. A reliable preparation is the English *succus conii*, which may be administered in doses of gr. i–ijss once daily. The fluid extract combined with bromide of potassium is an efficient remedy in chorea and epilepsy. *Bromide of potassium* is also a spinal depressant, and diminishes reflex excitability. It is indicated in morbid excitement of the lumbar part of the spinal cord, as in nymphomania and satyriases. The vomiting of pregnancy, that after the inhalation of ether, and the spasmodic state of various sphincters, are cured by it. In epilepsy it is our chief reliance.

Tonics and Restoratives. Two remedies of this class are especially applicable to the nervous system—*phosphorus* and *fats*. The various phosphates, if they do good, do so very slowly, and are of little service in serious disease. Phosphorus itself may be administered in Thompson's formula (see Index), or as phosphide of zinc. Some organizations are very susceptible to this remedy. In cerebral mal-nutrition, in neuralgia, in spinal irritation, in hysteria, and in varieties of paralysis, this drug is of the highest value. *Fatty food*, as cod liver oil, etc., is demanded in these same conditions. Of other tonics may be named strychnia, arsenic, zinc, iron, quinia, and cold. *Cold* may be applied by sea-bathing, the shower-bath, etc. *Strychnia* as a tonic must be given in small doses for a long period. *Arsenic* is often marvelous in chorea. In the nervous state produced by alcohol, *oxide of zinc* is almost as satisfactory as arsenic in chorea. It is best when combined with *nux vomica*. *Quinia*, appears to exert a direct restora-

tive effect on the nerve-centers. *Iron* does not especially affect the nervous system.

Two specific modes of medication may be mentioned here, which relieve many cases of nervous disease by acting on the pathological conditions which cause them. One is the treatment of constitutional syphilis by iodide of potassium; the other the specific medication in malaria. Malarial neuralgia is a frequent affection. A rapid removal of this disease is secured by using quinia in the shape of hypodermic injection over the affected nerve.

Counter-Irritants. The alternate application of cold and heat does much to improve the nutrition of paralyzed parts. The *actual cautery* is a means of the utmost value in the treatment of nervous affections. A deep burn is not needed, surface irritation being all that is necessary. The cautery, heated to a white heat, should be rapidly drawn over the part selected, four or more strokes, of from one to six inches in length, being made. If properly performed, the pain is not great, sores never result, and even blisters but seldom. *Setons* have gone out of fashion, and properly, as there are grave objections to their use. *Blisters* are an invaluable means. They should not produce suppuration. Repeat the blisters and heal the blebs as soon as possible. *Dry cupping* is beneficial in many spinal affections; their number should be large—from ten to twenty. It is not necessary to apply them immediately over the seat of disease. *Wet cups* are still used in some inflammatory affections, but it is doubtful if they act better than the dry, and the irritation is greater. *Sulphur baths* act as a counter-irritant by their effect on the skin. The *sulphuret of potassium* may be dissolved in water for a bath; or sulphur may be vaporized while the patient is covered, except his head, by a rubber cloth. Ten or twenty minutes of this exposure produces a great deal of cutaneous irritation. In locomotor ataxia a series of such baths have done great good. They are also useful in lead-poisoning.

Electricity. All three forms of electricity may be used—franklinic, galvanic and faradic. Within certain limits, all of these have useful applications in nervous diseases.

Hygienic Means. The principal measures are a supporting diet, the relief of constipation, moderate exercise short of exhaustion, etc. Pneumonic affections and bed-sores must be guarded against. With regard to coffee, alcoholic drinks and tobacco, in dealing with those accustomed

to their use, we should require moderation rather than abstinence, as the nervousness consequent upon giving up an established habit is worse than the effects of moderate indulgence.

APOPLEXY AND CEREBRAL HYPERÆMIA.

DR. EDWARD C. MANN.

Cerebral Hyperæmia; Congestion of the Brain.—This diseased condition is a common one of American life. Its treatment, as laid down by Dr. MANN (*Psychological Medicine*, 1884) is to warn the patient against great elevations and alternations of temperature, exposure to the sun, alcohol, and sleeping with the head too low. *Dry cups* to the back of the neck, hot mustard water to the feet, ice to the head and the nape of the neck, purgatives, fluid extract of *ergot* ʒss to ʒj, thrice daily, combined with sodium bromide, and the use of cerebral electrization, are all to be used as circumstances demand.

Where there are a feeble pulse and pale surfaces, *stimulants* should be used in moderate quantities. Tonics are also demanded, and there is none better in the case of overworked professional and business men than "Warburg's tincture" (see Index) in ʒj doses before breakfast daily. It is disagreeable, but should be taken clear on an empty stomach, in capsules, if too unpleasant. Static electricity is excellent in these cases. Change of climate is desirable, and the prompt cessation of prolonged study and overstrain are demanded. When there is any mechanical impediments to the proper return of blood from the brain, we must do all in our power to keep the heart and lungs working normally. Rest for body and mind is imperative.

DR. F. RUNGE.

Cerebral Pressure; Kopfdruck.—This is described by the above author (*Journal of Mental Science*, 1878) as presenting such symptoms as a feeling of tightness or weight in the head, occasional headache, slight vertigo, difficulty of fixing the attention, and loss of power to do "hard work."

For the treatment of this, he recommends rest of the affected organ, rest from mental exertion, and freedom from disquietude. The patient

should lead a quiet and idle life for two or three months. Change of air and scene is useful, though sometimes the annoyances and discomforts of traveling do harm; but Dr. Runge has found residence in high situations, as in some health resorts of Switzerland, to be contra-indicated. The diet should be non-stimulating; wine, beer, and condiments should be abstained from; and the author has known cases where two or three cups of strong coffee brought back the distress for a week at a time.

Dr. Runge says that many of the vegetarians whom he had met had suffered from Kopfdruck, and preferred an unstimulating diet, because it relieved their malady. He further recommends cold baths to the legs and lower part of the body, which, he thinks, have a more permanent action in calling blood to the surface than hot baths. He observes that parts exposed to the cold are habitually red. He does not recommend cold baths applied to the head, but sometimes uses bromide of potassium where there is a tendency to melancholia. He has tried electricity, but only occasionally found benefit from it.

PROF. HUGO ENGEL, M. D., OF PHILADELPHIA.

Apoplexy. To be able successfully to treat a case of apoplexy, the following points have to be kept in view: There are four varieties—*a* the congestive, *b* the hemorrhagic, *c* the embolic, and *d* the serous variety.

In the congestive variety and in the hemorrhagic variety, when in the latter the same symptoms show themselves, as flushed face, injected eyes, throbbing carotids, stertorous breathing, total unconsciousness, slow, full pulse, decrease of temperature and hemiplegia—the most successful treatment is:

Application of 12 to 16 leeches to anus.

Three to four drops of croton-oil, diluted with same quantity of olive-oil, rubbed into the back of the tongue.

Loosening of tight clothing, perfect rest, position of head higher than the body.

This is the only treatment which has any effect in these two varieties. As soon as the patient revives, means are employed to diminish the arterial flow of blood to the brain, to limit the inflammation ensuing, and to cause absorption in case of clot. These objects are best achieved—

By perfect rest in the recumbent position (the head higher) in a large, well-ventilated room, which is kept dimly lighted. No visitors are allowed; the patient is not permitted to talk on business affairs, all excitement is kept from him, as well as all and every exciting or depressing influence, and he is regularly fed on a low fluid, and concentrated, but bland diet.

The action of the skin is slightly stimulated by daily sponging with tepid water. A bladder with ice is applied to the head, until all acute symptoms have vanished. The bowels are kept open by salines, of which Glauber-salt, administered on the empty stomach in the morning, one hour before breakfast, and in a dose sufficient to insure three motions, is the best.

Fifteen to twenty grains of bromide (sodium or lithium) are administered three times daily, two hours after meals, largely diluted with water.

In case of throbbing pulse and overacting heart, two drops of the tincture of the root of aconite are given every three hours, until active symptoms have ceased; but in case of weak, rapid and irregular pulse, one drachm of infusum of digitalis every four hours is substituted for the aconite, and continued until the pulse becomes normal.

As soon as the active symptoms have all ceased—never before—the above treatment is gradually abolished, and the following adopted:

Gradually increasing doses of corrosive sublimate are continued for about three to four weeks. The following prescription is to be recommended, a tablespoonful representing one-sixteenth of a grain:

- | | | |
|-------|----------------------------|----------------------|
| 1. R. | Hydrarg. bichlor. corros., | gr. i |
| | Sodii chlorid., | gr. xx |
| | Aquæ destillat., | f. $\frac{3}{4}$ vij |
| | Glycerinæ, | f. $\frac{3}{4}$ j. |

M. S. First, a dessert-spoonful in water three times daily, immediately after meals; then three teaspoonfuls (second week), and lastly, one tablespoonful (third week). If then there are still left any residual symptoms (disturbances of special senses, sensation or motion), the hair is shaved off on top of the head, and a large fly-blister (five by six inches) applied to the head, while internally iodide of potassium is administered in gradually increasing doses, beginning with three grains three times daily in water two hours after meals, and continuing up to fifteen, twenty or more grains. At the same time electricity and manipulation may now be applied to the paretic limbs. Until all sequelæ have vanished, the treatment with corrosive sublimate and iodide of potassium is alternately employed, but the general attention to the secretions is kept up.

If the symptoms are such as denote the existence of minute aneur-

isms, or if the hemorrhage is of an oozing character, fluid extract of ergot hypodermically employed is beneficial.

Most cases of the embolic variety evince almost no loss of consciousness; but the patient, a victim of chronic valvular lesion, is generally at once seized with paralysis. It is a real paralytic stroke. Here the face usually is pale, the pulse irregular, the heart excited. It is natural that in such a case the treatment greatly varies from the foregoing.

Such a patient is kept at perfect rest, the same as in the varieties just described. Attention is paid to the secretions, which should not be specially stimulated, but neither be allowed to be locked up.

The physician should first try to prevent a repetition of the accident. The procedure will depend but upon one fact: whether there still is compensatory action in the heart or not. If there is, the patient should be put under the influence of aconite until the excitement is passed. In most cases, however, compensation has ceased or very nearly so, and infusion of digitalis, f.5j, four times daily is the treatment indicated. Besides, the patient should be put on carbonate of ammonia, a remedy counteracting more than any other the tendency of the blood to clot formation; ten to twenty grains three times daily usually are sufficient.

The next object to be achieved is the absorption of the embolus. This is done in the same manner as in the case of a clot from the hemorrhagic variety. The treatment of the paralysis also is the same as of that following the first two kinds of apoplexy.

The picture of the serous variety is very similar to that presented in cases of large hemorrhagic effusions at the base. They mostly are cases of chronic Morbus Brightii, or of dilated heart, or of general fatty degeneration. The patient looks pale, the pupils are widely dilated, the pulse is rapid, feeble and irregular, and the patient is utterly insensible to all external impressions, and there is total want of motion. No treatment is the best, for all these cases die, and never regain consciousness.

Convulsions, if happening in apoplexy, can be stopped with hypodermic injections of morphia.

The last question to decide, but often the most important regarding treatment of apoplexy, is: Bleeding or not bleeding? If a patient dies, and we have not bled him, people say he ought to have been bled; if he dies and was bled, they say he ought not to have been bled. The following rules experience has proven to be the safest:

Bleed, if the pulse is regular and below 80.

Bleed, if the face is flushed and the eyes are injected.

Bleed, if there is hypertrophy or overaction of the heart.

Bleed, if it is a case of congestion in a high liver.

Never bleed if the pulse is irregular and rapid.

Never bleed if the face is pale.

Never bleed if there is valvular lesion of the heart or dilatation of it.

Never bleed if there are convulsions.

Persons with a tendency to apoplexy, or such having had one attack, should observe the following: Never take a cold or hot bath (tepid baths or sponge-baths they may take); never allow the bowels to be constipated longer than twenty-four hours; never strain at stool; keep the circulation quiet.

There is one other form of apoplexy demanding special notice. If a person below the age of thirty-five or forty, with no organic lesion of the heart, not the victim of Morbus Brightii, and not subject to epilepsy or to petit-mal, is suddenly seized with unconsciousness, accompanied by convulsions or not, and simulating apoplexy, the attack is caused by syphilis. Mercurial inunction (daily one drachm) or internal administration of calomel until salivation sets in, followed by a course of iodide of potassium in gradually increasing doses (beginning with ten and increasing up to eighty grains three times daily), until the patient for one year has had no similar attack, and later every spring and every fall the administration of fifteen to twenty gr. doses of the iodide for about three weeks, will prevent further seizures. If the attack was accompanied by convulsions, bromide of sodium in twenty gr. doses three times daily has to be administered, until the specific treatment has made a sufficient impression. During the attack itself, absolutely no treatment is by far the most successful method.

DR. WILLIAM A. HAMMOND.

Apoplexy. In his treatise on *Diseases of the Nervous System*, Dr. Hammond observes that the general treatment of apoplexy should be directed, *first*, to the prodromic stage of cerebral congestion; *secondly*, to the apoplectic seizure; *thirdly*, to its consequences.

1. In active cerebral congestion, *local bleeding*, as cups to the nape of the neck, or leeches to the temples or just inside the nostrils, often affords marked relief. *Cold* is another very useful agent, applied as

ice or cold water to the cranium and nape of the neck. During sleep the head should be elevated. The clothing should be loose about the neck. Sinapisms to the feet and mustard to the epigastrium, are often of service. Internally, the *bromides* of potassium, sodium, calcium or lithium, are indicated.

2. R. Potassii bromidi, ℥j
 Aquæ, f. ℥iv. M.
 A teaspoonful three times a day, in a little water.

This, continued for about ten days, is usually sufficient to relieve the head symptoms. The *bromide of lithium* often acts more promptly than either of the others. Dose, gr. xxx. *Ergot* contracts the cerebral vessels, and is of great value in the treatment of active cerebral congestion, in all its forms, especially in the first or hyperæmic stage. An excellent formula is :

3. R. Sodii bromidi, ℥j
 Ergotæ extracti fluidi, f. ℥iv. M.
 A teaspoonful three times a day.

Or the *ergotin* of Bonjean may be given in pill form, gr. iij-x, *ter die*. When the pain in the head is a prominent feature, prompt relief is often given by

4. R. Extr. fl. paulliniæ, f. 3 ss.-j, p. r. n.

In conjunction with these remedies, the *oxide of zinc*, gr. ij thrice daily, is a powerful agent in relieving congestion and giving tone to the nervous system. It should be taken after meals to avoid nausea.

After the symptoms of congestion have disappeared under this treatment, it becomes expedient to give tonics and restoratives. Among these, strychnia, phosphorus and cod-liver oil stand first. The following extemporaneous mixture is preferable to any of the manufactured syrups or elixirs, with like ingredients :

5. R. Strychniæ sulphatis, gr. j
 Ferri pyrophosphatis, āā 3 j
 Quiniæ sulphatis, āā f. ℥ij. M.
 Acidi phosphor. diluti, āā
 Syrupi zingiberis,
 A teaspoonful three times a day in a little water.

Or the following :

- | | | | |
|-------|----------------------------------|----------|----|
| 6. R. | Strychniæ sulphatis, | gr. j | |
| | Quiniæ sulphatis, | ʒi | |
| | Ferri bromidi, | ʒ ss | |
| | Acid. hydrobrom. (Fothergill's.) | f. ʒiv | |
| | Glycerinæ, | f. ʒ ij. | M. |
- Teaspoonful in water after each meal.

Phosphorus almost always acts well in such cases. It may be given in the form of the phosphorated oil, or the phosphorated resin; or in one of the following formulæ:

- | | | | |
|-------|-------------------|----------|----|
| 7. R. | Olei phosphorati, | f. ʒ ss | |
| | Mucil. acaciæ, | f. ʒ j | |
| | Olei bergamii, | gtt. xi. | M. |
- Make an emulsion. Dose, fifteen drops thrice daily.
-
- | | | | |
|-------|------------------|---------|--|
| 8. R. | Zinci phosphidi, | gr. iij | |
| | Rosar. conserv., | q. s. | |
- Make thirty pills. Dose, one three times a day.

Arsenious acid, in doses of one-fiftieth of a grain, three times daily, after eating, has a favorable action, especially in those forms of cerebral congestion which have been the result of mental exertion or anxiety.

The bowels should be kept in a soluble condition, severe muscular exertion avoided, the stomach should not be overloaded, alcoholic stimulants should be shunned, and the mind maintained tranquil.

When the case demands abstraction of blood, leeches should be applied inside the nostrils. "In no other way can the blood be so advantageously drawn from the brain."

2. During the attack, when there is coma and the ordinary symptoms of the apoplectic condition, there is nothing to be done in the way of medication which can afford the slightest prospect of relief. Blood-letting, purgation and the iodide of potassium, do no good at all, and the first two probably harm. The patient should be kept perfectly quiet, with the head well elevated, and in a room with the temperature about 60°, and well ventilated. The bowels, if not moved naturally every day, may be emptied by an enema of warm water; the urine should be drawn off with a catheter, if required; the strength, if feeble, as indicated by the pulse, should be kept up by the cautious use of stimulants; and if the patient is restless and does not sleep well, some one of the bromides should be administered. The food should be of the most

nutritious character and taken frequently. If symptoms of inflammation make their appearance, cold to the scalp and blisters to the neck are called for.

3. In about two weeks after the stroke it will be proper to take action to restore the power of motion. The means are passive motion, strychnia, phosphorus and electricity. The muscles of the affected limb should be kneaded and the joints flexed and extended, and the surface rubbed every day for five or ten minutes at a time. Strychnia should be administered, preferably by hypodermic injection, in doses of $\frac{1}{2}$ of a grain once a day. In many old cases of hemiplegia, this has an excellent effect. Phosphorus may be employed as follows:

9.	R.	Zinci phosphidi,	gr. iiij	
		Extracti nucis vomicæ,	gr. x.	M.

Make thirty pills. One three times a day.

No agent is so valuable, however, as electricity. The induced current is usually sufficient for recent cases; but old cases may require the primary current. The poles, terminated by wet sponges, should be applied to the skin covering the paralyzed muscles, or to the nerves. The interrupted current should be applied. The same agent is valuable to restore sensibility to the parts.

J. HUGHLINGS JACKSON, M. D., LONDON.

Apoplexy.—Pending the seizure the chief thing is to keep the patient quiet. Two drops of *croton oil* should be put on the tongue and the urine drawn off, if it does not pass freely without help. The application of blisters to the back of the neck, or of mustard plasters to the calves of the legs, is a common practice, with no evidence to show that these applications are of service. Still, in certain cases of chronic cerebral disease, blisters relieve the patient of severe pain in the head; and it is not possible to deny that they may be of service when the circulation of the brain is quickly disturbed after a clot. Were our author to use blisters, which he never does, he would only employ them where there is secondary disturbance of the circulation. It is hard to believe that mustard plasters to the calves of the legs can be of any service or disservice. Dr. JACKSON never prescribes any medicine, except croton oil, by the mouth.

If the patient is found in, or has passed into, a condition in which the face is flushed, the temperature above 100, the veins prominent, and the

respiration and circulation largely disturbed, treatment, beyond, perhaps, purgatives, is of little use. If, however, the pulse be not very rapid, little more than 100, and if it and the respiration be regular, it is not unreasonable to think that the system may be relieved—though the ultimate size of the clot may not be influenced—by *bleeding*. The practice of bleeding in cerebral hemorrhage, deprecated by TROUSSEAU and TODD, must be exceeding rare in England, as in the whole course of Dr. JACKSON's life he has seen but one person bled for cerebral hemorrhage. If the pulse be very high, 120, 130 or 150, or if it be weak, venesection must not be thought of.

This is about all of the little that can be done for patients who are in the apoplectic condition. When this condition is passed, we are practically unable to help in the immediate difficulty of infusion of blood in nervous tissue. We have still, as in many other diseases, to improve the general health. The proper care of a patient who has a clot of blood in his brain, and who is liable to have more effused, consists in attending to his diet, excretions, sleep and exercise. Yet there is nothing in this peculiar to the treatment of cerebral hemorrhage. For the immediate lesion, blood in nervous tissue, our author knows of no direct treatment.

DR. ALLEN M. HAMILTON, OF NEW YORK,

In his work on *Nervous Diseases*, speaks very highly of *hydrobromic acid* in cerebral hyperæmia. He gives the following formula:

10.	R.	Sol. acidi hydrobromici,		
		Elixiris simplicis,	āā	f. ʒij.
				M.

A teaspoonful before each meal.

As a prophylactic measure, he has patients apply cold (ice-bag, etc.,) to the back of the neck for fifteen minutes, night and morning, "and finds it succeed admirably." It is of great importance to improve the cutaneous circulation by the flesh-brush, cold, the Turkish bath, etc.

His treatment of the attack and sequelæ agrees substantially with that given by HAMMOND.

DR. ROBERTS BARTHOLOW, OF PHILADELPHIA.

To allay cerebral irritation and excitement, *prussic acid* has been employed with benefit. We should use from two to five minims of Scheele's dilute acid, giving it at first at short intervals (every quarter

of an hour) and, when effects are produced, every hour or two. It is also employed subcutaneously, in five-minim doses.

NOTES ON REMEDIES.

Acidum Arseniosum is of benefit in strong plethoric subjects with a tendency to apoplectic congestion, but it is not applicable to old, weakly persons. It is supposed to act by reducing the excess of the red globules of the blood. HAMILTON esteems it most when the circulation is sluggish, and suggests.

- | | | | |
|--------|-----------------------|----------|----|
| 11. R. | Sol potass. arsenit., | f. ʒ ij | |
| | Quiniæ sulph., | ʒ ss | |
| | Acidi sulph. aromat. | q. s | |
| | Aquæ anisi, | f. ʒ iv. | M. |
- A teaspoonful every four hours.

Acidum Hydrobromicum. Fothergill's hydrobromic acid is of great value in cerebral hyperæmia. Dr. D. C. WADE, of Detroit, combines it with ergot as follows :

- | | | | |
|--------|----------------------------|----|---------|
| 12. R. | Ergotæ extracti fluidi, | | |
| | Syrupi simplicis. | āā | f. ʒ ss |
| | Acidi hydrobromici diluti, | | f. ʒ j. |
- M. A teaspoonful in water as required.

He says of this that it surpasses any other combination he has ever tried, rapidly reducing symptoms of plethora and the determination of blood to the brain.

Aconitum. In simple apoplexy in plethoric persons, with full pulse, hot skin and flushed face, Dr. C. D. F. PHILLIPS thinks aconite "decidedly the best" remedy that can be employed, ℥j. of the tincture of the root every half hour. It should be avoided when the heart is weak and the face pale. (*Materia Medica*, 1879.)

Barii Chloridum in doses gr. i-ij, is favorably spoken of by BARTHOLOW in cerebral hemorrhage.

Bromides. These are recommended by all recent writers for diminishing arterial tension and cerebral blood pressure. They may be combined with ergot (F. 3), with aconite if cardiac action is violent, or with digitalis if the heart is weak ; as

- | | | | |
|--------|-------------------------|-------------|----|
| 13. R. | Tinct. aconit. radicis, | f. ʒ j-ʒ ij | |
| | Sodii bromidi, | ʒ iss | |
| | Aquæ menth. piper, | ad f. ʒ iv. | M. |
- A teaspoonful three times a day. (HAMILTON.)

Colchicum is a serviceable remedy in acute cerebral congestion in plethoric subjects.

Digitaline—granules (one sixtieth of a grain bis die) will, not unfrequently, permanently relieve the congestive form of hemicrania.

Ergota is a powerful cerebral anæmiant. It may be given in 5-gr. doses of the watery extract thrice daily, alone or combined. Dr. N. S. FOSTER (*Lancet*, September, 1878,) uses ergotine, gtt. xij of a solution gr. x, to aquæ f3j, injecting it deeply into the muscles, to avoid suppuration. When the patient was seen early, his success was striking.

**Hydrargyri Chloridum Corrosivum*. Dr. HEADLAND recommends for the threatenings of apoplexy in old age. (*e. g.*, vertigo, confusion of ideas, and general embarrassment of the mental faculties,) corrosive sublimate in doses of gr. $\frac{1}{2}$ in solution, three times a day, for three or four weeks.

Hydrargyri Chloridum Mite. Administered as recommended by Dr. J. CORLAND, (grs. x-xv, mixed with a few grains of gamboge, rubbed up with butter, and placed at the root of the tongue,) calomel, although a speedy and effectual purge, is not unattended with danger. Alarming ptyalism has followed this treatment.

Potassii Iodidum has been recommended in cerebral apoplexy, on account of its eliminative power, after the acute symptoms have subsided.

Sinapis. In cases of apoplexy from over-distention of the stomach, mustard, in doses of a tablespoonful or less in a tumbler of warm water, is a good and quick emetic, which, by removing the exciting cause, is of essential benefit.

Strychnia. Dr. MARSHALL HALL has derived great benefit from the use of acetate of strychnia in the threatenings of apoplexy. He advises the following formula :

14. R.	Strychniæ acetatis,	gr. j
	Acidi acetici,	mxx
	Alcoholis,	f. 3 ij
	Aquæ,	f. 3 vj.

Dose, ten drops, combined with a rigid system of mental discipline, diet, etc.

Veratrum Viride. Dr. ROBERTS BARTHOLOW says there can be no doubt that this drug renders valuable service in acute parenchymatous congestion of the brain.

GENERAL REMEDIES.

Blood letting is no longer the routine practice in apoplexy. M. CONSTANTIN PAUL however, believes that bleeding may be of service in apoplexy, not as affecting the cerebral lesion, but as acting on the apoplectic condition, on the asphyxia and stasis of blood in the veins. He has seen the condition of such patients very markedly improved. It is now generally recognized that there is less danger in not bleeding at all, than in taking blood in doubtful cases, in which there may be some of the indications for bleeding.

The *contra-indications* of bleeding in apoplexy are an age over sixty ; a feeble, very frequent, intermitting, slow or large pulse, or one inclined to double beat ; a respiration labored and accompanied with *cold* per-

spiration; great mobility of the nervous system, with weak muscles, whether the body be thin or corpulent; an attack soon after a full meal, or after great mental or bodily fatigue.

The *indications* for bleeding are a quick, wiry, resisting pulse; flushed countenance; warm perspirations; noisy breathing; a tendency to spasmodic muscular contraction; and an age under sixty.

Cathartics. In all cases of cerebral plethora, the bowels should be kept soluble. Aloetic preparations are said by some writers to be specially derivative; others prefer the regular use of Friederichshall, Hunyadi Janos, or other similar mineral water. In an apoplectic attack, purgatives are indicated. One to two drops of croton oil on the back of the tongue, will bring on purgation when the patient is unconscious. When conscious, $\frac{1}{2}$ grain of elaterium with $\frac{1}{4}$ gr. nux vomica makes an efficient pill.

* *Diffusible stimulants* may be given with advantage, particularly when the pulse has a double beat, in atonic cases where bleeding is contra-indicated. *Aqua ammoniæ* is one of the best of these: ℞ij-xv may be administered in water, and the vapor also applied to the nostrils. Or f. 3 ss-j of the *aromatic spirit of ammonia* may be given in water or camphor mixture. Or grs. v of the carbonate of ammonium may be ordered.

Enemata. Terebinthinate and other enemata often afford relief, not only by removing scybala, but by their powerful revulsive action.

EXTERNAL REMEDIES.

Counter-irritation. *Capsicum cataplasms* to the feet are powerful and excellent revulsives, which will, however, cause vesication if kept on too long. *Sinapisms* to which have been added powdered capsicum, or oil of turpentine, may, in most instances, be applied with evident benefit, to the soles of the feet and the inner parts of the calves and thighs. *Blisters* applied to the calves of the legs and sometimes to the nape of the neck, are of benefit in *simple* or "*serous*" apoplexy, but of little advantage in *sanguineous* apoplexy. An open blister at the nape of the neck (or better, a seton or issue there,) is often very useful when there is a tendency to apoplexy. *Turpentine stupes* to the extremities are frequently productive of benefit.

Electricity is considered by Dr. HAMMOND an invaluable agent for the relief of the sequelæ of an apoplectic attack, (p. 28.) It will generally be found to restore the sensibility of the part before removing the motor paralysis. The electric wire brush and the wet sponge are the appliances most frequently demanded. In recent cases this will almost invariably prove effectual. Hyperæsthesia, which is sometimes a troublesome legacy of the cerebral extravasation, may be similarly managed.

Galvanism.—According to Dr. ROBERT BARTHOLOW, galvanization of the brain and of the cervical sympathetics is one of the measures to be resorted to in acute active or passive congestion of the brain.

Galvanization in cerebral hyperemia has been found of much efficacy by Dr. C. H. HUGHES, of St. Louis. He says that an essential property of the

constant descending galvanic current in induced cerebral hyperæmia is that of a tranquilizer of irritable nerve tissue, secondarily contributing to the contraction of over-distended vessels. It acts on the irritable brain like bromides, hyoseyamin and chloral, vaso-motor results being secondarily induced when there is over vascular distention as well as primarily accomplished. (*Alienist and Neurologist*, 1884.)

Ice applied to the head while the feet are immersed in hot water, is sometimes productive of benefit, but must be employed with great caution in debilitated or old subjects. Dr. BARTHOLOW is of opinion that ice is often used without due discrimination. When the face is pale, the surface cool and the circulation depressed, cold applications to the head are harmful. On the other hand, Dr. JAMES M. GOODHART, of Guy's Hospital, London, believes that the indication to restrain the hemorrhage is so imperative, that the application of cold locally is very generally called for, although its employment seems rather contradictory to the general course of treatment otherwise required.

* *Leeches*. In threatened apoplexy arising from the suppression of an habitual discharge, as from piles, leeches applied to the verge of the anus often afford prompt relief. If the threatened attack arise from suppression of the menses, the leeches should be applied to the inner side of the thighs. Sir HENRY HOLLAND states that he knows of no mode in which a given quantity of blood can be removed with equal good effects. Leeches to the verge of the anus give more immediate benefit than three times the number to the temples or elsewhere.

CEREBRAL ANÆMIA.

DR. EDWARD C. MANN.

According to this author (*Psychological Medicine*, 1884), the first essential of treatment in this disease is to obtain *rest* of both body and mind. Alcohol, preferably in the form of old rye whisky, is useful. The constant galvanic current is a valuable remedy. *Strychnia*, in doses of gr. $\frac{1}{32}$, is one of the most certain drugs. Cod-liver oil and phosphorus are also indicated.

Where *spinal anæmia* is also present, the first great means of cure resides in the judicious employment of counter-irritation to the affected portions of the spine. The compound mustard liniment is one of the most suitable agents for this purpose. It should be applied on flannel and the whole covered with oil silk.

A full nourishing diet is necessary, and as soon as the system is some-

what improved, change of scene, traveling, and cheerful society are to be recommended.

NOTES ON REMEDIES.

Arseniosum Acidum, in the form of Fowler's solution, gtt. ii-ij., thrice daily, combined with the same quantity of laudanum, three times a day, has been exhibited with excellent results in some cases by French physicians.

Auri Chloridum is stated by Dr. BARTHOLOW to have a high degree of utility when there is present the condition of cerebral anæmia.

Ferrum. As in all cases of anæmia, the preparations of iron are of use. The chalybeate waters offer an excellent method of administering it. Where there are epileptoid seizures, BARTHOLOW recommends the following :

15. R.	Potassii bromidi,	℥j
	Ferri bromidi,	gr. iv
	Aquæ,	℥ij
	Syr. simplicis,	℥vj.

A tablespoonful twice daily.

Nitro-glycerine has a remarkable power to develop cerebral hyperæmia, and is therefore of value in transitory cerebral anæmia. The dose is gtt. $\frac{1}{100}$. (HAMMOND.)

Phosphorus and the Phosphates are applicable in these cases through their stimulating action on the nutrition of the brain substance.

CHOREA.

DR. EDWARD C. MANN.

The treatment adopted by this author he reports as having given most satisfactory results (*Psychological Medicine*, 1884). He endeavors to give the nervous system *rest* and *nutrition*, the former by avoidance of excitement, early hours, and warm baths at bed time; the latter by using *phosphates*, *cod-liver oil*, or the oil in connection with the phosphide of zinc, gr. $\frac{1}{10}$, thrice daily. Gentle gymnastic exercises are valuable and should not be neglected.

The specific, however, is *arsenic*, in the shape of Fowler's solution, *used hypodermically*. Three minims, mixed with an equal amount of water to avoid irritation, are injected subcutaneously for a week every other day, increasing two minims each week, and in a month or six

weeks a cure will be obtained, though in old cases sixty or seventy days may elapse before a cure is complete. Marked improvement is always noticed from the first by this plan of treatment. The gastric disturbances which are noticed when the medicine is given by the mouth are avoided, and the good results are obtained much more rapidly.

G. H. BARLOW, M. D., LONDON.

In ordinary cases the exhibition of purgatives to keep the bowels freely open, and the *sulphate of zinc*, in doses gradually increased from gr. j-xij-xv-xx, or even more, will effect a cure; when, however, the sulphate has been used in these large doses, its sudden discontinuance seems to be felt by the system, and a return of the symptoms ensues; the best rule, therefore, for its exhibition is as follows:

The bowels being kept open, the sulphate should be commenced in doses of gr. j, for a child æt. twelve years, and this should be increased by the addition of gr. j to each dose daily, until it either causes sickness, or there is an obvious diminution of the choreic movements. In the former case, the dose should be diminished by at least one-half, and so continued for several days, with a view of establishing a tolerance; but if, on the other hand, there be marked improvement, it should be no further increased, but continued without alteration until either the improvement ceases—in which case it should be again gradually increased—or the disease has altogether subsided. When the latter is the case, the dose should be diminished day by day, rather than discontinued suddenly, as by following the former course, we have less reason to dread a relapse. When anæmia is present, *iron* has more control over the disease than zinc. The sulphate of iron may be combined with the zinc.

PROF. BOUCHUT, OF PARIS.

This eminent writer classifies the forms of chorea and their treatment as follows:

1. *Verminal chorea*, from worms. Santonine, or if tænia be suspected, male fern, or kousso.

2. *Dental chorea*, from irritation of teeth. Remove the offending tooth.

3. *Anæmic chorea*, seen in convalescence. Iron, quinine, arseniate of iron and soda, &c.

4. *Chorea from moral cause.* Isolation; rest, antispasmodics, firm government.

5. *Electric chorea*, where the affection is partial, limited to a single limb and monoplegic. Subcutaneous injections of morphia, and sulphate of eserine, tincture of gelseminum, &c.

6. *Organic chorea*, that is, resulting from the organic nature of the malady. All remedies which have power to disperse the congestion of the spinal cord, as ether-spray, douches, sulphur baths, vesicatories, cauterization, &c. M. Bouchut prefers dotted cauterization with *fusain*, *euonymus europæus*, and daily hypodermic injections of morphia, or valerianate of ammonia. The drugs he employs are opium, emetics, arseniate of soda, chloral hydrate. He lays great stress on procuring sleep, and says that he has made children sleep fifteen and eighteen hours a day for eight and ten consecutive days, by administering *chloral* to them. After breakfast at eight o'clock he gives forty-five grains of chloral. The child is roused for its dinner, two hours after which a similar dose is given; it wakes towards evening, and on being fed, enjoys a natural sleep. He has never seen any ill effects from this remedy in children.

THOMAS KING CHAMBERS, M. D., LONDON.

This author recommends Fowler's solution, with cod-liver oil and iron, if the general indications demand them. He adds that injudicious management of patients afflicted with chorea frequently protracts the case. One of the most common forms of injudicious management is the fixing the attention of patients upon their infirmity, by telling them how bad they are, offering unnecessary help, etc. They should be encouraged to make every exertion to direct the movements of the limbs; as by slow walking to music, carrying trays and crockery, and other things that demand care. In order that their attention may be withdrawn from their deficiencies, looking-glasses and the distressing sight of other choreics should be avoided. They should be got away from home as soon as possible. Sent under the care of a judicious person to the seaside, or anywhere else for an excuse, children often recover rapidly; whereas, had they remained at home, they would have continually relapsed.

PROF. J. M. DA COSTA, M. D., PHILADELPHIA.

16. R. Zinci valerianatis,
Cinchoniæ sulphatis,
For twenty pills; one thrice daily.

℞ij
℞j.

M.

The *bromide of iron* has been employed by Dr. DA. COSTA with excellent effects. It should be given in increasing doses, beginning with never less than five grains for a child, and running up the dose rapidly to twenty grains three times a day. He likes it better than any other one article in the treatment of chorea. It may be given in plain syrup and water, in the form of a pill, or better, in an effervescing powder. It not only affects the chorea, but also influences the nervous system as a sedative, quieting it and giving the patient rest. It is a valuable agent in treating the incontinence of urine in children, and it was in a case of this kind, complicating chorea, that he first observed its value; being surprised to see that, as the symptom which led to its administration improved, the chorea also diminished, and soon disappeared. Since then he has used it almost continuously. In answer to the question whether it is the bromide or the iron that benefits, he thinks it is the combination; that neither *alone* accomplishes the result; for it will benefit cases that have previously taken iron without improvement; and, as regards the other bromides, we certainly cannot claim for them any especial value in chorea.

17. R. Cupri ammoniati, gr. $\frac{1}{4}$.
In pill thrice daily; to be gradually increased to gr. j.

18. R. Extracti cimicifugæ, gtt. xx.
For one dose, three times a day.

PROF. WM. A. HAMMOND, M. D., NEW YORK.

Of the preparations of zinc, Dr. HAMMOND prefers:

19. R. Zinci bromidi, $\overline{3j}$
Syr. simplicis, f. $\overline{3j}$. M.
Ten drops three times a day, increased as rapidly as the stomach can bear it.

After the disappearance of the choreic symptoms, the doses should be diminished in the same gradual manner they were increased. Dr. HAMMOND's main reliance, however, is upon *strychnia*.

20. R. Strychniæ sulph., gr. ij
Aquæ, f. $\overline{3j}$. M.
Five drops three times a day to a child from ten to fifteen years of age.

This dose should be increased one drop daily until the physiological

effects of the medicine are manifested, as evidenced by stiffness of the legs and neck. When this takes place, the dose should be at once reduced to five drops, and increased again as before. TROUSSEAU also recommends this drug in high terms.

He also found *ether-spray to the spine* of unequivocal value. The whole spine is exposed, and the ether-spray is thrown upon it from the occiput to the sacrum for about ten minutes every day or every other day.

DR. A. W. HAMILTON, NEW YORK CITY.

This writer gives the results of his observation with various drugs in the *New York Medical Journal*, February, 1877. He has found *strychnia* serviceable carried up to the point where stiffness of the spinal muscles is arrived at. Next to this comes *arsenic*. It must be given in large doses. Occasionally, digestive troubles are produced very quickly by this drug, and then *strychnia* may be substituted. Cold to the spine cannot be overestimated as a plan of treatment; either the ether-spray, or ice-bags every day, allowing them to stay on about ten minutes. PERROUD, who has used the ether-spray, makes applications from four to eight minutes in duration every day. In thirty-five cases treated with the ether-spray, from fifteen to twenty applications produced permanent benefit. The spray should be directed to the upper part of the cord, over the upper cervical vertebræ. *Eserine* has been recommended, and BOUCHUT has given the results of four hundred and thirty-seven cases, two hundred and five who took it in pilular form, and two hundred and thirty-two hypodermically. The average dose was from two to five milligrammes. He obtained temporary benefit, which seemed to wear off; but when the drug was repeatedly administered, he accomplished many cures. He reports twenty-three cures by an average of seven injections. It is a dangerous remedy, however, and produces severe gastric symptoms. *Conium* is occasionally efficacious, but its effects are temporary. *Phosphorus*, with cod-liver oil, is a most valuable curative agent, and in cases where everything else failed, it has succeeded. In some instances, nothing does good. Put them in a dark room, and keep them perfectly quiet. The diet should be regulated with judgment. Plenty of fresh air and sleep come next, and absolute mental rest must be enforced. The school-books and the school-room are to be parted from, and agreeable

diversions planned. An excellent auxiliary to medication is the salt-bath. A handful of rock-salt in the water, and the energetic use of the rough towel, will infuse a tone and vigor that will soon become apparent. Decided medication is useless in these patients when their personal habits are not looked after.

THOMAS HILLIER, M. D., LONDON.

- | | | | |
|--------|------------------------------|----------|----|
| 21. R. | Liquoris potassii arsenitis, | mij | |
| | Potassii bicarbonatis, | gr. iij | |
| | Potassii iodidi, | gr. ij | |
| | Aquæ camphoræ, | f. ʒ ss. | M. |

For one dose, thrice daily, to children aged five, for aggravated chorea, attend with severe pains in the limbs, and rheumatic persistent swellings.

Arsenic in full doses is a valuable remedy in a fair proportion of cases, but in some instances it entirely fails. *Iodide of potassium* is useful when the patient is subject to chronic rheumatism.

Occasionally purgatives and tonics, especially *iron*, are attended with much success. *Strychnia*, so highly recommended by TROUSSEAU, seems, to our author, to be really injurious in the acuter stages of the disease; in the more chronic form, and where there is a tendency to paralysis, it is of service. Iron and strychnia may be combined thus:

- | | | | |
|--------|-------------|----------|----|
| 22. R. | Strychniæ, | gr. ½ | |
| | Vini ferri, | f. ʒ iv. | M. |

A dessertspoonful, thrice daily, to a child ten years of age.

Narcotics, such as opium, belladonna, cannabis indica, or conium, are of little or no use. Antispasmodics, such as valerian and assafœtida, are also useless.

Our author has seen good results from the employment of *baths of sulphuret of potassium*:

- | | | |
|--------|----------------------|--------|
| 23. R. | Potassii sulphureti, | ʒ iv |
| | Aquæ (90°F.), | o xxx. |

For a bath; the patient to remain in it for an hour daily.

This bath is also recommended by Dr. FELIX VON NIEMEYER, when there is anæmia.

Gymnastic exercises, shampooing and passive movements, are of service. As many muscles as possible should be exercised, without fatigu-

ing any of them. *Shower baths* are useful in the latter stages, when the patient is not timid, or too much excited by them.

Dr. NIEMEYER quotes BENEDIKT, who declares that out of more than twenty cases of chorea, treated by him by the *constant galvanic current*, not one has failed to recover. The current which he employs is just strong enough for the patient to feel it distinctly, and he applies it along the spine, the patient standing erect. Painful currents aggravate the symptoms.

PROF. WILLIAM AITKEN, M. D., EDINBURGH.

The indications of cure are: 1. To remove, if possible, all morbid states of the body which may tend to aggravate the disease, such as constipation, anæmia, amenorrhœa, worms. 2. By well-regulated purgative medicines to subdue any cerebral congestion. 3. To sustain the strength and improve the vigor of the nervous system by tonic and stimulant medicines, by food, and by the cold bath.

24. R. Camphoræ, ℥v
Syrupi, q. s. M.

Divide into twenty pills. One three times a day. Useful after discharges have become healthy by the action of the purgatives.

25. R. Spiritûs ætheris nitrosi, f. ℥j
Aquæ camphoræ, f. ℥iij. M.

Tablespoonful three times a day.

Many young women, who attribute the attack to fright, get well under this treatment.

DR. GEORGE T. STEVENS, OF NEW YORK.

The novel view is advanced by this writer, (in the *Transactions of the New York Academy of Medicine*, 1876,) that chorea arises from irritation dependent upon anomalous refraction of the eye, and in a very large proportion of cases upon hypermetropia. Hence in regard to treatment the first and great indication is to correct the faulty refraction by the use of proper glasses. This will often relieve the patient at once. If glasses for any reason cannot be supplied or used, the child's eyes may be covered, for it is a well-known fact that when the patient with chorea sleeps, the choreic movements often cease; this is doubtless because the eyes are at rest.

The *Calabar bean* has of late years been found of value in the treatment of chorea; and, when we remember that its peculiar and charact-

eristic effect is to stimulate the ciliary muscles, we shall see that its use is rational.

As children and others suffering from chorea usually also suffer from general want of vigor, tonics, chalybeates, and arsenic are generally indicated.

The success which Dr. STEVENS has met with in a number of cases, goes to show that, in some instances at least, this view is well founded.

HÔPITAL DES ENFANTS, PARIS.

The means of treating chorea more particularly employed at this hospital, are the *tartar emetic* plan advocated by GILLETTE, and still employed by ROGER; *gymnastics*, advocated by BLACHE; and *sulphurous baths*. The tartar emetic plan is this: Three days' treatment: during the first day, three to six grains of the remedy, (according to age;) during the second, five to ten grains; during the third, eight to twelve grains; then three days' rest; then three days' treatment; commencing the new series with a dose one grain stronger than in the first series—to end with a third turn, often attended by good effects. Gymnastics are highly beneficial in milder cases of chorea, and in the declining stage of the severer forms. As much may be said of sulphurous baths. The other means employed at the hospital are: *Opium*, *chloroform inhalations*, *salt baths*, *hydrotherapy*, *tonics*, etc. Strychnia, so warmly advocated by Prof. TROUSSEAU, is scarcely ever employed at the Hôpital des Enfants.

TREATMENT BY ENFORCED REST.

A number of writers have reported favorably of the treatment of chorea major by absolute and prolonged rest in bed.

Dr. W. F. WADE, (*British Medical Journal*, December 28, 1872,) gives the patient a sufficiency of good plain wholesome food; keeps her in bed, in order that the muscular and nervous systems may have the greatest attainable repose, and thus prevent wear and tear. This also secures a uniform warm temperature. He prescribes two tablespoonfuls of compound decoction of aloes twice or thrice a day, if there is constipation. This is all that is necessary in many cases; but if improvement is not going on satisfactorily, he gives sulphate of zinc, in doses of two or three grains, three times a day, in water, and increases the dose until each dose produces slight nausea.

In the *Chicago Journal of Nervous Diseases*, April, 1877, Dr.

RANSOM DEXTER relates a case of chorea major in which many remedies had proved inefficient, and which was much improved by such absolute rest as was obtained by closing all means of external irritation. The patient was kept in bed, the room was darkened, the eyes bound up, and the ears plugged with cotton. The result of this procedure is very satisfactory.

Dr. JOHN VAN BIBBER, of Baltimore, has tried a modified plan of this treatment with success, (*Transactions of the Medical and Surgical Faculty of Maryland*, 1878.) He states that mere confinement in bed in a hospital ward is not sufficient. The presence of other patients, the passing to and fro of nurses, the bustle of administration, and numerous distractions, would give to a choreic patient much cause for restlessness. Unless the rest is without interruption or irritation, it will be found, if not useless, of little benefit. Hence, it is necessary to provide an apartment for the patient where the blinds can be closed to avoid irritation by the eyes, absence of noise to avoid excitement by the ears, absence of any conversation to render the mind as quiet as possible. Add to these requisites an attendant who can be still and patient, the advantage of good nourishment and necessary medicines, and this is a system which is essentially good and curative. To prevent the patient being injured by too long confinement, he employs massage, once or twice daily.

NOTES ON REMEDIES.

**Acidum Arseniosum* is highly recommended by numerous authors. Dr. A. FLINT says it has stronger testimony than any other drug. Dr. BEGBIE, who, in an experience of thirty years, states that he has never known it to fail, prescribed gtt. v. of Fowler's solution twice daily, after a meal, and added one drop to the dose every day, until the specific effects appeared, when he suspended it for a time. Dr. EUSTACE SMITH, of London, remarks that the tolerance of children for arsenic is a matter of common observation, and this tolerance is especially marked in the case of a non-febrile disease, such as chorea, where there is no increased irritability of the digestive organs. To a child between the ages of five or six and twelve, the subject of this complaint, Fowler's solution may be given in doses of 10 minims three times a day, directly after meals. The influence of this treatment upon the disorder is seen almost immediately, and it is rare for any of the physiological effects of the drug to be observed. By this means, cases of the disease which had resisted smaller doses of arsenic, may be cured in a few days, and even severe cases seldom last longer than a fortnight or three weeks.

Anilin, first employed by Dr. JAMES TURNBULL, is also recommended by Dr. F.

E. ANSTIE. The dose is gr. i-ij. thrice daily, of the anilin sulphate. It is a white powder, easily taken.

Antimonii et Potassii Tartras is recommended by Dr. C. WEST, in acute cases, given in full doses. Dr. THOMAS HILLIER, however, condemns its use (which he has found inutile), as a depressing remedy, ill adapted to a disease like chorea, of weakly, badly-fed children. In French practice, it is advocated by GILLETTE and ROGER. (p. 41.)

Apomorphia has had a limited but favorable trial in this disease.

Argenti Nitras is much less employed than formerly, before the discovery of other remedies of greater efficiency.

**Arsenic.* Prof H. NOTHNAGEL recommends the following—

26. R. Liq. Fowleri,	f. ʒi
Aq. destill.,	f. ʒ iv.

M. S.—Five drops to be given in a tumbler of water immediately after each meal, and the dose to be increased by three drops every day until it reaches thirty drops, after which it is to be slowly diminished.

Dr. C. E. ARMAND SEMPLE reports two cases cured by the following formula :

27. R. Liquoris arsenicalis,	℥ iij
Mist. ferri perchloridi,	ʒ ij ter die.

Belladonna, though formerly much employed, is characterized by Dr. THOMAS HILLIER as of "little or no service," and such seems to be the general modern verdict.

Brominium, see *Potassii Bromidum*.

Camphora is a remedy in which many physicians have considerable confidence. The *monobromide* is said by Dr. LAMANDE, of Paris to cure some cases with great rapidity.

Cannabis Indica has been found useful in some cases. In the sleeplessness attendant upon severe chorea, the tincture has proved an excellent hypnotic, in full doses.

Chloral, by its hypnotic effect and the consequent ameliorating influence of the sound sleep it produces, has been found of benefit in many cases.

Chloroformum Purificatum. Chloroform inhalations are not approved of by Dr. C. B. RADCLIFFE, unless alcoholic stimulants are given before the inhalation. Friction along the course of the spine, night and morning, with a liniment composed of equal parts of chloroform and oil of almonds, is beneficial in some cases.

**Cimicifuga* is highly recommended by many physicians. Prof. A. STILLÉ considers it as one of the most valuable remedies in this disease in pure uncomplicated cases, given in doses of sufficient strength to develop its specific effects. Dr. SYDNEY RINGER, however, finds it only effectual in cases of rheumatic origin, and even in these he prefers arsenic.

**Conium* is very highly lauded by Dr. JOHN HARLEY. He gives the succus in full doses, and asserts that, apart from its effects on the motor centre, conium possesses no direct influence on the circulatory, nutritive or secretory functions, and that its use can be prolonged with safety, it being entirely destitute of any cumulative action.

Conii Succus. To insure good effects from this drug, Mr. J. F. W. SIEK, (*Lancet*, 1883,) says that it must be given in large doses, and that its action must be sustained by frequent repetitions of the dose at short intervals. The uncertainty of the action of given specimens of succus conii necessitates great care in its administration, and militates against its general adoption. But cases in which neither chloral nor morphia have any effect may arise, and in which, as in the above, succus conii may prove efficacious. Drs. CLIFFORD ALLBUTT, EDDISON and CHURTON have obtained good results in the treatment of chorea, with violent movements, by large doses of succus conii. The patients took from ʒij every hour to ʒss every four hours during two or three days. It was given sometimes alone, sometimes in combination with morphia or bromide of potassium. It seems that the best results are obtained by large doses at first, until the system is thoroughly under its influence, when smaller doses, frequently repeated, will keep its action. No toxic effects were ever noticed in these cases.

Cupri Sulphas has been thought useful, but is doubtless of less value than conium, and a number of other remedies.

Cuprum Ammoniatum once enjoyed a high reputation in the purely nervous form of chorea. (F. 17.)

Dracontium. Dr. H. C. WOOD, Jr., (*New York Medical Journal*, December, 1877,) tested the use of dracontium, or skunk cabbage, in chorea. He used sixty to ninety drop doses of a tincture of the fresh root three times a day, and had good results.

Electricity. Prof. H. NOTHNAGEL highly recommends the constant current, combined with tepid bathing or the application of ice-bags to the spine.

**Ferri Bromidum* is much used by Prof. DA COSTA.

Ferri Carbonas is often a useful remedy.

Ferri Oxidum Hydratum in large doses, (ʒss-j increased to ʒiii-iv, every six hours,) is highly recommended by Drs. ELLIOTSON and WILKS, conjoined with the use of active purgatives. Children readily take ½ drachm doses in treacle.

Ferri et Quiniæ Citras has also been found useful. HAMILTON says digitalis often increases the benefit of iron preparations.

Hyoscyamin in doses of ʒ_{ss} gr., at first twice daily, increased in frequency, is praised by Drs. OULMENT and LAURENT.

Hypophosphites are recommended by Dr. RADCLIFFE, combined with cod-liver oil.

**Morrhue Oleum* is very favorably spoken of by Drs. F. E. ANSTIE, C. B. RADCLIFFE, and THOMAS HILLIER, in cases of chorea in scrofulous

children or those of spare habit. Dr. RADCLIFFE generally gives the oil in conjunction with hypophosphite of soda (gr. v-viij), making the draught containing the hypophosphite the vehicle for the oil. Sometimes he adds carbonate of ammonia to the solution of the hypophosphite; and sometimes dissolves camphor in the cod-liver oil, thus masking the taste of the oil and making the stomach more tolerant of it.

Moschus has been employed with benefit, but generally fails.

Nux Vomica was employed by ROUGIER in minute doses gradually increased. TROUSSEAU preferred strychniæ sulphas (which see).

Oleum Terebinthinæ is sometimes given with the view of obtaining its anthelmintic, purgative and stimulant effects.

Opium is tolerated in very large doses in this disease, but Dr. RADCLIFFE's experience is not favorable to its use.

Phosphorus is favorably reported upon by a number of physicians, as a remedy in this disease.

Physostigma has been used with success by Drs. HARLEY and OGLE. The latter gives:

28. R. Pulveris physostigmatis, $\mathfrak{z}\text{j}$
Alcoholis, f. $\mathfrak{z}\text{j}$. M.

Begin with 20 minims, thrice daily, and increase, by 10 minims a dose, to f. $\mathfrak{z}\text{j}$.

**Potassii Sulphuretum*. Drs. HILLIER and VON NIEMEYER recommend baths of this salt (F. 23.)

Potassii Bromidum has proved a failure in the hands of Dr. RAMSKILL, but has been favorably reported upon by a number of French physicians.

Potassii Iodidum is considered useful by Dr. HILLIER in cases of chorea occurring in children subject to chronic rheumatism. It is also beneficial in patients tainted by syphilis or scrofula.

Quiniæ Sulphas has been employed in some cases with advantage.

**Santonin* proves often valuable, by disclosing the unsuspected cause of the choreic movements, viz., worms.

Sodii Arsenias is useful in those cases in which other arsenical preparations are badly borne.

Sodii Salicylas has been found of some value by Dr. S. WEIR MITCHELL; and it has been well spoken of by Dr. DRESCH. (*Bull. Gen. de Ther.*, 1879.)

Stanni Chloridum has been employed in doses of gr. $\frac{1}{5}$ to $\frac{1}{4}$ three times a day in pill, or dissolved in hydrochloric ether. If it increases the symptoms at first, this is regarded as a good omen. If it be followed by gastrointestinal irritation or dryness of the throat, it must be stopped or the dose lessened.

Strychniæ Sulphas is recommended by TROUSSEAU. Its effects should be carefully watched, and its administration confined to obstinate chronic cases.

Valeriana is recommended by some writers.

Vinum. Dr. WILLIAM STRANGE (*Brit. Med. Jour.*, 1881) gives a purge to clear out the primæ viæ and then administers f. $\frac{3}{4}$. iij-vj of port wine daily. This simple treatment cures most cases.

Zinci Oxidum is generally considered inferior to the sulphate of zinc.

* *Zinci Sulphas* is very highly recommended by some physicians, (BARLOW, GOLDING BIRD and WEST,) but in the hands of others, (STONE, HILLIER, etc.,) it has not fulfilled expectations. Some authorities say that it only does good up to a given time, which is shown by the patient's complexion assuming a bright and clear color; also, that when this point is reached, the zinc not only becomes injurious, but if continued, marked anæmia shows itself. Should this state of things happen, iron combined with the zinc sometimes does good; as a rule, however, iron and quinine act much better.

According to BRISTOW, the rule to be followed in giving sulphate of zinc, is to begin with doses of a grain or two three times a day, which are slowly increased by successive increments until from 20 to 40 grains are given at a time. In the less acute type, valerianate of zinc will be found of especial use, particularly when the attack has with it some of the characteristics of hysteria. TROUSSEAU says: "Zinc does best with florid, and iron with pallid children."

EXTERNAL REMEDIES.

Baths. Simple cold baths and affusions are always of service. An excellent auxiliary is the salt-bath. In anæmia, sulphur-baths are beneficial.

Electricity is principally used in cases dependent upon deranged catamenial function. Dr. GOLDING BIRD recommends it in the form of sparks taken in the course of the spinal column, every alternate day, for about five minutes at each time. Dr. HAMMOND states he has used both the primary galvanic and induced currents in many cases, but he is of opinion that they are inefficacious, except in that form where there is distinct paralysis.

Ether-spray. The application of atomized ether along the spine has proved of service in relieving the spasms, especially in recent cases and those originating in fright. It is said also to be of benefit in chronic cases and in rheumatic chorea.

Galvanization of the Brain. Dr. CHARLES L. DANA contributes an article on this subject to the *Med. News*, 1883, in which he claims for anodal galvanization that it is a most valuable adjunct in the treatment of chorea; that given daily for a week or ten days with arsenic, and probably without, it will materially shorten the duration of the disease in most cases of chorea in children. The method of application is as follows: A large sponge-electrode of flexible brass, four by two inches, is thoroughly moistened with salt water. The hair of the patient is also thoroughly wetted, and the electrode applied over the side of the head, above the ear. In hemi-chorea it need only be applied over the side

opposite to the one affected. The other electrode is placed in hand of the affected side. The electrode upon the scalp is made positive, and a stable current of three to six Stöhrer's, four to eight Daniell's cells, is passed for from three to six minutes.

Hygienic Measures. Fresh air, change of scene, regulated movements, good food, and friction to the spine, are of the utmost importance in all cases. Gymnastics are especially recommended in the milder forms of chorea, and during convalescence in the severe cases. Drs. E. B. GRAY and H. M. TUCKWELL have published a large number of cases, (*Lancet*, November, 1876,) to show that the hygienic plan alone is sufficient to cure chorea, and quite as promptly as by any drug. They trust to judicious nursing and good diet, and merely give a placebo of sugared water. As early as 1871, they showed that the average duration of chorea treated on the expectant plan was, as far as their observations had then gone, from ten to eleven weeks; and that the average duration of chorea treated with arsenic in gradually increasing doses, according to BEGBIE's plan, was likewise from ten to eleven weeks; the difference, if any, being slightly in favor of the expectant plan. They also compared these results with the estimate of Dr. HILLIER, who, in his work on *Diseases of Children*, gives the mean duration of thirty cases treated with arsenic, etc., as "about ten weeks;" and with SÉE's estimate of "sixty-nine days," as the average duration of one hundred and seventeen cases treated with various medicines. An isolated ward; a good nurse; a large crib, well padded round, and walled in with pillows; plenty of nutritious food, without stimulants—on these they feel inclined to rely in the treatment of severe chorea, till we obtain more conclusive evidence than has yet been adduced that the disease can, in the slightest degree, be favorably influenced by any medicine as yet discovered.

Inhalations of chloroform arrest the convulsive movements so long as the patient is under their influence, but the progress of the disease is not checked by their administration, and they can only be regarded as palliatives in severe cases.

EPILEPSY.

Since our last edition not much of importance has been added to the treatment of epilepsy, the bromides still remaining on the whole the most effective remedies.

PROF. C. E. BROWN-SÉQUARD, M. D., PARIS.

The formula which this distinguished neurologist recommends in idiopathic epilepsy, is:

dose already given in the twenty-four hours is so large that any increase of it causes great sleepiness in the daytime, a decided lack of will and of mental activity, dullness of the senses, drooping of the head, considerable weakness of the body, and a somewhat tottering gait.

4. That it is never safe for a patient taking either of the bromides or both, and receiving benefit therefrom, to be even one day without his medicine, so long as he has not been at least fifteen or sixteen months quite free from attacks.

5. That the debilitating effect of the bromides in patients already weak, as are most epileptics, ought to be prevented or lessened by the use of strychnia, arsenic, the oxide of silver, ammonia, or cod-liver oil, cold douches or shower baths, and, of course, wine and a most nourishing diet. In making use of strychnia or arsenic, it must be kept in mind that not only the bad influences of the bromides, but also their favorable influence against epilepsy, can be diminished by these powerful agents (especially strychnia), and that it is therefore necessary, when these agents are used, to increase the dose of the bromides.

6. That *iron* and *quinine*—which are generally injurious to epileptics, except in cases in which the nervous affection is caused, or at least aggravated by chlorosis, anæmia, or malarial cachexia—are more particularly injurious in cases in which the bromides are taken.

7. That a gentle purge every five or six weeks usually gives a new impulse to the usefulness of the bromides against epilepsy.

DR. E. C. SEGUIN, OF NEW YORK.

Dr. SEGUIN's method of prescribing the bromides in the treatment of a case of "idiopathic" epilepsy is the following :

Two solutions are employed.

30. R.	Potassii bromidi,	$\frac{m}{3}$ j ss vij.	M.
	Ammonii bromidi,		
	Aquæ fontanæ,		
S.	To be given by the teaspoonful.		

And,

31. R.	Sodii bromidi,	$\frac{m}{3}$ j ss vij.	M.
	Ammonii bromidi,		
	Aquæ font.,		
S.	To be given by the teaspoonful.		

The quantity administered is, as a rule, so divided as to give by far

the largest dose in the evening. The bromide is cautiously increased, still keeping the nocturnal dose the largest, until slight bromism is produced. It is usually necessary to maintain slight bromism for months, but just as little is to be given as will prevent the attacks. The precise quantity required must be studied in each case. Children tolerate the bromides, as well as the iodides, in relatively large doses. It is regarded as important, thoroughly to dilute the bromides in order to facilitate their absorption—the dose to be taken in a wineglassful or half a tumblerful of water. Under no circumstances should the bromides be discontinued; they may be *diminished*, but not *stopped* until the word *cure* can be pronounced. They should be continued at least three years after the last attack.

The adjunct treatment consisted in the use of measures to prevent the acne to a certain extent, such as the occasional use of arsenic, sulphur-ointment, mercurial plaster, and alkaline lotions; to correct the general debility or slight paresis, by the use of strychnia, nux vomica, oxide of zinc and quinia; to relieve the dizziness by the inhalation of nitrite of amyl, by stimulants and by quinia; regulating the patient's diet and hygiene, and the use of cream, cod-liver oil, iron, quinine, phosphorus, strychnia, with nitro-muriatic acid, wine, beer, or whisky. In certain cases such medicines as act more directly upon the morbid state of the nervous centres are associated with the bromides, and the favorite among these is belladonna. In the treatment of cases of epilepsy in which a definite causative lesion can be made out, the bromides are used simply to combat the habit.

Regarding the method of using the bromine salts in the treatment of epilepsy and other neuroses, he makes the following remarks:

1. The prolonged use of bromides is contra-indicated by congenital feebleness.

2. The bromides are well borne by persons of fairly full habit and good nervous power.

3. The bromides are indicated in cases of abnormally great irritability of the nervous system in its motor (muscular and vaso-motor) and ideational tracts.

4. The contra-indications above named, are to be much less regarded in the management of that formidable neurosis, epilepsy.

5. Epilepsy is the only disease which justifies the deliberate production of a degree of bromism for its cure.

From extensive and very elaborately conducted experiments, to determine the precise effects of bromide of potassium in epilepsy and its proper dose, Dr. T. S. CLOUSTON, of Edinburgh, has found that the diminution of the fits, and all the other good effects of the medicine, reach their maximum in adults at thirty-grain doses thrice daily; while ill effects are manifested when thirty-five-grain doses thrice daily are reached.

In regard to the other bromides, Dr. ERLÉNMEYER (*Correspondenz-Blatt*, 1877,) has experimented with *bromide of lithium*, recommended by Dr. LEVI, of Paris, on account of having no action on the heart. From his own experiments, he is disposed to believe that it is inferior to bromide of potassium as a remedy against epilepsy. He has also tried *bromide of quinine*, and found it had a more hypnotic action than bromide of potassium; he finds it useful in treating periodic fits of insanity and hypochondria, but is not inclined, on the whole, to give it the preference to the more commonly used drug in epilepsy.

In combination with the bromides, *atropia* has been favorably mentioned by Dr. L. P. YANDELL, jr., of Louisville.

32. R. Potassii bromidi, Atropiæ sulphatis, Aquæ,	gr. xv gr. $\frac{1}{60}$ q. s.	M.
---	---------------------------------------	----

For one dose, three times a day.

It is said to act more promptly than the bromide alone, (*Medical and Surgical Reporter*, 1878.) *Atropia* has been used successfully alone by Dr. SVETLIN, of Vienna, and later by Dr. LEIDESDORF. The latter found it to act beneficially if given to the extent of $\frac{1}{60}$ grain of the sulphate daily. The trial was made in consequence of the accepted fact that small doses diminish the action of the reflex nerve-centres, while large doses produce an opposite effect. Cases of motor epilepsy of not long standing recovered rapidly, and some old cases, complicated with mental derangement, also got well while taking the drug. In other cases the attacks were rendered less frequent. The observation was confirmed by experiments on animals. (*Medical Examiner*, No. 34, 1877.)

BROWN-SÉQUARD recommends

33. R. Morphiæ sulphatis, Atropiæ sulphatis,	gr. $\frac{1}{4}$ gr. $\frac{1}{60}$.	M.
---	---	----

For one *hypodermic injection*, in a few minims of distilled water. He succeeded in a case of epilepsy by the use of this injection alone.

A combination of the bromides with *chloral hydrate* has been used with advantage by Dr. J. C. SHAW, of Brooklyn, and various New York physicians. The usual formula is:

34. R.	Potassii bromidi,	$\frac{3}{4}$ j	
	Chloral hydratis,	$\frac{3}{4}$ ss	
	Aquæ,	f. $\frac{3}{4}$ vij.	M.

Four to six teaspoonfuls a day.

This seems to bring about the good—*i. e.*, antispasmodic—action of the bromide without so much cutaneous eruption, general physical deterioration, and mental depression, as we too often see during the ordinary treatment of chronic cases of epilepsy.

With reference to these untoward constitutional effects,

PROF. WILLIAM A. HAMMOND, M. D., NEW YORK,

Enumerates them as follows, in the usual order of their occurrence: 1. Contraction of the pupils. 2. Drowsiness. 3. Weakness of the arms and legs. 4. Depression of mind. 5. Failure of memory. 6. Delusions. The first three of these are the usual accompaniments of a dose of the medicine capable of producing any influence over epilepsy. In adults they never follow less doses than ten grains. Doses of five grains produce no effect. The ordinary dose of either one of the bromides—of potassium, sodium or calcium—is gr. xv three times a day, in solution. After two months, this must be increased by one-half, and more if paroxysms have intervened.

Dr. H. has used the *bromide of zinc* with very beneficial results: His formula is:

35. R.	Zinci bromidi,	$\frac{3}{4}$ j	
	Syrupi simplicis.	f. $\frac{3}{4}$ j.	M.

Ten drops three times a day, gradually increased to thirty or more, largely diluted with water.

There is no danger of bromism in using this agent, and it exerts a positive curative influence.

In the nocturnal form of epilepsy, *strychnia* is sometimes remarkably efficacious. A good formula for its administration is:

36. R.	Strychniæ sulphatis,	gr. ij	
	Aquæ destillatæ,	f. $\frac{3}{4}$ j.	M.

Dose, eight drops three times a day for the first two weeks, then nine drops for the next two weeks, increasing thus a drop every two weeks for a year, and perhaps longer.

In cases produced by gastric derangement, this will also be found a useful treatment.

The *nitrite of amyl* acts well as inhibiting the paroxysms, and preventing individual convulsions, but as a curative agent, exerts no power.

Epileptiform seizures due to syphilitic infection, require the iodide of potassium.

Of surgical measures, the excision of any cicatrix which, by entangling a nerve, may be the source of reflex irritation, is occasionally a useful measure. In epilepsy following injuries of the skull, trephining often relieves. A seton, or the actual cautery to the nape of the neck, has at times proved decidedly beneficial.

The hygienic treatment is important. Moderate exercise in the open air, nutritive and easily-digested food, fresh air, etc., should be insisted upon. Causes of eccentric irritation, as intestinal worms, disordered menstruation, and local irritations, must be removed.

The *alkaloids of nux vomica* appear to have a well-merited reputation in this disease. Thus:

WALTER TYRRELL, M. D., LONDON,

States that he has watched the effects of *strychnia* upon various forms of epilepsy, and has no hesitation in affirming that in a large majority of cases, its effects are most beneficial. He found but three cases in which it produced no favorable result, and no cases in which it produced an unfavorable effect. He gives a medium quantity as a dose, for a lengthened period, rather than carrying the dose too high at first. The best results are obtained from gr. $\frac{1}{10}$ to gr. $\frac{1}{8}$, twice a day, in solution, the system appearing to regain its nervous strength under the continued use of the medicine.

The following was a favorite prescription with the late

MARSHALL HALL, M. D., LONDON:

37. R.	Strychniæ acetatis,	gr. j	
	Acidi acetici,	℥xx	
	Alcoholis,	f. 3 ij	
	Aquæ destillatæ,	f. 3 vj.	M.

Ten drops (gr. $\frac{1}{30}$) to be taken in water thrice daily.

Another alkaloid, *brucia*, has been tried by

J. SPENCE RAMSKILL, M. D., LONDON.

38. R.	Bruciae,	gr. iv	
	Alcoholis,	f. ʒ ij	
	Aquæ destillatæ,	f. ʒ vj.	M.

Ten minims to be taken, diluted with water, twice daily; every third day, an addition of five minims should be made to the dose, until from a third to a half grain is reached, in the treatment of stomachal epilepsy.

If any stiffness of the jaws or other toxic symptoms appear, the dose is to be diminished five minims, and continued until any new objectionable symptom is manifested; then it is again lessened. No benefit will be derived until a full dose is reached; often the reverse effect. (*London Lancet*, January 13th, 1869.) As a rule, patients will take twice as much brucia as strychnia, without any necessity for diminishing the dose. After the continuous administration of brucia for a month, it is well to suspend its use for some days, and then again resume it. Great satisfaction will be obtained by giving the bromide of potassium in large doses, at bedtime, and at the same time ordering brucia twice daily, thus insuring the sedative influence of the bromide and the tonic effect of the brucia on the whole nervous system.

DR. ALLEN M'LANE HAMILTON, NEW YORK.

In regard to the bromides, this writer prefers the *sodic bromide*, administered on an empty stomach, to the extent of ʒj in the twenty-four hours. He has observed that a heavy dose at night is apt to do more good than if the amount prescribed is equally divided up through the day. In a great many patients the attacks occur at the waking hour, probably owing to the sudden change in the cerebral circulation. A mild diffusive stimulant has overcome this, and in many cases warded off the attack. He directs his patients who have their convulsions at this time to keep a glass of sherry or a small quantity of spts. ammoniæ aromaticus near at hand, to be taken before rising. Cold douches to the head are valuable. If the attacks be irregular, it will be found necessary to divide up the dose.

Belladonna and its alkaloids are of great value when the seizures occur in the daytime, or are of the variety known as *petit mal*. He has injected the sulphate of atropia, in $\frac{1}{4}$ gr. doses, beneath the skin at the back of the neck, with good effect. It should be administered until dryness of the throat is obtained, and should be given a patient trial. A remedy of his own suggesting is *nitro-glycerine*. He says it is as

powerful a medicinal agent as it is an explosive, $\frac{1}{10}$ of a drop producing a rapid cerebral hyperæmia. It is safer than amyl nitrite, and produces a more lasting congestion, and acts better as an abortant of the attack.

39. R. Nitro-glycerinæ, gtt. v.
Alcoholis, f. $\frac{3}{4}$ j. M.

Ten drops of this contain about $\frac{1}{10}$ of a drop of nitro-glycerine, which amount is the proper dose. Mixed with alcohol it is non-explosive.

Dr. HAMILTON has found a combination of the various anti-epileptic remedies to act better than any one singly, and for many years has derived much satisfaction from the following formula:

40. R. Strychniæ sulphatis, gr. j
Fl. ext. ergotæ, $\frac{3}{4}$ iss
Sol. potass. arsenitis, $\frac{3}{4}$ ij
Sodii bromidi, $\frac{3}{4}$ iss
Tr. digitalis, $\frac{3}{4}$ ij
Aquæ menth. pip., ad $\frac{3}{4}$ iv. M.

S.—A teaspoonful before eating, in a half tumblerful of water.

If the attacks be of the form known as *petit mal*, he thinks either *ergot* or *belladonna* are our best agents. With either form of treatment it may be found often necessary to use auxiliary general treatment. The syrup of the combined phosphates, or the syrup of the lacto-phosphate of lime, is a good adjunct; and salt baths, cold head douches, regular food, early hours, and the breaking off of bad habits, will often cure the disease, even when it has lasted for many years.

As a last resort, should continued medication prove useless, the *actual cautery* or a deep *seton* at the back of the neck, will occasionally arrest these bad cases.

The *oxide of zinc* is an old and standard remedy. From recent trials, Dr. AUGUSTE VOISIN states that though it acts more slowly than the bromide of potassium, its effects are more certain; and, after having used both these two medicines for more than twenty years, he habitually gives the preference to the oxide of zinc, used after the method of HERPEN, of Geneva, that is, commencing with small doses, gr. ij–ijj, combined with valerian, thrice daily, and slowly increasing them to the maximum, gr. xv. thrice daily, and continuing for a long period, three to six months. The same high opinion of the zinc is expressed by

J. WARING CURRAN, M. D., DUBLIN.

41. R. Zinci oxidi, gr. ijss
 Extracti glycyrrhizæ, q. s. M.
 For one pill. One or two thrice daily.

This, together with the bromide of potassium in mixture, forms, he believes, a method of treatment not to be equaled in epilepsy, when assisted by the occasional application of *Chapman's spinal ice-bag*. Neither remedy succeeds so well alone; the one is essential to the other.

EDWARD C. SPITZKA, OF NEW YORK.

This neurologist has found *ergot* and *conium* valuable drugs in epilepsy, (*Physician and Pharmacist*, 1879.)

42. R. Extr. ergotæ fluidi, ʒ ss-j
 Extr. conii fluidi, gtt. v. M.
 This for a dose, three times daily.

Respecting ergot, his theory of its action is that in small and frequently repeated doses it reduces the overtone of the vaso-motor centre by repeated excitation, and thus strikes at the root of the disease. Conium, he has found, experimentally, to be one of the most reliable cerebral sedatives in the market, and whose protracted administration is followed by no bad results. Although, as a matter of routine, he has given both drugs together, he should formulate their indications differently. When acts of violence replace the convulsive attack, conium will produce the desired effect, while ergot, the bromides and all other drugs will fail. He has found that in convulsions due to cortical disease, as *remollissement rouge*, abscess and tumor, the same drug, in combination with bromides, exerts the best influence. On *petit mal*, vertigo and the epileptic attack as such, ergot alone exercises an immediate effect, and herein ranks far above the bromides. For some of the epileptic phenomena, as headache, the bromides are useful; but as to the influencing of epilepsy *per se*, the weight of authority seems to be favorable to the view that they do very little good, and much harm.

Ergot has this advantage, shared by conium, that its use may be commenced and abandoned with suddenness without precipitating an epileptic explosion, as the sudden interruption of the bromide administration is sure to do.

INHALATIONS.

GEORGE JOHNSON, M. D., *King's College Hospital, London*, speaks in high terms of *chloroform* in connection with bromide of potassium in this affection. He thinks that the action of chloroform inhalation in warding off a threatened fit, and in cutting short a violent and prolonged paroxysm, is as uniform and certain as the action of anæmia in exciting convulsions.

The same has been said of the use of inhalations of *amyl nitrite*. It is considered by Dr. CRICHTON BROWNE to be invaluable in many cases, in not only postponing, but in altogether preventing epileptic seizures. A vinaigrette, or small-stoppered bottle, containing a sponge soaked in nitrite of amyl, should be carried in the pocket, and the fumes inhaled on the first symptoms of the aura. Even in the advanced stage of the *status epilepticus*, Dr. B. believes its inhalation very beneficial in most cases. (*West Riding Lunatic Asylum Reports*, 1873.)

MECHANICAL AND PHYSICAL MEANS.

Dr. BROWN-SÉQUARD has found:

1. That it is not necessary to apply an irritation (by a ligature, pinching, etc.,) on the very limb from which an aura seems to start, as the same means applied elsewhere may succeed; but the chance of success is much greater by the former than by the latter way.

2. That a constant or a frequent irritation (by a blister, an issue, a seton, the actual cautery, etc.,) on the place from which an aura seems to start, may not only prevent fits, but, by some change of nutrition locally (if the aura is really of peripheric origin), and in the nervous centres, may reduce, or even destroy altogether, the tendency to fits, and lead to a complete cure.

3. That as a circular ligature may procure a temporary good effect, so a narrow *circular blister* applied all around a limb, a toe or a finger, or a circular cauterization with a white-hot iron, may cure epilepsy in cases with a distinct aura.

4. That even in cases in which there is no aura, felt or unfelt, ligatures, pinching, and other means of irritation, may prevent the occurring of expected fits.

When an attack of epilepsy is followed by a comatose state, or even a sleep with heavy breathing, it is of the greatest importance to place the head of the patient in such a position that the tongue, which is then paralyzed, will not fall on the larynx and cover its aperture.

PROF. J. M. DA COSTA, M. D., PHILADELPHIA.

43. R. Zinci valerianatis, gr. iij
 Extracti belladonnæ, gr. $\frac{1}{2}$
 Pulveris digitalis, gr. $\frac{1}{4}$. M.

For one pill. To be taken thrice daily in cases of epilepsy, associated with irregularity of the heart.

HOSPITAL OF DISEASES OF THE CHEST, LONDON.

44. R. Potassii bromidi, gr. x
 Tincturæ conii, ℥xxx
 Tincturæ valerianæ ammoniatæ, ℥x
 Aquæ camphoræ, f. ʒj. M.

For one dose, thrice daily.

HOSPITAL OF UNIVERSITY COLLEGE, LONDON.

45. R. Potassii bromidi, gr. x
 Spiritus chloroformi, ℥xviiij
 Infusi quassiae, f. ʒj. M.

For one dose, thrice daily.

NOTES ON REMEDIES.

Acidum Hydrobromicum. *Hydrobromic Acid* is preferred to the bromides by Dr. C. L. DANA (*N. Y. Med. Jour.*, 1883). He has seen the greatest benefit from it. He uses it in preference because it is agreeable to take and does not produce an eruption or bromism. It is argued by Dr. H. C. WOOD (*Med. News*, 1884) that the failure of this acid is owing to the small doses given. He has tried ʒij–iij. daily with decidedly beneficial results. Each dose was given after meals, with an equal quantity of ginger syrup and diluted with a full half pint of water. These doses were well borne and there was little bromism. Dr. W. C. WADE, (*Detroit Medical Journal*, December, 1877) gives the following prescription :

46. R. Stramonii extract, fluidi, gtt. 160
 Acidi hydrobromici diluti, āā f. ʒj. M.
 Syrupi,

One-half teaspoonful in water, the dose to be increased until the specific effects of the stramonium are marked, and there to be maintained.

He adds : "I will simply say of this combination, that its effects in this disease are remarkable, and I think I have reason to consider it superior to any other plan of medication."

Ammonii Bromidum is often prescribed in epilepsy alone, or in combination with bromide of potassium. (F. 30.)

Ammonii Carbonas is a valuable palliative, though possessed of no curative influence in this disease. Dr. ANSTIE and others assert that, if there be time to administer a dose previous to an impending attack, it will often

avert it. PEREIRA recommends it in large doses (gr. x-xx), in the hysterical form of the disease.

Ammonii Formias has been found of service in some forms of epilepsy, hurtful in others. Dr. RAMSKILL advises a further trial of this remedy.

Ammonia Aqua is recommended by Dr. HOPE and others, in epileptic congestion of the brain, arising from debility. The following formula is the most efficacious :

47. R. Aquæ ammoniæ, ℥ xij
 Aquæ menthæ viridis, f. ʒ iss. M.

For one dose. If taken at the first warning of an attack of this character, it seldom fails to arrest its supervention. The inhalation of ammoniacal vapor immediately after the first warning of an attack, is recommended by PEREIRA, PINEL and others, as often averting its occurrence.

Ammonii Valerianas, although very deliquescent, and therefore uncertain in its operation, is favorably reported upon by a number of writers as a remedy in epilepsy.

Amyl Nitrite. See Nitrite of Amyl.

Anilin is recommended by Drs. TURNBULL and ANSTIE. The latter finds that large doses aggravate the fits, but that one grain thrice daily, with an additional grain on the occurrence of any prodromata of a fit, delays or mitigates the paroxysm, or even averts it for a considerable time.

Argenti Nitras is no longer given in prolonged courses, at the risk of the patient's turning blue, a fate so frequently reserved for epileptics in former times. Other remedies of equal or greater efficacy, and less danger, have now, to a great extent, displaced it.

Arseniosum Acidum is employed in epilepsy, but it is much less effective in this disease than in chorea. Dr. RADCLIFFE employs it hypodermically and endermically, in the same manner as for chorea, (which see.)

Arsenici Bromidum. Dr. CLEMENS, of Frankfort-on-the-Main, has employed for a number of years a *liquor arsenici bromidi*, as follows :

48. R. Pulv. arsenici albi,
 Potassæ carbonatis, āā ʒj
 Aquæ destillatæ, Oss.

Boil and add,

Aquam, ad f. ʒ xij
 Brominii, f. ʒ ij.

After cooling and filtration, it should be a colorless liquid. Dose, one or two drops in a glass of water once or twice daily. This dose need not be increased. All his cases have been decidedly improved by this, but he claims only two complete cures in fifteen years.

Assafætida is recommended by Dr. POLLOCK, (*Lancet*, August 21st, 1869,) as capable of giving satisfactory results in the following combination :

49. R. Tincturæ assafætidæ, ℥ xxx
 Ammonia carbonatis, grs. iij. M.

For one dose, to be repeated thrice daily. In cases of epileptic seizures, due to the presence of worms in the alimentary canal, *assa-fœtida* is a useful agent.

Atropia. See above, (p. 51.)

Belladonna, though praised by TROUSSEAU, WILKS and HARLEY, (particularly in cases arising from emotional excitement, when it should be combined with zinc,) is superseded in practice by *atropia*.

Bismuthi Subnitras has been employed by Dr. COPLAND (the bowels being kept freely open) alone and in combination with tonics and antispasmodics, with good results.

Brucia. (See F. 38.)

Camphora seems to be useful in cases associated with hysteria or uterine disorders. It should be given in combination with tonics and antispasmodics. The *mono-bromide* is a desirable form.

Cannabis Indica has recently had testimony advanced in its favor by Dr. WHARTON SINKLER. (*Philadelphia Medical Times*, October, 1878.) FLINT doubts its value.

Castoreum is a very old remedy in this disease, having been recommended by CELSUS.

Centaurea Benedicta. Dr. J. G. WESTMORELAND (*Acology and Therapeutics*, p. 320,) observes that the "blessed thistle" has proved a useful remedy in epilepsy. A decoction of the leaves is used, \mathfrak{z} j to water Oj, the dose being \mathfrak{z} iij-iv, daily.

Cerii Oxalas is recommended by Prof. SIMPSON.

**Chloroformum*. Chloroform inhalations are recommended (p. 57,) both during the paroxysm and in the interval. Dr. BROWN-SÉQUARD considers this method of treatment particularly valuable in cases partaking of a hysterical character.

Cinchona is rarely successful in true epilepsy, but of great benefit in the epileptiform paroxysms of intermittent fever.

Conium has been found by Dr. HARLEY to cause rapid and decided improvement in epilepsy arising from sexual abuse, or from the irritation of dentition, but in that arising from peripheral disorder of sensation, from menstrual irregularity, or from emotion, it fails to exercise any beneficial influence.

Cupri Sulphas Ammoniatum produces positive benefit, according to AUGUSTE VOISIN.

Curare has been recommended by C. F. KUNZE, as follows :

50.	R.	Curaræ,	gr. vijss
		Aquæ destil.,	℥lxxv
		Acid. hydrochlor.,	℥vij.

For hypodermic injection, eight drops every five or six days.

Hydrargyri Iodidum Rubrum has been recommended by Dr. FULLER (*Medical*

Times, February 14th, 1857,) in cases where there is reason to suppose thickening of the dura mater.

Hyoscyamus in occasional full doses (f. ʒ iv-vij of the succus, or f. ʒ iv-vij of the tincture,) has been found by Dr. HARLEY very serviceable in epilepsy arising from emotional disturbances, but in other varieties of this disease it has proved useless in his hands.

Moschus is sometimes useful. Dr. A. T. THOMSON says that it diminishes the violence of the paroxysms of idiopathic epilepsy, and greatly lengthens the intervals, when administered as follows :

51. R. Moschi, ʒj.

In a bolus for one dose, every eight hours.

Nickel Bromide, has been found by Dr. J. M. DA COSTA (*Med. News*, 1884,) to be decidedly active, more so than other bromides. The dose is gr. v-x, repeated as necessary. It has a marked effect on epileptic seizures.

Nitrite of Amyl by inhalation, has been found eminently useful by Dr. S. WEIR MITCHELL and others.

Nux Vomica. In epilepsy following the disappearance of the menstrual discharge, COPLAND recommends *nux vomica* combined as follows :

52. R. Extracti nucis vomicæ, gr. x
Pilulæ aloes cum myrrha, ʒ ij. M.

For thirty-six pills. From one to two, night and morning. Its employment demands caution.

Opium is now rarely employed, since the discovery of the value of bromide of potassium, in this affection.

Oxygen Inhalations are recommended by Dr. RAMSKILL. (*Medical Times*, July 4th, 1863.) On the accession of giddiness, or other uncomfortable symptoms, they should be stopped.

Phosphorus is regarded by Dr. ANSTIE as well worthy of further trial in this disease, as it improves the patient's general health and relieves the sense of languor and depression.

****Potassii Bromidum** is the remedy *par excellence* in epilepsy. See opinions already given of Drs. BROWN-SÉQUARD, CLOUSTON, HAMMOND, etc. It cures in very many cases, and rarely fails to diminish the number of attacks.

Quinæ Sulphas is sometimes prescribed. Prof. LUDWIG MEYER, of the Berlin Charité Hospital, has long recommended a full dose of quinine, gr. xx-xxiv, at the first symptoms of the aura. In some instances, he has succeeded in obtaining a permanent cure by continuing this treatment. Dr. NOTHNAGEL has witnessed the same favorable result from a large dose of common salt, (*Allgemeine Medicinische Central Zeitung*, November 19th, 1876.)

Santonin is of service when the epilepsy is due to the presence of worms in the alimentary canal.

Sinapis is a useful emetic in epilepsy from over-distention of the stomach.

Sodii Chloridum. See under *Quiniæ sulphas*.

Stanni Chloridum has been employed in doses of gr. $\frac{1}{8}$ to $\frac{1}{4}$ thrice daily, in pill, or dissolved in hydrochloric ether.

Strychnia is strongly recommended by WALTER TYRRELL, and in the threatenings of epilepsy, by MARSHALL HALL.

Tabacum is supposed to have a curative influence in epilepsy produced by onanism, as the use of tobacco allays sexual lust.

Terebinthinæ Oleum, though little used in epilepsy, is strongly recommended by Sir THOMAS WATSON, who states that if he were called upon to name any single drug from which, in ordinary cases, he should hope for relief, he should mention turpentine; and by Dr. HEADLAND, who prescribes it as follows:

53. R. Olei terebinthinæ, f. ʒ ss.
Thrice daily in capsule or emulsion.
54. R. Olei terebinthinæ,
Olei ricini, āā f. ʒ iij. M.
For one dose at occasional intervals.

Valeriana is highly praised as a remedy for epilepsy, by BRISBANE, FOTHERGILL, HALLER, WILLIS, and others, but regarded as of little value by CULLEN, HEBERDEN, and HOME.

Veratrum Viride has been found by Dr. E. F. MORDOUGH, of New York, to diminish the severity and frequency of the convulsions. His prescription is:

55. R. Morph. sulph., gr. iss.
Tinct. verat. virid.,
Aquæ, āā f. ʒ ss. M.
Twenty minims hypodermically during or before a convulsion. (*New York Medical Record*, September, 1878.)

Zinci Lactas was first recommended in epilepsy by Dr. HERPIN, (*Bull. Gén. de Thérap.*, November, 1876,) who regards it as of equal medicinal value with the oxide, and superior to it in taste and tolerance. He gives it, at first, in 2-grain doses, in pill, thrice daily, and gradually increases the quantity until 10 grains are taken during the day.

* *Zinci Oxidum* is regarded by Dr. RUSSELL REYNOLDS as superior to the sulphate of zinc, in epilepsy. He gives it in doses of grs. iii-v, thrice daily. Dr. WARING CURRAN strongly urges it as an adjunct to bromide of potassium. (See further, p. 56.)

Zinci Sulphas is a long-used remedy in epilepsy, of occasional value.

Zinci Valerianas is considered by many as the best salt of zinc in epilepsy. It is specially valuable in cases of a hysterical character. The dose should be small at first, one grain, gradually increased as the stomach will bear it. Dr. DA COSTA sometimes prescribes it with belladonna and digitalis. (F. 43.)

Emetics, given when an attack is imminent, will often, according to Dr. MARSHALL HALL, ward it off.

Stimulants. A draught of wine or of some diffusible stimulant may put off an attack.

EXTERNAL REMEDIES.

Cold Shower Baths are useful in epilepsy when the patient is not too timid nor too much excited by them. Methodical hydropathic treatment often benefits, but, according to NOTHNAGEL, is never sufficient alone to effect a cure.

Turkish Baths are sometimes employed, but great caution is required in the use of so powerful an agent.

**Blisters*. A narrow circular blister applied above or at the starting point of the aura, will sometimes prevent or modify a paroxysm.

Dry Cupping, sometimes, according to Dr. GRAVES, averts a paroxysm of epilepsy, where previous headaches or other premonitory symptoms advertise its approach.

Electricity has given unsatisfactory results in epilepsy. Dr. COPLAND says that "the safest method of employing electricity is to place the patient on the insulating stool and subject him to the electric bath, and to draw sparks from different parts, when thus insulated and placed in connection with the prime conductor."

Forcible Extension of the muscles where the aura is seated, or the application of pressure between the starting point of the aura and the trunk, may avert an attack. BROWN-SÉQUARD has stated that forcible extension of the *great toe* will arrest an attack, as he has verified in many cases.

Hypodermic Injections. *Atropia*, subcutaneously, in this disease, was proposed by BROWN-SÉQUARD, combined with morphia. (F. 33.) Prof. BARTHOLOW has also employed this combination. **Morphia* alone, or in combination with atropia, has been found, when injected hypodermically, not only to relieve quickly the paroxysms in violent cases, but to afford permanent benefit by diminishing the number, frequency and severity of succeeding attacks, and, therefore, to dispute with the bromide of potassium for the foremost position among the remedies against epilepsy. As one may succeed when the other fails, Prof. BARTHOLOW points out that bromide of potassium is most effective in cases of *grand mal*, in which the paroxysms occur frequently, with great violence, and during the daytime, and less effective in those which occur chiefly at night; while the hypodermic injection of morphia is preferable in epilepsy, the paroxysms of which occur at night, in the *petit mal*, and in convulsive tic, but not proper, as a general rule, in cases of epileptoid character dependent upon cerebral lesion. Prof. B. advises, when the paroxysms succeed each other rapidly, and are violent, that the injection be made during an attack, and without loss of time, but states that ordinarily two or three times a week will suffice, (dose, gr. $\frac{1}{4}$,) the onset of an expected attack being anticipated whenever practicable.

The Actual Cautery. This is a most valuable resource in epilepsy. BROWN-SÉQUARD'S rules are as follows: 1. The instrument has to be heated to whiteness. 2. It must be shaped like an olive, (blunt end.) 3. It must not be allowed to burn the skin, but is passed rapidly along the space to be acted upon, so that after the operation is over nothing is seen but a *white line*, very much resembling a scratch with a finger nail. So applied, it is never painful. It can be repeated every other day, as there are no scars and no irritation left. The necessity for observing these rules is obvious, as the object of the operation is to act on the terminations of nerve-fibres in the skin, but not to destroy them; hence it is clear that the method of first applying ether-spray on the spot to be cauterized, so as to deaden the pain of the cautery, mars the good effects of the application of the instrument, because the ether-spray is itself *more painful* than the actual cautery, when the latter is used after the manner above stated; and, moreover, it anæsthetizes the nerve-ends upon which it is desired to act, so that the cauterization becomes useless.

The Seton. Dr. THOMAS J. GRIFFITHS has strongly urged the employment of the seton in epilepsy. (*Report of the Supervising Surgeon, U. S. Marine Hospital Service, 1876.*) He introduces a large seton, fifteen or twenty strands of silk or flax thread, and allows it to remain for months, if necessary, removing only when the patient seems to be permanently improved or has recovered, unless too great irritation be produced by its presence, when it is removed temporarily, to allow the irritation to subside. The place of election for its introduction is the back of the neck. Of five cases of epilepsy, four were cured or greatly benefited. No other treatment was used except tonics. The disturbance of the mental faculties, which is the common attendant in such cases, rapidly improves after the insertion of the seton.

Trephining. In an examination of 145 cases of epilepsy, due to injury or disease of the skull, Dr. M. G. ECHEVERRIA found 93 cured and 18 benefited after trephining. (*Arch. Gen. de Med.*, December, 1878.) Such results should restore this operation to the position of a proper one in these cases.

HEADACHES.

Headache as a symptom is found associated with many varying conditions, as periostitis of the cranium, disease of the teeth, ear and eye, astigmatism, cerebral affections, neuralgia and rheumatism of the scalp, quininism, uræmia, alcoholic poisoning, etc. Beyond these cases, where the treatment is strictly etiological, there is a variety of idiopathic forms which require special treatment.

BILIOUS OR DYSPEPTIC HEADACHE.

Dr. THOMAS HAWKES TANNER recommends in this form:

56. R. Quiniæ sulphatis, gr. xxiv
Pulveris rhei, gr. xxxvj
Glycerinæ, q. s. M.

Divide into twelve pills, and order one to be taken at night.

Often of service in curing *bilious headaches*; the patient should take daily exercise in the open air, and avoid too much sleep.

Dr. HENRY G. WRIGHT states that when in this variety, the pain comes on directly after a meal, and when it can be traced to indigestible articles of food, and the patient is tolerably strong, an emetic is useful.

57. R. Pulveris ipecacuanhæ, gr. xxv
Ammonii carbonatis, gr. v
Aquæ menthæ viridis, f. ʒ iss. M.

Take at one dose, and follow by some warm fluid.

Where the pain ensues some hours after taking food, a warm draught, with the following formula, is generally beneficial:

58. R. Pulveris rhei, ʒ ijss
Magnesii carbonatis, ʒ ij
Spiritus ammoniæ aromatici, f. ʒ ij
Syrupi zingiberis, f. ʒ ss
Aquam menthæ piperitæ, ad f. ʒ ij. M.

A tablespoonful in water.

Such a headache may often be warded off by the following:

59. R. Pulveris rhei, gr. xvij
Pulveris capsici, gr. v
Sodii carbonatis exsiccatae,
Pulveris aloes,
Saponis castilliensis, āā gr. xij. M.

Divide into twelve pills, one to be taken before the meal, as a dinner pill.

60. R. Pilulæ hydrargyri,
Pilulæ rhei compositæ, āā gr. iv
Extracti hyoseyami, gr. ij. M.

Divide into two pills. To be taken at night, in cases of headache depending upon *dyspepsia*.

Or:

61. R. Pulveris ipecacuanhæ, gr. j
Pilulæ colocynthidis compositæ, gr. vij
Extracti gentianæ, gr. ij
Olei carui, ℥ ½. M.

Divide into two pills. To be taken at bedtime.

When the system is debilitated, it is often more advisable to employ the following, in place of the night pills :

- | | | | | |
|--------|-------------------------------|----|---------|----|
| 62. R. | Infusi sennæ, | | | |
| | Infusi rhei, | āā | f. 3v | |
| | Tincturæ cardamomi compositæ, | | f. 3j | |
| | Syrupi, | | f. 3ss. | M. |
- To be given as a draught in the morning.

Great benefit is generally derived from the combination of a bitter with an alkali, as in :

- | | | | | |
|--------|------------------------------|----|----------|----|
| 63. R. | Sodii carbonatis, | | 3ij | |
| | Spiritûs ammoniæ aromatici, | | f. 3vj | |
| | Tincturæ aurantii, | | | |
| | Syrupi aurantii corticis, | āā | f. 3ijj | |
| | Tincturæ gentianæ compositæ, | | f. 3iss. | M. |
- A dessertspoonful twice a day.

If the stomach be very irritable, with excessive flatulence, the following mixture is more suitable :

- | | | | | |
|--------|-------------------------------|----|------------|----|
| 64. R. | Bismuthi subnitratis, | | | |
| | Sacchari, | | | |
| | Pulveris acaciæ, | āā | 3j | |
| | Tincturæ cardamomi compositæ, | | | |
| | Tincturæ zingiberis, | āā | f. 3ss | |
| | Aquæ anethi, | | f. 3ijjss. | M. |
- A tablespoonful taken twice a day.

The same medicine may be combined with soda and capsicum, thus :

- | | | | | |
|--------|------------------------------|----|-----------|----|
| 65. R. | Bismuthi subnitratis, | | | |
| | Sodii carbonatis exsiccatae, | āā | 3ijss | |
| | Pulveris capsici, | | gr. viij. | M. |

For eight powders. One to be taken twice a day, in dyspepsia with much acidity, with loss of appetite and general want of tone.

- | | | | |
|--------|--------------------|-----------|----|
| 66. R. | Argenti oxidi, | gr. xij | |
| | Pulveris capsici, | gr. iv | |
| | Extracti gentianæ, | gr. xxiv. | M. |

Divide into eight pills. One to be taken twice a day.

In atonic dyspepsia, when the tongue is pale at the tips and edges, and the system weakly, few recipes prove so invariably successful as the above.

CONGESTIVE OR PLETHORIC HEADACHE.

In this form, Dr. WRIGHT states that the employment of medicines should, as far as possible, be dispensed with. They should only be re-

sorted to when the necessities of business prevent, or the solicitations of indolence interfere with, a strict control over the diet and regimen. In these cases, a saline diuretic should be ordered, such as :

- | | | | |
|--------|------------------------------|------------|----|
| 67. R. | Potassii acetatis, | ℥ij | |
| | Potassii nitratis, | ℥j | |
| | Spiritus juniperi compositi, | f. ℥xj | |
| | Aquam menthæ piperitæ, | ad f. ℥iv. | M. |

A teaspoonful twice a day, together with an occasional aperient at night, viz. :

- | | | | |
|--------|---------------------------------|--------|----|
| 68. R. | Pilulæ colocynthidis compositæ. | gr. l | |
| | Saponis castilliensis, | gr. ix | |
| | Olei anethi, | ℥ij. | M. |

Divide into twelve pills. Two to be taken at bedtime, and followed by a Seidlitz powder in the morning.

Persons subject to plethoric headaches should not partake of animal food more than once a day; should never indulge the appetite to satiety; should avoid beer, spirits, coffee, and all stimulating beverages; should bathe the head freely at night, and lie with it elevated on a hard pillow during sleep; should have an airy bed-room, and rise so soon as fairly awake, for otherwise, activity of the thoughts in a recumbent position will congest the head and cause it to ache.

When the patient has been exposed to cold, and the headache comes on at night, with the head hot and the skin harsh and dry, the following sudorific, taken at bed-time, is often of great service, the body being kept warm during its action :

- | | | | |
|--------|----------------------------------|-------------------|----|
| 69. R. | Antimonii et potassii tartratis, | gr. $\frac{1}{6}$ | |
| | Pulveris Jacobi veri, | gr. v | |
| | Pulveris potassii nitratis, | gr. v. | M. |

To be taken at night.

In the plethoric headache of pregnancy, relief is afforded by the use of saline medicines, as

- | | | | |
|--------|-------------------------------|----|---------|
| 70. R. | Magnesii sulphatis, | | |
| | Sodii sulphatis, | āā | ℥j |
| | Acidi sulphurici diluti, | | f. ℥ij |
| | Tincturæ cardamoni compositæ, | | f. ℥iss |
| | Syrupi aurantii corticis, | | f. ℥ss |
| | Aquæ cinnamomi, | | f. ℥j. |
| | | | M. |

A dessertspoonful twice a day.

Fluids should be avoided as far as possible. Sea air and sponging the body with tepid salt water generally prove beneficial.

In the treatment of *congestive headaches*, the aperients so frequently required should be cordial and saline, such as

71. R. Extracti sennæ fludi, f. ʒ ijss
 Magnesii sulphatis, ʒ ij
 Acidi sulphurici aromatici, f. ʒ ij
 Syrupi aurantii corticis, f. ʒ j
 Infusi rhei, f. ʒ ijss. M.

Dose, a tablespoonful in the morning.

There are no medicines so invariably useful in cases of congestive headache, attended with debility, as the preparations of *iron*. If the patient be of stout phlegmatic habit, the tonic may be combined with a cordial and saline, according to the following formula:

72. R. Ferri sulphatis, gr. xxxij
 Magnesii sulphatis, ʒ x
 Acidi sulphurici diluti, f. ʒ ij
 Tincturæ cardamomi compositæ, f. ʒ ij
 Syrupi, āā
 Aquæ pimentæ, āā f. ʒ j. M.

A dessertspoonful in water twice a day.

73. R. Tincturæ ferri chloridi, f. ʒ ij
 Acidi muriatici diluti, f. ʒ ss
 Tincturæ cinnamomi, f. ʒ iss
 Syrupi, āā
 Aquæ cinnamomi, āā f. ʒ vj. M.

A dessertspoonful in water, twice a day, about half an hour after food.

In the congestive headache of females past the middle period of life, especially when these headaches accompany alterations of the whole system at the great climacteric period, Prof. J. M. DA COSTA finds the use of saline cathartics is of service. Also, a mustard foot bath every night. The application of a *hot salt bag* to the back of the neck often affords relief. This form of headache is frequently associated with cardiac enlargement.

GOUTY HEADACHE.

In *gouty* headaches, according to Dr. WRIGHT, colchicum may be employed with greater freedom than in ordinary gout, care being taken that the bowels are freely open during its administration. An actual attack is best relieved by a brisk aperient followed by an effervescing mixture, containing an excess of potash, viz.:

74. R. Potassii carbonatis, ʒ iv
 Ammonii carbonatis, ʒ ij
 Tincturæ serpentariæ, f. ʒ ss
 Aquæ camphoræ, f. ʒ ijss. M.

Two tablespoonfuls to be added to a tablespoonful each of water and lemon juice, and to be taken effervescing twice or three times a day.

HYSTERICAL HEADACHE.

A pill of the following character is recommended by Dr. THOMAS H. TANNER :

- | | | | |
|--------|----------------------|--------------|----|
| 75. R. | Zinci valeriantis, | gr. xij-xxiv | |
| | Extracti belladonnæ, | gr. iij-vj | |
| | Extracti gentianæ, | gr. xxiv. | M. |
- Divide into twelve pills. One to be taken three times a day.

Useful in *hysterical headache*, especially when there is habitual constipation.

- | | | | |
|--------|---------------------------|--------------|----|
| 76. R. | Zinci phosphatis, | ʒj-ij | |
| | Acidi phosphorici diluti, | f. ʒ iss | |
| | Tincturæ cinchonæ, | f. ʒ vj | |
| | Aquam menthæ piperitæ, | ad f. ʒ iij. | M. |

Tablespoonful in half a wine glass of water three times a day, in *hysterical headache associated with general debility*.

In *clavus hystericus* and hysterical forms of migraine, Dr. ANSTIE found *muriate of ammonia*, grs. xx-xxx, repeated several times, of much benefit.

MIGRAINE, HEMICRANIA, PERIODICAL OR SICK HEADACHE.

This frequent and distressing form can best be met by *cannabis indica*, as recommended by Dr. RICHARD GREENE, (*Practitioner*, p. 267, 1872,) and since by Dr. E. C. SEGUIN, of New York. (*Opera Minora*, 1884.) The latter gives the details of his plans as follows :

First.—The *treatment of the patient* consists in removing all relievable exciting causes, and more especially in correcting acidity. For this purpose he gives nitro-muriatic acid and alkalies, greatly reducing the saccharine and amylaceous foods of the patient. In cases attended by debility, anæmia, and imperfect nutrition, it may be necessary to resort to tonics, including cod-liver oil.

Second.—*Treatment of the attack*. The first thing to be done is to place the patient under circumstances which secure quiet and semi-darkness. The attempt to “fight out” a sick-headache is nearly always vain, and may be injurious. It is better not to allow the patient any food, not even liquids, until toward the close of the attack, or even not till next day ; by this, nothing is lost, and much wretchedness is avoided. Ice, or ice washed in brandy, is grateful.

If the patient have a warning (aura of migraine) before nausea or pain, much can be done to cut short the attack or diminish its severity

by the use of guarana, caffeine or croton chloral hydrate, (grs. fifteen or twenty every hour.) In his hands, guarana or the powder of the seeds of *paullinia sorbilis*, has proved very efficacious.

Third.—*Treatment of the disease.* The principle of the treatment is to keep the nervous system steadily under a slight influence of cannabis for a long period of time. He gives to adult females $\frac{1}{2}$ of a grain of the alcoholic extract of *cannabis indica* before each meal, increasing the dose after a few weeks to $\frac{1}{2}$ grain. Males can generally begin with $\frac{1}{2}$ grain, and it is well to give them $\frac{3}{4}$ grain in two or three weeks. These doses must be taken with the greatest regularity, just as faithfully and regularly as bromides in epilepsy. Indeed, when beginning such treatment, he usually obtains a promise from the patient that he will regularly take the pills for a period of three months.

As a rule, no appreciable immediate effect is produced by the above doses, though lightness of the head and slight confusion of mind may result from an initial dose of $\frac{1}{2}$ grain three times a day.

Under this apparently and essentially simple plan of treatment, results are obtained; that is, the majority are greatly relieved, some entirely cured.

PROF. HARVEY L. BYRD, M. D., BALTIMORE.

In the treatment of *sick headache*, Prof. HARVEY L. BYRD, M. D., of Baltimore, recommends the use of an emetic of ipecacuanha, and after its action, the use of the following:

77. R. Chloralis,	ʒi.	
Aquæ,	f. ʒij.	M.

A tablespoonful every hour until sleep is induced.

DR. A. SEURE, OF PARIS,

strongly recommends (*Bull. de Ther.*, 1879,) the administration of *chloral* in an *enema*, on the occurrence of the paroxysm of migraine, having found it almost infallible, and exempt from the disadvantages attendant on other modes of giving this substance, and far preferable to the employment of morphia, quinine, etc., for the same purpose. He gives, according to the sex or robustness of the patient, from one and one-half to three grammes, in a glass of tepid water, the effect of the chloral being rendered still more prompt by the addition of a tablespoonful of brandy. If a sense of burning in the rectum is excited, this may be prevented by adding the yolk of an egg or substituting tepid milk

for the water. When, as in some persons, there is difficulty in retaining the enema, the quantity of the liquid may be diminished, and a drop or two of laudanum added.

PROF. AUSTIN FLINT, NEW YORK.

Periodical headaches, as regards successful treatment, belong among the opprobria of medical art. If patients be not unpleasantly affected by opiates, an attack may sometimes be warded off, or its severity much lessened, by a full dose of opium or one of its alkaloids. The carbonate of ammonium and a saline purgative are sometimes effective at the commencement of an attack. Various palliative measures may be resorted to, such as *inhalation of chloroform*, *evaporating lotions to the head*, (alcohol, spirits, vinegar, ether,) etc. In some cases a towel or napkin wrung out in water as hot as can be borne, and wound around the head, is more efficient than cold applicaitons. *Warm stimulating pediluvia*, strong *coffee* or *tea*, and the application of the *galvanic* or the *electro-galvanic current*, are useful in some cases. During the intervals the remedies which are sometimes of service by way of prophylaxis are *nux vomica* or *strychnia*, in small doses, *arsenic*, small doses of *quinia*, *belladonna*, and the preparations of *zinc*, more especially the valerianate. They may be tried in succession. Hygienic measures are important, and the avoidance of everything which experience shows, in individual cases, to act as exciting causes.

GEORGE KENYON, M. D., HARROWGATE, ENGLAND.

78. R. Carbonis bisulphidi.

f. $\bar{3}$ ij.

As a local application in periodical and hysterical headache, and even in many cases of dyspeptic cephalalgia.

About two drachms of the bisulphide of carbon is poured upon cotton wool, with which a small glass-stoppered bottle is half filled. The mouth of the bottle is applied *closely* to the temple or behind the ear, or as near as possible to the seat of the pain, and so held from three to five or six minutes. In a minute or two, a sensation as of several leeches biting the part is felt, and in three or four minutes more, the smarting and pain become rather severe, but subside almost immediately after the removal of the bottle. (*Medical Times and Gazette*, July, 1868.) It is very seldom any redness of the skin is produced. The effect of the application is generally immediate; it may be re-

peated, if necessary, three or four times a day. The sedative vapor of the bisulphide is probably absorbed through the skin, and acts upon the superficial nerves of the part to which it is applied.

DR. J. E. LOCKRIDGE, OF INDIANAPOLIS.

This writer (*American Practitioner*, March, 1877,) gives a formula which he has found of great service in every variety of headache, almost. It is:

79. R.	Potassii bromidi,	℥ij	
	Tinct. radice aconit.,	f. 3j	
	Aquæ destillatæ,		
	Syrupi simplicis,	āā	f. 3ij. M.

Take a dessertspoonful in some water every hour until relieved.

This is a large dose of the principal ingredients, but it rarely has to be repeated, and he believes a less dose is not to be depended upon.

Dr. WM. H. DAY states that in dyspeptic, bilious, or sick headache, one drop of tincture of nux vomica or of tincture of aconite, given in a teaspoonful of water, will often, after a very few doses, give entire relief, especially where the extremities are cold and the pulse small.

NERVOUS HEADACHE.

PROF. R. J. GRAVES, M. D., DUBLIN.

80. R.	Olei terebinthinæ,	f. 3j-ij.
--------	--------------------	-----------

For one dose, to be given in cold water.

Our author places much reliance on this medicine in the hysterical and nervous headaches of young women. Some will bear and derive benefit from two or three of these doses in the day, experiencing from its use a diminution of headache, the removal of flatulence, together with a moderate action on the bowels and kidneys. The turpentine may also be given in the form of an enema. Occasionally it causes dysuria, and cannot be persevered in.

Dry cupping at the nape of the neck, between the shoulders, and below the clavicles, is often of great service in hysterical headaches. Six cups should be applied and allowed to remain on for ten or fifteen minutes.

PROF. WILLIAM A. HAMMOND, M. D., NEW YORK.

81. R.	Zinci oxidi,	℥ij-v	•
	Confectionis rosæ,	q. s.	M.

Divide into twenty pills. One to be taken three times a day, after meals.

This formula is of great value. The minimum dose (gr. ij) should be commenced with, gradually increasing to the maximum (gr. v), if necessary.

82. R.	Extracti nucis vomicæ,	gr. v	
	Ferri redacti,	℥j	
	Quinise sulphatis,	gr. x	
	Syrupi,	q. s.	M.

Divide into twenty pills. One to be taken three times a day, after meals.

Nux vomica is preferable to strychnia.

83. R.	Bismuthi subcarbonatis,	℥ij	
	Confectionis rosæ,	q. s.	M.

Divide into twenty pills. One after each meal.

These pills will often take the place of those of oxide of zinc. They are particularly useful when there is gastric disturbance.

Bromide of potassium is serviceable when the nervous system has been irritated; when exhausted, it does harm.

Bromide of ammonium is similar to the bromide of potassium in its action, but the dose need not be so large. Our author often uses both combined.

Opium and its preparations are rarely of value in this disorder. If used, the hypodermic method is the best.

Narcein has, Dr. HAMMOND still thinks, a decided hypnotic effect when given in large doses.

Nitrite of amyl has been used by Dr. R. A. DOUGLAS LITHGOW, with much success. He places two drops on the palm of the patient's hand, and quickly diffusing these with his finger, over the palmar surface, tells her to cover her mouth and nose with her hand, and to inspire deeply and quietly. No time should be lost after the nitrite is dropped on the hand, as it evaporates rapidly. The patient should be seated while inhaling, as the peculiar effects of the nitrite are produced almost instantaneously, and may occasionally alarm a very nervous or hysterical female. Fortunately, these symptoms last a very short time—generally less than two or three minutes—and with their cessation, the pain almost invariably ceases. Two drops may be given as a draught in water, instead of by inhalation, but the latter mode is much more satisfactory.

Phosphorus is beneficial in all the forms of nervous headache. It is, however, difficult of administration, and leaves an unpleasant odor about

the person. The best results are obtained from the following method of administration:

84. R. Acidi phosphorici diluti, f. ʒvj
Syrupi phosphati compositi, f. ʒij.

A dessertspoonful, in water, three times a day.

Arsenic, as a nerve tonic, stands next in value to zinc. Granules of arsenious acid (gr. $\frac{1}{60}$) are preferable to FOWLER'S solution.

Galvanism is highly praised by some, and severely condemned by others, in this affection. The brain cannot be acted upon to any considerable extent by the induced current or by the reflex action. Our author advises, always, the *constant current*; being careful to avoid too great intensity, lest amaurosis be produced. (*Half-yearly Compendium of Medical Science*, July, 1868.)

THOMAS H. TANNER, M. D.

85. R. Acidi nitro-muriatici diluti, f. ʒij
Strychniæ, gr. $\frac{1}{2}$
Spiritus chloroformi, f. ʒvj
Tincturæ zingiberis, f. ʒij
Aqua ad f. ʒij. M.

A teaspoonful in water, three times a day, *in nervous headache*.

Holding the arms high above the head produces a marked effect upon the cerebral circulation, and will frequently relieve the severity of that peculiar morning headache with which some persons constantly awake.

Compression of the temporal arteries with a couple of pads and a bandage, may sometimes be of service.

Cold lotions, eau de cologne, etc., to the head, dry cupping or blisters, or setons to the nape of the neck; the removal of decayed teeth or stumps from the mouth, and change of air, are occasionally indicated.

Dr. WRIGHT states that in nervous headache, the combinations of hyoscyamus with camphor, with chloroform, or with a diffusible stimulant, generally afford great relief. The following formulæ may be used:

86. R. Extracti hyoscyami, āā ʒijss M.
Pulveris camphoræ,

Divide into twenty pills. Two to be taken when the pain is severe.

87. R. Chloroformi, f. ʒiss
Tincturæ hyoscyami,
Tincturæ cardamomi compositæ, āā f. ʒss
Olei limonum, ℥xvj
Sacchari,
Pulveris acaciæ, āā ʒss
Aquæ camphoræ, f. ʒij.

Dose, a tablespoonful.

88. R. Tincturæ hyoscyami,
Spiritus ammoniac aromatici, āā f. ℥ss
Syrupi aurantii corticis, f. ℥j
Aquæ menthæ piperitæ, f. ℥ij. M.

Dose, a tablespoonful.

- | | | | | | |
|-----|----|---------------------|----|---------|----|
| 89. | R. | Pulveris camphoræ, | | | |
| | | Quiniæ sulphatis, | āā | gr. x | |
| | | Extracti aloes, | | gr. xij | |
| | | Extracti hyoscyami, | | 3 ss | |
| | | Mucilaginis acaciæ, | | q. s. | M. |

Divide into eighteen pills. Two to be taken twice a day, in case of nervous headache when there is great debility and sluggishness of the system.

NEURALGIC HEADACHE.

True neuralgic headache is recognized by its complete intermittence, by its plainly superficial character, and by its following the course of recognizable nerve branches, which often present one or more painful spots in their course. The *stimulant narcotics* are the most efficient drugs to combat it, especially the preparations of opium. Dr. WRIGHT gives :

- | | | | | |
|-----|----|---------------------|----------|----|
| 90. | R. | Morphiæ muriatis, | gr. j | |
| | | Pulveris camphoræ, | gr. xxiv | |
| | | Mucilaginis acaciæ, | q. s. | M. |

Divide into six pills. One pill to be taken when the headache is very intense. Five hours are to be allowed to elapse before repeating the dose.

WILLIAM HENRY DAY, M. D., LONDON.

This author, in his work on *Headaches, their Nature, Causes and Origin*, states that the subjects of neuralgic headache are usually anæmic and require nutritious diet and rest. Frequently taints of gout, syphilis or mercury are present in very obstinate cases, and must receive proper treatment before a cure can be effected. Locally, we may use a chloroform and aconite liniment, or ointment of belladonna or veratria. The following is useful :

91. R. Sodii hypophosphitis, 3 ss
 Infusum columbæ, ad f. 3vj. M.
 Two tablespoonfuls thrice daily.

Arsenic is a remedy of the greatest value ; it must not be abandoned as soon as constitutional symptoms appear, but only suspended for a day or two and recommenced.

92. R. Tinct. cinchonæ, ℥^{xx}
 Liq. potass. arsenitis, ℥^v
 Mist. camphoræ, f. ʒj. M.
 This amount thrice daily.

In this form of headache as well as in some others,

DR. JOHN KENT SPENDER

speaks well of the local use of *chloroform* or of *bisulphide of carbon*, applied as follows: A small quantity of either is poured upon cotton wool, with which a small, wide-mouthed, glass-stoppered bottle is filled. The mouth of the bottle is then applied as near as possible to the seat of pain, and held there for five minutes. It produces a smarting and burning sensation on the skin, which is promptly followed by relief of the pain.

Another neat and efficient plan to relieve this and similar forms of headache is to steep a piece of blotting paper in *liquor atropiæ*, (gr. j to f. ʒj,) and apply it to the brow or painful part.

Heat and *cold* are valuable aids in the relief of the pain. A cloth wrung out in water as hot as the hands can bear it and laid on the brow, or a bladder of ice—whichever the patient seems to prefer—will often effect a speedy dispersion of the pain for the time being.

RHEUMATIC HEADACHE.

DR. HENRY G. WRIGHT, LONDON.

93. R. Linimenti chloroformi, āā f. ʒ^{iss}
 Linimenti belladonnæ, f. ʒj. M.
 Tincturæ opii,

For external application, *in rheumatic headaches*.

Mustard plasters applied to the neck are also exceedingly useful as a means of counter-irritation. In such cases (rheumatic headache) the following aperient is of advantage, viz.:

94. R. Pilulæ colocynthidis compositæ, gr. xv
 Extracti colchici acetici, gr. iij
 Olei carui, ℥j. M.

Divide into four pills. Two to be taken at bedtime, and one on consecutive nights. These pills should be followed, in persons of a costive habit, by a morning purgative, as follows:

95. R. Magnesiæ, ℥iv
 Liquoris potassæ, ℥xlv
 Extracti sennæ fluidi, f. ʒij
 Syrupi zingiberis, āā f. ʒss.
 Tincturæ aurantii, f. ʒj. M.
 Aquæ,
 Dose, a tablespoonful.

The administration of an alkaline medicine containing potash, if continued with regularity, will generally be followed by rapid amelioration of the pain and tenderness. The following may be used:

96. R. Potassii bicarbonatis, ℥iv
 Potassii chloratis, ʒiss
 Tincturæ cinnamomi, āā f. ʒvj
 Tincturæ aurantii, f. ʒiss. M.
 Syrupi aurantii corticis,

A dessertspoonful to be taken twice or three times a day.

If imprudent exposure to cold has produced an aggravation of the headache, and particularly if the patient be subject to catarrh, it is advisable to administer a sudorific at bedtime, such as

97. R. Pulveris ipecacuanhæ compositi, gr.xij
 Pulveris camphoræ, āā gr.iv. M.
 Pulveris gualiaci,

For one powder. To be taken about bedtime.

NOTES ON REMEDIES.

BILIOUS AND DYSPEPTIC HEADACHES.

Ammonii Acetatis Liquor is recommended by Prof. STILLÉ, who considers few remedies so successful in sick headaches as a teaspoonful or two of this solution.

Ammonii Carbonas will sometimes offer almost instantaneous relief.

**Ammonii Chloridum* is useful in bilious, as well as hysterical headaches. According to Dr. ANSTIE, gr. x-xx, if given early enough, seldom fails to cut short and greatly mitigate the attack. It is given with great benefit when the pain is most intense, in any form of headache.

Argenti Nitras is recommended by Dr. J. JOHNSON (in combination with minute doses of compound colocynth pills), in the habitual stomach headache to which delicate and literary men are so subject.

Bismuthi Subnitras is frequently advised, variously combined, in dyspeptic headache, by Dr. WRIGHT. (F. 64, 65.)

Carbonis Bisulphidum is recommended by Dr. KENNION, in sick headache, applied locally. (P. 55, 78.)

Hydrargyri Chloridum Mite, or other mercurial, is of service in headaches arising

ing from biliary derangement, or a torpid state of the bowels. It should be followed, at a short interval, by a saline or other purgative.

Magnesia, in cases accompanied with constipation, affords great relief.

Magnesii Citratis Liquor. A dose of the effervescing citrate is often very effectual in dyspeptic headache.

Podophyllum is prescribed with benefit in many forms of bilious headache.

Potassii Bromidum has been recommended for the prompt relief of the paroxysm.

CONGESTIVE OR PLETHORIC HEADACHE.

Asarum Europæum, gr. ij-iv of the powdered root, snuffed into the nostrils, has proved useful in cases of severe chronic headache originating in inflammation of the frontal sinuses.

Aloetic purgatives are beneficial in congestive headaches.

**Leeches* may be applied with advantage to the inner nostril, temples, to the crown of the head, or to the verge of the anus.

Saline aperients are recommended by Drs. DA COSTA and WRIGHT.

Sinapisms, to which have been added powdered capsicum or oil of turpentine, are often applied with benefit to the soles of the feet and the inner parts of the calves and thighs.

**Water*, locally applied, *hot*, is often more effectual than the cold lotions usually employed. Diligently steeping the temples, forehead, occiput, and nape of the neck with water, as hot as can be borne, in many cases will be found a useful remedy. The hot foot-bath is often of signal service.

GOUTY HEADACHE.

Colchicum may be employed with greater freedom in gouty headache than in ordinary gout, according to Dr. WRIGHT.

HYSTERICAL AND NERVOUS HEADACHES.

Ammonii Chloridum is advised by Dr. ANSTIE, to be administered in the same manner as for bilious headache (which see above.)

Ammonii Bromidum is recommended by Prof. HAMMOND, alone or combined with the bromide of potassium.

Argenti Nitras is highly prized by Dr. GRAVES, administered in considerable doses, *i. e.*, gr. $\frac{1}{2}$ five or even six times daily, for five or six days at a time, when the paroxysm has abated. If there be constipation, the nitrate should be combined with minute doses of compound colocynth pill.

Bismuthi Subcarbonas is often prescribed by Dr. HAMMOND, particularly when there is gastric disturbance.

Cajuputi Oleum is used both externally and internally, in nervous, rheumatic, and neuralgic headaches.

Camphora is useful, combined with hyoscyamus.

**Chloroformi Spiritus*, in the dose $\mathfrak{m}\text{xv}$ -xxx, often affords prompt relief in nervous and hysterical headaches.

Cimicifuga is useful in the headaches attendant on uterine derangement.

Guarana is particularly efficacious in the relief of sick headache, in doses of half a drachm to a drachm.

Hyoscyamus is strongly recommended by Dr. WRIGHT, in nervous headache. It may be combined with camphor, with chloroform, or with a diffusible stimulant.

Lavandula. The compound tincture, administered according to the following formula, makes an agreeable stimulant :

98. R. Tincturæ lavandulæ compositæ,
 Spiritûs chloroformi,
 Spiritûs ætheris compositi, āā f. ʒj
 Aquæ camphoræ, f. ʒ iij. M.

Dose, two tablespoonfuls in water. Opium may be added if indicated.
 This is a very effectual recipe for the headache of hysteria.

Narcein is recommended as a hypnotic by Prof. HAMMOND. (P. 62.)

Nitro-muriaticum Acidum is recommended by Dr. TANNER. (F. 85.)

Phosphorus is recommended by Prof. HAMMOND in all forms of nervous headache.

Quiniæ Sulphas is beneficial in hysterical headaches associated with anæmia.

Rosmarinus. The infusion is a mild stimulant, sometimes of service.

Stramonium has been prescribed with benefit by Dr. COPLAND.

Strychnia. Prof. HAMMOND considers nux vomica as preferable to strychnia in nervous headaches.

Terebinthinæ Oleum is recommended by Dr. GRAVES in hysterical headache.

* *Valeriana* is of especial benefit in hysterical headache. Dr. ASHWELL recommends the following formula :

99. R. Tincturæ valerianæ,
 Spiritûs ætheris compositi,
 Spiritûs lavandulæ compositi, āā ℥xxx
 Tincturæ hyoscyami, ℥xx
 Aquæ camphoræ, f. ʒ x. M.

For one dose.

Dr. CONNOLLY recommends the follownig:

100. R. Tincturæ valerianæ ammoniatæ,
 Spiritûs ætheris compositi, āā ℥xxx
 Aquæ camphoræ, f. ʒ iss. M.

Or,

101. R. Tincturæ valerianæ ammoniatæ, f. ʒj
 Acidi sulphurici diluti, ℥xv
 Aquæ camphoræ, f. ʒ iss.

Make a draught to be given twice or thrice a day.

Zinci Oxidum is regarded as of great value in nervous headache by Prof. HAMMOND.

Zinci Phosphas is recommended by Dr. TANNER, in hysterical headache, associated with debility.

Zinci Valerianas is also recommended by Dr. TANNER, combined with belladonna.

EXTERNAL REMEDIES.

Æther to the forehead, on a very thin compress, to avoid irritation of the skin, by thus favoring its free evaporation, is an excellent application.

Camphora, locally applied, is a valuable remedy in nervous headaches. The following formula may be employed :

102. R.	<i>Camphoræ,</i>	℥j	
	<i>Aceti,</i>	℥j	
	<i>Aquæ,</i>	℥j-ij.	M.

Another excellent local application is what is known in France as "L'Eau Sedative de Raspail," which is made as follows :

103. R.	<i>Aquæ ammoniæ,</i>	100 parts.	
	<i>Camphoræ,</i>	2 "	
	<i>Sodii chloridi,</i>	20 "	
	<i>Aquæ,</i>	900 "	
	<i>Olei rosæ,</i>	q. s.	M.

Dry Cupping is recommended by Dr. GRAVES.

Oleum Cajuputi. A few drops rubbed upon the painful part are often effectual.

Oleum Menthæ applied to the forehead and the base of the occiput often gives immediate relief. It is best dissolved in alcohol.

Shower Baths are often of great advantage in hysterical headache.

NEURALGIC HEADACHE.

Ammonii Chloridum, given when the pain is most intense, in the dose of gr. x-xx, sometimes affords prompt relief.

Cajuputi Oleum is used, both externally and internally, by Dr. COPLAND and others.

Camphora, in the form of "Raspail's sedative water," (F. 103,) is sometimes very useful.

Caffein. The citrate has been recommended in doses of gr. j. every hour, for some time before the expected attack. Coffee is much used by Belgian physicians. The juice of a lemon makes a valuable addition to a cup of strong coffee in these cases. Caffein may also be employed hypodermically,

Nitrite of Amyl, by inhalation, has been successfully employed in neuralgic headache, associated with a diminished supply of blood in the brain.

PERIODIC HEADACHE.

Arsenici Iodidum has been found, by Dr. WALSH, to give relief in the most violent cases of periodic headache. It should be given in doses of gr.

$\frac{1}{16}$ – $\frac{1}{12}$, twice a day, two hours after eating, and continued for a month or two.

Ammonii Carbonas and a saline purgative, at the commencement of an attack, are recommended by Prof. FLINT. (P. 51.)

Belladonna is sometimes of service as a prophylactic.

Caffea is a popular and efficient remedy.

Cannabis Indica. (See above, p. 50.)

Nectandra is a useful remedy in periodical headaches.

Potassii Bromidum often rapidly relieves headache of a paroxysmal character, with heat of the head and flushing of the face.

Quiniæ Sulphas, in small doses, is recommended, by way of prophylaxis, by Prof. FLINT.

Sulphate of Nickel was used successfully by Sir JAMES Y. SIMPSON. (*Medical Press and Circular*, December, 1868.)

Zincum. The various preparations of this metal, more especially the valerianate, may be employed with advantage in the intervals of the attacks.

RHEUMATIC HEADACHE.

Aconitum in small doses, either alone or in combination with other remedies, is often of service. Aconitia ointment is serviceable, especially in brow ague.

* *Sulphur* internally is often of great benefit in rheumatic headaches, and those which occur in women about the period of the cessation of the menses.

Alkaline treatment, if continued with regularity, will generally be followed by satisfactory results.

HYPOCHONDRIASIS, OR MELANCHOLIA.

DR. C. H. HUGHES, OF ST. LOUIS.

This able specialist calls attention (*Med. Review*, 1883,) to the association of visceral disorders with mental derangement of a melancholic character, and especially of the co-existence of glycosuria and hypochondria. He is satisfied that there is an intimate relationship between this symptom and nervous depression.

His belief is that the source of both disorders is hepatic, but behind that nervous; and that neurotic tranquilization, tonicity and reconstruction are the indications, and that *codeia* and *electricity* are the best special therapeutics of the trouble, with such other neurotic auxiliaries as the deranged nervous states mentioned may demand.

A. O. KELLOGG, M. D.

In the treatment of the early stages of this form of mental alienation, this writer (*Am. Jour. of Med. Sciences*, Jan., 1878,) says that the digestion and sleep, especially the latter, are to be improved. For this purpose, he has found the following drugs to be most useful, in the order given: *Hydrate of chloral*, gr. x-lx; *tincture of hyoscyamus*, f. ʒj-iiij; *liq. morphiae sulphatis*, f. ʒj-f. ʒiiij; or two or more may be combined, especially the two first mentioned. Opium often aggravates the insomnia. Traveling, so often recommended, is worse than useless. In the majority of cases, overwork or exhaustion from innutrition, lactation, etc., has led to the condition, and *rest* is imperatively demanded. The patient must be carefully nourished and sustained by a full, generous diet. From fear of biliousness or indigestion, he has probably fasted, abstained from food at least, and perhaps from drink; but as to the last, it is not improbable that he may have indulged excessively in stimulating beverages. These, in moderation, however, are not without their use; for in all such cases, there is generally deficiency of nerve and brain force, which, next to good, nutritious food, is supplied by wine, eggs, milk, malt liquors, iron, quinine, strychnine, arsenic, etc., as, in the judgment of the physician, may seem indicated.

W. W. GULL, M. D., LONDON.

The *constitutional treatment* is to be directed towards improving the general nutrition. The administration of *iron* is doubtless of great use to some anæmic patients, and *sea-bathing* frequently appears to exercise a very beneficial influence; but the first of these remedies is generally the most efficacious when taken in the form of the chalybeate waters of some foreign spa, and there is good reason to doubt whether both mineral waters and sea-bathing do not owe most of their apparent power to the moral influences of travel and change of scene and mode of life. The more specific nervous tonics, such as strychnia, quinia, or phosphorus, seem to exercise but a doubtful and exceptional influence.

While it is desirable to avoid concentrating the patient's attention on parts which are the apparent seat of morbid sensations, it is important to relieve him of the distress caused by real (though mere functional) disorders of the digestive system. Decided acidity of the stomach should be counteracted by the use of antidotes, of which none is more efficacious than *magnesia* in ten-grain doses thrice daily, with

gentian or cascarilla. The excessive or too prolonged use of alkalies is, of course, to be avoided. The distressing flatulence which is often one of the earliest, and also one of the most annoying symptoms, is greatly relieved by *creasote* (one drop in a pill two or three times daily) or the *infusion of valerian*. Alcoholic tinctures should be cautiously employed, if at all, for there is real danger of the patient getting to appreciate the comforting sensations given by the spirit so highly that he gradually takes to drink; this is particularly apt to occur in hypochondriacal women, and in patients exhausted by masturbation, or other venereal indulgence. The constipation, frequently so obstinate and troublesome, should be remedied, if possible, without drugs, so as to avoid stimulating the patient's love for self-doctoring in the direction of the habitual use of purgatives. Fruit, green vegetables, etc., and active bodily exercise, are desirable means of accomplishing this object.

Physical exercise should be amusing, and never produce severe fatigue. A short continuance of the malpractice of carrying exercise to the fatigue point, will usually suffice to produce a profound deterioration of the vigor of the nervous system, and an aggravation of the hypochondriacal fancies.

PROF. FELIX VON NIEMEYER, M. D., TUBINGEN.

The only way to cure a hypochondriac, is to rid him of his morbid sensations; but it is useless to dispute with him and try to convince him of the error of his ideas. In the first place, any existing derangement of the system which would cause a felling of illness in a sane person, should be corrected. The benefit derived from various *natural mineral springs* is no doubt due to their happy effect upon the gastric disorders which so frequently give rise to this disease. Preparations of *iron*, *sea-bathing*, and *cold foot-baths*, are frequently appropriate. Drastic *cathartics* must be used with caution, though they cannot be altogether dispensed with, and generally afford temporary relief. Patients should be warned against over-dosing. This also applies to the use of carminatives, for which they nearly always beg.

The *psychical* treatment should have in view the diversion of the attention from the sensory to the motor and intellectual spheres. This object will not be attained in educated patients by ordering long walks, wood-sawing, gymnastics, and mechanical pursuits, because the attention

is not diverted by such action out of the sphere of sensation. A hypochondriac merchant sawing wood is always busy with the thought that he is doing so because he is sick. General rules are of little value, as the directions must always be in accordance with the capacity, education and means of the patient.

NOTES ON REMEDIES.

Assafœtida, in combination with bitter tonics and mild aperients, may often be given with advantage in hypochondriasis attended with dyspepsia.

Cimicifuga is alleged by PHILLIPS (*Materia Medica*, 1879,) to exercise special powers in nervous and sexual hypochondriasis.

Oreasotum, one drop, in pill, two or three times daily, will greatly relieve the distressing flatulence of hypochondriasis.

Fel Bovinum Purificatum is advised by Dr. COPLAND in hypochondriasis attended with torpor of the intestines :

104.	R.	Fellis bovini purificati,			
		Pilulæ assafœtidæ compositæ,	āā	gr. xxx	
		Extracti aloes,		gr. xx	
		Saponis,		gr. x	
		Pulveris ipecacuanhæ compositi.		gr. viij.	M.

For thirty pills. One or two to be taken before dinner.

Hyoscyamus has been found of service by Dr. HARLEY, in occasional full doses, (f. ʒ iv-vij) of the succus.

Moschus, in the dose of one grain every two hours, often proves successful in the sleeplessness of hypochondriasis.

Narcotics. (See above, p. 82.)

Potassii Bromidum is an uncertain remedy in this disease, affording great relief in some cases, and proving inert in others.

Sumbul Radix. The resinous extract of this root, in doses of gr. j-ij, thrice daily, has proved beneficial in the hands of Russian physicians.

INSANITY.

DR. EDWARD C. MANN, NEW YORK.

In his treatise on *Psychological Medicine* (1884), this author sums up as follows the general treatment of the insane :

Primarily, give plenty of nutritive food, and give barks, wine and iron, or malt liquors, when indicated.

Warm clothing and bedding, and a moderately warm and dry atmos-

phere, and plenty of occupation and exercise—a *sine qua non* in the treatment of the insane, and change of scene and surroundings.

- | | | |
|---------|-----------------------|------------|
| 105. R. | Fl. ext. hyoscyami, | gtt. x |
| | Am. tinct. lupulin, | ℥j |
| | Camphoræ (Neergaard), | gr. v to x |
| | Syr. aurantii, | ℥ij |
| | Aquæ camph, | ℥vi. |

M. et ft. haust.

Useful in asthenic cases as narcotic and sedative.

A good sedative pill in general paralysis of the insane, and in incipient mental disorder, with some excitability, is the following:

- | | | |
|---------|--------------------|---------|
| 106. R. | Zinci valerianat., | ℥j |
| | Ext. belladonnæ, | gr. ij. |
- M. et ft. pil. No. 30. Sig.—Pill every two hours.

Narcotic friction of scalp with Ungt. Bellad. Comp. is also old, but useful at times.

- | | | |
|---------|------------------|-----|
| 107. R. | Ext. belladonnæ, | ℥j |
| | Morphiæ sulph., | ℥j. |
- M. et ft. ungt.

To subdue inordinate excitement, use the prolonged warm bath, with cold effusions to head, protracted an hour or two, and followed, if necessary, by pill of camphor, turpentine and hyoscyamus.

Narcotic baths of two pounds of henbane, belladonna, hemlock, and cherry laurel leaves, infused in hot bath, are excellent, even though old.

Tinct. digitalis, carried to sixty minims *ter die*, after a mercurial cathartic, produces sleep, quiet, and restoration to health in subacute mania.

In delirium tremens $\frac{1}{2}$ ounce doses of *digitalis*, or

- | | | |
|---------|----------------|------|
| 108. R. | Chloral hyd., | ℥ij |
| | Tr. hyoscyami, | ℥ss |
| | Syrupi, | ℥iv. |
- M. et sig. Dose, tablespoonful, to be repeated as necessary.

Monobromide of camphor in puerperal mania, after ten grains of calomel, followed by a saline, is a specific.

In nymphomania, four grains camph. monobromide every three hours after regulating secretion, will soon cure the patient.

If nymphomania is accompanied by clitoric irritation, we may, with advantage, remove the clitoris.

In uterine irritation in insanity, which keeps up sleeplessness, use suppositories of one grain *stramonium* at bed-times. It quiets ganglionic nerves of uterus efficiently, and have displacement and flexion remedied at once.

In dysmenorrhœa in insanity, use Scheiffelin's fluid *extract viburnum prunifolium* (black haw), \mathfrak{z} ss to \mathfrak{z} j, every hour until relief is experienced. Very valuable in ordinary dysmenorrhœa.

In hysteria and hysterical mania, the introduction of a little pill of ungt. belladonnæ comp. into the os uteri acts like magic.

109.	R.	Ext. belladonnæ,	\mathfrak{z} j
		Morph. sulph.,	\mathfrak{z} j.
	M.	et ft. ungt.	

A combination of iron, phosphorus, zinc and strychnia, antagonizes cerebral degeneration in insanity.

110.	R.	Ferri redact.,	gr. ii
		Zinci phosphidi,	gr. $\frac{1}{15}$ to $\frac{1}{5}$
		Strychniæ sulph.,	gr. $\frac{1}{60}$ to $\frac{1}{32}$.
	M.	et ft. pil. No. 1, t. i. d.,	after meals.

With a moist, relaxed state of the skin, cold extremities, shrivelled surface, pale complexion, and feeble circulation, use stimulants and full diet, and centric galvanization, carb, am. or old whiskey or port.

In great depression, melancholia, especially if religious or suicidal, a course of prolonged warm baths, with gradually increasing doses of morph. hydrochlorate, will antagonize the morbid psychic state and cure the patient quickly.

Enveloping the whole person in a sheet wrung out of hot mustard water, and putting the patient to bed, is a valuable sedative measure, with dry cupping at back of neck.

Use all your mental resources to weaken and dissipate false ideas, a happy mixture of kindness and firmness.

Remember, that hallucinations may be troublesome and disturb repose, but if the individual acknowledges their falsity, he or she is sane. Prolonged dry cupping may relieve this symptom.

We can control those tearing everything to pieces, leaving themselves naked, which is due to hyperæsthesia of the body, by monobromide of camph., Clin's capsules (four grs.) ter die.

Fluid extract *ergot* (Squibb's) in \mathfrak{J} ss. to \mathfrak{J} j. thrice daily, is the physiological antagonist to states of recurrent mania and chronic mania with lucid intervals, and also to epileptic mania; may be combined with the bromides.

In states of great motor excitement with homicidal impulses, fluid extract *conium* in $\mathfrak{J}\frac{1}{4}$ to $\mathfrak{J}\frac{1}{2}$ doses, or \mathfrak{J} j, may be used to quiet motor brain centres; produces quiet and muscular relaxations.

Puerperal mania is generally due to the absorption of the retained products of conception, which causes septicæmia, and we therefore give a mercurial cathartic, followed by salines, and four grains monobromide camphor ter die.

In hallucination with excitement, dry cups at back of neck, prolonged warm baths, add thirty grains sodium bromide, with thirty minims tinct. cannabis indica in combination, ter die, will produce quiet and tranquility. In dementia can. indica has no effect.

Nourishment.—The nourishment of a sane man is generally insufficient for an insane. Full feeding means tranquillity; insufficient diet, excitement.

Diuretics.—When the urine of the insane is scanty and high-colored, use diuretics, such as sp. ether., nit., liq. am. acetat., or inf. of digitalis.

Carefully examine daily the state of the pulse and skin, the head, the extremities, the tongue, the bowels, the urine, and regulate the catamenia. If the action of the liver is sluggish, give dilute nitro-muriatic acid, fifteen gtt. in \mathfrak{J} j of tinct. gentian. comp., ter die.

Restraint.—The entire disuse of restraint will soon be possible, as insanity is gradually assuming in our country a milder type than formerly. The disuse of restraint should be the rule and system.

Occupation and exercise in the open air for the insane cannot be insisted on too strongly. This, with full feeding and warm baths, together with change of scene, will cure many patients.

Epileptic and suicidal patients should always be watched throughout the night by a night attendant.

Sleep.—We should remember that sleep is the first of medicines as a physiological antagonist to the states of an inordinate degree of excitement and over-activity of nervous system or mental anxiety.

Massage and electricity are of inestimable service in the neurasthenia, which is often the incipient stage of insanity, and rest for the over-worked brain; the *galvanic current* antagonizes congestive states of the brain.

Proper diet and exercise, change of air and scene, and useful and agreeable occupation of the mind, are the most valuable means at our command in the treatment of insanity.

Disorders of the digestive organs, indicated by coated tongue, constipation, flatulence and morbid appetite, may be regulated by *mist. rhei, sodæ and gentian*, or the triplex pill, when well made.

In recent cases, rapid recovery may follow copious evacuations of the bowels and free diuresis, followed by a course of prolonged warm baths. The latter calms general mental excitement wonderfully.

The subsidence of excitement is not always the immediate precursor of convalescence. Having removed from your patient all moral causes of excitement, do not administer narcotics in large doses, but be content to apply remedies calculated to correct the disorders of the other bodily functions, and *time*, with good hygiene, will often cure your patient.

Sleeplessness, accompanied by high nervous excitement, is more safely combated by prolonged warm baths, with cold to the head, than by medicine, and tranquility and uninterrupted rest will often follow.

Insanity is a disease generally accompanied by bodily disorders, and these we must remedy. When the digestive organs have resumed their healthy condition, when the pulse becomes natural, and there is no unusual heat or dryness of the skin, we must rely on time and moral treatment to cure our patient.

Importance of Good Nurses.—Have no nurse for your sick insane patient, either male or female, with whom you could not place your wives, brothers or sisters, and always treat your patient with the utmost attention, delicacy and respect. They *always* respond to such treatment.

When the mind is weak, but slowly recovering its vigor, a good nurse will keep a patient ordinarily cheerful and happy, while a poor one will throw the patient into a state of anxiety and depression. A good nurse will have the *confidence, respect and affection* of her patient.

We must remember that a suicidal patient is a recoverable case, but they are *always* on the look-out for means of self-destruction, and the nurse must exercise unremitting, vigilant watching; when the suicidal act has once been meditated, *it is an imperative duty* to watch such a patient.

While we can often trust the veracity of the insane, as to going and coming punctually, it is not safe to do this when the hazard is that of

life in a suicidal patient. Such patients often throw the inexperienced and unskilled off their guard and commit suicide.

We can inculcate self-respect and self-control more often in mental disease than many imagine. Hardly any insane man or woman is beyond the reach of a motive, if rightly presented.

The word of a violent patient to be quiet, industrious, and orderly, is often to be relied on in hospital treatment, and we thus avoid the use of restraint.

DR. S. RIFAS.

The value of *digitalis* in many cases of mental disease is highly extolled by this writer (*El sigla medico*, 1883). It must, however, be confined to certain classes of cases. He defines the indications and counter indications as follows:—

Digitalis is indicated:—1. In acute mania from anæmia, where there is dilatation of the pupil, and feeble, rapid heart's action; 2. In chronic tranquil mania, especially those cases having hallucinations of sight and hearing; 3. In hypochondriacal mania, especially when there exists disease of the lungs, heart, or great vessels; 4. In "*manie anxieuse*;" 5. In suicidal melancholia; 6. In hypochondriacal melancholia; 7. In religious ecstasy; 8. In the melancholic form of general paralysis, particularly if there be an anæmia or a suicidal or hypochondriacal tendency; 9. In simple dementia.

Contra-indications to the use of *digitalis* are present:—1. In acute delirium in general, and in restless, violent, or incoherent delirium; 2. In chronic, following acute manias; 3. In acute or chronic hallucinations, which coincide with other mental affections caused apparently by over-nutrition of the brain; 4. In simple and chronic melancholia; 5. In cataleptic ecstasy; 6. In most forms of general paralysis.

DR. EDWARD C. SEGUIN.

In the treatment of insane and melancholic patients at home, this writer (*Opera Minora*, 1884) states that the first requisite is to secure a good nurse, and create a cheerful moral atmosphere. Many such patients will need building up with tonics and nutritious food. Phosphorus in the shape of "Thompson's solution" (see Index), iron, arsenic, strychnia and alcohol, are demanded. To produce sleep, *chloral* or *opium* are preferable to the bromides. *Cannabis indica* is also useful. Galvanization of the spinal cord has been used in the treatment of melancholia with apparent success.

The hygiene of the patient must be carefully watched, the bowels regulated, cold sponging advised, and the patient occupied and amused.

Where there is a decidedly suicidal impulse, or where food is persistently refused, it is better to place the invalid in an asylum.

NOTES ON REMEDIES.

Auri et sodii chloridum has produced excellent results in various forms of mental disorder. It is especially indicated where there is anemia, but is not suitable for plethoric subjects. The dose is gr. $\frac{1}{10}$ – $\frac{1}{20}$, in pill or dissolved in mint water.

Paraldehyde has been recently advocated by *Morselli* and others in dementia and acute mania. The ordinary dose is twenty to forty minims, but *Bartholow* recommends that the maximum dose, f. \mathfrak{z} ij–iij, be administered in such cases. It is a colorless liquid, of ethereal odor, and unpleasant taste.

Physostigma has led to remarkable improvement in some cases of dementia paralytica, but in others it has been administered without advantage.

Potassii iodidum proves of service in many cases of insanity, because they arise from syphilitic poison. In twenty-one cases of dementia paralytica, Dr. SNELL found signs of syphilitic infection in eleven. (*Schmidt's Jahrbücher*, 1883).

INSOMNIA OR SLEEPLESSNESS.

DR. G. EICKHOLT.

In a study of this disease (*Deutsche Med. Woch.*, Nov. 1883,) this writer divides it into three forms:

1. Insomnia from mental or physical over-exertion, including that from the excessive use of tea, coffee or opium.
2. Insomnia from disease, as neurasthenia, chronic gout, neuralgia, etc.
3. Insomnia from functional or organic disease of the brain.

The treatment begins with establishing careful hygienic regulation of the habits and diet. Tea and coffee must not be taken in the evening. Warm baths, a half hour in duration, before retiring, are beneficial. Hyperæmia of the brain may be reduced by wrapping the head in cloths wrung out of cold water. When anæmia of the brain is present, a glass of beer or hot punch acts as a hypnotic.

Of medicines, *bromide of potash* or *soda* stands first in neurasthenic and irritative cases. A full dose, about eighty grains, should be taken, well diluted, say in a tumbler of weak mint tea, about an hour before retiring. *Opium* may answer in anæmic, but is contra-indicated in uræmic individuals. *Chloral* is generally useful, but should be avoided where the heart is weak, where there is arterial disease, and in old age. *Cannabis indica* and *tannate of cannibine* are useful in neurasthenia and melancholia, but are to be avoided in cases of cerebral excitement. *Paraldehyd* should not be exhibited on account of its unpleasant taste and smell; and the same is still more true of *acetal*.

PROF. WILLIAM A. HAMMOND, M. D., NEW YORK.

The principles which should prevail in the treatment of wakefulness may be arranged into two classes:

1st. Those which, by their tendency to soothe the nervous system, or to distract the attention, diminish the action of the heart and blood vessels, or correct irregularities in their function, and thus lessen the amount of blood in the brain.

2d. Those which directly, either mechanically or through a specific effect upon the circulatory organs, produce a similar effect.

In slight cases, the measures belonging to the first class often prove effectual. Among them are music, monotonous sounds, gentle frictions of the surface of the body, soft undulatory movements, the repetition by the insomnolent of a series of words till the attention is diverted from the existing emotion which engages it, and many others of similar character. In persistent insomnia, however, these are nugatory.

Chief among the means embraced under the second head are those which tend to improve the general health of the patient.

In regard to food, while it is an error to suppose, as is generally done, that a moderately full meal, eaten shortly before bedtime, is necessarily productive of wakefulness, there is no doubt that this condition is induced by an excessive quantity of irritating or indigestible food. A hearty supper of plainly cooked and nutritious food rather predisposes to sleep. This is due to the fact that the process of digestion requires an increased amount of blood in the organs which perform it, and consequently the brain receives a less quantity. This hypnotic effect is neutralized, however, when the food is immoderate in amount or irritative in quality, as it then, either by the pressure upon the abdominal

vessels or through a reflex action on the heart, augments instead of diminishes the quantity of blood circulating in the brain. Attention should, therefore, be paid to the diet of the insomnolent. As a rule, people are under-fed. This is especially true of women. The tone of the system is thus lowered, and local congestions of different parts of the body are produced. If the brain be one of these, wakefulness results.

Most of the cases of insomnia in women are of the passive variety, and require not only nutritious food, but *stimulants*. Whisky is generally to be preferred to brandy and many kinds of wine. Nothing can be better, as a good stimulant, and at the same time tonic, than *Tarragona wine*, drank at dinner, to the extent of a glass or two. Next must be ranked good *lager beer*.

There are cases in which *coffee* induces sleep. Our author mentions several in which passive wakefulness was entirely and speedily cured by a cup of strong coffee, taken for three or four nights in succession, at bedtime. In females of languid circulation and a consequent tendency to internal congestions, it is particularly useful.

The employment of stimulants is only of service in the asthenic or passive form of insomnia; in the sthenic or active form they would, of course, increase the difficulty.

Physical exercise in the open air, extended to the point of inducing a slight feeling of fatigue, is productive of good effects.

The *warm bath* calms nervous irritability and determines blood from the head. Putting the feet in water of the temperature of 100° F. will often induce sleep, particularly in children, after other means have failed.

Cold water (32°F.) applied directly to the scalp, has a good influence in those cases in which the individual is strong, the heart beating with force and frequency, and the mental excitement great. It is not admissible in the asthenic form of wakefulness.

In insomnia dependent upon severe and long-continued mental exertion, all means will fail if the individual will not consent to use his brain in a rational manner. Proper intervals of relaxation must be insisted upon, and in some cases mental rest. Travel is always of the greatest advantage in such cases.

Among the purely medicinal agents, *bromide of potassium* holds the first rank. It diminishes the amount of blood in the brain, and allays any excitement which may be present in the sthenic form of insomnia.

The flushed face, the throbbing of the carotids and temporals, the suffusion of the eyes, the feeling of fullness in the head, all disappear as if by magic under its use.

C. HANDFIELD JONES, M. D., OF LONDON,

recommends, when sleeplessness depends upon hyperæsthesia, to give not only a *stimulant*, but also some *digestible nourishment*, about the time of going to rest, or even in the course of the night, when debility to a serious extent exists. A *mustard poultice* to the epigastrium is also an excellent hypnotic in some cases, probably owing to its stimulating a languid solar plexus. Among the various soporifics, it is doubtful whether any are more potent, especially for the weakly and hyperæsthetic, than prolonged exposure to the *cold open air*. This should be so managed as not to cause great fatigue, and be followed by a sufficient meal. The effects of a *monotonous voice* in inducing sleep are well known, particularly when the speaker or reader is dictating some matter not without a considerable tinge of dullness. The old monk's prescription for sleeplessness, viz., to tell your beads, was sound advice. *Hyoscyamus* frequently proves itself a really valuable remedy. It is well given in enema, in a dose not exceeding that for the mouth.

H. V. SWERINGEN, M. D., OF INDIANA.

Very much depends upon the cause of sleeplessness in the choice of remedies for it. In the sleeplessness of the exhaustive fevers, a glass of champagne, or whisky or brandy, or ale or a strong cup of coffee, or a dish of soup, will often induce a most refreshing sleep, when all other remedies would fail. And, again, there are times in which these would do harm, or when the following would be useful :

111. R.	Potassii bromidi,		
	Chloral hydratis,	āā	ʒ iij
	Morph. sulphatis,		gr. j
	Syr. Tolutani,		
	Aquæ menth. pip.,	āā	f. ʒ iij.

A tablespoonful in water, every hour or half hour, until sleep is induced.

DR. WILLEMIN, OF FRANCE,

In the *Archives Générales de Médecine*, June, 1877, finds most cases of insomnia to be produced either by (1) anæmia, or by (2) irritation. Hygienic measures should never be neglected, and are often sufficient.

Of hypnotics the opium alkaloids are best in insomnia from pain. Of these, *codeia* leaves the least *malaise*. *Bromide of potassium* is excellent where there is excitement, but is contra-indicated in anæmia. *Hydrate of chloral* is very efficacious, but must be given with great caution where there is dyspnœa, heart disease, or great debility. *Quinine* relieves cerebral congestion with good effect. In aged and debilitated persons, tonics, stimulants and hydropathy are better than anodynes.

SOME NEW HYPNOTICS.

Several hypnotics have within the last few years (1882-4) been introduced to the profession, and merit attention.

Jamaica dogwood, or the bark of *Piscidia erythrina*, L., the fluid extract, in doses of ʒss-ʒj, in many cases produces the hypnotic effect of opiates without their ill effects. Various preparations of this drug, also known as Jamaica Dogwood, are manufactured by Messrs. Parke, Davis & Co., of Detroit, to which energetic house the introduction of this agent is due.

Recent medical authorities report that the action of *piscidia erythrina*, when taken internally, appears to be a direct sedative to the nerve-centres; that it causes sleep without producing the cerebral hyperæmia which often follows opium and its active principles. The sleep is tranquil and refreshing; it is said to soothe bronchial cough, and to moderate the paroxysms of asthma and nervous coughs. In view of the fact that opium is often not well borne when an anodyne hypnotic is required, it seems that the drug in question deserves further study.

Napelline has been strongly urged as a hypnotic by Dr. LABORDE, of Paris, (*Jour. Therapeut.*, 1883). He asserts that it possesses physiological properties which, while approaching in their fundamental characteristics to those of crystallized aconitia, sensibly differ from this, both in their much less activity, and by hypnotic and somniferous effects which do not belong to it. Its effects are chiefly manifested in local painful affections—especially in neuralgias—and in excessive excitability of the nervous system, with persistent insomnia.

Codeia phosphate. In the *Memorabilien*, 1883, Dr. FRONMULLER recommends this preparation as superior to morphia, milder in its action, more soluble, and admirably suited to hypodermic injection in neuralgic pains, etc.

Tannate of Cannabine. In a report in 1883, Dr FRONMULLER speaks of tannate of cannabine as a very efficient medicament to induce sleep without any disagreeable after effects. The dangers of the abuse of morphine are so great that any other agent capable of inducing similar hypnotic effects, would be a veritable boon to the practitioner.

Mr. HILLER, of London, has experimented with the medicament, and has found that it gives good results, particularly in the milder forms of sleeplessness. Its effects are less marked in serious cases of delirium tremens, in mania, and in subjects already habituated to narcotics. It may be ordered in powders:

112. R. Tannate of cannabine,	gr. xv
Sacch. alb.,	gr. xx.
M. et divid. in cht. No. iv.	
S.—One or more at bedtime.	

Dr. EICHHOLT (*Deut'sch. Med. Woch.* 1883,) puts the dose at not over one grain, and recommends it especially in neurasthenic insomnia.

Paraldehyd. This is a modification of aldehyd, and has the formula $C^6H^{12}O^3$. It is a colorless liquid, soluble in eight volumes of water, and is both anæsthetic and hypnotic.

Dr. VICENZO CERVELLO, of Palermo, says that paraldehyde in doses of thirty to forty-five grains induces a quiet, reparative sleep, not followed by headache, loss of appetite, or mental confusion, as after chloral. It is very useful in the periodical mania of progressive paralysis, in acute and chronic mania, delirium tremens, etc.

Dr. MORSELLI, in cases of mental excitement, finds this drug an efficient hypnotic without depressing the circulation. On the disease itself, it has no effect. In cases of depression, as in melancholia, it is less efficacious.

In idiocy with agitation, in the excitement of general paresis and of hysteria, the drug is useful, but only because of its hypnotic effect, which makes it useful in the insomnia of fever, rheumatism, gout, prurigo, chloro-anæmia, etc.

Diethylacetal. This chemical compound ($C^6H^{14}O^2$) has been presented by M. VON MERING (*Berliner Klin. Wochen.*, 1883,) as an excellent substitute for chloral. It is a substance of a bitter, slightly burning taste, soluble in eighteen times its volume of water, and freely in alcohol.

At the dose of from ten to twelve grams, it produced sleep lasting an

entire day in six subjects, and induced a marked attenuation in subjects suffering from severe injuries.

In a case of dementia, where continual agitation and persistent insomnia were the main features, an entire night's rest was obtained after the administration of five grams of this medicament, and the patient remained calm all the following day. This salutary effect continued, and an extended use in a series of cases confirmed the general good effects. Others have found it to derange digestion and induce nausea.

According to Dr. THOMAS LEGARI, of Charleston, S. C., the best treatment for insomnia is to be found in the following points (*Med. Rep.*, 1884):

1. Retire early to bed.
2. Eat little, and always some hours before going to bed; cold food only should be taken for supper.
3. The cares and burdens of the mind should be put aside; never read or study in bed.
4. The bed-chamber should contain pure, sound air, and be roomy and high, if possible, and the windows should be always kept open except in the night-time.
5. When in bed, endeavor to lie horizontally, with the head slightly raised. If there is any forced or constrained posture, making the body form an angle, circulation in the stomach is checked, and a free and uninterrupted circulation of the blood is defeated.
6. It is improper to have a light burning in the bed-chamber during the night; our senses should not be acted upon by external impressions.
7. Endeavor to sleep not less than six or more than eight hours in the twenty-four.

NOTES ON REMEDIES.

Aconitum. Dr. ANSTIE says, "I have often seen insomnia accompanied by a harsh, dry skin, yield to the administration of aconite in repeated small doses." One drop of the tincture in a teaspoonful of water every fifteen minutes.

Alcohol. Dr. JAMES SAWYER observes (*Lancet*, June, 1878,) "To many people a 'nightcap' of toddy is a superfluous and hurtful luxury. But it can give, perhaps, better than anything else, rest and sleep to the exsanguine and worried brain."

Atropia. This alkaloid is combined by Dr. ROBERTS BARTHOLOW with morphia to procure sleep, care being taken that there is no excess of atropia, or an amount of atropia sufficient to antagonize the cerebral effects of the

morphia. They should be used in the proportions of gr. $\frac{1}{120}$ — $\frac{1}{96}$ of atropia to gr. $\frac{1}{4}$ — $\frac{1}{2}$ of morphia. As the susceptibility to atropia varies immensely, the precise quantity to be employed in any case must be regulated accordingly.

Cannabis Indica is often prescribed in sleeplessness, alone or combined with hyoscyamus. Dr. FULLER gives it in chronic rheumatism, to induce sleep, and Dr. OXLEY, (*Liverpool Medical and Surgical Reporter*, 1868,) in chorea, in doses of $\mathfrak{m}\text{x}$ of the tincture, for a child of seven, and so on. Dr. TILT gives one-grain doses.

* *Chloral* is a valuable remedy for sleeplessness in exhaustion of the brain through severe mental application or excitement of feeling, in convalescence from acute febrile diseases, and in insanity. It should only be used as a temporary remedy, when it is necessary that we should at once secure a fair amount of sleep. The patient ought never to be allowed to swallow this dangerous but valuable drug whenever he feels disposed; he ought only to take it upon the special prescription of his doctor.

Hyoscyamus is very variously esteemed by different writers. It is an uncertain drug, most useful where the insomnia is purely psychic. It acts better combined with quinia. (TILT.)

Ignatia, in small doses, is highly praised by H. G. PIFFARD, in insomnia from nervous erethism. In such doses, he says it is “*par excellence*, the controller of functional phenomena of the cerebro-spinal axis.” (Note to PHILLIPS’ *Mat. Med.*, 1879.)

Lupulin is often convenient in the insomnia of the aged.

Moschus has proved eminently successful in the hands of Dr. GRAVES, in persons of irritable and nervous disposition. He administers it in one-grain doses every two hours during the day, either alone or with assafoetida, in order to procure sleep at night.

Opium and its alkaloids must be used cautiously in simple wakefulness. Morphia is the most hypnotic. Next, narceia and codeia.

* *Potassii Bromidum* is a powerful hypnotic, especially in cases of overtaxed mental energy or emotion, and in the sleeplessness of convalescence from fever, at the termination of acute diseases, or after surgical operations. In the well-nourished, it is, says Dr. JAMES SAWYER, by far the best hypnotic. It soothes the irritated and irritable cerebral cells; it is a direct and absolutely safe brain sedative, and it is marvelously powerful in producing nervous calm. But it must be given in full doses, thirty to sixty grains at bedtime. It is well to conjoin with it some drug which will favor the contraction of weakened cerebral vessels; for this purpose we may give tincture of ergot or tincture of digitalis, one or both. In many cases of chronic wakefulness, arising from mental strain, the patient is distinctly anæmic. Unless the anæmia be remedied, the insomnia cannot be cured. (*Lancet*, June, 1878.)

Sumbul, gr. xx, may be tried, but this drug seems to have disappointed expectations.

EXTERNAL REMEDIES.

Cold and Warm Baths, etc. Sleep may often be induced by the temporary application of cold to the general surface of the body. A person who has been lying awake will often fall asleep at once after getting out of bed and sousing his head, neck, and hands in cold water, or after standing at the bedside until he feels chilly, and turning over, shaking up, and cooling his pillows and the bed-clothes. So also warm sponging and warm foot-baths, before retiring, will often induce sleep. Dr. ANSTIE makes the important observation that the action of all narcotics is greatly aided by heat to the extremities, and a hot mustard foot-bath, on going to bed, will enable the patient to sleep with one-third of his ordinary hypnotic dose.

Electricity. When sleeplessness results from an overworked brain and consequent paresis of the vaso-motor nerves, the stimulus of electricity has been resorted to. ALTHAUS recommends this treatment. Two large pads are used with a Weiss' constant battery of from ten to fifteen cells. One pad is placed over the nape of the neck, the other, which can be conveniently made of an old reflector, and covered with chamois leather, is placed over the stomach. The anode is applied to the back, the cathode to the stomach, for about half an hour at a time.

NEURALGIA.

DR. E. C. SEGUIN.

In the treatment of trigeminal neuralgia Dr. SEGUIN employs *aconitia* in the following formula:

113.	R.	Aconitiæ (Duquesnel's),	.006
		Glycerinæ,	
		Spts. vini rect.,	ââ. 4.
		Aq. menth. pip.	ad 62.

A teaspoonful two or three times a day on an empty stomach.

In some cases he has used .008 or even .01 aconitia in the same formula. The susceptibility of patients to aconitia varies greatly. (*Opera Minora*, 1884.)

He has obtained "remarkable results" with this formula, giving relief in some most severe cases of epileptiform trigeminal neuralgia. It

must be remembered however that in the treatment of chronic neuralgia, and many neuroses, it is necessary to obtain the physiological effect of the drug employed in order to do good. The drugs most esteemed by Dr. S., in this disease, are aconitia, arsenic, iodide of potassium, gelsemium, belladonna, quinia, morphia, and phosphorus. The last mentioned is particularly efficacious in

THOMPSON'S SOLUTION :

114. R.	Phosphori,	.06
	Alcoholis,	20.
Dissolve with heat.		
	Glycerinæ,	48.
	Alcoholis,	8.
	Essentiæ menth. piper.,	1.2
Mix the two solutions.		
Dose, one teaspoonful—about gr. $\frac{1}{20}$ of phosphorus.		

Full doses should be given, about one teaspoonful every three or four hours. Dr. SEGUIN has cured a severe chronic facial neuralgia in twenty-four hours with this preparation.

Where there is nervous prostration, cerebral anæmia or melancholia, he combines it as follows :

115. R.	Sol. phosphori (Thompson),	1 part
	Olei morrhuæ,	4-6 parts.
A teaspoonful after each meal.		

FRANCIS E. ANSTIE, M. D., LONDON.

This skillful practitioner, in his classic work, *Neuralgia and its Counterfeits*, emphasizes strongly the importance of an abundant and nutritive diet, defective nutrition being nearly always present. His general plan of treatment is as follows :

1. CONSTITUTIONAL TREATMENT.—(a) *Nutritive remedies* : fat, cod-liver oil, butter, etc. (b) *Preparations of iron* are useful (the carbonate, especially,) only where there is actual anæmia. (c) The *nerve tonics* (quinine, arsenic, and zinc,) are beneficial in some cases, and not so in others. Sulphate of quinine is useful in malarial neuralgia, and also, in small doses, in some cases of non-malarial neuralgia, especially in ophthalmic branches of the fifth pair. Arsenic is useful, and more widely so than quinine, in both the above classes of neuralgia. Valerianate of zinc has also been employed with benefit. (d) *Constitu-*

tional remedies directed against a real or presumed depravation of the blood by some special poison; as, iodide of potassium and corrosive sublimate against syphilitic neuralgia; colchicum against gouty neuralgia; and iodide of potassium against rheumatic neuralgia.

2. NARCOTIC STIMULANT REMEDIES FOR NEURALGIA.—At the head stands *opium*, represented by *morphia*. Next, *belladonna*, or more exactly, *atropia*. The *subcutaneous injections* of these substances are of considerable benefit. Best medium, hypodermic dose of morphia gr. $\frac{1}{6}$; of atropia gr. $\frac{1}{20}$ – $\frac{1}{30}$. Belladonna (gr. $\frac{1}{6}$ – $\frac{1}{2}$ of the extract) given by the mouth, has a special influence on painful affections of the pelvic organs. *Indian hemp*, internally, (in pills of gr. $\frac{1}{4}$ – $\frac{1}{2}$, one every night,) is very effective in certain forms of neuralgia, especially migraine and clavus. *Chloride of ammonium* (doses of gr. 10–20) is useful in migraine, clavus, intercostal and hepatic neuralgias. *Sulphuric ether* is supremely useful in certain visceral neuralgias, as gastralgia, uterine, or ovarian neuralgia. *Turpentine* pearls are beneficial in sciatica. *Aconitia* is, ordinarily, too depressing. *Alcohol* is most useful taken at meals, or especially for the relief of pain.

3. EXTERNAL REMEDIES.—Flying *blisters*, of extreme and general use; *mustard plasters*, liniments, and ointments, (best is chloroform, diluted with seven parts of oil liniment;) and *electricity* (continuous low tension current from BUNSEN'S or DANIELL'S battery), are recommended.

PROF. C. E. BROWN-SÉQUARD, M. D., OF PARIS.

116. R.	Extracti belladonnæ,	gr. $\frac{1}{6}$.
	Extracti stramonii,	gr. $\frac{1}{5}$.
	Extracti cannabis indicæ,	gr. $\frac{1}{4}$.
	Extracti aconiti,	gr. $\frac{1}{3}$.
	Extracti opii,	gr. $\frac{1}{2}$.
	Extracti hyoseyami,	gr. $\frac{2}{3}$.
	Extracti conii,	gr. j.
	Pulveris glycyrrhizæ,	q. s.

For one pill.

According to circumstances, BROWN-SÉQUARD gives, without producing any great constitutional disturbance, three, four, and even five pills in a day, and sometimes in about eight or ten hours, for the relief of neuralgic or other pains. There must be, therefore, some influence exerted by some of these substances upon the others, diminishing their bad and not their good effects. He recommends the following *hypodermic injection*:

117. R. Morphiae sulphatis, gr. $\frac{1}{2}$ - $\frac{3}{4}$
 Atropiae sulphatis, gr. $\frac{1}{25}$.
 For one injection, in gtt. xx of distilled water.

The doses BROWN-SÉQUARD at first employed were gr. $\frac{1}{2}$ of the sulphate of morphia to gr. $\frac{1}{50}$ of the sulphate of atropia. He now employs those given above, the antagonistic effects of morphia and atropia on the brain rendering it possible, while securing the good effects against pain of the two remedies to use safely, or at least without great or lasting cerebral or cardiac disturbance, large doses of these narcotics.

In this connection, we may give the doses employed by other authorities in administering morphia and atropia subcutaneously in neuralgia.

DR. CHARLES HUNTER, of London, says, as a rule, never to use, in the first injection, in any case, more than one-half the stomachic dose of these alkaloids for males, and not more than a third for females.

Dr. RUPANER, of New York, places the minimum dose of morphia at gr. $\frac{1}{8}$; the maximum, gr. $\frac{3}{4}$; the minimum dose of sulphate of atropia, gr. $\frac{1}{50}$; the maximum, gr. $\frac{1}{10}$.

Dr. TILT, of London, states that the initial hypodermic dose of morphia (used alone) for a woman, should not exceed gr. $\frac{1}{6}$ (the acetate being the salt he prefers), and that of atropia should not exceed (used alone) gr. $\frac{1}{10}$. In combination, gr. $\frac{1}{6}$ of the sulphate of morphia with gr. $\frac{1}{50}$ of the sulphate of atropia.

DR. ROBERTS BARTHOLOW, PHILADELPHIA.

The *deep injection of chloroform* for the relief of neuralgia has been practiced by this physician and others with success. He gives the following directions regarding it:

1. *The Method.* The injection consists of five to fifteen minims of chloroform, thrown into the part by a hypodermatic syringe. This amount is sufficient to produce the best results. The needle is passed into the part deeply enough to reach the neighborhood of the affected nerve. The chloroform is then injected slowly, so that it may diffuse into the tissues immediately adjacent to the sheath of the nerve. The action is local, and not systemic. He has heard with amazement of the subcutaneous use of chloroform, as morphia is employed. Thrown into the subcutaneous connective tissues, not only is the pain great, but there is danger of producing abscess and sloughing.

2. *The Site of the Injection.* This method is applicable only to su-

perforial neuralgia. It is essential to success that the chloroform reach the tissues about the trunk of the nerve. In the case of tic-douloureux, the chloroform must be deposited near the foramina at which the various divisions of the nerve emerge; in the case of sciatica, about the ischiatic notch. The same rule obtains in regard to other forms of neuralgia.

3. *Theory of its Mode of Action.* The local effect of chloroform is to produce anæsthesia and analgesia. It accomplishes this result in part by suspending the power of the nerve to transmit impressions, in part by the local swelling which acts mechanically on the nerve-trunk. For the production and perception of pain, it is necessary, first, that an irritation be made at the periphery; second, that end-organs exist for the reception of the painful impression; third, that conducting fibres are in a position to transmit the pain to the centres of consciousness; fourth, that these same conducting fibres are in a condition to refer outwardly the painful impression to the point at which it originated. It follows that, when the nerve-trunk is rendered incapable of transmitting the painful impression, the pain ceases to exist, or, what is equivalent, ceases to reach the centres of conscious impressions.

The same writer extols a combination of anodyne and hypnotic remedies as follows:

118. R.	Chloral hydratis,		
	Camphoræ pulv.,	āā	℥j
	Morphiæ sulphatis,		gr. ij
	Chloroformi,		℥xl.

The morphia will dissolve more readily if aided by gentle heat. Atropia may be added to this solution, if desired. The proportion of atropia, as also of morphia, is determined by the uses to which the remedy is to be devoted. To the 160 minims of the above solution, $\frac{1}{2}$ of a grain of atropia may be added. Twenty minims of this solution will contain $7\frac{1}{2}$ (seven and a-half) grains of chloral, the same quantity of camphor, $\frac{1}{4}$ of a grain of morphia, and $\frac{1}{96}$ (one ninety-sixth) of a grain of atropia.

This combination, Dr. B. extols as of wide application in all forms of neuralgic suffering. Locally, it may be painted on the surface, in superficial neuralgia, in *coëcydynia*, in *acute rheumatism*, *cancer*, *ulcers*, etc.

Internally, it should be administered in capsules, or on a lump of

sugar, not mixed with water. As an anodyne, the dose is gtt. x-xxx. Whenever pain or convulsive nervous action is to be allayed, Dr. B. believes no combination of remedies will be found equal to this one in general usefulness.

PROF. J. M. DA COSTA, M. D. PHILADELPHIA.

119. R.	Aconitiæ,	gr. iv	
	Veratriæ,	gr. xv	
	Glycerinæ,	f. ʒ ij	
	Cerati,	ʒ vj.	M.

To be rubbed over the painful parts, care to be taken to see that there is no abrasion of the skin.

(Prof. GROSS sometimes employs veratria ointment in cases of neuralgia, of the strength of ʒj to ʒj.)

120. R.	Potassii bromidi,	ʒ ss	
	Succi conii,	f. ʒ x	
	Aquam cinnamomi,	ad f. ʒ iij.	M.

A desertspoonful to be taken three times a day.

Useful in epileptiform neuralgia.

121. R.	Strychniæ sulphatis,	gr. ss	
	Quiniæ sulphatis,	gr. xv	
	Cinchoniæ sulphatis,	ʒ j	
	Pulveris rhei,	ʒ ss	
	Extracti gentianæ,	q. s.	M.

Divide into thirty pills; one to be taken four times a day, in intercostal neuralgia of malarial origin, associated with constipation.

DR. NATHAN S. DAVIS, OF CHICAGO,

states in his *Clinical Lectures on Important Diseases*, 1874, that in many severe and obstinate cases of neuralgia, sciatica, tic-douleureux, etc., careful examination will reveal traces of inherited or acquired syphilis. In such, the only hope of recovery must be founded on an effort to change the diathesis. The means best calculated to effect this are:

1. Regular and judicious exercise in the open air.
 2. A plain nutritious diet, chiefly of milk, farinaceous articles and fruits, with absolute exclusion of all alcoholic drinks, tobacco, and *strong* tea and coffee.
 3. The use of alteratives which will not depress the system.
- To effect the last mentioned, small doses of the iodides and mercury are well to begin with, as

122. R. Sodii iodidi, ℥ iij
 Hydrarg. chloridi corrosivi, gr. j
 Extracti conii fluidi, f. ℥ j
 Syrupi simplicis,
 Aquæ menthæ, āā f. ℥ j. M.

A teaspoonful before each meal and at bedtime.

As soon as this preparation has had a visible effect on the gums or the Schneiderian membrane (if the latter has been involved), it should be discontinued, and the following given in its place :

123. R. Potassii iodidi, ℥ iij
 Potassii bromidi, ℥ vj
 Extracti conii fluidi, ℥ j
 Aquæ menthæ, ℥ iij. M.

A teaspoonful three or four times a day.

If, under this treatment, which must be continued six or eight weeks, the patient becomes debilitated, some direct tonic must be given. The best Dr. DAVIS has used in such cases is the *syrupus ferri pyrophosphatis* f. ℥ j half an hour before each meal ; or, a pill as follows :

124. R. Ferri citratis, gr. ij
 Strychniæ, gr. $\frac{1}{30}$. M.

For one pill. One thrice daily before meals.

In many cases of uncomplicated neuralgia, Dr. DAVIS has derived great advantage from *belladonna*, in such combinations as the following :

125. R. Ætheris nitrici, f. ℥ iij
 Tincturæ belladonnæ, f. ℥ ij. M.

A teaspoonful in sweetened water every three or four hours.

126. R. Chloroformi,
 Tincturæ belladonnæ, āā f. ℥ iij
 Syrupi aurantii corticis, f. ℥ iij. M.

A teaspoonful every two hours until the pupil becomes slightly dilated, when the interval between the doses should be doubled.

127. R. Tincturæ belladonnæ,
 Tincturæ gelsemii, āā f. ℥ ss. M.

Twenty drops every two or three hours.

PROF. S. D. GROSS, M. D., PHILADELPHIA.

128. R. Quiniæ sulphatis, ℥ ij
 Morphiæ sulphatis, gr. j
 Strychniæ, gr. $\frac{1}{3}$
 Acidi arseniosi, gr. j
 Extracti aconiti, gr. x. M.

Divide into twenty pills. One to be taken three or four times a day. Add to the recipe, ferri sulphatis ℥ ij, if the system be anæmic.

This pill is useful in a great variety of cases of neuralgia. Its effects should, of course, be carefully watched.

PROF. WILLIAM A. HAMMOND, M. D., NEW YORK.

129. R. Extracti belladonnæ, gr. v.
Divide into twenty pills. One thrice daily, the dose to be increased as necessary.

The use of belladonna is chiefly to change the habits of the system. This drug, although at one time much overlauded, is very efficient in the treatment of neuralgia. Our author has not used atropia often, as the dose is difficult to graduate.

Hypodermic injections of morphia may be used during the paroxysms of pain. In their use, avoid the face; a good point is the inside of the arm.

130. R. Tincturæ aconiti, f. ℥ ss.
Rub with a rag upon the painful part, until a sense of pricking is felt.

This is next in value to the subcutaneous use of morphia. The action is often very powerful. Dr. H. once caused temporary paralysis of the arm, in a lady, by the too free application of the tincture.

Chloroform may be used externally, internally, or by inhalation, not carried to insensibility. Repetitions of the inhalation may break up the paroxysm.

Hypophosphites are useful; may be given in doses of from ten to twenty grains. They act by setting free phosphorus in the stomach.

Galvanism.—The direct galvanic current may be successfully used both for the relief of the paroxysm and for breaking up the habit of the disease. For the first, apply the poles so that the position is near the seat of the greatest pain, and pass the current continuously for several minutes. To change the habit of the system, apply one pole to the nape of the neck, and the other over the course of each sympathetic nerve, moving it along the neck. DUCHENNE says that neuralgia can generally be relieved by faradization. Dr. H. has not been so uniformly successful, but often effects a cure by the application mentioned.

The use of tea generally aggravates neuralgia, while coffee, on the contrary, does not, but if strong, is often of service.

In *neuralgia of the fifth nerve*, various authorities have recently re-

ported favorably of *croton-chloral*. Dr. E. P. HURD, of Massachusetts, has employed it in various cases of intractable neuralgia, in the following formula :

- | | | | |
|---------|-------------------|----|-----------|
| 131. R. | Croton-chloralis, | 3j | |
| | Glycerinæ, | | |
| | Syrupi aurantii, | āā | f. 3j. M. |
- A teaspoonful as required.

Dr. JULES WORMS, after several experiments made with a view of rendering this drug less repugnant to the taste, recommends the following formula :

- | | | | |
|---------|--------------------|-----------|----|
| 132. R. | Croton-chloral, | 1 gram. | |
| | Glycerine, | 60 grams. | |
| | Water, | 60 grams. | |
| | Oil of peppermint, | 3 drops. | |
| | Simple syrup, | 25 grams. | M. |
- This amount at one dose.

Intercostal neuralgia or pleurodynia, a complaint which is often a painful and annoying one, may generally be relieved completely by applying two or three strips of plaster firmly round the side over the seat of pain. The emb. roborans is a good form of plaster for these cases. It is desirable that the practitioner should himself see that the application is properly made, and that it is not left to the patient, as the result entirely depends on its being accomplished in an effectual manner.

- | | | | |
|---------|----------------------|----------|----|
| 133. R. | Extracti belladonnæ, | 3 ss | |
| | Chloroformi, | m℥ | |
| | Glycerinæ, | f. 3 ss. | M. |

Anoint with liniment, several times a day, the affected parts in neuralgia of the anus, when there exists neither fistula nor any lesion of the rectum.

GUY'S HOSPITAL, LONDON.

- | | | | |
|---------|------------------------------|----|---------|
| 134. R. | Liquoris plumbi subacetatis, | | |
| | Tincturæ opii, | | |
| | Mellis, | āā | f. 3 ij |
| | Confectionis rosæ, | | 3j. M. |
- Make a liniment.

This is known in the pharmacopœia of the hospital as *linimentum plumbi opiatum*, and is much used.

NOTES ON REMEDIES.

**Acidum Arseniosum* is a remedy of a deservedly high reputation in neuralgia,

having a wider range of application than quinine. In malarial cases it should be given in larger doses (℥x-xxx, thrice daily, of Fowler's solution) than in non-malarial. In the latter, ℥v of Fowler, or gr. $\frac{1}{16}$ of the *arsenate of sodium*, in pill, with extract of lupulin, will prove effectual in the cases amenable to this remedy. It may also be given hypodermically and endermically.

** *Aconitia*. (See F. 128.) Dr. E. C. SEGUIN states that Duquesnel's aconitia, in doses of $\frac{1}{84}$ to $\frac{1}{200}$ of a grain, three times daily, is a powerful agent for relieving and curing trigeminal neuralgia. Form for administration :

135. R.	<i>Aconitiæ</i> , (Duquesnel's,)	gr. $\frac{1}{16}$	
	Glycerinæ,	3j	
	Alcohol,	3j	
	Aq. menth. pip.,	ad 3ij.	M.

One teaspoonful (one hundred and fortieth of a grain) two or three times a day, on an empty stomach.

Prof. GUBLER also obtained some brilliant successes with this substance. He said, indeed, he had never met a neuralgia of the fifth pair which resisted it. He used the *nitrate of aconitine*, in doses of gr. $\frac{1}{40}$, in solution, or Hottot's aconitine, dose, gr. $\frac{1}{80}$, pushing it up slowly to gr. $\frac{1}{2}$, if the disease did not yield. He disapproved of pills, as they may not be dissolved.

Aconitiæ Oleas. This preparation has been found by Dr. H. A. KELLY, of Philadelphia, (*Med. News*, 1883,) very effective in severe trigeminal neuralgia. Three drops of the two per cent. solution is gently rubbed into the skin over the painful part. Complete relief is afforded in from three to five minutes.

* *Aconitum* is a remedy of great value in chronic cases of neuralgia, when the secretions are in order, and the general health not much impaired. Its use should, therefore, be often preceded by purgatives and tonics. In acute cases it is not indicated before all signs of active disease have been subdued by other means. Its external use should usually precede its internal administration, which need only be resorted to in case of failure of the former. The combination of quinine with aconite seems to increase its efficacy, and add permanency to the relief afforded by it. This combination is especially indicated in anemic or debilitated patients, where there is no apparent nerve lesion or exciting cause.

Æsculus Hippocastanum. The alkaloid of horse chestnut, *esculin*, has been given in doses of gr. xv, twice daily, with success, by French physicians, in the treatment of periodic neuralgia.

Æther. In simple neuralgia, particularly of the nerves of the head, Prof. STILLÉ recommends combining the anodyne effect of ether with its revulsive effects, by applying it on small compresses covered with some solid substance, immediately over the points where the painful nerve emerges from its bony canal, or where it in any other way becomes superficial.

Alcohol. Dr. ANSTIE has several times observed, in acute neuralgia, that after large doses of various narcotics had been tried in vain, the first real and substantial relief was obtained by the use of a moderate dose of alcohol; the consequence of overdosing the remedy is, however, nearly always disastrous.

Ammonii Chloridum is often of signal benefit, in doses of twenty to thirty grains.

Anthemis. Chamomile, in substance, in doses of gr. lx, or in a very concentrated infusion, is administered with success in France, in neuralgia of the fifth nerve.

Apiol is said to be of service in intermittent neuralgia.

Beberia Sulphas has proved successful in some cases, after the failure of quinine. In doses of gr. x, night and morning, it is more effective than in smaller doses.

**Belladonna*, whether used externally or internally, ranks among the first of the sedatives and anodynes of service in neuralgia. (P. 100.) It occasionally fails of effect, however. BROWN-SÉQUARD combines it with six other narcotics, for the relief of neuralgic pains. (F. 116.) *Atropia* is preferred by some to the preparations of belladonna, but the dose is more difficult to graduate. The sulphate is the most soluble.

Caffea is much employed by Belgian physicians. Prof. HAMMOND finds that it is often of service, but that tea generally aggravates neuralgia.

Cannabis Indica is ranked as a neuralgic remedy next in value to opium and belladonna and their alkaloids.

Carbolicum Acidum, by hypodermic injection, gtt. ss., has been shown to be efficacious by Dr. SCHRUMPF, of Strasbourg (*Gaz. Med. de Strasbourg*, 1883). He has never found any ill results, and usually four or five injections will cure any idiopathic case of sciatica, lumbago, pleurodynia, etc.

Chloroformum always acts as a palliative in, but rarely cures, neuralgia.

Conium is inferior to opium and belladonna.

Croton Chloral Hydrate, in doses of gr. ij every hour, has been found promptly to mitigate paroxysms of neuralgic pain.

Cupri Sulphas Ammoniatum. The value of ammoniacal sulphate of copper in the treatment of this affection has been asserted by M. FERÉOL, in a communication, in April, 1879, to the Académie de Médecine. He states that in cases in which every treatment has failed, even the administration of gelsemium and of aconitia, a cure or remarkable relief may be obtained to the most severe symptoms by this drug. The dose is gr. j three times daily, and may safely be increased to gr. v.

Digitalis has proved of service in the hands of Dr. FULLER, in cases in which the pain is intermittent. He gives ℥x-xx of the tincture three or four times a day.

Ergota is, according to PHILLIPS, (*Mat. Med.*, 1879,) most useful in visceral neuralgias, especially *gastralgia*.

Ferrum is useful in cases associated with anæmia; in others it fails.

Galbanum has been found of service in neuralgia dependent upon uterine derangement, given as follows:

136. R.	Pilulæ galbani compositæ,	gr. iij-vij	
	Pilulæ hydrargyri,	gr. iij.	M.
One pill to be taken every other night.			

Gelsemium is an important addition to the therapeutics of neuralgia, especially in non-congestive forms. According to Dr. E. HEROGUELLE, of Paris, (*Medical Examiner*, December, 1877,) it acts favorably in cases of dental neuralgia of the fifth pair, of the frontal, temporal, supra, and infra-orbital nerves, the brachial plexus, the intercostal and ilio-lumbar nerves. Sciatic neuralgia appears to resist, rather more than other neuralgias, the calming effects of the drug. Two to five drops of the fluid extract is the dose. According to Prof. MASSINI, the cases in which gelsemium produces most benefit are those of simple rheumatic neuralgia of the alveolar branches of the trigeminus; in those it rarely fails. It also sometimes relieves the pain remaining after the stopping of a carious tooth. Where there is any inflammatory affection of the bone or periosteum, no good can be expected from the remedy.

J. CHIRON uses the following formula:

137. R.	Tinct. Gelsemii,	m c
	Syrup. simplic.,	f. 3 x
	Aquæ destillat.,	q. s. ad f. 3 viij.

M. S.—From three to five tablespoonfuls daily, either half an hour before or three hours after the mid-day meal.

Glonoïn (nitro-glycerine) in very small doses (one drop diluted with 100 of rectified spirit), has been recommended by HAMILTON.

Guaiacum is of service in forms of neuralgia of a rheumatic character. From ℥xx to ℥xl of the ammoniated tincture may be given every five hours, until relief is obtained.

Hyoscyamus is considered by Dr. HARLEY as more efficacious than belladonna, in neuralgic affections of the internal viscera.

Hyoscyamin. In *Le Progres Medical*, M. VERNEUIL reports a case of obstinate neuralgia cured by this drug after all other means had failed.

Hyperosmic Acid. Prof. BILLROTH, of Vienna, has had success in some obstinate cases of sciatica, by the hypodermic injection of a one per cent. solution of hyperosmic acid. Within a day or two the pain was greatly relieved, and finally disappeared. (*Med. Compend.*, 1884.)

Ipecacuanha. Dover's powder at night often affords much comfort. An eighth or a quarter of a grain of morphia may be added when indicated.

* *Morphia*, hypodermically injected, affords speedy and often permanent relief. Dr. ANSTIE has cured cases by three or four injections of gr. $\frac{1}{8}$, which would have been very tedious under the old plan of treatment. Many cases yield in a week or ten days, to hypodermic injections twice daily.

Napelline. Dr. GROGNOT (*Bull. Gen. de Therap.*, 1884,) reports a case of trigeminal neuralgia, rebellious to a variety of treatment, that finally yielded to napelline in doses of $\frac{1}{800}$ of a grain every two hours. This dose seems like infinitesimalism, but it is that which is reported.

Oleum Cajuputi may be employed with advantage, either externally or internally, in cases unconnected with inflammatory action.

Oleum Terebinthinæ is not unfrequently of benefit. It is equally appropriate to the inflammatory and non-inflammatory states of neuralgia, and it is said relapses are less frequent after it than after most other remedies.

Oleum Tigllii has been used with success in some cases, in which it has seemed to exercise a special influence apart from its purgative action.

Opium, both internally and externally used, is of great service in certain cases; but ordinarily, far inferior to morphia injected subcutaneously.

Physostigma has exerted local anodyne effects in several cases of neuralgia reported.

Piscidia Erythrina. The Jamaica dogwood has been proved to be an excellent agent in many forms of neuralgia. It is a calmative and hypnotic, well suited for those who cannot tolerate opium. The credit of its introduction is due to the house of PARKE, DAVIS & Co., of Detroit. The dose is \mathfrak{z} ss-j. of the fluid extract. An efficient combination is:

138. R.	Potassii bromidi,	\mathfrak{z} ss
	Fld. extr. piscidiæ,	f. \mathfrak{z} ij
	Syrupi simplicis,	ad f. \mathfrak{z} iv.

A teaspoonful hourly till relieved.

**Plumbi Acetas* is said to be a safe and effective remedy in many cases of superficial neuralgia, and even in neuralgia of the abdominal viscera.

**Potassii Bromidum*, in full doses, often effectually relieves some forms of neuralgia.

Potassii Chloras, in doses of gr. xxx a day, has, in three weeks' time, it is reported, produced wonderful effects in facial neuralgia.

Potassii Iodidum is of service in neuralgic affections dependent upon an inflammatory state of the nerve coverings.

Pyrethrum, the root, chewed, will frequently relieve facial and dental neuralgia.

**Quiniæ Sulphas* is of great value in all neuralgias of malarial origin, and in a certain number of non-malarial cases. In the former, it should be given in full doses (gr. v-xx), shortly before the expected attack; in the latter, doses of gr. ij-ijj thrice daily, are as large as are likely to benefit. Dr. ANSTIE says that he knows of no circumstances which indicate beforehand that quinine will be useful in non-malarial cases, excepting that in neuralgia of the ophthalmic branches of the fifth, it is more effective than in other non-malarial neuralgias. Prof. GROSS frequently combines quinine with morphia, strychnia, arsenic, and aconite, in the treatment of neuralgia. Dr. TANNER combines it with belladonna and camphor, and with the chloride of arsenic.

139. R. Quiniæ sulphatis, ℞j
 Liquoris arsenici chloridi, f. ʒ iij-iv
 Acidi sulphurici aromatici, f. ʒ ij
 Syrupi zingiberis, ad f. ʒ iij. M.

One teaspoonful in two tablespoonfuls of water, directly after breakfast, dinner, and tea, in severe neuralgia.

Salicin has lately been spoken of highly in this disease.

Sodii Arsenias is of service in neuralgia where other arsenical preparations are badly borne.

Sodii Carbonas often, in full doses, speedily arrests neuralgia connected with acidity of the stomach.

Sodii Hypophosphis, in drachm doses, given in beef tea, thrice daily, has proved successful.

Stramonium is only found of benefit by Dr. FULLER where the symptoms shift from limb to limb, but never when the disease is stationary.

Strychnia is of use in epileptiform neuralgia and locally.

Valeriana exerts a favorable influence in neuralgia associated with hysteria. The ammoniated tincture may then be given, combined with guaiacum.

Zinci Valerianas is a serviceable remedy in those cases in which the disease is purely nervous, and in those which accompany uterine derangement.

Zinci Phosphas. Dr. CAMILLE TROUBERT (*Gazette des Hôpitaux and Courier Médicale*, 1878,) mentions a number of cases in which this treatment, first proposed by ASHBURTON THOMSON, was successful when other means failed. Two or three granules, of four milligrammes each, were given daily. It is important that the granules shall be used, as other preparations are uncertain.

* *Stimulants*. Dr. ANSTIE has found in some cases of acute neuralgia (particularly of the fifth nerve), that after the failure of various narcotics, a moderate dose of alcohol has at once given relief. He also states that, in certain chronic neuralgias of the aged, where the power of digesting ordinary food is almost suspended by reflex irritation, an almost exclusively alcoholic diet, continued for some time, occasionally works wonders.

EXTERNAL REMEDIES.

** *Aconitum and Aconitia*, locally applied over the seat of pain, is a most powerful palliative in neuralgic affections. Dr. FULLER, having observed that it fails so long as signs of active disease exist, restricts its use to purely chronic cases. Dr. DA COSTA sometimes prescribes it with veratria, in ointment. (F. 119.)

Acupuncture is of use after inflammation has subsided, in those cases in which there is effusion within the sheath of the nerve. It then promises speedy and effectual relief, but is useless, and may be prejudicial, in all other cases.

Æther. The local application of atomized ether is now often resorted to. In facial neuralgia, Dr. A. M. CARTLEDGE (*Med. Herald*, May, 1883,) pro-

teets the eye with a piece of oil silk, and directs a spray of ether upon the part affected until its temperature is down to the freezing point of water—say for eight minutes—which will generally suffice. To generate the spray he uses Richardson's atomizer, which is a simple addition to the ordinary perfume atomizer.

Aqua Ammonia. The following method of applying strong water of ammonia in neuralgia, is recommended by Prof. STILLÉ: Take small disks of cloth, about a half or three quarters of an inch in diameter, and apply them, moistened with the ammonia, over such points in the course of the affected nerve as are tender upon pressure, or are the seat of spontaneous pain. Cover the pieces of cloth with coins, buttons, or other convenient bodies, to prevent evaporation. Cotton contained in a thimble and saturated with the caustic liquid, forms a still more convenient arrangement. As soon as slight vesication occurs, the pain is relieved, yet not so thoroughly as when the same effect has been obtained by means of cantharides. The addition of morphia to the vesicating liquid renders its effect more prompt and certain.

Argenti Nitras, administered hypodermically, five or six drops of a strong solution (one to four) has been used with prompt results by M. LUTON. (*La France Médical*, June, 1877.) Intense pain is caused at first.

Belladonna is much used externally. TROUSSEAU highly recommends the external application of belladonna when the neuralgia is superficial. In such cases, may also be employed a compress saturated with a solution of atropia. Dr. FLINT uses atropia in preference to belladonna, in liniments and ointments. These applications, however, are all inferior to the hypodermic injection of atropia. Dr. A. WALLER, of Geneva, recommends that this and similar extracts and alkaloids should be dissolved in chloroform for external use, as they are more easily absorbed.

* *Cantharis*. Blisters should be tried in cases of any severity; if the hypodermic injection of morphia and atropia has failed, flying blisters should be employed. Dr. ANSTIE states that, in numerous early cases, one or two flying blisters, applied successively over the different points in the course of the painful nerve, have at once and permanently arrested the disease.

Chloroformum may be applied pure (especially serviceable when the pain is localized in a small space) or in liniments. (F. 133.) "Gelatinized chloroform," made by incorporating one part of chloroform and two parts of white of egg, is a useful form for application.

Electricity seems to be of service only in cases unconnected with irritation in distant parts of the body; then the continuous galvanic current is often of essential service. If employed before the pain has subsided, the interrupted current very generally aggravates the patient's sufferings. (See HAMMOND'S remarks, p. 105.)

Franklinization. Dr. V. J. DROSDERFF (*Med. Record*, 1883,) has employed franklinic electricity in neuralgia or rheumatism. Immediately after a sitting, pain and tenderness disappear, and, during the remaining part

of the day, the patients feel much better than before a sitting. Each successive franklinization brings a further decrease of intensity of neuralgia. In cases of short standing, three or four sittings suffice to completely remove the agonizing pains. Ten to fifteen sittings cured even cases of neuralgia of twelve years' standing, which had obstinately resisted galvanization, faradization, and all possible therapeutic means. In each of the author's cases, either complete cure or very considerable improvement followed. There is no necessity to undress a franklinized patient, as the therapeutic effects of this excellent agent remain unchanged—a circumstance of practical importance in female cases.

Frigus. Ice applied to the spine, by means of his spinal bag, is recommended by Dr. CHAPMAN.

Hydrargyri Iodidum Viride. The following ointment has been strongly recommended :

140.	R.	Hydrargyri iodidi viridis,	℞ij	
		Adipis,	℥j.	M.

****Hypodermic Injections.** Dr. ANSTIE says that the discovery of the hypodermic method has instituted a new era in the treatment of severe neuralgias: Atropia, or morphia or the two combined, may be used. The face should be avoided in making the injection. Prof. BARTHOLOW says. "The neuralgias are best treated by the combined morphia and atropia solution. There are several reasons for this: much larger doses of morphia may in this way be injected without danger to the patient; and the larger the quantity, as BROWN-SÉQUARD has shown, the greater the curative power. Morphia and atropia combined are more effective than either singly. The systemic effects during the time of maximum narcosis, and also after the narcosis has disappeared, are much less unpleasant and depressing when the two agents are combined than when morphia is used alone. Sometimes atropia is better borne than morphia, and *vice versa*: in this case, the agent whose effects are least unpleasant should be in excess." He recommends the following proportions: $\frac{1}{48}$ to $\frac{1}{120}$ of a grain of atropia, $\frac{1}{4}$ to $\frac{1}{2}$ of a grain of morphia. The physiological effects of atropia here predominate, but the toxic effects are guarded by the morphia. For the facial neuralgia of pregnancy, he uses morphia alone. This treatment promptly cures the affection, a fact first pointed out by Dr. H. R. STORER, of Boston. "These cases, as is well known, are extremely obstinate under the old methods of treatment, and those who have suffered from them on former occasions are exceedingly grateful for the relief so promptly and permanently afforded by the hypodermic method."

Iodoform has been recommended for local use in neuralgias, in the form of saturated solution in chloroform or ether.

* *Leeches* applied along the course of the nerve often afford immediate relief after the failure of anodynes and other local measures. They are particularly indicated when there is inflammatory action or much vascular excitement.

Mentha and Monarda. Oil of horsemint, in liniment, with camphor and laudanum, sometimes affords prompt relief. Oil of peppermint is at times very prompt and efficient. An alcoholic solution is used.

Menthol, or Peppermint Camphor, has been recently introduced for similar purposes. One or other of these substances is included in most popular "pain-killers."

Neuroteny, (nerve stretching,) *Neurotomy*, (division of the nerve,) and *Neurectomy*, (exsection of a piece of the nerve,) are surgical resources in very obstinate cases.

Rectal Suppositories of belladonna, hyoscyamus and opium, are sometimes employed.

Stramonium is sometimes employed locally in the same manner as belladonna, to which, however, it is inferior.

* *Tuning-Fork.* Dr. RASORI (*Cinn. Lan. and Clin.* 1884) applies the tuning-fork, while vibrating, over the course of the painful nerve. The sitting usually lasts about half an hour.

Veratria, when there are no active or inflammatory symptoms present, is often beneficial, externally applied, the ointment being rubbed in diligently until it causes a sense of heat and tingling. It is inferior, however, to aconitia.

NEURASTHENIA.

DR. ROBERTS BARTHOLOW.

The division of neurasthenic cases into two main types, the *congestive* and the *anemic*, is advocated by this writer (*Med. and Surg. Reporter*, 1884). The latter are greatly more numerous. The treatment of both varieties is chiefly symptomatic and hygienic. In congestive cases the remedies consist in a careful regulation of the diet, in baths, exercise, in a reduction of the hours devoted to work, but not the cessation of work; in the use of a laxative quantity of sodium phosphate daily, and in the administration of the aqueous extract of ergot, with the chloride of gold and sodium, and a minute quantity of bichloride of mercury.

It is of course of the greatest importance that the habits of excess, of work and of worry, which have given rise to the diseased condition, be checked. This is often one of the most difficult points of the management.

DR. S. WEIR MITCHELL.

In his work entitled "Fat and Blood" (3d ed., 1884), this author

gives directions for a treatment of certain forms of Neurasthenia by a combination of entire rest and excessive feeding, made possible by passive exercise obtained through the steady use of massage and electricity. Most of the cases—though not all—to whom this method is applicable, are nervous women with little fat and a lack of blood. Success by this method can only be obtained by the punctilious observance of a great number of minute details of treatment, and under circumstances and surroundings which are usually out of the power of the ordinary practitioner to command.

NOTES ON REMEDIES.

Cannabinum Tannicum, or tannate of cannabine, in doses of gr. $\frac{1}{4}$ -j, is stated by Dr. EICHHOLT (*Deutsche Med. Woch.*, 1883.) to be especially useful in neurasthenic insomnia. It does not derange the digestion, and soothes effectively.

Napelline, a new product from aconite, is stated by Dr. LABORDE (*Jour. de Therap.*, 1883,) to allay the excessive excitability of nervous exhaustion with excellent effect. He employs it in subcutaneous injections of from one to four centigrammes.

Strychnia. In exhausted nerve function Dr. J. C. THOROWGOOD (*Med. Times*, 1883), has obtained excellent results from small doses of strychnia, long continued, as gr. $\frac{1}{30}$, two or three times a day. He thinks larger doses rather increase the nervous irritation.

Fattening Food. Dr. CLOUSTON (*Lectures*, 1883,) advises the free use of fattening foods in nervous diseases. He writes: "All acute mental diseases, like most nervous diseases, tend to thinness of body, and therefore all foods and all medicines and all treatments that fatten are good. To my assistants and nurses and patients, I preach the gospel of fatness as the great antidote to the exhausting tendencies of the disease we have to treat; and it would be well if all people of nervous constitution would obey this gospel."

PARALYSIS.

The forms of paralysis are numerous, and treatment must always be guided by the character of the changes in the cord, or other pathology of the disease.

In *general paralysis* a plausible plan of treatment has been suggested (*Berliner Klinische Wochenschrift*, No. 21, 1877,) by

L. MEYER, M. D., BERLIN.

On the theory that it is a chronic encephalitis, he applies over the great fontanelle, on a space as large as a dollar, some antimonial ointment. He repeats this operation at the end of twenty-four hours, using an amount of ointment the size of a pea, and a little piece of linen. The second friction is managed with care, so as to save the skin. Generally these two frictions are sufficient; if not, a third; a small quantity of the ointment is applied without friction, and a piece of linen to which it is applied may be left on the surface.

In the course of three or four days the tumefaction of the scalp extends to the forehead and even to the face, more rarely to the occiput or the nucha. The pustular eruption is insignificant; the pustules are small, concrete and dry rapidly.

After the swelling has become intense, it may be covered with warm poultices. In a few days the suppuration is so ample that it completely separates the integuments, which slough off spontaneously. Too resistant adherences are destroyed with caustics or the scissors. To reach this point usually fifteen days are required. From that time the suppuration is treated for two or three months with basilicon ointment. The patient may take during the same time moderate doses of iodide of potassium, and should have a substantial diet. Furthermore, he should be in the open air as much as possible, employed in gardening, etc., only taking care to avoid exposure to extreme heat. Baths should be forbidden, as conducing to cerebral congestion.

During fifteen years this treatment has been employed by MEYER, in the cases of seventeen general paralytics, two of whom remain under treatment. Eight of the fifteen were cured, and their cases are given in his memoir. One of them, however, suffered a relapse after two years.

The periosteum may be affected and there may occur exfoliations, more or less considerable, of the bone. The author met with this accident a number of times, but attaches to it no special importance. The loss of hair is readily concealed.

STRYCHNIA AND ITS ADMINISTRATION.

Of the numerous drugs which have been used in paralysis, none equals *strychnia* in efficaciousness.

Dr. F. W. HEADLAND, of London, states that it is only successful in

paralysis in cases where the injury to the nervous centre has healed up, and when the limb continues paralyzed merely because the motor nerves have lost the power of transmitting the necessary impulse, from having been so long unaccustomed to discharge this office. On the other hand, when the lesion of the nervous centre is of recent occurrence, or when it has been of so serious or extensive a nature as to admit of no repair in the course of time, the remedy will be ineffectual.

Sir THOMAS WATSON lays down the general rule that strychnia can be of benefit in these cases only when the cord is free from disease.

Prof. WILLIAM A. HAMMOND has employed it very successfully by a new and improved method, in treating *facial paralysis*. (*St. Louis Clinical Record*, May, 1878.) The improvement consists in the administration of strychnia in increasing doses to the point of rapidly—as rapidly, in fact, as is consistent with prudence—bringing the patient under its full physiological influence.

For this purpose he makes use of a solution of the sulphate of strychnia in the proportion of one grain to the ounce of water.

Every ten minims of such a solution contain $\frac{1}{8}$ of a grain of the medicine. Generally, he begins with ten minims of this solution three times a day for the first day; the next day eleven minims are given three times; the next twelve, and so on till the patient experiences a sensation of cramp or rigidity in the legs, or in muscles of the back of the jaw. Usually the cramp is first felt in the calves of the legs. The further administration is now stopped, and, if necessary, on the following day the solution is given as before, in doses of ten minims, and the doses are again run up to the extent of producing the muscular cramp. Of course, with this, electricity, passive exercise, and support of the affected side of the face by a little hook placed in the angle of the mouth, and attached to the ear by an elastic band, are advantageously combined.

ELECTRICITY IN PARALYSIS.

DR. G. VIVIAN POORE, OF LONDON,

lays down the following rules as to its application to paralysis: 1. *If the paralysis to the will remain absolute, and if the contractility of the muscles be perfect, we do no good by persevering with electrical treatment.* This condition is often met with in hemiplegia. The patient is absolutely helpless on one side, although the muscles are in

no degree wasted, and their irritability remains normal. 2. *If the paralysis to the will remain absolute, and if the irritability of the muscles be diminished, then electricity is useful, in so far as it helps to improve the nutrition of the muscles, and restore their normal degree of irritability.* The normal degree of irritability being restored, (the paralysis to the will remaining absolute,) then electrical treatment may be discontinued. The irritability of muscles may be diminished from mere want of use, and electricity may serve to restore this irritability. 3. We should bear in mind the advice of Sir THOMAS WATSON, that "our aim should be to preserve the muscular part of the locomotive apparatus in a state of health and readiness, until, peradventure, that portion of the brain from which volition proceeds having recovered its functions, or the road by which its messages travel having been repaired, the influence of the will shall again reach and re-animate the palsied limbs." 4. Whenever we meet with the *degenerative reactions*, we should employ that form of current to which the muscles most readily respond, which is invariably the galvanic. We must persevere doggedly, and we may find our perseverance rewarded by the gradual return of the normal muscular irritability. 5. If the irritability to both forms of current has completely disappeared, we are not justified in persevering too long, nor in holding out delusive hopes to the patient. Nevertheless, treatment should not be abandoned without a patient trial. In treating paralysis with electricity, an exact diagnosis is, above all things, necessary, and we should be able to bring our knowledge to decide as to whether or not the paralyzing lesion is of a permanent nature, or one that, according to our pathological knowledge, is or is not removable or susceptible of improvement. If not, then our efforts to cure must of necessity be unavailing. It is in those cases where the paralyzing lesion has disappeared, or has improved, and we have only to combat with the effects produced by it, that our efforts are the most successful. *Hysterical paralysis* is especially susceptible to electrical treatment, and with it we occasionally work apparent miracles. When a patient, who has no will to move her muscles, sees that her muscles can be made to move in spite of her will, it is no wonder that such a discovery should exercise a wholesome moral influence over her mental condition.

Other authors state that it should never be resorted to in recent cases, if there be reason to suppose that effused blood or coagulum remains

unabsorbed, nor so long as there are any febrile or inflammatory symptoms present. It is of no benefit in cases with organic lesion of the nervous system. Its greatest utility lies in cases of purely *local* paralysis. Great and terrifying shocks are rarely beneficial where small and repeated shocks have failed. Both faradization and galvanism have been employed with success. Electro-puncture has also its advocates.

NOTES ON REMEDIES.

Ammonii Formias is recommended by Dr. RAMSKILL as a stimulant in cases of chronic paralysis accompanied by general torpor.

Argenti Nitras, in doses of gr. $\frac{2}{5}$ -j daily, has been employed with success by BOUCHUT in paraplegia and progressive general paralysis, resulting from sexual or alcoholic excess.

Arseniosum Acidum in paralysis from cerebro-spinal sclerosis, is well spoken of by FLINT.

Cantharis. Sir THOMAS WATSON recommends tincture cantharides in paraplegia. Generally, when it does good, it acts as a diuretic.

Cocculus Indicus. According to PHILLIPS, this is a very valuable remedy in partial palsies accompanied by giddiness, as also in hysterical, choreic and epileptic hemiplegia. The dose of the tincture (\mathfrak{z} j to \mathfrak{Z} j) is gtt. ij-x; of *picROTOXIN*, the active principle of the berry, gr. $\frac{1}{12}$ - $\frac{1}{60}$.

Ergota is recommended in those forms of paralysis arising from spinal congestion, in paraplegia complicated with menstrual irregularity, (MEADOWS,) and in the constipation of the paralytic, after the most powerful cathartics have failed. (WARING CURRAN.)

Ferri et Strychniæ Citras is serviceable in paralysis associated with anæmia.

Hydrargyrum. Dr. MAUDSLEY says: "To administer mercury systematically in general paralysis, as has been done, is as unaccountable in theory as it undoubtedly is pernicious in practice."

Oleum Tiglii has been very highly commended in some obscure cases—drops 1-2 every two hours until the bowels are acted upon, and then an opiate to relieve the griping. (J. W. DAVIS, *Nashville Jour. of Med. and Surg.*, May, 1878.)

Phosphorus has been successfully employed in some cases in the form of the oil, (made by gently warming two parts of phosphorus in 100 parts of almond oil, in a closed flask, for about twenty minutes, and decanting off, when cool, the oil from the undissolved phosphorus. Dose, gtt. iij-vij, in emulsion, or with cod-liver oil.)

Picrotoxin. See *Cocculus Indicus*.

Potassii Iodidum proves successful in syphilitic paralysis. It should be given in doses of \mathfrak{z} j and more, combined with sublimate, gr. $\frac{1}{20}$.

**Strychnia* and *Nux Vomica*. (See p. 116.)

Sulphur gives very good results, according to Dr. GRAVES, in paralysis of an

asthenic or chronic character, when employed as a sequel to a course of strychnia. He uses it in the form of an electuary and in baths.

Toxicodendron is recommended by TROUSSEAU in paraplegia from concussion of the spinal marrow, administered as follows : gr. v the first day of treatment, increased daily in the same amount until gr. lx a day are reached. It is asserted that it strengthens rather than enfeebles digestion, and causes no inconvenience other than occasionally slight strangury.

EXTERNAL REMEDIES.

Armoraciæ Radix. Poultices of fresh horse-radish root form a ready counter-irritant in paralytical affections.

Cajuputi Oleum makes an excellent embrocation for paralytic limbs. Diligently rubbed in, it stimulates the parts and relieves pain when present.

Delphina is believed by Dr. TURNBULL to be more useful in paralysis than veratria, from its property of exciting the circulation in the affected part.

Mucuna. The setæ maintained on the affected limb by a bandage have been employed as a local stimulant in paralysis.

Myristica. The oil, diluted with soap liniment, forms a useful stimulating embrocation.

Sulphuricum Acidum. The following ointment, perseveringly employed, has resulted beneficially :

141. R.	Acidi sulphurici,	f. ʒj	
	Adipis,	ʒj.	M.
It acts as a powerful irritant.			

Veratria ointment, diligently used, has occasionally been followed by great improvement.

Cold Shower Baths, in cases of paralysis consequent on severe inflammatory attacks of the brain and spinal column, are often attended with benefit, but sometimes fail, and may even aggravate the symptoms.

Cantharis. Blisters to the spine are useful when there is reason to expect effusion within the theca.

Electricity. (See above, p. 117.)

Issues and Setons are sometimes employed with advantage.

Leeches at the verge of the anus are beneficial where there is suppression of an habitual discharge, as from piles.

Metaliotherapy. It has been positively shown in the Parisian hospitals that the application of metal rings and plaques to the paralyzed limbs restores sensation and motion, under certain circumstances, and this when the elements of both faith and fraud are carefully eliminated.

PROGRESSIVE LOCOMOTOR ATAXIA (POSTERIOR SPINAL SCLEROSIS.)

ALLEN M. HAMILTON, M. D., NEW YORK.

The treatment is entirely empirical.

In the early stages ergot and bromide of sodium control the pain.

142. R. Extr. ergotæ fl.,	f. $\frac{3}{4}$ ij	
Sodii bromidi,	f. $\frac{3}{4}$ iss	
Aquæ camphoræ.	ad f. $\frac{3}{4}$ iv.	M.

A teaspoonful every four hours.

A combination of belladonna and turpentine, as suggested by TROUSSEAU, not only relieves the pain, but helps the vesical trouble, if present.

143. R. Extr. belladonnæ,	gr. iv	
Olei terebinth.,	f. $\frac{3}{4}$ ij	
Butyri cocoæ,	q. s.	M.

Make twelve capsules. One thrice daily.

No one remedy has had so much said in its favor as nitrate of silver. It is well to begin with a one-fourth-grain dose, and increase to gr. $\frac{1}{2}$ or j thrice daily. It may be combined.

144. R. Argenti nitratis,	gr. vi-viiij	
Extr. nucis vomicæ,	gr. xij.	M.

Make twenty-four pills. One after each meal.

Or:

145. R. Argenti nitratis,		
Extr. belladonnæ,	āā	gr. vi-viiij
Extr. gentianæ.		q. s.

Make twenty-four pills. One after each meal.

A salt foot-bath every night, is beneficial, and the sulphur-bath has worked wonders in cases of long standing, and deserves a faithful trial. A small lump of sulphide of potassium is to be thrown into the water in which the patient bathes, and he is to be well rubbed down afterwards.

DR. EDWARD C. MANN.

This author (*Psychological Medicine*, 1884,) is of opinion that some cases of this disease are curable, especially if seen in their early

147. R. Potassii iodid., \mathfrak{z} vi-viiij
 Ferri am. citrat, \mathfrak{z} ij
 Syr. simp.,
 Tr. aurant. cort., $\mathfrak{a}\mathfrak{a}$ \mathfrak{z} iij
 Aqua menth. pip., q. s. ad f. \mathfrak{z} iv. M.
 Ft. sol. et. Sig.—A teaspoonful in water about an hour after each meal.

PROF. TROUSSEAU, PARIS.

This author advises not to employ setons, moxas, etc., along the spine, as they cause great pain and irritation, and do no possible good. *Electricity* is the great remedy which he recommends, combined with *flagellations* prudently employed, with *belladonna* to abate pain, with *sulphurous baths*, which have been very successful. *Hydrotherapy* may be of great service in the constitutional treatment, as well as other appropriate remedies of a like action. Preparations of *mercury* and *iodide of potassium* are of great good where there exists a syphilitic taint. TROUSSEAU recommends his "compresseur" of the prostate in the spermatorrhœa which sometimes attends the outset of the disease. *Turpentine* is useful in the vesical manifestations of the malady. Hygienic rules are to be strictly observed.

NOTES ON REMEDIES.

**Argenti Nitras*, according to Dr. ALTHAUS, is useful in a large proportion of cases. It should be given in doses of gr. $\frac{1}{16}$ — $\frac{1}{2}$, two or three times daily. Its combination with hypophosphite of soda seems to add to its efficacy. After its use has been continued for four or six weeks consecutively, its administration should be interrupted for two or three weeks, and a slightly aperient mineral water given. The nitrate may then be resumed for a month or two. The gums should be carefully watched, and, on the appearance of the slightest blue coloration, the remedy should be discontinued. Nitrate of silver has been thus employed in ataxia with variable success.

Argenti Oxidum. In the *Med. and Surg. Reporter*, Dec. 18th, 1875, Dr. CHARLES CARTER, of Philadelphia, reports two cases in which the oxide of silver, gr. $\frac{1}{2}$, t. d., increased to gr. ij, t. d., controlled the neuralgia and apparently arrested the progress of the disease. This drug, also, does not discolor the skin.

Argenti Phosphas, gr. $\frac{1}{8}$, is preferred to the nitrate by Dr. A. M. HAMILTON.

Belladonna and *Cannabis Indica* are both said, by Dr. AITKIN, to give great relief to the pains.

Nitro-muriaticum Acidum is an excellent tonic in these cases.

Phosphorus is recommended by Dr. AITKIN and Dr. A. FLINT. The administration of the phosphates of the metals and salts may be conjoined with the use of dilute phosphoric acid as a drink, in the daily allowance of water, and with phosphate of soda as an aperient.

- | | | | |
|------|---|----------|----|
| 150. | R. Salicin, | gr. xij. | |
| | This amount every six hours, in acute rheumatic sciatica. | | |
| 151. | R. Tinct. colchici, | ℥x | |
| | Pulv. opii, | gr. j. | M. |
| 152. | R. Tinct. aconit. rad., | ℥x | |
| | Potassii iodidi, | gr. x. | M. |
| | Thrice daily in rheumatic and gouty forms. | | |
| 153. | R. Quiniæ sulphatis, | gr. ij-v | |
| | Tinct. nucis vomicæ, | ℥v. | M. |

In pure neuralgic sciatica:

- | | | |
|------|---------------------------|----------|
| 154. | R. Tinct. actææ racemosæ, | f. ʒj-ij |
| | Potassii bromidi, | ʒj. |
| | This amount thrice daily. | |

DR. HENRY LAWSON, ENGLAND.

Out of some eighty cases of sciatica, lumbago, and brachialgia, treated by this physician, some of them excessively bad cases, in not more than three did the *subcutaneous injection of morphia* fail to give relief, in most cases complete. By means of this relief, the patient was enabled to eat and drink with comfort, and by help of perchloride of iron, and in some cases by cod-liver oil, he was enabled to put on flesh rapidly, and to repair the waste of tissue under which he had been laboring. The salt he uses is the muriate. Five grains are dissolved, by the aid of heat, in one drachm of distilled water; it must be injected warm, as the solution solidifies on cooling. The place of injection should be chosen as near as possible to the seat of pain, and the point of the needle should not be more than half an inch long. Dr. LAWSON does not object to the puncture of a minute vein, as thereby the medicine is more rapidly absorbed, while the alarming symptoms which ensue are generally very transitory. The patient should have eaten a hearty meal about half an hour previous to the injection; this prevents, to some degree, the soporific effect of the drug, while it heightens its effect in allaying pain. (*Medical Times and Gazette*, January, 1877.)

C. G. COMEGYS, M. D., CINCINNATI.

Following MALGAIGNE, this physician has frequently succeeded in promptly curing refractory sciaticas by touching, with a red-hot wire, a

certain spot on the ear, on the same side as the one affected, to wit, the anterior part of the helix, just at its entrance into the concha.

Another very efficient remedy is to inject fifteen to twenty drops of common ether (in the ordinary superficial manner) behind the great trochanter. The pain is sharp, but not lasting, and the relief often astonishingly prompt. It should be repeated daily, if required. (*Cincinnati Lancet and Clinic*, August, 1878.)

PROF. J. M. DA COSTA, M. D., PHILADELPHIA.

155. R. Emplastri epispastici, 1½ in. x 5 in.
To be applied over the affected part. Let it draw for five or six hours, poultice it, and then remove the cuticle and dress with:
156. R. Morphiae sulphatis, gr. ¼
Pulveris marantæ, gr. ij. M.
For one powder. Also, ten grains of Dover's powder, to be taken at night.

DR. LABORDE, FRANCE.

157. R. Olei terebinthinæ, f. 3 viij
Chloroformi, āā
Tincturæ opii, f 3 ij. M.
A useful liniment in sciatica. To be rubbed on by means of a piece of flannel, several times a day.
158. R. Aquæ ammoniæ, f. 3 v
Olei terebinthinæ, āā
Olei amygdalæ dulcis, f. 3 j. M.
Rub on three times a day, in obstinate sciatica.

PROF. FELIX VON NIEMEYER, M. D., TUBINGEN.

159. R. Olei terebinthinæ, f. 3 j
Mellis, f. 3 j.
For an electuary. A tablespoonful twice daily.

VOGT'S METHOD OF SUBCUTANEOUS NEUROTENSION.

In the *Centralzeitung*, 1883, Dr. VOGT describes a method of nerve-stretching without a cutting operation. This is by forcible and extreme flexion of the limb. In sciatica the leg is extended at the knee, and forcibly flexed at the hip-joint upon the body, while the healthy leg, the pelvis and the body, are retained in a fixed position. By manœuvres of this kind, properly adjusted and repeated, the nerve of the suffering part is firmly extended, and a similar result obtained as when it is lifted from its sheath and stretched. Very favorable results are reported.

ROMBERG also speaks very highly of this as a specific remedy in the treatment of sciatica.

H. V. SWERINGEN, M. D., OF INDIANA.

160. R. Morph. sulph, grs. iij
Ft. ch. No. ix et Sig.—Inject one hypodermically night and morning, over the seat of pain; let one be taken per orem in the interim if necessary.

161. R. Quin. sulph, $\overline{3}^{ss}-\overline{3}^j$
Tr. ferri chlor, f. $\overline{3}^{ij}$
Spts. æth. nit., f. $\overline{3}^{iv}$
Syr. simp., f. $\overline{3}^{ij}$
Vin. Xerici, q. s. ad f. $\overline{3}^{iv}$. M.
Ft. sol et Sig —A tablespoonful every three or four hours.

EDWARD WAKES, M. D., LONDON.

162. R. Extracti ergotæ fluidi, f. $\overline{3}^{ij}$
Aquæ cinnamomi, f. $\overline{3}^{ij}$.
A dessertspoonful in water every four hours.

Tinctura ferri chloridi may be added if indicated. When ergot is likely to be useful, its good effects commence immediately.

PROF. M. ROSENTHAL, M. D., OF VIENNA.

This author advises wet cups or leeches in very painful, acute cases, followed by flying blisters to the painful points, and the inunction of opiated ointments. When turpentine can be given internally, without causing digestive troubles, it should be administered thus, but otherwise only by frictions. Electricity is of little value in sciatica of central origin, but when from peripheral irritation, the secondary current should be applied to the painful points for five to ten minutes, by means of dry electrodes, using weak currents at first and gradually increasing their intensity. A hydrotherapeutic plan, applicable to all cases, and often successful, is the use of moist packs, followed by cool affusions, or half baths at about 65° Fah., with friction to the limbs. Hot vapor baths, ice-bags, cold douches and sweats, advocated in many hydrotherapeutic establishments, should be confined to robust individuals, and to the peripheral forms of sciatica, in which their success is often gratifying.

NOTES ON REMEDIES.

Aconitum is, in the opinion of Dr. FULLER, not only a valuable and efficient remedy in many obstinate forms of this disease, but is also safe and manageable, when properly administered and carefully watched.

Ammonii Chloridum is recommended by Dr. ANSTIE in the milder cases of sciatica occurring in the young.

**Atropia*, hypodermically, is a successful form of treatment now much employed.

**Belladonna* is one of the most valuable sedatives and anodynes we possess in this affection.

**Cannabis Indica* ranks next in value to belladonna and opium.

Conium is sometimes of service, and may be resorted to when belladonna and opium fail or are contra-indicated.

Creasotum has been employed internally by Dr. ELLIOTSON, with apparent benefit.

Digitalis is recommended by Dr. FULLER as a reliable remedy, when the pain is intermittent and of a purely neuralgic character. He gives either the powdered leaves (gr. j-iss,) or the tincture, (℥x-xx) thrice daily.

**Morphia*, given hypodermically, affords speedy and often permanent relief. It may also be employed endermically. (F. 156.)

Nitro-muriaticum Acidum, in full doses (℥vj-x,) is an effectual remedy in sciatica accompanied with malaria. Its action should be aided by an occasional brisk purgative, the cold shower bath, or a cold douche down the spine, followed by friction, and change of air and scene.

Oleum Tigllii has been employed with success, internally.

Opium, although of great occasional service, is less used since the introduction of the more effective treatment by the hypodermic injection of morphia.

Potassii Iodidum is strongly advised by Dr. GRAVES, in doses of ʒj in the course of the day, dissolved in a decoction of sarsaparilla, in cases of a subacute or chronic character, and where the patient has to follow his out-door avocations.

Strychnia is inadmissible in acute or inflammatory cases, but in others, nuxvomica has been used with occasional success.

**Terebinthinae Oleum* is considered by Dr. ANSTIE as a remedy worthy at least a trial, in doses of ℥x thrice daily, in the more obstinate cases of sciatica. It is, however, usually very disagreeable to the patient. Dr. NIEMEYER also recommends it in electuary, and Dr. ROMBERG speaks of it almost as a specific. Dr. W. A. JAMIESON (*Edinburgh Med. Jour.*, March, 1877,) gives the following formula :

163. R.	Olei terebinth.,	f. ʒij
	Olei ricini,	f. ʒiv
	Tinct. cardam. comp.,	f. ʒj
	Mucilaginis,	
	Aquam,	aa q. s. ad f. ʒij.

Take at one dose.

He adds, the patients who derive most benefit from oil of turpentine in sciatica present the following appearances: Usually, middle-aged;

they look older than they ought, and have hair, as ANSTIE so well phrases it, *prematurely and permanently gray*. The complexion may be pale or fresh, but it nearly always exhibits a peculiar *ashy gray* or *stone-colored* aspect.

EXTERNAL REMEDIES.

The Actual Cautery. Prof. MICHEL PETER, of Paris, (*Lancet*, 1878,) prefers the cautery to any other means. He recommended a number of superficial cauterizations by an olive-headed cautery along the course of the sciatic nerve and its divisions, from the trochanteric region to the outer malleolus. M. PETER considers this treatment preferable to blistering, because of being enabled by it to follow the whole course of the nerve, whilst it does not produce suppuration, or lead to any vesical trouble. It may also be repeated, if necessary, with impunity. (See, also, p. 125.)

Aconite and Aconitia, in liniment or ointment, are powerful local sedatives.

Acupuncture is limited to purely chronic cases, and to those in which there is effusion within the sheath of the nerve. In appropriate cases, it promises speedy and effectual relief.

Atropia has been strongly recommended by Dr. ROBERTS BARTHOLOW. He says: "The principal triumphs of atropia over neuralgia have been in cases of sciatica. It is now admitted that atropia is one of the best remedies for this disease." It must be injected at the seat of pain. In severe cases, gr. $\frac{1}{25}$ of sulphate of atropia may be used; but this quantity will excite very severe symptoms in susceptible subjects. Generally, gr. $\frac{1}{48}$ will produce decided atropinism. Cessation of the pain is not immediate upon the systemic effects; indeed, the pain is often at first increased; but improvement takes place after a variable interval, and is often more permanent than after the morphia injection. When atropia fails, *Morphia* may be tried, or both may be employed together.

Cantharis. Great and immediate relief sometimes results from applying a blister to the heel. Flying blisters along the nerve are approved by many.

Chloroformum is advised externally by Dr. FULLER, who says that a strip of flannel, soaked in chloroform, placed along the course of the nerve, and covered with oiled silk, to prevent evaporation, proves sometimes extremely serviceable.

Dry Cups along the nerve are frequently beneficial.

Electricity in the form of galvanization of the affected nerve, does great good in most cases.

Extension. Dr. JOHN R. BUCHAN reports in the *Weekly Medical Review*, June 23, 1883, five cases of obstinate sciatica, eventually relieved by extension, and only one case in which it failed. About eight pounds was the weight used. This method of treatment is rational, and in all cases deserves a fair trial.

Pix Burgundica. Enveloping the whole thigh in a Burgundy pitch plaster has, according to TROUSSEAU, cured sciatica after the failure of blisters and the endermic use of morphia.

Sulphur, externally, is highly praised by Dr. FULLER, who finds it very generally efficacious and worthy of a trial in every case, when all active symptoms have been subdued or are absent. He sprinkles a piece of new flannel, thickly, with precipitated sulphur, encases the whole limb in this, from the foot upward, and covers it with oiled silk or gutta-percha, which has the effect not only of increasing the warmth and confining the vapor of the sulphur, but of obviating the odor, annoying to the patient. It should be kept applied night and day. Absorption takes place rapidly, and the breath, urine, stools, and perspiration, unmistakably attest its presence in the system

SPINAL IRRITATION.

PROF. WILLIAM A. HAMMOND, M. D., NEW YORK.

This eminent neurologist believes that that important group of symptoms which, since the essay of Dr. C. BROWN, of Glasgow (1828), have been subsumed under the term "spinal irritation," are due to anæmia of the posterior column of the spinal cord. The principles of its treatment are four:

1. To remove any manifest cause of irritation.
2. To improve the general tone of the system.
3. To improve the special nutrition of the cord.
4. To set up counter-irritation in the vicinity of the disordered region of the cord.

In regard to improving the general system, Dr. HAMMOND recommends distilled liquors, the oxide of zinc, and cod-liver oil. The third indication is easily filled by strychnia, phosphorus, and opium. Thus:

164. R.	Strychniæ sulphatis,	gr. $\frac{1}{32}$	
	Acidi phosphorici diluti,	f. 3 ss	
	Syr. aurant. cort.,		
	Aquæ,	ââ	f. 3 ij. M.

This amount several times a day.

Or a pill of the phosphide of zinc (gr. $\frac{1}{15}$) and nux vomica (gr. $\frac{1}{2}$.) Where there is contraction of the limbs, *opium* is very beneficial, as by hypodermic injections of morphia or in the form of suppositories, gr. $\frac{1}{2}$ of the aqueous extract with sufficient cacao butter. Water, as hot as it can be borne, applied to the spine by one of Chapman's bags, is an admirable adjuvant. But the most efficacious means to summon blood

to the cord, is the direct *galvanic current*. Either pole may be placed above and the other below the seat of pain. The current should not be passed at a *séance* for more than fifteen minutes, and no one application should last longer than three or four minutes. For the relief of the spinal tenderness, one pole should be applied directly to the painful part, and the other to a point distant laterally from it a few inches.

The fourth indication is very important. In slight cases, counter-irritation alone will effect a cure. Dr. HAMMOND prefers *blisters*, applied immediately over the painful part of the spine. *Tartar emetic ointment* is useful, but painful. *Dry cups* almost always do good. Leeches and blood-letting are always prejudicial. When patients feel more comfortable lying down, they should be permitted to do so, but, at the same time, they should be encouraged to take regular gentle exercise daily.

Special attention should be given to the stomach when there is persistent vomiting, and everything is rejected. The diet must then be of the simplest possible character, and only very small quantities taken at a time. The smallest quantity that can be kept down, is better than a larger quantity thrown up. Warm liquids are often more apt to be retained than cold ones. Among other medicines, the best is *valerianate of caffeine*, in doses of from three to five grains, as often as it may be required. Its only objection is its great cost. Cerium is inefficacious in the vomiting of spinal irritation.

DR. NATHAN S. DAVIS, OF CHICAGO.

Regarding spinal irritation as simple chronic inflammation, involving the roots of the nerves of sensation, or motion, or both, what is the appropriate treatment? Dr. DAVIS answers, that internally, an alterative plan like the following:

165. R.	Tincturæ cimicifugæ,	f. $\frac{3}{4}$ ij	
	Tincturæ stramonii,	f. $\frac{3}{4}$ ss	
	Potassii iodidi,	$\frac{3}{4}$ ij ss	
	Syrupi simplicis,	f. $\frac{3}{4}$ jss.	M.
A teaspoonful three or four times a day.			

166. R.	Hydrargyri chloridi mitis,	gr. j
	Potassii nitratis,	
	Pulv. ipecac. et opii,	āā gr. viij.

This amount, in a powder, three times a day until the mercurial is perceptible in the breath or gums, when it should be discontinued.

Externally, dry cupping along the spine, (for which purpose common tumblers are well adapted,) followed by belladonna plasters, or hypodermic injections of atropine, are required. These means, with rest in the horizontal position, will be likely to remove the disease in from four to six weeks.

ALLAN M. HAMILTON, M. D., NEW YORK.

In three-quarters of the female cases of this disease, it depends upon some ovarian or uterine derangement, which must be sought out and treated, or the physician's efforts are fruitless. When there is general anæmia, one of the best tonics is:

167. R.	Ferri et ammon. citratis,	℥ij
	Tinct. gentianæ,	f. ℥iv.
A teaspoonful in water after eating.		

Dr. THOMPSON'S solution of phosphorus does excellent service. It is:

168. R.	Phosphori,	gr. ss-iss
	Alcohol. absolut.,	q. s. to dissolve
	Ess. menth. pip.,	q. s.
	Glycerinæ,	ad f. ℥iv. M.
A teaspoonful after eating.		

Nutritious food and stimulants are required. Strychnia often does harm, while opium in small doses is generally of great and immediate value. Galvanization of the cervical sympathetic is an important auxiliary; so also are Chapman's spinal bags, filled alternately with ice and hot water.

NOTES ON REMEDIES.

Conium. According to Dr. BROWN-SÉQUARD, as an agent affecting the circulation in the brain and spinal cord, and as a paralyzant of voluntary muscles by its effect upon the afferent spinal nerves, conium has not yet attained the high place in general practice which it deserves. It must be given in full doses of the fluid extract, and so administered is very efficient in spinal irritation.

Counter-irritants are, according to Dr. McCALL ANDERSON, (*Clinical Lectures*, 1878,) the most efficient means of cure, and rarely fail. Blisters and leeches should be applied directly to the spot of spinal tenderness.

Digitalis is often of service.

Electricity in short sittings, every other day, will relieve uncomplicated cases.

Leeches, said by Dr. HAMMOND to be hurtful, have been employed by others with advantage. Dr. McCALL ANDERSON orders half a dozen over the

painful spot, repeats them if called for, and follows with a blister. (*Glasgow Medical Journal*, August, 1870.) Dr. ARMAINGAUD, of Paris, also recommends this treatment when there is pain or pressure over one or more of the vertebral spinal processes. (*Chicago Journal of Mental and Nervous Disease*, January, 1874.)

Opium in small doses is favored by HAMILTON.

Phosphorus is often of value. (F. 168.)

Picrotoxine, in doses of $\frac{1}{30}$ to $\frac{1}{100}$ of a grain, has been given with advantage by Prof. HAMMOND.

Sinapis, applied steadily to the spine, is lauded very highly in this disease by Dr. ARTHUR GAMGEE. (*Practitioner*, February, 1877.)

Strychnia is rather a dangerous drug in these cases.

Zinc in the following combination forms an admirable revulsive. The formula is of Dr. A. ROTHROCK, of Pennsylvania :

169.	R.	Zinci chloridi,	3j
		Zinci cyanureti,	3ij
		Cerati simplicis,	3j.

For a plaster. Apply for an hour or two daily.

SPASMODIC DISEASES.

DR. W. H. ERB, OF HEIDELBERG.

This contributor to Ziemssen's *Cyclopædia* considers spasmodic diseases among the most unsatisfactory objects of medical treatment. Whatever treatment there is, is purely empirical. Tonic and dietetic measures form an important part of it. Bathing, message, travel, and the movement cure, have all their applications. *Electricity* occupies a prominent place in curative measures, but definite rules for its use cannot be given. Several methods must often be applied in succession before the most effective one is discovered. Narcotics and nervines are the most valuable classes of drugs, and all of these have been tried with varying success.

In *mimetic facial spasm* or *convulsive tic*, brilliant results have sometimes been obtained by neurotomy. Of other means, electricity is of greatest avail. The subcutaneous injection of morphia is important, but satisfactory results are rarely obtained.

PROF. WM. A. HAMMOND, M. D., OF NEW YORK.

The class of paralyses or spasms produced by the habitual use of a

set of muscles for a long time, as in *writers' cramp* or *scriveners' palsy*, etc., has been called by Professor HAMMOND *anapeiratic paralysis*. He believes rest from the labor causing the disease is indispensable in the treatment, and sometimes sufficient in itself. The constant galvanic current has proved the most effectual agent in his hands. In incipient stages, *bromide of zinc* is valuable for restoring tone to the nervous system. It should be used in gradually increasing doses. Phosphide of zinc and *nux vomica* (F. 8,) are also valuable.

DR. REUBEN A. VANCE, OF OHIO.

This physician has derived great advantage in *writers' cramp* from the hypodermic use of *atropia*, combined with rest to the part.

The best place in which to inject the remedy is beneath the skin, in the loose cellular tissue over the pronator muscles of the forearm. Either extremity can be used, but when the treatment is to be continued for a length of time, it is better to inject in each arm alternately. The hypodermic syringe should be small, and armed with a gold-pointed needle. The solution of *atropia* should be concentrated, so that but a small amount of liquid need be injected at a time. The following is a convenient formula :

170. R.	Atropiæ sulphat.,	gr. j	
	Acid. sulph. dil.,	q. s.	
	Aquam,	ad ʒj.	M.
Dose, one minim.			

The injections should be administered about three times a week, and when attention is paid to the cleanliness of the needle, and the due concentration and freshness of the solution, no danger of producing abscesses need be feared, if the operation is performed with a reasonable degree of dexterity.

Dr. F. RUNGE has called attention to some causes of local irritation, which produce these cramps. In one case he found a chronic periostitis at the external condyle of the humerus. This node was extremely sensitive to pressure, and this pressure evoked spasmodic contractions of the extensor muscles, originating there, of precisely the same character as those suffered from when the patient persisted in writing. The actual cautery was applied over the inflamed spot, and this, with six weeks' entire rest, cured the patient, who was still well twelve months afterwards. In the second case, an ingrowing nail (of the right thumb

of the patient) was the original cause of the cramps, for it compelled the patient to alter the mechanism—in other words, the position—of his hand in writing. When this was healed, and the matrix of the nail no longer tender, the cramps all ceased, and the patient was well. (*Berlin Klin. Wochenschrift*, No. 21, 1873.)

Nitrate of Strychnia, hypodermically, deserves trial.

Dr. E. HERTZKA reports a case of cure of *piano-players' cramp*, in the *Petersburger Medicinische Zeitung*, 1876, by the use of eight drops of *tinctura gelsemii* three times a day. The treatment lasted three weeks; hydrotherapeutic and electric methods of treatment had proved insufficient.

Dr. G. VIVIAN POORE, of London, has derived better results in *writers' cramp*, from *galvanism* than from any other means. (*Lancet*, January, 1875.) His words are: "For genuine 'writers' cramp,' dependent upon chronic fatigue, I have found the refreshing effects of the galvanic current of the greatest possible service, and I have had four extreme cases of patients who had exhausted all the usual remedies, and had suffered for very lengthy periods, in whom rest and the use of the galvanic current, combined with a rhythmical exercise of the affected muscles, was productive of rapid and marked amelioration of the objective and subjective symptoms."

In *hiccough*, *singultus*, or *spasm of the diaphragm*, violent mental impressions will often check the nervous action. Hot fomentations, sinapisms, or the electric brush applied to the diaphragmatic region, have the same effect. Internally, a very effectual remedy is snow, pulverized ice, or a plate of ice cream. Either of these will relieve ordinary cases. Of drugs, the most effectual are the narcotics and anti-spasmodics. The following combination will often give immediate relief in obstinate cases:

171. R.	Chloralis hydratis,		
	Potassii bromidi,		
	Potassii bicarbonatis,	āā	3j
	Liq. morph. sulphatis,		f. 3j
	Aquam menth.,		ad f. 3vj.
			M.

Two tablespoonfuls every three hours.

The internal use of mustard has been highly commended; one teaspoonful of flour of mustard is to be infused in half a pint of boiling water, the infusion filtered, and to be given the patient at one draught.

In chronic hiccough from no assignable cause, the following is useful:

172. R. Tinct. physostigmatis, ℥x
 Pot. carb., gr. x
 Mist, acaciæ, ʒj. M.
 S. One dose, *ter die*.

No vegetables should enter into the diet. The food should be non-stimulant, making use of milk, eggs, etc. In some cases, tinct. of Calabar bean alone answers very well. Sometimes a hypodermic injection of morphia cures or relieves.

The following combination has been used with excellent effect by Dr. F. PEYRE PORCHER, of South Carolina, in troublesome hiccough :

173. R. Olei succini, f. ʒj
 Spts. ætheris nitrosi, f. ʒss
 Olei menthæ piper., f. ʒj
 Potassii bicarbonatis, ʒij
 Spts. ætheris compos., f. ʒij
 Mucilaginis, f. ʒij. M.

A tablespoonful every half hour.

The other remedies to which we must have recourse in obstinate cases, are: *Chloroform*, combined with small doses of morphia (RINGER), *morphia*, *hypodermically*, *camphor*, *chloral*, *ether*, *jaborandi*, or *pilocarpine*, *belladonna*, *quinine* in full doses, *musk*, (which has been extolled as a specific in this affection,) and *oil of amber* (which is said to rank next to musk). Dr. FAVIER reports a case of very obstinate hiccough that had lasted without interruption for fifty days, and was finally cured by firm *compression of the epigastrium* with a tourniquet; and firm pressure with the end of the finger in the sulcus, at the base of the nasal septum, in an always convenient and frequently effective means. Several cases are reported in the *Compend. of Med.*, 1884, where a hypodermic injection of gr. $\frac{1}{2}$ of pilocarpine checked at once cases of hiccough which had resisted all other remedies.

In the condition of tonic muscular contraction, known as *tetany*, which is closely akin to hysterical contractions, Dr. N. B. EMERSON has made successful use of *conium*, which is a powerful relaxant. (*New York Medical Journal*, July, 1876.)

Cramp of the muscles, especially of the gastrocnemius, is a common affection of obscure origin. Women with child are very often subject to it. Rubbing the part, or jumping from the bed and standing on the affected leg, are the usual means employed. If these fail, a hypodermic injection of morphia will give the most prompt relief. When hereditary, these attacks are extremely obstinate.

Dr. FREDERICK JAMES BROWN has called attention to the importance, in all convulsive disorders, of placing the patient in the sinistro-lateral posture. (*Practitioner*, May, 1877). He is certain that by attention to this, important results may often be attained. He gives the following example: "Recently, a man aged fifty-six years, in impaired health from chronic catarrh, was seized with unilateral (right) convulsions. His consciousness and power of speech were intact. He had been convulsed for ten minutes when I entered the house, and he was growing worse. I turned him over upon his left side, and the convulsions ceased in about ten or fifteen seconds."

Spasmus nictitans. In the *Med. Record*, 1883, Dr. Friedr. Betz relates the history of a boy, fourteen years of age, who had suffered for several years with a bilateral nictitating spasm of the lids. Several physicians had attempted to relieve him, but in vain. The boy was healthy, and his eyesight good. He had, however, quite long eyelashes, and the edges of the lids were reddened. Betz performed epilation of numerous lashes above and below on both eyes. The result was immediately successful.

TIC DOULEUREUX (FACIAL NEURALGIA).

FRANCIS E. ANSTIE, M. D., LONDON.

In the treatment of this terrible kind of neuralgia, to which TROUSSEAU gave the name of *epileptiform*, Dr. A. recommends (*Half-Yearly Compendium of Medical Science*, July, 1869,) the following plan in the earlier stages of the malady:

1. *Counter-irritation*, applied, not to the branches of the fifth, but to those of the occipital nerve, at the nape of the neck. A blister in the former situation is often as useful; in the latter, it is sometimes strikingly effective in gaining a short respite.

2. *Nutritive Tonics.* The assiduous use of cod-liver oil, or of some fatty substitute for it, should be insisted on from the first, and is of the highest consequence.

3. *Subcutaneous Injection* of morphia, or of atropia, according to circumstances. Commence with the use of $\frac{1}{6}$ of a grain of morphia twice daily, increasing this, if necessary, to $\frac{1}{4}$ and $\frac{1}{2}$ of a grain, and in

Prof. NUSSBAUM, of Munich, in the *Ärtzl. Int. Blatt*, 1883, this great surgeon recommends a trial with *salicylic acid* before proceeding to stretching or to resection of the nerve. In all the recently-sent cases he first tried this remedy, and he found it in every one a radical cure; not only a palliative effect, but really an utter disappearance of the painful disease was the result in every case. Especially in cases of rheumatic nature, N. is positive of having discovered in salicylic acid a specific for tic douleureux. He administered the drug in the following manner:

178. R. Acidi salicylici, 0.2, gr. $3\frac{1}{4}$
 Sodii salicylatis, 2.0, gr. 32. M.
 Ft. pulv. Within twenty-four hours the patient takes from four to six of such powders.

EDWARD WAKES, M. D., LONDON.

179. R. Potassii bicarbonatis, \mathfrak{z} iss
 Extracti ergotæ fluidi, f. \mathfrak{z} j
 Infusi ergotæ, f. \mathfrak{z} vj. M.
 Two tablespoonfuls every four hours.

B. M. WALKER, M. D., PLYMOUTH, NORTH CAROLINA.

Our author reports (*American Journal of the Medical Sciences*, April, 1874,) a case of tic douleureux treated by *ice*. The patient, a lady aged sixty, had been treated in various ways during five years, no remedy securing a longer period of relief than a few weeks. She was directed to use ice upon the part, which was over the fifth pair of nerves, and to continue the application as long as she could bear it, at the same time holding brandy in the mouth. The pain, for a while, was more intolerable than ever before; then it began to subside, and perfect quiet was obtained. The pain did not return the next day, as was customary, nor was there any subsequent return nor necessity for treatment.

NOTES ON REMEDIES.

Acidum Arseniosum is sometime prescribed by Prof. DA COSTA. (F. 174.)

Aconitia is employed by Prof. GUBLER, as follows:

180. R. Aconitiæ crystal, gr. $\frac{1}{8}$
 Alcoholis, q. s
 Aquam, q. s., ad f. \mathfrak{z} ij.

One teaspoonful every eight hours. He says he has never known a case to resist this heroic remedy.

Ammonii Chloridum, in doses of \mathfrak{z}_{ss} , four times daily, often succeeds. If the pain does not yield after five doses, no benefit need be expected from this remedy.

Aqua Ammoniae has been found of benefit in some cases, in doses of \mathfrak{m}_{xx} -xl in a cupful of thick gruel, at bedtime.

Argenti Nitras has been found, by ROMBERG, of great, but not permanent benefit, in doses of gr. j several times daily.

**Atropia*, hypodermically, is a valuable remedy, in this, as in other forms of neuralgia.

Belladonna ranks among the first of sedatives and anodynes, in this, as in other neuralgic affections.

Beberiae Sulphas, (in doses of gr. x, night and morning,) has been recommended by Prof. J. Y. SIMPSON and others.

Cannabis Indica is a capricious remedy, but one which (\mathfrak{m}_{xij} -xx of the tincture) often allays pain after the failure of other sedatives.

Caffein. One grain doses of the citrate every hour, for sometime before the expected paroxysms, have been recommended.

Ergota has been found useful by D. E. WAKES. (*British Medical Journal*, August 8th, 1868.)

Ferrum is of service in anæmic cases.

Hyoscyamus, combined with oxide of zinc, has an excellent reputation in Germany. (F. 177.)

**Morphia*, hypodermically injected, is a speedy and powerful remedy. The injections should not be made under the skin of the face.

Oleum Tigllii has been employed with benefit, internally.

Piscidia Erythrina, the Jamaica dogwood, is reported as a valuable means of relief by Prof. J. A. LARRABEE, of Louisville, (PARKE, DAVIS & Co., *Working Bulletin*, 1883.)

Stramonium is said by PHILLIPS, often to give very decided relief when exhibited in large and steadily continued doses.

EXTERNAL REMEDIES.

**Aconitum*, locally, often subdues the disease, attention being paid, at the same time to the state of the bowels, and to the general health.

Aqua Ammoniae has been found, by a number of French physicians, to rapidly cure some obstinate cases, when applied, with a camel-hair brush, to the palate and gums, so as to cause a profuse discharge of tears and saliva.

Counter-Irritation at the nape of the neck is recommended by Dr. ANSTIE. Dr. DA COSTA sometimes orders a hot salt-bag to the back of the neck.

Delphinia, applied externally, is stated by Dr. TURNBULL to be very efficient. It may be employed in the form of a solution:

181. R. Delphiniae,
Alcoholis,

\mathfrak{z}_{ij}
f. \mathfrak{z} xij. M.

Or, in ointment :

182.	R.	Delphiniaë,	gr. xxx	
		Olei olivæ,	℥. ʒj	
		Adipis,	ʒj.	M.

Hydrargyri Iodidum Viride is recommended in ointment, of the strength of forty grains to the ounce of lard.

Physostigma. A solution of the extract of Calabar bean (one in forty) applied to the eye has been used successfully.

Veratria is of great service, applied in ointment.

Electro-puncture, employed during the intermission of pain, is often a valuable method of treatment.

Hypodermic Injections. Dr. ANSTIE employs injections of morphia or of atropia, according to circumstances. Dr. RUPPNER reports a number of cases treated with success by the injection of liquor opii compositus, ℥x. In general, in regard to the use of hypodermic injections, the same is true as in other forms of neuralgia.

Neurotomy. Prof. NIEMEYER quotes BRUNS in support of division of the nerve, or excision of part of it, in proper cases. He shows, after a careful examination of the cases where it had been done, that, after excluding those where its want of benefit depended on error of diagnosis or operation, and after excluding the cases where the recurrence of pain was to be regarded as a new attack, not as a relapse, there was a considerable number of cases where the neurotomy caused either a complete and permanent benefit, or at least a temporary one for a few months or years. The operation is indicated when the pain is very limited, when it is probable that its exciting cause is situated at a point beyond which the nerve is accessible to the knife, when other treatment has proved fruitless, and when the pain renders the patient unfit for business. There is also an indication to operate when we cannot hope to divide the nerve between the point of disease and the brain, but when the pain never occurs spontaneously, being always the result of some external irritation acting on the peripheral terminations of the nerve. In such cases, the operation may, to some extent, protect the patients from the exciting causes of their attacks of pain. The temporary *compression* of the affected nerve, and the artery supplying it, has also sometimes proved an excellent palliative, and deserves a trial in suitable cases.

VERTIGO.

DR. EDWARD C. MANN.

Vertigo arising from disordered digestion, this author states (*Psychological Medicine*, 1884,) may be effectually relieved by the follow

ing treatment: The patient should take a cold bath on rising, and use free friction with a Turkish towel or flesh-brush. No malt liquor must be taken. The diet must be plain and regular. Vichy water, with perhaps a little brandy, may be used as a drink. The patient must retire early and sleep in a well-aired room. The following may be taken before meals:

183. R.	Pulv. rhei.,		f. 3j
	Sodii bicarbonat.,		
	Pulv. gentianæ,	āā	3 ij
	Aquæ menth. pip.,		
	Aquæ destillatæ,	āā	f. 3 iij.

Vt Sig.—Tablespoonful thrice daily before meals.

Instead of this, five drops of the tincture of nux vomica may be given before meals.

Many cases of vertigo arise from alcoholic indulgence and mental anxiety. They will not improve until these causes are removed. In others, an examination of the urine will reveal oxalate of lime, which produces oxaluria, and gives rise to vertigo. This can readily be relieved by fifteen-drop doses of dilute nitro-muriatic acid given in water thrice daily before meals. Other cases are characterized by an abundance of lithic acid or lithates in the urine. In these the vertigo is severe and often repeated daily or twice daily, accompanied with confusion of mind and loss of memory. The patient must be placed on a strict diet, using only white meats; he must rest, exercise freely, and use alkaline purgative waters, as the Saratoga, Vichy and Poland. *Citrate of lithia* is an appropriate medicine, and a course of small doses of arsenic is advisable.

PROF. AUSTIN FLINT, M. D., NEW YORK.

The first and most important measure in the treatment of vertigo consists in the removal of the cause or causes at work. All measures, such as blood-letting, counter-irritation and purgation, addressed to cerebral congestion, are hurtful. On the contrary, tonic remedies, nutritious diet and other hygienic means of invigorating the general health, are useful. The practitioner should bear in mind the importance of assuring the patient positively of the absence of all danger. *Quinia* is the most useful drug Dr. FLINT has tried; it must be given in full doses.

DR. J. M. DA COSTA.

In regard to gastric vertigo, this teacher states that it is very often necessary for the patient to make an entire change in his habits and surroundings—*e. g.*, to take a sea voyage or a trip of some duration. After due regulation of the diet, he thinks that the best treatment consists in the administration of *bitters* and *alkalies*, the bitters being given before meals, and alkaline waters, etc., after meals. Later on, when the way has been well paved by the regular administration of these, iron, combined with $\frac{1}{16}$ of a grain of *strychnia*, given three times a day, is to be preferred. Another plan consists in the administration of small doses of *corrosive sublimate*, while paying strict attention to the bowels and diet. This exerts a most happy influence upon the gastric and cerebral phenomena. Iron should not be given early in the case, when it may not be well digested, and may cause constipation.

DR. S. WEIR MITCHELL, OF PHILADELPHIA.

This able neurologist, at the close of an exhaustive study of the causes of vertigo (*Medical and Surgical Reporter*, June, 1877), states that there is no drug which holds to it the same relation that the bromides do to epilepsy. The nearest approach to specific medication lies in the use of vascular tonics like *digitalis*; moderate doses of *ergot*; and where pallor occurs, inhalations of nitrate of amyl as a means of relief for the immediate onsets. Cold douches are valuable; but, above all, the use, once in five days, of the *actual cautery* on the neck, is the most certain agent. Freeze the part, first, with a piece of ice sprinkled with salt, then press on the insensible part the point of a gas cautery, until it is just felt and no more.

In *gastric vertigo*, when all other means fail, the thorough adoption of *milk diet* seldom disappoints.

Aural vertigos demand points of cautery behind and in front of the ear, alternately. Ocular vertigos require appropriate glasses. *Tobacco* is a common cause of vertigo, and in all cases, its use must be prohibited.

H. V. SWERINGEN, M. D., OF INDIANA.

Vertigo is very often dependent upon imperfect digestion; in consequence of this fact, I seldom fail to afford relief by the following:

184. R. Potass. bicarb, 3 ij
 Liq. lactopeptin, f. 3 j
 Tr. nucis vom. f. 3 ss
 Tr. card. co., f. 3 ij
 Syr. simp., f. 3 ij
 Aqua Menth. pip. q. s. ad. f. 3 iv. M.
 Ft. sol. et. Sig.—A tablespoonful in water every three or four hours.

For plethoric persons I sometimes add to the above the following:

185. R. Potass. bitart. 3 vj
 Jalapa pulv., 3 ij. M.
 Sig.—A tablespoonful in milk, syrup or molasses every two or three hours until it operates freely on the bowels.

C. HANDFIELD JONES, M. D., LONDON.

186. R. Hydrargyri chloridi corrosivi, gr. j
 Glycerinæ, f. 3 j
 Tincturæ cinchonæ compositæ, f. 3 ij
 Olei menthæ piperitæ, mxxv. M.

A teaspoonful in a wineglassful of water three times a day, in the *vertigo of old persons*, which occurs sometimes paroxysmally, as a single symptom, unassociated with any special state that can account for it. Dr. THOMAS HAWKES TANNER also recommends this treatment for the attacks of temporary dizziness to which the aged are liable.

NOTES ON REMEDIES.

Ammonii Bromidum has been found, by Dr. RAMSKILL, of great help, when given in an effervescing form, with cascarilla, in cases of vertigo from overwork in the well-fed, when there is usually restlessness, insomnia, and depression of spirits, with a sense of impending evil.

Cubeba, in doses of gr. 1, three times a day, at meal time, is recommended by Dr. DEBOUT, in vertigo occasioned by excessive mental emotion or intellectual labor, and associated with dyspeptic derangement.

* *Ferri et Strychniæ Citras* is recommended by Dr. RAMSKILL in essential vertigo (*i. e.*, that unaccompanied with any other head symptoms). The patient should be placed on a long course of it, given in an effervescing form, and alternating, month by month, with tincture of larch and small doses of digitalis.

* *Hydrargyri Chloridum Corrosivum* is recommended in small doses, with bark, by Drs. C. HANDFIELD JONES and T. H. TANNER, in the temporary vertigo of old persons.

Nux Vomica and *Strychnia* are available in many cases.

Potassii Bicarbonas is very serviceable in dyspeptic vertigo.

* *Potassii Bromidum* is, according to Dr. RUSSELL REYNOLDS, often of service in vertigo, especially in cases which occur paroxysmally, without the co-existence of obvious spasm or organic disease of the brain.

Tonic remedies are recommended by Prof. AUSTIN FLINT.

Valeriana is very serviceable in that form of vertigo, with cardiac palpitation, which is not an unfrequent accompaniment of the menopause, or of the first establishment of the menses, in which the blood seems to rush into the head and then suddenly back upon the heart.

Zinci Valerianas, in doses of gr. i-ij, thrice daily, associated with extract of valerian, is recommended by Dr. DELIOUX in the treatment of nervous vertigo.

II. DISEASES OF THE RESPIRATORY SYSTEM.*

Aphonia—Asthma—Bronchitis, Acute—Bronchitis, Chronic—Catarrh—Coryza and Post-Nasal Catarrh—Emphysema—Empyema—Hæmoptysis—Hay Fever, Hay Asthma, Rose Cold, and Autumnal Catarrh—Laryngitis—Phthisis—Pleurisy, Acute, Subacute and Chronic—Pneumonia.

APHONIA.

The forms of aphonia usually encountered are (1) the catarrhal, (2) the hysterical, nervous or spastic, and (3) paralytic aphonia.

In *catarrhal aphonia* and excessive hoarseness,

DR. J. M. DA COSTA, PHILADELPHIA,

states that the local application of the *sulphate of zinc* is often productive of permanent good.

187. R. Zinci sulphatis,	gr. j-vj	
Aquæ,	f. $\frac{3}{4}$ j.	M.
Use as an inhalation.		

Muriate of ammonia is another efficient agent.

188. R. Ammonii chloridi,	f. $\frac{3}{4}$ ss	
Syrupi prun. virgin.,	f. $\frac{3}{4}$ iij.	M.
A teaspoonful thrice daily.		

This can be advantageously supported by $\frac{1}{8}$ of a grain of strychnia thrice daily, to give tone to the relaxed laryngeal muscles.

Nitre and *Benzoin* are efficient drugs in this variety.

DR. H. NICHELL, OF BUFFALO,

has found great advantages from the following combination in catarrhal aphonia.

*The subjects Stomatitis, Pharyngitis, Tonsillitis, Tonsillar Hypertrophy, Ozæna and Rhinitis are treated in Napheys' *Surgical Therapeutics*. Diphtheria, Croup, etc., will be found in this volume, under Diseases of Children.

189. R. Potassii bromidi, ℥ ij
 Extracti aconiti, gr. ij
 Ant. et pot. tartrat. gr. j
 Syr. pruni virgin,
 Aquæ destill., āā ℥ ij.

A dessertspoonful every three hours until relieved.

Dr. H. V. SWERINGEN, of Indiana, has used with advantage a combination of *muriate of ammonia* and *cubebæ*, ℥ij of the former and gtt. xxx of the tincture of the latter in a ℥vj mixture, of which a tablespoonful should be taken every three hours.

N. GALLOIS, M. D., PARIS.

190. R. Immerse a piece of thick paper in a solution of nitrate of potassium, and dry it; then cover it over with the compound tincture of benzoin, cut it up into pieces $3\frac{1}{2}$ by $1\frac{1}{2}$ inches, and make into cigarettes.

These cigarettes are highly praised in aphonia.

PROF. O. REVEIL, M. D., PARIS.

191. R. Pulveris benzoini, ℥ ss.
 Place a portion on some live coals, and inhale the vapor, by deep inspiration, in cases of aphonia and hoarseness.

Common salt is often beneficial.

192. R. Sodii chloridi, gr. v
 Aquæ destil., f. ℥j. M.
 Use with an atomizer.

M. ROZIERE, PARIS.

193. R. Balsami tolutani, f. ℥v
 Æther. sulphuric., f. ℥ iss. M.

Put in a wide-mouthed bottle and inhale for a few minutes every half hour, in aphonia and excessive hoarseness.

The German physicians use largely inhalations of *oil of turpentine*, one to five drops to the ounce of water. It is also highly spoken of by TROUSSEAU, COHEN and others.

In *hysterical or nervous aphonia*, the inhalation of an anæsthetic sometimes gives prompt relief. Dr. RUSSELL REYNOLDS has found benefit in blistering the larynx. But undoubtedly the most effective treatment is the electrical. Both kinds of electric currents have given satisfactory results; and when local applications fail, Dr. JOH. SCHNITZLER, of Vienna, has obtained prompt cures from galvanization of the nerve centres. (*Medicinische Presse*, No. 22, 1875.)

Locally, one may employ

194. R. Ferri chloridis, gr. iij
Aque, f. ʒj. M.
For an inhalation.

195. R. Argenti nitratis, gr. j-ij
Aque, ʒj. M.
For an inhalation.

Air, exercise, diet, and the daily shower-bath, are important auxiliaries, and will often of themselves restore the voice. Subjecting the patient to complete anæsthesia will sometimes at once break up the neurosis.

Paralytic aphonia may also be treated by inhalation. Prof. JOSEPH PANCOAST has used, with success, *chlorine vapor*, evolved by dropping slowly dilute muriatic acid into a solution of chloride of lime.

Turpentine and *tannin*, in inhalation, may also be used.

But the correct treatment of this form is undoubtedly by *electrization*. Drs. BEARD and ROCKWELL remark that there are few local disorders that yield more uniformly to any mode of treatment than aphonia to electrization. The current may be applied externally; but the proper course is to use the faradic current, and make the application directly to the cords by means of laryngeal electrodes.

NOTES ON REMEDIES.

Alumen. BENATI, formerly physician at l'Opéra Italien, Paris, ordered the following prescription to restore the natural tones of the voice:

196. R. Aluminis, ʒj
Decocti hordei, f. ʒ ijss. M.

To be used as a *gargle*, the strength to be gradually increased. The general health should be, at the same time, carefully looked after, and the exercise of the voice regulated. The internal administration of alum (gr. x a day, in divided doses, gradually increased to gr. xlv), has been also found very beneficial. Alum lozenges are indicated to complete the cure, and are efficient alone, in milder cases.

**Ammonia.* The inhalation of ammoniacal vapor, disengaged from a mixture of a solution of chloride of ammonia and carbonate of potassium, has given excellent results in aphonia in consequence of exposure to cold. The utility of inhalations of chloride of ammonium vapor, in these cases, has been tested by Dr. BEIGEL in a number of cases. (*Practitioner*, August, 1868.

**Atropine*, Dr. ROBERTS BARTHOLOW says, is almost sure to cure hysterical aphonia, if given to produce constitutional symptoms.

Acidum Nitricum is very effective in the hoarseness of singers, when the aphonia is reflex from stomach disorders, and in fatigue of vocal cords. It should be given well diluted, in sweetened water.

Benzoinum, in vapor, is recommended by French physicians. (F. 190.)

Chlorinum. Chlorine inhalations are serviceable in aphonia following an ordinary cold, without organic lesion.

Iodinium. Iodine inhalations, and the application of the tincture of iodine to the fore part of the neck, are worthy of trial in cases arising from chronic thickening, or from debility and relaxation.

Quiniæ Sulphas. In all cases where the aphonia is intermittent in character, malarial poisoning may be suspected, and the free administration of quinine will be appropriate.

Sodii Boras. According to Dr. J. W. CORSON, (*Medical Record*, January, 1873,) relief will often be given when the voice has been suddenly lost, by slowly dissolving in the mouth and partially swallowing a piece of borax, grs. iv-v.

Zinci Sulphas is of use, either locally applied, in solution, to the cords, or inhaled (F. 187.)

Zingiber. A ginger gargle, or a piece of the root employed as a masticatory, is often useful in aphonia from relaxation of the parts.

ASTHMA.

Many drugs are more or less beneficial in asthma; but what succeeds admirably in one case, may fail completely in the next one, and the physician must resort to something else. Whenever possible, treatment should be directed to the cause; and in all instances, hygienic rules should be insisted upon, especially *diet*:

DR. JOHN C. THOROWGOOD, OF LONDON.

Diet in asthma. This writer calls attention, in the *Medical Press and Circular*, 1884, to the importance of carefully regulating the *diet* in cases of asthma and bronchitis. An overly distended stomach, by pressing on the diaphragm, may offer a mechanical impediment to free respiration, and may also interfere with the proper action of the heart. He mentions a gentleman who gained a great reputation for his ability to cure bronchitis and bronchial asthma by diet, and says that the great secret of this diet cure was judiciously starving the patient for a time.

The stomach may be, to a certain extent, sluggish, so that it will not at the time feel any inconvenience from the excess of food forced upon it, but in an hour or two dyspepsia in some form will indicate the outrage that has been perpetrated. The food used should be nutritious, without being bulky. A lightly-boiled egg, the wing of a chicken, or some fish, with one cup of tea, coffee or cocoa, and toast, will form three alternatives of breakfast fare.

Soups should be avoided at dinner, because they tend to distend the stomach. Vegetables should be used sparingly. Malt liquors, pastry, cheese and dessert should be rigidly eschewed, and for drink, claret with water, or lemon juice and water, may be allowed. Effervescing drinks of all kinds are not to be commended. By this arrangement of diet we keep up nutrition, while we avoid overloading the stomach, and so oppressing the diaphragm. The heart and lungs are enabled to act freely, and the danger of sudden stoppage of the first-named organ is put at a distance.

M. HUCHARD, OF FRANCE,

uses the following when the symptoms of bronchial catarrh are added to the attacks of asthma:

197. R.	Distilled water,	300 parts	
	Iodide of potassium,		
	Tincture of lobelia,		
	Tincture polygala,	āā 10 parts	
	Extract thebaic,	℥ ¹ / ₈ part.	M.

A teaspoonful to be taken night and morning.

PROF. J. M. DA COSTA, M. D., PHILADELPHIA.

In the treatment of the paroxysms of asthma, all nauseants have a certain influence in relaxing the bronchial spasms. *Lobelia inflata* is particularly serviceable, because it acts both as a nauseant and expectorant.

198. R.	Tincturæ lobeliæ,		
	Tincturæ hyoscyami,		
	Spiritūs ætheris compositi,		
	Syrupi tolutanti,	āā f. ʒj.	M.

A teaspoonful in water every half hour, during the paroxysm, until some effect is produced on the breathing, and then every hour or two.

Or one of the following combinations:

- | | | | | | |
|------|----|-----------------------------|----|--------|----|
| 199. | R. | Spiritus ætheris compositi, | | | |
| | | Extracti valerianæ fluidi, | āā | f. 3j | |
| | | Tincturæ lobeliæ, | | f. 3ss | |
| | | Potassii chloratis, | | 3iss | |
| | | Syrupi tolutani, | | f. 3j. | M. |

A dessertspoonful in water three times a day.

Direct, also, the following stimulating liniment :

200. R. Chloroformi, f. $\frac{3}{3}$ ss
Olei terebinthinæ, f. $\frac{3}{3}$ j
Spiritus rosmarini, f. $\frac{3}{3}$ iss. M.

To be rubbed on the chest several times a day.

- | | | | | |
|------|----|--------------------|-------------------|-----------|
| 201. | R. | Potassii iodidi, | 3 ij | |
| | | Morphiæ sulphatis, | gr. $\frac{3}{4}$ | |
| | | Tincturæ scillæ, | | |
| | | Tincturæ lobeliæ, | | |
| | | Syrupi. | āā | f. 3j. M. |

A teaspoonful thrice daily, in asthma with emphysema and chronic bronchitis.

In order to prevent the recurrence of the paroxysms, too great care cannot be taken to inquire into the state of all the functions. In perhaps one-half of the cases, asthma is not a disease of the lungs, but a reflected trouble. In cases of stomach origin :

202. R. Pilulæ hydrargyri, gr.ij.

At night, to be followed by one-half an ounce of Rochelle salts in the morning. Afterwards direct arsenic, combined with gentian or belladonna; as,

203. R. Liquoris potassii arsenitis, f. ʒj
Tincturæ gentianæ compositæ, f. ʒ iij. M.

A dessertspoonful thrice daily.

Opium and *belladonna*, or their alkaloids, *morphia* and *atropia*, have been extensively used. Dr. J. G. OLIVER, of London, combines them in subcutaneous injection.

204. R. Morphiae acetatis, gr. $\frac{1}{4}-\frac{1}{3}$
Atropiae sulphatis, gr. $\frac{1}{100}$
Aquæ, q. s. M.

For one hypodermic injection in spasmodic asthma.

The conclusions he reaches, are :

1. Morphia and atropia are superior to morphia alone; the good effect is more speedy and complete, and they produce no depressing gastric disturbance.
2. The very frequent use of the injection does not appear to injure the general health.

2. The very frequent use of the injection does not appear to injure the general health.

3. Speedy relief. The effect is generally declared in five minutes, in the form of comfortable sleep and quiet breathing. The most intense attack has given way completely in from fifteen to twenty minutes. (*The Practitioner*, January, 1876.)

Dr. C. HANDFIELD JONES, London, recommends :

205. R. Liguoris opii sedativi, (Battley,) $\mathfrak{m}\text{x}$.
For one subcutaneous injection into the left front of the chest.
206. R. Atropiæ sulphatis, $\text{gr. } \frac{1}{15}$
Aquæ, $\mathfrak{m}\text{x}$. M.
For one hypodermic injection in the vicinity of the left vagus nerve.
207. R. Extracti cannabis indicæ, gr. v .
For ten pills.

This drug occasionally proves very useful, one pill immediately checking the spasm.

DR. LECLERC, FRANCE.

208. R. Extracti belladonnæ, gr. xv
Myrrhæ pulveris,
Ipecacuanhæ pulveris, $\text{āā } 3\text{ ss.}$ M.
Divide into thirty-six pills. One thrice a day, in nervous asthma. Nitre paper to be burned in the chamber of the patient, and stramonium leaves to be smoked.

The following cigarettes are those employed with good effect by Dr. GEO. A. THOMAS, of Indiana :

209. R. Pot. nitratis, $\text{gr. } \frac{1}{4}$
Tr. tabaci, $\mathfrak{m}\text{x}$
Olei anisi, $\mathfrak{m} \frac{1}{8}$
Opil, $\text{gr. } \frac{1}{4} - \frac{1}{8}$.
Swedish filtering paper $2\frac{1}{2} \times 4$ inches. This incorporated in the filtering paper and smoked, at intervals as necessary, by asthmatics, in his hands has given marked relief.

N. GALLOIS, M. D., PARIS.

210. R, Extracti opii, $\text{gr. } \frac{3}{4} - \text{iss}$
Belladonnæ foliæ, 3 j.
Dissolve the extract in a small quantity of water, and then moisten the belladonna leaves with this solution ; dry them, and roll into cigarettes.

TROUSSEAU substitutes for the belladonna leaves, stramonium leaves. Such cigarettes are useful in nervous asthmas and dry coughs.

DR. R. B. FAULKNER, OF NEW YORK.

In pure *spasmodic asthma* this writer (*Med. Record*, 1883,) advo-

cates a treatment by painting a strip of iodine over the course of the pneumogastric nerves in the neck. He gives three cases of pure spasmodic asthma, which were relieved of their attacks by this means, after having resisted every other remedy of which he could think.

DR. ROBERT SAUNDBY, of England, subsequently tried this method, and reported excellent results from it in the British *Med. Journal*.

PROF. FELIX VON NIEMEYER, GERMANY.

Among medicaments especially in repute for the prevention of new paroxysms, and for the radical cure of asthma, *quinine* stands first. The shorter and more regular the intervals of the attack, so much the more is to be expected from this drug. It is unsuitable when the pauses between the seizures are very long or irregular in their occurrence. In such cases we must have recourse to other remedies from the list of the so-called nervines.

As a rule, the metallic nervines are to be preferred before tincture of valerian, assafoetida, castor or camphor. The following may be used:

- | | | |
|--|------------|----|
| 211. R. Ferri carbonatis, | ℥j | |
| Syrupi, | q. s. | M. |
| For twenty pills, One thrice daily. | | |
| 212. R. Zinci oxidi, | gr. xxv | |
| Syrupi, | q. s. | M. |
| For twenty pills. One thrice daily. | | |
| 213. R. Argenti nitratis, | gr. iij-iv | |
| Confectionis rosæ, | q. s. | M. |
| For twenty-four pills. One thrice daily. | | |

The *iodide of potassium* was a favorite with

PROF. ARMAND TROUSSEAU, PARIS.

- | | | |
|--------------------------|----------|----|
| 214. R. Potassii iodidi, | ℥ ijss | |
| Spiritûs vini gallici, | f. ℥ iss | |
| Decocti senegæ, | f. ℥ iij | |
| Syrupi papaveris, | f. ℥ j. | M. |

In essential asthma, give a tablespoonful twice a day, an hour before the meals, in a little sweetened water.

During the attack, the tincture of lobelia may be administered, in doses of from twenty to thirty drops, from half-hour to half-hour. The respiration of chloroform is also useful; also stramonium cigarettes.

PROF. GERMAIN SÉE

(*L'Union Med.*, March 6, 1883), regards asthma as made up of three elements: Nervous, catarrhal, and emphysematous. When the latter condition is unduly prominent, we have permanent asthma.

His method of treatment, for which he claims excellent results, consists in the exhibition of full doses (thirty to forty-five grains per diem) of *iodide of potassium*. Under its continued use for some length of time, the asthmatic exacerbations disappear, and the permanent dyspnoea is greatly lessened, if not entirely controlled. It is sometimes useful to add from three-fourths to one and one-half grain of opium to the daily dose of the potassium salt, to prevent iodism.

During the paroxysm, however, M. Sée employs the *iodide of ethyle*, a substance discovered in 1825, by GAY-LUSSAC, and composed of iodine and ether, the new compound possessing the respective properties of both these substances. He administers it by inhalation, and he has often found that a single dose of five or six drops has been sufficient to cut short a paroxysm. The breathing once relieved, he then trusts to the iodide of potassium to effect a cure. The above treatment has been found useful in all cases of asthma, whatever their origin.

DR. A. TRASTOUR, OF PARIS.

Catarrhal Asthma.—In those asthmatic cases which are complicated with catarrh, this practitioner (*Bull. de Therapeutique*, 1883) lays down the following rules of treatment:

1. To modify the circulation. *Bleeding*, or one or more applications of leeches, repeated p. r. n.

2. To diminish morbid secretions. Persevering application of *blisters*, at longer or shorter intervals, on the chest. Daily frictions of the entire surface of the body with a hair mitten or brush. An occasional wet pack or douche. Tar, eucalyptol, turpentine, carbolic acid, or, as he prefers, creosote.

3. To combat the chronic inflammation, and bring about trophic changes, he advises *sulphur*, but principally *iodine*; and of this he thinks the best form is the *iodide of calcium*.

4. As to antispasmodics and sedatives, he has no preference; without any special indication, he is guided in his choice by personal experimentation on the patient. *Morphine*, however, is by far the best.

PROFESSOR DUJARDIN BEAUMETZ, OF PARIS,

recommends the use of *bromide of potassium* in those cases of intermittent dyspnœa, which are really a cardiac asthma, and which are frequently attendant on aortic affections. Small doses of pure chlorhydrate or bromhydrate of cicutine may be added to it.

THOMAS HAWKES TANNER, M. D., LONDON.

- | | | | |
|---------|------------------------------|----------|------|
| 215. R. | Potassii iodidi, | ℥ijss-iv | |
| | Spiritus ammoniæ aromatici, | f. ̄j | |
| | Tincturæ belladonnæ, | ℥l-f. ̄j | ijss |
| | Tincturæ cinchonæ compositæ, | f. ̄j | ij |
| | Aquæ menthæ piperitæ, | f. ̄j. | M. |
- A dessertspoonful, in water, three times a day.

In some cases of asthma, the author has found remarkable benefit from this formula. It requires to be persevered with for some weeks, the patient being watched, lest it impoverish the blood and produce purpura or boils, or even a carbuncle. If there be constipation, order

- | | | | |
|---------|------------------------|----|--------|
| 216. R. | Pilulæ rhei compositæ, | | |
| | Extracti conii, | āā | gr. v. |
- For two pills. To be taken at bedtime.

DR. WILLIAM M. WELSH, OF PHILADELPHIA,

has found the following compound of great benefit in asthmatic cases:

- | | | |
|---------|------------------------|----------|
| 217. R. | Potassii nitratis, | 2½ parts |
| | Belladonnæ fol. pulv., | ½ part |
| | Stramonii fol. pulv., | 5 parts |
| | Sacch. alb., | ½ part. |

Dissolve the nitre in just enough water to make a saturated solution, mix with the leaves, dry the mass into a coarse powder, and add the sugar. A small quantity is to be placed on a tin plate and ignited, and the smoke inhaled.

J. S. MONELL, M. D., NEW YORK.

Our author recommends (*New York Medical Record*, August 15th, 1866,) *forced expiration and inspiration, for the relief of spasmodic asthma.*

He directs that all the air be expired that it is possible for the patient to do, and not to inspire until it is found absolutely necessary. Then carry inspiration to its fullest capacity, and retain, with great effort, for many seconds. This act of forced expiration, waiting, thorough inspiration, and again waiting, should be continued for some fifteen minutes,

when it will be found that the spasm is relieved. It requires great exertion on the part of the patient to perform this act. The first attempt at retaining the inspired air, during the asthmie attack, will cause the patient to think he cannot continue it, but perseverance will soon delight him with relief from the spasm.

A general estimate of the value of the various remedies in asthma, is given in his classical treatise by

HYDE SALTER, M. D., LONDON.

Coffee relieves about two-thirds of the cases in which it is tried. The relief is very unequal, often merely temporary, sometimes very slight, sometimes complete and permanent. The following are valuable hints on its administration; 1. It cannot be given too strong. 2. It is best given without sugar and milk—pure *café noir*. 3. It should be given on an empty stomach. 4. It seems to act better if given hot—very hot.

Lobelia is given by Dr. S., in the form of the tincture, in repeated doses, every half hour, increasing the dose $\text{m} \vee$ each time, till either the physiological effects (faintness, sickness,) are manifested, with or without relief, or that relief is obtained without such symptoms. In either case, the medicine is to be at once discontinued. Generally, no relief takes place until symptoms of lobelia poisoning show themselves. Any amount may be given, short of producing the effects, without any relief at all. The quantity required differs in each case—generally, Ml – lx suffice; in others, $\text{f.}\overline{3}$ iss fail to produce an effect. When once the necessary dose has been ascertained in this way, the patient should start with it on the next occasion, which generally has the desired effect at once.

Iodide of potassium entirely fails in a great many cases. Sometimes, however, the most striking results attend its use. It often requires to be continued for some time before it begins to manifest its effects. It should be used in every case in which it has not been tried, because of its occasional great value.

Quinine is the best of all tonics in asthma; next to it is *iron*. The two may be combined with a mineral acid and given with good results.

Stramonium is, in some cases, *the* remedy; in the majority of cases, as ordinarily used, it does some good: and in some few it is positively injurious. Its value in any case can only be ascertained by personal experience. When the leaves and stems fail, substituting the seeds

may prove effectual. They appear to be more powerful in their action. The inhalation of the smoke, and swallowing the saliva, may be advantageously combined with the ordinary method of smoking. The earlier in a paroxysm it is had recourse to, the more likely will it be to relieve the spasm; over one that has been long established, it has but little power. As the great thing is to give it in time, an asthmatic should keep his pipe all ready filled, with the means of lighting it by his bedside, so that on waking with dyspnœa, he may immediately use it. A good plan is to adopt the habit of smoking a pipe of it the last thing at night, whether a paroxysm is threatened or not. The *extract* (gr. $\frac{1}{4}$, gradually increased to gr. j,) sometimes produces excellent effects, or the *tincture* (℞-xx every four hours) may be substituted.

Tobacco smoking affords, in some cases, remarkable ease. The habit of smoking has rendered powerless, in a large number of cases, what may, without any qualification, be called its most potent remedy.

Chloroform inhalations, *short of anæsthesia*, are often of benefit, and may be used in all cases.

DR. ROBERTS BARTHOLOW, OF PHILADELPHIA.

Asthma, when purely spasmodic, is usually quickly checked by inhalations of *amyl nitrite*. The paroxysms of difficult breathing which accompany emphysema and cardiac disease are not relieved in this way; indeed, the author has known the most serious distress to be produced by the inhalation under these circumstances. Some cases of spasmodic asthma are much relieved by nitro-glycerine. One minim of a one per cent. solution is the dose.* In uræmic asthma and chronic bronchitis, with asthmatic breathing, *quebracho* has afforded very considerable relief. It is much less useful, if not injurious, in the dyspnœa from valvular disease of the heart, and has had no effect in the dyspnœa of old persons, due to atheroma of the vessels. The preparation most used is the extract, and the adult dose is five grains. When asthma is accompanied with catarrh of the mucous membrane, *ethyl iodide* gives good results. Fifteen or twenty drops are put on a handkerchief and inhaled as may be necessary—every two, three or four hours.

In asthma with profuse expectoration (humid asthma) *pilocarpine* has been very successful. According to BERKHART, when injected subcutaneously, it will arrest the most violent paroxysms of spasmodic asthma, and the relief persists for some time.

* It should not be forgotten that the alcoholic solution will explode if struck with a wet hammer.

In humid asthma Dr. B. highly lauds the following:

218. R.	Tinct. sanguinariae,	3j	
	Tinct. lobeliae,	3j	
	Ammonii iodidi,	3j	
	Syr. toluani,	3vj.	M.

S.—A teaspoonful every two, three or four hours. The same formula may be used in spasmodic asthma.

NOTES ON REMEDIES.

Acetum. As a preventive, bathing the chest every morning with vinegar and water has been advised.

Æther by inhalation is sometimes resorted to with benefit in asthma. The following formulas have been recommended :

219. R.	Ætheris sulphurici,	30 parts	
	Acidi benzoici,	15 "	
	Balsami Peruviani,	8 "	M.

Or the following :

220. R.	Ætheris sulphurici,	2 parts	
	Spiritus terebinthinæ,	15 "	
	Acidi benzoici,	15 "	
	Balsami Peruviani,	8 "	M.

Place the mixture in a vessel having a large opening ; the warmth of the hand is sufficient to volatilize the materials, and inhalations may be used four or more times a day, as occasion demands.

Ammonii Carbonas has proved (in doses of gr. x-xv) more successful than any other remedy, in the hands of Dr. J. HOPE, in cases of asthma arising from, or connected with, *disease of the heart*. In a very obstinate case, the following afforded much relief:

221. R.	Ammonii carbonatis,	gr. vij	
	Antimonii et potassii tartratis,	gr. ¼	
	Aquæ anisi,	f. 3 iss.	M.

A draught to be taken every alternate hour.

Ammoniacum is of great service in both *spasmodic* and *hysterical* asthma. In the former case it should be combined with tincture of squill and a small portion of hyoscyamus or conium ; in the latter, with equal parts of assafoetida. Plasters of ammoniacum applied to the chest often prove useful.

Aqua Ammonia (four parts to one of water) applied to the velum palati, has proved of service in simple idiopathic asthma. This application, which may prove dangerous, is only to be made, if at all, with great caution, so as not to touch the back part of the pharynx. The inhalation of the vapor of aqua ammonia is useful.

Argenti Chloridum. Dr. CURCI, of Venice, highly recommends in nervous asthma:

222. R.	Argenti chloridi,	part j
	Sodii hyposulphitis,	" iij
	Aquæ destillatæ,	" xxx. M.

It may be administered by the mouth or hypodermically, and is not irritating.

Arsenicum is recommended by Dr. HANDFIELD JONES and others, but is inadmissible in organic disease of the lungs and heart. Dr. MARTELLI, of Florence, claims excellent results from hypodermic injection of Fowler's solution, one part to two of water. Dr. THOROWGOOD (*Med. Times*, 1883,) says: "I know of no drug so generally useful in the treatment of asthma as arsenic, and could give, from fifteen years' experience, a great many cases of spasmodic asthma that have seemed to me to get quite well under the influence of small doses of arsenic. I have never exceeded the dose of three minims of either Fowler's solution or of liquor sodæ arsenitis three times in the day, and have never seen any symptoms of a dangerous character."

Aspidosperma Quebracho. The good effects of this new drug in spasmodic asthma appear to be well established by the observations of Dr. PENZOLDT and others. Dyspnœa in all forms is often relieved by it. The drug has been difficult to obtain heretofore in this country, but at present several preparations of it are manufactured by PARKE, DAVIS, & Co., Detroit. The dose of the fluid extract is ʒ ss.

Aspidospermine. This is the alkaloid of quebracho bark. Dr. MARAGLIANO (*Lond. Med. Record*, 1884,) uses gr. $\frac{1}{4}$ for hypodermic injection in cases of asthmatic dyspnœa. Relief is felt in about five minutes.

Assafœtida, in combination with other antispasmodics and with narcotics, is often very beneficial in spasmodic asthma.

223. R.	Tincturæ assafœtidæ,	f. ʒ ss
	Tincturæ opii,	m _{xx}
	Spiritûs ætheris compositi,	f. ʒ ss
	Aquæ camphoræ,	f. ʒ iss. M.

For one draught.

* *Belladonna* is a valuable remedy in spasmodic asthma, administered in the manner directed by Dr. HYDE SALTER, viz., in one full dose, sufficiently large to produce the full physiological effects of the drug, every night at bedtime. This plan he regards as infinitely superior to the system of divided doses during the day. As there is a marked difference in the tolerance of belladonna in individuals, the proper dose may be ascertained by commencing with small doses and gradually increasing them until the requisite full dose is ascertained.

Bromides relieve in purely spasmodic asthma, but soon lose their effect.

Camphora is recommended by Dr. COPLAND as one of the most generally beneficial in any of the class of narcotics or antispasmodics. In *nervous* and *spasmodic* forms of asthma it should be given in large doses, (gr. iij-x,) and combined with musk, castor, assafœtida, etc., or with sedatives.

Cannabis Indica is said to occasionally prove very beneficial, but in the hands of Dr. HYDE SALTER it has been a complete failure in the few cases in which he employed it. Dr. EMIL KOLLER has, however, found it valuable in nocturnal asthma, in doses of gtt. v-vj of the tincture every three or four hours.

Castoreum, combined with ipecacuanha (equal parts of the tincture of castor and the wine of ipecac) is recommended by Dr. GRAVES.

* *Caffea* is an efficacious remedy. For its administration, see Dr. HYDE SALTER's directions (above, p. 156.)

Chloroformum by inhalation is an excellent and prompt remedy for spasmodic asthma.

Datura Tatula is said to be of great efficacy smoked in cigarettes.

Digitalis has been employed, but is inferior in efficacy and safety to other remedies.

Dracontium sometimes succeeds (in doses of gr. xxx-l of the powdered root during the paroxysm) in obstinate cases of spasmodic asthma.

Euphorbia pilulifera. This plant, obtained in Queensland, may be taken in decoction or extract, and is said to relieve severe attacks of chronic bronchial asthma. (Beck, *Therapent*, 1884.)

Grindelia Robusta. The plants and flowers of this native of the Pacific coast have been much lauded of late in spasmodic asthma. One drachm of the fluid extract may be taken when the paroxysm comes on, or half a drachm three times a day as a prophylactic. *Grindelia cigarettes* are made as follows: Impregnate well-nitred tobacco with an alcoholic fluid extract of *grindelia*; let dry and use in cigarettes. Owing to the nitre, they will continue to glow and to develop smoke.

Hyoscyamus is, according to Dr. HARLEY, the appropriate remedy in *cardiac* and *pulmonary* asthma. It is best administered subcutaneously.

Iodinium. In cases of dry asthma, of constitutional character, without obvious exciting cause, Dr. C. A. L. REED, of St. Louis, recommends, in the *Clinical Record*, 1879, the following prescription:

224. R. Liquor iodinii comp.,	f. ʒj	
Mellis,	f. ʒij.	M.
Teaspoonful every three hours.		

He finds that the use of this is promptly (after the third dose) followed by a restoration of the secretion, diminution in the frequency and severity of the paroxysms from the start, an improvement in the appetite, and an increase in the tone and vigor of the general system.

* *Ipecacuanha*, in full emetic doses (not less than gr. xx), is often effectual in arresting a paroxysm of spasmodic asthma, if given at or near the outset of the attack. Proportionate doses to children may be given with safety and advantage. Small doses of the wine (℥v) repeated every ten to thirty minutes for two or three hours, unless relief comes much sooner, in which case the dose may be diminished and taken less frequently, is

a method of treatment recommended by Dr. C. D. PHILLIPS, (*Practitioner*, Nov., 1869,) in *nocturnal spasmodic* asthma, when there is no organic disease of the heart or lungs.

* *Lobelia*. Dr. HYDE SALTER reports favorably of the tincture. (P. 156.) Dr. RINGER, while regarding it as a remedy of great efficacy, considers that it should be given with caution where heart disease exists, as the pulse may become irregular and very weak. It should also be remembered that some persons exhibit great intolerance of its action, poisonous effects following the smallest doses, and that different specimens differ considerably in strength.

Moschus, in doses regulated according to the severity of the attack, is often productive of benefit in *purely spasmodic* or *nervous* asthma. It has also been applied endermically (gr. vj-xv) with advantage.

Nitrite of Amyl. Cases have been reported in which the inhalation of five drops of this liquid has arrested very distressing attacks of spasmodic asthma.

Nitrogen. Dr. EDGAR A. HOLDEN recommends nitrogen inhalations. (*American Jour. of the Med. Sciences*, Oct., 1872.) The following is the method adopted with the protoxide : The gas, fresh and pure, is passed through a jar containing a fluid extract of conium, Calabar bean, or belladonna, according to the necessities of the case ; and from this jar is inhaled by the patient once daily, before retiring, and for four or five consecutive days, the quantity varying according to the condition of the patient, but not exceeding five to eight gallons, never to anæsthesia, the sitting occupying at least ten minutes. Relief almost always follows the first inhalation, and produces an uninterrupted night's rest. Upon the disappearance of the nightly paroxysm and the subsidence of all oppression in breathing, the remedy must be discontinued, but must be instantly resorted to upon the first evidence of return.

Opium is of questionable value in asthma. Dr. HYDE SALTER says : "Prefer any other sedative to opium ; and unless there is some special complication that indicates it, never give it at all."

Oxygen Inhalations have been tried with reported "wonderful" results in nervous asthma.

Petroleum. Dr. H. V. SWERINGEN, of Indiana, reports that in asthma he has derived the *most* benefit from the administration in half teaspoonful doses, gradually increased, of an emulsion of petroleum.

Pilocarpine is highly recommended by BARTHOLOW in spasmodic asthma. (See above.)

Potassii Bromidum, in full doses, at night, has been recommended in spasmodic asthma. In the *spasmodic asthma of childhood*, given in syrup, in doses of gr. vj every two hours, it has proved very successful after the failure of other remedies.

Potassii Iodidum is an uncertain remedy, which sometimes, however, proves strikingly successful.

Potassii Nitratis. The inhalation of the fumes of paper impregnated with nitre,

is often of striking benefit. The following formula is given in the *Journal de Pharmacie et de Chimie*: Take four ounces of white paper, and allow it to macerate in warm water, until reduced to a uniform paste. Then press out the greater portion of water, and mix the residue in a mortar, with the following powder :

225. R.	Potassii nitratis,	℥ ij
	Myrrhæ,	
	Olibani,	āā 3 ijss
	Belladonnæ,	
	Stramonii,	
	Digitalis,	āā gr. x.

When a uniform mass has been formed, roll into sheets a line or so thick, dry, and cut into strips. This paper is said to burn less quickly than the ordinary nitre paper, and to be more effective. On account of the unpleasant odor of the burning paper, Dr. R. E. THOMPSON (*Practitioner*, April, 1879,) recommends the following formula as a basis for a *medicinal cigarette*:

226. R.	Swedish filtering paper, size,	4 in. x 2½ in.
	Potassii nitratis,	gr. ¼
	Tinct. tabaci,	℥ x
	Olei anisi,	℥ ½.

The tincture of tobacco is made with ℥ 2½ of the leaf to a pint of spirit. A solution of any drug which is to be experimented on, can then be prepared, and the paper, having been floated through the solution in a flat dish, when dry, can be cut into a certain size, and the dose thus accurately measured.

Quinia Sulphas is highly recommended by Drs. NIEMEYER and HYDE SALTER. (See above.) FLINT says that gr. xx-xxx at one dose, will often break up a paroxysm.

Quebracho is recommended by Dr. ROBERTS BARTHOLOW. (See above.)

Sanguinaria has been prescribed with good effects. Dr. L. B. ANDERSON, of Virginia, gives it as in croup, until vomiting occurs. It rarely fails to give prompt and grateful relief. As a preventive, he continues it in doses of 20 drops of the tincture three times a day. (*Southern Medical Record*, February, 1876.)

Scilla. Benefit is sometimes derived from squill, combined with hemlock or henbane. The following formula is often useful :

227. R.	Tincturæ scillæ,	gtt. xv	
	Extracti hyoscyami,	gr. iij	
	Acidi nitrici diluti,	℥ xxx	
	Aquæ,	f. 3 iss.	M.

For one dose.

Spiritus Ætheris Compositus. Hoffman's anodyne, though favorably spoken of by some writers, has proved entirely useless in the hands of Dr. HYDE SALTER, and in some cases, has seemed to increase the spasm.

**Stramonium*. (See above.)

Strychnine is indicated when there is weakness of the respiratory muscles.

Sulphur, in doses of about gr. viij daily, to be given while fasting, and persevered in for a long period, has been recommended by French physicians.

Tabacum. The palliative effects of tobacco fumes, in nervous asthma, are well known.

Veratrum Viride has been advised in nervous asthma. The following formula is said, by Dr. J. L. COOK, to give admirable results when other remedies fail :

228. R.	Tinct. veratri viridis,	gtt. xxxvj	
	Morphiæ sulphatis,	gr. j	
	Syrupi ipecac.,	f. ʒvj.	M.

A teaspoonful every three hours, if necessary. (*Louisville Medical News*, April, 1876.)

Zinci Oxidum is strongly recommended by Dr. SYMONDS, (*British Medical News*, June 13th, 1868,) who states that, whether for abating the morbid susceptibility of the respiratory nerves which engenders the spasmodic paroxysms, or for diminishing the liability to catarrhs, he knows nothing to surpass long courses of the oxide of zinc, with or without extract of conium for the one indication, and quinine for the other. It is also recommended by Dr. NIEMEYER.

Zinci Sulphas has been employed, with the result of diminishing the frequency and force of the attacks.

Zinci Valerianas is sometimes prescribed.

EXTERNAL MEASURES.

**Cold Bathing*, employed in the intervals of the attacks of spasmodic asthma, is stated to lessen the morbid sensibility to the impression of cold atmospheric changes, and to give tone and vigor to the system. Simply *sponging the body* is preferred by some to immersion; salt (ʒj, ad aquam Oj) may be added, and a rough towel or flesh-brush employed. The best time is immediately after getting out of bed in the morning. The *shower-bath* is preferred by Sir T. WATSON.

Electricity has been employed in some cases, but Dr. HYDE SALTER condemns, altogether, its use. Galvanization of the pneumogastric has been tried by BARTHOLOW with doubtful results.

PROFESSOR MUNTER, of Russia, recommends *excursions of three or four hours daily upon salt water*, as of the greatest value.

ACUTE BRONCHITIS.

JOHN SYER BRISTOWE, M. D., LONDON.

The treatment is, (1) *Hygienic*; an equable temperature, about 65°; moist atmosphere; hot foot-baths; regulated diet. (2) *Local*; mustard plasters, or dry cupping to the upper chest; inhalations of steam, simple or medicated. (3) *Medicinal*. The expectorant or nauseating drugs, combined with the narcotics and sedatives. A combination of ipecac., squills, and small doses of opium, is very serviceable in earlier stages. Later, when the bronchial secretion becomes abundant and muco-purulent, this may be replaced with a more stimulating mixture. That recommended by Dr. STOKES is very efficacious, namely, carbonate of ammonia, senega, and opium. (F. 268.) The balsams and gums and resins are also valuable in this stage. In protracted cases and during convalescence, tonics are called for.

PROF. M. CHARTERIS, M. D., GLASGOW.

In his *Hand-book of the Practice of Medicine*, 1878, this writer states that during the early or catarrhal stage, bronchitis may be aborted by a full dose of opium in wine whey. If the fever is too intense for this, give a hot bath, followed by a weak saline mixture. As an expectorant:

229. R.	Oxymel. scillæ,	3 ss	
	Spir. æth. nitrici dule.,		
	Tinct. camph. comp.,	āā	f. ʒj
	Mist. amygdal.,		f. ʒj. M.

This amount every six hours.

If a gouty dyscrasia is present, add colchicum to this. For local applications, sinapisms, hot water and turpentine stupes, and jacket poultices of linseed meal, are valuable. The following is useful in children.

230. R.	Vini ipecac.,	f. ʒ iss	
	Syrupi simp.,	f. ʒ ss	
	Tinct. camph. comp.,	f. ʒ ij	
	Liq. ammon. citrat.,	f. ʒ ss	
	Aquam,	ad f. ʒ ij.	M.

A teaspoonful every two hours, in the febrile stage.

The *wine whey*, above referred to, may be made as follows:

234. R. Morphiae acetatis, gr. j
 Potassii acetatis, ʒ iij
 Liquoris ammonii acetatis, f. ʒ iij
 Syrupi tolutani, f. ʒ j. M.
 A dessertspoonful every third hour. A useful diaphoretic alkaline mixture.
235. R. Ammonii carbonatis, gr. xvj
 Spiritus ætheris compositi, f. ʒ iss
 Syrupi tolutani, āā f. ʒ j. M.
 Aquæ, āā f. ʒ j. M.
 A teaspoonful every two hours; a stimulating expectorant for a child a year old, affected with bronchitis of two weeks' standing.

Counter-irritation to be applied to the chest by means of weak mustard plasters (one part of mustard to four of Indian meal). Also, if the child be much debilitated, 15 gtt. of brandy every four hours. When the child is seen frequently, so that the effect may be watched, there is no better treatment than relieving the lung mechanically by emetics. Hoffman's anodyne, in the above recipe, acts as a diaphoretic and quieting agent, which latter influence would not be obtained from sweet spirits of nitre.

Where there is obvious bronchial catarrh:

236. R. Tr. verat. viridis, ʒ j
 Ammon. mur., ʒ iij
 Mist. glycyrrh. comp., ʒ iij
 Syr. pruni. virg., f. ʒ vij. M.
 S. Teaspoonful four times a day, in water.
 After taking this for a day or two, the dose ought to be reduced. Use, also, mustard to the chest.

PROF. GEORGE JOHNSON, M. D., LONDON.

Acute bronchitis is an exaggerated catarrh; the two diseases are essentially the same, and they require the same principle of treatment, only modified according to the character of the symptoms. (*British Medical Journal*, Oct. 23d, 1869.)

In the early stage of acute bronchitis, when the mucous membrane is dry and swollen, the hot-air bath or the wet packing may be employed once or oftener with advantage. Another very useful remedy in this stage is tartar emetic, in doses of $\frac{1}{6}$ of a grain, combined with liquor ammonii acetatis:

237. R. Antimonii et potassii tartratis, gr. j $\frac{1}{4}$
 Liquoris ammonii acetatis, f. ʒ ij
 Syrupi tolutani, āā f. ʒ j. M.
 Aquæ, āā f. ʒ j. M.
 A tablespoonful thrice daily. The inhalation of steam, repeated several times in the course of the day, is often very soothing and beneficial.

In the second stage, when free secretion has been established, antimony and acetate of ammonium are to be discontinued. At this period, a combination of carbonate of ammonium, with spirit of chloroform, is useful as a stimulating expectorant or antispasmodic.

- | | | | |
|---------|--|--|----|
| 238. R. | Ammonii carbonatis,
Spiritus chloroformi,
Syrupi acaciæ,
Aquam, | $\overline{3}^{\text{iss}}$
$\text{f. } \overline{3}^{\text{v}}$
$\text{f. } \overline{3}^{\text{ij}}$
ad $\text{f. } \overline{3}^{\text{iv}}$. | M. |
|---------|--|--|----|
- A dessertspoonful thrice daily.

Brandy or wine, in moderate quantities, may now also be required to sustain the strength. When, in the advanced stage, there is a profuse secretion, with copious perspiration, the ammonia mixture may be replaced by the following:

- | | | | |
|---------|---|---|----|
| 239. R. | Quiniæ sulphatis,
Zinci sulphatis,
Acidi sulphurici aromatici,
Aquæ menthæ piperitæ, | gr. xvj
$\overline{3}^{\text{j}}$
$\text{f. } \overline{3}^{\text{v}}$
$\text{f. } \overline{3}^{\text{ijss}}$. | M. |
|---------|---|---|----|
- A dessertspoonful thrice daily, in water.

This combination often checks, very rapidly, the excessive secretion from the bronchial mucous membrane. The *stimulating expectorants* are sometimes useful at this stage of the disease—senega, squill, ammoniac, and the compound tincture of benzoin. If, as sometimes happens, the stimulating expectorants suddenly check secretion, tighten the breath, and increase dyspnœa, their employment must at once be discontinued. When the secretions accumulate and threaten suffocation, the patient being blue, and cold, and drowsy, and the cough nearly or quite ceasing, an emetic of sulphate of zinc is often wonderfully efficacious in clearing the air-passages.

Dr. JOHNSON gives an especial warning with regard to *opium*. Never order an opiate for a bronchitic patient who has the slightest blueness of the lips.

DR. ROSS C. RUSS, OF OHIO.

This writer says, in the *Cincinnati Lancet and Observer* (February, 1877,) that the medicinal treatment of acute bronchial inflammation should be commenced with *wine of ipecacuanha*, given with the view to unload the bronchi of the excessive secretion, and to allay the irritability of the vagus nerve, but should only be given in the first stage. After the subsidence of the acute stage, he has witnessed the best results from the following:

240. R. Quiniæ sulphatis, ℥ iij
 Acid. phos. diluti, ss
 Syr. tolutani, ss
 Aquæ destillatæ, ℥ iij. M.
- A dessertspoonful every four hours.

The dose should be increased or diminished according to age of the patient. The object which is to be obtained, is to produce sedation over the turgid and relaxed capillaries of the mucous tissue of the bronchi, at the same time increasing the tonicity of the part.

When the disease assumes a still more grave form, solidification and ultimately softening of the lobules of the lung tissue are threatened. The following should be then given :

241. R. Spts. vini gallici, ℥ vj
 Glycerinæ, ℥ iij
 Tinct. hyoseyami, ℥ iij. M.
- Teaspoonful every four or five hours. The dose to be regulated according to age of the patient.

These medicinal agents soothe the harassing cough, aid digestion, assist assimilation, and prevent undue tissue change.

DR. MAIN, OF SCOTLAND.

With the view of promoting free secretion from the bronchial mucous membrane, Dr. MAIN has found nothing more useful, both for adults and children, than the following :

242. R. Potass. bicarb., ℥ iij
 Tr. hyosey., ℥ iij-℥ iv
 Spt. æth. nitrosi, ss
 Spt. chlorof., ℥ iij-℥ iij
 Aq., ℥ xij. ad. M.

And,

243. R. Acidi citrici, ℥ iij-℥ iij.
 Aq., ℥ vj. ad. M.

Sig.—Two tablespoonfuls of the former mixture to be taken with one of the latter during effervescence every three or four hours (for an adult).

If the secretion be profuse and the heart's action weak, he has often found the following mixture useful:

244. R. Acidi nitrici dil., ℥ iij
 Tr. bellad., ℥ iij
 Spt. chloroformi, ℥ iij
 Aq., ℥ xij. ad. M.

Sig.—Two tablespoonfuls every four hours (for an adult.)

"In dealing with children, it is well to bear in mind that, if the amount of secretion be excessive, and embarrassing the breathing, a timely simulating emetic, such as carbonate of ammonia, or mustard, often proves invaluable. This now brings us to the stage approaching convalescence, in which such drugs as quinine, vegetable bitters, steel, nux vomica, and the dilute mineral acids, all have their uses; and when convalescence has become established, I am of opinion that if we can get our patient persuaded to take cod-liver oil for a month or two, it has the effect of preventing a fresh attack."

DR. ROBERTS BARTHOLOW, PHILADELPHIA.

Muscarine ought to afford prompt relief at the beginning of an acute bronchitis, as it produces strong contraction of the pulmonary capillaries:

245. R. Muscarinæ,	gr. j	
Aquæ,	℥j.	M.
S.—A teaspoonful every three hours.		

The best results can be obtained by the use of aconite in doses of from half a drop to one drop every half hour, until an impression is made on the fever movement, and then every hour or two.

NOTES ON REMEDIES.

Ammonii Acetatis Liquor makes a useful diaphoretic alkaline vehicle for antimony, morphia, &c.

Ammonii Carbonas may be given with benefit, combined with ipecacuanha, squills when expectoration is scanty or difficult, or with tincture of lobelia if much spasm be present. Five grains, given at bed-time, when the first symptoms of a common cold express themselves, often at once subdue the disease.

Ammonii Bromidum has been recommended.

Amyl-nitrite. Dr. D. H. KITCHEN recommends for the dyspnœa of acute bronchitis, the inhalation of gtt. v-xv of this substance. (*American Journal of Insanity*, October, 1873.)

**Ammonii et Potassii Tartras* is of service in small doses (gr. $\frac{1}{15}$ - $\frac{1}{4}$), in the earliest stages of acute bronchitis in young plethoric subjects, and where much febrile action is present. When the phlegm accumulates in the bronchial tubes, an antimonial emetic will readily remove it. The nauseating and depressing effects of antimony should not, however, be long kept up in this disease, as they interfere with proper nutrition, which is of much importance in these cases.

Apomorphia.

246. R. Apomorphiæ hydrochloratis, gr. $\frac{1}{2}$
 Infusi senegæ, f. $\overline{3}$ v
 Syrupi simplicis, f. $\overline{3}$ v
 Aquæ destillatæ, f. $\overline{3}$ iv. M.

A teaspoonful every half hour in severe cough in children.

Chloroformum by inhalation is said to palliate the cough and excitement, and directly favor the resolution of the extravasated fibrin, and prevent the deposit of more. A drachm of chloroform should be inhaled in the course of fifteen or twenty minutes, and the inhalation repeated from two to twelve times in the twenty-four hours.

* *Ciimcifuga* is an excellent expectorant.

247. R. Ext. cimicifugæ fl., $\overline{3}$ ss
 Tinct. opii deod., $\overline{3}$ j
 Syr. Tolu, $\overline{3}$ x. M.

S.—A teaspoonful every four hours.

This is very useful in acute bronchitis, after the more active symptoms have subsided.

Colchicum is recommended in acute stages of *gouty bronchitis* by Dr. GREENHOW, who gives the following with great success :

248. R. Potassii iodidi, $\overline{a}\overline{a}$ gr. iv
 Ammonii carbonatis, $\overline{a}\overline{a}$ \overline{m} x
 Vini colchici, $\overline{a}\overline{a}$ \overline{m} xx
 Tincturæ scillæ, $\overline{a}\overline{a}$ q. s.
 Tincturæ hyoscyami,
 Aquæ camphoræ,

Make a draught, to be taken thrice daily.

* *Ipecacuanha* is a valuable expectorant and diaphoretic, and a useful adjunct to other remedies of the same class.

Monarda. The warm infusion of horsemint makes an excellent diaphoretic.

Oleum Ricini seems, more than any other medicine of its class, to mitigate the inflammation and promote resolution by expectoration.

Opium, or morphia, given in full doses at the very outset of a common cold, often averts the attack. But an opiate should never be given to a bronchitic patient with the slightest blueness of the lips.

Oxygen. In the dyspnœa of capillary bronchitis, the inhalation of oxygen affords relief, and has been the means of preserving life.

Phellandrium Aquaticum is considered by French physicians a specific sedative to the bronchial mucous membrane.

Pilocarpine is often serviceable, if given at the beginning.

Potassii Nitræs is frequently administered. Dr. GRAVES speaks highly of the following combination :

249. R. Potassii nitratis, 3 ij
 Antimonii et potassii tartratis, gr. j
 Tincturæ opii camphoratæ, f. 3 ss
 Misturæ amygdalæ, f. 3 xii.

A tablespoonful every hour.

Sanguinaria. Dr. BARTHOLOW speaks highly of this formula :

250. R. Tinct. sanguinariæ, 3 j
 Tinct. lobeliæ, 3 j
 Vin. ipecac, 3 ij
 Syrup. tolutan, 3 ss. M.

S.—A tablespoonful every three hours.

Senega in decoction is a valuable remedy in the treatment of acute bronchitis in the aged.

Veratrum Viride is often of service in the early stages.

Zinci Sulphas is used in the advanced stages, when there is profuse secretion.

Counter-irritants to the chest are often of service. For this purpose, dry-cupping, *hot turpentine stupes, or mild sinapisms may be employed. Turpentine embrocations are also serviceable.

Stimulants are often useful, and sometimes necessary, as in the *bronchitis of the aged*, attended with profuse secretion, threatened apnœa and a weak pulse. At the very outset of a common cold, a full supper, with a moderate alcoholic stimulant, will frequently abort the attack.

Water. The *vapor of hot water* is often of signal benefit. Wet packing is also of service in the early stages.

CHRONIC BRONCHITIS.

PROF. G. P. ANDREWS, M. D., DETROIT.

Dr. ANDREWS holds that this disease depends upon organic changes, and a cure cannot be looked for. Expectorants are useless, except during acute exacerbations. Asthmatic symptoms can be most speedily relieved by inhalations of *nitrite of amyl*. Sedatives may be called for when the cough is very harassing; but a certain amount of cough is desirable, to rid the lungs of mucus. Patients should be instructed to restrain the impulse to cough, as it is very much under the control of the will, or, at least, can be brought under by practice. When there is much emphysema and dyspnœa, the *inhalation of compressed air* has been very efficacious in giving prompt relief.

The patient should live out of doors as much as possible, in a dry,

cool, equable climate, and give careful attention to hygiene. (*Detroit Review of Medicine*, September, 1878.)

DR. GEORGE ST. GEORGE, OF DUBLIN.

This author highly recommends the following formula in chronic bronchitis. (*Dublin Journal Medical Sciences*, 1883.)

251. R.	Tr. scillæ,	℥ xv	
	Sp. æther nit.,	℥ j	
	Liq. amm. acet.,	℥ j	
	Tr. hyoscyam,	℥ ss	
	Aq. comp.,	ad. ℥ j.	M.

S.—Taken three times a day.

As soon as the urgent symptoms have subsided, substitute the following :

252. R.	Acid nitro-hydrochl. dil.,	℥ x	
	Tr. hyoscyam,		
	Tr. loricis,	āā ℥ ss	
	Inf. gent. C,	ad. ℥ j.	M.

S.—Taken three times a day.

For bronchitis caused by the inhalation of dust mixed with fine particles of flax, or other irritating substances, the following :

253. R.	Ammon. carb.,	gr. v	
	Tr. scillæ,		
	Tr. camph C.,	āā ℥ xv	
	Inf. senegæ,	ad. ℥ j.	M.

S.—Taken three times a day.

After improvement has set in, give the patient two drachms of cod liver oil after each meal, and as a tonic before meals the following :

254. R.	Acid nitro-hydroch. dil.,		
	Tr. camph. c.,	āā ℥ x	
	Vin. ipecac,	℥ vj	
	Inf. gent. co.,	ad. ℥ j.	M.

PROF. WILLIAM AITKEN, M. D., EDINBURGH.

In chronic cases of bronchitis, especially in patients who have made considerable progress in the journey of life, remedies which tend to invigorate the general system are indicated. Besides the selection of a beneficial climate and the use of nutritious, easily-digested food, stimulating embrocations are useful. The following liniment, employed at the Meath Hospital at Dublin, is highly recommended for this purpose :

255. R. Spiritus terebinthinæ, f. $\bar{3}$ iij
 Acidi acetici, f. $\bar{3}$ ss
 Vitelli ovi, j
 Aquæ rosæ, f. $\bar{3}$ iijss
 Olei limonis, f. $\bar{3}$ j. M.

To be rubbed, morning and evening, not only over the chest, before and behind, but also along the sides of the neck. It generally reddens the skin and produces small pimples.

Of the foetid gums, ammoniac in particular is a useful remedy. From the following combination (formula of Prof. EASTON, of Glasgow,) decided beneficial results are obtained:

256. R. Ammoniaci, $\bar{3}$ ij
 Acidi nitrici diluti, f. $\bar{3}$ ij
 Aquæ, f. $\bar{3}$ xij.

Two tablespoonfuls in gruel, thrice daily.

It is often advantageous to administer astringent remedies, as:

257. R. Acidi tannici, \emptyset j- $\bar{3}$ j.
 For twenty pills. One thrice daily.

Or,

258. R. Olei cubebæ, gtt. x.
 For one dose, three or four times a day, on a piece of sugar.

In the protracted bronchitic affections of the aged, diuretics are of great service. The following formulæ are recommended by our author, and by Drs. MACLACHLAN and STOKES, as well suited in a variety of cases of *senile chronic catarrh*:

259. R. Potassii nitratis, gr. xxxvj
 Tincturæ conii, f. $\bar{3}$ ss
 Spiritus ætheris nitrosi, āā f. $\bar{3}$ vj
 Oxymellis scillæ, ad f. $\bar{3}$ vj. M.
 Decoctum senegæ,

A tablespoonful thrice daily.

260. R. Potassii acetatis, $\bar{3}$ ss
 Aceti scillæ, āā f. $\bar{3}$ vj
 Spiritus ætheris nitrosi, f. $\bar{3}$ ss
 Tincturæ opii camphoratæ, f. $\bar{3}$ iijss
 Liquoris ammonii acetatis, f. $\bar{3}$ vj. M.
 Syrupi aurantii corticis,

A tablespoonful thrice daily.

DR. BEIGEL, LONDON.

261. R. Pulveris aluminis, gr. ij
 Tincturæ opii, m_x
 Aquæ destillatæ, f. $\bar{3}$ j. M.

For inhalation by means of atomizer, night and morning, in obstinate cases of chronic bronchitis.

The following inhalation also affords relief in severe cases :

262. R. *Acidi tannici*, gr. iij
Extracti hyoscyami, gr. ij
Aquæ destillatæ, f. $\frac{3}{4}$ j. M.

To be inhaled night and morning. On account of the taste, sulphate of iron (gr. iv to f. $\frac{3}{4}$ j) may be substituted for the tannic acid.

PROF. J. M. DA COSTA, M. D., PHILADELPHIA.

263. R. *Ammonii chloridi*, $\frac{3}{4}$ ij
Misturæ glycyrrhizæ compositæ, f. $\frac{3}{4}$ ij. M.
 A dessertspoonful three times a day.

Chloride of ammonium, in order to be effective, should be given in ten-grain doses. In the bronchitis of patients affected with phthisis, it may be combined as follows :

264. R. *Ammonii chloridi*, $\frac{3}{4}$ ss
Morphiæ muriatis, gr. j
Extracti pruni virginianæ fluidi, f. $\frac{3}{4}$ ij. M.
 A teaspoonful three or four times a day.

265. R. *Ammonii chloridi*, $\frac{3}{4}$ ij
Potassii chloratis, $\frac{3}{4}$ j
Tincturæ hyoscyami, f. $\frac{3}{4}$ ss
Extracti pruni virginianæ fluidi, f. $\frac{3}{4}$ ijss. M.
 A dessertspoonful thrice daily, when there is a tenacious secretion.

Chlorate of potash thins the secretion and promotes expectoration ; it is useful in both acute and chronic bronchitis.

266. R. *Potassii chloratis*, $\frac{3}{4}$ ij
Tincturæ scillæ, f. $\frac{3}{4}$ ss
Misturæ glycyrrhizæ compositæ, f. $\frac{3}{4}$ ijss. M.
 A dessertspoonful three or four times a day, when there are dry râles.

267. R. *Vini picis liquidæ*, f. $\frac{3}{4}$ ij.
 A dessertspoonful thrice daily, in cases bronchial catarrh, together with

268. R. *Plumbi acetatis*, $\frac{3}{4}$ ij
Extracti glycyrrhizæ, q. s. M.
 For twenty pills. One three times a day.

269. R. *Potassii carbonatis*, $\frac{3}{4}$ ss
Vini ipecacuanhæ, f. $\frac{3}{4}$ ij
Tincturæ opii camphoratæ,
Syrupi tolutani, āā f. $\frac{3}{4}$ iss. M.
 A teaspoonful three times a day, as an alkaline expectorant mixture, to thin the secretion.

E. HEADLAM GREENHOW, M. D., LONDON.

270. R. Vini ipecacuanhæ,
 Acidi nitro-muriatici diluti, āā ℥x
 Tincturæ hyoscyami, ℥xx
 Tincturæ gentianæ compositæ, f. ʒij M.

For one dose. To be taken in water three times a day, in chronic bronchitis.

In almost all cases of chronic bronchitis, a time arrives when expectorants cease to be useful. The expectoration has become of the nature of an habitual flow from the bronchial membrane. Treatment of a tonic character is then required. In these cases, Dr. GREENHOW has long been accustomed to prescribe, with great advantage, the *mineral acids*, especially the nitro-muriatic, in combination with a vegetable bitter, as above. In chronic cases attended by very copious expectoration, he adds to each dose of the above mixture, twenty minims of the *tincture of larch*, which has the effect not only of lessening the expectoration, and with it the cough and dyspnœa, but also, apparently, of restoring the debilitated membrane to a more healthy tone, and of rendering patients less liable to catarrhal attacks at every change of the weather or season.

JOHN FORSYTH MEIGS, M. D., PHILADELPHIA.

271. R. Acidi carbolici fluidi, ℥x-xv
 Aquæ, Oss. M.

To be placed in an ordinary inhaling bottle, and used three or four times a day, in bronchial catarrh, offensive secretion from the bronchial tubes, etc.

L. PARISEL, PARIS.

272. R. Pulveris cinchonæ flavæ,
 Sulphuris loti, āā ʒss
 Syrupi althææ, q. s. M.

This solution should be inhaled by means of an atomizer: useful both in the simple form of chronic bronchitis and in that which usually complicates the advanced stages of phthisis. It diminishes the secretion, and thereby lessens the waste of substance and the exhaustion occasioned by repeated and urgent coughing.

PROF. ALFRED STILLE, M. D., PHILADELPHIA.

273. R. Acidi carbolici fluidi, gtt. xv-xx
 Tincturæ conii, f. ʒj-ij
 Aquæ destillatæ, Oij. M.

For inhalation.

PROF. WILLIAM STOKES, M. D., DUBLIN.

274. R. Potassii iodidi, \mathfrak{z} iij
 Tincturæ opii camphoratae, f. \mathfrak{z} ss
 Decocti senegæ, f. \mathfrak{z} iv
 Syrupi tolutani, f. \mathfrak{z} iss. M.

Two teaspoonfuls a day, in chronic bronchitis.

275. R. Tincturæ opii camphoratae, f. \mathfrak{z} ss
 Ætheris,
 Tincturæ tolutani, āā f. \mathfrak{z} ij. M.

A teaspoonful in a small quantity of warm drink, to quiet the cough, in bronchitis.

276. R. Tincturæ opii camphoratae,
 Tincturæ scillæ, āā f. \mathfrak{z} iss
 Ammonii carbonatis, gr. xvij
 Decocti senegæ, f. \mathfrak{z} iv
 Syrupi tolutani, f. \mathfrak{z} ss. M.

Two or three teaspoonfuls a day, in chronic bronchitis, after the inflammatory symptoms have lost a little of their intensity.

277. R. Acidi benzoici,
 Ammoniaci, āā \mathfrak{z} ss
 Saponis, q. s.

Divide into twenty pills. From four to eight in the twenty-four hours, in chronic pulmonary catarrh. Repeated sinapisms to the chest.

JAMES TURNBULL, M. D., LIVERPOOL.

278. R. Ipecacuanhæ, \mathfrak{z} ij
 Olei olivæ, f. \mathfrak{z} ij
 Adipis, \mathfrak{z} ss. M.

This pomade is to be spread upon the skin once or twice a day, when it is desired to produce an eruption or a rubefacient effect in bronchitis.

Or:

279. R. Emetinæ, gr. xv
 Adipis, \mathfrak{z} ss. M.

The emetine is to be dissolved in a small quantity of alcohol, and the solution thus obtained is thoroughly mixed with the lard.

Rubbed on the skin for a few minutes, once or twice a day, this ointment produces a copious crop of small pustules, unattended with pain, which remain out for many days, and leave no scars. In this last point, it has an advantage over tartar emetic ointment, for which Dr. T. proposes it as a substitute.

SIR THOMAS WATSON, LONDON.

In cases of chronic bronchitis, which are especially remarkable on account of the great abundance of the bronchial secretion, our author

has derived great advantage from the administration of the *sulphate of iron* in two or three-grain doses, in the *compound infusion of rose*, thrice daily. When there is any fever present, these remedies are apt to augment it; but when the pulse is quiet and the skin cool, they are often of the greatest service.

Our author has also seen the excessive expectoration diminished, and the patient gain strength, under the use of the balsams, the *compound tincture of benzoin*, for example, a form of medicine much employed formerly, and too much neglected at present.

JOHN WILLIAMS, M. D., LONDON.

280. R.	Ammoniaci,		
	Ammonii carbonatis,	āā	gr. xv
	Ipecacuanhæ,		gr. iv
	Morphiæ muriatis,		gr. iss
	Mucilaginis acaciæ,		q. s. M.

Divide into ten pills, which coat with a solution of balsam of tolu in chloroform. One pill, morning and evening, in chronic bronchitis, especially when the secretion is thick and the expectoration difficult.

J. MILNER FOTHERGILL, M. D., LONDON.

281. R.	Sp. chloroformi,	℥xx
	Acidi hydrobromici,	ʒss
	Syr. scillæ,	ʒj
	Aquæ,	ad ʒj.

For one dose, three times a day.

The dose, of course, is reduced for children. Any other acid in this mixture is very agreeable, but the hydrobromic acid, from the effect of bromine upon reflex mechanism, allays the cough often so troublesome. It possesses much the same action as opium, without the ill effects upon the digestive organs or on the bronchial secretion.

In some recent articles Dr. FOTHERGILL dwells at some length on the great value of *strychnia* as an expectorant in bronchitis. By its action on the respiratory centre, it proves useful when increase of respiratory power is needed for the expulsion of mucus gathered in the air-tubes. He gives it either alone or in combination with the ordinary cough mixture. On other expectorants, Dr. F. gives these directions: As long as the skin is dry, and the bronchial lining membrane tumid, and secretion arrested, ipecacuanha with acetate of ammonia is indicated; or, a little antimony may be added with advantage. When the skin is once thrown into action, and the bronchial secretion also estab-

lished, then acids with syrup of squills are suitable measures. But it is not a successful plan to administer squill with acids until the skin is moist. When there is a tendency to the free action of the skin, this latter combination, in full doses, is a useful plan of treatment. Neither is the union of carbonate of ammonia and senega, in severe cases, indicated until the secretions alike of the skin and bronchial lining membrane are thoroughly established.

BELLEVUE HOSPITAL, NEW YORK CITY.

282. R.	Ammon. carb.,	gr. xxxij	
	Ext. fl. senegæ,		
	Ext. fl. scillæ	ââ	f. 3j
	Tr. opii camph.,		f. 3vj
	Aquæ,		f. 3ss
	Syrup. tolut.,	q. s. ad	f. 3iv. M.

Teaspoonful at a dose.

NOTES ON REMEDIES.

Acidum Carbolicum, by atomized inhalation, is of service when the bronchial secretions acquire a repulsive fœtor, and indeed quite generally.

Allium. After the febrile stage has passed, syrup of garlic, or the domestic onion syrup, is a very useful expectorant, particularly for children.

Alumen. Dr. F. L. ANDREWS, of London, recommends alum in acute as well as in chronic bronchitis, both in children and adults, when the expectoration is ropy. To children of three years, he gives one grain in solution, every five or six hours; to adults, six to eight grains, at like intervals.

Ammonii Carbonas is an excellent stimulant expectorant, of benefit in the advanced stages of bronchitis, when there is no febrile action present. The following formula may be employed :

283. R.	Ammonii carbonatis,	gr. iv	
	Spiritûs chloroformi,	℥xx	
	Aquæ camphoræ,	f. 3x.	M.

To be taken every three or four hours.

Infusion of senega may often replace, with advantage, the camphor water in this recipe. It is of especial benefit in the *asthenic bronchitis of old age*.

**Ammonii Chloridum* is a most valuable remedy in chronic bronchitis, but must be given in ten-grain doses in order to be effective. The following is a useful formula :

284. R.	Ammonii chloridi,	3ij
	Spiritûs ætheris compositi,	f. 3ij
	Extracti glycyrrhizæ,	3j
	Decoctum senegæ,	ad f. 3vj.

Tablespoonful every two or three hours. Ipecacuanha or squill may be added, if indicated.

Angelica is useful in the chronic bronchial catarrh of aged and feeble persons, under circumstances when senega is usually prescribed.

Apomorphia. Dr. WERTNER, of Vienna, has used this substance internally as an expectorant with the most pleasing success. His prescription is :

285. R. Apomorphiæ, gr. i-iii
Aquæ destillatæ, f. ʒ xij. M.

Tablespoonful every two hours for an adult. It is an excellent cough sedative, especially useful in bronchial catarrh in the exudation stage. (*Dobell's Reports*, 1876.)

Arsenicum is recommended by Dr. THOROWGOOD, (*Lancet*, November 13th, 1869,) in chronic bronchitis where copious expectoration and much emaciation are present. At times, he finds it to act quite like a specific in improving the breathing. He gives ℥ij of Fowler's solution in alkaline infusion of columbo thrice daily.

Assafetida is of service when the cough is spasmodic, and the expectoration difficult, on account of general debility. It is therefore, often useful in the chronic bronchitis of old persons.

Balsamum Peruvianum. The balsams, both of Peru and of Tolu, are useful stimulant expectorants, and vehicles for squill and other remedies of the same kind, in chronic bronchitis. The vapor, obtained by placing them upon a piece of hot iron or hot coals, is also of benefit.

Benzoinum. The compound tincture of benzoin is a useful stimulant expectorant; the tincture of benzoin may often be employed with benefit, by means of a steam atomizer. Benzoin must be used only in cases not injuriously influenced by stimulation.

Calcis Phosphas, in doses of gr. i-ij several times a day, is recommended by Dr. RINGER as of great value in checking profuse discharges in chronic bronchitis.

Cerii Oxalas. Dr. THOMAS CLARK (*Practitioner*, April, 1878.) considers this drug to be purely sedative, and therefore a great desideratum in treatment of lung diseases, inasmuch as it does not disturb the digestive tract—the only unpleasant subjective feature of its use being occasional dryness of the mouth. In gr. v doses he has found that it will relieve many harassing coughs, irrespective of the pathological conditions which cause them. Dyspnœa is usually relieved at the same time. He claims that relief for a period of twenty-four hours often follows a single dose taken before rising in the morning.

Cicutin. The hydrobromate of *cicutin* has been found an efficient sedative in cough by M. DELIOUX. (*Dobell's Reports*, 1877.)

Cimicifuga is especially useful in the chronic bronchitis and coughs of old age.

Cinchona. The acidulated cold infusion of bark is very useful when there is rapid wasting of the system, under the influence of profuse expectoration and night sweats.

Conium, in the form of the compound pill (gr. v-x), is frequently useful. Squill increases its efficacy. Vapor coniæ may be inhaled with benefit.

Copaiba often restrains and modifies the bronchial secretion in chronic bronchitis attended with profuse expectoration, when there is no fever and not much vascular irritability.

Creasotum is sometimes given by inhalation and by the stomach, in the bronchorrhœa of old persons.

Creta Præparata is recommended by Dr. COPLAND in chronic bronchitis in the advanced stages, particularly when colliquative sweats and diarrhœa are present. He employs the following mixture, which, even when the bowels are regular, is by no means productive of constipation :

286. R.	Misturæ cretæ,	f. $\frac{3}{4}$ vjss	
	Vini ipecacuanhæ,	f. $\frac{3}{4}$ iss	
	Tincturæ opii camphoratae,	f. $\frac{3}{4}$ vj	
	Syrupi tolutani,	f. $\frac{3}{4}$ iij.	M.

Tablespoonful three or four times daily.

Cubeba, in small and often-repeated doses, is of service in the chronic bronchitis of old age, attended with profuse secretion and much debility of constitution.

Digitalis. The tincture is exceedingly useful in the bronchitis of elderly persons with deficient circulation and dyspnœa. It may be combined with ammonia.

Yerba Santa. The leaves of the *Erioduction Californicum*, known by this name, have been long used as a popular expectorant on the Pacific coast. A fluid extract is now manufactured by Parke, Davis & Co., of Detroit. Dose, f. $\frac{3}{4}$ j, as required. An appropriate combination is :

287. R.	Fld. extr. yerbæ santæ,		
	Syr. pruni virgin.,	ââ	f. $\frac{3}{4}$ ij
	Glycerinæ,		f. $\frac{3}{4}$ j.

A dessertspoonful thrice daily.

It is a tonic expectorant, agreeable to the palate, and without nauseating properties.

Ferri Mistura Aromatica is highly praised as calculated to improve the general health and check superabundant secretion, by Dr. GRAVES. He orders:

288. R.	Misturæ ferri aromaticæ.	f. $\frac{3}{4}$ iss-ijj	
	Aquæ menthæ piperitæ,	f. $\frac{3}{4}$ vj.	M.

Tablespoonful thrice daily.

BARTHOLOW prefers the *Misturæ Ferri Compositæ*.

Gelsemium is an efficient sedative when the cough is of the habitual nervous variety.

Helenin in doses of gr. $\frac{1}{5}$ thrice daily in pill has been found by Dr. F. VALENZUELA very efficient to allay cough. (*Sig. Med.*, October, 1883.)

* *Helenine* as an expectorant, is highly recommended by Dr. F. VALENZUELA, of Spain. The dose is five drops of the tincture, three times a day.

Ipecacuanha is a useful expectorant, usually given combined with other remedies. Dr. WILLIAM MURRELL, of London, uses ipecacuanha spray in

winter cough. The spray has proved most successful in those cases of winter cough in which the dyspnoea is the prominent symptom. The ipecacuanha wine was used either pure or variously diluted, the most usual strength being one part of the wine to one or two of water. The use of undiluted wine occasionally produces dryness of the throat and nausea, the latter symptom most frequently occurring in those cases in which the patient is not careful to expectorate the mixture of saliva and condensed spray which accumulates in the mouth. The quantity of wine used at each inhalation varied from a half to two drachms.

* *Koumiss*. BARTHOLOW says that koumiss possesses great value in the treatment of chronic bronchitis.

Larix Europæa is recommended by Dr. GREENHOW, to check profuse passive expectoration in chronic bronchitis.

Lobelia is a serviceable sedative, allaying spasm and facilitating expectoration. It is usually given in combination with other remedies.

Marrubium. The expressed juice of the fresh plant, taken with milk, has been strongly recommended.

Myrrha, when there is no fever or irritability present, is a valuable medicine in long-standing chronic bronchitis, particularly after a course of expectorants.

Naphthaline, $C_{10}H_8$, reduced to powder and united with a little gum acacia, in gr. j-ij, or advantageously combined with a small quantity of codeia, not only slightly lessens expectoration, but alleviates cough without interfering with the other secretions.

Oenothera Biennis. Dr. F. H. DAVIS, of Chicago, (*Half-Yearly Compendium*, July, 1879,) considers this an exceedingly valuable drug in cases where much local catarrhal inflammation is present. His medication is :

289. R.	Acid carbolic. cryst.,	gr. vj
	Glycerinæ,	3ss
	Ext. œnothæræ bi. fl.,	3iiss
	Aquæ,	3ij.

S.—Teaspoonful before each meal.

In addition, the following :

290. R.	Pil. hydrargyri,	gr. x
	Pulv. aloes,	gr. v
	Zinc. oxidî,	3j.

S.—One pill to be taken at bedtime, or after breakfast and at bedtime.

Opium, though often of much service in allaying bronchitic cough, may prove dangerous if there is much accumulation in the bronchi.

Penthorum Sedoides. This plant, *vulgó*, the Virginia stone-crop, has been highly praised in bronchial catarrh, by Dr. G. L. TINKER. (*Half-Yearly Compendium*, January, 1879.)

Pilocarpin. In chronic dry catarrh, as well as in acute bronchitis, Dr. O. KAH-

LER, of Prague, has witnessed much benefit from this alkaloid. (*Centralblatt Med. Wis.*, April, 1878.)

Pix Liquida is of some value in bronchial catarrh. The following formula was recommended by the late Prof. DUNGLISON :

291. R.	<i>Picis liquidæ,</i>	f. $\frac{3}{4}$ j
	<i>Aquæ destillatæ,</i>	Oij.

Digest for eight days, and decant. Dose, from eight to twelve table-spoonfuls, daily, in milk.

Dr. SYDNEY SNYDER gives tar in two-grain doses, every three or four hours, in winter cough.

**Plumbi Acetas* frequently proves useful when there is profuse mucous secretion. (F. 255.)

Potassii Chloras is recommended by Prof. DA COSTA in both acute and chronic bronchitis, to thin the secretion and promote expectoration. (F. 252, 253.) FLINT gives $\frac{3}{4}$ ss of the saturated solution, two or three times a day.

**Potassæ Liquor*, \mathfrak{m} x-xv, added to an ordinary expectorant mixture, is often attended with good results when the sputa is scanty, thick, and viscid.

***Potassii Iodidum* acts beneficially in the advanced stages of chronic bronchitis, in restraining inflammatory action and promoting absorption. (F. 261.) FLINT says of it, that it exerts more curative influence over the disease than any other drug.

Scilla is of service after all active inflammation has subsided, in mild cases unattended with purulent expectoration. It is best combined with an alkali and a small quantity of opium. (F. 247.)

**Senega* is a highly serviceable remedy, particularly when combined with carbonate of ammonium. (F. 267.)

**Sodii Hypophosphis*, or *potassii hypophosphis*, is useful in very chronic cases, with thick, fœtid expectoration. Camphor water is a flood vehicle.

Sodii Hyposulphite. Dr. LEVIEZ has collected (*Jour. de Med. Prat.*, 1883,) a considerable number of observations upon this drug, and confirms the favorable opinion of M. Lancereaux, who cured seventeen cases by this remedy. One drachm of the salt must be given daily during five or six weeks. The good effect is generally visible after about a week.

Stramonium is recommended, combined with expectorants, by Dr. HEADLAND, in the form of the tincture (\mathfrak{m} x), in the chronic bronchitis of the aged, where there is great dyspnoea, accompanied by a dry state of the tubes.

Strychnia, or *nux vomica*, in small doses, with iron, is advocated by Dr. THOROWGOOD, especially when there is constant dyspnoea with prolonged expiration. (*Lancet*, November 13th, 1869.)

Zinci Sulphas is recommended by Dr. BARLOW. He employs the following combination for moderating the cough and expectorations :

292 R.	<i>Zinci sulphatis,</i>	gr. j	
	<i>Extracti conii,</i>	gr. iv.	M.

One pill thrice daily.

Emetics are recommended in chronic bronchitis, by Dr. GRAVES, when there is no fever, no remarkable dyspnoea, nor acceleration of the pulse, but when the secretion is very copious. In such cases, an emetic every night, for two or three nights, is productive of great advantage, freeing the stomach and lungs from a large amount of mucus, rendering the expectoration easier, clearing the tongue, and improving the appetite.

Inhalations are frequently employed in the treatment of chronic bronchitis. The following are the principal drugs used: *Alum*; *Balsams of Peru and Tolu*; *Carbolic Acid*; *Chlorine*; *Chloroform*, a few whiffs, $\mathfrak{m}\mathfrak{v}\mathfrak{j}$ -l, often of signal, but only temporary benefit; *Conium Vapor*; *Creasote*; *Iodine*; *Olibanum Fumigations*, the gum rosin placed on burning coals; *Tar Vapor*, generated by heating tar over a spirit lamp, with the addition of a small quantity of potash, to neutralize the irritant fumes; *Nitre Paper Fumes*; *Sulphurous Acid Spray or Fumigation*; *Tannic Acid*; *Turpentine*; *Hot-Water Vapor*; *Sulphate of Zinc*.

EXTERNAL REMEDIES.

Counter-Irritants. The following combination of *aqua ammoniac*, applied to the chest, is an efficient counter-irritant:

293. R.	<i>Aquæ ammoniac</i> ,	f. $\mathfrak{z}\mathfrak{ss}$ -j
	<i>Olei amygdala.</i> ,	f. $\mathfrak{z}\mathfrak{ss}$
	<i>Olei rosmarini</i> ,	f. $\mathfrak{z}\mathfrak{j}$
	<i>Aquæ</i> ,	f. $\mathfrak{z}\mathfrak{i}\mathfrak{j}$.

Tartar-Emetic ointment is occasionally employed as an irritant to the chest, but other irritants are generally preferred. *Croton Oil* liniment is sometimes very useful. *Turpentine* liniments are highly serviceable. The following combination, supposed to be an imitation of that used by ST. JOHN LONG, is recommended by Dr. GRAVES:

294. R.	<i>Olei terebinthinæ</i> ,	f. $\mathfrak{z}\mathfrak{i}\mathfrak{j}$
	<i>Acidi acetici</i> ,	f. $\mathfrak{z}\mathfrak{ss}$
	<i>Ovi vitelli</i> ,	j.
	<i>Olei limonis</i> ,	f. $\mathfrak{z}\mathfrak{j}$
	<i>Aquæ rosmarinæ</i> ,	f. $\mathfrak{z}\mathfrak{i}\mathfrak{j}\mathfrak{ss}$.

To be well rubbed in over the chest, the nape of the neck, the epigastrium, and the course of the cervico-spinal and pneumogastric nerves generally. *Sinapisms* to the chest are sometimes of service.

Electricity. A fine primary induction current, of low intensity, passed through the chest from the back and neck, stimulating the sympathetic and spinal nerves, has been found productive of great benefit.

Nitro-muriatic Acid lotions sponged over the chest are recommended for checking profuse expectoration, and mitigating the severity of the symptoms.

Plasters. A *Burgundy pitch* plaster often proves of service as a protective against atmospheric changes, and as a rubefacient. An *opiate* plaster is sometimes of benefit.

**Poultices.* Large, light, soft linseed-meal poultices, made to cover the whole of the chest, often afford great relief in chronic bronchitis. If it be desirable to stimulate the skin, *Mustard* (one part to four) is applicable.

CATARRH, CORYZA, AND POST-NASAL CATARRH.

The inflammation of the outer air passages commonly called "a cold," is known as a *coryza*, or "cold in the head," when it begins in the mucus membrane of the nose, and a *catarrh* when it attacks first the membrane of the mouth, fauces and upper air passages.

THE ABORTIVE TREATMENT OF A COLD.

A successful method is that used by

DR. HORACE DOBELL, LONDON.

His plan is as follows :

1. Order:

295. R.	Ammonii carbonatis, Morphiæ sulphatis, Misturæ amygdalæ,	℞j gr. $\frac{1}{6}$ f. $\frac{3}{4}$ iv.	M.
---------	--	---	----

Two tablespoonfuls every three hours.

2. At night give a tablespoonful of spirit of mindererus in a tumbler of cold water, after the patient has got into bed and been covered up with several extra blankets ; cold water to be drunk freely during the night should the patient be thirsty.

3. In the morning, the extra blankets should be removed, so as to allow the skin to cool down before getting up.

4. Let him get up as usual, and take his usual diet, but continue the ammonia and morphia mixture every five hours.

5. At bedtime, the second night, give a compound colocynth pill. No more than twelve doses of the mixture, from the first to the last, need be taken, as a rule ; but should the catarrh seem disposed to come back after leaving off the medicine for a day, another six doses may be taken, and another pill. During the treatment the patient should live a little better than usual.

As everything depends upon the promptitude of the treatment, persons who are subject to catarrh, especially if it inclines to the influenza character, should be provided with a prescription for the medicine, and full instructions how to manage themselves when a cold sets in.

An addition to this plan is needed in persons whose colds seize at

once upon the bronchial mucous membrane. Besides the manner of proceeding just described, they should use the following inhalation :

296. R.	Extracti conii,	gr. x	
	Tincturæ benzoini,	f. ʒj	
	Spiritus ammoniæ aromatici,	f. ʒss	
	Aquæ (170° F.),	Oj.	M.

Inhale the steam for fifteen minutes at bedtime each night ; put a mustard plaster on the front of the chest one night, and between the shoulders the next ; and, unless the weather is warm, wear a respirator out-doors until all signs of the cold have disappeared.

DR. C. J. B. WILLIAMS, LONDON,

has found the "dry plan" eminently successful ; and it does not oblige the patient to leave his business. His directions are :

1. Begin with a sharp aperient in the solid form of pills, swallowing them with as little water as possible.

2. The food should be rather less in quantity and more digestible than usual, and at first should be dry ; later on, the moister forms of food are more easily swallowed and digested.

3. No fluid should be taken whatever.

4. As much exercise as possible should be taken, in warm clothing, so as to promote the action of the skin and bowels.

Twenty-four hours of this treatment will give immense relief ; thirty-six to forty-eight hours will effect a cure ; and rarely even a trace of bronchial irritation remains.

T. J. MACLAGAN, M. D., LONDON.

This writer (*Practitioner*, November, 1877,) prefers, as an abortant in commencing cold, a full dose of *quinine*, (gr. x,) or several doses of *salicin* (gr. xv-xx, every one or two hours, until ʒii-ij are taken).

As a prophylactic for those cases who are constantly "catching cold," a writer in the *Practitioner*, February, 1875, recommends filling the ears with *cotton-wool*. Not only does this prevent them, but both sore throats and colds are much benefited during their continuance by this simple device.

Dr. FERRIER advocates the use of this snuff:

297. R.	Morph. sulphatis,	gr. j	
	Bismuth. subnitrat.,	ʒ ij	
	Pulv. acaciæ,	ʒj.	M.

Use by insufflation at the commencement of a catarrh.

DR. J. S. PROUT, NEW YORK,

advocates the use of the tincture of the *chloride of iron*, ℥xx-xxx, taken directly after the cold is caught. The improvement should be decided in half an hour; if not, the dose ought to be repeated two or three times. A convenient vehicle for the iron is glycerine. (*Medical Record*, January, 1874.)

One full dose of *opium* or *morphia* will often succeed in promptly dissipating the symptoms. So also will twenty or thirty drops of *tincture of belladonna*.

The following powder is highly recommended by the editor of the *Archives of Neurology*, 1877 :

298. R.	Camphoræ,	ʒv	
	Ætheris sulphurici,	q. s.	
	Dissolve to the consistence of cream, and add :		
	Ammonii carbonatis,	ʒiv	
	Pulveris opii,	ʒj.	M.
	Keep in a tightly-corked bottle.		

The dose is regulated by the opium, and ranges from gr. iij-xv. This much may be taken in a little water just before retiring, or at any hour of the day when there is a suspicion of having caught cold; and the dose may be repeated if necessary. The taste is not disagreeable, and the effect is "singularly and inexplicably efficacious." It is well and safe to keep it on hand and take about as much as one can lay on the finger-nail, whenever one becomes badly chilled.

DR. BRAND, VIENNA.

299. R.	Acidi carbolici,		
	Liq. ammonii fortis,	āā	f.ʒv
	Alcoholis,		f.ʒij.
	Keep in a stoppered dark-glass bottle.		M.

When a catarrh is commencing, a few drops are placed on three or four layers of blotting-paper; the patient holding this in his hand, and closing his eyes, inhales deeply from it as long as any smell is perceptible. The remedy should be applied every two hours.

PROF. M. CHARTERIS, M. D., GLASGOW.

300. R.	Liq. morphiæ (B. Ph.),	℥x	
	Vin. antimon.,	℥viij.	M.

This much at a dose, given twice at an interval of three hours, will effectually stop a commencing cold.

In a note to PHILLIPS' *Materia Medica*, Dr. H. G. PIFFARD, of New York, says that Mij-v of the imported tincture of eye-bright, *Euphrasia officinalis*, repeated every two or three hours, will often abort acute nasal catarrh.

Dr. J. SOLIS COHEN has found that the inhalation of chloroform to the induction of anæsthesia, will often abort acute coryza attended by intense pain and tension in the frontal region, and obstruction to breathing.

GENERAL TREATMENT OF CATARRH.

PROF. FELIX VON NIEMEYER, M. D., TUBINGEN.

In the treatment of *acute* nasal catarrh, the production of active diaphoresis is the only procedure worthy of confidence. A Russian bath should be advised where practicable. In most cases, all that is necessary is to direct the patient to confine himself to his room for a few days; to keep the head and feet warm; to swallow some hot drinks frequently during the day; to use linen pocket-handkerchiefs, not silken or cotton, and to change them frequently; and to smear the upper lip with salve, to protect it from the acrid secretion. In the latter stages, a long walk in the open air, or even an occasional pinch of snuff, hastens the cure. In infants at the breast, the nostrils should be cleared by syringing them with warm water, and they should be fed by the spoon or bottle so long as the obstacle to sucking continues.

In the treatment of *chronic* nasal catarrh, the cachexia, whatever it may be, should be attended to. Local remedies are also of the utmost importance. The most effective is penciling the swollen mucous membrane with the following solution:

301. R.	Argenti nitratis,	gr. i-xxx	
	Aquæ destillatæ,	f. ʒj.	M.

Or cauterizing with the lunar caustic in substance. The following, to be used as a snuff, is in great repute:

302. R.	Hydrargyri chloridi mitis,		
	Hydrargyri oxidi rubri,	āā	gr. xij
	Sacchari,		ʒss.
			M.

For one powder.

When the discharge is offensive, and the above treatment fails, the following may be tried:

303. R. Iodinii, gr. ij-iv
 Potassii iodidi, gr. iv-vij
 Aquæ, f. ʒvj. M.

To be used as an injection into the nostrils.

To diminish the irritation of the mucous membrane, a variety of *snuffs* have been suggested. A simple one, and yet effective, is that recommended by Dr. BROWN-SÉQUARD. It is *pulverized white sugar*, used *ad libitum*.

Dr. E. C. MANN, of New York, modifies this to :

304. R. Pulv. sacchari albi,
 Pulv. camphoræ, āā q. s. M.
 For a snuff.

Prof. TROUSSEAU recommends, in chronic coryza, a snuff of finely pulverized *tannin*. Prof. STILLÉ also recommends a snuff containing tannic acid and white sugar, reduced to a very fine powder and forcibly inspired twice a day or oftener, according to the exigencies of the case ; or, a glycerine of tannin, applied with a brush, after the parts have been cleaned by the usual *douche* or otherwise.

DR. ROBERTS BARTHOLOW, PHILADELPHIA.

A solution of quinine will sometimes, when applied to the nares, arrest an attack of summer catarrh. The preparation most suitable for this purpose is an aqueous solution of the hydrochlorate (gr. iv-vij to ʒj). This should be applied by a large camel's hair brush or spray-producer, to the nares and fauces. The utility of quinine in this peculiar disease will be determined by the extent to which the local trouble has proceeded ; it can be useful only when the irritation is confined to the nares and fauces. An attack of acute catarrh may often be entirely aborted by a full dose (ten grains) of quinine and morphine (one-half a grain) if given at the incipiency of the attack.

A variety of *inhalations* are applicable :

L. WALDENBERG, M. D., BERLIN.

305. R. Ammonii chloridi, gr. iv
 Aquæ destillatæ, f. ʒj. M.

For inhalation by atomizer, in that form of coryza known as dry snuffles. When the mouth is closed and the proper inclination given to the head, the nebulized spray can readily be inspired into the nostrils. Solutions of common salt may also be employed in the same manner.

at once destroy the stench from the nose, and to substitute for it the odor of the oil of thyme, which is rather pleasant than otherwise. The patient is directed to cleanse his nose with a solution of two drachms of bicarbonate of soda to one quart of tepid water morning and evening, and to throw a spray of the Listerine solution into the nostrils after the cleansing. The oil of thyme and of eucalyptus, besides being disinfectants, have also a stimulating effect upon the mucous membrane, and thus aid in the second portion of the treatment.

The stimulation of the serous glands to a normal action may be brought about by a variety of remedies, such as astringents in various strengths; but the insufflation into the anterior nasal cavities of finely powdered *nitrate of silver*, diluted with starch powder, has given the best results. Where there is complete absence of the lower turbinated bones, the introduction of a wad of absorbent cotton, which is to remain until washed out, and then be re-introduced by the patient himself, often aids in the stimulation by continually irritating the mucous surface with which it is in contact. Next in effectiveness to the silver powders, is a weak solution of *ferric alum* in the form of a spray thrown into the nasal cavities, and the natural iron water of Cresson springs is peculiarly adapted for these cases.

A treatment such as this carried out for several months has given most satisfactory results.

The internal administration of small doses of bromide and iodide of potassium in combination, on account of their influence upon the nasal mucous membrane, will greatly aid the local treatment. At the same time we must look to the general health of the patient, and administer tonics when their use is indicated.

DR. S. BARUCH.

This author (*Proceedings South Carolina Medical Association*, 1883,) very wisely urges that every diseased part must be seen by the eye of the attendant before medication, and again examined to ascertain if the remedies have been applied to the entire surface. The anterior nares and a small portion of the posterior surfaces can be best reached by the atomizer; the posterior nares and vault of the pharynx by the post-nasal syringe.

He uses the following formula:

307. R.	Acid carbolie,	gr. j	
	Sodii bicarbonat.,		
	Sodii biborat.,	āā	gr. ij
	Glycerinæ,		ss
	Aquæ,		℥j.
			M.

This is used to thoroughly cleanse the parts before applying the curative medication. *Not until every diseased and healthy part presents itself clearly*, should medication be thought of. He then recommends finely-powdered arrow root as a vehicle, and nitrate of silver, zinc sulphate, tannin, alum, and bismuth as the medicinal agents. By dusting the parts well, the powder mixes with the mucus and forms a firmly adhering plasma, which slowly dissolves in the secretion, thus permitting a thorough action on the subjacent tissue. By attention to the thorough cleansing of the parts, many obstinate cases of catarrh can be made to yield to medication. Many cases of catarrh are maintained by the habit of nose-cleaning, now so much in vogue. The delicate lining of the nasal cavities will not brook the frequent scouring.

THE RADICAL TREATMENT OF NASAL CATARRH.

DR. P. M. RIXEY, U. S. NAVY.

According to this writer, (*Proceedings Naval Medical Society*, vol. 1,) the radical treatment consists in the use of either chemical or surgical measures for the partial or entire removal of hypertrophy. The first consists in the use of various caustics, such as acetic acid, London paste, nitric acid or the nitrate of silver. Nitric acid must be used very carefully, owing to the difficulty in controlling its action. Each application removes only a superficial slough, consequently numerous applications have to be made at intervals of from five to ten days, and in the meantime sprays or douches have to be used daily to keep the parts clean. When the hypertrophy is great, surgical treatment must be resorted to. This consists in the use of forceps, curette, snares, actual or electric cautery, and electrolysis. Jarvis' snare works admirably when the enlargement is considerable, and the galvano-cautery is very efficient.

DR. BEVERLY ROBINSON, NEW YORK.

According to this writer, (*New York Medical Journal*, November, 1875,) no method of treatment in post-nasal catarrh is curative in every case, and this results, in great part, from the fact that it is

impossible to apply remedies to every portion of the diseased membrane. The results of his observation are to the effect that constitutional treatment proves a most important element in the case. If the patient is anæmic, *iron*, *quinine*, and *cod-liver oil* prove of decided advantage; and if there is malaria present, it has to be combated by a sufficient amount of quinine. Of the agents which have a local effect on the naso-pharyngeal and pharyngeal mucous membranes, he has experimented with balsam-copaiba, cubebs, guaiacum, ammoniacum, and sulphur, and is led to the conclusion that the three most important are *cubebs*, *sulphur*, and *copaiba*, and of these, he is in favor of *cubebs*. At first, he used the oleo-resin of cubebs, but found that it did not possess the virtues of cubebs itself. A form that has proved satisfactory, is :

308. R. Cubebæ,	℥ ij	
Syr. aurantij,	f. ℥ iij	
Aquæ menth. piper.,	f. ℥ viij.	M.
Half a teaspoonful every three hours.		

It is necessary to continue the cubebs for one or two months, and by its use the sensation of stuffiness in the head and hawking mucus may be relieved.

Diarrhoea or an herpetic eruption may appear during the use of the cubebs, and, if so, it is judicious to interrupt its administration for a few days.

The *nasal douche* is frequently used in this affection, often, it is believed, to the detriment of the patient. The following simple douche is recommended by

DR. L. MARIEOSKY, OF RUSSIA.

A thin tube of India-rubber, about twenty-eight inches long, is connected on one end with the narrow part of a glass funnel, and the other end with a wooden or bone nose-piece fitting into one of the nostrils; the solution is poured into the funnel held above the level of the patient, and thus conducted to the nostrils; the patient is told to open his mouth and slowly inspire, when the fluid will flow out of the other nostril, and thus wash out all the parts within the nasal cavities. Dr. M. reports numerous successful cases of acute and chronic catarrhs by this means. He uses a continuous current of weak solutions of permanganate of potash, salicylic acid, or sulpho-carbolate of zinc. (*Dobell's Reports*, 1876.)

DR. J. W. BARD, OF OHIO.

This practitioner has had very good success in recurrent post-nasal catarrh from the following:

309. R. Pulv. cubebæ, $\bar{3}j$
Flor. anthemis nob., $\bar{3}$ ss. M.

Mix thoroughly. To be put into a common tobacco pipe and smoked, puffing the smoke through the nostrils. It should be discontinued if a fresh attack of nasal catarrh set in, and resumed as soon as the inflammatory symptoms commence to subside. (*Medical and Surgical Reporter*, April 6th, 1878.)

MR. LENNOX BROWNE, LONDON,

mentions *iodoform* as a therapeutical agent, and recommends it as a local application in subacute and chronic inflammation of the nasopharynx. He applies it in two ways; first, as an ethereal solution (one to ten or twelve), with a brush, sponge, or with cotton-wool, the throat being first treated, and then the remedy applied to the nasal passages. Only momentary discomfort is produced during the evaporation of the ether. After the application, which should be repeated twice or thrice a week, the mucous membrane appears paler and is covered with a thin film of iodoform. After about six repetitions, the swelling of the membrane will have been much diminished. The second method, consists in the admixture of the remedy with unguentum petrolei:

310. R. Unguenti petrolei, $\bar{3}j$
Iodoformi, gr. v-vij
Æth. sulph., q. s. M.

Dissolve the iodoform in the ether and add to the unguent. Apply night and morning with a brush.

NOTES ON REMEDIES.

Acetum. Bathing the chest every morning with vinegar and water, often diminishes the liability to an attack in persons subject to repeated catarrhs.

Aconitum is highly recommended by Dr. RINGER, at the outset of febrile catarrh:

311. R. Tincturæ aconiti radiceis, $m\text{viiij}-xvj$
Aquæ, f. $\bar{3}ij$. M.

A teaspoonful every ten or fifteen minutes, for two hours, and afterwards repeated every hour. If there be much prostration, and the pulse be feeble and weak, the dose should be diminished. So soon as the sensation of "pins and needles" is produced, the aconite, which has then been pushed to its full extent, should be stopped.

Ammoniacum is a useful remedy in the chronic catarrh of old age.

312. R. Ammoniaci, ℥ ij
 Acidi nitrici diluti, f. ℥ ij
 Misturæ acaciæ, f. ℥ viij. M.

Two tablespoonfuls, in any bland fluid, every two hours.

Ammonii Carbonas is a very useful remedy in this affection. For Dr. DOBELL's method of employing it to stop a cold, see above.

Antimonii et Potassii Tart. BARTHOLOW recommends it in doses of from $\frac{1}{12}$ to $\frac{1}{10}$ grain in the first stage, when the mucous membrane is red and swollen.

Arsenicum is of benefit, when the affection is of a local character, and when there is an absence of inflammatory action and febrile disturbance, which states contra-indicate its use.

Atropine Sulphate at the outset of a coryza acts promptly and efficiently. Dose, one milligramme or about four drops of the officinal *liq. atropiæ*. After the coryza is established and in chronic cases, it does no good.

Bismuthi Oxidum. The London *Med. Record*, 1880, recommended a powder of oxide of bismuth, powdered gum acacia, and a small quantity of muriate of morphia. It should be well mixed, and then, if used as a snuff in severe coryza or post-nasal catarrh, it acts in a most charming manner. Cases of great severity and long duration have yielded to its influence after three or four days.

Cannabis Indica is highly praised by Dr. WARING CURRAN (*Medical Press*, September 9, 1868,) as a sedative and expectorant. He orders the following mixture :

313. R. Tincturæ cannabis indicæ, f. ℥ ij
 Pulveris tragacanthæ compositi, ℥ j
 Ætheris chlorici, f. ℥ iss
 Aquæ anisi, f. ℥ vj. M.

Two tablespoonfuls every second hour.

Cheken. This new drug, derived from the *Myrtus cheken*, of Chili, has been introduced by Messrs. Parke, Davis & Co., Detroit. It is a tonic expectorant, with diuretic qualities. Dr. MURRELL employed it in a number of cases in the London Hospital for Diseases of the Chest, and reports that in all cases the patients derived some benefit, and in most instances the relief was very marked, the cough lessening promptly. The dose of the fluid extract is f.℥ij-iv.

Hydrargyrum ammoniatum. In chronic coryza with irritating secretions, Dr. GUERSANT, of Paris, recommends :

314. R. Hydrargyri ammoniati, ℥ iss
 Pulveris althææ, ℥ j. M.

Use as a snuff.

Ipecacuanha. A full dose of Dover's powder (gr. x) taken at bedtime, and at the very outset of the attack, often arrests it.

**Opium*. A full opiate at bedtime, followed by a laxative in the morning, will, if given at the outset, frequently cut short a common attack of catarrh.

Pulsatilla is adapted to the treatment of acute inflammation of the nasal mucous membrane, but its use is contra-indicated when there is gastro-intestinal disturbance.

Potassii Chloras. Dr. L. SEDGWICK (*British Medical Journal*, 1873,) speaks of the value of potassium chlorate in catarrh. If taken early and frequently, it will stop many a cold. The best form is the lozenge, eight or ten or more of which may be taken in the twenty-four hours. These should be sucked very slowly, for its action is chiefly if not altogether local. It always quickly relieves the stuffing of the nose, the rawness of the throat, the thickness of the voice, and if begun soon enough, speedily cures the cold.

Sanguinaria. The regular employment of small doses of the tincture of this root often affords relief in protracted catarrhal affections, assuming the character of incipient phthisis. It should be persevered in for several weeks.

Scilla is useful after all active inflammation has subsided, before which time it should never be employed. It unloads the air-passages and relieves congestion and dyspnoea. Dr. C. J. B. WILLIAMS states that squill is principally used in mild cases, unattended with purulent expectoration. He combines it with an alkali and a small portion of opium, to prevent its too rapid passage by the kidneys.

EXTERNAL REMEDIES.

Argenti Nitras. Penciling the swollen mucous membrane with a solution of nitrate of silver is recommended by Prof. NIEMEYER. (F. 301.)

Electricity. In the acute stage, a constant current from six to eight cups, applied by means of a nasal bath, produces good effects. In chronic coryza nearly all cases, it is asserted, may be cured by galvanic and Faradic stimulation.

Inhalations and Fumigations. Prof. TROUSSEAU highly recommends *benzoin* fumigations in chronic catarrh. The air of the patient's apartment should be impregnated with the vapor of benzoin, the drug being thrown upon burning coals; or it may be inhaled from a common inhaler, the balsam being placed in barley water. *Benzoin and Opium* fumes combined are of service in cases complicated with catarrhal neuralgia; *iodine vapor* is useful. Inhaling the fumes of burning *turmeric* is a common East Indian remedy. A small piece of burning turmeric should be placed under a small funnel, and the vapor, as it passes through the small aperture, drawn up into the nostrils. *Oxygen* inhalations have been advised in catarrh complicated with emphysema. *Sulphurous acid* fumigation (sprinkle from time to time, on a few red-hot cinders, a small portion of sulphur, so as to fill the room not inconveniently with the fumes,) or spray, is highly recommended.

M. ST. MARTIN, of Paris, recommends the following :

315. R. Acid. carbolie,	5 parts
Amm. pur. liq.,	6 "
Aquæ dest.,	10 "
Alcohol,	15 "

Soak some cotton wool with this mixture, and breathe the vapor from a wide-mouthed bottle.

The following drugs may be used in the form of spray. The dose mentioned is to be added to one ounce of water:

Acidum carbolicum,	1-2 grs
Acidum sulphurosum,	2-8 fl. drs.
Acidum tannicum,	3-12 grs.
Alumen exsiccatum,	3-20 grs.
Argenti nitras,	1-3 grs.
Borax,	5-20 grs.
Extractum belladonnæ,	$\frac{1}{4}$ -1 gr.
Extractum conii,	5-10 grs.
Extractum opii,	$\frac{1}{4}$ -2 grs.
Ferri ammonio-sulphas,	3-6 grs.
Hydrargyri perchloridum,	$\frac{1}{8}$ - $\frac{1}{4}$.
Liquor arsenicalis,	3-8 mins.
Oleum terebinthinæ,	1-5 mins.
Potassii chloras,	5-10 grs.
Potassii permanganas,	2-4 grs.
Potassii bromidum,	2-10 grs.
Potassii iodidum,	2-10 grs.
Sodii chloridum,	5-40 grs.
Tinctura ferri perchloridi,	5-30 mins.
Tinctura iodi,	1-15 mins.
Tinctura opii,	3-20 mins.
Zinci sulphas,	3-15 grs.

EMPHYSEMA.

DR. ROBERTS BARTHOLOW, PHILADELPHIA.

Arsenic, long continued, in ordinary medicinal doses, is capable of effecting considerable amelioration. Besides the stomachic administration, it is used with advantage by the process of fumigation. The following is TROUSSEAU's formula for arsenical cigarettes:

316. R. Arsenite of potassa,	15 grains
Distilled water,	1 ounce.

Unsize white paper is thoroughly moistened with this solution, dried and cut into twenty equal parts, and each part rolled into a cigarette. Two or three of these are smoked daily. The arseniate of soda may be used in the same way, and under the same conditions. If continued for a sufficient length of time, cod-liver oil is of the greatest service.

The lacto-phosphates, the phosphates, and the hypophosphites, are undoubtedly useful. The utility of these preparations depends on their power to promote constructive metamorphosis. If they improve the appetite, promote digestion and increase the body-weight, they do good; if they disagree with the stomach, they do harm. Not infrequently the syrup of the hypophosphites gives rise to distressing tormina. This may be obviated by combining it with dilute phosphoric acid—a combination very effective, therapeutically.

317. R. Syr. hypophos. co, $\bar{3}$ iiss
 Acid. phosphor. dil., $\bar{3}$ ss. M.
 S.—Teaspoonful three times a day.

Strychnine will be found serviceable.

318. R. Strychniæ, gr. j
 Acid muriat. dil., $\bar{3}$ j. M.
 S.—Five to ten drops in water three times a day.

DR. J. M. DA COSTA, PHILADELPHIA.

The treatment in cases of emphysema should be a double one; to modify the bronchial trouble which keeps up the affection, and to alleviate the difficulty of breathing, which may at times rise into paroxysms of attacks of asthma. In point of radical treatment, there are no means which will cause the distended air-vessels to resume their natural size. As a matter of absolute experience, Dr. DA COSTA has thought that he has seen, in cases of pure emphysema, not associated with bronchial symptoms, a long course of *iodide of potassium* (three to five grains, thrice daily, for several months,) favorably influence the disease, the respiratory murmur becoming fuller and freer, and the prominence of the chest-walls less visible. Good results are also obtained from persistent *counter-irritation*. Small flying blisters at various portions of the chest are serviceable. A number of cases are reported in which the constant employment of the continuous *galvanic current* has led to a diminution in the size of the chest.

Strychnia and nux vomica are useless. *Chlorate of potassium*, as well as iodide, is of service.

319. R. Potassii chloratis, $\bar{3}$ iss
 Tincturæ belladonnæ, f. $\bar{3}$ iss
 Extracti pruni virginianæ fluidi, āā f. $\bar{3}$ ij M.
 Tincturæ cinchonæ compositæ, āā
 A dessertspoonful four times a day, in emphysema with chronic bronchitis and loss of appetite. Also, dry cups applied to the chest morning and evening.

320. R. Potassii iodidi, ℥ ij
 Extracti senegæ fluidi, f. ℥ j
 Syrupi pruni virginianæ, f. ℥ ij. M.

A teaspoonful thrice daily.

Or:

321. R. Potassii iodidi, ℥ ij
 Syrupi ipecacuanhæ,
 Tincturæ scillæ, āā f. ℥ ss
 Syrupi, f. ℥ ij. M.

A tablespoonful thrice daily.

Together with counter-irritation to the chest.

322. R. Chloroformi, f. ℥ ss
 Linimenti ammoniæ,
 Linimenti saponis, āā f. ℥ iss. M.

To be rubbed on morning and evening, and to be placed on flannel, and allowed to remain against the skin for fifteen minutes.

The iodide is also favored by

E. HEADLAM GREENHOW, M. D., LONDON.

323. R. Potassii iodidi, gr. xxxvj
 Ammonii carbonatis, ℥ j
 Tincturæ scillæ,
 Tincturæ hyoseyami, āā f. ℥ ss
 Aquæ camphoræ, f. ℥ v. M.

A tablespoonful thrice daily. Also direct the patient to smoke a stramonium cigarette so soon as he feels the commencement of an asthmatic attack.

PROF. FELIX VON NIEMEYER, M. D., TUBINGEN.

The symptomatic indications in emphysema (our author considers we are totally unable to fulfill the indications of the disease itself, the nutritive alterations upon which it depends being irreparable,) are first, the proper treatment of bronchial catarrh, which almost always accompanies this affection, and greatly adds to the distress of the patient. Habitual wearing of flannel next the skin, stimulants to the chest, warm baths of water or vapor, and the alkaline muriatic mineral springs (especially the thermal springs of Ems), are of signal benefit.

The next symptomatic indication is to moderate the habitual shortness of breath and the attacks of severe dyspnoea. Sending the patient, during the summer, to the pine-wood region, and particularly to places where there is a heavy fall of dew, will allay the persistent oppression of the chest. The inhalation of compressed air, for the same reason,

is an excellent palliative. To avert the asthmatic attacks, a strict diet, the avoidance of food likely to induce flatulence, light suppers, and the keeping of the bowels open, are required.

During the attacks of emphysematous asthma, the narcotics should be used with caution, unless called for by bronchial spasm. The more suitable remedies (beside the emetics, which are very appropriate,) are the stimulants, camphor, musk, benzine, and

324. R. Vini portensis, f. $\overline{3}$ j-iss
Every three hours.

When these fail, use

325. R. Olei terebinthinæ, f. $\overline{3}$ j-iv
Aquæ menthæ piperitæ, f. $\overline{3}$ iv
Sacchari,
Pulveris acaciæ, āā $\overline{3}$ j. M.
A tablespoonful every three hours.

For the dropsy complicating the affection, vigorous diaphoresis will give excellent results whenever it depends upon a capillary bronchitis. Later in the disease, when it arises from failure of the heart to compensate for the circulatory derangement of the lungs, it may be relieved for a time, by

326. R. Pulveris digitalis, $\overline{3}$ ss-j
Aquæ, f. $\overline{3}$ vj. M.
For an infusion. A tablespoonful thrice daily.

Where digitalis fails, squills may be employed.

327. R. Aceti scillæ, f. $\overline{3}$ j
Aquæ destillatæ, f. $\overline{3}$ vj
Potassii carbonas, ad saturandum. M.
A tablespoonful every two hours.

MEREDITH CLYMER, M. D., NEW YORK.

For the relief of the asthmatic fits of emphysema, nothing is so sure as a full dose of *opium* with sulphuric or chloric ether.

328. R. Tincturæ opii, f. $\overline{3}$ j
Ætheris, f. $\overline{3}$ ij. M.
Sixty drops every twenty minutes.

To each dose may be added:

Tincturæ lobeliæ æthereæ,

gtt. xx.

Lobelia is also extolled by

THOMAS KING CHAMBERS, M. D., LONDON.

329. R.	Tincturæ ferri chloridi,	f. ℥ ijss	
	Tincturæ lobeliæ æthereæ,	f. ℥ ij	
	Aquæ camphoræ,	f. ℥ iv.	M.

A tablespoonful, in water, thrice daily.

The object of the iron is to try and restore its full vital powers to the creative arterial blood, so that it may renew the pulmonary membrane, that it may form healthy elastic tissue, instead of the imperfectly elastic degenerated tissue. The lobelia is ordered as a substitute for a more powerful medicinal agent, *tobacco*. Nothing calms the distressing asthma so well as a few whiffs of strong Virginia.

PNEUMATIC TREATMENT.

Several German writers have reported cures of emphysematous cases by expiration of rarefied air, and inspiration of condensed air rich in oxygen. The object is to remove as much as possible of the "residual air" which remains in the superior parts of the lungs after each inspiration, and to introduce highly-oxygenized fresh air, without the alveoli enduring a higher pressure than the surface of the body. Dr. JOSEPHSON, of Hamburg; Dr. PIRCHER, of Meran, and others, have reported very considerable success by these methods, systematically carried out. (*Dobell's Report*, 1877, pp. 375-6.)

NOTES ON REMEDIES.

Aspidosperma quebracho. The troublesome dyspnœa accompanying emphysematous attacks can often be relieved by the fluid extract of this plant, f. ℥ ss, *pro re nata*. It has, however, little effect on aged subjects. (*Therap. Gaz.*, 1880.)

Æther Sulphuricus is a valuable anti-spasmodic. Dr. M. CHARTÉRIIS recommends in the paroxysmal cough of emphysema:

330. R.	Tinct. lobeliæ,	f. ℥ ij	
	Spts. æth. sulph.,	f. ℥ ij	
	Tinct. conii,	f. ℥ ij	
	Mist. amygdal.,	ad f. ℥ vj.	M.

A tablespoonful every three hours.

Grindelia is recommended as an effective remedy for the bronchitis of emphysema.

Oleum Terebinthinæ is advised by Prof. NIEMEYER for attacks of emphysematous asthma.

**Opium* is of decided value in asthmatic emphysema, especially combined with ether. (F. 327.)

Stramonium is undoubtedly useful in many cases.

Tabacum, according to Dr. CHAMBERS, smoked, relieves the bronchial spasm.

EXTERNAL REMEDIES.

Respirators. Various attempts have been made to devise means to facilitate expiration. A method suggested by Dr. BERKHART to effect this purpose consists in adjusting an apparatus to the mouth and nose, from which a tube proceeds to a suction-pump, which is worked with each expiration, so as to rarefy the air.

Elastic Respirator. Dr. BAZILE FÉRIS (*Bull. Gén. de Thérap.*, 1883), considering that dyspnœa in emphysema is produced by loss of elasticity of the alveolar walls, has designed an apparatus to facilitate expiration, and thus restore to the lung its lost elasticity and abolish the dyspnœa. The apparatus designed is extremely simple, and can be easily worn. It resembles exactly a double hernia-truss: from a pad situated between the shoulders, the two limbs of the truss pass round under the armpits to the terminal pads in front, which are rather larger and thinner than those of a hernia-truss. Light straps passing over the shoulders keep the pads in position. The parts at which these pads should be made to exercise pressure are the least resistant parts of the thoracic wall, usually the upper and anterior part of the chest, over the cartilages and the ends of the ribs. The first effect of the pressure is to drive the air out from the chest, thus aiding expiration; inspiration follows naturally, the inspiratory muscles being increased in volume. Tracings show increased movement of the chest when the apparatus is worn.

Compression. Dr. GERHARDT has adopted, with some advantage, an old method, consisting in making pressure, during expiration, on the chest walls and on the abdomen. The results of this method are favorable in proportion to the youth of the patient and the flexibility of the cartilages of the ribs. It is very serviceable in cases in which some of the air-passages are blocked up by viscid secretion. While Dr. GERHARDT has found very advantageous results from this expedient, methodically applied, he has recognized two dangers in its application, neither of which has, however, had any untoward result. These are, firstly, hæmoptysis, which, no doubt, was the result of the increased pressure of the blood, brought about by the external compression; and, secondly, that muscular twitching which was observed to supervene during the process.

EMPHYEMA.

DR. JOHN G. BLAKE, BOSTON.

This writer states that the treatment of empyema is now chiefly surgical. (*Medical and Surgical Reports of the Boston City Hospital*, Second Series, 1877.) His method of operating is to make an incision from one to two inches long, parallel with the ribs, between the seventh and eighth, a little inside the scapula, and toward the axillary line. This point possesses the advantage of being the most dependent, in the position easiest to the patient, and of allowing perfect drainage. Usually, a catheter or tube has to be maintained in the incision to keep it open. No general anæsthetic should be administered, as experience shows that patients requiring this operation bear even ether badly.

The after-treatment depends a good deal on the character of the pus. When it is free from odor, it is not desirable to begin syringing the cavity at once. Such a procedure cannot fail to be irritating, and sudden deaths have been attributed to it. It is time to begin washing out the cavity when the pus becomes offensive. It may then be done once or several times a day, with a solution of carbolic acid of appropriate strength, as:

331. R.	Acidi carbolici,	gr. j-iiij	
	Glycerinæ,	f. $\frac{3}{4}$ j	
	Aquæ,	f. $\frac{3}{4}$ j.	M.

For an injection.

Or salicylic acid may be used:

332. R.	Acidi salicylici,	$\frac{3}{4}$ j	
	Sodii phosphatis,	$\frac{3}{4}$ iiij	
	Aquæ,	f. $\frac{3}{4}$ viiij.	M.

Dilute with an equal quantity of water, and use for a dressing and injection.

Or iodine may be employed, as:

333. R.	Potassii iodidi,	gr. xxx	
	Iodinii,	gr. x	
	Aquæ,	f. $\frac{3}{4}$ viiij.	M.

In reference to the class of cases in which the operation is indicated, Dr. BLAKE adopts the conclusions laid down by Dr. HENRY I. BOWDITCH, as follows:

First. Children with empyema are more likely to recover than adults ; nearly all recover.

Second. Adults with recent trouble are in a more favorable condition than those in whom the disease has lasted many months. When the disease is recent, the lungs expand rapidly, and improvement begins from the time of operation ; when chronic, expansion is slower, may be only partial, or, possibly, may not occur at all.

A permanent opening may be made in two ways, either by a large trocar and canula, and the insertion of a drainage tube afterwards, or by making an incision one or two inches long between the ribs, and keeping it open. The latter is more painful, but it is the better operation.

Do not make a permanent opening in cases where there is well-marked phthisis.

If the pus is sweet, injections are not required. If there are signs of purulent absorption, use carbolic-acid solution (one or two grains to the ounce of warm water is generally sufficient), wash out the chest twice a day, or oftener, if necessary, *and be sure you have constant drainage.*

DR. BEVERLY ROBINSON, NEW YORK.

This writer has carefully examined the statistics of death after *aspiration* for fluids in the pleural cavity, and maintains, as the result of his researches, that one of the most positive advances of our day, in the therapeutics of pleurisy with effusion, is thoracentesis by means of the pneumatic aspirator. Inasmuch as it is proved that puncture of the chest walls, with a capillary needle attached to the improved aspirator, whenever performed with due precautions against the entrance of air into the pleura, is a perfectly simple and harmless operation, and, further, that any appreciable amount of liquid, irrespective of its nature, is, by its presence, pernicious, and may become dangerous : therefore, he holds that, *in all cases of pleuritis, in which fluid is present, we should, without hesitation, make use of the aspirator to withdraw the morbid effusion.*

To this law he only affixes *one* limitation and *one* exception.

The limitation is : Whenever very large or excessive quantities of fluid are present, it is wiser to puncture the chest on two successive occasions, so that all risk of acute oedema of the lung on the affected side shall be avoided.

The exception is: If the patient be very much enfeebled, and the effusion be small or moderate, we may, with advantage, delay the operation, during a brief period, until his forces have been somewhat re-established.

By proceeding after this manner, all danger of *fatal syncope* will be obviated. (*New York Medical Record*, Nos. 325, 326.)

Meanwhile, of course, if the effusion from small, or moderate, rapidly becomes large or excessive, the formal and imperative indication is to operate so soon as possible.

DR. HENRY W. FULLER, LONDON.

This experienced physician thinks there is a tendency to have recourse to tapplings or incisions too early in pleuritic effusions. (*British Medical Journal*, January, 1872.) When called to a patient whose chest is full of fluid, we should use an aspirator, or small grooved needle, to ascertain the character of the fluid. If it is simply serum, other measures should precede tapping. Diuretics, diaphoretics, and mercury, are generally of no value. In many instances, *blisters* prove very beneficial; so, also, the constant application of a weak solution of iodine.

334. R. Tinct. iodinii compos.,
Glycerinæ,
Aquæ,

f. $\frac{z}{3}$ j
f. $\frac{3}{4}$ ij
f. $\frac{3}{4}$ v. M.

For external use.

This weak solution has the advantage of possessing all the absorbent properties of iodine, without damaging the cuticle, and thus destroying the absorbing surface. In addition to these, the *dry method* of treatment may be employed. This is denying the patient all fluid food; it is useful where the pleural membrane is not yet much diseased. If, after a month of treatment, marked improvement is not visible, tapping should be resorted to. The fear of bad results from the admission of air into the pleural sac, is unfounded.

DR. BALTHAZAR FOSTER, LONDON.

The respective indications for aspiration, tapping, and incision, in cases of pleuritic effusion, have recently been considered by this writer. Experience having shown that both the expectant and the active medicinal treatment are alike untrustworthy, the surgical is our only resource. Dr. FOSTER divides cases of pleuritic effusion into several

classes: 1. *Acute cases.* In these, the use of the aspirator is forced upon us by certain urgent symptoms, especially dyspnœa. Here, the operation is a means of saving life, not of treatment; for a chest full of fluid, dyspnœa being absent, is not an indication for aspiration. Certain adynamic cases, however, occur where the fluid soon becomes purulent; here, a preliminary opening should be made, and, if pus be present, aspiration is the most appropriate treatment. 2. *Cases of over three weeks' duration, where fever has abated.* Here, the operation should be performed to relieve the morbid condition and expedite recovery. In such cases, the lung becomes covered with a fibrinous deposit, and absorption of fluid is impeded. No dependence, therefore, can be placed upon the use of diuretics, blisters, and other general remedies. 3. *Chronic pleurisy, with exacerbations.* Here, if the chest be from one-half to two-thirds full of fluid, and the disease have lasted over two months, aspiration is decidedly indicated. 4. *Empyema.* Pus being present, the rule should be evacuation. Dr. FOSTER compares the various methods employed for emptying the pleural cavity. He recommends in children, repeated aspirations, with, in some cases, washing out the pleural cavity with carbolized fluid. In other cases, and especially in adults, he prefers two incisions through the chest-wall and the use of drainage tubes; and, if this is not sufficient, a free opening and a thorough washing out of the pleural cavity. Some cautions are necessary in aspirating. First, the operation must not be too rapidly performed; second, the quantity of fluid allowed to escape must be carefully watched, and must depend upon the condition of the lung and the state of the patient.

MOUNT SINAI HOSPITAL, NEW YORK CITY.

The method employed in a large number of cases in this hospital, with good success, is to make two openings in the chest, one anterior, and the other lateral and slightly posterior. The pus is entirely removed, and *warm water*, of a temperature of 100° , either plain or very slightly medicated with carbolic or salicylic acid, or iodine, is carried in at the upper opening and out at the lower one, until all trace of pus has disappeared from the discharge. Pieces of oil-silk are placed over the lower opening, and so fastened with collodion that the bed-clothes are not soiled. In order to secure a free circulation for the water, the catheter conducting it in should be inserted well into the anterior open-

ing. In some cases, where a counter-opening may seem to be contra-indicated, it becomes necessary to carry a gum-elastic catheter in for its full length, and even then, all trace of pus may not be removed. It has been found by experiment that, when a double canula is introduced for the entrance and exit of the fluid, a current is caused between the two openings of the canula, and only an ounce or two of pus is removed; and, as before suggested, the next best plan to that of counter-openings is to make use of a long catheter and allow discharge to take place at the opening made in the chest. (*New York Medical Journal*, August, 1876.)

NOTES ON REMEDIES.

Acidum Carbolicum, when used as an injection, is said by Dr. DICKENSON to cause serious irritation. (*British Medical Journal*, July 8th, 1876.) But it is approved by Dr. BLAKE (F. 331) and Dr. WINIWARTER; the latter washes the pleural cavity with a one per cent. solution.

Acidum Salicylicum has been used. (F. 332.)

Iodinium is a favorite injection. Dr. DICKENSON recommends :

335. R. Liquoris iodinii, Aquaë,	f. $\frac{3}{5}$ ij Oj.	M.
-------------------------------------	----------------------------	----

EXTERNAL MEASURES.

Aspiration is often a successful method of withdrawing the accumulation.

HÆMOPTYSIS.

DR. SEYMOUR TAYLOR.

According to this author (*Lancet*, June 14, 1884), there should be no undue precipitancy in employing many of the astringents usually advocated in the early stages of pulmonary consumption. He would rather attend to the patient's general health and strength. He would not restrict him from gentle exercise, but he would, at the same time, make a distinction between exercise and over-exertion. In the later stages he strongly recommends warm applications to the thorax. He believes that we have but few drugs that possess any value as controllers of hemorrhage. Opium (and its derivatives) acts like a charm. It should not be given, if administered internally, with any other remedy likely to retard or interfere with its full action. It should be given in such

doses as to produce its physiological effects. If the hemorrhage be very profuse, digitalis may be added, but also in large doses, such as fifteen to twenty minims of the tincture, till its specific action on the heart is manifest, and the frequency of the pulse materially diminished. As a result the patient is calmed, his excited circulation controlled, and he falls into a much-needed sleep.

There is yet another method of administering the sedative—viz., by hypodermic injection—which he has found most useful and efficacious. Four minims of the injectio morphiæ hypodermica are introduced into the subcutaneous tissue of the arm, or into the subcutaneous tissue of the chest immediately over the presumed seat of the hæmorrhage, and repeated if requisite. The advantage of this method of exhibiting the remedy is its speedy action; the momentary and trivial discomfort of the operation is more than counter-balanced by the good results which ensue. He considers the above method of treatment happier in its immediate results than the administration of astringents and so-called styptics, which mainly affect the intestinal tract. There are two other drugs which he has found extremely useful when opium is contra-indicated, viz.: oil of turpentine and the liquid extract of ergot. The former may be given by the mouth, the latter either by the mouth or by subcutaneous injection. He has seen excellent results from both in a few cases. The objection to turpentine is its nauseating effect, but its action in controlling hemorrhage is undoubted. His experience of ergot has not been extensive, but he found it a powerful remedy in four cases of severe and continued blood-spitting.

DR. MARIUS CARRE, PARIS.

This writer, (*Archives Gen. de Médecine*, January–March, 1877,) remarks on the treatment of the nervous variety of hæmoptysis. Blood-letting, useful in the congestive variety, is injurious here. Anti-spasmodics, narcotics and revulsives are of little value. The use of general baths is decidedly efficient. With these should be combined the constitutional use of arsenic and quinine :

336. R. Acidi arseniosi,
Quiniæ sulphatis,

gr. $\frac{1}{40}$ — $\frac{1}{60}$
gr. i-ij. M.

For one pill. Twice or thrice daily.

M. GUBLER has shown that quinine stimulates capillary contractility directly by contact, and indirectly by its action on the spinal cord.

Attacks of profuse hemorrhage call for the temporary use of astringents and acids.

DR. BARTLETT, NEW YORK.

This practitioner states that for ten years he has used the following formula in hæmoptysis, and knows of no combination that at all approaches it in efficiency (*Buffalo Medical Journal*, September, 1878):

337. R.	Tr. digitalis,	℥ iss	
	Ol. terebinth.,	℥ ij	
	Ol. menth. pip.,	gtt. x	
	Acidi sulph. ar.,	℥ ij	
	Alcoholis,	q. s. ad ℥ ij.	M.

Dose, forty to sixty drops well mixed with sugar, to which one or more tablespoonfuls of water may be added, every two, three or four hours, according to urgency of the hemorrhage.

DR. J. M. DA COSTA, PHILADELPHIA.

The following are the agents generally employed by this practitioner:

338. R.	Acidi gallici,	gr. xx.	
For one powder. In acute hæmoptysis. To be repeated every ten minutes until hemorrhage ceases.			

339. R.	Cupri sulphatis,	gr. $\frac{1}{2}$	
	Ferri sulphatis,	gr. ij	
	Extracti hyoscyami,	gr. j.	M.
For one pill, thrice daily.			

In persistent slight pulmonary hemorrhage. Sulphate of copper in such cases seems to control the circulation within the lungs, and to arrest a tendency to bleeding. This action is not due to its nauseating effect, for, in this dose, it does not nauseate. The administration of cod-liver oil should be postponed until the hæmoptysis is gotten rid of.

340. R.	Ferri chloridi,	gr. ij-x	
	Aquæ destillatæ,	f. ℥ j.	M.
For inhalation by atomizer.			

341. R.	Liquoris ferri subsulphatis,	℥ x-xl	
	Aquæ destillatæ,	f. ℥ j.	M.
For inhalation by atomizer.			

342. R.	Acidi tannici,	gr. x-xx	
	Aquæ destillatæ,	f. ℥ j.	M.
For inhalation by atomizer.			

343. R. Pulveris aluminis, gr. xxx
 Aquæ destillatæ, f. ʒj. M.
 For inhalation by atomizer.

HORACE DOBELL, M. D., LONDON.

344. R. Extracti ergotæ fluidi, f. ʒij
 Tincturæ digitalis, f. ʒij
 Acidi gallici, ʒj
 Magnesii sulphatis, ʒv
 Acidi sulphurici diluti, f. ʒj
 Infusi rosæ compositi, f. ʒvj. M.

Two tablespoonfuls every three hours until the hemorrhage is arrested.

In spite of the fashionable outcry against complicated prescriptions, Dr. DOBELL recommends the above as the most efficacious and the most rational combination of remedies for a case of profuse tubercular pulmonary hemorrhage. In any given case, either of the ingredients may be omitted, if the symptoms indicate that it is not required, or that it has already done its duty. The object of the ergot is to contract the vessels; of the digitalis, to steady the heart; of the gallic acid, to clot the blood; of the epsom salts, to relieve the congestion; and of the dilute sulphuric acid, to assist the rest.

PROF. FELIX VON NIEMEYER, M. D., TUBINGEN.

345. R. Copaibæ,
 Syrupi,
 Aquæ menthæ piperitæ,
 Alcoholis, aa f. ʒj
 Spiritûs ætheris nitrosi, f. ʒss. M.

A dessertspoonful every two to four hours.

A formula much in use in very obstinate hæmoptysis:

346. R. Aluminii et potassii sulphatis, ʒj
 Morphiæ muriatis, gr. ʒ
 Sacchari, ʒj. M.

Divide into twelve powders. One every hour, in hæmoptysis. Cold compresses to the chest, absolute rest, and silence.

In the first place, the physician should seek to calm the spirits of the patient, which are always much excited; and, inasmuch as the attacks are almost always repeated several times, it is well to save him from further agitation, by straitway informing him that there is more blood to come, while, at the same time, we should absolutely deny the possibility of his bleeding to death. Indeed, we are certainly warranted

here in deceiving the patient, by affecting to make light of the affair, and even to represent the hemorrhage as a salutary process. With a little tact, the physician may leave his patient in a state of comfort and peace of mind—a success of no slight importance. The chamber must be kept cool, all hot drinks forbidden, and all food eaten cold. Conversation should be interdicted, and the patient urged to resist, stoutly, the provocation to cough. Coughing in hæmoptysis is quite as hurtful as is snuffing and wiping the nose in epistaxis.

NOTES ON REMEDIES.

Aceticum Acidum Dilutum is employed both externally and internally, but is a remedy of minor importance.

Acidum Gallicum. When it occurs in relaxed and debilitated conditions—is passive in character—we may use :

347. R.	Acid. gallici,	3j	
	Ergotine (aq. ex.,)		
	Digitalis,	āā	℞j. M.

Ft. Pil. No. xx. S.—One every four hours.

Alumen is serviceable in purely atonic hæmoptysis. It may be given in solution, by inhalation, or be allowed to dissolve slowly in the mouth, in substance or in lozenge. In moderate but persistent tubercular hæmoptysis, Dr. SYMONDS recommends (*British Medical Journal*, June 13th, 1868,) the following :

348. R.	Aluminis,	℞ij	
	Acidi sulphurici diluti,	f. 3 ijss-iv	
	Magnesii sulphatis,	℞iv	
	Aquæ,	f. 3 iv.	M.

A tablespoonful in water, every three hours, or oftener, as required.

Ammonii Chloridum is recommended by Dr. COPLAND, in combination with muriatic acid, in passive cases of hæmoptysis, when the vital powers are depressed. The following is his formula :

349. R.	Ammonii chloridi,	3 iss	
	Acidi muriatici,	f. 3 ss	
	Decocti hordei compositi,	Oj.	M.

A small wineglassful every two or three hours.

Argenti Oxidum, in doses of gr, ss-j, thrice daily, has been recommended, in atonic hæmoptysis.

Barium Chloride, says BARTHOLOW, will probably prove useful.

Chloroform. Several cases are reported by Dr. A. M. WEIR, of Edinburgh, (*Lancet*, January, 1877,) in which the hæmoptysis was promptly checked, after other remedies had failed, by applying a flannel pad soaked in chloroform, over the lung from which the blood flows.

The pad should be covered with towels or oiled silk. Its effect is said to be immediate and lasting.

Cnicus Lanceolatus. The milk-thistle is said by Dr. LESSENWITCH to be serviceable for the relief of hæmoptysis, he having tried it in five cases, using the tincture in doses of fifteen to twenty drops in a teaspoonful of water every two hours.

Copaiba is much employed by Prof. NIEMEYER.

Creasotum fails to give satisfactory results.

Cupri Sulphas is highly recommended by Prof. DA COSTA, in cases of persistent slight hemorrhage.

**Digitalis* is advised by Dr. WALSH, when there is irritable heart and moderate hæmoptysis. By some, it is regarded as the best remedy for hemorrhage from cavities in the advanced stages of phthisis.

Ergota is strongly recommended by Dr. DOBELL, who, however, usually associates it with other medicines. (F. 331.) Dr. J. M. WILLIAMSON (*Lancet*, November, 1875,) gives it in $\mathfrak{m}\mathfrak{x}$ doses of the fluid extract every half hour. Prof. J. HIRSCHFELD, of Vienna, believes the very best remedy in hæmoptysis is hypodermic injections of *ergotin* in concentrated solution (one part in ten of glycerine.) *Dobell's Reports*, 1877.)

BARTHOLOW uses the following :

350. R.	Ext. ergotæ fl.,	f. \mathfrak{z} iij	
	“ ipecac fl.,		
	Tinct. opii deodor.,	āā f. \mathfrak{z} ss.	M.

S.—A teaspoonful every half hour or hour.

When the sputa are heavily charged with blood, but there is no defined hemorrhage, he uses :

351. R.	Ergotinæ (aq. ex.,)	℥j	
	Pulv. ipecac,	gr. x	
	Acid. gallic.,	℥j.	M.

Ft. Pil. No. xx. S.—One every hour or two.

Ferrum is particularly of service in hæmoptysis connected with amenorrhœa. The preparations most employed are the tincture of the chloride, the solution of the nitrate, (in large doses, $\mathfrak{m}\mathfrak{xxx}$ – $\mathfrak{l}\mathfrak{x}$,) the sulphate, and the ammonio-sulphate.

***Gallicum Acidum* is highly recommended by Profs. DA COSTA, WALSH, and others as one of the best remedies in hæmoptysis, but it must be given in large doses (F. 338), gr. x–xxx, repeated as necessary.

Hamamelis. Dr. HUGO ENGEL (*Med. and Surg. Rep.*, May, 1879,) states that this drug is far superior to any other in hæmoptysis. In violent cases, he gives two teaspoonfuls of the fluid extract of hamamelis right away, and repeats the dose every half hour till the severe bleeding stops, and continues it later, in the dose of one or half a teaspoonful three times daily, until all signs of the spitting of blood have disappeared.

Hydrocyanicum Acidum seems to possess some power in controlling hæmoptysis when the system is irritable and spasmodic symptoms are present.

**Ipecacuanha* is recommended by Prof. GRAVES and other practitioners, as a remedy of considerable value. From one to two grains are given every fifteen or thirty minutes, until nausea is felt, when the benefit is generally evident, and the medicine is discontinued. Many cases are reported, attesting the efficacy of this treatment in arresting the hemorrhage and restoring heat and life to patients who were in a state of collapse from excessive loss of blood. Although it is better to avoid vomiting, no ill effects, it is said, are produced by it. The hæmoptysis of the early stages of phthisis may often be readily arrested by this remedy.

Matico is sometimes employed. In slight cases, the infusion alone is often sufficient. It is advantageously combined with ergot.

Oleum Terebinthinæ is often efficient.

Phosphoricum Acidum Dilutum, in doses of gtt. x-xxx in mucilage, thrice daily, is reported upon favorably by M. HOFFMAN, (*The Practitioner*, August, 1868,) who considers it superior to other mineral acids in hæmoptysis.

Pyrogallicum Acidum. Dr. A. VESEY has found this of the greatest service, in doses of gr. j every half hour. It may advantageously be combined with ergot (*Dublin Jour. of Med. Science*, December, 1875.) It has the advantage of not disturbing the stomach.

**Plumbi Acetas*, in combination with morphia and acetic acid, or with opium, is a remedy much used. Dr. PEREIRA considers that the action of sugar of lead and opium in hæmoptysis is rendered more certain and speedy by being given with a draught containing nitric acid, nitrate of lead being thus formed in the stomach.

Potassii Nitras is useful where there is much excitement and fever, but is not to be relied upon alone. It may be given in doses of gr. viij-x, several times daily, largely diluted, or combined with tartar emetic or digitalis.

Sodii Chloridum is recommended by Prof. WALSHE, in drachm doses, either in powder or dissolved in water. It is sometimes promptly successful, even when it produces emesis.

**Sulphuricum Acidum Dilutum* is a useful remedy, though less certain in its action than gallic acid, and is a valuable adjunct to other treatment.

Tannicum Acidum, in combination with opium and ipecacuanha, has been highly recommended.

Veratrum Viride in full medicinal doses, will sometimes control active hemorrhage, occurring in the plethoric.

EXTERNAL MEASURES.

Cold. Ice to the chest, to the spine, or over the heart, will sometimes instantaneously arrest the hemorrhage.

Dry Cups. In the *Allg. Wien. Med. Zeit.*, May, 1878, Dr. H. SCHNEIDER expresses the opinion that the best means for the relief of congestion, or

a hyperæmic condition about the lungs, is an old remedy which has fallen into disuse of late years—the application of cups. The number of cups to be applied must depend on the amount of congestion, and the effect which it is desired should be produced. A considerable amount of blood can be withdrawn from the general circulation, and will be kept in the outer wall of the chest. In comparatively mild cases, dry cups are applied to the thorax on the side corresponding with the congested lung, while if the bleeding is severe, wet cups should be placed on the affected side.

Ligation of the limbs by a tourniquet, or the Esmarch bandage, so as to prevent the venous flow, is of service in severe cases. “Jounod’s boots” have been used for the same purpose.

HAY FEVER—HAY ASTHMA—ROSE COLD—AUTUMNAL CATARRH.

PROFESSOR HARRISON ALLEN, OF PHILADELPHIA.

The conclusions to be drawn from the study of a number of cases reported by Dr. ALLEN, in the *American Journal of the Medical Sciences*, January, 1884,) may be summarized briefly as follows :

1. That the treatment of all conditions of obstruction in the nasal chambers, no matter from what cause arising, can be successfully carried out by destroying the causes of obstruction. If the cause be an overgrowth of bone-tissue, it must be filed, sawed or drilled away. If it be caused by a deviated cartilaginous portion of the septum, such portion must be re-set in a new place. If, as is often the case, it is due to periodic turgescence of the mucous membrane, or the resulting secondary hypertrophies, such growths must be destroyed, either by the galvano-cautery, by the snare, or by caustic acids.

2. That the treatment of hay-fever and allied periodically-recurring nasal affections in no way differs from the treatment of other nasal diseases accompanied by obstruction, and that the treatment may be conducted during an attack as well as in the intervals between any two attacks.

MR. J. B. HANNAY, OF NEW BRUNSWICK,

writes to *Nature* that he has succeeded in finding a means for the prevention of Hay Fever. He found that the only thing required was to prevent the entrance of pollen grains into the nose. For this purpose

he uses a piece of brass or steel ribbon, bent double, and having only sufficient spring to close the nostrils without undue pressure. When going among hay a further precaution should be taken, viz., plugging the lachrymal ducts. He uses for this purpose dumb-bell shaped pieces of glass, which were easily slipped into the ducts and could be removed when wanted.

Many authorities, prominently among them Dr. CARL SEILER, of Philadelphia, claim that an hypertrophy of the post-nasal mucous membrane predisposes to hay fever, and that the tendency cannot be eradicated until this hypertrophied tissue is removed. This can be accomplished with the galvano-ecraseur.

Dr. CHARLES E. SAJOLS, of Philadelphia, has cured some severe cases by the topical application of *glacial acetic acid*, but owing to the pain produced by this agent, he prefers the galvanic cautery. (*Med. and Surg. Reporter*, Dec., 1883.)

DR. GEORGE M. BEARD, NEW YORK.

According to this writer, there is no specific for hay fever in general, but individuals may be able to find a specific for their own cases. Many measures afford great relief in all cases. A sea voyage rarely fails, as well as a residence in elevated regions, and in certain localities.

The medical treatment he divides into general and local. Of internal remedies he considers *quinine* the most efficient. When given to prevent the disease, gr. j–ij, two or three times a day, for several weeks before the time of the attack, is the proper dose. *Arsenic*, especially when combined with belladonna, is efficient.

352. Liq. potassii arsenitis,
Tinct. belladonnæ,

f. $\bar{3}$ j.
f. $\bar{3}$ ij. M.

From five to ten drops after meals.

Phosphide of zinc, gr. $\frac{1}{10}$ – $\frac{1}{4}$ acts well in many cases. *Opium* in any form will palliate the symptoms. One of the best preparations for the purpose, is the "cold powder," as follows:

353. R. Camphoræ,
Ether. sulphuric.,

$\bar{3}$ v
q. s.

Dissolve to the consistency of cream, and add:

Ammonia carbonatis,
Pulveris opii,

$\bar{3}$ iv
 $\bar{3}$ j. M.

Keep tightly corked.

The dose is to be regulated by the quantity of opium borne.

Locally, many substances can be applied by means of the atomizer. Solutions of quinine are perhaps the most effectual of all. The following is commended by Dr. R. P. LINCOLN, of New York:

- | | | | |
|---------|-------------------|---------|----|
| 354. R. | Quiniæ sulphatis, | gr. vj | |
| | Aquæ camphoræ, | f. ʒvj | |
| | Sodii chloridi, | ʒss | |
| | Aquæ, | f. ʒiv. | M. |
- Use as an atomizer or snuff from the hand.

The *pinus canadensis* has been used in the same way.

- | | | | |
|---------|-----------------------------|-------|----|
| 355. R. | Fld. ext. pinus canadensis, | f. ʒj | |
| | Aquæ, | Oj. | M. |

The following is well spoken of:

- | | | | |
|---------|---------------------|---------|----|
| 356. R. | Potassii chloratis, | ʒj | |
| | Morphiæ sulphatis, | gr. xij | |
| | Aquæ, | f. ʒvj. | M. |
- Use with the atomizer.

Inhalations of aquæ ammoniæ relieve some cases.

Smoking stramonium leaves, nitre paper, etc., benefits some. Dr. MORRILL WYMAN recommends:

- | | | | |
|---------|----------------------------|--------|----------|
| 357. R. | Belladonnæ fol., | gr. ix | |
| | Hyoscyami fol., | | |
| | Stramonii fol., | āā | gr. ivss |
| | Phellandrii aquatici fol., | | gr. iss |
| | Opii, | | gr. ¼. |
| | | | M. |

Make two cigarettes by rolling in paper. Only two to be used during an attack.

DR. E. C. SEGUIN, NEW YORK.

- | | | | |
|---------|------------------|--------|----|
| 358. R. | Ammonii bromidi, | ʒi-ij | |
| | Aquæ, | f. ʒj. | M. |

Use as a gargle frequently.

- | | | | |
|---------|------------------|-----------|--|
| 359. R. | Ammonii bromidi, | gr. x-xxx | |
| | Aquæ, | f. ʒj. | |

Use by an atomizer, or inhale into the nostrils.

This has given marked relief to patients. (*New York Medical Record*, November, 1876.)

DR. J. S. HOLDEN, ENGLAND.

360. R. Quiniæ sulphatis, gr. xij
 Acidi hydrochlorici diluti, ℥xxx
 Aquæ, f. ℥vj. M.

Add

Sodii bicarbonatis, q. s.

To make a neutral solution, add one ounce of hot water to two ounces of this lotion, and douche the nasal passages three times a day for a few days at the beginning of the attack. (*Lancet*, May, 1876.)

The *grindelia robusta* is said to give prompt relief.

361. R. Fld. extracti grindeliæ, f. ℥iv
 Fld. extracti rhei,
 Fld. extracti sennæ, āā f. ℥j. M.

A dessertspoonful every half hour during the spasm, and afterwards at intervals of three hours.

DR. HORACE DOBELL, LONDON.

362. R. Chloral hydratis,
 Camphoræ pulv., āā gr. xvj
 Acidi carbolic, ℥j
 Morphiæ, gr. xij
 Acidi oleici, gr. vij
 Olei ricini, f. ℥vij. M.

Rub well together to make a lotion. Apply by means of a little ivory or hard-wood club to the interior of the nostril.

This author states that the tincture of *eucalyptus globulus* is valuable in hay fever where there is much prostration.

DR. JOHN O. ROE, OF NEW YORK,

contributes an article on this subject to the *N. Y. Med. Jour.*, May 19, 1883, in which he reaches the following conclusions :

1. That the treatment during the attack can only be palliative, such as to soothe the inflamed parts and to quiet the systematic disturbance which may be occasioned.

2. That in most cases the only effective relief during the attack consists in going to a seaport or mountainous region, or to any locality where the air is free from the substance which produces the irritation.

3. That curative measures can only be adopted when the individual is free from the attack.

4. That the removal of the diseased tissue in the nasal passages removes the susceptibility of the individual to future attacks of hay fever.

For the removal of this tissue, Jarvis' snare is the most efficient means of removing that at the posterior end of the turbinated bone, while that covering the middle and anterior portion is most reliably removed by the galvano-cautery.

DR. EDWARD R. MAYER, PENNSYLVANIA.

363. R. Quiniæ sulphatis, gr. iv
Tinct. gelsemii concent., gtt. vj.

This amount in water every four waking hours. Regarded by Dr. EDWARD R. MAYER as the best remedy for and preventive of this disease. Cures in forty-eight hours. (*Hints on Specific Medication*, p. 21.)

DR. ROBERTS BARTHOLOW, PHILADELPHIA.

364. R. Potassii iodidi, \bar{z} j
Liq. potassii arsenitis, f. \bar{z} j
Aquæ, f. \bar{z} iv. M.
A teaspoonful every four or six hours.

In addition to this internal medication, apply locally the following with a post-nasal syringe:

365. R. Tinct. iodinii, f. \bar{z} j
Acidi carbolici, gtt. x
Aquæ destillatæ, f. \bar{z} iv. M.

Arsenical cigarettes, as recommended by TROUSSEAU, may be used with advantage.

366. R. Potassii arsenitis, gr. xv
Aquæ, f. \bar{z} j. M.

Moisten unsized white paper with this solution, cut and roll into cigarettes. Two or three may be smoked daily.

NOTES ON REMEDIES.

Acid Carbolic. BARTHOLOW says that inhalations of carbolic spray are highly useful. A solution in water, to the proportion of one per cent., can be used.

Acidum Sulphurosum. According to Dr. JAMES DEWAR, of Ireland, the symptoms of hay fever are often checked at once by inhaling sulphurous acid in the form of the fumes of burning sulphur. (*Dublin Medical Press*, September, 1866.)

Aconitum has been recommended by Dr. RINGER.

Ammonia Murias, taken internally, is suggested by Dr. BEARD.

Anthemis Nobilis. Chamomile flowers smoked, or their fumes inhaled, often act very well.

Arsenicum is an important auxiliary.

Atropine is useful when the secretion is profuse.

Belladonna, either the leaves smoked or in doses of the tincture, is occasionally efficient. Mr. W. F. PHILLIPS (*Brit. Med. Jour.*,) 1884, uses:

367. R. Succi belladonnæ, \mathfrak{m}_{xxiv}
 Aquam, ad \mathfrak{z} ij.
 M.—A teaspoonful to be taken every hour till relief is obtained.

“The medicine was taken without the production of any undesirable effect, and with very marked advantage indeed—an advantage that became still more evident and unmistakable, both to the patient and myself, when the dose was increased from one minim to one and a quarter, (half a drachm in three ounces).”

* *Bromine* is highly recommended,

368. R. Bromine, \mathfrak{z}^{ss}
 Alcoholis, \mathfrak{z} iv. M.
 S.—For inhalation.

A small quantity of this solution may be placed in a wide-mouthed vial and vaporized by the heat of the hand. The vapor should be snuffed into the nose.

Caffea. Strong hot coffee is a useful palliative.

Camphora is a useful local application. It may be placed dry on the pillow at night.

Chloroform, by inhalation, will quickly relieve the spasm.

Creasotum. Dr. WALSHE recommends a trial of creasote inhalations once or twice daily, having seen remarkably satisfactory effects from them in two instances.

Euphrasia Officinalis, vulgó, eyebright, gtt. ij-v of the tincture, as often as required, is reported upon very favorably by Dr. H. G. PIFFARD, of New York.

Guarana is useful in relieving the headache.

Grindelia Robusta. In the asthmatic attacks consecutive on hay asthma, great relief has been experienced from the fluid extract of wild sunflower. It may be administered as follows:

369. R. Grindeliæ rob. ex. fld., (Parke, Davis & Co.,) f. \mathfrak{z} j
 Syrupi simplicis, f. \mathfrak{z} ij.
 A teaspoonful at a dose.

Iodides. The iodides carried to iodism afford great relief; they may be combined with arsenic.

Iodinium, in inhalation, benefits some cases. Dr. J. G. WESTMORELAND re-

commends placing one grain in a wide-mouthed vial, heating till it fumes, and inhaling. Repeat every day. (*Atlanta Medical and Surgical Journal*, October, 1870.)

Muscarine will probably prove useful in the asthmatic stage if the membrane is dry.

Oleum Menthe Piperitæ. Dr. JAMES E. BELL, of Georgia, himself a sufferer, states (*Medical and Surgical Reporter*, December, 1877,) that the remedy which he has found superior to all others to allay the irritation and to quiet the cough, is the oil of peppermint, applied locally, in vapor or in substance.

Opium is a valued palliative. The hypodermic injection of morphia gives great relief. But there is danger of inducing the opium habit by a resort to these palliatives.

**Quiniæ Sulphas* is the most generally useful remedy yet found. (See above.)

Salicin, gr. xx-xxx, twice daily, is extolled by Dr. T. J. MACLAGAN.

Stramonium Leaves, smoked, are efficient to allay the paroxysm.

Tabacum. The smoking of tobacco, to those who have not been much accustomed to it, has proven beneficial.

Tannin, in solution, may be used with the atomizer.

Zinci Sulphas. Dr. FULLER reports ten cases effectually cured by means of sulphate of zinc, administered internally, in gradually increasing doses, whilst zinc lotion was applied to the eyes, and a lotion composed of eight grains of sulphate of zinc and an ounce of glycerine to the lining membrane of the nostrils.

Mechanical Measures. Wearing a thick veil, or a respirator, or keeping two fine sponges in the nostrils, moistened with a glycerine solution, and learning to breathe constantly through the open mouth, are procedures which have relieved some cases.

LARYNGITIS.

For clinical purposes, the varieties of the disease may be classed as follows:

Acute Laryngitis.

Chronic Laryngitis. { Follicular or glandular laryngitis.
Aphonia clericorum, clergyman's sore throat.
Tubercular laryngitis.
Syphilitic laryngitis.

The tubercular and syphilitic forms are characterized by ulcers, which are not present in the other varieties.

ACUTE LARYNGITIS.

The rules of treatment of this common form, which so often attends an ordinary catarrh, are thus laid down by

JOHN SYER BRISTOWE, M. D., LONDON.

The patient should be kept in a warm atmosphere, free from exposure to draughts, apply hot fomentations (a sponge wrung out in hot water) or mustard plasters to the throat, and use a gargle of *warm milk*, or some slightly astringent substance. The faucial discomfort may be relieved by the inhalation of steam (in which the patient should be taught *not* to take deep, but ordinary inspirations). Diaphoresis should be encouraged by warm foot-baths, etc. Opium, in small doses (Dover's powder), greatly relieves the discomfort. In cases severe from the beginning, leeching externally and scarification within are of essential importance for the œdematous condition, which may even demand tracheotomy, an operation which it is better to perform early, before the patient is moribund. No patient should be permitted to die of this disease without having this chance placed in his favor.

DR. KRISHABER, GERMANY.

When the laryngitis is of the dangerous *œdematous* variety, this writer recommends the following treatment: If it is markedly inflammatory, antiphlogistic treatment should be practiced; one or two general bleedings, and wet cupping on the neck, will give considerable relief, diminishing the inflammation, and lessening the engorgement of the tissues. By means of a spray apparatus, water impregnated with tannin or alum may be applied to the back of the throat. In the absence of a spray-producer, a simple irrigator filled with astringent fluid may be used. Recourse may also be had to cauterization with nitrate of silver, insufflations of tannin or alum, and scarifications and divisions of the ary-epiglottic folds. These last two methods are somewhat difficult to practice, but often, under the influence of spray irrigation and local bleedings, the inflammation diminishes and disappears. When the œdema of the glottis is due to tuberculosis or cancer, art is powerless against the diathesis and the laryngitis.

As an effective combination in acute febrile attacks of laryngitis,

DR. T. WHIPHAM, ENGLAND,

has derived great benefit from the following (*Med. Times and Gazette*, 1877):

370. R. Vin. antimon.; f. ʒ ss
 Vin. ipecac., ℥_x
 Potass. acet., ʒj
 Spts. æth. nit., f. ʒj
 Potass. nitrat., ʒss. M.
- S.—To be given every four hours.

DR. LENNOX BRÔWNE, LONDON.

Inhalation is a valuable means in acute laryngitis.

371. R. Tinct. benzoini comp., f. ʒ iij
 Chloroformi, ℥_{xxv}. M.
- A teaspoonful in a pint of water, at 140° F., for each inhalation. A valuable sedative in acute inflammations of the pharynx and larynx.

372. R. Ammonii chloridi, ʒj
 Tincturæ opii, ℥_v
 Decocti cinchonæ, ad f. ʒj. M.
- For one dose, to be taken at 11 and 4, between meals. Used in catarrhal conditions, and in commencing head-colds.

Other inhalations are :

THOMAS HAWKES TANNER, M. D., LONDON.

373. R. Acidi hydrocyanici diluti, ℥_{xv}
 Spiritus chloroformi, f. ʒ iij
 Aquæ bullientis, f. ʒ viij. M.
- For inhalation.

L. WALDENBERG, M. D. BERLIN.

374. R. Sodii chloridi, gr. iv-x
 Aquæ destillatæ, f. ʒj. M.
- For inhalation in acute laryngeal catarrh, by means of nebulizer.

LONDON HOSPITAL.

375. R. Tincturæ benzoini compositæ, f. ʒj
 Aquæ bullientis, f. ʒ x. M.
- Let the vapor be inhaled frequently.

376. R. Tincturæ iodinii, ℥_{xl}
 Aquæ bullientis, f. ʒ x. M.
- To be inhaled frequently.

CHRONIC LARYNGITIS.

PROF. M. CHARTERIS, M. D., GLASGOW.

For chronic laryngitis, *chloride of zinc* has been specially recommended :

377. R. Zinci chloridi, gr. xxx
 Aquæ destil., f. $\frac{3}{4}$ j. M.
 Apply locally, every day, for a week, and afterwards on alternate days.

If the secretion is excessive, turpentine or creasote makes the best inhalation; and when the laryngeal irritation is the principal feature inhalations of hops are soothing.

378. R. Creasoti, $\frac{3}{4}$ ij
 Magnes. carbon., $\frac{3}{4}$ iss
 Aquam, ad f. $\frac{3}{4}$ ij. M.

Or:

379. R. Acidi carbolici, f. $\frac{3}{4}$ j
 Aquæ, f. $\frac{3}{4}$ vij. M.
 A teaspoonful in a pint of water, at 150° F., for each inhalation.

For touching or painting the inflamed surface, use

380. R. Acidi tannici, $\frac{3}{4}$ ss
 Glycerinæ, $\frac{3}{4}$ iv.

Or, when ulceration is present:

381. R. Sodii biborat., $\frac{3}{4}$ j
 Glycerinæ, $\frac{3}{4}$ ij. M.

DR. J. H. GARNER, CANADA.

In all varieties of laryngitis, acute and chronic, in pharyngitis, commencing tonsillitis, etc., this practitioner uses *guaiacum*. (*Canada Lancet*, July, 1861.) In clergyman's sore throat, for instance, he would prescribe:

382. R. Tinct. guaiaci ammon., āā f. $\frac{3}{4}$ ij
 Liq. potassæ, f. $\frac{3}{4}$ ij
 Tinct. opii, ad f. $\frac{3}{4}$ vij. M.
 Aquam cinnamomi,
 Make a gargle. Use every hour. The liquor potassæ keeps the gum dissolved.

In the hoarseness and sore throat in young ladies, so commonly connected with menstrual troubles, this gargle does excellent service; or the pharynx and upper larynx may be swabbed with the pure ammoniated tincture. This produces a momentary asphyxia, but as soon as the patient gives a cough it is gone.

DR. ISAMBERT, PARIS.

In clergyman's sore throat, oratory, tobacco, and all alcoholic drinks,

must be interdicted. Gargles are of little avail. Pulverizers and inhalers are not much more useful. Direct local treatment to the walls of the larynx by means of a sponge, is the true plan. For this purpose, nothing equals solutions of zinc chlorides:

383. R. Zinci chloridi, parts $\frac{1}{2}$ -1
Aquæ, parts 100. M.

MR. LENNOX BROWNE, LONDON.

384. R. Potassii chloratis, gr. ij
Eucalypti extracti, gr. j
Pulveris cubebæ, gr. $\frac{1}{4}$
Confectionis fructus (fruit paste), q. s. M.

For one lozenge; has a joint astringent, sialogogue and expectorant action; valuable as a voice lozenge.

385. R. Morphiæ muriatis, gr. $\frac{1}{36}$
Ipecacuanhæ pulveris, gr. $\frac{1}{12}$
Confectionis, q. s. M.

For one lozenge; useful in allaying irritable cough, and assisting expectoration in laryngeal and bronchial catarrh.

In *syphilitic laryngitis* the specific treatment for syphilis, especially the mercurial treatment, is essential to success. Locally, the ulcers may be touched with dilute solutions of nitrate of silver.

In all cases of *syphilitic and tuberculous laryngitis*, the local application of a saturated solution of *nitrate of silver* in glycerine once in ten days has been recommended. The object is to substitute an acute for a chronic inflammation, and the inflammation caused by nitrate of silver recovers much quicker than that caused by most of the other caustics. Then use a spray or gargle of common salt-water three or four times a day. Occasionally an antiseptic should be added, and the best are said to be oil of cinnamon, winter-green, pepper, etc.

NOTES ON REMEDIES.

Acidum Carbolicum. The inhalation of pulverized solutions of carbolic acid (gtt. j-v to f. $\bar{3}$ j. aq., with a few drops of glycerine added,) is frequently of service.

Alumen is valuable as a gargle (℞j-iiij to aquæ f. $\bar{3}$ j) on the decline of the inflammatory symptoms in the acute, and generally in the chronic forms.

Ammonii Chloridum. The following is used by Dr. DA COSTA:

386. R. Ammonii chloridi, gr. x-xx
Aquæ destillatæ, f. $\bar{3}$ j. M.

Use with any form of steam atomizer, throwing a fine spray, in *laryngeal catarrh*, acute as well as chronic. The dose best borne is not above ten grains to the ounce, although as much as two drachms to the ounce have been employed.

Argenti Nitras may be used in inhalation (gr. v-x to f. $\bar{3}$ j) as an abortant in the early stages. In the ulcerated forms of tubercular or syphilitic origin, a strong solution, $\bar{3}$ j to water f. $\bar{3}$ j, is the most efficient of all applications.

Balsamum Peruvianum. The recommendation originally made by HOFFMAN, in chronic inflammation of the larynx, of diffusing the vapors of the balsam of Peru, or of benzoin or tolu, throughout the air of the patient's room, by throwing the drug upon hot coals, is an excellent one. Or, the vapor may be inhaled through the inhaler.

Benzoinum is of service in strumous laryngitis. The following formula may be ordered :

387. R.	Tincturæ benzoini compositæ,		
	Mucilaginis acaciæ,		
	Syrupi papaveris,	āā	f. $\bar{3}$ j
	Aquæ cinnamomi,		f. $\bar{3}$ vj. M.

For one dose.

Or :

388. R.	Tint. benz. comp.,		f. $\bar{3}$ ij
	Glycerinæ,		f. $\bar{3}$ ss
	Aquam,		ad f. $\bar{3}$ iv. M.

For a gargle.

Bismuthi Subnitrates has been found highly efficacious, applied by means of insufflation.

Catechu is an excellent ingredient for lozenges. As the pure gum is not unpleasant, a small piece dissolved in the mouth will be found efficacious in relaxed conditions of the membrane.

Collinsonia Canadensis, the stone root, is esteemed by eclectic practitioners a specific in clergyman's sore throat. (J. H. SCUDDER.)

Ferri Perchloridum is an extremely useful addition to inhalations and gargles in the follicular form.

Guaiacum may be regarded as the best of remedies in many inflammatory cases

Potassii Chloras. Saturated solutions of chlorate of potash are valuable gargles in all forms of laryngitis and pharyngitis. Some also recommend full doses internally in severe cases.

Quinine is recommended by Dr. E. BRIAND, of France, in a form of laryngitis which he has observed to be due to malarial poison.

Sodii Chloridum. Solutions of common salt, either as a gargle or inhalation, are of considerable service.

**Turpeth Mineral* (Hydrargyri Sulphas Flava) is highly extolled by Dr. C. F. MELSHEIMER (*Medical and Surgical Reporter*, Nov. 11, 1882).

Veratrum Viride. Dr. HANDFIELD JONES has used this drug very successfully in severe acute laryngitis. He gives \mathfrak{m} iv of the tincture every hour until the symptoms abate or vomiting is produced. In the latter case, it may be renewed after a few hours. (*Medical Times and Gazette*, January, 1875.)

Zinci Chloridum. (See p. 177.)

EXTERNAL MEASURES.

Counter-irritants are of benefit in chronic laryngitis. The following may be employed: *Tartar-emetic ointment* (not much used); *Subclavicular Setons* (three or four silk threads); pustulations of the skin of the throat by *Croton oil*.

Dilatation in laryngeal stenosis has been successfully employed by Prof. SCHROTTER, of Vienna. He has devised dilators for the purpose, (*Dobell's Reports*, 1876.)

Inhalations are largely employed. The following are the principal preparations used: *Ammonii Chloridum*; *Balsams of Peru and Tolu* (vapor generated by placing them upon hot iron); *Benzoinum* (vapor generated by throwing the drug upon hot coals, and thus impregnating the patient's apartment, or from a common inhaler); *Hydrocyanicum Acidum Dilutum*; *Iodinium*; *Olibanum* (vapor generated by placing the gum resin on burning coals); *Sodii Chloridum*; *Sulphurosum Acidum* (spray or fumigation); *Vapor of hot Vinegar*; *Steam of boiling water* (as much as possible by the nostrils).

Insufflations of tannin or alum are serviceable in the œdematous forms.

Tracheotomy should be performed without hesitation in threatening œdema of the larynx or glottis.

PHTHISIS.

This formidable disease usually presents itself in one of two forms: (1) acute miliary tuberculosis, or galloping consumption; (2) the insidious and much slower varieties known as chronic phthisis.

The treatment of the two kinds is broadly distinct. Acute phthisis, if cured at all, must be by immediate and active medication; while the management of chronic phthisis is largely hygienic and dietetic, and the exhibition of drugs plays but a subordinate part.

ACUTE PHTHISIS.

DR. MCCALL ANDERSON, GLASGOW.

This practitioner claims to have treated, with success, several cases of this most terrible form of the disease. (*Lectures on Clinical Medicine*, 1877.) He feeds the patient every hour or half hour with milk, soup, or other light nourishment, and gives four to six ounces of

brandy daily. The night sweats he checks by the hypodermic injection of $\frac{1}{16}$ grain of atropia nightly. As a stimulating expectorant:

388. R.	Ammonii chloridi,	$\mathfrak{z}\text{j}$	
	Syr. scillæ,	f. $\mathfrak{z}\text{ iij}$	
	Infus. senegæ,	f. $\mathfrak{z}\text{ vj}$.	M.

A tablespoonful three times a day.

Should the febrile symptoms continue unabated, the following, which is "Niemeyer's antipyretic powder," may be ordered:

389. R.	Quiniæ sulphatis,		
	Pulv. digitalis,	$\bar{\text{a}}\bar{\text{a}}$	gr. xij
	Pulv. opii,		gr. vj. M.

Make twelve powders. One every four or six hours.

The temperature can also be promptly lowered by the application of iced cloths to the abdomen for half an hour, from time to time. To do this properly, the abdomen should be well exposed, and a double blanket placed under the patient. Two pieces of flannel are folded each into four layers, large enough to cover the whole front and sides of the abdomen. One of these is wrung out of ice water, and covered with a piece of dry flannel, while the other is left to soak in the ice water. The pieces are changed every one or two minutes. Two nurses are employed to do this. Of course, if there is any tendency to coldness or collapse, the procedure must be stopped.

CHRONIC PHTHISIS.

We shall first give some general plans of treatment, and later, that by special methods.

PROF. AUSTIN FLINT, M. D., NEW YORK.

After an analysis of 670 cases under his own care, in private practice, in this country, Dr. FLINT sums up the hygienic treatment as follows:

1. Many patients are benefited by change of climate; but the benefit depends more on the incidental circumstances of traveling than on any special climatic agency.

2. The most important hygienic measure is to change from indoor, sedentary habits, to outdoor, active ones.

3. Sea voyages are often of great service, but here, too, the benefit is from the accessory circumstances.

4. The appetite and taste of the patient should suggest the food, and also the use of alcoholic drinks.

5. Daily spongings, followed by frictions and moderate gymnastic exercises, are useful.

6. *Medicines.* Cod-liver oil benefits about a fourth of the cases; the hypophosphites about the same; alcohol checks the disease more or less, in a third of the cases. *Pancreatic emulsion* is useful where the dejections contain floating fat. It is better tolerated than pure fat or oil.

PROF. VON NIEMEYER.

In the treatment of phthisis, fever is the symptom which principally demands attention whenever it persists at all severely, in spite of the remedies directed against the main disease. *Digitalis* and *quinia* have a well-merited reputation as means of arresting the abnormal calorification, and reducing the animal heat, in spite of the continuation of the disease.

390. R.	Pulveris digitalis,	gr. x	
	Pulveris ipecacuanhæ,		
	Pulveris opii,	āā gr. v	
	Extract. helenii,	q. s.	M.

For twenty pills. One three times a day.

Add quiniæ sulphatis, ℥j, to the above prescription when the type assumed by the fever becomes periodical, the evening exacerbations severe, and the chills by which they are ushered in pronounced.

Our author is so much in the habit of using this (known as Heim's) pill, with or without quinine, in consumption, when the fever proves refractory to other remedies, that it has become a very common prescription at his clinic. The exhibition is suspended whenever a distinct reduction of the temperature and of the frequency of the pulse becomes apparent, and is resumed so soon as the effect subsides. Patients pretty soon learn to judge for themselves, when it is time to stop the pills and when to resume them.

A serviceable means of distinction between the cases in which *cod-liver oil* is indicated and those in which nothing is to be expected from it, is afforded by the symptoms of the torpid and erethritic forms of *crofula*. When the patient's slender frame, the lack of fat beneath his skin, and his accelerated pulse, warrant the belief that his nervous system is in a state of over-activity, cod-liver oil is generally of most signal benefit. Under its use, the plumpness of the body increases, while the general susceptibility of the system, and the diseases consequent upon it, subside. These are the cases to which this article owes

its name as an anti-scorfulous remedy. But if the patient be clumsy and thick-set; if the nose and upper lip be enlarged, and the adipose layer over the rest of the body be strongly developed; if the action of the heart be retarded rather than accelerated; if the irritability of the nervous system seem unusually obtuse; in short, should there be reason to suppose that the waste of the system is diminished rather than increased, we cannot hope to relieve the disease by means of the oil.

DR. L. BURNEY YEO, M. D., M. R. C. P.,

Assistant Physician to the Brompton Hospital for Diseases of the Chest, England. This writer, whose opportunities for the study of phthisis are unusually large, has lately given his experience on some modern remedial measures in this disease; to wit:

- a. The use of the *alkaline hypophosphites*.
- b. The use of antiseptic inhalations, and of the sulpho-carbolates internally.
- c. The treatment of phthisis by rest (local rest).
- d. The removal to mountain climates.

In regard to the *hypophosphites*, he believes they have *no claim whatever* to be regarded as specific remedies for tubercular disease; but they are of considerable value in a certain limited class of cases.

In advanced phthisis, with both lungs involved, the hypophosphites are of no use. Nor are they of use in less advanced cases which are obviously running a rapid course, and are attended with much fever and cachexia. In short, in those cases which we have always been accustomed to regard as hopeless and beyond the reach of all remedies, the hypophosphites are of no avail. They are less successful with dark persons of sanguine or bilious temperament than with fair persons of phlegmatic temperament. They are much more useful in young than in older persons. The greatest benefit arises from their use, in the first place, in children in all forms of chronic lung disease; and, in the second place, in young adults of fair complexion, not very cachectic or emaciated, and with disease limited to the upper part of one lung. In the cases in which they do good, their beneficial effect is generally noticed almost immediately. The patients usually say they feel very much better "in themselves;" they feel stronger, in better spirits, are more active, eat better and sleep better. The night-sweats disappear, the cough sometimes disappears also; but it will often be observed

that, while there is this general amelioration and a subjective feeling of improvement, the cough will continue unrelieved, and even occasionally become more troublesome. Notwithstanding the evident improvement in general health, the physical signs often remain the same, and even may increase in extent, while the patient has been feeling so much better that he has resumed his occupation, and declared that he felt "as well as ever he did in his life." The improvement which follows the use of the hypophosphites is frequently only of temporary duration; this is especially the case with hospital out-patients. It is much less so with private patients, who can be more carefully watched and more thoroughly protected from adverse circumstances.

Sometimes all the symptoms of phthisis disappear during the use of the hypophosphites, and the general condition as well as the physical signs undergo rapid amelioration. The patient has been, apparently, quite well, but an imprudent visit to a theatre has resulted in a fresh catarrh, and the disease has again advanced with unusual rapidity; the former remedies are now quite useless, and, in a few months, the case ends fatally. It has, indeed, seemed that, in some of these cases where the hypophosphites led to temporary arrest of the disease, the subsequent advance has been unusually rapid and quickly fatal. Patients who are mending under the influence of these remedies should be protected with more than usual care against all those conditions which may possibly lead to a relapse.

Of the *antiseptic inhalations*. LEMAIRE's observations on the inhalation of carbolic acid may be briefly summarized as follows. It must be remembered that he gave it also internally in aqueous solution. He found, he says, very remarkable effects follow its use.

There was diminution of cough after twenty-four hours, in some cases almost a complete disappearance after a few days. The expectoration was diminished or almost suppressed, and, if the sputa were offensive, their fœtor disappeared. In many the physical condition of the respiratory organs was ameliorated. Two were cured; in others, there was a subsidence or disappearance of *râles*, and parts became pervious to air which had previously been impervious. In cases in the second stage, he had noticed increase of strength, return of appetite and sleep, increased freedom of breathing, and general exhilaration. The amendment, however, did not persist in all cases.

Dr. SANSOM also offers his testimony as to the "real value" of the

“dry inhalation of carbolized air” in phthisis. This excellent observer has placed on record many cases in which the internal administration of the sulpho-carbolate appeared to be of great service. These antiseptic inhalations have received new impetus by the recent discoveries of KOCH, which will be mentioned on a later page.

The inhalation of balsamic vapors—frankincense, turpentine, storax, etc.—in phthisis, is of very ancient date, and its value was from time to time strongly insisted upon by the older physicians. Again and again has the dilute vapor of tar been advocated as of much use in pulmonary consumption.

There is a peculiar nauseating odor in the breath of many phthisical patients, even before the development of marked physical signs. Of course, if antiseptic inhalations are to be of use, they must be frequently employed, and means must be adopted which will insure the antiseptic vapor reaching the whole of the diseased portion of the pulmonary tissues. The vapor must be evenly diffused through the atmosphere the patient breathes.

In regard to *rest*, Dr. YEO wholly doubts that enforced inactivity of the respiratory function is advantageous in phthisis generally. When complicated with an acute attack of pneumonia or pleurisy, limitation of the respiratory movements of the chest-wall may be of temporary advantage; but those who advocate the establishment of a sort of pulmonary anæmia, as a means of checking the advance of tubercular phthisis, are resting on a false analogy and an erroneous pathology.

As for mountain air, there is no doubt it benefits a large class of patients in the earlier stages. But cases of very advanced disease, of profound cachexia, cases with active febrile symptoms, are unsuited for such migration. All who have seen much of phthisis know well enough that a great number of cases never do so well as in the quiet and comfort of home. But a restlessness seizes on many consumptive patients, even in advanced stages of the disease, and they *will not* remain at home. For such cases, a southern watering-place is better suited than a mountain valley. The rarefied air of elevated regions is quite unsuited to cases in which large tracts of lung tissue are disorganized or hopelessly disabled.

DR. MARY HALLOWELL, OF PENNA.

This writer (*Med. and Surg. Reporter*, 1884,) makes some useful

suggestions for treating the symptoms of the advanced stages of phthisis.

If associated with dyspnœa, arising from tenacious catarrhal products in the tubes, ammon. carb. is indicated, given as follows:

391. R.	Ammon. carb.,	gr. v	
	Vin. picis liq.,	gtt. xv	
	Syr. acaciæ,	gtt. xx	
	Aq. anisi,	q. s. ad. f. ʒj.	M.

S.—Take every three hours.

The syrup of acacia disguises the carbonate of ammonia, and also has a beneficial influence on the mucous membrane.

The syrup of wild cherry may be substituted for the tar when a tonic and sedative is indicated, in place of the more stimulating tea. If the expectoration be scanty and the cough spasmodic, troches of morph. and ipecac (containing gr. $\frac{1}{8}$ of morph.,) generally afford marvellous results.

Oxalate of cerium is beneficial when violent coughing with free expectoration exists with decided gastric disturbance.

It not only quiets the stomach and lessens expectoration, but also exerts a striking influence in diminishing the severity of the cough.

It does not operate well when expectoration is scanty, and because of its tendency to dry the air passages cannot be continued with advantage longer than a few days at a time.

Inhalations are sometimes serviceable, and with some physicians are the sole remedy employed.

The list of agents which can thus be administered possesses the advantage of medicating the air-passages directly, without disordering the stomach in passing through the general circulation, and whose virtues are not to be measured by their merely antiseptic qualities.

Of these we have lime and belladonna (gr. j.—ʒiv.), or if the secretions be fetid, carbolic acid may be added, after the combination used by Dr. COGHILL, of London, which is composed of

392. R.	Creasoti,	
	Acid carbolicæ,	āā ʒj
	Tr. iodid. co.,	f. ʒ ij
	Alcohol,	f. ʒ iss.

This diminishes the quantity of expectoration, allays cough, lessens, or entirely destroys, fetor of breath.

It may be administered by an extemporized respirateur, made of a cylindrical box, with a perforated cap, in which is placed the absorbent cotton on which the solution is dropped. To be inhaled once or twice daily, 10 to 15 minutes.

In the later stages, when the cough is long-continued and interferes with sleep, acetate of morph. may be given with tr. sanguinariae.

The pains in the chest may be dissipated by counter-irritation in the form of tr. iodine and croton oil, or mustard or belladonna plaster. Sometimes a flannel bandage worn around the waist affords a relief not to be obtained from drugs.

The fever requires special treatment. This symptom will not entirely disappear until the substances which cause it are entirely expelled from the body, and for the purpose of abating it, quinia and digitalis are the main treatment; but neither should be given continuously, on account of their injurious influence on the stomach.

Our main reliance, therefore, should be diet and rest of body. Food should consist principally of milk, boiled grains, and stewed fruit, with a little bread.

Salicin, in doses of gr. xx., two hours before the onset, sometimes acts like a charm.

Sponging the surface with whisky and alum is often beneficial. Tablespoonful of whisky at bedtime and another at period of awakening may be given with advantage. If medicines are to be employed, we have a favorite combination in ext. bellad., gr. $\frac{1}{4}$, in pill-form, with oxide of zinc, gr. ij. Give one at bedtime.

PROFESSOR JACCOUD, PARIS.

The following review of the drugs most lauded in phthisis is given by this distinguished authority:

Arsenic.—In chronic phthisis, as soon as the phenomena attributed to anæmia have been remedied by iron, or in the absence of amelioration after a delay of a couple of months, the author abandons ferruginous preparations, and has recourse to arsenical medication. He employs exclusively granules of arsenious acid, of gr. $\frac{1}{16}$ each, and administers them at the commencement of each of the two principal meals. He commences with two a day, and every week increases two a day, until eight or ten a day are given. When this maximum dose is attained, he maintains it indefinitely, unless phenomena of intolerance make their

appearance, such as cramps, inflammation of the eyes, cutaneous eruptions, vomitings, diarrhoea. Then, however, he does not stop the medicine, but diminishes temporarily the dose, and returns, as soon as possible, to the maximum of toleration.

Arsenic acts powerfully upon the nutritive processes in chronic pulmonary phthisis. It quiets the nervous excitement, and possesses an anti-febrile action sufficiently marked to effectually combat the evening intermittent fever.

So long as the disease remains chronic, and does not develop acute attacks of pseudo-continuous fever, the arsenical medication ought to be maintained at the maximum of toleration, conjointly with the special regimen, cod-liver oil, cinchona and alcoholized raw meat.

Cod-liver Oil.—When the cod-liver oil is badly supported alone, in chronic pulmonary phthisis, the author adds to it brandy, rum, kirsch or whisky, in the proportion of two parts of the oil to one of the spirit, and recommends the patient to close the nostrils at the moment of swallowing it. He commences by small doses of the oil and spirit mixture, then two tablespoonfuls a day, and advances to five or six fluid ounces a day.

Iron.—The author reserves the employment of ferruginous preparations for those cases where the patients present the certain signs of globular anæmia. He gives, in preference, the syrup of the iodide of iron to young children, and the pills of the iodide of iron in other cases. From four to eight of these pills are to be taken a day, at the commencement of the meals. He advises not to administer the iron where there has been recent hæmoptysis.

Raw Meat.—In the treatment of patients with whom the physician is obliged to dissimulate, he may order the pulp of raw fillet of beef, well deprived of all fibrous matter, in some unleavened bread, under the pretext of administering medicinal boluses. When he can act openly, the author prefers to mix the pulp of the raw meat, properly prepared with brandy, rum or whisky, in such a manner as to obtain a dough-like mass, which the patient takes with sugar or salt, according to taste, in tablespoonful doses during the day. This mixture is very palatable, and does not produce that rapid intolerance which is often observed with other meat preparations.

Counter-irritation.—The author, in acute phthisis, attacks the pulmonary disorder by means of large flying blisters, renewed without in-

terruption. Instead of the ordinary dressing, he covers the vesicated surface, and envelops it in every direction, a finger's breadth, by diachylon plaster. On removing the protecting plaster at the end of four days, the cicatrization is found complete.

In commencing chronic phthisis, the author applies, under the clavicle of one or both sides, cauteries of Vienna paste, of the size of a sixpence at the maximum, and repeats these punctiform cauterizations as long as any favorable influence is noticed.

COD-LIVER OIL TREATMENT.

In addition to what has been said above, we quote from various specialists their directions for the exhibition of this important remedy.

DR. C. J. B. WILLIAMS, LONDON,

gives the following minute rules for administering the oil: 1. The *best time* for administering it is immediately after, or to those who prefer it, at or before, a solid meal. 2. In cases of *peculiar weakness of stomach*, with tendency to retching or vomiting, strychnia (gr. $\frac{1}{2}$ — $\frac{1}{4}$) proves a valuable adjunct. So does salicine. Neither of these, though a powerful tonic, has any of the heating properties of quinine or iron. If their bitter taste is objected to, they may be given in the form of pill, after or before the oil. 3. It may be advantageously *combined with a mineral acid*. Nitric acid is best in inflammatory cases, and in those attended with much lithic deposit in the urine, but its tendency to injure the teeth is an objection to its long-continued use. Sulphuric acid is more eligible where there is liability to hæmoptysis, profuse sweats, or diarrhœa. But in most cases, for long continuance, diluted phosphoric acid is preferable to either. 4. The *bulk of the whole dose* of the oil and vehicle should be so small that it may be swallowed at a single draught; therefore, the vehicle should not exceed a tablespoonful, with at first a teaspoonful of the oil, to be gradually increased to a tablespoonful. 5. The *dose* of the oil should rarely exceed a tablespoonful, twice or thrice daily. Larger quantities either derange the stomach or liver, or some of it passes unabsorbed by the bowels. 6. The *diet* requires attention. With some persons the oil agrees so well, and so much improves their digestive powers, that they require few or no restrictions in diet; but this is not the case with the majority. The richness of the oil proves more or less a trial, sooner or later, to some persons; therefore it becomes proper to omit or reduce all other rich and

greasy articles of food. All pastry, fat meat, rich stuffing, etc., should be avoided, and a great moderation observed in the use of butter, cream, and very sweet things. Even new milk in any quantity is not generally well borne during a course of oil, and we may find malt liquor too heavy, increasing the tendency to bilious attacks. A plain nutritious diet of bread, fresh meat, poultry, game, with a fair proportion of vegetables, and a little fruit, and only a moderate quantity of liquid at the earlier meals, commonly agrees best. 7. Should a bilious *attack* come on, indicated by nausea, headache, furred tongue, offensive eructations, high-colored urine, etc., it is necessary to suspend the oil, lighten the diet, and give an aperient.

In regard to taking the oil,

DR. J. M. DA COSTA, PHILADELPHIA,

uses the following combination :

393. R.	Olei morrhuae,	f. $\frac{N}{2}$ j	
	Aquæ menthæ piperitæ,	f. $\frac{N}{2}$ ss	
	Tincturæ aurantii,	f. $\frac{N}{2}$ ss	
	Misturæ acaciæ,	f. $\frac{N}{2}$ iijss	
	Olei gaultheriæ,	m x.	M.

A dessertspoonful three times a day. This formula disguises somewhat the taste of the cod-liver oil.

Or :

394. R.	Olei morrhuae,	f. $\frac{3}{4}$ ss.
---------	----------------	----------------------

For one dose ; to be taken three times a day in *carbonic acid water*.

Place in a tumbler a small amount of any preferred syrup, (orgeat or sarsaparilla is the best adapted to disguise the taste of the oil,) and fill up with carbonic acid water, from a bottle furnished with a siphon for table use ; then, while it is still foaming, put in a tablespoonful of the oil. It is astonishing how perfectly the taste is concealed in this manner. Other modes of taking the oil are, floating on ice water, in lemon juice, and in the froth of porter. Less than a tablespoonful is not worth taking. The best time for its administration is between meals, after the process of digestion is pretty well finished. Persons are exceptionally met with who take it in preference just before meals, and thus avoid the disagreeable eructations. Nobody likes to take it the instant after meals.

DR. THOMAS H. TANNER

suggests that in cases of phthisis where the stomach will not tolerate any form of cod-liver oil, resort may be had to *cod-liver oil embrocations*:

395. R. Olei morrhuae, f. $\frac{3}{4}$ ijss
 Spiritus ammoniæ aromatici, f. $\frac{3}{4}$ j
 Tincturæ opii, f. $\frac{3}{4}$ ss
 Olei lavandulæ, ℥xxx. M.
 One-half to be rubbed over the chest and abdomen, night and morning.

Or:

396. R. Olei morrhuae, f. $\frac{3}{4}$ j
 Olei cajuputi, f. $\frac{3}{4}$ j. M.
 To be rubbed over the chest at bedtime, and applied by means of lint well saturated with it. The cajuput oil well disguises the smell of the embrocation.

It seems to be well settled that the assimilation of the oil is materially hastened by combining it with *ether*. The following is the prescription of

PROF. BALTHAZAR W. FOSTER, M. D., LONDON.

397. R. Ætheris, f. $\frac{3}{4}$ ijss
 Olei morrhuae, f. $\frac{3}{4}$ iv. M.
 Dessertspoonful thrice daily, before meals.

He gives many cases where this combination acted much more efficiently than the oil alone. (*Lectures on Clinical Medicine*, London, 1874.)

TREATMENT BY HYPOPHOSPHITES.

This special treatment is closely connected with the name of

J. F. CHURCHILL, M. D., PARIS.

His own formulæ, as published, are:

SOLUTION OF HYPOPHOSPHITE OF SODA.

398. R. Sodii hypophosphitis, 1 to 5 parts
 Aquæ destillatæ, 150 parts.
 Dissolve and filter. Dose, a teaspoonful three or four times a day, in sweetened water.

SYRUP OF HYPOPHOSPHITE OF SODA.

399. R.	Sodii hypophosphitis,	5 parts
	Sodii carbon. crystal.,	1 part
	Syrupi simplicis,	500 parts.

Dissolve, keep in a dark place. Dose, one to four tablespoonfuls four times a day.

SYRUP OF HYPOPHOSPHITE OF LIME.

400. R.	Calcii hypophosphitis,	1 part
	Aquæ destillat.,	30 parts
	Sacchari albi,	64 parts
	Aquæ calcis,	6 parts.

Dissolve with gentle heat; filter; keep in a dark place. Dose, teaspoonful three to twelve times a day.

Dr. CHURCHILL'S usual method of treatment is to give a tablespoonful of the syrup of hypophosphite of soda or lime (of the strength of six grains of the salt to the fluid ounce,) once or twice a day, in the absence of any inflammatory symptoms—sometimes a teaspoonful of syrup of hypophosphite of lime in the morning, and the same dose of syrup of hypophosphite of soda at night. For females, particularly persons of nervous temperament or delicate constitution, leading a sedentary life, and unused to physical exertion, the dose should be decreased one-half. For children from seven to fifteen, the dose is the same as for females. From two to seven years of age, one or two teaspoonfuls daily, of the syrups, are sufficient. For infants of a more tender age, the dose should seldom exceed a quarter or half a teaspoonful in twenty-four hours.

The syrups may be taken alone or in a wineglass of water, and should, in general, be exhibited at meal time, either before or after. They have no medicinal taste whatever.

Contra-indications. 1. The use of the hypophosphites ought not to be commenced during an acute inflammatory state. 2. Their employment should be temporarily suspended, if, during the course of the treatment the patient be attacked by any inflammatory complication but resumed as soon as acute complication shall have disappeared. 3. They should not be given during an active hæmoptysis of a passive form. 4. Their effects should be carefully watched, in the case of patients affected with disease of the heart. 5. The dose should not be too high, nor the administration too prolonged. Great mischief may result from incautiously pressing the remedy. Among the signs which

indicate that the hypophosphites have been given in excessive doses, are lassitude, wandering pains, headache, buzzing in the ears, vertigo, sudden loss of appetite and strength, and, above all, bleeding at the nose, however slight the loss of blood may be.

The following is the usual prescription of

JOHN C. THOROWGOOD, M. D., LONDON :

401. R. Sodii hypophosphitis, gr. v
Glycerinæ,
Aquæ, āā f. ʒss.
For one dose, thrice daily.

He believes that it is in cases of catarrhal pneumonia, limited to one lung, that we may expect satisfactory results from the hypophosphites. (*British Medical Journal*, May, 1877.)

Dr. J. G. S. COGHILL (*Lancet*, September 6, 1879,) recommends the following formula as suiting most patients well:

402. R. Sodii hypophos.,
Calcii hypophos., aa gr. viij
Infusi cascarillæ, ʒj.
To be taken twice a day, after meals.

Dr. C. denies that these salts exert any specific effect, but thinks they are valuable tonics. They are contra-indicated in advancing stages, with fever, but when the disease is arrested, they are often of decided advantage.

CALCIFICATION OF THE LUNG.

The process of calcification which the tubercles undergo in cases of spontaneous cure, have led various investigators to the endeavor to bring about this change in progressive cases.

Dr. JULIUS BOYER, of Paris, reported some years ago, (*Phthisie Pulmonaire*, Paris, 1874; *Dobell's Reports*, 1875,) several cases of alleged cure in this way by the administration of a salino-calcary powder, composed of phosphate and carbonate of lime, and of bicarbonate of soda.

The same plan has been advocated by Dr. CARL BOTH, of New York. But no satisfactory reports of this treatment have been published.

THE SPECIFIC OR ANTI-PARASITIC TREATMENT OF TUBERCULOSIS.

Since the presence of the *bacillus tuberculosis* was demonstrated by KOCH and others, and this fatal disease attributed to its action, many experiments have been made with a view to check the malady by the administration of germitoxic agents.

Some of the most carefully conducted observations were those of Dr. HILLER, of Berlin. He placed ninety-one patients on this treatment, all of them being cases which still held out some hopes of recovery.

There were employed, (1) corrosive sublimate, (2) iodoform, (3) bromine, (4) ethylic alcohol, (5) methylic alcohol, (6) sulphuretted hydrogen, (7) arsenious acid, (8) boric acid, (9) salicylic acid. These remedies were given as inhalations in the form of gas or vapor or a medicated spray, by parenchymatous and subcutaneous injections, and by the mouth. Whatever the drug used, and whatever way administered, the tubercular process was not checked, nor did the bacilli disappear from the sputa.

Treatment by *corrosive sublimate* gave the best results; but the patients thus treated were benefited by nutritious food, cod-liver oil, brandy, and acid phosphate of lime. In every case the medicine was administered hypodermically, in daily doses of .001—.0015. Forty-four cases were thus treated; and, with more than half, the treatment had to be given up, either because the patient dreaded the pain caused by the injection, or because the medication caused accidents to be spoken of later.

Of nineteen patients where the treatment could be fully carried out,—i. e., forty injections in the course of forty-five days,—five died in a few weeks. The other twelve showed an improvement, which HILLER called apparent only, but which allowed them to leave the hospital. He treated chiefly tuberculous patients with lesions of the apices. The general condition, satisfactory to start with, was maintained, and in some cases improved. The stethoscopic signs improved, and the expectoration diminished. The tubercular bacilli became less numerous in the sputum.

In some cases, injections of sublimate are badly borne, causing stomatitis, salivation, indigestion, a bad taste in the mouth, diarrhoea, tremors, insomnia, prostration, etc. In advanced cases, bedridden and feverish patients, sublimate is deleterious, produces prostration, and hastens the fatal termination.

In two cases of tuberculous laryngitis, with very slight implication of the lungs, HILLER used inhalations of an atomized sublimate solution, with no good result.

In three other cases, he practised injections of sublimate into the pulmonary parenchyma. With a syringe he injected each day 2 cc. of a one-tenth-per-cent. solution through an intercostal space. Sublimate is exceedingly irritating to the lung; and the inevitable contact of the medicated solution with the bronchial mucous membrane sets up violent fits of coughing, followed in one case by serious hæmoptysis.

Dr. ALBRECHT,* of Berlin, also submitted his patients to methodical inhalations of *pure oxygen*, without changing their usual diet, and he claims to have checked the progress of the disease; the patients stop losing weight, and perhaps even gain; and KOCH's bacilli become less numerous in the sputum. Guinea-pigs inoculated with tubercle were submitted to inhalations of oxygen as soon as the disease appeared developed. Those having oxygen lived six months; those not having it, four months.

Of germicides for this purpose, *thymol* has been advocated by some. It may be employed in the form of

WARREN'S INHALATION.

403. R.	Thymol,	gr. 8
	Sodii boratis,	gr. 300
	Glycerini,	fl. oz. 1 $\frac{1}{4}$
	Aquæ camphoræ,	fl. oz. 2 $\frac{1}{2}$
	Aquæ picis,	fl. oz. 7.

To be used as an inhalation, by means of the atomizer.

The use of *aluminium* as a bacteriacide has been advocated by Dr. JULIUS PICK (*Med. Record*, 1883), who claims to have effected some cures with it. He administers about gr. iss daily, continuing it with inunction of vegetable oils and supporting diet.

A most experienced specialist, Dr. C. THEODORE WILLIAMS, writes of this plan of treatment in the *Lancet*, 1883. He refers to the great importance of first studying the conditions of life of these low organisms outside of the human body, and then of testing the effects of various antiseptic drugs on them. In this way we may discover substances fatal to the bacillus, and then proceed to devise methods of applying

* *Deut. Med. Woch.*, No. 29, 1883.

them, so as to reach and attack the enemy in his strongholds of the human body. But this knowledge can only be attained by long and careful investigation and the work of many patient observers.

Another mode of antiseptic treatment is by hot water or steam inhalations of various kinds, some form of inhaler being used, and the different drugs kept at a temperature suitable to promote their vaporization. In this way, we at any rate do not get the drug inhaled to a certain extent. The best form of this treatment seems to be jets of steam spray charged by means of capillary tubes with necessary medicaments. The patients receive a good deal of the drug into their bronchial tubes and lungs in a short time, owing to the force of the steam current; but the objection lies in the damp and hot atmosphere which it causes, and the inexpediency of subjecting the patient for any length of time to such strong measures.

The third method is by diffusing through the air of a chamber medicated vapors. In this way consumptive cases can be kept under the influence of special drugs for long periods. In this method we only imitate some varieties of climate, such as those of the sea coasts, of pine woods, and of sulphur springs, or in the neighborhood of volcanoes. Rooms are set apart for this purpose, and are kept specially impregnated with the vapor of iodine or chlorine.

CLIMATES AND TRAVEL.

The benefit of change of climate is most perceptible in cases in overworked persons of sedentary life, who leave their indoor habits at home when they travel, and in hemorrhagic cases.

In ordinary chronic phthisis a dry climate is best; in phthisis of catarrhal origin, a moist air is preferable. (Dr. C. T. WILLIAMS.)

The most important of all considerations is that the air shall be pure, *i. e.*, containing plenty of oxygen and free from organic impurities.

As a general rule, according to Dr. MADDEN (*Health Resorts of Europe and Africa*, 1876), in the first and early second stage of ordinary cases of phthisis a dry and stimulating climate is preferable. When the disease is of a chronic, bronchial type, with irritative symptoms, a moist, equable, unstimulating climate will be most beneficial. All patients in the third stage should stay at home.

It is the advice of Dr. A. L. LOOMIS, (*Diseases of the Respiratory Organs*, 1875,) to be governed in the choice of a climate by inquiring

of the patient whether his health has usually been better in warm or cold weather, in a dry or moist atmosphere ; and to direct his travels accordingly, without much reference to the particular form of his disease.

Two general rules should be observed: (1) In whatever locality a phthisical patient finds himself improving, he should remain there. (2) Whenever he finds his health deteriorating in a locality, he should leave it.

Special praise has of late years been given to *altitude* as a curative agent in phthisis. Dr. LOMBARD, of Geneva, even claims that the disease cannot arise in localities over 5,000 feet above the sea level. Certain it is that not only are residents of lofty plateaux singularly exempt, but patients seeking these elevated regions are often singularly improved. This has led to the establishment of mountain sanatoria in Europe and America, the benefit of a residence in which has been amply demonstrated.

In regard to *sea voyages*, Dr. RODERICK MACLAREN (*Practitioner*, 1877,) states that when the infiltration of the lung is localized and chronic, when it is confined to the one side, and there is absence of evidence of disease elsewhere, it may be looked on as a favorable case ; and the further the departure from this condition, the less are the patient's chances of benefit. It is stated that cases in which hæmoptysis is the first, or at least a very early symptom, do especially well. Cases are unsuitable for a voyage in which there are signs of extensive progressive excavation, and all in which there is great muscular weakness. When we have evidence of the disease being general, as shown by glandular, bowel, or laryngeal disease, our recommendation should certainly be withheld. The bowels in particular should be in good trim, for the sedentary life and peculiar diet of a ship are apt, in healthy people, to produce intestinal disturbance, (constipation and subsequent diarrhœa, or either of these alone.) In a case which persistently grows worse under favorable circumstances of habitation, etc., on shore, the prospect of benefit would be small. A phlegmatic habit of body should also be looked upon, to a considerable extent, as a counter-indication for this remedy.

The following list embraces the most important health resorts in phthisis :

Mountain Sanatoria.

Europe: Davos, Moritz, Lamaden, in Switzerland; Hohwald, Wangenburg, Soultzmatt, in the Vosges; Chaumont, Chateau, Ranz, in the Jura; Gerbersdorf, in Silesia.

America: The Andes of Bolivia; the higher levels in Colorado; the Appalachian plateau about Asheville, N. C.

Lower Dry Climates.

Europe, Asia and Africa: Montpellier, Cannes, (warm, moderately exciting); Nice, (variable, exciting); Mentone and the Riviera, (cold and changeable); Upper Egypt, (the best tonic, dry, warm climate); the Kirghis Steppes.

America: The Costa Rican and Mexican plateau; Minnesota, (dry and exciting); Colorado about Denver; the Adirondack region, (cool, moderately dry, very good at all seasons); Aiken, S. C., and northern Georgia; Santa Barbara, Cal., (dry, warm.)

Lower Damp Climates.

Europe, etc.: Madeira, (very moist, equable); Arcachon; Southern Italy, (now condemned); Algiers, (warm, moist, equable.)

America: The Bermudas, (doubtful); the Bahamas, (moist and equable); Florida, (moist, but changeable); Isle of Pines, Santa Cruz, etc., (very equable and moist.)

TREATMENT OF COMPLICATIONS.

THE ANÆMIA AND DEBILITY.

According to Dr. SYMES THOMPSON (*Practitioner*, September, 1868,) the *tincture of the chloride of iron* is far superior to other chalybeates. The improvement of appetite, diminution of flatulence, etc., which occur under its administration, are often remarkable, cod-liver oil and other fats, refused previously, being digested without discomfort. It both checks diarrhœa and relieves constipation, by giving tone to the feeble muscular fibres of the bowels; it lessens night sweats, though these often call for oxide of zinc in addition, and is a valuable remedy in hæmoptysis. Many believe the *iodide of iron* one of the best chalybeate tonics which can be employed in the anæmia of phthisis, if it do not prove too stimulating. The syrup (℞xx-f.ʒj, thrice daily,) is the best form.

The *iodide of calcium* has been extolled as an excellent tonic by Dr. A. MALET, of Rio de Janeiro. (*Bull. Gén. de Thér.*, Aug., 1868.) He administers gr. viij in solution, in divided doses, daily, and says it awakens the appetite, regulates the digestion, renders the respiration freer and deeper, diminishes the cough and modifies the expectoration, increases the strength and lessens the perspiration. If constipation ensue in consequence of its long use, it will often disappear of itself; otherwise the dose may be diminished, or the remedy temporarily stopped.

Prof. FURTER, of Montpellier, employs, with good results, raw mutton or beef, given in conjunction with strong or diluted alcohol, in small doses.

405. Take some finely-minced raw mutton or beef, and roll it up in sugar or in a saccharine electuary. Give in teaspoonful doses to the amount of 3-10 ounces per day.

406. R. Alcoholis, f. ℥j
Syrupi, f. ℥iij. M.

A teaspoonful dose every hour. The dose and frequency of administration are to be modified by the patient's individuality.

Dr. J. WARING CURRAN praises the *oxide of zinc*.

407. R. Zinci oxidi, gr. ij
Extracti conii, gr. j. M.

For one pill; to be taken three times a day. The quantity of the oxide of zinc is gradually to be increased.

In the latter stages of phthisis, where profuse sweating and colliquative diarrhœa harass the patient and rapidly lower the vital capacity, this combination is very effective. It is of great value, also, in the earlier stages of the disease. It seems to steady the nervous system, and act as a sedative to the wandering pains.

THE COUGH.

Anodynes are urgently demanded when the cough is troublesome. The following is recommended by Dr. SUNDERLIND:

408. R. Ammonii chloridi, gr. xxx
Opii pulveris, gr. ix
Digitalis pulveris, ââ
Scillæ pulveris, gr. xv. M.

Divide into thirty pills. One every six hours, to quiet the cough and facilitate the expectoration in the early stage of phthisis.

409. R. Morphiae sulphatis, gr. j
Syrupi ferri iodidi, f. $\frac{3}{4}$ ss
Glycerinae, f. $\frac{3}{4}$ ijss.

A teaspoonful two or three times a day, to quiet the cough and retard the emaciation in phthisis.

The following is in use in most chronic pulmonary affections, in the Charity Hospital, New York:

410. R. Potassii bromidi, āā $\frac{3}{4}$ iss
Potassii chloratis, f. $\frac{3}{4}$ iv. M.
Ammoniae muriatis,
Syrupi tolutani,

A teaspoonful every few hours.

A writer in the *Canada Lancet*, March, 1879, says the following formula is almost a specific in phthisical cases with troublesome coughing.

411. R. Atropiae sulph., gr. j
Morphiae sulph., gr. viij
Acid. sulph. aromat., $\frac{3}{4}$ ij
Aquam menth. pip., ad $\frac{3}{4}$ j. M.

Dose, five drops thrice daily and at bedtime.

The Consumption Hospital, London, employs, as a house prescription:

412. R. Morphiae muriatis, gr. ss
Acidi hydrocyanici diluti, ℥xv
Acidi muriatici diluti, ℥ijss
Oxymellis scillae, f. $\frac{3}{4}$ ss
Aquam, q. s. ad f. $\frac{3}{4}$ j. M.

Dose, one to two drachms.

Combinations like the following are employed by Dr. J. M. DA COSTA:

413. R. Morphiae acetatis, gr. ij
Potassii cyanidi, gr. j
Acidi acetici, f. $\frac{3}{4}$ j
Extracti pruni virginianae fluidi, āā f. $\frac{3}{4}$ ij. M.
Misturae acaciae,

A teaspoonful four or six times a day, as a sedative mixture, for the cough of phthisis.

414. R. Liquoris morphiae sulphatis, f. $\frac{3}{4}$ j
Extracti pruni virginianae fluidi, f. $\frac{3}{4}$ ij
Acidi sulphurici diluti, f. $\frac{3}{4}$ ij. M.

A teaspoonful three or four times a day, when night sweats and cough are troublesome.

415. R. Extracti opii, gr. ss
Aquaë, f. ʒj. M.

For one inhalation twice a day, by means of any form of steam atomizer throwing a fine spray. In the irritative cough of phthisis, causing gastric irritability.

Dr. J. H. HUTCHINSON, of Philadelphia, recommends:

416. R. Acidi sulphurici dil.,
Tinct. opii deodoratæ, āā gtt. iij
Syr. pruni virginianæ,
Elixir. cinchonæ, āā ʒ ss. M.

Signa. Take at a dose.

Locally, various methods to allay the irritation, may be employed. The inhalation of warm water, impregnated with emollient herbs, such as *althæa officinalis*, or with narcotic extracts, palliates cough, dryness of throat, and laryngeal irritation. Inhalations of *tar*, *creasote*, and *carbolic acid*, simple, or in association with sedative extracts, have occasionally produced remarkable improvement in both the local and constitutional state. Also, hyoscyamus, gr. v of the extract to Oj of boiling water.

Dr. RINGER strongly recommends the application of the *glycerine of tannin*, with a small quantity of morphia added, to the throat. When the cough is troublesome, a good night's rest may often be obtained by applying it at bedtime.

For the distressing night cough, Dr. YEO finds *croton chloral* an efficient agent. It may be given in doses as high as sixty grains.

Dry cupping on the chest, under the clavicles, often gives great relief in distressing dyspnoea and cough.

Dr. SANDBY, of London, (*Brit. Med. Jour.*, April 12th, 1879,) has found nothing so good as *codeia*. It allays the cough without disturbing the digestive system. He prescribes gr. j., dissolved in syrup of tolu.

THE DIARRHŒA.

The complication of diarrhœa is not very frequent, but is exceedingly exhaustive. When caused by the presence of irritating ingesta, Dr. BARLOW uses:

417. R. Olei ricini, āā f. ʒ ij
Tincturæ rhei, miv
Tincturæ opii, f. ʒ iv. M.
Aquaë cinnamomi,

To be taken at a draught.

This may be followed by :

418. R.	Acidi nitrici diluti,	℥xij	
	Tincturæ opii,	℥v-x	
	Syrupi,	f. 3j	
	Aquæ cinnamomi,	f. 3x.	M.

Make a draught to be taken every fourth or sixth hour.

The logwood is a pleasant astringent. Dr. F. W. PARRY combines it thus :

419. R.	Extracti hæmatoxyli,	3ij	
	Misturæ cretæ,	3xij	
	Vini ipecacuanhæ,	f. 3ij	
	Vini opii,	f. 3j.	M.

A tablespoonful twice, thrice, or oftener daily.

Dr. GRAVES has recommended as one of the best remedies *nitrate of silver*, gr. j, in pill, three or four times daily. He also speaks well of :

420. R.	Acidi sulphurici diluti,	℥xv-xx	
	Tinct. opii,	℥x-xv	
	Tinct. cinchon. comp.,	ad f. 3j.	M.

For one dose.

THE DYSPNŒA.

The external use of dry cups or croton oil liniment to the surface of the chest and under the clavicles will sometimes relieve the distressing dyspnœa. When it is greatly aggravated by coughing spells, the inhalation of chloroform will often check these. Inhalations of hyoscyamus vapor, gr. v of the extract to Oj of warm water, is also recommended.

THE HÆMOPTYSIS.

The hæmoptysis of phthisis has already been considered under "Hæmoptysis," which see.

THE LARYNGITIS.

Tuberculous laryngitis has been mentioned under the general heading "Laryngitis," which see.

THE NIGHT SWEATS.

Nothing so effectually controls these as *belladonna* or *atropia*, as suggested by RINGER, FOTHERGILL and BARTHOLOW. The average dose of *atropia* is gr. $\frac{1}{16}$, which may be given in pill, or in solution ; or gr. $\frac{1}{16}$ hypodermically. It can very advantageously be combined with morphia. The aqueous solutions soon spoil, so the pill form is better.

Dr. A. H. HASSALL, of the Ventnor, England, Hospital for Consumption, gives :

421. R. Atropiæ sulphatis, gr. $\frac{1}{80}$ - $\frac{1}{60}$,
 Extracti gentianæ, gr. ij. M.
 For one pill.

Dr. J. T. NAIRNE (*British Medical Journal*, February, 1878,) uses the ordinary tincture of belladonna externally in these cases :

422. R. Tinct. belladonnæ,
 Spiritûs frumenti, ââ f. ʒ ij. M.

This is usually sufficient to cover the body and produce the desired effect.

Dr. THOMAS H. TANNER uses one of the following for night sweats :

423. R. Acidi gallici, ʒ ij
 Extracti cannabis indicæ, gr. v
 Confectionis rosæ, gr. x. M.
 For ten pills. One to be taken every night at bedtime.

Or :

424. R. Zinci oxidi, gr. xij
 Extracti conii
 vel
 Extracti hyoscyami, gr. xvij. M.
 For six pills. One to be taken every night at bedtime.

An excellent combination is :

425. R. Extracti belladonnæ,
 Zinci sulphatis,
 Extracti gentianæ, ââ gr. j.
 For one pill at night.

Or :

426. R. Zinci sulphatis, gr. j
 Extracti hyoscyami, gr. ij. M.
 For one pill at night.

Dr. FRAENTZEL recommends (*Wiener Med. Blätter*, July 5,) *hydrate of hyoscin* in the treatment of the night-sweating of phthisis, in subcutaneous doses of half a milligramme ($\frac{1}{40}$ gr.), or in pill, beginning with the same dose. He does not find it so universally beneficial as atropine, and it sooner exhibits a narcotic action; but he has seen

good results from it in cases where atropine has proved useless, or has failed to act after being in use for some time.

Speaking of the unsatisfactory results of all the remedies recommended for this condition, Dr. J. R. FORREST writes to the *Lancet*, October 27, 1883, that he has found the following a most efficacious lotion :

427. R.	Sulphate of zinc,	gr. iv
	Tincture of belladonna,	ʒj
	Water,	ʒj.

The body to be sponged with the lotion at bedtime.

Dr. HAYDEN believes that Dover's powder, if judiciously given is more efficient than belladonna. If the sweating occurs towards morning, as is usually the case, gr. v of Dover's powder, once or twice during the night, will check it. Tepid sponging with vinegar and water is a useful auxiliary. (*Dublin Journal of Med. Science*, April, 1877.)

Quinine is another useful remedy. It proves of most avail when there is a considerable rise of temperature at some period of the day. It is frequently given in two-grain doses, but five grains are much more likely to succeed. A large dose (8 or 10 grains) administered at once or in portions repeated hourly, is a good form.

Nitrite of amyl is a good remedy for night-sweats, but for promptness of action is decidedly inferior to atropia and other remedies.

Agaric, gr. v-vj, at night, is occasionally effectual.

Externally, sponging the surface with hot water (130° Fah.) is recommended by Dr. DRUITT; or with a strong solution of common salt. Aromatic vinegar and water is also useful, but has its disadvantages. Dr. ELLIOTSON speaks well of a mixture of sulphuric acid and water—a drachm to the pint—as a wash. The application of belladonna is useful for local sweating, but when the sweating is general, the internal administration of atropia is to be preferred.

Duboisia was reported upon favorably by the late Prof. GUBLER. In the Philadelphia Hospital, *jaborandi* has been found very effective:

428. R.	Extracti jaborandi fluid.,	f. ʒss	
	Tinct. cardamomi,		
	Syrup. pruni virginian.,	āā	f. ʒij. M.

One teaspoonful in half a wineglassful of water the first night, then half a teaspoonful every following night until cessation of sweats.

It will rarely be necessary to give the medicine oftener than four

consecutive nights. if, after some weeks, the sweats should return, which is frequently the case, one or two of the smaller doses will be sufficient to stop them again.

Picrotoxin, gr. $\frac{1}{15}$ – $\frac{1}{10}$, in solution, has been used with success in many cases. (*Practitioner*, Oct., 1878.)

Pilocarpine, the alkaloid of jaborandi, has been given in the form of pill made up with sugar of milk, and in one-twentieth grain doses three or four times a day; equivalent to about eight drops of the fluid extract of jaborandi.

NOTES ON REMEDIES.

Acidum Carbolicum. The inhalation of pulverized solutions of carbolic acid is of frequent benefit in the advanced stages of phthisis, by diminishing the secretion and lessening the cough and exhaustion, and by correcting fœtor of the expectorations. A solution of drops xv–xx, in two pints of water, with the addition of one or two ounces of tincture of conium, or of alcohol, may be used. Lately, Dr. J. SCHNITZLER, of Vienna, has claimed “remarkable success” in the treatment of phthisis by hypodermic injections of carbolic acid, using a one or two per cent. solution twice daily. (*Dobell's Reports*, 1877.)

Acidum Tannicum. Prof. STILLÉ says that this medicine appears sometimes to remove the crepitating rhonchi so generally observed around crude tubercles at the apex of the lung during the first stage of phthisis, while it diminishes the dyspnœa, cough and expectoration. At a late period, it sometimes so manifestly palliates all the symptoms as to inspire a belief that a cure has been accomplished, converting moist rhonchi, or gurgling, into signs of solidification merely, or of a dry cavity. The more nearly the disease resembles the acute form of phthisis, the less appropriate will the medicine become.

Æther has been recommended to promote the assimilation of cod-liver oil by Dr. FOSTER. The objections to his preparation are its disagreeable taste, and the eructations it occasions.

* *Alcohol*. Dr. ANSTIE says that there is a variety of pulmonary phthisis, especially common in persons with delicate skin and slight frame, with a marked tendency to colliquative sweating, and a notable inability to assimilate either ordinary food or fatty matter of any kind, in which the tolerance for large and long-continued doses of alcohol is very remarkable, and the benefit produced by such treatment very great. Prof. FLINT also recommends strongly the alcoholic treatment in cases in which stimulants are well borne.

Ammonii Iodidum is recommended in incipient phthisis by Dr. RICHARDSON. It is also advised by Dr. TANNER, in cases in which cod-liver oil is not tolerated.

* *Arsenicum* is often a remedy of great value, but it is frequently badly borne by

the stomach. Prof. DA COSTA recommends it as often beneficial in slow consumption. It is highly advocated by a number of French physicians.

Arseniosum Acidum. Dr. A. JACOBI, of New York (*Med. News*, 1884), says that in phthisis arsenic is the best alterative and nutritive which we can employ. He does not use it while hectic fever is high, but when the temperature falls arsenic in small doses is a very valuable agent in doses of from one-fifteenth to one-sixth of a grain, with a little opium. *Digitalis* is also valuable.

Calcii Chloridum is valuable in strumous phthisis. Dr. A. H. MELLERSH says many cases improve upon it. (*Medical and Surgical Reporter*, Feb., 1878.)

Caoutchouc, dissolved in turpentine and made into a confection, has been given in many cases by Dr. VARICK, of Jersey City, and has been found very useful where cod-liver oil cannot be tolerated. (*Med. Record*, Nov., 1873.)

Cinchona. The acidulated cold infusion of bark is an excellent remedy for the profuse perspiration, night sweat and hectic fever.

Creasotum has lately been recommended by Dr. HUGUES, of Paris, in a thesis. He tried it with good effect in twenty-seven cases, in the following formula :

429. R.	Creasote,	3j	
	Alcohol,		
	Water,	āā	f. ʒiv. M.

Of this, a dessertspoonful twice daily. The creasote must be pure and of wood-tar, not the impure stuff usually found in the shops.

Digitalis enjoyed formerly a higher reputation than it now possesses, although it is still highly spoken of. Dr. NIEMEYER considers it, in combination with quinine, as an admirable agent for arresting the abnormal calorification. It is often very useful in hæmoptysis, especially in the advanced stages of the disease.

Ferrum. Almost any of the preparations of iron are useful in the early stages, the choice of which should depend on the patient's idiosyncrasies.

Glycerinæ. In the non-febrile stage of phthisis Dr. TISNE, of Paris, uses the following as a stimulant to the digestive organs :

430. R.	Glycerinæ,	ʒiv	
	Spiritûs vini,	ʒss	
	Essentiæ menthi.,	gtt. j.	M.

In three doses each day. It is well to begin with half this quantity daily, though the above amount can soon be tolerated. It should be suspended if excitability of the heart is present.

Hydrogenii Peroxidum is said to greatly improve digestion in the first stage of phthisis, and aid the action of iron ; in the last stage, it unquestionably affords relief to the breathlessness and oppression, acting like an opiate without producing narcotism.

Iodinium is a valuable counter-irritant. Dr. FULLER recommends :

431. R.	Hydrargyri chloridi mitis,	gr. viij	
	Iodinii,	3 ss	
	Alcoholis,	f. 3 iss	
	Unguenti,	3j.	M.

Rub in a portion over the affected lung, morning and evening, until a pustular eruption comes out.

Iodoformum. Prof. CHIARMELLI, of Milan, states that even at an advanced period of the disease, with the presence of cavities, iodoform gives excellent results. "Iodoform," he says, "diminishes the fever and affects the expectoration, which it not only diminishes in quantity, but alters in character, preventing the putrefaction of its albuminoid elements. I am also convinced that the contents of the cavities in the lung exercise a powerful influence upon the production of hectic fever." He claims that it arrests the march of this malady and prolongs the life of the sufferer. He also holds that in cases where caseous pneumonia is commencing, iodoform administered for a time proves efficacious in arresting the progress of the disease. With many individuals affected with chronic bronchitis and emphysema, it renders great service. The formula which is employed is as follows :

432. R.	Iodoform,	gr. iss
	Powdered lycopodium,	gr. j
	Ext. of gentian,	q. s.

Make into ten pilules. Take three to five in the day.

If the dose is increased gastric disorders supervene, and it is better to continue the above dose for a considerable time.

Juglans. Infusion of the leaves of the *Juglanda regia*, or an extract, is much used in Germany when cod-liver oil cannot be tolerated, or in the hot season when the oil is apt to disagree.

Koumiss, the fermented milk of mares or cows, is a popular remedy, and is now prepared by druggists in our leading cities. A "cure" requires 12 to 15 lbs. of koumiss daily, taken every half hour, conjoined with plenty of open-air exercise. The Russian method of preparing koumiss is as follows : Take two teacupfuls of wheat-flour dough, two spoonfuls of millet flour, one spoonful of honey, one of beer yeast ; mix with milk to a thin paste, and put in a warm place to ferment. When fermented, put in a linen bag and hang it in a jar with 16 lbs. of fresh milk ; cover, and let stand until the milk is acidulous ; skim, decant and agitate for an hour, then bottle and place in a cold place.

The following is an American formula for koumiss :

433. R.	Fresh milk,	5 gallons
	Spring water,	13 pints
	Baker's yeast,	2 "
	Simple syrup,	1½ "

Mix, and skim off the top. Then put the liquid into bottles, which are to be laid on their side, from one to four weeks, in a cool place,

the temperature of which must be below 70°. The different stages of the preparation are as follows : After one week the koumiss is *fresh* ; after two weeks, *medium* ; after three or four weeks, *old*. A champagne tap is very convenient for drawing the liquid.

A preparation called *ferruginous koumiss* is prepared by daily adding ordinary doses of lactate of iron to two or three bottles of common koumiss. The iron salt easily dissolves, without changing the taste of the koumiss. This preparation is very well borne, even by patients who cannot take iron administered in pills, powders, etc.

*****Morrhue Oleum*** is deserving of the highest confidence. The earlier it is employed in the disease the better. In regard to its mode of action and method of administration, see opinions of Dr. AITKEN, of Prof. DA COSTA, of Dr. FOSTER, of Prof. JACCOUD, of Dr. NIEMEYER, of Dr. WALSHE, and of Dr. C. J. B. WILLIAMS (above.)

Muriaticum Acidum, alone or in combination with other mineral or vegetable tonics, is an important auxiliary to treatment.

Nitrogen Gas. Inhalations of this gas have been recommended by Prof. STEINBRUCK, of Vienna. In the first and second stages, especially in young persons, they give great relief and sometimes cure ; but in the third stage they are injurious. Their immediate effect is to allay nervous irritability and lower the circulation. (*Dobell's Reports*, 1876.)

Oleum Tiglii, as a counter-irritant to the chest in the earlier stages, is undoubtedly beneficial.

Opium, by the mouth or enema, is often a valuable palliative in advanced stages of phthisis, to relieve the cough, lessen the expectoration, check diarrhœa, and afford comfort to the patient.

****Pancreatic Emulsion***, introduced by Dr. HORACE DOBELL, is an excellent remedy, either with or without cod-liver oil, and is well borne by persons unable to retain the oil. He says it is best to give the *Pancreatic Emulsion* from one to two hours after a full meal, such as breakfast or dinner, particularly avoiding a longer interval, and also avoiding warm drinks for two or three hours afterwards. When cod-liver oil agrees, give a tablespoonful of oil directly after breakfast, to supply olein to the blood through the portal system, and a tablespoonful of emulsion, in a cup of milk or water, two hours after dinner, to supply the blood with pancreatized solid fats through the lacteal system. If cod-liver oil cannot be taken, give the emulsion two hours after breakfast, and two hours after dinner.

Plumbi Acetas is very useful in diminishing expectoration, checking hemorrhage and diarrhœa, and, to a less extent, the profuse perspirations of phthisis ; but its administration is objectionable, because of the anorexia which often speedily results.

Potassii Chloras is recommended in the painful deglutition of advanced phthisis.

Potassii Iodidum is of value, given with iron and cod-liver oil, when there is a syphilitic taint.

Prunus Virginiana improves the appetite and palliates the cough.

Quiniæ Sulphas is a valuable tonic in phthisis, given in conjunction with iron and cod-liver oil.

Saccharum. Dr. SYMONDS recommends (*British Medical Journal*, June 13th, 1868,) sugar as an article of diet in phthisis, taken to the extent of a quarter of a pound or more daily. He gives it with eggs as a succedaneum for cod-liver oil, when the latter disagrees.

Sanguis. The fresh blood of animals, drunk while warm, is a nutritive drink, and has proved of value in many phthisical cases.

Sanguis Bovinus Exsiccatus. This nutritive preparation has been found of great use in combating the exhaustion of phthisis. Dr. F. W. PUTNAM states that it is the most satisfactory of any such substance he has used. (*Therap. Gazette*, 1883.)

Sevum Præparatum, Suet, is a popular and useful remedy, boiled in milk, and taken in the morning.

Silphium Cyrenaicum. In Italy, Dr. ANTONIO LOMBARDO assigns the following effects to the treatment of phthisis with *Silphium Cyrenaicum*: The expectoration is much facilitated; the dyspnœa is diminished, and the circulation becomes more regular; while no digestive disturbance results from its administration. (*Dobell's Reports*, 1877.)

Sodii Sulpho-Carbolas has been recommended by Dr. SANSOM. (*Practitioner*, July, 1869.)

Sulphur is useful in the costiveness of phthisis, and may also be employed against profuse perspirations.

Verbascum Thapsus, the common mullein, has been praised for its effect in phthisis by Dr. F. J. B. QUINLAN. In the *Brit. Med. Jour.*, December 8, 1883, this gentleman reports a case of pre-tubercular phthisis, in which the patient gained twelve pounds in weight in one month under the use of mullein. He considers that it possesses all the advantages and none of the drawbacks of cod-liver oil.

Vitis Vinifera. The grape cure is of benefit in chronic catarrhal cases, but is of doubtful propriety in tuberculosis cases.

Zinci Oxidum is strongly recommended by Dr. WARING CURRAN, as a tonic.

INHALATIONS.

Arsenicum. TROUSSEAU and others have recommended arsenious fumigation by means of cigarettes—a mode of treatment which should be conducted with caution.

Carbolicum Acidum. Dr. MARCET recommends (*Practitioner*, November, 1868,) the atomization of the following solution:

434. R.	Acidi carbolici,	gr. ss-iss	
	Aquæ,	f. ʒj.	M.

This inhaled in the form of spray acts beneficially in the first stages of the disease, by improving the circulation in the lungs, but in the advanced stages it should be withheld.

Chloroformum. A few whiffs (m.vj-x) are often useful in the cough and dyspnoea of phthisis, but the effect is only temporary.

* *Iodinium.* In cases of troublesome cough and abundant expectoration, benefit sometimes results from employing, night and morning, vapor of iodine.

435. R. Tincturæ iodinii, f. ʒj
Aquæ, f. ʒj. M.

Apply gentle heat, and let the vapor that arises be inhaled. Or, the dilute compound tincture of iodine may be atomized.

Iodoformum. The inhalation of the dilute vapor obtained by mixing iodoform with starch, spreading it on paper and allowing it to volatilize, has been recommended.

* *Opium*, by inhalation, is an excellent means of soothing irritative cough.

Oxygen inhalations can only be looked upon as palliative, and seem, in some cases, after affording immediate relief, to invite fresh and more intense exacerbations.

* *Pix Liquida.* Tar vapor is now little used, having lost the reputation it once enjoyed in phthisis.

EXTERNAL REMEDIES.

Aceticum Acidum Dilutum. The application of diluted vinegar to the chest and upper part of the body, to allay the profuse perspirations of phthisis, is recommended by Dr. WALSHE, Sir C. SCUDAMORE, and others, as a measure greatly conducing to the comfort and well-being of the patient. The mixture of one part of vinegar, one of eau de cologne, and two of water, is employed by Sir C. SCUDAMORE.

* *Ammonii Iodidum*, in ointment, is recommended by Dr. TANNER.

* *Baths.* The *Turkish Bath* is sometimes of service in incipient phthisis. The *Compressed Air Bath* may often be employed with great advantage.

* *Counter Irritation.* The counter-irritants most frequently resorted to in phthisis, are the following: *Aqua Ammonia*; *Tartar Emetic Ointment*; *Calomel and Iodine Ointment*; *Croton Oil Liniment*. * *Tincture of Iodine* (painted over the chest every night) also does good, by the iodine which is inhaled, from the air around the patient being slightly impregnated by the evaporation of the tincture; *Burgundy Pitch Plaster*. * *Thapsia Plaster*; the following plaster is recommended by Dr. BARLOW, in the first stage of phthisis:

436. R. Emplastri picis com., ʒ iss
Antimonii et potassii tartratis, gr. x. M.
Make a plaster.

Electricity. As far back as 1863, Dr. BASTINGS, of Brussels, announced the cure of twenty-five cases of phthisis out of thirty-seven patients, by means of electricity. (*Traite de la Phthisie*, 1863.) The *Archives Med. Belges*, 1879, contains the details of a case of well-marked phthisis, added to

his successes. Dr. B. electrizes the muscles of the chest and arms by means of an electro-galvanic machine of Gaiffe, employing such a strength that the resulting contractions correspond to those natural to respiration. One or two sittings of five minutes daily, are sufficient.

Setons. In the early stages, a sub-clavicular seton of three or four silk threads, sometimes retards the progress of the disease.

PLEURISY.

JOHN SYER BRISTOWE, M. D., LONDON.

In mild cases of so-called dry pleurisy, the chest should be irritated by a mustard plaster, and then swathed in a broad flannel roller, so as to restrain motion, while small doses of opiates should be commenced.

When there is manifest fever and commencing effusion, it is often beneficial to apply half a dozen to a dozen leeches to the surface of the chest, followed up by poultices, or flannels wrung out in hot water. Opiates are here, also, of extreme value, and some febrifuge medicine may be employed.

If the effusion still increases, diuretics and purgatives, mercury, digitalis, and iodide of potassium, have been employed. All these Dr. B. rejects as useless. He would prefer tonics, especially iron and quinine; counter-irritants, especially repeated small blisters; and, finally, paracentesis thoracis, or aspiration. (See "Empyema.")

ALFRED L. LOOMIS, M. D., NEW YORK.

In mild as well as severe cases of pleurisy, all the treatment required for the successful management of the disease is to put the patient quietly to bed (very important), apply an anodyne poultice to the affected side, and administer hypodermic injections of morphia in quantities sufficient to relieve all pain. Ordinarily, the patient can sit up in four or five days, and in two weeks resume his business. But if the exudation is abundant, he should keep his room and avoid exertion. If the patient is anæmic, the syrup of the iodide of iron should be given, f.ʒj, three or four times a day. Stimulants should never be given until the second or third week of the disease; and all so-called antiphlogistics and counter-irritants should be avoided.

Iron is the remedial agent which has the greatest power to promote

absorption. When the case is sub-acute, the most nutritious food and free stimulation are in place. The nutritive processes should be raised to the highest point. In any case, when the fluid accumulation remains stationary for one week, or is increasing after the cavity is half filled with fluid, *aspiration* should be resorted to.

PROF. J. M. DA COSTA, M. D., PHILADELPHIA.

- | | | | |
|---------|---------------------------|----------|----|
| 437. R. | Potassii acetatis, | gr. xv | |
| | Spiritus ætheris nitrosi, | f. 3 ss | |
| | Vini ipecacuanhæ, | gtt. iij | |
| | Syrupi tolutani, | f. 3 ss. | M. |

For one dose, four times a day. Useful in sub-acute pleurisy.

- | | | | |
|---------|-----------------------------|-----------|----|
| 438. R. | Tincturæ veratri viridis, | ℥xxiv | |
| | Potassii acetatis, | 3 ss | |
| | Morphiæ acetatis, | gr. ss | |
| | Liquoris potassii citratis, | f. 3 ijss | |
| | Syrupi tolutani, | f. 3 ss. | M. |

A dessertspoonful every three hours, in dry pleurisy.

Locally, apply, two or three times a day, turpentine stupes.

- | | | | |
|---------|---------------------------|----------|----|
| 439. R. | Potassii acetatis, | 3 j | |
| | Tincturæ digitalis, | f. 3 ij | |
| | Extracti cinchonæ fluidi. | f. 3 j | |
| | Aquæ, | f. 3 ij. | M. |

A teaspoonful thrice daily, in pleuritic effusions.

- | | | | |
|---------|----------------------------|----------|----|
| 440. R. | Potassii iodidi, | 3 ij | |
| | Tincturæ scillæ, | f. 3 vj | |
| | Tincturæ opii camphoratae, | f. 3 iss | |
| | Misturæ acaciæ, | f. 3 vj. | M. |

A teaspoonful four times a day, in chronic pleurisy with consolidation of the lung. Also, a good, nourishing diet; either whisky or gin, half an ounce three times a day; and counter-irritation by means of tincture of iodine.

TREATMENT BY COMPRESSION OF THE THORAX.

Professor CONCATO, of Bologna, has introduced a new treatment of unilateral pleurisy with effusion, which has been received with much approbation in Italy. It is by manual compression performed with both hands applied to the *healthy* side twice a day, during five to fifteen minutes each time. The patients experience rapid benefit from this treatment, the dyspnœa disappearing and rapid absorption taking place. According to Dr. SANTE ALBERTOZZI, of Florence, it takes the place of thoracentesis, and is, of course, safer. No remedies are required, and the treatment is assisted only by nourishing diet. To suc-

ceed well, fever should be absent and the effusion recent. (*Dobell's Reports*, 1876, 1877.) See further, Dr. GERHARDT's method under "Emphysema."

On the other hand, Dr. J. C. GLEASON, of Boston, recommends strapping *the affected* side with adhesive strips, so as to limit motion and secure the greatest possible rest in all cases of acute pleurisy, pleuro-pneumonia, etc., in which severe pain, resulting from the respiratory movements, comes in as a leading symptom. (*Boston Medical and Surgical Journal*, December, 1877.)

NOTES ON REMEDIES.

**Aconite*. Bartholow considers aconite and opium as very effective in acute pleurisy before the stage of effusion and highly recommends the following :

441. R.	Tinct. aconiti rad.	℥ ij	
	Tinct. opii. deod.	℥ vj.	M.

S.—Eight drops in water every hour or two.

If the pain is severe, a large dose of opium should be given, when the effect can be maintained by the quantity directed in the above prescription.

Antimonii et Potassii Tartras, in doses gr. $\frac{1}{8}$ — $\frac{1}{4}$, sometimes proves useful in the earliest stages of acute pleurisy in young, plethoric subjects. It is, according to LÆNNEC, and the older writers, in general, well supported in pleurisy, and contributes powerfully to subdue the inflammatory tendency; but, nevertheless, when the pain in the side and fever have ceased, it loses further power over the disease; at least, it does not appear to promote the removal of the fluid effused, so that its use must generally be abandoned as soon as the acute symptoms have passed away.

Digitalis is an antiphlogistic, adapted to the pre-exudative stage.

Chloral is recommended when there is restlessness or delirium.

Pilocarpine aids in the absorption of the exudation.

Iodinium. In effusion, after the febrile action, the *compound tincture of iodine*, in doses of *twenty minims freely diluted*, is a valuable medicine.

Jaborandi has proved very serviceable in several cases of pleuritic effusion, and in a late number of *La France Medical*, Dr. MICHOU reports two cases, the happy termination of which certainly appears to be due to the action of this drug. In both cases the patients were rather bad subjects, the symptoms severe, and other remedies unsuccessful. Five grammes (75 gr.) of jaborandi leaves were given in infusion, and in both cases this dose was soon followed by copious expectoration, salivation, and diaphoresis, together with marked absorption of the effusion, and speedy relief to all the symptoms.

**Morphia*. Nothing can be more satisfactory than the treatment of pleurisy in its early stage by the hypodermic injection of morphia. It relieves at once the pain, and arrests or diminishes the morbid process.

442. R. Morphiae sulphatis, gr. xvj
Aqua destillatae, f. ʒj. M.
Dissolve and filter. Dose, for a hypodermic injection, five to ten minims.

Potassii Iodidum, variously combined according to indications, is considered by many of the highest value in restraining inflammatory action and promoting absorption. (F. 440.)

Quinia Sulphas, in the advanced stages, when the case assumes a typhoid character, especially when the patient is old or debilitated, is of great service.

Scilla is sometimes given, combined with digitalis and mercury.

Tannicum Acidum. Dr. DUBOUE (in the *Gazette Hebdomadaire*, December, 1872,) relates the remarkably good effects which he has obtained from the use of tannic acid. The exclusive use of tannic acid in eleven grave cases of purulent pleurisy, attended with spontaneous evacuation of pus, was entirely successful in a very short time in eight cases; the ninth case was under treatment for a long time, with only partial success; two cases ended in death. He usually administered the tannin in pills:

443. R. Acidi tannici, gr. ij
Confectionis rosae, q. s.
For one pill. Four to eight daily, one-half in the morning, the remainder in the evening.

Dr. DUBOUE was guided, of course, by the condition of the stomach, which bore the drug remarkably well as long as there was a source of suppuration; as also by the effects obtained, and not only the return of appetite, diminution of expectoration, sweats, etc.

PNEUMONIA.

THE ABORTIVE TREATMENT OF PNEUMONIA.

The fact that acute pneumonia sometimes aborts spontaneously leads to the hope that this termination may be brought about by art. With this aim, various means have been used.

Blood-letting, free and early, in young and vigorous subjects, especially in healthy country localities, is an old and approved means. As, however, death in pneumonia often comes from exhaustion, instead of venesection we may apply a number of cups to the chest, and follow with hot fomentations and poultices.

Internally, *ergot* has been used with marked reported success, f. 5ss of the fluid extract every two hours, by Dr. J. B. SCEARCE (*Med. and Surg. Reporter*, March, 1878,) and Dr. J. B. YEAMAN (*St. Louis Clin. Rec.*, Feb., 1878).

Dr. SIDNEY RINGER and others speak very strongly in favor of *aconite*, gtt. i-ij of the tincture of the root every half hour, for two hours, and then less frequently. It must be begun early, or its effects are inconsiderable. It has no power over the consolidated portions.

That *quinine* aborts the disease in a certain number of cases, Dr. A. FLINT says there is no doubt. He gives to an adult gr. xx-xl, either in a single dose or in two or three doses during the first day. He thinks it should always be given, as it favorably modifies when it fails to arrest the disease. (*Clin. Med.*, 1879.)

Belladonna in the first stage is spoken well of by PHILLIPS. (*Mat. Med. and Ther.*, 1878.) Doses of ℥xv every hour or two may be given until the physiological effects on the eye and throat are produced.

Persistent *inhalation* of the steam from hot water, plain, or medicated with some of the above agents, is stated by COHEN to have had long repute as an abortant of pulmonary congestion in its earliest stages.

Prof. A. B. PALMER remarks in the *Trans. of the Am. Med. Association*, 1876, that for twenty years he had taught and verified in practice the fact that if a patient with the ordinary acute form of pneumonia be seen within a few hours after the attack, and a free dose of an *opiate* be given, and once or twice repeated, so that its full anodyne effect is produced and for some hours maintained, in a large majority of cases the progress of the disease will be arrested. After the effect of the narcotic has subsided, a mild cathartic and eliminative will complete the cure. Within a few years he had modified this treatment by combining the opiate with quinine as follows: When called to a patient within twenty-four hours of the chill he immediately gives,

444. R.	Morphiæ sulphatis,	gr. $\frac{3}{4}$ - $\frac{1}{2}$	
	Quiniæ sulphatis,	gr. vj-x.	M.
For one dose.			

In about an hour this induces free perspiration. The quinine alone is then repeated, gr. v-x, every two or three hours, until 5ss-j has been taken. No attention should be paid to the ringing in the ears, etc., as these symptoms are transitory and of no moment. The effect

to be produced is the checking of the fever, and this nearly invariably occurs. If the treatment is commenced later, after consolidation is present, the morphine must be omitted, but the quinine administered as above.

The use of *salicylic acid* as an abortant of the disease has been recommended by Dr. L. L. SILVERTHORN. (*Chicago Med. Jour. and Exam.*, Feb., 1878.) He administers gr. xx. every two hours until four or five doses are taken, when free diaphoresis should follow, with resolution.

GENERAL TREATMENT.

PROFESSOR BAMSLER, OF THE UNIVERSITY OF FRIEBURG,

directs his efforts chiefly towards sustaining the patient's strength until the disease leaves him. The pyrexia being a chief cause of exhaustion, the endeavor is to keep down the body-heat, which he does by cold baths, wet packing and quinia in 15 to 20 gr. doses, in the evening, or grains 60 to 80 of salicylate of soda within an hour in the middle of the night.

The patient's diet must receive careful attention. See to it that he is sufficiently nourished, as by broths, beef tea, milk, and a half to a pint of light wine, in twenty-four hours.

When there are *pleuritic pains*, an *ice-bag* is applied to the chest.

Restlessness, great pain or diarrhœa, is to be met by morphia or Dover's powder.

If bronchial catarrh is a prominent symptom, ipecacuanha in infusion is administered. He never employs sweet spirits of nitre in pneumonia.

M. RIEBE, OF PARIS,

has tested clinically the iodide of potassium in pneumonia. He used it exclusively in thirty-seven cases, in twelve of which there was double pneumonia, while three cases were complicated with pleurisy. He administers every second hour a tablespoonful of the following solution:

445. R. Potass. iod.,	$\frac{5}{3}$ iss	
Aquæ,	$\frac{5}{3}$ viii.	M.

A bladder containing ice was placed over the region of the thorax, corresponding to the affected portion of the lung. Of the thirty-seven cases, but one died, and this was a case of double pneumonia.

PROFESSOR SILVESTRINI, OF ITALY,

has been using *nitrite of amyl* in pneumonia, and he thus formulates his conclusions:

1. In the pre-organic stage of pneumonia, the nitrite of amyl is of prompt and effective service.

2. One may repeat with impunity the inhalation of this remedy during several successive days, and in doses relatively enormous. (He has administered as much as fifty grammes of the nitrite in five days, the inhalation being carried on for five minutes every half hour.)

3. In cases which have a fatal issue, whether from extension of the pneumonia or from complications, these inhalations may retard death.

PROF. NATHAN S. DAVIS, M. D., CHICAGO.

In the treatment of active sthenic pneumonia in its first stage, when the patient has been strong and healthy, a free bleeding from the arm (oz. 16-20) during the first day or two, followed by arterial sedatives, just enough to control circulation, united with just enough anodynes to lessen irritability and pain, constitutes the most efficient treatment.

When the disease is associated with a typhoid condition of the system, as is usually the cases in cities, bleeding is not well borne. In this class, Dr. DAVIS has succeeded best by giving, during the first two or three days, the two following prescriptions:

446. R.	Ammoniāe muriatis, Antim. et. pot. tartratis, Morphiāe sulphatis, Syrupi glycyrrhizæ,	℥ iij gr. ij gr. iij f. ℥ iv.	M.
A teaspoonful every four hours.			

447. R.	Hydrarg. chloridi mitis, Ipecac. pulveris, Opii pulveris, Sacchari albi,	āā gr. vj gr. iij gr. xxx.	M.
---------	---	----------------------------------	----

For six powders. One powder every four hours alternately with the preceding prescription. At the same time cover the chest with emollient poultices.

At the end of twenty-four hours the powders should be omitted, and if the bowels have not been moved, a mild laxative should be given; but the muriate of ammonia mixture may be continued every three or four hours for several days. If the symptoms are not favorably modified by the third or fourth day, a blister from three to six inches square should be drawn on the side of the chest most affected. If the pulse

becomes soft and frequent, the breathing abdominal, and the lips of a leaden hue, the following may be given with advantage between the doses of the other medicine :

448. R.	Quiniæ sulphatis,	gr. ij	
	Ammoniæ carbonatis,	gr. iv.	M.
For one dose.			

If delirium or morbid vigilance becomes troublesome, add chloroform, ℞-xij, to each dose of the ammonia mixture. This will generally procure sleep, or greatly improve the condition of the patient. Where a diurnal remission of the fever indicates the presence of malarious influence, fair anti-periodic doses of quinine should be given during the remission.

Dr. DAVIS reports recently (*Journal of the American Medical Association*, 1884,) two cases of pneumonia, in which certain alarming features of the disease were promptly relieved by the exhibition of ergot. He remarks, that there is no fixed or routine treatment which is adapted to all cases of pneumonia, simply because the morbid processes which constitute the different stages of pneumonic inflammation, are liable to be much modified by the previous condition of the patient, and the coincident meteorological and sanitary influences that surround him. The benefit to be derived from ergot is when there is a demand for more steadiness and tone to the action of the heart. Given in connection with digitalis under those circumstances, it is, perhaps, the most reliable agent we possess for resisting the over-engorgement of the pulmonary vessels, during the first stage of the disease, in which the condition of the heart referred to is found to exist.

ALFRED L. LOOMIS, M. D., NEW YORK.

This writer discards blood-letting, antimony and calomel, as well as veratrum viride and aconite, as equally useless or dangerous in nearly all cases of acute lobar pneumonia. Rest in bed, in a well-ventilated atmosphere of about 70° Fah., and abundance of liquid nutritious food, are the first steps to be taken. The pain, if severe, should be allayed by morphia subcutaneously; the cough by small doses of opium or chloral. This is often enough.

In severe types, with high temperature, the German system of cold compresses or baths is of temporary avail and sometimes dangerous. The best antipyretic is *sulphate of quinine*, in doses of gr. xx-xxx per

day. This lowers the temperature, shortens the febrile stage, and hastens resolution.

Nearly all the deaths from pneumonia result from *failure of heart power*. The most serviceable remedy against this is *alcohol*. But it must not be used indiscriminately; as a rule, a feeble pulse, running 120 or 130 per minute, demands stimulants. If they act favorably, a change for the better will be noted in a few hours. The quantity must be graded to the needs; f. 5 vi-viiij of brandy in twenty-four hours is about an average.

Narcotics must be used cautiously and in small doses. *Counter-irritants* are useless and distressing. *Carbonate of ammonia*, so much used by some, is inferior to *champagne* as a diffusible stimulant. When œdema is detected in the unaffected portion of the lung, *dry cups* give marked relief.

DR. E. H. SHOLL, OF ALABAMA.

The cold water pack. This procedure has been used and is advocated by the above writer (*Med. and Surg. Rep.*, Dec., 1883). The following description of its application in the case of a young girl will show the method and its results when properly indicated:

“A lounge was prepared on which a heavy double blanket, well soaked with the coldest water to be had, was laid. In this, the clothing being removed, she was carefully and thoroughly rolled and packed by careful nurses, and over this another pair of wet blankets was placed. Immediately the shock gave rise to violent paroxysms of coughing, with abundant expectoration of rusty-colored sputa. In less than thirty minutes the temperature had fallen a degree, the restless, tossing girl had become quiet, and in one hour she was sleeping a gentle and undisturbed sleep. The mistress of the house, an intelligent lady, made repeated observations of the temperature, had the cold water poured on as freely as was necessary, watching the pulse, one hand being left out, and at the end of seven hours, according to my directions, she was taken out of the pack, carefully dried, put back in bed to rest, and from this time on she continued to steadily improve. The temperature never came back to its old height; convalescence was as rapid as could be expected, thus happily terminating a case about as far removed from the possibilities of recovery as is ever seen.”

DR. A. T. H. WATERS, LIVERPOOL.

The treatment employed by Dr. W., with a loss of only one case in forty-four, consisted mainly (apart from the occasional use of antimony, as above,) of *carbonate of ammonia*, combined with *chloroform* (449), *chloric ether*, *opium*, a *generous, nutritious diet*, beef tea, milk, etc., with a judicious use of *alcoholic stimulants* from an early period. *Turpentine stupes*, *sinapisms*, and *linseed meal poultices*, were employed in the early stages of the disease, and *blisters* in the more advanced.

449. R.	Ammonii carbonatis,	gr. iv	
	Spiritus chloroformi,	℥xx	
	Aquæ camphoræ,	f. ʒx.	M.

To be given every three or four hours ; may be given either with or without alcoholic stimulants.

Dr. W. considers that too high a value has been placed upon *mercury* as a remedy in the stage of hepatization, for which it has been considered peculiarly applicable. He finds that in these cases it possesses no special properties for promoting absorption of the effused matters. As a purgative, it is very useful, as it tends to relieve the portal system, often overloaded in pneumonic inflammation ; but if given in frequent doses, or with a view of producing salivation, its effects, save in a few exceptional cases, will be more or less prejudicial. It was only administered in one of the above-mentioned forty-four cases, and then it in no wise hastened, in the opinion of Dr. W., the recovery.

Opium is very desirable in many cases, often relieving pain, allaying the distressing cough, and procuring sleep. The pain in the side which so frequently accompanies pneumonia, and for which depletion was formerly prescribed, may generally be relieved by a dose of opium. It is believed of great value in arresting the delirium so apt to supervene in this disease. On the first appearance of the indications of approaching delirium, *e. g.*, want of sleep, restlessness, slight staring of the eyes, and slight tremor of the hands, a few doses of opium, and the exhibition of stimulants and nourishment, may at once arrest the attack. At the same time, all depletory measures should be avoided ; the aim should be to improve the nutrition of the brain and give vigor to the circulation, for when these results are accomplished the delirium will disappear.

Few cases are now met with which are benefited by general *bleeding*, and the abstraction of blood by cupping or leeching is not often necessary.

The pulse affords an excellent indication as to the need for *stimulants*. As a rule, it may be said that when the pulse is below 100 the case is not a grave one, and will yield to treatment of a simple character; but when it rises to 110, 120, and upward, the case is more serious. *The more frequent the pulse, the greater is the need for stimulants*, and it is very remarkable to see the effect they produce on the pulse when it is abnormally frequent, in some cases of pneumonia. Intermission of the pulse is an invariable indication for the free use of stimulants. In the pneumonia of aged people, stimulants may be given without hesitation.

NOTES ON REMEDIES.

Aconitum is a potent remedy in the early stages, (F. 441.)

Ammonii Carbonas. See above. (F. 449.)

**Ammonium Iodide* is highly recommended to prevent caseation of the inflammatory products of catarrhal and fibrinous pneumonia. To lessen the effect of this remedy on the tissue changes, arsenic should be combined with it, and every means used to support the body nutrition.

Antimonii et Potassii Tartras is little given at present.

Belladonna is highly spoken of by Dr. HARLEY, who found that under its use the grave symptoms rapidly subsided, and convalescence was speedily established.

Bryonia. In mild cases, especially of pleuro-pneumonia, Dr. O. T. SCHULZ, (*Amer. Pract.*, Sept., 1879,) believes that tincture of bryonia, gtt. j every two hours is beneficial in the stage of hepatization.

Camphora is recommended by Dr. COPLAND, as one of the most valuable remedies in *æsthenic* pneumonia, in doses of gr. ij-vij, every four, five, or six hours, combined with calomel and opium, or with antimony and henbane, according to the character of the attack; the latter being preferred when the inflammation approaches the sthenic form, in which case the doses of the camphor may be lessened.

Chloroformum, by inhalation (eight or ten drops at a time, three or four times a day,) is said to relieve the stitch in the side and palliate the cough and oppression.

Vinchona. See Quiniæ Sulphas.

Cuprum. KISSEL regards the salts of copper as curative in pneumonia, and the preparation which he prefers is the tincture of the acetate (*Phar. Ger.*). The mortality under this treatment was only 4.3 per cent.

Digitalis has been much employed of late years. M. RONY-SAUCEROTTE re-

gards it (*Practitioner*, March, 1869,) as one of the best means of relieving the febrile and other grave symptoms; less active as an antipyretic than veratrum, but more easily managed and less offensive; acting more slowly than leeches, but producing more durable results.

**Ergot.* Ergot in powder has been used successfully in *croupous pneumonia* by Dr. WYCESK (*New Remedies*, No. 2). He would not advise it, however, in large infiltrations, in emphysema, in cases of rupture of cerebral vessels, or in exhausted and decrepit individuals, as it might in such cases prove hazardous.

Ether by subcutaneous injection is remarkably effective when there is an adynamic condition. Fifteen to twenty minims should be injected three or four times a day according to the degree of the adynamia.

Eucalyptus. When pneumonia passes into *pulmonary gangrene*, Dr. BUCQUOY, of l'Hôpital Cochin, Paris, states that no drugs have given him better results than eucalyptus globulus. He uses it in the form of an alcoholate—two grammes—(half a drachm) daily in a mixture of water, gum, orange-flower water, and syrup.

Helenine has been used with satisfaction in chronic broncho-pneumonia by Dr. VALENZUELA (*Bul. Gen. de Therapeut.*, 1883). He gives $\frac{1}{8}$ gr. ten times daily.

Kairin has been strongly recommended by ERLANGEN. It certainly reduces the temperature, but it is apt to be followed by prostration and vomiting. Hence it is condemned as useless or dangerous by KORACH (*Deutsche Med. Zeitung*, Jan., 1884).

**Muscarine.* BARTHOLOW considers that muscarine ought to be very useful in incipient congestion of the lungs, because it produces strong contraction of the pulmonary capillaries. It ought to be especially effective when combined with digitalis, for while digitalis will aid the curative action on the pulmonary vessels, it will at the same time antagonize the cardiac depression caused by muscarine. The dose is from $\frac{1}{8}$ to $\frac{1}{3}$ of a grain.

Nitrite of Amyl. Prof. SILVESTRINI, in *La France Medicale*, 1883, states that he has been using nitrite of amyl in pneumonia, and he thus formulates his conclusions: 1. In the pre-organic stage of pneumonia, the nitrite of amyl may be of prompt and effective service. 2. One may repeat with impunity the inhalation of this remedy during several successive days, and in doses relatively enormous. (He has administered as much as fifty grammes of the nitrite in five days, the inhalation being carried on for five minutes every half hour.) 3. In cases which have a fatal issue, whether from extension of the pneumonia, or from complications, these inhalations may retard death.

**Opium* is frequently a valuable remedy in arresting delirium, relieving pain and cough, and inducing sleep.

Pilocarpin. M. DUGROUX asserts that pilocarpin by hypodermic injection, in the dose of one-sixth to one-third of a grain, has proven successful in some

forms of polyuria. It brought about complete cure in two cases of azoturic polyuria, the one essential and the other symptomatic of interstitial nephritis; in this last case it also caused the disappearance of very marked amblyopia. In two cases of simple polyuria it caused a notable amelioration of the general symptoms. (*Thèse de Paris*, 1883.)

Plumbi Acetas is highly recommended in *asthenic* pneumonia, occurring in debilitated subjects. It may be combined with quinine (gr. j of each every two hours,) or, if the cough be very violent, opium may replace the quinine. Dr. STROHL (*Journal de Médecine*, February, 1882,) considers it to be the best of the internal remedies for pneumonia. It is preferable, he remarks, to tartar emetic, to digitalis, and to veratria, because its action is more certain, more prompt, and more free from inconveniences. Its action is incontestably superior in the pneumonia of old people. About five grains may be taken per diem in solution, in divided doses. He has never observed the slightest indication of saturnine poisoning in the course of this treatment. Far from producing constipation, it rather occasions diarrhoea. It can be administered at all ages. It does not interfere with any of the phenomena concomitant to the critical resolution, as expectoration, diaphoresis, etc. Under its action the pulse rapidly diminishes in frequency; the febrile symptoms and the temperature fall in the course of six days. The use of the lead may be intermitted as soon as the fever has abated and resolution has fairly set in.

Potassii Bicarbonas. Dr. JOHN PAPHAM, of Cork, recommends:

450.	R.	Potassii bicarbonatis,	℥j-vj	
		Misturæ acaciæ,	f. ℥iij.	M.

A dessertspoonful in water, four, six or eight times in the twenty-four hours.

The evidence of the good effects of this alkaline treatment appears on the second or third day (*British Medical Journal*, December 28th, 1869). It acts as a sedative by delaying the cough and abating the state of congestion on which it depends.

Potassæ Liquor, in doses of ℥x-xv, added to an ordinary expectorant mixture, is often beneficial in the advanced stages of pneumonia when the sputa are scanty, thick and viscid.

Quiniæ Sulphas is of the highest value in *asthenic* pneumonia, and in the advanced stages of pneumonic inflammation of the old and debilitated. Under the administration of gr. v every third hour, the pulse becomes slow and steady, and the respiration free. BARTHOLOW considers quinine an important remedy under two conditions: in large doses during congestion and in small tonic doses as a stimulant when depression comes on.

Sanguinaria is recommended in *typhoid* pneumonia, when respiration is very difficult, the extremities livid, and the pulse full, soft and compressible. The dose should be large in proportion to the violence of the disease, and repeated until it excites vomiting or relieves symptoms.

**Senega*, in decoction, is recommended by Dr. C. J. B. WILLIAMS, when the inflammatory symptoms have subsided, and any amount of debility, with weak pulse, cool skin, cough, and dyspnœa, remains. Cases which have resisted other remedies, often almost immediately improve by its use, the pulse becoming slower, the breathing freer, the tongue cleaner, and the strength greater.

Serpentaria is frequently prescribed as a stimulant expectorant combined with carbonate of ammonia, in pneumonia of low grade.

Turpentine. This is an important remedy in acute pneumonia. Dr. R. E. POWER writes in the *British Medical Journal*, 1876, that for many years he has adopted the following treatment with great success: First, a hot terebinthinate stupe is applied until the skin is well reddened; then a little plain oil of turpentine sprinkled over the affected part; finally, a blanket wrung out of boiling water, covered with a dry blanket. He has had patients delirious and gasping for breath, with sordes on the lips, fall asleep as the last blanket was applied, and awake out of danger. The internal remedies subsequently used were quinine and tincture of perchloride of iron. Diet: milk and water, beef-tea, lemonade *ad libitum*, occasionally wine. The application of the turpentine to little children must, of course, be modified to suit the age. In all cases, keep the swathe three or four days or more uninterruptedly. As a rule, the active treatment need not be pursued very long, the patient being generally out of danger in twenty-four or forty-eight hours. The sequelæ of pneumonia are much modified, frequently altogether escaped.

Valeriana. The ammoniated tincture, combined with camphor, is sometimes employed in the advanced stages of pneumonia, but is inferior to musk.

Veratria has been employed with great advantage. It may be given in pills or in solution (if the stomach be irritable, with an effervescing draught). Prof. VOGT, of Berne, commences with gr. $\frac{1}{50}$, every two or three hours, until it produces vomiting or diminution of the pulse.

**Veratrum Viride* is much employed in pneumonia. The plan recommended by Dr. NORWOOD, is to begin with gtt. viij of the tincture every third hour, and add one drop to each successive dose, until the pulse is sufficiently reduced or nausea supervenes; the subsequent doses are then to be regulated so as to sustain the depressed state of the circulation, with as little disturbance of the stomach as possible, any excess of nausea being controlled by a little morphia. With the reduction of the pulse the inflammatory symptoms decline, and convalescence ensues. Many physicians have reported very favorably upon this remedy; but for all that, its propriety is doubtful.

EXTERNAL REMEDIES.

Baths. On the theory that the danger in pneumonia arises from enfeeblement of the heart, Dr. T. JURGENSEN, of Kiel, has advocated the withdrawal of heat from the body by cold baths, as often as the temperature reaches 104°. The duration of the bath ought to be from seven to twenty-five minutes, and with the old and weak, tepid water may be

used. Before the use of the bath, a stimulant ought always to be administered, on account of the increased effort which the bath entails on the heart for the time being. In addition to the lowering of the temperature by the bath, JURGENSEN recommends the administration of quinia in doses amounting to thirty grains, dissolved in water by the aid of acid. This quantity to be administered every second evening, between six and eight o'clock. There is no danger, he says, in even doubling this dose. He gives a nutritious diet, with a moderate amount of wine. The wet pack is also highly recommended. As a rule, a hot wet pack gives more relief than a cold one, but the feelings of the patient are a proper guide. When the organs within the chest are inflamed, it is good practice to wrap the whole chest tightly in a pack to limit the motion of the chest walls. The method of proceeding is as follows: Wring out in cold or hot water a large towel, fold it and place over the affected side or part; have in readiness a bandage or large towel sufficient to encircle the chest, and confine the wet pack by pinning as tightly as possible around the chest the bandage or towel. Spongipiline is an excellent material for making these hot or cold applications.

Blisters are not beneficial in the early, but may be useful in the advanced stages of pneumonia.

Lukewarm Baths. Dr. BOZZOLO has tried warm instead of cold baths (*Lond. Med. Record*, 1884). He believes that they exert a decidedly beneficial effect. They are of one to three hours duration.

Counter-irritants may be employed in the early stages of the disease. The principal applications employed are *tartar-emetic ointment*; *croton-oil*; **tincture of iodine*; **turpentine stupes*.

Poultices are especially useful in the early stages of the inflammation. Flaxseed poultices, made so as to envelop the whole chest, give great relief, and exercise a salutary influence on the inflammatory action.

Venesection. In regard to bleeding, in pneumonia, we may quote the opinions of three good authorities. Dr. SAMUEL WILKS, F. R. S., says: "In great engorgement of the lungs or of the right side of the heart, bleeding will afford relief," and should be practised. (*Guy's Hospital Reports*, 1874.) Dr. A. L. LOOMIS also says: "There is one condition in which a patient with pneumonia may be bled to advantage, and that is when there is evidence that the heart is engorged with blood, accompanied by the evidences of sudden pulmonary congestion and œdema." In his *Clinical Lectures* (p. 280), Dr. N. S. DAVIS, of Chicago, says: "Bleeding and direct sedatives must be limited to the first three days of the disease. That a prompt and free bleeding in the first or congestive stage of active pneumonia, followed judiciously by sedatives and anodynes, is capable of greatly lessening the tendency to exudation, and favorably modifying the whole subsequent tendency of the disease, even to the extent of materially shortening its duration, I have no doubt. I have seen it demonstrated so frequently, that to doubt is to distrust the evidence of my senses." These remarks of Dr. DAVIS are not intended by him to apply to pneumonia as seen in cities or malarious districts, where it nearly always partakes of an asthenic character.

III. DISEASES OF THE CIRCULATORY SYSTEM.

General Therapeutics of Heart Disease—Angina Pectoris—Cardiac Dilatation—Cardiac Hypertrophy—Carditis—Fatty Degeneration of the Heart—Functional Palpitation and other Derangements—Irritable Heart—Valvular Disease of the Heart.

GENERAL THERAPEUTICS OF HEART DISEASE.

PROF. GERMAIN SÉE, OF PARIS.

In the last edition of his *Clinical Lectures*, (Paris, 1883,) this distinguished teacher sums up the principles he adopts in treating cardiac disease as follows :

In cardiac dyspnœa he prescribes belladonna, datura, jusquiamine, which paralyze the nerves of the heart; anæsthetics and oxygen in inhalations, which cause congestion of the lungs; venesection, which weakens the heart. The application of sinapisms, cups, and blisters, is useful when not harmful.

He advises *iodide of potash*. With the iodide he also sometimes gives digitalis, convallaria, or chloral hydrate. For the paroxysms, he gives inhalations of the *iodide of ethyl*, injections of morphine, and extract of convallaria daily, dry cups, sometimes inhalations of chloroform.

In uræmic dyspnœa, the preceding treatment cannot be counted on.

In the dyspnœa of hæmoptysis, he advises ʒj of the aqueous extract of *ergot* in twenty-four hours, with gr.j of the extract of opium; simultaneously, turpentine in capsules, and perhaps digitalis, which by strengthening the heart prevents stasis of the blood.

In cardiac dropsies he give diuretics (milk 3–4 quarts, or convallaria) and non-emetic purgatives.

Opium and belladonna are proscribed, as they check intestinal and renal secretion.

Diaphoretics (as *jaborandi*) can be given with milk and the drastics. When the cardio-nephritic dropsy accompanies the uræmic dyspnœa,

he thinks the subcutaneous use of morphine dangerous. In such cases, iodide of potash should be added to milk diet and digitalis; and, if the dropsy persists, recourse should be had to scarifications.

Cardiac irregularity he treats by digitalis, convallaria, bromides, and chloral. In intermission he prohibits tea, coffee, alcohol, tobacco; and prefers convallaria to digitalis.

In palpitation from hypertrophy, he prefers iodide of potash; in anæmic palpitations from spermatorrhœa, he rejects iron, douches, bromides, and anti-spasmodics, and prescribes iodide of potash twice a day, with meals; a pill of aqueous extract of ergot gr. j, alcoholic extract of digitalis, gr. $\frac{1}{8}$, three times a day; a farinaceous diet, with fat meat and fish; astringent red wines; no mineral waters, beer, milk, coffee, nor tea; tepid baths every morning; intellectual work and moderate physical exercise; absolute continence in sexual matters.

In chlorosis, iron is generally successful, if aided by hydrotherapy and a hearty diet.

Hysterical palpitations he treats with digitalis, bromides, veratrine, and hydrotherapy.

General contra-indications to the treatment of diseases of the heart are hydrotherapy, mineral waters, ordinary and mineral baths, venesection, cauterization, and blistering, iron and tonics (quinine).

General hygienic prescriptions are: A climate of medium and uniform temperature, rather moist, and with little wind. Mountain air suits some, sea air others, but more choose the temperate climate of valleys and plains, protected against winds. No severe or prolonged exercise (hunting, gymnastics, fencing), no climbing, no excessive study. The diet should be "anti-obésique" and nitrogenous,—soup, fresh vegetables, lean meat, cheese, fruit, milk, *eau rouge*, beer; coffee, if no palpitations are present; no heavy wines, liquor, white wine, tea, aerated and mineral waters.

DR. S. O. HABERSHON, LONDON.

This physician lays down seven rules of treatment applicable in all cases of heart disease. (*Guy's Hospital Reports*, 1887.)

The *first* is to *lessen the work of the heart*; this may be done by mechanical rest, by the recumbent position, and by avoiding stimulants and sudden changes of temperature.

The *second* is to *insure regularity of action*, by avoiding mental ex-

citement, by guarding against indigestion, and by never allowing constipation to continue.

The *third* is to *lessen distension*, especially of the right side of the heart, by purgatives, diuretics, and by mechanically diminishing the quantity of fluid in circulation.

The *fourth* is the prevention of syncope. With this view, sudden muscular movements must be avoided; stimulants may be required, as ammonia, brandy, etc.; and sedatives must be withheld or cautiously administered.

The *fifth* is to strengthen the muscular fibres of the heart by suitable nourishment; a bracing air, if other conditions allow; chalybeate medicines; and if the patient be exhausted by want of sleep, this symptom must, if possible, be relieved.

The *sixth* is to prevent fibrination of the blood. For this purpose, carbonate of ammonia will often be useful; other alkalies, as potash, soda, and other salts, may be beneficial; but, if long continued, in considerable doses, they depress the action of the heart. The acetate and iodide of potash may be advantageously combined with the carbonate of ammonia, or perhaps the hydrochlorate of ammonia.

The *seventh* is to prevent secondary complications, and to relieve them when produced. These complications are: 1. Broncho-pneumonia and pleuritic effusion. 2. Pulmonary apoplexy and other hemorrhages. 3. Visceral engorgement, as hepatic and renal congestions, with ascites and anasarca. By freely acting on the bowels, the portal congestion is greatly diminished, and the liver is enabled to act in a normal manner. Thus a free mercurial purge is of great value. The kidneys may be excited to a more vigorous action by a combination of mercurial medicine, with squill and with digitalis, when the latter can be borne. Salivation should be avoided. Diuretics are useful. An effectual way of diminishing the anasarca is by puncturing the skin on the thighs. The pulmonary engorgement is sometimes greatly reduced by applying cupping-glasses between the shoulders, or by the application of a blister to the chest.

J. MILNER FOTHERGILL, LONDON.

Treatment of Primary Heart Disease. It is of the utmost importance, in primary disease of the heart, to reduce to a minimum the calls upon its action. Rest in bed is desirable, and light labor alone must

be attempted. The tendency usually is for the veins to become too full of blood, while the arteries remain insufficiently filled. Much relief is afforded by unloading this venous congestion. Brisk catharsis, say pulv. jalapæ comp. ℥ij, at frequent intervals, does much good. Diaphoresis is also valuable. The form of diaphoretic best adapted to heart cases, is the application of heat externally. The bath, however, should not include the head, hence Turkish baths are objectionable. In the attacks of *dyspnœa*, so common in diseases of the heart, large, hot poultices of linseed meal, faced with mustard, and applied over the front of the chest and between the shoulders, are most serviceable. Of all drugs in cardiac adynamy, *digitalis* takes the lead. Its cumulative action has been greatly exaggerated, if it exists at all. The continuous use of small doses is the best plan of treating chronic cases. In cardiac dropsy, full doses must be given. Its effects are most gratifying when it produces a full flow of urine. In simple cardiac debility the following is a pleasant combination:

- | | | | |
|---------------------------|---------------------------|--------|----|
| 451. R. | Tinct. digitalis, | ℥x | |
| | Spiritus ætheris nitrici, | f. ʒss | |
| | Infusi buchu, | f. ʒj. | M. |
| This amount thrice daily. | | | |

At other times potash in any of its forms, but especially the citrate, may be added to the mixture. Where atonic gout is combined with the disease, the following is constantly used by Dr. F.:

- | | | | |
|---|-------------------|--------|----|
| 452. R. | Potassii bicarb., | gr. x | |
| | Fer. pot. tart., | gr. v | |
| | Tinct. digitalis, | ℥x | |
| | Infusi buchu, | f. ʒj. | M. |
| This amount thrice daily, to be followed by a good draught of water, and taken half an hour before meals. | | | |

Where there is palpitation, acute anæmia and debility, the following may be given:

- | | | | |
|--|-------------------------|--------|----|
| 453. R. | Tinct. digitalis, | ℥x | |
| | Ferri ammonio-citratis, | gr. v | |
| | Ammoniae carbonatis, | gr. v. | M. |
| This amount three or four times daily. | | | |

In the complex condition of cardiac debility, gastric catarrh, copious eructations of wind, and inactivity of the bowels, so commonly found together in chronic heart disease, *digitalis* may best be given in pill, as in this combination:

454. R. Pulv. digitalis, gr. xxx
 Ferri sulph. exsic., gr. xv
 Pulv. capsici, gr. xl
 Pulv. aloes et myrrhæ, ʒ ij. M.
 Make sixty pills. One twice daily.

In advanced dropsy, poulticing the abdomen and thighs with digitalis leaves has been found to produce excellent results.

TREATMENT OF SECONDARY AFFECTIONS OF THE HEART.

High arterial tension is usually the result of continued hypertrophy, lithiasis (suppressed gout), and spasm of the arterioles. In lithiasis the various natural waters which contain potash are very valuable. When drugs are prescribed, it is a matter of first importance that each dose be accompanied by large draughts of water; this makes them much more effective. A good form of mixture is furnished by the following combination:

455. R. Potassii iodidi, gr. v
 Potassii bicarbonatis, gr. xv
 Infusi buchu, f. ʒj. M.
 This amount, three or four times a day on an empty stomach, washed down with a tumblerful of water.

If the pulse be strong and hard, and the first sound of the heart good, then vini colchici, gtt. x, may be added to the above with advantage. If the opposite condition of the heart exist, with a compressible pulse and dilatation, then instead of the colchicum, tinct. digitalis, ℥x, may be added. This will maintain the vigor of the heart under the depressing action of the alkalies.

DR. THOMAS KING CHAMBERS, LONDON.

Dietetics in Heart-Disease. Dr. CHAMBERS points out (*A Manual of Diet in Health and Disease*) the fact that in disease of the heart the most remarkable change in respect to digestion is the slowness with which liquids are absorbed by the stomach, owing to the weakening of the circulation. The fluid in excess remains in the viscus and impedes digestion. The observation of a *dry diet* contributes greatly to the comfort of the patient, and more particularly so where dilatation of the heart's walls exists, and where, therefore, distension of the stomach would interfere with the laborious action of the heart; also, in cases of cardiac disease complicated with obesity.

The dietary should be at least as nitrogenous as in health. There is danger of atrophic degeneration of the cardiac muscle, which must be guarded against by generous diet. Alcohol should be used either not at all, or only in sufficient quantity to facilitate the assimilation of food; as soon as it diminishes assimilation, it is injurious. Indeed, we may say that alcohol is rather indicated during the times of acute failure than as a permanent part of the dietary of heart affections. The use of tea and tobacco, both of which are neurotic poisons when taken in quantities, must be rigidly restricted.

Climate in Heart Disease. Dr. CHAMBERS points out that in England degenerative heart disease is the rule and acute the exception; while in Italy, degenerative disease is the exception and acute disease the rule. Aneurism is almost unknown in Italy. The severe winters of the north of Europe and the northern United States are particularly trying on patients with heart disease. A low temperature, by contracting the capillaries, raises the blood pressure, and makes additional demands on the weakened heart. The diminished pressure of the atmosphere in elevated regions, as of our western plains, leads to excitability and hypertrophy of the heart. As a rule, cardiac diseases are less common and less fatal near the sea, where the extremes of temperature are less marked, and the atmospheric pressure the greatest. The considerations offer suggestions for the proper climatic treatment of cardiac affections.

DR. T. CLIFFORD ALLBUT, M. A., F. L. S., ETC., LONDON.

To subdue the distress in nervousness, the dyspnœa, the pain and night terrors of advanced heart disease, this writer urges (in the *Practitioner*, December, 1869,) the value of the *Prunus Virginiana*, and especially the *hypodermic use of morphia*. He justly says there is scarcely any disease more distressing than the latter stages of these maladies, when the patient, with a clear head, but with gasping breath, a striving heart, and swollen limbs, hardly finds a moment's rest from torment. Opium cannot be given, on account of its effect on the kidneys. His experience convinces him of the utility of the wild cherry in many of the milder cases of cardiac distress, and in such, he urges its claim warmly; but it is impotent in the more severe forms. Here he employs *morphia*, hypodermically, especially in angina with diseased coronary arteries, in neuralgic distress from intra-thoracic tumors, in

mitral regurgitation, and in aortic disease, where the heart is big and pumping.

He uses the hydrochlorate of morphia in doses varying from one-tenth to one-third of a grain; seldom half a grain, except in such a case as intense pain from angina or intra-thoracic tumor. He always begins with one-eighth or one-sixth; and in ordinary cases, he finds a quarter of a grain the proper dose—in a case, say, of mitral regurgitation with pulmonary congestion in an otherwise healthy adult. The dose is best given in the evening, and should always be followed by perfect quiet in the room. The last is an important element in all cases of injection of morphia. The urine should be examined and the drug withheld, or given with caution, if albumen be found. There need be no great fear of it during albuminuria only secondary to the heart disease, unless there be reason to suppose that excessive renal congestion be present, and head symptoms at hand. He scarcely ever remembers that morphia caused disturbance of the stomach or other inconvenience. As in cases of severe pain, so it seems that in cardiac disturbance, the intolerable distress “carries off” the troublesome qualities of the drug. Its effects, again, are not merely comforting, but are curative so far as such cases are to be cured. An injection of morphia three or four times a week, by tranquilizing the heart and allowing the circulation to recover its freedom, sets free, also, the organs which are oppressed, and the system can once more find something like its balance. Thus relieved, the patient may, with incessant care, enjoy some respite from his malady, and the occasional use of morphia may put off the day of its return.

HORACE DOBELL, M. D., LONDON.

DR. DOBELL lays down certain “aphorisms” in reference to *pain in the region of the heart*, which should materially influence the treatment adopted. (*On Affections of the Heart*, London, 1876.) Thus, such pain does not necessarily indicate heart disease; a neglected dyspepsia, especially that characterized by an excess of acidity, may produce it; also, an erratic neuralgia of the gastric branches of the pneumogastric; deteriorated blood in anæmia; and local interference with the circulation. But if the pain at the heart is accompanied by pain in the left arm, it is an important symptom of cardiac disease; also, if it is excited by exercise on an empty stomach. Of course, in various dangerous forms of heart disease, pain is entirely absent.

In treatment, the patient should be instructed to take the warning given by nature, and stop exercise or exertion when called upon to do so by pain at or about the heart; yet a rigorous and persistent restriction of exercise, necessary in the early part of the treatment, must not be carried to the extent of weakening the muscular action of the heart. For dyspeptic pain, alkaline mixtures and vegetable tonics are demanded. A favorite one with Dr. D. is:

456. R.	Sodii bicarbonatis,	gr. v-x	
	Pulv. calumbæ,	gr. x-xx	
	Pulv. zingiberis,	gr. x.	M.

Take in water before each meal.

DR. LAYCOCK, DUBLIN.

457. R.	Potassii iodidi,	gr. v	
	Potassii bromidi,	gr. xv.	M.

For one dose; to be taken at night.

In the distressing *dyspnœa* of heart disease, Dr. L. has found the above combination superior to any other, believing it to be both palliative and curative. Of opiates, he prefers the hypodermic use of morphia, or else ten to fifteen minims of the solution of the bimeconate of morphia by the mouth; although he is satisfied that in any form opium tends to increase the albuminuria and dropsy which accompany cardiac lesions. Should the gastric vagus be implicated, hydrocyanic acid is the most efficacious remedy. (*Dublin Medical Journal*, December, 1873.)

DR. C. M. DURRANT, LONDON.

Hygienic Rules in Incurable Heart Disease. The first rule to impress upon patients, says this writer, (*British Medical Journal*, Feb., 1875,) is the careful avoidance of sudden, hurried or violent motion. Rapid walking, lifting weights, prolonged traveling, must be shunned.

The second rule, and one not to be neglected without great risk, is the avoidance of partaking at any one time of a large, distending meal. The sudden death of heart disease is very apt to come shortly after or during a hearty meal. The food, therefore, should be easily assimilable, and taken in small quantities at frequent intervals.

The third rule is to avoid taking much liquid, especially cold liquids. By exciting nerve irritation, they produce cardiac syncope and sudden death.

Lastly, a patient, laboring under serious heart disease, should, as a rule, be absolutely forbidden to indulge in sexual intercourse.

DR. BEVERLY ROBINSON.

Blisters in Chronic Heart Disease. Dr. BEVERLY ROBINSON recently told the Practitioners' Society, of New York, (*Med. Record*, December 22, 1883,) that he had been using blisters over the heart in certain cases of chronic cardiac disease, with enfeebled action and accompanying dyspnœa. They were cases where the ordinary methods failed to give relief. The blistering seemed to strengthen the heart's action.

The Use of Thermal Baths. There is in Germany a summer resort by the name of Bad-Nauheim, to which for now almost a century thousands of patients, suffering from organic diseases of the heart, have been sent by their physicians. For many years this was done, simply because the experience of many had proven that all such cases vastly improved. Of late, especially by the laborious researches of Benedict, of Vienna, the causes inducing such beneficial changes in diseases, which, especially in their last stages, are usually very little amenable to treatment, have been scientifically investigated. It has been demonstrated that it is the carbonic acid in these waters and the temperature of the latter (70–80°) which exerts such a wholesome influence. With the spirometer, as well as with Waerenburg's pneumatic apparatus, it has been found that the capacity of the lungs to inhale and exhale air greatly increases by the use of these baths, and that they also have a similar tonic effect upon the main organ of circulation, as we know of digitalis that it possesses.

Venous stasis in any form, if induced by organic affection of the heart, and, therefore, also dropsies, usually disappear without any other treatment within two weeks after beginning of the cure. That this is not brought about by increased diaphoretic or diuretic functions, but in reality by giving the heart greater tone, has been proven by experiments and observations carefully made, beyond a doubt.

ANGINA PECTORIS.

DR. M. HUCHARD, OF PARIS.

The treatment of this writer (*Revue de Médecine*, 1883,) is *preventive* and *immediate*, and *constitutional*. To *prevent* attacks, he advises strict hygienic treatment: to lead a quiet life in a dry atmosphere; avoid climbing, and any fatiguing physical exercise; dispense with exciting foods; have many meals a day to avoid over-filling the stomach; an occasional period of milk-diet; abstinence from strong and spirituous liquors, moderation with tea and coffee; no tobacco, no sexual excesses, no fatigue of any sort, and no extensive movements of the left arm, which provoke attacks in some patients.

He relies, for the *immediate* treatment of the attack, on inhalations of *amyl nitrite*; and, second, hypodermic injections of *morphine*. He says treatment by chloral, electricity, or metallo-therapy is unwarrantable, because of their slowness and unreliability.

When syncope seems imminent, he advises the use of diffusible stimulants,—Hoffman's anodyne, ether, ammonia, caffeine, punctate cauterization, hot applications, and perhaps the faradic brush.

He thinks *amyl nitrite* acts by stimulating the intra-myocardic circulation, by producing relaxation of the spasm which obliterates the coronary arteries. It dilates the peripheral arteries, and increases the strength and working-power of the heart.

For the *constitutional* treatment of the disease, he places the greatest reliance on the continued administration of the *iodide of potassium*.

While not pretending to know the *modus operandi* of the drug in these cases, he is inclined to think with BROWN-SÉQUARD that it relieves by improving the nutrition of the nervous centres, as good nutrition lessens their reflex excitability.

DR. ALFRED L. LOOMIS, NEW YORK.

This author states that the only remedies he has found of service in delaying and rendering less severe the paroxysms of this disease are *iron*, *strychnine* and *arsenic*, administered daily in small doses. During the paroxysm, rest and free administration of *digitalis* are of the greatest service. He adds that opiates, chloroform, and nitrite of *amyl* should not be used.

DR. T. H. TANNER, LONDON.

This author has found the following mixture exceedingly valuable in angina :

458. R.	Spiritus ætheris compositi,	f. ℥j	
	Liquoris opii sedativi (Battey),	gtt. x	
	Misturæ camphoræ,	f. ℥ss.	M.

This amount as required.

(BATTEY'S sedative solution of opium is a secret preparation much used in England ; said to be made as follows :

459. R.	Extracti opii (hard),	℥ iij
	Aquæ dest. bul.,	℥ iij.

Dissolve and filter. When cold, add

	Spirit. vini rect.,	℥ vj
	Aquæ q. s. to make	℥ iij.

The above are English imperial measures. The dose is half that of laudanum.)

The patient should keep the above medicine by him, in order that it may be taken on the least threatening of an attack. A belladonna plaster worn constantly over the præcordial region may do good. The return of the seizure is to be guarded against by improving the general health, by great attention to diet, and by the avoidance of stimulants, strong exercise, and all mental excitement.

DR. F. W. MOINET, SCOTLAND.

During the paroxysm of angina, the treatment most efficacious in relieving the symptoms is that by diffusible stimulants and opium, hot brandy and water, sinapisms to the feet, and placing the hands in hot water. Galvanism, carefully and at first gently applied, is of service even in organic angina, by giving tone and stimulus to the heart's muscular fibre. When the circulation is languid and stagnant, bleeding is called for. Nitrite of amyl is occasionally useful. (*Edinburgh Medical Journal*, Feb., 1871.)

DR. HENRY HARTSHORNE, PHILADELPHIA.

Mustard plasters should be applied between the shoulders and over the chest, and the patient's feet placed in a warm foot-bath. Stimulants and anodynes are indicated, as

460. R. Sodii bicarbonatis, ℥iv
 Spts. ammon. aromat., f. ℥j
 Spts. ætheris compos., f. ℥j
 Tinct. cardamom. comp., f. ℥iij
 Aquæ camphoræ,
 Mucilag. acaciæ, āā f. ℥x. M.

A dessert or tablespoonful at a dose.

461. R. Tinct. rhei et sennæ, f. ℥iss
 Syrupi zingiberis, f. ℥iij
 Tinct. opii, f. ℥j. M.

A teaspoonful in hot water.

462. R. Chloroformi,
 Spts. ammon. aromat., āā f. ℥ij
 Spts. ætheris compos.,
 Tinct. opii camphor.,
 Mucilag. acaciæ, āā f. ℥ss. M.

A teaspoonful at a dose.

DR. G. W. BALFOUR.

In the treatment of angina pectoris, Dr. BALFOUR relies chiefly on the inhalation of *chloroform*; it not only relieves pain, but regulates the circulation. He has also a very high opinion of *arsenic*, stating that in all forms of cardiac angina it acts almost as a specific, and is a special tonic to the cardiac muscle. It has this effect even in very small doses. The preparation he uses is the liquor arsenicalis. (*On Diseases of the Heart and Aorta*, London, 1876.)

DR. FAUCONNET, PARIS.

463. R. Sulphureti antimonii aurati,
 Sulph. ferri exsic.,
 Sulph. quiniæ, āā gr. x. M.

Make twelve pills. One thrice daily.

Dr. FAUCONNET, from personal experience, recommends the *golden sulphuret of antimony*, in cardiac neuroses of various forms. Professor RICHARD MCSHERRY, of Baltimore, states that he has employed it largely, with generally satisfactory results. As organic heart diseases are nearly always attended with neurotic disorders, it is valuable as a palliative, even in incurable cases. Most patients bear well gr. j three times a day.

PROFESSOR G. SÉE, PARIS.

M. SÉE has no faith in nitrite of amyl, chloroform or belladonna, in this disease. Electricity is dangerous. During the paroxysm he repeats twice or three times hypodermic injections of *morphia* in full

strength, following it immediately by an enema of *chloral hydrate*, gr. xxx-xlv, aquæ, f. ʒv. Sometimes *acetate of ammonia*, which is an energetic cardiac stimulant, will be found greatly to facilitate respiration. After the attack the morphia should be used daily for a few days to prevent a recurrence; and as a prophylactic treatment, the only one of much value is the frequent use of bromide of potassium and digitalis, so as to regulate the circulation. (*Gazette des Hôpitaux*, June, 1876.)

DR. T. L. BRUNTON, LONDON.

On the other hand, Dr. BRUNTON (*British Medical Journal*, July 13th, 1872,) highly extols the *amyl nitrite*, and in the same journal (November 28th, 1874,) and elsewhere instances of its great value are given. About two or three drops, used by inhalation, are generally sufficient to afford relief. Fatty degeneration contra-indicates it.

DR. A. MUHLBERGER, OF HERRENALT,

believes that in concussion we have an excellent means of keeping in check the violence and duration of the sterno-cardiac attacks. He further notes that kneading and rubbing have long been tried and approved remedies against muscular cramp, and that as angina pectoris is nothing more or less than cramp of the cardiac muscle, it seemed to him likely a similar result would be obtained in parallel cases by the like means.

He finally sounds a note of not unheeded warning in the *Deutsch. Med. Zeitung*, against the too bold use of remedies in this affection, and we may add in others also, and recalls the case of an elderly medical man who was relieved of his angina only to die of the morphia that procured the relief.

DR. LOWENFELD

relates a case of angina pectoris, in which galvanization proved beneficial. The patient, a man aged forty-seven, was subject to attacks of the disease occurring every month or two. These were characterized by excited respiration, oppression, small, frequent pulse, sternal pain radiating to the left arm, and convulsive tremors of the limbs, and lasted about one hour. The heart was normal. The constant current was applied for one minute to each side of the neck, along the course of the pneumogastric. The sense of oppression was immediately relieved.

Ten such applications in the course of three weeks were followed by complete freedom from the attacks for more than two years.

DR. N. GALLOIS, PARIS.

464. R. Tincturæ digitalis,
Tincturæ belladonnæ,
Tincturæ valerianæ,
Spiritus ætheris compositi, āā f. ʒj. M.

Dose, ten to twenty drops during the access of pain; stimulating frictions over the sternum; and if the attack prolongs itself, hypodermic injections of atropia.

465. R. Quiniæ sulphatis, gr. xxx
Acidi arseniosi, gr. ss
Extracti valerianæ, q. s.

Divide into thirty pills. From two to four a day for persons subject to attacks, in order to lessen their frequency.

NOTES ON REMEDIES.

Aceticum Acidum Dilutum. Bathing the chest every morning with vinegar and water is said to lessen the liability to attacks.

Ammonii Bromidum. This substance, in doses of gr. xv-xx, three or four times a day, has been very successfully used to check paroxysms of angina by Dr. R. K. HINTON. (*Medical and Surgical Reporter*, March, 1875.)

Amyl Nitrite (see above).

Argenti Nitras, gr. j daily, in the form of a pill, is favorably spoken of by Dr. COPLAND.

**Arsenic.* Dr. ANSTIE says he has seen the most remarkable relief afforded by this remedy. It may be given in the form of Fowler's solution; in pill of arsenious acid; or (gr. $\frac{1}{8}$) of arseniate of soda, with extract of hop; by subcutaneous injection; or, by inhalation of the smoke of arsenical cigarettes. Dr. ANSTIE states that it will lessen the severity of the attacks, reducing them in time to mere "tightness of the chest."

Assafœtida, combined with opium and ether, is sometimes of service.

Camphor, in combination with opium or hydrocyanic acid, sometimes acts favorably.

Ether is praised as a palliative by Dr. ANSTIE, who considers it, however, as inferior to arsenic or strychnia.

Ethyl Bromide.

466. R. Ethyli bromidi, 1 part
Aquæ destill, 200 parts.

Take two to four tablespoonfuls. This is recommended by Dr. WILLIAM SQUIRE, of London, in anginal attacks (*Compend. Med. Sci.*, 1884.)

**Morphia*, by hypodermic injection, is praised by Profs. BARTHOLOW and SÉE.

Nitro-Glycerine acts similarly to amyl nitrite but more slowly and is more man-

ageable. The dose is one or more minims of a one per cent. alcoholic solution.

* *Strychnia*, by hypodermic injection (gr. $\frac{1}{120}$ – $\frac{1}{60}$) twice daily for several weeks, is highly lauded by Dr. ANSTIE, who has seen remarkable benefit attending its use in presence of a threatening paroxysm.

Quiniæ Sulphas has been advised in intermittent angina pectoris.

Sodium Nitrite, in doses of gr. iij–v, has been used by Dr. HAY and M. W. COLLIER, (*Compend. of Med.*, 1884,) and its effects in relieving the spasms were prompt and satisfactory. As it is a much cheaper drug than the nitrite of amyl capsules, it is a proper substitute in hospital practice.

Zinci Sulphas sometimes proves of great benefit, when persevered in during the intermissions.

Stimulants. The relief of pain is better accomplished by the use of ether, or of morphia or atropia hypodermically, than by large doses of spirits.

* *Turpentine* stupes, applied to the chest, often mitigate the severity of the paroxysm, particularly in the old and debilitated.

Issues. An issue at the nape of the neck sometimes acts favorably in mitigating the symptoms.

CARDIAC DILATATION.

DR. ALFRED L. LOOMIS, NEW YORK.

The indications are: 1. To maintain nutrition. 2. To avoid violent cardiac action. For the first, our author commends an exclusively milk diet, the fluid taken in small quantities at short intervals. "As a rule, it is always safe to daily administer *iron* to a patient with dilated heart." Slightly stimulating baths, flannel next to the skin, the avoidance of emotion, and the arrest of all exhausting discharges, are important. A dry, bracing atmosphere is most suitable. The most important remedy is *digitalis*, in sufficiently large doses to regulate the heart's action. As it loses its controlling power by continued use, it should not be exhibited indiscriminately.

J. M. DA COSTA, M. D., PHILADELPHIA.

467. R.	Pulveris digitalis,	gr. v	
	Extracti belladonnæ,	gr. j	
	Ferri redacti,	ʒij.	M.
For twenty pills. One thrice daily.			

468. R. Emplastri belladonnæ, 4x4.
To be worn over the cardiac region.

Belladonna is one of the best agents that can be employed to overcome irregularity of the action of the heart, and to relieve pain. *Digitalis* is also useful for the same purpose, especially where the action of the heart is feeble; it is the only sedative which will reduce the frequency and not the force of the heart. It may be combined as follows:

469. R. Ferri lactatis, ʒss
Pulveris digitalis, gr. v. M.
For twenty pills. One thrice daily.

470. R. Tincturæ digitalis, f. ʒss.
Ten drops thrice daily, in cases of dropsy caused by cardiac dilatation.

Also,

471. R. Pulveris jalapæ compositi, gr. x.
For one dose at night.

In such cases, a tablespoonful of lemon juice, three times a day, acts as a diuretic and stomachic. Baths, also, are advantageous. Dry cups applied to the chest relieve the pulmonary congestion. It is more important to start the secretions and relieve internal congestions, than to give tonics and iron, which find their appropriate place in after-treatment.

DR. ROBERTS BARTHLOW, OHIO.

472. R. Ferri redacti,
Quiniæ sulphatis, āā ʒj
Pulv. digitalis, gr. x. M.
Pulv. scillæ,
Make twenty pills. One three or four times a day.

Recommended as extremely serviceable in dilatation of the heart and mitral regurgitation, accompanied by cough, difficult breathing and general dropsy.

Another valuable remedy in dilatation, without valvular lesion, is *ergot*. It may be given with *digitalis*.

473. R. Extracti ergotæ fluidi, f. ʒijss
Tincturæ digitalis, f. ʒss. M.
A teaspoonful three times a day.

Ergot slows the action of the heart and produces a contraction of the smaller vessels.

DR. FELIX VON NIEMEYER, PROF. UNIVERSITY OF TUBINGEN.

Our author has convinced himself, by a great number of observations, that digitalis is a very efficient means of extemporarily strengthening the heart's contractile power, and of thus allaying dropsy and cyanosis. In dilatation of the heart, digitalis, combined with an exclusively milk diet, is an invaluable remedy. Dr. VON N. has repeatedly succeeded in obtaining complete removal of dropsical effusions of great magnitude, and producing considerable temporary relief, by this mode of treatment.

Iron, which, fortunately, no longer has the reputation of being "heating," should always be prescribed when the patient shows any signs of anæmia or hydræmia.

DR. WALSHE.

When dropsy appears in cases of dilatation of the heart, the diuretics which yield most relief are the acetate, nitrate, iodide, and bitartrate of potassium, nitrous ether, compound tincture of iodine, the infusion and spirits of juniper and gin. Hydragogue cathartics, elaterium, gamboge, citrate of potassa, and the compound jalap powder, also aid in subduing the dropsical effusion. The following formula is a useful one for the administration of elaterium :

474. R.	Extracti elaterii,	gr. $\frac{1}{8}$ -j	
	Extracti creasoti,	gr. ij	
	Extracti hyoscyami,	gr. ij.	M.
For one pill.			

NOTES ON REMEDIES.

Adonia Vernalis. This is a popular remedy in Russia for heart disease and dropsy. It stimulates the motor and inhibitory ganglia of the heart, and is not cumulative in its action.

**Belladonna* is prescribed by Prof. DA COSTA to overcome the irregularity of the heart's action, and to relieve pain.

Casca Bark in doses of five minims of the tincture (one to ten) thrice daily has proved serviceable in the hands of Dr. McCALL ANDERSON, of Scotland.

**Convallaria Majalis* is highly recommended in similar cases as digitalis has usually been prescribed for.

**Digitalis* is recommended in dilatation, by Prof. DA COSTA and Prof. NIEMEYER (see above), Drs. GULL, WILKS, FULLER, HANDFIELD JONES, TROUSSEAU, and others. The conclusions of Dr. MURRAY are as follows : 1. That digitalis will stimulate and strengthen a weak heart,

and that the weaker are the muscular tissues of the heart, the safer will be the administration of the medicine. 2. That in hypertrophied heart, it will fail to reduce the pulse either in frequency or strength, and in such cases, will prove dangerous. 3. That in a weak organ, acting, on account of its weakness, with great rapidity, it will reduce the number of its contractions, and, as it were, strengthen or tone then down. To strengthen and quicken the action of a weak, slowly-acting heart, and to reduce the number of the rapid strokes of a full heart, is, according to ANSTIE, to do the work of a true stimulant; bringing action up to the normal standard, on the one hand, and reducing it to a level on the other. (*Brit. For. Med. Chir. Rev.*, July, 1865.)

Ergot is a useful remedy in case of enlarged heart, (dilated cavities) *without valvular lesion*. It may be given with *digitalis*.

475. R. Ext. ergotæ fluidi,	$\frac{3}{2}$ ijss	
Tinct. digitalis,	$\frac{3}{2}$ ss.	M.
S.—A teaspoonful three times a day.		

Nitrite of Amyl, in dilated heart, has been found of value. Dr. E. HOLDEN has reported a striking case in the *New York Medical Record*, April 27th, 1878.

Opium, in small doses, (five minims of the tincture) in combination with *ergot* or *digitalis*, is considered by BARTHOLOW as a valuable tonic to a weak and dilated heart.

Strychnia Sulphate in doses of two or three milligrammes three times daily gives great relief.

Stropanthus. Dr. LOGAN, of Liverpool, (*Med. Compend.*, 1884,) has expressed the opinion that this agent is superior to *digitalis* as a cardiac tonic.

Dry Cups to the chest are useful to relieve the pulmonary congestion.

CARDIAC HYPERTROPHY.

DR. HENRY HARTSHORNE, PHILADELPHIA.

When there is dilatation with hypertrophy, the main principle is to avoid exciting causes, especially violent exercise, alcohol, and venery. Moderate venesection at long intervals, is appropriate for robust patients; occasional leeching or cupping over the heart may be done. As an astringent and cardiac sedative, *acetate of lead* is worthy of trial.

476. R. Plumbi acetatis,	$\frac{3}{4}$ ss	
Pulveris opii,	gr. v	
Confect. rosæ,	q. s.	M.
Make twenty pills. One thrice daily.		

Where abnormal rapidity of the heart is present, *digitalis* may be used unhesitatingly, but moderately. In violent acceleration of the pulse, we may, however, expect more from *veratrum viride* as a sedative and palliative (gtt. ij-v of Norwood's tincture every three or four hours, diminishing the dose if nausea follows). Benefit is obtained, also, in such cases, from the use of *wild-cherry bark*.

DR. LEOPOLD SCHROETER, VIENNA.

In idiopathic hypertrophy, two remedies, in particular, are useful. The first is *cold*, used persistently and regularly. It is best applied by the temporary or continuous use of ice-bags laid over the chest or worn suspended around the neck. It often quiets, in a wonderful manner, the over-activity of the heart. The second is *digitalis*. It very decidedly diminishes the frequency of the heart's action, and since it is usually required but temporarily, is quite in place in this disease. *Veratrum viride* and tartar emetic, recommended by some, our author considers are objectionable.

For the acute dilatation, which occurs in the course of inflammatory diseases, our object will be simply to maintain the energy of the heart's contractions for a short period; and for this purpose stimulants will be of great assistance, and amongst these he considers *tea* of special value.

ALFRED L. LOOMIS, M. D., NEW YORK.

This author condemns the practice of blood-letting, to which the full pulse and symptoms of cerebral hyperæmia have prompted some practitioners in this disease. Of all remedial agents to diminish the force of the heart's action, he has found *aconite* the best. His prescription is:

477. R. Tincturæ aconiti rad. (Fleming's), q. s.
Two to three drops every three or four hours.

"No drug so fully and promptly relieves the vertigo and other painful sensations that attend cardiac hypertrophy." It is contra-indicated if the dilatation overbalances the hypertrophy of the cardiac walls. *Digitalis* is contra-indicated unless there is degeneration of the hypertrophied walls.

DR. CONSTANTIN PAUL.

In *La Presse Medicale*, 1883, this writer states that in cases of hypertrophy of the heart, following lesions of the arch of the aorta

(endarteritis), or more remote morbid conditions, as that pathological state of the capillaries frequently preceding and always accompanying chronic cases of morbus Brightii, he found for the anæmia which characterizes such cases, no medicine as effectual as the following:

478. R.	Syrup simplic,	260 (℥. ⅔ viij)
	Syrup flor. aurant,	60 (℥. ⅓ xv)
	Ferr. citr. ammon. pyrophosph,	3 (gr. xlvij)
	Solut. Fowleri,	1.50 (℥. xxiv.)
M. S. Dose.—Tablespoonful.		

J. M. DA COSTA, M. D., PHILADELPHIA.

In the treatment of hypertrophy of the heart, as much rest as possible should be insisted upon. The patient must be directed to lie down for several hours each day. The sinking of the pulse, which naturally occurs in the recumbent position, makes this posture as potent a cardiac sedative as we possess. All stimulants to the action of the heart should be removed. This includes the removal of any dyspeptic symptoms which may be present, and of any other disordered function which can react upon the heart.

There are only two drugs in which our author has any confidence; one is *aconite*, or its active principle, *aconitia*, and the other *veratria*. These are the only medicines which directly and positively control the element of muscular power of the heart. *Digitalis* does not compare as a pure sedative with either *aconite* or *veratrum viride*. *Gelsemium* is useless; it has a false reputation. It is time lost to employ it. Hydrocyanic acid is often a useful and pleasant adjunct when there is gastric disturbance; alone it is of no value.

These principles apply both in the treatment of simple hypertrophy, and in that complicated with valvular disease. A certain amount of hypertrophy with valvular disease is beneficial, and judgment must be exercised in order to determine when to interfere, and when not.

479. R.	Tinct. veratri viridis,	℥. ⅓ iss	
	“ aconiti radiceis,	℥. ⅓ ss	
	“ zingiberis,	℥. ⅓ vss.	M.

Fifteen drops thrice daily, two hours after meals, in water.

The addition of the tincture of ginger causes the *veratrum viride* to be better borne by the stomach. Our author also frequently orders:

480. R.	Tinct. aconiti radiceis,	gtt. j
Thrice daily for many months, its effects being watched.		

In a large number of cases this remedy thus employed prevents the further growth of the heart, and in some it lessens the already existing bulk of the organ.

NOTES ON REMEDIES.

**Aconite* is one of the two drugs (the other being *veratrum*) in which most authorities have any confidence in hypertrophy of the heart.

Asparagin. This derivative from asparagus is said by Dr. MARY J. MATLACK (*Med. and Surg. Rep.*, July, 1879,) to be particularly serviceable in hypertrophy of the heart. The following formula is of value :

481. R.	Asparagin,	grs. x	
	Potass. bromidi.,	3 ij.	
	Sacch. alb.,	3 ij.	M.

Make ten powders, one thrice daily.

Camphor has been recommended, in doses of from grs. iij-xij, daily, as rendering regular the most tumultuous palpitations, and removing the dyspnoea attendant on hypertrophy of the heart with dilatation.

**Digitalis*. The best effects of *digitalis* in hypertrophy are obtained from small doses, large doses being regarded by many as hurtful, if not dangerous, in this affection. Drs. GULL, WILKS, FULLER, W. MURRAY, and HANDFIELD JONES regard *digitalis* as useless, and even dangerous, in simple hypertrophy, and would restrict its use to cases of dilatation and enfeeblement of the heart.

Fel Bovinum Purificatum has been recommended in certain forms of hypertrophy.

Plumbi Acetas has been recommended in the incipient stages of hypertrophy, but it is of doubtful value other than in relieving, in some cases, violent palpitations connected with hypertrophy.

Potassii Bromidum has been employed, but its utility is not established.

Potassii Iodidum, in small doses, *long continued*, seems to exercise, in some cases, a beneficial influence.

**Veratrum Viride* is one of the only two drugs (the other being *aconite*) in which Prof. DA COSTA has any confidence. (F. 479.)

Diuretics are strongly recommended by Dr. J. HOPE in hypertrophy, not only when there is dropsy, but also when there is none, as powerful derivatives.

CARDITIS, ENDOCARDITIS AND PERICARDITIS.

DR. CONSANTIN PAUL.

In his last work (*Mal. du Cœur.*, 1884), this writer disapproves of cupping and mercurials. He relies largely on *blisters*, but they must be made large, so as to cover the whole cardiac space. Their prompt action can be observed by the stethoscope. The *ice-bag* is also valuable where there are local pains, tumultuous action of the heart, and anxiety. It may be applied from one to three hours at a time. The pulse and temperature should be watched and the bag removed as soon as they return to the normal. *Digitalis*, usually in small doses, is indicated by the rapid action of the heart.

The pain may be relieved by opium internally or by subcutaneous injections of morphine. Insomnia may be combated by *chloral* in doses of grs. xx-3j. The depression of the vital energies must be met by wine, brandy punch, and quinine. The chilly sensations sometimes felt may be counteracted by subcutaneous injections of ether or camphor.

When effusion has taken place, blistering may be tried, but it usually disappoints. The same may be said of diuretics, and hydragogue cathartics, such as squills, nitrate and acetate of potash, jalap, colocynth, etc.

Paracentesis is a last resort. Few of the patients recover, but when suffocation is threatened, it should be performed. The place of election is the fifth intercostal space, to the left of the sternum, care being taken to avoid the internal mammary artery. A preliminary exploration should be made with the hypodermic needle. The trocar can then be inserted safely. After the fluid has been withdrawn, the following may be injected :

482. R. Tinct. iodinii,
Potassii iodidi,
Aquæ destill.,

f. ʒss
gr. xv
f. ʒij. M.

As a rule the patient experiences immediate relief, but it is rarely permanent, as the fluid is reproduced.

DR. BERNHARD KRAUS, VIENNA.

The principles of treatment of pericarditis embrace absolute rest in the recumbent position, cold applications, as ice bladders or cloths

wrung out in ice water and laid over the præcordial region, and a mineral acid. (*Diagnose und Therapie der Krankheiten des Menschen*. Wien, 1877.) *Digitalis* should be given promptly, but after two or three days, be suspended, as it not only is alleged to have a cumulative action, but according to SKODA, a debility of the contractile power of the heart appears, which must be met by quinine, opiates and alkalies. Collapse is to be combated by wine, coffee, musk, carbonate of ammonia, and other stimulants.

As a diuretic :

483. R. Extracti digitalis, gr. xv
 Aquæ petroselinæ, f. $\frac{3}{4}$ vj
 Oxymel. scillæ, $\frac{3}{4}$ vj. M.
 A dessertspoonful every two hours.

Or ten drops of the liquor ammoniæ acetatis in a cup of warm infusion every two hours. Should there be much discharge into the pericardium (hydropericardium,) the diet should be nourishing, tonics be exhibited, and reliance be placed upon diuretics as :

484. R. Infusi bacarum junip.,
 Liquoris potassæ acetatis, āā f. $\frac{3}{4}$ iv
 Liquoris ammoniæ succinici, f. $\frac{3}{4}$ ss. M.
 A dessertspoonful every two hours.

The external application of "Hahn's diuretic liniment," has been found of use. It is :

485. R. Olei terebinth., f. $\frac{3}{4}$ j
 Vitelli ovi, No. ij
 Aq. menth. piper., f. $\frac{3}{4}$ xx. M.
 For external use. To be rubbed over the kidneys two or three times daily. Flannel around the loins.

If the dyspnœa increases, paracentesis of the pericardium is indicated. It is performed in the fourth intercostal space on the inner side of the line of the mamma. Frequently it results in permanent improvement.

DR. HAYDEN, DUBLIN.

Our author treats acute pericarditis in young and vigorous subjects with a saline aperient followed by leeches—ten, twenty or thirty, as the case may demand—following them with a warm poultice. If the impulse continues strong and abrupt, the rhythm regular, the sounds

normal and sharp, and the præcordial pain and oppression persistent, he would not hesitate to repeat the leeching. This should be followed by

486. R. Hydrarg. chlor. mitis, gr. ss
Pulv. antimonialis, gr. j. M.

One such powder every hour until slight salivation is produced.

When the improvement has well commenced, frequently-repeated moderate doses of wine are given to remove the cardiac debility and relaxation of the coronary arteries. To ensure sleep at night a moderate dose of some opiate is exhibited. The diet should be bland and nutritious. When the acute inflammatory stage is subdued, generally sustaining measures should be adopted, with iodide of potassium and the application of a blister. If the heart becomes tumultuous and irregular, digitalis is useful. When extreme effusion defies remedial agents, paracentesis must be resorted to. The point of election is that of maximum dullness, minimum impulse, and most distant cardiac sounds; usually a point one or two inches to the left of the sternum in the fourth or fifth interspace. Nearer to the sternum than one inch endangers the internal mammary artery. For acute endocarditis he relies on the same treatment.

DR. AUSTIN FLINT, NEW YORK.

On the subject of pericarditis, the author says blood-letting should be employed with great circumspection. Depletion by means of saline purgatives, with restricted diet, is indicated in the early stage. For some years the author has treated all cases of pericarditis without mercury, and he has found no reason to be dissatisfied with its omission. Opium should be administered. Locally, mild, revulsive and soothing applications. In the second stage, if the amount of effusion be sufficient to enfeeble the heart, it is highly important to remove the liquid as speedily as possible. For this end, the præcordia may be painted daily with the tincture of iodine, and small blisters may be applied, removing them as soon as vesication begins. Hydragogue cathartics and diuretics as in pleurisy, taking care not to depress the vital powers. The strength in this stage should be supported by alcoholic stimulants, tonics and nourishment.

ALFRED L. LOOMIS, M. D., NEW YORK.

Whatever the conditions which develop *endocarditis*, the patient

should be kept in bed. Opium, in small doses, is proper. The temperature of the room should always be above 70° Fah. The chest should be covered with flannel, and not exposed. If the heart becomes feeble, digitalis is indicated. "Rest, opium, iron, and the most nutritious diet, with occasional use of stimulants, constitute the most serviceable agents to be employed in the treatment of this affection."

In the *acute ulcerative form* of endocarditis, large doses of quinine and stimulants are our only dependence.

In *pericarditis*, the same cautions about rest, etc., are to be observed. Blisters, blood-letting, diuretics, and cathartics, our author considers useless; digitalis of doubtful value. "Opium, iron, moderate stimulation, rest in the recumbent posture, anodyne poultices, and concentrated nutrition, are about the only measures which he has found of service in the treatment of acute pericarditis." Aspiration he commends only when it can be positively determined that pus is in the pericardium.

DR. N. O. PARKS, OF RHODE ISLAND.

This physician stated to the Rhode Island State Medical Society, 1884, that he had employed *topical cardiac blistering* in acute rheumatic pericarditis in seven cases, and in one of them no other measures whatever were used; the result being in this, as in the other cases, a prompt and complete cure. In the remaining cases, salicin, or one of its allied compounds, and ammonium carbonate, were given internally. *Oleum gaultheriæ* was used in some instances—internally in doses of fifteen drops once in two hours, and applied locally to the inflamed joints, combined with an equal amount of lin. camphoræ comp. The series described included cases of acute articular rheumatism, and rheumatic gout, so-called. In some of the cases an endocardial murmur was present. Usually, one blistering sufficed, but in case of relapse it was repeated. The writer expressed his belief that we have, in the method under consideration, the most valuable adjunct to our means of successfully treating rheumatic carditis.

DR. SHINGLETON SMITH, LONDON,

regards (*Lancet*, August 22d, 1874,) the use of *aspiration* in copious effusion into the pericardium, as of great value. From ten to fifteen ounces of fluid may be withdrawn, and the relief is immediate. The needle may be introduced into the space between the fourth and fifth rib, about two inches to the left of the middle line of the trunk.

NOTES ON REMEDIES.

Aconite is recommended by Dr. RINGER in pericarditis when the heart throbs violently, and thus produces extreme pain.

Antimonii Ses-iodidum is sometimes employed.

**Digitalis*, combined with hyoseyamus tincture (ââ ℥xv-xx), is recommended by Dr. J. HOPE in pericarditis, after the acute symptoms have been subdued. Dr. BEVERLY ROBINSON, of New York, considers it *the remedy* indicated for the treatment of the pulmonary complications of acute heart disease. (*Medical Record*, November, 1877.)

Hydrargyrum is considered by Dr. A. T. H. WATERS as rarely, if ever, desirable in pericarditis, except as a purgative. He prefers to the mercurial treatment and venesection recommended by Drs. LATHAM, HOPE, and others, the milder but more effectual remedies of opium, bicarbonate of potash, the moderate and cautious use of stimulants, large linseed poultices to the chest, and blisters in the advanced stage, with good nourishment throughout the disease.

**Opium*, in grain doses, every three, four, six, or eight hours, is recommended by Dr. A. T. H. WATERS in pericarditis. The great relief to pain which often follows the local abstraction of blood in this disease, may be equally obtained by the administration of opium.

Potassii Iodidum, in doses of gr. iij-v ter die, is sometimes given with advantage in chronic pericarditis, appearing to favor the absorption of effused fluid.

Salicin. Dr. MACLAGAN, of London, endorses the views of those who hold that salicin possesses the power of preventing cardiac complication in rheumatism. He considers small doses ineffectual, and recommends twenty to forty grains every hour for six hours, or until relief from pain is experienced; by its *early* use he thinks we may *prevent* valvular inflammation.

Veratrum Viride is recommended by Dr. WARING-CURRAN (*The Practitioner*, August, 1868,) in pericarditis. He gives it in the form of an extract made by inspissating the juice of the root, and combines with it gr. j of calomel, every two hours, carefully watching the effect. A combination of opium and veratrum has been highly extolled in pericarditis and valvular disease by Dr. J. L. LYNCH, of Baltimore. It avoids all the unpleasant gastric effects usually attendant on veratrum, and is palatable. It is:

487. R.	Tinct. verat. vir.,		
	Tinct. opii,	ââ	f. ʒj
	Sodii bicarb.,		ʒij
	Sacc. alb.,		ʒiv
	Aquam q. s.,	ad	ʒvj. M.

S.—Tablespoonful every two or three hours.

For children, the quantity of opium should be reduced one-half, and the dose graduated for their ages.

Blisters are hurtful in the early stages of cardiac inflammation, but useful in the advanced stages.

**Poultices*. Large linseed-meal poultices are of signal service in the earlier stages of inflammatory action.

**Turpentine* stupes, which can be employed at a far earlier period than blisters, are often of manifest advantage in relieving pain and distress, and arresting inflammatory action.

Issues in the præcordial region frequently give relief in chronic inflammation of the heart or its membranes.

RHEUMATIC CARDITIS.

**Colchicum*, with alkalies, etc., is, according to Dr. FULLER, necessary in rheumatic carditis, to counteract and get rid of the materies morbi, without the removal of which it is difficult to conceive that a cure can be effected.

**Hydrargyrum*, combined with opium, and carried to the extent of slight salivation, is strongly recommended by Dr. FULLER in rheumatic inflammation of the heart, occurring in young, robust subjects, but he does not advocate its use in weakly, irritable and unhealthy constitutions.

**Opium*, in large and repeated doses, is strongly recommended in rheumatic pericarditis by Drs. WALSHE, FARRE (*St. Bartholomew's Hospital Report*, 1867,) and others. If the surface has been blistered, *morphia* may be used endermically; and often, when opium internally is inadmissible, benefit is obtained from *morphia* hypodermically. Dr. FULLER says that in rheumatic carditis, opium in *full doses* is indispensable in every case. In the weak and irritable, where mercury is of little service, it proves particularly valuable. It is, of all remedies, that which comes most powerfully in aid of blood-letting and mercury. If, after all active symptoms have subsided, much irritability of the heart remains, occasional doses of opium and *digitalis* should be given, and an opium or belladonna plaster applied to the chest, while the general health is being attended to.

Quinina Sulphas, in doses of gr. ijss, combined with gr. xx of carbonate of potash, given in mucilage every four hours, was found by Dr. WINN, (*Lancet*, November 14th, 1868,) highly beneficial in a case of rheumatic pericarditis attended with delirium.

Salicin, in scruple doses, every four hours, has been found by Dr. G. P. MAY of great use. (*British Medical Journal*, January, 1877.)

Stimulants. Alcoholic stimulants are often very beneficial in rheumatic pericarditis, especially when the attack is characterized by great pain, sleeplessness, and jactitating movements of the limbs. In this form of the disease, Dr. ANSTIE regards alcohol as superior to opium as an anodyne and specific. It should be given in repeated small doses, the production of even the minor signs of intoxication being carefully avoided.

Blood-letting. Dr. GARROD states that so much relief follows the application of leeches to the cardiac region in rheumatic pericarditis, and in endocar-

ditis, that he has no hesitation in recommending the measure, the bleeding never being allowed to produce any appreciable weakening of the patient. Generally, from three to twelve leeches are sufficient. The slow loss of blood by leeching is preferable to the more expeditious abstraction by cupping.

**Blisters* are recommended by Dr. FULLER in the advanced stages of rheumatic carditis, when effusion has taken place, as, of all local remedies, the most serviceable, their virtue being most unequivocally displayed when the amount of fluid is greatest.

FATTY DEGENERATION OF THE HEART.

MR. JOHN MARTIN, DUBLIN.

This writer (*Dublin Journal of Medical Science*, February, 1875,) believing that the fatty change is owing to an excess of carbonate of lime in the system, recommends the use of the *nitric* and *hydrochloric acids* in combination with iron. They should be taken in small quantities and for a long time. The diet should be one avoiding compounds of lime.

DR. A. T. H. WATERS, LIVERPOOL.

Ferri tinctura chloridi is highly recommended by Dr. A. T. H. WATERS, who believes that if it be given in small doses for a long time, it will often restore to an enfeebled and fatty heart a good deal of its vigor, and possibly of its structure. But its administration must be continued for months, and even years, being omitted from time to time as the digestive organs may demand. A chalybeate mineral water may be substituted for the tincture of chloride of iron, if the latter cannot be taken.

NOTES ON REMEDIES.

Arseniosum Acidum, is, according to Dr. LOCKIE, of great value in fatty degeneration, and this in spite of the fact that experiments tend to show that fatty degeneration of the heart is one of the results of feeding animals with arsenical preparations.

Cimicifuga, is, according to BARTHOLOW, more efficient and safer, in fatty heart, to relieve the symptoms, than *digitalis*.

Digitalis, according to Dr. A. L. LOOMIS, will sometimes afford relief when there is yielding of the cardiac walls. By some, fatty degeneration is held to contra-indicate *digitalis*; but such an authority as BALFOUR states that it may be given without danger and with advantage.

Ergot is recommended by Dr. WARING CURRAN, as of material use in severe cases.

Ferrum in any of the officinal forms will frequently prove of use.

Strychnia. PHILLIPS says this will sometimes do good in cases of fatty heart, when all other remedies have failed ; but it must be pushed cautiously, and stopped if signs of nervous worry appear. (*Mat. Med.*, 1879.)

FUNCTIONAL PALPITATION AND OTHER DERANGEMENTS.

DR. H. BERNHEIM, PARIS.

With regard to the treatment of functional deficiency of the heart, when there is a presumption that the cardiac tissue is healthy, and when at the same time the obstacle which produces asystole is not insuperable, *digitalis* is indicated. When either of these conditions is absent, *digitalis* is contra-indicated. Thus in those cases of palpitation in which strong, full pulse, arterial plethora, and abundant urinary secretion point to exaggerated cardiac activity, these drugs which, like *muscarin* and *aconitia*, abate cardiac velocity and arterial tension, should be prescribed, to the exclusion of *digitalis*. So also in the functional disturbances accompanying cyanosis, dropsy, œdema, etc., with high arterial tension, the use of *digitalis* is attended with danger. (*Leçons de Clinique Médicale*, 1877.)

J. MILNER FOTHERGILL, M. D., LONDON.

This writer commends (*Medical Times*, April, 1876,) as the most satisfactory prescription, the following, or an appropriate modification of it, in palpitation :

488. R.	Potassii bromidi,	gr. xv	
	Tincturæ digitalis,	gtt. v.	M.
This amount thrice daily.			

If there is anæmia present, the *potassio-tartrate* of iron will be found to combine nicely with this prescription. A belladonna plaster should be worn over the region of the heart. Of course, in the treatment of such a neurosal affection, it is well to cut off every form of irritation, and everything which tends to create nervous excitement. The mind and body should both be kept as calm and tranquil as is pos-

sible. The alimentary canal should be attended to; the condition of the reproductive system ought to be carefully scrutinized, and the habits of life taken into consideration. Where there is a lack of tone in the nervous system, it is well to give the digitalis with hydrobromic acid, in which quinine readily dissolves, so as to furnish a tonic in the mixture. Where there is anæmia, the bromide and the digitalis may be combined with the potassio-tartrate of iron, or a few drops of some preparation of iron may be taken twice a day, after food, with advantage.

J. M. DA COSTA, M. D., PHILADELPHIA.

Inquiry should always be made as to the cause, for the first step in the treatment is its removal. The cause may be found to be drinking of coffee, chewing of tobacco, smoking, alcoholic drinks, masturbation, etc.

In all cases of functional disorder of the heart, attended with palpitation, *digitalis* is very serviceable, more so than aconite. If there be masturbation as the exciting cause, the following is a useful combination:

489. R.	Potassii bromidi, Tincturæ digitalis, Infusi cascarillæ,	$\overline{3}$ vss $\overline{f. \frac{3}{2}}$ ijss $\overline{f. \frac{3}{2}}$ iv.	M.
---------	--	---	----

A dessertspoonful two or three times a day.

HENRY M. BOWDITCH, BOSTON.

In the *Boston Med. and Surg. Jour.*, 1883, Dr. BOWDITCH highly praises the following:

490. R.	Pulv. digitalis, Pulv. colchici sem., Sodii bicarbonatis,	gr. x gr. xx gr. xxx.
---------	---	-----------------------------

M. et div. in pil. No. 20.

These are to be taken three or four times daily at first; subsequently to be reduced until only one is taken at bedtime; the treatment to be continued for three to nine months.

He has used it for twenty-five years, and has found it to relieve even the most serious cardiac affections.

ROBERTS BARTHOLOW, M. D., PHILADELPHIA.

The violent and irregular actions of the heart occurring in hysterical subjects are immediately relieved by the use of the hypodermic syringe.

Morphia alone is used. In cases of dyspnœa, dependent upon dilated right cavities, pulmonary œdema and mitral disease, advantage is derived from the following:

HYPODERMIC INJECTION.

491. R. Morphiæ sulphatis, gr. xvj
 Atropiæ sulphatis, gr. j
 Aquæ destillatæ, f. ʒj. M.
- Filter. Dose, five minims (equal to one-sixth of a grain of morphia, and one ninety-sixth of a grain of atropia.)

DR. BOUCHUT, PARIS.

492. R. Assafœtidæ, gr. xxxvj
 Digitalis pulveris, gr. iij
 Extracti valerianæ, gr. vij. M.
- Divide into eighteen pills; one morning and evening, to remedy nervous palpitations of the heart. If the blood be impoverished, a ferruginous natural mineral water is a useful drink with the meals, mixed with wine.

NOTES ON REMEDIES.

Aconite, in small doses, often controls nervous palpitations of the heart.

Belladonna, in the form of a plaster, frequently does good.

The Bromides are often of service. According to Dr. ANGRISANI, of Bologna, bromide of potash has no effect on the muscular fibres or arteries of the heart, but is the most beneficial drug for purely functional palpitation, intermittence, irregularity, etc. (*Dobell's Reports*, 1877.)

Cactus Grandiflora is extolled by eclectic practitioners as an unequalled heart tonic in functional disorders.

Camphor, in doses of gr. iij-xij daily, is recommended in tumultuous palpitation.

Cimicifuga, in palpitation with dilated or fatty heart, is particularly serviceable.

Digitalis is recommended by Prof. DA COSTA as more serviceable than aconite, in all cases of functional disorder of the heart.

Eucalyptus. In the palpitations frequent in women about the change of life, this is an efficient drug.

Hyoscyamus, in large doses, is especially indicated in functional disturbance of the heart arising from emotion. Dr. HARLEY recommends, in these cases, the hypodermic injection of gr. $\frac{1}{8}$ of sulphate of hyoscyamia.

Morphia, hypodermically, is recommended by Prof. BARTHOLOW in violent palpitation produced by emotion or reflex irritation.

**Potassii Bromidum*, in doses of gr. v-x ter die, is recommended by Dr. RUSSELL REYNOLDS in palpitation or fluttering of the heart, arising from disturbances of the vaso-motor system.

Senega, in combination with hyoscyamus and ammonia, often proves useful in *hysterical palpitations*.

Veratrum Viride is efficient to reduce the heart's action.

EXTERNAL MEASURES.

Emplastra of belladonna or of veratria are powerful local sedatives. A caoutchouc bag, filled with pounded ice, suspended around the neck so that it rests over the heart, is an admirable calmant.

Posture. In many instances, moderate palpitations, not depending on organic disease, may be almost immediately arrested by bending the head downward, and allowing the arms to hang pendant. The effect is still more rapidly produced by holding the breath a few seconds, while the body is in this bent position.

Pressure on the Carotids. This is an excellent measure for palpitations of a nervous character, not dependent on organic disease. The compression should be exercised on both carotid arteries at once, and maintained only a few seconds. The relief should be instantaneous.

Pressure on the Pneumogastric. This measure has been urged by Dr. J. F. A. ADAMS, of Boston, as giving results very striking and suggestive. In one, violent nervous disturbance of the heart, with hysterical symptoms, came on a lady suffering from an attack of acute rheumatism. Compression was made, first of one vagus and afterwards of both. In a few minutes, the agitation had subsided, the heart's action became quieter and more regular, and, finally, all the nervous symptoms subsided. A few minutes afterwards, she sank into a steady sleep, and, on awakening, was perfectly quiet, and the cardiac symptoms never re-appeared.

IRRITABLE HEART.

PROF. J. M. DA COSTA, M. D., PHILADELPHIA.

This common condition of the heart, frequently the result of over-exertion, the demands of modern "high-pressure" life, the excessive consumption of meat and stimulants, etc., demands, in the first instance, *rest*, and the cessation of the habits which have been the exciting cause. The patients should lie down several hours daily. *Digitalis*, in doses of gtt. viij-x of the tincture, is the most useful drug. Where the impulse is forcible, however, aconite may be employed in place of *digitalis*. The dose is tinct. rad. aconit. (U. S. Ph.) m-i-j, ter die. In a few cases, this dose requires to be doubled. In cases of decided increase of the organ, aconite most shows its influence. No medicine

can then be compared to it. It may be used steadily, in moderate doses, for months. *Belladonna* is an efficient agent, especially in instances of irregular action. Where no irregularity exists, but merely rapid action, it rarely shows itself of decided use. It may be advantageously combined with *digitalis* and iron. Opium has some quieting action, but only of a temporary character. Hypodermic injections of morphia are excellent to relieve the cardiac pain occasionally present. Tonics are often excellent adjuncts to the treatment, but are insufficient to control the disorder. Zinc, tincture of the chloride of iron, and gentian, may be mentioned. Where the attacks of palpitation are violent, Hoffman's anodyne and rest in the recumbent position are called for. Hypodermic injections of morphia and atropia relieve the pain. A belladonna plaster, persistently worn, is often, also, productive of good. Counter-irritation is valueless. The treatment is never a short one.

VALVULAR DISEASE OF THE HEART.

ALFRED L. LOOMIS, M. D., NEW YORK.

All valvular diseases require rest, the avoidance of strong emotions, a nutritious diet, and temperance. If anæmia is present, iron is called for. *Digitalis* should be employed carefully; "it should never be resorted to in aortic regurgitation while the hypertrophy compensates for the dilatation." Dr. LOOMIS thinks the best mode of administering it is by infusion.

The following tonic combination is recommended by BARTHOLOW:

493.	R.	Pulveris digitalis,	℥ij	
		Ferri redacti,		
		Quiniæ sulphatis,	āā	℥j. M.

To make twenty pills. One to be taken two or three times a day.

There is considerable diversity of opinion in the books as to the use of *digitalis* in aortic cases, though in other forms of heart disease opinions are pretty well agreed. Some authorities would regard aortic insufficiency as a nearly absolute contra-indication to the use of the drug. Dr. G. W. BALFOUR regards it as the most valuable remedy in this lesion. Mr. THOMAS HAYDEN says: "Given as the tincture, and combined with iron and ether, I have rarely known *digitalis* to cause

unpleasant symptoms; on the contrary, in numerous instances it has been followed by the most favorable and satisfactory results. I have in a few instances been under the necessity of suspending the use of digitalis, owing to threatened syncope; but in the cases referred to there was fatty degeneration of the heart, in some instances with, and in others without, disease of the aortic valves." Mr. HAYDEN's prescription is: 10 minims each of tincture of digitalis, tincture of chloride of iron, and spirits of nitric ether, in an ounce of infusion of quassia, thrice daily.

Dr. BALFOUR, however, will not allow the fatty degeneration to be the explanation, as he quotes a very good case in which the drug was most useful, and says, speaking of fatty heart, "I have never seen any reason to withhold digitalis, and have always hitherto been rewarded by considerable and well-marked improvement." He prefers the tincture, and orders ℥v-xxx every four hours, to be suspended if there is any faltering of the pulse or nausea on movement. (*Edinburgh Medical Journal*, Feb., 1875.)

PROF. S. S. ROSENSTEIN, M. D., LEYDEN.

The indication in valvular disease is to develop and maintain compensatory action of the heart, and to moderate over-compensation. The diet should be nourishing, but not stimulating; coffee and tea should be weak; distilled liquors avoided; the exercise be moderate. Lukewarm baths followed by a cold rub-down attract the blood to the surface, and thus act favorably. Mentally, an "Olympian calm" should be warmly recommended. To allay the irritability of the cardiac muscle, cold to the præcordial region ranks first. The patient may carry, for a few hours daily, a gutta-percha bag filled with ice suspended over the heart. Counter-irritants are injurious. The sovereign remedy in every stage of valvular disease is *digitalis*. Dr. R. has found this combination excellent:

494. R.	Tinct. digitalis,			
	" valerian.,	āā	f. $\frac{2}{3}$ j	
	" ferri acetat. æther.,		f. $\frac{2}{3}$ iss.	M.
Twenty-five drops thrice daily.				

(The ethereal tincture of the acetate of iron, G. Ph., is made as follows:

495. R. Liq. ferri acetat., 9 parts
 Alcoholis, 2 "
 Ætheris acetic., 1 part. M.)

The patient should take this a fortnight, but no longer.

The swelling of the liver and gastric catarrh, which is apt to arise, may be met by the use of quassia, gentian, or rhubarb, and gentle aperients, like the saline mineral waters.

When dropsy occurs, which is usually first noticed about the ankles, acetate of potash is called for:

496. R. Potass. acetatis, 3 ij
 Tinct. ferri acetat. ether., ℥xlvi
 Syrupi rheædos, f. 3v
 Aquam, ad f. 3vj. M.)

A tablespoonful thrice daily.

Squill may be reckoned a tried diuretic in such cases.

497. R. Aceti scillæ, f. 3j
 Potassii bicarb., ad saturandum
 Aquæ cinnamon., f. 3viij. M.)

One to two tablespoonfuls thrice daily.

Respiration of compressed air sometimes gives great though not lasting relief, especially in mitral affections with stenosis. Digitalis combined with quinia is often also of great value; it should be given in continued small doses.

In the fainting and dizzy fits so often exhibited by patients with aortic stenosis, the prompt application of volatile stimulants, as HOFFMAN'S anodyne, wine, or ether, is urgently necessary; the head should be placed low, while the body lies horizontal. The active congestion of the brain which occurs in insufficiency of the aortic valves, must be combated by sinapisms or dry cupping to the neck, or small bleedings over the mastoid process.

PROFESSOR GERHARDT.

This writer (*Deutsches Archiv für Klinische Medizin*, Bd. V.) has treated fibrinous deposits and valvular diseases of the left side of the heart by the *inhalation of alkaline solutions*. He uses:

498. R. Potassii bicarbonatis, gr. v-viij
 Aquæ bullientis, f. 3j. M.)

For use as an inhalation.

He believes the frequent inhalation of this alkaline vapor prevents

fibrinous deposits, and being carried by the pulmonary veins directly to the left side of the heart, may dissolve already existing vegetations. He claims good results from its employment.

DR. B. W. RICHARDSON, LONDON.

In cases of fibrinous deposit, especially *in extremis*, this physician recommends that depressing influences be at once withdrawn. The distress present suggests opium, but opium is fatal. The body should be placed in absolute rest, concentrated food should be given, the temperature should be 60°–70°, and the body kept warm by external dry applications. Internally he administers concentrated food and liquor ammoniæ, ℞, in ice water every hour, with potassii iodidi, gr. v, every alternate hour. This treatment has been followed by a degree of success he had never anticipated.

PROFESSOR SÉE, PARIS.

This distinguished teacher recommends (*Union Médicale*, May, 1875,) the following combined treatment in the œdema and anasarca which so frequently complicate cardiac affections. He claims to have effected their speedy disappearance:

499. R.	Extracti scillæ,	gr. xv	
	Pulv. rad. scillæ,	gr. iss.	M.
For ten pills. Six to ten of these a day.			

Also,

500. R.	Potassii bromidi,	3j.
This amount to be taken daily, in divided doses.		

For continuous cardiac dyspnœa he has seen much benefit from iodide of potassium:

501. R.	Potassii iodidi,	gr. xxx	
	Chloral hydratis,	ʒi	
	Gum julep,	f. ʒiv.	M.
Take as required every two hours during the day.			

The Action of Convallaria. The general action of this recently introduced heart tonic may be discussed in this connection. The preparation employed may be the fluid extract, such as is prepared by Messrs. Parke, Davis & Co., of Detroit, the dose of which is fifteen to twenty drops to an adult every three hours until its effect is produced.

According to the observations of Prof. E. MARAGLIANO, of the diseases in which the good effects of the convallaria preparations were obtained, heart diseases occupy the first place. It acted well in valvular diseases, especially of the mitral orifice, with insufficient activity of the cardiac muscle. In nine of thirteen cases the preparations were decidedly advantageous. They increased the amount of urine, and regulated the heart's action. In many cases convallaria is preferable to digitalis. In four cases in which it failed, digitalis also failed. Convallamarin also acts most powerfully in cardiac disease.

Dr. CONSTANTIN PAUL considers it a true heart tonic and adds that it must be employed in large doses. He gives one teaspoonful of the tincture at once.

Dr. E. T. BRUEN (*Med. and Surg. Reporter*, 1883), says that in valvular disease it is most useful in cases of mitral obstruction. It loses its effectiveness in proportion as fatty degeneration is present. In advanced valvular heart-disease or cardiac failure in the catarrhal nephritic forms of Bright's disease, convallaria has not proven as useful as digitalis.

The drug is of great service in cases where palpitation and dyspnœa, rather than deficient cardiac systole, are the prominent features. Broadly speaking, cases of phthisis or asthma in which palpitation and dyspnœa are prominent symptoms, before cardiac failure from advanced fatty change ensues, are much benefited by this drug.

Its effect is often brilliant in purely functional heart disorder, especially palpitation with irregular cardiac action dependent on general debility, and in cases of anæmia or hysteria. It is sometimes an efficient remedy in the irregular action of the heart caused by tobacco.

ROBERT J. GRAVES, M. D., DUBLIN.

- | | | | |
|---------|---------------------|------------------|----|
| 502. R. | Potassii nitratis, | ℥j-ij | |
| | Tincturæ digitalis, | ℥ _{xxv} | |
| | Tincturæ hyoscyami, | ℥ _{xv} | |
| | Misturæ amygdalæ, | f. ℥ x. | M. |
- A tablespoonful from hour to hour in the œdema which accompanies disease of the heart.

DR. BOUCHUT, PARIS.

- | | | | |
|---------|---------------------|-------|----|
| 503. R. | Digitalis pulveris, | ℥iv | |
| | Morphiæ muriatis, | gr. v | |
| | Camphoræ pulveris, | ℥ss | |
| | Confectionis rosæ, | q. s. | M. |
- Divide into forty pills. One to be administered morning and evening, in organic affections of the heart.

504. R. Tincturæ digitalis, f. ʒivss
 Vini colchici, f. ʒ iss
 Potassii iodidi, ʒ ijss
 Syrupi sarsaparillæ, l. ʒ iss
 Aquæ destillatæ, f. ʒ ijss. M.

A teaspoonful three or four times a day in the anasarca of heart disease. The patient should be purged every third day with compound jalap powder.

505. R. Tincturæ aloes, f. ʒ j
 Tincturæ scillæ,
 Tincturæ digitalis, āā ℥xv. M.

For one dose every second or third day, for the dropsies of heart disease.

NOTES ON REMEDIES.

Aconite is contra-indicated or to be used with the utmost caution, in organic disease, where the power or force of the heart is impaired.

Arseniosum Acidum. Arsenic is a valuable cardiac stimulant, alone and as an adjunct to digitalis, where there is a failure of compensation.

Belladonna is recommended by Dr. A. T. H. WATERS in form of plaster or lotion over the heart, in palpitation combined with valvular disease, the patient at the same time being kept at rest in bed; rest alone is often efficacious. In severe cases belladonna is superior to digitalis (which see).

Casca Bark is thought by Dr. T. LAUDER BRUNTON to be a more powerful heart tonic than digitalis, and to be capable of relieving cases of advanced mitral disease where digitalis fails. (*British Medical Journal*, March, 1877.)

* *Digitalis* is recommended by Dr. A. T. H. WATERS in valvular disease, especially when there is hypertrophy of the heart. Dr. E. MACKEY (*British Medical Journal*, May 31st and July 11th, 1868,) thinks that neither *mitral regurgitation* nor *mitral obstruction* contra-indicates the use of digitalis, and gives numerous cases in which it was employed with advantage; but he regards its employment in *aortic regurgitation* and in *fatty degeneration* as contra-indicated or of doubtful safety. In any case where the heart's action is very feeble, Dr. M. considers that the commencing dose of the tincture should be ℥v-x every four or six hours; this dose may be increased to ℥xv-xx, or even ℥xxx, with benefit; but, as a general rule, such a dose as ℥xxx should not be repeated more than twice in the twenty-four hours. An equivalent dose is f. ʒss of the fresh infusion, or gr. j of the powdered leaves. The patient, before taking such a dose, must be kept recumbent at least half an hour, and kept so for two hours after. Dr. M. hesitates to prescribe a larger dose than ℥xv of the tincture to any patient following his occupation, and believes that half-drachm doses of the tincture, repeated several times at short intervals, especially in the upright position, may dangerously interfere with the circulation, whether by overstimulation or partial paresis. But, given in the manner directed, no accumulative action has been observed.

Jaborandi. Prof. RIENZI has exhibited powdered jaborandi about ʒj daily in divided doses in cases of aortic insufficiency with dyspnœa and painful palpitations. After a few days' treatment, the sounds were lessened, the pulse was less frequent, and the painful palpitations had ceased. (*Compend. of Med.*, 1884.)

**Morphia* has been employed hypodermically by Dr. CLIFFORD ALLBUT (*Practitioner*, December, 1869,) with marked advantage in diseases of the heart. He gives the injection in the evening, and enjoins perfect quiet afterward. His commencing dose is gr. $\frac{1}{8}$ or $\frac{1}{6}$ of the muriate, increased, in ordinary cases, to gr. $\frac{1}{4}$ for an adult. He finds it valuable in *mitral regurgitation*, but less valuable in mitral obstruction and in disease of the aortic valves; but in aortic disease, when the heart is big and pumping, it gives much ease. The urine, however, should always be examined, and the morphia given with caution or withheld, if albumen be found. Prof. BARTHOLOW also recommends morphia, hypodermically, in *mitral disease*, but is opposed to its employment in narrowing or obstruction of the aortic orifice.

Quebracho Aspidosperma, is reported to act potently in cardiac dyspnœa, Dr. MARIANA Y. LANION gives the following results from his study of it (*London Med. Record*, 1883). 1. Quebracho diminishes the frequency of the respirations and cardiac contractions. 2. Its action appears to be principally directed to the heart, strengthening and regulating its contractions, either directly or by the influence of the nervous system. 3. This action is evident and immediate. 4. It is the only remedy which has a manifest antidyspnœic action. Other reports to the same effect are given in *The Newer Materia Medica*, 1884, issued by PARKE, DAVIS & Co., Detroit, who prepare the drug for medical purposes.

**Senega*, combined with some other diuretic, is recommended by Dr. BARLOW in valvular disease of the heart, to promote the action of the kidneys and relieve the palpitation, especially the palpitation arising from aortic disease. He employs the following formula :

506. R. Spiritus ætheris compositi,	f. ʒij	
Tincturæ hyoscyami,	f. ʒ iss	
Decocti senegæ,	f. ʒ iij	
Misturæ camphoræ,	q. s., ad f. ʒ iv.	M.

Take one-fourth thrice daily.

Tinctura Ferri Chloridi is recommended in chronic valvular disease by various writers; given in long-continued small doses, as directed for fatty degeneration.

Veratrum Viride is often employed with benefit to the pulse.

Zinci Sulphas is favorably spoken of by Dr. BARLOW in *mitral disease*. He employs :

507. R. Zinci sulphatis,	gr. iij	
Extracti lupuli,	gr. vj.	M.

Make six pills. Take one thrice daily. The dose of the zinc should be gradually increased to gr. iij. When there is much palpitation, he advises its combination with camphor.

508. R. Zinci sulphatis, gr. j-ij
Camphoræ, gr. j
Extracti hyoseyami, gr. ij. M.

Make into pills. Take thrice daily. When there is much irritability of the stomach, the iodide should be substituted for the sulphate.

Diuretics are found beneficial by Dr. HOPE, in every stage of valvular disease of the heart. They draw off the serous portions of the blood, diminish the quantity without deteriorating the quality of that fluid, and thus relieve palpitation and dyspnœa, and obviate infiltration, without materially reducing the strength.

IV. DISEASES OF THE DIGESTIVE SYSTEM.*

Appetite, Loss of — *Biliary Calculi* — *Cholera Asiatica* — *Colic* (*Flatulence, Ileus, Colica Pictonum*) — *Constipation, Habitual* — *Diarrhœa* (*acute, chronic*) — *Dysentery* (*acute, chronic*) — *Dyspepsia* — *Gastritis and Gastro-Enteritis* — *Gastric Ulcer* — *Hematemesis* — *Hepatitis* (*acute, chronic*) — *Intestinal Worms* — *Jaundice* — *Liver: Functional Disorders of* — *Pyrosis* — *Vomiting*.

APPETITE, LOSS OF.

Although this is merely a symptom, and not a disease, the physician is constantly consulted with reference to it.

Hygienic measures are those which are indicated in simple anorexia, and will generally suffice.

Most of the "bitters," as calumba, gentian, quassia, chiretta, cascarilla, etc., are popular agents to increase the appetite. They exert a slight irritant action on the stomach, and in this indirect way assist both appetite and digestion. They should be taken a short time before food, as their effect soon wears off.

Cinchona or quinine is an excellent sharpener of the appetite, and also assists digestion. It is particularly useful in the case of elderly people and inhabitants of towns, who have been "run down" by confinement, sedentary occupations and impure air.

DR. N. GALLOIS, PARIS.

509. R.	Tincturæ nucis vomicæ,	gtt. v	
	Extracti gentianæ,	gr. xv	
	Syrupi aurantii corticis,	f. $\frac{3}{4}$ iss	
	Vini cinchonæ,	f. $\frac{3}{4}$ v.	M.

One-half to be taken half an hour before each of the two principal meals, in order to awaken the appetite.

* NOTE.—Diseases of the Teeth, Oral Cavity and Pharynx, including Tonsillitis, Pharyngitis, Caries, Odontalgia, Goitre, etc., are treated of in NAPHEYS' *Surgical Therapeutics*, Chap. X.

510. R. Rhei, gr. xxxv
 Aloës socotrinæ, gr. xxv
 Myrrhæ,
 Saponis, āā ʒj
 Spiritūs menthæ piperitæ, ℥ij
 Extracti taraxaci, ʒij. M.

Divide into forty-five pills; and order from one to three at a dose, as laxatives, and to stimulate the functions of the stomach.

511. R. Aloës socotrinæ, gr. xvj
 Ipecacuanhæ, gr. viijss
 Zingiberis, gr. xxv
 Syrupi, q. s. M.

Divide into sixteen pills; order one a day, before dinner, to stimulate the appetite and regulate the system.

512. R. Fel bovini purificati, ʒiv
 Ferri carbonatis, ʒss.
 Rhei,
 Extracti gentianæ, āā ʒiv. M.

Divide into one hundred and twenty pills; order six or ten a day, to combat a want of appetite.

DR. REECE, FRANCE.

513. R. Sodii carbonatis exsiccata, ʒj
 Extracti gentianæ, ʒij
 Zingiberis, gr. xij. M.

Divide into thirty-six pills, and order two morning and evening, as absorbents and stomachics.

PROF. FONSSAGRIVES, MONTPELLIER, FRANCE.

514. R. Extracti cinchonæ flavæ, ʒss
 Tincturæ nucis vomicæ, gtt. v
 Syrupi aurantii corticis, f. ʒ iss
 Claret wine, f. ʒ viij. M.

Take several tablespoonfuls at the commencement of the meals, to stimulate the appetite.

LONDON HOSPITAL.

515. R. Pulveris anthemidis, ʒss
 Pulveris rhei,
 Pulveris zingiberis, āā ʒij. M.

Divide into sixteen powders. Take one a day, an hour before or after meals, to stimulate the appetite and facilitate digestion.

DR. HUSS, FRANCE.

516. R. Pulveris nucis vomicæ, gr. xv-xx
 Pulveris quassia, āā ʒss. M.
 Cretæ præparatæ,

Divide into twenty powders, and order one three times a day, half an hour before each meal, to stimulate the appetite and favor digestion.

BILIARY CALCULUS.

DR. GEORGE HARLEY, OF LONDON.

This author states that it is very easy to prevent the formation of gall stones, because in the majority of cases cholesterin is their chief ingredient, and the abnormal formation of cholesterin can readily be prevented by foods and medicines. "Let the patient avoid an excess of fatty and fat-forming foods, either liquid or solid." Sugar and starch are to be especially avoided. Alkalies, especially carbonate of soda, will prevent the formation of gall stones.

In connection with the warm bath (already referred to) to favor the expulsion of gall stones, Dr. HARLEY recommends opium and belladonna, given every two hours until their physiological effects are produced. Belladonna may be used in form of liniment, plaster or suppository. Gentle friction with the warm hand from right to left, but not from left to right, over the seat of acute pain, together with kneading pressure downwards from gall bladder towards navel, may also be had recourse to; and during the whole time this treatment is being carried out, let the patient drink freely of alkaline warm water. He usually recommends a teaspoonful of bicarbonate of soda to the tumbler of warm water or milk and water. The administration of a brisk alkaline mercurial purgative ought under no circumstances to be omitted. He uses

517. R.	Pulv. hydrarg. c. creta,	gr. viij	
	Pulv. rhei,	gr. iv	
	Magnesia,	℥i.	M.

When head symptoms supervene he gives repeated doses of

518. R.	Ammonii chloridi,	gr. xx	
	Pulv. antimonialis,	gr. iij	
	Aq. sambuci,	℥ ss.	M.

DR. JOHN A. OCTERLONY, KENTUCKY.

This practitioner records his observations in thirty-five cases of cholelithiasis. (*Transactions of the Kentucky State Medical Society*. 1877.) The indications are two: first, to relieve the pain during the attack; and, secondly, to prevent its recurrence.

1. For the immediate relief of the patient, hypodermic injections of

morphia are most appropriate. Opiates should never be given by the mouth nor in solid form for the relief of gall-stone colic. In some cases where the pain is so atrocious that morphia seems to be inadequate, chloroform or ether may be given by inhalation. *Chloroform* has also been given internally, f. ʒj every hour, while the pain lasted, and as much after each meal for five days longer. It is asserted to melt away gall-stones of considerable size. Such large doses of chloroform may, however, produce alarming symptoms, and must be used with caution.

The general prolonged hot bath is sometimes of signal benefit. Surrounding the patient's waist with a sheet wrung out of hot water is a good substitute. Leeches to the sides are sometimes effectual when other measures fail.

Emetics ought to be rigorously proscribed.

2. To prevent the recurrence of the paroxysm, any gall-stones remaining in the gall-bladder must be dissolved, and their formation prevented.

The *diet* must be carefully watched. Highly-seasoned food, malt liquors, fats, sweets, and rich viands must be forbidden. The meals should be at *short intervals*, three or four a day. As a remedial measure to prevent the formation of the stones, Dr. BUCKLER has proposed succinic acid and peroxide of iron, on account of the large amount of oxygen contained in both of them, which is almost wanting in cholesterine. These can be combined in the *hydrated succinate of peroxide of iron*, and should be taken continuously for six months according to the following formula:

519.	R.	Hydrated succinate of peroxide of iron,	℥ ^{iss}	
		Distilled water,	ʒvjss.	M.
A teaspoonful after each meal.				

In almost every case Dr. OCTERLONY has used this salt with complete success. (Dr. BUCKLER's article is in the *American Journal of the Medical Sciences*, July, 1867.)

Carlsbad, Vichy, or Marienbad waters, and their counterparts in this country, are useful.

A remedy much employed and with good results by the late Dr. LEWIS ROGERS, of Louisville, is:

520.	R.	Ammoniae muriatis,	āā	ʒ ^{ss}	
		Extracti taraxaci,		f. ʒvj.	M.
		Aquæ,			
A dessertspoonful three times a day.					

DR. JAMES E. JACKSON, PENNSYLVANIA.

This practitioner states that the following combination forms the most active solvent for gall-stones, biliary secretions, etc., he has ever tried. In connection with its use the patient should avoid meat and fats, and confine himself to a milk and vegetable diet:

521. R.	Chloroformi,		
	Ætheris sulphurici,	āā	f. ʒss
	Olei terebinthinæ,		f. ʒj
	Sacchari albi,		ʒij
	Mucilaginis acaciæ,		f. ʒij.
			M.

A teaspoonful three times a day.

DR. DURAND, PARIS.

522. R.	Olei terebinthinæ,	f. ʒijj	
	Ætheris sulphurici,	f. ʒij.	M.

Half a teaspoonful night and morning.

This is the celebrated "solvent" of DURAND, endorsed by TROUSSEAU. It has unquestionable utility as a preventive of the recurrence of an attack. Alkaline drinks, such as Vichy water, or, as Dr. PROUT recommends, bicarbonate of soda, ʒij, in warm water, Oj, aid its action.

DR. M. SCHIFF, GENEVA.

The *cholate of soda* is recommended (*L'Imparziale*, Feb. 16th, 1873,) by our author in the treatment of biliary calculus. He says that biliary calculi are generally produced, not in consequence of an excessive formation of cholesterine, but because there is a deficiency of its solvents in the bile—the cholinates and cholates of soda and potash. He suggests the administration of these salts in cases of gall-stone; not to remove the cholesterine already deposited, which he does not think possible, but to hold it in solution and prevent further deposit. It has been proved that bile or a solution of biliary salts, taken into the stomach or intestine, is conveyed to the liver, and that the bile discharged is richer and more dense. He advises that the cholate of soda should be given in doses of $7\frac{1}{2}$ grains twice a day, until symptoms of disturbance of digestion or of circulation are noticed. Saturation of the system with the medicine is indicated by irregularity of the pulse, which becomes very slow during rest, and is sensibly accelerated by movement of the body or by slight excitement. When this occurs, the medicine should not be interrupted, but the

dose must be diminished. No good effect is to be expected unless the remedy be continuously administered for some time.

NOTES ON REMEDIES.

**Belladonna*. Dr. MURCHISON highly recommends the extract of belladonna, in half-grain doses, every two or three hours, in the passage of gall-stones.

**Benzoic Acid*.

523. R. Acidi benzoici, 3ij
Potassæ causticæ, ʒ ss (vel. liq. p. ʒss)
Aquæ destil., ʒ vj. M.
Dissolved with heat, and a tablespoonful taken in water thrice a day.

The above is highly recommended by Dr. HARLEY, in the treatment of impacted gall-stones.

Chioral is frequently employed with advantage, on account of its producing muscular relaxation.

**Chloroform* by the mouth, or in the form of inhalation, relieves pain, diminishes spasm, and does not interfere with the onward propulsion of the stone. It is also believed to prevent the formation of stones. D. BOUCHUT gives :

524. R. Chloroformi, f. ʒ ijss
Alcoholis diluti, f. ʒ ijss
Syrupi acaciæ, f. ʒ viij. M.
A small wineglassful thrice daily, to persons subject to hepatic colic.
Abstinence from fats, oils, fatty food, and strong wine.

Dr. TOURASSE prescribes :

525. R. Chloroformi, f. ʒ ss
Olei amygdalæ dulcis, f. ʒ j
Syrupi acaciæ, f. ʒ jss. M.
To be well shaken. A tablespoonful every quarter or half hour, in hepatic colic.

Ether, internally, has been recommended on account of its power of dissolving cholesterine, as well as because of its anti-spasmodic properties. A combination of equal parts of turpentine and ether constitutes the well-known remedy of DURAND for the solutions and cure of biliary calculi.

Olive Oil. Olive oil is frequently prescribed with good effect to facilitate the passage of gall-stones through the intestines.

**Opium* in full doses (gr. ij or gtt. xl of laudanum) by the mouth or rectum, repeated in half an hour, if necessary, generally relieves the intense pain attending the passage of biliary calculi better than any other remedy. It should be combined with the use of the hot-bath. The administration of ether or chloroform may also be advantageously resorted to. The *hypodermic injection of morphia* (gr. ¼), repeated, if necessary, is particularly useful from the rapidity of its action.

Podophyllin, gr. $\frac{1}{8}$ daily, is recommended by Prof. BUFALINI. It should be continued for many months.

Sodii Phosphas, in doses of \mathfrak{Dj} – \mathfrak{Zj} , administered for several months, before each meal, has been found by BARTHOLOW to be extremely efficacious in preventing the recurrence of attacks of hepatic colic. It is believed to act by relieving the catarrh of the duodenum which is present in most of these cases, and, by transference to the gall-bladder, forms a nucleus of mucus and bile, around which the cholesterine crystallizes.

Saline Mineral Waters, such as Carlsbad, especially the Sprudel Spring there, are extremely useful, if taken steadily, and supported by a strict regimen. All fat, butter, cheese, rich meats, spices, salt fish and meat, strong tea, coffee, and alcoholic beverages, must be strictly excluded. "Sprudel Salts" may be used instead of the natural water.

Nitro-Muriatic Acid Baths are a valuable remedy. The right hypochondrium may be repeatedly bathed with a solution of the acid, f. \mathfrak{Z} iij to a gallon of water, and the whole person rubbed down with it. Or a flannel roller, ten or twelve inches wide, and long enough to encircle the body twice, may be moistened with the fluid and worn around the waist.

CHOLERA ASIATICA.

DR. R. G. JACK, OF ENGLAND.

This author (*Lancet*, August 4, 1883,) is sure that the great failure in all treatment arises from the fact that all absorption is arrested and the medicine put into the stomach is returned to the basin. He relies mainly on morphia and Fowler's solution of arsenic; as follows:

526. R. *Liquor arsenicalis*, gtt. lxxx
Aquæ, ad f. \mathfrak{Z} ij. M.

S.—Teaspoonful doses every fifteen minutes until some effect is produced.

At the same time, a blister may be made over the epigastrium, with nitrate of silver. He would not omit the use of hot flannels and the free drinking of iced water. Having seen a great number of cases of cholera while in charge of troops between Marseilles and the Crimea, and afterwards in hospital and private practice in China, he bears painful testimony to the small effect produced by medicines given by mouth or rectum. Arsenic in large doses he has seen do some good, as also chlorodyne, but very little.

THE MEDICAL, STAFF U. S. ARMY.

The classical work on *The Cholera Epidemic of 1873, in the United States*, mainly prepared by Dr. ELY McCLELLAN, U. S. A., contains the well-digested experience of practitioners during the last cholera epidemic in this country. We quote some of the most successful measures adopted.

Dr. R. W. MITCHELL, of Memphis, Tenn., writes: "When satisfied that I had cholera to contend with, as indicated by rice-water discharges, vomiting, cramps, and shrinkage of extremities, I ceased medication by the stomach, and used the formula given, hypodermically:

527. R.	Acidi sulphurici,	3 ss	
	Morphiæ sulphat.,	gr. $\frac{1}{3}$	
	Spiritus vini gallici,	5 iss	
	Aquæ destillatæ,	3 iij.	M.

"The above was injected under the skin of the arms, legs, and over the stomach, every hour until the symptoms of the disease were relieved."

Dr. G. B. THURSTON, of Memphis, Tenn., writes: "The general treatment that I adopted was about as follows: Calomel and opium in alterative doses, say in two of the former to one of the latter, repeated every hour for several hours. In conjunction with this, used the following:

528. R.	Extracti cannabis indicæ,	gr. xvj	
	Camphoræ,	3 ss	
	Chloroformi,	3 ss	
	Olei terebinthinæ,	3 ij	
	Mucilag. acaciæ,		
	Syrupi simplicis,	ââ 3 ss	
	Aquæ cinnamomi,	3 j.	M.

Dose, a teaspoonful every one or two hours.

Dr. J. T. JONES, of Nashville, Tenn., reports that, in the cold stage of cholera, he has used *creasote* with greater success than any other article of the materia medica, and suggests the following formula:

529. R.	Creasoti,	gtt. j	
	Aquæ camphoræ,		
	Infusi gentianæ compositi,	ââ 3vj.	M.

At a dose, and repeated every two hours.

Dr. S. T. CHANDLER, of Campbellsville, Ky., who treated, during the epidemic of cholera in 1854, some sixty to seventy cases of the disease in its various stages, records his experience as having been

largely in favor of the use of *dilute sulphuric acid*, in full doses, and repeated every fifteen to thirty minutes, until the vomiting and purging are arrested. In the epidemic of the year 1873, the same plan of treatment was adopted, and with like results. "The acid relieves the nausea, arrests the vomiting like a charm, and gradually the dejections are stopped."

Dr. McCLELLAN adds, that the necessity of instituting treatment at the earliest moment after the occurrence of a diarrhœa, was most generally recognized throughout the area of infection. Whenever physicians, during the progress of the epidemic, ceased the attempt to diagnose between cholera, cholera-morbus and diarrhœa, and treated all cases that occurred as cases of cholera, the lists of mortality were reduced; while, on the other hand, increasing mortality was shown whenever too great reliance was placed upon diagnostic powers.

The evidence is conclusive that the exhibition of opium, followed by alterative doses of calomel, and absolute rest in the recumbent posture, almost invariably arrested the disease when in the premonitory stage. In the advanced stages, the entire range of the pharmacopœia seems to have been brought into use, with no better results than had been obtained in previous epidemics.

The experience of the writer is strongly corroborative of the beneficial results which may be obtained from the use of *sulphate of iron* and *dilute sulphuric acid* as prophylactics during an epidemic of cholera; and, further, that the most successful treatment of the disease is to be found only in all that is implied in the word sanitation.

Dr. JOHN M. WOODWORTH, Supervising Surgeon U. S. Marine Hospital Service, records the experience of his department in the following language: "Beginning with the year 1814, the cholera literature down to the present time abounds in proofs, clinical, physiological, pathological and meteorological, of the efficacy of *sulphuric acid*; and there can be no doubt, despite the dicta of the last International Sanitary Conference, that we possess in the mineral acids a certain means of prophylaxis against cholera. The lessons of the epidemic of 1873 point strongly to the value of sulphuric acid even as a therapeutic agent against the disease. According to Dr. McCLELLAN, mortality among cholera patients treated with acids was only eight per cent., while the lowest mortality rate where other remedies were used was twenty-three per cent., and the highest fifty-nine per cent."

DR. WILLIAM STEVENS, LONDON.

This author, in two epidemics, claims to have had very successful results from the *saline* treatment, in his hospital and prison patients. Cases presenting premonitory symptoms—diarrhoea and vomiting—were removed into an observation ward, where an even temperature was constantly maintained. A seidlitz powder was immediately administered; if sinking was felt, without purging, three or four teaspoonfuls of Epsom salts were added to the powder. On these agents acting, plenty of thin beef tea, well seasoned with salt, was given; if there was any pain, a sinapism was applied to the gastric region; and thirst was relieved with seltzer water, soda, or pure water *ad libitum*. Most of the cases were thus cured. If, however, cramps, coldness or shrinking of the pulse came on, the patients were considered as cholera cases in the second degree. The following was then administered:

530. R.	Sodii chloridi,	ʒ ij	
	Sodii carbonatis,	ʒ ij	
	Potassii chloratis,	ʒ ij	
	Aquæ,	f. ʒvj.	M.

Two tablespoonfuls in water about every half-hour.

If there was much irritability of stomach, a large sinapism was applied; if much heat or burning pain, an additional quantity of carbonate of soda was added to the mixture. In cases in the stage of collapse, a strong solution of the same salts, dissolved in hot water (100° F.) was thrown into the bowels, and repeated every two or three hours. Sinapisms were also applied to the stomach, between the shoulders, etc.; and in the cold stage, frictions with warm towels were used. A pure air for the patient to breathe was considered of the greatest importance.

In mild cases, the "wet-sheet-envelope" will favor reaction, but it has only proved mischievous in severe instances. When the vomiting is severe, or the thirst insatiable, nothing gives more relief than permitting ice to be continually sucked. The patient is also to be isolated as far as possible. He is to have plenty of fresh air; and care must be taken that the water he drinks is uncontaminated—particularly, that it has not been drawn from a well near any sewer. Moreover, his excreta should be received in a pan containing a disinfecting fluid, and then at once thrown away.

The greatest caution will subsequently be required for many days as to diet; not a few deaths have occurred from the too early use of ani-

mal food. As a rule, broths and farinaceous substances only should be allowed, without any solids whatever, until the renal secretion has been fully re-established, and all the symptoms have vanished.

DR. FLEMING, QUEEN'S HOSPITAL, BIRMINGHAM.

531. R.	Plumbi acetatis,	gr. xxiv	
	Liquoris morphiae acetatis,	f. ℥j	
	Acidi acetici diluti,	℥xij	
	Aquæ destillatæ,	f. ℥vj.	M.

A tablespoonful every two hours (an hour before or an hour after food is taken) in a tablespoonful of water; beef tea and milk, alternately, every four hours; warm negus in moderation.

The value of acetate of lead and opium in diarrhoea is well known. Dr. FLEMING calls attention to the *mode of administration*. The astringent action of both lead and morphia is in consequence of their absorption and conveyance in the blood to the affected part. Hence, the marked advantage of giving them in a thorough solution in distilled water, which promotes their passage into the blood. This is further secured by giving the medicine on an empty stomach before meals, and so avoiding, as far as possible, precipitation of the lead by contact with the gastric fluids. In the ordinary lead and opium pill, more or less lead is probably converted into the meconate; or the pill dissolving slowly in the stomach, in contact with the gastric secretions, the lead runs much risk of conversion into the insoluble chloride. Our author has adopted this mode of administration for many years, and speaks strongly of its efficiency. In the diarrhoea of children the same mixture, according to the following formula, gives most valuable results:

532. R.	Plumbi acetatis,	gr. xij	
	Liquoris morphiae acetatis,	℥xij	
	Acidi acetici diluti,	f. ℥j	
	Aquæ destillatæ,	f. ℥ij.	M.

A teaspoonful every five, six or eight hours, to a child one year of age.

WILLIAM SEDGWICK, M. R. C. S., ETC., LONDON.

Our author strongly recommends (the *Lancet*, August 19, 1871,) *dilute phosphoric acid* in half-drachm doses, combined with syrup of orange, and largely diluted with ice-water.

In cases of choleraic diarrhoea, and in the early stage of confirmed cholera, this remedy appears to be capable of checking the progress of the disease with greater certainty than sulphuric acid.

GEORGE JOHNSON, M. D., F. R. C. P.

Professor of Medicine in King's College, London; Physician to King's College Hospital, etc. In the treatment of cholera and choleraic diarrhœa, which is, in fact, cholera in a mild form, the main principle to bear in mind is, that the discharges are as essentially curative as in the eruption of small-pox. The discharges are not to be abruptly stopped by opiates. Experience has abundantly proved that this is a pernicious practice. Neither are they to be permitted to accumulate in the alimentary canal. There is one remedy which is almost universally applicable in all forms and stages of the disease, and that is an abundant supply of *cold water*, to flush the intestinal sewer and to wash out the poisonous discharges. A copious imbibition of pure cold water will suffice for the cure of most curable cases.

Palpation and percussion of the abdomen reveal the fact that there occurs not unfrequently a painful and sometimes a paralyzing over-distention of the bowel by rapidly effused morbid secretion. This, if not promptly relieved, may even go to the extent of causing a fatal obstruction. More especially is this likely to happen when the sensibility of the bowel has been deadened by opium. The plan to prevent and to remove this accumulation is to give some quickly-acting yet unirritating evacuant dose. For this purpose, *castor oil* is, on the whole, better suited than any other remedy. He directs, as early in the premonitory diarrhœa as possible, a tablespoonful of castor oil in a mixture of orange or lemon juice and water, or cold water, or other convenient vehicle. If this be vomited, it should be again given, and the patient directed to lie still and take no more liquid for half an hour, by which time the oil will have passed from the stomach into the bowels. Within an hour or two, the oil will usually have acted freely, when a tablespoonful of brandy, in thin arrowroot or gruel, is given; and if there be much feeling of irritation, with a sense of sinking, gtt. v-x of tincture of opium, in cold water. In this manner, most cases of choleraic diarrhœa are promptly arrested. Rhubarb (gr. xv of the powder, or f. ʒss of the tincture) may be employed, if objections exist to castor oil. If vomiting be present, it should be encouraged by copious draughts of tepid water; and if nausea exists without vomiting, especially if the stomach is supposed to contain undigested or unwholesome food, or morbid secretions, an emetic is advised (gr. xx of ipecacuanha, or a teaspoonful of mustard, or a tablespoonful of common salt).

The time to give *opium*, if at all, is in small doses, to soothe the bowel, after the expulsion of the poisonous secretions. Opiates are useless, or even dangerous, when the blood is poisoned, or when the bowel contains offensive morbid secretions. Opiates, in the early stages of diarrhœa and cholera, would be more frequently and decidedly injurious, were it not for the fact that their absorption is prevented by the rapid current of liquid which is being passed from the blood into the alimentary canal; therefore, they are quickly expelled, together with the morbid secretions, and they are powerless to arrest the discharges. (*The British Medical Journal*, August 19th, 1871.)

Sir THOMAS WATSON, Bart., M. D., etc., states, in the last edition of his *Practice*, that he firmly believes that the doctrines advanced by Dr. JOHNSTON are well founded, and approves his rules of treatment. No doubt the true indication of treatment is to stop the flow as soon as possible; but this may sometimes be best effected (as also in crapulous diarrhœa, and in the summer cholera of Sydenham,) by carrying off the offending matter.

JOHN MURRAY, M. D.,

Inspector-General of Hospitals, Indian Medical Service. When irritating or indigestible food in the bowels is the cause, as is frequently the case, of the early diarrhœa, it should be removed, if it has not previously been discharged in the evacuations, and a recurrence of the looseness guarded against. For this purpose, our author recommends the following cholera pill:

533. R. Pulveris opii,
Piperis,
Assafœtidæ,

To make one pill.

gr. j
gr. ij
gr. iiij. M.

It appears to check the looseness and stimulate the secretions, and does no harm if needlessly exhibited. It should be repeated should the looseness continue. It will cure most cases, and in all, restrain the symptoms until regular medical advice can be procured. It may be distributed to every house, and be available in a few minutes, whereas, the delay of a few hours may allow the disease to advance beyond control. These pills have been distributed in tens of thousands, in the towns and villages of India, with most satisfactory results. Some surgeons prefer red to black pepper, and others add camphor to the opium.

and assafoetida, and report favorably of the combination. They are distributed in the dispensaries and placed in the charge of the police in India. (*The British Medical Journal*, August 12th, 1871.)

In a discussion at the thirty-ninth annual meeting of the British Medical Association, between Drs. JOHN MURRAY and GEORGE JOHNSON, Dr. M. stated that Dr. JOHNSON seemed to confound elimination with purgation, and that nature was best assisted by restraining the further advance of the disease, of which the most dangerous symptom is looseness, whether naturally induced by indigestion, by improper food, or by purgative remedies. Dr. JOHNSON said that the main difference between Dr. M. and himself was with regard to the treatment of the early stage of cholera, Dr. M. considering evacuants dangerous and opiates safe, while he (Dr. J.) held an opposite opinion with regard to both classes of remedies. (*The British Medical Journal*, August 26th, 1871.)

DR. EDWARD R. SQUIBB, BROOKLYN.

534. R. Tincturæ opii, depurat.,
 Spirit. camphoræ,
 Tinct. capsici, āā f. $\frac{3}{4}$ j
 Chloroformi purificat., f. $\frac{3}{4}$ ij
 Alcohol, 95 per cent., q. s. to f. $\frac{3}{4}$ v. M.

Each fluid drachm, or teaspoonful, contains about one hundred drops, consisting of twelve minims of each of the first three ingredients, and four and a half minims, or eighteen drops, of the chloroform. Dose, for persons over eighteen years of age, a teaspoonful; for persons fourteen to eighteen years of age, a small teaspoonful; for persons ten to fourteen years of age, half a teaspoonful; for persons six to ten years of age, thirty drops; for persons two to six years of age, ten to thirty drops; for infants, one to ten drops, according to age. To be taken in water.

In time of epidemic cholera or diarrhœa, when any person has two movements of the bowels more than natural within twenty-four hours, the second one should be followed by a dose of this mixture; the dose to be repeated after every movement that follows. If the movements increase in frequency, or in copiousness, after the second dose of the medicine has been taken, a physician should be sent for at once, and a double dose be taken after each movement, until he arrives. Immediately after taking the first dose, the person should go to bed, and remain there for twelve hours after diarrhœa has entirely ceased.

Made a little more dilute, to adapt it better to extended popular use, it was largely applied in the last epidemic of cholera by the Metropolitan Board of Health of New York, and by the Boards of Health of

some other cities, and appears to have accomplished all that could be reasonably expected from any arbitrary formula.

HAMLIN'S CHOLERA MIXTURE.

No. 1.

- | | | | | | |
|------|----|--------------------|----|----------|----|
| 535. | R. | Tincturæ opii, | | | |
| | | Tincturæ camphoræ, | āā | 1 part | |
| | | Tincturæ rhei, | | 2 parts. | M. |

No. 2.

- | | | | | | |
|------|----|-------------------------|----|--------------|----|
| 536. | R. | Tincturæ opii, | | | |
| | | Tincturæ capsici, | | | |
| | | Tincturæ cardamom. co., | | | |
| | | Ginger, | āā | equal parts. | M. |

RUSCHENBERGER'S CHOLERA MIXTURE.

- | | | | | | |
|------|----|---------------------|----|--------------|----|
| 537. | R. | Tincturæ zingiberi, | | | |
| | | Tincturæ capsici, | | | |
| | | Tincturæ piperitæ, | | | |
| | | Tincturæ opii, | āā | equal parts. | M. |

CHLORODYNE.

- | | | | | | |
|------|----|-------------------------|----|-----------|----|
| 538. | R. | Sulphate of morphia, | | grs. lxxx | |
| | | Hydrocyanic acid. dil., | | | |
| | | Glycerine, | | | |
| | | Caramel, | āā | f. ʒ ss | |
| | | Extr. Indian hemp, | | ʒ ij | |
| | | Oil or peppermint, | | ʒ ss | |
| | | Oleo-resin of capsicum, | | gtt. xv | |
| | | Chloroform. pur., | | f. ʒ vj | |
| | | Alcohol, | | f. ʒ j. | M. |

DR. JOHN SULLIVAN, M. D. BRITISH INDIA.

In his recent work on *The Endemic Diseases of Tropical Climates*, (London, 1877,) this writer altogether rejects the treatment by elimination. The plan he has found most successful is: First, to prescribe complete cessation from all labor and fatigue out of doors, and the adoption of the horizontal position on the very first appearance of diarrhœa. Secondly, arrest of diarrhœa, "whether it exists in the preliminary state, or whether it takes the form of rice-water evacuations;" and for this purpose he prefers the employment of opium or laudanum, by means of suppositories and enemata. And, thirdly, he endeavors to supply the loss of fluid, by giving *ad libitum* "iced or cold water or tea;" and occasionally some moderately astringent cordial.

NOTES ON REMEDIES.

Acid Sulphuric. A combination of aromatic sulphuric acid with opium, is, according to BARTHOLOW, one of the most effective remedies we have in the treatment of cholera.

Aquæ Ammoniæ, when inhaled or taken internally, largely diluted, is a useful stimulant and restorative in the collapse of cholera.

Argenti Nitras has been recommended injected into the colon, by means of a long, flexible tube.

The following solution is thus injected:

539. R. Argenti nitratis, gr. xvj
Aquæ destillatæ, f. $\frac{3}{4}$ iv. M.

Ten minutes afterward the following enema is to be administered.

540. R. Tincturæ opii, f. $\frac{3}{4}$ iv
Gruel, f. $\frac{3}{4}$ vj. M.

Dr. WARING considers this treatment worthy of trial. In an epidemic of cholera at Assam, in 1853, Dr. BARRY used nitrate of silver internally with great success, giving one grain after each stool.

Arsenic. Fowler's solution with opium is sometimes valuable.

Atropiæ Sulphas, hypodermically, in dose gr. $\frac{1}{100}$ — $\frac{1}{40}$, is recommended by Dr. HARLEY as one of the most appropriate and useful means of resuscitation in the collapse of cholera. It stands at the head of all our stimulants in such cases, for there is no medicine which approaches belladonna in its simple, direct, immediate, and powerful influence in exalting the force and rapidity of the heart's action.

Calomel in small doses, ($\frac{1}{20}$ to $\frac{1}{12}$ of a grain every half hour) will sometimes check the vomiting.

**Camphor* is recommended by Dr. SYDNEY RINGER, from personal experience. He gives six drops of a strong alcoholic solution of camphor every ten minutes, until the symptoms have abated, and then less frequently. By this treatment, he generally at once controlled the vomiting and diarrhœa, and often altogether checked them, removed the cramps, and restored warmth to the extremities. An Italian physician, Dr. RUBINI, of Naples, is said to have treated with success nearly six hundred patients by a solution of camphor in alcohol at 60° over proof. The commencing dose was gtt. iv, every five minutes (in severe cases, gtt. xx, or more,) preserved in until reaction set in, the patient being well wrapped up in blankets. It should be given in sugar, and not in water, as in the latter the camphor solidifies and loses its power.

Cannabis Indica has been highly spoken of, ℥x-xxx of the tincture being administered in repeated doses.

Carbo Ligni, in drachm doses, has been employed with repeated benefit.

Carbolic Acid has, it is stated, (*Medical Press*, January 22, 1868,) proved successful.

Chloral is highly effective by subcutaneous injection; usually combined with morphia.

* *Chloroform*, in doses of ℥v-vij every hour or half-hour, often arrests the vomiting and relieves the spasms and cramps in the early stages of cholera. When the vomiting is excessive, a little chloroform, on lint, placed over the epigastrium and covered with oiled silk or gutta percha, will frequently relieve it. Chloroform liniment, alone or with turpentine, diligently used, is very effectual against the spasms and cramps. The spasms are also quickly relieved by chloroform inhalations, stopped short of complete anæsthesia. In giving chloroform internally, some prefer to combine it with the carbonated alkalies, soda or potash (grs. xl-lx) in solution; others, with the oil of turpentine (℥xx-xxx.) Dr. T. M. LOWNDES, of the Bombay army, advises (*British Medical Journal*, August 29th, 1868,) the following formula, which he has used for years in the treatment of choleraic diarrhœa, with the best results:

541. R.	Chloroformi,	℥xv-xx	
	Tincturæ opii,	℥x-xv	
	Spiritus vini,	f. ʒj	
	Aquæ,	f. ʒj.	M.

Chloroform, as a remedy in cholera, has the advantage that, being very volatile, it does not, like calomel and solid opium, accumulate in the stomach, and thus produce serious effects when reaction is established.

Cotoin. The *Med. Press*, 1883, states that cotoin was recommended in Egypt. For internal administration the use of an emulsion containing fifteen grains of cotoin in about four ounces was recommended, a tablespoonful to be taken every quarter or half an hour; or cotoin may be given as a powder in doses of one-third of a grain every half hour or hour.

Hydrargyri Chloridum Mite has, as Dr. WARING remarks, been given in cholera for upward of half a century, in every variety and stage of the disease, in every gradation of dose, from one grain to sixty, in almost every possible form of combination, with the view, by turns, of obtaining its purgative, or its cholagogue, or its stimulant, or its sedative action, and the reported results have been of the most diversified and unsatisfactory character. Dr. AYRE has reported great success from the following plan of treatment, which, probably, deserves the preference, if the mercury be employed at all; grs. j-ij of calomel, with ℥j-v of laudanum, are given every five, ten or fifteen minutes, omitting the latter when the dose has reached sixty or eighty drops. Perhaps a great measure of the success recorded of this treatment is due to the patient being allowed to drink freely of cold water, the colder the better. During the stage of collapse, the fact must not be lost sight of, in administering calomel and other powerful drugs, that the stomach has lost its power of absorption, exposing the patient to all the dangers of an accumulated drug when reaction sets in. *General mercurial inunction* is another plan that has had its advocates.

Sodii Chloridum was formerly given in large draughts of cold water, with the view of inducing emesis, and at the same time supplying the blood with the saline constituents supposed to be carried out of the system in the evacuations.

Strychnia has been used with success.

Sulphur was first suggested as a remedy by Dr. A. BLACKLOCK, of the Madras army, in 1848. He recommended, also, as a prophylactic, during epidemics, sulphur internally, and a diet rich in sulphuretted ingredients. Dr. J. GROVE (in his work *On Epidemic Cholera*, 1865,) highly praises this remedy, which he prescribes as follows :

544. R.	Sulphuris præcipitati,		
	Sodii bicarbonas,	āā	℥iv
	Spiritus lavandulæ compositi,		f. ℥xxiv
	Aquæ,		f. ℥lxxij. M.

The soda and sulphur are to be first thoroughly triturated together, in a mortar, the spirit of lavender gradually added, till the whole is well mixed, when the water is added.

Dose, two teaspoonfuls, in a little water, every two, three or four hours, in simple choleraic diarrhœa; but if the case is urgent, every ten or fifteen minutes. In sudden or severe attacks, ℥x-xxx of laudanum are given with the first dose.

GENERAL MEASURES.

**Ice* in lumps and iced drinks, given *ad libitum*, are grateful to the patient and highly beneficial in relieving the burning heat at the pit of the stomach and the intolerable thirst, arresting the vomiting and exciting reaction.

Stimulants. Champagne, ammonia and turpentine are among the best, but weak brandy and water is sometimes retained better than any other stimulant.

Turpentine Stupes applied hot successively to the abdomen, the cardiac region and along the spine and extremities, and turpentine frictions and enemata, are valuable aids to the other treatment.

Intra-Venous Injections. Some remarkable results have been obtained by the injection of a saline solution in the veins in cases of the *collapse of cholera*. Unfortunately, the appearances of improvement, which are very remarkable, are not usually sustained. Hodder, of Canada, reports three cases wherein he transfused *milk* in *cholera collapse*, and two recovered. When milk is used for transfusion, it should be fresh and directly from the cow, if possible, and its temperature should be 100° Fah. Not more than four to six ounces should be injected at one time, lest the heart be paralyzed by over-distention.

COLIC.

CHARLES MURCHISON, M. D., F. R. S., LONDON.

Flatulent colic will often be relieved by the various ethers and the essential oils of peppermint, anise and cajuput, by vegetable charcoal, galbanum and assafœtida. When, however, it is due to decomposition, from deficient or deteriorated bile, those remedies will be found most useful which act by checking decomposition, such as creasote, turpentine or carbolic acid.

545.	R.	Acidi carbolici fluidi,		
		Spiritus chloroformi,	āā	f. $\frac{3}{4}$ -iij
		Aquæ menthæ piperitæ,		f. $\frac{3}{4}$ j-iij. M.

Dose—a tablespoonful.

Or, a pill containing one drop of creasote.

Flatulence, and other dyspeptic symptoms arising from want of bile in the bowels, are also greatly relieved by the use of purified bile from the ox or pig, which may be given in doses of from three to six grains, about two hours after meals. As it is not desirable that the bile should come in contact with the stomach, it is well to give it inclosed in capsules, or in pills coated with a solution of tolu in ether.

The following recipe will be found useful for the same purpose :

546.	R.	Sodii chloratis,	gr. x	
		Aquæ menthæ piperitæ,	f. $\frac{3}{4}$ ss.	M.

For one dose.

DR. D. L. PHARES,

In the *Transactions of the Mississippi State Medical Association*, 1878, directs attention to the *mechanical treatment* of colic. This consists in simply supporting the patient in an inverted position—in other words, in standing him on his head. In some instances, cases that have for hours or days resisted all ordinary treatment, have by this simple means been relieved and permanently cured in from one to five minutes. Cases attended with most intense pain, vomiting and other phenomena of so-called “bilious colic,” have been thus cured.

MR. JONATHAN HUTCHINSON, LONDON.

In all cases of unusually severe colic, the possibility of *intestinal obstruction* must be considered. In such cases the following rules for treatment should be observed :

1. In all early stages, and in all acute cases, abstain entirely from giving either food or medicine by the mouth.

2. Use anæsthetics promptly. Put the patient under the full influence of ether; examine the abdomen and rectum carefully before tympanites has concealed the conditions; administer large enemata in the inverted position of the body; and, if advisable, practice abdominal taxis. If you do not succeed at first, do it repeatedly.

3. Copious enemata, aided perhaps by the long tube, are advisable in almost all cases, and in most should be frequently repeated.

4. Fluid injections may be sometimes replaced by insufflation of air in cases of invagination, since air finds its way upwards better, and is more easily retained. It is, however, somewhat dangerous, and has, perhaps, no advantages over injections with the trunk inverted.

5. Insufflation is to be avoided in all cases of suspected stricture, since the air may be forced above the stricture, and there retained.

6. Saline laxatives are admissible in certain cases where impaction of fæces is suspected, and in cases of stricture where fluidity of fæces is advisable.

7. Opium (or morphia) must be used in proportion to the pain which the patient suffers. It should be administered by the rectum or hypodermically, and should be combined with belladonna. If there be not much pain or shock, it is better avoided, since it increases constipation and may mask the symptoms.

8. A full dose of opium administered hypodermically will put a patient in a favorable condition for bearing a prolonged examination under ether, and attempts at abdominal taxis.

9. In cases of uncertain diagnosis, it is better to trust to the chance of spontaneous cure, or relief by repeated abdominal taxis, than to resort to exploratory operation.

DR. NATHAN S. DAVIS, CHICAGO.

In obstinate *intestinal obstruction*, not dependent upon invagination, but produced by irregular contraction of the muscular coat of the intestines, this physician has long used, with gratifying results, *tobacco enemata*. One drachm of chewing tobacco is put into one pint of boiling water, and when cool enough, one-half of this amount is injected into the rectum. The powerfully relaxing effect of the tobacco relieves the intestinal contractions, and soon produces copious evacuations. It

is, however, a remedy of extreme power, inducing faintness and great temporary prostration, and should be used with corresponding caution.

DR. EDWARD MONTGOMERY, ST. LOUIS.

This writer (*Half-Yearly Compendium*, January, 1878,) recommends in cases of faecal impaction, hot poultices to the abdomen, warm water and saline injections, and after the evacuation of the lower portion of the alimentary canal by enemata, purgatives by the mouth. In young, robust patients, at the early period of the disease, one-tenth of a grain of tartar emetic and one drachm of sulphate of magnesia, every hour, will very often aid in breaking up and removing the obstruction.

As it is very common for these cases of intestinal obstruction to be associated with conditions of torpor, or even paresis of portions of the bowel, the administration of the following pill will be found of great service:

547. R.	Strychniæ,	gr. j	
	Ext. belladonnæ,		
	Podophyllin,	āā	gr. jv
	Pulv. ferri sulphat.,		
	Aloes socot,	āā	gr. xx
	Syrupi simp.,	q. s.	M.

Divide into twenty pills; one to be taken every eight hours.

Even after the removal of the impacted mass, this pill will be found valuable in preventing a relapse, and in promoting regularity of the bowels afterward.

NOTES ON REMEDIES.

Ammonii Carbonas, in doses of gr. v-viij in some aromatic water, frequently proves useful, especially in the flatulence or atonic dyspepsia of hysterical females.

Anthemis, in strong infusion, or gtt. ii-ij of the volatile oil, frequently affords relief after the failure of other remedies.

Assafatida, in enema, is of especial value in the flatulent colic of hysteria.

**Belladonna* is probably the most efficient remedy in *intestinal obstruction*; gr. j-ij of the extract every hour, together with opiate fomentations to the belly and warm applications. The remedy should take effect in six or eight hours.

Carbo Ligni, inclosed freshly prepared in gelatine capsules, each containing gr. x of heavy vegetable ivory charcoal (dose, two, *i. e.*, gr. xx,) is a valuable remedy, from its gas-absorbent property. If administered in this manner, the large nauseous doses so frequently prescribed are avoided.

Caryophyllum in infusion, or the volatile oil (gtt. iiij-v) is useful.

Cinnamomum is a pleasant and excellent carminative.

Creasotum, in doses of gtt. j in pill thrice daily, will greatly relieve the distressing flatulence of hypochondriasis.

Fel Bovinum. Dr. W. MURRAY says in the *Lancet*, September, 1878: Were I called to a desperate case of obstruction of the bowels, I should first of all push the belladonna treatment to its fullest extent, and while the patient is fully under the influence of the belladonna, I should administer, by enema, eight ounces of ox-gall diluted with eight ounces of water containing a few crystals of washing soda.

**Nux Vomica* is valuable in obstinate flatulence. Dr. BARLOW recommends the following useful formula:

548. R.	Extracti nucis vomicæ,	gr. jss	
	Argenti nitratis,	gr. ij	
	Extracti lupuli,	gr. xij.	M.

Make six pills. Take one thrice daily. This formula is particularly serviceable when there is also pyrosis and heartburn.

Oleum Carui is often of benefit, in doses of gtt. ij-v on sugar.

Oleum Menthæ Piperitæ, with or without a few drops of laudanum, is a pleasant carminative.

Oleum Myristicæ, in doses of gtt. ij-v, affords relief. Grated nutmeg may be given in a little warm brandy and water.

Oleum Terebinthinæ. Dr. J. COPLAND remarks that, notwithstanding constant or even feculent vomiting, advantage will sometimes be derived from one to two tablespoonfuls of the unrectified oil of *turpentine*, taken with aromatics. He has seen the vomiting cease and the distention of the abdomen rapidly subside immediately after this dose, which should be repeated if rejected at first.

Potassii Permanganas. Dr. R. BARTHOLOW has found marked advantage from the use of this substance in the flatulence and dyspepsia so constantly attendant on obesity, as follows:

549. R.	Potassii permanganatis,	gr. vij	
	Aquæ destillatæ,	f. ʒij.	M.

Keep in a glass-stoppered bottle. Dose, a teaspoonful thrice daily.

Rheum.

550. R.	Magnesii carbonatis,	f. ʒj
	Pulveris rhei,	gr. xxx
	Tincturæ rhei,	f. ʒ ij
	Tincturæ opii,	℥ xv
	Spiritus anisi,	gtt. vj
	Spiritus menthæ piperitæ,	f. ʒ ij
	Aquæ destillatæ,	f. ʒ vj.

This is a popular remedy in the west of England, in intestinal pains. From three to six tablespoonfuls a day.

Spiritus Etheris Compositus is useful, particularly in the flatulence of hysterical females.

Spiritus Etheris Nitrosi, in doses of a teaspoonful in a cupful of any convenient vehicle, is a popular and efficacious remedy.

Zinci Sulphas has been recommended in flatulent affections of the bowels, especially of the colon. The following formula has been employed :

551. R.	Zinci sulphatis,	gr. xvijj	
	Pulveris opii,	gr. iij	
	Mucilaginis,	q. s.	M.

Make six pills. Take one four or five times a day, after a meal.
When the stomach can dispense with the opium, substitute extract of gentian or rhubarb.

Galvanism is often very serviceable in obstruction. Drs. BEARD and ROCKWELL recommended that the positive pole be applied along the spine, while the negative pole is passed gently over the abdomen, following the direction of the ascending colon. A case, accompanied with fæcal vomiting, is reported (*Dublin Quarterly Journal of Medicine*, November, 1864,) as successfully treated by the application of galvanism to the mucous surface of the intestine. A sponge with the metallic handle to which it was attached, was passed up the rectum two or three inches, whilst the other sponge was applied to the abdominal walls. The effect was immediate, the constipation being at once relieved and the patient recovering from an apparently hopeless condition.

CONSTIPATION—HABITUAL.

The habit of constipation may be broken up by *massage*, *change of diet*, *enemata*, *laxative mineral waters*, or *drugs*.

In regard to *massage*, or the movement cure, the following little operation executed by the patient upon himself, when properly performed, is valuable :

Place the tips of the fingers of the right hand exactly over the cæcal region, with *very slight* pressure ; carry them upward, along the ascending colon, to the right hypochondrium ; continue the movement *without any intermission* over the region of the transverse colon to the angle of junction with the descending colon ; stop not, but proceed downward, gently and steadily, to the iliac region of that side ; instead of the previously very gentle pressure, the finger must now be pressed firmly and deeply (without pain) into the pelvic cavity, and there retained for about fifteen seconds ; then remove the hand altogether, rest a few seconds, and repeat the procedure. This may be continued for the period of from a few minutes to a quarter of an hour or more.

Some little care and tactile dexterity are needed to do this properly; and where the hand is dry, or the cuticle thick and hard, it is advisable to slightly moisten the ends of the fingers. When the right hand is tired, the left can be used, and so alternately, but it is better not to alternate them too rapidly. If there be failure, it must not necessarily be given up. Invalids themselves will often fail; almost invariably, if their bowels are *extremely intractable*. But now the aid of a friend for passive movements may be invaluable. The medical adviser can give instructions regarding the precise anatomical relations of the parts involved, and the method of performance. The proceeding should usually extend over a period of from five or six minutes to occasionally twenty-five minutes.

For occasional use, in the treatment of *constipation in old age*, the following pill affords an excellent formula:

552. R.	Extracti colocynthidis compositi,	gr. viij	
	Extracti hyoseyami,	gr. ij.	M.
For one pill.			

DR. ROBERTS BARTHOLOW, PHILADELPHIA.

A proper regulation of the diet is of great importance in the treatment of *habitual constipation*. This usually depends on deficient secretion, or torpor (a paretic state) of the muscular layer of the intestines. Corn-bread, cracked wheat, oatmeal, bread of unbolted flour, fruits, and such vegetables as green corn, tomatoes, and celery, are indicated. Those troubled with habitual constipation, to a moderate extent, may overcome it by the daily use at dessert of a few almonds and raisins, about six of each. Hemorrhoids, due to congestion of the portal vein, or to constipation, are much benefited by the grape-cure or a diet of fruits and succulent vegetables.

J. M. DA COSTA, M. D., PHILADELPHIA.

553. R.	Podophyllin,		
	Extracti belladonnæ,	āā	gr. j
	Capsici,		gr. v
	Pulveris rhei,		ʒj.
			M.
For twenty pills. One three times a day.			

Belladonna is undoubtedly a stimulant to the muscular fibres of the intestines. It acts on them as it acts on the bladder; it stimulates to contraction. It also increases the action of purgatives, enabling the

physician to get along with smaller quantities of purgative medicine. Podophyllin is useful in torpor of the upper portion of the bowels, to increase the secretion of the liver.

554. R. Tincturæ gentianæ compositæ, f. ℥iij
 Tincturæ rhei dulcis, f. ℥j
 Tincturæ belladonnæ, f. ℥ iss. M.
 Dessertspoonful thrice daily.

555. R. Extracti gentianæ, ℥j
 Extracti nucis vomicæ, āā gr. ijss
 Podophyllin, gtt. xx. M.
 Olei cajaputi,
 For twenty pills. One twice a day, as a tonic for chronic constipation.

556. R. Extracti belladonnæ, gr. iss
 Pulveris rhei, gr. xxiv
 Pulveris zingiberis, gr. xij. M.
 For twenty-four pills. One four times a day.

DR. J. MORTIMER GRANVILLE, OF LONDON.

The following are recommended by Dr. J. MORTIMER GRANVILLE in the *British Medical Journal*, May 26, 1883:

When there is a lax and torpid condition of the muscular coat of the alimentary canal, evidenced by flatulence, etc., he uses the following:

557. R. Sodii valerianatis, gr. xxxvj
 Tincturæ nucis vomicæ, ℥lx
 Tincturæ capsici, ℥xlviij
 Syrupi aurantii, ℥ iss
 Aquâ ad., ℥vj. M.

Fiat mistura, cajúsumatur cochleare magnum ex aquâ ter die semihorâ ante cibum.

When there is a deficiency of glandular secretions, generally, throughout the intestine, manifested by a peculiarly dry and earthy character of the dejecta when the bowels *do* act, he gives;

558. R. Aluminis, ℥iij
 Tincturæ quassiæ, ℥j
 Infusi quassiæ, ℥ vij. M.

Fiat mistura, cujúsumantur cochlearia duo magna ter quotidie, post cibum.

The third form, which depends chiefly on interruption of the natural habit of periodic discharge, often results from repeated failure to move the bowels, in consequence of one or other of the two preceding forms of this trouble. This may generally be relieved by directing a perfectly regular attempt to go to stool, and by the use of the following

draught, taken the first thing after *rising* from bed—not on awaking—in the morning, as nearly as possible at the same hour. It will be observed that it is not an aperient in the ordinary sense of the term. It is, as a rule, neither necessary nor desirable to continue it for longer than a fortnight. In most instances, it will be found to re-establish the normal habit in a week or less.

559. R. Ammoniae carbonatis, 3j
 Tincturae valerianae, 3j
 Aquae camphorae, 3v. M.
 Fiat mistura, capiat partem sextam in modo dicto.

S. S. BIRCH, M. D., LONDON.

560. R. Extracti rhei alcoholici, 3ss
 Extracti taraxaci, gr. xxiv
 Quiniae sulphatis, gr. ij. M.
 Divide into twelve pills.

One should be taken either on rising in the morning or at dinner-time, or even at both periods, when the constipation is very obstinate. This is a very gentle stomachic and tonic evacuant, particularly useful for the delicate. In addition, when there is torpor of the liver, deficiency or perversion of the biliary secretion, the patient should be ordered:

561. R. Hydrargyri cum cretâ, gr. ¼j
 Sacchari albi, gr. v. M.
 For one powder.

This is a sufficient dose, when given alone, at bed-time, for two or three successive nights, or, in very sensitive persons, every second or third night. But the hydrargyrum cum cretâ is often prepared imperfectly, and then causes considerable annoyance and dissatisfaction to the practitioner.

PROF. WILLIAM THOMPSON, M. D., MEW YORK.

This writer observes (*Medical Record*, May, 1877,) that obstinate constipation is owing to deficient action of, 1, the small intestine; or, 2, the large intestine.

1. Deficient action of the small intestine is often owing to deficient secretion. Constipation from this cause is common in the Northern States, as a sequel to diarrhoea, and is often met with in the Southern States, as the result of malarial poisoning. There is, in these cases, no extraordinary accumulation and impaction, but a sluggish action of the

bowels, medicine being required to produce the stools in four or five days, which are, even then, moderate in amount and quite dry. In most cases, there is present a dull pain, or, rather, an uncomfortable sensation, at the back of the head, which is best relieved by a free discharge of bile. The tongue is not large and flabby, but usually small, with a little redness at the tip and along the edges. The secretions of the mouth are viscid, indicating the condition of defective secretion that prevails all along the canal.

Mild cathartics, in these cases, are injurious, and active purgatives still more so. What is wanted, is the presence, in the intestine, of a small increase of lubricating liquid, and this is obtained by causing the patient to take a great deal more water than customary—drinking on rising, for example, two tumblerfuls of water; and, as a rule, those who drink a considerable quantity of water are not constipated. Its laxative action may be insured by the addition of some saline, as carbonate of soda or salt, the water to which this is added not passing so readily through the mucous membrane into the general system, and thus more readily exciting peristaltic action. A half drachm of *sulphate magnesia*, dissolved in a pint of water, will, for this reason, act, sufficiently as a cathartic. A curious fact is, that the addition of a small dose of quinine to salines, increases their power of acting on the intestinal canal.

562. R.	<i>Magnesiae sulphatis,</i>	3j	
	<i>Quiniae sulphatis,</i>	gr. j.	M.
To be taken in a tumbler of water, every morning.			

This dose rarely fails to produce all the laxative effect required in every form of deficient secretion, as, for instance, in constipation following fever, where we wish to obtain free evacuations. Patients should be told not to expect much effect for a week or two; but, if they can be induced to persist in the daily use of large quantities of water, great benefit will almost always follow.

When flatulence is present, and defective innervation, the following pill is excellent:

563. R.	<i>Pulveris assafoetidae,</i>	gr. iv	
	<i>Extracti nucis vomicae,</i>	gr. ss	
	<i>Saponis,</i>	gr. ix.	M.
For three pills. To be taken daily.			

Such a combination is suitable in persons of a sedentary habit, with a tendency to headache and a dull, sallow color.

2. *Deficient Action of the Large Intestine.* This also may depend upon deficient secretion, or defective innervation; but far more frequently on the latter. One of the worst forms of this is when the constipation depends upon deficient nerve-power in the rectum alone, and which, if overlooked, may give rise to rectal abscess. The patients have little knowledge that they should have a movement, and when the sensation does occur, they have little or no power of expelling the fæcal accumulation.

One of the most common causes of this condition is chronic inflammation set up about hæmorrhoids. The relaxed condition which follows the defective innervation renders prolapsus of the rectum very probable. These patients are remarkable for being generally low-spirited, so that even insanity may be induced by such a state of the rectum, disappearing when this has been relieved. In treating these cases the first indication is to keep the rectum empty; and when accumulations do take place, they are best removed by enemata. These, however, should never be prescribed as a regular treatment, for if the habit is acquired of emptying the bowels daily by their aid they can never be dispensed with. When the accumulation which has caused the enema to be employed, has been thoroughly removed, other means should be restored to for restoring the innervation of the bowel; and in these cases the injection of *strychnia* into the submucous tissue is an exceedingly valuable specific. If necessary, a fold of the mucous membrane may be drawn down, and the injection inserted. It will frequently cure the worst forms of prolapsus of the rectum, as well as that condition in which there is simple debility with hypertrophy of the mucous membrane.

In other forms of constipation there are accumulations of scybalous matters in the upper part of the rectum and in the transverse colon. When these last are dislodged they come down into the sigmoid flexure; and mineral waters, such as the Kissingen, are useful in effecting such dislodgment, loosening the scybalæ without depressing the patient. The water may be taken in the morning; a suppository of *stramonium* or *belladonna* (sufficient to cause a little dryness of the throat and slight dilatation of the pupil) may be introduced in the evening. Constipation may be met with during the recovery from pneumonia and other

febrile affections, arising from accumulations at various parts of the canal. Compound jalap powder is a very effective means for preventing such accumulations.

THOMAS HAWKES TANNER, M. D., F. L. S., LONDON.

564. R. Zinci valerianatis, gr. xij-xxiv
 Extracti belladonnæ, gr. iij-vj
 Extracti gentianæ, gr. xxiv M.

Make twelve pills and silver them. One to be taken three times a day, in nervous cases of habitual constipation, and in spasmodic contraction of the sphincter ani.

PROF. W. H. VAN BUREN, M. D., NEW YORK.

565. R. Extracti aloës, 3 ss
 Extracti nucis vomicæ, gr. vj
 Extracti hyoscyami, ʒj
 Pulveris ipecacuanhæ, gr. j. M.

Divide into twenty pills. One to be taken at night.

This recipe, termed the "*Pil. Salutis*," is of special value in the constipation of females.

DR. METTAUER'S ALKALINE MIXTURE OF ALOES.

566. R. Aloes, 2½ ounces
 Bicarbonate of soda, 6 ounces
 Compound spirit of lavender, 2 fluid ounces
 Water, 4 pints. M.

Macerate for two weeks, and filter. Dose—from one fluid drachm to one fluid ounce half an hour after meals, for persons of a costive habit.

It has been observed that *aloes* is useful in habitual constipation due to atony of the colon, the result of fever and debilitating diseases, old age or sedentary habits. It may be combined as indicated, with aromatics, hyoscyamus, ipecacuanha, nux vomica or quinine. Dr. SPENDER recommends (*Medical Times and Gazette*, February 19, 1870,) the following pill:

567. R. Extracti aloes aq., gr. ss-j
 Ferri sulphatis, gr. jss-ij. M.

For one pill, to be taken thrice daily, immediately after the principal meals, and continued regularly for a few days till the occurrence of the first loose motion, after which one only is to be taken night and morning. At the end of a week or two, another pill may be omitted, and within a month a single pill once or twice a week suffices.

Dr. W. CRAIG, of Edinburgh, prefers *aloin*. He has never seen any bad effects follow its use; and, from its small dose, absence of

gripping, and certain action, he holds it should always be used in preference to the crude drug. He gives :

568. R. Aloinæ, gr. j $\frac{1}{4}$
 Ferri sulphat. exsic., gr. jss
 Quiniæ sulphatis, gr. j
 Extr. nucis vomicæ, gr. ss
 Extr. gentianæ, q. s. M.
 For one pill.

DR. F. PEYRE PORCHER.

569. R. Extracti hyoscyami, gr. j
 Podophyllin, gr. $\frac{1}{4}$ $\frac{1}{6}$
 Ferri sulphatis, gr. ij
 Extracti gentianæ, q. s. M.
 For one pill, to be used as a mild laxative in torpid bowels. One at night, to be repeated at intervals as required.

PROF. ROBLEY DUNGLISON.

570. R. Magnesii sulphatis, $\overline{3j}$
 Potassii bitartratis, $\overline{3j}$
 Ferri sulphatis, gr. x. M.
 For one powder. Add to a quart of water and take a wineglassful, on rising every morning.

The recipe was frequently recommended in his lectures by the late distinguished professor of physiology.

JOHN FORSYTH MEIGS, M. D.

571. R. Confectionis sennæ, $\overline{3j}$
 Potassii bitartratis, $\overline{3ij}$
 Sulphuris præcipitati, $\overline{3j}$
 Ferri subcarbonatis, āā q. s. M.
 Mellis despumati,
 Make an electuary. Teaspoonful after meals.

PROF. METCALF, M. D., NEW YORK.

572. R. Extracti aloës (purif.), āā $\overline{3j}$
 Extracti hyoscyami, gr. xij
 Extracti nucis vomicæ, gtt. x. M.
 Olei anisi,
 Divide into sixty pills. One to be taken after each meal, particularly for constipation in women.

HYPODERMIC ADMINISTRATION OF CATHARTICS.

DR. N. HILLER.

This author in *Zeitschrift für Klin. Med.*, iv. 40, has verified and repeated a number of experiments that go to prove that cathartics can

be administered subcutaneously, and bring about catharsis. He has noticed that from $\frac{1}{4}$ to 1 grain of aloin will produce a copious watery evacuation in from 4 to 6 hours. He uses an alcoholic watery solution of glycerine as the menstruum.

Elaterium, extracts as well as pure elaterine, are partially inefficient, and some are not applicable for other reasons.

Leptandrin, a glucoside insoluble in water, from *leptandra Virginiana*, caused an increased peristalsis when injected in doses of one grain without causing diarrhœa.

FOR NERVOUS CONSTIPATION.

573. R.	Pulv. ferri sulphatis,	gr. $1\frac{1}{2}$	
	Aloes socotrinæ,	gr. $\frac{3}{4}$	
	Ext. Belladonnæ,	gr. $\frac{1}{15}-\frac{1}{6}$	M.
Ft. Pil. One to four pills daily.			

NOTES ON REMEDIES.

Aleuritis Triloba. The oil of the aleuritis nut has been highly recommended as a substitute for castor oil. It has the further advantage of not being disagreeable to take, having the flavor of the hazel-nut. An adult is easily purged with a dose of fifteen grammes. It is an excellent aperient, and its effects on the intestine are identical with those of castor oil. About three hours are required for its action, which takes place without pains or colic.

574. R.	Olei aleuritis,	f. $\frac{3}{5}$ ss	
	Sacchari albi,	$\frac{3}{5}$ ss	
	Pulv. acaciæ,		
	Aquæ,	aa	$\frac{3}{4}$ iij M.
For one dose.			

In rebellious cases of constipation and in abdominal pains, good results have been obtained by making frictions on the abdomen with the following liniment :

575. R.	Olei aleuritis,	f. $\frac{3}{5}$ ss	
	Tincturæ cantharidis,	f. $\frac{3}{5}$ iij	
	Ammoniac carbonatis,	$\frac{3}{5}$ iij.	M.
For a liniment.			

Aloes and Aloin. The following formula is highly recommended.

576. R.	Ext. aloes soc. aquos.,	gr. ij	
	“ Nucis vomicæ,	gr. ss	
	Ipecac pulv.,	gr. ss.	M.
Ft. Pil. To be taken each day with dinner.			

Arseniosum Acidum is said by Dr. ISNARD, of Marseilles, to be particularly useful in the constipation of aged people and sedentary persons.

**Belladonna* is especially called for where there is atony or paralysis of the muscular coat of the intestines; gtt. v-x, thrice daily, of the tincture, will often overcome the most obstinate constipation from this cause.

Benzoini Tinctura, in doses of gtt. xx, thrice daily, is said, in cases of constipation apparently depending upon stricture of the colon, to keep up the peristaltic actions without irritating the bowels.

Calcis Saccharas. Dr. J. CLELAND, of Edinburgh, first introduced this aperient through the *Edinburgh Medical Journal*, August, 1859, to the profession. He prepares it as follows:

577. R.	Slake of quicklime,	$\overline{3}$ viij	
	Rub up with it of white sugar	$\overline{3}$ v	
	Add of water,	f. $\overline{3}$ xx.	M.

Stir and filter. The product should be perfectly clear, with only a slightly yellowish tinge. Each ounce, by weight, should contain 18 grains of lime. Dose, \mathfrak{m} xv-xxx, or f. $\overline{3}$ j.

Dr. C. finds this saccharate of lime a very valuable means of overcoming gradually that chronic constipation which so frequently accompanies dyspepsia, and that persons who for years have been in the constant habit of using aperient medicines have been able to abandon them, in a great measure, after taking this remedy for some time.

Camphora. In constipated persons, who suffer much from *difficult defecation*, the following enema is very highly recommended: Take a tumbler and fill it half full of water, at the temperature of the room; pour in a few drops of tincture of camphor—just enough to give the water a slight sapidity—then fill the glass with water. Inject this slowly into the rectum, till about sixty or eighty grammes have been introduced. At first, no effect is perceived, but in about ten minutes, the desire to defecate becomes irresistible. The effect becomes energetic in proportion to the quantity of tincture of camphor added. After the defecation, it is well to repeat the injection of a small quantity of the same mixture, and retain it in the rectum, which can readily be done, so as to prevent constipation on the following day.

Cascara Sagrada, otherwise known as the *Rhamnus Purshiana*, is a well-established, generally efficient remedy in habitual constipation, especially when dependent on deficient biliary secretion. Of the fluid extract, f. $\overline{3}$ ss daily is a dose; of the cordial $\overline{3}$ ss-j. Both are prepared by the house of Parke, Davis & Co., Detroit, who have introduced the drug to the profession.

Colchicum, in doses of gtt. v of the tincture of the root several times daily, has been highly recommended in obstinate constipation.

Colocynth. Dr. CRICHTON recommends (*British Medical Journal*, November 28th, 1868,) the Prussian tincture of colocynth, in doses of \mathfrak{m} v-x in a little water, taken about an hour before breakfast, as an efficient remedy. The formula of the Prussian tincture is as follows:

578. R.	Colocynthidis pulpæ,	$\overline{3}$ j	
	Illicii anisati,	$\overline{3}$ j	
	Alcohol,	f. $\overline{3}$ xx.	M.

Dr. BIRCH recommends colocynth combined with hyoscyamus, in the constipation of old age.

Fel Bovinum Purificatum. When there is torpidity of the liver and deficiency of the biliary secretion, ox-gall may prove useful by supplying the deficiency. Either of the following formulæ may be used :

579.	R.	Fel bovini purificati,	℥ ij	
		Olei carui,	℥ x	
		Magnesiæ carbonatis,	q. s.	M.

Make thirty-six pills. Take two twice daily.

Or :

580.	R.	Fel bovini purificati,		
		Pilulæ assafœtidæ compositæ,	āā	gr. xxx
		Extracti aloes,		gr. xx
		Saponis duris,		gr. x
		Pulveris ipecacuanhæ,		gr. viij
				M.

Make thirty pills. Take one to two daily, before dinner.

Ferri Carbonas has not, in the opinion of Dr. GRAVES, been duly appreciated as a remedy in habitual constipation. He has succeeded in curing with it alone, a patient who had long been subject to extreme constipation, and who had been reduced to the necessity of taking an immense dose of purgatives almost every week.

Magnesiæ Sulphas, in small, frequently-repeated doses, with the addition of a few drops of diluted sulphuric acid, will sometimes subdue obstinate constipation after the failure of other means.

Muscarine. BARTHOLOW recommends the following :

581.	R.	Muscarinæ,		gr. iv
		Ext. belladonnæ,		
		Ext. nucis vomicæ,	āā	gr. iij
		Euonymin,		℥ ss.
				M.

Ft. Pil. No. xii.

S.—One pill morning and evening.

Muscarine is strongly indicated when the digestion of starches and fats is imperfect.

Nitricum Acidum, in the opinion of Dr. GRAVES, is a very reliable remedy in habitual constipation, and combines tonic with aperient qualities.

Pitzahoic acid. This acid, obtained from the root of the pipitzahuat, a Mexican plant, is stated to be an excellent and safe purgative. The remedy was first introduced in the *Annalen der Chemie*, 1855, and in 1883 into Germany more prominently.

Pix Liquida, persevered in for some time, often effectually removes constipation.

Physostigma. BARTHOLOW recommends the following :

582.	R.	Tinct. physostigmatis,		
		Tinct. nucis vomicæ,		
		Tinct. belladonna,	āā	℥ ij.
				M.

S.—Thirty drops in water, morning and evening.

Or,

583. R. Extract physostigmatis,
Extract belladonnæ,
Extract nucis vomicæ, āā gr. v. M.

Ft. Pil. No. x.

S.—One pill at bed-hour.

Or,

584. R. Ext. physostigmatis,
Resinæ podophylli, āā gr. iij. M.

Ft. Pil. No. vi.

S.—One pill at bed-hour.

Podophyllin is useful when there is torpor of the upper portion of the bowel. It may be continued indefinitely, in small doses. Dr. HORACE DOBELL recommends the following formula as one of the most satisfactory he has ever tried :

585. R. Podophylli, gr. ij
Essentiæ zingib., ʒ ij
Alcoholis, ʒ ij. M.

A teaspoonful at bed-time, in a wineglass of water, nightly, or every second or third night.

Rhamnus Frangula and *Rhamnus Purshiana* (*Cascara Sagrada*) are valuable as laxatives in constipation from dyspepsia.

Rheum. Dr. F. W. PAVY recommends the following pill in habitual constipation :

586. R. Pilulæ rhei comp., gr. ij-ijj
Pulveris capsici, gr. j. M.

To be taken *with the food*, and repeated daily, as occasion requires.

In the constipation of anæmic females, Dr. ASHWELL recommends the following mixture :

587. R. Pulveris rhei,
Magnesii carbonatis, āā ʒ ss
Confectionis aromaticæ, gr. xx
Aquæ cinnamomi, f. ʒ iij
Tincturæ cardamomi compositæ, f. ʒ j. M.

Make a drink. To be taken at bed-time, every three or four days.

Ricini Oleum is one of the best purgatives which can be employed in habitual constipation, acting speedily, without much griping, and causing less subsequent constipation than any other purgative given to remedy this state.

Senna, particularly in the form of confection, is a mild, efficient purgative, causing little excitement, and leaving no subsequent constipation.

Stramonium. Prof. ARMOR (*American Practitioner*) has been in the habit of using stramonium in the form of suppository, and is pleased with the results. Half or three parts of a grain of the extract may be combined

with a sufficient quantity of cocoa butter. It is admirably adapted to obstinate constipation of nervous females, who suffer, at the same time, from pelvic irritations from various causes.

Tabaci Vinum. In habitual constipation, due to a relaxed state of the muscular layer of the bowel, five minims of the wine of tobacco, administered at bed time will, not infrequently, afford relief.

**Nux Vomica* or *Strychnia* is often of equal benefit. Dr. COPLAND recommends the following pills :

588.	R.	Extracti nucis vomicæ,	gr. iij	
		Extracti colocynthis compositi,	℥ ij	
		Saponis castiliensis,	gr. xij.	M.

Make twelve pills. Take one every night, for a few weeks.

It is also indicated in those cases in which there is inaction of the muscular layer, and consequently great fecal accumulations. It may be most advantageously combined with purgatives. BARTHOLOW recommends the following :

589.	R.	Tinct. aloes et myrrhæ,	℥ vi	
		Tinct. nucis vomicæ,	℥ ij.	M.

S.—Fifteen to thirty drops two or three times a day.

Veratrum Viride, in doses of gtt. iij of the tincture four or five times daily, is said (*Practitioner*, November, 1868,) to be useful in some cases of obstinate constipation.

Electricity. Dr. ALTHAUS and Drs. BEARD and ROCKWELL have reported a number of cases of habitual constipation which yielded readily to faradization of the bowels after the failure of other remedies.

DIARRHŒA, ACUTE AND CHRONIC.

As the term diarrhœa is applied to a condition characterized by fluid fæcal discharges accompanied with intestinal pains, its treatment must always have reference to the exciting cause of the discharge. Frequently it is not advisable to check it, as it is a natural effort to carry off irritating matter from the intestinal canal. Indeed, it is often better to assist this effort by gentle laxatives and judicious diet, as for instance on the plan recommended by

DR. ROBERTS BARTHOLOW, PHILADELPHIA.

The diarrhœa and dysentery of infants (ileo-colitis) is frequently treated with minute doses of calomel or hydrargyrum cum creta. When

there are much straining and bloody mucus, it is said that small doses of corrosive chloride prove very effective, but the author is convinced that mercurials are much abused in these affections. Children are quickly poisoned by mercurials, though they are not easily salivated. The spinach-colored stools which so frequently occur in the summer complaint of children, and which are, by ignorant practitioners, supposed to be produced by the mercury administered, really belong to cases of ileo-colitis, and may, by their persistence and profuseness, signify an increased irritation of the intestinal mucous membrane, due to the remedies given. While the author believes that other remedies are more useful in the ileo-colitis of children, he is convinced of the utility of minute doses of calomel ($\frac{1}{20}$ to $\frac{1}{12}$ of a grain every half hour), when there is much irritability of the stomach. Mercurials are contraindicated in the diarrhœa and dysentery of adults, as a rule. Sulphuric acid is more decidedly astringent than nitric and muriatic, and is, therefore, as a rule to be preferred in diarrhœa.

BENCE JONES places them, as regards their actions, thus: Hydrochloric more promotes digestion; nitric acid, secretion; and sulphuric, astringency. Nitric and nitro-hydrochloric, are, according to this view, better suited to stomach and hepatic disorders, characterized by deficient secretion; and sulphuric is more appropriate for the relief of a relaxed state of the mucous membrane. A combination of aromatic sulphuric acid with opium is one of the most effective remedies we possess in the treatment of summer diarrhœa and cholera. When the mineral acids do not quickly improve the discharges and lessen their frequency, and when they increase the tormina and tenesmus, they should be suspended.

The *grape-cure*, which is a treatment much in vogue in some parts of France and Germany, consists, according to CARRIÈRE, of a diet exclusively of grapes. They are taken many times a day, to repletion. It is usual to commence with a pound, and progressively to increase the amount to two, three, six and eight pounds, a limit which is not exceeded. The first grape repast, which may be the most abundant, is in the early morning, but not, as are the others, eaten in the vineyard. Another is taken at the time of the morning meal (corresponding to our breakfast); the next after the morning walk, at noon, consisting of bread and water (because there is not a requisite amount of nitrogenous matter in grapes); another before the usual dinner hour

(evening), and finally, before retiring. The treatment is continued during the five or six weeks of the duration of the grape crop.

A method of treating diarrhoea, long practiced in Russia, consists in the use of a pulp of raw meat. A bit of fillet of beef is deprived of all fat and aponeurotic fibre, minutely divided, and beaten in a mortar until all trace of fibre has disappeared. It is then pressed through a fine seive and mixed with sugar, conserve of roses, or some suitable aromatic, or seasoned with salt and pepper to the taste. It may be administered in this form, with fruit jelly, or spread on thin pieces of bread. A beefsteak, hastily broiled on a hot fire, so as to retain its juices, may be treated by the same method; or the raw beef scraped to a pulp, rejecting the fibre, may be thrown on to a hot skillet for a few seconds, to give an odor and appearance of cooked meat. This method, which has been used especially in the treatment of diarrhoeal diseases of early life, is equally efficacious in the chronic diarrhoea of adult life. The chief objection to this mode of alimentation is the great frequency with which tape-worm follows.

A milk diet has proved very efficacious.

DAVID YOUNG, M. D., FLORENCE, ITALY.

This writer (*Practitioner*, March, 1875; December, 1876,) states that in nearly every form of diarrhoea he trusts almost exclusively to diet and one or two forms of castor-oil emulsion. For instance:

590. R.	Olei ricini,	℥xxiv	
	Spt. chloroformi,	℥jss	
	Sol. morph. mur.,	℥j	
	Pulv. gum acaciæ,	℥ijss	
	Syrupi,	℥ss	
	Aquam,	ad ℥iv.	M.

A small dessertspoonful every hour and a half till the bowels are quieted.

He adds the following rules:

1. When the diarrhoea is chronic, and the stools contain mucus, he increases the dose of castor-oil to from four to six drops.

2. If the pain is very severe, six drops of morphia (Sol. B. Ph.) may be given with each dose, but he has never had occasion to give more.

3. If the mixture is carefully prepared it is pleasant and readily taken, and the taste of the oil is so completely covered that in only two or three cases of the large number in which he had given it, was the mixture suspected to taste like castor oil.

4. The mixture does not keep well, especially in warm weather, but the addition of four grains of quinine to a three-ounce bottle will keep it fresh for several weeks.

In muco-enteritis, or dysenteric diarrhœa, Dr. YOUNG has derived decided advantage from the employment of *arsenic*. It is especially indicated where the stools are slimy, contain much mucus and blood, and are attended with great prostration of strength. Two drops of the *liquor arsenicalis* (B. Ph.) are given every two or three hours. If the movements occur soon after food, the arsenic should be taken before meals; if not, in the usual way, after food.

DR. THOMAS HILLIER, LONDON.

591. R. Olei ricini, f. ℥ ij
 Pulveris acaciæ, ℥ j
 Tincturæ opii, ℥ viij
 Syrupi, f. ℥ ij
 Aquæ carui, q. s. ad f. ℥ ij. M.

Dose—A tablespoonful for a child six years old.

A useful oleaginous mixture in dysenteric diarrhœa.

W. KEMPSTER, M. D., UTICA, N. Y.

592. R. Acidi carbolicæ, gr. j
 Aquæ, f. ℥ j. M.

This is the strength of the standard house solution in the State Lunatic Asylum at Utica, New York. Of this a dessertspoonful is given thrice daily, in case of sluggishness of the bowels accompanied by offensive breath. Diarrhœa produced by eating unripe fruit, or other articles which promote fermentation, is relieved by combining a drachm or two of the solution with the usual remedies. When a foetid smell emanates from the cutaneous surface, order a warm bath, and then wash the surface with a solution, gr. v-f. ℥ j.

Poultices and hot fomentations help to relieve the pain.

Mr. R. P. WHITE, of Dublin, speaks in the highest terms of a *turpentine* emulsion in all cases of severe diarrhœa, even those in typhoid fever.

593. R. Olei terebinthinæ, āā f. ℥ ij
 Liquoris potassæ, f. ℥ iv
 Mucilaginis acaciæ, āā f. ℥ viij
 Syrupi papaveris, āā ad f. ℥ viij. M.
 Syr. flor. aurantii,
 Aquam camphoræ,

Shake well, and take a tablespoonful every four hours.

After the acute stage of the attack is over, the time has come for opiates and astringents.

DR. WILLIAM AITKEN, M. D., EDINBURGH,

says that one general rule may be acted on in the cure of diarrhœa, which is, that in the adult, whatever be the form of the diarrhœa, if the stools be dark at first and then become light-colored, purgative medicines are no longer beneficial, and in no instance ought they to be continued longer than is sufficient to remove any irritative substance accumulated in the alimentary canal. Often a tonic is required:

594. R. Salicin, gr. v-x.
For one powder. To be taken every four or six hours.

In cases of diarrhœa with clear tongue, which will not yield to opiates, astringents, or stimulants, either singly or combined, and which probably, depend on a want of tone in the intestine, the above recipe has often stopped a discharge that appeared fast hurrying the patient to his grave.

DR. J. M. DA COSTA, M. D., PHILADELPHIA.

In cases in which there exists persistent irritability of the bowels, influenced by the taking of much food which cannot be digested, and in which there are gastric symptoms in connection with the dysenteric affection, the *subnitrate of bismuth* will be found very serviceable. In order that it shall produce an effect, it is necessary that it shall be administered in sufficiently large doses, not less than 20 grains. The dose may gradually be increased to a drachm.

595. R. Bismuthi subnitratis, ʒj
Acidi tannici,
Pulveris ipecacuanhæ compositæ, āā gr. iij. M.
For one powder. To be taken three times a day, in chronic dysenteric diarrhœa.

SIR J. FAYRER, F. R. C. P., BRITISH INDIA.

This author believes that in the treatment of chronic diarrhœa, *diet* is the most important element—more so than drugs. He gives the following rules:

All irritating or indigestible or solid food must be at first prohibited, and only that which is most easily absorbed and assimilated allowed. Milk alone, or, better, diluted with about one-fourth or one-third part of lime-water, given in small quantities and at frequent intervals, say a wineglassful every second or third hour, in some cases more fre-

quently, will generally be found to answer well, and may be continued for a time to the exclusion of all other food. Milk, undiluted, will seldom agree, as may be seen by its causing increased irritation, frequency of action, and the passage of undigested caseine. Beef tea, raw beef juice, or other plain animal broth, free from all extraneous matters; a raw egg beaten up with milk, to which a teaspoonful of brandy may be added, will sometimes be tolerated; arrowroot, tapioca, or other plain farinaceous food, will sometimes, but not always answer.

Tea and coffee, as a general rule, disagree, and should be avoided. Stimulants, especially for those who have long been habituated to their use, are often needed; the best are a little whisky or brandy diluted with Vals or Vichy or potash water; but these should be laid aside for plain water if they, as they sometimes do, increase the action of the bowels. Or a little very good port wine may be tried; as a general rule, all other wines are unsuitable.

Regularity in the times of administration and in the quantity of nourishment given is most essential. The greatest care should be taken not to give too much of anything at a time, and at once to discontinue whatever appears to disagree.

It is necessary that the patient should be kept warm, and at an equable temperature day and night. The body should be covered with flannel or woolen next the skin, and a flannel bandage should surround the abdomen. Chills and damp are especially to be avoided; exposure to them may seriously aggravate the mischief; during cold weather the patient should not leave the house.

JOHN FORSYTH MEIGS, M. D.

596. R.	Pulveris opii,	gr. vj	
	Extracti nucis vomicæ,	gr. iij	
	Cupri sulphatis,	gr. j.	M.

Divide into twelve pills. One three times a day, in chronic diarrhœa.

The value of this pill has been proved by army surgical experience.

CHARLES MURCHISON, M. D., F. R. C. S., ETC.

597. R.	Acidi tannici,	℥iv	
	Tincturæ opii,	℥xl	
	Glycerinæ,	f. ̄ss	
	Aquæ menthæ piperitæ,	ad f. ̄iv.	M.

A tablespoonful in a tablespoonful of water every four hours.

After the diarrhœa is checked, order:

598. R. Acidi nitro-muriatici, f. ʒi ss
 Tincturæ opii, m̄xl
 Syrupi, f. ʒss
 Aquæ, ad f. ʒiv. M.

A tablespoonful in water, four times a day.

DRS. BURKART AND RICKER, STUTTGART, GERMANY.

These writers have lately called attention to what they deem the extraordinary powers of *coto bark*, and its active principle, *cotoïn*, in diarrhoea, intestinal catarrh, and dysenteric disease. The remedy was employed in the form of powder, tincture, and the active principle, *cotoïn*. The taste of the powdered bark and tincture is particularly disagreeable, increasing the flow of the saliva in a marked degree. In moderate doses, they generally produce a burning sensation in the stomach, and, very frequently, eructations and vomiting, making both these forms of exhibition extremely undesirable. The employment of *cotoïn* is free from these objections, for very small doses produce the desired effect, without inducing secondary disturbances or disagreeable sensations of any kind.

599. R. Cotoïnæ, gr. j
 Aquæ destillatæ, f. ʒiv
 Alcoholis, gtt. x.
 Syrupi, f. ʒj. M.

A tablespoonful every hour.

Some of the cases were of long standing, some were severe attacks of cholera morbus, and others had been but little benefited by tannin, opium, or lead acetate. The above mixture produced speedy improvement—generally in a few hours—and complete recovery in from twelve hours to six days.

DR. EDWARD R. SQUIBB, BROOKLYN.

The following is a favorite prescription of this physician, in diarrhoea:

600. R. Tincturæ opii, āā f. ʒj
 Tincturæ camphoræ, f. ʒiij
 Tincturæ capsici, ad f. ʒv. M.
 Chloroformi purificati,
 Alcoholis,

A teaspoonful as required.

NOTES ON REMEDIES.

ACUTE DIARRHŒA.

Acetum. Pure cider vinegar has been recommended by Dr. T. C. STELWAGEN, of Philadelphia, as a safe and efficacious remedy in all forms of diarrhoea. It should be taken without water, an adult dose being about

two oz. It can also be given to children in chronic forms of the disease with advantage (*Reporter*, 1884).

Acidum Nitrosi. When there are painless, watery stools, of a light color, and alkaline in reaction, Hope's mixture has long been used with advantage. The following is the formula :

601. R.	Acid. nitrosi,	℥j
	Tinct. opii,	gtt. xl
	Aquæ camphoræ,	℥ viij.

S.—One-fourth to be taken every three or four hours.

Camphor is regarded by Dr. RINGER as one of the most efficacious of remedies in summer diarrhœa, but it must be employed at the very commencement of the disease, or it will fail of its effect.

Capsicum is regarded by Dr. COPLAND as almost a specific in diarrhœa arising from putrid matter in the intestines, and especially when occasioned by fish.

Carbolicum Acidum has been recommended in the diarrhœa caused by eating unripe fruit, etc. (F. 592.)

Deuberry Root, (*Rubus Procumbens*) in teaspoonful doses of the fluid extract, has been found very useful by Dr. JOHN S. LYNCH, (*Virginia Medical Monthly*, January, 1884.)

Ipecacuanha is often of benefit after the other remedies have proved useless.

Leptandrin is valuable when the stools are destitute of bile and the mucous membrane is irritable. In such cases, the stools, under its use, are said to assume their natural color and consistence.

Magnesii Sulphas. A dose at the outset often brings prompt relief.

Oleum Ricini. (F. 590, 591.)

**Opium* is a most valuable remedy in diarrhœa. In cases due to the ingestion of indigestible food, etc., an aperient should be first given, to remove the irritating matter ; then, if the diarrhœa persist, opium, in mixture, or pill, or enema, will often arrest it. Dr. WARING recommends, in the so-called summer or autumnal diarrhœa, the combination of laudanum (℥x-xx) with dilute sulphuric acid (℥xv-xx), as often successful.

**Rheum* is often of great benefit, by its property of first removing irritant matters from the bowels, and then acting as an astringent and tonic. The following is an excellent formula for general use, which by itself, will often effect a cure:

602. R.	Pulveris rhei compositi,	℥j
	Sodii carbonatis,	gr. xx
	Tincturæ opii,	℥x-xv
	Aquæ menthæ piperitæ,	f. ℥x.

Make a drink. If, after its operation, the diarrhœa continues, chalk mixture and opium may be given.

Sodii Bicarbonas, in full dose (℥ij-℥j), combined with tincture of ginger (f. ℥ss-f. ℥j) and aromatic spirits of ammonia (f. ℥ss-f. ℥j) and a little laudanum, is of great service in diarrhœa attended with colic and gas-

tric irritability and flatulence, especially when this is consequent on the ingestion of unwholesome food.

Tannic Acid and vegetable astringents containing it, especially rubus, kino, and catechu, are much used in simple diarrhœa.

Emetics. An ipecacuanha emetic, repeated once or twice, according to circumstances, often arrests a diarrhœal discharge.

Enemata are often of great service, either mucilaginous or containing opium, ipecacuanha, or astringents. *Cold water enemata* are highly recommended by WINTERITZ.

SUB-ACUTE AND CHRONIC DIARRHŒA.

Alum, in doses of \mathfrak{z} ss- \mathfrak{z} j, daily, or in enema (\mathfrak{z} j-vj), is often of benefit in diarrhœa dependent upon a relaxed condition of the intestinal mucous membrane. The following formula is useful :

603.	R.	Aluminis,	gr. x	
		Pulveris kino compositi,	gr. v	
		Confectionis rosæ,	q. s.	M.

Make pill. Take every sixth hour.

Argenti Nitras is of service in chronic diarrhœa when the stools are watery or mucous.

Arsenic is highly spoken of by Dr. RINGER in chronic diarrhœa. He gives one to two minims of Fowler's solution before each meal. BARTHOLOW recommends it, with tincture of opium, when undigested food is passed soon after meals, and in chronic diarrhœa.

Bael. A teaspoonful of the liquid extract, twice daily, is recommended by Dr. J. K. SPENDER, in painless chronic diarrhœa. FAYRER and MURCHISON also praise it highly.

**Bismuthi Subnitratis*, (See F. 595) Professor ALONZO CLARKE recommends as particularly excellent in chronic diarrhœa:

604.	R.	Bismuthi subnitratis,	gr. v	
		Morphiæ sulphatis,	gr. $\frac{1}{12}$.	M.

This much, two, three or more times daily.

Calcis Carbolas, in one-grain doses, combined with henbane, has been successfully employed by Dr. HABERSHON (*Lancet*, January 4th, 1868,) in diarrhœa attended with flatulent distention of the colon.

Calcis Phosphas, in doses of gr. j-ij, several times a day, is praised by Dr. RINGER as of great value in chronic diarrhœa, effecting both a general and local improvement.

Carbazotate of Ammonia, in grain doses, thrice daily, has proved successful in very chronic cases.

Cascarilla is much used in Germany in the advanced stages of diarrhœa.

**Catechu*, when there is no inflammatory action, is useful (in doses of gr. x-xv of the compound powder, or f. \mathfrak{z} j-ij of the infusion) in diarrhœa de-

water is added to it, till the whole amounts to thirty ounces. The liquid ought to be transparent and of a beautiful dark-red color.

Ferri et Potassii Tartras. The following formula is recommended by Dr. E. GOODEVE, in chronic diarrhœa attended with anæmia :

609. R.	Ferri et potassii tartratis, Tincturæ opii, Aquæ cinnamomi,	℥j-ij f. ℥ iss-iiij f. ℥iv.	M.
---------	---	-----------------------------------	----

A tablespoonful in water thrice daily.

Galla, in doses of gr. x-xx of the powder, several times a day, is occasionally useful in chronic diarrhœa. Dr. J. K. SPENDER recommends ℥j of the tincture thrice daily in painless diarrhœa.

Granati Fructus Cortex, in decoction, is recommended by Dr. WARING in chronic diarrhœa unattended by inflammatory action. It is particularly useful in diarrhœa consequent on debilitating diseases.

Hæmatoxyllum, in decoction, in the form of an extract (gr. x-xxx,) is a useful astringent and tonic after active inflammation has subsided.

Hydrargyri Chloridum Corrosivum. In intestinal catarrh. especially in children, the use of the following prescription, with a milk diet, often produces "marvelous" results :

610. R.	Hydrarg. chlor. corrosivi, Aquæ,	gr. $\frac{1}{8}$ - $\frac{1}{2}$ f. ℥iv.	M.
---------	-------------------------------------	--	----

A teaspoonful every two or three hours.

* *Kino*, in doses of gr. xij-xx, with or without opium, is a valuable astringent in chronic diarrhœa, especially that connected with follicular derangement. It is also frequently prescribed even when febrile symptoms are present.

Krameria is much employed by Spanish physicians, in chronic diarrhœa, and is especially useful when the stools are mucous and slimy, and when there is no inflammatory action.

Matico, in infusion, is usually of benefit in chronic diarrhœa.

Muriatic Acid, combined with tincture of nux vomica, and given before meals, is found of benefit, by Dr. WILLIAM FOX, in the nervous disorder of digestion, associated with diarrhœa.

Myristica, in mild cases of asthenic diarrhœa, is recommended by Dr. PEREIRA as a substitute for opium, with warm brandy and water (if not contra-indicated) as a vehicle.

Nitricum Acidum Dilutum, in doses of ℥v-xv, combined with laudanum (℥xv-xx), in a little vegetable infusion, has been recommended in chronic diarrhœa.

Nux Vomica is highly praised by Prof. GRAVES, in obstinate chronic diarrhœa, which resists ordinary remedies, and in that form, especially, which accompanies exhaustion after fevers. It may be combined with opium and sulphate of copper.

* *Opium* is frequently given, alone or in combination with other remedies. Dr. SPENDER says it is now and then absolutely necessary, and it should always be prescribed in comparatively small and frequent doses, so as to obtain the least physiological with the most medicinal effect. Let the wine of opium be given to an adult, in the quantity of three or four minims (with an ounce of chloroform water), five or six times in the twenty-four hours; and the remedy ought, invariably, to be left off by degrees. In ordinary *choleraic diarrhœa* or *cholera morbus* of adults, no remedy will so promptly relieve the cramps and pain, and check the purging, as the hypodermic administration of *morphiæ sulphas*, gr. ss.

Plumbi Acetas, by the mouth or in the form of an enema, is often an effectual remedy.

Potassii Chloras is recommended by Dr. COPELAND in chronic mucous diarrhœa, with whitish, grayish, or mucilage-like stools, arising from the absence of bile, etc. Professor BONFIGLI lately urges it in vaso-paralytic diarrhœas. In grave cases of cachexia, connected with great nervous depression, the diarrhœa diminishes, but does not completely cease. In obstinate cases, there are alterations of the vascular parietes (amyloid or fatty degeneration), or ulcerative lesions of the mucous membrane. These require an energetic and prolonged use of the remedy. It is of no use if the diarrhœa is kept up by an active process of the mucous membrane, catarrhal enteritis, etc.

Quercus Cortex, in decoction, is sometimes serviceable. Acorns, roasted and powdered, have also been used.

Salicin, in cases of chronic diarrhœa dependent on want of tone, is recommended by Dr. AITKEN. It should be administered in a dose of five or six grains, perhaps combined with a grain of ipecacuanha. Let them be mixed into a couple of pills, and taken three or four times a day. This plan seldom fails to appease an obstinate diarrhœa.

Sarracenia Flava, is a popular agent in chronic diarrhœa. One physician reports uniform success with it. (*Therapeutic Gazette*, 1880.)

Tannin, with opium, sometimes proves of surprising efficacy in cases of chronic diarrhœa which have resisted all ordinary treatment, and which are not dependent on obstructive disease of the heart or liver. It is especially useful where there is an irritable, weakly mucous membrane. It may be given in the form of a pill or mixture.

Terebinthinæ Oleum in Stupes, applied to the whole abdomen and allowed to remain on as long as the patient will endure them, is often of great benefit. Internally, see F. 593.

Zinci Oxidum, in doses of gr. iij-v, thrice daily, has been recommended by Dr. WARING-CURRAN. Dr. BONAMY, of Paris, has employed the method indicated by M. GUBLER, that is to say, 20 grains of oxide of zinc, mixed with $7\frac{1}{2}$ grains of bicarbonate of soda, and divided into three or four doses, to be taken every three hours. M. BONAMY has not observed any vomiting. He sums up the advantages of the treatment thus: 1st. *Superiority of action*; since all the cases of diarrhœa in which

oxide of zinc was employed with success, had resisted other means of treatment; and since the diarrhœa re-appeared as soon as the oxide of zinc was replaced by another agent. 2d. *Rapidity of action*; diarrhœa dating for several months was favorably modified after the administration of the first dose.

Zinci Sulphas, combined with ipecacuanha and opium, occasionally proves of advantage.

Blisters. A small blister to the abdomen is often very effectual in obstinate sub-acute diarrhœa.

Enemata of cold water have recently been found very useful by Dr. M. J. B. MESSLER. (*American Journal Medical Science*, July, 1878.)

DYSENTERY, ACUTE AND CHRONIC.

ACUTE DYSENTERY.

AUSTIN FLINT, M. D., NEW YORK.

In *sporadic* dysentery, if we have reason to conclude that fæces are retained in the large intestine, it is a rational procedure to first take steps to secure a complete evacuation of such an accumulation. Castor oil is a remedy which has long been employed, and is one well suited to meet the indication in these cases. Salines have been used for the same purpose, and perhaps are to be preferred, because they are far more easily taken, and afford more relief. The next indication is to keep the inflamed intestine perfectly quiet, which is best accomplished by the use of *opium*. Opium may be administered by the mouth, by the rectum, or hypodermically. Perhaps we succeed better in securing quietude of the large intestine by the use of anodyne enemata than by the use of opium, either by the mouth or hypodermically—the effect upon the large intestine seems to be more direct. After a day or two it may be found that the alimentary canal is again more or less loaded, and, if deemed necessary, the salines or the oil can be repeated. A bland diet is necessary, and cold water enemata for the tenesmus.

On the other hand, in *epidemic* dysentery, purgatives are to be avoided. Salines, which operate by producing a more or less abundant watery transudation, are contra-indicated. So far as medical treatment is concerned, our chief reliance must be placed upon opium. Administer opium early and persistently, and to the extent of absolutely quiet-

ing the intestines, but at the same time avoiding the risk of narcotism. It is a noteworthy fact that the quantity of opium which can be administered in these cases, without exposing the patient to danger from over-use of the drug, is sometimes very large.

Astringents may be administered, provided they are well tolerated by the stomach, with a certain amount of benefit—not marked, however—but they should never displace the use of opium.

Supporting measures must also be employed, and with regard to alcoholics, the same is true as with reference to opium—there is an increased tolerance. We cannot go too far, in severe cases of epidemic dysentery, in the use of alcohol, if we do not carry it beyond its supporting effect, and the life of the patient may depend upon its use. *The persistent use of opium and alcoholics is the most essential feature of the treatment of epidemic dysentery.* If the disease be associated with other affections, additional indications may be derived from the latter. For example, if the disease be associated with malaria, the use of *quinine* is indicated, and other indications may be developed by complications with other diseases.

DR. F. RAWLE, OF ENGLAND,

recommends the following treatment in the *British Medical Journal*, January 27, 1883.

First, having placed the patient between warm blankets, he proceeds to inject a pint and a half of warm water, at a temperature of 90° Fahr. This is seldom retained longer than a few minutes, but is pronounced very grateful to the patient. When the water has soothed the mucous membrane of the colon and rectum, and brought away any *effete* matter, he then proceeds to administer a small injection of two ounces, by measure, with a gum-elastic bottle. The form he administers is the following:

611. R.	Quiniæ disulphat.,	gr. x	
	Tinct. camphoræ comp.,	f. ʒ iv	
	Decoctum amyli ad.	f. ʒ ij.	M.

And when about milk-warm, inject.

It is generally retained, but if ejected, it may be repeated after an hour or two. This he has found of great service, and very grateful to the patient. He does not stop to inquire how it acts, but the effect is like magic. If griping pains be felt over the region of the epigastrium, he administers half-drachm doses of chlorodyne, in some aromatic water,

mint, caraway, or aniseed. The diet, of course, should be of the most soothing kind: jellies, isinglass, linseed, toast and barley water, *ad libitum*. Ipecacuanha is of little service. He has used with advantage warm turpentine stupes on warm flannels, over the hypogastrium.

DR. CARL SCHWALBE, COSTA RICA.

In the *Deutsches Archiv für Klinische Medizin*, March, 1875, this writer gives a very judicious treatment of acute dysentery as it occurs in hot climates, by pursuing which he had the good fortune not to lose a case. He advises, whenever a diarrhoea sets in during a heated term, or immediately after a marked fall in the temperature, to suspect the possibility of dysentery, and at once to administer a large enema of *lukewarm water*, 100° Fah., after it has been thoroughly boiled. This should be repeated until one and a half to two gallons have been thrown up, and the colon thoroughly emptied. After the onset of the dysenteric symptoms, these copious clysters should be repeated thrice daily. Against the tenesmus, when very annoying, Dr. SCHWALBE uses:

612. R. Atropiæ sulphatis, gr. j
Aquæ destillatæ, f. ʒj. M.

Two or three drops every half hour, in water, until the pupil enlarges and the throat feels dry.

When there is much intestinal pain the following recipe proves of value:

613. R. Acidi muriatici diluti, f. ʒij
Morphiæ sulphatis, gr. ij.
Aquæ destillatæ, f. ʒiij. M.

A teaspoonful three or four times daily.

Mustard plasters should be applied to the abdomen. The diet should be fluid and nutritious. White of egg water, made by shaking thoroughly the whites of four eggs in a pint of boiled and cooled water, is especially serviceable.

Dr. JOHN G. EARNEST, of Atlanta, Ga., adds his testimony to the efficacy of hot water injections. Beginning with tepid water, the temperature is gradually increased until the limit of toleration is reached. In conjunction with the hot water treatment, the author uses ipecac in full doses (twenty to thirty grains) and small doses of morphia hypodermically.

PROF. W. C. MACLEAN, M. D., ENGLAND.

Ipecacuanha should be administered in large doses as early in the disease as possible. From grs. xxv to xxx should be given, in as small a quantity of fluid as possible. Some advise the administration half an hour previously of ℥xxx of laudanum, or a few drops of chloroform. The patient should be kept perfectly still in bed, and abstain from fluid for at least three hours. If thirsty, he may suck a little ice, or may have a teaspoonful of cold water. Under this management, nausea is seldom excessive, and vomiting rarely troublesome, usually not setting in until two hours after the medicine has been taken. *Sinapisms* or *turpentine stupes* should be affixed to the abdomen. In from eight to ten hours, according to the urgency of the symptoms and the effect produced by the first dose, *ipecacuanha*, in a reduced dose, should be repeated, with the same precautions as before. The effects of this treatment are surprising, and soon manifest; the tormina and tenesmus subside, the motions quickly become feculent, blood and slime disappear, and often, after profuse action of the skin, the patient falls into a tranquil sleep, and awakens refreshed. The treatment may require to be continued for some days, the medicine being given in diminished doses, care being taken to allow a sufficient interval to admit of the patient taking some mild nourishment suited to the stage of the disease. As the disease abates, the dose should be reduced, but it is well to administer grs. x-xij at bedtime for a night or two, after the stools are, to all appearance, healthy.

Fomentations, *turpentine stupes*, or *chloroform liniment*, to the abdomen, lessen tormina and diminish suffering. If a little diarrhoea without the dysenteric odor remain, it may be checked with a small quantity of an astringent mixture; but *astringents*, in any shape, during the acute stage, are not only useless but dangerous.

The *hot bath* is of great service in the mild forms of dysentery. It should be brought to the bedside, and kept at a high temperature, the patient remaining in until he feels faint. He should then be quickly and carefully dried, put to bed, and given *ipecacuanha* (gr. xv-xx,) which may require to be repeated in eight or ten hours, the patient abstaining from all fluid for some hours after taking the medicine, and *turpentine stupes* being applied to the abdomen. This simple form of treatment, if it be resorted to early, will suffice in a great many cases of the mild form of dysentery which follows chills, without much charging the system with malaria.

DR. NATHAN S. DAVIS, CHICAGO.

This teacher states that in the treatment of acute dysentery he has never succeeded with the evacuant treatment, by laxative salines. Nor does he think highly of the method by ipecacuanha in large doses, although in some cases he states that it does produce the happiest effects.

The plan he usually follows in the first stage is to give, if the patient has not had free evacuations at the beginning of the attack, gr. iv-v of calomel, either with or without gr. ij-ijj of ipecac., followed in five or six hours by a laxative of castor oil, sulphate of magnesia, or Rochelle salts, as it is safe to assume in such cases that there is more or less fæcal matter retained in the alimentary canal.

The patient is then placed upon some combination that is sufficiently anodyne to overcome the pain and reduce the frequency of the discharges. *Turpentine* has some property that diminishes inflammatory action of the mucuous membrane, especially after the first stage. It is not merely astringent or tonic, but it possesses an alterative influence that is valuable in the peculiar condition of the vessels that belongs to these cases after the acute stage.

614. R.	Olei terebinthinæ,		
	Tincturæ opii,	āā	f. ʒ ij
	Acaciæ,		
	Saccharæ albæ,	āā	ʒ ij
	Aquæ menthæ,		f. ʒ ij. M.

One teaspoonful every four hours.

When this emulsion begins to nauseate it should be suspended. In many instances it may profitably be alternated with an anodyne carbolic acid solution, as :

615. R.	Acidi carbolicæ crystal.,	gr. vj	
	Tincturæ opii camphoratæ,	f. ʒ iss	
	Glycerinæ,	f. ʒ ss	
	Aquæ,	f. ʒ ij.	M.

A teaspoonful every four hours.

DR. LIDDELL.

This writer (*Am. Jour. Med. Sci.*, Jan., 1877,) believes that *saline purgatives* are of great value in the following cases: (1) Those in which the bowels are habitually constipated, for example, females and people of sedentary habits, for such cases are generally burdened with intestinal obstructions, which can be got rid of only by repeated doses

of purgative medicine. (2) Cases of bilious dysentery. (3) Cases where portal congestion is present. (4) Cases complicated with malarial fever, or with malarial intoxication and ague-cake; and (5) Cases of chronic dysentery. In all cases the doses should be repeated until bilious and feculent matter appears in the stools. In such cases, he exhibits *cream of tartar*, in doses of three and four drachms every three or four hours, until large watery motions, tinged with bile, are produced, and the relief, as to all the distressing and dangerous symptoms of the disease, was usually "something wonderful."

CHRONIC DYSENTERY.

The treatment of this disease can be conducted on two different principles. (1) By medication by the mouth. (2) Medication by the rectum.

We shall first consider the former:

DR. B. F. THOMPSON, OF THE SEAMAN'S HOSPITAL, GREENWICH, after trying many drugs, finds them all inferior to *ipecacuanha*. (*British Medical Journal*, January, 1876.) He believes the disease is best treated by rigidly keeping the patient at rest in bed, in a supine position; by carefully regulating the temperature of the room to about 62° Fahr.; by restricting the diet to few and simple foods, chiefly milk and mutton; and by administering, at frequent short intervals, every three hours, small doses (three to five grains) of the powdered *ipecacuanha*. If nausea is produced, the dose is diminished or omitted for a time, as it is very desirable to avoid inducing any disinclination to food. Alcohol, in any form, very decidedly aggravates the symptoms, and it must be always strictly forbidden.

W. E. WHITEHEAD, M. D., ASSISTANT SURGEON, U. S. A., also recommends *ipecacuanha* in chronic dysentery. He directs the following treatment:

A *hot salt-bath* daily, or several times a week, just before going to bed; such nourishing food as best agrees; flannel next the skin; a well-ventilated, dry and warm sleeping-room; and

616. R. Pulveris ipecacuanhæ,

3 iij.

For twelve powders. One to be taken morning and evening, on an empty stomach.

As improvement takes place lessen the dose of ipecacuanha to ten grains twice a day, and then to eight grains at bedtime. After the ipecacuanha is stopped, it may be followed with advantage by ten drops of the *muriated tincture of iron* morning and night. Other testimony to the value of ipecacuanha in chronic dysentery has been recently given by Dr. A. A. WOODHULL, U. S. A., and others.

MR. R. DONALDSON, BRITISH INDIA.

This writer, in the *Indian Medical Gazette*, June, 1876, recommends the *compound tincture of benzoin* as a most efficient remedy in dysenteric affections. He says that, in Burmah, dysentery is a very common affection, and in the European, as well as in the native, exhibits a marked tendency to become chronic. In many of these cases, ipecacuanha appears to have little or no effect; any persistence in the treatment by large doses of this drug, far from being productive of good, is fruitful of positive mischief. The stomach is rendered so irritable by it, that the patient is unable to retain nourishment; and he then suffers from exhaustion, the combined effect of the disease and in-nutrition. In these cases, the tincture of benzoin, given in combination with astringents—notably with logwood—has been found extremely useful; often, indeed, acting like a charm; and it may be truly said of it, that its powers of healing diseased mucous membranes equal its performances when applied externally to wounds. The formula recommended is:

617. R.	Tincturæ benzoini comp.,	f. ʒ ss	
	Tincturæ catechu comp.,	f. ʒ j	
	Tincturæ opii,	℥ x	
	Extracti hæmatoxyli,	gr. x	
	Aquam,	ad f. ʒ j.	M.

For a draught: to be given thrice daily. If necessary, the remedy may be administered by the rectum.

LUCIUS M'GUIRE, M. D., SAN FRANCISCO.

Our author recommends the use of *iron* in chronic dysentery:

618. R.	Ferri subsulphatis,	ʒ ij
	Glycerinæ,	f. ʒ j
	Aquæ,	f. ʒ vij

One-half to be given as an injection every two hours.

A single injection will often arrest the discharges almost immediately.

The doctor has had occasion to try the Monsell salt a number of times as a *dernier resort* in dysentery, and with uniform success.

MEDICATION BY THE RECTUM.

Some excellent results in chronic dysentery have been reported by

DR. RICHARD B. MAURY, MEMPHIS,

in the *Atlanta Medical and Surgical Journal*, December, 1872, and later in the *New York Medical Journal*, March, 1876. The patient is placed in the left lateral position, used for making uterine examinations, and the interior of the rectum is exposed to view by means of a small Sims', or other appropriate speculum. This reveals one or more superficial ulcerations of the mucous surface. These are carefully cleansed with cotton-wool, and a solution of nitrate of silver, $\mathfrak{z}\text{ij}$ to $\text{f.}\mathfrak{z}\text{ij}$ of water, is thoroughly and carefully applied to the ulcerated surface. The treatment causes considerable pain, which should be relieved by morphia hypodermically, and the patient be kept in bed. The improvement is immediate, the movements become less frequent, and the sanguineous discharges ceases. In from four days to a week the caustic application should be repeated, which is often sufficient to effect a permanent cure. The following specific directions are given :

Proceed deliberately and carefully, as in any other surgical procedure. Always etherize the patient, and always stretch the sphincter ani. When properly done, these are both perfectly safe measures. The patient being etherized, the operator is enabled to explore the rectum, and make his applications deliberately and thoroughly. Paralyzing the sphincter is not only necessary to complete and satisfactory explorations, but the quieting influence secured thereby to the rectum can hardly be over-estimated. Put the patient in Sims' position for uterine examinations, and use Sims' vaginal speculum for examining the anterior and lateral walls of the rectum, and the bivalve with hinges on one side for examining the posterior wall.

If much disease is discovered on examination, the surgeon may use nitric acid. If ulceration is superficial, and the mucous membrane not much hypertrophied, the nitrate of silver does well. The patient should be required to remain in bed for a week or two, on a milk and meat diet.

DR. H. C. WOOD, JR., PHILADELPHIA,
recommends large rectal injections of nitrate of silver:

619. R. Argenti nitratis,
Aquæ,
For one injection.

ʒj
Oijj. M.

No force should ever be used. The patient should be brought to the edge of a hard bed, his buttocks resting upon a hard pillow in such a way as to elevate the pelvis, and cause the injected fluid naturally to flow downward and inward. A well-oiled, smooth, somewhat flexible, hard tube, with openings in the sides, (an œsophageal tube will answer well,) and with a closed end, must then be gently and slowly introduced from eight to twelve inches into the rectum. The free outer end of this may be fastened to a flexible india-rubber tube, in the end of which a funnel is inserted. This being elevated five or six feet, the water warmed to the temperature of the body is poured in, and by its own weight, with irresistible gentleness, forces its way into the gut. Other writers assert that this extensive irrigation with simple warm water is all-sufficient. (ALOIS MONTI.)

The securing of an even and easily-regulated hydrostatic pressure is an essential feature of this method. *Still more essential is the distention of the rectum with fluid before attempting to pass the tube through the sigmoid flexure.* This precaution secures the smoothing out of the folds of mucous membrane, and straightens the curves of the flexure, thus rendering the passage of the tube perfectly safe and easy.

DR. B. H. WASHINGTON, TENNESSEE.

This physician, whose experience in obstinate chronic dysentery has been considerable, is convinced that the treatment by enema and scarification fails unless the ulcerations are strictly confined to the rectum. (*Nashville Journal of Medicine and Surgery*, 1874.)

He directs his patients to use a wet bandage covering the whole stomach and bowels. This is covered with a dry bandage on the outside, so attached that both lie smooth on the person. It should be wet anew in cold water when it grows dry or too warm. Every other day the patient is dry-cupped down the whole length of the spinal column. As the skin is almost always dry and torpid, the patient should be washed down daily with hot water and rubbed with a coarse cloth,

until the action of the skin is well stimulated; after which such a bath twice weekly will be sufficient.

He also encourages the patient to drink as freely as possible of pure water, containing gr. xxx–xl of bicarbonate of potash to the pint. Under this treatment, he has found some exceedingly obstinate cases recover completely.

Dr. ROBERTS BARTHOLOW, Philadelphia, lays great stress upon diet. (See his remarks on diet in diarrhoea, pp. 347, 348.)

NOTES ON REMEDIES.

ACUTE DYSENTERY.

Aconitum, gtt. ij of the tincture of the root, every hour, in acute sthenic cases, has been much praised.

Argenti Nitrates, in doses of gr. ss–iss daily, reduced to fine powder, and conjoined with Dover's powder in the form of pill, is highly recommended by Dr. WARING and others, in the advanced stages of acute dysentery.

Belladonna, both internally and externally, is often extremely efficacious.

Bismuthi Subcarbonas, ʒj–ij, suspended in mucilage, ʒ i–ij with laudanum, ʒ ss, forms an excellent enema. The lower bowel should first be gently washed out with warm water.

Carbolicum Acidum. Considerable success in the treatment of dysentery is reported by Dr. A. M. RAGLAND, of Texas, (*Va. Med. Monthly*, October, 1876,) by the use of the following.

620. R.	Acidi carbolicæ,	gtt. x	
	Syrupi rhei aromatici,	f. ʒj	
	Olei limonis,		
	Olei sassafras,	āā	gtt. v. M.

A teaspoonful every two or three hours.

If tormina and tenesmus are present, add ten drops of laudanum to each dose.

Creasotum has been found by Dr. J. R. CUSHING, of Chicago, a valuable anti-dysenteric remedy.

621. R.	Creasoti,	gtt. xx	
	Acidi acetici,	gtt. xl	
	Morphiæ,	gr. ij	
	Aquæ,	f. ʒ ij.	M.

A teaspoonful every two hours until relieved.

In connection with this, he was accustomed to cover the bowels with cloths saturated with the following liniment:

622. R.	Aquæ ammoniæ,		
	Olei sassafras,	āā	f. ʒ ss
	Tinct. opii,		f. ʒ ii
	Olei olivæ,		
	Olei petrolei,	āā	f. ʒ ij M.

For a liniment. (*Chicago Medical Journal*, January, 1875.)

The same agent is strongly recommended by Dr. H. VON SWERINGEN, of Indiana, (1884.)

Ergot. BARTHOLOW recommends the following :

623.	R.	Ext. ergotæ fluidi,	f. $\frac{3}{4}$ ij. ss	
		Tinct. opii deod.,	f. $\frac{3}{4}$ ss.	M.

S.—Teaspoonful three times a day.

624.	R.	Ergotinæ or (aq. ex.),	℥j	
		Ext. nucis vom,	gr. v	
		Ext. opii,	gr. v	M.

Ft. pil. No. xx. S.—One every four or six hours.

Ergotin, in doses of gr. vj, in emulsion, or in bland enema (gr. xij–xv), has been used with excellent results by Dr. GROSS. (*Practitioner*, November, 1868.) One of the effects noted under this treatment, was the speedy reduction in the quantity of blood voided.

Gambogia. MALGAIGNE and BETZ have found minute doses of gamboge (gr. $\frac{1}{4}$, three times a day,) valuable in dysentery.

Hydrargyrum, formerly always resorted to in large doses, in acute dysentery, has now fallen into comparative disuse, except as an occasional aperient, and ipecacuanha has taken its place. Particularly in the asthenic and scorbutic forms of dysentery, mercury, in every form, should be avoided. If, in exceptional cases of acute dysentery, it be employed, the following formula, for many years very generally used in Southern India, may be given :

625.	R.	Pilulæ hydrargyri,	gr. ii–ijj	
		Pulveris ipecacuanha radicis,	gr. i–ij	
		Opii,	gr. $\frac{1}{4}$.	M.

Make pill. Take one every four or five hours.

The following is also an effectual combination :

626.	R.	Hydrargyri chloridi mitis,	gr. ss	
		Morphiæ sulphatis,	gr. $\frac{1}{4}$	
		Quiniæ sulphatis,	gr. ij.	M.

Make pill. Take one every two or three hours.

Iodinium. Dr. PARIS recommends :

627.	R.	Iodinii,	gr. ix	
		Potassii iodidi,	gr. xv	
		Aquæ destillatæ,	f. $\frac{3}{4}$ ij.	M.

To be employed as an enema in dysentery. It is sometimes necessary to administer it twice in the course of the twenty-four hours, and to continue its use several days. One of the principal effects of the iodine enema is to relieve, promptly, the tenesmus.

**Ipecacuanha*, justly called “radix anti-dysenterica,” is of the highest value in dysentery. Although, from its introduction, (by Piso, in 1658,) its power in this disease has been more or less recognized, it was not—in consequence of the practice of conjoining it with other remedies sup-

posed to possess similar virtues—until 1858 that its real value was demonstrated by Mr. DOCKER, (*Lancet*, July and August, 1858,) who re-introduced the original plan of PISO and HELVETIUS, of giving it alone, in large and effective doses. The following facts now seem to be established: that acute dysentery can be successfully and speedily treated by large doses of ipecacuanha; that this remedy is more effectual in the acute than in the chronic forms; that large doses (gr. xxv-xxx) may be given with perfect safety, without fear of hyperemesis or other ill effects. The mode of treatment, in the main, now generally practiced, is that detailed by Dr. W. C. MACLEAN, in the article on dysentery, in REYNOLDS' *System of Medicine*, a condensed statement of which has already been given. Dr. T. J. HAPPEL (*Medical and Surgical Reporter*) has had good results with the following formula:

628. R. Pulv. ipecac co, ℞j
 " ipecac, ℞iv. M.
 Ft. Chart. No. iv. S.—One after each action till relieved.

Ixora Danduca. This East India drug comes well recommended. It acts best when cases are seen early. It possesses the advantage of not creating nausea, while it has an aromatic and agreeable taste. It has also been given in 15 to 30-grain doses, three or four times a day, and may be administered to patients of all ages.

Limonum Succus should form an essential part of the treatment when the state of the gums or other circumstances lead to the belief in a scorbutic taint. A mixture of equal parts of lime juice, decoction of cinchona, and port wine, was employed with good results in the British army, in the first Burmese war.

* *Opium* fulfills three important indications: 1. Allaying pain and muscular excitement; 2. Moderating the peristaltic motion of the intestines; and, 3. Promoting the cutaneous secretion. It is considered by Dr. WARING as, on the whole, however, inferior, especially at the onset of an acute attack, to ipecacuanha. A preliminary dose of opium is often of great service in enabling the stomach to retain ipecacuanha, and in preventing its emetic operation. The dose of solid opium may be increased from gr. j to gr. iij-iv, three or four times a day, not only without inconvenience, but with benefit, there being a marked tolerance of opium in this disease. The signs that its administration has been carried as far as is consistent with safety, are nausea and vomiting, tympanitic distention of the abdomen, and scanty stools. *Tormina* and *tenesmus* are best relieved by opiate enemata. Frequently *morphia*, hypodermically, acts much better than by the mouth, relieving the vomiting, pain, and tenesmus, as if by magic.

Plumbi Acetas, in doses of gr. iij-iv, in combination with Dover's powder (gr. iv-v), may be given with advantage in the advanced stages of acute dysentery, especially when the stools contain much mucus. The following *enema* may often be advantageously employed:

629. R.	Plumbi acetatis,	gr. x	
	Acidi acetici diluti,	℥x	
	Morphiæ acetatis,	gr. ½	
	Aquæ tepidæ,	℥iv.	M.

Potassii Bitartras. In the advanced stages of acute dysentery, when laxatives are indicated, and the stools abound with thick, viscid mucus, cream of tartar seems to exercise a special influence in diminishing the mucous intestinal secretion and in unloading the portal system. It may be given with equal parts of sulphur, or with jalap, in the form of the approved pulvis jalapæ compositus.

Potassii Chloras. Injections of this salt have been strongly advocated :

630. R.	Potassii chloratis,	℥ss	
	Glycerinæ,	℥ss	
	Aquæ,	℥iij-iv.	M.

For an injection. Two or three a day, the patient to retain them as long as possible.

Quiniæ Sulphas, in full doses (not less than ℥j, in solution,) should be given in malarial dysentery, and repeated until cinchonism is induced. *Ipecacuanha* should then be given as directed.

Sodæ Chlorinatæ Liquor. Dr. E. M. MORSE, (*California Medical Gazette*, September, 1868,) has met with marked success by throwing up into the rectum and colon from two to five pints of Labarraque's solution of chlorinated soda, largely diluted, in chronic simple uncomplicated dysentery, by which are meant those cases not kept up by organic disease of the heart, or phthisis pulmonalis, nor dependent on immediate obstruction of the liver or spleen. The right strength for the first enema is twenty parts of water to one of Labarraque's solution.

Sulphur appears to exercise a favorable influence in some cases of chronic dysentery.

Tannin, in combination with opium and ipecacuanha, is often of service.

Turpentine. BARTHOLOW recommends the following in chronic dysentery :

631. R.	Ol. terebinthinæ,	℥j	
	Ol. amygdal. express,	℥ss	
	Tinct. opii,	℥ij	
	Mucil. acaciæ,	℥v	
	Aquæ laur-cerasi,	℥ss.	M.

S.—A teaspoonful every three, four or six hours. The same formula is useful in acute dysentery, after the subsidence of the more acute symptoms.

Zinci Oxidum, in doses of gr. iij-v thrice daily, is recommended by Dr. WARING-CURRAN.

Zinci Sulphas, combined with ipecacuanha and opium, has been employed with success.

Water Belt. Dr. W. MACLEAN recommends the use of a water belt over the

abdomen for some hours daily. It acts as a fomentation, and the steady, uniform pressure it maintains seems to favor the absorption of the fibrine effused between the intestinal coats. If there be much uneasiness about the anus, a water compress over the part often affords more relief than an opiate enema. The *cold hip-bath* daily, for a few months, is of service.

Mineral Waters. The Oak Orchard Water, of Lockport, New York, and the Rock Alum and similar springs of Virginia, possess astringent and tonic properties often of very decided benefit in cases of chronic dysentery.

EXTERNAL MEASURES.

Turpentine Stupes are highly serviceable in lessening tormina and tenesmus.

**Leeches* to the verge of the anus often afford great relief to the tormina and tenesmus, and, by unloading the portal and hemorrhoidal veins, exercise a favorable influence in the course of the disease.

Cathartics. When given judiciously, mild cathartics are often beneficial, but ill-chosen or ill-timed, they do much mischief. In sporadic cases a full dose of *sulphate of Magnesia* or *bitartrate of potassa* at the outset will often cut short the attack.

Bandage. A large flannel roller around the bowels acts in preventing the peristaltic action of the intestines.

Enemata. Large enemata (four or six pints thrown into the transverse colon by means of a long flexible tube per rectum) are sometimes given. Small enemata (f. $\frac{3}{4}$ iss-ij) containing opium and ipecacuanha, are often useful in allaying tormina and tenesmus. Hot and cold water enemata have each their advocates.

Trichlorphenol. This has been tried with success by Prof. V. N. POPOFF, of St. Petersburg, in dysentery by enemata of 2 per cent. solution. (*Med. Compend.*, 1884.) It is a parasiticide of remarkable power.

Position. The tenesmus is frequently relieved by placing the patient in the prone position, with pillows under the abdomen, so as to elevate the buttocks and rectum.

Suppositories. Dr. J. H. CARSTENS (*Detroit Review of Medicine*, February, 1875,) considers the treatment of dysentery by suppositories "the most rational and scientific," and instances many successes. His prescription is :

632. R.	Pulveris ipecacuanhæ,	3 ss	
	Pulveris ergotæ,	gr. xv	
	Quiniæ sulphatis,	gr. iv	
	Olei theobromæ,	q. s.	M.

Make six suppositories. One every two hours..

CHRONIC DYSENTERY.

Acidum Nitro-muriaticum Dilutum (in lotion of f. $\frac{3}{4}$ j to aquæ Oj,) applied to the abdomen, is of benefit, especially when there is hepatic disease.

Alum is much less employed in chronic dysentery now than formerly. It may be given in combination with Dover's powder.

Ammonii Chloridum. Dr. WILLIAM STEWART (*Madras Monthly Journal of Medical Science*) has found chronic dysentery, associated with chronic disease of the liver, yield to a few twenty-grain doses of the chloride of ammonium, after ipecacuanha and other remedies had failed.

**Argenti Nitras* is of great service in chronic dysentery. It should be given in the same manner as directed for the advanced stage of the acute affection.

**Arsenic*. When chronic dysentery is dependent upon the changes induced by chronic malarial infection, arsenic is often beneficial.

Benzoinum has proved successful in the hands of Dr. CHIPPERFIELD, of Madras, (*Madras Quarterly Medical Journal*, January, 1867,) given as follows :

633. R.	Tincturæ benzoini compositæ,	f. ℥j-ij	
	Tincturæ opii,	℥xx-xl	
	Pulveris acaciæ,	gr. xxx	
	Aquam carui,	ad f. ℥vj	M.

Dose—One-fourth part every four hours.

**Bismuthi Subnitras* is highly praised in subacute and chronic dysentery by TROUSSEAU and RÉCAMIER.

Carbo Ligni was recommended by Prof. NATHANIEL CHAPMAN, who found it, internally administered, to remove the acrid and offensive character of the stools.

**Cupri Sulphas*, combined with opium, is often highly serviceable in chronic dysentery.

Eserine, in doses of gr. $\frac{1}{60}$ by hypodermic injection, has controlled obstinate chronic diarrhœas in the hospitals of Berlin (*Neurol. Centralblatt*, 1883).

Fermentum has been recommended for removing the fœtor of the stools and diminishing their frequency :

634. R.	Fermenti,	f. ℥x	
	Camphoræ,	gr. xxx	
	Spiritûs ætheris nitrosi,	f. ℥ss.	M.

A tablespoonful every two or three hours, according to the severity of the symptoms.

Hæmatoxyllum, in decoction or extract (gr. x-xxx,) is an excellent tonic astringent, but inadmissible so long as any active inflammation exists.

Hydrargyri Chloridum Corrosivum, gr. j to water Oj, f. ℥j *ter die*, will often be found to relieve and cure the most obstinate cases.

**Ipecacuanha*, in the form of Dover's powder, is often of the greatest benefit in chronic dysentery, given at bedtime in full dose, (gr. x-xij,) or in small, frequently-repeated doses, in combination with nitrate of silver.

Liquor Ferri Chloridi, in doses of gtt. ij-vij, well diluted with water and syrup, is frequently employed in France. The same doses in enema, combined with opium, are often serviceable.

**Liquor Ferri Nitratis* is of great benefit, particularly in those anæmic from loss of blood and the depraving influence of malaria.

Liquor Ferri Subsulphatis, diluted, in enema, has been recommended.

Nickel Sulphate. In obstinate diarrhœa, Dr. J. M. DA COSTA has obtained good results from this substance in doses of gr. i-ij thrice daily (*Med. News*, 1884). He considers it worthy of more attention.

**Opium* is of great value in chronic dysentery, and may be given combined with nitrate of silver, sulphate of copper, or sugar of lead.

Plumbi Acetas, though less effectual perhaps than nitrate of silver or sulphate of copper, may be employed at an earlier stage than either of those salts.

Potassii Bitartras is one of the best purgatives which can be employed when laxatives are indicated. M. GUBLER has found this most useful in the diarrhœa of phthisis, and whenever ulceration of the uterus is suspected. He gives it in powders in the following form: Oxide of zinc, thirty grains; bicarbonate of soda, ten grains; in four powders, two or three daily.

DYSPEPSIA.

JOHN SYER BRISTOWE, M. D., LONDON.

The first thing to be done is to seek the origin of the dyspeptic symptoms. See that the *teeth* are in good order; that the patient masticates well; that he gives ample time to eating; that the hours of meals are regular; that the food is wholesome, plain and digestible. Note, also, the condition of the stomach, and decide whether the dyspepsia is primary or secondary to some other affection.

In all cases of chronic dyspepsia, hygienic treatment, moderate exercise, regular hours, ventilated rooms, and change of air and scene, are of extreme importance.

DR. C. F. KUNZE, GERMANY.

The use of narcotics and alcoholics in excess must be abandoned. The patient should take only a small quantity of food at a time. The injurious habit of drinking large quantities of water must be abolished. If there is deficiency of hydrochloric acid in the gastric juice, give:

635. R.	Acidi hydrochlorici,	gtt. x	
	Aquæ,	f. $\frac{3}{4}$ iv	
	Syr. simplicis,	f. $\frac{3}{4}$ j.	M.
A tablespoonful every two hours.			

If there is fermentation of the contents of the stomach, they should be removed by an emetic or stomach-pump, and the patient take:

636. R.	Sodii sulphitis,	℥ ij	
	Aquæ,	f. ℥ iv.	M.
A tablespoonful three or four times a day.			

Or:

637. R.	Benzinæ.	gtt. xx	
	Aquæ,	f. ℥ iij	
	Syrupi simplicis,	f. ℥ j.	M.
For one dose.			

DR. HENRY GIBBONS, SAN FRANCISCO.

This physician also strongly commends the use of five drops of *chlorhydric acid* in a tablespoonful of water, after eating, when the stomach feels burdened. Not only in his own case, but in a very large proportion of dyspeptic affections among patients, he has found it a most valuable remedy. From three to five drops is the proper quantity, and it may be taken either regularly before or after each meal, or on special occasions—*pro re nata*. It is the best thing he has ever given for cardialgia, or any affection of the stomach arising from excessive acidity. It should not be too much diluted, nor should it be sweetened.

DR. I. BURNEY YEO, LONDON.

In regard to the *flatulent* form of dyspepsia, this experienced practitioner considers that, in the first place, it is necessary to maintain for some time a lax condition of the bowels. In order to get rid of the tendency to the accumulation of gas in the upper part of the intestinal canal, it is absolutely necessary that not even a temporary block should be allowed to exist in the lower regions; hence the value in such cases of a mild course of laxative mineral waters, as Carlsbad, etc., which secures two or three loose evacuations before any food is taken. Even in cases of organic constriction, it often proves of the greatest service. There are some patients with whom no aperient produces so much relief as *castor oil*, the comfort of each day depending on a dessertspoonful of this medicine, taken in the early morning. The Hunyadi water also answers well, and is often efficient in small quantities. No breakfast should be taken until the bowels have been freely relieved.

With regard to food and drink, tea, coffee, cocoa, and all hot and sweet beverages, must be entirely banished from the dietary. In such

conditions, tea or coffee will almost immediately disturb the heart's action. Malt and sherry are also under the ban. Sound claret and water, or milk mixed with some alkaline effervescing water, are the safest beverages; but much fluid of any kind is hurtful. Oatmeal porridge is still better. Animal food should be restricted to roast or boiled mutton, the lighter kinds of fish, and a little broiled fat bacon. Fruit, jellies, and marmalade, should take the place of butter.

Of drugs, he has found great use in a pill of creasote, carbonate of magnesia, and extract of hop, taken immediately after every meal; and when this has failed, he has seen excellent results from a few grains of ox-gall, made into a pill with powered rhubarb, and taken after breakfast and dinner.

One of the finest preparations in dyspepsia with flatulence, is the following:

638. R. Sodii sulphitis, $\overline{3}$ ij
 Tincturæ nucis vomicæ, f. $\overline{3}$ v
 Aquæ, f. $\overline{3}$ iv. M.
 A teaspoonful thrice daily, after meals.

PROF. T. GAILLARD THOMAS, NEW YORK.

639. R. Magnesiæ sulphatis, $\overline{3}$ ij
 Ferri sulphatis, gr. xvj
 Acidi sulphurici diluti, f. $\overline{3}$ j
 Aquæ, Oj. M.

Two tablespoonfuls in a tumbler of ice water, every morning upon rising, when a ferruginous tonic, combined with a saline, is indicated.

640. R. Potassii et sodii tartratis, $\overline{3}$ ij
 Vini ferri amari, f. $\overline{3}$ ij
 Acidi tartarici, $\overline{3}$ ij
 Aquæ, f. $\overline{3}$ xij. M.

Two tablespoonfuls in a tumbler of ice water, before breakfast. Should this dose be not sufficient, two or three may be taken daily, for the result will prove tonic and reparative, as well as cathartic.

641. R. One rennet, washed and chopped,
 Vini rubri, Oj. M.

Macerate for twelve days, and then decant, filter, and add:

- Acidi nitro-muriatici diluti, f. $\overline{3}$ ij
 Tincturæ nucis vomicæ, f. $\overline{3}$ ij
 Bismuthi subnitratæ, $\overline{3}$ ij. M.

One tablesspoonful in a quarter of a tumbler of water, before each meal, as a digestive tonic.

This prescription embraces the tonic properties of nux vomica, and the peculiar restorative influence of bismuth, with a fluid which resem-

bles the gastric juice. In many cases of habitual indigestion, our author has obtained from it the best results.

642. R.	Quiniae sulphatis,	℞ ij	
	Ferri sulphatis,	℞ j	
	Acidi sulphurici aromatici,	gtt. x	
	Mucilaginis acaciae,	q. s.	M.

Divide into twenty pills. One to be taken three times a day, before each meal.

DR. WILLIAM BRINTON, LONDON.

This writer gives the following useful hints in regard to the *salts of iron*, in dyspepsia. They are generally beneficial, not only in proportion to the anæmia and general derangement of nutrition present, but in proportion to the patient's deprivation of a due share of light, air, and exercise, which is the cause of so much of the dyspepsia of civilized life. Hence, iron is more useful in females than in males; more in the sedentary dyspeptic than in the florid and over-fed. Where great nausea and irritability exist, it should generally be avoided. It should always be given immediately after a meal, excepting after tea. The *effervescing citrate* will generally be borne by the most delicate stomach. Neither this nor any other salt of iron should be given in doses sufficiently large to cause nausea or flatulence. The *carbonate*, *phosphate*, *sulphate*, and *chloride*, form a scale of increasingly irritative preparations, and therefore demand not only a diminished dose, but an increased dilution. Thus, while we may give gr. x of the citrate as a common dose, gr. ij of the sulphate, and ℥vij-vij of the tincture of the chloride, with half a wineglassful of water, are as much as most dyspeptics will safely bear. The *oxide* and other comparatively insoluble preparations should be avoided. The efficacy of the salts of iron, in dyspepsia, often appears to be increased by conjoining them with the *sulphate* or *oxide of zinc*.

Cathartics, according to Dr. W. BRINTON, are of great service in dyspepsia in emergencies; of little service, almost hurtful, as prominent features in treatment, which ought always to aim at such a tonic effect as is incompatible with habitual purgation. They should be given in the smallest doses, on an empty stomach, either before dinner, in the shape of a slowly-acting pill, or early in the morning, in that of a more rapidly-acting liquid. The pill form is preferable, as a rule, especially when the aperient is often repeated, the liquid aperients, especially the saline, often acting injuriously on an irritable stomach. Castor oil, however, is, in many constitutions, an exception to this rule.

DR. ARTHUR LEARED, LONDON.

This author has brought forward some new views on the subject of dyspepsia, (*Brit. Med. Jour.*, May, 1879,) to wit, that the cause of most cases is not defective or impaired secretions, but a diminution of the contractile movements of the stomach. He recommends that the principal meal should be taken early in the day, before the nervous system has been exhausted by nervous or bodily exertion. *Strychnia* is of great value in impaired gastric peristalsis, but care must be used in its administration, and a dose of $\frac{1}{16}$ of a grain should be but rarely exceeded. It should never be given in the form of pill.

Thymol has been found useful in checking flatulence by hindering fermentation, and charcoal capsules are recommended for the absorption of the gases when the stomach is unable to expel flatulency in consequence of temporary paralysis from over-distention.

M. CHARTERIS, M. D., GLASGOW.

Of the many symptoms of dyspepsia, the more prominent may be met as follows:

For the loss of appetite, a bitter infusion, as:

643. R.	Ammon. sesquicarb.,	℥j	
	Potass. bicarb.,	℥ iss	
	Infusum chiretæ,	ad f. ℥ vj.	M.
A tablespoonful thrice daily.			

For the flatulence, belching, and wind on the stomach, we may order a warm carminative, as:

644. R.	Tinct. cardamom. comp.,	f. ℥iv	
	Acid. hydrocyan. dilut.,	℥xl	
	Spts. ammon. aromat.,	f. ℥ ij	
	Tinct. zingib.,	f. ℥ iiij	
	Spt. chloroform,	f. ℥ ij	
	Aquæ carui,	ad f. ℥ vj.	M.
A tablespoonful occasionally.			

When the belching is accompanied by a "rotten-egg" flavor, showing the evolution of sulphuretted hydrogen, charcoal biscuits should be ordered, and creasote or carbolic acid.

For simple heartburn, black sugar is efficacious, or eating an apple, or some liquor bismuthi. When there is gastrodynia, or cramp of the stomach, a brisk purgative, followed by carminatives with a mustard-poultice, gives relief.

Waterbrash or pyrosis is often brought on by some article of diet, as oatmeal. As a remedy, pulv. kino comp. can be recommended, and Friederichshall or Pullna waters.

CEREBRAL DYSPEPSIA.

DR. JOHN S. MAIN, OF ENGLAND.

The cerebral form of dyspepsia is well seen, in many cases, where a healthy man, with a good appetite, suddenly receives bad news when sitting down to a meal. But of all conditions acting on the brain in this manner, and through the brain on the stomach, no one is more injurious, or more jarring to the cerebral elements, than uncertainty, and the worry caused by the same, more particularly in preternaturally irritable subjects. The mind, in such cases, preys upon itself; the cerebral elements seem to get jarred and out of gear; and with this condition the stomach sympathizes. Such cases are most commonly met with amongst those who are engaged in the hottest part of the "battle of life," or "struggle for existence;" and, again, amongst those chiefly whose business or profession leads to much anxiety, uncertainty, or overstretching of the mental powers. He has found that the only treatment capable of doing these cases any permanent good, is a change, in the wide sense of the term—a relaxation from business or study; and as regards medicines, not such as are meant to act on the stomach directly, but those meant to act on the cerebrum. Amongst these, he has found the most useful to be the bromide of ammonium or bromide of potassium—preferably the former—given in a sufficient dose at bedtime to secure a good night's sleep, this being often very indifferent, and so tending to complicate the case; and, combined with this, to be taken three or four times during the day, such medicines as are known to have a building-up effect on the nervous system—amongst these, the most useful being phosphorus, or the hypophosphites, and cod-liver oil. Arsenic and quinine are often also useful, and a generous diet is always indicated. Unless the stomach has passed into a state of disease (which it may do, if overtasked when in this weakened state), any of these medicines are generally well borne. It will be well to bear in mind, however, that if the mucous membrane of the stomach be in a state of irritation, quinine, arsenic, phosphorus, the hypophosphites, and sometimes even cod-liver oil, are generally inadmissible.

CLIMACTERIC DYSPEPSIA.

MR. PRANGLEY, OF ENGLAND.

This usually occurs in women about the change of life. The treatment consists in the administration of bismuth, bicarbonate of potash, and ammonia, adding valerian, if the nervous symptoms predominate, followed by quinine, strychnia, and dilute nitro-muriatic acid.

The following is the treatment adopted at

THE DEMILT DISPENSARY,

New York, as described by Dr. D. LEWIS, in the *New York Medical Journal* :

When there is constipation, they have found the rhubarb and soda mixture most useful :

645. R.	Pulv. rhei,	℥j	
	Sodii bicarb.,	℥ iss	
	Ol. menth. vir.,	gtt. iv	
	Aquæ,	f. ℥iv.	M.

A tablespoonful before meals.

The alkaline mixture probably owes its efficacy to its stimulating action upon the gastric glands—a property of alkalies which has been amply demonstrated by many experimenters. When an additional laxative is necessary, a compound rhubarb pill is ordered at bedtime, or, what is preferable in many cases, the pill of aloes, belladonna, and strychnia :

646. R.	Ext. aloës,	gr. ijss	
	Ext. belladonnæ,		
	Ext. nucis vom.,	āā gr. ¼.	M.

One at bedtime.

In contrast with such cases are those patients who are anæmic, and complain of the symptoms common to that condition—loss of appetite, palpitation of the heart, intercostal neuralgia, and headache. In some instances, this condition is a natural sequence of prolonged dyspepsia, but is more commonly dependent upon other causes, such as bad hygiene, overwork, or malarial influences. Tonic treatment is here indicated, and the following prescription is usually effective :

647. R.	Quiniæ sulph.,	gr. xij	
	Tr. ferri chloridi,	f. ℥ijss	
	Aquæ,	f. ℥iv.	M.

A teaspoonful in a wineglass of cold water, half an hour after meals. An aloes and belladonna pill is occasionally required at bedtime.

When there is irritability of the stomach (probably gastritis) with nausea and vomiting, a bismuth mixture is often ordered :

648. R.	Bismuth. subnit.,	℥ iv	
	Acid. nitric. dil.,	℥ iij	
	Tr. nucis vom.,	℥ iss	
	Aq. menth. pip.,	f. ℥ iv.	M.

A teaspoonful after meals. Shake well before using.

Since it has been pretty clearly demonstrated that bismuth acts mechanically by adhering to the mucous coat of the stomach, it is evident that a large dose should be administered. But the very large doses given by LUSANNE, MENNERET, and others (who gave ℥j per diem), no doubt hinder the excretion of gastric juice, thereby causing the cachetic symptoms which those observers found to follow its prolonged use.

NOTES ON REMEDIES.

Aloes, in doses of gr. ij-iv, combined with ipecacuanha (gr. j-ij,) two or three times a week, is useful in the dyspepsia of persons of relaxed habit, or those debilitated by long illness, especially if it be probable that the duodenum is implicated. (F. 646.)

Aqua Calcis, in doses of f. ℥ ij in milk, is useful in dyspepsia connected with acidity of the stomach ; also when the urine shows a strong acid reaction, and when vomiting is a prominent symptom.

Aqua Chloroformi, in the proportion of one part of chloroform to 2000 parts distilled water, has been found by Professor LASÉGUE to be eminently suitable in painful digestion arising from dilatation of the stomach and similar conditions (*Gaz. des Hopitaux*, 1883).

**Argenti Nitras* is useful in many obstinate cases of dyspepsia attended with morbid sensibility and hypochondriasis. The following formula may be used :

649. R.	Argenti nitratis,	gr. ss	
	Extracti hyoseyami,	gr. ij-iv.	M.

Make one pill. Take every night.

The quantity of the nitrate may be gradually increased to grs. ij-ijj daily, and this may be continued with safety for six or eight weeks, but not beyond that time. It may sometimes be advantageously combined with small doses of quinine. The diet should be carefully directed. Or the nitrate may be given in solution :

650. R.	Argenti nitratis crystallini,	gr. vj	
	Aquæ destillatæ,	f. ℥ iij.	M.

A tablespoonful to be taken at bedtime, on an empty stomach, and repeated every night, or every second, third or fourth night, according to the severity of the symptoms.

Argenti Oxidum was much employed by Dr. GOLDING BIRD in gastralgia, gastrodynia, and in those forms of dyspepsia attended with irritable stomach and pain after food. He regarded it as a valuable sedative and tonic, possessing all the virtues without the inconveniences of the nitrate. Dr. J. EYRE used it with advantage in pyrosis. It must be remembered that it cannot, in these cases, be combined with creasote, for violent decomposition, attended with heat, and even flame, follows the mixture of these substances.

**Arsenic* is a valuable remedy in *gastrodynia*, particularly when of a neuralgic character. Fowler's solution should be given after a meal, commencing with small doses (℥iij), and gradually increasing them, adding a few drops of laudanum when large doses are reached, to prevent purging. Dr. RINGER recommends Fowler's solution (℥j-ij,) shortly before each meal, in many forms of chronic dyspepsia. He finds it very effectual in checking the morning vomiting of drunkards.

Aurantii Amari Cortex, in tincture or infusion, is an elegant tonic in dyspepsia.

Assafætida, in combination with bitter tonics and mild aperients, is useful in dyspepsia attended with hypochondriasis and other mucous affections.

**Bismuthi Subnitratis* is of great value in atonic dyspepsia. Dr. W. BRINTON finds it particularly useful in the "morbid sensibility of the stomach" of old writers. Dr. WILSON FOX especially recommends it in the milder forms of gastric inflammation. He finds that in *pyrosis*, when combined with opium, it seldom fails to relieve, but that in order to effect a cure a more direct astringent, like kino, is required. Professor R. J. GRAVES considers it one of the best remedies we possess in *gastrodynia*.

Boldo and *boldine*, derived from a Brazilian plant, have been found to act as a stimulant to digestion and exert a favorable influence on torpid liver.

**Calumba*, in infusion, is a valuable vegetable bitter tonic in atonic dyspepsia. It may be given at an earlier period than other remedies of the same class, and forms an excellent vehicle for other appropriate remedies.

Capsicum is especially useful in the atonic dyspepsia of hard drinkers and long residents in hot climates. The following formula may be employed :

651.	R.	Pulveris capsici,	gr. ij-iiij	
		Pilulæ rhei compositæ,	gr. v	
		Pulveris ipecacuanhæ radicis,	gr. ss.	M.

Make two pills. To be taken daily, an hour before dinner.

**Carbo Ligni*, enclosed freshly prepared in gelatine capsules, is a valuable remedy in dyspepsia attended with obstinate flatulence and gastrodynia.

Cardamomum, in the form of the compound tincture, is highly serviceable in the dyspeptic affections of old persons.

Caryophyllum, in infusion, or volatile oil (gtt. iiij-v), is of benefit in atonic dyspepsia attended with a sense of coldness in the stomach and much flatulence.

Cascara Sagrada. In the constipation so frequently associated with dyspepsia, one of the most efficient remedies is the fluid extract (dose f. ʒss) of this drug, or its cordial (dose, f. ʒss-j). A tonic dyspepsia with inefficient hepatic action are conditions which especially indicate its use. The cascara is prepared by PARKE, DAVIS & Co., of Detroit. A formula recommended is :

652. R. Fluid extract cascarae, f. ʒj
Syrupi simplicis, f. ʒiij. M.
A teaspoonful three or four times a day.

Cerii Oxalas was much employed by the late Prof. SIMPSON in irritable dyspepsia, attended with gastrodynia, pyrosis and chronic vomiting.

Colocynth is useful in some forms of dyspepsia and gastrodynia, when no inflammatory symptoms are present. The following formula may be employed :

653. R. Extracti colocynthidis compositi, ʒij
Pilulae rhei compositae, ʒj
Saponis, gr. vj
Olei caryophylli, gtt. iv. M.
Make sixteen pills. Take one or two every half hour.

Cubebæ. Dr. W. T. SEELEY, of New York, has for many years found great advantage in the following prescription in dyspepsia :

654. R. Bismuthi subnitrat., ʒiij
Cubebæ pulv., ʒij
Magnesiæ, ʒiij
Rhei pulv., ʒj
Zingib. pulv., ʒiss. M.
Half a teaspoonful in a wineglass of water after each meal.

Eucalyptus is one of the most useful of the so-called stomachics, in atonic dyspepsia.

Ferrum. (See above, page 379.)

Gallicum Acidum has been recommended in dyspepsia arising from a relaxed state of the mucous membrane of the stomach. It is useful in *pyrosis*, unaccompanied by organic disease of the stomach, or of the liver.

Gentian, in the form of tincture in an aromatic water, is a valuable stomachic in atonic dyspepsia and in the dyspepsia of gouty subjects.

Hops, as a stomachic tonic, is quite as serviceable as many more rare and costly medicines.

Hydrargyri Pilula, in doses of gr. ij-iv, is useful in dyspepsia attended with hepatic derangement, or when the duodenum is supposed to be the seat of the disease. It may be combined with ipecacuanha (gr. j-ij) and with a sedative. Ptyalism must be avoided.

Hydrargyri Chloridum Mite is sometimes productive of marked benefit in dyspepsia characterized by hepatic derangement. But its habitual use,

even as a purgative, is not without detriment, and its constitutional effects are mischievous.

Hydrastis is very useful as a stomachic tonic, and may take the place of calumba in the treatment of atonic dyspepsia. A few drops of the tincture or fluid extract (five to fifteen) taken before meals, daily, for some time, will often cure chronic gastric catarrh, and remove the distressing headache which frequently accompanies this disease. It is one of the best remedies for the stomach catarrh of chronic alcoholism, and is, probably, the best substitute, if given in sufficient doses, for the alcoholic stimulant, when its habitual use is to be abandoned.

Hydrocyanic Acid, in doses of mij-iv , combined with infusion of columbo, is often signally useful in dyspepsia attended with gastric irritation. Dr. WILSON FOX states that in the irritative dyspepsia of phthisis it is a most effectual remedy.

Ipecacuanha, in doses of gr. ss-j, in the form of pill, with gr. ij-iv of rhubarb, taken before meals, often proves serviceable in atonic dyspepsia, especially when the biliary secretion is deficient or vitiated.

Kino is useful in the incipient stages of follicular dyspepsia. It does not appear to constipate the bowels, if diarrhoea be not present. Sir T. WATSON recommends it highly in *pyrosis*, in the form of pulvis kino compositus:

655. R.	Pulv. kino,	ʒ iv	
	Pulveris opii.	gr. xv	
	Cinnamomi,	f. ʒj.	M.

Divide into thirty powders. One thrice daily.

Dr. WILSON also regards it as the best remedy in *pyrosis*.

Kreoehyle. This preparation of meat, made on the formula of Prof. BARFF, is said to be a valuable dietetic remedy in aggravated dyspepsia. The *Birmingham Medical Review* concludes that kreoehyle is superior to beef tea, as it contains not only the extracts, but a large amount of albumen. It is pleasant to the taste, and has proved valuable in the hands of many practitioners.

**Lactic Acid*, first introduced as a remedy in dyspepsia by MAGENDIE, has been employed with success by Dr. C. HANDFIELD JONES in cases of irritative dyspepsia, when the digestion has been for a long time painful and imperfect. Its use is not advised at the commencement of a severe case, but only after the irritation is somewhat reduced. The dose is ʒxv-xx , in a tablespoonful of water, taken at meal-times. Its use may be extended to all cases where it is desirable to improve the tone and power of the stomach. But it is essential, in order to obtain good effects, that the acid should be pure and of good quality, which unfortunately is frequently not the case with that which is generally dispensed.

Liquor Potassæ.

656. R.	Inf. calumbæ,	f. ʒiv	
	Liq. potassæ,	f. ʒss.	M.

S.—A dessert to a tablespoonful three times a day, before meals.

Magnesia Sulphas, in small doses, has been found very effectual in dyspepsia accompanied by costiveness. The following formula may be used :

657. R.	Magnesiae sulphatis,	℥j	
	Infusi quassiae,	℥iss	
	Spiritus ammoniae aromatici,	f. ℥ij.	M.

Drink a wineglassful of this every morning, fasting.

Or the following may be employed :

658. R.	Magnesiae sulphatis,	℥vj	
	Magnesiae carbonatis,	℥iss	
	Vini aloës,	f. ℥vj	
	Tincturae humuli,	f. ℥ij	
	Acidi hydrocyanici diluti,	℥xv	
	Infusi cascarrillae,	f. ℥vij.	M.

Take three tablespoonfuls thrice daily.

Manganesii Oxidum Nigrum, in doses of gr. x-xv thrice daily, has been recommended in certain irritable conditions of the stomach, and forms of dyspepsia.

Mineral Acids. BARTHOLOW highly recommends muriatic acid after meals, with or without pepsin, in atonic dyspepsia. In an excess of acid of stomach-juice, the mineral acid before meals. In excess of uric acid as well as of stomach acid, nitric acid before meals.

Midzu Ame. This is a malted extract of rice, prepared in Japan, and is considered to possess nutrient and analeptic qualities of an unusually high order. It has been advocated by Dr. J. C. BERRY (*Med. News*, 1884,) in indigestion from nervous exhaustion, as an adjuvant to food medicines, and as a restorative in exhausting diseases.

Morphia, subcutaneously injected, in doses of one-fourth grain, has been used with excellent success by Dr. CLIFFORD ALBUTT (*Practitioner*, June, 1869), in those forms of dyspepsia characterized by irritability or erethism, or such as is often associated with hysteria, or which arises from great mental exertion or depression.

Muriaticum Acidum Dilutum, in doses of ℥xv-xx, properly diluted, given immediately before, during, or directly after a meal, is highly recommended by Dr. WILSON FOX in atonic dyspepsia. It may be combined with vegetable bitters. In nervous disorder of digestion associated with diarrhoea, he combines the acid with tincture of nux vomica, and gives before meals.

* *Pancreatine* is highly recommended in conjunction with pepsin.

Pepsine is especially indicated in dyspepsia connected with deficient secretion of gastric juice. It is particularly useful in gastric disturbances following the use of animal food, often enabling a patient who has not dared attempt it, to eat of it with impunity. The severest cases of *gastralgia* are relieved by it. If it fails after three or more doses, it is probable that the dyspepsia does not arise from a defect of the gastric secretion. Dr. WILSON FOX strongly recommends it, not only in atonic dyspepsia,

but in irritative states of the gastric mucous membrane. It may be advantageously combined with muriatic acid, at meal times. It may be combined with other medicines which do not impede its therapeutic action; thus, with muriate of morphia, to relieve violent pain of the stomach; with strychnia, to stimulate peristaltic movement; with sub-nitrate of bi-smuth, lactate or iodide of iron, etc.

Potassii Sulphuretum, in doses of gr. x, alone or combined with an aromatic, is useful when the mucous follicles are supposed to be implicated.

Potassii Sulphas, in doses of gr. xv- \mathfrak{z} ij, alone or combined with rhubarb (gr. v-x) or aloes, is a safe and efficient aperient in dyspeptic affections.

Potassii Tartras is sometimes employed as a mild purgative in dyspeptic affections, attended with slight febrile action.

* *Potassii Iodidum*, in small doses (gr. j-ij), with bicarbonate of potash (gr. viij-x), is recommended by Dr. W. BRINTON, in flatulent dyspepsia, in which, whether from a too starchy diet, deficient or hasty mastication, decayed teeth, the abuse of tobacco, or other causes, the salivary excretion seems either deficient in quantity, or faulty (*i. e.*, acid) in quality. Two or three administrations will often induce a marked change.

Quassia, in infusion, combined with carbonate of soda, is recommended by TROUSSEAU, in dyspepsia attended by acidity of the stomach, sour eructations, and marked especially by vertigo and a tendency to syncope. It is peculiarly useful in the dyspepsia of drunkards.

* *Rheum* is often highly serviceable, as it is warm and carminative in its nature, speedy, and neither stimulant nor drastic in its action. Eight grains made into pills with soap, and taken every night at bedtime, together with some mild, bitter infusion, and an alkali in the daytime, is a very effectual mode of treatment. The following mixture is useful :

659. R.	Aloës,	\mathfrak{z} j	
	Rhei,		
	Glycyrrhizæ,	āā	\mathfrak{z} ss
	Spiritus lavandulæ compositi,		f. \mathfrak{z} ss
	Aquæ calcis,		f. \mathfrak{z} viij. M.

Infuse for twelve hours, and strain. Dose, two tablespoonfuls, two or three times daily.

The following formula is of benefit when acidity is a prominent symptom. :

660. R.	Rhei,	\mathfrak{z} iss	
	Potassii carbonatis,	\mathfrak{z} ij	
	Aquæ ferventis,	\mathfrak{z} xij.	M.

Macerate for twelve hours, strain, and add :

Tincturæ cinnamomi compositæ,	f. \mathfrak{z} ij.
-------------------------------	-----------------------

Dose, a tablespoonful to a wineglassful.

The old plan of chewing a piece of solid rhubarb is an excellent one, as by this means no more is swallowed than what is dissolved in the

saliva, and this is frequently sufficient to keep the bowels regular, and impart tone to the digestive organs without producing subsequent constipation.

**Ricini Oleum*, in doses of a teaspoonful daily, given at bedtime, is one of the most valuable remedies in dyspepsia attended with inflammation or vascular excitement of the pylorus and duodenum. The nature of the morbid condition of the mucous membrane of the stomach, is indicated by the effects of castor oil. In atonic dyspepsia, it is borne with the greatest difficulty, causing nausea and vomiting; in irritable dyspepsia, it acts severely, and with much griping; but in vascular excitement of the mucous membrane of the stomach, it soothes most admirably, and produces the best effects when it has no aperient action. It may be combined with liquor potassæ, or administered in emulsion, in coffee, or in a mild carminative water.

Sapo, in combination with rhubarb and essential oil, is useful in dyspepsia attended with acidity; also in *pyrosis*, given either alone or with opium.

Senna. A mixture of equal parts of the infusions of senna and gentian is often beneficial in atonic dyspepsia, especially when attended by constipation.

Serpentaria is recommended by Dr. A. T. THOMPSON, in dyspepsia, when the skin is hot and dry.

Sinapis. White mustard seeds, in doses of two or three teaspoonfuls, two or three times a day, have been advised in some forms of dyspepsia.

**Sodii Bicarbonas*, in full doses (℥ij-ii℥), combined with tincture of ginger (℥xxx) and aromatic spirits of ammonia (f.℥ss-j), is very useful in dyspeptic attacks attended with colic and gastric irritability, especially when consequent on the ingestion of unwholesome food. Either in atonic or catarrhal dyspepsia, it answers admirably combined with a bitter; *e. g.* :

661. R.	Sodii bicarbonatis,	gr. x	
	Tinct. cort. aurantii,	℥xxx	
	Infusum calumbæ,	ad f. ℥j.	M.

Dissolve and mix for a draught.

Sodii Hyposulphis is praised by Sir W. JENNER and Dr. W. BRINTON, in flatulent dyspepsia. In doses of gr. x-xv, in infusion of quassia, it has proved successful in *sarcina ventriculi*.

**Strychnia* or *Nux Vomica* is useful in chronic dyspepsia, in preventing distention by flatus. It is also serviceable in *pyrosis* and *heart-burn*.

Tannic Acid, which may advantageously be combined with dilute nitric acid, is often very serviceable in dyspepsia, the symptoms disappearing under its use, the appetite improving, and the bowels, far from being constipated, often becoming more free.

Taraxacum is much used where the liver is implicated, and in the indigestion of gouty subjects. The following formula may be employed :

662. R.	Extracti taraxaci,	3j	
	Potassii nitratis,	3 ^{ss}	
	Spiritus ætheris nitrosi,	f. 3j	
	Infusi aurantii,	f. 3vj.	M.

Make two tablespoonfuls thrice daily.

Wild Cherry is an excellent stomach tonic and may be substituted for calumba in the class of cases in which the latter is considered especially applicable.

Zinci Sulphas is recommended by Dr. GOLDING BIRD in dyspepsia associated with irritability of the nervous system.

Zinci Oxidum has, according to Dr. WILSON FOX, the advantage over the sulphate in irritable conditions of the nervous system associated with dyspepsia, when given at bedtime, in doses of gr. ij-ijj, for procuring sleep.

GASTRITIS, GASTRO-ENTERITIS AND ENTERITIS.

These forms of intestinal inflammation are closely associated, both anatomically and with reference to symptoms and treatment. They are marked off from inflammations of the lower bowel (colitis, dysentery,) by the usual presence of vomiting and constipation, the absence of hemorrhagic discharge and tenesmus.

JOHN SYER BRISTOWE, M. D., LONDON.

In the treatment of typical acute gastritis, local measures are of great importance. Mustard poultices or hot fomentations to the epigastrium will answer in milder cases; but in severer ones, from twelve to twenty leeches should be applied. Food should be reduced to the minimum, and small pieces of ice should be swallowed, or iced milk be sipped. *Opiates* in large doses are called for, either hypodermically, which is best, or combined with bismuth or magnesia. Lime-water and minute doses of hydrocyanic acid are soothing. Stimulants are not desirable.

The treatment is much the same for the milder cases of enteritis. As regards the more severe forms, two principles of treatment may be considered well established. They are: (1) To relieve pain, and prevent the movements of the bowels, by opium. (2) To avoid the administration of purgatives. Constipation for a few days, or even a week or two, is of little consequence. The opiate is best given by hypodermic injection. Leeches, early in the disease, followed by hot

fomentations, will be a valuable measure. For the prostration which rapidly ensues, stimulants and concentrated food are required, but they must be given *by enema*, as the stomach is generally too sensitive to bear them.

DR. JOHN M. WHITE, ALABAMA.

In a number of cases of acute gastritis this practitioner (*Medical and Surgical Reporter*, March, 1877,) has used the following combination with entire satisfaction :

663. R.	Tinct. veratri viridis,	gtt. xv	
	Liq. morph. sulphatis (Magendie's,)	gtt. xxv	
	Acidi carbolici,	gtt. iv	
	Aquæ menth. piper.,	f. ʒj.	M.

A teaspoonful every three hours.

Of course, the employment of ice, leeches, and fomentations should supplement this when possible.

HENRY HARTSHORNE, M. D., PHILADELPHIA.

In the chronic forms of gastritis counter-irritation of the epigastrium with blisters will be useful. Internally, the most valuable remedy is *nitrate of silver* :

664. R.	Argenti nitratis,	gr. v	
	Pulveris opii,	gr. ijss.	M.

For twenty pills. One thrice daily.

Begin with gr. $\frac{1}{4}$ of the silver thrice daily, and prudently raise in a few weeks to gr. $\frac{1}{2}$. Another useful drug is *subnitrate of bismuth*, gr. x-xx, thrice daily. The diet must be bland, and the food taken in small quantities and at short intervals. The patient should not drink much water, but quench his thirst with small pieces of broken ice. When other means fail, an absolute *skim-milk* diet will at times succeed.

DR. MILLET, FRANCE.

For the relief of some of the symptoms of chronic gastritis, this author recommends :

665. R.	Extracti hyoseyami,	ʒij	
	Argenti nitratis,	gr. vj	
	Bismuthi subnitratis,	gr. xxx.	M.

Divide into forty pills. One morning and evening, in the variety of gastralgia which intervenes frequently during the course of chronic gastritis. Counter-irritation to the epigastrium.

666. R. Extracti belladonnæ, gr. v
 Quiniæ sulphatis, gr. xxx
 Extracti valerianæ, q. s. M.
 Divide into fifteen pills. One thrice daily, in gastralgia.

DR. DEBOVE, OF PARIS,

has had skimmed milk evaporated (cream being but slightly digestible) and the residue to a fine powder. If this be dissolved in hot water, two or three litres of milk may be injected in the same volume as a single litre. He claims excellent results from this milk powder in gastric affections.

NOTES ON REMEDIES.

Acacia. Solutions of gum arabic are valuable emollients.

Argenti Nitrates, in chronic gastritis, is praised by many authorities. (F. 664.)

Arsenic. The vomiting of chronic gastric catarrh, especially the alcoholic form, is relieved by one or two drops of FOWLER'S solution taken before meals.

Bismuthi Subnitrates has a remarkable effect on relieving pain, vomiting, and intestinal irritation. Dr. WILLIAM BRINTON, of London, recommends :

667. R. Bismuthi subnitratis, gr. x-xx
 Pulv. kino compos., gr. v-x. M.
 For one dose every four to eight hours.

Calceis Aqua is an important dietetic adjuvant.

Hydrargyri Chloridum Mite. In acute gastritis, Dr. AITKEN recommends three to five grains of calomel, followed by a dose of castor-oil.

Opium and its derivatives are indispensable in the treatment of gastro-enteritis. (See above, p, 387.)

Sodii Bicarbonas is often called for as an antacid.

Tannicum Acidum is an excellent astringent to prevent excessive purging. A convenient formula is :

668. R. Aquæ camphoræ. f. ʒ ij
 Morphine acetatis, gr. ij
 Acidi tannici, gr. x-xv. M.
 A teaspoonful every two hours.

Veratrum Viride has been used. (F. 663.)

EXTERNAL MEASURES.

Bandaging. A wide flannel bandage around the abdomen acts as a sort of splint to the part, and sometimes gives much relief.

Blisters. A small blister on the stomach or abdomen often gives prompt relief.

Cups, dry or wet, are often of service.

Fomentations. The continuous use of hot water fomentations, *as hot* as the patient can bear them, over the region of the stomach, is of great benefit in acute cases, according to Dr. WILLIAM BRINTON.

Leeches are the most convenient means of abstracting blood when they can be had.

GASTRIC ULCER.

DR. DONALD W. HOOD, OF LONDON.

This practitioner, in the *Med. Press*, 1883, makes the point that all writers on the stomach treat ulcer as a chronic disease, but none of them advise such treatment as would be adopted in regard to an ulcer occurring on the surface of the body.

In treating assumed ulcer, he thinks it imperative that the patient be kept in a recumbent position, in order that the stomach walls might be in a state of rest. When there is much enlargement of the stomach he generally uses a counter-irritant, and he freely prescribes the various preparations of opium in those cases where there is no doubt that the dyspepsia does not arise from hepatic engorgement, as he finds that it not only stimulates the bowels, but also makes the patient intolerant of restraint. Such treatment in the early stages of dyspepsia is sure to result well.

Where the presence of ulceration is well marked, such treatment should be resorted to as would be insisted on if the ulcer occurred on the surface of the body. He had found this method of treatment productive of the best results in the case of a lady who, at the time of first seeing him, was afraid to take the simplest food lest its ingestion should cause her pain. On making a careful examination of the abdomen he failed to find enlargement, but in the epigastric region there was a spot most tender on pressure. He ordered her to bed, recommended rich milk and light broth as diet, and prescribed six drops of laudanum every four hours. Iodine was suggested as a counter irritant. A simple soap-and-water enema was to be used every other day. From the first the patient's progress was satisfactory. Pain ceased, there was no vomiting, and the bowels acted regularly. She left her bed in fourteen days, and recently he received a letter stating that she was quite well. He regarded the case as representing ulceration or excoriation at that

stage in which great good can be done by a decided course of treatment.

DR. H. LEBERT, VEVEY.

A careful study of the treatment of gastric ulcer is given by this writer in the *Transactions of the International Medical Congress*, Philadelphia, 1876. His rules are :

1. Absolute rest in bed.

2. A milk diet, about two quarts per day in divided amounts every three hours, except at night. If it is badly borne, add one-fourth part lime-water or a little bicarbonate of soda. The milk diet should be kept up about six weeks. After this it may gradually be combined with light, nutritious diet.

For the pain, opiates by enema, or subcutaneously are required. Hæmatemesis can be controlled by ice externally and internally, hourly doses of ergot, or of tincture of chloride of iron (gtt. v,) or acetate of lead or alum whey. Iced milk is the best anti-emetic; or effervescent powders with bismuth; or small doses of opiates. Often, when other means fail, *iodide of potassium* will check the vomiting, as :

669. R.	Potassii iodidi,	ʒi	
	Tinct. cinchonæ,	f. ʒj.	M.
Ten to twenty drops several times daily.			

Constipation should be prevented by cold enemata, or aloes pills. Perforation demands the most absolute rest and full doses of opiates, (subcutaneously) with ice to the epigastrium. This mode of treatment has occasionally proved successful in this very dangerous complication.

As the patient manifests improvement, a mild ferruginous tonic is indicated. The tincture of the *malate of iron* is one of the best; or the lactate; or the following :

EFFERVESCENT FERRUGINOUS POWDER.

670. R.	Ferri lactatis,	gr. xlv	
	Acidi tartarici,	ʒij	
	Sodii bicarbonatis,	ʒiij.	M.

Keep in a wide-mouthed bottle, well stoppered. Dose, from a half to one teaspoonful in a small wineglassful of water, two or three times a day.

PROF. H. ZIEMSEN, ERLANGEN.

This writer (*Medicinische Central Zeitung*, April, 1872,) recommends the use of alkalies, especially Carlsbad water, continued several

months. Most vegetable, and fatty, and saccharine matters are injurious. A *milk diet* has proved of benefit in numerous instances; but veal, chicken, ham, white bread, and light claret, are harmless. Morphia, hypodermically, must be employed to combat the cramps in the stomach.

As regards the consequences of gastric ulcer, especially of stricture of the orifices, the indications are, return of the gastric catarrh, retention of the contents of the stomach with acid fermentation, vomiting after meals, constipation, sinking of the hypogastrium, dilatation of the stomach, which is capable of physical proof, etc. These symptoms require the due application of sounds and of the stomach-pump. The so-called system of dry diet may be adopted with advantage, for the obstinate constipation so commonly remaining, even for years, as the consequence of gastric ulcers. M. ZIEMSEN recommends the use of rhubarb before going to bed, to which, if any pain follows its employment, or if great atony exists, he adds a tenth of a grain of extract of belladonna, or of the extract of *nux vomica*.

J. M. DA COSTA, M. D., PHILADELPHIA.

671. R.	Argenti nitratis,	gr. v	
	Extracti opii,	gr. x.	M.
For twenty pills. One thrice daily.			

The following is an excellent combination for the pain and vomiting, when opium is inadmissible:

672. R.	Bismuthi subnitratis,	ʒ ij	
	Acidi hydrocyanici diluti,	f. ʒ ss	
	Mucilag. acaciæ,		
	Aquæ menthæ piperitæ,	āā	f. ʒ ij. M.
A tablespoonful thrice daily.			

DR. ROBERTS BARTHOLOW, OF PHILADELPHIA.

This author states that it has been proposed to treat ulcer of the stomach by absolute rest of the organ and the introduction of foods by the rectum. It should not be forgotten that the rectum is not an organ of digestion; hence nutrient enemata must contain the materials for artificial digestion. Furthermore, the mucus and fluids of the rectum are alkaline in reaction. To secure rapid osmosis, therefore, the enemata should have an acid reaction. The following formula is suitable for the purpose:

673.	R.	Beef-tea,	℥iv	
		Hydrochloric acid,	℥x	
		Glycerole of pepsin, (Scheffer's)	℥ij.	M.

If the rectum is irritable, ten to twenty drops of the tincture of opium may be added to the injection. If stimulants are indicated, brandy may also be added. The rectum soon becomes intolerant of injections; hence, the greatest care should be used in practicing them, to avoid sudden distention of the bowel and frequent introduction of nutrient materials should be avoided. Five times in the twenty-four hours should be the maximum—for artificial digestion is much slower than normal stomach digestion. LEUBE recommends the following as a nutrient injection: "Take about five ounces of finely-scraped meat; chop it still finer, add to it one and a half ounces of finely chopped pancreas free from fat, then add about three ounces of lukewarm water, and stir to the consistence of a thick pulp."

Or the following peptonized formula may be used:

"A nutritive enema should be prepared in the usual way—of milk—or of milk with beef-tea or eggs—or of milk-gruel. To half a pint of the warm enema, a tablespoonful of the liquor pancreaticus and thirty grains of the bicarbonate of soda should be added. The enema can then be administered at once."

PEPTONIZED MILK.

"Fresh milk is diluted with water in the proportion of three parts of milk to one part of water. A pint of this mixture is heated to boiling, and then poured into a covered jug. When it has cooled down to about 140° Fah., one or two teaspoonfuls of the liquor pancreaticus, and a small pinch of bicarbonate of soda (in solution) are mixed therewith. The jug is then placed under a 'cozy,' in a warm situation, for one hour. At the end of this time the product is again boiled for a couple of minutes. It can then be used like ordinary milk."

PEPTONIZED MILK-GRUEL.

"Half a pint of well-boiled gruel is added, while still boiling hot, to half a pint of cold milk in a covered jug. The mixture will have a temperature of about 125° Fah. The liquor pancreaticus and the bicarbonate of soda are then added in the same proportion as in the preceding process. The jug is placed under a 'cozy,' and kept warm for an hour and a half. The contents are then boiled for a couple of minutes,

and the product is ready for use. By this second method, the use of the thermometer is dispensed with."

SUPPLEMENTARY RECTAL ALIMENTATION.

Under this designation, Dr. A. H. SMITH, of New York, describes a method of rectal alimentation with defibrinated blood, which seems in a high degree useful. He ascertained that "three to four ounces of blood administered at night, would be so completely absorbed in the course of eight to ten hours that no trace of it could be found in the morning evacuation." To retain the blood fluid, it must be defibrinated at the moment it is drawn, which may be done by stirring it with a bundle of twigs as it flows away. In chronic cases, three to six ounces may be thrown into the rectum morning and evening; in acute cases, every two to three hours. It may be used cold, but it is better to raise it to the temperature of the rectum. Constipation usually results, and, in some instances, the body exhales a rather fetid odor, and the stools are offensive. Another objection may be urged against this method: sometimes a foul-smelling and tenacious material coats the surface of the mucous membrane and prevents absorption. For this reason, and to promote a favorable disposition of the blood, the bowel should be irrigated with water once or twice a week to clear away any retained or adherent matter. If the rectum is irritable, a little laudanum may be added to each blood enema.

DR. FENWICK, LONDON.

This author states (*Diseases of the Stomach and Duodenum*) that in ulcer of the stomach, whenever the pain is very severe and is attended by frequent vomiting, we may be sure that inflammation is going on, and the case must be treated as one of subacute gastritis. Perfect rest should be enjoined. A few leeches to the epigastrium are sometimes required; but, generally, the use of poultices, turpentine fomentations, or dry cupping, is alone necessary.

The main point in the treatment is the regulation of the diet. Usually, small quantities of farinaceous food may be given at frequent intervals. Arrow-root, milk, and gruel, agree the best. The frequent use of ice often relieves the pain, and obviates the necessity of larger quantities of liquids. Sometimes, as when dangerous hæmatemesis has previously followed a severe attack of pain, the patient ought to be supported by an enema.

For the more acute stages of the complaint, *morphia* and *hydrocyanic acid*, combined with small doses of nitrate of potassium—or, if there be much acidity, with magnesia or soda—are the best medicines.

As the pain lessens and the vomiting becomes more rare, the patient may be allowed to take a little exercise. Rather more liberal quantities of food may be given, but of a liquid nature. Blisters to the epigastrium generally alleviate the sufferings, and the raw surface may be dressed with *morphia*. When the pain is moderate, bismuth with magnesia and *morphia*, and lime-water with milk, are beneficial. As the case improves, tonics may be given. If the urine deposits phosphates, the mineral acids, with calumba, may be prescribed. Later, *iron*—the citrate or the reduced, and afterwards the sulphate—may be given. Dr. FENWICK has often prescribed the *nitrate* or *oxide of zinc*, and the salts of copper or zinc, with very good results. During convalescence, a little brandy or pale sherry may be given. In very obstinate cases, residence abroad sometimes affords relief.

DR. DEBOVE, OF PARIS.

During the first few days, this distinguished clinician (*Boston Medical and Surgical Journal*) washes out the stomach to free it of acid matters it may contain. This operation has never provoked hæmatemesis, and is considered safe. He uses a soft flexible tube, so that even if it does hit against the ulcer it can do no harm. He then administers, three times a day, three-fourths of an ounce of powdered meat, suspended in milk and mixed with one hundred and fifty grains of bicarbonate of soda. This mixture is administered at meal time and by means of the stomach-tube, as it is very disagreeable to take in the ordinary way as food. The patient is given, besides, a quart of milk a day, rendered alkaline by lime water. The patients have been confined to this diet for several months at a time, and this treatment has never given rise to any of the symptoms which characterize what has been called the “alkaline cachexia.”

NOTES ON REMEDIES.

Argenti Nitras, though generally considered of value in ulcer of the stomach, is believed by Dr. BRINTON to be, as generally prescribed, absolutely inert, so far as the gastric ulcer is concerned. He prefers bismuth. Professor DA COSTA sometimes prescribes the nitrate with opium. (F. 671.) Notwithstanding the strong opinion which BRINTON has given adversely to the use of salts of silver in the stomach, BAR-

THOLOW agrees with FOX, that these agents are, in this affection, next in value to bismuth. The oxide or nitrate may be given in pill form, or the nitrate in solution.

674. R.	Argenti nitrat.,	gr. xv
	Aq. destil.,	q. s.
	Ext. belladonnæ,	gr. x
	Ol. caryophilli,	gtt. x
	Rad. gentian pulv.,	
	Ext. gentianæ,	āā
		q. s. et fit. pil. No. xl.

S.—One pill three times a day.

Arsenic is recommended by Dr. RINGER, who gives one to two minims of FOWLER'S solution before each meal.

Atropine and morphine arrest pain and vomiting, even in very minute quantity.

Bismuthi Subnitras is highly spoken of by all authorities. It relieves pain and vomiting, as well as diarrhœa, when present.

Eucalyptus Globulus has been prescribed with great success by Dr. BENJAMIN BELL, of Edinburgh, in formidable cases of gastric ulcer. (*Edinburgh Medical Journal*, February, 1878.)

Cannabis, in extract, often answers admirably as a sedative, and would replace opium entirely, were its effects more uniform.

* *Opium* is highly recommended by Dr. W. BRINTON in ulcer of the stomach. In his opinion, its efficacy in this affection is not due to its sedative or anodyne properties, but rather to its supporting the strength, buoying the nervous system, and checking the waste or expenditure of the tissues generally. When vomiting is excessive, and resists a combination of this powder with bismuth, opium is best borne in a solid form. Thus given, its effects are often very striking. Large and frequently-repeated doses are of great service in perforating ulcer of the stomach.

Plumbi Acetas often allays the severity of the symptoms.

Potassii Iodidi. (F. 669.) Dr. W. BRINTON gives this formula as of the best effect in the flatulent dyspepsia of gastric ulcer :

675. R.	Potassii iodidi,	gr. j
	Potassii bicarbonatis,	gr. xv
	Tincturæ aurantii,	f. ʒ ss
	Infusi ca'umbæ,	f. ʒ vijss. M.

For one dose an hour after eating.

Ricini Oleum, in a moderate dose, may often be given in the ordinary way without increasing the pain or vomiting. When the latter is frequent, however, the oil is best given in enema.

* *Turpentine* stupes often relieve the pain. When, however, the strength is exhausted by constant vomiting, *dry cupping* is preferred by Dr. W. BRINTON, as a more effectual means of relieving the pain. In comparatively recent cases in the young and well-nourished, *blisters* are to be preferred.

**Enemata*. Life may be prolonged, and even in some cases preserved, in those ulcerous conditions of the stomach in which it is unadvisable or impossible to give food by the mouth, by the persevering use of enemata of beef-tea, milk, raw eggs, cod-liver oil, wine, and, in extreme cases, even diluted brandy or opium. The quantity of each should be as small as possible, and should rarely be repeated in less than three or four hours.

**Ice*, in small lumps and raspings, greatly relieves the pain and vomiting.

Leeches to the epigastrium often relieve the pain, but their use is of doubtful propriety in the weak and cachectic.

Milk Diet. This is referred to above (p. 391); ass' and mare's milk are best, but not often procurable. The milk should always be obtained from the same animal, and used warm, (95° Fah.,) but not boiled. Half a pint every four hours is a fair average. It should not be taken at a draught, but by spoonfuls, slowly, to prevent curdling in the stomach. Crusts of bread or fresh crackers may be soaked in it, or lime-water or soda added to it. Apollinaris water is a pleasant addition. Where the irritability of the stomach is not excessive, arrow-root may be mixed with the milk and boiled to a thin pulp, to be taken cool. Sago and tapioca are generally less easily borne. Rice flour may be used in the same way.

HÆMATEMESIS.

DR. JACCOUD, PARIS.

There is no better treatment than that by *ice*, internally and externally. The mineral acids or alum may be given, but their efficiency is neither as rapid nor certain as that obtained from ice. Dry cups on the limbs and trunk are useful when the hemorrhage is copious.

FREDERICK T. ROBERTS, M. D., M. R. C. P., LONDON.

The judicious rules laid down by this author, in his treatise on *Practice*, are; bodily rest, and nutrition by enema, if the case is severe; small lumps of ice should be swallowed at intervals, and a light ice poultice to the epigastrium, followed by sinapisms, if the vomiting is violent. It is of the utmost importance to check the emetic efforts, and if the means just mentioned are insufficient, recourse should be had to a few drops of dilute hydrocyanic acid in cold mucilage; morphia, internally or by subcutaneous injection; or an enema containing tincture of opium. Should there be signs of sinking, stimulants are

called for, and should be given by enema. The most efficient medicines are gallic acid or acetate of lead, in full doses, combined with opium; or oil of turpentine.

DR. A. HILLER, OF GERMANY.

Hæmatemesis is sometimes due to visceral syphilis, when specific treatment must be instituted.

NOTES ON REMEDIES.

Alum, in doses of gr. viij, in combination with opium, thrice daily, is useful, though inferior to sugar of lead.

Ammonii Chloridum is sometimes employed with advantage.

Argenti Oxidum, in doses of gr. ss-j, thrice daily, is highly recommended by Sir J. EYRE.

**Ergot* and *Ergotine* have been successfully employed by many practitioners.

**Ferri Nitratis Liquor*, in large doses, is very efficacious.

Ferri Chloridi Tinctura, gtt. v, in water, every half hour or hour, is recommended by Dr. LEBERT.

**Gallicum Acidum*. Dr. W. BRINTON recommends the following formula :

676.	R.	Acidi gallici,	ʒij	
		Acidi sulphurici diluti,	℥xl	
		Aquæ,	f. ʒij.	M.

Dose, a tablespoonful in a little water.

**Ipecacuanha* often appears to exercise a powerful influence. It should be given in the manner recommended in hæmoptysis.

Magnesiae Sulphas. Dr. BARLOW considers that, in hæmatemesis and melæna, it is advisable to clear out the bowels before resorting to astringents. For this purpose, he employs the following formula :

677.	R.	Magnesiae sulphatis,	ʒj-ij	
		Acidi sulphurici diluti,	℥x	
		Aluminis,	gr. x	
		Syrupi papaveris,	f. ʒ ss	
		Infusi rosæ compositi,	f. ʒ xj.	M.

Make a drink. Take every four hours.

Matico has been advised.

**Plumbi Acetas*, in doses of gr. ij-ijj, combined with acetate of morphia (gr. $\frac{1}{4}$ -ss), is very useful in controlling hemorrhage from the stomach, whether idiopathic or from ulceration. The auxiliaries are the external application of cold, acidulated drinks, perfect rest, and antiphlogistic diet.

**Sulphuricum Acidum Dilutum*, alone or combined with gallic acid, is a very efficient remedy.

Tannicum Acidum, in doses of gr. iii-vj, three or four times a day, is a useful astringent and tonic, only contra-indicated by the presence of inflammatory symptoms.

Turpentine deserves a trial after the failure of less nauseous remedies.

**Ice*, in lumps, coming in contact with the bleeding vessels, acts as a powerful astringent.

HEPATITIS, HEPATIC ABSCESS.

DR. GEORGE HAWLEY, LONDON.

The treatment of hepatic abscess, whether it be of the idiopathic, traumatic, metastatic, or pyæmic varieties, in so far as the local suppuration is concerned is always the same. But the constitutional treatment varies according to the cause of the suppuration.

1. If called early, when suppuration is threatened, but before pus has actually formed, try to arrest the onward progress of the disease by leeching, cupping, and the application of a freezing mixture of pounded ice and salt over the most pronounced seat of pain. Not only is the freezing process to be continued until the subjacent parts are frozen quite hard, but until actual blistering of the skin subsequently takes place, as then, and then only, is the cold communicated to the deep tissues of the liver sufficiently intense to abort the suppurating process.

2. In the incipient stages avoid hot fomentations and poultices: they favor the formation of pus.

3. Administer a brisk mercurial purgative. Enjoin strict rest of body and mind. Put the patient on a low diet, and keep the room well ventilated and of a temperature of not more than 66° Fahr.

4. Prescribe germicides, in the form of salicylic, carbolic or mineral acids, and quinine. Do not give alkalies.

5. If the case is not seen until pus has formed, all we can do is to use iodine liniment, mustard poultices, or blistering, in the vain hope of arresting further progress.

When it is desirable to favor the suppuration towards bursting, the temperature of the room should be raised to from 65 to 70° Fah. Sleep must be encouraged by chloral, bromide of ammonium and such like, but not by opiates. The diet should be milk and eggs, animal and vegetable soups, and no solids, except white fish, such as whiting,

haddock, sole, turbot, cod, etc; no mackerel, eels or salmon, lobsters or crabs. No brandy or whiskey unless their use be distinctly indicated; no port wine or dry sherry; no heavy indigestible malt liquors. But in their place light nutritious alcoholic drinks. Good sound claret, hock or Moselle. Effervescing drinks in moderation.

PROF. WILLIAM A. HAMMOND, NEW YORK.

This eminent clinician is of opinion that obscure abscesses of the liver are far more common than are usually supposed. (*St. Louis Clinical Record*, June, 1878.) They may exist without local symptoms or disturbance of the general health. The treatment, in all such cases, is to open them at the earliest possible moment, and without waiting for adhesions to form between the liver and the abdominal wall. This can readily be done, without any danger, by *aspiration*. There is no risk of peritonitis. The proper place for performing the operation of aspiration, is in one of the intercostal spaces, usually the tenth, a little behind a line let fall from the middle of the right axilla.

DR. WILLIAM AITKEN.

In acute parenchymatous inflammation of the liver, this writer prescribes antimonii et potassii tartras (gr. $\frac{1}{8}$ – $\frac{1}{4}$) every two or three hours, providing it can be given within the first three days, when there is much vascular excitement and a full, bounding, unyielding pulse, with a dry, hot skin, and scanty urine. General blood-letting may be adopted, if there be evidence of obstruction to venous blood-flow through the right side of the heart. Local depletion, by leeches over the liver and also round the anus, must be at once resorted to, followed by fomentations and linseed-meal poultices, with or without mustard or laudanum, over the hepatic region. Saline purgation and alkalies, with or without colchicum, are also to be freely administered. Hot *turpentine epithems* are most beneficial.

In the chronic form, *iodide of potassium* with *taraxacum* is of great service. Alcoholic stimulants and fermented drinks must be absolutely forbidden, and the diet restricted to mild nutriment, such as milk, beef-tea, and farinaceous food generally.

DR. J. KINGSTON FOWLER, OF ENGLAND,

considers that the trocar is preferable to incision for evacuating hepatic abscesses, for the following reasons :

1. The risk from hemorrhage is much less; for when the liver is incised there is often a violent gush of blood. This, it is true, soon ceases; but these patients are usually not in a condition to bear a loss of blood which a distinguished surgeon, Mr. Lister, describes as "alarming."

2. There is less danger from septic absorption along the track of the wound, as the pus flows through a canula or drainage tube.

3. As a trocar of any diameter may be used, the opening into the sac may be of any size that is considered desirable. For an exploratory puncture he prefers one having a diameter of one-eighth of an inch; but if there is certain evidence of the presence of pus, it is very important to employ an instrument of at least three-eighths of an inch or half an inch in diameter. If a smaller one be used, it is liable to get blocked by the solid shreds of liver tissue which these abscesses so frequently contain.

PROF. W. C. MACLEAN, M. D., LONDON.

This distinguished surgeon strongly opposes the usual treatment of acute hepatitis by venesection, mercurials, and purgatives, given above by Dr. AITKEN, and recommends, in place of it, *ipecacuanha*, in large doses (gr. xx-xxx), repeated every few hours, until the symptoms subside:

678. R.	Pulveris ipecacuanbæ,	℞j	
	Mucilaginis acaciæ,	q. s.	M.

Make four pills. The whole to be taken every sixth hour, day and night.

DR. ALFRED A. WOODHULL, U. S. A., has recently published some cases corroborating the value of this treatment. Dr. MACLEAN considers *ipecacuanha*, in hepatitis, almost as valuable as in dysentery.

DR. GEORGE HARLEY, LONDON.

The first thing in a case of acute hepatic congestion is to enjoin strict rest. The second is to place the patient on low diet. The third, to freely clear out the bowels. The fourth, to relieve the slight discomfort by the application of hot, thick, and large linseed poultices. The fifth, if there be signs of acute inflammation, to apply a freezing mixture of ice and salt or leeches, or even cupping glasses over the painful hepatic region. Mercury is here our sheet-anchor, both as a purgative and an antiphlogistic.

J. M. DA COSTA, M. D., PHILADELPHIA.

In chronic hepatitis, a certain amount of drain should be kept up from the portal circulation. For this purpose, very small doses of *podophyllin*, *cream of tartar*, or *Rochelle salts*, now one and now another, are useful. A very good pill is the following:

679. R.	Podophyllin,	gr. vj	
	Capsici,	gr. iv	
	Pulyeris rhei,	gr. xij.	M.

For twelve pills; one on alternate nights.

In order to reduce the state of induration or enlargement of the liver, the *mineral acids* may be employed. Or, when the case is not chronic, the salts of soda, as in the following formula:

680. R.	Sodii bicarbonatis,	$\bar{3}$ ss	
	Infusi gentianæ,	i. $\bar{3}$ vj.	M.

A tablespoonful three times a day, after or between meals.

Should this fail, employ *nitro-muriatic acid baths*. *Sulphur-baths* are of advantage; or those of *sulphuret of potassium*; or the use of *sulphur ointment*, followed by warm baths.

SIR RANALD MARTIN, LONDON.

Nitro-muriaticum acidum is employed, both externally and internally, with great advantage in chronic hepatitis, and in the advanced stages of acute hepatitis and congestion of the liver. It is most serviceable in the form of a *bath*, for which the following directions are given (*Lancet*, December 9th, 1865,) by Sir RANALD MARTIN: 1. The proportions of acid are given by the following formula:

681. R.	Acidi muriatici,	f. $\bar{3}$ iij	
	Acidi nitrici,	f. $\bar{3}$ ij	
	Aquæ,	f. $\bar{3}$ v.	M.

2. Two gallons of water (about ten bottles) may suffice for a bath. 3. To each gallon of water, add f. $\bar{3}$ iij of the above acid mixture. 4. The bath thus prepared will keep in use for three days, by adding f. $\bar{3}$ ss of the acid mixture and Oj of water, morning and evening, to make up for the waste by evaporation. 5. A portion only of the bath to be heated for use, after which it is to be added to the remainder, so as to make the whole of a comfortable warmth (96°–98°.) 6. Let both feet be placed in the bath, while the inside of the legs and thighs,

the right side (over the liver,) and inside of both arms, are sponged alternately. This should be continued for ten or fifteen minutes, morning and evening. 7. While using the bath, a gentle aperient should be taken every morning. 8. Earthenware or wooden vessels should be preferred for foot-baths, and all the sponges and towels should be kept in cold water, as the acid corrodes them. 9. In urgent cases, a general bath, to envelop the whole body, may be used. 10. If the acid bath create much irritation of the skin, the quantity of the acid may be diminished. 11. The influence of the acid is not in the least degree counteracted by opium, even when exhibited in the largest doses.

NOTES ON REMEDIES.

Ammonii Chloridum has for many years been esteemed in hepatic abscess. According to Dr. WILLIAM STUCKERT, of the British Indian Army, (*Medical Times*, April, 1878,) in active congestion of the liver, the special and characteristic action of the medicine will be found to be more marked than in cases of chronic hepatitis. The dose necessary for its full therapeutic effect is gr. xx, and the only condition which contraindicates its use in cases of hepatitis is the existence of a dry and hot skin. Under such circumstances, its use should not be commenced till the skin is rendered moist and perspirable by the administration of some simple diaphoretic mixture in repeated small doses, say \mathfrak{z} ij liq. amm. acet. with \mathfrak{m} v tinct. hyoscyam. in each dose, every half hour. In congestion of the liver (or acute hepatitis, when the skin has been made moist or perspirable by the above means) the chloride should be at once commenced in twenty-grain doses, twice or thrice daily.

Colchicum is advised by Dr. COPLAND, to increase the biliary secretion in chronic inflammation or enlargement of the liver, and to promote the resolution of the former and the diminution of the latter state.

Hydrargyri Bromidum has been administered with relief in chronic hepatitis with enlargement.

Hydrargyri Iodidum Rubrum, in ointment, is recommended by Dr. W. MACLEAN, in malarial enlargements of the liver and spleen, to be rubbed in for ten minutes, in the morning, and the patient exposed to the powerful heat of a fire as long as he can endure it.

Hydrargyri Chloridum Mite. Mercury is much less employed than formerly in chronic hepatic affections, it having been superseded by nitro-muriatic acid and iodide of potassium.

Iodine and its preparations are beneficial in chronic hepatic affections, but their indiscriminate use is to be avoided.

Nitricum Acidum, with sarsaparilla or taraxacum, is often productive of good in chronic hepatitis, especially of old persons, where mercurials have been largely used. Nitro-muriatic fomentations and baths may be employed at the same time.

**Potassii Iodidum*, combined with taraxacum, is of value in chronic inflammation and enlargement of the liver. Dr. COPLAND has found that it is chiefly when enlargement, obstruction or torpor of the liver occurs after periodic fevers, or in the scrofulous diathesis, that iodide of potassium, employed internally or externally, or both, and in conjunction with other deobstruents, as liquor potassæ, or alternated with purgatives, is of most benefit.

**Taraxacum* is an efficient remedy in chronic inflammation of the liver; where mercury is advisable, rendering it more efficient, and under certain circumstances, taking its place advantageously. Sir RANALD MARTIN has found, in indolent enlargements of the liver, accompanied with torpid action of the viscus, that mercury is of little service, and that he has derived more advantage from the following formula, conjoined with a spare diet and the external application of cantharides :

682.	R.	Extracti taraxaci,	gr. xxxvj	
		Extracti aloës,	gr. xij	
		Extracti acetici colchici,		
		Pulveris ipecacuanhæ,	āā	gr. vj. M.

Make twelve pills. Take two every night.

INTESTINAL WORMS.

TÆNIA—TAPE WORMS.

The varieties of tape worms are often so difficult to dislodge, that the therapeutic resources of the physician are severely tasked. The principal remedies are mentioned below.

PROF. ARNOLD HELLER, KIEL.

Before commencing treatment the intestinal tract should be cleared out by mild laxatives and enemata for two days. The diet should be limited and concentrated. As onions and salt herrings are unpleasant to the worm, the patient may partake freely of these. On the morning of the third day he should take ʒv of *koosso*, and two hours after this an ounce of castor oil. Instead of the bulky *koosso* itself, its alkaloid, *koosin*, may be used; of this, a dose is gr. xxx.

DR. J. G. BROOKS, OF KENTUCKY.

683.	R.	Ext. male fern.,	f. ʒj	
		Emul. ol. ricini,	(50 pr. ct.) f. ʒiij.	M.

S.—All to be taken at once, after twenty-four hours fast.

DR. CARRE, FRANCE.

684. R.	Pulv. kousso,	℥vij
	Olei ricini (hot),	℥iss.
	Strain, and pour on the residuum	
	Aquæ bullientis,	℥iss.

Filter and combine the two percolates by means of yolk of egg in an emulsion, and add gtt. xl ether sulphuric. Sweeten and aromatize. For one dose.

DR. TILLESEN, GREIFSWALD.

This writer (in the *Deutsche Medicinische Wochenschrift*, Nos. 46 and 47, 1876,) speaks of the method of treating tænia in the Greifswald medical clinic. It consists essentially in clearing out the intestine with castor oil and enemata, followed by the exhibition of liquids containing substances supposed to be distasteful to the parasite, such as herring-brine, and so forth. In addition, a considerable amount of water is given, which is supposed to distend the tænia and separate it from the intestinal wall. The increased weight of the parasite, together with the peristaltic action of the intestine, suffices to remove the worm. Several cases are reported, and in each the head was brought away.

PROF. MOSLER.

The chief characteristic of this practitioner's treatment is the injection of large quantities of *warm water* into the colon, after the administration of the anthelmintic. The diet is first regulated, food being given which is supposed to be distasteful to the tape-worm—bilberry tea, herrings, sour cucumber, salted meats. The intestine having been, as far as possible, emptied by laxatives, a dose of the extract of pomegranate bark is administered, prepared from the fresh bark, and then a large quantity of warm water is injected into the rectum. The theory is that the worm, previously brought down into the colon, is prevented by the water from attaching himself to the wall, and is brought away by the liquid on its escape. It is asserted that in every case in which this treatment was adopted, the head of the worm was removed.

DR. M'PHAIL, VA.

After keeping the patient on a light diet for the preceding twenty-four hours, give from ℥ss-j of the oil of male fern in ℥ij of thin mucilage, and ten minutes afterwards, a goblet of sweet milk. Two hours

after the above dose, give $\mathfrak{z}\text{j}$ castor-oil, combined in mucilage, with $\mathfrak{z}\text{j}$ —iv of spts. turpentine. This always does the work, and with remarkably little discomfort to the patient. (*Virginia Medical Monthly*, 1876.)

SIR THOMAS WATSON, M. D.

This eminent physician gives the preference to *oil of turpentine*, in large doses— $\text{f.}\mathfrak{z}\text{ss}$ —ij, in combination with castor-oil; or castor-oil may be administered afterwards, to assist its purgative effect. The patient should take it in the morning fasting, and no drink should be admitted into the stomach until the medicine begins to operate, lest sickness and vomiting should be provoked. The worm generally is voided dead within an hour or two. The bowels should be kept open with castor-oil, so long as the urine retains the violet smell which indicates the presence of turpentine in the circulation.

MR. A. J. SCHAFIRT, WASHINGTON, D. C.

In a number of cases, this pharmacist employed no preliminary provisions beyond forbidding the patient to breakfast the day of treatment, and giving him a full dose of Rochelle salts the preceding night. At ten o'clock in the morning, the following was taken at one dose:

685. R.	Granati radiceis,	$\mathfrak{z}\mathfrak{ss}$.	
	Peponis seminum,	$\mathfrak{z}\text{j}$	
	Extract. filicis ether.,	$\text{f.}\mathfrak{z}\text{j}$	
	Pulv. ergotæ,	$\mathfrak{z}\text{ij}$	
	Pulv. acaciæ,	gtt. ij.	M.

The pomegranate-bark and pumpkin-seed were thoroughly bruised, and, with the ergot, boiled in eight ounces of water for fifteen minutes, then strained through a coarse cloth. The croton oil was first well rubbed up with the acacia and extract of male fern, and then formed into an emulsion with the decoction. In each case, the worm was expelled alive and entire within two hours. No unpleasant effects followed. In each case, the worm was passed with the head firmly fastened to the side of its body, at about the widest part, from which it was with difficulty removed, and the worm was twisted and doubled into various knots.

DR. FORBES DICK, OF ENGLAND,

says that the male shield fern, properly administered, will expel not

only the segments but also the head of the tapeworm. He says that the adult dose of the Pharmacopœial liquid extract, as an efficient vermicide in cases of *tænia*, is not, as laid down in the *British Pharmacopœia*, from fifteen to thirty minims; nor, as usually administered to soldiers, one drachm; nor even, as Squire and others allow, eighty or ninety minims, but two drachms. The mode of administration should be as follows: A full mid-day meal may be eaten, and a little bread, with tea, at 5 p. m. At 10 or 11 p. m. a binder is to be applied to steady the stomach; and on lying down in bed the following is to be taken. (Neither prolonged fasting nor filling the big bowel with fluid seems to be necessary.

686. R.	Extracti filicis liq., Spiritus chloroformi, Pulveris acaciæ, Aquæ,	$\bar{3}$ ij \mathfrak{m}_{xxv} gr. xx ad $\bar{3}$ ij.	M.
---------	--	--	----

In the morning half an ounce of turpentine beaten up with the yolk of an egg should follow. The above large dose of the male shield fern is occasionally vomited, but he has failed to discover any deleterious effect contra-indicating its use. When vomiting happens, which is rare, on the following evening the same quantity should be taken in two doses, at half an hour's interval. The half an ounce of turpentine is more liable, especially in the tropics, to produce irritant symptoms. If active exercise be taken under a tropical sun after this dose, strangury follows, which there indicates the propriety of rest, to limit perspiration and prevent concentration of urine, and the drinking of bland fluids.

NOTES ON REMEDIES.

Alianthus Glandulosa. The powdered bark (gr. vij-xx) is an active anthelmintic in tape-worm.

Alumen, given in milk, is a remedy of some avail.

Aspidium Marginale. Dr. CRESSLER has obtained excellent results from the use of the oleo-resin of *Aspidium marginale* as a *tænifuge*. The oleo-resin is put in capsules of about grs. x each, and administered after a total abstinence of twelve hours from food. A dose of castor oil is administered about one or two hours afterward. The *Aspidium marginale* is related to the male fern.—*Unido Medica*, 1883.

Benzine has been tried with success by Dr. MOSLER and other German physicians. Dose, sixty to one hundred drops, in capsules. The after-effects are sometimes unpleasant.

**Brayera*, if well preserved, is highly recommended by Dr. J. MARTIN (*Medical and Surgical Reporter*, Oct. 20, 1883). The patient fasts for about twenty hours, and then takes one ounce of the herb, well boiled in one and a half pints of skimmed milk, in two equal doses, two hours apart.

Carbolicum Acidum has been used by Dr. RUMBOLD, of St. Louis.

687.	R.	Acidi carbolici,	gr. xx
		Aquæ destillatæ,	f. ℥viij.
One-third every two hours.			

Others have not found the worm affected by this agent.

Copaiva. Dr. S. CARO has recorded a case (*N. Y. Med. Jour.*, May, 1876,) of the expulsion of a worm by the balsam copaiva, when nearly all other reputed remedies had failed.

Creasotum. This is recommended by Dr. HENRY BRICKWELL, of England, in doses of gtt. j-ij, three times a day for a week, followed by a brisk purge.

Eupatorium Perfoliatum. The decoction of this plant is reported by Dr. H. S. WILKINS (*Medical and Surgical Reporter*, April, 1874,) to have expelled, promptly, a tape worm.

Ether Sulphuricus. Dr. LORTET, in 1867, suggested ether (f. ℥v at a dose), followed, in two hours, by olei ricini, f. ℥j. He and various observers since, report successful cases.

Filix Mas is an old and esteemed remedy. The fresh root must be used.

688.	R.	Extr. filicis maris æth.,	
		Pulv. rad. filicis,	āā gr. xxx.
Make thirty pills. Take ten at night, ten the next morning, and an hour after, the remainder. (KRAUS.)			

TROUSSEAU's method was to place the patient on a milk diet for a day; the next morning he prescribed ℥j of the oleo-resin, in four equal doses, a quarter of an hour apart; the third day, the same quantity, followed by a brisk purgative.

Granati Radicis Cortex. The pomegranate enjoys a wide reputation against tape worm.

689.	R.	Granati rad. cort.,	℥iij	
		Sodii bicarb.,	℥j	
		Aquæ,	Oj.	M.

Macerate six hours, and boil to Oss.

Take in the morning, fasting, in three equal doses, following the last with olei ricini, ℥j. Or:

690.	R.	Extr. granati rad. cort.,	℥v	
		Extr. fil. maris æther.,	℥ij.	M.

Make thirty capsules. Take them within two hours.

The bark of the pomegranate root is objectionable, on account of the abdominal pain it causes in many persons.

Kameela. This product of the *Rottlera tinctoria* is used in doses of ʒj-ij. It is an effective remedy, but liable to gripe.

691. R.	Extracti filicis maris,	ʒ iss	
	Pulveris kamalæ,	ʒ ij	
	Mucilaginis acaciæ,		
	Syrupi simplicis,	āā	f. ʒ ij
	Aquæ cinnamomi,		ad f. ʒ ij. M.

S.—Half to be taken at bed time, and the other half early in the morning.

Mr. J. B. LAWSON reports good results from this in the *Glasgow Med. Jour.*, January, 1884.

Kousso (*Brayera anthelmintica*). The reputation first enjoyed by this tæniifuge has somewhat diminished of late. It is disagreeable to take, and produces much intestinal distress. The dose is about ʒ ss. According to Dr. ALOIS MONTI, of Vienna, the compressed pastilles of kousso flowers, from Erlangen, are unsurpassed in usefulness. He has found no difficulty in their employment; and ten grammes he thinks infallible. The alkaloid, *koossine*, if quite pure, is more convenient, because the dose required is smaller; two grammes to five grammes, according to the age of the child, are sufficient. With these two, there is no need for any preparatory treatment; and if the head of the tænia does not come away the first time, they may be repeated the next day. Children take them as if they were sugar-plums; and they do not cause nausea or vomiting, nor are they followed by persistent diarrhœa. Dr. LABOULBENE, of Paris, attributes failure to the smallness of the dose. (1884.)

Nux Vomica and its alkaloid, *Strychnia*, has been reported upon favorably, both in Europe and America. Dr. A. R. KILPATRICK, of Texas, uses the following:

692. R.	Strychniæ sulphatis,		
	Arsenici,		
	Hydrarg. chlor. corrosiv.,	āā	gr. ʒ i.
			M.

For one pill. One three times a day.

Dr. MASSE publishes in the *Montpelier Medical*, 1877, some observations on a case in which he procured the expulsion of three tape worms by the administration of the tincture of *nux vomica*.

**Pelletierine* is highly recommended.

Peponis Semina, pumpkin-seeds, are unquestionably both a powerful and a pleasant remedy in tape worm. From 2 to 4 ounces of seeds should be pounded in a mortar, then mixed with milk or water, strained, and the fluid taken on an empty stomach. If the bowels do not act in a few hours, castor oil should be given. The dose should be repeated daily until the worm is expelled.

Pepsina. Full doses of pepsin have been found, by M. BOUCHUT, eminently successful in dislodging and destroying tape worms in children. (*Medical Examiner*, 1878.)

Potassii Picronitras. This has been recommended by some German physicians.

693. R. Potassii picronitratis, gr. xxviii
 Pulv. jalapæ, ʒ iʒ
 Extr. glycyrrhizæ, q. s. M.
 Make 30 pills. One three times a day.

It is efficient, but apt to be followed by general malaise.

Saoria, the ripe fruit of the *mæsa picta*, an Abyssinian tree, has long enjoyed a good reputation as a tænicide. The dose is ʒ ss-j, as :

694. R. Saoriæ, ʒj
 Zingiberis, ʒ iʒ
 Cassiæ, gr. xv
 Aquæ, Oj. M.

Strychnia. See *Nux Vomica*.

Terebinthinæ Oleum. Recommended by Dr. WATSON (see above) in large doses. Others prefer ʒ ss, twice a day, continued for some time. Sometimes it is combined with castor oil :

695. R. Olei terebinth.,
 Olei ricini, āā f. ʒ ss
 Mucilag. acaciæ, q. s. M.
 Make an emulsion. For one dose, fasting.

ASCARIS LUMBRICOIDES—LUMBRICI—ROUND WORMS.

DRS. MEIGS AND PEPPER, PHILADELPHIA.

These writers consider that as the diagnosis of lumbrici is nearly always doubtful, it is best never to risk the irritating vermifuges unless worms have actually been passed. In slight and doubtful cases they prefer small quantities of *worm-seed oil* to anything else. It may be given in doses of gtt. iv, to children of two years, three times a day for three days, followed on the fourth day by some castor oil or syrup of rhubarb. The following is a good formula to disguise its taste :

696. R. Olei chenopodii, gtt. lx-f. ʒj
 Mucilag. acaciæ, ʒ iʒ
 Syrupi simplicis, f. ʒj
 Aquæ cinnamomi, f. ʒ iʒ. M.
 Give a dessertspoonful three times a day for three days, and repeat after several days.

In giving the *pink root*, it is best to combine it with some cathartic substances, as in the following formula of Dr. G. B. Wood :

- | | | | | | |
|------|----|-------------------|----|-----------------|----|
| 697. | R. | Spigeliæ, | āā | 3 ^{ss} | |
| | | Sennæ, | | 3 ^{ij} | |
| | | Magnesiae sulph., | | 3 ^j | |
| | | Fœniculi, | | Oj. | |
| | | Aquæ ferventis, | | | M. |

Macerate for two hours in a covered vessel. Dose, a tablespoonful, to a child two years old, once or twice a day, or every other day, so as to procure two or three evacuations in the twenty-four hours.

The *fluid extract of spigelia and senna* is also a very acceptable combination.

PROF. J. LEWIS SMITH, M. D., NEW YORK.

The long-popular American remedy for worms is the pink root,
spigelia marilandica:

698. R. Extracti spigeliæ fluidi, f. $\frac{3}{4}$ j
 Extracti sennæ fluidi, f. $\frac{3}{4}$ ss. M.
 One teaspoonful to a child of three to five years.

Or:

699. R. Extracti spigeliæ et sennæ fluidi, f. ʒj
Santonini, gr. viij. M.
One teaspoonful to a child of five years.

“This is probably the best anthelmintic that can be employed for the destruction of the round worm in uncomplicated cases, and it is also very useful in treating the *ascaris vermicularis*.”

In some cases of protracted intestinal disease, attended by an increased and vitiated secretion, there is nothing so good as *turpentine*. The following is Dr. CONDIE'S formula:

- | | | | | |
|------|----|---------------------------|----|----------|
| 700. | R. | Mucilaginis acaciæ, | | f. ʒij |
| | | Sacchari albi, | | ʒj |
| | | Olei terebinthinæ, | | |
| | | Spiritûs ætheris nitrosi, | āā | f. ʒ iij |
| | | Magnesiae, | | ʒj |
| | | Aquæ menthæ, | | f. ʒj. |
- A teaspoonful to a child of five years.

DR. EUGENE BOUCHUT, PARIS.

- | | | | |
|---------|----------------------------|---------|----|
| 701. R. | Hydrargyri chloridi mitis, | gr. ij | |
| | Santonini, | gr. jss | |
| | Sacchari lactis, | gr. xv. | M. |

To be given in one dose, as a vermifuge, in a teaspoonful of honey, to an infant two years old.

DR. GIORDANO, FLORENCE.

702. R. Sulphuris loti,
Magnesiæ,

℥ss
℥j. M.

Divide into four powders. One to be given morning and evening to children who have passed worms, in order to prevent their reproduction.

DR. G. CALDERWOOD, ENGLAND.

Of all remedies for lumbrici, undoubtedly the best is *santonin*. In regard to its administration, this writer remarks (*British Medical Journal*, February, 1875,) that it should always be given for three or four days continuously before a purgative is prescribed. To give it with, or only once before, a cathartic, does not accomplish the work completely. It acts toxically, and must have time to do its work. The worms are passed like other fæcal matter, when dead; and so long as they are alive, no amount of purging will bring them all away.

DR. F. T. ROBERTS, ENGLAND.

This author prescribes an aperient, limits the diet to liquids for a day, and then employs santonin in doses of from one to three grains every morning for two or three days, in sugar or syrup, or made up into lozenges, or with gingerbread. It is more efficacious when mixed with castor oil; and Küchenmeister advises that from two to four grains be dissolved in ℥j of the oil, and ℥j taken every hour until it acts. Troches of santonin, each containing half a grain of the principle, can now be obtained from all druggists.

DR. ROBERTS BARTHOLOW.

This author says that the infusion of quassia is one of the most effective injections for the destruction of the ascarides vermicularis, which infest the rectum. The stomach administration of simple bitters undoubtedly hinders the development of intestinal worms, probably by correcting a morbid state of the mucous membrane. In the treatment of intestinal parasites, therefore, much good is derived from the use of bitters, administered with the view of restoring normal digestion. *Eucalyptus* may be used by injection. *Carbolic acid* is an efficient, but unsafe parasiticide, when injected into the rectum. A solution of the *tincture of iron* is one of the numerous remedies used to destroy this form of parasite. As their development is favored by the anæmic state, it is good practice to conjoin with any local treatment the internal

use of iron, notably the syrup of the iodide. He uses *santonin* as follows: A laxative in the morning, fasting through the day, a dose of santonin and calomel at bedtime, a senna draught on the following morning. He sometimes uses troches of santonin:

703. R. Santonin, ℥ ss
 Sugar,
 Tragacanth,
 Orange flower water, āā q. s. M.
 To make four hundred and eighty troches. Each troche contains half a grain of santonin.

The enema of aloes is very efficient.

704. R. Aloës, ℥ ij
 Potass. carbonat., gr. xv
 Mucil. amyli vel decoct. hordei, ℥ x.

Considerable medicinal doses of hydrocyanic acid are very fatal to round worms.

Salicylic acid has been used successfully in the removal of ascarides, by local application and by internal administration. An enema of water should precede the salicylic acid solution, which may be composed of borax and salicylic acid with some glycerin:

705. R. Acid. salicylic, ℥ ss
 Sodii biborat., ℥ ss
 Aquæ, Oj. M.
 S.—Warm, and administer the whole amount at one time. For a young child, this quantity should be reduced one-half or more.

Myrol, a new antiseptic, has been used successfully.

DR. SCHILDOWSKY, OF RUSSIA,
 has used *iodoform* successfully. He gives, to an adult, one grain with ten grains of bicarbonate of soda three times a day, and a quarter of a grain to a child.

ASCARIS VERMICULARIS—OXYURIS—THREAD WORMS— SEAT WORMS.

DR. T. SPENCER COBBOLD, LONDON.

This helminthologist calls attention to the true residence of the seat worm. (*British Medical Journal*, February, 1874.) He maintains that the entire length of the colon is the territory inhabited by the threadworm, the cæcum itself constituting the parasite's true head.

quarters. Knowing this, therefore, to be the case, and likewise understanding, as we now do, how these entozoa originally gain access to their human bearers, it is clear that we are in a position to treat the threadworm helminthiasis on thoroughly rational principles. Injections cannot be conveniently employed to wash out the contents of the arch of the colon and cæcum, and therefore we must, in the main, rely upon the operation of medicines taken by the mouth. He therefore employs injections only for the purpose of dislodging such oxyurides as have been driven down to the lower bowel.

When it comes to the question of the choice of drugs, a great variety of medicinal agents may be used with good effect. Some seem to answer better in particular cases than in others. Speaking generally, he gives a preference to active *saline cathartics* repeated for several days in succession, followed by the use of cold water enemata. Small doses of chloric ether and sulphate of iron are eminently serviceable additions, and the same may be said of aloes, with or without assafoetida. In cases where these drugs are objected to, he has employed various active mineral waters with good results. Bitters of all kinds are useful, and patients who object to salines will swallow any reasonable amount of the infusion of gentian and other pure vegetable tonics. As a rule, oxyurifuges should be administered in the form of copious draughts; and the oftener they are repeated, for a short interval, the more effective will they prove in the end. The rapid passage of the drugs through the bowel, will, if several times repeated, carry most of the parasites sufficiently low down to be within reach of the clysters; and the prevention of the return of the parasites to the upper part of the colon, is one of the practical points worth looking to. The employment, therefore, of enemata, after the exhibition of remedies by the mouth, cannot fail to prove beneficial.

MR. WILLIAM DATE, LONDON.

706.	R.	Infusi quassiae,	Oj	
		Sodii chloridi,	ʒij.	M.
Use as an injection, once a day.				

When this fails to dislodge them completely, give :

707.	R.	Ferri sulphatis,		
		Quiniæ sulphatis,		
		Pil. aloës cum myrrhâ,		
		Pil. galbani compos.,	āā	gr. j.
One such pill thrice daily.				

This treatment, in a fortnight, will completely destroy the parasites.

PROF. W. H. VAN BUREN, M. D. NEW YORK.

708. R.	Acidi carbolici, Glycerinæ, Potassii chloratis, Aquæ,	gtt. x-xx ℥j ad saturandum f. ℥ viij. M.
---------	--	---

Use as an enema, in thread worms, *oxyuris vermicularis*.

It is sometimes necessary to increase the amount of carbolic acid.

DR. F. WEBB, MICHIGAN.

This practitioner has stated (*Michigan Medical Journal*, November, 1871,) that in cases of ascarides, the *hyposulphite of soda*, in doses large enough to produce slight catharsis, given three times a day, has availed to expel ascarides when other vermifuges and various enemata have failed.

Along with whatever other treatment is used for thread-worms, it is important to anoint, with mercurial or other parasiticide ointment, the anus and adjacent parts, from time to time, in order to destroy the ova of the worm, which are frequently deposited externally to the rectum.

DR. GUICHON, PARIS.

709. R.	Santonini pulveris, Resinæ jalapæ, Chocolate,	℥j gr. ij ℥j. M.
---------	---	------------------------

Divide into thirty lozenges. Give one in the morning, on an empty stomach, to an infant two years old; two or three to older children.

710. R.	Aloës barbadensis, Potassii carbonatis, Decocti amyli,	℥ ss gr. xv f. ℥ x. M.
---------	--	------------------------------

To be given as an injection, in ascarides of the rectum.

DR. AITKEN.

711. R.	Pulv. scammon. comp., Pulv. aromatici,	gr. iv gr. v. M.
---------	---	---------------------

For one dose, at night.

The day after taking this purgative, the patient ought to be kept on low diet, with solid food. On the day succeeding the purgative, from 5 to 10 grains of the ethereal extract of santonin, or from 3 to 4 grains of santonin itself, may be given. Injections of infusion of quassia, or

of steel and quassia and aloes, or a solution of common salt in gruel, or a similar quantity of lime-water, are useful in the expulsion of the *ascaris vermicularis*. Half an ounce of the compound decoction of aloes, taken in the morning, fasting, once or twice a week, and 3 ounces of infusion of quassia, may be taken every morning that the aloes are not taken.

NOTES ON REMEDIES.

Adeps. Anointing the anus and sphincter with lard destroys the ova, and is a useful adjuvant, as it is believed that the ova are only set free in the rectum near the anus.

Asarum Canadense. As an internal remedy, the Canada snake-root has been found valuable by Dr. S. S. BOND, of Philadelphia, (*Medical Times*, September, 1874,) and others. It may be given in infusion, or syrup, or children may chew it as they would liquorice-root.

Aqua Calcis, in injections, has been commended.

Argenti Nitras, gr. j-iiij to aquæ \mathfrak{z} j, as an enema.

Ferri Chloridi Tinctura, \mathfrak{z} ss to aquæ Oj, is said to be a very efficient enema.

Fuligo Ligni. An infusion of soot has been strongly recommended in *ascarides*, by Dr. HEWSON, of Philadelphia. It is made by adding a cupful of soot to a pint of boiling water, and straining. Half or a fourth of this quantity should be thrown up the rectum daily, for one or two weeks.

Hydrargyri Chloridum Corrosivum, gr. j to aquæ Oj, has been used in obstinate cases by Dr. NIEMEYER.

Hydrargyri Chloridum Mite. M. VALLEIX was accustomed to anoint the anus and lower bowel with :

712.	R.	Hydrarg. chlor. mitis,	\mathfrak{z} iv	
		Axungiae,	\mathfrak{z} vj.	M.

Dr. E. F. WALKER (*New York Medical Journal*, 1878,) says that a large dose of calomel (gr. x-xxx) gives better results than any of the ordinary drugs in dislodging seat-worms.

Hydrargyri Unguentum. Anointing the anus and sphincter with mercurial ointment is very useful in destroying the worms there located and their ova. It may also be introduced on a bougie.

Potassii Sulphuretum, in the proportion of \mathfrak{z} ss-j to aquæ Oj, is an efficient injection.

Sulphur. A dose of sulphur taken every morning aids in dispelling the parasites.

Terebinthinæ Oleum, \mathfrak{z} ss to milk Oj, may be used as an enema.

TRICHINÆ, TRICHINOSIS.

The general treatment of trichinosis is symptomatic, as up to the

present we have discovered no specific vermicide for the *trichina spiralis*. The suggestion of *ergot* has been made by Dr. RHODE, of Berlin. He had a case of trichinosis in which severe bleeding of the nose occurred, and in which he prescribed extract of *secale cornutum* as a styp-tic. The hemorrhage was immediately arrested, and with this, rapid improvement of the general symptoms also occurred. This result led him to prescribe *ergot* in other cases of the disease, and in all instances distinct improvement followed. He believes therefore that we have, perhaps, in *ergotin*, a means of treatment which, without having any marked effect on the human economy, may prove fatal to *trichinae* and their offspring.

Picric and carbolic acid, as well as benzine, which have all, at times, been recommended, have proved entirely useless in pigs, and probably would be equally so in the human subject. Probably the best treatment in the early stage, to cleanse the intestines from the parasite, would be purgative doses of calomel, $\mathfrak{D}\text{i}-\text{ij}$, on the plan recommended above for seat-worms by Dr. E. F. WALKER.

Dr. FERRER (*Gazz. de les Hospital Valencia*) has cured a case of trichinosis with alcohol. He commenced with six and increased to nine ounces daily, in sugared water, in the intervals of feeding. The cure was effected in eighteen days.

JAUNDICE.

CATARRHAL JAUNDICE.

PROFESSOR HUGO ENGEL, OF PHILADELPHIA.

If a case has an acute and violent beginning, and the diagnosis can be early made, he administers to the patient a fever-mixture, as, for instance:

713. R.	Spiriti mindereri, Spiriti ætheris nitrosi, Tinctur. radicis aconit,	f. $\frac{3}{4}$ v f. $\frac{3}{4}$ j ℥xxiv.	M.
---------	--	--	----

Of this, he lets the patient take a tablespoonful every three hours in half a tumblerful of sweetened orange-water, lemonade, Seltzer or Apollinaris water, until the fever has ceased. Occasionally the dose

of the aconite has to be diminished. If the patient is very plethoric, if the fever is high, the pulse full and somewhat hard, and the face looks flushed, he bleeds him at the arm, taking from twelve to sixteen ounces of blood. If there is much nausea, or even if there is little, but the tongue heavily coated, or if there is occasional vomiting, he administers a tablespoonful of mustard in a bowl of tepid water, or a hypodermic injection of apomorphia. He cautions against tartar-emetic, to be used for the same purpose. If the fever is intermittent (very, very rare), or remittent (frequently), and above 101° , he prescribes quinine, a few large doses (twenty grains) once every twenty-four hours.

If the case has an insidious beginning, he puts the patient at once upon the same treatment that he employs in the acuter cases after the fever has ceased.

But no matter how the case is ushered in, as soon as he is positive as regards the diagnosis, he orders the local application of from six to twelve European (or twenty-four to forty-eight American) leeches. If there is no special point douloureux or local tenderness (if there is, they are applied as near as possible to it), he has the leeches applied in the hypogastric region, just above the umbilicus and slightly more to the right side of it.

The diet consists of milk, beef tea (bouillon), oatmeal gruel (strained), bread soup, and the like, given at regular intervals. With the bowels or the excretion of the kidneys (save as by the medicine above referred to), he interferes only when necessity demands such a procedure. Either oleum ricini or Glauber salt answers in such a case the first, and acetate of potash the second purpose.

When the acute symptoms have ceased, or have never been present, his treatment is as follows: After the leech-bites have healed, he knows of no better way to assist the system to remove the obnoxious bile than by acting on skin and kidneys, and by applying a large fly-blister 7x8 inches over the seat of the lesion. Often he applies a second, and sometimes a third blister, always waiting for the skin first to heal; but more frequently, after the second or third blister, he has half a drachm of mercurial ointment rubbed into the abdominal wall every day or every second day, certainly taking care not to cause salivation. But it is amazing, the quantity of such an inunction that the skin will gradually absorb in these cases, as well as in others, where absorption is our aim, without bringing about salivation. He is convinced that many a patient has derived incalculable benefit from this treatment.

Then he prescribes two remedies: acetate of potash, one drachm four times daily in a large quantity of water, his usual prescription being:

714. R.	Potassii acetatis,	f. ℥ iij	
	Tinctur. nucis vomic.,	℥ xlviij	
	Glycerin.		
	Syrup rubi idari,	āā	q. s. ad. f. ℥ vj. M.

Two teaspoonfuls of this medicine, to which the glycerin and the raspberry syrup are added, only to improve the taste of it, and the nux vomica to allay any irritability of the stomach, contain exactly one drachm of the acetate of potash; this dose is administered in a large tumblerful of water (if possible, Apollinaris) four times daily.

Further, he orders two teaspoonfuls of Carlsbad salt to be taken every morning on an empty stomach. But to insure the real effect of this almost specific remedy, certain conditions have to be fulfilled. First, for this purpose he never uses the effervescing salt; he employs either the imported, genuine, or, with still greater preference, the artificial Carlsbad salt. As it is difficult to dissolve, as it should be taken lukewarm, in a large quantity of water and in a certain manner, he advises his patients as follows: They are to take a goblet of the size of a large beer-glass; into this they put the two heaped teaspoonfuls of artificial salt, leaving the metal spoon in the glass to prevent it from cracking; then the glass is *slowly* filled with boiling water, and the solution continually stirred. If these last two precautions are not observed, a part of the mixed salt will recrystallize, and not dissolve in the quantity of water prescribed. Then the glass is covered with a saucer, and kept over night in the bedroom of the patient, who in the morning on awakening, say, for sake of illustration, seven o'clock, once more stirs his Carlsbad water, and then drinks the sixth part of it. Ten minutes later he takes the second sixth, and, if he is now able to leave his bed, as most are, he rises, and while washing and dressing himself, he sips the third sixth. Then he goes into his yard, or in inclement weather into the largest and best-aired room in the house, and walking up and down he drinks every ten minutes another sixth, so that within about half an hour the Carlsbad water has been swallowed. The patient must then wait half an hour for his breakfast.

To assist nature still more, he has the patient take a steam-bath every night on going to bed. The room in which the patient is to take the bath is brought to a temperature of 74°, as determined by the thermo-

meter—to prevent chilling. When the pain has ceased and the fæces have had a green color for a week, without interruption, he gives the patient every fifth or sixth day, in the evening, from five to ten grains of blue mass, followed in the morning by the usual dose of Carlsbad salt.

ROBERTS BARTHOLOW, M. D., PHILADELPHIA.

Excellent results have been obtained by this practitioner in malarial and catarrhal jaundice by the use of *manganese*:

715. R. Chinoidin, ʒj
 Manganesii sulphatis exsiccatae, ʒij. M.
 For twenty pills. One three times a day, in malarial jaundice.

716. R. Fel bovini purificati, ʒj
 Manganesii sulphatis exsiccatae, ʒij
 Resinae podophylli, gr. v. M.
 For twenty pills. One three times a day, in catarrhal jaundice.

In similar cases he also finds the following of great use:

717. R. Quiniæ sulphatis, ʒij
 Ferri sulphatis exsiccatae, ʒj
 Acidi arseniosi, gr. j. M.
 To make twenty pills. One thrice daily, in malarial jaundice.

PROF. WILLIAM PEPPER, M. D., PHILADELPHIA.

In simple jaundice—acute congestion of the liver—the indications are to check vomiting, to relieve the congestion, and to start the bile. To accomplish the first, give hydragogue cathartics, elaterium, bitartrate of potash, etc.; or, if the stomach is too irritable, administer an enema of croton oil, gtt. ij, in emulsion. The congestion of the stomach and liver can be relieved by counter-irritation, as by a large blister over the epigastrium. This is very effective to arrest the vomiting. By the mouth, give the following:

718. R. Hydrargyri chloridi mitis, gr. iij
 Pulveris opii, gr. ij
 Bismuthi subnitrat, ʒijss. M.
 Make six powders. One to be given every three hours.

This combination powerfully aids the mercurial in relieving the hepatic congestion, and stimulating the gastro-intestinal glands. The diet in such cases should be restricted to milk. (*Medical and Surgical Reporter*, 1875.)

DR. GEORGE HARLEY, LONDON.

In *pyæmic* or *septicæmic* jaundice, great caution must be used not to fatally depress the already broken-down system. If a mercurial is indicated, it should be given in the form of grey powder, and when the alimentary canal has been unloaded it is advisable to give tonic germicides, such as quinine and its congeners. If the patient is sufficiently strong, salicylic, benzoic and carbolic acids may be substituted. In *congenital jaundice*, the mercurial had better be given to the mother or wet-nurse than to the baby. In *contagious jaundice*, after clearing out the stomach and bowels by an active emetic followed by a rapidly acting purgative, germicides and quinine should be given. Iron should be avoided. For the head symptoms use chloride of ammonium, bromide of ammonium, acetate of ammonia and the carbonate of ammonia, either given separately or combined in pairs dissolved in camphor mixture. For *malarial* jaundice he uses mercurials freely, in combination with quinine, salicine or ipecacuanha. But under the most favorable influences, many of the cases that have long been subjected to some of the worst forms of malarial influences, succumb, from the constitution having become, as it were, thoroughly and irretrievably undermined by the poison. In *catarrhal* jaundice the free administration of emetics is recommended. In jaundice from *permanent obstruction*, pig's bile will supply an important element to the system, and tend to ward off the fatal end. Fresh bile, taken directly from the gall bladder of the freshly-killed pig, is filtered through very porous filter-paper to free it from mucus. It is then as rapidly as possible evaporated to dryness at a temperature not exceeding 160° Fah. As soon as dry it is ready for use. Five grains are equal to about one hundred grains of liquid bile, and ten grains is the dose, to be given at the end of stomachal digestion.

PROF. A. GUBLER, FACULTÉ DE MEDECINE, PARIS.

719. R. Aloës socotrinæ,
 Gambogiæ,
 Hydrargyri chloridi mitis, āā gr. xv
 Syrupi, q. s. M.

Divide into ten pills. Order one or two a week, to keep the bowels soluble, in icteric disease of the liver.

MIDDLESEX HOSPITAL, LONDON.

720. R. Pilulæ hydrargyri, gr. xxx
 Digitalis pulveris,
 Scillæ pulveris, āā gr. v. M.

Divide into ten pills. One morning and evening, in icterus, to eliminate the coloring matter of the bile from the blood.

NOTES ON REMEDIES.

Acidum Benzoicum has recently attracted attention as a remedy in jaundice due to suppression of the biliary secretion.

Acidum Nitro-Muriaticum Dilutum is useful both internally and externally, in the form of a bath. It is contra-indicated if there be inflammatory action present.

Aloes has been used with benefit in jaundice.

Ammonii Chloridum, in doses of gr. xx every five hours, has, in the hands of Dr. ANSTIE, restored the biliary secretion, in a few doses, when it has been suppressed by a powerful nervous shock or mental perturbation. In his opinion, it is one of the most powerful of all biliary functional restoratives.

Colchicum, in small doses, combined with mild mercurials, or soap, or alkalies, or with magnesia or the neutral salts, according to circumstances, has been prescribed by Dr. COPLAND with benefit. It must be continued for a considerable time, and carefully watched. If there be much debility, or if it produce depression, it should be combined with camphor.

Copaiba. Dr. B. J. MASSIAH (*British Medical Journal*, April, 1879,) says that the cholagogue action of this drug is not sufficiently recognized. In hepatic torpidity, cirrhosis and jaundice, it is a valuable aid. The dose may be ℥j thrice daily.

Ether, from its power of dissolving cholesterine and its anti-spasmodic properties, has been recommended as an internal remedy in jaundice depending on the presence of gall-stones.

Euonymin, in chronic hepatic affections, has many supporters. In 1884 it was highly spoken of at the Paris Society of Therapeutists.

Hydrargyri Chloridum Mite. Jaundice sometimes yields rapidly to a calomel purgative (gr. v,) followed by a saline, which, however, prove useless when the affection is connected with obstruction by gall-stones, or organic diseases of the liver.

Magnesii Sulphas. The following formula is sometimes useful in severe forms of jaundice :

721.	R.	Magnesii sulphatis,	℥ ss-j	
		Magnesii carbonatis,	gr. xv	
		Spiritus ammonii aromatici,	℥ xxx	
		Aquæ,	f. 3 x.	M.
Make a drink. Take thrice daily.				

**Podophyllin* is often a very reliable remedy in jaundice.

Potassii Chloras, combined with carbonate of soda, is useful in chronic or asthenic jaundice, accompanied by a torpid state of the liver.

Potassii Sulphas is recommended by Dr. A. T. THOMPSON, as more useful than other saline purgatives in jaundice.

Sanguinaria, in the form of the powder (gr. ij-v,) or the infusion, has been employed with advantage in jaundice and torpor of the liver.

Sodii Benzoate. In contagious jaundice (the violent form) Dr. HARLEY recommends this drug, combined with salicylate of quinine, or quinine itself, when the temperature is high.

Taraxacum, combined with small doses of colchicum, has been found serviceable in jaundice depending upon hepatic disease.

**Cathartics*. Dr. GROSS states that in jaundice depending upon hepatic derangement, after bilious evacuations have been produced, an active aperient, every second or third day, for the space of ten days or a fortnight, should never be omitted, so as to carry off the remains of the disease, and prevent a relapse. This is a simple but successful practice which should not be neglected.

Enemata. Dr. KRULL, of Mecklenburg, extols enemata of cold water, 60° Fah., as an excellent remedy in catarrhal jaundice (*Berliner Klin. Wochenschrift*, No. 12, 1877). He uses one to two litres once a day, to be retained as long as possible.

TORPIDITY AND FUNCTIONAL DERANGEMENT OF THE LIVER.

CHARLES MURCHISON, M. D., LL.D., F. R. S., LONDON.

In the "Croonian Lectures" of 1874, this eminent authority discussed the functional derangements of the liver and their treatment. Commencing with *diet*, he recommends one chiefly of stale bread, plainly cooked mutton, white fish, poultry, game, eggs, a moderate amount of vegetables, and weak tea, cocoa, or coffee. In severe cases, potatoes, rice, sago and fruits have to be given up; milk suits some, not all. The principal meal should be in the morning, and the quantity not beyond the needs of the system. All alcoholic drinks should be prohibited. *Cholagogues*: Great advantage is derived from aperient medicines; salines are the best, as sulphate of magnesia, sulphate of soda, tartrate of potash and soda, Friederichshall, Pullna, or other saline water; as to the benefit of mercury, our author says that the skepticism of the most doubting physician would be removed should he unfortunately find it necessary to test its efficacy in his own person. Calomel is the form he prefers. Podophyllin he regards as less certain in its action and more liable to cause griping. Colchicum is a useful adjunct to other aperients. *Taraxacum*, he thinks, has no specific

action, but is useful as a mild aperient, and may be advantageously combined either with alkalies or mineral acids. *Alkalies*: These are next in importance. One grain of carbonate of lithia, or of carbonate of ammonia, nearly equals a grain and a half of carbonate of soda, or two grains of carbonate of potash. It is well to suspend their use occasionally as, when long continued, they derange gastric digestion. *Chloride of ammonium* has a great and well-deserved reputation in hepatic congestion and other functional derangements of the liver:

722. R. Ammonii chloridi purificati, $\bar{\zeta}^{ss}$
 Aquæ menthæ piperitæ, f. $\bar{\zeta}$ iij. M.
 A dessertspoonful three times a day.

It may be combined with either alkalies or mineral acids. The last-mentioned remedies, our author considers of minor value, in these complaints. *Tonics* are apt to disagree in many cases of functional hepatic derangement. Neither iron nor quinine acts well. Gentian, chiretta, cascarilla, serpentaria, and nux vomica, are preferable to preparations of iron and cinchona. *Arsenic* suits some patients. *Opium* is contra-indicated, except in those functional hepatic troubles which arise in diabetes.

NOTES ON REMEDIES.

Ammonii Chloridum is worthy of trial when there is want of biliary action.

Euonymin, from the *Euonymus atropurpureus*, and *Iridin*, from the *Iris versicolor*, have been reported upon by Prof. RUTHERFORD, of Edinburgh. He gives of the former, gr. j-ij; of iridin, gr. iv, in pill form. There is little difference in their action, both being feeble intestinal stimulants, increasing the functional activity of the liver, and acting mildly and without depression. (*Brit. Med. Jour.*, June, 1879.) For ordinary and simple cases, the following is a convenient form:

723. R. Ex. euonymi fluid., f. $\bar{\zeta}$ iij
 Syr. aurantii, f. $\bar{\zeta}$ j. M.
 S.—One teaspoonful before breakfast.

If the debility and torpor are somewhat extreme, a dose may be taken before each meal, in which case, half a teaspoonful is almost always amply sufficient.

Hydrargyri Pilulæ. ABENERTHY'S practice, in bilious derangements, of giving, almost indiscriminately, 5 grains of blue pill at night and a senna draught in the morning, is still continued by many practitioners.

**Ipecacuanha*, in small and long-continued doses, is frequently productive of great benefit in functional derangement of the liver.

Iridin is a cholagogue aperient, gentler in its action than podophyllin. It is very useful in a sluggish state of the bowels, arising from torpidity of the liver, or when the stools are pale, particularly in the interval of overt attacks in gouty persons.

Leptandrin is useful in torpidity of the liver, in which case it is said to be a valuable adjunct to podophyllin and iridin.

Nux Vomica and *Strychnia* are valuable hepatic stimulants :

724. R. Extracti nucis vomicæ, gr. viij
 Extracti colocynthidis comp., gr. xlvij. M.
 For twenty-four pills. One three times a day, in chronic hepatic congestion.

Oleum Olivæ. Dr. E. P. TOWNSEND (*Country Practitioner*, October, 1879,) states that, for many years, he has used olive oil, ℥j, two or three times daily, in cases of chronic hepatitis and torpidity of the liver, with or without jaundice, with undoubted good effect.

**Podophyllin* is now much employed in torpor of the liver.

Sanguinaria has been used with advantage in torpidity of the liver, the powder (gr. ij-v), or infusion, being preferred to the tincture.

Senna, in infusion (f. ℥j-ijj), either alone or combined with sulphate of magnesia (℥ij-iv), or tartrate of potash (℥ij-℥ij), is an efficient purgative in bilious derangements.

**Taraxacum* is a valuable remedy in torpor of the liver.

Turkish Baths are sometimes of great advantage as excitants of the liver.

PYROSIS.

Pyrosis, or water-brash, is not unfrequently connected with organic disease of the stomach; but its most typical form is as a functional disorder, often due to the use of certain ingesta, among which, oat-meal, cheese, and tobacco, may be mentioned. One of the most useful medicines is *bismuth*, which may be combined with compound kino powder.

The following are some of the prescriptions of

PROF. PIORRY, OF PARIS.

725. R. Sodii bicarbonatis, ℥ iss
 Aquæ destillatæ, āā f. ℥j
 Syrupi aurantii florum, gtt. j. M.
 Olei anisi,
 For one dose, in pyrosis.

726. R. Quiniæ sulphatis, gr. xxij
 Pepsinæ, ℥vss
 Extracti absinthii, q. s. M.

Divide into forty pills. Two before each meal, in functional derangements of the stomach, particularly pyrosis, flatulence, and gastralgia, after eating.

727. R. Sodii bicarbonatis, gr. xiv
 Tincturæ columbæ, f. ʒj
 Aquæ destillatæ, f. ʒiij
 Syrupi gentianæ, f. ʒj. M.

To be taken during the day, in the same manner as the above formula.

728. R. Magnesiae, gr. iss
 Bismuthi subnitratæ, gr. iv-vij
 Opii pulveris, gr. ss. M.

For one dose; to be taken before eating, in acid dyspepsia and with pyrosis.

NOTES ON REMEDIES.

Acidum Carbolicum has been employed with success by Dr. PODMORE JONES (*Practitioner*, November, 1868,) in the following formula :

729. R. Acidi carbolici, gr. j
 Alcoholis, f. ʒj. M.

Twenty-five drops in a wineglassful of water, an hour before each meal.

Acidum Gallicum is of benefit in cases accompanied by organic disease of the stomach, or by disease of the liver.

Acidum Sulphurosum, in the hands of Dr. LAWSON (*Practitioner*, September, 1868), has produced the best effects, in doses of f. ʒss-j, thrice daily, shortly before meals, in plain distilled water.

Alum, in doses of ℥j thrice daily, in the form of an electuary, has been strongly recommended.

Argenti Nitras is an excellent remedy.

Argenti Oxidum has been employed with marked benefit. It must not, in these cases, be combined with creasote, as the combination causes violent decomposition and the development of great heat, and even flame.

**Benzoin*. A fluid drachm of the tincture, with mucilage, is said to be efficacious in many cases.

Bismuthi Subnitras, combined with opium, seldom fails to afford relief.

**Kino Pulvis Compositus* is said by Sir T. WATSON to be an admirable remedy in this affection, in doses of gr. x, thrice daily.

Liquor Ferri Nitratis, in doses of gtt. x-xv, thrice daily, is useful in those cases where tonics and astringents are indicated.

Lupuli Extractum has been found serviceable.

Magnesiae Carbonas is occasionally an effectual remedy.

**Nux Vomica* is very useful in many cases. Dr. BARLOW recommends the following formula :

730. R.	Extracti nucis vomicæ, Argenti nitratis, Extracti lupuli,	gr. issj gr. ij gr. xij.	M.
---------	---	--------------------------------	----

Divide into six pills. One thrice daily.

**Opium*, in combination with gentian, frequently affords speedy relief.

VOMITING.

In the treatment of this distressing symptom, it is of the utmost importance to distinguish between vomiting from *cerebral*, from *uterine*, and from *gastric* irritation, as the management of the three is essentially different. That from uterine sympathy is not considered here. The principal anti-emetics will be found in the notes on remedies:

DR. S. G. WEBBER, OF MASSACHUSETTS.

Sometimes only a large amount of food taken at one time excites vomiting: then it is sufficient to resort to frequent feeding, giving a very small quantity each time, a mouthful or a spoonful every fifteen or thirty minutes; thus the stomach never contains a large mass of food requiring considerable muscular exertion to roll it about, and by its weight or bulk exciting the reflex irritability of the nerve centres. Many times, however, this is not enough; the stomach requires more complete rest, and the best treatment is to withhold all food and medicine; sometimes a few hours' rest is enough, again it requires two or three days; then it will be necessary to use nutrient enemata. Where there has been much vomiting, thirst may be very annoying to the patient; small lumps of ice held in the mouth will relieve this, and generally do not cause vomiting. After the stomach has had sufficient rest, it is best to commence feeding by the mouth, with caution, giving a little frequently. Milk and lime water, equal parts, a teaspoonful every half hour, should be first tried; if well borne, the amount can be increased gradually. It is a mistake to increase the quantity too rapidly.

NOTES ON REMEDIES.

Acidum Arseniosum, in the form of Fowler's solution, is advised by Dr. RINGER, in the morning vomiting of drunkards.

**Acid Carbolic* has remarkable sedative effects, and will arrest the vomiting of acute stomachal and intestinal disorders; it may be advantageously

combined with bismuth, in an emulsion. BARTHOLOW recommends the following formula :

731. R.	Acidi carbolici,	gr. iv	
	Bismuthi subnitrat.,	ʒ ij	
	Mucil. acaciæ,	ʒ j	
	Aq. menth. pip,	ʒ iiij.	M.

S.—A tablespoonful every two, three or four hours.

Acid Carbonic, in the form of an effervescing draught, is often an effective remedy.

Acid Sulphurous, in doses of ℥xxx, thrice daily, has been employed by Dr. DRYSDALE, with success (*Lancet*, July 24, 1869,) in constant vomiting and eructation.

Acidum Hydrocyanicum, given in an effervescing draught, is a very efficient remedy in vomiting arising from any morbid conditions.

Acidum Tartaricum, in effervescing draughts, sometimes succeeds in allaying vomiting.

Ammonii Citras has been recommended in the vomiting attendant on ulcer of the stomach, but Dr. W. BRINTON considers that the salts of ammonia do more harm than good in these cases.

Ammonii Chloridum is said to relieve the vomiting attendant on scirrhus of the stomach more promptly and uniformly than any other remedy.

Bismuthi Subnitrates, especially when combined with hydrocyanic acid, often speedily allays vomiting in connection with stomach diseases.

Bromides are serviceable in cerebral vomiting, and in cholera infantum in children, and in some cases of reflex vomiting.

Calcis Aqua sometimes succeeds when hydrocyanic acid and other remedies fail. It has less influence on sympathetic vomiting than on that connected with functional derangement of the stomach.

Calomel, in very minute doses, will stop vomiting in cholera infantum and in other intestinal disorders in children.

**Chloral* is highly useful in vomiting of sea-sickness, cholera, and reflex vomiting.

**Chloroform*, in doses of a few drops, is often very useful in obstinate vomiting.

Cinnamomum, in aqueous infusion, is often of benefit in continued nausea and vomiting.

Cocculus Indicus, according to PHILLIPS, is the great rival of nux vomica in nervous vomitings, when attended with a dull and heavy pain in the head, and intolerance of light and sound.

Creasotum is considered by many as an overrated remedy in vomiting. Dr. RINGER considers the principal cause of its failure is due to its being given in too large doses. He obtains the best effects by adding just sufficient creasote to water to make it taste distinctly but not strongly of the medicine, and administering of this a dessertspoonful frequently. He finds it effectual, given in this manner, in allaying nausea and retching, though some consider it to have less power over actual vomiting.

**Ipecacuanha* is considered by Dr. RINGER to have few equals in checking some forms of vomiting. In numerous instances he has found drop doses of the wine, administered every hour or thrice daily, according to the urgency of the case, check vomiting of pregnancy, the morning vomiting of drunkards, the vomiting which accompanies general weakness after acute diseases, and that of acute catarrh of the stomach in children. That form of vomiting after meals in which there is no nausea or pain, or even discomfort, is quickly stopped by these small doses of ipecacuanha. Even in vomiting from cancer of the stomach, he has sometimes succeeded with it after the failure of other remedies. Dr. C. D. PHILLIPS states (*Practitioner*, November, 1869,) that its beneficial operation is clearly discernible in most cases of continued and obstinate retching or vomiting where the stomach is not primarily affected, but disturbed by sympathy with some other organ or part of the body, whether the primary affection be acute or chronic.

Iodine. Dr. T. I. GAUNT (*Am. Jour. Med. Sci.*, April, 1883,) has for a number of years been employing the compound tincture of iodine in drop doses in nearly all forms of emesis, and he reports thirteen cases of the most varied character, in all of which vomiting was promptly arrested by its use.

Koumiss, wineglassful doses every half hour, has been found effectual in obstinate cases.

Nitrite of Amyl and *Nitro-glycerine* are said to be very effective in the vomiting of sea-sickness.

Nux Juglandis. Dr. EDWARD MACKEY has found (*Practitioner*, December, 1878,) a tincture of walnuts very efficacious in hysterical vomiting, that from dyspepsia, pregnancy and anomalous cases. His formula is :

732. R.	Fresh walnuts,	$\bar{3}$ xxx	
	Alcohol,	$\bar{3}$ xij	
	Water,	q. s.	M.
Distill $\bar{3}$ xvi. Dose, f. $\bar{3}$ j every four hours.			

Sodii Carbonas is beneficial in vomiting arising from acidity of the stomach.

Nux Vomica, in the form of the tincture, is recommended in the vomiting of pregnancy. Dr. DOUGLAS POWEL (*Practitioner*, November, 1868,) says that the tincture, in doses of \mathfrak{m} x, will often arrest the vomiting of phthisis. Dr. BARLOW recommends, in vomiting connected with malignant disease of the stomach, the following formula :

733. R.	Extracti nucis vomicæ,	gr. j	
	Extracti conii,	gr. xij.	M.
Make six pills. Take one three times a day.			

Blisters. A small blister over the epigastric region frequently effectually checks vomiting arising from functional or nervous disorder of the stomach, and also that consequent on fevers.

Ice in lumps, or minute quantities of iced brandy and water, or champagne,

given every hour, is useful in severe vomiting attendant on pregnancy or uterine disease.

Leeches. Dr. TILT states that in vomiting connected with uterine affections, after the failure of ordinary measures, he has, on several occasions, seen it suddenly checked by the application of six leeches to the pit of the stomach, although there was no sign of inflammation there, and the patient's debility was such as not to justify loss of blood. The same procedure sometimes succeeds in idiopathic vomiting.

V. DISEASES OF THE URINARY SYSTEM.

Albuminuria — Bright's Disease — Diabetes Insipidus — Diabetes Mellitus — Dropsy (Ascites, Anasarca) — Functional Urinary Disorders (Incontinence) — Hæmaturia — Nephralgia — Nephritis — Uræmia.

ALBUMINURIA.

Recent researches have demonstrated very conclusively that albuminuria may and very frequently does exist when there is no disease of the kidney. In fact, some authorities go so far as to say that there is a "physiological albuminuria," noted in perfectly healthy persons.

DR. CALVIN ELLIS

enumerates among the various causes capable of producing albuminuria 1. The irritant poisons. 2. Disorders of the circulatory system, as affections of the heart, various fevers, as intermittent or yellow fever, peritonitis, pneumonia or pleurisy. 3. Blood poisoning with or without fever, as ischæmia, acute rheumatism, typhoid or scarlet fever, diphtheria, variola, measles. 4. Affections or diseases of the nervous system or brain, as convulsions, mania, affection of the spine or spinal cord. 5. Changes in the blood, as in suppuration, syphilis, scurvy, cholera. 6. Lead poisoning, tumors or cancerous affections, goitre, interference with the functions of the skin by cold, burns, from varnishing the skin with gum arabic.

DR. CHATEAUBOURG, OF PARIS,

concludes that, 1st. Albumen is found in the urine of the majority of healthy persons, more or less abundantly, and transient in its character. 2. Rest in bed has a clearly-marked influence in diminishing the amount of albumen excreted. 3. Bodily fatigue greatly influences the production of physiological and transient albuminuria. 4. Intellectual labor augments with most people the quantity of albumen in the urine. 5. Cold bathing exerts considerable influence in increasing physiological

albuminuria. 6. Sexual excitement and menstruation manifestly affect albuminuria in the healthy. 7. Albuminuria is as frequent in children as in adults, but the quantity of albumen excreted is less. 8. Digestion, if accompanied by rest, does not exert much influence upon physiological albuminuria.

DR. F. DE HAVILLAND HALL, LONDON.

In the treatment of acute albuminuria, such as is a frequent sequela of scarlet fever, this author condemns the usual procedure (in England) of "hard purging and sweating" (*Practitioner*, August, 1876.) He has achieved much better results from the following plan:

Directly any albumen was detected in the urine, the patient was ordered the *perchloride of iron*, and was allowed no solid food except a little bread and milk, and as much water as he liked to drink; this treatment, together with keeping the skin gently acting, sufficed in the majority of cases, but in a certain number the urine was almost suppressed, and in some there were uræmic symptoms. Whenever either of these contingencies occurred, he forbade all food for twelve hours, the child to have nothing but water and a drink made as follows:

734.	R.	Acid tartrate of potash,	$\overline{3j}$	
		Lemon syrup,	f. $\overline{3j}$	
		Water,	Oj.	M.

If, at the end of this time, the kidneys were beginning to act, he allowed a little milk, but not more than a pint in the twenty-four hours; if, however, the uræmia continued with little or no urinary secretion, he persevered with the water and bitartrate of potash; and in severe cases, nothing else has been given for thirty-six hours. Dry cupping, mustard poultices over the loins, and a purgative, were the only additional remedies employed.

The explanation of the good effects of abstention from solid food, and especially meat, during the course of acute desquamative nephritis, is, that if a patient is entirely deprived of nitrogenous food, the work of the kidneys is lessened and the urine is rendered less irritating, and the mild diuretic action of the bitartrate of potash seems to be useful.

He sums up the treatment of acute Bright's disease in the following words:

1. Milk and water with arrowroot, no solid food.
2. Mild diuretics, such as the citrate or bitartrate of potash with a free supply of water.
3. The skin kept just moist.
4. A daily evacuation of the bowels.

DR. ROBERTS BARTHOLOW, OF PHILADELPHIA.

Some important additions to the therapeutics of albuminuria were suggested by this writer in a paper before the Phila. Co. Med. Soc., in 1883.

He would use *nitro-glycerine*, in doses of one minim of a centesimal solution (one minim of the pure drug to 100 minims of alcohol), every three or four hours, in acute cases immediately after the subsidence of the acute symptoms, and in chronic cases at all periods. *Chloride of gold and sodium* is indicated in the sub-acute and chronic cases, especially the latter. The earlier it is given the better, if structural changes are to be prevented or arrested. The good effects to be expected from it will depend necessarily on the extent of the damage already inflicted on the kidneys.

The usual dose is one-twentieth grain, twice a day, but this may be much increased, if necessary. At the outset, one-tenth grain may be given; in a week the dose should be lowered to one-fifteenth grain, and after a month the regular dose of one-twentieth grain should be steadily pursued, with occasional intermissions. Indigestion, gastralgia, and colic pains, nausea or diarrhoea, are occasionally caused by it; and, if so, the quantity administered must be reduced. It is usually borne without any discomfort.

DR. T. LAUDER BRUNTON, LONDON.

This practitioner states that where true albuminuria is present, the indications are:

1. To remove the venous congestion of the kidneys. When this depends on cardiac disease, digitalis is called for; on pregnancy, etc., the prone position lessens it.
2. To diminish the flow of blood to the kidneys. To fulfill this, warm clothing, warm baths, diuretics and purgatives are required.
3. The anæmia consequent on the drain of albumen must be removed by iron, and light, digestible food.

WILLIAM AITKEN, M. D., LONDON.

735. R. Acidi gallici,
Acidi sulphurici diluti,
Tincturæ lupuli,
Infusi lupuli,

3 ij
f. 5 ss
f. 3 j
f. 5 vj. M.

A tablespoonful three times a day.

Said by the author to check the waste of albumen, especially in the more acute and recent forms of albuminuria.

J. M. DA COSTA, M. D., PHILADELPHIA.

736. R.	Ammoniae muriat.,	℥ ij	
	Tincturae ferri chloridi,	f. ℥ iss	
	Aquae,	f. ℥ vj.	M.

A tablespoonful thrice daily, in anæmic albuminoid degeneration of the kidneys or liver.

This author considers the *muriate of ammonia* one of the best agents we possess to counteract the tendency to albuminoid degeneration in any organ.

NOTES ON REMEDIES.

Acidum Gallicum, in doses of gr. x, thrice daily, has been successfully employed.

Acidum Tannicum, in from gr. xxx-lx per day, has been advised in the anasarca accompanying albuminuria.

**Antimonii et Potassii Tartras* is recommended by Dr. BARLOW in the acute forms of albuminuria. He considers it indicated by the nature of the affection, and calculated to equalize the circulation, subdue the inflammatory action, and restore the functions of the skin. But he does not employ it to the exclusion of other remedies, as moderate venesection, hydragogue cathartics, the warm bath, and the application of large linseed-meal poultices to the loins.

Aqua. Dr. W. N. DICKINSON, of London, recommends drinking large quantities of pure water, as the safest and best diuretic in this disease. (*Medical Times and Gazette*, June, 1876.)

Arsenicum. In some cases of albuminuria, seemingly dependent on imperfect digestion, Dr. T. L. BRUNTON has derived excellent results from Fowler's solution, gtt. iij, with meals. (*Practitioner*, June, 1877.)

Belladonna will, in the opinion of Dr. HARLEY, prove very serviceable in chronic albuminuria, provided that the kidney has not passed into the degenerative state bordering on fatty degeneration.

Cantharis, internally, has, it is said, been employed with benefit in purely chronic cases, but must be used with great caution.

Chloral. Dr. THOS. WILSON says in the *Brit. Med. Jour.*, 1882: "The attention of the profession does not yet seem to have been generally drawn to the effect of chloral in removing albumen from the urine, and also the existing œdema." He adds some striking cases.

Digitalis is indicated where there is cardiac complication, but must be used cautiously.

Eucalyptus Globulus. In albuminuria and post-scarlatinal nephritis, Dr. WILLIAM ANDERSON, of Brooklyn, has used with good results the elixir

and fluid extract of eucalyptus (*Proceedings of Kings Co. Medical Society*, Aug., 1879.) He expresses himself very decidedly as to the benefit he has witnessed from it. He gives gtt. x of the fluid extract, in water, every four or six hours.

**Ferri Iodidum* is often beneficial, particularly in old, broken-down constitutions.

Ferri Chloridi Tinctura is one of the most generally useful preparations of iron in albuminuria, possessing diuretic as well as chalybeate properties.

Fuchsin or *Rosanilin Hydrochloride*, has been found to check the secretion of albumen promptly, by M. BOUCHET, of Paris, (*Gaz. des Hop.* 1879.) He gives gr. j, thrice daily, in aromatic water; in one case gr. iss daily for five days was followed by complete recovery.

Oxygen, in inhalation, has attracted considerable attention, but the results have been unsatisfactory.

Potassii Bitartras in acute and chronic cases is an efficient diuretic, $\bar{3}$ j to aquæ Oj.

Potassii Sulphas, gr. x-xx, two or three times a day, is said by Dr. DICKINSON to be sometimes invaluable as a habitual laxative.

Potassii Sulphuretum has been recommended in the following formula, to establish free diaphoresis :

737. R.	Potassii sulphureti,	gr. v	
	Tincturæ guaiaci ammoniatæ,	f. $\bar{3}$ ss	
	Liquoris ammoniæ acetatis,	f. $\bar{3}$ ss	
	Aquæ,	f. $\bar{3}$ j.	M.

For one dose, to be taken at bedtime, followed by a pint of hot whey.

Scoparius. In scarlatinal albuminuria, according to Dr. S. D. BELL (*Transactions Pennsylvania State Medical Society*, 1876,) scoparius has in his hands yielded "invariably the most flattering results." He uses it in the form of decoction, made by boiling half an ounce of the tops in a pint and a half of water down to one pint. Of this a tablespoonful to a wineglassful was given every four or six hours, according to the age and severity of the symptoms.

Sodii Carbonas has been recommended in albuminuria, on the ground that alkalescence is a necessary condition of the blood; that the free alkali is soda; that when this constituent is deficient, the result is coagulation of blood in the capillary vessels and ensuing inflammation, and the potash or soda (uncombined or as carbonates,) taken into the stomach, have the power of rendering the urine alkaline and dissolving fibrine. The following formula may be employed :

738. R.	Sodii carbonatis,	$\bar{3}$ ij	
	Liquoris potassæ,	f. $\bar{3}$ ij	
	Decocti chondri crispi,	f. $\bar{3}$ viiij.	M.

A tablespoonful every two hours, in milk. Tartrate of iron may be added when anæmia is very marked.

Tiglii Oleum is one of the best hydragogue cathartics which can be given in albuminuria, as it produces copious evacuations with less sickness and discomfort than elaterium.

BRIGHT'S DISEASE.

ALFRED L. LOOMIS, M. D., NEW YORK.

This author is convinced that the diaphoretic and cathartic plan of treatment is wrong, and its apparently good results deceptive, and prefers the following :

In the acute stage of parenchymatous nephritis, whatever may have been the exciting cause, the patient must be kept in bed, in a large, well-ventilated apartment, with a temperature of 75° Fah.; milk should be his only article of diet. Dry cups should be applied over the kidneys, followed by poultices of digitalis leaves; and the infusion of digitalis leaves should be administered at the rate of half an ounce every two hours, for twenty-four hours, after which it may be continued at longer intervals, so long as uræmic symptoms are urgent. When the flow of urine commences, the digitalis should be discontinued, and diluent drinks freely administered to keep up the diuresis. If renal secretion is not restored within twenty-four hours after the commencement of this treatment, warm baths, or hot-air baths, should be used in addition; and if the symptoms remain urgent, hydragogue cathartics may be given.

In the latter stages, those commonly known as chronic Bright's disease, the plan of treatment is essentially tonic. Iron and cod-liver oil are the two great remedial agents, and should be daily administered, if they do not disturb the stomach. Milk should be the principal article of diet, and the patient should seek a warm and equable climate. The milk may be taken cold or hot, from a half pint to a pint at a time. Wine, in moderation, may be taken with the food. No depletory remedies should be employed in this stage, except in times of an emergency, when, from some sudden renal congestion, the remaining healthy portion of the kidney is impaired, and active uræmic symptoms present themselves. At such times, a hot-air bath, hydragogue cathartics, or stimulating diuretics are to be employed, but with great caution.

J. M. DA COSTA, M. D., PHILADELPHIA.

For the treatment of an acute case, following scarlatina:

739. R. Pulveris ipecacuanhæ compositi,	gr. iij	
Potassii nitratis,	gr. v.	M.

For one powder. To be taken at night.

And,

740. R. Liquoris ammoniæ acetatis,	f. ʒ iij	
Spiritus ætheris nitrosi,	℥ xx	
Syrupi tolutani,	℥ xxxvij	
Tincturæ digitalis,	℥ iij.	M.

For one dose, thrice daily.

The skin should be made to act freely by means of hot baths and a few drachms of Rochelle salts, administered on alternate days. The diet should be mild and unirritating, all stimulants being avoided.

741. R. Tincturæ ferri chloridi,	f. ʒ ss	
Acidi acetici,	f. ʒ ss	
Mix and add,		
Liquoris ammoniæ acetatis,	f. ʒ v	
Curacoæ,	ʒ ij.	M.

Teaspoonful thrice daily.

This recipe, a modification of BASHAM'S mixture, is useful in chronic Bright's disease, and in all those cases in which the union of a tonic and diuretic effect is desired.

DR. N. M. BASKELL, OF MISSOURI.

According to this author (*Medical and Surgical Reporter*, March 29, 1884,) mercurials are very useful in acute Bright's Disease. He always uses them, and has never yet salivated a patient. His usual plan, if there is much œdema, with impeded heart-action, is to give the following recipe:

742. R. Hydrarg. chlor. mit.,	gr. ss	
Sodii bicarb.,	gr. ijss.	M.
Ft. chart No. j.		

S.—One powder twice a day.

In conjunction with this he gives night and morning:

743. R. Potass. bitart.,	ʒ i. vel jss	
Jalapæ pulv.,	gr. viij	
Gambogiæ pulv.,	gr. j.	M.
Ft. chart No. j.		

Should this not cause an elimination of the watery elements rapid enough, and there is danger of uremic coma or apoplexy from retardation of the blood, he adds :

744. R. Pilocarpin muriat., Sacchari lactis, Ft. chart No. x.	gr. j grs. x. M.
---	--------------------------

S.—One three times per day.

For the action of the heart and system, he uses the following tonic, varying the ingredients to suit the exigencies of the case :

745. R. Potass. iodid.,
Tinct. digitalis,
Syr. squills,
Liq. ammon. acetat.,
Fl. ext. phytolac. dec.,
Syr. sarsap. comp.

These ingredients, given according to the age and condition of the patient and the effect desired.

Under this treatment his acute cases have always improved rapidly. He does not continue the mercurials long after the dropsy is subdued. But he has frequently found it necessary to return to them when the symptoms manifested themselves, and always with benefit.

In the chronic condition, the most that can be done is to keep the bowels regular and build up the system with tonics. For this purpose, he finds nothing better than the comp. jalap powder, the alterative mixture mentioned above, and a teaspoonful of the ferri nitrogeniz. comp. This to be taken three times per day, after meals. Occasionally he gives the powders of pilocarpin three or four days in succession.

DR. FELIX VON NIEMEYER, PROF. UNIVERSITY OF TUBINGEN.

The loss of albumen from the blood being the immediate cause of most symptoms of the disease, the most important task of the physician is to cover the loss of albumen by a diet rich in protein substances and by appropriate medication. Soft-boiled eggs, milk, strong meat broths and roast beef, in as large quantity as the patient is able to digest, are probably the best preventives of the dropsy. Besides this, a moderate quantity of beer or good wine should be prescribed, as by the use of these the waste of tissue is retarded and nutrition promoted. Quinine and iron are the most suitable medicines. Our author has obtained most brilliant results, where all other treatment had failed, by putting

the patients upon an exclusive diet of *milk*. They took no medicine whatever, but drank five or six pints of cow's milk daily. After the "cure" had been continued in this manner for about five weeks, some of the patients, who, prior to the treatment, had been in the most wretched condition, had got rid of their dropsy, recovered an appearance of health, and regained so much strength as even to be able to perform hard labor.

If the above measures fail in averting or allaying the dropsy, active diaphoresis is strongly to be recommended. Patients in an advanced state of dropsy often rid themselves of it completely in a few weeks by the daily use of a hot bath, of a temperature of 80° to 100° F., followed by sweating for two hours in woolen blankets. Debilitated patients sometimes, however, suffer so much from this treatment as to compel its discontinuance.

Whatever the theoretical objections to the employment of diuretics may be, yet, in desperate cases, recourse should always be had to them. Squills and other stimulating diuretics must not be employed without the utmost caution. But there are certain salts, particularly *cream of tartar*, which are decidedly beneficial in their effects. The free use of buttermilk, conjoined with the employment of cream of tartar and small doses of Dover's powders, will prove serviceable.

The drastic cathartics should only be resorted to in cases of extreme need, since the patient is liable to be affected by them, and since, by their persistent use, the digestion becomes impaired. Those most frequently employed are colocynth and

746. R. Elaterii,

gr. $\frac{1}{8}$ -ss.

For one dose, as occasion requires.

FRANCIS SIBSON, M. D., D. C. L., ETC., LONDON.

In a review of the therapeutics of Bright's disease, (*British Medical Journal*, February, 1877,) this eminent practitioner lays down the following principles:

The employment of *narcotics* in any form, in cases of Bright's disease, directly adds poison to poison, of the like kind, and narcotics are, therefore, to be strenuously forbidden in every case of Bright's disease. This axiom does not rest upon physiological considerations only, which, if rightly interpreted, never can lead us wrong, and are, therefore, the true peculiar guide in the treatment of disease; for we have, also, di-

rect evidence of the deadly effect of narcotics in Bright's disease. Death has occurred again and again, from the subcutaneous injection of moderate doses of morphia in those affected with Bright's disease. Neither opium, chloral, chloroform, hyoscyamus, belladonna, nor stramonium, should ever be exhibited.

For the extreme restlessness, we must give the *bromides* of ammonium and potassium, and Hofmann's anodyne ether; and a few drops of ether may safely be inhaled, as well as in the cases with excessive distress of breathing. The bromide may be pushed to large and extreme doses, and their effect ought to be carefully watched, so that if drowsiness be induced by it, in addition to the poison of the disease, iodide of potassium, coffee, and rousing and anti-narcotic means, may be freely resorted to.

The danger to combat is from the narcotic poison in the blood, which is not eliminated by the kidneys. To meet this indication, the patient should be encouraged to take exercise in the open air, and to keep his faculties active; they should be roused by strong coffee and tea, by the society of friends, and by the business of every day life.

When the respiration is greatly distressed and suspended by turns, the proper treatment is steadily to eliminate the poison, and, at the same time, to rouse the powers, either by a small quantity of tea, strong coffee, or whisky; by stimulants externally; and by maintaining the power by means of iron and food. These cases are very formidable, and generally die; but not unfrequently, the patients recover from the emergency, and even live for years. When coma and sopor are present, the hot air bath and external stimulants, the strongest coffee, and a treatment analogous to that of deep opium poisoning, but differing according to the necessities of the case, must be pursued; and this sometimes succeeds.

BELLEVUE HOSPITAL, NEW YORK.

The following is a summary of a plan of treatment recommended at Bellevue Hospital, New York:

Diet. This class of patients should abstain, as much as possible, from meat. The opinion has been expressed that the excessive animal diet accounts for the great prevalence of the disease in America. Milk should be substituted for meat, and should be associated with lime. Butter may be used; eggs, if they agree, and fresh fish in the morn-

ing. Fried fats should be carefully excluded, but cream may be taken without stint. Vegetables and fruits are *always* good, but those should be selected which contain the least amount of woody fibre. Rice and potatoes, therefore, may be used; but asparagus, turnips, cabbage, and, notably, beans, which contain woody fibre in large quantities, should be assiduously avoided. Onions may be eaten with impunity, and are rather beneficial.

For the Anæmia. Iron should be administered from first to last, and, by preference, the tincture of the chloride. This preparation is assimilated with difficulty; hence should not be given alone, but combined with nux vomica, and to this, spirits of nitre may be added, to assist the determination towards the kidneys. For example, ten drops of the tincture of the chloride of iron, ten drops of tincturæ nucis vomicæ, and one drachm of sweet spirits of nitre, may be given three times a day. Cod-liver oil increases the red corpuscles of the blood, because it is digested by the liver, and the product enters into them as an ingredient. The irritability of the stomach may make it troublesome to take, but it should be relied upon as much as in the treatment of phthisis.

To Combat the Disease Itself. We have one agent which may be regarded as a specific against increase of connective tissue in the body, wherever the interstitial inflammation may occur, and that is the bichloride of mercury. It should be given in small doses, $\frac{1}{20}$ of a grain is the usual amount, and should be combined with a diuretic, to make it act upon the kidneys. For example, $\frac{1}{20}$ of a grain of the bichloride, one grain of digitalis, and one grain of quinine, may be given three times a day, with the result of producing a specific action upon the kidneys, and will *raise* the specific gravity of the urine.

Attention to the Condition of the Skin will materially assist the embarrassed kidneys, and to do this, we may have recourse to two things. If excessive œdema is present, the pressure produced shuts off the circulation to a great extent, and prevents removal of the fluid by diaphoresis. It is much better then to make punctures in the distended skin of the legs, and let the water drain away at once. No apprehension need be had with reference to this trifling operation, if the limb, when the punctures have been made, is wrapped with cloths wet in a solution of carbolic acid in water, to which has been added essence or oil of cinnamon. The latter is to correct the smell of the carbolic acid,

and is also equally antiseptic. The second thing is, to rub the patient all over once a day with sweet oil. If extra diaphoresis is desirable, it can be best obtained by placing a blanket in an empty bucket, and pouring hot water upon it, for in this way much less water is required, and then wringing it out and quickly applying it around the body and covering it with a dry blanket. The skin should be well oiled before the blanket is applied.

NOTES ON REMEDIES.

Aconitum. Dr. RINGER recommends that after scarlet fever, the temperature be taken night and morning, and on the slightest rise being noted, aconite be given to prevent nephritic inflammation.

Aqua. On the use of water in the early stages of Bright's disease, see (p. 432.)

Arsenicum. According to Dr. S. V. PAP, (*Wiener Medicinische Presse*, No. 13, 1875,) arsenic is useful in most cases. Its beneficial influence is manifested in the lighter forms more rapidly, certainly, and distinctly, than in the more severe forms, with a greater proportion of sugar than four per cent. In the former, the sugar may entirely disappear, which is but rarely the case in the latter. The diminution, or the disappearance of the sugar secretion, also takes place with a mixed diet. The maxim remains undisputed, however, that the amylaceous foods should, at least temporarily, be limited, as far as possible. After the cessation of the use of arsenic, months may pass before the disease returns.

Auri Chloridum. Dr. BARTHOLOW calls especial attention to the use of the salts of gold in chronic Bright's disease, granular and fibroid kidney. He has observed remarkable improvement follow the use of the chloride of gold in these affections. They are best given in pill form and in small doses, $\frac{1}{30}$ — $\frac{1}{20}$ of a grain three times a day. They are, of course, not adapted to the acuter forms of Bright's disease.

Cannabis Indica is spoken of by Dr. RINGER, as having a high reputation in acute and chronic Bright's disease, especially when the urine is bloody.

Elaterium is a useful purgative when there is much dropsy, but it is best to combine it with hyoscyamus, as it sometimes causes exhausting diarrhoea, especially if given after symptoms of uræmic poisoning have set in.

Euphorbia Corollata ("spurge," "wandering milk-weed," "go quick," etc.,) is highly recommended by Dr. RANSOM (*Proceedings Illinois State Medical Society*, 1884,) in the dropsy of Bright's disease, to remove the anasarca.

Ferri Chloridi Tinctura. As a chalybeate diuretic, this form of iron is not surpassed by any.

Gallicum Acidum. Dr. H. C. WOOD, Jr., states in certain forms of Bright's disease, when there was an abnormally large secretion of highly albuminous urine, he has found gallic acid to lessen very materially the albumen.

Guaiacum, in tincture or decoction, is regarded by Dr. COPLAND as the best diuretic in Bright's disease, especially when the skin is cool as well as dry.

Ipecacuanha, in the form of Dover's powder, (gr. v-viij, thrice daily,) is not only a useful diaphoretic in Bright's disease, but also allays pain and irritability, especially if assisted by a warm bath every other evening, or oftener.

Jaborandi has been suggested on theoretical grounds in this disease. A largely increased elimination of urea by the skin is one of the results of its administration.

Potassii Iodidum is said by Dr. RINGER to prove sometimes of signal service, even when the dropsy has been extensive and the urine very scanty.

Scoparius, in decoction, combined with nitric ether, is recommended by Dr. BARLOW, when the renal secretion is very deficient. A pint of the infusion may be given.

Tannicum Acidum has been proved by experiment to diminish the loss of albumen in Bright's disease. Its value, however, in a remedial sense, is yet uncertain.

Diuretics. Drs. BRIGHT, OSBORNE, WATSON, and others, condemned the use of diuretics, particularly those which are stimulating, in inflammatory dropsy attendant on granular disease of the kidneys. Dr. CHRISTISON believes, however, that they can be used without risk, and advises the following formula :

747. R.	Pulveris digitalis,	gr. j-ij	
	Potassii bitartras,	ʒj-ij.	M.
	Ter in die, ex aq.		

Diuretics may sometimes be employed, with good results, by the *endemic* method.

Skimmed Milk. From six to eight pints should be used daily, and the diet restricted to it. The good effect should be seen inside of a week.

EXTERNAL MEASURES.

Camphorated Vapor Baths. At the Queen's Hospital, Birmingham (England), these baths are used with good effects for the relief of Bright's disease. They are also said to produce very free perspiration, and are given every evening, in the following manner : The patient is seated upon a cane-bottomed chair, with a large blanket pinned around his neck ; half an ounce of camphor is placed upon a tin plate, under the chair, and above the flame of a small spirit-lamp, by the heat of which the camphor is slowly vaporized. This plan of treatment is also employed in some cases of convalescence from acute or sub-acute rheumatism, when the action of the skin is defective, and when some pain and stiffness of the joints remain.

Electricity. Dr. H. J. PRATT, of Colorado, reports a case of Bright's disease,

where the galvanic current applied over the dropsical abdomen, and general faradization, resulted in relief of the dropsy, and in a diminution of the amount of the albumen.

DIABETES INSIPIDUS, POLYURIA. POLYDIPSIA.

PROF. A. BOUCHARDAT.

The Hygienic Treatment. This is always essential to success. The excretions should be regular and complete; the bladder thoroughly emptied every six hours, at least. Exercise should be taken regularly, and to a considerable amount; chilliness, after it, being carefully avoided. A sponge-bath should be taken every morning, followed by long and quick flesh rubbing. The following may be added to the bath, to increase the action of the skin:

748. R. Potassii carbonatis, Essentiæ lavendulæ, Tincturæ benzoin.,	3 ijss gtt. xxx f. 3j. M.
---	--------------------------------------

The diet is of the utmost importance.

Food should be taken sparingly, well cut, and thoroughly masticated. Tomatoes should be forbidden, and also asparagus and beans, if their use cause renal pain or slight deposit in the urine. Meats of all kinds may be allowed, but only moderately; still more sparingly must we allow eggs, fish, lobsters, shrimps, shell-fish, and old cheese; milk is often useful. Almost all seasonable vegetables are good, particularly spinach, endive, lettuce, artichokes, celery, carrots, parsnips, and potatoes. The last-named are useful, and should take the place of some of the bread with the meal. Radishes may be advantageously used daily. The daily use of cresses or salad is indicated. All fruits, if the stomach bear them well, may be served daily. Olives, almonds, walnuts, hazelnuts, and pistachio nuts, must be served moderately. Good chocolate is allowable. If coffee provokes urinary secretion, it may be advised. Brandy and liquors should be avoided; very little beer allowed; and for alcoholic drink, a light white or red wine, diluted freely with water. White, sparkling wines are contra-indicated, as well as gaseous drinks, as soda-water. On waking, going to bed, and with meals, sufficient watery drink should be taken to give about three pints of urine. This

watery drink should consist of pure water, or, better still, a litre of water in which is dissolved one or two teaspoonfuls of powdered Seignette salt (tartrate of potash and soda.)

DR. C. P. B. CLUBBE

recommends the use of electricity. He reports the case of a woman who passed from eighteen to twenty pints of urine per diem. It was light colored, very low specific gravity, and contained no sugar. All the drugs recommended for diabetes were tried, without result. She was then ordered electricity (faradaism) to be applied over the region of her kidneys every day, for about twenty minutes at a time. Under this treatment she improved very much, and the improvement continued.

DR. J. M. DA COSTA, OF PHILADELPHIA.

One of the very few cases of complete cure from this generally fatal disorder, is reported by this author in the *Transactions of the College of Physicians of Philadelphia*, 1875. The treatment was by *ergot*, f.ʒj of the fluid extract being given at the outset, thrice daily, increased, in a few days, to f.ʒij, three times a day. The cure was accomplished in about two months, and was permanent.

In the volume of the *Transactions* for the following year, Dr. JAMES TYSON reported a case in which he administered *ergot* with temporary benefit, but not with actual success. He then prescribed *gallic acid* (gr. xv) four times a day, with rapid and he believed permanent improvement.

PROF. A. TROUSSEAU, PARIS.

In some instances, Professor TROUSSEAU derived advantage from *valerian*; it should be administered in full doses, *extracti valerianæ*, gr. xv, thrice daily.

Dr. BOUCHARD considers that the action of *valerian* is to spare waste of tissue; and in support of this view, he mentions that certain Indians of Lower California and Mexico are accustomed to go through a course of it for a month, before they enter upon a severe expedition, so that they may be better able to bear fatigue. He gives it in frequent small doses, and gradually increases the amount taken, until he has, in some cases, reached a dose of thirty grammes in twenty-four hours. (*Medical Times and Gazette*, May 23d, 1874.)

DR. W. MULLER.

The assistant of Prof. ECKSTEIN, in the clinic at Kiel, has published some important observations on the use of salicylate of soda in the treatment of diabetes. It appears that full doses of the salt cause the sugar to disappear entirely, but this is not a permanent result, for, after a while, the sugar returns. A very great tolerance of the remedy exists in this disease, and large doses may be taken with impunity. BOUCHARDAT, in commenting on these observations, states that he has not had good results from this treatment. BARTHOLOW concludes, however, that since the remedy is well tolerated, and during its exhibition in considerable doses the sugar disappears, it may be used with advantage at critical times to stop the waste.

DR. HENRY KENNEDY, LONDON.

This gentleman (*Practitioner*, February, 1878,) relates five cases all substantially cured by *dilute nitric acid*. He commences with f.ʒj to a quart of water, this amount to be taken in divided doses daily; and if necessary increases the acid to the daily amount of f.ʒvj. He has had no failures with it.

NOTES ON REMEDIES.

Atropia has been tried, but though it diminishes the urine, the thirst continues. (Dr. RENDUER, France.)

Auri Chloridum is considered by BARTHOLOW to be a promising remedy. The dose is from $\frac{1}{30}$ to $\frac{1}{15}$ grain.

Belladonna. Dr. JOHN M. SCUDDER, of Cincinnati, (*Eclectic*,) claims that belladonna merits the name of a specific in this disease.

Creasotum has been employed with some advantage.

Ergota has been highly praised by Dr. DA COSTA (above).

Gallicum Acidum, combined with opium, is spoken of by Dr. H. C. WOOD, Jr., as the most generally successful remedy in this disease.

Jaborandi. This drug was used in this disease by Professor T. LAYCOCK, of Edinburgh, in 1875. Its efficacy is doubtful.

Lactic Acid has been used with varying success in the treatment of diabetes. Dr. FOSTER reports some cases apparently decidedly benefited, and Dr. OGLE gives an account of two cases in which no good results were attained. Cases have been lately reported in which the patients were improved by the use of lactic acid, but on the whole the utility of this agent in diabetes must be held to be as yet *sub judice*. Disappointment in the use of lactic acid is frequently experienced from the poor quality of the drug.

Milk. Cures by the milk diet have been reported ; in consequence of the larger proportion of lactic acid which it contains, buttermilk is more especially indicated in diabetes.

Muscarine has been used with apparent success.

Nitricum Acidum. Almost a specific (see above.)

Opium, with tannic or gallic acid, is valuable.

Oxygen is indicated, and has been successfully employed. The internal administration of chalybeate medicines, or mineral waters, should accompany the inhalations of oxygen. Pure oxygen is not necessary ; an admixture with three parts of air will suffice, and the inhalation should be made morning and evening.

Rhus Aromatica has been introduced by Messrs. PARKE, DAVIS & CO., Detroit, on the recommendation of a number of practitioners of its effects in checking polyuria. The dose is f. ʒss-ʒ of the fluid extract in sweetened water, several times a day. It is stated to act promptly in arresting the excessive secretion of urine.

Valeriana was introduced by TROUSSEAU. Its effects are valuable. (P. 445.) The doses must be large and increasing to ʒj-ij of the powder thrice daily. Failures are attributed to its too timid use.

DIABETES MELLITUS.

DR. AUSTIN FLINT, JR., OF NEW YORK.

This author gives his treatment in a paper before the last meeting of the American Medical Association (1884). He places almost sole reliance on the use of a diet from which starches and sugar have been excluded. This should be carried out absolutely. Systematic daily muscular exercise should be enforced, but fatigue should be scrupulously avoided. All alcoholic excesses and the use of sweet fruits must be avoided. In cases where the sugar persists, he uses a solution of the bromide of arsenic, in doses of three drops three times a day. This may be continued for weeks or months without unpleasant effects. He continues the rigid diet for at least two months, even in the mildest cases. The return to ordinary diet must be gradual, and the urine during this time should be examined for sugar every five or six days.

PROF. A. BOUCHARDAT, PARIS.

The Dietetic Treatment. This is indispensable to success. Some writers are exceedingly stringent, but M. BOUCHARDAT has proved, by

the comparison of the articles of food which a diabetic may take without prejudice, that it is quite possible to keep up a sufficient degree of variety in the character of the meals. The list of articles which it permits is about as follows: All sorts of meat, roasted, stewed, and even dressed with spices, but not with flour; fresh-water fish and marine fish, in eating which the want of bread is less felt than in eating meat; oysters, mussels, crabs, lobsters, etc.; eggs in all forms known to the culinary art; rich, good cream, but no milk; of vegetables, spinach, artichokes, asparagus, green beans, the different varieties of cabbage; of the salads, water cresses, endives, lettuce; of fruits, strawberries and peaches. Although the loss of sugar is augmented by a very free supply of liquid, and decreased by a prevention of it, yet it is not advisable to forbid the patient to quench his thirst.

J. M. DA COSTA, M. D., PHILADELPHIA.

749. R. Pulveris opii, gr. ss.
For one pill, thrice daily.

The *opium treatment*, Dr. DA COSTA has found to be productive of the most good in his hands. He cannot explain the *rationale*, but he does know that it has the most influence in lessening the thirst, the excretion of sugar and the general waste of flesh. The only objection to its use is the risk which is run of accustoming the patient to taking this powerful drug.

750. R. Ammoniae carbonatis, ℥ ij-iv
Aquae cinnamomi, f. ℥vj. M.

A tablespoonful three or four times a day. This, in addition to a moderately restricted diet, forms a very good treatment.

751. R. Tincturae ferri chloridi, gtt. xx-lx.
For one dose, in water, thrice daily.

This will often do good when nothing else proves of service. The *pepsin* and *rennet treatments* amount to nothing. Dr. DA COSTA has tried them faithfully, and merely lost time.

Diet. Although substances containing a great deal of sugar, and still more those containing a great deal of starch—which is readily converted into sugar—are poisonous to diabetic patients, yet it is not advisable to put them on a strictly animal diet. Such a regimen is irksome in the extreme, and will not be followed out. After all, it is not

the saccharine substances which go into the body that cause diabetes ; they merely add to it. It is simply impossible to avoid giving food capable of being converted into sugar.

Bran bread is perfectly unobjectionable ; but a small quantity of ordinary bread should be permitted. Cakes made from *almond-flour* are favorites with English physicians. Fruits should be interdicted, and potatoes sparingly used.

Bran-baths (two or three handfuls of bran in the bath) are very serviceable in relieving the dryness of the skin, which is so troublesome.

752. R. Infusi cascarillæ, Oj.
A wineglassful three or four times a day.

The use of this light bitter lessens the thirst. Of course, water must be allowed beside ; but the infusion will take the place of a good deal of water.

PROF. S. D. GROSS, M. D., PHILADELPHIA.

A very successful treatment suggested by this eminent surgeon is recorded in the *Medical and Surgical Reporter*, August 5, 1871, and March 16, 1872. The drugs used are opium, tannic acid, and ergot internally, and the application of ointment of veratria to the spine. The following prescriptions may serve as illustrations of the manner in which they may be used :

753. R. Acidi tannici, gr. v
Opium pulveris, gr. ss. M.
To be taken three times a day, between meals, and tincturæ ergotæ, f. ʒj, in water, before each meal.

754. R. Veratriæ, ʒj
Unguenti cetacei, ʒj. M.
For an ointment. A piece the size of a cherry, to be well rubbed along the spine morning and evening.

The patient should be directed to abstain as much as possible from animal food, to confine himself to a simple vegetable diet, and to procure lager beer or ale to drink instead of water. If either of these becomes nauseous, he can use weak tea or whiskey and water.

The tannic acid should be increased gradually until twenty or thirty grains, three or four times a day, are administered. The quantity of

opium and ergot need not, however, be increased in like proportion. The combination may be :

- | | | | |
|---------|------------------|-----------|----|
| 755. R. | Acidi tannici, | ℥j | |
| | Opii pulveris, | gr. j | |
| | Tincturæ ergotæ, | f. ʒ iij. | M. |
- This quantity three or four times a day.*

Under the judicious use of this treatment, even severe cases of this very grave disease have recovered in three or four months, and have remained free from relapses.

DR. J. HOAG, INDIANA.

This writer claims, in the *Journal of Materia Medica*, September, 1874, to have achieved uniform success with the following treatment: The patient is forbidden all fruits and vegetables save bread of unbolted wheat flour, and this in the most limited quantities. Tea, coffee, liquors, and all drinks except pure water and milk diluted with lime-water, at meals, are also forbidden. The diet should be fresh meats, cream, cheese, butter, soft eggs, fish, oysters and small quantities of bran bread. The entire body is sponged daily with an alkaline bath, and the following ordered :

- | | | | |
|---------|-------------------------------|--------|----|
| 756. R. | Antimonii sulphureti, | ℥j | |
| | Pulveris ipecacuanhæ et opii, | ℥ij | |
| | Ammonia carbonatis, | ʒj | |
| | Pulveris opii, | gr. x. | M. |
- Make twenty powders. One to be taken morning and night.

Also :

- | | | | |
|---------|-------------------------------|---------|----|
| 757. R. | Tincturæ cinchonæ, | f. ʒj | |
| | Tincturæ ferri chloridi, | f. ʒ iv | |
| | Tincturæ digitalis, | f. ʒj | |
| | Spiritûs lavandulæ compositi, | f. ʒj. | M. |
- A teaspoonful daily, at noon.

DR. ORSON MILLARD, MICHIGAN.

- | | | | |
|---------|--------------------------|-----------|----|
| 758. R. | Tincturæ ferri chloridi, | f. ʒ ij | |
| | Olei morrhuæ, | f. ʒ iv | |
| | Acidi carbolicî, | f. ʒ iss. | M. |

A teaspoonful before each meal. Said by this writer (*Michigan University Medical Journal*, January, 1872,) to bring about the disappearance of the sugar promptly in uncomplicated cases.

THE MILK DIET TREATMENT.

This is either absolute or modified. Dr. A. S. DONKIN, of Dublin, (*Dublin Journal of Medical Science*, November, 1875,) prefers the former as the only one of real value. He uses *skim-milk*, and his rule is that the skim-milk *regimen must be exclusive*; that the quantity of skim-milk—*properly prepared by the careful removal of the cream*—beginning with four, five, or six pints on the first day, must be increased more or less gradually, according to circumstances, to eight, nine, ten, eleven, or twelve pints in the twenty-four hours, according to the age, sex, size, and condition of the patient. No rule as to quantity can be laid down to suit individual cases; it must be regulated to suit the requirements of each, *but in no instance should it exceed twelve pints*. Not more than seven or eight pints should be taken in the natural fluid condition. When a larger quantity is necessary, the surplus should be made into curd by the essence of rennet, and taken at separate meals. The skim-milk may be taken cold, or warmed to a temperature of 106° or so, but it must not be boiled, because a temperature of 212° alters the physical properties of caseine, and greatly impairs its therapeutic properties. The specific gravity of the skim-milk used should never be below 1035; that of the best quality is 1040. The daily allowance must be divided into regular meals. The constipation which this diet generally produces must be carefully remedied by the frequent administration of castor oil, or of some mild saline aperient. The author does not parade this method of treatment as a panacea; his experience of it has led him to the conclusion that if there is no progressive reduction of the specific gravity of the urine and of the sugar it contains after the expiration of a week (all rules having been observed,) it may as well be discontinued.

On the other hand, Dr. GEORGE W. BALFOUR gives pure milk. From two to three months is the period usually stated as sufficient to cause the complete disappearance of sugar from the urine; and then the diet may, if desired, be rendered more solid by the detraction of some of the milk, and the addition of butchers' meat and green vegetables. It is doubtful, however, whether diabetic patients, even though apparently cured, can ever return to ordinary diet. Present experience of such cases is too limited to afford any data at all in regard to this. When the milk diet is employed in the cure of other forms of ill-health, it is often conjoined, after the first few days, with a small amount of solid

food; and what such patients relish most is plain bread with salt, or a salt herring, the mawkish sweet taste of milk inclining them this way. In diabetes, the thirst present usually makes it better to give the patient his full allowance of milk, and let him divide it over the day as he pleases.

NOTES ON REMEDIES.

Alkalies and Alkaline Waters have, according to Professor SEEGEN, proved themselves the best remedies in diabetes—the natural waters, such as Carlsbad, in Europe; in the United States, Bedford Springs, Perry Springs, Congress Springs, Bethesda or Waukesha, etc.

Ammonii Carbonas is regarded by Dr. BARLOW, (*Guy's Hospital Reports*, vol. x,) in doses of gr. v–viij, with a few drops of tincture of opium, in a light, bitter infusion, every six hours, as a most efficient remedy. Animal food, together with cruciferous vegetables, as greens, brocoli, turnip tops, etc., should, at the same time, be freely taken.

Ammonii Citras, conjoined with the use of Dover's powder or ipecacuanha, is considered by Dr. W. PROUT as one of the best of diaphoretics in diabetes.

Ammonii Phosphas is recommended by Dr. BASHAM (*British Medical Journal*, April 10, 1869,) combined as follows:

759. R.	Ammonii phosphatis,		
	Ammonii carbonatis,	āā	gr. x
	Spiritūs ammoniæ aromatici,		℥xxx
	Aquæ,		f. ℥j. M.

Add this to the juice of a fresh lemon, and order to be taken thrice daily. Its use must be persevered in.

Arsenicum renders good service when the patient is thin, and assimilation defective, but in fat subjects it should not be exhibited. (BARTHOLOW.) It may be combined with the phosphates, thus:

760. R.	Syr. calcii lacto-phos.,	℥iv	
	Liq. potass. arsenit.,	f. ℥j	M.

S.—A dessertspoonful ter die.

**Arsenite of Bromine* is highly recommended by Dr. R. H. GILLIFORD (*Medical and Surgical Reporter*, June 9, 1883).

Belladonna, in full doses, in connection with sulphur baths, has been recommended by Prof. GUENEAU DE MUSSY.

Calcii Sulphidum has been recommended in diabetes by Dr. RINGER, gr. $\frac{1}{2}$ thrice daily. It has also been very favorably reported on by other physicians, and deserves a careful trial. Dr. C. M. CAULDWELL (*N. Y. Med. Jour.*, April 5, 1884,) has used calcium sulphide in three cases; in one it produced no effect whatever, in the others improvement began and recovery took place during the administration of the remedy. The dose is $\frac{1}{4}$ to $\frac{1}{2}$ grain three to five times daily. Dr. C. concludes that although calcium

sulphide is certainly not a specific in diabetes, yet it seems worthy of a trial in persistent cases of this distressing disease.

Carbolicum Acidum has been used successfully in Germany.

761. R. Acidi carbolici, gr. xv
Aquæ menthæ piper., f. ʒx. M.

One-third of this quantity in divided doses, each day, increased gradually to one-half the quantity daily.

Codeia is preferred by Dr. F. W. PAVY (*Guy's Hospital Reports*, 1870,) to opium and morphia, as being equally effectual in controlling the disease without exerting the same narcotic effect. The proper commencing dose is gr. $\frac{1}{2}$, thrice daily, which may be gradually increased to gr. ij-ijj.

Creasote has been recommended by Sir T. WATSON and Dr. T. H. TANNER. It will, in some instances, remove the extreme thirst, especially when the quantity of sugar is small.

Cuprum. Finely pulverized copper, gr. $\frac{1}{6}$ - $\frac{1}{3}$, twice daily, with meals, has been found by Dr. GRAZZINI to diminish the saccharine excretion. (*London Medical Record*, December, 1878.)

Ferrum. Preparations of iron are nearly always useful :

762. R. Tincturæ ferri chloridi, f. ʒvj
Acidi nitro-muriatici, gtt. lxxx. M.
Twenty drops three times a day.

Glycerina has been administered in Germany.

763. R. Glycerinæ puræ, f. ʒ vj-viiij
Acidi citrici, ʒ iss
Aquæ, Oij. M.
To be drunk at intervals through the day.

Hydrogen Peroxide has been tried successfully by Dr. JOHN DAY. (*Lancet*, January, 1868.)

764. R. Tinct. hydrogen. perox. ether., f. ʒ ss
Aquæ, f. ʒj. M.
This amount three times a day.

**Iodoform* is highly recommended by Professor MOLESCHOTT, of Rome. He uses the following formula :

765. R. Iodoformi, gr. xv
Ext. lactuc. sat., gr. xv
Cumarin, gr. $\frac{1}{15}$. M.
Ft. pil. No. xx. S.—One pill from twice to eight times a day.

Jaborandi has, of late, been employed with success, in some cases of this disease.

Nux Vomica, gr. j, increased to gr. iv of the extract, has been found successful by Dr. E. ZARZANA. (*Gaz. Med. di Roma*, December, 1878.)

Opium and its preparations are well borne, and very valuable in diabetes. From gr. vj-xij, daily, are required to produce a decided impression. Yet it cannot be said to be curative. Of the various alkaloids, Dr. F. W. PAVY (*Guy's Hospital Reports*, vol. xv,) finds that opium, morphia, and codeine, all possess the power of checking the elimination of sugar in the urine. Of the three, he considers codeine to be the best. Dr. PAVY begins with $\frac{1}{2}$ grain of the last, and gives it, by gradually increasing doses, to the extent of 10 grains three times a day. He says it is superior to all other medicinal agents in its controlling influence over the disease, none other giving the same immediate and striking effects. It should be given in full doses, and persevered in, diabetic patients being very tolerant of its use. It produces its best effects in old cases occurring in the aged.

Phosphoric Acid, largely diluted, lessens the inordinate thirst more effectually than any other acidulated drink.

Potassii Bromidum. Dr. AUSTIN FLINT (*American Practitioner*, January, 1870,) relates three cases treated by bromide of potassium, in doses of fifteen to twenty grains three times a day, combined with ordinary diabetic diet. In each, there was a rapid diminution of thirst, a decrease in the specific gravity of the urine, and an improvement in the general health. In a paper before the Paris Academy, in 1883, Mr. DUJARDIN BEAUMETZ said that bromide of potash is essentially an adjuvant in the treatment of diabetes; it has its advantages and disadvantages. In cases of medium intensity, and sometimes in grave cases, in conjunction with dietetic means and the alkaline treatment, it may cause the sugar to disappear from the urine, but at the expense of a loss of strength on the part of the patient. The general impression of the members of the Academie appeared to be that this medicament might be used when the sugar did not disappear under the use of proper dietetic means, but that it was not well to employ it primarily.

Potassium Iodidum has cured cases of syphilitic origin.

Quiniae et Morphiae Bromidum. This combination has been used in diabetic phthisis, by Dr. BENJAMIN W. RICHARDSON, of London. He states that, under its influence, the quantity of sugar and of urine notably decreases, the cough is lessened, the appetite and digestion are improved, and the recurrent hectic is held in abeyance more certainly, than by any other remedy with which he is conversant.

Quiniae Sulphas will diminish the sugar in most instances.

Rheum is praised by Dr. COPLAND, not only as a useful aperient in diabetes, but as a promoter of the digestive and assimilative processes.

Rhus Aromatica, as supplied by Messrs. PARKE, DAVIS & Co., of Detroit, has been found of marked efficacy by Dr. P. T. HEARTT, of New York, and others.

Sodii Citras. Recent journals have called attention to a recommendation by M. GUYOT DARMECY, of citrate of soda in the treatment of diabetes,

given in daily doses of one-half to one drachm. Analysis has shown that sugar disappears from the urine when this salt is used with the food instead of common salt. In the *Detroit Lancet*, 1883, Dr. O. C. KNIGHT recommends the soda citrate as "an excellent remedy."

Sodii Phosphas is regarded by Dr. W. PROUT as one of the few saline purgatives admissible in this disease.

Sodii Salicylas. Dr. MULLER WARNER, in the *Berlin Klinische Wochenschrift*, January, 1877, gives the following conclusions: 1. Salicylate of soda can completely remove the symptoms of diabetes mellitus, yet its action does not appear to be lasting in many cases. 2. The symptoms of diabetes are made to disappear the quicker by salicylate of soda, the greater the dose in which it is used and the longer it is borne by the patient. 3. The first action of medium daily doses (ʒij-ʒijss), on diabetes, appears gradually to abate, while large doses (ʒiv in a day) act on patients progressively. 4. Salicylate of soda can be borne in chronic diabetes mellitus, in large daily doses, for a long time, without special disturbance of the general condition.

Bethesda Water. The *Bethesda* water, from Waukesha, Wis., has a wide and well-deserved reputation in this disease. Its use should not be neglected.

DROPSY.

FREDERICK T. ROBERTS, M. D., M. R. C. P., LONDON.

This esteemed authority remarks that far too little heed is usually given to the question of *rest* and *position*, in the treatment of dropsy. The part affected should be maintained in an elevated posture, the legs, for instance, if anasarcaous, placed higher than the body. *Pressure* should be applied by bandages or adhesive strips, and is a very valuable adjunct. All external dropsical parts should be kept clean and dry. A decided diminution of drinks and liquid articles of food is adapted to dropsical accumulations in serous cavities. The only *diaphoretic* he considers of much practical value, is some form of bath which promotes perspiration, such as the warm, vapor, hot-air, or Turkish bath. They are most valuable in renal dropsy. The best *purgatives* are extract of elaterium, (gr. $\frac{1}{6}$, gradually increased to gr. ss,) jalap (ʒj to ʒj), and cream of tartar (ʒj to ʒij.) For a diuretic, the following:

766. R. Extracti elaterii, gr. v
 Pulveris scillæ,
 Pulveris digitalis, āā gr. xx
 Extracti hyoseyami, gr. xxx.

To make twenty pills. One to be given every other night.

PROF. NATHAN S. DAVIS, M. D., CHICAGO.

It is well ascertained that, in a pathological sense, dropsy is not a disease, but a symptom directly dependent either on inflammation, mechanical obstruction of blood vessels, or altered composition of the blood. Its rational treatment must, therefore, be guided by an appreciation of these causative changes. Where, as in many cases, we are called upon to strengthen a weak heart and produce efficient action of the kidneys, we can accomplish this by combining digitalis and scutellaria, as :

767. R. Extracti fluidi scutellariæ, f. $\frac{3}{4}$ ijss
 Tincturæ digitalis, f. $\frac{3}{4}$ j
 Tincturæ hyoseyami, f. $\frac{3}{4}$ ss
 Potassii nitratis, $\frac{3}{4}$ ij. M.

A teaspoonful every three hours.

When there is dyspnœa from effused fluid, a blister to the chest aids in relieving this symptom. When diuretic remedies do not act efficiently in such cases, the system will respond to their action promptly by the use of a proper nerve tonic, as :

768. R. Strychniæ sulphatis, gr. $\frac{1}{16}$
 Ferri citratis, gr. ij. M.

For one pill. One three times a day.

DR. WALSHE.

This author believes that the abstraction of a small quantity of blood, either by venesection or by cupping of the chest, facilitates the action of hydragogue remedies when a clogged condition of the heart's cavities co-exists with pulmonary congestion and anasarca of more or less acute course, and the general vigor is as yet unimpaired to any serious degree. Of hydragogue purgatives, elaterium, gamboge, bitartrate of potass., and the pulvis jalapæ comp., are the most valuable of the class. The following is a useful formula :

769. R. Extracti elaterii, gr. $\frac{1}{8}$ — $\frac{1}{2}$
 Creasoti, gtt. j
 Extracti hyoseyami, gr. ij. M.

For one pill. To be repeated as required.

Diuretics, uncertain in their effects, are often rendered much more active by preliminary cupping or dry-cupping of the renal regions. Of the numerous diuretic agents, the *infusion of digitalis* has appeared to Dr. WALSHE the most active, and no amount of apparent weakness of the heart is, *per se*, an absolute contra-indication to its use. Hydragogue purgatives often prevent the action of more direct diuretics; and small doses of blue pill occasionally, at bedtime, are very useful adjuvants.

Diaphoretic drugs are of very little utility, but the free diaphoresis produced by the vapor or hot-air bath is sometimes strikingly beneficial. If there be any tendency to bronchitis, bronchial discharge should be encouraged by the free use of expectorants. Removal of anasarca fluid by mechanical means becomes necessary, if medicines fail in controlling its increase; and if inflammatory changes in the skin appear imminent, scarification is dangerous, being not unoften followed by erysipelas, while *acupuncture* over the insteps, or at the upper and inner parts of the thighs, is not attended with the same danger.

C. MURCHISON, M. D., F. R. S., ETC., LONDON.

770. R.	Pulveris scillæ,	gr. iss	
	Pulveris digitalis,	gr. ss	
	Pilulæ hydrargyri,	gr. ij.	M.

For one pill, two or three times a day.

This is a pill which has enjoyed a long and merited reputation for treatment of dropsy, in the Middlesex Hospital.

Diuresis will also sometimes be induced by fomenting the abdomen with an infusion of digitalis of about four times the usual strength.

771. R.	Potassii acetatis,	gr. xx	
	Spiritûs ætheris nitrosi,	f. $\frac{3}{4}$ ss	
	Decocti scoparii,	f. $\frac{3}{4}$ iss.	M.

To be given with each dose of the above pills.

At the same time, an ointment composed of equal parts of blue ointment and ointment of belladonna is to be applied over the abdomen. This treatment is a successful one in cases of ascites due to cirrhosis.

ROBERT J. GRAVES, M. D., DUBLIN.

772. R. Potassii nitratis,
 Potassii carbonatis, āā 3j
 Tincturæ scillæ,
 Tincturæ digitalis, āā gr. xxx
 Mellis despumati, f. 3ij. M.

For an electuary. To be taken in teaspoonful doses, in the course of three or four days, in order to increase the activity of the kidneys, in various forms of dropsy.

773. R. Oxymellis scillæ, f. 3 iss
 Potassii bitartratis, 3ij. M.

Dose, a tablespoonful, as a diuretic, in various forms of dropsy.

DR. GUIBERT, PARIS.

774. R. Tincturæ scillæ,
 Tincturæ digitalis,
 Tincturæ colchici, āā f. 3 iij
 Linimenti camphoræ, f. 3 vj
 Aquæ ammoniæ, f. 3 iss. M.

To be rubbed well into the abdomen and thighs in dropsy.

DR. PORCHER, PARIS.

775. R. Sodii sulphatis,
 Potassii bitartratis, āā 3j
 Spiritûs ætheris nitrosi, f. 3 ijss
 Syrupi, f. 3j
 Aquæ, f. 3 vss.

A tablespoonful twice daily, in dropsy accompanied with an active, full circulation. It procures an abundant alvine evacuation, and often, at the same time, a copious secretion of urine.

776. R. Tincturæ cantharidis, f. 3 iss
 Spiritûs ætheris nitrosi, f. 3 vj
 Syrupi aurantii corticis, f. 3j
 Aquæ fœniculi, f. 3 viij. M.

A tablespoonful two or three times a day, in different forms of dropsy.

PROF. A. GUBLER, FACULTÉ DE MÉDECINE DE PARIS.

777. R. Tincturæ digitalis,
 Extracti ergotæ fluidi, āā f. 3 ss
 Acidi gallici, 3j
 Potassii bromidi, 3ij
 Aquæ lauro-cerasi, f. 3 ij
 Syrupi, f. 3 iij
 Oxymellis scillæ, f. 3 iv. M.

A tablespoonful in water, or a diuretic infusion, two or three times a day, in various forms of dropsy.

urine increases. In *inflammatory dropsy attendant on granular disease of the kidneys*, the use of diuretics, particularly those of a stimulating character, is generally condemned. The administration of diuretics by the *endermic method* is sometimes effectual in removing dropsies. The diuretics usually employed in the treatment of dropsies are digitalis and digitalin, sweet spirits of nitre, juniper, acetate of potash, nitrate of potash, cream of tartar, squills, broom tops. The following formulæ have been recommended :

783. R. Digitalin, gr. $\frac{3}{4}$
 Pulveris scillæ,
 Pulveris scammon., āā gr. lxxv
 Mucilaginis, q. s. M.
 Make one hundred pills. Of these two, then four, and lastly six, are to be given daily, according to the effect produced.

784. R. Potassii nitratis, ℥ iij
 Baccar. juniper., ℥ xv
 Vini albi, Oiss.
 Dose, f. ℥ iij daily.

785. R. Potassii bitartratis,
 Juniper baccar. contus., āā ℥ ss
 Aquæ ferv., Oj. M.

This infusion is considered by Prof. STILLÉ to have few superiors in the idiopathic forms of general dropsy.

786. R. Scoparii (seeds), ℥ iij
 Alcoholis, f. ℥ viij. M.
 Macerate for ten days. Dose, f. ℥ j-ij thrice daily.

If it cause diarrhœa, five or six drops of laudanum may be added ; if the patient is much debilitated, iron or quinine may be added. This formula is of service in all forms of dropsy excepting ovarian.

Cathartics, particularly elaterium, gamboge, and cream of tartar, are of great service in dropsical affections, but they are to be used with great caution when the patient is nearly debilitated, and, as a rule, to be avoided in dropsy connected with heart disease. Dr. CHRISTISON recommends gamboge, in doses of gr. v-vij, every other day, finely powdered and combined with cream of tartar, as a certain and easily-managed cathartic in reducing dropsy. *Colocynth* may be employed in dropsical affections connected with disease of the liver, but is inferior to elaterium. *Croton oil* is sometimes useful when given with great caution, in small and only gradually-increased doses, but is inadmissible

if the patient is old or debilitated. *Elaterium* is a very efficient cathartic in dropsy, anasarca, and œdema, arising in connection with disease of the heart. Dr. J. HOPE gives it in the form of a pill, in the doses of gr. $\frac{1}{8}$ – $\frac{1}{4}$, combined with capsicum, to prevent griping, and with a grain of calomel, to prevent its emetic effects. One of these pills should produce six or eight watery evacuations, and may be repeated two or three mornings in succession, or every second or third morning, according to the patient's strength. The dose of elaterium may be carried, with caution, to two grains. The effects of this remedy, according to Dr. HOPE, are sometimes truly astonishing, removing an extreme universal anasarca in three or four days; but, as it is apt to be violent in its operation, it should be given to strong subjects only. The following formulæ are also valuable:

787. R. Elaterii, gr. ij
 Extracti gentiani, gr. xx. M.
 Divide into four pills; order one every hour, till it causes free evacuations. Repeat every two or three days, for a fortnight, and then suspend for a short time.

788. R. Elaterii, gr. j
 Extracti colocynthidis compositi, gr. xl
 Extracti hyoscyami, gr. xij. M.
 Divide into twelve pills. Take one every night.

789. R. Pulveris jalapæ compositi, gr. xx-xxx
 Potassii bitartratis, gr. v-x-xv
 Olei carui, gtt. ij
 Aquæ, f. ʒiss.
 For one dose; to be repeated, so as to keep up an unremitting discharge from the bowels.

This formula is recommended by Dr. CHAPMAN.

NOTES ON REMEDIES.

Ammoniac Murias and *Benzoas* are useful in dropsy dependent upon hepatic disease, and also in ovarian dropsy. Dr. COPLAND recommends these combinations with warm, diuretic effusions.

Arlentin. This alkaloid of uva ursi is a potent diuretic, and has been commended by Dr. MENSCHÉ (*Lond. Med. Record*, 1883,) and others.

**Asclepias Syriaca*. This plant, *vulgò* milk-weed, has a specific power in dispelling dropsical effusions. It was first brought forward by Dr. JEWETT, in an article in the *New York Medical Society's Transactions* for 1869, and later, Dr. HUTCHINS reported to the Kings County Society, very remarkable results from its use. For instance, in forty-eight hours, a man described as frightfully distended, was reduced to a

skeleton, and any disposition to re-accumulation of water was rapidly relieved by an infusion of this drug. If not carefully given, it may produce vomiting or diarrhoea, both of which are to be avoided. Dr. JEWETT's mode is to put $\frac{1}{2}$ ounce in a pint of gin, add an infusion of 2 ounces of juniper berries in a pint of water, and give from 1 to 4 teaspoonfuls, as the stomach will tolerate it, every four hours, until the effect is obtained. He expresses unbounded confidence in it, after an experience of thirty years; but he warns us that, although the *root* is officinal in the United States Dispensatory, it is only the *bark of the root* which is active. A fluid extract is prepared, the dose of which is 10 drops. It has been erroneously referred to, in various articles, as *Apocynum Cannabinum*.

Armoracia Radix. Horse-radish was much employed as a diuretic, by SYDENHAM and the older physicians, in dropsical affections.

Blatta Orientalis. A derivative obtained from the cockroach, has been introduced from Russia. In doses of gr. ij-iv, its diuretic action is well marked, and it has secured the testimony of a number of observers.

Caffein. The citrate of caffein, gr. iv-viiij, has been used by Prof. GUBLER with good results, in cardiac dropsies.

Cannabis Indica has been used (℥xx of the tincture every four hours) as a diuretic in dropsy.

Chenopodium Anthelminticum. Dr. SAMEUL R. BURROUGHS, of Houston, Texas, in a report on the "Indigenous Medical Resources of Texas," in the *Transactions of the Texas State Medical Association*, 1877, makes the following note: "The chenopodium anthelminticum root, split or cut to pieces, and put into an ordinary bottle filled with vinegar, to which has been added 2 or 3 drachms of carbonate of iron, will relieve anasarca when a sequel to scarlatina, or when dependent on any functional derangement."

Copaiba. Dr. S. WILKS (*Lancet*, March 21st, 1873,) speaks with the greatest confidence of the resin of copaiba as a diuretic. It is particularly useful in cases of dropsy from cardiac or hepatic disease. Fifteen or twenty grains in mucilage and flavored water should be given, three or four times a day.

**Digitalis* is much employed in dropsical affections, alone or combined with squills, mercury, carbonate of ammonia, or tincture of the chloride of iron. Sir H. HOLLAND recommends a combination of digitalis and tincture of the chloride of iron in the dropsy following scarlet fever. The same combination is useful in many other forms of dropsy attended with anæmia. Diuresis is sometimes induced by the external application of digitalis to the abdomen, either by fomentation with an infusion of about four times the usual strength, or by frictions with an embrocation of equal parts of tincture of digitalis and soap liniment, or by poulticing with a warm linseed poultice, to which a fluid ounce of the tincture has been added.

Ferri et Potassii Tartras has been found useful as a tonic and diuretic in cases of anasarca connected with disturbed action of the heart, in which it would be dangerous to give any stimulating tonic.

**Jaborandi* is very valuable in renal dropsy, when the secretion of urine is much decreased or suppressed.

Phosphoricum Acidum Dilutum, diluted with oil and acid, as an embrocation over the abdomen, has been employed with benefit in some cases of dropsy.

Rheum, in small doses, or in infusion as a vehicle for the saline diuretics, squills, juniper or colchicum, is regarded by Dr. COPLAND as a useful medicine in dropsies.

Sinapis is sometimes beneficial, because of its diuretic and stimulant qualities. It is best given in the form of whey, made as follows :

790. R. Bruised mustard seeds, ℥ss.
Boil in Oj of milk, and strain. The whole to be taken daily in divided doses.

Terebinthina Oleum, in doses sufficient to act freely on the bowels and kidneys, is occasionally useful in ascites, unconnected with renal disease or with great gastric irritability or inflammatory action.

Urtica Dioica. An infusion of the root of the stinging nettle is considered by Dr. MARIS, of Philadelphia, the very best diuretic he is acquainted with.

EXTERNAL MEASURES.

Acupuncture is the introduction of needles into the body to relieve or cure disease, and is often needful in dropsical affections. It is performed by running five or six fine steel needles, two or three inches long, right into the dropsical part. The needles are furnished at the blunt end with a knob of sealing-wax or ivory. They should be introduced by a slight pressure combined with a semi-rotary motion between the thumb and forefinger, and withdrawn with the same motion. Ordinarily, no fluid escapes on the withdrawal of the needles, but occasionally a drop of blood follows. Gentle pressure will suffice to restrain any hemorrhage, which exceptionally may occur. The pain is quite trifling, and often scarcely noticed. The needles may be introduced into muscular, aponeurotic and tendinous parts, but not into serous cavities. They should be inserted to the depth of one-fourth of an inch to two or three inches, according to the thickness of the muscles.

In regard to the period of sojourn of the needles in the part, Dr. ELLIOTSON remarks that one needle allowed to remain an hour or more is often more efficacious than several speedily withdrawn. The pain sometimes ceases instantly. The operation may require repetition several times ; usually, however, twice is sufficient.

Abdominal Compression. When there is any tendency to absorption in ascites, it will be greatly accelerated by tightly bandaging the abdomen with a flannel bandage.

FUNCTIONAL URINARY DISORDERS.

INCONTINENCE OF URINE.

This affection is most frequent in children, although adults are not free from it. One of the most popular remedies is *belladonna*, or its alkaloid, *atropia*, administered in sufficient doses to exert the physiological effect of the drug.

PROF. S. D. GROSS, PHILADELPHIA.

In the nocturnal incontinence of children, there should be a bland and unirritant diet, late suppers should be avoided, no drink taken after supper, and the bladder emptied on retiring. The position in bed should be upon the side or belly, to prevent the urine from irritating the neck of the bladder. The internal remedies of most use are as follows :

791. R.	Strychniæ,	gr. j	
	Pulveris cantharidis,	gr. ij	
	Morphiæ sulphatis,	gr. iss	
	Ferri pulveris,	℥j.	M.

Make forty pills. Give one three times a day to a child ten years old.

The cold shower-bath once or twice a day is a valuable auxiliary.

DR. BRUGELMANN, COLOGNE.

Our author relates (*Berliner Klinische Wochenschrift*, February 10, 1873,) a case of well-marked incontinence of urine in a girl aged thirteen, very nervous, and distinctly anæmic, where the administration of syrup of *iodide of iron* afforded marked relief in eight days. After persevering with it for another fortnight, she was completely cured.

DR. ROBERT FARQUHARSON, LONDON.

This writer, in a study of the action of drugs on incontinence, (*Practitioner*, July, 1879,) divides cases of incontinence into three classes.

In the first class, debility and anæmia are the exciting causes. The patients are pale and weak. In such, small doses of *iron* will act better than anything else, and will effect a cure in the lighter cases.

The second class includes those in which the debility is generally congenital and dependent on an excessive irritability of the bladder.

The drugs most potent in these cases are *ergot* and *belladonna*. Of these, Dr. F. prefers the latter. It must, however, be vigorously pushed, up to the full development of its physiological effects, which, when this drug is given to young persons, need never excite any apprehension. He has given the tincture in doses of f.ʒij, to a boy of nine, f.ʒijss, to a girl of thirteen, etc.

The third class may be called a sort of choreic condition of the bladder, sometimes associated with skin diseases, often more or less intermittent. In such, sulphate of zinc, arsenic, strychnia and iron may be used, singly or in combination, with possible good results, but on the whole the treatment is far from satisfactory. Galvanism and blisters to the spine occasionally are efficacious. The dietetic plan, recommended by some, of withholding all meat, is rejected as useless by our author. Santonin, praised by RINGER, he has tried without effect.

In children who sleep heavily, Professor FONNSAGRIVES has occasionally observed that a cup of strong coffee in the evening prevented them from wetting the bed; he believes that this, and also belladonna, act simply by causing them to sleep less profoundly, and then they do not lose control of the sphincter muscles of the bladder.

DR. SAMUEL S. ADAMS, OF GEORGETOWN, D. C.

According to this author (*American Journal of Obstetrics*, June, 1884), *belladonna* is the remedy *par excellence* in the treatment of those cases believed to be associated with a tonic spasm of the bladder. In order to derive benefit from the drug, it should be given in large doses at bed-time, which should be increased drop by drop, daily, until improvement results or its physiological effects are obtained. If there is relaxation of the sphincter vesicæ, or paresis of the muscles of the bladder, strychnia is indicated. If the patients are puny or in ill-health, the general health must be improved while we are giving the above remedies.

He does not advocate circumcision as the certainly indicated remedy in all cases, nor does he believe that every boy who has incontinence, with an elongated prepuce, should be compelled to undergo the operation. If the prepuce cannot be retracted, he would advise operating; and while sometimes the opening in the prepuce is large enough, yet it cannot be retracted owing to adhesions. In such cases, if these adhesions are broken up, favorable results will follow.

He performs the operation of circumcision in the following manner: The prepuce is drawn forward and Henry's clamp tightly applied; the end of the prepuce is then cut off with scissors and the clamp left on the stump for several minutes to check bleeding; a director is then pushed along the upper surface of the glans, and the mucous membrane divided beyond the corona. The membrane is then turned back to meet the retracted skin, and made fast by five silk sutures. In about ten days the patient is well. He removes only that part of the mucous membrane that is cut off with the end of the prepuce; for by leaving a long membrane it can be turned back and thereby hide the cicatrix, which is a source of mortification to many parents.

DYSURIA.

PROF. J. B. FONNSAGRIVES, PARIS.

The forms of dysuria of functional origin are derived from neuralgia, paralysis or torpor of the muscles of the bladder and its neck. Hysterical subjects are quite liable to them. In nervous affections of the neck, the plan adopted by CIVIALE is the best; this includes catheterism of the urethra, vesical injections at first of warm water, then of cold, douches on the hypogastrium of warm and cold water, followed by revulsives to the same spot, and light purges.

Where the dysuria is owing to nervous atony or partial paralysis, Prof. F. believes the best results are to be had from *electricity* and *strychnia*. The former may be employed in either of the three methods recommended by DUCHENNE, (described in most works on medical electricity.) When these means fail, he would resort to injections of very weak solutions of *tincture of cantharides*, gtt. j-iiij, in half a tumbler of warm water, a method successfully employed by LISFRANC.

NOTES ON REMEDIES.

Arlentin. This new alkaloid of uva ursi has been extolled by German physicians in dysuria.

Benzoates. The benzoates of lithium or potassium have come into favor as agents to increase the quantity of the urinary secretion, and also to cause the disappearance of uric acid from the urine. They are well spoken of for this purpose by Dr. A. B. GARROD, of London (*Lumleian Lectures*, 1883).

Cantharides. In cases of temporary scanty or suppressed urine, as in hard drinkers, single drop doses hourly of the tincture of cantharides will often re-establish the secretion.

Circumcision. Dr. WM. MUIR reports (*Glasgow Medical Journal*, 1884,) the case of a boy suffering from incontinence in whom the foreskin was unusually long and a little inflamed. Circumcision was performed, and in two months the boy was quite cured.

Rhus Aromatica. The introduction of *rhus aromatica* has supplied the physician with an efficient remedy against that very annoying condition of the bladder which gives rise to incontinence of urine. The mode of its action seems to be through its specific tonic influence on the bladder, and particularly on the sphincters of that viscus. The dose is gtt. x or xx of the fluid extract every three hours, or on going to bed.

Stigmata Maidis. The stigmata of maizes exert an action on the urinary organs both anæsthetic and diuretic, and promises to have a wide applicability in their functional disorders. About a scruple of the extract is a dose.

HÆMATURIA.

DR. JAMES TYSON, OF PHILADELPHIA,

read a paper on "*Malarial Hæmaturia*" before the Pennsylvania State Medical Society in 1883, in which he said that if the diagnosis is accurately made, the results of treatment are often brilliant. It is that for malaria. Quinine, in doses of three to five grains every three hours when the hemorrhage is continuous, until it ceases. Or, if intermittent, sixteen to twenty grains, sufficiently anticipating the paroxysm, should be given. Mercurials may be combined with quinine. Iron, arsenic, and astringents have not been found of much service. The natural waters containing iron and alum may be expected to be of service, such as the Orchard Acid Springs of New York, and Bath Alum and other springs in Virginia, and some have a reputation in hæmaturia.

The "malignant" form of malarial hæmaturia, of which many more cases have occurred in the Southern United States during the past fifteen years than previously, is much more serious and more fatal. The hæmaturia is here sometimes ushered in with a chill or two, or the bloody urine may appear at once. Large quantities of bloody urine are passed, it is said one or two gallons in twenty-four hours. The urine is porter-like, and the sediment sometimes almost tarry in consistence. There is obstinate nausea, and vomiting of bilious and dark matter resembling black vomit. Intense jaundice rapidly supervenes;

sometimes in an hour the whole body is stained yellow. The jaundice is hæmatogenetic. There may be fever, hot skin, and temperature of 104° to 106° , but the strength rapidly declines, and the patient often dies in from twenty-four to sixty hours, and if he recovers, convalescence is prolonged. The negro seems to be exempt.

Autopsies show the same intense coloration of the internal organs, and sometimes the spleen is enlarged.

The only curative treatment is by quinine aided by mercurials. Morphia and carbolic acid have controlled the vomiting. Stimulants are necessary.

SAMUEL FENWICK, M. D., LONDON.

Hæmaturia occasionally appears without obvious cause, or follows a slight chill or other exposure. When the urine is decidedly tinged, an ice-bag may be applied to the loins. The patient must be kept at rest, and if severe he is better in bed. Sound port wine is one of the most useful astringents. Ergotin may be injected subcutaneously. The acetate of lead is of but slight efficacy in these cases. Internally, gallic acid may be prescribed, or one of the following combinations:

792.	R.	Extracti ergotæ fluidæ,	℥xiv	
		Acidi gallici,	gr. x	
		Aquæ cassiæ,	℥j.	M.
For one dose.				

793.	R.	Aluminis,	℥ iss	
		Syrupi simplicis,	℥vj	
		Infusum rosæ acidi,	ad ℥viij.	M.
One to two tablespoonfuls as required.				

The discharge of bloody urine is a symptom of so many diseases as well as local injuries, that it can only be referred to here in general terms. Internally, one of the most efficient remedies is *ergot*.

DR. CHARLES HORION, PARIS.

794.	R.	Ergotæ pulveris,	gr. xv	
		Acidi tannici,	gr. iij	
		Digitalini,	gr. ̄.	M.
Divide into ten pills. Give five a day, in hæmaturia. Prolonged cold injections into the bladder, cold compresses to the perineum and pubis.				

DR. W. LANGE, ERLANGEN.

795. R. Extracti ergotæ fluidi, ℥xxv
 Acidi tannici, gr. xxx
 Aquæ destillatæ, f. ʒvj
 Syrupi, f. ʒj. M.

The whole to be given in tablespoonful doses, in the course of the twenty-four hours in hæmaturia. Cold compresses to the hypogastrium, ice by the mouth, cold enemata.

PROF. S. D. GROSS, M. D., D. C. L., PHILA.

796. R. Olei terebinthinæ, āā f. ʒj
 Acidi sulphurici diluti, gr. xxx
 Acidi gallici, f. ʒss
 Mucilaginis acaciæ, f. ʒj. M.
 Aquæ,

A dessertspoonful every three hours.

In all cases, ice to the hypogastrium and in the rectum or vagina is serviceable; rest in the recumbent posture, the pelvis elevated, and the room cool, are important accessories.

Dr. W. ATLEE gives the fluid extract of ergot in doses f. ʒss-j, and continues it daily at bedtime for a considerable period.

NOTES ON REMEDIES.

Alumen injected into the bladder, in a solution of the strength of gr. xx to aquæ Oj, sometimes arrests the discharge in hæmaturia, after the failure of ordinary remedies. As an internal remedy in hæmaturia, alum is less effectual than gallic acid.

Ammonii Chloridum, in five grain doses, four times daily, has been successfully used by Dr. W. JONE MORRIS, of England.

Argenti Nitras, in very weak solution, injected into the bladder, in hæmaturia arising in connection with vesical tumor, is recommended by Sir H. THOMPSON.

Benzoin Odoriferum. Dr. R. M. NICHOLLS, of Louisiana, says, in the *Medical and Surgical Reporter*, February 3, 1883, that *quinine*, with the infusion of spice wood, *benzoin odoriferum*, as a drink in the place of water, will cure hæmaturia.

Chian Turpentine with three drops of arsenical solution, three times daily, and a four-grain dose of quinine every day at noon, has proved beneficial in the hands of Dr. RALFE, (*Lancet*, November 17, 1883.)

Digitalis, according to BARTHOLOW, has an undoubted power to arrest hemorrhage; it may be combined with other remedies that are synergistic, thus:

797. R. Infus. digitalis, f. ʒij
 Tinct. krameriæ,
 Ext. ergotæ fl. āā f. ʒj. M.

S.—A tablespoonful pro re nata.

Ferri Chloridi Tinctura, in doses of ℥x-xx, several times a day, is a valuable remedy in hemorrhage from the bladder.

* *Gallicum Acidum*, alone or combined with sulphuric acid, opium or digitalis, is an excellent astringent in hæmaturia. Dr. W. BRINTON recommends the following formula :

798. R.	Acidi gallici,	gr. x	
	Acidi sulphurici diluti,	℥x	
	Aquæ,	f. ℥j.	M.

For one dose.

Guaiaci Tinctura, ℥j, twice daily, has been used with success by Dr. JOHN N. UPSHUR, of Va. (*Half-Yearly Compend.*, July, 1879.)

* *Ipecacuanha* is a valuable remedy in hæmaturia. It should be given in the manner recommended for hæmoptysis.

Krameria, in scruple doses of the extract, is highly commended by Sir T. WATSON, after all ordinary remedies have failed.

* *Matico*, in infusion, doses ℥ij, every two or three hours, is regarded by Sir H. THOMPSON as equal to gallic acid or lead in hæmaturia.

* *Plumbi Acetas*, in doses of gr. ij-iv, combined with gr. ss-j of opium, is a well-known and effectual remedy.

* *Quinine* is highly effective in the intermittent trouble.

Tannicum Acidum, in the form of a pill, is an effectual astringent in hemorrhage from the bladder.

Terebinthinæ Oleum is employed with benefit where there is debility, relaxation and anæmia. Prof. BARTHOLOW commends the following formula :

799. R.	Olei terebinthinæ,	f. ℥ iij	
	Extracti digitalis fluidi,	f. ℥ j	
	Mucilaginis acaciæ,	f. ℥ ss	
	Aquæ menthæ piperitæ,	f. ℥ j.	M.

A teaspoonful every three hours.

NEPHRALGIA, (RENAL COLIC.)

Nephralgia, as a symptom, may be regarded as expressive of a gravelly or lithic acid diathesis ; it arises from the irritation of gravelly particles of lithic acid (amorphous), or of crystalline grains of lithic acid or lithates, detained in the renal tubes, and, till washed out by the urinary stream, giving rise to a distinctive group of symptoms.

Bearing this pathological fact in mind, the remedial agency of *diluents*, of alkaline *citrates* and *tartrates*, or any agent which, while it dilutes the urine and increases its quantity, also renders the uric acid

more readily excreted, will readily be admitted. Nephralgia, arising from the above causes, is recognized without much difficulty, and its temporary relief easily and promptly obtained. The potash salts, as effervescing citrates, mild mercurial alteratives and saline purgatives, especially the Carlsbad or Friederichshall waters, speedily effect the desired relief.

As nephralgia is symptomatic of a gravelly irritation of the urinary passages, so, if the gravelly condition continues unrelieved, nephralgia becomes the sign of a calculous disease of the kidney. The general treatment of *Lithiasis* is a question which is considered under "Surgical Therapeutics."

The immediate treatment of nephralgia must be by *opium* in large doses, preferably either by enema or by the hypodermic injection of morphia. Dr. F. KING, of Georgia, has observed that *ipecacuanha*, gr. xxx, at one dose, will sometimes relieve both the irritability of the stomach and the pain in such cases. (*Half-yearly Compend.*, Jan., 1878.)

Active counter-irritants over the seat of pain also aid in giving relief. See further in "Surgical Therapeutics," under "Lithiasis."

PAROXYSMAL HÆMATINURIA.

DR. STEPHEN MACKENZIE, OF LONDON.

According to this author (*Lancet*, Jan., 26, 1884), syphilis has been shown to be an antecedent in several cases, and the disease has been cured in some such by anti-syphilitic treatment. In others, malaria is the cause. Syphilis should be inquired for; and when present, treated. In the majority of cases quinine is the remedy of greatest service, even in cases not evidently malarial. Probably some patients who get well when taking quinine would have done so without it. Most patients find by experience that the only way to prevent attacks is to avoid exposure to cold by clothing and habits. But it was possible, as suggested by Dr. Barlow, that gradual accustoming to cold might be at once the most philosophical and successful mode of treatment, and this is decidedly worthy of trial.

NEPHRITIS.

DR. AUFRECHT, OF MAGDEBURG,

who has paid much attention to the pathology of nephritis, advocates most forcibly the adoption of an expectant line of treatment, in which dietetics play an important rôle. He absolutely discountenances the prescription of diuretic and diaphoretic drugs, and regards it as not good practice to encourage diaphoresis by hot baths or heated air. His plan consists in the administration of neutral salines, as bicarbonate of soda, which may be replaced by iron in the later stages, when the anæmia evoked by the albuminuria appears. The diet at first must, he says, be wholly vegetable, in the form of vegetable soups, and starchy and saccharine foods; even milk is to be avoided till after the first or the second week of the acute attack. This rigid regimen is necessary to diminish the special function of the kidney—elimination of nitrogenous matters. He quotes Lichtheim and Senator in support of this doctrine, which recognizes in the defective elimination of the kidney a much graver incident than the outpouring of albumen, and which, therefore, emphasizes the necessity of reducing to a minimum the ingestion of proteids. Aufrecht details one case (of scarlatinal nephritis) where suppression of urine lasted for eighty hours, and where this expectant treatment was followed by recovery. He mentions another, where the anuria lasted for fourteen hours; and he asserts that under the diuretic and diaphoretic plan of treatment cases such as these would almost certainly have succumbed. To avoid misconception, we should add that he prescribes a warm bath occasionally in such cases, but never to the extent of inducing profuse diaphoresis, and that he makes no mention of free purgation.

ALFRED L. LOOMIS, M. D., NEW YORK.

Incipient nephritis, or active renal congestion, can, in most cases, be speedily relieved by the following treatment: First, place the patient in bed, in a room with a temperature above 75° Fah., then apply a dozen dry or wet cups over the lumbar region. Let the patient drink freely of diluent drinks, and administer one or two drastic purgatives. Induce moderate diaphoresis, and avoid carefully all stimulants.

In passive renal congestion, which is generally associated with some cardiac disease, *digitalis* is the most serviceable remedial agent.

DR. HENRY B. MILLARD, NEW YORK.

This author strongly recommends rest, and the avoidance of highly nitrogenized food. Light animal broths can be given, while oysters and milk may be partaken of freely. Fruits, vegetables and cereals are suitable, while light acid wines are not objectionable. Diaphoresis he considers indispensable, and this he procures by hot air baths. After the bath, he directs affusion with moderately cold water, followed by rubbing with equal parts of alcohol and water, which removes the cutaneous sensitiveness left by the bath. The hot air bath should be employed daily or on alternate days, until permanent relief is established or so long as it seems to be of use. He uses the mild chloride and the corrosive chloride of mercury. He gives the former by combining or triturating 1 part of the drug with 99 parts of sugar of milk, the adult dose being 5 to 10 grains every two or three hours, taking care not to give it in such doses as to affect the gums or relax the bowels. Of corrosive sublimate he uses 1 part of the drug to 10,000 parts of sugar of milk; giving 8 to 10 grains at a dose. An important general distinction in the selection of remedies in acute and chronic nephritis, is that whereas a certain class of irritant and stimulating diuretics, as squills, iron, cantharides, turpentine, etc., are sometimes not only useless but dangerous in acute inflammation and recent congestion of the kidneys, in proportion as these conditions recede from an acute or recent character they will be found appropriate and serviceable. Cases of chronic interstitial nephritis *are cured or recover*, but in attempting to accomplish a cure, the practitioner will err if he rely solely upon the wise and appropriate selection of drugs. It will be fortunate if his patient be so situated that he can have the advantage of rest mentally and bodily. The influence of the latter has been shown to be of the greatest value in diminishing albuminous exudation from the kidneys. Much benefit will often be derived by keeping the patient in bed for a few days until the albumen disappears under appropriate treatment. Silk or all wool undergarments should be worn next to the skin the entire year. The milk diet is highly recommended. The best alcoholic beverages are light Rhine wines and light claret. In nephritis accompanying or caused by syphilis, iodide of potassium has proved useful. It is also useful in the nephritis produced by lead poisoning, and in the gouty kidney or the nephritis of rheumatic gout it has served a good purpose. Arsenic has sometimes proved benefi-

cial in diminishing albuminuria and in relieving headaches and nausea. *Turpentine* might be found of use where there is a feeble state of the renal circulation; it should be given in drop doses, to an adult, two or three times daily. Various *saline mineral waters* are recommended. An essential feature of chronic interstitial nephritis is that not only are anasarca and oedema usually absent, but the flow of urine is, on an average, in excess of that of a healthy kidney; consequently the occasions for the use of diuretics are rare. In enfeebled conditions of the system, however, and of the heart, they are sometimes needed; *broom*, *juniper* and simple *water* increase the amount of urea, and they become endowed with increased value in proportion as there is a diminution of urea in the urine. The carbonate, bitartrate and acetate of potash and urea should be of use as diuretics, when the secretion of urea falls below the normal.

DR. AUSTIN FLINT, NEW YORK.

In cases of acute tubal nephritis, this practitioner has derived great advantage from the free use of *water*, or milk and water, when there is a demand for nourishment. His plan is to administer the fluid in small quantities very frequently, a teaspoon or a tablespoonful every few minutes, so that from four to six ounces are taken hourly, or more if it can be given without disturbing the stomach.

A similar plan is highly recommended by Dr. W. H. DICKINSON. He combined with copious draughts of water, repeated doses of digitalis (inf. digit. f. ʒj to f. ʒiv *ter die*,) the real object being to increase the secretion of water at the kidney itself, and thus flush out the choked-up tubules. In the majority of cases thus treated, the urine will increase, the dropsy diminish, and the patient pass into convalescence without the occurrence of the various secondary evils which tend to swell the mortality of the disease.

NOTES ON REMEDIES.

Acidum Nitricum, alone or in conjunction with the proto or bi-chloride of mercury, has been sometimes found of great service.

800. R. Acidi nitrici puri, ʒ ss
 Aq. destill., ʒ ss. M.
 Three to six drops three times daily.

Acidum Phosphoricum, dilute, has also proved of value, especially after the subsidence of the most acute symptoms.

Belladonna is recommended by Dr. HARLEY in acute nephritis.

Colchicum, conjoined with magnesia, is useful in the nephritis of gouty subjects.

Convallaria Majalis. (Lily of the Valley.) Dr. MILLARD says that convallaria is likely to be of great use in nephritis with insufficient power of the left ventricle of the heart.

**Digitalis* is highly recommended by Dr. HENRY B. MILLARD.

801. R.	Tinct. digitalis,	f. $\frac{7}{8}$ ss	
	Vini scillæ,	f. $\frac{3}{4}$ iss	
	Spts. aeth. nit.,	f. $\frac{3}{4}$ ij.	M.

S.—A teaspoonful every three or four hours.

Or :

802. R.	Junip. contus,	$\overline{3}^x$
	Pulv. digitalis,	$\overline{3}^{ij}$
	Pulv. scillæ,	$\overline{3}^j$
	Vin Xerici,	Oj.

Macerate for four days and add :

Potass. acetatis,	$\overline{3}^{ij}$
-------------------	---------------------

Express and filter.

S.—A tablespoonful three times a day for an adult.

Ergot has been recommended.

**Euonymus Atropurpureus*. Dr. WM. H. HOLCOMBE, of New Orleans, describes several cases in which he used with perfect success euonymine, the alkaloid of euonymus ("Wahoo").

Enemata of three or four quarts of water, at blood heat, will, it is said, sometimes cut short an attack of nephritis. The first injection usually brings away with it much faecal matter; the second, to be given immediately after the return of the first, is usually retained without difficulty.

Helonias Dioica (unicorn plant) has been accorded curative properties.

**Iron*. Dr. MILLARD says that the chloride of iron has been of invaluable service to him in aiding to diminish the excretion of albumen, either alone or in combination with digitalis where there has been feebleness of the heart's action.

**Jaborandi*. *Jaborandi* or its alkaloid *pilocarpine* are remedies from which, in nephritis, particularly if dropsy or œdema be present, we should expect to derive benefit. Of *jaborandi*, the fluid extract is the most reliable preparation; the average dose to an adult is 30 drops every two hours. *Pilocarpine* can be given hypodermically in doses of from $\frac{1}{10}$ to $\frac{1}{20}$ of a grain, or even, very guardedly, to an adult $\frac{1}{8}$ of a grain.

Nitro-Glycerine is recommended by Drs. MAYO ROBSON and ROBERTS BARTHOLOW. The dose is one minim of a one per cent. solution (alcoholic) every half hour.

Opium is a remedy of great value, when assisted by diluents, the hip-bath, and large flaxseed poultices over the lumbar region.

Scoparius in infusion is of some value as a hydragogue diuretic.

Sodium Tannate, in doses of 10 to 20 grains three or four times daily, well diluted with water, has been used to advantage.

EXTERNAL REMEDIES.

**Baths*. The hot bath, or the hot hip-bath, is a useful soothing remedy.

Counter-Irritants over the region of the kidneys have been considered of value.

Dry Cupping has been considered beneficial.

Local Abstraction of Blood has been thought useful by several writers.

**Poultices*. Large light flaxseed poultices not only afford present relief, but arrest the progress of the disease.

URÆMIA.

DR. ROBERTS BARTHOLOW, OF PHILADELPHIA.

Probably the most conspicuous good results from the administration of *pilocarpin* have been obtained in the eclampsia of albuminuria. The powerful action on the sudoriparous glands, and the excretion of the urinary solids by the sweat, are the causes of the extraordinary relief obtained from this remedy in suitable cases. It should be borne in mind that pilocarpine tends to excite uterine action, and hence its administration may, under some circumstances, be improper, but in most cases this effect is desired. There are, however, two important contraindications; a weak heart, from thinning and atrophy of the walls of the organs, or from fatty degeneration, and a tendency to pulmonary congestion and œdema. In some instances, in a few minutes after its hypodermic use, the respiration becomes embarrassed, by congestion and odema of the lungs and by enormous bronchial secretion. It may be well to mention that atropine, subcutaneously, in one case in which it was administered, removed these symptoms and saved the patient's life.

PROF M. CHARTERIS, M. D., GLASGOW.

The objects of treatment are to increase the flow of urine, and awaken the action of the skin. Hence, diuretics and diaphoretics are in order, as:

803. R.	Tinct. scillæ,	f. ʒij	
	Liq. ammon acetat.,	f. ʒij	
	Decoctum scoparii,	ad f. ʒvj.	M.
Two tablespoonfuls thrice daily.			

And:

804. R. Liq. ammon. acetat.,
Spts. ætheris nitrosi,
Tinct. hyoseyami,
Aquam camphoræ,

f. ℥j
f. ℥ss
f. ℥ iij
ad. f. ℥vj.

M.

A tablespoonful every three hours.

During the convulsions, chloroform should be inhaled; or chloral may be injected hypodermically. In sudden attacks in plethoric persons, as sometimes in pregnancy, free venesection has much to recommend it, and should not be dismissed as absurd simply because it was the panacea of our forefathers.

DR. ALFRED L. LOOMIS, NEW YORK.

Of diuretics in acute uræmia *digitalis* stands first; but it must be given in large doses, f. ℥ss of the infusion of the leaves every three hours for twenty-four hours, or until the diuretic effect is produced. During the convulsions, chloroform should be given by inhalation. When they cease, *morphia*, hypodermically, is demanded. This will control the muscular spasms and aid the processes of elimination. The injections may be given every two hours until these effects are produced, both in puerperal and non-puerperal cases. The dread which has so often been expressed of opium in uræmic conditions is groundless; as are also the alleged "accumulative" effects of *digitalis*.

DR. LIONEL S. BEALE, ENGLAND,

lays down the rule that in blood-poisoning occurring in the course of renal disease, or when there appears any tendency to this condition, we must never give sedatives. A very little opium may destroy life in chronic renal affections. The treatment that must be adopted is free purgation and the administration of sudorifics, and where there is reason to think the kidneys will respond, diuretics, such as certain tartrates, citrates, acetates, carbonates, or nitrates; but irritating diuretics, such as cantharides, juniper, and the infusion of broom, should not be given. The frequent use of the *hot-air bath* will often keep off renal coma for a long time. This may be given by placing the patient inside of a frame, with his head out, covering the frame with blankets, and heating the interior by means of a spirit lamp, the chimney of which can be covered with an iron tube emerging between the blankets.

Headache and nausea are often early symptoms of uræmia, and should

at once be treated with sudorifics and purgatives, as elaterium, gr. $\frac{1}{4}$; this drug acts satisfactorily, but must be given with caution for fear of hypercatharsis. The vomiting may be relieved by small doses of creasote, hydrocyanic acid, ice, or carbonate of soda. All patients showing any tendency to uræmia require careful watching, and should be visited frequently.

PROF J. M. DA COSTA, M. D., PHILADELPHIA.

This teacher remarks (*Med. and Surg. Reporter*, April, 1879,) that his observation has taught him that in many cases uræmic poisoning is present in spite of the fact that the secretion of urine is quite free. In all cases of Bright's disease, where the patient is attacked with persistent vomiting, with a clean tongue, accompanied by headache, dullness, dilated pupils, and diarrhœa, there is probably incipient uræmic poisoning, no matter what the quantity of the secretion may be.

In such cases, an excellent agent is *benzoic acid*; gr. xx may be given every third hour, largely diluted with water. This acid prevents the accumulation of the urinary salts in the blood; it is a diuretic in the sense that it influences the excretion of solid material.

Such cases also require free purgation. Pills containing one drop each of croton oil, may be conveniently used for this purpose. The *vapor-bath* is also valuable, which may be given in bed by placing hot bricks, wrapped in damp towels, around the patient, under the covers; or a warm bath may be substituted if the patient is able to take it. Dry cups to the loins should not be neglected, as they aid in stimulating the kidneys.

When the benzoic acid begins to disagree with the stomach, as it probably will after three or four days, some other acid may be substituted, as the muriatic, gtt. x, three or four times a day, in an appropriate vehicle.

NOTES ON REMEDIES.

Bleeding. Dr. HIRAM CORSON has found bleeding from the arm highly beneficial.

**Chloral Hydrate.* Dr. TYSON highly recommends it.

Chloroform is considered by many an important adjuvant, pushed to its anæsthetic effects.

Morphia. In uræmic convulsions, Prof. A. L. LOOMIS recommends hypodermic injections of half a grain or more of morphia.

VI. DISEASES OF THE BLOOD.

Anæmia—Cerebro-Spinal Meningitis—Dengue (Break-bone Fever) —Gout — Hemorrhagic Malarial Fever — Intermittent Fever (Ague, Chills)—Milk Sickness—Purpura Hemorrhagica—Rheumatism, Acute—Rheumatism, Chronic—Typhoid Fever—Typho-Malarial Fever—Variola—Yellow Fever.

ANÆMIA.

DR. ROBERTS BARTHOLOW, PHILADELPHIA.

As a large consumption of the phosphate of lime takes place during suckling, the anemia of the nursing mother may be most advantageously treated with the *lacto-phosphate of lime*. The usual remedies for anemia may be much assisted by central galvanization and local applications to the vegetative organs. Beard & Rockwell employ the method termed by them "general electrization," which consists in faradic applications to the surface of the body, "one pole, usually the negative, being placed at the feet or the coccyx, while the other is applied all over the surface of the body."

In anæmia and chlorosis, *wines* render an important service by increasing digestion and assimilation. To aid in this process, red wines with a good deal of sugar and extractives are most necessary.

When wines produce headache, and the digestion is disordered by them, and the appetite impaired, they are not serviceable in these maladies. Moreover, for the nervous and hypochondriacal, wines must be prescribed with caution, for the habit of indulgence is quickly acquired by such subjects.

JULIUS POLLOCK, M. D., M. R. C. P., LONDON.

In a large number of cases, especially in women, this writer (*Medical Times and Gazette*, 1877,) extols the following combination:

805.	R.	Ferri ammonio-citratis,	gr. v	
		Rhei extracti fluidi,	gtt. v	
		Infusi quassiaë,		
		Aquæ menth. piper.,	ââ	f. 3 ij. M.

For one dose, thrice daily.

Dr. P. insists on the rhubarb, as it greatly assists the action of the iron when the stomach is out of order. Bromide of potassium, gr. x, can be added to the mixture if the patient is nervous. The diet should be light and simple. Pepsin with the meals may often be employed with advantage. A moderate amount of exercise out of doors, when the weather permits, should be insisted upon, but anything like fatigue must be avoided. A tepid bath in the morning and a rub-down afterwards with a rough towel is a good thing. By-and-by, in a few weeks, more or less, the steel and rhubarb mixture may be left off, and fifteen drops of the solution of perchloride of iron given after each meal in a wineglass of water.

Under the very best and most careful treatment, the improvement in anæmia is not unfrequently slow and tedious, and, to prevent disappointment, it is as well to caution patients and their friends that they are not to expect, as a rule, any very rapid change. Once get the patient on the road to health, however, and time and perseverance will do the rest.

DR. HUGUENIN, PARIS.

In *pernicious anæmia*, when iron given internally can no longer be absorbed, Dr. HUGUENIN (*Gazette des Hôpitaux*, August 26, 1876,) makes use of the subcutaneous method of its administration. The formula for the injection of iron is as follows:

806. R.	Ferri pyrophosph.,	
	Ammon. sulph.,	āā 3j½
	Aquæ destillat.,	3j½ M.

So that a subcutaneous syringe holds half a grain of iron. Soon after the injection, redness of the skin, œdema, and often even heart-symptoms are seen, but they soon disappear.

Dialyzed iron, fer Bravais, has claimed a high position in the treatment of anæmia, both by the mouth and by hypodermic injection. Dr. J. M. DA COSTA has used a daily injection of ℥xv-xxx, at first diluted, later undiluted, if no irritation of the skin is produced. With the best preparations of the so-called dialyzed iron, there is not much danger of abscess. French physicians have expressed doubts as to its value, believing it is not absorbed.

M. PERRONCITO, OF FRANCE,

describes a form of anæmia occurring among miners, which is due to a

parasite, the *anchylostoma duodenale*. The larvæ do not develop in the intestine, but in the fæcal substances after dejection, and penetrate the organism by the air or water after their development. The larvæ are quickly killed by a concentrated solution of chloride of sodium in sulphuric or hydrochloric acid, in ethylic alcohol, or in one to five per cent. solutions of carbolic acid, and in one half per cent. solution of thymic acid, and in ethereal extract of male fern. By any of these means the fæcal larvæ can readily be destroyed. The male fern, given internally, is always effectual, even in a single dose. Thymic acid may also be given by the stomach for the same purpose.

PROF. JOHN B. BIDDLE, M. D., PHILADELPHIA.

807. R. Quiniæ sulphatis, gr. ij
 Ferri sulphatis, gr. j
 Strychniæ, gr. $\frac{1}{60}$.
 For one pill, thrice daily.

An excellent tonic combination, frequently prescribed by our author.

TONIC BATH.

THOMAS K. CHAMBERS, M. D., LONDON.

808. R. Acidi muriatici, f. $\bar{3}$ -ij
 Aquæ calidæ, Cxxx.

In a wooden bath, the patient to remain in it for from ten to twenty minutes.

Dr. CHAMBERS has found this bath to confer muscular strength even when employed alone. When used in cases of anæmia, while iron is being given internally, it gives an impetus to the improvement of the patient; more iron is taken up, and the blackening of the fæces ceases.

TONIC INHALATION.

J. M. DA COSTA, M. D., PHILADELPHIA.

809. R. Ferri lactatis, gr. j-ij
 Aquæ destillatæ, f. $\bar{3}$ j.

For one inhalation. To be administered (by means of any form of steam atomizer throwing a fine spray) two or three times a day. Useful in anæmia, when iron given by the stomach is not assimilated. Also in cases of gastric ulcer, when the constitutional effect of this agent is called for, while it is desirable to spare the stomach.

810. R. Ferri pyrophosphatis, $\bar{3}$ j
 Aquæ bullientis, f. $\bar{3}$ ss.
 Mix and add,

Extracti gentianæ fluidi,	f. $\frac{3}{4}$ ss	
Curacoæ,	f. $\frac{3}{4}$ iss	
Vinum,	q. s. ad f. $\frac{3}{4}$ iv.	M.

Teaspoonful thrice daily.

This preparation, known as *elixir gentianæ ferratæ*, is much used and highly esteemed in this city.

DR. GUERDER, OF PARIS.

This author (*Therapeutic Gazette*, Feb'y 15, 1884,) highly recommends desiccated defibrinated blood. The mode of preparation is described as follows: The blood, taken from well-selected animals, is received at the moment of slaughter into perfectly clean vessels, is immediately defibrinated by whipping, cooked from one to two hours over a sea-water bath at about the boiling point (212° F.); it then has a reddish-grey aspect, but being exposed to the air recovers the bright red arterial color, which proves that the hemoglobin has retained all its affinity for oxygen. The cooked blood is then dried in a special apparatus. Cooking is indispensable to destroy any morbid germs which may be contained in the blood; it does not, moreover, alter the digestibility of the preparation any more than that of meat or eggs. Blood powder well prepared is of a beautiful scarlet color. It has no odor when it is sufficiently dried, and may be kept any length of time. Dissolved in water it has a slightly sickish taste, which is easily corrected.

In a glass tumbler place a teaspoonful of the powder of blood and add a teaspoonful of sugar; pour on cold water, stirring all the while till as complete a solution as possible is effected. Add a teaspoonful of some kind of liquor—rum, brandy, anise cordial, or wine—according to the liking of the patient. It may also be given in the form of an emulsion with milk, syrup, coffee, etc.

The dose should be large enough to obtain a definite result. The medium quantity is from three to four teaspoonfuls a day, representing from one-half an ounce to an ounce of the powder. In many cases this dose may be exceeded. In the usual dose the patient will absorb as food the equivalent of four to five ounces of fresh blood or of meat, from a quarter to a half drachm of soluble chlorides and phosphates, and hemoglobin in sufficient quantity to give from one to two grains of metallic iron.

PROF. EASTON, UNIVERSITY OF GLASGOW.

The following formula of our author has become very popular in

England since its first publication in AITKEN'S "Practice of Medicine," as a general tonic in anæmia and cachexia. It is known as the *Syrupus Ferri, Quiniæ et Strychniæ Phosphatum*:

811. R.	Ferri sulphatis,	℥v
	Sodii phosphatis,	℥vj-℥j
	Quiniæ sulphatis,	gr. ccxij
	Acidi sulphurici diluti,	q. s.
	Aquæ ammoniæ,	q. s.
	Strychniæ,	gr. vj
	Sacchari albi,	℥xiv
	Acidi phosphorici diluti,	f.℥xiv.

Dissolve the sulphate of iron in one ounce of boiling water, and the phosphate of soda in two ounces of boiling water. Mix the solution, and wash the precipitated phosphate of iron till the washings are tasteless. With sufficient dilute sulphuric acid, dissolve the sulphate of quinia in two ounces of water. Precipitate the quinia with ammonia water and carefully wash it. Dissolve the phosphate of iron and quinia thus obtained, as also the strychnia, in the diluted phosphoric acid; then add the sugar and dissolve the whole, and mix without heat.

The above syrup contains about one grain phosphate of iron, one grain phosphate of quinia, and $\frac{1}{2}$ of a grain of phosphate of strychnia in each drachm. The *dose* might, therefore, be a teaspoonful three times a day.

The amount of phosphate of quinia might be increased according to circumstances; and if eight grains of strychnia were employed in place of six, as in the above, the phosphate of strychnia would be in the proportion of $\frac{1}{2}$ of a grain in every fluid drachm of the syrup. A much larger dose should scarcely be ventured upon.

PROF. S. D. GROSS, M. D., PHILADELPHIA.

812. R.	Tincturæ ferri chloridi,	f.℥j	
	Quiniæ sulphatis,	gr. xx.	M.

Twenty drops thrice daily, in sweetened water, through a tube.

He prefers the tincture of chloride to all other preparations of iron.

Prof. CHARLES D. MEIGS considered *reduced iron* (ferrum redactum) to be the most efficient of the chalybeates, in two-grain doses three times a day, after each meal, on a full stomach.

DR. JOHN FORSYTH MEIGS, M. D., PHILADELPHIA.

813. R.	Ferri et quiniæ citratis,	℥iv	
	Extracti gentianæ fluidi,		
	Spiritûs lavandulæ compositi,	āā	f.℥iij
	Alcoholis,		f.℥vj
	Aquæ,		f.℥ivss.
			M.

A tablespoonful thrice daily.

PROF. ELLERSLIE WALLACE, M. D., PHILADELPHIA.

814. R. Ferri pyrophosphatis, ℥ij
 Curacoæ, f. ̄ss
 Aquæ, f. ̄ijss. M.
 A teaspoonful four times a day.

NOTES ON REMEDIES.

Arsenicum. In *progressive pernicious anæmia*, Dr. BYROM BRAMWELL (*Med. Times and Gazette*, Sept., 1877,) has succeeded after the failure of iron, cod-liver oil, etc., with liquor arsenicalis, ℥ij, three times a day.

Ferrum, in all its forms, has been largely employed, and numerous formulæ are given above. A few are here added.

Ferri Mistura Aromatica. The following formula is an excellent one :

815. R. Pale cinchona bark, powdered, ℥ij
 Calumba root, in coarse powder, ss
 Cloves, bruised, ℥ij
 Fine iron wire, ss.

Macerate in peppermint water, f. ̄xij, for three days, agitating occasionally. Filter, add peppermint water, q. s. ad f. ̄xijss; then add compound tincture of cardamoms, f. ̄ij; and tincture of orange-peel, f. ̄ss, and preserve in a well-stoppered bottle. Dose, f. ̄j-ij.

Ferri Peroxidum Hydratum. Dr. S. ASHWELL recommends the following formula as one which will often agree when other ferruginous compounds cause irritation :

816. R. Ferri peroxidi hydrati, gr. viij
 Pulveris ipecacuanhæ radiceis, gr. j
 Hydrargyri cum cretâ, gr. ij. M.
 Make one powder. Take one twice a day.

Ferri Sulphas is a very effective salt of iron in anæmia. It may be combined with gentian; with quinine and strychnia; with a sedative, as conium; with an aperient, as rhubarb or aloes; with hops, as advised by Dr. S. ASHWELL.

817. R. Ferri sulphatis, gr. j-ij
 Extracti humuli, gr. ij-ijj. M.
 Make pill. Three to be taken daily.

Albuminate of Iron by sub-peritoneal injection is recommended by Prof. A. VACHETTA (*Gazz. degli Ospitali*).

Manganesii Oxidum Nigrum has been employed with success in simple anæmia, instead of iron.

* *Oleum Morrhuæ* is a valuable remedy in all forms of anæmia, its efficacy being increased by combination with a ferruginous salt.

Oxygen. Mr. HAYEM, of Paris, has observed that, in anæmia, when oxygen is inhaled in the quantity of ten litres daily, after two or three inhalations,

there is a marked improvement in the appetite, digestive troubles disappear, and in a week or two the patient takes a substantial diet, and gains weight. But it is necessary that this treatment be backed by iron tonics, to alter the quality of the blood corpuscles and make the improvement permanent. (*Lancet*, September 6th, 1879.)

Pepsin and Pancreatine act well, by aiding digestion.

Phosphorus is valuable in combination.

Potassæ Liquor, in *pernicious anæmia*, sometimes succeeds when all other means fail. If improvement is not prompt, it should not be pushed.

Sanguis Bovinus Exsiccatus. This is a preparation of dried blood manufactured by Messrs. Parke, Davis & Co., from a formula by Dr. STEWART. It presents the nutritive constituents of healthy beef's blood in a readily assimilable form, and has been shown by an extensive experience to be admirably adapted to most cases of anæmia, and in wasting diseases.

EXTERNAL MEASURES.

Hypodermic Injections of iron have been used with advantage. Dr. J. M. DA COSTA prefers a double salt produced by the addition of pyrophosphate of iron to a solution of citrate of sodium. Two grains of the salt, in this form, are given every day, varying the points of puncture, but generally administering it under the skin of the extremities; in this form no abscesses have been observed. With other solutions of iron, including dialyzed iron, abscesses were quite common, even with every precaution as to the cleanliness of the syringe.

Transfusion of Saline Solutions. In cases of acute anæmia, Dr. M. SCHWARZ (*Berlin Klin. Woch.*, 1882,) says that if ordinary means fail, recourse may be had to a very innocent method, yet one that has proven extremely sure and active; this is direct injection of feeble alkaline solutions (6 per cent.) of chloride of sodium into the circulatory system. The action of this transfusion on cardiac activity, blood pressure, respiration, and all the other vital functions, has shown itself with surprising rapidity in rabbits and dogs who had lost from one-half to two-thirds of the entire quantity of blood contained in the body. The minimum quantity of liquid to be injected in the adult should be about five hundred cubic centimeters. This species of transfusion is also indicated in the serious collapse coming on during operations on the abdomen. Since the above conclusion appeared in the inaugural thesis of the author, five cases of transfusion of chloride of sodium solutions have been reported by BISCHOFF, KUSTNER, KOCHER and KUMMEL.

CEREBRO-SPINAL MENINGITIS.

This disease, known also as "spotted fever," "epidemic meningitis," etc., is more frequent in children than adults, but by no means confined to those of tender years. In its epidemic forms, it is frequently very fatal at all ages. The following is the treatment pursued by

DR. NATHAN S. DAVIS, CHICAGO.

818. R.	Tincturæ physostigmatis,	f. $\frac{3}{4}$ j	
	Extracti fluidi ergotæ,	f. $\frac{3}{4}$ iss.	M.
Half a teaspoonful every two hours.			

The *Calabar bean* is used very freely by Dr. DAVIS, in this disease, and with more apparent effect in controlling it, than any other remedy he has tried. The pulse becomes slower, the respiration more regular, the patient quieter, the pain less intense. If the disease assumes a chronic form, with transient and wandering, but severe neuralgic pains, a fretful condition of mind, variable appetite, debility, and disturbed sleep, he has obtained very decided and permanent relief from

819. R.	Tincturæ physostigmatis,	f. $\frac{3}{4}$ j	
	Tincturæ opii camphoratæ,	f. $\frac{3}{4}$ iss.	M.
Half a teaspoonful three times a day; and at bedtime, a moderately full dose of Dover's powder and quinia.			

DR. D. C. RAMSEY, OF ST. LOUIS.

The *salicylic acid* treatment is strongly advocated by this practitioner in the *St. Louis Med. and Surg. Journal*, Feb., 1884. He states that this drug exerts a direct influence for good over the inflammation itself, and can be taken in frequent large doses without bad effect: having given a boy 15 years of age half-drachm doses every four hours for three or four days, with the only result of a great benefit in all the symptoms connected with the disease, is, he thinks conclusive evidence of its harmlessness.

Its good effects are soon apparent, and it does not interfere with the use of other measures of relief, as ice, blisters, etc.

The best mode of using the remedy is to administer large doses frequently. For adults begin on doses of 15 grs. repeated every two hours, and increase the dose as may be found necessary to obtain the

desired effect, to \mathfrak{D} ij., at intervals of two hours, if need be. When the disease is under control, which will be determined by the reduction in temperature, relief of pain, and placid countenance, decrease the dose, give at longer intervals, but still continue the use of it in small doses as long as the least symptom is present indicative of the disease.

PROF. FELIX VON NIEMEYER, M. D., TUBINGEN.

In the treatment of *acute meningitis*, it is not generally proper to bleed from the arm, but *leeches* may be applied to the brow and behind the ears, and, if the strength of the patient permit, the application may be repeated. The shaven head may be covered with *cold compresses*, and an active *purge* of calomel and jalap administered, viz.:

820. R.	Hydrargyri chloridi mitis,	gr. ij	
	Extracti jalapæ,	gr. viij.	M.
Divide into two pills, and order both to be taken.			

In the latter stages of the disease, if there be coma and other signs of cerebral palsy, apply a large *blister* to the nape of the neck, and rub the following pustulating ointment on the head:

821. R.	Olei tiglii,	$\mathfrak{m}_{\text{xxv}}$	
	Adipis,	\mathfrak{z}_{ss}	M.

One-fourth part to be rubbed into the skin every eight hours, until an abundant eruption is produced.

Still more efficacious than these derivatives, are *douche-baths*—pouring cold water over the head from a pitcher held some distance above it. The patient almost always recovers consciousness as this is being done, but it must be repeated at intervals of a few hours, to secure a permanent result; with each successive employment, the number of pitcherfuls is to be increased. Frictions with *mercurial ointment*, and continued doses of *calomel*, are much employed.

PROF. J. LEWIS SMITH, M. D., NEW YORK.

This author, in speaking of this disease, especially as it appears in children, disapproves of blood-letting, even by leeches. The one drug he relies upon, is *bromide of potassium* in doses of gr. v-vj every two, three, or four hours, to a child of five years, according to the urgency of the case. After the first week it should be given less frequently, and finally omitted. Small doses of hydrate of chloral are best to relieve the pain. Quinia is of no service. Bismuth, in large doses,

controls most effectively the nausea. Frequent counter-irritation along the spine by dry cups or an irritating liniment is useful from the first, and also vesication of the nucha later in the disease. Sustaining measures should be commenced early.

PROF. WILLIAM FULLER, M. D., MONTREAL,

urges strongly (*Canada Medical Record*, September, 1877,) *hot water* to the head, instead of cold, and the use of *morphia*. The plan he adopts when he suspects meningitis is: 1. To regulate the temperature, that is, keep the patient cool, but equally warm all over. 2. Unload the bowels by a mild purgative, remove worms, etc., if any cause of irritation is present. 3. Quietude and soothing medicines, the use of which is indicated by flushings and irregular cutaneous circulation, which are the premonitory symptoms of spasms or coma; should the latter symptoms, spasms or coma, supervene, he increases the dose. 4. Frequent bathing the head with warm water. The points he insists upon are: 1. That coma is, in most instances, not due to the pressure of effusion, but to irritation. 2. That *opiates* are not contra-indicated in meningitis, even when coma is present, or threatened, if there are irregularities of cutaneous circulation or spasms, and that opium actually, by relieving irritation, dissipates the coma. 3. That warm water is more agreeable, more soothing, and more efficacious than ice-caps.

DR. J. B. HAMILTON, NEW YORK.

In the *New York Medical Journal*, February, 1875, this writer gives his treatment as follows: For the first twenty-four or forty-eight hours:

822. R.	Extracti ergotæ fluidi, Spiritus ammoniæ aromatici,	℥j ℥ij	M.
A teaspoonful in a little water every four hours.			

823. R.	Potassii acetatis, Aquæ camphoræ,	℥xij ℥vj.	M.
A tablespoonful every two hours, until diuresis is produced.			

In addition to these remedies, direct a warm bath, followed by wrapping in flannel; or rubbing with dry mustard, every three, four, or six hours, according to the urgency of the case. Stimulants may be necessary from the outset.

As soon as the symptoms show any amelioration, the ergot mixture

may be diminished in frequency, and at the third or fourth day discontinued. Quinine in large doses will then be found of advantage, and a more stimulant diuretic, as spiritus ætheris nitrosi, may be substituted for the potassa acetate; and for the sequelæ nothing has proved more serviceable than *iodide of potassium*, as occasion demanded. This writer considers the bromide of potassium treatment "utterly worthless."

DR. WILLIAM READ, BOSTON, MASS.

One of the most rational and also successful methods of treatment of this disease is described by Dr. READ (*Medical and Surgical Reporter*, vol. xxx, page 469). He usually commences with a full dose of castor oil, or other mild laxative, and then prescribes:

824.	R.	Ergotinæ (Bonjean's),	gr. x	
		Extracti belladonnæ,	gr. j-ij.	M.
Make ten pills. One every three hours.				

Croton oil is applied to the nucha until a copious eruption is produced, and if the spine is tender it is blistered. Concentrated nourishment is ordered.

PROFESSOR VON ZIEMSEN, GERMANY.

The treatment of this physician comprises leeches behind the ears, and ice-bags to the head, neck and back; the inunction of fifteen to thirty grains of mercurial ointment twice daily; frequently rinsing the mouth with chlorate of potash solution, to prevent stomatitis; and especially *morphine*, in full doses, gr. $\frac{1}{3}$ —ss, hypodermically in adults. Quinine is useful only in those rare cases where the temperature ranges very high. It is then required in doses of gr. xv—xxx. In a later period of the disease, *iodide of potassium* is of great service in producing absorption of the exudation.

NOTES ON REMEDIES.

Aconite. BARTHOLOW says that in acute inflammation of the cerebral and spinal meninges, and in cerebro-spinal meningitis before effusion has taken place, aconite is as serviceable as in other acute inflammations. It is generally advisable to combine opium with it, especially in cerebro-spinal meningitis.

Antimonii et Potassii Tartras has been exhibited in repeated fractional doses with benefit. It is contra-indicated if the patient be very young, if there have been prodromic symptoms, and if the inflammatory condition be not well marked.

826. R. Tinct. cinchonæ,
Tinct. gentianæ,
Tinct. calumbæ, āā f. ℥ij. M.
Tablespoonful every two hours, in a little water, until four doses are
taken, every morning.

If, on the contrary, the pulse is hard and full, and the fever sthenic, tincturæ aconiti fol., gtt. v–xv, may be given every few hours. When the tonsils are enlarged, and the mouth and fauces show a papulous eruption, the following is useful :

827. R. Tinct. capsici,
Tinct. guaiaci, āā f. ʒj.
A teaspoonful in a cup of water, as a gargle.

Relapses are common, and convalescence usually tedious. The above-mentioned tonic is advisable in such cases.

SIR J. FAYRER, M. D., K. C. S., ETC.

This distinguished practitioner, speaking of the treatment of dengue, as he has observed it in India, (*The Practitioner*, Dec., 1876,) expresses his opinion that neither emetics nor active purgatives are necessary. They do no good, but increase the weakness and aggravate the suffering by the muscular movements necessarily induced. Moderate action of the bowels, followed by a warm carminative aperient, with an occasional dose of calomel, rhubarb, or colocynth, especially if they remain confined, to which there does not appear to be any peculiar tendency, though the alvine evacuations are dark, and often slimy and confined, at the outset; salines, such as the acetate of ammonia, citrate of potash, with nitrous ether, combined with aconite, are good in the pyrexia. In cases of very high temperature, 105°, 106°, 107°, cold sponging is beneficial.

Belladonna seems to confer great relief in this disease ; 10 to 15 drops of the tincture may be given, and two or three such doses, given at intervals of an hour, will sometimes produce excellent effects, and give much relief. The extract may be given, if prepared in doses of $\frac{1}{3}$ of a grain ; or the juice, in similar doses to those of the tincture.

For the pains and nocturnal restlessness, *morphia* or Dover's powder may be given; and, as external applications, liniments containing opium, belladonna, and chloroform, are serviceable, rubbed on the spine, back, and joints.

G O U T .

DR. HENRY W. FULLER, LONDON.

For the purpose of clinical instruction, Dr. FULLER divides cases of acute gout into two classes, namely: 1. Cases in which the excretory organs are originally sound and functionally active—cases in which the attack of gout is due principally to excess and indiscretions of diet; and 2. Cases in which the excretory organs are in some way disordered, and fail in performing their eliminatory functions—cases in which the patient is not necessarily guilty of indiscretions of diet, but in which the liver and kidneys fail in their action, either as the result of functional disorder, or of organic change in their structure.

The first class of cases correspond with those which pass under the name of asthenic gout; the tongue is usually furred, the urine loaded, and the bowels are commonly torpid. In these cases, until the acute symptoms have subsided, Dr. FULLER restricts the diet to liquids, administers a saline draught containing sulphate and carbonate of magnesia, and a few drops of colchicum wine; occasionally gives an aperient pill containing calomel, aconite, and opium, and wraps the joints in finely-carded wool, or in flannels steeped in a solution of soda and laudanum. As the acute symptoms subside, a more generous diet is permitted, and some light bitter tonic, such as tincture of gentian or calumba, is added to the mixture.

The second class of cases have more affinity with what is termed atonic gout; the tongue is often clean and the urine clear—sometimes of low specific gravity—and the bowels are regular. In these cases, Dr. FULLER does not restrict the diet to the same degree; he allows a little meat without vegetables, and also, if desired, a glass of sherry or a little spirits and water. He acts freely on the skin by means of the hot-air bath; administers an aperient in the morning, containing taraxacum and sulphate of magnesia, and, during the day, he gives a warm stomachic draught, containing ammonia and a few grains of soda in a light bitter infusion. Occasionally, a dinner pill is prescribed, containing rhubarb and a grain of colchicum; and in some instances—characterized by pale, clear urine—a draught containing quinine, the mineral acids, and taraxacum, is substituted for the mixture just referred to. In these cases as the acute symptoms have subsided, a drachm

of the syrup of phosphate of iron is given each morning before breakfast.

DR. H. SALTER, LONDON.

The *local treatment* of this author is all that is peculiar. It consists in the application of the following lotion:

830. R.	Potassii iodidi,	ʒj	
	Potassii bicarbonatis,	ʒj	
	Aquæ bullientis,	Oj.	M.

To this a little tincture of opium may be advantageously added.

Double lint, saturated with this lotion, is applied to the part affected, and covered with oil-silk; to that is put a layer of cotton-wool, and the whole swathed in a flannel bandage. The lint should be taken off from time to time, and re-dipped in the lotion. The relief that the patients experience from this application is very great. With or without this lotion, there are three other things on which Dr. SALTER insists, in the local treatment of a gouty joint—perfect physical rest, protection, and preventing the part affected being too dependent.

PROF. S. D. GROSS, PHILADELPHIA.

831. R.	Vini colchici radicis,	f. ʒj	
	Morphiæ sulphatis,	gr. j.	M.

For one dose, at bedtime, in gouty affections of the joints.

This treatment should be preceded by purgation or venesection, if indicated, and be followed in the morning by a gentle laxative. These doses are recommended by our author as the most efficient, and as seldom disappointing the most sanguine expectations. Where there is a full, bounding pulse, indicating excessive arterial action, then the following will come into play:

832. R.	Tincturæ aconiti radicis,	f. ʒj	
	Morphiæ sulphatis,	gr. ij	
	Antimonii et potassii tartratis,	gr. j	
	Aquæ,	f. ʒjss	
	Syrupi zingiberis,	f. ʒ ss.	M.

A teaspoonful every three hours.

Veratrum viride may be substituted for the aconite, in the same, or double the dose. The action of these potent remedies should, of course, be carefully watched, and kept within proper limits. Together with the above means, the following should be employed, to neutralize the acid state of the blood:

833. R. Potassii bicarbonatis, $\bar{3}j$
 Sodii bicarbonatis, $\bar{3}ij$. M.
 For six powders. One to be taken every six hours, in a wineglassful of water.

As a local application, nothing will be found better than :

834. R. Tincturæ opii, f. $\bar{3}j$
 Linimenti saponis, f. $\bar{3}ij$. M.
 To be rubbed in twice a day, and constantly kept in contact with the affected joint, by means of a piece of flannel covered with oiled silk. A fly-blister may be used, if the disease manifests a disposition to linger.

PROF. A. B. GARROD, KING'S COLLEGE, LONDON.

835. R. Lithii carbonatis, gr. iv
 Aquæ rosæ, f. $\bar{3}vj$. M.

This solution is to be warmed, and then imbibed by means of charpie or a piece of sponge, and applied on the seat of pain, and covered by a sheet of gutta-percha. Two or three times a day, the application is to be wet with the solution, in order to keep it always moist. Internally, the carbonate of lithia, in doses of gr. $ij-vj$, or the citrate, in doses of gr. $v-x$, in aerated water.

836. R. Fraxini foliæ, $\bar{3}ss$
 Aquæ, Oj.

The leaves are to be boiled in the water during ten or fifteen minutes, then strained and sweetened. To be taken in fractional doses during the day, an hour before the meals, in chronic gout.

The leaves of common ash, though much extolled in France and Germany, in the treatment of gout and rheumatism, have not succeeded in the hands of our author, in acute gout; when taken for a long time, and in large quantities, in the form of decoction of the leaves, they have, probably, some influence in keeping off the attacks in chronic gout.

DR. HENRY G. PIFFARD, NEW YORK.

In the treatment of the gouty dyscrasia, the two principal indications are, 1, to depurate the blood, and 2, to promote oxidation.

1. Depuration of the blood is to be effected by calling into more vigorous action, either the kidneys, bowels, or skin. If the trouble is due to defective renal activity, a point which may be determined by careful estimation of the amount of nitrogen daily discharged in the urea, uric acid, etc., we must treat these organs with some leniency,

and be careful not to urge them too much, since, by undue stimulation, we may increase the difficulties under which they are laboring. We shall be obliged, therefore, to depend upon the skin and bowels to fulfill the first indication. The functional activity of the skin is increased by exercise, bathing, and warmth, and is most rapidly and vigorously influenced by the hot-air or Turkish bath.

If the bowels are to be stimulated, we may employ various cathartics, the most useful in this connection being the ordinary *senna* and *salts*, given in sufficient doses to produce one or two loose evacuations daily. Professor HARDY, of Paris, praises very highly an infusion of *wild pansy*, *viola tricolor*, combined with *senna*, in about the following proportions:

837. R.	<i>Violæ tricoloris</i> ,	$\bar{3}j$	
	<i>Sennæ</i> ,	$\bar{3}^{ss}$	
	<i>Aquæ bullientis</i> ,	<i>Öij.</i>	M.
For an infusion. Filter when cold.			

One-quarter to one-half of this quantity is to be taken daily, and the amount to be diminished gradually, as the necessity for its employment lessens. Professor HARDY states that he has given this purgative two or three months at a time without ill effect.

In place of these remedies we may employ some of the natural mineral waters, as those of Seidlitz, Pullna, or of Friederichshall. The native waters which seem to approach most nearly to these in composition, are the Estill and Crab Orchard Springs, in Kentucky.

It is, however, but rarely that active purgation is required. If the kidneys are perfectly healthy, we may call upon them to perform the work, and leave the bowels alone. To increase the renal activity we have recourse to diuretics. The ones especially serviceable in this connection, are *vinum colchici*, *infusum digitalis*, *balsam of copaiva*, *propylamine*, *carbonate of lithia*, Vichy water and *wine of green coffee*. The latter valuable preparation is prepared as follows:

VINUM CAFFEEÆ VIRIDIS.

838. R.	Ground, unburned coffee,	One pound
	Good sherry wine,	One quart.
Digest for two weeks. Filter through flannel, press the residue, and add wine to make a quart. Dose, f. $\bar{3}j$ -ij per diem.		

These remedies, one and all, exert a marked influence in increasing the action of the kidneys, and are among our chief reliances in general

gouty and rheumatic conditions. They may be used singly or combined, and for a considerable period. The alkalies, however, have a tendency to impoverish the blood. A good plan, if we anticipate a prolonged use of them, is to intermit their employment for one or two weeks out of each month, giving iron, if necessary, in the intervals.

If incomplete oxidation is owing to an excessive amount of nitrogenized food, we must rely chiefly on dieting. Meat should be cut off to a greater or less extent, and in place of it a larger quantity of bread, vegetables and fat should be consumed.

If, on the other hand, the patient has been accustomed to but a moderate quantity of meat, it will be necessary to institute measures specially designed to promote oxidation. The use of preparations of *iron* here takes the first place. But in addition to this we must endeavor to supply a full amount of oxygen by exercise in the open air, a well-ventilated bed-room, and the like. In addition, we may attempt to furnish oxygen directly to the blood, by inhalations of the pure gas, or partly oxonized. Further, we may employ certain medicines which contain oxygen largely, and are believed to be capable of giving it up to the blood, as for instance, the *chlorate of potassium*. Finally, if the liver be torpid, we may have recourse to the occasional or frequent use of those drugs which have the reputation of hepatic stimulants, as *mercury*, *podophyllin*, etc.

MINERAL WATERS.

The prolonged and moderate use of the natural mineral waters has been spoken of as one of the most efficient corrections of the arthritic dyscrasia. They must be used with a proper discrimination, if good effects from them are expected.

The waters chiefly employed for this purpose contain either alkaline carbonates, chlorides, or sulphates.

The *Vichy* waters, rich in carbonate and bicarbonate of soda, appear to be well adapted for strong subjects in whom the functions of the liver and digestive organs are at fault; but are contra-indicated when there is a tendency to the formation of chalky concretions about the joints, and in debilitated constitutions.

Waters containing an abundance of chlorides, such as those of *Wiesbaden*, are less debilitating and more stimulating than those of *Vichy*; they are consequently better adapted for cases in which the circulation

is sluggish, and the secretions deficient; also, in cases of rigidity from thickening of the textures.

The springs, rich in sulphates, a typical example of which is those of *Carlsbad*, act with considerable force on both the bowels and kidneys. They are often very useful in cases accompanied by deficient action of the bowels with a congestive state of the liver; but should be avoided by weakly patients.

In addition to these, the waters containing *lithia* are claimed to have a peculiar and almost specific power in removing the chalky concretions of urate of soda so frequent in this diathesis. In Europe, the springs of Baden-Baden, and in the United States, the *Gettysburg* water, have acquired particular reputation in this direction.

In all cases the waters should be commenced cautiously, small quantities being taken at a time, with care not to oppress the stomach by the presence of too much liquid; and whenever there are symptoms of exciting febrile disturbance in the system, their use should be suspended.

NOTES ON REMEDIES.

Aconite. Professor GROSS recommends tincture of aconite root, with morphia and antimony, when there is excessive arterial action.

Ammonii Phosphas has been recommended, in ʒj dose daily, highly diluted. Dr. GARROD has derived much benefit from the long-continued use of this salt in chronic conditions of the gouty habit.

Arsenic. Prof. DA COSTA teaches that arsenic, in small doses, is very useful in chronic gout.

Arnica. In rheumatic gout very decided curative effects are sometimes procured from arnica; very probably owing to the fact it contains trimethylamine. (BARTHOLOW.)

Cajuputi Oleum, in doses of gtt. v-vj, frequently repeated, is serviceable in retrocedent gout.

** *Colchicum*, in gout, has been compared to cinchona in ague, but unfortunately, unlike the latter remedy, it does not prevent a return of the disease. Dr. A. B. GARROD states that often an almost magical change is produced by a single large dose, without the least increase in any of the secretions, the effect showing itself in the rapid decrease of the pain, and other symptoms of the joint inflammation. Dr. N. S. DAVIS, of Chicago, recommends forty drops of an equal mixture of the acetated tincture of opium and wine of colchicum seeds to control acute paroxysms of gout. This dose may be repeated in an hour, if necessary. Oftentimes, one or two doses will abort what threatens to be a very severe attack. When the paroxysm is under control, the same

remedies may be continued in smaller doses, three or four times daily, if any gout remains.

Frazini Folia. Ash leaves, in infusion, have been used by Dr. A. B. GARROD with advantage, but, as he remarks, the amount of the infusion given daily being considerable, the action of the diluent must not be forgotten. (F. 836.)

Guaiacum is considered by Dr. A. B. GARROD as especially useful in the asthenic gout of old subjects. It may be given continually, for a long time, without injury. This remedy is much employed in Germany.

Lithii Carbonas is a valuable remedy. So, also, are the natural lithia waters of Baden-Baden. (F. 835.)

**Lithii Benzoas.* The benzoate is probably superior to all other preparations of lithium in the treatment of the uric acid diathesis. It is readily soluble in water, and the benzoic acid, by its transformation into the nitrogenous hippuric acid, counteracts the secretion of the uric acid. Instead of the rather insoluble urates, which are generally regarded as the cause of the gouty affection, appear the readily soluble hippuric alkalies and earths, which are excreted from the organism with the urine.

Lithii Iodidi. Iodide of lithium is said, by a writer in the *British Medical Journal*, 1883, to be the most useful of the lithium salts in the treatment of gout. Not only is it serviceable in well-developed cases, but also in the varieties of eczema and dyspepsia which are of gouty origin.

Magnesii Carbonas, dissolved in an excess of carbonic acid, is an excellent manner of exhibiting magnesia, which is often of great service in chronic gout.

Manganese. Cachectic states arising from gout are successfully treated with manganese in combination with the syrup of the iodide of iron.

Manna is an excellent aperient, in the following combination :

839. R.	Mannæ,	3 iss	
	Vini colchici,	℥xx	
	Potassii tartratis,	3 ij	
	Tincturæ cardamomi compositæ,	f. 3j	
	Infusi sennæ compositi,	f. 3 x.	M.

For one draught.

Muriaticum Acidum Dilutum is strongly recommended by Dr. J. F. DUNCAN (*Medical Press*, May 2d, 1866,) as acting more directly upon the primary cause of the disease, the presence of lithic acid in the blood, and holding out a greater prospect of effecting a real cure than any antacids. He directs the following mixture :

840. R.	Acidi muriatici diluti,	f. 3 iss	
	Spiritus chloroformi,	f. 3 ij	
	Tincturæ colchici,	f. 3j	
	Infusi cascarillæ,	f. 3 vj.	M.

Two tablespoonfuls every three hours.

Oleum Olivæ is strongly recommended by Dr. J. SPENCER RAMSKILL, of London. *Potassii Iodidum* is frequently indicated in chronic gout. When fluid effused into the joint cavities is slow of absorption, the iodide often acts to great advantage.

Potassii Nitræs is recommended by Prof. BENNETT, of Edinburgh, in acute gout.

Quiniæ Sulphas. From his experience in a half a dozen cases, Dr. C. W. SCHÖNEMAN (*Pacific Medical Journal*, May, 1879,) believes that quinine will cut short an attack of gout. He gives :

841. R. Quiniæ sulphatis, gr. iij
Sodii bicarb., gr. xij. M.

This amount every two hours during the day.

No other medication is needed, though the joint may be painted with tincture of iodine.

**Rheum*, taken regularly between the attacks, often wards off a return. At the period of an impending paroxysm, Professor GRAVES, of Dublin, advises the following :

842. R. Pulveris rhei, ℥j
Aurantii corticis, ℥ij
Pulveris aloës c. canella, ℥ij
Spiritus vini gallici, Oiv. M.

A tablespoonful of the strained liquor, in water, night and morning.

**Salicylic Acid* is very useful in acute gout.

Sulphur Waters and Baths are useful in gout.

EXTERNAL APPLICATIONS.

Aconitia, applied locally, over the seat of the pain, is an excellent palliative.

Argentæ Nitræs. Dr. FROMMULLER, for gouty pains, rubs a solid stick of nitrate of silver over the moistened surface for about thirty seconds; when dry, the skin is covered with a compress of salicylated cotton, which is removed once or twice a day until the blister is healed. In order to be efficient, the irritation should be sufficient to cause an exudation beneath the epidermis.

Alcohol. Sir C. SCUDAMORE recommends a tepid lotion of one part of alcohol and three of camphor mixture.

Cajuputi Oleum, externally, is of much benefit.

Colchicum, locally, to the affected part, is highly praised by Dr. LAYCOCK.

Iodoform, externally applied, has succeeded, in the hands of some practitioners, in relieving the pain.

**Iodine*, locally used, is a very valuable remedy. The following formula is said to afford, sometimes, almost instant relief :

843. R. Iodinii, ℥ij
Alcoholis, f. ℥j
Aquæ, f. ℥vj-f. ℥j. M.

Dr. J. PEREIRA says that no external remedy gives such relief as iodine, but that the skin of different individuals is most unequally susceptible to its influence; in some it excites great pain, while in others it produces scarcely any. It is especially useful when effusion into the synovial membranes or sheaths has taken place.

Morphia. A solution of morphia, applied by means of lint, often affords great relief:

844. R. Morphiae muriatis, gr. viij
Aquaë, f. ʒj. M.

Pressure. A number of recent writers have borne strong testimony to the value of gentle and equable pressure over the affected limb, in gout. Some diachylon should be spread on felt, and over this, a flannel bandage be firmly and equably applied. The great art in this treatment is to have the pressure evenly distributed.

Sinapisms to the inflamed part often afford speedy relief. They are also useful in retrocedent gout, in inviting a return of the disease to its original seat.

Sodii Phosphas. The finely-powdered bibasic phosphate of soda, sprinkled over moistened spongeo-piline, and applied so as to envelop the whole of the affected part, affords immediate relief in many cases.

Sulphur. The exposure of the bed-clothes to strong sulphur fumes, before they are spread over the patient, is recommended to induce copious perspiration and relief of pain.

Tabaci Folia. The application of moistened tobacco leaf often affords great relief.

Veratria, in ointment, is of occasional service, and is recommended by Sir C. SCUDAMORE:

845. R. Veratriæ, gr. xxx
Unguenti, ʒj. M.

Zingiber. A ginger plaster placed over the epigastrium, relieves the pain of gout in the stomach.

HEMORRHAGIC MALARIAL FEVER.

DR. R. D. WEBB, ALABAMA.

The indications of treatment are—to arrest the paroxysms; to relieve the nausea and restlessness, so as to facilitate the administration of medicines, and add to the comfort of the patient; to sustain the patient by appropriate food and stimuli; to assist the emunctory func-

tions of the liver and skin; and, in the latter part of the disease, to promote the action of the kidneys, so as to avoid the accumulation of deleterious matters in the blood, and to prevent a recurrence.

He usually commences with a mercurial purge, and follows with a full amount of quinine, gr. xxx, in four doses, one every two hours, if the stomach will retain it. If it will not, the drug should be used hypodermically. Chloral hydrate and bromide of potassium, by enema, will lessen the gastric irritability; or small doses of morphine may be given for that purpose. Of the objections to quinine, on the ground that it increases the blood in the urine, Dr. WEBB says: "I have ever looked upon the mere fact of blood in the urine as of minor importance, and have never directed remedies specially to the arrest of this symptom. I believe it to be dependent upon the condition of the nervous system, and, by directing my remedies to the source of the evil, I expect to remove the *cause*, and the *effect* will cease. This, I believe, is best accomplished by the use of quinine; and if, in its administration, the hæmaturia is temporarily increased—as I have no doubt it is, at certain times—I do not, upon this account, hesitate to use it. I regard it here, notwithstanding this temporary increase, as the great hæmostatic, by its power of removing or counteracting the condition upon which the hæmaturia, in common with other symptoms, is dependent." (*Hemorrhagic Malarial Fever—A Monograph*, Selma, 1876.)

DR. THOMAS J. PUGH, LOUISIANA.

This writer states that he became dissatisfied with quinine and mercury in this troublesome form of malarial poisoning, and now uses instead:

846.	R.	Vini ergotæ,	f. ʒj	
		Tinct. ferri chloridi,	gtt. xx	
		Olei terebinthinæ,	gtt. v	
		Spiritus etheris nitrosi dulcis,	gtt. xx.	M.

This amount, in a gill of water, every two hours, until the urine clears.

After this omit the ergot and oil of turpentine, but continue the iron and nitre, and add to them a preparation of nux vomica—the fluid extract—in five-drop doses, every four hours, until the patient is safe from a recurrence of the disease, and then direct it every eight hours, or before meals. The danger in this disease is not from an excessive

flow of bloody urine, but rather from a suppression of urine. (*Atlantic Med. and Surg. Jour.*, Jan., 1877.)

For *Notes on Remedies*, see under "Intermittent Fever."

INTERMITTENT FEVER (AGUE, CHILLS).

THE PERSONAL PROPHYLAXIS OF MALARIA.

The general belief that the cinchona alkaloids are reliable prophylactics against the malarial poison, is so far from correct that it should be renounced. It is even dangerous, for it has been well established by the observations of the English surgeons in West Africa and elsewhere, that when long used as a preventive, it not only loses its prophylactic, but also its therapeutic powers. (See Surgeon GORE's *Medical History of the West African Campaign*, page 164.) The best prophylactics are a generous diet, plenty of work, mental and physical, but not to exhaustion, avoidance of night air and wet, proper clothes, and a cheerful disposition. The eucalyptus, in the form of infusion or extract, and infusion of box leaves, have been lately advocated on respectable authority.

The free use of pepper in the food, and the habit of taking a small cup of strong coffee on rising, may be favorably mentioned. During the night, doors and windows should be closed, and an open fire should be lighted in the evening and early morning. The drinking water should be filtered, or thoroughly boiled before using.

From the experience of English surgeons in India, (*Indian Medical Gazette*, March, 1871,) *quinoidine* is the alkaloid of bark that ranks highest in prophylactic qualities. Three grains daily was the amount prescribed. Small and regular doses of *strychnia* have been highly praised as a prophylactic by Dr. C. W. WAYLEN, of the East India Service.

PROF. S. M. BEMISS, M. D., NEW ORLEANS.

This writer (*New Orleans Medical and Surgical Journal*, Sept., 1876,) states that in the treatment of malarial fever, three objects must be held in view: 1. To restore to the blood its normal constituents; 2. To rid it of impurities; and 3. To ward off malarial paroxysms.

pound extract colocynth, and castile soap; make two pills, and give at one dose. Restorative and eliminant remedies may be combined, as:

851. R.	Magnesii sulphatis,	\mathfrak{z} j-ij	
	Ferri sulphatis,	\mathfrak{z} ij	
	Acidi sulphurici diluti,	f. \mathfrak{z} j	
	Syrupi aurant. cort.,	f. \mathfrak{z} vij	
	Aquam,	ad f. \mathfrak{z} vj.	M.

A teaspoonful, in water, twice daily.

The third indication is best subserved by the preparations of cinchona. These may be given habitually, as, for example, from 3 to 5 grains of quinine in 1 or 2 ounces of coffee daily.

The alimentation of anæmic malarial patients, is a most important consideration. They should be liberally supplied with carefully-selected and well-prepared animal food, to which such vegetables and fruits may be added as are regarded suitable to their condition. Digestion may be aided by acid solutions of pepsin, or by mineral acids, diluted in some bitter infusion. The list of promoters of digestion is incomplete if we omit alcoholic drinks, or, for many patients, coffee and tea. A good prescription is the following:

852. R.	Tinct. cinchonæ comp.,	f. \mathfrak{z} iv.
	Dessertspoonful, in water, after each meal.	

Whenever considered indicated, from 5 to 8 drops of nitro-muriatic acid may be added to each dose. Alcohol may also be given, in the form of wine or beer.

In violent forms of *congestive chill*, or *algid pernicious malarial fever*, Dr. BEMISS states that opium, chloroform, belladonna, chloral hydrate, and bromide of potash, prove more or less valuable, according to the idiosyncrasy of the patient, or the circumstances under which they were used. Opium is the most valuable of the agents, and may be combined with other drugs, as an adjuvant. It is better to use it in moderate doses, and repeat these at short intervals. Twenty drops of laudanum, with $\frac{1}{2}$ teaspoonful of chloroform, is an efficient prescription. One-sixth of a grain of morphia, with $\frac{1}{40}$ of a grain of atropia, may be injected subcutaneously. Nitrite of amyl is highly recommended. These remedies may be used at any stage of the chill, without fear of prejudicing the subsequent career of the case.

DR. ROBERTS BARTHOLOW, OF PHILADELPHIA.

In chronic malarial infection, important changes have been produced in the intestinal canal, liver, spleen, kidneys, cerebro-spinal axis: the paroxysms of fever occur irregularly; various abnormal manifestations of the infection take place (dumb ague, enlarged spleen, etc.) Under these circumstances, quinine is less curative than when the infection is recent, and the paroxysms will recur from time to time notwithstanding its use, unless these structural alterations are corrected. In chronic malarial disease, salicylate of quinine and salicylate of cinchonidine are especially effective:

858. R. Cinchonidinæ salicylat., $\overline{3}$ ij
 Acidi arseniosi, gr. j
 Ferri sulph. exsic., $\overline{9}$ j. M.
 Ft. Pulv. (Wafers) No. xx. S.—One three times a day.

Also:

859. R. Chinoidin, $\overline{3}$ ij
 Quiniæ salicylat., $\overline{2}$ j
 Ferri sulph. exsic., $\overline{9}$ j. M.
 Ft. Pil. No. lxxx. S.—Two pills three times a day.

Or:

860. R. Quiniæ sulph.,
 Chinoidinæ, $\overline{a}\overline{a}$ $\overline{3}$ j
 Res, podophylli, gr. v
 Ferri sulph. exsic., $\overline{3}$ ss. M.
 Ft. pil. No. lx. S.—Two pills three times a day.

As an antipyretic this author prefers the benzoate of sodium to the salicylate of the same, because it is safer. In the convalescence from malarial fever and in chronic malarial poisoning, the various bitters are useful in promoting constructive metamorphosis. Dogwood possesses the most positive antiperiodic properties, and is considered by the physicians of the southern States as next to quinine in efficiency. An excellent tonic combination of decided utility in chronic malarial disease is the following:

861. R. Dogwood bark,
 Calumba,
 Poplar (liriodendron),
 Wild cherry, $\overline{a}\overline{a}$ 6 ounces
 Boneset (eupatorium),
 Cayenne pepper, $\overline{a}\overline{a}$ 4 ounces.

Mixed and sifted. Of the mixture, a teaspoonful in warm or cold water, three or four times a day. A useful tincture, to serve the purpose of a tonic, and as a remedy in malarial affections, may be prepared from the above combination of bitter tonics.

The following is a valuable restorative in the malarial cachexia :

862. R. Quiniae sulph.,
 Ferri sulph. exsicc.,
 Manganesii sulph. exsicc., āā Ḑj. M.
 Ft. pil. No. xx.
 S.—One pill three times a day.

When the malaria has produced jaundice, the author has seen excellent effects from the following :

863. R. Chinoidin, ḩj
 Manganesii sulph. exsicc., Ḑij. M.
 Ft. pil. No. xx. S.—One three times a day.

Recent experience has conclusively shown the curative power of iodine in malarial fevers. Thus three hundred consecutive cases were, with few exceptions, promptly arrested (ANDERSON). In some cases, no paroxysm occurred after the medicine was administered ; sometimes the attacks persisted for two or three days, but then the relief was complete. The official compound tincture of iodine may be prescribed—for adults, ten to fifteen minims, three times a day, well diluted and before meals. Carbolic acid, itself having considerable antiperiodic power, may be combined with the compound tincture of iodine :

864. R. Tinct. iodinii comp., ḩij
 Acid. carbolic, ḩj. M.
 S.—Ten to fifteen minims three or four times a day.

The officinal tincture of iodine is given by many practitioners. If quinine be necessary to break up the paroxysms, iodine is highly useful in the interim, as has been indicated.

ANTI-QUININE TREATMENT OF INTERMITTENT FEVER.

DR. BEARDSLEY.

At a recent meeting of the Medico-Chirurgical Society of Montreal, Dr. BEARDSLEY gave the results of fifty years' experience in treating intermittent fever without quinine. Briefly, it is as follows :

First, brisk alterative purgatives, aloes, blue-mass, and capsicum, in

equal parts, made into pill-form, or calomel in place of the blue-mass. An alterative purgative was to be taken at the outset, and repeated as the nature of the case demands. This was to be followed by an aromatic bitter, and perhaps an alkali with it, or in combination with boneset tea, drunk very freely, was an element in the cure not to be overlooked. He stated that this method had been very satisfactory in his and others' hands, and was often successful where quinine had utterly failed, and that, too, where it had been pushed; indeed, as far as twenty-grain doses three times a day, or even oftener.

DR. A. R. HALL, INDIA.

The use of *emetics*, in intermittent fever, should not be neglected. Often, the anti-periodic will not act until the stomach is thus emptied. Dr. HALL also states (*The Practitioner*, May, 1875,) that, in India, it is customary to give an emetic at the commencement of the cold stage, when shivering first begins. *Sulphate of zinc* (gr. xxx), followed by copious draughts of hot water—say three or four pints, if the patient can be induced to swallow them—will generally induce free vomiting, and cut short the paroxysm at once. *Ipecacuanha* may also be employed, but the depression it leaves behind it is much greater than the zinc.

DR. J. B. SCRIVEN, LAHORE, INDIA.

This English surgeon reports very favorably of the *nitric-acid bath*, in malarial disorders. (*Lancet*, November, 1876.) He has used it with the best effect, in a vast number of cases. For children, it is an extremely pleasant remedy, though it often produces a slight irritation of the skin. They generally sleep soundly after it, and rapidly improve in health. It is always necessary to continue the baths daily for at least a fortnight—often for a month or six weeks. With adults, of course, it is less manageable, but not less efficacious.

As pure nitric acid is very expensive, it may be well to add that he has found the impure acid, which is very cheap, and readily procurable, quite good enough for the acid bath. It contains a little hydrochloric acid and iron. A wooden tub should, of course, be used in preference to a metal one, or, in the absence of this, an earthen vessel.

PROF. M'LEAN, M. D., INDIA.

This distinguished authority speaks in the highest terms of *Warburg's tincture*. His words are :

"I have treated remittent fevers of every degree of severity, contracted in the jungles of the Deccan and Mysore, at the base of mountain ranges in India, on the Coromandel Coast, in the pestilential highlands of the northern division of the Madras Presidency, on the malarial rivers of China, and in men brought to Netley Hospital from the swamps of the Gold Coast ; and I affirm that I have never seen quinine, when given alone, act in the manner characteristic of this tincture. And, although I yield to no one in my high opinion of the inestimable value of quinine, I have never seen a single dose of it, given alone, to the extent of $9\frac{1}{2}$ grains, suffice to arrest an exacerbation of remittent fever, much less prevent its occurrence, while nothing is more common than to see the same quantity of the alkaloid in Warburg's tincture bring about similar results."

The following is the formula, as made known by Professor McLEAN:

865. R.	Aloës (socotr.,)		lb.j
	Rad. rhei (East India,)		
	Sem. angelicæ,		
a.	Confect. damocratis,	āā	℥iv
	Rad. helenii (s. enulæ,)		
	Croci sativi,		
	Sem. foenicul.,		
b.	Cret. præparat.,	āā	℥ij
	Rad. gentianæ,		
	Rad. zedoariæ,		
	Pip. cubeb.,		
	Myrrh. elect.,		
	Camphoræ,		
c.	Bolet. laricis,	āā	℥j.

The above ingredients are to be digested with 500 oz. proof spirits in water-bath for twelve hours ; then expressed and 10 ounces of disulphate of quinine added ; the mixture to be replaced into the water-bath till all the quinine be dissolved. The liquor, when cool, is to be filtered, and is then fit for use.

a. This confection, which consists of an immense variety of aromatic substances, was once officinal, and is to be found in the London Pharmacopœa, 1746.

b. Dr. WARBURG states that this ingredient was added to correct the otherwise extremely acrid taste of the tincture. Many other substances were tried, but none answered so well as prepared chalk.

c. This is the Polyporus laricis (P. officinalis, Boletus purgans, or larch-agaric), "formerly," says PEREIRA, "used as a drastic purgative, and still kept by the herbalist."

The tincture is of a deep brown color, has an aromatic and slightly terebinthinate odor, and an intensely bitter and warm aromatic taste: a noteworthy point about it is that there is nothing spirituous in either taste or smell, and it seems as if the alcohol employed in its manufacture were entirely saturated, and, so to speak, extinguished, by the substances taken up. The tincture evaporates readily, leaving a copious brown residue.

DR. J. MILNER FOTHERGILL, LONDON.

The great remedy in malarial attacks is *quinine*. It is best given in an acid solution. The following is an agreeable dose:

866. R.	Quiniæ sulphatis,	gr. x	
	Acidi sulphurici diluti,	℥x	
	Syrupi aurant.,	f. ʒ ij	
	Aquam,	ad f. ʒ j.	M.

This amount three times a day.

In some cases where the quinine seems powerless to break up the chill, the administration of an emetic each morning brings the disease at once under the control of the drug.

Some patients manifest a marked intolerance of quinine, especially those who have resided in the tropics. In such patients, arsenic, strychnine, berberin and the eucalyptus must be resorted to.

The following combination is highly extolled by a writer in the *American Medical Bi-weekly*, March, 1877, in order to prevent the return of the chills after they have been broken by quinine:

867. R.	Quiniæ sulphatis,	ʒ ij	
	Tinct. ferri chloridi,	f. ʒ ij	
	Liq. potassii arsenitis,	f. ʒ iss	
	Tinct. opii,	f. ʒ ss	
	Strychniæ sulphatis,	gr. ij	
	Alcoholis,	f. ʒ viij	
	Aquæ,	f. ʒ xxiv.	M.

One tablespoonful three times daily.

An experienced practitioner says of this prescription that when administered after the liver has been acted on by a mercurial, it will cure any case of enlarged spleen or chills, if persisted in.

PROF. HARVEY L. BYRD, M. D., BALTIMORE.

In the treatment of that much-dreaded and often rapidly-fatal condition, the "Congestive Chill," Prof. BYRD knows nothing of equal value to *chloroform* and *opium*, used as follows:

873. R. Chloroformi, q. s.
Inhale to anæsthesia, or to the restoration of the circulation in the extremities.

The doctor often prescribes inhalation of chloroform to a limited extent, to mitigate or curtail the *cold stage* in intermittent fevers.

Or:

874. R. Pulveris opii, gr. ij
Pulveris capsici, ʒj. M.
Make powders. One powder every hour, in severe cases, until the re-establishment of the circulation.

He is accustomed to give between the powders 10 to 15 drops of aromatic sulphuric acid, in a little water.

When reaction of the circulation is established in "congestive chill," the case should be treated in the usual manner.

DR. BENJAMIN LEE, PHILADELPHIA.

This practitioner recommends a full dose of quinine about an hour before the chill, and half an hour before it the following:

875. R. Chloral hydratis, ʒj
Aquæ camphoræ, ʒj. M.
For one dose.

876. R. Quiniæ sulphatis, āā
Ferri redacti, ʒj
Acidi arseniosi, gr. ij
Piperinæ, ʒ iss. M.
Make twenty pills. One three or four times a day. A favorite prescription, in obstinate chills, with some Southern physicians.

DR. BABIEFF.

This author regards (*Mediz. Obozr.*, Feb., 1882) systematic *faradization of the spleen* as one of the best adjuvant means in the treatment of intermittent fever, and adduces some cases of his own in which the usual antiperiodic remedies had remained unsuccessful until the electricity had been added. The faradization not only diminished the splenic tumor, but also acted beneficially on the malarial process. This

favorable influence the author attempts to explain as follows: 1. Faradization gives rise to contractions of the splenic vessels, and, in a reflex way, acts tonically on the vaso-motor centres. 2. It counteracts blood-stagnation, and possibly accumulation of miasmatic products in the spleen. 3. It acts indirectly on the neighboring kidney, increasing correspondingly the secretion of urine. 4. It prevents the formation of infarcts and ruptures in the spleen. The editor of the *Mediz. Obozr.*, Dr. V. F. SPRIMON, also testifies to the high value of faradization and galvanization in cases of chronic malarial tumor of the spleen. He saw its prompt disappearance, as well as a great improvement of the general state, in four out of five of his patients treated by this method. In four of the eight of Dr. SPRIMON'S cases which presented intermittent of recent standing, and were treated by faradization alone, the fever disappeared and never returned after five to ten sittings (half an hour daily).

DR. JAMES H. HUTCHINSON, OF PHILADELPHIA,

considers that while quinia arrests the chills, it is not sufficient in itself to improve the character of the blood. He therefore gives iron in the form of the tincture of the chloride. He knows of no better preparation. It may be given in much larger doses than are set down in the books. In anæmia and erysipelas, he frequently gives thirty drops every three hours, without producing any bad effects. The only precautions which it is necessary to observe are to dilute it sufficiently, and to direct that it shall be taken through a tube, in order to prevent its acting upon the teeth. This medicine has another advantage over many of the other forms of iron, in that it is cheap.

THE HYPODERMIC USE OF QUININE IN INTERMITTENT.

The subcutaneous injection of quinine in malarious fevers has been frequently tested at home and abroad, the efficacy of the drug in controlling fever, and the saving effected in consequence of the smaller quantity required, being sufficiently important to fully justify the share of attention which the subject has received. When rapid action is called for, or when, from irritability of the stomach, idiosyncratic or symptomatic, quinine is not well borne; or, again, when its exhibition *per viam naturalem* is resisted, the hypodermic injection of the drug may be advisable.

By this plan, there is a great saving in the quantity used, the cases being usually conducted to a satisfactory conclusion, so far as fever is concerned, with a very moderate expenditure of quinine, say 3 grains or so. On the double ground of checking the fever and economy in the consumption of the drug, this mode of treatment would seem to be superior to the old method; but, unfortunately, a drawback presents itself in inflammation at the point of insertion of the syringe. This may be avoided, to some extent, by a careful solution and administration.

Surgeon Major GEORGE Y. HUNTER, (*Lancet*, March, 1876,) of Bombay, uses:

877. R.	Quiniæ sulphatis,	gr. lxxx	
	Acidi citrici,	gr. ccc	
	Aquæ destillatæ,	f. ʒj.	M.

Dissolve with the aid of heat. ℥vj contain one grain of quinine.

The formula of Dr. LEORUST, of Bremen (which is recommended by Dr. AUSTIN FLINT), is:

878. R.	Quiniæ sulphatis,	gr. xxxiij	
	Acidi sulphurici,	gtt. xj	
	Aquæ destillatæ,	f. ʒj.	M.

Mix the acid and water, and gradually add the quinine, stirring with a glass rod; if the solution is not clear, it may require one, possibly two drops more of the acid. Filter through paper, and it is ready for use. Ten drops of this mixture are equivalent to fifteen *minims*, which contain one grain of quinine. One grain, hypodermically, is equivalent to three or four by the mouth.

As regards the best place for insertion, it is stated that the region of the collar-bones is the most preferable. Any spot will do that is at all suitable for hypodermic injections; but when it is inserted where the muscles are in constant use just beneath the skin and fascia, it will be apt to occasion soreness and swelling. Abscesses are rare when the solution is kept free from particles and sediment. If care is not taken, the instrument used may be spoiled by the acid acting on the piston and other parts; to avoid this, it should be washed after using, first in clear water, then in a solution of soda (a teaspoonful in a glass of water), then in clear water again. Ether, perhaps, will answer for the same purpose, but is not always convenient.

There is considerable pain on forcing out the solution, for the acid is irritating to the tissues, and there are frequently hard, indurated spots,

where the fluid is forced out, lasting a few weeks, or even months, but they always gradually disappear.

Dr. J. M. FOTHERGILL recommends as not liable to cause ulceration:

879.	R.	Quiniæ sulphatis,	gr. x	
		Acidi sulphurici diluti,	℥x	
		Aquæ destillatæ,	gtt. c.	M.

The hypodermic method does not produce ringing in the ears, deafness, or other unpleasant effects of quinine taken by the mouth. It ought, as a rule, to be used only in those instances in which the ordinary method is inapplicable, as the following: 1st. Those in which quinine by the mouth has failed to stop the fever. 2d. Those in which there is great irritability of stomach, or great disgust for the remedy. 3d. Those in which the fever, though controlled by quinine in the ordinary method, yet requires large and repeated doses. 4th. Those in which the effect on the head or nervous irritability produced by the drug is distressing. 5th. Cases in which death seems to be imminent, and in which it is necessary to produce a decided and rapid effect upon the fever. The following formula, containing gr. iv to f. ʒj, is an appropriate one:

880.	R.	Quiniæ sulphatis,	gr. xxxij	
		Alcoholis diluti,	f. ʒ vijss	
		Aquæ ammoniæ,	f. ʒ ss.	M.

Dr. LENTE's formula is:

881.	R.	Quiniæ sulphatis,	gr. l	
		Acidi sulph. diluti,	℥c	
		Acidi carbolic liq.,	℥v	
		Aquæ,	f. ʒj.	M.

Dissolve the quinine by the aid of heat, and, after filtration, add the carbolic acid.

All acid solutions, however, cause considerable pain, and it is probably better to dissolve the pure alkaloid quinia in ether.

Prof. W. BERNATZIK recommends:

882.	R.	Quiniæ,	gr. vij	
		Etheris sulphurici,	℥xv.	M.

The alkaloid can be obtained by precipitation from a solution of the sulphate by ammonia.

THE TREATMENT OF ENLARGED SPLEEN OR AGUE CAKE.

According to some authorities, *mercury*, in any form, is contra-indicated when malarial enlargement of the spleen exists.

The *hyposulphite of soda* is recommended by Dr. THOMAS HILL, of Missouri. (*Louisville Medical and Surgical Journal*, 1872.)

883. R.	Quiniæ sulphatis,	gr. ix	
	Sodii hyposulphitis,	gr. xvj	
	Acidi sulph. aromat.,	gtt. vj	
	Aquæ,	f. 3j.	M.

A teaspoonful every two hours to a child. Ten grains of the soda, thrice daily, may be given to an adult.

In the enlarged spleen, often associated with leucocythemia, which so frequently follows malarial disease, the *iodide of ammonium*, gr. v three times a day, with inunctions of the red iodide of mercury, has been found efficient. Dr. DA COSTA prefers, however, the treatment by the hypodermic administration of *ergotine*, gr. iij-v daily, which has certainly produced excellent results. Dr. J. B. C. GAZZO, of Louisiana (*Medical and Surgical Reporter*, March, 1877,) reports favorably of throwing the injection into the tissues of the spleen itself. He uses:

884. R.	Fld. extr. ergotæ ether.,	f. 3ij	
	Glycerinæ,	f. 3j.	M.

This solution is injected into the spleen, two syringefuls at once, and repeated after twenty-four hours. The more deeply the needle is driven the less is the sensibility. Constipation must be avoided.

Professor MOSLER, of Switzerland, has also employed parenchymatous injections to reduce the spleen. He employed *carbolic acid* in a two per cent. solution, and *liquor potassii arsenitis*, one part to ten of distilled water. The former caused so much pain that he did not continue it; the latter, however, was repeated a number of times, and brought about a very decided diminution of the organ. (EULENBURG, *Die Hypodermatische Injection*, Berlin, 1875.)

According to Dr. R. WILLIAMS the *bromide of potassium* has remarkable powers in the reduction of splenic hypertrophy. It is given internally, in full doses.

Dr. R. BARTHOLOW believes that small doses of the *iodide of ammonium*, combined with arsenic, will reduce an enlarged spleen from malarial causes; as:

885. R.	Ammonii iodidi,	ʒj	
	Liq. potassii arsenitis,	f. ʒss	
	Tincturæ calumbæ,	f. ʒss	
	Aquæ,	f. ʒiiss.	M.

A teaspoonful three times a day before meals. Paint the side with tinct. iodin. compos.

The *iodide of lead*, gr. $\frac{1}{2}$ thrice daily, has testimony in its favor.

Very excellent results have been reported by Dr. L. McGUIRE, of California, from the use of *strychnia* or *nux vomica* (*Pacific Medical Journal*, Feb., 1873,) in ague cake. He gives $\frac{1}{16}$ of a grain of the former, or ℞xx, of the tinct. nucis vomicæ, three times daily, combined, if need be, with iron or quinine.

In splenic and hepatic enlargement, Dr. EVERS (*London Medical Record*, June, 1875,) reports the best results from the milky juice of the unripe fruit of the paw-paw, *carica papaya*. About a teaspoonful of the juice is mixed with sufficient sugar and divided into three boluses, one to be taken thrice daily. If it causes symptoms of gastric irritation, a little opium may be added. From twenty to twenty-five days is sufficient for a cure in nearly all cases of ague cake.

Of *external remedies*, blistering, painting with the compound tincture of iodine, inunction of ointment of biniodide of mercury, local abstraction of blood, etc., have each been lauded, but it is doubtful whether any of them is particularly useful. In Germany, the "dry-packing" method is employed at hydropathic establishments, and is said to reduce the spleen at a rapid rate without any other medication. It is certainly a simple and excellent derivative. B. SKOREZEWSKY (*Wiener Med. Wochensch.*, 1876, No. 21,) believing the enlargement of the spleen in ague to be due in chief to a depressed action of the vaso-motor nerves, employed *faradization* to diminish the size of the gland. He placed the electrodes, one over the anterior, the other over the posterior border of the spleen, and gradually increased the intensity of the current. Each sitting lasted fifteen to twenty minutes. The results he arrived at are as follows: The induction current produced a decided effect in reducing the size of the tumor; the first application was generally followed by the most marked alteration; the softer the spleen, the more striking was the effect of the electricity. The spleen became hard on palpation, and the pain and uneasiness in the side were alleviated. The reduction of the gland, which was always best marked in the first few days of this treatment, was in rare cases noticed as early

as twenty-four hours after its application. No relapses occurred in any recent case of ague with a swollen spleen in which faradization had been carried out. The benefits of the electrical treatment were essentially aided by the administration of quinine.

In India, *puncture* of the spleen with long, sharp stilets of considerable thickness, is a common and successful practice with native physicians. It is not followed by bad results, and has been commended by some English physicians.

Dr. W. T. GADBURY, of Mississippi, speaks in the highest terms of a solution of *oxy-sulphate of iron*. He says that in an extensive practice for many years he has never had a case of malarial splenic hypertrophy which resisted it. (*Medical and Surgical Reporter*, April, 1866.)

886. R. Ferri sulphatis,	℥j	
Acidi nitrici,	f. ℥j.	M.
Stir until it ceases to effervesce, then add		
Aquæ,	f. ℥iv.	

This is used as a part of the following combination :

887. R. Liquoris oxy-sulphatis ferri, (above),	f. ℥ijj	
Aquæ cinnamomi,	f. ℥viiij	
Quiniæ sulphatis,	℥ss	
Potassii nitratis,	℥ijss.	M.

One tablespoonful three or four times a day.

In obstinate cases, an iodine ointment or some other counter-irritant should be rubbed in over the enlarged viscus, and a roller worn round the body, so as to make compression on all the organs supplied by the coeliac axis. In cases of children suffering from lumbrici, it is well to commence with anthelmintics.

In the *Atlanta Medical and Surgical Journal*, 1876, Dr. D. L. PHARES, of Mississippi, highly recommends the root of the *Ceanothus Americanus* as a remedy for enlarged spleen. He exhibits it in form of a tincture, made by filling a bottle loosely with the root and root bark, pouring whiskey upon it, and allowing maceration to continue for a week. The dose is f.℥ss-j *ter die*. The same tincture is rubbed over the spleen twice daily.

NOTES ON REMEDIES.

Aconitum. Dr. S. LOVING (*Ohio Med. Jour.*, Aug., 1878,) says that he knows of nothing which promotes the comfort of patients in the hot stage more than a combination of aconite and morphia, in small doses.

Alstonia. Several species of this Australian tree are used for their antiperiodic effects. Their value is moderate or doubtful.

Ammonii Carbazotas. It is stated by Dr. DUJARDIN-BEAUMETZ that this agent acts very efficaciously in intermittent fever. The daily dose is gr. ss-j, in pill form. Dr. SLANE, of Goalpara, India, finds carbazotate of ammonia to be a very efficient remedy. It does not increase the appetite, but like quinine, after a few doses, it produces tinnitus aurium and slight deafness. The urine becomes tinged of a deep orange color, and occasionally the skin and conjunctiva acquire a jaundiced hue, but the discoloration soon disappears. It may be given in pill, with extract of gentian, and combination with a quarter of a grain of extract of nuxvomica seems to increase its efficacy. Being perfectly soluble in water, it may be given in solution, but it has an intensely bitter taste. The largest quantity Dr. SLANE ever gave was eight grains in twelve hours. (*Indian Medical Gazette*, April, 1876.) It has also been favorably reported upon in this country, especially on the Pacific coast.

Ammonii Chloridum, in doses of ʒij daily, as an average, was once very popular in ague, and is still employed by some.

Amyl Nitrite. In intermittent fever, amyl has aborted the chill, but only shortens the latter stages: it may be given even after the algid stage has fairly set in. Some bold exhibitions have been tried, even to the extent of thirty drops, with good rather than bad effects, in this form of fever. The ordinary dose has been six drops.

Arsenicum undoubtedly stands second only to quinine in the treatment of malarial poisoning; its especial field is the chronic forms. It may usefully be combined with iron:

888.	R.	Pill. ferri carbonatis,	ʒij	
		Acidi arseniosi,	gr. j.	M.
For twenty pills. One three times a day.				

Or:

889.	R.	Quinise sulphatis,	ʒij	
		Ferri sulph. exsic.,	ʒj	
		Acidi arseniosi,	gr. j.	M.
For twenty pills. One three times a day.				

When used alone, ten drops of Fowler's solution may be given three times a day after meals. The diet should be abundant and nourishing.

Atropia. Some writers state that, in the cold stage of intermittent or congestive fevers, atropia injected under the skin, will bring about reaction with more certainty than any other means commonly used in such cases. It will abort the chill, if applied at the onset, in most cases.

Berberin. This alkaloid, from the *Hydrastis Canadensis*, *Berberis vulgaris*, etc., is highly esteemed in chronic malarial poisoning.

Buxinæ Sulphas. The alkaloid of the *Buxus Sempervirens*, in doses of gr. xv, has achieved some reputation in Italy.

Camphora. A tablespoonful of the tincture of camphor, about the time the chill is expected, will occasionally prevent it.

Capsicum. Prof. WM. H. THOMPSON says that either capsicum, ginger, or other aromatics, combined with quinia, will diminish the amount required of the latter. "A good dose of capsicum, combined with 20 grains of quinine, will act as well as 30 grains of quinine without capsicum."

Carbolicum Acidum. This has rather disappointed expectations. It may be given in doses of gr. iv, in infusion of gentian, and no doubt occasionally acts well. M. DÉCLAT, of Paris, has used it hypodermically, with success, in obstinate cases. (*Comptes Rendus*, No. lxxv.)

Chloroformum, in doses of f. ʒss-j, in a small glass of milk, every two hours, about the time of the expected chill, will frequently abort it.

Chinoidine. This is the *cheapest* of the cinchona alkaloids. From gr. xv-xx, daily, for five or six days, and then gradually reduced, will cure mild intermittents very satisfactorily.

Chloral Hydratis has been highly recommended by Dr. P. BAILEY, of South Carolina, (*Charleston Medical Journal and Review*, October, 1873,) and others. When the fever heat is intense, its refrigerant and calmative influence is undoubted; and while it quiets the nervous system, it rather promotes the secretions—in this particular, being far superior to the opiates. In the malarial fevers of children, with a tendency to convulsion, or when this unfortunate complication has supervened, it can scarcely be excelled. Although the first impression on the gastric surface is rather to excite emesis, nothing can be more demonstrable than its promptness in arresting the vomiting, in these fevers, when all other efforts have been fruitless.

890. R.	Chloral hydratis,	ʒiss	
	Potassii bicarbonatis,	ʒij	
	Spts. æth. nitrosi,	f. ʒss	
	Syrupi tolutani,	f. ʒijss	
	Aquæ,	f. ʒj.	M.

A tablespoonful every hour or two.

Cimicifuga Racemosa. Dr. EDWARD R. MAYER, of Wilkesbarre, Pa., states that the tincture of cimicifuga, alone or combined with tincture of *kalenia*, is one of the most efficient remedies known for the malarious neuralgia of the first branch of the fifth pair, commonly called *brow ague*. (*Hints in Specific Medication*, 1876, p. 17.)

Cinchonia. A cheap and useful alkaloid, about half as strong as quinine. It has the great advantage that it can be rendered quite tasteless without impairing its efficiency. Dr. SAMUEL ASHHURST, of Philadelphia, combines it thus:

891. R.	Cinchoniæ,	gr. j	
	Sacchari lactis,	gr. iv	
	Sodii bicarbonatis,	gr. ʒ.	M.

the chill, is a useful domestic remedy. A decoction of corn shucks or fodder is popular in the South.

Ferri Persulphatis Liquor. This has been successfully used in obstinate intermittents, in doses of gtt. viij–xvj, every four or six hours. Almost all the preparations of iron are valuable as tonics in prolonged malarial cases. The taste of the patient and the convenience of their administration, should be the guides in selecting from them.

Furfurine Nitrate. The nitrate of furfurine, obtained from bran, sawdust, and similar substances, has been recently brought forward as an energetic antiperiodic.

Gelsemium. Several writers have praised, very highly, the jasmine as an antiperiodic. The dose recommended by Professor WILLIAM W. MURRAY, of Baltimore (*Medical and Surgical Reporter*, January, 1873), is mv of the tincture every hour, until it produces heaviness of the eyelids, dilated pupils, or double vision, the same course to be pursued for four or five days successively. He adds that it is superior to quinia, in preventing the recurrence of the attacks.

Grindelia Squarrosa. The fluid extract of this plant has been used successfully in hypertrophied spleen. The dose is $\text{f}\text{.}\text{ʒj}$ four times a day.

Hydrastis. In obstinate recurring intermittents, Dr. EDWARD R. MAYER has found the following recipe more effective than quinia :

895. R. Hydrastiae sulphatis,
Ammoniae carbazotatis, $\bar{\text{a}}\bar{\text{a}}$ gr. $\frac{1}{4}$.

According to Dr. BARTHOLOW, "hydrastia ranks next to quinia as a remedy for intermittents."

Hydrocyanicum Acidum. "Peach leaf tea" is a popular remedy in intermittents in Maryland and Delaware. On this hint, Dr. JAMES H. PEABODY, of Omaha, has used with great success hydrocyanic acid. In chronic intermittents it has served him when cinchona and its alkaloids have entirely failed. A favorite prescription is :

896. R. Acidi hydrocyanici dil., gtt. xlvij
Tr. cinchonae comp.,
Tr. gentianae comp.,
Syrupi pruni virginianae, $\bar{\text{a}}\bar{\text{a}}$ ʒij . M.
Two teaspoonfuls four times daily.

Iodinium. Recommended by Dr. DE WILLEBRAND as curative, (F. 870,) but its especial and great value is after the ague has been checked by quinine, to prevent its recurrence. For this purpose, tinct. iodinii, gtt. v–x, three times a day, in a spoonful of water, before meals, is said by Dr. JAMES Y. SHEARER, of Pennsylvania, (*Medical and Surgical Reporter*, January, 1874,) to surpass all other agents yet suggested. Nothing disguises the taste of iodine so well as sweet milk ʒj to gtt. x of the tincture.

Ipecacuanha. In both intermittents and remittents, accompanied by congestion

of the portal circulation, ipecacuanha is often very serviceable. It is probably the most suitable emetic which can be administered at the outset of a malarial attack, when such an agent is called for, and where the consequent depression is not to be feared.

Leptandra Virginica. This is said to be an invaluable remedy to prevent the return of intermittents after they have been broken up by quinia. It is a favorite ingredient of secret remedies for this purpose, and should be combined with the cinchona alkaloid, and continued some weeks. (Dr. EDWARD R. MAYER, *Hints in Specific Medication*, 1876.)

Narcotine, in doses of gr. j-ijj, possesses antiperiodic powers, and by some practitioners has been asserted to equal quinine.

Nitricum Acidum. In old and obstinate cases, says Dr. LYMAN, (*Chicago Medical Journal and Examiner*, December 7, 1861,) we want to administer something powerfully oxidizing in character, to wash or cleanse the body of the remains of the disease. We should administer first a diuretic. The acetate or the citrate of potassa is useful if quinine alone has failed. But if they are not, a course of nitric acid may relieve for years.

Opium, in full dose, three hours before the expected chill, will frequently stop its appearance. *Morphia* may nearly always be advantageously combined with quinine. Dr. MERIWETHER LEWIS, of Tennessee, lays down these aphorisms in the *American Journal of Medical Science*, July, 1878: 1st. The combination of an opiate with quinia insures an earlier and more complete cure of intermittent and remittent fevers, than quinia alone; 2d. Less quinia is required, and hence there is less cerebral disturbance; and, 3d. Any paroxysm of intermittent or remittent fever may be aborted by the use, at the proper time, of a *decided dose of morphia and quinia*, with reasonable certainty.

Piper. In doses of gr. v-xx, pepper has been found of considerable value, especially for combining with quinia when the stomach is not susceptible to the latter's action.

Potassii Nitras. This has been highly praised. It is administered in doses of gr. x in brandy and water, or dry on the tongue. Dr. SAWYER, of St. Louis, states he has never failed to arrest with it an uncomplicated paroxysm of ague.

Quineti Sulphas, a salt that represents all the alkaloids, is recommended in doses of gr. x-xv, by Dr. HOLLIS. (*Brit. Med. Jour.*, May, 1879.)

Quiniae Bromohydras. Professor GUBLER, of Paris, (*Journal de Therapeutique*.) points out that the proportion of quinine in the neutral bromhydrate is more considerable than in its analogue, the neutral sulphate of quinine. It is also six times as soluble in water. Thus its greater solubility and richness in alcohol give it superior advantages for the purposes of hypodermic injection. The new compound also is alleged to offer the valuable combination of the properties, in part synergic, of quinine and of the bromated preparations. M. GUBLER orders about $1\frac{1}{2}$ grains, in pill, not exceeding six grains daily, as an ordinary dose, and rarely,

if ever, giving as much as 15 grains a day. Its effects are those of quinine, producing ultimately headache, ringing in the ears and deafness, or those of slight bromism, muscular languor, accompanied by a strong tendency to sleep. M. GUBLER has sometimes employed with success hypodermic injections of bromhydrate of quinine in a solution of the strength of $\frac{1}{10}$, in water slightly alcoholized. Notwithstanding the presence of a small proportion of alcohol, the bromhydrate of quinine, in subcutaneous injection, has always shown itself inoffensive to the tissues, which is contrary to that which occurs with sulphate of quinine. Dr. W. E. FORREST, of New York, has combined hydrobromic acid with sulphate of quinine, and finds the distressing symptoms of quininism are thus wholly avoided.

897. R. Quiniæ sulphatis, ℥j
 Acidi hydrobromici,
 Aquæ, āā f. ℥iss. M.

Two teaspoonfuls contain 5 grains of quinine.

The hydrobromic acid used must not be that usually sold, (Merck's,) but that prepared according to Dr. J. MILNER FOTHERGILL's formula :

898. R. Potassii bromidi, ℥xj
 Aquæ, Oiv.

Dissolve, and add :

Acidi tartaraci, ℥xiiij.

The hydrobromic acid remains in solution, and potassa bitartrate is precipitated.

M. LATOUR's formula is :

899. R. Quiniæ sulphatis, gr. x
 Acid. sulph. diluti, ℥xx
 Aquæ, f. ℥j.

To this, add

Potassii bromidi, gr. xx.

This forms a neutral bromohydrate.

Quiniæ Salicylas. The salicylate and carbolate of quinia have been recommended on theoretical grounds.

Quiniæ Sulphas. This is the standard remedy, the typical specific, in all phases of malarial diseases. In regard to the *form* of its exhibition, the most active is in solution. Its bitter taste is best concealed by liquorice, glycyrrhizin, chocolate, tannic or malic acid. Sugar-coated pills and chocolate caramels, containing quinia, may readily be obtained. Frequently, a clear solution, free from turbidity, is a *desideratum*. With aromatic sulphuric acid, we get a passable solution, but the acid is often objectionable, if not absolutely contra-indicated. Dr. ISAAC SMELT, Jr., (*N. Y. Medical Journal*, August, 1876,) finds that, in practice, the

sweet spirits of nitre is all that is desired. One ounce of it will dissolve about two drachms of quinine, giving a transparent solution. The following is said to disguise the taste well :

900. R. Quiniæ sulphatis, ℥j
 Cocoæ, ℥j
 Syr. aurant. cort., f. ʒxx.
 Triturate well the cocoa, after which add the quinine and orange peel. Mix thoroughly.

Sweetened hot tea, drawn strong, will almost entirely conceal the bitterness so much complained of. The tannic acid is the efficient agent, which may be utilized as follows :

901. R. Quiniæ sulphatis, gr. xxiv
 Syr. sarsap. comp., f. ʒiij
 Acidi tannici, gr. iij
 Olei menth. piper., gtt. v. M.
 Milk, in the proportion of f. ʒj to gr. j of quinine, will conceal its taste tolerably.

In giving quinine to children, it is frequently better done by enema :

902. R. Quiniæ sulphatis, gr. ij-x
 Acidi tannici, gr. ij-vj
 Aquæ, f. ʒj. M.
 For one injection. The tannin aids the retention.

Or the *endermic* method may be very successfully employed :

903. R. Quiniæ sulphatis, ʒss-j
 Alcoholis, f. ʒss
 Adipis, ʒviij.
 Rub well into the skin. (Dr. AUGUSTE NONAT.)

Often, the action of quinine is greatly aided by stimulant and aromatic adjuvants, as alcohol, ether, capsicum, pepper, ginger, etc. Dr. FULLER has found that when there is splenic enlargement, the addition of colocynth and rhubarb greatly increases the energy of the alkaloid.

Dr. ROBERTS BARTHOLOW (*Prize Essay on the Use of Quinine*,) lays down the rules : 1. Quinine is equally effective, whether administered in the interval or during the seizure. 2. Its maximum effect is five hours after taking ; hence, it should be administered about this long before the expected paroxysm. 3. It is better to give the whole amount in a single dose, than in a succession of small doses. In ordinary intermittents, gr. x in the sweating stage, and the same quantity five hours before the time of the next paroxysm, is the most effective plan.

An important practical point in the administration of *quinine* has been noted by Dr. SCHILLING, (*Glasgow Med. Jour.*, 1883). This is, that the ringing in the ears and deafness produced by large doses may be counteracted by combining *ergot* with the cinchona salt, in equal

quantities or about two to one. This association is said also to prevent the amblyopia which sometimes supervenes after large doses of quinine.

Some useful suggestions are made by Dr. AUGUSTE NONAT. (*Practitioner*, September, 1871.) The older the case, the larger the dose required; whenever the spleen is enlarged a double dose is required; quartan fever requires a much larger dose than quotidian or tertian. The total time required to eradicate the disease will generally be three weeks; the first week, the maximum dose must be given daily; the second week, one-third less, and the third week, two-thirds less; even after this, small doses, from time to time, are needed.

Salicin was extensively used during the Civil War, by the Confederate surgeons, and has recently been revived; gr. xxx, three or four times a day, is an average dose. On account of its alterative effect on mucous surfaces, it is deemed particularly applicable in malarial hæmaturia.

Sierra Salvia, four drop doses of the fluid extract every two hours, combined with tinct. veratrum, or tinct. nux vomica, has been successfully used in remittent fever by Dr. FRANCIS A. EVANS.

Resorcin, in doses of gr. xxx-xl, or larger, has been found by Dr. BASSI, of Venice, (*Compend. of Med. Sci.*, 1884) to check intermittent cases with almost the certainty of quinine. No harm resulted from these large doses, but sometimes there was temporary deafness and ringing in the ears. The rapid absorption of resorcin, which was noticed in all cases, makes it necessary to give the remedy half an hour to one hour before the expected access; if given at a longer interval it seems to exhaust its special action, and fails to influence the access. Resorcin seems to have the same value as quinine in this disease and in the same cases; where one fails, the other fails. It has the advantage of being comparatively much cheaper.

Sodii Bisulphis. This has been employed with good effect. Dose, ℥j in water, every two hours. It is only available in mild cases.

Sodii Chloridum. Dr. BROOKES recommends the following (*Zeitschrift für Praktische Medicin*, 1876, No. 33): Heat table salt in a pan until it turns brown; dissolve a tablespoonful in a glass of hot water; take the whole at one draught, on an empty stomach, the day after the chill, or, in the quotidian type, two hours after the fever. Avoid drinking water, and diet low. One such dose is usually enough. In doses of ʒ viij-x, in the apyrexia, it was recommended as far back as 1854, by J. C. HUTCHINSON, of New York.

Strychnæ Sulphas. In India, cases of intermittent have been successively treated with this agent alone. It is also freely used in this country, in numerous anti-periodic combinations. (F. 867, 872.)

Tela Araneæ. In the *Cincinnati Lancet and Observer*, 1877, Dr. L. M. JONES reports that having failed in treating a case of chronic intermittent, he adopted the suggestion of Dr. JACKSON, in the U. S. Dispensatory, to use spider-web. A bunch of spider-web, collected from a dark cellar,

(as it is the web of a species of spider which inhabits dark places that possesses medicinal properties,) about the size of a large hulled walnut, was put into 4 ounces of whiskey and allowed to macerate forty-eight hours, when it was filtered. The patient took a teaspoonful four hours before the expected chill, and at hourly intervals until she had taken four doses, and then took a like quantity before each meal and at bedtime, until all was taken. Although the anticipated chill came, it was very light and was the last one experienced up to the time of the report, a period of four months.

Urea, Artificial. The *Jour. d'Hygiene*, 1883, reports that Dr. BELVOUSOFF, of Charkow, Russia, has used artificial urea (carbamide) as a remedy for intermittent fever in place of quinine. It is almost tasteless, and does not depress the nervous system. In Southern Russia, the peasants have used urine as a febrifuge for centuries; this has suggested the rational use of urea to Dr. B.

MILK SICKNESS.

DR. J. C. PATTON, INDIANA.

This practitioner describes (*Indiana Journal of Medicine*, August, 1875,) a treatment which, he says, "almost deserves the character of a specific:"

904. R. Infus. sennæ foliarum, Oj
 Magnesiæ sulphatis, $\frac{3}{4}$ xij. M.

Of this give from one to two ounces, which will be vomited; then immediately give a second dose, and so continue until it is retained.

Large injections of cold whiskey and water should be thrown well up the rectum. For milder cases, sulphur, in free doses, until the bowels are moved, followed by diffusible stimulants, is sufficient.

DR. FULTON, OF OHIO.

This writer states (*Medical and Surgical Reporter*, April 12, 1884), that the treatment consists in sedatives, relaxants, cathartics, enemata, etc., judiciously and perseveringly applied. "Each practitioner will have his preferences in making his selection of articles from these different classes of medicines. Some may be more fortunate than others in their choice. I have my preferences. And I may say that in a thirty years' practice among it, I have been so fortunate as not to have lost a single case of my own attending.

"I said above, that the means must be *perseveringly* applied. If one dose is rejected, another must be given, and that 'not until seven times only, but until seventy times seven.' "

DR. W. H. PHILLIPS, OHIO.

This writer, in an essay on this disease, published in 1877, throws doubt on the value of alcohol in its treatment. He says the practical difficulty is that a large proportion of patients cannot tolerate it. The plain indications of treatment are to care for and sustain, by every proper means, the strength of the patient while he is passing through the disease. Dr. P. is not in favor of any plan of treatment that will create active perturbations of the system, as they inwardly do mischief; the means which will allay the nausea, and procure quietude and rest, are always the most satisfactory. Patients treated upon a severely expectant method, recovered in quite as short a time, and with less discomfort than those treated by the more heroic methods.

According to his experience, *sulphate of strychnia* accomplishes more good than any other remedy. By its use he has been able, in some measure, to overcome the atonic condition of the muscles, especially the muscular coats of the stomach and bowels. He also thinks that by virtue of its action upon the vaso-motor nerve centres, it is capable of improving the circulation, and in that manner invigorating the functionary power of the system.

He is in the habit of using from $\frac{1}{4}$ to $\frac{1}{8}$ of a grain of strychnine, every four to six hours as indicated, with the results above stated. As a palliative, some agreeable effervescent mixture is usually well borne, as the citrate of magnesia, or the granulated nitrate of cerium, with a little fruit syrup and fresh water. This serves to allay the thirst and nausea. He has also used the bitartrate of potassa, with the view of increasing the secretion of urine, and thus saving the patient from danger arising from retention of effete matter, with very satisfactory results in some cases.

PURPURA HÆMORRHAGICA.

DR. THOMAS HAWKES TANNER.

This writer says the treatment should consist in the exhibition of full doses of sulphate of soda with sulphuric acid, or of castor oil, until the bowels are thoroughly cleared. Then quinine, iron, the mineral acids, and arsenic, as in the following combination :

905. R.	Quiniæ sulphatis,	gr. ix	
	Acidi phosphorici diluti,		
	Tincturæ ferri chloridi,	āā	f. ʒ iss
	Liquoris arsenici hydrochlorici,		℥ xv-xl
	Syrupi zingiberis,		f. ʒ vj
	Infusum quassiae,	ad	f. ʒ viij. M.

One-sixth part directly after breakfast, dinner and supper.

The patient should be given nourishing diet, fresh fruit or vegetables, and a fair allowance of malt liquor or wine. The oil of turpentine, or gallic and aromatic sulphuric acids, where there is internal hemorrhage, are often quite effective.

DR. ROBERTS BARTHOLOW, OF PHILADELPHIA,

says that in purpura and the hæmorrhagic diathesis, digitalis is useful when given conjointly with restorative medicines ; but, as a dyscrasia exists on which the extravasations of blood depend, it is obviously necessary to correct this state of things, in order that the patient shall be benefited by a remedy which gives tone to the heart and vascular system. Gallic acid is useful and the following mixture is highly recommended :

906. R.	Acidi gallici,	ʒ ss	
	Acid. sulphur. dil.,	f. ʒ j	
	Tinct. opii deod.,	f. ʒ j	
	Inf. rosæ comp.,	f. ʒ iv.	M.

S.—A tablespoonful every four hours or oftener.

PROF. N. S. DAVIS, CHICAGO.

This experienced observer says that he places more reliance on *digitalis* and *ergot*, internally, aided by the solution of the persulphate of iron applied to the bleeding surfaces, than on any other remedies. During a period of actual bleeding, the remedies should be given in pretty full and frequent doses, aided by entire rest and mild diet. After the

bleeding ceases, he continues the same remedies, in smaller doses, three times a day, from three to six months, interrupting them occasionally for three or four days at a time. Careful attention should also be given to keeping the digestive and excretory organs in good order, and there should not be too much haste in promoting the reformation of blood by the use of iron and rich food.

NOTES ON REMEDIES.

Ergot should be combined with tonics, and will be found of value.

Ferrum is always useful. The following prescription is a good one :

907. R.	Ferri sulphatis exsiccatae,	℥ij
	Quiniæ sulphatis,	℥j
	Strychniæ sulphatis,	gr. ss.
Make twenty pills; one thrice daily.		

Sulphuricum Acidum sometimes acts happily in purpura.

Terebinthinæ Oleum is an important agent in the treatment of purpura. It should be given in emulsion.

RELAPSING FEVER.

CITY HOSPITAL, RIGA.

In the decade 1865-75, two thousand cases of relapsing fever were treated in this hospital. The therapeutics adopted are described in the *Deutsches Archiv für Klinische Medicin*, Bd. XIX, (March, 1877,) by Dr. X. GIRGENSOHN. After frequent attempts to abort or shorten the disease by quinine, aqua chlorinii, acids, etc., all of which failed, a pure symptomatic plan was adopted. During the febrile stadia, acids, in mucilaginous decoctions, were exhibited, and in the intermissions, some preparation of iron. Wine and brandy, in moderate quantities, were given throughout, and also a simple but nourishing diet, consisting of milk, bouillon, eggs, flesh, etc. For the vomiting, ice pills were found most efficacious, as also subnitrate of bismuth, in doses of gr. iij-v. For the attacks of diarrhoea—which, however, often proved a favorable indication—opium was at first used, but later, usually, tannin, gr. iij-v. For the profuse hemorrhage—sometimes of decided gravity—ice water injections, ice to the breast and back of the neck, and large hypodermic injections of ergot, were occasionally

called for. The muscular pains which the patients complained of, especially during convalescence, were treated with the warm bath, or rubbing with a mixture of oil and chloroform. The severe pain occasionally felt in the splenic region, could usually be allayed with a bladder of ice to the part. Morphia injections were rarely employed. The delirium was usually mild, and if not, was easily controlled by the ice-bladder and chloral. The last-mentioned drug was also very satisfactory in overcoming insomnia, in doses of gr. v–xiv. The icterus and gastric symptoms rarely called for special attention. The parotitis and abscesses occurring during convalescence, demanded a prompt use of the knife to give exit to the pus. The neuralgia which sometimes remained after the disease, yielded most rapidly to a judicious application of electricity.

DR. R. MURCHISON, ENGLAND.

This author states his opinion that we may prevent the occurrence of uræmic intoxication, which is one of the main causes of death in uncomplicated cases, by keeping up the action of the kidneys from the first. He prescribes:

908.	R.	Spiritus ætheris nitrosi dulcis,	f. ʒ ij	
		Acidi nitrici diluti,	f. ʒ j	
		Syrupi simplicis,	f. ʒ j	
		Decocti hordei,	Oij.	M.

To be taken in the twenty-four hours. Should symptoms of jaundice manifest themselves, hydrochloric acid should be substituted for the nitric.

DR. HENRY HARTSHORNE, PHILADELPHIA.

Relapsing fever was very prevalent in Philadelphia and New York, in 1870, with a mortality of a little over fifteen per cent. of the cases. From the experience of this epidemic, Dr. H. recommends, after a mild cathartic at the beginning—and if headache be severe—a few cups or leeches to the back of the neck. Cooling diaphoretics may be given, as solution of citrate of potash or acetate of ammonia. After the crisis, which occurs from the fifth to the seventh or eighth day, quinine may be given in moderate doses, at least until the relapse. Many persons will require support, especially in the third week, by beef tea, alcoholic stimulants, etc. It has been proved that any amount of quinine will not prevent the relapse, and it is therefore useless to administer it with this object in view.

ACUTE RHEUMATISM.

SALICIN AND SALICYLIC ACID TREATMENT.

DR. T. J. MACLAGAN, DUBLIN.

This author (*Lancet*, March, 1876,) praises the action of *salicin* in acute rheumatism. Relief is afforded in forty-eight hours or less.

909. R. Salicin,

gr. xv.

This amount in water every three hours.

Later observers corroborate this statement; gr. x every two hours, is preferred by some.

Decoction of the inner bark of the willow (*salix fragilis*, etc.,) which contains the active principle, *salicin*, has also been used with excellent results.

Salicylic acid, now produced by synthesis, was formerly derived from salicin. Dr. L. TRAUBE, of Berlin, and many following him, give it high praise in acute and subacute rheumatism. It disperses the fever and soothes the pain. Grains xv to xxx, in powder, in a capsule, may be given every hour. Or:

910. R. Acidi salicylici,
Sodii boratis,
Glycerinæ,
Aquæ menthæ,

$\overline{3}$ ss
 $\overline{3}$ j
f. $\overline{3}$ j
f. $\overline{3}$ v. M.

A tablespoonful as occasion may require.

The borax makes the acid more soluble.

Dr. N. B. KENNEDY, of Texas, remarks in the *Med. and Sur. Reporter*, 1883, that he has for a number of years used the following formula, with the happiest effects. It is quite pleasant to take.

911. R. Sodii bicarbonatis,
Acidi salicylici,
Glycerinæ,
Aquæ puræ,

$\overline{3}$ ij
 $\overline{3}$ ij
āā f. $\overline{3}$ ij. M.

S.—One teaspoonful every 4 hours.

The carbonic acid gas is set free, and the sodium uniting with the salicylic acid forms salicylate of sodium, which is held in suspension by the glycerine.

In administering the salicylates, lasting troubles with audition are

often found to remain, owing to thickened tympanum, brought about by the vaso-paralytic action of these remedies. Dr. SCHILLING, (*Glasgow Med. Jour.*, 1883,) has discovered that in most cases—at least in three-fourths—this may be prevented by combining ergot with the salicylate. He prescribes :

912. R.	Ergotinæ,	gr. xv
	Sodii salicylatis,	ʒ ijss
	Aquæ,	f. ʒvj.

A tablespoonful every hour.

Dr. STRICKER, of Berlin, in a *résumé* of his experience with salicylic acid, says :

1. If the salicylic acid act thoroughly in the manner expected of it, there should be, during the treatment of articular rheumatism by a certain quantity given internally, no return of the disease in a hitherto unaffected joint, nor any secondary inflammation of the endocardium or other serous membranes.

2. In order to prevent a relapse, it is necessary to continue the use of the salicylic acid in small doses for some days after the end of the principal treatment.

3. Salicylic acid is of doubtful use in chronic articular rheumatism.

4. It is not likely to be useful in gonorrhœal and dysenteric rheumatism, or in septicæmic inflammation of the joints.

The following is an eligible formula, which forms the salicylate of soda :

913. R.	Acidi salicylici,	ʒj	
	Spts. eth. nit.,	f. ʒvj	
	Sodii bicarb.,	gr. lxx	
	Spts. lavand. comp.,	f. ʒij	
	Aquæ,	f. ʒij	
	Syr. aurantii corticis,	q. s. ad f. ʒvj.	M.

One teaspoonful every three or four hours.

In preparing this prescription, mix the acid and the spirits of ether in a bottle, then add the soda, and afterward the water, gradually, and after effervescence ceases, the spirits of lavender and syrup. This prescription agrees well with the stomach, and is pleasant to take.

In large doses of salicylic acid, there is danger of producing severe cerebral symptoms, sick stomach and rapid collapse. Dr. E. PRIDEAUX (*Practitioner*, Sept., 1878,) has found this can be obviated by giving salicylate of ammonia. He prefers to produce a combination of salicylate of soda and ammonia, as in the following prescription :

914. R. Sodii bicarb., Ammon. carb., Acid. salicylic, Aquam,	āā gr. v gr. xx ad f. 3j.	M.
---	---------------------------------	----

For one dose.

This avoids successfully the unpleasant symptoms alluded to.

DR. ROBERTS BARTHOLOW, OF PHILADELPHIA.

In summing up the testimony, it is clear that *salicylate of soda* is a most valuable remedy in acute rheumatism, but various unpleasant, even dangerous symptoms arise in some cases. Violent headache, vertigo and *tinnitus aurium* are common; wild delirium is an occasional symptom; weakness of the heart, obscuration of the first sound and profound anæmia, are especially noted by GREENHOW; very great gastro-intestinal disturbance is now and then produced. EMPIS narrates a fatal case of acute rheumatism which was treated with salicylic acid. Great depression of the powers of life, in rare instances approaching collapse, has been observed. Sufficient is now known of the danger attending the salicylic medication of acute rheumatism, to justify the author in urging circumspection on the attention of his readers. The robust and vigorous rheumatics are proper subjects for this treatment, whereas in the pale, feeble and cachectic, especially those with weak hearts, this treatment must be pursued cautiously or must not be undertaken at all. Relapses are frequent, probably because of the very rapid elimination of the remedy, as suggested by Prof. SÉE, and hence its administration must be continued for some time after all of the local and systemic symptoms have subsided. The mineral acids, especially the hydrochloric, have lately been proposed as remedies for acute rheumatism. It is highly probable that the mineral acids check the formation of lactic acid in the blood. Whatever may be the nature of the action, good results from the treatment have been reported.

LEMON OR LIME JUICE TREATMENT.

One of the recent advocates of this practice,

DR. A. H. CHANDLER, NEW BRUNSWICK,

gives the following as the details of his plan, (*Canada Lancet*, Nov., 1877): Without regard to the condition of the bowels, unless previously much constipated, he usually begins with at least 10 ounces of lime-juice, increasing rapidly up to 18 or 24, in the twenty-four hours

—from $\frac{1}{2}$ an ounce to an ounce or more every hour, with not less than double or treble the quantity of cold, soft water—usually diluted and sweetened, however, to the patient's taste. Very often, on the second day, the amendment is decided, and the disease, in acute cases more particularly, sthenic or asthenic, generally subsides on the fourth or fifth day of treatment. One grain of opium is usually given, with or without lead, and tannin, night and morning, in order to restrain the bowels, which the juice has a tendency to relax. The first effects of such heavy doses is the rapid diminution of joint swelling, the diminished perspiration, together with steady falling of pulse, the latter often quite slow with a slight tendency to syncope. The majority of the cases require quinine, and supporting food about the sixth or seventh day, when convalescence advances rapidly.

THE ALKALINE TREATMENT.

DR. H. W. FULLER, LONDON.

The treatment of rheumatic fever adopted by Dr. FULLER, at St. George's Hospital, is *essentially alkaline*, and consists not only in administering salines and small doses of alkalies, but in pushing alkalies, as rapidly as possible, to the point of producing alkalinity of the secretions. He usually prescribes :

915. R.	Sodii bicarbonatis,	3 ^{iss}	
	Potassii acetatis,	3 ^{ss}	
	Liquoris ammoniæ acetatis,	f. 3 ^{ijj}	
	Aquæ,	f. 3 ^{iss} .	M.

For one dose. To be taken in a state of effervescence, in combination with

916. R.	Acidi citrici,	3 ^{ss}	
	Aquæ,	f. 3 ^{ij} .	M.

The quantity of the fluid not only takes off from the nauseous quality of the medicine, but promotes its absorption, and thereby facilitates its action. As soon as the urine manifests an alkaline re-action, the dose is repeated three times only in twenty-four hours; and on the following day, if the urine still remains alkaline, twice only. After three days, two doses only of this mixture in twenty-four hours usually suffice to keep the urine alkaline; and then Dr. FULLER adds two grains of quinine to each dose; or, if quinine be not well borne, he substitutes :

917. R.	Sodii bicarbonatis,	āā	3 ^{ss}	
	Potassii acetatis,		f. 3 ^{iss}	
	Tincturæ cinchonæ,		f. 3 ^{iss}	
	Decocti cinchonæ flayæ,		f. 3 ^{iss} .	M.

For one dose.

Day by day, as the tongue cleans, and the other symptoms subside, the quantity of alkali is cautiously diminished, until a simple quinine draught is taken; but the condition of the urine is constantly watched, with a view to the immediate administration of a small quantity of alkali, should the least acidity call for its use. The diet is another point on which Dr. FULLER lays particular stress. He gives beef tea or broth throughout, and, if stimulants appear to be needed, does not object to their being administered cautiously; but he withholds solid food until the tongue is quite clean, and has often proved to his class in the wards that a too early recourse to meat will induce a relapse, and prevent convalescence.

Under this treatment, Dr. FULLER says the pains commonly subside in five or six days, and the patients are seldom ten days in bed. Moreover, the heart may be regarded as safe from attack; for in two instances only, in the whole course of his hospital experience, has inflammation, either of the endocardium or pericardium, arisen after the patient has been twenty-four hours under treatment.

DR. FINCHAW, WESTMINSTER HOSPITAL, LONDON.

918. R. Potassii bicarbonatis, ℥j
 Potassii nitratis, ℥ij
 Liquoris ammoniæ acetatis,
 Aquæ, āā f. ℥ij. M.

A tablespoonful, in water, every four hours, with a full opiate at night, if the pain be severe.

He does not think it advisable to continue the alkaline treatment for any lengthened period; but to give quinine, in doses of two or three grains, every six hours, when the urgent symptoms begin to yield, especially if the sweating is over-profuse. He believes that, by giving quinine earlier than is generally the custom, convalescence is less tedious, and there is less chance of relapse. As regards cardiac complications, if pericarditis supervene, and there be sharp, catching pain, he applies a few leeches, followed by linseed poultices; should the pain be slight or absent, he omits the leeches. In all cases he applies, subsequently, one or more blisters. As to medicines, he continues the alkalies, giving at the same time a grain of opium every four or six hours. Should endocarditis manifest itself, he contents himself with the alkaline treatment, as he cannot satisfy himself that local remedies have any effect.

DR. F. J. FARRE, LONDON.

Dr. FARRE's ordinary mode of treating acute rheumatism, is the "alkaline." He usually commences with three five-grain doses of calomel, followed by :

919. R.	Magnesiae sulphatis,	ʒij	
	Tincturae sennae,	f. ʒ ij	
	Spiritus ammoniae aromatici,	℥xx	
	Infusi sennae compositi,	f. ʒ iss.	M.

For one dose.

He repeats this daily till the evacuations are natural. At the same time, he gives the bicarbonate or acetate of potash, in 20 or 30-grain doses, every four or six hours, according to the severity of the attack, generally using the former, but preferring the latter when there is synovial effusion.

When there is little or no perspiration, or when the heart is much excited, he adds 10 or 15 grains of nitrate of potash. He not unfrequently, also, gives 1 grain of opium every night. For local treatment, especially of the smaller joints, he relies chiefly on the tincture or liniment of iodine, using the tincture for women and children, the liniment for the robust; one or the other of these is used in almost every case, and with nearly certain relief. When, however, there is synovial effusion, Dr. FARRE uses either mustard or cantharides plaster. Mustard is always useful, especially when applied to the larger joints, as the shoulder. The appetite being always faulty, Dr. FARRE gives milk diet until the tongue is clean, or cleaning. Meat given before it can be digested, immediately brings back pain in the joints. He keeps the patient between blankets. As soon as the pain has gone, and the tongue is clean, he gives bitter tonics, omitting or diminishing the alkali if the urine is alkaline or neutral.

DR. FLEMING, BIRMINGHAM, ENGLAND.

920. R.	Potassii bicarbonatis,	gr. xxx	
	Aquæ,	f. ʒ ij.	M.

For one dose, before each meal. Add $\frac{1}{2}$ an ounce of fresh lemon juice, and take the mixture during effervescence.

If there be high fever, from 1 to 3 minims of *Fleming's tincture of aconite* are added to each draught. If there be much pain in the muscles, in place of aconite, from 5 to 10 minims of *tincture of hemlock* are

added to each dose of the alkaline. If, on the other hand, the pericardium be affected, from 2 to 6 grains of the *iodide of potassium* are given. To relieve pain and secure sleep, Dr. FLEMING orders, at bedtime, a full draught of morphia and Indian hemp; as a drink, potassa water, or lemonade, freely. If necessary, colocynth and hyoscyamus pill is given to relieve the bowels. Cotton wadding is applied to the affected joints. Active and repeated counter-irritation, and poultices over the heart, are employed in cardiac inflammation. During convalescence, warm clothing, full diet, with quinine and iron. After considerable experience, Dr. FLEMING has found that this treatment has furnished very good results, and that the number of those attacked with cardiac inflammation *after* their admission into the hospital, is undoubtedly small. Placing the patient between blankets materially promotes perspiration, and prevents chills.

DR. W. C. BOTELER, OF MISSOURI.

This practitioner writes (1884): "I will ask that you give fuller credit to the following treatment in acute and subacute inflammatory rheumatism than is, as yet, given in any of our works:

- | | | | | | |
|------|----|---|----|--------|----|
| 921. | R. | Ammon muriat., | | | |
| | | Pot. brom., | āā | ℥ss | |
| | | Huxham's tinct., | | f. ℥ij | |
| | | Syr. zingiberis, | | | |
| | | Aquæ destil., | āā | f. ℥j. | M. |
| | | Ft. S.—℥j every two hours." | | | |
| 922. | R. | Deodorized iodoform, | | ℥iss | |
| | | Vaseline, | | ℥j. | M. |
| | | Ft. S.—Apply well e. i. l. to inflamed and swollen parts. | | | |

The great superiority of this internal and *external* treatment, has been attested to me by innumerable cases annually. Just what, the "modus medendi" of the iodoform locally is, I cannot positively say, but, I never have cases even of the worst and some *very chronic*, under treatment longer than twelve days, with this combination. I have used it with immediate relief on patients who had for twenty-nine weeks resisted the "alkalies" and "salicylates." Trying either alone I find more virtue in the ointment.

THOMAS KING CHAMBERS, M. D., ETC., LONDON.

This author calls rheumatic fever "a pleasant disease for the doctor to treat, though not for the patient to bear," and gives a very simple,

uniform plan of treatment, which, he states, hardly ever requires modification.

Bedding.—The patient's bed is made in a peculiar fashion. No linen should touch the skin. A slight calico shift or shirt may be allowed; but if the patients possess underclothing only of the prohibited sort, they are better naked. Even a linen front to the shirt is dangerous. The sheets should be removed, and the body carefully wrapped in blankets, the newest and fluffiest that can be got. The head is to be carefully protected from currents of air.

Fomentations.—Those joints or limbs which are swollen, red, or painful, are to be wrapped up in flannels, soaked either in hot water or a decoction of poppy-heads, with one-half ounce of carbonate of soda to each pint.

Curative Drugs.—If the skin is red, swollen, or painful about the joints, if motion is impossible or the cause of exquisite suffering, and especially if these phenomena are metastatic, then the "alkaline treatment" is employed, as follows:

923. R. Potassii carbonatis,
Aque camphoræ,

℞j
f. 5ij. M.

For one dose. To be repeated every three hours, day or night, when awake.

If, however, the above symptoms are insignificant, and the pain is felt more in the bones, being intensified by pressure rather than by motion, and fixed, not metastatic, then two grains of iodide of potassium are to be added to each dose. So soon as the symptoms take a favorable turn, the alkali is to be omitted altogether, and only the iodide of potassium given.

Palliatives.—Opium is to be administered in amounts proportionate to the subjective sensation of pain—from 1 to 2 grains at a dose. Immediately upon the relief of the pain, the quantity is diminished. Pure opium, in the pill and the tincture, are the best preparations. If the pain remains fixed in one joint after it has left the other places, leeches are to be applied there, and the part kept poulticed. Bruised laurel leaves may be mixed with the poultice. If the heart becomes affected, leeches and poultices are to be applied to the cardiac region.

Diet.—The food is to be varied to some extent, by the social and personal state of the patients. If they have been hearty and well-to-do persons before the attack, simple diet is proper, *i. e.*, bread and but-

ter, gruel and tea. If they have been ill-nourished, a pint of broth or beef tea is added.

DR. CARPANI, OF FRANCE,

thus formulates his experience in the treatment of acute rheumatism :

1. Salicylate of soda is useful in cases of acute febrile polyarthrititis, attended by highly localized articular manifestations. It is contra-indicated by the concomitance of cardiac affections, nervous troubles, or gastro-intestinal or renal difficulties.

2. Bisulphate of quinine is indicated in cases where rheumatism is a manifestation of, or associated with, malarial infection.

3. Benzoic acid should only be used when nephritis is a complication of acute articular rheumatism, attended by fevers.

4. Blisters are the most reliable agents for curing mono-articular rheumatism, or that form in which but few articulations are interested.

DR. J. MORTIMER GRANVILLE, ENGLAND.

This writer (*Brit. Med. Jour.*, 1883,) says: "In acute rheumatism, as early as possible in the case, give the mixture described below, in the diluted form in which I have prescribed it. Do nothing else, except to pack the painful joints in wraps of very loose cotton-wool, covered with light flannel; not oil-silk or any other vapor-proof material."

924. R.	Tinct. aconiti (B. Ph.),	℥xij
	Ammonii sulphidi,	℥xvj
	Aquæ menth. destil.,	f. ℥vj.

A fourth part every fourth hour till the fever is abated.

The mixture should not be prescribed in larger quantity than will suffice for four doses, on account of the tincture of aconite, and, more especially, the tendency of the sulphide of ammonium to decompose and deposit sulphur.

J. M. DA COSTA, M. D., PHILADELPHIA.

925. R.	Ammonii bromidi,	℥ss	
	Tincturæ aurantii corticis,	f. ℥ss	
	Aquæ,	f. ℥ijss.	M.

A dessertspoonful every three hours, except at night. In acute rheumatic fever, the beneficial influence of the bromide of ammonium is undoubted.

Over the pains and aches of acute rheumatism, this remedy exerts an unquestionable control; but in chronic rheumatism, it is decidedly

inferior to iodide of potassium. But slight amelioration follows its use, in persistent swelling of the joints of rheumatic origin, and none in rheumatism due to a venereal taint.

926. R. Atropiæ sulphatis, gr. $\frac{1}{10}$
Aquæ destillatæ, ℥x. M.

For one injection. To be thrown under the skin, in certain forms of muscular rheumatism, particularly wry neck, over or into the rigid parts. To be repeated once a day. Sometimes, the most marked and speedy relief follows this treatment, after ordinary remedies have failed.

927. R. Potassii iodidi, ʒj
Vini colchici radicis, f. ʒ ij
Morphiæ sulphatis, gr. iij
Syrupi, f. ʒ j
Aquæ, f. ʒ ij. M.

A teaspoonful three times a day, after meals, in muscular rheumatism.

928. R. Potassii carbonatis, ʒ ij
Potassii acetatis, ʒ iij
Vini colchici seminis, f. ʒ i
Syrupi tolutani, f. ʒ iss
Aquæ, f. ʒ ss. M.

A teaspoonful thrice daily, in lumbago, the following liniment to be rubbed in night and morning :

929. R. Chloroformi, f. ʒ ss
Linimenti ammoniæ, f. ʒ iss
Linimenti saponis, f. ʒ ij. M.

In a week or ten days, after the pain has subsided, apply :

930. R. Emplastri picis cum cantharide,
Emplastri logani, āā equal parts. M.

The quickest way of treating lumbago is by throwing $\frac{1}{16}$ of a grain of *atropia* under the skin, near the affected muscles. This will sometimes dissipate the attack as if by magic.

931. R. Tincturæ guaiaci, f. ʒ ij
Vini colchici radicis, f. ʒ ij
Potassii bicarbonatis, ʒ ss
Syrupi aurantii corticis, f. ʒ ij. M.

A dessertspoonful in water, thrice daily, in rheumatic arthritis. Also have the joints wrapped in cotton, and painted, morning and evening, with equal parts of tincture of iodine and alcohol, until they become sore. Sulphur baths are beneficial. Small blisters are also of service in the neighborhood of the joints, frequently repeated and dressed with:

932. R. Morphiæ acetatis, gr. $\frac{1}{4}$
Pulveris marantæ, gr. j. M.

Arsenic is an excellent remedy in rheumatic arthritis. It may be administered as follows :

933. R. Liquoris potassii arsenitis, f. ʒij
 Potassii iodidi, ʒij
 Syrupi, f. ʒiij. M.

A teaspoonful thrice daily, between meals, in water.

934. R. Vini colchici seminis, f. ʒj
 Potassii acetatis, ʒvj
 Spiritus ætheris nitrosi, āā f. ʒj. M.
 Syrupi,

A teaspoonful thrice daily, in pleurodynia.

Also use a stimulating liniment, and keep the parts warm.

935. R. Potassii iodidi, ʒij
 Vini colchici radicis, f. ʒss
 Extracti cinchonæ fluidi, āā f. ʒiss. M.
 Elixir cinchonæ,

A dessertspoonful thrice daily, in muscular rheumatism.

Cinchona disguises the taste of the iodide of potassium. Quinine is often serviceable alone in those cases in which the joints are not affected, and in which there is pallor and loss of strength.

936. R. Potassii iodidi, ʒij
 Tincturæ belladonnæ, f. ʒiss
 Syrupi aurantii corticis, f. ʒss
 Aquam, q. s. ad f. ʒiij. M.

A teaspoonful thrice daily, in muscular rheumatism associated with vesical incontinence.

In case of swelling and stiffness of the joints following acute rheumatism, nothing does so much good as *sulphur-baths*. When practicable, send the patient to the sulphur-baths of Virginia, to get rid of these remnants of the acute attack. Artificial sulphur-baths are also useful. When these are not practicable, great benefit will be obtained from the iodide of potassium and compound tincture of guaiacum.

937. R. Potassii iodidi, ʒij
 Tincturæ guaiaci compositæ, f. ʒvj. M.

A dessertspoonful thrice daily.

This prescription is very efficient, though not an elegant one. It is very unpleasant to take. There are few who will go on with it for a long enough time to get its full therapeutical effects.

In the treatment of the *subacute form of muscular rheumatism*, nitrate of potash is very valuable. This is an old remedy, but the advance of therapeutics has shown no other, in this form of rheumatism, of which our author thinks so highly. It may be conjoined with Dover's powder.

938. R. Potassii nitratis, gr. xv
 Pulveris ipecacuanhæ compositi, gr. iij.
 For one powder, to be taken every fourth hour.

NOTES ON REMEDIES.

Aconite is, according to Dr. REITH, (*Edinburgh Medical Journal*,) ordinarily given in too large doses in acute rheumatism. He obtains good results in mild cases by its agency alone in small and often-repeated doses, ($\text{m}\frac{1}{4}$ of the tincture every two hours.) In some cases he combines it with the full alkaline or blister treatment.

Chloral has been extolled in cerebral rheumatism by M. BOUCHUT.

Cimicifuga was brought forward as a remedy in acute rheumatism by Dr. F. N. JOHNSON, of New York. It is less employed in the acute than in the chronic form.

**Colchicum* relieves the symptoms and shortens their duration, when combined with small doses of calomel, ipecacuanha, alkalies and opium. It is less beneficial, according to Dr. FULLER, in the weak and nervous than in the robust; less in purely fibrous rheumatism than in that complicated by synovial inflammation; less in proportion as the fever exceeds the articular swelling and as the urine is less highly charged with the lithates. During its use, its operation must be carefully watched, the bowels kept regulated, and if the lithates disappear from the urine, if the pulse become weak, if faintness, or nausea, or purging supervene, it must be discontinued. The acetous extract, or the inspissated juice, (gr. j-iss,) or the wine (mxv - xx ,) may be given twice or thrice daily.

Ergota, in full doses internally, is a very efficacious remedy in many cases of acute rheumatism.

Ferri Chloridi Tinctura has been employed by Dr. RUSSELL REYNOLDS, of London, in doses of about mxxx every six hours, with the effect of shortening the duration of the disease, and relieving the joint affection.

Kairin is said by Dr. H. MENCHE, of Berlin, (*Blatt. Klin. Med.*, 1883,) to act more quickly in reducing the temperature in acute inflammatory rheumatism than the salicylic compounds. He gave large doses, as much as ʒj a day, in divided amounts every two or three hours.

Limonum Succus. See page 535.

Lithium Bromide. GARROD first introduced the lithium salts into medical practice for the treatment of rheumatism, and in his recent Lumleian lectures has demonstrated anew their exceptional value. Recently the compound of lithium and salicylic acid has been brought forward as a more effective remedy in the rheumatic diseases. The sub-acute and chronic cases, and the so-called rheumatic gout, are the forms of disease in which the lithium salts are most serviceable.

Manaca. This is a preparation of the root of *Franciscea uniflora*, a Brazilian shrub, introduced by Messrs PARKE, DAVIS & Co., Detroit. It is official in the Brazilian dispensaries, as an anti-syphilitic. The dose of the fluid extract is gtt. x-xx, of the powdered root, gr. x. Administered

in chronic rheumatism, it is stated to have exerted a favorable influence on most cases. In some cases of acute rheumatic arthritis it is also said to have shortened the attacks. Dr. GEORGE HERSCHELL, of England, has had excellent results in gonorrhœal rheumatism from five-minim doses, every three hours, of the fluid extract. He has also used it successfully in simple acute rheumatism.

Methyl Salicylate, is said by a writer in the *Ed. Med. Jour.*, 1884, to give prompt relief in many cases of rheumatic pain. The prescription is :

939. R. Methyl salicylatis,
Olei olivæ, equal parts.
Apply externally to joints affected by acute rheumatic pains.

**Opium*, in large and repeated doses, is given by SIR D. CORRIGAN, of Dublin. He commences with gr. j every two or three hours, and increases the dose, both in frequency and quantity, until the patient feels decided relief, when it should be kept at that dose until the disease is steadily declining. The average quantity required in twenty-four hours, is about gr. xij, and even that amount does not affect the head; occasionally, it produces diarrhœa, when astringents are required to check it. The tolerance of opium, in this disease, is remarkable.

Potassii Acetas is much employed. The dose usually given is about gr. xxx every four hours.

**Potassii Bicarbonas* is a favorite remedy. Dr. A. B. GARROD, of London, gives gr. xxx. in solution, every four hours, until the joint symptoms and febrile disturbance have completely disappeared. Dr. FULLER, of London, considers that \bar{z} iss is the minimum dose, which will suffice to overcome the acid condition of the blood during the first twenty-four or forty-eight hours of treatment, and in sthenic cases, \bar{z} ij are often needed. Small doses he regards as useless.

Potassii Bitartras, in doses of gr. xv-xx, every four hours, given in mint-water, alone or in combination with tincture of hyoscyamus (m_{xxx}-xl), is said to produce good results.

Potassii Nitras. Dr. BASHAM, who instituted a series of experiments to ascertain the real value of nitre in acute rheumatism, arrived at the following conclusions: 1. That it is a most valuable remedy in doses of \bar{z} ij-ij, in Oiv of barley-water, daily. 2. That it acts by restoring the saline constituents of the blood, and by lessening the excess of fibrine. 3. That there is a certain amount of exemption from cardiac complication, and that cardiac inflammation, when it supervenes, is more amenable to remedies. 4. That the tendency to collapse is much diminished, and the acute or inflammatory symptoms usually give way on the third or fourth day.

Potassii Permanganas. Dr. FINN (*Canada Medical Journal*, January, 1868,) writes in the highest terms of the efficacy of the permanganate of potash, in the treatment of rhematism, gr. $\frac{1}{2}$ thrice daily, which he believes

to be due to the large proportion of oxygen which it contains. This salt promotes the transformation of lactic into carbonic acid.

Propylamine. See *Trimethylamine*.

Quinæ Sulphas, according to Dr. FULLER, should, ordinarily, only be used as a restorative when the febrile paroxysm is beginning to abate, and not as a cure during the active stages of the disease.

Salicin and Salicylic Acid. See above, page 533.

Trimethylamine has, at times, been popular as a remedy in rheumatism. Its favorable action is thought to be on account of its alkaline properties. There can be no doubt that it reduces the temperature, diminishes the pulse, and removes the pain in some cases of acute rheumatism and gout, and that in a very short time—from six to thirty-six hours after its use. The more *acute* the disease, the more marked is the effect of the remedy. In chronic cases, there is little or no benefit from it. The dose is gr. iij every two hours. It is often confounded with *Propylamine*, which is a different substance.

Zinci Cyanuretum. This remedy was suggested by LUTON, of Rheims, some time since. Lately, DESCHAMPS has used it with success, and recommends it highly. He employs the following formula :

940. R.	Zinci cyanid.,	gr. $\frac{1}{12}$	
	Pulv. acaciæ,		
	Sacch. lactis,	āā	q. s. M.

For one pill. To be taken to the number of ten in twenty-four hours.

In recent acute cases, the relief is immediate.

LOCAL APPLICATIONS.

Baths. Dr. R. J. BUTLER (*Medical Press*, August 5th, 1868,) has found great benefit in acute rheumatism, from baths containing *valerian*. He directs :

941. R.	Valerianæ radicis,	℥bj
	Aquæ,	C.j.

Boil together for a quarter of an hour, strain the liquor, and add to an ordinary bath, at 98°, the patient remaining in it for twenty or thirty minutes.

For any remaining inflammation about the joints, he applies linseed-meal poultices, prepared with a strong decoction of valerian. In the arthritic form of the disease, he considers valerian of extraordinary efficacy in subduing the pain and inflammation.

Prof. DA COSTA, of Philadelphia, recommends sulphur-baths, artificial or natural, for the swelling and stiffness of the joints following acute rheumatism.

**Blisters.* The blister treatment has been prominently advocated. Dr. HERBERT DAVIS directs armlets, wristlets, and even fingerlets of blister-plaster, to be applied at the time the inflammation is most acute, lin-

seed-meal poultices being subsequently applied to keep up the serous discharge. He places these blisters entirely around the affected limbs, and in the case of the knees, orders them at least three inches wide, regarding any slight strangury which may arise as of little importance, compared with the benefit afforded by the free vesication. This treatment may be aided by a *mild* course of alkali, but he does not regard this as essential, excepting when the rheumatic virus attacks the heart in the first instance, and seems to hesitate in fixing itself upon the joints. Dr. RINGER, of London, considers all the alleged good effects may be obtained by the use of "flying blisters." Dr. A. FLEISCHMAN (*Lancet*, May 1st, 1869,) looks upon a blister about two inches square, placed an inch and a half below the left clavicle, as an almost certain preventive of cardiac mischief.

Cold Packing. Applying a mixture of pounded ice and salt to the swollen joint for not longer than five minutes at a time, will sometimes relieve the pain promptly. Or, the cold pack may be employed. Two dry blankets are laid on a bed, and over these a sheet, wrung out as dry as possible with cold water, is laid so that when folded it will completely embrace the affected limb or limbs. The patient is placed upon the sheet, which is tightly folded over the rheumatic parts and covered immediately with the blankets. This may be renewed every two or three hours.

Hot Packing. This is much esteemed by some practitioners. As employed in Mount Sinai Hospital, New York City, the method is as follows: The bed is prepared by spreading over it a sheet of rubber cloth, and over this a blanket. Hot blankets are then wrung out of water of a temperature about as hot as the hand can bear, and with these the patient is enveloped. Two or three thicknesses of dry blankets are superimposed, and the whole retained in position till the patient is free from pain—a time varying from two to six hours. When the pack is applied, the patient sweats very profusely, as might be supposed, and after its removal, no danger has been found to occur. If the pain is confined to one joint, that joint is enveloped by the hot blanket, which is removed as soon as the pain disappears. One case of chronic rheumatism, extending over a period of seven months, was treated by means of local packings, and after slight relapses, occurring at intervals for five weeks, was discharged perfectly cured. This method presents the advantage in private practice of not shocking the friends of the patient as much as the ice and ice-water packings.

CHRONIC RHEUMATISM.

PROF. FRANCIS DELAFIELD, M. D., NEW YORK.

Cases of this disease are apt to be extremely obstinate. A fair general plan of treatment is to give increasing doses of *iodide of potassium*, commencing with gr. x, thrice daily, together with counter-irritation over the affected muscles. The very best form of counter-irritation, and one almost sure to produce decided relief, is *massage*, shampooing or kneading, by a skilled manipulator. Where the services of such a person cannot be obtained, the next best plan is to apply a succession of blisters, together with painting the part for a long time with tincture of iodine.

DR. ROBERTS BARTHOLOW, OF PHILADELPHIA,

says that the alkaline springs have long had a deserved reputation for the cure of chronic rheumatism. With the internal use of the waters should be conjoined baths, douches, etc. Rheumatic affections of internal organs are equally amenable to the same treatment. The Turkish bath is highly useful. In the so-called rheumatic gout, colchicum with alkalies is extremely serviceable. Attacks, without decided pain and inflammation, of *soreness of joints*, which have been the seats of gouty attacks, or about which nodosities have been deposited, are relieved by colchicum. Manganese is useful in rheumatic patients of gouty antecedents.

DR. HENRY HARTSHORNE, PHILADELPHIA.

This author considers the treatment of chronic rheumatism largely experimental. Local treatment generally does more for it than medicine. He has found a liniment containing turpentine, oil of sassafras, ammonia, and laudanum, diluted with soap liniment, of great use. Where pain is considerable, chloroform or aconite liniment is better. Blisters may be applied in bad cases. For rigidity of the joints, and for pain in them, *pouring hot water* continuously over the parts affords great relief. The hot or vapor-bath, as well as the hot dry-air bath (130° to 200°) will sometimes prove very effective.

DR. FULLER, LONDON.

In attacks of *lumbago*, this author recommends brisk purgatives at the commencement of the attack, when the bowels are costive and the urine scanty, high-colored, or loaded; but when the urine is abundant and light-colored, the bowels regular, and the dejections natural, the value of purgatives is not so apparent. Cupping on the loins is of essential service—dry cupping when it is not thought desirable to extract blood. Baths rendered alkaline by carbonate of soda or potash; hot fomentations formed by placing across the loins flannels wetted with an alkaline and opiate solution, and covered with a piece of gutta percha. In obstinate cases, stimulating and opiate embrocations. As excellent liniments in these cases he gives:

942. R.	Olei cajuputi,		
	Tincturæ opii,	āā	f. ʒ ij
	Olei terebinthinæ,		f. ʒ iv
	Linimenti ammoniæ,		f. ʒ j.

M.

For a liniment.

Or:

943. R.	Liquoris ammoniæ,		℥ ʒ j
	Tincturæ opii,		f. ʒ ij
	Tincturæ cantharidis,		f. ʒ iij
	Linimenti saponis,		f. ʒ x.

M.

For a liniment.

Guaiacum has an old repute in chronic rheumatism. Dr. WILLIAM AITKEN gives the following formula:

944. R.	Pulveris guaiaci,		ʒ j
	Pulveris rhei,		ʒ ij
	Potassii bitartratis,		
	Sulphuris sublimati,	āā	ʒ j
	Pulveris nucis moschatæ,		ʒ ij
	Mellis,		℔ j.

M.

Of this compound, two large spoonfuls are to be taken night and morning.

It is used in some large hospitals, both civil and military, in the treatment of old chronic cases. The beneficial effects of *guaiacum* are obtained in those cases which are unaccompanied by perspiration, and in which the secreting organs are gently excited by this drug. When the diaphoresis is free, it does little good. The following is used in the Philadelphia hospitals:

945. R. Pulv. guaiaci resin.,
 Potass. iodidi, āā gr. x
 Tinct. colchici semin., f. ʒss
 Aq. cinnamomi,
 Syrupi, āā q. s. ad f. ʒj. M.

A dessertspoonful to a tablespoonful thrice daily.

946. R. Fl. ext. polymniæ uvedaliæ, ʒij
 Adipis, ʒj. M.

It should be applied twice daily. (*Med. and Surg. Reporter*, 1878.)

NOTES ON REMEDIES.

Cajuputi Oleum. In doses of gtt. j-ij, in pill form, this has been given internally, in chronic rheumatism, with alleged good effects. It is also an ingredient in liniments.

Cimicifuga. In some cases of this disease, this drug exerts a prompt and very positive healing action, but, as it frequently fails, it is impossible to point out the indications for its use.

Guaiacum has long been esteemed in this disease; f. ʒj of the tincture may be administered thrice daily. (Or F. 944.)

Iodoform has been recommended by Dr. KNOLL.

947. R. Iodoformi,
 Ferri redacti, āā gr. xliij
 Extr. glycyrrhizæ, q. s. M.
 Make sixty pills. Two to be taken thrice daily.

Kerosene. See *Petroleum*.

Morrhua Oleum is one of the most efficient remedies in chronic rheumatism with dyspepsia and general exhaustion.

Opium and its alkaloids are especially valued in the form of hypodermic injections, as additions to liniments, etc.

Petroleum has been used with much advantage, locally, well rubbed into the affected parts. Internally, a teaspoonful of kerosene, in water, every other night, has cured some cases. (*British Medical Journal*, May, 1872.)

Phytolacca Decandra. The poke-root has quite an extended reputation in this disease. No doubt it cures certain cases with astonishing promptness. The indications for its use are, however, not definitely known.

Polymnia Uvedalia, in ointment, is highly esteemed. (F. 946.)

Potassii Iodidum is the most generally successful of all remedies, in this disease. SIR JAMES PAGET has pointed out that the addition of carbonate of ammonia greatly increases its therapeutic action. Five grains of the iodine may be combined with three of the carbonate.

Rhus Toxicodendron. Drs. PHILLIPS and PIFFARD give high praise to this plant, in sub-acute muscular or tendinous rheumatism. The former says it is "positively invaluable." The latter has found it most useful

in non-syphilitic cases, where the pain is worse at night. He gives $\text{℥}\frac{1}{8}-\frac{1}{4}$ of a tincture made by macerating one part of fresh leaves to two of alcohol.

Sassafras. This is stated to be an efficacious remedy when the circulation is languid, the skin dry, and the kidneys, liver, and bowels, inactive. It has the advantage over guaiacum, of not being purgative. Dr. FULLER, of London, recommends the following formula :

948. R.	Sassafras radicis,	$\text{℥}\text{ijss}$	
	Mezerel,	$\text{℥}\text{iv}$	
	Taraxaci radicis,	$\text{℥}\text{iiij}$	
	Aquæ ferventis,	Oj.	M.

Dose, f. $\text{℥}\text{ijss}$, conjoined with a plentiful use of diluents.

Sulphur, internally, is often of benefit.

Terebinthinæ Oleum is a valuable addition to liniments and stimulating embrocations. Dr. BREMOND, of Paris, recommends the employment of baths of the vapor of turpentine, which may be administered by placing the body of the patient in a wooden chest, the head remaining free. Into this chest, the vapor of water is conducted, loaded with a fine spray of oil of turpentine. The temperature of this vapor-bath is 105° – 110° F., and the time during which a patient may be exposed to it, is about fifteen minutes.

Xanthoxylum. The bark of the prickly ash has long enjoyed a popular reputation in chronic rheumatism.

Electricity, in its various forms, has been very extensively tried in chronic rheumatism. Sometimes, the results are most satisfactory ; but Drs. BEARD and ROCKWELL state that, on the whole, in cases of long standing, they have been disappointed in its effects.

TYPHOID FEVER.

PROF. ALONZO CLARK, M. D., NEW YORK.

An average case of typhoid fever needs no medication whatever, except for the relief of symptoms.

For the *diarrhæa* he gives :

949. R.	Bismuth. subnit.,	$\text{℥}\text{j}$
	Morphiæ sulph.,	gr. j.

M. et div. in chart. No. xii. One to four a day.

The common astringents, tr. kino and tr. catechu, may be employed ; and the decoction of blackberry root is sometimes very serviceable.

In some cases, it requires the moderately free use of opium to restrain the diarrhœa.

For the *cough*, the tonic expectorants will be most likely to do good. Perhaps one of the best that can be used, is the compound *tincture of benzoin*, in doses of ten drops on sugar, once in three or four hours. A very good combination is the tincture of the balsam of tolu and the *mistura guaiaci*:

950. R. Mist. guaiaci, ℥j-℥ss
Tr. balsam tolu, gtt. vj-x. M.

This can be repeated every two, three or four hours. Sometimes the inhalation of the vapor of warm water seems to be required for one or two hours each day.

Restlessness is one of the prominent features of the disease. This will very frequently be entirely quieted by sponging the surface of the body with warm or cold water. If the temperature is high, cold water is better than warm; and in some cases a Dover's powder will be required.

In regard to the *temperature*, if it be so high that the temperature of the body is 105° F., the patient may be placed in a bath having a temperature of 95° F.; then some of the warm water can be removed, and be replaced by cold water until the bath has been reduced to 80° F. If the patient is permitted to remain in such a bath twenty minutes, the temperature is usually reduced one, two, three, four or even five degrees. He is then removed from the bath, put back into bed, and it will be several hours, usually, before the temperature will rise as high as it was before using the bath. When it rises, another bath is to be given, and in that manner repeated as often as may be necessary to keep the temperature below the point of danger.

For the *hemorrhage from the bowels* there is but little that can be done, unless, in addition to absolute rest, the fluid extract of ergot be administered.

For *perforation*, full doses of opium offer the best and only chance.

The *diet* must be liquid, and nutritive. Milk, beef essence, etc., are demanded. Fresh air and free ventilation must never be neglected.

DR. ETHAN SPENCER, OF INDIANA.

This author (*Therapeutic Gazette*, April, 1884,) has found the *hydropathic treatment* almost the only successful treatment in cases of

typhoid fever, where there is much involvement of the bowels, and where almost all food or medicines irritates and distresses the patient. The surface in these cases is dry, the tongue dark and dry, the energies of the system fail and the functions are suspended; the temperature need not be excessive. Here the tepid pack should be applied. Great care should be observed, by means of warmth to the extremities, to aid the system to react and thus to relax the surface and bring about a natural perspiration. The patient will often fall into a quiet sleep and rest for twelve hours. The pack should then be repeated. In three days the tongue usually becomes moist, the dark color disappears, and every symptom becomes more favorable.

BOSTON CITY HOSPITAL.

The treatment at this hospital, as practiced under the surveillance of Drs. STEDMAN and DRAPER, is a modification of the bathing process, occupying an intermediate position between the full carrying out of cold-water bathing and the expectant treatment. On entrance, patients are immediately put to bed, and their diet restricted to milk, of which they are expected to drink about two and a half quarts in twenty-four hours. This is perhaps the most important part of the treatment; at first they may object to milk, but on becoming accustomed to it they take it freely. To keep up the strength is the chief aim, and if insufficient nourishment is taken, by giving a small amount at regular intervals, the end sought for is accomplished. This diet is continued till the temperature has been normal for several days, when light soups and easily-digested farinaceous articles are given; in a few days more, a gradual return to regular, though still somewhat restricted, diet is allowed.

To lower the temperature, sponge-baths, of equal parts of cold water and alcohol, are quickly applied to the surface of the body, which is immediately dried.

In mild cases of diarrhoea nothing is done, two or three loose dejections per diem not being considered harmful; when, however, it becomes excessive, it is quickly checked by a pill composed of a grain of opium and a quarter of a grain of sulphate of copper.

In delirium and wakefulness, when the patient can be made to tolerate it, the ice-cap is of great benefit.

The frequent use of a mouth-wash, of equal parts of glycerine and

rose-water, is very pleasant to the patient, relieving the foul taste and dryness of the mouth.

In very mild cases, 20 drops of dilute nitro-muriatic acid may be given every two hours.

Excepting the medicines above mentioned, no drugs are given, but every attention possible is shown to the comfort, nourishment, sleep and general welfare of the patients.

PROF. WILLIAM PEPPER, M. D., PHILADELPHIA.

This practitioner (*Boston Medical and Surgical Journal*, Oct., 1877,) recommends, in the second week of the disease, when the abdominal symptoms of pain and diarrhoea have fully set in, the use of *nitrate of silver*.

951. R.	Argenti nitratis,	gr. $\frac{1}{4}$	
	Extracti belladonnæ,	gr. $\frac{1}{4}$	
	Extracti opii aquosi,	gr. $\frac{1}{8}$ - $\frac{1}{2}$	M.
For one pill. Three times a day after food.			

Under this treatment, diarrhoea and tenderness diminish, and patients make very rapid recoveries. In most cases very little stimulant is used. Milk and beef tea are the only articles of food he allows. Quinia is given with other tonics. Fever is reduced by frequent spongings of the skin of the entire body. When the high fever resists sponging, cool baths are employed. Indiscriminate bathing in typhoid fever is often extremely injurious. The best time for the use of the cold bath is in the early stage, during the first week or ten days, in cases where the temperature rises above 103° , and is not controlled by frequent spongings, large doses of quinia, diaphoretics, etc. When the fever in subsequent stages runs high, it is of the nature of a sympathetic fever, largely dependent on the amount of intestinal lesion; hence, cold baths are less valuable at that time and attended with more risk. Nitrate of silver is used, both with the hope of limiting the amount of specific follicular catarrh of the intestines, and with the intention of favorably modifying the secondary sympathetic symptoms. Dr. PEPPER has cured *thirty-nine* out of the *forty* cases of typhoid fever in which it has been employed, by this nitrate of silver treatment.

PROF. KARL LEIBERMEISTER, M. D., TUBINGEN.

This writer, whose essay on typhoid fever forms part of ZIEMSEN'S

Cyclopædia, believes it possible that both *iodine* and *calomel* may exert a specific influence in cutting short an attack of typhoid fever, if given early. He gave iodide of potassium, ℥j-℥j, in the twenty-four hours; and when he employed calomel, gr. viij, three or four times in the same period. He prefers the latter, and recommends it in the stage of the disease before the ninth day.

His main reliance, however, is *the antipyretic treatment*. This is to be carried out by cold baths and quinine. For adults, the full-length cold bath 60° to 68° Fahr., is to be preferred. The duration of the bath should be ten minutes; but, if the patient is feeble, seven or five minutes. After the bath, the patient is to be wrapped in a dry sheet; put to bed, and given a glass of wine. In very severe cases, the bath should be repeated every two hours; the average cases require four to eight baths daily. *Contra-indications* to the baths are hemorrhage and perforation of the bowels, weakness of the heart's action, and extreme debility.

Quinine comes next as an antipyretic. But it must be given without timidity, and in a definite manner, as follows: from gr. xxij to gr. xlv *must positively be taken within the space of half an hour, or, at most, an hour*. Six or eight grains every ten minutes is a good rule. If this is boldly done, cinchonism is produced, and a material fall of temperature follows. This dose is repeated once every forty-eight hours; or, if the remission is not marked, once in twenty-four hours, the quantity being increased. When the fever spontaneously manifests a tendency to remissions, the quinine is of little value. This dose should reduce the temperature in the rectum to 100° Fahr. Of the two antipyretics mentioned, our author would generally rely on the quinine, if it is properly given.

When the fever is of such obstinancy that additional antipyretic effects are required, he usually gives from 11 to 22 grains of powdered *digitalis*, extended over a period of about thirty-six hours; this should be followed by a full dose of quinine, say gr. xl, in one hour, as above directed.

The diarrhœa he controls with opium, in small and repeated doses; meteorism, with cold compresses to the abdomen, changed every half hour, cold enemata, and frictions with turpentine; hemorrhage from the bowels, with opium (gr. ij, in the twenty-four hours,) ice to the abdomen, and perfect quiet.

PROFESSOR S. JACCOUD, PARIS.

This able clinical teacher, like most French and English physicians, discredits the cold bath treatment of typhoid. He abjures from the commencement all debilitating measures, such as purging. If needful, a mild laxative of Seidlitz water is given. During each twenty-four hours the patient takes a cordial mixture containing 2 or 3 scruples of extract of cinchona bark, 2 drachms of tincture of cinnamon, an ounce of syrup of orange peel, 3 ounces of red Bordeaux wine, and an ounce of old brandy, which may be increased, as occasion requires, to 3 or 4 ounces. The main constituent of the food is beef soup, given at least twice daily; about 8 ounces of old Bordeaux wine are allowed, and this is taken alternately with the stimulant mixture just mentioned. The ordinary beverage is lemonade, containing one-fourth of red wine. If it appear needful, all this is supplemented at the end of the second week by nutritive enemata twice daily. This mode of alimentation is stated to be well tolerated by the stomach and bowels.

When the patient's temperature reaches 102° Fahr., with a rise in the evening not above 103° , M. JACCOUD directs that twice during the day the whole body should be well sponged with aromatic vinegar. If the temperature rises higher, three, or even four, of these applications are made daily. They are continued, more or less frequently, until the close of the fever, and both comfort and benefit are derived from their refreshing and cooling effects.

Should any threatenings of bronchial or pulmonary implication arise, they are met by dry cupping.

DR. ROBERTS BARTHOLOW, OF PHILADELPHIA.

Influenced by the germ theory of disease, *carbolic acid* has been much prescribed in typhoid fever. Whether the theory be true or false, there can be no doubt of the good effects of creosote and of carbolic acid in this disease. PÉCHOLIER, of *Montpelier*, has given creosote in sixty cases of typhoid, set apart for study of the action. Good effects were obtained; the temperature kept down; the delirium and insomnia were much less; the intestinal disturbance declined, and thus the violence of the disease was distinctly lessened. M. CHAPELLE maintains that it cuts short an attack of typhoid. Recently ROTHE has treated a number of cases of typhoid fever, with great success, by a mixed antiseptic and antipyretic method, in which he employs carbolic

acid and iodine administered in infusion of digitalis. The author has, for several years, treated the cases of typhoid fever which have been in his hands with a drop each of tincture of iodine and liquefied carbolic acid every two or three hours, with results which were certainly most favorable. As ROTHE has observed, this antiseptic treatment lowers the heat, lessens the diarrhœa, improves the mental state, and indeed diminishes the severity of the disease remarkably. The dose of carbolic acid for internal use ranges from half a grain to two grains, or of the liquefied acid, from half a minim to two minims. It may be given in mint-water which covers the odor somewhat. This quantity may be repeated every hour or two, if necessary, without producing ill effects, if not too long continued. BOUCHARD and GIMBERT have sometimes reached as high as fifteen grains a day.

From REISS, who regards salicylate of soda as a specific in typhoid fever, to FILATOW, who holds that it is worse than useless, there are various shades of opinion, with the weight of authority decidedly in favor of the view that it is a remedy of real value. Since REISS, no one has claimed in this remedy a specific for typhoid, but it is generally admitted that it renders the course of the disease milder by keeping the temperature within safe limits (TOMKINS, HALLOPEAU, BUTT, IMMERMANN, SENATOR, etc). WEISS concludes, after an experience of ninety-six cases of typhoid in children, treated with salicylate of soda, that it is a powerful antipyretic in the typhoid of children, and that, while it does not shorten the course of the disease, it renders it much milder. He finds that the results are better, when the typhoid of children is treated with salicylate of soda, than have hitherto been obtained from quinine and cold baths. A still more recent English experience is to the same purport as the German.

952. R. Acid salicylic,
Sodii bicarbonat.,
Aquæ,
Syrup. simpl.,

℞iv
℞j
℞jss
℞ss. M.

A tablespoonful every two hours until the temperature is reduced to the proper point, at which it may be held by half the dose at the same interval, unless the occurrence of an exacerbation requires the exhibition of a larger quantity. It should be borne in mind that this remedy is not safe if the heart is very weak or if albuminuria is present.

THE TURPENTINE TREATMENT OF TYPHOID.

The oil of turpentine has been very largely used in this disease, but in different manners. The typical turpentine treatment is thus described by Dr. A. MAFFITT, in the *Lancet*, February, 1876 :

As soon as the disease is diagnosed, oil of turpentine, in one-half-drachm doses, made up with mucilage of eggs, is given four times a day, and continued throughout its course ; and nourishment in the liquid state, such as beef tea, milk, chicken broth, eggs beaten up, etc., is given freely, and a stimulant, generally brandy, administered as the symptoms indicate.

Dr. R. P. WHITE, of the Meath Hospital, Dublin, (*British Medical Journal*, December, 1877,) reports flattering success with this method. He gives the following turpentine mixture :

953. R.	Olei terebinthinæ,		
	Liquoris potassæ,	āā	3 ij
	Mucilaginis acaciæ,		3 iv
	Syrupi papaveris albi,		
	Syrupi floris aurantii,	āā	3 viij
	Aquæ camphoræ,	q. s. ad f.	3 viij. M.

A tablespoonful to be taken every fourth hour, the bottle being first shaken.

Since he commenced this treatment, he has never lost any case of typhoid, from either bronchitis or diarrhœa, or from its sequelæ of ulceration or hemorrhage. He does not know any symptom which contra-indicates the adoption of this plan.

According to the majority of writers, there are two conditions in typhoid when turpentine is of the utmost value. About the end of the second week, the tongue sometimes becomes very dry, red, chapped, perhaps coated in the centre with a brownish fur, and at the same time, marked meteorism develops. Ten drops of turpentine every two hours during the day, and every three hours during the night, will, in the majority of cases, remove these threatening symptoms. Again, when the diarrhœa persists after the acute stage of the fever is passed, when the convalescence is protracted, and, evidently, the ulcers of Peyer's patches are slow to heal, turpentine acts almost as a specific. It has been recommended in 5 to 10-drop doses, frequently repeated, for the intestinal hemorrhage, and in doses of f. 3j, every six hours, when there are tympanites and prostration, but no diarrhœa.

THE ANTISEPTIC TREATMENT OF TYPHOID.

Although this method of treatment cannot be considered as established, there has been enough testimony in its favor to commend it to careful consideration. In 1870, Dr. G. WILKES, of England, recommended the administration of *sulphurous acid*, *acidum sulphurosum*, in enteric fever; and in this country, Dr. J. WESLEY BOTKIN, of Illinois, (*Medical and Surgical Reporter*, May, 1876,) and others, have testified to its efficacy. The dose is from gtt. x-xx, every three or four hours, in lemonade. It can be combined with opium or digitalis. It should be continued until the tongue is perfectly clean.

In the *Practitioner*, September, 1873, Mr. STEPHEN SKINNER reports very excellent results from the *sulpho-carbolate of soda*. He generally commenced with gr. xx every four hours, gradually increasing, during the next four days, to the full dose of gr. xxx. A larger dose than this may produce cerebral symptoms. When the diarrhœa was excessive, opium was combined with the salt. He also claims that, by its early use, the premonitory symptoms of the disease are dispersed, and the threatened attack aborts.

THE SPECIFIC OR ANTI-PARASITIC TREATMENT OF TYPHOID FEVER.

This method of treatment, based upon the theory of the origin of the disease from germs, has found favor with some recent observers. The following is the plan pursued by

DR. J. C. WILSON, OF PHILADELPHIA,

as set forth in the *Med. and Surg. Reporter*, 1884. He orders the patient to bed, and gives fluid food in small amounts at intervals of 2 or 3 hours. At night he gives gr. vj-vij of calomel. This dose is repeated every second evening until three, or rarely four, doses have been given in the course of the first six or eight days.

It is given alone, or in connection with sodium bicarbonate. There is commonly a slight increase of diarrhœa, if it be present, without aggravation of other symptoms; and in some instances the tendency of the temperature at this time to steadily rise appears to be controlled. If no diarrhœa had been present, the calomel usually produces two or three large evacuations on the following day. If the case does not come under observation until after the tenth day, one only, or at most two, doses of calomel are given.

In only one case in sixteen was there observed mercurial fetor, and slight swelling of the gums. Excessive diarrhoea is controlled by opium.

The author lays down the invariable rule of maintaining a horizontal posture. From the beginning of the attack, one, two, or three drops of the following mixture are regularly administered in a wineglassful of ice-water, after food every two or three hours, during the day and night:—

954. R.	Carbolic acid,	gtt. ij
	Tincture of iodine,	gtt. iv.

Sponging with aromatic vinegar and tepid or cold water is advisable, because cleanly, grateful to the patient, and to a slight degree refrigerating. When the axillary temperature reaches 40° , quinine in 1.6 to 2. doses should be given on a falling temperature, and repeated in from forty-eight to seventy-two hours if necessary.

Alcohol is not recommended in the early part of the sickness, unless the patient be used to it. The average duration of the disease was twenty-eight days.

As salicylic acid is of all remedies known one of the most poisonous to bacteria, and as bismuth is indicated for the bowel affection, Dr. H. DESPLATS, of Lille, has prepared a *salicylate of bismuth*, and found that it may be given in large quantities with good results (*New Remedies*, 1883). The following working process will furnish a good product:

955. R.	Subnitrate of bismuth,	4 parts
	Salicylate of sodium,	7 parts.
	Nitric acid,	
	Water,	each a sufficient quantity.

Mix the subnitrate of bismuth, in a flask, with twenty-five parts of water, and gradually add nitric acid, constantly stirring, until the subnitrate is dissolved, avoiding an excess as much as possible. Dissolve the salicylate of sodium in two hundred and fifty parts of water. Then, while stirring the latter, pour the solution of nitrate of bismuth gradually into it. Collect the precipitate on a filter, wash it thoroughly, and dry it. Average yield: 3.75 parts, or about 94 per cent. of the subnitrate.

NOTES ON REMEDIES.

Acida. The mineral acids have been, at times, much prescribed in typhoid fever. Dr. GEORGE JOHNSON, of London, however, discountenances them, on the ground that they must irritate the ulcerated mucous membrane of the intestines, cause pain and griping, and increase the diarrhœa.

Acidi Phenicum is highly commended as an anti-febrifuge by M. VAN OYE, of France. He commences with a dose of seven and a half grains, *per rectum*, and progressively increases, until he reaches half a drachm, six times daily.

Alcohol. The value of alcoholic stimulants, in typhoid, has been much discussed. The following judicious rules are those laid down by Dr. ALFRED L. LOOMIS, of New York:

First. They should never be administered indiscriminately—that is, never give a patient stimulants simply because he has typhoid fever.

Second. When there is reasonable doubt as to the propriety of giving or withholding stimulants, it is safer to withhold them, at least until the signs which indicate their use become more marked.

Third. In every case—but especially when stimulants are not clearly indicated—watch, carefully, the effect of the first few doses. There are few whose experience in the treatment of typhoid fever is such as to enable them to positively determine, from the appearance of the patient, when the administration of stimulants should be commenced.

Should you commence the administration of stimulants, it is necessary to see your patient every two hours, and note carefully the effect produced. If you find the tongue becoming dry, the patient more restless, the delirium more active, the temperature ranging higher, and the pulse more and more rapid, you may be certain that stimulants are contra-indicated. If, on the other hand, the pulse becomes fuller and more regular, if the first sound of the heart is more distinctly heard, or, if having been absent, it has returned, if the restlessness and delirium are less marked, the tongue more moist, and the patient more intelligent, you may be certain that the time for the administration of stimulants has arrived. When you have commenced their use, it is of the greatest importance that you administer them at stated intervals, especially during the night.

In a severe case of typhoid fever, a free administration of stimulants, just at a critical period, (which may not last more than twenty-four hours,) will often be followed by a refreshing sleep, and your patient may rapidly pass from an apparently hopeless condition to one of convalescence.

Alumen may be employed to check the diarrhœa.

Ammonium Acetas and *Carbonas* are often exhibited, the former as a febrifuge, the latter as a stimulant. Prof. BARTHOLOW, however, considers that all salts of ammonium are distinctly contra-indicated by the excess of that element in the blood in this disease.

Caffea. Dr. PIGNE-DUPUYTREN, of Cal., speaks in the highest terms of coffee in this disease. (*Pacific Med. and Surg. Jour.*, April, 1879.) His rule is as follows: Give (to an adult) two or three tablespoonfuls of strong, black coffee (without chicory) every two hours, to be alternated with one or two spoonfuls of good claret or Burgundy wine. Make the patient take, in the course of the day, a few wineglasses of lemonade or citrate of magnesia; then sulphate of quinia. Dr. RADCLIFFE, of London, has used it for years in this disease as a stimulant, in the place of alcohol.

Camphora is frequently excellent in allaying the nervous agitation.

Chloral is often serviceable, in small doses, to induce sleep.

Digitalis. Dr. WUNDERLICH has recommended this drug to lower the temperature and slacken the pulse. Other German writers have also urged its claims, sometimes for its alleged antipyretic effects, sometimes as a roborant of the heart; but as yet no positively favorable opinion can be pronounced for its exhibition in this disease.

Ergota. This is of great value in the hemorrhage of typhoid. A common prescription in St. Thomas Hospital, London, is:

956. R.	Tincturæ opii,		
	Olei terebinthinæ,		
	Spts. chloroformi,	āā	℥x-xx
	Acidi tannici,		gr. x
	Mucilaginis et aquæ,		q. s. M.

For one dose.

When this is not retained by the stomach, the following may be tried:

957. R.	Acidi tannici,	gr. x	
	Extracti ergotæ fluidi,	ʒ ss-j.	M.

For one dose every hour.

Or, gr. v of ergotin may be used hypodermically. In 1883 Dr. DUBONÉ, of Pau, published a brochure in Paris claiming that sound fresh ergot (preferably in powder) would reduce the mortality of the severe type of typhoid to less than six per cent. The dose is gr. x-xx, thrice daily. Concentrated and abundant nourishment fills the remaining indication.

Eucalyptus Globulus, a teaspoonful of the tincture, well diluted, at regular intervals of three or four hours, is highly recommended by Dr. BENJAMIN BELL, of Edinburgh.

Iodinium. In St. Francis Hospital, London, iodine is given internally in all cases of typhoid fever. It is thought to lessen the troublesome gastric irritability so often present, and to diminish the number of evacuations from the bowels.

The following formula is usually employed:

958. R.	Iodinii,	3j	
	Potassii iodidi,	3ij	
	Aquæ dest.,	3x.	M.

Gtt. iij in a wineglassful of water every three hours.

LUGOL's solution, the liquor iodinii comp. of the Pharmacopœia, may be used in doses of six drops every three hours.

The cold bath, quinine and iodine, with close attention to the diet and the exhibition of stimulants, when indicated, constitute there the main features in the treatment of typhoid fever.

Hgdrargyrum. The old practice of giving small doses of calomel at the commencement of the disease has largely fallen into desuetude. It is, however, recommended on the high authority of the late Dr. PARKES. The gums should not be touched. Sometimes the diarrhœa will yield to gr. $\frac{1}{100}$ of corrosive sublimate every two or three hours.

Kairin has been tried as an antipyretic, but is condemned as dangerous in this disease by Dr. F. W. DRAPER, of Mass. (1884.)

Ænothera Biennis. Dr. J. F. SULLIVAN says of this remedy, in the *Pacific Medical and Surgical Journal*, January, 1878: "I am convinced that it essentially modifies the inflammatory condition which precedes ulceration of Peyer's patches, and that its use may frequently prevent ulceration. The usual dose in typhoid fever is from fifteen to thirty drops every three hours."

Opium is of great value to allay the nervous jactitation and insomnia. One-eighth of a grain of morphia in an ounce of camphor water, is an admirable draught. It may also be given in the diarrhœa. Nevertheless, it must be used cautiously. Dr. GEORGE JOHNSON has pointed out that although the direct effect of opium is soothing, yet that, when given in sufficient doses to render the intestines torpid, it acts indirectly as an irritant, by retaining the putrid morbid secretions until they decompose and give off noxious gases, which distend and irritate the bowel, and thus increase and prolong the diarrhœa.

Potassii Bromidum is one of the best agents to combat the restlessness.

Quinia. The use of quinia as an antipyretic and its administration has been discussed above. (P. 555.) Its value is far from proven. As a tonic in the latter stages, it is highly and justly esteemed.

Plumbi Acetas is sometimes useful in checking the diarrhœa.

Salicin and Salicylic Acid have been largely employed for their antipyretic and antizymotic properties. The general result, up to the present date, is favorable to their use. (See above.)

Serpentaria. Dr. BARTHOLOW considers that in typhoid fever, serpentaria is indicated, and is unquestionably serviceable when much depression exists. It is apt to increase the diarrhœa, however, and must be given with caution when the intestines are very irritable. The fluid extract is an appropriate form.

Terebinthinæ Oleum. (See above p. 558.)

Veratrum Viride has been lately urged by Dr. A. W. NELSON, of Connecticut, as of much value in typhoid as a cardiac sedative. His method of treatment is to administer the officinal tincture in doses of one to two drops every hour, from the commencement of the disease to convalescence. As *veratrum viride* is eliminated rather rapidly from the system, he estimates that these patients were usually under the influence of from three to twelve drops continuously. His conclusion is that: "A tendency of the typhoid ferment to exhaust itself at about fourteen days, the *veratrum viride* emphasizes, so that very many cases determine at twelve days, some at fourteen or fifteen, a smaller number at three weeks. Very few determine indefinitely, as do a large proportion in other treatments." (*Compend. Med. Sci.*, 1883.)

TYPHO-MALARIAL FEVER.

DR. ROBERTS BARTHOLOW, OF PHILADELPHIA,

has seen excellent results from small doses of FOWLER'S solution three times a day in typho-malarial fever. When there is much diarrhœa, a few drops of tincture of opium should be added to each dose of arsenic. In doses of half a drop to one drop of FOWLER'S solution, the tongue cleans, the skin becomes moist, and the delirium lessens, in a most remarkable manner, sometimes.

DR. JEFFERSON D. WILLIAMS.

This writer, practicing in Mississippi, relates an extensive experience in this disease. (*Virginia Medical Monthly*, March, 1877.) In its treatment, he condemns quinine, believing that it aggravates the symptoms. The most essential point he has found, is to control the diarrhœa, tympanites, and pain in the bowels. To this end, he administers *subnitrate of bismuth*, gr. xxx; *pulvis ipecac. et opii*, gr. x, every few hours; this failing, and the diarrhœa becoming excessive, he checks it by injections of opium, acetate of lead, subnitrate of bismuth, and glycerine. When the red and dry tongue, the tympanites and pain, denote active inflammation of the intestinal tract, he trusts to frequent doses of *chlorate of potash*, a teaspoonful of a saturated solution, every one, two, or three hours. Turpentine he has not found to be of value. Abundant, easily-digestible food and moderate stimulation, are required throughout the disease.

DR. ALONZO HARLOW, DETROIT, MICHIGAN.

In typho-malarial fever, this physician states (*Peninsular Medical Journal*, December, 1873,) that he has used, with much satisfaction, the following :

959. R. Quiniæ sulphatis,	gr. xxx	
Olei piperi nigri,	gtt. xxiv	
Acidi sulphurici aromatici,	ʒj	
Syrupi simplicis,	ʒiv	
Alcoholis,	ʒj.	M.

Tablespoonful every hour, until half the quantity has been used ; afterward, every other hour.

If the chill recur, repeat in same or less quantity ; milk-punch to be used freely. He used aconite, or veratrum viride, or gelsemium, for the fever, and did not believe in too general stimulation with alcoholic stimulants. At a later stage, and in some cases, diffusible stimulants, as ammonium carbonate and brandy, are necessary.

TYPHUS FEVER.

DR. JAMES TYSON, OF PHILADELPHIA.

The treatment is exceedingly simple. In the first place, we know that the greatest danger arises from asthenia, especially during the eruptive stage, which we might say is the most important period. To guard against this, we must use stimulants freely. If there is one disease in which the free use of alcohol is especially indicated, it is typhus fever. The quantity to be used should, of course, be governed by the condition of the patient. In some cases it may be necessary to give an ounce every hour. In addition to the use of alcohol, quinine is absolutely indispensable. Digitalis should also be given, both as an antipyretic and as a heart-strengthenener. When the temperature becomes high, sponging of the body is necessary. After the crisis, which, in this disease, is strikingly well marked, it is simply necessary to treat symptoms as they arise. Of the complications, one of the most frequent is bronchitis, which, of course, should be treated as any other bronchitis. Diarrhœa is an unusual symptom.

HENRY HARTSHORNE, M. D., PHILADELPHIA.

The treatment should commence with a mild laxative, as citrate of magnesia or rhubarb. Small doses of oil or enemata may be needed throughout the attack.

To combat the depression of the middle stage, the mineral acids have considerable reputation. Either dilute sulphuric, nitric, or nitro-muriatic acid, may be used :

960.	R.	Acidi nitrici,	gtt. xl	
		Aquæ,	f. $\frac{3}{4}$ iv.	M.

One or two tablespoonfuls every three hours.

961.	R.	Acidi nitro-muriatici,	f. $\frac{5}{8}$ ss	
		Spiritus ætheris nitrosi dulcis,	f. $\frac{5}{8}$ ss	
		Aquæ camphoræ,	f. $\frac{5}{8}$ vss.	M.

A tablespoonful every two or three hours.

Quinine acts well in 1 or 2-grain doses, four or five times daily, after defervescence has begun. Dr. H. is satisfied that the plan of treating typhus early with large doses of quinine is not only futile, but unsafe. The *diet*, at first, may be of gruel, toast water, etc.; but very soon, must milk and beef tea, or chicken or mutton broth (or an alternation of these) be given to support the strength. Before the first week is out, half the cases will need wine in moderation; some, brandy or whiskey. In the second and third week, more than half the cases will require steady support of a positive kind. In such cases, the proper routine is, a tablespoonful of brandy or whiskey punch, (one part of spirit to three, two, or one of milk,) every two, three, or four hours, and the alternate hours, a tablespoonful or two of beef essence or beef tea.

The great point of skill will be to determine when and how far to stimulate. Delirium favors the probability of its being needed, especially a low, muttering delirium. Of course a very feeble pulse indicates it. On trial, when the pulse grows slower, the skin more moist, and the restlessness or delirium is quieted, the stimulus has done good, and should be continued. If, on the contrary, a more hurried or a *harder* pulse follows, with heat of head and dryness of skin, and wilder delirium or deeper stupor, it should be stopped, for a while, at least; or, if given, be diminished in amount.

Hypostatic pneumonia, in typhus, cannot be treated actively. Even abstraction of blood by cups, is hardly ever to be ventured upon. *Dry*

cups, between the shoulders, and a blister upon the breast, will be as far as we can go.

PROF. M. CHARTERIS, M. D., GLASGOW.

This author approves of the plan pursued by the late Dr. J. HUGHES BENNETT, of giving an emetic in the early, incubatory stage of the disease, which, he believes, will often abort the attack. When the fever once sets in, he confines himself to treating symptoms. He gives a purgative at the outset, of *rhei pulveris*, gr. xxx-lx; subsequent constipation he relieves by enemata of tepid water. The head should be shaved, and cold lotions applied. Stimulants are called for when there is rapid, feeble pulse, and weakness of the first sound of the heart. If the debility is marked, the patient must be roused to take them. When there is much nervous irritability, with sleeplessness, a sedative, such as the following, is required :

962. R.	Potassii bromidi,	3 iij	
	Chloral hydratis,	3 i	
	Aquæ,	f. 3	viss. M.

A tablespoonful every two hours.

H. MCNAUGHTON JONES, M. D., DUBLIN.

This practitioner has derived great advantage, in typhus fever, from *digitalis*. (*Dublin Journal of Medical Science*, April, 1875.) His usual prescription is :

963. R.	Tincturæ digitalis,		
	Ætheris chlorici,	āā	gtt. xx. M.

In water, every three or four hours.

This has a well-marked effect in strengthening the pulse, lowering the temperature, preventing delirium, and improving the secretions. He believes cases to do better, under this drug, without alcohol in any form, and formulates the following conclusions :

1. That in the treatment of fever, typhus and other forms, too much reliance has been placed on alcoholic stimulants, and that fashion, rather than reason, has swayed many in their indiscriminate employment.

2. That the percentage of cases requiring such stimulants is a low one, and that while our administration of them, as regards quantity and kind, must depend entirely on the condition of the patient, still, the utmost caution is required.

3. That in *digitalis* we have a powerful cardiac stimulant, which, while it gives force to the heart, does not do so at the expense of the system, but rather is a conservative agent, which controls expenditure and limits waste of vital action.

4. That *digitalis* appears to be indicated in the early periods of many cases of typhus, in which we have a rapid pulse and high temperature range, regulating our administration by its effects on both, using it rather with the object of guiding the patient up to a certain point, than of curing the disease.

TREATMENT BY COLD BATHS.

The cold-bath treatment has been strongly urged by Professor MOSLER. He gives the following manner of applying it: The patient is placed, as far as the neck, in a water-bath at a temperature of 64° Fah., or of one or two degrees lower than this. When the temperature of the water is raised by the heat of the body, it is important to restore it to its original state—that is, to 64°. During the bath, the cold water should be poured over the head of the patient; this douche should not be warmer than the water in the bath, but sometimes even a little colder. This plan of treatment is to be carried out whenever the thermometer placed in the patient's axilla, marks 103° Fah. The duration of the bath must be regulated according to the nature of the sensations of the patient, and should be suspended whenever complaints are made of intense cold. The ordinary duration is from ten to thirty minutes; the patient is then to be dried with great care, and placed in a well-warmed bed. It is a good plan to administer to the patient, after the bath, some strong red wine.

Dr. SCHOLZ, of Bremen, has carried this plan out very successfully, in both typhus and typhoid, with some modifications. He uses a bath from 50° to 70°, and considered it indicated whenever the axillary temperature rose to 102°. (*Medicinische Rundschau*, 1872.) When immersed, the patient either moved himself or was lightly rubbed. The duration of the bath was from ten to fifteen minutes, or, in many cases, not more than five or six minutes. He was placed in bed without being dried, and was covered with a woollen coverlet. When the brain was much affected, cold douches were directed against the head whilst the patient was in the bath, and ice-caps were occasionally used. Cloths dipped in cold water were placed, at intervals of a quarter or

half an hour, upon the chest and belly. These appeared to exert a favorable influence upon the diarrhoea, and were felt to be agreeable to the patient. Scarcely any drugs were administered. The diet was light, but nutritive. In very severe cases, where there is no material augmentation of the animal heat, Dr. SCHOLZ does not think it right to use the baths, but brings the skin into action by powerful thermic and mechanical irritation (frictions). Dr. SCHOLZ distinguishes two periods in the treatment of the disease—in the first, or early period, the patient must not be bathed at night, and not more than five times during the day; whilst, in the second or later period, the number of baths may be sufficient to keep down the temperature of the body to near its normal degree.

NOTES ON REMEDIES.

Alcohol, in some form, is recommended by most authors. The indications for its use are given above.

Belladonna is recommended by Dr. GRAVES, when stupor with contraction of the pupil supervenes.

Digitalis has been highly praised by Dr. H. McNAUGHTON JONES. (F. 963.)

Erythroxylon Coca. From the ascertained fact that this drug diminishes the amount of urea secreted, Dr. SAMUEL McBEAN has recommended it in typhus and typhoid fever. (*British Medical Journal*, March, 1877.) He gives f. ʒj of the tincture every four hours, combined with a little ginger if it causes flatulence.

Moschus. Dr. MURCHISON considers that musk is a stimulant of very great value, which has fallen into unmerited neglect. It may be given in the following formula:

964. R.	Pulveris moschi,	gr. x	
	Mucilaginis acaciæ,		
	Syrupi aurantii,	āā	f. ʒij
	Aquæ camphoræ,		f. ʒss. M.

This amount at a dose, every five or six hours.

Opium. The use of opium in typhus varies greatly. Some authorities consider it always dangerous. Dr. MURCHISON discriminates between three forms or stages of the disease; when the delirium is active and the pulse of good strength, he would combine opium with a depressant and push it; *e. g.*:

965. R.	Liq. opii sedativi,	f. ʒj	
	Antim. tartratis,	gr. j	
	Aquæ camphoræ,	f. ʒvj.	M.

A tablespoonful every hour until sleep is induced.

When the pulse is feeble, and the delirium wandering, the opium should be combined with stimulants; *e. g.*:

966. R. Liq. opii sedativi, f. $\frac{3}{4}$ ss
 Spiritus ætheris, f. $\frac{3}{4}$ j
 Aquæ camphoræ, f. $\frac{3}{4}$ iiss. M.

Two tablespoonfuls every hour until sleep is induced.

Finally, when there is dyspnœa, a livid face, nervous congestion and a contraction of the pupil, opium is wholly contra-indicated.

Potassium Nitras may be given alone or in combination, to stimulate the secretions. Dr. MURCHISON recommends *nitre whey*, prepared by boiling $\frac{3}{4}$ j of nitre in Oj of milk, and straining.

Quiniæ Sulphas, is given by some physicians in large doses, gr. xx-xxx at the outset, to abort the fever and as an antipyretic. The effect of this treatment has not been generally satisfactory. The same may be said of the administration of an equal amount in the twenty-four hours, in divided doses. Small doses at regular intervals are more beneficial, as :

967. R. Quiniæ sulphatis, gr. $\frac{1}{2}$ -j
 Acidi sulphurici diluti, gtt. xx-xxx
 Ætheris sulphurici, gtt. xx
 Syrupi, f. $\frac{3}{4}$ j
 Aquam, ad f. $\frac{3}{4}$ j. M.

For one dose, every three or four hours.

Sulphuricum Acidum, and the other mineral acids, are largely used and favorably spoken of. (F. 967.)

VARIOLA—SMALL-POX.

W. H. BARLOW, M. D., MANCHESTER, ENGLAND.

Dr. BARLOW has no doubt that the exclusion of light exerts a retarding influence upon the progress of the pustules in their later stages, so that, instead of acuminating in the usual way, they shrink and shrivel about the sixth or seventh day of the eruption. But some of his cases of confluent small-pox have led him to believe that death may have resulted from the exclusion of light, the poison having, by the abstraction of the accustomed stimulus, been prevented from taking its usual course to the skin, just as patients die from the suffusion of the exanthem of scarlatina or measles. This view, which he supports by the history of a number of cases, leads him to the conclusion that the mode of treatment of small-pox by the exclusion of light, although very useful in the lighter cases of unmodified, and in all cases of modified small-pox, should be employed with caution in the graver cases of a confluent character. (*Lancet*, July 1, 1871.)

DR. W. R. CLARIDGE, OF PENN.

The use of *salicylic acid* as an abortive is highly lauded by this writer (*Med and Surg. Reporter*, June, 1883). The dose was gr. x, every three hours. In the case described, on the third day of its administration the pustules began to abort, their contents absorbed gradually, leaving a *pale, thin scab*, or scale, much resembling a minute fish-scale. These scales separated so rapidly that large quantities were removed each morning when changing the bed linen.

In two weeks the patient was convalescent, after which the remedy was continued one week longer in same dose three times a day.

DR. E. PRIDEAUX, ENGLAND.

This writer gives in the *Lancet*, 1877, the treatment of small-pox which he has employed with much success. On entering the small-pox hospital under his charge, the patients were ordered:

968. R. Acidi salicylici, gr. xx
Sodii bicarbonatis,
Ammonii carbonatis, āā gr. iv. M.

This amount in water every two or four hours, according to the severity of the case.

In the later stage of the disease, gr. v of citrate of iron and ammonia were added to this mixture.

In the first stage of the disease the patients were placed in baths of 90° and cooled down to 70°, they being allowed to remain in them about fifteen or twenty minutes. These baths were administered every day, and in severer cases twice daily. After each bath they were painted over with a mixture of glycerine, gelatine and carbolic acid, in the following proportion:

969. R. Acidi carbolic, f. ℥j
Glycerinæ, f. ℥vj
Gelatinae, f. ℥j
Aquæ, f. ℥xxvj. M.

For local use.

As soon as the pustules were full, those on the face were picked, if necessary, and the gelatine mixture painted on frequently. All the patients had at their own desire a pot of the mixture, with a brush, in order that they might paint their faces frequently. They said it relieved the pain, and entirely prevented any itching. It kept the skin

soft and moist, acting as an antiseptic, and preventing any unpleasant smell. In this stage of the disease each patient had a warm bath daily. Later on, soda in considerable quantity was put into each bath, and the patients well washed with carbolic soap, to aid the desquamation and so expedite their removal from the hospital.

The results were very gratifying, in that there was an almost entire absence of pitting, whilst the marks left even in the most severely confluent cases were such as would almost disappear after a short time; and the number of confluent cases was rendered much smaller by the eruption being checked both by the local and constitutional measures employed.

DR. W. HITCHMAN, ENGLAND,

states (*British Medical Journal*, Feb., 1879,) that he has attended hundreds of cases of small-pox without a single fatal issue, which he attributes solely to treating it with full doses of *belladonna* or *atropia*. He adds that he has suggested to several professors in France, Germany, Holland, and Italy, the value of *atropia* in small-pox, all of whom have communicated to him their high opinion of its therapeutic value. In fact, the medicine, when appropriately given, neutralizes the disease, and no morbid remedial augmentation, however temporary, is usually witnessed, unless the doses have been unnecessarily, and therefore injuriously large.

NOTES ON REMEDIES.

Ammonium Carbonas is indicated if there is manifest depression, especially during the suppuration period.

Belladonna. See above.

Camphor is advantageously employed, particularly in confluent or malignant small-pox, when the vital power is greatly depressed. It may be given alone or in combination with opium.

* *Carbolic Acid* has been administered with marked benefit. It is more useful at an early stage of the disease, although, given afterward, it much modifies the symptoms, and hastens the curative process.

Chloral is highly useful and necessary when there are high temperature, wakefulness and delirium.

970. R.	Acidi carbolici,	āā	f. ʒj-iss	
	Acidi acetici,			
	Tincturæ opii,			
	Spiritus chloroformi,	āā	f. ʒj	
	Aquam,		ad. f. ʒviiij.	M.

A tablespoonful every four hours, till the fever subsides. After the

first dose, the mixture is rather grateful to the patient than otherwise. SCHWIMMER recommends the local application of carbolic acid and thymol. He prescribes as follows :

971.	R.	Acid. carbol.,	℥j	
		Ol. oliv.,	℥viij	
		Cretæ prep.,	℥iss.	M.

Or:

972.	R.	Thymol,	℥j	
		Ol. leni,	℥viij	
		Cret. prep.,	℥iss.	M.

Mustard. Dr. E. S. LYNDON, of Athens, Ga., (*Medical and Surgical Reporter*) believes that a mustard plaster applied to any part of the body will bring out the eruption twenty-four to thirty-six hours earlier than usual, so that a diagnosis can be made on the first day of the fever.

**Opium* often fails to procure rest in the early stage of small-pox, but in the advanced stage of the disease, patients who are wakeful, but otherwise doing well, are benefited by it, given once or twice. But neither it, nor any other anodyne, is to be given when there is copious salivation and mucous expectoration, as the patient may die asphyxiated during sleep, from the secretion accumulated in the air-passages.

Quinine is useful, in small doses as a tonic, in full doses for the suppuration, and in antipyretic doses if the temperature is high.

Salicylic Acid is indicated from a theoretical standpoint.

Sodii Sulphis. The sulphites are recommended by Professor POLLI, of Milan, as mitigating the febrile action, rendering the confluent and malignant forms benignant, shortening the course of the milder forms, and in all cases accelerating convalescence.

Sulphuricum Acidum Dilutum. In confluent small-pox, treatment by dilute sulphuric acid, ℥xv-xx, with gr. j of quinine, every three or four hours, together with stimulants and nutritives, is an excellent one.

TO PREVENT PITTING.

Amylum. A thick mucilage of starch has been used as a local application to prevent pitting, with excellent effect. The entire surface of the body should be first sponged with tepid water.

Argenti Nitras. VELPEAU and other French writers recommended the practice of opening each vesicle on the third or fourth day of the eruption, and touching each with a stick of the nitrate scraped to a point. This, however, is a tedious and painful process, not unattended with danger. It has been recommended to wash the face on the fourth or fifth day with a strong solution of the nitrate.

Calcis Aqua. The application of cotton-wool, soaked in a mixture of equal parts of lime-water and olive oil, is said to be an efficacious one.

Carbolicum Acidum. Dr. H. YATES reports a case (*Lancet*, January 25th, 1868,) treated with success by the following ointment:

973. R. Acidi carbolici, ℥ ʒ ij
 Adipis, ʒ ij. M.

The ointment was colored with lamp-black, and applied thickly spread on wadding. It was changed every second day, the parts first washed with soap and warm water, and then with warm water impregnated with carbolic acid.

Glycerina. The following application is recommended by J. F. MARSON :

974. R. Glycerinæ, 1 part
 Aquæ rosæ, 2 parts. M.

Apply (after the pustules have discharged, and the discharge has begun to dry) freely for a few days, until the scabs begin to loosen.

Hydrargyri Unguentum. Inunction of mercurial ointment has been advised by Prof. BENNETT and others, but excessive salivation may follow its use. The following ointment is employed in the Children's Hospital, Paris :

975. R. Unguenti hydrargyri, ʒ xxv
 Cerae flavæ, ʒ x
 Picis nigræ, ʒ vj. M.

Iodinium. The tincture, applied once or twice a day, is a remedy which has been long used. Although it does not absolutely prevent pitting, it usually renders the marks and cicatrices less evident.

Zinci Carbonas. Prof. BENNETT, of Edinburgh, prefers the following to all other ointments :

976. R. Zinci carbonatis, ʒ iij
 Zinci oxidi, ʒ j
 Olei olivæ, q. s. M.

The same object is said to be obtained by treating the whole surface thickly with powdered calamine. This application has a very soothing effect.

EXTERNAL MEASURES.

Cold Baths. Dr. CURSHMANN, in ZIEMSEN'S *Cyclopædia*, maintains that baths are only useful in the premonitory and eruptive stages. But considering that the greatest danger in small-pox is during the suppuration fever, Dr. CLEMENT, of Lyons, thinks that it is to this period of the disease that treatment should be directed. The most noticeable effects produced by baths on small-pox patients are, a fall in the temperature, often sudden and of some degrees ; cessation of the delirium soon after the first bath ; and a great reduction of the pulse-rate and of the respirations. The first baths should always be of a temperature from 80° to 85° F. The effect of the bath must be carefully gauged by the thermometer. The patient should be submerged beneath the water up to his shoulders ; and when he is put to bed he should be enveloped in a sheet, and covered only with a simple woolen covering. Two or three baths may be given in the twenty-four hours.

YELLOW FEVER.

DR. DOMINGOS FREIRE, OF BRAZIL.

As he found in the dejections, the saliva and the blood of yellow fever patients, numbers of microscopic organisms, bacteria and cryptococci, he comes to the conclusion that the disease is of an infectious nature. And as the *salicylate of soda* has been much recommended in analogous affections, such as septicæmia, variola and typhoid fever, he determined to investigate its effects in yellow fever. He has used it, hypodermically, in doses of from $2\frac{1}{2}$ to 22 grains, according to the intensity of the fever, in the first period of the malady. In the second period, the dose should be smaller, as it then acts as an antiseptic; it is of no service in the third period, when black vomiting has commenced. He reports several cases with favorable issue, under this treatment.

DR. DE LACAILLE, OF RIO DE JANEIRO,

claims to have cured thirty-eight consecutive cases of yellow fever by the use of Déclat's preparations of *phenic* and *sulpho-phenic acids*, and in grave cases, the *phenate of ammonium*. In the early stages he gives the remedies by the mouth, but in the advanced stages the hypodermic method is necessary. He contrasts very favorably his recent experience with his former sad failures without these drugs.

GREENSVILLE DOWELL, M. D., GALVESTON, TEXAS.

This experienced physician directs that as soon as the first symptoms come on, the patient should be put to bed at once, and comfortably covered. If the stomach is full, it should be emptied by an emetic of ipecac, or mustard. If such is not the case, a moderate dose of castor oil or other mild laxative should be administered. Should inceptive chilliness be followed by perspiration, the patient should not be disturbed; but if the skin is dry and there is much thirst, warm teas of orange leaf or sage may be given.

Should there be vomiting at the outset, neither emetics nor warm drinks should be given, but instead of them, sinapisms should be applied to the epigastrium, and small pieces of ice be swallowed. Should these measures not check the vomiting, the patient should be cupped freely over the stomach.

When the fever is high and the pain in the head and back is great, he prescribes the following:

977. R. Hydrargyri chloridi mitis,
 Quiniae sulphatis,
 Pulveris opii et ipecacuanhæ, āā gr. xij. M.
 Make four powders. One every three hours.

This is to be continued with more or less frequency as long as the fever lasts.

If the fever continues high and the skin hot, it may be reduced by:

978. R. Tinct. aconiti fol., gtt. x
 Spts. æth. nitrici dulcis, ʒj. M.
 This amount in a wineglassful of water, every three or four hours.

In forty or fifty hours after the commencement of the disease, the fever subsides and the stage of calm comes on. If this is associated with much exhaustion, stimulants are demanded; if with marked restlessness, valerianate of zinc, gr. v-x, or morphia, gr. $\frac{1}{4}$, will be called for, repeated as necessary. When there is retching and vomiting, the following mixture may often be used with the best results:

979. R. Morphiæ sulphatis, gr. iv
 Creasoti, ʒj
 Spiritus vini gallici, f. ʒiv. M.
 A tablespoonful every three hours, as needed.

The stomach, at this stage, is generally swollen and sore, which should be met by mustard or blisters. If black vomit appears, the above brandy and creasote mixture should be alternated with tincture of chloride of iron, gtt. xv, or solution of the perchloride of iron or tannic acid. In the stage of calm, Dr. DOWELL never gives quinine. When there is suppression of urine, nitre or buchu is called for.

The patient should have light and nutritious food, regularly administered, and should be rigidly kept in bed, and prevented from any but the most necessary movements. Relapses are easily provoked, and are far more dangerous than the first attack. Embolism is a frequent cause of death, and must be guarded against by complete quiet.

DR. CHARLES BELOT, HAVANA.

This writer, Director of the Hospital for Yellow Fever at Havana, considers *local blood-letting*, at the outset of the disease, indispensable

in most cases. Cuppings are preferable to leeches. Eight cups on the nape of the neck, ten on the loins, and eight on the stomach, generally suffice to alleviate the symptoms; and if, four hours after the first application, there is not sensible amelioration, let it be repeated.

For the pains of the head and loins, sinapisms and mustard foot-baths are needed. *Tincture of aconite*, given in doses of six drops in twelve ounces of water, administered by spoonfuls every hour, should never be neglected in the first or congestive period.

If the disease has commenced with vomiting, if the tongue is charged with mucous deposits, let an emetic be given, composed of an infusion of thirty-six grains of ipecac. in four ounces of warm water. When this commences to produce its effect, the vomiting will be assisted with warm water, and continue until the ejections are clear as water.

If the tongue is not charged with mucous deposits, if the patient complains of great pain in the epigastric region, vomiting must be avoided, as its consequences would be grave.

About four hours after the vomiting, the patient will have less fever, the pulse will be less frequent and strong, the skin will be moist, or there will be general transpiration. It is necessary to take advantage of this moment, to administer a purgative of castor-oil or of sulphate of magnesia. If this remedy fails to act after two or three hours, its action should be promoted by injections of warm sea-water, or of sweetened water and olive-oil.

With free ventilation, quiet, and warm drinks to aid transpiration, this treatment will often disperse the alarming symptoms in three days. When, however, the disease continues its progress, the symptoms of the second period are presented.

The headache should be resisted with flying blisters on the nape of the neck, with compresses of brandy with camphor and belladonna on the forehead. To soothe the oppression of the stomach, apply compresses of brandy with camphor and belladonna, and give internally tincture of *nux vomica*, in very small doses. When, despite the compresses, the pain in the stomach persists, a flying blister is applied at this point. If nausea occurs, *bicarbonate of soda* alternated with *nux vomica* is the most useful remedy. If the disease advances and vomitings occur, they must be treated according to their character. If they are white, foamy, acid, holding in suspension flakes of mucous membrane, black vomit is not far off. It is often arrested by bicar-

bonate of soda in small doses repeated every half hour, alternated with carbonate of magnesia. If the vomitings are bilious, nux vomica and arsenical solution will be employed in preference, and will be continued if black vomit occurs. A little fresh water, slightly sugared, may be allowed, and thirst may be appeased with small pieces of ice.

Sometimes hemorrhages occur without black vomit. They constitute of themselves a grave condition. Limited hemorrhages, for example buccal and nasal hemorrhages, are combated with tonic applications, ice, and astringents, as dilute sulphuric acid, etc.

When hemorrhages have not taken place, if the symptoms invade the brain, cold water to the head, blisters to the nape of the neck, and calomel internally, are remedies to which recourse should be had. But this will be frequently useless, because the disease progresses with frightful rapidity.

The remittent and intermittent types begin with or without chills. This is a very important therapeutic indication. This first symptom (the chill) coming on perhaps after a meal, perhaps in the morning between two and three o'clock, an emetic, administered at once, disembarrasses the stomach of ingesta and saburral deposits, and determines a favorable transpiration. After the emetic, the patient should be allowed repose and sudorific drinks. If the symptoms call for an application of cupping to the nape of the neck, to the loins, or the abdomen, according to the indications of local congestion, it is rarely that twelve hours pass without the patient showing marked relief. If the congestive symptoms do not yield to the first cupping, they will be overcome on the second application, and by completing the effect of the emetic with a purgative. In most instances, after this treatment, the remission will be found to come, and this is the most important moment for the physician. This remission often occurs after the first twenty-four hours—in other instances, it is delayed two or three days. When, after the emetic and purgative, a general relaxation is observed, but insufficient to bring on transpiration, use should be made of diaphoretics, and preferably of *Dover's powder*, taken every hour in doses of a grain, until sweat is presented.

In this remission, *sulphate of quinine* must be administered in a dose of thirty-six grains, taken at once in a half-cup of black coffee without sugar. When the intermission is complete, its action is marvelous; the disease is immediately moderated. When the disease commences

with chills, followed by abundant sweats after the emetic and purgative, there is assurance that there will be another remission, and the quinine is in order.

In cases where sulphate of quinine cannot be employed, *calomel* is an excellent remedy, especially when, in the absence of remission, the tongue shows itself humid, loaded, white, large, the gums engorged, the stools difficult, or when there is bilious diarrhœa. Dr. B. administers from twenty-four to thirty grains, in two-grain doses every half hour, until the characteristic stools appear. Very often, after the administration of calomel, remission of the fever and of the congestive symptoms takes place; the skin becomes moist, and sulphate of quinine, the effect of which will be more sure in proportion to the distinctness of the remission, may then be appropriately used. Its effect is assisted by oil and by emollient injections. If there was no chill in the commencement, *aconite* and *tincture of digitalis* will be pressed. These are ordinarily sufficient to bring the patient into full convalescence. Dr. B. disapproves of combining calomel and quinine.

DR. F. PEYRE PORCHER, CHARLESTON, S. C.

This writer sums up the treatment of yellow fever as follows, (*Transactions of the South Carolina Medical Association*, 1872): 1. The treatment must begin *early*. 2. The first step should be a large mercurial purge, followed by a saline cathartic, thus emptying the bowels, and disgoring the liver and the glandular apparatus. 3. Simultaneously, and from the very beginning, use revulsives to the surface of the abdomen, hot stimulating baths to the lower extremities—with the assiduous and protracted application of ice-cold water to the head, hands and arms, *as long as there is a continuance of abnormal heat*. 4. All purgatives, all active depressing agents, must now be discontinued absolutely, and give place to a mild alkaline diuretic and diaphoretic, containing morphia. Cold sponging, and the use of sinapisms, and hot mustard pediluvia, are to be continued so long as headache, morbid heat of surface, with high thermometric range, indicate the existence or progress of combustion and destructive tissue metamorphosis—*fever*, in other words. Then, too, the recuperative powers of nature are to be trusted to, aided, if need be, by mild tonics, moderate stimulants, and nourishing diet.

EDWARD FOWLER, M. D., NEW YORK.

This writer (*Medical Record*, September, 1878,) from an experience of several severe epidemics in cities of the southern states, distrusts the line of treatment usually pursued, as laid down by most authors. He does not believe that the perspiration sought to be produced at all helps the case; mercury, he regards in all cases of true yellow fever as "hurtful to an extreme degree," and only valuable when the fever is mild and blended with an intermittent; he is wholly adverse to blisters; quinine, in the therapeutics of the true disease, has no place whatever.

Having thus discarded the traditional treatment, he adopted one which he states to have been much more satisfactory.

During the first hours of the attack, he recommends hot acid drinks in teaspoonfuls at a time, to allay thirst. Small quantities of broths at regular intervals will prove the best nutriment. For the costiveness which usually attends the disease, Seidlitz powders are unobjectionable. For the tympanitic condition of the abdomen, enemata containing carbonate of soda, gum *foetida*, and minute quantities of aloes, act not only promptly and delightfully, but often snatch the patient out of a state of great danger, and establish improvement.

For the prominent and distressing symptom of nausea, poultices to the epigastrium are demanded, and the following mixture, which should be given throughout the entire disease, beginning with the second day:

980. R. Potassii iodidi, gr. ij.
Liquoris potassii arsenitis, gtt. ij. M.

This amount for a dose every two or three hours, in any convenient vehicle.

The effects of this combination are uniform in relieving the tenderness of the stomach and keeping black vomit at bay.

Another important feature of Dr. F.'s treatment is to give a full dose of *opium* at the *commencement* of the disease, as much as morphine sulphas gr. i-ij. Of the use of this remedy he says: "It is a drug that depends for its remedial action on the stage of the fever in which it is given. Administered with the accession of the fever, and when the sufferer is writhing in universal pain, and given in sufficient quantity, it will not only relieve and give gentle rest, but clear up the troubled atmosphere of the brain, and roll back every darkening cloud from the mind, and leave no vestige of opium stupor or even drowsiness. But if given to check vomiting or diarrhoea, or restlessness after

the first twenty four hours, and then in small quantities even, it induces new symptoms, and endangers the patient."

NOTES ON REMEDIES.

Arsenicum. Dr. CHARLES BELOT says that, toward the end of the second period, when the vomiting cannot be arrested, when the patient has continual nausea, when the vomit contains bile or mucosities filled with blackish or sanguinolent streaks—in a word, when the characteristic signs of pronounced yellow fever are developed—there is no better remedy than arsenic. It is given by him as arsenious acid dissolved in water. It will be seen above, that Dr. FOWLER uses it throughout the disease.

Belladonna. According to Dr. BELOT, compresses of camphorated alcohol and belladonna, placed upon the epigastric region, diminish beating of the cœliac trunk, the epigastric pain, and the vomiting. Laid at the bottom of the abdomen, they quiet the colic pains and facilitate the passage of urine.

Camphora, combined with belladonna, finds its use, internally, in combating hiccough, and camphor alone is especially useful in the typhoid period of the disease.

Hydrargyrum Chloridum Mite. There is great discrepancy, not only as to the value, but even as to the safety of giving calomel in yellow fever. The safest plan is to confine its exhibition to those cases where obvious evidence of malarial poison co-exists, and to use moderate doses. For the various opinions, see above.

Ipecacuanha. As an emetic, at the outset of the disease, this drug is preferred by most physicians.

Nux Vomica has been found by Dr. BELOT to act very favorably on the epigastric distress.

Opium, in all forms and doses, is discarded by many. Dr. AITKIN especially warns against it when there is tendency to suppression of urine. For Dr. FOWLER's employment of it, see above. It will be noticed that he is directly at issue with Dr. DAGNINO, in the use of it for the latter symptoms.

Plumbi Acetas. For the gastritis, Dr. G. B. WOOD knows nothing better than the acetate of lead—1 or 2 grains every hour or two, and continued until from 30 to 40 grains have been taken; the astringent properties of this remedy may also render it useful in preventing the black vomit, which is now admitted to be a sort of hemorrhage.

Quinia Sulphas is given by Dr. M. DAGNINO (*Monograph on Yellow Fever*, New York, 1878,) in small and repeated, by Dr. BELOT in large and rare, and by Dr. DOWELL in medium doses. Various practitioners aim to establish cinchonism promptly, in all cases; others, and the more recent observers of the complaint, would confine the early exhibition of quinine to cases with obvious malarial symptoms, and employ the drug

later, merely for its tonic properties. In the continuous form of the fever, quinine certainly exerts little or no effect.

Potassii Chloras. Dr. H. R. Frost (*Charleston Medical Journal and Review*, vol. viii,) has found the chlorate of potash an excellent remedy in the febrile stage of the disease. After having freely evacuated the bowels, he gives the chlorate in the quantity of 3 or 4 drachms, in divided doses, during the twenty-four hours, employing, at the same time, when necessary, leeches to the temples and iced water to the head.

Potassii Iodidum is employed with marked success against the nausea, by Dr. FOWLER. (F. 980.)

Ricini Oleum is generally given the preference, as a laxative.

Salcylicum Acidum. Dr. HARTWIG BUNZ, of Savannah, states (*Berliner Klinische Wochenschrift*, September 2d, 1878.) that in the epidemic in that city, of 1876, finding that the fever was of an intermittent type, he resolved to make an experiment with salicylic acid. He gave adults a dose of $1\frac{1}{2}$ drachms, either in solution, in capsules, or rubbed up in sugar; and if the stomach rejected it, he gave a double dose per rectum. The result was excellent. Of one hundred and seventy-nine patients, only four died. He regards the acid as the most powerful antipyretic against yellow fever, both of the intermittent and remittent type, but has no experience of it in the continuous form.

Sulphuricum Acidum is advantageously used in small, dilute doses, in threatened black vomit.

Tannicum Acidum. Dr. BELOT recommends its employment where nitric acid reveals the commencement of albuminous deposit in the urine. Its use must be suspended if the albumen persists or increases. Tannin is administered every hour, in grain doses, in a spoonful of water. When the twelfth grain has been given, and it works no favorable change, it is replaced by arsenic, in his method.

Turpentine is indicated when there is cardiac weakness, depression of the vasomotor nervous system, and a dissolved state of the blood.

VII. TOXIC DISEASES.

Alcoholism (Inebriety; Delirium Tremens)—Arsenicism (Chronic Arsenical Poisoning)—Hydrargism (Chronic Mercurial Poisoning)—Opiumism (Opium Eating; the Opium Habit)—Plumbism (Chronic Lead Poisoning).

ALCOHOLISM (INTOXICATION; DELIRIUM TREMENS; CHRONIC ALCOHOLISM.)

The therapeutics of alcoholism embraces the treatment of this form of poisoning as it is presented in three distinct forms: 1. *Drunkenness*, including the gastritis, headache, etc., it leaves behind it. 2. *Acute Alcoholism*, known as delirium tremens, mania à potu, etc., and 3. *Chronic Alcoholism*, including the numerous and grave structural changes brought about by the prolonged consumption of alcoholic fluids.

1. DRUNKENNESS.

In cases of drunkenness or coma from alcoholic intoxication, the first indication is to empty the stomach. This may sometimes be accomplished by tickling the fauces with a feather; by the free administration of warm water, or mustard and water, if the patient can swallow; or, in some instances, by the hypodermic injection of apomorphia, or the internal use of brown emetine, gr. ij–iv, which acts very promptly. These failing, the stomach pump is the next resort.

Cold affusion, especially cold water, poured from a height of five or six feet upon the head, is a powerful revulsive, and may often advantageously be employed. Galvanism or electricity may also be resorted to. It is generally important to promote warmth, especially when the face is pale and the extremities cold. Sinapisms to the feet, bottles of hot water around the limbs, and mustard to the epigastric and cardiac regions, are then required. It is a great mistake to take a man who is partially intoxicated into the cold air under the impression that it will restore him. The change and sudden checking of the cutaneous transpiration are very apt to render him completely drunk.

When the stomach can retain fluids, the patient may take very

strong, hot coffee, without milk or sugar, in tablespoonful doses every five minutes. Or one of the following mixtures:

981. R. Ammonii acetatis, ℥ ij
 Syrupi aurantii florum, f. ℥ iss. M.
 To be taken in tablespoonful doses, in a small cup of tea, at quarter of an hour intervals.

982. R. Ammonii acetatis, ℥ ijss
 Sodii chloridi, ℥ j
 Syrupi, f. ℥ iss. M.
 To be given in a small cup of strong coffee in two doses at quarter of an hour interval.

The following is recommended by Dr. JAMIESON, of the Mercer Hospital, Dublin, as very efficacious in dispelling drunkenness:

983. R. Sesqui-carbonate of ammonia, ℥ j
 Table vinegar, f. ℥ j. M.
 This amount for one dose; to be repeated in half an hour if necessary.

This is a cheap extemporaneous plan of preparing spirits of mindererus.

The nausea, vomiting and irritable stomach consequent upon alcoholic excess, may be treated with iced milk diluted one-half, with Vichy, Apollinaris, or Saratoga water, or with lime water, a tablespoonful every five or ten minutes. A bladder of ice to the epigastrium is also efficient.

In the persistent retching and vomiting that sometimes follow a debauch, *liquor potassæ arsenitis*, in one-drop doses every hour or half hour, either alone or combined with a few grains of capsicum, is extremely efficacious.

For the nervous prostration, sense of anxiety and of impending misfortune, *nux vomica*, gtt. v-xv of the fluid extract, has been highly commended. This amount may be taken three or four times a day.

Dr. ALFRED L. LOOMIS, recommends the following in cases of "rum stomach," with gaseous distention of the stomach and bowels:

984. R. Tr. nucis vomicæ, gtt. v-xv
 Tr. gentianæ comp., āā
 Tr. columbæ comp., ℥ j. M.
 Take before meals.

The bowels are kept soluble by an aloetic and mercurial purge.

"Bitters" of different kinds are very popular as restoratives. *Quassia* is supposed to be peculiarly efficacious, but almost all the

vegetable bitters are employed. The following is a palatable combination :

985. R. Liquoris ammoniæ acetatis,
Tincturæ aurantii amari,
Syrupi aurantii amari, āā gtt. xx
Aquæ, f. ʒj. M.

For one dose ; to be repeated every half hour or hour.

For the headache and wakefulness which follow a debauch, Dr. A. McLANE HAMILTON prescribes *monobromide of camphor*, finding that it possesses peculiar hypnotic power.

986. R. Camphoræ monobrom., ʒj
 Confect. rose, q. s. M.
Make twelve pills. One or two as required.

Dr. A. A. SMITH, of New York, states that the first indication is to remove the alcohol from the intestinal canal. For this give of rhubarb and magnesia calcined, each a half drachm, and then give the following :

987. R. Spiritûs ammoniæ aromat, ʒij
 Tincturæ camph., ʒi ss
 Tincturæ hyoscyami, ʒij ss
 Spiritûs lavandulæ comp., q. s. ad ʒij. M.

ʒj every hour until the headache is relieved, and then give capsicum, gr. ij, and quinine, gr. iij. before each meal, for several days. If there be sleeplessness, give :

988. R. Sodii bromidi, ʒss
 Chloral hydrat., ʒij ss
 Syrupi aurantii cort., ʒss
 Aquæ, ʒij ss. M.

ʒss at night ; repeat in two hours if necessary to produce sleep. (*N. Y. Medical Record.*)

The following "quieting mixture" has been long used with the greatest success by Dr. F. PEYRE PORCHER, of Charleston, S. C., and is confidently recommended :

989. R. Chloral hydratis, ʒi ss
 Potassii bromidi, ʒij
 Spts. ætheris compos., f. ʒij
 Tincturæ valerianæ, f. ʒij
 Aquæ, f. ʒvj. M.

A tablespoonful every two, three or four hours in the nervousness following a debauch, commencing delirium tremens, excessive nervous excitement, etc.

Dr. J. H. NOWLIN, of Arkansas, has found the following of great value (*Louisville Medical News*, July, 1878:)

J. WARING CURRAN, M. D., DUBLIN.

993. R. Zinci oxidi, ℥j-ij
 Confectionis rosæ, q. s. M.
 For ten pills; one thrice daily.

When morphia or cannabis has done its duty in this disease, the after treatment by *oxide of zinc* is something to be observed rather than described; the constant dread, restlessness, and disturbed sleep, are quickly overcome by the bracing agency of the drug. (*London Lancet*, October 24, 1868.) In administering the oxide of zinc, care must be taken not to give it upon an empty stomach, as it produces nausea and a dislike for the medicine.

PROF. R. J. GRAVES, M. D., DUBLIN.

994. R. Antimonii et potassii tartratis, gr. iv
 Tincturæ opii, ℥. 3j
 Camphoræ, gr. xv
 Alcoholis, ℥. 3ss
 Aquæ destillatæ, ℥. 3viij. M.

Powder the camphor by the aid of the alcohol, add the water, pass through a piece of fine linen and then add the tartar emetic and the laudanum. Give a tablespoonful every two hours.

DR. LYONS, HARDWICKE HOSPITAL, DUBLIN.

995. R. Pulveris capsici, gr. xx-xxx
 Mellis rosæ, q. s. M.
 Make a bolus. For one dose.

This usually suffices to produce quietude and sleep. In exceptional instances, however, a second and even a third dose is required before full tranquillity is secured. The drug is well borne, and quiets the stomach in cases in which irritability and vomiting are present. Our author sums up (*British Medical Journal*, November 7, 1869,) his experience as follows: 1st. Capsicum is a valuable and reliable drug when opium fails, or is for any cause contra-indicated. 2. It is a safe drug for general employment in delirium tremens, and as such may be confidently recommended. 3d. It is not open to the objection which attaches to the continued use of opium, which, when it fails to tranquilize and produce sleep, adds to the state of excitement, and when pursued beyond a certain limit may induce opium coma. 4th. Capsicum has been employed in the delirium of fever when opium has failed to cause sleep, and with marked success in certain cases.

DR. GEORGE W. BALFOUR, EDINBURGH.

This writer (*Lancet*, February 1, 1879,) speaks in the most decided terms of the superior efficacy of *chloral hydrate*, in delirium tremens. He recommended it in all cases, from the lightest to the most severe. After a full trial of it for nine years, he pronounces it most satisfactory. He states, however, that there are very few cases, indeed, which yield to a less dose than fifty grains, and a considerable number which require a great deal more; those cases requiring the largest doses being those ushered in by the *status epilepticus*, which chloral arrests as rapidly and safely as it does delirium tremens itself. But, even in these cases, he has not found it necessary to give more than 120 grains of Liebreich's chloral, in divided doses; and this dose, though large, is not a dangerous one. From the irritated condition of the mucous lining of the stomach of a drunkard, it is probable that the absorption of ingested fluids is not so rapid as usual; it is but fair, therefore, for that reason, also, to allow a moderate interval between the doses, so as to avoid, as far as possible, any risk of giving more than enough. At the same time, we must shun the opposite extreme of giving doses in themselves too small to have any decided effect, and which have any possible cumulative effect destroyed by too long an interval being permitted to elapse between the giving of each dose. Acting upon these principles, he is in the habit of treating cases of delirium tremens by giving gr. xl chloral hydrate every hour, for three times, if necessary. Sometimes, but rarely, the first dose is enough; most commonly, two doses are required, and it is only in the very rarest instances that the third dose is necessary. If the attack be ushered in by the *status epilepticus*, he shortens the intervals between the doses to half an hour, as, in these cases, time is of the utmost importance, and a large dose is sure to be required. Should the heart be feeble, he gives each dose of chloral in half an ounce or an ounce of the infusion of digitalis. Under this treatment, the results are flattering. Unquestionably, fatal cases must occasionally occur under this, as well as under other modes of treatment, but the number of them are much decreased, because, from the rapidity with which a cure is brought about, many dangerous risks are averted. Thus we avoid all the risks arising from a long continuance of maniacal excitement, or from a suicidal state of mind, all risk from the exhaustion following persistent sleeplessness, or defective nutrition, the result of long-continued insufficiency of food, etc.

PROF. JOHN CURNOW, M. D., LONDON.

In the opinion of this teacher, (*Lancet*, February, 1878,) delirium tremens must be differently treated in the young and in the old. In first attacks in young subjects, complete abstention from alcohol, light and easily-assimilated food (milk diet), moderate purgation, and, occasionally, tartrated antimony, in doses of from $\frac{1}{8}$ to $\frac{1}{4}$ of a grain, very carefully watched, are required. If the patient is restless for two or three nights in succession, 30 grains of bromide of potassium, or 20 grains of chloral hydrate, may be given every four hours, for two or three doses; but, as the disease in young people is spontaneously curable, sedatives must not be pushed. One or more experienced attendants should be always present, but no form of mechanical restraint is permissible, for it always shows a lack of proper attendance.

In older cases, a mild purge should begin the treatment, and light but very nourishing food should be administered at short intervals. Milk, beef tea, raw eggs beaten up with milk, strong soup, and such articles, are to be given freely, and very small quantities of stimulants may be occasionally put into them, so as to coax the patient to take them, when, by careful management and good nursing, a very severe attack may be tided over, and natural sleep will return in from three to seven days. The early administration of sedatives is to be deprecated; but, should the restlessness persist, in spite of careful and assiduous feeding, a full dose of *laudanum* (30 to 40 minims), at bedtime, is of great value. In the absence of albuminuria, lung complications, or any sign of failure of the heart's action, this drug is preferable to other sedatives. If the opium alone fails, its combination with an alcoholic stimulant, (brandy, whiskey, and especially stout,) often succeeds. If there be any tendency to syncope, if pneumonia should come on, and in cases complicated with shock, as in surgical injuries, a free use of stimulants is imperative.

III. CHRONIC ALCOHOLISM.

PROF. WILLIAM A. HAMMOND, M. D., NEW YORK.

In the treatment of chronic alcoholism, the physician should peremptorily insist upon entire cessation from the use of alcoholic liquors. A mild purgative should commence the treatment, the bowels being always more or less deranged:

996. R. Pulv. aloës,
 Ext. fel. bovis exsic., āā gr. xv
 Resinæ podophylli, gr. ij. M.
 For five pills. One every alternate day.

For the special treatment of the condition, the *oxide of zinc* has been highly commended, and certainly possesses great power, (gr. ij–ijj, *ter die*.) But the bromides are yet more efficacious, especially the *bromide of zinc*:

997. R. Zinci bromidi, gr. ij
 Syrupi simplicis,
 Aquæ, āā f. ʒj. M.

This amount three or four times a day, gradually increased as rapidly as the stomach will permit, to two or three times the quantity.

Digitalis is an important adjunct to the treatment. It is the most active agent we possess as an eliminant of alcohol through the kidneys, and is also a heart tonic. It may be given in the infusion, f. ʒss thrice daily; or of the tincture gtt. xv–xxx, as often.

In the acute form of alcoholism, that which follows the sudden cessation of the wonted stimulus, the main indication is to procure sleep as soon as possible. No means in such cases is so effectual as the hypodermic injection of a full dose of morphia, gr. $\frac{1}{4}$ – $\frac{1}{2}$, as often as may be required, combined with the internal administration of brandy or whiskey in moderate quantities.

When however, the affection has come on *during* a debauch, these substances add fuel to the flames. In such cases, the bromides, in large doses, combined with digitalis, are the most effective remedies:

998. R. Potassii bromidi, ʒj
 Infusi digitalis, ʒss. M.

This amount every hour or two until the patient sleeps.

The monobromide of camphor, gr. iv in capsule, every hour, has been well spoken of.

With the medical treatment of either form of alcoholism, the strength should be supported by beef tea, and after convalescence, quinine, iron and strychnia will prove of service.

DR. T. D. CROTHERS, HARTFORD, CONN.

In the practice of this able specialist, the liquor is taken from the patient *at once*, and the following given:

999. R. Ammonii bromidi, gr. xl
 Tincturæ zingiberis, ʒj
 Aquæ, ʒj. M.

This, with the addition of tinct. cinchonæ, is often given every three hours.

The first and second night the following is used :

1000. R. Chloral hydratis, gr. xv
 Tinct. zingiberis, ʒj
 Aquæ or syrupi, ʒj. M.

If this acts well it is used until natural sleep is restored.

In some cases the following is prescribed :

1001. R. Spiritus ætheris sulphurici compositus, ʒj
 Syrupi cinchonæ, ʒj. M.

This is often a substitute for liquor, and if it does not disturb the stomach may be freely used. *Congress* and *mineral waters* are given freely.

1002. R. Ammonii bromidi, gr. xxx
 Infusi quassiae, ʒij. M.

Is often used with good results.

Phosphoric acid and *cinchona* form the basis of all the tonics given. The *iodide* and *bromide of potassium* are valuable, but are not certain, and often disagree with the stomach. Bathing and electricity are always good, although the latter is not well borne in many cases. Remedies that derange the stomach are avoided if possible. The first want of the disordered system is rest. Neuralgia and obstinate insomnia follow in nearly every case. Of the iron preparations, the *citrate* is the most *useful*, with *cinchona*; *quinine*, unless indicated by the presence of malaria, frequently causes cinchonism. No effort is made to cause sleep until night, then one of the above remedies is given. Fluid extr. lupulin is excellent with bromide of ammonium.

All *tinctures* and *alcoholic* extracts are carefully avoided after the first few days. Infusions and other preparations are used. Fruits of all kinds are recommended. Bathing and reclining, with some of the above remedies, will often suspend the depression complained of. As a gentle tonic, nothing is more useful, after the system recovers from the immediate effect of liquor, than :

1003. R. Potassii chloridi, gr. x
 Infusi calumbæ, ʒj. M.

Taken four times a day for a long time.

After the patient has been under treatment a few days the following cathartic is excellent:

1004. R. Hydrargyri pilulæ, gr. x.
With Seidlitz powder in the morning.

The treatment varies somewhat with each case. In some instances, no medicines are tolerated, and only external means, with exact hygienic rules, can be applied. In others, general alterative plans of treatment give good results. Narcotics are always transient in their effects, and more or less uncertain; and stimulants are frequently irritating and complicate the trouble. In delirium tremens, *constant nursing, nutritious food, chloral bathing* at night, with the *bromides, mineral waters*, etc., are the most effectual means. Occasionally, Dr. C. gives *milk-punch* in protracted delirium, but does not think it the most valuable remedy. A steam-vapor bath is equally as good. *Milk*, and *tea* and *coffee*, are given *ad libitum*. Beef tea, etc.

DR. N. KEELER MORTON,

physician to the New York Inebriate Asylum. This physician gives the following plan of treating the partial nausea, loss of appetite, nervous irritation, prostration and headache, which follow prolonged excessive alcoholic indulgence, symptoms to which the Germans apply the significant name, *Catzenjammer*. (*Medical and Surgical Reporter*, April, 1875.)

If, in such cases, we continue the exhibition of alcohol, (brandy and soda, etc.,) we temporarily alleviate the suffering, but are surely adding fuel to the fire. Experience has shown that, if we can substitute the stimulus of nourishing food for liquor, we can accomplish our end without running any risk to our patient. But, to do this, we must create an appetite for the food, and cause it to be retained and assimilated. For this purpose, Dr. MORTON universally pursues the following plan:

A double Seidlitz powder is administered in as large a quantity of water as possible, for the purpose of producing a rapid and thorough evacuation of the stomach and bowels. This is followed by:

1005. R.	Etheris chlorici,	gtt. xv	
	Glycerinæ,	f. ʒ ss	
	Tincturæ capsici,	gtt. iij	
	Syrupi zingiberis.		
	Tincturæ cinchonæ comp.,	āā f. ʒ ss	
	Aquæ,	f. ʒ iss.	M.

For one dose.

This is only given two or three times, once each morning, followed, during the day, by a teaspoonful of the following, every three hours:

1006. R. Lactopeptini, \bar{z} iij
 Acidi hydrochlorici, gtt. xxx
 Glycerinæ, f. \bar{z} ss
 Aquæ aurantii, f. \bar{z} iij. M.
 For a mixture.

The result of this treatment is almost always satisfactory.

DR. S. B. MARKEL, PENNSYLVANIA.

1007. R. Quiniæ valerianatis, gr. v
 Ferri sulphatis, gr. x
 Spiritus myristicæ, f. \bar{z} ss
 Aquæ menthæ piperitæ, f. \bar{z} iij. M.
 A teaspoonful whenever the craving for drink is felt. (Tilden's *Journal of Materia Medica*, May, 1877.)

The following formula, similar to the above, has been considerably employed for the same purpose:

1008. R. Ferri sulphatis, gr. v
 Magnesiæ, gr. x
 Aquæ menth. pip., \bar{z} xj
 Spir. myristicæ, \bar{z} j. M.

This amount twice daily, to appease the craving for liquor, is said to remove it entirely, and prevent the physical and mental prostration that accompany the absolute cessation.

Or the following:

1009. R. Pulv. calumbæ,
 Pulv. pruni virginianæ, āā \bar{z} iij
 Pulv. capsici, \bar{z} j. M.
 Mix well and sift. A teaspoonful in a glass of cold or hot water, when needed to allay the desire for stimulants.

1010. R. Tinct. capsici, f. \bar{z} iij
 Tinct. digitalis, f. \bar{z} j
 Aquam camphoræ, ad f. \bar{z} vj. M.
 A teaspoonful, as required.

1011. R. Tinct. capsici, f. \bar{z} ij
 Tinct. nucis vomicæ,
 Acidi nitro-hydrochlorici diluti, āā f. \bar{z} j
 Infusum gentianæ, ad f. \bar{z} xij. M.
 Two tablespoonfuls, as required.

Either of the above preparations is stated by writers in the *Lancet*, (June, 1877,) to be very useful in allaying the craving for alcoholic

stimulus, and removing the sense of sinking and faintness so much complained of by intemperate persons who have ceased indulgence in their wonted stimulus.

The *oxide of zinc* has also proved of much service in these cases. Dr. ROBERTS BARTHOLOW recommends that it be combined with pepper:

1012.	R.	Zinci oxidi,	ʒj	
		Piperinæ,	ʒj.	M.
For twenty pills. One three or four times a day.				

These, he says, relieve the trembling and the gastric catarrh, and diminish the appetite for stimuli. When commencing the administration of oxide of zinc, it should be given on a full stomach, otherwise it is apt to produce nausea.

MR. C. O. GROOM NAPIER, ENGLAND.

This gentleman pointed out, in a paper read before the British Association for the Advancement of Science, in 1875, that a vegetable diet materially decreases the desire for alcohol in those addicted to its use. Most people find that they can take wine with animal food, but not with farinaceous or amylaceous food. Hence, vegetarianism is a valuable aid in checking habits of inebriety.

Mr. NAPIER states that certain articles of vegetable diet are especially antagonistic to the taste for alcohol. They are: Macaroni, haricot beans, green dried peas and lentils, soaked for twenty-four hours and well boiled, onions, celery, rice, and highly glutinous bread. It is well known that nations who consume but little meat, subsisting principally on fruits and vegetables, are conspicuously temperate; and individuals who, for other reasons, adopt an exclusively vegetable diet, soon lose their desire for even an accustomed stimulant.

NOTES ON REMEDIES.

Alcohol. Most practitioners used to believe that in acute alcoholism in feeble subjects, moderate doses of alcohol are indicated. The latest authorities generally deem it needless.

Ammonii Bromidum has been employed in delirium tremens.

Ammonii Carbonas is valuable in inebriety. (See above.)

Ammonii et Potassii Tartras is much employed by German physicians, in this affection, but little used in America or England, excepting in combination with opium, when it often induces sleep after the failure of opium alone.

Arsenicum. Fowler's solution, in 1-drop doses, proves serviceable in the vomiting after a debauch, and in the morning vomiting of chronic alcoholism, either alone or combined with from 3 to 5 grains of capsicum.

Asarum Europæum. The leaves of this herbaceous plant enjoy a great repute in Russia as a remedy for the deranged state of health consequent on habits of intoxication. They are used in infusion, (℥ij-iv to aquæ Oj), or powder, (gr. iij, in pill.)

Atropia, hypodermically, is recommended by BARTHOLOW when there is obstinate insomnia with great restlessness, weakened action of the heart, coldness of the surface, clammy sweat, and a failure of nutrition with bromide of potassium, chloral and hypodermic injections of morphia.

Camphora is recommended by Dr. LAYCOCK in delirium tremens in persons of a nervous habit, where the exhaustion is great, and morphia inadmissible. The *monobromide* of camphor has been praised also.

Cannabis Indica is regarded by Dr. ANSTIE as preferable to opium when there is any reason, from the quality of the pulse, to believe the circulation much enfeebled. He prescribes gr. $\frac{1}{4}$ — $\frac{1}{2}$ of a good extract. TYRRELL records (*Medical Press*, March 13th, 1867,) a case of delirium tremens, in which, after three doses of ℥xx of the tincture of cannabis indica, every third hour, the benefit was marked after the failure of capsicum, and when opium was contra-indicated.

* *Capsicum*, in large doses, is strongly recommended by Dr. LYONS and others.

Chloral has been successfully combined with bromide of potassium in the indescribable nervousness, uneasiness and malaise which supervene in chronic alcoholism:

1013. R.	Chloral hydratis,	gr. x	
	Potassii bromidi,	℥j	
	Syrupi simplicis,	f. ℥j	
	Aquæ,	ad f. ℥j.	M.

This amount every hour.

Very much larger doses have been given by some physicians, (℥ij of chloral at a dose, Dr. F. BRADNACK, in the *Buffalo Medical and Surgical Journal*, September, 1877.) But the practice is undoubtedly dangerous. It has been shown beyond reasonable doubt, by Dr. MADISON MARSH, of Louisiana, and later by Dr. ERNEST MAGNAN of Paris, that drunkards do not bear chloral at all well. Its use by them even in moderate doses is liable to be followed by sudden death. The latter physician abjures it entirely in alcoholic cases. (See, however, p. 588.)

Cinchona, in strong infusion, is said to act very happily in inebriety and in the tremulousness which affects habitual drunkards.

Conium is recommended by Dr. HARLEY, in combination with opium, thus:

1014. R.	Succi conii,	f. ℥iv-vj-viij	
	Tincturæ opii,	℥xx-xxx.	M.

For one dose.

When the patient is old, and when there is much depression, the opium may be advantageously combined with carbonate of ammonia (gr. iij-iv), or quinine (gr. ij-iiij), or with camphor. Prof. STILLÉ says the best method of administering opium in delirium tremens, is to begin with gr. $\frac{1}{4}$, or its equivalent, and progressively augment the dose by small and hourly additions, until sleep is produced, or a sufficient degree of tranquillity obtained.

Phosphorus is a very useful remedy in the treatment of chronic alcoholism, according to Dr. D'ANCONA. (*Jour. d'Hygiène*, February 21st, 1878.) It is perfectly tolerated in doses which no one has dared to give heretofore—10 centi-grammes (nearly $1\frac{1}{2}$ grains) a day, for many weeks. He says the remedy gives to drinkers a feeling of comfort and strength, and furnishes the force necessary to carry on their organic functions, which they have been accustomed to get from alcoholic liquors.

* *Potassii Bromidum* is recommended as of conspicuous benefit, by removing delusions, calming the delirium, and procuring sleep, in the earlier stages of mania-à-potu, before the delirium has become furious. It is also of service in removing any delusions that may remain after the attack has been partially subdued.

* *Quiniæ Sulphas* is the nervine tonic in which Dr. ANSTIE has the most confidence, in grain doses, two or three times daily. It should be given from the first, if possible, being, if the stomach is very irritable, administered in effervescence with bicarbonate of potash and citric acid.

Sinapis. In that stage when a timely emetic will cut short the attack, none is so appropriate as mustard.

Strychnia. On account of its excellent powers as a nerve tonic, strychnia is often given to restore the tone of the system after alcoholic excesses. It is said to be an ingredient in several of the "bitters" used as "pick-me-ups," among *habitués* of saloons. Dr. H. C. MOREY relates a case of a drunkard who, after long habit, would take a number of grains at once, to restore his tone. (*Pacific Medical Journal*, 1875.)

Zinci Oxidum is a most valuable nervine sedative, in the tremor and exhaustion from alcoholic excess. (See above.)

ARSENICISM (CHRONIC ARSENICAL POISONING).

Chronic arsenical poisoning is probably often overlooked. It is produced by the dust thrown off by many kinds of wall papers, from the consumption of candies, etc., colored with arsenical pigments, among laborers in arsenical manufactures, etc. Varying degrees of fever attend the attack, the digestion is deranged, the appetite impaired,

bowels irritable, tongue white, furred and sore, mouth and throat dry and sore, and conjunctivitis frequent. Pain and weakness in the back and limbs, (arsenical paralysis) are not uncommon, as well as other nervous symptoms.

DR. GUENEAU DE MUSSY, PARIS.

It is well to commence the treatment with free bathing and a moderate purge, so as to eliminate any remains of the poison deposited on the skin or mucous membranes. This done, the *phosphide of zinc* should be administered in regular doses. It has been found to act as an efficient antidote in these cases, and the arsenical reaction, which is usually quite marked in the patients' urine, will, under the influence of the zinc, rapidly diminish. The phosphide may be given in pill form, combined or not with *nux vomica*. Its beneficial effects should be perceptible in a week's time. In some instances, the nervous symptoms disappear with astonishing rapidity under its use.

DR. A. S. TAYLOR, LONDON.

When arsenic has been taken into the system in a gradual manner no reliance can be placed on the chemical antidotes usually recommended. The treatment should begin with emetics, purgatives and baths, so as to remove as much of the poison from the surface as possible. ORFILA has recommended that diuretics be employed, in order to promote the secretion of urine, and thus favor the more speedy elimination of the poison from the system. In some cases the urinary secretion is deficient or suppressed, when proper measures are required to restore it.

OPIISM (OPIUM EATING; THE OPIUM HABIT).

SIR ROBERT CHRISTISON, EDINBURGH.

After considerable experience in the management of opium-eaters, this author doubts if success can be obtained by the gradual reduction of the amount consumed. He believes the drug can safely be left off abruptly, even after many years' indulgence. The nervous irritability which ensues he treats with full doses of bromide of potassium; and the insomnia with chloral. For the first three days after the cessation

of the habit, the patient suffers from great mental depression, nausea and vomiting; but by the fourth night these symptoms usually disappear; the patient obtains a refreshing sleep, after which the progress is satisfactory. The diarrhoea apt to occur he treats with suppositories of morphia.

He acknowledges, however, that there is great danger of a relapse after apparent cure.

DR. T. D. CROTHERS, HARTFORD, CONN.

This physician is accustomed to reduce the amount of opium gradually in some cases, rapidly in others. The more rapid the reduction, the more marked the insomnia, anæmia and nervous exhaustion; but if these can be combated successfully, the final recovery is more positive and permanent.

After the opium is discontinued, the following nerve-tonic is often used with good effect.

1016. R.	Tincturæ nucis vomicæ,	gtt. xij	
	Acidi phosphorici diluti,	gtt. xx	
	Syrupi pruni virginiani,	f. ʒss.	M.

This amount, at one dose, twice a day.

Another very excellent tonic for the nerve depression, always present in these cases, is:

1017. R.	Acidi phosphorici diluti,	gtt. xx	
	Tincturæ cinchonæ,	f. ʒ ss	
	Syrupi simplicis,	f. ʒss.	M.

This amount three or four times a day.

The following is also a favorite for the nerve depression:

1018. R.	Ammonii bromidi,	gr. xxx	
	Tincturæ cinchonæ,	f. ʒss	
	Syrupi,	f. ʒj.	M.

To be taken every three or four hours.

Also the following, which is varied from time to time, combining with it fluid ext. lupulin, or *valerian*, or some of the bitter tonics:

1019. R.	Chloral hydratis,	gr. xx	
	Syrupi cinchonæ,	f. ʒj.	M.

Taken every three hours.

The ferri et quiniæ citratis is given in many cases, but cinchona and

some of the vegetable tonics seem to work best. In addition to medicines, he orders a hot or tepid bath every day, enjoins out-door exercise, light, nutritious diet, and the galvanic current over the spine.

For the *insomnia*, he only gives remedies at night, after a warm bath, usually one of the following:

1020. R.	Chloral hydratis,	gr. xxv	
	Syrupi,	℥ss.	M.

To be repeated in two hours, if it does not act well.

If this does not work well, but is followed by nausea, or other disturbances, next day the following may be used:

1021. R.	Ammonii bromidi,	gr. xl	
	Tincturæ zingiber.,	℥j	
	Aquæ,	℥ij.	M.

This may also be repeated.

No other remedies in his experience seem more certain than these. *Electricity* frequently increases the insomnia. *Valerian*, *belladonna*, and *lupulin* occasionally work well, but are not alike in their effects, and derange the stomach. *Tea*, *coffee*, and *port wine* act well in many cases, taken just before bed-time. Rubbing and bathing and perfect quiet, in good surroundings, and the entire confidence of the patient, are essentials. The diarrhœa seldom requires particular attention; if it does, the following is usually sufficient:

1022. R.	Bismuthi subnitratis,	℥j.	
	Acidi tannici,	gr. x.	M.

For one powder, every three hours.

The excessive sweating will usually yield to acid mixtures, of which the dilute phosphoric acid is always the base. In both these conditions, the free use of subacid fruits, such as lemons, oranges, limes and grapes, is extremely advantageous.

Special attention must be given to the mental condition, as melancholia and brooding are prominent symptoms, and require to be met by light occupation, cheerful conversation, amusements, varied society, etc.

HYDRARGISM (CHRONIC MERCURIAL POISONING.)

PROF. WILLIAM A. HAMMOND, M. D.

The earlier symptoms of chronic mercurial poisoning are salivation with swollen gums, foetid breath, loosening teeth, and ulcerations. Mental and physical weakness, paralysis and convulsions, and caries and necrosis of the bones, are results which manifest themselves later.

The specific treatment, in all cases, is by the internal administration of *iodide of potassium*, which as MELSENS has shown, separates mercury from its combination with the tissues of the body, forming with it the iodide of mercury, which is eliminated with the urine. The iodide of potassium must be given in full doses, say gr. xxx, in divided doses, every day. The urine can be readily tested for the mercury eliminated under its influence, by placing a few drops, excreted during the second day, on a bright copper plate, and adding a drop of hydrochloric acid. A bright metallic stain will be found on the plate, if mercury is present; the iodide of mercury is decomposed, and the metallic mercury is precipitated.

Of course, while under treatment, the patient must not be subjected to continued poisoning by mercury, either medicinally or through his avocation.

Tonics, such as iron, quinine, and strychnia, are useful adjuncts. The prognosis, under favorable circumstances, is generally good, even in severe cases.

JAMES E. GARRETSON, M. D., D. D. S.

Mercurial ulitis, when confined alone to these parts, will generally disappear on the cessation of the drug. If this should not prove the case, nothing better can be done than to scarify and paint the gums with tincture of iodine; as a rule, the officinal tincture, diluted one-half. Chlorate of potash as a wash, and also used internally, may be employed; to an adult, gr. x in a tablespoonful of water, four or five times a day, is an appropriate dose. As a local application, ʒj to aquæ f. ʒj, is a very good strength. The bowels are to be kept in a lax condition by saline cathartics.

When the mercurial poisoning passes to that stage which results in the breaking down of the tissue, the treatment must be more active,

and directed quite as much to the constitution as to the mouth itself. Locally, the fœtor will urgently demand attention; and one of the best means to correct this will be found in the permanganate of potash in solution, gr. ij-x to aquæ f.℥j. Constitutionally vigorous tonic medicines are required. If hemorrhage intervenes, tincture of *Erigeron Canadensis*, gtt. j-ij, repeated occasionally until such bleeding is controlled. Locally, *cobweb* saturated in alum water may be used. Laid carefully upon the bleeding part, and retained in position, it will seldom be found to fail. Monsel's solution, etc., should not be used, owing to the great liability to sloughing.

Incisions may still be made, but few in number and cautiously, and the tincture of iodine applied should be diluted with two-thirds water. The dilute aromatic sulphuric acid is also a happy local remedy in such cases, say f.℥j to aquæ f.℥viij. A combination for local use, of beneficial effect, is:

1023. R.	Potassii chloratis,	℥ss
	Sodii biboratis,	
	Aluminis pulveris,	āā
	Potassii permanganatis,	℥ij
	Aquæ coloniae,	gr. xxv
	Tincturæ cinchonæ,	f.℥j
	Tincturæ myrrhæ,	f.℥ij
	Tincturæ capsici,	f.℥j
	Tincturæ kramerizæ,	f.℥j
	Aquæ,	f.℥viij. M.

For a mouth-wash.

If, in defiance of all that is done, the parts should slough, one of three things occurs—the disease will seem to have exhausted itself, and the parts, after a little rest, give evidence of recovery; or the bone follows the ulcerative action, and sloughs likewise; or the patient dies from irritation and exhaustion.

In the second case, it is important that the dead bone be left to detach itself, and not be removed by the surgeon.

Mercurial glossitis and ulceration of the tongue are also conditions frequently demanding treatment. In rapid swelling of the tongue, it may be necessary to leech the throat vigorously, to sweat the patient thoroughly, and follow this with a saline cathartic. These measures, combined with an arterial depressant, aconite, digitalis, or veratrum, will usually relieve from immediate danger. Great lassitude and depression are apt to follow.

Locally, sage tea, saturated with chlorate of potash; common table

tea; and the internal use of bromide of potassium, gr. xx, three or four times a day, are grateful measures.

For the mercurial ulcer, equal parts of bismuth, red bark and borax act sometimes very happily. Chlorate of potash, tannic acid and glycerine form another excellent combination. Solutions of sulphate of copper or zinc, gr. j to aquæ f. ʒj, are cleansing and stimulating applications.

PROF. JAMES SYME, SCOTLAND.

Persons whose constitutions have been injured by the use of mercury, frequently suffer from sores, either superficial and confined to the integuments, or deep-seated and affecting the periosteum or bones. In both cases the condition concerned is distinguished by characters which a practical eye readily recognizes, and proves very obstinate under ordinary treatment.

For these ulcers Prof. JAMES SYME recommends *blisters*, which shall not extend much beyond the ulcerated surface, in connection with small doses of *iodine*, gr. ij, three times a day, in simple solution. This simple and powerful mode of treatment has, in his hands, completely restored limbs which had been condemned to amputation by other surgeons, for ulcerations and disease of the bones. (*Surgical Works*, 1866, p. 916.)

Dr. JUKES STYRAP, physician extraordinary to the Salop Infirmary, has found the following prescription exceedingly efficacious in controlling ptyalism—in fact, a positive “specific” for it:

1024. R.	Sulphur præcip.,	ʒij-iv	
	Potassii chloratis,	ʒij-ʒj	
	Liq. morphiæ,	f. ʒj-iss	
	Mist. amygdalæ,	f. ʒ viij.	
Mix well.	Two tablespoonfuls every three or four hours. Shake well.		

Especial care must be taken to diminish the quantity if any relaxation of the bowels supervene. In thirty-six hours the secretion and tenderness of the gums become very sensibly diminished.

PROF. S. D. GROSS, PHILADELPHIA.

1025. R.	Liquoris plumbi subacetatis,	f. ʒj	
	Aquæ,	f. ʒ viij.	M.
To be used as a mouth-wash every hour or two, in cases of <i>mercurial stomatitis</i> .			

The only objection to this lotion is, that it discolours the teeth, which effect, however, quickly disappears. At the same time, internally, the chlorate of potassa should be administered. Fifteen to thirty grains are to be taken *ter die*, in mucilage or lemonade.

Dr. FAHNESTOCK, of Pennsylvania, has stated that an infusion of the inner bark of the root of the sumac, *Rhus glabrum*, is almost a specific for the sore mouth attending inordinate mercurial salivation.

The hypodermic injection of the *sulphate of atropia* is a powerful agent in arresting mercurial salivation. Dr. EBSTEIN, of Breslau, arrested, by this means, in seven minutes, a case which had continued a month; on doubling the dose, the secretion was arrested for twelve hours. This action of the drug is explained through its inhibitory influence on the permanent irritation of the secretory fibres of the salivary glands.

Pilocarpin has been found useful to relieve the system of the drug.

PLUMBISM—LEAD POISONING.

PROF. WILLIAM A. HAMMOND, M. D.

In the numerous forms of chronic lead poisoning—whether this assumes the character of lead colic, paralysis, encephalopathy, hyperæsthesia, or anæsthesia—the specific treatment consists in the administration of the *iodide of potassium* in full doses. Although some claim that the resulting compound—the iodide of lead—is very poisonous Dr. H. has never seen the least untoward result, and he has always used the potassic salt in large doses from the very beginning.

In the treatment of lead encephalopathy, the free administration of the iodide of potassium, combined with the bromide of potassium, affords the best prospect of success.

In lead colic, the hypodermic injection of morphia, in doses sufficient to keep the pain in check while the iodide of potassium is doing its work, with an occasional purgative, will generally be all the treatment required.

But, in lead paralysis, in addition to the iodide, we must use direct measures. Chief among those is electricity. The induced current, if

it will cause the muscles to contract, is to be preferred. Each paralyzed muscle must be acted on for two or three minutes every alternate day. In ordinary cases, two months will suffice to effect a cure.

When the electric contractility of the muscles is completely abolished, the primary interrupted current must be used until contractility is established. Frictions, kneading of the muscles and passive exercise are also useful.

PROFESSOR WILLIAM PEPPER, M. D., PHILADELPHIA.

This practitioner remarks that the treatment of lead colic should have several points in view. First, the removal of lead from the intestinal canal; second, its elimination from the tissues; and third, the treatment of special manifestations.

For the removal of lead from the intestinal canal he prefers the *sulphate of magnesium* with *dilute sulphuric acid*. This converts it into an insoluble salt, and acts also as a purgative. Internally, to eliminate the lead from the tissues, he gives the iodide of potassium in full doses. This forms the iodide of lead, which is excreted by the skin and kidneys. With regard to the treatment of special symptoms, if there is severe colicky pain it must be relieved by anodynes or anæsthetics. The following is a preparation which acts well:

1026. R.	Ext. colocynth. comp.,	gr. ij	
	Ext. opii,	gr. ss	
	Ext. belladonnæ,	gr. $\frac{1}{2}$.	M.

This pill repeated until the pain is relieved and the bowels opened.

The palsy may be relieved by the internal administration of *strychnia*, and locally by the application of the faradic current to the affected muscles.

The anæmia is best treated by the iodide of potassium and the salts of iron.

PARIS HOSPITALS.

Sulphuric acid is much employed in the Paris hospitals in the treatment of saturnine poisoning, with the most marked success. Mild cases yield in about three days, severe ones in six or seven days.

1027. R.	Acidi sulphurici diluti,	f. ʒij	
	Aquæ,	Oij.	M.

This quantity is given during the day. The stomach soon becomes

accustomed to it, even when the first dose or two is rejected. The abdominal pains are relieved after the first or second day, the constipation yielding soon after. This treatment should be combined with the repeated use of sulphur baths.

DR. LUTZ, PARIS.

1028. R. Sulphuris loti,
Mellis despumati, āā ʒiv. M.

Give an ounce and a half of this electuary three consecutive days in lead colic, then give in doses gradually lessened. On the third day the acute pain has disappeared.

1029. R. Chloroformi, f. ʒss
Alcoholis diluti, f. ʒss
Mix and add,
Aquæ, f. ʒ viij. M.

To be given as an enema in lead and nervous colics.

1030. R. Podophyllin,
Extracti nucis vomicæ,
Extracti belladonnæ, āā gr. iij. M.

Divide into ten pills. One twice or thrice a day, to relieve the painful constipation of workers in lead. At the same time, sulphur baths are to be administered.

PROFESSOR NIEMEYER.

1031. R. Olei ricini, f. ʒij
Olei tiglii, gtt. iij. M.

A tablespoonful every two hours in lead colic. The patient also to take in the twenty-four hours three opium pills of from gr. $\frac{1}{4}$ – $\frac{3}{4}$. Warm baths, narcotic fomentations, enemata, sometimes purgative, sometimes narcotic.

GEORGE HAY, M. D., PHILADELPHIA.

This writer (*Medical Times*, 1878,) objects to the exhibition of potassium iodide in cases of chronic lead poisoning on the ground that the iodide of lead is more insoluble than the chloride of the same metal. Preferably he would exhibit *sodium chloride* in drachm doses, three times a day, because, as he says, the kidneys eliminate the chloride of lead freely, or an oxide of lead dissolved in a solution of a chloride. In addition to this a tepid bath ought to be employed daily, for three reasons, viz., to keep the skin active, to remove whatever lead has been excreted in that way during twenty-four hours, and to prevent its re-absorption. The use of a soft flesh-brush would be a great assistance to the bath by removing mechanically from the skin any lead, in whatever form, adhering to its surface. Sponging of the body might be

substituted if the patient is too weak to bear the fatigue of bathing. During the course of recovery from chronic lead poisoning, to improve the nutrition and build up the strength of the patient, analeptics, nerve tonics, and nutrients are indicated, such as *tinctura ferri chloridi*, *liquor strychniæ* (not the sulphate), or *tinctura nucis vomicæ*, *tinctura cinchonæ*, or the alkaloid quinia (not its sulphate), together with animal broths, milk, eggs, etc.

JOHN SYER BRISTOWE, M. D.

This author advises, in the treatment of lead colic, that the pain and discomfort be relieved by opiates and fomentations, and that no purgatives whatever be administered, but the bowels be left to act of themselves, which they usually do in a few days. At most, copious enemata of warm water or warm gruel may be given. The recommendation of Dr. PEREIRA was baths medicated with potassium sulphide.

1032.	R.	Potassii sulphidi,	℥ij	
		Aquæ,	Cxv.	M.

The theory is that the lead escaping from the surface of the skin is converted into an insoluble sulphide. Dr. BRISTOWE, however, doubts whether the baths promote the escape of the lead in any important degree.

For the restoration of the paralyzed and wasting muscles, galvanism is the only effectual remedy. The most effectual is the slowly interrupted constant current, used three times a week, ten or fifteen minutes at each sitting. Each muscle should be separately galvanized.



INDICES.

I. INDEX OF AUTHORS.

- Abernethy, John, England, 424.
 Adams, J. F. A., United States, 302.
 Adams, Samuel S., United States, 465.
 Ainstie, F. E., England, 99, 113, 137, 250.
 Aitken, Prof. Wm., Scotland, 40, 123, 172, 350, 400, 405, 433.
 Albertozzi, Sante, Italy, 257.
 Allbutt, T. Clifford, England, 276.
 Allen, Prof. Harrison, United States, 213.
 Althaus, Dr., England, 123.
 Anderson, L. B., United States, 162.
 Anderson, McCall, Scotland, 225.
 Anderson, William, United States, 434.
 Andrews, F. L., England, 178.
 Andrews, Prof. G. P. United States, 171.
 Angrisani, Dr., Italy, 301.
 Armangaud, Dr., France, 133.
 Armor, Professor, United States, 345.
 Ashhurst, Samuel, United States, 521.
 Ashwell, S., England, 484.
 Atkinson, W. B., United States, 17.
 Atlee, W., United States, 469.
 Aufrecht, Dr., Germany, 472.
 Ayre, Joseph, England, 327.
 Babieff, Dr., 513.
 Bailey, P., United States, 521.
 Balfour, G. W., Scotland, 282, 451.
 Bamsler, Professor, Germany, 261.
 Bard, J. W., United States, 193.
 Barlow, G. H., England, 35.
 Barlow, W. H., England, 182, 398, 426, 452, 570.
 Barry, Dr., India, 326.
 Bartholow, Roberts, United States, 29, 101, 114, 157, 169, 188, 196, 217, 286, 300, 335, 346, 392, 412, 420, 433, 476, 479, 507, 530.
 Bartlett, Dr., United States, 208.
 Baruch, S., United States, 190.
 Basham, Dr., England, 452.
 Baskell, N. M., United States, 436.
 Bassi, Dr., Italy, 527.
 Bastings, Dr., Belgium, 255.
 Bates, W., England, 328.
 Battley, Dr., England, 281.
 Beale, Lionel S., England, 477.
 Beard, George M., United States, 214.
 Beardsley, Dr., Canada, 508.
 Begbie, J. W., Scotland, 42.
 Beigel, Dr., England, 173.
 Bell, Benjamin, Scotland, 396.
 Bell, James E., United States, 219.
 Bell, O. D., United States, 435.
 Belot, Dr., France, 582.
 Belot, Charles, Cuba, 576.
 Belvounsoff, Dr., Russia, 528.
 Bemiss, Professor S. M., United States, 503.
 Benate, Dr., France, 148.
 Benedikt, M., Germany, 40.
 Bennett, Professor, Scotland, 500, 574.
 Berkhart, Dr., Germany, 201.
 Bernatzek, Professor, 516.
 Bernheim, H., France, 299.
 Berry, J. C., United States, 384.
 Betz, Dr., Germany, 368.
 Biddle, John B., United States, 481.
 Birch, S. S., England, 337.
 Bird, Golding, England, 380.
 Blache, Dr., France, 41.
 Blacklock, A., India, 329.
 Blake, John G., United States, 202.
 Bonamy, Dr., France, 357.
 Bond, S. T., United States, 416.
 Bonfigli, Professor, Italy, 357.
 Bonjean, Dr., France, 26.
 Boteler, W. C., United States, 539.
 Both, C., United States, 238.
 Botken, J. Wesley, United States, 559.
 Bowditch, Henry M., United States, 300.
 Bouchardat, A., France, 444, 447.

- Bouchut, Professor, France, 35, 301, 307, 409.
 Boyer, Julius, France, 238.
 Bramwell, Byrom, England, 484.
 Brand, Dr., Austria, 186.
 Bremond, Dr., France, 551.
 Brickwell, Henry, England, 408.
 Brinton, William, England, 376.
 Brisbane, Dr., England, 62.
 Bristowe, John Syer, England, 164, 220, 256, 373, 387.
 Brookes, Dr., Germany, 527.
 Brooks, J. G., United States, 404.
 Brown, C., Scotland, 130.
 Brown, F. J., England, 137.
 Brown-Séquard, C. E., France, 47, 57, 100.
 Browne, C. England, 57.
 Browne, Lennox, England, 193, 221, 223.
 Bruen, E. T., United States, 307.
 Brugelmann, Dr., Germany, 464.
 Bruns, Dr., Germany, 141.
 Brunton, T. Lauder, England, 283, 433.
 Buckler, Dr., United States, 314.
 Bucquoy, Dr., France, 267.
 Bufalini, Professor, Italy, 317.
 Bunz, H., United States, 582.
 Burkart, Dr., Germany, 352.
 Burroughs, Samuel R., United States, 462.
 Butler, R. J., England, 546.
 Byrd, Harvey L., United States, 70, 513.
 Calderwood, G., England, 412.
 Caro, S., United States, 408.
 Carpani, Dr., France, 541.
 Carre, Marius, France, 207, 405.
 Carstens, J. H., United States, 371.
 Carter, Charles, United States, 123.
 Cauldwell, C. M., United States, 452.
 Celsus, 60.
 Cervello, Vibenzo, Italy, 96.
 Chambers, Thomas King, England, 36, 200, 275, 481, 539.
 Chandler, A. H., New Brunswick, 535.
 Chandler, S. T., United States, 318.
 Chapman, Professor N., United States, 372.
 Chapman, J., England, 56.
 Chapman, E. N., United States, 461.
 Chaoteris, Professor M., Scotland, 164, 186, 221, 377, 476, 567.
 Chateaubourg, Dr., France, 431.
 Chiarmelli, Professor, Italy, 252.
 Chipperfield, Dr., India, 372.
 Christison, Robert, Scotland, 598.
 Churchill, J. F., France, 236.
 Civile, Dr., France, 466.
 Claiborne, J. H., United States, 512.
 Claridge, W. R., United States, 571.
 Clark, Thomas, England, 179.
 Clarke, Professor Alonzo, United States, 354, 551.
 Cleland, J., Scotland, 343.
 Clemens, Dr., Germany, 59.
 Clement, Dr., France, 574.
 Clouston, T. S., Scotland, 51.
 Clubbe, C. P. B., 445.
 Clymer, Meredith, United States, 199.
 Cobbold, T. Spencer, 413.
 Cohen, J. S., United States, 147, 187, 260.
 Coghill, J. G. S., England, 238.
 Comegys, C. G., United States, 125.
 Concato, Professor, Italy, 257.
 Condie, C. F., United States, 411.
 Connolly, Dr., England, 79.
 Cook, J. L., United States, 163.
 Copland, J., England, 31, 61, 63, 159, 210, 266, 333, 346, 461.
 Corrigan, D., Ireland, 545.
 Corson, Hiram, United States, 478.
 Corson, J. W., United States, 149.
 Craig, W., Scotland, 340.
 Crandall, J. B., United States, 355.
 Cressler, Dr., Spain, 407.
 Crichton, Dr., England, 343.
 Cross, J., England, 490.
 Crothers, T. D., United States, 590, 599.
 Curci, Dr., Italy, 158.
 Curnow, John, England, 589.
 Curran, J. Waring, Ireland, 56, 244.
 Cushing, J. R., United States, 367.
 Dagnino, Dr., Italy, 581.
 Da Costa, J. M., United States, 36, 58, 103, 126, 138, 143, 146, 150, 165, 174, 197, 208, 235, 245, 257, 285, 290, 300, 302, 335, 350, 392, 402, 434, 478.
 Darmecy, Guyot, France, 454.
 Date, William, England, 414.
 Davies, H., England, 546.
 Davis, F. H., United States, 181.
 Davis, J. W., United States, 119.
 Davis, N. S., United States, 103, 131, 262, 331, 362, 456, 486, 530.
 Day, John, England, 453.
 Day, William Henry, England, 75.
 Debove, Dr., France, 395.
 Declat, M., France, 521.
 Delafield, F., United States, 548.
 Delieux, Dr., France, 145, 179.
 Denton, Ashley N., United States, 138.
 Deschamps, Dr., France, 560.
 De-plats, H., France, 560.
 Dewar, J., Ireland, 217.
 De Willebrand, Felix, Finland, 512.

- Dexter, Ransom, United States, 42.
 Dick, Forbes, England, 406.
 Dickinson, W. H., England, 474.
 Dickinson, W. N., England, 434.
 Dobell, Horace, England, 184, 209, 216, 277.
 Doboué, Dr., France, 259, 389.
 Docker, Wm., England, 369.
 Donaldson, R., British India, 364.
 Donkin, A. S., Ireland, 451.
 Dowell, T. G., United States, 490, 506, 575.
 Draper, J. R., United States, 553.
 Draper, F. W., United States, 563.
 Dresch, Dr., Germany, 45.
 Drinkhard, Dr., United States, 124.
 Druitt, Dr., England, 249.
 Duchenne, Dr., France, 466.
 Drysdale, Dr., United States, 429.
 Dujardin-Beaumetz, Professor, France, 155, 520.
 Duncan, J. F., England, 499.
 Dunglison, Professor Robley, United States, 341.
 Durand, Dr., France, 315.
 Durant, C. M., England, 278.
 Earnest, John G., United States, 360.
 Easton, Professor, Scotland, 482.
 Echeverria, M. G., United States, 64.
 Eickholt, G., Germany, 90.
 Elliotson, Dr., England, 44, 249.
 Ellis, Calvin, England, 431.
 Emerson, N. B., United States, 136.
 Engel, Professor Hugo, United States, 22, 417.
 Erb, W. H., Germany, 133.
 Erlenmeyer, Dr., Germany, 51.
 Eschmarch, Professor, Germany, 213.
 Eulenberg, Dr., Germany, 517.
 Evers, Dr., England, 518.
 Eyre, Sir J., England, 381.
 Farquharson, Robert, England, 404.
 Farre, F. J., England, 538.
 Fauconnet, Dr., France, 282.
 Faulkner, R. B., United States, 152.
 Favier, Dr., France, 136.
 Fayrer, J., British India, 350, 491.
 Fenwick, Dr., England, 394, 468.
 Fereol, M., France, 108.
 Feris, Brazile, France, 201.
 Ferrer, Dr., Spain, 417.
 Ferrier, Dr., London, 185.
 Finchaw, Dr., England, 537.
 Finn, Dr., Canada, 545.
 Fleischmann, A., England, 547.
 Fleming, Dr., England, 321, 538.
 Flint, A., United States, 42, 71, 142, 226, 260, 294, 358, 474.
 Fonssagrives, Professor, France, 312, 466.
 Forrest, W. E., United States, 525.
 Foster, Balthazar, 204, 236.
 Foster, N. S., England, 31.
 Fothergill, J. Milner, England, 177, 273, 298, 511.
 Fowler, J. Kingston, England, 400.
 Fowler, Edward, United States, 580.
 Fox, William, 356.
 Fox, Wilson, England, 381.
 Fraentzel, Dr., Austria, 248.
 Freire, Domingos, Brazil, 575.
 Frisbie, C. W., United States, 512.
 Frost, H. R., United States, 582.
 Fuller, Henry W., England, 204, 493, 536.
 Fuller, William, Canada, 488.
 Fulton, Dr., United States, 528.
 Furter, Professor, France, 244.
 Gadbury, W. T., United States, 519.
 Gallois, N., France, 147, 152, 284, 311.
 Gamgee, A., England, 133.
 Garner, J. H., Canada, 222.
 Garrettson, J. E., United States, 601.
 Garrod, A. B., England, 466, 545.
 Gaunt, T. J., United States, 429.
 Gazzo, J. B. C., United States, 517.
 Gerhardt, Professor, 305.
 Germain-Sée, Professor, France, 154, 271, 282, 306.
 Gibbons, Henry, United States, 374.
 Gillette, Dr., France, 41.
 Gilliford, R. H., United States, 452.
 Giordan, Dr., Italy, 412.
 Girgensohn, X., Germany, 531.
 Gleason, J. G., United States, 258.
 Goodeve, E., 356.
 Goodhart, J. M., England, 33.
 Gore, Surgeon, England, 503.
 Granville, J. M., England, 122, 336, 541.
 Graves, Professor, R. J., Ireland, 72, 247, 307, 500.
 Gray, E. B., England, 47.
 Grazzini, Dr., Italy, 453.
 Greene, R., England, 69.
 Greenhow, E. H., England, 175, 198.
 Griffiths, T. J., United States, 64.
 Gross, Dr., England, 368.
 Gross, S. D., United States, 104, 449, 464, 494.
 Grove, J., England, 329.
 Gubler, A., France, 421, 524.
 Guersant, Dr., France, 194.
 Guibert, Dr., France, 458.
 Guerder, Dr., France, 482.
 Guichon, Dr., France, 415.
 Gull, W. W., England, 82.

- Habershon, S. O., England, 272.
 Hahn, Dr., Germany, 293.
 Hall, A. R., India, 509.
 Hall, F. De Haviland, England, 432.
 Hall, Marshall, England, 53.
 Haller, Dr., Germany, 62.
 Hallowell, Mary, United States, 230.
 Hamilton, Allen M., United States, 29, 54, 121, 132.
 Hamilton, A. W., United States, 38.
 Hamilton, J. B., United States, 488.
 Hamlin, Dr., United States, 325.
 Hammond, William A., United States, 25, 37, 52, 72, 91, 105, 130, 133, 400.
 Hannay, J. B., United States, 213.
 Happel, T. J., United States, 369.
 Hardy, Professor, France, 496.
 Hare, J. C., England, 165.
 Harley, George, England, 313, 399, 401, 421.
 Harlow, Alonzo, United States, 565.
 Hartshorne, Henry, United States, 281, 288, 388, 532, 566.
 Hassall, A. H., England, 248.
 Hay, George, United States, 606.
 Hayden, Dr., Ireland, 293.
 Headland, F. W., England, 116.
 Heartt, P. T., United States, 454.
 Heberden, Dr., England, 62.
 Heller, Arnold, Germany, 404.
 Helvetius, Dr., Germany, 369.
 Heroguelle, Dr., France, 109.
 Herpin, Dr., Geneva, 55, 62.
 Herschell, George, England, 545.
 Hertzka, E., Germany, 135.
 Hewson, A., United States, 416.
 Hill, Thomas, United States, 517.
 Hiller, N., Germany, 341, 398.
 Hillier, Thomas, England, 39, 349.
 Hinton, R. K., United States, 284.
 Hirschfeld, J., Vienna, 211.
 Hitchman, W., England, 572.
 Hoag, J., United States, 450.
 Hoffman, Dr., Germany, 224.
 Hoffman, M., England, 212.
 Holcombe, W. H., United States, 475.
 Holden, E., United States, 288.
 Holden, J. S., England, 216.
 Holland, H., England, 462.
 Hollis, Dr., England, 524.
 Home, Dr., England, 62.
 Hood, Donald W., England, 390.
 Hope, J., England, 59, 158, 291, 459.
 Horion, Charles, France, 468.
 Huchard, M., France, 150, 280.
 Hughes, C. H., United States, 81.
 Huguenin, Dr., France, 480.
 Hugues, Dr., France, 251.
 Hunter, Charles, England, 101.
 Hunter, George Y., India, 515.
 Hurd, E. P., United States, 106.
 Huss, Dr., France, 312.
 Hutchins, Dr., United States, 461.
 Hutchinson, J. H., United States, 246, 514.
 Hutchinson, Jonathan, England, 330.
 Isambert, Dr., France, 222.
 Isnard, Dr., France, 342.
 Jaccoud, Professor, France, 232, 397, 556.
 Jack, R. G., England, 317.
 Jackson, J. A., United States, 315.
 Jackson, J. H., England, 28.
 Jackson, S., United States, 527.
 Jacobi, A., United States, 251.
 Jamieson, W. A., Scotland, 128.
 Jenner, W., England, 386.
 Jewett, Dr., United States, 461.
 Johnson, F. N., United States, 544.
 Johnson, George, England, 57, 166, 322.
 Johnson, James, England, 77.
 Jones, C. Handfield, England, 93, 144, 152.
 Jones, L. M., United States, 527.
 Jones, Podmore, England, 426.
 Jones, H. McNaughton, Ireland, 567.
 Jones, J. T., United States, 318.
 Josephson, Dr., Hamburg, 200.
 Jounod, Dr., France, 213.
 Jurgensen, T., Germany, 269.
 Kahler, O., Germany, 181.
 Kellogg, A. O., United States, 82.
 Kempster, W., United States, 349.
 Kennedy, Henry, England, 446.
 Kennedy, N. B., United States, 533.
 Kenyon, George, England, 71.
 Kerr, Dr., England, 355.
 Kilpatrick, A. R., United States, 409.
 King, F., United States, 471.
 Kitchen, D. H., 169.
 Knight, O. C., United States, 455.
 Knoll, Dr., Germany, 550.
 Koller, Dr. E., Germany, 160.
 Kraus, Bernhard, Austria, 292.
 Kraus, Emil, Germany, 409.
 Krishaber, D., Germany, 220.
 Krull, Dr., Germany, 423.
 Kuchenmeister, Dr., Germany, 412.
 Kunze, C. F., Germany, 373.
 Laborde, Dr., France, 126.
 Lacaille, Dr. De, Brazil, 575.
 Laennec, Dr., France, 258.
 Lambert, W., United States, 124.
 Lange, W., Germany, 469.

- Lanion, Dr. Mariana Y., 309.
 Latham, Dr., England, 296.
 Latour, M., France, 525.
 Laurent, Dr., France, 44.
 Lawson, Henry, England, 125.
 Lawson, J. B., England, 409.
 Laycock, Dr., Ireland, 278.
 Laycock, T., Scotland, 446.
 Leared, Arthur, England, 377.
 Lebert, H., Switzerland, 391.
 Leclerc, Dr., France, 152.
 Lee, Benjamin, United States, 513.
 Legari, Thomas, United States, 95.
 Leibermeister, Karl, Germany, 554.
 Leidesdorf, Dr., Germany, 51.
 Lemaire, Dr., France, 229.
 Lente, F. D., United States, 516.
 Leourst, Dr., Germany, 515.
 Levi, Dr., France, 51.
 Lewis, D., United States, 379.
 Lewis M., United States, 519.
 Liddell, Dr., United States, 362.
 Lincoln, R. P., United States, 215.
 Lisfranc, Dr., France, 466.
 Lithgow, R. A. D., England, 73.
 Lockie, Dr., England, 298.
 Lockridge, J. E., United States, 72.
 Logan, Dr., England, 288.
 Lombard, Dr., Switzerland, 189.
 Lombardo, Antonio, Italy, 254.
 Long, St. John, England, 183.
 Loomis, A. L., United States, 241, 256,
 263, 280, 285, 289, 294, 302, 436.
 Lorlet, Dr., France, 408.
 Loving, S., United States, 519.
 Lowenfeld, Dr., 283.
 Lowndes, T. M., India, 327.
 Lusanne, Dr., France, 380.
 Luton, M., France, 112.
 Lyman, Henry M., United States, 124.
 Lutz, Dr., France, 605.
 Lynch, John S., United States, 353.
 Lyndon, E. G., United States, 573.
 Mackenzie, Stephen, England, 471.
 Mackey, Edward, England, 429.
 MacLagan, T. J., Ireland, 185, 533.
 MacLaren, R., England, 242.
 Maclean, Professor W. C., England,
 361, 401.
 Madden, Dr., England, 241.
 Maffitt, A., England, 558.
 Main, John S., Scotland, 168, 378.
 Malet, A., Brazil, 244.
 Malgaigne, Professor, France, 125.
 Mann, E. C., United States, 21, 33, 34,
 84, 121, 141.
 Marcet, Dr., England, 254.
 Marieosky, L., Russia, 192.
 Maris, D., United States, 463.
 Markel, S. B., United States, 593.
 Marshall, John S., United States, 586.
 Martelli, Dr., Italy, 159.
 Martin, John, Ireland, 298.
 Martin, Ranald, 402.
 Masse, Dr., France, 409.
 Massiah, B. J., England, 422.
 Massini, Professor, Italy, 109.
 Matlack, Mary J., United States, 291.
 Maudsley, Henry, England, 119.
 Maury, R. B., United States, 365.
 May, G. P., England, 297.
 Mayer, E. R., United States, 217, 521.
 McBean, S., England, 569.
 McClellan, Ely, United States, 318.
 McGuire, L., United States, 364, 518.
 McLean, Professor, India, 510.
 McPhail, Dr., United States, 405.
 McSherry, Richard, United States, 282.
 Meadows, Dr., England, 118.
 Meigs, J. F., United States, 175, 341,
 351, 410.
 Menneret, Dr., France, 380.
 Mellersh, A. H., United States, 251.
 Menche, H., Germany, 544.
 Messler, M. J. B., United States, 358.
 Metcalf, Professor, United States, 341.
 Mettauer, Dr., 340.
 Meyer, L., Germany, 116.
 Michou, Dr., France, 258.
 Millard, Henry B., United States, 473.
 Millard, Orson, United States, 450.
 Millet, Dr., France, 388.
 Mitchell, R. W., United States, 318.
 Mitchell, S. Weir, United States, 114,
 143.
 Moinet, F. W., Scotland, 281.
 Moleschott, Professor, Italy, 453.
 Monell, J. S., United States, 155.
 Montgomery, Edward, United States,
 332.
 Monti, Alois, Austria, 409.
 Moorman, J. W., United States, 355.
 Mordough, E. F., United States, 62.
 Morris, W. J., England, 469.
 Morse, E. M., United States, 370.
 Morton, N. K., United States, 592.
 Mosler, Professor, Switzerland, 405,
 407, 517.
 Muhlberger, A., Germany, 283.
 Muir, Wm., Scotland, 467.
 Muller, W., Germany, 446.
 Munro, W., England, 138.
 Murchison, Dr., England, 316, 330,
 351, 423, 457, 532.
 Murray, John, India, 323.
 Murray, W. W., United States, 523.
 Murrell, William, England, 180.
 Mussy, Professor Gueneau de, France,
 452.

- Nairne, J. T., England, 248.
 Napier, C. O. G., England, 594.
 Nelson, A. W., United States, 564.
 Newton, R. C., United States, 522.
 Nichell, H., United States, 146.
 Nicholls, R. M., United States, 469.
 Nonat, Auguste, 527.
 Norwood, Dr., United States, 269.
 Notlmagel, Professor H., 43.
 Nowlin, J. H., United States, 585.

 Oeterlony, J. A., United States, 313.
 O'Daniel, W., United States, 512.
 Ogle, Dr., England, 45.
 Oliver, J. G., England, 151.
 Osborne, Dr., England, 443.
 Oulment, Dr., France, 44.
 Oxley, Dr., England, 97.

 Paget, Sir James, England, 550.
 Palmer, Professor A. B., United States, 260.
 Pancoast, J., United States, 148.
 Papham, John, Ireland, 268.
 Parisel, L., France, 175.
 Pap, S. V., Austria, 442.
 Paris, Dr., France, 368.
 Parks, N. O., United States, 295.
 Parry, F. W., United States, 247.
 Patton, J. C., United States, 528.
 Paul, Constantin, France, 289, 292.
 Pavy, F. W., England, 247, 356, 453.
 Peabody, J. H., United States, 523.
 Pepper, Professor W., United States, 410, 420, 554.
 Pereira, Dr., England, 356.
 Perroncito, M., France, 480.
 Peter, M., France, 129.
 Phillips, C. D., England, 429.
 Phillips, W. H., United States, 529.
 Pick, Julius, United States, 240.
 Pigne, Dupuytren, United States, 562.
 Pinel, Dr., France, 59.
 Piorry, Professor, France, 425.
 Percher, Dr., Germany, 200.
 Phares, D. L., United States, 330, 519.
 Piffard, H. G., United States, 495.
 Piso, Professor, Italy, 368.
 Polli, Professor, Italy, 573.
 Pollock, Julius, England, 479.
 Poore, G. V., England, 117, 135.
 Popoff, Professor V. N., Russia, 571.
 Porcher, Dr., France, 458.
 Porcher, F. Peyre, United States, 341, 492, 579.
 Powel, Douglas, England, 429.
 Power, R. E., England, 269.
 Prangley, Mr., England, 379.
 Pratt, H. J., United States, 443.
 Prideaux, E., France, 534, 571.

 Prout, J. S., United States, 186.
 Prout, W., England, 315.
 Pugh, T. J., United States, 502.

 Quinlan, F. J. B., Ireland, 254.

 Ragland, A. M., United States, 367.
 Ralfe, Dr., England, 469.
 Ramsey, D. C., United States, 486.
 Ramskill, J. S., England, 54, 500.
 Raspail, Dr., France, 80.
 Rawle, F., England, 359.
 Read, W., United States, 489.
 Recamier, Dr., France, 372.
 Reece, Dr., France, 312.
 Reed, C. A. L., United States, 160.
 Reith, Dr., Scotland, 544.
 Renduer, Dr., France, 446.
 Reveil, Professor O., France, 147.
 Reynolds, Russell, England, 301, 544.
 Rhode, Dr., Germany, 417.
 Richardson, B. W., England, 306.
 Ricker, Dr., Germany, 352.
 Riebe, M., France, 261.
 Rienzi, Professor, Italy, 309.
 Rifas, D. S., —, 89.
 Ringer, Sidney, England, 260.
 Rirey, P. M., United States, 191.
 Roberts, F. T., England, 397, 412, 455.
 Robinson, Beverly, United States, 191, 203, 279.
 Rockwell, A. D., United States, 124.
 Roe, John O., United States, 216.
 Roger, Dr., France, 41.
 Rogers, Lewis, United States, 314.
 Romberg, Dr., Germany, 127.
 Rony-Saucerotte, Dr., Paris, 266.
 Rosenstein, Professor S. S., Germany, 304.
 Rosenthal, Professor M., Austria, 127.
 Rothrock, A., United States, 133.
 Rougier, Dr., France, 45.
 Roziere, M., France, 147.
 Rubini, Dr., Italy, 326.
 Rumbold, Dr., United States, 408.
 Runge, Dr. F., Germany, 21.
 Ruppaner, A., United States, 101.
 Ruschenberger, W. S. W., United States, 325.
 Russ, Ross C., United States, 167.
 Rutherford, Professor, Scotland, 424.

 Salter, Hyde, England, 156, 494.
 Sansom, Dr., England, 254.
 Sanydby, Dr., England, 246.
 Sawyer, Dr., United States, 524.
 Sawyer, James, England, 96.
 Scearce, J. B., United States, 260.
 Schafirt, A. J., United States, 406.
 Schenck, W. L., United States, 590.

- Schiff, M., Switzerland, 315.
 Schildowsky, Dr., Russia, 413.
 Schilling, Dr., Scotland, 526.
 Schneider, H., Vienna, 212.
 Schnitzler, Dr. Johann, Austria, 147, 250.
 Schoonemann, C. W., United States, 500.
 Scholz, Dr., Germany, 569.
 Schroeter, Leopold, Austria, 289.
 Schulz, O. T., United States, 266.
 Schwalbe, Carl, Costa Rica, 360.
 Schwarz, M., Germany, 485.
 Scriven, J. B., India, 509.
 Scudamore, Sir C., England, 255.
 Scudder, John M., United States, 446.
 Sedgwick, William, England, 321.
 Sedgwick, L., England, 195.
 Seegen, Professor, Germany, 452.
 Seeley, W. T., United States, 382.
 Sée, Professor Germain, Paris, 154, 271, 282, 306.
 Seguin, E. C., United States, 17, 49, 69, 89, 98, 215.
 Seiler, Carl, United States, 189.
 Semple, Armand, England, 43.
 Seure, A., France, 70.
 Shaw, J. C., United States, 52.
 Shearer, J. Y., United States, 523.
 Sholl, E. H., United States, 264.
 Sibson, F., England, 439.
 Silverthorne, United States, 261.
 Silvestrini, Professor, Italy, 262.
 Simpson, Professor, Scotland, 382.
 Sinkler, Wharton, United States, 522.
 Skoda, Dr., Germany, 293.
 Skinner, Stephen, England, 559.
 Skorejewsky, B., Russia, 518.
 Slane, Dr., India, 520.
 Smelt, Isaac, United States, 525.
 Smith, A. H., United States, 394.
 Smith, J. Lewis, United States, 411, 487.
 Smith, Shingleton, England, 295.
 Spencer, Ethan, United States, 552.
 Spender, J. K., England, 76, 356.
 Spitzka, E. C., United States, 56.
 Squibb, E. R., United States, 324, 352.
 Stedman, C. E., United States, 553.
 Steinbruck, Dr., Germany, 253.
 Stelwagen, T. C., United States, 352.
 Stevens, George T., United States, 40.
 Stevens, William, England, 320.
 Stewart, F. E., United States, 485.
 Stewart, William, India, 372.
 St. George, George, Ireland, 172.
 Stillé, Professor Alfred, United States, 112, 175, 250.
 Stokes, Professor William, Ireland, 176.
 Stone, Dr., England, 46.
 Storer, H. R., United States, 113.
 Stricker, Dr., Germany, 534.
 Strohl, Dr., France, 268.
 Stuckert, William, England, 403.
 Sullivan, John, British India, 325.
 Sunderland, Dr., United States, 244.
 Sunderland, J. F., United States, 563.
 Svetlin, Dr., Vienna, 51.
 Sweringen, H. V., United States, 93, 122, 127, 143, 147.
 Sydenham, Dr., England, 462.
 Syme, James, Scotland, 603.
 Symonds, J. A., England, 163.
 Tanner, Thomas Hawkes, England, 65, 74, 155, 221, 236, 248, 281, 340, 530.
 Taylor, Seymour, England, 206.
 Taylor, A. S., England, 598.
 Thomas, T. Gaillard, United States, 375.
 Thompson, A. T., England, 61.
 Thompson, Ashburton, England, 111.
 Thompson, B. F., England, 363.
 Thompson, Sir H., England, 470.
 Thompson, Symes, England, 243.
 Thompson, William, United States, 337.
 Thompson, William H., United States, 521.
 Thorowgood, John C., England, 149, 238.
 Thurston, G. B., United States, 318.
 Tillesen, Dr., Germany, 405.
 Tilt, Edward J., England, 97.
 Tinker, G. L., United States, 181.
 Tisne, Dr., France, 251.
 Todd, J. F., England, 291.
 Tourasse, Dr., France, 316.
 Townsend, E. P., United States, 425.
 Trastour, A., France, 154.
 Traube, L., Germany, 533.
 Troubert, C., France, 111.
 Trousseau, Professor, France, 123, 153, 445.
 Tuckwell, H. M., England, 47.
 Turnbull, James, England, 176.
 Tyrrell, Walter, England, 53.
 Tyson, James, United States, 467, 565.
 Upsher, J. N., United States, 470.
 Vachetta, Professor A., Italy, 484.
 Valleix, M., France, 416.
 Van Bibber, John, United States, 42.
 Van Buren, Professor W. H., United States, 340, 415.
 Vance, Reuben A., United States, 134.
 Vansant, Dr., United States, 522.
 Varick, Dr., United States, 251.
 Velpeau, Dr., France, 573.

- Vesey, A., England, 212.
 Vogt, Dr., Germany, 126.
 Voison, Aug., France, 55.
 Von Niemeyer, Felix, Germany, 83,
 126, 138, 153, 187, 198, 209, 227, 438,
 487.
 Von Ziemssen, Professor, Germany,
 489.
- Waakes, Edward, England, 127, 139.
 Wade, W. F., England, 41.
 Waldenberg, L., Germany, 188, 221.
 Walker, B. M., United States, 139.
 Walker, E. F., United States, 416.
 Wallace, E., United States, 484.
 Walshe, Dr., England, 287, 456.
 Waller, A., Geneva, 112.
 Warburg, Dr., India, 510.
 Waring, Dr., England, 356.
 Warner, Muller, Germany, 455.
 Washington, B. H., United States, 366.
 Waters, A. T. H., England, 265, 298.
 Waters, Eben, Scotland, 124.
 Watson, Thomas, Scotland, 176, 406.
 Waylen, C. W., India, 503.
 Webb, F., United States, 415.
 Webb, R. D., United States, 501.
 Webber, S. G., United States, 427.
 Weir, A. M., Scotland, 210.
 Welsh, W. M., Scotland, 155.
- Wertner, Dr., Vienna, 179.
 West, Charles, England, 43.
 Westmoreland, J. G., United States,
 60.
 Whipham, T., England, 220.
 White, John M., United States, 388.
 White, R. P., Ireland, 349, 558.
 Whitehead, W. E., United States, 363.
 Wilkes, G., England, 559.
 Wilkins, H. S., United States, 408.
 Wilks, Samuel, England, 270.
 Willebrand, Dr. F., Russia, 512.
 Willemen, Dr., France, 93.
 Williams, C. J. B., England, 185, 234.
 Williams, John, England, 177.
 Williams, J. D., United States, 564.
 Wilson, J. C., United States, 559.
 Wilson, Thomas, England, 434.
 Wood, G. B., United States, 410.
 Wood, H. C., United States, 366.
 Woodworth, J. M., United States, 319.
 Wright, Henry G., 65, 66, 76
- Yeaman, J. B., United States, 260.
 Yeo, L. Burney, England, 228, 374.
 Young, David, Italy, 348.
- Zarzana, E., Italy, 453.
 Ziemssen, Professor Von, Germany,
 391.

II. INDEX OF REMEDIES AND REMEDIAL MEASURES.

N. B.—Articles of the *Materia Medica* should be looked for under their *pharmacopœial* name. References are confined to passages where some *special directions* are given for their exhibition or preparation.

- Abdominal compression, 463.
 Abortive treatment of pneumonia, 259.
 Acacia, 389.
 Acetum, 158.
 Acida, in typhoid fever, 561.
 in typhus fever, 566.
 Acidum aceticum, 210.
 Acidum arseniosum, 30, 34, 42, 59, 106,
 119, 251.
 in constipation, 342.
 in fatty heart, 298.
 in vomiting, 428.
 Acidum benzoicum, in biliary calculi,
 316.
- Acidum benzoicum, in jaundice, 421.
 in uræmia, 478.
 Acidum carbolicum, 108, 178, 217, 250.
 in colic, 330.
 in diabetes, 453.
 in diarrhœa, 353.
 in jaundice, 421.
 in pyrosis, 426.
 in tape worm, 408.
 in typhoid fever,
 556.
 in variola, 571.
 in vomiting, 428.
 Acidum carbonicum, in vomiting, 428.

- Acidum chlor-hydricum*, in dyspepsia, 374.
Acidum gallicum, 210, 382.
 in albuminuria, 435.
 in Bright's disease, 442.
 in pyrosis, 426.
Acidum hydrobromicum, 29, 30, 58.
Acidum hydrocyanicum, 212, 383, 523.
 in vomiting, 428.
Acidum hyperosmicum, 109.
Acidum lacticum, 383.
 in diabetes, 446.
Acidum muriaticum, 356.
 nitricum, in constipation, 344.
 in fatty heart, 298.
 in hepatitis, 403.
 nitro-muriaticum, 79, 123, 175, 356.
 in diabetes, 446.
 in fatty heart, 298.
 in hepatitis, 402.
 nitrosi, 353.
 phenicum, in typhoid fever, 561.
 in yellow fever, 575.
 phosphoricum, 212.
 in cholera, 321.
 in diabetes, 454.
 pitzaholicum, in constipation, 344.
 pyrogallicum, 212.
 salicylicum, 139.
 in cerebro-spinal meningitis, 486.
 in jaundice, 421.
 in rheumatism, 533.
 in typhoid fever, 563.
 in variola, 571.
 in worms, 413.
 in yellow fever, 582.
 succinicum, in biliary calculi, 314.
 sulpho-phenicum in yellow fever, 575.
 sulphuricum, 120.
 in cholera, 319.
 in typhus fever, 570.
 in variola, 573.
 in yellow fever, 582.
 in plumbism, 605.
 sulphurosum, 217.
 in pyrosis, 427.
 tannicum, 212, 250, 259, 354.
 in albuminuria, 435.
 in diabetes, 449.
 in yellow fever, 582.
 tartaricum, in vomiting, 429.
- Aconitum*, 30, 81, 96, 98, 107, 193, 258, 289.
 in Bright's disease, 442.
 in catarrhal jaundice, 417.
 in rheumatism, 544.
 in yellow fever, 577.
 Actual cautery, 143.
 Acupuncture, 111, 463.
 Adeps, in seat worms, 416.
Adonia vernalis, in cardiac dilatation, 287.
Æsculus hippocastanum, 107.
Agaricum, 249.
 Air, condensed, 200.
 Alcohol, 33, 96, 108, 250.
 in pneumonia, 264.
 in trichinosis, 417.
 in typhoid fever, 561.
 in typhus fever, 565.
Aleuritis triloba, in constipation, 342.
Alianthus glandulosa, in tape worm, 407.
 Alimentation, rectal, 394.
 Alkalies, in diabetes, 452.
 in torpidity of liver, 424.
 in rheumatism, 536, 548.
 Alkaline hypophosphites, in phthisis, 228.
 Allium, 178.
 Almond flower cakes, in diabetes, 449.
 Aloes, 78.
 in constipation, 340, 432.
 Aloin, 340.
 Alstonia, 520.
 Alumen, 148, 178, 210, 354.
 in hæmaturia, 469.
 in pyrosis, 426.
 in tape-worm, 407.
 in typhoid fever, 561.
 Ammoniacal vapor, 148.
 Ammoniacum, 158, 167, 193.
 Ammonium acetatis, 77, 421.
 in typhoid fever, 561.
 Ammonium bromidum, 58, 73, 421.
 Ammonium carbazotatis, 354, 550.
 Ammonium carbonas, 48, 58, 71, 77, 158, 169, 421.
 in diabetes, 452.
 in typhoid fever, 561.
 in variola, 572.
 Ammonium chloridum, 77, 108, 174, 421.
 in dysentery, 372.
 in hepatic congestion, 403, 424.
 in vomiting, 427.
 Ammonium citratis, in diabetes, 452.
 in vomiting, 427.
 Ammonium formias, 59, 119.
 Ammonium iodidum, 250.

- Ammonium muriaticum, 69, 146.
 in albuminuria, 434.
 in biliary calculi, 314.
 Ammonium phosphatis, in diabetes, 452.
 Ammonium valerianatis, 59.
 Amyl nitrite, 53, 59, 73, 157, 171, 249, 262, 280, 520.
 in vomiting, 429.
 Amylum, in variola, 573.
 Angelica, 179.
 Anilin, in chorea, 42.
 Anise, in flatulent colic, 330.
 Anodyne, Hoffman's, 162.
 Anthemis, in colic, 108, 332.
 Antimonial ointment, 116.
 Antimonium et potassium tartratis, 194, 258, 400.
 in albuminuria, 434.
 in chorea, 41, 43.
 Antimonium ses-iodidum in heart disease, 296.
 Antimonium sulphuretum, 282.
 in diabetes, 450.
 Anti-parasitic treatment of tuberculosis, 239.
 typhoid fever, 559.
 Antiseptic inhalations, 229.
 powder of Niemeyer, 226.
 treatment of typhoid fever, 559.
 Apiol, 108.
 Apomorphia, 43, 179.
 in jaundice, 418.
 Aquam, in albuminuria, 434.
 Aquam ammonium, 59, 112, 158.
 calcium, in seat worms, 416.
 in variola, 573.
 in vomiting, 428.
 Argenti chloridum, 158.
 Argenti nitratis, 43, 77, 112, 119, 121, 354, 388.
 in dysentery, 366.
 in seat worms, 416.
 in variola, 573.
 oxidum, 66, 210.
 in pyrosis, 426.
 Arlentin, 461, 466.
 Armoraciæ radix, 120, 462.
 Arsenicum, 19, 27, 34, 43, 71, 159, 196, 232, 520.
 in albuminuria, 434.
 in anæmia, 484.
 in Bright's disease, 442.
 in cholera, 317.
 in chorea, 39.
 in diabetes, 452.
 in diarrhœa, 349.
 in yellow fever, 581.
 bromidum, 59, 78.
 Arsenicum, bromidum in diabetes, 447.
 Aisenite of bromine, in diabetes, 452.
 Arsenical cigarettes, 196, 217.
 Arsenicum chloro-phosphidum, 122.
 Asarum canadense, in seat-worms, 416.
 " Europæum, 78, 595.
 Asclepias, syriaca, 461.
 Asparagin, 291.
 Aspidium marginale, in tape-worm, 407.
 Aspidosperma quebracho, 159, 200.
 Aspidospermine, 159.
 Aspiration, 206.
 Assafoetida, 59, 84, 159.
 in colic, 332.
 Asthma cigarettes, 215.
 Astringents, in hemorrhage, 208.
 Atropia, 96.
 Atropia sulphatis, 194, 226.
 in variola, 572.
 Auri chloridum, 34, 90.
 in albuminuria, 433.
 in Bright's disease, 442.
 in diabetes, 446.
 Bael, 354.
 Balsams, 177.
 Balsamum peruvianum, 179.
 Bandages, in dysentery, 366.
 Barii chloridum, 30, 210.
 Baths, 92, 98, 120, 121, 269, 279, 361.
 bran, in diabetes, 449.
 camphorated vapor, in Bright's disease, 443.
 in chorea, 46.
 cold in typhus fever, 568.
 cold in variola, 574.
 cold shower, in incontinence of urine, 464.
 in nephritis, 473.
 nitric acid, in intermittent fever, 509.
 nitro-muriatic, in biliary calculi, 317.
 nitro-muriatic, in hepatitis, 402.
 in rheumatism, 546.
 of potassium sulphide, in plumbism, 607.
 shower, 63.
 steam, in jaundice, 419.
 sulphur, in hepatitis, 402.
 tonic, 481.
 Turkish, 63.
 in torpidity of liver, 425.
 Belladonna, 43, 54, 60, 71, 85, 104, 108, 121, 151, 286.
 in albuminuria, 434.
 in biliary calculi, 313.
 in constipation, 335.
 in diabetes, 446.

- Belladonna*, in incontinence of urine, 464.
in intestinal obstruction, 332.
in typhus fever, 569.
in variola, 572.
in yellow fever, 581.
plasters, in irritable heart, 302.
- Benzinae*, in dyspepsia, 374.
in tape-worm, 407.
- Benzoinum*, 146.
in constipation, 343.
in dysentery, 364.
in pyrosis, 426.
odoriferum, in hæmaturia, 469.
- Berberiæ sulphas*, 108.
- Berberin*, 520.
- Bethesda water*, 455.
- Bile*, ox or pig, 330.
- Bismuthum* in pyrosis, 425.
- Bismuthum oxidum*, 194.
subcarbonas, 78.
subnitras, 60, 66, 77.
in diarrhœa, 350.
in dyspepsia, 380.
in typho-malarial fever, 564.
- Blatta orientalis*, 462.
- Blisters*, 63, 269.
in cholera, 317.
in heart disease, 279, 295.
in jaundice, 418.
in rheumatism, 546.
in vomiting, 429.
- Blood*, in anæmia, 482.
- Boldo*, 381.
- Boldine*, 381.
- Bran bread* in diabetes, 449.
- Brayera*, in tape worm, 407.
- Bromine*, 218.
arsenite of, in diabetes, 452.
- Bryonia*, 266.
- Buttermilk*, in Bright's disease, 439.
- Buxinæ sulphas*, 520.
- Cactus grandiflora*, 301.
- Caffea*, 108, 160, 462.
in typhoid fever, 562.
- Cajuputi oleum*, 78, 110, 120.
in flatulent colic, 330.
in rheumatism, 550.
- Calcium carbolas*, 354.
chloridum, 251.
iodidum, 154, 244.
lacto-phosphatis, in anæmia, 479.
phosphatis, 179, 354.
saccharatis, in constipation, 343.
sulphidum, in diabetes, 452.
- Calumba*, 381.
- Camphora*, 60, 78, 266.
in cholera, 326.
in chorea, 40.
in constipation, 343.
in diarrhœa, 353.
in typhoid fever, 562.
in variola, 572.
in yellow fever, 581.
mono-bromidum, 85.
- Cannabine tannate*, 95.
- Cannabis indica*, 60, 69, 87, 91, 108, 160, 194, 462.
in Bright's disease, 442.
in chorea, 43.
- Cantharis*, 112, 119, 466.
in albuminuria, 434.
- Caoutchouc*, 251.
- Carbo ligni*, in colic, 332.
- Carbon bisulphide*, 71, 76.
- Cardiac blistering*, 295.
- Carlsbad salts*, 419.
- Caryophyllum*, in colic, 333.
- Casca bark*, 287.
- Cascara sagrada*, in constipation, 343.
in dyspepsia, 381.
- Cascarilla*, 354.
- Castoreum*, 60, 160.
- Catarrh snuff*, 188.
- Catechu*, 224, 354.
- Cathartics*, 68, 371, 376.
in apoplexy, 32.
in Bright's disease, 439.
in constipation, 338.
in dropsy, 460.
hypodermic use of, 341.
in jaundice, 423.
- Cautery actual*, 64.
- Centaurea benedicta*, 60.
- Cerii oxalas*, 60, 179, 231, 382.
- Charcoal*, in flatulent colic, 330.
in dyspepsia, 377.
- Cheken*, 194.
- Chenopodium anthelminticum*, 410, 462.
- Chian turpentine*, 469.
- Chinoidin*, 507.
- Chloral hydrate*, 19, 36, 70, 91, 102, 283.
in albuminuria, 434.
in chorea, 43.
in delirium tremens, 588.
in rheumatism, 544.
in typhoid fever, 562.
in sea-sickness, 428.
in variola, 572.
- Chlorine vapor*, 148.
- Chlorodyne*, 329.
- Chloroformum*, 43, 60, 71, 78, 101, 108, 210.
in angina pectoris, 282.
in biliary calculi, 315.
in cholera, 327.

- Chloroformum in dysentery, 361.
 in obstinate chills, 512.
 in uræmia, 477.
 in vomiting, 428.
 Cholagogues in torpidity of liver, 423.
 Cholera mixtures, 329.
 Cicutin, 179.
 Cigarettes of arsenic, 197.
 for asthma, 215.
 of nitre, 152.
 Cimicifuga, 37, 79, 84, 179, 301, 521.
 in fatty heart, 298.
 in rheumatism, 544, 560.
 Cinchona, 60, 179, 251.
 Cinchonidiæ iodidum, 522.
 Cinchonidinæ salicylatis, 507.
 Cinnamomum, in colic, 333, 355.
 in vomiting, 428.
 Circumcision, for incontinence, 467.
 Climate in heart disease, 276.
 and travel in phthisis, 241.
 Cocculus indicus, 119.
 in vomiting, 428.
 Codeia, 246.
 in diabetes, 453.
 in gout, 496.
 Codeia phosphas, 94.
 Coffee, 92, 156.
 Colchicum, 30, 68, 78.
 in constipation, 343.
 in gout, 498.
 in hepatitis, 403, 422.
 in rheumatic carditis, 297.
 in rheumatism, 544.
 Cold water pack, in pneumonia, 264.
 in rheumatism, 547.
 Collinsonia canadensis, 224.
 Colocyth, 343, 382.
 Compressed air, 305.
 Compression, 201.
 abdominal, in ascites, 463.
 of thorax, in pleurisy, 257.
 Condensed air, 200.
 Conium, 19, 44, 56, 60, 87, 108, 132.
 Convallaria majalis, in cardiac dilatation, 287, 306.
 Copaiba, 180, 462.
 in jaundice, 422.
 tape worm, 408.
 Cotoin, in cholera, 327.
 in diarrhœa, 351.
 Counter-irritation, 32, 33, 122, 132, 166, 233.
 in epilepsy, 57.
 in jaundice, 420.
 Creasotum, 83, 180, 251, 355.
 in cholera, 318.
 in colic, 330.
 in diabetes, 453.
 in tape worm, 408.
 Creasotum in vomiting, 428.
 Creta præparata, 180.
 Cricus lanceolatus, 211.
 Croton chloral, 106, 108, 246.
 Cubeba, 144, 180, 382.
 Cupping, 21, 63, 72, 87, 127, 212.
 in albuminuria, 432.
 in Bright's disease, 436.
 in typhus fever, 566.
 Cuprum, in pneumonia, 266.
 in diabetes, 453.
 ammoniatum, 37, 60, 108.
 sulphas, 208, 355.
 Curare, 60.
 Cuspariæ cortex, 355.
 Datura tatula, 160.
 Delphina, 120.
 Diaphoretics, in dropsy, 455.
 Diet, 88.
 in albuminuria, 432.
 in biliary calculi, 314.
 in Bright's disease, 436, 438, 440, 473, 474.
 in constipation, 335.
 in diabetes, 448, 449, 450.
 in diarrhœa, 350.
 in dysentery, 367.
 in incontinence of urine, 464.
 in jaundice, 418.
 in malaria, 505.
 milk, in diabetes, 451.
 in rheumatism, 540.
 in torpidity of liver, 423.
 Dietetics in diabetes, 447.
 in heart disease, 275.
 in vomiting, 427.
 Diethylacetal, 96.
 Digitaline, 30.
 Digitalis, 85, 108, 132, 160, 251, 462.
 in albuminuria, 434.
 in Bright's disease, 436, 472.
 in cardiac dilatation, 285.
 in nephritis, 474.
 in purpura hemorrhagica, 530.
 in typhoid fever, 562.
 in typhus fever, 568.
 Diuretic liniment of Hahn, 293.
 Diuretics, 87, 173.
 in Bright's disease, 438.
 in cardiac dilatation, 287.
 in dropsy, 455, 459.
 Douche, nasal, 192.
 Dover's powder, 249.
 Dracontium, 160.
 in chorea, 44.
 Drytreatment of pleuritic effusions, 204.
 Duboisia, 249.
 Elastic respirator, 201

- Elaterium, in Bright's disease, 439.
 in jaundice, 420.
 Electricity, 32, 44, 71, 87, 98, 112, 117,
 121, 132, 148.
 in Bright's disease, 443.
 in constipation, 346.
 in diabetes, 445.
 in dysuria, 466.
 in rheumatism, 551.
 Elimination in cholera, 325.
 Embrocations of cod liver oil in phthi-
 sis, 236.
 Emetics, 63, 165.
 in diarrhoea, 354.
 in intermittent fever, 509.
 in jaundice, 421.
 Enemata, 32, 354, 359, 371.
 in jaundice, 423.
 in nephritis, 475.
 in seat worms, 414.
 of tobacco, in intestinal ob-
 struction, 331.
 Ergota, 31, 56, 87, 108, 119, 121, 143,
 211, 355.
 in cardiac dilatation, 286.
 in diabetes, 445, 449.
 in hæmaturia, 468.
 in rheumatism, 544.
 in trichinosis, 417.
 in typhoid fever, 562.
 Erigeron canadensis, 355.
 Erythroxylon coca, in typhus fever, 569.
 Eserine, 372.
 Ether, 80, 107.
 in tape worm, 408.
 in spray, 38.
 Ethereal tincture of acetate of iron,
 305.
 Ethyle iodide, 154.
 Eucalyptus globulus, 216, 301, 382, 503.
 in albuminuria, 434.
 in pneumonia, 267.
 in typhoid fever, 562.
 in worms, 412.
 Euonymus atropurpureus, 475.
 Euonymin, in jaundice, 422, 424.
 Eupatorium perfoliatum, in tape worm,
 408.
 Euphorbia pilulifera, 160.
 corollata, 442.
 Euphrasia officinalis, 218.
 Exercise, 87, 92.
 Expiration, forced, 155.
 Faradization of the spleen, 513.
 Farina maidis, 522.
 Fattening food, 115.
 Fel bovinum purificatum, 84, 344.
 in intestinal obstruction, 333.
 Felix mass, 405, 408.
 Fermentoleum solani, 596.
 Fermentum, 372.
 Ferric alum, 190.
 Ferrum, 34, 68, 86, 109, 233, 251.
 in diabetes, 453.
 in dysentery, 364.
 in dyspepsia, 376.
 in malaria, 514.
 in pernicious anemia, 480.
 albuminatum, 484.
 bromidum, 37.
 carbonas, 344.
 chloridum, tinctura, 186, 224.
 in albuminuria, 432.
 in fatty heart, 298.
 in seat-worms, 416.
 dialysatum, 480.
 iodidum, 435, 464.
 malate, 391.
 mistura aromatica, 180.
 nitratis, 426.
 oxidum hydratum, 44.
 oxysulphatum, 519.
 pernitratis, liquor, 355.
 peroxidum, in biliary calculi,
 314.
 potassio-tartrate, 299, 356.
 pyrophosphatis, 104.
 sulphas, 177.
 Fomentations, in dysentery, 361.
 Food, fattening, 115.
 Fowlers' solution, in typhus fever, 564.
 Franklinization, 112.
 Fraxini foliæ, in gout, 495.
 Frigus, 112.
 Fuchsin or rosanilin hydrochloride, in
 albuminuria, 435.
 Fuligo ligni, in seat-worms, 416.
 Fumigations, 195.
 Furfurine nitras, 523.
 Galbanum, 109.
 Galla, 356.
 Galvanic cautery, 214.
 Galvanism, 32, 74.
 in intestinal obstruction,
 334.
 in plumbism, 607.
 Gambogia, 368.
 Gelsemium, 109, 180, 523.
 Gentian, 305, 382.
 Germicides, in hepatitis, 399.
 Glonoin, 109.
 Glycerinæ, 251.
 in diabetes, 453.
 in variola, 574.
 Glycerine of tannin, 246.
 Granati fructus cortex, 355.
 Granati radicis cortex, in tape-worm,
 408.

- Grape cure, in phthisis, 254.
in diarrhœa, 347.
- Grindelia robusta, 160, 216.
squamosa, 523.
- Guaiacum, 109, 222.
in Bright's disease, 443.
in rheumatism, 549.
- Guarana, 79, 218.
- Hæmatoxylum, 356.
- Hahn's diuretic liniment, 293.
- Hamamelis, 211.
- Hamlin's cholera mixture, 325.
- Heim's pill, 227.
- Helenin, 180, 267.
- Helonias dioica, 475.
- Herba serpylli, 596.
- Hoffman's anodyne, 162.
in valvular heart disease, 305.
- Hope's mixture, 351.
- Hops, 382.
- Hot packing in rheumatism, 547.
- Hydrargyrum, 65, 119, 382.
in Bright's disease, 437.
in dysentery, 368.
in jaundice, 420.
in nephritis, 473.
in typhoid fever, 563.
ammoniatum, 194.
bromidum, in hepatitis, 403.
chloridum corrosivum, 31.
in apoplexy, 23.
in Bright's disease, 441.
in diarrhœa, 356.
in seat-worms, 416.
chloridum mite, 31, 77, 382.
in cholera, 318.
in dengue, 492.
in diarrhœa, 346.
in seat-worms, 416.
in trichinosis, 417.
in yellow fever, 581.
cum creta, 313.
iodidum rubrum, 60.
in malarial enlargement of spleen, 403.
viride, 113.
sulphas flava, 224.
unguentum, in variola, 574.
- Hydrastis, 383, 523.
- Hydriodate of hyoscin, 248.
- Hydrogenum peroxidum, 251.
in diabetes, 453.
- Hydrotherapy, 123.
- Hygiene of chorea, 46.
- Hygiene of diabetes, 444.
" incurable heart disease, 278.
- Hyoscin, hydriodate, 248.
- Hyoseyamus, 61, 79, 84, 85, 93, 109, 160, 301.
- Hypodermic injections, 63, 105.
use of cathartics, 341.
quinine, 514.
- Hypophosphites, in phthisis, 236.
alkaline, in phthisis, 228.
- Ice, 139, 397.
in apoplexy, 33.
in cholera, 329.
in hæmaturia, 469.
in vomiting, 429.
- Ignatia, 97.
- Inhalations, 149, 161.
alkaline, 305.
antiseptic, 229.
in catarrh, 186.
in chorea, 46.
in epilepsy, 57.
tonic, 481.
Warren's, 240.
- Injections, hypodermic, 63.
intra-venous, in cholera, 329.
in seat worms, 414.
- Inspiration, forced, 155.
- Insufflations, 225.
- Iodinium, 149, 160, 252, 403, 523.
in dysentery, 368.
in malarial fever, 512.
in typhoid fever, 562.
in variola, 574.
in vomiting, 429.
- Iodoformum, 113, 193, 252.
in diabetes, 453.
in rheumatism, 550.
in worms, 413.
- Ipecacuanha, 65, 109, 160, 167, 212, 401.
in Bright's disease, 443.
in cholera, 328.
in dysentery, 361.
in torpidity of liver, 424.
in vomiting, 428.
in yellow fever, 581.
- Iridin, 425.
- Ixora danduca, 369.
- Jaborandi, 249, 268, 309, 475.
in Bright's disease, 443.
in diabetes, 453.
- Jalapæ, pulveris comp., 340.
- Jamaica dogwood, 94.
- Jarvis' snare, 217.
- Juglans, 252.
- Kairin, 267.

- Kairin in rheumatism, 544.
in typhoid fever, 563.
Kameela, in tænia, 409.
Kerosene, in rheumatism, 550.
Kino, 356, 383.
in pyrosis, 426.
Koosso, in tænia, 404, 409.
Koumiss, 181, 252.
in vomiting, 429.
Krameria, 356.
Kreochyle, 383.
- Larix Europæa, 181.
Laudanum, in alcoholism, 589.
Lavandula, 79.
Leeches, 22, 33, 78, 113, 120, 132.
in hepatitis, 399.
in jaundice, 418.
in vomiting, 430.
Lemon juice, in rheumatism, 535.
Leptandra virginica, 524.
Leptandrin, in constipation, 342.
in diarrhœa, 353.
in torpidity of liver, 425.
Lime juice, in rheumatism, 535.
Limonum succus, 369.
Liniment, diuretic, 293.
stimulating, 173.
Linimentum plumbi opiatum, 106.
Listerine, 189.
Lithium bromidum, in rheumatism, 544.
citras, 142.
Lobelia inflata, 150.
Lupulin, 97, 426.
- Magnesium, 78.
sulphas, 344, 353, 422.
in plumbism, 605.
- Male fern, 405.
Manaca, in rheumatism, 544.
Manganese oxidum nigrum, 384.
sulphas, in jaundice, 420.
Marrubium, 181.
Massage, 87.
in constipation, 334.
Matco, 212, 356.
Meat, raw, in phthisis, 233, 244.
in diarrhœa, 348.
Mechanical remedies for hay fever, 219.
Medication by the rectum, 365.
Mentha, 114.
Menthol, 114.
Metallotherapy, 120.
Methyl salicylatis, in rheumatism, 545.
Midzu ame, 384.
Milk diet, 143, 388, 391.
in diabetes, 447, 451.
in diarrhœa, 348.
peptonized, 393.
warm, as a gargle, 220.
- Mineral acids, in dyspepsia, 384.
in hepatitis, 402.
waters, in dysentery, 370.
in gout, 497.
- Monarda, 114.
Morphia, 109.
Morphia acetas, in angina pectoris, 283.
bromidum, in diabetes, 454.
hydrochloras, 277.
Moschus, 61, 84, 97, 161.
in typhus fever, 569.
Moxas, 123.
Mucuna, 120.
Muscarine, 169, 267.
in constipation, 344.
in diabetes, 447.
- Myristica, 120, 356.
Myrrha, 181.
- Napelline, 94, 110.
Naphthaline, 181.
Narcein, 73.
Narcotics, in Bright's disease, 439.
Narcotine, 524.
Nasal douche, 192.
Nectandra, 81.
Nerve stretching, 126.
vibration, 122.
Neurectomy, 114.
Neuroteny, 114.
Neurotomy, 114, 141.
Nickel bromidum, 61.
sulphas, 81, 373.
Niemeyer's antipyretic powder, 226.
Nitre, 146.
cigarettes, 152.
paper, 147.
- Nitric acid bath, in intermittent fever, 509.
Nitrite of amyl, 50, 61, 73, 157, 171, 249, 262.
in angina pectoris, 280.
in vomiting, 429.
- Nitrogen, 161.
Nitrogen gas, 253.
Nitro-glycerine, 34.
in albuminuria, 433.
in nephritis, 475.
in vomiting, 429.
- Nursing, 88.
Nux juglandis, in vomiting, 429.
Nux vomica, 61, 311, 356.
in colic, 333.
in diabetes, 453.
in pyrosis, 426.
in tape-worm, 409.
in vomiting, 429.
in yellow fever, 577.
- Occupation, 87.

- Oenothera biennis*, 181.
 in typhoid fever, 563.
Oleum cajuputi, 78, 110, 120.
 in rheumatism, 550.
carui, 332.
menthæ piperitæ, as a carminative, 333.
morruæ, 33, 169, 233.
 in Bright's disease, 436.
 in chorea, 44.
 by embrocation, 236.
 in phthisis, 234.
 in rheumatism, 550.
myristicæ, in colic, 333.
olivæ, in biliary calculi, 316.
 in chronic hepatitis, 425.
ricinum, in cholera, 322, 386.
 in diarrhœa, 348.
 in dyspepsia, 374.
 in yellow fever, 582.
terebinthinæ, 62, 79, 110, 333.
 in rheumatism, 551.
 in round worms, 411.
 in tænia, 406.
tiglii, 22, 110, 119.
 in albuminuria, 436.
 in jaundice, 420.
Opium, 19, 36, 61, 73, 91, 110, 151, 199, 353.
 in biliary calculi, 313.
 in cholera, 318.
 in diabetes, 448, 449, 454.
 in malarial fever, 505.
 in nephralgia, 471.
 in rheumatism, 545.
 in typhoid fever, 563.
 in typhus fever, 569.
 in variola, 573.
 in yellow fever, 581.
Ox bile, 330.
Oxygen inhalations, 61, 161, 435.
 in diabetes, 447.
Packing, in rheumatism, 547.
Pancreatic emulsion, 227, 253.
Paracentesis thoracis, 203, 292.
Paracotoin, in cholera, 328.
Paraldehyde, 90.
Pelletierine, in tape worm, 409.
Penthorum sedoides, 181.
Peponis semina, in tape worm, 409.
Peppermint, in flatulent colic, 330.
Pepsina, 384.
 in tape worm, 409.
Peptonized formula for gastric ulcer, 393.
 milk, 393.
Peptonized milk-gruel, 393.
Petroleum, 161.
 in rheumatism, 550.
Phellandrium aquaticum, 170.
Phosphorus, 19, 34, 38, 61, 73, 86, 119.
Physostigma, 90, 110, 124, 138.
 in cerebro-spinal meningitis, 486.
 in chorea, 40.
 constipation, 344.
Phytolacca decandra, in rheumatism, 550.
Picrotoxin, 119, 132, 250.
Pig's bile, in jaundice, 421.
Pill of Heim, 227.
Pilocarpine, 157, 181, 250.
 in Bright's disease, 437.
 in uræmia, 476.
Pinus canadensis, 215.
Piper nigrum, in cholera, 328.
Piscidia erythrina, 94, 110.
Pix Burgundica, 129.
 liquida, 182.
 in constipation, 344.
Plasters, 183.
Plumbum acetas, 110, 182, 268, 369.
 in cardiac hypertrophy, 288.
 in cholera, 328.
 in rheumatism, 545.
 in typhoid fever, 563.
 in yellow fever, 581.
 iodidum, 518.
Podophyllum, 78.
 in biliary calculi, 317.
 in constipation, 336.
Podophyllin, in jaundice, 422.
Polymnia uvedalia, in rheumatism, 550.
Position, in colic, 330.
 in dropsy, 455.
 in hæmaturia, 469.
 in incontinence of urine, 464.
Potassium acetas, in jaundice, 419.
 bicarbonas, in rheumatism, 545.
 bitartras, 370.
 in albuminuria, 435.
 in rheumatism, 545.
 bromidum, 19, 48, 58, 73, 78, 84, 91, 110, 155.
 in diabetes, 454.
 in typhoid fever, 563.
 chloras, 110, 182, 195, 357, 370, 422.
 in typho-malarial fever, 564.
 in yellow fever, 582.
 iodidum, 31, 45, 90, 110, 119, 153, 280, 391, 400.
 in Bright's disease, 443.
 in chronic hepatitis, 404.
 in diabetes, 454.

- Potassium in hydrargysm, 601.
 in pneumonia, 261.
 in rheumatism, 548.
 in yellow fever, 582.
 nitras, 161, 212.
 in rheumatism, 545.
 in typhus fever, 570.
 permanganas, 333.
 in rheumatism, 545.
 picronitras, in tape worm, 410.
 sulphas, 422.
 in albuminuria, 435.
 sulphuretum, 385.
 in albuminuria, 435.
 in chorea, 39.

 Poultices, 116, 183.
 Pressure on the carotids, 302.
 in gout, 501.
 on the pneumogastric, 302.
 Prophylamine, 546.
 Prunus virginiana, 254, 276.
 Pulsatilla, 195.
 Pulveris doveri, 249.
 Pyrethrum, 110.

 Quassia, 305, 385.
 for worms, 412.
 Quebracho, 157, 309.
 Quercus cortex, 357.
 Quinia, in diabetes, 454.
 in hæmaturia, 467.
 in heart disease, 295.
 in jaundice, 418, 421.
 in typhoid fever, 563.
 anti, treatment of intermittent fever, 508.
 bromidum, in diabetes, 454.
 bromohydras, 524.
 hydrochloras, 188.
 hypodermically, in intermittent fever, 514.
 salicylas, 525.
 sulphas, 61, 65, 71, 110, 142, 153, 370.
 in diabetes, 454.
 in rheumatism, 546.
 in typhus fever, 570.
 in yellow fever, 581.

 Quineti sulphas, 524.
 Quinoidine, 503.

 Rectal alimentation, 394.
 Rectal medication, 365.
 Rectal suppositories, 114.
 Resorcin, 527.
 Respirators, 201.
 elastic, 201.
 Rest, 41.
 Restraint, 87.

 Rhamnus frangula, in constipation, 345.
 Rheum, 65, 305, 353, 385.
 in anæmia, 480.
 in cholera, 322.
 in colic, 333.
 in diabetes, 454.
 Rhus aromatica, 454, 467.
 toxicodendron, in rheumatism, 550.
 Rochelle salts, 402.
 in Bright's disease, 437.
 Rosmarinus, 79.
 Rubus procumbens, in diarrhœa, 353.
 Ruschenberger's cholera mixture, 325.

 Saccharum, 254.
 Salicin, 111, 232, 527.
 in diarrhœa, 350.
 in rheumatism, 533.
 in typhoid fever, 563.
 Salines, in cholera, 320,
 in dysentery, 365.
 in seat-worms, 414.
 Saline mineral waters, in biliary calculi, 317.
 solutions, transfusion of, in anæmia, 485.
 Salutis pilulæ, in constipation, 340.
 Sanguinaria, 158, 171, 268, 423.
 in torpidity of liver, 425.
 Sanguis, 254.
 bovinus exsiccatus, 254.
 Santonin, 61.
 in chorea, 45.
 in lumbrici, 412.
 Saoria, in tape-worm, 410.
 Sapo, 386.
 Sarracenia flava, 357.
 Sassafras in rheumatism, 551.
 Scilla, 162, 182, 195.
 Scoparius, in albuminuria, 435.
 Scutellaria, 456.
 Sea voyages in phthisis, 242.
 Senega, 167, 171, 269, 309.
 Senna, 66, 386, 411, 425.
 Serpentaria, 269, 386.
 in typhoid fever, 563.
 Setons, 64, 120.
 Sevum præparatum, 254.
 Sierra salviæ, 527.
 Silphium cyrenaicum, 254.
 Sinapis, 31, 62, 93, 133, 386, 418.
 in variola, 573.
 Sleep, 87.
 Snuff for catarrh, 188.
 Sodium arsenias, 36, 111.
 benzoatum, 423.
 bromidum, 26, 54, 121.
 carbonas, 66, 111.

- Sodium carbonas in vomiting, 429.
 bicarbonas, 352, 386.
 in yellow fever, 577.
 in albuminuria, 435.
 chloridum, 62.
 in plumbism, 606.
 chlorinatum, 370.
 cholinatum, in biliary calculi,
 315.
 citras, 454.
 hypophosphis, 111, 182.
 hyposulphis, 182, 386.
 in worms, 415.
 phosphas, in biliary calculi,
 317.
 in diabetes, 455.
 salicylas, in chorea, 45.
 in diabetes, 446, 455.
 in rheumatism, 535.
 in typhoid fever, 557.
 sulphis, in dyspepsia, 374.
 in variola, 573.
 in yellow fever, 575.
 sulpho-carbolas, 492.
 Specific treatment of tuberculosis, 239.
 Spigelia, 411.
 Spiritus aetheris compositus, 162.
 Stanni chloridum, 62.
 Stigmata maidis, 467.
 Stimulants, 21, 63, 92, 111.
 in cholera, 329.
 Stimulating liniment, 173.
 Stramonium, 79, 86, 111, 152.
 in constipation, 339.
 Stropanthus, in cardiac dilatation, 288.
 Strychnia, 19, 31, 33, 37, 62, 71, 86,
 111, 116, 162.
 in apoplexy, 26.
 in constipation, 339.
 in dyspepsia, 377.
 in epilepsy, 52, 53.
 in incontinence of urine, 464.
 in milk sickness, 529.
 in plumbism, 605.
 Suet, 254.
 Sulphur, 81, 119, 129, 163, 254.
 baths, in hepatitis, 402.
 in cholera, 329.
 in rheumatism, 551.
 in seat-worms, 416.
 Sumbul radix, 84, 98.
 Suppositories, rectal, 114, 371.
 Tabacum, 62, 157, 200.
 vinum, in constipation, 346.
 Tannin, 357.
 glycerine of, 246.
 Taraxacum, 386, 400, 423.
 in chronic hepatitis, 404.
 Tela araneæ, 527.
 Terebinthinæ, oleum, 62, 79, 110, 121,
 269, 357,
 in colic, 330.
 in diarrhœa, 349.
 in round worms, 411.
 in taenia, 406.
 in typhoid fever, 558,
 563.
 in yellow fever, 582.
 stupes of, in cholera, 329.
 in dysentery, 361.
 Thermal baths, in heart disease, 279.
 Thymol, 377.
 Tiglli, oleum, 110, 119.
 Tobacco, enemata of, in intestinal ob-
 struction, 331.
 Tonic bath, 481.
 inhalation, 481.
 Toxicodendron, 120.
 Tracheotomy in laryngitis, 225.
 Travel and climate in phthisis, 241.
 Treatment of enlarged spleen, 517.
 typhoid fever, 551.
 Trephining in epilepsy, 64.
 Trichlorphenol, 371.
 Trimethylamine in rheumatism, 546.
 Tuning fork, 114.
 Unguentum antimonium, 116.
 hydrargyrum, in jaundice,
 418.
 in seat-worms, 415.
 veratria, 120.
 in diabetes, 449.
 Urea, in intermittent fever, 528.
 Urtica dioica, 463.
 Valeriana, 62, 79, 111, 145.
 in diabetes, 445.
 Vegetable bitters, 311.
 Veratria, 114.
 in diabetes, 449.
 ointment, 121.
 Veratrum viride, 31, 62, 163, 269.
 in cardiac hypertrophy, 289.
 in constipation, 346.
 in typhoid fever, 564.
 Verbascum thapsis, 254.
 Venesection, in apoplexy, 31.
 in dropsy, 456.
 in jaundice, 418.
 in pneumonia, 262.
 in uræmia, 478.
 Viburnum prunifolium, 86.
 Violæ tricoloris, in gout, 496.
 Vitis vinifera, 254.
 Warburg's tincture, 510.
 Warner's inhalation, 240.
 Water, 78, 92.

- Water, cold, in cholera, 322.
 in pneumonia, 264.
 belt, 370.
 Wild cherry bark, in cardiac hypertrophy, 289.
 Wine in anæmia, 479.
 whey, 465.
 Worm seed oil, 410.
 Xanthoxylum, in rheumatism, 551.
 Yerba santa, 180.
 Zincum bromidum, 37.
 in epilepsy, 52.
 Zincum chloridum, 221.
 carbonas, in variola, 574.
 cyanuretum, in rheumatism, 546.
 lactas, 62.
 oxidum, 19, 55, 62, 80, 163, 254, 357, 594.
 phosphas, 80, 86, 111.
 phosphidum, 214.
 in arsenicism, 598.
 sulphas, 35, 62, 146, 163.
 in colic, 333.
 valerianas, 36, 58, 69, 80, 85, 111.
 Zingiber, 149.

III. INDEX OF DISEASES.

- Abscess, hepatic, 399.
 Acidity of the stomach, (see Pyrosis).
 Acute rheumatism, 533.
 Ague, 503.
 Albuminuria, 481.
 Alcoholism, 583, 589.
 Anæmia, 479.
 of the brain, 33.
 of phthisis, 243.
 Angina pectoris, 280.
 Aphonia, 146.
 nervous, 147.
 paralytic, 148.
 Apoplexy, 22, 25, 28.
 Appetite, loss of, 311.
 Arsenicism, 597.
 Ascaris lumbricoides, 410.
 Ascares vermicularis, 413.
 Asthma, 149.
 cardiac, 155.
 catarrhal, 154.
 in children, 161.
 humid, 157.
 spasmodic, 152, 157.
 uræmic, 157.
 Autumnal catarrh, 213.
 Bladder, tonic spasm of, 465.
 Biliary calculi, 313.
 Bright's disease, 436.
 Bronchitis, acute, 164.
 of the aged, 171.
 chronic, 171.
 of the aged, 173.
 Bronchitis, gouty, 170.
 suffocative, 165.
 Calculi, biliary, 313.
 Cardiac derangements, functional, 299.
 dilatation, 285.
 disease, valvular, 303.
 hypertrophy, 288.
 Carditis, 292.
 Catarrh, 184.
 chronic senile, 173, 178.
 post-nasal, 189.
 Catarrhal jaundice, 417.
 Cerebral anæmia, 33.
 hyperæmia, 21.
 pressure, 21.
 Cerebro-spinal meningitis, 486.
 Chill, congestive, 505.
 Cholera asiatica, 317.
 Chorea, 34.
 anæmic, 35.
 dental, 35.
 electric, 36.
 from moral causes, 36.
 organic, 36.
 verminal, 35.
 Colic, 330.
 renal, 470.
 Congenital jaundice, 421.
 Congestive chill, 505.
 Constipation, 334.
 Contagious jaundice, 421.
 Convulsif tic, 133.
 Coryza, 184.

- Cramp, muscular, 136.
piano-player's, 135.
- Debility of phthisis, 243.
- Delirium tremens, 583.
- Dengue, 490.
- Diabetes insipidus, 444.
mellitus, 447.
- Diaphragm, spasm of, 135.
- Diarrhoea, acute, 346.
chronic, 354.
of phthisis, 244, 246.
- Dilatation, cardiac, 285.
- Dropsy, 455.
- Dysentery, acute, 358.
chronic, 363.
- Dyspepsia, 373.
atonic, 383, 384.
cerebral, 378.
climacteric, 379.
flatulent, 375.
irritative, 383.
- Dyspnœa of phthisis, 247.
- Dysuria, 466.
- Emphysema, 196.
- Empyema, 202.
- Endocarditis, 292.
- Enlargement of spleen, 517.
- Enteritis, 387.
- Epilepsy, 47.
- Facial neuralgia, 137.
- Fatty degeneration of heart, 298.
- Fever, break-bone, 490.
hay, 213.
hemorrhagic malarial, 500.
intermittent, 503.
pernicious malarial, 505.
relapsing, 531.
typhoid, 551.
- Functional cardiac derangements, 299.
derangement of liver, 423.
urinary disorders, 464.
- Gastric ulcer, 390.
- Gastritis, 387.
- Gastro-enteritis, 387.
- Glossitis, mercurial, 602.
- Gout, 493.
- Hæmatemesis, 397.
- Hæmatinuria, paroxysmal, 471.
- Hæmaturia, 467.
- Hæmoptysis, 206.
- Hay asthma, 213.
- Hay fever, 213.
- Headaches, 64.
bilious or dyspeptic, 65.
congestive or plethoric, 66.
- Headaches, gouty, 68.
hysterical, 69.
nervous, 72.
neuralgic, 75.
periodical, 69.
rheumatic, 76.
sick, 69.
- Heart-burn, 386.
- Heart, fatty degeneration of, 298.
functional derangements of, 299.
irritable, 302.
valvular disease of, 303.
- Hemicrania, 69.
- Hemorrhagic malarial fever, 501.
- Hepatic abscess, 399.
- Hepatitis, 399.
- Hiccough, 135.
- Hydrargysm, 600.
- Hyperæmia of the brain, 21.
- Hypertrophy, cardiac, 288.
- Hypochondriasis, 81.
- Hysterical paralysis, 118.
- Insanity, 84.
- Insomnia, 88, 90.
- Intercostal neuralgia, 106.
- Intermittent fever, 503.
- Intestinal worms, 404.
- Intoxication, 583.
- Irritable heart, 302.
- Irritation, spinal, 130.
- Jaundice, catarrhal, 417, 421.
congenital, 421.
contagious, 421.
malarial, 421.
pyæmic, 421.
simple, 420.
- Laryngitis, 219.
chronic, 221.
syphilitic, 223.
tuberculous, 223.
- Lead poisoning, 604.
- Liver, functional derangement of, 423.
torpidity of, 423.
- Locomotor ataxia, 121.
- Lumbrici, 410.
- Malarial fever, hemorrhagic, 501.
jaundice, 421.
- Melancholia, 81.
- Meningitis, cerebro-spinal, 486.
- Migraine, 69.
- Milk sickness, 528.
- Mimetic facial spasms, 133.
- Muscles, cramp of, 136.
- Nephralgia, 470.
- Nephritis, 472.

- Neuralgia, 98.
 , facial, 187.
 of the fifth nerve, 105.
 , intercostal, 106.
 Neurasthenia, 114.
 Nictitans, spasmus, 137.
 Night sweats, 247.

 Oxyuris, 413.

 Pain in cardiac region, 277.
 Palpitation of the heart, 299.
 Paralysis, 115.
 hysterical, 118.
 Paroxysmal hæmatinuria, 471.
 Pernicious malarial fever, 505.
 Pericarditis, 292.
 Phthisis, 225.
 chronic, 226.
 Piano-player's cramp, 135.
 Pleurisy, 256.
 Pleurodynia, 106.
 Plumbism, 604.
 Pneumonia, 259.
 Polydipsia, 444.
 Polyuria, 444.
 Purpura hæmorrhagica, 530.
 Pyæmic jaundice, 421.
 Pyrosis, 386, 425.

 Relapsing fever, 531.
 Renal colic, 470.
 Rheumatism, acute, 533.
 chronic, 548.
 Rose cold, 213.
 Round worms, 410.

 Sciatica, 103, 124.
 Sclerosis, spinal, 121.
 Scrivener's palsy, 134.
 Sea-sickness, 428.
 Seat worms, 413.
 Senile catarrh, chronic, 173, 178.

 Singultus, 135.
 Sleeplessness, 88, 90.
 Small pox, 570.
 Spasm of the bladder, 465.
 diaphragm, 135.
 Spasmodic diseases, 133.
 Spasmus nictitans, 137.
 Spinal irritation, 130.
 Spinal sclerosis, 121.
 Spleen, enlargement of, 517.
 Sweats, night-, 247.

 Tænia, 404.
 Tape worm, 404.
 Thread worms, 413.
 Tic convulsif, 133.
 Tic douloureux, 103, 137.
 Torpidity of liver, 423.
 Trichinæ, 416.
 Trichinosis, 416.
 Typhoid fever, 551.
 Typho-malarial fever, 564.
 Typhus fever, 565.

 Ulcer, gastric, 390.
 Uritis, mercurial, 601.
 Uræmia, 476.
 Urinary disorders, functional, 464.

 Valvular heart disease, 303.
 Variola, 570.
 Vertigo, 141.
 aural, 143.
 gastric, 143.
 of old persons, 144.
 Vomiting, 427.

 Worms, intestinal, 404.
 round, 410.
 seat, 413.
 thread, 413.

 Yellow fever, 575.

