

**Contributions to the history, diagnosis and treatment of croup / by John Ware.**

**Contributors**

Ware, John, 1795-1864.  
Royal College of Physicians of Edinburgh

**Publication/Creation**

Boston : David Clapp, printer, 1850.

**Persistent URL**

<https://wellcomecollection.org/works/n3de6kk4>

**Provider**

Royal College of Physicians Edinburgh

**License and attribution**

This material has been provided by This material has been provided by the Royal College of Physicians of Edinburgh. The original may be consulted at the Royal College of Physicians of Edinburgh. where the originals may be consulted.

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>



CONTRIBUTIONS

TO THE

HISTORY, DIAGNOSIS AND TREATMENT

OF

# C R O U P .

BY JOHN WARE, M. D.

---

From the Boston Medical and Surgical Journal.

---

B O S T O N :

DAVID CLAPP, PRINTER....184 WASHINGTON STREET.

1850.

THE following papers were originally communicated to the Boston Society for Medical Improvement, and to the Suffolk District Medical Society. The first of them was published some years since in the New England Medical and Surgical Journal, but, as it is closely connected with the subsequent ones, it seemed desirable to reprint it with them. The whole substance of these papers might have been easily condensed and presented in the form of a single essay. As they were, however, prepared at different times, and in the course of a continued series of observation and inquiry, I preferred offering them to the profession in the form originally given to them.

R55270

## ON CROUP.

---

I.—*Contributions to the History and Diagnosis of Croup.*—Read before the Boston Society for Medical Improvement, in 1842.

EVERY physician who has much practical acquaintance with disease, will have observed that there are great differences of character among the cases to which he finds it convenient, in accordance with the custom of medical men, to give the general name of *croup*. He finds that a certain portion of these cases—and by far the larger portion—yield readily to the means which he employs, and very often to the ordinary domestic remedies of mothers and nurses. He has indeed reason to believe that a considerable number of them would spontaneously subside if left to themselves. On the other hand, he finds that there are some cases, fortunately but few in proportion to the whole, which exhibit throughout their course a character of obstinacy that bids defiance to treatment, and which, with few exceptions, pass on to a fatal termination uninfluenced by any remedies he can employ.

Different views may be taken of the nature of these cases. It is believed by some that the former are not, for the most part, essentially different from the latter; that the difference is more in degree than in kind, or that the difference in the severity and result depends on difference of management; that the favorable character and course of the former are mainly owing to early and judicious treatment, and the fatal event of the latter to the inefficient or too tardy application of remedies. A long, and I trust a faithful examination of this disease has, however, satisfied me that this opinion is not correct. I have been led to believe that there is an original and essential difference in these cases; that those of the first kind are pathologically different from the second; that the former, even if they terminate fatally, which happens in some rare instances, do not terminate in the same way, or at least do not

exhibit the same morbid conditions ; and that no variety or deficiency of treatment will cause a case of the one kind to assume the character of the other.

I do not, however, mean to imply that all the cases to which I refer, are capable of being classed under two varieties. Among those which I have characterized as the more mild and tractable sort, we still find great differences in the mode of attack, course, and mode of termination, and also in the degree in which they appear to be influenced by remedies. The object of this paper is to endeavor to contribute something towards determining the nature and extent of the distinctions referred to. With this view I have made an examination of all the cases of croup of every kind which have occurred during the last twelve and a half years, in my own practice, and of this examination I now submit the results. Upon certain points relating to the severer form of the disease, I have included the examination of a number of other cases, extending over a period of twenty-five years, witnessed partly in my own practice, partly at dissections, and partly in consultations.

It should be first observed, that, in noting cases in order to this inquiry, I have set down as croup, all those which in the common language of the profession are included under this name—viz., all those which, at any stage of their progress, present a fair question of diagnosis ; all those in which is heard that shrill, sharp, ringing cough, which is regarded as the cough of croup, accompanied by a distinct embarrassment of respiration, however slight, and by some affection of the voice. It follows, of course, that many very slight cases must have been included among those on which these remarks are founded—cases which yielded or subsided almost at once. Yet it is right that these should form part of the materials of our examination. When we are in search of means of diagnosis, our attention should be directed to all those cases which have, at any period of their progress, exhibited symptoms that give rise to a well-grounded suspicion of their character. Although many cases which excite the apprehension of severe croup on their first attack, pass away very readily, and by their result show themselves to have been of very moderate severity ; yet, on the other hand, it is to be recollected, that many cases, which at last terminate fatally, do not, at their beginning, exhibit symptoms at all more severe, or excite apprehensions at all more serious, than those which have so readily subsided.

Of the cases to which this inquiry relates, occurring during the period extending from January, 1830, to July, 1842, the number is 131. For the convenience of examination, these may be divided into four classes.

I do not intend by this arrangement to express the opinion that they constitute four distinct diseases. I would not even be understood to assert positively, with our present amount of knowledge, that they are not different manifestations of the same disease. The purpose now is to speak of them as groups of cases distinguished by certain differences in their symptoms and course, which may or may not be connected with an essential difference in their nature. These classes may be designated, with a view to their probable character and for the purpose of referring to them more intelligibly, by the terms membranous, inflammatory, spasmodic and catarrhal. Of the whole number there were :—

	Cases.	Deaths.
Of membranous Croup,	22	19
Inflammatory “	18	0
Spasmodic “	35	0
Catarrhal “	56	0
	<hr/> 131	<hr/> 19

In the first class are included those cases in which there is reason to believe that a false membrane has been actually formed lining the larynx and trachea.

In the second class, those cases in which the symptoms are for the most part of the same character as in the first, but in which there is reason to believe that no membrane has been formed. The grounds for the opinion formed of the nature of these two classes will be stated subsequently.

The terms applied to the third and fourth classes, require no particular explanation.

The symptoms on which we depend for the diagnosis of croup, relate to the cough, the voice and the respiration.

In the early stage of the first form of croup, the cough is by no means peculiar. In the advanced, it assumes a somewhat different character. In the early period it is sharp, shrill, ringing; it does not vary from that which we hear in the other forms, except perhaps that in some of the less formidable cases it is much louder and more violent at their beginning, than it is in those which prove ultimately more alarming. In the latter period it becomes less loud and ringing, but is equally sharp—it often becomes almost inaudible, bearing the same relation to a common cough, that a whisper does to the common voice. The cough, then, affords no certain means of distinguishing this form of croup at that period of it in which the diagnosis would be most valuable.

Of the state of the voice, nearly the same remark may be made. In the advanced stage of a case it is sufficiently characteristic. It becomes

a sharp, and almost inaudible whisper. But early in the disease it is not always affected at all; and, if it be, cannot with certainty be distinguished from the hoarse voice of common catarrh.

The condition of the respiration affords us far more important information. In the early period of the disease, however, when we most need means of diagnosis, it is not a symptom which always attracts attention, even from the physician; much less from others who are around the patient. The common description of the breathing in croup, does not apply well to the beginning of the membranous variety. It seems rather taken from cases of a less dangerous kind, in which the breathing is from the first, loud, harsh, suffocative; attended with great efforts, and much loud coughing; creating great alarm, and calling at once for efficient means of relief. But the breathing in membranous croup does not excite attention in the very commencement of the disease. It is comparatively quiet and unobtrusive. Its true character is not at once to be detected, but only by a careful and accurate observation. The patient has not the ordinary aspect of difficult breathing; in fact, the breathing is not difficult at the very first. He probably experiences no distress. There is no real deficiency in the performance of the function, and no obvious embarrassment. There is only a little more effort in drawing in the air, and a little more force exercised in its expulsion, whilst the amount of air admitted and expelled is fully equal to the necessities of life. This perhaps would not be noticed on a casual glance at the patient, but will be at once perceived on attending to the muscular movements subservient to the function, which are—to use an expressive French term—somewhat exalted. It is indicated very soon, also, by a slight dilatation of the nostrils, and a little whiz or buzz accompanying the passage of air through the rima glottidis. This sound is distinguished either by placing the ear near the mouth of the patient, or by applying the stethoscope on the back of the neck, or directly upon the upper part of the larynx.

This at its very beginning is the essential respiration of membranous croup, and it affords far more aid in diagnosis than either the cough or the voice. It is not, however, always found as pure as has been described. It is often mingled with, and obscured by, other sounds. Thus the disease is often attended by paroxysms of irregular and spasmodic breathing, accompanied by violent muscular efforts and great distress, and of course producing other and more obvious sounds than those described. There is often also present in the air passages, either above or below the glottis, a quantity of mucus, giving rise to a constant or occasional rattling, which seems to mask the proper sound of croup. These adventitious sounds,

being also as frequently heard in the other forms of croup, are therefore of no service in diagnosis. Generally there are intervals of relief from these superadded symptoms, especially immediately after vomiting or bleeding, but the essential breathing of the disease will be found to be unchanged and unmitigated in these intervals of ease; although the apparent relief may be so considerable as to give rise to strong, but fallacious hopes of recovery.

We occasionally hear, in cases of considerable enlargement of the tonsils, a kind of breathing which closely resembles the early breathing of croup. Usually in such patients the respiration is loud, sonorous, unequal and irregular, but in a few it is quiet, steady, with a muscular effort occasioned by a mechanical obstruction like that in croup. The distinction between them can, however, be readily made, by attending carefully to the seat of the obstruction, which is above the rima glottidis in the one case, and at it in the other; by the sound of the cough and voice, which are not croupy, and by the fact that the obstruction varies in degree and sometimes vanishes, with change of position.

I have endeavored to describe this respiration as it exists in its slightest appreciable degree, at the earliest period of its manifestation. As the disease advances, it becomes very strongly marked, whilst the condition on which its peculiar character depends, viz. a mechanical narrowing of the orifice through which the air passes, becomes much more obvious.

The muscular effort, in the latter stage, becomes very strong, both in inspiration and expiration. During inspiration, whilst all the muscles concerned in it are in the highest state of activity, the mechanical impediment against which they act is often strikingly displayed by the falling in of the soft parts about the neck and clavicles, at the epigastrium, and between and along the lower edge of the ribs—the air not passing in through the narrowed opening of the glottis so rapidly as the dilatation of the chest by the increased muscular effort would render necessary. The expiration is chiefly characterized by the amount of force employed to expel the air. In health the expiration is easy, and accompanied by little effort. Where there is no unusual obstruction, the mere tendency to collapse of the lungs would be sufficient for the expulsion of the air, as we see in the dead body; so that the walls of the chest have merely to follow up this contraction, without adding to its force by any muscular effort. But in croup, this is not enough; and we often find that the air is blown out forcibly against the mechanical resistance occasioned by the disease. We find the same strong contraction of the muscles concerned, especially of the abdominal muscles, which is observed when air is blown out forcibly through a narrow passage.



This is the proper breathing of croup; becoming more and more intense as the disease approaches its termination, till the whole life of the individual seems, as it were, to concentrate itself in this one effort. The patient in this extreme condition seeks, by a multitude of changes of place and position, to find some alleviation of his agony; the cough, and with it the voice, have become nearly extinct; and his inarticulate appeals and beseeching looks for relief to those from whom he is accustomed to look for it, constitute one of the most touching scenes which we are called upon to witness in the practice of medicine. Happily the extreme suffering usually, though not always, subsides towards the close of life, and death takes place at last with comparative ease.

In the advanced stage of croup, the breathing is often modified by circumstances other than the mere mechanical obstruction at the upper part of the larynx. After a certain period the false membrane is in some places separated from its adhesion to the mucous surface, by the secretion of pus. The passage of air to and fro, and the efforts of coughing, detach it partially from its adhesion, and break it up more or less into shreds, which however still adhere at one of their ends. These ragged portions of membrane, mingled with the pus, move up and down the air passages, causing some variety in the sounds and also in the actual difficulty of breathing. Death is sometimes very suddenly produced by a collection of this material into a mass which becomes impacted in, and thus plugs up, either the upper or lower part of the larynx. This at least, from the state in which the parts are found on dissection, would appear to be the mode in which death takes place.

The respiration may also be modified in croup from a congestion or inflammation of the lungs, which occasionally supervenes. The embarrassment of respiration has also sometimes appeared to be increased by an accumulation of air in the lungs, which arises from a deficient balance between inspiration and expiration. Owing to the greater ease with which we can make extraordinary and continued efforts of inspiration than we can of expiration, a greater quantity is admitted than can be readily expelled, before the suffocative feeling of the patient impels him to a new effort for relief.

But although there may be a combination of the respiration of this disease with that produced by other affections of the throat or lungs, yet the respiration of croup is in its nature and character essentially distinct from them. In them the difficulty of breathing and the unusual muscular effort may arise from a variety of causes, producing great varieties in the modes of dyspnoea; in croup the one essential condition is the mechani-

cally contracted state of the passage through which the air passes, and all the peculiarities of the dyspnœa proceed from this condition. In one particular the breathing of asthma resembles that of croup, viz., in the intensity of the effort by which the current of air is made to move in both directions against a mechanical resistance ; but the point of the resistance and consequently the other circumstances of the function prevent the resemblance from extending to other points.

The *first* form of croup, then, is distinguished by the cough, the voice, and by a peculiarity of the respiration, which I have attempted to describe, and which, for the sake of distinguishing it in this essay, may be called *intense*.

In the cases of the inflammatory croup, which constitute the *second* form of the disease, the condition of the voice, cough and breathing are precisely the same as in the cases of the first class. There is no certain way by which, so far as these symptoms are concerned, cases of the one kind are to be distinguished from those of the other. The cases enumerated among the second class were of all degrees of severity, but none of them were fatal. Cases, however, of croup which terminated fatally, and in which no membrane was found on dissection, are recorded upon the best authority. To these we shall have occasion to advert hereafter. In addition to the symptoms proceeding from the character of the cough, voice and respiration, I have noted, in a few examples of this form of the disease, a tenderness of the larynx on pressure.

As cases of this class are then usually favorable in their termination, whilst those of the first are usually fatal, the diagnosis between them, in the early stages especially, becomes of very great importance, both as regards prognosis and treatment. Of the means by which this distinction may probably be made, and of the grounds for believing these two to be essentially distinct diseases, and not different states or conditions of the same disease, I shall take occasion to speak, after considering the other two classes which have been enumerated.

The *third* includes certain cases which are generally designated as *spasmodic croup*, and sometimes as *spasmodic asthma*. The attack is always sudden, and usually occurs after the subject has been, for some time, asleep. Very often it occurs in the evening, during the first sleep of the child, before its parents have retired to bed ; but perhaps as frequently at a later hour of the night, or very early in the morning. The patient wakes in great distress for breath. His inspiration is attended with great effort ; it is loud, ringing, shrill, somewhat resembling the hooping inspiration of hooping cough, but louder and more sonorous. The

expiration is comparatively quiet and easy. The voice, at the same time, is hoarse and broken, and there is a loud, hoarse, barking cough, which closely resembles that of the preceding kinds, and indeed alone, would not serve as a mark of distinction from them. These cases seem occasionally to rise from indigestion; but more frequently we can trace their occurrence to cold, especially as they have been often preceded for a few days by symptoms of catarrh. When left to themselves, they will usually subside spontaneously, but from their suddenness and violence, they cause great alarm, and call for immediate assistance. They rarely fail to yield to an emetic or venesection, leaving behind them for a longer or shorter period, rarely for more than twenty-four hours, some hoarseness and some degree of the croupy sound of the cough, with a little huskiness or stuffiness of breathing. At no period is there any proper *intensity* of respiration.

These cases, from their suddenness, the time of the attack, the great violence of the first symptoms, and the consequent alarm which they create, produce a stronger impression on the minds of common observers and even of many practitioners, than those of the other kinds. This mode of attack is most closely associated in their minds with the term croup; and it is regarded as tending, if not checked, to terminate in the same state of things with cases of the first class. So far as the cases before us are concerned, however, this never happens, and of the whole number included under this examination, no one proved fatal.

The fourth class includes cases not falling under either of the above, and yet frequently presenting a very close resemblance to them. The subjects usually exhibit at first the symptoms of common catarrh. After a few days the voice becomes hoarse; the cough becomes croupy, and there is tightness, oppression, and some approach to the croupy sound of respiration; there is, however, no intense or exalted action of the respiratory muscles, and no indication of that mechanical impediment to the current of air which exists at the rima glottidis in the two first forms of the disease. Still the resemblance is sometimes quite close enough to cases of these forms, in their earliest stage, to occasion some anxiety, and there is also sometimes a sudden attack of dyspnoea, with loud, shrill and sonorous breathing, which imitates the symptoms of the third form, and is perhaps to be regarded as an attack of the same kind.

The cases of this form yield gradually, the croupy character wearing off in a few days, and leaving behind simply catarrhal symptoms. I suppose them, from the mode in which they come on and go off, to be properly a catarrhal inflammation of the mucous membrane covering the organs of

voice. We frequently observe that the catarrhal affection of the same membrane which occurs in the first stage of measles, is accompanied by the same croupy symptoms as those which have been now described—going off with the other catarrhal symptoms. In a few instances the attacks of this form of croup have terminated in severe bronchitis, or in inflammation of the lungs themselves. But among the 56 cases included above, there was no one fatal.

Having thus described these several forms of this disease, and stated in general what seemed to be their nature, the question now arises as to the justice of the distinction which has thus been assumed to exist. Is there any sufficient ground for such a distinction? Are these different cases different diseases? Are not the favorable ones, which constitute so large a proportion of the whole number, similar in their nature to the more severe; but either of less severity in their origin, or else modified and controlled in their course, by the influence of treatment. These questions it is obviously of great importance, to the prognosis and treatment of the cases in question, to be able to answer correctly. If we can with regard to a large proportion of them confidently predict from the outset a favorable issue, the practitioner and the friends will be saved much unnecessary anxiety, and the patient many annoying and debilitating remedies.

I proceed, therefore, to state the grounds for a belief that the first form of croup is a disease essentially distinct from all the others, and that it depends on a peculiar pathological condition to which they have no tendency. Whether there be any equally marked distinction between the other forms, it is not of the same practical importance to determine; and as we have no sufficient materials for a satisfactory inquiry into this question, our attention will be confined to the evidence for the distinct character of the first form.

Every physician is familiar with an affection of the throat, both in adults and children, consisting in an inflammation of the mucous membrane, of that peculiar character which produces the effusion of a layer of coagulable lymph, or false membrane. The connection of this affection of the throat with croup was long since pointed out; and it is well known to practitioners among us, that this complaint, known familiarly, though inaccurately, under the name of "ulcerated sore throat," often accompanies or is followed by croup, and that croup thus connected is peculiarly fatal in its character. This circumstance in the history of croup was many years since strongly impressed upon my mind by an eminent practitioner in this neighborhood.\* I was in consequence led, in all cases of

---

\* Dr. William J. Walker.

croup, subsequently to this period, to make a careful examination of the fauces, with the view of determining exactly the extent to which this visible affection of the throat was connected with the more important disease.

Two causes prevent the completeness of these observations. We are very apt, in making record of cases, especially of those which appear of a slight degree of severity, to omit the *noting* of negative facts, even when they have been actually the objects of attention. Hence, although I have very rarely failed to examine the fauces in any case of supposed croup, I have often in the lighter cases, and sometimes in the severer, failed to note their condition. The second cause of incompleteness is the impossibility in some patients, from their terror and consequent resistance, of getting such a view of the parts as would authorize us to pronounce decidedly what their state is. Notwithstanding these circumstances, the state of the throat has been noticed and recorded in a sufficient number of cases to afford very fair materials for inference.

With a view to this examination, I may include a considerable number of other cases, besides those which constitute the particular subjects of inquiry in this paper, which have been noticed at other times, or in the practice of my friends. Including these cases with the 22 above referred to, I have memoranda, more or less complete, of 39 cases of what I have denominated membranous croup. The state of the fauces was observed and noted in 33, and of these, in 32 a false membrane was present; most frequently, and sometimes only on the tonsils, sometimes on other parts also, as the palate, uvula and pharynx. In one case no such membrane was present; but it was found to exist in the larynx after death. In 3 of these 33 cases, recovery took place; all the others were fatal. In 14, an examination was made after death, and the usual appearances were found to exist in all of them.

On the other hand, I have memoranda of 109 cases of what I have classed as the other forms of croup, and of these the state of the tonsils and fauces was noted in 45. In no one was there such a condition of the parts as was found to exist in the membranous form. In 3 cases there was indeed a thin, slight exudation on the tonsils, of the color and appearance of starch, like that which is sometimes seen on the edges and surface of the tongue. This I apprehend to be a formation of an entirely different nature from that which exists in the other class of cases. Of the 45, 12 were of the second, 11 of the third, and 22 of the fourth class.

From this statement, it seems probable that the appearance of a false membrane upon the tonsils or other visible part of the throat, in a case of

croup, may be regarded as a pretty certain diagnostic sign that it is the membranous form of the disease; and its absence as a pretty certain indication that it is one of the other forms. Still there will be exceptions. There will be cases in which the membrane is formed in the larynx, although it has not appeared in the throat; and there may be those in which a membrane exists in the throat, unaccompanied by a similar condition of the air passages. Of the former I have recorded one example; of the latter, none. How frequent such exceptions will be, must be determined by more extensive observation. If they are not more frequent than they have been among the cases here recorded, the observation of this symptom will afford a sufficiently safe guide, since of 75 cases in which it was looked for and the result noticed, it failed as a diagnostic sign in but a single instance.

The question now presents itself, what are the grounds for believing that the two forms of the disease which I have distinguished as membranous and inflammatory, are not the same in different degrees or in different stages? and may not pass one into the other? The grounds are—

1. The very great preponderance of fatal results in the membranous croup and a similar preponderance of recoveries in the inflammatory, and the evidence which exists that in the few cases of recovery from the former, the membrane has been formed, and in the few cases on record of death from the latter, that a membrane has not been formed—afford strong reason for believing that the diseases are essentially different.\*

---

\* No fatal cases having occurred of inflammatory croup under my own notice, I am happy to be able to avail myself, in support of the views above taken, of an account of four such cases, contained in the first volume of the *New England Journal of Medicine and Surgery*, by James Jackson, M.D., formerly Professor of Theory and Practice of Physic in Harvard University. The symptoms in all these cases were unquestionably those of croup. In one of them bronchotomy was performed.

In the first case, "the mucous membrane of the larynx was much inflamed, and smeared over with a quantity of loose mucus, but without any false membrane. The inflammation extended into the trachea as far as could be examined without opening the chest."

In the second case, "the appearances in the larynx were the same. The lungs were more full of blood than usual."

In the third case, "there was not any coagulable lymph, the mucous membrane was highly inflamed and swollen, and the rima glottidis was thus very much narrowed. The membrane was smeared over with a thick mucus."

The fourth case I give at length in the words of the author.

"I was called to this on Sunday, July 5, 1812, at 3 o'clock, P.M. The disease had commenced 20 hours before, and was very strongly marked. The symptoms were considerably mitigated after vomiting. I tried in vain to take blood; the child was very fat, and the veins were all hidden, even the external jugular. The respiration grew bad again before morning, but the patient lived till the next morning, the 7th, so that the disease continued two days and a half, or 60 hours. In 8 hours after death, Dr. Bigelow examined the body, and the following is his report of the appearances. 'The trachea with the larynx was removed. The whole tube was pervious as usual, excepting the presence of a large quantity of mucus of the ordinary consistence. On dividing the

2. The formation of a false membrane does not seem to require either an advanced stage or a very intense degree of the inflammation from which it proceeds. It is rather the result of a peculiarity in the kind of inflammation, than of any period or degree of it. It appears to be a very early product of the inflammation, if it be not indeed almost contemporaneous with it. It resembles in this respect the similar effusion taking place on the serous membranes, which in them occurs very early, and has even been supposed to be the first act of inflammation. In the common inflammation of the tonsils which is accompanied by this symptom, a layer of lymph is observed to be effused over the surface of the part as soon as any signs of disease exist.

3. The circumstances attending recovery from simple inflammatory croup differ materially from those which accompany recovery from membranous croup. In the former the amendment is rapid and speedily complete. There is left behind only a moderate soreness of the larynx, and, in the worst cases, some hoarseness. There is at no time any copious or solid expectoration. In the latter, recovery is slow, unequal, and accompanied by phenomena which must necessarily attend the separation of the membrane, and the process through which the diseased mucous surface must go in order to its restoration to a healthy condition. The natural cure of the disease takes place by the occurrence of the suppurative inflammation upon the diseased surface, by which the false membrane is thrown off, and the mucous membrane then gradually returns to its natural state. In examinations after death, we usually find that this process has begun in the trachea, the membrane being there separated and often broken up into shreds, whilst the inflamed surface is covered by a layer of pus. Above, in the upper part of the larynx, around the glottis, the false membrane usually remains closely adherent. It is obvious that recovery might always take place could the parts be spared

---

larynx and trachea at the posterior side, and exposing the internal surface, the mucus being removed, a number of distinct red spots were discovered, of considerable size, on the lining membrane. One of these was immediately below the glottis. Between the mucus and the lining membrane there was no factitious substance whatever, nor any appearance the least resembling the membranes which I have seen formed in some other cases of croup. The lungs were not examined."

"In the other cases I had thought it possible that the disease had not continued long enough to allow the effusion to take place, as the patients all died in less than 48 hours from the attack. But in this last case such a supposition cannot be admitted; for I have in my possession a preparation in which the false membrane is exhibited in great perfection, and this came from a patient of Dr. Channing which I had seen with him, and in which death had occurred in about 30 hours after the seizure."

The history of these cases, especially with the authority upon which they are recorded, affords very satisfactory evidence of the existence of a class of cases like those which have been above described, of a disease with the symptoms of croup, but without the formation of a false membrane either in the air passages or upon the visible parts of the throat.

long enough from their functions to go through the necessary steps—and it is also obvious when it does take place, that it must be accompanied by a copious expectoration of pus, and of the membrane either in pieces, if firm enough, or else broken up and partially dissolved by the pus. Now these appearances do not accompany recovery from even the severest cases of the inflammatory croup, whilst they do accompany recovery from well-marked cases of the membranous form.

Of the three cases of membranous croup which are noted as having recovered, there are but two of which I have such an account as would justify me in presenting them as fair examples of the processes through which the parts pass in recovery. These were both of the most decided character, and had arrived at that stage of the disease in which we expect a fatal event to occur almost from hour to hour. In the first of them, six days elapsed before any sensible mitigation of the symptoms, and even then the progress to recovery was very slow and apparently doubtful. Improvement was attended by a copious muco-purulent expectoration, in which it is true no large pieces of membrane were ever detected, but of such a consistence and appearance as would favor the belief that the membrane had escaped in a comminuted or partially dissolved state. After the probable removal of the membrane, there was for some days a bloody expectoration, the voice did not return, and it was indeed many weeks before it resumed its natural tone.

In the second case, a considerable portion of the membrane was spit up in a tubular form, after a violent fit of suffocative cough, and this was followed by the rejection of smaller pieces, mixed with a muco-purulent, at first, and then a bloody expectoration. There continued an entire loss of voice for more than a week, and for at least ten weeks after recovery it had not regained its natural tones.

The contrast is very striking between the protracted character of these recoveries, and the speedy return to health of all those who labored only under the other forms of the disease, however severe.

The observations to which the preceding remarks relate, were all made in this city and its immediate neighborhood; how far they correspond to the disease as it appears in other places, must be left to others to judge. So far as they go, they appear to me to justify the following conclusions.

1. That the only form of croup attended with any considerable danger to life, is that which is distinguished by the presence of a false membrane in the air passages.

2. That the existence of this membrane in the air passages is in a very large proportion of instances indicated by the existence of a similar membrane in the visible parts of the throat.



3. That this affection differs not in stage or degree, but in kind, from all the other cases which are commonly known by the same name, and that the latter have no tendency to become converted into or to terminate in the former.

As my intention has not been to write a complete history of croup, I have omitted all such notices of the symptoms, cause, morbid anatomy, &c. of the disease as have no direct bearing on that point in its character which it was my desire to illustrate. It may not be amiss, however, to record, in connection with this paper, a few circumstances with regard to its history, which have been incidentally determined from an examination of the cases before us.

Croup is often regarded as a disease which attacks suddenly and violently. This is only true of the milder forms. Genuine or membranous croup is commonly rather gradual in its approach, and consequently often insidious. It supervenes often on the common sore throat of children; and in such cases, though its development is frequently rapid and apparently sudden, yet a careful examination of the past history of such a case will generally satisfy us, that although it may have had a sudden outbreak of violence at the time it was supposed to begin, yet that it had really been coming on for several days. Of 30 cases in which I have had an opportunity of determining the mode of attack, in only two could it in any proper sense be called sudden, although in many, the attention of friends was called to it quite unexpectedly, by a rapid increase in the violence of the symptoms. A sudden and violent attack is, therefore, to be regarded as affording a favorable indication of the character of the case in which it occurs. The unexpected manner in which croup sometimes steals upon the common sore throat of children, should lead always to the careful inspection and watching of such cases. It is true that but a very small proportion of them do terminate in this way; but as it is the only considerable source of danger, and the only way in which they are likely to have a fatal termination, the possibility of such a course of things should not be overlooked. No case of this kind can be regarded as entirely safe from such a result. The danger is even not confined to childhood. Two of the above-named cases of fatal croup occurred in females of 12 years of age, in which it had supervened on this affection of the throat.

The membranous croup also sometimes occurs as a sequel to the affection of the throat in scarlatina. The most common primary affection of the throat in this disease, is of the same kind with that denominated the ulcerated sore throat, viz., an inflammation, with an effusion of false membrane upon the parts inflamed. When croup supervenes upon this,

the case is usually very rapid and inevitably fatal. Of the cases above enumerated, two were of this character. A third occurred to me, not enumerated among them, in which there were no symptoms of croup during life, the patient apparently dying from affection of the brain, but in which the usual appearances of croup were found after death. The subject of this was a young man 17 years of age. These cases all occurred between eight and ten years since. None have been observed during the more recent periods of the prevalence of scarlatina.

Croup varies considerably in its duration ; I mean its duration after its characteristic symptoms are fairly developed and there is reason to believe that the membrane is formed. Of 23 cases,

	1	continued 1 day from distinct croupy symptoms.
6	"	2 to 2½
9	"	3 to 3½
3	"	4
1	"	5
1	"	9
1	"	11
1	"	19

Nineteen cases, or more than three-fourths, therefore, were of four days duration or less.

Croup, in this form, rarely attacks children under two years of age. Of 30 deaths and 3 recoveries, of which the ages were known,

Deaths.	Recoveries.		
1	0	took place at	12 months.
1	0	"	18 do.
5	0	"	2 to 2½ years.
3	0	"	3 to 3½ do.
8	0	"	4
6	1	"	5
2	0	"	6
0	2	"	7
1	0	"	8
2	0	"	12
1	0	"	17

Twenty-two, or two-thirds of the cases, occurring between the ages of 2 and 5.

It will be seen, by the following statement of the ages of 95 patients affected with croup of the other varieties, that the tendency to the disease in them exists at a much earlier age. Whilst but 1 case in 16 occurred under two years of age in the first class of cases, 23 out of 95, or about 1 in 4, happened under the same period among the others.

Age.	Second form.	Third.	Fourth.	Total.
Under 1	1	1	3	5
1 to 2	2	5	11	18
2 to 3	6	6	11	23
3 to 4	2	3	5	15
4 to 5	1	6	3	10
5 &c.	1	3	3	7
6	1	1	2	4
7	1	2	5	8
8	0	0	2	2
9	0	1	0	1
10	1	0	0	1
11	0	1	0	1
	16	34	45	95

In cases of the first kind, the tendency to the disease seems to be about equal in the two sexes. Of the 22 cases embraced in this inquiry, the number of each sex was precisely the same—11 males and 11 females. Adding to them 12 other cases in which the sex is noted, we still have numbers too nearly equal to indicate any peculiar tendency to the disease in either sex, viz., 16 males and 18 females.

In cases of the other forms of croup, the difference seems too great and too uniform to be merely accidental. In

18	cases of 2d class,	11	males,	7	females.
35	" 3d "	25	" 10	" "	"
56	" 4th "	33	" 23	" "	"
		69		40	

As it is of some interest to observe the degree of influence which season is capable of having on disease, I subjoin a table containing a statement of the numbers of the cases referred to in this paper, occurring in the different months. As the number of cases, however, is too small of the first class to afford any very satisfactory result, I have added in another column the number of deaths from croup, occurring in the several months, out of 263 cases, drawn from the bills of mortality for this city. I have made the table to begin with November, for the sake of comparing more easily the cases and deaths of the colder half with those of the warmer half of the year.

	Membranous Croup.	Other Forms.	Deaths from Croup.
November	2	11	31
December	2	11	31
January	9	14	31
February	2	18	22
March	2	9	33
April	2—19	17—30	18—166
May	4	9	14
June	3	6	11
July	1	3	13
August	2	0	13
September	2	5	16
October	3—15	7—30	30—97

I should observe that the several years vary very much in the amount of mortality of the several months. Thus in the month of January, there was in one year 13 deaths, in another year only 1, and a similar though less remarkable inequality in other months. Still the results are upon the whole too uniform to leave any doubt of the greater tendency to these diseases in certain periods of the year.

II.—*Treatment of Croup.* Read before the Boston Society for Medical Improvement, Nov. 11, 1844.

The history of the case of croup reported at the last meeting,\* which I had an opportunity of witnessing during its progress, has confirmed me in an opinion I have, for some time, been disposed to entertain, that the methods of treating this disease in common use require a careful re-consideration. This opinion is connected with, or perhaps has proceeded from, certain views concerning the distinctive character of various forms of disease which ordinarily are included under this one common appellation, and which I have formerly communicated to the Society. It is not too much to say, that the received mode of treating these cases, which, so far as I know, is very much the same for all their varieties, has come down to us by a sort of tradition from our predecessors in the profession, and varies but little, if at all, from that which was originally adopted when the disease first became the object of attention. It is true that in single cases and by particular individuals, there have been occasional variations from the established practice; still, in the main, emetics and bleeding, blisters and calomel, have been the principal remedies. The depleting, reducing and perturbing method is that on which dependence has been chiefly placed.

That this treatment may be applicable to a very considerable proportion of the cases which pass under the common denomination of croup, I am not prepared to deny. Those which in a preceding communication have been classed as inflammatory, spasmodic and catarrhal, certainly recover under its influence, and apparently with a greater speed than if left entirely to the resources of nature. So far as my experience has gone, however, it has appeared to produce no impression upon those in which there is satisfactory evidence that a membrane has been formed.

These cases, I should repeat the opinion expressed in the paper just

---

\* This was the case of a child with membranous croup, communicated by my brother Dr. Charles Ware, of this city, in which the anodyne treatment was mainly employed, and in which the membrane was separated and thrown off. Everything promised favorably for recovery so far as croup was concerned, but the patient died ultimately by the rapid supervention of inflammation of the lungs.

referred to, are essentially of a distinct nature from the others, and constitute but a small proportion of those which are usually regarded as croup. They are not aggravated cases of the same kind as the others—cases which have gone on to an ulterior stage of disease—but in their origin and conception different. The inflammation which is essential to them is peculiar in its character; the effusion of false membrane is not the result of an advanced stage of it, but is one of its early results—is perhaps the first visible act of its existence; as there is much reason to believe that it is of serous membranes. It has been common to describe the stage of effusion, in croup, as preceded by one of longer or shorter duration—a formative stage. If I am right in the views taken of the character of the disease, this distinction is made by making up its history from different sets of cases—going to one for the history of the first stage, and to another for the history of the second. The same confusion of diagnosis has given also an apparent success to means used for treatment. Where all the different cases which have been referred to, are grouped together as examples of the same disease in different stages or degrees, the proportion of recoveries will not appear discouragingly small. If we were to class together, as cases of consumption, all those in which there was cough and expectoration, as is done by those who profess to cure this malady, we should have no reason to be disheartened with regard to its curability; and, in the same way, so long as we class all cases together as croup which have a croupy cough and some difficulty of breathing, the amount of mortality will not be greater than in other acute diseases of children. A more accurate diagnosis will, I am convinced, put an end to our complacency on this point. Membranous croup unquestionably does sometimes come to a favorable termination; but recovery is comparatively so rare, it forms so much the exception, that, admitting the distinctive character of the disease, it is difficult to conceive that the treatment has anything to do with the recovery. Where, under any given method of treatment, but one case out of six or eight recovers, one must be very sanguine indeed to attribute much influence upon the result to the remedies.

The question then properly arises—if the mode of treating croup commonly adopted does no good, are we sure that it does no hurt? This is a question we are far too unwilling to put to ourselves. What will happen if nothing be done? This should always be the first thought of the physician, in each individual case. Till he knows this, he cannot know with certainty what effect his treatment has; and just in proportion to the amount of his knowledge of the natural history of disease, and of the time and mode of its natural termination in recovery or

death, will be his power of judging of the influence of treatment upon the result.

Now when we examine the cases of recovery of membranous croup which actually take place, and compare them with the condition of the parts in those which are examined after death, we find very clear evidences of a tendency in the disease to go through a certain course of changes which will terminate in health. The false membrane is effused, and, at the same time, the mucous membrane is thickened and congested. After a time, a process of suppuration is established upon the surface of the mucous membrane, underneath the false membrane, which of course separates the latter from the former, so that it lies loosely upon it, whilst between them is a layer of pus. If the membrane thus thrown off be thick and strong, it is expectorated in distinct pieces, sometimes of a considerable size; if it be thin and less firm, it is either converted partially into pus, or else is broken up into smaller shreds and mixed with the pus so as not to be distinguished from it, except by very careful examination, and thus it is all gradually thrown up. The diseased membrane does not free itself from the false membrane over its whole surface at once. Those portions from which the false membrane has separated are left in an inflamed and irritable state—the expectorated membrane and pus are often tinged with blood, probably from the fact that by the violent effort of coughing some portions are torn off from the mucous surface before the purulent process has effected a complete separation. The cough, then, with more or less expectoration, and a hoarseness, in some cases amounting to an incapacity for speaking except in a whisper, continue for some time—the affection of the voice for several weeks. The parts are at length, however, perfectly restored.

In cases which prove fatal, we find evidences that the same succession of changes is taking place; that an effort has been making to bring about the same result. It is in fact from the examination of the progress which has been made in fatal cases, that we are enabled to judge what is the exact condition of the parts, and what the processes through which they go, in those which recover. Thus in some portions of the organ affected, we find the false membrane very closely adhering to the mucous, whilst the latter is reddened and thickened. This especially occurs at the top of the larynx. Lower down the false membrane is more or less extensively loosened from its adhesion—usually irregularly so—whilst a layer of pus lies between it and the mucous membrane. In some places the effused coat has been entirely separated, and has been either spit up, or else is found loose, enveloped in pus, in some part of the passage; whilst the surface to which it adhered is red,

swollen and besmeared with pus. Thus we trace everywhere distinctly the existence of a process the tendency of which is obviously to bring about recovery ; but death has taken place before it has been completed. It takes place in different steps of the process. Sometimes quite early, before any separation has taken place, the patient apparently dying from the diminished aperture of the air passages from spasm and inflammation. Sometimes later, when the separation has taken place below, but not at the top of the larynx. At other times the membrane separates in considerable quantities, becomes collected into considerable masses, and produces suffocation by being wedged in at the bifurcation of the trachea or at the very top of the larynx. There are other cases in which recovery is also obviously taking place from croup, but in which death occurs from the supervening of secondary disease in the lungs.

Croup, when once established, can then only be recovered from, by going through with this regular course of changes. These are essential to it. When once this process has begun ; when the false membrane has been fairly effused, the parts can no more recover without them than the eruption of smallpox can be cut short in its progress. A rational method of treatment, then, is that which will promote the necessary changes. And what do we need? 1. To prolong life, to prevent suffocation, in order to give time for the required process to be completed by the efforts of the organs themselves ; and, 2. To use means which will promote and hasten this process—which will aid the system in the work which she is aiming to perform.

Now are the usual means likely to answer these purposes? Have they answered these purposes? That emetics and bleeding sometimes relieve violent turns of dyspnoea, must be admitted ; yet that they actually prevent suffocation in many cases, admits of very great doubt. But do they contribute at all to those changes upon which alone we can depend for actual recovery? There is no evidence that they do ; whilst on the contrary there is reason to fear that they may interfere with them, may retard them, may prevent them. If, then, these remedies be at best of doubtful efficacy, is it not right, in so formidable a disease, to make the trial whether other measures may not be more successful? At any rate, if other means are not more successful, they may at least be less tormenting to the patient, and inflict a less amount of unnecessary suffering.

It is to be remarked of the case which has suggested these observations, that the subject of it rejected all remedies, so that it was in fact a case left very much to the resources of nature. Still, so far as the morbid condition in which croup consists is concerned, recovery was very fairly taking place, and would have been complete, except for the occur-

rence of a secondary affection. I may say also of the very few cases which I have seen completely recover by the expectoration of the membrane, that they were not the subjects of very active perturbing treatment, especially after the first stages had gone by, but were left a good deal to palliatives—to mild, soothing applications. It would seem worth while, therefore, to make the attempt of treating the disease without the persevering use of the heroic remedies by which it has been ordinarily encountered; that we should—not perhaps leave the disease wholly to nature—but trust it at least to such remedies as will not interfere with that regular course by means of which nature is always attempting to give relief.

III.—*Further Remarks on the Treatment of Croup.* Read before the Boston Society for Medical Improvement, Feb. 20, 1845.

Some remarks were presented to the Society, a few months since, on the treatment of croup, including suggestions concerning the management of that form of the disease which is attended by the formation of a false membrane in the larynx and trachea. A case of the disease has since occurred to me, which seems to be worthy of notice in connection with those remarks.

The subject was a male, 5½ years of age; of pale and delicate aspect, and slender habit. He had not been perfectly well since an attack of scarlatina, two years ago; since then, he had been frequently liable to colds, with severe coughs. He had enlargement of the submaxillary glands and of the tonsils.

He was first seen on Sunday eve, Feb. 9, 1845. The account given by his parents was, that he had had a cough with a croupy sound—a sound with which they were familiar—for ten days past; but with it no trouble in breathing; that to-day, however, his voice had become hoarse, and that he had had several turns of hard, suffocative breathing. The cough and respiration were at this time distinctly those of croup, though at the time of the visit there was no distress. There was false membrane on the tonsils. He had taken an emetic of ipecacuanha and a dose of castor oil.

He was directed to take, once in three hours, 1½ grains of Dover's powder and ½ a grain of calomel—to sponge the neck frequently with warm water, and to apply to it this liniment—R. Olei oliv., ℥j.; aquæ potass., ℥ij.; ungu. hyd. fort., ℥j. M.

Feb 10th.—The night had been easy upon the whole, though there had been several turns of distress. During one of these he took two



drachms of wine of ipecac., with free vomiting. The symptoms of membranous croup were perfectly well-marked, but there was no distress. The liniment was continued, a flax-seed poultice was applied to the neck, and the powders continued every two hours; to be suspended, however, if he became fully opiated.

During the day the voice became quite extinct; and the cough lost the loud and ringing sound which it presents in the early period of this disease. The breathing became more labored, and was accompanied by greater muscular effort both in inspiration and expiration. Still he was not distressed, owing apparently to the influence of opium. The air entered the lungs well. There was much sound of loose secretions in the larynx and trachea, but no expectoration, except of a little frothy mucus. It having been found difficult to keep the poultices in contact, the parents substituted boiled mullen leaves, which were assiduously applied. At the same time the patient was made constantly to inhale the vapor from a boiling decoction of the same plant, and this was persevered in uninterruptedly for several days.

It is not necessary to follow up a detailed history of the case. These measures were continued without change for several days, i. e., the poultices, the liniment, the inhalation, and the calomel and opium in sufficient quantities to keep him under a moderate narcotism.

On Feb. 12, Wednesday, there had been no distress of breathing; but its croupy character still continued; there had been no return of natural voice; but the sound of the cough had changed, and was like that of common catarrh—quite loose. Through Wednesday and Thursday, there was much rattling of loose matter in the larynx and trachea, and it was coughed up in considerable quantities. Portions of the sputa were mixed with blood, and false membrane was detected in detached pieces enveloped in mucus and pus. One portion of it was of considerable size and distinctly tubular. The fits of coughing, especially when masses of false membrane were ejected, were suffocative, and the sputa were dislodged with difficulty. On Thursday there was still a large thick patch of false membrane on the tonsils. He was occasionally delirious. The pulse were about 120; the respiration varied from 12 to 20, and continued distinctly croupy, though without any distress. He was extremely prostrated.

On Saturday, the respiration had lost the croupy character, but there was still a loose rattling sound in the air-passages, and the voice was unchanged. This day, for the first time, he manifested a little appetite, and his tongue became clean. He had continued occasionally to throw up pieces of false membrane.

On Monday, Feb. 17, he appeared perfectly well except as to strength and voice. By considerable exertion he could make a slight approach to proper voice, but for the most part he spoke in a whisper.\*

The important point to determine in connection with this case, is, how far recovery depended upon the treatment. The treatment consisted—

1. In the absence of all reducing, depleting, and disturbing remedies.
2. Keeping the patient under the full influence of opium combined with calomel.
3. Constant external application of warmth and moisture, and of a mercurial liniment slightly stimulating.
4. Constant inhalation of watery vapor.

It is too much to say that the recovery in this case was to be attributed, with anything like certainty, to the mode of treatment employed. It may have been only one of those coincidences which so frequently mislead us in studying the effects of remedies. Still, as the expectoration of the false membrane has not been a very common occurrence under my observation, and recovery not universal even where it has taken place, it will be at least useful to notice the circumstances which have accompanied a favorable case.

On the supposition that the successful result may have been connected in some degree with the treatment, I should be disposed to attribute it to the following circumstances.

1. To the absence of all such measures as tend to irritate the parts inflamed, and thus to interfere with the natural process of restoration—especially vomiting. That vomiting gives relief to the paroxysms of bad breathing in croup, will not be doubted; and so does it give temporary relief to the distress of an inflamed stomach. But relief of a symptom is not the cure of disease, and does not always tend to its cure. It is not in accordance with what we know of the effects of remedies in other inflamed parts, that concussion, motion, &c., should allay their inflamed condition. Vomiting relieves inflammation of some parts, and some kinds of inflammation; but in this case the parts inflamed are mechanically disturbed by the act, and it has, so far as we can judge, no probable influence upon that peculiar condition which constitutes the disease.

2. To the absence of all depressing and debilitating remedies—as bleeding, purging and vomiting, considered in their effects upon the sys-

---

\* This patient has had no return of the disease to the present time, March, 1850. His voice was not perfectly restored for many weeks.

tem. Such means may be beneficial when we expect resolution of an inflammation. But where the successful issue of the disease depends upon its going through with a certain course of changes, as in croup, they are as likely to interfere with as to promote them.

3. To the relief of the spasmodic contraction of the rima glottidis, which seems more or less to accompany its mechanical diminution by the effused membrane, and to aggravate very much the difficulty of breathing. It is probably upon the suspension of this spasmodic condition that the temporary relief produced by vomiting chiefly depends, and especially vomiting by means of tobacco.

4. To the influence of external warmth and moisture in promoting the suppurative process, by which alone the false membrane can be safely separated.

5. To the constant inhalation of watery vapor. This may have promoted the separation of the false membrane by keeping it from becoming dried by the constant passage of air—and by rendering it pliable and soft, so as to be easily managed and expelled by the organs in the act of coughing.

These considerations lead to the belief that this method of treating croup is at least worthy of trial. But even should it not prove more successful, it is certainly vastly more comfortable than the ordinary method. The patient, whose case has been recorded, suffered very little after the first day, even before the extrication of the membrane. Indeed, taking the disease altogether, it was not attended by more distress than accompanies the average of the acute affections of children.

#### IV.—*Additional Remarks on the Treatment of Croup.* Read before the Suffolk District Medical Society, March, 1850.

Since the occurrence of the case described in the foregoing paper, I have had, from various circumstances, fewer opportunities of witnessing cases of croup than in former years, and only five of this form of the disease have fallen under my notice. The three first of these were treated in the method pursued in the case above related.

The first case was that of a male, 4 years old, who was taken with membranous sore throat accompanied by high constitutional irritation, Oct. 14, 1845. No croupy symptoms occurred till Oct. 18, when they were manifested in a perfectly distinct manner. On the 20th and 21st, patches of false membrane with bloody sputa were raised—and one piece of four inches in length. The raising of the latter was accompanied by a severe and suffocative paroxysm of coughing. On the 22d

he died, eight days from the commencement of the disease, and four from the access of croup. The suffering in this case was very considerable, but far less than I have been accustomed to witness in cases of croup treated according to the ordinary method.

The second was that of a female, 4 years of age, taken with croup on the 8th of Nov. 1845. No depleting or reducing remedies were employed. Patches of membrane, and one piece of considerable size, were brought up on the 10th and a few following days. She never suffered much, improved steadily, and on the 15th seemed well in all respects except the voice, so that on the 16th I did not see her. On the 17th there was a return of all the croupy symptoms, including the appearance of lymph upon the tonsils, and she died on the night of the 19th, eleven days after her first seizure. During no part of the disease was the suffering from dyspnoea very intense for any continued period.

On dissection, the usual appearances were found, and in one lung the false membrane extended for some distance into the bronchi in the substance of the organ.

The third case was a female, 6 years of age, who was seized with the disease Oct. 31, 1847. The onset of the disease was gradual, yet quite distinct. Nov. 2d, the symptoms had become quite severe, and Nov. 3d there was bloody expectoration and pieces of membrane were spit up. Pieces of membrane continued to be found in the sputa for several days, and she was very comfortable and breathed with tolerable ease, yet never losing the distinct croupy sound of respiration and voice. She retained some appetite, and sat up and amused herself as usual. On the 8th she became rapidly worse, but without distress, and died on the 9th, quite easily, ten days from the first attack of the disease.

It will be admitted, I think, that these cases, especially the two last, exhibited certain differences from the common course of this disease, which indicated a favorable influence from difference of treatment.

In all of them the membrane was thrown up in considerable quantities.

In all of them the disease was attended by very much less distress than is usual in croup, and, in two, there was so decided a mitigation of symptoms following the separation of the membrane, as to lead to considerable hope of a favorable termination.

In two, at least, the disease was prolonged to at least twice its average duration under the usual treatment.

In the two other cases, to which reference was made, the same general course of treatment was followed, with the addition of the introduction of a sponge wet with a solution of the nitrate of silver into the larynx.

In each of these cases the application was made as early in the disease as I became satisfied of its distinct character. It was repeated morning and evening. It decidedly gave relief to the breathing soon after each application, and both cases ultimately recovered perfectly. For the suggestion and adoption of this valuable addition to our means of treating this formidable disease, we are indebted, as is well known, to the enterprise of Dr. Horace Green, of New York. The profession, I think, owe to him a large debt of gratitude, for the energy and perseverance manifested in the introduction of this remedy, and I am the more disposed to render this tribute to him because so many attempts have been made to detract from his merit in relation to it.

I am well satisfied from what I have now seen of this method of treating croup, as compared with that which has been followed for so many years, that it has the advantages which were pointed out in one of the preceding papers. It is a disease which I would treat without depletion—except perhaps by a few leeches—without vomiting, without purging, without blisters, without antimonials, ipecac., and all those other nauseous remedies which have been usually resorted to. I would trust to opiates, perhaps calomel, emollients, and the local application of the nitrate of silver.

I ought to add that many of my friends in the profession have informed me of cases in their practice, treated on these principles, which have recovered in a favorable manner. Among them I would refer to Dr. Fisher, Dr. Henry G. Clark, Dr. E. H. Clarke,\* Dr. Buckingham,

---

\* For the following case of croup, treated successfully by cauterization, opium and calomel, occurring in the practice of Dr. M. S. Perry, I am indebted to Dr. E. H. Clarke, of this city.

The patient was a male child, three years and eight months old, with brown hair and dark eyes, and rather stout. He had been a very healthy boy, up to the time of the present attack. This was preceded for several days by a cough, hoarseness and slight difficulty of breathing, which his parents supposed to be an ordinary cold. On the noon of Jan. 6, 1847, they noticed a "white patch" on back side of his throat, which they regarded as a very trivial matter. At night, however, his breathing became whistling and impeded, and they called a physician.

At 3½, P. M., the patient was in the following condition. Asleep; skin natural; pulse 80, intermitting once in 20 beats; extremities warm; respiration laborious, accompanied with a wave-like motion of laryngeal muscles; slight tumefaction about throat externally. On awaking, respiration became easier, but was still labored; cough hoarse and ringing (croupy); tone of voice altered; fauces swollen; left tonsil enlarged and red; right one, coated with lymph, and also epiglottis, as far as visible. The tonsils, fauces, epiglottis, and other parts as far down as possible, were cauterized with nitrate of silver. As the child was too young to steam its throat, the room was kept filled constantly with aqueous vapor by throwing water over heated bricks. A fifteenth of a grain of opium and a grain of calomel was exhibited every second hour. This quantity of opium was sufficient to keep the patient quiet, but not opiated.

At one o'clock of the morning of the 7th, five hours from the commencement of treatment, a gentle emetic was administered. This was followed by free emesis, and a dozen or fifteen shreds of membrane were thrown up, varying in length from 1-3 to 1-4 of an inch, and some of them a

and my brother (Dr. Charles Ware) of this city, Dr. Cotting of Roxbury, and Dr. Spooner of Dorchester.

line in thickness. Respiration became easier, and cough looser and less croupy. There was a decided amelioration of all the symptoms.

At eight in the morning, cough was again hoarse and ringing; respiration whistling and sharp; respiratory movements impeded, and there was much agitation of laryngeal muscles. The left tonsil and epiglottis were found to be again coated with lymph. The right tonsil free. Cauterization was again practised. Steam, opium and calomel were exhibited as before. During the previous night and this day, the pulse varied but little from eighty. At 11, A. M., an emetic similar to the last was given, which was followed by the rejection of a number of shreds of false membrane. These shreds were white, and less tenacious than those first thrown off. At 3, P. M., there was a dejection (from castor oil), which contained considerable muco-filamentous matter, tinged green. At 6, there was another dejection. At 8, P. M., there was a return of the bad symptoms. Respiration was labored; cough ringing and brassy, and tone of voice shrill. On examination, both tonsils and the epiglottis were discovered to be coated with fresh lymph. Cauterization was again practised with the solid nitrate of silver, which was carried as far down the throat and upon the epiglottis as it was possible to do it. The caustic was taken out, completely enveloped in pieces of tough, elastic membrane, which were entangled by it. The room was kept filled with steam as before. Calomel was given once in five hours, and opium in sufficient quantity to keep the child quiet. Croton oil was applied externally about the region of the tonsils.

Jan. 8, A. M., thirty-six hours from commencement of treatment. Patient had passed a comfortable night; had one dejection towards morning. Slight eruption from croton oil. There was a circular slough upon each tonsil, but no appearance of recent lymph. Cough loose, with an occasional ringing sound; respiration easier; pulse 90; tongue coated white; voice shrill. A mixture of tolu, squills, spirits of nitre, with opium enough to allay irritation, was now ordered. Air of room was kept moist but not filled with vapor.

Jan. 9.—Patient was about the same.

Jan. 10.—Had slept well; cough was still hoarse, and slough visible on each tonsil; tongue cleaning at edges; some aphthæ about mouth; is able to sit up and play. On the 11th, there was some yellowness of conjunctivæ and skin, for which a grain of calomel and 1-16 of a grain of opium was exhibited in the evening. After taking the powder, vomited a quantity of thick, stringy, purulent-matter. On the 12th, had slept well; countenance better and less yellowness. Cries to eat; pulse 80, feeble. Bowels have been opened regularly. Continue previous mixture. In the evening, child's throat seemed to be plugged up with a mass of muco-purulent matter, which was cleared out by means of a linen rag, attached to a stick and dipped in a mixture of chloride of soda and tinct. cinchonæ. Previous medicine was omitted, and opium and camphor were ordered, and also wine whey every two hours.

Jan. 13.—Pulse 120, feeble. Throat was again washed out, and much purulent matter brought away. Wine whey and chicken broth were given every hour or two. From this day he steadily improved, and on the 21st, fifteen days from the commencement, was discharged well. His voice, however, did not fully regain its natural tone for several weeks.

Dr. Perry has lately informed me of at least four similar cases of membranous croup, treated in the same manner, within the last year, which have recovered.

E. H. CLARKE.

April, 1850.

and the history of the city, the coming of the  
people and the growth of the city.

The history of the city is a story of growth and progress. It is a story of the people who have lived here and the things they have done.

The city has grown from a small settlement to a large and important center. It has been a place of many firsts and many achievements.

The people of the city have been brave and bold. They have faced many hardships and they have overcome them.

The city has been a place of many great men and women. They have made the city what it is today.

The city has a rich and varied history. It is a history that is full of interest and excitement.

The city is a place where the past and the present meet. It is a place where history is alive and well.

The city is a place of many wonders and many mysteries. It is a place that is full of life and energy.

The city is a place where the future is bright and full of promise. It is a place that is always growing and always changing.

The city is a place where the best of the past and the best of the future meet. It is a place that is truly unique.

The city is a place where the people are proud of their history and their future. It is a place that is truly special.

The city is a place where the people are always looking forward. It is a place that is always moving forward.

The city is a place where the people are always working together. It is a place that is truly united.

The city is a place where the people are always striving for the best. It is a place that is truly excellent.

The city is a place where the people are always making a difference. It is a place that is truly impactful.

The city is a place where the people are always creating a better future. It is a place that is truly hopeful.

The city is a place where the people are always building a brighter future. It is a place that is truly inspiring.

The city is a place where the people are always making a better world. It is a place that is truly beautiful.





