The occurrence of aluminium in vegetable products, animal products, and natural waters: A contribution to the bibliography of the subject / by C.F. Langworthy, PH. D., and Peter T. Austen, PH. D.

Contributors

Langworthy, C. F. 1864-1932. Austen, Peter T. 1852-1907.

Publication/Creation

New York: J. Wiley & sons; [etc., etc.], 1904.

Persistent URL

https://wellcomecollection.org/works/qcuyh2sk


License and attribution

The copyright of this item has not been evaluated. Please refer to the original publisher/creator of this item for more information. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use.

See rightsstatements.org for more information.




Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org





Med K23022



Digitized by the Internet Archive in 2016

8:

THE



OCCURRENCE OF ALUMINIUM

IN

VEGETABLE PRODUCTS, ANIMAL PRODUCTS,

AND

NATURAL WATERS.

A CONTRIBUTION TO THE BIBLIOGRAPHY
OF THE SUBJECT.

BY
C. F. LANGWORTHY, Ph.D.,
AND
PETER T. AUSTEN, Ph.D.

FIRST EDITION.
FIRST THOUSAND.

NEW YORK:


JOHN WILEY & SONS.

LONDON: CHAPMAN & HALL, LIMITED.
1904.

Copyright, 1904, BY JOHN WILEY & SONS.

WEL	LCOME INSTITUTE
Coll.	welMOmec
Call	
No.	WA

ROBERT DRUMMOND, PRINTER, NEW YORK.



INTRODUCTION.

THE material included in this contribution to the bibliography of aluminium deals only with the occurrence of this element in vegetable products, animal products, natural waters, and a few miscellaneous materials, such as edible earths. The general purpose has been to include only reference to such articles as report separate determinations of aluminium or some of its salts, and to omit the very large number in which iron and aluminium are reported together. In a few cases analyses have been cited which report "traces" of aluminium. but the bulk of this material also has been omitted. In older investigations, particularly those dealing with the mineral constituents of plants, data regarding aluminium are more abundant than in later works, and doubtless some of the aluminium reported came from impure reagents, from dirt contaminating the sample, or some similar cause. Such a criticism would not be limited to the constituent under consideration, but applies more or less, in principle at least, to many of the determinations included in early analytical work. The greater part of the material included in the compilation does not seem open to that objection, for, as time has progressed, analytical methods and chemical manipulations have improved, and there is no reason why determinations of aluminium made within recent years should not be fairly good.

No attempt has been made to comment on the value of individual analyses cited, as the object of this bibliography was the collection of data rather than the critical examination of them.


In collecting the data, a systematic search has been made of the files of the Journal of the London Chemical Society,

Copyright, 1904, BY JOHN WILEY & SONS.

114781511

WEI	LIBRARY
Coll.	welMOmec
Call	
No.	WA

ROBERT DRUMMOND, PRINTER, NEW YORK.



INTRODUCTION.

THE material included in this contribution to the bibliography of aluminium deals only with the occurrence of this element in vegetable products, animal products, natural waters, and a few miscellaneous materials, such as edible earths. The general purpose has been to include only reference to such articles as report separate determinations of aluminium or some of its salts, and to omit the very large number in which iron and aluminium are reported together. In a few cases analyses have been cited which report "traces" of aluminium. but the bulk of this material also has been omitted. In older investigations, particularly those dealing with the mineral constituents of plants, data regarding aluminium are more abundant than in later works, and doubtless some of the aluminium reported came from impure reagents, from dirt contaminating the sample, or some similar cause. Such a criticism would not be limited to the constituent under consideration, but applies more or less, in principle at least, to many of the determinations included in early analytical work. The greater part of the material included in the compilation does not seem open to that objection, for, as time has progressed, analytical methods and chemical manipulations have improved, and there is no reason why determinations of aluminium made within recent years should not be fairly good.

No attempt has been made to comment on the value of individual analyses cited, as the object of this bibliography was the collection of data rather than the critical examination of them.

In collecting the data, a systematic search has been made of the files of the Journal of the London Chemical Society,

The American Journal of Pharmacy, The Analyst, Jahresbericht der Thier-Chemie, Jahresbericht der Agricultur-Chemie, Just's Botanischer Jahresbericht, Chemical News, Zeitschrift für Untersuchung der Nahrungs- und Genussmittel, Experiment Station Record, the later volumes of the Comptes Rendus, de l'Académie des Sciences, Paris, Zeitschrift für Physiologische Chemie, the bulletins and other publications of the United States Geological Survey, and the reports of the Geological Survey of Canada, as well as numerous scientific journals, bulletins and reports of the Agricultural Experiment Stations, reports of State Boards of Geology and of Agriculture, and miscellaneous volumes on chemistry, mineral waters, foods, and other topics, including such works as Wolff's "Aschen-Analysen von Landwirtschaftlichen Producten, Fabrik-Ausfällen, und Wildwachsenden Pflanzen," König's "Chemie der Menschlichen Nahrungs- und Genussmittel," etc.

In a great many instances the data found in a periodical or work of reference have been verified in the original publication and so cited. All possible precautions have been taken to insure accuracy, but those who have engaged in similar work know how difficult it is to eliminate all error.




TABLE OF CONTENTS.

	PAGE
Introduction.	iii
ALUMINIUM IN VEGETABLE PRODUCTS	1
ALUMINIUM IN ANIMAL PRODUCTS	49
ALUMINIUM IN NATURAL WATERS	51
ALUMINIUM IN MISCELLANEOUS MATERIALS	131
INDEX	133
v	



ALUMINIUM IN VEGETABLE PRODUCTS.

Aderholdt (Ann. Chem. u. Pharm., 82 (1852), p. 111) reports 6.1 per cent ash in the dry matter of a club moss (*Lycopodium chamaecyparissus*) gathered in March, of which 51.85 per cent was said to be Al₂O₃, and 57.364 per cent Al₂O₃ in the ash (4.5 per cent total) of a sample gathered in November. A club moss (*L. clavatum*) was found to contain in dry matter 4.7 per cent ash, 26.65 per cent being alumina. No alumina was found in other plants of the same region (near Bonn), such as oak, fir, and beech.

Allen, A. H. (Analyst, 13 (1888), p. 41; Jour. Chem. Soc. London, 54 (1888), p. 631), concedes on the statements of analysts that aluminium is present in minute proportions as a normal constituent of wheat, the amount corresponding to about 8 grains of alum in a four-pound loaf of bread. A method for determining the alumina present is proposed.

Andreasch, R. (Jour. Prakt. Chem., n. ser., 18 (1878), pp. 204-207), studied the ash of different parts of the garden pink and rose. The following is quoted:

ALUMINIUM IN CERTAIN PLANTS.

	Total Ash.	Al ₂ O ₃ .
Garden pink (Dianthus caryophyllus):	Per Cent.	Per Cent.
Root.	5.64	2.56
Stem	5.26	trace
Leaves	4.44	
Flower	5 - 59	
Garden rose (Rosa remontana):		
Root	2.04	trace
Stem	2.31	
Leaves	9.43	
Flowers	6.27	

Apoiger (Vierteljahressch. Prakt. Chem., 6, p. 481; Jahresb. Chem., 1857, p. 530) found 7.76 per cent ash in the seed of Maesa pic:a dried at 100°. This contained 0.98 per cent Al₂O₃.

Athenstaed, W. (Ber. Deut. Bot. Gesell., 3 (1885), p. 57; Just's Bot. Jahresbericht, 1885, pt. 1, p. 81), reports the ash analyses of a mixture of leaves, flowers, and fruit of Labrador tea (*Ledum palustre*) according to the method of Grandeau and Fresenius. The crude ash constituted 3.95 per cent of the total dry matter and of this 1.17 per cent was Al₂O₃. The pure ash constituted 2.77 per cent of the total dry matter and of this 1.67 per cent was Al₂O₃.

Baer, W. (Arch. Pharm., 2d ser., 66, p. 285; Pharm. Centbl., 1851, p. 826; Jahresb. Chem., 1851, p. 710, Tab's C and C, p. 708), reports in rape-seed (dry material) 6.98, 5.97, 5.93, and 4.58 per cent ash respectively, containing 0.56, 1.02, 0.60, and 0.49 per cent alumina respectively. He also reports 4.47 and 4.41 per cent total ash in rape-straw (dry material) with respectively 0.63 and 0.22 per cent alumina.

Bailey, E. H. S. (Trans. Kansas Acad. Sci., 11 (1887-8), p. 49), found that the ash of corn-cobs contained a "little over one per cent of ash," 2.02 per cent of this being Al₂O₃.

Bastin, E. S., and H. Trimble (Amer. Jour. Pharm., 69 (1897), pp. 90-97), in an article on North American Coniferæ, report that the bark of hemlock (*Tsuga canadensis*) (air-dry) contains 1.42 per cent ash. Alumina was a consituent of this; the amount not reported.

Baudrimont, E. (Jour. Pharm., 3d ser., 42, p. 388; Jahresb. Chem., 1862, p. 512), reports 23.28 per cent ash in eel-grass (Zostera marina). Of this 0.26 per cent was Al₂O₃.

Béchamp, A. (Compt. Rend. Acad. Sci. Paris, 73, p. 337; Jour. Chem. Soc. London, 24 (1871), p. 855), notes a trace of alumina in the ash of yeast the total ash in the dry yeast being 9.730 per cent.

Bell, J. Carter (Analyst. 4 (1879), pp. 126-133), reports that in four analyses of flour called No. 2 Crown and ground from a mixture of English and California wheat, he found 0.021 and 0.017, 0.020 and 0.024 per cent respectively of aluminium phosphate. Five pounds was made into bread, the crumb containing, according to analyses, 0.011 per cent aluminium phosphate. A Russian and a Ghirka flour, which the author states were "coarse grains flours such as a good baker would not like to use," contained respectively 0.58 and

o.592 per cent ash, with o.o16 per cent aluminium phosphate in each case. Bread was made from 2.75 pounds of the Russian flour. The coarse brown loaf contained o.o10 per cent of aluminium phosphate. Other samples of flour were examined as follows (the ash content and aluminium phosphate content only being quoted)

ALUMINIUM PHOSPHATE IN FLOUR.

Kind of Flour.	Ash	Aluminium Phosphate.
	Per Cent	Per Cent.
r. One Crown flour	0.628	0.018
2. Two Crown flour	.604	.0204
3. Three Crown flour	.52	.000
4. Four Crown flour	.448	.007
5. Five Crown flour	.488	.005
6. White English wheat, 1877	.484	.004
7. Red English wheat, 1877	.438	.010
8. American spring wheat	.492	.013
o. American red winter wheat	.380	.007
o. Empress Hungarian	.368	.002
1. Residue flour, from 1, 2, 3 mixture of Crown	.476	.010
2. Bran flour	.488	.012
3. Exhaust flour	.610	.011
4. Russian flour	.580	.0162
5. Egyptian flour	.700	.0059
6. Ghirka flour	.592	.0163
7. Semolina flour	.440	.0062
8. Flour, English and foreign wheat. & English,	.440	.0002
å foreign, principally Californian	.400	.001
9. Straws, made in Salford	.672	.0023
o. Coarse flour, or seconds, containing bran	1.44	.0058
White flour made in Salford	.680	.0018
2. Peerless flour made by Banaman, Sherman	.000	.0010
& Co., Rochester, New York	.500	.001
3. Gilt Edge, made by Chase, Bristol and Bide,	.500	.001
Rochester	.516	.0008
4. Californian flour.	.480	.0078
5. White English, 1878	.368	12.5
6. Purchased in Salford		.0049
7. Purchased in Salford		.0015
8. Two Crown flour, total produce of wheat—		.009
40 per cent Red English, 30 per cent		
Californian White 30 per cent Canadian		
	260	000=
White	.260	.0087
	.76	.016
o. Two Crown flour, less 5 per cent bran flour	.26	.0106
I. P. A. Campbell, San Francisco	.300	.004
2. J.F., Salem Mills, United States	.448	.004
3. Albany City Mills, United States	.520	.001

Thirty-two samples of bread purchased from bakers in Salford were analyzed, the crumb being used. The amount of moisture and aluminium phosphate follows:

ALUMINIUM PHOSPHATE IN BREAD.*

Kind of Bread.	Moisture.	Aluminium Phosphate.
	Per Cent.	Per Cent.
Sample No. 1	46.10	0.0030
Sample No. 2	45.00	.0067
Sample No. 3	45.50	.0037
Sample No. 4	40.30	.0022
Sample No. 5	40.50	.0030
Sample No. 6	46.00	.0046
Sample No. 7	44.50	.0034
Sample No. 8	46.00	.0082
Sample No. 9	46.00	.0040
Sample No. 10	45.50	.0031
Sample No. 11	46.50	.0052
Sample No. 12	46.00	.0050
Sample No. 13	46.00	.0057
Sample No. 14	46.00	.0026
Sample No. 15	46.20	.0028
Sample No. 16	47.50	.0045
Sample No. 17	46.50	.0043
Sample No. 18	49.50	.0032
Sample No. 19	-47.50	.0064
Sample No. 20	46.50	.0037
Sample No. 21	47.20	.0042
Sample No. 22	46.50	.0068
Sample No. 23	45.50	.0064
Sample No. 24	46.00	.0066
Sample No. 25	45.30	.0052
Sample No. 26	46.20	.0044
Sample No. 27	44.50	.0038
Sample No. 28	44.00	.0058
Sample No. 29	45.50	.0044
Sample No. 30	44.50	.0048
Sample No. 31	44.00	.0042
Sample No. 32	46.00	.0058

The amount of aluminium phosphate found in some samples of bread which did not contain alum was as follows: 0.005, 0.004, 0.005, 0.003, 0.010, 0.010, 0.009, 0.010, 0.009, and 0.008 per cent.

^{*}The inference is that these breads contained no added alum.

In the following table the amount of total moisture and aluminium phosphate in a number of samples of bread known to contain no added alum is quoted from more complete analyses reported by Bell.

ALUMINIUM PHOSPHATE IN BREAD.

Bread made from	Ash Moisture.	Aluminium Phosphate.
	Per Cent.	Per Cent.
One Crown flour	45.00	0.003
Two Crown flour	45.00	.011
Three Crown flour	40.00	.000
Four Crown flour	38.00	.004
Five Crown flour	36.50	.0022
White English wheat	36.90	.004
Red English wheat	39.50	.0053
Russian Flour	42.70	.0053
50 per cent English Red, 50 per cent Canadian		
White	42.50	.0082
Pure White Canadian	42.50	.0035
Canadian Semolina	43.90	.0032
50 per cent English Red, 50 per cent Semolina	43.00	.009
California flour	43.50	.0100
One Crown flour	46.50	.0031
Four Crown flour.	45.00	.0053

The author states that in view of these results he thought it would be of interest to trace the aluminium from the wheat through the various milling products. He, therefore, had flour, bran, etc., especially prepared from wheat which he procured. From the analyses reported the ash content and aluminium phosphate are quoted as follows:

ALUMINIUM PHOSPHATE IN WHEAT AND ITS MILLING PRODUCTS.

Whe at and its Milling Products.	Ash.	Aluminium Phosphate.
	Per Cent.	Per Cent.
English wheat grown upon chalk soil	1.720	0.013
Bran (18.5 per cent of the wheat)	5.640	.016
Sharps (8.5 per cent of the wheat)	2.000	.017
One Crown flour (26 per cent of the wheat) Four Crown flour (45 per cent of the wheat)	.500	.007
Four Crown flour (45 per cent of the wheat)	. 368	.006

Bell, J. Carter (Analyst, 6 (1881), pp. 197-201), reports analyses of eleven grape-juices and of various samples of unfermented and other wines. Among other constituents, the aluminium phosphate was determined. The amount of this

constituent and the total ash found in the different samples follows.

ALUMINIUM IN GRAPE-JUICE, FERMENTED AND UNFER-MENTED WINES.

	Total Ash in 100 cc.	Aluminium Phosphate.
Pure Grape-juices.	Per Cent.	Per Cent.
Black English hot-house grapes	0.356	0.084
White English hot-house grapes	.331	.051
Almeira, 1879	.311	.160
Almeira, 1880	.258	.251
duction of ordinary wine	.273	.512
Bordeaux, 1880. Mixture of Carbenet Sauvignion,	.252	.433
Malbec, and Verdat	. 298	-359
Douro	.261	1.020
Pineau (Champagne grape). From the Cot d'or	.289	.653
Folly Blanc. (Cognac grape)	. 266	1.178
Blanquette	. 284	.760
Grenach No. 1	.291	.789
Grenach No. 2	.289	1.795
Grenache	. 305	3.850
Clairette	. 348	2.582
Congress (from Vineland, N. J.).	-395	,220
Madeira Videilho	. 267	.642
Madeira Tinta	.318	.431
Fermented and Unfermented Wines.		
"Bell's unfermented juice of the vine," pure,		
uncolored fruit	.033	2.085
Unfermented sherry	.030	2.216
Unfermented port	.034	3.956
" new wine," free from alcohol;	54	3.75
guaranteed fruit	.190	.518
vice.	.261	1.002
"Pure and unfermented fruit of the vine "	.342	1.244
Castle Tent, an unfermented sweet wine	. 593	1.341
Castle Rota Tent. Similar to Castle Tent "Unfermented wine, free from alcohol, and unin-	.570	1.360
toxicating"	.290	.492
Greek wine from island of Scio, "unfermented".	. 368	.642
White grape wine, fermented, from island of Scio.	.158	-377
Deidesbeimer, pale alcoholic wine	.185	.971
Deidesheimer Aucolee, a pale alcoholic wine	.194	.684
Italian juice, from Palmi, Calabria	.347	.998

Bergstrand, C. F. ("Ofversicht K. Svenska Vetensk. Akad. Förhandl., 1875, pp. 27-37), studied the effect of Swedish soils,

containing alum, on vegetation. When the soil contained 0.5 per cent alum, the common cultivated plants would not grow. Mountain raspberry (Rubus articus), however, grew and produced fine-tasting fruit. The entire plant when thus grown contained 4.68 per cent pure ash with 3.47 to 5.59 per cent Al₂O₃.

Berthelot and G. André (Compt. Rend. Acad. Sci. Paris, 120 (1895), p. 288, and Ann. Chim. Phys., 7th ser., 5 (1895), p. 429) report, among other data regarding the existence of alumina in plants, the following determinations: Carefully washed alfalfa roots, 0.127 gm. or 0.5 per cent of pure alumina, second sample 0.48 per cent; convolvulus roots, 0.0596 gm. or 0.4 per cent; Bermuda-grass roots, 0.011 gm. or 0.12 per cent; lupine leaves, 0.013 gm. or 0.037 per cent; and linden leaves, 0.0012 gm. or 0.0025 per cent.

"These results indicate that alumina is present in considerable amounts in plants with extensive root systems, but that it remains largely in the roots and is found in only minute quantities in the leaves."

Bevan, E. J., and C. F. Cross (Chem. News, 42 (1880), pp. 77, 91; Jahresb. Chem., 1880, p. 1064) report 0.6 to 2.0 per cent ash in jute fiber; of this, on an average, 5.51 per cent was Al₂O₃.

Bitto, B. von (Landw. Vers. Stat., 42 (1893), 369; Jour. Chem. Soc. London, 64 (1893), II, p. 546), notes 6.76 per cent ash in the whole fruit of red peppers (Capsicum), and 5.66 per cent in the husk of the ripe fruit; these contained respectively a trace and 0.22 per cent Al₂O₃.

Bitto, B. von (Landw. Vers. Stat., 46 (1895), p. 327), reports traces of Al₂O₃ in the ash of red-pepper fruit, "paprika."

Block, H. (Arch. Pharm., 226 (1888), p. 953), reports the ash content of the root, stem, and leaves of English ivy to be 6.34, 4.92, and 12.6 per cent, respectively, of the dry matter. Of this 0.371, 0.637, and 0.312 per cent, respectively, was made up of Al₂O₃. In this and a majority of similar analyses the other ash constituents are reported also.

Blythe, A. W. (Foods: Composition and Analysis. London, 1888, p. 177), is of the opinion that properly cleaned wheat contains no alumina. He believes, however, that particles of clay and sand from millstones find their way into flour, "and

there is no second-class flour in commerce which does not contain some small percentage of alumina." He is of the opinion that the aluminium is present as silicate.

Bobierre (Barral. Jour. Agr., 1876, p. 175; Jahresb. Agr. Chem., 1875-76, p. 133) reports 0.35 per cent Al₂O₃ in the material remaining after breaking hemp.

Bondurant, C. S. (Amer. Jour. Pharm., 59(1887), pp. 122-124), finds that the hydrangea (*Hydrangea arborescens*) contains 3.41 per cent ash, one of the constituents being aluminium (amount not stated).

Bondurant, C. S. (Amer. Jour. Pharm., 59 (1887), pp. 340-342), states that the leaves of coltsfoot (*Tussilago farfara*) contain 17.10 per cent ash, one of its constituents being aluminium (amount not stated).

Brandes, R. (Arch. Pharm., 2d ser., 75 (1853), p. 269; Pharm. Centbl., 1853, p. 739; Jahresb. Chem., 1853, p. 581), notes in roots of Russian, Chinese, old Austrian, and new Austrian rhubarb (air-dry) 18.2, 8.82, 5.8, and 5.54 per cent ash, containing respectively 0.008, 0.015, 0.060, and 0.015 per cent alumina. The Austrian roots were grown near Bilitz.

Brandt (Wittstein's Vierteljahressch., 13, p. 322; Jahresber. Agr. Chem., 1865, p. 108) finds that the best Rheinpfalz to-bacco leaves contain 20.24 per cent ash. Of this 0.216 per cent was found to be Al₂O₃.

Bromeis (Ann. Landw., etc., 14, p. 2; Jour. Prakt. Chem., 48, p. 447; Pharm. Centbl., 1849, pp. 753, 769; Jahresb. Chem., 1849, p. 667 and Tab. B, p. 656) notes 2.10 per cent ash in peas (dry matter) from Frankenfelde. Of this 0.17 per cent was alumina.

Brown, L. P. (Amer. Chem. Jour., 11 (1889), p. 37; Jour. Chem. Soc. London, 56 (1889), p. 543), notes in the ash of tobacco screenings, composed of the small fragments of stems and leaves which, with the dust, are sifted from tobacco during manufacture 43.40 per cent total mineral matter. This contained 0.47 per cent Al₂O₃.

Browne, C. A., Jr. (Pennsylvania Dept. Agr. Bul., 58; Jour. Amer. Chem. Soc., 23 (1901), p. 869), reports 0.30 per cent ash in the flesh of ripe apple. This contained 0.80 per cent alumina.

Bunge, G. von (Ztschr. Biol., 41 (1901), p. 155), gives a detailed analysis of honey. It contains (when fresh) 0.0046 per cent Al₂O₃.

Clarkson, P. S. (Amer. Jour. Pharm., 59 (1887), pp. 277-278), reports upon an analysis of cocoa shells. The total ash amounted to 9.07 per cent. The author states that in addition to the usual constituents, it contained aluminium. He says:

"This element has not been reported in some analyses of the ash, but was found by Wanklyn in the ash of cacao."

Clinch, J. H. M. (Amer. Jour. Pharm., 56 (1884), p. 131), reports an analysis of the leaves of New Jersey tea (Ceanothus americanus). The dry leaves contained 10.9 per cent moisture and 5.31 per cent ash. This was made up, according to the author, of chlorides, sulphates, phosphates, and carbonates of potassium, calcium, magnesium, aluminium, etc. with silicates.

Colby, G. E. (California Station Rpt., 1898–1901, pt. II, p. 252), reports analyses of hops grown in California. Ukiah hops contained 5.80 per cent ash and Wheatland hops 8.65 per cent. The ash contained respectively 2.15 and 2.16 per cent Al₂O₃.Fe₂O₃. The Wheatland hop soil (fine earth) contained from 6.24 to 15.42 per cent Fe₂O₃ and from 2.57 to 9.79 per cent Al₂O₃ according to analyses made by F. J. Snow.

Coppola, M. (Gaz. Chim. Ital., 10, p. 9; Jour. Chem. Soc. London, 37 (1880), p. 382), found 11.16 per cent ash in Stereo-caulon vesuvianum. Of this 11.13 per cent was Al₂O₃.

Chassevent, A., and C. Richet (Compt. Rend. Acad. Sci. Paris, 117 (1893), p. 673; Jour. Chem. Soc. London, 66 (1894), II, p. 63), in a paper on the influences of metallic salts on lactic fermentation, found that the order of toxicity of the metals studied was as follows: Mg, Li, Ca, Sr, Ba, Al, Mn, Fe, Pb, Zn, Cu, Cd, Pt, Hg, Ni, Au, and Co.

Chatin, A. (Compt. Rend. Acad. Sci. Paris, 110 (1890), p. 376; Jour. Chem. Soc. London, 58 (1890), p. 659), states that truffles, contain from 20.84 to 24.26 per cent dry matter with 5.62 to 9.88 ash. This contained traces of aluminium.

Church, A. H. (Chem. News, 30 (1874), p. 137, in a paper on the occurrence of aluminium in certain cryptogams), reports investigations on the aluminium content of a number of varieties. A summary follows.

ALUMINIUM IN CRYPTOGAMS.

Kind of Cryptogam.	Ash in Dry Plant.	Al ₂ O ₃ in Ash.
	Per Cent.	Per Cent.
Club moss (Lycopodium alpinum	3.68	33.50
Club moss (L. clavatum)	2.80	15.24
Club moss (L. selago)	3.20	7.29
Selaginella martensii	11.66	0.26
S. spinulosa	3.44	none
Horsetail (Equisetum maximum)	20.02	none
Adder's tongue (Ophiglossum vulgatum)	8.25	none
Psilotum triquetrum	5.06	trace?

Church, A. H. (Jour. Botany, n. ser., 4 (1875), p. 169), reports that the ash of a club moss (*Lycopodium billardieri*) (5.46 per cent of the dry matter) contained no alumina. This he considered the first instance in which alumina has not been found in lycopodium ash.

Church, A. H. (Proc. Royal Soc. London, 44 (1888), p. 121), in an article on the occurrence of aluminium in certain vascular cryptogams discusses at length the occurrence of aluminium in plants and reports a number of analyses. The results follow.

ALUMINIUM IN CRYPTOGAMS.

Kind of Plant.	Ash in Dry Plant.	Al ₂ O ₃ in Ash.
	Per Cent.	Per Cent.
Club moss (Lycopodium cernuum)		16.00
Club moss (Lycopodium phlegmaria)	4.08	0.45
Club moss (L. billardieri)	5.46	trace
Salvinia natans	16.82	1.86
Marsilea quadrifoliata	11.66	0.54
Tree fern, New Zealand		19.65
Moss (Cyathea serra)	2.70	0.20
Fontinalis antipyretica	4.76	2.82

Quotations from the author's discussion follow.

"So far as the materials at one's disposal warrant any definite conclusions, it may, perhaps, be permissible to say that aluminium is a characteristic and abundant constituent of the ash of many, if not of all, the species of terrestrial Lycopodia; that it is absent from Selaginella and from a number of other allied vascular cryptogams; that it is present in notable quantity in at least one species of tree-fern though practically absent from others; and that it occurs in insignificant amount (like many other elements) in almost every plant in which its presence has been carefully sought for. As to the state of combination in which alumina exists in those plants in which it occurs in mere traces, we have very little information, but in the cereal grains and pulses it is probably in combination with phosphoric acid. In Lycopodia John states that aluminium acetate occurs, Ritthausen speaks of the malate, Arosenius of the tartrate. Anyhow, it is easy to extract abundance of an organic salt of aluminium by exhausting dried and pulverized Lycopodium alpinum with boiling water. So, in some cases, at least, the alumina present in these plants does not exist, as silica does in Equisetum and other highly silicious vegetable structures, in an insoluble form. As to the physiological function, if any, of this element, it is rash to offer an opinion. It is just possible that it may serve to some extent to neutralize the abundant organic acids of the plants in which it occurs, and thus assist, like the cognate element magnesium, in the metabolic processes of vegetation."

In a postscript to his paper he notes the occurrence of a very large quantity of alumina in the ash from the caudex of a tree-fern (Alsophila australis), from Tasmania; also more than mere traces of alumina in the ash of the caudex of Dicksonia squarrosa.

Councier, C. (Landw. Vers. Stat., 29 (1883), p. 241), compared the ash of the leaves of box elder (*Acer negundo*) grown by water-culture method and in soil. The former contained in dry matter 21.29 per cent ash; the latter, 13.29 per cent, with 0.0 and 4.00 per cent respectively of Al₂O₃.

Councier (Bot. Centralblatt, 40 (1889), p. 97, 129) reported the ash analyses of leaves of broad-leaved maple or sycamore (Acer Pseudoplatanus), gathered after they had fallen, lilac (Syringa vulgaris), beech-leaves, gentian (Gentiana ciliata), bloody eye (blut-auge) (Adonis aestivalis), club moss (Lycopodium annotinum), adder's-tongue (Ophioglossum vulgatum),

carrot peel, winter rye seed, winter wheat seed, potatoes, stems, leaves, and fruit of mistletoe (Viscum album), twigs of Scotch pine (Pinus sylvestris), and pine twigs affected with mistletoe. In only one of these, namely, Lycopodium annotinum, was any aluminium found. The amount of this in the ash was 18.1 per cent.

Coupin, H. (Compt. Rend. Acad. Sci. Paris, 132 (1901), p. 645; Jour. Chem. Soc. London, 80 (1901), II, p. 335), in a study of the sensibility of higher plants to very feeble amounts of a number of mineral salts, notes that 1/50,000 Al₂ (SO₄)₃ in distilled water was toxic to wheat roots, the roots being injured by this and the other salts tested but not killed.

Crawfurd (Vierteljahressch. Prakt. Pharm., 6, p. 361; Chem. Centbl., 1857, p. 604; Jahresb. Chem., 1857, p. 530) notes 2.6 per cent ash in the seed of corn cockle (Agrostemma githago). Of this 1.17 per cent was Al₂O₃.

Cugini (Pharm. Jour. Trans., 3d ser., 7, p. 616; Jahresb. Chem., 1878, p. 940) regards alumina as one of the elements accidentally present in plants, and not as necessary for their growth.

Dafert, F. W. (Relatorio annual. Inst. Agron. Estado Sao Paulo et Campenia [Brazil], 1893; Jahresb. Agr. Chem., 1894, pp. 223-225), reports the ash analyses of a number of grasses. One of the panic grasses (Panicum monostachyum), cut just before blooming, contains 20.14 per cent pure ash in dry matter; of this 0.57 per cent was Al₂O₃. Tickle-grass (Panicum capillare) cut after blooming contained in the dry matter 8.52 per cent pure ash, with 2.23 per cent Al₂O₃ in it. The Al₂O₃ was not reported in the thirteen other varieties analyzed, but was included generally with the Fe₂O₃.

Damarçay, E. (Compt. Rend. Acad. Sci. Paris, 130 (1900), p. 91; Jour. Chem. Soc. London, 78 (1900), II, p. 235), states that the ash of Scotch fir, silver fir, vine, oak, poplar, and horn beam, when treated with boiling water, gives a solution from which hydrogen sulphide precipitates a brown material soluble in hydrochloric acid. When examined with a spectroscope such solutions showed the presence of several metals including a trace of aluminium.

Dambergis, A. K. (Oesterr. Chem. Ztg., 1 (1898), p. 479).

reports the ash analyses of a number of sorts of Greek tobacco. The following is quoted:

ASH OF GREEK TOBACCO.

Kind of Tobacco.	Total Ash.	Al ₂ O ₃ .
l'obacco from	Per Cent.	Per Cent
Aghias	18.75	0.187
Almyro	18.23	.177
Argos.	16.02	.132
Arta	14.58	.151
Carditza	16.30	.100
Corinth	12.41	.102
Epidaure-Limira	16.37	.083
Curytanie	16.03	.131
Kalavrita	13.70	.008
Larissa	15.70	.114
Locrides	17.15	.156
Messi-Corfou	13.33	.148
Messolonghi	20.78	.177
Nauplia	21.03	.151
Oros	15.43	.140
Parnassus	18.55	.076
Pharsala	21.91	.134
Phthiotis	19.83	.180
Trikala	12.65	.105
Trichonia	24.69	.253
Tyrnavo	17.08	.142
Volto.	16.29	.119
Volos.	19.38	.177
Vonitza.	20.61	.223
Xyrochorion	22.79	.188
ımbek (Nicotina persica) from		
Argos.	18.85	.212
Messolonghi	16.92	.154
Nauplia,	18.34	.152
Phthiodis	13.13	.100
Trichonia	20.54	.169

Daubrawa (Vierteljahressch. Prakt. Pharm., 3, p. 337; Pharm. Centbl., 1854, p. 731; Jahresb. Chem., 1854, p. 659) found 12.12 per cent ash, containing 25.8 per cent carbon in shepherds' purse (Capsella bursa pastoris) (air-dry plant); without carbon, the Al₂O₃ content was 0.14 per cent.

Dieterich, E. (Vierteljahressch. Prakt. Pharm., 15, p. 196; Ztschr. Chem., 1866, p. 286; Chem. Centbl., 1867, p. 271; Bul. Soc. Chim., 2d ser., 7, p. 165; Jahresb. Chem. Will, 1866, p. 706), reports 3.02 per cent ash in the dry flesh of sweet chestnuts and 1.845 per cent in the shells, dried at 110°; of the former 0.090 per cent being Al₂O₃ and of the latter 5.793 per cent.

Doolittle, R. E., and W. H. Hess (Jour. Amer. Chem. Soc., 22 (1900), p. 218) note that the ash of cider vinegar should amount to not less than 0.25 per cent and that it contains a trace of aluminium.

Dunnington, F. P. (Jour. Amer. Chem. Soc., 2 (1880), p. 24), reports the ash constituents of eleven weeds. The plants were just about to bloom when gathered. They are as follows:

A T TTTT	TATTTTAE TA	T TITTETT TO CO
AN I I I I I		VALUE IN THE
TILLUMI	TIVE CIVE II	WEEDS.

Kind of Plant.	Pure Ash.	Al ₂ O ₃ .	
No. 1. Broom sedge (Andropogon scoparius) No. 2. Wiregrass (Eleusine Indica). No. 3. Blue thistle (Echium vulgare). No. 4. Potato weed (Solanum carolinense) No. 5. Purslane (Portulaca oleracea). No. 6. Sumach (Rhus glabra) No. 7. Sassafras (S. officinale) No. 8. Ragweed (Ambrosia artemisiæfolia) No. 9. Mullein (Verbascum thapsus) No. 10. Dock (Rumex obtusifolius)*	Per Cent. 2 . 24 2 . 45 4 . 69 2 . 17 1 . 52 1 . 48 1 . 41 1 . 93 1 . 01 1 . 60	Per Cent. 0.20 0.22 0.56 0.11 0.49 1.15 1.26 0.00 1.15 0.45	

The SiO₂, Al₂O₃, Fe₂O₃, and Mn₃O₄ reported are regarded as due to dust adhering to the plants analyzed.

Dupré, A. (Analyst, 4 (1879), p. 1), in a paper on the detection and estimation of aluminium in wheat flour, reports the examination of twelve samples of flour purchased from reputable bakers, in which alum could not be detected. A four-pound sample of each of these flours contained aluminium as follows:

No. 1, 0.63 grains; No. 2, 1.26 grains; No. 3, 1.80 grains; No. 4, 1.41 grains; No. 5, 2.30 grains; No. 6, 1.07 grains; No. 7, 1.62 grains; No. 8, 2.79 grains; No. 9, 2.79 grains; No. 10, 2.50 grains; No. 11, 2.05 grains, and No. 12, 3.72 grains.

Ebert, A. E. (Vierteljahressch. Prakt. Pharm., 17, p. 438; Jahrb. Chem. Will., 1868, p. 815), reports 51.476 per cent ash in

^{*}An abstract of this investigation (Jahresber. Agr. Chem., 1882, p. 141) gives figures for the ash analysis of stickweed (*Verbesina siegesbeckia*), 3.53 per cent being the pure ash content and 0.15 per cent the Al₂O₃. No analyses of stickweed appear in the original publication.

the air-dry seeds of anatto (Bixa orellana). This contained 0.808 per cent alumina.

England, J. W. (Amer. Jour. Pharm., 55 (1883), p. 246), studied a myrtle (*Myrtus cheken*) and found 4.84 per cent ash in stems and 8.4 per cent in leaves. Both contained aluminium. The amount was not stated.

Enz (Vierteljahressch. Prakt. Pharm., 5, p. 196; Chem. Centbl., 1856, p. 393; Jahresb. Chem., 1856, p. 691) notes 1 per cent ash in the fresh berries of honeysuckle (Lonicera xylosteum). Of this o.1 per cent was Al₂O₃.

Eylerts, C. (Vierteljahressch. Prakt. Pharm., 9, p. 529; Jahresb. Chém., 1860, p. 561), found 6.57 per cent ash in air-dry "German worm seed" (Semen cinæ), i.e. the unexpanded flower heads of wormwood (Artemisia maritima var. Stechmanniana). Of this 2.33 per cent was Al₂O₃.

Eylerts, C. (Arch. Pharm., 2d ser., 109, p. 105; Viertel-jahressch. Prakt. Pharm., 11, p. 517; Chem. Centbl., 1862, p. 154; Jahresb. Chem., 1862, p. 512), reports the following in red beets:

Beets from Munich dried at 110°:

Root, total ash, 13. 4 per cent, Al₂O₃ in ash, 0.28 per cent; leaves, total ash, 20.86 per cent, Al₂O₃ in ash, 0.12 per cent.

Beets from Weyenstephen dried at 110°:

Root, total ash, 11.27 per cent, Al₂O₃ in ash, 0.25 per cent; leaves, total ash, 15.57 per cent, Al₂O₃ in ash, 1.99 per cent.

Falco, C. (Vierteljahressch. Prakt. Pharm., 15, p. 509; Chem. Centbl., 1867, p. 142; Jahresb. Chem. Will., 1866, p. 709), reports 8.3 per cent ash in the bark of *Petalostigma quadrilo-culare*, containing 0.047 per cent Al₂O₃.

Fraas, H. (Vierteljahressch. Prakt. Pharm., 15, p. 338; Jahresb. Chem. Will., 1866, p. 710), reports 3.149 per cent ash in the dry leaves of *Gastrolobium bilobum*, a leguminous plant of West Australia reported to be poisonous. Of this 1.145 per cent was Al₂O₃.

Frankforter, G. B. (Amer. Jour. Pharm., 69 (1897), p. 134), reports that poke-weed (*Phytolacca decandra*) root contains 13.38 per cent ash. Of this 1.62 per cent was Al₂O₃.

Fritsch, K. (Arch. Pharm., 3d ser., 27 (1889), p. 193), studied the ash constituents of several plants. A boletus (Boletus edulis) contained 9.20 per cent dry matter, of which 7.32 per

cent was pure ash. Al₂O₃ constituted 0.02 per cent of the latter. The amount of pure ash in the dry matter (the amount not stated) of *Polysaccum pisocarpium* was 5.28 per cent. Of this Al₂O₃ constituted 1.35 per cent.

Gabba, L. (Trattato di analysi chimica, Vol. II, p. 514), in discussing the aluminium content of wine, reports that it does not occur in excess of o.o. per cent. Ricciardi notes that he has demonstrated the occurrence of alumina in wine and has found o.13 gm. in 479 gms. grape seed (equal to o.0027 per cent).

Gasparin, P. v. (Barral. Jour. Agr., 2 (1875), p. 410; Jahresb. Agr. Chem., 1875–76, p. 139; Centbl. Agr. Chem., 8 (1875), p. 249), reports 1.375 per cent Al₂O₃ in soil and none in alfalfa.

Gaze, R. (Apoth. Ztg., 5 (1900), p. 9; Just's Bot. Jahresbericht., 1890, I, No. 1, p. 51), reports that the root of golden seal (*Hydrastis canadensis*), dried at 100°, contains 0.3259 per cent Al₂O₃.

Glenk, R. (Amer. Jour. Pharm., 63 (1891), p. 328), reports a study of water-hemlock (*Cicuta maculata*). The fruit dried at 110° lost 10 per cent in weight. The ash amounted to 6 per cent and contained 20 per cent Al₂O₃.

Gmelin (Anorganische Chemie, 2, 1, p. 608) notes that the ash of wild plants often contains alumina, and cites the following: eel-grass (Zostera marina) 0.27 per cent; alga (Cladophora glomerata) 0.31 per cent; wormwood (Artemisia maritima) 1.5 per cent, chickweed (Arenaria media) 1.03 per cent; chickweed (Arenaria rubra) 1.92 per cent; plantain (Plantago media) 0.63 per cent, corydalis (Corydalis bulbosa) 3.88 per cent; heather (Erica vulgaris) 2.3 per cent; "Weigmann" 0.51 to 0.84 per cent, morel (Morchella esculenta) 1.32 per cent; "Birkenschwamm" 3.73 per cent; edible mushroom (Chlorangium jussuffii) 11.9 per cent; turf-moss (sphagnum) 3.0 to 8.0 per cent, Lycopodium clavatum 26.65 and Lycopodium chamae-cyparissus 51.85 per cent alumina.

Other Cryptogams contain alumina, notably Gyrophora, Cladonia, Usnea, and Cetrarid.

Gobley (Jour. Pharm., 3d ser., 37, p. 19; Jour. Chim. Med., 4th ser., 6, p. 129; Rev. Chim. Appliquée, 2, p. 6; Vierteljahressch, Prakt. Pharm., 9, p. 567; Chem. Centbl., 1860, p. 587;

Chem. News, 2, p. 119; Jahresb. Chem., 1860, p. 550) notes the occurrence of alumina in the root of Kava root (*Piper methysticum*).

Godeffroy, R. (Arch. Pharm., 3d ser., 10 (1877), p. 297; Jahresb. Agr. Chem., 1877, p. 118), reports a trace of Al_2O_3 in the ash of cockle-burr (Xanthium spinosum).

Griffiths, A. B. (Compt. Rend. Acad. Sci. Paris, 131 (1900), p. 422), reports the percentage composition of a number of medicinal plants. Several contained alumina as follows: Sarsaparilla, o.1 per cent; Cardamon, o.1 per cent; "Chine," o.13 per cent, and Rhatany, o.1 per cent.

Guerrieri, F. (Staz. Sper. Agr. Ital., 34 (1901), p. 338), in an article on the value of grapevine branches as a feed for farm animals, reports a total of 2.9053 per cent ash. Of this 0.0650 was found to be aluminium sesquioxid. The proportions in hay are given as total ash, 8.0259, aluminium sesquioxid, 0.2340; in straw, total ash, 6.0584, aluminium sesquioxid, 0.2778 per cent.

Hadelich, W. (Chem. Centbl., 1881, p. 394; Jour. Chem. Soc. London, 42 (1882), p. 121), found in white wine from Erfurt 0.281 per cent ash which contained a trace of Al₂O₃.

Harms, E. (Arch. Pharm., 2d ser., 88 (1856), p. 165; Chem. Centbl., 1856, p. 944; Jahresb. Chem., 1856, p. 690), notes 6.05 per cent ash in air-dry Kousso (*Brayera anthelmintica*) containing 1.97 per cent Al₂O₃.

Harms, E. (Arch. Pharm., 2d ser., 1858, p. 137; Chem. Centbl., 1858, p. 443; Jahresb. Agr. Chem., 1858-59, pp. 64, 65), reports the following:

ALUMINIUM IN CERTAIN PLANTS.

Kind of Plant.	Water.	Ash in Fresh Material.	Aluminium Phosphate in Ash.	
Chickweed (Arenaria media) whole plant in bloom. Plantain (Plantago maritima) green portion. Plantain (Plantago maritima) seed. Chickweed (Arenaria rubra) entire plant	83.48 79.52	Per Cent. 4.60 3.91 5.04 2.56	Per Cent. 2.01 1.13 0.69 4.05	

The ash was said to be not quite free from the soil on which the plants were grown. The aluminium content, therefore, is regarded as too large.

Harper, H. W. (Amer. Jour. Pharm., 53 (1881), p. 209), in a study of fragrant sumach (*Rhus aromatica*), identified aluminium as one of the constituents of the ash of the bark. The total ash was 13.85 per cent.

Hébert, A. (Bul. Soc. Chim., 3d ser., 13, p. 927; Jour. Chem. Soc. London, 70 (1896), II, p. 494), notes the occurrence of alumina in the sap of the banana tree, *Musa paradisica*.

Heller, C. F. (Amer. Jour. Pharm., 59 (1887), p. 68), reports 5.5 per cent ash in Bryony root, containing 7.5 per cent moisture. The ash was found to contain aluminium in addition to other mineral constituents, amount not stated.

Herapath, T. J. (Jour. Roy. Agr. Soc. England, 11, pt. 1, p. 93; Quart. Jour. Chem. Soc., 3, p. 193; Ann. Chem. Pharm., 76, p. 383; Jahresb. Chem., 1850, p. 647, 666, 671, Tab. A, B and D, p. 661), notes the occurrence of a trace of alumina in the ash of wheat, the total ash in dry material being 2.30 per cent, and 0.07 per cent alumina in the ash of oats, the total ash in the dry material being 3.06 per cent. He also reports a trace of alumina in the ash of horse bean (Vicia faba) seed, total ash, 4.3 per cent of dry material and traces in ash of eddoes or taro (Arum (Colocasia) esculentum) and yams (sweet potato) (Convolvulus batatas) containing respectively 1.647 per cent ash and 1.51 per cent ash in the fresh materials.

Hermann, J. C. (Viertel. Prakt. Pharm., 18, p. 481; Jahresb. Chem. Will., 1869, p. 796), notes 0.5851 per cent alumina in the ash of ergot (Secale cornutum); the amount of total ash is not stated.

Herrmann, J. C. (Viertel. Prakt. Pharm., 1869, p. 481; Amer. Jour. Pharm., 42 (1870), p. 143), reports 2.201 per cent ash in ergot. Aluminium phosphate is noted as a constituent of the ash.

Hertwig (Chevallier et Baudrimont's Dictionnaire des alterations, Paris, 1878, p. 1132) notes the occurrence of 9.04 per cent calcium, magnesium, iron, manganese phosphates and aluminium phosphates in Havana tobacco and 17.95 per cent of calcium, magnesium, iron, manganese, and aluminium phosphates in the ash of Hanover tobacco.

Herzberg, W. (Dingler's Polytech. Jour., 277, p. 478 (Ausz); Jour. Soc. Chem. Ind., 9, p. 758 (Ausz); Chem. Centbl., 1890, II, p. 186; Jahresb. Chem., 1890, p. 2877), reports 17.47 per cent ash in the fibers of the baobab tree (Andansonia digitata) used for paper making. Of this 0.14 per cent was alumina.

Hoeven, J. (Amer. Jour. Pharm., 55 (1888), p. 253), in a study of cheken leaves, reports 9.48 per cent ash. The portion insoluble in H₂O, but soluble in HCl, amounted to 5.61 per cent. This contained aluminium; amount not stated.

Höhn, H. (Arch. Pharm., 2d ser., 139 (1869), p. 213; Jahresb. Chem. Will., 1869, p. 771), reports 3.24 per cent ash in the stem and 6.61 in the leaf of *Ophelia chirata*; of this 1.47 and 4.35 per cent was Al₂O₃.

Hollandt, H. (Vierteljahressch. Prakt. Pharm., 10, p. 321; Jahresber. Agr. Chem., 1861-62, p. 60), reports 1.2035 per cent Al₂O₃ in the ash of the bark of *Millingtonia hortensis*.

Hooper, D. (Pharm. Jour. and Trans., 1887, Jan. 8; Amer. Jour. Pharm., 59 (1887), p. 86), reports a study of the ash of cinchona bark. As shown by the average of three hundred determinations, such bark (from India), contained 3.42 per cent ash. The ash of *C. officinalia* and *C. succirubra* grown on the Nilgiris contained respectively 2.70 and 4.24 per cent Al₂O₃.

Huber, L. (Arch. Pharm., 3d ser. 7 (1875), p. 394; Jahresb. Agr. Chem., 1875-76, p. 136), reports 0.250 per cent Al₂O₃ in the ash of the rind of the root of elder (Sambucus nigra), the ash being equal to 11.717 per cent.

Irby, J. R. McD., and J. A. Cabell (Chemical News, 1874, p. 117, Centbl. Agr. Chem., 6 (1874), p. 366) note the presence of some Al₂O₃ with the Fe₂O₃ in six sorts of Virginia tobacco.

Jackson, D. D. (Jour. Soc. Chem. Ind., 21 (1902), p. 681), reports the results of a study of the precipitation of iron, manganese and aluminium from natural waters by the action of bacteria, viz., Crenothrix kühniana, C. manganifera, and C. ochracea, respectively. Dr. Jackson states that each of the three species seems to exercise a selective power in precipitating the special oxid ascribed to it, and that about one-third of the dry organism is composed of the oxid selected, whether iron, manganese, or aluminium. He reports analyses of the precipitate due to the three species in three different waters:

PRECIPITATE	CAUSED BY	CRENOTHRIX	SP. FROM
	DIFFERENT	WATERS.	

	Fe ₂ O ₃ .	Mn ₂ O ₃ .	Al ₂ O ₃ .	SiO ₂ .	Alumin- ium Sili- cate.	Unde- termin- able Mineral Matter.	Organic Matter.
	Per Ct.	Per Ct.	Per Ct.	Per Ct.	Per Ct.	Per Ct.	Per Ct.
Crenothrix kühniana from filter gallery, I p s w i c h River, Mass	31.6				10.4		37.0
Charles River, Newton, Mass Crenothrix ochracea, driven wells at	14.4	33.9	1.5	12.5	8.0	1.8	27.9
Oconee, Brooklyn, New York			33 · 3	5.9	7.6	2.3	36.2

The waters in which the crenothrix species grew were analyzed. They contained alumina (parts per million) as follows: Water from filter gallery, shore Ipswich River, Reading, Massachusetts, o.6; water from filter gallery, shore Charles River, Newton, Massachusetts, o.5, and water from driven wells, Oconee, Brooklyn, N. Y., 2.0.

Jessler, A. (Vierteljahressch. Prakt. Pharm., 17, p. 266, Jahresber. Chem. Will., 1868, p. 807), reports 13.18 per cent ash in the rhizome of dog-bane (*Cynanchum monspeliacum*). This is reported to contain alumina; the amount is not stated.

Hassall, J. (Food Adulteration, p. 617), reports 4.80 per cent ash in anatto. This contained 0.20 per cent alumina (with a trace of iron).

Kane (Phil. Mag., 3d ser., 31, pp. 36, 105; Jour. Prakt. Chim., 41, p. 434; 32, p. 354; Jahresb. Chem., 1847–48, p. 1085) reports total ash as follows in flax stems (treated in different ways) from Ireland, Holland, Belgium, etc., 4.237, 5.434, 3.670, 5.151, and 5.000 per cent. Of this 0.44, 0.44, 0.72, 1.44, and 6.08 per cent respectively was alumina.

Karmrodt, C. (Ztschr. Landw. Ver. Rheinpreussen, 1864, p. 427; Jahresb. Agr. Chem., 1864, p. 98), determined the constituents of the ash of a number of Coniferæ. Larch needles (*Pinus larix*) contained 0.264 per cent; Scotch pine needles

(P. sylvestris) 0.125 per cent, and fir needles (Abies excelsa) 1.427 per cent Al₂O₃.

Kayser, R. (Repert. Analyt. Chem., 1883, p. 289; Jahresb. Agr. Chem., 1883, p. 349), in an article on fruit and fruit juices, reports traces of Al₂O₃ in the juice of red and white currants (containing respectively 0.38 and 0.5 per cent ash) and of strawberries.

Kayser, R. (Repert. Analyt. Chem., 1883, p. 182; Jahresb. Chem., 1883, p. 1407), analyzed dry blueberries, reporting in them 0.005 per cent alumina.

Kayser (Repert. Analyt. Chem., 1882, p. 242) finds that the presence of 0.3-0.4 grain of alumina per liter in wine cannot safely be taken as a proof that alum has been added because it takes into solution during its course of manufacture some alumina. This is especially the case when the wine has been cleared with kaolin (silicate of alumina), as some wines dissolve small amounts of the kaolin, and hence contain alumina in the form of a soluble salt.

Kebler, L. F. (Amer. Jour. Pharm., 67 (1895), p. 398), in a study of the determination of morphine in opium, reports that the ash of crude morphine (percentage amount not given) contained 0.43 per cent aluminium phosphate.

Keim, W. (Ztschr. Analyt. Chem., 30 (1891), p. 401), reports the following amounts of pure ash in cherry (*Prunus cerasus*) fruit at different times:

May 21, 0.516 per cent; May 28, 0.646 per cent; June 19 (ripe), 0.739 per cent. Of these, 2.55, 0.80, and 0.81 per cent respectively was Al₂O₃.

Khittel (Vierteljahressch. Prakt. Pharm., 7, p. 348; Jahresb. Chem., 1858, p. 530) found 7.91 per cent ash in leaves of poisonivy (*Rhus toxicodendron*) (dried at 100°). Of this 0.49 per cent was Al₂O₃.

Knop (Kreislauf des Stoffes, 1868, p. 263) states that the roots of the Lycopodiæ secrete besides carbonic acid strong organic acids, which dissolve the alumina of soils, and this solution enters the root. The lichens also contain in their structures peculiar crystallizable acids which, by the action of reagents, break up into oxalic acid and a new organic acid. The oxalic acid thus formed dissolves the alumina and iron oxid.

and thus the mosses corrode the surfaces on which they grow, the hard rock as well as the dust that clings to it. In this way the dissolved alumina passes into the interior of the moss. It would seem a principle that plants secreting strong organic acids, which act on surrounding material, will contain alumina and show it in their ash.

Knop, W. (Landw. Vers. Stat., 7 (1865), pp. 436-450), reports 31.01 per cent ash (including 15.00 per cent sand) in Chlorangium jussuffii; this included 1.90 per cent Al₂O₃.

Other ash analyses of lichens which he reports are quoted in part as follows:

ALUMINA IN CERTAIN LICHENS.

Kind of Lichen.	Pure Ash in Dry Matter.	Al ₂ O ₃ .
Gyrophora pustulata	Per Cent. 1.666 2.346	Per Cent. 0.344 0.031

Kresling, K. (Arch. Pharm., 3d ser., 229 (1891), p. 389; Jour. Chem. Soc. London, 62 (1892), II, p. 232), in the pollen of Scotch pine (*Pinus sylvestris*), reports 5.51 per cent crude ash, equivalent to 3.0 pure ash. This contained 1.86 per cent Al₂O₃.

Krutzsch, H. (Tharander forstl. Jahrb., 6 (1850), p. 88), as cited by Wolff (Aschen Analysen, Berlin, 1880, pt. 2, pp. 87, 130), determined the ash constituents of freshly fallen fir needles, and of dry twigs, dry bark, dry cones, and moss, found in the litter. The pure ash in the several materials was equal to 1.601, 1.043, 1.275, 0.477, and 2.884 per cent respectively. Of the pure ash 4.20, 8.33, 7.86, 9.20, and 20.5 per cent respectively was Al₂O₃. The Al₂O₃ is regarded as due to earth present in the material. Wolff states that on an average (based on fourteen analyses) fir-needle litter contains 1.410 per cent pure ash; from 0 to 9.98 per cent or on an average 3.58 per cent of this being Al₂O₃. In this and all the other analyses cited from Wolff the complete ash analyses are given.

Langer, A. (Ueber Bestandtheile der Lycopodiumsporen. Inaug. Diss. Erlangen, 1889; Just's Bot. Jahresb., 1889, I, p. 39), reports an analysis of the spores of a club moss (Lycopodium clavatum). These contained on an average 1.155 per cent of mineral matter with 24.41 per cent of Al₂(PO₄)₂.

Langer, A. (Arch. Pharm., 3d ser., 27 (1889), p. 289; Jahresb. Agr. Chem., 1889, p. 383), reports 1.15 per cent ash in commercial lycopodium powder (*L. clavatum*) which was 96.67 per cent pure. Of this 15.30 per cent was Al₂O₃.

LaWall, C. H. (Amer. Jour. Pharm., 69 (1897), p. 137), in a study of the ash of various drugs, reports aluminium as a constituent of aconitum root, belladonna leaves, calumba roots, castanea leaves, chimaphila leaves, cinchona, calisaya bark, digitalis leaves, frangula bark, gelsemium root, geranium rhizome, glycyrrhiza root, guarana seed (crushed), hops, hydrangea root, hydrastis rhizome, hyoscyamus leaves, jalap tuberous root, kola-nut, Piscidia erythrina bark, cho ke-cherry (Prunus virginiana) bark, cascara (Rhamnus purshiana) bark, ruta herba, sabadilla seed, bloodroot (Sanguinaria) rhizome, dandelion (Taraxicum) root, red clover (Trifolium pratense) flowering tops, elm (Ulmus) bark, hellebore (Veratrum viride) rhizome rootlets, and prickly ash (Xanthoxylum) bark. The amount of this and other constituents was not determined. The total ash was generally determined.

Lermer, J. (Dingler's Polytech. Jour., 179, p. 71; Bul. Soc. Chim., 2d ser., 6, p. 429; Jahresb. Chem. Will., 1866, p. 882), reports 6.91 per cent ash in Hungarian malt (barley) sprouts and 6.19 per cent in German. The ash contained respectively 1.06 and 0.45 Al₂O₃.

Lermer (Wittstein Viertel. Jahressch., 13, p. 182; Jahresb. Agr. Chem., 1865, p. 117) reports 7.135 per cent ash in air-dry hops; 0.763 per cent of this was found to be Al₂O₃.

Lewitsky, J. (Arbeiten der dritten Versammlung russischer Naturforscher zu Kiew.— Vereinigte Sectionen der Botanik und Chemie; Just's Bot. Jahresber., 2 (1874), II, p. 866), reported a series of studies of the relation of the phosphates of iron, calcium, and aluminium to plant growth. Investigations have shown that the addition of calcium phosphate to the soil results

in the formation of ferric phosphate and aluminium phosphate, so that after a time, even under the most favorable conditions, practically only a trace of calcium phosphate remains. On the basis of this conclusion, the author believes that under normal conditions the roots of plants have access only to iron phosphate and aluminium phosphate and must derive their phosphates from these salts. To test the theory, culture experiments were made with barley by Hellriegel's method (Ann. Landw., 38, p. 296). Briefly stated, the grain was sown in quartz sand to which nutrient solutions were added. These were made up of KNO3 (or under certain conditions, Ca(NO₃)₂), CaCO₃, NaCl, MgSO₄, and Fe₂O₃, in what were deemed the proper proportions. With such a basal mixture, Fe2(PO4)2, Ca3(PO4)2, and PO4KH2 were added in the first test of the experiments, while in a control experiment, which is described in less detail, Al₂(PO₄), was also used. The plants were watered with water containing CO2. Their growth and seed production is described in detail. In the principal test, the best growth was made by the barley grown with a nutrient mixture containing Fe₂(PO₄)₂. The next best results were obtained with KH2PO4; the least satisfactory, with Ca2(PO4)2. The remaining test was not regarded as entirely satisfactory and gave results somewhat at variance with those already obtained; but in general it showed, according to the author, that Al₂(PO₄), had practically the same effect as a corresponding iron salt and that the corresponding calcium salt was not as satisfactory for the plants as the iron salt.

L'Hote (Compt. Rend. Acad. Sci. Paris, 104 (1887), p. 853), reported an investigation dealing with the detection and estimation of alumina in wine and grapes. Wines from seven localities were found to contain alumina per liter as follows:

From Bourgogne, 0.02 gm.; Cher, 0.036 gm.; Touraine, 0.000 gm.; Roussillon, 0.032 gm.; Spain, 0.016 gm.; Sicily, 0.012 gm.; and from Aude, 0.016 gm.

When prepared in the laboratory, wine from grapes from Huesca contained per liter 0.012 gm.

From 479 gms. of red grapes 0.013 gm. alumina was obtained, while the stalks weighed 6.482 gms. and contained 0.003 gm. alumina.

Light, W. W. (Amer. Jour. Pharm., 56 (1884), p. 3), reports the analysis of the fruit of prickly pear (*Opuntia vulgaris*). The ripe fruit contained 68.2 per cent moisture and 1.76 per cent ash, and contained phosphate of aluminium.

Lindenmeyer, G. (Vierteljahressch. Prak. Pharm., 17, p. 290; Jahrb. Chem. Will., 1868, p. 814), reports 3.5878 per cent ash in China bark (*Cinchona*) from Puerto Cabello, containing 0.488 per cent alumina.

Huschke, O. (Vierteljahressch. Prak. Pharm., 17, p. 284; Jahresb. Chem. Will., 1868, p. 814), reports 1.21 per cent ash in orange peel. This contained 0.069 per cent alumina.

Loew, O. (The Physiological Rôle of Mineral Nutrients: U. S. Dept. Agr., Division of Vegetable Physiology and Pathology Bul. 18 (1899), p. 9). The possibility of the occurrence of aluminium salts in plants is frequently spoken of by botanical writers. Thus the author cited says regarding mineral compounds found in living organisms: "Occasionally there are present in plants small quantities of titanic and boracic acids, lithia, and alumina, and of the oxids of lead, zinc, and copper."

Luca, S. de (Compt. Rend. Acad. Sci. Paris, 51 (1860), p. 176), reported 6 per cent ash in Spanish moss (*Tillandsia dianthoidea*), an aerial parasite. This contained a trace of Al₂O₃.

Ludwig, H. (Arch. Pharm., 2d ser., 52 (1847), p. 61; Pharm. Centbl., 1848, p. 669; Jahresb. Chem., 1847-48, p. 1095), found 2.36 per cent ash in so-called "Caraccas" Sarsaparilla root, purchased in Honduras. Of this 5.11 per cent was alumina.

Ludwig, H. (Arch. Pharm., 3d ser., 1 (1872), p. 482; Jour. Chem. Soc. London, 26 (1873), p. 525), notes Al₂O₃ in the ash of different parts of the coffee tree as follows: Root of young tree in bearing, 7.85; root of old tree, 1.59; leaves, 9.11; pulp of pericup, trace; parchment-like coating of coffee bean, 4.19; coffee beans grown in gneiss soil, 2.78 per cent, and coffee beans grown on limestone soil, a trace.

MacDonald, J. W. (Chem. News, 37, p. 127; Jour. Chem. Soc. London, 1878, II, p. 624), examined the ash of cane and beet sugar accumulated from the analyses made in a year in a large sugar refinery, reporting among other constituents, 6.90 per cent ferric oxid and alumina in cane-sugar ash and 0.28 per cent in beet-sugar ash.

Maiden, J. H., and H. G. Smith (Jour. and Proc. Roy. Soc. New South Wales, 29 (1895), p. 325) report, in a silky oak log (Grevillea robusta), a large amount of a white substance having the appearance of whiting. Some six ounces were secured, but it is stated that a large proportion of the substance must have been lost and that there is no means of knowing the total quantity originally present in the cavity in the log. Analysis showed this material to consist of basic aluminium succinate. The authors regard it probable that the aluminium was originally present in the sap of the tree as a malate. No inorganic acid was detected in the sample analyzed.

Malaguti, F. J., and J. Durocher (Ann. Chim. et Phys., 3d ser., 54 (1858), p. 257, tables I to 4) report detailed studies of the ash of one hundred and fifteen wild plants of different families growing on a variety of soils in France. The oxids of iron, aluminium, and manganese are reported together. This work is of especial interest historically.

Mariani, G. (Studi chimico-agrari sugli equiseti, Studi e ricerche istituite nel R. Laboratorio di Chim. Agrar. di Pisa, 7 (1888), p. 69; Staz. Sper. Agrar. Ital., 14 (1888), p. 355; Just's Bot. Jahresber., 1888, pt. 1, p. 58), reports an analysis of a horse tail (Equisetum telmateja) grown in Pisa. This had a water content of 78.033 per cent. The CO₂ free ash (amount not given) contained 0.963 per cent Al₂O₃.

Matusow, H. (Amer. Jour. Pharm., 69 (1897), p. 341), reports aluminium as one of the constituents of the ash of broad-leaved laurel (Kalmia latifolia) root; amount not stated. The total ash was 1.24 per cent.

Mayer, A., (Lehrbuch der Agricultur Chemie 3d Ed., 1886, p. 280) states that aluminium is found in a seriesof plants as a regular constituent, and frequently in not subordinate amounts. In Lycopodium complanatum acetate of aluminium occurs in such quantities that a lye made from the plant can be used directly in dyeing as a mordant. This same salt has been found in wine grapes; and it is an assured fact that various parts of the grapevine on incineration leave in the residuum aluminium compounds. Vauquelin states that he found acetate of aluminium in the sap of the birch, which at best must be considered as an anomaly.

Minas, M. (Zur Frage über den Einfluss pasteurisierten Traubensaftes auf den allgemeinen Stickstoffumsatz, das Körpergewicht und die Darmfäulniss beim gesunden Menschen bei gemischter Kost. Inaug. Diss. Dorpat, 1900; Ztschr. Untersuch. Nahr. u. Genussmtl., 4 (1901), p. 204), analyzed a pasteurized grape juice (prepared by a Russian firm for medicinal use) and found that it contained 0.004 per cent of Al₂O₃.

Mitscherlich (Berlin. Acad. Ber., 1847, Nov., p. 430; Jour. Prak. Chem., 43, p. 158; Pharm. Centbl., 1848, p. 337; Inst., 1848, p. 186; Jahresb. Chem., 1847–48, p. 832) found in one of the marine Algæ (Conferva glomerata) (dried at 130°), 12.27 per cent ash; of this 0.42 per cent was alumina.

Moritz, J. (Ann. Oenologie. 4, p. 153; Just's Bot. Jahresber., 1874, II, p. 855), studied the draft on soil made by grapes of the variety "Sylvaner." The fresh wood contained 50 per cent more ash than the fresh fruit. He calculated that the grape plants withdrew 35.343 pounds mineral matter per morgen (0.63 acre) from the soil; of this 0.05 pound was Al₂O₃.

Muck (Vierteljahressch. Prak. Pharm., 5, p. 544; Chem Centbl., 1856, p. 848; Jahresb. Chem., 1856, p. 690) notes 12.5 per cent ash in the leaves of belladonna (Atropa belladonna) (dried at 100°) containing 0.01 per cent Al₂O₃.

Munroe, W. R. (Amer. Jour. Pharm., 70 (1898), p. 489), in an analysis of the rhizome of an aralia (Aralia californica), reports 2.22 per cent ash in which aluminium was present in the form of phosphate.

Nettlefold, P. (Chem. News, 55, p. 191; Jahresb. Chem., 1887, p. 2304), reports 0.571 per cent ash in an edible fungus (Bovista gigantaea). Of this 15.66 per cent was alumina.

Parodi, D. (Pharm. Jour. Trans., 3d ser., 10, p. 667; Jour. Chem. Soc. London, 37 (1880), p. 721), in a cucurbitaceous plant, tayuya (*Trianos perma ficifolia*), found 1.23 per cent iron and aluminium.

Payen (Compt. Rend. Acad. Sci. Paris, 28, p. 613; Pharm. Centbl., 1849, p. 516; Jahresb. Chem., 1849, p. 481) notes 0.12 per cent insoluble, and 0.16 per cent soluble ash in sugar cane. One of the ash constituents was alumina.

Peckolt, T. (Pharm. Jour. Trans., 3d ser., 10, pp. 343, 383; Ztschr. Oesterr. Apoth. Verein, 1879, pp. 361, 373; Jahresb.

Chem., 1879, p. 931), in a study of the fruit of paw-paw (Carica papaya), states that the dry flesh of the fruit of one variety, Mamao femea, contained 8.457 per cent ash. Of this 3.857 per cent was alumina.

Peckolt, T. (Wiener. Akad. Ber. 54 (2. Abt.) p. 462; Jahresb. Chem. Will., 1866, p. 708), reports 10.19 per cent ash in shells of seeds of Guarana (*Paulina sorbilis*); 1.704 per cent in seeds without shells and 2.6 per cent in Guarana paste; the last two contained respectively 1.06 and 0.82 per cent Al₂O₃.

Penney, M. D. (Chem. News; 39, p. 80; Jour. Chem. Soc. London, 36 (1879), II, p. 556), studied the occurrence of alumina in flour and wheat, finding, he states, larger quantities than are generally supposed to be present. Part of this he attributes to careless packing of the wheat. The Egyptian wheat noted below was largely contaminated with clay owing to the manner of packing in Nile boats.

In six samples of flour, he found alumina as follows, the amount representing grains in four pounds: No. 1, 24.30; No. 2, 21.75; No. 3, 21.25; No. 4, 17.00; No. 5, 12.40, and No. 6, 6.34.

The aluminium phosphate (milligrams per 100 grams) found in twenty-two samples of wheat was as follows: No. 1, Calcutta, 24.30; No. 2, Calcutta, 21.00; No. 3, Calcutta, 18.50; No. 4, Kourish, 27.20; No. 5, Kourish, 31.00; No. 6, Russian, 17.20; No. 7, Russian, 24.45; No. 8, Russian, 13.10; No. 9, Russian, 16.35; No. 10, Chicago, 4.00; No. 11, Oregon, 4.00; No. 12, English, 5.12; No. 13, English, 6.40; No. 14, English, 7.30; No. 15, English, 3.80; No. 16, Stein, 16.33; No. 17, Stein, 13.24; No. 18, California, 3.00; No. 19, Mixed (eleven varieties), 15.10; No. 20, Egyptian, as imported, 167.00; No. 21, Egyptian, hand-picked, 49.49; No. 22, Egyptian, washed, 14.10.

Percival, J. (Agricultural Botany, New York, 1900, p. 167), notes the occasional presence of alumina in small amounts in certain plants. In sea-weeds bromine and iodine, he states, are present, and many other elements such as aluminium, zinc, and copper have been occasionally discovered in small quantities in certain species of plants.

Petter, K. (Vierteljahressch. Prak. Pharm., 11, p. 545; Jahresb. Chem., 1862, p. 511), found 23.15 per cent ash in one of the common Algæ (Cladophora glomerata) (dried at 110°). Of this 0.225 per cent was Al₂O₃.

Pfeffer, W. (Pflanzen-Physiologie, 2d ed., 1897, Vol. I, p. 432), in discussing the nutrients of plants, makes in effect the following statement: Aluminium, although it is so generally distributed, occurs in the ash of most plants only in small amounts. In the case of Lycopodium chamaecyparissus and alpinum, it constitutes 22 to 27 per cent of the ash. It is also abundantly present in Chlorangium jussuffii according to Wolff (Aschen Analysen, 1871, pp. 134, 136), while Church found only traces of it in Lycopodium phlegmaria, in Selaginella, etc. Whether or not aluminium occurs in Lycopodium, as Arosenius states in the form of tartrate, is not definitely known.

Pizzi, A. (Staz. Sper. Agr. Ital., 16 (1888), p. 737; 17 (1889), p. 1; Centbl. Agr. Chem., 19 (1890), p. 353; Jahresb. Agr. Chem., 1890, pp. 287, 288), reports 1.80 per cent ash in white truffle (*Tuber magnatum*) from the Apennines, of which 6.9 per cent was Al₂O₃. In black truffle (*T. melanosporum*), from the same region, he found 2.09 per cent ash of which 5.3 per cent was Al₂O₃.

Pizzi, A. (Staz. Sper. Agr. Ital., 17, p. 167; Jahresb. Chem., 1889, p. 2109), reports 1.342 per cent ash in the morel (Morchella esculenta). Of this 3.120 per cent was Al₂O₃.

Ramdohr (Arch. Pharm., 2d ser., 91 (1857), p. 129; Ztschr. Pharm., 1857, p. 70; Chem. Centbl., 1857, p. 705; Jahresb. Chem., 1857, p. 515) notes 3.62 per cent ash in air-dry ergot (Secale cornutum) from cheat or chess (Bromus secalinus) and 2.91 per cent in ergot from rye (Secale cereale). These contained respectively 1.09 and 0.33 per cent Al₂O₃.

Reichardt, E. (Arch. Pharm., 2d ser., 73 (1853), p. 257; Pharm. Centbl., 1853, p. 267; Jahresb. Chem., 1853, p. 581), examined the leaves, bark, and wood of willow (Salix vitellina) in autumn and spring, reporting (per 1000 parts fresh material) 0.05 Al₂O₃ in autumn leaves, and 0.03 in autumn bark and wood, 0.03 in spring leaves, and 0.06 in spring bark and wood. He found also (Arch. Pharm., 2d ser., 75 (1853), p. 19; Pharm. Centbl., 1853, p. 566; Jahresb. Chem., 1853, p. 584) 0.39 and 0.46 Al₂O₃.PO₅ per 1000 parts dry matter in thick

(old) and thin (young) bark, from a willow, about five years old, gathered near the end of October, 1852.

Reichardt, E. (Chem. Bestandtheile der Chinarinden, Braunschweig, 1855; Pharm. Centbl., 1855, p. 631; Ztschr. Pharm., 1856, p. 25; Jahresb. Chem., 1855, p. 720), reports ash in the following drugs: China flava fibrosa (this and all the others dried at 100°) 1.6338 per cent; C. rubra, 1.6607 per cent; C. huanuco, 2.5134 per cent; C. regia sine epid., 1.2236 per cent; C. regia cum epid., 1.6449 per cent. He reports aluminium phosphate in these as follows: 0.0473, 0.0528, 0.0467, 0.0465, and 0.0403, or re-calculated and expressed in percentage of total ash (according to abstracts for Jahresb. Chem.) 1.203, 1.324, 0.771, 1.580, and 1.017 per cent alumina.

Reithner (Vierteljahressch. Prakt. Pharm., 4, p. 382; Jahresb. Chem., 1855, p. 723) notes 4.3 per cent ash in the leaves of holly (*Ilex aquifolium*) dried at 100°. Of this 0.66 per cent was Al₂O₃.

Reithner (Vierteljahressch. Prakt. Pharm., 4, p. 481; Pharm. Chem., 1855, p. 834; Jahresb. Chem., 1855, p. 724) notes 4.05 per cent ash in the dry flowers, without anthers, of colchicum or autumn crocus (*Colchicum autumnale*) and 4.15 per cent in the dry anthers. The ash contained respectively 0.27 per cent and a trace of Al₂O₃.

Ricciardi, L. (Gaz. Chim. Ital., 19 (1889), p. 150), studied the aluminium content of the ash of different portions of several plants. He found that the ash of grape-vines grown on clay soil contained 0.85 per cent alumina; on soil containing an abundance of lime, o.81 per cent; and on soil with a medium lime content, 1.14 per cent. He also found 0.218 per cent alumina present in the ash of the branches and twigs, 0.093 in the skins and seeds, and 0.022 in the leaves of the mandarin orange. The ash of Indian fig fruit, "prickly pear" from Bari, contained 0.063 per cent; from Catania, 0.092. The ash of the rind of the prickly-pear fruit from Bari contained 0.148 per cent; from Catania, 0.167. The ash from the flesh of St. John's breadfruit (Ceratonia siliqua) from Bari contained 0.503 per cent; from Catania, 0.607; the ash of the seeds from Bari, 0.062; from Catania, 0.064. The ash of almond shells grown near Bari contained 0.695 per cent Al₂O₃, the ash of the kernels

o.138. The ash of tobacco leaves contained 2.151 per cent alumina. The seeds and skin of mandarin oranges grown near Cara contained in the ash o.121 per cent Al₂O₃, and white lupine ash o.042 per cent Al₂O₃. Further, wine was found to contain o.022 gm. Al₂O₃ per liter. The author concludes that the assimilation of alumina does not depend on the percentage in the soil, and that, generally speaking, alumina is most abundant in the trunk and branches, less so in the husks and seeds, and least of all in the leaves.

Ringer, F. A., and E. Brooke (Amer. Jour. Pharm., 64 (1892), p. 255) report 4.29 and 1.32 per cent ash in roots of true and commercial pareira respectively. Both contained Al₂O₃; the amount not stated.

Ritthausen, H. (Jour. Prakt. Chem., 53 (1851), p. 413) reports 36.250 per cent Al₂O₃ in the pure ash of a club-moss (*Lycopodium complanatum*).

Ritthausen, H. (Jour. Prakt. Pharm., 58, p. 133; Pharm. Centbl., 1853, p. 531; Jahresb. Chem., 1853, p. 586) reports 39.07 and 20.69 per cent Al₂O₃ in ash of club-mosses (*Lycopodium chamaecyparissus* and *L. clavatum*). The amount of total ash was not stated.

Rode, C. (Vierteljahres. Prakt. Pharm., 9, p. 530; Jahresb. Agr. Chem., 1861-62, p. 61), reports 0.15 per cent Al₂O₃ in the ash of the leaves of cuckoo-pint (*Arum maculatum*). The leaves contained a total of 88 per cent water; the water-free material, 10.12 per cent ash.

Rotondi, E. (Rel. lav. esequiti. lab. Chim. R. Staz. enol. Asti, 1878, p. 111; Centbl. Agr. Chem., 8 (1879), p. 530), studied the ash constituents of grape must, branches, and leaves from two vineyards near Asti. In every case the vine and leaves were selected after the grape harvest from vines which had borne fruit. The branches of Pinot grapes from Galletria d'Asti contained 2.91 per cent pure ash in dry matter, of which 0.13 per cent was Al₂O₃. Branches of Barbara grapes from Costigliole d'Asti contained 3.37 per cent pure ash in dry matter; of this 0.02 per cent was Al₂O₃. Traces of alumina are also reported in branches and leaves of Barbara and Grignolino grapes from Galleria d'Asti and in must of Barbara and must and branches from Fresia grapes from Costigliole d'Asti.

Ruge, G. (Apoth. Ztg., 1891, p. 208; Jahresb. Agr. Chem., 1891, p. 244), states that floating buttercup (Ranunculus fluitans) contains 14.349 per cent ash in the dry matter; 0.33 per cent of this being Al₂O₃.

de Saussure, T. (Recherches Chimiques sur la Végétation, Paris, 1804) discusses the ash constituents of plants, and in tables at the end of the volume reports the ash constituents of a considerable number. From these tables the following is quoted: In the leaves and other portions of a large number of trees and plants the amount of aluminium in the ash was less than 0.01 per cent. The leaves of a rhododendron (Rhododendron ferrugineum) contained 3 per cent ash in the dry matter; of this 0.12 per cent was alumina. Another sample contained 2.5 per cent ash in the dry matter, 0.12 per cent of this also being alumina. The branches of the same plants contained 0.8 per cent dry matter; of this 0.12 per cent was alumina. Saussure's investigation is interesting chiefly from an historical standpoint.

Sayre, L. E. (Amer. Jour. Pharm., 69 (1897), p. 543), reports a study of the composition of dandelion (*Taraxacum*) root. A sample dried at 100° contained 11.13 per cent ash. Of this 18.07 per cent was Al₂O₃.

Schäzler, M. (Vierteljahressch. Prakt. Pharm., 11, p. 270; Jahresb. Chem., 1862, p. 514), found 5.12 per cent ash in the finest Ceylon cinnamon bark (dried at 110°). Of this 0.55 per cent was Al₂O₃.

Schlegel, C. E. (Amer. Jour. Pharm., 57 (1885), p. 426), reports the analysis of the fruit of star anise (*Illicium anisatum*). The seed capsules contained 10.36 per cent moisture and 3.5 per cent ash. This was made up, according to the author, of potassium, sodium, iron, aluminium, hydrochloric acid, and phosphoric acid.

Schreiner (Vierteljahressch. Prakt. Pharm., 5, p. 207; Jahresb. Chem., 1856, p. 691), reports 0.72 per cent ash in the fresh fruit of sloe or black thorn (*Prunus spinosa*). Of this 0.5 per cent was Al₂O₃.

Schriddl, P. (Arch. Pharm., 1873, p. 375; Centbl. Agr. Chem., 5 (1874), p. 50), reports the percentage of Al₂O₃ in the ash of four samples of tea leaves and two of wood as follows: 0.810, 1.015,

2.226, 2.072, 4.093, and 2.834. They were from Java and in the case of Nos. 3, 4, and 6 were affected with a tea disease.

Sestini, F. (Gaz. Chim. Ital., 4 (1874), p. 182; Jahresb. Chem., 1874, p. 906), notes a trace of alumina in the ash of a seaweed (*Posidonia oceanica*) which is used as a fertilizer.

Sestini, F. (Gaz. Chim. Ital., 15 (1885), p. 107; Landw. Vers. Stat., 32 (1886), p. 197; Jahresb. Chem., 1885, p. 1848), studied the relation between the atomic weight and the physiological functions of the elements. He says in effect regarding certain elements, "Many elements which are not yet regarded as necessary for plants are found in plant ash, as aluminium, lithium, and fluorine." He believes that no element with an atomic weight over 56 enters directly into the formation of living material.

Sestini, F. (Stud. e Ricer. Inst. Lab. Chim. Agr. Pisa, 6, p. 87; Chem. Centbl., 1887, p. 939; Jahresb. Chem., 1887, p. 2647), determined the aluminium content of a number of wines. He found per 1000 cc. in ordinary red wine, from Peccioli (1884), 0.017 gm.; in ordinary red wine from Ghezzano (1884), 0.034 gm.; in red wine from Calci (1884), 0.038 gm.; from Collesalvetti (1883), 0.006 gm.; from Collesalvetti (1884), 0.002 gm.; from S. Giuliano (1884), 0.005 gm.; and from S. Giuliano (1885), 0.009 gm. He concludes, therefore, that a natural wine does not contain more than 0.03 gm. alumina per liter and that alumina will never exceed 0.2 per cent of the pure ash.

Shepard, C. U. (Trans. N. Y. State Agr. Soc., 1844, p. 343), reports traces of alumina in the ash of rice straw. He also reports traces in the ash of corn (grains) as quoted by J. H. Salisbury (Trans. N. Y. State Agr. Soc., 1848, p. 726).

Smith, H. G., in an article entitled "Aluminium, the Chief Inorganic Element in a Proteaceous Tree, and the Occurrence of Aluminium Succinate in Trees of this Species" (Chem. News, 88 (1903), p. 135), reports the occurrence of a large deposit of basic aluminium succinate of the formula Al₂(C₄H₄O₄)₃Al₂O₃ in the center of a silky oak tree 3 feet in diameter (Orites excelsa) from Queensland. The ash of the wood farthest from the deposit contained 79.61 per cent alumina. The large deposit was regarded as a natural effort to get rid of a surplus of alumina not needed by the tree.

Three other samples of the same kind of tree from northern New South Wales were examined and in the ash of all a large amount of alumina was found, the quantity ranging from 36 to 43 per cent. Much of the alumina in the ash was present as a potash salt soluble in water, and as no carbonate of potash was found "it is supposed that the potassium aluminate was originally present in the tree as such."

As no alumina was found in the ash of *Grevillea robusta*, G. helliana, and G. striata, the author believes that it is probable that the tree reported to contain it in an earlier article by J. H. Maiden and himself (see page 26) was O. excelsa and not G. robusta, as stated at the time.

Smith, A. Percy (Chem. News, 28, p. 261, 324; Jahresber. Chem., 1873, p. 851), reports (by difference) 1.459 per cent alumina and lithium carbonate in the ash of Havana tobacco.

Smith, Watson (Jour. Chem. Soc. London, 37 (1880), II, p. 416), analyzed the ash of two varieties of eucalyptus wood. *E. rostrata* contained 2.25 per cent ash and *E. globulus* 2.01 per cent. The former contained 0.78 per cent ferric and aluminic phosphate, the latter 1.07 per cent. In the latter a trace of Al₂O₃ was also noted.

Snyder, H. (Chemistry of Plant and Animal Life, 1903, Easton, Pa., p. 168), says: "Aluminium is found in the ash of many plants, as wheat, peas, beans, and rice, although it occurs in very small amounts and, so far as known, is not essential for plant growth. Most soils contain traces of soluble silicates of aluminium, and hence plants cannot well be free from it."

Solms-Laubach, Graf (Ann. Chem. u. Pharm., 100 (1856), p. 297), reports 2 per cent Al₂O₃ in the ash of a lycopod (*Lycopodium denticulatum*).

Speiss, E. (Wittstein's Vierteljahresschrift, 9, p. 392; Jahresber. Agr. Chem., 1860–61, p. 59), analyzed the ash of the bark of the root of the pomegranate and the root-stalks of male fern. The ash of the former contained a trace of Al₂O₃ and that of the latter 0.071 per cent.

Staffel, E. (Preisschrift. Arch. Pharm., 2d ser., 64, I, p. 129; Ann. Chem. Pharm., 76, p. 379; Pharm. Centbl., 1850, p. 897, 1851, p. 146; Jahresb. Chem., 1850, p. 661 and Table D, p. 661),

in a study of the question whether the inorganic constituents of different organs of a green plant varied at different seasons, reports alumina in ash as follows: Horse-chestnut wood (autumn), 0.23 per cent; bark, 0.18 per cent; leaves (spring), 0.41 per cent; leaves (autumn), 0.51 per cent. The total ash in the above (dry material) was 3.38, 6.57, 7.69, and 7.52 per cent respectively. He also reports alumina in ash of English walnut bark (autumn), 0.29 per cent; leaves (spring), 0.18 per cent; leaves (autumn), 0.06 per cent. The total ash in the above (dry material) was 6.403, 7.719, and 7.005 per cent respectively.

Stoklasa, J. (Ann. Agron., 23 (1897), p. 588), in connection with a study of the condition of phosphoric acid in the soil made laboratory experiments with salts of known composition which led to the conclusion that the water-soluble portion of superphosphates never contains acid ferrous phosphate, this salt changing almost immediately upon its formation into di-tri-ferriphosphates of varying composition insoluble in water. The addition of ferrous salts to soluble phosphates results in the formation of di-tri-ferriphosphates unless an excess of free phosphoric acid is present. Acid ferric phosphate may be found in superphosphates only when there is at least 30 per cent of free phosphoric acid present. If this is not the case, the acid ferric phosphate may be transformed into mono-di-tri-ferriphosphate, Fe₂O₃(P₂O₅)₂.8H₂O.

"It is thus seen that the retrogression of phosphoric acid in superphosphate is very largely dependent upon the free phosphoric acid present. Aluminium salts do not behave like iron salts in superphosphate, but like the salts of lime and magnesia.

"Pot experiments with barley on a fertile soil containing o.63 per cent calcium carbonate and 2 per cent of humus and treated with different phosphates of calcium, aluminium, and iron showed that the effect of the acid phosphates of these elements was almost the same. The tribasic phosphates were about one-half as effective as the acid phosphates. The effect of the normal reverted phosphates was about one-half greater than that of the tribasic phosphates."

Other conclusions regarding phosphates were drawn.

Strawinski, J. Frank (Amer. Jour. Pharm., 70 (1898), p. 189), finds that the ash of the rhizome and rootlets of plantain (*Plantago major*) amounted to 24.70 per cent and contained alumina in addition to other constituents.

Strohecker, R. (Arch. Pharm., 2d ser., 145, p. 131; Jour. Chem. Soc. London, 24 (1871), p. 428), in an article on chemical substitution in plants discusses analyses by Wittstein of water from the Ohe and Isar rivers and of the ash of Fontinalis antipyretica taken from these waters. The river waters contained respectively 0.016 and 0.2250 per cent total solids; in that of the Ohe water 0.108 per cent was Al₂O₃. The plants from the two rivers contained 22.60 and 9.88 per cent ash respectively, of which 9.272 and 1.616 per cent was Al₂O₃.

Struckmann (Ann. Chim. Pharm., 97, p. 143; Jour. Prakt. Chem., 68, p. 379; Chem. Centbl., 1856, p. 187; Jahresb. Chem., 1856, p. 687) found 9.2 per cent ash in the root (Wedel) of male fern (Aspidium filix mas) (dried at 100°), and 8.1 per cent in that of asplenium (A. filix femina), the ash containing respectively 2.40 and 2.20 per cent Al₂O₃.

Teller, G. L. (Arkansas Station Bulletin 42, pp. 75, 77), reports complete analyses of whole wheat, wheat flour, and other milling products, the samples being all obtained from the same lot of winter wheat grown in Arkansas. The data regarding alumina follow:

ALUMINA IN ASH OF WINTER WHEAT AND ITS MILLING PRODUCTS.

		Straight Flour.		Dust- room.	Ship- stuff.	Bran.	Wheat.
Total ash		Per Ct. 0.40	2012				
	.41	.15	. I 2	.04	.18	.07	.11

The presence of alumina in wheat has been attributed to the wearing down of the millstones. On this point Professor Teller says:

"This could not have been a source of the material in these mill products, as the wheat was crushed entirely by iron rollers, and an examination of the amounts of alumina found in the mill products and in the whole grain indicate that it is no more foreign to the true ash than any of the other constituents named. To bring further proof on this point, 100 gms. of the unground wheat was carefully washed with distilled water and, after drying, was burned without being pulverized. The same amounts of both alumina and zinc were found as in the wheat which had not been washed. It seems a little remarkable that the zinc should have accumulated to the greatest extent in the ash of the bran, while the alumina and silica should have reached their largest proportion in the ash of the finer flours. Alumina is found to be of frequent occurrence in the mineral waters of this State."

To ascertain whether alumina is present in the ash of wheat grown on a very sandy soil, a sample of wheat was obtained which had grown on a sandy soil in Michigan. This was examined for alumina and none was found

Thielau (Vierteljahresschr. Prakt. Pharm., 4, p. 537; Pharm. Centbl., 1855, p. 811; Jahresb. Chem., 1855, p. 728) found 3.33 per cent ash in ergot (Secale cornutum) dried at 100°. Of this 0.29 per cent was Al₂O₃.

Thomas, Mason B. (Proc. Indiana Acad. Science, 1893, p. 239), in a report on the study of the ash of trees makes the following statements:

"The substances usually found in the ashes of all trees when burned at a low temperature are potash, soda, lime, magnesia, and iron (K₂O, Na₂O, CaO, MgO, Fe₂O₃) in combination with phosphoric acid (P₂O₅), sulphuric acid (SO₃), chlorine (Cl), carbon dioxide (CO₂), and silica (SiO₂); iodine (I), aluminium (Al), and manganese (Mn) are often present. One portion of these mineral constituents exists in solution in the sap and the other in the tissue of the plant in the solid form."

The author reports the ash constituents usually determined, but does not report the amount of aluminium present. His experiments were made with common forest and fruit trees.

Tod, W. (Arch. Pharm., 2d ser., 78 (1854), p. 136; Jour. Prakt. Chem., 62, p. 503; Pharm. Centbl., 1854, p. 452; Jahresb. Chem., 1854, p. 665), found that the flesh of plum (*Prunus domestica*) made up 93 per cent of the fruit and the pit 7 per

cent. The fresh flesh contained 0.407 per cent ash, 0.003 per cent being Al₂O₃.

Tollens (Jour. Landw., 50 (1902), p. 231) states that aluminium is always present in very small quantities in grain.

Trimble, H., and H. J. Schuchard (Amer. Jour. Pharm., 57 (1885), p. 21) report the analysis of the tops and leaves of water pepperwort (*Polygonum hydropiper*). The ash constituted 7.4 per cent of the total plant and contained aluminium phosphate.

Tschirch, A. (Gartenzeitung, 1883, p. 34), reports in hyacinth (Hyacinthus orientalis) 10.11 per cent dry matter and 8.5787 per cent pure ash. The Al₂O₃ in the ash was 0.8871 per cent.

Tucker, G. M., and B. Tollens (Ber. Deut. Chem. Gesellsch., 32 (1899), p. 2575; Jour. Chem. Soc. London, 78 (1900), II. p. 35) note Al₂O₃. Fe₂O₃ in plane-tree leaves.

Vielguth (Vierteljahresseh. Prakt. Pharm., 5, p. 187; Chem. Centbl., 1856, p. 423; Jahresb. Chem., 1856, p. 690) notes 7.6 per cent ash in bed straw (Galium mollugo) (whole plant). This contained 0.3 per cent Al₂O₃.

Völcker, A. (Rpt. 19th Meeting British Association; Notes and Abstracts, p. 43, Jahresb. Chem., 1850, p. 672 and Table D, p. 661), notes 1.97 per cent alumina in the ash of Armeria maritima (green plant?). The amount of total ash was not stated.

Wallace (Analyst, 3 (1878), p. 243) reports that the ash of sugar made from cane growing near the seacoast in Demerara contained, in addition to other constituents enumerated, 0.65 per cent aluminium. The amount of total ash is not stated.

Walz (Jahrb. Prakt. Pharm., 15, p. 65; Jahresb. Chem., 1847-48, p. 1083; and Table B, p. 1074) reports ash as follows in grapevine and leaf: Clevner, 6.19 per cent; Riesling, 7.74 per cent; Drollinger, 6.00 per cent. Of this 1.15, 0.91, and 1.00 per cent respectively was aluminium phosphate.

Wanklyn, J. A., and W. J. Cooper (Bread Analysis, London, 1886, pp. 22, 24, 25) state that according to their analyses the ash of flour contains 1.3 per cent Fe₂O₃ and Al₂O₃ together. In eight samples of flour analyzed, 100 gms. of material yielded from 0.01 to 0.02 gm. of phosphate of iron and alumina. In

eight samples of bread, the ash in 100 gms. of material ranged from 0.890 gm. to 1.742 gms.; the phosphates of iron and alumina from 0.006 gm. to 0.014 gm.

Warden, C. J. H. (Chem. News, 38, p. 146; Jahresber. Agr. Chem., 1878, p. 106), reports a trace of Al₂O₃ in the ash of opium from Behar, India.

Warden, C. J. H. (Chem. News, 39, p. 27; Jahresb. Chem., 1879, p. 927), reports 0.9701 per cent Al₂O₃ in the ash of poppy leaves or petals used to wrap up opium. The total amount of ash is not stated.

Warden, C. J. H. (Chem. News, 64 (1891), p. 161), reports 24.334 per cent ash in the leaves (dried at 100°) of prickly chaff flower (*Achyranthes aspera*), an Indian weed. Of this 2.0651 per cent was Al₂O₃. The author says that there was considerable soil which it was not possible to separate from the leaves. This would, of course, increase the Al₂O₃ content.

Wasowicz, D. v. (Pharm. Jour. Trans., 3d ser., 10, pp. 301, 341, 463; Arch. Pharm., 3d ser., 14 (1879), p. 193; Jahresb. Chem., 1879, p. 927), notes the presence of aluminium in the ash of the root of an aconite (Aconitum heterophyllum). The total ash was equal to 2.331 per cent. The root of another aconite (Aconitum japonicum) contains 2.799 per cent ash. Aluminium is reported as a constituent of this also.

Watts, H. (Phil. Mag., 3d ser., 32, p. 54; Jahresb. Chem., 1847-48, p. 1077; and Table A, p. 1074), reports 6.5 per cent ash in hop blossoms; of this 1.18 per cent was alumina.

Way, J. T., and G. H. Ogston (Jour. Roy. Agr. Soc., 11, pt. 2, p. 497; Jahresb. Chem., 1850, p. 666, Table A, p. 660) note 1.90 per cent ash in rye (dry material). Of this 0.50 per cent was alumina.

Weber, R., and E. Ebermayer ("Lehre der Waldstreu," Berlin, 1876. Cited by Wolff in "Aschen Analysen," Berlin, 1880, pt. 2, pp. 87, 130) determined the Al₂O₃ in fir litter, i.e. needles, etc., under fir trees. The pure ash in eleven analyses ranged from 1.07 to 2.00 per cent. From 1.09 to 9.98 per cent of the total pure ash was found to be Al₂O₃.

Weber and Ebermayer ("Lehre der Waldstreu," Berlin, 1876. Cited by Wolff in "Aschen Analysen," Berlin, 1880, pp. 100, 131) also give figures showing that the pure ash of white pine litter ranges from 1.99 to 5.27 per cent. Of this 1.44 to 7.02 per cent was found to be Al₂O₃.

Wefers-Bettink, H. (Rev. Trav. Chim. Pays-Bas, 2, p. 126; Jahresb. Chem., 1883, p. 1496), analyzed "Lěgèn," a Japanese drug, which has been said to consist largely of the excrement of an insect, "Dendang." He concluded that it was of vegetable origin and probably prepared from the seed of some strychnin-yielding plant. Lěgèn contained 16.88 per cent ash, of which Al₂O₃ was one of the constituents.

Weigelt, C. H. (Jour. Prakt. Chem., 106 (1869), p. 193), reports 10.5 per cent pure ash in one specimen of *Parmelia scruposa*; of this 28.171 per cent was found to be Al₂O₃.

Wellborn, G. (Pharm. Jour. Trans., 3d ser., 9, p. 181; Jour. Chem. Soc. London, 1878, II, p. 1009), in a paper on the detection of alum in flour states that too much allowance has been made for alumina in wheat, as it is not an invariable constituent.

Wheeler, C. Gilbert (Erdman's Journal, 94, p. 385; Jahresb. Agr. Chem., 1865, p. 114), reports traces of Al₂O₃ in the ash of nine varieties of Bavarian hops.

Will and Fresenius (Chevallier et Baudrimont, "Dictionnaire des alterations," Paris, 1878, p. 1131) note the occurrence of alumina in the ash of Hungarian tobacco.

Winternitz (Vierteljahressch. Prakt. Pharm., 4, p. 542; Pharm. Centbl., 1855, p. 820; Jahresb. Chem., 1855, p. 722) found 8.88 per cent ash in dandelion (*Leontodon taraxacum*) (whole plant without root, dried at moderate temperature). Of this 0.402 per cent was Al₂O₃.

Witting (Keller and Tiedemann's Nord. American Monatsber. Pharm. Chem., 1851, p. 404; Jahresb. Chem., 1851, p. 712, Table C, p. 708) reports in ash of birch "No. 76" 1.38 per cent alumina and in "No. 74" 0.42 per cent, and in ash of beech "No. 75" 0.05 per cent. The amount of total ash is not stated.

Wittstein (Repert. Pharm., 2d ser., 46, p. 329; Pharm. Centbl., 1847, p. 739; Jahresb. Chem., 1847–48, p. 1082; and Table A, p. 1074) reports 1.56 per cent ash in green leaves of Virginia creeper (Vitis hederacea); of this 0.07 per cent was alumina.

Wittstein (Vierteljahressch. Prakt. Pharm., 3, p. 10; Pharm. Centbl., 1854, p. 12; Jahresb. Chem., 1853, p. 579) reports 1.13, 1.98, and 2.92 per cent ash in the bark of pine trees (Scotch

pine) (*Pinus silvestris*), 220, 172, and 135 years old, respectively, which grew on the sandy soil of the "Hauptmoor," near Bamberg, the ash containing respectively 10.12, 4.49, and 3.08 per cent Al₂O₃. The wood from the trunks of these trees contained 0.45, 0.58, and 0.42 per cent ash with respectively 0.92, 0.72, and 0.26 per cent Al₂O₃.

Wittstein (Vierteljahressch. Prakt. Pharm., 4, p. 525; Jahresb. Chem., 1855, p. 722) reports two analyses (the first by Nutzinger; the second by Thielau) of the ash of heather (Calluna vulgaris) (whole plant, without root, in bloom): sample "a," grown on moorland, contained, air-dry, 2.876 per cent ash with 0.844 per cent Al₂O₃, and sample "b," grown on sandstone, contained, air-dry, 3.324 per cent ash with 0.513 per cent Al₂O₃.

Wittstein, G. C. (Arch. Pharm., 2d ser., 111, p. 14; Jahresb. Chem., 1862, p. 509), reports the following:

ALUMINA IN CERTAIN PLANTS.

Kind of Plant.	Total Ash.	Al ₂ O ₃ in Ash.
Reindeer moss (Cladonia rangiferina) dried at	Per Cent.	Per Cent.
160°; from Riefs, near Passau	1.18	1.805
berge	1.325	1.694
hood of Sterz Lichen (Variolaria dealbata) from Bärenstein-	0.905	1.948
leithe, near Grafenau; not well cleaned	18.20	7 - 495
Old man's beard (Usnea barbata) from Grafenau	1.426	1.653
Lichen (Gyrophora pustulata) from Veitsberge Iceland moss (Cetraria islandica) from Bavarian	3.00	4.069
forest	0.8	4.348
Upper Bavaria. White birch (Betula alba) from Bavarian forest;	2.014	2.834
wood dried at 100°	0.864	0.663
leaves dried at 100°	3.958	0.287
Beech (Fagus sylvatica) from Bavarian forest; wood	0.74	0.508
leaves	4.3	0.162
wood pumilio) from Bavarian forest;	0.284	0.124
wood. Dwarf pine (Pinus pumilio) from Bavarian forest; bark.	1.375	0.240

These plants are discussed in relation to the material (rock or soil) on which they grew.

Wittstein (Vierteljahressch. Prakt. Pharm., 13, p. 364; Jahresb. Agr. Chem., 1865, p. 108) quotes an analysis showing that fresh fig leaves from plants grown on calcareous soil contain 0.286 per cent ash, 0.03 per cent of this being Al₂O₃.

Wittstein, G. C. (Ann. Chem. u. Pharm., 108, p. 203; Chem. Centbl., 1858; Jahresb. Agr. Chem., 1858-59, p. 121), reports the ash content of the different parts of a primrose (*Primula farinosa*). The following figures are quoted, being for material dried at 100°:

Root, 1.617; leaves, 0.955; stem, 0.539; flower heads, 1.145; and entire plant, 0.832 per cent Al₂O₃ respectively; or, differently stated:

Of the 0.832 per cent Al₂O₃ in the entire plant, 0.175 per cent was in the root, 0.198 per cent was in the leaves, 0.294 per cent was in the stem, and 0.165 per cent was in the flower head.

Wittstein, G. C. (Vierteljahressch. Prakt. Pharm., 16, p. 81; Jahresb. Chem., 1867, p. 769), reports alumina as follows in leaves and flowers of lilac (Syringa vulgaris), all dried at 110°.

ALUMINA IN DIFFEREN	NT PARTS OF LILAC.
---------------------	--------------------

Part of Plant.	Ash.	A12O3.
Leaves (variety with white blossoms) Leaves (variety with purple blossoms)	Per Cent. 4.389 4.922	Per Cent. 0.195 .188
Flowers (without calyx, variety with white blossoms)	5.76	.135

Wittstein, G. C. (Arch. Pharm. 3d ser., 8, p. 341; Jahresb. Agr. Chem., 1875-76, p. 139; Jour. Chem. Soc. London, 1877, I, p. 487), reports 5.936 per cent total ash in a spurge (Euphorbia amygdoloides) (whole plant) grown on soil rich in silicate and 4.85 per cent in another sample from similar soil. The ash of the first contained 1.057 and of the latter 1.325 per cent Al₂O₃.

Wittstein, G. C. (Arch. Pharm., 3d ser., 7 (1875), p. 394; Jour. Chem. Soc. London, 1876, I, p. 736), finds that the ash of the elder tree (Sambucus nigra) contains 0.250 per cent Al₂O₃.

Wittstein also reports (Arch. Pharm., 3d ser., 8 (1876), p. 229; Jahresb. Agr. Chem., 1875-76, pp. 132, 138) 94 per cent total dry matter in sunflower seeds (*Helianthus annuus*), the ash containing 0.23 per cent Al₂O₃ and 1.9 per cent ash, with 0.28 per cent Al₂O₃ in the whole plant.

Wittstein, G. C. (Arch. Pharm., 3d ser., 8 (1876), p. 342; Jahresb. Agr. Chem., 1875-76, p. 139), reports 7.132 per cent ash in *Herniaria glabra* (dry plant) grown in silicious soil and 6.622 per cent in plants grown on dolomite soil. The former contained 1.321 and the latter 1.755 per cent Al₂O₃.

Wolff, E. (Aschen Analysen, pt. 2, p. 128), cites figures showing that heather contains 2.08 per cent pure ash. It is stated that on an average 0.33 per cent of the total ash is Al₂O₃, the range in eleven analyses being from 0 to 2.3 per cent. Wolff aslo states (Aschen Analysen, p. 128) that in moss (the variety not given) the pure ash constituent is equal to 2.74 per cent on an average. Of this Al₂O₃ constitutes 2.35 per cent, the range in eleven analyses being from only 0 to 7.99 per cent. Wolff also states (Aschen Analysen, p. 128) that the pure ash content of lycopodium is 5.10 per cent, 39.17 per cent of this being Al₂O₃, the range of this constituent in six analyses being from 22.2 to 57.36 per cent.

Some of the data cited by Wolff were undoubtedly from early analyses, interesting chiefly from an historical standpoint.

Wolff, J. (Vierteljahressch. Prakt. Pharm., 3, p. 1; Jahresb. Chem., 1853, p. 562), found 1.2 per cent ash in air-dry birch fungus (*Birken schwamms*). Of this 3.14 per cent was Al₂O₃. The plant analyzed was without doubt the rough boletus (*Boletus scaber*), sometimes called Birken-pilz.

Wypfel, M. (23 Jahresb. Niederöster. Landes Realgym. Waldhofen, 1892, p. 22; Just's Jahresb. Bot., 20 (1892), I, 1, p. 425), studied the effect of chlorids on plant growth. Seedlings of corn (Zea), bean (Phaseolus), pea (Pisum), cucurbit (Cucurbita), sunflower (Helianthus), beet (Beta), onions, and Hartwegia comosa were watered daily with 0.5 to 2.0 per cent solutions of different chlorids. Among the conclusions drawn were the following: Chlorids which are widely distributed and which contain important material for plant growth (magnesium, calcium, potassium, and aluminium chlorid) at first had a favorable effect. Later, when they became more concentrated in the soil, they hindered growth and finally killed

the plants. Magnesium chlorid was the least harmful, potassium and aluminium chlorids most harmful.

Yardley, H. B. (Chem. News, 79 (1899), p. 122; Jour. Chem. Soc. London, 76 (1899), II, p. 793), found 4.19 per cent ash in cardamon seeds and husks. Of this 1.53 per cent was Al₂O₃.

Yoshida, H. (Jour. Chem. Soc. London, 43 (1883), p. 472), reports a chemical study of lacquer (*Urushi*), the milky secretion of lac (*Rhus vernicefera*). Gum constitutes from 3 to 8 per cent of the original juice. A sample weighing 0.5267 gm. gave 0.0267 gm. ash. The ash was found to contain in addition to other constituents 7.85 per cent Al₂O₃.

Yoshida (Jour. Chem. Soc. London, 51 (1887), pp. 748-750) takes exception to the idea that alumina is not a normal constituent of flowering plants, as stated by Allen (Commercial Organic Analysis, vol. 1, p. 38). In the Japanese lacquer tree (Rhus vernicefera) alumina is present, apparently as an arabate.

According to the author the soil of the plain of Musashi, on which Tokyo is situated, is of volcanic origin and is remarkable for the large proportion of alumina in it which is soluble in hydrochloric acid. This gave promise that here, if anywhere, aluminium should certainly be found in flowering plants. He therefore examined a number of plants and flowers grown on the farm lands of the Imperial College of Agriculture at Komaba, near Tokyo. Each one was carefully picked over and all imperfect and soaked grains rejected.

Great care was taken to clean the grains and parts so as to remove all dirt. They were ground and incinerated, care being taken to avoid all possible contamination with clay or charcoal. The materials examined included soy bean (Soja hispida) (which the author calls "pea"), whole seed, cotyledons, and hull or skin, red bean (Adzuki) (Phaseotus radiatus), hill and paddy rice, wheat, barley, two sorts of millet, and buck-wheat. The analytical methods are described in detail. The table beyond reports the results of the analyses.

Of this work the author says: "It will be seen that I have found alumina in every case except that of the pea [soja], while in the hull or skin of the pea, one of the largest amounts of alumina occurs. The results here recorded may at least serve to indicate the propriety of reconsidering the accuracy of the

dictum that aluminium is not a constituent of flowering plants."

ASH	CONSTITUENTS	OF A	NUMBER	OF PLANTS
-----	--------------	------	--------	-----------

Kind of Plant.	Ash in	In Ash.			
Isling of Figure.	Air-dried Sample.	Alumina.	Phosph. Acid.	Silica, etc.	
	Per Cent.	Per Cent.	Per Cent.	Per Cent.	
"Pea" (whole) (Soja hispida)		0.053	33.48		
"Pea" (cotyledons) (Soja hispida)	4.22	.000		0.50	
"Pea" (hull or skin) (Soja hispida)		.268	5.66	3.60	
Red bean (Adzuki) (Phaseolus ra-					
diatus)	2.60	.096	32.89	0.25	
Rice (hill)		.161	51.33	9.36	
Rice (paddy)	0.56	.189	52.79	10.99	
Wheat	2.62	.106	65.55	1.31	
Barley	1.09	.140	33.19	1.19	
Millet (Awa) (Panicum italicum)	1.68	.272	40.43	8.91	
Millet (Hiyé) (P. cruscorvi)	0.94	.185	39.87	8.62	
Buckwheat	1.72	.113	1.94	0.81	

Young, W. C. (Analyst, 2 (1878), p. 13), in an article on the estimation of alum in bread, reported that from bread containing no added alum he obtained 0.07 grain Al. PO₄ per 1000 grains.

Young, W. C., also reports (Analyst, 12 (1887), p. 29), in an article on "Sour Bread and the Logwood Test," that he found aluminium phosphate equivalent to 7.8 grains of alum in 4 pounds of bread which he subsequently learned contained no added alum. A second sample of the same bread yielded aluminium phosphate equivalent to 8 grains of alum in 4 pounds of bread. In his investigation he found that the logwood test as applied by him indicated the presence of aluminium in the samples of flour or bread examined which were known to contain no added alum, provided the sample underwent acid fermentation or was acidulated with acetic acid. In a discussion of his paper, W. Blythe states that all bread contains more or less alumina.

Young, W.C. (Analyst, 12 (1887), p. 145), reports later a continuation of his investigation. He found that the aluminium normally present in the flours he examined was contained in the gluten of the wheat and not in the starch, and was present

in the form of aluminium phosphate. The following quotation is of interest in this connection:

"Another conclusion which may be fairly drawn from my results is that the alumina naturally present in flour is combined with the gluten. This being so, I thought it would be interesting to see if gluten would absorb alumina from a solution of alum. With this object I prepared some gluten, thoroughly dried it, and found the ash equal to 1.28 per cent. A portion of this gluten finely powdered was placed in a 2 per cent solution of alum and kept for some hours at a temperature of about 180° F.; it was then thrown on to a filter, thoroughly washed, and dried. The ash then amounted to 1.32 per cent. Taking into consideration the great difficulty of washing the gluten free from excess of alum, I do not think this slight increase of ash is due to absorption of alumina. Although the gluten does not appear to have the property of absorbing alumina from a solution of alum, vet its properties are affected in a marked degree by contact with such a solution, as it may then be kept moist in a warm place without giving any indication of decomposition, whilst ordinary gluten, as is well known, darkens in color, swells, and rapidly decomposes under similar circumstances."

Young, W. C. (Analyst, 13 (1888), p. 5), reports the determination of alumina in the best quality of Vienna flour, containing 0.7 per cent ash and about 8 per cent gluten. From 100 gms. of flour he obtained 0.0075 gm. of phosphate of alumina.

Of his experiment he says:

"The gluten was separated by washing in a muslin bag in the usual way, and when dried contained 1.26 per cent ash; 20 gms. of this dried gluten, finely powdered, was then treated with about 250 cc. of a mixture of equal volumes of acetic acid and water and heated in the water-bath for about twentyeight hours. By this time the mass had become quite liquid, the gluten having lost its firmness in the same way that gelatin does under similar circumstances. After standing a short time the liquid was poured off and the sediment further treated with weak acetic acid twice, and the three portions of liquid evaporated to dryness, the sediment being rejected. In this way I think that any extraneous earthy matter present in the gluten was separated, and, therefore, only the natural alumina retained.

"The dried residue was then burnt to a perfect ash, the ash dissolved in dilute hydrochloric acid and filtered, the insoluble matter being well washed and weighed. The insoluble matter thus obtained weighed only 0.009 gm., and of this .0075 was silica.

"The insoluble matter was then fused with about twice its bulk of mixed alkaline carbonates, dissolved in dilute hydrochloric acid, and filtered. This filtrate was added to the acid solution of the ash, evaporated to dryness, redissolved in a small quantity of dilute hydrochloric acid, and filtered. The filtrate was then boiled and cautiously added to 25 cc. of a saturated solution of pure caustic soda, also boiling, and the whole kept boiling for a few minutes. It was then filtered, and the precipitate washed, the filtrate made slightly acid with hydrochloric acid, about 5 cc. of a saturated solution of sodium phosphate added, and finally a slight excess of ammonia. After boiling for about ten minutes, the precipitate of phosphate of alumina was collected and weighed. . . .

"In this way I obtained .o185 gm. of phosphate of alumina from 20 gms. of gluten. Now, as the flour contained 8 per cent of gluten, and gave originally .o075 per cent of phosphate of alumina, 20 gms. of gluten would be equivalent to 250 of the flour, which would yield .o1875 of phosphate of alumina. So that practically I obtained the whole of the alumina of the flour in the gluten. As in the process of washing the starch from the gluten a large proportion of any foreign earthy matter that may have been present must have been separated, and any remaining eliminated by dissolving the gluten in acetic acid, there can be no doubt that the alumina obtained in this experiment was present as a natural constituent of the flour, and I think further that the interesting fact is established that the bulk of it is associated with the gluten."

Young, W. C. (Analyst, 15 (1890), p. 83), in a report of an investigation of the solubility of phosphate of alumina in acetic acid undertaken with special reference to the estimation of alumina in flour, bread, etc., makes the following statement: "I may mention that rye bread, of which I frequently have

samples for analysis from the east end of London, invariably gives a strong reaction with a logwood test, although I have never found more than .008 gm. phosphate of alumina in 100 gms."

Yvon, M. (Jour. Pharm. et Chim., 4th ser., 25 (1877), p. 588), reports the ash analyses of *Thapsia garganica* from Hegeria and *T. sylphium* * from Asia, the former with 7.52 per cent and the latter with 5.746 per cent in the fresh plant. Of this 0.3 and 0.433 per cent respectively was said to be Al₂O₃.

Zeyer, N. (Vierteljahressch. Prakt. Pharm., 10, p. 504; Jahresb. Chem., 1861, p. 769), found 4.05 per cent ash in the bark of Atherosperma moschatum dried at 100°. Of this 0.191 per cent was Al₂O₃.

^{*} There is an evident error in the names of the plants in the original tables as published. The names as quoted above have been corrected.

ALUMINIUM IN ANIMAL PRODUCTS.

Dieterich, E. (Vierteljahressch. Prakt. Pharm., 16, p. 56; Chem. Centbl., 1867, p. 287; Jahresb. Chem., 1867, p. 822), reports 3.2 per cent ash in cochineal (silver-gray Honduras variety); of this 1.390 was Al₂O₃.

Harrison and Kinnicutt, according to Rotch (Pediatrics, Philadelphia, 1896, p. 175), found 0.40 per cent iron oxid and alumina in the ash of woman's milk, the total ash ranging from 0.1 to 0.2 per cent. From the data at hand, the proportion of the ash constituents was recalculated, the value for iron oxid and alumina being given as 0.37 per cent. Silica was also found, and it is stated that this and alumina have not been noted in any previous analysis.

Data are also quoted by Rotch (loc. cit., p. 218) regarding the ash of cow's milk which it is said were reported by König, Forster, and others. The value given for total ash is 0.70 per cent, of which 0.44 per cent was iron oxid and alumina.

Heyl (Ann. Chem. Pharm., 62, p. 87, Repert. Pharm., 2d ser., 47, p. 231; Pharm. Centbl., 1847, p. 591: Jahresb. Chem., 1847–48, p. 938) notes 29.18 per cent aluminium silicate in a sample of officinal burned sponge (Schwammkohle).

Liebermann, C. (Ber. Deut. Chem. Gesellsch., 1885, p. 1969; Jahresb. Chem., 1885, p. 1847), estimates that cochineal yields 9 to 10 per cent pure coloring matter. In good cochineal carmine, he found 7 per cent ash. This contained a trace of Al₂O₃. Carmine is regarded as an aluminium-calcium-protein compound of the coloring matter present.

Mayrhöfer, J. (Chem. Centbl., 1891, I, p. 175; Jahresb. Chem., 1891, p. 2820), notes a trace of alumina in the ash of that portion of an abnormal leather which was soluble in cold water.

McMurtrie, W. (American Chemist, 4, p. 339; Watts' Dictionary, 1879, vol. 8, pt. 1, p. 765), finds in the excrements of the common bat 19.88 per cent of alumina.

Müntz, A. (Ann. Chim., 4th ser., 20, p. 309; Compt. Rend. Acad. Sci., Paris, 69, p. 1309; Bul. Soc. Chim., 2d ser., p. 379; Dingler's Polytech. Jour., 195, p. 466; Jahresb. Chem., 1869, p. 1149), in a study of the chemical changes in tanning, reports 0.6693 and 0.4670 per cent ash respectively in fresh, rather thin hide from a young cow, and very thick ox hide. The ash contained respectively 0.0930 and 0.0704 per cent Al₂O₃.Fe₂O₃.

Schütze, R. (Chem. Centbl., 1889, II, p. 588; Jahresb. Chem., 1889, p. 2156), found in the ash of the mantle of *Phallusia mammillaris* (after extraction with ether) 9.52 per cent Al₂O₃. The amount of ash is not stated.

Staffel (Arch. Pharm., 2d ser., 64 (1850), p. 148; Pharm. Centbl., 1851, p. 162; Jahresb. Chem., 1850, p. 573) found in veal (breast) free from fat and skin 77.64 per cent water, and in the dry material 3.1 per cent ash. This contained according to his analysis a trace of alumina. He also found a trace in the ash of beef. The fresh material contained 72.63 per cent water; the dry 2.2 per cent ash.

Theile, R. (Jenaische. Ztschr. Med., 3, p. 147; Ztschr. Chem., 1868, p. 125; Chem. Centbl., 1867, p. 296, 305; Vierteljahressch. Prakt. Pharm., 17, p. 197; Bul. Soc. Chim., 2d ser., 10, p. 153; Jahresb. Chem., 1867, p. 772), reports 2.3 per cent ash in egg albumen, 15.04 per cent of this being Al₂O₃. PO₅.

Thezard (Compt. Rend. Acad. Sci., Paris, 120 (1895), p. 1126; Jour. Chem. Soc., London, 68 (1895), II, p. 456) analyzed a very white, porous, and very brittle tibia from an adult human mummy from an Egyptian tomb of unknown age. In addition to other constituents, he reports 0.534 per cent Al₂O₃.

Walter, G. (Ztschr. Physiol. Chem., 13, p. 464; Jahresb. Chem., 1889, p. 2156), reports 28.165 per cent ash in the shell-like, outer covering of *Protopterus annectens* (air-dry). This contained 5.491 per cent Al₂O₃.

ALUMINIUM IN NATURAL WATERS.

Abbéne (Jour. Pharm., 3d ser., 12, p. 412; Pharm. Centbl., 1848, p. 47; Jahresb. Chem., 1847-48, p. 1009) notes traces of alumina in the water of the Prè Saint Didier upper and lower mineral springs at Courmayeur.

Adams, F. D. (Geol. and Nat. Hist. Survey, Canada, 1885. Chemical Contributions, p. 15, M), reports 0.0005 parts alumina per 1000 in water from a spring at Halowell Grant, near Antigonish, Nova Scotia. This water is used for medicinal purposes.

Adams, F. D. (Geol. and Nat. Hist. Canada, 1888-9. Chemical Contributions, p. 17, R.), in an analysis of water from a boring in the west half of Lot 26, in the fourth range of Otanabee, Peterborough County, Ontario, reports 0.0008 part alumina per 1000.

Agrestini, A. (Gaz. Chem. Ital.; Jour. Chem. Soc. London, 64 (1893), II, p. 175), found in the hepatic mineral spring water of the Valle del Gallo, near Urbino, 0.0005 AlPO₄ per 1000 gms.

Aillaud (Compt. Rend. Acad. Sci., Paris, 95, p. 104; Jahresb. Chem., 1882, p. 1627) reports 0.007 gm. Al₂S₃O₁₂ per 1000 cc. in high water from the Rio Grande which was used for the water supply of Panama.

Aillaud (Compt. Rend. Acad. Sci. Paris, 95, p. 104; Jahresb. Chem., 1882, p. 1637) reports 0.020 gm. Al₂O₃ per 1000 cc. in well-water from Emperador, Panama, and 0.011 gm. Al₂O₃ in water from a second well in the same locality.

Allemann, H. (Wiener Akad. Ber., 56 (2. Abt.), p. 47; Wiener Akad. Anz., 1867, p. 135; Jour. Prakt. Chem., 101, p. 317; Chem. Centbl., 1867, p. 423; Inst. 1867, p. 341; Jahresb. Chem. 1867, p. 1039), in an analysis of the Sauerbrunn at Ebriach in Karnthen, reports 0.034 part Al₂O₃ and 0.015 parts aluminium phosphate per 10,000.

Andouard (Jour. Chim. Med., 3d ser., 5, p. 466; Jahresb. Chem., 1849, p. 617) found 0.003 gm. aluminium phosphate and 0.006 gm. alumina per 1000 cc. in mineral water from Villecelle, near Lamalou (Hérault).

Andouard, A. (Jour. Pharm., 4th ser., 9, p. 336; Jahresb. Chem., 1869, p. 1291), reports two analyses of Beaupréau spring-water from Dep. Maine-et-Loire. It contained 0.022 and 0.010 gm. Al₂O₃ per 10,000 gms. The second analysis was made with a sample taken after heavy rain.

Ashley, J. M., E. T. Bennett, and T. J. Herapath are quoted (Jahresber. Chem., 1849, p. 620) as reporting traces of aluminium in Thames water near London Bridge, Thames water at Greenwich, and mineral water from Kingswood, near Bristol, respectively.

Avequin (Jour. Pharm., 3d ser., 32, p. 288; Jahresb. Chem., 1857, p. 729) notes 1.753 grains alumina per gallon (=58,372 grains) in Mississippi River water taken at Carrollton, a few miles above New Orleans, Louisiana.

Bailey, E. H. S., in an extended treatise entitled "Special Report on Mineral Waters" (University Geol. Survey of Kansas, vol. 7), gives analytical data regarding the composition of a large number of waters in Kansas. The following table summarizes the data concerning alumina in these waters.

ALUMINA IN KANSAS MINERAL WATERS.

	Analyst.	Page.	Alumina per Liter.
The Chlorid Croup			
The Chlorid Group.	B W C B "		Grams.
Arkansas City Well	E. H. S. Bailey	131	0.0266
Eureka Mineral Well	E. H. S. Bailey	137	.0017
Geuda Spring, No. 4	Bailey and Franklin	147	.0158
Bromo-magnesium Well	E. H. S. Bailey E. H. S. Bailey	150	trace
	D. II. S. Daney	157	.0178
The Sulphate Group.			
Burr Oak	G. H. Failyer and	1.60	
	C. M. Breese	169	.1175
Capioma Mineral Well	E. H. S. Bailey	170	.0103
Carbondale Spring	E. H. S. Bailey	172	.0096
Centralia Gypsum Well	E. B. Knerr	173	.0070
Conway	Failyer and Breese	173	.0104
Chlor-Sulphate Group.			
Merrill Spring	A. Merrill	196	0024
Great Bend Mineral Well	E. H. S. Bailey	197	trace
Great Spirit Spring, No. 2	Bailey and Rice	205	.0166
Lincoln Springs, No. 1	Bailey and Franklin	207	.0005
Lincoln Springs, No. 2	Bailey and Franklin	208	.0047
Lincoln Springs, No. 3	Bailey and Franklin	209	.0034
Carbonate Group.			
Dixon Spring	E. B. Knerr	200	
Kickapoo Springs	E. H. S. Bailey	219	trace
	D. II. O. Daney	237	1100.
Sulfid Group.			
Cherokee City Well	Bailey and Hull	260	.0037
Columbus Well	Failyer and Willard	262	trace
Fort Scott Sulfo-magnesian Well .	E. H. S. Bailey	266	.0144
Moss Springs Well	G. H. Failyer	268	.0241
wakeneld Sulfur Well	Bailey and Porter	271	.0530
Chalybeate (Iron) Group.			
Atchison Electric Light Well	E. B. Knerr	280	.0567
McDuff's Spring	E. B. Knerr	281	.0120
Special Group.			
	E-11 1 W. 1	The same of	
Lithium Spring, Omio	ranger and Willard	307	.1940
	J. H. Banks	308	trace
Soft-water Group.			
Atchison Parker's Spring	E. B. Knerr	311	.0032
			1
New York and the Control of the Cont	THE RESERVE OF THE PARTY OF THE		

Bailey quotes analyses of a number of waters for purposes of comparison, but does not cite the original place of publication. The following table shows data regarding the occurrence of alumina in such as have not been noted from other sources.

ALUMINA IN A NUMBER OF WATERS.

	Analyst.	Page.	Alu- minium Chlorid per Gallon.	Alu- minium Phos- phate per Gallon.	Alumina per Gallon.	Alu- minium Sul- phate per Gallon.
Oranien Quelle, Kreuznach, Rhen- ish Prussia Dead Sea	Liebig	160	Grains.	.760	Grains.	Grains.
Carabana, Spain Friedrichshall, Saxe-Meiningen, Germany	A. Proust Bauer	192			.029	
Apollinaris, Germany. Sulphur Spring, Aix- les-Bains, France.	Bischoff Bonjeau	246				3.200
Excelsior Sp'gs, Mo. { Sparta Artesian Well,	Woodward and Rob'tson	} 301				3
Wisconsin	J. M. Hirsch	301		.048		

Bailey, E. H. S., and Mary A. Rice (Trans. Kansas Acad. Sci., 14 (1896), p. 40; Jour. Chem. Soc. London, 72 (1897), II, p. 109) report 1.66 parts Al₂O₃ per 100,000 in water from a mineral spring in Mitchell County, Kansas, near Cawker City.

Baker, W. H. (Dingler's Polytech. Jour., 218, p. 267; Jour. Chem. Soc. London, 1876, I, p. 890), notes 55.128 grains Al₂(SO₄)₃ per gallon in the water of a *swallet* in the Empire mine of the Luzerne Company.

Ballo, M. (Ber. Deut. Chem. Gesellsch., 1878, p. 1900; Jahresb. Chem., 1878, p. 1306), notes 0.0017 Al₂O₃ (with a trace of P₂O₅) per 1000 in a thermal spring at the foot of the Blocksberg.

Ballo, M. (Ber. Deut. Chem. Gesellsch., 1878, p. 1902; Jahresb. Chem., 1878, p. 1305), in two samples of Ofener Bitter Wasser reports 0.229 and 0.089 parts Al₂O₃ (with P₂O₅) per 10,000, respectively.

Ballo, M. (Russ. Ztschr. Pharm., 22, p. 68; Jahresb. Chem., 1883, p. 1945), notes 0.0229 part Al₂O₃ per 1000 parts in the bitter water of the Victoria well at Ofen, Hungary.

Ballo, M. (Ber. Deut. Chem. Gesellsch., 1884, p. 673; Jahresb. Chem., 1884, p. 2035), reports 0.00952 gm. Al₂O₃ per 1000 cc. in Borhegyer Sauerwasser from Bibarczfalva, Austria-Hungary.

Barth, L. v., and R. Wegscheider (Wiener. Klin. Wochensch., 1891, No. 8; Jahresb. Chem., 1891, p. 2620) report 0.3575 gm.

Al₂O₃ per 10,000 gms. in mineral spring water from Mitterbad, Ultenthale, Tirol.

Barzilowski, J. (Ber. Deut. Chem. Gesellsch., 1884, p. 183; Jahresb. Chem., 1884, p. 2037), notes 0.0055 part Al₂O₃ per 1000 parts in Tokiew mineral water; and 0.103 part Al₂O₃ per 1000 parts in Alexandro-Jermolowski mineral water, both from the Caucasus.

Bauck, G. (Analyse der Salzsoolen von Colberg, Inaug. Diss. Göttingen, 1860; Jahresb. Chem., 1860, p. 830), reports alumina (parts per 10,000) in Colberg, Germany, saline mineral water as follows: Salinensoole, 0.011; Zillenbergsoole, 0.008, and Marktsoole, 0.006.

Bechamp, A. (Compt. Rend. Acad. Sci. Paris, 62, p. 1088; Jour. Pharm., 4th ser., 3, p. 448; Jour. Prakt. Chem., 98, p. 189; Chem. Centbl., 1866, p. 864; Jahresb. Chem., 1866, p. 998), found 0.0052 gm. alumina per 1000 cc. in Thérèse sulphur spring at Fumades (Arrondissement d'Alais), and (Compt. Rend. Acad. Sci. Paris, 62, p. 1034; Bul. Soc. Chim., 2d ser., 6, p. 9; Inst. 1866, p. 173; Jour. Pharm., 4th ser., 3, p. 444; Jour. Prakt. Chem., 98, p. 190; Chem. Centbl., 1866, p. 864; and Compt. Rend. Acad. Sci. Paris, 63, p. 559; Jahresb. Chem., 1866, p. 998) also reports 0.00106, 0.0008, and 0.0011 gm. alumina per liter respectively in Dulimbert, Bouillants, and Granier mineral springs at Vergèze, Dép. du Gard.

Becker, G. F. (U. S. Geol. Survey Mon. 13, p. 265), quotes an analysis of water from Borax Lake, Cal., by Melville (Geol. Survey Cal., 1, p. 98) which showed 0.0029 gm. Al₂O₃ per liter. Becker (pp. 347 and 349) also gives an analysis of the water of Steamboat Springs, Nevada. The Al₂O₃ present was 0.0025 gm. per 10 liters.

Bell (Chem. News, 21, p. 83, from Amer. Jour. Pharm.; Jahresb. Chem. Naumann, 1870, p. 1392) notes traces of Al₂O₃ in a mineral spring in New York in the Adirondack region.

Bender, R. (Arch. Pharm., 3d ser., 11 (1877), p. 50; Jour. Chem. Soc. London, 1878, II, p. 18), notes that Fresenius found 0.00013 parts AlPO₄ per 1000 in Heilbrunnen mineral spring water, a spring situated near Lake Taach, and in the water of the Stahlbrunnen at Wassenach, 0.00045 AlPO₄.

Bender, R. (Arch. Pharm., 2d ser., 185 (1868), p. 5; Jahresb. Chem., 1868, p. 1035), reports 0.005, 0.182, and 0.045 Al₂O₃ per 10,000 parts respectively in the Victoria, Angustus, and Cold Spring III. These waters are all from the thermal springs of Neuenahr.

He also reports per 10,000 parts 0.176 Al₂O₃ in the old or small Sprudel spring No. I and 0.19 Al₂O₃+Fe₂O₃ in the new or large Sprudel No. II. The Marien Sprudel contained 0.06 Al₂O₃+Fe₂O₃.

Bensemann, R. (Chem. Centbl. Ausz., 1882, p. 186; Jahresb. Chem., 1882, p. 1629), reports 0.0026 Al₂O₃ per 100 in a Soolquelle at Kammin, Germany.

Berlin, N. J. (Oefversigt. K. Vetenskops. Akad. Förhl., 1863, p. 221; Jahresb. Chem., 1865, p. 939), notes 0.00794 part alumina per 10,000 in Torpasaltkälle at Lilla Edet in Sweden.

Bertoni, G. (Gaz. Chem. Ital., 14 (1884), p. 232; Jahresb. Chem., 1884, p. 2035), in an analysis of the Acquarossa Spring at Biasca, Italy, notes 0.0485 gm. Al₂O₃ per 1000 cc.

Billot, E. (Jour. Chim. Med., 3d ser., p. 569; Jahresb. Chem., 1853, p. 716), reports alumina (gm. per 1000 cc.) in water from Besançon, France, as follows: Doubs, 0.002; spring-water from the Grand Rue, 0.0094.

Bizio (Wiener. Acad. Ber., 41, p. 335, Rep. Chim. Pure. 2, p. 390; Jahresb. Chem., 1860, p. 837) reports (parts per 10,000) 0.0371 Al₂O₃ in the St. Gothard Spring water from Ceneda, Italy.

Bizio, G. (Atti. Imp. Reg. Inst. Venato Sci. Let. ed Arti., 3d ser., 11; Jahresb. Chem., 1865, p. 940), notes 7.7507 alumina per 10,000 in mineral water from Civillina.

Bizio, G. (Gaz. Chim. Ital., 10 (1880), p. 43; Jahresb. Chem., 1880, p. 1529), reports new analyses of the springs of Civillina, Italy. The old basin contained 1.28414 gms. aluminium sulphate; the new basin, 1.50764 gms. aluminium sulphate per 1000 gms.

Blondeau (Compt. Rend. Acad. Sci. Paris, 30, p. 481: Pharm. Centbl., 1850, p. 910; Jahrb. Chem., 1850, p. 621) notes the occurrence of aluminium salts as among the ordinary constituents of water of brooks, rivers, and springs. Water

containing 0.4 to 0.5 gm. total mineral matter per liter is regarded as satisfactory for household purposes.

Blondeau (Compt. Rend. Acad. Sci. Paris, 35, p. 147; Inst. 1852, p. 238; Jour. Prakt. Chem., 57, p. 244; Jahresb. Chem., 1852, p. 756) reports 0.0016 gm. alumina (per 1000 gms.?) in water from Salles-la-Source, Aveyron.

Bobierre and Moride (Compt. Rend. Acad. Sci. Paris, 32, p. 376; Inst. 1851, p. 90; Jour. Pharm., 3d ser., 20, p. 244; Jahresb. Chem., 1851, p. 666) note 0.401 gm. total solids per 1000 cc. in mineral water from Kirouars, France; this contained a trace of alumina. The same authors (Compt. Rend. Acad. Sci. Paris, 33, p. 322; Inst. 1851, p. 306; Pharm. Centbl., 1851, p. 748; Jahresb. Chem., 1851, p. 666) noted 0.350 gm. total solids per 1000 cc. in mineral water from La Bernerie, France. This contained 0.6 per cent alumina.

Bolley, P., and F. Schweizer (Ann. Chem. Pharm., 106, p. 237; Chem. Centbl., 1858, p. 600; Jahresb. Chem., 1858, p. 800) found 0.0103 (gm.?) alumina per 1000 cc. in the sulphur water from Schinznach, Switzerland.

Bolley, P., and Brigel (Schweiz. Polytech. Ztschr., 1865, p. 47; Ztschr. Chem., 1865, p. 639; Jahresb. Chem. 1865, p. 936) report 0.0130 alumina per 1000 parts in the sulphur spring A at Lostorf in the Jura.

Bosshard, E. (Chem. Centbl., 1892, II, p. 1039; Jahresb. Chem., 1892, p. 2689), reports 0.00057 gm. Al₂O₃ per 10,000 gms. in the Old Spring water at St. Moritz in the Engadine and 0.00040 gm. Al₂O₃ per 10,000 gms. in Paracelsus spring water from the same locality.

Boussingault (Compt. Rend. Acad. Sci. Paris, 78, pp. 453, 526, 593, Ann. Chim. Phys., 5th ser., 2, p. 76; Arch. Pharm., 3d ser., 5, p. 354; Jahresb. Chem., 1874, p. 1337) reports 0.4028 gm. Al₂O₃ per liter in the water of the Rio Vinagre, near the falls of San Antonio, Colombia, a volcanic region, and 0.500 gm. per liter in a hot spring at the foot of the Ruiz volcano.

Boutel, G. F., examined the mineral waters of Saint Nectaire (Ann. Chim. Phys., 6th ser., 7, p. 536; Jour. Chem. Soc., London, 50 (1886), II, p. 858) and reported per liter 0.0330 gm. Al₂O₃ in Source Rouge and 0.0239 gm. in Source du Mont Cornador.

Braconnot (Jour. Chim. Med., 3d ser., 7, p. 737; Jahresb.

Chem., 1851, p. 665) notes 0.0080 gm. silica and alumina per 1000 cc. in the Luxeuil Spring.

Brignone, G. (Gaz. Chem. Ital., 14 (1884), p. 42; Jahresb. Chem., 1884, p. 2036), found in water from the island of Pantelleria (Italy), 0.00245 gm. Al₂O₃. P₂O₅ per 1000 cc.

Buchner, A. (Jour. Prak. Chem., 102, p. 209; Neue Repert. Pharm., 16, p. 481; Neue Jahrb. Pharm., 28, p. 292; Jahrb. Chem. Will., 1867, p. 1036), found 0.00104 gm. alumina per liter in potable water from the spring at Neumarkt, Oberpfalz, Bavaria.

Buchner, L. A. (Neue Rep. Pharm., 17, p. 357; Jour. Chem., 104, p. 360; Jahresb. Chem. Will., 1868, p. 1037), reports traces of alumina in the sulphur spring at Oberdorf in Allgäu.

Buchner, M. (Wiener. Akad. Ber., 2. Abt., 71, p. 309; Jahresb. Chem., 1875, p. 1296), notes 0.0274 parts aluminium phosphate per 10,000 in Moriz spring-water from Sauerbrunn, near Rohitsch, Austria.

Buchner, M. (Chem. Centbl., 1876, p. 789; Jour. Chem. Soc. London (1877), II, p. 176), in the water of the Tempelbrunnen at Rohitsch, notes 0.0095 parts (= AlPO₄) per 100,000.

Buchner, M. (Chem. Centbl., 1881, p. 567; Jahresb. Chem., 1882, p. 1631), found 0.0249 parts Al₂O₃.P₂O₅ per 10,000 in the Lindenbrunnen at Zlatten, Austria-Hungary.

Bull (Silliman's Amer. Jour., 2d ser., 4, p. 385; Pharm. Centbl., 1848, p. 319; Jahresb. Chem., 1847-48, p. 999) reports 0.14 gm. alumina per 10,000 gm. in spring-water from Hartford, Connecticut.

Bunsen (Neue Jahresb. Pharm., 2, pp. 190, 194; Pharm. Centbl., 1854, p. 931; Jahresb. Chem., 1854, p. 758) examined a number of mineral waters, reporting aluminium phosphate (parts per 1000) as follows: Sophienquelle, 0.0033; Petersquelle, 0.0071, and Salzquelle, 0.0035, all at Petersthal.

Bunsen, according to Riegel (Neue Jahresb. Pharm., 9, p. 301; Jahresb. Chem., 1858, p. 794), found 0.007 part alumina per 10,000 parts in water from the Elizabethenquelle at Rothenfels, Baden.

Bunsen (Beiträge zur Statistik der inneren Verwaltung des Grossherzthums Baden, 11, Nos. 30, 43, and 56; Jahresb.

Chem., 1861, p. 1090) in Baden mineral springs notes alumina as follows (parts per 10,000): Ursprung, 0.011; Judenquelle, 0.011; Brühquelle, 0.009; Murquelle, trace, and Rothenfelser Mineralquelle, 0.007.

Bunsen and Kirchoff (Poggendorf's Ann., 113, p. 358; Phil. Mag., 4th ser., 22, p. 344; Jahresb. Chem., 1861, p. 1092) note 0.001 alumina (parts per 10,000) in the Umgemach thermal spring (Baden-Baden) and note 0.00020 in saline water from Dürkheim.

Burrell, B. A. (Jour. Chem. Soc. London, 69 (1896), p. 536), in an analysis of the water of the dropping well at Knaresborough in Yorkshire, notes a trace of alumina.

Carius, L. (Ann. Chem. Pharm., 137, p. 106; Jahresb. Chem. Will., 1866, p. 987), notes in the Ottilienquelle, Badequelle, and Marienquelle from the Inselbad near Paderborn, 0.009, 0.009, and 0.004 grams alumina per 10,000 respectively.

Carnelley, T. (Chem. News, 31, p. 27; Jahresb. Chem., 1875, p. 1298), reports in a spring (iron) near Trefriw, Wales, 233.3 parts aluminium per million. An earlier analysis by Hassal (Chem. News, 31, p. 27; Jahresb. Chem., 1875, p. 1298) gives 112.4. These values correspond to 1358.9 aluminium sulphate and 3.2 aluminium phosphate, and 707.7 aluminium sulphate, and 0.0 aluminium phosphate per million respectively.

Carnot, A. (Ann. Mines, 9th ser., 6 (1894), p. 355), reports a series of analyses of French mineral waters made at the École du Mines, containing data for the period from January, 1885, to June, 1894. The following contained alumina (grams per liter): "Sources de Fraysse, No. 1," Commune de Cransac, Département l'Aveyron, 0.0016; "Sources de Fraysse, No. 2," 0.0840, which were calculated to be equal to 0.0087 and 0.2790 gm. Al₂(SO₄)₃ respectively. The "Sources Roques" and "Rouquette," Cransac, Département l'Aveyron, contained respectively 0.0093 and 0.0071 Al₂O₃ equal to 0.0300 and 0.0230 aluminium sulphate. The "Sources No. 2," Cransac, contained 0.1340 Al₂O₃=0.4462 aluminium sulphate. Water from the Briscous salt-wells, Commune Briscous, Département Basses-Pyrénées, 0.0050 Al₂O₃ (with traces of Fe₂O₃). (See also Compt. Rend. Acad. Sci. Paris, 111, p. 192.)

Carnot, A. (Ann. Mines, 9th ser., 16 (1899), p. 33), reports a later series of analyses of mineral waters made at the laboratory of the École Nationale superieure des Mines, July, 1894–June, 1899. In the following, alumina was reported (grams per liter): "Source Saint-Louis," Département de la Drome, 0.1650 (=0.5505 aluminium sulphate); spring, Canton et Arrond. de Bagnères-de-Bigorre, Département Hautes-Pyrénées, 0.00060.

Casali, A. (Staz. Sper. Agr. Ital., 19, p. 509; Jahresb. Chem., 1890, p. 2658), notes traces of aluminium in potable water from Bologna, Riggio-Emilia, Ferrara, and Ancona.

Casselmann, W. (Jahrb. Ver. Naturk. Herzogthum Nassau, No. 15, p. 139; Jour. Prak. Chem., 83, p. 385; Chem. Centbl., 1861, p. 874; Jahresb. Chem., 1861, p. 1094), reports alumina (per cent by weight) as follows in Nassau mineral water: Soolsprudel, 0.00062; Soolsprudel (sample taken in 1859), 0.00012; No. IV (1857), 0.00054; No. VII (1858), 0.00005; No. I (1859), 0.00016; No. X (1859), 0.000023; No. III (1859), 0.00016; and Neuenhainerquelle (1860), trace.

Cassels, J. L. (Amer. Chem., 4 (1873), p. 169; Jahresb. Chem., Naumann, 1873, p. 1246), reports 0.333 gm. Al₂O₃ per 10,000 cc. in spring-water from Castalia, Ohio.

Chandler, C. F., F. A. Cairns, and S. P. Sharpless (Ann. Chem., 2d ser., 3, pp. 93, 164, and 202; Jahresb. Chem., Naumann, 1872, p. 1189) found in the Saratoga waters "Glacier Spouting," "Empire," and "Triton," 0.08 gm., 0.07 gm., and a trace of alumina per 10 liters.

Chandler, C. F., and F. A. Cairns (Amer. Chem., 4 (1873), p. 86; Jahresb. Chem., Naumann, 1873, p. 1246) note 0.055 gm. Al₂O₃ per 10 liters in Saratoga union spring-water (0.324 grains per gallon).

Chandler, C. F., and F. A. Cairns (Amer. Chem., 6, p. 241; Jahresb. Chem., 1876, p. 1309) report 0.1002 gm. Al₂O₃ per 10,000 cc. in St. Leon water from Canada, and (Amer. Chem., 6, p. 370; Jahresb. Chem., 1876, p. 1309) 0.0220 gm. Al₂O₃ per 10,000 cc. in artesian water from Sheboygan, Wisconsin.

Chatard, T. M. (U. S. Geol. Survey, Bul. 9, p. 23), found in water from Humboldt River, Nevada, 0.3615 gm. total solids per liter, of which 0.0013 gm. or 0.37 per cent was Al₂O₃.

Chatard, T. M. (U. S. Geol. Survey, Bul. 9, p. 24), in water

from Hot Springs, Hot Springs Station, Nevada, found 2.4924 gms. total solids per liter, 0.0010 gm. or 0.04 per cent being aluminium.

Chatard, T. M. (U. S. Geological Survey, Bul. 9, p. 27), found in warm spring-water, Mono Basin, California, 2.0850 gms. total solids, containing 0.0018 gm. or 0.09 per cent Al₂O₃.

Chatard, T. M. (U. S. Geological Survey, Bul. 9, p. 29), found in water from City Creek, Utah (collected above reservoir which supplied Salt Lake City), 0.2400 gm. total solids per liter, 0.0010 gm. or 0.041 per cent being Al₂O₃.

Chatard, T. M. (U. S. Geol. Survey, Bul. 55, p. 93), found in water from Owens Lake, California, 0.023 gm. Al₂O₃ per kilo.

Chatard, T. M. (U. S. Geological Survey, Bul. 64, pp. 59, 60), notes 0.0005 and 0.0012 gm. Al₂O₃ per liter respectively in artesian wells A and B, St. Augustine, Florida; also 0.0077 gm. Al₂O₃ per liter in water from a well four miles northwest of Clinton, Miss.

Chevallier, A. (Jour. Chem. Med., 3d ser., 7, p. 193; Jahresb. Chem., 1851, p. 666), found 0.125 gm. alumina and silica per liter in mineral water from Wattweiler.

Clark (Chem. Soc. Quar. Jour., 1, p. 155; Pharm. Centbl., 1848, p. 650; Jahresb. Chem., 1847-48, p. 998) notes a trace of alumina in the Thames water at Twickenham.

Clark, F. W. (U. S. Geol. Survey, Bul. 55, p. 92; Jahresb. Chem., 1891, p. 2622), reports 0.0020 gm. Al₂O₃ per 1000 cc. in Potash Sulphur Spring water from Garland County, eight miles southeast from Hot Springs.

Cloëz, C. (Compt. Rend. Acad. Sci. Paris, 98, p. 1282; Jahresb. Chem., 1884, p. 2036), found 0.0710 gm. Al₂O₃,SiO₂ per 1000 c.c. in a mineral spring near Brucourt, in the neighborhood of Dives, France.

Crawfurd (Vierteljahressch. Prakt. Pharm., 6, p. 161; Chem. Centbl., 1875, p. 267; Jahresb. Chem., 1857, p. 722) notes 0.023421 Al₂O₃. SiO₃ per 1000 parts in mineral water from Kellberg near Passau, Germany.

Crook, J. K. (Mineral Waters of the United States and their Therapeutic Uses, Philadelphia, 1899), made an extensive compilation of analyses of mineral waters of the United States which was based, the author says, on treatises on mineral waters, issued by the various States, National Geological reports, railroad guides, etc., as well as information received in reply to inquiries addressed to all spring resorts and commercial springs of the United States which could be reached. Personal visits were also paid to many of the more important localities. In quoting analytical data he has expressed the results in grains per United States gallon. The whole subject is discussed with special reference to the use of mineral waters in the treatment of disease. It has been the author's purpose to include all the springs of the United States whose waters are actually used at the present time. The following statements are made regarding aluminium in discussing the value of the different constituents of mineral waters:

"This substance is found in springs in the form of the oxide, or alumina, and of the sulphate. It occurs in variable quantities, ranging from a mere trace to sixty or eighty grains per gallon, as seen in some of the Virginia alum springs. The sulphate is almost always present in the sulphureted chalybeate waters, and, as stated above, in the acid springs. internal use of alum waters is generally governed by these associated ingredients. Some of the alum springs have acquired a considerable reputation in scrofulous diseases and in chronic diarrhœa and dysentery. The iron-alum waters are beneficial in passive hemorrhages and in exhausting night-sweats. Locally their astringent action calls them into service in much the same class of cases as are benefited by the acid waters. have produced valuable results in conjunctivitis, stomatitis, chronic vaginitis, and other relaxed or inflammatory states of those portions of the mucous surfaces accessible to local treatment. They have also been found to act as a useful auxiliary in the treatment of ulcerated surfaces, abrasions, etc. In large quantities the alum waters have a laxative influence, but they are seldom used in virtue of this action."

The following table summarizes the data quoted from Dr. Crook's report:

ALUMINIUM IN SPRING-WATERS OF THE UNITED STATES.

Spring.	Analyst.	Page.	Aluminium Sulphate per U. S. Gallon.	Alumina per U. S. Gallon.
Alabama.			Grains.	Grains.
	F. A. Smith,			
Matchless Mineral Wells	E. A. Smith, J. B. Little	91		3.65
Arkansas.				
Arkansas Lithia Springs	Muehler	98		0.92
California.				
Agua de Vida Lower Spring.	Anderson	110		.37
Agua de Vida Upper Spring.	Anderson	110		.40
Alum Rock Alkaline Saline	A 1	2007		
Spring	Anderson Hatch	112		6.45
Alum Rock Chalybeate Sp'g Anderson Iron Spring	G. E. Colby	112		.15
Anderson Sour Spring	G. E. Colby	115	7.11	-93
Blodgett's Springs	Anderson	120		4.13
Calistoga Springs	Anderson	126		.47
Calistoga Swimming Pool	Anderson	126		.27
El Paso de Robles, Main			The state of	
Sulphur Spring	Anderson	130		.25
El Paso de Robles, Soda		100		
Spring.	Anderson W. D. Johnson	131		.85
Eureka Springs Felt's Mineral Springs	W. D. Johnson	131		1.30
California Geysers, Lemon-		132		a
ade Spring	T. Price	138	32.02	
California Geysers, Witches'		-30	3	
Caldron	T. Price	138	2.04	.27
California Geysers, Acid Sp'g	T. Price	138	20.62	
CaliforniaGeysers, Alum Sp'g	T. Price	138	63.82	
California Geysers, Iron Gey-	m n .			
ser Creek	T. Price	138		.89
California Geysers, Spring on hill	T. Price	0		
California Geysers, Iron Sp'g	T. Price	138	.20	
California Geysers, Indian	1.11100	130		.17
Spring	T. Price	138		.18
California Geysers, Mud In-		-		
dian Spring	T. Price	138	22.78	
California Geysers, Spring				
little above Indian Spring	T. Price	138	118.78	
California Geysers, hot sul-				
phur-water, above bath-	T. Price		0.00	
house	1. Fince	138	2.39	
Teakettle	T. Price	138	31.16	
Glen Alpine Springs	Anderson	142		1.43
Gordon Springs	Anderson	142		3.55
Harbin Hot Sulphur Spring.		144		1.60
Harbin Chalybeate Spring		144		.73
		the same		a de la constitución de la const

a. Amount not stated. Qualitative analysis.

Spring.	Analyst.	Page.	Aluminium Sulphate per U.S. Gallon.	Alumina per U. S. Gallon.
			Grains.	Grains.
Highland Seltzer	W. Anderson	146		
Highland Dutch or Ems	W. Anderson	The second second		1.75
Hot Porete	The second secon	146		.12
Hot Borate		147		2.04
Howard Springs, Excessor,	W (D W 11	- 0	1	
No. 1	W. T. Wenzell	148		0.03
Howard Springs, The Twins,	m			
No. 2	W. T. Wenzell	148		0.15
Howard Springs, Eureka,				
No. 3	W. T. Wenzell	148		0.10
Howard Springs, Neptune,		69.5		
No. 4	W. T. Wenzell	148		0.19
Howard Springs, Soda, No. 5	W. T. Wenzell	148		0.13
	San Francisco Re-)		
Lane Mineral Springs	fining and Analyti-	- 150		2.01
	cal Association	1		
Litton Seltzer Springs	W. Anderson	151		6.8r
Mark West Springs	TT. IIIIdeibon	152		a
Mono Lake	W. Anderson	153		26.63
Pagoda Spring (Napa Soda).	W. Anderson	The state of the s	CO. TO SOME STREET, ST	
Newsom's Arroyo Grande	W. Anderson	155		.57
	W. Anderson	6		
Springs	W. Anderson	156		.33
Pacific Congress Springs	W. Anderson	158		4.50
Great Paraiso Hot Soda				
Springs		160		.56
Piedmont White Sulphur		and the second		
"Iron Springs"	W. Anderson	161		.45
Santa Barbara Hot Springs	W. Anderson	162	2.90	
Santa Rosa White Sulphur				
Springs	W. Anderson	163		.93
Santa Ysabel Sulphur				
Springs, "Main Warm"	W. Anderson	164		-73
Santa Ysabel Sulphur				.,,
Springs, Cold Sulphur No. 1	W. Anderson	164		.83
Santa Ysabel Sulphur	W. IIII derboil			.03
Springs, Cold Sulphur No.2	W. Anderson	164		.84
Santa Ysabel Warm Sulphur	W. Hinderson	104		.04
3.5 1	W. Anderson	165		.65
Summit Soda Springs	W. Anderson	168		
	W. Anderson	0.0000000000000000000000000000000000000		1.13
Thermal Acid (Caso Range)	W. Anderson	169	7404.41?	
Tolenas Springs		170		1.10
Tuscan (or Lick) Springs	F. W. Hatch	170		a
Wilbur Springs	W. Anderson	174		3.93
Witter's Mineral Springs		174		1.65
Young's Natural-gas Well	717		1 - 1 - 1	
and Mineral Springs	W. Anderson	175		5.18
Colorado.	and the second second	- 2-1		
Manitou Soda Springs,				
Manitou Soda Springs,	E. Waller	187		0.4
Maintou	D. Waner	10/		.07

a. Qualitative analyses. Amount not reported.

Spring.	Analyst.	Page.	Aluminium Sulphate per U. S. Gallon.	Alumina per U. S. Gallon.
Manitou Soda Springs, Navajo Poncho Hot Springs Connecticut.	E. Waller	187	Grains.	Grains. .10 5.20
Stafford Spring	L. Norton	197		.11
Bowden Lithia Upper Spring Bowden Lithia Lower Spring Bowden Lithia Lower Spring Franklin Spring	Pratt Doremus	206 206 206 212	1.33 2.61 .53 a	
Hughes' Mineral Well	H. C. White	214	1.02	a
Trentham Spring	H. C. White	219		.46
Indan-Ha Spring	C. F. Chandler	222		.05
Illinois. Kane County Magnesia Spring	W. S. Hains	226		.09
Indian Alkaline Saline Spring	E. T. Cox W. A.Noyes	231 234	.82	.17
Iowa. Fry's Mineral Spring Lineville Mineral Springs	W. S. Hains A. E. Woodward	239 239		.06b
Geuda Spring No. 1	Bailey Bailey Bailey Bailey	244 244 244 244 244 247		.06 .01 .01 .01 .01
Kentucky. Big Bone Springs		250	a	
Maine. Keystone Spring Maryland.	R. C. Stanley	265		.850
Mardela Spring	P. B. Wilson	272		-34

a. Qualitative analyses. Amount not given.

b. Heinrichs, in an analysis of this water, reports .29 grains "Silica Alumina" per U. S. gallon.

c. With silica.

Spring.	Analyst.	Page.	Aluminium Sulphate per U. S. Gallon.	Alumina per U.S. Gallon.
Michigan.				
			Grains.	Grains.
Clark's Riverside Mineral	S. P. Duffield	286		
Springs	S. P. Dumeid	280		13.41
Eastman's Springs, King David		286		
Eastman's Springs, Bimini		287		.11
Eastman's Springs, Golden		201		.03
Fountain		288		.01
Owosso Mineral Water		293		.62a
St. Clair Mineral Spring	S. P. Duffield	296		830.00
Minesota.	MALE DE LA PROPERTIE DE LA PRO			
Indian Medical Spring	C. W. Drew	200		
White Mineral Spring	W. A. Noyes	299		.03
	W. A. Noyes	299		.14
Mississippi.			-	
Brown's Wells, Spring No. 1.	J .R. Chilton	302	8.11	
Brown's Wells, Spring No. 2.	G. Little	302		ь
Castilian Mineral Spring,	D W III		,	
No. 1	E. W. Hilgard	303	b	
Missouri.				
B. B. Mineral Water	P. Schweitzer	300	18.31	
Blue Lick Springs	P. Schweitzer	310		b
Blue Lick Springs		311		10.23
Excelsior Springs		312		6
Nevada.				
Walley's Hot Springs	J. W. Phillips	325		.02
	J 2	3-3		
New Hampshire.				
Bradford Mineral Spring	Jackson, also	327		6
Londonderry Lithia Spring	Richards (H. Halvorson	228	- 0-	
	II. Halvoison	328	5.05	
New Mexico.				
Las Vegas Hot Spring,				
No. 6	W. S. Hains	334		.10
New York.		1000		
Geneva Lithia Water	A. A. Cunningham	350	8.75	
Deer Lick Spring	C. F. Chandler	354		-34
North Carolina.		00.		
Thompson's Bromine-ar- senic Spring	H. Froehling	25.4		
seme opring	11. Proening	374		C

a With silica.

b. Amount not given. Qualitative analyses.
 c. Contains .12 grain aluminium phosphate per U. S. gallon.

Spring.	Analyst.	Page.	Aluminium Sulphate per U. S. Gallon.	Alumina per U.S. Gallon.
			Grains.	Grains.
North Carolina Hot Springs.	S Chandler and	376		.04
	A CALCII		and the state of the state of	
Park's Springs	A. R. Ledoux	378		3.50
Pennsylvania.				
Bedford Magnesia Spring	V. G. Bloede	400		.II
Bedford Bowling Alley				
Spring	V. G. Bloede	400		.09
Gaylord & Gulick Mineral				
Springs.	F. A. Genth	406	6.58a	
Minnequa Springs	C. M. Cresson	409		.126
Tuscarora Lithia Spring	W. A. Middleton?	416		С
South Dakota.				
South Dakota Hot Springs,				
Mammoth Mineral Spring.	C. B. Gibson?	430		.27
Lakatah Springs	C. B. Gibson?	431		.02
Tennessce.		11811		
Dixie Mineral Water	J. W. Slocum	425		20
Red Boiling Springs, No. 2	J. T. Anderson	435		.30
	(I M Sofford and	444		
Upper Red Boiling Spring	J. C. Wharton	446	.15	
Texas.	()	444	3	
Burdett Mineral Wells	W. H. Johnson	449		bc
Overall Mineral Well No. 1	E. Everhart	452	81.71	
Overall Mineral Well No. 2 .	E. Everhart	452	4.99	Budie
Utah.				
Midway Warm Springs	A. Meacham	460	.56	
Utah Hot Springs	S. F. Baird	461		.25
Virginia.				
Harris Antidyspeptic and		1777		
Tonic Springs, No. 1	E. T. Fristoe	476		.16b
Blue Ridge Springs	H. Froehling	480		d
Cold Sulphur Springs		482	2.46	
Crocket's Arsenic - Lithia	TT Description	.0-		HE PLAN
Springs	H. Froehling	485		6
Glenola (Wayland) Spring Hunter's Pulaski Alum	M. B. Hardin	489		f
Springs	W. H. Taylor	107	16.40	HOTE
-p.m.go	W. II. Taylor	497	10.40	,

a. Practically the same analyses as quoted by Peale (U. S. Geological Survey Bulletin 32, page 49) for Blassburg (Pa.) springs.

b. The value given refers to aluminium.

c. Qualitative analyses. Amount not reported.

d. Contains .02 grain aluminium phosphate and .15 grain aluminium silicate per U. S. gallon.

e. Contains trace of aluminium phosphate and .12 grains aluminium silicate per U. S. gallon.

f. Contains .04 grain aluminium phosphate and .20 grain aluminium silicate per U. S. gallon.

Spring.	Analyst.	Page.	Aluminium Sulphate per U. S. Gallon.	Alumina per U. S. Gallon.
Iron Lithia Springs	H. Froehling J. W. Mallet G. B. M. Zerr H. Froehling	497 499 500 506	Grains. 8.05 b	Grains. a 0.13 .26
West Virginia. Hart Well	S. C. Wells. W. W. Daniels	531		d
Fort Crawford Mineral Water	G Bode	535		.66
Fountain Geyser, Yellow- stone National Park		552		.96e

a. With .11 grain aluminium phosphate per U. S. gallon in addition.

Dambergis, A. K. (Ber. Deut. Chem. Gesellsch., 20, p. 3328; in Jour. Chem. Soc. London, 54 (1888), II, p. 238), in the water of one of the sulphur springs of the peninsula of Methana on the east side of the Chelona range near the village of Wromolimni, reports 0.019 parts Al₂O₃ per 10,000.

Dambergis (Ber. Deut. Chem. Gesellsch., 19, p. 2538; Jour. Chem. Soc. London, 52 (1887), II, p. 23) reports 0.0200 alumina in water from Ægina Springs and 0.06000 in water from Andros, grams per 10,000 cc. in both cases.

Davis, R. H. (Jour. Chem. Soc. London, 39, p. 19; Jahresb. Chem., 1881, p. 1449), notes 12.78 gms. Al₂O₃ per 10,000 cc. in the Harrowgate, England, "Old Alum Well."

Darton, N. H. (U. S. Geological Survey Bul. 138), in his report on artesian well prospects on the Atlantic coastal plain region, quotes a number of analyses of water in which iron and alumina are reported together.

b. With trace of aluminium phosphate in addition.

c. With .02 grain aluminium phosphate per U. S. gallon in addition.

d. Contains .03 grain aluminium phosphate and .09 grain aluminium and iron carbonate per U. S. gallon. See also analyses quoted by Peale, U. S. Geological Survey Bulletin 32, page 73.

e. Chemical combinations calculated by E. E. Smith from analytical data.

De Luca (Compt. Rend. Acad. Sci. Paris, 67, p. 909; Inst. 1868, p. 354; Jahresb. Chem. Will., 1868, p. 1040) reports traces of alumina in the "Solfatare" of Puzzuoli.

De Luca, S. (Compt. Rend. Acad. Sci. Paris, 70, p. 408; Jour. Pharm., 4th ser., 12, p. 33; Inst. 1870, p. 49; Chem. Centbl., 1870, p. 180; Chem. News, 21, p. 119; Jahresb. Chem., Naumann, 1870, p. 1389), reports 0.335 gm. Al₂O₃ per 1000 cc. in the "Solfatare" water from Pozzuoli.

De Negri, A. and E. (Gaz. Chim. Ital., 8 (1878), p. 120; Jour. Chem. Soc. London (1878), II, p. 715), note 0.000417 gm. aluminium phosphate per liter in mineral water from Casteggio, Italy.

Devarda, A. (Chem. Centbl., 1891, II, p. 365; Jahresb. Chem., 1891, p. 2619), in mineral water from Costalta (Southern Tyrol), notes 0.01614 gm. Al₂O₃. 3SiO₂ per 1000 cc.

Deville (Ann. Chim. Pharm., 3d ser., 23, p. 32; Arch. Pharm., 2d ser., 55, p. 301; Jahresb. Chem., 1847–48, p. 996) analyzed a number of waters, finding alumina as follows (grams per 100,000 cc.): Seine River at Bercy, 0.0005; Rhine at Strasburg, 0.0025; Loire at Orleans, 0.0071; Rhone at Geneva, 0.0039; Doubs at Rivotte, 0.0021; Mouillère spring-water, 0.0043; Billecul Spring, 0.0043; Arcier Spring, 0.0090; Bregille Spring at Besançon, 0.0065; Suzon Spring at Dijon, 0.0010; Arcueil at Paris, 0.0053; well in the Grand Rue, 0.0094; well Rue de la Préfecture, 0.0062; and at the Faculté des Sciences at Besançon, 0.0039.

Dewar, J. (Chem. News, 24, p. 171; Jour. Chem. Soc. London, 25 (1872), p. 60), found 1.8 grains Al₂O₃ per gallon in the acidulous chalybeate water from Melrose.

Dietrich, H. (Min. Petr. Mitth., 2d ser., 3, p. 439; Jahresb. Chem., 1880, p. 1527), reports 0.005, 0.006, and 0.005 gm. Al₂O₃ per 10,000 gms., respectively, in three mineral springs at Krynica, Austria-Hungary.

Dietrich, H. (Jahrb. K. K. Geol. Reichsanst., 43 (1893), p. 275; Jour. Chem. Soc. London, 70 (1896), II, p. 435), found 0.00335 Al₂O₃ per 1000 parts in the water of the Klebelsberg Spring, Ischl, Austria.

Dormoy (Ann. Min., 6th ser., 12, p. 461; Jahrb. Chem. Will., 1867, p. 1040) reports alumina with iron oxid or iron and

manganese oxid in fourteen waters from Luxeuil, the amount ranging from 0.00157 to 0.01486 gm. per liter.

Drown, T. M. (State Board Health Mass. Rpt., 24 (1892), p. 345), reported analyses of a number of waters from Massachusetts. The data regarding alumina follow:

ALUMINIUM IN MASSACHUSETTS WATERS.

Kind of Water.	Alumina. Parts per 100,000.
Mansfield, well	0.0173
Framingham, Spring northeast of sewage fields	.0444
Stoughton, well	.0475
Everett, Spring	.0325
Malden, tubular wells	.0125
ramingham, Spring northwest of sewage field	.0250
Ivde Park, Neponset River	.0062
Ivde Park, tubular wells near river	.0440
Iyde Park, starch-factory well near river	.0175
Voburn, Horn Pond	.0400
Voburn, filter-gallery	.0143
Vayland, reservoir	.0360
Vayland, filter-gallery	.0475
Vestborough, Insane Hospital tubular wells	.0250
Reading, filter-gallery	.0493
Bradford, Well No. 7	.0250
Bradford, Well No. 12	.0100
Marblehead Water Co., Swampscott, large wells and tubular	
wells	.0125
farblehead, town supply, large well and tubular wells	.0110
awrence, sewage filtered through filter-paper	.3850
awrence, effluent from intermittent filtration of sewage.	
Tank No. 4	.0266
awrence, effluent from intermittent filtration of sewage.	
Tank No. 9	.0457

Du Mênil (Arch. Pharm., 2d ser., 69 (1852), p. 1; Pharm. Centbl., 1852, p. 229; Jahresb. Chem., 1852, p. 754) found 0.500 grain alumina (with silica) per 3 pounds in sulphur-water from Seebruch near Vlotho.

Du Ponteil (Ann. Chem. Pharm., 96, p. 193: Jour. Prakt. Chem., 67, p. 249; Chem. Centbl., 1856, p. 4; Ann. Chim. Phys., 3d ser., 46, p. 233; Jahresb. Chem., 1855, p. 831) found 0.3546 Al₂O₃·3SO₃ per 100 parts in the clear yellow very acid water of a hot lake on the volcanic White Island, Plenty Bay, New Zealand.

Eakins, L. G. (U. S. Geol. Survey Bul. 60, p. 172), found in water from a 1000-foot well in Lebanon, Missouri, 0.0032 gm. Al₂O₃ per liter.

Effenberger, A. (Wiener Akad. Ber. 51, 2. Abt., p. 252; Chem. Centbl., 1865, p. 848; Jahresber. Chem. Will., 1865, p. 934), notes per 10,000 parts, 0.006 part alumina in the medicinal springs at Müllaken in Upper Austria.

Egger, E. (Chem. Centbl., 1881, p. 664; Jahresb. Chem., 1881, p. 1443), reports an analysis of the Adelheidquelle at Heilbrunn, Germany. This water contained 0.0010 gm. Al₂O₃ per 1000 cc.

Egger, E. (Chem. Centbl., 1882, p. 187; Jahresb. Chem., 1882, p. 1629), in a sulphur spring at Seon, Germany, reports 0.0018 gm. aluminium phosphate per 1000 cc.

Emerson, B. K. (U. S. Geol. Survey Mon. 29, p. 750), quotes an analysis made by S. D. Hayes of the water of Mount Mineral Spring, Shutesbury, Massachusetts. A trace of alumina is reported.

Emerson, B. K. (U. S. Geol. Survey Bul. 159, p. 91), reports 12.80 Al₂O₃ parts per 1,000,000 in water from an artesian well in Dalton, Massachusetts.

Emmons, S. F. (U. S. Geol. Survey Rpt., 17 (1895-96), pt. II, p. 411), in an article on the mines of Custer County, Colorado, reports a number of analyses by Hillebrant. Vadose water from the Geyser mine, 500-foot level, contained 0.8 Al₂O₃·P₂O₅ parts per 1,000,000 (p. 461); deep water from the Geyser mine, 2000-foot level, 1.06 Al₂O₃ parts per 1,000,000 (p. 462). These values refer to the theoretical composition of the water before sediment was deposited.

Essner, J. C. (Bul. Soc. Chim., 3d ser., 6, p. 148; Jahresb. Chem., 1891, p. 2615), reports 0.240 gm. aluminium sulphate (+18H₂O) per 1000 cc. in a subterranean water from the neighborhood of Port Vendres.

Essner, J. C. (Bul. Soc. Chim., 3d ser., 7, p. 480; Jahresb. Chem., 1892, p. 2688), reports 0.0240 gm. (?) Al₂O₃ per 1000 cc. in water from the iron sulphur spring at Roufaque.

Fehling (Württemberg. Naturwissensch. Jahreshefte, 13, p. 113, Jahresb. Chem., 1857, p. 720) found traces of alumina

in the following artesian mineral water from the Stuttgart bath near Berg: Haupt-Trinkquelle and Westliche Quelle.

Fehling (Württemberg. Naturw. Jahreshefte, 16, pp. 106, 129; Neues Jahrb. Pharm., 14, pp. 286, 295; Jahresb. Chem., 1860, pp. 833, 834) reports alumina as follows (grams per 100,000) in Württemberg mineral waters: Wildbad Warm Springs; Trinkquelle No. 10, Trinkhalle, 0.055; Quelle No. 19, Katherienbad, 0.059; mixture of water from ten springs, 0.070; Teinach mineral springs, Hirschquelle, 0.126; Bachquelle, trace; and Dintenquelle, 0.071.

Fehling, H. (Württemb. Naturw. Jahresb., 22, pp. 129, 147, 159; Jahresb. Chem. Will., 1867, p. 1035), notes an average of 0.055 parts of alumina per 100,000 in waters from Wildbad; 0.038 in water from Liebenzell; traces in Bachquelle from Teinach, and 0.126 in Hirschquelle from Teinach.

Feliciani, G. (Gaz. Chim. Ital., 26 (1896), I, p. 281; Jour. Chem. Soc. London, 70 (1896), II, p. 615), notes a trace of alumina in the water of the acid spring at Ponte Molle near Rome.

Fellenberg reports (Untersuchung d. Schwefelwasser d. Gurnigelbades, Bern, 1849; Jahresber. Chem. Will., 1850, p. 623) 0.0051 part aluminium silicate per 10,000 gms. in one water analyzed, designated No. A. Further, 0.08 alumina and iron oxid in another, designated No. B. The author also analyzed the water from two other sulphur springs from the same neighborhood. No aluminium was reported in either. In the sediment of the Stockquelle (l. c., p. 624) he found 14.47 per cent alumina (with calcium phosphate). The Stockquelle water was one of those mentioned which contained no alumina.

Ferstl (Jahrb. K. K. Geol. Reichsanstalt, 1853, No. 4, p. 683; Jahresb. Chem., 1853, p. 712) reports in the Luhatschowitz (Austria) mineral water, Vincenzbrunnen, Amandbrunnen, Johannisbrunnen, and Luisenquelle, aluminium phosphate (parts per 1000) as follows: 0.0047, 0.0048, 0.0041, and 0.0086, respectively.

Figuier and Mialhe (Jour. Pharm., 3d ser., 13, p. 401; Jour. Chim. Med., 3d ser., 4, p. 635; Pharm. Centbl., 1848, p. 662; Jahresb. Chem., 1847–48, p. 1005) note a trace of alumina in Niederbronn (Alsace) and alumina (grams per 1000 cc.)

as follows in mineral water from Bourbonne: Source de la Place, 0.030; Source de l'intérieur de l'établissement, 0.029. They also note (Jour. Pharm., 3d ser., 11, p. 338; Jour. Prakt. Chem., 42, p. 465; Pharm. Centbl., 1847, p. 431; Jahresb. Chem., 1847-48, p. 1006) a trace of aluminium in mineral water from Rieumajou near Salvétat, Dept. de l'Hérault.

Filhol (Jour. Pharm., 3d ser., 20, p. 81; Jahresb. Chem., 1851, p. 664) reports aluminium silicate (gms. per 1000 cc.) in Bagnères-de-Luchon springs as follows: Bayen, a trace; Azémar, 0.0237; Richard (upper spring), 0.0292; Grotte supérieure, 0.0109; Blanche, 0.0101; Ferras (upper spring No. 2), trace; Pré No. 1 and Bordeu No. 4, each 0.0073; Grotte inférieure, 0.0141, and La Reine, 0.0274. Ferras also contained 0.0022 gm. alumina per 1000 cc., and La Reine a trace.

Filippuzzi (Wiener. Acad. Ber. 21, p. 561; Chem. Centbl., 1856, p. 937; Jahresb. Chem., 1856, p. 773) reports 0.01927 aluminium phosphate and 0.36004 aluminium sulphate per 10,000 parts in mineral water from Valdagno, Italy.

Finckh, C. (Neue. Jahrb. Pharm., 34, p. 13; Chem. Centbl., 1870, p. 615; Jahresb. Chem. Naumann, 1870, p. 1382), notes traces of aluminium in Ochsenhausen mineral water from Bieberach, Germany.

Flückiger, F. A. (Mittheil. Naturf. Gesellsch. Bern, 1862, p. 17; Arch. Pharm., 2d ser., 111 (1862), p. 111; Vierteljahressch. Prakt. Pharm., 11, p. 342; Jahresb. Chem., 1862, p. 820), analyzed a sample of acid water taken by Stöhr and Zollinger in 1858 at the first waterfall of the brook Sungi Paït flowing from the lake in the crater of Idjen volcano in Java. It contained per 100 gms. 0.150 Al₂O₃.

Folberth (Verhandl. und Mittheil. Siebenbürg. Vereins Naturwissensch., 1855, No. 7; Jahresb. Chem., 1855, p. 844) reports 0.0222 basic aluminium phosphate per 1000 parts in the saline water from the Felsenquelle near Bassen, Hungary.

Folberth, F. (Verhandl. u. Mittheil. Siebenbürg. Vereins Naturw. Hermannstadt, 11, p. 78; Jahresb. Chem., 1861, p. 1102), notes Al₂O₃ (grams per 10,000) as follows in mineral water: Pokolsár, 0.142, and Czifra-víz, 0.402.

Fresenius (Untersuchungen d. Mineralwasser d. Herzogthums Nassau, I, Wiesbaden, 1850; Jahresb. Chem., 1850, p. 622) notes 0.0051 gm. aluminium silicate per 10,000 gms. in the Wiesbaden Kochbrunnen water.

Fresenius (Jahrb. Vereins. Naturk. Herzogthum Nassau, 7, pt. p. 145; Untersuchungen der Mineralwasser des Herzogthums Nassau, II, Wiesbaden, 1851; see also Ann. Chem. Pharm., 82, p. 249; Jahresb. Chem., 1851, p. 652) examined the mineral waters of Ems, reporting per 1000 parts aluminium phosphate as follows: Kesselbrunnen, 0.00125 part; Krähnchen, 0.00042 part; Fürstenbrunnen, 0.00044 part; Neue Quelle, 0.00142 part.

Fresenius (Chem. Unters. Mineralwasser. Herzogth. Nassau, III, Ann. Chem. Pharm., 83, p. 252; Pharm. Centbl., 1853, p. 45; Jahresb. Chem., 1852, p. 753) notes a trace of alumina in the warmest Schlangenbad mineral water.

Fresenius (Jour. Prakt. Chem., 58, p. 156; Arch. Pharm., 2d ser., 75, p. 301; Pharm. Centbl., 1853, p. 405; Jahresb. Chem., 1853, p. 709) in the Bernhard spring and the Johann-Georgen spring at Krankenheil-Tölz, Bavaria, found 0.002034 and 0.002782 gm. aluminium silicate per 1000 gms. respectively.

Fresenius (Jahrb. Ver. Naturk. Herzogthums Nassau, No. 11; Jour. Prakt. Chem., 70, p. 1; Neue. Jahrb. Pharm., 7, p. 7; Chem. Centbl., 1857, p. 49; Jahresb. Chem., 1856, p. 770) found 0.000133 gm. aluminium phosphate per 1000 gms. in water from the sulphur spring at Weilbach, Nassau, Germany.

Fresenius (Jour. Prakt. Chem., 72, p. 1; Chem. Centbl., 1857, p. 913; Jahresb. Chem., 1857, p. 720) notes a trace of alumina in the mineral spring at Geilnau in Nassau, Germany.

Fresenius, R. (Jour. Prakt. Chem., 98, p. 321; Jahresb. Chem. Will., 1866, p. 989), reports 0.000254 aluminium phosphate per 1000 parts in the potable spring-water (Trinkquelle) of Driburg and 0.000335 aluminium phosphate in the Herster mineral spring near the same place.

Fresenius, R. (Jour. Prakt. Chem., 95, p. 151; Chem. Centbl., 1865, p. 728; Vierteljahresschr. Prakt. Pharm., 15, p. 208; Jahresber. Chem. Will., 1865, p. 929), notes aluminium phosphate per 1000 parts in three springs at Pyrmont as follows: Stahlbrunnen, 0.000084; Brodelbrunnen, 0.000295; and Klos-

teralleequelle, 0.000091. The water of the first is used for drinking, the second for bathing.

Fresenius, R. (Jour Prakt. Chem., 97, p. 1; Chem. Centbl., 1866, p. 335; Jahresb. Chem. Will., 1866, p. 990), reports 0.000102 part aluminium phosphate per 1000 in Felsenquelle No. 2 at Ems (Bad Ems).

Fresenius, R. (Jahrb. nassau. Verein Naturkunde, pt. 19, 20 (1864), pp. 453, 488; Jour. Prakt. Chem., 103, pp. 321, 425; Chem. Centbl, 1868, pp. 703, 800; Jahrb. Min., 1868, p. 629; Jahrb. Chem. Will., 1867, p. 1034), reports 0.000430 aluminium phosphate per 1000 parts in a mineral water from Niederselters.

Fresenius, R. (Jour. Prakt. Chem., 106, pp. 193, 206; Chem. News, 20, p. 213; Jahresb. Chem. Will., 1869, p. 1281), found in Tönnisteiner Heilbrunnen, Tönnisteiner Stahlbrunnen, and Lamscheider Mineralbrunnen, 0.00013, 0.00045, and 0.000460 part Al₂O₃P₂O₄ per 1000 respectively. In the latter 0.000034 part alumina united with silica is also reported.

Fresenius, R. (Jahrbücher nassau. Ver. Naturkunde, 25 and 26, pp. 347, 361; Jahresb. Chem., 1871, p. 1226), notes 0.000134 part aluminium phosphate per 1000 in Victoria and 0.000120 part in Römerquelle water from Ems.

Fresenius, R. (Jour. Prakt. Chem., 2d ser., 9, p. 368; Jour. Chem. Soc. London, 27 (1874), p. 968), notes 0.000193 part aluminium phosphate per 1000 in water from the warm mineral spring in the bath-house of the Royal William Medical Establishment at Wiesbaden.

Fresenius, R. (Jour. Prakt. Chem., 2d ser., 6, p. 53; Jahresb. Chem. Naumann, 1872, p. 1181), reports new analyses of the mineral springs at Ems. These were Kränchen, Fürstenbrunnen, Kesselbrunnen, and Neue Badequelle. They contained respectively, per 1000 parts, 0.000116, 0.000117, 0.000200, and 0.000209 part aluminium phosphate.

Fresenius, R. (Jour. Prakt. Chem., 2d ser., 7, p. 191; Jahresb. Chem. Naumann, 1873, p. 1238), notes 0.000021 part aluminium phosphate per 1000 parts in Carlsquelle water from Bad Helmstedt.

Fresenius, R. (Jour. Prakt. Chem., 2d ser., 9, p. 368; Jahresb. Chem., 1874, p. 1325), reports 0.000193 part aluminium phosphate per 1000 parts in the mineral water of Wilhelmsheilanstalt, Wiesbaden.

Fresenius, R. (Jour. Prakt. Chem., 2d ser., 25, p. 310; Jahresb. Chem., 1882, p. 1628; Jour. Chem. Soc. London, 40 (1882), p. 1178), notes traces of aluminium in the Oberbrunnen at Salzbrunn, Germany. In this same water Valentiner in 1866 noted 0.0005 gm. alumina and phosphoric acid per 1000.

Fresenius, R. (Jour. Prakt. Chem., 2d ser., 45, p. 287; Jahresb. Chem., 1892, p. 2684; Jour. Chem. Soc. London, 62 (1892), p. 796), reports per 1000 gms. in Julianenbrunnen and Georgenbrunnen water from Bad Eilsen 0.000126 and 0.000136 alumina respectively, in addition to 0.000155 and 0.000060 aluminium phosphate.

Fresenius, C. R. (Jahrb. nassau. Ver., 46 (1893), p. 1; Jour. Chem. Soc. London, 70 (1896), II, p. 315), notes a trace of alumina in the water of the Victoria Spring at Oberlahnstein.

Fresenius, H. (Jour. Prakt. Chem., 2d ser., 35, p. 237; Jahresb. Chem., 1887, p. 2532), found 0.000334 gm. aluminium phosphate and 0.000401 gm. aluminium silicate per 1000 gms. Schützenhofquelle water, Wiesbaden, Germany.

Fresenius, H. (Jahrb. nassau. Ver. Naturk., 51 (1898), p. 1; abs. Jour. Chem. Soc. London, 76 (1899), II, p. 114), finds in water from a new boring at Selters near Weilburg on the Lahn a trace of alumina.

Garrigau (Compt. Rend. Acad. Sci. Paris, 84, p. 963; Jahresb. Chem., 1877, p. 1387) notes 0.0096 part Al₂O₃ per 1000 in the Cliff Spring at Saint-Nectaire le Haut, France.

Genth (Keller and Tiedemann's Nord. Amer. Monatsber., 1852, June, p. 246; Pharm. Centbl., 1852, p. 588; Jahresb. Chem., 1852, p. 758) notes a trace of alumina in mineral water from Bristol near Philadelphia, Pennsylvania.

Giles (Pharm. Jour. Trans., 7, p. 75; Jahresb. Chem., 1847–48, p. 998) notes 0.260 gm. alumina per 10,000 gms. in springwater from Wolverton.

Gintl, W. F. (Jour. Prakt. Pharm., 2d ser., 20, p. 356; Jahresb. Chem., 1879, p. 1264), in Ferdinand Spring, Marienbad, Bohemia, reports 0.06334 gm. basic aluminium phosphate per 10,000 gms.

Gintl, W. F. (Jour. Prakt. Chem., 2d ser., 24, p. 25; Jahresb. Chem., 1881, p. 1445), notes 0.05256 Al₂P₂O₈ per 10,000 parts in Ambrosius water from Marienbad, Bohemia.

Gintl, W. F. (Separate; abstracted in Jahresb. Chem., 1882, p. 1632), notes 0.0744 part basic aluminium phosphate in the West spring and 0.1019 part in the East spring at Langenbruck near Franzensbad, Austria-Hungary.

Girardin (Jour. Chim. Med., 3d ser., 4, p. 643; Jour. Pharm., 3d ser., 15, p. 113; Jahresb. Chem., 1847–48, p. 1006) found a trace of aluminium sulphate in St. Paul spring-water from Rouen.

Glasel, E. (Jahrb. Geol. Reichsanst., 19, p. 295; Chem. News, 20, p. 190; Jahresb. Chem. Will., 1869, p. 1290), reports 0.0028 gm. aluminium per 1000 in mineral water from Rajec-Töplitz.

Gläser, M., and W. Kalmann (Ber. Deutsch. Chem. Gesellsch., 21 (1637); Jour. Chem. Soc. London, 54 (1888), p. 796), in the Roncegno (South Tyrol) water, report 0.4343 gm. Al₂O₃ per liter.

Godeffroy, R. (Sep. from Ztschr. Allgm. Oesterr. Apothekervereins; Jahresb. Chem., 1882, p. 1623), found in water from the middle of Gmunden Lake 0.100 part Al₂O₃ per 100,000.

Gooch, F. A., and J. E. Whitfield (U. S. Geol. Survey Bul. 47, pp. 36-81, and table facing page 82) report, in analyses of waters of the Yellowstone National Park, the following, the quantities in every case being grams per kilogram of water. In every case the author gives the calculated amount of the aluminium compound present, as well as the amount of alumina determined:

Cleopatra Spring, 0.0049 Fe.Al calculated = 0.0093 Al₂O₃; Hot River, 0.0051 Fe.Al = 0.0097 Al₂O₃; Gardiner River, sample taken Oct. 12, 1883, above Hot River, 0.0042 Fe.Al = 0.0079 Al₂O₃; Gardiner River, sample taken Sept. 26, 1884, at Mammoth Hot Springs, 0.0010 Fe.Al = 0.0019 Al₂O₃; Watersupply at Mammoth Hot Springs, 0.0011 Fe.Al = 0.0021 Al₂O₃; Soda Spring, 0.0004 Al = 0.0008 Al₂O₃; Fearless Geyser, 0.0002 Al = 0.0004 Al₂O₃; Pearl Geyser, 0.0031 Al = 0.0059 Al₂O₃; Constant Geyser, 0.0048 = 0.0304 Al₂(SO₄)₃; Coral Spring, sample taken Aug. 20, 1884, 0.0029 Al = 0.0143 Al₂Cl₆; Coral Spring, sample taken Oct. 11, 1886, 0.0077 Al = 0.0139 Al₂O₃; Echinus Spring, 0.0027 Al = 0.0171 Al₂(SO₄)₃; Schlammkessel, 0.0081 Fe.Al = 0.0513 Al₂(SO₄)₃.Fe₂(SO₄)₃; Fountain Geyser,

0.0057 Al=0.0108 Al₂O₃; Great Fountain Geyser, 0.0021 Al = 0.0040 Al₂O₃; Hygeia Spring, 0.0036 Al = 0.0068 Al₃O₃; Firehole River, 0.0031 Al=0.0059 Al₂O₃; Excelsior Geyser, 0.0012 Al=0.0023 Al2O3; Old Faithful Geyser, 0.0009 Al= 0.0017 Al₂O₃; Splendid Geyser, sample taken Sept. 10, 1885, 0.0027 Al=0.0051 Al2O3; Splendid Geyser, sample taken Aug. 28, 1884, 0.0034 Al = 0.0064 Al₂O₃; Giantess Geyser, 0.0049 Al = 0.0093 Al₂O₃; Beehive Geyser, 0.0029 Al = 0.0055 Al₂O₃; Grotto Geyser, 0.0036 Al=0.0068 Al2O3; Turban and Grand Geysers, 0.0032 Al=0.0061 Al₂O₃; Artemisia Geyser, 0.0079 Al = 0.0150 Al₂O₃; Taurus Geyser, 0.0040 Al = 0.0075 Al₂O₃; Asta Spring, 0.0059 Fe.Al=0.0112 Al2O3; Bench Spring, 0.0145 Fe.Al = 0.0066 Al2(SO4)3+0.0263 Al2O3; Firehole River, 0.0029 Al=0.0055 Al2O3; Yellowstone Lake, 0.0021 Al=0.0040 Al2O3; Alum Creek, 0.0025 Al=0.0158 Al2(SO4)3; Chrome Spring, 0.0043 Fe.Al=0.0082 Al₂O₃; Mush Pot Spring, 0.0045 Al = 0.0285 Al₂(SO₄)₃; Devil's Ink Pot, 0.0037 Al = 0.0234 Al₂(SO₄)₃; Soda Butte Spring, 0.0069 Fe.Al=0.0131 Al₂O₃.

Gorup-Besanez (Ann. Chem. Pharm., 79, p. 50; Pharm. Centbl., 1851, p. 718; Jahresb. Chem., 1851, p. 653) found alumina (unweighable amount) in mineral water from Stebin, Bavaria.

Gossart (Jour. Pharm., 4th ser., 11, p. 292; Chem. News, 21, p. 214; Jahresb. Chem. Naumann, 1870, p. 1389) notes traces of Al₂O₃ in a sulphur spring near Meurchin, Pas-de-Calais.

Göttl (Oester. Ztschr. Pharm., 1853, pp. 253, 266; Jahresb. Chem., 1853, p. 711) found 0.0040 gm. alumina per 1000 gms. in the Karlsbad Schlossbrunnen.

Göttl (Vierteljahressch. Prakt. Pharm., 4, p. 192; Arch. Pharm., 2d ser., 84, p. 179; Pharm. Centbl., 1855, p. 286; Jahresb. Chem., 1855, p. 841) notes 0.022 alumina per 10,000 parts in Giesshübler water from the Rodisfort(Germany) acid spring.

Göttl (Vierteljahressch. Prakt. Pharm., 5, p. 161; Jahresb. Chem., 1856, p. 772) notes 0.028 alumina per 1000 parts in the Karlsbad Sprudel water.

Gottlieb (Wiener. Acad. Ber., 30, p. 191; Chem. Centbl., 1858, p. 612; Jahresb. Chem., 1858, p. 796) notes 0.019 basic aluminium phosphate per 10,000 parts in Marienbrunnen water from Gabernegg, Austria-Hungary.

Gottlieb (Wiener. Akad. Ber., 56, 2. Abt., p. 836; Jour. Prakt. Chem., 102, p. 472; Jahresb. Chem. Will., 1867, p. 1038) reports 0.0147 aluminium phosphate per 10,000 parts in water from Emmaquelle at Gliechenberg in Steiermark.

Gottlieb, J. (Wiener. Akad. Ber., 60, 2. Abt., pp. 349, 357; Jahresb. Chem. Will., 1869, p. 1287), found in "old" and "new" Johannes water, and Hauptquelle from Neuhaus (all in Steiermark), 0.0233, 0.0481, and 0.0026 part aluminium phosphate per 10,000 respectively.

Gottlieb, J. (Wiener Akad. Ber., 62 2. Abt., p. 780; Jahresb. Chem. Naumann, 1870, p. 1386), notes in the Königsbrunnen from Kostreinitz in Steiermark 0.0213 part aluminium phosphate per 10,000.

Graham, T., W. A. Miller, and A. W. Hofmann (Rpt. Govt. Commission on the Chemical Quality of the Supply of Water to the Metropolis, London, 1851; Quart Jour. Chem. Soc., 4, p. 375; Jahresb. Chem., 1851, p. 656) note traces of alumina in water supplied by New River Water Co., East London Water Co., Kent Water Co., and Hampstead Water Co. In the second the alumina reported equals 0.47 grains per gallon with iron and phosphoric acid salts. The waters were not from the Thames. Thames water and five spring-waters from the Hindhead (Surrey) district, Farnham and Gravesend, were also found to contain alumina with iron and phosphate.

Grandeau, L. (Ann. Chim. Phys., 3d ser., 60, p. 479; Jahresb. Chem., 1860, p. 839), reports 0.0408 alumina per 1000 cc. in mineral water from Pont-a-Mousson, Dept. Meurthe, France.

Grange (Ann. Chim. Phys., 3d ser., 24, p. 496; Jahresb. Chem., 1847-48, p. 996) reports in the Isère water at Grenoble 0.0035 gm. alumina per 100,000 cc.

Grange (Jahresber. Chem., 1850, p. 622) studied the water of the Isèrethal. He regarded the earthy taste of some springwater as due to alumina held in solution by CO₂.

Griffin, M. L. (unpublished data), found 0.718 gm. aluminium sulphate per liter in water from a spring running through the new engine-pit of the Boston and Maine Car-shops in Mechanics-ville, New York.

von Gümbel, C. W. (Chem. Centbl., 1891, II, p. 566: Jahresb.

Chem., 1891, p. 2618), notes analyses by A. Schwager of Pliniusquelle and Ostgothenquelle thermal water from Bormio and the Ortler region. The former contained 0.0137 gm. Al₂O₃ per 1000 cc.; the latter 0.0042 gm.

Günsberg, R. (Wiener. Acad. Ber., 43, 2. Abt., p. 197; Jahresb. Chem., 1861, p. 1103), reports 0.0012 gm. aluminium phosphate per 1000 gms. in Siegwasser from the Bronislaw spring at Truskawice, Galicia.

Guyot, P. (Compt. Rend. Acad. Sci. Paris, 77, p. 1384; Jahresb. Chem. Naumann, 1873, p. 1242), notes 0.010 gm. SiO₂.Al₂O₃ per liter in Saint-Thiebaut water from Nancy.

Hamberg, N. P. (Jour. Prakt. Chem., 80, p. 385: Chem. Centbl., 1860, p. 955; Ztschr. Ges. Naturw., 17, p. 71; Jahresb. Chem., 1860, p. 842), reports aluminium sulphate (parts per 10,000) in the water of medicinal springs at Ronneby. Sweden, as follows: Eckholzquelle, 15.038230, and Alte Quelle, 3.834603.

Hardin, M. B. (Amer. Chemist, 4, p. 247, Arch. Pharm., 3d ser., 5 (1874), p. 180; Jahresb. Chem., 1874, p.1336), made analyses of four alum springs in Rockbridge County, Virginia, and reports 5.35961, 7.30856, 7.53946, 12.41395. and 3.25892 gms. alumina per 1000 cc. The water was collected under different conditions as regards rainfall. The residue obtained by evaporating the water contained 9.490 per cent alumina.

von Hauer (Jahrb. K. K. Geol. Reichsanstalt, 1853, p. 154; Jahrsb. Chem., 1853, p. 712) found in mineral water from Roggendorff, Hungary, 0.033 parts aluminium sulphate per 1000 parts.

von Hauer (Jahrb. K. K. Geol. Reichsanstalt, 1858, p. 165; Jahresb. Chem., 1858, p. 798) notes 0.0013 alumina per 1000 parts in the thermal sulphur-water from Warasdin-Töplitz, Croatia.

von Hauer (Jahrb. K. K. Geol. Reichsanstalt, 1861, p. 57: Jahresb. Chem., 1861, p. 1100) notes 0.023 grains alumina per pound (=7680 grains) in acid water from Suliguli near Visco in Marmarosch.

Hehner, O. (Chem. News, 38, p. 249; Jahresb. Chem., 1878. p. 1314), in brown water from a spring 32 kilometers from Cape Town notes 525.19 parts Al₂S₃O₁₂ per 100,000.

Heller, F. (Wiener. Acad. Ber., 19, p. 363; Jahresb. Chem., 1856, p. 772), notes 0.030 alumina per 10,000 parts in water from the Franz Joseph Quelle at Füred on the Plattensee.

Henry (Jour. Pharm., 3d ser., 13, p. 5; Pharm. Centbl., 1848, p. 264; Jahresb. Chem., 1847-48, p. 1007) reports aluminium silicate (grams per 1000 cc.) as follows: in Source Grande Grille, 0.230; Source nouvelle, 0.233; Source Prè Salé, 0.070 (Vichy); Source de l'Hôpital, 0.120, and Source du Puits at Cusset, 0.080.

Henry, O. (Jour. Chim. Med., 3d ser., 6, p. 314; Jahresb. Chem., 1850, p. 628), notes 0.368 parts alumina per 1000 parts in the Fraysse mineral water from Cransac, Dept. Aveyron. Poumerède (Jahresb. Chem., 1850, p. 628) found in the same water 0.369 part alumina per 1000 parts. Henry (Jour. Pharm., 3d ser., 17, p. 161; Jahresb. Chem., 1850, p. 628) also notes aluminium sulphate and aluminium ammonium sulphate in Source-haute (Richard) and Source-basse (Richard) mineral water from Cransac.

Henry, O. (Jour. Pharm., 3d ser., 19, p. 104; Jahresb. Chem., 1851, p. 663), notes aluminium silicate and lithium silicate in mineral water from Sail-lès Chateaumorand as follows: Hamel or Saule, d'Urfé, Roman, Sulphur and Iron Sulphur springs: also alumina with calcium sulphate and silica in the Bellety (steel) spring from the same place.

Henry, O. (Jour. Pharm., 3d ser., 20, p. 161; Pharm. Centbl., 1851, p. 748; Jahresb. Chem., 1851, p.664), reports alumina and silica (parts per 1000) as follows: in spring-water from Saint-Denis near Blois, Dept. Loire and Cher, Medicis, 0.007; Renaulme, 0.007; and Saint-Denis, 0.044.

Henry, O. (Jour. Pharm., 3d ser., 21, p. 401; Jahresb. Chem., 1852, p. 757), in mineral water from Saint-Honoré, Dept. Nièvre, notes 0.023 part aluminium silicate per 1000 parts.

Henry, O. (Jour. Pharm., 3d ser., 30, pp. 172, 246; Jahresb. Chem., 1856, p. 774), reports 0.050 gm. alumina and silica per 1000 cc. in mineral water from Saxon, Canton Wallis, Switzerland.

Henry, O., Sr. according to Grasset (Compt. Rend. Acad, Sci. Paris, 46, p. 182; Inst., 1858, p. 37; Jahresb. Chem., 1858. p. 802), found 0.128 gm. alumina and silica per 1000 cc. in water from Bondonneau, Dept. Drôme, France.

Henry, O., and L'Héritier (Jour. Pharm., 3d ser., 28, pp. 333, 408; Jahresb. Chem., 1855, p. 834) found alumina (grams per 1000 cc.) in mineral water from Plombières as follows: Source du Crucifix, 0.0120; Sources des Dames, 0.0100; Source de Sainte-Catherine, 0.0110; Bain Romain, 0.0130; Bain tempère, 0.0110; Source du Savonneuse, 0.01400; and Source ferrugineuse de Bourdeille, 0.00750.

Hessert, J. (Ann. Chem., 176, p. 241; Neue. Repert. Pharm., 24, p. 541 (see under J. Volhard): Jahresb. Chem., 1875, p. 1302), reports 0.0024 part alumina per 1000 parts in sulphurwater from Spring Bir Keraui, after removal of the sediment which separated on standing. Including sediment there was 0.0236 alumina.

Hidegh, C. (Wiener Akad. Ber., 53 (2. Abt.), p. 395; Jahresb. Chem. Will., 1866, p. 993), reports in the spring at Johannisbad near Vienna 0.008 part alumina (and P₂O₅) per 10,000 parts.

Hillebrand, W. F. (U. S. Geol. Survey Bul. 113, p. 50), found 2.5 parts Al₂(SO₄)₃ per 1,000,000 in East Spring, and 3.2 in West Spring at Joplin, Missouri, both zinc-bearing springs.

Hillebrand, W. F. (U. S. Geol. Survey Bul. 113, p. 114), found in water from Ojo Caliente, a thermal spring near Taos, N. M., o.5 part Al₂O₃ per million parts.

Hoffmann, J. (Analysen der beiden Bohrquellen zu Homburg, Homburg, 1856; Neue Jahrb. Pharm., 7, p. 52; Chem. Centbl., 1856, p. 821; Jahresb. Chem., 1856, p. 770), notes traces of alumina in the water from two bored wells at Homburg, Germany.

Hruschauer (Ann. Chem. Pharm., 63, p. 229; Jour. Prakt. Pharm., 42, p. 466; Pharm. Centbl., 1847, p. 829; Jour. Pharm., 3d ser., 13, p. 49; Jahresb. Chem., 1847–48, p. 1002) reports 0.163 gm. basic aluminium phosphate per 10,000 gms. in Kostreiniz (Steiermark) mineral water.

Hübener (Chem. Ztg., 14, p. 1410; Chem. Centbl., 1890, II, p. 846) notes 0.002500 gm. alumina (with a trace of phosphoric acid per liter in water from a bored well in Westerland.

The well from which this water was obtained had been recently bored. The water contained a much larger proportion of iron than many of the longer-known steel-wells. Westerland is on the Island of Sylt, Prussia.

Huppert (Chem. Centbl., 1877, p. 137; Jour. Chem. Soc. London, 1878, II, p. 209) reported analyses of the water of the new spring and Josefsquelle at Bilin; they contained respectively 0.0056, and 0.0057 Al₂O₃.P₂O₅, presumably parts per 10,000. He quotes an analysis of the Josefsquelle water made in 1845 by Redtenbacher, reporting 0.084 Al₂O₃.P₂O₅.

Hunt, T. S. (Silliman's Amer. Jour., 2d ser., 8, p. 364; Jahresb. Chem., 1849, p. 621), reports 0.4681 part alumina per 1000 parts in Tuscarora Sour spring-water from the neighborhood of Brantford, Canada.

Hunt, T. S. (Silliman's Amer. Jour., 2d ser., 11, p. 174; Jahresb. Chem., 1851, p. 669), reports a trace of alumina and phosphate in water from the "outer spring" at Varennes; a trace of alumina in the "inner spring" water from Varennes; o.o14500 part alumina per 1000 parts in St. Leon and o.o0500 in Caxton water; all mineral springs in Canada.

Hunt, T. S. (Phil. Mag., 4th ser., 13, p. 239; Chem. Centbl., 1857, p. 683; Jahresb. Chem., 1857, p. 728), found a trace of alumina in Ottawa River water taken at St. Anne Lock, near Montreal, and St. Lawrence River water, taken on the south side of the Pointe-des-Cascades near Vaudreuil.

Husemann, A. (Neue Jahrb. Pharm., 33, p. 197; Chem. Centbl., 1870, p. 392; Amer. Chemist, 2d ser., 1, p. 119; Jahresb. Chem., Naumann, 1870, p. 1383), notes traces of aluminium in the Belvedra Spring near Chur.

Husemann, A. (Neue Jahrb. Pharm., 38, p. 257; Jahresb. Chem., Naumann, 1872, p. 1184), reports 0.0022, 0.0021, 0.0018, and 0.0018 part alumina per 10,000 parts in four samples of mineral water from Tarasp.

Husemann, A. (Jahrb. Pharm., 39, p. 200, 315; Jahresb. Chem., Naumann, 1873, p. 1240), reports 0.0011 part alumina per 10,000 parts in Neue Belvedraquelle, and traces in three other springs, all near Chur, Switzerland.

Husemann, A. (Arch. Pharm., 3d ser., 6, pp. 97, 395; 7, p. 204; Jahresb. Chem., 1875, p. 1291), reports respectively in the "old" and "new" St. Moritz (Switzerland) Eisen Säuerling, 0.00050 and 0.00030 part alumina per 10,000 parts. Three Tarasp (Lower Engadine) mineral springs contained per 10,000 parts respectively 0.0025, 0.0019, and 0.0021 part alumina.

Two Val Sinestra springs (Lower Engadine) contained per 10,000 parts respectively 0.0020 and 0.0021 part alumina.

Jackson, D. D., says in unpublished data regarding the natural occurrence of aluminium sulphate in waters: "Many of our driven wells along the southern shore of Long Island contain aluminium sulphate, but in no case does an entire well-plant contain it.

"Alum is noticeable in amount in individual wells at Oconee, Clear Stream, Wantagh, Merrick, and Matowa. One of the wells recently analyzed at Merrick contained 268 parts per million of aluminium sulphate. This is equivalent to 15.6 grains per gallon. In other words, the water would remove the color and impurities from fifteen to twenty times its volume of impure water.

"This analysis was made after the water had been actively drawn upon for two weeks, and at this time the taste was strong of alum. During periods of rest the alum is considerably greater in amount.

"At the same time the examination of the entire Merrick plant revealed the fact that no alum was present, but that the water was milky with aluminium hydrate which had been precipitated by the carbonate in the adjoining wells. The alkalinity of the entire supply from the plant was only 3 parts per million. This shows that the carbonate had been used up in the precipitation of the alum.

"The driven wells at Oconee gave 2.0 parts per million of Al₂O₃, and those at Clear Stream gave 1.6 parts per million. This refers to the entire plant during active operation in each case."

Janeček, G. (Chem. Centbl., 1887, p. 172; (Ausz.) Jahresb. Chem., 1887, p. 2535), reports 0.00272 gm. Al₂O₃ per 1000 gms. in the Jamnicer-alkalisch-muriatischen Säuerlings water.

John, C. v. (Verhandl. Geol. Reichsanst., 1876, p. 114; Jahresb. Chem., 1876, p. 1302), notes 0.1490 part Al₂O₃ per 10,000 parts in well-water from Ločendol, Steiermark.

John, C. v. (Jahrb. Geol. Reichsanst., 31, p. 509; Jahresb. Chem., 1881, p. 1447), reports in spring-water from Ločendol, Austria-Hungary, 0.1490 part Al₂O₃ per 10,000 parts.

John, C. v. (Chem. Centbl., 1891, II, p. 881; Jahresb.

Chem., 1891, p. 2621), found 0.0139 gm. alumina per 10,000 gms. in Friedrichsquelle water from Zeidelweid, Bohemia.

John, C. v. (Jahrb. Geol. Reichsanst. Wien, 48, 375; Chem. Centbl., 70 (1899), II, p. 1047), reported analyses of mineral waters from a number of localities in Eastern Bohemia. The data regarding Al₂O₃ follow:

ALUMINIUM IN SOME MINERAL WATERS FROM EASTERN BOHEMIA.

Kind of Water.	Dry Matter per Liter at 180° C.	Al ₂ O ₃ in Dry Matter.
	Grams.*	Grams.
Lukorna	62.4400	0.0040
Mickňovka	60.6400	.0093
avůrka	6.6430	.0023
Bučina	3.1002	.0040
Straschov	3.9840	.0400
Bohdaneč Kapelle	1.8080	.0020
Bohdaneč, St. George	3.4720	.0050

^{*} Probably grams per liter-not stated in the abstract cited.

John, C. v., and H. B. v. Foullon (Chem. Centbl., 1890, II, p. 772; Jahresb. Chem., 1890, p. 2657) report analyses of four potable springs at Luhatschowitz, namely, Vincenzbrunnen, Amandbrunnen, Johannbrunnen, and Louisenquelle. These contained respectively, 0.004, 0.005, 0.004, and 0.001 alumina (probably parts per 1000).

John, C., and C. v. Hauer (Verh. Geol. Reichsanst., 1876, p. 355; Jahresb. Chem., 1876, p. 1302) report 0.0415 part Al₂O₃ per 10,000 parts in Ranigsdorf Säuerling.

Johnstone, W. (Chem. News, 31, p. 15; Jahresb. Chem., 1875, p. 1300), reports 0.1970 gm. Al₂S₃O₁₂ per 1000 cc. in Hertfell Spring near Moffat, Scotland.

Johnstone, W. (Chem. News, 39, p. 259; Jahresb. Chem., 1879, p. 1269), reports in St. Dunstan's Well, Melrose, Scotland, 0.019503 gm. aluminium phosphate per 1000 cc.

Johnstone, J. W. (Analyst, 12, p. 90; Jour. Chem. Soc. London, 52 (1887), II, p. 1087), notes in Flitwick water, rising through a ferruginous peat-bed in Flitwick Moor, per 1000 parts, 0.0044 part alumina.

Jolles, A. (Ztschr. Nahrungsmittel u. Hyg., 1892, p. 373; Jahresb. Chem., 1892, p. 2684), found 0.064 gm. alumina per 10,000 gms. in water from the Kärtner Römerquelle in Prevali near Gutenstein.

Jones, W. Black (British Med. Jour., 1903, p. 1055), quotes from the Lancet, 1894, an analysis of Llangammarch mineral water, reporting 3.340 grain alumina and silica per gallon.

Kachler, J. (Wiener. Akad. Ber. (2. Abt.), 70, p. 654; Jahresb. Chem., 1875, p. 1296), reports respectively 0.0743 and 0.0042 part alumina per 10,000 parts in two sour springs near Poschitz, Bohemia.

Kalecsinszky, A. (Ungar. Naturw. Ber., 1, p. 370; Jahresb. Chem., 1886, p. 2321), reports 0.0052 part H₆Al₂O₆ per 10,000 parts in iron-water from Rosenau, Comitat Gömör, Hungary.

Kemper (Arch. Pharm., 2d ser., 108, p. 163; Jahresb. Chem., 1861, p. 1096) found a trace of alumina in artesian water from Gosling's Garden at Osnabrück.

Kenrick, E. B. (Geol. and Nat. Hist. Survey, Canada, Chemical Contributions, 1886, p. 13 T), reports "a very small quantity" of alumina in a qualitative analysis of water from Dougherty's so-called carbonic acid spring located in the mountains between Clinton and Carguiles, British Columbia. The total solid matter per 1000 parts in the filtered water was 1.442 parts.

In the same publication (p. 14 T) the author reports a similar analysis of water from a spring at the foot-hills of Western Butte, Sweet Grass Hills, District of Alberta, Northwest Territory. The total solids in the filtered water amounted to 0.857 part per 1000 parts. According to a qualitative analysis it contained a "very small quantity" of alumina.

Kersting (Ann. Chem. Pharm., 90, p. 158; Jour. Prakt. Chem., 63, p. 125; Pharm. Centbl., 1854, p. 589; Jahresb. Chem., 1854, p. 771), in a sulphur-spring water from Schöneck near Segewold, Russia, notes 0.0017 part alumina per 1000 parts.

Knerr, E. B. (Trans. Kansas Acad. Sci., 15 (1895-96), p. 88), notes a trace of alumina in a spring-water from Atchison, reputed to possess medicinal virtues. In the water from a well two and a half miles north of Centralia he notes 3.7 parts alumina per 1,000,000.

Kofler, L. (Vierteljahressch. Prakt. Pharm., 15, p. 161; Jahresb. Chem. Will., 1866, p. 992), reports in the Voralberg springs, Rothenbrunnen, Eisenquelle von Uebersaxen, Eisenwasser des Bad Reuthe, Eisenwasser Bad Andelsbuch, Schwefelwasser Bades Hopfreben, and Quelle zu Raggal, 0.0322, 0.031, 0.0081, 0.0476, 0.0514, and 0.0107 grain Al₂O₃ per pound respectively.

Kónya, S. (Wiener Akad. Ber., 61, 2. Abt., p. 7; Viertel-jahresb. Prakt. Pharm., 19, p. 373; Chem. Centbl., 1870, p. 132; Inst., 1870, p. 88; Jahresb. Chem., Naumann, 1870, p. 1387), notes 0.008 part alumina per 10,000 parts in the Bitterwasser from Weilutza near Jassy.

Kosmann (Jour. Pharm., 3d ser., 17, p. 43; Jour. Prakt. Chem., 50, p. 49; Pharm. Centbl., 1850, p. 141; Jahresb. Chem., 1850, p. 627) notes a trace of alumina in mineral water from Niederbronn, Germany.

Kyle, J. J. (Chem. News, 38, p. 28; Jahresb. Chem., 1878, p. 1295), reports in Rio de la Plata water taken 8 kilometers above Buenos Ayres 0.0060 part Al₂O₃ per 1000 ccm., and in the Parana water 8 kilometers above its union with the La Plata 0.0030 part Al₂O₃ per 1000 parts.

Lahache, E. (Jour. Pharm. et. Chim., 6th ser., 9 (1899), p. 477), reports "alumina, silica, etc.," as follows, in a number of potable waters from new wells, and old wells recently cleaned, in the region of Tougourt-Ouargla in the Sahara: Tougourt, o.080; Bled et Amax, o.088; El Fetir, o.043; H. Messaoud, o.035; El-Hadjira, o.068; H. Dahane, o.058; H. Debiche et Strifigi, o.034; Donionidi, o.049; N'Gonça, o.005; and Ouargla, o.008 gm. per liter.

Lane, A. C. (U. S. Geol. Survey, Water-supply and Irrigation Papers, 31, pp. 18-93), in a study of the mineral water of Lower Michigan, cites a number of analyses. The data relating to aluminium are quoted below.* In every case the values are parts per 1000 unless otherwise stated.

In Detroit River water (p. 18) S. H. Douglas (Biennial Rpt. Mich. State Geologist, 1861, p. 204) reports 0.0105 Al₂O₃.

The Dearborn Drug and Chemical Works, Chicago (p. 18),

^{*} Some of the analyses given were cited by Peale (U. S. Geol. Survey Bulletin 32, see p. 103); all such are omitted.

analyzed water from Muskegon Lake, Manistee Lake, Traverse Bay, and Detroit River, finding in the first o.ooi (FeAl)₂O₃, and in the others traces of (FeAl)₂O₃.

In Lake Superior water (p. 26) W. F. Jackman (Proc. Mich. Pharm. Assoc., 5th year, p. 110) found 0.90 part Al₂O₃ per 1,000,000 parts.

In Shiawassee River water at Owasso, the Dearborn Drug and Chemical Works (p. 27) found 0.001 (AlFe)₂O₃ and traces in Battle Creek city water, usually taken from Lake Goguac, Grand River water at Jackson, and water from the creek at Ann Arbor (Huron River?).

G. A. Kirchmaier (p. 27) (An. Rpt. Saginaw Board Water Com., 1892, Sup., pp. 31–33) reports (AlFe)₂O₃ as follows: Shiawassee River, 0.0061; Cass River, 0.0070; West Side Pumping Station, Saginaw, 0.007; East Side Pumping Station, 0.0611; and Tittabawassee River, 0.0843. J. E. Graves (p. 27) notes in Chippewa River water (Midland Water Supply) 0.0094 (AlFe)₂O₃.

A number of waters from wells and springs in the unconsolidated deposit (Pleistocene) follow (pp. 31-33). Welcome Island lithia water taken three miles north of Pontiac in the center of Lake Angelus, analyzed by J. E. Clark, 0.0014 Al₂O₃; King David Spring, Benton Harbor, analyzed by W. S. Haines, 0.0020 Al₂O₃; Nochemo Spring, Reed City, analyzed by R. Fischer and A. B. Prescott, 0.001 Al₂O₃; and well, 18 feet deep, Ypsilanti, analyzed by Dearborn Drugand Chemical Co., trace Al₂O₃.

In the wells included under the group "Miscellaneous Analyses" (p. 57) are the following: Owasso City supplywells, analyzed by Dearborn Drug and Chemical Works, trace of Al₂O₃, and Hudson Zauberwasser, analyzed by A. B. Prescott, 0.00043 Al₂O₃.

In water of the Midland Mineral Spring (p. 60), said to be 400 feet deep, 0.0247 Al₂(P₂O₅)₃ is reported by S. P. Duffield. The analysis is probably the same as that quoted by Peale (U. S. Geol. Survey Bul. 32, see p. 103).

S. P. Duffield (p. 65) reports an analysis of Butterworth's Grand Rapids Magnetic Spring, 261-274 feet deep, with 0.007 Al₂O₃.

Chilton (p. 66) (Biennial Rpt. State Geologist, Michigan, 1860, pp. 171, 186) notes 0.245 Al₂O₃+SiO₂ in an East Saginaw well 649 feet deep.

In the group entitled "Water from Devonian or Silurian Limestones" (pp. 72-85) are the following: S. P. Duffield found in wells at Clark's Riverside Bath-house, Detroit, 0.230 Al₂O₃. J. H. Long noted 0.005 Al₂O₃ in water from Excelsior Well, 1400 feet deep, Benton Harbor. In water from Coldwater Prescott notes 0.643 aluminium sulphates. In water from the St. Clair Spring, Oakland House Well, 1250 feet deep, S. P. Duffield reports 12.00 Al₂O₃, and also reports 0.421 Al₂O₃ in the Mount Clemens Original Well, which has also been known as the Avery and Soolbad. T. Tonnelé found 0.044 Al₂O₃ in the water of the Mount Clemens Clementine Well, 1060 feet deep. S. P. Duffield reports 0.601 Al,O. in the Mount Clemens Media Well. C. F. Chandler and C. E. Pellew (Geol. Survey Michigan, 5, pt. 2, p. 46) in Alma wells Nos. 1, 2, and 3, report 5.0325, 3.905, and 4.0823 grains Al₂O₃ per gallon (?), respectively.

A number of miscellaneous analyses are included in the appendix (pp. 91, 93). H. F. Northrup and G. W. Rafter (Special Water-supply Committee of the City Council, Travers City, Rpt., p. 20), note a trace of Al₂O₃ in Lake Michigan water at Chicago. C. G. Wheeler notes a trace of Al₂O₃ in water of the Mount Clemens Pagoda Spring. In water from the Ypsilanti Sanitarium J. E. Clark found 0.15 Al₂O₃.

Lang, E. E. (Verhandl. Ver. Naturkunde Presburg, 2, 1857, pt. 2, p. 1; Jahresb. Chem., 1858, p. 798), reports alumina (parts per 1000) in water from Trentschin Töplitzer Bad, Hungary, as follows: Brünnlein (Urquelle), 0.009 and Spiegelbad, No. 1, 0.017.

Langer, T. (Arch. Pharm., 3d ser., 2, p. 304; Jahresb. Chem., Naumann, 1872, p. 1186), notes traces of alumina in a mineral spring at Mattigbad, Austria.

Lambert, E. G. (Ann. Chim. et Phys., 4th ser., 12, p. 309; Jahrb. Chem. Will., 1867, p. 1043), notes in the sulphur spring of Monterey (Mexico) 0.027 gm. aluminium silicate per liter. The water is used internally and for baths. The occurrence

of aluminium is also noted in potable water, sulphur-water, etc., of Maria Island.

Laminne (Jour. Pharm., 3d ser., 13, p. 354; Jour. Chim. Med., 3d ser., 4, p. 461; Pharm. Centbl., 1848, p. 512; Jahresb. Chem., 1847, 1848, p. 1009) found 0.0020 part alumina per 1000 parts in mineral water from Tongern, Belgium.

De Launay (Ann. Mines, 9th ser., 5 (1894), p. 139) quotes an analysis of the mineral waters of Pfaefers-Ragatz, Canton St. Gall, Switzerland, made by Planta-Reichenau. In this water 0.00091 gm. aluminium phosphate per liter is reported, which was calculated to be equivalent to 0.00038 Al₂O₃.

Leber (Pharm. Centbl., 1849, p. 791; Jahresb. Chem., 1849, p. 614) found 0.1157 alumina per 1000 parts in the "new" mineral spring-water from Salzschlirf near Fulda.

Lefort (Jour. Pharm., 3d ser., 16, p. 14; Jahresb. Chem., 1849, p. 617) notes 0.017 gm. aluminium silicate per 1000 cc. in Enclos des Célestins mineral water from Vichy.

Lefort, J. (Jour. Pharm., 3d ser, 21, p. 340; Jahresb. Chem., 1852, p. 757), notes 0.009, 0.008, and 0.005 gm. alumina per 1000 gms. in the "right," "left," and "middle" mineral springs, respectively, at Jenzat, Dept. Allier.

Lefort, J. (Jour. Pharm., 3d ser., 31, p. 84; Jahresb. Chem., 1857, p. 725), found traces of alumina in the following mineral waters from Royat and Chamalières, Dept. Puy-de-Dôme, France; Royat Spring, Cäsarbad, Saint-Mart, and Roches (Beaurepaire) Spring.

Lefort, J. (Jour. Pharm., 3d ser., 32, p. 264; Jahresb. Chem., 1857, p. 726), notes a trace of alumina in mineral water from Neyrac, Dept. d'Ardèche, France.

Legrip (Jour. Chim. Med., 3d ser., 4, p. 83; Pharm. Centbl., 1848, p. 431; Jahresb. Chem., 1847–48, p. 1006) found 0.0074 gm. alumina per 1000 cc. in mineral water from Doulaux, Dept. Creuse.

Le Grip (Jour. Chim. Med., 3d ser., 5, p. 514; Jahresb. Chem., 1849, p. 618) notes 1.75 per cent alumina in total solids of mineral water from Chaumaix, Dept. Creuse, the total solids being equal to 5.80 gms. per 10,000 cc.

Leverett, F. (U. S. Geol. Survey Rpt., 17 (1895-96), pt. II,

pp. 701-828), in an article on the water resources of Illinois, gives a number of analyses of water from springs, shallow wells, artesian wells, etc., a number of which, it is stated, were quoted from D. W. Mead's "Hydrography of Illinois." Some of these have been noted from Peale's compilation (U. S. Geol. Survey Bul. 32); others follow. G. H. Ellis (p. 824) reported 0.13 gm. Al₂O₃ per U. S. gallon in Bushnell Railroad Well; W. S. Haines (p. 824) notes 0.08 grains per U. S. gallon in Woodstock "drift" well.

In "St. Peter's" water from Macomb, Ill., G. Steiger (p. 826) reports 0.0013 gm. Al₂O₃ per 1000 cc.

The alumina (grains per U. S. gallon) found in a number of artesian-well waters by different analysts (pp. 827, 828) follows:

ALTIMINA	IN CERTAIN	ARTESIAN	WATERS

Artesian Wells.	Analyst.	Al ₂ O ₃ per U. S. Gal.
Auditorium, Chicago, Ill. Munger's laundry, Chicago, Ill. Davenport, Iowa, glucose factory. Dekalb, Ill., water-works Dixon, Ill., water-works. Galena, Ill., water-works Geneseo, Ill., water-works Jerseyville, Ill., water-works Lagrange, Mo., Wyaconda well. Macomb, Ill., water-works Monmouth, Ill., water-works Montezuma, Ind. Peru, Ill., water-works Rockford, Ill. Rockford, Ill., water-works Sterling, Ill.	E. G. Smith E. G. Smith E. G. Smith E. Guteman G. M. Davidson E. G. Smith W. Simpson D. M. Stanner E. G. Smith ? G. Steiger E. G. Smith W. A. Noyes E. G. Smith	Grams07 .03 .36 .69 .12 .06 8.55 .06 .09 .075 .10 .07 .04 .13 .06 .05

In several others a trace of alumina is noted.

Lengyel, B. v. (Földtani Közlöny, 23 (1893), p. 293; Jahrb. Min., 1895, I, Ref. 66; Jour. Chem. Soc. London, 68 (1895), II, p. 118), finds that the Kolop Sulphur Spring, situated near Tisza Süly in the great Hungarian plain, contains o.o1o1 gm. Al₂O₃ per kilogram.

Lepsius, B. (Ber. Deut. Chem. Gesellsch., 21, p. 552; Jour. Chem. Soc. London, 54 (1888), p. 435), notes 0.00017 gm. Al₂(PO₄)₃ per liter in the water of the Tönnissteiner medicinal

spring and quotes an earlier analysis by Fresenius, reporting 0.00013 gm.

Lewy (Compt. Rend. Acad. Sci. Paris, 24, p. 449; Jahresb. Chem., 1847–48, p. 1011) reports 1.66 aluminium sulphate per 1000 parts in thermal-spring water from Paramo de Ruiz, New Granada. See also Boussingault (Compt. Rend. Acad. Sci. Paris, 24, p. 397; Ann. Chim. Phys., 3. ser., 20, p. 109; Iour. Pharm., 3. ser., 11, p. 487; Jour. Prakt. Chem., 40, p. 438; Ann. Chem. Pharm., 64, p. 292; Poggendorf's Ann., 71, p. 444; Pharm. Centbl., 1847, p. 414).

L'Hôte, L. (Chem. Centbl., 1891, I, p. 207; Jahresb. Chem., 1891, p. 2620), found 0.0012 gm. alumina in the 2.216 gms. total solids obtained from 1000 cc. of mineral water from Penon de los Banos, Mexico.

Liebig (Ann. Chem. Pharm., 63, p. 127; Jour. Prakt. Chem., 42, p. 463; Pharm. Centbl., 1847, p. 828; Jour. Pharm., 3d ser., 13, p. 65; Jahresb. Chem., 1847–48, p. 1002) notes a trace of aluminium oxid in bitter water from Friedrichshall at Hildburghausen.

Liebig (Ann. Chem. Pharm., 79, p. 94; Pharm. Centbl., 1851, p. 916; Vierteljahressch. Prakt. Pharm., 1, p. 218; Jour. Pharm., 3d ser., 20, p. 315; Jahresb. Chem., 1851, p. 650) examined the sulphur waters of Aachen, reporting traces of aluminium phosphate in Kaiserquelle, Corneliusquelle, Rosenquelle, and Quirinusquelle.

Liebig (Ann. Chem. Pharm., 98, p. 145; Vierteljahressch. Prakt. Pharm., 5, p. 547; Jour. Prakt. Chem., 69, p. 28; Chem. Centbl., 1856, p. 390; Jahresb. Chem., 1856, p. 765) notes the occurrences of traces of aluminium phosphate in the following Kissingen mineral waters: Racoczy, Pandur, and Maxbrunnen.

Liebig (Ann. Chem. Pharm., 98, p. 350; Jour. Prakt. Chem., 69, p. 331; Chem. Centbl., 1856, p. 350; Jahresb. Chem., 1856, p. 766) notes the occurrence of traces of aluminium phosphate in the following mineral waters: Bitter water from Mergentheim, Bonifaciusquelle, Marienquelle, Elizabethenquelle, and Hermannsquelle, the last four at Neuhaus near Neustadt, Germany.

Limouzin-Lamothe (Jour. Chim. Med., 3d ser., 9, pp. 716, 763; Jahresb. Chem., 1853, p. 716) in mineral water from the

Dept. Aveyron, viz., from Prugnes and from Cayla (Princess Spring, Magdalen Spring, and Rose Spring), found alumina as follows: 0.035, 0.050, 0.055, and 0.050 gm. per 1000 cc. respectively.

Liversidge, A. (Chem. News, 42, p. 324; Jahresb. Chem., 1880, p. 1536; Jour. Chem. Soc., London, 40 (1881), p. 564), notes 12.86 parts aluminium chlorid per 100,000 in addition to 4.17 alumina with a trace of ferric oxid in a spring on Kantavu, one of the Fiji Islands.

Liversidge, A. (Chem. News, 62, p. 264; Jour. Chem. Soc. London, 60 (1891), p. 280), in water from a hot spring in Savo Island, notes the occurrence of aluminium.

Liversidge, A. W. Skey, and G. Gray (Rpt. Assoc. Adv. Sci. Australasia, 7 (1898), p. 87) have collected analyses of the mineral waters of Australasia. The data regarding aluminium are given on page 94.

In addition to the above traces of alumina are reported by W. Skey (p. 105) in eighteen Te Aroha mineral waters, Queensland, Aluminium and iron are reported together in a number of other waters.

Lord (Geol. Survey Ohio, 6(1888); U. S. Geol. Survey Rpt. 8, pt. II, p. 621) analyzed a brine from a well drilled at Lorain, Lorain County, Ohio, which contained 0.040 part alumina per 1000 parts.

Losanitsch, S. M. (Ber. Deutsch. Chem. Gesellsch. 20, p. 1114; Jour. Chem. Soc. London, 52 (1887), p. 648), reports in mineral waters from Servia Al₂O₃ (parts per 1000) as follows: Wrnjačka banja, 0.00105; Bukowik, 0.00271; Palanka, 0.00199; Alexinačka banja, 0.00279; Ribarska banja, 0.0010; Wranjska banja, 0.0005, and Brestowačka banja, 0.00140.

Löwe, J. (Jahresb. Phys. Ver. Frankfurt a. M., 1853-54, p. 55; Jahresb. Chem., 1854, p. 759), notes in the Kronthal, Germany, Salzquelle, 0.000550 part aluminium silicate per 1000 parts.

Löwe, J. (Jahresb. Phys. Ver. Frankfurt a. M., 1854-55, p. 58; Jahresb. Chem., 1856, p. 770), notes 0.001269 gm. aluminium silicate per 1000 gms. in the Kronthal, Nassau, Germany, Stahlquelle water.

ALUMINIUM IN AUSTRALASIAN MINERAL WATERS.

	Analyst.	Page.	Alu- minium Sul- phate per Gallon.	Alu- mina per Gallon.	Alu- minium Phos- phate per Gallon.	Alu- minium Chlo- ride per Gallon.	Alu- minium Silicate per Gallon.
			Grains.	Grains.	Grains.	Grains.	Grains.
New Zealand.							
Alkaline water, Auck-							
land	W. Skey	88		.30			
Springs Hanmer,			1				
Plains, Nelson	J. v. Haast	89			trace		
Chlorinated water, Helensville	W. Skey	80		. 45	1		
Chlorinated water,		09		.43			
Kopuowhara Mahia	W. Skey	90			-37	16.42	
Acid alum water, Mo- tuhora (Whale Isl-							
and, Bay of Plenty .	J. A. Pond	90	48.48				
Acid aluminous water,			45.45				
Ohaeawai, Auckland	J. Hector	92	a				
Alum water Onetapu, Desert Auckland	W. Skey	02	ь				
Otira Gorge, sulphur-		,-					
etted and siliceous	C C	1					
Pahua Wairarapa,	G. Gray	92		. 21			
chlorinated water	W. Skey	93			.64		
Chlorinated water,	TIT CL						
Tologo Bay Chlorinated water, No.	W. Skey	94		2.150			
I, Waimate Block.	W. Skey	95		trace			
Chlorinated water, No.	W C1			- 06			
2, Waimate Block Chlorinated water, No.	W. Skey	95		2.86			
3. Waimate Block	W. Skey	95		trace			
Weak chalybeate Wai-			The same of				
rongoa, North Taiera Otoga	A. G. Kidson- Hunter	95		. 25			
Alkaline saline water, Waiwera, Auckland.	W. Skey	96		trace			
Chlorinated water, A,		90					
Wanganui	W. Skey	96				.91	
Wanganui	W. Skey	96				1.22	
Chlorinated water, C							
Wanganui	W. Skey	96				.21	The state of
water, Whangarei	J. A. Pond	97			traces		
Acid water, Lake							
White Island or Whakaari	T Wester	1	0 -				
	J. Hector	97	80.5			1703.1d	
New South Wales.						1	-
Alkaline water, Balli-						1	
more	J. C. H. Mingaye	98		trace			
Water Milparinka	A. H. Jackson	98		little			
Queensland.		1					
Te Pupunitanga							
Priets's Bath, Ro- turua District	W. Skey	100	21.67			- 10	
	oney	100		Same and			
							-

a. Temperature 60°-116° F. Deposits alum and sulphur (amount not stated) on cooling. Total solids, 134.4 grains per gallon.

b. Total solids, 456 grains per gallon, mainly potash, alum and magnesia, and ferrous chlorides.

c. With trace of iron.

d. Sesquichloride.

ALUMINIUM ?	N AUSTRALASIA	AN MINERAL	WATERS Continued.
-------------	---------------	------------	-------------------

	Analyst.	Page.	Alu- minium Sul- phate per Gallon.	Alu- mina per Gallon.	Alu- minium Phos- phate per Gallon.	Alu- minium Chlo- ride per Gallon.	Alu- minium Silicate per Gallon.
	The state of the s		Grains.	Grains.	Grains.	Grains.	Grains.
Waikupapapa Saddler's Bath, Roturua Dis- trict	W. Skey	100	.32	Grains.	Grains.	Grains.	Grains.
Bath, Roturua Dis- trict	W. Skey	100	33.22				
trict	W. Skey	100	1.34				
turua District Te Mimi Okakahi, Ro-	W. Skey	100	.68				
turua District Ti Kute Great Spring.	W. Skey	100	trace				
Roturua District Sulphur Bay Spring,	W. Skey	100	11.22				11(4), 21
Rotorua District Perekari, Rotorua	W. Skey	100	trace				
District	W. Skey	100	trace		8488		a grade
South Australia.							
Wyly's Well Farina	G. A. Goyder	107					.91
Spring, The Peake	G. A. Goyder	107					. 26
Spring, Cadna-owie Spring, Andrawilla,	G. A. Goyder	107					.31
Eleanor River Healy Springs, Indul-	G. A. Goyder	107	• • • • • • • • • • • • • • • • • • • •				. 42
kana Springs Oolarinna Soakage	G. A. Goyder	107					.10
Well	G. A. Goyder	107					. 25
Strangways Sulphur Spring, Strang-	G. A. Goyder	107					-45
ways	G. A. Goyder	107					.75
Nilpinna Spring	G. A. Goyder	107					. 32
Weedina Spring	G. A. Goyder	107					.15
Lake Harry Bore	G. A. Goyder	107					.30
Spring Alton Downs,					1/1/11		1 5
Herbert River, North Territory	G. A. Goyder	107					.68

Ludwig, E. (Wiener Akad. Ber., 52, 2. Abt., p. 264; Jahresber. Chem. Will., 1865, p. 934), notes 0.005 aluminium phosphate per 10,000 parts in the thermal spring at Tobelbad, near Gratz, in Steiermark.

Ludwig, E. (Wiener. Akad. Ber., 50, 2. Abt., p. 247; Chem. Centbl., 1865, p. 702; Jahresber. Chem. Will., 1865, p. 935), notes traces of aluminium phosphate in three mineral springs at Mähren, viz., Johannisbrunnen, Neue Quelle, and Paulaquelle.

Ludwig, E. (Min. Petr. Mittheil., 2d ser., 4, p. 519; Jahresb. Chem., 1881, p. 1447), found 0.0053 Al₂O₃ per 10,000 parts in the Sauerquelle from Apatovac, Austria-Hungary.

Ludwig, E. (Min. Petr. Mittheil. 6. p. 150; Jahrseb. Chem., 1884, p. 2035), notes 0.0010 Al₂O₃ per 10,000 parts in Maria Theresia Quelle at Andersdorf, Austria-Hungary.

Ludwig, E. (Die Mineralquellen Bosniens, abs. Chem. Centbl., 1889, II, p. 264; Jahresb. Chem., 1889, p. 2633), reports in the following mineral waters from Bosnia: (1) Thermal spring at Ilidze near Sarajewo; (2) Säuerling from Kiseljack; (3a) Säuerling from Bistrica; (3c) Säuerling from Ljeskovica (Giftquelle); (4) Rjecicaquelle at Maglaj; (6) Säuerling at Dragunje; (7) Säuerling at Kiseljack near D. Tuzla; and (8) iodine spring at Navioci near Han Sibosica; Al₂O₃ as follows: 0.012, 0.002, 0.006, 0.004, 0.0017, 0.005, 0.001, and 0.021 (probably parts per 10,000).

Ludwig, E. (Tschermak's Mineral. Mittheil. 1890, Jahresb. Chem., 1890, p. 2662), reports 1.319 aluminium sulphate in Cervena-Rjeka-Quelle and 0.146 aluminium sulphate in the Quelle an der Strasse (presumably parts per 10,000 cc.). Both these "vitriol" springs are at Srebrenica, Bosnia. He also reports (Tschermak's Mineral. Mittheil., 1890; Jahresb. Chem., 1890, p. 2663) 13.468 gms. aluminium sulphate per 10,000 cc. in a mineral spring at Büdös (Bálványos) in Siebenbürgen, 0.007 gm. alumina per 10,000 cc. in Karlsquelle; 0.0025 gm. aluminium per 10,000 cc. in Fidelisquelle; and 0.961 part aluminium sulphate per 10,000 parts in the alum spring, all on the Büdös (Stinkberg).

Ludwig, E. (Die Mineralquellen Bosniens, abs. Chem. Centbl., 1890, II, pp. 468. 846; Jahresb. Chem., 1890, p. 2666), reports analyses of mineral-spring waters from Bosnia. The following waters contained alumina (grams per 10,000 in every case): No. 1a, saline, Dolnj-Tuzla, 0.005; No. 1b, saline, Dolnj-Tuzla, 0.009; No. 2, thermal spring at Gradačac, 0.007; No. 5, sulphur spring, Raso, 0.001; No. 6, Säuerling at Jasenica, 0.006; No. 7, Säuerling at Dubnica, 0.002; No. 8, Säuerling at Tesanj, 0.019; No. 9, thermal spring, Vručica, 0.008; No. 10, thermal spring, Kulasi, 0.007; No. 11, sulphur spring, Smodelac, 0.003; No. 12, thermal spring, Gorni Sehar, 0.001; No. 13, thermal spring, Slatina Ilidze, 0.005; No. 14, Säuerling spring, Slatina Ilidze, 0.005; No. 15, thermal spring at Gata, 0.002;

No. 16, thermal spring at Fojnica, o.oo1; and No. 17, thermal spring at Banja Visegrad, o.oo1.

Other springs contained aluminium sulphate as follows (grams per 10,000): Waters from Srebrenica (Haldenwasser), No. 4a, arsenic vitriol Črni Guber, 2.227; No. 4b, arsenic vitriol Mala Kiselica, 1.482; and No. 4c, arsenic vitriol Velika Kiselica, 0.384.

Ludwig, E. (Tsch. Min. Mittheil., 16 (1896), p. 133; Jour. Chem. Soc. London, 72 (1897), II, p. 110), notes in a carbonated water from a spring at Seifersdorf, Austrian Silesia, 0.001 parts Al₂O₃ per 10,000, and also notes (Tsch. Min. Mittheil., 16 (1896), p. 140; Jour. Chem. Soc. London, 72 (1897), II, p. 110) 0.002 part Al₂O₃ per 10,000 parts in water from the Constantine Spring in Gleichenberg, Styria.

Ludwig, E. (Wien. Klin. Wochensch., 10 (1897), p. 56; Chem. Centbl., 1897, I, p. 718; Jour. Chem. Soc. London, 74 (1898), II, p. 392), reports 0.004 Al₂O₃ parts per 10,000 in water from the iodine spring at Wels, Upper Austria.

Ludwig, E., and V. Ludwig (Wien. Klin. Wochensch., 11 (1898), 207) studied the composition of the acid iron springs at Johannisbrunn in Silicia, reporting per 10,000 parts by weight 0.001 part alumina in Marienquelle and 0.002 part in Paulaquelle.

Ludwig, E., and J. Mauthner (Min. Petr. Mittheil., 2d ser., 2, p. 269; Jahresb. Chem., 1880, p. 1524) report analyses of the Carlsbad thermal springs. These contained Al₂O₃ per 10,000 cc. as follows: Sprudel, 0.004 gm.; Marktbrunnen, 0.007 gm.; Schlossbrunnen, 0.005 gm.; Mühlbrunnen, 0.005 gm.; Neubrunnen, 0.006 gm.; Theresienbrunnen, 0.005 gm.; Elizabethquelle, 0.006 gm.; Felsenquelle, 0.003 gm.; Kaiserbrunnen, 0.005 gm. The article (loc. cit., p. 1526) contains a reference to Lerch's analyses made in 1847. He reported 0.005 gm. Al₂O₃ per 10,000 cc. in Orchesterquelle, and 0.006 gm. in Elizabethquelle.

Ludwig, E., and T. Panzer (Wiener Klin. Wochensch., 13 (1900), p. 617) note a trace of alumina in the water of the Gastein, Austria, Hauptquelle or Elizabethquelle. They also quote data obtained by earlier analysts. Thus Hünefeld reported 0.016 Al₂O₃ in 1828; Saltmann, 0.027 in 1837;

Wolff, 0.022 in 1845; and Ullik, 0.003 in 1845, parts per 10,000 in every case.

Ludwig, H. (Arch. Pharm., 2d ser., 93 (1858), pp. 129, 257; Jahresb. Chem., 1858, p. 795), notes 0.0054 Al₂O₃.SiO₃ per 1000 parts in the cold sulphur-water from Tennstädt, Thuringia.

Ludwig, H. (Arch. Pharm., 2. ser., 133 (1868), p. 1; Jahresb. Chem. Will., 1868, p. 1037), reports per 1000 cc. 0.0033 Al₂O₃, 2SiO₂ in Friedensquelle from Rostenberg in Sachsen-Weimer-Eisenach.

Lunge, G., and L. Landolt (Sep. from Corresp. Schweizer-Aerzt.; Jahresber. Chem., 1885, p. 2317) report 0.0259 gm. Al₂Cl₆ per 1000 gms. in the Schweizerhalle saline spring at Basel.

Lunge, G., and R. E. Schmidt (Zeit. Analyt. Chem., 25, p. 309; Jour. Chem. Soc. London, 50 (1886), p. 996), in water of the St. Lorenz hot spring at Leuk, note per kilogram 0.00051 gm. alumina.

Magerstein, V. T. (Verhandl. Geol. Reichsanst., 1879, p. 191; Jahresb. Chem., 1879, p. 1265), reports a trace of Al₂O₃ in two mineral waters from Zuckmantel and Einsiedel, Austria.

Maier, P. J. (Naturk. Tijdschrift. Nederland Indië, 21, p. 1; 22, p. 44; 23, pp. 46, 378; Jahresb. Chem, 1861, p. 1113), found Al₂O₃ (grams per 100) in mineral water from Dutch East India as follows: Spring at Koeningan near the river Tjisangarong, 0.01730; Sissipan Spring, near the river Tjisangarong, 0.01050; spring northeast from the Palimang mountain, 0.00039; Prajan Spring at Res. Madioen, trace; hot spring at the foot of Kaba mountain, 0.00059; and thermal spring on Pengalengen plateau, 0.00038.

Mangini, F. (Gaz. Chim. Ital., 17 (1887). p. 517; Jour. Chem. Soc. London, 54 (1888), p. 1261), in an analysis of the chalybeate water of Raffanelo in the commune of Canale Monterano, reports 0.1350 gm. Al₂O₃ per liter.

Marchand, E., and E. Leudet (Jour. Pharm., 37, p. 328; Jahresb. Chem., 1860, p. 838) report aluminium salts as follows (grams per 1000 gm.) in mineral water from Blèville, France: 0.0031 Al₂O₃3SO₃; 0.0151 Al₂O₃.NH₄O.4SO₃: 0.0145 Al₂O₃.KO.4SO₃; 0.0022 Al₂O₃PO₅; and 0.0493 CaO.Al₂O₃.4SiO₃.

Mason, W. P. (Chem. News, 61, p. 123; Jour. Chem. Soc., London, 58 (1890), p. 854), reports an analysis of the water of Excelsior Springs near Kansas City, Missouri, which showed 2.10 parts Al₂O₃ per 1,000,000.

Maumené (Compt. Rend. Acad. Sci., Paris, 31, p. 270; Pharm. Centbl., 1850, p. 754; Jahresb. Chem., 1850, p. 626) notes 0.00112, 0.00119, 0.00140, and 0.00197 gm. alumina per 1000 cc., respectively, in water from the Vesle taken on January 23, 1849, at Château-d'Eau; Vesle water taken at the same place June 18, 1849; Vesle water taken June 18, 1849, at Saint-Brice, and water from the Suippe.

Meyer, R. (Ber. Deut. Chem. Gesell., 11 (1878), p. 1521; Jour. Chem. Soc. London, 36 (1879), II, p. 33), in water from the mineral spring at Tenninger Bad, Somvirex Tobel, Grisons, notes 0.0008 part Al₂O₃ and H₃PO₄ per 10,000.

Mingaye, J. C. H. (Jour. Proc. Roy. Soc. New South Wales, 26 (1892), p. 73), reports a number of analyses in a study of some of the well, spring, mineral, and artesian waters of New South Wales and their probable value for irrigation. A potable water from Yarrangobilly caves and one from Jenolan caves each contained a trace of alumina. Other waters contained alumina (parts per 1000) as follows: Water from Jenkins's quarry, 0.0082; from Kidwell's quarry, 0.0104; from Marden's quarry, 0.0126; from Walsh's quarry, 0.0096; water from brewery, Broken Hill, o.0044; Silverthorne's well, Broken Hill, 0.0030; water from Portion 86, taken from a bore in a limestone quarry, Broken Hill, o.0004; Wilcannia, water from Tarella, 0.0064; water from Barragan, near Mudgee, trace; Belabula River, Clifden Run, trace; artesian water from Cuttabura, Bourke private bore, and Bourke Corrella Station, trace; and mineral water from Jarvisville, near Picton, Ballinore, Tarbragar River, near Dubbo (artesian), and Rock-Flat Spring, near Cooma, Monara District, each a trace. In one or two other cases alumina with iron is noted.

Mingaye, J. C. (Rpt. Australian Assoc. Adv. Science, 6 (1895), p. 265), in a second paper, which deals with analyses of the artesian waters of New South Wales and their value for irrigation and other purposes, reports alumina as follows (parts per thousand): Kelly's Camp bore, 0.0028 (with a trace of

Fe₂O₃), water from Barringun bore, 0.0036 (with a trace of Fe₂O₃); Corella bore No. 1 (private), trace; Waroo Springs bore, 0.0036 (with a trace of Fe₂O₃); Cuttaburra bore and Dungle Ridge bore, each a trace. In several other waters alumina and ferric oxid are reported together.

Miteregger, J. (Jahrb. Naturhist. Landesmuseums Kärnthen, 1861, No. 5, p. 1; Jahresb. Chem., 1861, p. 1099), notes aluminium (parts per 10,000 cc.) as follows, in water from Radlbad, near Gmünd, 0.820; Säuerling Bades Vellach in Vellathale Hauptquelle No. IV, 0.130; Säuerling Bades Vellach in Vellathale Quelle No. II, 0.189; Säuerling Bades Vellach in Vellathale Quelle No. III, 0.080; Säuerling, Bades Vellach in Vellathale Quelle No. V, 0.100, and Säuerling from Ebriach, near Eisenkappel, 0.130.

Mitteregger, J. (Jahrb. Naturhist. Landesmuseums Kärnthen, 1862, p. 109; Jahresb. Chem., 1862, p. 813), notes alumina per 10,000 gms. in Kärnthen medicinal waters as follows: Katharien-Bad at Klein-Kirchheim, 0.050; Preblauer Sauerbrunnen, 0.040; Sauerbrunnen at Weissenbach in the Lavantthale, 0.140; and Kleininger Sauerbrunnen in the Lavantthale, 0.240.

Moissenet (Ann. Min., 5th ser., 17, p. 7; Jahresb. Chem., 1860, p. 839) reports Al₂O₃ (grams per 1000 cc.) in French mineral waters as follows: Water from Sylvanès, Dept. Aveyron, Source des Moines, 0.0300; Source des Petites Baignoires, 0.0750; Source des Petites-Eaux, 0.0200; Source des Bains Nouveaux, 0.0200; water from Plombières, Sources des Dames, 0.0075, and Source ferrugineuse, 0.1433; and a trace of Al₂O₃ in water from Vittel, near Contrexeville, Dept. Vosges, Source Marie, Source des Demoiselles, and Grand Source.

Moitessier, A. (Compt. Rend. Acad. Sci. Paris, 51. p. 636; Jahresb. Chem., 1860, p. 839), notes 0.0030 gm. alumina per 1000 in the water of the warm spring about 2 km. north of Montpellier, France.

Morelli (Chem. Ztg., 1881, p. 815; Jahresb. Chem., 1881, p. 1448) found 0.0299 part aluminium sulphate per 1000 in one spring and 0.0326 part aluminium sulphate in another at Ceresole Reale, Italy.

Morin, P. (Jour. Pharm., 3d ser., 21, p. 5: Arch. Phil. Nat.,

18, p. 224; Jahresb. Chem., 1851, p. 667), found 0.0162 gm. aluminium silicate per 1000 gms. in mineral water from Coëse (Savoy).

Morin, P. (Jour. Pharm., 3d ser., 40, p. 183; Jahresb. Chem., 1861, p. 1106), found 0.0027 gm. alumina per 1000 gms. in mineral water from the Guillot spring at Evian, Savov.

Muck, F. (Jour. Prakt. Chem., 96, p. 459; Chem. Centbl., 1866, p. 384; Jahresb. Chem., 1865, p. 931), reports in 1000 parts Natronsäuerling from Nassau 0.00060 gram alumina.

Mügge, O. (Jahrb. Min., Beilageband, 4, p 576: Jahresb. Chem., 1886, p. 2324), reports an analysis by C. Pieper of a hot spring south of Naiwaschasee, Massailand, East Africa. This contained 0.779 gm. Al₂S₃O₁₂ per 1000 cc.

Müller, E. (Arch. Pharm., 2d ser., 186 (1868), p. 16; Jahresb. Chem., 1868, p. 1038), reports per 500 gms. 0.0027 Al₂O₃PO₅ and 0.0150 Al₂O₃.3HO in the water of the sulphur spring at Seebruch.

Müller, G. (Neue. Jahrb. Pharm., 3, p. 205; Pharm. Centbl., 1855, p. 526; Jahresb. Chem., 1855, p. 840), reports 0.0340 alumina per 1000 gms. in the Bergstrasse Stahlquelle water from Weinheim.

Müller, G. (Wiener. Akad. Ber., 2. Abt., 58, p. 101; Inst., 1868, p. 384; Jahresb. Chem., 1868, p. 1035), reports traces of Al₂O₃ in the Ida spring at Biloves in Bohemia.

Muspratt, J. S. (Pharm. Jour. Trans., 11, p. 151; Jahresb. Chem., 1851, p. 653), notes traces of alumina in the water of the principal spring at Baden-Baden.

Nasini, R., and F. Anderlini (Gaz. Chim. Ital., 24 (1894), I, p. 327; Jour. Chem. Soc., London, 66 (1894), II, p. 422) report 0.0015 Al₂O₃ part per 10,000 by weight in the water of Hot Springs at Monte Irone, Albane.

Netwald (Unters. des Mineralwasser zu Hall bei Kremsmünster, Linz, 1853; Jahresb. Chem., 1853, p. 711) notes in mineral water from Hall near Kremsmünster, Austria, 0.0038 gm. aluminium silicate per 1000 gms.

Northcote, A. B. (Phil. Mag., 4th ser., 14, p. 457; Jahresb. Chem., 1857, p. 727), notes a trace of alumina in the following

saline waters from Cheshire, England: Anderton, Marston, Winsford, and Wheelock.

Nuricsan, J. (Földtani Közlöny, 23 (1893), p. 296; Jahrb. Min., 1891, I, Ref. 67; Jour. Chem. Soc. London, 68 (1895), II, p. 118), notes 0.0017 gm. Al₂O₃ per kilogram in the water of the Roman brine spring and 0.0064 in the Shaft spring at Torda, Hungary.

Nuricsany and Spängler (Wiener Acad. Ber., 14, p. 121; Pharm. Centbl., 1855, p. 78; Jahresb. Chem., 1854, p. 771) note in mineral water from Roggendorff (Hungary) 0.019 gm. alumina per 1000 gms.

Ochsenius, C. (Ztschr. Geol. Ges., 34, p. 357; Jahresb. Chem., 1882, p. 1626), notes traces of Al₂O₃ in water from the Great Salt Lake, Utah.

Oppermann (Mém. Mus. Strassburg, 4, 1853, p. 21; Jahrb. Min., 1854, p. 184; Jahresb. Chem., 1853, p. 717) notes 0.006250 gm. alumina per 1000 cc. in mineral water from Sulzbach, Alsace.

Orlow, N. A. (Farmaz. Jour., 40, 1901, pp. 2, 26; Ztschr. Untersuch. Nahr. u. Genussmtl., 5 (1902), p. 133), reports 0.0022 gm. Al₂O₃ per 1000 cc. in the Director spring water from Staraja-Russa.

Orlowsky, A. (Ber. Deut. Chem. Gesell., 1883, p. 978 (corresp.); Jahresb. Chem., 1883, p. 1947), notes 0.001904 part Al₂O₃ per 1000 parts in Bolschoi water (used for baths) and 0.001784 part Al₂O₃ per 1000 parts in Kasimer water (used for drinking), both springs in Slawinsk, Poland.

Orosi (Gaz. Toscana, 1847, p. 99; Pharm. Centbl., 1847, p. 413; Jour. Prakt. Chem., 42, p. 468; Jahresb. Chem., 1847–48, p. 1009) found 0.0049 part alumina per 1000 in Mofetta di S. Quirico, water from Tuscany.

Osnaghi (Wien. Acad. Ber., 17, p. 443; Chem. Centbl., 1856, p. 40; Jahresb. Chem., 1855, p. 842) notes 0.010 part alumina per 1000 parts in mineral water from Galdhof, near Seelowitz, Austria.

Otto, R., and H. Beckurts (Arch. Pharm., 3d ser., 18, p. 115; Jahresb. Chem., 1881, p. 1444) report 0.0030000 part Al₂O₃ per 1000 parts in Hroswitha water and 0.0035000 part Al₂O₃ in

Wilhelmsquelle; both waters from Herzogludolfsbad, near Gandersheim, Germany.

Parmentier, F. (Compt. Rend. Acad. Sci. Paris, 115, p. 125; Jahresb. Chem., 1892, p. 2687), writing of the occurrence of alumina in mineral water, reports the following in springs at Vichy, Saint-Yorre (Allier) Frobert, 0.008; Saint-Louis No. 1, 0.007; Précieuse, 0.006; Jeanne d'Arc, 0.004; Sévigné, 0.003. In Vichy (Allier) springs: Dubois, 0.015; Vincent, 0.010. In springs at Hauterive (Allier): Bayard, 0.006; Amélie d'Hauterève, 0.001. In Pougues-les-Eaux (Nièvre) springs: Grande Source, 0.014; Jeanne d'Arc, 0.006; Saint-Léon, 0.002; and in the Chatelguyon (Puy-de-Dôme) spring, Yvonne, 0.009 gm. per 1000 cc.

Parmentier, F. (Compt. Rend. Acad. Sci. Paris, 132, 1901, p. 1332; Ztschr. Untersuch. Nahr. u. Genussmtl., 5 (1902), p. 138), states that aluminium occurs in most of the springwaters previously analyzed by him (for instance, Vichy), although it was not reported.

In Puits Chomel he reports 0.0117 gm. per 1000 cc., and in Grande Grille he reports 0.0075 gm. per 1000 cc. These waters contained flocculent particles of aluminium hydrate.

Parmentier, F. (Compt. Rend. Acad. Sci. Paris, 121 (1895), p. 644; Jour. Chem. Soc. London, 70 (1896), II, p. 195), in a bituminous water from the Grassion spring at Clermont, France, found 0.004 gm. Al₂O₃ per liter.

Passy, A. (Compt. Rend. Acad. Sci. Paris, 68, p. 171; Jahresb. Chem., 1869, p. 1290), notes 0.002 and 0.010 gm. Al₂O₃ per liter, respectively, in spring-water and Aube water from Etufs, France. The analyses were made by H. Magnon.

Paternò, E., and G. Mazzara (Gaz. Chim. Ital., 9 (1879), p. 71; Jahresb. Chem., 1879, p. 1266) note a trace of Al₂O₃ in Termini-Imerese water from Palermo, Sicily.

Paternò, E. (Gaz. Chim. Ital., 21 (1891), II, p. 40; Jahresb. Chem., 1891, p. 2619), notes traces of aluminium phosphate in thermal-spring water from Sclafani.

Peale, A. C., of the United States Geological Survey (U. S. Geol. Survey Bulletin No. 32, p. 235), published some years ago a very extended compilation of analyses of waters of

mineral springs in the United States. The author states that the information was derived primarily from various State Geological Reports, State guide-books and handbooks, government geological reports, etc., and various other scientific publications. Much additional material was obtained from members of the Geological Survey, and in answer to a special set of inquiries sent throughout the United States. Efforts were made by the compiler to verify in each instance the matter presented in regard to the various localities by direct correspondence. Some eight thousand springs are included in the list, of which about eight hundred had been analyzed. The tables, with a few exceptions, do not include brine springs and wells which are used in the production and manufacture of salt, since such waters are not applied to the ordinary uses of mineral springs. The results are arranged by States and Territories, and the analytical data are in the main incorporated in the tables as given in the sources from which they were taken, except that where originally expressed in grains per fraction of a gallon they were recalculated to grains per gallon. Unless otherwise stated the gallon mentioned is the standard United States gallon or wine gallon of 231 cubic inches. In a number of cases the iron and aluminium were reported together. Such figures have been omitted in preparing the data here given, nor is reference made to those spring-waters in which a trace of alumina was reported. The pages of Dr. Peale's report on which the analyses occur, from which the data regarding aluminium are quoted, are given, as well as the name of the analyst when it was stated. References to the original place of publication are not given, as they were not quoted by Dr. Peale. The following table records the data regarding the occurrence of alumina (Al,O3):

AMOUNT OF ALUMINA IN THE WATER OF A NUMBER OF MINERAL SPRINGS IN THE UNITED STATES.

Name of Spring.	Analyst.	Page.	Alumina per Gallon.
Maine.			Grains.
Poland Star Spring	F. L. Bartlett	15	.32 a
Roscierucian Spring		16	.03
Hartford Cold Springs		16	.24
New Hampshire.			
Iron Spring, Milford	G. E. Sewell	17	1.04 b
Birchdale Springs, Concord	G. B. bewen	-1	1.04 0
Springs	C. F. Chandler	18	.12
Unity Springs, Iron Spring	S. Dana Hayes	18	.04
Vermont.			
Middletown Springs, No. 1	Peter Collier	20	.67 €
New York.			
Artesian Lithia Spring, Ball-			
ston, Spa	C. F. Chandler	33	.08
Franklin Artesian Well, Balls-		33	
Washington Lithia Well (old	C. F. Chandler	33	.26
Washington Lithia Well (old			
Conde Dentoman), Banston,			
Spa	C. F. Chandler	33	.40
Springe Sulphur, Chittellango	C. F. Chandler	22	.08
Springs	C. F. Chandier	33	.00
Springs	C. F. Chandler	33	.22
Cherry Valley Phosphate Spring	C. F. Chandler	34	.14
Lebanon Thermal Spring	H. Dussance	35	.45
Richfield Sulphur Spring	T. Deecke	37	.10
Saratoga Champion Spouting			
Spring.	C. F. Chandler	38	.46
Saratoga Crystal Springs	C. F. Chandler	38	.31
Saratoga Congress Spring Saratoga Empire Spring	J. R. Chilton C. F. Chandler	38 38	.32
Saratoga Eureka Spring	Allen	-39	.42
Saratoga Flat Rock Spring	C. F. Chandler	39	.23
Saratoga Hathorn Spring	C. F. Chandler	39	.13
Saratoga High Rock Springs	C. F. Chandler	39	1.22
Saratoga Pavilion Spring	C. F. Chandler	40	-33
Saratoga Pavilion Spring	J. R. Chilton	40	.42
Saratoga Putnam Spring	J. R. Chilton	40	.56
Saratoga New Putnam Spring	C. F. Chandler	40	.22
Saratoga A or Alum Spring Saratoga Seltzer Spring	J. G. Pohle C. F. Chandler	40	.38
Saratoga Union Spring	C. F. Chandler	40	.37
Saratoga United States Spring	C. F. Chandler	41	.09
Saratoga Vichy Spring	C. F. Chandler	41	.48
New Jersey.			
Schooley's Mountain Spring	C. McIntyre, Jr.	42	7.4
Warwick Spring		43	.14
	or a r community	43	.,,

a. With iron. b. With soda. c. Grains per cubic foot.

ALUMINA IN MINERAL SPRINGS-Continued.

Name of Spring.	Analyst.	Page.	Alumina per Gallon.
Pennsylvania.			Grains.
Cresson Magnesia Spring	F. A. Genth	47	.01
Gettysburg Lithia Spring	F. A. Genth	48	.02
Mineral Springs, Hanover	Hollenbush	48	.016
Kane Sulphur Spring	F. A. Genth	1000	
Maryland.	1.11. 001111	49	.03
Strontia Mineral Spring	W. Simon	F 2	1.86 a
Virginia.	W. Dimon	53	1.00 4
Bath Alum Spring, No. 1	Hayes	58	10 20
Bath Alum Spring, No. 3	Hayes	58	10.29 12.29 b
Clifton Spring, No. 1	J. L. Campbell	58	
Blue Ridge Spring	F. A. Genth		.07 C
Farmville Lithia Springs, No. 2	E. T. Fristoe	59	.14
Mineral Springs, near Farmwell	13. 1. 1 113000	00	2.52
Station, No. 1	R. B. Riggs	60	.0070 d
Mineral Springs, near Farmwell	D D D'		
Station, No. 3 Boiler Bath (Old Ladies' Boiler	R. B. Riggs	60	.0105 d
Bath Hot Spring)	F. W. Clarke	61	.0020 d
Hot Spout Bath	F. W. Clarke	61	.0025 d
Octagon Bath	F. W. Clarke	61	.0035 d
New Hot Spring	F. W. Clarke	61	.0060 d
Sulphur Bath or Ladies' Sulphur			
Bath	F. W. Clarke	61	.0065 d
Jordan Alum Chalybeate Spring.	J. W. Mallet	62	.05
Jordan White Sulphur Springs	T. Antisell	63	.01
Massanetta Mineral Springs	J. W. Mallet	63	.16
Rawley Springs, Main Fountain Rockbridge Alum Springs, Chaly-	J. W. Mallet	63	.05 C
beate Spring	J. W. Mallet	64	.06
Rockbridge Alum Spring, No. 1	A. A. Hayes	64	14.76
Rockbridge Alum Spring, No. 2	A. A. Hayes	65	17.91
Rockbridge Alum Spring, No. 3	M. B. Hardin	65	43.95
Rockbridge Alum Spring, No. 4	A. A. Hayes	65	24.09
Rock Enon Springs	Gale and Mew	67	.80
Roanoke Red Sulphur Springs	M. B. Hardin	67	.01
Shenandoah Alum Springs	J. W. Mallet	67	12.39 C
West Virginia.			
Capon Main Spring	J. W. Mallet	71	.02 6
Capon Beauty Spring	J. W. Mallet	71	.02 C
Greenbrier White Sulphur Sour	3	1	
Spring	W. B. Rogers	72	16.20 €
Salt Sulphur Iodine Spring	D. Stewart	73	.18
	- Della Lande S	, ,	

a. Parts per 100,000.

b. Also aluminium phosphate. See p. 113.

c. Grains per imperial gallon.

d. Grams per liter.

e. Grains in 100 cubic inches.

ALUMINA IN MINERAL SPRINGS .- Continued.

Name of Spring.	Analyst.	Page.	Alumina per Gallon.
North Carolina.			Grains.
Greensborough Spring	W. C. Kerr, a	77	.18
Panacea Spring, near Littleton	W. C. Kerr, a	78	.32
Park's Alkaline Mineral Spring	A. R. Ledoux	78	3.50
Georgia.			
Helicon Springs	H. C. White	85	.05 b
Alabama.			AND THE RESERVE
Johnson's Wells	W. C. Stubbs	93	12.41
Talladega Spring	W. C. Stubbs	93	1.45
Mississippi.			
Lauderdale Springs	L. Harper	97	.00005 6
Tennessee.		1	3
	A. Dove	102	2.00
Austin's Springs	J. M. Safford	103	.06
Galbraith's Springs	W. A. Noyes	103	.04
Hurricane Springs	J. M. Safford	104	.29
Montvale Springs	J. B. Mitchell	104	.50
Tilford's Mineral Well	J. M. Safford Troost	105	.06
White Cliff Springs	110051	100	.08
Kentucky.			
Estill Springs, White Sulphur	D D	1	,
Spring	R. Peter R. Peter	111	.02 d, e
Olympian White Sulphur Spring .	R. Peter	116	.0006 f
Kuttawa Mineral Springs	J. P. Barnum	118	1.20 f
Arkansas.			
	E. H. Larkin	122	-45
Hot Springs	Wright and	122	.49
	Merrill		
Texas.			
Palo Pinto Well		127	1.54
Wootan Well, No. 1	C. F. Chandler W. M. Mew	128	1.22
Wootan Well, No. 4	W. M. Mew	128	1.56 3.46
Ohio.	***************************************	120	3.40
	O N Staddard		-0
Green Mineral Spring	O. N. Stoddard E. S. Wayne	133	.98
Cedar Springs, Washington	D. O. Wayne	134	
Spring	A. Fennel	134	.22
Indiana.			
Greencastle, North or Daggy		1	
Spring	E. T. Cox	138	.19 b

a. Reported in Geol. North Carolina, 1 (1875). e. With trace of phosphates.

b. Grains per imperial gallon.

c. Parts per 100.

d. Grains per 1000.

f. Parts per 1000.

g. With iron and manganese carbonates.

ALUMINA IN MINERAL SPRINGS .- Continued.

Name of Spring.	Analyst.	Page.	Alumina per Gallon.
Indiana.			Grains.
Greencastle, Middle or Dewdrop			Oranio.
		138	.09 a
Spring		139	.35 a
West Saratoga Spring, No. 1	E. T. Cox	140	.22 a
West Saratoga Spring, No. 2	E. T. Cox	140	.43 a
Illinois.			
Alcyone Mineral Springs	Wheeler and		
Alcyone Mineral oprings	Blaney	144	1.99 a
Glen Flora Springs	I V Z Blaney	TAA	
Versailles Mineral Spring, No. 1.	J. V. Z. Blaney J. V. Z. Blaney	144	.15 .73a, b
Rockford Artesian Well, No. 1	E. G. Smith	144	.0009 6
		-44	.00090
Michigan.			
Riverside Magnetic Spring, No. 2		147	.51 d
Butterworth's Magnetic Spring,	S. P. Duffield	147	.49 a
Grand Haven Mineral Spring	C. G. Wheeler	148	.30
Mount Clemens Mineral Well	S. P. Duffield	148	29.47
Mount Clemens Media Spring Mount Clemens Soolbad Spring	S. P. Duffield S. P. Duffield	148	29.00
Warner's Spring	S. P. Dumeid	148	11.21
		150	. 25
Wisconsin.			
Arctic Spring.	W. W. Daniels	153	.15
Bethesda Springs, Waukesha	C. F. Chandler	153	. I 2
Black Earth Mineral Springs	A. C. Barry	153	1.00 €
Buckhert's Fountain	G. Bode	153	. 16
Eureka Springs	G. Bode	154	.19 C
Fountain Springs	J. V. Z. Blaney	154	.00
Gibson Springs	G. Bode F. C. Chandler	154	.13 6
Glen Springs	F. C. Chandler	154	.05
Hacket's Spring	G. Bode	154	.13
Horeb Springs	G. Bode	154	.23
Hygeia Springs	A. Thiel	154	.72
Iodo-Magnesian Springs	F. C. Chandler	154	.06
Jacob's Artesian Well	G. Bode	154	.19
Jordan's Mineral Well	G. Bode	154	.68
Palmyra Springs, Zenobia's Foun-			
tain	G. Bode	155	.22
Palmyra Springs, Eye Springs	G. Bode	155	.19
St. Croix Mineral Spring	J. V. Z. Blaney	155	.49
Shealtiel Mineral Springs	G. Bode	156	.09
Sheboygan Mineral Spring	G. Bode	156	1.10
Sheboygan Mineral Spring	C. F. Chandler	156	.13
Sheridan Springs	G. Bode	156	.05
Silurian Spring	W. S. Haines G. Bode	156	.59
Vesta Spring	G. Bode	156	.13

a. Grains per imperial gallon.

b. With trace of iron.

c. Parts per 1000.

d. With calcium phosphates.

c. Apparently parts per 100 of total ash.

f. With iron.

ALUMINA IN MINERAL SPRINGS .- Continued.

Name of Spring.	Analyst.	Page.	Alumina per Gallon.
Wisconsin.			Grains.
White Rock Spring	I. Campbell Browne C. D. Marsh	157	.75 a .000411 b
Chien	G. Bode Daniels	157 157	.0610
Minnesota. Owatonna Vichy Spring Owatonna Mineral Spring (name	Enno Sander	159	.10
unknown)	J. A. Dodge	159 159	. 28 2 . 38
Dakota. Artesian Well at Aberdeen Mineral Well at foot of bluff at	Erastus G. Smith	161	.0816
Pierre	G. A. Mariner	161	.01
Iowa Acid Spring	J. H. Siebel	162	226.41
Springs	J. W. Draper G. Bode	163	.29
Missouri. Bowsher Mineral Spring	P. Schweitzer	168	.6484
Columbia Chalybeate Springs	P. Schweitzer	168	.79
Climax Springs	H. W. Wiley P. Schweitzer	168	5.08 c .67
Akesion Spring	C. P. Williams	169	.17
Sweet Spring	C. P. Williams P. Schweitzer	169 169	.09
Kansas.			
Flowing Spring	Barnes and Sim Barnes and Sim E. H. Failyer	175 175 175	1.12 1.4083
Montana. Hunter's Hot Spring	W. A. Noyes	180	.12 d, f
Utah. Utah or Bear River Hot Springs	F. W. Clarke	187	.0040 €
Colorado. Hartsel Hot Mineral Springs	G. E. Patrick	192	.65 d

a. Grains per imperial gallon.

b. Parts per 100.

c. With iron oxid.

d. Parts per 100,000.

e. Parts per 1000.

f. Aluminium.

ALUMINA	IN	MINERA	L SPRIN	GS-C	Continued.
---------	----	--------	---------	------	------------

Name of Spring.	Analyst.	Page.	Alumina. per Gallon
California. El Paso de Robles Hot Sulphur			Grains.
Spring	Price and Hewston	210	.22 a
SpringSkagg's Hot Springs	E. W. Hilgard	211	.01
Highland Seltzer Spring	Rising	212	1.56
Highland Dutch Spring	Rising	212	.11
Highland Magic Spring	Rising	212	.17
Mono Basin Warm Springs	T. M. Chatard	212	.0018 b
Summit Soda Spring	J. F. Rudolph	214	1.75
Tolenas Springs	J. Hewson, Jr.	214	.96
Washington.			
Medical Lake	G. A. Mariner	218	.175

a. Grains per imperial gallon.

b. Grams per liter.

The following table reports data on the occurrence of aluminium sulphate in mineral waters of the United States:

AMOUNT OF ALUMINIUM SULPHATE IN THE WATER OF A NUMBER OF MINERAL SPRINGS IN THE UNITED STATES.

Name of Spring.	Analyst.	Page.	Aluminium Sulphate per Gallon.
New York. Oak Orchard Acid Springs, No. 1 Oak Orchard Acid Springs, No. 1 Oak Orchard acid water " " " " Pennsylvania.	Silliman and Norton J. R. Clinton Porter H. Erni W. J. Craw	37 37 37 37 37 37	Grains. 21.69 9.68 6.41 .37 a .32 a
Cresson Iron Spring	F. A. Genth F. A. Genth F. A. Genth F. A. Genth S. A. Lattimore	47 47 48 49 49	1.60 21.20 1.53 6.58 .55
Bath Alum, No. 2	W. H. Taylor W. H. Taylor M. B. Hardin W. Gilham W. P. Tonry W. P. Tonry W. P. Tonry	58 58 59 59 59 59 59	29.99 .02 24.18 7.24 8.18 b 9.07 b 3.04 b

a. Parts per 1000.

b. Grains per imperial gallon.

ALUMINIUM SULPHATE IN MINERAL SPRINGS-Continued.

ALUMINIUM SULFITATE IN	MINERAL SI KI	1100	Communea.
Name of Spring.	Analyst.	Page.	Aluminium Sulphate per Gallon.
		114	Grains.
Cold Culphur Springs			
Cold Sulphur Springs Church Hill Alum Springs	J. C. Booth	59	.25
	J. C. Booth	00	72.93 a
Jordan Alum Springs alum	W E V Ailson	60	0
spring	W. E. K. Aiken	62	25.38
Jordan Alum Springs, No. 2	J. W. Mallet	62	11.20
Jordan Alum Springs, No. 3	J. W. Mallet J. W. Mallet	62	6.88
Jordan Alum Springs, No. 4	J. W. Mallet	62	81.05
Jordan Alum Springs, No. 5	J. W. Mallet	62	26.11
Jordan Alum Springs, No. 6	J. W. Mallet	62	27.85
Pulaski Alum Spring	J. L. Campbell	63	21.59
Orkney Healing Spring	J. W. Mallet	64	.03
Orkney Powder Spring	J. W. Mallet	64	.02
Orkney Bear Wallow Spring	J. W. Mallet	64	.06
Rockbridge Alum Spring, No. 1	M. B. Hardin	64	31.25
Rockbridge Alum Spring, No. 2	M. B. Hardin	65	42.61
Rockbridge Alum Spring, No. 4	M. B. Hardin	65	72.37
Rockbridge Alum Spring, No. 4	M. B. Hardin	65	19.00
Rockbridge Alum Spring, No. 5	J. W. Mallet	65	11.20
Rockbridge Alum Spring, No. 6	J. W. Mallet	65	6.88
Rockbrigde Alum Spring, No. 7	J. W. Mallet	65	81.05
Rockbridge Alum Spring, No. 8	J. W. Mallet	65	26.11
Rockbridge Alum Spring, No. 9	J. W. Mallet	65	27.85
Stribling or Augusta Alum			
Spring	J. L. Campbell	66	16.68 a
Spring, No. 4	D. K. Little	66	16.69
Stribling or Augusta Alum			
Spring, No. 5	D. K. Little	66	17.95
Stribling or Augusta Alum			
Spring, No. 6	D. K. Little	66	38.41
Seven Springs, Abingdon	J. W. Mallet	67	15.22 b
Variety Springs, Alum Spring	W. Gilham	67	34.43
Wallawhatoola Alum Springs	J. L. Campbell	68	72.10
Wallawhatoola Alum Springs	C. F. Chandler	68	137.889 C
Yellow Sulphur Springs	W. Gilham	68	3.18
West Virginia.			
Greenbrier White Sulphur Springe			
Greenbrier White Sulphur Springs	W P Posses	-	
(name unknown)	W. B. Rogers	72	.02
Irondale Springs	Breneman	73	11.34
Georgia.			
Catoosa Spring, No. 9, "White			
Sulphur"	W. J. Land	83	2.47
Catoosa Spring, No. 10, "Buffalo"	W. J. Land	83	2.38
falo"	W. J. Land	83	.50
Catoosa Spring, No. 2, "Red	W. J. Land	83	.66
a Risulphate	h Parts per	100	

a. Bisulphate.

b. Parts per 100.

c. Also 4.201 grains potassium and aluminium sulphate and 3.867 grains sodium and aluminium sulphate.

ALUMINIUM SULPHATE IN MINERAL SPRINGS-Continued.

Name of Spring.	Analyst.	Page.	Aluminium Sulphate per Gallon.
Catoosa Spring, No. 4, "Chaly-			Grains.
beate". Catoosa Spring, No. 3, "Cos-	WIIIand	84	.67
metic	W lond	84	2.36
Catoosa Spring, No. 5, "Magnesia". Catoosa Spring, No. 6, "Congress".	W. J. Land	84	.69
gress"	W. J. Land W. J. Land	84 84	1.10
Sulphur"	W. J. Land W. J. Land	84 84	1.42 -43
Mississippi.			
Cooper's Well	J. Lawrence Smith	97	6.12
Tennessee.			
Raleigh's Mineral Springs, Box Spring	Enno Sander	105	.153 a
Spring	Enno Sander	105	.086 a
Kentucky.			
Estill Springs, Black Sulphur			
Spring.	R. Peter	III	.02 b
Kentucky Alum Springs Hickman's Springs, No. 1, Alum	L. D. Kastenbine R. Peter	114	40.26 1.25 b
Hickman's Springs, No. 2, Alum Hickman's Springs, No. 6,	R. Peter	114	.33 b
Sweet	R. Peter	114	.35 b
Louisville Artesian Well Paroquet Springs	J. Lawrence Smith J. Lawrence Smith	115	1.80 6
Davis Mineral Well	E. S. Wayne	118	83.16
Arkansas.			
Ravenden Spring	Wright and Merrill	122	2.36
Texas.			5.40 d
Fairview Springs.	F. W. Mallet	127	45.52
Sour Lake Mineral Spring, No. 7 Sour Lake Mineral Spring, No. 9	F. W. Mallet	128	31.28
Weaver Well	J. K. Wright	128	23.24 100.08 e
Sour Springs, Caldwell County	H. H. Dinwiddie	128	100.00 6
Ohio.			
Cuyahoga Lithia and Magnesia Springs, Bitter Water		133	515.51
Indiana.			A CONTRACTOR OF THE PARTY OF TH
French Lick Springs, Pluto's Well	E. T. Cox	138	5.98 f
	1 7 1 11	41000	

a. Grams per 10,000.
b. Grains per 1000.
c. Also 1.21 grains aluminium chloride.

d. Including alumina.
 e. Including potassium sulphate.
 f. Grains per imperial gallon.

ALUMINIUM S	ULPHATE IN	MINERAL S	SPRINGS—Continued.
-------------	------------	-----------	--------------------

Name of Spring.	Analyst.	Page.	Aluminium Sulphate per Gallon.	
Indian Spring	E. T. Cox E. T. Cox E. T. Cox	139 139 141	Grains. .99 11.67 a 5.41 a	
Missouri. Bratton Spring	P. Schweitzer Wright and Merrill	168	52.45 b 5.20	
Hot Springs at Hot Springs Station	T. M. Chatard	202	.0063 <i>c</i>	
Thermal Acid Spring	O. Loew	210	127.00 d	

a. Grains per imperial gallon.b. Tersulphate.

The table following reports the occurrence of aluminium phosphate in the mineral waters of the United States:

AMOUNT OF ALUMINIUM PHOSPHATE IN MINERAL WATERS OF THE UNITED STATES.

Name of Spring.	Analyst.	Page.	Aluminium Phosphate per Gallon.
Virginia.	TENTAL STATE		Grains.
Alleghany Springs	F. A. Genth	58	.03 a
Bath Alum, No. 3	Hayes	58	3.15b
Wolf Trap Lithia Springs	M. B. Hardin	68	.04
West Virginia.		14000	
Borland Mineral Well	T. G. Wormley	71	.23
Hart Well	S. C. Wells	73	.23
Michigan.			4.00
Midland Magnetic Well	S. P. Duffield	148	1.73 €
Wisconsin.			
Sparta Mineral Magnetic Well	J. M. Hirsh	156	.06 €
oparta Amerai Magnetic Well	J. M. IIIIon	130	.000

Also aluminium silicate. See p. 114.
 Also alumina. See p. 106.
 Grains per imperial gallon.

Several authors report the occurrence of aluminium silicate. Such data are summarized in the following table:

c. Grams per liter.
d. Persulphate, parts per 100,000.

AMOUNT OF ALUMINIUM SILICATE IN MINERAL WATERS OF THE UNITED STATES.

Name of Spring.	Analyst.	Page.	Aluminium Silicates per Gallon.
Virginia. Alleghany Springs	F. A. Genth	58	Grains.
Cullum's Ferruginous Spring Illinois.	Abequin	92	3.320
Perry Spring, No. 3, Lower Sulphur	H. Engelman	144	. 27

Also aluminium phosphate. See p. 113.

In addition to the above it should be noted that an analysis quoted by Dr. Peale (p. 133) reports in Cuyahoga Lithia and Magnesia Springs, Lithia Well, Ohio, 17.67 aluminium chloride (grains per gallon).

Pettenkofer (Ann. Chem. Pharm., 77, p. 183; Jahresb. Chem., 1851, p. 653) in the Adelheid spring at Heilbrunn, Bavaria, reports 0.1424 grain alumina per pound (7680 grains).

Philippi (Jahrb. Vereins Naturk. Nassau, Pt. 8, 1852, p. 90; Pharm. Centbl., 1853, p. 12; Jahresb. Chem., 1852, p. 753) notes a trace of alumina in Wiesbaden Faulbrunnen.

Phillips, J. A. (Chem. News, 27, p. 62; Proc. Roy. Soc. London, 21, p. 132; Phil. Mag., 4th ser., 46, p. 26; Jahresb. Chem., Naumann, 1873, p. 1243), reports in two analyses of a saline spring at Camborne, Cornwall, 0.3456 and 0.3460 gram alumina per liter.

Pistor, C. (Ber. Deut. Chem. Gesell., 17 (1884), p. 2894; Jahresb. Chem., 1884, p. 2033), reports 0.0450 gm. Al₂O₃ per 1000 cc. in Römerbrunnen from Echzell in Wetterau.

Planta (Ann. Chem. Pharm., 115, p. 330; Chem. Centbl., 1860, p. 1010; Jahresb. Chem., 1860, p. 837) notes 0.00004 gm. aluminium phosphate per 1000 gms. in the medicinal spring at Bormio, Italy.

Planta and Kekulé (Ann. Chem. Pharm., 87, p. 364; Pharm. Centbl., 1854, p. 59; Jahresb. Chem., 1853, p. 713) found in sulphur-water from Serneus, Switzerland, Canton Graubünden, 0.0012 part alumina per 1000 parts.

Planta and Kekulé (Ann. Chem. Pharm., 90, p. 316; Pharm. Centbl., 1854, p. 642; Jour. Prakt. Chem., 63, p. 61; Jahresb. Chem., 1854, p. 763) examined the mineral waters of St. Moritz, Canton Graubünden, Switzerland, finding alumina as follows: Large (old) spring 0.0003 and Small (new) spring 0.0004 (parts per 1000).

Planta-Reichenau, A. v. (Die Heilquellen von Alveneu, Tiefenkasten, Solis in Graubünden, etc., 1865; Ann. Chem. Pharm., 136, p. 145; Jahresber. Chem. 1865, p. 935), notes 0.0047 part aluminium phosphate per 1000 parts in the sulphur spring at Alveneu and 0.0068 part in the Donatusquelle in Albulathale. A trace of alumina is noted also in St. Petersquelle at Tiefenkasten.

Planta-Reichenau, A. v. (Ann. Chem. Pharm., 155, p. 161; Chem. Centbl., 1870, p. 581; Amer. Chemist, 2d ser., 1, p. 228; Jahresb. Chem., Naumann, 1870, p. 1383), reports 0.0091 part aluminium phosphate per 10,000 parts in the Ragaz-Pfäffers thermal spring.

Planta-Reichenau, A. v. (Ber. Deut. Chem. Gesell. (1878), p. 1793; Jahresb. Chem., 1878, p. 1306; Jour. Chem. Soc. London, 1879, II, p. 126), reports 0.074, 0.068, and 0.013 part aluminium phosphate per 10,000 parts in Natural Soda-water from Passug, Iodine Donatus from Solis, and in St. Petersquelle from Tiefenkasten, respectively.

Plumert, A. (Chem. Centbl., 1884, p. 423 (Ausz.); Jahresb. Chem., 1884, p. 2037), reports 0.206 part aluminium sulphate per 10,000 parts in the thermal sulphur-spring water from Tschéhirghé, and 0.918 part aluminium sulphate per 10,000 parts in similar water from Bad Jeni-Kaplidja, both in Anatolia.

Poggiale and Lambert (Compt. Rend. Acad. Sci. Paris, 54, p. 1062; Jour. Pharm., 41, p. 337; Instit., 1862, p. 172; Rép. Chim. Appliquée, 4, p. 214; Jahresb. Chem., 1862, p. 817) report 0.001 gm. Al₂O₃ per 1000 gms. in warm water from an artesian well at Passy, France.

Pohl, J. J. (Wien. Acad. Ber., 15, p. 303; Jahresb. Chem., 1855, p. 842), found traces of alumina in a spring-water from Vienna (Josefstadt).

Poleck, T. (Jour. Prakt. Chem., 52, p. 353; Pharm. Centbl., 1851, p. 540; Jahresb. Chem., 1851, p. 654), notes 0.00960 grain

aluminium phosphate per pound (7680 grains) in mineral water from Nieder-Langenau, and also reports an analysis by Duflos of the same water in which the occurrence of a trace of aluminium phosphate is noted.

Poleck, T. (Ber. Deut. Chem. Gesellsch., 1879, p. 1902; Jahresb. Chem., 1879, p. 1263), reports 0.0087 gm. Al₂P₂O₈ per 10,000 cc. in the Oberbrunnen at Flinsberg, Silesia.

Poleck, T. (Kronenquelle zu Salzbrunn, Breslau, 1882; Jahresb. Chem., 1884, p. 2034), notes 0.00047 Al₂O₃ and 0.00036 part Al₂P₂O₈ per 1000 parts in the Kronenquelle at Salzbrunn.

Poumarède (Rev. Sci. Ind., 38, p. 21; Jahresb. Chem., 1850, p. 627) reports 0.028 gm. alumina per 1000 cc. in mineral water from Vilaine-Saint-Aubin, Dept. Loiret.

Power, F. B. (Jour. Prakt. Chem., 2d ser., 19, p. 223; Jour. Chem. Soc. London, 36 (1879), II, p. 698), notes 0.0019 Al₂O₃ part per 10,000 parts in mineral water of Rosheim, Alsace.

Pozziale (Jour. Pharm., 3d ser., 24, p. 277; Jahresb. Chem., 1853, p. 716) notes 0.006 gm. alumina per 1000 gm. in mineral water from Orezza in Corsica.

Pozziale (Jour. Chim. Med., 3d ser., 9, p. 81; Jour. Pharm., 3d ser., 23, p. 114; Jahresb. Chem., 1853, p. 718) found 0.0150 and 0.018 part alumina per 1000 parts in Bullicame sulphurwater and the iron-water, respectively, from Viterbo.

Pribram, R. (Chem. Untersuchung der Arsenquelle zu Dorna-Sara Czernowitz; Jahresb. Chem., 1885, p. 2318), notes o.00112 part Al₂O₃ per 1000 in the Roumanian arsenic spring at Dorna-Sara.

v. Radziszewski (Arch. Pharm., 3d ser., 13 (1878), p. 459; Jahresb. Chem., 1879, p. 1265) notes a trace of aluminium in Carlsquelle and Ameliaquelle water from Iwoniez, Austria-Hungary.

Rasenack, P. (Arbt. K. Gsndhtsamt., 5, p. 370; Chem. Centbl., 1889, II, p. 166; Jahresb. Chem., 1889, p. 2632), notes a trace of aluminium in mineral water from the neighborhood of Bimbia, Africa.

Reade, T. M. (Silliman's Amer. Jour., 3d ser., 29, p. 290; Jahresb. Chem., 1885, p. 2315), reports analysis of river-water. Mississippi water contained 3.13 parts Al₂O₃ per 100,000 (1.753)

grains per gallon); Amazon River water, 0.38 part Al₂O₃ (with Fe₂O₃) per 100,000.

Reibenschuh, A. F. (Wiener. Akad. Ber., 62, 2. Abt., p. 786; Jahresb. Chem., Naumann, 1870, p. 1385), notes in water from the Johannis spring which is near Stainz in Steiermark 0.0269 part aluminium phosphate per 10,000 parts.

Reibenschuh, A. F. (Mitt. Ver. Steiermark, 30 (1894), p. 358; Jour. Chem. Soc. London, 70 (1896), II, p. 435), in water from the Radein (Styria) Sauerbrunnen found 0.0092 part Al₂O₃ per 10,000 parts.

Reibenschuh, A. F. (Mitt. Ver. Steiermark, 29 (1893), p. 262; Jour. Chem. Soc. London, 70 (1896), II, p. 435), notes 0.05371 part Al₂O₃ per 10,000 parts in water from the Semlitsch spring at Radein.

Reibenschuh, A. F. (Mitt. Ver. Steiermark, 34 (1898), p. 177; Jour. Chem. Soc. London, 76 (1899), II, p. 308), notes in water from the Neubrunnen at Radein, Styria, 0.10191 gm. Al₂O₃ per 10,000 gms.

Reichardt, E. (Arch. Pharm., 3d ser., 2 (1873), p. 124; Jahresb. Chem., Naumann, 1873, p. 1237), reports in Neue Stahlquelle from Lobenstein, Agnesquelle from Lobenstein, and spring-water from Steben, respectively, 0.00397, 0.00122, and 0.00386 part alumina per 1000 parts.

Reichardt, E. (Arch. Pharm., 3d ser., 16 (1880), p. 208; Jour. Chem. Soc. London, 40 (1881), p. 29), in the water of the Grosslüder mineral spring at Salzschlirf reports 0.0045 gm. Al₂O₃ per 1000 gms.

Reiner, M. (Wien. Akad. Ber., 56, 2. Abt., p. 456; Wiener Akad. Anz., 1867, p. 178; Jour. Prakt. Chem., 102, p. 58; Chem. Centbl., 1868, p. 384; Inst., 1868, p. 30; Jahresb. Chem., 1867, p. 1038), reports 0.088 part aluminium phosphate per 10,000 parts in the Wiener-Neustadt Sauerbrunn.

Reinsch (Jahrb. Prakt. Pharm., 21, pt. 3; Arch. Pharm., 2d ser., 67 (1851), p. 47; Jahresb. Chem., 1851, p. 666) notes 0.009457 gm. aluminium chloride per 1000 cc. in mineral water from Sulz (Alsace).

Riggs, R. B. (U. S. Geol. Survey Bul. 42, p. 147), reports analysis of two springs one mile from Farmwell Station, Lou-

doun County, Virginia. Spring A contained 0.0070 and Spring B, 0.0105 gm. Al₂O₃ per liter.

Riggs, R. B. (U. S. Geol. Survey Bul. 42, p. 148), in water from Beck's Hot Springs, near Salt Lake City, Utah, reports 0.0090 gm. Al₂O₃ per liter.

Riggs, R. B. (U. S. Geol. Survey Bul. 55, p. 91), in artesian-well water from Albany, Georgia, and Fitzpatrick, Alabama, reports 181.0 and 271.0 parts total solids per million, 1.0 and 2.4 being Al₂O₃. In similar wells at Smithville, Americus, and Montezuma, Ga., he reports traces of Al₂O₃.

Rivier and Fellenberg (Arch. Phil. Natur., 21, p. 59; Jahresb. Chem., 1853, p. 714) note a trace of alumina in mineral water from Saxon (Canton Wallis), Switzerland. It is stated that Morin (Arch. Phil. Nat., 22, p. 52; Jour. Pharm., 3d ser., 23, p. 188; Jour. Prakt. Chem., 58, p. 483; Pharm. Centbl., 1853, p. 325; Jahresb. Chem., 1853, p. 714) found 0.0005 part alumina per 1000 parts in the same water. Heidepriem and Poselger (Jour. Prakt. Chem., 58, p. 473; Pharm. Centbl., 1853, p. 324; Jahresb. Chem., 1853, p. 714) also note a trace of alumina in this water, as did Braun in two analyses reported by Rivier and Fellenberg (Jour. Prakt. Chem., 59, p. 303; Jahresb. Chem., 1853, p. 715).

Robinson, A. E., and C. F. Mabery (Jour. Amer. Chem. Soc. 18 (1896), p. 915; Jour. Chem. Soc. London, 72 (1897), II, p. 510), report in bittern water from a boring 2667 feet deep at Conneautsville, Crawford County, Pennsylvania, 31.1 parts AlCl₃ per 100,000.

Roux (Compt. Rend. Acad. Sci. Paris, 73, p. 910; Jour. Chem. Soc. London, 24 (1871), p. 1181)₂ reports 0.00720 gm. aluminium silicate per liter in the water of an artesian well at Rochefort.

Rupp, G. (Ztschr. Ang. Chem., 1891, p. 448; Jahresb. Chem., 1891, p. 2618), in a steel spring at Bad Griesbach found 0.003 gm. alumina per 1000 cc.

Russell, I. C. (U. S. Geol. Survey Bul. 108, p. 93, notes that T. M. Chatard found in Owens Lake (California) 0.023 part Al₂O₃ per 1000 parts.

Russell, I. C. (U. S. Geol. Survey Bul. 52, p. 38), quotes an analysis of James River, Va., water reported by W. H. Taylor

(An. Rpt., Richmond (Va.) Board of Health, 1876). The account of Al₂O₃ noted is 0.00041 part per 1000 parts.

Russell, I. C. (U. S. Geol. Survey Mon. 11, Tables, p. 176), quotes a number of analyses of American rivers, inclosed lakes and springs. Some of these have been noted already from other publications. Others follow, the values given in every case representing parts per 1000. Hydrant (river) water from Los Angeles, California, according to W. J. Jones (Rpt. Cal. State Board Health, 1878) contains 0.00171 part Al₂O₃ and hydrant (river) water from Sacramento, California, 0.00120 part Al₂O₃. E. R. Horsford (Geol. of New Jersey, 1868, p. 703) notes 0.01342 part Al₂O₃ in Pasaic River water taken four miles above Newark. T. M. Chatard reports 0.0013 Al₂O₃ in Humboldt River water taken at Battle Mountain, Nevada (see also Mon. 11, p. 41).

Of the analyses of American springs, two, which should be noted, contained alumina. These are the following: artesian well, "Glacier Spouting Spring," Saratoga, N. Y., analyzed by F. A. Cairns and C. F. Chandler (Amer. Chemist, 1872, Nov., p. 164), containing 0.00770 part Al₂O₃, and artesian well, Sheboygan, Wis., analyzed by C. F. Chandler (Amer. Chemist, 1876, p. 370), containing 0.0022 part per 1000 parts in both cases.

Russell states also that O. Loew (Ann. Rpt. Chief Engineers, 1876, Appendix JJ, p. 410) reports an analysis of the water of Owens Lake, California, noting a trace of aluminium.

According to data quoted by Russell (loc. cit.) Terreil (Larted, Geological Explorations of Dead Sea, p. 278) found that the Dead Sea water contains traces of alumina.

Say, M. (Wien. Acad. Ber., 13, pp. 298 and 457; Pharm. Centbl., 1854, p. 923; Jahresb. Chem, 1854, p. 771), reports (grams per 1000 gms.) in Hildegarde spring-water from Ofen 0.047, and in mineral water from Lippa 0.0157, both towns in Hungary.

Scherer (Ann. Chem. Pharm., 99, p. 257; Vierteljahressch. Prakt. Pharm., 6, p. 81; Jour. Prakt. Chem., 70, p. 151; Chem. Centbl., 1856, p. 810; Jahresb. Chem., 1856, p. 767) found c.0002 gm. alumina per 1000 cc. in Wernarzer mineral water,

and a trace in the sweet potable water near the Sinnbergerquelle, both waters from Brückenau, Germany.

Scherer (Neue. Jahrb. Pharm., 7, p. 309; Jahresb. Chem. 1857, p. 721) notes a trace of alumina in the saline water from Philippsquelle at Orb, Germany.

Scherfel, A. (Ungar. Naturw. Ber., 1, p. 195; Jahresb. Chem., 1886, p. 2321), notes 0.002349 part AlPO₄ and 0.019317 part Al₂O₃ per 1000 parts in water from a spring at Kirchdrauf in der Zips, Szepesváralja, Hungary.

Scherfel, A. (Ungar. Naturw. Ber., 1, p. 230; Jahresb. Chem., 1886, p. 2322), notes 0.014722 part AlPO₄ and 0.017778 part Al₂O₃ per 10,000 parts in mineral water from Czeméte near Eperies; Hungary.

Schmidt, C. (Arch. Naturk. Liv. Esth. Kurlands., 1st ser., 1, p. 293; Jahresb. Chem., 1854, p. 771), notes a trace of alumina in the bored saline-spring water from Staraja-Russa, Russia.

Schmidt, C. (Mélanges Phys. et. Chim. tirées Bul. Akad. Imp. Sci. St. Petersburg 7, p. 427; Petersburg Acad. Bul. 12, p. 1; Jahresb. Chem., 1867, p. 1042), notes 0.021 part alumina per 10,000 parts in iron water from Stolypin, Russia. Alumina was not noted in the deposit from the water.

Schneider, F. C. (Wiener. Akad. Ber., 2. Abt., 69, p. 55; Jahresb. Chem., 1874, p. 1330), reports 0.010, 0.006, 0.015, and 0.011 part alumina per 10,000 parts, respectively, in four St. Helena thermal springs at Battaglia, Italy.

Schröckinger (Verhandl, Geol. Reichsans., 1878, p. 89; Jahresb. Chem., 1878, p. 1301), reports 0.0044 part aluminium phosphate per 10,000 parts in water from a bored well 127 meters deep, near Brüx, Bohemia. The analysis was made by Gintl.

Schoeller, R. (Ber. Deut. Chem. Gesellsch., 20 (1887), p. 1784; Jahresb. Chem., 1887, p. 2537), reports analyses of streams affecting the Rio de la Plata water. That designated Uruguay water contained 0.0018 gm. alumina per 1000 gm.

Schwarz, E. (Wiener Akad. Ber., 55, 2. Abt., p. 35; Wiener Akad. Anz., 1867, p. 7; Chem. Centbl., 1867, p. 558, Inst., 1867, p. 151; Jahresb. Chem., 1867, p. 1037), notes 0.007 part aluminium phosphate per 10,000 parts in mineral water from Mödling near Vienna.

Schweissinger (Chem. Centbl., 1890, I, p. 875; Jahresb. Chem., 1890, p. 2662) notes a trace of alumina in the Bad Marienborn iron-sulphur water.

Shepard, C. U. (Silliman's Amer Jour., 2d ser., 47, p. 357; Jahresb. Chem., 1869, p. 1293), reports 0.0000093 part calcium, iron, and aluminium phosphate per 100 parts in artesian water from Charleston, South Carolina.

Siegmund, H., and P. Juhász (Wiener Akad. Ber., 54, 2. Abt., p. 216; Wiener Akad. Anz., 1866, p. 162; Inst., 1866, p. 400; Jahresb. Chem., 1866, p. 996) found 0.002 part aluminium phosphate per 10,000 parts in the Vöslau mineral water.

Simmler, T. (Jour. Prakt. Chem., 71, p. 1; Chem. Centbl., 1857, p. 498; Jahresb. Chem., 1857, p. 723), found 0.0046 gm. alumina with phosphoric acid per 1000 gm. in sulphur-water from Stachelberg, Canton Glarus, Switzerland.

Smith (Jour. Chem. Soc. London, 15 (1862), p. 57; Jour. Prakt. Chem., 89, p. 186; Chem. News, 5, p. 96; Rép. Chim. Appliquée, 4, p. 216; Jahresb. Chem., 1862, p. 822) found in the water of the Te Tarata hot spring on the coast of Rotomahana, north New Zealand, 0.32 grain alumina per gallon.

Smith (Jour. Prakt. Chem., 89 (1863), p. 186; U. S. Geol. Survey Rpt. 9, p. 655) notes 0.005 gm. alumina per kilogram in White Terrace Geyser water from New Zealand.

Smith, L. (Silliman's Amer. Jour., 2d ser., 12, p. 10; Jahresb. Chem., 1851, p. 668), reports aluminium sulphate in mineral water from Brusa (Asia Minor) as follows: Kukurtlu 0.0043 and Bademli Baghtsche 0.0020 gm. per 1000 cc., and Gueuzayasma a trace. He reports further (Silliman's Amer. Jour., 2d ser., 12, p. 366; Jour. Prakt. Chem., 55, p. 110; Jahresb. Chem., 1851, p. 668) a trace of aluminium sulphate in the Yalova hot spring, and 0.0221 gm. aluminium sulphate per 1000 cc. in the Touzla Spring water.

Smoot, L. E. (Amer. Chem. Jour., 19 (1897), p. 234; Jour. Chem. Soc. London, 72 (1897), II, p. 329), notes 0.891 gm. Al₂O₃ per liter in an alum-water from Lee County, Virginia.

Soubeiran, E. (Jour. Pharm., 3d ser., 32, p. 19; Jahresb.

Chem., 1857, p. 726), found 0.054 gm. alumina and silica per 1000 cc. in Vic-sur-Cère mineral water, Cantal, France.

Späth, E. (Chem. Centbl., 1889, II, p. 896; Jahresb. Chem., 1889, p. 2636), notes traces of Al₂O₃,P₂O₅ in Temple and Wiesen Spring water from Bad Steben at Oberfranken.

Spengler quotes F. Mohr's analysis of the Felsenquelle of Ems (Arch. Balneologie, 3 (1864), p. 136; Viertelj. Prakt. Pharm., 14, p. 234; Chem. Centbl., 1865, p. 608; Jahresber. Chem., 1865, p. 931). A pound (7680 grains) contained 0.0960 grain alumina.

Spica, P. (Gaz. Chim. Ital., 12 (1882), p. 555; Jahresb. Chem., 1882, p. 1623), noted traces of Al₂O₃ in two samples of water from Lake Derkol near Constantinople.

Spica, P. (Gaz. Chim. Ital., 22 (1892), I, p. 354; Jahresb. Chem., 1892, p. 2686), notes 0.001999 gm. aluminium phosphate per 1000 cc. in mineral water from Burge Malo.

Spurr, J. E. (U. S. Geol. Survey Mon. 31, p. 212), quotes an analysis by C. F. Chandler of water from Yampa Spring, Glenwood Springs, Colorado. This contained a trace of alumina.

Stackmann, A. (Russ. Ztschr. Pharm., 24, pp. 129, 145, 161, 209, 225, 273, 289, 321, 337; Jahresb. Chem., 1885, p. 2319), reports analyses of mineral water from Psekoup or Gorjatschy Kljutsch. Data are quoted for (1) Alexanderquelle, (2) Olgaquelle, (3) Marienquelle, (4) Karmalinquelle, (5) Michailquelle, (6) Kamenewquelle, (7) Südliche Drainagequelle, (8) spring without particular name, and (9) mixture of above eight waters. These contained respectively 0.0010, 0.0005, 0.0005, 0.0005, 0.0005, 0.0007, 0.0009, 0.0010, 0.0005, and 0.0012 Al₂O₃ (presumably grams per 1000 cc.). The author also reports 0.0050 gm. Al₂O₃ per 1000 cc. in a sulphur spring and 0.0390 gm. Al₂O₃ per 1000 cc. in a saline spring analyzed by him.

Steiger, G. (U. S. Geol. Survey Bul. 113. p. 113), in water from American Carlsbad Spring, Nashville, Illinois, found 3.60 parts Al₂O₃ per 1,000,000.

Stein, W. (Schmidt's Jahrb. Ges. Med., 70. p 142; Jahresb. Chem., 1851, p. 654), notes traces of alumina in the medicinal waters of Elster. Saxony, as follows: Gasquelle, Salzquelle or Augenquelle, and Trinkquelle or Stahlbrunnen.

Steinbrück (Arch. Pharm., 2d ser., 145 (1871), p. 97; Jahresb. Chem., 1871, p. 1228) reports 0.0212 and 0.0134 gm. Al₂O₃ per liter in two springs at Neuragoczi near Halle.

Stieren, E. (Vierteljahressch. Prakt. Pharm., 10, p. 365; Jahresb. Chem., 1861, p. 1112), notes 0.040830 part alumina per 1000 parts in artesian water from Peterson's saline water, Tarentum, Pennsylvania.

Strecker, A. and H. (Chem. Lab. Univ. Christiana, 1854, p. 67, Ann. Chem. Pharm., 95, p. 177; Jahresb. Chem., 1854, p. 770), in the sulphur-water of Sandefjord found 0.0068 part alumina per 1000 parts, and 0.0033 parts in near-by seawater.

Streit, S., and W. Holeček (Wiener Akad. Ber., 53, 2. Abt., p. 371; Jahresb. Chem., 1866, p. 995) note 0.021 and 0.0067 part aluminium phosphate per 10,000 parts, respectively, in Töplitz and Someraubad mineral water from Mähren.

Thresh, J. C. (Chem. News, 46, p. 226; Jahresb. Chem., 1882, p. 1635), found 10.416 gms. Al₂S₃O₁₂ and 0.065 gm. AlPO₄ per 10,000 cc. in the Orchard Alum Spring water, England.

Thresh, J. C. (Jour. Chem. Soc. London, 41 (1882), p. 117), in Buxton thermal water reports 0.000683 Al₂O₃.Ca₃(PO₄)₂ per 10,000 gms. He notes that in 1852 Playfair reported 0.034 part Al₂O₃.Fe₂O₃ per 10,000 parts in the same water.

Thörner, W. (Repert. Analyt. Chem., 3, p. 22; Jahresb. Chem., 1883, p. 1944), reports in a bored well (Soolquelle) at Melle, Germany, 0.000102 part Al₂O₃ per 1000 parts.

Thörner, W. (Ztschr. Angew. Chem., 1889, p. 309; Jahresb. Chem., 1889, p. 2631), reports 0.0020 gm. alumina per 1000 cc. in water from the Germania Spring at Schwalheim, Germany.

Thörner, W. (Chem. Ztg., 17, p. 1411; Jour. Chem. Soc. London, 66 (1894), II, p. 195), reports Al₂O₃ as follows in some German mineral-spring waters: Stahl 0.0132, Angelika at Tönnisstein 0.0056, Wilhelms 0.0679, New Spring at Melle 0.0045, and Sulphur Spring at Levern 0.0020 gm. per liter.

Thorpe, T. E. (Jour. Chem. Soc. London, 65 (1894), p. 772), reported analyses of a number of mineral waters from Chelten-

ham, England. The Chadnor Villa well contained 0.00015 part aluminium phosphate per 1000 parts, and the Lansdowne well and the Pittville wells, Nos. 1, 2, and 3, each traces.

Tissandier, G. (Compt. Rend. Acad. Sci. Paris, 80, p. 58; Jahresb. Chem., 1875, p. 1284), reports a trace of Al₂O₃ in melted snow collected in Paris and in the country.

Tournaire (Ann. Minn., 5th ser., 17, p. 65; Jahresb. Chem., 1860, p. 841), reported 0.165 gm. Al₂O₃.SiO₂ in mineral water from Roddes (France), near Ambert.

Tournaire (Ann. Minn., 5th ser., 17, p. 66; Jahresb. Chem., 1860, p. 841) found 0.055 gm. Al₂O₃.SiO₃ per 1000 cc. in a mineral water from Ceyssat, Dept. Puy-de-Dôme, France.

Treadwell, F. P. (Arch. Pharm., 231 (1893), p. 579; Jour. Chem. Soc. London, 66 (1894), II, p. 323), in the upper spring of Gyrenbad, Canton Zürich, found 0.000200 part AlPO₄ per 10,000 parts.

Trillich, H. (Chem. Ztg. II. Rept., p. 211; Chem. Centbl., 1887, p. 1239; Jahresb. Chem., 1887, p. 2534), notes 0.00109 gm. aluminium phosphate per 1000 gms. in the sulphur-water of Bad Wemding at Ries.

Trillich, H. (Chemische Analyse des Hauptbrunnens zu Münster a. Stein, Munich; Chem. Centbl., 1889, II, p. 896; Ausz.; Jahresb. Chem., 1889, p. 2631), notes 0.00168 part Al₂O₃ per 1000 parts in water from the Hauptbrunnen at Münster, Germany.

Tschermak (Jahrb. K. K. Geol. Reichsanstalt, 1858, p. 297; Jahresb. Chem., 1858, p. 796) found 0.0092 part aluminium phosphate per 1000 parts in mineral water from Kondran near Regensburg, Germany.

Tschermak, T. (Min. Petr. Mitth., 2d ser., 3, p. 315; Jahresb. Chem., 1880, p. 1528), quotes analyses made by P. Schnell and Stenner in 1855 of water from an iron spring at Slanik, Roumania. This contained 0.07 gm. Al₂O₃ per 10,000 gms.

Turner, T. (Chem. News, 49, p. 186; Jahresb. Chem., 1884, p. 2036), reports 0.0740 gm. Al₂O₃ (with some Fe₂O₃) per 1000 cc. in water from the salt wells of Dudley, England.

Valentiner, W. (Jour. Prakt. Chem., 99, p. 91; Jahresb. Chem., 1866, p. 994), in Oberbrunnen and Mühlbrunnen

mineral waters from Ober-Salzbrunn reports per 1000 parts, 0.005 and 0.0003 part aluminium with phosphoric acid, respectively.

Vierthaler, A. (Wiener Akad. Ber., 56, 2. Abt., p. 463; Wiener Akad. Anz., 1867, p. 220; Jour. Pharm. Chem., 102, p. 381; Jahrb. Chem., 1867, p. 1040), reports traces of aiumina in the Cattani No. 2 and San Francesco springs of Spalato.

Vohl, H. (Dingler's Polytech. Jour., 199, p. 311; Chem. Centbl., 1871, p. 269; Arch. Pharm., 2d ser., 146 (1871), p. 199; Jahresber. Chem., 1871, p. 1223), reports 0.0008 gm. Al₂O₃ per 10,000 cc. in one sample and 0.010 gm. in five samples of Rhine water taken under different circumstances near Cologne.

Vohl, H. (Ber. Deut. Chem. Gesellsch., 1875, p. 611; Arch. Pharm., 3d ser., 7, p. 134 (see under R. Bender); Jahresb. Chem., 1875, p. 1290), notes 0.0556 gm. alumina per 10,000 cc. in Birresborn mineral water.

Vohl, H. (Ber. Deut. Chem. Gesellsch., 1876, p. 20; Jahresb. Chem., 1876, p. 1298), reports 0.3925 gm. alumina per 10,000 cc. in Birresborn mineral water taken after the spring had been cleaned.

Vohl, H. (Ber. Deut. Chem. Gesellsch., 11 (1878), pp. 605, 877: Jahresb. Chem., 1878, p. 1299; Jour. Chem. Soc. London, 34 (1878), II, p. 714), notes in Gerolsteiner Schlossbrunnen, located near Pelm, Germany, 0.0002 part Al₂O₃ per 1000 parts, and in Birresborn spring-water from the same locality traces of alumina.

Vohl, H. (Ber. Deut. Chem. Gesell., 11 (1878), p. 1678; Jahresb. Chem., 1878, p. 1304), reports 0.3067 part Al₂O₃ per 10,000 parts in Ofener Rakoczy Bitterwasser. He quotes C. R. C. Tichborn's analysis, which gave 0.27 part Al₂O₃ per 10,000 parts, and J. Mólnar's, which gave 0.4840 part Al₂O₃ per 10,000 parts. The water varies in composition at different seasons, etc.

von der Marck, W. (Arch. Pharm., 2d ser., 102 (1860), p. 1; Vierteljahressch. Prakt. Pharm., 9, p. 161; Chem. Centbl., 1860, p. 485; Ztschr. Chem. Pharm., 1860, p. 170; Jahresb. Chem., 1860, p. 831), reports 0.00676 gm. alumina per 1000 in the Stahlbrunnen, one of the Hermannsborn mineral springs near Pömbsen, Germany.

Wait, F. G. (Rpt. Section of Chem. and Mineral., Geol. Survey Canada, 1898, p. 48 R), reports 0.345 part alumina per 1000 in the filtered water from Lake Goodenough, a soda lake in British Columbia. It was calculated that this represented 0.825 part aluminium phosphate or 64.016 grains per imperial gallon.

Wait (Geol. Survey Canada, Chem. Contributions, 1892-93, p. 78 R) notes 0.0002 part alumina with a little ferrous oxide per 1000 parts in water from a bored well 1943 feet deep near Deloraine, Manitoba.

Wallace (Analyst, 1880, p. 79; Jahresb. Chem., 1880, p. 1535) notes 8.0 parts Al₂O₃ per 1,000,000 in an English artesian water.

Wallace (Rept. British Assoc., 1862, Notices and Abstracts, p. 94; Jahresb. Chem., 1862, p. 818) found alumina and phosphate as follows: Loch Katrine water (thirty-five miles from Glasgow) 0.10, and the same water at Glasgow 0.16 grain per gallon.

Walton, G. E., in his volume on mineral waters of this country and Canada (The Mineral Springs of the United States and Canada, New York, 1883) quotes a number of analyses in which alumina was reported. Such data (unless noted from other sources) are included in the table on the next page.

Walz, G. F. (Neue. Jahrb. Pharm., 6, p. 265; Jahresb. Chem., 1856, p. 769), found 0.0010 gm. alumina per 1000 gm. in sulphur-spring water from Weisloch, Baden, the sample being taken at the spring-house. In a sample taken after flowing one-eighth mile through a pipe the water contained 0.0009 gm. alumina per 1000 gms.

Wandesleben (Neue Jahrb. Pharm., 3, p. 123; Pharm. Centbl., 1855, p. 361; Jahresb. Chem., 1855, p. 840) notes 0.0013 gm. alumina per 1000 gms. in Oestringer sulphur-water from Langenbrücken, Baden.

Weed, W. H., and L. V. Pirsson (U. S. Geol. Survey Bul. 139, p. 150) quote an analysis of White Sulphur Springs, Montana, which records a trace of alumina.

MINERAL WA'	TERS OF	THE	UNITED	STATES	AND	CANADA.
-------------	---------	-----	--------	--------	-----	---------

Spring.	Analyst.	Page.	Al ₂ O ₃ per Pint.	Aluminium Sulphate per Pint.
California Seltzers Spring		148	Grains.	Grains.
Versailles, Ill., Curry Spring	J. V. Z. Blaney	149	.091 a	
Caledonia, Ontario, Gas Spring	T. Sterry Hunt	164	.032	
Caledonia, Ontario, Saline Spring	T. Sterry Hunt	164	trace	
mittent Spring Caledonia, Ontario, sulphur-	T. Sterry Hunt	164	trace	
water	T. Sterry Hunt	183	.019	
Big Bone Kentucky springs		193		Ь
Berkshire, Mass., Soda spring.		335	b	
Caxton, Quebec, spring	T. Sterry Hunt	389	.036	
Slatersville, N. Y., spring	B. Hitchcock	399	.036	
Excelsior, Mo., springs		418	b	
South Park, Col., spring	G. E. Patrick	450	.047	

a. With trace of iron.

Weidel, H., and G. Goldschmiedt (Wiener Acad. Ber., 2. Abt., 74, p. 391; Jahresb. Chem., 1877; p. 1386) report 0.0063 part Al₂O₃ per 10,000 parts in O Tura Säuerling water from Hungary.

Wertheim, T. (Wiener Acad. Ber., 42, p. 479; Rép. Chim. pure, 3, p. 131; Jahresb. Chem., 1860, p. 836) notes 0.02899 part alumina per 10,000 parts in Franz-Joseph-Bad water from Tüffer, Austria.

Weselsky, P., and A. Bauer (Wiener Acad. Ber., 29, p. 585; Chem. Centbl., 1858, p. 652; Jahresb. Chem., 1858, p. 797) note 0.00227 part alumina (with phosphoric acid?) per 1000 parts in water from the König Ferdinand Eisenbad, Weidritzthal near Presburg, Hungary.

Wiggers (Bal. Ztg. 1, p. 4; Pharm. Centbl., 1854, p. 934; Jahresb. Chem., 1854, p. 762) notes that a potable water from a spring at Hofgeismar, Hesse-Nassau, Germany, contained a trace of alumina too small to weigh.

Wiggers (Arch. Pharm., 2d ser., 102, p. 215; Jahresb. Chem., 1860, p. 832) found 0.001920 grain alumina per pound (=7680 grains) in the Driburg, Germany, mineral water.

b. Qualitative analysis—amount not stated.

Wildenstein, R. (Jour. Prakt. Chem., 85, p. 100; Chem. Centbl., 1862, p. 200; Jahresb. Chem., 1862, p. 810), notes 0.00180 part aluminium phosphate per 10,000 parts in water from a hot mineral spring a Burtscheid, Germany.

Will (Ann. Chem. Pharm., 61, p. 181; Pharm. Centbl., 1847, p. 507; Jahresb. Chem., 1847–48, p. 1002) reports in water from Rippoldsau as follows: Josephsquelle 0.0953, Wenzelsquelle 0.0840, and Leopoldsquelle 0.0822 gm. alumina per 10,000 gms.

Will (Ann. Chem. Pharm., 81, p. 73; Pharm. Centbl., 1852, p. 237; Jahresb. Chem., 1851, p. 650) found 0.00119 part aluminium phosphate per 1000 parts in sulphur-water from Weilbach.

Willm, E. (Compt. Rend. Acad. Sci. Paris, 86, p. 543; Bul. Soc. Chim., 2d ser., 29, p. 291; Jahresb. Chem., 1878, p. 1309), notes in Aix-les-Bains (Savoy) sulphur spring 0.0013 gm. aluminium per liter, and in the alum-spring water 0.0003 gm. Al₂S₃O₁₂ per liter.

Willm, E. (Compt. Rend. Acad. Sci. Paris, 86, p. 613; Jour. Chem. Soc. London, 34 (1878), II, p. 560), analyzed mineral water from Challes, Savoy, reporting 1.21851 and 0.3306 gm. per liter soluble matter in Source principale and Petite Source, the former containing 0.0059 gm. alumina and the latter 0.0232 gm. alumina and sand together.

Willm, E. (Compt. Rend. Acad. Sci. Paris, 90, p. 547; Jahresb. Chem., 1880, p. 1534), reports an analysis of the Richard spring at Cransac, France. April 15, 1879, the water contained 0.02800 gm. Al₂S₃O₁₂, and July 14, 1879, 0.1760 gm. per 1000 cc.

Willm, E. (Bul. Soc. Chem., 2d ser., 31, p. 3; Jahresb. Chem., 1879, p. 1266), reports 0.0027, 0.0024, and 0.0012 gm. alumina per 10,000 cc., respectively, in three Auvergne mineral waters, and traces in three others.

Willm, E. (Compt. Rend. Acad. Sci. Paris, 90, p. 630; Jahresb. Chem., 1880, p. 1533), analyzed the Bussang springs in the Vosges. Salmade contained 0.0012, Ober Quelle 0.011, and Quelle Marie 0.0010 gm. Al₂O₃, respectively, per 1000 cc.

Willm, E. (Compt. Rend. Acad. Sci. Paris, 103, p. 416;

Jour. Chem. Soc. London, 50 (1886), p. 997, notes 0.0003 gm. alumina per liter in the Bosquet water of Bagnères de Luchon, Haute Garonne.

Willm, E. (Compt. Rend. Acad. Sci. Paris, 113, p. 87; Jahresb. Chem., 1891, p. 2617), notes per 1000 cc. in Madeleine water No. 1 0.0644, in Madeleine water No. 2 0.0248, in Cercle water 0.0154, in Damont water 0.0963, and in Des demoiselles spring 0.0399 aluminium sulphate. These springs are at Rennes-les-Bains (Aude). The first three belong to a company; the last two to the town.

Wittstein (Vierteljahressch. Prakt. Pharm., 10, p. 342; Jahresb. Chem., 1861, p. 1097) reports Al₂O₃ in water as follows: Ohe River 0.00017, Höhenbrunnerfilz 0.00076, Schliessheimer moor water 0.00029, Isar River 0.00030, and Brunnthaler water 0.00013 gm. per 1000 gms.

Wolff, J. (Wiener Akad. Ber., 56, 2. Abt., p. 55; Wiener Akad. Anz., 1867, p. 137; Jour. Prakt. Chem., 101, p. 318; Chem. Centbl., 1867, p. 424; Inst., 1867, p. 391; Jahrb. Chem., 1867, p. 1039), reports 0.0064 part alumina and phosphoric acid per 10,000 parts in water from the mineral spring at Siebenbürg.

Wreden, F., and A. Fuchs (Ber. Deut. Chem. Gesellsch., 1874, p. 1147; Jahresb. Chem., 1874, p. 1335) note in "5 per cent Soljanka" mineral water and water from an artesian well, both at Ciechocinek, Poland, 0.0035 and 0.0101 part Al₂(HO)₆ per 1000 parts, respectively. A salt obtained from the "5 per cent Soljanka" water was analyzed and found to contain 0.0314 per cent Al₂O₃. Dried at 110° the salt contained 1.005 per cent material which was not soluble in water.

Wroblewsky, E. (Bul. Soc. Chim., 2d ser., 30, p. 436; Ber. Deut. Chem. Gesell., 1878, p. 1848 (corresp.); Jahresb. Chem., 1878, p. 1314), reports 0.02119 part Al₂O₃ per 1000 parts in a spring near Ekaterinodar in the Caucasus.

Wurtz, H. (Silliman's Amer. Jour., 2d ser., 22, pp. 124, 301; Jahresb. Chem., 1857, p. 729), notes a trace of alumina in Delaware River water taken near Trenton, New Jersey.

Zaleski, S. S. (Chem. Ztg., 1892, p. 594; Jahresb. Chem.,

1892, p. 2683), notes o.oo18 part Al₂O₃ per 1000 parts in Ingol Lake water, Gouv. Jenissejsk.

Zäugerle, M. (Neue. Repter. Pharm., 14, p. 9; Chem. Centbl., 1865, p. 798; Jahresber. Chem., 1865, p. 933), reports traces of alumina in the Schillingsforst spring, Bavaria.

ALUMINIUM IN MISCELLANEOUS MATERIALS.

Corney, B. G., D. Guthrie, and F. B. Guthrie (Jour. and Proc. Roy. Soc. New South Wales, 33 (1900), p. 224; Jour. Chem. Soc. London, 78 (1900), II, p. 569) note the presence of 41.53 per cent Al₂O₃ in an edible earth from Fiji. The earth is regarded as a silicate with the formula Al₂O₃(SiO₂)₂(H₂O)₂, with Fe₂O₃ present as an impurity.

Hebberling, M. (Vierteljahressch. Prakt. Pharm., 18, p. 558; Dingler's Poly. Jour., 194, p. 88; Chem. News, 20, p. 249; Jahresb. Chem., 1869, p. 1119), analyzed earth from Berbek, Java, which is eaten by the natives. He found 25.939 per cent alumina.

Love, E. G. (Chem. News, 41, p. 187; Jahresber. Chem., 1880, p. 1118), reports the analysis of a Japanese edible earth. This contained 13.61 per cent alumina.

Muir, M. M. P. (Chem. News, 36, p. 202; Jour. Chem. Soc., London, 34 (1878), II, p. 120), notes 17.97 per cent Al₂O₃ in an edible clay from Mackenzie County (South Island), New Zealand. He states that this clay is largely eaten by sheep.

Schmidt, C. (Ann. Chim. Phys., 4th ser., 26, p. 535; Jour. Chem. Soc. London, 26 (1873), p. 151), in an article on Eatable Earths of Lapland and Southern Persia, notes 40.797 per cent alumina in earth eaten in Lapland. The specimen came from the village of Ponoï. Alumina was not reported in the earth from Kirman (called G'hel i G'iveh) in Southern Persia.

Stark, J. F. (Chem. News, 23, p. 199; Jour. Chem. Soc. London, 24 (1871), p. 425), reports 4.17 per cent Al₂O₃ in an earthball from a horse. The ball weighed over two pounds.

The principal constituent of the earth-ball was ammonium-magnesium phosphate, which amounted to 83.2 per cent.

Thompson, W. Gilman (Practical Dietetics, New York, 1902, p. 281), notes the occasional use of alum and aluminium chloride as preservatives. This use of aluminium compounds has also been noted by other writers.

INDEX.

Abies excelsa, 21. Acer negundo, 11. Achyranthus aspera, 39. Aconite roots, 23, 39. Aconitum heterophyllum, 39. Aconitum japonicum, 39. Acquarossa Spring, Biasca, Italy, 56. Adder's tongue, 10. Adelheidquelle, Heilbrunn, Germany, 71, 114. Adirondacks, New York, spring, 55. Ægina Springs, Greece, 68. Agnesquelle, Lobenstein, Germany, 117. Agrostemma githago, 12. Agua de Vida Lower Spring, Cal., 63. Agua de Vida Upper Spring, Cal., 63. Aix-les-Bains, France, alum spring, 128. Aix-les-Bains, France, sulphur spring, 54, 128. Albany, Ga., artesian well, 118. Alcyone Mineral Springs, Illinois, 108. Alexanderquelle, Psekoup, 122. Alexandro-Jermolowski, Caucasus, spring, 55. Alexinačka banja, Servia, spring, 93. Alfalfa soil, 16. Alfalfa roots, 7. Alga, 16. Alleghany Springs, Virginia, 113, 114. Allouez Mineral Spring, Wisconsin, 68. Alma, Mich., wells, 89. Almond shells, 30. Almonds, 30. Alsophila australis, 11. Alte Quelle, Ronneby, Sweden, 8o. Alum Creek, Vellowstone National Park, 78. Alum Rock Alkaline Saline Spring, California, 63. Alum Rock Chalybeate Spring, California, 63.

I34 INDEX.

Alum Spring, Saratoga, N. Y., 105.

Alveneu Sulphur Spring, 115.

Amandbrunnen, Luhatschowitz, Austria-Hungary, 72, 85.

Amazon River, 117.

Ambrosia artemisiæfolia, 14.

Ambrosius Spring, Marienbad, Austria-Hungary, 76.

Ameliaquelle, Iwoniez, Austria-Hungary, 116.

Amélie d'Hauterève, Hauterève, France, spring, 103.

American Carlsbad Spring, Nashville, Ill., 122.

Americus, Ga., artesian well, 118.

Anatto, 20.

Anatto seeds, 15.

Ancona, Italy, spring, 60.

Andansonia digitata, 19.

Anderson Iron Spring, California, 63.

Anderson Sour Spring, California, 63.

Anderton, England, spring, 102.

Andropogon scoparius, 14.

Angelika, Tönnisstein, Germany, spring, 124.

Angustus Spring, Neuenahr, 56.

Ann Arbor (Huron River?), creek, 88.

Apollinaris Spring, Germany, 54.

Apples, ripe, 8.

Araliarhizomes, 27.

Aralia californica, 27.

Arcier Spring, Besançon, France, 69.

Arctic Spring, Wisconsin, 108.

Arcueil Spring, Paris, France, 69.

Arenaria media, 16, 17.

Arenaria rubra, 16.

Arkansas City Well, Kans., 53.

Arkansas Lithia Springs, 63.

Armeria maritima, 38.

Artemisia Geyser, Yellowstone National Park, 78.

Artemisia maritima, 16.

Artesian Lithia Spring, Ballston Spa, New York, 105.

Artesian Mineral Well, Prairie du Chien, Wis., 109.

Artesian Well, Aberdeen, S. D., 109.

Artesian Well, England, 126.

Artesian Well, Louisville, Ky., 112.

Artesian Well, No. 1, Rockford, Ill., 108.

Arum (Colocasia) esculentum, 18.

Arum maculatum, 31.

Aspidium filix femina, 36.

Aspidium filix mas, 36.

Asta Spring, Vellowstone National Park, 78.

Atchison Electric Light Well, Kans., 53.

Atchison, Kans., spring, 86.

Atchison Parker's Spring, Kansas, 53.

Atherosperma moschatum, 48.

Atlantic coastal plain, United States, artesian wells, 68.

Atropa belladonna, 27.

Aube River, Etufs, France, 103.

Auditorium, Chicago, Ill., artesian well, 91.

Augenquelle, Elster, Germany, 122.

Austin's Springs, Tennessee, 107.

Australasian mineral waters, 94, 95.

Auvergne, France, springs, 128.

Azémar Spring, Bagnères-de-Luchon, France, 73.

Bachquelle, Teinach, Germany, 72. Bad Griesbach, Germany, spring, 118. Bad Jeni-Kaplidja, Anatolia, Asia Minor, spring, 118. Bad Marienborn, spring, 121. Bad Wendig, Ries, spring, 124. Bademli Baghtsche, Brusa, Asia Minor, spring, 121. Baden-Baden, Germany, spring, 101. Badequelle, Paderborn, Germany, 59. B. B. Mineral Spring, Mo., 66. Bagnères-de-Bigorre, spring, France, 60. Bain Romain, Plombières, France, 82. Bain tempère, Plombières, France, 82. Banana-tree sap, 18. Banja Visegrad, Bosnia, 97. Baobab fibers, 19. Barley, 45. Barley sprouts, 23. Barragan, New South Wales, spring, 99. Barringun, New South Wales, artesian well, 100. Bat manure, 50. Bath Alum Spring, No. 1, Virginia, 106. Bath Alum Spring, No. 2, Virginia, 106, 110. Bath Alum Spring, No. 3, Virginia, 113. Bayard, Hauterive (Allier), France, spring, 103. Bayen Spring, Bagnères-de-Luchon, France, 73. Beans, 34. Bear River Hot Springs, Utah, 109. Beaupréau Spring, France, 52. Beck's Hot Spring, Salt Lake City, Utah, 118. Bed straw, 38. Bedford Bowling Alley Spring, Pennsylvania, 67. Bedford Iron and Alum Springs, Virginia, 110. Bedford Magnesia Spring, Pennsylvania, 67. Beech, 40, 41.

Beef, 50.

Beehive Geyser, Yellowstone National Park, 78.

Beet sugar, 25.

Beets, 14.

Belabula River, New South Wales, 99.

Belladonna leaves, 23, 27.

Bellety Steel Spring, Sail-lès Chateaumorand, France, 81.

Belvedra Spring, Chur, Switzerland, 83.

Bench Spring, Yellowstone National Park, 78.

Bergstrasse Stahlquelle, Weinheim, Germany, 101.

Bermuda-grass roots, 7.

Bernhard Spring, Krankenheil-Tölz, Germany, 74.

Bethesda Springs, Waukesha, Wis., 108.

Betula alba, 41.

Big Bone Springs, Kentucky, 65, 127.

Billecul Spring, Besançon, France, 69.

Bimbia, Africa, spring, 116.

Birch, 40,

Birch fungus, 43.

Birch sap, 26.

Birchdale Spring (Concord Spring), New Hampshire, 105.

Birken schwamms, 16, 43.

Birresborn, Pelm, Germany, spring, 125.

Bitter Wasser, Ofen, Germany, 54.

Bitterwasser, Weilutza, Roumania, 87.

Bixa orellana, 15.

Black Earth Mineral Springs, Wisconsin, 108.

Black thorn, 32.

Black truffle, 29.

Blanche, Bagnères-de-Luchon, France, spring, 73.

Bled et Amax, Sahara, Africa, well, 87.

Blèville, France, spring, 98.

Blocksberg Mountains, Germany, spring, 54.

Blodgett's Springs, California, 63.

Bloodroot rhizome, 23.

Blossburg Spring, Pennsylvania, 110.

Blueberries, dried, 21.

Blue Lick Springs, Mo., 66.

Blue Ridge Springs, Virginia, 67, 106.

Blue thistle, 14.

Boiler Bath (Old Ladies' Boiler Bath Hot Spring), Virginia, 106.

Boletus edulis, 15.

Bologna, Italy, spring, 60.

Bolschoi, Slawinsk, Poland, spring, 102.

Bondonneau, France, spring, 81.

Bonifaciusquelle, Neuhaus, Germany, 92.

Bouillants Spring, Vergéze, France, 58.

Borax Lake, California, 55.

Bordeu No. 4 Spring, Bagnereres-de-Luchon, France, 73.

"Borhegyer Sauerwasser," Austria-Hungary, 54.

Borland Mineral Well, West Virginia, 113.

Bormio, Italy, spring, 114.

Bosquet, Bagnères-de-Luchon, France, spring, 129.

Bourke Corrella Station, New South Wales, artesian well, 99.

Bourke, New South Wales, artesian well, 99.

Bovista gigantæa, 27.

Bowden Lithia Springs, Georgia, 65.

Bowsher Mineral Spring, Missouri, 109.

Box elder leaves, II.

Bradford Mineral Spring, New Hampshire, 66.

Bradford, Mass., wells, 70.

Bratton Spring, Missouri, 113.

Brayera anthelmintica, 17.

Bread, 3, 4, 5, 39, 45.

Bregille Spring, Besançon, France, 69.

Brestowačka banja, Servia, 93.

Briscous Salt Wells, France, 59.

Bristol, Pa., spring, 76.

Bromus secalinus, 29.

Broom sedge, 14.

Brucourt, France, spring, 61.

Bryony root, 18.

Brodelbrunnen, Pyrmont, Germany, 74.

Broken Hill Brewery, New South Wales, well, 99.

Broken Hill Quarry, New South Wales, artesian well, 99.

Bromo-magnesium Well, Kansas, 53.

Bronislaw Spring, Truskawice, Galicia, 80.

Brown's Wells, Springs, Mississippi, 66.

Brühquelle, Baden, Germany, 59.

Brünnlein (Urquelle), Trentschin Töplitzer Bad, Austria-Hungary, 89.

Brünthaler, 129.

Brüx. Austria-Hungary, artesian well, 120.

Buckhert's Fountain, Wisconsin, 108.

Buckwheat, 45.

Budös (Bálványos), Siebenbürgen, spring, 96.

Büdös (Stinkberg), Bosnia, spring, 96.

Buffalo Lithia Springs, Virginia, 110.

Bukowik, Servia, spring, 93.

Bullicame, Viterbo, Italy, springs, 116.

Burdett Mineral Wells, Texas, 67.

Burge Malo, spring, 122.

Burr Oak Spring, Kansas, 53.

Burtscheid, Germany, spring, 128.

Bushnell Railroad Well, Illinois, 91.

Butterworth's Magnetic Spring, Grand Rapids, Mich., 88, 108. Buxton, England, spring, 123.

Cacao, o. Caledonia Springs, Ontario, 127. California Geysers, Acid Spring, 63. California Geysers, Alum Spring, 63. California Geysers, Devil's Teakettle, 63. California Geysers, Hot Sulphur Springs, 63. California Geysers, Indian Spring, 63. California Geysers, Iron Spring, 63. California Geysers, Iron Spring, Geyser Creek, 63. California Geysers, Lemonade Spring, 63. California Geysers, near Indian Spring, 63. California Geysers, Mud Indian Spring, 63. California Geyser Spring, near river, 63. California Geysers, Witches' Caldron, 63. California Geysers, Seltzer Spring, 127. Calisaya bark, 23. Calistoga Springs, California, 63. Calistoga Swimming Pool, California, 63. Calluna vulgaris, 41. Calumba roots, 23. Camborne, Cornwall, England, spring, 114. Cane sugar, 25, 38. Cape Town, Africa, spring, 8o. Capon Beauty Spring, West Virginia, 106. Capon Main Spring, West Virginia, 106. Capioma Mineral Well, Kansas, 53. Capsella bursa pastoris, 13. Capsicum, 7. Carabana, Spain, spring, 54. Carbondale Spring, Kansas, 53. Cardamon, 17. Cardamon seeds and husks, 44. Carica papaya, 28. Carlsquelle, Bad Helmstedt, Germany, 75. Carlsquelle, Iwoniez, Austria-Hungary, 116. Cäsarbad, France, spring, 90. Cascara bark, 23. Cass River, Michigan, 88. Castalia, Ohio, spring, 6o. Castanea leaves, 23. Casteggio, Italy, spring, 69. Castilian Mineral Spring, Mississippi, 66. Catoosa Springs, Georgia, 111, 112.

Cave Springs, Virginia, 110.

Cawker City, Kans., spring, 54.

Caxton, Canada, springs, 83, 127.

Ceanothus americanus, 9.

Cedar Springs, Washington Spring, Ohio, 107.

Centralia Gypsum Well, Kansas, 53.

Centralia, Kans., well, 86.

Ceratonia siliqua, 30.

Cercle, Rennes-les-Bains, France, spring, 129.

Ceresole Reale, Italy, spring, 100.

Cervena-Rjeka-Quelle, Srebrenica, Bosnia, 96.

Cetraria islandica, 41.

Cetrarid, 16.

Ceyssat, France, spring, 124.

Chadnor Villa, Cheltenham, England, well, 124.

Chalybeate Springs, Georgia, 112.

Champion Spouting Spring, Saratoga, N. Y., 105.

Charleston, S. C., artesian well, 121.

Chatelguyon, Puy-de-Dôme, France, spring, 103.

Chaumaix, France, spring, 90.

Cheken leaves, 19.

Cherokee City Well, Kans., 53.

Cherokee Magnetic Mineral Spring, Iowa, 109.

Cherries, 21.

Cherry Valley Phosphate Spring, New York, 105.

Cheshire, England, springs, 102.

Chestnuts, 13.

Chickweed, 16, 17.

Chimaphila leaves, 23.

China flava fibrosa, 30.

China huanuco, 30.

China regia cum epid., 30.

China regia sine epid., 30.

China rubra, 30.

"Chine," 17.

Chippewa River, Michigan, 88.

Chittenango Cave Spring, New York, 105.

Chittenango White Sulphur Springs, New York, 105.

Chlorangium jussuffii, 16, 22, 29.

Choke-cherry bark, 23.

Chrome Spring, Yellowstone National Park, 78.

Chur, Switzerland, springs, 83.

Church Hill Alum Springs, Virginia, 111.

Cicuta maculata, 16.

Cider vinegar, 14.

Ciechocinek, Poland, artesian well, 129.

Cinchona, 19, 23.

Cinchona bark, 25.

Cinchona officinali, 19.

Cinchona succirubra, 19.

Cinnamon bark, 32.

City Creek, Utah, 61.

Civillina, Italy, spring, 56.

Cladonia, 16.

Cladonia rangiferina, 41.

Cladophora glomerata, 16, 29.

Clark's Riverside Mineral Springs, Detroit, Mich., 66, 89.

Clay, edible, from New Zealand, 131.

Cleopatra Spring, Yellowstone National Park, 77.

Cliff Spring, Saint-Nectaire le Haut, France, 76.

Clifton Spring, No. 1, Virginia, 106.

Climax Springs, Missouri, 109.

Clinton, Miss., well, 61.

Club moss, I, 10, 12, 23, 31.

Cochineal, 49.

Cockle-burr, 17.

Cocoa shells, 9.

Coëse, France, spring, 101.

Coffee beans, 25.

Coffee tree, 25.

Colchicum, 30.

Colchicum autumnale, 30.

Cold Spring III, Neuenahr, 56.

Cold Sulphur Springs, Virginia, 67, 111.

Coldwater, Mich., well, 89.

Columbus Well, Kansas, 53.

Coltsfoot, leaves, 8.

Columbia Chalybeate Spring, Missouri, 109.

Conferva glomerata, 27.

Congress Spring, Saratoga, N. Y., 105.

Conneautville, Pa., artesian well, 118.

Constant Geyser, Yellowstone National Park, 77.

Constantine Spring, Gleichenberg, Austria-Hungary, 97.

Convolvulus batatas, 18.

Convolvulus roots, 7.

Conway Springs, Kans., 53.

Cooper's Well, Mississippi, 112.

Coral Spring, Yellowstone National Park, 77.

Corella, New South Wales, artesian well, 100.

Corncobs, 2.

Corn cockle, 12.

Corn kernels, 33.

Corneliusquelle, Aachen, Germany, 92.

Corydalis, 16.

Corydalis bulbosa, 16

Costalta, Southern Tyrol, spring, 69. Cattani Spring, No. 2, Spolato, 125. Cow's milk, 49. Crenothrix manganifera, 20. Crenothrix kühniana, 20. Crenothrix ochracea, 20. Cresson Alum Spring, Pennsylvania, 110. Cresson Iron Spring, Pennsylvania, 110. Cresson Magnesia Spring, Pennsylvania, 106. Crisp Springs, Tennessee, 107. Črni Guber, Srebrenica, Bosnia, spring, 97-Crocket's Arsenic-Lithia Springs, Virginia, 67. Crocus, autumn. (See Colchicum.) Crystal Springs, Saratoga, N. V., 105. Cuckoo-pint leaves, 31. Cullum's Ferruginous Spring, Alabama, 114. Curry Spring, Versailles, Ill., 127. Cuttaburra, New South Wales, artesian well, 99, 100. Cuyahoga Lithia and Magnesium Springs, Ohio, 112, 114. Cyathea serra, 10. Cynanchum monspeliacum, 20. Czeméte, Austria-Hungary, spring, 120. Czifra-viz, spring, 73.

Dalton, Mass., artesian well, 71. Damont, Rennes-les-Bains, France, spring, 129. Dandelion, 32, 33, 40. Davenport, Ia., glucose factory, artesian well, 91. Davis Mineral Well, Kentucky, 112. Dead Sea, 54, 119. Deer Lick Spring, New York, 66. Dekalb, Ill., water-works, artesian well, 91. Delaware River, New Jersey, 129. Deloraine, Manitoba, artesian well, 126. Des demoiselles, Rennes-les-Bains, France, spring, 129. Detroit River, Michigan, 87, 88. Devil's Ink Pot, Yellowstone National Park, 78. Dianthus caryophyllus, 1. Dicksonia squarrosa, II. Digitalis leaves, 23. Dintenquelle, Teinach, Germany, 72. Director Spring, Starja-Russa, 102. Dixie Mineral Spring, Tennessee, 67. Dixon, Ill., water-works, artesian well, 91. Dixon Spring, Kansas, 53. Dock, 14. Dog-bane, 20.

Dolnj-Tuzla, Bosnia, springs, 96.
Donatusquelle, Albulathale, 115.
Donionidi, Sahara, Africa, well, 87.
Dorna-Sara Arsenic Spring, Roumania, 116.
Doubs River, France, 56, 69.
Doulaux, France, spring, 90.
Dougherty's Carbonic Acid Spring, British Columbia, 86.
Driburg, Germany, spring, 127.
Dropping Well, Knaresborough, England, 59.
Dubois (Allier), Vichy, France, spring, 103.
Dudley, England, wells, 124.
Dulimbert Spring, Vergèze, France, 55.
Düngle Ridge, New South Wales, artesian well, 100.
Durkheim, Germany, spring, 59.
Dwarf pine, 41.

Earth ball from a horse, 131. Earth, edible, from Berbek, Java, 131. Earth, edible, from Fiji, 131. Earth, edible, from Japan, 131. Earth, edible, from Ponoi, Lapland, 131. Eastern Bohemia, mineral waters, 85. Eastman's Springs, Michigan, 66. East Saginaw, Mich., well, 89. East Spring, Joplin, Mo., 82. East Spring, Langenbruck, Austria-Hungary, 77. Echium vulgare, 14. Echinus Spring, Yellowstone National Park, 77. Eckholzquelle, Ronneby, Sweden, 80. Eddoes. (See Taro.) Eel grass, 2, 16. Egg albumen, 50. Einsiedel, Austria-Hungary, spring, 98. Eisenquelle von Uebersaxen, Voralberg, Austria-Hungary, 87. Eisenwasser Bad Andelsbuch, Voralberg, Austria-Hungary, 87. Eisenwasser des Bad Reuthe, Voralberg, Austria-Hungary, 87. Ekaterinoder, in the Caucasus, spring, 129. Elder, 19, 42. Eleusine indica, 14.

El Feter, Sahara, Africa, well, 87.

El-Hadjira, Sahara, Africa, well, 87.

Elizabethquelle, Carlsbad, Austria-Hungary, 97.

Elizabethenquelle, Neuhaus, Germany, 92.

Elizabethenquelle, Rothenfels, Germany, 58.

Elm bark, 23.

El Paso de Robles Hot Sulphur Spring, California, 110.

El Paso de Robles Main Sulphur Spring, California, 63.

El Paso de Robles Soda Spring, California, 63. Emmaquelle, Gliechenberg, Austria-Hungary, 79. Emperador, Panama, well, 51. Empire Mine, swallet, 54. Empire Spring, Saratoga, N. Y., 60, 105. Enclos des Célestins, Vichy, France, spring, 90. English ivy, 7. English walnut bark, 35. Equisetum maximum, 10. Equisetum telmateja, 26. Ergot, 18, 29, 37. Erica vulgaris, 16. Estill Springs, Black Sulphur Spring, Kentucky, 112. Estill Springs, White Sulphur Springs, Kentucky, 107. Etufs, France, spring, 103. Eucalyptus, 34. Eucalyptus globulus, 34. Eucalyptus rostrata, 34. Euphorbia amygdoloides, 42. Eureka Mineral Well, Kansas, 53. Eureka Springs, California, 63. Eureka Spring, Saratoga, N. Y., 105. Eureka Springs, Wisconsin, 108. Everett Spring, Massachusetts, 70. Excelsior Geyser, Yellowstone National Park, 78. Excelsior Springs, Missouri, 54, 66, 99, 127. Excelsior Well, Benton Harbor, Mich., 89.

Faculté des Sciences, Besançon, France, well, 69. Fagus sylvatica, 41. Fairview Springs, Texas, 112. Farmville Lithia Springs, No. 2, Virginia, 106. Farmwell Station, Va., springs, 118. Farnham, England, spring, 79. Fearless Geyser, Yellowstone National Park, 77. Felsenquelle, Bad Ems, Germany, 75, 122. Felsenquelle, Carlsbad, Austria-Hungary, 97. Felsenquelle, near Bassen, Austria-Hungary, 73. Felt's Mineral Springs, California, 63. Ferdinand Spring, Marienbad, Austria-Hungary, 76. Fern root-stalks, 34. Fern roots, 36. Ferrara, Italy, spring, 60. Ferras Spring, Bagnères-de-Luchon, France, 73. Fidelisquelle, Bosnia, 96. Fig leaves, 42. Fir bark, dry, 22.

Fir cones, dry, 22. Fir moss, 22. Fir needle litter, 22, 39. Fir needles, 21, 22. Fir twigs, 22. Firehole River, Yellowstone National Park, 78. Fitzpatrick, Ala., artesian well, 118. Flat Rock Spring, Saratoga, N. Y., 105. Flax stems, 20. Flitwick Moor, spring, 85. Floating buttercup, 32. Flour, 2, 3, 14, 28, 38, 46. Flowing Spring, Kansas, 109. Fojnica, Bosnia, spring, 97. Fontinalis antipyretica, 10, 36. Fort Crawford Mineral Spring, Wisconsin, 68. Fort Scott Sulpho-magnesian Well, Kansas, 53. Fountain Geyser, Yellowstone National Park, 68, 78. Fountain Springs, Wisconsin, 108. Framingham, Massachusetts, springs, 70. Frangula bark, 23. Franklin Artesian Well, Ballston Spa, N. Y., 105. Franklin Spring, Georgia, 65. Franz-Joseph-Bad, Tüffer, Austria-Hungary, 127. Franz Joseph Quelle, Füred, Austria-Hungary, 81. Fraysse, Cransac, France, spring, 81. French Lick Springs, Pluto's Well, Indiana, 112. Friedensquelle, Rostenberg, Germany, 98. Friedrichshall, Germany, spring, 54. Friedrichshall, Germany, spring, 92. Friedrichsquelle, Zeidelweid, Austria-Hungary, 85. Fry's Mineral Spring, Iowa, 65. Fulbrunnen, Wiesbaden, Germany, spring, 114. Fungus, edible, 27. Fürstenbrunnen, Ems, Germany, 74, 75.

Galbraith's Spring, Tennessee, 107.
Galdhof, Seelowitz, Austria-Hungary, spring, 102.
Galena, Ill., water-works, artesian well, 91.
Galium mollugo, 38.
Garden pink, 1.
Garden rose, 1.
Gardiner River, Yellowstone National Park, 77.
Gasquelle, Elster, Germany, 122.
Gastrolobium bilobum, 15.
Gata, Bosnia, spring, 96.
Gaylord and Gulick's Mineral Spring, Pennsylvania, 67, 110.
Geilnau, Nassau, Germany, spring, 74.

Gelsemium root, 23.

Geneseo, Ill., water-works, artesian well, 91.

Geneva Lithia Spring, New York, 66.

Georgenbrunnen, Bad Eilsen, Germany, 76.

Geranium rhizome, 23.

Gerlosteiner Schlossbrunnen, Pelm, Germany, 125.

Germania Spring, Schwalheim, Germany, 123.

Gettysburg Lithia Spring, Pennsylvania, 106.

Geuda Springs, Kansas, 53, 65.

Geyser Mine, Custer County, Colorado, 71.

Geyser Mineral Well, Kansas, 53.

Giantess Geyser, Yellowstone National Park, 78.

Gibson Springs, Wisconsin, 108.

Giesshübler, Rodisfort, Germany, spring, 78.

Glacier Spouting Spring, Saratoga, N. Y., 60, 119.

Glen Alpine Springs, California, 63.

Glen Flora Springs, Illinois, 108.

Glen Springs, Wisconsin, 108.

Glenola (Wayland) Spring, Virginia, 67.

Glycyrrhiza root, 23.

Gmunden Lake, Austria-Hungary, 77.

Golden seal root, 16.

Gombert's Well, Wisconsin, 108.

Gordon Springs, California, 63.

Gorni Sehar, Bosnia, spring, 96.

Gosling's Garden, Osnabrück, artesian well, 86.

Gradačac, Bosnia, spring, 96.

Grand Haven Mineral Spring, Michigan, 108.

Grand River, Jackson, Mich., 88.

Grand Rue Spring, Besançon, France, 56, 69.

Grande Grille, Vichy, France, spring, 103.

Grande Source, Pougues-les-Eaux, Nièvre, France, 103.

Grande Source, Vittel, France, 100.

Granier Spring, Vergèze, France, 55.

Grape branches, 31.

Grape juice, 6, 27.

Grape leaves, 31, 38.

Grape must, 31.

Grape seeds, 16.

Grape wood, 27.

Grapes, 24, 26.

Grapevines, 17, 30, 38.

Grassion Spring, Clermont, France, 103.

Gravesend, England, spring, 79.

Great Bend Mineral Well, Kansas, 53.

Great Fountain Geyser, Yellowstone National Park, 78.

Great Paraiso Hot Soda Springs, California, 64.

Great Salt Lake, Utah, 102. Great Spirit Spring, No. 2, Kansas, 53. Greenbrier White Sulphur Springs, West Virginia, 106, 111. Green Mineral Spring, Ohio, 107. Greensborough Spring, North Carolina, 107. Grevillea robusta, 26. Grosslüder Spring, Salzschlirf, Germany, 117. Grotte inférieure, Bagnères-de-Luchon, France, spring, 73. Grotte supérieure, Bagnères-de-Luchon, France, spring, 73. Grotts Geyser, Yellowstone National Park, 78. Guarana paste, 28. Guarana seed, 23, 28. Guarana shells, 28. Gueuzayasma, Brusa, Asia Minor, spring, 121. Guillot Spring, Evian, France, 101. Gyrenbad, Switzerland, spring, 124. Gyrophora, 16 Gyrophora pustulata, 22, 41.

Hacket's Spring, Wisconsin, 108. Haddon Mineral Well, Kansas, 109. Hall, Austria-Hungary, spring, 101. Halowell Grant Spring, Nova Scotia, 51. Harbin Chalybeate Spring, California, 63. Harbin Hot Sulphur Spring, California, 63. Hart Well, West Virginia, 68, 113. Hartford Cold Springs, Maine, 105. Hartford, Conn., spring, 58. Harris Antidyspeptic and Tonic Springs, Virginia, 67. Hartsel Hot Mineral Springs, Colorado, 109. Hartwegia comosa, 43. Hathorn Spring, Saratoga, N. Y., 105. Hauptbrunnen, Münster, Germany, 124. Hauptquelle, Gastein, Austria-Hungary, 97. Hauptquelle, Neuhaus, Austria-Hungary, 79. Haupt-Trinkquelle, Stuttgart, Germany, 72. H. Dahne, Sahara, Africa, well, 87. H. Debiche et Strifigi, Sahara, Africa, well, 87. Heather, 16, 41, 43. Heilbrunnen, near Lake Taach, 55. Helianthus annuus, 43. Helicon Springs, Georgia, 107. Hellibore rhizome rootlets, 23. Hemlock bark, 2. Hemp, refuse, 8. Hermannsquelle, Neuhaus, Germany, 92. Herniaria glabra, 43.

Herster, Driburg, Germany, spring, 74. Hertfell Spring, Scotland, 85. Hickman's Springs, Kentucky, 112. Hides, 50. High Rock Spring, Saratoga, N. Y., 105.

Highland Dutch or Ems Spring, California, 64, 110.

Highland Magic Spring, California, 110.

Highland Seltzer Spring, California, 64, 110.

Hildegarde Spring, Ofen, Austria-Hungary, 119.

Hindhead, England, springs, 79.

Hirschquelle, Teinach, Germany, 72.

H. Messaoud, Sahara, Africa, well, 87.

Höhenbrunnerfilz, 129.

Holly leaves, 30.

Homburg, Germany, artesian well, 82.

Honey, 9.

Honeysuckle berries, 15.

Hop blossoms, 39.

Hops, 9, 23, 40.

Horeb Springs, Wisconsin, 108.

Horse bean, 18.

Horse chestnut, 35.

Horsetail, 10, 26.

Hosea Saline Sulphur Spring, Indiana, 113.

Hot Borate Spring, California, 64.

Hot River, Yellowstone National Park, 77.

Hot Spout Bath, Virginia, 106.

Hot Spring, Savo Island, 93.

Hot Springs, Arkansas, 107.

Hot Springs, Hot Springs Station, Nevada, 61, 113.

Hot Springs, Monte Irone, Albane, Italy, 101.

Howard Springs, California, 64.

Hroswitha, Herzogludolfsbad, Germany, spring, 102.

Hughes' Mineral Well, Georgia, 65.

Human milk, 49.

Humboldt River, Nevada, 60, 119.

Humboldt's Salt Well, Minnesota, 109.

Hunter's Hot Spring, Montana, 109.

Hunter's (Pulaski) Alum Springs, Virginia, 67.

Hurricane Springs, Tennessee, 107.

Hyacinth, 38.

Hyacinthus orientalis, 38.

Hyde Park, Mass., artesian wells, 70.

Hyde Park, Mass., wells, 70.

Hydrangea, 8, 23.

Hydrangea arborescens, 8.

Hydrastis canadensis, 16.

Hydrastis rhizome, 23. Hygeia Springs, Wisconsin, 108. Hygeia Spring, Yellowstone National Park, 78. Hyoscyamus leaves, 23.

Iceland moss, 41. Ida Spring, Biloves, Austria-Hungary, 101. Idjen volcano, Java, lake, 73. Ilex aquifolium, 30. Ilidze, Bosnia, spring, 96. Illicium anisatum, 32. Indian fig. (See Prickly pear.) Indan-Ha Spring, Idaho, 65. Indian Medical Spring, Minnesota, 66. Indian Spring, Indiana, 65, 113. Ingol, Lake, Russia, 130. Iodine Donatus, Solis, 115. Iodo-Magnesian Springs, Wisconsin, 108. Iowa Acid Spring, 109. Irondale Springs, West Virginia, 111. Iron Lithia Springs, Virginia, 68. Iron Spring, Milford, N. H., 105. Iron-Sulphur Spring, Sail-lès-Chateaumorand, France, 81. Isar River, Germany, 36. Isère River, France, 79.

Jacob's Artesian Well, Wisconsin, 108. Jalap, 23. James River, Richmond, Va., 118. Jamnicer-alkalisch-muriatischen Säuerlings, 84. Jarvisville, New South Wales, spring, 99. Jeanne d'Arc, Pougues-les-Eaux, Nièvre, France, spring, 103. Jeanne d'Arc (Allier), Vichy, France, spring. 103. Jenkins's Quarry, New South Wales, spring, 99. Jenolan Caves, New South Wales, spring, 99. Jenzat, France, springs, 90. Jerseyville, Ill., waterworks, artesian well, 91. Johann-Georgen Spring, Krankenheil-Tölz, Germany, 74. Johannis Spring, Stainz, Austria-Hungary, 117. Johannes Springs, Steiermark, Austria-Hungary, 79. Johannesbad, Vienna, Austria-Hungary, spring, 82. Johannbrunnen, Luhatschowitz, Austria-Hungary, 72, 85. Johannisbrunnen, Mahren, Germany, 95. Johnson's Wells, Alabama, 107. Jordan AlumSprings, Virginia, 106, 111. Jordan White Sulphur Springs, Virginia, 106. Jordan's Mineral Well, Wisconsin, 108.

149

Josefsquelle, Bilin, Austria-Hungary, 83. Josephsquelle, Rippoldsau, 128. Judenquelle, Baden, Germany, 59. Julianenbrunnen, Bad Eilsen, Germany, 76. Jute fiber, 7.

Kaba Mountain, East India, spring, 98.

Kaiserbrunnen, Carlsbad, Austria-Hungary, 97.

Kaiserquelle, Aachen, Germany, 92.

Kalmia latifolia, 26.

Kamenewquelle, Psekoup, 122.

Kane County Magnesia Spring, Illinois, 65.

Kane Sulphur Spring, Pennsylvania, 106.

Kantavu, Fiji Islands, spring, 93.

Karlsbad Sprudel, Austria-Hungary, 78.

Karlsquelle, Bosnia, 96.

Karmalinquelle, Psekoup, 122.

Kärtner Römerquelle, Prevali, Austria-Hungary, 86.

Kasimer Slawinsk, Poland, spring, 102.

Katharien-Bad, Kärnthen, Austria-Hungary, 100.

Kava root, 17.

Kellberg, Germany, spring, 61.

Kelly's Camp, New South Wales, artesian well, 99.

Kentucky Alum Springs, 112.

Kesselbrunnen, Ems, Germany, 74, 75.

Keystone Spring, Maine, 65.

Kickapoo Springs, Kansas, 53.

Kidwell's Quarry, New South Wales, spring, 99.

King David Spring, Benton Harbor, Mich., 88.

Kingswood, England, spring, 52.

Kirchdrauf in der Zips, Austria-Hungary, spring, 120.

Kirouars, France, spring, 57.

Kittanning Mineral Spring, Pennsylvania, 110.

Klebelsberg Spring, Ischl, Austria-Hungary, 69.

Kleininger Sauerbrunnen, Kärnthen, Austria Hungary, 100.

Klosteralleequelle, Pyrmont, Germany, 75.

Kochbrunnen, Wiesbaden, Germany, 74. Koeningan, East India, spring, 98.

Kola-nut, 23.

Kolop Sulphur Spring, Austria-Hungary, 91.

Kondran, Germany, spring, 124.

König Ferdinand Eisenbad, Weidritzthal, Austria-Hungary, 127.

Königsbrunnen, Krostreinitz, Austria-Hungary, 79.

Kostreiniz, Steiermark, Austria-Hungary, spring, 82.

Kousso, 17.

Krähnchen, Ems, Germany, 74, 75.

Kronenquelle, Salzbrunn, Germany, 116.

I 50 INDEX.

Krynica, Austria-Hungary, spring, 69. Kukurtlu, Brusa, Asia Minor, spring, 121. Kulasi, Bosnia, spring, 96. Kuttawa Mineral Springs, Kentucky, 107.

La Bernerie, France, spring, 57. La Reine Spring, Bagnères-de-Luchon, France, 73. Labrador tea, 2. Lacquer (Urushi), 44. Lagrange, Mo., Wyaconda, artesian well, 91. Lakatah Springs, South Dakota, 67. Lake Derkol, Turkey, 122. Lake Goguac, Michigan, 88. Lake Goodenough, British Columbia, 126. Lake Michigan, Chicago, Ill., 89. Lake Superior, Michigan, 88. Lamscheider, Mineralbrunnen, Germany, 75. Landreth's Mineral Well, Missouri, 100. Lane Mineral Springs, California, 64. Lansdowne Well, Cheltenham, England, 124. Larch needles, 20. Las Vegas Hot Spring, New Mexico, 66. Lauderdale Springs, Mississippi, 107. Laurel root, 26. Lawrence, Mass., sewage, 70. Leather, 49. Lebanon, Mo., artesian well, 71. Lebanon Thermal Spring, New York, 105. Ledum palustre, 2. Lee County, Virginia, spring, 121. "Lěgèn," 40. Leontodon taraxacum, 40. Leopoldsquelle, Rippoldsau, 128. Lichen, 41. Liebenzell, Württemberg, Germany, 72. Lilac flowers and leaves, 42. Linden leaves, 7. Lindenbrunnen, Zlatten, Austria-Hungary, 58. Lincoln Springs, Kansas, 53. Lineville Mineral Springs, Iowa, 65. Lippa, Austria Hungary, spring, 119. Lithium Spring, Omio, Kans., 53. Litton Seltzer Springs, California, 64. Llangammarch, spring, 86. Ločendol, Austria-Hungary, well and spring, 84. Loch Katrine, Scotland, 126. Loire River, Orleans, France, 69.

London, England, water supply, 79. Londonderry Lithia Spring, New Hampshire, 66. Long Island, southern shore, wells, 84. Lonicera xylosteum, 15. Lorain, Ohio, artesian well, 93. Los Angeles, Cal., river, 119. Louisenquelle, Luhatschowitz, Austria-Hungary, 72, 85. Lupine leaves, 7. Luxeuil Springs, France, 58, 70. Lycopodium, 43. Lycopodium powder, 23. Lycopodium alpinum, 10, 29. Lycopodium annotinum, 12. Lycopodium billardieri, 10. Lycopodium cernuum, 10. Lycopodium chamaecyparissus, 1, 16, 29, 31. Lycopodium clavatum, 1, 10, 16, 23, 31. Lycopodium complanatum, 26, 31. Lycopodium denticulatum, 34. Lycopodinm phlegmaria, 10, 29. Lycopodium selago, 10.

Macomb, Ill., water-works, artesian well, 91. Madeleine Spring, No. 1, Rennes-les-Bains, France, 129. Madeleine Spring, No. 2, Rennes-les-Bains, France, 129. Maesa picta, 1. Magdalen Spring, Cayla, France, 93. Magnetic Mineral Spring, Indiana, 65. Magnetic Spring No. 2, Riverside, Mich., 108. Mala Kiselica, Srebrenica, Bosnia, spring, 97. Malden, Mass., tubular wells, 70. Mammoth Hot Springs, Vellowstone National Park, 77. Mandarin orange, 30, 31. Manistee Lake, Michigan, 88. Manitou Soda Springs, Colorado, 64, 65. Mansfield, Mass., well, 70. Marblehead, Mass., Wells, 70. Marblehead, Water Company, Swampscott, Mass., wells, 70. Mardela Spring, Maryland, 65. Marden's quarry, New South Wales, spring, 99. Maria Island, spring, 90. Maria Theresia Quelle, Andersdorf, Austria-Hungary, 96. Marien Sprudel Spring, 56. Marienbrunnen, Gabernegg, Austria-Hungary, 78. Marienquelle, Johannisbrunn, Austria-Hungary, 97. Marienquelle, Neuhaus, Germany, 92. Marienquelle, Paderborn, Germany, 59.

Marienquelle, Psekoup, 122. Marine Algæ, 27, 29. Marion Artesian Well, Indiana, 108. Mark West Springs, California, 64. Marktbrunnen, Carlsbad, Austria-Hungary, 97. Marktsoole, Colberg, Germany, 55. Marsilea quadrifoliata, 10. Marston, England, spring, 102. Massanetta Mineral Springs, Virginia, 68, 106. Matchless Mineral Wells, Alabama, 63. Mattigbad, Austria-Hungary, spring, 89. Maxbrunnen, Kissengen, Germany, 92. McDuff's Spring, Kansas, 53. Mechanicsville, N. Y., spring, 79. Medical Lake, Washington, 110. Medicis Spring, Saint-Denis, France, 81. Melle, Germany, artesian well, 123. Melle, Germany, spring, 123, Melrose, Scotland, spring, 69. Mergentheim, Germany, spring, 92, Meriweather County Warm Springs, Georgia, 65. Merrill Spring, Kansas, 53. Meurchin, Pas-de-Calais, France, spring, 78. Michailquelle, Psekoup, 122. Midland Magnetic Well, Michigan, 113. Midland Mineral Spring, Michigan, 88. Middle or Dewdrop Spring, Greencastle, Ind., 108. Middletown Springs, No. 1, Vermont, 105. Midway Warm Springs, Utah, 67. Milboro Sulphur Spring, Virginia, 68. Millet, 45. Millingtonia hortensis, 19. Mineral Spring, No. 1, Versailles, Ill., 108. Mineral Springs, Farmwell Station, Va., 106. Mineral Springs, Hanover, Pa., 106. Mineral Well, Pierre, S. Dak., 109. Minnequa Springs, Pennsylvania, 67. Mississippi River water, 52, 116. Mitterbad, Tirol, spring, 55. Modling, Austria-Hungary, spring, 120. Mofetta di S. Quirico, Tuscany, Italy, spring, 102. Monmouth, Ill., water-works, artesian well, 91. Mono Basin, California, spring, 61. Mono Basin Warm Springs, California, 110. Mono Lake Spring, California, 64. Monterey, Mexico, spring, 89. Montezuma, Ga., artesian well, 118. Montezuma, Ind., artesian well, 91.

Montpellier, France, spring, 100. Montvale Springs, Tennessee, 107. Morchella esculenta, 16, 29. Morel, 16, 29. Moresville Mineral Springs, Missouri, 113. Moriz Spring, Rohitsch, Austria-Hungary, 58. Morphine, 21. Moss, 10, 43. Moss Springs Well, Kansas, 53. Mouillère, Besançon, France, spring, 69. Mountain raspberry, 7. Mount Clemens Clementine Well, Michigan, 89. Mount Clemens Media Spring, Michigan, 108. Mount Clemens Media Well, Michigan, 89. Mount Clemens Mineral Well, Michigan, 108. Mount Clemens Original Well, Michigan, 89. Mount Clemens Soolbad Spring, Michigan, 108. Mount Mineral Spring, Shutesbury, Mass., 71. Mühlbrunnen, Carlsbad, Austria-Hungary, 97. Mühlbrunnen, Ober-Salzbrunn, Austria-Hungary, 125. Müllaken, Upper Austria, springs, 71. Mullein, 14. Munger's laundry, Chicago, Ill., artesian well, 91. Murquelle, Baden, Germany, 59. Musa paradisica, 18. Mushroom, edible, 15. Muskegon Lake, Michigan, 88. Mush Pot Spring, Yellowstone National Park, 78. Myrtle leaves and stems, 15. Myrtus cheken, 15.

N'Gonça, Sahara, Africa, well, 87. Naiwaschasee, Massailand, East Africa, spring, 101. Natronsäuerling, Nassau, Germany, 101. Natural soda-water, Passug, 115. Navioci, Bosnia, spring, 96. Neponset River, Hyde Park, Mass., 70. Neubrunnen, Carlsbad, Austria-Hungary, 97. Neubrunnen, Radein, Austria-Hungary, 117. Neue Badequelle, Ems, Germany, 75. Neue Belvedraquelle, Switzerland, 83. Neuenhainerquelle, Nassau, Germany, 60. Neue Quelle, Ems, Germany, 74. Neue Quelle, Mahren, Germany, 95. Neue Stahlquelle, Lobenstein, Germany, 117. Neumarkt, Germany, spring, 58. Neuragozi, Germany, spring, 123.

New Baden Spring, Missouri, 109.
New Hot Spring, Virginia, 106.
New Jersey tea, 9.
New Johannes Spring, Neuhaus, Austria-Hungary, 79.
New Putnam Spring, Saratoga, N. Y., 105.
New Spring, St. Moritz, Switzerland, 83.
Newsonis Arroyo Grande Springs, California, 64.
Neyrac, France, spring, 90.
Niederbronn, Germany, 72, 87.
Nieder Langenau, spring, 116.
Niederselters, Germany, spring, 75.
Nochemo Spring, Reed City, Mich., 88.
North Carolina Hot Springs, 67.
North or Daggy Spring, Greencastle. Ind., 107.

O Tura Säuerling, Austria-Hungary, 127. Oak Orchard Acid Springs, New York, 110. Oak Orchard Acid Water, New York, 110. Oats. 18. Oberbrunnen, Ober-Salzbrunn, Germany, 76, 125. Oberbrunnen, Flinsberg, Silesia, 116. Oberquelle, Bussang, France, 128. Ochsenhausen, Bieberach, Germany, spring, 73. Octagon Bath, Virginia, 106. Oestringer, Langenbrücken, Germany, spring, 127. Ofener Rakoczy Bitterwasser, 125. Ohe River, 36, 129. Ohio Magnetic Spring, 107. Ojo Caliente, N. M., spring, 82. Old Alum Well, Harrowgate, England, 68. Old Faithful Geyser, Yellowstone National Park, 78. Old Johannes Spring, Neuhaus, Austria-Hungary, 79. Old man's beard, 41. Old Spring, St. Moritz, Switzerland, 57, 83. Olgaquelle, Psekoup, 122. Olympian Salt Sulphur Spring, Kentucky, 107. Olympian White Sulphur Spring, Kentucky, 107. Ophelia chirata, 19. Ophiglossum vulgatum, 10. Opium, 39. Opuntia vulgaris, 25. Orange peel, 25. Oranien Quelle, Kreuznach, Germany, 54. Orchard Alum Spring, England, 123. Orchesterquelle, Carlsbad, Austria-Hungary, 97. Orezza, Island of Corsica, spring, 116. Orites excelsa, 33.

Orkney Bear Wallow Spring, Virginia, 111.
Orkney Healing Spring, Virginia, 111.
Orkney Powder Spring, Virginia, 111.
Ostgothenquelle, Bormio, Italy, 80.
Ottawa River, Montreal, Canada, 83.
Ottilienquelle, Paderborn, Germany, 59.
Ouargla, Sahara, Africa, well, 87.
Overall Mineral Wells, Texas, 67.
Owasso City, Mich., well, 88.
Owasso Mineral Spring, Michigan, 66.
Owatonna Mineral Spring, Minnesota, 109.
Owatonna Vichy Spring, Minnesota, 109.
Owens Lake, California, 61, 118, 119.

Pacific Congress Springs, California, 64. Pagoda Spring, Mount Clemens, Mich., 89. Pagoda (Napa-Soda) Spring, California, 64. Palanka, Servia, spring, 93. Palmyra Springs, Eye Springs, Wisconsin, 108. Palmyra Springs, Zenobia's Fountain, Wisconsin, 108. Palo Pinto Well, Texas, 107. Panacea Spring, Littleton, N. C., 107. Pandur, Kissingen, Germany, spring, 92. Panic grass, 12. Panicum capillare, 12. Panicum cruscorvi, 45. Panicum italicum, 45. Panicum monostachyum, 12. Pantelleria Island, Italy, 58. "Paprika," 7. Paracelsus Spring, St. Moritz, Switzerland, 57. Paramo de Ruiz, New Granada, spring, 92. Parana River, above union with Rio de la Plata, 87. Pareira, 31. Paris, France, melted snow, 124. Park Spring, Wisconsin, 109. Park's Springs, North Carolina, 67, 107. Parmelia scruposa, 40. Paroquet Springs, Kentucky, 112. Passaic River, New Jersey, 119. Passy, France, artesian well, 115. Paulaquelle, Mahren, Germany, 95. Paulaquelle, Johannisbrunn, Austria-Hungary, 97. Paullina sorbilis, 28. Pavilion Spring, Saratoga, N. Y., 105. Paw-paw, 28. Pearl Geyser, Yellowstone National Park, 77.

Peas, 8, 34, 45.

Pengalengen Plateau, East India, spring, 98.

Peninsula of Methana, spring, 68.

Penon de los Banos, Mexico, spring, 92.

Pepperwort, 38.

Perry Spring, No. 3, Lower Sulphur, Illinois, 114.

Peru, Ill., water-works, artesian well, 91.

Petalostigma quadriloculare, 15.

Peterborough County, Ontario, artesian well, 51.

Petersquelle, Petersthal, Germany, 58.

Petite Source, Challes, France, 128.

Pfaefers-Ragatz, Switzerland, spring, 90.

Pfister's Spring, Kansas, 109.

Phallusia mammilaris, 50.

Phaseolus radiatus, 44, 45.

Philippsquelle, Orb, Germany, 120.

Phytolacca decandra, 15.

Piedmont White Sulphur Iron Springs, California, 64.

Pinus larix, 20.

Pinus pumilio, 41.

Pinus sylvestris, 21, 22, 41.

Piper methysticum, 17.

Piscidia erythrina, bark, 23.

Pittsville Wells, Cheltenham, England, 124.

Plane tree leaves, 38.

Plantago major, 36.

Plantago maritima, 17.

Plantago media, 16.

Plantain, 16, 36.

Plantain seed, 17.

Pliniusquelle, Bormio, Italy, 8o.

Plums, 37.

Poison ivy, 21.

Poke weed root, 15.

Pokolsár, spring, 73.

Poland Star Spring, Maine, 105.

Polygonum hydropiper, 38.

Polysaccum pisocarpium, 16.

Pomegranate root bark, 34.

Poncho Hot Springs, Colorado, 65.

Pont-à-Mousson, France, spring, 79.

Ponte Molle, Italy, spring, 72.

Poppy petals, 39.

Portion, New South Wales, spring, 99.

Portulaca oleracea, 14.

Port Vendres, France, subterranean, 71.

Poschitz, Austria-Hungary, spring, 86. Posidonia oceanica, 33. Potash Sulphur Spring, Arkansas, 61. Potato weed, 14. Prajan Spring, Res. Madioen, East India, 98. Pré No. 1 Spring, Bagnères-de-Luchon, France, 73. Prè Saint Didier Springs, Courmayeur, Italy, 51. Précieuse (Allier), Vichy, France, spring, 103. Preblauer Sauerbrunnen, Kärnthen, Austria-Hungary, 100. Preservatives, 132. Prickly ash bark, 23. Prickly chaff flower, 39. Prickly pear, 25, 30. Primrose, 42. Primula farinosa, 42. Princess Spring, Caylus, France, 93. Protopterus annectus, 50. Providence Mineral Well, Kansas, 53. Prugnes, France, spring, 93. Prunus cerasus, 21. Prunus domestica, 37. Prunus spinosa, 32. Psekoup Springs, 122. Psilotum triquetrum, 10. Puits Chomel, Vichy, France, spring, 103. Pulaski Alum Spring, Virginia, 111. Purslane, 14. Putnam Spring, Saratoga, N. Y., 105.

Quelle an der Strasse, Srebrenica, Bosnia, 96. Quelle Marie, Bussang, France, 128. Quelle, No. 19, Katherienbad, Württemberg, Germany, 72. Quelle zu Raggel, Voralberg, Austria-Hungary, 87. Quirinusquelle, Aachen, Germany, 92.

Racoczy, Kissingen, Germany, spring, 92.
Raffanelo, Italy, spring, 98.
Ragaz-Pfäffers, spring, 115.
Ragweed, 14.
Rajecz-Töplitz, Austria-Hungary, spring, 77.
Raleigh's Mineral Spring, Box Springs, Tennessee, 112.
Raleigh's Mineral Spring, Marble Springs, Tennessee, 112.
Ramalina fraxinea, 22.
Ranigsdorf Säuerling, 85.
Ranunculus fluitans, 32.
Rape seed, 2.

Rape straw, 2.

Raso, Bosnia, spring, 96.

Ravenden Spring, Arkansas, 112.

Rawley Springs, Main Fountain, Virginia, 106.

Reading, Mass., filter gallery, 70.

Red bean, 45.

Red Boiling Springs, Tennessee, 67.

Red clover tops, 23.

Red currant juice, 21.

Red grape stalks, 24.

Red grapes, 24.

Red peppers, husk, 7.

Reindeer moss, 41.

Renaulme Spring, Saint-Denis, France, 81.

Rhatany, 17.

Rhine River, Germany, 69, 125.

Rhododendron, 32.

Rhododendron ferrugineum, 32.

Rhone River, Geneva, Switzerland, 69.

Rhubarb roots, 8.

Rhus aromatica, 18.

Rhus glabra, 14.

Rhus toxicodendron, 21.

Rhus vernicefera, 44.

Ribarska banja, Servia, 93.

Rice, 34, 45.

Rice straw, 33.

Richard Spring, Bagnères-de-Luchon, France, 73.

Richard Spring, Cransac, France, 128.

Richfield Sulphur Spring, New York, 105.

Rieumajou, France, spring, 73.

Riggio-Emilia, Italy, spring, 60.

Rio de la Plata, 87, 120.

Rio Vinagre, San Antonio, Colombia, 57.

Rjecicaquelle, Maglaj. Bosnia, 96.

Roanoke Red Sulphur Springs, Virginia, 68, 106.

Rochefort, France, artesian well, 118.

Roches (Beaurepaire), France, spring, 90.

Rock Enon Springs, Virginia, 106.

Rock Flat Spring, New South Wales, 99.

Rockbridge Alum Springs, Virginia, 80, 106, 111.

Rockford, Ill., artesian well, 91.

Rockford, Ill., water-works, artesian well, 91.

Roddes, France, spring, 124.

Roggendorff, Austria-Hungary, spring, 80, 102.

Roman Brine Spring, Torda, Austria-Hungary, 102.

Roman Spring, Sail-lès Chateaumorand, France, 81.

Römerbrunnen, Echzell, Wetterau, 114.

Römerquelle, Ems, Germany, 75.

Roncegno, South Tyrol, spring, 77.

Rosa remontana, 1.

Rose Spring, Caylus, France, 93.

Rosenau, Austria-Hungary, spring, 86.

Rosenquelle, Aachen, Germany, 92.

Roscicrucian Spring, Maine, 105.

Rosheim, France, spring, 116.

Rothenbrunnen, Voralberg, Austria-Hungary, 87.

Rothenfelser Mineralquelle, Baden, Germany, 59.

Roufaque, spring, 71.

Royal William Medical Establishment, Wiesbaden, Germany, spring, 75.

Royat Spring, France, 90.

Rubus articus, 7.

Rue de la Préfecture, Besançon, France, well, 69.

Ruiz Volcano Hot Spring, 57.

Rumex obtusifolius, 14.

Ruta herba, 23.

Rye, 39.

Sabadilla seed, 23.

Sacramento, Cal., river, 119.

Saginaw, Mich., water-supply, 88.

Saint-Denis Spring, France, 81.

Saint-Honoré, France, spring, 81.

Saint-Léon, Pougues-les-Eaux, Nièvre, France, 103.

Saint-Louis No. 1 (Allier), Vichy, France, spring, 103.

Saint-Mart, France, spring, 90.

Saint-Thiebaut, France, spring, 8o.

Saint-Yorre (Allier), Frobert, Vichy, France, spring, 103.

Salinensoole, Colberg, Germany, 55.

Salix vitellina, 29-

Salles-la-Source, Aveyron, France, 57.

Salmade, Bussang, France, spring, 128.

Salt Sulphur Iodine Spring, Virginia, 106.

Salvinia natans, 10.

Salzquelle, Kronthal, Germany, 93.

Salzquelle, Petersthal, Germany, 58.

Salzschlirf, Fulda, Germany, spring, 90.

Sambucus nigra, root, 19, 42.

San Francesco Spring, Spolato, 125.

Sandefjord, Sweden, spring, 123.

Santa Barbara Hot Springs, California, 64.

Santa Rosa White Sulphur Springs, California, 64.

Santa Ysabel Sulphur Springs, California, 64.

Sarsaparilla, 17, 25.

Sassafras, 14.

Sassafras officinale, 14.

Sauerbrunn, Ebriach, Austria-Hungary, 52.

Sauerbrunnen, Kärnthen, Austria-Hungary, 100.

Sauerbrunnen, Radein, Austria-Hungary, 117.

Säuerling Bades Vellach Quelle, No. II, Germany, 100.

Säuerling Bades Vellach Quelle, No. III, Germany, 100.

Säuerling Bades Vellach Quelle, No. IV, Germany, 100.

Säuerling Bades Vellach Quelle, No. V, Germany, 100.

Sāuerling, Bistrica, Bosnia, 96.

Säuerling, Ebriach, Germany, 100.

Säuerling, Dragunje, Bosnia, 96.

Săuerling, Dubnica, Bosnia, 96.

Säuerling, Jasenica, Bosnia, 96.

Säuerling, Kiseljack, Bosnia, 96.

Säuerling, Ljeskovica (Giftquelle), Bosnia, 96.

Säuerling, Slatina Ilidze, Bosnia, 96.

Säuerling, Tesanj, Bosnia, 96.

Säuerquelle, Apatovac, Austria-Hungary, 95.

Saule d'Urfe Spring, Sail-lès-Chateaumorand, France, 81.

Saxon, Switzerland, spring, 81, 118.

Schillingsforst Spring, Germany, 130.

Schlammkessel, Yellowstone National Park, 77.

Schlangenbad, Germany, spring, 74.

Schliefheiner, moor, 129.

Schlossbrunnen, Carlsbad, Austria-Hungary, 78, 97.

Schöneck, Russia, spring, 86.

Schooley's Mountain Spring, New Jersey, 105.

Schützenhofquelle, Wiesbaden, Germany, 76.

Schwefelwasser Bades Hopfreben, Voralberg, Austria-Hungary, 87.

Schweizerhalle, Basel, Switzerland, spring, 98.

Sclafani, Italy, spring, 103.

Scotch pine, 41.

Scotch pine needles, 20.

Scotch pine pollen, 22.

Sea weed, 28, 33.

Secale cereale, 29.

Secale cornutum, 18, 29, 37.

Seebruch, Austria-Hungary, spring, 70, 101.

Seifersdorf, Austria-Hungary, spring, 97.

Seine River, Bercy, France, 69.

Selaginella, 29.

Selaginella martensii, 10.

Selaginella spinulosa, 10.

Selters, Germany, artesian well, 76. Seltzer Spring, Saratoga, N. Y., 105. Semen cinæ, 15. Semlitsch Spring, Radein, Austria-Hungary, 117. Seon, Germany, spring, 71. Serneus, Switzerland, spring, 114. Seven Springs, Abingdon, Va., 111. Sévigné (Allier), Vichy, France, spring, 103. Shaft Spring, Torda, Austria-Hungary, 102. Shealtiel Mineral Springs, Wisconsin, 108. Sheboygan Mineral Springs, Wisconsin, 108. Sheboygan, Wis., artesian well, 60, 119. Shenandoah Alum Spring, Virginia, 106. Shepherd's purse, 13. Sheridan Springs, Wisconsin, 108. Shiawassee River, Owasso, Mich., 88. Siebenbürg, Austria-Hungary, spring, 129. Siloam Springs, Iowa, 109. Silurian Spring, Wisconsin, 108. Silverthorne's Well, Broken Hill, New South Wales, 99 Sinnbergerquelle, Brückenau, Germany, 120. Sissipan, East India, spring, 98. Skagg's Hot Springs, California, 110. Slanik, Roumania, spring, 124. Slatersville, N. Y., spring, 127. Slatina Ilidze, spring, 96. Sloe. (See Black thorn.) Smithville, Ga., artesian well, 118. Smodelac, Bosnia, spring, 96. Soda Butte Spring, Yellowstone National Park, 78. Soda Spring, Berkshire, Mass., 127. Soda Spring, Yellowstone National Park, 77. Soja hispida, 45. Solanum carolinense, 14. Solfatare, Puzzuoli, Italy, 69. Soljanka, Ciechocinek, Poland, spring, 129. Someraubad, Mähren, Germany, 123. Soolquelle, Kammin, Germany, 56. Soolsprudel, Nassau, Germany, 60. Soolsprudel I, Nassau, Germany, 60. Soolsprudel III, Nassau, Germany, 60. Soolsprudel IV, Nassau, Germany, 60. Soolsprudel VII, Nassau, Germany, 60. Soolsprudel X, Nassau, Germany, 60. Sophienquelle, Petersthal, Germany, 58. Sour Lake Mineral Springs, Texas, 112.

Sour Springs, Texas, 112.

Source de la Place, Bourbonne, France, 73.

Source de l'hôpital, France, 81.

Source de l'intérieur de l'établissement, Bourbonne, France, 73.

Source de Sainte-Catherine, Plombières, France, 82.

Source des Bains Nouveaux, Sylvanès, France, 100.

Source des Demoiselles, Vittel, France, 100.

Source des Moines, Sylvanès, France, 100.

Source des Petites Baignoires, Sylvanès, France, 100.

Source des Petites-Eaux, Sylvanès, France, 100.

Source du Crucifix, Plombières, France, 82.

Source du ferrugineuse de Bourdeille, Plombières, France, 82.

Source du Mont Cornador, Saint Nectaire, 57.

Source du Puits, Cusset, France, 81.

Source du Savonneuse, Plombières, France, 82.

Source ferrugineuse, Plombières, France, 100.

Source Grande Grille, France, 81.

Source Marie, Vittel, France, 100.

Source nouvelle, France, 81.

Source Prè Salé, Vichy, France, 81.

Source principale, Challes, France, 128.

Source Roques, France, 59.

Source Roquette, France, 59.

Source Rouge, Saint Nectaire, 57.

Source Saint-Louis, France, 60.

Source-basse Richard, Cransac, France, 81.

Source-haute Richard, Cransac, France, 81.

Sources de Fraysse, France, 59.

Sources des Dames, Plombières, France, 82, 100.

South Dakota Hot Springs, Mammoth Mineral Springs, 67.

South Park, Col., spring, 127.

Spanish moss, 25.

Sparta Artesian Well, Wisconsin, 54.

Sparta Mineral Magnetic Well, Wisconsin, 113.

Sphagnum cuspidatum, 41.

Sphagnum moss, 41.

Spiegelbad No. 1, Trentschin Töplitzer Bad, Austria-Hungary, 80.

Splendid Geyser, Yellowstone National Park, 78.

Sponge, 49.

Spring Bir Keraui, 82.

Sprudel, Carlsbad, Austria-Hungary, 97.

Sprudel Spring I, 56.

Sprudel Spring II, 56.

Spurge, 42.

St. Augustine, Fla., artesian well, 61.

St. Clair Mineral Spring, Mich., 66.

St. Clair Spring, Oakland House Well, Mich., 89.

St. Croix Mineral Spring, Wisconsin, 108.

St. Dunstan's Well, Melrose, Scotland, 85.

St. Gothard Spring, Ceneda, Italy, 56.

St. Helena, Battaglia, Italy, springs, 120.

St. John's breadfruit, 30.

St. Lawrence River, Pointe-des-Cascades, Canada, 83.

St. Léon, Canada, spring, 60, 83.

St. Lorenz Hot Spring, Leuk, Switzerland, 98.

St. Moritz, Switzerland, springs, 115.

St. Paul Spring, Rouen, France, 77.

St. Peter's, Macomb, Ill., spring, 91.

St. Petersquelle, Tiefenkasten, 115.

Stachelberg, Switzerland, spring, 121.

Stafford Spring, Connecticut, 65.

Stahl, Germany, spring, 124.

Stahlbrunnen (Hermannsborn), Pömbsen, Germany, 126.

Stahlbrunnen, Pyrmont, Germany, 74.

Stahlbrunnen, Wassenach, 55.

Stahlquelle, Nassau, Germany, 93.

Star anise, 32.

Staraja-Russa, Russia, artesian well, 120.

Steamboat Springs, Nevada, 55.

Steben, Germany, spring, 78, 117.

Stereocaulon vesuvianum, 9.

Sterling, Ill., artesian well, 91.

Stickweed, 14.

Stolypin, Russia, spring, 120.

Stoughton, Mass., well, 70.

Strawberry juice, 21.

Stribling or Augusta Alum Springs, Virginia, III.

Strontia Mineral Spring, Maryland, 106.

Südliche Drainagequelle, Psekoup, 122.

Sugar cane, 27.

Suippe River, France, 99.

Suliguli, spring, 80.

Sulphur Bath or Ladies' Sulphur Bath, Virginia, 106.

Sulphur Spring A, Lostorf, Switzerland, 57.

Sulphur Spring, Livern, Germany, 124.

Sulphur Spring, Oberdorf, Germany, 58.

Sulphur Spring, Sail-lès-Chateaumorand, France, 81.

Sulphur Spring, Schinznach, Switzerland, 57.

Sulz. Germany, spring, 117.

Sulzbach, Germany, spring, 102.

Sumach, 14, 18.

Summit Soda Springs, California, 64, 110.

Sunflower seeds, 43.
Suzon Spring, Dijon, France, 69.
Sweet potato, 18.
Sweet Springs (Akesion), Missouri, 109.
Sweet Springs (Sweet), Missouri, 109.
Syringa vulgaris, 42.

Talladega Spring, Alabama, 107. Tarasp, Switzerland, springs, 83. Tarbragar River, near Dubbo, New South Wales, artesian well, 99. Tarentum, Pa., artesian well, 123. Taro, 18. Tarrella, New South Wales, spring, 99. Taurus Geyser, Yellowstone National Park, 78. Tayuya, 27. Te Aroha, Queensland, springs, 93. Te Tarata Hot Spring, New Zealand, 121. Tea, 32. Tempelbrunnen, Rohitsch, Germany, 58. Temple Spring, Oberfranken, 122. Tenninger Bad, Grisons, Switzerland, 99. Tennstadt, Thuringia, Germany, spring, 98. Termini-Imerese, Palermo, Sicily, 103. Thames River, Greenwich, 52. Thames River, London, 52, 79. Thames River, Twickenham, England, 61. Thapsia garganica, 48. Thapsia sylphium, 48. Thérèse Sulphur Spring, Fumades, France, 55. Theresienbrunnen, Carlsbad, Austria-Hungary, 97. Thermal Acid Spring, California, 64, 113. Thompson's Bromine-arsenic Spring, North Carolina, 66. Tibia from Egyptian mummy, 50. Tickle grass, 12. Tilford's Mineral Well, Tennessee, 107. Tillandsia dianthoidea, 25. Tittabawassee River, Michigan, 88. Tobacco, 13, 18, 19, 34, 40. Tobacco leaves and screenings, 8, 31. Tobelbad, Steiermark, Austria-Hungary, spring, 95. Tokiew, Caucasus, spring, 55. Tolenas Springs, California, 64, 110. Tongern, Belgium, spring, 90. Tönnissteiner Heilbrunnen, Germany, 75, 91. Tönnissteiner Stahlbrunnen, Germany, 75.

Topeka Mineral Wells, Kansas, 65.

Töplitz, Mähren, Germany, spring, 123. Torpasaltkälle, Lilla Edet, Sweden, 56. Tougourt, Sahara, Africa, well, 87. Touzla Spring, Asia Minor, 121. Traverse Bay, Michigan, 88. Tree fern, New Zealand, 10, 11. Trefriw, Wales, spring, 59. Trentham Spring, Georgia, 65. I rianosperma ficifolia, 27. Trinkquelle, Driburg, Germany, 74. Trinkquelle, Elster, Germany, 122. Trinkquelle No. 10, Trinkhalle, Württemberg, Germany, 72. Triton Spring, Saratoga, N. Y., 60. Truffles, 9. Tschéhirghé, Anatolia, Asia Minor, spring, 115. Tsuga canadensis, 2. Tuber magnatum, 29. Tuber melanosporum, 29. Turban and Grand Geysers, Yellowstone National Park, 78. Turf moss, 16. Tuscan (or Lick) Springs, California, 64. Tuscarora Lithia Spring, Pennsylvania, 67. Tuscarora Sour Spring, Brantford, Canada, 83. Tussilago farfara, 8.

Umgemach, Baden-Baden, Germany, spring, 59.
Union Spring, Saratoga, N. Y., 60, 105.
United States Spring, Saratoga, N. Y., 105.
Unity Springs, (Iron Spring), New Hampshire, 105.
Ursprung, Baden, Germany, 59.
Usnea, 16.
Usnea barbata, 41.
Utah Hot Springs, 67.
Utley Mineral Spring, Wisconsin, 109.

Val Sinestra, Switzerland, springs, 84.
Valdagno, Italy, spring, 73.
Valle del Gallo, near Urbino, spring, 51.
Varennes, Canada, springs, 83.
Variety Springs, (Alum Spring), Virginia, 111.
Variolaria dealbata, 41.
Veal breast, 50.
Velaine-Saint-Aubin, France, spring, 116.
Velika Kiselica, Srebrenica, Bosnia, spring, 97.
Verbascum thapsus, 14.
Verbesina siegesbeckia, 14.

Vesle River, France, 99. Vesta Spring, Wisconsin, 108. Vichy Spring, Saratoga, N. Y., 105. Vicia faba, 18. Victoria Spring, Ems, Germany, 75. Victoria Spring, Neuenahr, 56. Victoria Spring, Oberlahnstein, 76. Victoria Well, Ofen, Austria-Hungary, 54. Vienna (Josefstadt), Austria-Hungary, spring, 115. Villecelle, France, spring, 52. Vincenzbrunnen, Luhatschowitz, Austria-Hungary, 72, 85. Vincent (Allier), Vichy, France, spring, 103. Virginia creeper, 40. Vic-sur-Cère, Cantal, France, spring, 122. Vitis hederacea, 40. Vittel, France, spring, 100. Voslau, spring, 121. Vruciča, Bosnia, spring, 96.

Wakefield Sulphur Well, Kansas, 53. Wallawhatoola Alum Springs, Virginia, 111. Walley's Hot Springs, Nevada, 66. Walsh's quarry, New South Wales, spring, 99. Warasdin-Töplitz, Austria-Hungary, spring, 80. Warm Springs, Arkansas, 107. Warner's Spring, Michigan, 108. Waroo Springs, New South Wales, artesian well, 100. Warwick Spring, New Jersey, 105. Washington Lithia Well, Ballston Spa, New York, 105. Water hemlock, 16. Wattweiler, Alsace, spring, 61. Wayland, Mass., filter gallery, 70. Wayland, Mass., reservoir, 70. Weaver Well, Texas, 112. "Weigmann," 16. Weilbach, Nassau, Germany, spring, 74, 128. Weisloch, Baden, Germany, spring, 126. Welcome Island Lithia Water, 88. Wels, Austria-Hungary, spring, 97. Wenzelsquelle, Rippoldsau, spring, 128. Wernarzer, Brückenau, Germany, spring, 119. West Baden Spring, No. I, Indiana, 113. West Saratoga Springs, Indiana, 108. West Spring, Joplin, Mo., 82. West Sprins, Langenbruck, Austria-Hungary, 77. Westborough Insane Hospital, Mass., wells, 70.

Westerland, Germany, artesian well, 82. Western Butte, Northwest Territory, spring, 86. Westliche Quelle, Stuttgart, Germany, 72. Wheat, 1, 18, 28, 34, 36, 45. Wheat flour, 8, 14, 36. Wheat, milling products, 5, 36. Wheelock, England, spring, 102. White birch, 41. White Cliff Sulphur Spring, Tennessee, 107. White currant juice, 21. White Island, New Zealand, hot lake, 70. White Mineral Spring, Minnesota, 66. White pine litter, 40. White Rock Springs, Wisconsin, 109. White Sulphur Springs, Montana, 126. White Terrace Geyser, New Zealand, 121. White truffle, 29. White wine, 17. Wiener-Neustadt Sauerbrunn, Austria-Hungary, 117. Wilcannia, Tarella, New South Wales, spring, 99. Wiesen Spring, Oberfranken, 122. Wilbur Springs, California, 64. Wilhelms, Germany, spring, 124. Wilhelmsheilanstalt, Wiesbaden, Germany, 75. Wilhelmsquelle, Herzogludolfsbad, Germany, 103. Willow, 29. Wine, 6, 16, 21, 24, 33. Winsford, England, spring, 102. Wire grass, 14. Witter's Mineral Springs, California, 64. Woburn, Mass., filter gallery, 70. Woburn, Mass., pond. 70. Wolf Trap Lithia Springs, Virginia, 113. Wolverton, England, spring, 76. Woodstock, Ill., drift well, or. Wootan Wells, Texas, 107. Worm seed, 15. Wormwood, 16. Wrnjčka banja, Servia, 93.

Xanthium spinosum, 17.

Wranjaska banja, Servia, 93.

Yalova Hot Spring, Asia Minor, 121. Yampa Spring, Glenwood Springs, Colorado, 122. Yams. (See Sweet potato.)
Yarrangobilly caves, New South Wales, spring, 99.
Yeast, 2.
Yellow Sulphur Springs, Virginia, 111.
Yellowstone Lake, Yellowstone National Park, 78.
Young's Natural-gas Well and Mineral Springs, California, 64.
Ypsilanti, Mich., well, 88.
Ypsilanti Sanitarium, Mich., spring, 89.
Yvonne, France, spring, 103.

Zauberwasser, Hudson, Mich., 88. Zillenbergsoole, Colberg, Germany, 55. Zostera marina, 2, 16. Zuckmantel, Austria-Hungary, spring, 98.







SHORT-TITLE CATALOGUE

OF THE

PUBLICATIONS

OF

JOHN WILEY & SONS,

NEW YORK.

LONDON: CHAPMAN & HALL, LIMITED.

ARRANGED UNDER SUBJECTS.

Descriptive circulars sent on application. Books marked with an asterisk are sold at net prices only, a double asterisk (**) books sold under the rules of the American Publishers' Association at net prices subject to an extra charge for postage. All books are bound in cloth unless otherwise stated.

AGRICULTURE.

Armsby's Manual of Cattle-feeding12mo,		
Principles of Animal Nutrition	4	00
Budd and Hansen's American Horticultural Manual:		
Part I.—Propagation, Culture, and Improvement	1	50
Part II.—Systematic Pomology	I	50
Downing's Fruits and Fruit-trees of America8vo,	5	00
Elliott's Engineering for Land Drainage12mo,	1	50
Practical Farm Drainage12mo,		
Green's Principles of American Forestry12mo,	1	50
Grotenfelt's Principles of Modern Dairy Practice. (Woll.)12mo,	2	00
Kemp's Landscape Gardening12mo,	2	50
Maynard's Landscape Gardening as Applied to Home Decoration12mo,	1	50
Sanderson's Insects Injurious to Staple Crops	1	50
Insects Injurious to Garden Crops. (In preparation.)		
Insects Injuring Fruits. (In preparation.)		
Stockbridge's Rocks and Soils8vo,	2	50
Woll's Handbook for Farmers and Dairymen	1	50
seren's Text-book on Iteld Fortification Small Byo 2 on		
ARCHITECTURE.		
Baldwin's Steam Heating for Buildings12mo,	2	50
Berg's Buildings and Structures of American Railroads4to,	5	col
Birkmire's Planning and Construction of American Theatres 8vo,	3	0
	2	E 0
Architectural Iron and Steel8vo,	-	
Architectural Iron and Steel	-	-
	2	co
Compound Riveted Girders as Applied in Buildings	3	50
Compound Riveted Girders as Applied in Buildings	3 3	co 50 co
Compound Riveted Girders as Applied in Buildings	2 3 3 4	50 60
Compound Riveted Girders as Applied in Buildings	2 3 3 4 4	50 co oo
Compound Riveted Girders as Applied in Buildings	2 3 3 4 4 3	50 00 00 00 50
Compound Riveted Girders as Applied in Buildings	2 3 3 4 4 3 2	co co co oo oo 50
Compound Riveted Girders as Applied in Buildings	2 3 3 4 4 3 2 2	50 00 00 50 50 50
Compound Riveted Girders as Applied in Buildings	2 3 3 4 4 3 2 1 1	50 00 00 50 50 50 50
Compound Riveted Girders as Applied in Buildings	2 3 3 4 4 3 2 1 1	50 00 00 50 50 50 50

Fletcher's Practical Instructions in Quantitative Assaying with the Blowpipe		
12mo, morocco,	I	50
Fowler's Sewage Works Analyses	2	00
Fresenius's Manual of Qualitative Chemical Analysis. (Wells.)8vo,		
Manual of Qualitative Chemical Analysis. Part I. Descriptive. (Wells.)		
8vo,	3	00
System of Instruction in Quantitative Chemical Analysis. (Cohn.)	-	
2 vols8vo,	12	50
Fuertes's Water and Public Health	T	50
Furman's Manual of Practical Assaying8vo,	2	00
Getman's Exercises in Physical Chemistry	2	00
Gill's Gas and Fuel Analysis for Engineers	-	25
Grotenfelt's Principles of Modern Dairy Practice. (Woll.)	-	-5
Hammarsten's Text-book of Physiological Chemistry. (Mandel.)8vo,		
Helm's Principles of Mathematical Chemistry. (Morgan.)12mo,		
Hering's Ready Reference Tables (Conversion Factors)16mo, morocco,		
Hinds's Inorganic Chemistry8vo,		
* Laboratory Manual for Students	3	
		75
Holleman's Text-book of Inorganic Chemistry. (Cooper.)8vo,		
Text-book of Organic Chemistry. (Walker and Mott.)		
* Laboratory Manual of Organic Chemistry. (Walker.)12mo,		
Hopkins's Oil-chemists' Handbook		
Jackson's Directions for Laboratory Work in Physiological Chemistry 8vo,		25
Keep's Cast Iron8vo,		
Ladd's Manual of Quantitative Chemical Analysis		
Landauer's Spectrum Analysis. (Tingle.)8vo,		
Lassar-Cohn's Practical Urinary Analysis. (Lorenz.)	1	00
Leach's The Inspection and Analysis of Food with Special Reference to State		
Control. (In preparation.)		
Löb's Electrolysis and Electrosynthesis of Organic Compounds. (Lorenz.) 12mo,	I	00
Mandel's Handbook for Bio-chemical Laboratory	I	50
* Martin's Laboratory Guide to Qualitative Analysis with the Blowpipe . 12mo,		60
Mason's Water-supply. (Considered Principally from a Sanitary Standpoint.)		
3d Edition, Rewritten 8vo,	4	00
Examination of Water. (Chemical and Bacteriological.)12mo,	100	25
Meyer's Determination of Radicles in Carbon Compounds. (Tingle.) 12mo,		00
Miller's Manual of Assaying		
Mixter's Elementary Text-book of Chemistry		50
Morgan's Outline of Theory of Solution and its Results		00
Elements of Physical Chemistry		
Morse's Calculations used in Cane-sugar Factories 16mo, morocco,		
		50
Mulliken's General Method for the Identification of Pure Organic Compounds.	-	
Vol. I Large 8vo,	5	00
Nichols's Water-supply. (Considered mainly from a Chemical and Sanitary		
Standpoint, 1883.)8vo,		
O'Brine's Laboratory Guide in Chemical Analysis8vo,		
O'Driscoll's Notes on the Treatment of Gold Ores	2	00
Ost and Kolbeck's Text-book of Chemical Technology. (Lorenz-Bozart.)		
(In preparation.)		
Ostwald's School of Chemistry. Part One. (Ramsey.) (In press.)		
* Penfield's Notes on Determinative Mineralogy and Record of Mineral Tests.		
8vo, paper,		
Pictet's The Alkaloids and their Chemical Constitution. (Biddle.)8vo,	5	00
Pinner's Introduction to Organic Chemistry. (Austen.)12mo,		
Poole's Calorific Power of Fuels		
Prescott and Winslow's Elements of Water Bacteriology, with Special Refer-	I	
ence to Sanitary Water Analysis	I	25
* Reisig's Guide to Piece-dveing		

Richards and Woodman's Air, Water, and Food from a Sanitary Standpoint . 8vo,	2	00
Richards's Cost of Living as Modified by Sanitary Science	I	00
Cost of Food a Study in Dietaries	I	00
Richards and Williams's The Dietary Computer	I	50
Ricketts and Russell's Skeleton Notes upon Inorganic Chemistry. (Part I		
Non-metallic Elements.)		75
Ricketts and Miller's Notes on Assaying8vo,	3	00
Rideal's Sewage and the Bacterial Purification of Sewage	3	50
Disinfection and the Preservation of Food8vo,	4	00
	2	00
Sabin's Industrial and Artistic Technology of Paints and Varnish. (In press.)		
Salkowski's Physiological and Pathological Chemistry. (Orndorff.)8vo,	2	50
	2	50
Essentials of Volumetric Analysis	I	25
Spencer's Handbook for Chemists of Beet-sugar Houses 16mo, morocco,	3	00
Handbook for Sugar Manufacturers and their Chemists . 16mo, morocco,	2	00
Stockbridge's Rocks and Soils8vo,	2	50
	1	50
		00
		00
		00
	-	00
		50
		00
Wassermann's Immune Sera: Hæmolysins, Cytotoxins, and Precipitins. (Bol-		-
	_	00
		50
Short Course in Inorganic Qualitative Chemical Analysis for Engineering	53	30
		50
		50
	-	50
		50
		50
Wulling's Elementary Course in Inorganic Pharmaceutical and Medical Chem-	•	20
		00
15th year of the second	1	00
CIVIL ENGINEERING.		
BRIDGES AND ROOFS. HYDRAULICS. MATERIALS OF ENGINEER	TR	TG.
RAILWAY ENGINEERING.		
	g.	
Baker's Engineers' Surveying Instruments		
Bixby's Graphical Computing Table		25
** Burr's Ancient and Modern Engineering and the Isthmian Canal. (Postage,	a	
	200	50
		50
		00
	I	50
		00
Folwell's Sewerage. (Designing and Maintenance.)	3	00
Folwell's Sewerage. (Designing and Maintenance.)	3	
Folwell's Sewerage. (Designing and Maintenance.)	3	00
Folwell's Sewerage. (Designing and Maintenance.)	3 3 2	50
Folwell's Sewerage. (Designing and Maintenance.)	3 3 2 1	50 50
Folwell's Sewerage. (Designing and Maintenance.). Svo, Freitag's Architectural Engineering. 2d Edition, Rewritten. Svo, French and Ives's Stereotomy. 8vo, Goodhue's Municipal Improvements. 12mo, Goodrich's Economic Disposal of Towns' Refuse. 8vo, Gore's Elements of Geodesy. 8vo,	3 3 2 1 3	50 50 75
Folwell's Sewerage. (Designing and Maintenance.). 8vo, Freitag's Architectural Engineering. 2d Edition, Rewritten 8vo, French and Ives's Stereotomy. 8vo, Goodhue's Municipal Improvements 12mo, Goodrich's Economic Disposal of Towns' Refuse 8vo, Gore's Elements of Geodesy 8vo, Hayford's Text-book of Geodetic Astronomy 8vo,	3 3 2 1 3 2	50 50 75 50
Folwell's Sewerage. (Designing and Maintenance.)	3 3 2 1 3 2 3	50 50 75 50 50
Folwell's Sewerage. (Designing and Maintenance.). 8vo, Freitag's Architectural Engineering. 2d Edition, Rewritten 8vo, French and Ives's Stereotomy. 8vo, Goodhue's Municipal Improvements 12mo, Goodrich's Economic Disposal of Towns' Refuse 8vo, Gore's Elements of Geodesy 8vo, Hayford's Text-book of Geodetic Astronomy 8vo, Hering's Ready Reference Tables (Conversion Factors). 16mo, morocco, Howe's Retaining Walls for Earth 12mo,	3 3 2 1 3 2 3 2	50 50 75 50 50
Folwell's Sewerage. (Designing and Maintenance.). Svo, Freitag's Architectural Engineering. 2d Edition, Rewritten Svo, French and Ives's Stereotomy. Svo, Goodhue's Municipal Improvements 12mo, Goodrich's Economic Disposal of Towns' Refuse Svo, Gore's Elements of Geodesy Svo, Hayford's Text-book of Geodetic Astronomy Svo, Hering's Ready Reference Tables (Conversion Factors). 16mo, morocco, Howe's Retaining Walls for Earth 12mo,	3 3 2 1 3 2 3 2 1	50 50 75 50 50 50

Kiersted's Sewage Disposal12mo,	1	25
Laplace's Philosophical Essay on Probabilities. (Truscott and Emory.) 12mo.	2	00
Mahan's Treatise on Civil Engineering. (1873.) (Wood.)8vo,	5	00
Descriptive Geometry8vo,	-	E0
Merriman's Elements of Precise Surveying and Geodesy	-	50
Elements of Sanitary Engineering	-	50
Merriman and Brooks's Handbook for Surveyors16mo, morocco,	2	00
Werranan and Brooks's Handbook for Surveyors10mo, morocco,	2	00
Nugent's Plane Surveying8vo,	3	50
Ogden's Sewer Design	2	00
Patton's Treatise on Civil Engineering	7	50
Reed's Topographical Drawing and Sketching4to,	5	00
Rideal's Sewage and the Bacterial Purification of Sewage8vo,	3	50
Siebert and Biggin's Modern Stone-cutting and Masonry	I	50
Smith's Manual of Topographical Drawing. (McMillan.)8vo,	2	50
Sondericker's Graphic Statics, with Applications to Trusses, Beams, and		30
Arches		
* Trautwine's Civil Engineer's Pocket-book16mo, morocco,	-	00
Trantwine's Civil Engineer's Pocket-book	5	00
Wait's Engineering and Architectural Jurisprudence8vo,		
Sheep,	6	50
Law of Operations Preliminary to Construction in Engineering and Archi-		
tecture8vo,	5	00
Sheep,	5	50
Law of Contracts8vo,		
Warren's Stereotomy-Problems in Stone-cutting		
Webb's Problems in the Use and Adjustment of Engineering Instruments.	-	30
	_	
16mo, morocco,		
* Wheeler's Elementary Course of Civil Engineering		
Wilson's Topographic Surveying	3	50
AND LONG THE RESERVE AND ADDRESS OF THE PARTY OF THE PART		
BRIDGES AND ROOFS.		
	2	00
Boller's Practical Treatise on the Construction of Iron Highway Bridges 8vo,		
Boller's Practical Treatise on the Construction of Iron Highway Bridges8vo, * Thames River Bridge		
* Boller's Practical Treatise on the Construction of Iron Highway Bridges8vo, * Thames River Bridge	5	00
Boller's Practical Treatise on the Construction of Iron Highway Bridges8vo, * Thames River Bridge	5	50
* Thames River Bridge	3 10	50
* Thames River Bridge	5 3 10 5	50 00
* Thames River Bridge	5 3 10 5 2	50 00 00 50
* Thames River Bridge	5 3 10 5 2 1	50 00 00 50 25
* Thames River Bridge	5 3 10 5 2 1	50 00 00 50 25
Boller's Practical Treatise on the Construction of Iron Highway Bridges 8vo, * Thames River Bridge	5 3 10 5 2 1 2	50 00 00 50 25 50
* Thames River Bridge	5 3 10 5 2 1 2 2	50 00 00 50 25 50
* Thames River Bridge	5 3 10 5 2 1 2 4	50 00 50 50 25 50 50
Boller's Practical Treatise on the Construction of Iron Highway Bridges 8vo, * Thames River Bridge	5 3 10 5 2 1 2 4	50 00 50 50 25 50 50
Boller's Practical Treatise on the Construction of Iron Highway Bridges 8vo, * Thames River Bridge	5 3 10 5 2 1 2 2 4 2	50 00 00 50 25 50 00 00
Boller's Practical Treatise on the Construction of Iron Highway Bridges 8vo, * Thames River Bridge	5 3 10 5 2 1 2 2 4 2	50 00 00 50 25 50 00 00
Boller's Practical Treatise on the Construction of Iron Highway Bridges 8vo, * Thames River Bridge	5 3 10 5 2 1 2 2 4 2	50 00 50 50 50 50 00 00
Boller's Practical Treatise on the Construction of Iron Highway Bridges 8vo, * Thames River Bridge	5 3 10 5 2 1 2 2 4 2 10 2	50 00 50 50 50 50 00 00
Boller's Practical Treatise on the Construction of Iron Highway Bridges 8vo, * Thames River Bridge	5 3 10 5 2 1 2 2 4 2 10 2 2	50 00 00 50 25 50 00 00 00
* Thames River Bridge	5 3 10 5 2 1 2 2 4 2 10 2 2 2	50 00 00 50 50 50 00 00 50 50 50 50 50 5
Boller's Practical Treatise on the Construction of Iron Highway Bridges 8vo, * Thames River Bridge	5 3 10 5 2 1 2 2 4 2 10 2 2 2	50 00 00 50 50 50 00 00 50 50 50 50 50 5
Boller's Practical Treatise on the Construction of Iron Highway Bridges 8vo, * Thames River Bridge	5 3 10 5 2 1 2 2 4 2 10 2 2 2	50 00 50 50 50 50 00 00 50 50 50 50
Boller's Practical Treatise on the Construction of Iron Highway Bridges. 8vo, * Thames River Bridge	5 3 10 5 2 1 2 2 4 2 10 2 2 2 10	50 00 50 50 50 50 00 00 50 50 50 50
Boller's Practical Treatise on the Construction of Iron Highway Bridges. 8vo, * Thames River Bridge	5 3 10 5 2 1 2 4 2 10 2 2 2 2 10 3	50 00 50 50 50 50 00 00 50 50 50 50 50 00
Boller's Practical Treatise on the Construction of Iron Highway Bridges. 8vo, * Thames River Bridge	5 3 10 5 2 1 2 2 4 2 10 2 2 2 2 10 3 10 10 10 10 10 10 10 10 10 10 10 10 10	50 00 00 50 50 50 00 00 50 50 50 50 50 5
* Thames River Bridge	5 3 10 5 2 1 2 2 4 2 10 2 2 2 2 10 3 10 10 10 10 10 10 10 10 10 10 10 10 10	50 00 00 50 50 50 00 00 50 50 50 50 50 5
Boller's Practical Treatise on the Construction of Iron Highway Bridges. 8vo, * Thames River Bridge. 4to, paper, Burr's Course on the Stresses in Bridges and Roof Trusses, Arched Ribs, and Suspension Bridges. 8vo, Du Bois's Mechanics of Engineering. Vol. II. 5mall 4to, Foster's Treatise on Wooden Trestle Bridges. 4to, Fowler's Coffer-dam Process for Piers. 8vo, Greene's Roof Trusses. 8vo, Bridge Trusses. 8vo, Arches in Wood, Iron, and Stone. 8vo, Arches in Wood, Iron, and Stone. 8vo, Design of Simple Roof-trusses in Wood and Steel 8vo, Jehnson, Bryan, and Turneaure's Theory and Practice in the Designing of Modern Framed Structures. 5mall 4to, Merriman and Jacoby's Text-book on Roofs and Bridges: Part I.—Stresses in Simple Trusses 8vo, Part III.—Bridge Design. 4th Edition, Rewritten. 8vo, Part IV.—Higher Structures. 8vo, Morison's Memphis Bridge. 4to, Waddell's De Pontibus, a Pocket-book for Bridge Engineers. 16mo, moroccc. Specifications for Steel Bridges. 12mo, Wood's Treatise on the Theory of the Construction of Bridges and Roofs. 8vo, Wright's Designing of Draw-spans:	5 3 10 5 2 1 2 2 4 2 10 2 2 2 10 3 1 2 10 10 10 10 10 10 10 10 10 10 10 10 10	50 00 00 50 25 50 00 00 50 50 50 50 50 50 50 50 50 50
Boller's Practical Treatise on the Construction of Iron Highway Bridges. 8vo, * Thames River Bridge	5 3 10 5 2 1 2 2 4 2 10 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	50 00 00 50 25 50 00 00 50 50 50 50 50 50 50 50 50 50
Boller's Practical Treatise on the Construction of Iron Highway Bridges. 8vo, * Thames River Bridge. 4to, paper, Burr's Course on the Stresses in Bridges and Roof Trusses, Arched Ribs, and Suspension Bridges. 8vo, Du Bois's Mechanics of Engineering. Vol. II. 5mall 4to, Foster's Treatise on Wooden Trestle Bridges. 4to, Fowler's Coffer-dam Process for Piers. 8vo, Greene's Roof Trusses. 8vo, Bridge Trusses. 8vo, Arches in Wood, Iron, and Stone. 8vo, Arches in Wood, Iron, and Stone. 8vo, Design of Simple Roof-trusses in Wood and Steel 8vo, Jehnson, Bryan, and Turneaure's Theory and Practice in the Designing of Modern Framed Structures. 5mall 4to, Merriman and Jacoby's Text-book on Roofs and Bridges: Part I.—Stresses in Simple Trusses 8vo, Part III.—Bridge Design. 4th Edition, Rewritten. 8vo, Part IV.—Higher Structures. 8vo, Morison's Memphis Bridge. 4to, Waddell's De Pontibus, a Pocket-book for Bridge Engineers. 16mo, moroccc. Specifications for Steel Bridges. 12mo, Wood's Treatise on the Theory of the Construction of Bridges and Roofs. 8vo, Wright's Designing of Draw-spans:	5 310 5 2 1 2 2 2 2 10 3 1 2 2 2	50 00 00 50 50 50 00 00 50 50 50 50 50 5

HYDRAULICS.

Bazin's Experiments upon the Contraction of the Liquid Vein Issuing from an		
Orifice. (Trautwine.)8vo,	2	00
Bovey's Treatise on Hydraulics8vo,		00
Church's Mechanics of Engineering8vo,	-	00
Diagrams of Mean Velocity of Water in Open Channelspaper,		50
Coffin's Graphical Solution of Hydraulic Problems 16mo, morocco,		50
Flather's Dynamometers, and the Measurement of Power12mo,		00
Folwell's Water-supply Engineering8vo,		00
Frizell's Water-power	-	00
Fuertes's Water and Public Health12mo,		50
Water-filtration Works12mo,		50
Ganguillet and Kutter's General Formula for the Uniform Flow of Water in	-	30
Rivers and Other Channels. (Hering and Trautwine.)8vo,	4	00
Hazen's Filtration of Public Water-supply8vo,	260	
Hazlehurst's Towers and Tanks for Water-works8vo,	370	50
Herschel's 115 Experiments on the Carrying Capacity of Large, Riveted, Metal	-	30
Conduits8vo,	2	00
Mason's Water-supply. (Considered Principally from a Sanitary Stand-		-
point.) 3d Edition, Rewritten		00
Merriman's Treatise on Hydraulics. oth Edition, Rewritten		
* Michie's Elements of Analytical Mechanics	-	
Schuyler's Reservoirs for Irrigation, Water-power, and Domestic Water-	4	00
supply	-	00
** Thomas and Watt's Improvement of Rivers. (Post., 44 c. additional), 4to,		
Turneaure and Russell's Public Water-supplies		
Wegmann's Design and Construction of Dams4to,		
Water-supply of the City of New York from 1658 to 1895		
Weisbach's Hydraulics and Hydraulic Motors. (Du Bois.)8vo,		
Wilson's Manual of Irrigation Engineering	4	00
Wilson's Manual of Irrigation EngineeringSmall 8vo. Wolff's Windmill as a Prime Mover8vo,	4 3	00
Wilson's Manual of Irrigation EngineeringSmall 8vo. Wolff's Windmill as a Prime Mover8vo, Wood's Turbines8vo,	4 3 2	00 00 50
Wilson's Manual of Irrigation EngineeringSmall 8vo. Wolff's Windmill as a Prime Mover8vo,	4 3 2	00 00 50
Wilson's Manual of Irrigation Engineering	4 3 2	00 00 50
Wilson's Manual of Irrigation EngineeringSmall 8vo. Wolff's Windmill as a Prime Mover8vo, Wood's Turbines8vo,	4 3 2	00 00 50
Wilson's Manual of Irrigation Engineering	4 3 2 3	00 00 50 00
Wilson's Manual of Irrigation Engineering	4 3 2 3 5	00 50 00
Wilson's Manual of Irrigation Engineering	4 3 2 3 5 5	00 50 00 00
Wilson's Manual of Irrigation Engineering	4 3 2 3 5 5 5	00 00 00 00
Wilson's Manual of Irrigation Engineering	4 3 2 3 5 5 5	00 50 00 00
Wilson's Manual of Irrigation Engineering	4 3 2 3 5 5 5 7	00 50 00 00 00 00 50
Wilson's Manual of Irrigation Engineering	4323 5557 7	00 50 00 00 00 50
Wilson's Manual of Irrigation Engineering	4323 5557 7	00 50 00 00 00 00 50
Wilson's Manual of Irrigation Engineering	4323 5557 75	00 50 00 00 00 50 50
Wilson's Manual of Irrigation Engineering	4323 5557 75 3	00 50 00 00 00 50 50 00
Wilson's Manual of Irrigation Engineering	4323 5557 75 36	00 50 00 00 00 50 50 00 00 00
Wilson's Manual of Irrigation Engineering	4323 5557 75 367	00 00 50 00 00 50 50 00 00 50
Wilson's Manual of Irrigation Engineering	4323 5557 75 3676	00 00 50 00 00 50 50 00 00 50
Wilson's Manual of Irrigation Engineering. Small 8vo. Wolff's Windmill as a Prime Mover. 8vo. Wood's Turbines. 8vo. Elements of Analytical Mechanics. 8vo. MATERIALS OF ENGINEERING. Baker's Treatise on Masonry Construction. 8vo. Roads and Pavements. 8vo. Black's United States Public Works. Oblong 4to. Bovey's Strength of Materials and Theory of Structures. 8vo. Burr's Elasticity and Resistance of the Materials of Engineering. 6th Edition, Rewritten. 8vo. Byrne's Highway Construction. 8vo. Inspection of the Materials and Workmanship Employed in Construction. Church's Mechanics of Engineering. Vol. I. Small 4to. Johnson's Materials of Construction. Large 8vo. Keep's Cast Iron. 8vo.	4323 5557 75 36762	00 50 00 00 00 50 00 00 50 00 50
Wilson's Manual of Irrigation Engineering. Small 8vo, Wolff's Windmill as a Prime Mover. 8vo, Wood's Turbines. 8vo, Elements of Analytical Mechanics. 8vo, MATERIALS OF ENGINEERING. Baker's Treatise on Masonry Construction. 8vo, Roads and Pavements. 8vo, Black's United States Public Works. Oblong 4to, Bovey's Strength of Materials and Theory of Structures. 8vo, Burr's Elasticity and Resistance of the Materials of Engineering. 6th Edition, Rewritten. 8vo, Byrne's Highway Construction. 8vo, Inspection of the Materials and Workmanship Employed in Construction. I6mo, Church's Mechanics of Engineering. Vol. I. Small 4to, Johnson's Materials of Construction. Large 8vo, Keep's Cast Iron. 8vo, Lanza's Applied Mechanics. 8vo,	4323 5557 75 367627	00 50 00 00 00 50 50 00 50 50 50 50
Wilson's Manual of Irrigation Engineering. Small 8vo. Wolff's Windmill as a Prime Mover. 8vo, Wood's Turbines. 8vo, Elements of Analytical Mechanics. 8vo, MATERIALS OF ENGINEERING. Baker's Treatise on Masonry Construction. 8vo, Roads and Pavements. 8vo, Black's United States Public Works. Oblong 4to, Bovey's Strength of Materials and Theory of Structures. 8vo, Burr's Elasticity and Resistance of the Materials of Engineering. 6th Edition, Rewritten 8vo, Inspection of the Materials and Workmanship Employed in Construction. Inspection of the Materials and Workmanship Employed in Construction. Inspection of the Materials and Workmanship Employed in Construction. Large 8vo, Du Bois's Mechanics of Engineering. Vol. I. Small 4to, Johnson's Materials of Construction Large 8vo, Keep's Cast Iron. 8vo, Lanza's Applied Mechanics. 8vo, Martens's Handbook on Testing Materials. (Henning.) 2 vols. 8vo,	4323 5557 75 3676277	00 00 50 00 00 50 50 00 00 50 50 50 50 5
Wilson's Manual of Irrigation Engineering	4323 5557 75 36762775	00 00 50 00 00 50 50 00 50 50 50 50 50 5
Wilson's Manual of Irrigation Engineering	4323 5557 75 367627754	00 00 00 00 00 50 00 00 50 00 50 50 50 5
Wilson's Manual of Irrigation Engineering	4323 5557 75 3676277541	00 00 50 00 00 50 50 00 50 50 50 50 50 5
Wilson's Manual of Irrigation Engineering	4323 5557 75 36762775412	00 00 00 00 00 50 50 00 00 50 50 50 00 0

Rockwell's Roads and Pavements in France 12mo, 1 25 Smith's Materials of Machines 12mo, 1 00 Snow's Principal Species of Wood 8vo, 3 50 Spalding's Hydraulic Cement 12mo, 2 00 Text-book on Roads and Pavements 12mo, 2 00 Thurston's Materials of Engineering 3 Parts 8vo, 8 00 Part I.—Non-metallic Materials of Engineering and Metallurgy 8vo, 2 00 Part II.—Iron and Steel 8vo, 3 50 Part III.—A Treatise on Brasses, Bronzes, and Other Alloys and their Constituents 8vo, 2 50 Thurston's Text-book of the Materials of Construction 8vo, 5 00 Tillson's Street Pavements and Paving Materials 8vo, 4 00 Waddell's De Pontibus. (A Pocket-book for Bridge Engineers.) 16mo, mor., 3 00
Specifications for Steel Bridges. 12mo, 1 25 Wood's Treatise on the Resistance of Materials, and an Appendix on the Preservation of Timber. 8vo, 2 00 Elements of Analytical Mechanics 8vo, 3 00 Wood's Rustless Coatings: Corrosion and Electrolysis of Iron and Steel 8vo, 4 00
RAILWAY ENGINEERING.
Andrews's Handbook for Street Railway Engineers. 3×5 inches. morocco, 1 25 Berg's Buildings and Structures of American Railroads
Railway and Other Earthwork Tables
Fisher's Table of Cubic Yards
bankments
Searles's Field Engineering
Embankments by the Aid of Diagrams
Webb's Railroad Construction. 2d Edition, Rewritten16mo. morocco, 5 oo Wellington's Economic Theory of the Location of RailwaysSmall 8vo, 5 oo
DRAWING.
Barr's Kinematics of Machinery. 8vo, 2 50 Bartlett's Mechanical Drawing. 8vo, 3 00 " " Abridged Ed. 8vo, 1 50 Coolidge's Manual of Drawing. 8vo, paper, 1 00 Coolidge and Freeman's Elements of General Drafting for Mechanical Engi-
neers. (In press.) Durley's Kinematics of Machines
8 The state of the

Hill's Text-book on Shades and Shadows, and Perspective8vo,	2	00
Jamison's Elements of Mechanical Drawing. (In press.)		
Jones's Machine Design:		
Part I.—Kinematics of Machinery8vo,	I	50
Part II.—Form, Strength, and Proportions of Parts8vo,	3	00
MacCord's Elements of Descriptive Geometry8vo,	3	00
Kinematics; or, Practical Mechanism8vo,	5	00
Mechanical Drawing4to,	4	00
Velocity Diagrams8vo,	I	50
Mahan's Descriptive Geometry and Stone-cutting8vo,		
Industrial Drawing. (Thompson.)8vo,		
Reed's Topographical Drawing and Sketching		
Reid's Course in Mechanical Drawing8vo,		
Text-book of Mechanical Drawing and Elementary Machine Design . 8vo.		
Robinson's Principles of Mechanism		
Smith's Manual of Topographical Drawing. (McMillan.)8vo,		
Warren's Elements of Plane and Solid Free-hand Geometrical Drawing . 12mo,		
Drafting Instruments and Operations		
Manual of Elementary Projection Drawing		50
Manual of Elementary Problems in the Linear Perspective of Form and		100
Shadow12mo,		
Plane Problems in Elementary Geometry	I	
Primary Geometry		75
Elements of Descriptive Geometry, Shadows, and Perspective8vo,	-	
General Problems of Shades and Shadows8vo,	3	00
	7	50
Problems, Theorems, and Examples in Descriptive Geometry8vo,	2	50
Weisbach's Kinematics and the Power of Transmission. (Hermann and		
Klein.)8vo,	5	00
Whelpley's Practical Instruction in the Art of Letter Engraving12mo,	2	00
Wilson's Topographic Surveying8vo,	3	50
Free-hand Perspective8vo,	2	50
Free-hand Lettering	1	00
Woolf's Elementary Course in Descriptive GeometryLarge 8vo,	3	00
too a compared that the control of t		
ELECTRICITY AND PHYSICS.		
ELECTRICITI AND PHISICS.		
Anthony and Brackett's Text-book of Physics. (Magie.)Small 8vo.	3	00
Anthony's Lecture-notes on the Theory of Electrical Measurements12mo,	7	00
Benjamin's History of Electricity8vo,	-	00
Voltaic Cell8vo,		
Classen's Quantitative Chemical Analysis by Electrolysis. (Boltwood.)8vo,	3	00
Crehore and Squier's Polarizing Photo-chronograph8vo,	3	00
Dawson's "Engineering" and Electric Traction Pocket-book . 16mo, morocco,	3	00
T. 1 . 1 . 1	5	00
	335	
Ende.)		
	4	00
Flather's Dynamometers, and the Measurement of Power12mo,	3	00
Gilbert's De Magnete. (Mottelay.)8vo,	2	50
Hanchett's Alternating Currents Explained	I	00
Hering's Ready Reference Tables (Conversion Factors)16mo, morocco,	2	50
Holman's Precision of Measurements8vo,	2	00
Telescopic Mirror-scale Method, Adjustments, and TestsLarge 8vo,		75
	3	00
Le Chatelier's High-temperature Measurements. (Boudouard-Burgess.)12mo,	M.	00
Löb's Electrolysis and Electrosynthesis of Organic Compounds. (Lorenz.) 12mo.		00
Lyons's Treatise on Electromagnetic Phenomena. Vols. I. and II. 8vo, each.	6	00
* Michie. Elements of Wave Motion Relating to Sound and Light, 8vo.	4	00

Niaudet's Elementary Treatise on Electric Batteries. (Fishback.)12mo,	2	50
Rosenberg's Electrical Engineering. (Haldane Gee-Kinzbrunner.)8vo,		
Ryan, Norris, and Hoxie's Electrical Machinery. Vol. I	2	50
Thurston's Stationary Steam-engines8vo,		
* Tillman's Elementary Lessons in Heat8vo,	I	50
Tory and Pitcher's Manual of Laboratory Physics Small 8vo,		
Ulke's Modern Electrolytic Copper Refining8vo,	3	00
on a contract of the contract and State of the contract of the contract of the		
LAW.		
* Davis's Elements of Law8vo,	2	50
* Treatise on the Military Law of United States8vo,		00
* Sheep,		
Manual for Courts-martial	í	50
Wait's Engineering and Architectural Jurisprudence8vo,		
		50
Law of Operations Preliminary to Construction in Engineering and Archi-	Ē	-
tecture 8vo,	5	00
Sheep,		
Law of Contracts		
Winthrop's Abridgment of Military Law12mo,		
The second secon		
MANUFACTURES.		
MANUFACIURES.		
Bernadou's Smokeless Powder-Nitro-cellulose and Theory of the Cellulose		
Molecule12mo,	2	50
Bolland's Iron Founder		
"The Iron Founder," Supplement12mo,		
Encyclopedia of Founding and Dictionary of Foundry Terms Used in the		1187
Practice of Moulding	3	00
Eissler's Modern High Explosives8vo,	4	00
Effront's Enzymes and their Applications. (Prescott.)8vo,	3	00
Fitzgerald's Boston Machinist18mo,	I	00
Ford's Boiler Making for Boiler Makers18mo,	1	00
Hopkins's Oil-chemists' Handbook8vo,	3	00
Keep's Cast Iron8vo,	2	50
Leach's The Inspection and Analysis of Food with Special Reference to State		
Control. (In preparation.)		
Metcalf's Steel. A Manual for Steel-users	2	00
Metcalfe's Cost of Manufactures-And the Administration of Workshops,		
Public and Private8vo,		
Meyer's Modern Locomotive Construction4to, 1		
Morse's Calculations used in Cane-sugar Factories 16mo, morocco,		-
* Reisig's Guide to Piece-dyeing		
Smith's Press-working of Metals	-	
Spalding's Hydraulic Cement		
Spencer's Handbook for Chemists of Beet-sugar Houses16mo, morocco,	-	
Handbook for Sugar Manufacturers and their Chemists 16mo, morocco,	2	00
Thurston's Manual of Steam-boilers, their Designs, Construction and Opera-	_	00
tion		
West's American Foundry Practice		
Moulder's Text-book		-
Wiechmann's Sugar Analysis		
Wolff's Windmill as a Prime Mover		
Woodbury's Fire Protection of Mills		
Wood's Rustless Coatings: Corrosion and Electrolysis of Iron and Steel 8vo.		

MATHEMATICS.

Bass's Elements of Differential Calculus		
Briggs's Elements of Plane Analytic Geometry		0
Compton's Manual of Logarithmic Computations	0	0
Dayis's Introduction to the Logic of Algebra		0
* Dickson's College Algebra Large 12mo, 1 * Answers to Dickson's College Algebra 8vo, paper,	-	
* Answers to Dickson's College Algebra	5	0
	5	0
	2	5
Introduction to the Theory of Algebraic Equations Large 12mo, 1		=
Halsted's Elements of Geometry8vo, x	7	5
Elementary Synthetic Geometry		
	7	5
• Johnson's Three-place Logarithmic Tables: Vest-pocket sizepaper,	I;	-
	0	0
Mounted on heavy cardboard, 8 × 10 inches,	2	5
	0	0
Elementary Treatise on the Integral Calculus Small 8vo, 1	5	0
	00	D
Treatise on Ordinary and Partial Differential EquationsSmall 8vo, 3	5	0
	5	0
	0	0
	0	0
* Ludlow and Bass. Elements of Trigonometry and Logarithmic and Other		
Tables8vo, 3	0	0
	0	0
	0	0
	1 0	0
	0	0
Merriman's Method of Least Squares8vo, 2	0	0
Rice and Johnson's Elementary Treatise on the Differential Calculus . Sm., 8vo, 3	0	0
Differential and Integral Calculus. 2 vols. in oneSmall 8vo. 2	5	0
Sabin's Industrial and Artistic Technology of Paints and Varnish. (In press.)		
Wood's Elements of Co-ordinate Geometry8vo, 2	0	0
Trigonometry: Analytical, Plane, and Spherical12mo, 1	0	0
MECHANICAL ENGINEERING.		
MATERIALS OF ENGINEERING, STEAM-ENGINES AND BOILERS.		
MATERIALS OF ENGINEERING, STEAM-ENGINES AND BOILERS.		
Baldwin's Steam Heating for Buildings		
Baldwin's Steam Heating for Buildings	5	0
Baldwin's Steam Heating for Buildings	5 0	0
Baldwin's Steam Heating for Buildings 12mo, 2 Barr's Kinematics of Machinery 8vo, 2 * Bartlett's Mechanical Drawing 8vo, 3 * " " Abridged Ed 8vo, 1	5 5	0
Baldwin's Steam Heating for Buildings 12mo, 2 Barr's Kinematics of Machinery 8vo, 2 * Bartlett's Mechanical Drawing 8vo, 3 * " " Abridged Ed 8vo, 1 Benjamin's Wrinkles and Recipes 12mo, 2	5 5 6 6	0 0 0
Baldwin's Steam Heating for Buildings 12mo, 2 Barr's Kinematics of Machinery 8vo, 2 * Bartlett's Mechanical Drawing 8vo, 3 * " " Abridged Ed 8vo, 1 Benjamin's Wrinkles and Recipes 12mo, 2 Carpenter's Experimental Engineering 8vo, 6	5 5 0	0 0 0 0
Baldwin's Steam Heating for Buildings 12mo, 2 Barr's Kinematics of Machinery 8vo, 2 * Bartlett's Mechanical Drawing 8vo, 3 * " " Abridged Ed 8vo, 1 Benjamin's Wrinkles and Recipes 12mo, 2 Carpenter's Experimental Engineering 8vo, 6 Heating and Ventilating Buildings 8vo, 4	5 5 0	0 0 0 0
Baldwin's Steam Heating for Buildings	5 5 0	0 0 0 0
Baldwin's Steam Heating for Buildings	5 5 0 0 0	00000
Baldwin's Steam Heating for Buildings	5 5 0 0 0	00000
Baldwin's Steam Heating for Buildings	5 5 0 0 0	00000
Baldwin's Steam Heating for Buildings	5 5 0 0 0	00000
Baldwin's Steam Heating for Buildings	5 5 0 0 0 0 0 0 0	00000
Baldwin's Steam Heating for Buildings	5 5 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	00000
Baldwin's Steam Heating for Buildings	5 5 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	00000
Baldwin's Steam Heating for Buildings	5 5 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	000000000000000000000000000000000000000

Gill's Gas and Fuel Analysis for Engineers	I	25
Hall's Car Lubrication12mo,	I	00
Hering's Ready Reference Tables (Conversion Factors)16mo, morocco,	2	50
Hutton's The Gas Engine8vo,	5	00
Jones's Machine Design:		
Part I.—Kinematics of Machinery8vo,		
Part II.—Form, Strength, and Proportions of Parts8vo,		
Kent's Mechanical Engineer's Pocket-book 16mo, morocco,		00
Kerr's Power and Power Transmission8vo,	2	00
MacCord's Kinematics; or, Practical Mechanism8vo,	5	00
	-	00
Velocity Diagrams	1	50
Mahan's Industrial Drawing. (Thompson.)8vo,	3	50
Poole's Calorific Power of Fuels		00
Reid's Course in Mechanical Drawing8vo.	2	00
Text-book of Mechanical Drawing and Elementary Machine Design 8vo,	3	00
	I	50
Robinson's Principles of Mechanism8vo,	3	00
	3	00
Thurston's Treatise on Friction and Lost Work in Machinery and Mill		
Work8vo,	3	00
Animal as a Machine and Prime Motor, and the Laws of Energetics. 12mo,	1	00
Warren's Elements of Machine Construction and Drawing8vo,	7	50
Weisbach's Kinematics and the Power of Transmission. Herrmann-		
Klein.)8vo,		
Machinery of Transmission and Governors. (Herrmann-Klein.)8vo,		
Hydraulics and Hydraulic Motors. (Du Bois.)8vo,		
Wolff's Windmill as a Prime Mover		
Wood's Turbines8vo,	2	50
TOOK & ALTONICO.		-
		-34
MATERIALS OF ENGINEERING.		Mo
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures		50
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures		
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7	
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 7	50
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 76	50
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 766	50 50 00
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 7662	50 50 00
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 7666277	50 50 00 50 50 50
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 7662774	50 00 00 50 50 50
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 76627741	50 00 00 50 50 50 00
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 7666277412	50 00 00 50 50 00 00 00
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 7662774121	50 00 00 50 50 00 00 00
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 76627741218	50 00 00 50 50 50 00 00 00 00
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 76627741218	50 00 00 50 50 50 00 00 00 00
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 766277412183	50 00 00 50 50 00 00 00 00 50
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 766 2 7 7 4 1 2 1 8 3 2	50 00 00 50 50 50 00 00 00 00 50
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 766 2 7 7 4 1 2 1 8 3 2	50 00 00 50 50 00 00 00 00 50
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 766277412183	50 00 00 50 50 50 00 00 00 50 50
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 766277412183 25 2	50 00 00 50 50 50 00 00 00 50 50 00 00 0
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 766277412183 25 23	50 00 00 50 50 50 00 00 00 50 50 00 00 0
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 766277412183 25 23	50 00 00 50 50 50 00 00 00 50 50 00 00 0
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 766277412183 25 23	50 00 00 50 50 50 00 00 00 50 50 00 00 0
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 7666 2 7 7 4 1 2 1 8 3 2 5 2 3 4	50 00 00 50 50 50 00 00 00 50 50 00 00 0
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 766 2 7 7 4 1 2 1 8 3 2 5 2 3 4	50 00 00 50 50 50 00 00 00 50 50 50 00 0
MATERIALS OF ENGINEERING. Bovey's Strength of Materials and Theory of Structures	7 766277412183 25 234	50 00 00 50 50 50 00 00 00 00 00 00 00 0

Hemenway's Indicator Practice and Steam-engine Economy	-	-
		00
	100	00
Heat and Heat-engines8vo,	5	00
Kent's Steam-boiler Economy8vo,	-	00
Kneass's Practice and Theory of the Injector	1	50
MacCord's Slide-valves8vo,		00
	IO	00
Peabody's Manual of the Steam-engine Indicator		50
Tables of the Properties of Saturated Steam and Other Vapors8vo,		00
	150	130
Thermodynamics of the Steam-engine and Other Heat-engines8vo,	-	00
Valve-gears for Steam-engines8vo,	2	50
Peabody and Miller's Steam-boilers	4	00
Pray's Twenty Years with the Indicator Large 8vo,	2	50
Pupln's Thermodynamics of Reversible Cycles in Gases and Saturated Vapors.		
(Osterberg.)12mo,		25
Reagan's Locomotives: Simple, Compound, and Electric		
		17 10
Rontgen's Principles of Thermodynamics. (Du Bois.)8vo,	-	00
Sinclair's Locomotive Engine Running and Management	2	00
Smart's Handbook of Engineering Laboratory Practice	2	50
Snow's Steam-boiler Practice8vo,	3	00
Spangler's Valve-gears8vo,	2	50
		00
Spangler, Greene, and Marshall's Elements of Steam-engineering8vo,	3	00
	- C	
	1	50
Manual of the Steam-engine 2 vols. 8vo, 1	0	00
	6	00
Part II,—Design, Construction, and Operation8vo,	6	00
Handbook of Engine and Boiler Trials, and the Use of the Indicator and		
the Prony Brake8vo	=	00
Stationary Steam-engines8vo,	E.	50
Steam-boiler Explosions in Theory and in Practice12mo		-
		50
Manual of Steam-boilers, Their Designs, Construction, and Operation . 8vo,	5	00
	5	00
	5	00
Wilson's Treatise on Steam-boilers. (Flather.)		
	2	50
		50 00
Wood's Thermodynamics Heat Motors, and Refrigerating Machines8vo,		
Wood's Thermodynamics Heat Motors, and Refrigerating Machines8vo, MECHANICS AND MACHINERY.	4	00
Wood's Thermodynamics Heat Motors, and Refrigerating Machines8vo, MECHANICS AND MACHINERY.	4	00
Wood's Thermodynamics Heat Motors, and Refrigerating Machines8vo, MECHANICS AND MACHINERY. Barr's Kinematics of Machinery8vo,	4	50
Wood's Thermodynamics Heat Motors, and Refrigerating Machines8vo, MECHANICS AND MACHINERY. Barr's Kinematics of Machinery	2 7	50 56
Wood's Thermodynamics Heat Motors, and Refrigerating Machines8vo, MECHANICS AND MACHINERY. Barr's Kinematics of Machinery	2 7 2	50 50 50
Wood's Thermodynamics Heat Motors, and Refrigerating Machines8vo, MECHANICS AND MACHINERY. Barr's Kinematics of Machinery	2 7 2 2	50 50 50 00
Wood's Thermodynamics Heat Motors, and Refrigerating Machines8vo, MECHANICS AND MACHINERY. Barr's Kinematics of Machinery	2 7 2 2	50 50 50
Wood's Thermodynamics Heat Motors, and Refrigerating Machines8vo, MECHANICS AND MACHINERY. Barr's Kinematics of Machinery	2 7 2 2 6	50 50 50 00
Wood's Thermodynamics Heat Motors, and Refrigerating Machines8vo, MECHANICS AND MACHINERY. Barr's Kinematics of Machinery	2 7 2 6 2	50 50 50 00
Wood's Thermodynamics Heat Motors, and Refrigerating Machines8vo, MECHANICS AND MACHINERY. Barr's Kinematics of Machinery	2 7 2 6 2 1	50 50 50 00 00
MECHANICS AND MACHINERY. Barr's Kinematics of Machinery	2 7 2 6 2 1 1	50 50 50 00 00 50
MECHANICS AND MACHINERY. Barr's Kinematics of Machinery 8vo, Bovey's Strength of Materials and Theory of Structures 8vo, Chase's The Art of Pattern-making 12mo, Chordal.—Extracts from Letters 12mo, Church's Mechanics of Engineering 8vo, Notes and Examples in Mechanics 8vo, Compton's First Lessons in Metal-working 12mo, Compton and De Groodt's The Speed Lathe 12mo, Cromwell's Treatise on Toothed Gearing 12mo,	2 7 2 6 2 1 1	50 50 50 00 00 50 50 50
MECHANICS AND MACHINERY. Barr's Kinematics of Machinery 8vo, Bovey's Strength of Materials and Theory of Structures 8vo, Chase's The Art of Pattern-making 12mo, Chordal.—Extracts from Letters 12mo, Church's Mechanics of Engineering 8vo, Notes and Examples in Mechanics 8vo, Compton's First Lessons in Metal-working 12mo, Compton and De Groodt's The Speed Lathe 12mo, Cromwell's Treatise on Toothed Gearing 12mo, Treatise on Belts and Pulleys 12mo,	2 7 2 6 2 1 1	50 50 50 00 00 50 50
MECHANICS AND MACHINERY. Barr's Kinematics of Machinery	2 7 2 6 2 1 1 1	50 50 50 00 00 50 50 50 50 50
MECHANICS AND MACHINERY. Barr's Kinematics of Machinery 8vo, Bovey's Strength of Materials and Theory of Structures 8vo, Chase's The Art of Pattern-making 12mo, Chordal.—Extracts from Letters 12mo, Church's Mechanics of Engineering 8vo, Notes and Examples in Mechanics 8vo, Compton's First Lessons in Metal-working 12mo, Compton and De Groodt's The Speed Lathe 12mo, Cromwell's Treatise on Toothed Gearing 12mo, Treatise on Belts and Pulleys 12mo, Dana's Text-book of Elementary Mechanics for the Use of Colleges and Schools 12mo,	2 7 2 6 2 1 1 1	50 50 50 00 00 50 50 50 50 50
MECHANICS AND MACHINERY. Barr's Kinematics of Machinery 8vo, Bovey's Strength of Materials and Theory of Structures 8vo, Chase's The Art of Pattern-making 12mo, Chordal.—Extracts from Letters 12mo, Church's Mechanics of Engineering 8vo, Notes and Examples in Mechanics 8vo, Compton's First Lessons in Metal-working 12mo, Compton and De Groodt's The Speed Lathe 12mo, Cromwell's Treatise on Toothed Gearing 12mo, Treatise on Belts and Pulleys 12mo, Dana's Text-book of Elementary Mechanics for the Use of Colleges and Schools 12mo, Dingey's Machinery Pattern Making 12mo,	2 7 2 6 2 1 1 1	50 50 50 00 00 50 50 50 50 50
MECHANICS AND MACHINERY. Barr's Kinematics of Machinery 8vo, Bovey's Strength of Materials and Theory of Structures 8vo, Chase's The Art of Pattern-making 12mo, Chordal.—Extracts from Letters 12mo, Church's Mechanics of Engineering 8vo, Notes and Examples in Mechanics 8vo, Compton's First Lessons in Metal-working 12mo, Compton and De Groodt's The Speed Lathe 12mo, Cromwell's Treatise on Toothed Gearing 12mo, Treatise on Belts and Pulleys 12mo, Dana's Text-book of Elementary Mechanics for the Use of Colleges and Schools 12mo,	2 7 2 6 2 1 1 1	50 50 50 00 00 50 50 50 50 50

Gerhard's Guide to Sanitary House-inspection	1	00
Goodrich's Economical Disposal of Town's Refuse Demy 8vo,	3	50
Hazen's Filtration of Public Water-supplies	3	00
Kiersted's Sewage Disposal12mo,	I	25
Leach's The Inspection and Analysis of Food with Special Reference to State	ñ	-
Control. (In preparation.)		
Mason's Water-supply. (Considered Principally from a Sanitary Stand-		
point.) 3d Edition, Rewritten8vo,		00
Examination of Water. (Chemical and Bacteriological.)12mo,		25
Merriman's Elements of Sanitary Engineering 8vo,		00
Nichols's Water-supply. (Considered Mainly from a Chemical and Sanitary	-	00
Standpoint.) (1883.)		
Ogden's Sewer Design		50
Prescott and Winslow's Elements of Water Bacteriology, with Special Reference	2	00
to Sanitary Water Analysis		
* Price's Handbook on Sanitation		25
Richards's Cost of Food. A Study in Dietaries		50
Cost of Living on Medified by Conitors Calendary		00
Cost of Living as Modified by Sanitary Science	1	00
Richards and Woodman's Air, Water, and Food from a Sanitary Stand-		
point	15	00
* Richards and Williams's The Dietary Computer		50
Rideal's Sewage and Bacterial Purification of Sewage		50
Turneaure and Russell's Public Water-supplies8vo,	100	00
	-	50
Woodhull's Notes and Military Hygiene16mo,	1	50
MICOPET ANDOMO		
MISCELLANEOUS.		
	TO	
Barker's Deep-sea Soundings8vo,	2	00
Barker's Deep-sea Soundings		
Barker's Deep-sea Soundings	1	50
Barker's Deep-sea Soundings	1 4	50
Barker's Deep-sea Soundings	1 4 2	50 00 50
Barker's Deep-sea Soundings	1 4 2 1	50 00 50 25
Barker's Deep-sea Soundings	1 4 2 1	50 00 50 25 00
Barker's Deep-sea Soundings	1 4 2 1 1 3	50 00 50 25 00
Barker's Deep-sea Soundings	1 4 2 1 1 3	50 00 50 25 00
Barker's Deep-sea Soundings	I 4 2 I I 3 2	50 00 50 25 00
Barker's Deep-sea Soundings	I 4 2 I I 3 2 3 2	50 00 50 25 00 00 00 50 50
Barker's Deep-sea Soundings	I 4 2 I I 3 2 3 2	50 00 50 25 00 00 00 50 50
Barker's Deep-sea Soundings	I 4 2 I I 3 2 3 2	50 00 50 25 00 00 00 50 50
Barker's Deep-sea Soundings	I 4 2 I I 3 2 3 2	50 00 50 25 00 00 00 50 50
Barker's Deep-sea Soundings	1 4 2 1 1 3 2 3 2 I	50 00 50 25 00 00 50 50 50
Barker's Deep-sea Soundings	1 4 2 1 1 3 2 3 2 I	50 00 50 25 00 00 50 50 50
Barker's Deep-sea Soundings	1 4 2 1 1 3 2 3 2 I	50 00 50 25 00 00 50 50 50
Barker's Deep-sea Soundings	1 4 2 1 1 3 2 3 2 I	50 00 50 25 00 00 50 50 50
Barker's Deep-sea Soundings	1 4 2 1 1 3 2 3 2 I	50 00 50 25 00 00 50 50 50 00
Barker's Deep-sea Soundings	1 4 2 1 1 3 2 3 2 1 t	50 00 50 25 00 00 50 50 50 25
Barker's Deep-sea Soundings. 8vo, Emmons's Geological Guide-book of the Rocky Mountain Excursion of the International Congress of Geologists. Large 8vo Ferrel's Popular Treatise on the Winds. 8vo Haines's American Railway Management. 12mo, Mott's Composition, Digestibility, and Nutritive Value of Food. Mounted chart. Fallacy of the Present Theory of Sound. 16mo Ricketts's History of Rensselaer Polytechnic Institute, 1824-1894. Small 8vo, Rotherham's Emphasized New Testament. Large 8vo, Steel's Treatise on the Diseases of the Dog. 8vo, Totten's Important Question in Metrology. 8vo The World's Columbian Exposition of 1893. 4to, Worcester and Atkinson. Small Hospitals, Establishment and Maintenance, and Suggestions for Hospital Architecture, with Plans for a Small Hospital. 12mo, HEBREW AND CHALDEE TEXT-BOOKS.	1 4 2 1 1 3 2 3 2 1 t	50 00 50 25 00 00 50 50 50 25
Barker's Deep-sea Soundings. 8vo, Emmons's Geological Guide-book of the Rocky Mountain Excursion of the International Congress of Geologists. Large 8vo Ferrel's Popular Treatise on the Winds. 8vo Haines's American Railway Management. 12mo, Mott's Composition, Digestibility, and Nutritive Value of Food. Mounted chart. Fallacy of the Present Theory of Sound. 16mo Ricketts's History of Rensselaer Polytechnic Institute, 1824-1894. Small 8vo, Rotherham's Emphasized New Testament. Large 8vo, Steel's Treatise on the Diseases of the Dog. 8vo, Totten's Important Question in Metrology. 8vo The World's Columbian Exposition of 1893. 4to, Worcester and Atkinson. Small Hospitals, Establishment and Maintenance, and Suggestions for Hospital Architecture, with Plans for a Small Hospital. 12mo, HEBREW AND CHALDEE TEXT-BOOKS. Green's Grammar of the Hebrew Language. 8vo, Elementary Hebrew Grammar 12mo, Hebrew Chrestomathy 8vo,	1 4 2 1 1 3 2 3 2 1 t	50 00 50 25 00 00 50 50 50 25
Barker's Deep-sea Soundings	1 4 2 1 1 3 2 3 2 1 1 3 1 2	50 00 50 25 00 00 50 50 00 25 00 00 25
Barker's Deep-sea Soundings	1 4 2 1 1 3 2 3 2 1 1 3 1 2 5	50 00 50 25 00 00 50 50 00 25 00 00 25 00
Barker's Deep-sea Soundings	1 4 2 1 1 3 2 3 2 1 1 3 1 2 5	50 00 50 25 00 00 50 50 00 25 00 25 00

